



(51) International Patent Classification:

A61K 36/14 (2006.01) C07C 61/35 (2006.01)
C07C 49/755 (2006.01)

(21) International Application Number:

PCT/US2020/052545

(22) International Filing Date:

24 September 2020 (24.09.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/905,253 24 September 2019 (24.09.2019) US

(71) Applicant: **K-GEN, INC.** [US/US]; 863 Mitten Road, Burlingame, California 94010 (US).

(72) Inventors: **JIANG, Chun**; 863 Mitten Road, Burlingame, California 94010 (US). **CHEN, Ruihong**; 863 Mitten Road, Burlingame, California 94010 (US).

(74) Agent: **OH, Euk Charlie** et al.; DECHERT LLP, Three Bryant Park, 1095 Avenue of the Americas, New York, New York 10036-6797 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available):

AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available):

ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: DITERPENOID COMPOUNDS THAT ACT ON PROTEIN KINASE C (PKC)




FIG. 1

(57) Abstract: This present disclosure relates to protein kinase C (PKC) modulating compounds, methods of treating a subject with cancer using the compounds, and combination treatments with a second therapeutic agent.



Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

DITERPENOID COMPOUNDS THAT ACT ON PROTEIN KINASE C (PKC)

1. CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application No. 62/905,253, filed September 24, 2019, the contents of which are incorporated herein in their entireties by reference thereto.

2. BACKGROUND


[0002] Diterpenoid Protein Kinase C (PKC) modulating compounds display anti-cancer and cytotoxic activities. Most studied of these compounds are tricyclic diterpenoids, such as phorbol esters and prostratin. The biological effects of these compounds are thought to be mediated by transactivation, translocation and suppression of PKC enzymes, which play important roles in regulating signaling pathways that regulate or modulate cellular structure and gene expression.

[0003] Association of PKC enzyme activation and its inhibitory effect on cancer cell growth is supported by studies of PKC mutations in human cancers, which found that most of the PKC mutations are loss-of function mutations (Antal et al., 2015, Cell 160:489–502). This presence of PKC loss-of-function mutations in various cancer types suggests that PKC enzymes may act as tumor suppressors. Studies with prostratin suggest that its anti-tumor effect occurs by activation of PKC enzymes that specifically target oncogene K-RAS (see Wang et al., 2015, Cell 163(5):1237–1251). A different tricyclic diterpenoid compound, phorbol myristate 13-acetate (PMA) also displays inhibitory effects on tumor growth by its action on PKC enzymes but also non-PKC involved mechanisms (Bond et al., 2007, Int. J. Cancer 121:1445-1454).

[0004] In view of the therapeutic potential of diterpenoid PKC modulating compounds, it is desirable to further develop these class of compounds in applications for treating diseases and disorders capable of being affected by modulation of PKC activity.

3. SUMMARY

[0005] The present disclosure provides novel protein kinase C (PKC) modulating compounds. The compounds are diterpenoid PKC modulating compounds displaying enhanced pharmacological properties, including among others, potent PKC enzyme binding activity and anti-proliferative activity against different cancer cells. The PKC modulating compounds also have substituents that can enhance solubility and pharmacokinetic profiles. In one aspect, the compounds have the structure of formula (I):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

A is -OH, -C(O)OR¹, or -NR¹³R^{13'};

R¹ is H or a M⁺ counterion;

R² is a C₁-C₄alkyl;


R³ is O double bonded to the ring carbon when (- - -) is a bond, or -OR^a; wherein R^a is H or -C(O)R^{a1}, wherein R^{a1} is C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R⁴ and R⁵ are each independently H or -OR^b, wherein R^b is H, C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R^{5'} and R^{6'} are H, or R^{5'} and R^{6'} form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R⁶ is OH, halo, or -OC(O)R^c, wherein R^c is -C₁-C₆alkyl, -C₁-C₆alkyl-(NR^{cl})₂ or -C₁-C₆alkylC(O)OR^k, R^{cl} is H, C₁-C₆alkyl, or two R^{cl} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R⁶ is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino

acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^{6'} and R^{7'} are H, or R^{6'} and R^{7'} form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R⁷ is H or OH;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹² is H, -OH, -OC(O)R^f, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;

each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;


R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

n is 0 or 1.

[0006] In some embodiments, A is -OH. In some embodiments, A is -C(O)OR¹, wherein R¹ is H or a M⁺ counterion. In some embodiments, A is -NR¹³R^{13'}, wherein R¹³ and R^{13'} are each independently H or C₁-C₄alkyl.

[0007] In some embodiments, the present disclosure provides a compound of formula (II):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^2 is a C_1 - C_4 alkyl;


R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

$R^{5'}$ and $R^{6'}$ are H, or $R^{5'}$ and $R^{6'}$ form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

$R^{6'}$ and $R^{7'}$ are H, or $R^{6'}$ and $R^{7'}$ form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R^7 is H or OH;

R^9 is OR^e , wherein R^e is H, C₁-C₆alkyl, or aryl;

R^{11} is C₁-C₄alkyl;

R^{12} is H, -OH, $-OC(O)R^f$, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R^{13} and $R^{13'}$ are each independently H or C₁-C₄alkyl;

R^{14} is H or OR^g ; wherein R^g is H or C₁-C₆alkyl;

R^{17} and R^{18} are each independently C₁-C₄alkyl or C₁-C₄alkyl- OR^h , wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R^{21} is H, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, 5 to 12 membered bridged bicycyl, adamantyl, or $-C(O)OR^k$, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, bridged bicycyl, or adamantyl is optionally substituted with 1 to 3 of J^1 , and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J^1 ;

R^k is H or M^+ counterion;

J^1 is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

n is 0 or 1.

[0008] In some embodiments, the compound has the structure of formula (IIc):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1-C_6$ alkyl, $-C_1-C_6$ alkyl- $(NR^{c1})_2$ or $-C_1-C_6$ alkyl $C(O)OR^k$, R^{c1} is H, C_1-C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^{12} is H, $-OH$, $-OC(O)R^f$, wherein R^f is C_1-C_{12} alkyl, C_2-C_{12} alkenyl, $-C_0-C_{12}$ aliphatic- C_3-C_7 cycloalkyl, $-C_0-C_{12}$ aliphatic-heterocycloalkyl, $-C_0-C_{12}$ aliphatic-aryl, or $-C_0-C_{12}$ aliphatic-heteroaryl;

R^{13} and $R^{13'}$ are each independently H or C_1-C_4 alkyl;

L is absent, C_1-C_{12} alkylene, or C_2-C_{12} alkenylene, wherein the C_1-C_{12} alkylene or C_2-C_{12} alkenylene is optionally substituted with C_1-C_4 alkyl;

R^{21} is H, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or $-C(O)OR^k$, wherein the C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J^1 , and wherein optionally 1 to 2 carbon atoms of the spiro C_5-

C₁₂cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each Rⁱ is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

n is 0 or 1.

[0009] In another aspect, the present disclosure provides a compound of formula (IV):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R² is a C₁-C₄alkyl;


R³ is O double bonded to the ring carbon when (---) is a bond, or -OR^a, wherein R^a is H or -C(O)R^{a1}, wherein R^{a1} is C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R⁴ and R⁵ are each independently H or -OR^b, wherein R^b is H, C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R^{5'} and R^{6'} are H, or R^{5'} and R^{6'} form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R⁶ is OH, halo, or -OC(O)R^c, wherein R^c is -C₁-C₆alkyl, -C₁-C₆alkyl-(NR^{c1})₂ or -C₁-C₆alkylC(O)OR^k, R^{c1} is H, C₁-C₆alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R⁶ is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R⁶, and R⁷, are H, or R⁶ and R⁷ form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R⁷ is H or OH;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹² is H, -OH, -OC(O)R^f, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is C₀-C₆alkylarylene, C₀-C₆alkylheteroarylene, C₀-C₆alkylC₃-C₇cycloalkylene, C₁-C₁₂alkylene or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with OH or C₁-C₄alkyl; and

R²¹ is H, -OH, -SH, -S(O)₂Rⁱ, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, 5 to 12 membered bridged bicycyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, bridged bicycyl, or adamantyl is optionally substituted with 1 to 3 of J¹; and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl or, when an N atom is present an N-protecting group;

each Rⁱ is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylarylyl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylarylyl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

R^k is H or M⁺ counterion.

[0010] In some embodiments, the compounds can be used in a method to activate protein kinase C, comprising contacting a mammalian cell with an effective amount of a compound disclosed herein.

In some embodiments, the mammalian cell is a cancer cell.

[0011] In some embodiments, the compounds are used for treating cancer by administering a therapeutically effective amount to a subject in need thereof a compound disclosed herein.

[0012] In some embodiments, the cancer for treatment is adrenocortical cancer, anal cancer, biliary cancer, bladder cancer, bone cancer, brain cancer, breast cancer, cervical cancer, colon cancer, endometrial cancer, esophageal cancer, head and neck cancer, intestinal cancer, liver cancer, lung cancer, oral cancer, ovarian cancer, pancreatic cancer, renal cancer, prostate cancer, salivary gland cancer, skin cancer, stomach cancer, testicular cancer, throat cancer, thyroid cancer, uterine cancer, vaginal cancer, sarcoma, or soft tissue carcinomas.

[0013] In some embodiments, the cancer for treatment is a hematological cancer, such as leukemia or lymphoma. In some embodiments, the hematological cancer is lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), lymphoma (e.g., Hodgkin's lymphoma, Non-Hodgkin's lymphoma, Burkitt's lymphoma), chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), Hairy Cell chronic myelogenous leukemia (CML), and multiple myeloma.

4. BRIEF DESCRIPTION OF THE DRAWINGS

[0014] **FIG. 1** shows PKC activation in A549 non-small cell lung cancer cells by diterpenoid compounds as assessed by measuring levels of phosphorylated PKC (p-PKC) and phosphorylated ERK1/2 proteins (p-ERK1/2).

[0015] **FIG. 2** shows PKC activation in A549 non-small cell lung cancer cells by diterpenoid compounds as assessed by measuring levels of phosphorylated PKC (p-PKC) and phosphorylated ERK1/2 proteins (p-ERK1/2).

[0016] **FIG. 3** shows PKC activation in A549 non-small cell lung cancer cells by diterpenoid compounds assessed by measuring levels of phosphorylated PKC (p-PKC) and phosphorylated ERK1/2 proteins (p-ERK1/2), with prostratin (K101) provided for comparison.

[0017] **FIG. 4A** shows PKC activation in A549 non-small cell lung cancer cells by diterpenoid compounds based on levels of phosphorylated PKC (p-PKC) and phosphorylated ERK1/2 proteins (p-ERK1/2).

[0018] **FIG. 4B** shows PKC activation in A549 non-small cell lung cancer cells by diterpenoid compounds assessed by measuring levels of phosphorylated PKD/PKC μ (p-PKC) and phosphorylated PKC δ .

[0019] **FIG. 5** shows effect of selected diterpenoid compounds on levels of phosphorylated CaMKii (p-CaMKii), a marker of K-Ras stemness pathway inhibition, in Panc1 pancreatic cancer cell line.

[0020] **FIGS. 6A-6D** show sphere formation by Panc1 pancreatic cancer cell line treated with different diterpenoid compounds.

[0021] FIG. 7 shows effect of intratumoral administration (7 daily injections) of diterpenoid compounds into Panc2.13 tumors in mice with Panc2.13 pancreatic cancer cell line xenografts.

5. DETAILED DESCRIPTION

[0022] As used in this specification and the appended claims, the singular forms “a”, “an” and “the” include plural referents unless the context clearly indicates otherwise. Thus, for example, reference to “a protein” includes more than one protein, and reference to “a compound” refers to more than one compound.

[0023] Also, the use of “or” means “and/or” unless stated otherwise. Similarly, “comprise,” “comprises,” “comprising” “include,” “includes,” and “including” are interchangeable and not intended to be limiting.

[0024] It is to be further understood that where descriptions of various embodiments use the term “comprising,” those skilled in the art would understand that in some specific instances, an embodiment can be alternatively described using language “consisting essentially of” or “consisting of.”

[0025] It is to be understood that both the foregoing general description, including the drawings, and the following detailed description are exemplary and explanatory only and are not restrictive of this disclosure. The section headings used herein are for organizational purposes only and not to be construed as limiting the subject matter described.

5.1. Definitions

[0026] In reference to the present disclosure, the technical and scientific terms used in the descriptions herein will have the meanings commonly understood by one of ordinary skill in the art, unless specifically defined otherwise. Accordingly, the following terms are intended to have the meanings as described below.

[0027] “Alkyl” refers to straight or branched chain hydrocarbon groups of 1 to 20 carbon atoms, particularly 1 to 12 carbon atoms (C_1 - C_{12} or C_{1-12}), and more particularly (C_1 - C_8 or C_{1-8}) carbon atoms. Exemplary “alkyl” includes, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, and s-pentyl.

[0028] “Alkenyl” refers to straight or branched chain hydrocarbon group of 2 to 20 carbon atoms, particularly 2 to 12 carbon atoms (C_2 - C_{12} or C_{2-12}), and most particularly 2 to 8 (C_2 - C_8 or C_{2-8}) carbon atoms, having at least one double bond. Exemplary “alkenyl” includes, but are not limited to, vinyl ethenyl, allyl, isopropenyl, 1-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 2-ethyl-1-butenyl, 3-methyl-2-butenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 4-methyl-3-pentenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl and 5-hexenyl.

[0029] “Alkynyl” refers to a straight or branched chain hydrocarbon group of 2 to 12 carbon atoms (C_2 - C_{12} or C_{2-12}), particularly 2 to 8 carbon atoms (C_2 - C_8 or C_{2-8}), containing at least one triple bond. Exemplary “alkynyl” includes ethynyl, 1-propynyl, 2-propynyl, 1-butynyl, 2-butynyl, 3-butynyl, 1-pentynyl, 2-pentynyl, 3-pentynyl, 4-pentynyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl and 5-hexynyl.

[0030] “Alkylene”, “alkenylene” and “alkynylene” refers to a straight or branched chain divalent hydrocarbon radical of the corresponding alkyl, alkenyl, and alkynyl, respectively. The “alkylene”, “alkenylene” and “alkynylene” may be optionally substituted, for example with alkyl, alkyloxy, hydroxyl, carbonyl, carboxyl, halo, nitro, and the like.

[0031] “Aliphatic” refers to an organic compound characterized by substituted or unsubstituted, straight or branched, and/or cyclic chain arrangements of constituent carbon atoms. Aliphatic compounds do not contain aromatic rings as part of the molecular structure of the compounds. Aliphatic compound can have 1-20 (C_1 - C_{20} or C_{1-20}) carbon atoms, 1-12 (C_1 - C_{12} or C_{1-12}) carbon atoms, or particularly 1-8 (C_1 - C_8 or C_{1-8}) carbon atoms.

[0032] “Lower” in reference to substituents refers to a group having between one and six carbon atoms.

[0033] “Cycloalkyl” refers to any stable monocyclic or polycyclic system which consists of carbon atoms, any ring of which being saturated. “Cycloalkenyl” refers to any stable monocyclic or polycyclic system which consists of carbon atoms, with at least one ring thereof being partially unsaturated. Examples of cycloalkyls include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, bicycloalkyls and tricycloalkyls (e.g., adamantyl).

[0034] “Heterocycloalkyl” or “heterocyclyl” refers to a substituted or unsubstituted 3 to 14 membered, mono- or bicyclic, non-aromatic hydrocarbon, wherein 1 to 3 carbon atoms are replaced by a heteroatom. Heteroatoms and/or heteroatomic groups which can replace the carbon atoms include, but are not limited to, -O-, -S-, -S-O-, -NR'-, -PH-, -S(O)-, -S(O)₂-, -S(O)NR'-, -S(O)₂NR'-, and the like, including combinations thereof, where each R' is independently hydrogen or lower alkyl. Examples include oxiranyl, oxetanyl, azetidynyl, oxazolyl, thiazolidinyl, thiazolyl, morpholinyl, pyrrolidinonyl, pyrrolidinyl, piperidinyl, piperazinyl, 2,3-dihydrofuranyl, dihydropyranyl, tetrahydrofuranyl, tetrahydropyranyl, dihydropyridinyl, tetrahydropyridinyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, azapanyl, and the like.


[0035] “Carbocycle,” “carbocyclyl,” and “carbocyclic,” as used herein, refer to a non-aromatic saturated or unsaturated ring in which each atom of the ring is carbon. The ring may be monocyclic, bicyclic, tricyclic, or even of higher order. Thus, the terms “carbocycle”, “carbocyclyl”, and “carbocyclic”, encompass fused, bridged and spirocyclic systems: Preferably a carbocycle ring contains from 3 to 14 atoms, including 3 to 8 or 5 to 7 atoms, such as for example, 6 atoms.


[0036] “Aryl” refers to a six- to fourteen-membered, mono- or bi-carbocyclic ring, wherein the monocyclic ring is aromatic and at least one of the rings in the bicyclic ring is aromatic. Unless stated otherwise, the valency of the group may be located on any atom of any ring within the radical, valency rules permitting. Examples of “aryl” groups include phenyl, naphthyl, indenyl, biphenyl, phenanthrenyl, naphthacenyl, and the like.

[0037] “Heteroaryl” refers to an aromatic heterocyclic ring, including both monocyclic and bicyclic ring systems, where at least one carbon atom of one or both of the rings is replaced with a heteroatom independently selected from nitrogen, oxygen, and sulfur, or at least two carbon atoms of one or both of the rings are replaced with a heteroatom independently selected from nitrogen, oxygen, and sulfur. In some embodiments, the heteroaryl can be a 5 to 6 membered monocyclic, or 7 to 11 membered bicyclic ring systems. Examples of “heteroaryl” groups include pyrrolyl, pyrazolyl, imidazolyl, pyrazinyl, oxazolyl, isoxazolyl, thiazolyl, furyl, thienyl, pyridyl, pyrimidyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, purinyl, benzimidazolyl, indolyl, isoquinolyl, quinoxalyl, quinolyl, and the like.

[0038] “Bridged bicyclic” refers to any bicyclic ring system, i.e., carbocyclic or heterocyclic, saturated or partially unsaturated, having at least one bridge. As defined by IUPAC, a “bridge” is an unbranched chain of atoms or an atom or a valence bond connecting two bridgeheads, where a “bridgehead” is any skeletal atom of the ring system which is bonded to three or more skeletal atoms (excluding hydrogen). In some embodiments, a bridged bicyclic group has 5 to 12 ring members and 0-4 heteroatoms independently selected from nitrogen, oxygen, and sulfur. Such bridged bicyclic groups include those groups set forth below where each group is attached to the rest of the molecule at any substitutable carbon or nitrogen atom. Unless otherwise specified, a bridged bicyclic group is optionally substituted with one or more substituents as set forth for aliphatic groups. Additionally or alternatively, any substitutable nitrogen of a bridged bicyclic group is optionally substituted.

Exemplary bridged bicyclics include:





[0039] “Fused ring” refers a ring system with two or more rings having at least one bond and two atoms in common. A “fused aryl” and a “fused heteroaryl” refer to ring systems having at least one aryl and heteroaryl, respectively, that share at least one bond and two atoms in common with another ring.

[0040] “Carbonyl” refers to $-C(O)-$. The carbonyl group may be further substituted with a variety of substituents to form different carbonyl groups including acids, acid halides, aldehydes, amides, esters, and ketones. For example, an $-C(O)R'$, wherein R' is an alkyl is referred to as an alkylcarbonyl. In some embodiments, R' is selected from an optionally substituted: alkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, arylalkyl, heteroaryl, and heteroarylalkyl.

[0041] “Halogen” or “halo” refers to fluorine, chlorine, bromine and iodine.

[0042] “Haloalkyl” refers to an alkyl substituted with 1 or more halogen atoms. Preferably, the alkyl is substituted with 1 to 3 halogen atoms.

[0043] “Hydroxy” refers to $-OH$.

[0044] “Oxy” refers to group $-O-$, which may have various substituents to form different oxy groups, including ethers and esters. In some embodiments, the oxy group is an $-OR'$, wherein R' is selected from an optionally substituted: alkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, arylalkyl, heteroaryl, and heteroarylalkyl.

[0045] “Acyl” refers to $-C(O)R'$, where R is hydrogen, or an optionally substituted alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, arylalkyl, heteroaryl, or heteroarylalkyl as defined herein. Exemplary acyl groups include, but are not limited to, formyl, acetyl, cyclohexylcarbonyl, cyclohexylmethylcarbonyl, benzoyl, benzylcarbonyl, and the like.

[0046] “Alkyloxy” or “alkoxy” refers to $-OR'$, wherein R' is an optionally substituted alkyl.

[0047] “Aryloxy” refers to $-OR'$, wherein R' is an optionally substituted aryl.

[0048] “Carboxy” refers to $-COO^-$ or $COOM$, wherein H or a M^+ counterion.

[0049] “Carbamoyl” refers to $-C(O)NR'R'$, wherein each R' is independently selected from H or an optionally substituted alkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, arylalkyl, heteroaryl, or heteroarylalkyl.

[0050] “Cyano” refers to $-CN$.

[0051] “Ester” refers to a group such as $-C(=O)OR'$, alternatively illustrated as $-C(O)OR'$, wherein R' is selected from an optionally substituted: alkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, arylalkyl, heteroaryl, and heteroarylalkyl.

[0052] “Silyl” refers to Si, which may have various substituents, for example $-SiR'R'R'$, where R' is as defined in the specification. For example, each R' is independently selected from alkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, arylalkyl, heteroaryl, and heteroarylalkyl. As defined herein, any heterocycloalkyl or heteroaryl group present in a silyl group has from 1 to 3 heteroatoms selected independently from O, N, and S.

[0053] “Thiol” refers to $-SH$.

[0054] “Sulfanyl” refers to $-SR'$, wherein R' is selected from an optionally substituted: alkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, arylalkyl, heteroaryl, and heteroarylalkyl. For example, $-SR$, wherein R is an alkyl is an alkylsulfanyl.


[0055] “Sulfonyl” refers to $-S(O)_2-$, which may have various substituents to form different sulfonyl groups including sulfonic acids, sulfonamides, sulfonate esters, and sulfones. For example, $-S(O)_2R'$, wherein R' is an alkyl refers to an alkylsulfonyl. In some embodiments of $-S(O)_2R'$, R' is selected from an optionally substituted: alkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, arylalkyl, heteroaryl, and heteroarylalkyl.

[0056] “Amino” or “amine” refers to the group $-NR'R'$ or $-NR'R'R'$, wherein each R' is independently selected from H and an optionally substituted: alkyl, cycloalkyl, heterocycloalkyl, alkyloxy, aryl, heteroaryl, heteroarylalkyl, acyl, alkyloxycarbonyl, sulfanyl, sulfinyl, sulfonyl, and the like. Exemplary amino groups include, but are not limited to, dimethylamino, diethylamino, trimethylammonium, triethylammonium, methylsulfonylamino, furanyl-oxy-sulfamino, and the like.

[0057] “Amide” refers to a group such as, $-C(=O)NR'R'$, wherein each R' is independently selected from H and an optionally substituted: alkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, arylalkyl, heteroaryl, and heteroarylalkyl.

[0058] “Spiroalkyl” as used herein refers to a monospiro compound having two alicyclic rings attached together through a single common carbon atom. In some embodiments, the spiro compounds have 5 to 12 total ring atoms (e.g., C_5-C_{12} or C_{5-12}). In some embodiments, one or more of the carbon atoms can be replaced with a heteroatom, such as oxygen, nitrogen or sulfur. Exemplary spiroalkyl compounds include, among others, spiro[3,3]heptyl, spiro[3,4]octyl, and spiro[3,5]decyl.

[0059] “Adamantyl” refers to a compound of structural formula:



where optional substitutions can be present on one or more of R^a , R^b , R^c , and R^d . Adamantyl includes substituted adamantyl, e.g., 1- or 2-adamantyl, substituted by one or more substituents, including alkyl, halo, OH, NH_2 , and alkoxy. Exemplary derivatives include methyladamantane, haloadamantane, hydroxyadamantane, and aminoadamantane (e.g., amantadine).

[0060] “N-protecting group” as used herein refers to those groups intended to protect a nitrogen atom against undesirable reactions during synthetic procedures. Exemplary N-protecting groups include, but is not limited to, acyl groups such as acetyl and t-butylacetyl, pivaloyl, alkoxy carbonyl groups such as methyloxycarbonyl and t-butyloxycarbonyl (Boc), aryloxycarbonyl groups such as benzyloxycarbonyl (Cbz) and fluorenylmethoxycarbonyl (Fmoc) and aroyl groups such as benzoyl. N-protecting groups are described in Greene's Protective Groups in Organic Synthesis, 5th Edition, P. G. M. Wuts, ed., Wiley (2014).

[0061] “Optional” or “optionally” refers to a described event or circumstance may or may not occur, and that the description includes instances where the event or circumstance occurs and instances where the event or circumstance does not. For example, “optionally substituted alkyl” refers to an alkyl group that may or may not be substituted and that the description encompasses both substituted alkyl group and unsubstituted alkyl group.

[0062] “Substituted” as used herein means one or more hydrogen atoms of the group is replaced with a substituent atom or group commonly used in pharmaceutical chemistry. Each substituent can be the same or different. Examples of suitable substituents include, but are not limited to, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, arylalkyl, heterocycloalkyl, heteroaryl, OR' (e.g., hydroxyl, alkyloxy (e.g., methoxy, ethoxy, and propoxy), aryloxy, heteroaryloxy, arylalkyloxy, ether, ester, carbamate, etc.), hydroxyalkyl, alkyloxycarbonyl, alkyloxyalkyloxy, perhaloalkyl, alkyloxyalkyl, SR' (e.g., thiol, alkylthio, arylthio, heteroarylthio, arylalkylthio, etc.), $\text{S}^+\text{R}'_2$, $\text{S}(\text{O})\text{R}'$, $\text{SO}_2\text{R}'$, $\text{NR}'\text{R}''$ (e.g., primary amine (i.e., NH_2), secondary amine, tertiary amine, amide, carbamate, urea, etc.), hydrazide, halo, nitrile, nitro, sulfide, sulfoxide, sulfone, sulfonamide, thiol, carboxy, aldehyde, keto, carboxylic acid, ester, amide, imine, and imide, including seleno and thio derivatives thereof, wherein each of the substituents can be optionally further substituted. In embodiments in which a functional group with an aromatic carbon ring is substituted, such substitutions will typically number less than about 10 substitutions, more preferably about 1 to 5, with about 1 or 2 substitutions being preferred.

[0063] “Stereoisomer” refers to a compound made up of the same atoms bonded by the same bonds but having different three-dimensional structures, which are not interchangeable. Thus, “stereoisomer

thereof” with respect to a compound includes any stereoisomer of the compound and mixtures of stereoisomers, and includes “enantiomers,” which refers to two stereoisomers whose molecules are nonsuperimposable mirror images of one another. A compound may have more than one chiral center such that the compound may exist as either an individual diastereomer or as a mixture of diastereomers.

[0064] “Tautomer” refers to a proton shift from one atom of a molecule to another atom of the same molecule. Thus, “tautomers thereof” with respect to a compound includes any tautomers of the compound.

[0065] “Prodrug” refers to a derivative of an active compound (e.g., drug) that requires a transformation under the conditions of use, such as within the body or appropriate *in vitro* conditions, to release the active drug. Prodrugs are frequently, but not necessarily, pharmacologically inactive until converted into the active drug. Prodrugs can be obtained by masking a functional group in the drug believed to be in part required for activity with a progroup to form a promoiety which undergoes a transformation, such as cleavage, under the specified conditions of use to release the functional group, and hence the active drug. The cleavage of the promoiety may proceed spontaneously, such as by way of a hydrolysis reaction, or it may be catalyzed or induced by another agent, such as by an enzyme, by light, by acid, or by a change of or exposure to a physical or environmental parameter, such as a change of temperature. The agent may be endogenous to the conditions of use, such as an enzyme present in the cells to which the prodrug is administered or the acidic conditions of the stomach, or it may be supplied exogenously.

[0066] Various progroups, as well as the resultant promoieties, suitable for masking functional groups in the active drugs to yield prodrugs can be used. For example, a hydroxyl functional group may be masked as a sulfonate, ester or carbonate promoiety, which may be hydrolyzed *in vivo* to provide the hydroxyl group. An amino functional group may be masked as an amide, carbamate, imine, urea, phosphanyl, phosphoryl or sulfenyl promoiety, which may be hydrolyzed, e.g., *in vivo* or under appropriate *in vitro* conditions, to provide the amino group. A carboxyl group may be masked as an ester (including silyl esters and thioesters), amide or hydrazide promoiety, which may be hydrolyzed *in vivo* to provide the carboxyl group. Included within the scope of prodrugs are, among others, “biohydrolyzable carbonate”, “biohydrolyzable ureide”, “biohydrolyzable carbamate”, “biohydrolyzable ester”, “biohydrolyzable amide”, and “biohydrolyzable phosphate” groups.

[0067] “Solvate” refers to a complex of variable stoichiometry formed by a solute, such as a PKC activator compound, and a solvent. Such solvents are selected to minimally interfere with the biological activity of the solute. Solvents may be, by way of example and not limitation, water, ethanol, or acetic acid.

[0068] “Hydrate” refers to a combination of water with a solute, such as a PKC activator compound, wherein the water retains its molecular state as water and is either absorbed, adsorbed or contained within a crystal lattice of the solute (e.g., PKC activating compound).

[0069] “Pharmaceutically acceptable salts” is meant to include salts of the active compounds which are prepared with relatively nontoxic acids or bases, depending on the particular substituents found on the compounds described herein. When compounds of the present invention contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired base, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable base addition salts include sodium, potassium, calcium, ammonium, organic amino, or magnesium salt, or a similar salt. When compounds of the present invention contain relatively basic functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, phosphoric, partially neutralized phosphoric acids, sulfuric, partially neutralized sulfuric, hydroiodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, maleic, malonic, benzoic, succinic, suberic, fumaric, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galacturonic acids and the like. Certain specific compounds of the present disclosure may contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts. Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 17th Ed., Mack Publishing Company, Easton, Pa., (1985) and Journal of Pharmaceutical Science, 66:2 (1977), each of which is incorporated herein by reference in its entirety.

[0070] “Pharmaceutically acceptable carrier” or “pharmaceutically acceptable excipient” refers to an excipient, carrier or adjuvant that can be administered to a subject, together with at least one therapeutic agent, and which does not destroy the pharmacological activity thereof and is generally safe, nontoxic and neither biologically nor otherwise undesirable when administered in doses sufficient to deliver a therapeutic amount of the agent..

[0071] “K-RAS” refers to Kirsten rat sarcoma viral oncogene homolog, a small GTPase and a member of the RAS family of proteins involved in signal transduction. Exemplary human *K-RAS* nucleic acid and protein sequences are provided in GenBank Nos. M54968.1 and AAB414942.1, respectively. “K-RAS” as used herein encompasses variants, including orthologs and interspecies homologs, of the human K-RAS protein.

[0072] “Mutant K-RAS polypeptide”, “mutant K- RAS protein” and “mutant K- RAS” are used interchangeably and refer to a K- RAS polypeptide comprising at least one K- RAS mutation as compared to the corresponding wild-type K- RAS sequence. Certain exemplary mutant K- RAS polypeptides include, but are not limited to, allelic variants, splice variants, derivative variants, substitution variants, deletion variants, insertion variants, and fusion polypeptides.

[0073] “N-RAS” refers to Neuroblastoma RAS Viral (V-RAS) oncogene homolog, a small GTPase and a member of the RAS family of proteins involved in signal transduction. Exemplary human *N-RAS* nucleic acid and protein sequences are provided in NCBI Accession No. NP_002515 and GenBank Accession No. X02751, respectively. “N-RAS” as used herein encompasses variants, including orthologs and interspecies homologs of the human N-RAS protein.

[0074] “Mutant N- RAS polypeptide”, “mutant N- RAS protein” and “mutant N-RAS” are used interchangeably and refer to an N-RAS polypeptide comprising at least one N- RAS mutation as compared to the corresponding wild-type N- RAS sequence. Certain exemplary mutant N- RAS polypeptides include, but are not limited to, allelic variants, splice variants, derivative variants, substitution variants, deletion variants, insertion variants, and fusion polypeptides.

[0075] “H- RAS” refers to Harvey Rat Sarcoma viral oncogene homolog, a small GTPase and a member of the RAS family of proteins involved in signal transduction. Exemplary human *H-RAS* nucleic acid and protein sequences are provided in NCBI Accession No. P01112 and GenBank Accession No. NM_176795, respectively. “H- RAS” as used herein encompasses variants, including orthologs and interspecies homologs of the human H- RAS protein.

[0076] “Mutant H-RAS polypeptide”, “mutant H-RAS protein” and “mutant H-RAS” are used interchangeably and refer to an H-RAS polypeptide comprising at least one H- RAS mutation as compared to the corresponding wild-type H- RAS sequence. Certain exemplary mutant H- RAS polypeptides include, but are not limited to, allelic variants, splice variants, derivative variants, substitution variants, deletion variants, insertion variants, and fusion polypeptides.

[0077] “Activating K- RAS” refers to a form of K- RAS that has increased activity compared to wild-type K- RAS. The activation of K- RAS activity can result from a mutation or in some embodiments, overexpression of the K- RAS protein.

[0078] “Activating N- RAS” refers to a form of N- RAS that has increased activity compared to wild-type N- RAS. The activation of N- RAS activity can result from a mutation, or in some embodiments, overexpression of the N- RAS protein.

[0079] “Activating H- RAS” refers to a form of H- RAS that has increased activity compared to wild-type H- RAS. The activation of H- RAS activity can result from a mutation, or in some embodiments, overexpression of the H- RAS protein.

[0080] “Mutation” or “mutant” refers to an amino acid or polynucleotide sequence which has been altered by substitution, insertion, and/or deletion. In some embodiments, a mutant or variant sequence can have increased, decreased, or substantially similar activities or properties in comparison to the parental sequence.

[0081] “Identified” or “determined” refers to analyzing for, detection of, or carrying out a process for the presence or absence of one or more specified characteristics.

[0082] “Wild-type” or “naturally occurring” refers to the form found in nature. For example, a naturally occurring or wild-type polypeptide or polynucleotide sequence is a sequence present in an organism that can be isolated from a source in nature and which has not been intentionally modified by human manipulation.

[0083] “Control” or “control sample” or “control group” refers to a sample or group that is compared to another sample or group, where generally the control sample or group are the same as a comparison group except for one or more factors being compared.

[0084] “Selecting” refers to the process of determining that a subject will receive an agent to treat the occurrence of a condition. Selecting can be based on an individual susceptibility to a particular disease or condition due to, for example, presence of an identifying cellular, physiological or environment factor or factors. In some embodiments, selecting can be based on determining or identifying whether that subject will be responsive to an agent, for example as assessed by identifying the presence of a biomarker and/or drug target marker that makes the subject sensitive, insensitive, responsive, or unresponsive to an agent or treatment.

[0085] “Biological sample” refers to any sample including a biomolecule, such as a protein, a peptide, a nucleic acid, a lipid, a carbohydrate or a combination thereof, that is obtained from an organism, particularly a mammal. Examples of mammals include humans; veterinary animals like cats, dogs, horses, cattle, and swine; and laboratory animals like mice, rats and primates. In some embodiments, a human subject in the clinical setting is referred to as a patient. Biological samples include tissue samples (such as tissue sections and needle biopsies of tissue), cell samples (for example, cytological smears such as Pap or blood smears or samples of cells obtained by microdissection), or cell fractions, fragments or organelles (such as obtained by lysing cells and separating their components by centrifugation or otherwise). Other examples of biological samples include blood, serum, urine, semen, fecal matter, cerebrospinal fluid, interstitial fluid, mucous, tears, sweat, pus, biopsied tissue (for example, obtained by a surgical biopsy or a needle biopsy), nipple aspirates, milk, vaginal fluid, saliva, swabs (such as buccal swabs), or any material containing biomolecules that is derived from a first biological sample. In particular embodiments, the biological sample is a “cell free sample”, such as cell free or extracellular polynucleotides, and cell free or

extracellular proteins. In some embodiments, cell free DNA or cfDNA refers to extracellular DNA obtained from blood, particularly the serum.


[0086] “Subject” as used herein refers to a mammal, for example a dog, a cat, a horse, or a rabbit. In some embodiments, the subject is a non-human primate, for example a monkey, chimpanzee, or gorilla. In some embodiments, the subject is a human, sometimes referred to herein as a patient.

[0087] “Treating” or “treatment” of a disease, disorder, or syndrome, as used herein, includes (i) preventing the disease, disorder, or syndrome from occurring in a subject, i.e., causing the clinical symptoms of the disease, disorder, or syndrome not to develop in an animal that may be exposed to or predisposed to the disease, disorder, or syndrome but does not yet experience or display symptoms of the disease, disorder, or syndrome; (ii) inhibiting the disease, disorder, or syndrome, i.e., arresting its development; and (iii) relieving the disease, disorder, or syndrome, i.e., causing regression of the disease, disorder, or syndrome. As is known in the art, adjustments for systemic versus localized delivery, age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition may be necessary, and will be ascertainable with routine experimentation by one of ordinary skill in the art, particularly in view of the guidance provided in the present disclosure.

[0088] “Therapeutically effective amount” refers to that amount which, when administered to an animal for treating a disease, is sufficient to effect such treatment for the disease, disorder, or condition.

5.2. Compounds

[0089] The present disclosure provides protein kinase C (PKC) modulating compounds. In particular, the compounds are diterpenoid PKC modulating compounds displaying potent PKC modulating activity, as well as enhanced solubility and pharmacokinetic profiles. The diterpenoid PKC modulating compounds show potent activity against tumor cells, and can be applied to treatment of cancer. In one aspect, the present disclosure provides a compound of formula (I):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

A is -OH, -C(O)OR¹, or -NR¹³R^{13'};

R¹ is H or a M⁺ counterion;

R² is a C₁-C₄alkyl;


R³ is O double bonded to the ring carbon when (- - -) is a bond, or -OR^a; wherein R^a is H or -C(O)R^{al}, wherein R^{al} is C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R⁴ and R⁵ are each independently H or -OR^b, wherein R^b is H, C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R^{5a} and R^{6a} are H, or R^{5a} and R^{6a} form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R⁶ is OH, halo, or -OC(O)R^c, wherein R^c is -C₁-C₆alkyl, -C₁-C₆alkyl-(NR^{cl})₂ or -C₁-C₆alkylC(O)OR^k, R^{cl} is H, C₁-C₆alkyl, or two R^{cl} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R⁶ is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^{6a} and R^{7a} are H, or R^{6a} and R^{7a} form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R⁷ is H or OH;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹² is H, -OH, -OC(O)R^f, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-

C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

n is 0 or 1.

[0090] In some embodiments of the compound of formula (I), the carbon atom marked with “*”




is chiral and thus can be present in *S* or *R* stereochemical configuration. In some embodiments, the stereochemical configuration is *S* isomer. In some embodiments, the stereochemical configuration is *R* isomer.

[0091] In some embodiments, A is -OH. In some embodiments, A is -C(O)OR¹, wherein R¹ is H or a M⁺ counterion. In some embodiments, A is -NR¹³R^{13'}, wherein R¹³ and R^{13'} are each independently H or C₁-C₄alkyl.

[0092] In the embodiments herein, M⁺ is a metal cation, an ammonium group, or a suitable organic cation. In some embodiments, M⁺ is a cation of an alkaline or alkaline earth metal, for example, K⁺,

Na^+ , Li^+ , or Ca^{+2} . In some embodiments, M^+ is an ammonium ion NH_4^+ , or an organic cation derived from an amine.

[0093] In some embodiments, the compound has the structure of formula (Ia):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

A is -OH;

R^2 is a C_1 - C_4 alkyl;


R^3 is O double bonded to the ring carbon when (- -) is a bond, or $-\text{OR}^a$, wherein R^a is H or $-\text{C}(\text{O})\text{R}^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-\text{OR}^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

$\text{R}^{5'}$ and $\text{R}^{6'}$ are H, or $\text{R}^{5'}$ and $\text{R}^{6'}$ form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R^6 is OH, halo, or $-\text{OC}(\text{O})\text{R}^c$, wherein R^c is $-\text{C}_1$ - C_6 alkyl, $-\text{C}_1$ - C_6 alkyl- $(\text{NR}^{\text{cl}})_2$ or $-\text{C}_1$ - C_6 alkyl $\text{C}(\text{O})\text{OR}^k$, R^{cl} is H, C_1 - C_6 alkyl, or two R^{cl} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^{6'} and R^{7'} are H, or R^{6'} and R^{7'} form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R⁷ is H or OH;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹² is H, -OH, -OC(O)R^f, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and


n is 0 or 1.

[0094] In some embodiments of the compound of formula (Ia), the carbon atom marked with “*”



is chiral and thus can be present in *S* or *R* stereochemical configuration. In some embodiments, the stereochemical configuration is *S* isomer. In some embodiments, the stereochemical configuration is *R* isomer.

[0095] In some embodiments, the compound has the structure of formula (Ib):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

A is $-\text{C}(\text{O})\text{OR}^1$;

R^1 is H or a M^+ counterion;

R^2 is a C_1 - C_4 alkyl;

R^3 is O double bonded to the ring carbon when (---) is a bond, or $-\text{OR}^a$, wherein R^a is H or $-\text{C}(\text{O})\text{R}^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;


R^4 and R^5 are each independently H or $-\text{OR}^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 -

C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R^{5'} and R^{6'} are H, or R^{5'} and R^{6'} form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R⁶ is OH, halo, or -OC(O)R^c, wherein R^c is -C₁-C₆alkyl, -C₁-C₆alkyl-(NR^{cl})₂ or -C₁-C₆alkylC(O)OR^k, R^{cl} is H, C₁-C₆alkyl, or two R^{cl} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R⁶ is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^{6'} and R^{7'} are H, or R^{6'} and R^{7'} form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R⁷ is H or OH;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹² is H, -OH, -OC(O)R^f, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J^l, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and

optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and


n is 0 or 1.

[0096] In some embodiments, of the compound of formula (Ib), the carbon atom marked with “*”



is chiral and thus can be present in *S* or *R* stereochemical configuration. In some embodiments, the stereochemical configuration is *S* isomer. In some embodiments, the stereochemical configuration is *R* isomer.

[0097] In some embodiments, the present disclosure provides a compound of formula (II):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^2 is a C_1 - C_4 alkyl;


R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$; wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

$R^{5'}$ and $R^{6'}$ are H, or $R^{5'}$ and $R^{6'}$ form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

$R^{6'}$ and $R^{7'}$ are H, or $R^{6'}$ and $R^{7'}$ form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R^7 is H or OH;

R^9 is OR^e , wherein R^e is H, C_1 - C_6 alkyl, or aryl;

R^{11} is C_1 - C_4 alkyl;

R^{12} is H, $-OH$, $-OC(O)R^f$, wherein R^f is C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, $-C_0$ - C_{12} aliphatic- C_3 - C_7 cycloalkyl, $-C_0$ - C_{12} aliphatic-heterocycloalkyl, $-C_0$ - C_{12} aliphatic-aryl, or $-C_0$ - C_{12} aliphatic-heteroaryl;

R^{13} and $R^{13'}$ are each independently H or C_1 - C_4 alkyl;

R^{14} is H or OR^g ; wherein R^g is H or C_1 - C_6 alkyl;

R^{17} and R^{18} are each independently C_1 - C_4 alkyl or C_1 - C_4 alkyl- OR^h , wherein R^h is H or C_1 - C_6 alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each Rⁱ is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;


J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

n is 0 or 1.

[0098] In some embodiments of the compound of formula (II), the carbon atom marked with “*”




is chiral and thus compound may be present in *S* or *R* stereochemical configuration. In some embodiments, the stereochemical configuration is *S* isomer. In some embodiments, the stereochemical configuration is *R* isomer. In some embodiments, the compound of formula (II) has the structure of formula (II')




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^2 , R^3 , R^4 , R^5 , $R^{5'}$, R^6 , $R^{6'}$, R^7 , $R^{7'}$, R^9 , R^{11} , R^{12} , R^{13} , $R^{13'}$, R^{14} , R^{17} , R^{18} , L and R^{21} are as defined for formula (II).

[0099] In some embodiments, the compound of formula (II) has the structure of formula (II''):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^2 , R^3 , R^4 , R^5 , $R^{5'}$, R^6 , $R^{6'}$, R^7 , $R^{7'}$, R^9 , R^{11} , R^{12} , R^{13} , $R^{13'}$, R^{14} , R^{17} , R^{18} , L and R^{21} are as defined for formula (II).

[0100] In some embodiments, the compound has the structure of formula (IIa):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein


R^2 is a C_1 - C_4 alkyl;

R^3 is O double bonded to the ring carbon when (- -) is a bond, or $-OR^a$; wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^7 is H or OH;

R^9 is OR^e , wherein R^e is H, C_1 - C_6 alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹² is H, -OH, -OC(O)R^f, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

n is 0 or 1.


[0101] In some embodiments, the compound of formula (IIa) has the structure for formula (IIa')



(IIa')


or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R², R³, R⁴, R⁵, R⁶, R⁷, R⁹, R¹¹, R¹², R¹³, R^{13'}, R¹⁴, R¹⁷, R¹⁸, L and R²¹ are as defined for formula (II).

[0102] In some embodiments, the compound of formula (IIa) has the structure for formula (IIa'')



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^9 , R^{11} , R^{12} , R^{13} , $R^{13'}$, R^{14} , R^{17} , R^{18} , L and R^{21} are as defined for formula (II).

[0103] In some embodiments, the compound has the structure of formula (IIb):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^2 is a C_1 - C_4 alkyl;


R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ -

C₆alkylC(O)OR^k, R^{c1} is H, C₁-C₆alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R⁶ is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹² is H, -OH, -OC(O)R^f, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and


n is 0 or 1.

[0104] In some embodiments, the compound has the structure of formula (IIb'):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^9 , R^{11} , R^{12} , R^{13} , $R^{13'}$, R^{14} , R^{17} , R^{18} , L and R^{21} are as defined for formula (II).


[0105] In some embodiments, the compound has the structure of formula (IIb''):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^9 , R^{11} , R^{12} , R^{13} , $R^{13'}$, R^{14} , R^{17} , R^{18} , L and R^{21} are as defined for formula (II).

[0106] In some embodiments of the compound of formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), and (IIb''), R^3 is $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl. In some embodiments, the aryl of C_0 - C_6 alkylaryl

is phenyl. In some embodiments, the aryl of C₀-C₆alkylaryl is optionally substituted with 1 to 3 of OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl. In some embodiments, R^{al} is selected from:



[0107] In some embodiments, R^{al} is selected from:



[0108] In some embodiments of the compound of formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), and (IIb''),

R³ is O double bonded to the carbon atom.

[0109] In some embodiments of the compound of formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), and (IIb''), one or more of R², R¹¹, R¹⁷, and R¹⁸ are -CH₃. In some embodiments, each of R², R¹¹, R¹⁷, and R¹⁸ is -CH₃.

[0110] In some embodiments of the compound of formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), and (IIb''), R⁴ and R⁵ are each independently H or -OH.


[0111] In some embodiments of the compound of formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), and (IIb''),

R², R¹¹, R¹⁷, and R¹⁸ are -CH₃;

R³ is O double bonded to the carbon atom; and

R⁴ and R⁵ are each independently H or -OH.

[0112] In some embodiments, the compound has the structure of formula (IIc):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1-C_6$ alkyl, $-C_1-C_6$ alkyl- $(NR^{c1})_2$ or $-C_1-C_6$ alkyl $C(O)OR^k$, R^{c1} is H, C_1-C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^{12} is H, $-OH$, $-OC(O)R^f$, wherein R^f is C_1-C_{12} alkyl, C_2-C_{12} alkenyl, $-C_0-C_{12}$ aliphatic- C_3-C_7 cycloalkyl, $-C_0-C_{12}$ aliphatic-heterocycloalkyl, $-C_0-C_{12}$ aliphatic-aryl, or $-C_0-C_{12}$ aliphatic-heteroaryl;

R^{13} and $R^{13'}$ are each independently H or C_1-C_4 alkyl;

L is absent, C_1-C_{12} alkylene, or C_2-C_{12} alkenylene, wherein the C_1-C_{12} alkylene or C_2-C_{12} alkenylene is optionally substituted with C_1-C_4 alkyl;

R^{21} is H, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_3-C_{12} cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or $-C(O)OR^k$, wherein the C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_3-C_{12} cycloalkyl, bridged bicyclyl, or adamantyl is

optionally substituted with 1 to 3 of J^1 , and wherein optionally 1 to 2 carbon atoms of the spiro C_5 - C_{12} cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C_1 - C_4 alkyl, or when an N atom is present an N-protecting group;


each R^i is independently C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkyl C_3 - C_7 cycloalkyl, C_0 - C_6 alkylheterocyclyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl, wherein the C_3 - C_7 cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J^1 ;

R^k is H or M^+ counterion;

J^1 is selected from OH, CN, halo, C_1 - C_4 alkyl, and halo C_1 - C_4 alkyl; and


n is 0 or 1.

[0113] In some embodiments, the compound has the structure of formula (IIc'):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^6 , R^{12} , R^{13} , $R^{13'}$, L and R^{21} are as defined for formula (IIc).

[0114] In some embodiments, the compound has the structure of formula (IIc''):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^6 , R^{12} , R^{13} , $R^{13'}$, L and R^{21} are as defined for formula (IIc).

[0115] In some embodiments, the compound has the structure of formula (II d):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1-C_6$ alkyl, $-C_1-C_6$ alkyl- $(NR^{c1})_2$ or $-C_1-C_6$ alkyl $C(O)OR^k$, R^{c1} is H, C_1-C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R¹² is H, -OH, -OC(O)R^f, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(Rⁱ)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and


n is 0 or 1.

[0116] In some embodiments, the compound has the structure of formula (IIId')




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^6 , R^{12} , R^{13} , $R^{13'}$, L and R^{21} are as defined for formula (II d).

[0117] In some embodiments, the compound has the structure of formula (II d'')




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^6 , R^{12} , R^{13} , $R^{13'}$, L and R^{21} are as defined for formula (II d).


[0118] In some embodiments of the compound of formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (IIc), (IIc''), (II d), (II d') and (II d''), R^{12} is $-OC(O)R^f$, wherein R^f is C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, $-C_0$ - C_{12} aliphatic- C_3 - C_7 cycloalkyl, $-C_0$ - C_{12} aliphatic-heterocycloalkyl, $-C_0$ - C_{12} aliphatic-aryl, or $-C_0$ - C_{12} aliphatic-heteroaryl. In some some embodiments, R^f is selected from



[0119] In some some embodiments, R^f is selected from:



[0120] In some embodiments, the compound has the structure of formula (III):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^2 is a C_1 - C_4 alkyl;


R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

$R^{5'}$ and $R^{6'}$ are H, or $R^{5'}$ and $R^{6'}$ form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^{6'} and R^{7'} are H, or R^{6'} and R^{7'} form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R⁷ is H or OH;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(R^l)₂, -Si(R^l)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each Rⁱ is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and


n is 0 or 1.

[0121] In some embodiments, the compound has the structure of formula (III³):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^2 , R^3 , R^4 , R^5 , $R^{5'}$, $R^{6'}$, R^6 , $R^{6'}$, $R^{7'}$, R^7 , R^9 , R^{11} , R^{13} , $R^{13'}$, R^{14} , R^{17} , R^{18} , L and R^{21} are as defined for formula (III).

[0122] In some embodiments, the compound has the structure of formula (III''):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^2 , R^3 , R^4 , R^5 , $R^{5'}$, $R^{6'}$, R^6 , $R^{6'}$, $R^{7'}$, R^7 , R^9 , R^{11} , R^{13} , $R^{13'}$, R^{14} , R^{17} , R^{18} , L and R^{21} are as defined for formula (III).

[0123] In some embodiments, the compound has the structure of formula (IIIa):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein


R^2 is a C_1 - C_4 alkyl;

R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^7 is H or OH;

R^9 is OR^e , wherein R^e is H, C_1 - C_6 alkyl, or aryl;

R^{11} is C_1 - C_4 alkyl;

R^{13} and $R^{13'}$ are each independently H or C_1 - C_4 alkyl;

R^{14} is H or OR^g ; wherein R^g is H or C_1 - C_6 alkyl;

R^{17} and R^{18} are each independently C_1 - C_4 alkyl or C_1 - C_4 alkyl- OR^h , wherein R^h is H or C_1 - C_6 alkyl;

L is absent, C_1 - C_{12} alkylene, or C_2 - C_{12} alkenylene, wherein the C_1 - C_{12} alkylene or C_2 - C_{12} alkenylene is optionally substituted with C_1 - C_4 alkyl;

R^{21} is H, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3 - C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5 - C_{12} cycloalkyl, 5 to 12 membered bridged bicycyl, adamantyl, or $-C(O)OR^k$, wherein the C_3 - C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5 - C_{12} cycloalkyl, bridged bicycyl, or adamantyl is optionally substituted with 1 to 3 of J^1 , and wherein optionally 1 to 2 carbon atoms of the spiro C_5 - C_{12} cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C_1 - C_4 alkyl, or when an N atom is present an N-protecting group;


each R^j is independently C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkyl C_3 - C_7 cycloalkyl, C_0 - C_6 alkylheterocyclyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl, wherein the C_3 - C_7 cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J^1 ;

R^k is H or M^+ counterion;

J^1 is selected from OH, CN, halo, C_1 - C_4 alkyl, and halo C_1 - C_4 alkyl; and


n is 0 or 1.

[0124] In some embodiments, the compound has the structure of formula (IIIa'):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^9 , R^{11} , R^{12} , R^{13} , $R^{13'}$, R^{14} , R^{17} , R^{18} , L and R^{21} are as defined for formula (III).

[0125] In some embodiments, the compound has the structure of formula (IIIa''):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^9 , R^{11} , R^{12} , R^{13} , $R^{13'}$, R^{14} , R^{17} , R^{18} , L and R^{21} are as defined for formula (III).

[0126] In some embodiments, the compound has the structure of formula (IIIb):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^2 is a C_1 - C_4 alkyl;


R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ -

C_6 alkylC(O)OR^k, R^{c1} is H, C₁-C₆alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R⁶ is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and


n is 0 or 1.

[0127] In some embodiments, the compound has the structure of formula (IIIb'):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^9 , R^{11} , R^{13} , $R^{13'}$, R^{14} , R^{17} , R^{18} , L and R^{21} are as defined for formula (III).

[0128] In some embodiments, the compound has the structure of formula (IIIb''):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^9 , R^{11} , R^{13} , $R^{13'}$, R^{14} , R^{17} , R^{18} , L and R^{21} are as defined for formula (III).

[0129] In some embodiments of the compound of formula (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb') and (IIIb''), R^3 is $-OR^a$; wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl. In some embodiments, the aryl of C_0 - C_6 alkylaryl is phenyl. In some embodiments, the aryl of C_0 - C_6 alkylaryl is optionally substituted with 1 to 3 of OH, CN, halo, C_1 - C_4 alkyl, and halo C_1 - C_4 alkyl. In some embodiments, R^{a1} is selected from:



[0130] In some embodiments, R^{a1} is selected from:



[0131] In some embodiments of the compound of formula (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb') and (IIIb''),

R^3 is O double bonded to the carbon atom.

[0132] In some embodiments of the compound of formula (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb') and (IIIb''), one or more of R^2 , R^{11} , R^{17} , and R^{18} are $-CH_3$. In some embodiments, each of R^2 , R^{11} , R^{17} , and R^{18} is $-CH_3$.

[0133] In some embodiments of the compound of formula (III), (III'), (IIIa), (IIIa'), (IIIb), and (IIIb'), R^4 and R^5 are each independently H or $-OH$.


[0134] In some embodiments of the compound of formula (III), (III'), (IIIa), (IIIa'), (IIIb), and (IIIb'),

R^2 , R^{11} , R^{17} , and R^{18} are $-CH_3$;

R^3 is O double bonded to the carbon atom; and

R^4 and R^5 are each independently H or $-OH$.

[0135] In some embodiments, the compound has the structure of formula (IIIc):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1-C_6$ alkyl, $-C_1-C_6$ alkyl- $(NR^{c1})_2$ or $-C_1-C_6$ alkylC(O)OR^k, R^{c1} is H, C_1-C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^{13} and $R^{13'}$ are each independently H or C_1-C_4 alkyl;

L is absent, C_1-C_{12} alkylene, or C_2-C_{12} alkenylene, wherein the C_1-C_{12} alkylene or C_2-C_{12} alkenylene is optionally substituted with C_1-C_4 alkyl;

R^{21} is H, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or $-C(O)OR^k$, wherein the C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J^1 , and wherein optionally 1 to 2 carbon atoms of the spiro C_5-C_{12} cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C_1-C_4 alkyl, or when an N atom is present an N-protecting group;


each R^j is independently C_1-C_6 alkyl, C_2-C_6 alkenyl, C_0-C_6 alkyl C_3-C_7 cycloalkyl, C_0-C_6 alkylheterocyclyl, C_0-C_6 alkylaryl, or C_0-C_6 alkylheteroaryl, wherein the C_3-C_7 cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J^1 ;

R^k is H or M^+ counterion;

J^1 is selected from OH, CN, halo, C_1-C_4 alkyl, and halo C_1-C_4 alkyl; and


n is 0 or 1.

[0136] In some embodiments, the compound has the structure of formula (IIIc'):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^6 , R^{13} , $R^{13'}$, L and R^{21} are as defined for formula (IIIc).

[0137] In some embodiments, the compound has the structure of formula (IIIc''):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^6 , R^{13} , $R^{13'}$, L and R^{21} are as defined for formula (IIIc).

[0138] In some embodiments, the compound has the structure of formula (IIIId):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1-C_6$ alkyl, $-C_1-C_6$ alkyl- $(NR^{c1})_2$ or $-C_1-C_6$ alkyl $C(O)OR^k$, R^{c1} is H, C_1-C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^{13} and $R^{13'}$ are each independently H or C_1-C_4 alkyl;

L is absent, C_1-C_{12} alkylene, or C_2-C_{12} alkenylene, wherein the C_1-C_{12} alkylene or C_2-C_{12} alkenylene is optionally substituted with C_1-C_4 alkyl;

R^{21} is H, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or $-C(O)OR^k$, wherein the C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J^1 , and wherein optionally 1 to 2 carbon atoms of the spiro C_5-C_{12} cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and

optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and


n is 0 or 1.

[0139] In some embodiments, the compound has the structure of formula (III d'):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R⁶, R¹³, R^{13'}, L and R²¹ are as defined for formula (III d).

[0140] In some embodiments, the compound has the structure of formula (III d''):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^6 , R^{13} , $R^{13'}$, L and R^{21} are as defined for formula (IIIc).

[0141] In some embodiments for the compound of any of the foregoing embodiments, n is 0.

[0142] In some embodiments for the compound of any of the foregoing embodiments, each of R^{13} and $R^{13'}$ is H.

[0143] In some embodiments, the compound has the structure of formula (IIIe):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1-C_6$ alkyl, $-C_1-C_6$ alkyl- $(NR^{c1})_2$ or $-C_1-C_6$ alkyl $C(O)OR^k$, R^{c1} is H, C_1-C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

L is absent, C_1-C_{12} alkylene, or C_2-C_{12} alkenylene, wherein the C_1-C_{12} alkylene or C_2-C_{12} alkenylene is optionally substituted with C_1-C_4 alkyl;

R^{21} is H, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl,


spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicycyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicycyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;

each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion; and


J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl.

[0144] In some embodiments, the compound has the structure of formula (IIIe'):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R⁶, L and R²¹ are as defined for formula (IIIe).

[0145] In some embodiments, the compound has the structure of formula (IIIe''):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^6 , L and R^{21} are as defined for formula (IIIe).

[0146] In some embodiments, the compound has the structure of formula (III f):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1-C_6$ alkyl, $-C_1-C_6$ alkyl- $(NR^{c1})_2$ or $-C_1-C_6$ alkyl $C(O)OR^k$, R^{c1} is H, C_1-C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

L is absent, C_1-C_{12} alkylene, or C_2-C_{12} alkenylene, wherein the C_1-C_{12} alkylene or C_2-C_{12} alkenylene is optionally substituted with C_1-C_4 alkyl;

R^{21} is H, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or $-C(O)OR^k$, wherein the C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J^1 , and wherein optionally 1 to 2 carbon atoms of the spiro C_5-


C₁₂cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;

each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion; and

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl.


[0147] In some embodiments, the compound has the structure of formula (III^f):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein R⁶, L and R²¹ are as defined for formula (III^f).

[0148] In some embodiments, the compound has the structure of formula (III^{f'}):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R⁶, L and R²¹ are as defined for formula (III^f).

[0149] In some embodiments of the compounds of (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''),

(IIIId), (IIIId'), (IIIId''), (IIIe), (IIIe'), (IIIe''), (IIIIf), (IIIIf'), and (IIIIf''), R²¹ is C₃-C₇cycloalkyl, wherein the C₃-C₇cycloalkyl is optionally substituted with 1 to 3 of J¹, wherein J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl. In some of the foregoing embodiments, the C₃-C₇cycloalkyl is selected from the group consisting of cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl.

[0150] In some embodiments of the compounds of (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (IIIIf), (IIIIf'), and (IIIIf''), R²¹ is a heterocyclyl, wherein the heterocyclyl is optionally substituted with 1 to 3 of J¹, wherein J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl. In some embodiments, the heterocyclyl is selected from oxiranyl, oxetanyl, azetidynyl, oxazolyl, thiazolidinyl, thiazolyl, morpholinyl, pyrrolidinonyl, pyrrolidinyl, piperidinyl, piperazinyl, 2,3-dihydrofuranyl, dihydropyranyl, tetrahydrofuranyl, tetrahydropyranyl, dihydropyridinyl, tetrahydropyridinyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, and azapanyl, wherein the heterocyclyl is optionally substituted with 1 to 3 of OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl.

[0151] In some embodiments of the compounds of (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (IIIIf), (IIIIf'), and (IIIIf''), R²¹ is aryl, wherein the aryl is optionally substituted with 1 to 3 of J¹, wherein J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl. In some of the foregoing embodiments, R²¹ is a phenyl or naphthyl, wherein the phenyl or naphthyl is optionally substituted with 1 to 3 of OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl.


[0152] In some embodiments of the compounds of (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (IIIIf), (IIIIf'), and (IIIIf''), R²¹ is heteroaryl, wherein the heteroaryl is optionally substituted with 1 to 3 of J¹, wherein J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl. In some of the foregoing embodiments, the heteroaryl is selected from the group consisting of pyrrolyl, pyrazolyl, imidazolyl, pyrazinyl, oxazolyl, isoxazolyl, thiazolyl, furyl, thienyl, pyridyl, pyrimidyl, benzoxazolyl, benzothiazolyl, purinyl, benzimidazolyl, indolyl, isoquinolyl, quinoxaliny, and quinolyl, wherein the heteroaryl is optionally substituted with 1 to 3 of OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl.

[0153] In some embodiments of the compounds of (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (IIIIf), (IIIIf'), and (IIIIf''), R²¹ is adamantyl, wherein the adamantyl is optionally substituted with J¹, wherein J¹ is selected from OH, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl.

[0154] In some embodiments of the compounds of (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (IIIf), (IIIf'), and (IIIf''), R^{21} is spiro C_5 - C_{12} cycloalkyl, wherein the spiro C_5 - C_{12} cycloalkyl has 0-2 carbon atoms replaced with 0-2 heteroatoms selected from N, O and S, and is optionally substituted with 1 to 3 of J^1 , wherein J^1 is selected from OH, CN, halo, C_1 - C_4 alkyl, and halo C_1 - C_4 alkyl, or when an N atom is present, is optionally substituted with an N-protecting group.

[0155] In some embodiments of the compounds of (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (IIIf), (IIIf'), and (IIIf''), R^{21} is 5 to 12 membered bridged bicycyl, wherein the bridged bicycyl has 0-2 carbon atoms replaced with 0-2 heteroatoms selected from N, O and S, and is optionally substituted with 1 to 3 of J^1 , wherein J^1 is selected from OH, CN, halo, C_1 - C_4 alkyl, and halo C_1 - C_4 alkyl, or when an N atom is present, is optionally substituted with an N-protecting group.

[0156] In some embodiments of the compounds of formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (IIIf), (IIIf'), and (IIIf''), R^{21} is selected from the following:




wherein J^1 is OH, CN, halo, C_1 - C_4 alkyl, and halo C_1 - C_4 alkyl, and n is 0-3. In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2. In some embodiments, halo C_1 - C_4 alkyl is $-CH_2F$, $-CHF_2$, or $-CF_3$.

[0157] In some embodiments, for any of the compounds herein, L is C_3 - C_{12} alkylene. In some embodiments, for any of the compounds herein, L is C_3 - C_6 alkylene. In some embodiments, for any of the compounds herein, L is C_1 - C_6 alkylene.


[0158] In some embodiments, for any of the compounds herein, L is C_3 - C_{12} alkenylene. In some embodiments, for any of the compounds herein, L is C_3 - C_6 alkenylene. In some embodiments, for any of the compounds herein, L is C_1 - C_6 alkenylene.

[0159] In some embodiments, L is C_1 - C_{12} alkylene, or C_2 - C_{12} alkenylene, wherein the C_1 - C_{12} alkylene or C_2 - C_{12} alkenylene is optionally substituted with C_1 - C_4 alkyl; and R^{21} is H.

[0160] In some embodiments of the compounds of formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (III f), (III f'), and (III f''), $-L-R^{21}$ is a C_2 - C_6 alkenyl selected from:




[0161] In some embodiments for any of the foregoing compounds, e.g., formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (III f), (III f'), and (III f''), wherein R^6 is



each occurrence of R^A is independently hydrogen (glycine), methyl (alanine), propan-2-yl (valine), propan-1-yl (norvaline), 2-methylpropan-1-yl (leucine), 1-methylpropan-1-yl (isoleucine), butan-1-yl (norleucine), phenyl (2-phenylglycine), benzyl (phenylalanine), p-hydroxybenzyl (tyrosine), indol-3-ylmethyl (tryptophan), imidazol-4-ylmethyl (histidine), hydroxymethyl (serine), 2-hydroxyethyl (homoserine), 1-hydroxyethyl (threonine), mercaptomethyl (cysteine), methylthiomethyl (S-methylcysteine), 2-mercaptoethyl (homocysteine), 2-methylthioethyl (methionine), carbamoylmethyl (asparagine), 2-carbamoylethyl (glutamine), carboxymethyl (aspartic acid), 2-carboxyethyl (glutamic acid), 4-aminobutan-1-yl (lysine), 4-amino-3-hydroxybutan-1-yl


(hydroxylysine), 3-aminopropan-1-yl (ornithine), 3-guanidinopropan-1-yl (arginine), or 3-ureido-propan-1-yl (citrulline);

each occurrence of R^B is independently H, or R^B together with the adjacent R_A and the N atom form a prolyl side chain:



p is 0, 1 or 2.

[0162] In some embodiments for any of the foregoing compounds, e.g., formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (IIIf), (IIIf'), and (IIIf''), wherein R^6 is




each occurrence of R^A is independently methyl (alanine), propan-2-yl (valine), 2-methylpropan-1-yl (leucine), imidazol-4-ylmethyl (histidine), hydroxymethyl (serine), 1-hydroxyethyl (threonine), carbamoylmethyl (asparagine), 2-carbamoylethyl (glutamine), 4-aminobutan-1-yl (lysine), carboxymethyl (aspartic acid), 3-guanidinopropan-1-yl (arginine), benzyl (phenylalanine), or 4-aminobutan-1-yl (lysine);

R^B is H; and

p 0, 1, or 2.

[0163] In some embodiments for any of the foregoing compounds, e.g., formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (IIIf), (IIIf'), and (IIIf''), wherein R^6 is



each occurrence of R^A is independently propan-2-yl (valine), 2-methylpropan-1-yl (leucine), carboxymethyl (aspartic acid), benzyl (phenylalanine), or 4-aminobutan-1-yl (lysine);


each R^B is H; and

p is 0, 1, or 2.

[0164] In some embodiments for any of the foregoing compounds, e.g., formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (IIIf), (IIIf'), and (IIIf''), p is 0.

[0165] In some embodiments for any of the foregoing compounds, e.g., formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (IIIf), (IIIf'), and (IIIf''), p is 1.

[0166] In some embodiments for any of the foregoing compounds, e.g., formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (IIIf), (IIIf'), and (IIIf''), wherein R⁶ is



and p is 1;

first of R^A is propan-2-yl (valine) and second of R^A is propan-2-yl (valine); and each of R^B is H (i.e., dipeptide Val-Val); or


first of R^A is 2-methylpropan-1-yl (leucine), and second of R^A is 2-methylpropan-1-yl (leucine); and each of R^B is H (i.e., dipeptide Leu-Leu); or

first of R^A is methyl (alanine) and second of R^A is methyl (alanine); and each of R^B is H (i.e., dipeptide Ala-Ala); or

first of R^A is 4-aminobutan-1-yl (lysine); second of R^A is 4-aminobutan-1-yl (lysine); and each of R^B is H (i.e., dipeptide Lys-Lys); or

first of R^A is hydrogen; second of R^A is 4-aminobutan-1-yl, and each of R^B is H (i.e., dipeptide Gly-Lys).


[0167] In some embodiments for any of the foregoing compounds, e.g., formula (I), (Ia), (Ib), (II), (II'), (II''), (IIa), (IIa'), (IIa''), (IIb), (IIb'), (IIb''), (III), (III'), (III''), (IIIa), (IIIa'), (IIIa''), (IIIb), (IIIb'), (IIIb''), (IIIc), (IIIc'), (IIIc''), (IIId), (IIId'), (IIId''), (IIIe), (IIIe'), (IIIe''), (IIIf), (IIIf'), and (IIIf''), wherein R⁶ is



















each of the α -carbon of the amino acid other than glycine is in the L or D configuration. In some embodiments, the α -carbon of the amino acid other than glycine is in the L configuration









[0168] In some embodiments, the compound is selected from the group consisting of the compounds or a pharmaceutical salt thereof in **Table 1**:









[0169] **Table 1**

















K101-C1359




K101-C135901




K101-C135902




K101-C1361




K101-C136101









K101-C136102





K101-C1364




K101-C1365





[0170] In some embodiments, for each of the compounds of **Table 1**, the -OH on the C20 carbon atom, where appropriate, is substituted with



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;


each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

[0171] In some embodiments of the amino acid moiety on the C20 carbon,

each occurrence of R^A is independently hydrogen (glycine), methyl (alanine), propan-2-yl (valine), propan-1-yl (norvaline), 2-methylpropan-1-yl (leucine), 1-methylpropan-1-yl (isoleucine), butan-1-yl (norleucine), phenyl (2-phenylglycine), benzyl (phenylalanine), p-hydroxybenzyl (tyrosine), indol-3-ylmethyl (tryptophan), imidazol-4-ylmethyl (histidine), hydroxymethyl (serine), 2-hydroxyethyl (homoserine), 1-hydroxyethyl (threonine), mercaptomethyl (cysteine), methylthiomethyl (S-methylcysteine), 2-mercaptoethyl (homocysteine), 2-methylthioethyl (methionine), carbamoylmethyl (asparagine), 2-carbamoylethyl (glutamine), carboxymethyl (aspartic acid), 2-carboxyethyl (glutamic acid), 4-aminobutan-1-yl (lysine), 4-amino-3-hydroxybutan-1-yl (hydroxylysine), 3-aminopropan-1-yl (ornithine), 3-guanidinopropan-1-yl (arginine), or 3-ureidopropan-1-yl (citrulline);

each occurrence of R^B is independently H, or R^B together with the adjacent R_A and the N atom form a prolyl side chain:



p is 0, 1 or 2.

[0172] In some embodiments, each occurrence of R^A is independently methyl (alanine), propan-2-yl (valine), 2-methylpropan-1-yl (leucine), imidazol-4-ylmethyl (histidine), hydroxymethyl (serine), 1-hydroxyethyl (threonine), carbamoylmethyl (asparagine), 2-carbamoylethyl (glutamine), 4-aminobutan-1-yl (lysine), carboxymethyl (aspartic acid), 3-guanidinopropan-1-yl (arginine), benzyl (phenylalanine), or 4-aminobutan-1-yl (lysine);

R^B is H; and

p 0, 1, or 2.

[0173] In some embodiments, each occurrence of R^A is independently propan-2-yl (valine), 2-methylpropan-1-yl (leucine), carboxymethyl (aspartic acid), benzyl (phenylalanine), or 4-aminobutan-1-yl (lysine);

each R^B is H; and

p is 0, 1, or 2.

[0174] In some embodiments, p is 0.

[0175] In some embodiments, p is 1.

[0176] In some embodiments,

p is 1;

first of R^A is propan-2-yl (valine) and second of R^A is propan-2-yl (valine); and each of R^B is H (i.e., dipeptide Val-Val); or

first of R^A is 2-methylpropan-1-yl (leucine), and second of R^A is 2-methylpropan-1-yl (leucine); and each of R^B is H (i.e., dipeptide Leu-Leu); or


first of R^A is methyl (alanine) and second of R^A is methyl (alanine); and each of R^B is H (i.e., dipeptide Ala-Ala); or

first of R^A is 4-aminobutan-1-yl (lysine); second of R^A is 4-aminobutan-1-yl (lysine); and each of R^B is H (i.e., dipeptide Lys-Lys); or

first of R^A is hydrogen; second of R^A is 4-aminobutan-1-yl, and each of R^B is H (i.e., dipeptide Gly-Lys).

[0177] In some embodiments, each of the α -carbon of the amino acid other than glycine is in the L or D configuration. In some embodiments, each of the α -carbon of the amino acid other than glycine is in the L configuration.

[0178] In another aspect, the present disclosure provides a compound of formula (IV):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^2 is a C_1 - C_4 alkyl;


R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

$R^{5'}$ and $R^{6'}$ are H, or $R^{5'}$ and $R^{6'}$ form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

$R^{6'}$ and $R^{7'}$ are H, or $R^{6'}$ and $R^{7'}$ form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R^7 is H or OH;

R^9 is OR^e , wherein R^e is H, C₁-C₆alkyl, or aryl;

R^{11} is C₁-C₄alkyl;

R^{12} is H, -OH, $-OC(O)R^f$, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R^{14} is H or OR^g ; wherein R^g is H or C₁-C₆alkyl;

R^{17} and R^{18} are each independently C₁-C₄alkyl or C₁-C₄alkyl- OR^h , wherein R^h is H or C₁-C₆alkyl;

L is C₀-C₆alkylarylene, C₀-C₆alkylheteroarylene, C₀-C₆alkylC₃-C₇cycloalkylene, C₁-C₁₂alkylene or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with OH or C₁-C₄alkyl; and

R^{21} is H, -OH, -SH, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicycyl, adamantyl, or $-C(O)OR^k$, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicycyl, or adamantyl is optionally substituted with 1 to 3 of J^1 ; and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl or, when an N atom is present an N-protecting group;


each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J^1 ;

J^1 is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

R^k is H or M^+ counterion.

[0179] In some embodiments, excluded from the compounds of formula (IV) are compounds in which:

-L- R^{21} is



R^2 and R^{11} are CH₃;

R^3 is =O;

R^4 is OH;

R^5 , R^5' , R^7 and R^{14} are H;

$R^{6'}$ and $R^{7'}$ together form a bond;

R⁹ is OH;

R¹² is H;


R¹⁷ and R¹⁸ are -CH₃;

R^A is propan-2-yl (valine);

R^B is H; and

p is 0.

[0180] In some embodiments, the compound has the structure of formula (IVa):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein


R² is a C₁-C₄alkyl;

R³ is O double bonded to the ring carbon when (- -) is a bond, or -OR^a, wherein R^a is H or -C(O)R^{al}, wherein R^{al} is C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R⁴ and R⁵ are each independently H or -OR^b, wherein R^b is H, C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R⁶ is OH, halo, or -OC(O)R^c, wherein R^c is -C₁-C₆alkyl, -C₁-C₆alkyl-(NR^{cl})₂ or -C₁-C₆alkylC(O)OR^k, R^{cl} is H, C₁-C₆alkyl, or two R^{cl} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R⁶ is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R⁷ is H or OH;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹² is H, -OH, -OC(O)R^f, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is C₀-C₆alkylarylene, C₀-C₆alkylheteroarylene, C₀-C₆alkylC₃-C₇cycloalkylene, C₁-C₁₂alkylene or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with OH or C₁-C₄alkyl; and

R²¹ is H, -OH, -SH, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicycyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicycyl, or adamantyl is optionally substituted with 1 to 3 of J¹; and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl or, when an N atom is present an N-protecting group;


each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

R^k is H or M⁺ counterion.

[0181] In some embodiments, excluded from the compounds of formula (IVa) are compounds in which:

-L-R²¹ is



R² and R¹¹ are CH₃;

R³ is =O;

R⁴ is OH;

R^5 , R^7 and R^{14} are H;

R^9 is OH;

R^{12} is H;


R^{17} and R^{18} are $-\text{CH}_3$;

R^A is propan-2-yl (valine);

R^B is H; and

p is 0.

[0182] In some embodiments, the compound has the structure of formula (IVb)



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein


R^2 is a C_1 - C_4 alkyl;

R^3 is O double bonded to the ring carbon when (---) is a bond, or $-\text{OR}^a$, wherein R^a is H or $-\text{C}(\text{O})\text{R}^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-\text{OR}^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^6 is OH, halo, or $-\text{OC}(\text{O})\text{R}^c$, wherein R^c is $-\text{C}_1$ - C_6 alkyl, $-\text{C}_1$ - C_6 alkyl- $(\text{NR}^{c1})_2$ or $-\text{C}_1$ - C_6 alkyl $\text{C}(\text{O})\text{OR}^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^7 is H or OH;

R^9 is OR^e , wherein R^e is H, C₁-C₆alkyl, or aryl;

R^{11} is C₁-C₄alkyl;

R^{12} is H, -OH, $-OC(O)R^f$, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R^{14} is H or OR^g , wherein R^g is H or C₁-C₆alkyl;

R^{17} and R^{18} are each independently C₁-C₄alkyl or C₁-C₄alkyl- OR^h , wherein R^h is H or C₁-C₆alkyl;

L is C₀-C₆alkylarylene, C₀-C₆alkylheteroarylene, C₀-C₆alkylC₃-C₇cycloalkylene, C₁-C₁₂alkylene or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with OH or C₁-C₄alkyl; and


R^{21} is H, -OH, -SH, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or $-C(O)OR^k$, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J^1 ; and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl or, when an N atom is present an N-protecting group;

each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J^1 ;


J^1 is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

R^k is H or M^+ counterion.

[0183] In some embodiments of the compound of formula (IV), (IVa), and (IVb), the C₂-C₆alkenyl of R^{a1} is independently selected from:




[0184] In some embodiments, R^{a1} is independently selected from:




[0185] In some embodiments, R^3 is $-OR^a$; wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1-C_6 alkyl, C_2-C_6 alkenyl, C_0-C_6 alkylaryl, or C_0-C_6 alkylheteroaryl. In some embodiments, the aryl of C_0-C_6 alkylaryl is phenyl. In some embodiments, the aryl of C_0-C_6 alkylaryl is optionally substituted with 1 to 3 of OH, CN, halo, C_1-C_4 alkyl, and halo C_1-C_4 alkyl.

[0186] In some embodiments of the compound of formula (IV), (IVa), and (IVb), R^3 is O double bonded to the carbon atom.

[0187] In some embodiments of the compound of formula (IV), (IVa), and (IVb), the C_2-C_{12} alkenyl of R^f is independently selected from:



[0188] In some embodiments, R^f is independently selected from:




[0189] In some embodiments of the compound of formula (IV), (IVa), and (IVb), one or more of R^2 , R^{11} , R^{17} , and R^{18} are $-CH_3$. In some embodiments, each of R^2 , R^{11} , R^{17} , and R^{18} is $-CH_3$.

[0190] In some embodiments of the compound of formula (IV), (IVa), and (IVb), R^4 and R^5 are each independently H or $-OH$.

[0191] In some embodiments of the compound of formula (IV), (IVa), and (IVb), R^2 , R^{11} , R^{17} , and R^{18} are $-CH_3$; R^3 is O double bonded to the carbon atom; and R^4 and R^5 are each independently H or $-OH$.

[0192] In some embodiments, the compound has the structure of formula (V):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^2 is a C_1 - C_4 alkyl;


R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

$R^{5'}$ and $R^{6'}$ are H, or $R^{5'}$ and $R^{6'}$ form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

$R^{6'}$ and $R^{7'}$ are H, or $R^{6'}$ and $R^{7'}$ form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R⁷ is H or OH;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is C₀-C₆alkylarylene, C₀-C₆alkylheteroarylene, C₀-C₆alkylC₃-C₇cycloalkylene, C₁-C₁₂alkylene or C₂-C₁₂alkenylylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylylene is optionally substituted with OH or C₁-C₄alkyl; and

R²¹ is H, -OH, -SH, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹; and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl or, when an N atom is present an N-protecting group;


each R^j is independently C₁-C₆alkyl, C₂-C₆alkenylyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

R^k is H or M⁺ counterion.

[0193] In some embodiments, excluded from the compounds of formula (V) are compounds in which:

-L-R²¹ is



R² and R¹¹ are CH₃;

R³ is =O;

R⁴ is OH;

R⁵, R^{5'}, R⁷ and R¹⁴ are H;

R^{6*} and R^{7*} together form a bond;

R⁹ is OH;


R¹⁷ and R¹⁸ are -CH₃;

R^A is propan-2-yl (valine);

R^B is H; and

p is 0.

[0194] In some embodiments, the compound has the structure of formula (Va):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein


R^2 is a C_1 - C_4 alkyl;

R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R⁷ is H or OH;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is C₀-C₆alkylarylene, C₀-C₆alkylheteroarylene, C₀-C₆alkylC₃-C₇cycloalkylene, C₁-C₁₂alkylene or C₂-C₁₂alkenylylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylylene is optionally substituted with OH or C₁-C₄alkyl; and

R²¹ is H, -OH, -SH, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹; and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl or, when an N atom is present an N-protecting group;


each R^j is independently C₁-C₆alkyl, C₂-C₆alkenylyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

R^k is H or M⁺ counterion.

[0195] In some embodiments, excluded from the compounds of formula (V) are compounds in which:

-L-R²¹ is



R² and R¹¹ are CH₃;

R³ is =O;

R⁴ is OH;

R⁵, R⁷ and R¹⁴ are H;

R⁹ is OH;


R¹⁷ and R¹⁸ are -CH₃;

R^A is propan-2-yl (valine);

R^B is H; and

p is 0.

[0196] In some embodiments, the compound has the structure of formula (Vb):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein


R^2 is a C_1 - C_4 alkyl;

R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$; wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^9 is OR^e , wherein R^e is H, C_1 - C_6 alkyl, or aryl;

R^{11} is C_1 - C_4 alkyl;

R^{14} is H or OR^g ; wherein R^g is H or C_1 - C_6 alkyl;

R^{17} and R^{18} are each independently C_1 - C_4 alkyl or C_1 - C_4 alkyl- OR^h , wherein R^h is H or C_1 -

C₆alkyl;

L is C₀-C₆alkylarylene, C₀-C₆alkylheteroarylene, C₀-C₆alkylC₃-C₇cycloalkylene, C₁-C₁₂alkylene or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with OH or C₁-C₄alkyl; and


R²¹ is H, -OH, -SH, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹; and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl or, when an N atom is present an N-protecting group;

each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;


J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

R^k is H or M⁺ counterion.

[0197] In some embodiments of the compound of formula (V), (Va), and (Vb), R³ is -OR^a; wherein R^a is H or -C(O)R^{a1}, wherein R^{a1} is C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl. In some embodiments, the aryl of C₀-C₆alkylaryl is phenyl. In some embodiments, the aryl of C₀-C₆alkylaryl is optionally substituted with 1 to 3 of OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl. In some embodiments, R^{a1} is selected from:



[0198] In some embodiments, R^{a1} is selected from:



[0199] In some embodiments of the compound of formula (V), (Va), and (Vb),

R³ is O double bonded to the carbon atom.

[0200] In some embodiments of the compound of formula (V), (Va), and (Vb), one or more of R², R¹¹, R¹⁷, and R¹⁸ are -CH₃. In some embodiments, each of R², R¹¹, R¹⁷, and R¹⁸ is -CH₃.

[0201] In some embodiments of the compound of formula (V), (Va), and (Vb), R⁴ and R⁵ are each independently H or -OH.


[0202] In some embodiments of the compound of formula (V), (Va), and (Vb),

R^2 , R^{11} , R^{17} , and R^{18} are $-\text{CH}_3$;

R^3 is O double bonded to the carbon atom; and

R^4 and R^5 are each independently H or $-\text{OH}$.

[0203] In some embodiments, the compound has the structure of formula (Vc):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^6 is OH, halo, or $-\text{OC}(\text{O})\text{R}^c$, wherein R^c is $-\text{C}_1\text{-C}_6\text{alkyl}$, $-\text{C}_1\text{-C}_6\text{alkyl}-(\text{NR}^{\text{cl}})_2$ or $-\text{C}_1\text{-C}_6\text{alkylC}(\text{O})\text{OR}^k$, R^{cl} is H, $\text{C}_1\text{-C}_6\text{alkyl}$, or two R^{cl} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

L is $\text{C}_0\text{-C}_6\text{alkylarylene}$, $\text{C}_0\text{-C}_6\text{alkylheteroarylene}$, $\text{C}_0\text{-C}_6\text{alkylC}_3\text{-C}_7\text{cycloalkylene}$, $\text{C}_1\text{-C}_{12}\text{alkylene}$ or $\text{C}_2\text{-C}_{12}\text{alkenylene}$, wherein the $\text{C}_1\text{-C}_{12}\text{alkylene}$ or $\text{C}_2\text{-C}_{12}\text{alkenylene}$ is optionally substituted with OH or $\text{C}_1\text{-C}_4\text{alkyl}$; and

R^{21} is H, $-\text{OH}$, $-\text{SH}$, $-\text{S}(\text{O})_2\text{R}^j$, $-\text{SR}^j$, $-\text{N}(\text{R}^j)_2$, $-\text{Si}(\text{R}^j)_3$, $\text{C}_3\text{-C}_7\text{cycloalkyl}$, heterocyclyl, aryl,

heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicycyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicycyl, or adamantyl is optionally substituted with 1 to 3 of J¹; and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl or, when an N atom is present an N-protecting group;


each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

R^k is H or M⁺ counterion.

[0204] In some embodiments, excluded from the compounds of formula (Vc) are compounds in which:

-L-R²¹ is




R^A is propan-2-yl (valine);


R^B is H; and

p is 0.

[0205] In some embodiments, specifically excluded from the compounds of formula (IV), (IVa), (V), (Va) and (Vc) are compounds of the following structure:



[0206] In some embodiments, the compound has the structure of formula (Vd):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1-C_6$ alkyl, $-C_1-C_6$ alkyl- $(NR^{c1})_2$ or $-C_1-C_6$ alkyl $C(O)OR^k$, R^{c1} is H, C_1-C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

L is C_0-C_6 alkylarylene, C_0-C_6 alkylheteroarylene, C_0-C_6 alkyl C_3-C_7 cycloalkylene, C_1-C_{12} alkylene or C_2-C_{12} alkenylene, wherein the C_1-C_{12} alkylene or C_2-C_{12} alkenylene is optionally substituted with OH or C_1-C_4 alkyl; and

R^{21} is H, -OH, -SH, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, 5 to 12 membered bridged bicycyl, adamantyl, or $-C(O)OR^k$, wherein the C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, bridged bicycyl, or adamantyl is optionally substituted with 1 to 3 of J^1 ; and wherein optionally 1 to 2 carbon atoms of the spiro C_5-C_{12} cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C_1-C_4 alkyl or, when an N atom is present an N-protecting group;

each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

R^k is H or M⁺ counterion.

[0207] In some embodiments of the compounds of (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc) and (Vd), R²¹ is C₃-C₇cycloalkyl, wherein the C₃-C₇cycloalkyl is optionally substituted with 1 to 3 of J¹. In some of the foregoing embodiments, the C₃-C₇cycloalkyl is selected from the group consisting of cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl.

[0208] In some embodiments of the compounds of (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc) and (Vd), R²¹ is a heterocyclyl, wherein the heterocyclyl is optionally substituted with 1 to 3 of J¹. In some embodiments, the heterocycloalkyl is selected from oxiranyl, oxetanyl, azetidynyl, oxazolyl, thiazolidinyl, thiazolyl, morpholinyl, pyrrolidinonyl, pyrrolidinyl, piperidinyl, piperazinyl, 2,3-dihydrofuranyl, dihydropyranyl, tetrahydrofuranyl, tetrahydropyranyl, dihydropyridinyl, tetrahydropyridinyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, and azapanyl.

[0209] In some embodiments of the compounds of (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc), and (Vd), R²¹ is aryl, wherein the aryl is optionally substituted with 1 to 3 of J¹. In some of the foregoing embodiments, R²¹ is a phenyl, wherein the phenyl is optionally substituted with 1 to 3 of OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl.


[0210] In some embodiments of the compounds of (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc), and (Vd), R²¹ is heteroaryl, wherein the heteroaryl is optionally substituted with 1 to 3 of J¹. In some of the foregoing embodiments, the heteroaryl is selected from the group consisting of pyrrolyl, pyrazolyl, imidazolyl, pyrazinyl, oxazolyl, isoxazolyl, thiazolyl, furyl, thienyl, pyridyl, pyrimidyl, benzoxazolyl, benzothiazolyl, purinyl, benzimidazolyl, indolyl, isoquinolyl, quinoxaliny, and quinolyl, wherein the heteroaryl is optionally substituted with 1 to 3 of OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl.

[0211] In some embodiments of the compounds of (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc), and (Vd), R²¹ is adamantyl, wherein the adamantyl is optionally substituted with OH, halo, or C₁-C₄alkyl.

[0212] In some embodiments of the compounds of (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc), and (Vd), R²¹ is spiroC₅-C₁₂ cycloalkyl, wherein the spiroC₅-C₁₂ cycloalkyl has 0-2 carbon atoms replaced with 0-2 heteroatoms selected from N, O and S, and is optionally substituted with 1 to 3 of OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl, or when an N atom is present an N-protecting group.

[0213] In some embodiments of the compounds of (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc) and (Vd), R^{21} is 5 to 12 membered bridged bicycyl, wherein the bridged bicycyl has 0-2 carbon atoms replaced with 0-2 heteroatoms selected from N, O and S, and is optionally substituted with 1 to 3 of OH, CN, halo, C_1 - C_4 alkyl, and halo C_1 - C_4 alkyl, or when an N atom is present an N-protecting group.

[0214] In some embodiments of the compounds of formula (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc) and (Vd), R^{21} is selected from the following:




wherein J^1 is OH, CN, halo, C_1 - C_4 alkyl, and halo C_1 - C_4 alkyl, and n is 0-3. In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2. In some embodiments, halo C_1 - C_4 alkyl is $-CH_2F$, $-CHF_2$, or $-CF_3$.

[0215] In some embodiments, for any of the compounds herein, L is C_3 - C_{12} alkylene. In some embodiments, for any of the compounds herein, L is C_3 - C_6 alkylene. In some embodiments, for any of the compounds herein, L is C_1 - C_6 alkylene.


[0216] In some embodiments, for any of the compounds herein, L is C_3 - C_{12} alkenylene. In some embodiments, for any of the compounds herein, L is C_3 - C_6 alkenylene. In some embodiments, for any of the compounds herein, L is C_1 - C_6 alkenylene.

[0217] In some embodiments of the compounds of formula (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc) and (Vd), $-L-R^{21}$ is a C_2 - C_6 alkenyl selected from:




[0218] In some embodiments for any of the foregoing compounds, e.g., formula (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc) and (Vd), wherein

R^6 is



each occurrence of R^A is independently hydrogen (glycine), methyl (alanine), propan-2-yl (valine), propan-1-yl (norvaline), 2-methylpropan-1-yl (leucine), 1-methylpropan-1-yl (isoleucine), butan-1-yl (norleucine), phenyl (2-phenylglycine), benzyl (phenylalanine), p-hydroxybenzyl (tyrosine), indol-3-ylmethyl (tryptophan), imidazol-4-ylmethyl (histidine), hydroxymethyl (serine), 2-hydroxyethyl (homoserine), 1-hydroxyethyl (threonine), mercaptomethyl (cysteine), methylthiomethyl (S-methylcysteine), 2-mercaptoethyl (homocysteine), 2-methylthioethyl (methionine), carbamoylmethyl (asparagine), 2-carbamoylethyl (glutamine), carboxymethyl (aspartic acid), 2-carboxyethyl (glutamic acid), 4-aminobutan-1-yl (lysine), 4-amino-3-hydroxybutan-1-yl (hydroxylysine), 3-aminopropan-1-yl (ornithine), 3-guanidinopropan-1-yl (arginine), or 3-ureidopropan-1-yl (citrulline);

each occurrence of R^B is H, or R^B together with the adjacent R^A and the N atom form a prolyl side chain:



p is 0, 1 or 2.

[0219] In some embodiments for any of the foregoing compounds, e.g., formula (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc) and (Vd),


each R^A is independently methyl (alanine), propan-2-yl (valine), 2-methylpropan-1-yl (leucine), imidazol-4-ylmethyl (histidine), hydroxymethyl (serine), 1-hydroxyethyl (threonine), carbamoylmethyl (asparagine), 2-carbamoylethyl (glutamine), 4-aminobutan-1-yl (lysine), carboxymethyl (aspartic acid), 3-guanidinopropan-1-yl (arginine), benzyl (phenylalanine), or 4-aminobutan-1-yl (lysine);

R^B is H; and

p 0, 1, or 2.

[0220] In some embodiments for any of the foregoing compounds, e.g., formula (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc) and (Vd), wherein

R⁶ is



each occurrence of R^A is independently propan-2-yl (valine), 2-methylpropan-1-yl (leucine), carboxymethyl (aspartic acid), benzyl (phenylalanine), or 4-aminobutan-1-yl (lysine);

each R^B is H; and


p is 0, 1, or 2.

[0221] In some embodiments for any of the foregoing compounds, e.g., formula (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc) and (Vd), p is 0.

[0222] In some embodiments for any of the foregoing compounds, e.g., formula (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc) and (Vd), p is 1.

[0223] In some embodiments for any of the foregoing compounds, e.g., formula (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc) and (Vd), wherein

R⁶ is



and p is 1;

first of R^A is propan-2-yl (valine) and second of R^A is propan-2-yl (valine); and each of R^B is H (i.e., dipeptide Val-Val); or

first of R^A is 2-methylpropan-1-yl (leucine), and second of R^A is 2-methylpropan-1-yl (leucine); and each of R^B is H (i.e., dipeptide Leu-Leu); or


first of R^A is methyl (alanine) and second of R^A is methyl (alanine); and each of R^B is H (i.e., dipeptide Ala-Ala); or

first of R^A is 4-aminobutan-1-yl (lysine); second of R^A is 4-aminobutan-1-yl (lysine); and each of R^B is H (i.e., dipeptide Lys-Lys); or

first of R^A is hydrogen; second of R^A is 4-aminobutan-1-yl, and each of R^B is H (i.e., dipeptide Gly-Lys).

[0224] In some embodiments for any of the foregoing compounds, e.g., formula (IV), (IVa), (IVb), (V), (Va), (Vb), (Vc) and (Vd), wherein





R^6 is




each of the α -carbon of the amino acid other than glycine is in the L or D configuration. In some embodiments, each of the α -carbon of the amino acid other than glycine is in the L configuration.


[0225] In some embodiments, the compound is selected from the group consisting of the compounds or a pharmaceutical salt thereof, in **Table 2**.

[0226] **Table 2**







K101-C1306




K101-C1308





K101-C1311




K101-C1312




K101-C1313





or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof.

[0227] In some embodiments, for each of the compounds of **Table 2**, the -OH on the C20 carbon atom is substituted with



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and


p is 0, 1, or 2.

[0228] In some embodiments of the amino acid moiety on the C20 carbon,

each occurrence of R^A is independently hydrogen (glycine), methyl (alanine), propan-2-yl (valine), propan-1-yl (norvaline), 2-methylpropan-1-yl (leucine), 1-methylpropan-1-yl (isoleucine), butan-1-yl (norleucine), phenyl (2-phenylglycine), benzyl (phenylalanine), p-hydroxybenzyl (tyrosine), indol-3-ylmethyl (tryptophan), imidazol-4-ylmethyl (histidine), hydroxymethyl (serine), 2-

hydroxyethyl (homoserine), 1-hydroxyethyl (threonine), mercaptomethyl (cysteine), methylthiomethyl (S-methylcysteine), 2-mercaptoethyl (homocysteine), 2-methylthioethyl (methionine), carbamoylmethyl (asparagine), 2-carbamoylethyl (glutamine), carboxymethyl (aspartic acid), 2-carboxyethyl (glutamic acid), 4-aminobutan-1-yl (lysine), 4-amino-3-hydroxybutan-1-yl (hydroxylysine), 3-aminopropan-1-yl (ornithine), 3-guanidinopropan-1-yl (arginine), or 3-ureido-propan-1-yl (citrulline);

each occurrence of R^B is independently H, or R^B together with the adjacent R_A and the N atom form a prolyl side chain:



p is 0, 1 or 2.

[0229] In some embodiments, each occurrence of R^A is independently methyl (alanine), propan-2-yl (valine), 2-methylpropan-1-yl (leucine), imidazol-4-ylmethyl (histidine), hydroxymethyl (serine), 1-hydroxyethyl (threonine), carbamoylmethyl (asparagine), 2-carbamoylethyl (glutamine), 4-aminobutan-1-yl (lysine), carboxymethyl (aspartic acid), 3-guanidinopropan-1-yl (arginine), benzyl (phenylalanine), or 4-aminobutan-1-yl (lysine);

R^B is H; and

p 0, 1, or 2.

[0230] In some embodiments, each occurrence of R^A is independently propan-2-yl (valine), 2-methylpropan-1-yl (leucine), carboxymethyl (aspartic acid), benzyl (phenylalanine), or 4-aminobutan-1-yl (lysine);

each R^B is H; and

p is 0, 1, or 2.

[0231] In some embodiments, p is 0.

[0232] In some embodiments, p is 1.

[0233] In some embodiments,

p is 1;

first of R^A is propan-2-yl (valine) and second of R^A is propan-2-yl (valine); and each of R^B is H (dipeptide Val-Val); or

first of R^A is 2-methylpropan-1-yl (leucine), and second of R^A is 2-methylpropan-1-yl (leucine); and each of R^B is H (dipeptide Leu-Leu); or

first of R^A is methyl (alanine) and second of R^A is methyl (alanine); and each of R^B is H (dipeptide Ala-Ala); or


first of R^A is 4-aminobutan-1-yl (lysine); second of R^A is 4-aminobutan-1-yl (lysine); and

each of R^B is H (dipeptide Lys-Lys); or

first of R^A is hydrogen; second of R^A is 4-aminobutan-1-yl, and each of R^B is H (dipeptide Gly-Lys).

[0234] In some embodiments, each of the α -carbon of the amino acid other than glycine is in the L or D configuration.

[0235] In some embodiments, compounds disclosed herein can be synthesized according to the general schemes shown outlined below in **Scheme 1** and **Scheme 2**, where suitable reagents can be purchased from commercial sources or synthesized via known methods or methods adapted from the example procedures provided herein:



[0236] In **Scheme 1**, protection of S1 (K101A shown as example) with trityl chloride (or triphenylmethyl chloride) provides S2 (K101-C20Tr-A shown as example). Hydrolysis of S2 (K101-C20Tr-A shown as example) provides S3 (K101-C20Tr-B, shown as example), which can then be

coupled with compound S4 under esterification conditions using 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC, or EDCI) as the carboxyl activating agent and 4-dimethylaminopyridine (DMAP) as the catalyst to provide S5. Deprotection of S5 followed by further separation and purification provides S6.

[0237] In **Scheme 2**, S7 is prepared by epoxidation of S5 with a peroxycarboxylic acid such as meta-chloroperoxybenzoic acid (m-CPBA). Further separation and purification of S7 provides S8.

[0238] Appropriate starting materials and reagents for use in **Scheme 1** and **Scheme 2** can be purchased or prepared by methods known to one of skill in the art.

[0239] In some embodiments of the methods of **Scheme 1** and **Scheme 2**, the various substituents on the starting compounds (e.g., compounds S1 and S3) are as defined for Formula I. However, it should also be appreciated that chemical derivatization and/or functional group interconversion, can be used to further modify of any of the compounds of **Scheme 1** and **Scheme 2** in order to provide the various compounds of Formula I.

[0240] In some embodiments, synthesis of the prodrugs are prepared by reacting protected amino acids (e.g., N-protected amino acids) with relevant compounds, e.g., compounds having an –OH group at the R⁶ position. Guidance is provided in Examples 63 and 64 illustrating synthesis of amino acid prodrugs as well as knowledge of general procedures available in the art for producing such prodrugs (see, e.g., Vale et al., 2018, *Molecules*. 23(9):2318; Beauchamp et al., 1992, *Antiviral Chemistry & Chemotherapy* 3(3):157-164; incorporated herein by reference).

[0241] Other compounds of the disclosure can be synthesized using the synthetic routes above and adapting chemical synthetic procedures available to the skilled in the art. Exemplary methods of synthesis are provided in the Examples. It is to be understood that each of the procedures describing synthesis of exemplary compounds are part of the specification, and thus incorporated herein into the Detailed Description of this disclosure.

5.3. Pharmaceutically Acceptable Salts

[0242] In some embodiments, the PKC modulating compounds are in free form or where appropriate as pharmaceutically acceptable salt. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al., describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 1977, 66, 1–19, incorporated herein by reference.

[0243] Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. In some embodiments, pharmaceutically acceptable salt of the compounds herein can be prepared during final isolation and purification of the compounds. For example, a pharmaceutically acceptable salt of the compounds herein can be prepared by (1) reacting the compound in free base form with a suitable organic or inorganic acid, and

(2) isolating the salt thus formed. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like.

[0244] Base addition salts can be prepared by (1) reacting the compound, such as the purified compound, in its acid form with a suitable organic or inorganic base, and (2) isolating the salt thus formed. Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_1-C_4\text{alkyl})_4$ salts. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

5.4. Methods of Use

[0245] In another aspect, the compounds described herein are used in a method of treating cancer. In some embodiments, the method of treating cancer comprises administering to a subject in need thereof a therapeutically effective amount any of the compounds described herein.

[0246] In some embodiments, the compounds can be used as monotherapy, or as further provided below, in a combination therapy with one or more therapeutic treatments, particularly in combination with one or more chemotherapeutic agents. In some embodiments, the compounds are used in combination with a second therapeutic agent, where the compounds are used at levels that sensitizes the cancer or cancer cell to the second therapeutic agent, for example at levels of the compound that do not cause significant cell death. In some embodiments, the compounds can be used in combination with radiation therapy, either to sensitize the cells to radiation therapy or as an adjunct to radiation therapy (e.g., at doses sufficient to activate cell death pathway).

[0247] In some embodiments, the cancer for treatment with the compound can be selected from, among others, adrenocortical cancer, anal cancer, biliary cancer, bladder cancer, bone cancer (e.g., osteosarcoma), brain cancer (e.g., gliomas, astrocytoma, neuroblastoma, etc.), breast cancer, cervical

cancer, colon cancer, endometrial cancer, esophageal cancer, head and neck cancer, hematologic cancer (e.g., leukemia and lymphoma), intestinal cancer (small intestine), liver cancer, lung cancer (e.g., bronchial cancer, small cell lung cancer, non-small cell lung cancer, etc.), oral cancer, ovarian cancer, pancreatic cancer, renal cancer, prostate cancer, salivary gland cancer, skin cancer (e.g., basal cell carcinoma, melanoma), stomach cancer, testicular cancer, throat cancer, thyroid cancer, uterine cancer, vaginal cancer, sarcoma, and soft tissue carcinomas.

[0248] In some embodiments, the cancer for treatment with the compound is pancreatic cancer. In some embodiments, the pancreatic cancer for treatment with the compounds is pancreatic adenocarcinoma or metastatic pancreatic cancer. In some embodiments, the cancer for treatment with the compounds is stage I, stage II, stage III, or stage IV pancreatic adenocarcinoma.

[0249] In some embodiments, the cancer for treatment with the compounds is lung cancer. In some embodiments, the lung cancer for treatment with the compounds is small cell lung cancer or non-small cell lung cancer. In some embodiments, the non-small cell lung cancer for treatment with the compounds is an adenocarcinoma, squamous cell carcinoma, or large cell carcinoma. In some embodiments, the lung cancer for treatment with the compounds is metastatic lung cancer.

[0250] In some embodiments, the cancer for treatment with the compounds is a hematologic cancer. In some embodiments, the hematologic cancer is selected from acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), lymphoma (e.g., Hodgkin's lymphoma, Non-Hodgkin's lymphoma, Burkitt's lymphoma), chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), Hairy Cell chronic myelogenous leukemia (CML), and multiple myeloma.

[0251] In some embodiments, the cancer for treatment with the compounds is a leukemia selected from acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), Hairy Cell chronic myelogenous leukemia (CML), and multiple myeloma.

[0252] In some embodiments, the cancer for treatment with the compound is a lymphoma selected from Hodgkin's lymphoma, Non-Hodgkin's lymphoma, and Burkitt's lymphoma).

[0253] In some embodiments, the cancer for treatment with the compound is a cancer characterized by mesenchymal features or mesenchymal phenotype. In some cancers, gain of mesenchymal features is associated with migratory (e.g., intravasation) and invasiveness of cancers. Mesenchymal features can include, among others, enhanced migratory capacity, invasiveness, elevated resistance to apoptosis, and increased production of extracellular matrix (ECM) components. In addition to these physiological characteristics, the mesenchymal features can include expression of certain biomarkers, including among others, E-cadherin, N-cadherin, integrins, FSP-1, α -SMA, vimentin, β -catenin, collagen I, collagen II, collagen III, collagen IV, fibronectin, laminin 5, SNAIL-1, SNAIL-2, Twist-1, Twist-2, and Lef-1. In some embodiments, the cancer selected for treatment with the compounds

herein include, among others, breast cancer, lung cancer, head and neck cancer, prostate cancer, and colon cancer. In some embodiments, the mesenchymal features can be inherent to the cancer type or induced by or selected for by treatment of cancers with chemotherapy and/or radiation therapy.

[0254] In some embodiments, the cancer for treatment with the compound is identified as having or determined to have an activating or oncogenic RAS activity. In some embodiments, the RAS is K-RAS, H-RAS or N-RAS. In some embodiments, the activating or oncogenic RAS is an activating or oncogenic RAS mutation.

[0255] In some embodiments, the cancer for treatment is identified as having or determined to have an activating or oncogenic K-RAS mutation. In some embodiments, the cancer selected for treatment is identified as having or determined to have an activating or oncogenic mutation in human K-RAS at one or more of codon 5, codon 9, codon 12, codon 13, codon 14, codon 18, codon 19, codon 22, codon 23, codon 24, codon 26, codon 33, codon 36, codon 57, codon 59, codon 61, codon 62, codon 63, codon 64, codon 68, codon 74, codon 84, codon 92, codon 95, codon 97, codon 110, codon 115, codon 117, codon 118, codon 119, codon 135, codon 138, codon 140, codon 146, codon 147, codon 153, codon 156, codon 160, codon 164, codon 171, codon 176, codon 185, and codon 188.

[0256] In some embodiments, the activating or oncogenic K-RAS mutation can be a mutation in which: codon 5 is K5E; codon 9 is V9I; codon 12 is G12A, G12C, G12D, G12F, G12R, G12S, G12V, or G12Y; codon 13 is G13C, G13D, or G13V; codon 14 is V14I or V14L; codon 18 is A18D; codon 19 is L19F; codon 22 is Q22K; codon 23 is L23R; codon 24 is I24N; codon 26 is N26K; codon 33 is D33E; codon 36 is I36L or I36M; codon 57 is D57N; codon 59 is A59E, A59G, or A59T; codon 61 is Q61H, Q61K, Q61L, or Q61R; codon 62 is E62G or E62K; codon 63 is E63K; codon 64 is Y64D, Y64H, or Y64N; codon 68 is R68S; codon 74 is T74P; codon 84 is I84T; codon 92 is D92Y; codon 97 is R97I; codon 110 is P110H or P110S; codon 115 is G115E; codon 117 is K117N; codon 118 is C118S; codon 119 is D119N; codon 135 is R135T; codon 138 is G138V; codon 140 is P140H; codon 146 is A146T or A146V; codon 147 is K147N; codon 153 is D153N; codon 156 is F156L; codon 160 is V160A; codon 164 is R164Q; codon 171 is I171M; codon 176 is K176Q; codon 185 is C185R or C185S; and codon 188 is M188V.

[0257] In particular, the cancer for treatment is identified as having or determined to have an oncogenic or activating K-RAS mutations at codon 12, codon 13 and/or codon 61. In some embodiments, the oncogenic or activating K-RAS mutation at codon 12 is G12A, G12C, G12D, G12F, G12R, G12S, G12V, or G12Y; at codon 13 is G13C, G13D, or G13V; and at codon 61 is Q61H, Q61K, Q61L, or Q61R. In some embodiments, the oncogenic or activating K-RAS mutation is a combination of oncogenic or activating K-RAS mutations at codon 12 and codon 13; codon 12 and codon 61; codon 13 and 61; or codon 12, codon 13 and codon 61.

[0258] In some embodiments, the cancer for treatment is identified as having or determined to have an activating or oncogenic N-RAS mutation. In some embodiments, the cancer is identified as having or determined to have an activating or oncogenic mutation in human N-RAS at one or more of codon 12, codon 13 and codon 61. In some embodiments, the activating or oncogenic N-RAS mutation at codon 12 is G12A, G12C, G12D, G12R, G12S, or G12V. In some embodiments, the activating or oncogenic N-RAS mutation at codon 13 is G13A, G13C, G13D, G13R, G13S, or G13V. In some embodiments, the activating or oncogenic N-RAS mutation at codon 61 is Q61E, Q61H, Q61K, Q61L, Q61P, or Q61R. In some embodiments, the oncogenic or activating N-RAS mutation is a combination of activating or oncogenic N-RAS mutations at codon 12 and codon 13; codon 12 and codon 61; codon 13 and 61; or codon 12, codon 13 and codon 61.

[0259] In some embodiments, the cancer for treatment is identified as having or determined to have an activating or oncogenic H-RAS mutation. In some embodiments, the cancer selected for treatment is identified as having an activating or oncogenic mutation in human H-RAS at one or more of codon 12, codon 13 and codon 61. In some embodiments, the activating or oncogenic H-RAS mutation at codon 12 is G12A, G12C, G12D, G12R, G12S, or G12V. In some embodiments, the activating or oncogenic H-RAS mutation at codon 13 is G13A, G13C, G13D, G13R, G13S, or G13V. In some embodiments, the activating or oncogenic H-RAS mutation at codon 61 is Q61E, Q61H, Q61K, Q61L, Q61P, or Q61R. In some embodiments, the oncogenic or activating H-RAS mutation is a combination of activating or oncogenic H-RAS mutations at codon 12 and codon 13; codon 12 and codon 61; codon 13 and 61; or codon 12, codon 13 and codon 61.

[0260] In some embodiments, the cancer for treatment can be a cancer having prevalence (e.g., at least about 10% or more, or about 15% or more of the cancers), of an activating or oncogenic RAS mutation, such as cancer of the biliary tract, cervix, endometrium, pancreas, lung, colon, head and neck, stomach (gastric), biliary tract, endometrium, hematologic (e.g., leukemia, lymphomas, etc.), large intestine, lung, ovary, pancreas, prostate, salivary gland, skin, small intestine, stomach thyroid, aerodigestive tract, urinary tract, and ovary, small intestine, and urinary tract.

[0261] Biological sample for the method herein include any samples are amenable to analysis herein, such as tissue or biopsy samples containing cancer cells, or any biological fluids that contain the material of interests (e.g., DNA), such as blood, plasma, saliva, tissue swabs, and intestinal fluids. In some embodiments, exosomes extruded by cancer cells and obtained from blood or other body fluids can be used to detect nucleic acids and proteins produced by the cancer cells.

[0262] General biological, biochemical, immunological and molecular biological methods applicable to the present disclosure are described in Sambrook et al., *Molecular Cloning: A Laboratory Manual* 2nd Ed. (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; *Current Protocols in Molecular Biology*, Ausubel et al., ed., John Wiley & Sons (2015); *Current Protocols in Immunology*,

Coligan, JE ed., John Wiley & Sons (2015); and *Methods in Enzymology*, Vol. 200, Abelson et al., ed., Academic Press (1991). All publications are incorporated herein by reference.

5.5. Combination Treatments

[0263] In some embodiments, the diterpenoid PKC modulating compounds are used in combination with one or more second therapeutic agents. In some embodiments, the second therapeutic agent is selected from a platinating agent, alkylating agent, antibiotic agent, antimetabolic agent (e.g., folate antagonists, purine analogs, pyrimidine analogs, etc.), topoisomerase inhibiting agent, antimicrotubule agent (e.g., taxanes, vinca alkaloids), hormonal agent (e.g., aromatase inhibitors), plant-derived agent and synthetic derivatives thereof, anti-angiogenic agent, differentiation inducing agent, cell growth arrest inducing agent, apoptosis inducing agent, cytotoxic agent, agent affecting cell bioenergetics, i.e., affecting cellular ATP levels and molecules/activities regulating these levels, anti-cancer biologic agent (e.g., monoclonal antibodies), kinase inhibitors and inhibitors of growth factors and their receptors.

[0264] In some embodiments, the second chemotherapeutic agent is selected from afatinib, afuresertib, alectinib, alisertib, alvocidib, amsacrine, amonafide, amuvatinib, axitinib, azacitidine, azathioprine, bafetinib, barasertib, bendamustine, bleomycin, bosutinib, bortezomib, busulfan, cabozantinib, camptothecin, canertinib, capecitabine, cabazitaxel, carboplatin, carmustine, cenisertib, ceritinib, chlorambucil, cisplatin, cladribine, clofarabine, crenolanib, crizotinib, cyclophosphamide, cytarabine, dabrafenib, dacarbazine, dacomitinib, dactinomycin, danusertib, dasatinib, daunorubicin, decitabine, dinaciclib, docetaxel, dovitinib, doxorubicin, epirubicin, epitinib, eribulin mesylate, erlotinib, etirinotecan, etoposide, everolimus, exemestane, floxuridine, fludarabine, fluorouracil, gefitinib, gemcitabine, hydroxyurea, ibrutinib, icotinib, idarubicin, idelalisib, ifosfamide, imatinib, imetelstat, ipatasertib, irinotecan, ixabepilone, lapatinib, lenalidomide, lestaurtinib, lomustine, lucitanib, masitinib, mechlorethamine, melphalan, mercaptopurine, methotrexate, midostaurin, mitomycin, mitoxantrone, mubritinib, nelarabine, neratinib, nilotinib, nintedanib, omacetaxine mepesuccinate, olaparib, orantinib, oxaliplatin, paclitaxel, palbociclib, palifosfamide tris, pazopanib, pelitinib, pemetrexed, pentostatin, plicamycin, ponatinib, poziotinib, pralatrexate, procarbazine, quizartinib, raltitrexed, regorafenib, ruxolitinib, seliciclib, sorafenib, streptozocin, sulfatinib, sunitinib, tamoxifen, tandutinib, temozolomide, temsirolimus, teniposide, theliatinib, thioguanine, thiotepa, topotecan, uramustine, valrubicin, vandetanib, vemurafenib (Zelborae), vincristine, vinblastine, vinorelbine, vindesine, and the like.

[0265] In some embodiments, the second therapeutic agent is selected from the group consisting of a phosphoinositol-3 kinase (PI3K) inhibitor, AKT inhibitor, mammalian target of rapamycin (mTOR) inhibitor, poly ADP ribose polymerase (PARP) inhibitor, platinum-based anti-cancer compound (PBAC), CBP/ β -catenin inhibitor, Tankyrase (TNKS) inhibitor, probable protein-cysteine N-palmitoyltransferase (PORCN) inhibitor, src kinase/bcr-abl kinase inhibitor, Smoothed (SMO)

inhibitor, anti-cancer nucleoside analog or anti-metabolite, histone deacetylase (HDAC) inhibitor, Bromodomain and Extra-Terminal motif (BET) inhibitor, all-trans-retinoic acid (ATRA), Bruton's tyrosine kinase (BTK) inhibitor, EGFR receptor inhibitor, and combinations thereof.

[0266] In some embodiments, the second therapeutic agent is selected from the group consisting of idelalisib, pictilisib, duvelisib, pilaralisib, alpelisib, copanlisib, voxtalisib, dactolisib, gedatolisib, apitolisib, perifosine, miltefosine, ipatasertib, sirolimus, everolimus, temsirolimus, tacrolimus, ridaforolimus, ridaforolimus, dactolisib, olaparib, veliparib, rucaparib, talazoparib, niraparib, cisplatin, carboplatin, oxaliplatin, dicycloplatin, nedaplatin, lobaplatin, heptaplatin, phenathriplatin, phosphaplatin, LA-12, ICG-001, PRI-724, XAV-939, G007-LK, LGK-974, ETC-159, staurosporine, nilotinib, imatinib, ponatinib, saracatinib, dasatinib, bosutinib, saracatinib, cyclopamine, vismodegib, glasdegib, SANT-1, sonidegib, saridegib, taladegib, GSK1210151A, GSK525762, CPI-0610, RVX-208, vorinostat (SAHA), entinostat, panobinostat, mocetinostat, belinostat, romidepsin, rocilinostat, abexinostat, resminostat, givinostat, quisinostat, pracinostat, kevetrin, CC-292, CNX-774, LFM-A13, CGI1746, trastuzumab, pertuzumab, ado-trastuzumab emtansine, cetuximab, panitumumab, nimotuzuma, mAb806, rindopepimut, lapatinib, erlotinib, gefitinib, afatinib, neratinib, osimertinib, rociletinib, canertinib, and dacomitinib.

5.6. Formulations and Administration

[0267] In some embodiments, the pharmaceutical compositions of the therapeutic agents can be formulated by standard techniques using one or more physiologically acceptable carriers or excipients. Suitable pharmaceutical carriers are described herein and in Remington: The Science and Practice of Pharmacy, 21st Ed. (2005). The therapeutic compounds and their physiologically acceptable salts, hydrates and solvates can be formulated for administration by any suitable route, including, among others, topically, nasally, orally, parenterally, rectally or by inhalation. In some embodiments, the compounds and pharmaceutical compositions thereof are administered by intradermal, subdermal, intravenous, intramuscular, intranasal, intracerebral, intratracheal, intraarterial, intraperitoneal, intravesical, intrapleural, intracoronary or intratumoral injection, such as with a syringe or other devices. Transdermal administration is also contemplated, as are inhalation or aerosol administration. Tablets, capsules, and solutions can be administered orally, rectally or vaginally.

[0268] For oral administration, a pharmaceutical composition can take the form of, for example, a tablet or a capsule prepared by conventional means with a pharmaceutically acceptable excipient. Tablets and capsules comprising the active ingredient can be prepared together with excipients such as: (a) diluents or fillers, e.g., lactose, dextrose, sucrose, mannitol, sorbitol, cellulose (e.g., ethyl cellulose, microcrystalline cellulose), glycine, pectin, polyacrylates and/or calcium hydrogen phosphate, calcium sulfate; (b) lubricants, e.g., silica, talcum, stearic acid, its magnesium or calcium salt, metallic stearates, colloidal silicon dioxide, hydrogenated vegetable oil, corn starch, sodium

benzoate, sodium acetate and/or polyethyleneglycol; (c) binders, e.g., magnesium aluminum silicate, starch paste, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, polyvinylpyrrolidone and/or hydroxypropyl methylcellulose; (d) disintegrants, e.g., starches (including potato starch or sodium starch), glycolate, agar, alginic acid or its sodium salt, or effervescent mixtures; (e) wetting agents, e.g., sodium lauryl sulphate, and/or (f) absorbents, colorants, flavors and sweeteners. The compositions are prepared according to conventional mixing, granulating or coating methods.

[0269] Tablets may be either film coated or enteric coated according to methods known in the art. Liquid preparations for oral administration can take the form of, for example, solutions, syrups, or suspensions, or they can be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations can be prepared by conventional means with pharmaceutically acceptable carriers and additives, for example, suspending agents, e.g., sorbitol syrup, cellulose derivatives, or hydrogenated edible fats; emulsifying agents, for example, lecithin or acacia; non-aqueous vehicles, for example, almond oil, oily esters, ethyl alcohol, or fractionated vegetable oils; and preservatives, for example, methyl or propyl-p-hydroxybenzoates or sorbic acid. The preparations can also contain buffer salts, flavoring, coloring, and/or sweetening agents as appropriate. If desired, preparations for oral administration can be suitably formulated to give controlled release of the active compound.

[0270] The therapeutic agents can be formulated for parenteral administration, for example by bolus injection or continuous infusion. Formulations for injection can be presented in unit dosage form, for example, in ampoules or in multi-dose containers, with an optionally added preservative. Injectable compositions can be aqueous isotonic solutions or suspensions. In some embodiments for parenteral administration, the therapeutic agents can be prepared with a surfactant, such as Cremaphor, or lipophilic solvents, such as triglycerides or liposomes. The compositions may be sterilized and/or contain adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure and/or buffers. Alternatively, the therapeutic agent can be in powder form for reconstitution with a suitable vehicle, for example, sterile pyrogen-free water, before use. In addition, they may also contain other therapeutically effective substances.

[0271] In some embodiments, the therapeutic agent, e.g., the diterpenoid PKC modulating compounds, are administered intratumorally. In some embodiments, the therapeutic agent is administered directly into the tumor, allowing for high local concentration of the therapeutic agent and in some embodiments, increased bioavailability of the therapeutic agent at the site of the tumor. Any formulation of the therapeutic agent suitable for intratumoral administration can be used in the embodiments herein. Intratumoral administration can be by injection of the therapeutic agent into the tumor (see, e.g., Celikoglu et al., 2008, *Cancer Therapy*, 6:545-552) or by intravenous administration to blood vessels feeding to the tumor. In some embodiments, the injection device has a porous

delivery channel (e.g., needle) for wider distribution or infusion of the therapeutic agent for treating tumors with large volume.

[0272] For administration by inhalation, the therapeutic agent may be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, for example, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide, or other suitable gas. In the case of a pressurized aerosol, the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, for example, gelatin for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base, for example, lactose or starch.

[0273] Suitable formulations for transdermal application include an effective amount of a therapeutic agent with a carrier. Preferred carriers include absorbable pharmacologically acceptable solvents to assist passage through the skin of the subject. For example, transdermal devices are in the form of a bandage or patch comprising a backing member, a reservoir containing the therapeutic agent optionally with carriers, optionally a rate controlling barrier to deliver the compound to the skin of the host at a controlled and predetermined rate over a prolonged period of time, and a means to secure the device to the skin. Matrix transdermal formulations may also be used.

[0274] Suitable formulations for topical application, e.g., to the skin and eyes, are preferably aqueous solutions, ointments, creams or gels well-known in the art. The formulations may contain solubilizers, stabilizers, tonicity enhancing agents, buffers and preservatives.

[0275] In some embodiments, the therapeutic agent can also be formulated as a rectal composition, for example, suppositories or retention enemas, for example, containing conventional suppository bases, for example, cocoa butter or other glycerides, or gel forming agents, such as carbomers.

[0276] In some embodiments, the therapeutic agent can be formulated as a depot preparation. Such long-acting formulations can be administered by implantation (for example, subcutaneously or intramuscularly) or by intramuscular injection. The therapeutic agent can be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil), ion exchange resins, biodegradable polymers, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[0277] In some embodiments, the carrier is a cyclodextrins, such as to enhance solubility and/or bioavailability of the compounds herein. In some embodiments, the cyclodextrin for use in the pharmaceutical compositions can be selected from α -cyclodextrin, β -cyclodextrin, γ -cyclodextrin, derivatives thereof, and combinations thereof. In particular, the cyclodextrin is selected from β -cyclodextrin, γ -cyclodextrin, derivatives thereof, and combinations thereof.

[0278] In some embodiments, the compounds can be formulated with a cyclodextrin or derivative thereof selected from carboxyalkyl cyclodextrin, hydroxyalkyl cyclodextrin, sulfoalkylether cyclodextrin, and an alkyl cyclodextrin. In various embodiments, the alkyl group in the cyclodextrin is methyl, ethyl, propyl, butyl, or pentyl.

[0279] In some embodiments, the cyclodextrin is α -cyclodextrin or a derivative thereof. In some embodiments, the α -cyclodextrin or derivative thereof is selected from carboxyalkyl- α -cyclodextrin, hydroxyalkyl- α -cyclodextrin, sulfoalkylether- α -cyclodextrin, alkyl- α -cyclodextrin, and combinations thereof. In some embodiments, the alkyl group in the α -cyclodextrin derivative is methyl, ethyl, propyl, butyl, or pentyl.

[0280] In some embodiments, the cyclodextrin is β -cyclodextrin or a derivative thereof. In some embodiments, the β -cyclodextrin or derivative thereof is selected from carboxyalkyl- β -cyclodextrin, hydroxyalkyl- β -cyclodextrin, sulfoalkylether- β -cyclodextrin, alkyl- β -cyclodextrin, and combinations thereof. In some embodiments, the alkyl group in the β -cyclodextrin derivative is methyl, ethyl, propyl, butyl, or pentyl.

[0281] In some embodiments, the β -cyclodextrin or a derivative thereof is hydroxyalkyl- β -cyclodextrin or sulfoalkylether- β -cyclodextrin. In some embodiments, the hydroxyalkyl- β -cyclodextrin is hydroxypropyl- β -cyclodextrin. In some embodiments, the sulfoalkylether- β -cyclodextrin is sulfobutylether- β -cyclodextrin. In some embodiments, β -cyclodextrin or a derivative thereof is alkyl- β -cyclodextrin, in particular methyl- β -cyclodextrin. In some embodiments using methyl- β -cyclodextrin, the β -cyclodextrin is randomly methylated β -cyclodextrin.

[0282] In some embodiments, the cyclodextrin is γ -cyclodextrin or a derivative thereof. In some embodiments, the γ -cyclodextrin or derivative thereof is selected from carboxyalkyl- γ -cyclodextrin, hydroxyalkyl- γ -cyclodextrin, sulfoalkylether- γ -cyclodextrin, and alkyl- γ -cyclodextrin. In some embodiments, the alkyl group in the γ -cyclodextrin derivative is methyl, ethyl, propyl, butyl, or pentyl. In some embodiments, the γ -cyclodextrin or derivative thereof is hydroxyalkyl- γ -cyclodextrin or sulfoalkylether- γ -cyclodextrin. In some embodiments, the hydroxyalkyl- γ -cyclodextrin is hydroxypropyl- γ -cyclodextrin.

[0283] When used in a formulation with the compound of the present disclosure, the cyclodextrin can be present at about 0.1 w/v to about 30% w/v, about 0.1 w/v to about 20% w/v, about 0.5% w/v to about 10% w/v, or about 1% w/v to about 5% w/v. In some embodiments, the cyclodextrin is present at about 0.1% w/v, about 0.2% w/v, about 0.5% w/v, about 1% w/v, about 2% w/v, about 3% w/v, about 4% w/v, about 5% w/v, about 6% w/v, about 7% w/v, about 8% w/v, about 9% w/v, about 10% w/v, about 12% w/v, about 14% w/v, about 16% w/v, about 18% w/v, about 20% w/v, about 25% w/v, or about 30% w/v or more.

[0284] The pharmaceutical compositions can, if desired, be presented in a pack or dispenser device that can contain one or more unit dosage forms containing the active ingredient. The pack can, for example, comprise metal or plastic foil, for example, a blister pack. The pack or dispenser device can be accompanied by instructions for administration.

5.7. Effective Amount and Dosing

[0285] In some embodiments, a pharmaceutical composition of the therapeutic agent is administered to a subject, preferably a human, at a therapeutically effective dose to prevent, treat, or control a condition or disease as described herein. The pharmaceutical composition is administered to a subject in an amount sufficient to elicit an effective therapeutic response in the subject. An effective therapeutic response is a response that at least partially arrests or slows the symptoms or complications of the condition or disease. An amount adequate to accomplish this is defined as “therapeutically effective dose” or “therapeutically effective amount.”

[0286] The dosage of therapeutic agents can take into consideration, among others, the species of warm-blooded animal (mammal), the body weight, age, condition being treated, the severity of the condition being treated, the form of administration, route of administration. The size of the dose also will be determined by the existence, nature, and extent of any adverse effects that accompany the administration of a particular therapeutic compound in a particular subject.

[0287] In some embodiments, the diterpenoid PKC activating compound, the compound can be administered in a dose in the range from about 0.001 mg per kg of subject weight (0.001 mg/kg) to about 1000 mg/kg. In some embodiments, the dose is in the range of about 0.001 mg/kg to about 500 mg/kg. In some embodiments, the dose is in the range of about 1 mg/kg to about 500 mg/kg. In some embodiments, the dose is about 2 mg/kg to about 250 mg/kg. In another embodiment, the dose is about 5 mg/kg to about 100 mg/kg. In another embodiment, the dose is about 5 mg/kg to about 100 mg/kg. In some embodiments, the dose is about 0.001 mg/kg, 0.01 mg/kg, 0.05 mg/kg, 0.1 mg/kg, 0.2 mg/kg, 0.3 mg/kg, 0.4 mg/kg, 0.5 mg/kg, 1 mg/kg, 2 mg/kg, 3 mg/kg, 4 mg/kg, 5 mg/kg, 10 mg/kg, 20 mg/kg, 40 mg/kg, 50 mg/kg, 100 mg/kg, 200 mg/kg or 500 mg/kg. In some embodiments, the dose can be administered once per day or divided into subdoses and administered in multiple doses, e.g., twice, three times, or four times per day.

[0288] In some embodiments, the diterpenoid PKC activator can be administered with one or more of the second therapeutic agent sequentially or concurrently, either by the same route or by different routes of administration. When administered sequentially, the time between administrations is selected to benefit, among others, the therapeutic efficacy and/or safety of the combination treatment. In some embodiments, the diterpenoid PKC activator can be administered first followed by a second therapeutic agent, or alternatively, the second therapeutic agent administered first followed by the diterpenoid PKC activator. By way of example and not limitation, the time between administrations

is about 1 hr, about 2 hr, about 4hr, about 6 hr, about 12 hr, about 16 hr or about 20 hr. In some embodiments, the time between administrations is about 1, about 2, about 3, about 4, about 5, about 6, or about 7 more days. In some embodiments, the time between administrations is about 1 week, 2 weeks, 3 weeks, or 4 weeks or more. In some embodiments, the time between administrations is about 1 month or 2 months or more.

[0289] When administered concurrently, the diterpenoid PKC modulator can be administered separately at the same time as the second therapeutic agent, by the same or different routes, or administered in a single composition by the same route.

[0290] In some embodiments, the amount and frequency of administration of the second therapeutic agent can use standard dosages and standard administration frequencies used for the particular therapeutic agent. See, e.g., Physicians' Desk Reference, 70th Ed., PDR Network, 2015; incorporated herein by reference.

[0291] In some embodiments, where administration of the therapeutic agent is to a localized site, for example, intratumoral injection, the dosages can be the dosages used for systemic administration, such as dosages used for intravenous, intramuscular, and intraperitoneal administration. In some embodiments, the dose for localized administration, e.g., intratumoral administration, is higher than those used for systemic administration. In some embodiments, the administered dose is sufficient for the intended effect, for example killing or necrotization of tumor tissue. In some embodiments, intratumoral administration is done once, twice, three times, four times, five times or up to six times or more, where each administration is separated in time, for example, until the desired outcome is achieved.

[0292] It to be understood that optimum dosages, toxicity, and therapeutic efficacy of such therapeutic agents may vary depending on the relative potency of individual therapeutic agent and can be determined by pharmaceutical procedures in cell cultures or experimental animals, for example, by determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and can be expressed as the ratio, LD₅₀/ED₅₀. Therapeutic agents or combinations thereof that exhibit large therapeutic indices are preferred. While certain agents that exhibit toxic side effects can be used, care should be used to design a delivery system that targets such agents to the site of affected tissue to minimize potential damage to normal cells and, thereby, reduce side effects.

[0293] The data obtained from, for example, cell culture assays and animal studies can be used to formulate a dosage range for use in humans. The dosage of such small molecule compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed and the route of administration. For any compounds used in the methods of the invention, the therapeutically effective


dose can be estimated initially from cell culture assays. A dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC_{50} (the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma can be measured, for example, by high performance liquid chromatography (HPLC).

[0294] The following examples are provided to further illustrate the methods of the present disclosure, and the compounds and compositions for use in the methods. The examples described are illustrative only and are not intended to limit the scope of the invention in any way.

6. EXAMPLES

Example 1: Synthesis Scheme of **K101-Epoxyde** and **K101-DI-OH**.

[0295] The scheme for synthesis of compounds **K101-Epoxyde** and **K101-DI-OH** are illustrated below.



[0296] **Preparation of Compound K101-C20Tr-A.** To a solution of **K101A** (1 g, 2.56 mmol, 1 eq) in pyridine (40 mL) was added Trityl chloride (2.14 g, 7.68 mmol, 3.00 eq). The mixture was stirred at 40°C for 14 hours (hr) to give a yellow solution. LC-MS showed desired mass was found, and **K101A** was remained. The mixture was stirred at 40°C for 12hr again. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture was concentrated by drumming N_2 to give the crude product. The crude product was purified by flash column (eluting with: petroleum ether (PE)/ethyl acetate=100%PE to 20%) to give **K101-C20Tr-A** (1.6 g, 2.53 mmol, 98.73% yield) as a white solid.

^1H NMR (400MHz, CDCl_3) δ 7.59 (s, 1H), 7.44-7.42 (m, 6H), 7.31-7.29 (m, 6H), 7.24-7.21 (m, 3H), 5.63 (brs, 1H), 3.51 (s, 2H), 3.28 (s, 1H), 2.93 (s, 1H), 2.49-2.41 (m, 2H), 2.09-2.06 (m, 5H), 2.09-2.03 (m, 7H), 1.99-1.94 (m, 1H), 1.78 (s, 3H), 1.20 (s, 3H), 1.07 (s, 3H), 0.89-0.81 (m, 4H).

[0297] Preparation of Compound Epoxide-Trt. To a solution of **K101-C20Tr-A** (30.00 mg, 47.41 μmol , 1.00 equivalent (hereafter as eq)) in dichloromethane (hereafter as DCM) (2.00 mL) was added NaHCO_3 (11.95 mg, 142.23 μmol , 5.53 μL , 3.00 eq), meta-chloroperoxybenzoic acid (m-CPBA) (14.44 mg, 71.11 μmol , 1.5eq, 85% purity), and the reaction mixture was stirred at 20°C for 2 hours (h) to give a suspension. LC-MS showed the reaction was complete, and the desired MS value was observed. Thin-layer chromatography (TLC) (petroleum ether/ethyl acetate mixture ration 2:1 (PE/EtOAc=2/1), SiO_2) analysis showed no new spots. The reaction mixture was mixed with DCM (5 mL) and brine (2 mL), and the organic layer was separated and concentrated under reduced pressure to give 35.5 mg of crude product as a colorless gum. The product was purified by preparative (prep)-TLC (PE/EtOAc=2/1, SiO_2) to give **Epoxide-Trt** (20.10 mg, 65.34% yield) as a colorless gum.

[0298] ^1H NMR (400MHz, CDCl_3) δ 7.63 (s, 1H), 7.37-7.30 (m, 6H), 7.30-7.18 (m, 11H), 5.54 (brs, 1H), 3.96-3.89 (m, 1H), 3.17 (d, $J=9.3$ Hz, 1H), 3.08 (d, $J=8.5$ Hz, 1H), 2.85-2.74 (m, 2H), 2.09 (s, 3H), 2.07-2.03 (m, 1H), 2.02 (s, 1H), 1.98 (s, 1H), 1.96-1.87 (m, 1H), 1.86-1.81 (m, 1H), 1.80-1.74 (m, 3H), 1.66-1.59 (m, 1H), 1.21 (s, 3H), 1.04 (s, 3H), 0.97 (d, $J=4.3$ Hz, 1H), 0.89 (d, $J=6.5$ Hz, 3H).

[0299] Preparation of Compound K101-Epoxide and K101-DI-OH. To a solution of **Epoxide-Trt** (10.00 mg, 15.41 μmol , 1.00 eq) in DCM (500.00 μL) was added trifluoroacetic acid (hereafter as TFA) (100.00 μL), and the reaction solution stirred at 0°C for 1h. TLC (PE/EtOAc=2/1, SiO_2) showed the reaction was complete. The reaction was quenched by saturated aqueous NaHCO_3 (2 mL) at 0°C, then extracted with DCM (5 mL x 2), dried over Na_2SO_4 , filtered, and concentrated under reduced pressure to give a colorless gum. The residue was combined with a second preparation of crude product and purified by prep-HPLC (column: Waters Xbridge 150 x 25 x 5 mm; mobile phase: A [A: water (0.05% ammonia hydroxide v/v)]; B [acetonitrile (ACN)]; gradient B%: 25%-55% in 10min to give **K101-Epoxide** (1.51 mg, 3.71 μmol , 24.11% yield, 100% purity) and **K101-DI-OH** (1.20 mg, 2.60 μmol , 16.90% yield, 92.1% purity), both as white solid after lyophilization.

[0300] K101-Epoxide: LC-MS (m/z): 429.2 $[\text{M}+\text{Na}]^+$


[0301] K101-Epoxide: ^1H NMR (400MHz, CD_3OD) δ 7.53 (s, 1H), 3.48 (d, $J=11.9$ Hz, 1H), 3.44-3.39 (m, 1H), 3.36 (s, 1H), 3.16 (d, $J=8.6$ Hz, 1H), 2.66 (d, $J=16.8$ Hz, 1H), 2.14-2.06 (m, 4H), 2.04-1.89 (m, 3H), 1.77-1.74 (m, 3H), 1.59 (dd, $J=11.2, 14.3$ Hz, 1H), 1.22 (s, 3H), 1.06 (s, 3H), 1.02 (d, $J=4.9$ Hz, 1H), 0.89 (d, $J=6.6$ Hz, 3H).

[0302] K101-DI-OH: LC-MS (m/z): 447.1 $[\text{M}+\text{Na}]^+$

[0303] **K101-DI-OH**: $^1\text{H NMR}$ (400MHz, CD_3OD) δ 7.70 (s, 1H), 3.89 (s, 1H), 3.72-3.68 (m, 1H), 3.53 (d, $J=2.6$ Hz, 2H), 2.45 (d, $J=7.9$ Hz, 1H), 2.21-2.07 (m, 2H), 2.03 (s, 3H), 1.85-1.79 (m, 1H), 1.79-1.74 (m, 3H), 1.60-1.48 (m, 2H), 1.14 (s, 3H), 1.08 (s, 4H), 0.92 (d, $J=6.8$ Hz, 3H).

Example 2: Synthesis Scheme of **K101-C13OH**.

[0304] The scheme for synthesis of compound **K101-C13OH** is illustrated below.




[0305] Preparation of Compound **K101-C13OH**. To a solution of **K101A** (40.00 mg, 102.44 μmol , 1.00 eq) in MeOH (20.00 mL) was added $\text{Ba(OH)}_2 \cdot 8\text{H}_2\text{O}$ (322.69 mg, 1.02 mmol, 10.00 eq). The mixture was stirred at 20°C for 4 hours to give a yellow suspension. LC-MS and TLC (eluting with: EtOAc=100%) showed the reaction was complete. The reaction mixture was quenched with saturated NH_4Cl (10 mL) and extracted with dichloromethane (DCM) (100 mL x 3). The organic layers were dried over Na_2SO_4 and concentrated to give the crude product. The crude product was purified by prep-TLC (eluting with: EtOAc=2/1) to give **K101-C13OH** (9.30 mg, 26.69 μmol , 26.05% yield, 100% purity) as a white solid.

[0306] LC-MS (m/z): 371.2 $[\text{M}+\text{Na}]^+$

[0307] $^1\text{H NMR}$ (400MHz, CD_3OD) δ 7.29 (s, 1H), 5.18 (s, 1H), 4.58 (s, 2H), 3.81-3.75 (m, 1H), 3.50-3.46 (m, 1H), 3.20-3.11 (m, 3H), 2.16-2.11 (m, 1H), 1.76-1.63 (m, 5H), 1.27 (m, 1H), 1.17 (m, 3H), 1.74-1.53 (m, 8H), 1.17 (s, 3H), 1.05 (s, 3H), 1.06-0.88 (m, 6H).

Example 3: Synthesis Scheme of **K101-C1301**.

[0308] The scheme for synthesis of compound **K101-C1301** is illustrated below.



[0309] Preparation of Compound K101-C20Tr-A. To a solution of **K101A** (500.00 mg, 1.28 mmol, 1.00 eq) in pyridine (10.00 mL) was added trityl chloride (**TrtCl**) (1.07 g, 3.84 mmol, 3.00 eq). The mixture was stirred at 20°C for 14 hours to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture was concentrated by N_2 to give the crude product. The product was purified by flash column (eluting with: EtOAc in PE 1% to 50%) to give **K101-C20Tr-A** (790.00 mg, 1.02 mmol, 79.78% yield, 81.795% purity) as a white solid.

[0310] Preparation of Compound K101-C20Tr-B. To a solution of **K101-C20Tr-A** (290.00 mg, 458.30 μmol , 1.00 eq) in MeOH (76.00 mL) was added $\text{Ba(OH)}_2 \cdot 8\text{H}_2\text{O}$ (1.44 g, 4.58 mmol, 10.00 eq) at 0°C. The mixture was stirred at 20°C for 3 hours to give yellow suspension. LC-MS showed the reaction was complete. The reaction mixture was quenched with saturated aqueous NH_4Cl (50 mL) and extracted with dichloromethane (**DCM**) (300 mL x 3). The organic layers were dried over Na_2SO_4 . The organic layers was filtered on silica gel and washed with EtOAc (50 mL). The organic layer was concentrated to give **K101-C20Tr-B** (260.00 mg, 425.57 μmol , 92.86% yield, 96.694% purity) as a white solid.

[0311] Preparation of Compound K101-C1301-A. To solution of **K101-C1301-B** (35.00 mg, 59.25 μmol , 1.00 eq) in DCM (500.00 μL) were added 3-cyclopentylpropanoic acid (10.11 mg, 71.10 μmol , 10.11 μL , 1.20 eq), N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride EDC (**EDC**) (22.72 mg, 118.49 μmol , 2.00 eq) and 4-Dimethylaminopyridine (**DMAP**) (14.48 mg, 118.50 μmol , 2.00 eq). The mixture was stirred at 20°C for 14 hours to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=4/1) showed the reaction was complete. The reaction mixture was combined with a second preparation of the compound, quenched with H_2O (10 mL) and extracted

with DCM (15 mL x 3). The combined organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=4/1) to give **K101-C1301-A** (32.00 mg, 44.76 μmol, 65.12% yield, 100% purity) as a white solid.


[0312] Preparation of Compound K101-C1301. To a solution of **K101-C1301-A** (32.00 mg, 44.76 μmol, 1.00 eq) in DCM (1.00 mL) was added TFA (385.00 mg, 3.38 mmol, 250.00 uL, 75.44 eq). The mixture was stirred at 20°C for 1 hour to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=1/1) showed the reaction was complete. The solvent was removed by N₂ to give a crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=1/1) to give **K101-C1301** (8.10 mg, 17.14 μmol, 38.29% yield, 100% purity) as a white solid.

[0313] LC-MS (m/z): 495.3 [M+Na]⁺

[0314] ¹H NMR (400MHz, CD₃OD) δ 7.53 (s, 1H), 5.59 (s, 1H), 4.54-4.49 (m, 2H), 3.95-3.88 (m, 2H), 3.14 (s, 1H), 3.04 (s, 1H), 2.53-2.43 (m, 2H), 2.35-2.31 (m, 2H), 2.10-1.90 (m, 2H), 1.77-1.72 (m, 6H), 1.62-1.52 (m, 7H), 1.15 (s, 3H), 1.05 (s, 3H), 0.89-0.83 (m, 4H).

Example 4: Synthesis Scheme of **K101-C1302**.

[0315] The scheme for synthesis of compound **K101-C1302** is illustrated below.



[0316] Preparation of Compound K101-C1302-A. To a solution of **K101-C20Tr-B** (40.00 mg, 67.71 μmol, 1.00 eq) in DCM (1.00 mL) were added 3-[4-(trifluoromethyl)phenyl]propanoic acid (**C13-02**) (17.73 mg, 81.25 μmol, 1.20 eq), DMAP (16.54 mg, 135.42 μmol, 2.00 eq) and EDC (25.96 mg, 135.42 μmol, 2.00 eq). The mixture was stirred at 20°C for 2 hours to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The mixture was combined with a second preparation of the compound, quenched with saturated NaHCO₃ (5 mL) and extracted with DCM (10 mL x 3). The organic layers were washed with H₂O (5 mL), dried over Na₂SO₄ and then concentrated to give the crude product. The product was purified by prep-TLC


(eluting with: PE/EtOAc=2/1) to give **K101-C1302-A** (34.00 mg, 42.45 μ mol, 53.72% yield, 98.736% purity) as a white solid.

[0317] Preparation of compound **K101-C1302**. To a solution of **K101-C1302-A** (28.00 mg, 35.40 μ mol, 1.00 eq) in MeOH (1.00 mL) was added HClO₄ (464.95 mg, 4.63 mmol, 280.09 μ L, 130.73 eq). The mixture was stirred at 0°C for 0.5 hour to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was combined with a second preparation of the compound and purified by prep-HPLC (column: Waters XSELECT C18 150 x 30mm x 5 μ m; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 33%-63%, 10 min) to give **K101-C1302** (10.60 mg, 18.56 μ mol, 52.42% yield, 96.032% purity) as a white solid.

[0318] LC-MS (m/z): 571.3 [M+Na]⁺

[0319] ¹H NMR (400MHz, CD₃OD) δ 7.58-7.52 (m, 2H), 7.43-7.41 (m, 2H), 5.55 (s, 1H), 3.95-3.87 (m, 2H), 3.29-2.99 (m, 4H), 2.72-2.68 (m, 2H), 2.47-2.37 (m, 2H), 2.05-1.96 (m, 2H), 1.72 (s, 3H), 1.43-1.40 (m, 1H), 1.01 (s, 6H), 0.85-0.83 (m, 3H), 0.75-0.73 (m, 3H).

Example 5: Synthesis Scheme of K101-C1303



[00253] Preparation of Compound **K101-C1303-A**: To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μ mol, 1.00 eq) in DCM (2.00 mL) were added (2S)-2-(tert-butoxycarbonylamino)-3-phenylpropanoic acid (C13-03) (26.95 mg, 101.57 μ mol, 2.00 eq), DMAP (24.82 mg, 203.13 μ mol, 4.00 eq) and EDC (19.47 mg, 101.57 μ mol, 2.00 eq). The mixture was stirred at 20°C for 48 hours to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture was quenched with saturated NaHCO₃ (5 mL) and extracted with DCM (10 mL*3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The crude product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1303-A** (30.00 mg, 34.85 μ mol, 68.63% yield, 97.349% purity) as a white solid.

[00254] ¹H NMR (400MHz, CDCl₃) δ 7.50 (s, 1H), 7.37-7.35 (m, 5H), 7.24-7.22 (m, 8H), 7.17-7.12 (m, 7H), 5.54 (s, 1H), 5.04 (m, 1H), 4.87-4.85 (m, 1H), 4.49 (m, 1H), 3.18 (s, 1H), 3.02 (m, 1H), 2.83

(m, 1H), 2.46-2.41 (m, 1H), 2.32 - 2.28 (m, 1H), 1.98-1.93 (m, 2H), 1.70 (m, 3H), 1.35 (s, 9H), 1.19 (s, 3H), 0.99-0.98 (m, 3H), 0.78-0.77 (m, 3H), 0.70-0.69 (m, 1H).


[00255] Preparation of Compound K101-C1303: To a solution of **K101-C20Tr-B** (30.00 mg, 35.80 μ mol, 1.00 eq) in DCM (1.00 mL) was added TFA (154.00 mg, 1.35 mmol, 100.00 μ L, 37.73 eq) at 0°C. The mixture was stirred at 0°C for 2hr to give a red-brown solution. The mixture was stirred at 20°C for 14 hours to give a red-brown solution again. The reaction mixture was quenched with H₂O (5 mL) and the organic layer was separated. The water layer was lyophilized. The organic layer was dissolved in MeOH (3.00 mL) and added HClO₄ (83.00 mg, 826.20 μ mol, 50.00 μ L, 23.08 eq) at 0°C. The mixture was stirred at 0°C for 2 hours to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was purified by prep-HPLC (column: Phenomenex Gemini 150*25mm*10um; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 18%-48%, 10 min) to give **K101-C1303** (4.50 mg, 6.90 μ mol, 19.27% yield, 93.438% purity, TFA salt) as a white solid.

[00256] LC-MS (m/z): 519.3 [M+H]⁺

[00257] ¹H NMR (400MHz, CD₃OD) δ 7.45 (s, 1H), 7.31-7.21 (m, 5H), 5.51 (s, 1H), 4.27-4.21 (m, 1H), 3.84 (s, 2H), 3.25 (m, 1H), 3.06-3.00 (m, 3H), 2.40-2.33 (m, 1H), 2.33-2.28 (m, 1H), 2.08 (m, 1H), 2.04 (m, 1H), 1.65 (s, 3H), 1.44-1.40 (m, 1H), 1.03 (s, 3H), 0.97 (s, 3H), 0.87-0.81 (m, 4H).

Example 6: Synthesis Scheme of **K101-C1304**.

[0258] The scheme for synthesis of compound **K101-C1304** is illustrated below.



[0259] Preparation of Compound K101-C1304-A. To a solution of **K101-C20Tr-B** (40.00 mg, 67.71 μ mol, 1.00 eq) in DCM (2.00 mL) were added 2-phenylacetic acid (**C13-04**) (11.06 mg, 81.25 μ mol, 10.24 μ L, 1.20 eq), DMAP (33.09 mg, 270.84 μ mol, 4.00 eq) and EDC (25.96 mg, 135.42 μ mol, 2.00 eq). The mixture was stirred at 20°C for 12 hours to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture was combined with a second preparation of the compound, quenched with H₂O (5 mL) and extracted with DCM (15 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1304-A** (40.00 mg, 56.43 μ mol, 65.78% yield) as a white solid.


[0260] Preparation of Compound K101-C1304. To a solution of **K101-C1304-A** (40.00 mg, 56.43 μmol , 1.00 eq) in MeOH (3.00 mL) was added HClO_4 (83.00 mg, 826.14 μmol , 50.00 μL , 14.64 eq) at 0°C . The mixture was stirred at 0°C for 0.5 hr to give a yellow solution. LC-MS showed the reaction was complete. The mixture was purified by prep-HPLC (column: Waters XSELECT C18 150 x 30mm x 5um; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 33%-63%, 10 min) to give **K101-C1304** (16.30 mg, 34.94 μmol , 61.91% yield) as a white solid.

[0261] LC-MS (m/z): 489.3 $[\text{M}+\text{Na}]^+$

[0262] ^1H NMR (400MHz, CD_3OD) δ 7.53 (s, 1H), 7.33-7.28 (m, 5H), 5.57 (s, 1H), 3.95-3.88 (m, 2H), 3.64 (s, 2H), 3.14-3.03 (m, 2H), 2.48-2.39 (m, 2H), 2.12-2.02 (m, 2H), 1.73 (s, 3H), 1.56-1.50 (m, 1H), 1.03-1.01 (m, 6H), 0.88-0.78 (m, 4H).

Example 7: Synthesis Scheme of **K101-C1305**.

[0263] The scheme for synthesis of compound **K101-C1305** is illustrated below.



[0264] Preparation of Compound K101-C1305-A. To a solution of **K101-C20Tr-B** (45.00 mg, 76.17 μmol , 1.00 eq) in DCM (2.00 mL) were added 2-tert-butoxycarbonyl-2-azaspiro [3.3] heptane-6-carboxylic acid (C13-05) (27.57 mg, 114.26 μmol , 1.50 eq), DMAP (37.22 mg, 304.68 μmol , 4.00 eq), N,N-diisopropylethylamine (DIEA) (19.69 mg, 152.34 μmol , 26.61 μL , 2.00 eq) and EDC (29.21 mg, 152.34 μmol , 2.00 eq). The mixture was stirred at 20°C for 48 hours to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture was combined with a second preparation of the compound, quenched with saturated NaHCO_3 (5 mL) and extracted with DCM (10 mL x 3). The organic layers were dried over Na_2SO_4 and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1305-A** (70.00 mg, crude) as a white solid.

[0265] Preparation of Compound K101-C1305. To a solution of **K101-C1305-A** (70.00 mg, 85.99 μmol , 1.00 eq) in DCM (1.00 mL) was added TFA (154.00 mg, 1.35 mmol, 100.00 μL , 15.71 eq) at 0°C . The mixture was stirred at 0°C for 2 hr give a red-brown solution. The mixture was stirred at


20°C for 16 hours (hr). LC-MS showed that the desired product was not present. The reaction mixture was quenched with H₂O (5 mL) and the organic layer was separated. LC-MS The organic layer was dissolved in MeOH (3.00 mL) followed by addition of HClO₄ (83.00 mg, 826.20 μmol, 50.00 uL, 9.61 eq) at 0°C. The mixture was stirred at 0°C for 2hr to give a yellow solution. LC-MS showed the reaction was complete. The mixture was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10um; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 10%-40%, 10min) to give **K101-C1305** (11.80 mg, 20.15 μmol, 23.43% yield, TFA) as a yellow solid.

[0266] LC-MS (m/z): 494.2 [M+Na]⁺

[0267] ¹H NMR (400MHz, CD₃OD) δ 7.56 (s, 1H), 5.63 (s, 1H), 4.11-4.09 (m, 4H), 4.00-3.96 (m, 2H), 3.19-3.07 (m, 3H), 2.60-2.46 (m, 6H), 2.16-2.12 (m, 2H), 1.77 (s, 3H), 1.76-1.53 (m, 1H), 1.17 (s, 3H), 1.09 (s, 3H), 0.93-0.89 (m, 4H).

Example 8: Synthesis Scheme of **K101-C1306**.

[0268] The scheme for synthesis of compound **K101-C1306** is illustrated below.



[0269] Preparation of Compound **K101-C1306-A**. To a solution of **K101-C20Tr-B** (40.00 mg, 67.71 μmol, 1.00 eq) in DCM (2.00 mL) were added 2-(2-tert-butoxycarbonyl-2-azaspiro[3.3]heptan-6-yl)acetic acid (25.93 mg, 101.56 μmol, 1.50 eq), DMAP (33.09 mg, 270.84 μmol, 4.00 eq), DIEA (26.25 mg, 203.13 μmol, 3.00 eq) and EDC (25.96 mg, 135.42 μmol, 2.00 eq). The mixture

was stirred at 20°C for 16hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The mixture was combined with a second preparation of the compound, quenched with Saturated NaHCO₃ (5 mL) and extracted with DCM (10 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1306-A** (40.00 mg, 47.90 μmol, 70.74% yield, 99.157% purity) as a white solid.

[0270] Preparation of compound K101-C1312. To a solution of **K101-C1306-A** (10.00 mg, 12.08 μmol, 1.00 eq) in MeOH (3.00 mL) was added HClO₄ (83.00 mg, 826.20 μmol, 50.00 uL, 68.39 eq) at 0°C, and the mixture stirred at 0°C for 0.5hr. The mixture was stirred at 20°C for an additional 15.5 hr. LC-MS showed the reaction was complete. The mixture was purified by prep-HPLC (column: Waters Xbridge Prep OBD C18 150 x 30 x 5um; mobile phase: [A: water (0.05% ammonia hydroxide v/v)-B: ACN]; B%: 45%-75%, 10min) to give **K101-C1306-A** (3.80 mg, 6.49 μmol, 53.71% yield) as a white solid.

[0271] LC-MS (m/z): 608.2 [M+Na]⁺

[0272] ¹H NMR (400MHz, CD₃OD) δ 7.56 (s, 1H), 5.62 (s, 1H), 4.60-4.58 (m, 3H), 3.99-0.96 (m, 4H), 3.92-3.82 (m, 2H), 3.19-3.18 (m, 2H), 2.52-2.37 (m, 7H), 2.10-1.94 (m, 4H), 1.77 (s, 3H), 1.76-1.46 (m, 1H), 1.44(s, 9H), 1.18 (s, 3H), 1.08 (s, 3H), 0.93-0.86 (m, 4H).


[0273] Preparation of compound K101-C1306. To a solution of **K101-C1312** (15.00 mg, 25.61 μmol, 1.00 eq) in THF (500.00 uL) was added TFA (288.72 mg, 2.53 mmol, 187.48 uL, 98.88 eq), and the mixture stirred at 20°C for 4hr to give a colorless solution. LC-MS showed the reaction was complete, but mass of P2 was found. The reaction mixture was concentrated by N₂, and the resultant product dissolved in MeOH (500.00 uL) /H₂O (50.00 uL). The mixture was stirred at 20°C for 14hr to give a colorless solution. LC-MS showed the reaction was complete. The mixture was combined with a second preparation of the compound and concentrated by N₂ to give the desired product. The product was lyophilized to give **K101-C1306** (4.20 mg, 7.00 μmol, 25.80% yield, TFA) as a yellow gum. The product (13.4 mg) was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10um; mobile phase: [water (0.1%TFA)-ACN]; B%: 20%-50%, 10min) to give **K101-C1306** (4.20 mg, 7.00 μmol, 25.80% yield, TFA) as a white solid.

[0274] LC-MS (m/z): 508.2 [M+Na]⁺

[0275] ¹H NMR (400MHz, CD₃OD) δ 7.55 (s, 1H), 5.62-5.60 (m, 1H), 4.14 (s, 2H), 4.00 (s, 2H), 3.95 (s, 2H), 3.18 (s, 1H), 3.07 (s, 1H), 2.52-2.46 (m, 7H), 2.10-2.02 (m, 4H), 1.77 (s, 3H), 1.54-1.52 (m, 1H), 1.18(s, 3H), 1.08 (s, 3H), 0.93-0.86 (m, 4H).

Example 9: Synthesis Scheme of **K101-C1311**.

[0276] The scheme for synthesis of compound **K101-C1311** is illustrated below.




[0277] Preparation of Compound K101-C1311. To a solution of **K101-C1305-A** (22.00 mg, 27.03 μmol , 1.00 eq) in MeOH (3.00 mL) was added HClO_4 (26.09 mg, 259.73 μmol , 15.72 μL , 9.61 eq) at 0°C , and the mixture stirred at 0°C for 1hr. LC-MS showed the reaction was complete. The mixture was purified by prep-HPLC (column: Waters Xbridge 150 x 25 x 5 μ ; mobile phase: [A: water (0.05% ammonia hydroxide v/v)-B: ACN]; B%: 40%-70%, 10min) to give **K101-C1311** (3.30 mg, 5.77 μmol , 21.36% yield, 100% purity) as a yellow solid.

[0278] LC-MS (m/z): 594.3 $[\text{M}+\text{Na}]^+$

[0279] ^1H NMR (400MHz, CD_3OD) δ 7.57 (s, 1H), 5.63 (s, 1H), 3.99-0.89 (m, 5H), 3.18-3.05 (m, 3H), 2.56-2.42 (m, 5H), 2.15-2.04 (m, 2H), 1.76 (m, 3H), 1.44 (s, 9H), 1.36-1.31 (m, 3H), 1.17 (s, 3H), 1.09 (s, 3H), 0.93-0.88 (m, 4H).

Example 10: Synthesis Scheme of **K101-C1312**.

[0280] The scheme for synthesis of compound **K101-C1312** is illustrated below.



[0281] Preparation of compound K101-C1312. To a solution of **K101-C1306-A** (10.00 mg, 12.08 μmol , 1.00 eq) in MeOH (3.00 mL) was added HClO_4 (83.00 mg, 826.20 μmol , 50.00 μL , 68.39 eq) at 0°C , and the mixture stirred at 0°C for 0.5 hr. The mixture was stirred at 20°C for an additional 15.5


hr to give a yellow solution. LC-MS showed the reaction was complete. The mixture was purified by prep-HPLC (column: Waters Xbridge Prep OBD C18 150 x 30 x 5u; mobile phase: [A: water (0.05% ammonia hydroxide v/v)-B: ACN]; B%: 45%-75%, 10 min) to give **K101-C1306-A** (3.80 mg, 6.49 μmol , 53.71% yield) as a white solid.

[0282] LC-MS (m/z): 608.2 [M+Na]⁺

[0283] ¹H NMR (400MHz, CD₃OD) δ 7.56 (s, 1H), 5.62 (s, 1H), 4.60-4.58 (m, 3H), 3.99-0.96 (m, 4H), 3.92-3.82 (m, 2H), 3.19-3.18 (m, 2H), 2.52-2.37 (m, 7H), 2.10-1.94 (m, 4H), 1.77 (s, 3H), 1.76-1.46 (m, 1H), 1.44 (s, 9H), 1.18 (s, 3H), 1.08 (s, 3H), 0.93-0.86 (m, 4H).

Example 11: Synthesis Scheme of **K101-C1313**.

[0284] The scheme for synthesis of compound **K101-C1313** is illustrated below.



[0285] Preparation of Compound **K101-C1313-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) in DMF (2.00 mL) were added (E)-3-[4-(trifluoromethyl)phenyl]prop-2-enoic acid (C13-13) (21.95 mg, 101.56 μmol , 2.00 eq), DIEA (19.69 mg, 152.34 μmol , 26.61 μL , 3.00 eq), DMAP (24.82 mg, 203.12 μmol , 4.00 eq) and Hexafluorophosphate Azabenzotriazole Tetramethyl Uronium (HATU) (38.62 mg, 101.56 μmol , 2.00 eq). The mixture was stirred at 20°C for 14 hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc =2/1) showed the reaction was complete. The mixture was quenched with saturated NaHCO₃ (5 mL) and extracted with methyl-tert-butyl ether (MTBE) (15 mL x 3). The organic layers were washed with (H₂O), dried over Na₂SO₄ and then concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1313-A** (21.00 mg, 26.62 μmol , 45.19% yield) as a white solid.

[0286] Preparation of compound **K101-C1313**. To a solution of **K101-C1313-A** (21.00 mg, 26.62 μmol , 1.00 eq) in MeOH (3.00 mL) was added HClO₄ (83.00 mg, 826.28 μmol , 50.00 μL , 31.04 eq) at 0°C. The mixture was stirred at 0°C for 1hr to give a yellow solution. LC-MS showed the reaction


was complete. The mixture was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10um; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 55%-85%, 10min) to give **K101-C1313** (4.40 mg, 6.10 μ mol, 22.90% yield, 91.518% purity, TFA) as a yellow solid.

[0287] LC-MS (m/z): 569.1 [M+Na]⁺

[0288] ¹H NMR (400MHz, CD₃OD) δ 7.74-7.68 (m, 3H), 7.63-7.58 (m, 2H), 7.48-7.16 (m, 1H), 6.60-6.56 (m, 1H), 5.60-5.50 (m, 1H), 3.89 -3.82 (m, 2H), 3.12-3.08 (m, 1H), 3.07-2.90 (m, 1H), 2.42-2.33 (m, 2H), 2.13-2.11 (m, 1H), 2.07-1.99 (m, 1H), 1.66 (s, 3H), 1.58-1.51 (m, 1H), 1.14 (s, 3H), 1.02 (s, 3H), 0.88-0.83 (m, 4H).

Example 12: Synthesis Scheme of **K101-C1315**.

[0289] The scheme for synthesis of compound **K101-C1315** is illustrated below.



[0290] Preparation of Compound **K101-C1315-A**. To a solution of **K101-C20Tr-B** (20.00 mg, 33.86 μ mol, 1.00 *eq*) and **C13-15** (15.35 mg, 101.58 μ mol, 3.00 *eq*) in DCM (1.00 mL) was added EDC (19.47 mg, 101.58 μ mol, 3.00 *eq*) and DMAP (20.68 mg, 169.30 μ mol, 5.00 *eq*). The reaction solution was stirred at 25 °C for 3 hours to give a brown solution. LC-MS showed the reaction was complete. The reaction solution was diluted with DCM (5 mL), washed with brine (2 mL), dried over anhydrous Na₂SO₄, filtered, and the concentrated under reduced pressure to give **K101-C1315-A** (28.70 mg, crude) as a brown gum, which was used directly in the next step without further purification.


[0291] Preparation of compound **K101-C1315**. To a solution of **K101-C1315-A** (25.00 mg, crude) in MeOH (1.00 mL) was added HClO₄ (30.00 μ L) at 25 °C. The reaction solution was stirred at 25 °C for 0.5 hour to give a brown solution. LC-MS showed the reaction was complete. The reaction solution was quenched by adding K₂CO₃ (34 mg) in water (1 mL) dropwise at 0 °C to adjust the pH to 9. The mixture was filtered and the filtrate purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10um; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 15%-45%, 10min) to give **K101-C1315** (3.60 mg, 97% purity, TFA salt) as a white solid after lyophilization.

[0292] LC-MS (m/z): 504.2 [M+Na]⁺

[0293] $^1\text{H NMR}$ (400MHz, CD_3OD) δ = 8.72 (d, $J=6.5$ Hz, 2H), 7.95 (d, $J=6.5$ Hz, 2H), 7.53 (s, 1H), 5.61-5.55 (m, 1H), 3.99-3.89 (m, 2H), 3.24 (t, $J=7.3$ Hz, 2H), 3.17-3.11 (m, 1H), 3.06-3.01 (s, 1H), 2.90 (t, $J=7.2$ Hz, 2H), 2.57-2.47 (m, 1H), 2.45-2.37 (m, 1H), 2.15-1.95 (m, 2H), 1.77-1.72 (m, 3H), 1.48 (dd, $J=10.5, 14.3$ Hz, 1H), 1.10 (s, 3H), 1.05 (s, 3H), 0.92-0.83 (m, 4H).

Example 13: Synthesis Scheme of **K101-C1316**.

[0294] The scheme for synthesis of compound **K101-C1316** is illustrated below.



[0295] Preparation of Compound **K101-C1316-A**. To a solution of **K101-C20Tr-B** (40.00 mg, 67.71 μmol , 1.00 eq) in DCM (2.00 mL) were added **C13-16** (51.18 mg, 338.55 μmol , 5.00 eq), DMAP (33.09 mg, 270.84 μmol , 4.00 eq), hydroxybenzotriazole (HOBt) (18.30 mg, 135.42 μmol , 2.00 eq) and EDC (25.96 mg, 135.42 μmol , 2.00 eq). The mixture was stirred at 20°C for 12h to give a black solution. LC-MS showed 43.779% of the desired mass, with 16.864% of reactant remaining. The mixture was quenched with saturated NaHCO_3 (10 mL) and extracted with DCM (15 mL x 3). The organic layers were dried over Na_2SO_4 and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1316-A** (27.00 mg, 37.30 μmol , 48.97% yield) as a white solid.


[0296] Preparation of Compound **K101-C1316**. To a solution of **K101-C1316-A** (27.00 mg, 37.30 μmol , 1.00 eq) in MeOH (2.00 mL) was added HClO_4 (83.00 mg, 826.20 μmol , 50.00 μL , 22.15 eq) at 0°C. The mixture was stirred at 0°C for 0.5 hr to give a yellow solution. LC-MS showed the reaction was complete. The mixture was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 20%-50%, 10min) to give **K101-C1316** (11.40 mg, 18.81 μmol , 50.43% yield, 98.3% purity, TFA salt) as a white solid.

[0297] LC-MS (m/z): 504.2 $[\text{M}+\text{Na}]^+$

[0298] $^1\text{H NMR}$ (400MHz, CD_3OD) δ 8.72-8.71 (d, $J= 5.2\text{Hz}$, 1H), 8.45-8.41 (t, $J= 8.0\text{Hz}$, 1H), 7.95-7.93 (d, $J= 8.0\text{Hz}$, 1H), 7.86-7.83 (t, $J= 6.8\text{Hz}$, 1H), 7.55 (s, 1H), 5.60-5.59 (m, 1H), 3.99-3.95 (m, 2H), 3.31-3.29 (m, 2H), 3.16 (s, 1H), 3.05 (s, 1H), 3.00-2.96 (m, 2H), 2.51-2.40 (m, 2H), 2.15-2.03 (m, 2H), 1.76-1.75 (m, 3H), 1.51-1.47 (m, 3H), 1.10 (s, 3H), 1.06 (s, 3H), 0.90-0.86 (m, 4H).

Example 14: Synthesis Scheme of **K101-C1317**.

[0299] The scheme for synthesis of compound **K101-C1317** is illustrated below.



[0300] Preparation of Compound **K101-C1317-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) in DCM (1.00 mL) were added (2S)-2-(tert-butoxycarbonylamino) hexanoic acid (C13-17) (23.49 mg, 101.57 μmol , 2.00 *eq*), DMAP (24.82 mg, 203.13 μmol , 4.00 *eq*) and EDC (19.47 mg, 101.57 μmol , 2.00 *eq*). The mixture was stirred at 20°C for 5 h to give a colorless solution. LC-MS and TLC showed the reaction was complete. The mixture was quenched with H₂O (15 mL) and extracted with DCM (15 mL x 5). The organic layers were dried over Na₂SO₄ and concentrated to give a yellow solid. The product was purified by prep-TLC (eluting with Petroleum ether: Ethyl acetate = 5/1) to give **K101-C1317-A** (35.00 mg, 43.53 μmol , 85.73% yield) as a white solid.


[0301] Preparation of Compound **K101-C1317**. To a solution of **K101-C1317-A** (46.00 mg, 57.21 μmol , 1.00 *eq*) in THF (2.00 mL) were added TFA (6.52 mg, 57.21 μmol , 4.23 μL , 1.00 *eq*) and Et₃SiH (6.65 mg, 57.21 μmol , 9.11 μL , 1.00 *eq*). The mixture was stirred at 20°C for 2 h to give a colorless solution, which was concentrated to give yellow oil. TFA (1 mL) was added to the yellow oil in DCM (2 mL), and the mixture stirred at 20°C for 0.5h. LC-MS showed the reaction was complete. The reaction mixture was concentrated, dissolved with MeOH (20 mL), and stirred at 20°C for 14 h to give a yellow liquid. The product was concentrated to give a yellow solid, which was then purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%:30%-60%, 8 min). The separated layers were lyophilized to give **K101-C1317** (7.00 mg, 12.16 μmol , 21.26% yield, 100% purity, TFA) as a white solid.

[0302] LC-MS (m/z): 584.2 [M+Na]⁺

[0303] ¹H NMR (400MHz, MeOD) δ 7.57 (s, 1H), 5.64 (d, *J*=4.5 Hz, 1H), 4.04 (t, *J*=6.4 Hz, 1H), 4.00-3.94 (m, 2H), 3.20-3.15 (m, 1H), 3.10-3.05 (m, 1H), 2.59 - 2.39 (m, 2H), 2.27 (dd, *J*=7.0, 14.8 Hz, 1H), 2.13- 1.93 (m, 2H), 1.90-1.80 (m, 1H), 1.80-1.70 (d, *J*=1.5 Hz, 3H), 1.60-1.50 (dd, *J*=10.4, 14.9 Hz, 1H), 1.53 - 1.37 (m, 4H), 1.20 (s, 3H), 1.11 (s, 3H), 1.05 - 0.91 (m, 7H).

Example 15: Synthesis Scheme of **K101-C1318**.

[0304] The scheme for synthesis of compound **K101-C1318** is illustrated below.



[0305] Preparation of Compound **K101-C1318-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) in DCM (2.00 mL) were added (2S)-2-(tert-butoxycarbonylamino)-4-phenylbutanoic acid (C13-18) (28.37 mg, 101.56 μmol , 2.00 eq), DMAP (24.82 mg, 203.12 μmol , 4.00 eq) and EDC (19.47 mg, 101.56 μmol , 2.00 eq). The mixture was stirred at 20°C for 12hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture combined a second preparation of the compound, the mixture quenched with H₂O (10 mL) and then extracted with DCM (15 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1318-A** (32.00 mg, 37.56 μmol , 63.38% yield) as a white solid.


[0306] Preparation of Compound **K101-C1318**. To a solution of **K101-C1318-A** (30.00 mg, 35.21 μmol , 1.00 eq) in THF (2.00 mL) were added TFA (308.00 mg, 2.70 mmol, 200.00 μL , 76.72 eq) and Et₃SiH (4.91 mg, 42.25 μmol , 6.73 μL , 1.20 eq). The mixture was stirred at 20°C for 4hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated by N₂, and the resultant residue dissolved in MeOH (20 mL). The mixture was stirred at 20°C for 12 hr. LC-MS showed the reaction was complete. The mixture was concentrated to give the crude product. The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 23%-53%, 10 min) to give **K101-C1318** (10.80 mg, 17.24 μmol , 48.98% yield, 99.58% purity, TFA) as a white solid.

[0307] LC-MS (m/z): 532.1 [M+Na]⁺

[0308] ¹H NMR (400MHz, CD₃OD) δ 7.58 (s, 1H), 7.36-7.33 (m, 2H), 7.27-7.23 (m, 3H), 5.65 (s, 1H), 4.07-4.03 (m, 1H), 3.97 (s, 2H), 3.18 (s, 1H), 3.08 (m, 1H), 2.83-2.81 (m, 2H), 2.53-2.41 (m, 2H), 2.10-2.09 (m, 2H), 1.77 (s, 3H), 1.64-1.61 (m, 1H), 1.21(s, 3H), 1.12 (s, 3H), 1.05-1.04 (m, 1H), 0.97-0.95 (m, 3H).

Example 16: Synthesis Scheme of **K101-C1319**.

[0309] The scheme for synthesis of compound **K101-C1319** is illustrated below.



[0310] Preparation of compound **K101-C1319-A**. To a solution of **K101-C20Tr-B** (200.00 mg, 338.55 μmol , 1.00 eq) and **C13-19** (119.18 mg, 406.26 μmol , 1.20 eq) in anhydrous DCM (2.00 mL) were added EDC (194.70 mg, 1.02 mmol, 3.00 eq) and DMAP (124.08 mg, 1.02 mmol, 3.00 eq). The reaction solution was stirred at 20°C for 16 hours to give a light brown solution. LC-MS showed the reaction was complete. The reaction solution was concentrated under reduced pressure to give the crude product, and the product purified by silica gel column chromatography (PE/EtOAc=3/1) to give **K101-C1319-A** (273.50 mg, 315.79 μmol , 93.28% yield) as a colorless gum.


[0311] Preparation of compound **K101-C1319**. To a solution of **K101-C1319-A** (273.00 mg, 315.21 μmol , 1.00 eq) in MeOH (5.00 mL) was added HCl/MeOH (4 M, 5.00 mL, 63.45 eq) at 0°C. The reaction solution was stirred at 0°C for 3.5 h to give a clear solution. LC-MS showed the reaction was not complete, and thus the reaction solution was stirred at 20°C for 1.5 h. LC-MS showed the reaction was complete. The reaction solution was bubbled with N₂ for 0.5 hour to remove HCl, and the residual solution cooled to 0°C and the solution adjusted to pH 7 with saturated aqueous NaHCO₃. The mixture was extracted with DCM (10 mL x 2), and the combined extract dried over Na₂SO₄ and concentrated under reduced pressure to give the crude product as a brown gum. The product was purified by prep-TLC (DCM/MeOH=10/1, SiO₂) to give **K101-C1319** (78.70 mg, 140.52 μmol , 44.58% yield, 93.5% purity) as a colorless gum. The product was lyophilized to give a white solid.

[0312] MS (m/z): 546.2 [M+Na]⁺

[0313] ¹H NMR (400MHz, CD₃OD) δ 7.55 (s, 1H), 7.29-7.23 (m, 2H), 7.21-7.13 (m, 3H), 5.60 (d, $J=5.0$ Hz, 1H), 4.00-3.88 (m, 2H), 3.47 (t, $J=5.8$ Hz, 1H), 3.19-3.14 (m, 1H), 3.10-3.03 (m, 1H), 2.65 (t, $J=7.0$ Hz, 2H), 2.57-2.48 (m, 1H), 2.48-2.38 (m, 1H), 2.17-1.99 (m, 2H), 1.80-1.62 (m, 7H), 1.51 (dd, $J=10.5, 14.3$ Hz, 1H), 1.15 (s, 3H), 1.07 (s, 3H), 0.90 (d, $J=6.3$ Hz, 4H).

Example 17: Synthesis Scheme of **K101-C1320**.

[0314] The scheme for synthesis of compound **K101-C1320** is illustrated below.



[0315] **Preparation of Compound K101-C1320-A.** To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) and **C13-20** (26.09 mg, 152.34 μmol , 3.00 eq) in DCM (2.00 mL) were added EDC (58.41 mg, 304.68 μmol , 6.00 eq) and DMAP (37.22 mg, 304.68 μmol , 6.00 eq). The reaction solution was stirred at 25 °C for 16 hours to give a brown solution. TLC (PE/EtOAc=2/1, SiO_2) showed the reaction was complete. The reaction solutions were combined 5 mg of **K101-C20Tr-B**, diluted with DCM (5 mL), and washed with brine (2 mL). The extracted layers were dried over anhydrous Na_2SO_4 , filtered, and then concentrated under reduced pressure to give the crude product as a brown gum. The product was purified by prep-TLC (PE/EtOAc=2/1) to give **K101-C1320-A** (27.30 mg, 72.26% yield) as a colorless gum.


[0316] **Preparation of Compound K101-C1320.** To a solution of **K101-C1320-A** (25.00 mg, 33.60 μmol , 1.00 eq) in MeOH (1.00 mL) was added HClO_4 (30.00 μL). The reaction solution was stirred at 25 °C for 0.5 hour to give a clear solution. LC-MS showed the reaction was complete. The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 25%-55%, 10min) to give **K101-C1320** (9.50 mg, 18.46 μmol , 54.95% yield, 97.5% purity) as a white powder after lyophilization.

[0317] LC-MS (m/z): 524.1 $[\text{M}+\text{Na}]^+$

[0318] ^1H NMR (400MHz, METHANOL- d_4) δ = 9.04 (s, 1H), 7.56-7.51 (m, 1H), 5.58 (d, $J=4.3$ Hz, 1H), 3.99-3.89 (m, 2H), 3.19-3.13 (m, 3H), 3.04 (t, $J=5.4$ Hz, 1H), 2.72 (dt, $J=1.6, 7.0$ Hz, 2H), 2.56-2.47 (m, 1H), 2.46-2.39 (m, 4H), 2.13-1.98 (m, 2H), 1.74 (dd, $J=1.3, 3.0$ Hz, 3H), 1.47 (dd, $J=10.4, 14.2$ Hz, 1H), 1.06 (d, $J=10.5$ Hz, 6H), 0.88 (d, $J=6.3$ Hz, 3H), 0.82 (d, $J=5.8$ Hz, 1H).

Example 18: Synthesis Scheme of **K101-C1321**.

[0319] The scheme for synthesis of compound **K101-C1321** is illustrated below.



[0320] Preparation of Compound **K101-C1321-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) and **C13-21** (76.68 mg, 304.68 μmol , 6.00 *eq*) in DCM (1.00 mL) were added EDC (58.41 mg, 304.68 μmol , 6.00 *eq*) and DMAP (37.22 mg, 304.68 μmol , 6.00 *eq*). The reaction solution was stirred at 25 °C for 2 hours to give a light brown solution. LC-MS shows improved conversion to product. The reaction solution was combined with a second preparation of the compound, and the mixture partitioned between water (2 mL) and DCM (2 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered, and then concentrated under reduced pressure to give the crude product as a brown gum. The product was purified by prep-TLC (PE/EtOAc=3/2) to give **K101-C1321-A** (32.70 mg, 39.67 μmol , 78.11% yield) as a colorless gum.


[0321] Preparation of Compound **K101-C1321**. To a solution of **K101-C1321-A** (32.70 mg, 39.67 μmol , 1.00 *eq*) in MeOH (1.00 mL) was added HClO₄ (30.00 μL) at 25 °C. The reaction solution was stirred at 25 °C for 0.5 hour to give a brown solution. LC-MS showed the reaction was complete. The reaction solution was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water(0.1%TFA)-B: ACN]; B%: 50%-80%, 10min) to give **K101-C1321** (12.50 mg, 52.08% yield, 96.2% purity) as a white solid after lyophilization.

[0322] LC-MS (m/z): 604.1 [M+Na]⁺

[0323] ¹H NMR (400MHz, CD₃OD) δ 7.69-7.65 (m, 2H), 7.54-7.50 (m, 1H), 7.48-7.41 (m, 3H), 5.56-5.52 (m, 1H), 3.95-3.87 (m, 2H), 3.20-3.12 (m, 3H), 3.06-3.01 (m, 1H), 2.94-2.89 (m, 2H), 2.55-2.47 (m, 1H), 2.45-2.38 (m, 1H), 2.13-1.98 (m, 2H), 1.73 (dd, *J*=1.3, 3.0 Hz, 3H), 1.60-1.52 (m, 1H), 1.12 (s, 3H), 1.04 (s, 3H), 0.89-0.83 (m, 4H).

Example 19: Synthesis Scheme of **K101-C1322**.

[0324] The scheme for synthesis of compound **K101-C1322** is illustrated below.



[0325] Preparation of Compound K101-C1322-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) in DCM (2.00 mL) were added 3-(1H-pyrrolo [2, 3-b] pyridin-3-yl)propanoic acid (C13-22) (19.32 mg, 101.56 μmol , 2.00 eq), DMAP (24.82 mg, 203.12 μmol , 4.00 eq), HOBT (13.72 mg, 101.56 μmol , 2.00 eq) and EDC (19.47 mg, 101.56 μmol , 2.00 eq). The mixture was stirred at 20°C for 12h to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=1/1) showed the reaction was complete. The mixture was quenched with H₂O (5 mL) and extracted with DCM (15 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=1/1) to give **K101-C1322-A** (30.00 mg, 39.32 μmol , 58.22% yield) as a white solid.


[0326] Preparation of Compound K101-C1322. To a solution of **K101-C1322-A** (30.00 mg, 39.32 μmol , 1.00 eq) in MeOH (2.00 mL) was added HClO₄ (83.00 mg, 826.11 μmol , 50.00 μL , 21.01 eq) at 0°C. The mixture was stirred at 0°C for 0.5h to give a yellow solution. LC-MS showed the reaction was complete. The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 22%-52%, 10min) to give **K101-C1322** (17.40 mg, 27.42 μmol , 69.73% yield, TFA salt) as a yellow solid.

[0327] LC-MS (m/z): 543.1 [M+Na]⁺

[0328] ¹H NMR (400MHz, CD₃OD) δ 8.19-8.18 (m, 1H), 8.06-8.04 (m, 1H), 7.55 (s, 1H), 7.24 (s, 1H), 7.15-7.12 (m, 1H), 5.55-5.54 (m, 1H), 3.99-3.95 (m, 2H), 3.16-3.10 (m, 3H), 3.01 (s, 1H), 2.78-2.73 (m, 2H), 2.50-2.46 (m, 2H), 2.01-2.96 (m, 2H), 1.42-1.41 (m, 1H), 1.00 (s, 3H), 0.91 (s, 3H), 0.86-0.85 (m, 1H), 0.63-0.62 (m, 3H).

Example 20: Synthesis Scheme of **K101-C1323**.

[0329] The scheme for synthesis of compound **K101-C1323** is illustrated below.



[0330] Preparation of compound K101-C1323-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) and **C13-23** (23.49 mg, 152.34 μmol , 3.00 eq) in DCM (1.00 mL) were added EDC (58.41 mg, 304.68 μmol , 6.00 eq) and DMAP (37.22 mg, 304.68 μmol , 6.00 eq). The reaction solution was stirred at 25°C for 2 h to give a brown solution. TLC (PE/EtOAc=1/1, SiO_2) showed the reaction was complete. The reaction solution was combined with a second preparation of the compound, diluted with DCM (2 mL), washed with water (1 mL), brine (1 mL), and dried over anhydrous Na_2SO_4 . The mixture was filtered and then concentrated under reduced pressure to give the crude product as a brown gum. The product was purified by prep-TLC (PE/EtOAc=1/1) to give 32.5 mg of **K101-C1323-A** as a colorless gum.


[0331] Preparation of Compound K101-C1323. To a solution of **K101-C1323-A** (32.00 mg, 44.02 μmol , 1.00 eq) in MeOH (1.00 mL) was added HClO_4 (30.00 μL) at 25°C. The reaction solution was stirred at 25°C for 0.5 hour to give a clear solution. LC-MS showed the reaction was complete. The reaction solution was directly purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1% TFA)-B: ACN]; B%: 25%-55%, 10min) to give **K101-C1323** (9.10 mg, 40.48% yield, 94.9% purity) as a white solid.

[0332] LC-MS (m/z): 507.2 $[\text{M}+\text{Na}]^+$

[0333] ^1H NMR (400MHz, CD_3OD) δ 7.56-7.53 (m, 1H), 7.46 (s, 1H), 7.36 (s, 1H), 5.61-5.57 (m, 1H), 3.98-3.90 (m, 2H), 3.84 (s, 3H), 3.18-3.14 (m, 1H), 3.07-3.01 (m, 1H), 2.81-2.76 (m, 2H), 2.63-2.57 (m, 2H), 2.55-2.47 (m, 1H), 2.46-2.39 (m, 1H), 2.12-1.98 (m, 2H), 1.74 (dd, $J=1.3, 2.8$ Hz, 3H), 1.47 (dd, $J=10.4, 14.2$ Hz, 1H), 1.08 (s, 3H), 1.05 (s, 3H), 0.89 (d, $J=6.3$ Hz, 3H), 0.79 (d, $J=5.8$ Hz, 1H).

Example 21: Synthesis Scheme of **K101-C1324**.

[0334] The scheme for synthesis of compound **K101-C1324** is illustrated below.



[0335] Preparation of Compound K101-C1324-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) and **C13-24** (25.01 mg, 152.34 μmol , 3.00 *eq*) in DCM (1.00 mL) were added EDC (58.41 mg, 304.68 μmol , 6.00 *eq*) and DMAP (37.22 mg, 304.68 μmol , 6.00 *eq*). The reaction solution was stirred at 25 °C for 2h to give a brown solution. TLC (PE/EtOAc=2/1, SiO₂) showed the reaction was complete. The reaction solution was diluted with DCM (2 mL), washed with water (1 mL), brine (1 mL), and dried over anhydrous Na₂SO₄. The mixture was filtered and then concentrated under reduced pressure to give the crude product as a brown gum. The product was purified by silica gel column chromatography (eluting with PE/EtOAc=5/1) to give **K101-C1324-A** (37.20 mg, 99.41% yield) as a colorless gum.


[0336] Preparation of Compound K101-C1324. To a solution of **K101-C1324-A** (37.20 mg, 50.48 μmol , 1.00 *eq*) in MeOH (1.00 mL) was added HClO₄ (30.00 μL), and the reaction solution stirred at 25 °C for 1 hour to give a clear solution. LC-MS showed the reaction was complete. The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1% TFA)-B: ACN]; B%: 55%-85%, 10min) to give **K101-C1324** (7.80 mg, 30.40% yield, 97.3% purity) as a white solid after lyophilization.

[0337] LC-MS (m/z): 517.2 [M+Na]⁺

[0338] ¹H NMR (400MHz, CD₃OD) δ 7.55 (s, 1H), 7.30-7.24 (m, 2H), 7.21-7.14 (m, 3H), 5.63-5.58 (m, 1H), 4.00-3.89 (m, 2H), 3.19-3.15 (m, 1H), 3.09-3.03 (m, 1H), 2.65 (t, *J*=7.7 Hz, 2H), 2.57-2.48 (m, 1H), 2.47-2.39 (m, 1H), 2.34 (t, *J*=7.3 Hz, 2H), 2.15-2.07 (m, 1H), 2.07-1.98 (m, 1H), 1.97-1.88 (m, 2H), 1.74 (dd, *J*=1.3, 2.8 Hz, 3H), 1.53 (dd, *J*=10.5, 14.3 Hz, 1H), 1.16 (s, 3H), 1.07 (s, 3H), 0.91 (d, *J*=6.3 Hz, 3H), 0.85 (d, *J*=5.5 Hz, 1H).

Example 22: Synthesis Scheme of **K101-C1325**.

[0339] The scheme for synthesis of compound **K101-C1325** is illustrated below.



[0340] Preparation of Compound K101-C1325-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) in DCM (5.00 mL) were added (2S)-2-(tert-butoxycarbonylamino)-3-cyclobutyl-propanoic acid (C13-25) (24.71 mg, 101.56 μmol , 2.00 *eq*), DMAP (24.82 mg, 203.12 μmol , 4.00 *eq*), HOBT (13.72 mg, 101.56 μmol , 2.00 *eq*) and EDC (19.47 mg, 101.56 μmol , 2.00 *eq*). The mixture was stirred at 20°C for 14h to give a yellow solution. LC-MS and TLC showed the reaction was complete. The reaction mixture was quenched with H₂O (15 mL) and extracted with DCM (15 mL x 5). The organic layers were dried over Na₂SO₄ and concentrated to give a yellow solid. The product was purified by prep-TLC (eluting with Petroleum ether: Ethyl acetate = 7/2) to give **K101-C1325-A** (33.50 mg, 41.05 μmol , 69.09% yield) as a white solid.


[0341] Preparation of Compound K101-C1325. To a solution of **K101-C1325-A** (33.50 mg, 41.05 μmol , 1.00 *eq*) in DMF (20.00 mL) in THF (2.00 mL) were added TFA (1.54 g, 13.51 mmol, 1.00 mL, 329.02 *eq*) and Et₃SiH (4.77 mg, 41.05 μmol , 6.53 μL , 1.00 *eq*). The mixture was stirred at 20°C for 5h to give a colorless solution, which was concentrated to give a yellow oil. The oil was dissolved with DCM (2 mL) followed by addition of TFA (0.5 mL). The mixture was stirred at 20°C for 1h to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated to give a yellow oil, which was then purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 20%-50%, 10min). The separated products were lyophilized to give **K101-C1325** (10.00 mg, 17.02 μmol , 41.46% yield, 94.5% purity, TFA) as a white solid.

[0342] LC-MS (m/z): 596.2 [M+Na]⁺

[0343] ¹H NMR (400MHz, MeOH) δ = 7.54 (s, 1H), 5.60 (d, *J*=4.0 Hz, 1H), 3.99-3.83 (m, 3H), 3.14 (s, 1H), 3.07-2.98 (m, 1H), 2.57-2.33 (m, 3H), 2.26-1.84 (m, 8H), 1.79-1.65 (s, 5H), 1.58 (dd, *J*=10.4, 14.8 Hz, 1H), 1.17 (s, 3H), 1.07 (s, 3H), 0.99-0.95 (m, 1H), 0.97 (d, *J*=6.0 Hz, 1H), 0.92 (d, *J*=6.6 Hz, 3H)

Example 23: Synthesis Scheme of **K101-C1326**.

[0344] The scheme for synthesis of compound **K101-C1326** is illustrated below.



[0345] Preparation of Compound K101-C1326-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) in DCM (5.00 mL) were added (2S)-2-(tert-butoxycarbonylamino)-4-methylsulfonyl-butanoic acid (C13-26) (30.00 mg, 106.64 μmol , 2.10 *eq*), DMAP (30.00 mg, 245.78 μmol , 4.84 *eq*), EDC (19.96 mg, 104.10 μmol , 2.05 *eq*), and HOBt (14.00 mg, 103.59 μmol , 2.04 *eq*). The mixture was stirred at 20°C for 19h to give a colorless solution. LC-MS and TLC showed the reaction was complete. The reaction mixture was quenched with H₂O (15 mL) and extracted with DCM (15 mL x 5). The organic layers were dried over Na₂SO₄ and concentrated to give a yellow oil. The product was purified by prep-TLC (eluting with Petroleum ether: Ethyl acetate = 3/2) to give **K101-C1326-A** (23.00 mg, 26.93 μmol , 53.03% yield, crude product) as a colorless solid.


[0346] Preparation of compound K101-C1326. To a solution of **K101-C1326-A** (25.00 mg, 29.27 μmol , 1.00 *eq*) in THF (2.00 mL) were added TFA (3.34 mg, 29.27 μmol , 2.17 μL , 1.00 *eq*) and Et₃SiH (3.40 mg, 29.27 μmol , 4.66 μL , 1.00 *eq*). The mixture was stirred at 20°C for 3h to give a colorless solution. LC-MS showed some of the **K101-C1326-A** remained. Accordingly, the reaction mixture was concentrated to give a yellow oil, which was then dissolved with DCM (2 mL) followed by the addition of TFA (2 mL). The mixture was stirred at 20°C for 2h to give a colorless solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated to give a yellow oil, which was then purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10um; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 10%-40%, 10min). The separated layers were lyophilized to give **K101-C1326** (4.20 mg, 6.04 μmol , 20.64% yield, 90% purity, TFA) as a yellow solid.

[0347] LC-MS (m/z): 534.1 [M+Na]⁺

[0348] ¹H NMR (400MHz, MeOD) δ = 7.58 (s, 1H), 5.66 (s, 1H), 4.62 (s, 1H), 4.16 (s, 1H), 3.97 (s, 2H), 3.23 - 3.11 (m, 1H), 3.06 (s, 4H), 2.61 - 2.35 (m, 3H), 2.27 (dd, *J*=7.2, 14.7 Hz, 2H), 2.06 (s, 1H), 1.77 (d, *J*=1.5 Hz, 3H), 1.62 (dd, *J*=10.8, 14.8 Hz, 1H), 1.40 - 1.28 (m, 2H), 1.21 (s, 3H), 1.12 (s, 3H), 1.05 (d, *J*=6.0 Hz, 1H), 0.96 (d, *J*=6.5 Hz, 3H).

Example 24: Synthesis Scheme of **K101-C1327**.

[0349] The scheme for synthesis of compound **K101-C1327** is illustrated below.



[0350] Preparation of Compound K101-C1327-A. To a solution of **K101-C20Tr-B** (200.00 mg, 338.55 μmol , 1.00 eq) in DCM (2.00 mL) were added **C13-27** (367.46 mg, 1.35 mmol, 4.00 eq), DMAP (330.89 mg, 2.71 mmol, 8.00 eq), HOBt (91.49 mg, 677.11 μmol , 2.00 eq) and EDC (259.60 mg, 1.35 mmol, 4.00 eq). The mixture was stirred at 20°C for 12h to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The mixture was quenched with H₂O (15 mL) and extracted with DCM (30 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1327-A** (180.00 mg, 213.25 μmol , 62.99% yield) as a white solid.

[0351] Preparation of Compound K101-C1327. To a solution of **K101-C1327-A** (180.00 mg, 213.25 μmol , 1.00 eq) in THF (3.00 mL) were added TFA (3.08 g, 27.01 mmol, 2.00 mL, 126.67 eq) and Et₃SiH (49.59 mg, 426.50 μmol , 67.93 μL , 2.00 eq). The mixture was stirred at 20°C for 24h to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated by N₂, and the resultant residue dissolved in MeOH (20 mL) and then stirred at 20°C for 70h. LC-MS showed the reaction was complete. The mixture was concentrated to give the crude product. The crude product was triturated with PE (30 mL x 3) to give the desired product. The product was dissolved in saturated NaHCO₃ (20 mL) and extracted with DCM (30 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the free desired product, which was then lyophilized to give **K101-C1327** (70.00 mg, 136.05 μmol , 63.80% yield, 97.5% purity) as a white solid.

[0352] LC-MS (m/z): 524.2 [M+Na]⁺

[0353] ¹H NMR (400MHz, CD₃OD) δ 7.57 (s, 1H), 5.64-5.63 (m, 1H), 4.00-3.93 (m, 2H), 3.53-3.49 (m, 1H), 3.19 (s, 1H), 3.09 (s, 1H), 2.57-2.47 (m, 2H), 2.19-2.17 (m, 2H), 1.80-1.77 (m, 8H), 1.73-1.60 (m, 2H), 1.48-1.47 (m, 2H), 1.29-1.26 (m, 3H), 1.21 (s, 3H), 1.10 (s, 3H), 1.02-1.01 (m, 2H), 0.95-0.92 (m, 3H).

[0354] Second Procedure for Preparation of Compound K101-C1327-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) in DCM (2.00 mL) were added (2S)-2-(tert-butoxycarbonylamino)-3-cyclohexyl-propanoic acid (**C13-27**) (27.56 mg, 101.56 μmol , 2.00 eq), DMAP (24.82 mg, 203.12 μmol , 4.00 eq) and EDC (19.47 mg, 101.56 μmol , 2.00 eq). The mixture was stirred at 20°C for 12hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The mixture was quenched with H₂O (10 mL) and extracted with DCM (15 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1327-A** (24.00 mg, 28.43 μmol , 47.97% yield) as a white solid.


[0355] Second Procedure for Preparation of Compound K101-C1327. To a solution of **K101-C1327-A** (24.00 mg, 28.43 μmol , 1.00 eq) in THF (2.00 mL) were added TFA (308.00 mg, 2.70 mmol, 200.00 μL , 95.02 eq) and Et₃SiH (3.97 mg, 34.12 μmol , 5.43 μL , 1.20 eq). The mixture was stirred at 20°C for 4hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated by N₂, and the resultant residue dissolved in MeOH (20 mL). The mixture was stirred at 20°C for 12hr. LC-MS showed the reaction was complete. The mixture was concentrated to give the crude product. The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 25%-55%, 10min) to give **K101-C1327** (5.10 mg, 7.95 μmol , 27.97% yield, 95.99% purity, TFA) as a white solid.

[0356] LC-MS (m/z): 524.2 [M+Na]⁺

[0357] ¹H NMR (400MHz, CD₃OD) δ 7.57 (s, 1H), 5.64 (s, 1H), 4.10-4.07 (m, 1H), 4.00-3.93 (s, 2H), 3.18 (s, 1H), 3.08 (s, 1H), 2.53-2.45 (m, 1H), 2.45-2.41 (m, 1H), 2.29-2.27 (m, 1H), 2.05 (m, 1H), 1.84-1.29 (m, 16H), 1.19(s, 3H), 1.11 (s, 3H), 1.02-1.01 (m, 2H), 0.96-0.94 (m, 3H).

Example 25: Synthesis Scheme of **K101-C1328**.

[0358] The scheme for synthesis of compound **K101-C1328** is illustrated below.



[0359] Preparation of Compound K101-C1328-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) in DCM (2.00 mL) were added (2S)-2-(tert-butoxycarbonylamino)-4, 4-dimethyl-pentanoic acid (**C13-28**) (24.92 mg, 101.56 μmol , 2.00 eq), DMAP (24.82 mg, 203.12 μmol , 4.00 eq), HOBT (13.72 mg, 101.56 μmol , 2.00 eq) and EDC (19.47 mg, 101.56 μmol , 2.00 eq). The mixture was stirred at 20°C for 12h to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The mixture was quenched with H₂O (10 mL) and extracted with DCM (15 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1328-A** (28.00 mg, 34.23 μmol , 67.40% yield) as a white solid.


[0360] Preparation of Compound K101-C1328. To a solution of **K101-C1328-A** (28.00 mg, 34.23 μmol , 1.00 eq) in THF (2.00 mL) were added TFA (770.00 mg, 6.75 mmol, 500.00 μL , 197.29 eq) and Et₃SiH (7.96 mg, 68.46 μmol , 10.90 μL , 2.00 eq). The mixture was stirred at 20°C for 12h to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated by N₂ and the resultant product purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 30%-70%, 10min) to give **K101-C1328** (9.50 mg, 15.88 μmol , 46.38% yield, 98.54% purity, TFA) as a white solid.

[0361] LC-MS (m/z): 498.2 [M+Na]⁺

[0362] ¹H NMR (400MHz, CD₃OD) δ 7.57 (s, 1H), 5.64-5.63 (m, 1H), 4.07-4.00 (m, 1H), 3.97-3.93 (m, 2H), 3.18 (s, 1H), 3.09 (s, 1H), 2.52-2.45 (m, 1H), 2.45-2.38 (m, 1H), 2.27-2.23 (m, 1H), 2.05-2.00 (m, 2H), 1.77(s, 3H), 1.64-1.61 (m, 2H), 1.21(s, 3H), 1.12 (s, 3H), 1.05 (s, 9H), 1.03-1.01 (m, 1H), 0.96-0.94 (m, 3H).

Example 26: Synthesis Scheme of **K101-C1329**.

[0363] The scheme for synthesis of compound **K101-C1329** is illustrated below.



[0364] Preparation of Compound K101-C1329-A. To a solution of **K101-C20Tr-B** (20.00 mg, 33.86 μmol , 1.00 eq) and **C13-29** (14.80 mg, 50.79 μmol , 1.50 eq) in DCM (2.00 mL) were added EDC (38.95 mg, 203.16 μmol , 6.00 eq) and DMAP (24.82 mg, 203.16 μmol , 6.00 eq). The reaction

solution was stirred at 25 °C for 16 hours to give a brown solution. LC-MS showed the reaction was complete. The reaction solution was diluted with DCM (10 mL), then washed with water (3 mL), 0.5 M HCl (2 mL), brine (2 mL), and dried over anhydrous Na₂SO₄. The product was filtered and then concentrated under reduced pressure to give crude **K101-C1329-A**. The crude **K101-C1329-A** was purified by prep-TLC (PE/EtOAc=3/1, SiO₂) to give 15.7 mg of **K101-C1329-A** as a colorless gum.

[0365] Preparation of Compound **K101-C1329**. To a solution of **K101-C1329-A** (15.70 mg, 18.17 μmol, 1.00 eq) in dioxane (400.00 μL) was added HCl/dioxane (4 M, 200.46 μL, 44.13 eq). The reaction mixture was stirred at 25 °C for 2 hours to give a light brown solution. LC-MS showed the reaction was not complete so the reaction solution was stirred at 25 °C for an additional 1 hour and then for an additional 16 hours. LC-MS showed the reaction was complete. A byproduct was also detected by the LC-MS analysis. The reaction solution was diluted with CH₃CN (1mL) and the solution adjusted with K₂CO₃ (55 mg) in water (0.5 mL) to basic conditions. The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10um; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 30%-60%, 10min) to give **K101-C1329** (2.00 mg, 3.62 μmol, 19.92% yield, 94.4% purity) and **K101-C1329-Cl** (2.70 mg, 4.47 μmol, 24.60% yield, 92.4% purity), both as white solids after lyophilization.

[0366] K101-C1329 MS (m/z): 544.1 [M+Na]⁺


[0367] K101-C1329 ¹H NMR (400MHz, CD₃OD) δ 7.53 (s, 1H), 7.26-7.20 (m, 2H), 7.20-7.14 (m, 2H), 5.63-5.57 (m, 1H), 4.22 (d, *J*=6.0 Hz, 1H), 3.99-3.89 (m, 2H), 3.23-3.13 (m, 3H), 3.10-2.97 (m, 4H), 2.57-2.47 (m, 1H), 2.45-2.36 (m, 1H), 2.05-1.93 (m, 2H), 1.75 (dd, *J*=1.3, 2.8 Hz, 3H), 1.34-1.23 (m, 1H), 1.18 (s, 3H), 1.06 (s, 3H), 0.95 (d, *J*=6.0 Hz, 1H), 0.86 (d, *J*=6.0 Hz, 3H).

[0368] K101-C1329-Cl LC-MS (m/z): 562.1 [M+Na]⁺

[0369] K101-C1329-Cl ¹H NMR (400MHz, CD₃OD) δ 7.52 (s, 1H), 7.26-7.20 (m, 2H), 7.19-7.14 (m, 2H), 5.81-5.76 (m, 1H), 4.22 (d, *J*=5.8 Hz, 1H), 4.17-4.02 (m, 2H), 3.22-3.11 (m, 3H), 3.08-2.97 (m, 4H), 2.68-2.41 (m, 2H), 2.04-1.93 (m, 2H), 1.78-1.73 (m, 3H), 1.30-1.22 (m, 1H), 1.22-1.18 (m, 3H), 1.10-1.04 (m, 3H), 0.98-0.93 (m, 1H), 0.85 (d, *J*=6.3 Hz, 3H).

Example 27: Synthesis Scheme of **K101-C1330**.

[0370] The scheme for synthesis of compound **K101-C1330** is illustrated below.



[0371] Preparation of Compound K101-C1330-A. To a solution of **K101-C20Tr-B** (20.00 mg, 33.86 μmol , 1.00 eq) and **C13-30** (46.57 mg, 203.16 μmol , 6.00 eq) in DCM (2.00 mL) were added EDCI (38.94 mg, 203.16 μmol , 6.00 eq) and DMAP (24.82 mg, 203.16 μmol , 6.00 eq). The reaction solution was stirred at 25 °C for 16 hours to give a brown solution. LC-MS showed the reaction was complete. The reaction solution was diluted with DCM (10 mL), then washed with water (3 mL), 0.5 M HCl (2 mL), brine (2 mL), and dried over anhydrous Na_2SO_4 . The product was filtered and concentrated under reduced pressure to give the crude product. The product was purified by prep-TLC (PE/EtOAc=3/1, SiO_2) to give 16.2 mg of **K101-C1330-A** as a colorless gum.


[0372] Preparation of Compound K101-C1330. To a solution of **K101-C1330-A** (10.00 mg, 12.47 μmol , 1.00 eq) in dioxane (400.00 μL) was added HCl/dioxane (4 M, 200.17 μL , 64.21 eq). The reaction mixture was stirred at 25 °C for 2 hours to give a light brown solution. LC-MS showed the reaction was not complete so 0.2 mL of HCl/dioxane (4 M) was added and the reaction solution stirred at 25 °C for an additional 1 hour. LC-MS showed the reaction was almost complete. The reaction solution was combined with another preparation of **K101-C1330-A** and concentrated under reduced pressure. The residue was diluted with CH_3CN (1 mL) and water (1 mL) and the solution adjusted to pH 8 with addition of solid K_2CO_3 (5 mg). The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1% TFA)-B: ACN]; B%: 25%-55%, 10min) to give **K101-C1330** (4.70 mg, 65.05% yield, 99.0% purity, TFA salt) as a white powder after lyophilization.

[0373] MS (m/z): 482.1 $[\text{M}+\text{Na}]^+$

[0374] ^1H NMR (400MHz, CD_3OD) δ 7.55 (s, 1H), 5.64-5.59 (m, 1H), 4.12 (dd, $J=5.3, 7.8$ Hz, 1H), 4.00-3.90 (m, 2H), 3.19-3.14 (m, 1H), 3.09-3.03 (m, 1H), 2.58-2.48 (m, 1H), 2.45-2.37 (m, 1H), 2.25 (dd, $J=6.8, 14.6$ Hz, 1H), 2.12-2.01 (m, 1H), 1.95-1.85 (m, 1H), 1.80-1.70 (m, 4H), 1.61 (dd, $J=10.4, 14.7$ Hz, 1H), 1.18 (s, 3H), 1.09 (s, 3H), 1.02-0.97 (m, 1H), 0.93 (d, $J=6.8$ Hz, 3H), 0.8-0.76 (m, 1H), 0.66-0.59 (m, 2H), 0.27-0.17 (m, 2H).

Example 28: Synthesis Scheme of **K101-C1331**.

[0375] The scheme for synthesis of compound **K101-C1331** is illustrated below.



[0376] Preparation of Compound K101-C1331-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) in DCM (2.00 mL) were added (E,2S)-2-(tert-butoxycarbonylamino)-5-phenylpent-4-enoic acid (**C13-31**) (29.59 mg, 101.56 μmol , 2.00 eq), DMAP (24.82 mg, 203.12 μmol , 4.00 eq), HOBt (13.72 mg, 101.56 μmol , 2.00 eq) and EDC (19.47 mg, 101.56 μmol , 2.00 eq). The mixture was stirred at 20°C for 12h to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The mixture was combined with a second preparation of **K101-C1331-A** and the mixture quenched with H₂O (10 mL) and then extracted with DCM (15 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1331-A** (32.00 mg, 37.03 μmol , 62.49% yield) as a white solid.


[0377] Preparation of Compound K101-C1331. To a solution of **K101-C1331-A** (32.00 mg, 37.03 μmol , 1.00 eq) in THF (2.00 mL) were added TFA (770.00 mg, 6.75 mmol, 500.00 μL , 182.37 eq) and Et₃SiH (8.61 mg, 74.06 μmol , 11.79 μL , 2.00 eq). The mixture was stirred at 20°C for 12h to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated by N₂, and the product purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 27%-57%, 10min) to give **K101-C1331** (10.00 mg, 15.73 μmol , 42.48% yield, 100% purity, TFA salt) as a white solid.

[0378] LC-MS (m/z): 524.2 [M+Na]⁺

[0379] ¹H NMR (400MHz, CD₃OD) δ 7.57 (s, 1H), 5.64 (s, 1H), 4.10-4.07 (m, 1H), 4.00-3.93 (s, 2H), 3.18 (s, 1H), 3.08 (s, 1H), 2.53-2.45 (m, 1H), 2.45-2.41 (m, 1H), 2.29-2.27 (m, 1H), 2.05 (m, 1H), 1.84-1.29 (m, 16H), 1.19 (s, 3H), 1.11 (s, 3H), 1.02-1.01 (m, 2H), 0.96-0.94 (m, 3H).

Example 29: Synthesis Scheme of **K101-C1332**.

[0380] The scheme for synthesis of compound **K101-C1332** is illustrated below.



[0381] Preparation of Compound K101-C1332-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) in DCM (5.00 mL) were added (2S)-2-(tert-butoxycarbonylamino)-3-cyclopentyl-propanoic acid (C13-32) (26.14 mg, 101.56 μmol , 2.00 *eq*), DMAP (24.82 mg, 203.12 μmol , 4.00 *eq*) and EDC (19.47 mg, 101.56 μmol , 2.00 *eq*). The mixture was stirred at 20°C for 5h to give a yellow solution. LC-MS showed the reaction was incomplete. Additional EDC (10mg) was added, and mixture stirred at 20°C for 14hr. LC-MS showed some of the **K101-C20Tr-B** still remained. A further amount of EDC (11 mg) was added, and the mixture stirred at 20°C for another 5hr. LC-MS and TLC showed the reaction was complete. The reaction mixture was quenched with H₂O (15 mL) and extracted with DCM (15 mL x 5). The organic layers were dried over Na₂SO₄ and concentrated to give a yellow solid. The product was purified by prep-TLC (eluting with Petroleum ether: Ethyl acetate = 7/2) to give **K101-C1332-A** (23.00 mg, 27.71 μmol , 54.57% yield) as a colorless solid.

[0382] Preparation of Compound K101-C1332. To a solution of **K101-C1332-A** (23.00 mg, 27.71 μmol , 1.00 *eq*) in THF (2.00 mL) were added Et₃SiH (3.22 mg, 27.71 μmol , 4.41 μL , 1.00 *eq*) and TFA (770.00 mg, 6.75 mmol, 500.00 μL , 243.71 *eq*). The mixture was stirred at 20°C for 1.5h to give a colorless solution. The reaction mixture was concentrated to give a yellow oil, which was dissolved in DCM (2 mL) followed by addition of TFA (0.5 mL). The mixture was stirred at 20°C for 1hr and concentrated to give a yellow oil. LC-MS showed the reaction was complete. The reaction mixture was concentrated, and the product purified by prep-HPLC (column: Phenomenex Gemini 150 x 25 mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 30%-60%, 10min). The separated layers were lyophilized to give **K101-C1332** (8.00 mg, 13.03 μmol , 47.03% yield, 98% purity, TFA salt) as a white solid.


[0383] LC-MS (m/z): 510.2 [M+Na]⁺

[0384] ¹H NMR (400MHz, CDCl₃) δ = 7.57 (s, 1H), 5.64 (d, *J*=4.5 Hz, 1H), 4.02 - 3.99 (m, 1H), 3.97 (s, 2H), 3.18 (s, 1H), 3.15-3.02 (m, 1H), 2.60 - 2.36 (m, 2H), 2.26 (dd, *J*=6.9, 14.7 Hz, 1H), 2.05 (dd, *J*=6.9, 13.2 Hz, 1H), 2.02 - 1.97 (m, 1H), 1.94-1.83 (m, 2H), 1.83 (t, *J*=7.8 Hz, 1H), 1.77 (d, *J*=1.5

Hz, 4H), 1.74 - 1.54 (m, 5H), 1.35 - 1.14 (m, 6H), 1.11 (s, 3H), 1.02 (d, $J=6.0$ Hz, 1H), 0.95 (d, $J=6.5$ Hz, 3H)

Example 30: Synthesis Scheme of K101-C1333.

[0385] The scheme for synthesis of compound **K101-C1333** is illustrated below.



[0386] Preparation of Compound **K101-C1333-A**: To a solution of **K101-C20Tr-B** (22.00 mg, 37.24 μ mol, 1.17 *eq*) and **DHA** (10.50 mg, 31.96 μ mol, 1.00 *eq*) in DCM (500.00 uL) was added EDCI (36.77 mg, 191.79 μ mol, 6.00 *eq*) and DMAP (23.43 mg, 191.79 μ mol, 6.00 *eq*). The reaction solution was stirred at 20 °C for 16 hours to give a brown solution. LCMS showed the desired MS value. The reaction mixture was combined with reaction mixture of ES5329-184 (5 mg of **K101-C20Tr-B** was used in this batch) and concentrated under reduced pressure. The residue was purified by prep-TLC (PE/EtOAc=5/1) to give **K101-C1333-A** (7.00 mg, 7.77 μ mol, 24.30% yield) as a colorless oil.

[0387] ¹H NMR (400MHz, CDCl₃) δ 7.58 (s, 1H), 7.46-7.38 (m, 6H), 7.31-7.26 (m, 6H), 7.24-7.18 (m, 3H), 5.60 (d, $J=3.5$ Hz, 1H), 5.42-5.27 (m, 12H), 3.51 (s, 2H), 3.31-3.25 (m, 1H), 2.96-2.90 (m, 1H), 2.90-2.67 (m, 10H), 2.57-2.47 (m, 1H), 2.44-2.30 (m, 5H), 2.13-1.91 (m, 7H), 1.77 (dd, $J=1.1, 2.9$ Hz, 3H), 1.59-1.51 (m, 1H), 1.19 (s, 3H), 1.07 (s, 3H), 0.97 (t, $J=7.5$ Hz, 3H), 0.87 (d, $J=6.3$ Hz, 3H), 0.79 (d, $J=5.3$ Hz, 1H).

[0388] Preparation of Compound **K101-C1333**: To a solution of **K101-C1333-A** (6.00 mg, 6.66 μ mol, 1.00 *eq*) in MeOH (200.00 uL) was added HClO₄ (9.96 mg, 99.17 μ mol, 6.00 uL, 14.89 *eq*) and Et₃SiH (774.15 ug, 6.66 μ mol, 1.06 uL, 1.00 *eq*). Then the reaction mixture was stirred at 0 °C for 1.5 hour to give a white suspension. TLC (DCM/MeOH=20/1, SiO₂) showed no starting material remained and a new spot was observed. The reaction mixture was concentrated under reduced pressure. The residue was purified by prep-TLC (PE/EtOAc=1/1) to give **K101-C1333** (3.65 mg, 79.85% yield, 96.0% purity) as a white solid after lyophilization.


[0389] MS (m/z): 681.3 [M+Na]⁺

[0390] ¹H NMR (400MHz, CD₃OD) δ 7.56-7.53 (m, 1H), 5.62-5.58 (m, 1H), 5.51-5.20 (m, 12H), 3.99-3.88 (m, 2H), 3.20-3.14 (m, 1H), 3.09-3.03 (m, 1H), 2.92-2.80 (m, 10H), 2.56-2.48 (m, 1H), 2.47-2.38 (m, 5H), 2.15-2.06 (m, 3H), 2.06-1.99 (m, 1H), 1.74 (dd, $J=1.3, 2.9$ Hz, 3H), 1.55 (dd,

$J=10.7, 14.4$ Hz, 1H), 1.18 (s, 3H), 1.07 (s, 3H), 0.97 (t, $J=7.5$ Hz, 3H), 0.90 (d, $J=6.4$ Hz, 4H), 0.86 (d, $J=5.5$ Hz, 1H).

Example 31: Synthesis Scheme of K101-C1334.

[0391] The scheme for synthesis of compound **K101-C1334** is illustrated below.



[0392] Preparation of Compound **K101-C1334-A**: To a solution of **K101-C20Tr-B** (35.00 mg, 59.25 μ mol, 1.00 *eq*) and **C13-34** (136.45 mg, 592.47 μ mol, 10.00 *eq*) in DCM (1.00 mL) was added EDCI (34.07 mg, 177.74 μ mol, 3.00 *eq*) and DMAP (21.71 mg, 177.74 μ mol, 3.00 *eq*), then the mixture was stirred at 20°C for 14 hours to give colorless solution. The reaction was complete detected by LCMS (ES6477-17-P1B). The reaction solution was diluted with H₂O (10 ml), extracted with DCM: MeOH = 10:1 (10 ml * 5). The combined organic layers were dried over anhydrous Na₂SO₄, filtered, concentrated under reduced pressure to give crude product. The crude product purified by prep-TLC (SiO₂, PE:EA = 2:1) to give **K101-C1334-A** (26.40 mg, 32.88 μ mol, 55.49% yield) as a white solid.

[0393] **K101-C1334-A** ¹H NMR (400MHz, CDCl₃) δ 7.59 (s, 1H), 7.44-7.39 (m, 6H), 7.31-7.27 (m, 5H), 7.24-7.18 (m, 3H), 5.61 (s, 1H), 5.30 (s, 1H), 3.50 (s, 2H), 3.27 (s, 1H), 2.97-2.86 (m, 3H), 2.58-2.47 (m, 1H), 2.38 (d, $J=18.8$ Hz, 3H), 2.28 (t, $J=7.4$ Hz, 4H), 2.08-1.90 (m, 3H), 1.76 (s, 3H), 1.60-1.49 (m, 3H), 1.25 (s, 16H), 1.20-1.16 (m, 1H), 1.18 (s, 3H), 1.06 (s, 3H), 0.92-0.75 (m, 5H).

[0394] Preparation of Compound **K101-C1334**: To a solution of **K101-C1334-A** (26.00 mg, 32.38 μ mol, 1.00 *eq*) in THF (3.00 mL) was added TFA (770.00 mg, 6.75 mmol, 500.00 μ L, 208.56 *eq*). Then the solution was stirred at 0°C for 18 hours to give colorless solution. The reaction was complete detected by LCMS (ES6477-18-P1B). The reaction was concentrated under ordinary pressure to give yellow oil. The residue was purified by prep-HPLC (column: Phenomenex Gemini 150*25mm*10 μ m; mobile phase: [water (0.1%TFA)-ACN]; B%: 50%-80%, 10 min). The separated layers were lyophilized to give **K101-C1334** (3.00 mg, 5.35 μ mol, 16.52% yield, 97.28% purity, Free) as a white solid.


[0395] **K101-C1334**: LC-MS (m/z): 583.1 [M+Na]⁺

[0396] ¹H NMR (400MHz, MeOD) δ 7.56 (s, 1H), 5.63-5.58 (m, 1H), 3.99-3.89 (m, 2H), 3.20-3.15 (m, 1H), 3.09-3.04 (m, 1H), 2.56-2.49 (m, 1H), 2.47-2.40 (m, 1H), 2.37-2.26 (m, 4H), 2.22-2.10 (m,

1H), 2.10-1.99 (m, 2H), 1.77-1.73 (m, 3H), 1.66-1.51 (m, 5H), 1.32 (s, 12H), 1.18 (s, 3H), 1.07 (s, 3H), 0.91 (d, $J=6.3$ Hz, 3H), 0.86 (d, $J=5.5$ Hz, 1H).

Example 32: Synthesis Scheme of K101-C1335.

[0397] The scheme for synthesis of compound **K101-C1335** is illustrated below.



[0398] Preparation of Compound **K101-C1335-A**: To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) in DCM (2.00 mL) were added tridecanedioic acid (62.04 mg, 253.90 μmol , 5.00 *eq*), DMAP (37.22 mg, 304.68 μmol , 6.00 *eq*) and EDCI (58.41 mg, 304.68 μmol , 6.00 *eq*). The mixture was stirred at 20°C for 14hr to give a colorless solution. The mixture was stirred at 20°C for 5hr to give a colorless solution. LCMS and TLC showed the reaction was completed. The reaction mixture was quenched with H₂O (15 mL) and extracted with DCM (15 mL * 5). The organic layers were dried over Na₂SO₄ and concentrated to give a yellow solid. The yellow solid was purified by prep-TLC (eluting with Petroleum ether: Ethyl acetate = 3/1) to give **K101-C1335-A** (24.00 mg, 29.37 μmol , 57.84% yield) as a white solid.

[0399] ¹H NMR (400MHz, CDCl₃) δ = 7.65-7.55 (s, 1H), 7.45-7.35 (m, 6H), 7.32 - 7.27 (m, 6H), 7.25 - 7.19 (m, 3H), 5.65-5.55 (m, 1H), 3.55-3.45 (m, 2H), 3.30-3.25 (m, 1H), 3.00-2.90 (m, 1H), 2.59 - 2.22 (m, 6H), 2.20 (s, 3H), 2.13 - 1.90 (m, 4H), 1.80-1.72 (m, 3H), 1.45-1.20 (m, 14H), 1.19 (s, 3H), 1.07 (s, 3H), 0.88 (d, $J=6.3$ Hz, 3H), 0.79 (d, $J=5.3$ Hz, 1H).


[0400] Preparation of Compound **K101-C1335**: To a solution of **K101-C1335-A** (24.00 mg, 29.37 μmol , 1.00 *eq*) in THF (10.00 mL) was added TFA (770.00 mg, 6.75 mmol, 500.00 μL , 229.94 *eq*). The mixture was stirred at 0°C for 14hr to give a yellow solution. LCMS showed **K101-C1335-A** was remained, then added TFA (0.1 mL). The mixture was stirred at 0°C for 14hr to give a colorless solution. LCMS showed the reaction was completed. The reaction mixture was concentrated to give yellow oil. The yellow oil was purified by prep-HPLC (column: Phenomenex Gemini 150*25mm*10um; mobile phase: [water (0.1%TFA)-ACN]; B%: 70%-70%, 10min). The separated layers were lyophilized to give **K101-C1335** (5.40 mg, 9.15 μmol , 31.16% yield, 97.4% purity, Free) as a white solid.

[0401] LC-MS (m/z): 597.3 [M+Na]⁺

[0402] ¹H NMR (400MHz, MeOD) δ = 7.60-7.50 (m, 1H), 5.62 (m, 1H), 4.03 - 3.84 (m, 2H), 3.25-3.14 (m, 1H), 3.12 - 3.01 (m, 1H), 2.61 - 2.41 (m, 2H), 2.41 - 2.23 (m, 4H), 2.19 - 1.99 (m, 2H), 1.83 - 1.71 (m, 3H), 1.71 - 1.48 (m, 5H), 1.45-1.35 (m, 14H), 1.19 (s, 3H), 1.09 (s, 3H), 0.93 (d, *J*=6.3 Hz, 3H), 0.87 (d, *J*=5.8 Hz, 1H)

Example 33: Synthesis Scheme of **K101-C1336**.

[0403] The scheme for synthesis of compound **K101-C1336** is illustrated below.



[0404] Preparation of Compound **K101-C1336-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) in DCM (5.00 mL) were added (2R)-2-(tert-butoxycarbonylamino)-3-phenylpropanoic acid (**C13-36**) (26.95 mg, 101.56 μmol , 2.00 *eq*), DMAP (30.00 mg, 245.78 μmol , 4.84 *eq*) and EDC (19.96 mg, 104.10 μmol , 2.05 *eq*). The mixture was stirred at 20°C for 19h to give a colorless solution. LC-MS and TLC showed the reaction was complete. The reaction mixture was quenched with H₂O (15 mL) and extracted with DCM (15 mL x 5). The organic layers were dried over Na₂SO₄ and concentrated to give a yellow solid. The product was purified by prep-TLC (eluting with Petroleum ether: Ethyl acetate = 3/1) to give **K101-C1336-A** (16.00 mg, 19.09 μmol , 37.60% yield) as a colorless solid.


[0405] Preparation of Compound K101-C1336. To a solution of **K101-C1336-A** 16.00 mg, 19.09 μmol , 1.00 eq) in DCM (2.00 mL) were added TFA (770.00 mg, 6.75 mmol, 500.00 uL, 353.76 eq) and Et_3SiH (2.22 mg, 19.09 μmol , 3.04 uL, 1.00 eq). The mixture was stirred at 20°C for 5h to give a colorless solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated and the product purified by prep-HPLC column: Phenomenex Gemini 150 x 25mm x 10um; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 15%-45%, 10min). The separated layers were lyophilized to give **K101-C1336** (6.00 mg, 9.84 μmol , 51.55% yield, 100% purity, TFA salt) as a white solid.

[0406] LC-MS (m/z): 518.2 [M+Na]⁺

[0407] ¹H NMR (400MHz, MeOD) δ = 7.59 (s, 1H), 7.48 - 7.29 (m, 5H), 5.63 (s, 1H), 4.37 (dd, $J=5.9, 8.2$ Hz, 1H), 4.04 - 3.89 (m, 2H), 3.42 - 3.37 (m, 1H), 3.25 - 3.00 (m, 3H), 2.61 - 2.35 (m, 2H), 2.23 (dd, $J=7.0, 15.1$ Hz, 1H), 2.05 (s, 1H), 1.78 (d, $J=1.5$ Hz, 3H), 1.64 (dd, $J=10.4, 14.7$ Hz, 2H), 1.08 (d, $J=7.8$ Hz, 6H), 1.01 - 0.90 (m, 4H)

Example 34: Synthesis Scheme of **K101-C1337**.

[0408] The scheme for synthesis of compound **K101-C1337** is illustrated below.



[0409] Preparation of Compound K101-C1337-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) in DCM (2.00 mL) were added (2R)-2-(tert-butoxycarbonylamino)-3-cyclohexylpropanoic acid (C13-37) (27.56 mg, 101.56 μmol , 2.00 eq), DMAP (24.82 mg, 203.12 μmol , 4.00 eq), HOBt (13.72 mg, 101.56 μmol , 2.00 eq) and EDC (19.47 mg, 101.56 μmol , 2.00 eq). The mixture was stirred at 20°C for 12 h to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The mixture was combined with a second preparation of **K101-C1337-A**, the mixture quenched with H_2O (10 mL) and then extracted with DCM (20mL x 3). The organic layers were dried over Na_2SO_4 and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1337-A** (18.00 mg, 21.32 μmol , 41.99% yield) as a white solid.


[0410] Preparation of Compound K101-C1337. To a solution of **K101-C1337-A** (18.00 mg, 21.32 μmol , 1.00 eq) in THF (2.00 mL) were added TFA (770.00 mg, 6.75 mmol, 500.00 μL , 316.75 eq) and Et_3SiH (7.44 mg, 63.96 μmol , 10.19 μL , 3.00 eq). The mixture was stirred at 20°C for 2h to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated by N_2 , and the product purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 25%-55%, 10min) to give **K101-C1337** (5.20 mg, 10.37 μmol , 48.62% yield, 100% purity) as a white solid.

[0411] LC-MS (m/z): 524.3 $[\text{M}+\text{Na}]^+$

[0412] ^1H NMR (400MHz, CD_3OD) δ 7.58 (s, 1H), 5.64-5.63 (s, 1H), 4.12-4.09 (m, 1H), 4.00-3.94 (m, 2H), 3.18 (s, 1H), 3.08 (s, 1H), 2.53-2.45 (m, 1H), 2.40-2.32 (m, 1H), 2.26-2.24 (m, 1H), 2.05-2.03 (m, 1H), 1.84-1.64 (m, 12H), 1.33-1.31 (m, 3H), 1.20 (s, 3H), 1.11 (s, 3H), 1.01-0.95 (m, 6H).

Example 35: Synthesis Scheme of **K101-C1338**.

[0413] The scheme for synthesis of compound **K101-C1338** is illustrated below.



[0414] Preparation of compound K101-C1338-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) in DCM (2.00 mL) were added (2S)-2-(tert-butoxycarbonylamino)-4-methylsulfanylbutanoic acid (**C13-38**) (25.32 mg, 101.56 μmol , 2.00 eq), DMAP (24.82 mg, 203.12 μmol , 4.00 eq), HOBt (13.72 mg, 101.56 μmol , 2.00 eq) and EDC (19.47 mg, 101.56 μmol , 2.00 eq). The mixture was stirred at 20°C for 12h to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The mixture was combined with as second preparation of **K101-C1338-A**, and the mixture quenched with H_2O (10 mL) and extracted with DCM (15 mL x 3). The organic layers were dried over Na_2SO_4 and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1338-A** (36.00 mg, 43.79 μmol , 64.84% yield) as a white solid.

[0415] Preparation of compound K101-C1338. To a solution of **K101-C1338-A** (36.00 mg, 43.79 μmol , 1.00 eq) in THF (2.00 mL) was added TFA (1.54 g, 13.51 mmol, 1.00 mL, 308.44 eq). The mixture was stirred at 20°C for 12h to give a yellow solution. LC-MS showed the reaction was


complete. The reaction mixture was concentrated by N_2 and the product purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10um; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 20%-50%, 10min) to give **K101-C1338** (13.20 mg, 20.95 μ mol, 47.83% yield, 94.2% purity, TFA salt) as a white solid.

[0416] LC-MS (m/z): 502.1 [M+Na]⁺

[0417] ¹H NMR (400MHz, CD₃OD) δ 7.57 (s, 1H), 5.65-5.64 (m, 1H), 4.26-4.23 (m, 1H), 4.00-3.94 (m, 2H), 3.18 (s, 1H), 3.08 (s, 1H), 2.72-2.69 (m, 2H), 2.53-2.45 (m, 1H), 2.45-2.41 (m, 1H), 2.30-2.24 (m, 2H), 2.15-2.11 (m, 5H), 1.77 (s, 3H), 1.76-1.75 (m, 1H), 1.20 (s, 3H), 1.03 (s, 3H), 0.96-0.94 (m, 4H).

Example 36: Synthesis Scheme of **K101-C1339**.

[0418] The scheme for synthesis of compound **K101-C1339** is illustrated below.



[0419] Preparation of Compound **K101-C1339-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μ mol, 1.00 eq) in DCM (2.00 mL) were added (2R)-2-((tert-butoxycarbonylamino)-3-methylsulfanyl)propanoic acid (**C13-39**) (23.90 mg, 101.56 μ mol, 2.00 eq), DMAP (24.82 mg, 203.12 μ mol, 4.00 eq), HOBt (13.72 mg, 101.56 μ mol, 2.00 eq) and EDC (19.47 mg, 101.56 μ mol, 2.00 eq). The mixture was stirred at 20°C for 12hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The mixture was combined with a second preparation, which was quenched with H₂O (10 mL) and then extracted with DCM (15 mL). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1339-A** (35.00 mg, 43.32 μ mol, 85.30% yield) as a white solid.

[0420] Preparation of Compound **K101-C1339**. To a solution of **K101-C1339-A** (35.00 mg, 43.32 μ mol, 1.00 eq) in THF (2.00 mL) was added TFA (4.94 mg, 43.32 μ mol, 3.21 μ L, 1.00 eq). The mixture was stirred at 20°C for 3h to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated by N_2 and the product purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10um; mobile phase: [A: water (0.1%TFA)-B: ACN];


B%: 18%-48%, 10min) to give **K101-C1339** (4.40 mg, 7.11 μmol , 16.41% yield, 93.63% purity, TFA salt) as a white solid.

[0421] LC-MS (m/z): 488.1 [M+Na]⁺

[0422] ¹H NMR (400MHz, CD₃OD) δ 7.57 (s, 1H), 5.65-5.64 (m, 1H), 4.33-4.30 (m, 1H), 4.00-3.97 (m, 2H), 3.19-3.14 (m 2H), 3.07 (s, 1H), 3.00-2.97 (m, 1H), 2.53-2.45 (m, 1H), 2.40-2.32 (m, 1H), 2.25-2.23 (m, 4H), 2.10-2.05 (m, 1H), 1.77 (s, 3H), 1.20 (s, 3H), 1.11 (s, 3H), 1.05-1.03 (m, 1H), 0.96-0.94 (m, 3H).

Example 37: Synthesis Scheme of **K101-C1340**.

[0423] The scheme for synthesis of compound **K101-C1340** is illustrated below.



[0424] Preparation of compound **K101-C1340-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) and **C13-40** in DCM (3.00 mL) were added EDC (19.47 mg, 101.56 μmol , 2.00 *eq*) and DMAP (37.22 mg, 304.68 μmol , 6.00 *eq*). The reaction solution was stirred at 20 °C for 14 hours to give colorless solution. TLC (PE/EtOAc=1/1, SiO₂) and LC-MS showed the reaction was complete. The reaction solution was combined with a second preparation, and then diluted with H₂O (5 ml x 2) followed by extraction with DCM (10 ml x 5). The combined organic layers were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to give crude product as a pale yellow solid. The product was purified by prep-TLC (PE/EtOAc=1/1, SiO₂) to give **K101-C1340-A** (20.30 mg, 23.15 μmol , 39.29% yield) as a white solid.


[0425] Preparation of Compound **K101-C1340**. To a solution of **K101-C1340-A** (20.00 mg, 22.80 μmol , 1.00 *eq*) in THF (2.00 mL) were added sequentially TFA (770.00 mg, 6.75 mmol, 500.00 μL , 296.19 *eq*) followed by Et₃SiH (2.65 mg, 22.80 μmol , 3.63 μL , 1.00 *eq*). The mixture was stirred at 20°C for 18 hours to give a pale yellow solution. The reaction was complete detected by LC-MS. The reaction was concentrated under reduced pressure to give a yellow solid, and the product purified by prep-HPLC (column: Phenomenex Gemini 150 x 25 mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 55%-95%, 10 min). The separated layers were lyophilized to give **K101-C1340** (3.80 mg, 7.11 μmol , 31.17% yield, 96.78% purity, TFA salt) as a pale yellow solid.

[0426] LC-MS (m/z): 557.1 [M+Na]⁺

[0427] ^1H NMR (400MHz, MeOD) δ 7.58 (d, $J=7.5$ Hz, 1H), 7.54-7.51 (m, 1H), 7.38 (d, $J=8.0$ Hz, 1H), 7.19 (s, 1H), 7.17-7.12 (m, 1H), 7.10 -7.05 (m, 1H), 5.59-5.54 (m, 1H), 4.58 (s, 2H), 4.17-4.09 (m, 1H), 3.99-3.90 (m, 2H), 3.17-3.11 (m, 1H), 3.02-2.96 (m, 1H), 2.55-2.46 (m, 1H), 2.45-2.36 (m, 1H), 2.04-1.92 (m, 2H), 1.78-1.71 (m, 3H), 1.43-1.32 (m, 1H), 1.04-0.97 (m, 6H), 0.83 (d, $J=6.0$ Hz, 3H), 0.78-0.74 (m, 1H).

Example 38: Synthesis Scheme of **K101-C1341**.

[0428] The scheme for synthesis of compound **K101-C1341** is illustrated below.



[0429] Preparation of Compound **K101-C1341-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) in DCM (1.00 mL) were added (*2R*)-2-(tert-butoxycarbonylamino)-3-(1H-indol-3-yl)propanoic acid (**C13-41**) (154.55 mg, 507.83 μmol , 10.00 *eq*), EDC (19.47 mg, 101.57 μmol , 2.00 *eq*) and DMAP (37.22 mg, 304.70 μmol , 6.00 *eq*). The mixture was stirred at 20°C for 5hr to give a colorless solution. LC-MS and TLC showed the reaction was complete. The reaction mixture was quenched with H₂O (15 mL) and extracted with DCM (15 mL x 5). The organic layers were dried over Na₂SO₄ and concentrated to give a yellow solid. The product was purified by prep-TLC (eluting with Petroleum ether: Ethyl acetate = 2/1) to give **K101-C1341-A** (35.00 mg, 39.91 μmol , 67.35% yield) as a white solid.

[0430] Preparation of Compound **K101-C1341**. To a solution of **K101-C1341-A** (35.00 mg, 39.91 μmol , 1.00 *eq*) in THF (2.00 mL) were added TFA (770.00 mg, 6.75 mmol, 500.00 μL , 169.21 *eq*) and Et₃SiH (4.64 mg, 39.91 μmol , 6.36 μL , 1.00 *eq*). The mixture was stirred at 20°C for 5hr to give a colorless solution. The reaction mixture was concentrated and the resultant yellow oil dissolved with DCM (2 mL) followed by addition of TFA (0.5 mL). The mixture was stirred at 20°C for 2hr to give a yellow solution. The reaction mixture was concentrated and dissolved with DCM (2 mL) followed by addition of TFA (0.5 mL). This reaction mixture was stirred at 20°C for 1hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated to give a yellow oil, which was then dissolved in MeOH (4 mL) and stirred at 20°C for 14hr to give a yellow liquid. The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 20%-50%, 10min). The separated layers


were lyophilized to give **K101-C1341** (4.00 mg, 5.59 μmol , 14.01% yield, 90.7% purity, TFA salt) as a white solid.

[0431] LC-MS (m/z): 557.1 [M+Na]⁺

[0432] ¹H NMR (400MHz, MeOD) δ = 7.61 (d, J =7.5 Hz, 1H), 7.57 (s, 1H), 7.42 (d, J =8.3 Hz, 1H), 7.26 (s, 1H), 7.23 - 7.10 (m, 2H), 5.60-5.50 (m, 1H), 4.33 (t, J =7.4 Hz, 1H), 4.06 - 3.92 (m, 2H), 3.60 - 3.45 (m, 1H), 3.22 - 3.13 (m, 1H), 3.05-2.95 (m, 1H), 2.59 - 2.38 (m, 2H), 2.17 (dd, J =6.8, 14.6 Hz, 1H), 2.04 (d, J =8.8 Hz, 1H), 1.77 (d, J =1.5 Hz, 3H), 1.63 (dd, J =10.5, 14.8 Hz, 1H), 1.40-1.25(m, 2H), 1.04 (s, 3H), 0.93 (d, J =6.5 Hz, 3H), 0.87 (s, 3H), 0.69 (d, J =5.8 Hz, 1H)

Example 39: Synthesis Scheme of **K101-C1342**.

[0433] The scheme for synthesis of compound **K101-C1342** is illustrated below.



[0434] Preparation of Compound **K101-C1342-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) and **C13-42** (25.39 mg, 76.17 μmol , 1.50 *eq*) in DCM (1.00 mL) were added EDCI (58.41 mg, 304.68 μmol , 6.00 *eq*) and DMAP (18.61 mg, 152.35 μmol , 3.00 *eq*). The mixture was stirred at 20°C for 14h to give a yellow solution. The reaction was complete as detected by LC-MS. The reaction solution was combined with a second preparation, and diluted with H₂O (10 mL) followed by extraction with DCM (10 mL x 3). The combined organic layers were dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give a yellow solid. The product was purified by prep-TLC (PE/EtOAc=2/1, SiO₂) to give **K101-C1342-A** (50.30 mg, 55.52 μmol , 93.44% yield) as a white solid.


[0435] Preparation of Compound **K101-C1342**. To a solution of **K101-C1342-A** (50.00 mg, 55.19 μmol , 1.00 *eq*) in MeOH (500.00 μL) was added HCl/MeOH (4 M, 500.00 μL , 36.24 *eq*). The reaction was stirred at 20°C for 3.5 hours to give a pale yellow solution. LC-MS showed the reaction was almost complete. The solution was adjusted to pH 8 with saturated aqueous NaHCO₃ and then extracted with DCM (2 ml x 3). The combined organic layers were concentrated under reduced pressure to give a yellow solid. The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25 mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 25%-55%, 10 min) to give **K101-C1342** (10.70 mg, 18.99 μmol , 34.40% yield, 98.84% purity, TFA salt) as a white solid.

[0436] LC-MS (m/z): 586.1 [M+Na]⁺

[0437] ^1H NMR (400MHz, MeOD) δ 7.71 (d, $J=8.0$ Hz, 2H), 7.56-7.51 (m, 3H), 5.62-5.57 (m, 1H), 4.47-4.40 (m, 1H), 3.94 (s, 2H), 3.47-4.40 (m, 1H), 3.25-3.18 (m, 1H), 3.18-3.13 (m, 1H), 3.07-2.99 (m, 1H), 2.57-2.47 (m, 1H), 2.45-2.36 (m, 1H), 2.17 (dd, $J=7.2, 14.7$ Hz, 1H), 2.07-1.98 (m, 1H), 1.78-1.73 (m, 3H), 1.53-1.43 (m, 1H), 1.10 (s, 3H), 1.06 (s, 3H), 0.96 (d, $J=5.8$ Hz, 1H), 0.91 (d, $J=6.5$ Hz, 3H).

Example 40: Synthesis Scheme of **K101-C1343**.

[0438] The scheme for synthesis of compound **K101-C1343** is illustrated below.



[0439] Preparation of Compound **K101-C1343-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) and **C13-43** were added EDC (58.41 mg, 304.68 μmol , 6.00 *eq*) and DMAP (37.22 mg, 304.68 μmol , 6.00 *eq*). The reaction solution was stirred at 25 °C for 1 hour to give a brown solution. TLC (PE/EtOAc=1/1, SiO₂) showed the reaction was complete. The reaction solution was diluted with DCM (2 mL), washed with brine (1 mL), and dried over anhydrous Na₂SO₄. The product was filtered and concentrated under reduced pressure to give the crude product as a brown gum. The product was purified by prep-TLC (PE/EtOAc=1/1, SiO₂) to give **K101-C1343-A** (33.30 mg, 78.25% yield) as a colorless gum.


[0440] Preparation of Compound **K101-C1343**. To a solution of **K101-C1343-A** (25.00 mg, 29.83 μmol , 1.00 *eq*) in MeOH (1.00 mL) was added HCl/MeOH (4 M, 1 mL), and the reaction solution stirred at 20°C for 2.5 hours to give a clear solution. LC-MS showed the reaction was almost complete so the reaction solution was stirred at 20°C for an additional 1hr. The reaction solution was combined with a second preparation of the compound, and then cooled to 0°C. The solution was adjusted to pH 8 with saturated aqueous NaHCO₃. The solution was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10um; mobile phase: [A: water (0.05% HCl)-B: ACN]; B%: 15%-45%, 10min) to give **K101-C1343** (7.60 mg, 47.02% yield, 98.2% purity, HCl salt) as a white solid.

[0441] LC-MS (m/z): 518.1 [M+Na]⁺

[0442] $^1\text{H NMR}$ (400MHz, CD_3OD) δ 7.53 (s, 1H), 7.50-7.43 (m, 5H), 5.57 (d, $J=4.2$ Hz, 1H), 4.73 (t, $J=7.3$ Hz, 1H), 4.00-3.88 (m, 2H), 3.18-3.04 (m, 3H), 3.01 (t, $J=5.5$ Hz, 1H), 2.56-2.46 (m, 1H), 2.44-2.36 (m, 1H), 2.11-1.93 (m, 2H), 1.78-1.71 (m, 3H), 1.38 (dd, $J=10.1, 14.3$ Hz, 1H), 1.01 (s, 3H), 0.96 (s, 3H), 0.90-0.80 (m, 4H).

Example 41: Synthesis Scheme of **K101-C1344**.

[0443] The scheme for synthesis of compound **K101-C1344** is illustrated below.



[0444] Preparation of Compound **K101-C1344-A**. To a solution of **K101-C20Tr-B** (35.00 mg, 59.25 μmol , 1.00 eq) and **C13-44** (41.37 mg, 148.12 μmol , 2.50 eq) in DCM (1.00 mL) were added EDC (68.15 mg, 355.48 μmol , 6.00 eq) and DMAP (21.71 mg, 177.74 μmol , 3.00 eq). The mixture was stirred at 20°C for 18hr to give a yellow solution. The reaction was complete as detected by LC-MS. The reaction solution was diluted with H_2O (10 mL), extracted with DCM (10 mL x 3), and the combined organic layers dried over Na_2SO_4 . The solution was concentrated under reduced pressure and purified by prep-TLC (SiO_2 , PE: EA = 3:1). Concentration under reduced pressure yielded **K101-C1344-A** (43.60 mg, 51.17 μmol , 86.36% yield) as a white solid.

[0445] Preparation of Compound **K101-C1344**. To a solution of **K101-C1344-A** (35.00 mg, 41.08 μmol , 1.00eq) in MeOH (500.00 μL) was added HCl/MeOH (4 M, 583.29 μL , 56.80 eq). The solution was stirred at 20°C for 18hr to give a yellow solution. The reaction was complete as detected by LC-MS. The reaction solution was diluted with H_2O (10 mL) and extracted with DCM (10 mL x 3). The combined organic layers were dried over Na_2SO_4 and concentrated under reduced pressure to give a yellow solid. The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 20%-50%, 10min). The separated layers were lyophilized to give **K101-C1344** (5.00 mg, 8.02 μmol , 19.52% yield, TFA salt) as a white solid.


[0446] LC-MS (m/z): 532.3 [$\text{M}+\text{Na}$] $^+$

[0447] $^1\text{H NMR}$ (400MHz, CD_3OD) δ 7.53 (s, 1H), 7.40-7.37 (m, 2H), 7.34-7.33 (m, 1H), 7.29-7.27(m, 2H), 5.59 (s, 1H), 3.98 (s, 2H), 3.88-3.85 (m, 1H), 3.15 (s, 1H), 3.06-3.03 (m, 2H), 2.95-2.93

(m, 1H), 2.72-2.64 (m, 2H), 2.55-2.44 (m, 2H), 2.12-1.98 (m, 2H), 1.75 (s, 3H), 1.59-1.55 (m, 1H), 1.30-1.25 (m, 3H), 1.14 (s, 3H), 1.06 (s, 3H), 0.91-0.89 (m, 3H), 0.84-0.83 (m, 1H).

Example 42: Synthesis Scheme of **K101-C1345**.

[0448] The scheme for synthesis of compound **K101-C1345** is illustrated below.



[0449] Preparation of Compound **K101-C1345-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) and **C13-45** (36.23 mg, 126.96 μmol , 2.50 eq) in DCM (1.00 mL) were added EDC (48.68 mg, 253.91 μmol , 5.00 eq) and DMAP (18.61 mg, 152.35 μmol , 3.00 eq). The reaction mixture was stirred at 20°C for 18hr. The reaction was complete as detected by LC-MS. The reaction solution was diluted with H₂O (10 mL) and extracted with DCM (8 mL x 5). The combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure to give pale yellow solution. The product was purified by prep-TLC (SiO₂, PE: EA = 3:1) to give **K101-C1345-A** (26.30 mg, 30.65 μmol , 60.36% yield) as a white solid.


[0450] Preparation of Compound **K101-C1345**. To a solution of **K101-C1345-A** (25.00 mg, 29.13 μmol , 1.00 eq) in MeOH (500.00 μL) was added HCl/MeOH (4 M, 7.28 μL , 1.00 eq). The solution was stirred at 20°C for 18 hr to give a yellow solution. The reaction was complete as detected by LC-MS. The reaction solution was diluted with H₂O (10 mL), neutralized with NaHCO₃ aqueous solution and then extracted with DCM (8 mL x 5). The combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure to give a yellow solid. The residue was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN];B%: 25%-55%,10min), and the separated layers lyophilized to give **K101-C1345** (4.30 mg, 8.34 μmol , 28.63% yield) as a white solid.

[0451] LC-MS (m/z): 538.3 [M+Na]⁺

[0452] ¹H NMR (400MHz, CD₃OD) δ 7.56 (s, 1H), 5.62 (s, 1H), 5.49 (m, 1H), 4.02-3.95 (m, 3H), 3.16-3.06 (m, 2H), 2.56-2.44 (m, 2H), 2.39-2.22 (m, 1H), 2.04-1.94 (m, 3H), 1.76-1.57 (m, 9H), 1.30-1.25 (m, 7H), 1.18 (s, 3H), 1.01 (s, 3H), 0.99-0.93 (m, 5H).

Example 43: Synthesis Scheme of **K101-C1346**.

[0453] The scheme for synthesis of compound **K101-C1346** is illustrated below.



[0454] Preparation of Compound **K101-C1346-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) in DCM (1.00 mL) were added (2S)-2-(tert-butoxycarbonylamino)-3,3-dimethyl-butanoic acid (C13-46) (70.47 mg, 304.68 μmol , 6.00 *eq*) DMAP (43.43 mg, 355.46 μmol , 7.00 *eq*) and EDC (58.41 mg, 304.68 μmol , 6.00 *eq*). The mixture was stirred at 20°C for 5hr to give a yellow solution. TLC showed the reaction was complete. The reaction mixture was quenched with H₂O (15 mL) and extracted with DCM (15 mL x 5). The organic layers were dried over Na₂SO₄ and concentrated to give a yellow solid. The product was purified by prep-TLC (eluting with Petroleum ether: Ethyl acetate = 3/1) to give **K101-C1346-A** (28.00 mg, 34.83 μmol , 68.58% yield) as a colorless solid.


[0455] Preparation of Compound **K101-C1346**. To a solution of **K101-C1346-A** (28.00 mg, 34.83 μmol , 1.00 *eq*) in MeOH (500.00 μL) was added HCl/MeOH (4 M, 8.71 μL , 1.00 *eq*). The mixture was stirred at 20°C for 19hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was adjusted to pH 6 with saturated NaHCO₃ and the resultant product purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 20%-50%, 10min). The organic layers were lyophilized to give **K101-C1346** (7.80 mg, 13.55 μmol , 38.91% yield, 100% purity, TFA salt) as a white solid.

[0456] LC-MS (m/z): 584.2 [M+Na]⁺

[0457] ¹H NMR (400MHz, MeOD) δ = 7.60-7.54 (m, 1H), 5.70-5.60 (m, 1H), 4.04 - 3.92 (m, 2H), 3.86 (s, 1H), 3.33-3.20 (m, 1H), 3.20-3.12 (m, 1H), 2.60 - 2.38 (m, 2H), 2.26 (dd, *J*=7.0, 14.6 Hz, 1H), 2.19 - 2.02 (m, 1H), 1.77 (d, *J*=1.5 Hz, 3H), 1.60 (dd, *J*=10.8, 14.6 Hz, 1H), 1.26 (s, 3H), 1.17 (s, 9H), 1.13 (s, 3H), 1.02 (d, *J*=5.8 Hz, 1H), 0.95 (d, *J*=6.3 Hz, 3H).

Example 44: Synthesis Scheme of **K101-C1347**.

[0458] The scheme for synthesis of compound **K101-C1347** is illustrated below.



[0459] Preparation of Compound K101-C1347-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 *eq*) in DCM (2.00 mL) were added (2S)-2-(1-adamantyl)-2-(tert-butoxycarbonylamino) acetic acid (C13-47) (94.27 mg, 304.68 μmol , 6.00 *eq*), EDC (58.41 mg, 304.68 μmol , 6.00 *eq*) and DMAP (43.43 mg, 355.46 μmol , 7.00 *eq*). The mixture was stirred at 20°C for 14h to give a yellow solution. LC-MS and TLC showed the reaction was complete. The reaction mixture was quenched with H₂O (15 mL) and extracted with DCM (15 mL x 5). The organic layers were dried over Na₂SO₄ and concentrated to give a yellow solid. The product was purified by prep-TLC (eluting with Petroleum ether: Ethyl acetate = 3/1) to give **K101-C1347-A** (13.00 mg, 14.74 μmol , 24.80% yield) as a white solid.




[0460] Preparation of Compound K101-C1347. To a solution of **K101-C1347-A** (13.00 mg, 14.74 μmol , 1.00 *eq*) in THF (1.00 mL) were added TFA (500.55 mg, 4.39 mmol, 325.03 μL , 297.89 *eq*) and Et₃SiH (1.71 mg, 14.74 μmol , 2.35 μL , 1.00 *eq*). The mixture was stirred at 20°C for 3h to give a colorless solution. The reaction mixture was concentrated, dissolved with DCM (2 mL) and then followed by addition of TFA (0.5 mL). The mixture was stirred at 20°C for 2h to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated to give a yellow oil, which was dissolved with MeOH (2 mL) and stirred at 20°C for 14h to give a buff liquid. The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 25%-55%, 10min). The separated layers were lyophilized to give **K101-C1347** (6.50 mg, 9.94 μmol , 67.44% yield, 100% purity, TFA salt) as a white solid.

[0461] LC-MS (m/z): 562.3 [M+Na]⁺

[0462] ¹H NMR (400MHz, MeOD) δ = 7.62-7.55 (s, 1H), 5.70-5.60 (m, 1H), 4.05 - 3.90 (m, 2H), 3.75 - 3.56 (m, 1H), 3.25-3.15 (m, 1H), 3.16-3.08 (m, 1H), 2.62 - 2.38 (m, 2H), 2.27 (dd, *J*=6.9, 14.7 Hz, 1H), 2.15-2.06 (m, 4H), 1.91 - 1.53 (m, 16H), 1.27 (s, 3H), 1.13 (s, 3H), 1.03 (d, *J*=5.5 Hz, 1H), 0.96 (d, *J*=6.5 Hz, 3H)

Example 45: Synthesis Scheme of **K101-C1348**.

[0463] The scheme for synthesis of compound **K101-C1348** is illustrated below.

**Preparation of C13-48:**

[0464] Preparation of compound C13-48-B. LiAlH₄ (413.84 mg, 10.91 mmol, 1.50 eq) was suspended in THF (10.00 mL) at 0°C, then **C13-48-A** (1.50 g, 7.27 mmol, 1.00 eq) in THF (10.00 mL) was added dropwise at 0°C. The mixture was allowed to stir at 20°C for 4hr to give a brown solution. LC-MS showed the reaction was complete. Following quenching with H₂O (0.5 mL), aqueous NaOH (0.5 mL, 15%) and H₂O (1.5 mL) were added. The mixture was filtered on Celite and the filtrate was concentrated to give the 7-phenylheptan-1-ol (1.30 g, 6.76 mmol, 92.99% yield) as a yellow oil.

[0465] Preparation of compound C13-48-C. To a solution of oxalyl dichloride (1.72 g, 13.52 mmol, 1.18 mL, 2.00 eq) in DCM (20.00 mL) was added dropwise DMSO (2.64 g, 33.80 mmol, 2.64 mL, 5.00 eq) at -78°C. The mixture was stirred at -78°C for 0.5hr. Compound **C13-48-B** (1.30 g, 6.76 mmol, 1.00 eq) in DCM (10.00 mL) was added at -78°C. The mixture was stirred at -78°C for 1hr,

and the Et₃N (3.42 g, 33.80 mmol, 4.68 mL, 5.00 eq) was added dropwise at -78°C. The mixture was allowed to stir at 20°C for 2.5hr to give a yellow suspension. LC-MS and TLC (eluting with: PE/EtOAc=5/1) showed the reaction was complete. The reaction mixture was quenched with H₂O (30 mL) and extracted with DCM (50 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by column chromatography on silica gel (eluting with: PE/EtOAc=100%PE to 5/1) to give **C13-48-C** (1.10 g, 5.78 mmol, 85.52% yield) as a yellow oil.

[0466] Preparation of compound C13-48-E. To a solution of **C13-48-C** (1.05 g, 5.52 mmol, 1.00 eq) in EtOH (10.00 mL) were added trimethylsilyl cyanide (TMSCN) (547.46 mg, 5.52 mmol, 692.99 uL, 1.00 eq) and NH₃.H₂O (851.01 mg, 6.07 mmol, 935.18 uL, 25% purity, 1.10 eq). The mixture was stirred at 20°C for 7hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was quenched with H₂O (30 mL) and extracted with DCM (50mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give **C13-48-E** (1.10 g, crude) as a yellow oil.

[0467] Preparation of compound C13-48-D. To a solution of **C13-48-E** (1.10 g, 5.09 mmol, 1.00 eq) in EtOH (5.00 mL) was added NaOH (610.21 mg, 15.26 mmol, 3.00 eq). The mixture was stirred at 20°C for 2hr followed by the addition of H₂O (1.00 mL). The mixture was stirred at 90°C for 2hr to give a yellow solution. LC-MS showed the reaction was complete.

[0468] Preparation of Compound C13-48. Boc anhydride (Boc₂O) (2.23 g, 10.20 mmol, 2.34 mL, 2.00 eq) was added to the preparation of **C13-48-E**, and the mixture stirred at 20°C for 2hr to give a yellow suspension. LC-MS and TLC (eluting with: 100%EtOAc) showed the reaction was complete. The mixture was combined with a second preparation, and the combined mixture was diluted with H₂O (20 mL) followed by extraction with PE (20 mL x 3). The water layer was adjusted to pH 5 with HCl (1N) and extracted with EtOAc (30 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by column chromatography on silica gel (eluting with: PE/EtOAc=1/1 to 100%EtOAc) to give **C13-48** (700.00 mg, 2.09 mmol, 40.92% yield) as a yellow oil.

[0469] Preparation of Compound K101-C1348-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol, 1.00 eq) in DCM (2.00 mL) were added **C13-48** (34.07 mg, 101.56 μmol, 2.00 eq), DMAP (24.82 mg, 203.12 μmol, 4.00 eq), HOBt (13.72 mg, 101.56 μmol, 2.00 eq) and EDC (19.47 mg, 101.56 μmol, 2.00 eq). The mixture was stirred at 10°C for 12hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The mixture was combined with a second preparation, and the combined mixture was quenched with saturated NaHCO₃ (10 mL) followed by extraction with DCM (20 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1348-A** (18.00 mg, 19.82 μmol, 39.03% yield) as a white solid.

[0470] ^1H NMR (400MHz, CDCl_3) δ 7.57 (s, 1H), 7.44-7.42 (m, 5H), 7.31-7.29 (m, 7H), 7.24-7.18 (m, 8H), 5.59-5.58 (m, 1H), 3.51-3.50 (m, 2H), 3.28 (s, 1H), 2.93 (s, 1H), 2.60-2.56 (m, 2H), 2.50-2.42 (m, 1H), 2.05-1.99 (m, 2H), 1.97-1.95 (m, 3H), 1.77 (s, 2H), 1.55-1.54 (m, 2H), 1.34 (s, 9H), 1.25-1.19 (m, 12H), 1.08 (s, 3H), 0.87-0.79 (m, 4H).

[0471] Preparation of Compound **K101-C1348** (as a mixture of **K101-C134801** and **K101-C134802**). To a solution of **K101-C1348-A** (48.00 mg, 52.85 μmol , 1.00 eq) in THF (2.00 mL) was added TFA (6.03 mg, 52.85 μmol , 3.91 μL , 1.00 eq). The mixture was stirred at 10°C for 12hr to give a yellow solution. LC-MS and TLC (eluting with: EtOAc/MeOH=10/1) showed the reaction was complete. The reaction mixture was concentrated by N_2 and the resultant product dissolved in MeOH (30 mL). The reaction mixture was stirred at 20°C for 12hr. After the mixture was concentrated, the product of **K101-C1348** was purified by prep-TLC (eluting with: EtOAc/MeOH=10/1) to give **K101-C1348** as a mixture of stereoisomers of **K101-C134801** (11.10 mg, 19.62 μmol , 37.12% yield, 100% purity) and **K101-C134802** (10.30 mg, 17.73 μmol , 33.55% yield, 97.4% purity), each as a white solid.

[0472] **K101-C134801**: LC-MS (m/z): 588.2 [$\text{M}+\text{Na}$] $^+$


K101-C134801: ^1H NMR (400MHz, CD_3OD) δ 7.53 (s, 1H), 7.30-7.10 (m, 5H), 5.65-5.55 (m, 1H), 4.00-3.90 (m, 2H), 3.50-3.45 (m, 1H), 3.25-3.20 (m, 1H), 3.15-3.05 (m, 1H), 2.65-2.55 (m, 2H), 2.55-2.40 (m, 2H), 2.20-1.95 (m, 2H), 1.807-1.55 (m, 8H), 1.40-1.25 (m, 6H), 1.20 (s, 3H), 1.10 (s, 3H), 0.95-0.85 (m, 4H).

[0473] **K101-C134802**: LC-MS (m/z): 588.3 [$\text{M}+\text{Na}$] $^+$

[0474] **K101-C134802**: ^1H NMR (400MHz, CD_3OD) δ 7.53 (s, 1H), 7.30-7.10 (m, 5H), 5.65-5.55 (m, 1H), 4.0-3.85 (m, 2H), 3.70-3.60 (m, 1H), 3.20-3.10 (m, 1H), 3.05-3.0 (m, 1H), 2.70-2.55 (m, 2H), 2.55-2.40 (m, 2H), 2.15-2.15 (m, 2H), 1.85-1.45 (m, 8H), 1.45-1.30 (m, 6H), 1.20 (s, 3H), 1.09 (s, 3H), 0.95-0.90 (m, 4H).

Example 45A: Synthesis Scheme of **K101-C134801**

[0475] The scheme for synthesis of compound **K101-C134801** is illustrated below.



[0476] Preparation of compound C134801-C. To a solution of **C1348-D** (2 g, 7.35 mmol, 1 eq) in DMA (2 mL) were added CuI (139.97 mg, 734.96 μmol , 0.1 eq) and **C134801-B** (4.06 g, 10.29 mmol, 1.4 eq) and Pd(dppf)Cl₂ (537.77 mg, 734.96 μmol , 0.1 eq) under N₂. The mixture was stirred under N₂ at 90°C for 5hr to give a black suspension. LCMS and TLC (eluting with: PE/EtOAc=3/1) showed the reaction was complete. The reaction mixture was quenched with H₂O (100 mL) and extracted with MBTE (40 mLx 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The crude product was purified by a flash column (eluting with: PE/EtOAc=100%PE to 20%) to give **C134801-C** (750 mg, 2.16 mmol, 29.37% yield) as a yellow oil.

[0477] ¹H NMR (400MHz, CDCl₃): δ 7.30-7.20 (m, 2H), 7.20-7.16 (m, 3H), 5.58-5.51 (m, 1H), 5.34-5.27 (m, 1H), 5.02-5.00 (m, 1H), 4.37-4.33 (m, 1H), 3.73 (s, 3H), 2.62-2.58 (m, 2H), 2.46-2.44 (m, 2H), 2.07-2.02 (m, 2H), 1.70-1.66 (m, 2H), 1.44 (s, 9H).

[0478] Preparation of compound C134801-D. To a solution of **C134801-C** (0.75 g, 2.16 mmol, 1 eq) in MeOH (15 mL) was added Pd/C (500 mg, 2.16 mmol, 50% purity, 1 eq) under N₂. The suspension was degassed under vacuum and purged with H₂ several times. The mixture was stirred under H₂ (15psi) at 20°C for 12 hours. LCMS showed the reaction was complete. The reaction mixture was filtered on celite. The filtrate was concentrated to give **C134801-D** (750 mg, 2.15 mmol, 99.42% yield) as a black oil, which was used for next step without further purification.

[0479] ¹H NMR (400MHz, CDCl₃): δ 7.29-7.26 (m, 2H), 7.19-7.16 (m, 3H), 4.99-4.84 (m, 1H), 4.29-4.28 (m, 1H), 3.73 (s, 3H), 2.61-2.51 (m, 2H), 1.78-1.60 (m, 4H), 1.45 (s, 9H), 1.33-1.28 (m, 6H).

[0480] Preparation of compound BB-C134801. To a solution of **C134801-D** (750 mg, 2.15 mmol, 1 eq) in THF (5 mL) /H₂O (1 mL) was added LiOH.H₂O (90.06 mg, 2.15 mmol, 1 eq) at 0°C. The mixture was allowed to stir at 20°C for 12hr to give a yellow solution. LCMS showed the reaction was complete. The reaction mixture was acidified to pH=4 with HCl (1N) and extracted with MBTE (20 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give **BB-C134801** (700 mg, 2.09 mmol, 97.24% yield) as a yellow oil, which was used for next step without further purification.

[0481] Preparation of compound C134801-B. Zinc (6 g) was treated with 1N HCl aqueous (30 mL) with stirring for 10 min. Then it was filtered and washed with water (30 mL), EtOH (30 mL) and toluene (30 mL) in sequence, dried in vacuum to afford the zinc powder for next step. A mixture of activated Zn (2.62 g, 40.11 mmol, 4 eq) and I₂ (127.24 mg, 501.32 μmol, 100.98 uL, 0.05 eq) in DMA (10 mL) was stirred at 20°C for 5 min. Then **C134801-A** (3.3 g, 10.03 mmol, 1 eq) in DMA (10 mL) was added dropwise. The reaction mixture was stirred at 20°C for 25 min to give a black suspension. The reaction mixture (about 0.5015 mmol/mL) was used for next step without further purification.

[0482] Preparation of compound K101-C134801-A. To a solution of **K101-C20Tr-B** (200 mg, 338.55 μmol, 1 eq) in DCM (3 mL) were added **BB-C134801** (193.06 mg, 575.54 μmol, 1.7 eq), DMAP (165.44 mg, 1.35 mmol, 4 eq), HOBt (50.32 mg, 372.41 μmol, 1.1 eq) and EDCI (110.33 mg, 575.54 μmol, 1.7 eq). The mixture was stirred at 20°C for 12hr to give a yellow solution. LCMS showed the desired mass was found, and **K101-C20Tr-B** was remained. The mixture was stirred at 20°C for 12hr again. LCMS showed the desired mass was found, and **K101-C20Tr-B** was remained. The mixture was stirred at 20°C for 12hr to give a yellow solution again. LCMS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture was quenched with H₂O (10 mL) and extracted with MBTE (15 mL x 3). The organic layers were dried over Na₂SO₄ and

concentrated to give the crude product. The crude product was purified by flash column (eluting with: PE/EtOAc=100% PE to 40%) to give **K101-C134801-A** (210 mg, 231.23 μ mol, 68.30% yield) as a yellow solid.

[0483] ^1H NMR (400MHz, CDCl_3) δ 7.58 (s, 1H), 7.44-7.42 (m, 6H), 7.31-7.26 (m, 7H), 7.26-7.17 (m, 7H), 5.60-5.59 (m, 1H), 5.26-5.17 (m, 1H), 5.95-5.93 (m, 1H), 4.28-4.27 (m, 1H), 3.52 (s, 2H), 3.27 (s, 1H), 2.94 (s, 1H), 2.61-2.43 (m, 5H), 2.09-2.05 (m, 1H), 2.04-1.99 (m, 1H), 1.78-1.77 (m, 4H), 1.58-1.56 (m, 1H), 1.54-1.52 (m, 1H), 1.44 (s, 9H), 1.33-1.28 (m, 6H), 1.19 (s, 3H), 1.08 (s, 3H), 0.88-0.78 (m, 4H).

[0484] Preparation of compound **K101-C134801** and **K101-C134801-C**. To a solution of **K101-C134801-A** (610.00 mg, 671.68 μ mol, 1.00 eq) in DCM (5 mL) was added TFA (2.76 g, 24.23 mmol, 1.79 mL, 36.08 eq). The mixture was stirred at 20°C for 1hr to give a yellow solution. LCMS and TLC (eluting with: EtOAc/MeOH=10/1) showed the reaction was complete. The reaction mixture was concentrated by purging with N_2 . The residue from concentration was dissolved in MeOH (50 mL). The reaction mixture was stirred at 40°C for 12hr. LCMS showed the reaction was complete. The reaction mixture was concentrated to give **K101-C134801** (158 mg, 255.88 μ mol, 38.10% yield, 91.62% purity) as a white solid, which was the final product, and **K101-C134801-C** (130 mg, 160.88 μ mol, 23.95% yield) as a yellow solid, which was an intermediate.

[0485] **K101-C134801**: LC-MS (m/z): 588.2 $[\text{M}+\text{Na}]^+$

[0486] **K101-C134801**: ^1H NMR (400MHz, CD_3OD) δ 7.56 (s, 1H), 7.30-7.10 (m, 5H), 5.65-5.55 (m, 1H), 4.00-3.90 (m, 2H), 3.50-3.45(m, 1H), 3.25-3.20 (m, 1H), 3.15-3.05 (m, 1H), 2.65-2.40 (m, 4H), 2.20-1.95 (m, 2H), 1.807-1.55 (m, 8H), 1.40-1.25 (m, 6H), 1.20 (s,3H), 1.10 (s,3H), 0.95-0.85 (m, 4H).

[0487] **K101-C134801-C**: ^1H NMR (400MHz, CDCl_3) δ 7.51 (s, 1H), 7.36-7.35 (m, 6H), 7.24-7.22 (m, 7H), 7.18-7.10 (m, 7H), 5.54 (s, 1H), 5.26 (brs, 1H), 3.45-3.42 (m, 2H), 3.36-3.34(m, 1H), 3.21 (s, 1H), 2.87 (s, 1H), 2.54-2.36 (m, 4H), 1.99-1.97 (m, 3H), 1.71 (s, 3H), 1.27-1.21 (m, 6H), 1.13 (s,3H), 1.01 (s,3H), 0.81-0.71 (m, 4H).

[0488] Preparation of compound **K101-C134801**. To a solution of **K101-C134801-C** (130 mg, 160.88 μ mol, 1 eq) in MeOH (5 mL) was added HClO_4 (32.32 mg, 321.76 μ mol, 19.47 μ L, 2 eq) at 0°C. The mixture was stirred at 0°C for 0.5hr to give a yellow solution. LCMS and TLC (eluting with: EtOAc=10/1) showed the reaction was complete. The reaction mixture was quenched with saturated NaHCO_3 (10 mL) and extracted with EtOAc (20 mL x 3). The organic layers were dried over Na_2SO_4 and concentrated to give the crude product. The crude


product was purified by prep-TLC (eluting with: EtOAc=10/1) to give **K101-C134801** (20.5 mg, 30.07 μ mol, 18.69% yield, 82.99% purity) as a white solid

[0489] LC-MS (m/z): 588.2 [M+Na]⁺

[0490] ¹H NMR (400MHz, CD₃OD): δ 7.56 (s, 1H), 7.30-7.10 (m, 5H), 5.65-5.55 (m, 1H), 4.00-3.90 (m, 2H), 3.50-3.45(m, 1H), 3.25-3.20 (m, 1H), 3.15-3.05 (m, 1H), 2.65-2.40 (m, 4H), 2.20-1.95 (m, 2H), 1.807-1.55 (m, 8H), 1.40-1.25 (m, 6H), 1.20 (s,3H), 1.10 (s,3H), 0.95-0.85 (m, 4H).

Example 45B: Synthesis Scheme of **K101-C134802**

[0491] The scheme for synthesis of compound **K101-C134802** is illustrated below.



[0492] Preparation of compound **C1348-E**. To a solution of **C1348-E** (20 g, 146.85 mmol, 20.00 mL, 1 eq) in DCM (200 mL) was added 2,4-lutidine (25.18 g, 234.96 mmol, 27.16 mL, 1.6 eq) at -78°C . Then triflic anhydride (Ti_2O) (45.58 g, 161.54 mmol, 26.65 mL, 1.1 eq) was added dropwise at -78°C . The mixture was stirred at -78°C for 0.5hr to give a yellow suspension. TLC (eluting with: PE/EtOAc=3/1) showed the reaction was complete. The reaction mixture was diluted with PE (40

mL). The mixture was poured into silica gel and washed with PE/EtOAc (4L, 4/1) to give **C1348-F** (30.5 g, 113.70 mmol, 77.42% yield) as yellow oil.

[0493] Preparation of compound C1348-G. To a solution of ethynyl(trimethyl)silane (15.25 g, 155.30 mmol, 21.51 mL, 1.49 eq) in THF (200 mL) was added dropwise n-BuLi (2.5 M, 51.67 mL, 1.24 eq) at -78°C. The mixture was warmed to 0°C for 30min. Then **C1348-F** (28 g, 104.38 mmol, 1 eq) was added dropwise at -78°C. The mixture was warmed to 0°C for 1hr to give a yellow solution. TLC (eluting with: PE/EtOAc=10/1) showed the reaction was complete. The reaction mixture was quenched with saturated NH₄Cl (200 mL) and extracted with MBTE (150 mLx 2). The organic layers were dried over Na₂SO₄ and concentrated to give **C1348-G** (25 g, crude) as a yellow oil, which was used for next step without further purification.

[0494] ¹H NMR (400MHz, CDCl₃): δ 7.30-7.18 (m, 5H), 3.80-3.70 (m, 1H), 2.80-2.69 (m, 2H), 2.26-2.22 (m, 2H), 1.89-1.84 (m, 2H), 0.23-0.09 (m, 9H).

[0495] Preparation of compound C1348-H. To a solution of **C1348-G** (25 g, 115.53 mmol, 1 eq) in THF (50 mL) was added tetra-n-butylammonium fluoride (TBAF) (1 M, 150.19 mL, 1.3 eq). The mixture was stirred at 20°C for 1hr to give a yellow solution. TLC (eluting with: PE/EtOAc=10/1) showed the reaction was complete. The reaction mixture was quenched with H₂O (300 mL) and extracted with EtOAc (150 x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The crude product was purified by a flash column (eluting with: PE/EtOAc=100%PE to 5%) to give **C1348-H** (9.5 g, 65.88 mmol, 57.02% yield) as a colorless oil.

[0496] ¹H NMR (400MHz, CDCl₃): δ 7.32-7.13 (m, 5H), 2.74-2.65 (m, 2H), 2.14-2.11 (m, 5H), 1.93 (s, 1H), 1.80-1.67 (m, 2H).

[0497] Preparation of compound C1348-D. To a solution of **C1348-H** (3 g, 20.80 mmol, 1 eq) in THF (50 mL) was added ZrCp₂HCl (8.88 g, 33.28 mmol, 1.6 eq) at 0°C. The mixture was stirred at 0°C for 2.5hr, then stirred at 20°C for 1hr. I₂ (6.34 g, 24.96 mmol, 5.03 mL, 1.2 eq) in THF (10 mL) was added at -78°C. The mixture was stirred at -78°C for 1hr. The mixture was allowed to stir at 0°C for 1hr to give a brown suspension. TLC (eluting with:PE=100%) showed the reaction was completed. The reaction mixture was quenched with HCl (0.1N, 200 mL). The mixture was extracted with EtOAc (30 mLx 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The crude product was purified by column silica (eluting with: PE=100%) to give **C1348-D** (3.8 g, 13.96 mmol, 67.13% yield) as a yellow oil.

[0498] ¹H NMR (400MHz, CDCl₃): δ 7.34-7.21 (m, 2H), 7.19-7.15 (m, 3H), 6.55-6.51 (m, 1H), 6.02-5.99 (m, 1H), 2.64-2.60 (m, 2H), 2.10-2.07 (m, 2H), 1.73-1.71 (m, 2H).

[0499] Preparation of compound C134802-D. To a solution of **C134802-C** (4.35 g, 11.02 mmol, 1.5 eq) in DMA (10 mL) were added CuI (139.97 mg, 734.96 μmol, 0.1 eq) and **C1348-D** (2 g, 7.35

mmol, 1 eq) and Pd(dppf)Cl₂ (537.77 mg, 734.96 μmol, 0.1 eq) under N₂. The mixture was stirred under N₂ at 90°C for 12hr to give a black suspension. LCMS and TLC (eluting with: PE/EtOAc=4/1) showed the reaction was complete. The reaction mixture was quenched with H₂O (200 mL) and extracted with MBTE (100mLx 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The crude product was purified by flash column (eluting with: PE/EtOAc=100%PE to 10%) to give **C134802-D** (1 g, 2.88 mmol, 39.16% yield) as a yellow oil.

[0500] ¹H NMR (400MHz, CDCl₃) δ 7.30-7.27 (m, 1H), 7.20-7.16 (m, 1H), 5.58-5.1 (m, 1H), 5.34-5.26 (m, 1H), 5.02-5.00 (m, 2H), 4.37-4.32 (m, 1H), 3.73 (s, 3H), 2.62-2.58 (m, 2H), 2.48-2.44 (m, 2H), 2.07-2.02 (m, 2H), 1.71-1.66 (m, 2H), 1.44 (s, 9H).

[0501] Preparation of compound C134802-E. To a solution of **C134802-D** (1 g, 2.88 mmol, 1 eq) in MeOH (20 mL) was added Pd/C (200 mg, 2.88 mmol, 50% purity, 1 eq) under N₂. The suspension was degassed under vacuum and purged with H₂ several times. The mixture was stirred under H₂ (15psi) at 20°C for 12 hours to give a yellow solution. LCMS and TLC(eluting with: PE/EtOAc=4/1) showed the reaction was complete. The reaction mixture was filtered on celite and washed with MeOH (60 mL). The filtrate was concentrated to give **C134802-E** (800 mg, 2.29 mmol, 79.54% yield) as a yellow oil, which was used for next step without further purification.

[0502] ¹H NMR (400MHz, CDCl₃): δ 7.31-6.98 (m, 5H), 5.05-4.98 (m, 1H), 4.31-4.26 (m, 1H), 3.73 (s, 3H), 2.61-2.57 (m, 2H), 1.76-1.62 (m, 4H), 1.44 (s, 9H), 1.33-1.22 (m, 4H).

[0503] Preparation of compound BB-C134802. To a solution of **C134802-E** (700 mg, 2.00 mmol, 1 eq) in THF (10 mL) /H₂O (1.4 mL) was added LiOH.H₂O (84.06 mg, 2.00 mmol, 1 eq) at 0°C. The mixture was allowed to stir at 25°C for 12hr to give a yellow solution. LCMS and TLC (eluting with: EtOAc=100%) showed the reaction was complete. The reaction mixture was extracted with MBTE (20 mL). The water layer was acidified to pH=3 with HCl (0.5N) and extracted with EtOAc(30 mLx 3). The organic layers were dried over Na₂SO₄ and concentrated to give **BB-C134802** (560 mg, 1.61 mmol, 80.54% yield, 96.63% purity, 98.7% ee%) as a yellow oil.

[0504] ¹H NMR (400MHz, CDCl₃): δ 7.28-7.16 (m, 5H), 7.05-7.03 (m, 1H), 3.87-3.81 (m, 1H), 2.57-2.55 (m, 2H), 1.61-1.53 (m, 4H), 1.37 (s, 9H), 1.32-1.18 (m, 6H).

[0505] Preparation of compound K101-C134802-A. To a solution of **K101-C20Tr-B** (200 mg, 338.55 μmol, 1.00 eq) in DCM (5 mL) were added **BB-C134802** (136.28 mg, 406.27 μmol, 1.2 eq), DMAP (165.45 mg, 1.35 mmol, 4.00 eq), HOBT (48.03 mg, 355.48 μmol, 1.05 eq) and EDCI (77.88 mg, 406.27 μmol, 1.2 eq). The mixture was stirred at 20°C for 12hr to give a yellow solution. LCMS showed desired mass was found and **K101-C20Tr-B** was remained. The reaction was stirred at 20°C for 16hr again to give yellow solution. LCMS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The mixture was quenched with saturated NaHCO₃ (50 mL) and extracted

with DCM (80 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The crude product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C134802-A** (200 mg, 220.22 μmol, 65.05% yield) as a yellow solid.



[0506] Preparation of compound **K101-C134802**. To a solution of **K101-C134802-A** (340.00 mg, 374.38 μmol, 1.00 eq) in DCM (5 mL) was added TFA (1.54 g, 13.51 mmol, 1 mL, 36.08 eq). The mixture was stirred at 20°C for 1hr to give a yellow solution. LCMS showed the reaction was complete. The reaction mixture was concentrated by purging with N₂. The residue was dissolved in MeOH (50 mL). The reaction mixture was stirred at 40°C for 12hr. LCMS showed the reaction was complete. The reaction mixture was concentrated to give **K101-C134802** (107 mg, 170.84 μmol, 45.63% yield, 90.33% purity) as a white solid.

[0507] LC-MS (m/z): 566.4 [M+H]⁺

[0508] ¹H NMR (400MHz, CD₃OD): δ 7.56 (s, 1H), 7.30-7.10 (m, 5H), 5.70-5.60 (m, 1H), 4.0-3.90 (m, 2H), 3.50-3.40 (m, 1H), 3.20-3.0 (m, 2H), 2.70-2.55 (m, 2H), 2.55-2.40 (m, 2H), 2.15-2.05 (m, 2H), 1.85-1.45 (m, 8H), 1.45-1.30 (m, 6H), 1.20 (s, 3H), 1.09 (s, 3H), 0.95-0.90 (m, 4H).

Example 46: Synthesis Scheme of **K101-C1349**.

[0509] The scheme for synthesis of compound **K101-C1349** is illustrated below.

**Preparation of C13-49:**

[0510] Preparation of Compound C13-49-B. To a solution of C13-49-A (1.00 g, 6.09 mmol, 775.19 μ L, 1.00 eq) in toluene (5.00 mL) were added 2-methylpropanenitrile (1.68 g, 24.36 mmol, 4.00 eq) and KHMDS (1 M, 9.14 mL, 1.50 eq). The mixture was stirred at 60°C for 12hr to give a black solution. LC-MS and TLC (eluting with: PE/EtOAc=5/1) showed the reaction was complete. The reaction mixture was quenched with Sat.NH₄Cl (50 mL) and extracted with EtOAc (30 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by column chromatography on silica gel (eluting with: PE/EtOAc=100%PE to 10/1) to give C13-49-B (1.10 g, 5.16 mmol, 84.72% yield) as a colorless oil.

[0511] Preparation of Compound C13-49-C. To a solution of **C13-49-B** (1.10 g, 5.16 mmol, 1.00 eq) in toluene (10.00 mL) was added dropwise diisobutylaluminium hydride (1 M, 6.71 mL, 1.30 eq) at -50°C. The mixture was stirred at -50°C for 0.5hr and then stirred at 0°C for 0.5hr to give a colorless solution. LC-MS showed the reaction was complete. The reaction mixture was quenched with H₂SO₄ (1.5M, 9 mL), stirred at 0°C for 3hr, and allowed to stand for 12hr. The mixture was extracted with MTBE (30 mL x 3), the combined organic layers dried over Na₂SO₄ and then concentrated to give **C13-49-C** (1.30 g, crude) as a colorless oil.

[0512] Preparation of Compound C13-49-F. To a solution of **C13-49-C** (1.60 g, 7.40 mmol, 1.00 eq) in H₂O (20.00 mL) were added (NH₄)₂CO₃ (1.42 g, 14.80 mmol, 1.58 mL, 2.00 eq) and KCN (481.92 mg, 7.40 mmol, 317.05 uL, 1.00 eq). The mixture was stirred at 110°C for 12hr to give a yellow suspension. LC-MS showed the reaction was complete. The reaction mixture was concentrated to give the crude product, which was washed with PE/H₂O (3:1, 50 mL) and filtered to give **C13-49-F** (1.40 g, 4.89 mmol, 66.09% yield) as a yellow solid.

[0513] Preparation of Compound C13-49-E. To a solution of **C13-49-F** (900.00 mg, 3.14 mmol, 1.00 eq) in EtOH (2.00 mL) was added NaOH (3 M, 5.23 mL, 5.00 eq). The mixture was stirred at 100°C for 12hr to give a yellow suspension. LC-MS showed the reaction was complete. The reaction mixture was used for next step without further purification.

[0514] Preparation of Compound C13-49. Boc₂O (1.37 g, 6.28 mmol, 1.44 mL, 2.00 eq) was added to the preparation of **C13-49-E** and the mixture stirred at 10°C for 12hr to give a yellow suspension. LC-MS showed the reaction was complete. The reaction mixture was diluted with H₂O (20 mL) and extracted with PE (30 mL x 3). The water layer was adjusted to pH 4 with HCl (1N) and extracted with EtOAc (30 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give **C13-49** (1.05 g, 2.91 mmol, 92.54% yield) as a yellow gum. The product was used for next step without further purification.

[0515] Preparation of Compound K101-C1349-A. To a solution of **K101-C20Tr-B** (50.00 mg, 84.64 μmol, 1.00 eq) in DCM (2.00 mL) were added **C13-49** (91.75 mg, 253.92 μmol, 3.00 eq), DMAP (41.36 mg, 338.56 μmol, 4.00 eq), HOBt (13.72 mg, 101.57 μmol, 1.20 eq) and EDC (32.45 mg, 169.28 μmol, 2.00 eq). The mixture was stirred at 10°C for 12hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture was combined with a second preparation, and quenched with saturated NaHCO₃ (20 mL). The mixture was extracted with DCM (20 mL x 3) and the combined organic layers dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1349-A** (60.00 mg, 64.23 μmol, 63.24% yield) as a white solid.

[0516] Preparation of Compound K101-C1349. To a solution of **K101-C1349-A** (60.00 mg, 64.23 μmol, 1.00 eq) in tetrahydrofuran (THF) (3.00 mL) was added TFA (1.54 g, 13.51 mmol, 1.00 mL,

210.28 eq), and the mixture stirred at 10°C for 12hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated, dissolved in MeOH (30 mL), and the stirred at 40°C for 12hr. The mixture was concentrated to give the crude product, which was then dissolved in saturated NaHCO₃ (10 mL) and extracted with EtOAc (20 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: EtOAc / MeOH=10/1) to give **K101-C134901** (11.50 mg, 18.45 μmol, 28.72% yield, 94.9% purity) and **K101-C134902** (10.60 mg, 15.28 μmol, 23.79% yield, 85.3% purity) as white solids.

[0517] K101-C134901 LC-MS (m/z): 614.3 [M+Na]⁺


[0518] K101-134901 ¹H NMR (400MHz, CD₃OD) δ7.65 (s, 4H), 7.53 (s, 1H), 5.55-5.53 (m, 1H), 3.99-3.92 (m, 2H), 3.68(s, 1H), 3.15 (s, 1H), 3.01 (s, 1H), 2.54-2.41 (m, 2H), 2.04-1.98 (m, 2H), 1.76-1.75 (m, 3H), 1.52-1.45 (m, 6H), 1.31-1.28 (m, 1H), 1.01(s, 3H), 0.89-0.86 (m, 6H), 0.57-0.55 (m, 1H).

[0519] K101-C134902 LC-MS (m/z): 614.3 [M+Na]⁺

[0520] K101-C134902 ¹H NMR (400MHz, CD₃OD) δ7.65 (s, 4H), 7.53 (s, 1H), 5.55-5.53 (m, 1H), 3.93-3.89 (m, 2H), 3.67(s, 1H), 3.16 (s, 1H), 3.00 (s, 1H), 2.48-2.40 (m, 2H), 1.94-1.93 (m, 1H), 1.75-1.69 (m, 4H), 1.49-1.48 (m, 6H), 1.04-1.01 (m, 4H), 0.97-0.96 (m, 3H), 0.84-0.80 (m, 4H).

Example 47: Synthesis Scheme of **K101-C1350**.

[0521] The scheme for synthesis of compound **K101-C1350** is illustrated below.



[0522] Preparation of Compound **K101-C1350-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol, 1.00 eq) in DCM (1.00 mL) were added **C13-50** (74.49 mg, 253.91 μmol, 5.00 eq), EDC (58.41 mg, 304.70 μmol, 6.00 eq) and DMAP (37.22 mg, 304.70 μmol, 6.00 eq). The mixture was stirred at 20°C for 14hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was quenched with H₂O (15 mL) and extracted with DCM (15 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give a yellow oil. The product was purified by prep-TLC (eluting with Petroleum ether: Ethyl acetate = 3/1) to give **K101-C1350-A** (25.00 mg, 28.87 μmol, 56.84% yield) as a white solid.


[0523] Preparation of Compound K101-C1350. To a solution of **K101-C1350-A** (25.00 mg, 28.87 μmol , 1.00 eq) in THF (1.00 mL) were added TFA (770.00 mg, 6.75 mmol, 500.00 μL , 233.92 eq) and Et_3SiH (3.36 mg, 28.87 μmol , 4.60 μL , 1.00 eq). The mixture was stirred at 20°C for 5hr and then concentrated to give a yellow oil. The product was dissolved with DCM (1 mL), followed by addition of TFA (0.5 mL). The mixture stirred at 20°C for 1hr to give a yellow oil. LC-MS showed the reaction was complete. Following concentration, the resultant yellow oil was dissolved with MeOH (2 mL), and the mixture stirred at 20°C for 14hr. The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 23%-53%, 10min), and the organic layers lyophilized to give **K101-C1350** (6.00 mg, 9.41 μmol , 32.59% yield, 100% purity, TFA) as a white solid.

[0524] LC-MS (m/z): 546.2 [M+Na]⁺

[0525] ¹H NMR (400MHz, CD₃OD) δ 7.56 (s, 1H), 7.35-7.22 (m, 5H), 5.64 (s, 1H), 4.00-3.96 (m, 2H), 3.59-3.56 (m, 1H), 3.19 (s, 1H), 3.08 (s, 1H), 2.87-2.86 (m, 1H), 2.77-2.74 (m, 3H), 2.52-2.46 (m, 2H), 2.19-2.18 (m, 1H), 2.03-1.99 (m, 3H), 1.77 (s, 3H), 1.68-1.64 (m, 1H), 1.19 (s, 3H), 1.10 (s, 3H), 0.98-0.93 (m, 3H).

Example 48: Synthesis Scheme of **K101-C1351**.

[0526] The scheme for synthesis of compound **K101-C1351** is illustrated below.



[0527] Preparation of Compound K101-C1351-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) and **C13-51** (72.31 mg, 152.34 μmol , 3.00 eq) in DCM (1.00 mL) were added EDC (38.94 mg, 203.12 μmol , 4.00 eq) and DMAP (24.82 mg, 203.12 μmol , 4.00 eq). The mixture was stirred at 20°C for 18h to give a yellow solution. The reaction was complete as detected by LC-MS. The reaction solution was diluted with H₂O (15 mL), and extracted with DCM (10 mL x 5). The combined organic layers were dried over anhydrous Na₂SO₄, filtered, and then concentrated under reduced pressure to give a white solid. The product was purified by prep-TLC

(PE/EtOAc=3/1, SiO₂) to give **K101-C1351-A** (30.30 mg, 34.98 μmol, 59.39% yield) as a pale yellow solid.


[0528] Preparation of Compound **K101-C1351**. To a solution of **K101-C1351-A** (30.00 mg, 34.64 μmol, 1.00 eq) in MeOH (500.00 uL) was added HCl/MeOH (4 M, 500.03 uL, 57.74 eq). The solution was stirred at 20°C for 14h to give a colorless solution. The reaction was complete as detected by LC-MS. The reaction solution was diluted with H₂O (10 mL), neutralized with saturated aqueous NaHCO₃, and then extracted with DCM (8 mL x 5). The combined organic layers were dried over anhydrous Na₂SO₄, filtered, and then concentrated under reduced pressure to give a yellow solid. The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25 mm x 10 um; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 23%-53%, 10 min) to give **K101-C1351** (3.80 mg, 7.26 μmol, 20.95% yield, 100% purity, TFA salt) as a white solid.

[0529] LC-MS (m/z): 546.1 [M+Na]⁺

[0530] ¹H NMR (400MHz, MeOD) δ 7.58-7.53 (m, 1H), 7.31-7.25 (m, 2H), 7.22-7.15 (m, 3H), 5.63-5.58 (m, 1H), 4.00-3.90 (m, 2H), 3.89-3.85 (m, 1H), 3.18-3.14 (m, 1H), 3.09-3.03 (m, 1H), 2.68 (t, *J*=6.8 Hz, 2H), 2.57-2.49 (m, 1H), 2.45-2.38 (m, 1H), 2.17 (dd, *J*=6.9, 14.7 Hz, 1H), 2.10-2.01 (m, 1H), 1.95-1.86 (m, 1H), 1.83-1.73 (m, 5H), 1.72-1.65 (m, 1H), 1.58 (dd, *J*=10.3, 14.6 Hz, 1H), 1.11 (s, 3H), 1.07 (s, 3H), 0.95-0.88 (m, 4H).

Example 49: Synthesis Scheme of **K101-C1352**.

[0531] The scheme for synthesis of compound **K101-C1352** is illustrated below.



[0532] Preparation of Compound **K101-C1352-A:** To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) in Py (1.00 mL) were added **C13-52** (179.04 mg, 507.80 μmol , 10.0eq) and DMAP (12.41 mg, 101.56 μmol , 2.00 eq). The mixture was stirred at 90°C for 14hr in sealed tube to give a brown solution. LCMS showed the reaction was completed. The reaction mixture was concentrated to give a black oil. The black oil was dissolved by DCM (5 mL) and was adjusted to pH=4 with HCl (0.1 M) to give a yellow liquid. The yellow liquid was dried over Na_2SO_4 and concentrated to give a yellow oil. The yellow oil was purified by prep-TLC (eluting with Dichloromethane: Methanol = 10/1) to give **K101-C1352-A** (15.00 mg, 15.90 μmol , 31.31% yield) as a white solid.

[0533] Preparation of Compound **K101-C1350:** To a solution of **K101-C1352-A** (15.00 mg, 15.90 μmol , 1.00 eq) in THF (2.00 mL) were added TFA (770.00 mg, 6.75 mmol, 500.00 μL , 424.73 eq) and Et_3SiH (1.85 mg, 15.90 μmol , 2.53 μL , 1.00 eq). The mixture was stirred at 20°C for 5hr to give a colorless solution. LCMS showed the mass of **K101-C1352-A** was remained, then DCM (1 mL) was added. The mixture was stirred at 20°C for 14hr to give a colorless solution. LCMS and TLC (eluting with Dichloromethane: Methanol =8/1) showed the reaction was completed. The reaction mixture was concentrated to give a yellow solid. The yellow solid was purified by prep-TLC (eluting with Dichloromethane: Methanol = 8/1). The organic layers were concentrated to give a white solid.


[0534] The white solid was dissolved with MeCN (1 mL) and H_2O (5 mL), then was lyophilized to give **K101-C1352** (2.00 mg, 2.11 μmol , 13.28% yield, 74% purity) as a white solid.

[0535] LC-MS (m/z): 724.1 [M+Na]⁺

[0536] ¹H NMR (400MHz, CD₃OD) δ 7.53 (s, 1H), 5.59 (s, 1H), 3.95 (s, 2H), 3.15 (s, 1H), 3.06 (m, 1H), 2.64-2.44 (m, 4H), 2.08-2.02 (m, 3H), 1.72 (s, 3H), 1.60-1.57 (m, 4H), 1.27 (s, 30H), 1.18 (s, 3H), 1.06 (s, 3H), 0.89-0.86 (m, 7H).

Example 50: Synthesis Scheme of **K101-C1353**.

[0537] The scheme for synthesis of compound **K101-C1353** is illustrated below.



[0538] Preparation of Compound **K101-C1353-A**: To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol, 1.00 eq), 2-benzylpropanedioic acid (59.17 mg, 304.68 μmol, 6.00 eq) and DMAP (6.20 mg, 50.78 μmol, 1.00 eq) in a mixed solvent of DCM (1.50 mL) and CH₃CN (1.50 mL) was added a solution of DCC (31.43 mg, 152.34 μmol, 3.00 eq) in DCM (1.50 mL) at 0°C dropwise. Then the reaction solution was stirred at 0°C for 15 minutes and 15 °C for 2 hours to give a brown solution. LCMS showed the reaction was completed and the desired MS was observed. The reaction solution was combined with ES5329-254 (10 mg of **K101-C20Tr-B** was used in this batch) and diluted with DCM (5 mL), washed with 0.1 M HCl solution (5 mL x 2), brine (1 mL), dried over anhydrous Na₂SO₄, filtered, concentrated under reduced pressure to give the crude product as a brown gum. The crude product was purified by prep-TLC (PE/EtOAc=1/1, SiO₂) to give **K101-C1353-A** (30.50 mg, 78.32% yield) as a colorless gum. The structure would be confirmed in the next step.

[0539] Preparation of Compound **K101-C1353**: To a solution of **K101-C1353-A** (25.00 mg, 32.60 μmol, 1.00 eq) in DCM (2.50 mL) was added TFA (500.00 uL) at 0°C. Then the reaction solution

was stirred at 0°C for 1 hour to give a clear solution. The reaction solution was quenched by water (2 mL) and then sat. aq. NaHCO₃ solution was added at 0°C to render pH to 6-7. Then the mixture was extracted with DCM (5 mL x 2). The combined extract was washed with brine (5 mL), dried over Na₂SO₄, filtered, concentrated under reduced pressure to give 22.1 mg of brown gum as the crude product. The crude product was purified by prep-TLC (DCM/MeOH=10/1) to give 15 mg of product.


[0540] The product was purified by prep-HPLC (column: Phenomenex Gemini 150*25mm*10um; mobile phase: [water (0.1% TFA)-ACN]; B%: 35%-65%, 10min) to give **K101-C1353** (5.30 mg, 30.74% yield, 99.2% purity) as a white solid after lyophilization. Note: the product is a mixture of two isomers as shown in the synthetic scheme with a ratio of about 1:1 based on NMR and HPLC.

[0541] LC-MS (m/z): 547.7 [M+Na]⁺


[0542] ¹H NMR (400MHz, CD₃OD) δ 7.57-7.50 (m, 1H), 7.33-7.18 (m, 5H), 5.62-5.51 (m, 1H), 3.99-3.87 (m, 2H), 3.79-3.68 (m, 1H), 3.26-3.07 (m, 3H), 3.06-2.96 (m, 1H), 2.56-2.37 (m, 2H), 2.07-1.89 (m, 2H), 1.77-1.71 (m, 2H), 1.77-1.70 (m, 1H), 1.49-1.27 (m, 1H), 1.07 (s, 1H), 1.01 (d, J=4.0 Hz, 3H), 0.92-0.80 (m, 5H), 0.59 (d, J=5.8 Hz, 1H).

Example 51: Synthesis Scheme of **K101-C1354**.

[0543] The scheme for synthesis of compound **K101-C1354** is illustrated below.



Preparation of **C13-54**:



[0544] Preparation of Compound C13-54-B: To a solution of methyl C13-54-A (499.15 mg, 2.77 mmol, 1.00 eq) in DCM (2.00 mL) was added triisopropylsilyl chloride (TIPSCl) (640.87 mg, 3.32 mmol, 712.08 μ L, 1.20 eq), imidazole (565.74 mg, 8.31 mmol, 3.00 eq) and DMAP (33.84 mg, 277.00 μ mol, 0.10 eq). The mixture was stirred at 20°C for 14hr to give a white suspension. LCMS and TLC showed the reaction was completed. The reaction mixture was combined with ES5890-138, then was quenched with H₂O (20 mL) and extracted with DCM (20 mL *5). The organic layers were dried over Na₂SO₄ and then was purified by flash chromatography (eluting with Petroleum ether: Ethyl acetate = 0/1-3/1) to give C13-54-B (900 mg, 2.67 mmol, 96.54% yield) as a colorless oil.

[0545] ¹H NMR (400MHz, CDCl₃) δ = 7.26 (s, 2H), 7.24 - 7.18 (m, 3H), 4.53 (dd, J=5.8, 7.0 Hz, 1H), 3.65 (s, 3H), 3.10 - 2.91 (m, 2H), 1.56 (s, 4H), 1.06 - 1.02 (m, 3H), 1.02 - 0.94 (m, 18H).

[0546] Preparation of Compound C1354: To a solution of C13-54-B (800.00 mg, 2.38 mmol, 1.00 eq) in THF (5.00 mL) and H₂O (1.00 mL) was added LiOH.H₂O (499.32 mg, 11.90 mmol, 5.00 eq). The mixture was stirred at 45°C for 14hr to give a white suspension. LCMS showed the reaction was completed. The reaction mixture was quenched with H₂O (20 mL) and extracted with PE (20 mL *3). The water layers were adjusted to pH 6 with HCl (1N), then was extracted with DCM (20 mL *3). The organic layer was dried over Na₂SO₄ and filtered on silica gel. The filtrate was concentrated to give C1354 (500 mg, 1.55 mmol, 65.14% yield) as a buff oil.

[0547] ¹H NMR (400MHz, CDCl₃) δ = 7.31 - 7.27 (m, 2H), 7.26 - 7.19 (m, 3H), 4.66 (t, J=5.0 Hz, 1H), 3.10 (d, J=5.0 Hz, 2H), 1.09 - 0.98 (m, 21H)

[0548] Preparation of Compound K101-C1354-A: To a solution of (2S)-3-phenyl-2-triisopropylsilyloxy-propanoic acid (87.35 mg, 270.84 μ mol, 4 eq) in DCM (1.00 mL) was added DCC (55.88 mg, 270.84 μ mol, 54.79 μ L, 4 eq). The mixture was stirred at 20°C for 0.5hr to give a solution. Then K101-C20Tr-B (40 mg, 67.71 μ mol, 1.00 eq) and DMAP (66.18 mg, 541.69 μ mol, 8 eq) in DCM (1.00 mL) were added. The mixture was stirred at 20°C for 14hr to give a white suspension. LCMS and TLC showed the reaction was completed. The reaction mixture was combined with ES5890-150, then was quenched with H₂O (15 mL) and extracted with DCM (15 mL *5). The organic layers were dried over Na₂SO₄ and concentrated to give yellow oil. The yellow oil was purified by prep-TLC (eluting with Petroleum ether: Ethyl acetate = 4/1) to give K101-C1354-A (15.00 mg, 15.90 μ mol, 31.31% yield) as a white solid.

[0549] ¹H NMR (400MHz, CDCl₃) δ = 7.58 (s, 1H), 7.45 (d, J=7.5 Hz, 6H), 7.33 - 7.27 (m, 6H), 7.25 - 7.15 (m, 8H), 5.65-5.55 (m, 1H), 5.42 (s, 1H), 4.61 (t, J=5.3 Hz, 1H), 3.52 (s, 2H), 3.30-3.20 (m, 1H), 3.05 (d, J=5.8 Hz, 2H), 2.86 (s, 1H), 2.53 - 2.28 (m, 2H), 1.99 - 1.82 (m, 3H), 1.77 (d, J=1.5 Hz, 3H), 1.03 - 0.96 (m, 21H), 0.91 - 0.80 (m, 6H), 0.55 (d, J=5.0 Hz, 1H)

[0550] Preparation of Compound K101-C1354-B: To a solution of **K101-C1354-A** (25 mg, 27.93 μmol , 1 eq) in THF (1 mL) was added TFA (770.00 mg, 6.75 mmol, 500.00 μL , 241.82 eq). The mixture was stirred at 20°C for 14hr to give a yellow solution. LCMS (ES5890-154-P1A) showed the reaction was completed. The reaction mixture was concentrated to give a yellow oil, the yellow oil was dissolved with MeOH (10 mL). The mixture was stirred at 40°C for 14hr to give a yellow solution. The yellow solution was concentrated to give **K101-C1354-B** as yellow oil. The yellow oil was used next step without further purification.



[0551] Preparation of Compound K101-C1354: To a solution of **K101-C1354-B** (18.23 mg, 27.92 μmol , 1 eq) in THF (1 mL) was added TBAF (1 M, 55.84 μL , 2 eq). The mixture was stirred at 10°C for 14 hr to give a yellow solution. LCMS and TLC showed the reaction was completed. The reaction mixture was concentrated to give a yellow oil. The yellow oil was purified by prep-TLC (eluting with Ethyl acetate: Petroleum ether = 3/1) to give **K101-C1354** (3.5 mg, 7.05 μmol , 25.24% yield, 100% purity) as a white solid. 3.5 mg was delivered.

[0552] LC-MS (m/z): 519.1 [M+Na]⁺

[0553] ¹H NMR (400MHz, MeOD) δ = 7.56 (s, 1H), 7.35 - 7.18 (m, 5H), 5.62 (s, 1H), 4.40 (dd, J=4.6, 7.7 Hz, 1H), 4.01 - 3.88 (m, 2H), 3.22 - 2.91 (m, 4H), 2.60 - 2.37 (m, 2H), 2.12 - 1.95 (m, 3H), 1.76 (d, J=1.5 Hz, 3H), 1.09 (d, J=18.1 Hz, 6H), 0.92 - 0.82 (m, 4H)

Example 52: Synthesis Scheme of **K101-C1355**.

[0554] The scheme for synthesis of compound **K101-C1355** is illustrated below.

**Preparation of C13-55:**

[0555] Preparation of Compound C13-55-B: To a solution of **C13-55-B** (100.00 mg, 601.79 μmol , 1.00 eq) in DCM (1.00 mL) was added imidazole (122.91 mg, 1.81 mmol, 3.00 eq), DMAP (73.52 mg, 601.79 μmol , 1.00 eq) and TIPSCl (139.23 mg, 722.15 μmol , 154.70 μL , 1.20 eq). The mixture was stirred at 10°C for 12hr to give a yellow suspension. LCMS and TLC (eluting with: PE/EtOAc=5/1) showed the reaction was completed. The reaction mixture was quenched with saturated NaHCO_3 (15mL) and extracted with DCM (20 mL*3). The organic layers were dried over Na_2SO_4 and concentrated to give the crude product. The crude product was purified by prep-TLC (eluting with: PE/EtOAc=5/1) to give **C13-55-B** (150.00 mg, 465.10 μmol , 77.29% yield) as a colorless oil.

[0556] ^1H NMR (400MHz, CDCl_3) δ 7.41-7.26 (m, 5H), 4.54-4.51 (t, $J=6.0\text{Hz}$, 1H), 3.65 (s, 3H), 3.09-2.95 (m, 2H), 1.06-0.98 (m, 21H).

[0557] Preparation of Compound C13-55: To a solution of **C13-55-B** (1.90 g, 5.65 mmol, 1.00 eq) in THF (10.00 mL) / H_2O (2.00 mL) was added $\text{LiOH}\cdot\text{H}_2\text{O}$ (1.19 g, 28.25 mmol, 5.00 eq). The mixture was stirred at 40°C for 12hr to give a yellow suspension. LCMS showed the reaction was completed. The reaction mixture was concentrated to give the residue. The residue was acidified to pH=4 with HCl (1N) and extracted with EtOAc (20 mL*3). The organic layers were dried over

Na₂SO₄ and concentrated to give **C13-55** (1.10 g, 3.41 mmol, 60.37% yield) as a yellow oil. Used for next step without further purification.

[0558] ¹H NMR (400MHz, DMSO): δ 7.29-7.19 (m, 5H), 4.59-4.13 (m, 1H), 3.01-2.80 (s, 2H), 3.09-2.95 (m, 2H), 1.22-1.02 (m, 1H), 1.01-0.84 (m, 20H).

[0559] Preparation of Compound **K101-C1355-A**: Preparation of Compound **K101-C1355-A**: To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol, 1.00 eq) in DCM (1.00 mL) were added **C13-55** (32.76 mg, 101.57 μmol, 2.00 eq) and DMAP (6.20 mg, 50.78 μmol, 1.00 eq), DCC (20.96 mg, 101.57 μmol, 20.55 uL, 2.00 eq). The mixture was stirred at 10°C for 12hr to give yellow solution. LCMS and TLC (eluting with: PE/EtOAc=2/1) showed 61.484% of desired mass was found, and 23.129% of **K101-C20Tr-B** was remained. The reaction mixture was combined with ES5350-274. The mixture was quenched with H₂O (10 mL) and extracted with DCM (20 mL*3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The crude product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1355-A** (0.03 g, 33.51 μmol, 65.99% yield) as a white solid.

[0560] ¹H NMR (400MHz, CDCl₃) δ 7.60 (s, 1H), 7.46-7.37 (m, 6H), 7.30-7.28 (m, 6H), 7.25-7.22 (m, 8H), 5.59 (s, 1H), 4.49-4.46 (m, 1H), 3.52 (s, 2H), 3.27 (s, 1H), 3.10-2.86 (m, 3H), 2.46-2.38 (m, 2H), 2.09-2.05 (m, 2H), 1.77 (s, 3H), 1.28-1.25 (m, 3H), 1.03-1.02 (m, 18H), 0.97-0.84 (m, 11H), 0.56-0.54 (m, 1H).

[0561] Preparation of Compound **K101-C1355-B**: To a solution of **K101-C1355-A** (0.03 g, 33.51 μmol, 1 eq) in THF (3 mL) was added TFA (3.82 mg, 33.51 μmol, 2.48 uL, 1 eq). The mixture was stirred at 10°C for 12hr to give yellow solution. LCMS showed the reaction was completed. The reaction mixture was concentrated to give the residue by purging with N₂. The residue was dissolved in MeOH (20 mL). The mixture was stirred at 40°C for 12hr. The mixture was concentrated to give **K101-C1355-B** (0.022 g, crude) as a yellow solid. Used for next step without further purification.

[0562] Preparation of Compound **K101-C1355**: To a solution of **K101-C1355-B** (0.022 g, 33.69 μmol, 1 eq) in THF (3 mL) was added TBAF (1 M, 67.39 uL, 2 eq). The mixture was stirred at 10°C for 12hr to give a yellow solution. LCMS and TLC (eluting with: EtOAc/PE=2/1) showed the reaction was completed. The reaction mixture was concentrated to give the crude product by purging with N₂. The crude product was prep-TLC (eluting with: EtOAc/PE=3/1) and lyophilized to give **K101-C1355** (5.3 mg, 10.67 μmol, 31.68% yield) as a white solid. 5.3 mg was delivered.


[0563] LC-MS (m/z): 519.1 [M+Na]⁺

[0564] ¹H NMR (400MHz, CD₃OD) δ 7.57 (s, 1H), 7.31-7.23 (m, 5H), 5.61 (s, 1H), 4.37-4.34 (m, 1H), 4.00-3.96 (s, 2H), 3.18 (s, 1H), 3.12-3.06 (m, 2H), 2.98-2.94 (m, 1H), 2.51-2.46 (m, 2H), 2.12-


2.03 (m, 2H), 1.77 (s, 3H), 1.57-1.54 (m, 1H), 1.06 (s, 3H), 1.00 (s, 3H), 0.93-0.91 (m, 3H), 0.78-0.76 (m, 1H).

Example 53: Synthesis Scheme of **K101-C1356**.

[0565] The scheme for synthesis of compound **K101-C1356** is illustrated below.



Preparation of **C13-56**:



[0566] **Preparation of Compound C13-56-B:** To a solution containing LiAlH_4 (469.66 mg, 12.38 mmol, 1.50 eq) in THF (10.00 mL) was added dropwise 3-[4-(trifluoromethyl)phenyl]propanoic acid (1.80 g, 8.25 mmol, 1.00 eq) in THF (10.00 mL) at 0°C . The mixture was allowed to stir at 10°C for 12hr to give a yellow suspension. LCMS showed the reaction was completed. The reaction mixture was quenched with H_2O (0.47mL), NaOH(15%, 0.47 mL) and H_2O (1.41 mL). The mixture was stirred at 0°C for 20 min. The mixture was filtered on celite. The filtrate was dried over Na_2SO_4 and

concentrated to give **C1356-B** (1.40 g, 6.86 mmol, 83.11% yield) as a yellow oil. Used for next step without further purification.

[0567] ^1H NMR (400MHz, CDCl_3): δ 7.55-7.43 (m, 2H), 7.33-7.28(m, 2H), 3.69-3.68 (m, 2H), 2.76-2.70 (m, 2H), 1.96-1.89 (m, 2H).

[0568] Preparation of Compound C13-56-C: To a solution of oxalyl dichloride (1.74 g, 13.72 mmol, 1.20 mL, 2.00 eq) in DCM (15.00 mL) was added dropwise at -78°C . The mixture was stirred at -78°C for 0.5hr. Then **C13-56-B** (1.40 g, 6.86 mmol, 1.00 eq) in DCM (15.00 mL) was added dropwise at -78°C . The mixture was stirred at -78°C for 1hr. Then Et_3N (3.47 g, 34.30 mmol, 4.75 mL, 5.00 eq) was added dropwise at -78°C . The mixture was stirred at -78°C for 2.5hr to give a yellow suspension. LCMS and TLC(eluting with: PE/EtOAc=2/1) showed the reaction was completed. The reaction mixture was quenched with H_2O (30 mL) and extracted with DCM (30 mL*3). The organic layers were dried over Na_2SO_4 and concentrated to give the crude product. The crude product was purified by column chromatography on silica gel (eluting with: PE/EtOAc=100%PE to 10/1) to give **C13-56-C** (1.10 g, 5.44 mmol, 79.31% yield) as a yellow oil.

[0569] ^1H NMR (400MHz, CDCl_3): δ 9.85 (s, 1H), 7.72-7.56 (m, 2H), 7.35-7.28 (m, 2H), 3.06-3.01 (m, 2H), 2.86-2.81 (m, 2H).

[0570] Preparation of Compound C13-56-D: To a solution of **C13-56-C** (100.00 mg, 494.63 μmol , 1.00 eq) in EtOH (2.00 mL) were added $\text{NH}_3\cdot\text{H}_2\text{O}$ (208.04 mg, 1.48 mmol, 228.62 μL , 25% purity, 3.00 eq) and TMSCN (58.89 mg, 593.56 μmol , 74.54 μL , 1.20 eq). The mixture was stirred at 10°C for 70hr to give yellow solution. LCMS showed the reaction was completed. The reaction mixture was quenched with H_2O (10 mL) and extracted with DCM (30 mL*3). The organic layers were dried over Na_2SO_4 and concentrated to give **C13-56-D** (0.11 g, 482.01 μmol , 97.45% yield) as a yellow oil. Used for next step without further purification.

[0571] ^1H NMR (400MHz, CDCl_3): δ 7.59-7.54 (m, 2H), 7.35-7.30(m, 2H), 3.66-3.58 (m, 1H), 2.99-2.86 (m, 2H), 2.15 -2.05 (m, 2H), 1.66-1.64 (m, 2H).

[0572] Preparation of Compound C13-56-E: To a solution of **C13-56-D** (0.11 g, 482.00 μmol , 1 eq) in EtOH (2 mL) were added NaOH (96.39 mg, 2.41 mmol, 5 eq). The mixture was stirred at 40°C for 2hr. Then H_2O (0.4 mL) was added. The mixture was stirred at 90°C for 3hr to give a yellow solution. LCMS showed the reaction was completed. The reaction mixture was used for next step without further purification.

[0573] Preparation of Compound C13-56: Boc_2O (194.22 mg, 889.92 μmol , 204.44 μL , 2 eq) was added the mixture of **C13-56-E** (0.11 g, 444.96 μmol , 1 eq) from ES5350-284. The mixture was stirred at 10°C for 12hr to give a yellow suspension. LCMS showed the reaction was completed. The reaction mixture was diluted with H_2O (10 mL) and extracted with PE (15 mL*3). The water layers

was acidified to pH=4 with HCl(1N) and extracted with EtOAc (20 mL*3). The organic layers were dried over Na₂SO₄ and concentrated to give **C13-56** (0.11 g, 316.70 μmol, 71.18% yield) as a yellow oil. Used for next step without further purification.

[0574] ¹H NMR (400MHz, DMSO): δ 12.50 (brs, 1H), 7.66-7.64 (m, 2H), 7.49-7.48 (m, 2H), 6.14 (brs, 1H), 3.71 (s, 1H), 2.72-2.71 (m, 2H), 1.95-1.85 (m, 2H), 1.40 (s, 9H).

[0575] Preparation of Compound **K101-C1356-A**: To a solution of **K101-C20Tr-B** (0.05 g, 84.64 μmol, 1 eq) in DCM (2 mL) were added **C13-56** (58.79 mg, 169.28 μmol, 2 eq), DMAP (41.36 mg, 338.55 μmol, 4 eq), HOBT (13.72 mg, 101.57 μmol, 1.2 eq) and EDCI (32.45 mg, 169.28 μmol, 2 eq). The mixture was stirred at 40°C for 12hr to give a yellow solution. LCMS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was completed. The reaction was combined with ES5350-289. The reaction mixture was quenched with H₂O (15 mL) and extracted with DCM (30 mL*3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The crude product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1356-A** (0.039 g, 42.39 μmol, 41.73% yield) as a white solid.

[0576] Preparation of Compound **K101-C1356**: To a solution of **K101-1356-A** (0.039 g, 42.39 μmol, 1 eq) in THF (3 mL) were added TFA (1.54 g, 13.51 mmol, 1 mL, 318.63 eq) and Et₃SiH (4.93 mg, 42.39 μmol, 6.77 uL, 1 eq). The mixture was stirred at 10°C for 12hr to give a yellow solution. LCMS and TLC (eluting with: EtOAc/MeOH=10/1) showed the reaction was completed. The reaction mixture was concentrated to give the residue by purging with N₂. The residue was dissolved MeOH (20 mL). The mixture was stirred at 40°C for 12hr. The mixture was concentrated to give the crude product. The crude product was purified by prep-TLC (eluting with: EtOAc/MeOH=10/1) and lyophilized to give **K101-C135601** (4.7 mg, 7.03 μmol, 16.59% yield, 86.45% purity) as a white solid and **K101-C135602** (3.4 mg, 5.24 μmol, 12.35% yield, 88.96% purity) as a white solid. 4.7mg and 3.4mg were delivered.

[0577] **K101-C135601** LC-MS (m/z): 600.2 [M+Na]⁺


[0578] **K101-C135601** ¹H NMR (400MHz, CD₃OD) δ 7.62-7.58 (m, 3H), 7.46-7.38 (m, 2H), 5.63 (s, 1H), 3.97 (s, 2H), 3.54-3.53 (m, 1H), 3.19-3.09 (m, 2H), 2.84-2.82 (m, 2H), 2.57-2.42 (m, 2H), 2.20-2.08 (m, 4H), 1.77 (s, 3H), 1.64-1.61 (m, 1H), 1.21 (s, 3H), 1.10 (s, 3H), 0.95-0.93 (m, 4H).

[0579] **K101-C135602** LC-MS (m/z): 600.0 [M+Na]⁺

[0580] **K101-C135602** ¹H NMR (400MHz, CD₃OD) δ 7.62-7.57 (m, 3H), 7.46-7.38 (m, 2H), 5.63-5.60 (m, 1H), 3.96 (s, 2H), 3.49-3.46 (m, 1H), 3.19 (s, 1H), 3.09 (s, 1H), 2.86-2.82 (m, 2H), 2.52-2.42 (m, 2H), 2.20-1.91 (m, 4H), 1.77 (s, 3H), 1.61-1.57 (m, 1H), 1.20 (s, 3H), 1.11 (s, 3H), 0.97-0.93 (m, 4H).

Example 54: Synthesis Scheme of **K101-C1357**.

[0581] The scheme for synthesis of compound **K101-C1357** is illustrated below.



[0582] Preparation of Compound **K101-C1357-A**. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μ mol, 1.00 eq) and **C13-57** (40.42 mg, 152.35 μ mol, 3 eq) in DCM (1 mL) were added EDC (58.41 mg, 304.70 μ mol, 6 eq) and DMAP (37.22 mg, 304.70 μ mol, 6 eq). The mixture was stirred at 20°C for 16 hours to give a yellow solution. The reaction was complete detected by LC-MS. The reaction solution was diluted with H₂O (20 mL) and extracted with DCM (10 mL x 5). The combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure to give a yellow solid. The product was purified by prep-TLC (PE/EtOAc=3/1, SiO₂) to give **K101-C1357-A** (16.00 mg, 19.09 μ mol, 37.60% yield) as a white solid.


[0583] Preparation of Compound **K101-C1357**. To a solution of **K101-C1357-A** (16.00 mg, 19.09 μ mol, 1 eq) in MeOH (0.50 mL) was added HCl/MeOH (4 M, 0.50 mL, 104.75 eq). The solution was stirred at 20°C for 12 hours to give a black solution. The reaction was complete as detected by LC-MS. The reaction solution was concentrated under N₂ to give yellow solid, which was then purified by prep-HPLC (column: Phenomenex Gemini 150 x 25 mm x 10 μ m; mobile phase: [A: water (0.1% TFA)-B: ACN]; B%: 20%-50%, 10 min) to give **K101-C1357** (1.2 mg, 1.97 μ mol, 10.31% yield, 91.34% purity, TFA salt) as a white solid.

[0584] LC-MS (m/z): 518.1 [M+Na]⁺

[0585] ¹H NMR (400MHz, CD₃OD) δ 7.53 (s, 1H), 7.40-7.33 (m, 2H), 7.33-7.27 (m, 3H), 5.67 (s, 1H), 5.57-5.58 (m, 1H), 5.22 (s, 1H), 4.29-4.32 (m, 1H), 3.92 (s, 2H), 3.16-3.04 (m, 3H), 3.01-3.08 (m, 1H), 2.54-2.45 (m, 1H), 2.41-2.33 (m, 1H), 2.17 (dd, J=7.0, 14.8 Hz, 1H), 2.06-1.95 (m, 1H), 1.71-1.72 (m, 3H), 1.64-1.54 (m, 1H), 1.03 (s, 3H), 1.01 (s, 3H), 0.93-0.86 (m, 5H).

Example 55: Synthesis Scheme of **K101-C1358**.

[0586] The scheme for synthesis of **K101-C1358** is illustrated below.



[0587] Preparation of Compound K101-C1358-A. To a solution of **K101-C20Tr-B** (30.00 mg, 50.78 μmol , 1.00 eq) and **C13-58** (35.46 mg, 126.96 μmol , 2.50 eq) in DCM (1.00 mL) were added EDCI (58.41 mg, 304.70 μmol , 6 eq) and DMAP (18.61 mg, 152.35 μmol , 3 eq). The mixture was stirred at 15°C for 18 hours to give a yellow solution. The reaction was complete as detected by LC-MS. The reaction solution was combined with a second preparation, which was then diluted with H₂O (20 mL) and extracted with DCM (10 mL \times 5). The organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a yellow solution. The product was purified by prep-TLC (PE/EtOAc=3/1, SiO₂) to give **K101-C1358-A** (39 mg, 45.77 μmol , 90.13% yield) as a yellow solid.


[0588] Preparation of Compound K101-C1358. To a solution of **K101-C1358-A** (39.00 mg, 45.77 μmol , 1.00 eq) in MeOH (0.5 mL) was added HCl/MeOH (4 M, 500 μL , 43.70 eq). The solution was stirred at 10°C for 16 hours to give a black solution. LC-MS showed the reaction was complete. The reaction solution was diluted with H₂O (25 mL) and extracted with DCM (10 mL \times 5). The combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure to give a yellow solid. The product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10 μm ; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 20%-50%, 10 min). The separated layers were lyophilized to give **K101-C1358** (5.00 mg, 9.81 μmol , 21.43% yield, 92.55% purity, TFA salt) as a light yellow solid.

[0589] LC-MS (m/z): 532.1 [M+Na]⁺

[0590] ¹H NMR (400MHz, CD₃OD) δ 7.54 (s, 1H), 7.40-7.36 (m, 2H), 7.34-7.31 (m, 1H), 7.30-7.27 (m, 2H), 5.61-5.59 (m, 1H), 4.10 (q, J=7.0 Hz, 1H), 3.94 (s, 1H), 3.88-3.85 (m, 1H), 3.16-3.15 (m, 1H), 3.06-3.02 (m, 2H), 2.98-2.92 (m, 1H), 2.77-2.71 (m, 1H), 2.68-2.61 (m, 1H), 2.55-2.49 (m, 1H), 2.44-2.38 (m, 1H), 2.16 (s, 1H), 2.01 (s, 1H), 1.75-1.74 (m, 3H), 1.55-1.48 (m, 1H), 1.30-1.22 (m, 3H), 1.12 (s, 3H), 1.06 (s, 3H), 0.92-0.89 (m, 4H).

Example 56: Synthesis Scheme of **K101-C1359**.

[0591] The scheme for synthesis of **K101-C1359** is illustrated below.



[0592] Preparation of Compound C13-59A. To a solution of **13A** (13.13 g, 28.72 mmol, 1 eq) in THF (40 mL) was added dropwise NaHMDS (1 M, 57.43 mL, 2 eq) at 0°C. The mixture was stirred at 0°C for 0.5°C, followed by addition of 4-(trifluoromethyl)benzaldehyde (5 g, 28.72 mmol, 3.85 mL, 1 eq) in THF (20 mL) at 0°C. The mixture was allowed to stir at 10°C for 15.5hr to give a yellow suspension. LC-MS and TLC (eluting with: EtOAc/PE=2/1) showed the reaction was complete. The reaction mixture was quenched with H₂O (40 mL) and extracted with PE (50 mL x 3). The water layer was adjusted to pH 4 with HCl (1N) and extracted with EtOAc (50 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by a flash column (eluting with: PE/EtOAc=10% to 50%) to give **C13-59A** (5.9 g, 21.67 mmol, 75.46% yield) as a yellow solid.

[0593] Preparation of Compound C13-59B. To a solution of **C13-59A** (5.9 g, 21.67 mmol, 1 eq) in MeOH (80 mL) was added Pd-C (10%, 590 mg) under N₂. The suspension was degassed under vacuum and purged with H₂ several times. The mixture was stirred under H₂ (15psi) at 10°C for 12 hours to give a black suspension. HPLC showed the reaction was complete. The reaction mixture was filtered on Celite, and the filtrate concentrated to give **C13-59B** (5.7 g, 20.78 mmol, 95.90% yield) as a yellow solid.

[0594] Preparation of Compound C13-59C. To a solution containing LiAlH₄ (1.58 g, 41.56 mmol, 2 eq) in THF (30 mL) was added dropwise **C13-59B** (5.7 g, 20.78 mmol, 1 eq) in THF (30 mL) at 0°C. The mixture was stirred at 10°C for 16hr to give a black suspension. LC-MS showed the reaction was complete. The reaction mixture was quenched with H₂O (1.58 mL) and aqueous NaOH (1.58 mL) followed by additional H₂O (4.74 mL). The mixture was filtered and concentrated to give **C13-59C** (4.7 g, 18.06 mmol, 86.89% yield) as a colorless oil.

[0595] Preparation of Compound C13-59D. To a solution of oxalyl dichloride (4.58 g, 36.11 mmol, 3.16 mL, 2 eq) in DCM (30 mL) was added dropwise DMSO (7.05 g, 90.28 mmol, 7.05 mL, 5 eq) at -78°C. The mixture was stirred at -78°C for 0.5hr followed by addition of **C13-59C** (4.7 g, 18.06 mmol, 1 eq) in DCM (20 mL) at -78°C. Following stirring of the mixture at -78°C for 1hr, Et₃N (9.14 g, 90.28 mmol, 12.57 mL, 5 eq) was added dropwise at -78°C. The mixture was allowed to stir at 20°C for 2.5hr to give a yellow suspension. TLC (eluting with: PE/EtOAc=5/1) showed the reaction was complete. The reaction mixture was quenched with H₂O (40 mL) and extracted with DCM (50 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by column chromatography on silica gel (eluting with: PE/EtOAc=100% PE to 5/1) to give **C13-59D** (2.6 g, 10.07 mmol, 55.75% yield) as a colorless oil.

[0596] Preparation of Compound C13-59E. To a solution of **C13-59D** (2.6 g, 10.07 mmol, 1 eq) in EtOH (30 mL) were added NH₃·H₂O (14.11 g, 100.67 mmol, 15.51 mL, 25% purity, 10 eq) and TMSCN (2.00 g, 20.13 mmol, 2.52 mL, 2 eq). The mixture was stirred at 10°C for 16hr to give a yellow solution. LC-MS showed the presence of desired mass but that some reactant remained. The mixture was stirred at 10°C for an additional 70hr. LC-MS showed the reaction was complete. The reaction mixture was quenched with H₂O (30 mL) and extracted with DCM (40 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give **C13-59E** (3.1 g, crude) as a yellow oil.

[0597] Preparation of Compound C13-59F. To a solution of **C13-59E** (3.1 g, 10.90 mmol, 1 eq) in EtOH (30 mL) was added NaOH (1.31 g, 32.71 mmol, 3 eq). The mixture was stirred at 40°C for 2hr followed by the addition of H₂O (6 mL). The mixture was stirred at 90°C for 4hr to give a yellow solution. LC-MS showed the reaction was complete.

[0598] Preparation of Compound C13-59. Boc₂O (4.76 g, 21.82 mmol, 5.01 mL, 2 eq) was added to a preparation of **C13-59E** (3.31 g, 10.91 mmol, 1 eq) in THF (20 mL), and the mixture stirred at 10°C

for 16hr to give a yellow suspension. LC-MS and TLC (eluting with: EtOAc/PE= 2/1, 50 uL AcOH) showed the reaction was complete. The reaction mixture was concentrated, and the resultant residue diluted with H₂O (30 mL) and extracted with PE/MTBE (5/1, 40 mL x 3). The water layer was adjusted to pH 4 with HCl (1N) and extracted with EtOAc (50 mL x 3). The organic layers were dried over Na₂SO₄, concentrated, and the product purified by a flash column (eluting with: PE/EtOAc = 100%PE to 40%) to give **C13-59** (1.6 g, 3.97 mmol, 36.35% yield) as a yellow oil.

[0599] Preparation of Compound **K101-C1359-A**. To a solution of **K101-C20Tr-B** (50 mg, 84.64 μmol, 1 eq) in DCM (3 mL) were added **C13-59** (68.29 mg, 169.28 μmol, 2 eq), DMAP (41.36 mg, 338.55 μmol, 4 eq), HOBT (13.72 mg, 101.57 μmol, 1.2 eq) and EDC (32.45 mg, 169.28 μmol, 2 eq). The mixture was stirred at 40°C for 12hr. LC-MS showed that **K101-C20Tr-B** remained. The mixture was stirred at 40°C for an additional 16hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The mixture was combined with a second preparation of **K101-C1359-A**, and the combined mixture quenched with saturated NaHCO₃ (15 mL). Following extraction with DCM (30 mL x 3), the organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1359-A** (53 mg, 54.29 μmol, 53.46% yield) as a white solid.

[0600] Preparation of Compound **K101-C1359**. To a solution of **K101-C1359-A** (53 mg, 54.29 μmol, 1 eq) in THF (3 mL) were added TFA (1.54 g, 13.51 mmol, 1 mL, 248.76 eq) and Et₃SiH (6.31 mg, 54.29 μmol, 8.67 uL, 1 eq). The mixture was stirred at 10°C for 12hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated by N₂. The resultant residue dissolved in MeOH (20 mL), and the mixture stirred at 40 °C for 12hr. LC-MS showed the reaction was complete. The mixture was concentrated, and the product dissolved in DCM (20 mL) and adjusted to pH 8 with saturated NaHCO₃. The water layer was extracted with DCM (20 mL x 3), and the organic layers dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: DCM/MeOH=10/1) to give 13mg of **P1** and 14 mg of **P2**. The **P1** and **P2** products were purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10um; mobile phase: [A: water(0.1%TFA)-B: ACN];B%: 30%-60%, 10min) to give **K101-C135901** (10.2 mg, 13.64 μmol, 25.12% yield, 100% purity, TFA) and **K101-C135902** (9.8 mg, 12.53 μmol, 23.07% yield, 95.59% purity, TFA) also as white solids. The preparation contained by-product: [(21R,22S,23S,24R,32S,33R,34R)-33,34-dihydroxy-26-(hydroxymethyl)-21,25, 31,31-tetramethyl-27-oxo-32-tetracyclopentadeca-10(26),11(25)-dienyl]2-amino-8-[4-(difluoromethyl)phenyl]octanoate (about 1.0mg, TFA) and [(21R,22S,23S,24R,32S,33R,34R)-33, 34-dihydroxy-26-(hydroxymethyl)-21,25,31,31-tetramethyl-27-oxo-32-tetracyclopentadeca-10(26),11(25)-dienyl] 2-amino-8-[4-(difluoromethyl)phenyl]octanoate (about 0.8mg, TFA) as yellow oils.

[0601] **K101-C135901**: LC-MS (m/z): 656.2 [M+Na]⁺


[0602] **K101-C135901**: ¹H NMR (400MHz, CD₃OD) δ7.58-7.56 (m, 3H), 7.40-7.38 (m, 2H), 5.62-5.61 (m, 1H), 3.99-3.92 (m, 2H) 3.58 (s, 1H), 3.18 (s, 1H), 3.09 (s, 1H), 2.74-2.71 (m, 2H), 2.52-2.42 (m, 2H), 2.19-2.05 (m, 2H), 1.77-1.67 (m, 7H), 1.40-1.39 (m, 6H), 1.19 (s, 3H), 1.10 (s, 3H), 0.93-0.90 (m, 4H).

[0603] **K101-C135902**: LC-MS (m/z): 656.2 [M+Na]⁺

[0604] **K101-C135902**: ¹H NMR (400MHz, CD₃OD) δ7.58-7.56 (m, 3H), 7.41-7.39 (m, 2H), 5.63-5.62 (m, 1H), 3.99-3.95 (m, 2H) 3.50 (s, 1H), 3.19-3.15 (m, 2H), 3.09 (s, 1H), 2.75-2.71 (m, 2H), 2.57-2.42 (m, 2H), 2.13-2.05 (m, 1H), 1.77-1.68 (m, 7H), 1.41-1.40 (m, 6H), 1.19 (s, 3H), 1.09(s, 3H), 0.96-0.90 (m, 4H).

Example 57: Synthesis Scheme of **K101-C1361**.

[0605] The scheme for synthesis of **K101-C1361** is illustrated below.



[0606] Preparation of Compound C13-61A. To a solution of **C13A** (13.63 g, 29.81 mmol, 1 eq) in THF (50 mL) was added NaHMDS (1 M, 59.62 mL, 2 eq) at 0°C. Following stirring at 0°C for 0.5hr, 3-phenylpropanal (4 g, 29.81 mmol, 3.92 mL, 1 eq) in THF (50 mL) was added dropwise at 0°C. The mixture was allowed to stir at 10°C for 64.5hr to give a yellow suspension. LC-MS and TLC (EtOAc:PE=2:1) showed the reaction was complete. The reaction mixture was quenched with H₂O (50mL) and extracted with PE (50mL x 3). The water layer was adjusted to pH 3 with HCl(1N) and extracted with EtOAc (50mL x 3). The organic layers were dried over Na₂SO₄, concentrated, and then purified by flash column (PE/EtOAc=5% to 40%) to give **C13-61A** (5.55 g, 23.90 mmol, 80.18% yield) as a yellow oil.

[0607] Preparation of Compound C13-61B. To a solution of **K13-61A** (5.55 g, 23.90 mmol, 1 eq) was added Pd/C (550 mg) under H₂ (48.28 mg, 23.90 mmol). The suspension was degassed under vacuum and purged with H₂ several times. The mixture was stirred under H₂ (15psi) at 10°C for 16 hours to give a black suspension. HPLC showed the reaction was complete. The reaction mixture was filtered on Celite and then concentrated to give **C13-61B** (5.27 g, 22.49 mmol, 94.09% yield) as a white gum.

[0608] Preparation of Compound C13-61C. To a solution containing LiAlH₄ (1.71 g, 44.98 mmol, 2 eq) in THF (100 mL) was added dropwise the preparation of **C1361-B** (5.27 g, 22.49 mmol, 1 eq) in THF (100 mL) at 0°C. The mixture was stirred at 10°C for 16hr to give a black suspension. LC-MS showed the reaction was complete. The reaction mixture was quenched with H₂O (1.7mL) and aqueous NaOH (1.7 mL), following by additional H₂O (5.1mL). The mixture was filtered and concentrated to give **C13-61C** (3.63 g, 16.47 mmol, 73.26% yield) as a colorless oil.

[0609] Preparation of Compound C13-61D. To a solution of (COCl)₂ (3.46 g, 27.23 mmol, 2.38 mL, 2 eq) in DCM (100 mL) was added dropwise DMSO (5.32 g, 68.07 mmol, 5.32 mL, 5 eq) at -78°C. The mixture was stirred at -78°C for 0.5hr. The preparation of **C1361-C** (3 g, 13.61 mmol, 1 eq) in DCM (100 mL) was added at -78°C, and the resultant mixture stirred at -78°C for 1hr. TEA (6.89 g, 68.07 mmol, 9.48 mL, 5 eq) was added dropwise at -78°C and the mixture was allowed to stir at 20°C for 2.5hr to give a yellow suspension. TLC (eluting with: PE/EtOAc=5/1) showed the reaction was complete. The reaction mixture was quenched with H₂O (40 mL) and extracted with DCM (40 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by column chromatography on silica gel (eluting with: PE/EtOAc=100%PE to 20/1) to give **C13-61D** as a yellow oil.

[0610] Preparation of Compound C13-61E. To a solution of **C13-61D** (1.7 g, 7.79 mmol, 1 eq) in EtOH (20 mL) were added NH₃.H₂O (10.91 g, 77.86 mmol, 11.99 mL, 25% purity, 10 eq) and TMSCN (1.16 g, 11.68 mmol, 1.46 mL, 1.5 eq). The mixture was stirred at 15°C for 12hr to give a yellow solution. LC-MS showed desired mass was found, but some **C13-61D** remained unreacted.

The mixture was stirred at 15°C for an additional 18hr. LC-MS showed the reaction was complete. The reaction mixture was diluted with H₂O (20 mL) and extracted with DCM (25 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give **C13-61E** (1.6 g, 6.55 mmol, 84.09% yield) as a yellow oil. The preparation was used without further purification.

[0611] Preparation of Compound C13-61F. To a solution of **C13-61E** (270 mg, 1.10 mmol, 1 eq) in EtOH (2 mL) was added NaOH (132.57 mg, 3.31 mmol, 3 eq) and the mixture stirred at 40°C for 2hr. Following addition of H₂O (0.1 mL), the mixture was stirred at 90°C for 4hr to give a yellow solution. LC-MS showed the reaction was complete. The preparation was used without further purification.

[0612] Preparation of Compound C13-61. Boc₂O (482.26 mg, 2.21 mmol, 507.64 uL, 2 eq) was added to the preparation of **C13-61F** (290.99 mg, 1.10 mmol, 1 eq) in THF (5 mL), and the mixture stirred at 15°C for 4hr to give a yellow solution. LC-MS and TLC (eluting with: EtOAc/PE=2/1) showed the reaction was complete. The reaction mixture was diluted with H₂O (10 mL) and PE (10 mL x 3), and the water layer adjusted to pH 3 with HCl (1N). The mixture was extracted with EtOAc (20 mL x 3), and the organic layers dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by a flash column (eluting with: PE/EtOAc=0% to 50%) to give **C13-61** (150 mg, 412.67 µmol, 37.35% yield) as a yellow oil.

[0613] Preparation of Compound K101-C1361-A. To a solution of **K101-C20Tr-B** (70 mg, 118.49 µmol, 1 eq) in DCM (2 mL) were added **C13-61** (86.14 mg, 236.99 µmol, 2 eq), DMAP (57.90 mg, 473.98 µmol, 4 eq), HOBT (16.01 mg, 118.49 µmol, 1 eq) and EDC (45.43 mg, 236.99 µmol, 2 eq). The mixture was stirred at 40°C for 12hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture was quenched with saturated NaHCO₃ (10 mL) and extracted with DCM (20 mL x 3). The organic layers were dried over Na₂SO₄, concentrated, and the resultant product purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1361-A** (62 mg, 66.22 µmol, 55.89% yield) as a white solid.

[0614] Preparation of Compound K101-C1361. To a solution of **K101-C1361-A** (62 mg, 66.22 µmol, 1 eq) in THF (3 mL) were added TFA (1.54 g, 13.51 mmol, 1 mL, 203.95 eq) and Et₃SiH (7.70 mg, 66.22 µmol, 10.58 uL, 1 eq). The mixture was stirred at 15°C for 12hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated by N₂, and the resultant residue dissolved in MeOH (20 mL). The mixture was stirred at 30°C for 12hr. LC-MS showed the reaction was complete. The reaction mixture was concentrated, dissolved in DCM (30 mL) and the solution adjusted to pH 8 with saturated NaHCO₃. The organic layer was separated, dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: DCM/MeOH=10/1) to give **K101-C136101** (12.2 mg, 20.19 µmol, 30.49% yield, 98.26% purity) and **K101-C136102** (15.6 mg, 25.77 µmol, 38.91% yield, 98.09% purity) as white solids.

[0615] **K101-C136101**: LC-MS (m/z): 616.2 [M+Na]⁺


[0616] **K101-C136101**: ¹H NMR (400MHz, CD₃OD) δ7.56 (s, 1H), 7.28-7.13 (m, 5H), 5.6-5.61 (m, 1H), 3.98-3.91 (m, 2H), 3.51-3.50 (m, 1H), 3.19 (s, 1H), 3.09 (s, 1H), 2.64-2.62 (m, 2H), 2.60-2.47 (m, 2H), 2.18-2.06 (m, 2H), 1.76-1.60 (m, 8H), 1.35-1.31 (m, 10H), 1.21 (s, 3H), 1.10 (s, 3H), 0.94-0.89 (m, 4H).

[0617] **K101-C136102**: LC-MS (m/z): 616.3 [M+Na]⁺

[0618] **K101-C136102**: ¹H NMR (400MHz, CD₃OD) δ7.56 (s, 1H), 7.28-7.13 (m, 5H), 5.6-5.61 (m, 1H), 3.99-3.92 (m, 2H), 3.46-3.45 (m, 1H), 3.19 (s, 1H), 3.09 (s, 1H), 2.64-2.62 (m, 2H), 2.60-2.47 (m, 2H), 2.14-2.03 (m, 2H), 1.77-1.61 (m, 8H), 1.36-1.26 (m, 10H), 1.21 (s, 3H), 1.10 (s, 3H), 0.95-0.92 (m, 4H).

Example 58: Synthesis Scheme of **K101-C1364**.

[0619] The scheme for synthesis of **K101-C1364** is illustrated below.



[0620] Preparation of Compound **K101-C1364-A**. To a solution of **K101-C20Tr-B** (30 mg, 50.78 μmol, 1 eq) and **C13-64** (36.23 mg, 126.96 μmol, 2.5 eq) in DCM (1 mL) were added EDC (48.68 mg, 253.92 μmol, 5 eq) and DMAP (18.61 mg, 152.35 μmol, 3 eq). The mixture was stirred at 20°C for 5hr to give a pale yellow solution. The reaction was complete as detected by LC-MS. The reaction solution was combined with **K101-C20Tr-B** (5 mg), diluted with H₂O (10 mL), and extracted with DCM (10 mL x 3). The combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure to give a yellow gum. The product was purified by prep-TLC (PE/EtOAc=3/1, SiO₂) to give **K101-C1364-A** (23.6 mg, 27.50 μmol, 54.16% yield) as a yellow solid.

[0621] Preparation of Compound **K101-C1364**. To a solution of **K101-C1364-A** (23.6 mg, 27.50 μmol, 1 eq) in MeOH (0.5 mL) was added HCl/MeOH (4 M, 0.5 mL, 72.72 eq), and the mixture stirred at 20°C for 2 hours to give a yellow solution. The reaction was complete as detected by LC-


MS. The reaction solution was diluted with H₂O (2 mL), neutralized with saturated aqueous NaHCO₃ until pH 7, and the extracted with DCM (5 mL x 5). The combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure to give a yellow solid. The product was purified by prep-TLC (DCM/MeOH=10/1, SiO₂) to give **K101-C1364** (3.5 mg, 6.79 μmol, 24.68% yield, 100% purity) as a white solid.

[0622] LC-MS (m/z): 538.1 [M+Na]⁺

[0623] ¹H NMR (400MHz, CD₃OD) δ 7.55 (s, 1H), 5.60-5.61 (m, 1H), 4.62 (s, 1H), 4.00-3.88 (m, 2H), 3.26 (t, J=7.4 Hz, 1H), 3.20-3.15 (m, 1H), 3.07-3.10 (m, 1H), 2.57-2.39 (m, 2H), 2.34 (s, 3H), 2.20-2.11 (m, 1H), 2.11-2.01 (m, 1H), 1.79-1.82 (m, 1H), 1.74-1.75 (m, 4H), 1.69-1.71 (m, 3H), 1.59-1.46 (m, 3H), 1.31-1.22 (m, 3H), 1.21 (s, 3H), 1.09 (s, 3H), 0.94-0.87 (m, 5H).

Example 59: Synthesis Scheme of **K101-C1365**.

[0624] The scheme for synthesis of **K101-C1365** is illustrated below.




[0625] Preparation of Compound **K101-C1365**. To a solution of **K101-C1327** (5.00 mg, 9.97 μmol, 1 eq) in CH₃CN (0.5 mL) were added formaldehyde (8.09 mg, 99.67 μmol, 7.42 uL, 10 eq) and AcOH (598.54 ug, 9.97 μmol, 0.57 uL, 1 eq). After stirred at 20°C for 5 minutes, NaBH₃CN (3.76 mg, 59.80 μmol, 6 eq) was added in portions, and the mixture stirred at 20°C for 14hr to give a colorless solution. LC-MS showed the presence of the desired MS species. The reaction solution was quenched with saturated NaHCO₃ (10 mL) and extracted with EtOAc (10 mL x 2). The combined organic layers were washed with brine (10 mL), dried over anhydrous Na₂SO₄, filtered, and then concentrated under reduced pressure to give crude product as a yellow solid. The product was purified by prep-TLC (SiO₂, DCM: MeOH = 10:1) to give a yellow solid. The product was further purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10um; mobile phase: [A: water (0.05% HCl)-B: ACN]; B%: 20%-50%, 10min) to give **K101-C1365** (1.5 mg, 2.83 μmol, 20.29% yield) as a white solid.

[0626] LC-MS (m/z): 552.2 [M+Na]⁺

[0627] ¹H NMR (400MHz, CD₃OD) δ 7.56 (s, 1H), 5.64 (s, 1H), 4.21-4.18 (m, 1H), 3.96 (s, 2H), 3.17 (s, 1H), 3.06 (s, 1H), 2.94 (s, 6H), 2.56-2.43 (m, 3H), 2.30-2.28 (m, 1H), 1.94-1.45 (m, 15H), 1.32-1.29 (m, 9H), 1.20 (s, 3H), 1.11 (s, 3H), 0.95-0.90 (m, 4H).

Example 60: Synthesis Scheme of **K101-C1370**.

[0628] The scheme for synthesis of **K101-C1370** is illustrated below.



[0629] Preparation of Compound **K101-C1370-A**. To a solution of **K101-C20Tr-B** (30 mg, 50.78 μmol , 1 eq) and **K101-C13-70** (29.36 mg, 76.17 μmol , 1.5 eq) in DCM (1 mL) were added DMAP (3.10 mg, 25.39 μmol , 0.5 eq) and EDC (19.47 mg, 101.57 μmol , 2 eq). The mixture was stirred at 10°C for 1hr to give a yellow solution. LC-MS showed the reaction was complete. DCM (3 mL) was

added and the mixture filtered and concentrated. The product was purified by prep-TLC (SiO₂, PE: EtOAc = 2:1) to give **K101-C1370-A** (42 mg, 43.83 μmol, 86.31% yield) as a yellow oil.

[0630] Preparation of Compound **K101-C1370**. To a solution of **K101-C1370-A** (40 mg, 41.75 μmol, 1 eq) in THF (1 mL) was added TFA (3.08 g, 27.01 mmol, 2 mL, 647.06 eq). The mixture was stirred at 10°C for 2 hr to give a yellow solution. . The mixture was concentrated with N₂ to give a brown gum. The residue was diluted with MeOH (20 mL) and the mixture stirred at 15°C for 72hr to give a yellow solution. The final mixture was concentrated, and the product was purified by prep-HPLC (column: Phenomenex Gemini 150 x 25mm x 10um; mobile phase: [A: water (0.1%TFA)-B: ACN]; B%: 35%-45%, 10min) to give **K101-C137001** (6.5 mg, 8.91 μmol, TFA salt) and **K101-C137002** as white solids. **K101-C137002** was purified by prep-TLC (eluting with: EtOAc/MeOH=10/1) to give **K101-C137002** (6.1 mg, 9.91 μmol) as a white solid.

[0631] K101-C137001 LC-MS (m/z): 638.3 [M+Na]⁺


[0632] K101-C137001 ¹H NMR (400MHz, CD₃OD) δ 7.56 (s, 1H), 7.46-7.44 (m, 2H), 7.33-7.31 (m, 2H), 6.87-6.59 (m, 1H), 5.61 (s, 1H), 3.96 (s, 2H), 3.54-3.53 (m, 1H), 3.18 (s, 1H), 3.09 (s, 1H), 2.71-2.67 (m, 2H), 2.57-2.42 (m, 2H), 2.18-2.06 (m, 3H), 1.77 (s, 3H), 1.67-1.58 (m, 4H), 1.39-1.29 (m, 6H), 1.19 (s, 3H), 1.10 (s, 3H), 0.93-0.89 (m, 4H).

[0633] K101-C137002 LC-MS (m/z): 638.6 [M+Na]⁺

[0634] K101-C137002 ¹H NMR (400MHz, CD₃OD) δ 7.56 (s, 1H), 7.46-7.44 (m, 2H), 7.33-7.31 (m, 2H), 6.87-6.58 (m, 1H), 5.63 (s, 1H), 3.99 (s, 2H), 3.54-3.50 (m, 1H), 3.19 (s, 1H), 3.09 (s, 1H), 2.71-2.67 (m, 2H), 2.52-2.47 (m, 2H), 1.76-1.39 (m, 11H), 1.20 (s, 3H), 1.09 (s, 3H), 0.95-0.90 (m, 4H).

Example 61: Synthesis Scheme of **K101-C1373**.

[0635] The scheme for synthesis of **K101-C1373** is illustrated below.



[0636] Preparation of Compound C1373-B. Compound **C1373-A** (1.5 g, 9.61 mmol, 1 eq) was dissolved in H₂O (40 mL), and the mixture purged with N₂. 1-(chloromethyl)-4-fluoro-1, 4-diazoniabicyclo [2.2.2] octane ditetrafluoroborate (6.13 g, 17.29 mmol, 1.8 eq) and AgNO₃ (326.39 mg, 1.92 mmol, 323.16 μ L, 0.2 eq) were added, and the mixture stirred at 55°C for 12hr under N₂ to give a brown suspension. TLC (eluting with: EtOAc=100%) showed the reaction was complete. The reaction mixture was filtered, extracted with MTBE (50 mL x 3), and the organic layers dried over Na₂SO₄ and concentrated to give **C1373-B** (680 mg, 5.23 mmol, 54.40% yield) as a yellow solid. The preparation was used without further purification.

[0637] Preparation of Compound C1373-C. To a solution of **C1373-B** (680 mg, 5.23 mmol, 1 eq) in DCM (10 mL) were added EtOH (481.51 mg, 10.45 mmol, 611.06 μ L, 2 eq), DIEA (945.61 mg, 7.32 mmol, 1.27 mL, 1.4 eq), DMAP (127.69 mg, 1.05 mmol, 0.2 eq) and EDC (1.30 g, 6.79 mmol, 1.3 eq). The mixture was stirred at 20°C for 12hr to give a yellow solution. TLC (eluting with: PE/EtOAc=3/1) showed the reaction was complete. The reaction mixture was washed with HCl (1N, 30 mL), and extracted with DCM (30 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give **C1373-C** (560 mg, 3.54 mmol, 67.75% yield) as a yellow oil.

[0638] Preparation of Compound C1373-D. To a solution of **C1373-C** (560 mg, 3.54 mmol, 1 eq) in toluene (5 mL) was added dropwise Diisobutylaluminium hydride (DIBAL-H) (1 M, 4.25 mL, 1.2 eq) at -70°C. The mixture was stirred at -70°C for 1.5hr to give a yellow solution. TLC (eluting with: PE/EtOAc=3/1) showed the reaction was complete. The reaction mixture was quenched with MeOH (5mL), and used for next step without further purification.

[0639] Preparation of Compound C1373-E. To (2R)-2-amino-2-phenyl-ethanol (582.77 mg, 4.25 mmol, 1.2 eq) was added C1373-D (404 mg, 3.54 mmol, 1 eq), and the mixture stirred at 0°C for 3hr. TMSCN (1.05 g, 10.62 mmol, 1.33 mL, 3 eq) was added, and the mixture stirred at 20°C for 12hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction was quenched with saturated KF (20 mL) and extracted with EtOAc (20 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by a flash column (eluting with: PE/EtOAc=1/2) to give C1373-E (480 mg, 1.84 mmol, 52.09% yield) as a yellow oil.

[0640] Preparation of Compound C1373-F. To a solution of C1373-E (480 mg, 1.84 mmol, 1 eq) in DCM (10 mL) /MeOH (10 mL) was added Pb(OAc)₄ (1.23 g, 2.77 mmol, 1.5 eq). The mixture was stirred at 0°C for 15min to give a yellow solution. TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture was quenched with saturated NaHCO₃ (20 mL) and extracted with DCM (15 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give C1373-F (400 mg, 1.75 mmol, 95.03% yield) as a yellow oil.

[0641] Preparation of Compound C1373-G. Compound C1373-F (0.4 g, 1.75 mmol, 1 eq) was dissolved in HCl (6 M, 292.06 uL, 1 eq), and the mixture stirred at 100°C for 12hr to give a yellow solution. LC-MS showed the reaction was complete. The product was concentrated to give C1373-G (342 mg, 1.75 mmol, 99.77% yield, HCl) as a yellow solid.

[0642] Preparation of Compound BB-C1373. To a solution of C1373-G (342 mg, 2.15 mmol, 1 eq) in H₂O (3 mL)/dioxane (5 mL) were added NaOH (859.46 mg, 21.49 mmol, 10 eq) and Boc₂O (703.45 mg, 3.22 mmol, 740.48 uL, 1.5 eq), and the mixture stirred at 20°C for 12hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was extracted with MBTE (20 mL x 2), and the water layer extracted with EtOAc (20 mL x 3). The combined organic layers were dried over Na₂SO₄ and concentrated to give BB-C1373 (200 mg, 771.39 μmol, 35.90% yield) as a yellow solid.

[0643] Preparation of Compound K101-C1373-A. To a solution of K101-C20Tr-B (60 mg, 101.57 μmol, 1 eq) in DCM (3 mL) were added BB-C1373 (39.50 mg, 152.35 μmol, 1.5 eq), DMAP (49.63 mg, 406.27 μmol, 4 eq) and EDC (29.21 mg, 152.35 μmol, 1.5 eq). The mixture was stirred at 20°C for 12hr to give a yellow solution. LC-MS showed the desired mass was found, but some K101-C20Tr-B remained unreacted. The mixture was stirred at 20°C for an additional 12hr. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture was quenched with H₂O (10 mL) and extracted with DCM (15 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by a flash column (eluting with: PE/EtOAc=100%PE to 50%) to give K101-C1373-A (70 mg, 84.13 μmol, 82.84% yield) as a brown solid.

[0644] Preparation of Compound K101-C1373. To a solution of **K101-C1373-A** (70 mg, 84.13 μmol , 1 eq) in DCM (5 mL) was added TFA (770.00 mg, 6.75 mmol, 500.00 μL , 80.26 eq). The mixture was stirred at 20°C for 12hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated by N_2 , dissolved in MeOH (20 mL) and the mixture stirred at 20°C for 12hr. LC-MS showed the reaction was complete. The reaction mixture was concentrated by N_2 and then dissolved in DCM (20 mL) and adjusted to pH 8 with saturated NaHCO_3 (10 mL). The organic layer was separated, and the water layer extracted with DCM (20mL x 3). The combined organic layers were concentrated to give the crude product. The product was purified by prep-TLC(eluting with: EtOAc/MeOH=10/1) to give **K101-C137301** (13.6 mg, 26.76 μmol , 31.81% yield, 96.33% purity) and **K101-C137302** (11.6 mg, 22.28 μmol , 26.48% yield, 94.03% purity) as white solids.

[0645] K101-C137301 LC-MS (m/z): 512.1 $[\text{M}+\text{Na}]^+$


[0646] K101-C137301 ^1H NMR (400MHz, CDCl_3) δ 7.51 (s, 1H), 5.60-5.59 (m, 1H), 5.23 (brs, 1H), 4.00-3.91 (m, 2H), 3.67(s, 1H), 3.20 (s, 1H), 2.95 (s, 1H), 2.48-2.40 (m, 2H), 2.21-2.18 (m, 1H), 2.00-1.95 (m, 8H), 1.71 (s, 3H), 1.46-1.45 (m, 1H), 1.15 (s, 3H), 1.01 (s, 3H), 0.83-0.76 (m, 4H).

[0647] K101-C137302 LC-MS (m/z): 512.2 $[\text{M}+\text{Na}]^+$

[0648] K101-C137302 ^1H NMR (400MHz, CDCl_3) δ 7.51 (s, 1H), 5.59-5.58 (m, 1H), 5.27 (brs, 1H), 3.99-3.90 (m, 2H), 3.65(s, 1H), 3.21 (s, 1H), 2.95 (s, 1H), 2.48-2.41 (m, 2H), 2.05-1.99 (m, 9H), 1.71 (s, 3H), 1.43-1.42 (m, 1H), 1.12 (s, 3H), 1.02(s, 3H), 0.83-0.79 (m, 4H).

Example 62: Synthesis Scheme of **K101-C1375**.

[0649] The scheme for synthesis of **K101-C1375** is illustrated below.



[0650] Preparation of Compound BB-C1375. To a solution of **C1375-A** (50 mg, 203.56 μmol , 1 eq, HCl) in THF (2 mL)/H₂O (1 mL) were added NaHCO_3 (42.75 mg, 508.90 μmol , 19.79 μL , 2.5 eq) and Boc_2O (53.31 mg, 244.27 μmol , 56.12 μL , 1.2 eq) at 0°C. The mixture was stirred at 20°C for 12hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was adjusted to pH 4 with HCl (1N) and extracted with MBTE (10 mL x 3). The organic layers were

dried over Na₂SO₄ and concentrated to give the crude product. The product was washed with PE (10 mL) to give **BB-C1375** (75 mg, crude) as a white solid.

[0651] Preparation of Compound **K101-C1375-A**. To solution of **K101-C20Tr-B** (40 mg, 67.71 μ mol, 1 eq) in DCM (2 mL) were added **BB-C1375** (41.88 mg, 135.42 μ mol, 2 eq), DMAP (33.09 mg, 270.84 μ mol, 4 eq), HOBT (10.06 mg, 74.48 μ mol, 1.1 eq), and EDC (25.96 mg, 135.42 μ mol, 2 eq). The mixture was stirred at 20°C for 12hr to give a yellow solution. LC-MS showed the desired mass was found, but some **K101-C20Tr-B** remained unreacted. The mixture was stirred at 20°C for an additional 12hr. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction was quenched with H₂O (5 mL) and extracted with DCM (10 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=2/1) to give **K101-C1375-A** (44 mg, 49.89 μ mol, 73.67% yield) as a white solid.


[0652] Preparation of Compound **K101-C1375**. To a solution of **K101-C1375-A** (44 mg, 49.89 μ mol, 1 eq) in DCM (3 mL) was added TFA (770.00 mg, 6.75 mmol, 0.5 mL, 135.37 eq) at 0°C, and the mixture stir at 20°C for 12hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture was concentrated by N₂, and the resultant residue dissolved in MeOH (20 mL). The mixture was stirred at 20°C for 12hr. LC-MS showed the reaction was complete. The mixture was concentrated to give the crude product, which was dissolved in MBTE (20 mL) and washed with saturated NaHCO₃ (10 mL). The organic layer was separated, and the water layer extracted with MBTE (20 mL x 3). The combined organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by prep-TLC (eluting with: PE/EtOAc=10/1) to give **K101-C1375** (11.7 mg, 20.18 μ mol, 40.45% yield, 93.05% purity) as a white solid.

[0653] LC-MS (m/z): 562.2 [M+Na]⁺

[0654] ¹H NMR (400MHz, CD₃OD) δ 7.58 (s, 1H), 5.64-5.63 (m, 1H), 4.00-3.92 (m, 2H), 3.60 (s, 1H), 3.21(s, 1H), 3.10 (s, 1H), 2.52-2.43 (m, 2H), 2.21-2.01 (m, 8H), 1.77 (s, 3H), 1.58-1.55 (m, 1H), 1.22 (s, 3H), 1.11 (s, 3H), 0.98-0.93 (m, 4H).

Example 63: Synthesis Scheme of **K101-C134801C2003**.

[0655] The scheme for synthesis of **K101-C134801C2003** is illustrated below.



[0656] **Preparation of Compound C134801-C.** To a solution of **C1348-D** (2 g, 7.35 mmol, 1 eq) in DMA (2 mL) were added CuI (139.97 mg, 734.96 μmol , 0.1 eq) and **C134801-B** (4.06 g, 10.29 mmol, 1.4 eq) and Pd(dppf)Cl_2 (537.77 mg, 734.96 μmol , 0.1 eq) under N_2 . The mixture was stirred under N_2 at 90°C for 5hr to give a black suspension. LC-MS and TLC (eluting with: $\text{PE/EtOAc}=3/1$) showed the reaction was complete. The reaction mixture was quenched with H_2O 100 mL) and extracted with MBTE (40 mL x 3). The organic layers were dried over Na_2SO_4 and concentrated to give the crude product. The product was purified by a flash column (eluting with: $\text{PE/EtOAc}=100\%\text{PE}$ to 20%) to give **C134801-C** (750 mg, 2.16 mmol, 29.37% yield) as a yellow oil.

[0657] Preparation of Compound C134801-D. To a solution of C134801-C (0.75 g, 2.16 mmol, 1 eq) in MeOH (15 mL) was added Pd/C (500 mg, 2.16 mmol, 50% purity, 1 eq) under N₂. The suspension was degassed under vacuum and purged with H₂ several times. The mixture was stirred under H₂ (15psi) at 20°C for 12 hours. After LC-MS showed the reaction was complete, the mixture was filtered on Celite and then concentrated to give C134801-D (750 mg, 2.15 mmol, 99.42% yield) as a black oil.

[0658] Preparation of Compound BB-C134801. To a solution of C134801-D (750 mg, 2.15 mmol, 1 eq) in THF (5 mL)/H₂O (1 mL) was added LiOH.H₂O (90.06 mg, 2.15 mmol, 1 eq) at 0°C. The mixture was allowed to stir at 20°C for 12hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was adjusted to pH 4 with HCl (1N) and extracted with MBTE (20 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give BB-C134801 (700 mg, 2.09 mmol, 97.24% yield) as a yellow oil.

[0659] Preparation of Compound C134801-B. Zinc (6 g) was treated with 1N HCl aqueous (30 mL) with stirring for 10 min. The mixture was filtered and washed with water (30 mL), EtOH (30 mL) and toluene (30 mL) in sequence, and the dried in vacuum to afford the activated zinc powder. A mixture of activated Zn (2.62 g, 40.11 mmol, 4 eq) and I₂ (127.24 mg, 501.32 μmol, 100.98 uL, 0.05 eq) in DMA (10 mL) was stirred at 20°C for 5 min and C134801-A (3.3 g, 10.03 mmol, 1 eq) in DMA (10 mL) added dropwise. The reaction mixture was stirred at 20°C for 25 min to give a black suspension.

[0660] Preparation of Compound K101-C134801-A. To a solution of K101-C20Tr-B (200 mg, 338.55 μmol, 1 eq) in DCM (3 mL) were added BB-C134801 (193.06 mg, 575.54 μmol, 1.7 eq), DMAP (165.44 mg, 1.35 mmol, 4 eq), HOBt (50.32 mg, 372.41 μmol, 1.1 eq) and EDC (110.33 mg, 575.54 μmol, 1.7 eq). The mixture was stirred at 20°C for 12hr to give a yellow solution. LC-MS showed the presence of the desired mass, but some K101-C20Tr-B remained unreacted. The mixture was stirred at 20°C for an additional 12hr followed by a second 12hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture was quenched with H₂O (10 mL) and extracted with MBTE (15 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by a flash column (eluting with: PE/EtOAc=100% PE to 40%) to give K101-C134801-A (210 mg, 231.23 μmol, 68.30% yield) as a yellow solid.

[0661] Preparation of Compound K101-C134801-B. To a solution of K101-C134801-A (210 mg, 231.23 μmol, 1 eq) in MeOH (5 mL) was added HClO₄ (46.46 mg, 462.47 μmol, 27.99 uL, 2 eq) at 0°C. The mixture was stirred at 0°C for 0.5hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was quenched with saturated NaHCO₃ (10 mL) and extracted with MBTE (15 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the

crude product. The product was purified by flash column (eluting with: PE/EtOAc=100% to 50%) to give **K101-C134801-B** (120 mg, 180.22 μmol , 77.94% yield) as a yellow solid.

[0662] Preparation of Compound **K101-C134801C2003-A**. To a solution of **K101-C134801-B** (120 mg, 180.22 μmol , 1 eq) in DMF (3 mL) were added Boc-L-valine (78.31 mg, 360.44 μmol , 2 eq), DMAP (88.07 mg, 720.88 μmol , 4 eq), DIEA (46.58 mg, 360.44 μmol , 62.78 μL , 2 eq) and HATU (137.05 mg, 360.44 μmol , 2 eq). The mixture was stirred at 20°C for 12hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=1/1) showed the reaction was complete. The reaction mixture was quenched with H₂O (15 mL x 3), the combined organic layers dried over Na₂SO₄ and then concentrated to give the crude product. The product was purified by a flash column (eluting with: PE/EtOAc=100% to 40%) to give **K101-C134801C2003-A** (120 mg, 138.71 μmol , 76.97% yield) as a yellow solid.


[0663] Preparation of Compound **K101-C134801C2003**. To a solution of **K101-C134801C2003-A** (120 mg, 138.71 μmol , 1 eq) in DCM (3 mL) was added TFA (770.00 mg, 6.75 mmol, 500.00 μL , 48.68 eq). The mixture was stirred at 0°C for 3hr to give a yellow solution. LC-MS showed the desired mass was found but some **K101-C134801C2003-A** remained unreacted. The mixture was allowed to stir at 20°C for an additional 12hr. LC-MS showed the reaction was complete. The reaction mixture was concentrated by N₂ and the resultant product dissolved in DCM (20 mL). Following adjustment of the solution to pH 8 with saturated NaHCO₃, the organic layer was separated, and the water layer extracted with DCM (20 mL x 3). The combined organic layers were dried over Na₂SO₄ and concentrated to give **K101-C134801C2003** (72.6 mg, 103.89 μmol , 74.89% yield, 95.14% purity) as a white solid.

[0664] LC-MS (m/z): 665.4 [M+H]⁺

[0665] ¹H NMR (400MHz, CD₃OD): δ 7.56 (s, 1H), 7.30-7.10 (m, 5H), 5.75 (d, 1H, *J*=3.6 Hz), 4.70-4.50 (m, 3H), 3.55-3.45 (m, 1H), 3.20-3.10 (m, 2H), 2.70-2.40 (m, 4H), 2.20-1.95 (m, 3H), 1.80-1.55 (m, 8H), 1.45-1.30 (m, 6H), 1.21 (s, 3H), 1.10 (s, 3H), 1.0-0.90 (m, 10H).

Example 64: Synthesis Scheme of **K101-C134802C2003**.

[0666] The scheme for synthesis of **K101-C134802C2003** is illustrated below.



[0667] Preparation of Compound C134802-D. To a solution of **C134802-C** (4.35 g, 11.02 mmol, 1.5 eq) in DMA (10 mL) were added CuI (139.97 mg, 734.96 μmol , 0.1 eq) and **C1348-D** (2 g, 7.35 mmol, 1 eq) and Pd(dppf)Cl_2 (537.77 mg, 734.96 μmol , 0.1 eq) under N_2 . The mixture was stirred under N_2 at 90°C for 12hr to give a black suspension. LC-MS and TLC (eluting with: $\text{PE/EtOAc}=4/1$) showed the reaction was complete. The reaction mixture was quenched with H_2O (200 mL) and extracted with MBTE (100 mL x 3). The organic layers were dried over Na_2SO_4 and concentrated to give the crude product. The product was purified by a flash column (eluting with: $\text{PE/EtOAc}=100\%\text{PE}$ to 10%) to give **C134802-D** (1g, 2.88 mmol, 39.16% yield) as a yellow oil.

^1H NMR (400MHz, CDCl_3) δ 7.30-7.27 (m, 1H), 7.20-7.16 (m, 1H), 5.58-5.1 (m, 1H), 5.34-5.26 (m, 1H), 5.02-5.00 (m, 2H), 4.37-4.32 (m, 1H), 3.73 (s, 3H), 2.62-2.58 (m, 2H), 2.48-2.44 (m, 2H), 2.07-2.02 (m, 2H), 1.71-1.66 (m, 2H), 1.44 (s, 9H).

[0668] Preparation of Compound C134802-E. To a solution of **C134802-D** (1 g, 2.88 mmol, 1 eq) in MeOH (20 mL) was added Pd/C (200 mg, 2.88 mmol, 50% purity, 1 eq) under N_2 . The suspension was degassed under vacuum and purged with H_2 several times. The mixture was stirred under H_2 (15psi) at 20°C for 12 hours. After LC-MS showed the reaction was complete, the mixture was filtered on Celite, washed with MeOH (60 mL), then concentrated to give **C134802-E** (800 mg, 2.29 mmol, 79.54% yield) as a yellow oil.

^1H NMR (400MHz, CDCl_3): δ 7.31-6.98 (m, 5H), 5.05-4.98 (m, 1H), 4.31-4.26 (m, 1H), 3.73 (s, 3H), 2.61-2.57 (m, 2H), 1.76-1.62 (m, 4H), 1.44 (s, 9H), 1.33-1.22 (m, 4H).

[0669] Preparation of Compound BB-C134802. To a solution of **C134802-E** (700 mg, 2.00 mmol, 1 eq) in THF (10 mL)/ H_2O (1.4 mL) was added $\text{LiOH}\cdot\text{H}_2\text{O}$ (84.06 mg, 2.00 mmol, 1 eq) at 0°C . The mixture was allowed to stir at 20°C for 12hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was extracted with MBTE (20 mL). The water layer was acidified to pH=3 with HCl (0.5N) and extracted with EtOAc(30 mL x 3). The organic layers were dried over Na_2SO_4 and concentrated to give **BB-C134802** (560 mg, 1.61 mmol, 80.54% yield, 96.63% purity, 98.7% ee%) as a yellow oil.

^1H NMR (400MHz, CDCl_3): δ 7.28-7.16 (m, 5H), 7.05-7.03 (m, 1H), 3.87-3.81 (m, 1H), 2.57-2.55 (m, 2H), 1.61-1.53 (m, 4H), 1.37 (s, 9H), 1.32-1.18 (m, 6H).

[0670] Preparation of Compound C134802-C. Zinc (6 g) was treated with 1N HCl aqueous (30 mL) with stirring for 10 min. The mixture was filtered and washed with water (30 mL), EtOH (30 mL) and toluene (30 mL) in sequence, and the dried in vacuum to afford the activated zinc powder. A mixture of activated Zn (3.18 g, 48.61 mmol, 4 eq) and I_2 (154.23 mg, 607.66 μmol , 122.40 μL , 0.05 eq) in DMA (90 mL) was stirred at 20°C for 5 min. Then **C134802-B** (4 g, 12.15 mmol, 1 eq) in DMA (30 mL) added dropwise. The reaction mixture was stirred at 20°C for 25 min to give a black suspension and used for the next step without further purification.

[0671] Preparation of Compound K101-C134802-A. To a solution of **K101-C20Tr-B** (200 mg, 338.55 μmol , 1 eq) in DCM (5 mL) were added **BB-C134802** (136.28 mg, 406.27 μmol , 1.2 eq), DMAP (165.44 mg, 1.35 mmol, 4 eq), HOBT (48.03 mg, 355.48 μmol , 1.05 eq) and EDC (77.88 mg, 406.27 μmol , 1.2 eq). The mixture was stirred at 20°C for 12 hr to give a yellow solution. LC-MS showed the presence of the desired mass, but some K101-C20Tr-B remained unreacted. The mixture was stirred at 20°C for an additional 16 hr to give a yellow solution. LC-MS and TLC (eluting with: PE/EtOAc=2/1) showed the reaction was complete. The reaction mixture was quenched with

saturated NaHCO₃ (50 mL) and extracted with DCM (80 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The product was purified by a preparative TLC (eluting with: PE/EtOAc=2/1) to give **K101-C134802-A** (200 mg, 220.22 μmol, 65.05% yield) as a yellow solid.

¹H NMR (400MHz, CDCl₃) δ 7.60 (s, 1H), 7.47-7.44 (m, 6H), 7.33-7.28 (m, 7H), 7.24-7.20 (m, 7H), 5.62-5.61 (m, 1H), 5.14 (m, 1H), 4.25 (brs 1H), 3.54 (s, 2H), 3.29 (s, 1H), 2.95 (s, 1H), 2.64-2.60 (m, 2H), 2.52-2.40 (m, 2H), 2.07-1.99 (m, 6H), 1.80 (s, 4H), 1.64-1.60 (m, 2H), 1.46 (s, 9H), 1.37-1.29 (m, 5H), 1.23 (s, 3H), 1.007(s, 3H), 0.90-0.86 (m, 4H).

[0672] Preparation of Compound K101-C134802-B. To a solution of **K101-C134802-A** (190 mg, 209.21 μmol, 1 eq) in MeOH (5 mL) was added HClO₄ (42.03 mg, 418.42 μmol, 25.32 uL, 2 eq) at 0°C. The mixture was stirred at 0°C for 0.5hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was quenched with saturated NaHCO₃ (4 mL) and extracted with DCM (20 mL x 3). The organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The crude product was purified by a flash column (eluting with: PE/EtOAc=100% to 50%) to give **K101-C134802-B** (150 mg) as a yellow solid.

¹H NMR (400MHz, CDCl₃) δ 7.58 (s, 1H), 7.30-7.27 (m, 2H), 7.19-7.16 (m, 3H), 5.66-5.65 (m, 1H), 4.94-4.92 (m, 1H), 4.30-4.24 (m, 1H), 4.10-4.02 (m, 1H), 3.27 (s, 1H), 3.00 (s, 1H), 2.62-2.48 (m, 4H), 2.24 (s, 1H), 2.05-1.95 (m, 2H), 1.90-1.85 (m, 1H), 1.78 (s, 4H), 1.49-1.44 (m, 10H), 1.34-1.26 (m, 6H), 1.21 (s, 3H), 1.07 (s, 3H), 0.88-0.87 (m, 4H).

[0673] Preparation of Compound K101-C134802C2003-A. To a solution of **K101-C134802-B** (150 mg, 225.27 μmol, 1 eq) in DMF (5 mL) were added Boc-L-valine (97.89 mg, 450.55 μmol, 2 eq), DMAP (110.09 mg, 901.10 μmol, 4 eq), DIEA (58.23 mg, 450.55 μmol, 78.48 uL, 2 eq) and HATU (170.31 mg, 450.55 μmol, 2 eq). The mixture was stirred at 20°C for 28hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was quenched with H₂O (30 mL), the organic layer was dried over Na₂SO₄ then concentrated to give the crude product. The product was purified by a flash column (eluting with: PE/EtOAc=100% to 30%) to give **K101-C134802C2003-A** (150 mg, 173.39 μmol, 76.97% yield) as a yellow solid.

¹H NMR (400MHz, CDCl₃) δ 7.61 (s, 1H), 7.32-7.29 (m, 2H), 7.21-7.18 (m, 2H), 5.75 (m, 1H), 5.03-5.00 (m, 2H), 4.59-4.52 (m, 2H), 4.25-4.16 (m, 2H), 3.29 (s, 1H), 3.01-3.00 (m, 1H), 2.64-2.38 (m, 4H), 2.21-2.07 (m, 2H), 2.00-1.95 (m, 1H), 1.81 (s, 3H), 1.63-1.47 (m, 21H), 1.37-1.24 (m, 6H), 1.10 (s, 3H), 1.03 (s, 3H), 0.98-0.96 (m, 5H), 0.90-0.87 (m, 7H).

[0674] Preparation of Compound K101-C134802C2003. To a solution of **K101-C134802C2003-A** (150 mg, 173.39 μmol, 1 eq) in DCM (5 mL) was added TFA (770.00 mg, 6.75 mmol, 500.00 uL, 38.95 eq). The mixture was stirred at 20°C for 1hr to give a yellow solution. LC-MS showed the reaction was complete. The reaction mixture was concentrated by N₂ and the resultant product

dissolved in MTBE (20 mL). Following adjustment of the solution to pH 8 with saturated NaHCO₃, the organic layer was separated, and the water layer extracted with MTBE (20 mL x 3). The combined organic layers were dried over Na₂SO₄ and concentrated to give **K101-C134802C2003** (85.2 mg, 123.43 μmol, 71.19% yield, 96.32% purity) as a yellow solid.

[0675] LC-MS (m/z): 687.4 [M+Na]⁺

[0676] ¹H NMR (400MHz, CD₃OD): δ 7.56 (s, 1H), 7.30-7.10 (m, 5H), 5.80-5.75 (m, 1H), 4.70-4.50 (m, 3H), 3.55-3.45 (m, 1H), 3.20-3.05 (m, 2H), 2.70-2.50 (m, 3H), 2.50-2.40 (m, 1H), 2.20-1.95 (m, 3H), 1.85-1.45 (m, 8H), 1.45-1.25 (m, 6H), 1.21 (s, 3H), 1.10 (s, 3H), 1.0-0.90 (m, 10H).

Example 65: Compound Binding to C1b Domain of PKC Isoforms

[0677] Compounds described in this disclosure were tested for binding affinity to C1b domain of PKC isoforms. Compound binding affinity to C1b domain of PKC isoforms was derived from competitive binding assay using radioactive tracer ³H-PDBu and recombinant GST-C1b or full-length of human PKC proteins based on modified procedures from Methods in Molecular Biology, vol. 233: Protein Kinase C Protocols Edited by: A. C. Newton, Humana Press Inc., Totowa, NJ (2003) and Beans et al., Proc Natl Acad Sci U S A. 2013, 110(29):11698-703.

[0678] Genes encoding C1b domains of human PKCα, δ, ε, θ, and PKD1 (PKCμ) were synthesized and inserted into pGEX-2TK or pGEX-4T1 vectors for expression and production of GST-C1b proteins in *E. coli* BL21(DE3) via IPTG induction. The target proteins were purified by glutathione sepharose 4B and followed by ion exchange chromatography if needed to achieve purity >70%. Aliquots of the purified proteins were stored at -80 degree for binding assay (storage buffer: 20 mM Tris pH 8.0, 200 mM NaCl, 10% glycerol, 1 mM DTT). Full-length PKCδ was purchased from Millipore (cat# 14-504).

[0679] Competitive binding assay was performed in 96-well plate (Agilent-5042-1385). 300μL reaction contains 60μL of 1mg/mL phosphatidylserine (*Sigma-P7769*), 1μL DMSO, testing compound (10-point 3-fold or 4-fold serial dilution) or PDBu (*Sigma-P1269*) (final concentration of 5 or 10μM as the nonspecific binding control), 19μL of assay buffer (50mM Tris-HCl pH7.4, 100mM KCl, 0.15mM CaCl₂ and 0.2% BSA), 200μL of protein (final concentration of 1-10nM), and 20μL ³H-PDBu (*ARC-ART 0485*) (final concentration of 1-10nM). The plate was incubated at 37⁰C on a shaker at 300rpm for 10 minutes and then put on ice for 20 minutes. The reaction mixtures were filtered through the GF/B filter plate (*PE-6005177*) and the plate was washed with the wash buffer (20mM Tris-HCl pH7.4, stored at 4⁰C) for 6 times before drying at 50⁰C for 1 hour. The bottom of the filter plate was sealed and 50μL of MicroScint-20 (*PerkinElmer*) cocktail was added to each well. ³H-PDBu trapped in each well was counted using *PerkinElmer MicroBeta2 Reader*. Data were analyzed using *GraphPad Prism5* with the model “log(inhibitor) vs. response-variable slope” to fit the IC₅₀, and then IC₅₀ was transformed to K_i using Cheng-Prusoff equation: K_i=IC₅₀/(1+added radio

ligand/Kd). Kd is the binding affinity of each protein preparation to ³H-PDBu, determined experimentally. Unrelated protein made in similar expression system did not have specific binding to ³H-PDBu. Ki values of Diterpenoid compounds for human PKC α , δ , ϵ , θ , and PKD1 (PKC μ) were reported in **Table 3**.

[0680] As shown in **Table 3**, in general, binding affinity of new diterpenoid compounds in this disclosure to C1b domain of novel PKC isoforms or PKD1 (PKC μ) is much higher than that to conventional PKC α (up to >100-fold difference). The Ki values of the diterpenoid compounds for C1b domain of human PKC δ , α , ϵ , θ , and PKD1 (PKC μ) are as follows: A = ≤ 5 nM, B = > 5 nM and ≤ 10 nM, C = > 10 nM and ≤ 30 nM; D = > 30 nM and ≤ 60 nM; E = > 60 nM and ≤ 100 nM, F = > 100 nM and ≤ 200 nM; G = > 200 nM and ≤ 400 nM; H = > 400 nM and ≤ 800 nM; I = > 800 nM and ≤ 1000 nM; J = > 1000 nM and ≤ 2000 nM, K = > 2000 nM and ≤ 4000 nM; L = > 4000 nM and ≤ 6000 nM; M = > 6000 nM and ≤ 8000 nM; N = > 8000 nM.)

[0681] Table 3

Compound	PKC δ	PKC α	PKC θ	PKC ϵ	PKD1/PKC μ
PBDu (Phorbol 12,13-dibutyrate)	A	G	A	A	A
K101A (or K101, Prostratin)	G	M	F	F	D
K103 (TPA)	A	B	A	A	A
K101-C130H	M	N	K	K	L
K101-C1301	B	F	C	B	A
K101-C1302	A	D	A	A	A
K101-C1303	D	H	D	F	C
K101-C1304	B	F	B	C	A
K101-C1305	H	N	G	J	F
K101-C1306	F	J	F	G	C
K101-C1308	C	I	D	C	B
K101-C1311	C	H	C	D	A
K101-C1312	B	F	C	C	A
K101-C1313	C	G	C	C	A
K101-C1315	F	N	NT	NT	D
K101-C1316	F	M	NT	NT	D
K101-C1317	C	I	NT	NT	A
K101-C1318	B	F	C	D	A
K101-C1319	A	E	B	D	A
K101-C1320	D	K	NT	NT	C
K101-C1321	A	E	NT	NT	A
K101-C1322	D	I	NT	NT	B
K101-C1323	G	N	NT	NT	F
K101-C1324	A	F	A	A	A
K101-C1325	D	L	NT	NT	D
K101-C1326	H	N	NT	NT	G

K101-C1327	A	E	B	C	A
K101-C1328	C	G	NT	NT	A
K101-C1329	A	D	B	C	A
K101-C1330	E	J	NT	NT	D
K101-C1331	B	F	NT	NT	A
K101-C1332	B	F	NT	NT	B
K101-C1333	A	E	A	C	A
K101-C1334	H	N	NT	NT	D
K101-C1335	G	N	NT	NT	E
K101-C1336	F	K	NT	NT	D
K101-C1337	C	J	NT	NT	B
K101-C1338	F	K	NT	NT	D
K101-C1339	G	L	NT	NT	E
K101-C1340	C	G	NT	NT	B
K101-C1341	D	J	NT	NT	C
K101-C1342	B	F	NT	NT	A
K101-C1343	D	J	NT	NT	C
K101-C1344	D	J	NT	NT	C
K101-C1345	A	D	C	C	A
K101-C1346	C	G	D	E	B
K101-C1347	A	C	A	B	A
K101-C134801	B	H	C	C	A
K101-C134802	A	D	A	C	A
K101-C134901	C	G	C	C	B
K101-C134902	A	D	B	C	A
K101-C1350	C	H	NT	NT	A
K101-C1351	D	J	NT	NT	C
K101-C1352	G	L	NT	NT	G
K101-C1353	H	N	NT	NT	L
K101-C1354	D	J	NT	NT	C
K101-C1355	F	NT	NT	NT	D
K101-C135601	D	J	NT	NT	C
K101-C135602	C	G	NT	NT	A
K101-C1357	G	N	NT	NT	D
K101-C1358	C	G	NT	NT	B
K101-C135901	D	J	D	E	C
K101-C135902	B	E	C	C	B
K101-C136101	C	H	C	C	B
K101-C136102	C	D	A	B	A
K101-C1364	C	G	NT	NT	NT
K101-C1365	F	L	NT	NT	D
K101-C137001	C	I	D	D	C
K101-C137002	C	G	C	D	B

TPA = 12-*O*-Tetradecanoylphorbol-13-acetate (TPA), also commonly known as tetradecanoylphorbol acetate, tetradecanoyl phorbol acetate, and phorbol 12-myristate 13-acetate (PMA).

NT = Not tested

Example 66: Western Blot to Assess Activation of PKC and Downstream Target Protein by the Diterpenoid Compounds

[0682] According to the methods described in patent (WO/2017/083783) A549 lung cancer cells (~3 million cells) were seeded in 10cm tissue culture dishes (or ~1 million cells in 6cm dish) and grown overnight. Cells were then treated with different drugs at indicated concentrations for 30 minutes. Cell lysate preparation, protein quantitation, SDS-PAGE, and Western blotting procedures are described (WO/2017/083783). β -actin or Vinculin was used as loading controls. Imagequant LAS4000 (GE) was used to scan membranes if secondary antibodies anti-mouse IgG HRP conjugate or anti-rabbit IgG HRP conjugate (dilutions from 1:2000 – 1:10000) were used. For ProteinSimple Wes system, manufacture's procedure was followed (ProteinSimple 12-230 kDa Wes Separation Module) for sample preparation, loading, running, and data analysis.

[0683] Results: 1 μ M Prostratin (K101) and 0.3 μ M the diterpenoid compounds disclosed herein, such as K101-C1319, -C1321, -C1327, and -C1329, induced high levels of phosphorylation of PKC μ (detected by phospho-specific antibodies P-PKD/PKC μ (S916), Cell Signaling Technology cat#2051 and P-PKD/PKC μ (S744/748), Cell Signaling Technology cat#2054), PKC δ (detected by phospho-specific antibody P-PKC δ (T505), Cell Signaling Technology cat#9374), and one of the PKC downstream targets Erk1/2 (detected by phospho-specific antibody P-Erk1/2 (T202/Y204), Cell Signaling Technology cat#9106) in A549 cells (**FIG. 1**). Phosphorylation of these sites has been linked to acute activation or catalytic activity of the PKC isoforms and Erk1/2. High level of phosphorylation indicates strong activation. Surprisingly, K101-C1337, an enantiomer of K101-C1327, induced much lower levels of phosphorylation on these proteins, indicating that the stereochemistry of the moiety on the C12 is very important in determining the potency of the compound. Other less potent compounds, such as K101-C1303, -C1315, -C1316, and -C1336, induced phosphorylation at 3 μ M.

[0684] A subset of the diterpenoid compounds with good binding affinity to novel PKC isoforms and PKC μ was tested at two different concentrations in A549 cells to assess activation of PKC μ , PKC δ , and Erk1/2 by Western blot analysis. In addition to the phospho-specific antibodies described above, an additional phospho-specific antibody P-PKC δ (S299) (Abcam cat#133456) was also used. This antibody has been shown in the literature (Durgan et al., FEBS Lett. 2007, 581(18):3377-81, Novel phosphorylation site markers of protein kinase C delta activation) to be able to detect activation of PKC δ . K-101 (prostratin, 1 μ M) was included as reference compounds.

[0685] The results are shown in **FIGS. 2, 3, 4A, and 4B**. In summary, all of these compounds demonstrated activation of PKC μ , PKC δ , and Erk1/2 in a dose-dependent manner in the concentration

range tested. The relative strength of cellular PKC activation correlated well with that of binding affinity of these compounds to PKC isoforms.

Example 67: Effect of Compounds on CaMKii Phosphorylation in PANC1 Cell Line

[0686] Panc1 (a pancreatic cancer cell line) cells (~5-8 millions) were seeded in 10 cm dishes (or ~2 millions in 6 cm dishes) and treated next day with prostratin (K-101, 0.2 μ M and 1 μ M) or 0.02 μ M and 0.1 μ M of K101-C1347, -C134801, and -C134802 for 48 hours. Cells were then collected/lysed for Western blot analysis. Protein quantification, SDS-PAGE, and Western blot procedures were performed as described in Example 65, except that a different lysis buffer (1ml of 10X TNE [20mL 1M Tris pH7.5; 30mL 5M NaCl; 2mL 0.5M EDTA; 48mL d2H₂O], 1ml of 10% NP40, 7.7mL of dH₂O, 100 μ L of 10x Protease inhibitors, 100 μ L of 10x Phosphatase inhibitors, and 10 μ L DTT) was used. To assess the effect of diterpenoid compounds on CaMKii phosphorylation, a phospho-specific antibody P-CaMKii (T286) (Abcam, cat# 32678) was used for detection. Phosphorylation of CaMKii at T286 is a marker for activation of CaMKii kinase in the Wnt/Ca²⁺ signaling pathway when and CaM are dissociated and thus a downstream marker for inhibition of K-Ras stemness (Wang et al., Cell. 2015, 163(5):1237-51). As shown in **FIG. 5**, all compounds induced phosphorylation of CaMKii at T286. This result indicated that PKC activators activate CaMKii via inhibition of K-Ras stemness.

Example 68: Effect of Compounds on Proliferation of A549

[0687] The diterpenoid compounds were tested in A549, a non-small cell lung cancer cell line harboring a K-Ras activating mutation. Briefly, A549 cells at a density of 1,000 cells/well were seeded in 96-well plates and incubated at 37°C for 24 hours. A series of different concentrations of compound stocks (500x) were prepared by 3-fold serial dilution in DMSO. These compounds were further diluted in culture media and then added to cells so that the final DMSO concentration was not exceeding 0.25%. After 96 hours of incubation, 50 μ L of CellTiter Glo reagent (Promega) was added to each well and luminescence was measured after 10 minutes using EnVision (PerkinElmer). Luminescence from cells treated with DMSO alone was set as Max and % of inhibition was calculated as follows: Inhibition% = (Max-Sample value)/Max x 100. Data was analyzed using XL-fit software (ID Business Solutions Ltd.) and IC₅₀, relative IC₅₀, and % of top inhibition was calculated. The IC₅₀ for growth inhibition of A549 lung cancer cell line is shown in **Table 4**. Some of the compounds in this disclosure were very potent in blocking proliferation of A549 lung cancer cells at low nM range. In addition, potency of inhibiting A549 proliferation correlated well with its binding affinity to PKCs for this series of compounds. In other words, higher affinity binders had stronger inhibitory effects on A549 proliferation. This further support the notion that activation, rather than inhibition, of certain PKC isoforms is required to block proliferation of cancer cells, some of which may harbor K-Ras mutations. This is in sharp contrast to previous literature and common

belief that inhibition of PKC is necessary to block cancer cell growth (Kang, 2014, New Journal of Science, 2014:1-36). Data presented here is in agreement with recent findings that many loss-of-function but not gain-of-function mutations of PKC isoforms have been identified in many human cancers (Antal et al., 2015, Cell, 160:489-502). The IC₅₀ values of the diterpenoid compounds against A549 cell line are as follows: A = ≤ 0.001 uM, B = >0.001 uM and ≤ 0.01 uM, C = >0.01 uM, and ≤ 0.03 uM, D = >0.03 uM and ≤ 0.06 uM, E = >0.06 uM and ≤ 0.1 uM, F = > 0.1 uM and ≤ 0.2 uM, G = >0.2 uM and ≤ 0.6 uM, H = >0.6 uM and ≤ 1.0 uM, I = >1.0 uM and ≤ 2.0 uM, J = >2 uM and ≤ 5 uM, K = > 5 uM and ≤ 10 uM, L = >10 uM and ≤ 30 uM, and M = >30 uM).

[0688] Table 4

Compound	IC ₅₀ μ M (A549 Cell Line)
K101 or K101A (prostratin)	D
K103 (TPA)	A
K101-C130H	M
K101-Epoxyde	M
K101-DI-OH	M
K101-C1301	D
K101-C1302	C
K101-C1303	H
K101-C1304	D
K101-C1305	M
K101-C1306	M
K101-C1308	E
K101-C1311	F
K101-C1312	D
K101-C1313	E
K101-C1315	I
K101-C1316	H
K101-C1317	H
K101-C1318	F
K101-C1319	E
K101-C1320	G
K101-C1321	D
K101-C1322	J
K101-C1323	J
K101-C1324	C
K101-C1325	I
K101-C1326	M
K101-C1327	E
K101-C1328	G
K101-C1329	E
K101-C1329-C1	E
K101-C1330	J
K101-C1331	E
K101-C1332	F
K101-C1333	B
K101-C1334	L
K101-C1335	L

K101-C1336	I
K101-C1337	G
K101-C1338	K
K101-C1339	K
K101-C1340	L
K101-C1341	L
K101-C1342	E
K101-C1343	I
K101-C1344	I
K101-C1345	C
K101-C1346	G
K101-C1347	B
K101-C134802	B
K101-C134801	C
K101-C134902	D
K101-C134901	G
K101-C1350	H
K101-C1351	G
K101-C1352	K
K101-C1353	I
K101-C1354	G
K101-C1355	J
K101-C135602	F
K101-C135601	G
K101-C1357	J
K101-C1358	I
K101-C135902	C
K101-C135901	E
K101-C136102	B
K101-C136101	C
K101-C1364	G
K101-C1365	H
K101-C137002	F
K101-C137001	F
K101-C137302	L
K101-C137301	L
K101-C1375	E
K101-C134801C2003	D
K101-C134802C2003	C

Note: 12-*O*-Tetradecanoylphorbol-13-acetate (TPA), also commonly known as tetradecanoylphorbol acetate, tetradecanoyl phorbol acetate, and phorbol 12-myristate 13-acetate (PMA).

Example 69: Effect of Compounds on Proliferation of Multiple Cancer Cell Lines

[0689] The compounds in this disclosure were tested for their potency in blocking proliferation of a few other cancer cells lines, including K-Ras mutant pancreatic cell lines Panc2.13 and KP-4, leukemia cell line HL-60, and lymphoma cell lines Namalwa and Mino. Similar procedures as in Example 67 were followed. Initial cell numbers seeded in 96-well plates were different for different cell lines: 3000 cells/well for Panc2.13; 800-1000 cells/well for KP-4; 5000 cells/well for HL-60; 5000-10000 cells/well for Namalwa and Mino. The IC₅₀ data are shown in **Table 5** below. A ratio

ratio of (IC₅₀ of K101)/(IC₅₀ of compound) (data in parenthesis in **Table 5**) was used to normalize the compound potency from different assay batches to a common comparator K101. The IC₅₀ and Ratio Data for Various Cell Lines are as follows: A = ≤0.001 uM, B = >0.001uM and ≤ 0.01 uM, C =>0.01 uM, and ≤ 0.03 uM, D=>0.03 uM and ≤0.06 uM, E = >0.06 uM and ≤0.1 uM, F => 0.1 uM and ≤0.2 uM, G = >0.2 uM and ≤0.6 uM, H = >0.6 uM and ≤1.0 uM, I = >1.0 uM and ≤ 2.0 uM, J = >2 uM and ≤ 5 uM, K = > 5 uM and ≤10 uM, L = >10 uM and ≤ 30 uM, and M = >30 uM

[0690] **Table 5**

Compound ID	IC ₅₀ in μM (ratio*)				
	Panc2.13 (Pancreatic)	KP-4 (Pancreatic)	HL-60 (Leukemia)	Namalwa (Lymphoma)	Mino (Lymphoma)
K101	H (1)	G (1)	G (1)	E (1)	E (1)
K101-C1301	NT	NT	D (9.23)	D (1.73)	B (11)
K101-C1302	NT	NT	B(72)	B (13.8)	A (75)
K101-C1304	NT	NT	C (18)	C (3.29)	B (15.6)
K101-C1321	NT	NT	E (5.71)	D (1.15)	C (5.36)
K101-C1324	NT	NT	C (32.7)	B (8.63)	B (37.5)
K101-C1303	I (0.75)	H (0.67)	G (0.9)	G (0.19)	F (0.57)
K101-C1308	F (6.29)	F (4.18)	E (5.29)	D (1.38)	C (6.82)
K101-C1318	G (4.28)	F (3.29)	F (3.21)	E (0.93)	C (3.18)
K101-C1319	G (3.65)	F (3.54)	D (1.79)	D (1.53)	C (6)
K101-C1327	F (7.1)	F (4.18)	D (2.32)	D (2.03)	B (7.33)
K101-C1329	F (5.82)	E (4.6)	D (1.79)	D (1.6)	C (5.5)
K101-C1333	B (157)	B (135)	B (80)	B (20.9)	A (82.5)
K101-C1342	F (6.79)	E (5.75)	D (6.43)	D (1.68)	C (6.2)
K101-C1345	D (20.5)	C (20.3)	C (27.7)	B (7.42)	B (33)
K101-C1347	C (71.7)	B (48.4)	B (16.2)	B (18.2)	B (52.2)
K101-C134801	E (13.4)	D (8.52)	C (4.43)	C (4.93)	B (21.3)
K101-C134802	C (57.3)	B (50)	B (12.1)	B (16.1)	B (52.2)
K101-C134901	NT	NT	F (2.75)	F (0.49)	C (3.88)
K101-C134902	NT	NT	C (12)	C (3.5)	B (15.7)

NT = Not tested

Example 70: Sphere Formation Assay to Assess Effect of Compounds on Cancer Stemness

[0691] A group of diterpenoid compounds of the present disclosure were tested to assess their effects on blocking formation of spheres from cancer cell lines such as PANC1, a K-Ras mutant pancreatic cell line. Briefly, PANC1 cells were harvested, re-suspended as single cell suspensions, counted and seeded into Ultra Low Attachment Culture 96-well plate (Corning, Cat#3474) at 100 cells/well in 100µl of complete media (with 10% FBS) containing 2% Matrigel (Corning, Cat#354234) and DMSO or compounds. Six replicates per condition were seeded. The seeded cells were placed in the 37°C tissue culture incubator. 10µl of low serum containing media (with 0.1% FBS) was added to each well every week. Spheres formed after 3-4 weeks were counted. Sphere forming efficiency was expressed as percentage of # of spheres/# of cells seeded.

[0692] The results are shown in **FIGS. 6A-6D**. All compounds showed dose-dependent inhibition of sphere formation from PANC1 single cell suspension. K101 or K101A is prostratin. Compounds exhibiting at least 50% inhibition on the sphere forming efficiency: K101-C1347 and K101-C134802 at a concentration of ≤ 50 nM; K101-C1308, K101-C1329, K101-C1345, and K101-C134801 at a concentration of ≤ 150 nM; K101 at a concentration of ≤ 500 nM; and K101-C1346 and K101-C1319 at a concentration of ≤ 1500 nM.

Example 71: Testing Compounds in Animals – Pharmacokinetic studies

[0693] Pharmacokinetic (PK) studies were conducted for compounds described in this disclosure, and the PK study of K101-C1327 administered by intravenous injection in rats is given as an example. Briefly, K101-C1327 was dissolved in a pH 5 buffer solution at the concentrations of 2.0 mg/mL, and diluted to 0.04 and 0.08 mg/mL to achieve the doses of 0.1 and 0.2 mg/kg with a dosing volume of 2.5 mL/kg, respectively. The K101-C1327 dosing solutions were administered via intravenous bolus injection in 3 rats (Sprague Dawley). Plasma samples were taken at 1 minute, 5 minutes, 15 minutes, 30 minutes, 1 hour, 2 hours, 4 hours, 7 hours, and 24 hours. The drug concentrations at each time point were determined by LC-MS. The results of drug concentrations and pharmacokinetic parameters are presented in the **Table 6** and **Table 7**, respectively, below.

[0694] **Table 6:** K101-C1327 Concentration in plasma after bolus injection in rats

K101-C1327 Concentration (ng/mL) in Rat Plasma after Bolus Injection				
	Dose 0.1 mg/kg		Dose 0.2 mg/kg	
Time post dose (hour)	Mean (ng/mL)	SD	Mean (ng/mL)	SD
0.0167	216.0	50.0	334.0	107.0
0.0833	49.2	9.7	102.0	31.0

0.25	28.7	7.7	65.0	17.0
0.5	19.4	3.7	43.6	19.9
1	14.6	0.6	37.6	17.1
2	5.0	0.8	29.1	17.8
4	1.0	NA	6.0	NA
7	< 1	NA	< 1	NA
24	< 1	NA	< 1	NA

[0695] Table 7: PK parameters for K101-C1327 in rat plasma after bolus injection

Dose	$t_{1/2}$ (hr)	C_0 (ng/mL)	AUC_{last} (hr*ng/mL)	V_z (L/kg)	V_{ss} (L/kg)	CL (mL/min/kg)	MRT_{Inf} (hr)
0.1 mg/kg	0.857	314	48.3	2.44	1.73	32.9	0.873
0.2 mg/kg	2.29	450	119	3.25	2.77	21.5	2.93

Example 71: Tumor clearance in xenograft model by intra-tumoral (IT) injected compounds

[0696] The antitumor efficacy of the compounds cited in this disclosure was tested in a xenograft model via intra-tumoral injection route. Immunodeficient (athymic) Nu/Nu mice were implanted at one flank subcutaneously with Panc2.13 pancreatic cancer cells. When tumors grew to 50-100 mm³, each group of three mice was treated with seven daily intratumoral (IT) injections of vehicle (50% dimethyl sulfoxide (DMSO)/50% Poly(ethylene glycol) PEG400), K101-C1347 (4mg/mL, 20μl), K101-C134801 (20mg/mL, 20μl), or K101-C134802 (4mg/mL, 20μl). All groups tolerated drug treatments very well. Ulceration of tumor (and the skin covering the tumor) was observed for all compound-treated mice after one or two IT injections. Scabs formed after 1-2 weeks and the skin recovered with minimal scarring after 3-4 weeks. Tumor growth/re-growth on or near the injection site was monitored for up to 55 days. FIG. 7 shows tumor growth curves of various treatment groups. IT injection of K101-C134801 resulted in complete clearance of tumors in mice. K101-C1347 cleared tumors from two out of three mice and K101-C134802 cleared tumors from one out of three mice.


[0697] All publications, patents, patent applications and other documents cited in this application are hereby incorporated by reference in their entireties for all purposes to the same extent as if each individual publication, patent, patent application or other document were individually indicated to be incorporated by reference for all purposes.

[0698] While various specific embodiments have been illustrated and described, it will be appreciated that various changes can be made without departing from the spirit and scope of the invention(s).

CLAIMS

What is claimed is:

1. A compound of formula (I):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

A is -OH, -C(O)OR¹, or -NR¹³R^{13'};

R¹ is H or a M⁺ counterion;

R² is a C₁-C₄alkyl;


R³ is O double bonded to the ring carbon when (- - -) is a bond, or -OR^a, wherein R^a is H or -C(O)R^{a1}, wherein R^{a1} is C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R⁴ and R⁵ are each independently H or -OR^b, wherein R^b is H, C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R^{5'} and R^{6'} are H, or R^{5'} and R^{6'} form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R⁶ is OH, halo, or -OC(O)R^c, wherein R^c is -C₁-C₆alkyl, -C₁-C₆alkyl-(NR^{c1})₂ or -C₁-C₆alkylC(O)OR^k, R^{c1} is H, C₁-C₆alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R⁶ is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-

natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^{6'} and R^{7'} are H, or R^{6'} and R^{7'} form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R⁷ is H or OH;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹² is H, -OH, -OC(O)R^f, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

n is 0 or 1.

2. The compound of claim 1, wherein A is -OH.
3. The compound of claim 1, wherein A is -C(O)OR¹, wherein R¹ is H or a M⁺ counterion.
4. The compound of claim 1 having the structure of formula (II):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^2 is a C_1 - C_4 alkyl;


R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$; wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

$R^{5'}$ and $R^{6'}$ are H, or $R^{5'}$ and $R^{6'}$ form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

$R^{6'}$ and $R^{7'}$ are H, or $R^{6'}$ and $R^{7'}$ form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R^7 is H or OH;

R^9 is OR^e , wherein R^e is H, C₁-C₆alkyl, or aryl;

R^{11} is C₁-C₄alkyl;

R^{12} is H, -OH, $-OC(O)R^f$, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, $-C_0$ -C₁₂aliphatic-C₃-C₇cycloalkyl, $-C_0$ -C₁₂aliphatic-heterocycloalkyl, $-C_0$ -C₁₂aliphatic-aryl, or $-C_0$ -C₁₂aliphatic-heteroaryl;

R^{13} and $R^{13'}$ are each independently H or C₁-C₄alkyl;

R^{14} is H or OR^g ; wherein R^g is H or C₁-C₆alkyl;

R^{17} and R^{18} are each independently C₁-C₄alkyl or C₁-C₄alkyl- OR^h , wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R^{21} is H, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, 5 to 12 membered bridged bicycyl, adamantyl, or $-C(O)OR^k$, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, bridged bicycyl, or adamantyl is optionally substituted with 1 to 3 of J^1 , and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J^1 ;

R^k is H or M^+ counterion;

J^1 is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and


n is 0 or 1.

5. The compound of claim 4, having the structure of formula (II'):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof.

6. The compound of claim 4, having the structure of formula (IIa):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R² is a C₁-C₄alkyl;


R³ is O double bonded to the ring carbon when (- -) is a bond, or -OR^a, wherein R^a is H or -C(O)R^{a1}, wherein R^{a1} is C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R⁴ and R⁵ are each independently H or -OR^b, wherein R^b is H, C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R⁶ is OH, halo, or -OC(O)R^c, wherein R^c is -C₁-C₆alkyl, -C₁-C₆alkyl-(NR^{c1})₂ or -C₁-C₆alkylC(O)OR^k, R^{c1} is H, C₁-C₆alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered

heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R⁶ is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R⁷ is H or OH;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹² is H, -OH, -OC(O)R^f, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

R¹⁴ is H or OR^g, wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(Rⁱ)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₃-C₁₂cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₃-C₁₂cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₃-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkyl-C₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and


n is 0 or 1.

7. The compound of claim 6, having the structure of formula (IIa')



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof.

8. The compound of claim 4, having the structure of formula (IIb):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^2 is a C_1 - C_4 alkyl;


R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl-(NR^{c1})₂ or $-C_1$ - C_6 alkylC(O)OR^k, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered

heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R⁶ is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹² is H, -OH, -OC(O)R^f, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

R¹⁴ is H or OR^g, wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₃-C₁₂cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₃-C₁₂cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₃-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkyl-C₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

n is 0 or 1.

9. The compound of claim 6, having the structure of formula (IIc):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1-C_6$ alkyl, $-C_1-C_6$ alkyl- $(NR^{c1})_2$ or $-C_1-C_6$ alkyl $C(O)OR^k$, R^{c1} is H, C_1-C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^{12} is H, $-OH$, $-OC(O)R^f$, wherein R^f is C_1-C_{12} alkyl, C_2-C_{12} alkenyl, $-C_0-C_{12}$ aliphatic- C_3-C_7 cycloalkyl, $-C_0-C_{12}$ aliphatic-heterocycloalkyl, $-C_0-C_{12}$ aliphatic-aryl, or $-C_0-C_{12}$ aliphatic-heteroaryl;

R^{13} and $R^{13'}$ are each independently H or C_1-C_4 alkyl;

L is absent, C_1-C_{12} alkylene, or C_2-C_{12} alkenylene, wherein the C_1-C_{12} alkylene or C_2-C_{12} alkenylene is optionally substituted with C_1-C_4 alkyl;

R^{21} is H, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or $-C(O)OR^k$, wherein the C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, bridged bicyclyl, or adamantyl is

optionally substituted with 1 to 3 of J^1 , and wherein optionally 1 to 2 carbon atoms of the spiro C_5 - C_{12} cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C_1 - C_4 alkyl, or when an N atom is present an N-protecting group;


each R^i is independently C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkyl C_3 - C_7 cycloalkyl, C_0 - C_6 alkylheterocyclyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl, wherein the C_3 - C_7 cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J^1 ;

R^k is H or M^+ counterion;

J^1 is selected from OH, CN, halo, C_1 - C_4 alkyl, and halo C_1 - C_4 alkyl; and

n is 0 or 1.

10. The compound of claim 4 having the structure of formula (III):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^2 is a C_1 - C_4 alkyl;


R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

$R^{5'}$ and $R^{6'}$ are H, or $R^{5'}$ and $R^{6'}$ form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl $(-NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

$R^{6'}$ and $R^{7'}$ are H, or $R^{6'}$ and $R^{7'}$ form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R^7 is H or OH;

R^9 is OR^e , wherein R^e is H, C₁-C₆alkyl, or aryl;

R^{11} is C₁-C₄alkyl;

R^{13} and $R^{13'}$ are each independently H or C₁-C₄alkyl;

R^{14} is H or OR^g ; wherein R^g is H or C₁-C₆alkyl;

R^{17} and R^{18} are each independently C₁-C₄alkyl or C₁-C₄alkyl- OR^h , wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R^{21} is H, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, 5 to 12 membered bridged bicycyl, adamantyl, or $-C(O)OR^k$, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, bridged bicycyl, or adamantyl is optionally substituted with 1 to 3 of J^1 , and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J^1 ;

R^k is H or M^+ counterion;

J^1 is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and


n is 0 or 1.

11. The compound of claim 10 having the structure of formula (III'):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof.

12. The compound of claim 10 having the structure of formula (IIIa):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^2 is a C_1 - C_4 alkyl;


R^3 is O double bonded to the ring carbon when (- -) is a bond, or $-OR^a$; wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered

heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R⁶ is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R⁷ is H or OH;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

n is 0 or 1.

13. The compound of claim 10 having structure of formula (IIIb):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein


R^2 is a C_1 - C_4 alkyl;

R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^9 is OR^e , wherein R^e is H, C_1 - C_6 alkyl, or aryl;

R^{11} is C_1 - C_4 alkyl;

R^{13} and $R^{13'}$ are each independently H or C₁-C₄alkyl;

R^{14} is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R^{17} and R^{18} are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R^{21} is H, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

n is 0 or 1.

14. The compound of claim 10 having the structure of formula (IIIc):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R⁶ is OH, halo, or -OC(O)R^c, wherein R^c is -C₁-C₆alkyl, -C₁-C₆alkyl-(NR^{cl})₂ or -C₁-C₆alkylC(O)OR^k, R^{cl} is H, C₁-C₆alkyl, or two R^{cl} together with the N atom form a 5 to 7 membered

heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R⁶ is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R¹³ and R^{13'} are each independently H or C₁-C₄alkyl;

L is absent, C₁-C₁₂alkylene, or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with C₁-C₄alkyl;

R²¹ is H, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹, and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl, or when an N atom is present an N-protecting group;


each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

n is 0 or 1.

15. The compound of claim 14 having the structure of formula (IIIe):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1-C_6$ alkyl, $-C_1-C_6$ alkyl- $(NR^{c1})_2$ or $-C_1-C_6$ alkyl $C(O)OR^k$, R^{c1} is H, C_1-C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with the adjacent R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

L is absent, C_1-C_{12} alkylene, or C_2-C_{12} alkenylene, wherein the C_1-C_{12} alkylene or C_2-C_{12} alkenylene is optionally substituted with C_1-C_4 alkyl;

R^{21} is H, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or $-C(O)OR^k$, wherein the C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J^1 , and wherein optionally 1 to 2 carbon atoms of the spiro C_5-C_{12} cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C_1-C_4 alkyl, or when an N atom is present an N-protecting group;

each R^j is independently C_1-C_6 alkyl, C_2-C_6 alkenyl, C_0-C_6 alkyl C_3-C_7 cycloalkyl, C_0-C_6 alkylheterocyclyl, C_0-C_6 alkylaryl, or C_0-C_6 alkylheteroaryl, wherein the C_3-C_7 cycloalkyl,

heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

R^k is H or M⁺ counterion; and

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl.

16. The compound of any one of claims 1 to 15, wherein R²¹ is C₃-C₇cycloalkyl, wherein the C₃-C₇cycloalkyl is optionally substituted with 1 to 3 of J¹.

17. The compound of claim 16, wherein the C₃-C₇cycloalkyl is selected from the group consisting of cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl, wherein the C₃-C₇cycloalkyl is optionally substituted with 1 to 3 of J¹.

18. The compound of any one of claims 1 to 15, wherein R²¹ is heterocyclyl, wherein the heterocyclyl is optionally substituted with 1 to 3 of J¹.

19. The compound of claim 18, wherein the heterocyclyl is selected from the group consisting of oxiranyl, oxetanyl, azetidynyl, oxazolyl, thiazolidinyl, thiazolyl, morpholinyl, pyrrolidinonyl, pyrrolidinyl, piperidinyl, piperazinyl, 2,3-dihydrofuranyl, dihydropyranyl, tetrahydrofuranyl, tetrahydropyranyl, dihydropyridinyl, tetrahydropyridinyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, and azapanyl, wherein the heterocyclyl is optionally substituted with 1 to 3 of J¹.

20. The compound of any one of claims 1 to 15, wherein R²¹ is aryl, wherein the aryl is optionally substituted with 1 to 3 of J¹.

21. The compound of claim 20, wherein R²¹ is a phenyl or naphthyl, wherein the phenyl or naphthyl is optionally substituted with 1 to 3 of J¹.

22. The compound of any one of claims 1 to 15, wherein R²¹ is heteroaryl, wherein the heteroaryl is optionally substituted with 1 to 3 of J¹.

23. The compound of claim 22, wherein the heteroaryl is selected from the group consisting of pyrrolyl, pyrazolyl, imidazolyl, pyrazinyl, oxazolyl, isoxazolyl, thiazolyl, furyl, thienyl, pyridyl, pyrimidyl, benzoxazolyl, benzothiazolyl, purinyl, benzimidazolyl, indolyl, isoquinolyl, quinoxaliny, and quinolyl, wherein the heteroaryl is optionally substituted with 1 to 3 of J¹.

24. The compound of any one of claims 1 to 15 wherein R²¹ is adamantyl, wherein the adamantyl is optionally substituted with OH, halo, or C₁-C₄alkyl.


25. The compound of any one of claims 1 to 15, wherein

R^{21} is spiro C_5 - C_{12} cycloalkyl, wherein the spiro C_5 - C_{12} cycloalkyl has 0-2 carbon atoms replaced with 0-2 heteroatoms selected from N, O and S, and is optionally substituted with 1 to 3 of J^1 , or when an N atom is present an N-protecting group.

26. The compound of any one of claims 1 to 15, wherein

R^{21} is 5 to 12 membered bridged bicycyl, wherein the bridged bicycyl has 0-2 carbon atoms replaced with 0-2 heteroatoms selected from N, O or S, and is optionally substituted with 1 to 3 of J^1 , or when an N atom is present an N-protecting group.

27. The compound of any one of claims 1 to 15, wherein R^{21} is selected from:



wherein J^1 is OH, CN, halo, C_1 - C_4 alkyl, and halo C_1 - C_4 alkyl, and n is 0-3.


28. The compound of any one of claims 1 to 27, wherein L is C_1 - C_6 alkylene, C_3 - C_6 alkylene, or C_3 - C_{12} alkylene.

29. The compound of any one of claims 1 to 27, wherein L is C_1 - C_6 alkenylene, C_3 - C_6 alkenylene, or C_3 - C_{12} alkenylene.

30. The compound of any one of claims 1-29, wherein n is 0.

30. The compound of claim 1 selected from the group consisting of the compounds of **Table 1**, or an amino acid prodrug thereof.

31. A compound of formula (IV):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^2 is a C_1 - C_4 alkyl;


R^3 is O double bonded to the ring carbon when (- - -) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

$R^{5'}$ and $R^{6'}$ are H, or $R^{5'}$ and $R^{6'}$ form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein

each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^{6'} and R^{7'} are H, or R^{6'} and R^{7'} form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R⁷ is H or OH;

R⁹ is OR^e, wherein R^e is H, C₁-C₆alkyl, or aryl;

R¹¹ is C₁-C₄alkyl;

R¹² is H, -OH, -OC(O)R^f, wherein R^f is C₁-C₁₂alkyl, C₂-C₁₂alkenyl, -C₀-C₁₂aliphatic-C₃-C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is C₀-C₆alkylarylene, C₀-C₆alkylheteroarylene, C₀-C₆alkylC₃-C₇cycloalkylene, C₁-C₁₂alkylene or C₂-C₁₂alkenylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylene is optionally substituted with OH or C₁-C₄alkyl; and


R²¹ is H, -OH, -SH, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J¹; and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl or, when an N atom is present an N-protecting group;

each R^j is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

R^k is H or M⁺ counterion.

32. The compound of claim 31, having the structure of formula (IVa):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein


R^2 is a C_1 - C_4 alkyl;

R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^7 is H or OH;

R^9 is OR^e , wherein R^e is H, C_1 - C_6 alkyl, or aryl;

R^{11} is C_1 - C_4 alkyl;

R^{12} is H, $-OH$, $-OC(O)R^f$, wherein R^f is C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, $-C_0$ - C_{12} aliphatic- C_3 -

C₇cycloalkyl, -C₀-C₁₂aliphatic-heterocycloalkyl, -C₀-C₁₂aliphatic-aryl, or -C₀-C₁₂aliphatic-heteroaryl;

R¹⁴ is H or OR^g; wherein R^g is H or C₁-C₆alkyl;

R¹⁷ and R¹⁸ are each independently C₁-C₄alkyl or C₁-C₄alkyl-OR^h, wherein R^h is H or C₁-C₆alkyl;

L is C₀-C₆alkylarylene, C₀-C₆alkylheteroarylene, C₀-C₆alkylC₃-C₇cycloalkylene, C₁-C₁₂alkylene or C₂-C₁₂alkenylylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylylene is optionally substituted with OH or C₁-C₄alkyl; and


R²¹ is H, -OH, -SH, -S(O)₂R^j, -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, 5 to 12 membered bridged bicycyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, bridged bicycyl, or adamantyl is optionally substituted with 1 to 3 of J¹; and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl or, when an N atom is present an N-protecting group;

each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

R^k is H or M⁺ counterion.

33. The compound of claim 31, having the structure of formula (IVb)



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein


R² is a C₁-C₄alkyl;

R³ is O double bonded to the ring carbon when (---) is a bond, or -OR^a, wherein R^a is H or -C(O)R^{a1}, wherein R^{a1} is C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R⁴ and R⁵ are each independently H or -OR^b, wherein R^b is H, C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1-C_6$ alkyl, $-C_1-C_6$ alkyl- $(NR^{c1})_2$ or $-C_1-C_6$ alkyl $C(O)OR^k$, R^{c1} is H, C_1-C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^7 is H or OH;

R^9 is OR^e , wherein R^e is H, C_1-C_6 alkyl, or aryl;

R^{11} is C_1-C_4 alkyl;

R^{12} is H, $-OH$, $-OC(O)R^f$, wherein R^f is C_1-C_{12} alkyl, C_2-C_{12} alkenyl, $-C_0-C_{12}$ aliphatic- C_3-C_7 cycloalkyl, $-C_0-C_{12}$ aliphatic-heterocycloalkyl, $-C_0-C_{12}$ aliphatic-aryl, or $-C_0-C_{12}$ aliphatic-heteroaryl;

R^{14} is H or OR^g , wherein R^g is H or C_1-C_6 alkyl;

R^{17} and R^{18} are each independently C_1-C_4 alkyl or C_1-C_4 alkyl- OR^h , wherein R^h is H or C_1-C_6 alkyl;


L is C_0-C_6 alkylarylene, C_0-C_6 alkylheteroarylene, C_0-C_6 alkyl C_3-C_7 cycloalkylene, C_1-C_{12} alkylene or C_2-C_{12} alkenylene, wherein the C_1-C_{12} alkylene or C_2-C_{12} alkenylene is optionally substituted with OH or C_1-C_4 alkyl; and

R^{21} is H, $-OH$, $-SH$, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or $-C(O)OR^k$, wherein the C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5-C_{12} cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J^1 ; and wherein optionally 1 to 2 carbon atoms of the spiro C_5-C_{12} cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C_1-C_4 alkyl or, when an N atom is present an N-protecting group;

each R^j is independently C_1-C_6 alkyl, C_2-C_6 alkenyl, C_0-C_6 alkyl C_3-C_7 cycloalkyl, C_0-C_6 alkylheterocyclyl, C_0-C_6 alkylarylene, or C_0-C_6 alkylheteroarylene, wherein the C_3-C_7 cycloalkyl, heterocyclyl, alkylarylene, or heteroarylene is optionally substituted with 1 to 3 of J^1 ;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and R^k is H or M⁺ counterion.

34. The compound of claim 31, having the structure of formula (V):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R² is a C₁-C₄alkyl;


R³ is O double bonded to the ring carbon when (---) is a bond, or -OR^a; wherein R^a is H or -C(O)R^{a1}, wherein R^{a1} is C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R⁴ and R⁵ are each independently H or -OR^b, wherein R^b is H, C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl;

R^{5'} and R^{6'} are H, or R^{5'} and R^{6'} form a bond or are bonded to a common O atom to form an epoxide ring as permitted by valency;

R⁶ is OH, halo, or -OC(O)R^c, wherein R^c is -C₁-C₆alkyl, -C₁-C₆alkyl-(NR^{c1})₂ or -C₁-C₆alkylC(O)OR^k, R^{c1} is H, C₁-C₆alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R⁶ is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each

occurrence of R^B is same or different; and

p is 0, 1, or 2;

$R^{6'}$ and $R^{7'}$ are H, or $R^{6'}$ and $R^{7'}$ form a bond or are bonded to a common O atom to form an epoxide ring, as permitted by valency;

R^7 is H or OH;

R^9 is OR^e , wherein R^e is H, C₁-C₆alkyl, or aryl;

R^{11} is C₁-C₄alkyl;

R^{14} is H or OR^g ; wherein R^g is H or C₁-C₆alkyl;

R^{17} and R^{18} are each independently C₁-C₄alkyl or C₁-C₄alkyl- OR^h , wherein R^h is H or C₁-C₆alkyl;

L is C₀-C₆alkylarylene, C₀-C₆alkylheteroarylene, C₀-C₆alkylC₃-C₇cycloalkylene, C₁-C₁₂alkylene or C₂-C₁₂alkenylylene, wherein the C₁-C₁₂alkylene or C₂-C₁₂alkenylylene is optionally substituted with OH or C₁-C₄alkyl; and


R^{21} is H, -OH, -SH, -S(O)₂ R^j , -SR^j, -N(R^j)₂, -Si(R^j)₃, C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or -C(O)OR^k, wherein the C₃-C₇cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J^1 ; and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C₁-C₄alkyl or, when an N atom is present an N-protecting group;

each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J^1 ;

J^1 is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

R^k is H or M⁺ counterion.

35. The compound of claim 31, having the structure of formula (Va):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

or an enantiomer or pharmaceutically acceptable salt thereof;

wherein


R^2 is a C_1 - C_4 alkyl;

R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$; wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl- $(NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^7 is H or OH;

R^9 is OR^e , wherein R^e is H, C_1 - C_6 alkyl, or aryl;

R^{11} is C_1 - C_4 alkyl;

R^{14} is H or OR^g ; wherein R^g is H or C_1 - C_6 alkyl;

R^{17} and R^{18} are each independently C_1 - C_4 alkyl or C_1 - C_4 alkyl- OR^h , wherein R^h is H or C_1 - C_6 alkyl;

L is C_0 - C_6 alkylarylene, C_0 - C_6 alkylheteroarylene, C_0 - C_6 alkyl C_3 - C_7 cycloalkylene, C_1 - C_{12} alkylene or C_2 - C_{12} alkenylene, wherein the C_1 - C_{12} alkylene or C_2 - C_{12} alkenylene is optionally substituted with OH or C_1 - C_4 alkyl; and

R^{21} is H, $-OH$, $-SH$, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3 - C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5 - C_{12} cycloalkyl, 5 to 12 membered bridged bicycyl, adamantyl, or $-C(O)OR^k$, wherein the C_3 - C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5 - C_{12} cycloalkyl, bridged bicycyl,


or adamantyl is optionally substituted with 1 to 3 of J^1 ; and wherein optionally 1 to 2 carbon atoms of the spiro C_5 - C_{12} cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C_1 - C_4 alkyl or, when an N atom is present an N-protecting group;

each R^i is independently C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkyl C_3 - C_7 cycloalkyl, C_0 - C_6 alkylheterocyclyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl, wherein the C_3 - C_7 cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J^1 ;

J^1 is selected from OH, CN, halo, C_1 - C_4 alkyl, and halo C_1 - C_4 alkyl; and

R^k is H or M^+ counterion.

36. The compound of claim 31, having the structure of formula (Vb):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein


R^2 is a C_1 - C_4 alkyl;

R^3 is O double bonded to the ring carbon when (---) is a bond, or $-OR^a$, wherein R^a is H or $-C(O)R^{a1}$, wherein R^{a1} is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^4 and R^5 are each independently H or $-OR^b$, wherein R^b is H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkylaryl, or C_0 - C_6 alkylheteroaryl;

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 alkyl $(-NR^{c1})_2$ or $-C_1$ - C_6 alkyl $C(O)OR^k$, R^{c1} is H, C_1 - C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

R^9 is OR^e , wherein R^e is H, C_1 - C_6 alkyl, or aryl;

R^{11} is C_1 - C_4 alkyl;

R^{14} is H or OR^g ; wherein R^g is H or C_1 - C_6 alkyl;

R^{17} and R^{18} are each independently C_1 - C_4 alkyl or C_1 - C_4 alkyl- OR^h , wherein R^h is H or C_1 - C_6 alkyl;

L is C_0 - C_6 alkylarylene, C_0 - C_6 alkylheteroarylene, C_0 - C_6 alkyl C_3 - C_7 cycloalkylene, C_1 - C_{12} alkylene or C_2 - C_{12} alkenylene, wherein the C_1 - C_{12} alkylene or C_2 - C_{12} alkenylene is optionally substituted with OH or C_1 - C_4 alkyl; and


R^{21} is H, -OH, -SH, - $S(O)_2R^j$, - SR^j , - $N(R^j)_2$, - $Si(R^j)_3$, C_3 - C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5 - C_{12} cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or - $C(O)OR^k$, wherein the C_3 - C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiro C_5 - C_{12} cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J^1 ; and wherein optionally 1 to 2 carbon atoms of the spiro C_5 - C_{12} cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C_1 - C_4 alkyl or, when an N atom is present an N-protecting group;

each R^j is independently C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_0 - C_6 alkyl C_3 - C_7 cycloalkyl, C_0 - C_6 alkylheterocyclyl, C_0 - C_6 alkylarylyl, or C_0 - C_6 alkylheteroaryl, wherein the C_3 - C_7 cycloalkyl, heterocyclyl, alkylarylyl, or heteroaryl is optionally substituted with 1 to 3 of J^1 ;

J^1 is selected from OH, CN, halo, C_1 - C_4 alkyl, and halo C_1 - C_4 alkyl; and

R^k is H or M^+ counterion.


37. The compound of claim 31, having the structure formula (Vc):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;
wherein

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1-C_6$ alkyl, $-C_1-C_6$ alkyl- $(NR^{c1})_2$ or $-C_1-C_6$ alkylC(O)OR^k, R^{c1} is H, C_1-C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

L is C_0-C_6 alkylarylene, C_0-C_6 alkylheteroarylene, C_0-C_6 alkylC₃-C₇cycloalkylene, C_1-C_{12} alkylene or C_2-C_{12} alkenylene, wherein the C_1-C_{12} alkylene or C_2-C_{12} alkenylene is optionally substituted with OH or C_1-C_4 alkyl; and


R^{21} is H, -OH, -SH, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, 5 to 12 membered bridged bicyclyl, adamantyl, or $-C(O)OR^k$, wherein the C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂ cycloalkyl, bridged bicyclyl, or adamantyl is optionally substituted with 1 to 3 of J^1 ; and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicyclyl is replaced with a heteroatom selected from N, O and S, and optionally substituted with C_1-C_4 alkyl or, when an N atom is present an N-protecting group;

each R^j is independently C_1-C_6 alkyl, C_2-C_6 alkenyl, C_0-C_6 alkylC₃-C₇cycloalkyl, C_0-C_6 alkylheterocyclyl, C_0-C_6 alkylarylyl, or C_0-C_6 alkylheteroarylyl, wherein the C_3-C_7 cycloalkyl, heterocyclyl, alkylarylyl, or heteroarylyl is optionally substituted with 1 to 3 of J^1 ;

J^1 is selected from OH, CN, halo, C_1-C_4 alkyl, and halo C_1-C_4 alkyl; and

R^k is H or M^+ counterion.

38. The compound of claim 31, having the structure of formula (Vd):




or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof;

wherein

R^6 is OH, halo, or $-OC(O)R^c$, wherein R^c is $-C_1-C_6$ alkyl, $-C_1-C_6$ alkyl- $(NR^{c1})_2$ or $-C_1-C_6$ alkylC(O)OR^k, R^{c1} is H, C_1-C_6 alkyl, or two R^{c1} together with the N atom form a 5 to 7 membered heterocyclyl containing 1 to 3 heteroatoms selected from N, O, and S; or

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and

p is 0, 1, or 2;

L is C_0-C_6 alkylarylene, C_0-C_6 alkylheteroarylene, C_0-C_6 alkylC₃-C₇cycloalkylene, C_1-C_{12} alkylene or C_2-C_{12} alkenylene, wherein the C_1-C_{12} alkylene or C_2-C_{12} alkenylene is optionally substituted with OH or C_1-C_4 alkyl; and

R^{21} is H, -OH, -SH, $-S(O)_2R^j$, $-SR^j$, $-N(R^j)_2$, $-Si(R^j)_3$, C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, 5 to 12 membered bridged bicycyl, adamantyl, or $-C(O)OR^k$, wherein the C_3-C_7 cycloalkyl, heterocyclyl, aryl, heteroaryl, spiroC₅-C₁₂cycloalkyl, bridged bicycyl, or adamantyl is optionally substituted with 1 to 3 of J^1 ; and wherein optionally 1 to 2 carbon atoms of the spiroC₅-C₁₂cycloalkyl or bridged bicycyl is replaced with a heteroatom selected from N, O and S,

and optionally substituted with C₁-C₄alkyl or, when an N atom is present an N-protecting group;

each R^l is independently C₁-C₆alkyl, C₂-C₆alkenyl, C₀-C₆alkylC₃-C₇cycloalkyl, C₀-C₆alkylheterocyclyl, C₀-C₆alkylaryl, or C₀-C₆alkylheteroaryl, wherein the C₃-C₇cycloalkyl, heterocyclyl, alkylaryl, or heteroaryl is optionally substituted with 1 to 3 of J¹;

J¹ is selected from OH, CN, halo, C₁-C₄alkyl, and haloC₁-C₄alkyl; and

R^k is H or M⁺ counterion.

39. The compound of any one of claims 31 to 38, wherein

R²¹ is C₃-C₇cycloalkyl, wherein the C₃-C₇cycloalkyl is optionally substituted with 1 to 3 of J¹.

40. The compound of claim 39, wherein the C₃-C₇cycloalkyl is selected from the group consisting of cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl, wherein the C₃-C₇cycloalkyl is optionally substituted with 1 to 3 of J¹.

41. The compound of any one of claims 31 to 38, wherein

R²¹ is heterocyclyl, wherein the heterocyclyl is optionally substituted with 1 to 3 of J¹.

42. The compound of claim 41, wherein the heterocyclyl is selected from the group consisting of oxiranyl, oxetanyl, azetidynyl, oxazolyl, thiazolidinyl, thiazolyl, morpholinyl, pyrrolidinonyl, pyrrolidinyl, piperidinyl, piperazinyl, 2,3-dihydrofuranyl, dihydropyranyl, tetrahydrofuranyl, tetrahydropyranyl, dihydropyridinyl, tetrahydropyridinyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, and azapanyl, wherein the heterocyclyl is optionally substituted with 1 to 3 of J¹.

43. The compound of any one of claims 31 to 38, wherein

R²¹ is aryl, wherein the aryl is optionally substituted with 1 to 3 of J¹.

44. The compound of claim 43, wherein

R²¹ is a phenyl or naphthyl, wherein the phenyl or naphthyl is optionally substituted with 1 to 3 of J¹.

45. The compound of any one of claims 31 to 38, wherein

R²¹ is heteroaryl, wherein the heteroaryl is optionally substituted with 1 to 3 of J¹.

46. The compound of claim 45, wherein the heteroaryl is selected from the group consisting of pyrrolyl, pyrazolyl, imidazolyl, pyrazinyl, oxazolyl, isoxazolyl, thiazolyl, furyl, thienyl, pyridyl, pyrimidyl, benzoxazolyl, benzothiazolyl, purinyl, benzimidazolyl, indolyl, isoquinolyl, quinoxalinyl, and quinolyl, wherein the heteroaryl is optionally substituted with 1 to 3 of J¹.

47. The compound of any one of claims 31 to 38, wherein

R^{21} is adamantyl, wherein the adamantyl is optionally substituted with 1 to 3 of J^1 .

48. The compound of any one of claims 31 to 38, wherein

R^{21} is spiro C_5 - C_{12} cycloalkyl, wherein the spiro C_5 - C_{12} cycloalkyl has 0-2 carbon atoms replaced with 0-2 heteroatoms selected from N, O and S, and is optionally substituted with 1 to 3 of J^1 , or when an N atom is present an N-protecting group.

49. The compound of any one of claims 31 to 38, wherein

R^{21} is 5 to 12 membered bridged bicycyl, wherein the bridged bicycyl has 0-2 carbon atoms replaced with 0-2 heteroatoms selected from N, O or S, and is optionally substituted with 1 to 3 of J^1 , or when an N atom is present an N-protecting group.


50. The compound of any one of claims 31 to 49, wherein L is C_1 - C_6 alkylene, C_3 - C_6 alkylene, or C_3 - C_{12} alkylene.

51. The compound of any one of claims 31 to 49, wherein L is C_1 - C_6 alkenylene, C_3 - C_6 alkenylene, or C_3 - C_{12} alkenylene.

52. The compound of claim 31 selected from the group consisting of the compounds of **Table 2** or an amino acid prodrug thereof.

53. The compound of any one of claims 1 to 52, wherein:

R^6 is



wherein,

each occurrence of R^A is independently selected from a side chain of a natural or non-natural amino acid, wherein each occurrence of R^A is same or different;

each occurrence of R^B is independently H, or R^B together with R^A and the N atom to which it is attached form a heterocyclic ring of a natural or non-natural amino acid, wherein each occurrence of R^B is same or different; and


p is 0, 1, or 2.

54. The compound of claim 53, wherein

each R^A is independently hydrogen (glycine), methyl (alanine), propan-2-yl (valine), propan-1-yl (norvaline), 2-methylpropan-1-yl (leucine), 1-methylpropan-1-yl (isoleucine), butan-1-yl

(norleucine), phenyl (2-phenylglycine), benzyl (phenylalanine), p-hydroxybenzyl (tyrosine), indol-3-ylmethyl (tryptophan), imidazol-4-ylmethyl (histidine), hydroxymethyl (serine), 2-hydroxyethyl (homoserine), 1-hydroxyethyl (threonine), mercaptomethyl (cysteine), methylthiomethyl (S-methylcysteine), 2-mercaptoethyl (homocysteine), 2-methylthioethyl (methionine), carbamoylmethyl (asparagine), 2-carbamoylethyl (glutamine), carboxymethyl (aspartic acid), 2-carboxyethyl (glutamic acid), 4-aminobutan-1-yl (lysine), 4-amino-3-hydroxybutan-1-yl (hydroxylysine), 3-aminopropan-1-yl (ornithine), 3-guanidinopropan-1-yl (arginine), or 3-ureido-propan-1-yl (citrulline);

each R^B is H, or R^B together with the adjacent R_A and the N atom form a prolyl side chain:



p is 0, 1 or 2.

55. The compound of claim 54, wherein:

each R^A is independently methyl (alanine), propan-2-yl (valine), 2-methylpropan-1-yl (leucine), imidazol-4-ylmethyl (histidine), hydroxymethyl (serine), 1-hydroxyethyl (threonine), carbamoylmethyl (asparagine), 2-carbamoylethyl (glutamine), 4-aminobutan-1-yl (lysine), carboxymethyl (aspartic acid), 3-guanidinopropan-1-yl (arginine), benzyl (phenylalanine), or 4-aminobutan-1-yl (lysine);

R^B is H; and

p 0, 1, or 2.

56. The compound of claim 54, wherein

each R^A is independently propan-2-yl (valine), 2-methylpropan-1-yl (leucine), carboxymethyl (aspartic acid), benzyl (phenylalanine), or 4-aminobutan-1-yl (lysine);

each R^B is H; and

p is 0, 1, or 2.

57. The compound of claim 54, wherein p is 0.

58. The compound of claim 54, wherein p is 1.

59. The compound of claim 54, wherein p is 1;

first of R^A is propan-2-yl (valine) and second of R^A is propan-2-yl (valine); and each of R^B is H (dipeptide Val-Val); or

first of R^A is 2-methylpropan-1-yl (leucine), and second of R^A is 2-methylpropan-1-yl (leucine); and each of R^B is H (dipeptide Leu-Leu); or

first of R^A is methyl (alanine) and second of R^A is methyl (alanine); and each of R^B is H (dipeptide Ala-Ala); or

first of R^A is 4-aminobutan-1-yl (lysine); second of R^A is 4-aminobutan-1-yl (lysine); and each of R^B is H (dipeptide Lys-Lys); or

first of R^A is hydrogen; second of R^A is 4-aminobutan-1-yl, and each of R^B is H (dipeptide Gly-Lys).

60. The compound of any one of claims 53 to 59, wherein each of the α -carbon of the amino acid other than glycine is in the L or D configuration.

61. A pharmaceutical composition comprising a compound of any one of claims 1 to 60, and a pharmaceutically acceptable carrier.

62. A method of activating protein kinase C, comprising contacting a mammalian cell with an effective amount of a compound of any one of claims 1 to 60.

63. The method of claim 62, wherein the cell is a cancer cell.

64. A method of treating cancer, comprising administering to a subject in need thereof an effective amount of a compound of any one of claims 1 to 60.

65. The method of claim 64, wherein the cancer is selected from the group consisting of adrenocortical cancer, anal cancer, biliary cancer, bladder cancer, bone cancer, brain cancer, breast cancer, cervical cancer, colon cancer, endometrial cancer, esophageal cancer, head and neck cancer, intestinal cancer, liver cancer, lung cancer, oral cancer, ovarian cancer, pancreatic cancer, renal cancer, prostate cancer, salivary gland cancer, skin cancer, stomach cancer, testicular cancer, throat cancer, thyroid cancer, uterine cancer, vaginal cancer, sarcoma, and soft tissue carcinomas.

66. The method of claim 64, wherein the cancer is a hematological cancer.

67. The method of claim 66, wherein the hematological cancer is a leukemia or lymphoma.

68. The method of claim 67, wherein the hematological cancer is selected from the group consisting of acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), lymphoma (e.g., Hodgkin's lymphoma, Non-Hodgkin's lymphoma, Burkitt's lymphoma), chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), Hairy Cell chronic myelogenous leukemia (CML), and multiple myeloma.




FIG. 1




FIG. 2




FIG. 3




FIG. 4A




FIG. 4B




FIG. 5




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 6D




FIG. 7

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 20/52545

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.: 28-29, 50-51, and 53-68
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
(see supplemental sheet)

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1 and 2

- Remark on Protest**
- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
 - The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
 - No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 20/52545

A. CLASSIFICATION OF SUBJECT MATTER

IPC - A61K 36/14; C07C 49/755; C07C 61/35 (2020.01)

CPC - A61K 31/194; A61K 31/341; A61K 31/365; A61K 36/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

See Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

See Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

See Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2017/156350 A1 (K-GEN, Inc.) 14 September 2017 (14.09.2017), especially para [0005] and para [0203]- p.42-44	1 and 2
A	PubChem CID 58108372, Create Date: 19 August 2012 (19.08.2012), especially p. 2 formula	1 and 2
A	US 2011/0014699 A1 (WENDER et al.) 20 January 2011 (20.01.2011), entire document	1 and 2
A	US 2018/0311209 A1 (ALKON) 01 November 2018 (01.11.2018), entire document	1 and 2
A	US 2014/0315990 A1 (BLANCHETTE ROCKEFELLER NEUROSCIENCES INSTITUTE) 23 October 2014 (23.10.2014), entire document	1 and 2

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"D" document cited by the applicant in the international application

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

23 November 2020

Date of mailing of the international search report

17 FEB 2021

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents

P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Lee Young

Telephone No. PCT Helpdesk: 571-272-4300

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.

PCT/US 20/52545

Box III (Observations where Unity is Lacking)

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I+: Claims 1-27 and 30b directed to a compound of formula (I). The compound of formula (I) will be searched to the extent that the compound encompasses the first species wherein A is -OH; R1 is H; R2 is C1 alkyl; R3 is O double bonded to the ring carbon; R4 is H; R5 is H; R5' is H; R6' is H; R6 is OH; R7' is H; R7 is H; R9 is ORe wherein Re is H; R11 is C1 alkyl; R12 is H; R14 is H; R17 is C1 alkyl; R18 is C1 alkyl; L is absent; R21 is H; and n is 0. It is believed that claims 1 and 2 read on this first named invention, and thus these claims will be searched without fee. Applicant is invited to elect additional method(s) wherein each additional method elected will require one additional invention fee. Applicants must specify the claims that encompass any additionally elected method. Applicants must further indicate, if applicable, the claims which encompass the first named invention, if different than what was indicated above for this group. Failure to clearly identify how any paid additional invention fees are to be applied to the '+' group(s) will result in only the first claimed invention to be searched. Additionally, an exemplary election wherein different actual variables are selected is suggested. An exemplary election would be a compound of formula (I) wherein A is -NR13R13'; R1 is H; R2 is C1 alkyl; R3 is O double bonded to the ring carbon; R4 is H; R5 is H; R5' is H; R6' is H; R6 is OH; R7' is H; R7 is H; R9 is Ore wherein Re is H; R11 is C1 alkyl; R12 is H; R13 is H; R13' is H; R14 is H; R17 is C1 alkyl; R18 is C1 alkyl; L is absent; R21 is H; and n is 0 (i.e., claims 1, 4, 5, 10, and 11).

Group II: Claims 31-49 and 52 direction to a compound of formula (IV).

The group of inventions listed above do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Special Technical Features:

Group I+ includes the technical feature of a unique compound of formula (I) which is not required by any other invention of Group I+ or Group II.

Group II includes the technical feature of a unique compound of formula (IV), which is not required by Group I+.

Common technical features:

The inventions of Groups I+ share the technical feature of a compound of formula (I). However, this shared technical feature does not represent a contribution over prior art, because the shared technical feature is obvious over WO 2017/156350 A1 to K-GEN, Inc. (hereinafter K-GEN) K-GEN teaches the diterpenoid protein kinase C (PKC) activator (para [0005]- "methods of treating cancer by using a combination of a protein kinase C (PKC) activating compound, particularly a diterpenoid PKC activator" and para [0203]- "PKC activator is a compound of structural formula (PII)") compound of formula (I) wherein A is -OH; R1 is H; R2 is C1 alkyl; R3 is O double bonded to the ring carbon; R4 is H; R5 is H; R5' is H; R6' and R7' form a bond; R6 is OH; R7 is H; R9 is ORe wherein Re is H; R11 is C1 alkyl; R12 is H; R14 is H; R17 is C1 alkyl; R18 is C1 alkyl; L is absent; R21 is H; and n is 0 (para [0217]- compound of formula (PIIId) wherein R32 is H; R33 is H; R34 is H; and R31 is optionally substituted alkylcarbonyl, which is interpreted as including -OH substituted methylcarbonyl (methyl 2-hydroxyacetate moiety)), but does not disclose the specific example/embodiment of the compound. Based on K-GEN's teachings it would have been obvious to one of ordinary skill in the art to identify the specific compound through routine experimentation.

As said compound was known in the art at the time of the invention, this cannot be considered a special technical feature, that would otherwise unify the inventions of Group I+.

The inventions of Group I+ and Group II share the technical feature of a compound comprising a diterpenoid core scaffold. However, this shared technical feature does not represent a contribution over prior art, because the shared technical feature is anticipated by K-GEN. K-GEN discloses the protein kinase C (PKC) activator (para [0005]- "methods of treating cancer by using a combination of a protein kinase C (PKC) activating compound, particularly a diterpenoid PKC activator" and para [0203]- "PKC activator is a compound of structural formula (PII)") compound comprising a diterpenoid core scaffold (para [0005]- "diterpenoid", para [0201]- compound of structural formula (PI), and para [0203]- compound of structural formula (PII)).

As said compound was known in the art at the time of the invention, this cannot be considered a special technical feature, that would otherwise unify the inventions of Group I+ and Group II.

The inventions of Groups I+ and II thus lack unity under PCT Rule 13.

NOTES:

There are two claim 30. For the purposes of completing this ISR, the first and the second claim respectively assumed as claim 30a and 30b respectively.

Claims 28-29, 30a, 50-51, and 53-68 are determined unsearchable because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Regarding claim 30b, the claim appears to contain a typographical error in the phrase "the compounds of Table 1" given that Table 1 does not list any compounds. For the purposes of the ISR we are interpreting the claim as intending to read "the compounds of Table 2".