

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 February 2008 (21.02.2008)

PCT

(10) International Publication Number
WO 2008/021933 A2

(51) International Patent Classification:

C07D 231/08 (2006.01) A61K 31/4155 (2006.01)
C07D 231/22 (2006.01) A61P 27/06 (2006.01)
C07D 409/06 (2006.01) A61P 17/14 (2006.01)

(21) International Application Number:

PCT/US2007/075550

(22) International Filing Date: 9 August 2007 (09.08.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/821,994 10 August 2006 (10.08.2006) US

(71) Applicant (for all designated States except US): **ALLERGAN, INC.** [US/US]; 2525 Dupont Drive, Irvine, CA 92612 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **OLD, David, W.** [US/US]; 13771 Typee Way, Irvine, CA 92620 (US).
DINH, Danny, T. [US/US]; 11531 College Avenue, Garden Gorge, CA 92840 (US).

(74) Agents: **JOHNSON, Brent, A.** et al.; C/o Allergan, Inc., 2525 Dupont Drive, Irvine, CA 92612 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

(54) Title: THERAPEUTIC N-ARYL OR N-HETEROARYL PYRAZOLIDINE AND PYRAZOLIDINONE DERIVATIVES




(57) Abstract: A compound having a structure and therapeutic methods, compositions, and medicaments related thereto are disclosed herein.

WO 2008/021933 A2

THERAPEUTIC N-ARYL OR N-HETEROARYL PYRAZOLIDINE AND PYRAZOLIDINONE DERIVATIVES

DESCRIPTION OF THE INVENTION

Disclosed herein is a compound having a structure



or a pharmaceutically acceptable salt thereof, or a prodrug thereof;

wherein a dashed line represents the presence or absence of a bond;

Y is an organic acid functional group, or an amide or ester thereof comprising up to 14 carbon atoms; or Y is hydroxymethyl or an ether thereof comprising up to 14 carbon atoms; or Y is a tetrazolyl functional group;

A is $-(CH_2)_6-$, *cis* $-CH_2CH=CH-(CH_2)_3-$, or $-CH_2C\equiv C-(CH_2)_3-$, wherein 1 or 2 carbon atoms may be replaced by S or O; or A is $-(CH_2)_m-Ar-(CH_2)_o-$ wherein Ar is interarylene or heterointerarylene, the sum of m and o is 1, 2, 3, or 4, and wherein 1 $-CH_2-$ may be replaced by S or O, and 1 $-CH_2-CH_2-$ may be replaced by $-CH=CH-$ or $-C\equiv C-$;


U^1 and U^2 are independently hydrogen, O, or S;

J^1 is H; O; OH; O-alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms; alkyl having 1, 2, 3, 4, 5, or 6 carbon atoms; F; Cl; Br; I;

CN; or CF_3 ; and

B is substituted aryl or substituted heteroaryl.

Also disclosed herein is a carboxylic acid or a bioisostere thereof, said carboxylic acid having a structure



or a pharmaceutically acceptable salt thereof, or a prodrug thereof,;

wherein a dashed line represents the presence or absence of a bond;

A is $-(CH_2)_6-$, *cis* $-CH_2CH=CH-(CH_2)_3-$, or $-CH_2C\equiv C-(CH_2)_3-$, wherein 1 or 2 carbon atoms may be replaced by S or O; or A is $-(CH_2)_m-Ar-(CH_2)_o-$ wherein Ar is interarylene or heterointerarylene, the sum of m and o is 1, 2, 3, or 4, and wherein 1 $-CH_2-$ may be replaced by S or O, and 1 $-CH_2-CH_2-$ may be replaced by $-CH=CH-$ or $-C\equiv C-$;

U^1 and U^2 are independently hydrogen, O, or S;

J^1 is H; O; OH; O-alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms; alkyl having 1, 2, 3, 4, 5, or 6 carbon atoms; F; Cl; Br; I;

CN; or CF_3 ; and

B is substituted aryl or substituted heteroaryl.


"Bioisosteres are substituents or groups that have chemical or physical similarities, and which produce broadly similar biological properties." Silverman, Richard B., The Organic Chemistry of Drug Design and Drug Action, 2nd Edition, Amsterdam: Elsevier Academic Press, 2004, p. 29.

5 While not intending to be limiting, organic acid functional groups are bioisosteres of carboxylic acids. An organic acid functional group is an acidic functional group on an organic molecule. While not intending to be limiting, organic acid functional groups may comprise an oxide of carbon, sulfur, or phosphorous. Thus, while not intending to limit the scope of the invention in any way, in certain compounds Y is a carboxylic acid, sulfonic acid, or phosphonic acid functional group.

10 Additionally, an amide or ester of one of the organic acids mentioned above comprising up to 14 carbon atoms is also contemplated. In an ester, a hydrocarbyl moiety replaces a hydrogen atom of an acid such as in a carboxylic acid ester, e.g. CO₂Me, CO₂Et, etc.

In an amide, an amine group replaces an OH of the acid. Examples of amides include CON(R²)₂, CON(OR²)R², CON(CH₂CH₂OH)₂, and CONH(CH₂CH₂OH) where R² is independently H, C₁-C₆ alkyl, phenyl, or
15 biphenyl. Moieties such as CONHSO₂R² are also amides of the carboxylic acid notwithstanding the fact that they may also be considered to be amides of the sulfonic acid R²-SO₃H. The following amides are also specifically contemplated, CONSO₂-biphenyl, CONSO₂-phenyl, CONSO₂-heteroaryl, and CONSO₂-naphthyl. The biphenyl, phenyl, heteroaryl, or naphthyl may be substituted or unsubstituted.

Han *et. al.* (Biorganic & Medicinal Chemistry Letters 15 (2005) 3487-3490) has recently shown that the
20 groups shown below are suitable bioisosteres for a carboxylic acid. The activity of compounds with these groups in inhibiting HCV NS3 protease was comparable to or superior to similar compounds where the group is replaced by CO₂H. Thus, Y could be any group depicted below.

Carboxylic acid bioisosteres according to Han *et. al.*




While not intending to limit the scope of the invention in any way, Y may also be hydroxymethyl or an ether thereof comprising up to 14 carbon atoms. An ether is a functional group wherein a hydrogen of an hydroxyl is replaced by carbon, e.g., Y is CH₂OCH₃, CH₂OCH₂CH₃, etc. These groups are also bioisosteres of a carboxylic acid.

"Up to 14 carbon atoms" means that the entire Y moiety, including the carbonyl carbon of a carboxylic acid ester or amide, and both carbon atoms in the -CH₂O-C of an ether has 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, or 14 carbon atoms.


Finally, while not intending to limit the scope of the invention in any way, Y may be a tetrazolyl functional group.

While not intending to be limiting, examples of compounds having the identified Y are depicted below. In these examples R is H or hydrocarbyl, subject to the constraints defined herein. Each structure below represents a specific embodiment which is individually contemplated, as well as pharmaceutically acceptable salts and prodrugs of


compounds which are represented by the structures. However, other examples are possible which may not fall within the scope of the structures shown below.

**Organic Acids**

Carboxylic Acid

Esters


Carboxylic Acid Ester

Amides


Carboxylic Acid Amide




Phosponic Acid




Phosphonic Acid Ester




Phosphonic Acid Amide




Sulfonic Acid




Sulfonic Acid Ester



Sulfonic Acid Amide





Y is hydroxymethyl




Ether


- 5 A tetrazolyl functional group is another bioisostere of a carboxylic acid. An unsubstituted tetrazolyl functional group has two tautomeric forms, which can rapidly interconvert in aqueous or biological media, and are thus equivalent to one another. These tautomers are shown below.



- 10 Additionally, if R² is C₁-C₆ alkyl, phenyl, or biphenyl, other isomeric forms of the tetrazolyl functional group such as the one shown below are also possible, unsubstituted and hydrocarbyl substituted tetrazolyl up to C₁₂ are considered to be within the scope of the term "tetrazolyl."




While not intending to limit the scope of the invention in any way, in one embodiment, Y is CO_2R^2 , $\text{CON}(\text{R}^2)_2$, $\text{CON}(\text{OR}^2)\text{R}^2$, $\text{CON}(\text{CH}_2\text{CH}_2\text{OH})_2$, $\text{CONH}(\text{CH}_2\text{CH}_2\text{OH})$, CH_2OH , $\text{P}(\text{O})(\text{OH})_2$, $\text{CONHSO}_2\text{R}^2$, $\text{SO}_2\text{N}(\text{R}^2)_2$, SO_2NHR^2 ,




- 5 wherein R^2 is independently H, $\text{C}_1\text{-C}_6$ alkyl, unsubstituted phenyl, or unsubstituted biphenyl.

According to Silverman (p. 30), the moieties shown below are also bioisosteres of a carboxylic acid.


Carboxylic acid bioisosteres according to Silverman




- 10 Orlek et al. (*J. Med. Chem.* **1991**, 34, 2726-2735) described oxadiazoles as suitable bioisosteres for a carboxylic acid. These ester replacements were shown to be potent muscarinic agonists having improved metabolic stability. Oxadiazoles were also described by Anderson et al. (*Eur. J. Med. Chem.* 1996, 31, 417-425) as carboxamide replacements having improved in vivo efficacy at the benzodiazepine receptor.

Carboxylic acid bioisosteres according to Orlek *et. al.*

Kohara *et al.* (*J. Med. Chem.* **1996**, 39, 5228-5235) described acidic heterocycles as suitable bioisosteres for a tetrazole. These carboxylic acid replacements were shown to be potent angiotensin II receptor antagonists having improved metabolic stability.

Tetrazole bioisosteres according to Kohara *et. al.*


Drysdale *et al.* (*J. Med. Chem.* **1992**, 35, 2573-2581) have described carboxylic acid mimics of non-peptide CCK-B receptor antagonists. The binding affinities of many of the bioisosteres are similar to the parent carboxylic acid.

Carboxylic acid bioisosteres according to Drysdale *et. al.*


A is $-(CH_2)_6-$, *cis* $-CH_2CH=CH-(CH_2)_3-$, or $-CH_2C\equiv C-(CH_2)_3-$, wherein 1 or 2 carbon atoms may be replaced by S or O; or A is $-(CH_2)_m-Ar-(CH_2)_o-$ wherein Ar is interarylene or heterointerarylene, the sum of m and o is 1, 2, 3, or 4, and wherein 1 $-CH_2-$ may be replaced by S or O, and 1 $-CH_2-CH_2-$ may be replaced by $-CH=CH-$ or $-C\equiv C-$.

Thus, A may be $-(CH_2)_6-$, *cis* $-CH_2CH=CH-(CH_2)_3-$, or $-CH_2C\equiv C-(CH_2)_3-$.

Alternatively, A may be a group which is related to one of these three moieties in that any carbon is replaced with S or O. For example, A may be a moiety where S replaces one or two carbon atoms such as one of the following or the like.




Alternatively, A may be a moiety where O replaces one or two carbon atoms such as one of the following or the like.



5

Alternatively, A may have an O replacing one carbon atom and an S replacing another carbon atom, such as one of the following or the like.



10

Alternatively, in certain embodiments A is $-(CH_2)_m-Ar-(CH_2)_o-$ wherein Ar is interarylene or heterointerarylene, the sum of m and o is 1, 2, 3, or 4, and wherein 1 $-CH_2-$ may be replaced by S or O, and 1 $-CH_2-CH_2-$ may be replaced by $-CH=CH-$ or $-C\equiv C-$. In other words,

in one embodiment A comprises:

- 1) a) 1, 2, 3, or 4 $-\text{CH}_2-$ moieties, or
b) 0, 1 or 2 $-\text{CH}_2-$ moieties and $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$; and

2) Ar;

5 e.g. $-\text{CH}_2-\text{Ar}-$, $-(\text{CH}_2)_2-\text{Ar}-$, $-\text{CH}=\text{CH}-\text{Ar}-$, $-\text{C}\equiv\text{C}-\text{Ar}-$, $-\text{CH}_2-\text{Ar}-\text{CH}_2-$, $-\text{CH}_2\text{Ar}-(\text{CH}_2)_2-$, $-\text{CH}_2\text{Ar}-\text{CH}=\text{CH}-$, $-\text{CH}_2\text{Ar}-\text{C}\equiv\text{C}-$, $-(\text{CH}_2)_2-\text{Ar}-(\text{CH}_2)_2-$, and the like;

in another embodiment A comprises:

- 1) a) O; and 0, 1, 2, or 3 $-\text{CH}_2-$ moieties; or
b) O; and 0 or 1 $-\text{CH}_2-$ moieties and $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$; and

10 2) Ar;

e.g., $-\text{O}-\text{Ar}-$, $-\text{Ar}-\text{CH}_2-\text{O}-$, $-\text{O}-\text{Ar}-(\text{CH}_2)_2-$, $-\text{OAr}-\text{CH}=\text{CH}-$, $-\text{O}-\text{Ar}-\text{C}\equiv\text{C}-$, $-\text{O}-\text{CH}_2-\text{Ar}-$, $-\text{O}-\text{CH}_2-\text{Ar}-(\text{CH}_2)_2-$, $-\text{O}-\text{CH}_2\text{Ar}-\text{CH}=\text{CH}-$, $-\text{O}-\text{CH}_2\text{Ar}-\text{C}\equiv\text{C}-$, and the like; or

in another embodiment A comprises:

- 1) a) S; and 0, 1, 2, or 3 $-\text{CH}_2-$ moieties; or
b) S; and 0 or 1 $-\text{CH}_2-$ moieties and $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$; and

15

2) Ar;

e.g., $-\text{S}-\text{Ar}-$, $-\text{Ar}-\text{CH}_2-\text{S}-$, $-\text{S}-\text{Ar}-(\text{CH}_2)_2-$, $-\text{SAr}-\text{CH}=\text{CH}-$, $-\text{S}-\text{Ar}-\text{C}\equiv\text{C}-$, $-\text{S}-\text{CH}_2-\text{Ar}-$, $-\text{S}-\text{CH}_2-\text{Ar}-(\text{CH}_2)_2-$, $-\text{S}-\text{CH}_2\text{Ar}-\text{CH}=\text{CH}-$, $-\text{S}-\text{CH}_2\text{Ar}-\text{C}\equiv\text{C}-$, and the like.

In another embodiment, the sum of m and o is 2, 3, or 4 wherein one CH_2 may be replaced with S or O and

20 1 $-\text{CH}_2-\text{CH}_2-$ may be replaced by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$.

In another embodiment, the sum of m and o is 3 wherein one CH_2 may be replaced with S or O and 1 $-\text{CH}_2-\text{CH}_2-$ may be replaced by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$.

In another embodiment, the sum of m and o is 2 wherein one CH_2 may be replaced with S or O or 1 $-\text{CH}_2-\text{CH}_2-$ may be replaced by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$.

25 In another embodiment, the sum of m and o is 4 wherein one CH_2 may be replaced with S or O and 1 $-\text{CH}_2-\text{CH}_2-$ may be replaced by $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$.

Interarylene or heterointerarylene refers to an aryl ring or ring system or a heteroaryl ring or ring system which connects two other parts of a molecule, i.e. the two parts are bonded to the ring in two distinct ring positions.

Interarylene or heterointerarylene may be substituted or unsubstituted. Unsubstituted interarylene or


30 heterointerarylene has no substituents other than the two parts of the molecule it connects. Substituted interarylene or heterointerarylene has substituents in addition to the two parts of the molecule it connects.

In one embodiment, Ar is substituted or unsubstituted interphenylene, interthienylene, interfurylene,

interpyridinylene, interoxazolylene, and interthiazolylene. In another embodiment Ar is interphenylene (Ph). In

another embodiment A is $-(\text{CH}_2)_2-\text{Ph}-$. Substituents of Ar each have from 0 to 4 carbon atoms, from 0 to 3 oxygen
35 atoms, from 0 to 2 sulfur atoms, from 0 to 2 nitrogen atoms, from 0 to 3 fluorine atoms, from 0 to 1 chlorine atoms, from 0 to 1 bromine atoms, from 0 to 1 iodine atoms, and from 0 to 10 hydrogen atoms.

In another embodiment A is $-\text{CH}_2\text{-Ar-OCH}_2-$. In another embodiment A is $-\text{CH}_2\text{-Ph-OCH}_2-$. In another embodiment, Ph is attached at the 1 and 3 positions, otherwise known as *m*-interphenylene, such as when A has the structure shown below.




5

In another embodiment A is $-(\text{CH}_2)_6-$, *cis* $-\text{CH}_2\text{CH=CH-(CH}_2)_3-$, or $-\text{CH}_2\text{C}\equiv\text{C-(CH}_2)_3-$, wherein 1 or 2 carbon atoms may be replaced with S or O; or A is $-(\text{CH}_2)_2\text{-Ph-}$ wherein one $-\text{CH}_2-$ may be replaced with S or O.

10 In another embodiment A is $-(\text{CH}_2)_6-$, *cis* $-\text{CH}_2\text{CH=CH-(CH}_2)_3-$, or $-\text{CH}_2\text{C}\equiv\text{C-(CH}_2)_3-$, wherein 1 or 2 carbon atoms may be replaced with S or O; or A is $-(\text{CH}_2)_2\text{-Ph-}$.

In one embodiment, Ar is thienyl.

In other embodiments, A has one of the following structures.



In another embodiment A is $-\text{CH}_2\text{OCH}_2\text{Ar-}$.

15

In another embodiment A is $-\text{CH}_2\text{SCH}_2\text{Ar-}$.

In another embodiment A is $-(\text{CH}_2)_3\text{Ar-}$.

In another embodiment A is $-\text{CH}_2\text{O(CH}_2)_4-$.

In another embodiment A is $-\text{CH}_2\text{S(CH}_2)_4-$.

In another embodiment A is $-(\text{CH}_2)_6-$.

20

In another embodiment A is *cis* $-\text{CH}_2\text{CH=CH-(CH}_2)_3-$.

In another embodiment A is $-\text{CH}_2\text{C}\equiv\text{C-(CH}_2)_3-$.

In another embodiment A is $-\text{S(CH}_2)_3\text{S(CH}_2)_2-$.

In another embodiment A is $-(\text{CH}_2)_4\text{OCH}_2-$.

In another embodiment A is *cis* $-\text{CH}_2\text{CH=CH-CH}_2\text{OCH}_2-$.

In another embodiment A is $-\text{CH}_2\text{CH}=\text{CH}-\text{CH}_2\text{OCH}_2-$.

In another embodiment A is $-(\text{CH}_2)_2\text{S}(\text{CH}_2)_3-$.

In another embodiment A is $-\text{CH}_2-\text{Ph}-\text{OCH}_2-$, wherein Ph is interphenylene,.

In another embodiment A is $-\text{CH}_2-\text{mPh}-\text{OCH}_2-$, wherein mPh is *m*-interphenylene.

5 In another embodiment A is $-\text{CH}_2-\text{O}-(\text{CH}_2)_4-$.

In another embodiment A is $-\text{CH}_2-\text{O}-\text{CH}_2-\text{Ar}-$, wherein Ar is 2,5-interthienylene.

In another embodiment A is $-\text{CH}_2-\text{O}-\text{CH}_2-\text{Ar}-$, wherein Ar is 2,5-interfurylene.

In another embodiment A is (3-methylphenoxy)methyl.

In another embodiment A is (4-but-2-ynloxy)methyl.

10 In another embodiment A is 2-(2-ethylthio)thiazol-4-yl.

In another embodiment A is 2-(3-propyl)thiazol-5-yl.

In another embodiment A is 3-(methoxymethyl)phenyl.

In another embodiment A is 3-(3-propylphenyl).

In another embodiment A is 3-methylphenethyl.

15 In another embodiment A is 4-(2-ethyl)phenyl.

In another embodiment A is 4-phenethyl.

In another embodiment A is 4-methoxybutyl.

In another embodiment A is 5-(methoxymethyl)furan-2-yl .

In another embodiment A is 5-(methoxymethyl)thiophen-2-yl.

20 In another embodiment A is 5-(3-propyl)furan-2-yl.

In another embodiment A is 5-(3-propyl)thiophen-2-yl.

In another embodiment A is 6-hexyl.

In another embodiment A is (Z)-6-hex-4-enyl.

25 Interarylene or heterointerarylene refers to an aryl ring or ring system or a heteroaryl ring or ring system which connects two other parts of a molecule, i.e. the two parts are bonded to the ring in two distinct ring positions.

Interarylene or heterointerarylene may be substituted or unsubstituted. Unsubstituted interarylene or heterointerarylene has no substituents other than the two parts of the molecule it connects. Substituted interarylene or heterointerarylene has substituents in addition to the two parts of the molecule it connects.

30 In one embodiment, Ar is substituted or unsubstituted interphenylene, interthienylene, interfurylene, interpyridinylene, interoxazolylene, and interthiazolylene. In another embodiment Ar is interphenylene (Ph). In another embodiment A is $-(\text{CH}_2)_2-\text{Ph}-$. While not intending to limit scope of the invention in any way, substituents may have 4 or less heavy atoms, wherein the heavy atoms are C, N, O, S, P, F, Cl, Br, and/or I in any stable combination. Any number of hydrogen atoms required for a particular substituent will also be included. A substituent must be stable enough for the compound to be useful as described herein. In addition to the atoms listed above, a
35 substituent may also have a metal cation or any other stable cation having an atom not listed above if the substituent is acidic and the salt form is stable. For example, $-\text{OH}$ may form an $-\text{O}^-\text{Na}^+$ salt or CO_2H may form a CO_2^-K^+ salt. Any cation of the salt is not counted in the "4 or less heavy atoms." Thus, the substituent may be hydrocarbyl having up to 4 carbon atoms, including alkyl up to C_4 , alkenyl, alkynyl, and the like;

hydrocarbyloxy up to C₃;

organic acid such as CO₂H, SO₃H, P(O)(OH)₂, and the like, and salts thereof;

CF₃;

halo, such as F, Cl, or Br;

5 hydroxyl;


NH₂ and alkylamine functional groups up to C₃;

other N or S containing substituents such as CN, NO₂, and the like;

and the like.

10 In one embodiment A is -(CH₂)_m-Ph-(CH₂)_o- wherein the sum of m and o is 1, 2, or 3, and wherein one CH₂ may be replaced with S or O.

In another embodiment A is -CH₂-Ar-OCH₂-. In another embodiment A is -CH₂-Ph-OCH₂-. In another embodiment, Ph is attached at the 1 and 3 positions, otherwise known as *m*-interphenylene, such as when A has the structure shown below.




15

In another embodiment A is -(CH₂)₆-, *cis* -CH₂CH=CH-(CH₂)₃-, or -CH₂C≡C-(CH₂)₃-, wherein 1 or 2 carbon atoms may be replaced with S or O; or A is -(CH₂)₂-Ph- wherein one CH₂ may be replaced with S or O.

20 In another embodiment A is -(CH₂)₆-, *cis* -CH₂CH=CH-(CH₂)₃-, or -CH₂C≡C-(CH₂)₃-, wherein 1 or 2 carbon atoms may be replaced with S or O; or A is -(CH₂)₂-Ph-.

In other embodiments, A has one of the following structures, where Y is attached to the aromatic or heteroaromatic ring.



25

In another embodiment A is -CH₂OCH₂Ar.

In another embodiment A is -CH₂SCH₂Ar.

In another embodiment A is -(CH₂)₃Ar.

30 In another embodiment A is -CH₂O(CH₂)₄.

In another embodiment A is $-\text{CH}_2\text{S}(\text{CH}_2)_4-$.

In another embodiment A is $-(\text{CH}_2)_6-$.

In another embodiment A is *cis* $-\text{CH}_2\text{CH}=\text{CH}-(\text{CH}_2)_3-$.

In another embodiment A is $-\text{CH}_2\text{C}\equiv\text{C}-(\text{CH}_2)_3-$.

5 In another embodiment A is $-\text{S}(\text{CH}_2)_3\text{S}(\text{CH}_2)_2-$.

In another embodiment A is $-(\text{CH}_2)_4\text{OCH}_2-$.

In another embodiment A is *cis* $-\text{CH}_2\text{CH}=\text{CH}-\text{CH}_2\text{OCH}_2-$.

In another embodiment A is $-\text{CH}_2\text{CH}\equiv\text{CH}-\text{CH}_2\text{OCH}_2-$.

In another embodiment A is $-(\text{CH}_2)_2\text{S}(\text{CH}_2)_3-$.

10 In another embodiment A is $-\text{CH}_2-\text{Ph}-\text{OCH}_2-$, wherein Ph is interphenylene,.

In another embodiment A is $-\text{CH}_2-\text{mPh}-\text{OCH}_2-$, wherein mPh is *m*-interphenylene.

In another embodiment A is $-\text{CH}_2-\text{O}-(\text{CH}_2)_4-$.

In another embodiment A is $-\text{CH}_2-\text{O}-\text{CH}_2-\text{Ar}-$, wherein Ar is 2,5-interthienylene.

In another embodiment A is $-\text{CH}_2-\text{O}-\text{CH}_2-\text{Ar}-$, wherein Ar is 2,5-interfurylene.

15 In another embodiment A is (3-methylphenoxy)methyl.

In another embodiment A is (4-but-2-ynyloxy)methyl.

In another embodiment A is 2-(2-ethylthio)thiazol-4-yl.

In another embodiment A is 2-(3-propyl)thiazol-5-yl.

In another embodiment A is 3-methoxymethyl)phenyl.

20 In another embodiment A is 3-(3-propyl)phenyl.

In another embodiment A is 3-methylphenethyl.

In another embodiment A is 4-(2-ethyl)phenyl.

In another embodiment A is 4-phenethyl.

In another embodiment A is 4-methoxybutyl.

25 In another embodiment A is 5-(methoxymethyl)furan-2-yl .

In another embodiment A is 5-(methoxymethyl)thiophen-2-yl.


In another embodiment A is 5-(3-propyl)furan-2-yl.

In another embodiment A is 5-(3-propyl)thiophen-2-yl.



In another embodiment A is 6-hexyl.

30 In another embodiment A is (Z)-6-hex-4-enyl.

Compounds according to the each of the structures depicted below, and pharmaceutically acceptable salts thereof, and prodrugs thereof, are contemplated as individual embodiments. In other words, each structure represents a different embodiment.



J1 is H; O; OH; O-alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms; alkyl having 1, 2, 3, 4, 5, or 6 carbon atoms; F; Cl; Br; I; CN; or CF₃. Thus, each structure depicted below represents a compound embodiment which is individually contemplated. Pharmaceutically acceptable salts and prodrugs of compounds according to the structures below are also contemplated.




5



U1 and U2 are independently hydrogen, O, or S. Thus, each structure depicted below represents a compound embodiment which is individually contemplated. Pharmaceutically acceptable salts and prodrugs of compounds according to the structures below are also contemplated.

10



Aryl is an aromatic ring or ring system such as phenyl, naphthyl, biphenyl, and the like.

5 Heteroaryl is aryl having one or more N, O, or S atoms in the ring, i.e. one or more ring carbons are substituted by N, O, and/or S. While not intending to be limiting, examples of heteroaryl include thienyl, pyridinyl, furyl, benzothienyl, benzofuryl, imidazolyl, indolyl, and the like.

A substituent of aryl or heteroaryl may have up to 20 non-hydrogen atoms each in any stable combination and as many hydrogen atoms as necessary, wherein the non-hydrogen atoms are C, N, O, S, P, F, Cl, Br, and/or I in any stable combination. However, the total number of non-hydrogen atoms on all of the substituents combined must also be 20 or less. A substituent must be sufficiently stable for the compound to be useful as described herein. In addition to the atoms listed above, a substituent may also have a metal cation or other stable cation having an atom not listed above if the substituent is acidic and the salt form is stable. For example, -OH may form an $-O^-Na^+$ salt or CO_2H may form a $CO_2^-K^+$ salt. Thus, while not intending to limit the scope of the invention in any way, a substituent may be:

hydrocarbyl, i.e. a moiety consisting of only carbon and hydrogen such as alkyl, alkenyl, alkynyl, and the like, including linear, branched or cyclic hydrocarbyl, and combinations thereof;

hydrocarbyloxy, meaning O-hydrocarbyl such as OCH_3 , OCH_2CH_3 , O-cyclohexyl, etc, up to 19 carbon atoms;

other ether substituents such as CH_2OCH_3 , $(\text{CH}_2)_2\text{OCH}(\text{CH}_3)_2$, and the like;

thioether substituents including S-hydrocarbyl and other thioether substituents;

hydroxyhydrocarbyl, meaning hydrocarbyl-OH such as CH_2OH , $\text{C}(\text{CH}_3)_2\text{OH}$, etc, up to 19 carbon atoms;

nitrogen substituents such as NO_2 , CN, and the like, including

5 amino, such as NH_2 , $\text{NH}(\text{CH}_2\text{CH}_3\text{OH})$, NHCH_3 , and the like;

carbonyl substituents, such as CO_2H , ester, amide, and the like;

halogen, such as chloro, fluoro, bromo, and the like

fluorocarbyl, such as CF_3 , CF_2CF_3 , etc.;

phosphorous substituents, such as PO_3^{2-} , and the like;

10 sulfur substituents, including S-hydrocarbyl, SH, SO_3H , SO_2 -hydrocarbyl, SO_3 -hydrocarbyl, and the like.

Substituted aryl or heteroaryl may have as many substituents as the ring or ring system will bear, and the substituents may be the same or different. Thus, for example, an aryl ring or a heteroaryl ring may be substituted with chloro and methyl; methyl, OH, and F; CN, NO_2 , and ethyl; and the like including any conceivable substituent or combination of substituent possible in light of this disclosure.

15 Substituted aryl or substituted heteroaryl also includes a bicyclic or polycyclic ring system wherein one or more rings are aromatic and one or more rings are not. For example, indanonyl, indanyl, indanoly, tetralonyl, and the like are substituted aryl. For this type of polycyclic ring system, an aromatic or heteroaromatic ring, not a non-aromatic ring, must be attached to the remainder of the molecule, i.e. the part of the molecule that is not B. In other words, in any structure depicting -B herein, where - is a bond, the bond is a direct bond to an aromatic ring.

20 In one embodiment, B is substituted aryl or heteroaryl.

In another embodiment B is substituted phenyl.

In another embodiment B has no halogen atoms.

In another embodiment B is 4-(1-hydroxy-2,2-dimethylpropyl)phenyl.

In another embodiment B is 4-(1-hydroxy-2-methylpropan-2-yl)phenyl.

25 In another embodiment B is 4-(1-hydroxy-2-methylpropyl)phenyl.

In another embodiment B is 4-(1-hydroxybutyl)phenyl.

In another embodiment B is 4-(1-hydroxyheptyl)phenyl.

In another embodiment B is 4-(1-hydroxyhexyl)phenyl.

In another embodiment B is 4-(1-hydroxypentyl)phenyl.

30 In another embodiment B is 4-(1-hydroxypropyl)phenyl.

In another embodiment B is 4-(3-hydroxy-2-methylheptan-2-yl)phenyl.

In another embodiment B is 4-(3-hydroxy-2-methyloctan-2-yl)phenyl.

In another embodiment B is 1-hydroxy-2,3-dihydro-1H-inden-5-yl.

In another embodiment B is 2,3-dihydro-1H-inden-5-yl.

35 In another embodiment B is 3-(hydroxy(1-propylcyclobutyl)methyl)phenyl.

In another embodiment B is 4-(1-hydroxy-5,5-dimethylhexyl)phenyl.

In another embodiment B is 4-(hydroxy(1-propylcyclobutyl)methyl)phenyl.

In another embodiment B is 4-tert-butylphenyl.

In another embodiment B is 4-hexylphenyl.

In another embodiment B is 4-(1-hydroxy-2-phenylethyl)phenyl.

In another embodiment B is 4-(1-hydroxy-3-phenylpropyl)phenyl.

In another embodiment B is 4-(1-hydroxycyclobutyl)phenyl.

5 In another embodiment B is 4-(2-cyclohexyl-1-hydroxyethyl)phenyl.


In another embodiment B is 4-(3-cyclohexyl-1-hydroxypropyl)phenyl.

In another embodiment B is 4-(cyclohexyl(hydroxy)methyl)phenyl.

In another embodiment B is 4-(cyclohexylmethyl)phenyl.


In another embodiment B is 4-(hydroxy(phenyl)methyl)phenyl.

10 Another embodiment is a compound according to the structure




or a pharmaceutical salt thereof, or a prodrug thereof,
wherein R is hydrogen or C₁₋₁₀ hydrocarbyl.

Another embodiment is a compound according to the structure




or a pharmaceutical salt thereof, or a prodrug thereof,
wherein R is hydrogen or C₁₋₁₀ hydrocarbyl.

Another embodiment is a compound according to the structure



or a pharmaceutical salt thereof, or a prodrug thereof,
wherein R is hydrogen or C₁₋₁₀ hydrocarbyl.

Another embodiment is a compound according to the structure



"C₁₋₁₀" hydrocarbyl is hydrocarbyl having 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 carbon atoms.

Hydrocarbyl is a moiety consisting of only carbon and hydrogen, and includes, but is not limited to alkyl, alkenyl, alkynyl, and the like, and in some cases aryl, and combinations thereof.

5 Alkyl is hydrocarbyl having no double or triple bonds including:

linear alkyl such as methyl, ethyl, propyl, n-butyl, n-pentyl, n-hexyl, and the like;

branched alkyl such as isopropyl, branched butyl isomers (i.e. sec-butyl, tert-butyl, etc), branched pentyl isomers (i.e. isopentyl, etc), branched hexyl isomers, and higher branched alkyl fragments;

cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, etc.; and

alkyl fragments consisting of both cyclic and noncyclic components, whether linear or branched, which may be

5 attached to the remainder of the molecule at any available position including terminal, internal, or ring carbon atoms.

Alkenyl is hydrocarbyl having one or more double bonds including

linear alkenyl, branched alkenyl, cyclic alkenyl, and combinations thereof in analogy to alkyl.

Alkynyl is hydrocarbyl having one or more triple bonds including linear alkynyl, branched alkynyl, cyclic alkynyl and combinations thereof in analogy to alkyl.

10 Aryl is an unsubstituted or substituted aromatic ring or ring system such as phenyl, naphthyl, biphenyl, and the like.

Aryl may or may not be hydrocarbyl, depending upon whether it has substituents with heteroatoms.












Arylalkyl is alkyl which is substituted with aryl. In other words alkyl connects aryl to the remaining part of the molecule. Examples are -CH₂-Phenyl, -CH₂-CH₂-Phenyl, and the like. Arylalkyl may or may not be hydrocarbyl, depending upon whether the aryl portion has substituents with heteroatoms.











15 Unconjugated dienes or polyenes have one or more double bonds which are not conjugated. They may be linear, branched, or cyclic, or a combination thereof.













Combinations of the above are also possible.

Thus, each of the structures below is contemplated. These structures, or pharmaceutically acceptable salts thereof, or prodrugs thereof, individually represent a compound which is an embodiment contemplated herein. In

20 other words, each structure represents a different embodiment.







5

In the above embodiments, x is 5, 6, or 7, and $y + z$ is $2x + 1$.











In one embodiment, x is 5 and $y + z$ is 11.











In another embodiment, x is 6 and $y + z$ is 13.











In another embodiment, x is 7 and $y + z$ is 15.

Hypothetical examples of useful compounds are shown below.

10









Compound examples:

The following are hypothetical examples of useful compounds:

- 5 **Compound Example 1.** A compound having a structure




or a pharmaceutically acceptable salt thereof, or a prodrug thereof;

wherein a dashed line represents the presence or absence of a bond;


- 5 Y is an organic acid functional group, or an amide or ester thereof comprising up to 14 carbon atoms; or Y is hydroxymethyl or an ether thereof comprising up to 14 carbon atoms; or Y is a tetrazolyl functional group; A is $-(CH_2)_6-$, *cis* $-CH_2CH=CH-(CH_2)_3-$, or $-CH_2C\equiv C-(CH_2)_3-$, wherein 1 or 2 carbon atoms may be replaced by S or O; or A is $-(CH_2)_m-Ar-(CH_2)_o-$ wherein Ar is interarylene or heterointerarylene, the sum of m and o is 1, 2, 3, or 4, and wherein 1 $-CH_2-$ may be replaced by S or O, and 1 $-CH_2-CH_2-$ may be replaced by $-CH=CH-$ or $-C\equiv C-$;
- 10 U^1 and U^2 are independently hydrogen, O, or S;
 J^1 is H; O; OH; O-alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms; alkyl having 1, 2, 3, 4, 5, or 6 carbon atoms; F; Cl; Br; I; CN; or CF_3 ; and
 B is substituted aryl or substituted heteroaryl.

- 15 **Compound Example 2.** The compound according to compound example 1 wherein Y is selected from CO_2R^2 , $CON(R^2)_2$, $CON(OR^2)R^2$, $CON(CH_2CH_2OH)_2$, $CONH(CH_2CH_2OH)$, CH_2OH , $P(O)(OH)_2$, $CONHSO_2R^2$, $SO_2N(R^2)_2$, SO_2NHR^2 ,



wherein R^2 is independently H, C_1 - C_6 alkyl, unsubstituted phenyl, or unsubstituted biphenyl.

- 20 **Compound Example 3.** The compound according to compound example 1 or 2 wherein B is substituted phenyl.
Compound Example 4. The compound according to compound example 1 or 2 having a structure




or a pharmaceutically acceptable salt thereof, or a prodrug thereof;

wherein R is hydrogen or C₁₋₁₀ hydrocarbyl.

Compound Example 5. The compound according to compound example 4 wherein R is alkyl.

5 **Compound Example 6.** The compound according to compound example 4 wherein R is arylalkyl.

Compound Example 7. The compound according to compound example any one of compound examples 1 to 6 having a structure



or a pharmaceutically acceptable salt thereof, or a prodrug thereof;

10 wherein R is hydrogen or C₁₋₁₀ hydrocarbyl.

Compound Example 8. The compound according to compound example 1 or 2 wherein A is (3-methylphenoxy)methyl.

Compound Example 9. The compound according to compound example 1 or 2 wherein A is (4-but-2-ynyloxy)methyl.

15 **Compound Example 10.** The compound according to compound example 1 or 2 wherein A is 2-(2-ethylthio)thiazol-4-yl.

Compound Example 11. The compound according to compound example 1 or 2 wherein A is 2-(3-propyl)thiazol-5-yl.

20 **Compound Example 12.** The compound according to compound example 1 or 2 wherein A is 3-methoxymethyl)phenyl.

Compound Example 13. The compound according to compound example 1 or 2 wherein A is 3-(3-propylphenyl.

Compound Example 14. The compound according to compound example 1 or 2 wherein A is 3-methylphenethyl.

Compound Example 15. The compound according to compound example 1 or 2 wherein A is 4-(2-ethyl)phenyl.

Compound Example 16. The compound according to compound example 1 or 2 wherein A is 4-phenethyl.

25 **Compound Example 17.** The compound according to compound example 1 or 2 wherein A is 4-methoxybutyl.

Compound Example 18. The compound according to compound example 1 or 2 wherein A is 5-(methoxymethyl)furan-2-yl .

Compound Example 19. The compound according to compound example 1 or 2 wherein A is 5-(methoxymethyl)thiophen-2-yl.

Compound Example 20. The compound according to compound example 1 or 2 wherein A is 5-(3-propyl)furan-2-yl.

Compound Example 21. The compound according to compound example 1 or 2 wherein A is 5-(3-propyl)thiophen-2-yl.

Compound Example 22. The compound according to compound example 1 or 2 wherein A is 6-hexyl.

Compound Example 23. The compound according to compound example 1 or 2 wherein A is (Z)-6-hex-4-enyl.

Compound Example 24. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(1-hydroxy-2,2-dimethylpropyl)phenyl.

Compound Example 25. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(1-hydroxy-2-methylpropan-2-yl)phenyl.

Compound Example 26. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(1-hydroxy-2-methylpropyl)phenyl.

Compound Example 27. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(1-hydroxybutyl)phenyl.

Compound Example 28. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(1-hydroxyheptyl)phenyl.

Compound Example 29. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(1-hydroxyhexyl)phenyl.

Compound Example 30. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(1-hydroxypentyl)phenyl.

Compound Example 31. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(1-hydroxypropyl)phenyl.

Compound Example 32. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(3-hydroxy-2-methylheptan-2-yl)phenyl.

Compound Example 33. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(3-hydroxy-2-methyloctan-2-yl)phenyl.

Compound Example 34. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 1-hydroxy-2,3-dihydro-1H-inden-5-yl.

Compound Example 35. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 2,3-dihydro-1H-inden-5-yl.

Compound Example 36. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 3-(hydroxy(1-propylcyclobutyl)methyl)phenyl.

Compound Example 37. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(1-hydroxy-5,5-dimethylhexyl)phenyl.

Compound Example 38. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(hydroxy(1-propylcyclobutyl)methyl)phenyl.

Compound Example 39. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-tert-butylphenyl.

Compound Example 40. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-hexylphenyl.

5 **Compound Example 41.** The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(1-hydroxy-2-phenylethyl)phenyl.

Compound Example 42. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(1-hydroxy-3-phenylpropyl)phenyl.

10 **Compound Example 43.** The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(1-hydroxycyclobutyl)phenyl.

Compound Example 44. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(2-cyclohexyl-1-hydroxyethyl)phenyl.

Compound Example 45. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(3-cyclohexyl-1-hydroxypropyl)phenyl.

15 **Compound Example 46.** The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(cyclohexyl(hydroxy)methyl)phenyl.

Compound Example 47. The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(cyclohexylmethyl)phenyl.

20 **Compound Example 48.** The compound according to any one of compound examples 1, 2, and 8-23 wherein B is 4-(hydroxy(phenyl)methyl)phenyl.

Compound Example 49. The compound according to any one of compound examples 1 to 48 wherein J¹ is H.

Compound Example 50. The compound according to any one of compound examples 1 to 48 wherein J¹ is O.

Compound Example 51. The compound according to any one of compound examples 1 to 48 wherein J¹ is OH.

25 **Compound Example 52.** The compound according to any one of compound examples 1 to 48 wherein J¹ is O-alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms.

Compound Example 53. The compound according to any one of compound examples 1 to 48 wherein J¹ is alkyl having 1, 2, 3, 4, 5, or 6 carbon atoms.

Compound Example 54. The compound according to any one of compound examples 1 to 48 wherein J¹ is F.

Compound Example 55. The compound according to any one of compound examples 1 to 48 wherein J¹ is Cl.

30 **Compound Example 56.** The compound according to any one of compound examples 1 to 48 wherein J¹ is Br.

Compound Example 57. The compound according to any one of compound examples 1 to 48 wherein J¹ is I.

Compound Example 58. The compound according to any one of compound examples 1 to 48 wherein J¹ is CN.

Compound Example 59. The compound according to any one of compound examples 1 to 48 wherein J¹ is CF₃.

Compound Example 60. The compound according to any one of compound embodiments 1 to 59 wherein U¹ is H.

35 **Compound Example 61.** The compound according to any one of compound embodiments 1 to 59 wherein U¹ is O.

Compound Example 62. The compound according to any one of compound embodiments 1 to 59 wherein U¹ is S.

Compound Example 63. The compound according to any one of compound embodiments 1 to 62 wherein U² is H.

Compound Example 64. The compound according to any one of compound embodiments 1 to 62 wherein U² is O.

Compound Example 65. The compound according to any one of compound embodiments 1 to 62 wherein U² is S. The following are hypothetical examples of compositions, kits, methods, uses, and medicaments employing the hypothetical compound examples.

Composition Example:

- 5 A composition comprising a compound according to any one of compound examples 1 to 65, wherein said composition is a liquid which is ophthalmically acceptable.

Medicament Examples:

Use of a compound according to any one of compound examples 1 to 65 in the manufacture of a medicament for the treatment of glaucoma or ocular hypertension in a mammal.

- 10 Use of a compound according to any one of compound examples 1 to 65 in the manufacture of a medicament for the treatment of baldness in a person.

A medicament comprising a compound according to any one of compound examples 1 to 65, wherein said composition is a liquid which is ophthalmically acceptable.

Method Example:

- 15 A method comprising administering a compound according to any one of compound examples 1 to 65 to a mammal for the treatment of glaucoma or ocular hypertension.

Kit Example:

- A kit comprising a composition comprising compound according to any one of compound examples 1 to 65, a container, and instructions for administration of said composition to a mammal for the treatment of glaucoma or ocular
20 hypertension.

- A "pharmaceutically acceptable salt" is any salt that retains the activity of the parent compound and does not impart any additional deleterious or untoward effects on the subject to which it is administered and in the context in which it is administered compared to the parent compound. A pharmaceutically acceptable salt also refers to any salt
25 which may form in vivo as a result of administration of an acid, another salt, or a prodrug which is converted into an acid or salt.

- Pharmaceutically acceptable salts of acidic functional groups may be derived from organic or inorganic bases. The salt may comprise a mono or polyvalent ion. Of particular interest are the inorganic ions lithium, sodium, potassium, calcium, and magnesium. Organic salts may be made with amines, particularly ammonium salts such as
30 mono-, di- and trialkyl amines or ethanol amines. Salts may also be formed with caffeine, tromethamine and similar molecules. Hydrochloric acid or some other pharmaceutically acceptable acid may form a salt with a compound that includes a basic group, such as an amine or a pyridine ring.

- A "prodrug" is a compound which is converted to a therapeutically active compound after administration, and the term should be interpreted as broadly herein as is generally understood in the art. While not intending to limit the
35 scope of the invention, conversion may occur by hydrolysis of an ester group or some other biologically labile group. Generally, but not necessarily, a prodrug is inactive or less active than the therapeutically active compound to which it is converted. Ester prodrugs of the compounds disclosed herein are specifically contemplated. An ester may be derived from a carboxylic acid of C1 (i.e. the terminal carboxylic acid of a natural prostaglandin), or an ester may be

derived from a carboxylic acid functional group on another part of the molecule, such as on a phenyl ring. While not intending to be limiting, an ester may be an alkyl ester, an aryl ester, or a heteroaryl ester. The term alkyl has the meaning generally understood by those skilled in the art and refers to linear, branched, or cyclic alkyl moieties. C₁₋₆ alkyl esters are particularly useful, where alkyl part of the ester has from 1 to 6 carbon atoms and includes, but is not limited to, methyl, ethyl, propyl, isopropyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *t*-butyl, pentyl isomers, hexyl isomers, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and combinations thereof having from 1-6 carbon atoms, etc.

Those skilled in the art will readily understand that for administration or the manufacture of medicaments the compounds disclosed herein can be admixed with pharmaceutically acceptable excipients which per se are well known in the art. Specifically, a drug to be administered systemically, it may be confectioned as a powder, pill, tablet or the like, or as a solution, emulsion, suspension, aerosol, syrup or elixir suitable for oral or parenteral administration or inhalation.

For solid dosage forms or medicaments, non-toxic solid carriers include, but are not limited to, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, the polyalkylene glycols, talcum, cellulose, glucose, sucrose and magnesium carbonate. The solid dosage forms may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the technique described in the U.S. Pat. Nos. 4,256,108; 4,166,452; and 4,265,874 to form osmotic therapeutic tablets for control release. Liquid pharmaceutically administrable dosage forms can, for example, comprise a solution or suspension of one or more of the presently useful compounds and optional pharmaceutical adjuncts in a carrier, such as for example, water, saline, aqueous dextrose, glycerol, ethanol and the like, to thereby form a solution or suspension. If desired, the pharmaceutical composition to be administered may also contain minor amounts of nontoxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like. Typical examples of such auxiliary agents are sodium acetate, sorbitan monolaurate, triethanolamine, sodium acetate, triethanolamine oleate, etc. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., 16th Edition, 1980. The composition of the formulation to be administered, in any event, contains a quantity of one or more of the presently useful compounds in an amount effective to provide the desired therapeutic effect.

Parenteral administration is generally characterized by injection, either subcutaneously, intramuscularly or intravenously. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol and the like. In addition, if desired, the injectable pharmaceutical compositions to be administered may also contain minor amounts of non-toxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like.

The amount of the presently useful compound or compounds administered is dependent on the therapeutic effect or effects desired, on the specific mammal being treated, on the severity and nature of the mammal's condition, on the manner of administration, on the potency and pharmacodynamics of the particular compound or compounds

employed, and on the judgment of the prescribing physician. The therapeutically effective dosage of the presently useful compound or compounds may be in the range of about 0.5 or about 1 to about 100 mg/kg/day.

A liquid which is ophthalmically acceptable is formulated such that it can be administered topically to the eye. The comfort should be maximized as much as possible, although sometimes formulation considerations (e.g. drug stability) may necessitate less than optimal comfort. In the case that comfort cannot be maximized, the liquid should be formulated such that the liquid is tolerable to the patient for topical ophthalmic use. Additionally, an ophthalmically acceptable liquid should either be packaged for single use, or contain a preservative to prevent contamination over multiple uses.

For ophthalmic application, solutions or medicaments are often prepared using a physiological saline solution as a major vehicle. Ophthalmic solutions should preferably be maintained at a comfortable pH with an appropriate buffer system. The formulations may also contain conventional, pharmaceutically acceptable preservatives, stabilizers and surfactants.

Preservatives that may be used in the pharmaceutical compositions of the present invention include, but are not limited to, benzalkonium chloride, chlorobutanol, thimerosal, phenylmercuric acetate and phenylmercuric nitrate. A useful surfactant is, for example, Tween 80. Likewise, various useful vehicles may be used in the ophthalmic preparations of the present invention. These vehicles include, but are not limited to, polyvinyl alcohol, povidone, hydroxypropyl methyl cellulose, poloxamers, carboxymethyl cellulose, hydroxyethyl cellulose and purified water.

Tonicity adjustors may be added as needed or convenient. They include, but are not limited to, salts, particularly sodium chloride, potassium chloride, mannitol and glycerin, or any other suitable ophthalmically acceptable tonicity adjustor.

Various buffers and means for adjusting pH may be used so long as the resulting preparation is ophthalmically acceptable. Accordingly, buffers include acetate buffers, citrate buffers, phosphate buffers and borate buffers. Acids or bases may be used to adjust the pH of these formulations as needed.

In a similar vein, an ophthalmically acceptable antioxidant for use in the present invention includes, but is not limited to, sodium metabisulfite, sodium thiosulfate, acetylcysteine, butylated hydroxyanisole and butylated hydroxytoluene.

Other excipient components which may be included in the ophthalmic preparations are chelating agents. A useful chelating agent is edetate disodium, although other chelating agents may also be used in place or in conjunction with it.

The ingredients are usually used in the following amounts:

<u>Ingredient</u>	<u>Amount (% w/v)</u>
active ingredient	about 0.001-5
preservative	0-0.10
vehicle	0-40
tonicity adjustor	1-10
buffer	0.01-10
pH adjustor	q.s. pH 4.5-7.5

antioxidant	as needed
surfactant	as needed
purified water	as needed to make 100%

5 For topical use, creams, ointments, gels, solutions or suspensions, etc., containing the compound disclosed herein are employed. Topical formulations may generally be comprised of a pharmaceutical carrier, cosolvent, emulsifier, penetration enhancer, preservative system, and emollient.

The actual dose of the active compounds of the present invention depends on the specific compound, and on the condition to be treated; the selection of the appropriate dose is well within the knowledge of the skilled artisan.


10 For treatment of diseases affecting the eye including glaucoma, these compounds can be administered topically, periocularly, intraocularly, or by any other effective means known in the art.

A person of ordinary skill in the art understands the meaning of the stereochemistry associated with the hatched wedge/solid wedge structural features. For example, an introductory organic chemistry textbook (Francis A. Carey, Organic Chemistry, New York: McGraw-Hill Book Company 1987, p. 63) states "a wedge indicates a bond coming from the plane of the paper toward the viewer" and the hatched wedge, indicated as a "dashed line", "represents a bond receding from the viewer."

Synthetic Methods

20

Scheme 1



(a) **a**, CuI, MeN(H)CH₂CH₂N(H)Me, K₂CO₃, MeCN; (b) NaH, **b**, DMF; (c) TFA, CH₂Cl₂; (d) 3-chloropropionyl chloride, K₂CO₃, DMF.

While there are many ways the compounds disclosed herein, one exemplary synthesis may begin with commercially available N-Boc hydrazine (**1**, also known as *t*-butyl carbazate, Aldrich Chemical Company, see


Scheme 1). N-arylation occurs to give **2** according to Buchwald's copper-catalyzed procedure (*Org. Lett.* **2001**, 3, 3803-3805) using a wide variety of substituted bromophenyl and other bromoaryl compounds **a**. The haloarenes **a** are either available commercially or may be made according to published literature procedures. For example, United States Patent Application No. 11/009,298, filed on December 10, 2004 and United States Provisional Patent
5 Application 60/742,779 filed on December 6, 2005, both of which are expressly incorporated by reference herein, disclose methods of making a number of useful substituted bromophenyl compounds. These procedures may also be readily adapted to other bromoaryl compounds such as substituted bromothieryl, substituted bromofuryl, substituted bromopyridinyl, substituted bromonaphthyl, substituted bromobenzothieryl, and the like. Intermediate **2** is then alkylated on N' using electrophiles **b** to provide intermediate **3**.

10 Examples of **b** include ethyl 7-bromoheptanoate (commercially available from Aldrich Chemical Company) and methyl 7-bromohept-5-ynoate (*Org. Synth.* **1993**, Collect. Vol. VIII,415-420). Examples of **b** also include electrophiles bearing aryl and heteroaryl groups (e.g. methyl 4-(2-bromoethyl)benzoate [available in one step from commercially available 4-(2-bromoethyl)benzoic acid] and methyl 5-(3-bromopropyl)-thiophene-2-carboxylate [see WO 2004/037786, incorporated by reference herein]). Other methods of preparing **b** are readily ascertained by those
15 of ordinary skill in the art based upon this disclosure.

Intermediate **3** is then deprotected to give intermediate **4** which is treated with 3-chloropropionyl chloride (using the method of Araldi, et al., US 2004/ 0254233) to give product **5**. Compound **5** may be the target compound, or may require deprotection(s) and/or functionalization (depending on the nature of B and Y) to arrive at the target compound.

20

Scheme 2




(a) **a**, Pd(OAc)₂, BINAP, NaOt-Bu, toluene; (b) TsOH, H₂O, EtOH; (c) BOC₂O, CH₂Cl₂; (d) 3-chloropropionyl chloride, K₂CO₃, DMF; (e) TFA, CH₂Cl₂; (f) NaH, **b**, DMF.

In another hypothetical example, benzophenone hydrazone (**6**, Aldrich chemical company) serves as the starting material (see Scheme 2). In this case, N'-arylation occurs to give **7** according to Buchwald's palladium-catalyzed procedure (*J. Am. Chem. Soc.* **1998**, *120*, 6621-6622) using a wide variety of substituted bromophenyl and other bromoaryl compounds **a**. Deprotection reveals the aryl hydrazine **8** which is protected to give N-Boc-N'-aryl hydrazine **9**. A large variety of substituted aryl hydrazines such as **8** are commercially available. Treatment of **9** with 3-chloropropionyl chloride (using the method of Araldi, et al., US 2004/ 0254233) gives intermediate **10** which is then deprotected and alkylated on nitrogen to afford product **12**. Compound **12** may be the target compound, or may require deprotection(s) and/or functionalization (depending on the nature of B and Y) to arrive at the target compound.

Intermediates **2** and **9** are regioisomers. In certain cases, the copper chemistry used to arrive at **2** will also afford compound **9** (see Buchwald, above, and Buchwald, *J. Am. Chem. Soc.* **2001**, *123*, 7727-7729). Palladium-catalyzed arylation may also afford mixtures of regioisomers (Wang, et al., *Tetrahedron Lett.* **1999**, *40*, 3543-3546), mainly favoring isomer **2**. This represents an alternative approach to compound **9**.

In another hypothetical example (scheme 3), intermediate **3** may be cyclized using malonyl chloride (or other activated source of malonate) to afford the pyrazolidine dione **13**. This may be the target compound, or may serve as an intermediate to compounds substituted at the carbon adjacent to both ketones such as **14**. The ring methylene of dione **13** is alkylated using alkyl electrophiles J by employing a mild base such as potassium carbonate using the method of Vennerstrom and Holmes, *J. Med. Chem.* **1987**, 30, 563-567. Chlorination (J=Cl) may be accomplished using the method of Pesin et al. *Z. Obshch. Khim.* **1958**, 20, 2816-2820. Other electrophiles known in the art can be envisioned to introduce other J groups (e.g. the Davis oxaziridine reagent or other suitable reagent for J=OH and Br₂ or NBS for J=Br). As an alternative, J may be introduced into the malonate derivative prior to reaction with compound **4**. As halogens such as Cl and Br are useful as nucleophilic substitution leaving groups, they provide a means for reaching other J groups using the nucleophile J- or a synthetic equivalent thereof. Compounds wherein J is OH may also be derivatized with triflate, tosylate, or the like to provide good leaving groups for nucleophilic substitution. For example, J = CN may be reached in this manner. These procedures are well known in the art.


Scheme 3



(a) CH₂(COCl)₂, Et₃N, dioxane; (b) **c**, K₂CO₃, DMF.

The carbon adjacent to the ketone in pyrazolidinones **5** and **12** may be alkylated using alkyl electrophiles J by employing LDA as the base (e.g. Hlasta et al., *J. Med. Chem.* **1991**, 34, 1560-1570) (Scheme 4). Other J groups may be introduced using LDA and the electrophiles discussed above for scheme 3. As an alternative, J may be introduced into the chloropropionate derivative prior to reaction with compound **4** or compound **9**.

Scheme 4




(a) **c**, LDA, THF.

A carbonyl oxygen may be replaced with sulfur using Lawesson's reagent or other methods. Thus, compounds wherein J¹, U¹ and/or U² are S may be prepared.

The steps of these synthetic procedures may also be rearranged. For example in Scheme 1, N'-alkylation may take place prior to amide N-arylation. It is also envisioned that chloro- or iodo-arenes may be employed in place of the bromoarenes described above. The requisite chloro- or iodo-arenes may be prepared by procedures analogous to those used to prepare the bromoarenes. Alternative starting materials to N-Boc hydrazine and benzophenone hydrazone include commercially available 3-pyrazolidinone HCl (see Scheme 5 and example 1) and 3,5-pyrazolidine dione.

Scheme 5



(a) K_2CO_3 , DMF, rt; (b) CuI, 1,10-phenanthroline, Cs_2CO_3 , DMF, 80 °C; (c) LiOH (aq.), THF, 40 °C.

5

Example 1

5-(3-(2-(4-(1-hydroxyhexyl)phenyl)-3-oxopyrazolidin-1-yl)propyl)thiophene-2-carboxylic acid (**25**)

Step 1. Alkylation of **20** with **21** to give **22**

Potassium carbonate (8.28 g, 60.0 mmol), methyl 5-(3-bromopropyl)thiophene-2-carboxylate (**21**, for preparation, see Billot, X. et al WO2004/037786, incorporated by reference herein, 5.66 g, 21.5 mmol), 3-pyrazolidinone HCl (**20**, 2.57 g, 21.0 mmol) and DMF (25 mL) were combined under argon and stirred at rt. After 3 d at rt, HPLC analysis indicated the reaction to be complete and EtOAc (100 mL) was added. The mixture was washed with water (2x75 mL). The combined aqueous phase was back-extracted with EtOAc (50 mL). The combined organic phase was washed with brine (75 mL), filtered through filter paper and concentrated in vacuo to afford 7.5 g of a crude oil.

Purification of the crude product by flash column chromatography on 75 g silica gel (50% CH_2Cl_2 /hexane $\rightarrow$ CH_2Cl_2 $\rightarrow$ 20% EtOAc/ CH_2Cl_2 $\rightarrow$ 100% EtOAc $\rightarrow$ 10% MeOH/EtOAc, gradient) afforded 3.2 g (57%) of **22** which solidified on standing.

Step 2. Arylation of **22** with **23** to give **24**

Cesium carbonate (4.89 g, 15.0 mmol), copper (I) iodide (19 mg, 0.10 mmol), 1,10-phenanthroline (180 mg, 1.0 mmol), amide **22** (2.88 g, 10.7 mmol) and 1-(4-iodophenyl)hexan-1-ol (**23**, prepared from 4-iodobenzaldehyde using the method described for the analogous bromoarene in Old and Dinh, WO2006/098918, 3.34 g 11.0 mmol) and DMF (25 mL) were combined into a sealable tube. Argon was bubbled through the mixture for 15 min, the tube was sealed, and the mixture was heated at 80 °C overnight. Upon cooling, HPLC analysis showed no reaction had taken place. More copper (I) iodide (60 mg, 0.32 mmol) and 1,10-phenanthroline (180 mg, 1.0 mmol) was added and the mixture was heated at 80 °C overnight. Upon cooling, HPLC analysis showed the reaction was nearly complete.

EtOAc (100 mL) was added and the mixture was filtered through 25 g silica gel, washing the pad with EtOAc (150

mL). The filtrate was concentrated in vacuo to afford 3.7 g of a crude oil. Purification of the crude product by flash column chromatography on 50 g silica gel (hexane $\rightarrow$ 25% CH₂Cl₂/hexane $\rightarrow$ CH₂Cl₂ $\rightarrow$ 5% EtOAc/CH₂Cl₂ $\rightarrow$ 25% EtOAc/CH₂Cl₂, gradient) afforded 0.29 g (6%) of **24** as a brown oil.

Step 3. Saponification of **24** to give **25**


- 5 Lithium hydroxide (0.22 mL of a 1.0 M aqueous solution, 0.22 mmol) was added to a solution of ester **24** (20 mg, 0.045 mmol) in THF (0.22 mL). The solution was heated at 40 °C for 18 h, then cooled to rt. The mixture was partitioned between 1.0 M HCl (1 mL) and EtOAc (5 mL). The phases were separated and the aqueous phase was extracted with EtOAc (2x5 mL). The combined extracts were washed with brine (5 mL), dried (Na₂SO₄), filtered and concentrated in vacuo. Purification of the crude residue by column chromatography on 4 g silica gel (CH₂Cl₂ $\rightarrow$ 20% MeOH/CH₂Cl₂) afforded 11 mg (57%) of the title compound.
- 10

Based upon this disclosure, numerous other ways of preparing the compounds disclosed herein will be apparent to a person of ordinary skill in the art.

15

In vitro testing


United States Patent Application Serial No. 11/553,143, filed on October 26, 2006, incorporated by reference herein, describes the methods used to obtain the in vitro data in the table below.

Structure	EP2 data			EP4 data		Other Receptors (EC50 in nM)					
	flpr EC50	cAMP EC50	Ki	flpr EC50	KI	hFP	hEP1	hEP3A	hTP	hIP	hDP
	79	1.8	251	NT	>10000	NA	NA	5068	NA	NA	NA

- 20 The foregoing description details specific methods and compositions that can be employed to practice the present invention, and represents the best mode contemplated. However, it is apparent for one of ordinary skill in the art that further compounds with the desired pharmacological properties can be prepared in an analogous manner, and that the disclosed compounds can also be obtained from different starting compounds via different chemical reactions. Similarly, different pharmaceutical compositions may be prepared and used with substantially the same result. Thus,
- 25 however detailed the foregoing may appear in text, it should not be construed as limiting the overall scope hereof; rather, the ambit of the present invention is to be governed only by the lawful construction of the claims.

What is claimed is:

1. A compound having a structure



or a pharmaceutically acceptable salt thereof, or a prodrug thereof;

- 5 wherein a dashed line represents the presence or absence of a bond;

Y is an organic acid functional group, or an amide or ester thereof comprising up to 14 carbon atoms; or Y is

hydroxymethyl or an ether thereof comprising up to 14 carbon atoms; or Y is a tetrazolyl functional group;

A is $-(CH_2)_6-$, *cis* $-CH_2CH=CH-(CH_2)_3-$, or $-CH_2C\equiv C-(CH_2)_3-$, wherein 1 or 2 carbon atoms may be replaced by S or

O; or A is $-(CH_2)_m-Ar-(CH_2)_o-$ wherein Ar is interarylene or heterointerarylene, the sum of m and o is 1, 2, 3, or 4, and

- 10 wherein 1 $-CH_2-$ may be replaced by S or O, and 1 $-CH_2-CH_2-$ may be replaced by $-CH=CH-$ or $-C\equiv C-$;


U¹ and U² are independently hydrogen, O, or S;

J¹ is H; O; OH; O-alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms; alkyl having 1, 2, 3, 4, 5, or 6 carbon atoms; F; Cl; Br; I;

CN; or CF₃; and

B is substituted aryl or substituted heteroaryl.

- 15 2. A compound which is a carboxylic acid or a bioisostere thereof, said carboxylic acid having a structure



or a pharmaceutically acceptable salt thereof, or a prodrug thereof,;

wherein a dashed line represents the presence or absence of a bond;

A is $-(CH_2)_6-$, *cis* $-CH_2CH=CH-(CH_2)_3-$, or $-CH_2C\equiv C-(CH_2)_3-$, wherein 1 or 2 carbon atoms may be replaced by S or

- 20 O; or A is $-(CH_2)_m-Ar-(CH_2)_o-$ wherein Ar is interarylene or heterointerarylene, the sum of m and o is 1, 2, 3, or 4, and

wherein 1 $-CH_2-$ may be replaced by S or O, and 1 $-CH_2-CH_2-$ may be replaced by $-CH=CH-$ or $-C\equiv C-$;


U¹ and U² are independently hydrogen, O, or S;

J¹ is H; O; OH; O-alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms; alkyl having 1, 2, 3, 4, 5, or 6 carbon atoms; F; Cl; Br; I;

CN; or CF₃; and


- 25 B is substituted aryl or substituted heteroaryl.

3. The compound of claim 1 wherein Y is selected from CO_2R^2 , $\text{CON}(\text{R}^2)_2$, $\text{CON}(\text{OR}^2)\text{R}^2$, $\text{CON}(\text{CH}_2\text{CH}_2\text{OH})_2$, $\text{CONH}(\text{CH}_2\text{CH}_2\text{OH})$, CH_2OH , $\text{P}(\text{O})(\text{OH})_2$, $\text{CONHSO}_2\text{R}^2$, $\text{SO}_2\text{N}(\text{R}^2)_2$, SO_2NHR^2 ,



wherein R^2 is independently H, $\text{C}_1\text{-C}_6$ alkyl, unsubstituted phenyl, or unsubstituted biphenyl.

4. The compound of claim 3 wherein A has a structure selected from:




5. The compound of claim 4 wherein A is 5-(3-propyl)thiophen-2-yl.

6. The compound of claim 3 wherein A is 6-hexyl.

7. The compound of claim 3 wherein A is (Z)-6-hex-4-enyl.

8. The compound of claim 4 wherein B is substituted phenyl.


9. The compound of claim 1 having a structure



or a pharmaceutically acceptable salt thereof, or a prodrug thereof;

wherein R is hydrogen or C_{1-10} hydrocarbyl.


10. The compound of claim 4 having a structure



or a pharmaceutically acceptable salt thereof, or a prodrug thereof;

wherein R is hydrogen or C₁₋₁₀ hydrocarbyl.

11. The compound according to any one of claims 1-10, wherein J is H.
- 5 12. The compound according to any one of claims 1-11, wherein U¹ is H.
13. The compound according to any one of claims 1-12 wherein U² is O.
14. The compound of claim 13 having the formula:



15. Use of a compound according to any one of claims 1-14 in the manufacture of a medicament for the treatment of
- 10 glaucoma or ocular hypertension in a mammal.
16. A method comprising administering a compound according to any one of claims 1-14 to a mammal for the treatment of glaucoma or ocular hypertension.
17. A kit comprising a composition comprising compound according to any one of claims 1-14, a container, and instructions for administration of said composition to a mammal for the treatment of glaucoma or ocular hypertension.
- 15 18. A composition comprising a compound according to any one of claims 1-14, wherein said composition is a liquid which is ophthalmically acceptable.