



US 20100209435A1

(19) **United States**

(12) **Patent Application Publication**  
**Boyle et al.**

(10) **Pub. No.: US 2010/0209435 A1**

(43) **Pub. Date: Aug. 19, 2010**

(54) **HUMAN ANTI-OPGL NEUTRALIZING ANTIBODIES AS SELECTIVE OPGL PATHWAY INHIBITORS**

(75) Inventors: **William J. Boyle**, Malibu, CA (US); **Eugene Medlock**, Westlake, CA (US); **John K. Sullivan**, Newbury Park, CA (US); **Robin L. Elliott**, Newbury Park, CA (US); **Frank Martin**, Newbury Park, CA (US); **Haichun Huang**, Fremont, CA (US)

Correspondence Address:  
**MCDONNELL BOEHNEN HULBERT & BERGHOFF LLP**  
**300 S. WACKER DRIVE, 32ND FLOOR**  
**CHICAGO, IL 60606 (US)**

(73) Assignees: **AMGEN INC.**, Thousand Oaks, CA (US); **MEDAREX, INC.**, Princeton, NJ (US)

(21) Appl. No.: **12/705,962**

(22) Filed: **Feb. 15, 2010**

**Related U.S. Application Data**

(63) Continuation of application No. 10/408,901, filed on Apr. 7, 2003, now Pat. No. 7,718,776.

(60) Provisional application No. 60/370,407, filed on Apr. 5, 2002.

**Publication Classification**

(51) **Int. Cl.**  
*A61K 39/395* (2006.01)  
*C07K 16/24* (2006.01)  
*C07H 21/04* (2006.01)  
*C12N 15/63* (2006.01)  
*C12N 5/10* (2006.01)  
*C12P 21/02* (2006.01)  
*A61P 19/10* (2006.01)  
*A61P 19/08* (2006.01)

(52) **U.S. Cl.** ..... **424/158.1**; 530/389.2; 536/23.53; 435/320.1; 435/358; 435/69.6

(57) **ABSTRACT**

Monoclonal antibodies and hybridomas producing them that interact with osteoprotegerin ligand (OPGL) are provided. Methods of treating osteopenic disorders by administering a pharmaceutically effective amount of antibodies to OPGL are also provided. Methods of detecting the amount of OPGL in a sample using antibodies to OPGL are further provided.

**Constant region Heavy Chain IgG1**

```

gcctccacca agggcccatc ggtcttcccc ctggcacccct cctccaagag cacctctggg      60
ggcacagcgg ccctgggctg cctggtcaag gactacttcc ccgaaccggg gacgggtgtcg      120
tggaaactcag gcgccctgac cagcggcgtg cacaccttcc cggctgtcct acagtcctca      180
ggacttact ccctcagcag cgtggtgacc gtgcccctcca gcagcttggg caccagacc      240
tacatctgca acgtgaatca caagcccagc aacaccaagg tggacaagaa agttgagccc      300
aaatcttgtg acaaaactca cacatgcccc ccgtgcccag cacctgaact cctgggggga      360
ccgtcagttt tcctcttccc cccaaaacc agggacacc tcatgatctc cgggacccct      420
gaggtcacat gcgtggtggt ggacgtgagc cacgaagacc ctgaggtcaa gttcaactgg      480
tacgtggacg gcgtggaggt gcataatgcc aagacaaagc cgcgggagga gcagtacaac      540
agcacgtacc gtgtggtcag cgtcctcacc gtctgcacc aggactggct gaatggcaag      600
gagtacaagt gcaaggtctc caacaaagcc ctcccagccc ccatcgagaa accatctcc      660
aaagccaaag ggcagccccg agaaccacag gtgtacaccc tgcccccatc ccgggatgag      720
ctgaccaaga accaggtcag cctgacctgc ctggtcaaaag gcttctatcc cagcgacatc      780
gccgtggagt gggagagcaa tgggcagccg gagaacaact acaagaccac gcctcccgtg      840
ctggactccg acggctcctt ctctctctat agcaagctca ccgtggacaa gaggagggtg      900
cagcagggga acgtctctc atgctccgtg atgcatgagg ctctgcacaa ccactacacg      960
cagaagagcc tctccctgtc tccgggtaaa tgataagtcg acatgcccctg aattctgcag      1020
atatccatca cactgggggc cgctcgagca tgcattctaga gggccc      1066
    
```

**Figure 1A**

ASTKGPSVFP	LAPSSKSTSG	GTAALGCLVK	DYFFPEPVTVS	WNSGALTSGV	HTFFPAVLQSS	60
GLYSLSSVVT	VPSSSLGTQT	YICNVNHHKPS	NTKVDKKVEP	KSCDKTHTTCP	PCPAPELLGG	120
PSVFLFPPKP	KDTLMISRTP	EVTCVVVDVDS	HEDPEVKFNW	YVDGVEVHNA	KTKPREEQYN	180
STYRVVSVLT	VLHQDWLNGK	EYKCKVSNKA	LPAPIEKTIS	KAKGQPREPQ	VYTLPPSRDE	240
LTKNQVSLTC	LVKGFYPSDI	AVEWESNGQP	ENNYKTTTPV	LDSDGSFFFLY	SKLTVDKSRW	300
QQGNVFSCSV	MHEALHNHYT	QKSLSLSPGK				330

Figure 1B

**Kappa chain constant region**

```

actgtggctg caccatctgt cttcatcttc ccgccatctg atgagcagtt gaaatctgga    60
actgcctctg ttgtgtgcct gctgaataac ttctatccca gagaggccaa agtacagtgg    120
aaggtaggata acgccctcca atcgggtaac tcccaggaga gtgtcacaga gcagggacagc    180
aaggacagca cctacagcct cagcagcacc ctgacgctga gcaaaagcaga ctacgagaaa    240
cacaagaagtct acgcctgcga agtcacccat cagggcctga gctcgcccgt cacaagagagc    300
ttcaacaggg gagagtgtta g                                     321
    
```

**Figure 2A**

```

TVAAPSVFIF PPSDEQLKSG TASVVCLLNN FYPREAKVQW KVDNALQSGN SQESVTEQDS    60
KDSTYLSLST LTLSKADYEK HKVYACEVTH QGLSSPVTKS FNRGEC                    106
    
```

**Figure 2B**

**22B3 Heavy Chain**

gaggtgcagc tgggtgcagtc tgggggaggc ttggtacatc ctgggggggtc cctgagactc 60  
 tcctgtgaag gctctggatt caccttcagt agcaatggtg tgcactgggt gcgcccagact 120  
 ccaggaaaag gtctggagt ggtatcaggt attggtactg ctggtggcac atactatgca 180  
 gactccgtga agggccgatt caccatttcc agagacaatg tcaagaagtc cttgtatcctt 240  
 caaatgaaca gcctgagagc cgaggacatg gctattttatt attgtgtaag aaaaaactgg 300  
 ggatggttcg acccctgggg ccagggagcc ctggtcaccg tctctagt 348

**Figure 3A**

EVQLVQSGGG LVHPGGSLRL SCEGSGFTFS SNGMHWRQT PGKGLEWVSG IGTAGGTYYA 60  
 DSVKGRFTIS RDNVKKSLYL QMNSLRAEDM AIYCVRKNW GWFDPWGQGA LVTVSS 116

**Figure 3B**

**22B3 Kappa Chain**

gaaattgtgc tgaccagtc tccagccacc ctgtctttgt ctccagggga aagagccacc 60  
ctctcctgca gggccagtc gagtgtaac agctacttag cctggttcca acagaaacct 120  
ggccaggctc ccagactcct catctatgat gcatccaaca gggccactgg catcccagcc 180  
aggttcagtg gcagtgggtc tgggacagac ttcaacttca ccatcagcag ccttgagcct 240  
gaagattttg caatttatta ctgtcagcag cgtagcaact ggcctccgtt cacttttggc 300  
caggggacca agctggagat caaacga 327

**Figure 4A**

EIVLTQSPAT LSLSPGERAT LSCRASQSVN SYLAWFQQKP GQAPRLLIYD ASNRATGIPA 60  
RFSGSGSGTD FTLTISLIEP EDFAIYYCQQ RSNWPPFTFG QGTKLEIKR 109

**Figure 4B**

**2E11 Heavy Chain**

```

gaggtgcagc  tgggtgcagtc  ggggggagac  ttggtacatc  ctgggggggtc  cctgagactc  60
tcctgtgtcg  gctctggatt  caccttcagt  cactatcctt  tgcaactgggt  tcgccaggct  120
ccaggaaaag  gtctggagtg  gataatcagg  attcatactg  gtggtggcac  atactataca  180
gactccgtga  agggccggtt  caccatctcc  agcgacaatg  ccaagaactc  cttatatctt  240
caatgaaca  ccctgagagc  cgaggacatg  gctgtgtatt  actgtgcaag  agggcgaaac  300
tcctttgact  actggggcca  gggaaacctg  gtcatcgtct  ctagt

```

**Figure 5A**

```

EVQLVQSGGD  LVHPGGSLRL  SCVSGGFTFS  HYPLHWVRQA  PGKGLEWISG  IHTGGGTYT  60
DSVKGRFTIS  SDNAKNSLYL  QMNTLRAEDM  AVYYCARGRN  SFDYWGQGTL  VIVSS  115

```

**Figure 5B**

**2E11 Kappa Chain**

gacatccaga tgaccagtc tccatcctca ctgtctgcat ctgtaggaga cagagtcacc 60  
atcacttgtc gggcgagtca ggtatttagc agctggttag cctggtatca gcagaaacca 120  
gagaaagccc ctaagtccct gatctatgct gcatccagtt tgcaaaagtgg ggtcccatca 180  
aggttcagcg gcagtgatc tgggacagat ttcaactctca ccatcagcag cctgcagtct 240  
gaagattttg caacttatta ctgccaacag tataatagtt accctcccac cttcggccaa 300  
gggacacgac tggagattaa acga 324

**Figure 6A**

DIQMTQSPSS LSASVGDTRVT ITCRASQGIS SWLAWYQQKP EKAPKSLIYA ASSIQSGVPS 60  
RFSGSGGTD FFLTISSLQS EDFATYYCQQ YNSYPPTFGQ GTRLEIKR 108

**Figure 6B**



**2D8 Heavy Chain**

```

gaggtgcagc  tgggtgcagtc  tgggggaggc  ttggtacatc  ctgggggggtc  cctgagactc  60
tcctgtgcag  gctctggatt  caccttcagt  agctatggga  tgcactgggt  tcgccaggct  120
ccaggaaaag  gtctggagtg  ggtatcaggt  attggtactg  gtggtggcac  atactatgca  180
gactccgtga  agggccgatt  caccatctcc  agagacaatg  tcaagaactc  cttgtatcct  240
caaatgaaca  gcctgagagc  cgaggacatg  gctgtgtatt  actgtgcaag  aaaaaactgg  300
ggatggtttg  actactgggg  ccaggggaacc  ctggtcaccg  tctctagt

```

**Figure 7A**

```

EVQLVQSGGG  LVHPGGSLRL  SCAGSGFTFS  SYGMHWVRQA  PGKGLEWVSG  IGTGGGTYYA  60
DSVKGRFTIS  RDNVKNSLYL  QMNSLRAEDM  AVYYCARKNW  GWFYWGQGT  LVTVSS      116

```

**Figure 7B**

**2D8 Kappa Chain**

gaaattgtgc tgaccagtc tccagccacc ctgtctttgt ctccagggga aagagccacc 60  
 ctctcctgca gggccagtca gattattagc agtacttag cctggtacca acagaaacct 120  
 gggcaggctc ccaggctcct catctatgat gcattccaaca gggccactgg catcccagcc 180  
 aggttcagtg gcagtgggtc tgggacagac ttcactctca ccatcagcag cctagagcct 240  
 gaagattttg cagtttatta ctgtcagcag cgtagcaaat ggcctccgta cacttttggc 300  
 caggggacca aactcgagat caaacga 327

**Figure 8A**

EIVLTQSPAT LSLSPGERAT LSCRASQSI SYLAWYQQKP GQAPRLLIYD ASNRATGIPA 60  
 RFSGSGSGTD FTLTISSLEP EDFAVYYCQQ RSKWPPYTFG QGTKLEIKR 109

**Figure 8B**

**18B2 Heavy Chain**

```

gaggtgcagc  tgggtgcagtc  tgggggaggc  ttggtacatc  ctgggggggtc  cctgagactc  60
tcctgtgtag  gctctagatt  caccttcagt  gcctatccta  tgcactgggt  tcgccaggct  120
ccaggaaaag  gtctggagtg  ggtatcaggt  attggttctg  gtggtggcac  aaactatgca  180
gactccgtga  agggccgatt  caccatctcc  agagacactg  ccaagaactc  cttgtatcct  240
caaatgaaca  gcctgagagc  cgaggacatg  gctgtgtatt  actgtgcaag  agggagggaat  300
tcttttgact  actggggcca  gggaaacctg  gtcaccgtct  ctagt

```

**Figure 9A**

```

EVQLVQSGGG  LVHFGGSLRL  SCVGSRFIFS  AYPMHVVRQA  PGKGLEWVSG  IGSGGGTNYA  60
DSVKGRFTIS  RDTAKNSLYL  QMNSLRAEDM  AVYICARGRN  SFDYWGQGTL  VTVSS

```

**Figure 9B**

**18B2 Kappa Chain**

gacatccaga	tgacccagtc	tccatcctca	ctgtctgcat	ctgtaggaga	cagagtcacc	60
atcacttgtc	gggcgagtca	gggtattagc	acctggttag	cctggtatca	gcagaaacca	120
gagaaagccc	ctaagtccct	gatctatgct	gcatccagtt	tgacagagtgg	ggtcccatcg	180
aggttcagcg	gcagtggatc	tgggacagat	ttcaactctca	ccatcagcag	cctgcagcct	240
gaagattttg	caacttatta	ctgccaacag	tataatagtt	accctccgac	gttcgggcaa	300
gggaccaagg	tggagatcaa	acga				324

**Figure 10A**

DIQMTQSPSS	LSASVGDVRT	ITCRASQGIS	TWLAWYQQKP	EKAPKSLIYA	ASSLQSGVPS	60
RFSGSGSGTD	FLLTISSSLQP	EDFATYYCQQ	YNSYPPTFGQ	GTKVEIKR		108

**Figure 10B**

**16E1 Heavy Chain**

```

gaggtccagc  tgggtgcagtc  tgggggaggc  ttggtacatc  ctgggggggtc  cctgagactc  60
tcctgtgcag  gctctggatt  caccttcagt  gccatgctt  tgcactgggt  tcgccaggct  120
ccaggaaaag  gtctggagtg  ggtatcaggt  attggtactc  atggtgggac  atactatgca  180
gactccgtga  agggccgatt  caccatctcc  agagacaatg  ccaagaactc  cttgtttctt  240
caaatgaaca  gcctgagcgc  cgaggacatg  gctgtgtatt  actgtacaag  aagaaactgg  300
ggacaatttg  actactgggg  ccaggggaacc  ctggtcaccg  tctctagt

```

**Figure 11A**

```

EVQLVQSGGG  LVHPGGSLRL  SCAGSGFTFS  GHALHWVRQA  PGKGLEWVSG  IGTHGGTYYA  60
DSVKGRFTIS  RDNAKNSLFL  QMNSLSAEDM  AVYYCTRRNW  GQFDYWGQGT  LVTVSS      116

```

**Figure 11B**

**16E1 Kappa**

gaaattgtgc tgactcagtc tccagccacc ctgtctttgt ctccagggga aagagccacc 60  
 ctctcctgca gggccagtca gagtgttagc agtacttag cctggtacca acagaaacct 120  
 gggcaggctc ccaggctcct catctatgat gcatccaaca gggccactgg catcccagcc 180  
 aggttcagtg gcagtgggtc tgggacagac ttcactctca ccatcagcag cctagagcct 240  
 gaagattttg cagtttatta ctgtcagcag cgtagcaact ggcctccgta cacttttggc 300  
 caggggacca agctggagat caaacga 327

**Figure 12A**

EIVLTQSPAT ILSLSPGERAT ISCRASQSVS SYLAWYQQKP GQAPRLLIYD ASNRATGIPA 60  
 RFSGSGSGTD FTLTISLEP EDFAVYQCQQ RSNWPPYTFG QGTKLEIKR 109

**Figure 12B**

**9H7 Heavy Chain**

```

gaggtgcagc  tgggtgcagtc  tggggaggc  ttggtacatc  ctgggggggtc  cctgagactc  60
tcctgtgaag  gctctggatt  caccttcagt  agcaatggta  tgcactgggt  gcgccagact  120
ccaggaaaag  gtctggagtg  ggtatcaggt  attggtactg  ctggtggcac  atactatgca  180
gactccgtga  agggccgatt  caccattcc  agagacaatg  tcaagaagtc  cttgtatcctt  240
caaatgaaca  gcctgagagc  cgaggacatg  gctatttatt  attgtgtaag  aaaagactgg  300
ggatggttcg  acccctgggg  ccaggagcc  ctggtcaccg  tctctagt

```

**Figure 13A**

```

EVQLVQSGGG  LVHPGGSLRL  SCEGSGFTFS  SNGMHWVRQT  PGKGLEWVSG  IGTAGGTYYA  60
DSVKGRFTIS  RDNVKKSLYL  QMNSLRAEDM  AIYVCVRKDW  GWFDPWGQGA  LVTVSS      116

```

**Figure 13B**

**9H7 Kappa Chain**

```

gaaattgtgc  tgaccagtc  tccagccacc  ctgtctttgt  ctccagggga  aagagccacc  60
ctctcctgca  gggccagtca  gagtattagc  agctacttag  cctggtagca  acagaaacct  120
ggccaggctc  ccaggctcct  catctatgat  gcatccaaca  gggccactgg  catccccagg  180
aggttcagtg  gcagtgggtc  tgggacagac  ttcactctca  ccatcagcag  cctagagcct  240
gaaagatttg  cagttatta  ctgtcagcag  cgtagcaaat  ggcctccgta  cacttttggc  300
caggggacca  agctgggat  caaacga

```

**Figure 14A**

```

EIVLTQSPAT  LSLSPGERAT  LSCRASQSI  SYLAWYQQK  PQQAPRLL  IYD  ASNRATG  IPA  60
RFSGSGSGTD  FTLTISGLEP  EDFAVYYC  QQ  RSKWPPY  TFG  QGTKLEIK  R  109

```

**Figure 14B**



# HEAVY CHAIN ALIGNMENT

	FR1	CDR1	FR2	CDR2
<b>16E1</b>	EVQLVQSGGG LVHPGGSLRL SCAGSGFTFS	<u>GHAEH</u> HWVRQA	PGKGLEWVVS <u>G</u>	<u>IGTHGG</u> TTY ADSVKGRFTI
<b>2E11</b>	EVQLVQSGGG LVHPGGSLRL SCVGS <u>GD</u> GFTFS	<u>HYEHE</u> HWVRQA	PGKGLEWVVS <u>G</u>	<u>IHTGGG</u> TTY ADSVKGRFTI
<b>18B2</b>	EVQLVQSGGG LVHPGGSLRL SCVGS <u>R</u> GFTFS	<u>AYEMH</u> HWVRQA	PGKGLEWVVS <u>G</u>	<u>IGSGGG</u> TTY ADSVKGRFTI
<b>2D8</b>	EVQLVQSGGG LVHPGGSLRL SCAGSGFTFS	<u>SYGMH</u> HWVRQA	PGKGLEWVVS <u>G</u>	<u>IGTGGG</u> TTY ADSVKGRFTI
<b>22B3</b>	EVQLVQSGGG LVHPGGSLRL SC <u>EGS</u> GFTFS	<u>SXGMH</u> HWVRQA	PGKGLEWVVS <u>G</u>	<u>IGTAGG</u> TTY ADSVKGRFTI
<b>9H7</b>	EVQLVQSGGG LVHPGGSLRL SC <u>EGS</u> GFTFS	<u>SXGMH</u> HWVRQA	PGKGLEWVVS <u>G</u>	<u>IGTAGG</u> TTY ADSVKGRFTI

	FR3	CDR3
<b>16E1</b>	SRD <u>NAKNSL</u> F LQM <u>NSL</u> RAED	MAVYYC <u>IR</u> RN <u>WGQ</u> FDY
<b>2E11</b>	<u>SD</u> NAKNSLY LQM <u>N</u> TLRAED	MAVYYC <u>ARG</u> NS-FDY
<b>18B2</b>	SRD <u>TA</u> KNSLY LQM <u>NSL</u> RAED	MAVYYC <u>ARG</u> NS-FDY
<b>2D8</b>	SRD <u>N</u> VKNSLY LQM <u>NSL</u> RAED	MAVYYC <u>ARK</u> NWGWEDY
<b>22B3</b>	SRD <u>N</u> VKNSLY LQM <u>NSL</u> RAED	MAVYYC <u>VR</u> K <u>NW</u> GWEDY
<b>9H7</b>	SRD <u>N</u> VKNSLY LQM <u>NSL</u> RAED	MAVYYC <u>VR</u> K <u>DW</u> GWEDY

Figure 15

# LIGHT CHAIN ALIGNMENT

	FR1	CDR1	FR2	CDR2
<b>16E1</b>	DIQMTQSPSS LSASVGDRTVIT <u>ITCRASQGLSSWLAWYQQKPEKAPKSLIY</u> <u>AASSLQSGYP</u> <u>SRFSGSGSGT</u>			
<b>2E11</b>	DIQMTQSPSS LSASVGDRTVIT <u>ITCRASQGLSTWLAWYQQKPEKAPKSLIY</u> <u>AASSLQSGYP</u> <u>SRFSGSGSGT</u>			
<b>18B2</b>	EIVLTQSPAT LSLSPGERAT <u>LSCRASQSISSYLAWYQQKPGQAPRLLIY</u> <u>DASNRATGIP</u> <u>ARFSGSGSGT</u>			
<b>2D8</b>	EIVLTQSPAT LSLSPGERAT <u>LSCRASQSISSYLAWYQQKPGQAPRLLIY</u> <u>DASNRATGIP</u> <u>ARFSGSGSGT</u>			
<b>22B3</b>	EIVLTQSPAT LSLSPGERAT <u>LSCRASQSI</u> <u>NSYLAWFQQKPGQAPRLLIY</u> <u>DASNRATGIP</u> <u>ARFSGSGSGT</u>			
<b>9H7</b>	EIVLTQSPAT LSLSPGERAT <u>LSCRASQSV</u> <u>SSYLAWYQQKPGQAPRLLIY</u> <u>DASNRATGIP</u> <u>ARFSGSGSGT</u>			
	FR3	CDR3		
<b>16E1</b>	DFTLTISSLQ <u>SEDFATYYCOQYNSYPPPT</u>			
<b>2E11</b>	DFTLTISSLQ <u>PEDFATYYCOQYNSYPPPT</u>			
<b>18B2</b>	DFTLTISSLE <u>PEDFAVYVYCOQRSKWPPYT</u>			
<b>2D8</b>	DFTLTISSLE <u>PEDFAVYVYCOQRSKWPPYT</u>			
<b>22B3</b>	DFTLTISSLE <u>PEDFATYYCOQRSNWPPPT</u>			
<b>9H7</b>	DFTLTISSLE <u>PEDFAVYVYCOQRSNWPPYT</u>			

Figure 16




Figure 17




Figure 18

**Cell Culture Process Flow Chart**




Figure 19




Figure 20





Figure 21

Figure 22





Mouse OPG ligand 199 T L S N G K L R V N Q D G F Y Y L Y A N I C F R H H E T S G S V P T D Y L Q L M V Y V V K T S I K I 248  
 Human OPG ligand 200 T S N G K L R V N Q D G F Y Y L Y A N I C F R H H E T S G D L A T E Y L Q L M V Y V V K T S I K I 249  
 Mouse OPG ligand/DE T L S N G K L R V N Q D G F Y Y L Y A N I C F R H H E T S G D L A T E Y L Q L M V Y V V K T S I K I

Figure 23




Figure 24




Figure 25

## HUMAN ANTI-OPGL NEUTRALIZING ANTIBODIES AS SELECTIVE OPGL PATHWAY INHIBITORS

[0001] This application is a continuation application of U.S. application Ser. No. 10/408,901, filed Apr. 7, 2003; which is a non-provisional application of, and claims priority, to U.S. Provisional Patent Application No. 60/370,407 filed on Apr. 5, 2002. The disclosure of each application is herein incorporated by reference.

[0002] The sequence listing is filed with this application in electronic format only and is incorporated herein by reference. The computer readable sequence listing text file "01\_1145\_A\_CONSeqLst.txt" was created on Feb. 15, 2010, and is 87,784 bytes in size.

### FIELD OF THE INVENTION

[0003] The invention relates to antibodies that bind osteoprotegerin ligand (OPGL). Compositions and methods for the treatment of bone diseases, such as osteoporosis, bone loss from arthritis, Paget's disease, and osteopenia, are also provided.

### BACKGROUND OF THE INVENTION

[0004] Living bone tissue exhibits a dynamic equilibrium between formation of bone, known as deposition, and breakdown of bone, known as resorption. These processes can be mediated by at least two cell types: osteoblasts, which secrete molecules that comprise the organic matrix of bone (deposition); and osteoclasts, which promote dissolution of the bone matrix and solubilization of bone salts (resorption). In certain individuals, such as post-menopausal women, the rate of resorption can exceed the rate of deposition, which may result in reduced bone mass and strength, increased risk of fractures, and slow or incomplete repair of broken bones.

[0005] Osteoprotegerin ligand (OPGL) is a member of the TNF family of cytokines and promotes formation of osteoclasts through binding to the receptor activator of NF- $\kappa$ B (RANK, also called osteoclast differentiation and activation receptor, or ODAR). Osteoprotegerin (OPG), on the other hand, inhibits the formation of osteoclasts by sequestering OPGL and preventing OPGL association with ODAR. Thus, the amount of OPGL associated with ODAR correlates with the equilibrium between bone deposition and resorption. Individuals who suffer from osteopenic diseases, such as osteoporosis, show a greater rate of bone resorption than deposition, which may result from increased levels or activity of OPGL. Thus, it would be useful to have molecules that can regulate the activity of OPGL in osteoclastogenesis. It would also be useful to be able to detect the amount of OPGL in a biological sample, such as a blood sample, to diagnose an osteopenic disorder relating to increased levels of OPGL.

### SUMMARY OF THE INVENTION

[0006] The invention provides monoclonal antibodies that bind to osteoprotegerin ligand (OPGL). Preferably, the antibodies inhibit binding of OPGL to an osteoclast differentiation and activation receptor (ODAR). Also provided by this invention are hybridoma cell lines that produce, and most preferably, secrete into cell culture media the monoclonal

antibodies of the invention. The antibodies of the invention are useful for treating various disorders associated with low bone density.

[0007] In certain aspects, the invention provides antibodies, preferably monoclonal antibodies, most preferably human antibodies, comprising a heavy chain and a light chain, wherein the heavy chain comprises an IgG<sub>1</sub>, IgG<sub>2</sub>, or an IgG<sub>4</sub> heavy chain constant region. Preferably, an antibody of the invention comprises an amino acid sequence of the IgG<sub>1</sub> heavy chain constant region as set forth in SEQ ID NO: 2 or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

[0008] The invention also provides antibodies, preferably monoclonal antibodies, most preferably human antibodies, comprising a heavy chain and a light chain, wherein the light chain comprises an amino acid sequence as set forth in SEQ ID NO: 4 or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

[0009] The invention relates specifically to human antibodies, most preferably monoclonal antibodies that specifically bind the D-E loop region of OPGL. The invention also relates to human antibodies, preferably monoclonal antibodies, that bind to a region of osteoprotegerin ligand (OPGL) that is outside the D-E loop region. In addition, the invention relates to human antibodies, preferably monoclonal antibodies, that bind to both a region of OPGL that is outside the D-E loop region and all or a portion of the D-E loop region. In one aspect, antibodies of the invention bind to a first region of OPGL that is outside the D-E loop region and then, while remaining bound to the first region, bind to a second region that is all or a portion of the D-E loop region. Such binding is referred to herein as consecutive. In another aspect, antibodies of the invention can bind to a first region of OPGL that is outside the D-E loop region and a second region that is all or a portion of the D-E loop region at the same time. Such binding is referred to herein as simultaneous.

[0010] In certain aspects, antibodies of the invention comprise a heavy chain and a light chain, wherein the variable region of the heavy chain comprises an amino acid sequence as set forth in any of SEQ ID NO: 6, SEQ ID NO: 14, SEQ ID NO: 22, or SEQ ID NO: 26, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof. In other aspects, the light chain variable region comprises an amino acid sequence as set forth in any of SEQ ID NO: 8, SEQ ID NO: 16, SEQ ID NO: 24, or SEQ ID NO: 28, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof. In additional aspects, the heavy chain comprises an amino acid sequence as set forth in any of SEQ ID NO: 30, SEQ ID NO: 38, SEQ ID NO: 46, or SEQ ID NO: 50, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof. In still further aspects, the light chain comprises an amino acid sequence as set forth in any of SEQ ID NO: 32, SEQ ID NO: 40, SEQ ID NO: 48, or SEQ ID NO: 52, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

[0011] The invention also provides antibodies that bind specifically to OPGL, wherein the heavy chain comprises a heavy chain variable region as set forth in SEQ ID NO: 6, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof, and the light chain comprises a light chain variable region comprising an amino acid

sequence as set forth in SEQ ID NO: 8, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0012]** In certain aspects, the invention provides antibodies, comprising a heavy chain and a light chain, (a) wherein the heavy chain comprises a first variable region, and wherein the first variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 6, and (b) wherein the light chain comprises a second variable region, and wherein the second variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 8, and (c) wherein the antibody interacts with OPGL.

**[0013]** In other aspects, the first variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 6, and the second variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 8.

**[0014]** In still other aspects, the first variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 6, and the second variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 8.

**[0015]** The invention further provides antibodies that bind specifically to OPGL, wherein the heavy chain comprises a heavy chain variable region as set forth in SEQ ID NO: 14, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof, and the light chain comprises a light chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 16, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0016]** In certain aspects, the invention provides antibodies, comprising a heavy chain and a light chain, (a) wherein the heavy chain comprises a first variable region, and wherein the first variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 14, and (b) wherein the light chain comprises a second variable region, and wherein the second variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 16, and (c) wherein the antibody interacts with OPGL.

**[0017]** In other aspects, the first variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 14, and the second variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 16.

**[0018]** In further aspects, the first variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 14, and the second variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 16.

**[0019]** The invention provides antibodies that bind specifically to OPGL, wherein the heavy chain comprises a heavy chain variable region as set forth in SEQ ID NO: 22, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof, and the light chain comprises a light chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 24, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0020]** In certain aspects, the invention provides antibodies, comprising a heavy chain and a light chain, (a) wherein the

heavy chain comprises a first variable region, and wherein the first variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 22, and (b) wherein the light chain comprises a second variable region, and wherein the second variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 24, and (c) wherein the antibody interacts with OPGL.

**[0021]** In particular aspects, the first variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 22, and the second variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 24.

**[0022]** In further aspects, the first variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 22, and the second variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 24.

**[0023]** In addition, the invention provides antibodies that bind specifically to the D-E loop region of OPGL, wherein the heavy chain comprises a heavy chain variable region as set forth in SEQ ID NO: 26, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof, and the light chain comprises a light chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 28, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0024]** In certain aspects, the invention provides antibodies, comprising a heavy chain and a light chain, (a) wherein the heavy chain comprises a first variable region, and wherein the first variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 26, and (b) wherein the light chain comprises a second variable region, and wherein the second variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 28, and (c) wherein the antibody interacts with OPGL.

**[0025]** In other aspects, the first variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 26, and the second variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 28.

**[0026]** In additional aspects, the first variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 26, and the second variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 28.

**[0027]** The invention also provides antibodies that bind specifically to OPGL, wherein the heavy chain comprises an amino acid sequence as set forth in SEQ ID NO: 30, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof, and the light chain comprises an amino acid sequence as set forth in SEQ ID NO: 32, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0028]** The invention also provides antibodies that bind specifically to OPGL, wherein the heavy chain comprises an amino acid sequence as set forth in SEQ ID NO: 38, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof, and the light chain comprises a light chain variable region comprising an amino acid

sequence as set forth in SEQ ID NO: 40, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0029]** The invention provides antibodies that bind specifically to OPGL, wherein the heavy chain comprises an amino acid sequence as set forth in SEQ ID NO: 46, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof, and the light chain comprises a light chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 48, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0030]** The invention provides antibodies that bind specifically to OPGL, wherein the heavy chain comprises an amino acid sequence as set forth in SEQ ID NO: 50, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof, and the light chain comprises a light chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 52, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0031]** In certain aspects, the invention provides antibodies that specifically bind OPGL and comprises a heavy chain and a light chain, wherein the heavy chain variable region comprises an amino acid sequence as set forth in SEQ ID NO: 10 or SEQ ID NO: 18, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof. In other aspects, the light chain variable region comprises an amino acid sequence as set forth in SEQ ID NO: 12 or SEQ ID NO: 20, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0032]** The invention also provides antibodies that specifically bind OPGL, wherein the heavy chain variable region comprises an amino acid sequence as set forth in SEQ ID NO: 34 or SEQ ID NO: 42, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof. In other aspects, the light chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 36 or SEQ ID NO: 44, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0033]** The invention further provides antibodies that specifically bind OPGL, wherein the heavy chain comprises a heavy chain variable region as set forth in SEQ ID NO: 10, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof, and the light chain comprises a light chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 12, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0034]** In certain aspects, the invention provides antibodies, comprising a heavy chain and a light chain, (a) wherein the heavy chain comprises a first variable region, and wherein the first variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 10, and (b) wherein the light chain comprises a second variable region, and wherein the second variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 12, and (c) wherein the antibody interacts with OPGL.

**[0035]** In further aspects, the first variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 10, and the second variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 12.

**[0036]** In other aspects, the first variable region comprises a sequence that has at least 99% identity to the amino acid

sequence set forth in SEQ ID NO: 10, and the second variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 12.

**[0037]** The invention also provides antibodies that specifically bind, wherein the heavy chain comprises a heavy chain variable region as set forth in SEQ ID NO: 18, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof, and the light chain comprises a light chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 20, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0038]** In certain aspects, the invention provides antibodies, comprising a heavy chain and a light chain, (a) wherein the heavy chain comprises a first variable region, and wherein the first variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 18, and (b) wherein the light chain comprises a second variable region, and wherein the second variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 20, and (c) wherein the antibody interacts with OPGL.

**[0039]** In other aspects, the first variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 18, and the second variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 20.

**[0040]** In still other aspects, the first variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 18, and the second variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 20.

**[0041]** The invention also provides antibodies that specifically bind OPGL, wherein the heavy chain comprises an amino acid sequence as set forth in SEQ ID NO: 34, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof, and the light chain comprises a light chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 36, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0042]** The invention provides antibodies that specifically bind OPGL, wherein the heavy chain comprises an amino acid sequence as set forth in SEQ ID NO: 42, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof, and the light chain comprises a light chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 44, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0043]** The invention also provides single chain antibodies, single chain Fv antibodies, Fab antibodies, Fab' antibodies, and (Fab')<sub>2</sub>.

**[0044]** In particular aspects, the invention provides a heavy chain comprising a variable region and a constant region, wherein the variable region comprises an amino acid sequence as set forth in any of SEQ ID NO: 6, SEQ ID NO: 10, SEQ ID NO: 14, SEQ ID NO: 18, SEQ ID NO: 22, or SEQ ID NO: 26, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0045]** In addition, the invention also provides a heavy chain comprising an amino acid sequence as set forth in any of SEQ ID NO: 30, SEQ ID NO: 34, SEQ ID NO: 38, SEQ ID

NO: 42, SEQ ID NO: 46, or SEQ ID NO: 50, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0046]** In certain aspects, the invention provides a light chain comprising a variable region and a constant region, wherein the variable region comprises an amino acid sequence as set forth in any of SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID NO: 16, SEQ ID NO: 20, SEQ ID NO: 24, or SEQ ID NO: 28, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0047]** In other aspects, the invention provides a light chain comprising an amino acid sequence as set forth in any of SEQ ID NO: 32, SEQ ID NO: 36, SEQ ID NO: 40, SEQ ID NO: 44, SEQ ID NO: 48, or SEQ ID NO: 52, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

**[0048]** The invention also relates to isolated human antibodies that specifically bind OPGL, wherein the antibody comprises: (a) human heavy chain framework regions, a human heavy chain CDR1 region, a human heavy chain CDR2 region, and a human heavy chain CDR3 region; and (b) human light chain framework regions, a human light chain CDR1 region, a human light chain CDR2 region, and a human light chain CDR3 region. In certain aspects, the human heavy chain CDR1 region can be the heavy chain CDR1 region of 16E1, 2D8, 22B3, or 9H7 as shown in FIG. 15 and the human light chain CDR1 region can be the light chain CDR1 region of 16E1, 2D8, 22B3, or 9H7 as shown in FIG. 16. In other aspects, the human heavy chain CDR2 region can be the heavy chain CDR2 region of 16E1, 2D8, 22B3, or 9H7 as shown in FIG. 15 and the human light chain CDR2 region can be the light chain CDR2 region of 16E1, 2D8, 22B3, or 9H7 as shown in FIG. 16. In still other aspects, the human heavy chain CDR3 region is the heavy chain CDR3 region of 16E1, 2D8, 22B3, or 9H7 as shown in FIG. 15, and the human light chain CDR3 region is the light chain CDR3 region of 16E1, 2D8, 22B3, or 9H7 as shown in FIG. 16.

**[0049]** The invention also relates to isolated human antibodies that specifically bind OPGL, wherein the antibody comprises: (a) human heavy chain framework regions, a human heavy chain CDR1 region, a human heavy chain CDR2 region, and a human heavy chain CDR3 region; and (b) human light chain framework regions, a human light chain CDR1 region, a human light chain CDR2 region, and a human light chain CDR3 region. In certain aspects, the human heavy chain CDR1 region can be the heavy chain CDR1 region of 2E11 or 18B2 as shown in FIG. 15 and the human light chain CDR1 region can be the light chain CDR1 region of 2E11 or 18B2 as shown in FIG. 16. In other aspects, the human heavy chain CDR2 region can be the heavy chain CDR2 region of 2E11 or 18B2 as shown in FIG. 15 and the human light chain CDR2 region can be the light chain CDR2 region of 2E11 or 18B2 as shown in FIG. 16. In still other aspects, the human heavy chain CDR3 region is the heavy chain CDR3 region of 2E11 or 18B2 as shown in FIG. 15, and the human light chain CDR3 region is the light chain CDR3 region of 2E11 or 18B2 as shown in FIG. 16.

**[0050]** In addition, the invention provides methods for treating an osteopenic disorder, comprising the step of administering a pharmaceutically effective amount of a monoclonal antibody of the invention or antigen-binding fragment thereof to an individual in need thereof.

**[0051]** The invention further relates to fusion proteins and other molecules capable of binding to a region of osteopro-

tegerin ligand (OPGL) that is outside the D-E loop region, or both a region of OPGL that is outside the D-E loop region and all or a portion of the D-E loop region, wherein binding is consecutive or simultaneous (together with the aforementioned antibodies, collectively referred to herein as "specific binding partners"), such as may be prepared using methods as described, for example, in WO 00/24782, which is incorporated by reference. Such molecules can be expressed, for example, in mammalian cells (e.g. Chinese Hamster Ovary cells) or bacterial cells (e.g. *E. coli* cells).

**[0052]** The invention also provides methods for detecting the level of OPGL in a biological sample, comprising the step of contacting the sample with a monoclonal antibody of the invention or antigen-binding fragment thereof. The anti-OPGL antibodies of the invention may be employed in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, immunoprecipitation assays and enzyme-linked immunosorbent assays (ELISA) (See, Sola, 1987, *Monoclonal Antibodies: A Manual of Techniques*, pp. 147-158, CRC Press, Inc.) for the detection and quantitation of OPGL. The antibodies can bind OPGL with an affinity that is appropriate for the assay method being employed.

**[0053]** Specific preferred embodiments of the present invention will become evident from the following more detailed description of certain preferred embodiments and the claims.

#### BRIEF DESCRIPTION OF THE FIGURES

**[0054]** FIGS. 1A-1B depict a cDNA sequence (FIG. 1A) encoding the anti-OPGL antibody heavy chain constant region (SEQ ID NO: 1) and the amino acid sequence (FIG. 1B) of the anti-OPGL antibody heavy chain constant region (SEQ ID NO: 2).

**[0055]** FIGS. 2A-2B depict a cDNA sequence (FIG. 2A) encoding the anti-OPGL antibody kappa chain constant region (SEQ ID NO: 3) and the amino acid sequence (FIG. 2B) of the anti-OPGL antibody kappa chain constant region (SEQ ID NO: 4).

**[0056]** FIGS. 3A-3B depict a cDNA sequence (FIG. 3A) encoding the 22B3 anti-OPGL antibody heavy chain variable region (SEQ ID NO: 5) and the amino acid sequence (FIG. 3B) of the 22B3 anti-OPGL antibody heavy chain variable region (SEQ ID NO: 6).

**[0057]** FIGS. 4A-4B depict a cDNA sequence (FIG. 4A) encoding the 22B3 anti-OPGL antibody kappa chain variable region (SEQ ID NO: 7) and the amino acid sequence (FIG. 4B) of the 22B3 anti-OPGL antibody kappa chain variable region (SEQ ID NO: 8).

**[0058]** FIGS. 5A-5B depict a cDNA sequence (FIG. 5A) encoding the 2E11 anti-OPGL antibody heavy chain variable region (SEQ ID NO: 9) and the amino acid sequence (FIG. 5B) of the 2E11 anti-OPGL antibody heavy chain variable region (SEQ ID NO: 10).

**[0059]** FIGS. 6A-6B depict a cDNA sequence (FIG. 6A) encoding the 2E11 anti-OPGL antibody kappa chain variable region (SEQ ID NO: 11) and the amino acid sequence (FIG. 6B) of the 2E11 anti-OPGL antibody kappa chain variable region (SEQ ID NO: 12).

**[0060]** FIGS. 7A-7B depict a cDNA sequence (FIG. 7A) encoding the 2D8 anti-OPGL antibody heavy chain variable region (SEQ ID NO: 13) and the amino acid sequence (FIG. 7B) of the 2D8 anti-OPGL antibody heavy chain variable region (SEQ ID NO: 14).

**[0061]** FIGS. 8A-8B depict a cDNA sequence (FIG. 8A) encoding the 2D8 anti-OPGL antibody kappa chain variable region (SEQ ID NO: 15) and the amino acid sequence (FIG. 8B) of the 2D8 anti-OPGL antibody kappa chain variable region (SEQ ID NO: 16).

**[0062]** FIGS. 9A-9B depict a cDNA sequence (FIG. 9A) encoding the 18B2 anti-OPGL antibody heavy chain variable region (SEQ ID NO: 17) and the amino acid sequence (FIG. 9B) of the 18B2 anti-OPGL antibody heavy chain variable region (SEQ ID NO: 18).

**[0063]** FIGS. 10A-10B depict a cDNA sequence (FIG. 10A) encoding the 18B2 anti-OPGL antibody kappa chain variable region (SEQ ID NO: 19) and the amino acid sequence (FIG. 10B) of the 18B2 anti-OPGL antibody kappa chain variable region (SEQ ID NO: 20).

**[0064]** FIGS. 11A-11B depict a cDNA sequence (FIG. 11A) encoding the 16E1 anti-OPGL antibody heavy chain variable region (SEQ ID NO: 21) and the amino acid sequence (FIG. 11B) of the 16E1 anti-OPGL antibody heavy chain variable region (SEQ ID NO: 22).

**[0065]** FIGS. 12A-12B depict a cDNA sequence (FIG. 12A) encoding the 16E1 anti-OPGL antibody kappa chain variable region (SEQ ID NO: 23) and the amino acid sequence (FIG. 12B) of the 16E1 anti-OPGL antibody kappa chain variable region (SEQ ID NO: 24).

**[0066]** FIGS. 13A-13B depict a cDNA sequence (FIG. 13A) encoding the 9H7 anti-OPGL antibody heavy chain variable region (SEQ ID NO: 25) and the amino acid sequence (FIG. 13B) of the 9H7 anti-OPGL antibody heavy chain variable region (SEQ ID NO: 26).

**[0067]** FIGS. 14A-14B depict a cDNA sequence (FIG. 14A) encoding the 9H7 anti-OPGL antibody kappa chain variable region (SEQ ID NO: 27) and the amino acid sequence (FIG. 14B) of the 9H7 anti-OPGL antibody kappa chain variable region (SEQ ID NO: 28).

**[0068]** FIG. 15 depicts the heavy chain alignment for anti-OPGL antibodies designated 16E1 (SEQ ID NO: 22), 2E11 (SEQ ID NO: 10), 18B2 (SEQ ID NO: 18), 2D8 (SEQ ID NO: 14), 22B3 (SEQ ID NO: 6), and 9H7 (SEQ ID NO: 26). CDRs are underlined, non-consensus amino acids are shaded and in bold type.

**[0069]** 16E1: heavy chain CDR1 is SEQ ID NO: 77, CDR2 is SEQ ID NO: 78, CDR3 is SEQ ID NO: 79.

**[0070]** 2E11: heavy chain CDR1 is SEQ ID NO: 80, CDR2 is SEQ ID NO: 81, CDR3 is SEQ ID NO: 82.

**[0071]** 18B2: heavy chain CDR1 is SEQ ID NO: 83, CDR2 is SEQ ID NO: 84, CDR3 is SEQ ID NO: 82.

**[0072]** 2D8: heavy chain CDR1 is SEQ ID NO: 85, CDR2 is SEQ ID NO: 86, CDR3 is SEQ ID NO: 87.

**[0073]** 22B3: heavy chain CDR1 is SEQ ID NO: 88, CDR2 is SEQ ID NO: 89, CDR3 is SEQ ID NO: 90.

**[0074]** 9H7: heavy chain CDR1 is SEQ ID NO: 88, CDR2 is SEQ ID NO: 89, CDR3 is SEQ ID NO: 91.

**[0075]** FIG. 16 depicts the light chain alignment for anti-OPGL antibodies designated 16E1 (SEQ ID NO: 24), 2E11 (SEQ ID NO: 12), 18B2 (SEQ ID NO: 20), 2D8 (SEQ ID NO: 16), 22B3 (SEQ ID NO: 8), and 9H7 (SEQ ID NO: 28). CDRs are underlined, non-consensus amino acids are shaded and in bold type.

**[0076]** 16E1: light chain CDR1 is SEQ ID NO: 92, CDR2 is SEQ ID NO: 93, CDR3 is SEQ ID NO: 94.

**[0077]** 2E11: light chain CDR1 is SEQ ID NO: 95, CDR2 is SEQ ID NO: 93, CDR3 is SEQ ID NO: 94.

**[0078]** 18B2: light chain CDR1 is SEQ ID NO: 96, CDR2 is SEQ ID NO: 97, CDR3 is SEQ ID NO: 98.

**[0079]** 2D8: light chain CDR1 is SEQ ID NO: 96, CDR2 is SEQ ID NO: 97, CDR3 is SEQ ID NO: 98.

**[0080]** 22B3: light chain CDR1 is SEQ ID NO: 99, CDR2 is SEQ ID NO: 97, CDR3 is SEQ ID NO: 100.

**[0081]** 9H7: light chain CDR1 is SEQ ID NO: 101, CDR2 is SEQ ID NO: 97, CDR3 is SEQ ID NO: 102.

**[0082]** FIG. 17 depicts a circular plasmid map of the pDSR $\alpha$ 19:9 H7 kappa chain expression vector.

**[0083]** FIG. 18 shows a circular plasmid map of the pDSR $\alpha$ 19:9 H7 heavy chain expression vector.

**[0084]** FIG. 19 depicts an exemplary cell culture process for producing anti-OPGL antibody.

**[0085]** FIG. 20 is a graph showing optical density versus anti-OPGL antibody concentration demonstrating OPGL antibody mediated inhibition of osteoclast formation.

**[0086]** FIG. 21 depicts graphs of serum concentrations of anti-OPGL antibodies following subcutaneous administration at 1.0 mg/kg in Cynomolgus monkeys.

**[0087]** FIG. 22 depicts graphs representing the percentage change in serum NTx from baseline following subcutaneous administration at 1.0 mg/kg of anti-OPGL antibodies in Cynomolgus monkeys.

**[0088]** FIG. 23 shows a comparison of murine (SEQ ID NO: 70), human (SEQ ID NO: 71), and murine DE variant (SEQ ID NO: 72) amino acid sequences in a region of OPGL between the D and E regions.

**[0089]** FIG. 24 depicts the results of an enzyme immunoassay showing six anti-OPGL antibodies of the invention binding murine OPGL (143-317).

**[0090]** FIG. 25 depicts the results of an enzyme immunoassay showing four of the anti-OPGL antibodies of the invention bind FLAG-murine OPGL/DE (158-316).

#### DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

**[0091]** The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described. All references cited in this application are expressly incorporated by reference herein for any purpose.

#### DEFINITIONS

**[0092]** Standard techniques were used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (e.g., electroporation, lipofection).

**[0093]** Enzymatic reactions and purification techniques were performed according to manufacturer's specifications or as commonly accomplished in the art or as described herein. The foregoing techniques and procedures were generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification. See e.g., Sambrook et al., 2001, MOLECULAR CLONING: A LABORATORY MANUAL, 3d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., which is incorporated herein by reference for any purpose. Unless specific definitions are provided, the nomenclature utilized in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art.



Standard techniques can be used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

**[0094]** As utilized in accordance with the present disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

**[0095]** The term “isolated polynucleotide” as used herein means a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, which by virtue of its origin the isolated polynucleotide (1) is not associated with all or a portion of a polynucleotide in which the isolated polynucleotide is found in nature, (2) is linked to a polynucleotide which it is not linked to in nature, or (3) does not occur in nature as part of a larger sequence.

**[0096]** The term “isolated protein” referred to herein means that a subject protein (1) is free of at least some other proteins with which it would normally be found, (2) is essentially free of other proteins from the same source, e.g., from the same species, (3) is expressed by a cell from a different species, (4) has been separated from at least about 50 percent of polynucleotides, lipids, carbohydrates, or other materials with which it is associated in nature, (5) is not associated (by covalent or noncovalent interaction) with portions of a protein with which the “isolated protein” is associated in nature, (6) is operably associated (by covalent or noncovalent interaction) with a polypeptide with which it is not associated in nature, or (7) does not occur in nature. Such an isolated protein can be encoded by genomic DNA, cDNA, mRNA or other RNA, of synthetic origin, or any combination thereof. Preferably, the isolated protein is substantially free from proteins or polypeptides or other contaminants that are found in its natural environment that would interfere with its use (therapeutic, diagnostic, prophylactic, research or otherwise).

**[0097]** The terms “polypeptide” or “protein” means molecules having the sequence of native proteins, that is, proteins produced by naturally-occurring and specifically non-recombinant cells, or genetically-engineered or recombinant cells, and comprise molecules having the amino acid sequence of the native protein, or molecules having deletions from, additions to, and/or substitutions of one or more amino acids of the native sequence. The terms “polypeptide” and “protein” specifically encompass anti-OPGL antibodies, or sequences that have deletions from, additions to, and/or substitutions of one or more amino acid of an anti-OPGL antibody.

**[0098]** The term “polypeptide fragment” refers to a polypeptide that has an amino-terminal deletion, a carboxyl-terminal deletion, and/or an internal deletion. In certain embodiments, fragments are at least 5 to about 500 amino acids long. It will be appreciated that in certain embodiments, fragments are at least 5, 6, 8, 10, 14, 20, 50, 70, 100, 110, 150, 200, 250, 300, 350, 400, or 450 amino acids long. Particularly useful polypeptide fragments include functional domains, including binding domains. In the case of an anti-OPGL antibody, useful fragments include but are not limited to a CDR region, a variable domain of a heavy or light chain, a portion of an antibody chain or just its variable region including two CDRs, and the like.

**[0099]** The term “immunologically functional immunoglobulin fragment” as used herein refers to a polypeptide fragment that contains at least the CDRs of the immunoglobulin heavy and light chains. An immunologically functional immunoglobulin fragment of the invention is capable of binding to an antigen. In preferred embodiments, the antigen is a ligand that specifically binds to a receptor. In these embodi-

ments, binding of an immunologically functional immunoglobulin fragment of the invention prevents binding of the ligand to its receptor, interrupting the biological response resulting from ligand binding to the receptor. Preferably, an immunologically functional immunoglobulin fragment of the invention binds specifically to OPGL. Most preferably, the fragment binds specifically to human OPGL.

**[0100]** The term “naturally-occurring” as used herein and applied to an object refers to the fact that the object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and that has not been intentionally modified by man is naturally occurring.

**[0101]** The term “operably linked” means that the components to which the term is applied are in a relationship that allows them to carry out their inherent functions under suitable conditions. For example, a control sequence “operably linked” to a coding sequence is ligated thereto so that expression of the protein coding sequence is achieved under conditions compatible with the transcriptional activity of the control sequences.

**[0102]** The term “control sequence” as used herein refers to polynucleotide sequences that can effect expression, processing or intracellular localization of coding sequences to which they are ligated. The nature of such control sequences may differ depending upon the host organism. In particular embodiments, control sequences for prokaryotes may include promoter, ribosomal binding site, and transcription termination sequence. In other particular embodiments, control sequences for eukaryotes may include promoters comprising one or a plurality of recognition sites for transcription factors, transcription enhancer sequences, transcription termination sequences and polyadenylation sequences. In certain embodiments, “control sequences” can include leader sequences and/or fusion partner sequences.

**[0103]** The term “polynucleotide” as referred to herein means single-stranded or double-stranded nucleic acid polymers of at least 10 bases in length. In certain embodiments, the nucleotides comprising the polynucleotide can be ribonucleotides or deoxyribonucleotides or a modified form of either type of nucleotide. Said modifications include base modifications such as bromuridine, ribose modifications such as arabinoside and 2',3'-dideoxyribose and internucleotide linkage modifications such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate and phosphoroamidate. The term “polynucleotide” specifically includes single and double stranded forms of DNA.

**[0104]** The term “oligonucleotide” as used herein includes naturally occurring, and modified nucleotides linked together by naturally occurring, and/or non-naturally occurring oligonucleotide linkages. Oligonucleotides are a polynucleotide subset comprising members that are generally single-stranded and have a length of 200 bases or fewer. In certain embodiments, oligonucleotides are 10 to 60 nucleotides in length. In certain embodiments, oligonucleotides are 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 nucleotides in length. Oligonucleotides may be single stranded or double stranded, e.g. for use in the construction of a gene mutant. Oligonucleotides of the invention may be sense or antisense oligonucleotides with reference to a protein-coding sequence.

**[0105]** The term “naturally occurring nucleotides” includes deoxyribonucleotides and ribonucleotides. The term “modified nucleotides” includes nucleotides with modified or sub-

stituted sugar groups and the like. The term “oligonucleotide linkages” includes oligonucleotides linkages such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoroamidate, and the like. See, e.g., LaPlanche et al., 1986, *Nucl. Acids Res.* 14: 9081; Stec et al., 1984, *J. Am. Chem. Soc.* 106: 6077; Stein et al., 1988, *Nucl. Acids Res.* 16: 3209; Zon et al., 1991, *Anti-Cancer Drug Design* 6: 539; Zon et al., 1991, OLIGONUCLEOTIDES AND ANALOGUES: A PRACTICAL APPROACH, (F. Eckstein, ed.), Oxford University Press, Oxford England, pp. 87-108; Stec et al., U.S. Pat. No. 5,151,510; Uhlmann and Peyman, 1990, *Chemical Reviews* 90: 543, the disclosures of each of which are hereby incorporated by reference for any purpose. An oligonucleotide can include a detectable label to enable detection of the oligonucleotide or hybridization thereof.

**[0106]** The term “vector” is used to refer to any molecule (e.g., nucleic acid, plasmid, or virus) used to transfer coding information to a host cell.

**[0107]** The term “expression vector” refers to a vector that is suitable for transformation of a host cell and contains nucleic acid sequences that direct and/or control the expression of inserted heterologous nucleic acid sequences. Expression includes, but is not limited to, processes such as transcription, translation, and RNA splicing, if introns are present.

**[0108]** The term “host cell” is used to refer to a cell which has been transformed, or that is capable of being transformed with a nucleic acid sequence and then of expressing a selected gene of interest. The term includes the progeny of the parent cell, whether or not the progeny is identical in morphology or in genetic make-up to the original parent, so long as the selected gene is present.

**[0109]** The term “transduction” is used to refer to the transfer of genes from one bacterium to another, usually by a phage. “Transduction” also refers to the acquisition and transfer of eukaryotic cellular sequences by retroviruses.

**[0110]** The term “transfection” is used to refer to the uptake of foreign or exogenous DNA by a cell, and a cell has been “transfected” when the exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are well known in the art and are disclosed herein. See, e.g., Graham et al., 1973, *Virology* 52: 456; Sambrook et al., 2001, *ibid.*; Davis et al., 1986, BASIC METHODS IN MOLECULAR BIOLOGY (Elsevier); and Chu et al., 1981, *Gene* 13: 197. Such techniques can be used to introduce one or more exogenous DNA moieties into suitable host cells.

**[0111]** The term “transformation” as used herein refers to a change in a cell’s genetic characteristics, and a cell has been transformed when it has been modified to contain a new DNA. For example, a cell is transformed where it is genetically modified from its native state. Following transfection or transduction, the transforming DNA may recombine with that of the cell by physically integrating into a chromosome of the cell, may be maintained transiently as an episomal element without being replicated, or may replicate independently as a plasmid. A cell is considered to have been stably transformed when the DNA is replicated with the division of the cell.

**[0112]** The term “naturally occurring” or “native” when used in connection with biological materials such as nucleic acid molecules, polypeptides, host cells, and the like, refers to materials which are found in nature and are not manipulated by man. Similarly, “non-naturally occurring” or “non-native”

as used herein refers to a material that is not found in nature or that has been structurally modified or synthesized by man.

**[0113]** The term “antigen” refers to a molecule or a portion of a molecule capable of being bound by a selective binding agent, such as an antibody, and additionally capable of being used in an animal to produce antibodies capable of binding to an epitope of that antigen. An antigen may have one or more epitopes.

**[0114]** The term “identity,” as known in the art, refers to a relationship between the sequences of two or more polypeptide molecules or two or more nucleic acid molecules, as determined by comparing the sequences thereof. In the art, “identity” also means the degree of sequence relatedness between nucleic acid molecules or polypeptides, as the case may be, as determined by the match between two or more nucleotide or two or more amino acid sequences. “Identity” measures the percent of identical matches between the smaller of two or more sequences with gap alignments (if any) addressed by a particular mathematical model or computer program (i.e., “algorithms”).

**[0115]** The term “similarity” is used in the art with regard to a related concept, but in contrast to “identity,” “similarity” refers to a measure of relatedness, which includes both identical matches and conservative substitution matches. If two polypeptide sequences have, for example, 10/20 identical amino acids, and the remainder are all non-conservative substitutions, then the percent identity and similarity would both be 50%. If in the same example, there are five more positions where there are conservative substitutions, then the percent identity remains 50%, but the percent similarity would be 75% (15/20). Therefore, in cases where there are conservative substitutions, the percent similarity between two polypeptides will be higher than the percent identity between those two polypeptides.

**[0116]** Identity and similarity of related and polypeptides can be readily calculated by known methods. Such methods include, but are not limited to, those described in COMPUTATIONAL MOLECULAR BIOLOGY, (Lesk, A. M., ed.), 1988, New York: Oxford University Press; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, (Smith, D. W., ed.), 1993, New York: Academic Press; COMPUTER ANALYSIS OF SEQUENCE DATA, PART 1, (Griffin, A. M., and Griffin, H. G., eds.), 1994, New Jersey: Humana Press; von Heinje, G., 1987, SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, New York: Academic Press; SEQUENCE ANALYSIS PRIMER, (Gribskov, M. and Devereux, J., eds.), 1991, New York: M. Stockton Press; and Carillo et al., 1988, *SIAM J. Applied Math.* 48:1073; and Durbin et al., 1998, BIOLOGICAL SEQUENCE ANALYSIS, Cambridge University Press.

**[0117]** Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity are described in publicly available computer programs. Preferred computer program methods to determine identity between two sequences include, but are not limited to, the GCG program package, including GAP (Devereux et al., 1984, *Nucl. Acid. Res.* 12:387; Genetics Computer Group, University of Wisconsin, Madison, Wis., BLASTP, BLASTN, and FASTA, Altschul et al., 1990, *J. Mol. Biol.* 215: 403-410). The BLASTX program is publicly available from the National Center for Biotechnology Information (NCBI) and other sources (BLAST Manual, Altschul et al. NCB/NLM/NIH Bethesda, Md.

20894; Altschul et al., 1990, supra). The well-known Smith Waterman algorithm may also be used to determine identity.

**[0118]** Certain alignment schemes for aligning two amino acid sequences may result in the matching of only a short region of the two sequences, and this small aligned region may have very high sequence identity even though there is no significant relationship between the two full-length sequences. Accordingly, in certain embodiments, the selected alignment method (GAP program) will result in an alignment that spans at least 50 contiguous amino acids of the target polypeptide.

**[0119]** For example, using the computer algorithm GAP (Genetics Computer Group, University of Wisconsin, Madison, Wis.), two polypeptides for which the percent sequence identity is to be determined are aligned for optimal matching of their respective amino acids (the “matched span”, as determined by the algorithm). In certain embodiments, a gap opening penalty (which is calculated as three times the average diagonal, wherein the “average diagonal” is the average of the diagonal of the comparison matrix being used; the “diagonal” is the score or number assigned to each perfect amino acid match by the particular comparison matrix) and a gap extension penalty (which is usually one-tenth of the gap opening penalty), as well as a comparison matrix such as PAM 250 or BLOSUM 62 are used in conjunction with the algorithm. In certain embodiments, a standard comparison matrix (see Dayhoff et al., 1978, *Atlas of Protein Sequence and Structure* 5:345-352 for the PAM 250 comparison matrix; Henikoff et al., 1992, *Proc. Natl. Acad. Sci. USA* 89: 10915-10919 for the BLOSUM 62 comparison matrix) is also used by the algorithm.

**[0120]** In certain embodiments, the parameters for a polypeptide sequence comparison include the following:

**[0121]** Algorithm: Needleman et al., 1970, *J. Mol. Biol.* 48:443-453;

**[0122]** Comparison matrix: BLOSUM 62 from Henikoff et al., 1992, supra;

**[0123]** Gap Penalty: 12

**[0124]** Gap Length Penalty: 4

**[0125]** Threshold of Similarity: 0

The GAP program may be useful with the above parameters. In certain embodiments, the aforementioned parameters are the default parameters for polypeptide comparisons (along with no penalty for end gaps) using the GAP algorithm.

**[0126]** As used herein, the twenty conventional amino acids and their abbreviations follow conventional usage. See IMMUNOLOGY—A SYNTHESIS, 2nd Edition, (E. S. Golub and D. R. Gren, Eds.), 1991, Sinauer Associates, Sunderland, Mass., which is incorporated herein by reference for any purpose. Stereoisomers (e.g., D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as  $\alpha$ -,  $\alpha$ -disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional amino acids may also be suitable components for polypeptides of the present invention. Examples of unconventional amino acids include: 4-hydroxyproline,  $\gamma$ -carboxyglutamate,  $\epsilon$ -N,N,N-trimethyllysine,  $\epsilon$ -N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine,  $\sigma$ -N-methylarginine, and other similar amino acids and imino acids (e.g., 4-hydroxyproline). In the polypeptide notation used herein, the left-hand direction is the amino terminal direction and the right-hand direction is the carboxy-terminal direction, in accordance with standard usage and convention.

**[0127]** Similarly, unless specified otherwise, the left-hand end of single-stranded polynucleotide sequences is the 5' end; the left-hand direction of double-stranded polynucleotide sequences is referred to as the 5' direction. The direction of 5' to 3' addition of nascent RNA transcripts is referred to as the transcription direction; sequence regions on the DNA strand having the same sequence as the RNA and which are 5' to the 5' end of the RNA transcript are referred to as “upstream sequences”; sequence regions on the DNA strand having the same sequence as the RNA and which are 3' to the 3' end of the RNA transcript are referred to as “downstream sequences”.

**[0128]** Naturally occurring residues may be divided into classes based on common side chain properties:

**[0129]** 1) hydrophobic: norleucine, Met, Ala, Val, Leu, Ile;

**[0130]** 2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;

**[0131]** 3) acidic: Asp, Glu;

**[0132]** 4) basic: His, Lys, Arg;

**[0133]** 5) residues that influence chain orientation: Gly, Pro; and

**[0134]** 6) aromatic: Trp, Tyr, Phe.

**[0135]** Conservative amino acid substitutions may encompass non-naturally occurring amino acid residues, which are typically incorporated by chemical peptide synthesis rather than by synthesis in biological systems. These include peptidomimetics and other reversed or inverted forms of amino acid moieties.

**[0136]** Non-conservative substitutions may involve the exchange of a member of one of these classes for a member from another class. Such substituted residues may be introduced into regions of the human antibody that are homologous with non-human antibodies, or into the non-homologous regions of the molecule.

**[0137]** In making such changes, according to certain embodiments, the hydropathic index of amino acids may be considered. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics. They are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (−0.4); threonine (−0.7); serine (−0.8); tryptophan (−0.9); tyrosine (−1.3); proline (−1.6); histidine (−3.2); glutamate (−3.5); glutamine (−3.5); aspartate (−3.5); asparagine (−3.5); lysine (−3.9); and arginine (−4.5).

**[0138]** The importance of the hydropathic amino acid index in conferring interactive biological function on a protein is understood in the art (see, for example, Kyte et al., 1982, *J. Mol. Biol.* 157:105-131). It is known that certain amino acids may be substituted for other amino acids having a similar hydropathic index or score and still retain a similar biological activity. In making changes based upon the hydropathic index, in certain embodiments, the substitution of amino acids whose hydropathic indices are within  $\pm 2$  is included. In certain embodiments, those which are within  $\pm 1$  are included, and in certain embodiments, those within  $\pm 0.5$  are included.

**[0139]** It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity, particularly where the biologically functional protein or peptide thereby created is intended for use in immunological embodiments, as in the present case. In certain embodiments, the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent

amino acids, correlates with its immunogenicity and antigen-binding or immunogenicity, i.e., with a biological property of the protein.

**[0140]** The following hydrophilicity values have been assigned to these amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0±1); glutamate (+3.0±1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5±1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5) and tryptophan (-3.4). In making changes based upon similar hydrophilicity values, in certain embodiments, the substitution of amino acids whose hydrophilicity values are within ±2 is included, in certain embodiments, those which are within ±1 are included, and in certain embodiments, those within ±0.5 are included. One may also identify epitopes from primary amino acid sequences on the basis of hydrophilicity. These regions are also referred to as “epitopic core regions.”

**[0141]** Exemplary amino acid substitutions are set forth in Table 1.

TABLE 1

Amino Acid Substitutions		
Original Residues	Exemplary Substitutions	Preferred Substitutions
Ala	Val, Leu, Ile	Val
Arg	Lys, Gln, Asn	Lys
Asn	Gln	Gln
Asp	Glu	Glu
Cys	Ser, Ala	Ser
Gln	Asn	Asn
Glu	Asp	Asp
Gly	Pro, Ala	Ala
His	Asn, Gln, Lys, Arg	Arg
Ile	Leu, Val, Met, Ala, Phe, Norleucine	Leu
Leu	Norleucine, Ile, Val, Met, Ala, Phe	Ile
Lys	Arg, Gln, Asn, 1,4 Diamine-butyric Acid	Arg
Met	Leu, Phe, Ile	Leu
Phe	Leu, Val, Ile, Ala, Tyr	Leu
Pro	Ala	Gly
Ser	Thr, Ala, Cys	Thr
Thr	Ser	Ser
Trp	Tyr, Phe	Tyr
Tyr	Trp, Phe, Thr, Ser	Phe
Val	Ile, Met, Leu, Phe, Ala, Norleucine	Leu

**[0142]** A skilled artisan will be able to determine suitable variants of the polypeptide as set forth herein using well-known techniques. In certain embodiments, one skilled in the art can identify suitable areas of the molecule that can be changed without destroying activity by targeting regions not believed to be important for activity. In other embodiments, the skilled artisan can identify residues and portions of the molecules that are conserved among similar polypeptides. In further embodiments, even areas that are important for biological activity or for structure may be subject to conservative amino acid substitutions without destroying the biological activity or without adversely affecting the polypeptide structure.

**[0143]** Additionally, one skilled in the art can review structure-function studies identifying residues in similar polypeptides that are important for activity or structure. In view of

such a comparison, the skilled artisan can predict the importance of amino acid residues in a protein that correspond to amino acid residues important for activity or structure in similar proteins. One skilled in the art may opt for chemically similar amino acid substitutions for such predicted important amino acid residues.

**[0144]** One skilled in the art can also analyze the three-dimensional structure and amino acid sequence in relation to that structure in similar polypeptides. In view of such information, one skilled in the art may predict the alignment of amino acid residues of an antibody with respect to its three dimensional structure. In certain embodiments, one skilled in the art may choose not to make radical changes to amino acid residues predicted to be on the surface of the protein, since such residues may be involved in important interactions with other molecules. Moreover, one skilled in the art may generate test variants containing a single amino acid substitution at each desired amino acid residue. The variants can then be screened using activity assays known to those skilled in the art. Such variants can be used to gather information about suitable variants. For example, if it was discovered that a change to a particular amino acid residue resulted in destroyed, undesirably reduced, or unsuitable activity, variants with such a change can be avoided. In other words, based on information gathered from such routine experiments, one skilled in the art can readily determine the amino acids where further substitutions should be avoided either alone or in combination with other mutations.

**[0145]** A number of scientific publications have been devoted to the prediction of secondary structure. See Molt, 1996, *Curr. Op. in Biotech.* 7: 422-427; Chou et al., 1974, *Biochemistry* 13: 222-245; Chou et al., 1974, *Biochemistry* 113: 211-222; Chou et al., 1978, *Adv. Enzymol. Relat. Areas Mol. Biol.* 47: 45-148; Chou et al., 1978, *Ann. Rev. Biochem.* 47: 251-276 and Chou et al., 1979, *Biophys. J.* 26: 367-384. Moreover, computer programs are currently available to assist with predicting secondary structure. One method of predicting secondary structure is based upon homology modeling. For example, two polypeptides or proteins which have a sequence identity of greater than 30%, or similarity greater than 40% often have similar structural topologies. The recent growth of the protein structural database (PDB) has provided enhanced predictability of secondary structure, including the potential number of folds within a polypeptide's or protein's structure. See Holm et al., 1999, *Nucl. Acid. Res.* 27: 244-247. It has been suggested (Brenner et al., 1997, *Curr. Op. Struct. Biol.* 7: 369-376) that there are a limited number of folds in a given polypeptide or protein and that once a critical number of structures have been resolved, structural prediction will become dramatically more accurate.

**[0146]** Additional methods of predicting secondary structure include “threading” (Jones, 1997, *Curr. Opin. Struct. Biol.* 7: 377-87; Sippl et al., 1996, *Structure* 4: 15-19), “profile analysis” (Bowie et al., 1991, *Science* 253: 164-170; Gribskov et al., 1990, *Meth. Enzym.* 183: 146-159; Gribskov et al., 1987, *Proc. Nat. Acad. Sci. USA* 84: 4355-4358), and “evolutionary linkage” (See Holm, 1999, *supra*, and Brenner, 1997, *supra*).

**[0147]** In certain embodiments, antibody variants include glycosylation variants wherein the number and/or type of glycosylation site has been altered compared to the amino acid sequences of the parent polypeptide. In certain embodiments, protein variants comprise a greater or a lesser number of N-linked glycosylation sites than the native protein. An

N-linked glycosylation site is characterized by the sequence: Asn-X-Ser or Asn-X-Thr, wherein the amino acid residue designated as X may be any amino acid residue except proline. The substitution of amino acid residues to create this sequence provides a potential new site for the addition of an N-linked carbohydrate chain. Alternatively, substitutions that eliminate or alter this sequence will prevent addition of an N-linked carbohydrate chain present in the native polypeptide. Also provided are rearrangements of N-linked carbohydrate chains wherein one or more N-linked glycosylation sites (typically those that are naturally occurring) are eliminated and one or more new N-linked sites are created. Additional preferred antibody variants include cysteine variants wherein one or more cysteine residues in the parent or native amino acid sequence are deleted from or substituted for another amino acid (e.g., serine). Cysteine variants are useful, inter alia when antibodies must be refolded into a biologically active conformation, for example, after the isolation of insoluble inclusion bodies. Cysteine variants generally have fewer cysteine residues than the native protein, and typically have an even number to minimize interactions resulting from unpaired cysteines.

**[0148]** In additional embodiments, antibody variants can include antibodies comprising a modified Fc fragment or a modified heavy chain constant region. An Fc fragment, which stands for "fragment that crystallizes," or a heavy chain constant region can be modified by mutation to confer on an antibody altered binding characteristics. See, for example, Burton and Woof, 1992, *Advances in Immunology* 51: 1-84; Ravetch and Bolland, 2001, *Annu. Rev. Immunol.* 19: 275-90; Shields et al., 2001, *Journal of Biol. Chem.* 276: 6591-6604; Telleman and Junghans, 2000, *Immunology* 100: 245-251; Medesan et al., 1998, *Eur. J. Immunol.* 28: 2092-2100; all of which are incorporated herein by reference). Such mutations can include substitutions, additions, deletions, or any combination thereof, and are typically produced by site-directed mutagenesis using one or more mutagenic oligonucleotide(s) according to methods described herein, as well as according to methods known in the art (see, for example, Maniatis et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 3rd Ed., 2001, Cold Spring Harbor, N.Y. and Berger and Kimmel, *METHODS IN ENZYMOLOGY*, Volume 152, Guide to Molecular Cloning Techniques, 1987, Academic Press, Inc., San Diego, Calif., which are incorporated herein by reference).

**[0149]** According to certain embodiments, amino acid substitutions are those that: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter ligand or antigen binding affinities, and/or (5) confer or modify other physicochemical or functional properties on such polypeptides. According to certain embodiments, single or multiple amino acid substitutions (in certain embodiments, conservative amino acid substitutions) may be made in the naturally-occurring sequence (in certain embodiments, in the portion of the polypeptide outside the domain(s) forming intermolecular contacts). In certain embodiments, a conservative amino acid substitution typically does not substantially change the structural characteristics of the parent sequence (e.g., a replacement amino acid should not tend to break a helix that occurs in the parent sequence, or disrupt other types of secondary structure that characterizes the parent sequence). Examples of art-recognized polypeptide secondary and tertiary structures are described in *PROTEINS, STRUCTURES*

AND MOLECULAR PRINCIPLES (Creighton, Ed.), 1984, W. H. New York: Freeman and Company; *INTRODUCTION TO PROTEIN STRUCTURE* (Branden and Tooze, eds.), 1991, New York: Garland Publishing; and Thornton et al., 1991, *Nature* 354: 105, each of which are incorporated herein by reference.

**[0150]** Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed "peptide mimetics" or "peptidomimetics". Fauchere, 1986, *Adv. Drug Res.* 15: 29; Veber and Freidinger, 1985, *TINS* p. 392; and Evans et al., 1987, *J. Med. Chem.* 30: 1229, which are incorporated herein by reference for any purpose. Such compounds are often developed with the aid of computerized molecular modeling. Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce a similar therapeutic or prophylactic effect. Generally, peptidomimetics are structurally similar to a paradigm polypeptide (i.e., a polypeptide that has a biochemical property or pharmacological activity), such as human antibody, but have one or more peptide linkages optionally replaced by a linkage selected from:  $-\text{CH}_2\text{NH}-$ ,  $-\text{CH}_2\text{S}-$ ,  $-\text{CH}_2-\text{CH}_2-$ ,  $-\text{CH}=\text{CH}-$  (cis and trans),  $-\text{COCH}_2-$ ,  $-\text{CH}(\text{OH})\text{CH}_2-$ , and  $-\text{CH}_2\text{SO}-$ , by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) may be used in certain embodiments to generate more stable peptides. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus sequence variation may be generated by methods known in the art (Rizo and Gierasch, 1992, *Ann. Rev. Biochem.* 61: 387), incorporated herein by reference for any purpose); for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

**[0151]** "Antibody" or "antibody peptide(s)" refer to an intact antibody, or a binding fragment thereof that competes with the intact antibody for specific binding. In certain embodiments, binding fragments are produced by recombinant DNA techniques. In certain embodiments, binding fragments are produced by enzymatic or chemical cleavage of intact antibodies. Binding fragments include, but are not limited to, Fab, Fab', F(ab')<sub>2</sub>, Fv, and single-chain antibodies.

**[0152]** The term "heavy chain" includes any immunoglobulin polypeptide having a heavy chain constant region and sufficient variable region sequence to confer specificity for an OPGL. The term "light chain" includes any immunoglobulin polypeptide having a light chain constant region and sufficient variable region sequence to confer specificity for an OPGL. A full-length heavy chain includes a variable region domain, V<sub>H</sub>, and three constant region domains, C<sub>H</sub>1, C<sub>H</sub>2, and C<sub>H</sub>3. The V<sub>H</sub> domain is at the amino-terminus of the polypeptide, and the C<sub>H</sub>3 domain is at the carboxyl-terminus. The term "heavy chain", as used herein, encompasses a full-length heavy chain and fragments thereof. A full-length light chain includes a variable region domain, V<sub>L</sub>, and a constant region domain, C<sub>L</sub>. Like the heavy chain, the variable region domain of the light chain is at the amino-terminus of the polypeptide. The term "light chain", as used herein, encompasses a full-length light chain and fragments thereof. A F(ab) fragment is comprised of one light chain and the C<sub>H</sub>1 and variable regions of one heavy chain. The heavy chain of a F(ab) molecule cannot form a disulfide bond with another

heavy chain molecule. A F(ab') fragment contains one light chain and one heavy chain that contains more of the constant region, between the C<sub>H</sub>1 and C<sub>H</sub>2 domains, such that an interchain disulfide bond can be formed between two heavy chains to form a F(ab')<sub>2</sub> molecule. The Fv region comprises the variable regions from both the heavy and light chains, but lacks the constant regions. Single-chain antibodies are Fv molecules in which the heavy and light chain variable regions have been connected by a flexible linker to form a single polypeptide chain that forms an antigen-binding region. Single chain antibodies are discussed in detail in WO 88/01649 and U.S. Pat. Nos. 4,946,778 and 5,260,203 incorporate by reference.

**[0153]** A bivalent antibody other than a “multispecific” or “multifunctional” antibody, in certain embodiments, is understood to comprise binding sites having identical antigenic specificity.

**[0154]** In assessing antibody binding and specificity according to the invention, an antibody substantially inhibits adhesion of a ligand to a receptor when an excess of antibody reduces the quantity of ligand bound to receptor by at least about 20%, 40%, 60%, 80%, 85%, or more (as measured in an in vitro competitive binding assay).

**[0155]** The term “epitope” includes any polypeptide determinant, preferably a polypeptide determinant, capable of specific binding to an immunoglobulin or T-cell receptor. In certain embodiments, epitope determinants include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl, or sulfonyl, and, in certain embodiments, may have specific three dimensional structural characteristics, and/or specific charge characteristics. An epitope is a region of an antigen that is bound by an antibody. In certain embodiments, an antibody is said to specifically bind an antigen when it preferentially recognizes its target antigen in a complex mixture of proteins and/or macromolecules. In certain embodiments, an antibody is said to specifically bind an antigen when the dissociation constant is  $\leq 10^{-8}$  M, in certain embodiments, when the dissociation constant is  $\leq 10^{-9}$  M, and in certain embodiments, when the dissociation constant is  $\leq 10^{-10}$  M.

**[0156]** The term “agent” is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials.

**[0157]** As used herein, the terms “label” or “labeled” refers to incorporation of a detectable marker, e.g., by incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotin moieties that can be detected by labeled avidin (e.g., streptavidin preferably comprising a detectable marker such as a fluorescent marker, a chemiluminescent marker or an enzymatic activity that can be detected by optical or colorimetric methods). In certain embodiments, the label can also be therapeutic. Various methods of labeling polypeptides and glycoproteins are known in the art and may be used advantageously in the methods disclosed herein. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (e.g., <sup>3</sup>H, <sup>14</sup>C, <sup>15</sup>N, <sup>35</sup>S, <sup>90</sup>Y, <sup>99m</sup>Tc, <sup>111</sup>In, <sup>125</sup>I, <sup>131</sup>I) fluorescent labels (e.g., fluorescein isothiocyanate or FITC, rhodamine, or lanthanide phosphors), enzymatic labels (e.g., horseradish peroxidase,  $\beta$ -galactosidase, luciferase, alkaline phosphatase), chemiluminescent labels, hapten labels such as biotinyl groups, and predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, or epitope

tags). In certain embodiments, labels are attached by spacer arms (such as (CH<sub>2</sub>)<sub>n</sub>, where n < about 20) of various lengths to reduce potential steric hindrance.

**[0158]** The term “biological sample”, as used herein, includes, but is not limited to, any quantity of a substance from a living thing or formerly living thing. Such living things include, but are not limited to, humans, mice, monkeys, rats, rabbits, and other animals. Such substances include, but are not limited to, blood, serum, urine, cells, organs, tissues, bone, bone marrow, lymph nodes, and skin.

**[0159]** The term “osteopenic disorder” includes, but is not limited to, osteoporosis, osteopenia, Paget’s disease, lytic bone metastases, periodontitis, rheumatoid arthritis, and bone loss due to immobilization. In addition to these bone disorders, certain cancers are known to increase osteoclast activity and induce bone resorption, such as breast and prostate cancer and multiple myeloma. These cancers are now known to produce factors that result in the over-expression of OPGL in the bone, and lead to increased osteoclast numbers and activity.

**[0160]** The term “pharmaceutical agent or drug” as used herein refers to a chemical compound or composition capable of inducing a desired therapeutic effect when properly administered to a patient.

**[0161]** As used herein, “substantially pure” or “substantially purified” means a compound or species that is the predominant species present (i.e., on a molar basis it is more abundant than any other individual species in the composition). In certain embodiments, a substantially purified fraction is a composition wherein the object species comprises at least about 50 percent (on a molar basis) of all macromolecular species present. In certain embodiments, a substantially pure composition will comprise more than about 80%, 85%, 90%, 95%, or 99% of all macromolecular species present in the composition. In certain embodiments, the species is purified to essential homogeneity (contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species.

**[0162]** The term “patient” includes human and animal subjects.

**[0163]** Unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular.

**[0164]** The invention provides antibodies, preferably monoclonal antibodies and most preferably human antibodies, that are immunologically specific for osteoprotegerin ligand (OPGL), a member of the tumor necrosis factor (TNF) family of cytokines that is involved in the formation of osteoclasts. Increased osteoclast activity correlates with a number of osteopenic disorders, including post-menopausal osteoporosis, Paget’s disease, lytic bone metastases, and rheumatoid arthritis. Thus, a reduction in OPGL activity may result in a decrease in osteoclast activity and may reduce the severity of osteopenic disorders. According to certain embodiments of the invention, antibodies directed to OPGL may be used to detect, diagnose, prevent and treat osteopenic disorders, including but not limited to, those mentioned above.

**[0165]** In certain embodiments of the present invention, there is provided a fully human monoclonal antibody against human OPGL. In certain embodiments, nucleotide sequences encoding, and amino acid sequences comprising, heavy and light chain immunoglobulin molecules, particularly sequences corresponding to the variable regions, are pro-

vided. In certain embodiments, sequences corresponding to complementarity determining regions (CDR's), specifically from CDR1 through CDR3, are provided. According to certain embodiments, a hybridoma cell line expressing such an immunoglobulin molecule and monoclonal antibody is also provided. In certain embodiments, the invention provides purified human monoclonal antibody against human OPGL.

**[0166]** The ability to clone and reconstruct megabase-sized human loci in yeast artificial chromosomes (YACs) and to introduce them into the mouse germline provides an approach to elucidating the functional components of very large or crudely mapped loci as well as generating useful models of human disease. Furthermore, the utilization of such technology for substitution of mouse loci with their human equivalents provides unique insights into the expression and regulation of human gene products during development, their communication with other systems, and their involvement in disease induction and progression.

**[0167]** An important practical application of such a strategy is the "humanization" of the mouse humoral immune system. Introduction of human immunoglobulin (Ig) loci into mice in which the endogenous Ig genes have been inactivated offers the opportunity to study mechanisms underlying programmed expression and assembly of antibodies as well as their role in B-cell development. Furthermore, such a strategy provides a source for production of fully human monoclonal antibodies (MAbs). Fully human antibodies are expected to minimize the immunogenic and allergic responses intrinsic to mouse or mouse-derivatized MAbs, and to thereby increase the efficacy and safety of the administered antibodies. Fully human antibodies can be used in the treatment of chronic and recurring human diseases, such as osteoporosis, inflammation, autoimmunity, and cancer, the treatment thereof requiring repeated antibody administration. Thus, one particular advantage of the anti-OPGL antibodies of the invention is that the antibodies are fully human and can be administered to patients in a non-acute manner while minimizing adverse reactions commonly associated with human anti-mouse antibodies or other previously described non-fully human antibodies from non-human species.

**[0168]** One skilled in the art can engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig loci so that the mice produce human but not mouse antibodies. Large human Ig fragments in mouse germline preserve variable gene diversity as well as the proper regulation of antibody production and expression. By exploiting the mouse machinery for antibody diversification and selection and the lack of immunological tolerance to human proteins, the reproduced human antibody repertoire in these mouse strains yield high affinity antibodies against any antigen of interest, including human antigens. Using hybridoma technology, antigen-specific human mAbs with the desired specificity can be produced and selected.

**[0169]** In certain embodiments, the skilled artisan can use constant regions from species other than human along with the human variable region(s) to produce chimeric antibodies.

#### Naturally Occurring Antibody Structure

**[0170]** Naturally occurring antibody structural units typically comprise a tetramer. Each such tetramer typically is composed of two identical pairs of polypeptide chains, each pair having one full-length "light" chain (in certain embodiments, about 25 kDa) and one full-length "heavy" chain (in certain embodiments, about 50-70 kDa). The amino-terminal

portion of each chain typically includes a variable region of about 100 to 110 or more amino acids that typically is responsible for antigen recognition. The carboxy-terminal portion of each chain typically defines a constant region that may be responsible for effector function. Human light chains are typically classified as kappa and lambda light chains. Heavy chains are typically classified as mu, delta, gamma, alpha, or epsilon, and define the antibody's isotype as IgM, IgD, IgG, IgA, and IgE, respectively. IgG has several subclasses, including, but not limited to, IgG<sub>1</sub>, IgG<sub>2</sub>, IgG<sub>3</sub>, and IgG<sub>4</sub>. IgM has subclasses including, but not limited to, IgM<sub>1</sub> and IgM<sub>2</sub>. IgA is similarly subdivided into subclasses including, but not limited to, IgA<sub>1</sub> and IgA<sub>2</sub>. Within full-length light and heavy chains, typically, the variable and constant regions are joined by a "J" region of about 12 or more amino acids, with the heavy chain also including a "D" region of about 10 more amino acids. See, e.g., FUNDAMENTAL IMMUNOLOGY, 2nd ed., Ch. 7 (Paul, W., ed.) 1989, New York: Raven Press (incorporated by reference in its entirety for all purposes). The variable regions of each light/heavy chain pair typically form the antigen-binding site.

**[0171]** The variable regions typically exhibit the same general structure of relatively conserved framework regions (FR) joined by three hypervariable regions, also called complementarity determining regions or CDRs. The CDRs from the two chains of each pair typically are aligned by the framework regions, which may enable binding to a specific epitope. From N-terminal to C-terminal, both light and heavy chain variable regions typically comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is typically in accordance with the definitions of Kabat Sequences of Proteins of Immunological Interest (1987 and 1991, National Institutes of Health, Bethesda, Md.), or Chothia & Lesk, 1987, *J. Mol. Biol.* 196: 901-917; Chothia et al., 1989, *Nature* 342: 878-883.

#### Bispecific or Bifunctional Antibodies

**[0172]** A bispecific or bifunctional antibody typically is an artificial hybrid antibody having two different heavy/light chain pairs and two different binding sites. Bispecific antibodies may be produced by a variety of methods including, but not limited to, fusion of hybridomas or linking of Fab' fragments. See, e.g., Songsivilai & Lachmann, 1990, *Clin. Exp. Immunol.* 79: 315-321; Kostelny et al., 1992, *J. Immunol.* 148: 1547-1553.

#### Preparation of Antibodies

**[0173]** The invention provides antibodies that specifically bind to human OPGL. In certain embodiments, the antibodies can be produced by immunization with full-length OPGL or fragments thereof. The antibodies of the invention can be polyclonal or monoclonal, and/or may be recombinant antibodies. In preferred embodiments, antibodies of the invention are human antibodies prepared, for example, by immunization of transgenic animals capable of producing human antibodies (see, for example, PCT Published Application No. WO 93/12227).

**[0174]** The complementarity determining regions (CDRs) of the light and heavy chain variable regions of anti-OPGL antibody may be grafted to framework regions (FRs) from antibodies from the same, or another, species. In certain embodiments, the CDRs of the light and heavy chain variable regions of anti-OPGL antibody may be grafted to consensus

human FRs. To create consensus human FRs, in certain embodiments, FRs from several human heavy chain or light chain amino acid sequences are aligned to identify a consensus amino acid sequence. In certain embodiments, the FRs of the anti-OPGL antibody heavy chain or light chain are replaced with the FRs from a different heavy chain or light chain. In certain embodiments, rare amino acids in the FRs of the heavy and light chains of anti-OPGL antibody are not replaced, while the rest of the FR amino acids are replaced. Rare amino acids are specific amino acids that are in positions in which they are not usually found in FRs. In certain embodiments, the grafted variable regions from anti-OPGL antibody may be used with a constant region that is different from the constant region of anti-OPGL antibody. In certain embodiments, the grafted variable regions are part of a single chain Fv antibody. CDR grafting is described, e.g., in U.S. Pat. Nos. 6,180,370, 5,693,762, 5,693,761, 5,585,089, and 5,530,101, which are hereby incorporated by reference for any purpose.

**[0175]** Antibodies of the invention are prepared using transgenic mice that have a substantial portion of the human antibody producing locus inserted in antibody-producing cells of the mice, and that are further engineered to be deficient in producing endogenous, murine, antibodies. Such mice are capable of producing human immunoglobulin molecules and antibodies and do not produce or produce substantially reduced amounts of murine immunoglobulin molecules and antibodies. Technologies utilized for achieving this result are disclosed in the patents, applications, and references disclosed in the specification herein. In certain embodiments, the skilled worker may employ methods as disclosed in International Patent Application Publication No. WO 98/24893, which is hereby incorporated by reference for any purpose. See also Mendez et al., 1997, *Nature Genetics* 15: 146-156, which is hereby incorporated by reference for any purpose.

**[0176]** The monoclonal antibodies (mAbs) of the invention can be produced by a variety of techniques, including conventional monoclonal antibody methodology, e.g., the standard somatic cell hybridization technique of Kohler and Milstein, 1975, *Nature* 256: 495. Although somatic cell hybridization procedures are preferred, in principle, other techniques for producing monoclonal antibodies can be employed, e.g., viral or oncogenic transformation of B-lymphocytes.

**[0177]** The preferred animal system for preparing hybridomas is the mouse. Hybridoma production in the mouse is very well established, and immunization protocols and techniques for isolation of immunized splenocytes for fusion are well known in the art. Fusion partners (e.g., murine myeloma cells) and fusion procedures are also known.

**[0178]** In a preferred embodiment, human monoclonal antibodies directed against OPGL can be generated using transgenic mice carrying parts of the human immune system rather than the mouse system. These transgenic mice, referred to herein as "HuMab" mice, contain a human immunoglobulin gene minilocus that encodes unrearranged human heavy ( $\mu$  and  $\gamma$ ) and  $\kappa$  light chain immunoglobulin sequences, together with targeted mutations that inactivate the endogenous  $\mu$  and  $\kappa$  chain loci (Lonberg et al., 1994, *Nature* 368: 856-859). Accordingly, the mice exhibit reduced expression of mouse IgM or  $\kappa$  and in response to immunization, the introduced human heavy and light chain transgenes, undergo class switching and somatic mutation to generate high affinity human IgG  $\kappa$  monoclonal antibodies (Lonberg et al., supra;

Lonberg and Huszar, 1995, *Intern. Rev. Immunol.*, 13: 65-93; Harding and Lonberg, 1995, *Ann. N.Y. Acad. Sci.* 764: 536-546). The preparation of HuMab mice is described in detail in Taylor et al., 1992, *Nucleic Acids Research*, 20: 6287-6295; Chen et al., 1993, *International Immunology* 5: 647-656; Tuailon et al., 1994, *J. Immunol.* 152: 2912-2920; Lonberg et al., 1994, *Nature* 368: 856-859; Lonberg, 1994, *Handbook of Exp. Pharmacology* 113: 49-101; Taylor et al., 1994, *International Immunology* 6: 579-591; Lonberg and Huszar, 1995, *Intern. Rev. Immunol.* 13: 65-93; Harding and Lonberg, 1995, *Ann. N.Y. Acad. Sci.* 764: 536-546; Fishwild et al., 1996, *Nature Biotechnology* 14: 845-851, the contents of all of which are hereby incorporated by reference in their entirety. See further U.S. Pat. Nos. 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,789,650; 5,877,397; 5,661,016; 5,814,318; 5,874,299; and 5,770,429; all to Lonberg and Kay, as well as U.S. Pat. No. 5,545,807 to Surani et al.; International Publication Nos. WO 93/1227, published Jun. 24, 1993; WO 92/22646, published Dec. 23, 1992; WO 92/03918, published Mar. 19, 1992, the disclosures of all of which are hereby incorporated by reference in their entirety. Alternatively, the HCo7 and HCo12 transgenic mice strains described in the Examples below can be used to generate human anti-OPGL antibodies.

**[0179]** According to certain embodiments, fully human monoclonal antibodies specific for OPGL are produced as follows. Transgenic mice containing human immunoglobulin genes are immunized with the antigen of interest. Lymphatic cells (such as B-cells) from the mice that express antibodies are obtained. Such recovered cells are fused with a myeloid-type cell line to prepare immortal hybridoma cell lines, and such hybridoma cell lines are screened and selected to identify hybridoma cell lines that produce antibodies specific to the antigen of interest. In certain embodiments, the production of a hybridoma cell line that produces antibodies specific to OPGL is provided.

**[0180]** In certain embodiments of the invention, the antibodies bind to OPGL with a dissociation constant ( $K_d$ ) of less than  $10^{-8}$  M. In certain embodiments, the antibodies of the invention bind to OPGL with a  $K_d$  of between approximately  $10^{-8}$  M and  $10^{-10}$  M.

**[0181]** In certain embodiments, the antibodies of the invention are of the IgG<sub>1</sub> isotype. In certain embodiments of the invention, the antibodies comprise a human kappa light chain and a human IgG<sub>1</sub> heavy chain. In certain embodiments, nucleic acid encoding the heavy and light chains comprising the antibodies of the invention were cloned for expression in mammalian cells. In certain embodiments, the variable regions of the antibodies are ligated to a constant region other than the constant region for the IgG<sub>1</sub> isotype.

**[0182]** In certain embodiments, conservative modifications to the heavy and light chains of anti-OPGL antibody (and corresponding modifications to the encoding nucleic acids) will produce anti-OPGL antibodies having functional and biochemical characteristics similar to those of anti-OPGL antibody. In contrast, substantial modifications in the functional and/or biochemical characteristics of anti-OPGL antibody may be achieved by creating substitutions in the amino acid sequence of the heavy and light chains that differ significantly in their effect on maintaining (a) the structure of the molecular backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulkiness of the side chain.



**[0183]** For example, a “conservative amino acid substitution” may involve a substitution of a native amino acid residue with a normative residue having little or no effect on the polarity or charge of the amino acid residue at that position. Furthermore, any native residue in the polypeptide may also be substituted with alanine, as has been previously described for “alanine scanning mutagenesis.”

**[0184]** Desired amino acid substitutions (whether conservative or non-conservative) can be determined by those skilled in the art at the time such substitutions are desired. In certain embodiments, amino acid substitutions can be used to identify important residues of anti-OPGL antibody, or to increase or decrease the affinity of the anti-OPGL antibodies described herein.

**[0185]** In alternative embodiments, antibodies of the present invention can be expressed in cell lines other than hybridoma cell lines. In these embodiments, sequences encoding particular antibodies can be used for transformation of a suitable mammalian host cell. According to these embodiments, transformation can be by any known method for introducing polynucleotides into a host cell, including, for example packaging the polynucleotide in a virus (or into a viral vector) and transducing a host cell with the virus (or vector) or by transfection procedures known in the art, as exemplified by U.S. Pat. Nos. 4,399,216, 4,912,040, 4,740,461, and 4,959,455 (which patents are hereby incorporated herein by reference for any purpose). Generally, the transformation procedure used may depend upon the host to be transformed. Methods for introduction of heterologous polynucleotides into mammalian cells are well known in the art and include, but are not limited to, dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, mixing nucleic acid with positively-charged lipids, and direct microinjection of the DNA into nuclei.

**[0186]** A nucleic acid molecule encoding the amino acid sequence of a heavy chain constant region, a heavy chain variable region, a light chain constant region, or a light chain variable region of an OPGL antibody of the invention is inserted into an appropriate expression vector using standard ligation techniques. In a preferred embodiment, the anti-OPGL antibody heavy chain or light chain constant region is appended to the C-terminus of the appropriate variable region and is ligated into an expression vector. The vector is typically selected to be functional in the particular host cell employed (i.e., the vector is compatible with the host cell machinery such that amplification of the gene and/or expression of the gene can occur). For a review of expression vectors, see METH. ENZ. 185 (Goeddel, ed.), 1990, Academic Press. Typically, expression vectors used in any of the host cells contain sequences for plasmid maintenance and for cloning and expression of exogenous nucleotide sequences. Such sequences, collectively referred to as “flanking sequences” in certain embodiments will typically include one or more of the following nucleotide sequences: a promoter, one or more enhancer sequences, an origin of replication, a transcriptional termination sequence, a complete intron sequence containing a donor and acceptor splice site, a sequence encoding a leader sequence for polypeptide secretion, a ribosome binding site, a polyadenylation sequence, a polylinker region for inserting the nucleic acid encoding the polypeptide to be expressed, and a selectable marker element. Each of these sequences is discussed below.

**[0187]** Optionally, the vector may contain a “tag”-encoding sequence, i.e., an oligonucleotide molecule located at the 5' or 3' end of the OPGL polypeptide coding sequence; the oligonucleotide sequence encodes polyHis (such as hexaHis), or another “tag” such as FLAG, HA (hemagglutinin influenza virus), or myc, for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as a means for affinity purification or detection of the OPGL antibody from the host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an affinity matrix. Optionally, the tag can subsequently be removed from the purified OPGL polypeptide by various means such as using certain peptidases for cleavage.

**[0188]** Flanking sequences may be homologous (i.e., from the same species and/or strain as the host cell), heterologous (i.e., from a species other than the host cell species or strain), hybrid (i.e., a combination of flanking sequences from more than one source), synthetic or native. As such, the source of a flanking sequence may be any prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequence is functional in, and can be activated by, the host cell machinery.

**[0189]** Flanking sequences useful in the vectors of this invention may be obtained by any of several methods well known in the art. Typically, flanking sequences useful herein will have been previously identified by mapping and/or by restriction endonuclease digestion and can thus be isolated from the proper tissue source using the appropriate restriction endonucleases. In some cases, the full nucleotide sequence of a flanking sequence may be known. Here, the flanking sequence may be synthesized using the methods described herein for nucleic acid synthesis or cloning.

**[0190]** Whether all or only a portion of the flanking sequence is known, it may be obtained using polymerase chain reaction (PCR) and/or by screening a genomic library with a suitable probe such as an oligonucleotide and/or flanking sequence fragment from the same or another species. Where the flanking sequence is not known, a fragment of DNA containing a flanking sequence may be isolated from a larger piece of DNA that may contain, for example, a coding sequence or even another gene or genes. Isolation may be accomplished by restriction endonuclease digestion to produce the proper DNA fragment followed by isolation using agarose gel purification, Qiagen® column chromatography (Chatsworth, Calif.), or other methods known to the skilled artisan. The selection of suitable enzymes to accomplish this purpose will be readily apparent to one of ordinary skill in the art.

**[0191]** An origin of replication is typically a part of prokaryotic expression vectors, particularly those purchased commercially, and the origin aids in the amplification of the vector in a host cell. If the vector of choice does not contain an origin of replication site, one may be chemically synthesized based on a known sequence, and ligated into the vector. For example, the origin of replication from the plasmid pBR322 (New England Biolabs, Beverly, Mass.) is suitable for most gram-negative bacteria and various origins (e.g., SV40, polyoma, adenovirus, vesicular stomatitis virus (VSV), or papillomaviruses such as HPV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression

vectors (for example, the SV40 origin is often used only because it contains the early promoter).

**[0192]** A transcription termination sequence is typically located 3' of the end of a polypeptide-coding region and serves to terminate transcription. Usually, a transcription termination sequence in prokaryotic cells is a G-C rich fragment followed by a poly-T sequence. While the sequence is easily cloned from a library or even purchased commercially as part of a vector, it can also be readily synthesized using methods for nucleic acid synthesis such as those described herein.

**[0193]** A selectable marker gene element encodes a protein necessary for the survival and growth of a host cell grown in a selective culture medium. Typical selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, tetracycline, or kanamycin for prokaryotic host cells; (b) complement auxotrophic deficiencies of the cell; or (c) supply critical nutrients not available from complex media. Preferred selectable markers are the kanamycin resistance gene, the ampicillin resistance gene, and the tetracycline resistance gene. A bacterial neomycin resistance gene can also be used for selection in both prokaryotic and eukaryotic host cells.

**[0194]** Other selection genes can be used to amplify the gene that will be expressed.

**[0195]** Amplification is a process whereby genes that cannot in single copy be expressed at high enough levels to permit survival and growth of cells under certain selection conditions are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Examples of suitable amplifiable selectable markers for mammalian cells include dihydrofolate reductase (DHFR) and promoterless thymidine kinase. In the use of these markers mammalian cell transformants are placed under selection pressure wherein only the transformants are uniquely adapted to survive by virtue of the selection gene present in the vector. Selection pressure is imposed by culturing the transformed cells under conditions in which the concentration of selection agent in the medium is successively increased, thereby leading to the amplification of both the selectable gene and the DNA that encodes another gene, such as an antibody that binds to OPGL polypeptide. As a result, increased quantities of a polypeptide such as an anti-OPGL antibody are synthesized from the amplified DNA. A ribosome-binding site is usually necessary for translation initiation of mRNA and is characterized by a Shine-Dalgarno sequence (prokaryotes) or a Kozak sequence (eukaryotes). The element is typically located 3' to the promoter and 5' to the coding sequence of the polypeptide to be expressed.

**[0196]** In some cases, for example where glycosylation is desired in a eukaryotic host cell expression system, various presequences can be manipulated to improve glycosylation or yield. For example, the peptidase cleavage site of a particular signal peptide can be altered, or pro-sequences added, which also may affect glycosylation. The final protein product may have, in the -1 position (relative to the first amino acid of the mature protein) one or more additional amino acids incident to expression, which may not have been totally removed. For example, the final protein product may have one or two amino acid residues found in the peptidase cleavage site, attached to the amino-terminus. Alternatively, use of some enzyme cleavage sites may result in a slightly truncated yet active form of the desired polypeptide, if the enzyme cuts at such area within the mature polypeptide.

**[0197]** The expression and cloning vectors of the present invention will typically contain a promoter that is recognized by the host organism and operably linked to nucleic acid encoding the anti-OPGL antibody. Promoters are untranscribed sequences located upstream (i.e., 5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control transcription of the structural gene. Promoters are conventionally grouped into one of two classes: inducible promoters and constitutive promoters. Inducible promoters initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, such as the presence or absence of a nutrient or a change in temperature. Constitutive promoters, on the other hand, initiate continual gene product production; that is, there is little or no experimental control over gene expression. A large number of promoters, recognized by a variety of potential host cells, are well known. A suitable promoter is operably linked to the DNA encoding anti-OPGL antibody by removing the promoter from the source DNA by restriction enzyme digestion or amplifying the promoter by polymerase chain reaction and inserting the desired promoter sequence into the vector.

**[0198]** Suitable promoters for use with yeast hosts are also well known in the art. Yeast enhancers are advantageously used with yeast promoters. Suitable promoters for use with mammalian host cells are well known and include, but are not limited to, those obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, retroviruses, hepatitis-B virus and most preferably Simian Virus 40 (SV40). Other suitable mammalian promoters include heterologous mammalian promoters, for example, heat-shock promoters and the actin promoter.

**[0199]** Additional promoters that may be of interest include, but are not limited to: the SV40 early promoter region (Bernoist and Chambon, 1981, *Nature* 290: 304-10); the CMV promoter; the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, et al., 1980, *Cell* 22: 787-97); the herpes thymidine kinase promoter (Wagner et al., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78: 1444-45); the regulatory sequences of the metallothionein gene (Brinster et al., 1982, *Nature* 296: 39-42); prokaryotic expression vectors such as the beta-lactamase promoter (Villa-Kamaroff et al., 1978, *Proc. Natl. Acad. Sci. U.S.A.*, 75: 3727-31); or the tac promoter (DeBoer et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80: 21-25). Also of interest are the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: the elastase I gene control region that is active in pancreatic acinar cells (Swift et al., 1984, *Cell* 38: 639-46; Ornitz et al., 1986, *Cold Spring Harbor Symp. Quant. Biol.* 50: 399-409; MacDonald, 1987, *Hepatology* 7: 425-515); the insulin gene control region that is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315: 115-22); the mouse mammary tumor virus control region that is active in testicular, breast, lymphoid and mast cells (Leder et al., 1986, *Cell* 45: 485-95); the albumin gene control region that is active in liver (Pinkert et al., 1987, *Genes and Devel.* 1: 268-76); the alpha-feto-protein gene control region that is active in liver (Krumlauf et al., 1985, *Mol. Cell. Biol.* 5: 1639-48; Hammer et al., 1987, *Science* 235: 53-58); the alpha 1-antitrypsin gene control region that is active in the liver (Kelsey et al., 1987, *Genes and Devel.* 1: 161-71); the beta-globin gene control region that is active in myeloid cells (Mogam et al., 1985, *Nature* 315: 338-40);

Kollias et al., 1986, *Cell* 46: 89-94); the myelin basic protein gene control region that is active in oligodendrocyte cells in the brain (Readhead et al., 1987, *Cell* 48: 703-12); the myosin light chain-2 gene control region that is active in skeletal muscle (Sani, 1985, *Nature* 314: 283-86); the gonadotropic releasing hormone gene control region that is active in the hypothalamus (Mason et al., 1986, *Science* 234: 1372-78); and most particularly the immunoglobulin gene control region that is active in lymphoid cells (Grosschedl et al., 1984, *Cell* 38: 647-58; Adames et al., 1985, *Nature* 318: 533-38; Alexander et al., 1987, *Mol. Cell. Biol.* 7: 1436-44).

**[0200]** An enhancer sequence may be inserted into the vector to increase the transcription of a nucleic acid encoding an anti-OPGL antibody of the present invention by higher eukaryotes. Enhancers are cis-acting elements of DNA, usually about 10-300 by in length, that act on promoters to increase transcription. Enhancers are relatively orientation and position independent. They have been found 5' and 3' to the transcription unit. Several enhancer sequences available from mammalian genes are known (e.g., globin, elastase, albumin, alpha-feto-protein and insulin). Typically, however, an enhancer from a virus will be used. The SV40 enhancer, the cytomegalovirus early promoter enhancer, the polyoma enhancer, and adenovirus enhancers are exemplary enhancing elements for the activation of eukaryotic promoters. While an enhancer may be spliced into the vector at a position 5' or 3' to a nucleic acid molecule, it is typically located at a site 5' from the promoter.

**[0201]** Expression vectors of the invention may be constructed from a convenient starting vector such as a commercially available vector. Such vectors may or may not contain all of the desired flanking sequences. Where one or more of the flanking sequences described herein are not already present in the vector, they may be individually obtained and ligated into the vector. Methods used for obtaining each of the flanking sequences are well known to one skilled in the art.

**[0202]** After the vector has been constructed and a nucleic acid molecule encoding an anti-OPGL antibody has been inserted into the proper site of the vector, the completed vector may be inserted into a suitable host cell for amplification and/or polypeptide expression. The transformation of an expression vector for an anti-OPGL antibody into a selected host cell may be accomplished by well-known methods including methods such as transfection, infection, calcium chloride, electroporation, microinjection, lipofection, DEAE-dextran method, or other known techniques. The method selected will in part be a function of the type of host cell to be used. These methods and other suitable methods are well known to the skilled artisan, and are set forth, for example, in Sambrook et al., supra.

**[0203]** A host cell, when cultured under appropriate conditions, synthesizes an anti-OPGL antibody that can subsequently be collected from the culture medium (if the host cell secretes it into the medium) or directly from the host cell producing it (if it is not secreted). The selection of an appropriate host cell will depend upon various factors, such as desired expression levels, polypeptide modifications that are desirable or necessary for activity (such as glycosylation or phosphorylation) and ease of folding into a biologically active molecule.

**[0204]** Mammalian cell lines available as hosts for expression are well known in the art and include, but are not limited to, many immortalized cell lines available from the American Type Culture Collection (ATCC), such as Chinese hamster

ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), and a number of other cell lines. In certain embodiments, cell lines may be selected through determining which cell lines have high expression levels and produce antibodies with constitutive OPGL binding properties. In another embodiment, a cell line from the B cell lineage that does not make its own antibody but has a capacity to make and secrete a heterologous antibody can be selected.

**[0205]** Antibodies of the invention are useful for detecting OPGL in biological samples and identification of cells or tissues that produce the protein. In certain embodiments, antibodies that bind to OPGL and block interaction with other binding compounds may have therapeutic use in modulating osteoclast differentiation and bone resorption. In certain embodiments, antibodies to OPGL may block OPGL binding to ODAR (RANK), which may result in a block in the signal transduction cascade and loss of NF- $\kappa$ B mediated transcription activation. Assays for measuring NF- $\kappa$ B-mediated transcription activation using, e.g., a luciferase reporter assay, are known to those skilled in the art.

**[0206]** In certain embodiments, antibodies to OPGL may be useful in treatment of bone diseases such as osteoporosis and Paget's disease. In certain embodiments, antibodies can be tested for binding to OPGL in the absence or presence of OPG and examined for their ability to inhibit OPGL-mediated osteoclastogenesis and/or bone resorption.

**[0207]** Anti-OPGL antibodies of the invention can be administered alone or in combination with other therapeutic agents, in particular, in combination with other cancer therapy agents. Such agents generally include radiation therapy or chemotherapy. Chemotherapy, for example, can involve treatment with one or more of the following: anthracyclines, taxol, tamoxifene, doxorubicin, 5-fluorouracil, and other drugs known to the skilled worker.

**[0208]** In addition, anti-OPGL antibodies can be administered to patients in combination with antibodies that bind to tumor cells and induce a cytotoxic and/or cytostatic effect on tumor growth. Examples of such antibodies include those that bind to cell surface proteins Her2, CDC20, CDC33, mucin-like glycoprotein and epidermal growth factor receptor (EGFR) present on tumor cells and induce a cytostatic and/or cytotoxic effect on tumor cells displaying these proteins. Examples of such antibodies include HERCEPTIN for treatment of breast cancer and RITUXAN for the treatment of non-Hodgkin's lymphoma. Also, combination therapy can include as cancer therapy agents polypeptides that selectively induce apoptosis in tumor cells, such as the TNF-related polypeptide TRAIL. Anti-OPGL or antigen binding fragments of the invention can be administered prior to, concurrent with, or subsequent to treatment with a cancer therapy agent. Anti-OPGL antibodies can be administered prophylactically to prevent or mitigate the onset of loss of bone mass by metastatic cancer or can be given for the treatment of an existing condition of loss of bone mass due to metastasis.

**[0209]** Anti-OPGL antibodies of the invention may be used to prevent and/or treat the growth of tumor cells in bone. Cancer that metastasizes to bone can spread readily as tumor cells stimulate osteoclasts to resorb the internal bone matrix. Treatment with an anti-OPGL antibody will maintain bone density by inhibiting resorption and decrease the likelihood of tumor cells spreading throughout the skeleton. Any cancer that metastasizes to bone may be prevented and/or treated with an anti-OPGL antibody.

**[0210]** In one embodiment, multiple myeloma may be prevented and/or treated with an anti-OPGL antibody or antigen binding fragment thereof. Multiple myeloma is localized to bone. Affected patients typically exhibit a loss of bone mass due to increased osteoclast activation in localized regions. Myeloma cells either directly or indirectly produce OPGL, which in turn activates osteoclasts resulting in local bone lysis surrounding the myeloma cells embedded in bone marrow spaces. The normal osteoclasts adjacent to the myeloma cell in turn produce IL-6, leading to growth and proliferation of myeloma cells. Myeloma cells expand in a clonal fashion and occupy bone spaces that are being created by inappropriate bone resorption. Treatment of an animal with an anti-OPGL antibody blocks activation of osteoclasts which in turn leads to a decrease in IL-6 production by osteoclasts, and a suppression of myeloma cell growth and/or proliferation.

**[0211]** Anti-OPGL antibodies may be used alone for the treatment of the above referenced conditions resulting in loss of bone mass or in combination with a therapeutically effective amount of a bone growth promoting (anabolic) agent or a bone anti-resorptive agent including but not limited to bone morphogenic factors designated BMP-1 to BMP-12, transforming growth factor- $\beta$  and TGF- $\beta$  family members, fibroblast growth factors FGF-1 to FGF-10, interleukin-1 inhibitors, TNF $\alpha$  inhibitors, parathyroid hormone, E series prostaglandins, bisphosphonates and bone-enhancing minerals such as fluoride and calcium. Anabolic agents include parathyroid hormone and insulin-like growth factor (IGF), wherein the latter agent is preferably complexed with an IGF binding protein. Preferred embodiments also include the combination of an anti-OPGL antibody with an interleukin-1 (IL-1) receptor antagonist or an anti-OPGL antibody with a soluble TNF receptor, such as soluble TNF receptor-1 or soluble TNF receptor-2. An exemplary IL-1 receptor antagonist is described in WO89/11540 and an exemplary soluble TNF receptor-1 is described in WO98/01555.

**[0212]** In preferred embodiments, the invention provides pharmaceutical compositions comprising a therapeutically effective amount of the antibodies of the invention together with a pharmaceutically acceptable diluent, carrier, solubilizer, emulsifier, preservative and/or adjuvant. In certain embodiments, pharmaceutical compositions comprising a therapeutically effective amount of anti-OPGL antibodies are provided.

**[0213]** Acceptable formulation materials preferably are nontoxic to recipients at the dosages and concentrations employed. The pharmaceutical composition may contain formulation materials for modifying, maintaining or preserving, for example, the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption or penetration of the composition. Suitable formulation materials include, but are not limited to, amino acids (such as glycine, glutamine, asparagine, arginine or lysine); antimicrobials; antioxidants (such as ascorbic acid, sodium sulfite or sodium hydrogen-sulfite); buffers (such as borate, bicarbonate, Tris-HCl, citrates, phosphates or other organic acids); bulking agents (such as mannitol or glycine); chelating agents (such as ethylenediamine tetraacetic acid (EDTA)); complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin or hydroxypropyl-beta-cyclodextrin); fillers; monosaccharides; disaccharides; and other carbohydrates (such as glucose, mannose or dextrans); proteins (such as serum albumin, gelatin or immunoglobulins); coloring, flavoring and diluting agents; emulsifying agents;

hydrophilic polymers (such as polyvinylpyrrolidone); low molecular weight polypeptides; salt-forming counterions (such as sodium); preservatives (such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid or hydrogen peroxide); solvents (such as glycerin, propylene glycol or polyethylene glycol); sugar alcohols (such as mannitol or sorbitol); suspending agents; surfactants or wetting agents (such as pluronics, PEG, sorbitan esters, polysorbates such as polysorbate 20, polysorbate 80, triton, tromethamine, lecithin, cholesterol, tyloxapal); stability enhancing agents (such as sucrose or sorbitol); tonicity enhancing agents (such as alkali metal halides, preferably sodium or potassium chloride, mannitol sorbitol); delivery vehicles; diluents; excipients and/or pharmaceutical adjuvants. (REMYN'TON'S PHARMACEUTICAL SCIENCES, 18<sup>th</sup> Edition, (A. R. Gennaro, ed.), 1990, Mack Publishing Company.

**[0214]** In certain embodiments, optimal pharmaceutical compositions will be determined by one skilled in the art depending upon, for example, the intended route of administration, delivery format and desired dosage. See, for example, REMYN'TON'S PHARMACEUTICAL SCIENCES, *ibid.* Such compositions may influence the physical state, stability, rate of in vivo release and rate of in vivo clearance of the antibodies of the invention.

**[0215]** The primary vehicle or carrier in a pharmaceutical composition may be either aqueous or non-aqueous in nature. For example, a suitable vehicle or carrier may be water for injection, physiological saline solution or artificial cerebrospinal fluid, possibly supplemented with other materials common in compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. Pharmaceutical compositions can comprise Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5, which may further include sorbitol or a suitable substitute therefor. Anti-OPGL antibody compositions may be prepared for storage by mixing the selected composition having the desired degree of purity with optional formulation agents (REMYN'TON'S PHARMACEUTICAL SCIENCES, *ibid.*) in the form of a lyophilized cake or an aqueous solution. Further, the anti-OPGL antibody product may be formulated as a lyophilizate using appropriate excipients such as sucrose.

**[0216]** Formulation components are present in concentrations that are acceptable to the site of administration. Buffers are advantageously used to maintain the composition at physiological pH or at a slightly lower pH, typically within a pH range of from about 5 to about 8.

**[0217]** The pharmaceutical compositions of the invention can be delivered parenterally. When parenteral administration is contemplated, the therapeutic compositions for use in this invention may be in the form of a pyrogen-free, parenterally acceptable aqueous solution comprising the desired anti-OPGL antibody in a pharmaceutically acceptable vehicle. A particularly suitable vehicle for parenteral injection is sterile distilled water in which the anti-OPGL antibody is formulated as a sterile, isotonic solution, properly preserved. Preparation can involve the formulation of the desired molecule with an agent, such as injectable microspheres, bio-erodible particles, polymeric compounds (such as polylactic acid or polyglycolic acid), beads or liposomes, that may provide controlled or sustained release of the product which may then be delivered via a depot injection. Formulation with hyalu-

ronic acid has the effect of promoting sustained duration in the circulation. Implantable drug delivery devices may be used to introduce the desired molecule.

**[0218]** The compositions may be selected for inhalation. In these embodiments, an anti-OPGL antibody is formulated as a dry powder for inhalation, or anti-OPGL antibody inhalation solutions may also be formulated with a propellant for aerosol delivery, such as by nebulization. Pulmonary administration is further described in PCT Application No. PCT/US94/001875, which describes pulmonary delivery of chemically modified proteins.

**[0219]** The pharmaceutical compositions of the invention can be delivered through the digestive tract, such as orally. The preparation of such pharmaceutically acceptable compositions is within the skill of the art. Anti-OPGL antibodies that are administered in this fashion may be formulated with or without those carriers customarily used in the compounding of solid dosage forms such as tablets and capsules. A capsule may be designed to release the active portion of the formulation at the point in the gastrointestinal tract when bioavailability is maximized and pre-systemic degradation is minimized. Additional agents can be included to facilitate absorption of the anti-OPGL antibody. Diluents, flavorings, low melting point waxes, vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders may also be employed.

**[0220]** A pharmaceutical composition may involve an effective quantity of anti-OPGL antibodies in a mixture with non-toxic excipients that are suitable for the manufacture of tablets. By dissolving the tablets in sterile water, or another appropriate vehicle, solutions may be prepared in unit-dose form. Suitable excipients include, but are not limited to, inert diluents, such as calcium carbonate, sodium carbonate or bicarbonate, lactose, or calcium phosphate; or binding agents, such as starch, gelatin, or acacia; or lubricating agents such as magnesium stearate, stearic acid, or talc.

**[0221]** Additional pharmaceutical compositions will be evident to those skilled in the art, including formulations involving anti-OPGL antibodies in sustained- or controlled-delivery formulations. Techniques for formulating a variety of other sustained- or controlled-delivery means, such as liposome carriers, bio-erodible microparticles or porous beads and depot injections, are also known to those skilled in the art. See, for example, PCT Application No. PCT/US93/00829, which describes the controlled release of porous polymeric microparticles for the delivery of pharmaceutical compositions. Sustained-release preparations may include semipermeable polymer matrices in the form of shaped articles, e.g. films, or microcapsules, polyesters, hydrogels, polylactides (U.S. Pat. No. 3,773,919 and EP 058,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman et al., 1983, *Biopolymers* 22: 547-556), poly (2-hydroxyethyl-methacrylate) (Langer et al., 1981, *J. Biomed. Mater. Res.* 15: 167-277) and Langer, 1982, *Chem. Tech.* 12: 98-105), ethylene vinyl acetate (Langer et al., *ibid.*) or poly-D(-)-3-hydroxybutyric acid (EP 133,988). Sustained release compositions may also include liposomes, which can be prepared by any of several methods known in the art. See e.g., Eppstein et al., 1985, *Proc. Natl. Acad. Sci. USA* 82: 3688-3692; EP 036,676; EP 088,046 and EP 143,949.

**[0222]** The pharmaceutical composition to be used for in vivo administration typically is sterile. In certain embodiments, this may be accomplished by filtration through sterile filtration membranes. In certain embodiments, where the

composition is lyophilized, sterilization using this method may be conducted either prior to or following lyophilization and reconstitution. In certain embodiments, the composition for parenteral administration may be stored in lyophilized form or in a solution. In certain embodiments, parenteral compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

**[0223]** Once the pharmaceutical composition of the invention has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or as a dehydrated or lyophilized powder. Such formulations may be stored either in a ready-to-use form or in a form (e.g., lyophilized) that is reconstituted prior to administration.

**[0224]** The invention also provides kits for producing a single-dose administration unit. The kits of the invention may each contain both a first container having a dried protein and a second container having an aqueous formulation, including for example single and multi-chambered pre-filled syringes (e.g., liquid syringes, lyosyringes or needle-free syringes).

**[0225]** The effective amount of an anti-OPGL antibody pharmaceutical composition to be employed therapeutically will depend, for example, upon the therapeutic context and objectives. One skilled in the art will appreciate that the appropriate dosage levels for treatment, according to certain embodiments, will thus vary depending, in part, upon the molecule delivered, the indication for which the anti-OPGL antibody is being used, the route of administration, and the size (body weight, body surface or organ size) and/or condition (the age and general health) of the patient. In certain embodiments, the clinician may titer the dosage and modify the route of administration to obtain the optimal therapeutic effect. Typical dosages range from about 0.1  $\mu\text{g}/\text{kg}$  to up to about 30  $\text{mg}/\text{kg}$  or more, depending on the factors mentioned above. In certain embodiments, the dosage may range from 0.1  $\mu\text{g}/\text{kg}$  up to about 30  $\text{mg}/\text{kg}$ ; or 1  $\mu\text{g}/\text{kg}$  up to about 30  $\text{mg}/\text{kg}$ ; or 5  $\mu\text{g}/\text{kg}$  up to about 30  $\text{mg}/\text{kg}$ .

**[0226]** Dosing frequency will depend upon the pharmacokinetic parameters of the anti-OPGL antibody in the formulation used. For example, a clinician will administer the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose, or as two or more doses (which may or may not contain the same amount of the desired molecule) over time, or as a continuous infusion via an implantation device or catheter. Further refinement of the appropriate dosage is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely performed by them. Appropriate dosages may be ascertained through use of appropriate dose-response data.

**[0227]** Administration routes for the pharmaceutical compositions of the invention include orally, through injection by intravenous, intraperitoneal, intracerebral (intra-parenchymal), intracerebroventricular, intramuscular, intra-ocular, intraarterial, intraportal, or intralesional routes; by sustained release systems or by implantation devices. The pharmaceutical compositions may be administered by bolus injection or continuously by infusion, or by implantation device. The pharmaceutical composition also can be administered locally via implantation of a membrane, sponge or another appropriate material onto which the desired molecule has been absorbed or encapsulated. Where an implantation device is used, the device may be implanted into any suitable tissue or

organ, and delivery of the desired molecule may be via diffusion, timed-release bolus, or continuous administration.

**[0228]** It also may be desirable to use anti-OPGL antibody pharmaceutical compositions according to the invention *ex vivo*. In such instances, cells, tissues or organs that have been removed from the patient are exposed to anti-OPGL antibody pharmaceutical compositions after which the cells, tissues and/or organs are subsequently implanted back into the patient.

**[0229]** In particular, anti-OPGL antibody can be delivered by implanting certain cells that have been genetically engineered, using methods such as those described herein, to express and secrete the polypeptide. In certain embodiments, such cells may be animal or human cells, and may be autologous, heterologous, or xenogeneic, or may be immortalized. In certain embodiments, the cells may be immortalized. In other embodiments, in order to decrease the chance of an immunological response, the cells may be encapsulated to avoid infiltration of surrounding tissues. In further embodiments, the encapsulation materials are typically biocompatible, semi-permeable polymeric enclosures or membranes that allow the release of the protein product(s) but prevent the destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissues.

#### EXAMPLES

**[0230]** The following examples, including the experiments conducted and results achieved are provided for illustrative purposes only and are not to be construed as limiting the present invention.

##### Example 1

##### Production of Human Monoclonal Antibodies Against OPGL

###### Transgenic HuMab Mice

**[0231]** Fully human monoclonal antibodies to OPGL were prepared using HCo7, HCo12, and HCo7+HCo12 strains of transgenic mice, each of which expresses human antibody genes. In each of these mouse strains, the endogenous mouse kappa light chain gene has been homozygously disrupted (as described in Chen et al., 1993, *EMBO J.* 12: 811-820) and the endogenous mouse heavy chain gene has been homozygously disrupted as described in Example 1 of PCT Publication WO 01/09187 (incorporated by reference). Each of these mouse strains carries a human kappa light chain transgene, KCo5 (as described in Fishwild et al., 1996, *Nature Biotechnology* 14: 845-851). The HCo7 strain carries the HCo7 human heavy chain transgene as described in U.S. Pat. Nos. 5,545,806; 5,625,825; and 5,545,807 (incorporated by reference). The HCo12 strain carries the HCo12 human heavy chain transgene as described in Example 2 of PCT Publication WO 01/09187 (incorporated by reference). The HCo7+HCo12 strain carries both the HCo7 and the HCo12 heavy chain transgenes and is hemizygous for each transgene. All of these strains are referred to herein as HuMab mice.

###### HuMab Immunizations:

**[0232]** To generate fully human monoclonal antibodies to OPGL, HuMab mice were immunized with purified recombinant OPGL derived from *E. coli* or CHO cells as antigen. General immunization schemes for HuMab mice are described in Lonberg et al. 91994, *Nature* 368: 856-859,

Fishwild et al., *ibid.*, and PCT Publication WO 98/24884 (the teachings of each of which are incorporated by reference). Mice were 6-16 weeks of age upon the first infusion of antigen. A purified recombinant preparation (50-100  $\mu$ g) of OPGL antigen (e.g., purified from transfected *E. coli* or CHO cells expressing OPGL) was used to immunize the HuMab mice intraperitoneally (IP) or subcutaneously (Sc).

**[0233]** Immunizations of HuMab transgenic mice were performed twice using antigen in complete Freund's adjuvant, followed by 2-4 weeks IP immunization (up to a total of 9 immunizations) with the antigen in incomplete Freund's adjuvant. Several dozen mice were immunized for each antigen. A total of 136 HuMab mice of the HCo7, HCo12, and HCo7+HCo12 strains were immunized with OPGL. The immune response was monitored by retroorbital bleeds.

**[0234]** To select HuMab mice producing antibodies that bound OPGL, sera from immunized mice was tested by ELISA as described by Fishwild et al. *supra*. Briefly, microtiter plates were coated with purified recombinant OPGL from CHO cells or *E. coli* at 1-2  $\mu$ L/mL in PBS and 50  $\mu$ L/well incubated at 4° C. overnight, then blocked with 200  $\mu$ L/well with 5% chicken serum in PBS/Tween (0.05%). Dilutions of plasma from OPGL-immunized mice were added to each well and incubated for 1-2 hours at ambient temperature. The plates were washed with PBS/Tween and then incubated with a goat anti-human IgG Fc-specific polyclonal reagent conjugated to horseradish peroxidase (HRP) for 1 hour at room temperature. After washing, the plates were developed with ABTS substrate (Sigma, A-1888, 0.22 mg/mL) and analyzed at OD of 415-495. Mice with sufficient titers of anti-OPGL human immunoglobulin were used to produce monoclonal antibodies as described below.

##### Generation of Hybridomas Producing Human Monoclonal Antibodies to OPGL

**[0235]** Mice were prepared for monoclonal antibody production by boosting with antigen intravenously 2 days before sacrifice, and spleens were removed thereafter. The mouse splenocytes were isolated from the HuMab mice and fused with PEG to a mouse myeloma cell line based upon standard protocols. Typically, 10-20 fusions for each antigen were performed.

**[0236]** Briefly, single cell suspensions of splenic lymphocytes from immunized mice were fused to one-quarter the number of P3X63-Ag8.653 nonsecreting mouse myeloma cells (ATCC, CRL 1580) using 50% PEG (Sigma Chemical Co., St. Louis, Mo.). Cells were plated at approximately  $1 \times 10^5$  cells/well in flat bottom microtitre plates, followed by about two week incubation in selective medium containing 10% fetal bovine serum, 10% P388D1 (ATCC, CRL TIB-63) conditioned medium and 3-5% origen (IGEN) in DMEM (Mediatech, CRL 10013, with high glucose, L-glutamine and sodium pyruvate) plus 5 mM HEPES, 0.055 mM 2-mercaptoethanol, 50 mg/mL gentamycin and 1xHAT (Sigma, CRL P-7185). After 1-2 weeks, cells were cultured in medium in which the HAT was replaced with HT.

**[0237]** The resulting hybridomas were screened for the production of antigen-specific antibodies. Individual wells were screened by ELISA (described above) for human anti-OPGL monoclonal IgG antibodies. Once extensive hybridoma growth occurred, medium was monitored usually after 10-14 days. Antibody secreting hybridomas were replated, screened again and, if still positive by ELISA for human IgG, anti-OPGL monoclonal antibodies were subcloned at least twice

by limiting dilution. The stable subclones were then cultured in vitro to generate small amounts of antibody in tissue culture medium for characterization.

Selection of Human Monoclonal Antibodies that Bind to OPGL

**[0238]** An ELISA assay as described above was used to screen for hybridomas that showed positive reactivity with OPGL immunogen. Hybridomas secreting a monoclonal antibody that bound with high avidity to OPGL were subcloned and further characterized. One clone from each hybridoma, which retained the reactivity of parent cells (as determined by ELISA), was chosen for making a 5-10 vial cell bank stored in liquid nitrogen.

**[0239]** An isotype-specific ELISA was performed to determine the isotype of the monoclonal antibodies produced as disclosed herein. In these experiments, microtitre plate wells were coated with 50  $\mu$ L/well of a solution of 1  $\mu$ g/mL mouse anti-human kappa light chain in PBS and incubated at 4° C. overnight. After blocking with 5% chicken serum, the plates were reacted with supernatant from each tested monoclonal antibody and a purified isotype control. Plates were incubated at ambient temperature for 1-2 hours. The wells were then reacted with either human IgG<sub>1</sub> or IgG<sub>3</sub>-specific horseradish peroxidase-conjugated goat anti-human polyclonal antisera and plates developed and analyzed as described above.

**[0240]** Monoclonal antibodies purified from six hybridoma supernatants that showed significant binding to OPGL as detected by ELISA were further tested for biological activity using in vitro receptor binding assays and human OPGL-dependent in vitro osteoclast forming assays (described in Example 6 below). The antibodies selected were designated 16E1, 2E11, 18B2, 2D8, 22B3, and 9H7. The heavy chain alignment for these anti-OPGL antibodies is shown in FIG. 15. The light chain alignment for the anti-OPGL antibodies is shown in FIG. 16. Non-consensus sequences are shown in bold and are shaded, and complementarity-determining regions (CDRs) are underlined.

### Example 2

#### Cloning the 9H7 Anti-OPGL Heavy and Light Chains

**[0241]** Cloning of the 9H7 anti-OPGL MAb light chain

**[0242]** Three anti-OPGL hybridoma light chain cDNAs (9H7, 16E1 and 18B2) were cloned into pDSR19 mammalian cell expression vector. The construction of a plasmid encoding the 9H7 kappa light chain is explicitly described; cloning of the other light chain species was performed using similar procedures. The anti-OPGL-9H7 kappa light chain variable region was obtained using polymerase chain reaction (PCR) amplification methods from first strand cDNA prepared from hybridoma 9H7 total RNA. First strand cDNA was prepared from 9H7 total RNA using a random primer with an extended 5'-adapter (5'-GGCCGGATAGGCCCTCACNNNNNNT-3', SEQ ID NO: 53) and the materials and methods provided by the Gibco SuperScript II™ Preamplification System for First Strand cDNA Synthesis kit (Catalogue No. 18089-011). The oligonucleotides below were used for the PCR:

5' GeneRacer™ (Invitrogen) primer  
(SEQ ID NO: 54):  
5'-GGA CAC TGA CAT GGA CTG AAG GAG TA- 3';

3' kappa RACE primer, 2310-03 (SEQ ID NO: 55):  
5'-GGG GTC AGG CTG GAA CTG AGG-3'.

**[0243]** The amplified DNAs were cloned into pCRII-TOPO (Invitrogen) and the resulting plasmids were sequenced. The kappa chain consensus sequence was used to design primers for PCR amplification of the variable region of the 9H7 kappa chain. To generate the signal sequence, a three-step PCR was performed. First, primers 2669-73 and 2708-53 (set forth below) were used with a 9H7 cDNA light chain clone template. Conditions used for the reaction were: 94° C. for 1 minute; 94° C. for 20 seconds, 42° C. for 30 seconds, 74° C. for 150 seconds for 2 cycles; 94° C. for 20 seconds, 56° C. for 30 seconds, 74° C. for 150 seconds for 25 cycles; and 74° C. for 7 minutes with Pfu polymerase and the appropriate buffer and nucleotides. The PCR product was then amplified with primers 2663-07 and 2708-53 followed by amplification with primers 2663-08 and 2708-53. These primers are shown below.

2663-08 (SEQ ID NO: 56)

HindIII XbaI Kozak  
5'-C AGC AG AAGCTTCTAGA CCACC ATG GAC ATG AGG GTG  
CCC GCT CAG CTC CTG GG-3';

2663-07 (SEQ ID NO: 57)

5'-CC GCT CAG CTC CTG GGG CTC CTG CTG TGG CTG  
AGA GGT GCC AGA T-3';

2669-73 (SEQ ID NO: 58)

5'-G TGG TTG AGA GGT GCC AGA TGT GAA ATT GTG CTG  
ACC CAG TCT CCA GCC ACC CTG TCT TTG TCT C-3';

2708-53 (SEQ ID NO: 59)

Sall  
5'-CTT GTC GAC TCA ACA CTC TCC CCT GTT GAA GCT  
C-3'.

**[0244]** The PCR reactions generated a 741 bp fragment encoding 238 amino acid residues (including the 22 amino acid signal sequence) that was purified using a QIAquick PCR Purification kit (Qiagen Cat. No. 28104), cut with XbaI and Sall, and Qiagen purified again. This fragment, containing the complete light chain with a 5'Kozak (translational initiation) site and the following signal sequence for mammalian expression:

MDMRVPAQLLGLLLLWLRGARC, (SEQ ID NO: 60)

was ligated into pDSR $\alpha$ 19 to generate plasmid pDSR $\alpha$ 19:9H7 kappa (FIG. 17). pDSR $\alpha$ 19 has been described previously (see International Application, Publication No. WO 90/14363, which is herein incorporated by reference for any purpose). Briefly, to make pDSR $\alpha$ 19, pDSR $\alpha$ 2 was modified in the following ways: the sequence containing the transcription termination/polyadenylation signal from the alpha sub-

unit of the bovine pituitary glycoprotein hormone alpha-FSH (follicle-stimulating hormone) was shortened by approximately 1400 base pairs, to 885 base pairs, and ends at the NdeI site after modification; the dihydrofolate reductase (DHFR) promoter contained 209 base pairs, having been shortened from the 5' end by approximately 1 kilobase; and an approximately 550 base pair BglII fragment from the DHFR polyA sequence was deleted.

[0245] The 9H7 kappa light chain expression clone was sequenced to confirm that it encoded the same peptide that was identified in the 9H7 hybridoma. The final expression vector, pDSR $\alpha$ 19:9 H7 kappa is 5479 by and contains the 7 functional regions described in Table 2.

cDNA Synthesis kit (Cat. No. 18089-011). The oligonucleotides below were used for the PCR:

5' heavy chain RACE primer, 2508-02  
(SEQ ID NO: 61):  
5'-(CG)AG GT(CG) CAG (CT)T(GT) GTG (CG)AG TC- 3';

3' heavy chain RACE primer, 2420-54  
(SEQ ID NO: 62):  
5'-CTG AGT TCC ACG ACA CC-3'.

[0248] Amplified DNA was cloned into pCRII-TOPO (Invitrogen) and the resulting plasmids were sequenced. The heavy chain consensus sequence was used to design primers

TABLE 2

Plasmid Base Pair Number:	
2 to 881	A transcription termination/polyadenylation signal from the $\alpha$ -subunit of the bovine pituitary glycoprotein hormone ( $\alpha$ -FSH) (Goodwin, et al., 1983, Nucleic Acids Res. 11: 6873-82; Genbank Accession Number X00004)
882 to 2027	A mouse dihydrofolate reductase (DHFR) minigene containing the endogenous mouse DHFR promoter, the cDNA coding sequences, and the DHFR transcription termination/polyadenylation signals (Gasser et al, 1982, Proc. Natl. Acad. Sci. U.S.A. 79: 6522-6; Nunberg et al., 1980, Cell 19: 355-64; Setzer et al., 1982, J. Biol. Chem. 257: 5143-7; McGrogan et al., 1985, J. Biol. Chem. 260: 2307-14)
2031 to 3947	pBR322 sequences containing the ampicillin resistance marker gene and the origin for replication of the plasmid in <i>E. coli</i> (Genbank Accession Number J01749)
3949 to 4292	An SV40 early promoter, enhancer and origin of replication (Takebe et al., 1988, Mol. Cell Biol. 8: 466-72, Genbank Accession Number J02400)
4299 to 4565	A translational enhancer element from the HTLV-1 LTR domain (Seiki et al., 1983, Proc. Natl. Acad. Sci. U.S.A. 80: 3618-22, Genbank Accession Number J02029)
4574 to 4730	An intron from the SV40 16S, 19S splice donor/acceptor signals (Okayama and Berg, 1983, Mol Cell Biol 3: 280-9, Genbank Accession Number J02400)
4755 to 5479	The 9H7D4 kappa light chain cDNA between the XbaI and SalI sites

#### Construction of pDSR19:hlgG1 C<sub>H</sub>

[0246] A pDSR19: rat variable region/human constant region IgG<sub>1</sub> plasmid was constructed using a three-piece ligation of a rat variable region sequence, the human constant region (CH1, hinge, CH2, and CH3 domains) and pDSR19. The linear pDSR $\alpha$ 19:hlgG1 C<sub>H</sub> plasmid was prepared by digesting the pDSR19:rat variable region/human constant region IgG<sub>1</sub> plasmid with restriction enzymes XbaI and BsmBI to remove the coding portion of the rat variable region. The resulting linear plasmid containing the 1.0 kbp human IgG<sub>1</sub> constant region domain (C<sub>H</sub>1, hinge, C<sub>H</sub>2 and C<sub>H</sub>3 domains) was gel isolated and used to accept hybridoma derived  $\alpha$ OPGL variable regions.

#### Cloning of the 9H7 Anti-OPGL MAb Heavy Chain

[0247] Three anti-OPGL hybridoma IgG<sub>1</sub> heavy chain cDNAs; 9H7, 16E1 and 18B2, were cloned into pDSR19 mammalian cell expression vector. The construction of a plasmid encoding the 9H7 IgG<sub>1</sub> heavy chain is explicitly described here; the other hybridoma heavy chains were cloned using similar procedures. The anti-OPGL-9H7 heavy chain variable region was obtained using PCR amplification methods from first strand cDNA prepared from hybridoma 9H7 total RNA. First strand cDNA was prepared from 9H7 total RNA using a random primer with an extended 5'-adapter (5'-GGCCGGATAGGCCTCACNNNNNNT-3', SEQ ID NO: 53) and the materials and methods provided by the Gibco SuperScript II<sup>TM</sup> Preamplification System for First Strand

for PCR amplification of the variable region of the 9H7 heavy chain. To generate the signal sequence, a three-step PCR was performed. First, primers 2512-98 and 2673-14 were used with a 9H7 heavy chain cDNA clone template. Conditions used for the reaction were: 94° C. for 1 minute; 94° C. for 20 seconds, 42° C. for 30 seconds, 74° C. for 150 seconds for 2 cycles; 94° C. for 20 seconds, 56° C. for 30 seconds, 74° C. for 150 seconds for 25 cycles; and 74° C. for 7 minutes with Pfu polymerase and the appropriate buffer and nucleotides. The PCR product was then amplified with primers 2663-07 and 2673-14 followed by amplification with primers 2663-08 and 2673-14. The primers are shown below.

2663-08 (SEQ ID NO: 63)

HindIII XbaI Kozak  
5'-C AGC AG AAGCTTCTAGA CCACC ATG GAC ATG AGG GTG  
CCC GCT CAG CTC CTG GG-3';

2663-07 (SEQ ID NO: 64)

5'-CC GCT CAG CTC CTG GGG CTC CTG CTG CTG TGG CTG  
AGA GGT GCC AGA T-3';

2512-98 (SEQ ID NO: 65)

5'-G TGG TTG AGA GGT GCC AGA TGT GAG GTG CAG CTG  
GTG CAG TCT-3';



-continued

2673-14

(SEQ ID NO: 66)

BsmBI

5'-GT GGA GGC ACT AGA GAC GGT GAC CAG GGC TCC CTG  
GCC CCA GGG GTC GAA-3'.

[0249] The PCR reactions generated a 443 by fragment encoding 138 amino acid residues (including the 22 amino acid signal sequence) that was purified using a QIAquick PCR Purification kit (Qiagen Cat. No. 28104), cut with XbaI and BsmBI, and Qiagen purified again. This fragment, con-

taining the heavy chain with a 5' Kozak (translational initiation) site and the following signal sequence for mammalian expression:

(SEQ ID NO: 60)

MDMRVPAQLLGLLLLWLRGARC,  
was ligated into pDSR $\alpha$ 19:hIgG1 C<sub>H</sub> to generate plasmid pDSR $\alpha$ 19:9 H7 IgG1 (FIG. 18).

[0250] The 9H7 IgG<sub>i</sub> heavy chain expression clone was sequenced to confirm that it encoded the same peptide that was identified in the 9H7 hybridoma. The final expression vector, pDSR $\alpha$ 19:rat variable region/human constant region IgG<sub>i</sub> is 6158 by and contains the 7 functional regions described in Table 3.

TABLE 3

Plasmid Base Pair Number:	
2 to 881	A transcription termination/polyadenylation signal from the $\alpha$ -subunit of the bovine pituitary glycoprotein hormone ( $\alpha$ -FSH) (Goodwin, et al., 1983, Nucleic Acids Res. 11:6873-82; Genbank Accession Number X00004)
882 to 2027	A mouse dihydrofolate reductase (DHFR) minigene containing the endogenous mouse DHFR promoter, the cDNA coding sequences, and the DHFR transcription termination/polyadenylation signals (Gasser et al, 1982, Proc. Natl. Acad. Sci. U.S.A. 79:6522-6 Nunberg et al., 1980, Cell 19355-64 Setzer et al., 1982, J. Biol. Chem. 257:5143-7; McGrogan et al., 1985, J. Biol. Chem. 260:2307-14)
2031 to 3947	pBR322 sequences containing the ampicillin resistance marker gene and the origin for replication of the plasmid in <i>E. coli</i> (Genbank Accession Number J01749)
3949 to 4292	An SV40 early promoter, enhancer and origin of replication (Takebe et al., 1988, Mol. Cell Biol. 8:466-72 Genbank Accession Number J02400)
4299 to 4565	A translational enhancer element from the HTLV-1 LTR domain (Seiki et al., 1983, Proc. Natl. Acad. Sci. U.S.A. 80:3618-22, Genbank Accession Number J02029)
4574 to 4730	An intron from the SV40 16S, 19S splice donor/acceptor signals (Okayama and Berg, 1983. Mol. Cell Biol. 3:280-9 Genbank Accession Number J02400)
4755 to 6158	The rVh/hCh1 heavy chain cDNA between the XbaI and SalI sites. The sequences of which follows (SEQ ID NO: 67): XbaI TCTAG ACCACCATGG ACATCAGGCT CAGCTTAGTT TTCCTGTCC TTTTCATAAA AGGTGTCCAG TGTGAGGTAG AACTGGTGGG GTCTGGGGGC GGCTTAGTAC AACCTGGAAG GTCCATGACA CTCTCCTGTG CAGCCTCGGG ATTCACTTTC AGAACCTATG GCATGGCCTG GGTCCGCCAG GCCCAACGA AGGGTCTGGA GTGGGTCTCA TCAATTACTG CTAGTGGTGG TACCACCTAC TATCGAGACT CCGTGAAGGG CCGCTTCACT ATTTTATAGG ATAATGCAAA AAGTACCCTA TACCTGCAGA TGGACAGTCC GAGGTCTGAG GACACGGCCA CTTATTTCTG TACATCAATT BsmBI TCGGAATACT GGGGCCACGG AGTCATGGTC ACCGTCTCTA GTGCCTCCACCAAGGGCCCA TCGGTCTTCC CCCTGGCACC CTCCTCCAAG AGCACCTCTGGGGCACAGC GGCCCTGGGC TGCCTGGTCA AGGACTACTT CCCGAACCG GTGACGGTGT CGTGGAATC AGGCGCCCTG ACCAGCGGCG TGACACCTT CCCCTGTGCT CTACAGTCTT CAGGACTCTA CTCCTCAGC AGCGTGGTGACCGTGCCTC CAGCAGCTTG GGCACCCAGA CCTACATCTG CAACGTGAATCACAAGCCCA GCAACACCAA GGTGGACAAG AAAGTTGAGC CCAAATCTTG TGACAAAAC CACACATGCC CACCGTGCCT AGCACCTGAA CTCCTGGGGG GACCGTCACT CTTCTCTTTC CCCCCAAAAC CCAAGGACAC CCTCATGATC TCCCGGACCC CTGAGGTGAC ATGCGTGGTG GTGGACGTGA GCCACGAAGACCTGAGGTC AAGTTCAACT GGTACGTGGA CGGCGTGGAG GTGCATAATG CCAAGACAAA GCCGCGGGAG GAGCAGTACA ACAGCACGTA CCGTGTGGTC AGCGTCCTCA CCGTCTGCA CCAGGACTGG CTGAATGGCA

TABLE 3-continued

Plasmid Base  
Pair Number:

```

AGGAGTACAAGTGCAAGGTC TCCAACAAAG CCCTCCCAGC
CCCCATCGAG AAAACCATCTCCAAGCCAA AGGGCAGCCC
CGAGAACCAC AGGTGTACAC CCTGCCCCCA TCCCGGGATG
AGCTGACCAA GAACCAGGTC AGCCTGACCT GCCTGGTCAA
AGGCTTCTAT CCCAGCGACA TCGCCGTGGA GTGGGAGAGC
AATGGGCAGCCGGAGAACAA CTACAAGACC ACGCTCCCG
TGCTGGACTC CGACGGCTCC TICTICICT ATAGCAAGCT
CACCGTGAC AAGAGCAGGT GGCAGCAGGG GAACGTCTTC
TCATGCTCCG TGATGCATGA GGCTCTGCAC AACCACTACA
CGAGAAGAG CCTCTCCCTG TCTCCGGGTA
      SalI
AATGATAAGT CGAC

```

## Example 3

## 9H7 Anti-OPGL MAb Expression in CHO Cells

[0251] Recombinant anti-OPGL antibodies are produced by Chinese hamster ovary cells, specifically CHO AM-1/D, as disclosed in U.S. Pat. No. 6,210,924 (incorporated by reference). DNA sequences encoding the complete heavy or light chains of each anti-OPGL antibody of the invention are cloned into expression vectors such as those described above. CHO AM-1/D cells are co-transfected with an expression vector capable of expressing a complete heavy chain and an expression vector expressing the complete light chain of the appropriate anti-OPGL antibody. For example, to generate the 22B3 antibody, cells are co-transfected with a vector capable of expressing a complete heavy chain comprising the amino acid sequence as set forth in SEQ ID NO: 30 and a vector capable of expressing a complete light chain comprising the amino acid sequence set forth in SEQ ID NO: 32. To generate the 2E11 antibody, cells are co-transfected with a

vector capable of expressing a complete heavy chain comprising the amino acid sequence as set forth in SEQ ID NO: 38 and a vector capable of expressing a complete light chain comprising the amino acid sequence set forth in SEQ ID NO: 40. To generate the 18B2 antibody, cells are co-transfected with a vector capable of expressing a complete heavy chain comprising the amino acid sequence as set forth in SEQ ID NO: 42 and a vector capable of expressing a complete light chain comprising the amino acid sequence set forth in SEQ ID NO: 44. To generate the 16E1 antibody, cells are co-transfected with a vector capable of expressing a complete heavy chain comprising the amino acid sequence as set forth in SEQ ID NO: 46 and a vector capable of expressing a complete light chain comprising the amino acid sequence set forth in SEQ ID NO: 48. To generate the 9H7 antibody, cells are co-transfected with a vector capable of expressing a complete heavy chain comprising the amino acid sequence as set forth in SEQ ID NO: 50 and a vector capable of expressing a complete light chain comprising the amino acid sequence set forth in SEQ ID NO: 52. Table 4 summarizes the complete heavy and complete light chains for the various OPGL antibodies.

TABLE 4

Antibody	Heavy Chain Variable Region + Heavy Chain Constant Region	Complete Heavy Chain
22B3	SEQ ID NO: 6 + SEQ ID NO: 2	SEQ ID NO: 30
2E11	SEQ ID NO: 10 + SEQ ID NO: 2	SEQ ID NO: 34
2D8	SEQ ID NO: 14 + SEQ ID NO: 2	SEQ ID NO: 38
18B2	SEQ ID NO: 18 + SEQ ID NO: 2	SEQ ID NO: 42
16E1	SEQ ID NO: 22 + SEQ ID NO: 2	SEQ ID NO: 46
9H7	SEQ ID NO: 26 + SEQ ID NO: 2	SEQ ID NO: 50
Antibody	Light Chain Variable Region + Light Chain Constant Region	Complete Light Chain
22B3	SEQ ID NO: 8 + SEQ ID NO: 4	SEQ ID NO: 32
2E11	SEQ ID NO: 12 + SEQ ID NO: 4	SEQ ID NO: 36
2D8	SEQ ID NO: 16 + SEQ ID NO: 4	SEQ ID NO: 40
18B2	SEQ ID NO: 20 + SEQ ID NO: 4	SEQ ID NO: 44
16E1	SEQ ID NO: 24 + SEQ ID NO: 4	SEQ ID NO: 48
9H7	SEQ ID NO: 28 + SEQ ID NO: 4	SEQ ID NO: 52

vector capable of expressing a complete heavy chain comprising the amino acid sequence as set forth in SEQ ID NO: 34 and a vector capable of expressing a complete light chain comprising the amino acid sequence set forth in SEQ ID NO: 36. To generate the 2D8 antibody, cells are co-transfected with a vector capable of expressing a complete heavy chain comprising the amino acid sequence as set forth in SEQ ID

[0252] Stable expression of the 9H7 anti-OPGL MAb was achieved by co-transfection of pDSR $\alpha$ 19:9 H7 IgG<sub>i</sub> and pDSR $\alpha$ 19:9 H7 kappa plasmids into dihydrofolate reductase deficient (DHFR<sup>-</sup>) serum-free adapted Chinese hamster ovary cells (CHO AM-1/D, U.S. Pat. No. 6,210,924) using the art-recognized calcium phosphate method. Transfected cells were selected in 96 well plates in medium containing dia-

lyzed serum but not containing hypoxanthine-thymidine to ensure the growth of cells expressing the DHFR enzyme. Over 5000 transfected clones were screened using assays such as HTRF (homogeneous time resolved fluorescence) and ELISA in order to detect expression of 9H7 anti-OPGL MAb in the conditioned medium. The highest expressing clones were selected for single cell cloning and creation of cell banks.

#### Example 4

##### Production of Anti-OPGL Antibodies

**[0253]** Anti-OPGL antibodies are produced by expression in a clonal line of CHO cells. For each production run, cells from a single vial are thawed into serum-free cell culture media. The cells are grown initially in a T-flask and are serially expanded through a series of spinner flasks until sufficient inoculum has been generated to seed a 20 L bioreactor. Following growth for 5-10 days, the culture is then used to inoculate a 300 L bioreactor. Following growth for an additional 5-10 days, the culture is used to inoculate a 2000 L bioreactor. Production is carried out in a 2000 L bioreactor using a fed batch culture, in which a nutrient feed containing concentrated media components is added to maintain cell growth and culture viability. Production lasts for approximately two weeks during which time anti-OPGL antibody is constitutively produced by the cells and secreted into the cell culture medium.

**[0254]** The production reactor is controlled at set pH, temperature, and dissolved oxygen level: pH is controlled by carbon dioxide gas and sodium carbonate addition; dissolved oxygen is controlled by air, nitrogen, and oxygen gas flows.

**[0255]** At the end of production, the cell broth is fed into a disk stack centrifuge and the culture supernatant is separated from the cells. The concentrate is further clarified through a depth filter followed by a 0.2  $\mu\text{m}$  filter. The clarified conditioned media is then concentrated by tangential flow ultrafiltration. The conditioned media is concentrated 15- to 30-fold. The resulting concentrated conditioned medium is then either processed through purification or frozen for purification at a later date. FIG. 19 depicts an exemplary cell culture process for producing an anti-OPGL antibody.

#### Example 5

##### Screening of Antibodies for Binding to OPGL by BIAcore

**[0256]** All experiments were performed on a BIAcore 2000 according to the manufacturer's instructions, with the following modifications. Experiments were performed at room temperature using a running buffer containing 10 mM Hepes (pH 7.4), 0.15M NaCl, 3 mM EDTA, and 0.005% Tween 20. Protein G at 50  $\mu\text{g}/\text{mL}$  in 10 mM acetate pH 4.5 was immobilized to a level of 1,600 response units (RU) onto a CM5 Research grade sensor chip (BIAcore, Inc.). Antibodies (8-20  $\mu\text{g}/\text{mL}$ ) were captured onto the Protein G chip at a level of 300-400 RUs. CHO human OPGL (hOPGL) 140 or *E. coli* mouse OPGL (mOPGL) 158 were passed over the immobilized antibodies at concentrations of 0.25-62 nM. Langmuir 1:1 model was used to determine binding kinetics. Protein G immobilized to 1600 RUs was used as a blank surface. A mouse monoclonal antibody was used as a positive control to show binding to hOPGL 140 and to monitor surface stability.

**[0257]** All anti-OPGL antibodies showed strong binding to CHO hOPGL 140. 22B3 appears to have a slower off rate than the other antibodies tested. No binding of *E. coli* mOPGL 158 was detected. The results are summarized in Table 5.

TABLE 5

Ab	hOPGL 140				off rate	
	ka (1/Ms)	kd (1/s)	KD (1/M)	KD	half life t $\frac{1}{2}$ (s)	mOPGL 158
9H7	1.27E+06	2.26E-04	1.93E-10	190 pm	3067	no binding
18B2	8.78E+05	1.86E-04	2.11E-10	210 pm	3726	no binding
2D8	1.97E+06	1.81E-04	9.20E-11	92 pm	3829	no binding
2E11	4.53E+05	1.32E-04	2.92E-10	290 pm	5251	no binding
16E1	2.16E+06	1.37E-04	6.33E-11	63 pm	5059	no binding
22B3	1.90E+06	6.39E-05	3.37E-11	34 pm	10847	no binding

#### Example 6

##### Anti-OPGL Antibody Neutralizing Activity

##### Inhibition of Osteoclast Formation

**[0258]** RAW 264.7 (Accession No. TIB-71, American Type Culture Collection, Manassas, Va.) is a murine macrophage cell line that was derived from an Abelson murine leukemia virus-induced tumor. RAW 264.7 cells will differentiate to osteoclast-like cells in the presence of OPGL. An assay for generation of osteoclasts in culture from RAW cells in the presence of OPGL has been described in detail by Hsu et al., 1999, *Proc. Natl. Acad. Sci. U.S.A.* 96:3540-3545, which is incorporated by reference herein.

**[0259]** RAW cells can be stimulated by OPGL ligand to differentiate into osteoclast-like cells, and the differentiation can be measured by tartrate-resistant acid phosphatase (TRAP) activity, a property of osteoclasts. This activity provides the basis for characterizing anti-OPGL antibodies produced according to the invention, by assaying the effect of said antibodies on osteoclastogenesis.

**[0260]** RAW cells were incubated for 4 days in the presence of a constant amount of OPGL (40 ng/mL) and varying amounts of anti-OPGL antibody (6.3 ng/mL to 200 ng/mL) in cell culture medium (DMEM, 10% FBS, 0.3 mg/mL L-glutamine, 100 units/mL penicillin G, 100  $\mu\text{g}/\text{mL}$  streptomycin sulfate). At the end of 4 days, the cells were stained for tartrate-resistant acid phosphatase (TRAP) activity by permeabilization and acidification, followed by treatment with p-nitrophenylphosphate (PNPP) for 5 minutes. Briefly, the media was aspirated from the cells, and 100  $\mu\text{L}$  of citrate buffer (having a formula of 410 mL 0.1M citric acid, 590 mL 0.1 M citrate, trisodium salt and 1 mL Triton X-100) was added to each well and the plates incubated for 3 to 5 minutes at room temperature. One hundred microliters of PNPP solution (having a formula of 157.8 mg acid phosphatase reagent (Sigma 104-100), 7.2 mL tartrate solution (Sigma Cat. No. 387-3), and 22.8 mL citrate buffer) was then added, and plates were incubated for 3 to 5 minutes at room temperature. The reaction was terminated by addition of 50  $\mu\text{L}$  0.5 M NaOH solution.

**[0261]** TRAP converts p-nitrophenylphosphate to para-nitrophenol, which can be quantitated by optical density measurement at 405 nm. The TRAP activity, which is a surrogate marker for osteoclast development, therefore correlates with the optical density at 405 nm. A plot of optical density versus

anti-OPGL antibody concentration is shown in FIG. 20, and demonstrates that anti-OPGL antibody inhibited osteoclast formation in this assay in a dose-dependent manner. IC<sub>50</sub> values were calculated using the forecast function, and are shown in Table 6. An alkaline phosphatase-linked rat polyclonal anti-human OPGL antibody (AP Ra-anti-HuOPGL Ab) with OPGL neutralizing activity was used as a positive control for the anti-huOPGL antibody neutralizing activity assay.

TABLE 6

Sample	IC <sub>50</sub> (ng/mL)
AP Ra-anti-HuOPGL Ab	112
9H7	129
18B2	80
2D8	611
2E11	77
16E1	352
22B3	146
IC <sub>50</sub> (ng/mL) Running Average AP Ra-anti-HuOPGL Ab	
Average	140
Stdev	35.8
CV	26%
Count	25

## Example 7

## Pharmacokinetics in Cynomolgus Monkeys

**[0262]** The in vivo activity and pharmacokinetics of the anti-OPGL antibodies of the invention were assayed using cynomolgus monkeys. Three female cynomolgus monkeys, not greater than 5 years of age and weighing 2 to 5 kg each received single subcutaneous (SC) doses of 1 mg/kg anti-OPGL antibody.

**[0263]** Animals were dosed with anti-OPGL antibody expressed from transfected Chinese hamster ovary (CHO) cells and serum samples were taken for determination of anti-OPGL antibody levels, anti-therapeutic antibody analysis, and analysis of the bone turnover marker serum N-telopeptide (serum N-Tx), alkaline phosphatase (ALP), and serum calcium (serum Ca).

**[0264]** The serum concentration-time profiles following SC administration are shown in FIG. 21. The serum N-Tx concentration-time profiles following SC administration are shown in FIG. 22.

## Example 8

## Identification of an Epitope for Antibodies on OPGL

## Production of Variant Murine OPGL

**[0265]** Human OPGL [143-317] was produced as described in Example 1 of WO 01/62932, published Aug. 30, 2001, which is hereby incorporated by reference in its entirety. Murine OPGL [158-316] containing amino acid residues 158 through 316 of murine OPGL (as shown in FIG. 1 of International Application, Publication No. WO98/46751, incorporated by reference) preceded by an introduced N-terminal methionine residue was produced in *E. coli*. Murine OPGL [158-316] was purified from the soluble fraction of bacteria as described previously (Lacey et al., 1998, *Cell* 93:165-176). FLAG-tagged murine OPGL [158-316] was

produced by introducing a nucleic acid encoding an N-terminal methionine followed by a FLAG-tag sequence fused to the N-terminus of residues 158-316 as shown in FIG. 1 of International Application, Publication No. WO98/46751 using conventional genetic engineering techniques. The FLAG-tagged OPGL [158-316] molecule was cloned into bacterial expression vector pAMG21 (pAMG21 was deposited with the American Type Culture Collection and having Accession No. 98113).

**[0266]** A FLAG-tagged murine OPGL [158-316] polypeptide variant was constructed in which amino acid residues SVPTD (SEQ ID NO: 68) at positions 229-233 (as shown in FIG. 1 of International Application, Publication No. WO98/46751) were substituted with corresponding amino acid residues DLATE (SEQ ID NO: 69) at positions 230-234 (as shown in FIG. 4 of International Application, Publication No. WO98/46751). The resulting construct referred to as "FLAG-murine OPGL [158-316]/DE" has the nucleic acid and protein sequence as shown in FIG. 23 (SEQ ID NO: 72) (which shows only where the mutations are located). The amino acid sequence changes are located in a region of OPGL between the D and E regions. FIG. 23 shows a comparison of murine (SEQ ID NO: 70), human (SEQ ID NO: 71), and murine DE variant (SEQ ID NO: 72) amino acid sequences in this region. The sequence changes in the murine variant are S231D, V232L, P233A and D235E with the T at position 234 unchanged. Flanking sequences in this region are virtually identical between murine and human OPGL.

**[0267]** This molecule was constructed using a two-step PCR reaction where the first step contained two separate PCR reactions, designated reaction A and reaction B. For both reaction A and reaction B, pAMG21-FLAG-murine OPGL [158-316] DNA was used as a template for PCR. Reaction A employed oligonucleotides #2640-90 and #2640-91 for PCR, whereas reaction B employed oligonucleotides #2640-92 and #2640-93.

#2640-90 (SEQ ID NO: 73):  
CCTCTCATATGGACTACAAGGAC;

#2640-91 (SEQ ID NO 74):  
AGTAGCCAGGTCTCCCGATGTTTCATGATG;

#2640-92 (SEQ ID NO: 75):  
CTGGCTACTGAATATCTTCAGCTGATGGTG;

#2640-93 (SEQ ID NO: 76):  
CCTCTCCTCGAGTTAGTCTATGTCC.

**[0268]** Conditions for reactions A and B were: 95° C. for 1 min; 95° C. for 20 seconds, 44° C. for 30 seconds, 72° C. for 45 seconds for 5 cycles; 95° C. for 20 seconds, 60° C. for 30 seconds, 72° C. for 45 seconds for 25 cycles; and 72° C. for 10 minutes with Pfu Turbo polymerase (Stratagene) and the appropriate buffer and nucleotides. After thermocycling was performed, PCR products from reactions A and B were purified from an agarose gel using conventional methods. The second step PCR reaction, designated reaction C, utilized purified reaction A and reaction B PCR products as a template and oligonucleotides #2640-90 and #2640-93 as primers. Conditions for reaction C were: 95° C. for 1 minute; 95° C. for 20 seconds, 37° C. for 30 seconds, 72° C. for 1 minute for 25 cycles; and 72° C. for 10 minutes with Pfu Turbo polymerase and the appropriate buffer and nucleotides. Following thermocycling, the product from reaction C was cloned into the pCRII-TOPO cloning vector (Invitrogen) and electroporated

into DH10b (Gibco) cells using methods provided by the manufacturer. Clones were selected and sequenced to confirm the amino acid sequence SVPTD (SEQ ID NO: 68) in murine OPGL [158-316] was changed to DLATE (SEQ ID NO: 69). The sequence-verified DNA was then digested with NdeI and XhoI, purified, and subcloned into bacterial expression vector pAMG21 giving rise to plasmid pAMG21-FLAG-murine OPGL[158-316]/DE.

**[0269]** *E. coli* host GM94 (deposited with the American Type Culture Collection under Accession No. 202173) containing plasmid pAMG21-FLAG-murine OPGL[158-316]/DE was grown in 2XYT media to an exponential growth phase and induced to express the FLAG-tagged murine OPGL[158-316]/DE protein by addition of *V. fischeri* synthetic autoinducer to 100 ng/mL. Approximately 3-6 hours after induction, the cells were pelleted and recombinant FLAG-murine OPGL[158-316]/DE protein was purified from the soluble fraction of *E. coli* using methods described in Lacey et al., *ibid*.

Binding of Anti-Human OPGL Antibodies to Human OPGL [143-317], Murine OPGL[158-316], and FLAG-Murine OPGL[158-316]/DE

**[0270]** Costar E.I.A./R.I.A. Plates (Flat Bottom High Binding, Cat# 3590) were coated with 100  $\mu$ L/well of either human OPGL[143-317] protein, murine OPGL[158-316] protein, or FLAG-tagged murine OPGL[158-316]/DE protein at 3  $\mu$ g/mL in PBS, overnight at 4 $^{\circ}$  C. with agitation. After overnight incubation, the protein solutions were removed from the plate and 200  $\mu$ L of 5% Chicken Serum (Gibco/BRL Cat# 16110-082) in PBST (PBS plus 0.05% Tween 20) was added to each well of the plate and plates were incubated at room temperature (RT) for 3 hours with agitation. After incubation and blocking, plates were washed 4 times with 1 $\times$  K-P wash solution in dH<sub>2</sub>O (Cat#50-63-00, Kirkegaard & Perry Laboratories) and dried. Purified anti-OPGL antibody or human OPGL [22-194]-Fc protein was serially diluted 1:1 from 2  $\mu$ g/mL to 1.953 ng/mL in 5% Chicken Serum in PBST and 100  $\mu$ L/well was added to appropriate wells of the microtiter plate coated with either human OPGL[143-317], murine OPGL[158-316], or FLAG-tagged murine OPGL[158-316]/DE protein. Plates were incubated for 2.25 hours at room temperature with agitation, washed four times with 1 $\times$  K-P wash solution and dried. Goat anti-human IgG (Fc) (Jackson ImmunoResearch, Cat# 109-036-098) was diluted 1:3000 in 5% Chicken Serum in PBST and 100  $\mu$ L was added to each well. Plates were incubated for 1.25 hours at room tempera-

ture with agitation, washed six times with 1 $\times$  K-P wash solution, and dried. 100  $\mu$ L of undiluted ABTS substrate (Kirkegaard & Perry; Cat#50-66-00) was added to each well and the dish was incubated at room temperature until sufficient blue-green color developed. Color development was stopped by addition of 100  $\mu$ L 1% SDS. Quantitation of color development was performed using a microtiter plate reader with detection at 405 nm.

**[0271]** The results of the enzyme immunoassay are shown in FIGS. 24 and 25. All six anti-OPGL antibodies of the invention bind to human OPGL[143-317]. However, only 22B3 antibody shows detectable binding to murine OPGL [158-316] over the concentration range tested (FIG. 24). While binding of 22B3 antibody to murine OPGL[158-316] occurs with a much lower affinity than to human OPGL[143-317], the 2D8, 9H7, 16E1, and 22B3 antibodies bind to FLAG-tagged murine OPGL[158-316]/DE (FIG. 25) almost as well as to human OPGL[143-317] under the assay conditions above. Thus, the amino acid changes in murine OPGL [158-316]/DE compared to murine OPGL[158-316] are important to the binding activity of antibodies 2D8, 9H7, 16E1, and 22B3. Antibodies 2E11 and 18B2 show no detectable binding to either murine OPGL[158-316] or murine OPGL[158-316]/DE.

**[0272]** The FLAG-murine OPGL[158-316]/DE was assayed for activity in a RAW cell assay as described in Example 6 and observed to have a similar ED50 for osteoclast formation as human OPGL[143-317], indicating that the DE variant is active in promoting osteoclast activity in vitro. Therefore, the binding of the anti-OPGL antibodies to murine OPGL[158-316]/DE is likely to inhibit osteoclast formation.

**[0273]** The epitope of the 2D8, 9H7, 16E1, and 22B3 anti-human OPGL antibodies is located to a region of human OPGL that includes at least amino acid residues DLATE (SEQ ID NO: 103) (residues 230 through 234 of human OPGL as shown in FIG. 4 of International Application, Publication No. WO98/46751) termed the D-E loop. The 2E11 and 18B2 anti-human OPGL antibodies do not bind to peptide fragments corresponding to the D-E loop region by itself. However, it will be recognized that in the native molecule these antibodies may bind to an epitope outside the D-E loop region or they may bind to all or a portion of the D-E loop region in combination with other portions of the molecule.

**[0274]** It should be understood that the foregoing disclosure emphasizes certain specific embodiments of the invention and that all modifications or alternatives equivalent thereto are within the spirit and scope of the invention as set forth in the appended claims.

---

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 103

<210> SEQ ID NO 1

<211> LENGTH: 1066

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

gcctccacca agggcccatc ggtcttcccc ctggcaccct cctccaagag cacctctggg 60

ggcacagcgg ccctgggctg cctggtcaag gactacttcc ccgaaccggg gacggtgtcg 120

-continued

---

```

tggaaactcag gcgccctgac cagcggcgctg cacaccttcc cggctgtcct acagtccctca 180
ggactctact ccctcagcag cgtgggtgacc gtgccctcca gcagcttggg caccagacc 240
tacatctgca acgtgaatca caagcccagc aacaccaagg tggacaagaa agttgagccc 300
aaatcttgtg acaaaactca cacatgcccc ccgtgcccag cacctgaact cctgggggga 360
ccgtcagtct tcctcttccc cccaaaaccc aaggacaccc tcatgatctc ccggaccct 420
gaggtcacat gcgtgggtgt ggacgtgagc cacgaagacc ctgaggtcaa gttcaactgg 480
tacgtggagc gcgtggaggt gcataatgcc aagacaaagc cgcgggagga gcagtacaac 540
agcacgtacc gtgtggtcag cgtcctcacc gtccctgcacc aggactggct gaatggcaag 600
gagtacaagt gcaaggtctc caacaaagcc ctcccagccc ccatcgagaa aaccatctcc 660
aaagccaaag ggcagcccgc agaaccacag gtgtacaccc tgccccatc ccgggatgag 720
ctgaccaaga accaggtcag cctgacctgc ctgggtcaaag gcttctatcc cagcgacatc 780
gccgtggagt gggagagcaa tgggcagccg gagaacaact acaagaccac gcctcccgtg 840
ctggactcgc acggctcctt ctctctctat agcaagctca cctgggacaa gagcaggtgg 900
cagcagggga acgtcttctc atgctccgtg atgcatgagg ctctgcacaa ccaactacag 960
cagaagagcc tctccctgtc tccgggtaaa tgataagtgc acatgccctg aattctgcag 1020
atatccatca cactggcggc cgctcgagca tgcattctaga gggccc 1066

```

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 330

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 2

```

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1          5          10          15
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20        25        30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35        40        45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50        55        60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65        70        75        80
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
85        90        95
Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
100       105       110
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
115       120       125
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
130       135       140
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
145       150       155       160
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
165       170       175
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
180       185       190

```

-continued

---

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn  
 195 200 205

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly  
 210 215 220

Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu  
 225 230 235 240

Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr  
 245 250 255

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn  
 260 265 270

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe  
 275 280 285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn  
 290 295 300

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr  
 305 310 315 320

Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys  
 325 330

<210> SEQ ID NO 3  
 <211> LENGTH: 321  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

actgtggctg caccatctgt cttcatcttc cggccatctg atgagcagtt gaaatctgga 60

actgcctctg ttgtgtgcct gctgaataac ttctatccca gagaggccaa agtacagtgg 120

aaggtggata acgcctcca atcgggtaac tcccaggaga gtgtcacaga gcaggacagc 180

aaggacagca cctacagcct cagcagcacc ctgacgctga gcaaagcaga ctacgagaaa 240

cacaaagtct acgcctgcga agtcacccat cagggcctga gctcgcccgt cacaaagagc 300

ttcaacaggg gagagtgtta g 321

<210> SEQ ID NO 4  
 <211> LENGTH: 106  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln  
 1 5 10 15

Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr  
 20 25 30

Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser  
 35 40 45

Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr  
 50 55 60

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys  
 65 70 75 80

His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro  
 85 90 95

Val Thr Lys Ser Phe Asn Arg Gly Glu Cys  
 100 105

-continued

<210> SEQ ID NO 5  
 <211> LENGTH: 348  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

```

gaggtgcagc tgggtgcagtc tgggggaggc ttggtacatc ctggggggtc cctgagactc      60
tcctgtgaag gctctggatt caccttcagt agcaatggta tgcactgggt gcgccagact      120
ccaggaaaag gtctggagtg ggtatcaggt attggtactg ctggtggcac atactatgca      180
gactccgtga agggccgatt caccatttcc agagacaatg tcaagaagtc cttgtatctt      240
caaatgaaca gctgagagc cgaggacatg gctatttatt attgtgtaag aaaaaactgg      300
ggatggttcg acccctgggg ccagggagcc ctggtcaccg tctctagt                      348
  
```

<210> SEQ ID NO 6  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

```

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly
1           5           10           15
Ser Leu Arg Leu Ser Cys Glu Gly Ser Gly Phe Thr Phe Ser Ser Asn
20           25           30
Gly Met His Trp Val Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
35           40           45
Ser Gly Ile Gly Thr Ala Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys
50           55           60
Gly Arg Phe Thr Ile Ser Arg Asp Asn Val Lys Lys Ser Leu Tyr Leu
65           70           75           80
Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Ile Tyr Tyr Cys Val
85           90           95
Arg Lys Asn Trp Gly Trp Phe Asp Pro Trp Gly Gln Gly Ala Leu Val
100          105          110
Thr Val Ser Ser
115
  
```

<210> SEQ ID NO 7  
 <211> LENGTH: 327  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

```

gaaattgtgc tgaccagtc tccagccacc ctgtctttgt ctccagggga aagagccacc      60
ctctcctgca gggccagtc gagtgtaac agctacttag cctggttcca acagaaacct      120
ggccaggctc ccagactcct catctatgat gcatccaaca gggccactgg catcccagcc      180
aggttcagtg gcagtgggtc tgggacagac ttcactetca ccatcagcag ccttgagcct      240
gaagattttg caatttatta ctgtcagcag cgtagcaact ggcctccgtt cacttttggc      300
caggggacca agctggagat caaacga                                           327
  
```

<210> SEQ ID NO 8  
 <211> LENGTH: 109



-continued

---

```

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1           5           10          15
Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Asn Ser Tyr
          20          25          30
Leu Ala Trp Phe Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
          35          40          45
Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
          50          55          60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65          70          75          80
Glu Asp Phe Ala Ile Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Pro
          85          90          95
Phe Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg
          100          105

```

```

<210> SEQ ID NO 9
<211> LENGTH: 345
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

gagggtgcagc tgggtgcagtc ggggggagac ttggtacatc ctggggggtc cctgagactc   60
tctctgtgtcg gctctggatt caccttcagt cactatcctt tgcactgggt tcgccaggct   120
ccaggaaaag gtctggagtg gatatcaggt attcactactg gtggtgccac atactataca   180
gactccgtga agggccgggt caccatctcc agcgacaatg ccaagaactc cttatatctt   240
caaatgaaca ccctgagagc cgaggacatg gctgtgtatt actgtgcaag agggcgaaac   300
tcctttgact actggggcca gggaaccctg gtcacgtctc ctagt                          345

```

```

<210> SEQ ID NO 10
<211> LENGTH: 115
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

Glu Val Gln Leu Val Gln Ser Gly Gly Asp Leu Val His Pro Gly Gly
1           5           10          15
Ser Leu Arg Leu Ser Cys Val Gly Ser Gly Phe Thr Phe Ser His Tyr
          20          25          30
Pro Leu His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile
          35          40          45
Ser Gly Ile His Thr Gly Gly Gly Thr Tyr Tyr Thr Asp Ser Val Lys
          50          55          60
Gly Arg Phe Thr Ile Ser Ser Asp Asn Ala Lys Asn Ser Leu Tyr Leu
65          70          75          80
Gln Met Asn Thr Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala
          85          90          95
Arg Gly Arg Asn Ser Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Ile
          100          105          110
Val Ser Ser

```

-continued

115

<210> SEQ ID NO 11  
 <211> LENGTH: 324  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

```

gacatccaga tgaccagtc tccatcctca ctgtctgcat ctgtaggaga cagagtcacc      60
atcacttgtc gggcgagtc ggggtattagc agctgggtag cctgggatca gcagaaacca    120
gagaaagccc ctaagtcct gatctatgct gcatccagtt tgcaaagtgg ggtcccatca    180
aggttcagcg gcagtgatc tgggacagat ttcactctca ccatcagcag cctgcagtct    240
gaagattttg caacttatta ctgccaacag tataatagtt accctcccac cttgggccaa    300
gggacacgac tggagattaa acga                                           324

```

<210> SEQ ID NO 12  
 <211> LENGTH: 108  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1           5           10          15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp
                20          25          30
Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys Ala Pro Lys Ser Leu Ile
                35          40          45
Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
                50          55          60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Ser
65          70          75          80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Tyr Pro Pro
                85          90          95
Thr Phe Gly Gln Gly Thr Arg Leu Glu Ile Lys Arg
                100         105

```

<210> SEQ ID NO 13  
 <211> LENGTH: 348  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

```

gaggtgcagc tgggtgcagtc tgggggagc ttggtacatc ctggggggtc cctgagactc      60
tctgtgagc gctctggatt caccttcagt agctatggga tgcactgggt tcgccagget    120
ccaggaaaag gtctggagtg ggtatcaggt attggtactg gtggtggcac atactatgca    180
gactccgtga agggccgatt caccatctcc agagacaatg tcaagaactc cttgtatctt    240
caaatgaaca gcctgagagc cgaggacatg gctgtgtatt actgtgcaag aaaaaactgg    300
ggatggtttg actactgggg ccaggaacc ctggtcaccg tctctagt                    348

```

<210> SEQ ID NO 14  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

-continued

&lt;400&gt; SEQUENCE: 14

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly  
 1 5 10 15  
 Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Ser Tyr  
 20 25 30  
 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45  
 Ser Gly Ile Gly Thr Gly Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys  
 50 55 60  
 Gly Arg Phe Thr Ile Ser Arg Asp Asn Val Lys Asn Ser Leu Tyr Leu  
 65 70 75 80  
 Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala  
 85 90 95  
 Arg Lys Asn Trp Gly Trp Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val  
 100 105 110  
 Thr Val Ser Ser  
 115

&lt;210&gt; SEQ ID NO 15

&lt;211&gt; LENGTH: 327

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 15

gaaattgtgc tgaccagtc tccagccacc ctgtctttgt ctccagggga aagagccacc 60  
 ctctcctgca gggccagtca gagtattagc agtacttag cctggtagca acagaaacct 120  
 ggccaggctc ccaggctcct catctatgat gcattccaaca gggccactgg catcccagcc 180  
 aggttcagtg gcagtgggtc tgggacagac ttcactctca ccatcagcag cctagagcct 240  
 gaagattttg cagtttatta ctgtcagcag cgtagcaaat ggctccgta cacttttggc 300  
 caggggacca aactcgagat caaacga 327

&lt;210&gt; SEQ ID NO 16

&lt;211&gt; LENGTH: 109

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 16

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly  
 1 5 10 15  
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Ile Ser Ser Tyr  
 20 25 30  
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile  
 35 40 45  
 Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly  
 50 55 60  
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro  
 65 70 75 80  
 Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Lys Trp Pro Pro  
 85 90 95  
 Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg  
 100 105

-continued

---

```

<210> SEQ ID NO 17
<211> LENGTH: 345
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

gaggtgcagc tgggtgcagtc tgggggaggc ttggtacatc ctgggggggc cctgagactc   60
tcctgtgtag gctctagatt caccttcagt gcctatccta tgcactgggt tcgccaggct   120
ccaggaaaag gtctggagtg ggtatcaggt attggttctg gtggtggcac aaactatgca   180
gactccgtga agggccgatt caccatctcc agagacactg ccaagaactc cttgtatctt   240
caaatgaaca gcttgagagc cgaggacatg gctgtgtatt actgtgcaag agggaggaat   300
tcttttgact actggggcca gggaaccctg gtcaccgtct ctagt                       345

```

```

<210> SEQ ID NO 18
<211> LENGTH: 115
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly
1      5      10      15
Ser Leu Arg Leu Ser Cys Val Gly Ser Arg Phe Thr Phe Ser Ala Tyr
20     25     30
Pro Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35     40     45
Ser Gly Ile Gly Ser Gly Gly Thr Asn Tyr Ala Asp Ser Val Lys
50     55     60
Gly Arg Phe Thr Ile Ser Arg Asp Thr Ala Lys Asn Ser Leu Tyr Leu
65     70     75     80
Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala
85     90     95
Arg Gly Arg Asn Ser Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr
100    105    110

Val Ser Ser
115

```

```

<210> SEQ ID NO 19
<211> LENGTH: 324
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

gacatccaga tgaccagtc tccatcctca ctgtctgcat ctgtaggaga cagagtcacc   60
atcacttgtc gggcgagtca gggattagc acctgggttag cctgggatca gcagaaacca   120
gagaaagccc ctaagtccct gatctatgct gcattccagtt tgcagagtgg ggtcccatcg   180
aggttcagcg gcagtggatc tgggacagat ttcactctca ccattcagcag cctgcagcct   240
gaagattttg caacttatta ctgccaacag tataatagtt acctccgac gttcggccaa   300
gggaccaagg tggagatcaa acga                                           324

```

```

<210> SEQ ID NO 20
<211> LENGTH: 108
<212> TYPE: PRT

```

-continued

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 20

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1      5      10      15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Thr Trp
20     25     30
Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys Ala Pro Lys Ser Leu Ile
35     40     45
Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50     55     60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65     70     75     80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Tyr Pro Pro
85     90     95
Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
100    105

```

&lt;210&gt; SEQ ID NO 21

&lt;211&gt; LENGTH: 348

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 21

```

gaggtccagc tgggtgcagtc tgggggaggc ttggtacatc ctggggggtc cctgagactc      60
tcctgtgcag gctctggatt caccttcagt ggccatgctt tgcactgggt tcgccaggct      120
ccaggaaaag gtctggagtg ggtatcaggt attggtactc atggtgggac atactatgca      180
gactccgtga agggccgatt caccatctcc agagacaatg ccaagaactc cttgtttctt      240
caaatgaaca gcttgagcgc cgaggacatg gctgtgtatt actgtacaag aagaaactgg      300
ggacaatttg actactgggg ccaggaacc ctggtcaccg tctctagt      348

```

&lt;210&gt; SEQ ID NO 22

&lt;211&gt; LENGTH: 116

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 22

```

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly
1      5      10      15
Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Gly His
20     25     30
Ala Leu His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35     40     45
Ser Gly Ile Gly Thr His Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys
50     55     60
Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Phe Leu
65     70     75     80
Gln Met Asn Ser Leu Ser Ala Glu Asp Met Ala Val Tyr Tyr Cys Thr
85     90     95
Arg Arg Asn Trp Gly Gln Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val
100    105    110
Thr Val Ser Ser
115

```

-continued

<210> SEQ ID NO 23  
 <211> LENGTH: 327  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 23

```

gaaattgtgc tgactcagtc tccagccacc ctgtctttgt ctccagggga aagagccacc   60
ctctcctgca gggccagtca gagtgtagc agctacttag cctggtacca acagaaacct   120
ggccaggctc ccaggctcct catctatgat gcatccaaca gggccactgg catcccagcc   180
aggttcagtg gcagtgggtc tgggacagac ttcactctca ccatcagcag cctagagcct   240
gaagattttg cagtttatta ctgtcagcag cgtagcaact ggctccgta cacttttggc   300
caggggacca agctggagat caaacga                                     327

```

<210> SEQ ID NO 24  
 <211> LENGTH: 109  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

```

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1           5           10          15
Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
                20          25          30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
          35          40          45
Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
          50          55          60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65          70          75          80
Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Pro
          85          90          95
Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg
          100         105

```

<210> SEQ ID NO 25  
 <211> LENGTH: 348  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

```

gaggtgcagc tgggtcagtc tgggggaggc ttggtacatc ctggggggtc cctgagactc   60
tctctgtaag gctctggatt caccttcagt agcaatggta tgcaactggg gcgccagact   120
ccaggaaaag gtctggagtg ggtatcaggt attggtactg ctggtggcac atactatgca   180
gactccgtga agggccgatt caccatttcc agagacaatg tcaagaagtc cttgtatctt   240
caaatgaaca gcctgagagc cgaggacatg gctatttatt attgtgtaag aaaagactgg   300
ggatggttcg acccctgggg ccagggagcc ctggtcaccg tctctagt                                     348

```

<210> SEQ ID NO 26  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 26

```

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly
1          5          10          15
Ser Leu Arg Leu Ser Cys Glu Gly Ser Gly Phe Thr Phe Ser Ser Asn
20          25          30
Gly Met His Trp Val Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
35          40          45
Ser Gly Ile Gly Thr Ala Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys
50          55          60
Gly Arg Phe Thr Ile Ser Arg Asp Asn Val Lys Lys Ser Leu Tyr Leu
65          70          75          80
Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Ile Tyr Tyr Cys Val
85          90          95
Arg Lys Asp Trp Gly Trp Phe Asp Pro Trp Gly Gln Gly Ala Leu Val
100         105         110
Thr Val Ser Ser
115
    
```

<210> SEQ ID NO 27

<211> LENGTH: 327

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

```

gaaattgtgc tgaccagtc tccagccacc ctgtctttgt ctccagggga aagagccacc    60
ctctcctgca gggccagtca gagtattagc agctacttag cctggtagca acagaaacct    120
ggccaggctc ccaggctcct catctatgat gcatccaaca gggccactgg catcccagcc    180
aggttcagtg gcagtgggtc tgggacagac ttcactctca ccatcagcag cctagagcct    240
gaagattttg cagtttatta ctgtcagcag cgtagcaaat ggcctccgta cacttttggc    300
caggggacca agctggagat caaacga                                     327
    
```

<210> SEQ ID NO 28

<211> LENGTH: 109

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28

```

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1          5          10          15
Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Ile Ser Ser Tyr
20          25          30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
35          40          45
Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
50          55          60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65          70          75          80
Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Lys Trp Pro Pro
85          90          95
Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg
100         105
    
```

-continued

---

<210> SEQ ID NO 29  
 <211> LENGTH: 1413  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 29

```

gagggtgcagc tgggtgcagtc tgggggaggc ttggtacatc ctggggggtc cctgagactc   60
tcctgtgaag gctctggatt caccttcagt agcaatgta tgcactgggt gcgccagact   120
ccaggaaaag gtctggagtg ggtatcaggt attggtactg ctggtggcac atactatgca   180
gactccgtga agggccgatt caccatttcc agagacaatg tcaagaagtc cttgtatctt   240
caaatgaaca gcctgagagc cgaggacatg gctatttatt attgtgtaag aaaaaactgg   300
ggatggttcg acccctgggg ccaggggacc ctggtcaccg tctctagtgc ctccaccaag   360
ggcccatcgg tcttccccct ggcaccctcc tccaagagca cctctggggg cacagcggcc   420
tgggctgcct ggtcaaggac tacttccccg aaccgggtgac ggtgtcgtgg aactcaggcg   480
ccctgaccag cggcgtgcac accttccccg ctgtcctaca gtctcagga ctctactccc   540
tcagcagcgt ggtgaccgtg cctccagca gcttggggcac ccagacctac atctgcaacg   600
tgaatcacia gcccagcaac accaaggtgg acaagaaagt tgagcccaaa tcttgtgaca   660
aaactcacac atgcccaccg tgcccagcac ctgaactcct ggggggaccg tcagtcttcc   720
tcttcccccc aaaaccaag gacaccctca tgatctcccg gaccctgag gtcacatgag   780
tgggtgtgga cgtgagccac gaagaccctg aggtcaagtt caactggtac gtggacggcg   840
tggaggtgca taatgccaa gcaaaagccgc gggaggagca gtacaacagc acgtaccgtg   900
tggtcagcgt cctcaccgtc ctgaccagg actggctgaa tggcaaggag tacaagtgca   960
aggtctccaa caaagccctc ccagccccc tgcagaaaac catctccaaa gccaaaaggc  1020
agccccgaga accacaggtg tacaccctgc cccatccccg ggatgagctg accaagaacc  1080
aggtcagcct gacctgctg gtcaaaggct tctatcccag cgacatgcc gtggagtggg  1140
agagcaatgg gcagccggag aacaactaca agaccacgcc tcccgtgctg gactccgacg  1200
gtcctctctt cctctatagc aagctcaccg tggacaagag caggtggcag caggggaacg  1260
tcttctcatg ctccgtgatg catgaggctc tgcacaacca ctacacgag aagagcctct  1320
cctgtctccc gggtaaatga taagtgcaca tgcctgaaat tctgcagata tccatcacac  1380
tggcggccgc tcgagcatgc atctagaggg ccc                               1413

```

<210> SEQ ID NO 30  
 <211> LENGTH: 446  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30

```

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly
1           5           10           15
Ser Leu Arg Leu Ser Cys Glu Gly Ser Gly Phe Thr Phe Ser Ser Asn
20          25          30
Gly Met His Trp Val Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
35          40          45
Ser Gly Ile Gly Thr Ala Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys
50          55          60
Gly Arg Phe Thr Ile Ser Arg Asp Asn Val Lys Lys Ser Leu Tyr Leu

```



-continued

65	70	75	80
Gln Met Asn Ser	Leu Arg Ala Glu Asp	Met Ala Ile Tyr Tyr	Cys Val
	85	90	95
Arg Lys Asn Trp	Gly Trp Phe Asp	Pro Trp Gly Gln Gly Ala	Leu Val
	100	105	110
Thr Val Ser Ser	Ala Ser Thr Lys Gly	Pro Ser Val Phe Pro	Leu Ala
	115	120	125
Pro Ser Ser Lys	Ser Thr Ser Gly Gly	Thr Ala Ala Leu Gly	Cys Leu
	130	135	140
Val Lys Asp Tyr	Phe Pro Glu Pro Val Thr	Val Ser Trp Asn Ser	Gly
	145	150	160
Ala Leu Thr Ser	Gly Val His Thr Phe	Pro Ala Val Leu Gln	Ser Ser
	165	170	175
Gly Leu Tyr Ser	Leu Ser Ser Val Val	Thr Val Pro Ser Ser	Ser Leu
	180	185	190
Gly Thr Gln Thr	Tyr Ile Cys Asn Val	Asn His Lys Pro Ser	Asn Thr
	195	200	205
Lys Val Asp Lys	Lys Val Glu Pro Lys	Ser Cys Asp Lys Thr	His Thr
	210	215	220
Cys Pro Pro Cys	Pro Ala Pro Glu Leu Leu	Gly Gly Pro Ser Val	Phe
	225	230	240
Leu Phe Pro Pro	Lys Pro Lys Asp Thr	Leu Met Ile Ser Arg	Thr Pro
	245	250	255
Glu Val Thr Cys	Val Val Val Asp Val	Ser His Glu Asp Pro	Glu Val
	260	265	270
Lys Phe Asn Trp	Tyr Val Asp Gly Val	Glu Val His Asn Ala	Lys Thr
	275	280	285
Lys Pro Arg Glu	Glu Gln Tyr Asn Ser	Thr Tyr Arg Val Val	Ser Val
	290	295	300
Leu Thr Val Leu	His Gln Asp Trp Leu	Asn Gly Lys Glu Tyr	Lys Cys
	305	310	320
Lys Val Ser Asn	Lys Ala Leu Pro Ala	Pro Ile Glu Lys Thr	Ile Ser
	325	330	335
Lys Ala Lys Gly	Gln Pro Arg Glu Pro	Gln Val Tyr Thr Leu	Pro Pro
	340	345	350
Ser Arg Asp Glu	Leu Thr Lys Asn Gln	Val Ser Leu Thr Cys	Leu Val
	355	360	365
Lys Gly Phe Tyr	Pro Ser Asp Ile Ala	Val Glu Trp Glu Ser	Asn Gly
	370	375	380
Gln Pro Glu Asn	Asn Tyr Lys Thr Thr	Pro Pro Val Leu Asp	Ser Asp
	385	390	400
Gly Ser Phe Phe	Leu Tyr Ser Lys Leu	Thr Val Asp Lys Ser	Arg Trp
	405	410	415
Gln Gln Gly Asn	Val Phe Ser Cys Ser	Val Met His Glu Ala	Leu His
	420	425	430
Asn His Tyr Thr	Gln Lys Ser Leu Ser	Leu Ser Pro Gly Lys	
	435	440	445

&lt;210&gt; SEQ ID NO 31

&lt;211&gt; LENGTH: 648

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapiens

-continued

&lt;400&gt; SEQUENCE: 31

```

gaaattgtgc tgaccagctc tccagccacc ctgtctttgt ctccagggga aagagccacc    60
ctctcctgca gggccagtca gagggttaac agctacttag cctggttcca acagaaacct    120
ggccaggctc ccagactcct catctatgat gcatccaaca gggccactgg catcccagcc    180
aggttcagtg gcagtgggtc tgggacagac ttcactctca ccatcagcag ccttgagcct    240
gaagattttg caatttatta ctgtcagcag cgtagcaact ggccctccgtt cacttttggc    300
caggggacca agctggagat caaacgaact gtggctgcac catctgtcct catcttcccg    360
ccatctgatg agcagttgaa atctggaact gcctctgttg tgtgctgct gaataacttc    420
tatcccagag aggccaaagt acagtggaag gtggataacg ccctccaatc gggtaactcc    480
caggagagtg tcacagagca ggacagcaag gacagcacct acagcctcag cagcacctcg    540
acgctgagca aagcagacta cgagaaacac aaagtctacg cctgcgaagt cacccatcag    600
ggcctgagct cgcccgtcac aaagagcttc aacaggggag agtgtagg                    648

```

&lt;210&gt; SEQ ID NO 32

&lt;211&gt; LENGTH: 215

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 32

```

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1             5             10            15
Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Asn Ser Tyr
          20            25            30
Leu Ala Trp Phe Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
          35            40            45
Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
          50            55            60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
          65            70            75            80
Glu Asp Phe Ala Ile Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Pro
          85            90            95
Phe Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala
          100           105           110
Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser
          115           120           125
Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu
          130           135           140
Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser
          145           150           155           160
Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu
          165           170           175
Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val
          180           185           190
Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys
          195           200           205
Ser Phe Asn Arg Gly Glu Cys
          210           215

```

-continued

<210> SEQ ID NO 33  
 <211> LENGTH: 1410  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 33

```

gagggtgcagc tgggtgcagtc ggggggagac ttggtacatc ctggggggtc cctgagactc   60
tcctgtgtcg gctctggatt caccttcagt cactatcctt tgcactgggt tcgccaggct   120
ccaggaaaag gtctggagtg gatatcaggt attcatactg gtggtggcac atactataca   180
gactccgtga agggccgggt caccatctcc agcgacaatg ccaagaactc cttatatctt   240
caaatgaaca ccctgagagc cgaggacatg gctgtgtatt actgtgcaag agggcgaaaac   300
tcctttgact actggggcca gggaaacctg gtcacgtctc ctagtgcctc caccaagggc   360
ccatcggtct tccccctggc accctcctcc aagagcacct ctggggggcac ageggcctgg   420
gctgcctggt caaggactac ttccccgaac cggtgacggt gtcgtggaac tcaggcgccc   480
tgaccagcgg cgtgcacacc ttccccggtg tcctacagtc ctcaggactc tactccctca   540
gcagcgtggt gaccctgccc tccagcagct tgggcaccca gacctacatc tgcaactgta   600
atcacaagcc cagcaacacc aaggtggaca agaaagtga gcccaaatct tgtgacaaaa   660
ctcacacatg cccaccgtgc ccagcacctg aactcctggg gggaccgtca gtcttcctct   720
tcccccaaaa acccaaggac accctcatga tctcccgac ccctgaggtc acatgcgtgg   780
tggtggagct gagccacgaa gaccctgagg tcaagttcaa ctggtacgtg gacggcgtgg   840
aggtgcataa tgccaagaca aagccgcggg aggagcagta caacagcacg tacctgtgtg   900
tcagcgtcct caccgtcctg caccaggact ggctgaatgg caaggagtac aagtgcaagg   960
tctccaacaa agccctccca gccccatcg agaaaacat ctccaagcc aaagggcagc  1020
cccgagaacc acaggtgtac accctgcccc catcccggga tgagctgacc aagaaccagg  1080
tcagcctgac ctgcctggtc aaaggcttct atcccagcga catgcctgtg gagggggaga  1140
gcaatgggca gccggagaac aactacaaga ccacgcctcc cgtgctggac tccgacggct  1200
ccttcttctc ctatagcaag ctcaccgtgg acaagagcag gtggcagcag gggaacgtct  1260
tctcatgctc cgtgatgcat gaggtctctg acaaccacta cacgcagaag agcctctccc  1320
tgtctccggg taaatgataa gtcgacatgc cctgaattct gcagatatcc atcacactgg  1380
cggccgctcg agcatgcatc tagagggccc                                     1410

```

<210> SEQ ID NO 34  
 <211> LENGTH: 445  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 34

```

Glu Val Gln Leu Val Gln Ser Gly Gly Asp Leu Val His Pro Gly Gly
1           5           10          15
Ser Leu Arg Leu Ser Cys Val Gly Ser Gly Phe Thr Phe Ser His Tyr
20          25          30
Pro Leu His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile
35          40          45
Ser Gly Ile His Thr Gly Gly Gly Thr Tyr Tyr Thr Asp Ser Val Lys
50          55          60
Gly Arg Phe Thr Ile Ser Ser Asp Asn Ala Lys Asn Ser Leu Tyr Leu

```

-continued

65	70	75	80
Gln Met Asn Thr	Leu Arg Ala Glu Asp Met Ala Val Tyr Tyr Cys Ala		
	85	90	95
Arg Gly Arg Asn Ser Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Ile			
	100	105	110
Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro			
	115	120	125
Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val			
	130	135	140
Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala			
	145	150	155
Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly			
	165	170	175
Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly			
	180	185	190
Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys			
	195	200	205
Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys			
	210	215	220
Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu			
	225	230	235
Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu			
	245	250	255
Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys			
	260	265	270
Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys			
	275	280	285
Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu			
	290	295	300
Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys			
	305	310	315
Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys			
	325	330	335
Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser			
	340	345	350
Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys			
	355	360	365
Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln			
	370	375	380
Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly			
	385	390	395
Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln			
	405	410	415
Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn			
	420	425	430
His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys			
	435	440	445

&lt;210&gt; SEQ ID NO 35

&lt;211&gt; LENGTH: 645

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapiens

-continued

&lt;400&gt; SEQUENCE: 35

```

gacatccaga tgaccagctc tccatcctca ctgtctgcat ctgtaggaga cagagtcacc    60
atcacttgtc gggcgagtcg gggattagc agctgggttag cctgggatca gcagaaacca    120
gagaaagccc ctaagtccct gatctatgct gcataccagtt tgcaaagtgg ggtcccatca    180
aggttcagcg gcagtggtgc tgggacagat ttcactctca ccatcagcag cctgcagtct    240
gaagattttg caacttatta ctgccaacag tataatagtt accctcccac cttcggccaa    300
gggacacgac tggagattaa acgaactgtg gctgcaccat ctgtcttcat cttcccgcga    360
tctgatgagc agttgaaatc tggaactgcc tctgttgtgt gcctgctgaa taacttctat    420
cccagagagg ccaaagtaca gtggaaggtg gataacgccc tccaatcggg taactcccag    480
gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg    540
ctgagcaaa gacactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc    600
ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gttag                      645

```

&lt;210&gt; SEQ ID NO 36

&lt;211&gt; LENGTH: 214

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 36

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1           5           10          15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp
                20           25           30
Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys Ala Pro Lys Ser Leu Ile
                35           40           45
Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50           55           60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Ser
65           70           75           80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Tyr Pro Pro
                85           90           95
Thr Phe Gly Gln Gly Thr Arg Leu Glu Ile Lys Arg Thr Val Ala Ala
100          105          110
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
115          120          125
Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
130          135          140
Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
145          150          155          160
Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
165          170          175
Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
180          185          190
Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
195          200          205
Phe Asn Arg Gly Glu Cys
210

```

-continued

---

```

<210> SEQ ID NO 37
<211> LENGTH: 1413
<212> TYPE: DNA
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 37
gagggtgcagc tgggtgcagtc tgggggagggc ttggtacatc ctgggggggc cctgagactc    60
tcctgtgcag gctctggatt caccttcagt agctatggga tgcactgggt tcgccaggct    120
ccaggaaaag gtctggagtg ggtatcaggt attggtactg gtggtggcac atactatgca    180
gactccgtga agggccgatt caccatctcc agagacaatg tcaagaactc cttgtatctt    240
caaatgaaca gcctgagagc cgaggacatg gctgtgtatt actgtgcaag aaaaaactgg    300
ggatggtttg actactgggg ccaggaacc ctggtcaccg tctctagtgc ctccaacaag    360
ggcccatcgg tcttccccct ggcaccctcc tccaagagca cctctggggg cacagcggcc    420
tgggctgctt ggtcaaggac tacttccccg aaccgggtgac ggtgtcgtgg aactcaggcg    480
ccctgaccag cggcgtgcac accttccccg ctgtcctaca gtctcagga ctctaactcc    540
tcagcagcgt ggtgaccgtg cctccagca gcttggggcac ccagacctac atctgcaacg    600
tgaatcacia gcccagcaac accaaggtgg acaagaaagt tgagccaaa tcttgtgaca    660
aaactcacac atgcccaccg tgcccagcac ctgaactcct ggggggaccg tcagtcttcc    720
tcttcccccc aaaaccaag gacaccctca tgatctcccg gaccctgag gtcacatgag    780
tgggtgtgga cgtgagccac gaagaccctg aggtcaagtt caactggtac gtggacggcg    840
tggaggtgca taatgccaa gcaaaagccgc gggaggagca gtacaacagc acgtaccgtg    900
tggtcagcgt cctcaccgtc ctgaccagg actggctgaa tggcaaggag tacaagtgca    960
aggtctccaa caaagccctc ccagccccc tgcagaaaac catctccaaa gccaaaaggc    1020
agccccgaga accacaggtg tacaccctgc cccatcccc ggatgagctg accaagaacc    1080
aggtcagcct gacctgctg gtcaaaggct tctatcccag cgacatgcc gtggagtggg    1140
agagcaatgg gcagccggag aacaactaca agaccacgcc tcccgtgctg gactccgacg    1200
gtcctctctt cctctatagc aagctcaccg tggacaagag caggtggcag caggggaaacg    1260
tcttctcatg ctccgtgatg catgaggctc tgcacaacca ctacacgagc aagagcctct    1320
cctgtctccc gggtaaatga taagtgcaca tgcctgaaat tctgcagata tccatcacac    1380
tggcggccgc tcgagcatgc atctagaggg ccc    1413

```

```

<210> SEQ ID NO 38
<211> LENGTH: 446
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 38
Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly
1           5           10          15
Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Ser Tyr
20          25          30
Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35          40          45
Ser Gly Ile Gly Thr Gly Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys
50          55          60
Gly Arg Phe Thr Ile Ser Arg Asp Asn Val Lys Asn Ser Leu Tyr Leu

```

-continued

65	70	75	80
Gln Met Asn Ser	Leu Arg Ala Glu Asp	Met Ala Val Tyr Tyr	Cys Ala
	85	90	95
Arg Lys Asn Trp	Gly Trp Phe Asp Tyr Trp	Gly Gln Gly Thr	Leu Val
	100	105	110
Thr Val Ser Ser	Ala Ser Thr Lys Gly Pro Ser	Val Phe Pro	Leu Ala
	115	120	125
Pro Ser Ser Lys	Ser Thr Ser Gly Gly Thr	Ala Ala Leu Gly	Cys Leu
	130	135	140
Val Lys Asp Tyr	Phe Pro Glu Pro Val Thr	Val Ser Trp Asn	Ser Gly
	145	150	155
Ala Leu Thr Ser	Gly Val His Thr Phe Pro	Ala Val Leu Gln	Ser Ser
	165	170	175
Gly Leu Tyr Ser	Leu Ser Ser Val Val Thr	Val Pro Ser Ser	Ser Leu
	180	185	190
Gly Thr Gln Thr	Tyr Ile Cys Asn Val Asn	His Lys Pro Ser	Asn Thr
	195	200	205
Lys Val Asp Lys	Lys Val Glu Pro Lys Ser	Cys Asp Lys Thr	His Thr
	210	215	220
Cys Pro Pro Cys	Pro Ala Pro Glu Leu Leu	Gly Gly Pro Ser	Val Phe
	225	230	235
Leu Phe Pro Pro	Lys Pro Lys Asp Thr Leu	Met Ile Ser Arg	Thr Pro
	245	250	255
Glu Val Thr Cys	Val Val Val Asp Val Ser	His Glu Asp Pro	Glu Val
	260	265	270
Lys Phe Asn Trp	Tyr Val Asp Gly Val Glu	Val His Asn Ala	Lys Thr
	275	280	285
Lys Pro Arg Glu	Glu Gln Tyr Asn Ser Thr	Tyr Arg Val Val	Ser Val
	290	295	300
Leu Thr Val Leu	His Gln Asp Trp Leu Asn	Gly Lys Glu Tyr	Lys Cys
	305	310	315
Lys Val Ser Asn	Lys Ala Leu Pro Ala Pro	Ile Glu Lys Thr	Ile Ser
	325	330	335
Lys Ala Lys Gly	Gln Pro Arg Glu Pro Gln	Val Tyr Thr Leu	Pro Pro
	340	345	350
Ser Arg Asp Glu	Leu Thr Lys Asn Gln Val	Ser Leu Thr Cys	Leu Val
	355	360	365
Lys Gly Phe Tyr	Pro Ser Asp Ile Ala Val	Glu Trp Glu Ser	Asn Gly
	370	375	380
Gln Pro Glu Asn	Asn Tyr Lys Thr Thr Pro	Pro Val Leu Asp	Ser Asp
	385	390	395
Gly Ser Phe Phe	Leu Tyr Ser Lys Leu Thr	Val Asp Lys Ser	Arg Trp
	405	410	415
Gln Gln Gly Asn	Val Phe Ser Cys Ser Val	Met His Glu Ala	Leu His
	420	425	430
Asn His Tyr Thr	Gln Lys Ser Leu Ser Leu	Ser Pro Gly Lys	
	435	440	445

&lt;210&gt; SEQ ID NO 39

&lt;211&gt; LENGTH: 648

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapiens

-continued

&lt;400&gt; SEQUENCE: 39

```

gaaattgtgc tgaccagctc tccagccacc ctgtctttgt ctccagggga aagagccacc    60
ctctcctgca gggccagtca gactattagc agctacttag cctggtagca acagaaacct    120
ggccaggctc ccaggctcct catctatgat gcatccaaca gggccactgg catcccagcc    180
aggttcagtg gcagtgggctc tgggacagac ttcactctca ccatcagcag cctagagcct    240
gaagattttg cagtttatta ctgtcagcag cgtagcaaat ggccctccgta cacttttggc    300
caggggacca aactcagatg caaacgaact gtggctgcac catctgtcct catcttcccg    360
ccatctgatg agcagttgaa atcttggaact gcctctgttg tgtgctgct gaataacttc    420
tatcccagag aggccaaagt acagtggaag gtggataacg ccctccaatc gggtaactcc    480
caggagagtg tcacagagca ggacagcaag gacagcacct acagcctcag cagcacctcg    540
acgctgagca aagcagacta cgagaaacac aaagtctacg cctgcgaagt cacccatcag    600
ggcctgagct cgcccgtcac aaagagcttc aacaggggag agtgtagg                    648

```

&lt;210&gt; SEQ ID NO 40

&lt;211&gt; LENGTH: 215

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 40

```

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1             5             10            15
Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Ile Ser Ser Tyr
          20            25            30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
      35            40            45
Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
      50            55            60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
      65            70            75            80
Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Lys Trp Pro Pro
          85            90            95
Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala
      100           105           110
Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser
      115           120           125
Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu
      130           135           140
Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser
      145           150           155           160
Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu
          165           170           175
Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val
      180           185           190
Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys
          195           200           205
Ser Phe Asn Arg Gly Glu Cys
      210           215

```



-continued

<210> SEQ ID NO 41  
 <211> LENGTH: 1410  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 41

```

gagggtgcagc tgggtgcagtc tgggggaggc ttggtacatc ctggggggtc cctgagactc   60
tcctgtgtag gctctagatt caccttcagt gcctatccta tgcactgggt tcgccaggct   120
ccaggaaaag gtctggagtg ggtatcaggt attggttctg gtggtggcac aaactatgca   180
gactccgtga agggccgatt caccatctcc agagacactg ccaagaactc cttgtatctt   240
caaatgaaca gcctgagagc cgaggacatg gctgtgtatt actgtgcaag agggagggaat   300
tcttttgact actggggcca gggaaacctg gtcacctctc ctagtgcctc caccaagggc   360
ccatcggtct tccccctggc accctcctcc aagagcacct ctggggggcac ageggcctgg   420
gctgcctggt caaggactac tccccgaac cggtgacggt gtcgtggaac tcaggcgccc   480
tgaccagcgg cgtgcacacc tccccggtg tcctacagtc ctcaggactc tactccctca   540
gcagcgtggt gaccctgccc tccagcagct tgggcaccca gacctacatc tgcaacgtga   600
atcacaagcc cagcaacacc aaggtggaca agaaagtga gcccaaatct tgtgacaaaa   660
ctcacacatg cccaccgtgc ccagcacctg aactcctggg gggaccgtca gtcttcctct   720
tcccccaaa acccaaggac accctcatga tctcccgac ccctgaggtc acatgcgtgg   780
tgggtggcgt gagccacgaa gaccctgagg tcaagttcaa ctggtacgtg gacggcgtgg   840
aggtgcataa tgccaagaca aagccgcggg aggagcagta caacagcacg tacctgtgtg   900
tcagcgtcct caccgtcctg caccaggact ggctgaatgg caaggagtac aagtgcaagg   960
tctccaacaa agccctccca gccccatcg agaaaacat ctccaagcc aaagggcagc  1020
cccgagaacc acaggtgtac accctgcccc catcccggga tgagctgacc aagaaccagg  1080
tcagcctgac ctgcctggtc aaaggcttct atcccagcga catgcctgtg gagggggaga  1140
gcaatgggca gccggagaac aactacaaga ccacgcctcc cgtgctggac tccgacggct  1200
ccttcttctc ctatagcaag ctcaccgtgg acaagagcag gtggcagcag gggaacgtct  1260
tctcatgctc cgtgatgcat gaggtctctg acaaccacta cacgcagaag agcctctccc  1320
tgtctccggg taaatgataa gtcgacatgc cctgaattct gcagatatcc atcacactgg  1380
cggccgctcg agcatgcatc tagagggccc                                     1410

```

<210> SEQ ID NO 42  
 <211> LENGTH: 445  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 42

```

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly
1           5           10           15
Ser Leu Arg Leu Ser Cys Val Gly Ser Arg Phe Thr Phe Ser Ala Tyr
20           25           30
Pro Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35           40           45
Ser Gly Ile Gly Ser Gly Gly Gly Thr Asn Tyr Ala Asp Ser Val Lys
50           55           60
Gly Arg Phe Thr Ile Ser Arg Asp Thr Ala Lys Asn Ser Leu Tyr Leu

```

-continued

65	70	75	80
Gln Met Asn Ser	Leu Arg Ala Glu Asp	Met Ala Val Tyr Tyr	Cys Ala
	85	90	95
Arg Gly Arg Asn Ser	Phe Asp Tyr Trp Gly	Gln Gly Thr Leu Val	Thr
	100	105	110
Val Ser Ser Ala Ser	Thr Lys Gly Pro Ser	Val Phe Pro Leu Ala	Pro
	115	120	125
Ser Ser Lys Ser Thr	Ser Gly Gly Thr Ala	Ala Leu Gly Cys Leu	Val
	130	135	140
Lys Asp Tyr Phe Pro	Glu Pro Val Thr Val	Ser Trp Asn Ser Gly	Ala
	145	150	155
Leu Thr Ser Gly Val	His Thr Phe Pro Ala	Val Leu Gln Ser Ser	Gly
	165	170	175
Leu Tyr Ser Leu Ser	Ser Val Val Thr Val	Pro Ser Ser Ser Leu	Gly
	180	185	190
Thr Gln Thr Tyr Ile	Cys Asn Val Asn His	Lys Pro Ser Asn Thr	Lys
	195	200	205
Val Asp Lys Lys Val	Glu Pro Lys Ser Cys	Asp Lys Thr His Thr	Cys
	210	215	220
Pro Pro Cys Pro Ala	Pro Glu Leu Leu Gly	Gly Pro Ser Val Phe	Leu
	225	230	235
Phe Pro Pro Lys Pro	Lys Asp Thr Leu Met	Ile Ser Arg Thr Pro	Glu
	245	250	255
Val Thr Cys Val Val	Val Asp Val Ser His	Glu Asp Pro Glu Val	Lys
	260	265	270
Phe Asn Trp Tyr Val	Asp Gly Val Glu Val	His Asn Ala Lys Thr	Lys
	275	280	285
Pro Arg Glu Glu Gln	Tyr Asn Ser Thr Tyr	Arg Val Val Ser Val	Leu
	290	295	300
Thr Val Leu His Gln	Asp Trp Leu Asn Gly	Lys Glu Tyr Lys Cys	Lys
	305	310	315
Val Ser Asn Lys Ala	Leu Pro Ala Pro Ile	Glu Lys Thr Ile Ser	Lys
	325	330	335
Ala Lys Gly Gln Pro	Arg Glu Pro Gln Val	Tyr Thr Leu Pro Pro	Ser
	340	345	350
Arg Asp Glu Leu Thr	Lys Asn Gln Val Ser	Leu Thr Cys Leu Val	Lys
	355	360	365
Gly Phe Tyr Pro Ser	Asp Ile Ala Val Glu	Trp Glu Ser Asn Gly	Gln
	370	375	380
Pro Glu Asn Asn Tyr	Lys Thr Thr Pro Pro	Val Leu Asp Ser Asp	Gly
	385	390	395
Ser Phe Phe Leu Tyr	Ser Lys Leu Thr Val	Asp Lys Ser Arg Trp	Gln
	405	410	415
Gln Gly Asn Val Phe	Ser Cys Ser Val Met	His Glu Ala Leu His	Asn
	420	425	430
His Tyr Thr Gln Lys	Ser Leu Ser Leu Ser	Pro Gly Lys	
	435	440	445

&lt;210&gt; SEQ ID NO 43

&lt;211&gt; LENGTH: 645

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapiens

-continued

&lt;400&gt; SEQUENCE: 43

```

gacatccaga tgaccagctc tccatcctca ctgtctgcat ctgtaggaga cagagtcacc    60
atcacttgtc gggcgagtcg gggattagc acctgggttag cctgggatca gcagaaacca    120
gagaaagccc ctaagtccct gatctatgct gcataccagtt tgcagagtgg ggtcccacg    180
aggttcagcg gcagtggtgc tgggacagat ttcactctca ccatcagcag cctgcagcct    240
gaagattttg caacttatta ctgccaacag tataatagtt accctccgac gttcggccaa    300
gggaccaagg tggagatcaa acgaactgtg gctgcaccat ctgtcttcat cttcccgcga    360
tctgatgagc agttgaaatc tggaactgcc tctgttgtgt gcctgctgaa taacttctat    420
cccagagagg ccaaagtaca gtggaaggtg gataacgccc tccaatcggg taactcccag    480
gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg    540
ctgagcaaa gacactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc    600
ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gttag                    645

```

&lt;210&gt; SEQ ID NO 44

&lt;211&gt; LENGTH: 214

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 44

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1           5           10          15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Thr Trp
                20           25           30
Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys Ala Pro Lys Ser Leu Ile
                35           40           45
Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
                50           55           60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65           70           75           80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Tyr Pro Pro
                85           90           95
Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
                100          105          110
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
                115          120          125
Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
                130          135          140
Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
145          150          155          160
Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
                165          170          175
Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
                180          185          190
Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
                195          200          205
Phe Asn Arg Gly Glu Cys
210

```

-continued

---

<210> SEQ ID NO 45  
 <211> LENGTH: 1413  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 45

```

gaggtcacgc tgggtgcagtc tgggggaggc ttggtacatc ctggggggtc cctgagactc   60
tcctgtgcag gctctggatt caccttcagt ggccatgctt tgcactgggt tcgccaggct   120
ccaggaaaag gtctggagtg ggtatcaggt attggtactc atggtgggac atactatgca   180
gactccgtga agggccgatt caccatctcc agagacaatg ccaagaactc cttgtttctt   240
caaatgaaca gcctgagcgc cgaggacatg gctgtgtatt actgtacaag aagaaactgg   300
ggacaatttg actactgggg ccaggaacc ctggtcaccg tctctagtgc ctccaaccag   360
ggcccatcgg ttttccccct ggcaccctcc tccaagagca cctctggggg cacagcggcc   420
tgggtgcctt ggtcaaggac tacttccccg aaccgggtgac ggtgtcgtgg aactcaggcg   480
ccctgaccag cggcgtgcac accttccccg ctgtcctaca gtctcagga ctctaactcc   540
tcagcagcgt ggtgaccgtg cctccagca gcttggggcac ccagacctac atctgcaacg   600
tgaatcacia gcccagcaac accaagggtg acaagaaagt tgagccaaa tcttgtgaca   660
aaactcacac atgcccaccg tgcccagcac ctgaactcct ggggggaccg tcagtcttcc   720
ttttcccccc aaaaccaag gacaccctca tgatctcccg gaccctgag gtcacatgag   780
tgggtgtgga cgtgagccac gaagaccctg aggtcaagtt caactggtac gtggacggcg   840
tggaggtgca taatgccaa gcaaaagccgc gggaggagca gtacaacagc acgtaccgtg   900
tggtcagcgt cctcaccgtc ctgaccagg actggctgaa tggcaaggag tacaagtgca   960
aggtctccaa caaagccctc ccagccccc tgcagaaaac catctccaaa gccaaagggc  1020
agccccgaga accacaggtg tacaccctgc ccccatcccc ggatgagctg accaagaacc  1080
aggtcagcct gacctgctg gtcaaaggct tctatcccag cgacatgcc gtggagtggg  1140
agagcaatgg gcagccggag aacaactaca agaccacgcc tcccgtgctg gactccgacg  1200
gtcctctctt cctctatagc aagctcaccg tggacaagag caggtggcag caggggaaacg  1260
ttttctcatg ctccgtgatg catgaggctc tgcacaacca ctacacgag aagagcctct  1320
cctgtctccc gggtaaatga taagtgcaca tgcctgaaat tctgcagata tccatcacac  1380
tggcggccgc tcgagcatgc atctagaggg ccc                               1413

```

<210> SEQ ID NO 46  
 <211> LENGTH: 446  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 46

```

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly
1           5           10          15
Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Gly His
20          25          30
Ala Leu His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35          40          45
Ser Gly Ile Gly Thr His Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys
50          55          60
Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Phe Leu

```

-continued

65	70				75				80				
Gln Met Asn Ser	Leu Ser Ala Glu Asp	Met Ala Val Tyr Tyr Cys Thr	85	90	95								
Arg Arg Asn Trp	Gly Gln Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val	100	105	110									
Thr Val Ser Ser	Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala	115	120	125									
Pro Ser Ser Lys Ser	Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu	130	135	140									
Val Lys Asp Tyr Phe	Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly	145	150	155									
Ala Leu Thr Ser	Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser	165	170	175									
Gly Leu Tyr Ser	Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu	180	185	190									
Gly Thr Gln Thr Tyr	Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr	195	200	205									
Lys Val Asp Lys Lys	Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr	210	215	220									
Cys Pro Pro Cys Pro	Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe	225	230	235									
Leu Phe Pro Pro	Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro	245	250	255									
Glu Val Thr Cys	Val Val Val Asp Val Ser His Glu Asp Pro Glu Val	260	265	270									
Lys Phe Asn Trp Tyr	Val Asp Gly Val Glu Val His Asn Ala Lys Thr	275	280	285									
Lys Pro Arg Glu Glu	Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val	290	295	300									
Leu Thr Val Leu His	Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys	305	310	315									
Lys Val Ser Asn Lys	Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser	325	330	335									
Lys Ala Lys Gly Gln	Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro	340	345	350									
Ser Arg Asp Glu Leu	Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val	355	360	365									
Lys Gly Phe Tyr Pro	Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly	370	375	380									
Gln Pro Glu Asn Asn	Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp	385	390	395									
Gly Ser Phe Phe Leu	Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp	405	410	415									
Gln Gln Gly Asn Val	Phe Ser Cys Ser Val Met His Glu Ala Leu His	420	425	430									
Asn His Tyr Thr Gln	Lys Ser Leu Ser Leu Ser Pro Gly Lys	435	440	445									

&lt;210&gt; SEQ ID NO 47

&lt;211&gt; LENGTH: 648

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapiens

-continued

&lt;400&gt; SEQUENCE: 47

```

gaaattgtgc tgactcagtc tccagccacc ctgtctttgt ctccagggga aagagccacc    60
ctctcctgca gggccagtca gagggttagc agctacttag cctggtacca acagaaacct    120
ggccaggctc ccaggctcct catctatgat gcatccaaca gggccactgg catcccagcc    180
aggttcagtg gcagtgggtc tgggacagac ttcactctca ccatcagcag cctagagcct    240
gaagattttg cagtttatta ctgtcagcag cgtagcaact ggccctccgta cacttttggc    300
caggggacca agctggagat caaacgaact gtggctgcac catctgtcct catcttcccg    360
ccatctgatg agcagttgaa atctggaact gcctctgttg tgtgctgct gaataacttc    420
tatcccagag aggccaaagt acagtggaag gtggataacg ccctccaatc gggtaactcc    480
caggagagtg tcacagagca ggacagcaag gacagcacct acagcctcag cagcacctg    540
acgctgagca aagcagacta cgagaaacac aaagtctacg cctgcgaagt cacccatcag    600
ggcctgagct cgcccgtcac aaagagcttc aacaggggag agtgtagg                    648

```

&lt;210&gt; SEQ ID NO 48

&lt;211&gt; LENGTH: 215

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 48

```

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1             5             10            15
Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
          20            25            30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
      35            40            45
Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
 50            55            60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65            70            75            80
Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Pro
      85            90            95
Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala
    100           105           110
Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser
    115           120           125
Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu
    130           135           140
Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser
    145           150           155           160
Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu
    165           170           175
Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val
    180           185           190
Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys
    195           200           205
Ser Phe Asn Arg Gly Glu Cys
    210           215

```

-continued

<210> SEQ ID NO 49  
 <211> LENGTH: 1413  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 49

```

gagggtgcagc tgggtgcagtc tgggggaggc ttggtacatc ctggggggtc cctgagactc   60
tcctgtgaag gctctggatt caccttcagt agcaatgta tgcactgggt gcgccagact   120
ccaggaaaag gtctggagtg ggtatcaggt attggtactg ctggtggcac atactatgca   180
gactccgtga agggccgatt caccatttcc agagacaatg tcaagaagtc cttgtatctt   240
caaatgaaca gcctgagagc cgaggacatg gctatttatt attgtgtaag aaaagactgg   300
ggatggttcg acccctgggg ccaggggacc ctggtcaccg tctctagtgc ctccaccaag   360
ggcccatcgg tcttccccct ggcaccctcc tccaagagca cctctggggg cacagcggcc   420
tgggctgcct ggtcaaggac tacttccccg aaccgggtgac ggtgtcgtgg aactcaggcg   480
ccctgaccag cggcgtgcac accttccccg ctgtcctaca gtctcagga ctctactccc   540
tcagcagcgt ggtgaccgtg cctccagca gcttggggcac ccagacctac atctgcaacg   600
tgaatcacia gcccagcaac accaaggtgg acaagaaagt tgagccaaa tcttgtgaca   660
aaactcacac atgcccaccg tgcccagcac ctgaactcct ggggggaccg tcagtcttcc   720
tcttcccccc aaaacccaag gacaccctca tgatctcccg gaccctgag gtcacatgag   780
tgggtgtgga cgtgagccac gaagaccctg aggtcaagtt caactggtac gtggacggcg   840
tggaggtgca taatgccaa gcaaaagccgc gggaggagca gtacaacagc acgtaccgtg   900
tggtcagcgt cctcaccgtc ctgaccagg actggctgaa tggcaaggag tacaagtgca   960
aggtctccaa caaagccctc ccagccccc tgcagaaaac catctccaaa gccaaagggc  1020
agccccgaga accacaggtg tacaccctgc ccccatcccg ggatgagctg accaagaacc  1080
aggtcagcct gacctgctg gtcaaaggct tctatcccag cgacatgcc gtggagtggg  1140
agagcaatgg gcagccggag aacaactaca agaccacgcc tcccgtgctg gactccgacg  1200
gtcctctctt cctctatagc aagctcaccg tggacaagag caggtggcag caggggaaacg  1260
tcttctcatg ctccgtgatg catgaggctc tgcacaacca ctacacgag aagagcctct  1320
cctgtctccc gggtaaatga taagtgcaca tgcctgaaat tctgcagata tccatcacac  1380
tggcggccgc tcgagcatgc atctagaggg ccc                               1413

```

<210> SEQ ID NO 50  
 <211> LENGTH: 446  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 50

```

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His Pro Gly Gly
1           5           10           15
Ser Leu Arg Leu Ser Cys Glu Gly Ser Gly Phe Thr Phe Ser Ser Asn
20           25           30
Gly Met His Trp Val Arg Gln Thr Pro Gly Lys Gly Leu Glu Trp Val
35           40           45
Ser Gly Ile Gly Thr Ala Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys
50           55           60
Gly Arg Phe Thr Ile Ser Arg Asp Asn Val Lys Lys Ser Leu Tyr Leu

```

-continued

65	70	75	80
Gln Met Asn Ser	Leu Arg Ala Glu Asp	Met Ala Ile Tyr Tyr	Cys Val
	85	90	95
Arg Lys Asp Trp	Gly Trp Phe Asp	Pro Trp Gly Gln Gly Ala	Leu Val
	100	105	110
Thr Val Ser Ser	Ala Ser Thr Lys Gly	Pro Ser Val Phe Pro	Leu Ala
	115	120	125
Pro Ser Ser Lys	Ser Thr Ser Gly Gly	Thr Ala Ala Leu Gly	Cys Leu
	130	135	140
Val Lys Asp Tyr	Phe Pro Glu Pro Val	Thr Val Ser Trp Asn Ser	Gly
	145	150	155
Ala Leu Thr Ser	Gly Val His Thr Phe	Pro Ala Val Leu Gln Ser	Ser
	165	170	175
Gly Leu Tyr Ser	Leu Ser Ser Val Val	Thr Val Pro Ser Ser	Ser Leu
	180	185	190
Gly Thr Gln Thr	Tyr Ile Cys Asn Val	Asn His Lys Pro Ser	Asn Thr
	195	200	205
Lys Val Asp Lys	Lys Val Glu Pro Lys	Ser Cys Asp Lys Thr	His Thr
	210	215	220
Cys Pro Pro Cys	Pro Ala Pro Glu Leu Leu	Gly Gly Pro Ser Val	Phe
	225	230	235
Leu Phe Pro Pro	Lys Pro Lys Asp Thr	Leu Met Ile Ser Arg	Thr Pro
	245	250	255
Glu Val Thr Cys	Val Val Val Asp Val	Ser His Glu Asp Pro	Glu Val
	260	265	270
Lys Phe Asn Trp	Tyr Val Asp Gly Val	Glu Val His Asn Ala	Lys Thr
	275	280	285
Lys Pro Arg Glu	Glu Gln Tyr Asn Ser	Thr Tyr Arg Val Val	Ser Val
	290	295	300
Leu Thr Val Leu	His Gln Asp Trp Leu	Asn Gly Lys Glu Tyr	Lys Cys
	305	310	315
Lys Val Ser Asn	Lys Ala Leu Pro Ala	Pro Ile Glu Lys Thr	Ile Ser
	325	330	335
Lys Ala Lys Gly	Gln Pro Arg Glu Pro	Gln Val Tyr Thr Leu	Pro Pro
	340	345	350
Ser Arg Asp Glu	Leu Thr Lys Asn Gln	Val Ser Leu Thr Cys	Leu Val
	355	360	365
Lys Gly Phe Tyr	Pro Ser Asp Ile Ala	Val Glu Trp Glu Ser	Asn Gly
	370	375	380
Gln Pro Glu Asn	Asn Tyr Lys Thr Thr	Pro Pro Val Leu Asp	Ser Asp
	385	390	395
Gly Ser Phe Phe	Leu Tyr Ser Lys Leu	Thr Val Asp Lys Ser	Arg Trp
	405	410	415
Gln Gln Gly Asn	Val Phe Ser Cys Ser	Val Met His Glu Ala	Leu His
	420	425	430
Asn His Tyr Thr	Gln Lys Ser Leu Ser	Leu Ser Pro Gly Lys	
	435	440	445

&lt;210&gt; SEQ ID NO 51

&lt;211&gt; LENGTH: 648

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapiens



-continued

&lt;400&gt; SEQUENCE: 51

```

gaaattgtgc tgaccagctc tccagccacc ctgtctttgt ctccagggga aagagccacc    60
ctctcctgca gggccagtca gagtattagc agctacttag cctggtagca acagaaacct    120
ggccaggctc ccaggctcct catctatgat gcatccaaca gggccactgg catcccagcc    180
aggttcagtg gcagtgggctc tgggacagac ttcactctca ccatcagcag cctagagcct    240
gaagattttg cagtttatta ctgtcagcag cgtagcaaat ggccctccgta cacttttggc    300
caggggacca agctggagat caaacgaact gtggctgcac catctgtcct catcttcccg    360
ccatctgatg agcagttgaa atctggaact gcctctgttg tgtgctgct gaataacttc    420
tatcccagag aggccaaagt acagtggaag gtggataacg ccctccaatc gggtaactcc    480
caggagagtg tcacagagca ggacagcaag gacagcacct acagcctcag cagcacctcg    540
acgctgagca aagcagacta cgagaaacac aaagtctacg cctgcgaagt cacccatcag    600
ggcctgagct cgcccgtcac aaagagcttc aacaggggag agtgtagg                    648

```

&lt;210&gt; SEQ ID NO 52

&lt;211&gt; LENGTH: 215

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 52

```

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1             5             10            15
Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Ile Ser Ser Tyr
          20            25            30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
      35            40            45
Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
 50            55            60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65            70            75            80
Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Lys Trp Pro Pro
          85            90            95
Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala
      100           105           110
Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser
      115           120           125
Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu
      130           135           140
Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser
      145           150           155           160
Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu
      165           170           175
Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val
      180           185           190
Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys
      195           200           205
Ser Phe Asn Arg Gly Glu Cys
      210           215

```

-continued

---

```

<210> SEQ ID NO 53
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for PCR
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (18)..(23)
<223> OTHER INFORMATION: n is a, c, t, or g

<400> SEQUENCE: 53
ggccgatag gcttcacnn nnnt                24

<210> SEQ ID NO 54
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for PCR

<400> SEQUENCE: 54
ggacactgac atggactgaa ggagta            26

<210> SEQ ID NO 55
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer PCR

<400> SEQUENCE: 55
ggggtcagc tggaactgag g                21

<210> SEQ ID NO 56
<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for PCR

<400> SEQUENCE: 56
cagcagaagc ttctagacca ccatggacat gagggtgcc gctcagctcc tggg            54

<210> SEQ ID NO 57
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for PCR

<400> SEQUENCE: 57
ccgctcagct cctggggctc ctgctgctgt ggctgagagg tgccagat                48

<210> SEQ ID NO 58
<211> LENGTH: 65
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for PCR

<400> SEQUENCE: 58
gtggttgaga ggtgccagat gtgaaattgt gctgacceag tctccagcca cctgtcttt            60
gtctc                65

```

---

-continued

---

<210> SEQ ID NO 59  
<211> LENGTH: 34  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: oligonucleotide primer for PCR  
  
<400> SEQUENCE: 59  
  
cttgctgact caaactctc ccctgttgaa gctc 34  
  
<210> SEQ ID NO 60  
<211> LENGTH: 22  
<212> TYPE: PRT  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: signal sequence for mammalian expression  
  
<400> SEQUENCE: 60  
  
Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Trp  
1 5 10 15  
  
Leu Arg Gly Ala Arg Cys  
20  
  
<210> SEQ ID NO 61  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: oligonucleotide primer for PCR  
  
<400> SEQUENCE: 61  
  
saggtscagy tkgtsagtc 20  
  
<210> SEQ ID NO 62  
<211> LENGTH: 17  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: oligonucleotide primer for PCR  
  
<400> SEQUENCE: 62  
  
ctgagttcca cgacacc 17  
  
<210> SEQ ID NO 63  
<211> LENGTH: 54  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: oligonucleotide primer for PCR  
  
<400> SEQUENCE: 63  
  
cagcagaagc ttctagacca ccatggacat gaggtgccc gtcagctcc tggg 54  
  
<210> SEQ ID NO 64  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: oligonucleotide primer for PCR  
  
<400> SEQUENCE: 64  
  
ccgctcagct cctggggctc ctgctgctgt ggctgagagg tgccagat 48

-continued

---

<210> SEQ ID NO 65  
 <211> LENGTH: 43  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: oligonucleotide primer for PCR  
  
 <400> SEQUENCE: 65  
  
 gtggttgaga ggtgccagat gtgaggtgca gctggtgcag tct 43

<210> SEQ ID NO 66  
 <211> LENGTH: 50  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: oligonucleotide primer for PCR  
  
 <400> SEQUENCE: 66  
  
 gtggaggcac tagagacggt gaccaggcct ccctggcccc aggggtcgaa 50

<210> SEQ ID NO 67  
 <211> LENGTH: 1409  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 67  
  
 tctagaccac catggacatc aggctcagct tagttttcct tgtccttttc ataaaagggtg 60  
 tccagtgtga ggtagaactg gtggagtctg ggggcggcctt agtacaacct ggaagggtcca 120  
 tgacactctc ctgtgcagcc tggggattca ctttcagaac ctatggcatg gectgggtcc 180  
 gccaggcccc aacgaagggt ctggagtggg tctcatcaat tactgctagt ggtggtacca 240  
 cctactatcg agactccgtg aagggccgct tcaactatctt tagggataat gcaaaaagta 300  
 ccctatacct gcagatggac agtccgaggt ctgaggacac ggccacttat tctgtacat 360  
 caatttcgga atactggggc cacggagtca tggtcaccgt ctctagtgcc tccaccaagg 420  
 gcccatcggt cttccccctg gcaccctcct ccaagagcac ctctgggggc acagcggccc 480  
 tgggtgcct ggtcaaggac tacttcccgg aaccgggtgac ggtgtcgtgg aactcaggcg 540  
 ccctgaccag cggcgtgcac acctcccgg ctgtcctaca gtcctcagga ctctactccc 600  
 tcagcagcgt ggtgaccgtg ccctccagca gcttggggcac ccagacctac atctgcaacg 660  
 tgaatcacia gccacgcaac accaagggtg acaagaaagt tgagccaaa tcttgtgaca 720  
 aaactcacac atgccaccg tgcccagcac ctgaactcct ggggggaccg tcagtcttcc 780  
 tcttcccccc aaaacccaag gacaccctca tgatctcccg gaccctgag gtcacatgcg 840  
 tgggtgtgga cgtgagccac gaagaccctg aggtcaagtt caactggtac gtggacggcg 900  
 tggaggtgca taatgccaa gcaaaagccg gggaggagca gtacaacagc acgtaccgtg 960  
 tggtcagcgt cctcacctgc ctgcaaccag actggctgaa tggcaaggag tacaagtgca 1020  
 aggtctccaa caaagccctc ccagccccc tgagaaaac catctccaaa gccaaaaggc 1080  
 agccccgaga accacagggt tacacctgc cccatcccgg ggtatgagctg accaagaacc 1140  
 aggtcagcct gacctgcctg gtcaaaggct tctatcccag cgacatcgcc gtggagtggg 1200  
 agagcaatgg gcagccggag aacaactaca agaccacgcc tcccgtgctg gactccgacg 1260  
 gctccttctt cctctatagc aagctcaccg tggacaagag caggtggcag caggggaacg 1320

-continued

---

tctttctcatg ctccgtgatg catgaggctc tgcacaacca ctacacgcag aagagcctct 1380

ccctgtctcc gggtaaatga taagtogac 1409

<210> SEQ ID NO 68  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 68

Ser Val Pro Thr Asp  
 1 5

<210> SEQ ID NO 69  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 69

Asp Leu Ala Thr Glu  
 1 5

<210> SEQ ID NO 70  
 <211> LENGTH: 50  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 70

Thr Leu Ser Asn Gly Lys Leu Arg Val Asn Gln Asp Gly Phe Tyr Tyr  
 1 5 10 15

Leu Tyr Ala Asn Ile Cys Phe Arg His His Glu Thr Ser Gly Ser Val  
 20 25 30

Pro Thr Asp Tyr Leu Gln Leu Met Val Tyr Val Val Lys Thr Ser Ile  
 35 40 45

Lys Ile  
 50

<210> SEQ ID NO 71  
 <211> LENGTH: 50  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 71

Thr Leu Ser Asn Gly Lys Leu Ile Val Asn Gln Asp Gly Phe Tyr Tyr  
 1 5 10 15

Leu Tyr Ala Asn Ile Cys Phe Arg His His Glu Thr Ser Gly Asp Leu  
 20 25 30

Ala Thr Glu Tyr Leu Gln Leu Met Val Tyr Val Thr Lys Thr Ser Ile  
 35 40 45

Lys Ile  
 50

<210> SEQ ID NO 72  
 <211> LENGTH: 50  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: murine DE variant amino acid sequence

<400> SEQUENCE: 72

-continued

---

Thr Leu Ser Asn Gly Lys Leu Arg Val Asn Gln Asp Gly Phe Tyr Tyr  
 1           5                   10                   15  
 Leu Tyr Ala Asn Ile Cys Phe Arg His His Glu Thr Ser Gly Asp Leu  
           20                   25                   30  
 Ala Thr Glu Tyr Leu Gln Leu Met Val Tyr Val Val Lys Thr Ser Ile  
           35                   40                   45  
 Lys Ile  
       50

<210> SEQ ID NO 73  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: oligonucleotide primer for PCR

<400> SEQUENCE: 73

cctctcatat ggactacaag gac 23

<210> SEQ ID NO 74  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: oligonucleotide primer for PCR

<400> SEQUENCE: 74

agtagccagg tctcccgatg tttcatgatg 30

<210> SEQ ID NO 75  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: oligonucleotide primer for PCR

<400> SEQUENCE: 75

ctggctactg aatatcttca gctgatggtg 30

<210> SEQ ID NO 76  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: oligonucleotide primer for PCR

<400> SEQUENCE: 76

cctctcctcg agttagtcta tgtcc 25

<210> SEQ ID NO 77  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 77

Gly His Ala Leu His  
 1           5

<210> SEQ ID NO 78  
 <211> LENGTH: 16  
 <212> TYPE: PRT

---

-continued

---

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 78

Gly Ile Gly Thr His Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys Gly  
1 5 10 15

<210> SEQ ID NO 79

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 79

Arg Asn Trp Gly Gln Phe Asp Tyr  
1 5

<210> SEQ ID NO 80

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 80

His Tyr Pro Leu His  
1 5

<210> SEQ ID NO 81

<211> LENGTH: 16

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 81

Gly Ile His Thr Gly Gly Gly Thr Tyr Tyr Thr Asp Ser Val Lys Gly  
1 5 10 15

<210> SEQ ID NO 82

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 82

Gly Arg Asn Ser Phe Asp Tyr  
1 5

<210> SEQ ID NO 83

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 83

Ala Tyr Pro Met His  
1 5

<210> SEQ ID NO 84

<211> LENGTH: 16

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 84

Gly Ile Gly Ser Gly Gly Gly Thr Asn Tyr Ala Asp Ser Val Lys Gly  
1 5 10 15

<210> SEQ ID NO 85

<211> LENGTH: 5

---

-continued

---

<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 85

Ser Tyr Gly Met His  
1 5

<210> SEQ ID NO 86  
<211> LENGTH: 16  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 86

Gly Ile Gly Thr Gly Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys Gly  
1 5 10 15

<210> SEQ ID NO 87  
<211> LENGTH: 8  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 87

Lys Asn Trp Gly Trp Phe Asp Tyr  
1 5

<210> SEQ ID NO 88  
<211> LENGTH: 5  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 88

Ser Asn Gly Met His  
1 5

<210> SEQ ID NO 89  
<211> LENGTH: 16  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 89

Gly Ile Gly Thr Ala Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys Gly  
1 5 10 15

<210> SEQ ID NO 90  
<211> LENGTH: 8  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 90

Lys Asn Trp Gly Trp Phe Asp Pro  
1 5

<210> SEQ ID NO 91  
<211> LENGTH: 8  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 91

Lys Asp Trp Gly Trp Phe Asp Pro  
1 5

<210> SEQ ID NO 92



---

-continued

---

<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 92

Arg Ala Ser Gln Gly Ile Ser Ser Trp Leu Ala  
1 5 10

<210> SEQ ID NO 93  
<211> LENGTH: 7  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 93

Ala Ala Ser Ser Leu Gln Ser  
1 5

<210> SEQ ID NO 94  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 94

Gln Gln Tyr Asn Ser Tyr Pro Pro Thr  
1 5

<210> SEQ ID NO 95  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 95

Arg Ala Ser Gln Gly Ile Ser Thr Trp Leu Ala  
1 5 10

<210> SEQ ID NO 96  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 96

Arg Ala Ser Gln Ser Ile Ser Ser Tyr Leu Ala  
1 5 10

<210> SEQ ID NO 97  
<211> LENGTH: 7  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 97

Asp Ala Ser Asn Arg Ala Thr  
1 5

<210> SEQ ID NO 98  
<211> LENGTH: 10  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 98

Gln Gln Arg Ser Lys Trp Pro Pro Tyr Thr  
1 5 10

-continued

---

```

<210> SEQ ID NO 99
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 99

Arg Ala Ser Gln Ser Val Asn Ser Tyr Leu Ala
1           5           10

```

```

<210> SEQ ID NO 100
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 100

```

```

Gln Gln Arg Ser Asn Trp Pro Pro Phe Thr
1           5           10

```

```

<210> SEQ ID NO 101
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 101

```

```

Arg Ala Ser Gln Ser Val Ser Ser Tyr Leu Ala
1           5           10

```

```

<210> SEQ ID NO 102
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 102

```

```

Gln Gln Arg Ser Asn Trp Pro Pro Tyr Thr
1           5           10

```

```

<210> SEQ ID NO 103
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 103

```

```

Asp Leu Ala Thr Glu
1           5

```

---

1. An isolated human antibody that specifically binds osteoprotegerin ligand (OPGL), comprising a heavy chain and a light chain, wherein the heavy chain comprises a heavy chain variable region comprising an amino acid sequence as set forth in any of SEQ ID NO: 6, SEQ ID NO: 14, SEQ ID NO: 22, or SEQ ID NO: 26, or an antigen-binding or an immunologically functional immunoglobulin fragment thereof.

2.-20. (canceled)

21. An isolated human antibody that specifically binds osteoprotegerin ligand (OPGL), wherein the antibody comprises:

a. a heavy chain having a heavy chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 6, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof, and a light chain having a light chain variable

region comprising an amino acid sequence as set forth in SEQ ID NO: 8, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof

b. a heavy chain having a heavy chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 14, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof, and a light chain having a light chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 16, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof

c. a heavy chain having a heavy chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 22, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment

- thereof, and a light chain having a light chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 24, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof;
- d. a heavy chain having a heavy chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 26, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof, and a light chain having a light chain variable region comprising an amino acid sequence as set forth in SEQ ID NO: 28, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof;
- e. a heavy chain comprising an amino acid sequence as set forth in SEQ ID NO: 30, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof, and a light chain comprising an amino acid sequence as set forth in SEQ ID NO: 32, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof;
- f. a heavy chain comprising an amino acid sequence as set forth in SEQ ID NO: 38, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof, and a light chain comprising an amino acid sequence as set forth in SEQ ID NO: 40, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof;
- g. a heavy chain comprising an amino acid sequence as set forth in SEQ ID NO: 46, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof, and a light chain comprising an amino acid sequence as set forth in SEQ ID NO: 48, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof; or
- h. a heavy chain comprising an amino acid sequence as set forth in SEQ ID NO: 50, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof, and a light chain comprising an amino acid sequence as set forth in SEQ ID NO: 52, an antigen-binding fragment thereof, or an immunologically functional immunoglobulin fragment thereof.
- 22.-49.** (canceled)
- 50.** The antibody of any of claims **1** or **21**, wherein the antibody inhibits binding of OPGL to an osteoclast differentiation and activation receptor (ODAR).
- 51.** A method of treating an osteopenic disorder in a patient, comprising administering to a patient a pharmaceutically effective amount of the antibody of claim **50**.
- 52.** A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of the antibody of claim **50**.
- 53.** A method of treating an osteopenic disorder in a patient; comprising administering to a patient the pharmaceutical composition of claim **52**.
- 54.** A method for detecting OPGL in a biological sample comprising:
- contacting the sample with the antibody of any of claims **1** or **21**; under conditions that allow for binding of the antibody to OPGL;
  - and measuring the level of bound antibody in the sample.
- 55.-73.** (canceled)
- 74.** An isolated nucleic acid molecule that encodes an antibody or antigen binding fragment thereof, that specifically binds osteoprotegerin ligand (OPGL), wherein the antibody or antigen binding fragment comprises: SEQ ID NO: 6; SEQ ID NO: 8; SEQ ID NO: 10; SEQ ID NO: 12; SEQ ID NO: 14; SEQ ID NO: 16; SEQ ID NO: 18; SEQ ID NO: 20; SEQ ID NO: 22; SEQ ID NO: 24; SEQ ID NO: 26; or SEQ ID NO: 28.
- 75.** An isolated nucleic acid molecule that encodes an antibody or antigen binding fragment thereof, that specifically binds osteoprotegerin ligand (OPGL), wherein the antibody or antigen binding fragment comprises: SEQ ID NO: 30; SEQ ID NO: 32; SEQ ID NO: 34; SEQ ID NO: 36; SEQ ID NO: 38; SEQ ID NO: 40; SEQ ID NO: 42; SEQ ID NO: 44; SEQ ID NO: 46; SEQ ID NO: 48; SEQ ID NO: 50; or SEQ ID NO: 52.
- 76.** An isolated nucleic acid molecule that encodes an antibody or antigen binding fragment thereof, that specifically binds osteoprotegerin ligand (OPGL), wherein the antibody or antigen binding fragment thereof comprises:
- SEQ ID NO: 77, SEQ ID NO: 78, SEQ ID NO: 79, SEQ ID NO: 92, SEQ ID NO: 93, and SEQ ID NO: 94;
  - SEQ ID NO: 80, SEQ ID NO: 81, SEQ ID NO: 82, SEQ ID NO: 95, SEQ ID NO: 93, and SEQ ID NO: 94;
  - SEQ ID NO: 83, SEQ ID NO: 84, SEQ ID NO: 82, SEQ ID NO: 96, SEQ ID NO: 97, and SEQ ID NO: 98;
  - SEQ ID NO: 85, SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 96, SEQ ID NO: 97, and SEQ ID NO: 98;
  - SEQ ID NO: 88, SEQ ID NO: 89, SEQ ID NO: 90, SEQ ID NO: 99, SEQ ID NO: 97, and SEQ ID NO: 100; or
  - SEQ ID NO: 88, SEQ ID NO: 89, SEQ ID NO: 91, SEQ ID NO: 101, SEQ ID NO: 97, and SEQ ID NO: 102.
- 77.** An isolated nucleic acid molecule that comprises a sequence that is complementary to the nucleic acid molecule of any of claims **74-76**.
- 78.** An expression vector comprising a nucleic acid molecule of any of claims **74-76**.
- 79.** An expression vector comprising a nucleotide sequence comprising two nucleic molecules of any of claims **74-75**.
- 80.** The expression vector of claim **79**, wherein the two nucleic acid molecules encode amino acid sequences comprising:
- SEQ ID NO: 6 and SEQ ID NO: 8;
  - SEQ ID NO: 10 and SEQ ID NO: 12;
  - SEQ ID NO: 14 and SEQ ID NO: 16;
  - SEQ ID NO: 18 and SEQ ID NO: 20;
  - SEQ ID NO: 22 and SEQ ID NO: 24;
  - SEQ ID NO: 25 and SEQ ID NO: 28;
  - SEQ ID NO: 30 and SEQ ID NO: 32;
  - SEQ ID NO: 34 and SEQ ID NO: 36;
  - SEQ ID NO: 38 and SEQ ID NO: 40;
  - SEQ ID NO: 42 and SEQ ID NO: 44;
  - SEQ ID NO: 46 and SEQ ID NO: 48; or
  - SEQ ID NO: 50 and SEQ ID NO: 52.
- 81.** An isolated host cell comprising the expression vector of claim **78**.
- 82.** An isolated host cell comprising the expression vector of claim **79**.
- 83.** An isolated host cell comprising a first expression vector and a second expression vector each according to claim **78**, wherein the first and second expression vectors each comprise a different sequence encoding amino acid sequences selected from the combinations:
- SEQ ID NO: 6 and SEQ ID NO: 8;

- (b) SEQ ID NO: 10 and SEQ ID NO: 12;
- (c) SEQ ID NO: 14 and SEQ ID NO: 16;
- (d) SEQ ID NO: 18 and SEQ ID NO: 20;
- (e) SEQ ID NO: 22 and SEQ ID NO: 24;
- (f) SEQ ID NO: 25 and SEQ ID NO: 28;
- (g) SEQ ID NO: 30 and SEQ ID NO: 32;
- (h) SEQ ID NO: 34 and SEQ ID NO: 36;
- (i) SEQ ID NO: 38 and SEQ ID NO: 40;
- (j) SEQ ID NO: 42 and SEQ ID NO: 44;
- (k) SEQ ID NO: 46 and SEQ ID NO: 48; or
- (l) SEQ ID NO: 50 and SEQ ID NO: 52.

**84.** The host cell of claim **81** wherein the host cell is a eukaryotic cell.

**85.** The host cell of claim **83** wherein the host cell is a eukaryotic cell.

**86.** The host cell of claim **85**, wherein the eukaryotic cell is a CHO cell.

**87.** A cell line comprising the expression vector of claim **78**.

**88.** A process of producing an antibody or antigen binding fragment thereof; comprising culturing the host cell of claim **81** under suitable conditions to express the antibody or antigen binding fragment thereof; and optionally isolating the antibody or antigen binding fragment thereof from the culture, wherein the antibody or antigen binding fragment thereof specifically binds human osteoprotegerin ligand (OPGL).

**89.** An antibody or antigen binding fragment thereof produced by the process of claim **87**.

\* \* \* \* \*