

9/1. 36708

26992

METHOD FOR STIMULATION OR REPAIR AND
REGENERATION OF INTESTINAL GUT CELLS IN
INFANTS AND ENHANCING THE IMMUNE RESPONSE OF T-CELLS

ABSTRACT

Nucleosides and/or nucleotides are added to infant formula, both milk and non-milk based, to provide a formula having enhanced physiological properties and also closely resemble human milk. Additionally, nutritionally balanced diet formulations are described having nucleosides and/or nucleotides incorporated therein.

✓

10
11
12
13
14

BAD ORIGINAL

26992

METHOD FOR STIMULATION OR REPAIR AND
REGENERATION OF INTESTINAL GUT CELLS IN
INFANTS AND ENHANCING THE IMMUNE RESPONSE OF T-CELLS

BACKGROUND OF THE INVENTION AND PRIOR ART

The present invention relates to the composition of, and processes for making, products for infant formulas and for nutritional products suitable for both infants and adults, and particularly for clinical nutrition. These products may be administered orally or by enteral feeding tubes. These products are enriched with specific nucleosides, nucleotides, or mixtures thereof.

Infant formulas are derived, to a large extent, from cow's milk. After being diluted, the cow's milk is enriched with whey proteins, diverse carbohydrates, such as lactose, dextrinmaltose and starches, different mixtures of vegetal and animal fats, vitamins and minerals. These components are present in suitable amounts to meet the requirements of low birth weight newborns or those of at term healthy infants during the first and second semester of life.

Sometimes, infant formulas contain isolated milk proteins, isolated vegetable proteins or protein hydrolyzates, from different origins such as casein, lactalbumin, soy and meat. Also, these infant formulas have one or more carbohydrates (sucrose, dextrinmaltose and starch), mixtures of diverse kind of fats, minerals and vitamins, to meet not only the healthy newborns' nutritional requirements, but also of infants and children with clinical symptoms of

lactose intolerance, protein intolerance and, in general, with diverse malabsorption nutrition syndromes.

The European Society of Pediatric Gastroenterology and Nutrition (ESPGAN), the American Academy of Pediatric (AAP),
5 the Codes Alimentarius Mundi, and the European Community Council, among other organizations, have given general rules for the composition of infant formulas (ESPGAN Committee on Nutrition, Acta Paed Scand, Supl 262. 1977: EPSGAN
Committee on Nutrition Acta Paed Scand, Supl 287, 1981:
10 ESPGAN Committee on Nutrition. Acta Paed Scand, Supl 302, 1982; ESPGAN Committee on Nutrition. Acta Paed Scand. Supl 330, 1987; AAP Committee on Nutrition, Pediatric Nutrition Handbook, 1979; AAP Committee on Nutrition, Pediatrics, 75, 976, 1985: EEC Council. 85/C 28/05 COM (84) 703 final, 15 1985; EEC Council, 86/C 124/06 COM/86 91 final, 1986 Codes Alimentarius Mundi, Codex Stan 72-1981).

As used herein, the term "infant formulas" is intended to refer to the well established understanding as defined by ESPGAN Committee on Nutrition, Acta Paed Scand, supl 262, pg
20 3, supra. and also the American Academy of Pediatrics (Pediatrics, Vol 57 no. 2, pg. 281, February 1976).

In general, infant formulas tend to have a composition qualitatively and quantitatively as similar as possible to human milk. Nevertheless, despite the efforts made by
25 several researchers, infant formulas still have a number of differences in their composition compared to human milk. This is because the latter has many substances, such as immunoglobulin, free amino acids, polyamines, nucleotides, polyunsaturated fatty acids, etc. which are not present in

cow's milk. Thus, it would be desirable that infant milk formulas have most of the substances present in human milk so as to produce the same physiological effects as human milk.

5 Nutritional products, such as those currently used in hospitals, special or for dietary purposes, are based on the utilization of diverse protein sources (casein, sodium and calcium caseinate, isolated soy protein, protein hydrolyzates and/or crystalline amino acids) mixtures of vegetables
10 and animal fats, carbohydrates (basically glucose polymers), vitamins and minerals to meet, at least, the dietary intakes recommended for healthy individuals (Committee on Dietary Allowances, Food and Nutrition Board, Nat Acad Sci, 9th Ed. 1980).

15 Protein energy malnutrition (PEM) is found in many patients admitted to hospitals. This happens not only in developing countries, but also in those with a high socio-economic level where the percentages of medicalsurgical patients vary between 40-50% (Bistran, et. al. JAMA, 235,
20 1567, 1976; G. Hill et al. Lancet, 1, 689, 1977; Gassull et al. Human Nutr: Clin Nutr, 38C, 419, 1984). Proper nutritional support for such patients, while not a primary mode of treatment is, nevertheless, an important factor for therapy and recovery. It is therefore important to adminis-
25 ter a nutritionally balanced diet given orally, enterally or parenterally, adequate to the needs of the patient. This is especially true for those patients where conventional feeding is contraindicated (gastroenterological patients) or is insufficient (hypercatabolic patients). The enteral

or oral mode of administration of foods is preferable to parenteral modes (E. Cabre and M. A. Gassull, J. Clin Gastroenterol Nutr. 1, 97, 1986) because of the lower morbidity, trophic effect upon the intestinal mucose, lower
5 necessity for instruments and lower costs.

Nutritional products for proper diets associated with parenteral administration should be formulated to meet the requirements of the individual needs in specific situations. Thus, complete balanced diets with an energy content between
10 130-150 Kcal/g nitrogen, are recommended for the preventive and repletive therapy in cases of PEM due to nervous anorexy, esophageal stenosis, maxillofacial surgery, chronic vasculo-cerebral disease, long evolution neurological syndromes, vascular surgery postoperative period,
15 malabsorption syndromes, preoperative period, complete intestinal occlusion, preparation of colon (surgery, radiology and endoscopy) and, in general, in all cases when it is necessary to take a balanced diet of nutrients. Diets with a high content of nitrogen (80-120 Kcal/g nitrogen) are
20 recommended for the nutritional therapy of burn patients or patients suffering cranial trauma, multiple trauma, open fractures, , Crohn disease, ulcerous colitis, digestive fistula, sepsis, oncology surgery, oncological radiotherapy and chemotherapy, pre-and postoperative periods, orthopedic
25 surgery, and, in general, for catabolic patients.

Diets containing protein hydrolyzates as a source of amino nitrogen are specially made for the nutritional support of patients with diverse malabsorption-malnutrition syndromes, such as short bowel, acute celiac disease, Crohn

disease, chronic pancreatic insufficiency, cystic fibrosis, intestinal fistulas, postoperative nutrition, and the like,

Furthermore, such products can be made as specific clinical diets for specific diseases, such as hepatopathies, chronic renal disease, and chronic obstructive pulmonary disease.

In addition, there is a variety of dietary products marketed to meet the nutritional needs of various individuals. For example, many individuals desirous of achieving varying degrees of weight loss, may benefit from the use of a special nutrition diet formulations to provide specific nutrients otherwise provided by a normal diet. Likewise, many people find it necessary to supplement their daily diet with additional nutrients due to age, allergy or physical afflictions.

As used herein, the term "nutritionally balanced diet formulations" is intended to refer to the above type of products.

Currently marketed nutrition products do not contain nucleic acids or their simpler compounds, either nucleosides and/or nucleotides, which are normally present in foods and carry out fundamental physiological functions, described further on.

In relation to the nutritional importance of nucleotides, some relevant aspects of these compounds such as their content in human milk, physiological effects in newborns, intestinal absorption, tissue utilization and effects upon cell immunity are shown below.

U.S. Pat. No. 4,544,559 teaches that human milk has a specific nucleotide content, very different from cow's milk. Human milk contains, at least; twelve different nucleotides, predominating cytidine monophosphate (CMP), adenosine monophosphate (AMP), uridine monophosphate (UMP), guanosine monophosphate (GMP), inosine monophosphate (IMP) and uridine derivatives, whereas cow's milk has very low amounts of CMP and AMP; it lacks the other nucleotides and has high amounts of orotic acid, which is absent in human milk.

Also, U.S. Pat. No. 4,544,559 teaches that a humanized milk enriched with nucleotides AMP, CMP, GMP, UMP and IMP in the same range as human milk, stimulates the development of *Bifidobacterium bifidum* Ti at the intestinal level. This bacterium comprises 80% of the total bifidobacteria present in the feces of breast-fed-newborns. Furthermore, this humanized milk promotes serum fatty acid profile very similar to that found in newborns fed with human milk.

Ziro et. al. U.S. Pat. No. 3,231,385 describes infant milk formulas supplemented with certain nucleotides to simulate human milk, improve the milk taste and lower the curd tension.

Nucleotides can be synthesized in most tissues by two processes: (a) de novo synthesis from the precursors which include pirophosphoribosilphosphate, glutamine, aspartate, glycine, formiate and carbon dioxide and (b) utilization of bases and the nucleosides liberated through the catabolism of nucleotides and nucleic acids contained in foods by the "salvage pathway" (S. Leleiko et al. *J Pediatr Gastroenterol Nutr.* , 313, 1983). This last way is an important alterna-

tive in the synthesis of nucleotides when the biosynthetic
de novo pathway is hindered by an insufficient supply of
precursors. Tissues such as bone marrow, intestine and the
liver are heavily dependent on the salvage pathway. The
5 activity of the "salvage pathway" has also been shown
demonstrated in kidney, brain and retina (P. Mandel,
Traite de biochemie generale, Polonovski, Boulanger,
Lemoigne, Wurmser, eds, Masson et Cie, Paris, 1972).

The intestinal mucosa needs a continuous supply of
10 nucleotides or their precursors from dietary origin, apart
from the hepatic supply by the vascular system, in order to
maintain continuous synthesis of RNA.

It has been confirmed in cuts made in the small
intestine of rats that the exogenous adenosine triphosphate
15 (ATP) increases the intracellular concentration of the
nucleotide and it has been observed that at temperature over
20°C the marked exogenous ATP is absorbed by everted sacs of
rat small intestine (J. Bronk and H Leese. J. Physiol.,
235,183, 1973; J. Blair, M. Lucas and A. Matty, J. Physiol.,
20 245, 333,1975). Also, it has been shown in rabbit's ileum
"in vitro" that, at low concentrations, the ATP as well as
the nucleoside adenosine are absorbed through a carrier
associated to the enterocyte membrane.

Since the carrier system works for ATP and adenosine,
25 it is likely that the system also works for other purine
nucleotides, because competitive inhibition measures have
proved that any compound with a purine ring united to a
ribose molecule is absorbed. (V. Harms and C. Stirling.
Am. J. Physiol., 233, E-47, 1977).

It has also been shown that the purines and pyrimidines in the RNA and DNA. present in the diet, are absorbed by mice, preferably as nucleosides. Between 2-5% of the nucleosides are used for nucleic acid synthesis in intestinal tissue, and cytosine nucleosides are used for DNA synthesis, specially in the spleen (F. Sonoda, M. Tatibana, Biochem. Biophys. Acta, 521, 55.1978).

It has also been shown that purine bases, such as adenine, guanine, hypoxanthine and xanthine are almost completely absorbed by rats. 4.5-6.5% being incorporated in tissues and in a greater proportion by the liver and intestine.

The absence of pyrimidine or purine derivatives in the diet is known to suppress the normal function of T lymphocytes (F. Rudolph et al. Adv. Exp. Med. Biol. 165, 175. 1984), and to increase the mortality in experimental animals by staphylococcus sepsis. The addition of pyrimidine and purine derivatives to the diet decreases the susceptibility of animals to infection (A. Kulkarni et al. JPEN, 10, 169,1986). Thus, the extent of purines and pyrimidines on the immune function can be a great importance in a number of clinic situations, such as transplants of organs in patients, malnutrition recovery, in diverse chemotherapeutic regimens and in the treatment of leukemias derived from T cells. Accordingly, one of the objects of the present invention is to provide improved nutritionally balanced diet formulations.

Another object of the present invention is to provide non-milk infant formulas which more closely resemble the

human milk of nursing mothers and also exhibit enhanced properties.

Still yet a further object of the present invention is to provide improved milk based infant formulas which not only closely resemble human milk, but which are more readily absorbed by the infant gut and enhance the infant's immune response.

These and other objects of the present invention will become more apparent from the discussion which follows.

10

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a range of compositions of infant formulas and clinical nutrition products enriched with nucleosides, nucleotides or mixes of these two classes of compounds and the processes for their preparation. The products are in a liquid ready to eat form, or concentrated liquid or powder.

According to the invention, adenosine, guanosine, cytidine, inosine, and uridine or their mixes are used as nucleosides, and adenosine phosphate, guanosine phosphate, cytidine phosphate, inosine phosphate and uridine phosphate are used as nucleotides.

The term uridine phosphate, guanosine phosphate, etc, is intended herein to refer collectively to the mono, di and/or tri phosphates as well as the sugar derivatives of the nucleotides mentioned. However, for various reasons which will be apparent to those knowledgeable in the art, the 5'-monophosphates are preferred.

The supplementation of nucleosides and/or nucleotides or their mixes to infant formulas and nutrition balanced diet formulations gives a better physiological fatty acid tissue membrane composition to newborns and adult patients, an improved cell immunity and a better intestinal repair in those patients with intestinal diseases.

Accordingly, one embodiment of the present invention provides for a nutritionally balanced diet formulation which comprises a source of amino nitrogen, carbohydrates, edible fats, minerals, vitamins and a nucleoside/nucleotide composition containing at least one of:

- (a) uridine, uridine phosphate or a mixture thereof;
 - (b) guanosine, guanosine phosphate or a mixture thereof;
 - (c) adenosine, adenosine phosphate or a mixture thereof;
 - (d) cytidine, cytidine phosphate or a mixture thereof;
- or
- (e) inosine, inosine phosphate or a mixture thereof.

Thus, the formulation must contain at least one of the fifteen different possible components in an amount (based on 100 grams of dry product) equal to 1 mg.

Generally, the diet formulation will contain (on a dry weight basis per 100 g) from 1 to 300 mg of components selected from (a), (b), (c), (d) and (e), with a preferred range being from about 50 to about 250 mg. The optimum amount appears to be about 150 mg per 100 grams product.

On a liquid basis, these ranges correspond to from about 0.2 to 60 mg/dl on a general basis, and preferably

about 10 to about 50 mg/dl, with the optimum being about 30 mg/dl.

A further embodiment of our invention provides for improved non-milk infant formulas. Such non-milk formulas are well known and generally comprise carbohydrates, a source of amino acids, vegetable oils, minerals and vitamins. According to this embodiment of the invention, there is added to such formulas at least one of uridine, uridine phosphate or mixtures thereof; guanosine, guanosine phosphate or mixtures thereof; adenosine, adenosine phosphate or mixtures thereof; cytidine, cytidine phosphate or mixtures thereof; or inosine, inosine phosphate or mixtures thereof. As a minimum, at least about 0.27 mg per 100 g of product of one of the components should be added to the infant formula. Generally, the non-milk infant formulas according to the present invention require on a dry weight basis in mg per 100 grams of total weight approximately the following additives:

- uridine and/or uridine phosphate..... 17.40-1.86 mg;
- guanosine and/or guanosine phosphate..... 3.32-0.27 mg;
- adenosine and/or adenosine phosphate..... 9.50-4.25 mg;
- cytidine and/or cytidine phosphate..... 10.16-3.52 mg, and
- inosine and/or inosine phosphate..... 1.92-0.00 mg.

On a liquid basis, per dl, these formulation correspond as follows:

- uridine and/or uridine phosphate..... 2.62-0.28 mg;
- guanosine and/or guanosine phosphate..... 0.50-0.04 mg;
- adenosine and/or adenosine phosphate..... 1.43-0.64 mg;
- cytidine and/or cytidine phosphate..... 1.53-0.53 mg, and
- inosine and/or inosine phosphate..... 0.29-0.00 mg.

For reasons discussed more fully below, it may be desirable to add small amounts of L-cistine and/or carnitine to the non-milk based infant formulas.

As yet a further embodiment of our invention there is provided an improved infant milk formula to which is added at least one nucleoside selected from the group consisting of uridine, guanosine, adenosine, cytidine and inosine. The added nucleosides must be present in an amount about 0.27 mg per 100 grams total product on a dry basis. To provide for a closer simulation of human breast milk and also enhance absorption by the infant gut, there should be added to the infant milk formula the following ingredients for each 100 g of total weight:

- uridine and/or uridine phosphate 17.40-1.86 mg;
- guanosine and/or guanosine phosphate 3.32-0.27 mg;
- adenosine and/or adenosine phosphate 3.75-0.00 mg;
- cytidine and/or cytidine phosphate 4.58-0.00 mg, and
- inosine and/or inosine phosphate 1.92-0.00 mg.

This of course correspond on a liquid basis (per dl) as follows:

- uridine and/or uridine phosphate 2.62-0.28 mg;
- guanosine and/or guanosine phosphate 0.50-0.04 mg;
- adenosine and/or adenosine phosphate 0.56-0.00 mg;
- cytidine and/or cytidine phosphate 0.69-0.00 mg, and
- inosine and/or inosine phosphate 0.29-0.00 mg.

Basically, infant formulas, according to the present invention have a composition adequate for meeting the requirements of low birth weight infants, at term infants,

children with lactose intolerance, children with cow's protein intolerance and/or malabsorption syndrome.

5 The infants and nutritionally balanced diet products of the present invention have been found to stimulate repair and regeneration of intestinal gut cells, enhance the immune response of T-cells and provide for specific fatty acid phospholipids profiles in re blood cell membranes.

10 The use of nucleosides herein is unique to the formulations of the present invention. These materials generally have been found to be at least as effective as their corresponding nucleotides, and even more effective in providing for enhanced absorption through use of the salvage pathway in the human body. This apparently may be due to the higher water solubility of nucleosides as compared to the corresponding nucleotides. Also, nucleoside stability when
15 used in the formulations of this invention is greater than that of the corresponding nucleotides.

20 When the nucleosides and/or nucleotides or their combinations are added to infant formulas in concentrations on the same range as human milk, according to this invention, they stimulate the conversion of essential fatty acids to their polyunsaturated derivatives (AGPI), which is reflected in the fatty acid composition of erythrocyte membrane both in the at-term newborn and in the preterm
25 newborn as well as in the fatty acids composition of plasma phospholipids.

In a study carried out by the inventors. 20 at-term newborns were fed exclusively on human milk. 19 with an infant formula and 19 with the same infant formula supple-

mented with nucleoside-5'-monophosphates according to this invention, in similar concentrations to those of human milk. The relative content of AGPI, of the w6 series, derived from linoleic acid, as well as w3 series, derived from linolenic acid, was significantly decreased, specially in phosphatidylethanolamine and phosphatidylserine of the erythrocyte membrane in infants fed milk formula with respect to infants fed nucleosides-5'-monophosphates supplemented milk formula or human milk. The same happened in the plasma phospholipids and cholesterol esters. The arachidonic (20:4w6) and docosahexaenoic (22:6w3) acids were the most increased fatty acids in infants fed nucleoside-5'-monophosphates supplemented milk formula, with respect to those fed milk formula.

In other study, 19 preterm infants were fed exclusively on human milk, 18 with an infant milk formula for pretermatures and 18 with the same milk formula supplemented with nucleosides -5'-monophosphates in concentrations similar to those of human milk, according to this invention. At one month of life, the relative contents of eicosatrienoic acid (20:3w6), arachidonic acid (20:4w6), docosatetraenoic acid (22:4w6) and docosapentaenoic acid (22:5w6) were significantly decreased in the erythrocyte membrane phospholipids in infants fed milk formula with respect to those fed nucleoside-5'-monophosphate supplemented milk formula or human milk. Also, infants fed nucleoside-5'-monophosphate supplemented milk formula showed an intermediate value of docosahexaenoic acid (22:6w3) between those fed human milk and those fed milk formula. The same results

were observed in the plasma phospholipids of preterm newborns.

The modulating effect of nucleosides and nucleotides of the diet upon cell immunity has been proved through the following method:

Six groups of BALB/C mice, constituted by 10 mice each, aged four weeks, weaning period, were feed with a conventional diet (Chow diet), a nucleosides and nucleotides free diet, a diet supplemented with nucleosides according to this invention, in the following proportions: 50 mg of uridine, 50 mg of guanosine, 50 mg of adenosine, 50 mg of citidine and 50 mg of inosine, a diet supplemented with nucleosides in proportions equivalent to mouse milk, a diet supplemented with 50 mg of the following nucleotides UMP, GMP, CMP, UMP according to this invention and a diet supplemented with nucleotides in proportions equivalent to mouse milk, respectively. The mice were feed during a period of four weeks, and with them we proceed with the testing of the cell immune response "in vitro" as response to allogeneic and syngeneic antigens using the lymphocyte mixed culture technique and quantifying the cell proliferation by the incorporation of ^3H -thymidine to DNA and secondly we proceed with the testing of the proliferation as response to phytohemagglutinin (mitogen agent) to quantify the state of lymphocyte reactivity, also with the incorporation of ^3H -thymidine.

The mice fed on the free nucleoside or nucleotide diets had an immune response mediated by T cells lower than the

other groups having a diet supplemented with these compounds.

The effect of nucleosides and nucleotides of the diet on the intestinal cell proliferation and on their enzymatic activity is proved as follows:

Two groups of Wistar mice, of 20 animals each, from the weaning (21 days of age), are fed during two weeks, the first of them on a diet (Diet A) containing 167 g of calcium caseinate. 489.5 of corn starch, 150 g of sugar, 50 g of cellulose, 100 g of soy oil, 3 g of DL-methionin, 1.1 g of choline chloride, 38.2 g of a mineral mixture and 1.2 g of a vitamin mixture, per Kg, to satisfy the nutritional requirements of these animals. The second group was fed with a similar diet, but with lactose instead of starch (Diet B). In this second group takes place an osmotic diarrhoea because of lactose intolerance giving rise to a malnutrition-malabsorption syndrome. Both groups are divided in two subgroups of 10 animals each, the first subgroup being fed on Diet A and the second with on Diet A supplemented with 50 mg of each of the following nucleosides: uridine, guanosine, adenosine, citidine and inosine, during 4 weeks or with 50 mg of each of the following nucleotides UMP, GMP, AMP, CMP and IMP, according to this invention.

The animals suffering malabsorption syndrome refed on the nucleoside or nucleotide supplemented diet, according to the invention, had ileal, jejunal and duodenal mucose weights significantly superior to those fed on a diet without such compounds. Also, the proportion of cells in a mitosis state, the mucose proteins content and the maltase

and sucrase enzymatic activities were significantly higher in animals fed on the nucleoside or nucleotide supplemented diet than in those fed on a free purine and pyrimidine diet.

Basic ingredients for infant formulas include cow's milk, protein, whey proteins, casein and its salts (i.e. calcium caseinate); soy protein isolates are substituted for milk derived proteins, and are used in the products made for infants with lactose intolerance and/or cow's protein intolerance. Protein hydrolyzates (i.e. casein and lactalbumin hydrolyzates) with low molecular weight, may also be used for the products.

The proportions of the diverse component nutrients are similar to those of human milk. Thus, the ratio of whey proteins to casein currently varies from 60:40 to 70:30 in infant formulas based on milk. The mixture of fats employed is made up of edible fats to provide an essential fatty acid profile. Lactose is used exclusively as the carbohydrate source for at-term newborns infants, except that dextrin-maltose is employed in products used for the treatment of lactose intolerance and malabsorption syndromes in infancy.

Infant formulas according to the invention contain minerals (including calcium, phosphorus, sodium, potassium, chloride, magnesium, iron, zinc, copper, manganese and iodine) and vitamins (including vitamin A, vitamin D₃, vitamin C, vitamin B₁, vitamin B₂, vitamin B₆, vitamin B₁₂, pantothenic acid, vitamin E, vitamin K, folic acid, biotin) adequate for the infants requirements. Also, in the products whose source of proteins is derived from soy or protein isolates or hydrolyzates, carnitine is included to

satisfy the nutritional requirement for this compound in infants with malabsorptive syndromes.

The inventors of the present compositions and processes have demonstrated that the amounts of cytosine, adenine, guanine, uracile and inosine derivatives in human milk vary between 1.53-0.54, 1.43-0.69, 0.50, 2.62 and 0.29-0.00 mg/dl respectively and the individual contents of CMP, AMP, GMP, UMP, and IMP oscillate between 1.37-0.53, 1.19-0.64, 0.21-0.04, 0.56-0.28, 0.29-0.00 mg/dl, respectively.

The content of nucleosides and/or nucleotides in the infant formulas of the present invention are in the range of those for human milk. An exemplary nucleoside and/or nucleotide mixture for infant formulas not containing cow's milk, according to the invention, is shown in Table I.

The amounts of adenosine and/or adenosine phosphate and cytidine and/or cytidine phosphate, inosine and/or inosine phosphate added to cow's milk based infant formulas, according to this invention, are lower than those shown in Table I, because cow's milk contains specific amounts of nucleosides and nucleotides.

In Table II an exemplary mixture of nucleosides and/or nucleotides for infant milk formulas containing cow's milk is shown.

The dietary products for balanced nutrition, according to the present invention, have a composition of nutrients adequate to the specific requirements of not only healthy human in need of a balanced nutritional product, but also those individuals in situations of energy-protein malnutri-

tion and in hypercatabolic states derived from traumatic, septic, surgical processes and malabsorption syndromes.

As nitrogenous sources, the following components are preferably employed: a mixture of dairy proteins (casein or sodium and calcium caseinates and lactose free lactalbumin) and protein hydrolyzates with low molecular weight (maximum molecular weight 1,000 daltons, average molecular weight, 500 daltons). As carbohydrate sources, glucose polymers are employed, such as dextrinmaltose with a different grade of dextrose equivalent degree, preferably between 10 and 30 DE. Fats are employed as a mixture of animal and one or more vegetable fats to meet the essential fatty acids requirements of patients.

Nutritional products according to the present invention provide mineral elements which include trace elements and vitamins in adequate proportions to satisfy the specific requirements of normal healthy individuals as well as those suffering malabsorption-malnutrition processes and in a hypercatabolic state.

The nutritional products are enriched with nucleosides and/or nucleotides in similar amounts of nucleotides to those present in foods.

An example of a nucleoside and/or nucleotide mixture for the enrichment of nutritional products is shown in Table III. On a dry weight basis, the amount of nucleosides and/or nucleotides may each vary from about 1 to about 300 mg per 100 grams of product, and preferably each ranges from about 50 to about 250 mg per 100 grams of product. On a liquid basis the amount may vary per deciliter of product

from about 0.2 to about 60 mg of each nucleoside and/or nucleotide, and preferably ranges from about 10 to about 50 mg.

TABLE I

5 Example of a characteristic mixture of nucleosides and/or nucleotides in infant formulas, not containing cow's milk.

10	Nucleosides and/or Nucleotides	Powdered product		Liquid product	
			Range mg/100g		Range mg/dl
15	Uridine/Uridine phosphate	3.42	17.40-1.86	0.51	2.62-0.28
	Guanosine/Guanosine phosphate	1.49	3.32-0.27	0.22	0.50-0.04
	Adenosine/Adenosine phosphate	6.90	9.50-4.25	1.03	1.43-0.64
	Cytidine/Cytidine phosphate	6.87	10.16-3.52	1.03	1.53-0.53
20	Inosine/Inosine phosphate	1.00	1.92-0.00	0.15	0.29-0.00

TABLE II

25 Example of a characteristic mixture of nucleosides and/or nucleotides in infant formulas based on cow's milk.

30	Nucleosides and/or Nucleotides	Powdered product		Liquid product	
			Range mg/100g		Range mg/dl
	Uridine/Uridine phosphate		17.40-1.86		2.62-0.28
	Guanosine/Guanosine phosphate		3.32-0.27		0.50-0.04
	Adenosine/Adenosine phosphate		3.75-0.00		0.56-0.00
	Cytidine/Cytidine phosphate		4.58-0.00		0.69-0.00
	Inosine/Inosine phosphate		1.92-0.00		0.29-0.00

TABLE III

35 Example of a characteristic mixture of nucleosides and/or nucleotides in nutritionally balanced diets.

40	Nucleosides and/or Nucleotides	Powdered product		Liquid product	
			Range mg/100g		Range mg/dl
	Uridine/Uridine phosphate	150	1-300	30	0.2-60
	Guanosine/Guanosine phosphate	150	1-300	30	0.2-60
	Adenosine/Adenosine phosphate	150	1-300	30	0.2-60
	Cytidine/Cytidine phosphate	150	1-300	30	0.2-60
	Inosine/Inosine phosphate	150	1-300	30	0.2-60

The invention also includes the process to obtain infant formulas, as well as specific diets to be used in good nutrition, enriched with nucleosides and or nucleotides (FIG. 1 to 4). The products can be prepared in liquid, ready to be used, concentrated to be diluted in water before its use, and in powder forms. This process comprises, in all cases, the preparation of a mixture containing water and non fat solids, except vitamins, minerals and nucleosides and or nucleotides, followed by a preheating to 75 -80 C., deaeration of the mixture, injection of the fat mixture, double homogenization at 70 -75 C. (usually 150 Kg/cm² in the first stage and 50 Kg/cm² in the second) cooling to 4 - 6 C. and storage in standardization tanks. The liquid products ready for consumption or concentrates to be disluted before use, are standardized in the said tanks, adapting the pH to values generally ranging from about 6.8 to about 7.1 and most preferably ranging from about 6.8 to 7.0 for infant formulas and from about 6.9 to 7.1 for nutritional products.

When the products are going to be UHT sterilized and aseptically packed in containers made of carton-aluminum-polyethylene, during the standardization, the vitamins, minerals and nucleotides or nucleosides mixtures are added as concentrated aqueous solutions and the content of mineral elements is adjusted by adding the required salts. The nucleoside and/or nucleotide solutions should be maintained preferably at pH 6-6.5 to avoid them to hydrolyze.

Once standardized, the products for consumption in liquid or concentrated forms, are sterilized through an UHT

system at 145 -150 C. for 2-4 seconds and can be either
aseptically packed or bottled in glass or polyethylene
bottles. In the latter case, products are standardized
prior to the UHT sterilization, only in their solids
5 contents, and the pH is adjusted to values equivalents as
noted above; immediately after they are sterilized,
refrigerated at 4°-6°C. and stored in standardization tanks,
the vitamins, minerals and the nucleoside and/or nucleotide
solutions are added: afterwards the products are reheated at
10 30°-70°C., packed in the polyethylene or glass bottles, and
sterilized in a continuous sterilizer at 120°-121°C. for 10
minutes.

In the case of powder products, after the phase of
concentrated solids recombination, preheating, deaeration,
15 fat mixture injection, homogenization, refrigeration final
pH standardization, concentration and addition of vitamins,
minerals nucleosides and/or nucleotides, the mixture is
reheated to 65°-70°C., homogenized at 100-150 Kg/cm² and
dried in a spray drier. Afterwards, the powdered product is
20 packed in polyethylene-aluminium containers or in cans,
internally coated with varnish, under inert atmosphere, or
in other acceptable containers.

25

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1 to 3 are block diagrams illustrating the stages
of the process for the preparation of the products of the

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1 to 3 are block diagrams illustrating the stages of the process for the preparation of the products of the invention when packaged aseptically, in bottles or in powder form.

FIG. 4 is a schematic view of the plant manufacturing process for preparing products of the present invention.

With reference to FIG 1, there is depicted a process for the preparation and aseptic packaging of a liquid infant formula or a liquid nutritionally balanced diet formulation comprising a composition as set forth above, said process comprising the steps of:

(a) mixing water and non-fat solids of said composition in the absence of any vitamins, nucleosides and nucleotides;

(b) preheating the mixture to a temperature ranging from about 75^o to about 80^o C. and then deaerating the heated mixture followed by adding any fats in the composition to the deaerated mixture;

(c) homogenizing the mixture under pressure followed by cooling;

(d) standardizing the mixture by addition of any vitamins, minerals, nucleosides, nucleotides and other components not added in step (a), and adjusting the pH in the range of about 6.8 to about 7.1;

(e) UHT sterilizing the standardized mixture and subsequently homogenizing the mixture under pressure;

(f) cooling the homogenized mixture and aseptically packaging same.

5 With reference to FIG. 2, there is depicted a process for the preparation and bottling of a liquid infant formula or a liquid nutritionally balanced diet formulation comprising a composition as set forth above, said process comprising the steps of:

10 (a) mixing water and non-fat solids of said composition in the absence of any vitamins, nucleosides and nucleotides;

(b) preheating the mixture to a temperature ranging from about 75° to about 85° C. and then deaerating the heated mixture followed by adding any fats in the composition to the deaerated mixture;

15 (c) homogenizing the mixture under pressure followed by cooling;

(d) adjust the pH of the mixture in the range of about 6.8 to about 7.1, followed by UHT sterilization;

20 (e) standardizing the UHT sterilized mixture by the addition of any vitamins, minerals, nucleosides, nucleotides and other components not added in step (a) and reheating the standardized mixture to a temperature ranging from about 30° to about 70° C. and bottling the heated mixture;

25 (d) sterilizing the bottled mixture a second time to obtain a final bottled product in liquid form.

With reference to FIG 3, there is depicted a process for the preparation in powder form of an infant formula or a nutritionally balanced diet formulation comprising a

composition as set forth above said process comprising the steps of:

(a) mixing water and non-fat solids of said composition in the absence of any vitamins, nucleosides and nucleotides;

5 (b) preheating the mixture to a temperature ranging from about 75° to about 80° C. and then deaerating the heated mixture followed by adding any fats in the composition to the deaerated mixture;

10 (c) homogenizing the mixture under pressure followed by cooling;

(d) standardizing the mixture by addition of any vitamins, minerals, nucleosides, nucleotides and other components not added in step (a), and adjusting the pH in the range of about 6.8 to about 7.1;

15 (e) reheating the standardized mixture to between about 65° C. and 70° C.

(f) homogenizing the reheated mixture and drying the homogenized mixture in a spray drier to obtain a final dry powder products; and

20 (g) packaging the dry powder product.

Referring to FIG. 4, the process is described in more detail below:

Through the plate heat exchanger 1, deionized water is fed to storage tanks 2, at a temperature between 60°-70° C. Through the centrifugal pump 3 and triblender 4 non fat solids (protein, carbohydrates and some minerals) are dissolved being maintained the temperature at 60°-65° C. by means of a plate heat exchanger 5.

The resulting mixture is fed through positive pump 6 to filters 7 and plate heat exchanger 8 to be heated to 75°-80° C. for 15-20 s to get the product pasteurized; being immediately deaerated in a vacuum deaerator 9, lowering the temperature to 70°-75° C. Afterwards, the deaerated product is fed through centrifugal pump 10 and mixed with fat through fat injector 15. The mixture of fats stored in tank 12 has been fed through positive pump 13 to plate heat exchanger 14 to be heated at 70°-75° C. before reaching fat injector 15. A retention valve 11 prevents the product which contains the non fat solids and fat to go back to the deaerator.

Immediately after fats are mixed to the non fat solids mixture, the product is homogenized at 16 and a temperature of about 70°-75° C. and 200-300 Kg/cm² of total pressure, preferably in two stages (1st 150-200 Kg/cm², 2nd 50-100 kg/cm²).

For liquid products which are to be aseptically packaged, after homogenization 16, they are cooled to 4° to 6° C. in plate heat exchanger 17 and fed to isothermal standardizing tanks 18 where the pH is adjusted to from about 6.8 to about 7.1 depending on the product desired. Vitamins, minerals, nucleotides and/or nucleosides in the required amounts are fed to 18 and the resulting mixture is fed by pumps 19 and 20 to a UHT sterilizer 21 which is at 145°-150° C., homogenized in 22 (preferably in a double stage at 80° C. and 200-250 Kg/cm²), then is cooled to 20° to 25° in heat exchanger and aseptically packaged in 36. The

packing is preferably brick type of cardboard, paperboard, aluminium and polyethylene.

For liquid products which are to be bottled, the process is the same as above through the cooling treatment 17. Then the pH is adjusted in tanks 18 to above noted values. The mixture is fed by pumps 19 and 20 for UHT sterilization at 21 and homogenization at 22. The sterilized mixture is cooled 23 and fed directly to standardizing tanks 32 where vitamins, minerals nucleosides and/or nucleotides as required are added. From tanks 32, and by means of pump 33, the mixture is fed to reheater 34 where the temperature is raised to about 30° C. (for polyethylene bottles) to 70 C. (for glass bottles). The mixture is bottled in a filling machine 35 and subjected to sterilization 37 at a temperature of about 120°-121° C. for about 10 to 15 minutes to give a bottled product.

For powder products, the process is the same as first described above through homogenization 16. As shown by the dotted line, the product is fed to heat exchanger 24 and cooled to about 4° about 6° C. and fed to isothermal standardizing tanks 25, where the pH is adjusted and the required vitamins, minerals, nucleosides and/or nucleotides are added. Then the standardized product is pumped 26 through filters 27 and fed to reheater 28 where the temperature is raised to about 65° to about 70° C., and then finally filtered 29 and homogenized 30 under a pressure of about 100 -200 Kg/cm². The homogenized products products is fed to a drying tower 31, spray dried and collected for packaging.

EXAMPLES

The invention is illustrated with the following examples, which are not to be constructed as limiting the scope of the invention.

5

EXAMPLE I

This example provides a product made to feed preterm and low-birth weight infants, enriched with nucleosides and/or nucleotides and according to the invention. Basically, the product is a mixture of cow's milk, demineralized serum protein, dextrinmaltose, fat mixture, mineral, vitamins and nucleosides and/or nucleotides mixtures.

The product has been adapted in the protein, fat, carbohydrates, minerals and vitamins contents to the ESP-GAN and AAP international recommendations as related to the feeding of low-birth weight infants (ESP-GAN, Committee on Nutrition, Acta Paediatr, Scand., 1987 (in press); AAP, Committee on Nutrition, Pediatrics, 1985).

TABLE IV

20

		For 100 g of powder	For 100 ml of liquid	
Ingredients:				
Water	----		85	%
Maltodextrines	28.91	%	4.33	%
25 Vegetable oils mixture	20.23	%	3.03	%
Skim milk (0.05% fat)	14.58	%	2.19	%
Lactalbumin	12.13	%	1.82	%
Lactose	11.92	%	1.79	%
Butterfat	6.45	%	0.97	%
Minerals	3.26	%	0.49	%
Calcium caseinate	1.97	%	0.296	%
Lecithin	0.41	%	0.061	%
Vitamins	0.12	%	0.018	%
Nucleosides and/or nucleotides	0.0078	%	0.0012	%
Ascorbile palmitate	0.006	%	0.0009	%
DL- α Tocopherol	0.001	%	0.0001	%

Nucleosides and/or nucleotides added:

	Uridine and/or uridine monophosphate	3.42 mg	515 µg
	Guanosine and/or guanosine monophosphate	1.49 mg	225 µg
5	Adenosine and/or adenosine monophosphate	1.32 mg	200 µg
	Cytidine and/or cytidine monophosphate	1.12 mg	170 µg
	Inosine and/or inosine monophosphate	0.45 mg	70 µg

Mineral salts added:

10	Calcium lactate	1.74 g	0.26 g
	Sodium phosphate dibasic	0.65 g	97 mg
	Calcium phosphate	0.36 g	54 mg
	Potassium chloride	0.23 g	34 mg
	Potassium phosphate dibasic	0.17 g	26 mg
	Ferrous lactate	51.7 mg	7.6 mg
	Magnesium sulfate	49 mg	7.3 mg
15	Zinc sulfate	7.3 mg	1.1 mg
	Cupric sulfate	1.9 mg	285 µg
	Sodium fluoride	1.5 mg	225 µg
	Potassium and chromium sulfate	510 µg	76 µg
	Sodium molybdate	265 µg	40 µg
	Sodium selenite	180 µg	27 µg
	Manganese sulfate	83 µg	12 µg
20	Potassium iodine	64 µg	10 µg

Vitamins added:

	Vitamin A	1.600 UI	240 UI
	Vitamin D	600 UI	90 UI
25	Vitamin E	5.5 mg	825 µg
	Vitamin K	60 µg	9 µg
	Tiamin	0.4 mg	60 µg
	Riboflavin	0.45 mg	67 µg
	Piridoxin	0.25 mg	37 µg
	Niacin	6.7 mg	1 mg
	Calcium pantothenate	5.5 mg	825 µg
	Vitamin B12	1.1 µg	0.16 µg
	Biotin	15 µg	2.2 µg
	Folic acid	350 µg	52 µg
	Vitamin C	100 mg	15 mg

EXAMPLE II

This example provides a milk formula made to feed at-term infants, during the first year of life, preferably for the 6 first months of lactation, supplemented with nucleosides and/or nucleotides in similar concentrations to those of human milk, according to the invention.

The product has been adapted in its composition and content of nutrients to the ESPGAN and AAP international recommendations for this kind of infants (ESPGAN. Committee on Nutrition, Acta Paediatr. Scand., suppl. 262, 1977; AAP Committee on Nutrition, Pediatric Nutrition Handbook, 1979)

15

TABLE V

EXAMPLE II

ADAPTED INFANT MILK FORMULA

Ingredients	For 100 g of powder	For 100 ml of liquid
Water	—	87%
Lactose	42.61%	5.54%
20 Powdered milk (26% M.G.)	25.47%	3.31%
Vegetable oils	13.37%	1.74%
Demineralized whey (65% of proteins)	9.28%	1.21%
Butterfat	7.77%	1.01%
Minerals salts*	1.11%	0.14%
Lecithin	0.31%	0.04%
Vitamins**	0.069%	0.009%
Nucleosides and/or nucleotides***	0.0078%	0.001%
25 DL- α Tocopherol	0.003%	0.0004%
Ascorbyle palmitate	0.001%	0.0001%

	<u>*Mineral salts added</u>	
	Tripotassium citrate	0.35 g 45 mg
	Tricalcium citrate	0.31 g 40 mg
	Dibasic potassium phosphate	0.24 g 31 mg
5	Calcium chloride	0.16 g 21 mg
	Ferrous lactate	39 mg 5.1 mg
	Zinc acetate	8.5 mg 1.1 mg
	Cupric sulfate	1.10 mg 143 µg
	Manganese sulfate	155 µg 20 µg
	Potassium iodine	65 µg 8.4 µg
	<u>**Vitamins added</u>	
10	Vitamin C	50 mg 6.5 mg
	Nicotinamide	6.7 mg 870 µg
	Vitamin E	5.5 mg 715 µg
	Calcium pantothenate	5.5 mg 715 µg
	Vitamin A	1,600 UI 208 UI
	Vitamin B ₂	450 µg 58 µg
	Vitamin B ₁	400 µg 52 µg
	Vitamin B ₆	300 µg 39 µg
15	Vitamin K ₁	60 µg 7.8 µg
	Folic acid	25 µg 3.2 µg
	Biotin	15 µg 1.9 µg
	Vitamin D ₃	300 UI 39 UI
	Vitamin B ₁₂	1.1 µg 0.14 µg
	<u>***Nucleosides and/or nucleotides added</u>	
20	Uridine and/or uridine monophosphate	3.42 mg 445 µg
	Guanosine and/or guanosine monophosphate	1.49 mg 195 µg
	Adenosine and/or adenosine monophosphate	1.32 mg 170 µg
	Cytidine and/or cytidine monophosphate	1.12 mg 145 µg
25	Inosine and/or inosine monophosphate	0.45 mg 58 µg

EXAMPLE III

This example provides an infant milk formula made to feed healthy infant from 4-5 months to one year of life,

supplemented with nucleosides and/or nucleotides, according to the invention. The product has been adapted in its composition content of nutrients to the ESPGAN recommendations for these infants ESPGAN Committee on Nutrition, Acta Paediatr. Scand. suppl. 287, 1981).

TABLE VI
EXAMPLE III
INFANT ADAPTED FOLLOW-UP MILK FORMULA

Ingredients	For 100 g of powder	For 100 ml of liquid
Water	—	85%
Full milk	46.61%	6.99%
Maltodextrines	23.18%	3.48%
Lactose	19.28%	2.89%
Vegetable oils	6.08%	0.91%
Demineralized whey	4.22%	0.63%
Mineral salts*	0.41%	0.061%
Lecithin	0.14%	0.021%
Vitamins**	0.069%	0.01%
Nucleosides and/or nucleotides***	0.0078%	0.0012%
DL- α Tocopherol	0.003%	0.0004%
Ascorbyle palmitate	0.001%	0.0001%
<u>*Mineral salts added</u>		
Monocalcium phosphate	0.36 g	54 mg
Ferrous lactate	39 mg	5.8 mg
Zinc acetate	8.5 mg	1.3 mg
Cupric sulfate	1.1 mg	165 μ g
Manganese sulfate	155 μ g	23 μ g
Potassium iodine	65 μ g	9.7 μ g
<u>**Vitamins added</u>		
As in Example II.		
<u>***Nucleosides and/or nucleotides added</u>		
Uridine and/or uridine monophosphate	3.42 mg	515 μ g
Guanosine and/or guanosine monophosphate	1.49 mg	225 μ g
Adenosine and/or adenosine monophosphate	1.32 mg	200 μ g
Cytidine and/or cytidine monophosphate	1.12 mg	170 μ g
Inosine and/or inosine monophosphate	0.45 mg	70 μ g

EXAMPLE IV

This example provides a lactose free infant formula containing protein from milk origin, supplemented with nucleosides and/or nucleotides to the same quantities as in human milk, according to this invention.

The product has been adapted in its composition and content of nutrients to the international recommendations mentioned before.

TABLE VII

LACTOSE FREE ADAPTED INFANT MILK FORMULA
CONTAINING COW'S PROTEIN

	For 100 g of powder	For 100 ml of liquid
<u>Ingredients</u>		
Water	—	85%
Dextrinomaltose	58.03%	2.7%
Calcium caseinate (supplemented with L-cystine)	16.7%	2.51%
Butterfat	11.96%	1.79%
Vegetable oils	10.35%	1.55%
Mineral salts*	2.18%	0.33%
Lecithin	0.69%	0.103%
Vitamins**	0.069%	0.01%
Carnitine	0.0089%	0.0013%
Nucleosides and/or nucleotides***	0.0078%	0.0012%
DL-α-tocopherol	0.003%	0.0004%
Ascorbide palmitate	0.001%	0.0001%

*Mineral salts added

	Dibasic potassium phosphate	588 mg	88 mg
	Tripotassium citrate	522 mg	78 mg
	Calcium lactate	272 mg	41 mg
5	Sodium chloride	389 mg	58 mg
	Magnesium chloride	260 mg	39 mg
	Calcium chloride	90 mg	13.5 mg
	Ferrous lactate	48 mg	7.2 mg
	Zinc acetate	11.2 mg	1.7 mg
	Cupric sulfate	1.15 mg	0.17 mg
	Manganese sulfate	107 µg	16 µg
10	Potassium iodine	65 µg	9.7 µg

**Vitamins added

As in Example II.

***Nucleosides and/or nucleotides and other substances added

	Uridine and/or uridine monophosphate	3.42 mg	515 µg
15	Guanosine and/or guanosine monophosphate	1.49 mg	225 µg
	Adenosine and/or adenosine monophosphate	3.32 mg	500 µg
	Cytidine and/or cytidine monophosphate	4.98 mg	750 µg
	Inosine and/or inosine monophosphate	1.00 mg	150 µg
20	L-cystine	0.1 g	15 µg
	Carnitine	2.9 mg	1.3 mg

EXAMPLE V

This example provides a lactose free adapted infant formula containing a protein isolate from vegetal origin, supplemented with nucleosides and/or nucleotides, according to this invention.

The product has been adapted, as in example IV, in its composition and content of nutrients to the suckling children and newborns.

TABLE VIII

LACTOSE-FREE ADAPTED INFANT FORMULA
CONTAINING VEGETAL PROTEIN

Ingredients	For 100 g of powder	For 100 ml of liquid
Water	—	85%
Dextrinemaltose	57.20%	8.58%
Soy protein isolate	16.67%	2.5%
Butterfat	11.96%	1.79%
Vegetable oils	10.35%	1.55%
Mineral salts*	3.04%	0.46%
Lecithin	0.69%	0.103%
Vitamins**	0.069%	0.01%
Carnitine	0.0089%	0.0013%
Nucleosides and/or nucleotides***	0.0072%	0.0012%
DL-ascorophorol	0.003%	0.0004%
Ascorbale palmitate	0.001%	0.0001%
<u>*Mineral salts added</u>		
Dibasic potassium phosphate	450 mg	67.5 mg
Tripotassium citrate	628 mg	94.2 mg
Calcium chloride	370 mg	55.5 mg
Calcium carbonate	400 mg	60 mg
Magnesium chloride	260 mg	39 mg
Calcium lactate	873 mg	131 mg
Ferrous lactate	48 mg	7.2 mg
Zinc acetate	11.2 mg	1.7 mg
Cupric sulfate	1.18 mg	0.18 mg
Magnesium sulfate	107 µg	25 µg
Potassium iodine	65 µg	9.7 µg
<u>**Vitamins added</u>		
As in Example II.		
<u>***Nucleosides and/or nucleotides and other substances added</u>		
Uridine and/or uridine monophosphate	3.42 mg	515 µg
Guanosine and/or guanosine monophosphate	1.49 mg	225 µg
Adenosine and/or adenosine monophosphate	3.32 mg	500 µg
Cytidine and/or cytidine monophosphate	4.98 mg	750 µg
Inosine monophosphate or inosine	1.00 mg	150 µg
Carnitine	8.9 mg	1.3 µg

EXAMPLE VI

This example provides a lactose-free infant formula which contains a mixture of lactalbumin and casein hydrolyzates with a low molecular weight, supplemented with nucleosides and/or nucleotides, as specified in the invention.

The composition and content of nutrients are adapted to the suckling children and newborns requirements, as in examples IV and V.

TABLE IX

HYPOALLERGENIC ADAPTED INFANT FORMULA

	For 100 g of powder	For 100 ml of liquid
<u>Ingredients</u>		
Water	—	85%
Maltodextrines	52.48%	7.87%
Vegetable oil mixtures	16.98%	2.55%
Lactalbumin enzymatic hydrolyzate	12.31%	1.85%
Casein enzymatic hydrolyzate	5.16%	0.77%
Corn starch	4.87%	0.73%
Butterfat	4.29%	0.64%
Minerals**	3.19%	0.48%
Emulsifier	0.60%	0.09%
Vitamins***	0.069%	0.01%
Lecithin	0.0231%	0.0035%
Carnitine	0.0089%	0.0013%
Nucleosides and/or nucleotides*	0.0078%	0.0012%
DL- α -Tocopherol	0.0038%	0.0006%
Ascorbyle palmitate	0.0015%	0.0002%
<u>*Nucleosides and/or nucleotides and other substances added</u>		
Uridine and/or uridine monophosphate	3.42 mg	515 μ g
Guanosine and/or guanosine monophosphate	1.49 mg	225 μ g
Adenosine and/or adenosine monophosphate	3.32 mg	500 μ g
Cytidine and/or cytidine monophosphate	4.98 mg	750 μ g
Inosine and/or inosine monophosphate	1.00 mg	150 μ g
Carnitine	8.9 mg	1.3 mg
<u>**Mineral salts added</u>		

	Calcium phosphate	0.86 g	0.13 g
	Tripotassium citrate	0.25 g	0.13 g
	Calcium chloride	0.49 g	73 mg
	Sodium phosphate dibasic	0.38 g	57 mg
	Potassium chloride	0.24 g	36 mg
	Magnesium sulfate	0.20 g	30 mg
	Potassium phosphate dibasic	0.12 g	18 mg
5	Ferrous lactate	39 mg	5.8 mg
	Zinc acetate	10 mg	1.5 mg
	Cupric sulfate	2.2 mg	330 µg
	Sodium fluoride	310 µg	46.5 µg
	Manganese sulfate	307 µg	46 µg
	Potassium chromium sulfate	115 µg	17 µg
	Sodium molybdate	83 µg	12 µg
	Potassium iodine	65 µg	9.7 µg
	Sodium selenite	37 µg	5.5 µg
	<u>***Vitamins added</u>		
10	As in Example II.		

The products in examples IV, V and VI contain carnitine in similar concentration to that found in human milk, to satisfy the newborns requirements of this compound.

The products in examples I to VI are presented as liquid products, ready to use, as liquid concentrate products, to be used with the addition of water and as powdered products.

EXAMPLE VII

Example VII provides a complete product and nutritionally balanced to be used in clinical nutrition orally or by feeding tubes, with an energy ratio of 146 Kcal/g nitrogen, enriched with nucleosides and/or nucleotides in agreement with the invention.

The composition and content of nutrients have been adapted to the specific nutritional requirements of ill adults suffering energy-protein malnutrition.

TABLE X
NORMOPROTEIN DIET FOR USE IN CLINICAL
NUTRITION

	For 100 g of powder	For 100 ml of liquid
<u>COMPOSITION</u>		
5	—	78.7%
	Water	52.13%
	Maltodextrines	12.1%
	Vegetable oils	11.63%
	Lactalbumin	10.05%
	Calcium caseinate	8.84%
	Butterfat	3.79%
	Minerals*	0.75%
	Nucleosides and/or nucleotides***	0.66%
	Soy lecithin	—
	Emulsifier	0.136%
10	Stabilizer	—
	Vitamins**	0.026%
	Ascorbile palmitate	0.0232%
	DL- α -Tocopherol	0.0008%
	*Mineral salts added	
	Sodium phosphate dibasic	1.1 g
	Potassium chloride	0.99 g
	Magnesium sulfate	1.014 g
	Calcium chloride	0.31 g
	Potassium phosphate dibasic	0.28 g
15	Sodium chloride	50 mg
	Ferrous lactate	21 mg
	Zinc acetate	14 mg
	Manganese sulfate	4 mg
	Cupric sulfate	3 mg
	Sodium fluoride	2.2 mg
	Potassium chromium sulfate	480 μ g
	Sodium molybdate	315 μ g
	Sodium selenite	166 μ g
	Potassium iodine	49 μ g
	**Vitamins added	
20	Vitamin C	15 mg
	Niacin	4.75 mg
	Vitamin E (mg, TE)	2.5 mg
	Pantothenic acid	1.75 mg
	Vitamin B ₆	550 μ g
	Vitamin B ₂	425 μ g
	Vitamin B ₁	375 μ g
	Vitamin A (mg Re)	250 μ g
	Folate	100 μ g
	Biotin	50 μ g
25	Vitamin K ₁	35 μ g
	Vitamin D	2.5 μ g
	Vitamin B ₁₂	0.75 μ g
	***Nucleosides and/or nucleotides added	
	Adenosine and/or adenosine monophosphate	150 mg
	Cytidine and/or cytidine monophosphate	150 mg
	Guanosine and/or guanosine monophosphate	150 mg
	Uridine and/or uridine monophosphate	150 mg
	Inosine and/or inosine monophosphate	150 mg

EXAMPLE VIII

This example provides a complete product and nutritionally balanced high protein content (91 Kcal/g nitrogen), enriched with nucleosides and/or nucleotides in agreement with the invention.

The composition and content of nutrients have been adapted to meet the specific nutritional requirements of ill adults in hypercatabolic state.

TABLE XI

HYPERPROTEIN DIET FOR USE IN CLINICAL NUTRITION		
	For 100 g of powder	For 100 ml of liquid
COMPOSITION		
HYPERPROTEIN DIET FOR USE IN CLINICAL NUTRITION		
	For 100 g of powder	For 100 ml of liquid
	—	77.28%
Water	50.6%	11.49%
Maltodextrines	15.96%	3.64%
Lactalbumin	13.08%	3.14%
Calcium caseinate	9.13%	1.99%
Vegetable oils	6.52%	1.49%
Butterfat	3.41%	0.68%
Minerals*	0.75%	0.15%
Nucleosides and/or nucleotides***	0.5%	—
Soy lecithin	—	0.11%
Emulsifier	—	0.02%
Stabilizer	0.026%	0.005%
Vitamins**	0.0232	0.0008%
Ascorbic palmitate	0.0008	0.0002%
DL- α -Tocopherol		
*Mineral salts added		
Potassium chloride	1.01 g	220 mg
Magnesium sulfate	1.014 g	203 mg
Sodium phosphate dibasic	0.88 g	180 mg
Potassium phosphate dibasic	0.19 g	30 mg
Sodium chloride	0.16 g	40 mg
Calcium chloride	0.11 g	5 mg
Ferrous lactate	21 mg	4 mg
Zinc acetate	14 mg	3 mg
Manganese sulfate	4 mg	760 μ g
Cupric sulfate	3 mg	640 μ g
Sodium fluoride	2.2 mg	442 μ g
Potassium chromium sulfate	480 μ g	96 μ g
Sodium molybdate	315 μ g	63 μ g
Sodium selenite	166 μ g	33 μ g
Potassium iodine	49 μ g	10 μ g
**Vitamins added		
As in Example VII		
***Nucleosides and/or nucleotides added		
Adenosine and/or adenosine monophosphate	150 mg	30 mg
Cytidine and/or cytidine monophosphate	150 mg	30 mg
Guanosine and/or guanosine monophosphate	150 mg	30 mg
Uridine and/or uridine monophosphate	150 mg	30 mg
Inosine and/or inosine monophosphate	150 mg	30 mg

EXAMPLE IX

This example provides a complete product and nutritionally balanced, with a high nitrogen contents, using as source of this element a protein hydrolyzate with a low molecular weight to make easier its absorption, enriched with nucleosides and/or nucleotides according to the invention. The energy ratio this product is 100 Kcal/g nitrogen.

This composition and content of nutrients have been adapted to satisfy specific nutritional requirements of ill adults suffering diverse malabsorption-malnutrition syndromes.

TABLE XII
EXAMPLE IX
PEPTIDE-MCT DIET FOR USE IN
CLINICAL NUTRITION

COMPOSITION	For 100 g of powder	For 100 ml of liquid
Water	—	77.83%
Maltodextrines	51.62%	11.43%
Casein hydrolyzate	25.80%	5.72%
Vegetable oils	12.44%	2.76%
Minerals*	5.02%	1.11%
Butterfat	3.62%	0.8%
Nucleosides and/or nucleotides***	0.75%	0.17%
Soy lecithin	0.50%	—
Emulsifier	—	0.11%
L-Cistine	0.20%	0.04%
Stabilizer	—	0.02%
Vitamins**	0.026%	0.0058%
Ascorbile palmitate	0.0232%	0.0051%
DL- α -Tocopherol	0.0008%	0.0002%
Sodium phosphate dibasic	1.05 g	233 mg
Magnesium sulfate	1.014 g	225 mg
Calcium chloride	0.84 g	186 mg
Tripotassium citrate	0.83 g	184 mg
Potassium phosphate dibasic	0.80 g	177 mg
Sodium chloride	0.44 g	97 mg
Ferrous lactate	21 mg	4.6 mg
Zinc acetate	14 mg	3.1 mg
Manganese sulfate	4 mg	888 μ g
Cupric sulfate	3 mg	665 μ g
Sodium fluoride	2.2 mg	488 μ g

Potassium chromium sulfate	430 µg	106 µg
Sodium molybdate	315 µg	70 µg
Sodium selenite	166 µg	37 µg
Potassium iodine	49 µg	11 µg
Adenosine and/or adenosine mono-phosphate	150 mg	30 mg
Cytidine and/or cytidine mono-phosphate	150 mg	30 mg
Guanosine and/or guanosine mono-phosphate	150 mg	30 mg
Uridine and/or uridine mono-phosphate	150 mg	30 mg
Inosine and/or inosine mono-phosphate	150 mg	30 mg
L-Cistine	200 mg	40 mg

5

*Mineral salts added

**Vitamins added

As in Example VII.

***Nucleosides and/or nucleotides and other substances added.

10

EXAMPLE X

This example provides a complete product and nutritionally balanced with a low protein content, supplemented with branched chain amino acids and enriched with nucleosides and/or nucleotides.

15

The composition and content of nutrients have been adapted to satisfy the specific nutritional requirements of ill adults suffering severe hepatopathy.

20

TABLE XIII

EXAMPLE X COMPLETE DIET FOR USE IN CLINICAL NUTRITION OF PATIENTS WITH LIVER DISEASE

COMPOSITION	For 100 g of powder	For 100 ml of liquid
Water	—	76.36%
Maltodextrines	72.13%	17.04%
Vegetable oils	7.48%	1.77%
Lactalbumin	7.26%	1.72%
Calcium caseinate	6.27%	1.48%
Minerals*	2.94%	0.69%
L-Leucine	1.16%	0.27%
L-Valine	0.87%	0.21%
L-Isoleucine	0.87%	0.21%
Nucleosides and/or nucleotides***	0.75%	0.18%
Soy lecithin	0.22%	—
Emulsifier	—	0.05%
Stabilizer	—	0.01%
Vitamins**	0.026%	0.006%
Ascorbic palmitate	0.0197%	0.005%

25

	DL- α -Tocopherol	0.0003%	0.00007%
	Magnesium sulfate	1.014 g	240 mg
	Potassium phosphate dibasic	0.67 g	158 mg
	Sodium phosphate dibasic	0.60 g	142 mg
5	Calcium chloride	0.49 g	116 mg
	Sodium chloride	0.12 mg	28 mg
	Ferrous lactate	21 mg	5 mg
	Zinc acetate	14 mg	3.3 mg
	Manganese sulfate	4 mg	946 μ g
	Cupric sulfate	3 mg	709 μ g
	Sodium fluoride	2.2 mg	520 μ g
	Potassium chromium sulfate	480 μ g	113 μ g
	Sodium molybdate	315 μ g	74 μ g
	Sodium selenite	166 μ g	39 μ g
10	Potassium iodine	49 μ g	11.6 μ g
	Adenosine and/or adenosine monophosphate	150 mg	30 mg
	Cytidine and/or cytidine mono- phosphate	150 mg	30 mg
	Guanosine and/or guanosine monophosphate	150 mg	30 mg
	Uridine and/or uridine mono- phosphate	150 mg	30 mg
	Inosine and/or inosine mono- phosphate	150 mg	30 mg
15	L-Leucine	1.16 g	274 mg
	L-Valine	870 mg	206 mg
	L-Isoleucine	870 mg	206 mg

*Mineral salts added

**Vitamins added

As in Example VII.

***Nucleosides and/or nucleotides and other substances added

20

25

EXAMPLE XI

This example provides a product considered as a nutritional supplement for the nutritional repletion of patients with chronic hepatopathy, constituted by a mixture of proteins from milk origin, supplemented with branched chain amino acids, carbohydrates, vitamins and minerals and enriched with nucleosides and/or nucleotides.

TABLE XIV

EXAMPLE XI
HYPERPROTENC DIET SUPPLEMENTED WITH
BRANCHED CHAIN AMINO ACIDS FOR USE IN
CLINICAL NUTRITION OF PATIENTS WITH
LIVER DISEASE

10

15

20

25

COMPOSITION	For 100 g of powder	For 100 ml of liquid
Water	—	80%
Maltodextrines	36.72%	7.32%
Lactalbumin	26.26%	5.25%
Sodium caseinate	21.95%	4.39%
L-Leucine	4.04%	0.81%
Minerals*	3.2%	0.64%
L-Valine	3.03%	0.61%
L-Isoleucine	3.03%	0.61%
Nucleosides and/or nucleotides***	0.75%	0.15%
Vitamins**	0.026%	0.005%
Magnesium sulfate	1.014 g	203 mg
Potassium chloride	0.89 g	178 mg
Calcium chloride	0.38 g	76 mg
Sodium phosphate dibasic	0.36 g	72 mg
Tripotassium citrate	0.34 g	68 mg
Potassium phosphate dibasic	0.17 g	34 mg
Ferrous lactate	21 mg	4.2 mg
Zinc acetate	14 mg	2.8 mg
Manganese sulfate	4 mg	800 µg
Cupric sulfate	3 mg	600 µg
Sodium fluoride	2.2 mg	440 µg
Potassium chromium sulfate	480 µg	96 µg
Sodium molybdate	315 µg	63 µg
Sodium selenite	166 µg	33 µg
Potassium iodine	49 µg	9.8 µg
Adenosine and/or adenosine mono-phosphate	150 mg	30 mg

TABLE XIV-continued

	Cytidine and/or cytidine mono-phosphate	150 mg	30 mg
	Guanosine and/or guanosine mono-phosphate	150 mg	30 mg
5	Uridine and/or uridine monophosphate	150 mg	30 mg
	Inosine and/or inosine monophosphate	150 mg	30 mg
	L-Leucine	4.04 g	810 mg
	L-Valine	3.03 g	610 mg
	L-Isoleucine	3.03 g	610 mg

*Mineral salts added

**Vitamins added

As to Example VII.

***Nucleosides and/or nucleotides and other substances added

10 The invention having been thus described, it will be appreciated by those in the art that variations can occur within the scope of the claims which follow.

We claim:

15 1. A method for the stimulation or repair and regeneration of intestinal gut cells in infants which comprises administering to an infant in need of such stimulation an amount effective for said stimulation of an infant formula free of cow's milk, said infant formula comprising a therapeutically effective amount of
20 carbohydrates, a source of amino acids, vegetable oils, minerals, vitamins, and at least one member selected from the group consisting of uridine, uridine phosphate and mixtures thereof; guanosine, guanosine phosphate and mixtures thereof; adenosine, adenosine phosphate and
25 mixtures thereof; cytidine, cytidine phosphate and mixtures thereof and inosine, inosine phosphate and mixtures thereof.

 2. The method according to claim 1 wherein said infant formula comprises per dl:

5

uridine	0.2-60 mg.
guanosine	0.2-60 mg.
adenosine	0.2-60 mg.
cytidine	0.2-60 mg. and
inosine	0.2-60 mg.

3. The method according to claim 2 wherein said formula comprises:

10

uridine	10-50 mg.
guanosine	10-50 mg.
adenosine	10-50 mg.
cytidine	10-50 mg. and
inosine	10-50 mg.

15

4. A method for the stimulation or repair and regeneration of intestinal gut cells in infants which comprises administering to an infant in need of such stimulation an amount effective for said stimulation of an infant formula in liquid form which comprises a therapeutically effective amount of a source of amino nitrogen, carbohydrates, edible fats, minerals, vitamins and a nucleoside and/or nucleotide composition comprising at least one member selected from the group consisting of:

20

(a) uridine, uridine phosphate and a mixture thereof;

25

(b) guanosine, guanosine phosphate and a mixture thereof;

(c) adenosine, adenosine phosphate and a mixture thereof;

(d) cytidine, cytidine phosphate and a mixture thereof; and

(e) inosine, inosine phosphate and mixture thereof;

wherein the total amount combined nucleoside and nucleotide content is in the range of between about 10 to 250 mg for each 100 g of said formula.

5
10
15
20
5. A method for enhancing the immune response of T-cells and for providing specific fatty acid phospholipids profiles in red blood cell membranes of infants, which comprises administering to an infant in need of enhanced immune response of T-cells an effective amount for enhancing said response to an infant formula free of cow's milk, said infant formula comprising a therapeutically effective amount of carbohydrates, a source of amino acids, vegetable oils, minerals, vitamins and at least one member selected from the group consisting of uridine, uridine phosphate and mixtures thereof; guanosine, guanosine phosphate and mixtures thereof; adenosine, adenosine phosphate and mixtures thereof; cytidine, cytidine phosphate and mixtures thereof and inosine, inosine phosphate and mixtures thereof.

6. The method according to claim 5, wherein said infant formula comprises per dl;

25

uridine	0.2-60 mg
guanosine	0.2-60 mg
adenosine	0.2-60 mg
cytidine	0.2-60 mg; and
inosine	0.2-60 mg.

7. The method according to claim 6 wherein said formula comprises:

5	uridine	10-50 mg.
	guanosine	10-50 mg.
	adenosine	10-50 mg.
	cytidine	10-50 mg. and
	inosine	10-50 mg.

8. A method for enhancing the immune response of T-cells and for providing specific fatty acid phospholipids profiles in red blood cell membranes of infants, which comprises administering to an infant in need of enhanced immune response to T-cells an effective amount for enhancing said responses an infant formula in liquid form which comprises a therapeutically effective amount of a source of amino nitrogen, carbohydrates, edible fats, minerals, vitamins and a nucleoside and/or nucleotide composition comprising at least one member selected from the group consisting of:

- 20 (a) uridine, uridine phosphate and a mixture thereof;
- (b) guanosine, guanosine phosphate and a mixture thereof;
- (c) adenosine, adenosine phosphate and a mixture thereof;
- 25 (d) cytidine, cytidine phosphate and a mixture thereof; and
- (e) inosine, inosine phosphate and mixture thereof;
- wherein the total combined nucleoside and nucleotide

content is in the range of between about 10 to 250 mg for each 100 g of said formula.

9. A method for the stimulation or repair and regeneration of intestinal gut cells in humans which comprises administering in need of such stimulation an amount effective for said stimulation of a formula free of cow's milk, said formula comprising a therapeutically effective amount of carbohydrates, a source of amino acids, vegetable oils, minerals, vitamins, and at least one member selected from the group consisting of uridine phosphate and mixtures thereof; guanosine, guanosine phosphate and mixtures thereof; adenosine, adenosine phosphate and mixtures thereof; cytidine, cytidine phosphate and mixtures thereof and inosine, inosine phosphate and mixtures thereof.

10. The method according to claim 9 wherein said formula comprises per dl:

20

uridine	0.2-60 mg;
guanosine	0.2-60 mg;
adenosine	0.2-60 mg;
cytidine	0.2-60 mg; and
inosine	0.2-60 mg.

25

11. The method according to claim 10 wherein said formula comprises:

uridine	10-50 mg;
guanosine	10-50 mg;
adenosine	10-50 mg;
cytidine	10-50 mg; and
inosine	10-50 mg.

12. A method for the stimulation or repair and regeneration of intestinal gut cells in humans which comprises administering to a human in need of such stimulation an amount effective for said stimulation of a formula in liquid form which comprises a therapeutically effective amount of a source of amino nitrogen, carbohydrates, edible fats, minerals, vitamins and a nucleoside and/or nucleotide composition comprising at least one member selected from the group consisting of:

- 10 (a) uridine, uridine phosphate and a mixture thereof;
 - (b) guanosine, guanosine phosphate and a mixture thereof;
 - (c) adenosine, adenosine phosphate and a mixture thereof;
 - 15 (d) cytidine, cytidine phosphate and a mixture thereof; and
 - (e) inosine, inosine phosphate and a mixture thereof;
- wherein the total combined nucleoside and nucleotide content is in the range of between about 10 to 250 mg for each 100 g of said formula.

20 13. A method for enhancing the immune response of T-cells and for providing specific fatty acid phospholipids profiles in red blood cell membranes of humans, which comprises administering to a human in need of enhanced immune response of T-cells an effective amount for enhancing said response of a formula free of cow's milk, said formula comprising a therapeutically effective amount of carbohydrates, a source of amino acids, vegetable oils, minerals, vitamins and at least one member selected from the

group consisting of uridine, uridine phosphate and mixtures thereof; guanosine, guanosine phosphate and mixtures thereof; adenosine, adenosine phosphate and mixtures thereof; cytidine, cytidine phosphate and mixtures thereof and inosine, inosine phosphate and mixtures thereof.

14. The method according to claim 14 wherein said formula comprises per dl:

uridine	0.2-60 mg;
guanosine	0.2-60 mg;
adenosine	0.2-60 mg;
cytidine	0.2-60 mg; and
inosine	0.2-60 mg.

15. The method according to claim 14 wherein said formula comprises:

uridine	10-50 mg;
guanosine	10-50 mg;
adenosine	10-50 mg;
cytidine	10-50 mg; and
inosine	10-50 mg.

16. A method for enhancing the immune response of T-cells and for providing specific fatty acid phospholipids profiles in red blood cell membranes of humans, which comprises administering to a human in need of enhanced immune response of T-cells an effective amount for enhancing said response of a formula in liquid form which comprises a therapeutically effective amount of a source of amino nitrogen, carbohydrates, edible fats, minerals, vitamins and a nucleoside and/or nucleotide composition comprising at least one member selected from the group consisting of:

- (a) uridine, uridine phosphate and a mixture thereof;

(b) guanosine, guanosine phosphate and a mixture thereof;

(c) adenosine, adenosine phosphate and a mixture thereof;

5 (d) cytidine, cytidine phosphate and a mixture thereof; and

(e) inosine, inosine phosphate and a mixture thereof;

10 wherein the total combined nucleoside and a nucleotide content is in the range of between about 10 to 250 mg for each 100 g of said formula.

ANGEL H. GIL
DANIEL V. MORALES
EDUARDO VALVERDE
Inventors