



US008569031B2

(12) **United States Patent**
Novick et al.

(10) **Patent No.:** US 8,569,031 B2
(45) **Date of Patent:** Oct. 29, 2013

(54) **CHEMICALLY MODIFIED CARBONIC ANHYDRASES USEFUL IN CARBON CAPTURE SYSTEMS**

(71) Applicant: **Codexis, Inc.**, Redwood City, CA (US)

(72) Inventors: **Scott J. Novick**, Palo Alto, CA (US);
Oscar Alvizo, Fremont, CA (US)

(73) Assignee: **Codexis, Inc.**, Redwood City, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/721,648**

(22) Filed: **Dec. 20, 2012**

(65) **Prior Publication Data**

US 2013/0189756 A1 Jul. 25, 2013

Related U.S. Application Data

(63) Continuation of application No. 13/174,253, filed on Jun. 30, 2011, now Pat. No. 8,354,262.

(60) Provisional application No. 61/360,040, filed on Jun. 30, 2010, provisional application No. 61/445,996, filed on Feb. 23, 2011, provisional application No. 61/492,758, filed on Jun. 2, 2011.

(51) **Int. Cl.**
C12N 9/96 (2006.01)
C02F 3/34 (2006.01)

(52) **U.S. Cl.**
USPC **435/188**; 435/197; 435/232; 435/262;
435/266

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,627,638 A 12/1971 Barker et al.
3,666,733 A 5/1972 Epton
3,702,804 A 11/1972 Barker et al.
5,900,364 A 5/1999 Visuri
6,143,556 A 11/2000 Trachtenberg
6,475,382 B2 11/2002 Parent
6,524,843 B1 2/2003 Blais et al.
6,908,507 B2 6/2005 LaLande et al.
7,132,090 B2 11/2006 Dziedzic et al.
7,176,017 B2 2/2007 Parent et al.
7,267,971 B2 9/2007 Thakur et al.
7,514,056 B2 4/2009 Fradette et al.
7,521,217 B2 4/2009 Daigle et al.
7,579,185 B2 8/2009 Parent et al.
7,596,952 B2 10/2009 Fradette et al.
7,641,717 B2 1/2010 Gal
7,642,076 B2 1/2010 Dziedzic et al.
7,740,689 B2 6/2010 Fradette et al.
7,803,575 B2 9/2010 Borchert et al.
7,820,432 B2 10/2010 Parent et al.
7,862,788 B2 1/2011 Gal et al.
7,867,748 B2 1/2011 DiCosimo et al.
7,964,170 B2 6/2011 Sing

2004/0259231 A1 12/2004 Bhattacharya
2006/0048517 A1 3/2006 Fradette et al.
2006/0128004 A1 6/2006 Ancil et al.
2006/0213224 A1 9/2006 Fradette et al.
2006/0246564 A1 11/2006 Parent et al.
2006/0257990 A1 11/2006 Daigle et al.
2007/0004023 A1 1/2007 Trachtenberg
2007/0128713 A1 6/2007 Parent et al.
2007/0256559 A1 11/2007 Chen et al.
2008/0003662 A1 1/2008 Trachtenberg
2008/0148939 A1 6/2008 Fradette et al.
2008/0296231 A1 12/2008 Schoevaart et al.
2009/0148930 A1 6/2009 Gal et al.
2009/0155889 A1 6/2009 Handagama et al.
2009/0202409 A1 8/2009 Smith et al.
2009/0227010 A1 9/2009 Daigle et al.
2010/0011956 A1 1/2010 Neumann et al.
2010/0047866 A1 2/2010 Borchert et al.
2010/0068784 A1 3/2010 Dziedzic et al.
2010/0086983 A1 4/2010 Gellett et al.
2010/0196244 A1 8/2010 Grauer et al.
2010/0203619 A1 8/2010 Fradette et al.
2010/0209997 A1 8/2010 Newman et al.
2010/0236408 A1 9/2010 Aspiron et al.
2010/0294131 A1 11/2010 Bade et al.

(Continued)

FOREIGN PATENT DOCUMENTS

EP 1521629 B1 2/2007
EP 1627041 B1 11/2009

(Continued)

OTHER PUBLICATIONS

Husic, David H. "Chemical Cross-Linking of Periplasmic Carbonic anhydrase from *Chlamydomonas reinhardtii*" in Current Research in Photosynthesis, vol. IV (1990), M. Baltscheffsky (Ed.), pp. 493-496.*

Alber, B.E., et al., "A carbonic anhydrase from the archaeon *Methinsarcina thermophila*", Proc. Natl. Acad. Sci. USA, 91:6909-6913, 1994.

Alber, B.E., et al., "Characterization of Heterologously Produced Carbonic Anhydrase from *Methanosarcina thermophila*", Journal of Bacteriology, 178(11):3270-3274, 1996.

(Continued)

Primary Examiner — Nashaat Nashed

(74) *Attorney, Agent, or Firm* — Codexis, Inc.

(57) **ABSTRACT**

The present disclosure relates to chemically modified carbonic anhydrase polypeptides and soluble compositions, homogenous liquid formulations comprising them. The chemically modified carbonic anhydrase polypeptides have improved properties relative to the same carbonic anhydrase polypeptide that is not chemically modified including the improved properties of increased activity and/or stability in the presence of amine compounds, ammonia, or carbonate ion. The present disclosure also provides methods of preparing the chemically modified polypeptides and methods of using the chemically modified polypeptides for accelerating the absorption of carbon dioxide from a gas stream into a solution as well as for the release of the absorbed carbon dioxide for further treatment and/or sequestering.

18 Claims, 1 Drawing Sheet

(56)

References Cited

U.S. PATENT DOCUMENTS

2010/0297723 A1 11/2010 Borchert et al.
 2010/0300894 A1 12/2010 Lin et al.
 2011/0100216 A1 5/2011 Kozak et al.
 2011/0100217 A1 5/2011 Soloveichik et al.

FOREIGN PATENT DOCUMENTS

EP 2354098 A1 8/2011
 EP 2354099 A1 8/2011
 GB 1365886 9/1974
 WO 9013635 A1 11/1990
 WO 9105840 A1 5/1991
 WO 9500171 A1 1/1995
 WO 9640414 B1 12/1996
 WO 9744445 A1 11/1997
 WO 9955310 A1 11/1999
 WO 0022157 A1 4/2000
 WO 2001090321 A2 11/2001
 WO 02070646 A2 9/2002
 WO 03066850 A1 8/2003
 WO 2004028667 A1 7/2004
 WO 2004056455 A1 7/2004
 WO 2004081208 A1 9/2004
 WO 2005066341 A1 7/2005
 WO 2006022885 A1 3/2006
 WO 2006089423 A1 8/2006
 WO 2006108532 A1 10/2006
 WO 2007036235 A1 4/2007
 WO 2008008179 A2 1/2008
 WO 2008041921 A1 4/2008
 WO 2008072979 A1 6/2008
 WO 2008095057 A2 8/2008
 WO 2008099252 A1 8/2008
 WO 2008110405 A2 9/2008
 WO 2008/037846 A2 11/2008
 WO 2009000025 A1 12/2008
 WO 2009016642 A1 2/2009
 WO 2009036145 A1 3/2009
 WO 2009059104 A1 5/2009
 WO 2009073422 A1 6/2009
 WO 2009076327 A1 6/2009
 WO 2009087060 A2 7/2009
 WO 2009087210 A2 7/2009
 WO 2009105419 A2 8/2009
 WO 2009117550 A1 9/2009
 WO 2010014773 A1 2/2010
 WO 2010014774 A2 2/2010
 WO 2010020017 A1 2/2010
 WO 2010037109 A2 4/2010
 WO 2010043459 A1 4/2010
 WO 2010053683 A1 5/2010
 WO 2010081007 A2 7/2010
 WO 2010108974 A1 9/2010
 WO 2010151787 A1 12/2010
 WO 2011014955 A1 2/2011
 WO 2011014956 A1 2/2011
 WO 2011014957 A1 2/2011
 WO 2011054107 A1 5/2011
 WO 2011066304 A2 6/2011
 WO 2011069857 A1 6/2011

OTHER PUBLICATIONS

Alber, B.E., et al., "Kinetic and Spectroscopic Characterization of the Gamma-Carbonic Anhydrase from the Methanoarchaeon *Methanosarcina thermophila*," *Biochemistry*, 38, 13119-13128, 1999.
 Cowan, R.M., et al., "CO₂ capture by means of an enzyme-based reactor", *Ann. N.Y. Acad. Sci.*, 984:453-469, 2003.
 Darde, V., et al., "Chilled ammonia process for CO₂ capture", *International Journal of Greenhouse Gas Control*, 4:131-136, 2010.
 Elleuche, S., et al., "Evolution of carbonic anhydrases in fungi", *Curr. Genet.*, 55:211-222, 2009.
 Salmon, S., Holmes, J., Saunders, P. et al., "Ultrasound-assisted Regeneration for the CO₂ Capture Processes," IPCOM00192838D, IPCOM Prior Art Database, 2010.

Iverson, T.M., et al., "A Closer Look at the Active Site of γ -Class Carbonic Anhydrases: High-Resolution Crystallographic Studies of the Carbonic Anhydrase from *Methanosarcina thermophila*", *Biochemistry*, 39:9222-9231, 2000.
 Kisker, C., et al., "A left-handed β -helix revealed by the crystal structure of a carbonic anhydrase from the archaeon *Methanosarcina thermophila*", *The EMBO Journal*, 15(10):2323-2330, 1996.
 Smith, K.S., et al., "A Plant-Type (β -Class) Carbonic Anhydrase in the Thermophilic Methanoarchaeon *Methanobacterium thermoautotrophicum*", *Journal of Bacteriology*, 181(20):6247-6253, 1999.
 Smith, K.S., et al., "Carbonic anhydrase is an ancient enzyme widespread in prokaryotes", *PNAS*, 96 (26):15184-15189, 1999.
 Trachtenberg, M.C., et al., "Carbon Dioxide Transport by Proteic and Facilitated Transport Membranes", *Life Support & Biosphere Science*, 6:293-302, 1999.
 Tripp, B.C., et al., "A Structure-Function Study of a Proton Transport Pathway in the γ -Class Carbonic Anhydrase from *Methanosarcina thermophila*", *Biochemistry*, 39:9232-9240, 2000.
 Tripp, B.C., et al., "Role of Arginine 59 in the γ -Class Carbonic Anhydrases", *Biochemistry*, 41:669-678, 2002.
 Wilbur, K.M., et al., "Electrometric and Colorimetric Determination of Carbonic Anhydrase", *The Journal of Biological Chemistry*, pp. 147-154, 1948.
 Zimmerman, S.A., "Understanding the Biochemistry and Physiology of Gamma Carbonic Anhydrases in *Methanosarcina thermophila*", The Pennsylvania State University, The Graduate School of Eberly College of Science, pp. 1-161, 2007.
 Genbank Accession No. U08885, Birgit, E.A., "*Methanosarcina thermophila* TM-1 carbonic anhydrase gene, complete cds," 1994.
 Genbank Accession No. YP_306494.1, Maeder, D. L. et al., Carbonic Anhydrase [*Methanosarcina barkeri* str. Fusaro], 2010.
 Genbank Accession No. NP_635112.1, Deppenmeier, U. et al., Carbonic Anhydrase [*Methanosarcina mazei* Go1], 2010.
 Genbank Accession No. NP_617439.1, Galagan, J.E., et al., Carbonate Dehydratase [*Methanosarcina acetivorans* C2A], 2010.
 Booterabi, F., et al., "Modification of carbonic anhydrase II with acetaldehyde, the first metabolite of ethanol, leads to decreased enzyme activity," *BMC Biochemistry*, vol. 9, No. 32, 2008.
 Nagelhus, E., et al., Carbonic Anhydrase XIV is enriched in specific membrane domains of retinal pigment epithelium, Muller cells, and astrocytes, *PNAS*, vol. 102, No. 22, pp. 8030-8035, 2005.
 Ulmasov, B. et al., "Purification and kinetic analysis of recombinant CA XII, a membrane carbonic anhydrase overexpressed in certain cancers," *PNAS*, vol. 97, No. 26. pp. 14212-14217, 2000.
 Moya, A. et al. "Carbonic Anhydrase in the Scleractinian Coral *Stylophora pistillata*," *Journ. Biol. Chem.*, vol. 283, No. 37, pp. 25475-25484, 2008.
 Khodayari, A., "Experimental and Theoretical Study of Carbon Dioxide Absorption into Potassium Carbonate Solution Promoted with Enzyme," Thesis, University of Illinois, Urbana-Champaign, 2010.
 Bhattacharya, S. et al., "CO₂ hydration by immobilized carbonic anhydrase," *Biotechnol. Appl. Biochem.*, 38,111-117, 2003.
 Epton, R., "A Study of Carbonic Anhydrase Covalently bound to a Poly(acryloylmorpholine) Network in Aqueous/Organic Solvents," *BiochemSocTrans.*, vol. 5, 274-276, 1977.
 Epton, R., "Water-Soluble Coloured Covalent Conjugates of Carbonic Anhydrase and N-(sym-Trinitroaryl) polyacrylamide/Acrylylhydrazide Co-Polymers," *BiochemSocTrans.*, vol. 5, 277-279, 1977.
 Epton, R., "Soluble polymer-protein conjugates: 1. Reactive-N-(sym-trinitroaryl) polyacrylamide/acrylylhydrazide copolymers and derived carbonic anhydrase conjugates," *Polymer*, vol. 18, 319-323, 1977.
 Farmer, T., "Assessing the Multimeric States of Proteins: Studies Using Laser Desorption Mass Spectrometry," *BiologicalMassSpectrometry*, vol. 20, 796-800m 1991.
 Gitlin, I., "Peracetylated Bovine Carbonic Anhydrase (BCA-Ac18) is Kinetically More Stable than Native BCA to Sodium Dodecyl Sulfate," *J. Phys. Chem B*, 110, 2372-2377, 2006.

(56)

References Cited

OTHER PUBLICATIONS

Gudiksen, K., "Influence of the Zn(II) Cofactor on the Refolding of Bovine Carbonic Anhydrase after Denaturation with Sodium Dodecyl Sulfate," *Anal. Chem.*, vol. 76, No. 24, 7151-7161, 2004.

Gudiksen, K., "Eliminating Positively Charged Lysine ϵ -NH₃⁺ Groups on the Surface of Carbonic Anhydrase Has No Significant Influence on Its Folding from Sodium Dodecyl Sulfate," *J. Am. Chem. Soc.*, 127, 4707-4714, 2005.

Gudiksen, K., "Increasing the Net Charge and Decreasing the Hydrophobicity of Bovine Carbonic Anhydrase Decreases the Rate of Denaturation with Sodium Dodecyl Sulfate," *Biophysical J.*, vol. 91, 298-310, 2006.

Hyyppä, M., "The Effects of Fixations on Carbonic Anhydrase Activity," *Histochemie* 12, 184-188, 1968.

Krishnamurthy, V., "Carbonic Anhydrase as a Model for Biophysical and Physical-Organic Studies of Proteins and Protein-Ligand Binding," *Chem. Rev.* 108, 946-1051, 2008.

Yang, J., "Synthesis of Monodisperse Polymers from Proteins," *J. Am. Chem. Soc.*, 125, 12392-12393, 2003.


Kabadi, V.N., "Heat of Dissolution Measurements for CO₂ in Mixed Alkanolamine Solvents," OSTI Final Technical Report, Oct. 8, 2007. Genbank Accession No. YP_002436805 dated May 7, 2009.

Krebs, J. F., et al., "Determinants of Catalytic Activity and Stability of Carbonic Anhydrase II as Revealed by Random Mutagenesis," *J. Biol. Chemistry*, 268, 2, 948, 1993.

Lopez-Alonso, Jorge P., et al. "Carbodiimide EDC Induces Cross-Links That Stabilize RNase A C-Dimer against Dissociation: EDC Adducts Can Affect Protein Net Charge, Conformation, and Activity", *Bioconjugate Chem.* 20, 1459-1473, 2009.

Husic, David H., et al. "Effect of dithiothreitol on the catalytic activity, quaternary structure and sulfonamide-binding properties of an extracellular carbonic anhydrase from *Chlamydomonas reinhardtii*", *Biochimica et Biophysica Acta*, 1078, 35-42, 1991.

* cited by examiner



1

CHEMICALLY MODIFIED CARBONIC ANHYDRASES USEFUL IN CARBON CAPTURE SYSTEMS

1. STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

This invention was made with Government support under DE-AR000071 awarded by the Department of Energy. The Government has certain rights in this invention.

2. CROSS REFERENCE TO RELATED APPLICATIONS

The present application claims priority of U.S. non-provisional application Ser. No. 13/174,253, filed Jun. 30, 2011, now issued as U.S. Pat. No. 8,354,262, and of U.S. provisional patent applications 61/360,040, filed Jun. 30, 2010, 61/445,996, filed Feb. 23, 2011, and 61/492,758, filed Jun. 2, 2011, each of which is hereby incorporated by reference herein.

3. TECHNICAL FIELD


The present disclosure relates to soluble compositions and formulations of chemically modified carbonic anhydrase polypeptides that exhibit increased activity and thermostability, and methods of using these polypeptides in carbon capture systems.

4. REFERENCE TO SEQUENCE LISTING, TABLE OR COMPUTER PROGRAM

The Sequence Listing concurrently submitted electronically under 37 C.F.R. §1.821 via EFS-Web in a computer readable form (CRF) as file name CX4-087US1_ST25.txt is herein incorporated by reference. The electronic copy of the Sequence Listing was created on Jun. 29, 2011 with a file size of 2,064,813 bytes.

5. BACKGROUND

The enzyme, carbonic anhydrase ("CA") (EC 4.2.1.1), catalyzes the reversible reactions depicted in Scheme 1:



In the forward or "hydration" reaction, CA combines carbon dioxide and water to provide bicarbonate and a proton, or depending on the pH, to provide carbonate (CO_3^{2-}) and two protons. In the reverse, or "dehydration" reaction, CA combines bicarbonate and a proton to provide carbon dioxide and water. Carbonic anhydrases are metalloenzymes that typically have Zn^{+2} in the active site. However carbonic anhydrases having e.g. Co^{+2} or Cd^{+2} in the active site have been reported. At least three classes of carbonic anhydrases have been identified in nature.

The α -class carbonic anhydrases are found in vertebrates, bacteria, algae, and the cytoplasm of green plants. Vertebrate α -class carbonic anhydrases are among the fastest enzymes known, exhibiting a turnover number (k_{cat}) (the number of molecules of substrate converted by an enzyme to product per catalytic site per unit of time) of 10^6 sec^{-1} . The β -class car-

2

bonic anhydrases are found in bacteria, algae, and chloroplasts, while γ -class carbonic anhydrases are found in *Archaea* and some bacteria. Although carbonic anhydrases of each of these classes have similar active sites, they do not exhibit significant overall amino acid sequence homology and they are structurally distinguishable from one another. Hence, these three classes of carbonic anhydrase provide an example of convergent evolution.

It has been proposed to use carbonic anhydrase as a biological catalyst to accelerate the capture of carbon dioxide produced by combustion of fossil fuels. See e.g., U.S. Pat. Nos. 6,143,556, 6,524,843 B2, 7,176,017 B2, 7,596,952 B2, 7,579,185 B2, 7,740,689 B2, 7,132,090 B2; U.S. Pat. Publ. Nos. 2009/0155889A1, 2010/0086983A1; PCT Publ. Nos. WO2006/089423A1, WO2010/014773A1, WO2010/045689A1. Naturally occurring carbonic anhydrases, however, are not well-suited for use under the process relevant conditions that are required for an economically viable carbon dioxide capture system. These process relevant conditions include: presence in solution with high concentrations of other CO_2 absorption mediating compounds (e.g., amines, ammonia, carbonate ions, amino acids); elevated temperatures (e.g., 40°C . or above, or 15°C . or below in NH_3), alkaline pHs (e.g., pH 8-12); presence of gas contaminants (e.g., high levels NO_x and SO_x); and extended periods of exposure to these challenging conditions (e.g., days to weeks). In addition, such carbonic anhydrases should also be stable to variations in these process conditions, e.g., stable not only at a relatively alkaline pH suitable for hydration and sequestration of carbon dioxide but also at a relatively acidic pH suitable for subsequent release and/or recapture of the hydrated and/or sequestered carbon dioxide.

Chemical conjugates of α -class carbonic anhydrases and some of their physical properties have been described in the following references: Epton et al. "Soluble polymer-protein conjugates: 1. Reactive N-(sym-trinitroaryl)polyacrylamide/acrylylhydrazide copolymers and derived carbonic anhydrase conjugates," *Polymer* 18: 319-323 (1977); Farmer et al., "Assessing the Multimeric States of Proteins: Studies Using Laser Desorption Mass Spectrometry," *Biol. Mass Spectrometry* 20, 796-800 (1991); Gitlin et al., "Peracetylated Bovine Carbonic Anhydrase (BCA-Ac₁₈) Is Kinetically More Stable than Native BCA to Sodium Dodecyl Sulfate," *J. Phys. Chem. B*. 110: 2372-2377 (2006); Gudiksen et al., "Eliminating Positively Charged Lysine e-NH₃⁺ Groups on the Surface of Carbonic Anhydrase Has No Significant Influence on Its Folding from Sodium Dodecyl Sulfate," *J. Am. Chem. Soc.* 127: 4707-4714 (2005); Gudiksen et al., "Increasing the Net Charge and Decreasing the Hydrophobicity of Bovine Carbonic Anhydrase Decreases the Rate of Denaturation with Sodium Dodecyl Sulfate," *Biophys. J.* 91: 298-310 (2006); Bootorabi et al., "Modification of carbonic anhydrase II with acetaldehyde, the first metabolite of ethanol, leads to decreased enzyme activity," *BMC Biochemistry* 9: 32 (2008); Trachtenberg et al., "Carbon Dioxide Transport By Proteic And Facilitated Transport Membranes," *Life Support & Biosphere Science* 6: 293-302 (1999); and Bhattacharya et al., "CO₂ hydration by immobilized carbonic anhydrase," *Biotechnol. Appl. Biochem.* 38: 111-117 (2003).

Accordingly, there is a need in the art for engineered and/or chemically modified carbonic anhydrases with further improved enzymatic properties that can effectively accelerate the absorption of carbon dioxide from a gas stream and/or accelerate desorption of carbon dioxide from a capture solution under process relevant conditions.

6. SUMMARY

The present disclosure provides soluble compositions and homogenous liquid formulations comprising a carbonic

anhydrase that is chemically modified by treatment with a cross-linking agent. The chemically modified carbonic anhydases of the present disclosure are not cross-linked or otherwise attached to a solid phase. The soluble compositions of the present disclosure are soluble in aqueous solvent, forming a homogenous liquid solution. For example, in one embodiment, the present disclosure provides a soluble composition having carbonic anhydrase activity comprising a carbonic anhydrase polypeptide chemically modified by treatment with a cross-linking agent, wherein the polypeptide amino acid sequence has at least 80% identity to SEQ ID NO:2. Similarly, the formulations of the present disclosure, which comprise a chemically modified carbonic anhydrase, a CO₂ absorption mediating compound, and an aqueous solvent, are also homogenous liquid solutions. For example, the homogenous liquid formulation can comprise an aqueous solution of the soluble composition of any of the chemically modified carbonic anhydrase polypeptides disclosed herein and a CO₂ absorption mediating compound.

A surprising advantage of the chemically modified carbonic anhydases of the present disclosure (and the soluble compositions and formulations comprising them) is that they have increased stability and/or increased carbonic anhydrase activity (e.g., at least 1.5-fold, at least 2-fold, at least 4-fold, or even at least 5-fold increased) relative to the same carbonic anhydrase that is not chemically modified under process relevant carbon capture conditions (e.g., high temperature and the presence of high concentrations of CO₂ absorption mediating compounds). More specifically, the chemically modified carbonic anhydases of the present disclosure are capable of improved acceleration (relative to the same carbonic anhydrase that is not chemically modified) of the absorption of carbon dioxide from a gas stream into a solution comprising a CO₂ absorption mediating compound (e.g., amines, ammonia, carbonate ion, amino acid) under suitable conditions useful for various carbon capture processes (e.g., flue-gas scrubbers). Thus, in various embodiments the present disclosure provides chemically modified carbonic anhydrase polypeptides, and compositions and formulations comprising them, that are capable of catalyzing the hydration of carbon dioxide to bicarbonate or the reverse dehydration of bicarbonate to carbon dioxide with increased activity relative to the same carbonic anhydases that are not chemically modified (and other known naturally occurring carbonic anhydases) after exposure to high concentrations of CO₂ absorption mediating compound and/or thermal (e.g., T>40° C.). For example, in some embodiments, the chemically modified carbonic anhydases have carbonic anhydrase activity in 4.2 M MDEA at 50° C. that is increased (e.g., at least 1.5-fold, at least 2-fold, at least 4-fold, or even at least 5-fold increased) relative to the activity of the same carbonic anhydrase polypeptide that is not chemically modified (i.e., unmodified). Similarly, in some embodiments, the chemically modified carbonic anhydrase is characterized by stability in 4.2 M MDEA at 75° C. that is increased (e.g., at least 1.5-fold, at least 2-fold, at least 4-fold, or even at least 5-fold increased) relative to the carbonic anhydrase polypeptide when it is not chemically modified.

Accordingly, the present disclosure also provides methods, processes, and bioreactors for using the disclosed chemically modified carbonic anhydases polypeptides, compositions, and formulations for carbon capture. In some embodiments, the chemically modified carbonic anhydrase polypeptides (and compositions and formulations comprising them) of the present disclosure are used in methods for removing carbon dioxide from a gas stream, e.g., flue gas produced by the combustion of fossil fuels. The methods for removing carbon

dioxide from a gas stream (e.g., capturing or extracting CO₂ gas) comprise the step of contacting the gas stream with a solution, wherein comprises a chemically modified carbonic anhydrase polypeptide of the disclosure having an improved property (e.g., increased activity, thermostability and/or solvent stability), whereby carbon dioxide from the gas stream is absorbed into the solution (e.g., CO₂ gas diffuses into solution and is hydrated to bicarbonate). In some embodiments, the present disclosure provides a method for removing carbon dioxide from a gas stream comprising the step of contacting the gas stream with a homogenous liquid solution under suitable conditions, wherein the solution comprises: (i) a carbonic anhydrase polypeptide chemically modified by treatment with a cross-linking agent; and (ii) a CO₂ absorption mediating compound; whereby the solution absorbs at least a portion of the carbon dioxide from the gas stream. The method can comprise further steps of isolating and/or separately treating the solution comprising the absorbed carbon dioxide according to known methods to further sequester and/or otherwise utilize the carbon dioxide. The methods of removing carbon dioxide from a gas stream using a chemically modified carbonic anhydrase polypeptide disclosed herein can be carried out in the presence of a range of CO₂ absorption mediating compounds, and under a range of suitable conditions disclosed herein including, but not limited to: polypeptide concentration (and polypeptide form—e.g., lysates, whole cells, or purified powder); solution temperature; solution pH; solution CO₂ loading (e.g., $\alpha=0$ to about 0.7); solvent composition; solution concentration of specified CO₂ absorption mediating compound—e.g., an amine compound, ammonia, and/or carbonate ion.

The present disclosure also provides methods, reagents, and conditions for preparing the chemically modified carbonic anhydases polypeptides having the improved properties of increased activity and/or stability that make them particularly useful in the carbon capture methods, processes and bioreactors. In some embodiments, the disclosure provides a method comprising contacting a solution of a carbonic anhydrase polypeptide (e.g., an α -class, β -class, γ -class, ξ -class (zeta-class), and/or recombinant or engineered carbonic anhydrase) with a solution of a cross-linking agent selected from the group consisting of a dialdehyde, a bis-imidate ester, a bis(N-hydroxysuccinimide) ester, a diacid chloride, and mixtures thereof.

The present disclosure provides a variety of carbonic anhydrase polypeptides and cross-linking agents useful for preparation and use of the chemically modified carbonic anhydrase polypeptides, compositions, and formulations exhibiting improved properties under carbon capture process conditions. The various cross-linking agents provided are selected from the group consisting of dialdehyde, a bis-imidate ester, a bis(N-hydroxysuccinimide) ester, a diacid chloride, and mixtures thereof and can include any one of the cross-linking agents: malondialdehyde, glutaraldehyde, dimethyl suberimidate, dimethyl pimelimidate, suberic acid bis(N-hydroxysuccinimide), and mixtures thereof.

In some embodiments, the carbonic anhydrase polypeptide that is chemically modified by treatment with a cross-linking agent is a naturally-occurring α -class, β -class, γ -class, or ξ -class (zeta-class) carbonic anhydrase, or a recombinant carbonic anhydrase derived therefrom. In some embodiments, the carbonic anhydrase is an α -class carbonic anhydrase comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 1298, 1300, 1302, 1304, 1306, and 1308, or a recombinant carbonic anhydrase derived therefrom. In some embodiments, the carbonic anhydrase that is chemically modified is a β -class carbonic anhydrase

comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 2, 1288, 1290, 1292, 1294, and 1296, or a recombinant carbonic anhydrase derived therefrom. In some embodiments, the carbonic anhydrase that is chemically modified is a recombinant or engineered carbonic anhydrase polypeptide that has improved enzymatic properties relative to a reference polypeptide—e.g., a naturally occurring carbonic anhydrase from which the engineered carbonic anhydrase was derived. Thus, the improved enzymatic properties associated with the engineered carbonic anhydrase can be further improved by chemical modification as described in the present disclosure. Accordingly, in one aspect, the chemically modified carbonic anhydrase polypeptides described herein can also have an amino acid sequence that has one or more amino acid differences as compared to a wild-type carbonic anhydrase or an engineered carbonic anhydrase that result in an improved property of the enzyme. Exemplary recombinant or engineered carbonic anhydrase polypeptides having an improved enzyme property can comprise an amino acid sequence selected from the polypeptide amino acid sequences summarized in Tables 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J, and disclosed in the accompanying Sequence Listing, specifically any one or more of the polypeptide amino acid sequences selected from the group consisting of the even-numbered sequence identifiers of SEQ ID NO: 4-1286.

Improvements of the chemically modified carbonic anhydrase polypeptides associated with the chemical modification by treatment with a cross-linking agent as disclosed herein can include increased carbonic anhydrase activity, and/or increased solvent or thermal stability of the carbonic anhydrase in the presence of compounds that mediate the absorption or sequestration of carbon dioxide, including, for example, ammonia, carbonate ions, or amine compounds (e.g., monoethanolamine (MEA), methyldiethanolamine (MDEA), 2-aminomethylpropanolamine (AMP), 2-(2-aminoethylamino)ethanol (AEE), triethanolamine (TEA), 2-amino-2-hydroxymethyl-1,3-propanediol (AHPD), piperazine, piperidine, mono- and diethanolamine). Accordingly, in some embodiments, the chemically modified carbonic anhydrase polypeptides, compositions and formulations comprising them, and methods of using them are characterized by at least 1.5-fold, at least 2-fold, at least 4-fold, or at least 5-fold increased carbonic anhydrase activity relative to the carbonic anhydrase polypeptide when it is not chemically modified, for example, when the activity is measured in 4.2 M MDEA at 50° C., or is measured in 2 M ammonia at 20° C. In some embodiments, the chemically modified carbonic anhydrase polypeptides (and compositions and formulations comprising them) are characterized by at least 1.5-fold, at least 2-fold, at least 4-fold, or at least 5-fold increased stability relative to the carbonic anhydrase polypeptide when it is not chemically modified, for example, when the stability is measured as residual carbonic anhydrase activity following 24 hours exposure to 4.2 M MDEA at 75° C.

The present disclosure also provides methods for preparing the chemically modified carbonic anhydrase polypeptides having improved properties relative to unmodified carbonic anhydrase polypeptides. In some embodiments of the method for preparing the chemically modified carbonic anhydrase polypeptides, the method comprising contacting in a solution: (i) a carbonic anhydrase polypeptide, wherein the polypeptide comprises an amino acid sequence having at least 80% identity to SEQ ID NO:2; and (ii) a cross-linking agent selected from the group consisting of a dialdehyde, a bis-imidate ester, a bis(N-hydroxysuccinimide) ester, a diacid

chloride, and mixtures thereof. The various cross-linking agents provided used in the method of preparing can include any one of the cross-linking agents: malondialdehyde, glutaraldehyde, dimethyl suberimidate, dimethyl pimelimidate, suberic acid bis(N-hydroxysuccinimide), or mixtures thereof. In embodiments of the method for preparing, the cross-linking agent is at a concentration of from about 0.05% to about 10%, from about 0.1% to about 5%, or from about 0.25% to about 2.5%, or at least about 0.05%, at least about 0.1%, at least about 0.25%, at least about 0.5%, at least about 1%, at least about 2%, or at least about 2.5%. In some embodiments of the method for preparing, the solution has a concentration of carbonic anhydrase polypeptide of from about 1 g/L to about 150 g/L, from about 10 g/L to about 100 g/L, from about 25 g/L to about 100 g/L, or at least about 1 g/L, at least about 5 g/L, at least about 10 g/L, at least about 25 g/L, at least about 50 g/L, at least about 75 g/L, or at least about 100 g/L.

7. BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 depicts plots of normalized carbonic anhydrase activity, $k_{1,CA}$ (s^{-1}) for a 1 g/L solution of a recombinant β -class carbonic anhydrase polypeptide of SEQ ID NO: 1152 that has been chemically modified by treatment with 0.25% glutaraldehyde (“GA modified CA”) and the same recombinant carbonic anhydrase polypeptide at 1 g/L that has not been chemically modified (“Unmod. CA”), both versus days of challenge by incubation in assay solution at 75° C. Assay was carried out in 4.2 M MDEA solution, unloaded with CO₂ at 50° C.

8. DETAILED DESCRIPTION

The present disclosure is directed to carbonic anhydrase polypeptides that are chemically modified by treatment with a cross-linking agent and that have improved properties, particularly improved carbonic anhydrase activity and/or stability as compared to the same carbonic anhydrase polypeptides that have not been chemically modified. The present disclosure also is directed to soluble compositions comprising these chemically modified carbonic anhydrase polypeptides, and homogenous liquid formulations of these chemically modified carbonic anhydrase polypeptides and CO₂ absorption mediating compounds. The present disclosure provides the chemically modified polypeptides, and methods of preparing these chemically modified polypeptides (and associated compositions and formulations) by treatment of unmodified naturally occurring α -class, β -class, γ -class, or ξ -class carbonic anhydrase polypeptides, or recombinant carbonic anhydrase polypeptides derived therefrom (which can include amino acid differences relative to a wild-type sequence) with any of a variety of cross-linking agents (e.g., malondialdehyde, glutaraldehyde, dimethyl suberimidate, dimethyl pimelimidate, suberic acid bis(N-hydroxysuccinimide)).

The present disclosure also provides methods for using such chemically modified carbonic anhydrase polypeptides, compositions, and formulations, in processes for the capture and sequestration of carbon dioxide e.g., generated by combustion of fossil fuel. The methods disclosed include the use of the chemically modified carbonic anhydrase polypeptides in combination with various CO₂ absorption mediating compounds (including amines, ammonia, carbonate ions), and under various reaction conditions including conditions comprising high concentrations of the CO₂ absorption mediating compounds including amines, ammonia, carbonate ions, and/

or temperatures that are significantly increased or decreased relative to ambient temperatures.

8.1. Definitions

The technical and scientific terms used in the descriptions herein will have the meanings commonly understood by one of ordinary skill in the art, unless specifically defined otherwise. Accordingly, the following terms are intended to have the following meanings.

“Carbonic anhydrase” and “CA” are used interchangeably herein to refer to a polypeptide having an enzymatic capability of carrying out the reactions depicted in Scheme 1. Carbonic anhydrase as used herein include naturally occurring (wild-type) carbonic anhydrases as well as non-naturally occurring, engineered, or recombinant carbonic anhydrase polypeptides generated by human manipulation.

“Protein”, “polypeptide,” and “peptide” are used interchangeably herein to denote a polymer of at least two amino acids covalently linked by an amide bond, regardless of length or post-translational modification (e.g., glycosylation, phosphorylation, lipidation, myristylation, ubiquitination, etc.). Included within this definition are D- and L-amino acids, and mixtures of D- and L-amino acids.

“Naturally occurring” or “wild-type” refers to the form found in nature. For example, a naturally occurring or wild-type polypeptide or polynucleotide sequence is a sequence present in an organism that can be isolated from a source in nature and which has not been intentionally modified by human manipulation.

“Recombinant” or “engineered” or “non-naturally occurring” when used with reference to, e.g., a cell, nucleic acid, or polypeptide, refers to a material, or a material corresponding to the natural or native form of the material, that has been modified in a manner that would not otherwise exist in nature, or is identical thereto but produced or derived from synthetic materials and/or by manipulation using recombinant techniques. Non-limiting examples include, among others, recombinant cells expressing genes that are not found within the native (non-recombinant) form of the cell or express native genes that are otherwise expressed at a different level.

“Percentage of sequence identity,” “percent identity,” and “percent identical” are used herein to refer to comparisons between polynucleotide sequences or polypeptide sequences, and are determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which either the identical nucleic acid base or amino acid residue occurs in both sequences or a nucleic acid base or amino acid residue is aligned with a gap to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. Determination of optimal alignment and percent sequence identity is performed using the BLAST and BLAST 2.0 algorithms (see e.g., Altschul et al., 1990, *J. Mol. Biol.* 215: 403-410 and Altschul et al., 1977, *Nucleic Acids Res.* 3389-3402). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information website.

Briefly, the BLAST analyses involve first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy

some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as, the neighborhood word score threshold (Altschul et al, supra). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff, 1989, *Proc Natl Acad Sci USA* 89:10915).

Numerous other algorithms are available that function similarly to BLAST in providing percent identity for two sequences. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith and Waterman, 1981, *Adv. Appl. Math.* 2:482, by the homology alignment algorithm of Needleman and Wunsch, 1970, *J. Mol. Biol.* 48:443, by the search for similarity method of Pearson and Lipman, 1988, *Proc. Natl. Acad. Sci. USA* 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the GCG Wisconsin Software Package), or by visual inspection (see generally, *Current Protocols in Molecular Biology*, F. M. Ausubel et al., eds., *Current Protocols*, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (1995 Supplement) (Ausubel)). Additionally, determination of sequence alignment and percent sequence identity can employ the BESTFIT or GAP programs in the GCG Wisconsin Software package (Accelrys, Madison Wis.), using default parameters provided.

“Reference sequence” refers to a defined sequence to which another sequence is compared. A reference sequence is not limited to wild-type sequences, and can include engineered or altered sequences. For example, a reference sequence can be a previously engineered or altered amino acid sequence. A reference sequence also may be a subset of a larger sequence, for example, a segment of a full-length gene or polypeptide sequence. Generally, a reference sequence is at least 20 nucleotide or amino acid residues in length, at least 25 residues in length, at least 50 residues in length, or the full length of the nucleic acid or polypeptide. Since two polynucleotides or polypeptides may each (1) comprise a sequence (i.e., a portion of the complete sequence) that is similar between the two sequences, and (2) may further comprise a sequence that is divergent between the two sequences, sequence comparisons between two (or more) polynucleotides or polypeptide are typically performed by comparing sequences of the two polynucleotides over a comparison window to identify and compare local regions of sequence similarity.

“Comparison window” refers to a conceptual segment of at least about 20 contiguous nucleotide positions or amino acids

residues wherein a sequence may be compared to a reference sequence of at least 20 contiguous nucleotides or amino acids and wherein the portion of the sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The comparison window can be longer than 20 contiguous residues, and includes, optionally 30, 40, 50, 100, or longer windows.

“Corresponding to”, “reference to” or “relative to” when used in the context of the numbering of a given amino acid or polynucleotide sequence refers to the numbering of the residues of a specified reference sequence when the given amino acid or polynucleotide sequence is compared to the reference sequence. In other words, the residue number or residue position of a given polymer is designated with respect to the reference sequence rather than by the actual numerical position of the residue within the given amino acid or polynucleotide sequence. For example, a given amino acid sequence, such as that of an engineered carbonic anhydrase, can be aligned to a reference sequence by introducing gaps to optimize residue matches between the two sequences. In these cases, although the gaps are present, the numbering of the residue in the given amino acid or polynucleotide sequence is made with respect to the reference sequence to which it has been aligned.

“Different from” or “differs from” with respect to a designated reference sequence refers to difference of a given amino acid or polynucleotide sequence when aligned to the reference sequence. Generally, the differences can be determined when the two sequences are optimally aligned. Differences include insertions, deletions, or substitutions of amino acid residues in comparison to the reference sequence.

“Derived from” as used herein in the context of engineered carbonic anhydrase enzymes, identifies the originating carbonic anhydrase enzyme, and/or the gene encoding such carbonic anhydrase enzyme, upon which the engineering was based.

“Amino acid residue” or “amino acid” or “residue” as used herein refers to the specific monomer at a sequence position of a polypeptide (e.g., D7 indicates that the “amino acid” or “residue” at position 7 of SEQ ID NO: 2 is an aspartic acid (D).)

“Amino acid difference” or “residue difference” refers to a change in the amino acid residue at a position of a polypeptide sequence relative to the amino acid residue at a corresponding position in a reference sequence. The positions of amino acid differences generally are referred to herein as “Xn,” where n refers to the corresponding position in the reference sequence upon which the residue difference is based. For example, a “residue difference at position X3 as compared to SEQ ID NO: 2” refers to a change of the amino acid residue at the polypeptide position corresponding to position 3 of SEQ ID NO:2. Thus, if the reference polypeptide of SEQ ID NO: 2 has a glutamine at position 3, then a “residue difference at position X3 as compared to SEQ ID NO:2” an amino acid substitution of any residue other than glutamine at the position of the polypeptide corresponding to position 3 of SEQ ID NO: 2. In most instances herein, the specific amino acid residue difference at a position is indicated as “XnY” where “Xn” specifies the corresponding position as described above, and “Y” is the single letter identifier of the amino acid found in the engineered polypeptide (i.e., the different residue than in the reference polypeptide). In some instances, the present disclosure also provides specific amino acid differences denoted by the conventional notation “AnB”, where A is the single letter identifier of the residue in the reference sequence, “n” is the

number of the residue position in the reference sequence, and B is the single letter identifier of the residue substitution in the sequence of the engineered polypeptide. For example, “D7S” would refer to the substitution of the amino acid residue, aspartic acid (D) at position 7 of reference sequence with the amino acid serine (S). In some instances, a polypeptide of the present disclosure can include one or more amino acid residue differences relative to a reference sequence, which is indicated by a list of the specified positions where changes are made relative to the reference sequence. The present disclosure includes engineered polypeptide sequences comprising one or more amino acid differences that include either/or both conservative and non-conservative amino acid substitutions.

“Conservative amino acid substitution” refers to a substitution of a residue with a different residue having a similar side chain, and thus typically involves substitution of the amino acid in the polypeptide with amino acids within the same or similar defined class of amino acids. By way of example and not limitation, an amino acid with an aliphatic side chain may be substituted with another aliphatic amino acid, e.g., alanine, valine, leucine, and isoleucine; an amino acid with hydroxyl side chain is substituted with another amino acid with a hydroxyl side chain, e.g., serine and threonine; an amino acids having aromatic side chains is substituted with another amino acid having an aromatic side chain, e.g., phenylalanine, tyrosine, tryptophan, and histidine; an amino acid with a basic side chain is substituted with another amino acid with a basis side chain, e.g., lysine and arginine; an amino acid with an acidic side chain is substituted with another amino acid with an acidic side chain, e.g., aspartic acid or glutamic acid; and a hydrophobic or hydrophilic amino acid is replaced with another hydrophobic or hydrophilic amino acid, respectively. Exemplary conservative substitutions are provided in Table 1.

TABLE 1

Residue	Possible Conservative Substitutions
A, L, V, I	Other aliphatic (A, L, V, I) Other non-polar (A, L, V, I, G, M)
G, M	Other non-polar (A, L, V, I, G, M)
D, E	Other acidic (D, E)
K, R	Other basic (K, R)
N, Q, S, T	Other polar
H, Y, W, F	Other aromatic (H, Y, W, F)
C, P	None

“Non-conservative substitution” refers to substitution of an amino acid in the polypeptide with an amino acid with significantly differing side chain properties. Non-conservative substitutions may use amino acids between, rather than within, the defined groups and affects (a) the structure of the peptide backbone in the area of the substitution (e.g., proline for glycine) (b) the charge or hydrophobicity, or (c) the bulk of the side chain. By way of example and not limitation, an exemplary non-conservative substitution can be an acidic amino acid substituted with a basic or aliphatic amino acid; an aromatic amino acid substituted with a small amino acid; and a hydrophilic amino acid substituted with a hydrophobic amino acid.

“Deletion” refers to modification of the polypeptide by removal of one or more amino acids from the reference polypeptide. Deletions can comprise removal of 1 or more amino acids, 2 or more amino acids, 5 or more amino acids, 10 or more amino acids, 15 or more amino acids, or 20 or more amino acids, up to 10% of the total number of amino acids, or up to 20% of the total number of amino acids making up the

polypeptide while retaining enzymatic activity and/or retaining the improved properties of an engineered carbonic anhydrase enzyme. Deletions can be directed to the internal portions and/or terminal portions of the polypeptide. In various embodiments, the deletion can comprise a continuous segment or can be discontinuous.

“Insertion” refers to modification of the polypeptide by addition of one or more amino acids to the reference polypeptide. In some embodiments, the improved engineered carbonic anhydrase enzymes comprise insertions of one or more amino acids to the naturally occurring carbonic anhydrase polypeptide as well as insertions of one or more amino acids to other improved carbonic anhydrase polypeptides. Insertions can be in the internal portions of the polypeptide, or to the carboxy or amino terminus. Insertions as used herein include fusion proteins as is known in the art. The insertion can be a contiguous segment of amino acids or separated by one or more of the amino acids in the naturally occurring polypeptide.

“Fragment” as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion, but where the remaining amino acid sequence is identical to the corresponding positions in the sequence. Fragments can typically have about 80%, 90%, 95%, 98%, and 99% of the full-length carbonic anhydrase polypeptide, for example the polypeptide of SEQ ID NO:2. The amino acid sequences of the specific recombinant carbonic anhydrase polypeptides included in the Sequence Listing of the present disclosure include an initiating methionine (M) residue (i.e., M represents residue position 1). The skilled artisan, however, understands that this initiating methionine residue can be removed by biological processing machinery, such as in a host cell or in vitro translation system, to generate a mature protein lacking the initiating methionine residue, but otherwise retaining the enzyme’s properties. Consequently, the term “amino acid residue difference relative to SEQ ID NO: 2 at position X_n” as used herein may refer to position “X_n” or to the corresponding position (e.g., position (X-1)_n) in a reference sequence that has been processed so as to lack the starting methionine.

“Improved enzyme property” or “improved property” as used herein refers to a functional characteristic of an enzyme that is improved relative to the same functional characteristic of a reference enzyme. Improved enzyme properties of the engineered carbonic anhydrase polypeptides disclosed herein can include but are not limited to: increased thermostability, increased solvent stability, increased pH stability, altered pH activity profile, increased activity (including increased rate conversion of substrate to product, or increased percentage conversion in a period of time), increased and/or altered stereoselectivity, altered substrate specificity and/or preference, decreased substrate, product, and side-product inhibition (e.g., CO₂, carbonate, bicarbonate, carbamate, or solvent-adducts thereof), decreased inhibition by a component of the feedstock (e.g. exhaust, flue gas components such as NO_x and SO_x compounds, etc.), decreased side-product or impurity production, altered cofactor preference, increased expression, increased secretion, as well as increased stability and/or activity in the presence of additional compounds reagents useful for absorption or sequestration of carbon dioxide, including, for example, amine solvents such as monoethanolamine, methyldiethanolamine, and 2-aminomethylpropanolamine.

“Stability in the presence of” as used in the context of improved enzyme properties disclosed herein refers to stability of the enzyme measured during or after exposure of the enzyme to certain compounds/reagents/ions (e.g., amine compound, ammonia, and/or carbonate ions) in the same

solution with the enzyme. It is intended to encompass challenge assays of stability where the enzyme is first exposed to the amine compound or ammonia for some period of time then assayed in a solution under different conditions.

“Solution” as used herein refers to any medium, phase, or mixture of phases, in which the carbonic anhydrase polypeptide is active. It is intended to include purely liquid phase solutions (e.g., aqueous, or aqueous mixtures with co-solvents, including emulsions and separated liquid phases), as well as slurries and other forms of solutions having mixed liquid-solid phases.

“Homogenous liquid solution” as used herein refers to a formulation that is uniformly liquid (e.g., a liquid that does not include a suspended solid phase).

“Soluble composition” as used herein refers to a composition capable of dissolving to form a homogenous liquid solution in an aqueous solvent.

“Thermostability” refers to the functional characteristic of retaining activity (e.g., more than 60% to 80%) in the presence of, or after exposure to for a period of time (e.g. 0.5-24 hrs), elevated temperatures (e.g. 30-100° C.) compared to the activity of an untreated enzyme.

“Solvent stability” refers to the functional characteristic of retaining activity (e.g., more than 60% to 80%) in the presence of, or after exposure to for a period of time (e.g. 0.5-24 hrs), increased concentrations (e.g., 5-99%) of solvent compared to the activity of an untreated enzyme.

“pH stability” refers to the functional characteristic of retaining activity (e.g., more than 60% to 80%) in the presence of, or after exposure to for a period of time (e.g. 0.5-24 hrs), conditions of high or low pH (e.g., pH 9 to 12) compared to the activity of an untreated enzyme.

“Increased enzymatic activity” or “increased activity” refers to an improved property of the engineered enzyme (e.g., carbonic anhydrase), which can be represented by an increase in specific activity (e.g., product produced/time/weight protein) or an increase in percent conversion of the substrate to the product (e.g., percent conversion of carbon dioxide to bicarbonate and/or carbonate in a specified time period using a specified amount of carbonic anhydrase) as compared to a reference enzyme under suitable reaction conditions. Exemplary methods to determine enzyme activity are provided in the Examples. Any property relating to enzyme activity may be affected, including the classical enzyme properties of K_m , V_{max} or k_{cat} , changes of which can lead to increased enzymatic activity. Improvements in enzyme activity can be from about 1.1-times the enzymatic activity of the corresponding wild-type carbonic anhydrase enzyme, to as much as 1.2-times, 1.5-times, 2-times, 3-times, 4-times, 5-times, 6-times, 7-times, or more than 8-times the enzymatic activity than the naturally occurring parent carbonic anhydrase. It is understood by the skilled artisan that the activity of any enzyme is diffusion limited such that the catalytic turnover rate cannot exceed the diffusion rate of the substrate, including any required cofactors. The theoretical maximum of the diffusion limit, or k_{cat}/K_m , is generally about 10^8 to 10^9 ($M^{-1} s^{-1}$). Hence, any improvements in the enzyme activity of the carbonic anhydrase will have an upper limit related to the diffusion rate of the substrates acted on by the carbonic anhydrase enzyme. Carbonic anhydrase activity can be measured by any one of standard assays used for measuring carbonic anhydrase, e.g., as provided in the Examples. Comparisons of enzyme activities are made, e.g., using a defined preparation of enzyme, a defined assay under a set of conditions, as further described in detail herein. Generally, when lysates are compared, the numbers of cells and the amount of protein assayed are determined as well as use of identical expression

systems and identical host cells to minimize variations in amount of enzyme produced by the host cells and present in the lysates.

“Conversion” refers to the enzymatic conversion of the substrate to the corresponding product. “Percent conversion” refers to the percent of the substrate that is reduced to the product within a period of time under specified conditions. Thus, the “enzymatic activity” or “activity” of a carbonic anhydrase polypeptide can be expressed as “percent conversion” of the substrate to the product.

“Isolated polypeptide” refers to a polypeptide which is substantially separated from other contaminants that naturally accompany it, e.g., protein, lipids, and polynucleotides. The term embraces polypeptides which have been removed or purified from their naturally-occurring environment or expression system (e.g., host cell or in vitro synthesis). The improved carbonic anhydrase enzymes may be present within a cell, present in the cellular medium, or prepared in various forms, such as lysates or isolated preparations. As such, in some embodiments, the improved carbonic anhydrase enzyme can be an isolated polypeptide.

“Substantially pure polypeptide” refers to a composition in which the polypeptide species is the predominant species present (i.e., on a molar or weight basis it is more abundant than any other individual macromolecular species in the composition), and is generally a substantially purified composition when the object species comprises at least about 50 percent of the macromolecular species present by mole or % weight. Generally, a substantially pure carbonic anhydrase composition will comprise about 60% or more, about 70% or more, about 80% or more, about 90% or more, about 95% or more, and about 98% or more of all macromolecular species by mole or % weight present in the composition. In some embodiments, the object species is purified to essential homogeneity (i.e., contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species. Solvent species, small molecules (<500 Daltons), and elemental ion species are not considered macromolecular species. In some embodiments, the isolated improved carbonic anhydrase polypeptide is a substantially pure polypeptide composition.

“Coding sequence” refers to that portion of a polynucleotide that encodes an amino acid sequence of a protein (e.g., a gene).

“Heterologous” polynucleotide refers to any polynucleotide that is introduced into a host cell by laboratory techniques, and includes polynucleotides that are removed from a host cell, subjected to laboratory manipulation, and then reintroduced into a host cell.

“Codon optimized” refers to changes in the codons of the polynucleotide encoding a protein to those preferentially used in a particular organism such that the encoded protein is efficiently expressed in the organism of interest. In some embodiments, the polynucleotides encoding the carbonic anhydrase enzymes may be codon optimized for optimal production from the host organism selected for expression.

“Control sequence” is defined herein to include all components, which are necessary or advantageous for the expression of a polynucleotide and/or polypeptide of the present disclosure. Each control sequence may be native or foreign to the polynucleotide of interest. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal peptide sequence, and transcription terminator.

“Operably linked” is defined herein as a configuration in which a control sequence is appropriately placed (i.e., in a

functional relationship) at a position relative to a polynucleotide of interest such that the control sequence directs or regulates the expression of the polynucleotide and/or polypeptide of interest.

“Cross-linking agent” as used herein refers to a compound or a mixture of compounds that causes or forms covalent or ionic bonds linking amino acid residues of one or more polypeptide molecules.

“Chemically modified polypeptide” as used herein in the context of “chemically modified carbonic anhydrase polypeptide” refers to a polypeptide molecule having one or more amino acid residues which have formed covalent or ionic bonds with a compound (e.g., a cross-linking agent such as glutaraldehyde).

“CO₂ absorption mediating compound” as used herein refers to a compound that increases the ability (e.g., kinetic and/or thermodynamic) of a solution in which it is present to absorb CO₂ gas. CO₂ absorption mediating compounds can include ammonia, carbonate salts, amino acids, and amine compounds, including but not limited to: 2-(2-aminoethylamino)ethanol (AEE), 2-amino-2-hydroxymethyl-1,3-propanediol (AHPD), 2-amino-2-methyl-1-propanol (AMP), diethanolamine (DEA), diisopropanolamine (DIPA), N-hydroxyethylpiperazine (HEP), N-methyldiethanolamine (MDEA), monoethanolamine (MEA), N-methylpiperazine (MP), piperazine, piperidine, 2-(2-tert-butylaminoethoxy) ethanol (TBEE), triethanolamine (TEA), triisopropanolamine (TIA), tris, 2-(2-aminoethoxy)ethanol, 2-(2-tert-butylaminopropoxy)ethanol, 2-(2-tert-amylaminoethoxy) ethanol, 2-(2-isopropylaminopropoxy)ethanol, and 2-(2-(1-methyl-1-ethylpropylamino)ethoxy)ethanol.

8.2. Chemically Modified Carbonic Anhydrase Polypeptides

The present disclosure provides carbonic anhydrase polypeptides that are chemically modified by treatment with a cross-linking agent. The disclosure also provides soluble compositions and homogenous liquid formulations comprising these chemically modified carbonic anhydrase polypeptides. These chemically modified carbonic anhydrases are not cross-linked or otherwise attached to a solid phase. The soluble compositions comprising them are soluble in aqueous solvent, forming a homogenous liquid solution. For example, in one embodiment, the present disclosure provides a soluble composition having carbonic anhydrase activity comprising a carbonic anhydrase polypeptide chemically modified by treatment with a cross-linking agent, wherein the polypeptide amino acid sequence has at least 80% identity to SEQ ID NO:2. Similarly, the present disclosure provides formulations comprising the chemically modified carbonic anhydrases, together with a CO₂ absorption mediating compound, and an aqueous solvent. These formulations are also homogenous liquid solutions. For example, these homogenous liquid formulations can comprise an aqueous solution of the soluble composition of any of the chemically modified carbonic anhydrase polypeptides disclosed herein and a CO₂ absorption mediating compound selected from ammonia, an amine compound, or carbonate ion.

A surprising advantage of these soluble compositions and homogenous liquid formulations comprising chemically modified carbonic anhydrases is that they have increased stability and/or increased carbonic anhydrase activity (e.g., at least 1.5-fold, at least 2-fold, at least 4-fold, or even at least 5-fold increased) under process relevant carbon capture conditions (e.g., high temperature and the presence of high concentrations of CO₂ absorption mediating compounds) relative

to the same carbonic anhydrase that is not chemically modified. Accordingly, the chemically modified carbonic anhydrases of the present disclosure (and their soluble compositions and homogenous liquid formulations) are capable of improved acceleration of the absorption of carbon dioxide from a gas stream into a solution comprising a CO₂ absorption mediating compound (e.g., amines, ammonia, carbonate ion, amino acid) under suitable conditions useful for various carbon capture processes (e.g., flue-gas scrubbers) relative to the acceleration of the same carbonic anhydrase that is not chemically modified. Thus, in various embodiments the present disclosure provides chemically modified carbonic anhydrase polypeptides, and compositions and formulations comprising them, that are capable of catalyzing the hydration of carbon dioxide to bicarbonate or the reverse dehydration of bicarbonate to carbon dioxide with increased activity relative to the same carbonic anhydrases that are not chemically modified (and other known naturally occurring carbonic anhydrases) after exposure to high concentrations of CO₂ absorption mediating compound and/or thermal (e.g., T>40° C.). For example, in some embodiments, the chemically modified carbonic anhydrases have carbonic anhydrase activity in 4.2 M MDEA at 50° C. that is increased (e.g., at least 1.5-fold, at least 2-fold, at least 4-fold, or even at least 5-fold increased) relative to the activity of the same carbonic anhydrase polypeptide that is not chemically modified (i.e., unmodified). Similarly, in some embodiments, the chemically modified carbonic anhydrase is characterized by stability in 4.2 M MDEA at 75° C. that is increased (e.g., at least 1.5-fold, at least 2-fold, at least 4-fold, or even at least 5-fold increased) relative to the carbonic anhydrase polypeptide when it is not chemically modified.

In some embodiments the present disclosure provides a soluble composition comprising a carbonic anhydrase polypeptide chemically modified by treatment with a cross-linking agent. In some embodiments of the soluble composition, the carbonic anhydrase polypeptide is a naturally occurring carbonic anhydrase selected from an α -class, γ -class, β -class, or ξ -class carbonic anhydrase, or a recombinant (or engineered) carbonic anhydrase derived from a naturally occurring α -class, γ -class, β -class, or ξ -class carbonic anhydrase. Carbonic anhydrase polypeptides, particularly engineered β -class carbonic anhydrase polypeptides, useful for chemical modification are described in greater detail below.

A wide-range of compounds useful for cross-linking proteins, particularly enzymes, are well-known in the art (see e.g., U.S. Pat. No. 4,101,380, which is hereby incorporated by reference herein) and commercially available (see e.g., catalog of "crosslinking reagents" available from Thermo Scientific, USA at www.piercenet.com). In some embodiments of the soluble composition, the cross-linking agent is selected from the group consisting of a dialdehyde, a bis-imidate ester, a bis(N-hydroxysuccinimide) ester, a diacid chloride, and mixtures thereof. In some embodiments, the specific cross-linking agent is selected from the group consisting of malondialdehyde, glutaraldehyde, dimethyl suberimidate, dimethyl pimelimidate, suberic acid bis(N-hydroxysuccinimide), and mixtures thereof.

In some embodiments of the soluble composition, the cross-linking agent is a dialdehyde optionally having one or more carbon atoms between the two aldehyde groups, for example wherein the dialdehyde is selected from the group consisting of glyoxal, succindialdehyde, malondialdehyde, glutaraldehyde, and mixtures thereof. In addition, the two dialdehyde groups can be linked by a polyethylene glycol group of varying lengths. In a particular embodiment, the cross-linking agent is glutaraldehyde.

In some embodiments of the soluble composition, the cross-linking agent is a bis-imidate ester, and in particular embodiments, a bis-imidate ester optionally having one or more carbon atoms between the two imidate ester groups. Useful imidate esters include bis-imidate esters optionally having one or more carbon atoms between the two imidate ester groups, including but not limited to: imidate esters (such as methyl or ethyl) of oxalimidate, malonimidate, succinimidate, glutarimidate, adipimidate, pimelimidate, and suberimidate. In addition, the two dialdehyde groups can be linked by a polyethylene glycol group of varying lengths.

The cross-linking of proteins using diacid chlorides is known in the art (see e.g., U.S. Pat. No. 4,101,380), and in some embodiments of the soluble composition, the cross-linking agent is a diacid chloride. Diacid chlorides useful in the chemically modified carbonic anhydrase polypeptides of the disclosure include those having structures analogous to the dialdehydes described herein. Accordingly, in some embodiments, the diacid chloride cross-linking agent can optionally have one or more carbon atoms between the two acyl chloride groups, and include, but are not limited to, diacid chloride compounds such as adipoyl chloride. In addition, the two acyl chloride groups can be linked by a polyethylene glycol group of varying lengths.

As shown in the Examples, carbonic anhydrase polypeptides modified with imidate esters may undergo a reversible cleavage reaction, whereby over time, the polypeptide loses the imidate ester chemical modification (i.e., modification undergoes an equilibrium cleavage reaction), and the improved activity and/or stability associated with it. Accordingly, in some embodiments of the soluble composition, the cross-linking agent is a bis(N-hydroxysuccinimide) ester of a di-carboxylic acid that forms an irreversible chemical modification of the polypeptide. Useful bis(N-hydroxysuccinimide) esters include those prepared from a di-carboxylic acid selected from the group consisting of oxalate, malonate, succinate, glutarate, adipate, pimelate, suberate, and mixtures thereof. Accordingly, in particular embodiments of the soluble composition, the cross-linking agent is a bis(N-hydroxysuccinimide) ester of a di-carboxylic acid selected from the group consisting of oxalate, malonate, succinate, glutarate, adipate, pimelate, suberate, and mixtures thereof. In addition, the two ester groups can be linked by a polyethylene glycol group of various length. Also, bis(N-hydroxysulfosuccinimide) esters of di-carboxylic acid as described above can be used. These have the advantage of being more water soluble than their bis(N-hydroxysuccinimide) ester counterpart due to the addition of a sulfonate group.

Various embodiments of preparing and using the carbonic anhydrase polypeptides chemically modified by treatment with cross-linking agents used in the soluble composition and homogenous liquid formulations are disclosed in greater detail below (see e.g., Examples). Generally, treatment comprises exposure of an unmodified carbonic anhydrase polypeptide (e.g., in an aqueous solution at a concentration of 10 g/L and 100 g/L) with the cross-linking agent also in the aqueous solution at a specified concentration. In some embodiments of the soluble composition, the treatment with a cross-linking agent comprises exposure of the carbonic anhydrase polypeptide to the cross-linking agent at a concentration of from about 0.025% to about 10%, from about 0.05% to about 5%, from about 0.1% to about 5%, or from about 0.25% to about 2.5%. In some embodiments, the treatment comprises exposure of the carbonic anhydrase polypeptide to the cross-linking reagent at a concentration of at least about 0.025%, at least about 0.1%, at least about 0.25%, at least about 0.5%, at least about 1%, at least about 2%, at least

about 2.5%, or at least about 5%. Generally, either percentage concentrations based on percent volume/volume (v/v) or weight/volume (w/v) can be used with the cross-linking agents disclosed herein without a significant difference in performance for the purposes disclosed herein. Typically, where the cross-linking agent is obtained as a liquid reagent, percent (v/v) is used. For example, as detailed in the Examples, glutaraldehyde is obtained from Sigma-Aldrich (St. Louis, USA) as a 25% solution and further diluted based on percentage (v/v) to the desired polypeptide treatment concentration. However, where the cross-linking agent typically obtained as a solid reagent a percent (w/v) solution can be used.

The structure of the soluble composition can vary depending on the specific carbonic anhydrase polypeptide amino acid sequence that is chemically modified. Generally, regardless of the specific sequence, the cross-linking agents disclosed herein result in chemical modification of one or more amino acid lysine residues, and preferably lysine residues that are present on the surface of the polypeptide or between subunits. Accordingly, in some embodiments of the soluble composition, the carbonic anhydrase polypeptide chemically modified by treatment with a cross-linking agent has at least one chemically modified lysine residue. In some embodiments, the treatment with a cross-linking agent results in the carbonic anhydrase polypeptide having at least one lysine residue cross-linked to another lysine residue on the same carbonic anhydrase polypeptide molecule. In some embodiments, the treatment with a cross-linking agent results in the carbonic anhydrase polypeptide having at least one lysine residue cross-linked to another lysine residue on a different carbonic anhydrase polypeptide molecule (i.e., at least one CA dimer).

In an embodiment of the soluble composition, the carbonic anhydrase polypeptide that is chemically modified is an α -class carbonic anhydrase polypeptide or a recombinant carbonic anhydrase polypeptide derived from an α -class carbonic anhydrase. In some embodiments, the α -class carbonic anhydrase that is chemically modified is an α -class carbonic anhydrase from human (*Homo sapiens*), rat (*Rattus norvegicus*), cow (*Bos taurus*), chicken (*Gallus gallus*), fish (*Cyprino carpio*), or the bacteria, *Neisseria gonorrhoeae*, or a recombinant carbonic anhydrase polypeptide derived from any one of these α -class carbonic anhydrase. In some embodiments, the α -class carbonic anhydrase that is chemically modified comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 1298, 1300, 1302, 1304, 1306, and 1308, or a recombinant carbonic anhydrase polypeptide derived from any one of these α -class carbonic anhydrase sequences.

In another embodiment of the soluble composition, the carbonic anhydrase polypeptide is a recombinant β -class carbonic anhydrase polypeptide derived from the wild-type *Desulfovibrio vulgaris* carbonic anhydrase comprising the amino acid sequence of SEQ ID NO: 2, or derived from a sequence homolog of SEQ ID NO: 2 selected from the group consisting of SEQ ID NO: 1288, 1290, 1292, 1294, and 1296. A wide range of engineered polypeptides useful in such an embodiment of the soluble composition are provided below in Tables 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J. In some embodiments, the carbonic anhydrase polypeptide amino acid sequence comprises an even-numbered amino acid sequence selected from any one of SEQ ID NO: 4-1286. In such embodiments, the carbonic anhydrase polypeptide amino acid sequence has surface lysine residues at the following positions (relative to SEQ ID NO: 2): X18, X37, X147, X156, X184, or X198. Accordingly, in some embodiments of the

soluble composition wherein the polypeptide is a recombinant β -class carbonic anhydrase having an amino acid sequence derived from SEQ ID NO: 2, the treatment with a cross-linking agent results in the carbonic anhydrase polypeptide having a chemically modified lysine residue at one or more of the following positions relative to SEQ ID NO: 2: X18, X37, X147, X156, X184, or X198. In some embodiments of the soluble composition, the carbonic anhydrase polypeptide amino acid sequence comprises at least the following amino acid residue difference relative to SEQ ID NO: 2: X56S. In some embodiments of the soluble composition, the carbonic anhydrase polypeptide amino acid sequence comprises at least the following amino acid residue difference relative to SEQ ID NO: 2: X30R, X40L, X56S, X84Q, X120R, and X139M. In some embodiments of the soluble composition, the carbonic anhydrase polypeptide amino acid sequence an amino acid sequence selected from any one of SEQ ID NO: 26, 190, 206, 238, 252, 270, 274, 284, 306, 318, 328, 332, 340, 354, 596, 606, 656, 678, 1080, 1110, 1148, 1152, 1156, and 1158.

In some embodiments, the present disclosure provides a soluble composition comprising chemically modified polypeptide having carbonic anhydrase activity characterized by an amino acid sequence having at least 80% identity to SEQ ID NO:2 and at least one residue chemically modified by treatment with a cross-linking agent selected from the group consisting of: glutaraldehyde, dimethyl suberimidate, dimethyl pimelimidate, suberic acid bis(N-hydroxysuccinimide), and mixtures thereof. In some embodiments, the at least one residue that is chemically modified by treatment with a cross-linking agent is a surface lysine residue at one or more of the following positions relative to SEQ ID NO: 2: X18, X37, X147, X156, X184, or X198.

As described in greater detail below, the recombinant carbonic anhydrase polypeptides derived from SEQ ID NO: 2 used in the soluble compositions typically have at least one improved enzyme property relative to the wild-type polypeptide of SEQ ID NO: 2. For example increased activity and/or stability in the presence of high concentrations of CO₂ absorption mediating compounds (e.g., >4 M MDEA or >2 M NH₃) and at increased temperatures (e.g., 40° C. or higher). Thus, in some embodiments of the soluble composition, the carbonic anhydrase polypeptide prior to chemical modification is a recombinant carbonic anhydrase polypeptide having an activity half-life (t_{1/2}) of at least 9 hours in 4 M MDEA at 50° C.

Generally, the embodiments of the soluble composition the chemically modified carbonic anhydrase polypeptide of the composition has an improved enzyme property of increased carbonic anhydrase activity and/or increased stability relative to the same carbonic anhydrase polypeptide that is not chemically modified. Thus, in some embodiments of the soluble composition, the carbonic anhydrase activity of the chemically modified carbonic anhydrase is increased relative to the carbonic anhydrase polypeptide when it is not chemically modified (i.e., unmodified), when measured in 4.2 M MDEA at 50° C. In some embodiments the carbonic anhydrase activity is increased at least 1.5-fold, at least 2-fold, at least 3-fold, at least 4-fold, or at least 5-fold.

In some embodiments of the soluble composition, the chemically modified carbonic anhydrase is characterized by the improved enzyme property of increased stability relative to the carbonic anhydrase polypeptide when it is not chemically modified (i.e., unmodified), when measured as residual carbonic anhydrase activity following 24 hours exposure to 4.2 M MDEA at 75° C. In some embodiments the carbonic

anhydrase stability is increased at least 1.5-fold, at least 2-fold, at least 3-fold, at least 4-fold, or at least 5-fold.

Due to their improved properties, the soluble compositions comprising chemically modified carbonic anhydrase polypeptides are particularly useful in methods for removing carbon dioxide from a gas stream. Generally, these methods, which are disclosed in greater detail below, comprise the step of contacting under suitable conditions the gas stream with a solution comprising a soluble composition of a chemically modified carbonic anhydrase polypeptide as disclosed herein, whereby the solution absorbs at least a portion of the carbon dioxide from the gas stream.

As mentioned above, the present disclosure also provides a homogenous liquid formulation comprising a carbonic anhydrase polypeptide chemically modified by treatment with a cross-linking agent and a CO₂ absorption mediating compound. These homogenous liquid formulations can comprise any of the carbonic anhydrase polypeptides chemically modified by treatment with a cross-linking agent disclosed elsewhere herein. The homogenous liquid formulations of the present disclosure can be prepared by dissolving any of the soluble compositions (disclosed elsewhere herein) in an aqueous solution also comprising the desired CO₂ absorption mediating compound. Accordingly, the present disclosure provides a homogenous liquid formulation comprising an aqueous solution of a soluble composition comprising (i) a carbonic anhydrase polypeptide chemically modified by treatment with a cross-linking agent and (ii) a CO₂ absorption mediating compound. In various embodiments of the homogenous liquid formulation, the CO₂ absorption mediating compound can be selected from the group consisting of an amine compound, ammonia, carbonate ion, and mixtures thereof.

In some embodiments, the CO₂ absorption mediating compound used in the homogenous liquid formulation is an amine compound selected from the group consisting of: 2-(2-aminoethylamino)ethanol (AEE), 2-amino-2-hydroxymethyl-1,3-propanediol (AHPD), 2-amino-2-methyl-1-propanol (AMP), diethanolamine (DEA), diisopropanolamine (DIPA), N-hydroxyethylpiperazine (HEP), N-methyldiethanolamine (MDEA), monoethanolamine (MEA), N-methylpiperazine (MP), piperazine, piperidine, 2-(2-tert-butylaminoethoxy) ethanol (TBEE), triethanolamine (TEA), triisopropanolamine (TIA), tris, 2-(2-aminoethoxy)ethanol, 2-(2-tert-butylaminopropoxy)ethanol, 2-(2-tert-amylaminoethoxy) ethanol, 2-(2-isopropylaminopropoxy)ethanol, 2-(2-(1-methyl-1-ethylpropylamino)ethoxy)ethanol, and mixtures thereof.

Due to their improved properties, the homogenous liquid formulations comprising a chemically modified carbonic anhydrase polypeptide and a CO₂ absorption mediating compound, as disclosed herein, also are particularly useful in methods for removing carbon dioxide from a gas stream. Such methods, which are disclosed in greater detail below, generally comprise a step of contacting the gas stream with the homogenous liquid formulation under suitable conditions, whereby the homogenous liquid formulation absorbs at least a portion of the carbon dioxide from the gas stream. In various embodiments, the concentration of the chemically modified carbonic anhydrase polypeptide and/or the CO₂ absorption mediating compound in the homogenous liquid formulation can be adjusted depending on the suitable conditions for the particular method of use. Various methods of use for carbon capture processes of the chemically modified carbonic anhydrase polypeptides, and the soluble compositions and homogenous liquid formulation that comprise them, are described in greater detail below, including suitable

conditions of polypeptide and CO₂ absorption mediating compound concentration, and temperature.

In some embodiments, the improved property of the chemically modified carbonic anhydrase polypeptides (and soluble compositions and homogeneous liquid formulations comprising them) disclosed herein is increased stability in the presence of compounds in the enzyme solution that improve the ability of the solution to absorb carbon dioxide (i.e., compounds that mediate the absorption of CO₂ by the solution). Such CO₂ absorption mediating compounds increase the amount of carbon dioxide that the solution can absorb, increase the rate at which carbon dioxide is absorbed, and/or improve the thermodynamic properties of the solution that control the carbon dioxide absorption or desorption. Accordingly, the chemically modified carbonic anhydrases, soluble compositions, and homogenous liquid formulations disclosed herein are advantageous for use in methods for carbon dioxide capture and sequestration that use solutions into which carbon dioxide is absorbed (i.e., captured by diffusing from gas stream into the liquid solution) and/or from which carbon dioxide is desorbed (i.e., extracted by diffusing from liquid solution into gas phase). Such compounds, solutions, and solvent systems for the absorption and/or desorption of carbon dioxide and the associated processes of using them for carbon dioxide capture from gas streams are described in e.g., U.S. Pat. Nos. 6,143,556, 6,524,843 B2, 7,176,017 B2, 7,596,952 B2, 7,641,717 B2, 7,579,185 B2, 7,740,689 B2, 7,132,090 B2; U.S. Pat. Publ. Nos. 2007/0256559A1, 2009/0155889A1, 2010/0086983A1; PCT Publ. Nos. WO2006/089423A1, WO2008/072979A1, WO2009/000025A1, WO2010/020017A1, WO2010/014773A1, WO2010/045689A1, each of which is hereby incorporated by reference herein.

In some embodiments, the improved property of the chemically modified carbonic anhydrase polypeptides, soluble compositions, and homogenous liquid formulations of the present disclosure is increased stability in the presence of an amine compound in the enzyme solution. In addition to increased stability to the presence of amine compound, in such embodiments the carbonic anhydrase can have increased thermostability, e.g., increased activity at temperatures above 40° C. The chemically modified carbonic anhydrase polypeptides disclosed herein having increased stability to amine compounds and increased solution temperature are particularly advantageous for use in methods for carbon dioxide capture and sequestration from flue gas streams using solutions comprising amine compounds (see e.g., U.S. Pat. No. 7,740,689 B2, and U.S. Pat. Publ. 2009/0155889 A1, each of which is hereby incorporated by reference herein) such as those amine compounds selected from the group consisting of: 2-(2-aminoethylamino)ethanol (AEE), 2-amino-2-hydroxymethyl-1,3-propanediol (AHPD), 2-amino-2-methyl-1-propanol (AMP), diethanolamine (DEA), diisopropanolamine (DIPA), N-hydroxyethylpiperazine (HEP), N-methyldiethanolamine (MDEA), monoethanolamine (MEA), N-methylpiperazine (MP), piperazine, piperidine, 2-(2-tert-butylaminoethoxy)ethanol (TBEE), triethanolamine (TEA), triisopropanolamine (TIA), tris, 2-(2-aminoethoxy)ethanol, 2-(2-tert-butylaminopropoxy)ethanol, 2-(2-tert-amylaminoethoxy)ethanol, 2-(2-isopropylaminopropoxy)ethanol, and 2-(2-(1-methyl-1-ethylpropylamino)ethoxy)ethanol.

In some embodiments, the improved property of the chemically modified carbonic anhydrase polypeptides, soluble compositions, and homogenous liquid formulations disclosed herein is increased stability in the presence of ammonia in the enzyme solution. In addition to increased

stability to the presence of ammonia, in such embodiments the carbonic anhydrase can have increased stability at increased or decreased temperatures (e.g., less than about 15° C.). The chemically modified carbonic anhydrases disclosed herein having increased stability to ammonia and/or increased thermostability are particularly advantageous for use in methods for carbon dioxide capture and sequestration from flue gas streams using solutions comprising ammonia, such as the chilled ammonia processes (see e.g., U.S. Pat. No. 7,641,717 B2, U.S. Pat. Publ. 2009/0155889 A1, each of which is hereby incorporated by reference herein).

8.3. Preparation of Chemically Modified Carbonic Anhydrase Polypeptides

The present disclosure also provides methods the carbonic anhydrase polypeptides that are chemically modified by treatment with a cross-linking agent, and the soluble compositions and homogenous liquid formulations comprising these chemically modified carbonic anhydrase polypeptides.

In some embodiments the present disclosure provides a method for preparing a chemically modified carbonic anhydrase comprising contacting in a solution: (i) a carbonic anhydrase polypeptide; and (ii) a cross-linking agent selected from the group consisting of a dialdehyde, a bis-imidate ester, a bis(N-hydroxysuccinimide) ester, a diacid chloride, and mixtures thereof.

In various embodiments of the method of preparation, the polypeptide used can comprise any of the wild-type or recombinant carbonic anhydrase polypeptides disclosed herein as useful for chemical modification. Thus, in some embodiments of the method of preparation, the polypeptide comprises an α -class carbonic anhydrase from human (*Homo sapiens*), rat (*Rattus norvegicus*), cow (*Bos taurus*), chicken (*Gallus gallus*), fish (*Cyprino carpio*), or the bacteria, *Neisseria gonorrhoeae*, or a recombinant carbonic anhydrase polypeptide derived from any one of these α -class carbonic anhydrase. For example, the polypeptide used in the method of preparation can comprise the α -class carbonic anhydrase comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 1298, 1300, 1302, 1304, 1306, and 1308, or a recombinant carbonic anhydrase polypeptide derived from any one of these α -class carbonic anhydrase sequences.

In other embodiments, the polypeptide used in the method of preparation can comprise a recombinant β -class carbonic anhydrase polypeptide derived from the wild-type *Desulfovibrio vulgaris* carbonic anhydrase comprising the amino acid sequence of SEQ ID NO: 2, or derived from a sequence homolog of SEQ ID NO: 2 selected from the group consisting of SEQ ID NO: 1288, 1290, 1292, 1294, and 1296. In some embodiments, the polypeptide used in the method of preparation comprises an amino acid sequence having at least 80% identity to SEQ ID NO:2. Exemplary engineered polypeptides useful in the methods of preparation are provide below in Tables 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J. In some embodiments of the method of preparation, the carbonic anhydrase polypeptide amino acid sequence comprises an even-numbered amino acid sequence selected from any one of SEQ ID NO: 4-1286. In some embodiments, the carbonic anhydrase polypeptide amino acid sequence comprises at least the following amino acid residue difference relative to SEQ ID NO: 2: X56S. In some embodiments, the carbonic anhydrase polypeptide amino acid sequence comprises at least the following amino acid residue difference relative to SEQ ID NO: 2: X30R, X40L, X56S, X84Q, X120R, and X139M. In some embodiments, the carbonic anhydrase

polypeptide amino acid sequence an amino acid sequence selected from any one of SEQ ID NO: 26, 190, 206, 238, 252, 270, 274, 284, 306, 318, 328, 332, 340, 354, 596, 606, 656, 678, 1080, 1110, 1148, 1152, 1156, and 1158.

In addition to using a range of polypeptides, the method of preparation of the chemically modified carbonic anhydrase polypeptides, soluble compositions, and homogenous liquid formulations disclosed herein can be carried using a range of cross-linking agents and associated reaction conditions.

In some embodiments of the methods of preparation, the cross-linking agent used is selected from the group consisting of a dialdehyde, a bis-imidate ester, a bis(N-hydroxysuccinimide) ester, a diacid chloride, and mixtures thereof. In some embodiments, the specific cross-linking agent is selected from the group consisting of malondialdehyde, glutaraldehyde, dimethyl suberimide, dimethyl pimelimidate, suberic acid bis(N-hydroxysuccinimide), and mixtures thereof.

In some embodiments of the methods of preparation, the cross-linking agent used is a dialdehyde having optionally one or more carbon atoms between the two aldehyde groups, for example wherein the dialdehyde is selected from the group consisting of glyoxal, succindialdehyde, malondialdehyde, glutaraldehyde, and mixtures thereof. In a particular embodiment, the cross-linking agent is glutaraldehyde.

In some embodiments of the methods of preparation, the cross-linking agent used is an imidate ester, and in particular embodiments, a bis-imidate ester having optionally one or more carbon atoms between the two imidate ester groups. Useful imidate esters include bis-imidate esters having optionally one or more carbon atoms between the two imidate ester groups, including but not limited to: imidate esters (such as methyl or ethyl) of malonimide, succinimide, glutarimide, adipimide, pimelimide, and suberimide.

In some embodiments of the methods of preparation, the cross-linking agent used is a bis(N-hydroxysuccinimide) ester of a di-carboxylic acid that forms an irreversible chemical modification of the polypeptide. Useful bis(N-hydroxysuccinimide) esters include those prepared from di-carboxylic acid selected from the group consisting of malonate, succinate, glutarate, adipate, pimelate, suberate, and mixtures thereof. Accordingly, in particular embodiments of the soluble composition, the cross-linking agent is a bis(N-hydroxysuccinimide) ester of a di-carboxylic acid selected from the group consisting of malonate, succinate, glutarate, adipate, pimelate, suberate, and mixtures thereof.

Various exemplary reaction conditions useful in the methods of preparing the chemically modified carbonic anhydrase polypeptides are disclosed in greater detail below (see e.g., Examples). Generally, the various embodiments of the methods for preparing comprise contacting in a solution the unmodified carbonic anhydrase polypeptide and the cross-linking agent in an aqueous solution each at a specified concentration. Typically, in the methods of preparation the concentration of the cross-linking agent used in the solution ranges from about 0.1% to about 5% and the concentration of the carbonic anhydrase polypeptide in the solution is from about 10 g/L to about 100 g/L. For example, in particular embodiment of the method of preparation, the concentrations of the cross-linking agent and polypeptide, respectively, are selected from: 0.25% (v/v) and 100 g/L; 0.25% (v/v) and 50 g/L; 0.25% (v/v) and 25 g/L; 0.50% (v/v) and 25 g/L; 0.75% (v/v) and 25 g/L; 1.0% (v/v) and 25 g/L, and 0.25% (v/v) and 10 g/L.

In some embodiments of the method of preparation, the concentration of cross-linking agent in the solution is from about 0.025% to about 10%, from about 0.05% to about 5%, from about 0.1% to about 5%, or from about 0.25% to about

2.5%. In some embodiments, the concentration of cross-linking agent in the solution is at a concentration of at least about 0.025%, at least about 0.1%, at least about 0.25%, at least about 0.5%, at least about 1%, at least about 2%, at least about 2.5%, or at least about 5%. As noted above, either percent volume/volume (v/v) or weight/volume (w/v) can be used with the cross-linking agents disclosed herein without a significant difference in performance of the methods for the purposes disclosed herein. Where the cross-linking agent typically is obtained as a liquid reagent, percent (v/v) is used. For example, as shown in the Examples, glutaraldehyde is obtained from Sigma-Aldrich (St. Louis, USA) as a 25% stock solution and this is further diluted based on percentage (v/v) to the desired concentration for the solution used in the method of preparation. However, where the cross-linking agent typically obtained as a solid reagent a percent (w/v) solution can be used.

In some embodiments of the method of preparation, the concentration of the carbonic anhydrase polypeptide in the solution is from about 0.1 g/L to about 100 g/L, from about 1 g/L to about 100 g/L, or from about 10 g/L to about 100 g/L. In some embodiments, the concentration of the carbonic anhydrase polypeptide in the solution is at least about 0.1 g/L, at least about 1 g/L, at least about 5 g/L, at least about 10 g/L, at least about 25 g/L, at least about 50 g/L, or at least about 100 g/L.

In some embodiments of the method of preparation, the cross-linking agent is glutaraldehyde, and the concentration of the cross-linking agent in the solution is about 0.25% and the concentration of the carbonic anhydrase polypeptide in the solution is from about 10 g/L to about 100 g/L. In particular embodiments of the method of preparation, the concentrations of glutaraldehyde cross-linking agent and polypeptide, respectively, are selected from: 0.25% (v/v) and 100 g/L; 0.25% (v/v) and 50 g/L; 0.25% (v/v) and 25 g/L; 0.50% (v/v) and 25 g/L; 0.75% (v/v) and 25 g/L; 1.0% (v/v) and 25 g/L, and 0.25% (v/v) and 10 g/L.

In some embodiments, the method of preparation can be carried out wherein the solution in prepared by adding a carbonic anhydrase polypeptide to an aqueous solution in the form of a powder. The powder may contain the polypeptide in a partially purified or a highly purified form prepared from cell extracts or cell lysates (e.g., shake-flask powder, or DSP powder). In some embodiments, the cell extracts or cell lysates used may be partially purified by precipitation (ammonium sulfate, polyethyleneimine, heat treatment or the like, followed by a desalting procedure prior to lyophilization (e.g., ultrafiltration, dialysis, and the like). Any of the cell preparations may be stabilized by crosslinking using known crosslinking agents, such as, for example, glutaraldehyde or immobilization to a solid phase (e.g., Eupergit C, and the like) or by the crosslinking of protein crystals or precipitated protein aggregate particles.

Other conditions used in the embodiments of the method of preparation include an incubation time of 1 h to 4 h, and an incubation temperature of about room temperature (e.g., about 25° C.), or about 20° C. to about 30° C. Additionally, the aqueous solution may further comprise buffer salts at concentrations typically used with the particular polypeptide (e.g., triethanolamine sulfate or sodium bicarbonate at about pH 7.7 to about pH 10).

8.4. Carbonic Anhydrase Polypeptides Useful for Chemical Modification

The present disclosure provides carbonic anhydrase polypeptides that are chemically modified by treatment with

a cross-linking agent and resulting in the surprising advantage of improved properties of increased carbonic anhydrase activity and/or increased stability under conditions useful for carbon capture process, e.g., in homogenous liquid formulations with high concentrations of CO₂ absorption mediating compounds and temperatures significantly increased or decreased above/below ambient temperature. The present disclosure contemplates that chemically modified carbonic anhydrase polypeptides can be prepared using any naturally occurring or recombinant (engineered) carbonic anhydrase polypeptide. Accordingly, in some embodiments of the compositions, formulations and methods of the present disclosure, the carbonic anhydrase that is chemically modified is selected from an α -class, γ -class, β -class, or ξ -class carbonic anhydrase.

In some embodiments, the present disclosure provides particular emphasis on the use of carbonic anhydrase polypeptides that already exhibit favorable properties of increased activity and stability under carbon capture process conditions. The α -class carbonic anhydrases, particularly the human carbonic anhydrase II (“HuCAII”), are among the fastest known enzymes, and generally have the highest specific carbonic anhydrase activity of the various classes. Accordingly, in some embodiments of the compositions, formulations and methods of the present disclosure, the carbonic anhydrase that is chemically modified is an α -class carbonic anhydrase polypeptide or a recombinant carbonic anhydrase polypeptide derived from an α -class carbonic anhydrase. In some embodiments, the α -class carbonic anhydrase that is chemically modified is an α -class carbonic anhydrase from human (*Homo sapiens*), rat (*Rattus norvegicus*), cow (*Bos taurus*), chicken (*Gallus gallus*), fish (*Cyprino carpio*), or the bacteria, *Neisseria gonorrhoeae*, or a recombinant carbonic anhydrase polypeptide derived from any one of these α -class carbonic anhydrase. In some embodiments, the α -class carbonic anhydrase that is chemically modified comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 1298, 1300, 1302, 1304, 1306, and 1308, or a recombinant carbonic anhydrase polypeptide derived from any one of these α -class carbonic anhydrase sequences.

Although naturally occurring β -class carbonic anhydrases have been found that exhibit relatively high thermostability (e.g., β -class carbonic anhydrase from *Methanobacterium thermoautotrophicum*), most β -class enzymes exhibit significantly lower specific activity in catalyzing the hydration of CO₂ than the α -class carbonic anhydrases such as the α -class HuCAII of SEQ ID NO: 1298. For example, in a bicarbonate dehydration assay at pH 8.0, 25° C., the β -class CA from *H. thermoautotrophicum* exhibits less than 4% of the specific activity of the α -class HuCAII. However, the β -class carbonic anhydrase from *Desulfovibrio vulgaris* exhibits a high specific activity that is comparable to an α -class enzymes and also exhibits high thermostability. For example, in the same bicarbonate dehydration assay comparison to the α -class HuCAII of SEQ ID NO: 1298, the wild-type β -class carbonic anhydrase from *D. vulgaris* of SEQ ID NO: 2 was exhibited 84% of the specific activity of HuCAII, and more than 20-fold the activity exhibited by β -class CA from *H. thermoautotrophicum*. Several naturally occurring β -class carbonic anhydrases have been identified that are sequence homologs having over 40% identity to the *D. vulgaris* enzyme of SEQ ID NO: 2. These include a β -class carbonic anhydrases from: *Desulfovibrio* sp. FW1012B (SEQ ID NO: 1288); *Desulfo- microbium baculatum* strain DSM 4028 (SEQ ID NO: 1290); *Desulfovibrio aespoeensis* (SEQ ID NO: 1292); *Des-*

ulfovibrio desulfuricans strain G20 (SEQ ID NO: 1294); and *Desulfovibrio magneticus* strain ATCC 700980 (SEQ ID NO: 1296).

Accordingly, in some embodiments of the compositions, formulations and methods of the present disclosure, the carbonic anhydrase that is chemically modified is a β -class carbonic anhydrase polypeptide or a recombinant carbonic anhydrase polypeptide derived from a β -class carbonic anhydrase. In some embodiments, the β -class carbonic anhydrase comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 2, 1288, 1290, 1292, 1294, and 1296. In some embodiments, the β -class carbonic anhydrase is a recombinant carbonic anhydrase polypeptide derived from a β -class carbonic anhydrase from *Desulfovibrio vulgaris*, and in some embodiments comprises an amino acid sequence having at least 80% identity to SEQ ID NO: 2.

As described in greater detail below, the β -class carbonic anhydrase from *Desulfovibrio vulgaris* of SEQ ID NO: 2 has been further engineered to provide recombinant carbonic anhydrase polypeptides having an improved property when compared with the naturally-occurring, wild type carbonic anhydrase enzyme obtained from *Desulfovibrio vulgaris* (SEQ ID NO: 2). These recombinant carbonic anhydrase polypeptides comprise one or more differences in their amino acid sequence (e.g., substitutions, insertions, and/or deletions) relative to a reference sequence (e.g., *Desulfovibrio vulgaris* CA polypeptide of SEQ ID NO: 2) that result in a carbonic anhydrase polypeptide having an improved property. The improved properties of these recombinant CA polypeptides include, but are not limited to, activity (e.g., hydration of carbon dioxide, or dehydration of bicarbonate), thermal stability, solvent stability, pH activity profile, refractoriness to inhibition or inactivation by other compounds in the solution with the enzyme, e.g. inhibition by bicarbonate, carbonate, amine compounds, ammonia, flue gas components (such as NO_x and SO_x compounds).

In some embodiments, the improved property of the engineered carbonic anhydrase polypeptide is with respect to an increase in its rate of conversion of the substrate to the product (e.g., hydration of carbon dioxide to bicarbonate). This improvement in enzymatic activity can be manifested by the ability to use less of the recombinant polypeptide as compared to a reference polypeptide and thereby reduce the amount of enzyme needed to convert the same amount of product.

In some embodiments, the improved property of the engineered carbonic anhydrase polypeptide is with respect to its thermostability. Accordingly, in some embodiments the recombinant carbonic anhydrase polypeptides have an improved property that comprises at least 1.2-fold, at least 1.3-fold, at least 1.5-fold, at least 2-fold, at least 3-fold, at least 4-fold, at least 5-fold, at least 10-fold, or at least 25-fold increased thermostability. In such embodiments, increased thermostability can be determined as increased activity relative to a reference polypeptide following exposure to thermostability challenge conditions—e.g., exposure to 30, 40, 50, or 60° C. solution for a defined time period, such as 24 h. In some embodiments, the carbonic anhydrase polypeptide has more than one improved property, such as a combination of increased enzyme activity and thermostability.

The present disclosure contemplates that any of these engineered carbonic anhydrase polypeptides having improved properties can be chemically modified by treatment with a cross-linking agent and used in the methods of carbon capture disclosed herein.

Exemplary recombinant carbonic anhydrase polypeptides useful for chemical modification according to the present disclosure include but are not limited to, the polypeptides that comprise the amino acid sequences corresponding to any one of SEQ ID NOS: 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28,

30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 176, 178, 180, 182, 184, 186, 188, 190, 192, 194, 196, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 228, 230, 232, 234, 236, 238, 240, 242, 244, 246, 248, 250, 252, 254, 256, 258, 260, 262, 264, 266, 268, 270, 272, 274, 276, 278, 280, 282, 284, 286, 288, 290, 292, 294, 296, 298, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 340, 342, 344, 346, 348, 350, 352, 354, 356, 358, 360, 362, 364, 366, 368, 370, 372, 374, 376, 378, 380, 382, 384, 386, 388, 390, 392, 394, 396, 398, 400, 402, 404, 406, 408, 410, 412, 414, 416, 418, 420, 422, 424, 426, 428, 430, 432, 434, 436, 438, 440, 442, 444, 446, 448, 450, 452, 454, 456, 458, 460, 462, 464, 466, 468, 470, 472, 474, 476, 478, 480, 482, 484, 486, 488, 490, 492, 494, 496, 498, 500, 502, 504, 506, 508, 510, 512, 514, 516, 518, 520, 522, 524, 526, 528, 530, 532, 534, 536, 538, 540, 542, 544, 546, 548, 550, 552, 554, 556, 558, 560, 562, 564, 566, 568, 570, 572, 574, 576, 578, 580, 582, 584, 586, 588, 590, 592, 594, 596, 598, 600, 602, 604, 606, 608, 610, 612, 614, 616, 618, 620, 622, 624, 626, 628, 630, 632, 634, 636, 638, 640, 642, 644, 646, 648, 650, 652, 654, 656, 658, 660, 662, 664, 666, 668, 670, 672, 674, 676, 678, 680, 682, 684, 686, 688, 690, 692, 694, 696, 698, 700, 702, 704, 706, 708, 710, 712, 714, 716, 718, 720, 722, 724, 726, 728, 730, 732, 734, 736, 738, 740, 742, 744, 746, 748, 750, 752, 754, 756, 758, 760, 762, 764, 766, 768, 770, 772, 774, 776, 778, 780, 782, 784, 786, 788, 790, 792, 794, 796, 798, 800, 802, 804, 806, 808, 810, 812, 814, 816, 818, 820, 822, 824, 826, 828, 830, 832, 834, 836, 838, 840, 842, 844, 846, 848, 850, 852, 854, 856, 858, 860, 862, 864, 866, 868, 870, 872, 874, 876, 878, 880, 882, 884, 886, 888, 890, 892, 894, 896, 898, 900, 902, 904, 906, 908, 910, 912, 914, 916, 918, 920, 922, 924, 926, 928, 930, 932, 934, 936, 938, 940, 942, 944, 946, 948, 950, 952, 954, 956, 958, 960, 962, 964, 966, 968, 970, 972, 974, 976, 978, 980, 982, 984, 986, 988, 990, 992, 994, 996, 998, 1000, 1002, 1004, 1006, 1008, 1010, 1012, 1014, 1016, 1018, 1020, 1022, 1024, 1026, 1028, 1030, 1032, 1034, 1036, 1038, 1040, 1042, 1044, 1046, 1048, 1050, 1052, 1054, 1056, 1058, 1060, 1062, 1064, 1066, 1068, 1070, 1072, 1074, 1076, 1078, 1080, 1082, 1084, 1086, 1088, 1090, 1092, 1094, 1096, 1098, 1100, 1102, 1104, 1106, 1108, 1110, 1112, 1114, 1116, 1118, 1120, 1122, 1124, 1126, 1128, 1130, 1132, 1134, 1136, 1138, 1140, 1142, 1144, 1146, 1148, 1150, 1152, 1154, 1156, 1158, 1160, 1162, 1164, 1166, 1168, 1170, 1172, 1174, 1176, 1178, 1180, 1182, 1184, 1186, 1188, 1190, 1192, 1194, 1196, 1198, 1200, 1202, 1204, 1206, 1208, 1210, 1212, 1214, 1216, 1218, 1220, 1222, 1224, 1226, 1228, 1230, 1232, 1234, 1236, 1238, 1240, 1242, 1244, 1246, 1248, 1250, 1252, 1254, 1256, 1258, 1260, 1262, 1264, 1266, 1268, 1270, 1272, 1274, 1276, 1278, 1280, 1282, 1284, and 1286.

Structure and function information for these exemplary recombinant carbonic anhydrase polypeptides of the present disclosure are shown below in Tables 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J. The odd numbered sequence identifiers (i.e., SEQ ID NOS) refer to the nucleotide sequence encoding the amino acid sequence provided by the even numbered SEQ ID NOS, and the sequences are provided in the electronic sequence listing file accompanying this disclosure, which is hereby incorporated by reference herein. The amino acid residue differences are based on comparison to the reference sequence of SEQ ID NO: 2, which is a wild type carbonic anhydrase from *Desulfovibrio vulgaris* str. "Miyazaki F" having GenBank accession ACL09337.1 GI:218758438.

Tables 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J also disclose the increased stability (solvent and/or thermostabil-

ity) in the presence of an amine compound (MDEA) and/or ammonia at various concentrations and temperatures relative to the reference polypeptide of SEQ ID NO: 2. Generally, increased stability was determined by measuring the relative rate of dehydrating bicarbonate to carbon dioxide in a high-throughput (HTP) assay following 24 h exposure to the specified solvent and temperature challenge conditions, and HTP activity assays were carried out in 96-well plate format assay using cell lysates containing the engineered polypeptides. General HTP challenge/assay conditions were as follows: 25 μL of cleared *E. coli* lysate added to 75 μL of challenge buffer solution (e.g., solution containing 4.0 M-6.66 M MDEA or NH₃) and incubated at the challenge temperature (e.g., 30°, 35°, 42°, 50° or 55° C.) for 24 h; followed by adding a 10 μL aliquot of the challenge solution to 190 μL of bicarbonate dehydration assay solution (200 mM KHCO₃, 400 μM phenolphthalein, pH 7 or 8) at 25° C. or 45° C., measuring carbonic anhydrase activity as slope of phenolphthalein indicator absorbance change at 550 nm over 20-30 minutes. Additional HTP assay details are provided in Example 1. As noted in Tables 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J, the measured level of increased activity of each engineered polypeptide relative to a reference polypeptide was classified as “+”, “++”, or “+++” for the different assays.

TABLE 2A

SEQ ID NO: (nt/aa)	Amino Acid Residue	Assay 1 (24 h/ 42° C./ 3M MDEA challenge/ 25° C. assay)	Assay 2 (24 h/ 50° C./ 3M MDEA challenge/ 25° C. assay)	Assay 3 (24 h/ 30° C./ 4.2M NH ₃ /CO ₂ challenge/ 25° C. assay)	Assay 4 (24 h/ 35° C./ 4.2M NH ₃ /CO ₂ challenge/ 25° C. assay)
3/4	K147E;	+++	+++		
5/6	T30R;	+++		++	+++
7/8	T139M;	+++	++		
9/10	G120R;	+++			
11/12	T30Q;	+++		++	++
13/14	T4F;	+++	+	+	++
15/16	A84Q;	+++	+++	++	++
17/18	Q119M;	+++			
19/20	L34H;	+++		++	++
21/22	T4M;	++		+++	+++
23/24	K147T;	++	+++		
25/26	A56S;	++	+++	++	++
27/28	Q32K;	++		+	++
29/30	V131L;	++	+	+	
31/32	Q15R;	++	++	+++	+++
33/34	N145W;	++			
35/36	R165;	++			
37/38	A40W;	++		+	+
39/40	N213E;	++			
41/42	H222C;	++		++	+++
43/44	E142L;	++			
45/46	G2T;	++	++		
47/48	R31P;	++	+		
49/50	S144L;	++			
51/52	E159H;	++			
53/54	T139Q;	++			
55/56	H148T;	++			
57/58	M170F;	++	+	+	
59/60	D86A				
61/62	A121K;	++	+	+	
63/64	N145F;	++			
65/66	Q32R;	++		++	+
67/68	A121W;	++	+	+	++
69/70	K37R;	++		+	++
71/72	A221C;	++		++	++
73/74	A84S;	++			
75/76	E200R;	++			
77/78	T139K;	++			

TABLE 2A-continued

SEQ ID NO: (nt/aa)	Amino Acid Residue	Assay 1 (24 h/ 42° C./ 3M MDEA challenge/ 25° C. assay)	Assay 2 (24 h/ 50° C./ 3M MDEA challenge/ 25° C. assay)	Assay 3 (24 h/ 30° C./ 4.2M NH ₃ /CO ₂ challenge/ 25° C. assay)	Assay 4 (24 h/ 35° C./ 4.2M NH ₃ /CO ₂ challenge/ 25° C. assay)
79/80	A95V;	++		+	+
81/82	A84N;	+		+	
83/84	Q43M;	+			
85/86	A121V;	+			
87/88	K147G;	+			
89/90	R223C;	+		++	++
91/92	T30A;	+		++	++
93/94	G2R;	+		+	+
95/96	A121H;	+		+	
97/98	A121Q;	+		+	+
99/100	A60C;	+		++	++
101/102	D96C;	+			
103/104	T30L;	+		++	+
105/106	A40L;	+	+	+	
107/108	H97F;	+			
109/110	E68A;	+		+	+
111/112	S42A;	+		+	+
113/114	V70I;	+		++	++
115/116	Q119T;	+			
117/118	D96E;	+		+	
119/120	S35A;	+		+	++
121/122	H124G;	+		+	
123/124	Q119K;	+			
125/126	V138L;	+			
127/128	D168E;	+			
129/130	T139H;	+			
131/132	A121T;	+		+	+
133/134	A121L;	+			
135/136	S144A;	+			
137/138	N145C;	+			
139/140	N213Q;	+			
141/142	D96K;	+			
143/144	A178G;	+			
145/146	H124R;	+			
147/148	D96A;	+			
149/150	S35R;	+		+	+
151/152	E159V;	+			
153/154	T47R;	+		+	
155/156	H148A;	+			
157/158	A84R;	+		++	++
159/160	Q43V;	+			
161/162	E159R;	+			
163/164	K147F;	+			
165/166	E68G;	+		+	+
167/168	V157A;	+		+	+
169/170	V138W;	+			
171/172	V138F;	+			
173/174	R223Q;	+			++
175/176	M207E;	+		+	
177/178	A84K;	+		++	+
179/180	A60V;	+		++	++
181/182	A40Q;	+		++	+
183/184	A22G;	+		++	
185/186	K143M;	+			
	M207N;	+			

60 Levels of increased activity were determined relative to the reference polypeptide of SEQ ID NO: 2 and defined as “+”, “++”, or “+++” for each of the four assays as follows:
 Assay 1: “+” = at least 1.3-fold but less than 2-fold increased activity; “++” = at least 2-fold but less than 3-fold increased activity; “+++” = at least 3-fold increased activity.
 Assay 2: “+” = at least 1.5-fold but less than 2-fold increased activity; “++” = at least 2-fold but less than 3-fold increased activity; “+++” = at least 3-fold increased activity.
 Assay 3: “+” = at least 1.3-fold but less than 1.5-fold increased activity; “++” = at least 1.5-fold but less than 2-fold increased activity; “+++” = at least 2-fold increased activity.
 Assay 4: “+” = at least 1.3-fold but less than 3-fold increased activity; “++” = at least 3-fold but less than 5-fold increased activity; “+++” = at least 5-fold increased activity.

29

TABLE 2B

SEQ ID NO: (nt/aa)	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 5	Assay 6	Assay 7	5
		(24 h/ 50° C./ 4M MDEA challenge/ 45° C. assay)	(24 h/ 50° C./ 5M MDEA challenge/ 25° C. assay)	(24 h/ 55° C./ 5M MDEA challenge/ 25° C. assay)	
187/188	T30R; R31P; A56S; A84Q;	++	+++	+++	10
189/190	A56S; A84Q; T139M;	++	+++	+++	
191/192	T30R; R31P; A40L; A56S; G120R;	+++	+++	+++	
193/194	R31P; A40L; A56S; G120R; T139M;	+++	+++	+++	
195/196	T30R; R31P; A56S; A84Q; Q119M;	+++	+++	+++	
197/198	R31P; A40L; A56S; A84Q;	+++	+++	+++	
199/200	T30Q; R31P; A56S; A84Q;	+++	+++	+++	
201/202	T30Q; L34H; A56S;	+++	+++	++	
203/204	T30R; R31P; A40L; A56S; K147T;	+++	+++	+++	
205/206	T30R; R31P; A56S; K147T;	+++	+++	+++	
207/208	T4F; A56S; A84Q;	++	++	++	
209/210	T30R; L34H; A56S;	++	++	++	
211/212	A56S; T139M;	+	+	+	
213/214	G2T; R31P; L34H; A40L; A56S; A84Q; T139M;	+	+	++	
215/216	T4F; L34H; A56S; G120R; K147E;	+	+	+	
217/218	A40L; A56S;	++	+	+	
219/220	R31P; A40L; A56S; Q119M; G120R;	++	+	++	
221/222	R31P; A56S; G120R; K147E;	++	+	+	
223/224	T4F; A40L; A56S; K147T;	++	+	+	
225/226	R31P; A40L; A56S;	++	+	+	
227/228	A56S; A84Q;	++	+	+	
229/230	T30R; A40L; A56S;	++	+	++	
231/232	T30Q; L34H; A56S; K147T;	++	+	++	
233/234	L34H; A56S;	++	+	+	
235/236	T30R; R31P; A56S;	+++	+	++	
237/238	T30R; A56S;	+++	+	++	
239/240	R31P; A56S; A84Q;	+++	+	++	
241/242	T4F; A56S;	+	+	+	
243/244	G2T; A56S; T139M;	+	+	+	
245/246	A56S; G120R; K147T;	+	+	+	
247/248	G2T; A56S; A84Q; T139M;	+	+	+	
249/250	A56S; Q119L; G120R;	+	+	+	
251/252	A40L; A56S; G120R;	+	+	+	
253/254	A56S; K147T;	+	+	+	
255/256	A40L; A56S; T139M; K147E;	+	+	+	
257/258	A40L; A56S; T139M;	++	+	+	
259/260	T4F; T30Q; A56S; G120R; T139M;	++	+	+	
261/262	L34H; A56S; A84Q; T139M;	++	+	++	
263/264	A56S; A84Q; K147E;	++	+	+	
265/266	A56S; A84Q; G120R;	++	+	+	
267/268	T30R; R31P; A56S; T139M;	++	+	++	

Levels of increased activity were determined relative to the reference polypeptide of SEQ ID NO: 26 (i.e., engineered polypeptide having A56S) and defined as "+", "++", or "+++" for each of the assays as follows:

Assay 5: "+" indicates at least 1.5-fold but less than 2.5-fold increased activity; "++" indicates at least 2.5-fold but less than 4-fold increased activity; "+++" indicates at least 4-fold increased activity.

Assay 6: "+" indicates at least 1.3-fold but less than 1.7-fold increased activity; "++" indicates at least 1.7-fold but less than 2-fold increased activity; "+++" indicates at least 2-fold increased activity.

Assay 7: "+" indicates at least 1.5-fold but less than 2.5-fold increased activity; "++" indicates at least 2.5-fold but less than 4-fold increased activity; "+++" indicates at least 4-fold increased activity.

30

TABLE 2C

SEQ ID NO: (nt/aa)	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 5		5
		(24 h/ 50° C./ 4M MDEA challenge/ 45° C. assay)	(24 h/ 50° C./ 5M MDEA challenge/ 45° C. assay)	
269/270	T30Q; A40L; A56S; A84Q;	+++	+++	10
271/272	A40L; A56S; A84Q; G120R; K147E;	+++	+++	
273/274	T30R; R31P; L34H; A40L; A56S; A84Q; G120R; T139M; K147T;	+++	+++	
275/276	R31P; A40L; A56S; Q119M; T139M; K147T;	+++	+++	
277/278	G2T; A40L; A56S; Q119M; G120R; K147T;	++	++	
279/280	G2T; T30Q; L34H; A56S; Q119M; K147T;	++	++	
281/282	T30R; A40L; A56S; A84Q; G120R; K147T;	+++	+++	
283/284	L34H; A56S; T139M; K147T;	+++	+++	
285/286	R31P; A40L; A56S; K147E;	++	++	
287/288	A40L; A56S; G120R; T139M; K147E;	++	++	
289/290	G2T; T30Q; A40L; A56S; A84Q; K147E;	++	++	
291/292	A56S; A84Q; G120R; T139M; K147E;	+++	+++	
293/294	T4F; T30Q; A56S; A84Q; T139M;	+++	+++	
295/296	T30Q; A56S; A84Q; Q119M; G120R; T139M; K147E;	+++	+++	
297/298	G2T; T4F; T30Q; R31P; A40L; A56S; A84Q;	+++	+++	
299/300	T4F; A40L; A56S; A84Q; T139M;	+++	+++	
301/302	G2T; T30R; A56S; Q119M; T139M;	++	++	
303/304	A56S; A84Q; G120R; T139M; K147T;	++	++	
305/306	T30R; R31P; A40L; A56S; T139M;	+++	+++	
307/308	T4F; A56S; A84Q; T139M;	+++	+++	
309/310	T30R; R31P; A56S; Q119M; G120R;	+++	+++	
311/312	G2T; T4F; A40L; A56S; Q119M; G120R; T139M;	++	++	
313/314	R31P; A56S; A84Q; Q119M;	++	++	
315/316	G2T; A40L; A56S; Q119M; G120R;	+	+	
317/318	R31P; A56S; A84Q; G120R; T139M;	+++	+++	
319/320	T4F; T30Q; R31P; A56S; G120R; K147T;	+++	+++	
321/322	T30Q; R31P; L34H; A56S; A84Q; T139M;	+++	+++	
323/324	T4F; T30Q; R31P; L34H; A40L; A56S; Q119M; T139M; K147T;	+++	+++	
325/326	L34H; A56S; G120R; K147E;	++	++	
327/328	G2T; A40L; A56S; A84Q; G120R; T139M; K147T;	++	++	
329/330	A56S; G120R;	+	+	
331/332	T30R; A40L; A56S; A84Q; G120R; T139M;	++	++	
333/334	T30R; A40L; A56S; A84Q; G120R; T139M; K147E;	+++	+++	
335/336	R31P; L34H; A56S; G120R;	+++	+++	
337/338	R31P; L34H; A40L; A56S; T139M; K147T;	+++	+++	
339/340	G2T; T30R; A40L; A56S;	++	++	
341/342	G2T; T30R; A56S; G120R; T139M; K147E;	+++	+++	
343/344	T30R; A56S; Q119M; G120R;	++	++	
345/346	T4F; T30Q; L34H; A40L; A56S; A84Q; G120R; T139M;	+++	+++	
347/348	T30R; R31P; L34H; A40L; A56S; A84Q; Q119M; K147T;	+++	+++	
349/350	T4F; A40L; A56S; Q119M; G120R; K147T;	+++	+++	
351/352	G2T; T4F; T30R; R31P; L34H; A56S; G120R; T139M;	++	++	
353/354	T4F; T30R; A40L; A56S;	+++	+++	
355/356	R31P; L34H; A56S; T139M;	+++	+++	
357/358	L34H; A40L; A56S; A84Q; Q119M; T139M;	+++	+++	
359/360	G2T; T4F; L34H; A56S; A84Q; Q119M; G120R; K147E;	++	++	
361/362	R31P; A40W; A56S; A95V; N145W; K147T;	++	++	
363/364	R31P; A40W; A56S; T139Q; N145L; E159V; A221C;	+	+	
365/366	R31P; A40W; A56S; A95V; T139Q; N145W; E159V; N213E;	++	++	
367/368	A40W; A56S; A95V; N213E;	++	++	
369/370	R31P; A56S; A95V; T139K; N145F; K147E;	++	++	
371/372	A40W; A56S; A95V; V131F; T139K; K147E; E159V; N213E;	+	+	
373/374	R31P; A56S; V131F; K147E; E159H; A221C;	+	+	
375/376	A56S; V131F; T139K; N145L; E159V; A221C;	++	++	
377/378	R31P; A40W; A56S; A95V; A121L; A221C;	++	++	
379/380	R31P; A40W; A56S; A95V; T139Q;	+	+	
381/382	A40W; A56S; A95V; V131F; E159V; N213E;	+	+	
383/384	R31P; A40W; A56S; A121L; V131F; T139Q; E159H; N213E;	+	+	
385/386	A40W; A56S; A95V; E159V; N213E;	+	+	
387/388	A40W; A56S;	+	+	
389/390	R31P; A56S; A95V; A121W; A221C;	++	++	

31

TABLE 2C-continued

SEQ ID	Assay 5 (24 h/ 50° C./ 4M MDEA challenge/ 45° C. assay)	5
NO: Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)		
391/392	A40W; A56S; T139K;	+
393/394	A56S; A121V; T139K; N213E; A221C;	+
395/396	A56S; A121K;	+
397/398	A56S; T139K; N145F; E159H; A221C;	+
399/400	R31P; A40W; A56S; A95V; A121V; N145W; K147E; N213E;	++
401/402	R31P; A40W; A56S; A95V; A121K;	+
403/404	R31P; A40W; A56S; A95V; T139K; N145F; N213E;	++
405/406	R31P; A56S; A95V; A121W; T139K; N145F; K147T;	++
407/408	R31P; A40W; A56S; A95V; A121V; K147T; N213E;	++
409/410	A56S; A121V; T139K; K147T;	++
411/412	R31P; A40W; A56S; N145F;	++
413/414	A56S; A121W;	++
415/416	R31P; A40W; A56S; A95V; A121W; T139Q; E159V; N213E;	++
417/418	R31P; A40W; A56S; A221C;	+
419/420	R31P; A40W; A56S; A95V; N145F;	++
421/422	A40W; A56S; A95V; A121K; V131F; K147T;	+
423/424	A40W; A56S; V131F; N145F; K147E;	+
425/426	A40W; A56S; T139K; N145F; A221C;	++
427/428	A40W; A56S; A121V;	++
429/430	R31P; A40W; A56S; T139Q;	+
431/432	R31P; A56S; A121W; T139Q;	+
433/434	A40W; A56S; A95V;	+
435/436	R31P; A56S; A95V; V131F; T139K; K147T; E159H;	+
437/438	A56S; N213E; A221C;	+
439/440	A40W; A56S; T139Q; K147T; A221C;	++
441/442	R31P; A56S; A95V; T139K; K147E;	++
443/444	R31P; A40W; A56S; V131F; T139Q;	++
445/446	R31P; A56S; A95V; A121W;	+
447/448	A40W; A56S; A95V; A121L; N213E;	+
449/450	R31P; A56S; T139K;	++
451/452	A56S; A95V; A121V; N145F; K147E; E159V;	+
453/454	A56S; A95V; A121L; E159V; N213E;	++
455/456	R31P; A40W; A56S; A121W; N145F; A221C;	++
457/458	A40W; A56S; N145F; K147T; N213E;	++
459/460	R31P; A56S; A95V; E159V;	++
461/462	R31P; A56S; A95V; A121K; E159H; A221C;	++
463/464	R31P; A40W; A56S; A121K; V131F; T139Q; N213E;	+
465/466	A56S; V131F; K147T;	+
467/468	A56S; E159V; N213E; A221C;	++
469/470	S42A; T47R; A56S; E68A; A95V; V138L; A221C;	+
471/472	A56S; E68A; H97F; V138L; S144L;	+
473/474	S35A; A56S; E68A; H97F; S144L; A219T; A221C;	+
475/476	S35R; A56S; E68A; S144L;	+
477/478	S42A; A56S; H97F; H124G; A219T; A221C;	+
479/480	S42A; T47R; A56S; A95V; H97F; H124R; A219T;	+
481/482	A56S; H124R; S144L;	+
483/484	S42A; T47R; A56S; A95V; V157A;	+
485/486	S42A; T47R; A56S; E68A; A95V;	+
487/488	A56S; A95V; H97F; H124G; S144L; V157A;	+
489/490	S42A; T47R; A56S; E68A; A95V; H124G; V157A;	+
491/492	S42A; A56S; V70I; A95V; A221C;	+
493/494	A56S; V157A;	+
495/496	S35R; T47R; A56S; E68A; V70I; A95V;	+
497/498	T47R; A56S; A95V; H124R; A221C;	+
499/500	A56S; E68A; A95V; H97F; H124R; V138L; S144L; V157A;	+
501/502	T47R; A56S; E68A; H97F; V138L; A219T;	+
503/504	A56S; V138L; S144L;	+
505/506	S35A; S42A; T47R; A56S; E68A; V70I; H97F; V138L; S144L; D168E; A219T;	+
507/508	T47R; A56S; E68A; V70I; H124R; V138L; A219T;	+
509/510	S42A; A56S; E68A; H97F;	+
511/512	A56S; E68A; A95V; A221C;	+
513/514	A56S; V70I; A95V; H124G; V138L; S144L;	+
515/516	A56S; A95V; H97F;	+
517/518	S35R; S42A; A56S; E68A; V70I; A95V; H97F; V157A;	+
519/520	T47R; A56S; E68A; A95V; H97F; D168E; A219T; A221C;	+

32

TABLE 2C-continued

SEQ ID	Assay 5 (24 h/ 50° C./ 4M MDEA challenge/ 45° C. assay)	5
NO: Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)		
521/522	T47R; A56S; A95V; S144L; V157A; A221C;	+
523/524	S42A; A56S; A95V; H124R; V138L; A219T;	+
525/526	T47R; A56S; H97F; H124R; V138L; S144L; A219T;	+
527/528	S42A; T47R; A56S; E68A; A95V; V138L; A219T;	++
529/530	A56S; A95V; V138L; S144L; A221C;	++
531/532	T47R; A56S; V157A; A219T;	+
533/534	S35A; T47R; A56S; E68A; H97F; H124G; V138L; S144L;	+
535/536	S42A; T47R; A56S; E68A; H97F; H124G; S144L; V157A; A221C;	+
537/538	S42A; A56S; E68A; H124G; A219T;	+
539/540	A56S; E68A; A95V; V138L; A219T; A221C;	++
541/542	S35R; A56S; H124R; V138L; S144L; A219T;	+
543/544	S42A; A56S; E68A; V70I; H97F; D168E; A221C;	+
545/546	A56S; E68A; A221C;	+
547/548	A56S; A95V; V138L; A219T; A221C;	+
549/550	T47R; A56S; E68A; V70I; H124R; A219T;	+
551/552	A56S; E68A; V70I; H97F; H124R; V157A; A221C;	++
553/554	S35R; A56S; A95V; V157A;	+
555/556	A56S; H97F;	+
557/558	S42A; T47R; A56S; E68A; A221C;	+
559/560	A56S; A95V; H97F; V138L;	+
561/562	S35R; A56S; V138L; S144L; A221C;	+
563/564	S35R; A56S; E68A; H124R; S144L; A221C;	++
565/566	S35R; T47R; A56S; E68A; V70I; S144L; V157A; D168E; A219T; A221C;	++
567/568	S35R; T47R; A56S; A95V; H97F; H124R; V138L; A219T; A221C;	+

Levels of increased activity were determined relative to the reference polypeptide of SEQ ID NO: 26 (i.e., engineered polypeptide having A56S) using Assay 5 and defined as follows:
 "+" indicates at least 1.5-fold but less than 2.5-fold increased activity;
 "++" indicates at least 2.5-fold but less than 4-fold increased activity;
 "+++" indicates at least 4-fold increased activity.

SEQ ID	Assay 8 (24 h, 65° C., 5M MDEA challenge/ 45° C., 1M MDEA pH 8.0 assay)	40
NO: Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)		
569/570	G2T; T30R; A40L; Q43M; A56S; V70I; A84Q; G120R; T139M; M170F; H222C; R223C;	+
571/572	G2T; T30R; Q32K; A40L; Q43M; A56S; A84Q; H97F; G120R; T139M; E200R; H222C;	+
573/574	G2T; T30R; R31P; A40L; Q43M; A56S; A84Q; G120R; T139M; E142L; H148T; M170F; H222C;	+
575/576	G2T; T30R; A40L; A56S; V70I; A84Q; H97F; G120R; T139M; E142L; H148T; E200R; H222C; R223C;	+
577/578	T30R; R31P; Q32R; A40L; A56S; A84Q; G120R; T139M; E142L; E200R; H222C;	+
579/580	T30R; A40L; Q43M; A56S; V70I; A84Q; D96E; G120R; T139M; M170F; H222C;	+
581/582	T30R; R31P; K37R; A40L; S42A; A56S; A60C; E68A; A84Q; Q119M; G120R; H124F; T139M; N213E; A219T;	+
583/584	T30R; Q32K; A40L; A56S; V70I; A84Q; D96E; G120R; A121L; T139M; E200R;	+
585/586	T30R; A40L; A56S; A84Q; G120R; A121L; T139M; H148T; E200R; R223C;	+
587/588	T30R; R31P; A40L; A56S; E68A; A84Q; G120R; T139M; N145W; N213E;	+
589/590	T30R; R31P; A40L; A56S; A84Q; G120R; T139M; E142L; H148T; M170F; H222C;	+

33

TABLE 2D-continued

SEQ ID NO: (nt/aa)	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 8 (24 h, 65° C., 5M MDEA challenge/ 45° C., 1M MDEA pH 8.0 assay)
591/592	T30R; R31P; A40L; A56S; V70I; A84Q; H97F; G120R; A121L; T139M; M170F; E200R;	+
593/594	T30R; R31P; A40L; A56S; A84Q; D96E; G120R; T139M; E142L; M170F; R223C;	+
595/596	T30R; R31P; A40L; A56S; A84Q; D96A; H97F; G120R; T139M; H148T; M170F; H222C;	+
597/598	T30R; K37R; A40L; S42A; A56S; E68A; A84Q; G120R; T139M; S144L; N145W; A219T; A221C;	+
599/600	T30R; A40L; A56S; A84Q; G120R; T139M; H148T; M170F;	+
601/602	T30R; R31P; A40L; A56S; A84Q; H97F; G120R; A121L; T139M; M170F; E200R;	+
603/604	T30R; R31P; A40L; A56S; A60C; A84Q; A95V; G120R; T139M; N145W; A219T; A221C;	+
605/606	T30R; Q32R; A40L; Q43M; A56S; V70I; A84Q; G120R; A121L; T139M; H148T; H222C;	+
607/608	T30R; K37R; A40L; A56S; A60C; E68A; A84Q; A95V; Q119M; G120R; T139M; A219T;	++
609/610	T30R; R31P; A40L; S42A; A56S; A84Q; G120R; H124R; T139M; S144L;	++
611/612	T30R; R31P; A40L; S42A; A56S; E68A; A84Q; G120R; H124F; T139M; S144L; N145F; A221C;	++
613/614	T30R; K37R; A40L; A56S; A60C; A84Q; Q119M; G120R; T139M; S144L; A219T; A221C;	++
615/616	T30R; A40L; S42A; A56S; A60C; E68A; A84Q; Q119M; G120R; T139M; N145W;	++
617/618	G2T; T30R; R31P; Q32K; A40L; Q43M; A56S; A84Q; G120R; T139M; H222C;	++
619/620	T30R; K37R; A40L; A56S; E68A; A84Q; G120R; H124R; T139M; V157A;	++
621/622	T30R; A40L; A56S; A84Q; Q119M; G120R; T139M; S144L; V157A; N213E;	++
623/624	T30R; Q32K; A40L; A56S; A84Q; G120R; T139M; E142L; H148T; M170F; E200R; R223C;	++
625/626	T30R; R31P; Q32K; A36T; A40L; A56S; V70I; A84Q; D96E; G120R; T139M; E142L; H148T;	++
627/628	T30R; A40L; A56S; A60C; A84Q; A95V; Q119M; G120R; T139M; V157A; N213E; A219T;	++
629/630	T30R; A40L; A56S; E68A; A84Q; G120R; T139M; S144L; N145W; V157A; N213E;	++
631/632	T30R; A40L; A56S; A60C; A84Q; Q119M; G120R; H124R; T139M; S144L; N145W; N213E; A221C;	++
633/634	T30R; R31P; A40L; A56S; A84Q; A95V; Q119M; G120R; H124R; T139M; S144L; N145W; V157A; A221C;	++
635/636	T30R; K37R; A40L; S42A; A56S; A60C; E68A; A84Q; Q119M; G120R; T139M; S144L; N145F; V157A; A221C;	++
637/638	T30R; R31P; A40L; S42A; A56S; A84Q; A95V; Q119M; G120R; H124R; T139M; A221C;	++
639/640	T30R; R31P; K37R; A40L; A56S; A60C; E68A; A84Q; A95V; G120R; H124F; T139M; S144L; N145F; N213E; A219T;	++

34

TABLE 2D-continued

SEQ ID NO: (nt/aa)	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 8 (24 h, 65° C., 5M MDEA challenge/ 45° C., 1M MDEA pH 8.0 assay)
641/642	T30R; A40L; S42A; A56S; A84Q; A95V; Q119M; G120R; H124R; T139M; S144L; A221C;	++
643/644	G2T; T30R; R31P; Q32K; A40L; Q43M; A56S; A84Q; D96A; H97F; G120R; T139M; M170F; E200R; H222C;	++
645/646	T30R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; T139M; S144L; N145F;	++
647/648	T30R; A40L; Q43M; A56S; V70I; A84Q; G120R; A121L; T139M; M170F; R223C;	++
649/650	T30R; R31P; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; H124F; T139M; S144L; V157A; N213E; A221C;	++
651/652	G2T; T30R; R31P; A40L; Q43M; A56S; V70I; A84Q; G120R; T139M; E142L; H148T; M170F; E200R; H222C;	++
653/654	T30R; K37R; A40L; A56S; A60C; E68A; A84Q; A95V; Q119M; G120R; H124R; T139M; N213E; A219T; A221C;	++
655/656	T30R; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; T139M; N145W; N213E; A219T;	++
657/658	T30R; A40L; A56S; A84Q; G120R; H124R; T139M; S144L; N145F; V157A; A219T; A221C;	++
659/660	T30R; K37R; A40L; S42A; A56S; A84Q; A95V; Q119M; G120R; H124R; T139M; S144L; V157A; N213E; A221C;	++
661/662	T30R; R31P; A40L; A56S; A84Q; A95V; G120R; T139M; S144L; N145F; V157A; A221C;	+++
663/664	T30R; R31P; A40L; A56S; A84Q; A95V; G120R; H124R; T139M; S144L; N145F; A219T; A221C;	+++
665/666	T30R; R31P; K37R; A40L; A56S; A60C; A84Q; A95V; Q119M; G120R; T139M; S144L; V157A; A219T;	+++
667/668	T30R; R31P; K37R; A40L; S42A; A56S; A60C; E68A; A84Q; A95V; Q119M; G120R; H124R; T139M; S144L; N145W; N213E; A219T; A221C;	+++
669/670	T30R; A40L; A56S; A60C; E68A; A84Q; Q119M; G120R; H124R; T139M; S144L; N145F; V157A;	+++
671/672	T30R; R31P; A40L; A56S; A60C; A84Q; A95V; Q119M; G120R; H124R; T139M; V157A; A221C;	+++
673/674	T30R; R31P; K37R; A40L; A56S; E68A; A84Q; A95V; G120R; T139M; S144L; N145F; V157A; N213E; A221C;	+++
675/676	G2T; T30R; R31P; A40L; Q43M; A56S; A84Q; H97F; G120R; T139M; M170F; E200R;	+++
677/678	T30R; R31P; A40L; S42A; A56S; A60C; E68A; A84Q; A95V; Q119M; G120R; T139M; N145F; N213E;	+++

Levels of increased activity were determined relative to the reference polypeptide of SEQ ID NO: 332 (i.e., engineered polypeptide having T30R, A40L, A56S, A84Q, G120R, and T139M) using Assay 8 and defined as follows:
 "+" indicates 1.5-fold but less than 1.7-fold increased activity;
 "++" indicates at least 1.7-fold but less than 2.0-fold increased activity;
 "+++" indicates at least 2.0-fold increased activity.

TABLE 2E

SEQ ID NO: (nt/aa)	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 9 (24 h, 44° C., 5.6M NH ₃ (α = 0.3) challenge/ 25° C., 0.28M NH ₃ assay)	Assay 10 (24 h, RT, 5.6M NH ₃ (α = 0.3) challenge/ 25° C., 0.28M NH ₃ assay)
679/680	Q15R; T30R; H124G; K156L;	+	
681/682	Q15R; T30R; H97F; V131F; V157A;	+	
683/684	T4F; Q15R; T30R; Q32K; S35R;	+	

TABLE 2E-continued

SEQ ID	NO: Amino Acid Residue Difference(s) (nt/aa) (relative to SEQ ID NO: 2)	Assay 9 (24 h, 44° C., 5.6M NH ₃ ($\alpha = 0.3$) challenge/ 25° C., 0.28M NH ₃ assay)	Assay 10 (24 h, RT, 5.6M NH ₃ ($\alpha = 0.3$) challenge/ 25° C., 0.28M NH ₃ assay)
685/686	Q15R; T30R; V131F; K156L; V157A; A219T;	+	
687/688	Q15R; T30R; Q32K; S35A; A221C;	+	
689/690	Q15R; T30R; K143R; H148T; N213E;	+	
691/692	Q15R; T30R; L34H; A56S; A60C; A221C; R223Q;	+	
693/694	Q15R; T30R; L34H; Y93W;	+	
695/696	Q15R; T30R; D96E; H124G; K156L; A219T;	+	
697/698	Q15R; T30R; V70I; Y93W;	+	
699/700	Q15R; T30R; V131F; V157A; A219T;	+	
701/702	Q15R; T30R; V138W;	+	
703/704	Q15R; T30R; V157A; E200R; A219T;	+	
705/706	Q15R; A22G; T30R; L34H; V70I; Y93W; A95V;	+	
707/708	T4F; Q15R; T30R; L34H; A221C;	+	
709/710	T4F; Q15R; T30R; Q32R; L34H; S35R; A56S; A60C; R223C;	+	
711/712	G2R; Q15R; T30R; N213Q; A219T;	+	
713/714	Q15R; T30R; A95V;	+	
715/716	Q15R; T30R; H124G; H148T; N213Q; A219T;	+	
717/718	Q15R; T30R; D96E; H124G; H148T; K156L; A219T;	+	
719/720	G2R; Q15R; T30R; D96E;	+	
721/722	Q15R; A22G; T30R; A40Q; V70I; Y93W; A95V;	+	
723/724	Q15R; T30R; Q32R; L34H; S35A;	+	
725/726	Q15R; T30R; S35A; A60C; A221C; R223C;	+	
727/728	Q15R; T30R; L34H; S35A; A60V; A221C;	+	
729/730	Q15R; T30R; S35R; K156L; V157A; E200R; A219T;	+	
731/732	Q15R; T30R; K37R; A40L; Y93W; A95V;	+	
733/734	G2R; Q15R; T30R; H124G; K156L; A219T;	+	
735/736	G2R; Q15R; T30R; N213Q;	+	
737/738	Q15R; T30R; Q32K; L34H; S35A; A221C; R223C;	+	
739/740	Q15R; T30K; R31P; S35R; H97F; E159V;	+	
741/742	Q15R; T30R; D96E; T139H; N145C; H148T; K156L; A219T;	+	++
743/744	Q15R; T30R; V70I; Y93W; A121Q;	+	
745/746	G2R; Q15R; T30R; T139H; N145C; N213Q; A219T;	+	+
747/748	T4F; Q15R; T30R; A56S; A221C;	+	
749/750	Q15R; T30R; A56S; A60V; A221C;	++	
751/752	Q15R; A22G; T30R; H222C;	++	
753/754	Q15R; T30R; L34H; A60V; A84R; A221C;	++	
755/756	T4M; Q15R; T30R; A60V;	++	
757/758	Q15R; T30R; T139Q; K143R; K156L; N213Q;	++	
759/760	Q15R; T30R; V131L; K156L; E200R;	++	
761/762	T4F; Q15R; T30R; Q32K; L34H; A56S; A60V; A221C;	++	
763/764	Q15R; T30R; D96E; H124G; T139Q; K143R; K156L;	++	
765/766	T4F; Q15R; T30R; A60V; A221C;	++	
767/768	G2R; Q15R; T30R; H124G; K156L;	++	
769/770	T4M; Q15R; T30R; Q32K; A56S; A60C; A221C; R223C;	++	
771/772	T4M; Q15R; T30R; L34H; S35R; A60V; A221C;	++	
773/774	T4M; Q15R; T30R; S35A; A60C; A221C;	++	
775/776	G2R; Q15R; T30R; E142L; N213Q; A219T;	++	
777/778	T4F; Q15R; T30R; L34H; A56S; A84N; A221C;	++	
779/780	G2R; Q15R; T30R; D96E; N145C; N213E;	++	+
781/782	Q15R; T30R; T47R; H97F; V131L; K156L; E159V; E200R; A219T;	++	
783/784	Q15R; T30R; T47R; H97F; K156L; V157A;	++	
785/786	Q15R; T30R; H97F; K156L; V157A; E200R; A219T;	++	
787/788	Q15R; T30R; Q32K; A56S; A60V; A84Q; A221C;	++	
789/790	Q15R; T30R; S42A; K156L; V157A; A219T;	++	
791/792	Q15R; T30R; Q32R; S35R; A56S;	++	
793/794	G2R; Q15R; T30R; T47R; K156L; V157A; E200R; A219T;	++	
795/796	Q15R; T30R; Q32R; A56S;	++	
797/798	Q15R; T30R; R31P; V131L; V157A; A219T;	++	
799/800	Q15R; T30R; L34H; S35A; A56S; A221C;	++	
801/802	T4M; Q15R; T30R; Q32R; L34H; R223C;	++	
803/804	Q15R; T30R; R31P; S42A; K156L;	++	
805/806	G2R; Q15R; T30R; T139Q; N145C; K156L; A219T;	++	++
807/808	Q15R; T30R; Y93W; A95V; A121Q; H222C;	++	
809/810	Q15R; T30R; K37R; A40Q;	++	
811/812	Q15R; T30R; Q32K; S35A; A56S; A84N; A221C;	++	+
813/814	Q15R; A22G; T30R; L34H; A40L;	++	
815/816	Q15R; T30R; H97F; V131L; K156L; V157A;	++	
817/818	Q15R; T30R; D96E; H124G; T139Q; N145C; H148C; N213E;	++	
819/820	Q15R; T30R; A40W; E68G; H222C;	++	
821/822	Q15R; T30R; D96E; T139H; K143R; N145C; H148T; N213Q;	++	++
823/824	Q15R; T30R; Q32K; L34H; A56S; A84Q;	++	

TABLE 2E-continued

SEQ ID	NO: Amino Acid Residue Difference(s) (nt/aa) (relative to SEQ ID NO: 2)	Assay 9 (24 h, 44° C., 5.6M NH ₃ (α = 0.3) challenge/ 25° C., 0.28M NH ₃ assay)	Assay 10 (24 h, RT, 5.6M NH ₃ (α = 0.3) challenge/ 25° C., 0.28M NH ₃ assay)
825/826	Q15R; A22G; T30R; L34H; A40W; V70I; A121T;	++	
827/828	G2R; Q15R; T30R; D96E; E142L; N145C; N213Q; A219T;	++	+
829/830	Q15R; T30K; R31P; S42A; H97F; V131L;	++	
831/832	Q15R; T30R; A56S; A60C;	++	
833/834	Q15R; T30R; K37R; A40L;	++	
835/836	T4F; Q15R; T30R; A56S; A84N; A221C; R223C;	++	
837/838	Q15R; T30R; L34H;	++	
839/840	Q15R; A22G; T30R; A40L; H222C;	++	
841/842	G2R; Q15R; T30R; H124G; K156L; N213E; A219T;	++	
843/844	G2R; Q15R; T30R; D96E; K156L; N213E;	++	
845/846	Q15R; T30R; V70I; Y93W; A95V; H222C;	++	
847/848	G2R; Q15R; T30R; R31P; S42A; V131L; V157A; A219T;	++	
849/850	Q15R; T30R; V131L; K156L; V157A;	++	
851/852	G2R; Q15R; T30R; D96E; H124G; T139H; E142L; N213E; A219T;	++	
853/854	Q15R; T30R; A40Q; V138W; H222C;	++	
855/856	Q15R; T30R; L34H; E68A; V70I;	++	
857/858	Q15R; T30R; V70I;	++	
859/860	Q15R; T30R; L34H; A40L;	++	
861/862	Q15R; T30R; A56S; A221C;	++	
863/864	Q15R; T30R; Q32R; S35R; A84Q; R223Q;	++	
865/866	G2R; Q15R; T30R; D96E; H148C; K156L; N213Q;	++	+
867/868	Q15R; T30R; S35A; A84N; A221C;	++	
869/870	T4F; Q15R; T30R; Q32K; L34H; S35A; A56S; A60V; A84K; A221C; R223C;	++	
871/872	Q15R; T30R; V131L; V157A; E200R;	++	
873/874	Q15R; T30A; R31P; S35A; T47R; H97F; V131L; K156L; V157A;	+++	
875/876	Q15R; T30R; A84R; A221C; R223C;	+++	
877/878	Q15R; T30A; R31P; S35R; H97F; V131L; V157A;	+++	
879/880	Q15R; T30R; H222C;	+++	
881/882	Q15R; T30R; R31P; S35A; V131L; K156L; V157A; A219T;	+++	
883/884	Q15R; T30R; L34H; K37C;	+++	
885/886	Q15R; T30R; L34H; S35R; A84R; A221C;	+++	
887/888	Q15R; T30R; L34H; K37R;	+++	
889/890	T4M; Q15R; T30R; Q32K; S35R; A56S; R223Q;	+++	
891/892	Q15R; T30R; Q32K; A56S; A60V; A84K; A221C; R223C;	+++	
893/894	G2R; Q15R; T30R; T139Q; K143R; N145C; K156L; N213Q;	+++	++
895/896	Q15R; T30R; A56S;	+++	
897/898	Q15R; T30R; D96E; H124G; T139H; K143R; N145C; N213E;	+++	++
899/900	Q15R; T30R; K37R; A40Q; E68V; H222C;	+++	
901/902	Q15R; A22G; T30R; E68A; H222C;	+++	
903/904	Q15R; L17X; T30R; A84N; R223C;	+++	
905/906	T4M; Q15R; T30R; S35R; A56S; A84Q;	+++	
907/908	Q15R; T30R; R31P; S35R; H97F; V131L; K156L; V157A; E159V; E200R;	+++	
909/910	Q15R; A22G; T30R; A40W; E68A; H222C;	+++	
911/912	Q15R; T30R; S35R; A56S; A60V; A84K; R223C;	+++	
913/914	Q15R; T30R; Q32K; A56S; A84Q; A221C;	+++	
915/916	T4M; Q15R; T30R; A56S;	+++	
917/918	Q15R; T30R; Q32R; L34H; A56S; A84N; A221C;	+++	
919/920	Q15R; T30R; K37R; A40L; V70I; A95V;	+++	
921/922	Q15R; T30R; V138F; H222C;	+++	
923/924	Q15R; T30R; A121K; H222C;	+++	
925/926	G2R; Q15R; T30R; H124G; K143R; N145C; H148T; N213Q;	+++	++
927/928	T4F; Q15R; T30R; Q32R; A56S; R223C;	+++	
929/930	Q15R; T30R; L34H; V138W; H222C;	+++	
931/932	Q15R; T30R; A56S; A84K; A221C;	+++	
933/934	Q15R; T30R; A56S; A84Q; A221C; R223C;	+++	
935/936	Q15R; T30R; A56S; A84R; A221C;	+++	
937/938	Q15R; T30R; L34H; A84Q; R223C;	+++	
939/940	Q15R; T30R; R31P; S35R; V131L; V157A;	+++	
941/942	Q15R; T30R; V138W; H222C;	+++	
943/944	Q15R; T30R; V70I; H222C;	+++	
945/946	T4M; Q15R; T30R; Q32R; L34H; A56S; A84K; A221C; R223C;	+++	
947/948	T4M; Q15R; T30R; L34H; A56S; R223C;	+++	
949/950	Q15R; T30R; L34H; H222C;	+++	
951/952	T4M; Q15R; T30R; Q32K; A84N; R223C;	+++	
953/954	T4M; Q15R; T30R; A84N; R223C;	+++	
955/956	G2R; Q15R; T30R; H124G; T139H; N145C; H148T; K156L;	+++	++
957/958	Q15R; T30R; S35R; A56S; A84R; R223C;	+++	
959/960	Q15R; T30R; L34H; A84R; R223C;	+++	
961/962	Q15R; T30R; A56S; A84R; A221C; R223C;	+++	

TABLE 2E-continued

SEQ ID	NO: Amino Acid Residue Difference(s) (nt/aa) (relative to SEQ ID NO: 2)	Assay 9 (24 h, 44° C., 5.6M NH ₃ ($\alpha = 0.3$) challenge/ 25° C., 0.28M NH ₃ assay)	Assay 10 (24 h, RT, 5.6M NH ₃ ($\alpha = 0.3$) challenge/ 25° C., 0.28M NH ₃ assay)
963/964	T4M; Q15R; T30R; Q32R; A56S; A84R; A221C; R223C;	+++	
965/966	Q15R; T30R; D96E; H124G; T139Q; N145C; K156L; N213Q;	+++	+++
967/968	G2R; Q15R; T30R; N145C;		+
969/970	Q15R; T30R; N145C; H148T; K156L;		++
971/972	Q15R; T30R; D96E; H124G; E142L; N145C;		++

Levels of increased activity were determined relative to the reference polypeptide of SEQ ID NO: 32 (i.e., engineered polypeptide having Q15R and T30R) and defined as follows:

Assay 9:

“+” indicates at least 1.5-fold but less than 1.8-fold increased activity;

“++” indicates at least 1.8-fold but less than 2.3-fold increased activity;

“+++” indicates at least 2.3-fold increased activity.

Assay 10:

“+” indicates at least 1.3-fold but less than 1.6-fold increased activity;

“++” indicates at least 1.6-fold but less than 2-fold increased activity;

“+++” indicates at least 2-fold increased activity.

TABLE 2F

SEQ ID	NO: Amino Acid Residue Difference(s) (nt/aa) (relative to SEQ ID NO: 2)	Assay 11 (24 h, 58° C., 8.4M NH ₃ ($\alpha = 0.3$) challenge/ 25° C., 1.37M NH ₃ assay)
973/974	Q15R; T30R; Q32K; S35A; A56S; V70I; A84N; V131L; T139H; V157A; A221C;	+
975/976	Q15R; T30R; R31P; Q32K; S35A; T47R; A56S; A84N; H97F; K156L; A221C;	+
977/978	T4M; Q15R; T30R; Q32K; S35A; A56S; A84K; V131L; T139Q; N145C; N213Q;	+
979/980	G2R; Q15R; T30R; R31P; Q32K; S35A; A56S; A84N; D96E; H97F; H148T;	+
981/982	G2R; Q15R; T30R; Q32K; S35A; A56S; A84N; D96E; H97F; V138F; K156L;	+
983/984	G2R; Q15R; T30R; R31P; Q32K; S35A; T47R; A56S; A84N; H97F; H148T;	+
985/986	Q15R; T30R; Q32K; L34H; S35A; A56S; A84N; A95V; H124G; V131L; T139Q;	+
987/988	T4M; Q15R; T30R; Q32K; S35A; A56S; A84N; A95V; V131L; N145C; A221C;	+
989/990	Q15R; T30R; Q32K; S35A; A56S; A84N; A95V; H124G; V131L; N145C;	+
991/992	G2R; Q15R; T30R; Q32K; S35A; A56S; A84N; D96E; H97F; A121Q; H148T;	+
993/994	T4M; Q15R; T30R; Q32K; S35A; A56S; V70I; A84K; A95V; H124G; V131L;	+
995/996	Q15R; T30R; R31P; Q32K; S35A; K37R; A56S; A84N; D96E; H97F; V138F;	+
997/998	T4M; Q15R; T30R; Q32K; S35A; A56S; A84N; H124G; V131L; N145C; A221C;	+
999/1000	T4M; Q15R; T30R; Q32K; L34H; S35A; A56S; A84N; A95V; H124G; V131L;	+
1001/1002	G2R; Q15R; T30R; R31P; Q32K; S35A; K37C; A40L; A56S; A84N; D96E;	+
1003/1004	Q15R; T30R; R31P; Q32K; S35A; A56S; A84N; D96E; H97F; K156L; A221C;	+
1005/1006	Q15R; T30R; Q32K; S35A; A56S; A84N; D96E; A121Q; V138F; H148T; K156L;	+
1007/1008	Q15R; T30R; Q32K; L34H; S35A; A56S; V70I; A84R; V131L; T139Q; N145C;	+
1009/1010	G2R; Q15R; T30R; Q32K; S35A; A56S; A84N; D96E; H97F; A121K; V138W;	+
1011/1012	Q15R; T30R; Q32K; L34H; S35A; A56S; A84R; V131L; A221C;	+
1013/1014	Q15R; T30R; Q32K; L34H; S35A; A56S; A84Q; H124G; V131L; T139Q; N145C;	+
1015/1016	T4M; Q15R; T30R; Q32K; L34H; S35A; A56S; V70I; A84Q; A95V; H124G;	+
1017/1018	Q15R; T30R; Q32K; S35A; A56S; A84K; A95V; V131L; N145C; V157A;	+

TABLE 2F-continued

SEQ ID NO: (nt/aa)	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 11 (24 h, 58° C., 8.4M NH ₃ (α = 0.3) challenge/ 25° C., 1.37M NH ₃ assay)
1019/1020	Q15R; T30R; Q32K; L34H; S35A; A56S; A84Q; H124G; V131L; T139H; N213E; A221C;	+
1021/1022	Q15R; T30R; Q32K; S35A; A56S; A84K; A95V; V131L; A221C;	+
1023/1024	Q15R; T30R; Q32K; S35A; A56S; A84Q; A95V; V131L; N213E; A221C;	+
1025/1026	T4M; Q15R; T30R; Q32K; L34H; S35A; A56S; V70I; A84Q; A95V; H124G; V131L; T139Q; N145C; A221C;	+
1027/1028	T4M; Q15R; T30R; Q32K; S35A; A56S; A84Q; A95V; V131L; V157A; A221C; H222C;	+
1029/1030	Q15R; T30R; Q32K; S35A; A56S; A84K; A95V; H124G; V131L; T139Q; N213Q; A221C; H222C; R223C;	+
1031/1032	Q15R; T30R; Q32K; S35A; A56S; A84Q; A95V; H124G; V131L; A221C;	+
1033/1034	T4M; Q15R; T30R; Q32K; S35A; A56S; V70I; A84Q; A95V; V131L; T139Q; A221C;	+
1035/1036	Q15R; T30R; Q32K; S35A; A56S; A84Q; H124G; V131L; A221C;	+
1037/1038	T4M; Q15R; T30R; Q32K; L34H; S35A; A56S; V70I; A84R; A95V; H124G; V131L; V157A; N213E; A221C; H222C; R223C;	+
1039/1040	Q15R; T30R; Q32K; S35A; A56S; V70I; A84Q; V131L; T139H; N213Q; A221C; R223C;	+
1041/1042	T4M; Q15R; T30R; Q32K; S35A; A56S; V70I; A84K; V131L; T139Q; A221C; H222C; R223C;	+
1043/1044	T4M; Q15R; T30R; Q32K; S35A; A56S; A84Q; A95V; H124G; V131L; N213E; A221C;	+
1045/1046	T4M; Q15R; T30R; Q32K; L34H; S35A; A56S; A84Q; H124G; V131L; N145C; V157A; N213E; A221C;	+
1047/1048	Q15R; T30R; Q32K; S35A; A56S; A84Q; A95V; V131L; T139H; N145C; V157A; A221C; H222C; R223C;	+
1049/1050	T4M; Q15R; T30R; Q32K; S35A; A56S; A84Q; A95V; H124G; V131L; V157A; A221C;	+
1051/1052	T4M; Q15R; T30R; Q32K; L34H; S35A; A56S; A84K; A95V; H124G; V131L; N145C; A221C; H222C; R223C;	++
1053/1054	T4M; Q15R; T30R; Q32K; S35A; A56S; A84K; A95V; H124G; V131L; V157A; N213Q; A221C; H222C; R223C;	++
1055/1056	T4M; Q15R; T30R; Q32K; S35A; A56S; V70I; A84Q; A95V; V131L; T139Q; V157A; A221C;	++

Levels of increased activity were determined relative to the reference polypeptide of SEQ ID NO: 812 (i.e., engineered polypeptide having Q15R, T30R, Q32K, S35A, A56S, A84N, and A221C) using Assay 11 and defined as follows:

“+” indicates 1.3-fold but less than 1.7-fold increased activity;

“++” indicates at least 1.7-fold but less than 2.0-fold increased activity;

“+++” indicates at least 2.0-fold increased activity.

TABLE 2G

SEQ ID NO: (nt/aa)	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 12 (24 h/70° C./ 5M MDEA challenge/ 45° C. assay/ 0.5M MDEA)
1057/1058	T30R; R31P; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; T139M; N145W; N213E; A219T;	+
1059/1060	T30R; R31P; Q32R; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; H124R; T139M; N145F; E200R; N213E; A219T;	+
1061/1062	T30R; R31P; Q32R; K37R; A40L; A56S; A60C; E68A; A84Q; A95V; H97F; Q119M; G120R; T139M; N145W; N213E; A219T; A221C; H222C;	+
1063/1064	T30R; Q32R; K37R; A40L; Q43M; A56S; E68A; A84Q; A95V; H97F; Q119M; G120R; H124R; T139M; N145F; H148T; V157A; M170F; E200R; N213E; A219T;	+
1065/1066	T30R; R31P; K37R; A40L; Q43M; A56S; A60C; E68A; A84Q; A95V; Q119M; G120R; T139M; N145W; N213E; A219T; A221C; H222C;	+
1067/1068	T30R; Q32R; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; H124R; T139M; N145W; E200R; N213E; A219T;	+
1069/1070	T30R; K37R; A40L; S42A; Q43M; A56S; E68A; A84Q; A95V; Q119M; G120R; T139M; N145W; H148T; V157A; E200R; N213E; A219T;	+
1071/1072	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; A84Q; A95V; D96A; H97F; Q119M; G120R; T139M; S144L; N145F; N213E; A219T;	+

TABLE 2G-continued

SEQ ID NO: (nt/aa)	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 12 (24 h/70° C./ 5M MDEA challenge/ 45° C. assay/ 0.5M MDEA)
1073/1074	T30R; R31P; Q32R; K37R; A40L; A56S; E68A; A84Q; A95V; H97F; Q119M; G120R; T139M; N145W; N213E; A219T;	+
1075/1076	T30R; Q32R; K37R; A40L; A56S; E68A; V70I; A84Q; A95V; D96E; Q119M; G120R; H124R; T139M; N145W; V157A; N213E; A219T;	+
1077/1078	T30R; K37R; A40L; A56S; A60C; E68A; A84Q; A95V; D96A; H97F; Q119M; G120R; H124F; T139M; N145W; H148T; M170F; N213E; A219T;	+
1079/1080	T30R; R31P; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; H124R; T139M; S144L; N145F; H148T; E200R; N213E; A219T;	+
1081/1082	T30R; R31P; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; H124R; T139M; S144L; N145W; H148T; V157A; M170F; E200R; N213E; A219T;	+
1083/1084	T30R; R31P; Q32R; K37R; A40L; A56S; E68A; V70I; A84Q; A95V; D96A; H97F; Q119M; G120R; T139M; N145W; N213E; A219T;	+
1085/1086	T30R; R31P; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; T139M; N145W; H148T; M170F; N213E; A219T;	+
1087/1088	T30R; Q32R; K37R; A40L; A56S; E68A; A84Q; A95V; D96A; H97F; Q119M; G120R; H124F; T139M; N145W; M170F; N213E; A219T; H222C;	+
1089/1090	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; A84Q; A95V; Q119M; G120R; H124R; T139M; S144L; N145W; H148T; M170F; N213E; A219T;	+
1091/1092	T30R; Q32K; K37R; A40L; A56S; E68A; A84Q; A95V; D96E; Q119M; G120R; T139M; S144L; N145W; H148T; M170F; N213E; A219T;	+
1093/1094	T30R; K37R; A40L; A56S; E68A; A84Q; A95V; H97F; Q119M; G120R; T139M; N145W; M170F; N213E; A219T; H222C;	+
1095/1096	T30R; R31P; K37R; A40L; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; T139M; N145W; H148T; M170F; N213E; A219T; A221C;	++
1097/1098	T30R; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; H124R; T139M; N145F; M170F; E200R; N213E; A219T;	++
1099/1100	T30R; R31P; Q32R; K37R; A40L; S42A; Q43M; A56S; E68A; A84Q; A95V; Q119M; G120R; T139M; N145W; N213E; A219T;	++
1101/1102	T30R; K37R; A40L; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; N145F; V157A; N213E; A219T;	++
1103/1104	T30R; K37R; A40L; A56S; E68A; A84Q; A95V; D96A; H97F; Q119M; G120R; T139M; N145F; H148T; M170F; N213E; A219T;	++
1105/1106	T30R; K37R; A40L; S42A; A56S; E68A; A84Q; A95V; H97F; Q119M; G120R; H124R; T139M; S144L; N145W; H148T; N213E; A219T;	++
1107/1108	T30R; K37R; A40L; S42A; A56S; E68A; A84Q; A95V; Q119M; G120R; H124R; T139M; S144L; N145W; V157A; N213E; A219T;	++
1109/1110	T30R; R31P; K37R; A40L; A56S; E68A; V70I; A84Q; A95V; D96A; H97F; Q119M; G120R; T139M; N145W; N213E; A219T; A221C; H222C;	++
1111/1112	T30R; R31P; K37R; A40L; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; T139M; N145F; V157A; N213E; A219T;	++
1113/1114	T30R; R31P; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; T139M; N145F; H148T; V157A; M170F; N213E; A219T;	++
1115/1116	T30R; K37R; A40L; A56S; E68A; V70I; A84Q; A95V; H97F; Q119M; G120R; T139M; S144L; N145F; H148T; V157A; N213E; A219T;	++
1117/1118	T30R; R31P; K37R; A40L; S42A; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; N145F; H148T; V157A; M170F; N213E; A219T;	++
1119/1120	T30R; R31P; K37R; A40L; Q43M; A56S; A60C; E68A; A84Q; A95V; Q119M; G120R; T139M; N145W; N213E; A219T; H222C;	++
1121/1122	T30R; R31P; Q32K; K37R; A40L; A56S; E68A; A84Q; A95V; D96E; H97F; Q119M; G120R; T139M; N145W; E200R; N213E; A219T;	++
1123/1124	T30R; R31P; Q32K; K37R; A40L; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; S144L; N145W; V157A; N213E; A219T;	++
1125/1126	T30R; R31P; K37R; A40L; A56S; E68A; A84Q; A95V; D96A; H97F; Q119M; G120R; T139M; N145W; H148T; M170F; N213E; A219T;	++
1127/1128	T30R; K37R; A40L; A56S; E68A; V70I; A84Q; A95V; H97F; Q119M; G120R; H124R; T139M; N145F; H148T; V157A; M170F; E200R; N213E; A219T;	++
1129/1130	T30R; R31P; K37R; A40L; A56S; E68A; A84Q; A95V; D96E; Q119M; G120R; H124R; T139M; N145W; V157A; M170F; N213E; A219T;	++
1131/1132	T30R; R31P; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; T139M; S144L; N145F; V157A; N213E; A219T;	++
1133/1134	T30R; R31P; K37R; A40L; A56S; E68A; V70I; A84Q; A95V; H97F; Q119M; G120R; T139M; N145W; H148T; M170F; N213E; A219T; H222C;	++
1135/1136	T30R; K37R; A40L; Q43M; A56S; E68A; A84Q; A95V; H97F; Q119M; G120R; T139M; S144L; N145F; V157A; N213E; A219T;	++
1137/1138	T30R; K37R; A40L; S42A; A56S; E68A; A84Q; A95V; Q119M; G120R; T139M; S144L; N145W; H148T; V157A; M170F; E200R; N213E; A219T;	++
1139/1140	T30R; R31P; Q32R; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; H124R; T139M; S144L; N145F; H148T; V157A; M170F; N213E; A219T;	++
1141/1142	T30R; R31P; Q32K; K37R; A40L; A56S; E68A; A84Q; A95V; D96E; Q119M; G120R; H124R; T139M; N145F; V157A; M170F; N213E; A219T;	++

TABLE 2G-continued

SEQ ID NO: (nt/aa)	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 12 (24 h/70° C./ 5M MDEA challenge/ 45° C. assay/ 0.5M MDEA)
1143/1144	T30R; K37R; A40L; S42A; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; T139M; S144L; N145F; M170F; E200R; N213E; A219T;	++
1145/1146	T30R; R31P; K37R; A40L; A56S; E68A; A84Q; A95V; H97F; Q119M; G120R; H124R; T139M; N145W; V157A; N213E; A219T;	++
1147/1148	T30R; R31P; Q32R; K37R; A40L; S42A; Q43M; A56S; E68A; A84Q; A95V; Q119M; G120R; T139M; S144L; N145F; N213E; A219T;	++
1149/1150	T30R; R31P; Q32K; K37R; A40L; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; T139M; N145F; V157A; M170F; N213E; A219T;	+++
1151/1152	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; N145F; H148T; V157A; M170F; N213E; A219T;	+++
1153/1154	T30R; K37R; A40L; A56S; E68A; A84Q; A95V; D96E; Q119M; G120R; H124R; T139M; S144L; N145F; N213E; A219T;	+++
1155/1156	T30R; R31P; K37R; A40L; S42A; A56S; E68A; A84Q; A95V; D96A; H97F; Q119M; G120R; T139M; N145W; V157A; N213E; A219T;	+++
1157/1158	T30R; R31P; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; H124R; T139M; N145W; V157A; M170F; N213E; A219T;	+++

Levels of increased activity were determined relative to the reference polypeptide of SEQ ID NO: 656 (i.e., engineered polypeptide having T30R; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; T139M; N145W; N213E; A219T) using Assay 12 and defined as follows:

“+” indicates at least 1.5-fold but less than 1.7-fold increased activity;

“++” indicates at least 1.7-fold but less than 2.0-fold increased activity;

“+++” indicates at least 2.0-fold increased activity.

TABLE 2H

SEQ ID NO: (nt/aa)	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 13 (24 h/70° C./ 8.4M ammonia challenge/ RT assay)
1159/1160	T4M; Q15G; T30R; Q32K; S35A; A56S; V70I; A84Q; A95V; D96E; H97F; V131L; T139Q; K156L; V157A; A221C;	+
1161/1162	T4M; Q15G; T30R; Q32K; S35A; A56S; V70I; A84Q; A95V; H97F; A121Q; V131L; T139Q; H148T; V157A; N213E; A221C;	+
1163/1164	T4M; Q15R; T30R; Q32K; S35A; A56S; V70I; A84Q; A95V; V131L; T139Q; H148T; V157A; A221C; R223C;	+
1165/1166	G2R; T4M; Q15R; T30R; Q32K; S35A; A56S; V70I; A84H; A95V; D96E; H97F; A121K; V131L; T139Q; V157A; N213E; A221C;	+
1167/1168	T4M; Q15G; T30R; Q32K; S35A; A56S; V70I; A84Q; A95V; H97F; V131L; T139Q; V157A; A221C;	+
1169/1170	G2R; T4M; Q15R; T30R; Q32K; S35A; A56S; V70I; A84Q; A95V; A121K; V131L; T139Q; V157A; N213E; A221C;	+
1171/1172	T4M; Q15R; T30R; Q32K; S35A; A56S; V70I; A84Q; A95V; V131L; T139Q; V157A; N213Q; A221C;	+

Levels of increased activity were determined relative to the reference polypeptide of SEQ ID NO: 1056 (i.e., engineered polypeptide having T4M; Q15R; T30R; Q32K; S35A; A56S; V70I; A84Q; A95V; V131L; T139Q; V157A; A221C) using Assay 13 and defined as follows:

“+” indicates at least 1.3-fold;

TABLE 2I

SEQ ID NO: (nt/aa)	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 14 (24 h/82.5° C./ 4.2M MDEA challenge/ 45° C. assay/ 960 mM MDEA)	Assay 15 (24 h/85° C./ 4.2M MDEA challenge/ 45° C. assay/ 960 mM MDEA)	Assay 16 (24 h/90° C./ 4.2M MDEA challenge/ 45° C. assay/ 960 mM MDEA)
1173/1174	T30R; R31P; A36L; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; N145F; H148T; V157A; M170F; N213E; A219T;	+		
1175/1176	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; M129F; T139M; N145F; H148T; V157A; M170F; N213E; A219T;	+		
1177/1178	T30R; R31P; K37R; A40L; Q43M; A56S; F66Y; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; N145F; H148T; V157A; M170F; N213E; A219T;	+		

TABLE 2I-continued

SEQ ID	Amino Acid Residue Difference(s) (nt/aa) (relative to SEQ ID NO: 2)	Assay 14 (24 h/82.5° C./ 4.2M MDEA challenge/ 45° C. assay/ 960 mM MDEA)	Assay 15 (24 h/85° C./ 4.2M MDEA challenge/ 45° C. assay/ 960 mM MDEA)	Assay 16 (24 h/90° C./ 4.2M MDEA challenge/ 45° C. assay/ 960 mM MDEA)
1179/1180	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; V123K; H124R; T139M; N145F; H148T; V157A; M170F; N213E; A219T;	+		
1181/1182	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; N145F; H148K; V157A; M170F; N213E; A219T;	+		
1183/1184	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; M129Y; T139M; S144R; N145F; H148I; V157A; M170F; N213E; A219T;	+++	+++	+++
1185/1186	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T;	++	+++	+
1187/1188	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119L; G120R; H124R; M129Y; T139M; N145C; H148T; V157A; M170F; N213E; A219T;	++	+++	++
1189/1190	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; M129F; T139M; N145C; H148T; V157A; M170F; N213E; A219T;	+++	+++	+
1191/1192	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; M129Y; T139M; N145C; H148T; V157A; M170F; N213E; A219T;	++	+++	
1193/1194	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119M; G120R; H124R; T139M; S144R; N145F; H148T; V157A; M170F; N213E; A219T;	++	+++	
1195/1196	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; M129Y; T139M; S144R; N145F; H148K; V157A; M170F; N213E; A219T;	++	++	+
1197/1198	T30R; R31P; A36L; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; N145C; H148K; V157A; M170F; N213E; A219T;	++	++	+
1199/1200	T30R; R31P; A36L; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119M; G120R; H124R; T139M; S144R; N145F; H148T; V157A; M170F; N213E; A219T;	++	++	
1201/1202	T30R; R31P; A36L; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; M129Y; T139M; N145C; H148I; V157A; M170F; N213E; A219T;	+	++	+
1203/1204	T30R; R31P; A36L; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; V123K; H124R; T139M; N145C; H148T; V157A; M170F; N213E; A219T;		++	++
1205/1206	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119M; G120R; H124R; M129Y; T139M; N145F; H148T; V157A; M170F; N213E; A219T;	++	++	+
1207/1208	T30R; R31P; K37R; A40L; Q43M; H44L; A56S; E68A; V70I; A84Q; A95V; Q119L; G120R; H124R; M129Y; T139M; S144R; N145F; H148T; V157A; M170F; N213E; A219T;	++	++	++
1209/1210	T30R; R31P; K37R; A40L; Q43M; H44L; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; M129F; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T;	++	++	+++
1211/1212	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119M; G120R; H124R; M129Y; T139M; N145C; H148T; V157A; M170F; N213E; A219T;		++	+
1213/1214	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; N145C; H148T; V157A; M170F; N213E; A219T;	++	++	
1215/1216	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; M129Y; T139M; N145F; H148T; V157A; M170F; N213E; A219T;	++	++	

TABLE 2I-continued

SEQ ID	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 14 (24 h/82.5° C./ 4.2M MDEA challenge/ 45° C. assay/ 960 mM MDEA)	Assay 15 (24 h/85° C./ 4.2M MDEA challenge/ 45° C. assay/ 960 mM MDEA)	Assay 16 (24 h/90° C./ 4.2M MDEA challenge/ 45° C. assay/ 960 mM MDEA)
1217/1218	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119L; G120R; H124R; T139M; S144R; N145C; H148I; V157A; M170F; N213E; A219T;	+	++	+
1219/1220	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; M129Y; T139M; S144R; N145F; H148T; V157A; M170F; N213E; A219T;	+	++	
1221/1222	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119L; G120R; H124R; T139M; S144R; N145F; H148I; V157A; M170F; N213E; A219T;	+	++	
1223/1224	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119M; G120R; H124R; T139M; N145C; H148T; V157A; M170F; N213E; A219T;	+	++	+
1225/1226	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119L; G120R; H124R; M129F; T139M; N145C; H148T; V157A; M170F; N213E; A219T;	++	++	+
1227/1228	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119M; G120R; H124R; M129Y; T139M; N145F; H148K; V157A; M170F; N213E; A219T;		++	
1229/1230	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119M; G120R; H124R; T139M; N145F; H148T; V157A; M170F; N213E; A219T;	+++	++	
1231/1232	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119L; G120R; H124R; M129F; T139M; N145C; H148T; V157A; M170F; N213E; A219T;		++	+
1233/1234	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119M; G120R; H124R; M129F; T139M; N145F; H148T; V157A; M170F; N213E; A219T;	++	++	
1235/1236	T30R; R31P; K37R; A40L; Q43M; H44L; A56S; E68A; V70I; A84Q; D86A; A95V; Q119M; G120R; H124R; M129Y; T139M; S144R; N145F; H148T; V157A; M170F; N213E; A219T;	++	++	+
1237/1238	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119L; G120R; H124R; T139M; S144R; N145F; H148I; V157A; M170F; N213E; A219T;	+	++	
1239/1240	T30R; R31P; A36L; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; S144R; N145F; H148T; V157A; M170F; N213E; A219T;	+	++	
1241/1242	T30R; R31P; A36L; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119L; G120R; H124R; M129Y; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T;		++	++
1243/1244	T30R; R31P; K37R; A40L; Q43M; H44L; A56S; E68A; V70I; A84Q; D86A; A95V; Q119L; G120R; H124R; M129Y; T139M; S144R; N145F; H148T; V157A; M170F; N213E; A219T;		++	+
1245/1246	T30R; R31P; A36L; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119M; G120R; V123K; H124R; M129F; T139M; N145C; H148I; V157A; M170F; N213E; A219T;		++	++
1247/1248	T30R; R31P; A36L; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119L; G120R; H124R; T139M; N145C; H148I; V157A; M170F; N213E; A219T;		+	
1249/1250	T30R; R31P; K37R; A40L; Q43M; H44L; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; V123K; H124R; M129Y; T139M; N145F; H148K; V157A; M170F; N213E; A219T;		+	++
1251/1252	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; S144R; N145F; H148T; V157A; M170F; N213E; A219T;	+	+	+
1253/1254	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119L; G120R; H124R; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T;		+	
1255/1256	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119L; G120R; V123K; H124R;		+	

TABLE 2I-continued

SEQ ID	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 14 (24 h/82.5° C./ 4.2M MDEA challenge/ 45° C. assay/ 960 mM MDEA)	Assay 15 (24 h/85° C./ 4.2M MDEA challenge/ 45° C. assay/ 960 mM MDEA)	Assay 16 (24 h/90° C./ 4.2M MDEA challenge/ 45° C. assay/ 960 mM MDEA)
	T139M; S144R; N145F; H148I; V157A; M170F; N213E; A219T;			
1257/1258	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119L; G120R; V123K; H124R; T139M; S144R; N145C; H148I; V157A; M170F; N213E; A219T;		+	
1259/1260	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; V123K; H124R; M129Y; T139M; N145C; H148T; V157A; M170F; N213E; A219T;		+	
1261/1262	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119L; G120R; V123K; H124R; M129F; T139M; S144R; N145C; H148I; V157A; M170F; N213E; A219T;		+	
1263/1264	T30R; R31P; K37R; A40L; Q43M; H44L; A56S; E68A; V70I; A84Q; A95V; Q119L; G120R; V123K; H124R; T139M; N145C; H148T; V157A; M170F; N213E; A219T;		+	++
1265/1266	T30R; R31P; K37R; A40L; Q43M; H44L; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; N145C; H148T; V157A; M170F; N213E; A219T;		+	
1267/1268	T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; D86A; A95V; Q119M; G120R; V123K; H124R; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T;		+	+
1269/1270	T30R; R31P; K37R; A40L; Q43M; H44L; A56S; E68A; V70I; A84Q; A95V; Q119L; G120R; V123K; H124R; M129F; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T;		+	+

Levels of increased activity were determined relative to the reference polypeptide of SEQ ID NO: 1152 (i.e., engineered polypeptide having T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; N145F; H148T; V157A; M170F; N213E; A219T) and defined as "+", "++", or "+++" for each of the assays as follows:

Assay 14:

"+" indicates at least 1.1-fold but less than 1.3-fold increased activity;

"++" indicates at least 1.3-fold but less than 1.5-fold increased activity;

"+++" indicates at least 1.5-fold increased activity.

Assay 15:

"+" indicates at least 1.1-fold but less than 1.5-fold increased activity;

"++" indicates at least 1.5-fold but less than 2-fold increased activity;

"+++" indicates at least 2-fold increased activity.

Assay 16:

"+" indicates at least 1.1-fold but less than 1.3-fold increased activity;

"++" indicates at least 1.3-fold but less than 1.4-fold increased activity;

"+++" indicates at least 1.4-fold increased activity.

TABLE 2J

SEQ ID	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 17 (24 h/87° C./ 4.2M MDEA challenge/ 45° C. assay/ 685 mM MDEA)
1271/1272	T30R; R31P; K37R; A40L; Q43M; H44L; Y49F; I52V; A56S; E68Q; V70I; A84Q; A95V; Q119M; G120R; H124R; S126N; M129F; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T;	++
1273/1274	T30R; R31P; K37R; A40L; Q43M; H44L; A56S; E68A; V70I; A84Q; A95V; D96E; Q119M; G120R; H124R; M129F; T139M; S144R; N145C; H148T; V157A; M170F; D196T; N213E; A219T;	+
1275/1276	T30R; R31P; K37R; A40L; Q43M; H44L; A56S; E68Q; V70I; A84Q; A95V; Q119M; G120R; H124R; S126N; M129F; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T;	+
1277/1278	T30R; R31P; K37R; A40L; Q43M; H44L; Y49F; I52V; A56S; E68Q; V70I; I76V; A84Q; A95V; Q119M; G120R; H124R; S126N; M129F; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T;	++
1279/1280	T30R; R31P; K37R; A40L; Q43M; H44L; I52V; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; S126N; M129F; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T;	++

TABLE 2J-continued

SEQ ID NO: (nt/aa)	Amino Acid Residue Difference(s) (relative to SEQ ID NO: 2)	Assay 17 (24 h/87° C./ 4.2M MDEA challenge/ 45° C. assay/ 685 mM MDEA)
1281/1282	T30R; R31P; K37R; A40L; Q43M; H44L; A56S; E68A; V70I; A84Q; A95V; D96E; Q119M; G120R; H124R; M129F; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T;	+
1283/1284	T30R; R31P; K37R; A40L; Q43M; H44L; Y49F; A56S; E68Q; V70I; A84Q; A95V; Q119M; G120R; H124R; S126N; M129F; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T;	+++
1285/1286	T30R; R31P; K37R; A40L; Q43M; H44L; Y49F; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; S126N; M129F; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T;	++

Levels of increased activity were determined relative to the reference polypeptide of SEQ ID NO: 1210 (i.e., engineered polypeptide having T30R; R31P; K37R; A40L; Q43M; H44L; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; M129F; T139M; S144R; N145C; H148T; V157A; M170F; N213E; A219T) and defined as "+", "++", or "+++" for each of the assays as follows:
Assay 17:

"+" indicates at least 1.1-fold but less than 1.3-fold increased activity;

"++" indicates at least 1.3-fold but less than 1.5-fold increased activity;

"+++" indicates at least 1.5-fold increased activity.

In addition to the exemplary polypeptides of Tables 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J in some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide having an improved enzyme property relative to a polypeptide of SEQ ID NO:2, and an amino acid sequence having at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to a reference amino acid sequence selected from any one of which comprise an amino acid sequence selected from the polypeptide amino acid sequences disclosed in the accompanying Sequence Listing, specifically any one of the polypeptide amino acid sequences of SEQ ID NO: 4-1286 (which correspond to the even numbered sequence identifier numbers from 4 to 1286, inclusive).

In some embodiments, the disclosure provides a recombinant carbonic anhydrase an improved enzyme property relative to a reference polypeptide of SEQ ID NO:2, said recombinant polypeptide comprising an amino acid sequence having a feature selected from one or more: (a) having at least 93.7%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identity to SEQ ID NO:1152 or to a fragment of SEQ ID NO: 1152, wherein the fragment has at least 90%, 95%, 98%, or 99% of the length of SEQ ID NO: 1152; (b) having 14 or fewer, 13 or fewer, 12 or fewer, 11 or fewer, 10 or fewer, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid residue differences relative to SEQ ID NO: 1152; (c) having at least a combination of amino acid residue differences relative to SEQ ID NO: 2 present in any one of the polypeptide sequences of SEQ ID NO: 270-568, 570-678, 1058-1158, or 1174-1286; (d) having at least 93.7%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identity to SEQ ID NO:1162 or to a fragment of SEQ ID NO: 1162, wherein the fragment has at least 90%, 95%, 98%, or 99% of the length of SEQ ID NO: 1162; (e) having 14 or fewer, 13 or fewer, 12 or fewer, 11 or fewer, 10 or fewer, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid residue differences relative to SEQ ID NO: 1162; and (d) having at least a combination of amino acid residue differences relative to SEQ ID NO: 2 present in any one of the polypeptide sequences of SEQ ID NO: 680-1056, or 1160-1172.

Each of the exemplary recombinant carbonic anhydrase polypeptides shown in Tables 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J comprises one or more amino acid residue differences as compared to SEQ ID NO: 2, and has at least 1.3-fold, at least 1.5-fold, at least 2-fold, at least 3-fold, or at least 5-fold increased stability relative to the polypeptide of SEQ

ID NO: 2. Specific amino acid differences are shown in Tables 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J and include one or more residue differences as compared to SEQ ID NO:2 at the following residue positions: X2; X4; X15; X16; X22; X30; X31; X32; X34; X35; X36; X37; X40; X42; X43; X44; X47; X49; X52; X56; X60; X66; X68; X70; X76; X84; X86; X93; X95; X96; X97; X119; X120; X121; X123; X124; X126; X129; X131; X138; X139; X142; X143; X144; X145; X147; X148; X156; X157; X159; X168; X170; X178; X196; X200; X207; X213; X219; X221; X222; and X223. Some of these positions appear in more than one polypeptide with different amino acid replacements. The specific amino acid residue differences found in the exemplary polypeptides having an improved property are: X2R; X2T; X4F; X4M; X15R; X16S; X22G; X30A; X30K; X30L; X30Q; X30R; X31P; X32K; X32R; X34H; X35A; X35R; X36L; X36T; X37C; X37R; X40L; X40Q; X40W; X42A; X43M; X43V; X44L; X47R; X49F; X52V; X56S; X60C; X60V; X66Y; X68A; X68G; X68Q; X68V; X70I; X76V; X84K; X84N; X84Q; X84R; X84S; X86A; X93W; X95V; X96A; X96C; X96E; X96K; X97F; X119K; X119L; X119M; X119T; X120R; X121H; X121K; X121L; X121Q; X121T; X121V; X121W; X123K; X124F; X124G; X124R; X126N; X129K; X129R; X131L; X131F; X138F; X138L; X138W; X139H; X139K; X139M; X139Q; X142L; X143M; X143R; X144A; X144L; X144R; X145C; X145F; X145L; X145W; X147E; X147F; X147G; X147T; X148A; X148C; X148K; X148T; X156L; X157A; X159H; X159R; X159V; X168E; X170F; X178G; X196T; X200R; X207E; X207N; X213E; X213Q; X219T; X221C; X222C; X223C; and X223Q.

It will be apparent to the skilled artisan that the residue positions and specific residue differences of the present disclosure which have been shown to improve stability in solutions comprising amine compounds and/or ammonia can be used to generate recombinant carbonic anhydrase polypeptides besides the exemplary polypeptides of Tables 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J. It is contemplated that additional recombinant carbonic anhydrase polypeptides having improved properties can be prepared comprising various combinations of the amino acid residue differences of the exemplary polypeptides of Tables 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J. This has been demonstrated by the recombinant carbonic anhydrase polypeptides of Tables 2B and 2C, which have improved stability in the presence of an amine compound, were prepared by combining the single amino

acid difference of SEQ ID NO: 26 (i.e., X56S) with various other amino acid differences from Table 2A to create the improved polypeptides of even-numbered SEQ ID NO: 188-568.

Similarly, the recombinant carbonic anhydrase polypeptides of Table 2D, which have increased stability in the presence of an amine compound relative to the polypeptides of Tables 2B and 2C, were prepared by combining the combination of amino acid differences of SEQ ID NO: 332 (i.e., X30R, X40L, X56S, X84Q, X120R, and X139M) with various other amino acid differences disclosed herein to create the improved polypeptides of even-numbered SEQ ID NO: 570-678; the recombinant carbonic anhydrase polypeptides of Table 2G, which have increased stability in the presence of an amine compound relative to the polypeptides of Table 2D, were prepared by combining the combination of amino acid differences of SEQ ID NO: 656 (i.e., T30R; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; T139M; N145W; N213E; A219T) with various other amino acid differences disclosed herein to create the improved polypeptides of even-numbered SEQ ID NO: 1058-1158; the recombinant carbonic anhydrase polypeptides of Table 2I, which have increased stability in the presence of an amine compound relative to the polypeptides of Table 2G, were prepared by combining the combination of amino acid differences of SEQ ID NO: 1152 (i.e., T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; M129F; T139M; T139M; N145F; H148T; V157A; M170F; N213E; A219T) with various other amino acid differences disclosed herein to create the improved polypeptides of even-numbered SEQ ID NO: 1174-1270; the recombinant carbonic anhydrase polypeptides of Table 2J, which have increased stability in the presence of an amine compound relative to the polypeptides of Table 2I, were prepared by combining the combination of amino acid differences of SEQ ID NO: 1210 (i.e., T30R; R31P; K37R; A40L; Q43M; H44L; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; M129F; T139M; 5144R; N145C; H148T; V157A; M170F; N213E; A219T) with various other amino acid differences disclosed herein to create the improved polypeptides of even-numbered SEQ ID NO: 1272-1286.

The recombinant carbonic anhydrase polypeptides of Table 2E, which have increased stability in the presence of ammonia relative to the polypeptides of Table 2A, were prepared by combining the combination of amino acid differences of SEQ ID NO: 32 (i.e., Q15R and T30R) with various other amino acid differences disclosed herein to create the improved polypeptides of even-numbered SEQ ID NO: 680-972; the recombinant carbonic anhydrase polypeptides of Table 2F, which have increased stability in the presence of ammonia relative to the polypeptides of Table 2E, were prepared by combining the combination of amino acid differences of SEQ ID NO: 812 (i.e., Q15R, T30R, Q32K, S35A, A56S, A84N, and A221C) with various other amino acid differences disclosed herein to create the improved polypeptides of even-numbered SEQ ID NO: 974-1056; and the recombinant carbonic anhydrase polypeptides of Table 2H, which have increased stability in the presence of ammonia relative to the polypeptides of Table 2F, were prepared by combining the combination of amino acid differences of SEQ ID NO: 1056 (i.e., T4M; Q15R; T30R; Q32K; S35A; A56S; V70I; A84Q; A95V; V131L; T139Q; V157A; A221C) with various other amino acid differences disclosed herein to create the improved polypeptides of even-numbered SEQ ID NO: 1160-1172.

Accordingly, in some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide hav-

ing at least 1.3-fold, at least 1.5-fold, at least 2-fold, at least 3-fold, or at least 5-fold increased stability relative to the polypeptide of SEQ ID NO: 2, comprises an amino acid sequence having at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to SEQ ID NO: 2, and further comprises the one or more amino acid residue differences as compared to SEQ ID NO: 2 of any one of amino acid sequences of the even-numbered SEQ ID NO: 4-1286. In some embodiments, in addition to the set of amino acid residue differences of any one of the recombinant carbonic anhydrase polypeptides comprising an amino acid sequence of even-number SEQ ID NO: 4-1286, the sequence of the recombinant polypeptide can further comprise 1-2, 1-3, 1-4, 1-5, 1-6, 1-7, 1-8, 1-9, 1-10, 1-11, 1-12, 1-14, 1-15, 1-16, 1-18, 1-20, 1-22, 1-24, 1-26, 1-30, 1-35, 1-40 residue differences at other amino acid residue positions as compared to the SEQ ID NO: 2. In some embodiments, the residue differences can comprise conservative substitutions and/or non-conservative substitutions as compared to SEQ ID NO: 2.

In some embodiments, any of the recombinant carbonic anhydrase polypeptides having an improved property relative to the polypeptide of SEQ ID NO: 2 and an amino acid sequence having at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to SEQ ID NO: 2 and one or more of the following amino acid residue differences relative to SEQ ID NO: 2: X2R; X2T; X4F; X4M; X15R; X16S; X22G; X30A; X30K; X30L; X30Q; X30R; X31P; X32K; X32R; X34H; X35A; X35R; X36L; X36T; X37C; X37R; X40L; X40Q; X40W; X42A; X43M; X43V; X44L; X47R; X49F; X52V; X56S; X60C; X60V; X66Y; X68A; X68G; X68Q; X68V; X70I; X76V; X84K; X84N; X84Q; X84R; X84S; X86A; X93W; X95V; X96A; X96C; X96E; X96K; X97F; X119K; X119L; X119M; X119T; X120R; X121H; X121K; X121L; X121Q; X121T; X121V; X121W; X123K; X124F; X124G; X124R; X126N; X129K; X129R; X131L; X131F; X138F; X138L; X138W; X139H; X139K; X139M; X139Q; X142L; X143M; X143R; X144A; X144L; X144R; X145C; X145F; X145L; X145W; X147E; X147F; X147G; X147T; X148A; X148C; X148K; X148T; X156L; X157A; X159H; X159R; X159V; X168E; X170F; X178G; X196T; X200R; X207E; X207N; X213E; X213Q; X219T; X221C; X222C; X223C; and X223Q.

The positions associated with the improved property of increased stability in the presence of an amine compound include: X2; X4; X15; X16; X30; X31; X32; X34; X35; X36; X37; X40; X42; X43; X44; X47; X49; X52; X56; X60; X66; X68; X70; X76; X84; X86; X95; X96; X97; X119; X120; X121; X123; X124; X126; X129; X131; X138; X139; X142; X144; X145; X147; X148; X159; X168; X170; X178; X196; X200; X213; X219; X221; X222; and X223. The specific amino acid residue differences associated with the improved property of increased stability in the presence of an amine compound include: X2R; X2T; X4F; X4M; X15R; X16S; X30A; X30K; X30L; X30Q; X30R; X31P; X32K; X32R; X34H; X35A; X35R; X36L; X36T; X37R; X40L; X40W; X42A; X43M; X43V; X44L; X47R; X49F; X52V; X56S; X60C; X68A; X68G; X68Q; X70I; X76V; X84N; X84Q; X84R; X84S; X86A; X95V; X96A; X96C; X96E; X96K; X97F; X119K; X119L; X119M; X119T; X120R; X121H; X121K; X121L; X121Q; X121T; X121V; X121W; X123K; X124F; X124G; X124R; X126N; X131F; X131L; X138L; X138W; X139H; X139K; X139M; X139Q; X142L; X144A; X144L; X144R; X145C; X145F; X145L; X145W; X147E; X147F; X147G; X147T; X148A; X148T; X159H; X159R; X159V;

X168E; X170F; X178G; X196T; X200R; X213E; X213Q; X219T; X221C; X222C; and X223C.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide in which the increased stability in the presence of an amine compound comprises at least 1.3-fold increased activity relative to the reference polypeptide of SEQ ID NO: 2 after 24 hours exposure to 4 M MDEA at 42° C. and the amino acid sequence comprises one or more of the following amino acid residue differences relative to SEQ ID NO: 2: X2R; X2T; X4F; X4M; X15R; X16S; X30A; X30K; X30L; X30Q; X30R; X31P; X32K; X32R; X34H; X35A; X35R; X36L; X36T; X37R; X40L; X40W; X42A; X43M; X43V; X44L; X47R; X49F; X52V; X56S; X60C; X68A; X68G; X68Q; X70I; X76V; X84N; X84Q; X84R; X84S; X86A; X95V; X96A; X96C; X96E; X96K; X97F; X119K; X119L; X119M; X119T; X120R; X121H; X121K; X121L; X121Q; X121T; X121V; X121W; X123K; X124F; X124G; X124R; X126N; X131F; X131L; X138L; X139H; X139K; X139M; X139Q; X142L; X144A; X144L; X144R; X145C; X145F; X145L; X145W; X147E; X147F; X147G; X147T; X148A; X148T; X159H; X159R; X159V; X168E; X170F; X178G; X196T; X200R; X213E; X213Q; X219T; X221C; X222C; and X223C.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide in which the increased stability in the presence of an amine compound comprises at least 1.5-fold increased activity relative to the reference polypeptide of SEQ ID NO: 2 after 24 hours exposure to 4 M MDEA at 50° C. and the amino acid sequence comprises one or more of the following amino acid residue differences relative to SEQ ID NO: 2: X2T; X4F; X31P; X40L; X56S; X84Q; X119M; X120R; X121K; X121W; X131L; X139M; X147E; X147T; and X170F.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide in which the increased stability in the presence of an amine compound comprises at least 2-fold increased activity relative to the reference polypeptide of SEQ ID NO: 2 after 24 hours exposure to 4 M MDEA at 50° C. and the amino acid sequence comprises one or more of the following amino acid residue differences relative to SEQ ID NO: 2: X2T; X56S; X84Q; X139M; X147E; and X147T.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide in which the increased stability in the presence of an amine compound in which the increased stability in the presence of an amine compound comprises at least 3-fold increased activity relative to the reference polypeptide of SEQ ID NO: 2 after 24 hours exposure to 4 M MDEA at 50° C. and an amino acid sequence comprising one or more of the following amino acid residue differences relative to SEQ ID NO: 2: X56S; X84Q; X147E and X147T.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide in which the improved enzyme property is increased stability in the presence of an amine compound and in which the amino acid sequence comprises the amino acid difference X56S and one or more of the following amino acid residue differences relative to SEQ ID NO: 2: X2R; X2T; X4F; X4M; X15R; X16S; X30A; X30K; X30L; X30Q; X30R; X31P; X32K; X32R; X34H; X35A; X35R; X36L; X36T; X37R; X40L; X40W; X42A; X43M; X43V; X44L; X47R; X49F; X52V; X60C; X68A; X68G; X68Q; X70I; X76V; X84N; X84Q; X84R; X84S; X86A; X95V; X96A; X96C; X96E; X96K; X97F; X119K; X119L; X119M; X119T; X120R; X121H; X121K; X121L; X121Q; X121T; X121V; X121W; X123K; X124F; X124G; X124R; X126N; X131F; X131L; X138L; X139H;

X139K; X139M; X139Q; X142L; X144A; X144L; X144R; X145C; X145F; X145L; X145W; X147E; X147F; X147G; X147T; X148A; X148T; X159H; X159R; X159V; X168E; X170F; X178G; X196T; X200R; X213E; X213Q; X219T; X221C; X222C; and X223C.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide in which the improved enzyme property is increased stability in the presence of an amine compound characterized by at least 1.5-fold increased activity relative to the reference polypeptide of SEQ ID NO: 332 after 24 hours exposure to 5 M MDEA at 65° C. In some embodiments, the amino acid sequence of the recombinant carbonic anhydrase polypeptide having increased stability in the presence of an amine compound comprises at least the following amino acid residue differences relative to SEQ ID NO: 2: X30R, X40L, X56S, X84Q, X120R, and X139M. In some embodiments, the amino acid sequence comprises at least a combination of amino acid residue differences relative to SEQ ID NO: 2 present in any one of the polypeptide sequences of SEQ ID NO: 570-678, 1058-1158, or 1174-1286. In some embodiments, the amino acid sequence comprises any one of the polypeptide sequences of SEQ ID NO: 570-678, 1058-1158, or 1174-1286.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide in which the improved enzyme property is increased stability in the presence of an amine compound characterized by at least 1.5-fold increased activity relative to the reference polypeptide of SEQ ID NO: 656 after 24 hours exposure to 5 M MDEA at 70° C. In some embodiments, the amino acid sequence of the recombinant carbonic anhydrase polypeptide having increased stability in the presence of an amine compound comprises at least the amino acid residue differences relative to SEQ ID NO: 2: X30R, X37R, X40L, X56S, X68A, X84Q, X95V, X119M, X120R, X139M, X145W, X213E, and X219T. In some embodiments, the amino acid sequence comprises at least a combination of amino acid residue differences relative to SEQ ID NO: 2 present in any one of the polypeptide sequences of SEQ ID NO: 1058-1158, or 1174-1286. In some embodiments, the amino acid sequence comprises any one of the polypeptide sequences of SEQ ID NO: 1058-1158, or 1174-1286.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide in which the improved enzyme property is increased stability in the presence of an amine compound characterized by retaining at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, or at least about 95% or greater residual activity after exposure to a solution comprising at least 4.2 M MDEA at 50° C. for period of at least about 1, 4, 7, 12, or 14 days. In some embodiments, the amino acid sequence of the recombinant carbonic anhydrase polypeptide having increased stability characterized by retention of at least 30% residual activity after exposure to a solution comprising about 4.2 M MDEA at 50° C. for at least about 14 days comprises at least the amino acid residue differences relative to SEQ ID NO: 2: X30R; X31P; X37R; X40L; X43M; X56S; X68A; X70I; X84Q; X95V; X119M; X120R; X124R; X139M; X144R; X145F; X148T; X157A; X170F; X213E; and X219T. In some embodiments, the amino acid sequence further comprises at least 2, at least 3, at least 4, or at least 5 lysine (K) residues and/or arginine (R) residues substituted at positions X121-X126 and/or at positions X144-X149 relative to SEQ ID NO: 2. In some embodiments, the at least 2, at least 3, at least 4, or at least 5 lysine (K) residue and/or arginine (R)

residue substitutions are selected from: X84K, X84R, X120R, X121K, X123K, X124R, X129K, X129R, X139K, X143R, X144R, and X148K. In some embodiments, the at least 2, at least 3, at least 4, or at least 5 lysine (K) residues and/or arginine (R) residues substitutions are selected from: X84R, X123K, X124R, X129K, X129R, X143R, X144R, and X148K. In some embodiments, the amino acid sequence comprises at least a combination of amino acid residue differences relative to SEQ ID NO: 2 present in any one of the polypeptide sequences of SEQ ID NO: 1058-1158, or 1174-1286. In some embodiments, the amino acid sequence comprises any one of the polypeptide sequences of SEQ ID NO: 1058-1158, or 1174-1286.

The positions associated with the improved property of increased stability in the presence of an ammonia include: X2; X4; X15; X22; X30; X32; X34; X35; X37; X40; X42; X47; X56; X60; X68; X70; X84; X86; X93; X95; X96; X121; X124; X138; X143; X156; X157; X170; X207; X219; X221; X222; and X223. The specific amino acid residue differences associated with the improved property of increased stability in the presence of ammonia include: X2R; X4F; X4M; X15R; X22G; X30A; X30K; X30L; X30Q; X30R; X32K; X32R; X34H; X35A; X35R; X37C; X37R; X40L; X40Q; X40W; X42A; X47R; X56S; X60C; X60V; X68A; X68G; X68V; X70I; X84K; X84Q; X86A; X93W; X95V; X96E; X121H; X121K; X121Q; X121T; X121W; X124G; X131F; X138F; X138W; X143M; X143R; X148C; X156L; 157A; X170F; X207E; X207N; X219T; X221C; X222C; and X223Q.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide in which the increased stability in the presence of ammonia comprises at least 1.3-fold increased activity relative to the reference polypeptide of SEQ ID NO: 2 after 24 hours exposure to ammonia at 35° C. and the amino acid sequence comprises one or more of the following amino acid residue differences relative to SEQ ID NO: 2: X2R; X4F; X22G; X30A; X30L; X30Q; X30R; X32K; X32R; X34H; X35A; X35R; X37R; X40Q; X40W; X56S; X60C; X60V; X68A; X68G; X70I; X84K; X84Q; X86A; X95V; X121Q; X121T; X121W; X157A; X221C; X222C; and X223Q.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide in which the increased stability in the presence of ammonia comprises at least 3-fold increased activity relative to the reference polypeptide of SEQ ID NO: 2 after 24 hours exposure to ammonia at 35° C. and the amino acid sequence comprises one or more of the following amino acid residue differences relative to SEQ ID NO: 2: X4F; X22G; X30A; X30Q; X30R; X32K; X34H; X35A; X37R; X56S; X60C; X60V; X70I; X84Q; X121W; X221C; X222C; and X223Q.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide having increased stability in the presence of ammonia in which the amino acid sequence comprises one or more of the amino acid residue differences selected from X15R and X30R, and further comprises one or more of the following amino acid residue differences relative to SEQ ID NO: 2: X2R; X4F; X4M; X22G; X30A; X30K; X30L; X30Q; X32K; X32R; X34H; X35A; X35R; X37C; X37R; X40L; X40Q; X40W; X42A; X47R; X56S; X60C; X60V; X68A; X68G; X68V; X70I; X84K; X84Q; X86A; X93W; X95V; X96E; X121H; X121K; X121Q; X121T; X121W; X124G; X131F; X138F; X138W; X143M; X143R; X148C; X156L; 157A; X170F; X207E; X207N; X219T; X221C; X222C; and X223Q. In some embodiments, the amino acid sequence comprises both X15R and X30R and further comprises one or more of the following amino acid residue differences relative to SEQ ID

NO: 2: X2R; X4F; X4M; X22G; X32K; X32R; X34H; X35A; X35R; X37C; X37R; X40L; X40Q; X40W; X42A; X47R; X56S; X60C; X60V; X68A; X68G; X68V; X70I; X84K; X84Q; X86A; X93W; X95V; X96E; X121H; X121K; X121Q; X121T; X121W; X124G; X131F; X138F; X138W; X143M; X143R; X148C; X156L; 157A; X170F; X207E; X207N; X219T; X221C; X222C; and X223Q.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide in which the improved enzyme property is increased stability in the presence of ammonia characterized by at least 1.5-fold increased activity relative to the reference polypeptide of SEQ ID NO: 32 after 24 hours exposure to 5.6 M NH₃ at 44° C. In some embodiments, the amino acid sequence of the recombinant carbonic anhydrase polypeptide having increased stability in the presence of ammonia comprises at least the amino acid residue differences relative to SEQ ID NO: 2: X15R, and X30R. In some embodiments, the amino acid sequence comprises at least a combination of amino acid residue differences relative to SEQ ID NO: 2 present in any one of the polypeptide sequences of SEQ ID NO: 680-1056, or 1160-1172. In some embodiments, the amino acid sequence comprises any one of the polypeptide sequences of SEQ ID NO: 680-1056, or 1160-1172.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide in which the improved enzyme property is increased stability in the presence of ammonia characterized by at least 1.3-fold increased activity relative to the reference polypeptide of SEQ ID NO: 812 after 24 hours exposure to 8.4 M NH₃ at 58° C. In some embodiments, the amino acid sequence of the recombinant carbonic anhydrase polypeptide having increased stability in the presence of ammonia comprises at least the amino acid residue differences relative to SEQ ID NO: 2: X15R, X30R, X32K, X35A, X56S, X84N, and X221C. In some embodiments, the amino acid sequence comprises at least a combination of amino acid residue differences relative to SEQ ID NO: 2 present in any one of the polypeptide sequences of SEQ ID NO: 974-1056, or 1160-1172. In some embodiments, the amino acid sequence comprises any one of the polypeptide sequences of SEQ ID NO: 974-1056, or 1160-1172.

In some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide in which the improved enzyme property is increased stability in the presence of ammonia characterized by at least 1.3-fold increased activity relative to the reference polypeptide of SEQ ID NO: 1056 after 24 hours exposure to 8.4 M NH₃ at 70° C. In some embodiments, the amino acid sequence of the recombinant carbonic anhydrase polypeptide having increased stability in the presence of ammonia comprises at least the amino acid residue differences relative to SEQ ID NO: 2: X4M, X15R, X30R, X32K, X35A, X56S, X70I, X84Q, X95V, X131L, T139Q, V157A, and X221C. In some embodiments, the amino acid sequence comprises at least a combination of amino acid residue differences relative to SEQ ID NO: 2 present in any one of the polypeptide sequences of SEQ ID NO: 1160-1172. In some embodiments, the amino acid sequence comprises any one of the polypeptide sequences of SEQ ID NO: 1160-1172.

As described in Tables 2A 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J and the Examples, the improved property of increased stability and/or increased activity are determined under suitable conditions. In some embodiments, improved property comprises at least 1.2-fold, at least 1.3-fold, at least 1.5-fold, at least 2-fold, at least 3-fold, at least 4-fold, at least 5-fold, at least 10-fold, or at least 25-fold increased activity of hydrating carbon dioxide or dehydrating bicarbonate under suitable

conditions. In some embodiments, the suitable conditions comprise a carbonic anhydrase polypeptide concentration of from about 0.1 g/L to about 10 g/L, about 0.25 g/L to about 7.5 g/L, about 0.5 g/L to about 5 g/L, less than 10 g/L, less than about 5 g/L, or less than about 2.5 g/L. In some embodiments, the suitable conditions comprise a loading of solution CO₂ of from about $\alpha=0.005$ to about $\alpha=0.4$, from about $\alpha=0.01$ to about $\alpha=0.3$, $\alpha=0.015$ to about $\alpha=0.25$, $\alpha=0.02$ to about $\alpha=0.2$, less than about $\alpha=0.3$, less than about $\alpha=0.25$, or less than about $\alpha=0.2$.

In some embodiments the improved property is activity measured after exposure of the carbonic anhydrase to thermal or solvent challenge conditions. Accordingly in some embodiments, the increased activity is determined after heating the recombinant carbonic anhydrase polypeptide and the reference polypeptide at a temperature of from about 30° C. to 60° C. for a period of time of about 60 minutes to about 1440 minutes. In such embodiments, the fold-increase in activity corresponds to the same fold-increase in thermostability or solvent stability—depending on the challenge conditions. Various other challenge conditions may be used as disclosed in the Examples and elsewhere herein.

In some embodiments the improved property is stability in the presence of an amine compound and the suitable conditions comprise a solution comprising an amine compound selected from the group consisting of: 2-(2-aminoethylamino)ethanol (AEE), 2-amino-2-hydroxymethyl-1,3-propanediol (AHPD), 2-amino-2-methyl-1-propanol (AMP), diethanolamine (DEA), diisopropanolamine (DIPA), N-hydroxyethylpiperazine (HEP), N-methyldiethanolamine (MDEA), monoethanolamine (MEA), N-methylpiperazine (MP), piperazine, piperidine, 2-(2-tert-butylaminoethoxy) ethanol (TBEE), triethanolamine (TEA), triisopropanolamine (TIA), tris, 2-(2-aminoethoxy)ethanol, 2-(2-tert-butylaminopropoxy)ethanol, 2-(2-tert-amylaminoethoxy) ethanol, 2-(2-isopropylaminopropoxy)ethanol, 2-(2-(1-methyl-1-ethylpropylamino)ethoxy)ethanol, and mixtures thereof. In some embodiments, the amine compound is selected from AMP, MEA, MDEA, TIA, and mixtures thereof. Further, in some embodiments the suitable conditions comprise an amine compound at a concentration of from about 1 M to about 10 M, from about 2 M to about 8 M, from about 2.5 M to about 6.5 M, from about 3 M to about 5 M, at least about 2 M, at least about 3 M, at least about 4.2 M, or at least about 5 M.

Solutions of amine compounds used for carbon dioxide absorption from gas streams typically are used at elevated temperatures. Accordingly, in some embodiments the improved property is stability in the presence of an amine compound and the suitable conditions comprise a solution temperature of from about 40° C. to about 110° C., from about 40° C. to about 90° C., from about 40° C. to about 80°, from about 40° C. to about 70° C., or from about 40° C. to about 60° C.

Solutions containing ammonia that are used for carbon dioxide absorption from gas streams can be used at either or both chilled temperatures (e.g., for absorption) and elevated temperatures (e.g., for desorption of carbon dioxide). Accordingly, in some embodiments, the improved property is stability in ammonia and the suitable conditions comprise a solution temperature of from about 0° C. to about 20° C., from about 0° C. to about 10° C., from about 5° C. to about 15° C., from about 8° C. to about 12° C., less than about 15° C., or less than about 10° C. Further, in some embodiments the suitable conditions comprise an ammonia concentration of about 1 M to about 8 M, from about 2 M to about 7 M, from about 3 M

to about 6 M, at least about 1 M, at least about 2 M, at least about 3 M, at least about 4 M, or at least about 5 M, or at least about 5.6 M.

Some solutions for the absorption of carbon dioxide from gas streams include high concentrations of carbonate ion (CO₃²⁻). Typically, the carbonate ion is provided in the form of potassium carbonate (K₂CO₃) or sodium carbonate (Na₂CO₃). Accordingly, in some embodiments of the recombinant carbonic anhydrase polypeptides, the improved property is increased stability in solution comprising carbonate ion under suitable conditions, wherein the suitable conditions comprise a solution comprising carbonate ion at a concentration of about 0.1 M CO₃²⁻ to about 5 M CO₃²⁻, from about 0.2 M CO₃²⁻ to about 4 M CO₃²⁻, or from about 0.3 M CO₃²⁻ to about 3 M CO₃²⁻.

The present disclosure also contemplates a recombinant carbonic anhydrase polypeptide having at least 1.3-fold, at least 1.5-fold, at least 2-fold, at least 3-fold, or at least 5-fold increased stability relative to the polypeptide of SEQ ID NO: 2, wherein the recombinant polypeptide comprises an amino acid sequence having at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to SEQ ID NO: 2, and further comprises a set of amino acid residue differences as compared to SEQ ID NO: 2, wherein the amino acid differences are based on locations or regions in the structure of reference polypeptide (e.g., SEQ ID NO: 2) and/or the associated functional properties. Accordingly, referring to Table 3, a recombinant carbonic anhydrase polypeptide of the present disclosure can include an amino acid substitution at a particular residue at a location in the structure of the reference polypeptide as identified in Table 3. Exemplary substitutions at relevant locations include those identified in Tables 2A 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J.

TABLE 3

Position	Structural Location
X2	Surface Exposed
X3	Surface Exposed
X4	Surface Exposed
X6	Surface Exposed
X7	Surface Exposed
X8	Surface Exposed
X12	Surface Exposed
X13	Buried
X14	Buried
X15	Surface Exposed
X16	Surface Exposed
X17	Buried
X18	Surface Exposed
X19	Surface Exposed
X20	Buried
X21	Buried
X22	Surface Exposed
X23	Surface Exposed
X24	Buried
X25	Surface Exposed
X26	Surface Exposed
X27	Surface Exposed
X28	Surface Exposed
X30	Surface Exposed
X36	Surface Exposed
X37	Surface Exposed
X38	Surface Exposed
X39	Buried
X41	Surface Exposed
X42	Buried
X43	Surface Exposed
X44	Surface Exposed
X46	Active Site - Outer Sphere
X47	Surface Exposed
X48	Buried
X50	Buried

63

TABLE 3-continued

Position	Structural Location
X51	Buried
X52	Buried
X53	Buried
X54	Active Site - Outer Sphere - Buried
X55	Metal Coordinating - Buried
X56	Active Site - Outer Sphere - Buried
X57	Metal Coordinating - Buried
X58	Active Site - Outer Sphere - Buried
X59	Active Site - Outer Sphere - Buried
X60	Active Site - Outer Sphere - Buried
X61	Buried
X62	Buried
X63	Buried
X64	Buried
X65	Buried
X66	Buried
X67	Buried
X68	Buried
X69	Buried
X70	Buried
X71	Buried
X72	Buried
X73	Buried
X74	Active Site - Outer Sphere - Buried
X75	Buried
X76	Active Site - Outer Sphere - Buried
X77	Active Site - Outer Sphere - Buried
X78	Active Site - Outer Sphere - Buried
X79	Active Site - Inner Sphere - Buried
X80	Active Site - Inner Sphere - Buried
X81	Active Site - Outer Sphere - Buried
X82	Active Site - Outer Sphere - Buried
X83	Buried
X84	Buried - Dimer-dimer interface region
X85	Buried - Dimer-dimer interface region
X86	Buried
X87	Buried
X88	Buried
X89	Buried
X90	Active Site - Outer Sphere - Buried
X91	Buried
X92	Buried
X93	Active Site - Outer Sphere
X94	Active Site - Outer Sphere - Buried
X95	Buried
X97	Active Site - Outer Sphere - Surface Exposed
X98	Active Site - Outer Sphere - Surface Exposed
X100	Buried
X101	Buried
X102	Buried
X103	Buried
X104	Buried
X105	Buried
X106	Active Site - Outer Sphere - Buried
X107	Active Site - Outer Sphere - Buried
X108	Metal Coordinating - Buried
X109	Active Site - Outer Sphere
X110	Active Site - Outer Sphere - Surface Exposed
X111	Metal Coordinating - Buried
X112	Active Site - Inner Sphere
X113	Active Site - Inner Sphere - Buried
X114	Active Site - Outer Sphere - Buried
X115	Active Site - Outer Sphere - Surface Exposed
X116	Active Site - Outer Sphere
X117	Active Site - Outer Sphere - Buried
X119	Surface Exposed
X120	Buried - Dimer-dimer interface region
X121	Buried - Dimer-dimer interface region
X122	Surface Exposed - Dimer-dimer interface region
X123	Surface Exposed - Dimer-dimer interface region
X124	Buried - Dimer-dimer interface region
X125	Buried - Dimer-dimer interface region
X126	Surface Exposed - Dimer-dimer interface region
X127	Dimer-dimer interface region
X128	Dimer-dimer interface region
X129	Surface Exposed - Dimer-dimer interface region
X130	Buried - Dimer-dimer interface region
X131	Buried - Dimer-dimer interface region

64

TABLE 3-continued

Position	Structural Location
X132	Buried - Dimer-dimer interface region
X133	Buried - Dimer-dimer interface region
X134	Dimer-dimer interface region
X135	Active Site - Outer Sphere - Buried - Dimer-dimer interface region
X136	Buried - Dimer-dimer interface region
X137	Buried - Dimer-dimer interface region
X138	Dimer-dimer interface region
X139	Buried - Dimer-dimer interface region
X140	Dimer-dimer interface region
X141	Dimer-dimer interface region
X142	Buried - Dimer-dimer interface region
X143	Buried - Dimer-dimer interface region
X144	Buried - Dimer-dimer interface region
X145	Dimer-dimer interface region
X146	Buried - Dimer-dimer interface region
X147	Surface Exposed - Dimer-dimer interface region
X148	Surface Exposed - Dimer-dimer interface region
X149	Surface Exposed - Dimer-dimer interface region
X150	Surface Exposed
X151	Buried
X152	Surface Exposed
X153	Surface Exposed
X154	Surface Exposed
X155	Surface Exposed
X156	Surface Exposed
X157	Buried
X158	Active Site - Outer Sphere - Buried
X160	Buried
X161	Active Site - Outer Sphere - Buried
X162	Buried
X163	Surface Exposed
X164	Dimer-dimer interface region
X165	Buried - Dimer-dimer interface region
X166	Buried - Dimer-dimer interface region
X167	Dimer-dimer interface region
X168	Dimer-dimer interface region
X169	Buried - Dimer-dimer interface region
X170	Dimer-dimer interface region
X171	Dimer-dimer interface region
X172	Buried - Dimer-dimer interface region
X173	Buried - Dimer-dimer interface region
X174	Dimer-dimer interface region
X175	Dimer-dimer interface region
X176	Buried - Dimer-dimer interface region
X177	Surface Exposed - Dimer-dimer interface region
X178	Surface Exposed
X181	Surface Exposed
X182	Surface Exposed
X184	Surface Exposed
X185	Buried
X186	Buried
X187	Buried
X188	Buried
X189	Buried
X190	Buried
X191	Buried
X192	Active Site - Outer Sphere - Buried
X193	Surface Exposed
X194	Active Site - Outer Sphere - Buried
X195	Surface Exposed
X196	Surface Exposed
X197	Buried
X198	Surface Exposed
X199	Buried
X200	Surface Exposed
X201	Surface Exposed
X202	Surface Exposed
X203	Buried
X204	Surface Exposed
X205	Surface Exposed
X207	Surface Exposed
X208	Surface Exposed
X209	Surface Exposed

65 "inner sphere" - residue has an atom within 4.5 angstroms of the bound metal at active site.
"outer sphere" - residue within 4.5 angstroms of an inner sphere residue.

In some embodiments, any of the recombinant carbonic anhydrase polypeptides having at least 1.3-fold, at least 1.5-fold, at least 2-fold, at least 3-fold, or at least 5-fold increased stability relative to the polypeptide of SEQ ID NO: 2 and an amino acid sequence having at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to SEQ ID NO: 2, can further comprise at least one amino acid residue difference selected from each of at least two of the following seven sets (i.e., (a) through (g)) of amino acid residue differences: (a) X2R; X2T; X4F; (b) X121H; X121K; X121L; X121Q; X121T; X121V; X121W; X144A; X144L; X178G; (c) X139H; X139K; X139M; (d) X30A; X30L; X30Q; X30R; X40L; X40W; X68A; X96A; X96C; X96E; X96K; X119K; X119L; X119M; X119T; X120R; (e) X35R; X124G; X147E; X147F; X147G; X147T; X159H; X159R; (f) X31P; and (g) X56S; X84N; X84Q; X84S. In some embodiments, the recombinant carbonic anhydrase polypeptide amino acid sequence comprises one amino acid residue difference selected from each of at least two, three, four, five, six, or all seven of the sets of amino acid residue differences.

Structural modeling and homology analysis indicate that the wild-type β -class carbonic anhydrase polypeptide from *D. vulgaris* of SEQ ID NO: 2 can form a dimer-of-dimers protein structure. The dimer-dimer interface regions occur at amino acid positions X84-X85, X120-X149, and X164-X177 of the wild-type polypeptide monomer of SEQ ID NO: 2. In particular, strong dimer-dimer interface region interactions can occur between amino acids at residue positions X121-X126 of one monomer with amino acids at residue positions X144-X149 of the opposite monomer of the dimer. It is a surprising discovery of the present disclosure based on the amino acid differences found in the exemplary engineered carbonic anhydrase polypeptides of Tables 2G, 2I, and 2J, that amino acid residue differences providing positively charged lysine (K) or arginine (R) residues in the positions of the dimer-dimer interface region provide increased stability in the presence of amine compounds, such as MDEA. Accordingly, in some embodiments, the present disclosure provides a recombinant carbonic anhydrase polypeptide having increased stability in the presence of amine compound relative to the wild-type β -class carbonic anhydrase of SEQ ID NO: 2, wherein the polypeptide comprises an amino acid sequence having at least 80% identity to SEQ ID NO: 1152 and at least 2, at least 3, at least 4, at least 5, or more lysine (K) and/or arginine (R) residues substituted at positions X84-X85, X120-X149, and/or X164-X177 relative to SEQ ID NO: 2. In some embodiments, the recombinant carbonic anhydrase polypeptide comprises at least 2, at least 3, at least 4, or at least 5 lysine (K) residues and/or arginine (R) residues substituted at positions X121-X126 and/or at positions X144-X149 relative to SEQ ID NO: 2. In some embodiments, the at least 2, at least 3, at least 4, or at least 5 lysine (K) residue and/or arginine (R) residue substitutions are selected from: X84K, X84R, X120R, X121K, X123K, X124R, X129K, X129R, X139K, X143R, X144R, and X148K. In some embodiments, the at least 2, at least 3, at least 4, or at least 5 lysine (K) residues and/or arginine (R) residues substitutions are selected from: X84R, X123K, X124R, X129K, X129R, X143R, X144R, and X148K. In some embodiments, the recombinant carbonic anhydrase polypeptide with increased stability comprising an amino acid sequence having at least 80% identity to SEQ ID NO: 1152 and lysine (K) residues or arginine (R) residues substituted at positions X121-X126 and/or at positions X144-X149 relative to SEQ ID NO: 2, further is characterized in having at least 30% residual activity following exposure to a solution comprising 4.2 mMDEA

at 50° C. for a period of time of at least about 1 day, 4 days, 7 days, 12 days, 14 days, or longer.

An analysis of the amino acid sequences of other naturally occurring β -class homologs which have more than 40% identity to SEQ ID NO: 2, shows that approximately 85% have a valine at position X60. The β -class carbonic anhydrase from *D. vulgaris* of SEQ ID NO: 2 has an alanine at position X60. Structurally, the alanine at position X60 of SEQ ID NO: 2 resides just outside the metal binding site but contacts three of the four zinc coordinating residues C55, D57, and H108. Without being bound by mechanism, the structure-function correlation between the alanine at position X60 so close to the metal binding site and increased beta-class specific activity suggests that the volumetric change resulting from alanine rather than valine at position X60 results in greater active site flexibility, which in turn results in the greater catalytic efficiency of the β -class carbonic anhydrase from *D. vulgaris* even at lower temperatures (e.g., 5° C. to 15° C.).

In some embodiments, the present disclosure provides a β -class carbonic anhydrase polypeptide capable of hydrating carbon dioxide in a solution comprising an amine compound or ammonia, wherein the polypeptide comprises an amino acid sequence at least about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to any one of beta carbonic anhydrases polypeptides of SEQ ID NO:2, 1174, 1176, 1178, 1180, or 1182, and has an alanine residue at position X60 relative to SEQ ID NO: 2. In some embodiments of the β -class carbonic anhydrase polypeptides the amino acid sequence can further comprise one or more of the following amino acid residue differences relative to SEQ ID NO: 2: X2R; X2T; X4F; X4M; X15R; X16S; X22G; X30A; X30K; X30L; X30Q; X30R; X31P; X32K; X32R; X34H; X35A; X35R; X36T; X37C; X37R; X40L; X40Q; X40W; X42A; X43M; X43V; X47R; X56S; X60C; X60V; X68A; X68G; X68V; X70I; X84K; X84N; X84Q; X84R; X84S; X86A; X93W; X95V; X96A; X96C; X96E; X96K; X97F; X119K; X119L; X119M; X119T; X120R; X121H; X121K; X121L; X121Q; X121T; X121V; X121W; X124F; X124G; X124R; X131L; X131F; X138F; X138L; X138W; X139H; X139K; X139M; X139Q; X142L; X143M; X143R; X144A; X144L; X145C; X145F; X145L; X145W; X147E; X147F; X147G; X147T; X148A; X148C; X148T; X156L; X157A; X159H; X159R; X159V; X168E; X170F; X178G; X200R; X207E; X207N; X213E; X213Q; X219T; X221C; X222C; X223C; and X223Q.

In addition to the residue positions specified above, any of the recombinant carbonic anhydrase polypeptides disclosed herein can further comprise other residue differences relative to SEQ ID NO:2 at other residue positions. Residue differences at these other residue positions provide for additional variations in the amino acid sequence without adversely affecting the ability of the recombinant carbonic anhydrase polypeptide to carry out the hydration of carbon dioxide to bicarbonate and/or increased stability relative to the polypeptide of SEQ ID NO: 2. In some embodiments, the polypeptides can have additionally 1-2, 1-3, 1-4, 1-5, 1-6, 1-7, 1-8, 1-9, 1-10, 1-11, 1-12, 1-14, 1-15, 1-16, 1-18, 1-20, 1-22, 1-24, 1-26, 1-30, 1-35, 1-40 residue differences at other amino acid residue positions as compared to the reference sequence. In some embodiments, the number of differences can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 18, 20, 22, 24, 26, 30, 35, and 40 residue differences at other residue positions. The residue difference at these other positions can include conservative changes or non-conservative changes. In some embodiments, the residue differences can comprise conser-

vative substitutions and non-conservative substitutions as compared to the wild-type carbonic anhydrase of SEQ ID NO: 2.

In some embodiments, the present disclosure provides recombinant carbonic anhydrase polypeptides that comprise deletions of the recombinant carbonic anhydrase polypeptides expressly described herein. Thus, for each and every embodiment comprising an amino acid sequence, there is another embodiment comprising a sequence having one or more amino acid deletions, 2 or more amino acid deletions, 3 or more amino acid deletions, 4 or more amino acid deletions, 5 or more amino acid deletions, 6 or more amino acid deletions, 8 or more amino acid deletions, 10 or more amino acid deletions, 15 or more amino acid deletions, or 20 or more amino acid deletions, up to 10% of the total number of amino acids deleted, up to 20% of the total number of amino acids deleted, as long as the functional activity of the polypeptide with respect to the hydration of carbon dioxide to bicarbonate with increased stability is present. In some embodiments, the deletions can comprise, 1-2, 1-3, 1-4, 1-5, 1-6, 1-7, 1-8, 1-9, 1-10, 1-11, 1-12, 1-14, 1-15, 1-16, 1-18, 1-20, 1-22, 1-24, 1-26, 1-30, 1-35, or 1-40 amino acid residues. In some embodiments, the number of deletions can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 18, 20, 22, 24, 26, 30, 35, or 40 amino acids. In some embodiments, the deletions can comprise deletions of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18, or 20 amino acid residues.

In some embodiments, the polypeptides can comprise fragments of the recombinant carbonic anhydrase polypeptides described herein. In some embodiments, the fragments can have about 80%, 90%, 95%, 98%, and 99% of the full-length polypeptide, e.g., the polypeptide of SEQ ID NO:2, as long as the functional activity of the polypeptide with respect to the hydration of carbon dioxide to bicarbonate with increased stability is present.

In some embodiments, the polypeptides of the disclosure can be in the form of fusion polypeptides in which the recombinant carbonic anhydrase polypeptides are fused to other polypeptides, such as, by way of example and not limitation, antibody tags (e.g., myc epitope), purification sequences (e.g., His tags for binding to metals), and cell localization signals (e.g., secretion signals). Thus, the recombinant carbonic anhydrase polypeptides described herein can be used with or without fusions to other polypeptides.

The polypeptides described herein are not restricted to the naturally-occurring genetically encoded L-amino acids but also include the D-enantiomers of the genetically-encoded amino acids. In addition to the genetically encoded amino acids, the polypeptides described herein may be comprised, either in whole or in part, of naturally-occurring and/or synthetic non-encoded amino acids that are known in the art (see, e.g., the various amino acids provided in Fasman, 1989, CRC Practical Handbook of Biochemistry and Molecular Biology, CRC Press, Boca Raton, Fla., at pp. 3-70 and the references cited therein, all of which are incorporated by reference). For example, conformationally constrained non-encoded amino acids of which the polypeptides described herein may be composed include: N-methyl amino acids (L-configuration); 1-aminocyclopent-(2 or 3)-ene-4-carboxylic acid; pipercolic acid; azetidine-3-carboxylic acid; homoproline (hPro); and 1-aminocyclopentane-3-carboxylic acid. Additionally, those of skill in the art will recognize that amino acids bearing side chain protecting groups may also comprise the polypeptides described herein—e.g., Arg(tos), Cys(methylbenzyl), Cys(nitropyridinesulfonyl), Glu(δ -benzylester), Gln(xanthyl),

Asn(N- δ -xanthyl), His(bom), His(benzyl), His(tos), Lys(finoc), Lys(tos), Ser(O-benzyl), Thr (O-benzyl) and Tyr(O-benzyl).

As described above the various modifications introduced into the naturally occurring polypeptide to generate an engineered carbonic anhydrase enzyme can be targeted to a specific property of the enzyme.

Any of the above-described carbonic anhydrase polypeptides useful for chemical modification can be prepared by the ordinary artisan using the polynucleotide sequences disclosed herein (e.g., in Tables and Sequence Listing) and standard molecular biology and biochemical techniques for further mutagenesis, preparation, isolation, purification, and manufacture of the enzymes. For example, the disclosed polynucleotides may be operatively linked to one or more heterologous regulatory sequences that control gene expression to create a recombinant polynucleotide capable of expressing the polypeptide. Expression constructs containing a heterologous polynucleotide encoding the engineered carbonic anhydrase can be introduced into appropriate host cells to express the corresponding carbonic anhydrase polypeptide. Manipulation of the isolated polynucleotide prior to its insertion into an expression vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotides and nucleic acid sequences utilizing recombinant DNA methods are well known in the art. Guidance is provided in Sambrook et al., 2001, Molecular Cloning: A Laboratory Manual, 3rd Ed., Cold Spring Harbor Laboratory Press; and Current Protocols in Molecular Biology, Ausubel, F. ed., Greene Pub. Associates, 1998, updates to 2006. Example 1 of the present disclosure provides exemplary techniques.

Additionally, methods for producing the above-described carbonic anhydrase polypeptides in host cells are well-known to the skilled artisan. For example, polynucleotides for expression of the carbonic anhydrase may be introduced into host cells by various methods known in the art. Techniques include among others, electroporation, biolistic particle bombardment, liposome mediated transfection, calcium chloride transfection, and protoplast fusion. In some embodiments, more than one copy of a polynucleotide sequence is inserted into a host cell to increase production of the gene product. An increase in the copy number of the nucleic acid sequence can be obtained by integrating at least one additional copy of the sequence into the host cell genome or by including an amplifiable selectable marker gene with the nucleic acid sequence where cells containing amplified copies of the selectable marker gene, and thereby additional copies of the nucleic acid sequence, can be selected for by cultivating the cells in the presence of the appropriate selectable agent.

An exemplary host cells for use in producing the recombinant carbonic anhydrase polypeptides of the present disclosure is *Escherichia coli* W3110 and *Escherichia coli* BL21. An expression vector encoding an improved carbonic anhydrase of the present disclosure can be created by operatively linking a polynucleotide into the plasmid pCK110900 (see, U.S. application publication 20040137585) operatively linked to the lac promoter under control of the lac repressor. The expression vector also contained the P15a origin of replication and the chloramphenicol resistance gene. Cells containing the subject polynucleotide in *Escherichia coli* W3110 can be isolated by subjecting the cells to chloramphenicol selection. Example 1 of the present disclosure provides exemplary techniques.

The carbonic anhydrase enzymes expressed in a host cell can be recovered from the cells and or the culture medium using any one or more of the well known techniques for

protein purification, including, among others, lysozyme treatment, sonication, filtration, salting-out, ultra-centrifugation, and chromatography. Suitable solutions for lysing and the high efficiency extraction of proteins from bacteria, such as *E. coli*, are commercially available under the trade name Cel-Lytic BTM from Sigma-Aldrich of St. Louis Mo. Additionally, due to their enhanced thermostability, the engineered carbonic anhydrase polypeptides of the present disclosure can be recovered, isolated and/or purified from other cellular protein components by heat purification. Typically, after heating the desired engineered carbonic anhydrase remains in solution due its increased thermostability, but all or nearly all of the other protein components in the solution denature and can be separated easily from the solution by e.g., centrifugation. Methods for recovery of thermostable proteins by heat purification are well-known in the art.

8.5. Methods of Using Chemically Modified Carbonic Anhydrase Polypeptides

The chemically modified carbonic anhydrase enzymes described herein can catalyze both the forward and reverse reactions depicted in Scheme 1. In certain embodiments, the chemically modified carbonic anhydrase of the present disclosure can be used to hydrate carbon dioxide in the form of bicarbonate and a proton, which in turn, will be converted to carbonate and/or a mixture of bicarbonate and carbonate at an elevated pH. In other embodiments, a chemically modified carbonic anhydrase of the disclosure can be used to dehydrate carbon dioxide by reaction at a relatively acidic pH, thereby catalyzing the release of hydrated CO₂ from solution.

Accordingly, in some embodiments the present disclosure provides methods for removing carbon dioxide from a gas stream (e.g., capturing or extracting CO₂ gas) comprising the step of contacting the gas stream with a homogenous liquid solution under suitable conditions, wherein the solution comprises: (i) a chemically modified carbonic anhydrase polypeptide of the disclosure (e.g., chemically modified polypeptide having improved property such as increased activity, thermostability and/or solvent stability); and (ii) a CO₂ absorption mediating compound (e.g., ammonia, or an amine compound such as MDEA); whereby carbon dioxide from the gas stream is absorbed into the solution (e.g., CO₂ gas diffuses into solution and is hydrated to bicarbonate).

In some embodiments, the methods of use can be carried out wherein the chemically modified carbonic anhydrase polypeptide used is capable of catalyzing the hydration of carbon dioxide to bicarbonate or the reverse dehydration of bicarbonate to carbon dioxide with increased activity relative to the same carbonic anhydrases that are not chemically modified (and other known naturally occurring carbonic anhydrases) after exposure to high concentrations of CO₂ absorption mediating compound and/or thermal (e.g., T>40° C.). For example, in some embodiments, a chemically modified carbonic anhydrase of the present disclosure is used having carbonic anhydrase activity in 4.2 M MDEA at 50° C. that is increased (e.g., at least 1.5-fold, at least 2-fold, at least 4-fold, or even at least 5-fold increased) relative to the activity of the same carbonic anhydrase polypeptide that is not chemically modified (i.e., unmodified). Similarly, in some embodiments of the methods, the chemically modified carbonic anhydrase used is characterized by stability in 4.2 M MDEA at 75° C. that is increased (e.g., at least 1.5-fold, at least 2-fold, at least 4-fold, or even at least 5-fold increased) relative to the carbonic anhydrase polypeptide when it is not chemically modified.

The chemically modified carbonic anhydrase polypeptides having these (and other) improved properties useful in the methods include those disclosed elsewhere herein, include those provided in the Examples. In some embodiments, the method of use can be carried out wherein the carbonic anhydrase polypeptide chemically modified by treatment with a cross-linking agent is a naturally occurring carbonic anhydrase selected from an α -class, γ -class, β -class, or ξ -class carbonic anhydrase, or a recombinant (or engineered) carbonic anhydrase derived from a naturally occurring α -class, γ -class, β -class, or ξ -class carbonic anhydrase. In some embodiments, the polypeptide is an α -class carbonic anhydrase that comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 1298, 1300, 1302, 1304, 1306, and 1308, or a recombinant carbonic anhydrase polypeptide derived from any one of these α -class carbonic anhydrase sequences.

In some embodiments of the methods, the carbonic anhydrase polypeptide is a recombinant β -class carbonic anhydrase polypeptide derived from the wild-type *Desulfovibrio vulgaris* carbonic anhydrase comprising the amino acid sequence of SEQ ID NO: 2, or derived from a sequence homolog of SEQ ID NO: 2 selected from the group consisting of SEQ ID NO: 1288, 1290, 1292, 1294, and 1296. Engineered polypeptides useful in embodiments of the method are provided in Tables 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, and 2J. In some embodiments, the carbonic anhydrase polypeptide amino acid sequence comprises an even-numbered amino acid sequence selected from any one of SEQ ID NO: 4-1286. In such embodiments comprising a polypeptide based on the β -class polypeptide of SEQ ID NO: 2, the carbonic anhydrase polypeptide amino acid sequence has surface lysine residues at the following positions (relative to SEQ ID NO: 2): X18, X37, X147, X156, X184, or X198. Accordingly, in some embodiments of the methods the polypeptide is a recombinant β -class carbonic anhydrase having an amino acid sequence derived from SEQ ID NO: 2, and the treatment with a cross-linking agent results in a carbonic anhydrase polypeptide having a chemically modified lysine residue at one or more of the following positions relative to SEQ ID NO: 2: X18, X37, X147, X156, X184, or X198. In some embodiments of the methods of use, the carbonic anhydrase polypeptide amino acid sequence comprises at least the following amino acid residue difference relative to SEQ ID NO: 2: X56S. In some embodiments, the carbonic anhydrase polypeptide amino acid sequence comprises at least the following amino acid residue difference relative to SEQ ID NO: 2: X30R, X40L, X56S, X84Q, X120R, and X139M. In some embodiments of the methods of use, the carbonic anhydrase polypeptide amino acid sequence an amino acid sequence selected from any one of SEQ ID NO: 26, 190, 206, 238, 252, 270, 274, 284, 306, 318, 328, 332, 340, 354, 596, 606, 656, 678, 1080, 1110, 1148, 1152, 1156, and 1158.

In some embodiments of the method of use, the polypeptide is characterized by an amino acid sequence having at least 80% identity to SEQ ID NO:2 and at least one residue chemically modified by treatment with a cross-linking agent selected from the group consisting of: glutaraldehyde, dimethyl suberimidate, dimethyl pimelimidate, suberic acid bis (N-hydroxysuccinimide), and mixtures thereof. In some embodiments, the at least one residue that is chemically modified by treatment with a cross-linking agent is a surface lysine residue at one or more of the following positions relative to SEQ ID NO: 2: X18, X37, X147, X156, X184, or X198.

In some embodiments, the methods of removing carbon dioxide from a gas stream using a chemically modified car-

bonic anhydrase can be carried out wherein the carbonic anhydrase polypeptide that is chemically modified comprises a naturally occurring β -class carbonic anhydrase polypeptide of any one of SEQ ID NO: 2, 1288, 1290, 1292, 1294, and 1296. In some embodiments, the methods can be carried out using the carbonic anhydrase polypeptide of SEQ ID NO: 2. In some embodiments, the methods can be carried out using a β -class carbonic anhydrase polypeptide, wherein the polypeptide comprises an amino acid sequence at least about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical SEQ ID NO:2 and has an alanine residue at position X60 relative to SEQ ID NO: 2.

In some embodiments, the methods of removing carbon dioxide from a gas stream using a chemically modified carbonic anhydrase can be carried out wherein the carbonic anhydrase polypeptide that is chemically modified comprises an amino acid sequence that is at least about 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence of SEQ ID NO:2, and comprises one or more amino acid differences relative to SEQ ID NO: 2 selected from the group consisting of: X2R; X2T; X4F; X4M; X15R; X16S; X22G; X30A; X30K; X30L; X30Q; X30R; X31P; X32K; X32R; X34H; X35A; X35R; X37R; X40L; X40Q; X40W; X42A; X43M; X43V; X47R; X56S; X60C; X60V; X68A; X68G; X70I; X84K; X84N; X84Q; X84R; X84S; X86A; X95V; X96A; X96C; X96E; X96K; X97F; X119K; X119L; X119M; X119T; X120R; X121H; X121K; X121L; X121Q; X121T; X121V; X121W; X124G; X124R; X131L; X138F; X138L; X138W; X139H; X139K; X139M; X139Q; X142L; X143M; X144A; X144L; X145C; X145F; X145W; X147E; X147F; X147G; X147T; X148A; X148T; X157A; X159H; X159R; X159V; X168E; X170F; X178G; X200R; X207E; X207N; X213E; X213Q; X219T; X221C; X222C; X223C; and X223Q. The foregoing carbonic anhydrase polypeptides may further comprise additional modifications, including substitutions, deletions, insertions, or combinations thereof. The substitutions can be non-conservative substitutions, conservative substitutions, or a combination of non-conservative and conservative substitutions. In some embodiments, these carbonic anhydrase polypeptides can have optionally from about 1-2, 1-3, 1-4, 1-5, 1-6, 1-7, 1-8, 1-9, 1-10, 1-11, 1-12, 1-14, 1-15, 1-16, 1-18, 1-20, 1-22, 1-24, 1-25, 1-30, 1-35 or about 1-40 mutations at other amino acid residues. In some embodiments, the number of modifications can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 18, 20, 22, 24, 26, 30, 35 or about 40 other amino acid residues.

In certain embodiments, the methods can be carried out using a chemically modified carbonic anhydrase polypeptide of the present disclosure, wherein the polypeptide comprises an amino acid sequence selected from the amino acid sequences of SEQ ID NO: 4-1286 (which correspond to the even-numbered sequence identifier numbers from 4 to 1286, inclusive).

In some embodiments of the methods of use, the cross-linking agent is selected from the group consisting of a dialdehyde, a bis-imidate ester, a bis(N-hydroxysuccinimide) ester, a diacid chloride, and mixtures thereof. In some embodiments, the specific cross-linking agent is selected from the group consisting of malondialdehyde, glutaraldehyde, dimethyl suberimide, dimethyl pimelimidate, suberic acid bis(N-hydroxysuccinimide), and mixtures thereof. In some embodiments, the cross-linking agent is a dialdehyde having optionally one or more carbon atoms between the two aldehyde groups, for example wherein the dialdehyde is

selected from the group consisting of glyoxal, succindialdehyde, malondialdehyde, glutaraldehyde, and mixtures thereof. In a particular embodiment, the cross-linking agent is glutaraldehyde. In some embodiments, the cross-linking agent is an imidate ester, and in particular embodiments, a bis-imidate ester having optionally one or more carbon atoms between the two imidate ester groups. Useful imidate esters include bis-imidate esters having at least 1 carbon atoms between the two imidate ester groups, including but not limited to: imidate esters (such as methyl or ethyl) of malonimide, succinimide, glutarimide, adipimide, pimelimide, and suberimide.

In some embodiments of the methods of use, the cross-linking agent is a bis(N-hydroxysuccinimide) ester of a dicarboxylic acid that forms an irreversible chemical modification of the polypeptide. Useful bis(N-hydroxysuccinimide) esters include those prepared from a di-carboxylic acid selected from the group consisting of malonate, succinate, glutarate, adipate, pimelate, suberate, and mixtures thereof. Accordingly, in particular embodiments of the soluble composition, the cross-linking agent is a bis(N-hydroxysuccinimide) ester of a di-carboxylic acid selected from the group consisting of malonate, succinate, glutarate, adipate, pimelate, suberate, and mixtures thereof.

In other embodiments, the methods of use of the present disclosure can comprise further steps of isolating and/or separately treating the solution comprising the absorbed carbon dioxide. In some embodiments, the carbon dioxide gas in the solution is desorbed (i.e., stripped) by contacting the isolated solution with protons (i.e., acidify) and a chemically modified carbonic anhydrase polypeptide, which may be the same or different than the polypeptide used in the absorption step, thereby converting the hydrated carbon dioxide to carbon dioxide gas and water. In some embodiments, the desorption of carbon dioxide from this separate solution can be carried out at significantly higher temperatures, and/or under lower pressure (e.g., vacuum) conditions that can require a carbonic anhydrase polypeptide (modified or unmodified) having different stability characteristics. Thus, it is contemplated that the solution can be removed from contact with the gas stream (e.g., isolated after some desired level of hydrated carbon dioxide is reached) and further treated with a chemically modified or unmodified carbonic anhydrase to convert the bicarbonate in solution into carbon dioxide gas, which is then released from the solution and sequestered, e.g., into a pressurized chamber.

In some embodiments, the methods for removing carbon dioxide from a gas stream of the present disclosure can comprise a further desorption step comprising exposing the solution comprising the chemically modified carbonic anhydrase polypeptide and absorbed carbon dioxide to suitable conditions for desorbing the carbon dioxide from the solution. In some embodiments, the suitable conditions for desorbing the carbon dioxide from the solution comprise heating the solution to an elevated temperature. In some embodiments, the suitable conditions for desorbing the carbon dioxide from the solution comprise exposing the solution to low pressure or a vacuum. (See e.g., Publ. U.S. Appl. No. 2007/0256559A1.) In some embodiments of the methods using the chemically modified carbonic anhydrase polypeptides of the present disclosure (which exhibit increased stability at elevated temperatures), the elevated temperatures for desorption can comprise a temperature of from about 40° C. to about 120° C., from about 50° C. to about 100° C., from about 50° C. to about 90° C., or at least about 40° C., at least about 50° C., at least about 60° C., at least about 70° C., at least about 80° C., or at least about 90° C.

In other embodiments, the further step of isolating the solution comprising the hydrated carbon dioxide is carried out and no further chemically modified carbonic anhydrase polypeptide is added to the solution. Instead the solution which is enriched in bicarbonate (i.e., hydrated carbon dioxide) can be used in processes that react with the bicarbonate to effectively sequester the carbon dioxide in another chemical form.

In some embodiments, the chemically modified carbonic anhydrases and associated methods for removing (e.g., extracting and sequestering) carbon dioxide from a gas stream disclosed herein can be used in existing systems that use a solution for absorbing carbon dioxide from e.g., flue gas. Equipment, processes, and methods for carbon dioxide capture and sequestration using solutions into which carbon dioxide is absorbed (i.e., captured by diffusing from gas stream into the liquid solution) and/or from which carbon dioxide is desorbed (i.e., extracted by diffusing from liquid solution into gas phase) are described in e.g., U.S. Pat. Nos. 6,143,556, 6,524,843 B2, 7,176,017 B2, 7,596,952 B2, 7,641,717 B2, 7,645,430 B2, 7,579,185 B2, 7,740,689 B2, 7,132,090 B2; U.S. Pat. Publ. Nos. 2007/0004023A1, 2007/0256559A1, 2009/0155889A1, 2010/0086983A1; PCT Publ. Nos. WO98/55210A1, WO2004/056455A1, WO2004/028667A1, WO2006/089423A1, WO2008/072979A1, WO2009/000025A1, WO2010/020017A1, WO2010/014773A1, WO2010/045689A1, each of which is hereby incorporated by reference herein.

Methods for linking (covalently or non-covalently) enzymes to solid-phase particles (e.g., porous or non-porous beads, or solid supports) such that they retain activity for use in bioreactors are known in the art. Methods for treating a gas stream using immobilized enzymes are described in e.g., U.S. Pat. No. 6,143,556, U.S. patent publication no. 2007/0004023A1, and PCT publications WO98/55210A1, WO2004/056455A1, WO2004/028667A1, WO2011/014955A1, WO2011/014956A1, and WO2011/014957A1, each of which is hereby incorporated by reference herein. Accordingly, in alternative embodiments, the methods for removing carbon dioxide from a gas stream can be carried out wherein a chemically modified carbonic anhydrase polypeptide of the present disclosure is immobilized on a surface, for example linked to the surface of a solid-phase particle (e.g., beads) in the solution. Such methods result in a biphasic (or heterogeneous) solution comprising the immobilized chemically modified carbonic anhydrase polypeptide and the solution comprising CO₂ and a CO₂ absorption mediating compound. In such embodiments, the methods using immobilized chemically modified carbonic anhydrase polypeptides can be carried out wherein the method further comprises a step of isolating or separating the immobilized chemically modified carbonic anhydrase polypeptide from the solution. After separating the immobilized chemically modified carbonic anhydrase from the solution, the solution can be treated to conditions that may inactivate the enzyme, e.g., desorption of CO₂ at high temperatures. Further, the separately retained immobilized enzyme can be added to another solution and reused.

In various embodiments, the methods of removing carbon dioxide from a gas stream using a chemically modified carbonic anhydrase polypeptide disclosed herein may be carried out under a range of suitable conditions. Suitable conditions can be determined by routine experimentation that includes, but is not limited to, contacting the solution containing the chemically modified carbonic anhydrase polypeptide with CO₂ at an experimental condition (e.g., amine concentration, temperature, CO₂ loading) and then detecting the relevant

activity (e.g., rate of CO₂ absorption), for example, using the methods described in the Examples provided herein.

The ordinary artisan also will recognize that certain suitable conditions can be selected that favor the absorption of carbon dioxide into a solution (e.g., via hydration of carbon dioxide to bicarbonate) and/or the desorption of carbon dioxide from a solution (e.g., via dehydration of bicarbonate to carbon dioxide and water). The chemically modified carbonic anhydrase polypeptides disclosed herein are biocatalysts having an improved property (e.g., increased activity or thermal stability) that allows them to accelerate the absorption of carbon dioxide gas into a solution and/or accelerate subsequent desorption from the solution under a range of conditions.

In some embodiments, the method can be carried out wherein the chemically modified carbonic anhydrase comprises the improved property at least 1.2-fold, at least 1.3-fold, at least 1.5-fold, at least 2-fold, at least 3-fold, at least 4-fold, at least 5-fold, at least 10-fold, or at least 25-fold increased activity of hydrating carbon dioxide or dehydrating bicarbonate under suitable conditions. Accordingly, in some embodiments, the suitable conditions used in the method can comprise a concentration of the chemically modified carbonic anhydrase polypeptide of from about 0.1 g/L to about 10 g/L, about 0.25 g/L to about 7.5 g/L, about 0.5 g/L to about 5 g/L, less than 10 g/L, less than about 5 g/L, or less than about 2.5 g/L.

The ability of the chemically modified carbonic anhydrase polypeptide to accelerate CO₂ absorption into or desorption from a solution can be affected by the mole ratio of CO₂ to other compounds already present in the solution, which is also referred to as the CO₂ loading of the solution and can be denoted by the mole ratio of CO₂ to the moles of the relevant CO₂ absorption mediating compound in the solution (e.g., amine compound, ammonia), which is denoted by the term "α." The carbonic anhydrase polypeptides of the present disclosure can be used under a range of loading conditions which can be varied depending on the particular CO₂ absorption mediating compound used in the solution. Accordingly, the methods of the disclosure can be carried out wherein the suitable conditions comprise a loading of solution CO₂ of from about α=0 to about α=0.7, from about α=0.01 to about α=0.6, from about α=0.02 to about α=0.5, from about α=0.05 to about α=0.4, from about α=0.1 to about α=0.4, from about α=0.2 to about α=0.3, less than about α=0.7, less than about α=0.5, or less than about α=0.3.

In some embodiments where the method is carried out in the presence of an amine compound (e.g., 4 M MDEA) the suitable conditions can comprise a loading of solution CO₂ of from about α=0 to about α=0.6, from about α=0.01 to about α=0.5, from about α=0.02 to about α=0.4, from about α=0.05 to about α=0.3, from about α=0.1 to about α=0.4, from about α=0.2 to about α=0.3, less than about α=0.4, less than about α=0.3, or less than about α=0.2.

In some embodiments where the method is carried out in the presence of ammonia (e.g., 10 wt % or 5.6 M NH₃) the suitable conditions can comprise a loading of solution CO₂ of from about α=0 to about α=0.7, from about α=0.1 to about α=0.7, from about α=0.1 to about α=0.5, from about α=0.1 to about α=0.3, from about α=0.4 to about α=0.7, from about α=0.5 to about α=0.7, less than about α=0.7, less than about α=0.5, or less than about α=0.3.

Additionally, the CO₂ loading of the solution can change from "lean" to "rich" during the process as the CO₂ is absorbed, and then desorbed. Typically, the initial condition of the solution used in the method is "lean loading" (e.g., α=0, or α=0.01 to 0.02), and as the absorption proceeds the solu-

tion condition becomes "rich loading" (e.g., $\alpha=0.2$ to 0.5 , or higher). As illustrated by the Examples herein, the acceleration of CO_2 absorption due to enzyme tends to be lower under "lean loading" conditions than under "rich loading" conditions. Further the loading conditions used for the method carried out in the presence of amine compounds tends to be lower than the loading used for the method carried out in the presence of ammonia. Accordingly, in some embodiments, the suitable conditions in the presence of an amine compound comprise a lean loading of solution CO_2 from about $\alpha=0$ to about $\alpha=0.02$ and a rich loading of solution CO_2 of from about $\alpha=0.2$ to about $\alpha=0.5$. However, in some embodiments, where the suitable conditions include the presence of ammonia, the loading can comprise a lean loading of solution CO_2 about $\alpha=0.1$ to about $\alpha=0.3$ and a rich loading of solution CO_2 of from about $\alpha=0.5$ to about $\alpha=0.7$.

Typically the gas streams from which CO_2 removal is desirable are at elevated temperatures, and upon contacting a solution, as in the method disclosed herein, heat is also transferred and the solution temperature also is elevated. This is particularly true in treating flue gas streams from coal-fired power plants. Accordingly, in some embodiments, the suitable conditions for carrying out the method comprise an elevated solution temperature. The presence of elevated temperature further underscores the importance of using thermostable carbonic anhydrase polypeptides such as those disclosed herein. Thus, in some embodiments the method is carried out wherein the suitable conditions comprise a solution temperature of from about 40°C . to about 110°C ., from about 40°C . to about 90°C ., from about 40°C . to about 80°C ., from about 40°C . to about 70°C ., or from about 40°C . to about 60°C .

The method of removing carbon dioxide disclosed herein involves contacting the gas stream with a solution comprising a chemically modified carbonic anhydrase polypeptide. The present disclosure has illustrated the use of the method in solutions comprising a high concentration of an amine compound, ammonia, and carbonate ion. A range of other solutions comprising other compounds known to facilitate the absorption of CO_2 from a gas stream, and it is contemplated that the present methods could be used with such solutions.

For capturing CO_2 from flue gas streams, solutions comprising a variety of different amine compounds are known. Such solutions comprising amine compounds that facilitate CO_2 absorption from a gas stream into a solution are described in e.g., PCT Publ. No. WO2006/089423A1, U.S. Pat. No. 7,740,689 B2, or U.S. Pat. Publ. No. 2009/0155889A1, each which is hereby incorporated by reference herein. Accordingly, in some embodiments, the methods of removing carbon dioxide from a gas stream can be carried out wherein the solution comprises an amine compound, preferably an amine compound that exhibits improved thermodynamic and kinetic properties for the absorption of CO_2 . Thus, in some embodiments of the methods, the suitable conditions comprise a solution comprising an amine compound, and the amine compound can be selected from the group consisting of: 2-(2-aminoethylamino)ethanol (AEE), 2-amino-2-hydroxymethyl-1,3-propanediol (AHPD), 2-amino-2-methyl-1-propanol (AMP), diethanolamine (DEA), diisopropanolamine (DIPA), N-hydroxyethylpiperazine (HEP), N-methyldiethanolamine (MDEA), monoethanolamine (MEA), N-methylpiperazine (MP), piperazine, piperidine, 2-(2-tert-butylaminoethoxy)ethanol (TBEE), triethanolamine (TEA), trisopropanolamine (TIA), tris, 2-(2-aminoethoxy)ethanol, 2-(2-tert-butylaminopropoxy)ethanol, 2-(2-tert-amylaminoethoxy)ethanol, 2-(2-isopropylaminopropoxy)ethanol, 2-(2-(1-methyl-1-ethylpropylamino)ethoxy)ethanol, and mixtures thereof. In

some embodiments, the amine compound is selected from the group consisting of: AMP, MEA, MDEA, TIA, and mixtures thereof. In one preferred embodiment the solution comprises the amine compound MDEA. Further, in the embodiments of the methods employing an amine compound in solution, the suitable conditions can comprise an amine compound concentration of from about 1 M to about 10 M, from about 2 M to about 8 M, from about 2.5 M to about 6.5 M, from about 3 M to about 5 M, at least about 2 M, at least about 3 M, at least about 4.2 M, or at least about 5 M.

Elevated temperatures are typically present when the method employs a solution comprising an amine compound are used to remove carbon dioxide from a gas stream. Thus, in some embodiments the method is carried out wherein the suitable conditions comprise a solution comprising an amine compound (e.g., MDEA) and a temperature of from about 40°C . to about 110°C ., from about 40°C . to about 90°C ., from about 40°C . to about 80°C ., from about 40°C . to about 70°C ., or from about 40°C . to about 60°C .

Another known process for capturing CO_2 from a gas stream (e.g., flue gas) uses a solution containing a high concentration of ammonia. Methods and conditions for capturing CO_2 using solutions comprising ammonia are described in e.g., WO2009/000025A1, WO2010/020017A1, and WO2010/045689A1, each which is hereby incorporated by reference herein. Due to the high volatility of ammonia vapor processes using ammonia are sometimes run at relatively low temperatures, in what is referred to as a "chilled ammonia" process. Methods and conditions of the chilled ammonia process for CO_2 capture from a flue gas stream are described in e.g., U.S. Pat. No. 7,641,717 B2, and U.S. Pat. Publ. No. 2009/0155889A1, each which is hereby incorporated by reference herein. Accordingly, in some embodiments of the methods of removing carbon dioxide disclosed herein, a solution containing ammonia is used to facilitate carbon dioxide absorption from the gas streams. Such ammonia solutions can be used under suitable conditions comprising an ammonia concentration of about 1 M to about 8 M, from about 2 M to about 7 M, from about 3 M to about 6 M, at least about 1 M, at least about 2 M, at least about 3 M, at least about 4 M, or at least about 5 M, or at least about 5.6 M. Further in some embodiments of the methods, the solution comprising ammonia can be used at chilled temperatures (e.g., for absorption) and/or elevated temperatures (e.g., for desorption of carbon dioxide). Accordingly, in some embodiments, the method using a solution comprising ammonia can be carried out wherein the suitable conditions comprise a solution temperature of from about 0°C . to about 20°C ., from about 0°C . to about 10°C ., from about 5°C . to about 15°C ., from about 8°C . to about 12°C ., less than about 15°C ., or less than about 10°C .

Some processes for CO_2 capture from a gas stream use contact with a solution comprising elevated concentration of carbonate ions (CO_3^{2-}). Various formulations and processes for CO_2 capture from gas streams using solutions comprising carbonate ions are known (see e.g., WO2011/014957A1). Typically, the carbonate ion is provided in the solution in the form of potassium carbonate (K_2CO_3) or sodium carbonate (Na_2CO_3). In such embodiments, the stability and activity of the chemically modified carbonic anhydrase in the presence of carbonate ions is an important functional characteristic. Accordingly, in some embodiments, the method of removing CO_2 from a gas stream can be carried out wherein the suitable conditions comprise a solution comprising carbonate ion at a concentration of about 0.1 M CO_3^{2-} to about 5 M CO_3^{2-} , from about 0.2 M CO_3^{2-} to about 4 M CO_3^{2-} , or from about

0.3 M CO_3^{2-} to about 3 M CO_3^{2-} , at least about 0.2 M Na_2CO_3 , at least about 0.4 M Na_2CO_3 , or at least about 1 M Na_2CO_3 .

In some embodiments, the chemically modified carbonic anhydrase polypeptide of the present disclosure can be used in processes for CO_2 capture from a gas stream that comprise contacting the gas stream with a solution comprising the polypeptide and elevated concentration of amino acid compounds. In some embodiments, the amino acid compound is a primary, secondary, or tertiary amino acid, or a derivative or salt thereof (e.g., a sodium salt). Exemplary amino acid compounds useful in the method include, but are not limited to, the twenty most prevalent naturally occurring α -amino acids (i.e., alanine, leucine, valine, isoleucine, glycine, methionine, aspartic acid, glutamic acid, lysine, arginine, asparagine, glutamine, serine, threonine, histidine, tyrosine, tryptophan, phenylalanine, cysteine, and proline), as well as, taurine, methyl taurine, dimethyl-glycine, diethyl-glycine, N-butyl-glycine, N-methyl-alanine, sarcosine, and mixtures thereof. Various formulations and processes for CO_2 capture from gas streams using solutions comprising amino acid compounds are known (see e.g., WO2011/014955A1). In one embodiment of the methods of CO_2 capture from a gas stream of the present disclosure, the suitable conditions comprise the presence of the amino acid compound sodium glycinate.

Generally, in the methods of the present disclosure, the solution comprises an aqueous solvent (water or aqueous co-solvent system) that may be pH buffered or unbuffered. Generally, the CO_2 absorption reaction via hydration of carbon dioxide can be carried out by the carbonic anhydrase polypeptides over a pH range of about pH 9 or above or at a pH of about pH 10 or above, usually in the range of from about 8 to about 12. During the course of both the hydration and the dehydration reactions, the pH of the reaction mixture may change. The pH of the reaction mixture may be maintained at a desired pH or within a desired pH range by the addition of an acid or a base during the course of the reaction. Alternatively, the pH may be controlled by using an aqueous solvent that comprises a buffer. Suitable buffers to maintain desired pH ranges are known in the art and include, for example, carbonate, HEPES, triethanolamine buffer, and the like. The ordinary artisan will recognize that other combinations of buffering and acid or base additions known in the art may also be used. In some embodiments, the methods can be carried out in a solution at a basic pH that thermodynamically and/or kinetically favors the solvation of CO_2 —e.g., from about pH 8 to about pH 12. Accordingly, in some embodiments, the rate is determined at a pH of from about pH 8 to about pH 12, from about pH 9 to about pH 11.5, or from about pH 9.5 to pH 11. In other embodiments, release (dehydration) of captured carbon dioxide (e.g., as bicarbonate) is carried out at a pH of about 9 or below, usually in the range of from about pH 5 to about pH 9, or about pH 6 to about pH 9. In some embodiments, the dehydration is carried out at a pH of about 8 or below, often in the range of from about pH 6 to about pH 8.

In some embodiments, the methods of removing carbon dioxide from a gas stream disclosed herein, the solution can comprise an aqueous co-solvent system. For example, certain co-solvents or compounds can be added to the aqueous solution to reduce their degradative or corrosive properties. In some embodiments of the method, the solution is an aqueous co-solvent system comprising a ratio of water to a co-solvent from about 95:5 (v/v) to about 5:95 (v/v), in some embodiments, from about 90:10 (v/v) to about 10:90 (v/v), in some embodiments, from about 80:20 to about 20:80 (v/v), in some embodiments, from about 70:30 (v/v) to about 30:70 (v/v), and in some embodiments, from about 60:40 (v/v) to about

40:60 (v/v). The solvent component of an aqueous co-solvent system may be miscible with the aqueous component, providing a single liquid phase, or may be partly miscible or immiscible with the aqueous component, providing two liquid phases. The co-solvent system may be pre-formed prior to addition to the reaction mixture, or it may be formed in situ in the reaction vessel.

Co-solvent systems used in the methods typically comprise a solvent or compound that thermodynamically and/or kinetically favors the solvation of CO_2 from a gas-solvent interface. In some embodiments, the co-solvent in the aqueous solution is an amine compound (e.g., AMP, MDEA, MEA, TEA, and/or TIA). In some embodiments of the methods disclosed herein, the solution can comprise a mixture or blend of amine compounds, and/or other compounds that facilitate the absorption of CO_2 into the solution, e.g., ammonia, carbonate ions, strong base (e.g., NaOH), and/or compounds such as dimethyl ether of polyethylene glycol (PEG DME).

In some embodiments, the aqueous co-solvent systems can have water and one or more organic solvents. In general, an organic solvent component of an aqueous co-solvent system is selected such that it does not completely inactivate the chemically modified carbonic anhydrase enzyme. Appropriate co-solvent systems can be readily identified by measuring the enzymatic activity of the specified chemically modified carbonic anhydrase enzyme in the candidate solvent system, utilizing an enzyme activity assay, such as those described herein.

9. EXAMPLES

Various features and embodiments of the disclosure are illustrated in the following representative examples, which are intended to be illustrative, and not limiting.

Example 1

Preparation and Screening of Carbonic Anhydrase Polypeptides Based on Wild-Type Gene from *Desulfovibrio Vulgaris*

This example illustrates designing and optimizing the wild-type carbonic anhydrase gene from *Desulfovibrio vulgaris*, as well as further optimization and functional screening of the gene to generate engineered polypeptides having increased solvent and thermostability under conditions suitable for CO_2 absorption from gas into a capture solvent.

Gene Acquisition, Synthesis, Cloning, and Expression:

The gene encoding a wild-type *Desulfovibrio vulgaris* carbonic anhydrase polypeptide of SEQ ID NO: 2 was codon-optimized for expression in *E. coli* as the nucleotide sequence of SEQ ID NO: 1. The codon-optimized gene of SEQ ID NO: 1 was synthesized using oligonucleotides, generally composed of 42 nucleotides, and cloned into the expression vector pCK110900 under the control of a lac promoter. This expression vector also contains the P15a origin of replication and the chloramphenicol resistance gene. Resulting plasmids containing the codon-optimized were transformed into *E. coli* W3110 using standard methods. The transformed wild-type gene sequence of SEQ ID NO: 1 was confirmed by standard sequencing techniques and the resultant expression of carbonic anhydrase activity by the transformed cells confirmed by high throughput activity assays as described below.

Preparation of Engineered Carbonic Anhydrase Library:

Using the codon-optimized wild-type gene of SEQ ID NO: 1 as the starting point, a library of engineered variant genes was synthesized that targeted every residue from position X2

to position X223 of SEQ ID NO: 2 with substitutions of all 19 amino acids. The resulting engineered carbonic anhydrases polypeptide sequences, specific amino acid differences, and relative level of improvement are listed in Tables 2A and 2B.

Cloning of Engineered Carbonic Anhydrase Genes:

As with the codon-optimized wild-type gene of SEQ ID NO: 1, the library of engineered variant genes was cloned into vector pCK110900 and expressed in *E. coli* W3110. Antibiotic resistant transformants were selected and processed to identify those expressing a CA with improved thermostability. Cell selection, growth, induced expression of CA variant enzymes and collection of cell pellets were as described below.

Picking:

Recombinant *E. coli* colonies carrying a gene encoding CA were picked using a Q-Bot® robotic colony picker (Genetix USA, Inc., Boston, Mass.) into 96-well shallow well microtiter plates containing in each well 180 μ L LB Broth, 1% glucose and 30 μ g/mL chloramphenicol (CAM). Cells were grown overnight at 37° C. with shaking at 200 rpm. A 10 μ L aliquot of this culture was then transferred into 96-deep well plates containing 390 μ L TB broth and 30 μ g/mL CAM. After incubation of the deep-well plates at 37° C. with shaking at 250 rpm for 2-3 hrs, recombinant gene expression within the cultured cells was induced by addition of IPTG to a final concentration of 1 mM, followed by addition of ZnSO₄ to a final concentration of 0.5 mM. The plates were then incubated at 37° C. with shaking at 250 rpm for 18 hrs.

Preparation of Clear Lysate for Assay:

Cells were pelleted by centrifugation (4000 RPM, 10 min, 4° C.), resuspended in 200 μ L lysis buffer and lysed by shaking at room temperature for 2 hours. The lysis buffer contained 25 mM HEPES buffer, pH 8, 1 mg/mL lysozyme, and 500 μ g/mL polymixin B sulfate (PMBS) and 1 mM dithiothreitol (DTT). After sealing the plates with aluminum/polypropylene laminate heat seal tape (Velocity 11, Menlo Park, Calif., Cat#06643-001), they were shaken vigorously for 2 hours at room temperature. Cell debris was pelleted by centrifugation (4000 RPM, 10 min., 4° C.) and the clear supernatant assayed directly or stored at 4° C. until use.

High-Throughput Screening for Improved Stability in Amine Solvent, MDEA:

Screening of the polypeptides encoded by the variant genes for carbonic anhydrases with improved stability in high concentrations of an amine solvent, MDEA, was carried out using the assays as follows. After lysis, 25 μ L of cleared *E. coli* lysate was added to 96-well Costar® shallow round bottom plate, followed by addition of 75 μ L of amine solvent challenge buffer (4 M MDEA, pH 10; pH adjusted using CO₂ gas) using a Biomek NXp robotic instrument (Beckman Coulter, Fullerton, Calif.). The resulting challenge solution MDEA solvent concentration was 3 M. Challenge buffers with increased MDEA concentrations of 5.33 M and 6.66 M were used to generate 4 M and 5 M MDEA challenge solutions, which also were similarly adjusted to pH 10 with CO₂. The plates were heat-sealed with aluminum/polypropylene laminate heat seal tape (Velocity 11, Menlo Park, Calif., Cat#06643-001) at 175° C. for 2.5 seconds. The challenge reactions were heated for 24 h at the challenge temperature (42° C., 50° C., or 55° C.). Control reactions were maintained at 25° C. for 24 h. After 24 h, the plates were centrifuged at 4° C. for 10 min to clarify the reaction mixtures. Carbonic anhydrase activity after challenge was measured using a bicarbonate dehydration assay as follows: 10 μ L of cleared reaction mixture was added to a 96-well NUNC™ polystyrene shallow flat bottom plate containing 190 μ L of a solution of 0.3 M MDEA, pH 8 (pH adjusted with CO₂ gas), 200 mM KHCO₃,

400 μ M phenolphthalein. The rate of the dehydration reaction was determined as the slope of absorbance change at 25° C. (or 45° C.) assay solution temperature monitored at 550 nm (phenolphthalein as indicator) over time (30 minutes) on a SpectraMax M2 reader (Molecular Devices, Sunnyvale, Calif.). Engineered carbonic anhydrase samples showing greater than 1.3-fold improvement in activity relative to the wild-type polypeptide of SEQ ID NO: 2 under the same challenge conditions (positive control) were retested in triplicate using the same conditions. As noted in Tables 2A, 2B, 2C, 2D, 2G, 2I, and 2J. HTP screening of engineered carbonic anhydrase polypeptides for amine solvent stability and thermostability has been carried out using at least seven different challenge conditions/assays. Assay 1: challenge for 24 h at 42° C. in 3 M MDEA solution followed by dehydration activity assay at 25° C.; Assay 2: challenge for 24 h at 50° C. in 3 M MDEA solution followed by dehydration activity assay at 25° C.; Assay 5: challenge for 24 h at 50° C. in 4 M MDEA solution followed by dehydration activity assay at 45° C.; Assay 6: challenge for 24 h at 50° C. in 5M MDEA solution followed by dehydration activity assay at 25° C.; Assay 7: challenge for 24 h at 55° C. in 5 M MDEA solution followed by dehydration activity assay at 25° C.; Assay 8: challenge for 24 h at 65° C. in 5 M MDEA solution followed by dehydration activity assay in 1 M MDEA, pH 8.0 at 45° C.; Assay 12: challenge for 24 h at 70° C. in 5 M MDEA solution followed by dehydration activity assay in 0.5 M MDEA at 45° C.; Assay 14: challenge for 24 h at 82.5° C. in 4.2 M MDEA solution followed by dehydration activity assay in 960 mM MDEA at 45° C.; Assay 15: challenge for 24 h at 85° C. in 4.2 M MDEA solution followed by dehydration activity assay in 960 mM MDEA at 45° C.; Assay 16: challenge for 24 h at 90° C. in 4.2 M MDEA solution followed by dehydration activity assay in 960 mM MDEA at 45° C.; and Assay 17: challenge for 24 h at 87° C. in 4.2 M MDEA solution followed by dehydration activity assay in 685 mM MDEA at 45° C. More stringent challenge conditions having higher amine solvent concentrations, and/or temperature, and/or additional reaction components (e.g., potential inhibiting impurities found in flue gas such as NO_x and SO_x compounds) are contemplated for screening further engineered carbonic anhydrase polypeptides having higher levels of stability and/or tolerance to the challenge conditions. High-throughput screening for improved stability in ammonia solvent: Screening of the engineered carbonic anhydrase polypeptides for improved stability in high concentrations of ammonia, was carried out using essentially the same HTP assay as for MDEA amine solvent described above but with the following changes. After lysis, 25 μ L of lysate was transferred into 96-well Costar® shallow round bottom plates containing 75 μ L of ammonia challenge buffer (5.6 M NH₃ (10 wt %) loaded with 0.3 molar equivalents of CO₂ gas). The resulting challenge solution ammonia concentration was 4.2 M (7.5 wt %). The challenge solutions were heated for 24 h at the challenge temperature (30° C. or 35° C.). Control solutions were maintained at 25° C. for 24 h. After 24 h under challenge conditions, carbonic anhydrase activity was measured using a bicarbonate dehydration assay as follows: 10 μ L of challenge (or control) solution was transferred to 190 μ L of buffer (100 mM HEPES buffer, pH 7; 200 mM KHCO₃, 400 μ M phenolphthalein). The rate of the dehydration reaction was determined as the slope of absorbance change at 25° C. assay solution temperature monitored at 550 nm (phenolphthalein is a color indicator) over time (20 minutes). Engineered carbonic anhydrase samples showing greater than 1.3-fold improvement in activity relative to the wild-type polypeptide of SEQ ID NO: 2 under the same

challenge conditions (positive control) were retested in triplicate using the same conditions.

As noted in Tables 2A, 2E, 2F, and 2H, HTP screening of engineered carbonic anhydrase polypeptides for ammonia solvent stability and thermostability has been carried out using at least six different challenge conditions/assays. Assay 3: challenge for 24 h at 30° C. in 4.2 M NH₃ solution containing 0.3 molar equivalents of CO₂ ($\alpha=0.3$), followed by dehydration activity assay at 25° C.; Assay 4: challenge for 24 h at 35° C. in 4.2 M NH₃ solution containing 0.3 molar equivalents of CO₂ ($\alpha=0.3$), followed by dehydration activity assay at 25° C.; Assay 9: challenge for 24 h at 44° C. in 5.6 M NH₃ solution containing 0.3 molar equivalents of CO₂ ($\alpha=0.3$), followed by dehydration activity assay in 0.28 M NH₃ at 25° C.; Assay 10: challenge for 24 h at 25° C. in 5.6 M NH₃ solution containing 0.3 molar equivalents of CO₂ ($\alpha=0.3$), followed by dehydration activity assay in 0.28 M NH₃ at 25° C.; Assay 11: challenge for 24 h at 58° C. in 8.4 M NH₃ solution containing 0.3 molar equivalents of CO₂ ($\alpha=0.3$), followed by dehydration activity assay in 1.37 M NH₃ at 25° C.; and Assay 13: challenge for 24 h at 70° C. in 8.4 M NH₃ solution containing 0.3 molar equivalents of CO₂ ($\alpha=0.3$), followed by dehydration activity assay at 25° C. More stringent challenge conditions having higher ammonia solvent concentrations, and/or higher or lower temperatures, and/or additional reaction components (e.g., potential inhibiting impurities found in flue gas such as NO_x and SO_x compounds) are contemplated for screening further engineered carbonic anhydrase polypeptides having higher levels of stability and/or tolerance to the challenge conditions.

Production of Recombinant Carbonic Anhydrase Shake-Flask Powder (SFP):

A shake-flask procedure was used to generate recombinant carbonic anhydrase polypeptide powders used in secondary screening assays or in the carbon capture processes disclosed herein. Shake flask powder (SFP) includes approximately 30% total protein and accordingly provide a more purified preparation of an engineered enzyme as compared to the cell lysate. A single microbial colony of *E. coli* containing a plasmid encoding a CA of interest was inoculated into 50 mL Luria Bertani broth containing 30 µg/mL chloramphenicol and 1% glucose. Cells were grown overnight (at least 16 hrs) in an incubator at 30° C. with shaking at 250 rpm. The culture was diluted into 250 mL 2XYT media containing 30 µg/mL chloramphenicol, in a 1 liter flask to an optical density at 600 nm (OD₆₀₀) of 0.2 and allowed to grow at 30° C. Expression of the CA gene was induced by addition of isopropyl β-D-thiogalactoside (IPTG) to a final concentration of 1 mM when the OD₆₀₀ of the culture was 0.6 to 0.8. ZnSO₄ was then added to a final concentration of 0.5 mM and incubation was then continued overnight (at least 16 hrs). Cells were harvested by centrifugation (5000 rpm, 15 min, 4° C.) and the supernatant discarded. The cell pellet was resuspended with an equal volume of cold (4° C.) 25 mM HEPES buffer, pH 8, and passed through a homogenizer twice at 33.6 kpsi while maintained at 4° C. Cell debris was removed by centrifugation (9000 rpm, 45 min., 4° C.). The clear lysate supernatant was collected and stored at -20° C. Lyophilization of frozen clear lysate provides a dry powder (shake flask powder) of recombinant carbonic anhydrase polypeptide.

Production of Recombinant Carbonic Anhydrase Downstream-Processed (DSP) Powder:

DSP powders contains approximately 80% total protein and accordingly provide a more purified preparation of the engineered carbonic anhydrase as compared to the cell lysate. Larger-scale (~100-120 g) fermentation of the engineered carbonic anhydrase for production of DSP powders can be

carried out as a short batch followed by a fed batch process according to standard bioprocess methods.

A single microbial colony of *E. coli* containing a plasmid with the recombinant carbonic anhydrase gene of interest was inoculated into 2 mL M9YE broth containing 30 µg/mL chloramphenicol and 1% glucose. Cells were grown overnight (at least 12 h) in an incubator at 37° C. with shaking at 250 rpm. After overnight growth, 0.5 mL of this culture was diluted into 250 mL M9YE Broth containing 30 µg/mL chloramphenicol and 1% glucose in 1 liter flask and allowed to grow at 37° C. with shaking at 250 rpm. When the OD₆₀₀ of the culture is 0.5 to 1.0, the cells were removed from the incubator and either used immediately, or stored at 4° C.

Bench-scale fermentations were carried out at 30° C. in an aerated, agitated 15 L fermentor using 6.0 L of growth medium consisting of: 0.88 g/L ammonium sulfate, 0.98 g/L of sodium citrate; 12.5 g/L of dipotassium hydrogen phosphate trihydrate, 6.25 g/L of potassium dihydrogen phosphate, 3.3 g/L of Tastone-154 yeast extract, 0.083 g/L ferric ammonium citrate, and 8.3 mL/L of a trace element solution containing 2 g/L of calcium chloride dihydrate, 2.2 g/L of zinc sulfate heptahydrate, 0.5 g/L manganese sulfate monohydrate, 1 g/L cuprous sulfate heptahydrate, 0.1 g/L ammonium molybdate tetrahydrate and 0.02 g/L sodium tetraborate. The vessel was sterilized at 121° C. and 15 PSI for 30 minutes, and ZnSO₄ was added to 0.5 mM post sterilization. The fermentor was inoculated with a late exponential culture of *E. coli* W3110 containing a plasmid encoding the CA gene of interest (grown in a shake flask as described above to a starting OD₆₀₀ of 0.5 to 1.0). The fermentor was agitated at 250-1250 rpm and air was supplied to the fermentation vessel at 0.6-25 L/min to maintain a dissolved oxygen level of 50% saturation or greater. The pH of the culture was maintained at 7.0 by addition of 20% v/v ammonium hydroxide. Growth of the culture was maintained by addition of a feed solution containing 500 g/L Cerelease dextrose, 12 g/L ammonium chloride and 5.1 g/L magnesium sulfate heptahydrate. After the culture reached an OD₆₀₀ of 70±10, expression of CA was induced by addition of isopropyl-β-D-thiogalactoside (IPTG) to a final concentration of 1 mM and fermentation is continued for another 18 hours. The culture was then chilled to 4° C. and maintained at that temperature until harvested. Cells were collected by centrifugation at 5000 G for 40 minutes in a Sorval RC12BP centrifuge at 4° C. Harvested cells were used directly in the following downstream recovery process or they may be stored at 4° C. or frozen at -80° C. until such use.

The cell pellet was resuspended in 2 volumes of 25 mM triethanolamine (sulfate) buffer, pH 7.5 at 4° C. to each volume of wet cell paste. The intracellular CA was released from the cells by passing the suspension through a homogenizer fitted with a two-stage homogenizing valve assembly using a pressure of 12000 psig. The cell homogenate was cooled to -20° C. immediately after disruption. A solution of 11% w/v polyethyleneimine pH 7.2 was added to the lysate to a final concentration of 0.5% w/v. A solution of 1 M Na₂SO₄ was added to the lysate to a final concentration of 100 mM. The lysate was then stirred for 30 minutes. The resulting suspension was clarified by centrifugation at 5000G in a Sorval RC12BP centrifuge at 4° C. for 30 minutes. The clear supernatant was decanted and concentrated ten-fold using a cellulose ultrafiltration membrane with a molecular weight cut off of 10 kD. The final concentrate was dispensed into shallow containers, frozen at -20° C. and lyophilized to provide the DSP powder. The recombinant carbonic anhydrase DSP powder was stored at -80° C.

Example 2

Acceleration of CO₂ Absorption by the Carbonic Anhydrase from *Desulfovibrio vulgaris* (SEQ ID NO: 2) in Presence of Various Amine Compounds and Carbonate Ions and Elevated Temperatures

This example illustrates the ability of the wild-type beta-class carbonic anhydrase from *Desulfovibrio vulgaris* (SEQ ID NO: 2) and the engineered carbonic anhydrase polypeptides identified from HTP screening to accelerate the absorption of CO₂ gas into solutions containing high concentrations of various amine compounds (e.g., MDEA), or Na₂CO₃, as well as the amine compound MDEA at various elevated temperatures.

Stirred Cell Reactor Apparatus:

A stirred cell reactor (SCR) was used to measure the acceleration of CO₂ absorption rate in the presence of carbonic anhydrase polypeptides of the present disclosure. The SCR consists of a hermetically-sealed cylindrical reactor vessel in which a gas and a liquid phase are mixed while their interface remains flat resulting in a mass transfer rate that is well known. The SCR allows the gas pressure and the gas and liquid temperatures to be controlled and monitored over time.

SCR Assay Method:

Carbonic anhydrase polypeptide shake-flask powder (DSP can also be used) and the CO₂ capture solution of interest (e.g., 4.2 M MDEA) are added to the reactor vessel. In some assays, the CO₂ capture solution is pre-loaded with a specific mole ratio of CO₂ per amine compound or ammonia defined by the term α . Pre-loading of a solution with CO₂ is carried out by first adding unloaded capture solution to the vessel, pressurizing the vessel with pure CO₂ gas and mixing the solution until the CO₂ pressure drops to a certain level. The difference between the highest pressure and lowest pressure is used (with the ideal gas law) to calculate α of the solution.

Following addition of enzyme and solution to the vessel, the pressure in the SCR is reduced until the boiling point is reached, and the system is allowed to equilibrate until the pressure and temperature no longer change. A reservoir containing CO₂ (pure or a mixture) is connected to the SCR and a connecting valve is opened briefly allowing CO₂ to enter the SCR. Typically, the valve is opened until there is a change in pressure of approximately 10 psi when pure CO₂ is used. After closing the connecting valve, the drop in pressure in the SCR, which corresponds to the capture of CO₂ in solution, is monitored over time along with the gas and liquid temperatures. A control assay without the enzyme is also carried out.

Calculation of Rate Acceleration:

The slope of the logarithm of the pressure drop in the SCR over time is used to calculate the overall pseudo-first order rate constant (k_{OV}) according to Eq. 1.

$$\text{slope} = \frac{\Delta \ln P_{CO_2}}{\Delta t} = - \frac{RT_G A}{V_G H e_{CO_2}} \sqrt{k_{OV} D_{CO_2}} \quad (\text{Eq. 1})$$

From k_{OV} , the second order rate constant, k_2 , can then be calculated according to Eq. 2.

$$r_{CO_2} = k_{OV}[CO_2], k_{OV} = k_{1,Base} + k_{1,CA} = k_{2,Base}[Base] + k_{2,CA}[CA] \quad (\text{Eq. 2})$$

The acceleration provided by a carbonic anhydrase polypeptide, or $E_{Cat, X g/L}$, is calculated by dividing the rate,

k_{OV} measured with a specified amount (X g/L) of the carbonic anhydrase by the rate, k_{OV} measured without enzyme, according to Eq. 3.

$$\text{Acceleration} = E_{Cat, X g/L} = \frac{k_{OV \text{ with } X g/L \text{ carbonic anhydrase}}}{k_{OV \text{ without carbonic anhydrase}}} \quad (\text{Eq. 3})$$

Certain equations and physical constants are used in calculating k_{OV} . For Eq. 1 and Eq. 2 to be valid, the reaction must be operated in the pseudo first order regime, which requires the following conditions: Hatta number ("Ha") > 2, and $E_{\infty}/Ha > 5$ (E_{∞} =infinite enhancement factor). The Hatta number, Ha, and infinite enhancement factor, E_{∞} , are determined according to Eq. 4 and Eq. 5, respectively.

$$Ha = \frac{\sqrt{k_{OV} \cdot D_{CO_2}}}{k_L} \quad (\text{Eq. 4})$$

$$E_{\infty} = \sqrt{\frac{D_{CO_2}}{D_{Base}}} + \sqrt{\frac{D_{Base}}{D_{CO_2}}} \cdot \frac{[Base] \cdot H_{CO_2}}{Z_{CO_2} \cdot P_{CO_2}} \quad (\text{Eq. 5})$$

The physical constants used for SCR assays in solutions containing MDEA are summarized in Table 4.

TABLE 4

Gas volume	325 mL
Liquid volume	175 mL
Interfacial area	$3.03 \times 10^{-3} \text{ m}^2$
Gas temperature	The average gas temperature during the part of the experiment where the slope is taken.
Liquid temperature	The average liquid temperature during the part of the experiment where the slope is taken.
Vapor pressure (P_{vap})	Taken from the average of the first 10 pressure readings before the CO ₂ valve is opened. Alternatively it can be calculated from: $133.3 * \text{EXP}(20.386 - 5130/T(K))$ assuming water is the only compound giving a vapor pressure.
Liquid side mass transfer coefficient (k_L)	determined experimentally to be $4.47 \times 10^{-5} \text{ m/s}$ (see e.g., Versteeg et al, Chem. Eng. Sci., 1987, 42, 1103-1119 for procedure).
Diffusivity of CO ₂ (D_{CO_2})	Calculated as a function of liquid temperature and mass fraction of MDEA by the correlation given in Sandall et al, J. Chem. Eng. data 1989, 34, 385-391.
Diffusivity of MDEA (D_{MDEA})	Calculated as a function of liquid temperature by the correlation given in Snijder et al., J. Chem. And Engi. Data, 1993, 38, 475-480.
Henry constant of CO ₂ (H_{CO_2})	Calculated as a function of liquid temperature and mass fraction of MDEA by the correlation given in Sandall et al, J. Chem. Eng. data 1989, 34, 385-391.
Stoichiometric coefficient of CO ₂ (Z_{CO_2})	1 for the MDEA system.

Results

As shown in Table 5, a loading of 1 g/L shake flask powder of the naturally occurring beta class carbonic anhydrase polypeptide of SEQ ID NO: 2 was capable of accelerating the absorption of CO₂ by solutions containing a range of amine solvents with no pre-loading of CO₂ ($\alpha=0$) at concentration ranges from 1 M up to 4.2 M. The observed amount of acceleration was greatest in the 1 M solutions and generally decreased with increasing amine concentration. However, even in 4.2 M MDEA, the acceleration relative to the rate without enzyme was 15.8.

85

TABLE 5

[Amine] (M)	Acceleration of CO ₂ absorption Acceleration ($k_{OR, cat@1 g/L} / k_{OR, no cat}$)			
	MDEA	AMP	TEA	TIA
1	52.3	4.15	49.8	95.1
2	27.1	1.54	43.3	85.1
3	22.4	1.23	14.5	16.1
4.2	15.8			

MDEA—Methyldiethanolamine
AMP—2-amino-2-methyl-1-propanol
TEA—Triethanolamine
TIA—Trisopropanolamine

As shown in Table 6, a loading of 1 g/L shake-flask powder of the naturally occurring carbonic anhydrase polypeptide of SEQ ID NO: 2 was capable of accelerating the absorption of CO₂ by a solution at 25° C. containing 1 M Na₂CO₃ (with no pre-loading of CO₂). The initial level of acceleration was 142-fold increased relative to the control solution without the biocatalyst. The enzyme maintained a high level of acceleration at least 65-fold increased relative to no biocatalyst even after 7 days in the solution at 25° C.

TABLE 6

1 g/L SEQ ID NO: 2, 1M Na ₂ CO ₃ , 25° C. (no CO ₂ pre-loaded)	
Time (h)	Acceleration
0	142
19.95	119
45.64	110
95	71
168	65

Further SCR assays of the naturally occurring carbonic anhydrase polypeptide of SEQ ID NO: 2 were carried out at 40° C. in solutions pre-loaded with CO₂ ($\alpha=0.1$) and containing 0.5 g/L of the polypeptide and 2.0 to 4.2 M MDEA. The assay solutions were monitored for up to 49 h. As shown by the results listed in Table 7, only 0.5 g/L of the polypeptide of SEQ ID NO: 2 was capable of initially accelerating the absorption of CO₂ in solutions at 40° C. containing 2 M to 4.2 M MDEA from about 11-fold to about 3-fold relative to the control solution without biocatalyst added. Further even after 16 h or more in the 2 M to 4.2 M MDEA solutions at 40° C., the polypeptide of SEQ ID NO: 2 was capable of still accelerating CO₂ absorption by at least 2-fold relative to the control solution.

TABLE 7

Assay Sample	Time (h)	k_{OR} (s ⁻¹)	Acceleration
2M MDEA (no enzyme)	0	10.8	1
2M MDEA + 0.5 g/L enzyme	0	116.9	10.9
2M MDEA + 0.5 g/L enzyme	20.4	30.2	2.8
2M MDEA + 0.5 g/L enzyme	49.2	14.5	1.3
2.5M MDEA (no enzyme)	0	12.6	1
2.5M MDEA + 0.5 g/L enzyme	0	112.6	9.0
2.5M MDEA + 0.5 g/L enzyme	21.1	38.4	3.1
2.5M MDEA + 0.5 g/L enzyme	48.25	22.2	1.8
3M MDEA (no enzyme)	0	15.0	1
3M MDEA + 0.5 g/L enzyme	0	103.5	6.9
3M MDEA + 0.5 g/L enzyme	19.92	37.5	2.5
3M MDEA + 0.5 g/L enzyme	44.33	20.3	1.4
4.2M MDEA (no enzyme)	0.0	12.4	1
4.2M MDEA + 0.5 g/L enzyme	0.0	40.7	3.3

86

TABLE 7-continued

Assay Sample	Time (h)	k_{OR} (s ⁻¹)	Acceleration
4.2M MDEA + 0.5 g/L enzyme	16.5	28.2	2.3
4.2M MDEA + 0.5 g/L enzyme	47.0	16.6	1.3

Example 3

Acceleration of CO₂ Absorption by Engineered Carbonic Anhydrase Polypeptides in the Presence of MDEA in Solution at Elevated Temperatures

This example illustrates the ability of engineered carbonic anhydrase polypeptides identified from HTP screening to accelerate the absorption of CO₂ gas into amine solvent (MDEA) solutions at elevated temperatures.

Assays measuring rate of CO₂ hydration catalyzed by engineered carbonic anhydrase polypeptides of SEQ ID NO: 6, 16, 26, 30, 42, 84, and 186 (and wild-type of SEQ ID NO: 2) in MDEA solvent at 40° C. and 50° C. were carried out using the SCR and methods as described above in Example 2. As shown in Table 8, the relative improvement in stability in MDEA solvent exhibited by the polypeptides was determined as fold-improvement in residual activity at various time points and also as half-life ($t_{1/2}$) of CO₂ hydration activity.

TABLE 8

Poly-peptide SEQ ID NO:	Fold Improvement relative to SEQ ID NO: 2					
	$t_{1/2}$ (h)	$t_{1/2}$	Activity at 0 h	Residual Activity at 24 h	Residual Activity at 48 h	Residual Activity at 72 h
0.5 g/L polypeptide, 3M MDEA, T = 40° C., $\alpha = 0.1$						
2	20.0	1	1	1	1	1
6	67.8	3.4	2.18	2.2	5.0	4.8
16	69.9	3.5	1.60	2.6	5.4	4.9
30	30.7	1.5	1.45	1.7	2.6	1.8
42	53.0	2.7	1.78	2.2	4.2	4.1
84	18.2	0.9	1.74	1.0	1.2	0.7
186	24.7	1.2	1.68	1.5	2.1	1.5
0.5 g/L polypeptide, 3M MDEA, T = 40° C., $\alpha = 0.02$						
2	32.0	1	1	1	1	1
6	132.1	4.1	1.20	2.0	2.7	2.8
16	153.7	4.8	1.02	2.0	2.7	2.6
30	82.8	2.6	1.03	1.8	2.7	2.6
42	88.6	2.8	1.13	1.4	2.2	2.3
84	27.6	0.9	1.10	0.9	1.0	0.8
186	45.5	1.4	1.06	1.2	1.6	1.1
0.5 g/L polypeptide, 3M MDEA, T = 50° C., $\alpha = 0.02$						
2	0.13	1.0				
16	9.0	71.0				
26	9.8	77.0				

The engineered carbonic anhydrase polypeptides of SEQ ID NO: 6, 16, 30, 42, and 186 exhibited 1.2-fold to 4.8-fold increased stability in a 3 M MDEA solution at the elevated temperature of 40° C. (with a CO₂ loading $\alpha=0.02$ or 0.1) when measured as $t_{1/2}$ for CO₂ absorption activity relative to the wild-type polypeptide of SEQ ID NO: 2. At the further elevated temperature of 50° C. (with a CO₂ loading of $\alpha=0.02$) the engineered polypeptides of SEQ ID NO: 16 and 26 exhibited over 70-fold increased $t_{1/2}$ for CO₂ absorption activity in a 3 M MDEA solution relative to the wild type polypeptide of SEQ ID NO: 2.

The engineered carbonic anhydrase polypeptides of SEQ ID NO: 6, 16, 30, 42, and 186, continued to maintain their improved stability even at 48 h at 40° C. Additionally, in the case of the assays at the higher CO₂ loadings ($\alpha=0.1$), the stability increased significantly relative to that of the wild-type polypeptide of SEQ ID NO: 2—e.g., for SEQ ID NO: 16 increased from 1.6-fold to 5.4-fold greater than SEQ ID NO: 2.

Example 4

Acceleration of CO₂ Absorption by the Carbonic Anhydrase from *Desulfovibrio Vulgaris* (SEQ ID NO: 2) in Presence of Ammonia in Solution at Chilled Temperatures

This example illustrates the ability of the beta-class carbonic anhydrase from *Desulfovibrio vulgaris* (SEQ ID NO: 2) to accelerate the absorption of CO₂ gas into a chilled ammonia solution.

Apparatus and Assay Method

To a Parr Series 5100 low pressure reactor system fitted with a mass flow meter, a digital pressure gauge, a septum-capped addition/sampling port, a thermal well, a cooling loop (used as baffles/agitator shaft support) and a 450 mL glass jacketed cylinder was added water and the water degassed via vacuum at room temperature for ~20-40 minutes (until no bubble formation was observed). The cylinder was detached under a gentle nitrogen flow and 30 wt % NH₃ solution was added to make up the desired NH₃ solution with a final volume of ~250 mL (e.g., 250 mL of 10 wt % NH₃ solution=166 mL of water and 83 mL of 30 wt % NH₃). The 450 mL glass jacketed cylinder with the NH₃ solution was reattached to the reactor under a nitrogen atmosphere and the internal temperature was adjusted to the desired level via an external heat exchanger/circulator.

The turbine propeller was positioned on the stirrer shaft such that it was slightly above the liquid level and was used to mix the gas phase. An egg-shaped stir bar was placed in the cylinder and was used to stir the liquid phase via an external stir plate situated underneath the cylinder. Typically, the gas phase was stirred at 1800-2000 rpm and the liquid phase was stirred at 900-1200 rpm (fastest rate such that the surface of the liquid remained relatively flat/ripple-free). The internal temperature of the gas phase, the internal temperature of the liquid phase, the internal gas phase pressure, the agitation rates and the jacket temperature were recorded via a data logger.

After the internal temperature and pressure had equilibrated/stabilized, CO₂ gas was introduced through the mass flow meter until the desired initial loading of CO₂ was obtained. Loading was denoted as “ α ” which corresponds to the mole ratio of CO₂ to NH₃ (e.g., $\alpha=0.3$ means 3 moles of CO₂ per 10 moles of NH₃). Generally, depending on process optimization in an industrial scale process for CO₂ capture using chilled ammonia solution it is contemplated that the solution will enter the flue gas absorber at a relatively “lean” loading, of about $\alpha=0.1$ -0.3 and after absorbing CO₂ will leave the absorber at a “rich” loading, dependent on equilibrium, of about $\alpha=0.5$ -0.7.

Biocatalyst was introduced as an aqueous solution through the addition port. For control reaction, no additional solution was introduced. Then, for both sample and control reactions, a quick burst of CO₂ was added to the reactor vessel such that the partial pressure of CO₂ in the reactor was 5-15 psig. The vessel then was sealed. The subsequent decrease in the partial pressure of CO₂ in the reactor over time was recorded. The

kinetic parameters were determined via analysis of the pressure versus time data under the prescribed reactor conditions. The composition of the solution in the reactor could also be monitored via samplings through the addition port. The acceleration in the rate of CO₂ absorption was calculated as described in Example 2.

Results

A set of assays were carried out at 10° C. in a solution containing 5.6 M NH₃ with and without 2 g/L of the naturally occurring beta class carbonic anhydrase of SEQ ID NO: 2, with the CO₂ loading of the solution varied from $\alpha=0.30$ to $\alpha=0.62$.

As shown in Table 9, the observed rate constants, k_{ov} , with and without enzyme decreased with increased CO₂ loading in the solution (i.e., increasing α), but k_{ov} increased as the CO₂ partial pressure in the gas phase decreased.

TABLE 9

Sample Loading	k_{ov} (s ⁻¹)			
	CO ₂ partial pressure drop (atm)			
	0.2 → 0.15	0.15 → 0.10	0.10 → 0.05	0.05 → 0.02
$\alpha = 0.30 + \text{enzyme}$	326	414	617	707
$\alpha = 0.30 \text{ control}$	110	137	179	193
$\alpha = 0.36 + \text{enzyme}$	165	223	342	537
$\alpha = 0.36 \text{ control}$	33.0	43.3	58.8	89.8
$\alpha = 0.41 + \text{enzyme}$	152	210	336	517
$\alpha = 0.41 \text{ control}$	14.4	17.4	23.6	32.4
$\alpha = 0.47 + \text{enzyme}$	95	131	223	432
$\alpha = 0.47 \text{ control}$	5.3	7.4	13.6	33.5
$\alpha = 0.53 + \text{enzyme}$	48	58	96	201
$\alpha = 0.53 \text{ control}$	3.1	3.5	3.5	2.5
$\alpha = 0.62 + \text{enzyme}$	61	64	73	72
$\alpha = 0.62 \text{ control}$	3.8	4.4	6.1	11.3

As shown in Table 10, the naturally occurring carbonic anhydrase polypeptide of SEQ ID NO: 2 exhibited significant CO₂ absorption acceleration in the chilled ammonia solution (5.6 M NH₃ at 10° C.).

TABLE 10

CO ₂ Loading (α)	Enzyme acceleration of CO ₂ absorption
0.30	3.0
0.36	5.5
0.41	13
0.47	17
0.53	22.5

The amount of acceleration by the presence of the polypeptide of SEQ ID NO: 2 increased linearly from a value of about 3.0, at a loading of $\alpha=0.30$, up to about 22.5, at a loading of $\alpha=0.53$. Above $\alpha=0.53$ the pseudo first order behavior of k_{ov} appeared to break down and the rate of acceleration could not be determined accurately.

Further assays were carried out at 10° C. in a solution containing 5.6 M NH₃, a solution CO₂ loading of the solution of $\alpha=0.30$ -0.40 and 2 g/L of a recombinant carbonic anhydrase from Table 2A. The recombinant carbonic anhydrases polypeptides had amino acid sequences of SEQ ID NO: 6, 26, 32, 60, and 124, and included the following amino acid residue differences relative to SEQ ID NO 2: X15R, X30R; X56S, X86A, and X119K. All of the assayed recombinant carbonic anhydrases polypeptides accelerated the CO₂ absorption by the 5.6 M NH₃ solution at 10° C. equivalent to the acceleration exhibited by wild-type of SEQ ID NO: 2. In contrast, the wild-type carbonic anhydrases of SEQ ID NO:

1174, 1176, and 1178, each of which has some amino acid sequence homology to SEQ ID NO: 2 exhibited no observable acceleration over baseline of the CO₂ absorption by the 5.6 M NH₃ solution at 10° C. Thus, wild-type carbonic anhydrase polypeptide from *D. vulgaris* of SEQ ID NO: 2, or one of the engineered carbonic anhydrase polypeptides comprising one or more of the amino acid differences X15R, X30R; X56S, and X119K, is capable of significantly accelerating carbon dioxide absorption by a solution under “chilled ammonia” process conditions of 5.6 M NH₃, α=0.3-0.4, 2 g/L polypeptide, and T=10° C.

Example 5

Acceleration of CO₂ Absorption by Engineered Carbonic Anhydrase Polypeptides in the Presence of MDEA in Solution at Elevated Temperatures

This example illustrates the ability of engineered carbonic anhydrase polypeptides identified from HTP screening to accelerate the absorption of CO₂ gas into amine solvent (MDEA) solutions at elevated temperatures.

Assays measuring rate of CO₂ hydration catalyzed by the engineered carbonic anhydrase polypeptides of SEQ ID NO: 26, 190, 206, 238, 252, 270, 274, 284, 306, 318, 328, 332, 340, 354, 596, 606, 656, 678, 1080, 1110, 1148, 1152, 1156, and 1158, in increasingly challenging conditions of MDEA solvent concentration and temperature, were carried out using the SCR and methods as described above in Example 2.

As shown in Tables 11-15, the relative improvement in stability in MDEA solvent exhibited by the polypeptides measured as half-life (t_{1/2}) of CO₂ hydration activity was determined as well as the fold-improvement in residual activity relative to a parent engineered polypeptide. For example, as shown in Table 15, the engineered carbonic anhydrase of SEQ ID NO: 1152 (which has the following residue differences relative to SEQ ID NO: 2: T30R; R31P; K37R; A40L; Q43M; A56S; E68A; V70I; A84Q; A95V; Q119M; G120R; H124R; T139M; N145F; H148T; V157A; M170F; N213E; and A219T) exhibited a 16-fold improvement in t_{1/2} over its parent engineered polypeptide of SEQ ID NO: 656 (which has the following residue differences relative to SEQ ID NO: 2: T30R; K37R; A40L; A56S; E68A; A84Q; A95V; Q119M; G120R; T139M; N145W; N213E; A219T), under the following conditions: 1.0 g/L polypeptide, 4.2 M MDEA, T=50° C. assay, 75° C. incubation, α=0.02. Similarly, as shown in Table 14, the engineered carbonic anhydrase of SEQ ID NO: 656 exhibited a 10-fold improvement in t_{1/2} over its parent engineered polypeptide of SEQ ID NO: 332 (which has the following residue differences relative to SEQ ID NO: 2: T30R, A40L, A56S, A84Q, G120R, and T139M), under the following conditions: 1.0 g/L polypeptide, 4.2 M MDEA, T=50° C. assay, 65° C. incubation, α=0.02. Hence, the results shown in this Example demonstrate the cumulative improvement for stability in the presence of an amine compound for the engineered carbonic anhydrase polypeptides through the addition of amino acid residue differences to the polypeptide sequences as disclosed herein.

TABLE 11

Polypeptide SEQ ID NO:	t _{1/2} (h)	Fold-improved (relative to SEQ ID NO: 26)
0.5 g/L polypeptide, 4M MDEA, T = 50° C., α = 0.02		
26	9.1	1.00
328	57.4	6.31

TABLE 11-continued

Polypeptide SEQ ID NO:	t _{1/2} (h)	Fold-improved (relative to SEQ ID NO: 26)
0.5 g/L polypeptide, 4M MDEA, T = 50° C., α = 0.02		
284	48.1	5.29
354	41.2	4.53
318	37.4	4.11
340	25.2	2.77
252	18.4	2.02
190	5.7	0.63
206	24.3	2.67

TABLE 12

Polypeptide SEQ ID NO:	t _{1/2} (h)	Fold-improved (relative to SEQ ID NO: 26)
1.0 g/L polypeptide, 4.2M MDEA, T = 53-55° C., α = 0.02		
26	0.59	1.0
332	10.3	17

TABLE 13

Polypeptide SEQ ID NO:	t _{1/2} (h)	Fold-improved (relative to SEQ ID NO: 26)
1.0 g/L polypeptide, 4.2M MDEA, T = 53-55° C., α = 0.02		
332	23	1.0
270	55	2.4
238	44	1.9
306	78	3.4
274	119	5.1

TABLE 14

Polypeptide SEQ ID NO:	t _{1/2} (h)	Fold-improved (relative to SEQ ID NO: 26)
1.0 g/L polypeptide, 4.2M MDEA, T = 50° C. assay, 65° C. incubation, α = 0.02		
332	2.8	1.0
656	28	10
596	21	7.6
606	81	29
678	27	9.8

TABLE 15

Polypeptide SEQ ID NO:	t _{1/2} (h)	Fold-improved (relative to SEQ ID NO: 26)
1.0 g/L polypeptide, 4.2M MDEA, T = 50° C. assay, 75° C. incubation, α = 0.02		
656	2.2	1.0
1152	35	16
1156	24	11
1110	21	9.7
1158	17	7.9
1148	12	5.5
1080	12	5.3

Example 6

Acceleration of CO₂ Absorption by Engineered Carbonic Anhydrase Polypeptides in the Presence of NO and SO_x Flue Gas Components

This example illustrates the ability of engineered carbonic anhydrase polypeptides identified from HTP screening to

91

accelerate the absorption of CO₂ gas into amine solvent (MDEA) solutions in the presence of NO_x and SO_x compounds that are typical flue gas components.

A 1 g/L solution of the engineered carbonic anhydrase polypeptide of SEQ ID NO: 332 (which has the following amino acid differences relative to SEQ ID NO: 2: T30R, A40L, A56S, A84Q, G120R, and T139M) was added to 100 mL of 4.2 M MDEA, preloaded with CO₂ at a mole ratio of $\alpha=0.02$, in the stirred cell reactor and allowed to equilibrate at 50° C. The enzyme activity was determined by pressurizing the system with pure CO₂ and measuring the rate of CO₂ pressure drop using the SCR and the overall rate constant k_{ov} was calculated as described above in Example 2. After this initial baseline assay without any NO_x or SO_x compound was performed, 1 mL of 100 g/L NaNO₃ (sodium nitrate) was added at a concentration of 1 g/L (or 1 part per thousand, ppt) NaNO₃ in the SCR. NaNO₃ at 1 ppt was used to simulate a typical NO_x compound flue gas component. The activity of the enzyme of SEQ ID NO: 332 was assayed as previously described. No loss of activity was observed due to the presence of NaNO₃. Then, 1 mL of 100 g/L NaNO₂ (sodium nitrite to simulate typical NO_x flue gas component) was added to the same solution and again assayed. Similarly, these assays were repeated sequentially with Na₂SO₃ (sodium sulfite) and Na₂SO₄ (sodium sulfate).

As shown in the Table 16 below, at no point did there appear to be a significant change in the activity of the engineered carbonic anhydrase polypeptide of SEQ ID NO: 332 after the addition of any of the salts of NO_x or SO_x. Hence, the improved activity of the engineered polypeptides of the present disclosure in accelerating the absorption of CO₂ in MDEA further exhibit resistance to inhibition by NO_x or SO_x compounds typically found as flue gas components.

TABLE 16

Sample	Rate k_{ov} (s ⁻¹)
No enzyme	44
SEQ ID NO: 332	145
SEQ ID NO: 332 + 1 ppt NaNO ₃	144
SEQ ID NO: 332 + 1 ppt NaNO ₃ + 1 ppt NaNO ₂	144
SEQ ID NO: 332 + 1 ppt NaNO ₃ + 1 ppt NaNO ₂ + 1 ppt Na ₂ SO ₃	133
SEQ ID NO: 332 + 1 ppt NaNO ₃ + 1 ppt NaNO ₂ + 1 ppt Na ₂ SO ₃ + 1 ppt Na ₂ SO ₄	141

Example 7

Acceleration of CO₂ Absorption by Engineered Carbonic Anhydrase Polypeptides in the Presence of Ammonia

This example further illustrates the ability of recombinant carbonic anhydrase polypeptides of the present disclosure to exhibit increased stability to ammonia and accelerate the absorption of CO₂ gas in solutions containing ammonia.

Uptake of CO₂ gas by solutions containing varying concentrations ammonia with and without enzyme were carried out in the stirred cell apparatus and using the assay protocol and general conditions described in Example 4.

As shown in Table 17, the recombinant carbonic anhydrase polypeptide of SEQ ID NO: 26 (which includes the amino acid difference A56S) accelerated the CO₂ gas uptake of a solution containing 8 M NH₃ at 5° C. The acceleration varied depending on the CO₂ loading (α) of the solution, ranging from about 1200 s⁻¹ at $\alpha=0.2$ loading, down to about 15 s⁻¹ at $\alpha=0.5$ loading.

92

As shown in Table 18, recombinant carbonic anhydrase polypeptides of SEQ ID NO: 32, 748, 788, 812, 962, 964, and 966, each of which have various amino acid residue differences relative to SEQ ID NO: 2, exhibit acceleration of CO₂ uptake relative to uncatalyzed solution in 10 wt % NH₃ ($\alpha=0.3$) at 10° C., even after high temperature challenge of 24 h at 44° C. or 2 h at 65° C.

TABLE 17

CO ₂ loading (α)	Acceleration ¹ [s ⁻¹]
0.20	1190.3
0.25	889.7
0.30	787.8
0.35	349.2
0.40	202.9
0.45	75.2
0.50	16.3
0.55	12.9

¹Acceleration¹ = k_{ov} , determined in stirred cell reactor using 2 g/L of polypeptide of SEQ ID NO: 26, 8M NH₃ at 5° C., over the CO₂ pressure drop range of from 0.15 atm to 0.10 atm.

TABLE 18

No challenge		After 24 h at 44° C.		After 2 h at 65° C.	
SEQ ID NO:	Acceleration ¹	SEQ ID NO:	Acceleration ¹	SEQ ID NO:	Acceleration ¹
966	3.1	966	2.3	748	1.4
32	3.0	962	2.3	962	1.3
812	3.0	964	2.0	964	1.3
962	3.0	812	1.8	966	1.2
964	2.9	788	1.7	788	1.2
788	2.7	748	1.6	812	1.1
748	1.7	32	1.4	32	1.0

¹Acceleration¹ = $k_{ov,cat}/k_{ov,uncat}$ where " $k_{ov,uncat}$ " refers to the baseline rate of CO₂ uptake without enzyme present in 10 wt % NH₃, ($\alpha=0.3$) at 10° C.

Example 8

Increased Acceleration of CO₂ Absorption by Glutaraldehyde-Treated α -Class, β -Class, and Engineered Carbonic Anhydrases in MDEA Solution

This example illustrates the preparation of chemically modified versions of the wild-type α -class human ("HuCAII") carbonic anhydrase polypeptide of SEQ ID NO: 1298, the wild-type β -class *Desulfovibrio vulgaris* carbonic anhydrase polypeptide of SEQ ID NO: 2, and the engineered β -class derived from *Desulfovibrio vulgaris* carbonic anhydrase polypeptides of SEQ ID NO: 656 and 1152 of the present disclosure, by treatment with glutaraldehyde. The example also illustrates ability of the glutaraldehyde-modified enzyme to exhibit equivalent or increased activity and stability in accelerating the absorption of CO₂ gas in a solution containing CO₂ absorption mediating compound MDEA.

Preparation of Chemically Modified Carbonic Anhydrase Polypeptides:

Shake-flask powder preparations of each of the carbonic anhydrase polypeptides were dissolved at 10 g/L concentration in 50 mM TEA-SO₄ buffer at pH 7.7, or in 50 mM Na₂CO₃ buffer at pH 10. A 25% aqueous solution of glutaraldehyde (Sigma-Aldrich Cat. #G6257; Sigma-Aldrich Corp., St. Louis, USA) was added directly to the carbonic anhydrase polypeptide solution to give the desired final glutaraldehyde concentration (e.g., 0.25% v/v). The polypeptide and cross-

linking agent solutions were mixed then allowed to incubate at room temperature without mixing for 1-4 h. The resulting solutions comprising the glutaraldehyde treated carbonic anhydrase polypeptide composition were slightly yellow in color and very slightly cloudy. Cloudiness was removed by centrifugation prior to assay.

Preparation of Chemically Modified Carbonic Anhydrase Formulations in MDEA and SCR Assay of Activity:

After incubation, 10 mL of the chemically modified carbonic anhydrase polypeptide solution was added to 90 mL of 4.66 M MDEA solution (not pre-loaded with CO₂). The resulting formulation of 1 g/L chemically modified carbonic anhydrase had a final MDEA concentration of 4.2 M. The formulation of chemically modified carbonic anhydrase polypeptide in 4.2 M MDEA was assayed for rate of CO₂ absorption at 50° C. in the SCR using the assay protocol and general conditions described in Example 2.

Briefly, the SCR assay was carried out as follows: the solution was heated until it reached 50° C.; the pressure in the SCR was reduced until the solution just started boiling, the valve to the vacuum pump was then closed; the temperature and pressure in the SCR was allowed to equilibrate; pure CO₂ was added to the SCR until it reached about 10 psia total pressure, after which the valve to the CO₂ source was closed; the pressure drop and gas and liquid temperatures were recorded; k_{OV} is calculated from the slope of $\ln P$ vs t . $k_{1,CA}$ can be calculated by subtracting k_{OV} without carbonic anhydrase from k_{OV} with carbonic anhydrase.

Results:

As shown in Table 19 below, the chemically modified carbonic anhydrase polypeptides resulting from treatment with cross-linking agent glutaraldehyde exhibited increased carbonic anhydrase activity (k_{OV}) relative to the same carbonic anhydrase polypeptides that were not treated with the cross-linking agent (i.e., "unmodified") when assayed in 4.2 M MDEA at 50° C. The human α -class carbonic anhydrase of SEQ ID NO: 1298 when chemically modified with glutaraldehyde (GA), exhibited the largest fold-improvement (14-fold) relative to its unmodified form. The *Desulfovibrio vulgaris* wild-type β -class carbonic anhydrase polypeptide of SEQ ID NO: 2, and the two engineered β -class carbonic anhydrase polypeptides of SEQ ID NO: 656 and 1152 all exhibited higher activity upon chemical modification by glutaraldehyde treatment, with significant improvements of 1.9-fold, 2.7-fold, and 5.0-fold, relative to the unmodified enzymes. Each of these β -class enzymes also exhibited greater overall activity in the assay than the α -class enzyme.

TABLE 19

Sample	Acceleration k_{ov} (s ⁻¹)	Fold-Improved (relative to unmodified)
Water	32	n/a
SEQ ID NO: 1298 (HuCAII) (unmodified)	38	n/a
SEQ ID NO: 1298 (HuCAII) + GA treatment	117	14
SEQ ID NO: 2 (unmodified)	151	n/a
SEQ ID NO: 2 + GA treatment	261	1.9
SEQ ID NO: 656 (unmodified)	208	n/a
SEQ ID NO: 656 + GA treatment	503	2.7
SEQ ID NO: 1152 (unmodified)	154	n/a
SEQ ID NO: 1152 + GA treatment	643	5.0

Example 9

Increased Stability of CO₂ Absorption Acceleration in MDEA by a Thermally Challenged Glutaraldehyde-Treated Engineered Carbonic Anhydrase

This example illustrates that a recombinant carbonic anhydrase polypeptide of the present disclosure that has been chemically modified by treatment with the cross-linking agent glutaraldehyde exhibits increased stability to thermal challenge in 4.2 M MDEA in its ability to accelerate the absorption of CO₂.

Assay for Increased Stability:

The engineered β -class carbonic anhydrase polypeptide of SEQ ID NO: 1152 was chemically modified by treatment with 0.25% glutaraldehyde and assayed in SCR to determine the k_{OV} for CO₂ uptake acceleration as described in Example 8. The heat challenge used to determine increased stability was carried out as follows. The sample was removed from the SCR and put into a bottle. The bottle was quickly heated to the stability challenge temperature of 75° C. and incubated in an oven set to this temperature. For the next assay point, the solution was quickly cooled to the 50° C. assay temperature and assayed in the SCR as above. This was repeated as necessary over a 13 day period to provide the time course for loss of stability at 75° C.

Results:

The results were plotted as time of heat challenge at 75° C. versus normalized activity (k_1) over the course of 13 days. As shown in FIG. 1, the plots both exhibited logarithmic decreases in activity over time but the rate of loss of activity in the chemically modified carbonic anhydrase was significantly decreased (indicating greater stability). The unmodified CA lost nearly 80% of its activity by the end of the first day, whereas the chemically modified CA had lost only to 25% of its activity. Based on a comparison of the line fits of the plots of all the data out to 12.5 days, the engineered β -class carbonic anhydrase polypeptide of SEQ ID NO: 1152 after chemical modification by glutaraldehyde treatment is about 4-fold more stable than the same engineered β -class carbonic anhydrase polypeptide of SEQ ID NO: 1152 that is unmodified.

Example 10

Increased Acceleration of CO₂ Absorption by a Glutaraldehyde-Treated Carbonic Anhydrase in the CO₂ Absorption Mediating Compound Solutions: AMP, MDEA, TEA, and Carbonate

This example illustrates the ability of the recombinant carbonic anhydrase polypeptides of the present disclosure that are chemically modified by treatment with glutaraldehyde to exhibit increased activity in accelerating the absorption of CO₂ by solutions containing CO₂ absorption mediating compounds other than MDEA including AMP, TEA, and carbonate ion.

The engineered carbonic anhydrase of SEQ ID NO: 1152 was chemically modified with 0.25% GA as described in Example 8. The chemically modified and unmodified polypeptides were then assayed in the SCR at 50° C. without pre-loading of CO₂ as described in Example 8, in solutions including the following concentrations of CO₂ absorption mediating compounds: 2 M AMP; 3 M TEA; and 2 M K₂CO₃.

Results:

As shown in Table 20 below. In 2 M AMP, the unmodified CA showed no activity in this solvent, in contrast, the GA modified CA did show activity and was ~4-fold higher compared to the solvent alone. In TEA, the GA modified CA was ~1.6-fold compared to TEA with unmodified CA. In 2 M K₂CO₃, no improvement was observed with the GA modified CA compared to the unmodified CA (significant precipitation was observed with the GA modified CA compared to the unmodified CA in K₂CO₃).

TABLE 20

Sample	Activity k _{ov} (s ⁻¹)	Fold-Improved (relative to unmodified)
2M AMP		
Water	1,400	
SEQ ID NO: 1152 (unmodified)	1,400	
SEQ ID NO: 1152 + GA treatment	4,200	~4
3M TEA		
Water	10	
SEQ ID NO: 1152 (unmodified)	71	
SEQ ID NO: 1152 + GA treatment	110	~1.6
2M K ₂ CO ₃		
Water	39	
SEQ ID NO: 1152 (unmodified)	110	
SEQ ID NO: 1152 + GA treatment	110	1.0
4.2M MDEA		
Water	32	
SEQ ID NO: 1152 (unmodified)	150	
SEQ ID NO: 1152 + GA treatment	640	~4.3

Example 11

Increased Acceleration of CO₂ Absorption by a
Glutaraldehyde-Treated Carbonic Anhydrase in the
CO₂ Absorption Mediating Compound Solution:
Ammonia

This example illustrates the ability of a wild-type β-class carbonic anhydrase polypeptide of that is chemically modified by treatment with glutaraldehyde to exhibit increased activity in accelerating the absorption of CO₂ by a solution containing the CO₂ absorption mediating compound, ammonia.

The wild-type *Desulfovibrio vulgaris* β-class carbonic anhydrase polypeptide of SEQ ID NO: 2 was chemically modified by treatment with 0.5% glutaraldehyde (GA) in 100 mM TEA sulfate buffer, pH 8.5 at 25 g/L CA concentration for 1-3 hrs. After the GA treatment, the solution was centrifuged to remove very slight precipitation that appeared in both GA modified and unmodified enzyme solutions. Using the stirred cell reactor (SCR) the kinetics of CO₂ absorption was measured without the enzyme present ("water"), with the carbonic anhydrase that was not chemically modified (unmodified), and with the GA-modified carbonic anhydrase. These SCR assays were performed at 22° C. in 1 M and 2 M ammonia with loading from 0.1 to 0.3. Activity was measured as pressure drop over time and calculated as the square of the

natural logarithm of the pressure drop with time, which is proportional to the pseudo-first order kinetic constant, k_{OP}.

Results:

As shown in Table 21 below, the GA modified CA exhibited 1.4 to 4.4-fold improved activity depending on the concentration and loading of CO₂ in the solution.

TABLE 21

Sample	Activity (lnAP/Δt) ² × 10 ⁴	Fold-Improved (relative to unmodified)
1M NH ₄ OH (α = 0.1)		
Water	0.952	
SEQ ID NO: 2 (unmodified)	2.21	
SEQ ID NO: 2 + GA treatment	4.55	2.9
2M NH ₄ OH (α = 0.1)		
Water	1.67	
SEQ ID NO: 2 (unmodified)	3.63	
SEQ ID NO: 2 + GA treatment	4.36	1.4
2M NH ₄ OH (α = 0.2)		
Water	2.50	
SEQ ID NO: 2 (unmodified)	3.31	
SEQ ID NO: 2 + GA treatment	6.02	4.4
2M NH ₄ OH (α = 0.3)		
Water	1.01	
SEQ ID NO: 2 (unmodified)	2.36	
SEQ ID NO: 2 + GA treatment	3.24	1.6

Example 12

Increased Acceleration of CO₂ Absorption in the
CO₂ Absorption Mediating Compound Solution
MDEA by a Recombinant Carbonic Anhydrase
Treated with Either of the Cross-linking Agents
Dimethyl Suberimidate or Dimethyl Pimelimidate

This example illustrates the ability of the recombinant carbonic anhydrase polypeptides of the present disclosure that are chemically modified by treatment with the cross-linking agents dimethyl suberimidate and dimethyl pimelimidate to exhibit increased activity in accelerating the absorption of CO₂ by a solution containing the CO₂ absorption mediating compound MDEA.

Preparation and Assay of Chemically Modified Carbonic Anhydrase Polypeptides:

Shake-flask powder preparations of the recombinant carbonic anhydrase of SEQ ID NO: 1152 were chemically modified by treatment with either of the cross-linking agents dimethyl suberimidate (Sigma-Aldrich Corp., St. Louis, USA) or dimethyl pimelimidate (Sigma-Aldrich Corp., St. Louis, USA) according to the same method used for glutaraldehyde treatment in Example 8, except that instead of glutaraldehyde either of dimethyl suberimidate or dimethyl pimelimidate was added as a solid to the polypeptide solution to give the desired final concentration of cross-linking agent. Two different treatment concentrations were used for each of dimethyl suberimidate and dimethyl pimelimidate: 0.25% and 2.5%.

Activity was determined by SCR assay of 1 g/L chemically modified enzyme in a solution 4.2 M MDEA, unloaded with CO₂ at 50° C. Stability of the chemically modified enzymes was also determined by measuring residual activity after 21 hours of incubation at 75° C. in the same assay solvent.

Results:

As shown in Table 22 below, for both the DM-suberimidate and DM-pimelimidate cross-linking agents under both treatment conditions, the chemically modified β-class carbonic

anhydrase polypeptide of SEQ ID NO: 1152 exhibited increased carbonic anhydrase activity relative to the same enzyme that was not chemically modified in an initial assay. After 21 h of incubation at 75° C. in the same assay solution, the enzymes chemically modified with dimethyl suberimidate and dimethyl pimelimidate no longer exhibited improved carbonic anhydrase activity compared to the unmodified enzyme. Thus, treatment with these cross-linking agents did not result in increased enzyme stability based on a 21 hour incubation at 75° C. This apparent lack of increased stability is not unexpected as both dimethyl suberimidate and dimethyl pimelimidate are known to result base labile cross-links that likely are cleaved during the challenge resulting in a loss of any stabilizing effect due to chemical modification that results in cross links.

TABLE 22

Sample	Initial Activity $k_{1,CA}$ (s^{-1})	Activity k_1 after challenge (21 h, 75° C.) (s^{-1})
SEQ ID NO: 1152 (unmodified)	193	113
SEQ ID NO: 1152 + 0.25% DM-Suberimidate treatment	501	131
SEQ ID NO: 1152 + 2.5% DM-Suberimidate treatment	231	134
SEQ ID NO: 1152 + 0.25% DM-Pimelimidate treatment	411	118
SEQ ID NO: 1152 + 2.5% DM-Pimelimidate treatment	303	113

Example 13

Increased Acceleration and Stability of CO₂ Absorption in the CO₂ Absorption Mediating Compound Solution MDEA by a Recombinant Carbonic Anhydrase Treated with the Base-Stable Cross-linking Agent Suberic Acid Bis(N-hydroxysuccinimide)

This example illustrates the ability of the recombinant carbonic anhydrase polypeptides of the present disclosure that are chemically modified by treatment with the base stable cross-linking agent suberic acid bis(N-hydroxysuccinimide) to exhibit increased activity and stability in accelerating the absorption of CO₂ by a solution containing the CO₂ absorption mediating compound MDEA.

The recombinant carbonic anhydrase of SEQ ID NO: 1152 was chemically modified by treatment with either 0.25% or 2.5% concentrations of the cross-linking agent suberic acid bis(N-hydroxysuccinimide). The chemically modified enzyme was assayed for carbonic anhydrase activity at 1 g/L in 4.2 M MDEA, unloaded with CO₂ at 50° C.

Preparation and Assay of Chemically Modified Carbonic Anhydrase Polypeptides:

Shake-flask powder preparations of the recombinant carbonic anhydrase of SEQ ID NO: 1152 were chemically modified by treatment with the cross-linking agent suberic acid bis(N-hydroxysuccinimide) (Sigma-Aldrich Corp., St. Louis, USA) according to the same method used for glutaraldehyde treatment in Example 8, except that suberic acid bis(N-hydroxysuccinimide) ("suberic-NHS") was added as a solid to the polypeptide solution and was mixed during the 1-4 hour incubation period. Two different treatment concentrations of suberic-NHS were used: 0.25% and 2.5%.

Results:

As shown in Table 23 below, there was a slight increase in the initial carbonic anhydrase activity for the sample solution of the carbonic anhydrase polypeptide of SEQ ID NO: 1152 modified with a 0.25% concentration of the cross-linking agent but no significant activity increase for the 2.5% sample. The sample modified with 2.5% cross-linking agent, however, exhibited no significant loss of activity after 23 h and 46 h incubation in the assay solvent at 75° C. In contrast, the same carbonic anhydrase polypeptide when unmodified showed 45 and 43% residual activity after 23 and 46 h (in the same solvent) and the 0.25% sample showed 58 and 41% residual activity in the same time frame and solvent. Thus, treatment with 2.5% of the suberic-NHS cross-linking agent results in a chemically modified enzyme with increased stability.

TABLE 23

Sample	Initial Activity $k_{1,CA}$ (s^{-1}) (% residual activity)	Activity $k_{1,CA}$ (s^{-1}) after 23 h challenge at 75° C. (% residual activity)	Activity $k_{1,CA}$ (s^{-1}) after 46 h challenge at 75° C. (% residual activity)	Activity $k_{1,CA}$ (s^{-1}) after 115 h challenge at 75° C. (% residual activity)
SEQ ID NO: 1152 (unmodified)	147 (100%)	66 (45%)	63 (43%)	16 (11%)
SEQ ID NO: 1152 + 0.25% Suberic NHS treatment	211 (100%)	122 (58%)	86 (41%)	40 (19%)
SEQ ID NO: 1152 + 2.5% Suberic NHS treatment	154 (100%)	157 (102%)	145 (94%)	36 (24%)

All publications, patents, patent applications and other documents cited in this application are hereby incorporated by reference in their entireties for all purposes to the same extent as if each individual publication, patent, patent application or other document were individually indicated to be incorporated by reference for all purposes.

While various specific embodiments have been illustrated and described, it will be appreciated that various changes can be made without departing from the spirit and scope of the invention(s).

SEQUENCE LISTING

The patent contains a lengthy "Sequence Listing" section. A copy of the "Sequence Listing" is available in electronic form from the USPTO web site (<http://seqdata.uspto.gov/?pageRequest=docDetail&DocID=US08569031B2>). An electronic copy of the "Sequence Listing" will also be available from the USPTO upon request and payment of the fee set forth in 37 CFR 1.19(b)(3).

What is claimed is:

1. A method for removing carbon dioxide from a gas stream comprising the step of contacting the gas stream with a homogenous liquid solution under suitable conditions, wherein the solution comprises:

- (i) an α -class carbonic anhydrase polypeptide chemically modified by treatment with a cross-linking agent; and
 - (ii) a CO₂ absorption mediating compound;
- whereby the solution absorbs carbon dioxide from the gas stream.

2. The method of claim **1**, wherein the chemically modified carbonic anhydrase has increased carbonic anhydrase activity in the presence of the CO₂ absorption mediating compound relative to the activity of the same carbonic anhydrase polypeptide that is not chemically modified.

3. The method of claim **1**, wherein the chemically modified carbonic anhydrase has increased carbonic anhydrase activity in 4.2 M N-methyldiethanolamine (MDEA) at 50° C. compared to the activity under the same conditions of the same carbonic anhydrase polypeptide that is unmodified.

4. The method of claim **1**, wherein the cross-linking agent is selected from the group consisting of a dialdehyde, a bis-imidate ester, a bis(N-hydroxysuccinimide) ester, a diacid chloride, and mixtures thereof.

5. The method of claim **1**, wherein the cross-linking agent is a dialdehyde having one or more carbon atoms between the two aldehyde groups.

6. The method of claim **1**, wherein the cross-linking agent is a bis-imidate ester having one or more carbon atoms between the two imidate ester groups.

7. The method of claim **1**, wherein the cross-linking agent is a bis(N-hydroxysuccinimide) ester of a di-carboxylic acid.

8. The method of claim **1**, wherein the α -class carbonic anhydrase polypeptide is a recombinant carbonic anhydrase polypeptide derived from an α -class carbonic anhydrase comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 1298, 1300, 1302, 1304, 1306, and 1308.

9. The method of claim **1**, wherein the CO₂ absorption mediating compound is an amine compound selected from the group consisting of: 2-(2-aminoethylamino)ethanol (AEE), 2-amino-2-hydroxymethyl-1,3-propanediol (AHPD), 2-amino-2-methyl-1-propanol (AMP), diethanolamine (DEA), diisopropanolamine (DIPA), N-hydroxyethylpiperazine (HEP), N-methyldiethanolamine (MDEA), monoethanolamine (MEA), N-methylpiperazine (MP), piperazine, piperidine, 2-(2-tert-butylaminoethoxy)ethanol (TBEE), triethanolamine (TEA), triisopropanolamine TIA), tris, 2-(2-aminoethoxy)ethanol, 2-(2-tert-butylaminopropoxy)ethanol, 2-(2-tert-amylaminoethoxy)ethanol, 2-(2-isopropylaminopropoxy)ethanol, 2-(2-(1-methyl-1-ethylpropylamino)ethoxy)ethanol, and mixtures thereof.

10. The method of claim **1**, wherein the CO₂ absorption mediating compound is MDEA and the suitable conditions

comprise an MDEA concentration of at least 3 M and a solution temperature of from 40° C. to 110° C.

11. The method of claim **1**, wherein the CO₂ absorption mediating compound is ammonia and the suitable conditions comprise an ammonia concentration of 1 M to 8 M and a solution temperature of from 0° C. to 20° C.

12. The method of claim **1**, wherein the CO₂ absorption mediating compound is carbonate ion and the suitable conditions comprise from 0.1 M CO₃⁻² to 5 M CO₃⁻².

13. The method of claim **1**, wherein the method further comprises exposing the homogenous solution comprising the chemically modified carbonic anhydrase polypeptide, the CO₂ absorption mediating compound, and absorbed carbon dioxide to suitable conditions for desorbing the carbon dioxide from the solution.

14. A homogenous liquid formulation comprising an aqueous solution of a CO₂ absorption mediating compound and a soluble composition having carbonic anhydrase activity comprising an α -class carbonic anhydrase polypeptide chemically modified by treatment with a cross-linking agent, wherein the chemically modified carbonic anhydrase has increased carbonic anhydrase activity in the presence of the CO₂ absorption mediating compound relative to the activity of the same carbonic anhydrase polypeptide that is not chemically modified.

15. The homogenous liquid formulation of claim **14**, wherein cross-linking agent is selected from the group consisting of a dialdehyde, a bis-imidate ester, a bis(N-hydroxysuccinimide) ester, a diacid chloride, and mixtures thereof.

16. The homogenous liquid formulation of claim **14**, wherein the α -class carbonic anhydrase polypeptide is a recombinant carbonic anhydrase polypeptide having an activity half-life (t_{1/2}) of at least 9 hours in 4 M MDEA at 50° C. prior to chemical modification and which is derived from an α -class carbonic anhydrase comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 1298, 1300, 1302, 1304, 1306, and 1308.

17. A method for removing carbon dioxide from a gas stream comprising the step of contacting the gas stream with a homogenous liquid formulation of claim **14** under suitable conditions, whereby the homogenous liquid formulation absorbs carbon dioxide from the gas stream.

18. A method for preparing a chemically modified α -class carbonic anhydrase comprising contacting in a solution: (i) a recombinant carbonic anhydrase polypeptide derived from an α -class carbonic anhydrase comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 1298, 1300, 1302, 1304, 1306, and 1308; and (ii) a cross-linking agent selected from the group consisting of a dialdehyde, a bis-imidate ester, a bis(N-hydroxysuccinimide) ester, a diacid chloride, and mixtures thereof.

* * * * *