



(86) Date de dépôt PCT/PCT Filing Date: 2012/06/19
(87) Date publication PCT/PCT Publication Date: 2013/01/10
(45) Date de délivrance/Issue Date: 2019/10/29
(85) Entrée phase nationale/National Entry: 2013/12/19
(86) N° demande PCT/PCT Application No.: FI 2012/050634
(87) N° publication PCT/PCT Publication No.: 2013/004895
(30) Priorité/Priority: 2011/07/06 (FI20115726)

(51) Cl.Int./Int.Cl. *A23C 9/00*(2006.01),
A23C 9/12(2006.01), *A23C 9/14*(2006.01),
A23C 9/142(2006.01), *A23C 9/15*(2006.01)
(72) Inventeurs/Inventors:
KALLIOINEN, HARRI, FI;
JARVIO, SOILE, FI
(73) Propriétaire/Owner:
VALIO LTD, FI
(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : FORMULATION A BASE DE LAIT
(54) Title: MILK-BASED FORMULATION

(57) Abrégé/Abstract:

The invention relates to a milk-based formulation with a reduced carbohydrate content, having a ratio of carbohydrates to protein of at most 1.1, a protein content of at least 5.4% on dry matter basis, and a ratio of ash to protein is substantially similar to that of a milk raw material used as a starting material, and a method for the preparation thereof. The milk-based formulation can be used in the preparation of a recombined milk product with a reduced lactose content, having a full taste of normal milk without any deficiencies in its organoleptic properties.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(10) International Publication Number

WO 2013/004895 A1

(43) International Publication Date
10 January 2013 (10.01.2013)

(51) International Patent Classification:

A23C 9/12 (2006.01) *A23C 9/146* (2006.01)
A23C 9/142 (2006.01) *A23C 9/15* (2006.01)

(21) International Application Number:

PCT/FI2012/050634

(22) International Filing Date:

19 June 2012 (19.06.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

20115726 6 July 2011 (06.07.2011) FI

(71) **Applicant** (*for all designated States except US*): **VALIO LTD** [FI/FI]; Meijeritie 6, FI-00370 Helsinki (FI).

(72) Inventors; and

(75) **Inventors/Applicants** (*for US only*): **KALLIOINEN, Harri** [FI/FI]; Nummisten tie 747, FI-04480 Haarajoki (FI). **JÄRVIÖ, Soile** [FI/FI]; Kiviniitynkuja 1 C, FI-02340 Masala (FI).

(74) Agent: **KOLSTER OY AB**; Iso Roobertinkatu 23, P.O.Box 148, FI-00121 Helsinki (FI).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:

— *with international search report (Art. 21(3))*



WO 2013/004895 A1

(54) Title: MILK-BASED FORMULATION

(57) **Abstract:** The invention relates to a milk-based formulation with a reduced carbohydrate content, having a ratio of carbohydrates to protein of at most 1.1, a protein content of at least 5.4% on dry matter basis, and a ratio of ash to protein is substantially similar to that of a milk raw material used as a starting material, and a method for the preparation thereof. The milk-based formulation can be used in the preparation of a recombined milk product with a reduced lactose content, having a full taste of normal milk without any deficiencies in its organoleptic properties.

MILK-BASED FORMULATION

FIELD OF THE INVENTION

The present invention relates to milk-based formulations. More particularly, the invention relates to milk-based formulations which are suitable for the preparation of recombined milk products with a reduced carbohydrate content.

BACKGROUND OF THE INVENTION

It is well recognized that recombined milk beverages prepared by recombining milk powder and water have several defects compared to normal milk. Deficiencies appear especially in the preparation of recombined milk beverages with reduced lactose contents. The deficiencies are both organoleptical and nutritional in nature. For instance, the recombined milk beverages are susceptible to form precipitates/sediments during heat treatment, and especially during storage of the beverages. The precipitated matter is mainly consisted of milk proteins. It is known that among milk proteins, whey proteins especially are very heat-sensitive. Certain minerals may also have an influence on the precipitation phenomenon of the recombined milk beverages. Currently, the problems associated with the sensitivity of precipitation of the recombined milk products, agglomeration has been avoided by using additives like stabilizers in the products.

Low lactose or lactose-free milk beverages can be prepared from recombined milk beverages by hydrolyzing the lactose thereof. However, the amount of natural lactose content is relatively high, which means that the hydrolysis of lactose produces a great amount of glucose and galactose imparting an uncommon sweet taste to the product.

One important factor causing various problems to the recombined milk products is the Maillard reaction which is a nonenzymatic browning reaction between reducing sugars and free amino groups of milk proteins. The Maillard reaction is a common problem especially in lactose-hydrolyzed milk products. Reducing sugars, glucose and galactose, produced in lactose hydrolysis are more reactive than lactose, thus causing stronger Maillard browning reaction. In hydrolyzed milk, the molar content of these reducing monosaccharides is almost double in comparison with that of regular milk lactose. It is further known that the Maillard reaction becomes even stronger during heat treatment of the lactose-hydrolyzed milk products.

The Maillard browning products cause an undesirable change in the organoleptic properties of the heat-treated milk, such as taste, colour, and struc-

ture. In addition, the Maillard reaction has a detrimental effect on the nutritional quality of milk. Bioavailability of lysine which is an important amino acid for the nutritional value is reduced. The Maillard reaction and destruction of lysine continues during the storage at a room temperature after the heat treatment of the 5 milk product.

The Maillard reaction can be monitored by furosine, which presents products formed in the reactions between free amino groups and reducing sugars, and resulting in a loss of their availability.

It is further known that drying of milk to powder changes the quality of 10 milk protein which can be seen in the recombined milk products. The Maillard reaction proceeds also during the production process of milk powders. Proteins also denature during drying.

Among many people, there is still much suspicion towards the use of 15 the recombined milk products obviously because of their defective organoleptic properties. Typically, the recombined milk products are not perceived as equating to normal milk.

There is thus a need for new milk products which are suitable for preparing recombined milk products with a reduced carbohydrate content and good 20 nutritional quality, and which reduce or eliminate sediment formation and minimize the taste and texture problems of the recombined milks.

It is very challenging to achieve recombined milk products with reduced carbohydrate contents that are completely flawless in taste and structure, that meet the consumers' expectations on an organoleptically competent milk product, and that are produced economically and simply.

25 BRIEF DESCRIPTION OF THE INVENTION

It has now been found a milk-based formulation which is suitable for preparing recombined milk products with a reduced carbohydrate content, having a good taste of normal milk, and which avoids the typical problems associated with the similar prior art products. The recombined milk products are completely flawless in their organoleptic properties, especially in taste. 30

In an aspect, the invention provides a milk-based formulation with a reduced carbohydrate content, having a ratio of carbohydrates to protein of at most 1.1, a protein content of at least 5.4% on dry matter basis, and a ratio of ash to protein is substantially similar to that of a milk raw material used as a 35 starting material.

It was surprisingly found that the defects in organoleptic properties, like precipitation and defects in taste, and nutritional quality of the recombined milk products can be avoided by reducing the lactose content of the milk-based formulation before concentrating it to a form suitable for recombination. As the 5 amount of the products produced in lactose hydrolysis of milk, involving in the Maillard reaction, are reduced, the Maillard reaction is significantly suppressed and the typical problems associated therewith are avoided.

Moreover, the shelf life of the recombined milk products of the inventions, especially at a room temperature, is prolonged.

10 In another aspect, the invention provides a process for producing a milk-based formulation of the invention, comprising removing carbohydrates from a milk raw material to provide the milk-based formulation with a reduced carbohydrate content.

15 In a still another aspect, the invention provides a process for producing a milk-based formulation of the invention, comprising

- a) separating components of a milk raw material into a protein fraction, carbohydrate fraction and mineral fraction,
- b) combining at least a portion of the protein fraction and of the mineral fraction to provide the milk-based formulation.

20 In a further aspect, the invention provides a use of milk powder having a ratio of carbohydrates to protein of at most 1.1 and a protein content of at least 5.4% on dry matter basis for the preparation of a recombined milk product with a reduced carbohydrate content.

25 The milk powder of the invention enables to prepare lactose-free milk with good taste without investments in equipment in the plant where the lactose-free milk is prepared. Lactose-free milk can be prepared from the milk powder of the invention and water and/or normal milk. Due to the favourable composition of the milk powder it can be dried to powder with reduced Maillard reactions. Quality of the recombined milk product prepared from the milk powder of the invention is thus improved.

30 In a still further aspect, the invention provides a process for producing a recombined milk product having a reduced carbohydrate content, comprising recombining milk powder having a ratio of carbohydrates to protein of at most 1.1 and a protein content of at least 5.4% on dry matter basis, and a liquid and 35 optional other ingredients to provide the recombined milk product with a reduced carbohydrate content. The composition of the recombined milk product can be

adjusted to correspond to that of normal milk except for lactose by combining the milk powder and liquid without any further additives, like mineral supplement.

The invention provides a process that is simple, economic, industrially applicable on a large scale, and does not cause additional cost.

The invention also provides a process which brings remarkable savings in transportation cost.

The invention further provides a process which allows to control the adverse effects of recombination in an economical, efficient and simple manner.

Taste, colour, and structural defects caused by natural milk enzyme activities and microbe-induced enzyme activities as well as the use of a lactase enzyme and side activities typical of commercial enzyme preparations in the heat-treated recombined milk products are avoided with the process of the invention. With the process of the invention, it is possible to improve the organoleptic properties, especially the stability of the taste properties and structure at room temperature, of the recombined milk products with reduced carbohydrate contents, as a result of which the shelf life of the product can be extended.

The claimed invention pertains to a milk-based powder with a reduced carbohydrate content, having a ratio of carbohydrates to protein of at most 1.1:1, and a protein content of at least 45% on dry matter basis, and wherein a ratio of ash to protein of the milk-based powder is substantially the same as that of a milk raw material used as a starting material.

The claimed invention also pertains to a process for preparing such a milk-based powder with a reduced carbohydrate content, comprising removing carbohydrates from said milk raw material to provide a milk-based formulation with a reduced carbohydrate content, and concentrating the milk-based formulation to form the milk-based powder.

The claimed invention also pertains to a process for producing a recombined milk product having a reduced carbohydrate content, comprising recombining milk powder having a ratio of carbohydrates to protein of at most 1.1:1 and a protein content of at least 45% on dry matter basis, and a liquid and optional other ingredients to provide the recombined milk product with a reduced carbohydrate content, wherein the milk powder has a ratio of ash to protein that is substantially the same as that of a milk raw material used as a starting material.

The claimed invention also pertains to use of milk powder having a ratio of carbohydrates to protein of at most 1.1:1 and a protein content of at least 45% on dry matter basis for the preparation of a recombined milk product with a reduced carbohydrate content, wherein the milk powder has a ratio of ash to protein that is substantially the same as that of a milk raw material used as a starting material.

The claimed invention also pertains to a process for preparing such a milk-based powder with a reduced carbohydrate content, comprising removing carbohydrates from said milk raw material to provide a milk-based formulation with a reduced carbohydrate content, and concentrating the milk-based formulation to form the milk-based powder.

DETAILED DESCRIPTION OF THE INVENTION

The term "recombined milk products" is used herein to mean milk products which are prepared by recombining a milk-based formulation with a liquid. Further ingredients, like milk fat (cream, butter) can be incorporated into the recombined milk product in order to obtain a desired fat content of the product. Formulation can also be produced with a desired fat, protein and mineral content.

The term "liquid" is used herein to mean water, a milk raw material, or plant (vegetable) origin material, or a combination of these. Accordingly, the liquid can be, for instance, side streams obtained from manufacturing process of milk products, such as rinsing waters (washing waters) derived from streams obtained from washing/rinsing of processing pipes, containers and vessels in dairy and plant (vegetable) product manufacturing plants. Typically, side streams include UF permeates, NF permeates, RO permeates, RO retentates, diawater, or mixtures thereof. Preferably, the liquid is water, skim milk or RO-retentate.

The term "milk raw material" is used herein to mean to milk, whey, and combinations of milk and whey as such obtained from an animal, such as a cow, sheep, goat, camel, mare or any other animal that produces milk suitable for human consumption, or milk that is pre-processed as desired, for example as

a concentrate. The milk may be supplemented with ingredients generally used in the preparation of milk products, such as fat, protein or sugar fractions, or the like. The milk may thus be, for instance, full-fat milk, cream, low-fat milk or skim milk, ultrafiltered milk, diafiltered milk, microfiltered milk or milk recombined from 5 milk powder, organic milk or a combination of these. The milk raw material can be, for instance, side streams obtained from manufacturing process of milk products. Preferably, the milk raw material is skim milk.

In an aspect, the invention provides a milk-based formulation with a reduced carbohydrate composition, having a ratio of carbohydrates to protein of 10 at most 1.1, a protein content of at least 5.4% on dry matter basis, and a ratio of ash to protein is substantially similar to that of a milk raw material used as a starting material.

In an embodiment, the formulation has a dry matter content of 8 to 60%.

In an embodiment, the formulation is powder. The dry matter content 15 of powder is typically in the range from 94 to 100%.

An ash to protein ratio of the formulation has a significant effect on the organoleptic properties, especially on taste, of the formulation. In an embodiment of the invention, ash is provided as milk-based minerals in the formulation.

In an embodiment, the ratio of carbohydrates to protein is at most 0.9. 20 In another embodiment, the ratio of carbohydrates to protein is at most 0.4. In still another embodiment, the ratio of carbohydrates to protein is at least 0.02.

In an embodiment, the milk-based formulation comprises 5.4 to 80% protein on dry matter basis.

In an embodiment, a ratio of the monovalent minerals of milk to protein 25 is substantially similar to that of the milk raw material.

In an embodiment, the milk-based formulation is powder.

It is generally known that the protein content of natural milk can vary within quite a wide range, depending on animal species, breed, feeding and season, etc. For example, the protein content of milk obtained from cows can 30 vary between 1.8 to 6.3%.

In an embodiment, the milk-based formulation comprises 5.4 to 65% protein, 4.6 to 41% carbohydrates, and 1.0 to 14% ash on non-fat total solid basis. In another embodiment, the milk-based formulation comprises about 48 to 60% protein, about 24 to 43% carbohydrates, and 10 to 13% ash.

The carbohydrate content of the milk raw material can be reduced by any manner known in the art. Carbohydrates present in natural milk are primarily lactose. Prior to lactose removal, the fat content of the milk raw material may be reduced.

5 In another aspect, the invention provides a process for preparing a milk-based formulation of the invention, with a reduced carbohydrate content, comprising removing carbohydrates from a milk raw material to provide the milk-based formulation with a reduced carbohydrate content.

10 In an embodiment, the milk-based formulation is concentrated to powder by an appropriate process, for example by spray drying.

15 In an embodiment, lactose is removed from the milk raw material by precipitation. The precipitation can be performed in a manner commonly known in the art. Precipitation of lactose allows to efficiently remove lactose from a milk raw material while all the other milk components are substantially retained in a desired manner.

20 In another embodiment, the lactose removal can be performed by means of enzymatic reactions. Conventional enzymatic processes for splitting lactose, or for conversion of lactose to derivatives, such as lactulose, lactitol, lactobionic acid and their degradation products, are generally known in the field. The process for splitting lactose comprising the step of adding lactase from fungus or yeast to milk in such a manner that lactose is split into monosaccharides, i.e. glucose and galactose, in over 80%. Enzymes typically catalyse several types of reactions, i.e. lactase both hydrolysis and transglucosylation reactions.

25 In still another embodiment, lactose is removed from the milk raw material by chromatographic separation. The milk raw material is eluted through a column filled with a cation exchange resin. The separation can be carried out so that a significant portion of the proteins and minerals are collected into a single fraction whereas lactose remains in the column.

30 In still another embodiment, lactose is removed from the milk raw material by means of a membrane filtration technique. Use of membranes with different cut-off values allows to efficiently separate the various milk components, i.e. proteins, carbohydrates and minerals, from each other into different fractions.

If desired, various techniques mentioned above for lactose removal can be combined in an appropriate manner.

Thus, in an aspect, the invention provides a process for preparing a milk-based formulation of the invention, with a reduced carbohydrate content, comprising

- a) separating components of a milk raw material into a protein fraction, carbohydrate fraction and mineral fraction,
- 5 b) combining at least a portion of the protein fraction and of the mineral fraction to provide the milk-based formulation.

In an embodiment, the milk-based formulation prepared above is concentrated to powder.

10 In dairy industry, ultrafiltration is typically used for separating the proteins and fat from lactose and minerals of milk. Prior to ultrafiltration, the fat content of milk can be standardized for example by separation. The milk proteins, and any fat, remain in the ultrafiltration retentate whereas lactose and minerals pass into a permeate. Ultrafiltration is typically carried out by a concentration 15 factor of 1 to 10.

20 Lactose and the monovalent minerals, mainly sodium and potassium, present in the ultrafiltration permeate can be separated from each other by nanofiltration. Lactose remains in a nanofiltration retentate, and the monovalent minerals pass into a nanofiltration permeate. The concentration factor of nanofiltration is typically in the range of 1 to 6.

The nanofiltration permeate solution including the monovalent minerals can be concentrated by reverse osmosis to provide a mineral concentrate for use in the milk-based formulation of the invention. Reverse osmosis is generally carried out by a concentration factor of 2 to 20.

25 In an embodiment, the dry matter content of the milk protein fraction and/or the milk mineral fraction obtained from membrane filtrations can be increased by evaporation, for example to about 17%, and use the evaporated fraction(s) for the preparation of the milk-based formulation.

30 As stated above, precipitation of lactose from the milk raw material, enzymatic lactose hydrolysis or conversion, and chromatographic lactose separation provide a milk-based formulation where all the other milk components except for lactose are retained in the composition. Membrane filtration technology further provides the possibility to separate lactose and milk minerals, mainly monovalent minerals, from each other and to separately compose an optimized 35 and tailor-made milk-based formulation of the invention from the protein and mineral fractions.

In an embodiment, one or more of the milk component fractions are separately concentrated to powder by an appropriate process and then used for preparing the milk-based formulation of the invention with a desired composition. In another embodiment, the milk-based formulation is first prepared from said fractions in an appropriate manner and then dried to powder.

In a further aspect, the invention provides a use of milk powder having a ratio of carbohydrates to protein of at most 1.1 and a protein content of at least 5.4% on dry matter basis for the preparation of a recombined milk product with a reduced carbohydrate content. In an embodiment, a ratio of ash to protein is substantially similar to that of a milk raw material used as a starting material. In an embodiment, the protein content of the milk powder is at least 37%. In another embodiment, the protein content is at least 45%.

In an embodiment, the recombined milk product is a milk beverage.

In an embodiment, the milk powder is the milk-based formulation of the invention, or prepared by the process of the invention described above.

The amount of the milk powder in the recombined milk product is 0.5 to 15% by weight, preferably 2 to 7% by weight.

In still a further aspect, the invention provides a process for producing a recombined milk product having a reduced carbohydrate content, comprising recombining milk powder having a ratio of carbohydrates to protein of at most 1.1 and a protein content of at least 5.4% on dry matter basis, and a liquid and optional other ingredients to provide the recombined milk product with a reduced carbohydrate content. In an embodiment, the protein content of the milk powder is at least 37%. In another embodiment, the protein content is at least 45%.

The process of the invention may be applied to an industrial large-scale production or to a small-scale production in households.

The lactose content of the recombined milk product is at most 3.1% by weight.

The milk powder can be recombined in a liquid which can be water, a milk raw material, plant (vegetable) origin material, or a mixture thereof. In an embodiment, the recombined milk is prepared so as correspond to composition of normal skim milk except for reduced carbohydrates. In an embodiment of the invention, an ash to protein ratio of the recombined milk product is similar to that of normal skim milk. In an embodiment, ash is provided as milk-based minerals in the recombined milk product. The composition of the recombined milk product

can be adjusted to correspond to that of normal skim milk except for reduced carbohydrates by combining the milk powder and liquid without any further additives, like mineral supplement. However, also milk-based minerals can be added to the recombined milk product.

5 The recombined milk product of the invention can be supplemented with ingredients typically used in milk product. Optional ingredients include edible fat, like milk fat or vegetable fat such as rapeseed oil, fractionated palm oil or coconut oil, vitamins, minerals, fibre, probiotics, flavours.

10 In an embodiment, the recombined milk product is prepared by recombining powder 1 of the invention with skim milk and water. In another embodiment, the recombined milk product is prepared by recombining 2.4% of powder 1, 54% of skim milk and 43.6% of water.

15 In an embodiment, the recombined milk product is subjected to a heat treatment. The heat treatment can be performed, for example, at a temperature of 153°C for 4 seconds.

20 If desired, any residual lactose present in the recombined milk product is subjected to lactose hydrolysis and/or conversion. The lactose hydrolysis and/or conversion can be performed with commercially available lactase enzymes in a manner known per se. In an embodiment, the lactose content of the recombined milk product is less than 1% by weight, generally designated as a low lactose milk beverage. In another embodiment, the lactose content is less than 0.01% by weight, generally designated as a lactose-free milk beverage.

25 Lactose hydrolysis can be performed on the recombined milk product which has been heat treated in a manner described above, or on the recombined milk product which is subsequently subjected to the heat treatment. In an embodiment, the lactose hydrolysis is performed after the heat treatment.

30 The furosine content of the recombined milk beverages prepared according to the invention were measured before and after the heat treatment of the beverages. The furosine content was compared to that of skim milk (beverage 6) and a recombined skim milk prepared by recombining conventional skim milk powder and water (beverage 7), which were used as references. The furosine contents of the beverages prepared from the milk compositions of the invention were lower than that of the reference recombined skim milk both before and after the heat treatment. Specifically, in milk beverages prepared by recombining the milk composition of the invention and water, or a mixture of skim milk and water, the furosine contents were significantly lower than that of the refer-

ence recombined skim milk. This indicates that the Maillard reaction is efficiently suppressed even in heat-treated recombined milk beverages. In an embodiment, the furosine content of the recombined milk product of the invention is at most 0.61 mg/g protein prior to heat treatment of the product. In another embodiment, 5 the furosine content of the recombined milk product of the invention is at most 0.92 mg/g protein after heat treatment of the product.

The organoleptic properties of the recombined milk product prepared according to the invention unexpectedly keep at room temperature even during a long storage. The process is easy to perform in production conditions without 10 significant extra costs.

The following examples are presented for further illustration of the invention without limiting the invention thereto.

Examples

Example 1

15 Skim milk was ultrafiltered with a GR61PP membrane (Dow, USA) at a temperature of 10°C and with a concentration factor of 4 to provide an ultrafiltration (UF) permeate and an ultrafiltration (UF) retentate. The UF permeate was further nanofiltered with a Desal DK membrane (Osmonics, USA) at temperatures from 10°C to 15°C and with a concentration factor of 4 to provide a nanofiltration (NF) permeate and a nanofiltration (NF) retentate. The NF permeate was 20 concentrated by reverse osmosis (RO) with a Filmtec RO-390-FF membrane (Dow, USA) and with a concentration factor of about 10 to provide an RO permeate and RO retentate.

The RO retentate was evaporated to a dry matter content of 17%.

25 The compositions of the milk raw material, i.e. skim milk, the UF retentate and permeate, the RO retentate, and the evaporated RO retentate are given in Table 1 below.

Table 1.

	Skim milk	UF ret	UF perm	RO ret	RO ret evapor.
Protein (%)	3.4	12.2	0.2	-	
Fat (%)	<0.1	0.2	0.0		
Lactose (%)	4.6	4.9	4.5	0.2	2.0
Ash (%)	0.8	1.6	0.5	1.7	13.8
Dry matter (%)	9.3	19.3	5.5	2.1	17
Na (mg/kg)	390	450	370	1 700	13 500
K (mg/kg)	1 700	1 990	1 600	6 300	50 200
Ca (mg/kg)	1 200	3 700	250	180	1 400
Mg (mg/mg)	120	250	70	50	420
P (mg/kg)	940	2 500	340	270	2 100
Na+K/protein (mg/g protein)		20			

The evaporated RO retentate (700 kg) and the UF retentate (10 000 kg) were mixed together and evaporated to a dry matter content of 33% to 40%.

5 The resultant mixture was dried to powder (powder 1) with a corresponding heat treatment of the conventional low-heat milk powder.

The UF retentate obtained from ultrafiltration of skim milk was evaporated and dried to powder (powder 2) with a corresponding heat treatment of the conventional low-heat milk powder.

10 The compositions of powder 1 and powder 2 are given in Table 2 below.

Table 2. Compositions of powder 1 and powder 2

	Powder 1	Powder 2
Protein (%)	57.5	60.7
Fat (%)	0.8	0.9
Lactose (%)	23.9	24.5
Ash (%)	12.6	7.7
Dry matter (%)	97	96.5
Na (mg/kg)	6 500	2 300
K (mg/kg)	25 900	10 000
Ca (mg/kg)	18 000	18 500
Mg (mg/kg)	1 300	1 200
P (mg/kg)	12 700	12 600
Na+K/protein (mg/g protein)	56	20

Example 2

Skim milk was ultrafiltered in a manner described in Example 1 except that the concentration factor was 1.6 to provide an UF retentate.

5 Cheese whey was nanofiltered with a Desal DK membrane (Osmonics, USA) at a temperature < 15°C and with a concentration factor of 4.5 to provide a nanofiltration (NF) permeate and a nanofiltration (NF) retentate. The NF permeate was concentrated by reverse osmosis (RO) as described in Example 1. The RO retentate obtained was evaporated to a dry matter content of 17.5%.

10 The compositions of the UF retentate, the cheese whey, the RO retentate, and the evaporated RO retentate are given in Table 3 below.

Table 3.

	UF ret	Cheese whey	RO ret	RO ret evapor.
Protein (%)	5.5	0.7	-	
Fat (%)	0.1	<0.1		
Lactose (%)	4.7	4.4	0.29	3.0
Ash (%)	1.0	0.52	1.41	14.6
Dry matter (%)	10.7	5.8	1.69	17.5
Na (mg/kg)	400	370	1 400	14 100
K (mg/kg)	1 800	1 500	5 100	52 500
Ca (mg/kg)	1 700	370	150	1 500
Mg (mg/mg)	150	80	45	470
P (mg/kg)	1 300	390	240	2 500

The UF retentate of skim milk (10 000 kg) and the evaporated RO retentate of cheese whey (120 kg) were mixed together and evaporated and dried to powder (powder 3) as described in Example 1. The composition of powder 3 is given in Table 4 below.

Table 4.

	Powder 3
Protein (%)	46.9
Fat (%)	0.7
Lactose (%)	40.9
Ash (%)	10.4
Dry matter (%)	97
Na (mg/kg)	5 000
K (mg/kg)	21 000
Ca (mg/kg)	15 000
Mg (mg/mg)	1 300
P (mg/kg)	11 600
Na+K/protein (mg/g protein)	55

Example 3

5 Skim milk was evaporated to provide a milk concentrate with a dry matter content of 30%. The milk concentrate was pumped to a chromatography column filled with a cation exchange resin. Skim milk eluted through the cation exchange resin was collected so that a significant portion of milk minerals and proteins were in the same fraction. When the collection of fractions was completed, a significant portion of lactose still remained in the chromatography column. The chromatographic separation was performed at a temperature of about 60°C.

10 The fraction obtained from the chromatographic separation was dried to powder (powder 4) as described in Example 1.

The composition of the collected fraction from chromatography and powder 4 are given in Table 5 below.

Table 5.

	Chromatographic fraction of skim milk	Powder 4
Protein (%)	5.47	64.1
Fat (%)	0.09	0.9
Lactose (%)	1.4	16.3
Ash (%)	1.2	13.5
Dry matter (%)	8.3	96.7
Na (mg/kg)	1 100	12 800
K (mg/kg)	2 200	25 700
Ca (mg/kg)	1 400	18 500
Mg (mg/kg)	180	2 100
P (mg/kg)	1 400	16 000
Na+K/protein (mg/g protein)		60

15

Example 4

Various milk beverages were prepared from the fractions and powders obtained in Examples 1 to 3. The specific recipes are given in Table 7. The numerical values illustrate percentages of each fraction in the recipe.

20 In addition to said fractions and powders, milk and low-heat milk powder were used in the preparation of the beverages. Their compositions are given in Table 6.

Table 6.

	Skim milk	Low-heat milk powder
Protein (%)	3.4	36.8
Fat (%)	<0.1	0.4
Lactose (%)	4.6	46.1
Ash (%)	0.8	8.2
Dry matter (%)	9.0	96.6
Na (mg/kg)	390	4 200
K (mg/kg)	1 700	17 000
Ca (mg/kg)	1 200	12 000
Mg (mg/kg)	120	1 200
P (mg/kg)	940	9 500
Na+K/protein (mg/g prot)	61	58

Table 7.

	Powder 1	Powder 2	Powder 3	Powder 4	Skim milk	Low heat milk powder	Water	RO ret	UF ret (Ex- ample 1)
Beverage 1	2.4				54.0		43.6		
Beverage 2		2.4			54.0		42.7	0.9	
Beverage 3			7.1				92.9		
Beverage 4		2.4			54.0		43.6		
Beverage 5	2.4					5.4	92.2		
Beverage 6 (reference)					100				
Beverage 7 (reference)						9.0	91.0		
Beverage 8				2.0	60.0		38.0		
Beverage 9					54.0		33.4	0.9	11.9

The beverages were pasteurized with a direct steam infusion UHT (ultra high temperature) equipment (APV, Denmark) at a temperature of 153°C for 4 seconds. The beverages were packed aseptically. Prior to packing, 0.03% by weight of lactase (Godo YNL2, Oenon, Japan) was added aseptically to the beverages. The milk beverages were lactose-free after one week storage. The composition of the above nine beverages are given in Table 8.

Table 8.

	Beverage 1	Beverage 2	Beverage 3	Beverage 4	Beverage 5	Beverage 6 (reference)	Beverage 7 (reference)	Beverage 8	Beverage 9
Protein (%)	3.3	3.3	3.3	3.3	3.4	3.4	3.3	3.3	3.3
Fat (%)	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
Lactose (%)	3.0	3.1	2.90	3.07	2.06	4.60	4.15	3.09	3.08
Ash (%)	0.75	0.77	0.81	0.64	0.82	0.77	0.81	0.76	0.76
Dry matter (%)	7.23	7.37	6.96	7.21	7.62	9.00	8.77	7.36	7.31
Na (mg/kg)	370	400	370	270	400	390	390	500	400
K (mg/kg)	1 500	1 600	1 500	1 100	1 500	1 700	1 500	1 500	1 600
Ca (mg/kg)	1 100	1 100	1 100	1 100	1 100	1 200	1 100	1 100	1 100
Mg (mg/kg)	100	100	90	90	100	120	110	110	100
P (mg/kg)	810	830	820	810	820	940	900	900	820
<i>After heat treatment</i>									
Furosine (mg/kg)	18	18	30	19	31	14	42	18	14
mg/g protein	0.55	0.54	0.90	0.57	0.92	0.46	1.27	0.54	0.43
<i>Prior to heat treatment</i>									
Furosine (mg/kg)	7	8	20	8	21	2	24	6	2
mg/g protein	0.23	0.23	0.59	0.23	0.61	0.06	0.73	0.19	0.06

The beverages having a temperature of about 15°C were evaluated in their organoleptic properties after about one week after preparation thereof. Results of the evaluation are given in Table 9. The organoleptic properties were compared to normal skim milk (beverage 6) including natural lactose content.

Table 9.

Beverage	
1	Similar to normal milk
2	Similar to normal milk
3	Similar to normal milk
4	Similar to normal milk, slightly watery
5	Similar to normal milk
6	Sweet
7	Sweet
8	Similar to normal milk
9	Similar to normal milk

All the beverages 1 to 9 were prepared so as to have a protein content to a desired level, i.e., about 3.3% (Table 8). Thus, the other results of the beverages are comparable and reliable. It can be seen from the results shown in Table 8 that the furosine content of reference beverage 7 which is a recombinant milk beverage produced from skim milk powder alone is significantly higher than that of the other beverages. The lowest furosine level is achieved in beverage 6 which is normal skim milk having a reduced fat content but a natural lactose content, and beverage 9 which is composed from skim milk, water, UF retentate and RO retentate. Surprisingly, only a slight increase in furosine contents of the other beverages compared to beverages 6 and 9 is detected although said other beverages were all prepared from protein-containing powders of milk. Also the furosine contents of beverages 3 and 5 are substantially lower than that of beverage 7 although beverages 3, 5 and 7 are all prepared from water and milk-based powders.

As to the taste of the beverages, the most similar to normal UHT-treated skim milk were beverages 1, 2, 3, 5, 8, and 9. Taste of beverage 4 was similar to that of normal milk except that it was slightly watery. The biggest deviations in taste were found in beverages 6 and 7 which were notably sweet and did not correspond in their properties to normal UHT-treated milk.

Moreover, each beverage 2 to 9 except 5 was further prepared by supplementing them with cream (38%) to provide a fat content of 1.5% to the beverage.

Powder 1 according to Example 1 was prepared from milk by ultrafiltering milk containing fat 3.1% and drying the UF-retentate to powder which was used in similar manner as powder 1 in the preparation of beverages 1 and 5. Beverages 1 and 5 had fat content of 1.5%.

The results obtained for each 1.5% fat containing beverage were similar to those shown in Table 9 except that the watery taste of beverage 4 could be partly masked by fattiness.

Quality of the beverages was controlled organoleptically for 6 months and no significant changes were obtained.

The results show that a lactose-free milk beverage of the invention, prepared from powder 1 produced in Example 1 has a better taste than the recombined milk prepared from normal skim milk powder (beverage 7; reference). Moreover, the furosine content of the beverage of the invention is substantially lower than that of the reference milk. Moreover, surprisingly, the taste and texture remained better in the beverage of the invention than in the normal UHT-treated skim milk.

It will be obvious to a person skilled in the art that, as the technology advances, the inventive concept can be implemented in various ways. The invention and its embodiments are not limited to the examples described above but may vary within the scope of the claims.

CLAIMS

1. A milk-based powder with a reduced carbohydrate content, having a ratio of carbohydrates to protein of at most 1.1:1, and a protein content of at least 45% on dry matter basis, and wherein a ratio of ash to protein of the milk-based powder is substantially the same as that of a milk raw material used as a starting material.

5 2. The milk-based powder according to claim 1, wherein the ratio of carbohydrates to protein is at most 0.9:1.

10 3. The milk-based powder according to claim 2, wherein the ratio of carbohydrates to protein is at most 0.4:1.

4. The milk-based powder according to any one of claims 1 to 3, wherein the ratio of carbohydrates to protein is at least 0.02:1.

15 5. The milk-based powder according to any one of claims 1 to 4, wherein a ratio of monovalent minerals to protein is substantially the same as that of the milk raw material.

6. The milk-based powder according to any one of claims 1 to 5, comprising 45 to 65% protein, 4.6 to 41% carbohydrates, and 1.0 to 14% ash on non-fat total solid basis.

20 7. The milk-based powder according to any one of claims 1 to 5, comprising 48 to 60% protein, 24 to 43% carbohydrates, and 10 to 13% ash on non-fat total solid basis.

8. The milk-based powder according to any one of claims 1 to 7, comprising a milk protein concentrate from ultrafiltration of the milk raw material.

25 9. The milk-based powder according to any one of claims 1 to 7, comprising a fraction comprising protein and minerals from chromatographic separation of the milk raw material.

10. The milk-based powder according to any one of claims 1 to 9, comprising a reverse osmosis concentrate of milk minerals obtained from a nanofiltration permeate derived from nanofiltration of an ultrafiltration permeate of the milk raw material.

30 11. A process for preparing a milk-based powder with a reduced carbohydrate content according to any one of claims 1 to 7, comprising removing carbohydrates from said milk raw material to provide a milk-based formulation with a reduced carbohydrate content, and concentrating the milk-based formulation to form the milk-based powder.

12. The process according to claim 11, wherein the removal of carbohydrates is performed by means of enzymes, chromatography, precipitation, membrane filtration technique or a combination thereof.

13. The process according to claim 11, comprising

5 a) separating components of a milk raw material into a protein fraction, carbohydrate fraction and mineral fraction,

b) combining at least a portion of the protein fraction and of the mineral fraction to provide the milk-based formulation.

10 14. The process according to claim 13, wherein the separation is accomplished by ultrafiltration, nanofiltration and reverse osmosis.

15 15. The process according to claim 14, wherein the milk raw material is subjected to ultrafiltration (UF) to provide an UF permeate, and the protein fraction as an UF retentate.

16. The process according to claim 15, wherein the ultrafiltration permeate is subjected to nanofiltration (NF) to provide a NF permeate, and the carbohydrate fraction as a NF retentate, and the NF permeate is subjected to reverse osmosis (RO) to provide the mineral fraction as an RO retentate.

20 17. The process according to any one of claims 13 to 16, wherein one or both of the protein fraction and the mineral fraction is concentrated to a dry matter content of at least 17%.

25 18. Use of milk powder having a ratio of carbohydrates to protein of at most 1.1:1 and a protein content of at least 45% on dry matter basis for the preparation of a recombined milk product with a reduced carbohydrate content, wherein the milk powder has a ratio of ash to protein that is substantially the same as that of a milk raw material used as a starting material.

19. The use according to claim 18, wherein the milk powder is a milk-based powder according to any one of claims 1 to 10 or prepared by a process according to any one of claims 11 to 17.

30 20. The use according to claim 18 or 19, wherein the recombined milk product is a milk beverage.

21. The use according to any one of claims 18 to 20, wherein the amount of the milk powder in the recombined milk product is 0.5 to 15% by weight.

22. The use according to claim 21, wherein the amount of the milk powder in the recombined milk product is 2 to 7% by weight.

23. A process for producing a recombined milk product having a reduced carbohydrate content, comprising

recombining milk powder having a ratio of carbohydrates to protein of at most 1.1:1 and a protein content of at least 45% on dry matter basis, and a liquid and optional 5 other ingredients to provide the recombined milk product with a reduced carbohydrate content.

wherein the milk powder has a ratio of ash to protein that is substantially the same as that of a milk raw material used as a starting material.

24. The process according to claim 23, wherein the milk powder is a milk-based powder according to any one of claims 1 to 10 or prepared by a process according to any one of claims 11 to 17.

25. The process according to claim 23 or 24, wherein the recombined milk is a beverage.

26. The process according to claim 23 or 24, wherein the carbohydrate content of the recombined milk product is at most 3.1% by weight.

27. The process according to any one of claims 23 to 26, wherein the recombined milk product is subjected to a heat treatment.

28. The process according to any one of claims 23 to 27, wherein any residual lactose of the recombined milk product is hydrolysed.

29. The process according to claim 26, wherein any residual lactose of the recombined milk product is hydrolysed after the heat treatment.

30. The process according to any one of claims 23 to 29, wherein the organoleptic properties of the recombined milk product is to the same as that of skim milk.