Office de la Propriété Intellectuelle du Canada
Un organisme d'Industrie Canada

Canadian Intellectual Property Office
An agency of Industry Canada

CA 2804311 C 2017/09/12

(11)(21) **2 804 311**

(12) **BREVET CANADIEN**
CANADIAN PATENT

(13) **C**

(86) **Date de dépôt PCT/PCT Filing Date:** 2011/06/22
(87) **Date publication PCT/PCT Publication Date:** 2011/12/29
(45) **Date de délivrance/Issue Date:** 2017/09/12
(85) **Entrée phase nationale/National Entry:** 2012/12/21
(86) **N° demande PCT/PCT Application No.:** US 2011/041366
(87) **N° publication PCT/PCT Publication No.:** 2011/163311
(30) **Priorités/Priorities:** 2010/06/22 (US61/357,317); 2010/06/22 (US61/357,314)

(51) **Cl.Int./Int.Cl.** ***C12N 15/85*** (2006.01), ***A01K 67/027*** (2006.01), ***C07K 16/00*** (2006.01), ***C07K 16/46*** (2006.01)

(72) **Inventeurs/Inventors:**
MACDONALD, LYNN, US;
STEVENS, SEAN, US;
GURER, CAGAN, US;
MURPHY, ANDREW J., US;
HOSIAWA, KAROLINA A., US

(73) **Propriétaire/Owner:**
REGENERON PHARMACEUTICALS, INC., US

(74) **Agent:** BLAKE, CASSELS & GRAYDON LLP

(54) **Titre : SOURIS EXPRIMANT UNE CHAINE LEGERE INCLUANT DES REGIONS VARIABLES LAMBDA HUMAINES ET DES REGIONS CONSTANTES MURINES**
(54) **Title: MICE EXPRESSING A LIGHT CHAIN WITH HUMAN LAMBDA VARIABLE AND MOUSE CONSTANT REGIONS**



(57) **Abrégé/Abstract:**
Genetically modified mice are provided that express human λ variable (hVλ) sequences, including mice that express hVλ sequences from an endogenous mouse λ light chain locus, mice that express hVλ sequences from an endogenous mouse λ light chain locus, and mice that express hVλ sequences from a transgene or an episome wherein the hVλ sequence is linked to a mouse constant sequence. Mice are provided that are a source of somatically mutated human λ variable sequences useful for making antigen-binding proteins. Compositions and methods for making antigen-binding proteins that comprise human λ variable sequences, including human antibodies, are provided.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
29 December 2011 (29.12.2011)

PCT

(10) International Publication Number
WO 2011/163311 A1

(51) International Patent Classification:
C12N 15/85 (2006.01) *C07K 16/00* (2006.01)
A01K 67/027 (2006.01) *C07K 16/46* (2006.01)

(21) International Application Number:
PCT/US2011/041366

(22) International Filing Date:
22 June 2011 (22.06.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/357,314 22 June 2010 (22.06.2010) US
61/357,317 22 June 2010 (22.06.2010) US

(71) Applicant *(for all designated States except US)*: **REGENERON PHARMACEUTICALS, INC.** [US/US]; 777 Old Saw Mill River Road, Tarrytown, NY 10591 (US).

(72) Inventors; and

(75) Inventors/Applicants *(for US only)*: **MACDONALD, Lynn** [US/US]; 16 Gedney Way, White Plains, NY 10605 (US). **STEVENS, Sean** [US/US]; 355 Berry Street, #413, San Francisco, CA 94158 (US). **GURER, Cagan** [TR/US]; 8 Pamela Lane, Valhalla, NY 10595 (US). **MURPHY, Andrew, J.** [US/US]; 10 Newton Court, Croton-on-Hudson, NY 10520 (US). **HOSIAWA, Karolina, A.** [CA/US]; 14 Church Street, 2nd Floor, Tarrytown, NY 10591 (US).

(74) Agents: **POBURSKY, Kevin, J.** et al.; Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, NY 10591 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*
— *with amended claims (Art. 19(1))*
— *with sequence listing part of description (Rule 5.2(a))*

(54) Title: MICE EXPRESSING ALIGHT CHAIN WITH HUMAN LAMBDA VARIABLE AND MOUSE CONSTANT REGIONS



(57) Abstract: Genetically modified mice are provided that express human λ variable (hVλ) sequences, including mice that express hVλ sequences from an endogenous mouse λ light chain locus, mice that express hVλ sequences from an endogenous mouse κ light chain locus, and mice that express hVλ sequences from a transgene or an episome wherein the hVλ sequence is linked to a mouse constant sequence. Mice are provided that are a source of somatically mutated human λ variable sequences useful for making antigen- binding proteins. Compositions and methods for making antigen-binding proteins that comprise human λ variable sequences, including human antibodies, are provided.

WO 2011/163311 A1

MICE EXPRESSING A LIGHT CHAIN WITH HUMAN LAMBDA VARIABLE AND MOUSE CONSTANT REGIONS

FIELD

[1] Genetically modified mice that comprise a mouse or human lambda variable (Vλ,) light chain sequence operably linked with a mouse or human light chain constant region (λ or kappa(κ)). Genetically modified mice that express epitope-binding proteins that comprise an immunoglobulin light chain comprising a variable domain derived from a human lambda variable (hVλ) gene segment, a human lambda J (hVλ) gene segment, and a mouse light chain constant (C_L) domain. Genetically modified mice, comprising an unrearranged immunoglobulin lambda (λ) light chain variable nucleic acid sequence at an endogenous mouse light chain locus. Mice capable of rearranging and expressing a chimeric human λ/mouse C_L light chain from an endogenous light chain locus that comprises a replacement of all endogenous mouse light chain variable region gene segments with one or more hVλ, gene segments and one or more hJλ gene segments. Somatically mutated antibodies comprising hVλ domains and mouse C_L domains.

BACKGROUND

[2] Mice that express antibodies that are fully human, or partly human and partly mouse, are known in the art. For example, transgenic mice that express fully human antibodies from transgenes containing human light and heavy chain immunoglobulin variable region genes have been reported. Genetically modified mice that comprise a replacement of the endogenous mouse heavy chain variable region (HCVR) gene segments and kappa (κ) light chain variable region (LCVR) gene segments with human HCVR and LCVR gene segments and that make chimeric antibodies with a chimeric human/mouse kappa chain are known as well.

[3] Antibody light chains are encoded by one of two separate loci; kappa (κ) and lambda (λ). Mouse antibody light chains are primarily of the κ type. The ratio of κ to λ light chain usage in humans is about 60:40, whereas in mice it is about 95:5. Biased usage of κ light chains in mice is reportedly sustained in genetically modified mice capable of expressing fully or partly human antibodies. Thus, mice that express fully or partly human antibodies appear to be constrained in lambda variable usage.

[4] There is a need in the art to generate lambda variable regions, whether mouse or human, for use in making epitope-binding proteins. There is a need in the art for mice that express fully or partly human antibodies, wherein the mice display an increased lambda variable (Vλ) usage.

[0005] There is a need in the art for mice that express fully or partly human antibodies, wherein the mice display an increased λ variable (Vλ) usage.

SUMMARY

[0006] Genetically modified mice, embryos, cells, tissues, as well as nucleic acid constructs for modifying mice, and methods and compositions for making and using them, are provided. Mice and cells that generate lambda (λ) variable regions (human or non-human) in the context of a kappa (κ) light chain are provided. Mice and cells that generate human λ variable regions in the context of a κ or a λ light chain, *e.g.*, from an endogenous mouse light chain locus, are also provided. Also provided are methods for making antibodies that comprise lambda variable regions. Methods for selecting heavy chains that express with cognate lambda variable regions are also provided.

[0007] Chimeric and human antigen-binding proteins (*e.g.*, antibodies), and nucleic acids encoding them, are provided that comprise somatically mutated variable regions, including antibodies that have light chains comprising a variable domain derived from a human Vλ and a human Jλ gene segment fused to a mouse light chain constant domain.

[0008] In one aspect, a mouse is provided that expresses a human λ variable region sequence on a light chain that comprises a mouse constant region. In one aspect, a mouse is provided that expresses a human λ variable region sequence on a light chain that comprises a κ constant region. In one aspect, a mouse is provided that expresses from an endogenous mouse light chain locus a light chain that comprises a human λ variable region sequence. In one aspect, a mouse is provided that comprises a rearranged light chain gene that comprises a human λ variable sequence linked to a mouse constant region sequence; in one embodiment, the mouse constant region sequence is a λ constant sequence; in one embodiment, the mouse constant region sequence is a κ constant sequence.

[0009] In one aspect, a genetically modified mouse is provided, wherein the mouse comprises an unrearranged human λ light chain variable gene segment (hVλ) and a human λ joining gene segment (hJλ). In one embodiment, the unrearranged hVλ and hJλ are at a mouse light chain locus. In one embodiment, the unrearranged hVλ and unrearranged hJλ are on a transgene and operably linked to a human or mouse constant region sequence. In one embodiment, the unrearranged hVλ and unrearranged hJλ are on an episome. In one embodiment, the mouse is capable of making an immunoglobulin that comprises a light chain that is derived from an unrearranged hVλ sequence and a hJλ sequence and a mouse light chain constant region (C_L) nucleic acid sequence. Methods and compositions for making and using genetically modified mice are also provided.

Antibodies are provided that comprise (a) a human heavy chain variable domain (hV_H) fused to a mouse heavy chain constant region, and (b) a human Vλ fused to a mouse C_L domain; including wherein one or more of the variable domains are somatically mutated, *e.g.*, during antibody or immune cell selection in a mouse of the invention. In one embodiment, the unrearranged hVλ and unrearranged hJλ are operably linked with a human or mouse κ constant region (Cκ). In one embodiment, the unrearranged hVλ and unrearranged hJλ are operably linked with a human or mouse λ constant region (Cλ).

[0010] In one aspect, a mouse is provided that comprises in its germline, at an endogenous mouse light chain locus, a human λ light chain variable region sequence, wherein the human lambda variable region sequence is expressed in a light chain that comprises a mouse immunoglobulin constant region gene sequence.

[0011] In one embodiment, the endogenous mouse light chain locus is a λ locus. In one embodiment, the endogenous mouse light chain locus is a κ locus.

[0012] In one embodiment, the mouse lacks an endogenous light chain variable sequence at the endogenous mouse light chain locus.

[0013] In one embodiment, all or substantially all endogenous mouse light chain variable region gene segments are replaced with one or more human λ variable region gene segments.

[0014] In one embodiment, the human λ light chain variable region sequence comprises a human Jλ sequence. In one embodiment, the human Jλ sequence is selected from the group consisting of Jλ1, Jλ2, Jλ3, Jλ7, and a combination thereof.

[0015] In one embodiment, the human λ light chain variable region sequence comprises a fragment of cluster A of the human light chain locus. In a specific embodiment, the fragment of cluster A of the human λ light chain locus extends from hVλ3-27 through hVλ3-1.

[0016] In one embodiment, the human λ light chain variable region sequence comprises a fragment of cluster B of the human light chain locus. In a specific embodiment, the fragment of cluster B of the human λ light chain locus extends from hVλ5-52 through hVλ1-40.

[0017] In one embodiment, the human λ light chain variable region sequence comprises a genomic fragment of cluster A and a genomic fragment of cluster B. In a one embodiment, the human λ light chain variable region sequence comprises at least one gene segment of cluster A and at least one gene segment of cluster B.

[0018] In one embodiment, more than 10% of the light chain naïve repertoire of the mouse is derived from at least two hVλ gene segments selected from 2-8, 2-23, 1-40, 5-45, and 9-49. In one embodiment, more than 20% of the light chain naïve repertoire of the

mouse is derived from at least three hVλ gene segments selected from 2-8, 2-23, 1-40, 5-45, and 9-49. In one embodiment, more than 30% of the light chain naïve repertoire of the mouse is derived from at least four hVλ gene segments selected from 2-8, 2-23, 1-40, 5-45, and 9-49.

[0019] In one aspect, a mouse is provided that expresses an immunoglobulin light chain that comprises a human λ variable sequence fused with a mouse constant region, wherein the mouse exhibits a κ usage to λ usage ratio of about 1:1.

[0020] In one embodiment, the immunoglobulin light chain is expressed from an endogenous mouse light chain locus.

[0021] In one aspect, a mouse is provided that comprises a λ light chain variable region sequence (Vλ) and at least one J sequence (J), contiguous with a mouse κ light chain constant region sequence.

[0022] In one embodiment, the mouse lacks a functional mouse Vκ and/or mouse Jκ gene segment.

[0023] In one embodiment, the Vλ is a human Vλ (hVλ), and the J is a human Jλ (hJλ). In one embodiment, the hVλ and the hJλ are unrearranged gene segments.

[0024] In one embodiment, the mouse comprises a plurality of unrearranged hVλ gene segments and at least one hJλ gene segment. In a specific embodiment, the plurality of unrearranged hVλ gene segments are at least 12 gene segments, at least 28 gene segments, or at least 40 gene segments.

[0025] In one embodiment, the at least one hJλ gene segment is selected from the group consisting of Jλ1, Jλ2, Jλ3, Jλ7, and a combination thereof.

[0026] In one embodiment, an endogenous mouse λ light chain locus is deleted in whole or in part.

[0027] In one embodiment, the mouse κ light chain constant region sequence is at an endogenous mouse κ light chain locus.

[0028] In one embodiment, about 10% to about 45% of the B cells of the mouse express an antibody that comprises a light chain comprising a human λ light chain variable (Vλ) domain and a mouse κ light chain constant (Cκ) domain.

[0029] In one embodiment, the human λ variable domain is derived from a rearranged hVλ/hJλ sequence selected from the group consisting of 3-1/1, 3-1/7, 4-3/1, 4-3/7, 2-8/1, 3-9/1, 3-10/1, 3-10/3, 3-10/7, 2-14/1, 3-19/1, 2-23/1, 3-25/1, 1-40/1, 1-40/2, 1-40/3, 1-40/7, 7-43/1, 7-43/3, 1-44/1, 1-44/7, 5-45/1, 5-45/2, 5-45/7, 7-46/1, 7-46/2, 7-46/7, 9-49/1, 9-49/2, 9-49/7 and 1-51/1.

[0030] In one embodiment, the mouse further comprises a human Vκ-Jκ intergenic region from a human κ light chain locus, wherein the human Vκ-Jκ intergenic region is

contiguous with the Vλ sequence and the J sequence. In a specific embodiment, the human Vκ-Jκ intergenic region is placed between the Vλ sequence and the J sequence.

[0031] In one aspect, a mouse is provided that comprises (a) at least 12 to at least 40 unrearranged human λ light chain variable region gene segments and at least one human Jλ gene segment at an endogenous mouse light chain locus; (b) a human Vκ-Jκ intergenic sequence located between the at least 12 to at least 40 human light chain variable region gene segments and the at least one human Jλ sequence; wherein the mouse express an antibody that comprises a light chain comprising a human Vλ domain and a mouse Cκ domain.

[0032] In one aspect, a mouse is provided that expresses an antibody comprising a light chain that comprises a λ variable sequence and a κ constant sequence.

[0033] In one embodiment, the mouse exhibits a κ usage to λ usage ratio of about 1:1.

[0034] In one embodiment, a population of immature B cells obtained from bone marrow of the mouse exhibits a κ usage to λ usage ratio of about 1:1.

[0035] In one aspect, a genetically modified mouse is provided, wherein the mouse comprises an unrearranged immunoglobulin Vλ and a Jλ gene segment operably linked to a mouse light chain locus that comprises a mouse C_L gene.

[0036] In one embodiment, the Vλ and/or Jλ gene segments are human gene segments. In one embodiment, the Vλ and/or Jλ gene segments are mouse gene segments, and the C_L is a mouse Cκ.

[0037] In one embodiment, the endogenous mouse light chain locus is a κ light chain locus. In one embodiment, the endogenous mouse light chain locus is a λ light chain locus.

[0038] In one embodiment, the unrearranged Vλ and Jλ gene segments are at an endogenous mouse light chain locus.

[0039] In one embodiment, the unrearranged immunoglobulin Vλ and Jλ gene segments are on a transgene.

[0040] In one embodiment, the mouse further comprises a replacement of one or more heavy chain V, D, and/or J gene segments with one or more human V, D, and/or J gene segments at an endogenous mouse heavy chain immunoglobulin locus.

[0041] In one embodiment, the mouse comprises an unrearranged immunoglobulin Vλ and a Jλ gene segment at an endogenous mouse κ light chain locus that comprises a mouse Cκ gene.

[0042] In one embodiment, the mouse comprises an unrearranged human immunoglobulin λ light chain variable gene segment (Vλ) and a λ joining gene segment (Jλ) at an endogenous mouse λ light chain locus that comprises a mouse Cλ gene.

[0043] In one embodiment, the light chain variable gene locus (the "V_L locus") comprises at least one human Vλ (hVλ) gene segment. In one embodiment, the V_L locus comprises at least one human Jλ (hJλ) gene segment. In another embodiment, V_L locus comprises up to four hJλ gene segments. In one embodiment, the V_L locus comprises a contiguous sequence comprising human λ and human κ genomic sequence.

[0044] In one embodiment, the κ light chain variable gene locus (the "κ locus") comprises at least one human Vλ (hVλ) gene segment. In one embodiment, the κ locus comprises at least one human Jλ (hJλ) gene segment. In one embodiment, the κ locus comprises up to four hJλ gene segments. In one embodiment, the κ locus comprises at least one hVλ and at least one hJλ, and lacks or substantially lacks a functional Vκ region gene segment and lacks or substantially lacks a functional Jκ region gene segment. In one embodiment, the mouse comprises no functional Vκ region gene segment. In one embodiment, the mouse comprises no functional Jκ region gene segment.

[0045] In one embodiment, the λ light chain variable gene locus (the "λ locus") comprises at least one hVλ gene segment. In one embodiment, the λ locus comprises at least one human Jλ (hJλ) gene segment. In another embodiment, the λ locus comprises up to four hJλ gene segments.

[0046] In one embodiment, the V_L locus comprises a plurality of hVλs. In one embodiment, the plurality of hVλs are selected so as to result in expression of a λ light chain variable region repertoire that reflects about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, or about 90% or more of the Vλ usage observed in a human. In one embodiment, the V_L locus comprises gene segments hVλ 1-40, 1-44, 2-8, 2-14, 3-21, and a combination thereof.

[0047] In one embodiment, the hVλs include 3-1, 4-3, 2-8, 3-9, 3-10, 2-11, and 3-12. In a specific embodiment, the V_L locus comprises a contiguous sequence of the human λ light chain locus that spans from Vλ3-12 to Vλ3-1. In one embodiment, the V_L locus comprises at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 hVλs. In a specific embodiment, the hVλs include 3-1, 4-3, 2-8, 3-9, 3-10, 2-11, and 3-12. In a specific embodiment, the V_L locus comprises a contiguous sequence of the human λ locus that spans from Vλ3-12 to Vλ3-1. In one embodiment, the V_L locus is at the endogenous κ locus. In a specific embodiment, the V_L locus is at the endogenous κ locus and the endogenous λ light chain locus is deleted in part or completely. In one embodiment, the V_L locus is at the

endogenous λ locus. In a specific embodiment, the V_L locus is at the endogenous λ locus and the endogenous κ locus is deleted in part or completely.

[0048] In one embodiment, the V_L locus comprises 13 to 28 or more hVλs. In a specific embodiment, the hVλs include 2-14, 3-16, 2-18, 3-19, 3-21, 3-22, 2-23, 3-25, and 3-27. In a specific embodiment, the κ locus comprises a contiguous sequence of the human λ locus that spans from Vλ3-27 to Vλ3-1. In one embodiment, the V_L locus is at the endogenous κ locus. In a specific embodiment, the V_L locus is at the endogenous κ locus and the endogenous λ light chain locus is deleted in part or completely. In another embodiment, the V_L locus is at the endogenous λ locus. In a specific embodiment, the V_L locus is at the endogenous λ locus and the endogenous κ locus is deleted in part or completely.

[0049] In one embodiment, the V_L locus comprises 29 to 40 hVλs. In a specific embodiment, the κ locus comprises a contiguous sequence of the human λ locus that spans from Vλ3-29 to Vλ3-1, and a contiguous sequence of the human λ locus that spans from Vλ5-52 to Vλ1-40. In a specific embodiment, all or substantially all sequence between hVλ1-40 and hVλ3-29 in the genetically modified mouse consists essentially of a human λ sequence of approximately 959 bp found in nature (*e.g.*, in the human population) downstream of the hVλ1-40 gene segment (downstream of the 3' untranslated portion), a restriction enzyme site (*e.g.*, PI-SceI), followed by a human λ sequence of approximately 3,431 bp upstream of the hVλ3-29 gene segment found in nature. In one embodiment, the V_L locus is at the endogenous mouse κ locus. In a specific embodiment, the V_L locus is at the endogenous mouse κ locus and the endogenous mouse λ light chain locus is deleted in part or completely. In another embodiment, the V_L locus is at the endogenous mouse λ locus. In a specific embodiment, the V_L locus is at the endogenous mouse λ locus and the endogenous mouse κ locus is deleted in part or completely.

[0050] In one embodiment, the V_L locus comprises at least one hJλ. In one embodiment, the V_L locus comprises a plurality of hJλs. In one embodiment, the V_L locus comprises at least 2, 3, 4, 5, 6, or 7 hJλ. In a specific embodiment, the V_L locus comprises four hJλ. In a specific embodiment, the four hJλs are hJλ1, hJλ2, hJλ3, and hJλ7. In one embodiment, the V_L locus is a κ locus. In a specific embodiment, the V_L locus is at the endogenous κ locus and the endogenous λ light chain locus is deleted in part or completely. In one embodiment, the V_L locus comprises one hJλ. In a specific embodiment, the one hJλ is hJλ1. In one embodiment, the V_L locus is at the endogenous κ locus. In a specific embodiment, the V_L locus is at the endogenous κ locus and the endogenous λ light chain locus is deleted in part or completely. In another embodiment,

the V_L locus is at the endogenous λ locus. In a specific embodiment, the V_L locus is at the endogenous λ locus and the endogenous κ locus is deleted in part or completely.

[0051] In one embodiment, the V_L locus comprises at least one hVλ, at least one hJλ, and a mouse Cκ gene. In one embodiment, the V_L locus comprises at least one hVλ, at least one hJλ, and a mouse Cλ gene. In a specific embodiment, the mouse Cλ gene is Cλ2. In a specific embodiment, the mouse Cλ gene is at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, 96%, 97%, 98%, or at least 99% identical to mouse Cλ2.

[0052] In one embodiment, the mouse comprises a replacement at the endogenous mouse κ locus of endogenous mouse Vκ gene segments with one or more hVλ gene segments, wherein the hVλ gene segments are operably linked to an endogenous mouse Cκ region gene, such that the mouse rearranges the human Vλ gene segments and expresses a reverse chimeric immunoglobulin light chain that comprises a human Vλ domain and a mouse Cκ. In one embodiment, 90-100% of unrearranged mouse Vκ gene segments are replaced with at least one unrearranged hVλ gene segment. In a specific embodiment, all or substantially all of the endogenous mouse Vκ gene segments are replaced with at least one unrearranged hVλ gene segment. In one embodiment, the replacement is with at least 12, at least 28, or at least 40 unrearranged hVλ gene segments. In one embodiment, the replacement is with at least 7 functional unrearranged hVλ gene segments, at least 16 functional unrearranged hVλ gene segments, or at least 27 functional unrearranged hVλ gene segments. In one embodiment, the mouse comprises a replacement of all mouse Jκ gene segments with at least one unrearranged hJλ gene segment. In one embodiment, the at least one unrearranged hJλ gene segment is selected from Jλ1, Jλ2, Jλ3, Jλ4, Jλ5, Jλ6, Jλ7, and a combination thereof. In a specific embodiment, the one or more hVλ gene segment is selected from a 3-1, 4-3, 2-8, 3-9, 3-10, 2-11, 3-12, 2-14, 3-16, 2-18, 3-19, 3-21, 3-22, 2-23, 3-25, 3-27, 1-40, 7-43, 1-44, 5-45, 7-46, 1-47, 5-48, 9-49, 1-50, 1-51, a 5-52 hVλ gene segment, and a combination thereof. In a specific embodiment, the at least one unrearranged hJλ gene segment is selected from Jλ1, Jλ2, Jλ3, Jλ7, and a combination thereof.

[0053] In one embodiment, the mouse comprises a replacement of endogenous mouse Vλ gene segments at the endogenous mouse λ locus with one or more human Vλ gene segments at the endogenous mouse λ locus, wherein the hVλ gene segments are operably linked to a mouse Cλ region gene, such that the mouse rearranges the hVλ gene segments and expresses a reverse chimeric immunoglobulin light chain that comprises a hVλ domain and a mouse Cλ. In a specific embodiment, the mouse Cλ gene is Cλ2. In a specific embodiment, the mouse Cλ gene is at least 60%, at least 70%, at least 80%, at

least 90%, at least 95%, or at least 98% identical to mouse Cλ2. In one embodiment, 90-100% of unrearranged mouse Vλ gene segments are replaced with at least one unrearranged hVλ gene segment. In a specific embodiment, all or substantially all of the endogenous mouse Vλ gene segments are replaced with at least one unrearranged hVλ gene segment. In one embodiment, the replacement is with at least 12, at least 28, or at least 40 unrearranged hVλ gene segments. In one embodiment, the replacement is with at least 7 functional unrearranged hVλ gene segments, at least 16 functional unrearranged hVλ gene segments, or at least 27 functional unrearranged hVλ gene segments. In one embodiment, the mouse comprises a replacement of all mouse Jλ gene segments with at least one unrearranged hJλ gene segment. In one embodiment, the at least one unrearranged hJλ gene segment is selected from Jλ1, Jλ2, Jλ3, Jλ4, Jλ5, Jλ6, Jλ7, and a combination thereof. In a specific embodiment, the one or more hVλ gene segment is selected from a 3-1, 4-3, 2-8, 3-9, 3-10, 2-11, 3-12, 2-14, 3-16, 2-18, 3-19, 3-21, 3-22, 2-23, 3-25, 3-27, 1-40, 7-43, 1-44, 5-45, 7-46, 1-47, 5-48, 9-49, 1-50, 1-51, a 5-52 hVλ gene segment, and a combination thereof. In a specific embodiment, the at least one unrearranged hJλ gene segment is selected from Jλ1, Jλ2, Jλ3, Jλ7, and a combination thereof.

[0054] In one aspect, a genetically modified mouse is provided that comprises a human Vκ-Jκ intergenic region sequence located at an endogenous mouse κ light chain locus.

[0055] In one embodiment, the human Vκ-Jκ intergenic region sequence is at an endogenous κ light chain locus of a mouse that comprises a hVλ and hJλ gene segment, and the human Vκ-Jκ intergenic region sequence is disposed between the hVλ and hJλ gene segments. In a specific embodiment, the hVλ and hJλ gene segments are capable of recombining to form a functional human λ light chain variable domain in the mouse.

[0056] In one embodiment, a mouse is provided that comprises a plurality of hVλ's and one or more hJλ's, and the human Vκ-Jκ intergenic region sequence is disposed, with respect to transcription, downstream of the proximal or 3' most hVλ sequence and upstream or 5' of the first hJλ sequence.

[0057] In one embodiment, the human Vκ-Jκ intergenic region is a region located about 130 bp downstream or 3' of a human Vκ4-1 gene segment, about 130 bp downstream of the 3' untranslated region of the human Vκ4-1 gene segment, and spans to about 600 bp upstream or 5' of a human Jκ1 gene segment. In a specific embodiment, the human Vκ-Jκ intergenic region is about 22.8 kb in size. In one embodiment, the Vκ-Jκ intergenic region is about 90% or more, 91% or more, 92% or more, 93% or more, 94% or

more, or about 95% or more identical with a human Vκ-Jκ intergenic region extending from the end of the 3' untranslated region of a human Vκ4-1 gene segment to about 600 bp upstream of a human Jκ1 gene segment. In one embodiment, the Vκ-Jκ intergenic region comprises SEQ ID NO:100. In a specific embodiment, the Vκ-Jκ intergenic region comprises a functional fragment of SEQ ID NO:100. In a specific embodiment, the Vκ-Jκ intergenic region is SEQ ID NO:100.

[0058] In one aspect, a mouse, a mouse cell (*e.g.*, a mouse embryonic stem cell), a mouse embryo, and a mouse tissue are provided that comprise the recited human Vκ-Jκ intergenic region sequence, wherein the intergenic region sequence is ectopic. In a specific embodiment, the ectopic sequence is placed at a humanized endogenous mouse immunoglobulin locus.

[0059] In one aspect, an isolated nucleic acid construct is provided that comprises the recited human Vκ-Jκ intergenic region sequence. In one embodiment, the nucleic acid construct comprises targeting arms to target the human Vκ-Jκ intergenic region sequence to a mouse light chain locus. In a specific embodiment, the mouse light chain locus is a κ locus. In a specific embodiment, the targeting arms target the human Vκ-Jκ intergenic region to a modified endogenous mouse κ locus, wherein the targeting is to a position between a hVλ sequence and a hJλ sequence.

[0060] In one aspect, a genetically modified mouse is provided, wherein the mouse comprises no more than two light chain alleles, wherein the light chain alleles comprise (a) an unrearranged immunoglobulin human Vλ and a Jλ gene segment at an endogenous mouse light chain locus that comprises a mouse C_L gene; and, (b) an unrearranged immunoglobulin V_L and a J_L gene segment at an endogenous mouse light chain locus that comprises a mouse C_L gene.

[0061] In one embodiment, the endogenous mouse light chain locus is a κ locus. In another embodiment, the endogenous mouse light chain locus is a λ locus.

[0062] In one embodiment, the no more than two light chain alleles are selected from a κ allele and a λ allele, two κ alleles, and two λ alleles. In a specific embodiment, one of the two light chain alleles is a λ allele that comprises a Cλ2 gene.

[0063] In one embodiment, the mouse comprises one functional immunoglobulin light chain locus and one nonfunctional light chain locus, wherein the functional light chain locus comprises an unrearranged immunoglobulin human Vλ and a Jλ gene segment at an endogenous mouse κ light chain locus that comprises a mouse Cκ gene.

[0064] In one embodiment, the mouse comprises one functional immunoglobulin light chain locus and one nonfunctional light chain locus, wherein the functional light chain locus comprises an unrearranged immunoglobulin human Vλ and a Jλ gene segment at an

endogenous mouse λ light chain locus that comprises a mouse Cλ gene. In one embodiment, the Cλ gene is Cλ2. In a specific embodiment, the mouse Cλ gene is at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% identical to mouse Cλ2.

[0065] In one embodiment, the mouse further comprises at least one immunoglobulin heavy chain allele. In one embodiment, the at least one immunoglobulin heavy chain allele comprises a human V_H gene segment, a human D_H gene segment, and a human J_H gene segment at an endogenous mouse heavy chain locus that comprises a human heavy chain gene that expresses a human/mouse heavy chain. In a specific embodiment, the mouse comprises two immunoglobulin heavy chain alleles, and the mouse expresses a human/mouse heavy chain.

[0066] In one embodiment, the mouse comprises a first light chain allele that comprises an unrearranged hVκ and an unrearranged hJκ, at an endogenous mouse κ locus that comprises an endogenous Cκ gene; and a second light chain allele that comprises an unrearranged hVλ and an unrearranged hJλ, at an endogenous mouse κ locus that comprises an endogenous Cκ gene. In a specific embodiment, the first and the second light chain alleles are the only functional light chain alleles of the genetically modified mouse. In a specific embodiment, the mouse comprises a nonfunctional λ locus. In one embodiment, the genetically modified mouse does not express a light chain that comprises a λ constant region.

[0067] In one embodiment, the mouse comprises a first light chain allele that comprises an unrearranged hVκ and an unrearranged hJκ, at an endogenous mouse κ locus that comprises an endogenous Cκ gene; and a second light chain allele that comprises an unrearranged hVλ and an unrearranged hJλ, at an endogenous mouse λ locus that comprises an endogenous Cλ gene. In a specific embodiment, the first and the second light chain alleles are the only functional light chain alleles of the genetically modified mouse. In one embodiment, the endogenous Cλ gene is Cλ2. In a specific embodiment, the mouse Cλ gene is at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% identical to mouse Cλ2.

[0068] In one embodiment, the mouse comprises six immunoglobulin alleles, wherein the first allele comprises an unrearranged immunoglobulin Vλ and Jλ gene segment at an endogenous mouse κ light chain locus that comprises a mouse Cκ gene, the second comprises an unrearranged immunoglobulin Vκ and Jκ gene segment at an endogenous mouse κ light chain locus that comprises a mouse Cκ gene, the third comprises an unrearranged immunoglobulin Vλ and Jλ gene segment at an endogenous mouse λ light

chain locus that comprises a mouse Cλ gene, the fourth and fifth each independently comprise an unrearranged V_H and D_H and J_H gene segment at an endogenous mouse heavy chain locus that comprises a mouse heavy chain gene, and the sixth comprises either (a) an unrearranged immunoglobulin Vλ and Jλ gene segment at an endogenous mouse λ light chain locus that comprises a mouse Cλ gene, (b) a λ locus that is nonfunctional, or (c) a deletion in whole or in part of the λ locus.

[0069] In one embodiment, the first allele comprises an unrearranged hVλ and hJλ. In one embodiment, the second allele comprises an unrearranged hVκ and hJκ. In one embodiment, the third allele comprises an unrearranged hVλ and hJλ. In one embodiment, the fourth and fifth each independently comprise an unrearranged hV_H and hD_H and hJ_H. In one embodiment, the sixth allele comprises an endogenous mouse λ locus that is deleted in whole or in part.

[0070] In one embodiment, the mouse comprises six immunoglobulin alleles, wherein the first allele comprises an unrearranged immunoglobulin Vλ and Jλ gene segment at an endogenous mouse λ light chain locus that comprises a mouse Cλ gene, the second comprises an unrearranged immumoglobulin Vλ and Jλ gene segment at an endogenous mouse λ light chain locus that comprises a mouse Cλ gene, the third comprises an unrearranged immunoglobulin Vκ and Jκ gene segment at an endogenous mouse κ light chain locus that comprises a mouse Cκ gene, the fourth and fifth each independently comprise an unrearranged V_H and D_H and J_H gene segment at an endogenous mouse heavy chain locus that comprises a mouse heavy chain gene, and the sixth comprises either (a) an unrearranged immunoglobulin Vκ and Jκ gene segment at an endogenous mouse κ light chain locus that comprises a mouse Cκ gene, (b) a κ locus that is nonfunctional, or (c) a deletion of one or more elements of the κ locus.

[0071] In one embodiment, the first allele comprises an unrearranged hVλ and hJλ gene segment. In one embodiment, the second allele comprises an unrearranged hVλ and hJλ gene segment. In one embodiment, the third allele comprises an unrearranged hVκ and hJκ gene segment. In one embodiment, the fourth and fifth each independently comprise an unrearranged hV_H and hD_H and hJ_H gene segment. In one embodiment, the sixth allele comprises an endogenous mouse κ locus that is functionally silenced.

[0072] In one embodiment, the genetically modified mouse comprises a B cell that comprises a rearranged antibody gene comprising a rearranged hVλ domain operably linked to a mouse C_L domain. In one embodiment, the mouse C_L domain is selected from a mouse Cκ and a mouse Cλ domain. In a specific embodiment, the mouse Cλ domain is derived from a Cλ2 gene. In a specific embodiment, the mouse Cλ domain is derived from

a Cλ domain that is at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% identical to mouse Cλ2.

[0073] In one aspect, a genetically modified mouse is provided that expresses a Vλ region on a C_L that is a Cκ. In one aspect, a genetically modified mouse is provided that expresses a hVλ region on a C_L selected from a human Cκ, a human Cλ, or a mouse Cκ. In one aspect, a genetically modified mouse is provided that expresses a hVλ region on a mouse Cκ.

[0074] In one embodiment, about 10-50% of the splenocytes of the mouse are B cells (*i.e.*, CD19-positive), or which about 9-28% express an immunoglobulin light chain comprising a hVλ domain fused to a mouse Cκ domain.

[0075] In a specific embodiment, about 23-34% of the splenocytes of the mouse are B cells (*i.e.*, CD19-positive), or which about 9-11% express an immunoglobulin light chain comprising a hVλ domain fused to a mouse Cκ domain.

[0076] In a specific embodiment, about 19-31% of the splenocytes of the mouse are B cells (*i.e.*, CD19-positive), or which about 9-17% express an immunoglobulin light chain comprising a hVλ domain fused to a mouse Cκ domain.

[0077] In a specific embodiment, about 21-38% of the splenocytes of the mouse are B cells (*i.e.*, CD19-positive), or which about 24-27% express an immunoglobulin light chain comprising a hVλ domain fused to a mouse Cκ domain.

[0078] In a specific embodiment, about 10-14% of the splenocytes of the mouse are B cells (*i.e.*, CD19-positive), or which about 9-13% express an immunoglobulin light chain comprising a hVλ domain fused to a mouse Cκ domain.

[0079] In a specific embodiment, about 31-48% of the splenocytes of the mouse are B cells *(i.e.*, CD19-positive), or which about 15-21% express an immunoglobulin light chain comprising a hVλ domain fused to a mouse Cκ domain. In a specific embodiment, about 30-38% of the splenocytes of the mouse are B cells (*i.e.*, CD19-positive), of which about 33-48% express an immunoglobulin light chain comprising a hVλ domain fused to a mouse Cκ domain.

[0080] In one embodiment, about 52-70% of the bone marrow of the mouse are B cells (*i.e.*, CD19-positive), or which about 31-47% of the immature B cells (*i.e.*, CD19-positive/B220-intermediate positive/IgM-positive) express an immunoglobulin light chain comprising a hVλ domain fused to a mouse Cκ domain.

[0081] In one embodiment, about 60% of the bone marrow of the mouse are B cells (*i.e.*, CD19-positive), or which about 38.3% of the immature B cells (*i.e.*, CD19-positive/B220-intermediate positive/IgM-positive) express an immunoglobulin light chain comprising a hVλ domain fused to a mouse Cκ domain.

[0082] In one embodiment, the mouse expresses an antibody comprising a light chain that comprises a variable domain derived from a human V and a human J gene segment, and a constant domain derived from a mouse constant region gene. In one embodiment, the mouse constant region gene is a Cκ gene. In another embodiment, the mouse constant region gene is a Cλ gene. In a specific embodiment, the Cλ region is Cλ2. In a specific embodiment, the mouse Cλ gene is derived from a Cλ gene that is at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% identical to mouse Cλ2. In a specific embodiment, the antibody further comprises a heavy chain comprising a variable domain derived from a human V, a human D and a human J gene segment, and a heavy chain constant domain derived from a mouse heavy chain constant region gene. In one embodiment, the mouse heavy chain constant region gene comprises a hinge-CH_2-CH_3 sequence of a heavy chain constant domain. In another embodiment, the mouse heavy chain constant region gene comprises a CH_1-hinge-CH_2-CH_3 sequence of a heavy chain constant domain. In another embodiment, the mouse heavy chain constant region gene comprises a CH_1-CH_2-CH_3-CH_4 sequence of a heavy chain constant domain. In another embodiment, the mouse heavy chain constant region gene comprises a CH_2-CH_3-CH_4 sequence of a heavy chain constant domain.

[0083] In one embodiment, the mouse expresses an antibody comprising a light chain that comprises a rearranged human Vλ-Jλ sequence and a mouse Cκ sequence. In one embodiment, the rearranged human Vλ-Jλ sequence is derived from a rearrangement of hVλ gene segments selected from a 3-1, 4-3, 2-8, 3-9, 3-10, 2-14, 3-19, 2-23, 3-25, 1-40, 7-43, 1-44, 5-45, 7-46, 1-47, 9-49, and a 1-51 gene segment. In one embodiment, the rearranged human Vλ-Jλ sequence is derived from a rearrangement of hJλ gene segments selected from Jλ1, Jλ2, Jλ3, and a Jλ7 gene segment.

[0084] In one embodiment, the mouse expresses an antibody comprising a light chain that comprises a rearranged immunoglobulin λ light chain variable region comprising a human VλJλ sequence selected from 3-1/1, 3-1/7, 4-3/1, 4-3/7, 2-8/1, 3-9/1, 3-10/1, 3-10/3, 3-10/7, 2-14/1, 3-19/1, 2-23/1, 3-25/1, 1-40/1, 1-40/2, 1-40/3, 1-40/7, 7-43/1, 7-43/3, 1-44/1, 1-44/7, 5-45/1, 5-45/2, 5-45/7, 7-46/1, 7-46/2, 7-46/7, 9-49/1, 9-49/2, 9-49/7 and 1-51/1. In a specific embodiment, the B cell expresses an antibody comprising a human immunoglobulin heavy chain variable domain fused with a mouse heavy chain constant domain, and a human immunoglobulin λ light chain variable domain fused with a mouse κ light chain constant domain.

[0085] In one aspect, a mouse is provided that expresses an antibody comprising (a) a heavy chain comprising a heavy chain variable domain derived from an unrearranged human heavy chain variable region gene segment, wherein the heavy chain variable

domain is fused to a mouse heavy chain constant (C_H) region; and, (b) a light chain comprising a light chain variable domain derived from an unrearranged hVλ and a hJλ, wherein the light chain variable domain is fused to a mouse C_L region.

[0086] In one embodiment, the mouse comprises (i) a heavy chain locus that comprises a replacement of all or substantially all functional endogenous mouse V, D and J gene segments with all or substantially all functional human V, D, and J gene segments, a mouse C_H gene, (ii) a first κ light chain locus comprising a replacement of all or substantially all functional endogenous mouse Vκ and Jκ gene segments with all, substantially all, or a plurality of, functional hVλ and hJλ gene segments, and a mouse Cκ gene, (iii) a second κ light chain locus comprising a replacement of all or substantially all functional endogenous mouse Vκ and Jκ gene segments with all, substantially all, or a plurality of, functional hVκ and hJκ gene segments, and a mouse Cκ gene. In one embodiment, the mouse does not express an antibody that comprises a Cλ region. In one embodiment, the mouse comprises a deletion of a Cλ gene and/or a Vλ and/or a Jλ gene segment. In one embodiment, the mouse comprises a nonfunctional λ light chain locus. In a specific embodiment, the λ light chain locus is deleted in whole or in part.

[0087] In one embodiment, the mouse comprises (i) a heavy chain locus that comprises a replacement of all or substantially all functional endogenous mouse V, D and J gene segments with all or substantially all functional human V, D, and J gene segments, a mouse C_H gene, (ii) a first λ light chain locus comprising a replacement of all or substantially all functional endogenous mouse Vλ and Jλ gene segments with all, substantially all, or a plurality of, functional hVλ and hJλ gene segments, and a mouse Cλ gene, (iii) a second λ light chain locus comprising a replacement of all or substantially all functional endogenous mouse Vλ and Jλ gene segments with all, substantially all, or a plurality of, functional hVλ and hJλ gene segments, and a mouse Cλ gene. In a specific embodiment, the mouse Cλ gene is Cλ2. In a specific embodiment, the mouse Cλ gene is derived from a Cλ gene that is at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% identical to mouse Cλ2.

[0088] In one embodiment, the mouse comprises a deletion of a Cκ gene and/or a Vκ and/or a Jκ gene segment. In one embodiment, the mouse comprises a nonfunctional κ light chain locus.

[0089] In one aspect, a genetically modified mouse that expresses an antibody is provided, wherein greater than 10%, greater than 15%, greater than 20%, greater than 25%, greater than 30%, greater than 35%, greater than 40%, greater than 60%, greater than 70%, greater than 80%, or greater than 90% of total IgG antibody produced by the

mouse comprises a λ-derived variable domain, and wherein the mouse expresses antibodies comprising a κ-derived variable domain fused with a mouse Cκ region. In specific embodiments, about 15-40%, 20-40%, 25-40%, 30-40%, or 35-40% of total antibody produced by the mouse comprises a λ-derived variable domain.

[0090] In one embodiment, the λ-derived variable domain is derived from a hVλ and a hJλ. In one embodiment, the λ-derived variable domain is in a light chain that comprises a mouse Cκ region. In a specific embodiment, the λ-derived variable region is in a light chain that comprises a mouse Cλ region. In another specific embodiment, the Cλ region is a Cλ2 region. In one embodiment, the κ-derived variable domain is derived from a hVκ and a hJκ, and in a specific embodiment is in a light chain that comprises a mouse Cκ region.

[0091] In one aspect, an isolated DNA construct is provided that comprises an upstream homology arm and a downstream homology arm, wherein the upstream and the downstream homology arms target the construct to a mouse κ locus, and the construct comprises a functional unrearranged hVλ segment and a functional unrearranged hJλ segment, and a selection or marker sequence.

[0092] In one aspect, an isolated DNA construct is provided, comprising, from 5' to 3' with respect to the direction of transcription, a targeting arm for targeting a mouse λ sequence upstream of mouse Vλ2, a selection cassette flanked 5' and 3' with recombinase recognition sites, and a targeting arm for targeting a mouse λ sequence 3' of mouse Jλ2. In one embodiment, the selection cassette is a Frt'ed Hyg-TK cassette. In one embodiment, the 3' targeting arm comprises mouse Cλ2, Jλ4, Cλ4, and mouse enhancer 2.4.

[0093] In one aspect, an isolated DNA construct is provided, comprising, from 5' to 3' with respect to the direction of transcription, a targeting arm for targeting the mouse λ locus 5' with respect to Vλ1, a selection cassette flanked 5' and 3' with recombinase recognition sites, and a 3' targeting arm for targeting a mouse λ sequence 3' with respect to mouse Cλ1. In one embodiment, the selection cassette is a loxed neomycin cassette. In one embodiment, the 3' targeting arm comprises the mouse λ 3' enhancer and mouse λ 3' enhancer 3.1.

[0094] In one aspect, an isolated DNA construct is provided, comprising from 5' to 3' with respect to the direction of transcription, a targeting arm for targeting the mouse λ locus 5' with respect to Vλ2, a selection cassette flanked 5' and 3' with recombinase recognition sites, and a 3' targeting arm for targeting a mouse λ sequence 3' with respect to mouse Jλ2 and 5' with respect to mouse Cλ2. In one embodiment, the selection cassette is a

Frt'ed hygromycin-TK cassette. In one embodiment, the 3' targeting arm comprises the mouse Cλ2-Jλ4-Cλ4 gene segments and mouse λ enhancer 2.4.

[0095] In one aspect, an isolated DNA construct is provided, comprising, from 5' to 3' with respect to the direction of transcription, a targeting arm for targeting the mouse λ locus 5' with respect to Vλ2, a selection cassette flanked 5' and 3' with recombinase recognition sites, a human genomic fragment comprising a contiguous region of the human λ light chain locus from hVλ3-12 downstream to the end of hJλ1, and a 3' targeting arm for targeting a mouse λ sequence 3' with respect to mouse Jλ2. In one embodiment, the selection cassette is a Frt'ed neomycin cassette. In one embodiment, the 3' targeting arm comprises the mouse Cλ2-Jλ4-Cλ4 gene segments and mouse λ enhancer 2.4.

[0096] In one aspect, an isolated DNA construct is provided, comprising a contiguous region of the human λ light chain locus from hVλ3-12 downstream to the end of hJλ1.

[0097] In one aspect, an isolated DNA construct is provided, comprising, from 5' to 3' with respect to the direction of transcription, a targeting arm for targeting the mouse λ locus 5' with respect to Vλ2, a selection cassette flanked 5' and 3' with recombinase recognition sites and a human genomic fragment comprising a contiguous region of the human λ light chain locus from hVλ3-27 downstream to the end of hVλ2-8. In one embodiment, the selection cassette is a Frt'ed hygromycin cassette. In one embodiment, the human genomic fragment comprises a 3' targeting arm. In a specific embodiment, the 3' targeting arm comprises about 53 kb of the human λ light chain locus from hVλ3-12 downstream to the end of hVλ2-8.

[0098] In one aspect, an isolated DNA construct is provided, comprising a contiguous region of the human λ light chain locus from hVλ3-27 downstream to the end of hVλ3-12.

[0099] In one aspect, an isolated DNA construct is provided, comprising, from 5' to 3' with respect to the direction of transcription, a targeting arm for targeting the mouse λ locus 5' with respect to Vλ2, a selection cassette flanked 5' and 3' with recombinase recognition sites, a first human genomic fragment comprising a contiguous region of the human λ light chain locus from hVλ5-52 downstream to the end of hVλ1-40, a restriction enzyme site, and a second human genomic fragment comprising a contiguous region of the human λ light chain locus from hVλ3-29 downstream to the end of hVλ82K. In one embodiment, the selection cassette is a Frt'ed neomycin cassette. In one embodiment, the restriction enzyme site is a site for a homing endonuclease. In a specific embodiment, the homing endonuclease is PI-SceI. In on embodiment, the second human genomic fragment is a 3' targeting arm. In a specific embodiment, the 3' targeting arm comprises about 27 kb of the human λ light chain locus from hVλ3-29 downstream to the end of hVλ82K.

[00100] In one aspect, an isolated DNA construct is provided, comprising a contiguous region of the human λ light chain locus from hVλ5-52 downstream to the end of hVλ1-40.

[00101] In one aspect, an isolated DNA construct is provided, comprising, from 5' to 3' with respect to the direction of transcription, a targeting arm for targeting the mouse κ locus 5' with respect to the endogenous Vκ gene segments, two juxtaposed recombinase recognition sites, a selection cassette 3' to the juxtaposed recombinase recognition sites, and a 3' targeting arm for targeting a mouse κ sequence 5' with respect to the κ light chain variable gene segments. In one embodiment, the juxtaposed recombinase recognition sites are in opposite orientation with respect to one another. In a specific embodiment, the recombinase recognition sites are different. In another specific embodiment, the recombinase recognition sites are a *lox*P site and a *lox*511 site. In one embodiment, the selection cassette is a neomycin cassette.

[00102] In one aspect, an isolated DNA construct is provided, comprising, from 5' to 3' with respect to the direction of transcription, a targeting arm for targeting the mouse κ locus 5' with respect to the mouse Jκ gene segments, a selection cassette, a recombinase recognition site 3' to the selection cassette, and a 3' targeting arm for targeting a mouse κ sequence 3' with respect to the mouse Jκ gene segments and 5' to the mouse κ intronic enhancer. In one embodiment, the selection cassette is a hygromycin-TK cassette. In one embodiment, the recombinase recognition site is in the same direction with respect to transcription as the selection cassette. In a specific embodiment, the recombinase recognition site is a *lox*P site.

[00103] In one aspect, an isolated DNA construct is provided, comprising, from 5' to 3' with respect to the direction of transcription, a first mouse genomic fragment comprising sequence 5' of the endogenous mouse Vκ gene segments, a first recombinase recognition site, a second recombinase recognition site, and a second mouse genomic fragment comprising sequence 3' of the endogenous mouse Jκ gene segments and 5' of the mouse κ intronic enhancer.

[00104] In one aspect, a genetically modified mouse is provided, wherein the genetic modification comprises a modification with one or more of the DNA constructs described above or herein.

[00105] In one aspect, use of an isolated DNA construct to make a mouse as described herein is provided. In one aspect, use of an isolated DNA construct as described herein in a method for making an antigen-binding protein is provided.

[00106] In one aspect, a non-human stem cell is provided that comprises a targeting vector that comprises a DNA construct as described above and herein. In one aspect, a

non-human stem cell is provided, wherein the non-human stem cell is derived from a mouse described herein.

[00107] In one embodiment, the non-human stem cell is an embryonic stem (ES) cell. In a specific embodiment, the ES cell is a mouse ES cell.

[00108] In one aspect, use of a non-human stem cell as described herein to make a mouse as described herein is provided. In one aspect, use of a non-human stem cell as described herein to make an antigen-binding protein is provided.

[00109] In one aspect, a mouse embryo is provided, wherein the mouse embryo comprises a genetic modification as provided herein. In one embodiment, a host mouse embryo that comprises a donor ES cell is provided, wherein the donor ES cell comprises a genetic modification as described herein. In one embodiment, the mouse embryo is a pre-morula stage embryo. In a specific embodiment, the pre-morula stage embryo is a 4-cell stage embryo or an 8-cell stage embryo. In another specific embodiment, the mouse embryo is a blastocyst.

[00110] In one aspect, use of a mouse embryo as described herein to make a mouse as described herein is provided. In one aspect, use of a mouse embryo as described herein to make an antigen-binding protein is provided.

[00111] In one aspect, a non-human cell is provided, wherein the non-human cell comprises a rearranged immunoglobulin light chain gene sequence derived from a genetically modified mouse as described herein. In one embodiment, the cell is a B cell. In one embodiment, the cell is a hybridoma. In one embodiment, the cell encodes an immunoglobulin light chain variable domain and/or an immunoglobulin heavy chain variable domain that is somatically mutated.

[00112] In one aspect, a non-human cell is provided, wherein the non-human cell comprises a rearranged immunoglobulin light chain gene sequence derived from a genetically modified mouse as described herein. In one embodiment, the cell is a B cell. In one embodiment, the cell is a hybridoma. In one embodiment, the cell encodes an immunoglobulin light chain variable domain and/or an immunoglobulin heavy chain variable domain that is somatically mutated.

[00113] In one aspect, use of a non-human cell as described herein to make a mouse as described herein is provided. In one aspect, use of a non-human cell as described herein to make an antigen-binding protein is provided.

[00114] In one aspect, a mouse B cell is provided that expresses an immunoglobulin light chain that comprises (a) a variable region derived from a hVλ gene segment and a hJλ gene segment; and, (b) a mouse C_L gene. In one embodiment, the mouse C_L gene is selected from a Cκ and a Cλ gene. In a specific embodiment, the Cλ gene is Cλ2. In a

specific embodiment, the mouse Cλ gene is derived from a Cλ gene that is at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% identical to mouse Cλ2. In one embodiment, the mouse B cell further expresses a cognate heavy chain that comprises (c) a variable region derived from a hV_H, a hD_H, and (d) a hJ_H segment. In one embodiment, the B cell does not comprise a rearranged λ gene. In another embodiment, the B cell does not comprise a rearranged κ gene.

[00115] In one aspect, a method for making an antibody in a genetically modified mouse is provided, comprising: (a) exposing a genetically modified mouse to an antigen, wherein the mouse has a genome comprising at least one hVλ and at least one hJλ at an endogenous light chain locus, wherein the endogenous light chain locus comprises a mouse C_L gene; (b) allowing the genetically modified mouse to develop an immune response to the antigen; and, (c) isolating from the mouse of (b) an antibody that specifically recognizes the antigen, or isolating from the mouse of (b) a cell comprising an immunoglobulin domain that specifically recognizes the antigen, wherein the antibody comprises a light chain derived from a hVλ, a hJλ and a mouse C_L gene. In a specific embodiment, the mouse C_L gene is a mouse Cκ gene.

[00116] In one embodiment, a method for making an antibody in a genetically modified mouse is provided, comprising: (a) exposing a genetically modified mouse to an antigen, wherein the mouse has a genome comprising at least one hVλ at an endogenous κ locus and at least one hJλ at the κ locus, wherein the κ locus comprises a mouse Cκ gene; (b) allowing the genetically modified mouse to develop an immune response to the antigen; and, (c) isolating from the mouse of (b) an antibody that specifically recognizes the antigen, or isolating from the mouse of (b) a cell comprising an immunoglobulin domain that specifically recognizes the antigen, wherein the antibody comprises a light chain derived from a hVλ, a hJλ and a mouse Cκ gene.

[00117] In one embodiment, the κ light chain constant gene is selected from a human Cκ gene and a mouse Cκ gene.

[00118] In one embodiment, a method for making an antibody in a genetically modified mouse is provided, comprising: (a) exposing a genetically modified mouse to an antigen, wherein the mouse has a genome comprising at least one hVλ at a λ light chain locus and at least one Jλ at the λ light chain locus, wherein the λ light chain locus comprises a mouse Cλ gene; (b) allowing the genetically modified mouse to develop an immune response to the antigen; and, (c) isolating from the mouse of (b) an antibody that specifically recognizes the antigen, or isolating from the mouse of (b) a cell comprising an immunoglobulin domain that specifically recognizes the antigen, or identifying in the mouse of B a nucleic acid sequence encoding a heavy and/or light chain variable domain that

binds the antigen, wherein the antibody comprises a light chain derived from a hVλ, a hJλ and a mouse Cλ gene.

[00119] In one embodiment, the λ light chain constant gene is selected from a human Cλ gene and a mouse Cλ gene. In one embodiment, the λ light chain constant gene is a human Cλ gene. In a specific embodiment, the human Cλ gene is selected from Cλ1, Cλ2, Cλ3 and Cλ7. In one embodiment, the λ light chain constant gene is a mouse Cλ gene. In a specific embodiment, the mouse Cλ gene is selected from Cλ1, Cλ2 and Cλ3. In a more specific embodiment, the mouse Cλ gene is Cλ2. In another specific embodiment, the mouse Cλ gene is derived from a Cλ gene that is at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% identical to mouse Cλ2.

[00120] In one aspect, a method for making a rearranged antibody gene in a genetically modified mouse is provided, comprising: (a) exposing a genetically modified mouse to an antigen, wherein the genetic modification comprises a hVλ and a hJλ at an endogenous light chain locus, wherein the endogenous light chain locus comprises a mouse C_L gene or functional fragment thereof; and, (b) identifying a rearranged immunoglobulin gene in said mouse, wherein the rearranged immunoglobulin gene comprises a λ light chain variable region gene segment and a C_L gene or functional fragment thereof.

[00121] In one embodiment, the method further comprises cloning a nucleic acid sequence encoding a heavy and/or light chain variable region from the mouse, wherein the heavy and/or light chain variable region is from an antibody that comprises a human Vλ and a mouse C_L.

[00122] In one embodiment, the mouse C_L gene or functional fragment thereof is selected from a human C_L gene and a mouse C_L gene, or functional fragment thereof.

[00123] In one embodiment, a method for making a rearranged antibody gene in a genetically modified mouse is provided, comprising: (a) exposing a genetically modified mouse to an antigen, wherein the genetic modification comprises a hVλ and a hJλ at a κ light chain locus, wherein the κ light chain locus comprises a mouse Cκ gene or functional fragment thereof; and, (b) identifying a rearranged immunoglobulin gene in said mouse, wherein the rearranged immunoglobulin gene comprises a λ light chain variable region gene segment and a Cκ gene or functional fragment thereof.

[00124] In one embodiment, the κ light chain constant gene or functional fragment thereof is selected from a human Cκ gene and a mouse Cκ gene, or a functional fragment thereof.

[00125] In one embodiment, the method further comprises cloning a nucleic acid sequence encoding a heavy and/or light chain variable region from the mouse, wherein the

heavy and/or light chain variable region is from an antibody that comprises a human Vλ and a mouse Cκ.

[00126] In one embodiment, a method for making a rearranged antibody gene in a genetically modified mouse is provided, comprising: (a) exposing a genetically modified mouse to an antigen, wherein the genetic modification comprises a hVλ and a hJλ at a mouse λ light chain locus, wherein the λ light chain locus comprises a mouse Cλ gene or functional fragment thereof; and, (b) identifying a rearranged immunoglobulin gene in said mouse, wherein the rearranged immunoglobulin gene comprises a λ light chain variable region gene segment and a Cλ gene or functional fragment thereof.

[00127] In one embodiment, the λ light chain constant gene or functional fragment thereof is selected from a human Cλ gene and a mouse Cλ gene, or a functional fragment thereof. In a specific embodiment, the λ light chain constant gene is a mouse Cλ gene, or a functional fragment thereof.

[00128] In one embodiment, the method further comprises cloning a nucleic acid sequence encoding a heavy and/or light chain variable region from the mouse, wherein the heavy and/or light chain variable region is from an antibody that comprises a human Vλ and a mouse Cλ.

[00129] In one aspect, a method for making an antibody is provided, comprising exposing a mouse as described herein to an antigen, allowing the mouse to mount an immune response that comprises making an antibody that specifically binds the antigen, identifying a rearranged nucleic acid sequence in the mouse that encodes heavy chain and a rearranged nucleic acid sequence in the mouse that encodes a cognate light chain variable domain sequence of an antibody, wherein the antibody specifically binds the antigen, and employing the nucleic acid sequences of the heavy and light chain variable domains fused to human constant domains to make a desired antibody, wherein the desired antibody comprises a light chain that comprises a Vλ domain fused to a C_L domain. In one embodiment, the Vλ domain is human and the C_L domain is a human or mouse Cλ domain. In one embodiment, the Vλ domain is mouse and the C_L domain is a human or mouse Cκ domain.

[00130] In one embodiment, a method for making an antibody is provided, comprising exposing a mouse as described herein to an antigen, allowing the mouse to mount an immune response that comprises making an antibody that specifically binds the antigen, identifying a rearranged nucleic acid sequence in the mouse that encodes a heavy chain and a rearranged nucleic acid sequence in the mouse that encodes a cognate light chain variable domain sequence of an antibody, wherein the antibody specifically binds the antigen, and employing the nucleic acid sequences of the heavy and light chain variable

domains fused to nucleic acid sequences of human constant domains to make a desired antibody, wherein the desired antibody comprises a light chain that comprises a Vλ domain fused to a Cκ domain.

[00131] In one embodiment, a method for making an antibody is provided, comprising exposing a mouse as described herein to an antigen, allowing the mouse to mount an immune response that comprises making an antibody that specifically binds the antigen, identifying a rearranged nucleic acid sequence in the mouse that encodes a heavy chain variable domain and a rearranged nucleic acid sequence that encodes a cognate light chain variable domain sequence of an antibody, wherein the antibody specifically binds the antigen, and employing the nucleic acid sequences fused to nucleic acid sequences that encode a human heavy chain constant domain and a human light chain constant domain to make an antibody derived from human sequences, wherein the antibody that specifically binds the antigen comprises a light chain that comprises a human Vλ domain fused to a mouse Cλ region.

[00132] In one embodiment, the mouse Cλ region is selected from Cλ1, Cλ2 and Cλ3. In a specific embodiment, the mouse Cλ region is Cλ2.

[00133] In one aspect, a method for making a rearranged antibody light chain variable region gene sequence is provided, comprising (a) exposing a mouse as described herein to an antigen; (b) allowing the mouse to mount an immune response; (c) identifying a cell in the mouse that comprises a nucleic acid sequence that encodes a rearranged human Vλ domain sequence fused with a mouse C_L domain, wherein the cell also encodes a cognate heavy chain comprising a human V_H domain and a mouse C_H domain, and wherein the cell expresses an antibody that binds the antigen; (d) cloning from the cell a nucleic acid sequence encoding the human Vλ domain and a nucleic acid sequence encoding the cognate human V_H domain; and, (e) using the cloned nucleic acid sequence encoding the human Vλ domain and the cloned nucleic acid sequence encoding the cognate human V_H domain to make a fully human antibody.

[00134] In one embodiment, a method for making a rearranged antibody light chain variable region gene sequence is provided, comprising (a) exposing a mouse as described in this disclosure to an antigen; (b) allowing the mouse to mount an immune response; (c) identifying a cell in the mouse that comprises a nucleic acid sequence that encodes a rearranged human Vλ domain sequence contiguous on the same nucleic acid molecule with a nucleic acid sequence encoding a mouse Cκ domain, wherein the cell also encodes a cognate heavy chain comprising a human V_H domain and a mouse C_H domain, and wherein the cell expresses an antibody that binds the antigen; (d) cloning from the cell a nucleic acids sequence encoding the human Vλ domain and a nucleic acid sequence

encoding the cognate human V_H domain; and, (e) using the cloned nucleic acid sequence encoding the human Vλ domain and the cloned nucleic acid sequence encoding the cognate human V_H domain to make a fully human antibody.

[00135] In one embodiment, a method for making a rearranged antibody light chain variable region gene sequence is provided, comprising (a) exposing a mouse as described herein to an antigen; (b) allowing the mouse to mount an immune response to the antigen; (c) identifying a cell in the mouse that comprises DNA that encodes a rearranged human Vλ domain sequence fused with a mouse Cλ domain, wherein the cell also encodes a cognate heavy chain comprising a human V_H domain and a mouse C_H domain, and wherein the cell expresses an antibody that binds the antigen; (d) cloning from the cell a nucleic acid sequence encoding the rearranged human Vλ domain and a nucleic acid sequence encoding the cognate human V_H domain; and, (e) using the cloned nucleic acid sequence encoding the human Vλ domain and the cloned nucleic acid sequence encoding the cognate human V_H domain to make a fully human antibody. In one embodiment, the mouse Cλ domain is mouse Cλ2. In a specific embodiment, the mouse Cλ domain is derived from a Cλ gene that is at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% identical to mouse Cλ2.

[00136] In one aspect, a genetically modified mouse is provided that expresses a human λ-derived light chain fused to an endogenous light chain constant region (C_L), wherein the mouse, upon immunization with antigen, makes an antibody comprising a human Vλ domain fused to a mouse C_L domain. In one embodiment, the mouse C_L domain is selected from a Cκ domain and a Cλ domain. In one embodiment, the mouse C_L domain is a Cκ domain. In one embodiment, the mouse C_L domain is a Cλ domain. In a specific embodiment, the Cλ domain is Cλ2. In a specific embodiment, the mouse Cλ domain is derived from a Cλ gene that is at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% identical to mouse Cλ2.

[00137] In one aspect, a genetically modified mouse comprising a modified endogenous κ or λ light chain locus as described herein, is provided that expresses a plurality of immunoglobulin λ light chains associated with a plurality of immunoglobulin heavy chains. In one embodiment, the heavy chain comprises a human sequence. In various embodiments, the human sequence is selected from a variable sequence, a C_H1, a hinge, a C_H2, a C_H3, and a combination thereof. In one embodiment, the plurality of immunoglobulin λ light chains comprises a human sequence. In various embodiments, the human sequence is selected from a variable sequence, a constant sequence, and a combination thereof. In one embodiment, the mouse comprises a disabled endogenous immunoglobulin locus and expresses the heavy chain and/or the λ light chain from a

transgene or extrachromosomal episome. In one embodiment, the mouse comprises a replacement at an endogenous mouse locus of some or all endogenous mouse heavy chain gene segments (*i.e.*, V, D, J), and/or some or all endogenous mouse heavy chain constant sequences (*e.g.*, C_H1, hinge, C_H2, C_H3, or a combination thereof), and/or some or all endogenous mouse light chain sequences (*e.g.*, V, J, constant, or a combination thereof), with one or more human immunoglobulin sequences.

[00138] In one aspect, a mouse suitable for making antibodies that have a human λ-derived light chain is provided, wherein all or substantially all antibodies made in the mouse are expressed with a human λ-derived light chain. In one embodiment, the human λ-derived light chain is expressed from an endogenous light chain locus. In one embodiment, the endogenous light chain locus is a κ light chain locus. In a specific embodiment, the κ light chain locus is a mouse κ light chain locus.

[00139] In one aspect, a method for making a λ-derived light chain for a human antibody is provided, comprising obtaining from a mouse as described herein a light chain sequence and a heavy chain sequence, and employing the light chain sequence and the heavy chain sequence in making a human antibody.

[00140] In one aspect, a method for making an antigen-binding protein is provided, comprising exposing a mouse as described herein to an antigen; allowing the mouse to mount an immune response; and obtaining from the mouse an antigen-binding protein that binds the antigen, or obtaining from the mouse a sequence to be employed in making an antigen-binding protein that binds the antigen.

[00141] In one aspect, a cell derived from a mouse as described herein is provided. In one embodiment, the cell is selected from an embryonic stem cell, a pluripotent cell, an induced pluripotent cell, a B cell, and a hybridoma.

[00142] In one aspect, a cell is provided that comprises a genetic modification as described herein. In one embodiment, the cell is a mouse cell. In one embodiment, the cell is selected from a hybridoma and a quadroma. In one embodiment, the cell expresses an immunoglobulin light chain that comprises a human λ variable sequence fused with a mouse constant sequence. In a specific embodiment, the mouse constant sequence is a mouse κ constant sequence.

[00143] In one aspect, a tissue derived from a mouse as described herein is provided.

[00144] In one aspect, use of a mouse or a cell as described herein to make an antigen-binding protein is provided. In one embodiment, the antigen-binding protein is a human protein. In one embodiment, the human protein is a human antibody.

[00145] In one aspect, an antigen-binding protein made by a mouse, cell, tissue, or method as described herein is provided. In one embodiment, the antigen-binding protein is a human protein. In one embodiment, the human protein is a human antibody.

[00146] Any of the embodiments and aspects described herein can be used in conjunction with one another, unless otherwise indicated or apparent from the context. Other embodiments will become apparent to those skilled in the art from a review of the ensuing description.

BRIEF DESCRIPTION OF THE FIGURES

[00147] **FIG. 1** shows a detailed illustration, not to scale, of the human λ light chain locus including the clusters of Vλ gene segments (A, B and C) and the Jλ and Cλ region pairs (J-C pairs)

[00148] **FIG. 2** shows a general illustration, not to scale, of a targeting strategy used to inactivate the endogenous mouse λ light chain locus.

[00149] **FIG. 3** shows a general illustration, not to scale, of a targeting strategy used to inactivate the endogenous mouse κ light chain locus.

[00150] **FIG. 4A** shows a general illustration, not to scale of an initial targeting vector for targeting the endogenous mouse λ light chain locus with human λ light chain sequences including 12 hVλ gene segments and hJλ1 gene segment (12/1-λ Targeting Vector).

[00151] **FIG. 4B** shows a general illustration, not to scale, of four initial targeting vectors for targeting the endogenous mouse κ light chain locus with human λ light chain sequences including 12 hVλ gene segments and hJλ1 gene segment (12/1-κ Targeting Vector), 12 hVλ gene segments and hJλ1, 2, 3 and 7 gene segments (12/4-κ Targeting Vector), 12 hVλ gene segments, a human Vκ-Jκ genomic sequence and hJλ1 gene segment (12(κ)1-κ Targeting Vector) and 12 hVλ gene segments, a human Vκ-Jκ genomic sequence and hJλ1, 2, 3 and 7 gene segments (12(κ)4-κ Targeting Vector).

[00152] **FIG. 5A** shows a general illustration, not to scale, of a targeting strategy for progressive insertion of 40 hVλ gene segments and a single hJλ gene segment into the mouse λ light chain locus.

[00153] **FIG. 5B** shows a general illustration, not to scale, of a targeting strategy for progressive insertion of 40 hVλ gene segments and a single hJλ gene segment into the mouse κ locus.

[00154] **FIG. 6** show a general illustration, not to scale, of the targeting and molecular engineering steps employed to make unique human λ-κ hybrid targeting vectors for construction of a hybrid light chain locus containing a human κ intergenic sequence, multiple hJλ gene segments or both.

[00155] **FIG. 7A** shows a general illustration, not to scale, of the locus structure for a modified mouse λ light chain locus containing 40 hVλ gene segments and a single hJλ gene segment operably linked to the endogenous Cλ2 gene.

[00156] **FIG. 7B** shows a general illustration, not to scale, of the locus structure for four independent, modified mouse κ light chain loci containing 40 hVλ gene segments and either one or four hJλ gene segments with or without a contiguous human Vκ-Jκ genomic sequence operably linked to the endogenous Cκ gene.

[00157] **FIG. 8A** shows contour plots of $Ig\lambda^+$ and $Ig\kappa^+$ splenocytes gated on $CD19^+$ from a wild type mouse (WT), a mouse homozygous for 12 hVλ and four hJλ gene segments including a human Vκ-Jκ genomic sequence (12hVλ-VκJκ-4hJλ) and a mouse homozygous for 40 hVλ and one hJλ gene segment (40hVλ-1hJλ).

[00158] **FIG. 8B** shows the total number of $CD19^+$ B cells in harvested spleens from wild type (WT), mice homozygous for 12 hVλ and four hJλ gene segments including a human Vκ-Jκ genomic sequence (12hVλ-VκJκ-4hJλ) and mice homozygous for 40 hVλ and one hJλ gene segment (40hVλ-1hJλ).

[00159] **FIG. 9A**, in the top panel, shows contour plots of splenocytes gated on singlets and stained for B and T cells ($CD19^+$ and $CD3^+$, respectively) from a wild type mouse (WT) and a mouse homozygous for 40 hVλ and four Jλ gene segments including a human Vκ-Jκ genomic sequence (40hVλ-VκJκ-4hJλ). The bottom panel shows contour plots of splenocytes gated on $CD19^+$ and stained for $Ig\lambda^+$ and $Ig\kappa^+$ expression from a wild type mouse (WT) and a mouse homozygous for 40 hVλ and four Jλ gene segments including a human Vκ-Jκ genomic sequence (40hVλ-VκJκ-4hJλ).

[00160] **FIG. 9B** shows the total number of $CD19^+$, $CD19^+Ig\kappa^+$ and $CD19^+Ig\lambda^+$ B cells in harvested spleens from wild type mice (WT) and mice homozygous for 40 hVλ and four Jλ gene segments including a human Vκ-Jκ genomic sequence (40hVλ-VκJκ-4hJλ).

[00161] **FIG. 9C** shows contour plots of splenocytes gated on $CD19^+$ and stained for immunoglobulin D (IgD) and immunoglobulin M (IgM) from a wild type mouse (WT) and a mouse homozygous for 40 hVλ and four Jλ gene segments including a human Vκ-Jκ genomic sequence (40hVλ-VκJκ-4hJλ). Mature (72 for WT, 51 for 40hVλ-VκJκ-4hJλ) and transitional (13 for WT, 22 for 40hVλ-VκJκ-4hJλ) B cells are noted on each of the contour plots.

[00162] **FIG. 9D** shows the total number of $CD19^+$ B cells, transitional B cells ($CD19^+IgM^{hi}IgD^{lo}$) and mature B cells ($CD19^+IgM^{lo}IgD^{hi}$) in harvested spleens from wild type mice (WT) and mice homozygous for 40 hVλ and four Jλ gene segments including a human Vκ-Jκ genomic sequence (40hVλ-VκJκ-4hJλ).

[00163] **FIG. 10A**, in the top panel, shows contour plots of bone marrow stained for B and T cells ($CD19^+$ and $CD3^+$, respectively) from a wild type mouse (WT) and a mouse homozygous for 40 hVλ and four Jλ gene segments including a human Vκ-Jκ genomic sequence (40hVλ-VκJκ-4hJλ). The bottom panel shows contour plots of bone marrow gated on $CD19^+$ and stained for $ckit^+$ and $CD43^+$ from a wild type mouse (WT) and a mouse homozygous for 40 hVλ and four Jλ gene segments including a human Vκ-Jκ genomic sequence (40hVλ-VκJκ-4hJλ). Pro and Pre B cells are noted on the contour plots of the bottom panel.

[00164] **FIG. 10B** shows the number of Pro ($CD19^+CD43^+ckit^+$) and Pre ($CD19^+CD43^-ckit^-$) B cells in bone marrow harvested from the femurs of wild type mice (WT) and mice homozygous for 40 hVλ and four Jλ gene segments including a human Vκ-Jκ genomic sequence (40hVλ-VκJκ-4hJλ).

[00165] **FIG. 10C** shows contour plots of bone marrow gated on singlets stained for immunoglobulin M (IgM) and B220 from a wild type mouse (WT) and a mouse homozygous for 40 hVλ and four Jλ gene segments including a human Vκ-Jκ genomic sequence (40hVλ-VκJκ-4hJλ). Immature, mature and pro/pre B cells are noted on each of the contour plots.

[00166] **FIG. 10D** shows the total number of immature ($B220^{int}IgM^+$) and mature ($B220^{hi}IgM^+$) B cells in bone marrow isolated from the femurs of wild type mice (WT) and mice homozygous for 40 hVλ and four Jλ gene segments including a human Vκ-Jκ genomic sequence (40hVλ-VκJκ-4hJλ).

[00167] **FIG. 10E** shows contour plots of bone marrow gated on immature ($B220^{int}IgM^+$) and mature ($B220^{hi}IgM^+$) B cells stained for Igλ and Igκ expression isolated from the femurs of a wild type mouse (WT) and a mouse homozygous for 40 hVλ and four Jλ gene segments including a human Vκ-Jκ genomic sequence (40hVλ-VκJκ-4hJλ).

[00168] **FIG. 11** shows a nucleotide sequence alignment of the Vλ-Jλ-Cκ junction of eighteen independent RT-PCR clones amplified from splenocyte RNA of mice bearing human λ light chain gene sequences at an endogenous mouse κ light chain locus. A6 = SEQ ID NO:57; B6 = SEQ ID NO:58; F6 = SEQ ID NO:59; B7 = SEQ ID NO:60; E7 = SEQ ID NO:61; F7 = SEQ ID NO:62; C8 = SEQ ID NO:63; E12 = SEQ ID NO:64; 1-4 = SEQ ID NO:65; 1-20 = SEQ ID NO:66; 3B43 = SEQ ID NO:67; 5-8 = SEQ ID NO:68; 5-19 = SEQ ID NO:69; 1010 = SEQ ID NO:70; 11A1 = SEQ ID NO:71; 7A8 = SEQ ID NO:72; 3A3 = SEQ ID NO:73; 2-7 = SEQ ID NO:74. Lower case bases indicate non-germline bases resulting from either mutation and/or N addition during recombination. Consensus amino acids within the Framework 4 region (FWR4) encoded by the nucleotide sequence of hJλ1 and mouse Cκ are noted at the bottom of the sequence alignment.

[00169] **FIG. 12** shows a nucleotide sequence alignment of the Vλ-Jλ-Cκ junction of twelve independent RT-PCR clones amplified from splenocyte RNA of mice bearing human λ light chain gene sequences including a contiguous human Vκ-Jκ genomic sequence at an endogenous mouse κ light chain locus. 5-2 = SEQ ID NO:87; 2-5 = SEQ ID NO:88; 1-3 = SEQ ID NO:89; 4B-1 = SEQ ID NO:90; 3B-5 = SEQ ID NO:91; 7A-1 = SEQ ID NO:92; 5-1 = SEQ ID NO:93; 4A-1 = SEQ ID NO:94; 11A-1 = SEQ ID NO:95; 5-7 = SEQ ID NO:96; 5-4 = SEQ ID NO:97; 2-3 = SEQ ID NO:98. Lower case bases indicate non-germline bases resulting from either mutation and/or N addition during recombination. Consensus amino acids within the Framework 4 region (FWR4) encoded by the nucleotide sequence of each human Jλ and mouse Cκ are noted at the bottom of the sequence alignment.

[00170] **FIG. 13** shows a nucleotide sequence alignment of the Vλ-Jλ-Cλ junction of three independent RT-PCR clones amplified from splenocyte RNA of mice bearing human λ light chain gene sequences at an endogenous mouse λ light chain locus. 2D1 = SEQ ID NO:101; 2D9 = SEQ ID NO:102; 3E15 = SEQ ID NO:103. Lower case bases indicate non-germline bases resulting from either mutation and/or N addition during recombination. Consensus amino acids within the Framework 4 region (FWR4) encoded by the nucleotide sequence of hJλ1 and mouse Cλ2 are noted at the bottom of the sequence alignment.

DETAILED DESCRIPTION

[00171] Although specific features of various embodiments are discussed in detail, the descriptions of the specific aspects, embodiments, and examples do not limit the subject matter of the claims; it is the claims that describe the scope of the invention. All terms and phrases used in this disclosure include the meanings normally ascribed to them in the art.

[00172] The term "contiguous" includes reference to occurrence on the same nucleic acid molecule, *e.g.,* two nucleic acid sequences are "contiguous" if they occur on the same nucleic molecule but are interrupted by another nucleic acid sequence. For example, a rearranged V(D)J sequence is "contiguous" with a constant region gene sequence, although the final codon of the V(D)J sequence is not followed immediately by the first codon of the constant region sequence. In another example, two V gene segment sequences are "contiguous" if they occur on the same genomic fragment, although they may be separated by sequence that does not encode a codon of the V region, *e.g.*, they may be separated by a regulatory sequence, *e.g.*, a promoter or other noncoding sequence. In one embodiment, a contiguous sequence includes a genomic fragment that contains genomic sequences arranged as found in a wild-type genome.

[00173] The phrase "derived from" when used concerning a variable region "derived from" a cited gene or gene segment includes the ability to trace the sequence back to a

particular unrearranged gene segment or gene segments that were rearranged to form a gene that expresses the variable domain (accounting for, where applicable, splice differences and somatic mutations).

[00174] The phrase "functional" when used concerning a variable region gene segment or joining gene segment refers to usage in an expressed antibody repertoire; *e.g.*, in humans Vλ gene segments 3-1, 4-3, 2-8, *etc.* are functional, whereas Vλ gene segments 3-2, 3-4, 2-5, *etc.* are nonfunctional.

[00175] A "heavy chain locus" includes a location on a chromosome, *e.g.*, a mouse chromosome, wherein in a wild-type mouse heavy chain variable (V_H) , heavy chain diversity (D_H), heavy chain joining (J_H), and heavy chain constant (C_H) region DNA sequences are found.

[00176] A "κ locus" includes a location on a chromosome, *e.g.*, a mouse chromosome, wherein in a wild-type mouse κvariable (Vκ), κ joining (Jκ), and κ constant (Cκ) region DNA sequences are found.

[00177] A "λ locus" includes a location on a chromosome, *e.g.*, a mouse chromosome, wherein in a wild-type mouse λ variable (Vλ), λ joining (Jλ), and λ constant (Cλ) region DNA sequences are found.

[00178] The term "unrearranged" includes the state of an immunoglobulin locus wherein V gene segments and J gene segments (for heavy chains, D gene segments as well) are maintained separately but are capable of being joined to form a rearranged V(D)J gene that comprises a single V,(D),J of the V(D)J repertoire.

Mice Expressing Human λ Variable Domains

[00179] Mice that express antibodies that are fully human, or partly human and partly mouse, have previously been reported. VELOCIMMUNE® genetically engineered mice comprise a replacement of unrearranged V(D)J gene segments at endogenous mouse loci with human V(D)J gene segments. VELOCIMMUNE® mice express chimeric antibodies having human variable domains and mouse constant domains (see, *e.g.*, US Pat. No. 7,605,237). Most other reports concern mice that express fully human antibodies from fully human transgenes in mice that have disabled endogenous immunoglobulin loci.

[00180] Antibody light chains are encoded by one of two separate loci: kappa (κ) and lambda (λ). Mouse antibody light chains are primarily of the κ type. Mice that make mouse antibodies, and modified mice that make fully human or chimeric human-mouse antibodies, display a bias in light chain usage. Humans also exhibit light chain bias, but not so pronounced as in mice; the ratio of κ light chains to λ light chains in mice is about 95:5, whereas in humans the ratio is about 60:40. The more pronounced bias in mice is

not thought to severely affect antibody diversity, because in mice the λ variable locus is not so diverse in the first instance. This is not so in humans. The human λ light chain locus is richly diverse.

[00181] The human λ light chain locus extends over 1,000 kb and contains over 80 genes that encode variable (V) or joining (J) segments (FIG. 1). Within the human λ light chain locus, over half of all observed Vλ domains are encoded by the gene segments 1-40, 1-44, 2-8, 2-14, and 3-21. Overall, about 30 or so of the human Vλ gene segments are believed to be functional. There are seven Jλ gene segments, only four of which are regarded as generally functional Jλ gene segments—Jλ1, Jλ2, Jλ3, and Jλ7.

[00182] The λ light chain locus in humans is similar in structure to the κ locus of both mice and humans in that the human λ light chain locus has several variable region gene segments that are capable of recombining to form a functional light chain protein. The human λ light chain locus contains approximately 70 V gene segments and 7 Jλ-Cλ gene segment pairs. Only four of these Jλ-Cλ gene segment pairs appear to be functional. In some alleles, a fifth Jλ-Cλ gene segment pair is reportedly a pseudo gene (Cλ6). The 70 Vλ gene segments appear to contain 38 functional gene segments. The 70 Vλ sequences are arranged in three clusters, all of which contain different members of distinct V gene family groups (clusters A, B and C; FIG. 1). This is a potentially rich source of relatively untapped diversity for generating antibodies with human V regions in non-human animals.

[00183] In stark contrast, the mouse λ light chain locus contains only two or three (depending on the strain) mouse Vλ region gene segments (FIG. 2). At least for this reason, the severe κ bias in mice is not thought to be particularly detrimental to total antibody diversity.

[00184] According published maps of the mouse λ light chain locus, the locus consists essentially of two clusters of gene segments within a span of approximately 200 kb (FIG.2). The two clusters contain two sets of V, J, and C genes that are capable of independent rearrangement: Vλ2-Jλ2-Cλ2-Jλ4-Cλ4 and Vλ1-Jλ3-Cλ3-Jλ1-Cλ1. Although Vλ2 has been found to recombine with all Jλ gene segments, Vλ1 appears to exclusively recombine with Cλ1. Cλ4 is believed to be a pseudo gene with defective splice sites.

[00185] The mouse κ light chain locus is strikingly different. The structure and number of gene segments that participate in the recombination events leading to a functional light chain protein from the mouse κ locus is much more complex (FIG. 3). Thus, mouse λ light chains do not greatly contribute to the diversity of an antibody population in a typical mouse.

[00186] Exploiting the rich diversity of the human λ light chain locus in mice would likely result in, among other things, a source for a more complete human repertoire of light chain V domains. Previous attempts to tap this diversity used human transgenes containing chunks of the human λ light chain locus randomly incorporated into the mouse genome (see, *e.g.*, US 6,998,514 and US 7,435,871). Mice containing these randomly integrated transgenes reportedly express fully human λ light chains, however, in some cases, one or both endogenous light chain loci remain intact. This situation is not desirable as the human λ light chain sequences contend with the mouse light chain (κ or λ) in the expressed antibody repertoire of the mouse.

[00187] In contrast, the inventors describe genetically modified mice that are capable of expressing one or more λ light chain nucleic acid sequences directly from a mouse light chain locus, including by replacement at an endogenous mouse light chain locus. Genetically modified mice capable of expressing human λ light chain sequences from an endogenous locus may be further bred to mice that comprise a human heavy chain locus and thus be used to express antibodies comprising V regions (heavy and light) that are fully human. In various embodiments. The V regions express with mouse constant regions. In various embodiments, no endogenous mouse immunoglobulin gene segments are present and the V regions express with human constant regions. These antibodies would prove useful in numerous applications, both diagnostic as well as therapeutic.

[00188] Many advantages can be realized for various embodiments of expressing binding proteins derived from human Vλ and Jλ gene segments in mice. Advantages can be realized by placing human λ sequences at an endogenous light chain locus, for example, the mouse κ or λ locus. Antibodies made from such mice can have light chains that comprise human Vλ domains fused to a mouse C_L region, specifically a mouse Cκ or Cλ region. The mice will also express human Vλ domains that are suitable for identification and cloning for use with human C_L regions, specifically Cκ and/or Cλ regions. Because B cell development in such mice is otherwise normal, it is possible to generate compatible Vλ domains (including somatically mutated Vλ domains) in the context of either Cλ or Cκ regions.

[00189] Genetically modified mice are described that comprise an unrearranged Vλ gene segment at an immunoglobulin κ or λ light chain locus. Mice that express antibodies that comprise a light chain having a human Vλ domain fused to a Cκ and/or Cλ region are described.

Sterile Transcripts of the Immunoglobulin κ Light Chain Locus

[00190] Variations on the theme of expressing human immunoglobulin λ sequences in mice are reflected in various embodiments of genetically modified mice capable of such expression. Thus, in some embodiments, the genetically modified mice comprise certain non-coding sequence(s) from a human locus. In one embodiment, the genetically modified mouse comprises human Vλ and Jλ gene segments at an endogenous κ light chain locus, and further comprises a human κ light chain genomic fragment. In a specific embodiment, the human κ light chain genomic fragment is a non-coding sequence naturally found between a human Vκ gene segment and a human Jκ gene segment.

[00191] The human and mouse κ light chain loci contain sequences that encode sterile transcripts that lack either a start codon or an open reading frame, and that are regarded as elements that regulate transcription of the κ light chain loci. These sterile transcripts arise from an intergenic sequence located downstream or 3' of the most proximal Vκ gene segment and upstream or 5' of the κ light chain intronic enhancer (Eκi) that is upstream of the κ light chain constant region gene (Cκ). The sterile transcripts arise from rearrangement of the intergenic sequence to form a VκJκ1 segment fused to a Cκ.

[00192] A replacement of the κ light chain locus upstream of the Cκ gene would remove the intergenic region encoding the sterile transcripts. Therefore, in various embodiments, a replacement of mouse κ light chain sequence upstream of the mouse Cκ gene with human λ light chain gene segments would result in a humanized mouse κ light chain locus that contains human Vλ and Jλ gene segments but not the κ light chain intergenic region that encodes the sterile transcripts.

[00193] As described herein, humanization of the endogenous mouse κ light chain locus with human λ light chain gene segments, wherein the humanization removes the intergenic region, results in a striking drop in usage of the κ light chain locus, coupled with a marked increase in λ light chain usage. Therefore, although a humanized mouse that lacks the intergenic region is useful in that it can make antibodies with human light chain variable domains (*e.g.*, human λ or κ domains), usage from the locus decreases.

[00194] Also described is humanization of the endogenous mouse κ light chain locus with human Vλ and Jλ gene segments coupled with an insertion of a human κ intergenic region to create a Vλ locus that contains, with respect to transcription, between the final human Vλ gene segment and the first human Jλ gene segment, a κ intergenic region; which exhibits a B cell population with a higher expression than a locus that lacks the κ intergenic region. This observation is consistent with a hypothesis that the intergenic region—directly through a sterile transcript, or indirectly—suppresses usage from the

endogenous λ light chain locus. Under such a hypothesis, including the intergenic region would result in a decrease in usage of the endogenous λ light chain locus, leaving the mouse a restricted choice but to employ the modified (λ into κ) locus to generate antibodies.

[00195] In various embodiments, a replacement of mouse κ light chain sequence upstream of the mouse Cκ gene with human λ light chain sequence further comprises a human κ light chain intergenic region disposed, with respect to transcription, between the 3' untranslated region of the 3' most Vλ gene segment and 5' to the first human Jλ gene segment. Alternatively, such an intergenic region may be omitted from a replaced endogenous κ light chain locus (upstream of the mouse Cκ gene) by making a deletion in the endogenous λ light chain locus. Likewise, under this embodiment, the mouse generates antibodies from an endogenous κ light chain locus containing human λ light chain sequences.

Approaches to Engineering Mice to Express Human Vλ Domains

[00196] Various approaches to making genetically modified mice that make antibodies that contain a light chain that has a human Vλ domain fused to an endogenous C_L (*e.g.* Cκ or Cλ) region are described. Genetic modifications are described that, in various embodiments, comprise a deletion of one or both endogenous light chain loci. For example, to eliminate mouse λ light chains from the endogenous antibody repertoire a deletion of a first Vλ-Jλ-Cλ gene cluster and replacement, in whole or in part, of the Vλ-Jλ gene segments of a second gene cluster with human Vλ-Jλ gene segments can be made. Genetically modified mouse embryos, cells, and targeting constructs for making the mice, mouse embryos, and cells are also provided.

[00197] The deletion of one endogenous Vλ-Jλ-Cλ gene cluster and replacement of the Vλ-Jλ gene segments of another endogenous Vλ-Jλ-Cλ gene cluster employs a relatively minimal disruption in natural antibody constant region association and function in the animal, in various embodiments, because endogenous Cλ genes are left intact and therefore retain normal functionality and capability to associate with the constant region of an endogenous heavy chain. Thus, in such embodiments the modification does not affect other endogenous heavy chain constant regions dependent upon functional light chain constant regions for assembly of a functional antibody molecule containing two heavy chains and two light chains. Further, in various embodiments the modification does not affect the assembly of a functional membrane-bound antibody molecule involving an endogenous heavy chain and a light chain, *e.g.*, a hVλ domain linked to a mouse Cλ region. Because at least one functional Cλ gene is retained at the endogenous locus,

animals containing a replacement of the Vλ-Jλ gene segments of an endogenous Vλ-Jλ-Cλ gene cluster with human Vλ-Jλ gene segments should be able to make normal λ light chains that are capable of binding antigen during an immune response through the human Vλ-Jλ gene segments present in the expressed antibody repertoire of the animal.

[00198] A schematic illustration (not to scale) of a deleted endogenous mouse Vλ-Jλ-Cλ gene cluster is provided in FIG. 2. As illustrated, the mouse λ light chain locus is organized into two gene clusters, both of which contain function gene segments capable of recombining to form a function mouse λ light chain. The endogenous mouse Vλ1-Jλ3-Cλ3-Jλ1-Cλ1 gene cluster is deleted by a targeting construct (Targeting Vector 1) with a neomycin cassette flanked by recombination sites. The other endogenous gene cluster (Vλ2-Vλ3-Jλ2-Cλ2-Jλ4-Cλ4) is deleted in part by a targeting construct (Targeting Vector 2) with a hygromycin-thymidine kinase cassette flanked by recombination sites. In this second targeting event, the Cλ2-Jλ4-Cλ4 endogenous gene segments are retained. The second targeting construct (Targeting Vector 2) is constructed using recombination sites that are different than those in the first targeting construct (Targeting Vector 1) thereby allowing for the selective deletion of the selection cassette after a successful targeting has been achieved. The resulting double-targeted locus is functionally silenced in that no endogenous λ light chain can be produced. This modified locus can be used for the insertion of human Vλ and Jλ gene segments to create an endogenous mouse λ locus comprising human Vλ and Jλ gene segments, whereby, upon recombination at the modified locus, the animal produces λ light chains comprising rearranged human Vλ and Jλ gene segments linked to an endogenous mouse Cλ gene segment.

[00199] Genetically modifying a mouse to render endogenous λ gene segments nonfunctional, in various embodiments, results in a mouse that exhibits exclusively κ light chains in its antibody repertoire, making the mouse useful for evaluating the role of λ light chains in the immune response, and useful for making an antibody repertoire comprising Vκ domains but not Vλ domains.

[00200] A genetically modified mouse that expresses a hVλ linked to a mouse Cλ gene having been recombined at the endogenous mouse λ light chain locus can be made by any method known in the art. A schematic illustration (not to scale) of the replacement of the endogenous mouse Vλ2-Vλ3-Jλ2 gene segments with human Vλ and Jλ gene segments is provided in FIG. 4A. As illustrated, an endogenous mouse λ light chain locus that had been rendered nonfunctional is replaced by a targeting construct (12/1-λ Targeting Vector) that includes a neomycin cassette flanked by recombination sites. The Vλ2-Vλ3-Jλ2 gene

segments are replaced with a genomic fragment containing human λ sequence that includes 12 hVλ gene segments and a single hJλ gene segment.

[00201] Thus, this first approach positions one or more hVλ gene segments at the endogenous λ light chain locus contiguous with a single hJλ gene segment (FIG. 4A).

[00202] Further modifications to the modified endogenous λ light chain locus can be achieved with using similar techniques to insert more hVλ gene segments. For example, schematic illustrations of two additional targeting constructs (+16-λ and +12-λ Targeting Vectors) used for progressive insertion of addition human hVλ gene segments are provided in FIG. 5A. As illustrated, additional genomic fragments containing specific human hVλ gene segments are inserted into the modified endogenous λ light chain locus in successive steps using homology provided by the previous insertion of human λ light chain sequences. Upon recombination with each targeting construct illustrated, in sequential fashion, 28 additional hVλ gene segments are inserted into the modified endogenous λ light chain locus. This creates a chimeric locus that produces a λ light chain protein that comprises human Vλ-Jλ gene segments linked to a mouse Cλ gene.

[00203] The above approaches to insert human λ light chain gene segments at the mouse λ locus, maintains the enhancers positioned downstream of the Cλ2-Jλ4-Cλ4 gene segments (designated Enh 2.4, Enh and Enh 3.1 FIG. 4A and FIG. 5A). This approach results in a single modified allele at the endogenous mouse λ light chain locus (FIG. 7A).

[00204] Compositions and methods for making a mouse that expresses a light chain comprising hVλ and Jλ gene segments operably linked to a mouse Cλ gene segment, are provided, including compositions and method for making a mouse that expresses such genes from an endogenous mouse λ light chain locus. The methods include selectively rendering one endogenous mouse Vλ-Jλ-Cλ gene cluster nonfunctional (*e.g.*, by a targeted deletion), and employing a hVλ and Jλ gene segments at the endogenous mouse λ light chain locus to express a hVλ domain in a mouse.

[00205] Alternatively, in a second approach, human λ light chain gene segments may be positioned at the endogenous κ light chain locus. The genetic modification, in various embodiments, comprises a deletion of the endogenous κ light chain locus. For example, to eliminate mouse κ light chains from the endogenous antibody repertoire a deletion of the mouse Vκ and Jκ gene segments can be made. Genetically modified mouse embryos, cells, and targeting constructs for making the mice, mouse embryos, and cells are also provided.

[00206] For the reasons stated above, the deletion of the mouse Vκ and Jκ gene segments employs a relatively minimal disruption. A schematic illustration (not to scale) of

deleted mouse Vκ and Jκ gene segments is provided in FIG. 3. The endogenous mouse Vκ and Jκ gene segments are deleted via recombinase-mediated deletion of mouse sequences position between two precisely positioned targeting vectors each employing site-specific recombination sites. A first targeting vector (Jκ Targeting Vector) is employed in a first targeting event to delete the mouse Jκ gene segments. A second targeting vector (Vκ Targeting Vector) is employed in a second, sequential targeting event to delete a sequence located 5' of the most distal mouse Vκ gene segment. Both targeting vectors contain site-specific recombination sites thereby allowing for the selective deletion of both selection cassettes and all intervening mouse κ light chain sequences after a successful targeting has been achieved. The resulting deleted locus is functionally silenced in that no endogenous κ light chain can be produced. This modified locus can be used for the insertion of hVλ and Jλ gene segments to create an endogenous mouse κ locus comprising hVλ and Jλ gene segments, whereby, upon recombination at the modified locus, the animal produces λ light chains comprising rearranged hVλ and Jλ gene segments operably linked to an endogenous mouse Cκ gene segment. Various targeting vectors comprising human λ light chain sequences can be used in conjunction with this deleted mouse κ locus to create a hybrid light chain locus containing human λ gene segments operably linked with a mouse Cκ region.

[00207] Thus, a second approach positions one or more human Vλ gene segments are positioned at the mouse κ light chain locus contiguous with a single human Jλ gene segment (12/1-κ Targeting Vector, FIG. 4B).

[00208] In various embodiments, modifications to this approach can be made to add gene segments and/or regulatory sequences to optimize the usage of the human λ light chain sequences from the mouse κ locus within the mouse antibody repertoire.

[00209] In a third approach, one or more hVλ gene segments are positioned at the mouse κ light chain locus contiguous with four hJλ gene sequences (12/4-κ Targeting Vector FIG. 4B).

[00210] In a third approach, one or more hVλ gene segments are positioned at the mouse κ light chain locus contiguous with a human κ intergenic sequence and a single hJλ gene sequence (12(κ)1-κ Targeting Vector, FIG. 4B).

[00211] In a fourth approach, one or more hVλ gene segments are positioned at the mouse κ light chain locus contiguous with a human κ intergenic sequence four hJλ gene sequences (12(κ)4-κ Targeting Vector FIG. 4B).

[00212] All of the above approaches to insert human λ light chain gene segments at the mouse κ locus, maintain the κ intronic enhancer element upstream of the Cκ gene

(designated Eκi, FIG. 4B and FIG. 5B) and the 3' κ enhancer downstream of the Cκ gene (designated Eκ3', FIG. 4B and FIG. 5B). The approaches result in four separate modified alleles at the endogenous mouse κ light chain locus (FIG. 7B).

[00213] In various embodiments, genetically modified mouse comprise a knockout of the endogenous mouse λ light chain locus. In one embodiment, the λ light chain locus is knocked out by a strategy that deletes the region spanning Vλ2 to Jλ2, and the region spanning Vλ1 to Cλ1 (FIG. 2). Any strategy that reduces or eliminates the ability of the endogenous λ light chain locus to express endogenous λ domains is suitable for use with embodiments in this disclosure.

Lambda Domain Antibodies from Genetically Modified Mice

[00214] Mice comprising human λ sequences at either the mouse κ or λ light chain locus will express a light chain that comprises a hVλ region fused to a mouse C_L (Cκ or Cλ) region. These are advantageously bred to mice that (a) comprise a functionally silenced light chain locus (*e.g.*, a knockout of the endogenous mouse κ or λ light chain locus); (b) comprise an endogenous mouse λ light chain locus that comprises hV and hJ gene segments operably linked to an endogenous mouse Cλ gene; (c) comprise an endogenous mouse κ light chain locus that comprises hVκ and hJκ gene segments operably linked to an endogenous mouse Cκ gene; and, (d) a mouse in which one κ allele comprises hVκs and hJκs; the other κ allele comprising hVλs and hJλs; one λ allele comprising hVλs and hJλs and one λ allele silenced or knocked out, or both λ alleles comprising hVλs and hJλs; and, two heavy chain alleles that each comprise hV_Hs, hD_Hs, and hJ_Hs.

[00215] The antibodies that comprise the hVλ domains expressed in the context of either Cκ or Cλ are used to make fully human antibodies by cloning the nucleic acids encoding the hVλ domains into expression constructs that bear genes encoding human Cλ. Resulting expression constructs are transfected into suitable host cells for expressing antibodies that display a fully hVλ domain fused to hCλ.

EXAMPLES

[00216] The following examples are provided so as to describe how to make and use methods and compositions of the invention, and are not intended to limit the scope of what the inventors regard as their invention. Unless indicated otherwise, temperature is indicated in Celsius, and pressure is at or near atmospheric.

Example I

Deletion of the Mouse Immunoglobulin Light Chain Loci

[00217] Various targeting constructs were made using VELOCIGENE® technology (see, *e.g.*, US Pat. No. 6,586,251 and Valenzuela *et al.* (2003) High-throughput engineering of the mouse genome coupled with high-resolution expression analysis, *Nature Biotech.* 21(6):652-659) to modify mouse genomic Bacterial Artificial Chromosome (BAC) libraries to inactivate the mouse κ and λ light chain loci.

[00218] **Deletion of the mouse λ light chain locus.** DNA from mouse BAC clone RP23-135k15 (Invitrogen) was modified by homologous recombination to inactivate the endogenous mouse λ light chain locus through targeted deletion of the Vλ-Jλ-Cλ gene clusters (FIG. 2).

[00219] Briefly, the entire proximal cluster comprising Vλ1-Jλ3-Cλ3-Jλ1-Cλ1 gene segments was deleted in a single targeting event using a targeting vector comprising a neomycin cassette flanked by *lox*P sites with a 5' mouse homology arm containing sequence 5' of the Vλ1 gene segment and a 3' mouse homology arm containing sequence 3' of the Cλ1 gene segment (FIG. 2, Targeting Vector 1).

[00220] A second targeting construct was prepared to precisely delete the distal endogenous mouse λ gene cluster containing Vλ2-Jλ2-Cλ2-Jλ4-Cλ4 except that the targeting construct contained a 5' mouse homology arm that contained sequence 5' of the Vλ2 gene segment and a 3' mouse homology arm that contained sequence 5' to the endogenous Cλ2 gene segment (FIG. 2, Targeting Vector 2). Thus, the second targeting construct precisely deleted Vλ2-Jλ2, while leaving Cλ2-Jλ4-Cλ4 intact at the endogenous mouse λ locus. ES cells containing an inactivated endogenous λ locus (as described above) were confirmed by karyotyping and screening methods (*e.g.*, TAQMAN®) known in the art. DNA was then isolated from the modified ES cells and subjected to treatment with CRE recombinase thereby mediating the deletion of the proximal targeting cassette containing the neomycin marker gene, leaving only a single *lox*P site at the deletion point (FIG. 2, bottom).

[00221] **Deletion of the mouse κ light chain locus.** Several targeting constructs were made using similar methods described above to modify DNA from mouse BAC clones RP23-302g12 and RP23-254m04 (Invitrogen) by homologous recombination to inactivate the mouse κ light chain locus in a two-step process (FIG. 3).

[00222] Briefly, the Jκ gene segments (1-5) of the endogenous mouse κ light chain locus were deleted in a single targeting event using a targeting vector comprising a hygromycin-thymidine kinase (hyg-TK) cassette containing a single *lox*P site 3' to the hyg-TK cassette (FIG. 3, Jκ Targeting Vector). The homology arms used to make this targeting

vector contained mouse genomic sequence 5' and 3' of the endogenous mouse Jκ gene segments. In a second targeting event, a second targeting vector was prepared to delete a portion of mouse genomic sequence upstream (5') to the most distal endogenous mouse Vκ gene segment (FIG. 3, Vκ Targeting Vector). This targeting vector contained an inverted *lox*511 site, a *lox*P site and a neomycin cassette. The homology arms used to make this targeting vector contained mouse genomic sequence upstream of the most distal mouse Vκ gene segment. The targeting vectors were used in a sequential fashion (*i.e.*, Jκ then Vκ) to target DNA in ES cells. ES bearing a double-targeted chromosome (*i.e.*, a single endogenous mouse κ locus targeted with both targeting vectors) were confirmed by karyotyping and screening methods (*e.g.*, Taqman™) known in the art. DNA was then isolated from the modified ES cells and subjected to treatment with Cre recombinase thereby mediating the deletion of endogenous mouse Vκ gene segments and both selection cassettes, while leaving two juxtaposed *lox* sites in opposite orientation relative to one another (FIG. 3, bottom; SEQ ID NO:1).

[00223] Thus, two modified endogenous light chain loci (κ and λ) containing intact enhancer and constant regions were created for progressively inserting unrearranged human λ germline gene segments in a precise manner using targeting vectors described below.

Example II

Replacement of Mouse Light Chain Loci with a Human λ Light Chain Mini-Locus

[00224] Multiple targeting vectors were engineered for progressive insertion of human λ gene segments into the endogenous mouse κ and λ light chain loci using similar methods as described above. Multiple independent initial modifications were made to the endogenous light chain loci each producing a chimeric light chain locus containing hVλ and Jλ gene segments operably linked to mouse light chain constant genes and enhancers.

[00225] **A human λ mini-locus containing 12 human Vλ and one human Jλ gene segment.** A series of initial targeting vectors were engineered to contain the first 12 consecutive human Vλ gene segments from cluster A and a hJλ1 gene segment or four hJλ gene segments using a human BAC clone named RP11-729g4 (Invitrogen). FIGs. 4A and 4B show the targeting vectors that were constructed for making an initial insertion of human λ light chain gene segments at the mouse λ and κ light chain loci, respectively.

[00226] For a first set of initial targeting vectors, a 124,125 bp DNA fragment from the 729g4 BAC clone containing 12 hVλ gene segments and a hJλ1 gene segment was engineered to contain a PI-SceI site 996 bp downstream (3') of the hJλ1 gene segment for

ligation of a 3' mouse homology arm. Two different sets of homology arms were used for ligation to this human fragment; one set of homology arms contained endogenous mouse λ sequences from the 135k15 BAC clone (FIG.4A) and another set contained endogenous κ sequence 5' and 3' of the mouse Vκ and Jκ gene segments from mouse BAC clones RP23-302g12 and RP23-254m04, respectively (FIG. 4B).

[00227] For the 12/1-λ Targeting Vector (FIG. 4A), a PI-SceI site was engineered at the 5' end of a 27,847 bp DNA fragment containing the mouse Cλ2-Jλ4-Cλ4 and enhancer 2.4 of the modified mouse λ locus described in Example 1. The ~28 kb mouse fragment was used as a 3' homology arm by ligation to the ~124 kb human λ fragment, which created a 3' junction containing, from 5' to 3', a hJλ1 gene segment, 996 bp of human λ sequence 3' of the hJλ1 gene segment, 1229 bp of mouse λ sequence 5' to the mouse Cλ2 gene, the mouse Cλ2 gene and the remaining portion of the ~28 kb mouse fragment. Upstream (5') from the human Vλ3-12 gene segment was an additional 1456 bp of human λ sequence before the start of the 5' mouse homology arm, which contained 23,792 bp of mouse genomic DNA corresponding to sequence 5' of the endogenous mouse λ locus. Between the 5' homology arm and the beginning of the human λ sequence was a neomycin cassette flanked by Frt sites.

[00228] Thus, the 12/1-λ Targeting Vector included, from 5' to 3', a 5' homology arm containing ~24 kb of mouse λ genomic sequence 5' of the endogenous λ locus, a 5' Frt site, a neomycin cassette, a 3' Frt site, ~123 kb of human genomic λ sequence containing the first 12 consecutive hVλ gene segments and a hJλ1 gene segment, a PI-SceI site, and a 3' homology arm containing ~28 kb of mouse genomic sequence including the endogenous Cλ2-Jλ4-Cλ4 gene segments, the mouse enhancer 2.4 sequence and additional mouse genomic sequence downstream (3') of the enhancer 2.4 (FIG. 4A).

[00229] In a similar fashion, the 12/1-κ Targeting Vector (FIG. 4B) employed the same ~124 human λ fragment with the exception that mouse homology arms containing mouse κ sequence were used such that targeting to the endogenous κ locus could be achieved by homologous recombination. Thus, the 12/1-κ Targeting Vector included, from 5' to 3', a 5' homology arm containing ~23 kb of mouse genomic sequence 5' of the endogenous κ locus, an I-CeuI site, a 5' Frt site, a neomycin cassette, a 3' Frt site, ~124 kb of human genomic λ sequence containing the first 12 consecutive hVλ gene segments and a hJλ1 gene segment, a PI-SceI site, and a 3' homology arm containing ~28 kb of mouse genomic sequence including the endogenous the mouse Cκ gene, Eκi and Eκ3' and additional mouse genomic sequence downstream (3') of Eκ3' (FIG. 4B, 12/1-κ Targeting Vector).

[00230] Homologous recombination with either of these two initial targeting vectors created a modified mouse light chain locus (κ or λ) containing 12 hVλ gene segments and a hJλ1 gene segment operably linked to the endogenous mouse light chain constant gene and enhancers (Cκ or Cλ2 and Eκi/Eκ3' or Enh 2.4/Enh 3.1) gene which, upon recombination, leads to the formation of a chimeric λ light chain.

[00231] **A human λ mini-locus with 12 human Vλ and four human Jλ gene segments.** In another approach to add diversity to a chimeric λ light chain locus, a third initial targeting vector was engineered to insert the first 12 consecutive human Vλ gene segments from cluster A and hJλ1, 2, 3 and 7 gene segments into the mouse κ light chain locus (FIG. 4B, 12/4-κ Targeting Vector). A DNA segment containing hJλ1, Jλ2, Jλ3 and Jλ7 gene segments was made by *de novo* DNA synthesis (Integrated DNA Technologies) including each Jλ gene segment and human genomic sequence of ~100 bp from both the immediate 5' and 3' regions of each Jλ gene segment. A PI-SceI site was engineered into the 3' end of this ~1 kb DNA fragment and ligated to a chloroamphenicol cassette. Homology arms were PCR amplified from human λ sequence at 5' and 3' positions relative to the hJλ1 gene segment of the human BAC clone 729g4. Homologous recombination with this intermediate targeting vector was performed on a modified 729g4 BAC clone that had been previously targeted upstream (5') of the human Vλ3-12 gene segment with a neomycin cassette flanked by Frt sites, which also contained an I-CeuI site 5' to the 5' Frt site. The double-targeted 729g4 BAC clone included from 5' to 3' an I-CeuI site, a 5' Frt site, a neomycin cassette, a 3' Frt site, a ~123 kb fragment containing the first 12 hVλ gene segments, a ~1 kb fragment containing human Jλ1, 2, 3 and 7 gene segments, a PI-SceI site, and a chloroamphenicol cassette. This intermediate targeting vector was digested together with I-CeuI and PI-SceI and subsequently ligated into the modified mouse BAC clone (described above) to create the third targeting vector.

[00232] This ligation resulted in a third targeting vector for insertion of human λ sequences into the endogenous κ light chain locus, which included, from 5' to 3', a 5' mouse homology arm containing ~23 kb of genomic sequence 5' of the endogenous mouse κ locus, an I-CeuI site, a 5' Frt site, a neomycin cassette, a 3' Frt site, a ~123 kb fragment containing the first 12 hVλ gene segments, a ~1 kb fragment containing hJλ1, 2, 3 and 7 gene segments, a PI-SceI site and a 3' homology arm containing ~28 kb of mouse genomic sequence including the endogenous the mouse Cκ gene, Eκi and Eκ3' and additional mouse genomic sequence downstream (3') of Eκ3' (FIG. 4B, 12/4-κ Targeting Vector). Homologous recombination with this third targeting vector created a modified mouse κ light chain locus containing 12 hVλ gene segments and four hJλ gene segments

operably linked to the endogenous mouse Cκ gene which, upon recombination, leads to the formation of a chimeric human λ/mouse κ light chain.

[00233] A human λ mini-locus with an integrated human κ light chain sequence. In a similar fashion, two additional targeting vectors similar to those engineered to make an initial insertion of human λ gene segments into the endogenous κ light chain locus (FIG. 4B, 12/1-κ and 12/4-κ Targeting Vectors) were engineered to progressively insert human λ light chain gene segments using uniquely constructed targeting vectors containing contiguous human λ and κ genomic sequences. These targeting vectors were constructed to include a ~ 23 kb human κ genomic sequence naturally located between human Vκ4-1 and Jκ1 gene segments. This human κ genomic sequence was specifically positioned in these two additional targeting vectors between human Vλ and human Jλ gene segments (FIG. 4B, 12(κ)1-κ and 12(κ)4-κ Targeting Vectors).

[00234] Both targeting vectors containing the human κ genomic sequence were made using the modified RP11-729g4 BAC clone described above (FIG. 6). This modified BAC clone was targeted with a spectinomycin selection cassette flanked by NotI and AsiSI restriction sites (FIG.6, top left). Homologous recombination with the spectinomycin cassette resulted in a double-targeted 729g4 BAC clone which included, from 5' to 3', an I-CeuI site, a 5' Frt site, a neomycin cassette, a 3' Frt site, a ~123 kb fragment containing the first 12 hVλ gene segments, a NotI site about 200 bp downstream (3') to the nonamer sequence of the hVλ3-1 gene segment, a spectinomycin cassette and an AsiSI site. A separate human BAC clone containing human κ sequence (CTD-2366j12) was targeted two independent times to engineer restriction sites at locations between hVκ4-1 and hJκ1 gene segments to allow for subsequent cloning of a ~23 kb fragment for ligation with the hVλ gene segments contained in the double targeted modified 729g4 BAC clone (FIG. 6, top right).

[00235] Briefly, the 2366j12 BAC clone is about 132 kb in size and contains hVκ gene segments 1-6, 1-5, 2-4, 7-3, 5-2, 4-1, human κ genomic sequence down stream of the Vκ gene segments, hJκ gene segments 1-5, the hCκ and about 20 kb of additional genomic sequence of the human κ locus. This clone was first targeted with a targeting vector containing a hygromycin cassette flanked by Frt sites and a NotI site downstream (3') of the 3' Frt site. The homology arms for this targeting vector contained human genomic sequence 5' and 3' of the Vκ gene segments within the BAC clone such that upon homologous recombination with this targeting vector, the Vκ gene segments were deleted and a NotI site was engineered ~133 bp downstream of the hVκ4-1 gene segment (FIG. 6, top right). This modified 2366j12 BAC clone was targeted independently with two targeting

vectors at the 3' end to delete the hJκ gene segments with a chloroamphenicol cassette that also contained either a hJλ1 gene segment, a PI-SceI site and an AsiSI site or a human λ genomic fragment containing four hJλ gene segments (*supra*), a PI-SceI site and an AsiSI site (FIG. 6, top right). The homology arms for these two similar targeting vectors contained sequence 5' and 3' of the hJκ gene segments. Homologous recombination with these second targeting vectors and the modified 2366j12 BAC clone yielded a double-targeted 2366j12 clone which included, from 5' to 3', a 5' Frt site, a hygromycin cassette, a 3' Frt site, a NotI site, a 22,800 bp genomic fragment of the human κ locus containing the intergenic region between the Vκ4-1 and Jκ1 gene segments, either a hJλ1 gene segment or a human λ genomic fragment containing hJλ1, Jλ2, Jλ3 and Jλ7, a PI-SceI site and a chloroamphenicol cassette (FIG. 6, top right). Two final targeting vectors to make the two additional modifications were achieved by two ligation steps using the double-targeted 729g4 and 2366j12 clones.

[00236] Double targeted 729g4 and 2366j12 clones were digested with NotI and AsiSI yielding one fragment containing the neomycin cassette and hVλ gene segments and another fragment containing the ~23 kb genomic fragment of the human κ locus containing the intergenic region between the Vκ4-1 and Jκ1 gene segments, either a hJλ1 gene segment or a genomic fragment containing hJλ1, Jλ2, Jλ3 and Jλ7 gene segments, the PI-SceI site and the chloroamphenicol cassette, respectively. Ligation of these fragments generated two unique BAC clones containing from 5' to 3' the hVλ gene segments, the human κ genomic sequence between the Vκ4-1 and Jκ1 gene segments, either a hJλ1 gene segment or a genomic fragment containing hJλ1, Jλ2, Jλ3 and Jλ7 gene segments, a PI-SceI site and a chloroamphenicol cassette (FIG. 6, bottom). These new BAC clones were then digested with I-CeuI and PI-SceI to release the unique fragments containing the upstream neomycin cassette and the contiguous human λ and κ sequences and ligated into a modified mouse BAC clone 302g12 which contained from 5' to 3' mouse genomic sequence 5' of the endogenous κ locus, an I-CeuI site, a 5' Frt site, a neomycin cassette, a 3' Frt site, hVλ gene segments (3-12 to 3-1), a NotI site ~200 bp downstream of Vλ3-1, ~23 kb of human κ sequence naturally found between the human Vκ4-1 and Jκ1 gene segments, either a hJλ1 gene segment or a genomic fragment containing hJλ1, Jλ2, Jλ3 and Jλ7 gene segments, the mouse Eκi, the mouse Cκ gene and Eκ3' (FIG. 4, 12hVλ-VκJκ-hJλ1 and 12hVλ-VκJκ-4hJλ Targeting Vectors). Homologous recombination with both of these targeting vectors created two separate modified mouse κ light chain loci containing 12 hVλ gene segments, human κ genomic sequence, and either one or four hJλ

gene segments operably linked to the endogenous mouse Cκ gene which, upon recombination, leads to the formation of a chimeric human λ/mouse κ light chain.

Example III

Engineering Additional Human Vλ Genes Segments Into a Human λ Light Chain Mini-Locus

[00237] Additional hVλ gene segments were added independently to each of the initial modifications described in Example 2 using similar targeting vectors and methods (FIG. 5A, +16-λ Targeting Vector and FIG. 5B, +16-κ Targeting Vector).

[00238] **Introduction of 16 additional human Vλ gene segments.** Upstream (5') homology arms used in constructing targeting vectors for adding 16 additional hVλ gene segments to the modified light chain loci described in Example 2 contained mouse genomic sequence 5' of either the endogenous κ or λ light chain loci. The 3' homology arms were the same for all targeting vectors and contained human genomic sequence overlapping with the 5' end of the human λ sequence of the modifications as described in Example 2.

[00239] Briefly, two targeting vectors were engineered for introduction of 16 additional hVλ gene segments to the modified mouse light chain loci described in Example 2 (FIG. 5A and 5B, +16-λ or +16-κ Targeting Vector). A ~172 kb DNA fragment from human BAC clone RP11-761I13 (Invitrogen) containing 21 consecutive hVλ gene segments from cluster A was engineered with a 5' homology arm containing mouse genomic sequence 5' to either the endogenous κ or λ light chain loci and a 3' homology arm containing human genomic λ sequence. The 5' mouse κ or λ homology arms used in these targeting constructs were the same 5' homology arms described in Example 2 (FIG. 5A and 5B). The 3' homology arm included a 53,057 bp overlap of human genomic λ sequence corresponding to the equivalent 5' end of the ~123 kb fragment of human genomic λ sequence described in Example 2. These two targeting vectors included, from 5' to 3', a 5' mouse homology arm containing either ~23 kb of genomic sequence 5' of the endogenous mouse κ light chain locus or ~24 kb of mouse genomic sequence 5' of the endogenous λ light chain locus, a 5' Frt site, a hygromycin cassette, a 3' Frt site and 171,457 bp of human genomic λ sequence containing 21 consecutive hVλ gene segments, ~53 kb of which overlaps with the 5' end of the human λ sequence described in Example 3 and serves as the 3' homology arm for this targeting construct (FIG. 5A and 5B, +16-λ or +16-κ Targeting Vectors). Homologous recombination with these targeting vectors created independently modified mouse κ and λ light chain loci each containing 28 hVλ gene segments and a hJλ1 gene segment operably

linked to endogenous mouse constant genes (Cκ or Cλ2) which, upon recombination, leads to the formation of a chimeric light chain.

[00240] In a similar fashion, the +16-κ Targeting Vector was also used to introduce the 16 additional hVλ gene segments to the other initial modifications described in Example 2 that incorporated multiple hJλ gene segments with and without an integrated human κ sequence (FIG. 4B). Homologous recombination with this targeting vector at the endogenous mouse κ locus containing the other initial modifications created mouse κ light chain loci containing 28 hVλ gene segments and hJλ1, 2, 3 and 7 gene segments with and without a human Vκ-Jκ genomic sequence operably linked to the endogenous mouse Cκ gene which, upon recombination, leads to the formation of a chimeric λ–κ light chain.

[00241] **Introduction of 12 additional human Vλ gene segments.** Additional hVλ gene segments were added independently to each of the modifications described above using similar targeting vectors and methods. The final locus structure resulting from homologous recombination with targeting vectors containing additional hVλ gene segments are shown in FIG. 7A and 7B.

[00242] Briefly, a targeting vector was engineered for introduction of 12 additional hVλ gene segments to the modified mouse κ and λ light chain loci described above (FIG. 5A and 5B, +12-λ or 12-κ Targeting Vectors). A 93,674 bp DNA fragment from human BAC clone RP11-22I18 (Invitrogen) containing 12 consecutive hVλ gene segments from cluster B was engineered with a 5' homology arm containing mouse genomic sequence 5' to either the endogenous mouse κ or λ light chain loci and a 3' homology arm containing human genomic λ sequence. The 5' homology arms used in this targeting construct were the same 5' homology arms used for the addition of 16 hVλ gene segments described above (FIG. 5A and 5B). The 3' homology arm was made by engineering a PI-SceI site ~3431 bp 5' to the human Vλ3-29P gene segment contained in a 27,468 bp genomic fragment of human λ sequence from BAC clone RP11-761I13. This PI-SceI site served as a ligation point to join the ~94 kb fragment of additional human λ sequence to the ~27 kb fragment of human λ sequence that overlaps with the 5' end of the human λ sequence in the previous modification using the +16-λ or +16-κ Targeting Vectors (FIG. 5A and 5B). These two targeting vectors included, from 5' to 3', a 5' homology arm containing either ~23 kb of mouse genomic sequence 5' of the endogenous κ light chain locus or ~24 kb of mouse genomic sequence 5' of the endogenous λ light chain locus, a 5' Frt site, a neomycin cassette, a 3' Frt site and 121,188 bp of human genomic λ sequence containing 16 hVλ gene segments and a PI-SceI site, ~27 kb of which overlaps with the 5' end of the human λ sequence from the insertion of 16 addition hVλ gene segments and serves as the 3'

homology arm for this targeting construct (FIG. 5A and 5B, +12-λ or 12-κ Targeting Vectors). Homologous recombination with these targeting vectors independently created modified mouse κ and λ light chain loci containing 40 hVλ gene segments and human Jλ1 operably linked to the endogenous mouse constant genes (Cκ or Cλ2) which, upon recombination, leads to the formation of a chimeric light chain (bottom of FIG.5A and 5B).

[00243] In a similar fashion, the +12-κ Targeting Vector was also used to introduce the 12 additional hVλ gene segments to the other initial modifications that incorporated multiple hJλ gene segments with and without an integrated human κ sequence (FIG. 4B). Homologous recombination with this targeting vector at the endogenous mouse κ locus containing the other modifications created a mouse κ light chain locus containing 40 hVλ gene segments and hJλ1, 2, 3 and 7 gene segments with and without a human Vκ-Jκ genomic sequence operably linked to the endogenous mouse Cκ gene which, upon recombination, leads to the formation of a chimeric λ–κ light chain.

Example IV

Identification of targeted ES cells Bearing Human λ Light Chain Gene Segments

[00244] Targeted BAC DNA made according to the foregoing Examples was used to electroporate mouse ES cells to create modified ES cells for generating chimeric mice that express human λ light chain gene segments. ES cells containing an insertion of unrearranged human λ light chain gene segments were identified by a quantitative TAQMAN® assay. Specific primers sets and probes were design for insertion of human λ sequences and associated selection cassettes (gain of allele, GOA), loss of endogenous mouse sequences and any selection cassettes (loss of allele, LOA) and retention of flanking mouse sequences (allele retention, AR). For each additional insertion of human λ sequences, additional primer sets and probes were used to confirm the presence of the additional human λ sequences as well as the previous primer sets and probes used to confirm retention of the previously targeted human sequences. Table 1 sets forth the primers and associated probes used in the quantitative PCR assays. Table 2 sets forth the combinations used for confirming the insertion of each section of human λ light chain gene segments in ES cell clones.

[00245] ES cells bearing the human λ light chain gene segments are optionally transfected with a construct that expresses FLP in order to remove the Frt'ed neomycin cassette introduced by the insertion of the targeting construct containing human Vλ5-52 – Vλ1-40 gene segments (FIG. 5A and 5B). The neomycin cassette may optionally be

removed by breeding to mice that express FLP recombinase (*e.g.*, US 6,774,279). Optionally, the neomycin cassette is retained in the mice.

Table 1

Primer	SEQ ID NO:	Probe	SEQ ID NO:
hL2F	2	hL2P	24
hL2R	3		
hL3F	4	hL3P	25
hL3R	5		
NeoF	6	NeoP	26
NeoR	7		
61hJ1F	8	61hJ1P	27
61hJ1R	9		
67hT1F	10	67hT1P	28
67hT1R	11		
67hT3F	12	67hT3P	29
67hT3R	13		
HygF	14	HygP	30
HygR	15		
MKD2F	16	MKD2P	31
MKD2R	17		
MKP8F	18	MKP8P	32
MKP8R	19		
MKP15F	20	MKP15P	33
MKP15R	21		
MK20F	22	—	—
MKP4R	23		
68h2F	34	68h2P	38
68h2R	35		
68h5F	36	68h5P	39
68h5R	37		
mL1F	75	mL1P	83
mL1R	76		
mL2F	77	mL2P	84
mL2R	78		
mL11F	79	mL11P	85
mL11R	80		

mL12F	81	mL12P	86
mL12R	82		

Table 2

Modification	Assay	Forward/Reverse Primer Set	Probe	Sequence Location
Insertion of 12 hVλ & hJλ1	GOA	hL2F/hL2R	hL2P	hVλ3-12 – hVλ3-1
		hL3F/hL3R	hL3P	
		61hJ1F/61hJ1R	61hJ1P	hJλ sequence
		NeoF/NeoR	NeoP	Neomycin cassette
	LOA	MK20F/MKP4R	—	*lox*511/*lox*P sequence of inactivated κ locus
		HygF/HygR	HygP	Hygromycin cassette from inactivated λ locus
		mL1F/mL1R	mL1P	Mouse Vλ1-Cλ1 Cluster
		mL2F/mL2R	mL2P	
		mL11F/mL11R	mL11P	Mouse Vλ2-Cλ2 Cluster
		mL12F/mL12R	mL12P	
	AR/LOA	MKD2F/MKD2R	MKD2P	Mouse sequence in 5' Vκ locus
		MKP15F/MKP15R	MKP15P	Mouse sequence in 3' Vκ locus
Insertion of 16 hVλ	GOA	67hT1F/67hT1R	67hT1P	hVλ3-27 – hVλ3-12
		67hT3F/67hT3R	67hT3P	
		HygF/HygR	HygP	Hygromycin cassette
	LOA	NeoF/NeoR	NeoP	Neomycin cassette
		mL1F/mL1R	mL1P	Mouse Vλ1-Cλ1 Cluster
		mL2F/mL2R	mL2P	
		mL11F/mL11R	mL11P	Mouse Vλ2-Cλ2 Cluster
		mL12F/mL12R	mL12P	
	AR	hL2F/hL2R	hL2P	hVλ3-12 – hVλ3-1
		hL3F/hL3R	hL3P	
	AR/LOA	MKD2F/MKD2R	MKD2P	Mouse sequence in 5' Vκ locus
		MKP15F/MKP15R	MKP15P	Mouse sequence in 3' Vκ locus

Insertion of 12 hVλ	GOA	68h2F/68h2R	68h2P	hVλ5-52 – hVλ1-40
		68h5F/68h5R	68h5P	
		NeoF/NeoR	NeoP	Neomycin cassette
	LOA	HygF/HygR	HygP	Hygromycin cassette
		mL1F/mL1R	mL1P	Mouse Vλ1-Cλ1 Cluster
		mL2F/mL2R	mL2P	
		mL11F/mL11R	mL11P	Mouse Vλ2-Cλ2 Cluster
		mL12F/mL12R	mL12P	
	AR	hL2F/hL2R	hL2P	hVλ3-12 – hVλ3-1
		hL3F/hL3R	hL3P	
		67hT1F/67hT1R	67hT1P	hVλ3-27 – hVλ3-12
		67hT3F/67hT3R	67hT3P	
	AR/LOA	MKD2F/MKD2R	MKD2P	Mouse sequence in 5' Vκ locus
		MKP15F/MKP15R	MKP15P	Mouse sequence in 3' Vκ locus

Example V

Generation of Mice Expressing Human λ Light Chains From an Endogenous Light Chain Locus

[00246] Targeted ES cells described above were used as donor ES cells and introduced into an 8-cell stage mouse embryo by the VELOCIMOUSE® method (see, *e.g.*, US Pat. No. 7,294,754 and Poueymirou *et al.* (2007) F0 generation mice that are essentially fully derived from the donor gene-targeted ES cells allowing immediate phenotypic analyses Nature Biotech. 25(1):91-99. VELOCIMICE® (F0 mice fully derived from the donor ES cell) independently bearing human λ gene segments were identified by genotyping using a modification of allele assay (Valenzuela *et al., supra*) that detected the presence of the unique human λ gene segments (*supra*).

[00247] **κ:λ light chain usage of mice bearing human λ light chain gene segments.** Mice homozygous for each of three successive insertions of hVλ gene segments with a single hJλ gene segment (FIG. 5B) and mice homozygous for a first insertion of hVλ gene segments with either a single hJλ gene segment or four human Jλ gene segments including a human Vκ-Jκ genomic sequence (FIG. 4B) were analyzed for κ and λ light chain expression in splenocytes using flow cytometry.

[00248] Briefly, spleens were harvested from groups of mice (ranging from three to seven animals per group) and grinded using glass slides. Following lysis of red blood cells (RBCs) with ACK lysis buffer (Lonza Walkersville), splenocytes were stained with fluorescent dye conjugated antibodies specific for mouse CD19 (Clone 1D3; BD

Biosciences), mouse CD3 (17A2; Biolegend), mouse Igκ (187.1; BD Biosciences) and mouse Igλ (RML-42; Biolegend). Data was acquired using a BD™ LSR II flow cytometer (BD Biosciences) and analyzed using FLOWJO™ software (Tree Star, Inc.). Table 3 sets forth the average percent values for B cells ($CD19^+$), κ light chain ($CD19^+Ig\kappa^+Ig\lambda^-$), and λ light chain ($CD19^+Ig\kappa^-Ig\lambda^+$) expression observed in splenocytes from groups of animals bearing each genetic modification.

[00249] In a similar experiment, B cell contents of the splenic compartment from mice homozygous for a first insertion of 12 hVλ and four hJλ gene segments including a human Vκ-Jκ genomic sequence operably linked to the mouse Cκ gene (bottom of FIG. 4B) and mice homozygous for 40 hVλ and one hJλ gene segment (bottom of FIG. 5B or top of FIG. 7B) were analyzed for Igκ and Igλ expression using flow cytometry (as described above). FIG. 8A shows the Igλ and Igκ expression in $CD19^+$ B cells for a representative mouse from each group. The number of $CD19^+$ B cells per spleen was also recorded for each mouse (FIG. 8B).

[00250] In another experiment, B cell contents of the spleen and bone marrow compartments from mice homozygous for 40 hVλ and four hJλ gene segments including a human Vκ-Jκ genomic sequence operably linked to the mouse Cκ gene (bottom of FIG. 7B) were analyzed for progression through B cell development using flow cytometry of various cell surface markers.

[00251] Briefly, two groups (N=3 each, 9-12 weeks old, male and female) of wild type and mice homozygous for 40 hVλ and four hJλ gene segments including a human Vκ-Jκ genomic sequence operably linked to the mouse Cκ gene were sacrificed and spleens and bone marrow were harvested. Bone marrow was collected from femurs by flushing with complete RPMI medium (RPMI medium supplemented with fetal calf serum, sodium pyruvate, Hepes, 2-mercaptoethanol, non-essential amino acids, and gentamycin). RBCs from spleen and bone marrow preparations were lysed with ACK lysis buffer (Lonza Walkersville), followed by washing with complete RPMI medium. $1x10^6$ cells were incubated with anti-mouse CD16/CD32 (2.4G2, BD Biosciences) on ice for 10 minutes, followed by labeling with a selected antibody panel for 30 min on ice.

[00252] Bone marrow panel: anti-mouse FITC-CD43 (1B11, BioLegend), PE-ckit (2B8, BioLegend), PeCy7-IgM (II/41, eBioscience), PerCP-Cy5.5-IgD (11-26c.2a, BioLegend), APC- B220 (RA3-6B2, eBioscience), APC-H7-CD19 (ID3, BD) and Pacific Blue-CD3 (17A2, BioLegend).

[00253] Bone marrow and spleen panel: anti-mouse FITC-Igκ (187.1, BD), PE-Igλ (RML-42, BioLegend), PeCy7-IgM (II/41, ebioscience), PerCP-Cy5.5-IgD (11-26c.2a,

BioLegend), Pacific Blue-CD3 (17A2, BioLegend), APC- B220 (RA3-6B2, eBioscience), APC-H7-CD19 (ID3, BD).

[00254] Following staining, cells were washed and fixed in 2% formaldehyde. Data acquisition was performed on a FACSCANTOII™ flow cytometer (BD Biosciences) and analyzed with FLOWJO™ software (Tree Star, Inc.). FIGs. 9A – 9D show the results for the splenic compartment of one representative mouse from each group. FIGs. 10A – 10E show the results for the bone marrow compartment of one representative mouse from each group. Table 4 sets forth the average percent values for B cells ($CD19^+$), κ light chain ($CD19^+Ig\kappa^+Ig\lambda^-$), and λ light chain ($CD19^+Ig\kappa^-Ig\lambda^+$) expression observed in splenocytes from groups of animals bearing various genetic modifications. Table 5 sets forth the average percent values for B cells ($CD19^+$), mature B cells ($B220^{hi}IgM^+$), immature B cells ($B220^{int}IgM^+$), immature B cells expressing κ light chain ($B220^{int}IgM^+Ig\kappa^+$) and immature B cells expressing λ light chain ($B220^{int}IgM^+Ig\lambda^+$) observed in bone marrow of wild type and mice homozygous for 40 hVλ and four hJλ gene segments including a human Vκ-Jκ genomic sequence operably linked to the mouse Cκ gene. This experiment was repeated with additional groups of the mice described above and demonstrated similar results (data not shown).

Table 3

Genotype	% B cells	% $Ig\kappa^+$	% $Ig\lambda^+$
Wild Type	46.2	91.0	3.6
12 hVλ+hJλ1	28.3	10.4	62.5
12 hVλ-VκJκ-hJλ1	12.0	11.0	67.5
12 hVλ-VκJκ-4hJλ	41.8	17.2	68.4
28 hVλ+hJλ1	22.0	13.3	51.1
40 hVλ+hJλ1	28.2	24.3	53.0

Table 4

Genotype	% B cells	% $Ig\kappa^+$	% $Ig\lambda^+$
Wild Type	49.8	91.2	3.5
40 hVλ-VκJκ-4hJλ	33.3	41.6	43.1

Table 5

Genotype	% B cells	% Mature B cells	% Immature B cells	% Immature Igκ$^+$ B cells	% Immature Igλ$^+$ B cells
Wild Type	62.2	9.2	12.0	79.0	8.84
40hVλ-VκJκ-4hJλ	60.43	2.59	7.69	38.29	43.29

[00255] **Human Vλ gene usage in mice bearing human λ light chain gene segments.** Mice heterozygous for a first insertion of human λ sequences (hVλ3-12 – hVλ3-1 and hJλ1, FIG. 5B) and homozygous for a third insertion of human λ sequences (hVλ5-52 – hVλ3-1 and hJλ1, FIG. 5B) were analyzed for human λ light chain gene usage by reverse-transcriptase polymerase chain reaction (RT-PCR) using RNA isolated from splenocytes.

[00256] Briefly, spleens were harvested and perfused with 10 mL RPMI-1640 (Sigma) with 5% HI-FBS in sterile disposable bags. Each bag containing a single spleen was then placed into a STOMACHER™ (Seward) and homogenized at a medium setting for 30 seconds. Homogenized spleens were filtered using a 0.7μm cell strainer and then pelleted with a centrifuge (1000 rpm for 10 minutes) and RBCs were lysed in BD PHARM LYSE™ (BD Biosciences) for three minutes. Splenocytes were diluted with RPMI-1640 and centrifuged again, followed by resuspension in 1 mL of PBS (Irvine Scientific). RNA was isolated from pelleted splenocytes using standard techniques known in the art.

[00257] RT-PCR was performed on splenocyte RNA using primers specific for human hVλ gene segments and the mouse Cκ gene (Table 6). PCR products were gel-purified and cloned into pCR2.1-TOPO TA vector (Invitrogen) and sequenced with primers M13 Forward (GTAAAACGAC GGCCAG; SEQ ID NO:55) and M13 Reverse (CAGGAAACAG CTATGAC; SEQ ID NO:56) located within the vector at locations flanking the cloning site. Eighty-four total clones derived from the first and third insertions of human λ sequences were sequenced to determine hVλ gene usage (Table 7). The nucleotide sequence of the hVλ-hJλ1-mCκ junction for selected RT-PCR clones is shown in FIG. 11.

[00258] In a similar fashion, mice homozygous for a third insertion of human λ light chain gene sequences (*i.e.* 40 hVλ gene segments and four hJλ gene segments including a human Vκ-Jκ genomic sequence, bottom of FIG. 7B) operably linked to the endogenous mouse Cκ gene were analyzed for human λ light chain gene usage by RT-PCR using RNA

isolated from splenocytes (as described above). The human λ light chain gene segment usage for 26 selected RT-PCR clones are shown in Table 8. The nucleotide sequence of the hVλ-hJλ-mCκ junction for selected RT-PCR clones is shown in FIG. 12.

[00259] In a similar fashion, mice homozygous for a first insertion of human λ light chain gene segments (12 hVλ gene segments and hJλ1, FIG. 4A & FIG. 5A) operably linked to the endogenous mouse Cλ2 gene were analyzed for human λ light chain gene usage by RT-PCR using RNA isolated from splenocytes (as described above). The primers specific for hVλ gene segments (Table 6) were paired with one of two primers specific for the mouse Cλ2 gene; Cλ2-1 (SEQ ID NO:104) or Cλ2-2 (SEQ ID NO:105).

[00260] Multiple hVλ gene segments rearranged to hλ1 were observed from the RT-PCR clones from mice bearing human λ light chain gene segments at the endogenous mouse λ light chain locus. The nucleotide sequence of the hVλ-hJλ-mCλ2 junction for selected RT-PCR clones is shown in FIG. 13.

Table 6

5' hVλ Primer	Sequence (5'-3')	SEQ ID NO:
VLL-1	CCTCTCCTCC TCACCCTCCT	40
VLL-1n	ATGRCCDGST YYYCTCTCCT	41
VLL-2	CTCCTCACTC AGGGCACA	42
VLL-2n	ATGGCCTGGG CTCTGCTSCT	43
VLL-3	ATGGCCTGGA YCSCTCTCC	44
VLL-4	TCACCATGGC YTGGRYCYCM YTC	45
VLL-4.3	TCACCATGGC CTGGGTCTCC TT	46
VLL-5	TCACCATGGC CTGGAMTCYT CT	47
VLL-6	TCACCATGGC CTGGGCTCCA CTACTT	48
VLL-7	TCACCATGGC CTGGACTCCT	49
VLL-8	TCACCATGGC CTGGATGATG CTT	50
VLL-9	TAAATATGGC CTGGGCTCCT CT	51
VLL-10	TCACCATGCC CTGGGCTCTG CT	52
VLL-11	TCACCATGGC CCTGACTCCT CT	53

3' Mouse Cκ Primer	Sequence (5'-3')	SEQ ID NO:
mIgKC3'-1	CCCAAGCTTA CTGGATGGTG GGAAGATGGA	54

Table 7

hVλ	Observed No. of Clones
3-1	2
4-3	3
2-8	7
3-9	4
3-10	3
2-14	1
3-19	1
2-23	7
3-25	1
1-40	9
7-43	2
1-44	2
5-45	8
7-46	3
9-49	6
1-51	3

Table 8

Clone	hVλ	hJλ
1-3	1-44	7
1-5	1-51	3
2-3	9-49	7
2-5	1-40	1
2-6	1-40	7
3b-5	3-1	7
4a-1	4-3	7
4a-5	4-3	7
4b-1	1-47	3
5-1	3-10	3
5-2	1-40	7
5-3	1-40	7
5-4	7-46	2
5-6	1-40	7
5-7	7-43	3
6-1	1-40	1
6-2	1-40	2
6-7	1-40	3
7a-1	3-10	7
7a-2	9-49	2
7a-7	3-10	7
7b-2	7-43	3
7b-7	7-46	7
7b-8	7-43	3
11a-1	5-45	2
11a-2	5-45	7

[00261] FIG. 11 shows the sequence of the hVλ-hJλ1-mCκ junction for RT-PCR clones from mice bearing a first and third insertion of hVλ gene segments with a single hJλ gene segment. The sequences shown in FIG. 11 illustrate unique rearrangements involving different hVλ gene segments with hJλ1 recombined to the mouse Cκ gene. Heterozygous mice bearing a single modified endogenous κ locus containing 12 hVλ gene segments and hJλ1 and homozygous mice bearing two modified endogenous κ loci containing 40 hVλ gene segments and hJλ1 were both able to produce human λ gene segments operably

linked to the mouse Cκ gene and produce B cells that expressed human λ light chains. These rearrangements demonstrate that the chimeric loci were able to independently rearrange human λ gene segments in multiple, independent B cells in these mice. Further, these modifications to the endogenous κ light chain locus did not render any of the hVλ gene segments inoperable or prevent the chimeric locus from recombining multiple hVλ and a hJλ (Jλ1) gene segment during B cell development as evidenced by 16 different hVλ gene segments that were observed to rearrange with hJλ1 (Table 7). Further, these mice made functional antibodies containing rearranged human Vλ–Jλ gene segments operably linked to mouse Cκ genes as part of the endogenous immunoglobulin light chain repertoire.

[00262] FIG. 12 shows the sequence of the hVλ-hJλ-mCκ junction for selected RT-PCR clones from mice homozygous for 40 hVλ and four hJλ gene segments including a human Vκ-Jκ genomic sequence. The sequences shown in FIG. 12 illustrate additional unique rearrangements involving multiple different hVλ gene segments, spanning the entire chimeric locus, with multiple different hJλ gene segments rearranged and operably linked to the mouse Cκ gene. Homozygous mice bearing modified endogenous κ loci containing 40 hVλ and four hJλ gene segments were also able to produce human λ gene segments operably linked to the mouse Cκ gene and produce B cells that expressed human λ light chains. These rearrangements further demonstrate that the all stages of chimeric loci were able to independently rearrange human λ gene segments in multiple, independent B cells in these mice. Further, these additional modifications to the endogenous κ light chain locus demonstrates that each insertion of human λ gene segments did not render any of the hVλ and/or Jλ gene segments inoperable or prevent the chimeric locus from recombining the hVλ and Jλ gene segments during B cell development as evidenced by 12 different hVλ gene segments that were observed to rearrange with all four hJλ gene segments (Table 8) from the 26 selected RT-PCR clone. Further, these mice as well made functional antibodies containing human Vλ–Jλ gene segments operably linked to mouse Cκ regions as part of the endogenous immunoglobulin light chain repertoire.

[00263] FIG. 13 shows the sequence of the hVλ-hJλ-mCλ2 junction for three individual RT-PCR clones from mice homozygous for 12 hVλ gene segments and hJλ1. The sequences shown in FIG. 13 illustrate additional unique rearrangements involving different hVλ gene segments, spanning the length of the first insertion, with hJλ1 rearranged and operably linked to the mouse Cλ2 gene (2D1 = Vλ2-8Jλ1; 2D9 = Vλ3-10Jλ1; 3E15 = Vλ3-1Jλ1). One clone demonstrated a nonproductive rearrangement due to N additions at the hVλ-hJλ junction (2D1, FIG.13). This is not uncommon in V(D)J recombination, as the

joining of gene segments during recombination has been shown to be imprecise. Although this clone represents an unproductive recombinant present in the light chain repertoire of these mice, this demonstrates that the genetic mechanism that contributes to junctional diversity among antibody genes is operating normally in these mice and leading to an antibody repertoire containing light chains with greater diversity.

[00264] Homozygous mice bearing modified endogenous λ loci containing 12 hVλ gene segments and hJλ1 were also able to produce human λ gene segments operably linked to an endogenous mouse Cλ gene and produce B cells that expressed reverse chimeric λ light chains containing hVλ regions linked to mouse Cλ regions. These rearrangements further demonstrate that human λ light chain gene segments placed at the other light chain locus (*i.e.*, the λ locus) were able to independently rearrange human λ gene segments in multiple, independent B cells in these mice. Further, the modifications to the endogenous λ light chain locus demonstrate that the insertion of human λ gene segments did not render any of the hVλ and/or hJλ1 gene segments inoperable or prevent the chimeric locus from recombining the hVλ and hJλ1 gene segments during B cell development. Further, these mice also made functional antibodies containing human Vλ–Jλ gene segments operably linked to a mouse Cλ region as part of the endogenous immunoglobulin light chain repertoire.

[00265] As shown in this Example, mice bearing human λ light chain gene segments at the endogenous κ and λ light chain loci are capable of rearranging human λ light chain gene segments and expressing them in the context of a mouse Cκ and/or Cλ region as part of the normal antibody repertoire of the mouse because a functional light chain is required at various checkpoints in B cell development in both the spleen and bone marrow. Further, early subsets of B cells (*e.g.*, pre-, pro- and transitional B cells) demonstrate a normal phenotype in these mice as compared to wild type littermates (FIGs. 9D, 10A and 10B). A small deficit in bone marrow and peripheral B cell populations was observed, which may be attributed to a deletion of a subset of auto-reactive immature B cells and/or a suboptimal association of human λ light chain with mouse heavy chain. However, the Igκ/Igλ usage observed in these mice demonstrates a situation that is more like human light chain expression than that observed in mice.

Example VI

Breeding of Mice Expressing Human λ Light Chains From an Endogenous Light Chain Locus

[00266] To optimize the usage of the human λ gene segments at an endogenous mouse light chain locus, mice bearing the unrearranged human λ gene segments are bred

to another mouse containing a deletion in the opposing endogenous light chain locus (either κ or λ). For example, human λ gene segments positioned at the endogenous κ locus would be the only functional light chain gene segments present in a mouse that also carried a deletion in the endogenous λ light chain locus. In this manner, the progeny obtained would express only human λ light chains as described in the foregoing examples. Breeding is performed by standard techniques recognized in the art and, alternatively, by commercial companies, *e.g.*, The Jackson Laboratory. Mouse strains bearing human λ light chain gene segments at the endogenous κ locus and a deletion of the endogenous λ light chain locus are screened for presence of the unique reverse-chimeric (human-mouse) λ light chains and absence of endogenous mouse λ light chains.

[00267] Mice bearing an unrearranged human λ light chain locus are also bred with mice that contain a replacement of the endogenous mouse heavy chain variable gene locus with the human heavy chain variable gene locus (see US 6,596,541, Regeneron Pharmaceuticals, the VELOCIMMUNE® genetically engineered mouse). The VELOCIMMUNE® mouse includes, in part, having a genome comprising human heavy chain variable regions operably linked to endogenous mouse constant region loci such that the mouse produces antibodies comprising a human heavy chain variable region and a mouse heavy chain constant region in response to antigenic stimulation. The DNA encoding the variable regions of the heavy chains of the antibodies can be isolated and operably linked to DNA encoding the human heavy chain constant regions. The DNA can then be expressed in a cell capable of expressing the fully human heavy chain of the antibody. Upon a suitable breeding schedule, mice bearing a replacement of the endogenous mouse heavy chain locus with the human heavy chain locus and an unrearranged human λ light chain locus at the endogenous κ light chain locus is obtained. Antibodies containing somatically mutated human heavy chain variable regions and human λ light chain variable regions can be isolated upon immunization with an antigen of interest.

Example VII

Generation of Antibodies From Mice Expressing Human Heavy Chains and Human λ Light Chains

[00268] After breeding mice that contain the unrearranged human λ light chain locus to various desired strains containing modifications and deletions of other endogenous Ig loci (as described above), selected mice are immunized with an antigen of interest.

[00269] Generally, a VELOCIMMUNE® mouse containing one of the single rearranged human germline light chain regions is challenged with an antigen, and lymphatic cells (such as B-cells) are recovered from serum of the animals. The lymphatic cells may be

fused with a myeloma cell line to prepare immortal hybridoma cell lines, and such hybridoma cell lines are screened and selected to identify hybridoma cell lines that produce antibodies containing human heavy chain and human λ light chain that are specific to the antigen used for immunization. DNA encoding the variable regions of the heavy chains and the λ light chains may be isolated and linked to desirable isotypic constant regions of the heavy chain and light chain. Due to the presence of the additional hVλ gene segments as compared to the endogenous mouse λ locus, the diversity of the light chain repertoire is dramatically increased and confers higher diversity on the antigen-specific repertoire upon immunization. The resulting cloned antibody sequences may be subsequently produced in a cell, such as a CHO cell. Alternatively, DNA encoding the antigen-specific chimeric antibodies or the variable domains of the light and heavy chains may be isolated directly from antigen-specific lymphocytes (*e.g.*, B cells).

[00270] Initially, high affinity chimeric antibodies are isolated having a human variable region and a mouse constant region. As described above, the antibodies are characterized and selected for desirable characteristics, including affinity, selectivity, epitope, *etc.* The mouse constant regions are replaced with a desired human constant region to generate the fully human antibody containing a somatically mutated human heavy chain and a human λ light chain derived from an unrearranged human λ light chain locus of the invention. Suitable human constant regions include, for example wild type or modified IgG1, IgG2, IgG3, or IgG4.

We Claim:

1. A method for making an antibody comprising a human lambda, λ, variable domain, the method comprising the steps of:

(a) exposing a genetically modified mouse to an antigen, wherein the genetically modified mouse has a genome that comprises an endogenous λ light chain immunoglobulin locus comprising:

i) a deletion of a first endogenous lambda variable-lambda joining-lambda constant, Vλ-Jλ-Cλ, gene cluster; and

ii) a replacement of a fragment of endogenous Vλ and Jλ gene segments in a second endogenous Vλ-Jλ-Cλ gene cluster with human Vλ and Jλ gene segments, wherein the human Vλ and Jλ gene segments comprise at least one human Vλ gene segment and at least one human Jλ gene segment, and wherein the human Vλ and Jλ gene segments are operably linked to a mouse λ constant (Cλ) gene;

(b) allowing the genetically modified mouse to develop an immune response to the antigen; and

(c) isolating from the mouse of (b) an antibody specific for the antigen, wherein the antibody comprises a human λ variable domain derived from the human Vλ and Jλ gene segments.

2. The method of claim 1, wherein the genetically modified mouse has a genome that comprises one or more human heavy chain V, D, and J segments.

3. The method of claim 2, wherein the one or more human heavy chain V, D, and J segments are at an endogenous mouse heavy chain locus.

4. The method of claim 1, wherein the genetically modified mouse has a genome comprising a replacement at the endogenous mouse heavy chain variable locus of one or more endogenous heavy chain V, D, and J segments with one or more human heavy chain V, D, and J segments.

5. The method of any one of claims 1 to 4, wherein the human Vλ and Jλ gene segments comprise a human Jλ1 gene segment.

6. The method of any one of claims 1 to 4, wherein the human Vλ and Jλ gene segments comprise four human Jλ gene segments.

7. The method of claim 6, wherein the four human Jλ gene segments are Jλ1, Jλ2, Jλ3 and Jλ7.

8. The method of any one of claims 1 to 4, wherein the human Vλ and Jλ gene segments comprise at least 12 human Vλ gene segments.

9. The method of claim 8, wherein the 12 human V gene segments include human Vλ3-1, Vλ4-3, Vλ2-8, Vλ3-9, Vλ3-10, Vλ2-11 and Vλ3-12.

10. The method of any one of claims 1 to 4, wherein the human Vλ and Jλ gene segments comprise at least 28 human Vλ gene segments.

11. The method of claim 10, wherein the 28 human Vλ gene segments include human Vλ2-14, Vλ3-16, Vλ2-18, Vλ3-19, Vλ3-21, Vλ3-22, Vλ2-23, Vλ3-25 and Vλ3-27.

12. The method of any one of claims 1 to 4, wherein the human Vλ and Jλ gene segments comprise at least 40 human Vλ gene segments.

13. The method of claim 12, wherein the 40 human V gene segments include human Vλ5-52, Vλ1-51, Vλ9-49, Vλ1-47, Vλ7-46, Vλ5-45, Vλ1-44, Vλ7-43 and Vλ1-40.

14. The method of any one of claims 1 to 13, wherein the endogenous mouse Vλ gene segments of the second endogenous Vλ-Jλ-Cλ gene cluster have been replaced with a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ3-12 gene segment to a human Vλ3-1 gene segment.

15. The method of any one of claims 1 to 13, wherein the endogenous mouse Vλ gene segments of the second endogenous Vλ-Jλ-Cλ gene cluster have been replaced with a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ3-27 gene segment to a human Vλ3-1 gene segment.

16. The method of any one of claims 1 to 13, wherein the endogenous mouse Vλ gene segments of the second endogenous Vλ-Jλ-Cλ gene cluster have been replaced with a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ3-27 gene segment to a human Vλ3-1 gene segment and a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ5-52 gene segment to a human Vλ1-40 gene segment.

17. A method for making a cell that expresses an antibody comprising a human lambda, λ, variable domain, comprising the steps of:

(a) exposing a genetically modified mouse to an antigen, wherein the genetically modified mouse has a genome that comprises an endogenous λ light chain immunoglobulin locus comprising:

i) a deletion of a first endogenous lambda variable-lambda joining-lambda constant, Vλ-Jλ-Cλ, gene cluster; and

ii) a replacement of a fragment of endogenous Vλ and Jλ gene segments in a second endogenous Vλ-Jλ-Cλ gene cluster with human Vλ and Jλ gene segments, wherein the human Vλ and Jλ gene segments comprise at least one human Vλ gene segment and at least one human Jλ gene segment, and wherein the human Vλ and Jλ gene segments are operably linked to a mouse λ constant (Cλ) gene;

(b) allowing the genetically modified mouse to develop an immune response to the antigen; and

(c) isolating from the mouse of (b) a cell that expresses an antibody specific for the antigen, wherein the antibody comprises a human λ variable domain derived from the human Vλ and Jλ gene segments.

18. The method of claim 17, further comprising fusing the cell with a myeloma cell to produce a hybridoma.

19. The method of claim 17, further comprising fusing the cell with a myeloma cell to produce a quadroma.

20. The method of claim 17, wherein the cell is a B cell.

21. The method of any one of claims 17 to 19, wherein the cell has a genome that comprises one or more human heavy chain V, D, and J segments.

22. The method of claim 21, wherein the one or more human heavy chain V, D, and J segments are at an endogenous mouse heavy chain locus.

23. The method of any one of claims 17 to 19, wherein the cell has a genome comprising a replacement at the endogenous mouse heavy chain variable locus of one or more endogenous heavy chain V, D, and J segments with one or more human heavy chain V, D, and J segments.

24. The method of any one of claims 17 to 19, wherein the human Vλ and Jλ gene segments comprise a human Jλ1 gene segment.

25. The method of any one of claims 17 to 19, wherein the human Vλ and Jλ gene segments comprise four human Jλ gene segments.

26. The method of claim 25, wherein the four human Jλ gene segments are Jλ1, Jλ2, Jλ3 and Jλ7.

27. The method of any one of claims 17 to 19, wherein the human Vλ and Jλ gene segments comprise at least 12 human Vλ gene segments.

28. The method of claim 27, wherein the 12 human V gene segments include human Vλ3-1, Vλ4-3, Vλ2-8, Vλ3-9, Vλ3-10, Vλ2-11 and Vλ3-12.

29. The method of any one of claims 17 to 19, wherein the human Vλ and Jλ gene segments comprise at least 28 human Vλ gene segments.

30. The method of claim 29, wherein the 28 human Vλ gene segments include human Vλ2-14, Vλ3-16, Vλ2-18, Vλ3-19, Vλ3-21, Vλ3-22, Vλ2-23, Vλ3-25 and Vλ3-27.

31. The method of any one of claims 17 to 19, wherein the human Vλ and Jλ gene segments comprise at least 40 human Vλ gene segments.

32. The method of claim 31, wherein the 40 human Vλ gene segments include human Vλ5-52, Vλ1-51, Vλ9-49, Vλ1-47, Vλ7-46, Vλ5-45, Vλ1-44, Vλ7-43 and Vλ1-40.

33. The method of any one of claims 17 to 32, wherein the endogenous mouse Vλ gene segments of the second endogenous Vλ-Jλ-Cλ gene cluster have been replaced with a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ3-12 gene segment to a human Vλ3-1 gene segment.

34. The method of any one of claims 17 to 32, wherein the endogenous mouse Vλ gene segments of the second endogenous Vλ-Jλ-Cλ gene cluster have been replaced with a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ3-27 gene segment to a human Vλ3-1 gene segment.

35. The method of any one of claims 17 to 32, wherein the endogenous mouse Vλ gene segments of the second endogenous Vλ-Jλ-Cλ gene cluster have been replaced with a contiguous sequence of the human immunoglobulin light chain locus that spans from a human Vλ3-27 gene segment to a human Vλ3-1 gene segment and a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ5-52 gene segment to a human Vλ1-40 gene segment.

36. A method for cloning a lambda, λ, light chain variable domain, comprising the steps of:

(a) obtaining a cell from a transgenic mouse whose genome comprises an endogenous λ light chain immunoglobulin locus comprising:

i) a deletion of a first endogenous lambda variable-lambda joining-lambda constant, Vλ-Jλ-Cλ, gene cluster; and

ii) a replacement of a fragment of endogenous Vλ and Jλ gene segments in a second endogenous Vλ-Jλ-Cλ gene cluster with human Vλ and Jλ gene segments, wherein the human Vλ and Jλ gene segments comprise at least one human Vλ gene segment and at least one human Jλ gene segment, and wherein the human Vλ and Jλ gene segments are operably linked to a mouse λ constant (Cλ) gene; and

(b) cloning a λ light chain variable domain derived from the human Vλ and Jλ gene segments.

37. The method of claim 36, further comprising fusing the cell with a myeloma cell to produce a hybridoma.

38. The method of claim 36, further comprising fusing the cell with a myeloma cell to produce a quadroma.

39. The method of claim 36, wherein the cell is a B cell.

40. The method of any one of claims 36 to 39, wherein the cell has a genome that comprises one or more human heavy chain V, D, and J segments.

41. The method of claim 40, wherein the one or more human heavy chain V, D, and J segments are at an endogenous mouse heavy chain locus.

42. The method of claim 40, further comprising cloning from the cell a human heavy chain variable (V_H) region nucleic acid sequence that is cognate with the human Vλ region nucleic acid sequence.

43. The method of claim 42, further comprising employing the nucleic acid sequences of the human V_H and human Vλ regions fused to human constant regions to make a human antibody.

44. The method of any one of claims 36 to 39, wherein the cell has a genome comprising a replacement at the endogenous mouse heavy chain variable locus of one or more endogenous heavy chain V, D, and J segments with one or more human heavy chain V, D, and J segments.

45. The method of any one of claims 36 to 44, wherein the human Vλ and Jλ gene segments comprise a human Jλ1 gene segment.

46. The method of any one of claims 36 to 44, wherein the human Vλ and Jλ gene segments comprise at least 12 human Vλ gene segments.

47. The method of claim 46, wherein the 12 human Vλ gene segments include human Vλ3-1, Vλ4-3, Vλ2-8, Vλ3-9, Vλ3-10, Vλ2-11 and Vλ3-12.

48. The method of any one of claims 36 to 44, wherein the human Vλ and Jλ gene segments comprise at least 28 human Vλ gene segments.

49. The method of claim 48, wherein the 28 human Vλ gene segments include human Vλ2-14, Vλ3-16, Vλ2-18, Vλ3-19, Vλ3-21, Vλ3-22, Vλ2-23, Vλ3-25 and Vλ3-27.

50. The method of any one of claims 36 to 44, wherein the human Vλ and Jλ gene segments comprise at least 40 human Vλ gene segments.

51. The method of claim 50, wherein the 40 human Vλ gene segments include human Vλ5-52, Vλ1-51, Vλ9-49, Vλ1-47, Vλ7-46, Vλ5-45, Vλ1-44, Vλ7-43 and Vλ1-40.

52. The method of any one of claims 36 to 44, wherein the human Vλ and Jλ gene segments comprise four human Jλ gene segments.

53. The method of claim 52, wherein the four human Jλ gene segments are Jλ1, Jλ2, Jλ3 and Jλ7.

54. The method of any one of claims 36 to 53, wherein the endogenous mouse Vλ gene segments of the second endogenous Vλ-Jλ-Cλ gene cluster have been replaced with a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ3-12 gene segment to a human Vλ3-1 gene segment.

55. The method of any one of claims 36 to 53, wherein the endogenous mouse Vλ gene segments of the second endogenous Vλ-Jλ-Cλ gene cluster have been replaced with a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ3-27 gene segment to a human Vλ3-1 gene segment.

56. The method of any one of claims 36 to 53, wherein the endogenous mouse Vλ gene segments of the second endogenous Vλ-Jλ-Cλ gene cluster have been replaced with a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ3-27 gene segment to a human Vλ3-1 gene segment and a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ5-52 gene segment to a human Vλ1-40 gene segment.

57. A method of making a transgenic mouse whose genome comprises an endogenous, lambda, λ, light chain immunoglobulin locus, the method comprising:

a) genetically engineering a mouse embryonic stem cell by:

i) a deletion of a first endogenous lambda variable-lambda joining-lambda constant, Vλ-Jλ-Cλ, gene cluster; and

ii) a replacement of a fragment of endogenous Vλ and Jλ gene segments in a second endogenous Vλ-Jλ-Cλ gene cluster with human Vλ and Jλ gene segments, wherein the human Vλ and Jλ gene segments comprise at least one human Vλ gene segment and at least one human Jλ gene segment, and wherein the human Vλ and Jλ gene segments are operably linked to a mouse Cλ gene;

b) introducing the genetically engineered stem cell into a mouse embryo; and

c) generating the transgenic mouse from the embryo.

58. The method of claim 57, wherein the human Vλ and Jλ gene segments comprise a human Jλ1 gene segment.

59. The method of claim 57, wherein the human Vλ and Jλ gene segments comprise four human Jλ gene segments.

60. The method of claim 59, wherein the four human Jλ gene segments are Jλ1, Jλ2, Jλ3 and Jλ7.

61. The method of claim 57, wherein the human Vλ and Jλ gene segments comprise at least 12 human Vλ gene segments.

62. The method of claim 61, wherein the 12 human Vλ gene segments include human Vλ3-1, Vλ4-3, Vλ2-8, Vλ3-9, Vλ3-10, Vλ2-11 and Vλ3-12.

63. The method of claim 57, wherein the human Vλ and Jλ gene segments comprise at least 28 human Vλ gene segments.

64. The method of claim 63, wherein the 28 human Vλ gene segments include human Vλ2-14, Vλ3-16, Vλ2-18, Vλ3-19, Vλ3-21, Vλ3-22, Vλ2-23, Vλ3-25 and Vλ3-27.

65. The method of claim 57, wherein the human Vλ and Jλ gene segments comprise at least 40 human Vλ gene segments.

66. The method of claim 65, wherein the 40 human Vλ gene segments include human Vλ5-52, Vλ1-51, Vλ9-49, Vλ1-47, Vλ7-46, Vλ5-45, Vλ1-44, Vλ7-43 and Vλ1-40.

67. The method of any one of claims 57 to 66, wherein the endogenous mouse Vλ gene segments of the second endogenous Vλ-Jλ-Cλ gene cluster are replaced with a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ3-12 gene segment to a human Vλ3-1 gene segment.

68. The method of any one of claims 57 to 66, wherein the replacement endogenous mouse Vλ gene segments of the second endogenous Vλ-Jλ-Cλ gene cluster are replaced with a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ3-27 gene segment to a human Vλ3-1 gene segment.

69. The method of any one of claims 57 to 66, wherein the replacement endogenous mouse Vλ gene segments of the second endogenous Vλ-Jλ-Cλ gene cluster are replaced with a contiguous sequence of the human immunoglobulin λ light chain locus that spans from a human Vλ3-27 gene segment to a human Vλ3-1 gene segment and a contiguous sequence of the

human immunoglobulin λ light chain locus that spans from a human Vλ5-52 gene segment to a human Vλ1-40 gene segment.


Human λ Light Chain Locus
0 kb
PPC2-like
top3B
4-69
8-61
thioesterase
Cluster C
250 kb
6-57
10-54
11-55
VPREB1
top3B.2
Cluster B
1-51
5-52
9-49
1-50
5-48
7-46
1-47
5-45
1-44
7-43
1-40
5-39
5-37
1-36
500 kb
Cluster A
500 kb
3-25
2-23
750 kb
ZNF279
ZNF280
PRAME
2-33
3-32
γ-glutamyltranspeptidase
BCRL 4
3-27
750 kb
3-22
3-21
3-19
2-18
3-16
2-14
3-12
3-10
2-11
3-9
2-8
4-3
3-1
Jλ-Cλ pairs
1000 kb

FIG. 1


Enh 3.1
Enh 3.1
Enh
Enh
V1
J3
C3
J1
C1
J3P
Targeting Vector 1
loxP
Neo
loxP
+ Cre
Enh 2.4
Enh 2.4
Mouse λ Light Chain Locus
J2
C2
C4P
J4P
V2
V3
Targeting Vector 2
Frt
Hyg-TK
Frt
Inactivated Mouse λ Light Chain Locus
Enh 3.1
Enh
loxP
Enh 2.4
C2
C4P
J4P
Frt
Hyg-TK
Frt

FIG. 2


Cκ
Eκi
Jκ
Vκ
Hyg-TK
loxP
Jκ Targeting Vector
+ Cre
Cκ
Eκi
loxP
511
Deleted Mouse κ Light Chain Locus
3 MB
Vκ
Neo
loxP
511
Vκ Targeting Vector
Mouse κ Light Chain Locus

FIG. 3


Inactivated Mouse λ Light Chain Locus
Enh 3.1
Enh
lox
C2-J4-C4
Enh 2.4
Frt
Hyg-TK
hVλ
3-12
hVλ
3-1
C2-J4-C4
Enh 2.4
hJλ1
PI-SceI
Frt
Neo
Frt
12/1-λ
Targeting Vector

FIG. 4A


Deleted Mouse κ Light Chain Locus
Cκ
Eκi
Eκ3'
lox P
511
hVλ 3-12
hVλ 3-1
hJλ1
PI-SceI
Frt
Neo
I-CeuI
hJλ 1,2,3,7
hVκ-Jκ
NotI
12/1-κ Targeting Vector
12/4-κ Targeting Vector
12(κ)1-κ Targeting Vector
12(κ)4-κ Targeting Vector

FIG. 4B


Inactivated Mouse λ Light Chain Locus
Enh 3.1
Enh
C2-J4-C4
Enh 2.4
Frt
Hyg-TK
Frt
lox
Frt
Neo
12/1-λ Targeting Vector
hVλ 3-12
hVλ 3-1
C2-J4-C4
Enh 2.4
hJλ1
PI-SceI
Frt
Hyg
Frt
+16-λ Targeting Vector
hVλ 3-12
hVλ 3-27
hVλ 1-40
PI-SceI
Frt
Neo
Frt
+12-λ Targeting Vector
hVλ 3-12
hVλ 3-1
hVλ 1-40 3-27
hVλ 5-52
PI-SceI
Frt
Neo
Frt

FIG. 5A


Deleted Mouse κ Light Chain Locus
Eκ3'
Cκ
lox P
511
Eκi
hVλ
3-12
Frt
Neo
I-CeuI
12/1-κ
Targeting Vector
hVλ
3-1
hJλ1
Eκi
PI-SceI
Cκ
3-12
3-27
Frt
Hyg
I-CeuI
+16-κ Targeting Vector
5-52
Frt
Neo
I-CeuI
1-40
PI-SceI
+12-κ Targeting Vector
Eκ3'
3-12
3-1
Jλ1
Eκi
PI-SceI
Cκ
1-40
3-27
PI-SceI
5-52
Frt
Neo
I-CeuI

FIG. 5B


hJκ
hJλ1
Chlor
AsiSI
PI-SceI
OR
hJλ 1,2,3,7
hVκ
Frt
Hyg
NotI
hVλ
Spec
Neo
I-CeuI
hVκ-Jκ
1. Digest with NotI & AsiSI
2. Ligate Fragments
Ligation of NotI – AsiSI Fragments

FIG. 6


Enh 3.1
Enh
lox
Enh 2.4
C2-J4-C4
hJλ1
PI-SceI
hVλ 3-1
hVλ 3-12
hVλ 1-40
hVλ 3-27
PI-SceI
hVλ 5-52
Frt
Neo
Frt

FIG. 7A


Eκi Cκ
Eκ3'
PI-SceI
hJλ1
hVλ 3-1
hVλ 3-12
NotI
hJλ 1,2,3,7
hVκ-Jκ
hVλ 1-40 3-27
hVλ 5-52
Frt
Neo

FIG. 7B



FIG. 8A


cells/spleen x 10^6
60
40
20
0
WT
12hVλ-VκJκ
40hVλ-1hJλ
2hVλ-4hJλ

FIG. 8B



FIG. 9A


% CD19+Igλ+
50
40
30
20
10
0
WT
40hVλ-Vκ Jκ-4hJλ
% CD19+Igκ+
100
80
60
40
20
0
WT
40hVλ-Vκ Jκ-4hJλ
% CD19+
60
40
20
0
WT
40hVλ-Vκ Jκ-4hJλ

FIG. 9B



FIG. 9C


Mature B cells
(CD19+ IgM^lo IgD^hi)
cells/spleen (x10^-6)
20
15
10
5
0
WT
40hVλ-
VκJκ-4hJλ
Transitional B cells
(CD19+ IgM^hi IgD^lo)
cells/spleen (x10^-6)
5
4
3
2
1
0
WT
40hVλ-
VκJκ-4hJλ
Total CD19+ B cells
cells/spleen (x10^-7)
3
2
1
0
WT
40hVλ-
VκJκ-4hJλ

FIG. 9D


40hVλ-VκJκ-4hJλ
WT
64
4
62
3.7
CD19
CD3
0.31
2.5
Pro
Pre
74
23
0.52
2
Pro
Pre
81
16
ckit
CD43

FIG. 10A


Pre B cells
(CD19+CD43-ckit-)
cells/femur (x10-6)
2.5
2.0
1.5
1.0
WT
40hVλ-VκJκ-4hJλ
Pro B cells
(CD19+CD43+ckit+)
cells/femur (x10-4)
20
15
10
5
0
WT
40hVλ-VκJκ-4hJλ

FIG. 10B


40hVλ-VκJκ-4hJλ
Immature
6.9
Mature
3.8
Pro/pre
64
B220
WT
Immature
13
Mature
11
Pro/pre
48
IgM

FIG. 10C


Mature B cells
(B220hi IgM+)
cells/femur (x10-5)
5
4
3
2
1
0
WT
40hVλ-
VκJκ-4hJλ
Immature B cells
(B220int IgM+)
8
6
4
2
0
WT
40hVλ-
VκJκ-4hJλ
cells/femur (x10-5)

FIG. 10D


Igλ
Igκ
Mature
Immature
WT
40hVλ-VκJκ-4hJλ
3.1
95
0.61
40
54
4.8
8.7
80
9.3
47
39
12

FIG. 10E

	3' Human Vλ	Human Jλ1	5' Mouse Cκ
A6	GCAACAATT	tcGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
B6	GCAACAATT	ATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
F6	GCAACAATT	ATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
B7	GCAACAATT	ATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
E7	GCAACAAT	GTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
F7	GCAACAATT	ATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
C8	GCAACAATTT	ATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
E12	CAAGTCGGTT	gTGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
1-4	TGAGTGCT	TATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
1-20	TGAGTGCg	gcttttTTtGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
3B43	CTGAATGGT	TATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
5-8	AGTGGTAAT	cATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
5-19	AGTGGTGCT	TATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
1010	AGCAGCACT	TATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATCTTC
11A1	AGCAGCGCT	TATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTA
7A8	GGTGGTGCT	TATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCC
3A3	AGTAGCACT	TATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATC
2-7	AGCAGCACT	TATGTCTTCGGAACTGGGACCAAGGTCACCGgCCTgG	GGGCTGATGCTGCACCA
FWR4		**F G T G T K V T V L G**	**A D A A P T V S I F**

FIG. 11

	3' Human Vλ	Human Jλ	5' Mouse Cκ
5-2	CAGCCTGAGTGGTTC	TGTGTTCGGAGGAGGCACCCGGCTGACCGCCCTCG	GGGCTGATGCTGCACCAACTGTATCCATC
2-5	CAGCCTGAGTGGTT	ATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATC
1-3	CAGCCTGAATGGT	GCTGTGTTCGGAGGAGGCACCCAGCTGACCGCCCTCG	GGGCTGATGCTGCACCAACTGTATCCATC
4B-1	CAGCCTGAGTGGTC	GGGTGTTCGGCGGAGGGACCAAGCTGACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATC
3B-5	CAGCAGCACTGC	TGTGTTCGGAGGAGGCACCCAGCTGACCGCCCTCG	GGGCTGATGCTGCACCAACTGTATCCATC
7A-1	CAGCAGTGGTAAT	GCTGTGTTCGGAGGAGGCACCCAGCTGACCGCCCTCG	GGGCTGATGCTGCACCAACTGTATCCATC
5-1	CAGCAGTGGTAATCATAG	GGTGTTCGGCGGAGGGACCAAGCTGACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATC
4A-1	CAGCCTGAGTGGTT	ATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATC
11A-1	CAGCAGCGCT	GTGGTATTCGGCGGAGGGACCAAGCTGACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATC
5-7	CTACTATGGTGGTGCTC	GGGTGTTCGGCGGAGGGACCAAGCTGACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATC
5-4	CTCCTATAGTGGTGCTCGa	GTATTCGGCGGAGGGACCAAGCTGACCGTCCTAG	GGGCTGATGCTGCACCAACTGTATCCATC
2-3	GAGCAACTTCGTGT	CTGTGTTCGGAGGAGGCACCCAGCTGACCGCCCTCG	GGGCTGATGCTGCACCAACTGTATCCATC
FWR4		**F G G G T K L T V L G**	**A D A A P T V S I**

FIG. 12

	3' Human Vλ	Human Jλ1	5' Mouse Cλ2
		F G T G T K V T V L G Q P K S S P T L	
2D1	GCAGGCAGCAACAATTTa	aGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GTCAGCCCAAGTCCACTCCCACTCTC
2D9	GACAGCAGTGGTAATCAT	TATGTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GTCAGCCCAAGTCCACTCCCACTCTC
3E15	GACAGCAGCACTGCc	GTCTTCGGAACTGGGACCAAGGTCACCGTCCTAG	GTCAGCCCAAGTCCACTCCCACTCTC
FWR4			

FIG. 13


Inactivated Mouse λ Light Chain Locus
C2-J4-C4
Enh 3.1
Frt
Hyg-TK
Frt
lox
Enh 2.4
Enh
hVλ
3-12
hVλ
3-1
C2-J4-C4
12/1-λ
Targeting Vector
Frt
Neo
Frt
hJλ1
Enh 2.4
PI-SceI