(19) DANMARK

(10) DK/EP 4087875 T3

(12) Oversættelse af europæisk patentskrift

Patent- og Varemærkestyrelsen

(51) Int.Cl.: *C 07 K 16/28 (2006.01)* *A 61 K 31/573 (2006.01)* *A 61 K 39/395 (2006.01)*
A 61 P 17/00 (2006.01) *A 61 P 37/06 (2006.01)* *C 07 K 16/18 (2006.01)*

(45) Oversættelsen bekendtgjort den: **2024-10-28**

(80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2024-08-28**

(86) Europæisk ansøgning nr.: **21701890.2**

(86) Europæisk indleveringsdag: **2021-01-08**

(87) Den europæiske ansøgnings publiceringsdag: **2022-11-16**

(86) International ansøgning nr.: **EP2021050275**

(87) Internationalt publikationsnr.: **WO2021140202**

(30) Prioritet: **2020-01-08 US 202062958543 P** **2020-01-13 US 202062960647 P**

(84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**

(73) Patenthaver: **Argenx BV, Industriepark Zwijnaarde 7, 9052 Gent, Belgien**

(72) Opfinder: **VERHEESEN, Peter, c/o Industriepark Zwijnaarde 7, 9052 Gent, Belgien**
DUPUY, Patrick, c/o Industriepark Zwijnaarde 7, 9052 Gent, Belgien

(74) Fuldmægtig i Danmark: **Holme Patent A/S, Valbygårdsvej 33, 2500 Valby, Danmark**

(54) Benævnelse: **Antagonister af human neonatal Fc-receptor (FcRn) til behandling af pemphigus-sygdomme**

(56) Fremdragne publikationer:
WO-A1-2020/236695
GOEBELER M. ET AL: "Treatment of pemphigus vulgaris and foliaceus with efgartigimod, a neonatal Fc receptor inhibitor: a phase II multicentre, open-label feasibility trial*", BRITISH JOURNAL OF DERMATOLOGY, vol. 186, no. 3, 28 November 2021 (2021-11-28), Hoboken, USA, pages 429 - 439, XP093055577, ISSN: 0007-0963, Retrieved from the Internet <URL:https://onlinelibrary.wiley.com/doi/full-xml/10.1111/bjd.20782> DOI: 10.1111/bjd.20782
PETER ULRICHTS ET AL: "Neonatal Fc receptor antagonist efgartigimod safely and sustainably reduces IgGs in humans", THE JOURNAL OF CLINICAL INVESTIGATION, vol. 128, no. 10, 24 July 2018 (2018-07-24), GB, pages 4372 - 4386, XP055552033, ISSN: 0021-9738, DOI: 10.1172/JCI97911
HOWARD JAMES F. ET AL: "Randomized phase 2 study of FcRn antagonist efgartigimod in generalized myasthenia gravis", NEUROLOGY, vol. 92, no. 23, 22 May 2019 (2019-05-22), US, pages e2661 - e2673, XP055789304, ISSN: 0028-3878, Retrieved from the Internet <URL:https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6556100/pdf/NEUROLOGY2018910505.pdf> DOI: 10.1212/WNL.0000000000007600
BALIGHI KAMRAN ET AL: "Comparing early and late treatments with rituximab in pemphigus vulgaris: which one is better?", ARCHIVES OF DERMATOLOGICAL RESEARCH, SPRINGER, INTERNATIONAL, BERLIN, DE, vol. 311, no. 1, 1 December 2018 (2018-12-01), pages 63 - 69, XP036671584, ISSN: 0340-3696, [retrieved on

Fortsættes ...

20181201], DOI: 10.1007/S00403-018-1881-1
NIH: "A Study to Evaluate the Safety, PD, PK and Efficacy of ARGX-113 in Patients With Pemphigus", 1 January 2017 (2017-01-01), XP055789389, Retrieved from the Internet <URL:https://clinicaltrials.gov/ct2/show/record/NCT03334058> [retrieved on 20210324]
KASPERKIEWICZ MICHAEL ET AL: "Pemphigus", NATURE REVIEWS DISEASE PRIMERS, vol. 3, no. 1, 11 May 2017 (2017-05-11), XP055790922, Retrieved from the Internet <URL:http://www.nature.com/articles/nrdp201726> DOI: 10.1038/nrdp.2017.26
NIH: "A Study to Assess the Long-term Safety and Efficacy of a Subcutaneous Formulation of Efgartigimod PH20 SC in Adults With Pemphigus (Vulgaris or Foliaceus) (ADDRESS+)", 22 October 2020 (2020-10-22), XP055790873, Retrieved from the Internet <URL:https://clinicaltrials.gov/ct2/show/NCT04598477> [retrieved on 20210329]

DESCRIPTION

Description

FIELD

[0001] The present disclosure relates to methods of treating pemphigus disorders, including but not limited to pemphigus vulgaris and pemphigus foliaceus. The methods involve use of an antagonist of human neonatal Fc receptor (FcRn), which in certain embodiments is efgartigimod.

BACKGROUND

[0002] Pemphigus is a group of chronic blistering epithelial diseases in which the production of IgG autoantibodies against extracellular domains of certain cell membrane proteins of keratinocytes results in acantholysis (loss of cell-cell adhesion between keratinocytes). Major forms of pemphigus include pemphigus vulgaris (PV) and pemphigus foliaceus (PF). Current treatment strategies in pemphigus vary in their capacity to induce disease control (DC) and durable clinical remission (CR). Corticosteroids rapidly affect PV symptoms but must be administered at high daily doses (e.g. oral prednisone at 1 to 1.5 mg/kg/day) to attain effectiveness. If DC is not achieved after 3 to 4 weeks, prednisone dose must be increased and in patients with very active disease, an intravenous (IV) bolus of corticosteroids (e.g., methylprednisolone) may be preferred, especially at treatment initiation. In the Ritux 3 study, the rituximab therapy in combination with low-dose corticosteroids allowed 89% of patients to achieve complete remission off therapy (CRoff) at 24 months, while systemic corticosteroids alone allowed only 34% of patients to achieve CRoff. Joly P. et al., Lancet 2017; 389:2031-2040. While treatment with rituximab offers pemphigus patients an improved therapeutic option, rituximab has a relatively slow onset of action requiring concomitant use of corticosteroids in moderate to high doses between 0.5 to 1.5 mg/kg/day, depending on disease severity, to induce DC. Moreover, patients receiving rituximab relapse in 25-60% of cases, requiring additional cycles of rituximab, and rituximab therapy poses an increased risk of infections as a consequence of systemic B-cell depletion. All these factors, along with the tremendous physical burden of pemphigus disease, necessitate the need for a fast-acting treatment that allows for a quick resolution of blisters, associated pain and physical discomfort, and permit quick steroid tapering to avoid steroid-associated side effects and toxicities. Therefore, there remains a need in pemphigus patients for a safer drug with a rapid onset of action that would achieve early disease control and maintain clinical remission, with or without a minimal dose of corticosteroids (e.g. prednisone 20 mg per day or 10 mg per day, or less).

[0003] Ulrichts et al., (2018) "Neonatal Fc receptor antagonist efgartigimod safely and sustainably reduces IgG in humans", J. Clin. Investig., 128(10), pp. 4372-4386, reports the results of *in vivo* studies carried out to investigate the ability of the FcRn antagonist efgartigimod to deplete IgGs. The studies reported in this paper were carried out in cynomolgus monkeys and human healthy volunteers.

[0004] James et al., (2019) "Randomized phase 2 study of FcRn antagonist efgartigimod in generalized myasthenia gravis", Neurology, 92(23), pp. 2661-2673, reports the results of a phase 2 study investigating use of efgartigimod in the treatment of generalized myasthenia gravis.

[0005] Balghi et al., (2018) "Comparing early and late treatments with rituximab in permphigues vulgaris; which one is better?", Arch. Dermatol. Res., 311(1), pp. 63-69, reviews the results of studies that have used rituximab in the treatment of pemphigus.

[0006] "A study to evaluate the safety, PD, PK and efficacy of ARGX-113 in patients with pemphigus" is a webpage from the ClinicalTrials.gov archive reporting the protocol for the NCT03334058 clinical study.

SUMMARY OF THE INVENTION

[0007] The instant disclosure is based on the discovery that human neonatal Fc receptor (FcRn) antagonists are highly effective in treating pemphigus and pemphigus-related disorders. Accordingly, the instant disclosure is broadly directed, at least in part, to methods for treating pemphigus and pemphigus-related disorders with FcRn antagonists.

[0008] The invention is directed to an FcRn antagonist for use in the treatment of pemphigus in a subject in need thereof, wherein

1. (i) the subject has
    1. (a) newly diagnosed pemphigus,
    2. (b) relapsing pemphigus, or
    3. (c) refractory pemphigus,
2. (ii) the FcRn antagonist consists of a variant Fc region, wherein said variant Fc region consists of two Fc domains which form a homodimer, and wherein the amino acid sequence of each of the Fc domains is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, and SEQ ID NO: 3, and
3. (iii) the FcRn antagonist is administered in combination with a corticosteroid and the corticosteroid is at a dose of ≤ 0.5 mg prednisone/kg/day or equivalent.

[0009] Described herein are methods of treating pemphigus, comprising administering to a

subject in need thereof an effective amount of a human neonatal Fc receptor (FcRn) antagonist, wherein the subject has (a) newly diagnosed pemphigus, (b) relapsing pemphigus, or (c) refractory pemphigus, thereby treating pemphigus in the subject.

[0010] In another aspect, the disclosure is directed to a method of treating pemphigus, comprising (i) selecting a subject that has (a) newly diagnosed pemphigus, (b) relapsing pemphigus, or (c) refractory pemphigus; and (ii) administering to the subject an effective amount of a human neonatal Fc receptor (FcRn) antagonist.

[0011] In certain aspects, the disclosure is also based on the surprising discovery that FcRn antagonists can achieve rapid disease control (e.g., inhibition of new lesion formation and beginning of healing of established lesions) of pemphigus. In certain embodiments, disease control can be obtained within about 1 month of administration of the initial dose of FcRn antagonist. In certain embodiments, disease control can be obtained within about 3 weeks, within about 2 weeks, or within about 1 week of administration of the initial dose of FcRn antagonist. In exemplary embodiments, disease control can be obtained within about 14-16 days of administration of the initial dose of FcRn antagonist. In certain embodiments, disease control can be obtained after 1-4 doses (e.g., infusions or subcutaneous administrations) of FcRn antagonist. In certain exemplary embodiments, the FcRn antagonists for use in accordance with the invention achieve disease control after 1 or 2 doses (e.g., infusions or subcutaneous administrations) of FcRn antagonist.

[0012] In yet other aspects, the disclosure is also based on the surprising discovery that FcRn antagonists, optionally when co-administered with a corticosteroid, can achieve disease control (e.g., rapid disease control) of pemphigus at much lower doses of corticosteroids than conventional therapies. Thus, in certain aspects the disclosure is based on the surprising discovery that the FcRn antagonists for use in accordance with the invention provide unprecedented levels of "corticosteroid-sparing."

[0013] For example, conventional therapies require very high weight-based dosing of corticosteroids (e.g., 1 or 2 mg/kg/day). By contrast, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of 0.5 mg/kg/day or less. In exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 0.4 mg/kg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 0.3 mg/kg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 0.25 mg/kg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 0.2 mg/kg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 0.1 mg/kg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 0.05 mg/kg/day.

[0014] In other embodiments, the invention can achieve disease control (e.g., rapid disease control) at much lower fixed doses of corticosteroids than conventional therapies. For example, while convention therapies require high fixed doses of corticosteroid (e.g., 100-150 mg/day), in certain embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of 20 mg/day or less. In exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 17.5 mg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 15 mg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 12.5 mg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 10 mg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 9 mg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 8 mg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 7 mg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 6 mg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 5 mg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 4 mg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 3 mg/day. In other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 2 mg/day. In still other exemplary embodiments, the invention can achieve disease control (e.g., rapid disease control) at corticosteroid doses of less than about 1 mg/day.

[0015] In certain embodiments, disease control is obtained within about 24 weeks, about 23 weeks, about 22 weeks, about 21 weeks, about 20 weeks, about 15 weeks, about 13 weeks, about 12 weeks, about 10 weeks, about 9 weeks, about 8 weeks, about 7 weeks, about 6 weeks, about 5 weeks, about 4 weeks, about 3 weeks, about 2 weeks, or about 1 week of administration of the initial dose of FcRn antagonist. In certain embodiments, disease control is obtained in 1 to 24 weeks, in 1 to 20 weeks, in 1 to 15 weeks, in 1 to 13 weeks, in 1 to 12 weeks, 1 to 10 weeks, in 1 to 6 weeks, in 1 to 4 weeks, in 1 week to 16 days, or in 1 to 2 weeks of administration of the initial dose of FcRn antagonist. In a specific embodiment, disease control is obtained in 1 to 13 weeks of administration of the initial dose of FcRn antagonist.

[0016] In other aspects, the disclosure is based on the surprising discovery that the FcRn antagonists for use in accordance with the invention achieve complete remission (e.g., no new lesions, and all established lesions completely healed) within about 4 months of administration of the initial dose of FcRn antagonist. In certain embodiments, complete remission is obtained within about 41 weeks, about 24 weeks, about 23 weeks, about 22 weeks, about 21 weeks, about 20 weeks, about 15 weeks, about 13 weeks, about 12 weeks, about 10 weeks, about 9

weeks, about 8 weeks, about 7 weeks, about 6 weeks, about 5 weeks, about 4 weeks, about 3 weeks, or about 2 weeks of administration of the initial dose of FcRn antagonist. In certain embodiments, complete remission is obtained in 2 to 41 weeks, in 2 to 24 weeks, in 2 to 20 weeks, in 2 to 15 weeks, in 2 to 13 weeks, in 2 to 12 weeks, in 2 to 10 weeks, or in 2 to 6 weeks of administration of the initial dose of FcRn antagonist. In a specific embodiment, complete remission is obtained in 2 to 41 weeks of administration of the initial dose of FcRn antagonist. In exemplary embodiments, complete remission is obtained following biweekly administration of the FcRn antagonist and about 0.5 mg/kg/day of corticosteroid or less (e.g., about 0.5 mg/kg/day, about 0.4 mg/kg/day, about 0.35 mg/kg/day, about 0.3 mg/kg/day, about 0.29 mg/kg/day, about 0.28 mg/kg/day, about 0.27 mg/kg/day, about 0.26 mg/kg/day, about 0.25 mg/kg/day, about 0.24 mg/kg/day, about 0.23 mg/kg/day, about 0.22 mg/kg/day, about 0.21 mg/kg/day or about 0.2 mg/kg/day). In exemplary embodiments, complete remission is obtained following weekly administration of the FcRn antagonist and about 0.5 mg/kg/day of corticosteroid or less (e.g., about 0.5 mg/kg/day, about 0.4 mg/kg/day, about 0.35 mg/kg/day, about 0.3 mg/kg/day, about 0.29 mg/kg/day, about 0.28 mg/kg/day, about 0.27 mg/kg/day, about 0.26 mg/kg/day, about 0.25 mg/kg/day, about 0.24 mg/kg/day, about 0.23 mg/kg/day, about 0.22 mg/kg/day, about 0.21 mg/kg/day or about 0.2 mg/kg/day). In exemplary embodiments, complete remission is obtained following biweekly administration of the FcRn antagonist and about 20 mg/day of corticosteroid or less (e.g., about 20 mg/day, about 17.5 mg/day about 15 mg/day, about 10 mg/day, about 5 mg/day, about 4 mg/day, about 3 mg/day, about 2mg/day or about 1mg/day). In exemplary embodiments, complete clinical remission is obtained following weekly administration of the FcRn antagonist and about 20 mg/day of corticosteroid or less (e.g., about 20 mg/day, about 17.5 mg/day about 15 mg/day, about 10 mg/day, about 5 mg/day, about 4 mg/day, about 3 mg/day, about 2mg/day or about 1mg/day).

[0017] In certain embodiments, the FcRn antagonists for use in accordance with the invention obtain Pemphigus Disease Area Index (PDAI) activity score improvement of at least about 50%, at least about 60%, at least about 70%, at least about 80%, or at least about 90%. In other embodiment, the FcRn antagonists for use in accordance with the invention result in an absolute PDAI activity score of 6 or less (e.g., an absolute PDAI activity score of 6, an absolute PDAI activity score of 5, an absolute PDAI activity score of 4, an absolute PDAI activity score of 3, an absolute PDAI activity score of 2, an absolute PDAI activity score of 1, or an absolute PDAI activity score of 0).

[0018] The disclosure is based on the surprising discovery that the FcRn antagonists for use in accordance with the invention achieve synergistic effects when FcRn antagonists are combined with low doses of corticosteroids. Without being bound to any particular theory, it is thought that the combination of FcRn antagonists and corticosteroids is capable of stimulating new basement membrane (e.g., via desmoglein 1 (i.e., DSG1 or Dsg-1) and/or desmoglein 3 (i.e., DSG3 or Dsg-3) synthesis) while also clearing pathogenic autoantibodies (e.g., anti-DSG1 or anti-DSG3 autoantibodies). Therefore, the FcRn antagonist is administered in combination with low doses of corticosteroid (e.g., about 0.2 or about 0.25 or about 0.5 mg/kg/day of oral prednisone or equivalents thereof). In preferred embodiments, the corticosteroid is administered orally in low doses. In certain embodiments, the corticosteroid is administered

topically to the skin. In certain embodiments, the corticosteroid is administered systemically by intravenous injection or infusion.

[0019] In still other aspects, the invention is based on the discovery that newly diagnosed pemphigus patients and pemphigus patients previously treated with corticosteroids may be treated with an FcRn antagonist and subsequently tapered off treatment with corticosteroids much earlier and/or at higher initial levels of corticosteroids than conventional therapies. For example, conventional therapies typically require sustained treatment of pemphigus with corticosteroids and gradual and minimal tapering of the corticosteroids once disease control is observed. By contrast, the invention can begin tapering of corticosteroids much earlier and with larger dose reductions than conventional therapies. Accordingly, an aspect of the disclosure is a method of treating pemphigus, comprising administering to a subject in need thereof an effective amount of a human neonatal Fc receptor (FcRn) antagonist and an initial tapering dose of corticosteroid beginning at ≤ 2 mg prednisone/kg/day or equivalent.

[0020] In certain embodiments, corticosteroid tapering can begin once disease control has been achieved. As noted above, a surprising finding is that pemphigus patients treated with an FcRn antagonist may begin corticosteroid tapering much earlier than with conventional therapies. In certain embodiments, corticosteroid tapering can begin within about 10 weeks, about 9 weeks, about 8 weeks, about 7 weeks, about 6 weeks, about 5 weeks, about 4 weeks, about 3 weeks and about 2 weeks of administration of the initial dose of FcRn antagonist. In other embodiments, corticosteroid tapering can begin following disease control or following complete clinical remission. In certain embodiments, the tapering of corticosteroids can begin within about 1 month, within about 3 weeks, within about 2 weeks or within about 1 week of administration of the initial dose of FcRn antagonist.

[0021] In still other embodiments, the invention allows for more rapid tapering of corticosteroids than conventional therapies. For example, whereas conventional therapies only allow for corticosteroid tapering as frequently as once a month, the FcRn antagonists of the invention may be tapered more frequently, provided that disease control is maintained. In certain exemplary embodiments, a reduction of the initial or subsequent tapering dose can occur every 3 weeks. In other exemplary embodiments, a reduction of the initial or subsequent tapering dose can occur every 2 weeks. In other exemplary embodiments, a reduction of the initial or subsequent tapering dose can occur every week. In certain embodiments, the tapering of corticosteroids can comprise a reduction of about 0.5 mg prednisone/kg/day or equivalent about every week, about every two weeks or about every month. In certain embodiments, the tapering of corticosteroids can comprise a reduction of about 0.25 mg prednisone/kg/day or equivalent about every week, about every two weeks or about every month. In certain embodiments, the tapering of corticosteroids can comprise a reduction of about 0.1 mg prednisone/kg/day or equivalent about every week, about every two weeks or about every month.

[0022] In accordance with each of the foregoing aspects:
The FcRn antagonist for use in accordance with the invention is capable of binding to FcRn

and preventing IgG recycling and/or causing a decrease in IgG level. The FcRn antagonist for use in accordance with the invention consists of a variant Fc region or domain and lacks an antigen binding site. Also described herein are FcRn antagonists, wherein the FcRn antagonist can be an antibody or antigen-binding fragment thereof that binds specifically to FcRn and inhibits binding of immunoglobulin to FcRn.

[0023] The FcRn antagonist for use in accordance with the invention is a human IgG1 antibody Fc fragment that has been engineered for increased affinity to FcRn. The FcRn antagonist for use in accordance with the invention consists of a variant Fc region, wherein said variant Fc region consists of two Fc domains which form a homodimer, wherein the amino acid sequence of each of the Fc domains is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, and SEQ ID NO: 3. In some embodiments, the amino acid sequence of each of the Fc domains is SEQ ID NO: 1. In some embodiments, the amino acid sequence of each of the Fc domains is SEQ ID NO: 2. In some embodiments, the amino acid sequence of each of the Fc domains is SEQ ID NO: 3.

[0024] In certain exemplary embodiments, the FcRn antagonist is efgartigimod.

[0025] In some embodiments, the pemphigus comprises pemphigus vulgaris (PV), pemphigus foliaceus (PF), or both PV and PF. In some embodiments, the pemphigus comprises pemphigus vulgaris (PV). In some embodiments, the pemphigus comprises pemphigus foliaceus (PF). In some embodiments, the pemphigus comprises both pemphigus vulgaris (PV) and pemphigus foliaceus (PF). In some embodiments, the pemphigus consists of pemphigus vulgaris (PV). In some embodiments, the pemphigus consists of pemphigus foliaceus (PF). In some embodiments, the pemphigus consists of both pemphigus vulgaris (PV) and pemphigus foliaceus (PF).

[0026] In certain embodiments, the pemphigus vulgaris (PV) is of the mucosal-dominant subtype. In certain embodiments, the pemphigus vulgaris (PV) is of the mucocutaneous subtype. In certain embodiments, the pemphigus vulgaris (PV) is of the cutaneous subtype.

[0027] In some embodiments, the subject has mild, moderate, or severe pemphigus as classified by Pemphigus Disease Area Index (PDAI).

[0028] In some embodiments, the subject has mild pemphigus as classified by PDAI. In exemplary embodiments, the pemphigus or pemphigus-related disorder is characterized by a PDAI score of less than 15.

[0029] In some embodiments, the subject has moderate pemphigus as classified by PDAI. In certain embodiments, the pemphigus-related disorder is moderate to severe pemphigus as classified by PDAI. In some embodiments, the subject suffering from pemphigus has a PDAI score ≥ 15 and < 45.

[0030] In some embodiments, the subject has severe pemphigus. In exemplary embodiments,

the subject suffering from severe pemphigus has a PDAI score ≥ 45.

[0031] In some embodiments, the subject has refractory pemphigus.

[0032] In certain embodiments, the pemphigus is newly diagnosed. In other embodiments, the pemphigus is relapsing pemphigus.

[0033] In certain embodiments, the FcRn antagonist is administered to the subject intravenously.

[0034] In certain embodiments, the FcRn antagonist is administered to the subject subcutaneously. In certain embodiments, the FcRn antagonist is co-formulated with hyaluronidase and administered subcutaneously.

[0035] In some embodiments, the FcRn antagonist is administered once weekly or more frequently (e.g., every 1, 2, 3, 4, 5, 6, or 7 days) until disease control.

[0036] In some embodiments, the FcRn antagonist is administered less frequently than once weekly (e.g., every 8, 9, 10, 11, 12, 13, or 14 days; or every 2, 3, 4, 5, 6 weeks) until disease control.

[0037] In some embodiments, the FcRn antagonist is administered once weekly or biweekly at a dose of about 10 mg/kg or about 25 mg/kg until disease control.

[0038] In some embodiments, the FcRn antagonist is co-administered with a corticosteroid dose of 10 mg/day or less.

[0039] In some embodiments, the invention further comprises administering a tapering dose of corticosteroid beginning at ≤ 0.5 mg prednisone/kg/day or equivalent. In certain embodiments, the tapering dose of corticosteroid is administered after disease control has been achieved.

[0040] In some embodiments, the corticosteroid is administered systemically.

[0041] In some embodiments, the FcRn antagonist is administered once weekly or more frequently (e.g., every 1, 2, 3, 4, 5, 6, or 7 days) until complete remission (e.g., a PDAI score of 0).

[0042] In some embodiments, the FcRn antagonist is administered once weekly or biweekly at a dose of about 10 mg/kg or about 25 mg/kg until complete remission (e.g., a PDAI score of 0).

[0043] In some embodiments, the FcRn antagonist is administered less frequently than once weekly (e.g., every 8, 9, 10, 11, 12, 13, or 14 days; or every 2, 3, 4, 5, or 6 weeks) until complete remission (e.g., a PDAI score of 0).

[0044] In some embodiments, complete remission is obtained with a corticosteroid dose of 20 mg/day or less.

[0045] In some embodiments, complete remission is obtained with a corticosteroid dose of 10 mg/day or less.

[0046] In some embodiments, a tapering dose of corticosteroid is administered after disease control or complete remission has been achieved.

[0047] In some embodiments, the FcRn antagonist is administered intravenously.

[0048] In some embodiments, the FcRn antagonist is administered intravenously once weekly.

[0049] In some embodiments, the FcRn antagonist is administered intravenously every two weeks.

[0050] In some embodiments, the FcRn antagonist is administered intravenously at a dose of 10 mg/kg to 30 mg/kg.

[0051] In some embodiments, the FcRn antagonist is administered intravenously at a dose of 10 mg/kg.

[0052] In some embodiments, the FcRn antagonist is administered intravenously at a dose of 25 mg/kg.

[0053] In some embodiments, the FcRn antagonist is administered subcutaneously.

[0054] In some embodiments, the FcRn antagonist is administered subcutaneously once weekly.

[0055] In some embodiments, the FcRn antagonist is administered subcutaneously every two weeks.

[0056] In some embodiments, the FcRn antagonist is co-formulated with hyaluronidase and administered subcutaneously.

[0057] In some embodiments, the FcRn antagonist is administered subcutaneously at a fixed dose of about 750 mg to about 3000 mg. In some embodiments, the FcRn antagonist is administered subcutaneously at a fixed dose of about 1000 mg or about 2000 mg. In some embodiments, the FcRn antagonist is administered subcutaneously at a fixed dose of about 1000 mg twice on the same day.

[0058] In some embodiments, the FcRn antagonist is administered subcutaneously once weekly or more frequently (e.g., every 1, 2, 3, 4, 5, 6, or 7 days) at a fixed dose of about 750

mg to about 1750 mg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly or more frequently (e.g., every 1, 2, 3, 4, 5, 6, or 7 days) at a fixed dose of about 1000 mg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly or more frequently (e.g., every 1, 2, 3, 4, 5, 6, or 7 days) at a dose of about 10 mg/kg to about 25 mg/kg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly or more frequently (e.g., every 1, 2, 3, 4, 5, 6, or 7 days) at a dose of about 10 mg/kg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly or more frequently (e.g., every 1, 2, 3, 4, 5, 6, or 7 days) at a dose of about 25 mg/kg.

[0059] In some embodiments, the FcRn antagonist is first administered subcutaneously at a fixed dose of about 1000 mg twice on the same day.

[0060] In some embodiments, the FcRn antagonist is administered in an induction phase and a consolidation phase.

[0061] In some embodiments, the FcRn antagonist is administered in an induction phase and a consolidation phase, wherein

1. (i) during the induction phase the FcRn antagonist is administered once weekly or biweekly and corticosteroid 0.5 mg prednisone/kg/day or equivalent is administered until disease control, and
2. (ii) during the consolidation phase the FcRn antagonist dose is reduced or the FcRn antagonist dosing interval is lengthened, and/or the corticosteroid dose is decreased or the corticosteroid dosing interval is lengthened, to an end-of-consolidation dose or dosing interval effective to prevent new lesions from appearing.

[0062] In some embodiments, during the induction phase, the FcRn antagonist dosing interval is once weekly or more frequently (e.g., a dosing interval of every 1, 2, 3, 4, 5, 6 or 7 days). In some embodiments, during the induction phase, the FcRn antagonist is administered subcutaneously at a fixed dose of about 750 mg to about 1750 mg. In some embodiments, during the induction phase, the FcRn antagonist is administered subcutaneously at a fixed dose of about 1000 mg. In some embodiments, during the induction phase, the FcRn antagonist is administered subcutaneously at a dose of about 10 mg/kg to about 25 mg/kg. In some embodiments, during the induction phase, the FcRn antagonist is administered subcutaneously at a dose of about 10 mg/kg. In some embodiments, during the induction phase, the FcRn antagonist is administered subcutaneously at a dose of about 25 mg/kg.

[0063] In some embodiments, during the induction phase, the FcRn antagonist is first administered subcutaneously at a fixed dose of about 1000 mg twice on the same day.

[0064] In some embodiments, during the consolidation phase, the FcRn antagonist dosing interval is once weekly or more frequently (e.g., a dosing interval of every 1, 2, 3, 4, 5, 6, or 7

days). In some embodiments, during the consolidation phase, the FcRn antagonist is administered less frequently than once weekly (e.g., a dosing interval of every 8, 9, 10, 11, 12, 13, or 14 days; or a dosing interval of every 2, 3, or 4 weeks) until disease control. In some embodiments, during the consolidation phase, the FcRn antagonist dosing interval is once weekly or every 2 weeks.

[0065] In some embodiments, the FcRn antagonist for use is administered in an induction phase and a consolidation phase, wherein (i) during the induction phase the FcRn antagonist is administered once weekly and with a corticosteroid dose of 0.5 mg prednisone/kg/day or equivalent until disease control, and (ii) during the consolidation phase the FcRn antagonist dose is decreased or the FcRn antagonist dosing interval is lengthened, and/or the corticosteroid dose is decreased or the corticosteroid dosing interval is lengthened, to an end-of-consolidation dose or dosing interval effective to prevent new lesions from appearing.

[0066] In some embodiments, the FcRn antagonist for use in accordance with the invention further comprise a maintenance phase, wherein (iii) during the maintenance phase the end-of-consolidation dose or dosing interval for the FcRn antagonist and/or the prednisone is continued until complete clearance of lesions. In some embodiments, during the maintenance phase the FcRn antagonist dosing interval is once weekly or more frequently (e.g., a dosing interval of every 1, 2, 3, 4, 5, 6, or 7 days). In some embodiments, during the maintenance phase the FcRn antagonist dosing interval is once weekly. In some embodiments, during the maintenance phase the FcRn antagonist dosing interval is less frequent than once weekly (e.g., a dosing interval of every 8, 9, 10, 11, 12, 13, or 14 days; or a dosing interval of every 2, 3, 4, 5, or 6 weeks). In some embodiments, during the maintenance phase the FcRn antagonist dosing interval is biweekly.

[0067] In some embodiments, during the induction phase the FcRn antagonist is administered intravenously at a dose of 10 mg/kg to 30 mg/kg. In some embodiments, during the induction phase the FcRn antagonist is administered intravenously at a dose of 10 mg/kg. In some embodiments, during the induction phase the FcRn antagonist is administered intravenously at a dose of 25 mg/kg.

[0068] In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of 750 mg to 3000 mg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of 1000 mg or 2000 mg. In some embodiments, during the induction phase the FcRn antagonist is first administered subcutaneously at a fixed dose of about 1000 mg twice on the same day.

[0069] In some embodiments, during the consolidation phase the FcRn antagonist dosing interval is once weekly, every 2 weeks or less frequently.

[0070] In some embodiments, during the maintenance phase the FcRn antagonist dosing interval is once weekly, every 2 weeks, every 4 weeks, or less frequently.

[0071] In some embodiments, the pemphigus goes into complete remission following treatment with FcRn antagonist. and a corticosteroid dose of ≤ about 0.5 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 0.3 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 0.2 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 0.1 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 0.2 mg prednisone/kg/day or equivalent to about 0.50 mg prednisone/kg/day or equivalent.

[0072] In some embodiments, the complete remission is achieved with a corticosteroid dose of about 20 mg prednisone/day (or equivalent) or less. In some embodiments, the complete remission is achieved with a corticosteroid dose of about 15 mg prednisone/day (or equivalent) or less. In some embodiments, the complete remission is achieved with a corticosteroid dose of about 10 mg prednisone/day (or equivalent) or less (e.g., about 10, about 9, about 8, about 7, or about 6 prednisone/day (or equivalent)). In some embodiments, the complete remission is achieved with a corticosteroid dose of about 5 mg prednisone/day (or equivalent) or less (e.g., about 5, about 4, about 3, about 2, about 1, or about 0.5 mg prednisone/day (or equivalent)).

[0073] In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 0.5 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 0.4 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 0.3 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 0.2 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 0.1 mg prednisone/kg/day or equivalent.

[0074] In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 10 mg prednisone/day or equivalent (e.g., about 10 mg, about 9 mg, about 8 mg, about 7 mg, or about 6 mg prednisone/day (or equivalent)). In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 5 mg prednisone/day or equivalent (e.g., about 5 mg, about 4 mg, about 3 mg, or about 2 mg prednisone/day (or equivalent)). In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 2 mg prednisone/day or equivalent. In some embodiments, the complete remission is maintained without corticosteroid.

[0075] In some embodiments, the subject has refractory pemphigus.

[0076] In some embodiments, the subject is rituximab-refractory.

[0077] In some embodiments, the subject is corticosteroid-intolerant.

[0078] In some embodiments, the FcRn antagonists for use further comprise administering a B-cell depleting agent (e.g., rituximab) to the subject.

BRIEF DESCRIPTION OF THE DRAWINGS

[0079]

Figure 1 depicts a schematic design of a completed Phase 2 trial (1701 study) of efgartigimod in patients with pemphigus vulgaris and foliaceus.

Figure 2 is a flow chart indicating patients' disposition for safety dataset analysis and efficacy dataset analysis and their assignments to cohorts in the Phase 2 trial (1701 study) of efgartigimod in patients with pemphigus vulgaris and foliaceus.

Figure 3 is a summary slide of the first interim results of a Phase 2 trial (1701 study) of efgartigimod in pemphigus patients. Cut-off date on the data from the Phase 2 trial is: 7 Nov 2019. The patient population included pemphigus vulgaris (PV) patients of the mucosal dominant (N=8), mucocutaneous (N=10), and cutaneous (N=1) subtypes, as well as pemphigus foliaceus (PF) patients (N=4). Patients exhibited a decrease in Pemphigus Disease Area Index (PDAI) activity which correlated with a reduction of pathogenic IgG4 autoantibodies (DSG-1 and DSG-3) following initial induction dosing of efgartigimod and subsequent weekly or biweekly maintenance dosing with efgartigimod.

Figures 4A-4D depict the clinical efficacy of efgartigimod as assessed by the Pemphigus Disease Area Index (PDAI) activity in 4 patient cohorts of the Phase 2 trial (initial interim results in 1701 study; cut-off date on the data from the Phase 2 trial is: 7 Nov 2019.). Cohort 1: 10 mg/kg i.v., induction of efgartigimod = 4 infusions (3 weeks), maintenance = 2 infusions (6 weeks). Cohort 2: Screening: allow suitable oral prednisone to stabilize disease activity at screening, to be continued at stable dose at induction (optional measure). Induction: Same dose (10 mg/kg i.v.) and frequency. Maintenance: two additional administrations, extend maintenance duration to 8 weeks (biweekly administration for 8 weeks). Follow-up: one additional follow-up visit (week 2) and prolongation to 10 weeks. Cohort 3: Screening: Allow suitable oral prednisone to stabilize disease activity at screening, may be further increased at any post-baseline visit (all patients). Induction: Same dose (10 mg/kg i.v.) and frequency. Maintenance: 2 additional administrations, extend maintenance duration to 12 weeks (dosing every two weeks for 12 weeks). Treatment-free follow-up 10 weeks. Figure 4A, cohort 1; Figure 4B, cohort 2; Figure 4C, cohort 3; and Figure 4D, cohort 4.
Most patients demonstrated strong PDAI score improvements: In cohort 1 there was a correlation in IgG reduction with anti-Dsg reduction with PDAI score improvement, early DC (mono/combo) and suboptimal efgartigimod dosing in maintenance; in cohort 2 there was improved maintenance with efgartigimod dosing every other week. In cohort 3 all patients ultimately associated with prednisone, maintenance further improved symptoms, and EoC/CR (end of consolidation/complete remission) was noted when associated with oral prednisone. In cohort 4, strong PDAI score improvements were noted, there was a high rate of EoC, and CR

when prednisone was not tapered before CR.

Figures 5A-5D depict clinical efficacy in moderate severity patients of the Phase 2 trial (PDAI 15-44 at baseline). Figure 5A, cohort 1; Figure 5B, cohort 2, Figure 5C, cohort 3; Figure 5D, cohort 4. Most patients demonstrated strong PDAI score improvements: cohort 1 comprised three moderate patients (3 PV, 0 pemphigus foliaceus (PF)); cohort 2 comprised two moderate patients (2 PV, 0 PF); cohort 3 comprised four moderate patients (4 PV, 0 PF); and cohort 4 comprised five moderate patients (1 PV, 4 PF). Data is from initial interim results in 1701 study of efgartigimod in pemphigus patients. Cut-off date on the data from the Phase 2 trial is: 7 Nov 2019.

Figures 6A-6E are graphs depicting PDAI activity scores over time in A) cohort 1, B) cohort 2, C) cohort 3, D) cohort 4 pemphigus vulgaris patients, E) cohort 4 pemphigus foliaceus patients. Data is from final results in the Phase 2 trial (1701 study) of efgartigimod in pemphigus patients. Cut-off date on the data from the Phase 2 trial is: 28 Oct 2020.

Figures 7A and 7B depict the clinical efficacy as assessed by the total IgG (Figure 7A) and PDAI activity (Figure 7B) per cohort. There is a correlation between IgG reduction, autoantibody reductions, and PDAI score improvements. Patients in cohorts 1 and 2 were treated with efgartigimod monotherapy or in combination with prednisone. All patients in cohort 3 were ultimately associated with prednisone. Patients in cohort 4 were being treated in combination with prednisone per protocol. Data is from initial interim results of the Phase 2 trial (1701 study) of efgartigimod in pemphigus patients. Cut-off date on the data from the Phase 2 trial is: 7 Nov 2019. FU, follow-up.

Figures 8A and 8B depict pharmacodynamic (PD) data for the reduction of total IgG in the Phase 2 trial (1701 study) of efgartigimod in pemphigus patients. PD profiles are shown for four weekly administrations of efgartigimod at 25 mg/kg. Figure 8A, PD profile in PV. Figure 8B, PD profiles in healthy volunteers (HV) and pemphigus vulgaris patients (PV). Similar PD profiles were observed between PV and HV, in line with expectations based on HV and modeling. Data is from initial interim results of the trial. Cut-off date on the data from the Phase 2 trial is: 7 Nov 2019.

Figures 9A and 9B depict pharmacodynamic (PD) data for the reduction of total IgG in PV, healthy volunteers (HV), and other indications. PD profiles are shown for four weekly administrations of efgartigimod at 10 mg/kg. Figure 9A, PD profile in PV. Figure 9B, PD profiles in HV, PV, myasthenia gravis (MG) and immune thrombocytopenia (ITP). Data is from initial interim results of the Phase 2 trial (1701 study) of efgartigimod in pemphigus patients. Cut-off date on the data from the Phase 2 trial is: 7 Nov 2019.

Figures 10A and 10B depict serum levels of total IgG in A) cohort 1-3, B) cohort 4. Data is from final results of the Phase 2 trial (1701 study) of efgartigimod in pemphigus patients. Cut-off date on the data from the Phase 2 trial is: 24 Jun 2020.

Figures 11A-11H depict serum levels of IgG subclasses IgG1, IgG2, IgG3, IgG4 in cohort 1-3 (A - D) and in cohort 4 (E - H). Data is from final results of the Phase 2 trial (1701 study) of

efgartigimod in pemphigus patients. Cut-off date on the data from the Phase 2 trial is: 24 Jun 2020.

Figures 12A-12D depict serum levels of anti-desmoglein 1 autoantibodies in cohorts 1-3 (Figure 12A) and cohort 4 (Figure 12B) and anti-desmoglein 3 autoantibodies in cohorts 1-3 (Figure 12C) and cohort 4 (Figure 12D), over time. Data is from final results of Phase 2 trial (1701 study) of efgartigimod in pemphigus patients.

Figure 13 is a diagram outlining the design of the Phase 3 clinical trial described in Example 12. ABQOL=Autoimmune Bullous Disease Quality of Life; CR=complete clinical remission; CRmin=complete remission on minimal therapy; DC=disease control; Efg=efgartigimod PH20 SC; EoC=end of consolidation; EoS=end of study; EQ-5D-5L=EuroQol 5-dimension 5-level; OLE=open-label extension; PD=pharmacodynamics; PDAI=Pemphigus Disease Area Index; PF=pemphigus foliaceus; PK=pharmacokinetics; PV=pemphigus vulgaris; SC=subcutaneous.

Figure 14 is a diagram outlining the design of the Phase 3 clinical trial described in Example 13. CR=complete clinical remission; CRmin=complete remission on minimal therapy; CRoff=complete remission off therapy; DC=disease control; Efg=efgartigimod PH20 SC; EoT=end of treatment; FU=follow-up; SC=subcutaneous; Trt Fail=treatment failure; W=week.

DETAILED DESCRIPTION OF THE INVENTION

[0080] The present disclosure provides engineered FcRn antagonists and methods for their use in treating pemphigus, including pemphigus vulgaris and pemphigus foliaceus. Advantageously, the methods disclosed herein permit faster disease control than achieved with current therapies, as well as the potential to taper and even discontinue corticosteroids after achieving clinical remission.

Definitions

[0081] Unless otherwise defined herein, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. The meaning and scope of the terms should be clear, however, in the event of any latent ambiguity, definitions provided herein take precedent over any dictionary or extrinsic definition. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclature used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well-known and commonly used in the art.

[0082] In order that the present invention may be more readily understood, certain terms are

first defined.

[0083] As used herein, the term "FcRn" refers to a neonatal Fc receptor. Exemplary FcRn molecules include human FcRn encoded by the FCGRT gene as set forth in RefSeq NM_004107. The amino acid sequence of the corresponding protein is set forth in RefSeq NP_004098.

[0084] As used herein, the term "FcRn antagonist" refers to any agent that binds specifically to FcRn and inhibits the binding of immunoglobulin to FcRn (e.g., human FcRn).

[0085] As used herein, in certain embodiments the term "FcRn antagonist" refers to any agent comprising or consisting of an Fc region (e.g., a variant Fc region disclosed herein) that binds specifically to FcRn through the Fc region and inhibits the binding of immunoglobulin to FcRn. The FcRn antagonist of the invention is not a full-length IgG antibody..

[0086] As used herein, the term "Fc region" refers to the portion of a native immunoglobulin formed by the Fc domains of its two heavy chains. A native Fc region is homodimeric.

[0087] As used herein, the term "variant Fc region" refers to an Fc region with one or more alteration relative to a native Fc region. Alteration can include amino acid substitutions, additions and/or deletions, linkage of additional moieties, and/or alteration of the native glycans. The term encompasses heterodimeric Fc regions where each of the constituent Fc domains is different. Examples of such heterodimeric Fc regions include, without limitation, Fc regions made using the "knobs and holes" technology as described in, for example, US 8216805. The term also encompasses single chain Fc regions where the constituent Fc domains are linked together by a linker moiety, as described in, for example, US 2009/0252729A1 and US 2011/0081345A1,.

[0088] As used herein, the term "Fc domain" refers to the portion of a single immunoglobulin heavy chain beginning in the hinge region just upstream of the papain cleavage site and ending at the C-terminus of the antibody. Accordingly, a complete Fc domain comprises at least a portion of a hinge (e.g., upper, middle, and/or lower hinge region) domain, a CH2 domain, and a CH3 domain.

[0089] As used herein the term "FcRn binding fragment" refers to a portion of an Fc region that is sufficient to confer FcRn binding.

[0090] As used herein, the term "EU position" refers to the amino acid position in the EU numbering convention for the Fc region described in Edelman, G.M. et al., Proc. Natl. Acad. USA, 63, 78-85 (1969) and Kabat et al, in "Sequences of Proteins of Immunological Interest", U.S. Dept. Health and Human Services, 5th edition, 1991.

[0091] As used herein, the term "CH1 domain" refers to the first (most amino terminal) constant region domain of an immunoglobulin heavy chain that extends from about EU

positions 118-215. The CH1 domain is adjacent to the VH domain and amino terminal to the hinge region of an immunoglobulin heavy chain molecule, and does not form a part of the Fc region of an immunoglobulin heavy chain.

[0092] As used herein, the term "hinge region" refers to the portion of a heavy chain molecule that joins the CH1 domain to the CH2 domain. This hinge region comprises approximately 25 residues and is flexible, thus allowing the two N-terminal antigen binding regions to move independently. Hinge regions can be subdivided into three distinct domains: upper, middle, and lower hinge domains (Roux et al., J. Immunol. 161: 4083 (1998)). The FcRn antagonists of the instant disclosure can include all or a portion of a hinge region.

[0093] As used herein, the term "CH2 domain" refers to the portion of a heavy chain immunoglobulin molecule that extends from about EU positions 231-340.

[0094] As used herein, the term "CH3 domain" includes the portion of a heavy chain immunoglobulin molecule that extends approximately 110 residues from N-terminus of the CH2 domain, e.g., from about position 341-446 (EU numbering system).

[0095] As used herein, the term "FcRn antagonist" may refer to an antibody or antigen-binding fragment thereof that binds specifically to FcRn via its antigen binding domain and inhibits the binding of the Fc region of immunoglobulin to FcRn.

[0096] An example of an antibody that binds specifically to FcRn and inhibits the binding of the Fc region of immunoglobulin to FcRn is nipocalimab, also known as M281. Nipocalimab is a full-length (150 kDa) "Fc dead" (aglycoslated and effectorless) IgG1 monoclonal antibody. Nipocalimab has been administered as an intravenous infusion in Phase 2 clinical trials for the treatment of myasthenia gravis (MG), warm autoimmune hemolytic anemia (WAIHA), and hemolytic disease of fetus and newborn (HDFN).
Nipocalimab is described in WO2019/118791 and comprises the following light (SEQ ID NO:4) and heavy (SEQ ID NO:5) chain sequences:

QSALTQPASVSGSPGQSITISCTGTGSDVGSYNLVSWYQQHPGKAPKLMIYGDSERPSGV
SNRFSGSKSGNTASLTISGLQAEDEADYYCSSYAGSGIYVFGTGTKVTVLGQPKAAPSVT
LFPPSSEELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNNKYAASS
YLSLTPEQWKSHKSYSCQVTHEGSTVEKTVAPTECS (SEQ ID NO:4)

EVQLLESGGGLVQPGGSLRLSCAASGFTFSTYAMGWVRQAPGKGLEWVSSIGASGSQTRY
ADSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCARLAIGDSYWGQGTMVTVSSASTK
GPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYS
LSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVF
LFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYASTYR
VVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN
QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN
VFSCSVMHEALHNHYTQKSLSLSPG (SEQ ID NO:5)

[0097] Another example of an antibody that binds specifically to FcRn and inhibits the binding of the Fc region of immunoglobulin to FcRn is rozanolixizumab, also known as UCB 7665.

Rozanolixizumab is a full-length (150 kDa) humanized IgG4 monoclonal antibody. Rozanolixizumab has been administered as a subcutaneous infusion in ongoing clinical trials for MG, immune thrombocytopenia (ITP), and chronic inflammatory demyelinating polyneuropathy (CIDP). Rozanolixizumab was first described in WO2014019727 and comprises the light chain of SEQ ID NO: 6 and the heavy chain of SEQ ID NO: 7.

DIQMTQSPSSLSASVGDRVTITCKSSQSLVGASGKTYLYWLFQKPGKAPKRLIYLVSTLD
SGIPSRFSGSGSGTEFTLTISSLQPEDFATYYCLQGTHFPHTFGQGTKLEIKRTVAAPSV
FIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSL
SSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:6)

EVPLVESGGGLVQPGGSLRLSCAVSGFTFSNYGMVWVRQAPGKGLEWVAYIDSDGDNTYY
RDSVKGRFTISRDNAKSSLYLQMNSLRAEDTAVYYCTTGIVRPFLYWGQGTLVTVSSAST
KGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY
SLSSVVTVPSSSLGTKTYTCNVDHKPSNTKVDKRVESKYGPPCPPCPAPEFLGGPSVFLF
PPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV
SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVF
SCSVMHEALHNHYTQKSLSLSLGK (SEQ ID NO:7)

[0098] A further example of an antibody that binds specifically to FcRn and inhibits the binding of the Fc region of immunoglobulin to FcRn is orilanolimab, also known as SYNT001. Orilanolimab is another full-length (150 kDa) humanized IgG4 monoclonal antibody. Orilanolimab has been administered as an intravenous infusion in Phase 2 clinical trials for treatment of WAIHA.

[0099] Yet another example of an antibody that binds specifically to FcRn and inhibits the binding of the Fc region of immunoglobulin to FcRn is batoclimab, also known as IMVT1401/RVT1401/HBM9161. Batoclimab is another full-length (150 kDa) "Fc dead" IgG1 monoclonal antibody. Batoclimab has been administered as a subcutaneous injection in ongoing Phase 2 clinical trials for treatment of MG and Graves' ophthalmopathy.

[0100] As used herein, the term "CD16" refers to FcγRIII Fc receptors that are required for Antibody-Dependent Cell-mediated Cytotoxicity (ADCC). Exemplary CD16 molecules include human CD16a as set forth in RefSeq NM_000569.

[0101] As used herein, the term "free cysteine" refers to native or engineered cysteine amino acid residue that exists in a substantially reduced form in a mature FcRn antagonist.

[0102] As used herein, the term "antibody" refers to immunoglobulin molecules comprising four polypeptide chains, two heavy (H) chains and two light (L) chains interconnected by disulfide bonds, as well as multimers thereof (e.g., IgM). Each heavy chain comprises a heavy chain variable region (abbreviated VH) and a heavy chain constant region. The heavy chain constant region comprises three domains, CH1, CH2 and CH3. Each light chain comprises a light chain variable region (abbreviated VL) and a light chain constant region. The light chain constant region comprises one domain (CL). The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDRs), interspersed

with regions that are more conserved, termed framework regions (FR).

[0103] As used herein, an "antigen-binding fragment" of an antibody generally comprises at least one VH paired with one VL, which together are capable of specifically binding to a particular antigen or epitope. Antigen-binding fragments can include, without limitation, Fv, Fab, Fab', and $F(ab')_2$ fragments, as well as engineered single-chain FV (scFV) fragments, diabodies, and the like.

[0104] As used herein the term "N-linked glycan" refers to the N-linked glycan attached to the nitrogen (N) in the side chain of asparagine in the sequon (i.e., Asn-X-Ser or Asn-X-Thr sequence, where X is any amino acid except proline) present in the CH2 domain of an Fc region. Such N-Glycans are fully described in, for example, Drickamer K, Taylor ME (2006). Introduction to Glycobiology, 2nd ed.

[0105] As used herein the term "afucosylated" refers to an N-linked glycan which lacks a core fucose molecule as described in US 8067232.

[0106] As used herein the term "bisecting GlcNAc" refers to an N-linked glycan having an N-acetylglucosamine (GlcNAc) molecule linked to a core mannose molecule, as described in US 8021856.

[0107] As used herein, the term "treat", "treating", and "treatment" refer to therapeutic or preventative measures described herein. In certain embodiments, the term "treat", "treating", and "treatment" refer to therapeutic measures described herein. The methods of "treatment" employ administration to a subject, an agent or combination of agents as described herein in order to prevent, cure, delay, reduce the severity of, or ameliorate one or more symptoms of a disease or disorder or recurring disease or disorder, or in order to prolong the survival of a subject beyond that expected in the absence of such treatment.

[0108] As used herein, the term "subject" includes any human or non-human mammal.

Pemphigus

[0109] Pemphigus is a group of chronic blistering epithelial diseases in which the production of IgG autoantibodies against extracellular domains of cell membrane proteins of keratinocytes results in acantholysis (loss of cell-cell adhesion between keratinocytes). Three major forms of pemphigus are: pemphigus vulgaris (PV), pemphigus foliaceus (PF), and paraneoplastic pemphigus (PP).

[0110] PV is caused by IgG autoantibodies against the desmosomal proteins, desmoglein-3 (Dsg-3) and/or desmoglein-1 (Dsg-1), on epidermal keratinocytes. Because the binding of the antibodies (IgG4 predominant, do not activate complement) to the extracellular domain of Dsg is sufficient to cause loss of keratinocyte adhesion and blister formation, they generate directly

the clinical manifestations of PV. Dsg-3 is expressed almost exclusively in the basal and parabasal cell layers of the skin and throughout the squamous layer of mucus membranes, whereas Dsg-1 prevails in the superficial layer of the skin and is nearly absent in the mucosa. Accordingly, mucosal PV lesions are mostly induced by anti-Dsg 3 antibodies, whereas cutaneous PV lesions are triggered by both anti-Dsg 3 and anti-Dsg 1 antibodies. Interestingly, disease activity has been shown to be closely correlated with serum levels of antibodies against Dsg-1 and, to a lesser extent, Dsg-3. Belloni-Fortina A et al., Clin Dev Immunol. 2009; 187864. doi: 10.1155/2009/187864; Abasq C et al., Arch Dermatol. 2009; 145(5):529-35.

[0111] PF is caused by antibodies against Dsg-1, which is expressed in the superficial layer of the epidermis and, therefore, involves the skin only. Two forms have been described: the non-endemic form, and the endemic form (also called "folgo selvagem"). Folgo selvagem has been observed in the Amazonian region, where a non-infectious protein (LJM11) residing in the salivary glands of the sand fly (*Lutzomyia longipalpis*) was suggested to cause a cross-reaction with Dsg-1. Otherwise, endemic and non-endemic PF share the same clinical, histological and immunological findings. Like in PV, anti-Dsg 1 autoantibodies are directly pathogenic.

[0112] PV and PF are rare. Pemphigus vulgaris is the most common subtype of pemphigus in Europe, the United States and Japan; it preferentially affects women, and most of the patients are 50-60 years of age at disease onset (Kasperkiewicz, Nat. Re. Dis. Primers, 2017 May 11;3:17026). Pemphigus foliaceus is the most common type observed in South America and North Africa owing to the endemic form, with sex predisposition differing among the regions and a preferential occurrence in young adults. PF is less common in North America and Europe (10-15% of pemphigus cases).

[0113] Both PV and PF are chronic and intractable, with a life-threatening potential. Clinically, PV presents as a mucosal-dominant, mucocutaneous or, less commonly, solely cutaneous type. Patients frequently shift from one type to another, typically from mucosal type at the beginning of the disease to become mucocutaneous later on. Mucocutaneous type tends to be a more severe disease, whereas diagnostic delay is common in the mucosal type. Although it may affect a wider range of age, its peak frequency ranges between 50 and 60 years of age. Women are slightly overrepresented in the PV population, in which the female-predominant thyroid diseases and rheumatoid arthritis are associated. Typically, lesions begin in the oral mucosa and may then extend to other mucosal areas and the skin. Mucosal involvement consists of flaccid blisters that rapidly rupture, leaving painful erosions. Mucosal lesions may also affect the pharynx, upper larynx, esophagus, nose and eyes, and genitals. They are usually associated with significant impairments, including difficulties of eating and swallowing, of having sexual intercourse, etc. They frequently lead to weight loss, malnutrition, and alteration of quality of life. Cutaneous lesions are characterized by flaccid blisters and erosions that easily ooze and become superinfected (crusty lesions). Although any skin area may be involved, lesions predominate in the head, upper trunk and groin. They are painful, especially in the folds. At examination, the epidermis can be detached when the finger rubs the skin at the periphery of the lesions (Nikolsky sign), which is highly indicative of PV diagnosis.

[0114] Diagnosis of PV follows an algorithm of different tools, which generally are performed in the presence of clinical manifestations suggestive of the disease. They include histology, direct immunofluorescence (DIF), and the detection of autoantibodies against Dsg-3 and/or Dsg-1 in serum either through indirect immunofluorescence (IIF) or enzyme-linked immunosorbent assay (ELISA) tests. Histopathology typically shows supra-basal acantholysis (keratinocytes floating in blister fluid). Acantholysis is highly suggestive of PV, as it is not seen in other autoimmune blistering diseases such as bullous pemphigoid (BP), but the diagnosis should be confirmed by the characteristic deposition of IgG and/or complement on the cell surface of keratinocytes by DIF. Today, while DIF is considered as the gold standard investigation for diagnosis, IIF and/or ELISA are necessary to confirm the diagnosis. Commercial ELISA assays are available for quantitative measurement of Dsg-1 and Dsg-3 autoantibodies in serum. They potentially offer advantages over IIF, such as increased sensitivity (>90%), but they are not helpful for excluding other self-antigens and autoimmune bullous dermatoses (AIBD). Therefore, IIF and ELISA may be considered complementary in the diagnostic investigation of PV. Taken together, current international guidelines recommend making a diagnosis of PV in patients with indicative clinical signs, confirmed by histopathology, positive DIF, and positive IIF and/or ELISA. Hertl M et al., J EurAcad Dermatol Venereol. 2015; 29(3):405-14.

[0115] PV is a chronic disease, with no tendency of spontaneous improvement. On the contrary, the disease typically worsens progressively and has a mortality rate three times higher than in the general population when untreated. Under treatment, the disease usually evolves in periods of remission and relapse. Eventually, it takes many years for achieving a definite cure. As treatments are part of the co-morbidity factors that are related to their high rate of serious side effects, mortality is still high, and severe infections remain today the main cause of death. Ren Z et al., J Eur Acad Dermatol Venereol. 2018; 32(10):1768-1776.

[0116] Because Dsg-1 is expressed in the superficial portion of the epidermis, PF patients typically present with itchy, scaly and crusted erosions of the cutaneous tissue. Blisters are uncommonly seen, owing to their superficial nature and easy rupture. Another clinical variant may be the development of squamous and crusty lesions on the face, scalp, chest and inter-scapular areas ("seborrheic pemphigus"). In more severe forms, desquamative erythroderma may be observed involving almost all of the skin surface. PF diagnosis follows the investigational tools which are used in PV. At histology, a cleft in the subcorneal or the superficial granular layer can be seen. DIF shows the intercellular deposition of IgG within the epidermis, and IIF reveals the presence of serum autoantibodies against intercellular components of the skin. ELISA tests quantify the positive level of anti-Dsg-1 antibodies in serum, whereas the search for anti-Dsg-3 antibodies is negative.

[0117] Like PV, PF does not tend to resolve spontaneously. Treatment is always needed, with the aim of healing existing lesions and preventing the appearance of new lesions as soon as possible.

[0118] Additional forms of pemphigus include pemphigus vegitans, pemphigus erythematosus, herpetiform pemphigus, and drug-induced pemphigus. Pemphigus vegitans is a variant of PV

characterized by fungoid vegetations and mediated by anti-Dsg 3 IgG autoantibodies. Pemphigus erythematosus is a variant of PF characterized by localized involvement, mainly on the face and upper part of the chest and back, and mediated by anti-Dsg 1 IgG autoantibodies. Herpetiform pemphigus is a subtype characterized by small vesicles and pustules, and mediated mainly by anti-Dsg 1 IgG autoantibodies.

Current Approaches to Treatment of Pemphigus

[0119] To help the practitioners in the therapeutic management of pemphigus patients, the following consensual definitions of the most important clinical endpoints among national and international experts, as well as regular guidelines for the diagnosis and treatment of these diseases are available (see Murrell et al., J Am Acad Dermatol. 2008; 58: 1043; Murrell et al., J Am Acad Dermatol. 2018 Feb 10; Harman et al., Br J Dermatol. 2017 Nov;177(5):1170-1201; Joly et al., Lancet. 2017 May 20;389(10083):2031-2040), Hébert et al., J Invest Dermatol. 2019 Jan;139(1):31-37):

1. (a) "Disease Control" (DC) - no new pemphigus lesions and established lesions begin to heal;
2. (b) "End of Consolidation Phase" (EoC) - no new pemphigus lesions for at least 2 weeks and about 80% established lesions are healed.
3. (c) "Complete Remission" (CR) - absence of new or established lesions, also referred to as "Complete Clinical Remission".
4. (d) "Complete Remission Off Therapy" (CRoff) - absence of new or established lesions while the patient is off all systemic therapy for at least 2 months.
5. (e) "Complete Remission On Therapy" (CRmin) - absence of new or established lesions while the patient is receiving Minimal Therapy.
6. (f) "Minimal Therapy" - prednisone (or the equivalent) at a dose of 10 mg/day or less and/or minimal adjuvant therapy for at least 2 months (for practical reasons during clinical trials this is shortened to 8 weeks).
7. (g) "Off Therapy" - no systemic drug for at least 2 months.
8. (h) "Relapse" or "Flare" (R) - appearance of 3 or more lesions a month that do not heal within 1 week, or extension of established lesions.
9. (i) "Treatment Failure" - failure to reach disease control with full therapeutic doses of systemic treatments (e.g., prednisone 1.5 mg/kg/day for 3 weeks).

[0120] In addition, validated scales for monitoring disease activity and extension (have been established (See, e.g., Rosenbach M et al., J Invest Dermatol. 2009; 129(10):2404-10; Rahbar Z et al., JAMA Dermatol. 2014; 150(3):266-72). Exemplary validated scores include the Pemphigus Disease Area Index (PDAI) and the Autoimmune Bullous Skin disorder Intensity Score (ABSIS). The pemphigus Disease Area Index (PDAI) is a well-established and widely used diagnostic tool used to assess and classify the severity of pemphigus in afflicted human

subjects. Using the PDAI, the severity of disease is scored on a scale from 0-263, where a score of 0 signifies no disease and a score of 263 signifies maximally severe disease.

[0121] Severity thresholds for mild, moderate, and severe pemphigus have been suggested in different papers with partially overlapping patient populations proposing PDAI 9 or 15 as cut-off values between mild and moderate/severe disease, and 25 or 45 as cut-off values between moderate and severe disease (Shimizu et al., J Dermatol. 2014; 41(11):969-973; Boulard et al., Br J Dennatol. 2016; 175(1):142-9). International guidelines concluded that it is currently premature to definitively state cut-off values to define mild, moderate, or severe disease (Murrell et al., J Am Acad Dermatol. 2020; 82(3):575-585).

[0122] In exemplary embodiments, "Mild Pemphigus" is characterized as a PDAI score of less than 15; "Moderate Pemphigus" is characterized by a PDAI score of 15 to less than 45 and "Severe Pemphigus" is characterized by a PDA Score of 45 or greater. The Autoimmune Bullous Disease Quality of Life (ABQOL) score was developed and validated for ascertaining the impact of the disease and its therapies on patients' daily lives. Sebaratnam DF et al., JAMA Dermatol. 2013; 149(10):1186-91.

[0123] According to the most recent international guideline (Ren Z et al., J Eur Acad Dermatol Venereol. 2018; 32(10):1768-1776), corticosteroids remain a first line therapy in pemphigus. They are the most rapidly acting form of treatment known today and provide disease control (no new lesions, established lesions starting to heal) in about 3 weeks when used at effective dose. Ratnam KV et al., Int J Dermatol. 1990; 29(5):363-7; Czernik A et al., Arch Dermatol. 2008; 144(5):658-61; Chaidemenos G et al., J Eur Acad Dermatol Venereol. 2011; 25(2):206-10. Oral prednisone is the most commonly used corticosteroid. The starting oral prednisone dose is high, ranging from 1-2 mg/kg daily, and may be reduced (0.5-1 mg/kg per day) if combined with rituximab or immunosuppressants. If disease control is not achieved after 3-4 weeks at the latest, the prednisone dose must be increased. In patients with very active disease, intravenous bolus of corticosteroids (e.g. methylprednisolone) may be preferred, especially at the treatment initiation.

[0124] As used herein, the term "prednisone equivalent dose" means a dose of prednisone or an equivalent dose of a systemic corticosteroid other than prednisone. Systemic corticosteroids are well-known and include compounds of various potencies and formulations. These are generally formulated as injectables or pills. Examples of commercially available systemic corticosteroids include, without limitation, betamethasone, cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisone, prednisolone, and triamcinolone.

[0125] In certain embodiments, a topical corticosteroid may also be administered. Topical corticosteroids are well-known and include compounds of various potencies and formulations. These are generally formulated as ointments, creams, oils, lotions, shampoos, foams, and/or gels. Examples of commercially available topical corticosteroids include alclometasone dipropionate, amcinonide, betamethasone dipropionate, betamethasone valerate, clobetasol propionate, clobetasone butyrate, desonide, desoximetasone, diflucortolone valerate,

diflorasone diacetate, fluocinolone acetonide, fluocinonide, flurandrenolide, fluticasone propionate, halcinonide, halobetasol propionate, hydrocortisone, hydrocortisone 17-butyrate, hydrocortisone acetate, hydrocortisone valerate, methylprednisolone aceponate, mometasone furoate, prenicarbate, and triamcinolone acetonide. Topical corticosteroids conveniently can be applied locally to affected areas or lesions in need of treatment, thereby limiting undesirable systemic effects of corticosteroid treatment.

[0126] The therapeutic management of patients with pemphigus is very challenging. Its primary principles are, on the one hand, to promptly stop the occurrence of new blisters and then achieve clinical remission and, on the other hand, to minimize the side effects of systemic therapy, e.g., corticosteroids. Corticosteroids have a high cumulative toxicity in these chronic diseases that must be mitigated by reducing their amount and duration of exposure as much as possible. Secondarily, the objective of the treatments is to prevent the development of new lesions either on minimal corticosteroid therapy, or even without any treatment. For this purpose, patients must be managed by referral centers, where they can be monitored carefully by experienced dermatologists. In many patients, a multi-disciplinary approach is necessary, including specialists in oral medicine, ophthalmologists, and gynecologists when mucosal lesions are present.

[0127] After disease control, oral prednisone is maintained at a stable dose for consolidating disease control. End of consolidation is defined as the absence of new lesions for at least 2 weeks and about 80% of established lesions already healed. This time point is chosen by a majority of practitioners as the inflexion time at which prednisone starts to be tapered. The duration of the consolidation period varies greatly between patients, mucosal erosions and extensive cutaneous lesions tending to be late to heal.

[0128] Once disease control is consolidated, a tapering schedule is put in place, during which the prednisone dose is gradually reduced. The goal of this phase is to reach complete clinical remission (no new lesions, all established lesions completely healed) and, at the same time, reach a minimal effective dose of prednisone (10 mg per day or less for at least 2 months), or even stop prednisone (off-therapy), in order to prevent side effects. Unfortunately, this double objective is hard to achieve and a majority of patients relapse under tapered dose of prednisone. In a prospective trial, 64% of PV patients with prednisone achieved a first clinical remission on minimal dose after 12 months, but relapse occurred frequently, i.e. in 45% of the cases within 6 months. Beissert S et al., J Invest Dermatol. 2010; 130(8):2041-8. Finally, the common course of PV is one of episodes of relapse and transient remission, and it takes several years of prednisone treatment before achieving permanent remission. In one study, 36% of patients with PV were treated with prednisone or equivalent for at least 10 years. Mimouni D et al., J Eur Acad Dermatol Venereol. 2010; 24(8):947-52. Meanwhile, the risk of corticosteroid-related side effects (osteoporosis, diabetes, hypertension, Cushing syndrome, cataracts, glaucoma, infections, etc.) increases with treatment duration. Accordingly, most PV patients need adjuvant therapies (e.g., rituximab, immunosuppressants) in order to maintain them under remission and reduce the cumulative prednisone dose.

[0129] As used herein, the term "initial dose" means a starting corticosteroid dose or initial dose of corticosteroids as administered to a pemphigus patient (before tapering).

[0130] As used herein, the term "tapering dose" means a reduced corticosteroid dose as administered to a pemphigus patient. The term "initial tapering dose" means the first, reduced corticosteroid dose as administered to a pemphigus patient, e.g. a dose of 0.5 mg/kg/day prednisone or equivalent. Further reduced tapering doses may be administered. The reduction of the initial and/or subsequent tapering dose can occur on a periodic basis, e.g. every 2 weeks.

[0131] The recent approval of rituximab in PV adults has dramatically changed the mainstay of therapy of the disease, and is now proposed as a first-line therapy in association with prednisone for the more severe cases. Ren Z et al., J Eur Acad Dermatol Venereol. 2018; 32(10): 1768-1776. Until recently, rituximab was primarily used in refractory PV patients, who do not respond well to other treatments. However, the lack of long-lasting remission and great number of serious adverse events associated with prednisone and immunosuppressants has led researchers to develop alternative first-line treatments. Rituximab as second-line and third-line treatment (1-2 g per cycle) was shown to induce long-term remission at a high rate, which could not be achieved by any other treatment (75% remission rate in PV after 1 year). Relapse rates after rituximab therapy ranged between 25% after 1 year up to 80% in long-term follow-up. Wang HH et al., Acta Derm Venereol. 2015; 95(8):928-32. Recent evidence has shown efficacy of a first cycle of rituximab (2 g) as first-line treatment followed by new cycles (0.5 g) after 12 months and 18 months, in combination with lower doses of prednisone (0.5-1 mg/kg per day according to disease severity). Joly et al., Lancet. 2017 May 20;389(10083):2031-2040). Using this first-line regimen, a higher proportion of complete remission (in this context what is meant is Complete Remission Off Therapy) after 2 years, i.e. 89%, was demonstrated. Relapse cases were observed to be lower (24% after 2 years), a majority occurring within 6 to 12 months after the first cycle. A strong prednisone sparing effect, i.e. by about two thirds, was demonstrated.

[0132] Nevertheless, rituximab, either used as first-line or second-/third-line treatment, has a late onset of action and does not result in an early achievement of disease control (mean time from first infusion: 6-7 weeks) and clinical remission (mean of 6-7 months on minimal prednisone therapy, 9 months off-therapy). Hébert V et al., J Invest Dermatol. 2019; 139(1):31-37; Wang HH et al., Acta Derm Venereol. 2015; 95(8):928-32. Furthermore, rituximab therapy carries the risk of inducing serious adverse events (e.g. 33% of infections). Late-onset neutropenia and hypogammaglobulinemia and the risk for patients to develop potentially fatal infections (e.g. *Pneumocystis carinii,* multifocal leukoencephalopathy, septicemia) require a long-term monitoring of the clinical and immunological status of the patients under rituximab.

[0133] Patients with PF typically follow the same type of management and monitoring of clinical outcome measures as PV patients, although rituximab has not been approved for this condition.

FcRn Antagonists

[0134] The neonatal Fc receptor (FcRn) influences serum levels and tissue distribution of IgG during all stages of life. FcRn is an intracellular trafficking, integral membrane receptor for IgG. Consequently, FcRn is a multifunctional molecule primarily involved in IgG transport and homeostasis. Challa DK et al., Curr Top Microbiol Immunol. 2014; 382:249-72; Roopenian DC et al., Nat Rev Immunol. 2007; 7(9):715-25. Following IgG uptake by pinocytosis in FcRn-expressing cells, the Fc part of IgGs binds FcRn with high affinity in early, acidic endosomes (pH <6.5). This binding spares IgGs from lysosomal degradation and drives them to the cell surface for recycling. At the near-neutral pH (pH ~7.4) of the extracellular space, IgGs are released from the complex with FcRn. This pH-dependent salvage pathway is accountable for the maintenance of high IgG concentrations in circulation and for the long half-life ($t_{1/2}$) of IgGs compared to other Igs, which are not recycled. Roopenian DC et al., J Immunol. 2003; 170(7):3528-33; Waldmann TA et al., J Clin Invest. 1990; 86(6):2093-2098; Wani MA et al., Proc Natl Acad Sci U S A. 2006; 103(13):5084-5089.

[0135] In one aspect, the invention provides an FcRn antagonist composition for use in the treatment of pemphigus in a subject, wherein

1. (i) the subject has
    1. (a) newly diagnosed pemphigus,
    2. (b) relapsing pemphigus, or
    3. (c) refractory pemphigus,
2. (ii) the FcRn antagonist consists of a variant Fc region, wherein said variant Fc region consists of two Fc domains which form a homodimer, and wherein the amino acid sequence of each of the Fc domains is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, and SEQ ID NO: 3; and the FcRn antagonist is administered in combination with a corticosteroid and the corticosteroid is at a dose of ≤ 0.5 mg prednisone/kg/day or equivalent.

[0136] In certain embodiments, these compositions comprise or consist of a variant Fc region, or FcRn-binding fragment thereof, that binds specifically to FcRn, particularly human FcRn, with increased affinity and reduced pH dependence relative to a native Fc region. The FcRn antagonist compositions described herein can comprise an antibody or antigen-binding fragment thereof that binds specifically to FcRn via its antigen binding domain and inhibits the binding of Fc region of immunoglobulin to FcRn. In general, these FcRn antagonists inhibit the binding of Fc-containing agents (e.g., antibodies and immunoadhesins) to FcRn *in vivo,* which results in an increased rate of degradation of the Fc-containing agents and, concomitantly, a reduced serum level of these agents.

[0137] An isolated variant Fc region (e.g., a variant Fc region comprising the amino acids Y, T,

E, K, F, and Y at EU positions 252, 254, 256, 433, 434, and 436 respectively) is a more efficacious FcRn antagonist *in vivo* than a full-length antibody comprising the same variant Fc region. Accordingly the FcRn antagonist compositions for use according to the invention are not full-length antibodies. The FcRn antagonist compositions for use according to the invention do not comprise an antibody variable domain and moreover do not comprise an antibody variable domain or a CH1 domain.

[0138] Any Fc region can be altered to produce a variant Fc region for use in the FcRn antagonist compositions disclosed herein. In general, an Fc region, or FcRn-binding fragment thereof, is from a human immunoglobulin. It is understood, however, that the Fc region may be derived from an immunoglobulin of any other mammalian species, including for example, a Camelid species, a rodent (e.g. a mouse, rat, rabbit, guinea pig) or non-human primate (e.g. chimpanzee, macaque) species. Moreover, the Fc region or portion thereof may be derived from any immunoglobulin class, including IgM, IgG, IgD, IgA and IgE, and any immunoglobulin isotype, including IgG1, IgG2, IgG3 and IgG4. In certain embodiments, the Fc region is an IgG Fc region (e.g., a human IgG region). In certain embodiments, the Fc region is an IgG1 Fc region (e.g., a human IgG1 region). In certain embodiments, the Fc region is a chimeric Fc region comprising portions of several different Fc regions. Suitable examples of chimeric Fc regions are set forth in US 2011/0243966A1. A variety of Fc region gene sequences (e.g. human constant region gene sequences) are available in the form of publicly accessible deposits. It will be appreciated that the scope of this disclosure encompasses alleles, variants and mutations of Fc regions.

[0139] An Fc region can be further truncated or internally deleted to produce a minimal FcRn-binding fragment thereof. The ability of an Fc-region fragment to bind to FcRn can be determined using any art recognized binding assay e.g., ELISA.

[0140] To enhance the manufacturability of the FcRn antagonists disclosed herein, it is preferable that the constituent Fc regions do not comprise any non-disulphide bonded cysteine residues. Accordingly, in certain embodiments the Fc regions do not comprise a free cysteine residue.

[0141] Any Fc variant, or FcRn-binding fragment thereof, that binds specifically to FcRn with increased affinity and reduced pH dependence relative to the native Fc region can be used in the FcRn antagonist compositions disclosed herein. In certain embodiments, the variant Fc region comprises amino acid alterations, substitutions, insertions and/or deletions that confer the desired characteristics. The variant Fc region or fragment for use according to the invention comprises the amino acids Y, T, E, K, F, and Y at EU positions 252, 254, 256, 433, 434, and 436, respectively. The amino acid sequences that can be used in variant Fc regions that are for use in accordance with the invention are set forth in **Table 1,** below.

Table 1. Amino acid sequences of non-limiting examples of variant Fc regions

SEQ ID NO:	Amino Acid Sequence
1	DKTHTCPPCPAPELLGGPSVFLFPPKPKDTL**Y**I**T**R**E**PEVTCVVV DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSV LTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQ VYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPEN NYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHE AL**K**F**H**YTQKSLSLSPG
2	DKTHTCPPCPAPELLGGPSVFLFPPKPKDTL**Y**I**T**R**E**PEVTCVVV DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSV LTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQ VYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPEN
	NYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHE AL**K**F**H**YTQKSLSLSPGK
3	CPPCPAPELLGGPSVFLFPPKPKDTL**Y**I**T**R**E**PEVTCVVVDVSHE DPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLH QDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLP PSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL**K**F**H** YTQKSLSLSPG
Amino acids at EU positions 252, 254, 256, 433, and 434 are bold	

[0142] An FcRn-antagonist described herein may consist of a variant Fc region, wherein the amino acid sequence of the Fc domains of the variant Fc region comprises the amino acid sequence set forth in SEQ ID NO: 1, 2, or 3. The amino acid sequence of the Fc domains of the variant Fc region described herein may comprise the amino acid sequence set forth in SEQ ID NO: 1. The amino acid sequence of the Fc domains of the variant Fc region described herein may comprise the amino acid sequence set forth in SEQ ID NO: 2. The amino acid sequence of the Fc domains of the variant Fc region described herein may comprise the amino acid sequence set forth in SEQ ID NO: 3.

[0143] In certain embodiments an FcRn-antagonist consists of a variant Fc region, wherein the amino acid sequence of the Fc domains of the variant Fc region consists of the amino acid sequence set forth in SEQ ID NO: 1, 2, or 3. In certain embodiments, the amino acid sequence

of the Fc domains of the variant Fc region consists of the amino acid sequence set forth in SEQ ID NO: 1. In certain embodiments, the amino acid sequence of the Fc domains of the variant Fc region consists of the amino acid sequence set forth in SEQ ID NO: 2. In certain embodiments, the amino acid sequence of the Fc domains of the variant Fc region consists of the amino acid sequence set forth in SEQ ID NO: 3.

[0144] An FcRn-antagonist for use according to the invention consists of a variant Fc region, wherein said variant Fc region consists of two Fc domains which form a homodimer, wherein the amino acid sequence of each of the Fc domains is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, and SEQ ID NO: 3. In certain embodiments an FcRn-antagonist consists of a variant Fc region, wherein said variant Fc region consists of two Fc domains which form a homodimer, wherein the amino acid sequence of each of the Fc domains is SEQ ID NO: 1. In certain embodiments an FcRn-antagonist consists of a variant Fc region, wherein said variant Fc region consists of two Fc domains which form a homodimer, wherein the amino acid sequence of each of the Fc domains is SEQ ID NO: 2. In certain embodiments an FcRn-antagonist consists of a variant Fc region, wherein said variant Fc region consists of two Fc domains which form a homodimer, wherein the amino acid sequence of each of the Fc domains is SEQ ID NO: 3.

[0145] In certain embodiments, the variant Fc region has altered (e.g., increased or decreased) binding affinity for an additional Fc receptor. The variant Fc region can have altered (e.g., increased or decreased) binding affinity for one or more of Fcγ receptors e.g., FcγRI (CD64), FcγRIIA (CD32a), FcγRIIB (CD32b), FcγRIIIA (CD16a), and FcγRIIIB (CD16b). Any art-recognized means of altering the affinity for an additional Fc receptor can be employed. In certain embodiments, the amino acid sequence of the variant Fc region is altered.

[0146] The variant Fc region described herein may comprise a non-naturally occurring amino acid residue at one or more positions selected from the group consisting of 234, 235, 236, 239, 240, 241, 243, 244, 245, 247, 252, 254, 256, 262, 263, 264, 265, 266, 267, 269, 296, 297, 298, 299, 313, 325, 326, 327, 328, 329, 330, 332, 333, and 334 as numbered by the EU index as set forth in Kabat. Optionally, the Fc region may comprise a non-naturally occurring amino acid residue at additional and/or alternative positions known to one skilled in the art (see, e.g., U.S. Pat. Nos. 5,624,821; 6,277,375; 6,737,056; PCT Patent Publications WO 01/58957; WO 02/06919; WO 04/016750; WO 04/029207; WO 04/035752 and WO 05/040217).

[0147] The variant Fc region described herein may comprise at least one non-naturally occurring amino acid residue selected from the group consisting of 234D, 234E, 234N, 234Q, 234T, 234H, 234Y, 234I, 234V, 234F, 235A, 235D, 235R, 235W, 235P, 235S, 235N, 235Q, 235T, 235H, 235Y, 235I, 235V, 235F, 236E, 239D, 239E, 239N, 239Q, 239F, 239T, 239H, 239Y, 2401, 240A, 240T, 240M, 241W, 241 L, 241Y, 241E, 241R. 243W, 243L 243Y, 243R, 243Q, 244H, 245A, 247V, 247G, 252Y, 254T, 256E, 262I, 262A, 262T, 262E, 263I, 263A, 263T, 263M, 264L, 264I, 264W, 264T, 264R, 264F, 264M, 264Y, 264E, 265G, 265N, 265Q, 265Y, 265F, 265V, 265I, 265L, 265H, 265T, 266I, 266A, 266T, 266M, 267Q, 267L, 269H, 269Y, 269F, 269R,

296E, 296Q, 296D, 296N, 296S, 296T, 296L, 296I, 296H, 269G, 297S, 297D, 297E, 298H, 298I, 298T, 298F, 299I, 299L, 299A, 299S, 299V, 299H, 299F, 299E, 313F, 325Q, 325L, 325I, 325D, 325E, 325A, 325T, 325V, 325H, 327G, 327W, 327N, 327L, 328S, 328M, 328D, 328E, 328N, 328Q, 328F, 328I, 328V, 328T, 328H, 328A, 329F, 329H, 329Q, 330K, 330G, 330T, 330C, 330L, 330Y, 330V, 330I, 330F, 330R, 330H, 332D, 332S, 332W, 332F, 332E, 332N, 332Q, 332T, 332H, 332Y, and 332A as numbered by the EU index as set forth in Kabat. Optionally, the Fc region may comprise additional and/or alternative non-naturally occurring amino acid residues known to one skilled in the art (see, e.g., U.S. Pat. Nos. 5,624,821; 6,277,375; 6,737,056; PCT Patent Publications WO 01/58957; WO 02/06919; WO 04/016750; WO 04/029207; WO 04/035752 and WO 05/040217).

[0148] Other known Fc variants that may be used in the FcRn antagonists disclosed herein include without limitations those disclosed in Ghetie et al., 1997, Nat. Biotech. 15:637-40; Duncan et al, 1988, Nature 332:563-564; Lund et al., 1991, J. Immunol., 147:2657-2662; Lund et al, 1992, Mol. Immunol., 29:53-59; Alegre et al, 1994, Transplantation 57:1537-1543; Hutchins et al., 1995, Proc Natl. Acad Sci USA, 92:11980-11984; Jefferis et al, 1995, Immunol Lett., 44:111-117; Lund et al., 1995, FASEB J., 9:115-119; Jefferis et al, 1996, Immunol Lett., 54:101-104; Lund et al, 1996, J. Immunol., 157:4963-4969; Armour et al., 1999, Eur J Immunol 29:2613-2624; Idusogie et al, 2000, J. Immunol., 164:4178-4184; Reddy et al, 2000, J. Immunol., 164:1925-1933; Xu et al., 2000, Cell Immunol., 200:16-26; Idusogie et al, 2001, J. Immunol., 166:2571-2575; Shields et al., 2001, J Biol. Chem., 276:6591-6604; Jefferis et al, 2002, Immunol Lett., 82:57-65; Presta et al., 2002, Biochem Soc Trans., 30:487-490); U.S. Pat. Nos. 5,624,821; 5,885,573; 5,677,425; 6,165,745; 6,277,375; 5,869,046; 6,121,022; 5,624,821; 5,648,260; 6,528,624; 6,194,551; 6,737,056; 6,821,505; 6,277,375; U.S. Patent Publication Nos. 2004/0002587 and PCT Publications WO 94/29351; WO 99/58572; WO 00/42072; WO 02/060919; WO 04/029207; WO 04/099249; WO 04/063351.

[0149] The variant Fc region described herein can also be a heterodimer, where the constituent Fc domains are different from each other. Methods of producing Fc heterodimers are known in the art (see e.g., US 8216805). The variant Fc region described herein could also be a single chain Fc region, where the constituent Fc domains are linked together by a linker moiety. Methods of producing single chain Fc regions are known in the art (see e.g., US 2009/0252729A1 and US 2011/0081345A1).

[0150] It is believed that pathogenic IgG antibodies observed in autoimmune diseases are either the pathogenic triggers for these diseases or contribute to disease progression and mediate disease through the inappropriate activation of cellular Fc receptors. Aggregated autoantibodies and/or autoantibodies complexed with self antigens (immune complexes) bind to activating Fc receptors, causing numerous autoimmune diseases (which occur in part because of immunologically mediated inflammation against self tissues) (see e.g., Clarkson et al., New Engl J Med 314(9), 1236-1239 (2013)); US 2004/0010124A1; US 2004/0047862A1; and US 2004/0265321A1). Accordingly, to treat antibody-mediated disorders (e.g. autoimmune diseases), it would be advantageous to both remove the deleterious autoantibodies and to block the interaction of the immune complexes of these antibodies with activating Fc receptors

(e.g., Fcγ receptors, such as CD16a).

[0151] Accordingly, in certain embodiments, the variant Fc region of the FcRn antagonist exhibits increased binding to CD16a (e.g., human CD16a). This is particularly advantageous in that it allows the FcRn antagonist to additionally antagonize the immune complex-induced inflammatory response of autoantibodies being targeted for removal by FcRn inhibition. Any art recognized means of increasing affinity for CD16a (e.g., human CD16a) can be employed. In certain embodiments, the FcRn-antagonist comprises a variant Fc-region comprising an N-linked glycan (e.g., at EU position 297). In this case it is possible to increase the binding affinity of the FcRn-antagonist for CD16a by altering the glycan structure. Alterations of the N-linked glycan of Fc regions are well known in the art. For example, afucosylated N-linked glycans or N-glycans having a bisecting GlcNAc structure have been shown to exhibit increased affinity for CD16a. Accordingly, in certain embodiments, the N-linked glycan is afucosylated. Afucosylation can be achieved using any art-recognized means. For example, an FcRn-antagonist can be expressed in cells lacking fucosyl transferase, such that fucose is not added to the N-linked glycan at EU position 297 of the variant Fc region (see e.g., US 8,067,232). In certain embodiments, the N-linked glycan has a bisecting GlcNAc structure. The bisecting GlcNAc structure can be achieved using any art recognized means. For example, an FcRn-antagonist can be expressed in cells expressing beta1-4-N-acetylglucosaminyltransferase III (GnTIII), such that bisecting GlcNAc is added to the N-linked glycan at EU position 297 of the variant Fc region (see e.g., US 8021856). Additionally or alternatively, alterations of the N-linked glycan structure can also be achieved by enzymatic means *in vitro.*

[0152] In certain embodiments, the instant disclosure provides FcRn-antagonist compositions wherein a portion of the FcRn-antagonist molecules contained therein comprise altered glycan structures. In certain embodiments, the FcRn-antagonist composition comprises a plurality of FcRn-antagonist molecules disclosed herein, wherein at least 50% (optionally, at least 60, 70, 80, 90, 95, or 99%) of the molecules comprise an Fc region or FcRn-binding fragment thereof having an afucosylated N-linked glycan. In certain embodiments, the FcRn-antagonist composition comprising a plurality of FcRn-antagonist molecules disclosed herein, wherein at least 50% (optionally, at least 60, 70, 80, 90, 95, or 99%) of the molecules comprise an Fc region or FcRn-binding fragment thereof comprising an N-linked glycan having a bisecting GlcNAc.

[0153] In certain embodiments, the variant Fc region does not comprise an N-linked glycan. This can be achieved using any art recognized methods. For example, the Fc variant can be expressed in a cell that is incapable of N-linked glycosylation. Additionally or alternatively, the amino acid sequence of the Fc variant can be altered to prevent or inhibit N-linked glycosylation (e.g., by mutation of the NXT sequon). Alternatively, the Fc variant can be synthesized in an acellular system (e.g., chemically synthesized).

[0154] In certain embodiments, FcRn-antagonist molecules may be modified, e.g., by the covalent attachment of a molecule (e.g., a binding or imaging moiety) to the FcRn-antagonist such that covalent attachment does not prevent the FcRn-antagonist from specifically binding

to FcRn. For example, but not by way of limitation, the FcRn-antagonist may be modified by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc.

[0155] In certain embodiments, the FcRn antagonist comprises a variant Fc region linked to a half-life extender. As used herein, the term "half-life extender" refers to any molecule that, when linked to an FcRn antagonist disclosed herein, increases the half-life of an FcRn antagonist. Any half-life extender may be linked (either covalently or non-covalently) to the FcRn antagonist. In certain embodiments, the half-life extender is polyethylene glycol or human serum albumin. In certain embodiments, the FcRn antagonist is linked to a binding molecule that specifically binds to a half-life extender present in a subject, such as a blood-carried molecule or cell, such as serum albumin (e.g., human serum albumin), IgG, erythrocytes, etc.

[0156] The FcRn antagonists disclosed herein have excellent manufacturability. For example, they can be expressed at high levels in mammalian cells (e.g., at 6g/L in CHO cells in a 10 L stirred tank bioreactor). Moreover, after Protein A purification, the resultant purified FcRn antagonist composition has a very high percentage of FcRn antagonist monomers, and contains an extremely low level of FcRn antagonist protein aggregates and degradation products. Accordingly, in certain embodiments, the instant disclosure provides an FcRn antagonist composition comprising a plurality of FcRn antagonist molecules as disclosed herein, wherein greater than 95% the of the FcRn antagonist molecules in the composition are monomers (e.g., greater than 95, 96, 97, 98, 99, 99.1, 99.2, 99.3, 99.4, 99.5, 99.6, 99.7, 99.8, 99.9 %). In certain embodiments, the instant disclosure provides an FcRn antagonist composition comprising a plurality of FcRn antagonist molecules disclosed herein, wherein less than 5% the of the FcRn antagonist molecules in the composition are present in aggregates, (e.g., less than 5, 4, 3, 2, 1, 0.9, 0.8, 0.7, 0.6, 0.5, 0.4, 0.3, 0.2, 0.1 %). In certain embodiments, the instant disclosure provides an FcRn antagonist composition comprising a plurality of FcRn antagonist molecules disclosed herein, wherein the composition is substantially free of FcRn antagonist molecule degradation products.

[0157] Methods for production of FcRn antagonists useful in the instant invention are disclosed in, for example, US 10,316,073.

[0158] The FcRn antagonist compositions for use in accordance with the invention are administered in combination with a corticosteroid. In some embodiments, the FcRn antagonist and corticosteroid can be used in combination with one or more additional therapeutic agents. In certain embodiments, the additional therapeutic agent is an anti-inflammatory agent. Any anti-inflammatory agent can be used in combination with the FcRn antagonists disclosed herein. Anti-inflammatory agents include corticosteroids, e.g., prednisone, prednisolone, methylprednisolone, cortisone, hydrocortisone, betamethasone, triamcinolone, and dexamethasone. In certain embodiments, the additional therapeutic agent is rituximab, daclizumab, basiliximab, muronomab-CD3, infliximab, adalimumab, omalizumab, efalizumab,

natalizumab, tocilizumab, eculizumab, golimumab, canakinumab, ustekinumab, or belimumab. In certain embodiments, the additional therapeutic agent is a leucocyte depleting agent (e.g., B-cell or T-cell depleting agent). Any leucocyte depleting agent can be used in combination with the FcRn antagonist compositions disclosed herein. In certain embodiments, the leucocyte depleting agent is a B-cell depleting agent. In certain embodiments, the leucocyte depleting agent is an antibody against a cell surface marker. Suitable cell surface markers include, without limitation, CD10, CD19, CD20, CD21, CD22, CD23, CD24, CD37, CD53, CD70, CD72, CD74, CD75, CD77, CD79a, CD79b, CD80, CD81, CD82, CD83, CD84, CD85, or CD86. In certain exemplary embodiments, the B-cell depleting agent is an antibody that binds CD20 (e.g., Rituxan). The FcRn antagonist and the additional therapeutic agent(s) can be administered to the subject simultaneously or sequentially, via the same or different route(s) of administration.

Efgartigimod

[0159] Efgartigimod (ARGX-113) is a modified human immunoglobulin (Ig) gamma (IgG) 1-derived Fc of the za allotype that binds with nanomolar affinity to human FcRn. Efgartigimod encompasses IgG1 residues D220-K447 (EU numbering scheme) and has been engineered using ABDEG™ technology to increase its affinity for FcRn at both physiological and acidic pH. Vaccaro C et al., Nat Biotechnol. 2005; 23(10):1283. See also U.S. Pat. No. 10,316,073. The increased affinity for FcRn of efgartigimod at both acidic and physiological pH results in a blockage of FcRn-mediated recycling of IgGs.

[0160] Efgartigimod has a molecular weight of about 54 kDa, which is about one-third the molecular weight of full-length IgG (MW ca. 150 kDa). Thus, 10 mg efgartigimod is about 185 nmol, such that a dose of 10 mg efgartigimod/kg body weight corresponds to about 185 nmol efgartigimod/kg body weight and a dose of 25 mg efgartigimod/kg of body weight corresponds to about 462.5 nmol efgartigimod/kg body weight. In contrast, a dose of 10 mg full-length IgG antibody/kg body weight corresponds to about 67 nmol/kg body weight. Furthermore, a 1000 mg fixed dose of efgartigimod corresponds to a fixed dose of about 18500 nmol of efgartigimod while a 2000 mg fixed dose of efgartigimod corresponds to a fixed dose of about 37000 nmol of efgartigimod.

[0161] Due to its increased affinity for FcRn at both acidic and neutral pH, efgartigimod blocks the FcRn/IgG complex from forming, which results in degradation of endogenous IgGs, including autoantibodies that cause IgG-mediated autoimmune diseases. This blocking of FcRn by efgartigimod results in a rapid and profound reduction in autoantibody levels, which underlies the therapeutic strategy for the treatment of autoimmune indications where IgG autoantibodies are expected to have a central role in the disease pathology, e.g., conditions such as pemphigus (PV and PF).

rHuPH20

[0162] Efgartigimod is under development for both the intravenous (IV) and subcutaneous (SC) administration route. For SC administration, in certain embodiments efgartigimod may be administered alone. Alternatively, for SC administration, in certain embodiments efgartigimod may be administered co-formulated with hyaluronidase, for example, in particular, rHuPH20. The co-formulated material will allow dosing of higher volumes.

[0163] rHuPH20 is the active ingredient of Halozyme's commercial product HYLENEX® recombinant (hyaluronidase human injection), referred to as HYLENEX®, which was approved by FDA for marketed use in the US in December 2005. HYLENEX® is a tissue permeability modifier indicated as an adjuvant in SC fluid administration for achieving hydration, to increase the dispersion and absorption of other injected drugs, and in SC urography, for improving resorption of radiopaque agents.

[0164] rHuPH20 is a recombinant enzyme human hyaluronidase produced by genetically engineered Chinese hamster ovary (CHO) cells containing a deoxyribonucleic plasmid encoding a soluble fragment of human hyaluronidase (posterior head protein 20 [PH20]).

[0165] The HZ202 rHuPH20 DS is currently registered in HYLENEX® and other biologic drug products co-formulated with rHuPH20 DS. As such, in certain embodiments HZ202 rHuPH20 DS is used in the efgartigimod / rHuPH20 co-formulated product for SC administration (i.e., efgartigimod PH20 SC).

[0166] SC injection volumes are typically limited to 2.5 mL due to concerns regarding injection pain associated with larger volumes. It has been demonstrated that rHuPH20 offers a solution to the volume limitation associated with fast SC injections. rHuPH20 acts locally and transiently to depolymerize hyaluronan, a gel-like substance found in the subcutaneous layer of the skin. This results in decreased resistance to fluid flow and may increase dispersion and absorption of injected medicines and fluids, allowing for larger volume to be injected with limited swelling or pain. It has been shown that rHuPH20 allows for the fast absorption of a relatively large volume (10 mL) when administered SC. Shpilberg O et al., Br J Cancer. 2013; 109(6):1556-1561. Very little injection site swelling was observed when 10 mL of IgG solution was administered SC using rHuPH20 at 2,000 U/mL, whereas a large injection site swelling was observed when 10 mL of IgG solution was injected without rHuPH20. Shpilberg O et al., Br J Cancer. 2013; 109(6):1556-1561.

[0167] rHuPH20 is transiently acting and is not systematically absorbed. It has been demonstrated to exert no long-term local effects. rHuPH20 has a half-life in the skin of less than 30 minutes. Hyaluronan levels in subcutaneous tissues return to normal within 24 to 48 hours because of the rapid natural turnover of hyaluronan.

[0168] rHuPH20 is approved for SC administration in co-formulations with other active

ingredients (RITUXAN HYCELA® /MABTHERA® SC [rituximab] for Non-Hodgkin's lymphoma (NHL) and chronic lymphocytic leukemia (CLL) and HERCEPTIN HYLECTA™ / HERCEPTIN® SC [trastuzumab] in the US and Europe with an enzyme concentration of 2,000 U/mL and an injectable volume that ranges from 5 to 13.4 mL.

Methods and therapeutic uses

[0169] Provided herein are methods for treating pemphigus using an FcRn antagonist. In accordance with the invention, the FcRn antagonists are for use in treating pemphigus in a subject. In certain embodiments, the pemphigus is pemphigus vulgaris (PV). In certain embodiments, the pemphigus is pemphigus foliaceus (PF). In certain embodiments, the pemphigus can include both PV and PF. In certain embodiments, the FcRn antagonist is efgartigimod. An important goal and feature of the methods and therapeutic uses disclosed herein is the reduction or even the elimination of the use of potentially toxic agents such as corticosteroids (e.g., prednisone) and Rituxan in the treatment of pemphigus. Another important goal and feature of the methods and therapeutic uses disclosed herein is rapid onset of disease control. Yet another important goal and feature of the methods and therapeutic uses disclosed herein is achievement of long-lasting complete remission on minimal treatment, preferably without the use of potentially toxic agents such as corticosteroids (e.g., prednisone) and Rituxan.

[0170] An aspect of the disclosure is a method of treating pemphigus, comprising administering to a subject in need thereof an effective amount of a human neonatal Fc receptor (FcRn) antagonist, wherein the subject has (a) newly diagnosed pemphigus, (b) relapsing pemphigus, or (c) refractory pemphigus.

[0171] An aspect of the invention comprises an FcRn antagonist for use in the treatment of pemphigus in a subject in need thereof, wherein

1. (i) the subject has
    1. (a) newly diagnosed pemphigus,
    2. (b) relapsing pemphigus, or
    3. (c) refractory pemphigus,
2. (ii) the FcRn antagonist consists of a variant Fc region, wherein said variant Fc region consists of two Fc domains which form a homodimer, and wherein the amino acid sequence of each of the Fc domains is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, and SEQ ID NO: 3, and
3. (iii) the FcRn antagonist is administered in combination with a corticosteroid and the corticosteroid is at a dose of ≤ 0.5 mg prednisone/kg/day or equivalent.

[0172] Relapsing pemphigus refers to the appearance of at least 3 new pemphigus lesions in a

4-week period that do not heal within a week, or extension of established lesions.

[0173] Refractory pemphigus refers to pemphigus that is not controlled on current therapy. In some embodiments, refractory pemphigus refers to pemphigus that is not controlled on corticosteroids. In some embodiments, refractory pemphigus refers to pemphigus that is not controlled on rituximab. In some embodiments, refractory pemphigus refers to pemphigus that is not controlled on rituximab plus corticosteroids. In some embodiments, refractory pemphigus refers to pemphigus that is not controlled on maximum corticosteroids. In some embodiments, refractory pemphigus refers to pemphigus that is not controlled on maximum rituximab. In some embodiments, refractory pemphigus refers to pemphigus that is not controlled on maximum rituximab plus maximum corticosteroids. In certain embodiments, refractory pemphigus refers to failure to reach disease control (see below) with full therapeutic doses of systemic treatments, e.g., prednisone 1.5 mg/kg/day for 3 weeks.

[0174] In some embodiments, the pemphigus comprises pemphigus vulgaris (PV), pemphigus foliaceus (PF), or both PV and PF. In some embodiments, the pemphigus comprises pemphigus vulgaris (PV). In some embodiments, the pemphigus comprises pemphigus foliaceus (PF). In some embodiments, the pemphigus comprises both pemphigus vulgaris (PV) and pemphigus foliaceus (PF). In some embodiments, the pemphigus consists of pemphigus vulgaris (PV). In some embodiments, the pemphigus consists of pemphigus foliaceus (PF). In some embodiments, the pemphigus consists of both pemphigus vulgaris (PV) and pemphigus foliaceus (PF).

[0175] In certain embodiments, the pemphigus may be characterized as mild, mild-to-moderate, moderate, severe (extensive), or moderate-to-severe pemphigus as classified by Pemphigus Disease Area Index (PDAI). In certain embodiments, the pemphigus may be characterized as mild pemphigus (e.g., a PDAI score of < 15). In other embodiments, the pemphigus may be characterized as moderate pemphigus (e.g., a PDAI score of 15 to < 45). In other embodiments, subject to be treated has severe pemphigus (e.g., a PDAI score of ≥ 45).

[0176] In some embodiments, the FcRn antagonist is administered once weekly until disease control. As used herein, "disease control" refers to no new lesions and established lesions beginning to heal.

[0177] In some embodiments, the FcRn antagonist is administered once weekly until complete remission. As used herein, "complete remission" refers to absence of new lesions and complete healing of established lesions (except for post-inflammatory hyperpigmentation or erythema from resolving lesions).

[0178] In some embodiments, the FcRn antagonist is administered intravenously. In some embodiments, the FcRn antagonist is administered intravenously once weekly at a dose of about 10 mg/kg to about 25 mg/kg. In some embodiments, the FcRn antagonist is administered intravenously once weekly at a dose of about 10 mg/kg. In some embodiments,

the FcRn antagonist is administered intravenously once weekly at a dose of about 15 mg/kg. In some embodiments, the FcRn antagonist is administered intravenously once weekly at a dose of about 20 mg/kg. In some embodiments, the FcRn antagonist is administered intravenously once weekly at a dose of about 25 mg/kg. In some embodiments, the FcRn antagonist is administered intravenously once weekly at a dose of 10 mg/kg to 25 mg/kg. In some embodiments, the FcRn antagonist is administered intravenously once weekly at a dose of 10 mg/kg. In some embodiments, the FcRn antagonist is administered intravenously once weekly at a dose of 15 mg/kg. In some embodiments, the FcRn antagonist is administered intravenously once weekly at a dose of 20 mg/kg. In some embodiments, the FcRn antagonist is administered intravenously once weekly at a dose of 25 mg/kg.

[0179] In some embodiments, the FcRn antagonist is administered subcutaneously. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a fixed dose of about 750 mg to about 1750 mg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a fixed dose of about 750 mg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a fixed dose of about 1000 mg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a fixed dose of about 1250 mg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a fixed dose of about 1500 mg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a fixed dose of about 1750 mg.

[0180] In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a fixed dose of 750 mg to 1750 mg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a fixed dose of 750 mg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a fixed dose of 1000 mg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a fixed dose of 1250 mg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a fixed dose of 1500 mg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a fixed dose of 1750 mg.

[0181] In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a dose of about 10 mg/kg to about 25 mg/kg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a dose of about 10 mg/kg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a dose of about 15 mg/kg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a dose of about 20 mg/kg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a dose of about 25 mg/kg.

[0182] In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a dose of 10 mg/kg to 25 mg/kg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a dose of 10 mg/kg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a dose of 15 mg/kg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a dose of

20 mg/kg. In some embodiments, the FcRn antagonist is administered subcutaneously once weekly at a dose of 25 mg/kg.

[0183] In some embodiments, the FcRn antagonist is administered in an induction phase and a consolidation phase. In certain embodiments, during the induction phase, the FcRn antagonist is administered once weekly or more frequently, e.g., twice a week or every other day. In certain embodiments, during the induction phase, the FcRn antagonist is administered less frequently than once weekly, e.g., once every other week. In certain embodiments, (i) during the induction phase the FcRn antagonist is administered once weekly and corticosteroid is administered at a dose of ≤ 0.5 mg prednisone/kg/day or equivalent is administered until disease control, and (ii) during the consolidation phase the FcRn antagonist dose is decreased and/or the FcRn antagonist dosing interval is lengthened, e.g., to once biweekly, and/or the corticosteroid dose is decreased and/or the corticosteroid dosing interval is lengthened, to an end-of-consolidation dose or dosing interval effective to prevent new lesions from appearing.

[0184] In certain embodiments, (i) during the induction phase the FcRn antagonist is administered once weekly and corticosteroid is administered at a dose of ≤ 0.5 mg prednisone/kg/day or equivalent is administered until disease control, and (ii) during the consolidation phase the FcRn antagonist dose is decreased and/or the FcRn antagonist dosing interval is lengthened, e.g., to once biweekly, to an end-of-consolidation dose or dosing interval effective to prevent new lesions from appearing.

[0185] In certain embodiments, (i) during the induction phase the FcRn antagonist is administered once weekly and corticosteroid is administered at a dose of ≤ 0.5 mg prednisone/kg/day or equivalent is administered until disease control, and (ii) during the consolidation phase the corticosteroid dose is decreased and/or the corticosteroid dosing interval is lengthened, to an end-of-consolidation dose or dosing interval effective to prevent new lesions from appearing.

[0186] In certain embodiments, (i) during the induction phase the FcRn antagonist is administered once weekly and corticosteroid is administered at a dose of ≤ 0.5 mg prednisone/kg/day or equivalent is administered until disease control, and (ii) during the consolidation phase the FcRn antagonist dose is decreased and/or the FcRn antagonist dosing interval is lengthened, e.g., to once biweekly, and the corticosteroid dose is decreased and/or the corticosteroid dosing interval is lengthened, to an end-of-consolidation dose or dosing interval effective to prevent new lesions from appearing.

[0187] In some embodiments, during the induction phase the FcRn antagonist is administered intravenously at a dose of about 10 mg/kg to about 25 mg/kg. In some embodiments, during the induction phase the FcRn antagonist is administered intravenously at a dose of about 10 mg/kg. In some embodiments, during the induction phase the FcRn antagonist is administered intravenously at a dose of about 15 mg/kg. In some embodiments, during the induction phase the FcRn antagonist is administered intravenously at a dose of about 20 mg/kg. In some embodiments, during the induction phase the FcRn antagonist is administered intravenously at

a dose of about 25 mg/kg.

[0188] In some embodiments, during the induction phase the FcRn antagonist is administered intravenously at a dose of 10 mg/kg to 25 mg/kg. In some embodiments, during the induction phase the FcRn antagonist is administered intravenously at a dose of 10 mg/kg. In some embodiments, during the induction phase the FcRn antagonist is administered intravenously at a dose of 15 mg/kg. In some embodiments, during the induction phase the FcRn antagonist is administered intravenously at a dose of 20 mg/kg. In some embodiments, during the induction phase the FcRn antagonist is administered intravenously at a dose of 25 mg/kg.

[0189] In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of about 750 mg to about 1750 mg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of about 750 mg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of about 1000 mg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of about 1250 mg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of about 1500 mg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of about 1750 mg.

[0190] In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of 750 mg to 3000 mg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of 750 mg to 1750 mg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of 750 mg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of 1000 mg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of 1250 mg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of 1500 mg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of 1750 mg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a fixed dose of 2000 mg.

[0191] In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a dose of about 10 mg/kg to about 25 mg/kg. In some embodiments, the induction phase the FcRn antagonist is administered subcutaneously at a dose of about 10 mg/kg. In some embodiments, the induction phase the FcRn antagonist is administered subcutaneously at a dose of about 15 mg/kg. In some embodiments, the induction phase the FcRn antagonist is administered subcutaneously at a dose of about 20 mg/kg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a dose of about 25 mg/kg.

[0192] In some embodiments, during the induction phase the FcRn antagonist is administered

subcutaneously at a dose of 10 mg/kg to 25 mg/kg. In some embodiments, the induction phase the FcRn antagonist is administered subcutaneously at a dose of 10 mg/kg. In some embodiments, the induction phase the FcRn antagonist is administered subcutaneously at a dose of 15 mg/kg. In some embodiments, the induction phase the FcRn antagonist is administered subcutaneously at a dose of 20 mg/kg. In some embodiments, during the induction phase the FcRn antagonist is administered subcutaneously at a dose of 25 mg/kg.

[0193] In some embodiments, during the consolidation phase the FcRn antagonist dosing interval is once weekly or less frequently. For example, in various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or 21 days; or once every 2, 3, 4, 5 or 6 weeks. In various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 7 to 8, 7 to 9, 7 to 10, 7 to 11, 7 to 12, 7 to 13, 7 to 14, 7 to 15, 7 to 16, 7 to 17, 7 to 18, 7 to 19, 7 to 20, or 7 to 21 days. In various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 8 to 9, 8 to 10, 8 to 11, 8 to 12, 8 to 13, 8 to 14, 8 to 15, 8 to 16, 8 to 17, 8 to 18, 8 to 19, 8 to 20, or 8 to 21 days. In various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 9 to 10, 9 to 11, 9 to 12, 9 to 13, 9 to 14, 9 to 15, 9 to 16, 9 to 17, 9 to 18, 9 to 19, 9 to 20, or 9 to 21 days. In various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 10 to 11, 10 to 12, 10 to 13, 10 to 14, 10 to 15, 10 to 16, 10 to 17, 10 to 18, 10 to 19, 10 to 20, or 10 to 21 days. In various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 11 to 12, 11 to 13, 11 to 14, 11 to 15, 11 to 16, 11 to 17, 11 to 18, 11 to 19, 11 to 20, or 11 to 21 days. In various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 12 to 13, 12 to 14, 12 to 15, 12 to 16, 12 to 17, 12 to 18, 12 to 19, 12 to 20, or 12 to 21 days. In various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 13 to 14, 13 to 15, 13 to 16, 13 to 17, 13 to 18, 13 to 19, 13 to 20, or 13 to 21 days. In various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 14 to 15, 14 to 16, 14 to 17, 14 to 18, 14 to 19, 14 to 20, or 14 to 21 days. In various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 15 to 16, 15 to 17, 15 to 18, 15 to 19, 15 to 20, or 15 to 21 days. In various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 16 to 17, 16 to 18, 16 to 19, 16 to 20, or 16 to 21 days. In various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 17 to 18, 17 to 19, 17 to 20, or 17 to 21 days. In various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 18 to 19, 18 to 20, or 18 to 21 days. In various certain embodiments, during the consolidation phase the FcRn antagonist dosing interval is once every 19 to 20, or 19 to 21 days.

[0194] In some embodiments, during the consolidation phase the FcRn antagonist dosing interval is once weekly or every 2 weeks.

[0195] In some embodiments, the FcRn antagonist for use in the treatment of pemphigus

further comprises a maintenance phase, wherein
(iii) during the maintenance phase the end-of-consolidation dose or dosing interval for the FcRn antagonist and/or the prednisone is continued until complete clearance of lesions.

[0196] In some embodiments, during the maintenance phase the FcRn antagonist dosing interval is once weekly or less frequently. For example, in various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or 21 days; or once every 2, 3, 4, 5, or 6 weeks. In various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 7 to 8, 7 to 9, 7 to 10, 7 to 11, 7 to 12, 7 to 13, 7 to 14, 7 to 15, 7 to 16, 7 to 17, 7 to 18, 7 to 19, 7 to 20, or 7 to 21 days. In various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 8 to 9, 8 to 10, 8 to 11, 8 to 12, 8 to 13, 8 to 14, 8 to 15, 8 to 16, 8 to 17, 8 to 18, 8 to 19, 8 to 20, or 8 to 21 days. In various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 9 to 10, 9 to 11, 9 to 12, 9 to 13, 9 to 14, 9 to 15, 9 to 16, 9 to 17, 9 to 18, 9 to 19, 9 to 20, or 9 to 21 days. In various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 10 to 11, 10 to 12, 10 to 13, 10 to 14, 10 to 15, 10 to 16, 10 to 17, 10 to 18, 10 to 19, 10 to 20, or 10 to 21 days. In various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 11 to 12, 11 to 13, 11 to 14, 11 to 15, 11 to 16, 11 to 17, 11 to 18, 11 to 19, 11 to 20, or 11 to 21 days. In various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 12 to 13, 12 to 14, 12 to 15, 12 to 16, 12 to 17, 12 to 18, 12 to 19, 12 to 20, or 12 to 21 days. In various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 13 to 14, 13 to 15, 13 to 16, 13 to 17, 13 to 18, 13 to 19, 13 to 20, or 13 to 21 days. In various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 14 to 15, 14 to 16, 14 to 17, 14 to 18, 14 to 19, 14 to 20, or 14 to 21 days. In various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 15 to 16, 15 to 17, 15 to 18, 15 to 19, 15 to 20, or 15 to 21 days. In various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 16 to 17, 16 to 18, 16 to 19, 16 to 20, or 16 to 21 days. In various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 17 to 18, 17 to 19, 17 to 20, or 17 to 21 days. In various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 18 to 19, 18 to 20, or 18 to 21 days. In various certain embodiments, during the maintenance phase the FcRn antagonist dosing interval is once every 19 to 20, or 19 to 21 days.

[0197] In some embodiments, during the maintenance phase the FcRn antagonist dosing interval is once weekly. In some embodiments, during the maintenance phase the FcRn antagonist dosing interval is once biweekly.

[0198] In some embodiments, the pemphigus goes into complete remission. Complete remission as described herein can be achieved at a corticosteroid dose of ≤ about 5 mg prednisone/kg/day or equivalent. Complete remission as described herein can be achieved at

a corticosteroid dose of ≤ about 3 mg prednisone/kg/day or equivalent. Complete remission as described herein can be achieved at a corticosteroid dose of ≤ about 2 mg prednisone/kg/day or equivalent. Complete remission as described herein can be achieved at a corticosteroid dose of ≤ about 1 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 0.5 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 0.4 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 0.3 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 0.2 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 0.1 mg prednisone/kg/day or equivalent.

[0199] Complete remission as described herein can be achieved at a corticosteroid dose of ≤ 5 mg prednisone/kg/day or equivalent. Complete remission as described herein can be achieved at a corticosteroid dose of ≤ 3 mg prednisone/kg/day or equivalent. Complete remission as described herein can be achieved at a corticosteroid dose of ≤ 2 mg prednisone/kg/day or equivalent. Complete remission as described herein can be achieved at a corticosteroid dose of ≤ 1 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ 0.5 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ 0.4 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ 0.3 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ 0.2 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ 0.1 mg prednisone/kg/day or equivalent.

[0200] In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 20 mg prednisone/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 15 mg prednisone/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 10 mg prednisone/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 5 mg prednisone/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 3 mg prednisone/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ about 2 mg prednisone/day or equivalent.

[0201] In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ 10 mg prednisone/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ 5 mg prednisone/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ 3 mg prednisone/day or equivalent. In some embodiments, the complete remission is achieved at a corticosteroid dose of ≤ 2 mg prednisone/day or equivalent.

[0202] Complete remission as described herein can be achieved without corticosteroid.

[0203] In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 20 mg prednisone/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 15 mg prednisone/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 10 mg prednisone/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 5 mg prednisone/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 2 mg prednisone/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 1 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 0.5 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 0.4 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 0.3 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 0.2 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ about 0.1 mg prednisone/kg/day or equivalent.

[0204] In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ 20 mg prednisone/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ 15 mg prednisone/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ 10 mg prednisone/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ 5 mg prednisone/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ 2 mg prednisone/day or equivalent. Complete remission as described herein can be maintained with a corticosteroid dose of ≤ 1 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ 0.5 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ 0.4 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ 0.3 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ 0.2 mg prednisone/kg/day or equivalent. In some embodiments, the complete remission is maintained with a corticosteroid dose of ≤ 0.1 mg prednisone/kg/day or equivalent.

[0205] In some embodiments, the complete remission is maintained without corticosteroid.

[0206] In some embodiments, the subject is rituximab-refractory.

[0207] In some embodiments, the subject is corticosteroid-intolerant.

EXAMPLES

[0208] The invention will be further understood with reference to the following non-limiting experimental examples.

Example 1 - Overview of a Phase 2 Clinical Trial of Efgartigimod in Treatment of Pemphigus Patients

[0209] This example describes (ARGX-113-1701) a completed open-label, non-controlled, adaptive-design Phase 2 trial to evaluate the safety, PD, PK, efficacy, and conditions of use (dosage, frequency of administration at maintenance) of efgartigimod (variant Fc fragment of human IgG1 capable of blocking the neonatal Fc receptor (FcRn) for the treatment of pemphigus (vulgaris or foliaceus).

[0210] As described in more detail below, thirty-four (34) patients suffering from mild to moderate pemphigus vulgaris or foliaceus were enrolled into the trial. In sequential cohorts efgartigimod was dosed at 10 mg/kg or 25 mg/kg intravenously with various dosing frequency, as monotherapy or as add-on to low-dose oral prednisone.

[0211] The study enrolled adult male or female patients with clinical diagnosis of PV or PF, that has been confirmed by positive direct immunofluorescence, and positive indirect immunofluorescence and/or enzyme-linked immunosorbent assay (ELISA) with mild to moderate disease severity (Pemphigus Disease Area Index [PDAI] <45). The serum levels of autoantibodies directed against Dsg-3 and/or Dsg-1 antigen were identified at screening. Patients were newly diagnosed patients or relapsing patients off therapy or patients who relapsed despite taking oral prednisone at tapered dose with or without a conventional immunosuppressant. Treatment during the study consisted of efgartigimod monotherapy or efgartigimod in combination with oral prednisone or equivalent.

[0212] Efgartigimod demonstrated a favorable safety and tolerability profile in pemphigus patients, consistent with previous studies of efgartigimod. A strong association was shown between serum IgG level reduction, autoantibody level reduction and improvement of pemphigus disease area index (PDAI) scores and clinical outcomes. Efgartigimod, either as monotherapy or combined with prednisone, demonstrated fast onset of action and resulted in disease control in 90% of patients with a median time of 16 days. Optimized prolonged treatment with efgartigimod with low dose corticosteroids at a dose range of 0.06 to 0.48 mg/kg/day led to complete clinical remission within 2 to 41 weeks in 59% of patients.

[0213] The results of the trial indicated that efgartigimod was safe and induced a fast onset of action on disease activity and clinical outcomes, overall supporting efgartigimod as a promising therapy for pemphigus.

Example 2: Study Design

[0214] PV and PF patients with mild to moderate disease severity (PDAI <45 at baseline) (Rosenbach, M, et al., J Invest Dermatol, 2009. 129(10): 2404-10) were enrolled in sixteen study centers in Europe and Israel. PV or PF diagnosis had to be confirmed by positive direct immunofluorescence and positive indirect immunofluorescence and/or Dsg-1/3 ELISA. Patients were either newly diagnosed, or relapsing. Patients having a treatment course of oral prednisone (or equivalent) and immunosuppressant at screening were includable, but the immunosuppressant had to be discontinued before baseline. Patients with a history of refractory disease to a second line therapy (e.g. IVIg, rituximab, plasma exchange/immunoadsorption) were excluded. In addition, use of therapies other than oral prednisone and conventional immunosuppressants, that could interfere with the clinical course of the disease (e.g. IV prednisolone bolus, dapsone, sulfasalazine, tetracyclines, nicotinamide, plasmapheresis/plasma exchange, immunoadsorption and IVIg) within 2 months prior to baseline visit were not permitted, neither the use of rituximab and other CD20-targeting biologics within 6 months prior to baseline visit. Patients with total IgG level <6 g/L in serum at screening were also excluded.

[0215] An open-label, non-controlled trial was conducted using an adaptive design with 4 sequential cohorts, in which efgartigimod, administered IV over a period of two hours, was tested at two different dosages (10 and 25 mg/kg body weight). At least 4 evaluable patients were to be enrolled in each cohort 1 to 3, and at least 10 evaluable patients in cohort 4. **Figure 1** presents a schematic of the design of this study. No pre-medication was required prior to efgartigimod infusion. Variable durations of the induction phase as defined by weekly infusions, and variable durations of the maintenance phase with a variable frequency of infusions (biweekly or four-weekly) were tested.

[0216] Overall, the trial included a 2-week screening period, a treatment period of 9 (cohort 1) to 34 weeks (cohort 4) as specified for each cohort, and a treatment-free follow-up period (8 weeks for cohort 1 and 10 weeks for cohorts 2 to 4). Before each administration, the total IgG serum level from the previous visit was verified, and no efgartigimod treatment was permitted if total IgG was < 1.2 g/L, until recovery above this level. An Independent Data Monitoring Committee (IDMC) provided recommendations on the changes in treatment regimen between cohorts, based on safety and efficacy data from the previous cohorts. A minimum of 4 evaluable patients was included in each cohort.

[0217] During the treatment period, eligible patients were to receive efgartigimod via IV infusions and dosing schedule as follows. Cohort 1 patients had a treatment period of 9 weeks by receiving 4 weekly 10 mg/kg infusions and two 10 mg/kg maintenance infusions with a 2- and 4-week interval, respectively. Cohort 2 patients had a treatment period of 11 weeks and received 4 weekly 10 mg/kg infusions and 4 biweekly (i.e., dosing every other week) 10 mg/kg maintenance infusions. Cohort 3 patients had a treatment period of 15 weeks and received 4 weekly 10 mg/kg infusions and 6 biweekly 10 mg/kg maintenance infusions. Patients in cohort 4 receive weekly 25 mg/kg infusions until reaching end of consolidation (EoC), with a minimum

of 5 weekly infusions, followed by biweekly 25 mg/kg maintenance infusions until the end of treatment at week 34. EoC is defined as the time at which no new lesions have developed for a minimum of 2 weeks and the majority (approximately 80%) of established lesions has healed.

[0218] In cohorts 1-3, efgartigimod was used from baseline either as monotherapy (in newly diagnosed patients and relapsing patients off therapy) or as add-on treatment to prednisone (in newly diagnosed patients on a first stable course of prednisone and in patients relapsing at a tapered prednisone dose). Under the latter circumstances, newly diagnosed patients received prednisone 20 mg/day and relapsing patients continued prednisone at the same tapered dose. In cohort 4, efgartigimod treatment was systematically initiated in association with prednisone, i.e., 20 mg/day in all newly diagnosed patients and relapsing patients off therapy, or at the tapered dose at which relapse occurred. In cohort 4, the oral prednisone dose could be tapered as of end of consolidation (EoC). Conversely, prednisone could be increased to 40 mg/day or at higher tapered doses according to clinical judgment, before study withdrawal in case of disease progression. No other systemic treatments for pemphigus were permitted during the study, whereas topical corticosteroids, analgesics, and supportive care for corticosteroid therapy (e.g., vitamin D, proton-pump inhibitors, specific diets) were allowed.

[0219] Descriptive statistical methods were used to analyze safety and efficacy data. Summaries were provided by cohort and/or efgartigimod dose.

[0220] As mentioned above, oral prednisone was the only treatment that was allowed in the study as concomitant treatment and rescue therapy. It consisted of addition of 20 mg/day of prednisone to efgartigimod treatment in newly diagnosed and off-therapy patients, and of an increase to the previous prednisone dose in relapsing patients. Patients under rescue treatment were kept in the study, unless disease progression led the patients to discontinue the study.

Example 3. Safety and Efficacy Assessment

[0221] The primary endpoint was safety, assessed throughout the course of the study, including the frequency and severity of treatment-emergent adverse events (TEAEs), serious adverse events (SAEs), vital signs, ECG (electrocardiogram) parameters, physical examination abnormalities and routine clinical laboratory assessments (hematology, biochemistry, urinalysis). As additional safety parameter, total IgG levels in serum were measured at each visit.

[0222] Efficacy endpoints included pemphigus disease activity index (PDAI) assessment; time to DC, defined as no new lesions, established lesions starting to heal; time until relapse, defined as the appearance of 3 or more new lesions a month that do not heal spontaneously within 1 week, or as the extension of established lesions, evaluated from any visit following DC; time to EoC, defined as the time at which no new lesions have developed for a minimum of 2 weeks, and approximately 80% of lesions have healed; time to complete CR, defined by the

absence of new lesions and established lesions completely healed (except for post-inflammatory hyperpigmentation or erythema from resolving lesions); and time to complete clinical remission under minimal therapy (CRmin), defined by a prednisone dose of 10 mg/day or less for at least 8 weeks.

[0223] Other secondary endpoints included the evaluation of the pharmacodynamic (total IgG and subtypes, anti-Dsg-1 and -3 autoantibodies) and pharmacokinetics parameters., and immunogenicity (incidence of anti-drug antibodies (ADA)). Measurement of serum levels of anti-Dsg-1 and anti-Dsg-3 autoantibodies was performed with anti-Desmoglein-1 and -3 IgG ELISA test kits (Euroimmun, Germany), respectively.

Example 4. Pemphigus Study Population and Patient Disposition

[0224] Fifty-three (53) patients were screened for eligibility, 35 were eligible to participate, and 34 were enrolled in the trial (one patient withdrew consent before baseline), which constituted the safety analysis set **(Figure 2).** Twelve study centers in 5 countries (Germany, Hungary, Israel, Italy, Ukraine) treated at least 1 patient for the study. Six patients were assigned to cohort 1, 5 patients were assigned to cohort 2, 8 patients were assigned to cohort 3, and 15 patients were assigned to cohort 4.

[0225] Results in this study include those from a first interim analysis (data cutoff November 7, 2019, report date February 21, 2020), those from a second (later) interim analysis (data cutoff March 25, 2020, report date July 30, 2020), a third (later) interim analysis (data cutoff June 24, 2020), and a final analysis (data cutoff October 28, 2020). Unless indicated otherwise, data presented in the Examples are from the final analysis.

Example 5. Interim Analyses

[0226] As summarized in **Figure 3,** an initial Interim analysis of results from this Phase 2 study (data cutoff Nov 7, 2019) indicated the following:

1. (i) 23 patients enrolled in adaptive trial evaluating 10mg/kg and 25mg/kg IV efgartigimod with multiple dosing regimens established a clear correlation between pathogenic IgG reduction and Pemphigus Disease Area Index score improvement;
2. (ii) 83% of patients overall achieved disease control, with 78% within four weeks;
3. (iii) Rapid disease control was associated with both monotherapy and combination with low-dose prednisone;
4. (iv) 70% (5/7) of patients who received optimized dosing of efgartigimod with concomitant corticosteroids achieved clinical remission (CR); five CRs were observed within 2-10 weeks; and 39% of all patients on combination therapy achieved CR; and
5. (v) Demonstrated potential to taper patients off concomitant corticosteroids after achieving CR.

A. Demographics and Baseline Characteristics

[0227] Interim patient demographics and baseline characteristics by analysis population are presented in Table 2A.

Table 2A. Demographics and Baseline Characteristics - Safety Analysis Set - Data from initial interim analysis (cut-off date on the data from the Phase 2 trial is: 7 Nov 2019)

Parameter	**Safety Analysis Set Locked Data**	**Safety Analysis Set All Available Data**	
	(N=16)	(N=26)	
	n (%)	n (%)	
Age at Screening (years)			
	n	16	26
	mean (std)	52.8 (14.21)	49.7 (14.58)
	median	51.0	49.0
	range (min, max)	29, 78	22, 78
Gender			
	Male	4 (25.0)	9 (34.6)
	Female	12 (75.0)	17 (65.4)
Pemphigus Vulgaris		15 (93.8)	21 (80.8)
	Mucosal-dominant	6 (40.0)	8 (38.1)
	Mucocutaneous	8 (53.3)	12 (57.1)
	Cutaneous	1 (6.7)	1 (4.8)
Pemphigus Foliaceus		1 (6.3)	5 (19.2)
Baseline PDAI severity			
Mild (PDAI <15)		5 (31.3)	9 (34.6)
Moderate (PDAI 15-44)		11 (68.8)	17 (65.4)
Newly diagnosed		6 (37.5)	11 (42.3)
Relapsing		10 (62.5)	15 (57.7)
ARGX-113 monotherapy at baseline		11 (68.8)	12 (46.2)
		5 (31.3)	14 (53.8)
Patient took prednisone at baseline			

[0228] Patient demographics and baseline characteristics for a second interim analysis are

presented in **Table 2B.** As of the cut-off date of June 24, 2020, 26 patients with PV and 8 patients with PF were enrolled, of which 22 were females and 12 males. PV subtypes included mucosal-dominant (n=9), mucocutaneous (n=14) and cutaneous (n=3). 12 patients with mild baseline PDAI (<15) and 22 with moderate baseline PDAI (15-44) entered the study. Newly diagnosed patients comprised 14 patients and relapsing patients accounted for 20 patients. 11 patients began the trial with efgartigimod monotherapy while 23 started in association with low-dose prednisone at baseline.

Table 2B. Demographics and Baseline Characteristics - Safety and Efficacy Analysis Sets - Data from final analysis (cut-off date on the data from the Phase 2 trial is: 24 Jun 2020)

Baseline characteristics		**Safety Analysis Set (N=34)**	**Efficacy Analysis Set (N=31)**
Age (mean ± SD)		51.5 ± 15.3	52.4 ± 15.5
Sex (n (%))			
	• Male	12 (35)	10 (32)
	• Female	22 (65)	21 (68)
Pemphigus vulgaris (n (%))		26 (77)	24 (77)
	• Mucosal-dominant	9 (35)	9 (38)
	• Mucocutaneous	14 (54)	12 (50)
	• Cutaneous	3 (11)	3 (12)
Pemphigus foliaceus (n (%))		8 (24)	7 (23)
Disease History (n (%))			
	• Newly diagnosed	14 (41)	12 (39)
	• Relapsing	20 (59)	19 (61)
Baseline PDAI severity (n (%))			
	• Mild (PDAI < 15)	12 (35)	12 (39)
	• Moderate (PDAI 15-44)	22 (65)	19 (61)
Baseline PDAI score (mean ± SD) (min, median, max score)			
	• Overall population	20.9 ± 11.5 (2.0, 20.4, 39.9)	20.1 ± 11.8 (2.0, 19.0, 39.9)
Treatment initiated at Baseline (n (%))			
	• Efgartigimod monotherapy	11 (32)	8 (26)
	• Efgartigimod + CS	23 (68)	23 (74)

B. Exposure

[0229] Extent of exposure (number of days of exposure to study drug and number of injections received by each patient) are presented by efgartigimod dose administered and overall for the Safety Analysis Set in **Table 3.**

[0230] In cohorts 1 to 3 patients received a dose of 10 mg/kg and in cohort 4 patients received a dose of 25 mg/kg.

Table 3. Trial ARGX-113-1701 Exposure - Safety Analysis Set - Data from initial interim analysis (cut-off date on the data from the Phase 2 trial is: 7 Nov 2019)

	Cohort 1 (10mg/kg) (N=6)	**Cohort 2 (10mg/kg) (N=5)**	**Cohort 3 (10mg/kg) (N=8)**	**Cohort 4 (25mg/kg) (N=7)**
Overall Exposure to study drug per patient (days)				
n	6	5	8	7
mean (SD)	38.3 (28.63)	54.8 (32.75)	100.1 (18.69)	84.9 (61.43)
median	39.0	78.0	106.0	85.0
range (min, max)	8, 66	16, 79	54, 110	8, 184
Overall Exposure to study drug per patient (mg)				
n	6	5	8	7
mean (SD)	2939.33	3979.20	6489.38	16213.16
median	(1616.196)	(1196.764)	(2274.694)	(9216.324)
range (min, max)	3025.00 1056.0, 5000.0	3456.00 2880.0, 5600.0	7132.50 2500.0, 9030.0	13704.50 3100.0, 28800.0
Overall Number of Infusions received per patient				
n	6	5	8	7
mean (SD)	4.0 (1.55)	6.2 (2.49)	9.3 (1.75)	9.3 (5.22)
median	4.0	8.0	10.0	10.0
range (min, max)	2, 6	3, 8	5, 10	2, 18
n=number of patients; SD=standard deviation				

C. Clinical Efficacy

[0231] Analyses of efficacy in this study were secondary objectives. All efficacy analyses were carried out using the efficacy analysis set. Efficacy assessments included an assessment of the extent of disease and evaluation of consensual clinical endpoints.

D. PDAI Activity Score

[0232] The change from baseline of the PDAI activity score at the end of each treatment phase (induction, maintenance, treatment-free follow up) for cohorts 1-3 is presented in **Table 4.** The changes from baseline of the PDAI activity at each visit by cohort is illustrated in **Figure 4** depicting the clinical efficacy assessed by the pemphigus disease area index (PDAI) activity in all 4 cohorts. Most patients demonstrated strong PDAI score improvements: In cohort 1 there was a correlation in IgG reduction with anti-Dsg reduction with PDAI score improvement, early DC (mono/combo) and suboptimal efgartigimod dosing in maintenance; in cohort 2 there was improved maintenance with efgartigimod dosing every other week. In cohort 3 all patients ultimately associated with prednisone, maintenance further improved symptoms, and EoC/CR was noted when associated with oral prednisone. In cohort 4, strong PDAI score improvements were noted, there was a high rate of EoC, and CR when prednisone was not tapered before CR.

[0233] **Figure 5** depicts clinical efficacy as assessed by PDAI activity in moderate severity patient cohorts 1-4. Most patients demonstrated strong PDAI score improvements: cohort 1 comprised three moderate patients (3 PV, 0 PF); cohort 2 comprised two moderate patients (2 PV, 0 PF); cohort 3 comprised four moderate patients (4 PV, 0 PF); and cohort 4 comprised five moderate patients (1 PV, 4 PF). The proposed patient population for the pemphigus Phase 3 trial is moderate to severe PV/PF.

[0234] **Figure 6** depicts PDAI activity scores over time in A) cohort 1, B) cohort 2, C) cohort 3, and D) cohort 4, as determined in the final analysis of the study.

Table 4. PDAI Activity Score Reduction: Percent Change from Baseline in Cohorts 1-3 - Efficacy Analysis Set - Data from initial interim analysis (cut-off date on the data from the Phase 2 trial is: 7 Nov 2019)

		Cohort 1 (10 mg/kg)	**Cohort 2 (10 mg/kg)**	**Cohort 3 (10 mg/kg)**
End of induction phase*1	N	3	3	7
	mean (std)	-53.77 (18.691)	88.20 (280.274)	-58.26 (65.130)
	median	-46.50	-54.40	-83.30
	range (min, max)	-75.0, -39.8	-92.1, 411.1	-100.0, 80.0

	95% CI	-100.2, -7.3	-608.0, 784.4	-118.5, 2.04
End of maintenance phase*2	N	3	3	7
	mean (std)	-23.83 (30.534)	-56.20 (43.139)	-15.49 (184.246)
	median	-25.50	-69.80	-91.70
	range (min, max)	-53.5, 7.5	-90.9, -7.9	-100.0, 400.0
	95% CI	-99.7, 52.0	-163.4, 51.0	-185.9, 154.94
End of treatment-free follow-up*3	N	3	2	6
	mean (std)	9.90 (80.158)	-76.30 (8.910)	-33.05 (130.366)
	median	-16.80	-76.30	-89.15
	range (min, max)	-53.5, 100.0	-82.6, -70.0	-100.0, 230.0
	95% CI	-189.2, 209.0	-156.3, 3.7	-169.9, 103.84

*1 End of induction phase cohorts 1 to 3 (visit 5)
*2 End of maintenance phase*2 (cohort 1 - M2; cohort 2-M4; cohort 3-M6)
*3 End of treatment-free follow-up (cohort 1 - FU2; cohorts 2 and 3 - FU3)
CI=confidence interval

[0235] Within the induction period comprising 4 weekly infusions (cohorts 1, 2 and 3), PDAI activity was reduced, ranging from median change -46.5% (range -39.8%, -75%) in cohort 1 to median change -83.3% (range -100%, +80.0%) in cohort 3 after 4 weeks. At the end of the maintenance period, the PDAI activity in cohort 1 (2 infusions with a 2-week then a 4-week interval) tended to be higher than at the end of induction (median change -25.5% at the end of maintenance and median change -46.5% at the end of induction). In cohort 2 (4 infusions every other week), the PDAI activity was decreased to a greater extent at the end of maintenance compared to the end of induction (median change -69.8% at the end of maintenance compared to median change -54.4% at the end of induction). In cohort 3 (6 infusions every other week), PDAI activity was further decreased compared to the preceding cohorts, with a median change of -91.7% at the end of maintenance (compared to end of induction median change -83.3%).

[0236] During the treatment-free follow up, the PDAI activity in cohort 1 tended to return to its baseline value (median change -16.8%), whereas in cohorts 2 and 3 the PDAI activity remained stable at a similar level compared to the end of maintenance (median change -76.30% and -89.2%, respectively).

[0237] In cohort 4, the treatment regimen consisting of an induction treatment period with weekly administration until end of consolidation and maintenance treatment period with administration every other week until week 34 was different as compared to cohorts 1-3.

Accordingly, no direct comparison of the periods with cohorts1-3 can be made. The time course of PDAI activity in cohort 4 is presented in **Figure 4D,** showing a median change of PDAI activity of -73.7% (range -51.5%, -80.3%) after 4 weeks.

E. Clinical Outcomes

[0238] Clinical outcomes include assessment of disease control (DC), complete clinical remission (CR), and relapse.

E.1. Disease Control

[0239] In the efficacy analysis population (n=31), 28 patients (90.3%) reached DC during the study and 24 patients (77.4%) had DC within the 4-week induction period (**Table 5**). DC distributed equally between patients of mild and moderate severity, newly diagnosed and relapsing, and under monotherapy and concomitant prednisone. The median time to DC was as early as 15.0 days (range: 8-30 days) for patients in cohorts 1 to 3 and 29 days (range 9-30 days) in cohort 4. Conversely, 2 patients in cohort 2 experienced disease progression, which resulted in early discontinuation in both cases.

Table 5. Trial ARGX-113-1701 – Incidence of Disease Control (Overall and During 4-weeks Induction Period) and Time to Disease Control – Efficacy Analysis Set – Data from initial interim analysis (cut-off date on the data from the Phase 2 trial is: 7 Nov 2019)

A

Subgroup	**Cohort 1 (10 mg/kg) n (%)**	**Cohort 2 (10 mg/kg) n (%)**	**Cohort 3 (10 mg/kg) n (%)**	**Cohort 4 (25 mg/kg) n (%)**	**Overall n (%)**
Overall	4	5	7	7	23
Achieved DC	4 (100.0)	3 (60.0)	7 (100.0)	5 (71.4)	19 (82.6)
No DC	0	2 (40.0)	0	2 (28.6)	4 (17.4)
Disease Severity at baseline					
Mild (PDAI <15)	1 (25.0)	2 (40.0)	3 (42.9)	2 (28.6)	8 (34.8)
Moderate (PDAI 15-44)	3 (75.0)	1 (20.0)	4 (57.1)	3 (42.9)	11 (47.8)
Disease history					
Relapsing Patients	4 (100.0)	1 (20.0)	5 (71.4)	3 (42.9)	13 (56.5)
Newly Diagnosed Patients	0	2 (40.0)	2 (28.6)	2 (28.6)	6 (26.1)
ARGX-113					

monotherapy at baseline	3 (75.0)	3 (60.0)	2 (28.6)	0	8 (34.8)
Patient took Prednisone at baseline	1 (25.0)	0	5 (71.4)	5 (71.4)	11 (47.8)

B

	Cohort 1 (10 mg/kg) n (%)	**Cohort 2 (10 mg/kg) n (%)**	**Cohort 3 (10 mg/kg) n (%)**	**Cohort 4 (25 mg/kg) n (%)**	**Overall n (%)**
Disease control during 4-weeks Induction period	4	5	7	7	23
Achieved DC	4 (100.0)	2 (40.0)	7 (100.0)	5 (71.4)	18 (78.3)
No DC	0	3 (60.0)	0	2 (28.6)	5 (21.7)

C

Summary Statistics for Time Variable Days to Control (days)	**Cohort 1 to 3 (10 mg/kg)**	**Cohort 4 (25 mg/kg)**
Median time to DC	15.0	22.5
Range	(8-92)	(15-30)

n=number of patients

E.2. Complete Clinical Remission (CR)

[0240] CR was assessed in cohorts 3 and 4. In cohort 3, CR was based on a PDAI activity score of 0, in cohort 4 CR was assessed by the investigator. CR was observed in cohorts 3 and 4 where efgartigimod treatment was prolonged (11 weeks in cohort 3, up to 34 weeks in cohort 4) compared to previous cohorts, and prednisone was associated in all patients. In these two cohorts, 7 patients achieved CR: 5 patients (71.4%) in cohort 3, and 7 (46.7%) patients in cohort 4 **(Table 6)**. Notably, 3 patients with a moderate disease severity reached CR in cohort 3.

[0241] Median time to CR was 36 days in cohort 3 (range: 13-93 days). In cohort 4, time to CR for the two patients was 44 and 72 days. The prednisone dose at the time of CR was found to be a contributing factor for achieving CR. Indeed, all patients reaching CR received a daily prednisone dose ranging from 0.06 to 0.48 mg/kg (median cohort 3: 0.27 mg/kg/day, median cohort 4: 0.28 mg/kg/day). Consequently, the combination of efgartigimod and a prednisone dose as low as 0.25-0.50 mg/kg per day may be sufficient for achieving CR.

Table 6. Trial ARGX-113-1701 - Incidence of CR and Concomitant Prednisone Equivalent Dose at the Time of CR - Cohorts 3 and 4 - Efficacy Analysis Set - Data from initial interim

analysis (cut-off date on the data from the Phase 2 trial is: 7 Nov 2019)

A			
Subgroup		**Cohort 3 (10 mg/kg) n(%)**	**Cohort 4 (25 mg/kg) n(%)**
Overall		7	7
	Achieved CR	5 (71.4)	2 (28.6)
	No Clinical Remission	2 (28.6)	5 (71.4)
Disease Severity at baseline			
	Mild (PDAI < 15)	2 (28.6)	2 (28.6)
	Moderate (PDAI 15-44)	3 (42.9)	0
Disease history			
	Relapsing Patients	3 (42.9)	1 (14.3)
	Newly Diagnosed Patients	2 (28.6)	1 (14.3)
B			
Parameter		**Cohort 3 (10mg/kg) (N=7)**	**Cohort 4 (25mg/kg) (N=7)**
Prednisone Equivalent Dose (mg/day) on day before CR			
	n	5	2
	mean (std)	17.00 (6.71)	20.00 (0.00)
	median	20.00	20.00
	range (min, max)	5.00, 20.00	20.00, 20.00
Prednisone Equivalent Dose (mg/kg/day) on day before CR			
N		5	2
	mean (std)	0.26 (0.15)	0.34 (0.04)
	median	0.27	0.340.31, 0.36
	range (min, max)	0.06, 0.48	
n=number of patients			

E.3. Relapse

[0242] Relapse is assessed in patients who reach DC (N=19) and was reported in 8 patients (42.1%; Table 7). It occurred more frequently early after DC (N=7) than later after CR (N=1). For patients in cohorts 1 to 3, median time to relapse after DC was 141 days (N=7; range 10-

169) and 1 patient in cohort 4 had a relapse 63 days after DC. At the time of relapse, 3 patients were treated without concomitant prednisone, and 5 patients were on prednisone treatment at the mean dose of 0.14±0.06 mg/kg per day (median 0.17 mg/kg/day; range 0.07-0.20 mg/kg/day).

[0243] Relapse, which may occur immediately after DC is reached, relapse was actually only observed under bi-weekly efgartigimod regimen (N=4) or during the efgartigimod treatment-free follow up (N=4), whereas no patient on weekly efgartigimod administration had a relapse. According to these findings, it can be hypothesized that a weekly dosage regimen of efgartigimod is optimal for preventing relapse.

Table 7. Trial ARGX-113-1701 – Incidence of Relapse and Concomitant Prednisone Equivalent Dose at the Time of Relapse – Efficacy Analysis Set – Data from initial interim analysis (cut-off date on the data from the Phase 2 trial is: 7 Nov 2019)

A

Subgroup	**Cohort 1 (10 mg/kg) n (%)**	**Cohort 2 (10 mg/kg) n (%)**	**Cohort 3 (10 mg/kg) n (%)**	**Cohort 4 (25 mg/kg) n (%)**	**Overall n (%)**
Overall	4	3	7	5	19
Patient with relapse	2 (50.0)	2 (66.7)	3 (42.9)	1 (20.0)	8 (42.1)
No relapse	2 (50.0)	1 (33.3)	4 (57.1)	4 (80.0)	11 (57.9)
Disease Severity at baseline					
Mild (PDAI < 15)	1 (25.0)	2 (66.7)	1 (14.3)	0	4 (21.1)
Moderate (PDAI 15-44)	1 (25.0)	0	2 (28.6)	1 (20.0)	4 (21.1)
Disease history					
Relapsing Patients	2 (50.0)	0	3 (42.9)	1 (20.0)	6 (31.6)
Newly Diagnosed Patients	0	2 (66.7)	0	0	2 (10.5)

B

Parameter	**Cohort 1-3 (10mg/kg) (N=14)**	**Cohort 4 (25mg/kg) (N=7)**
Prednisone Equivalent Dose (mg/day) on day before Relapse		
n	7	1
mean (std)	5.86 (6.96)	10.00 (NA)
median	5.00	10.00
range (min, max)	0.00, 16.00	10.00, 10.00
Prednisone Equivalent Dose (mg/kg/day) on day before Relapse		

N	7	2
mean (std)	0.07 (0.08)	0.46 (0.13)
median	0.07	0.46
range (min, max)	0.00, 0.18	0.36, 0.55

n=number of patients

Efficacy Conclusions on Interim Clinical Outcomes

[0244] In summary, the pharmacodynamic effects (effects on IgG, anti Dsg-1 and -3 autoantibodies), clinical effects on the activity (PDAI), and clinical outcomes of the disease (disease control, clinical remission) support a potential beneficial effect on the treatment of PV and PF, in combination with low dose prednisone.

[0245] The therapeutic dose regimen to reach the optimal pharmacodynamic effect and clinical efficacy was weekly infusions 10 mg/kg efgartigimod (which translates into 1000 mg efgartigimod PH20 SC).

[0246] In conclusion, in this Phase 2 study, efgartigimod was observed to be a potential effective and safe treatment for PV and PF.

Example 7 -Pharmacodynamics in Pemphigus Patients from Phase 2 Pemphigus Trial (ARGX-113-1701)

(a) Total IgG

[0247] Mean percent change from baseline of total IgG levels versus study days is presented in **Figure 7** and a summary of total IgG level reductions at the end of the induction phase, maintenance phase, and treatment-free follow-up phase is shown in **Table 8.**

[0248] Within the induction period comprising 4 weekly infusions in cohorts 1, 2 and 3, total IgG levels were reduced ranging from mean change from baseline of -55.0% to -67.9%. At the end of the maintenance period, total IgG levels in cohort 1 (2 infusions with a 2-week then a 4-week interval) had almost returned to baseline with a mean change from baseline of - 4.37%. In cohorts 2 (4 infusions every other week) and 3 (6 infusions every other week) IgG levels remained suppressed to mean changes from baseline of -50.67% and -49.21% at the end of the maintenance phase, respectively. This indicated a more efficient suppression of total IgG levels after infusions every other week compared to the 2-week and 4-week interval of cohort 1.

[0249] At the end of the treatment-free follow up, the total IgG levels in cohort 1 (8 weeks after last dosing) were increased to +23.30% from baseline, and had almost returned to baseline to -9.65% and -6.25% from baseline respectively in cohorts 2 and 3 (both 10 weeks after last dosing).

[0250] In cohort 4, the treatment regimen consisting of an induction treatment period with weekly administration until end of consolidation and maintenance treatment period with administration every other week until week 34 was different as compared to cohorts 1-3. Accordingly, no direct comparison of the periods with cohorts 1-3 can be made. The time course of total IgG reduction in cohort 4 is presented in **Figure 7A,** showing a mean change of -69.85% after 4 weeks. There is an apparent correlation between total IgG levels depicted in **Figure 7A** and the PDAI activity score, depicted in **Figures 7B** and **7C**.

[0251] The reduction of total IgG in PV in the Phase 2 study is shown in Figure 8. PD profiles are shown for four weekly administrations of efgartigimod at 25 mg/kg. **Figure 8A** shows the PD profile in PV, and **Figure 8B** the PD profiles in healthy volunteers (HV) and PV. Similar PD profiles were observed between PV and HV, in line with expectations based on HV and modeling.

[0252] Figures 9A and **9B** depict PD profiles for four weekly administrations of efgartigimod at 10 mg/kg. Figure 9A shows the PD profile in PV, and Figure 9B shows PD profiles in HV, PV, myasthenia gravis (MG) and ITP.

[0253] Figures 10A and 10B depict serum levels of total IgG in A) cohort 1-3, B) cohort 4. Data is from results of the Phase 2 trial (1701 study) of efgartigimod in pemphigus patients, date cut-off June 24 2020

[0254] Figures 11A-11H depict serum levels of IgG subclasses IgG1, IgG2, IgG3, IgG4 in cohort 1-3 (A - D) and in cohort 4 (E - H). Data is from Phase 2 trial (1701 study) of efgartigimod in pemphigus patients, date cut-off June 24 2020.

Table 8. Total IgG Reduction: Percent Change from Baseline in Cohorts 1-3 - Efficacy Analysis Set.

		Cohort 1 (10 mg/kg)	**Cohort 2 (10 mg/kg)**	**Cohort 3 (10 mg/kg)**
End of induction phase[a]	n	2	3	7
	mean (SD)	-55.00 (1.980)	-67.90 (5.205)	-63.44 (8.445)
	median	-55.00	-66.70	-60.70
	range (min, max)	-56.4, -53.6	-73.6, -63.4	-74.4, -53.8
	95% CI	-72.8, -37.2	-80.8, -55.0	-71.3, -55.6
End of maintenance phase[b]	n	3	3	7
	mean (SD)	-4.37 (8.372)	-50.67 (5.501)	-49.21

		Cohort 1 (10 mg/kg)	Cohort 2 (10 mg/kg)	Cohort 3 (10 mg/kg)
				(10.006)
	median	-9.10	-52.90	-50.50
	range (min, max)	-9.3, 5.3	-54.7, -44.4	-63.5, -36.8
	95% CI	-25.2, 16.4	-64.3, -37.0	-58.5, -40.0
End of treatment-free follow-up[c]	n	3	2	4
	mean (SD)	23.30 (22.761)	-9.65 (30.476)	-6.25 (22.809)
	median	29.10	-9.65	-7.45
	range (min, max)	-1.8, 42.6	-31.2, 11.9	-30.0, 19.9
	95% CI	-33.2, 79.8	-283.5, 264.2	-42.5, 30.0

[a] End of induction phase for cohorts 1 to 3 at Visit 5.
[b] End of maintenance phase (cohort 1 - M2; cohort 2-M4; cohort 3-M6)
[c] End of treatment-free follow-up (cohort 1 - FU2; cohorts 2 and 3 - FU3)
n=number of patients; CI=confidence interval

[0255] In summary, pemphigus patients treated with efgartigimod exhibited an approximate 40% reduction in total serum IgG level following the first infusion as compared to baseline. The median pharmacodynamic (PD) effect at 10 mg/kg following 4 weekly infusions at day 29 was 62% reduction of total IgG, while for the 25 mg/kg dose it was 66%.

(b) Anti-Dsg Antibodies

[0256] Mean levels for anti-Dsg-3 and anti-Dsg-1 levels versus study days is presented in **Figure 12A-D**. Serum levels of anti-Dsg-1- and -Dsg-3-specific IgG, the pathogenic autoantibodies in pemphigus which are predominantly of the IgG4 subclass, decreased in a similar fashion as total IgGs. A rapid clearance of pathogenic antibodies was observed and reached a median 61% reduction from baseline for anti-Dsg1 and 49% reduction for anti-Dsg3 antibodies at the end of the induction phase in all patients in the efficacy analysis set.

[0257] A clear association between pathogenic anti-Dsg-1/-3 autoantibody level reduction and improvement in the PDAI score was observed throughout the trial. At the end of 4 weeks of fixed dosing, the median reduction in PDAI activity scores was 75% **(Figures 6A-6E).** These observations were consistent regardless of pemphigus subtypes or disease history suggesting a potential broad utility of efgartigimod in pemphigus.

[0258] Due to variable numbers and frequencies of maintenance visits in cohorts 1 to 3, the median PD effect varied. The median PD effect in cohort 1 at the second maintenance visit was 9% IgG level reduction, in cohort 2 at the fourth maintenance visit 53% reduction, and in cohort 3 at the sixth maintenance visit 51% reduction. Patients in cohort 4 who achieved EoC and switched to biweekly dosing of efgartigimod had a sustained IgG level reduction at the level of approximately 50% for as long as biweekly infusions were maintained.

[0259] Similarly, the suppression of pathogenic antibodies could be maintained, albeit more heterogenous for anti-Dsg 3 antibodies. Extended treatment with efgartigimod in cohorts 3 and 4 translated into sustained PDAI activity reduction. 7 patients from cohort 3 completed the study with a median 78% PDAI activity score reduction (ranging between increase of 230 and reduction of 100) from baseline, and 4 patients from cohort 4 demonstrated a median 93% reduction (range 79, 100) from baseline at the end of study.

[0260] At the end of the treatment-free follow up (week 8 or week 10 post last drug administration), IgG levels rose back to normal, ranging between 39% reduction and 43% increase from baseline. For anti-Dsg1 autoantibodies the median change from baseline was 80% reduction and for anti-Dsg3 antibodies 43% reduction, overall depicting longer suppression of pathogenic antibodies as opposed to total IgG levels.

Example 8: Efgartigimod Monotherapy

[0261] Eight patients who received efgartigimod monotherapy at baseline exhibited a median 61% reduction in PDAI activity score (ranging between increase of 411 and reduction of 100) at the end of 4 weeks of fixed dosing as compared to a median 92% drop (ranging between increase of 50 and reduction of 100) in patients who additionally received prednisone at baseline. While concomitant low dose prednisone appeared to afford deeper PDAI activity score reduction, the evident effect of efgartigimod monotherapy suggested a large contribution of efgartigimod alone to clinical efficacy and a potential synergy of efgartigimod and low-dose corticosteroids.

Example 9 - Final Clinical Outcomes

[0262] Fast onset of efgartigimod effects resulted in a rapid DC in 28 out of 31 patients (90%) with a median time to DC of 16 days (range 6, 92) **(Table 9).** DC was achieved in comparable proportions in PV and PF, in mild and moderate disease, in newly diagnosed and relapsing patients, and in patients receiving efgartigimod monotherapy or efgartigimod with concomitant prednisone at baseline. Prolonged maintenance therapy led to 13 complete clinical remissions (5 patients in cohort 3 and 8 patients in cohort 4). Occurrence of relapse was assessed in patients who reached DC (n=28) and was reported in 10 patients (36%).

Table 9. Incidence of disease control, clinical remission, and relapses in overall population and

by subgroups from the efficacy analysis set.

	Disease Control	Clinical Remission	Relapse (from DC)
Overall, n (%)	31	22	28
yes	28 (90)	13 (59)	10 (36)
no	3(10)	9 (41)	18 (64)
Median time to (range)	16 (6 -92)	43 (13 - 287)	87 (10-211)
Pemphigus Vulgaris, n/N (%)	22/24 (92)	9/15 (60)	8/22 (36)
Pemphigus Foliaceus, n/N (%)	6/7 (86)	4/7 (57)	2/6 (33)
Disease history, n/N (%)			
Relapsing patients	18/19 (95)	7/13 (54)	7/18 (39)
Newly Diagnosed patients	10/12 (83)	6/9 (67)	3/10 (30)

[0263] Fast onset of action of efgartigimod and high rate of EoC, triggering steroid tapering, overall led to a low prednisone consumption. Baseline median prednisone equivalent dose was 0.26 mg/kg/day (range 0.06, 0.54) for cohort 1-3 patients and 0.31 mg/kg/day (range 0.06, 0.61) for cohort 4 patients. At the time of EoC, only assessed in cohort 4, median concomitant prednisone dose was 0.28 mg/kg per day (range 0.22, 0.40 mg/kg/day, n=11). The median daily prednisone dose at the time cohort 3 and 4 patients achieved CR was 0.27 mg/kg/day (range 0.06-0.48).

Example 10 - Pharmacokinetic Parameters

[0264] Pharmacokinetic parameters in patients treated IV with 10 mg/kg or 25 mg/kg of efgartigimod were in line with PK data in healthy volunteers dosed with 10 or 25 mg/kg [Ulrichts, P., et al., Neonatal Fc receptor antagonist efgartigimod safely and sustainably reduces IgGs in humans. J Clin Invest, 2018. 128(10): p. 4372-4386.] and with PK data from other studies with 10 mg/kg [Howard, J.F., Jr., et al., Randomized phase 2 study of FcRn antagonist efgartigimod in generalized myasthenia gravis. Neurology, 2019. 92(23): p. e2661-e2673 and Newland, A.C., et al., Phase 2 study of efgartigimod, a novel FcRn antagonist, in adult patients with primary immune thrombocytopenia. Am J Hematol, 2020. 95(2): p. 178-187]. No clear deviation from proportionality could be observed. Negligible accumulation of efgartigimod was apparent after weekly or biweekly dosing as shown by consistent C_{max} and C_{trough}. After the first weekly infusion of 10 or 25 mg/kg, mean C_{max} observed at the end of the infusion was 178 µg/mL and 586 µg/mL, and median C_{trough} observed at the end of the dosing interval was 9.34

µg/mL and 26.0 µg/mL, respectively.

Example 11 - Onset of Action, Depth of Response, and Conditions of Use for Achieving Complete Clinical Remission

[0265] Data presented in this example are from the second interim analysis of Study ARGX-113-1701.

[0266] Clinical efficacy in pemphigus can be interpreted from PDAI score improvement, achieving defined clinical outcomes, the prednisone dose that is required to achieve or sustain clinical improvements, and prevention of relapse.

[0267] Rapid PDAI activity score improvements and achieving early DC (i.e., within 4 weeks) indicate fast onset effects of treatment with efgartigimod. Whereas, overall, 90% (28 of 31) of patients across treatment cohorts achieved DC and 77% (24 of 41) of patients across treatment cohorts achieved DC within 4 weeks, it was also noted that for a majority of the patients (N=16) DC was achieved after only 1 or 2 drug administrations, i.e., observed before or at visit 3. The median time to achieve DC was 15 days for patients treated with efgartigimod alone or 22 days in patients treated with efgartigimod in combination with oral prednisone.

[0268] An overall summary of clinical outcomes observed in each cohort is presented in **Table 10.** Clinical outcomes that were assessed in cohorts 1-3 included DC and relapse. First observed in cohort 3, patients also achieved complete healing of all lesions. A complete healing of all lesions was defined as CR and corresponds to a PDAI activity score of 0. Since in cohorts 1-3, according to the protocol, CR was not a clinical outcome recorded by the investigators, the occurrences of CR in cohort 3 are reported based on the recorded PDAI activity score. For cohort 4, the assessment of EoC and CR were added as clinical outcomes.

Table 10. Summary of Clinical Outcomes in Cohorts 1-4

Outcome	**Cohort 1 (N = 4)**	**Cohort 2 (N = 5)**	**Cohort 3 (N = 7)**	**Cohort 4 (N = 15)[a]**
DC	4 (100%)	3 (60%)	7 (100%)	14 (93%)
EoC	NA	NA	NA	11 (73%)
CR[b]	NA	NA	5 (71%)	7 (47%)

CR=complete clinical remission; DC=disease control; EoC=end of consolidation; N=number of patients; NA=not applicable.
[a] Nine patients with an ongoing treatment at the time of interim analysis.
[b] CR calculated for cohort 3, investigator-assessed for cohort 4.

[0269] Since the prednisone dose at the time of CR was found to be a contributing factor for achieving CR, the rate of achieving CR was analyzed in patients receiving at least biweekly

efgartigimod treatment in combination with intermediate oral prednisone dose (i.e., between 0.25 and 0.50 mg/kg/day). In total, 10 patients were treated with this dose regimen of which 70% (7 out of 10 patients) achieved CR; 3 patients in cohort 3 and 4 patients in cohort 4. Of note, 6 of the 7 patients who achieved CR on the treatment combination of at least biweekly efgartigimod dosing and oral prednisone equivalent dose between 0.25 and 0.50 mg/kg/day had received a stable non-tapered prednisone dose until CR was achieved.

[0270] In cohort 3, 3 patients treated with at least biweekly efgartigimod and prednisone 0.25-0.50 mg/kg/day achieved CR. The other 4 patients were treated without prednisone during the efgartigimod treatment phase (N=1), with prednisone 0.06-0.14 mg/kg/day (N=2), or prednisone was tapered while the patient was on weekly efgartigimod treatment (N=1; prednisone taper from 0.54 mg/kg/day to 0.13 mg/kg/day in 4 steps in 30 days; PDAI activity improvement).

[0271] In cohort 4, with a prolonged treatment duration and more standardized prednisone use, 93% (14 out of 15 patients) achieved DC, 73% (11 out of 15 patients) achieved EoC, and 47% (7 out of 15) achieved CR at the time of interim analysis. In addition to the 4 patients who achieved CR with at least biweekly efgartigimod and prednisone 0.25-0.50 mg/kg/day treatment combination, 3 patients achieved CR with slightly lower prednisone doses between 0.19 and 0.24 mg/kg/day.

[0272] CR on minimal therapy was defined as the absence of new or established lesions while the patient was receiving minimal therapy. Minimal therapy was defined as prednisone (or equivalent) at a dose of 10 mg/d or less and/or minimal adjuvant therapy for at least 2 months). CR on minimal therapy was the next disease outcome parameter after achievement of DC, EoC, and CR at any prednisone dose. For that reason, the daily prednisone doses and duration were analyzed for patients who had achieved CR. At the time of the second interim analysis, 3 patients had achieved CR on minimal therapy (one patient in cohort 3 and two patients in cohort 4). In addition, other patients had a sustained CR with a prednisone dose higher than 10 mg/day or for less than 8 weeks, had achieved EoC without achieving CR and prednisone was tapered to 10 mg/day or below, or had achieved EoC with a prednisone dose higher than 10 mg/day. A small number of patients had only achieved DC, had an insufficient clinical improvement, or discontinued the trial early.

Example 12 - Discussion and Summary of Phase 2 Clinical Trial

[0273] During the efgartigimod induction phase, a sharp decrease of total serum IgG, IgG subclasses, and anti-Dsg1/3 autoantibodies was observed, by about 70% after a weekly treatment course of 2-3 weeks. In contrast, the B-cell depleting antibody rituximab demonstrates a slow and progressive decline in autoantibody titers within months, illustrating the critical differences in the modes of action between treatments. Blockade of FcRn with an FcRn antagonist, e.g., efgartigimod, causes an immediate degradation of autoantibodies responsible for keratinocyte detachment and keratinocyte destruction, while removal of

autoantibody-producing B cells with a B-cell depleting agent, e.g., rituximab, has no immediate impact on circulating autoantibodies that have a typical half-life of about 3 weeks.

[0274] The utility of IVIg to saturate FcRn and thereby eliminate pathogenic antibodies has been rationalized in several studies. Grando, SA, et al., Int Immunopharmacol, 2020. 82: 106385; Amagai, M, et al., J Am Acad Dermatol, 2009. 60(4): 595-603. Moreover, because FcRn-deficient mice are resistant to experimental pemphigus (Li, N, et al., J Clin Invest, 2005. 115(12): 3440-50) and expression of FcRn in keratinocytes has been documented (Cauza, K, et al., J Invest Dermatol, 2005. 124(1): 132-9), it is plausible that protection from pathogenic autoantibodies via FcRn inhibition is mediated not only via induction of autoantibody degradation but also via direct blockade of FcRn in keratinocytes. Thus overall, because the disease activity is directly linked to the presence of pathogenic antibodies with multitudes of pathogenic actions, strategies precisely aiming at efficient depletion of pathogenic antibodies can have a profound impact on patients' response to therapy. In line with that, in all cohorts of the study, DC was achieved as fast as within 1-4 weeks in the vast majority of patients, with or without concomitant prednisone therapy. DC was similarly observed in patients with PV and PF, newly diagnosed and relapsing, and mild and moderate pemphigus.

[0275] The adaptive nature of the trial has led to the clear observation that the lowering of serum IgG was well maintained while on biweekly maintenance phase of efgartigimod, whereas the four-weekly dosing was insufficient to keep pathogenic autoantibodies suppressed enough and led to a recovery of IgG serum levels after 4 to 8 weeks. In the cohorts with prolonged efgartigimod treatment (cohort 3: 11 weeks; cohort 4: 34 weeks), CR could be reached within weeks (median: 6 weeks, range: 2 to 41 weeks), providing encouraging and supporting data to test the rate of CR and time to CR in a larger cohort of pemphigus patients in a phase 3 trial.

[0276] Notably, CR occurred optimally when associated with prednisone doses ranging from 0.06 to 0.48 mg/kg/day (median: 0.28 mg/kg/day), indicating an additive effect of prednisone to efgartigimod. Corticosteroids are known to regulate gene expression and broadly suppress many inflammatory genes acting on leukocyte movement, leukocyte function, and humoral factors as well as cytokine signaling. Moreover, beyond its well-described immunosuppressive effects, prednisone has been shown also to up-regulate expression of genes encoding keratinocyte adhesion molecules such as E-cadherin and desmogleins. Nguyen, VT, et al., J Biol Chem, 2004. 279(3): 2135-46. Altogether, additive effects of prednisone in pemphigus patients receiving efgartigimod could be diversified.

[0277] Whereas anti-Dsg-1 and anti-Dsg 3 autoantibody reductions generally followed the course of total IgG level reductions, anti-Dsg 3 antibody levels were found to correlate less with IgG serum levels in some patients, without leading to increases in PDAI. This is in line with previous reports indicating that circulating anti-Dsg 3 antibody levels do not necessarily correlate with disease activity. Colliou, N, et al., Sci Transl Med, 2013. 5(175): 175ra30. Treatment discontinuation resulted in a progressive return to normal IgG levels within 4 to 6 weeks, although many patients remained in disease remission or with a disease of low grade.

Intriguingly, in several patients, autoantibody levels remained suppressed in the treatment-free follow-up while total IgG surged up as expected.

[0278] In this Phase 2 clinical trial, relapses occurred in 36% of the patients; they mainly consisted of early relapses, i.e., occurring before CR, and were observed during increased administration intervals of 2 or more weeks. Late relapses, i.e., after CR, occurred at biweekly dosage or during treatment-free follow-up. In contrast, no relapse occurred when patients were maintained at weekly efgartigimod dosage, suggesting that a weekly administration of efgartigimod beyond CR achievement may support prevention of the relapses.

Example 13 - Rationale for Phase 3 Trials with Subcutaneous administration

[0279] As pemphigus is a chronic disease, current therapies require chronic administration. In Phase 3 trials described in below, efgartigimod is administered with rHuPH20 via the subcutaneous (SC) route in a flat dose in a new highly concentrated formulation, which is more convenient for patients.

[0280] In order to allow for a convenient SC administration with efgartigimod to achieve the targeted exposure and PD effect, efgartigimod is co-formulated with rHuPH20. This compound is being used in co-formulations with approved therapeutic antibodies to facilitate SC injection with volumes larger than 2 mL. The use of rHuPH20 in combination with other therapeutic proteins typically increases the absorption rate of these proteins, albeit to different extents increased rate of absorption of efgartigimod by rHuPH20 is expected to increase the overall exposure to efgartigimod after SC administration allowing administration of a SC dose in an acceptable volume and duration of administration that targets an exposure that results in a close to the maximal PD effect.

[0281] A dose of 1000 mg efgartigimod SC is comparable to 10 mg/kg efgartigimod administered as IV infusion with respect to effect on IgG levels. The 10 mg/kg IV dose: 1) resulted in transient clinical efficacy in a Phase 2 trial in patients with ITP and a prolonged clinical effect in a Phase 2 trial in patients with generalized myasthenia gravis (gMG) following 4 weekly infusions; 2) was shown to result in close to saturated PD effect: Dosing higher or more frequently than weekly is not expected to result in an improved PD effect (i.e., further lowering of autoantibodies) and/or clinical effect and may be associated with a less optimal risk/benefit ratio. Dosing lower is expected to result in a lower PD effect and thus is likely to result in a less consistent and/or incomplete clinical response, which is undesirable given the serious and chronic manifestations of pemphigus; and 3) demonstrated a favorable safety profile in Phase 2 studies in patients with gMG and patients with ITP.

Example 14 - Design of Phase 3 Clinical Trial with Subcutaneous Administration - ARGX-113-1904

[0282] ARGX-113-1904 is a randomized, double-blinded, placebo-controlled trial to investigate the efficacy, safety, and tolerability of efgartigimod PH20 SC in adult patients with pemphigus (vulgaris or foliaceus). The trial is given the acronym ADDRESS.

[0283] Efgartigimod PH20 SC is administered on day 1 and day 8 at a dose of 2000 mg, followed by weekly administrations of 1000 mg until complete remission on minimal therapy (CR on minimal therapy). Patients will also be administered placebo with the same regimen.

[0284] This trial intends to demonstrate that efgartigimod PH20 SC with an add-on therapy of low doses of oral prednisone is a possible treatment modality for PV and PF, the administration of which will lead to early disease remission at minimal prednisone dose. The efficacy, safety, patient outcome measures, tolerability, immunogenicity, PK, and PD of efgartigimod PH20 SC is evaluated in patients with PV or PF.

Overall Design

[0285] This is a prospective, multicenter, randomized, double-blinded, placebo-controlled trial to investigate the efficacy, safety, patient outcome measures, tolerability, immunogenicity, PK, and PD of efgartigimod PH20 SC in adult patients aged 18 to 80 years with PV or PF. See **Figure 13** for a diagram of the trial design. Enrolled patients are either those who are newly diagnosed or experiencing flare as follows:

- Moderate to severe (PDAI activity score ≥15) newly diagnosed and naive to treatment.
- Moderate to severe (PDAI activity score ≥15) newly diagnosed while receiving a first course of oral prednisone (or equivalent). According to clinical judgment, the patient has shown no significant improvement of PV or PF signs for at least 2 weeks before baseline and is considered fit to start prednisone treatment at 0.5 mg/kg daily (qd) at baseline.
- Moderate to severe (PDAI activity score ≥15), experiencing flare, and off prednisone therapy ± a conventional immunosuppressant (e.g., azathioprine, cyclophosphamide, methotrexate, mycophenolate mofetil). Conventional immunosuppressants are discontinued before baseline.
- Moderate to severe (PDAI activity score ≥15), experiencing flare, while receiving a tapered dose of oral prednisone (or equivalent), provided that prednisone (or equivalent) has been given at stable dose ± a conventional immunosuppressant (e.g., azathioprine, cyclophosphamide, methotrexate, mycophenolate mofetil) and patients are fit to start prednisone treatment at 0.5 mg/kg qd at baseline. Conventional immunosuppressants are discontinued before baseline.

[0286] The trial comprises a screening period of up to 3 weeks, a treatment period of up to 30 weeks, and an 8-week follow-up period for patients who do not enroll into the open-label extension (OLE) trial ARGX-113-1905 **(Example 15)**.

[0287] After confirmation of eligibility, patients are randomized in a 2:1 ratio to receive efgartigimod PH20 SC or placebo, as follows:

- Efgartigimod PH20 SC is administered on day 1 and day 8 at a dose of 2000 mg, followed by weekly administrations of 1000 mg until CR on minimal therapy is observed. Efgartigimod PH20 SC is administered at on-site visits until complete remission (CR), with a minimum of 6 weekly on-site visits after baseline. After achieving CR, efgartigimod PH20 SC is administered at on-site visits or at home by a nurse until CR on minimal therapy is achieved.
- Placebo (vehicle with 2000 U/mL of rHuPH20) SC is administered using the same regimen.

[0288] CR is defined as the absence of new lesions and complete healing of established lesions (except for post-inflammatory hyperpigmentation or erythema from resolving lesions). CR on minimal therapy is defined as the absence of new lesions and complete healing of established lesions while the patient is receiving minimal prednisone therapy of ≤10 mg/day for at least 8 weeks. Randomization is stratified by disease status (experiencing flare and newly diagnosed), disease severity (PDAI activity score <30 and PDAI activity score ≥30), and body weight (≤77.5 kg and >77.5 kg) at baseline. Patients with severe PV or PF (PDAI activity score ≥45) will comprise a maximum of 30% of the overall trial population.

Concomitant Pemphigus Therapy

[0289] All patients, regardless of treatment assignment, concomitantly receive oral prednisone (or equivalent such as prednisolone) at a starting dose of 0.5 mg/kg qd. Except for oral prednisone (or equivalent), no other systemic therapies (e.g., immunosuppressants, IVIg, dapsone, immunoadsorption, anti-CD20 biologics) are permitted during the trial.

[0290] Patients visit the clinic weekly, until CR and for a minimum of 6 weeks, to receive efgartigimod PH20 SC and to be evaluated for disease activity and disease outcome. After CR, efgartigimod PH20 SC is administered at on-site visits or at home by a nurse until CR on minimal therapy. At any post-baseline visit before disease control (DC) is achieved, the prednisone dose is adjusted by incrementing dosage by 1 or 2 steps according to clinical judgment in case of disease progression or insufficient clinical change. Disease progression and insufficient clinical change are judged as follows:

- Disease progression: increase of at least 5 in PDAI activity score compared to baseline score, observed at any post-baseline visit before DC.
- Insufficient clinical change: absence of DC after 3 to 4 weeks of the patient being treated at the starting baseline prednisone (or equivalent) dose or after 3 to 4 weeks of any new incremented dose of prednisone.

[0291] The prednisone dose escalation rules based on starting dose are as follows:

- Stepwise escalation of daily prednisone dose occurs in 1 or 2 steps according to clinical judgment, with a recommendation of 1 step for moderate progression and 2 steps for severe progression.
- Adjustment by incrementing dosage by 1 step in case of insufficient clinical change.
- Possible further escalation from the previous step by 1 or 2 steps (e.g., 0.75 mg/kg to 1 or 1.25 mg/kg qd or from 1 mg/kg to 1.25 or 1.5 mg/kg qd), according to clinical judgment and under the same recommendation as above.
- Maximum escalation to 1.5 mg/kg qd for 3 weeks.

[0292] If, after a minimum of 3 weeks of oral prednisone at 1.5 mg/kg qd, DC is not attained, then the patient is considered a treatment failure.

[0293] For patients achieving DC with a daily prednisone dose of 0.5 mg/kg, prednisone is maintained at 0.5 mg/kg qd until CR and 2 weeks thereafter. For patients achieving DC with an escalated prednisone dose (i.e., >0.5 mg/kg qd), the prednisone dose is maintained until 2 weeks after achieving DC, then tapering is performed according to the following stepwise procedure: dose reductions by 0.25 mg/kg qd every 2 weeks until the starting dose is reached (i.e., 0.5 mg/kg qd). Then the starting dose (0.5 mg/kg qd) is maintained until a sustained CR is achieved for 2 weeks, or prednisone tapering may be initiated in case of sustained end of consolidation (EoC, defined by the time at which no new lesions have developed for a minimum of 2 weeks and approximately 80% of lesions have healed) for at least 4 weeks. Further tapering is performed thereafter, as long as CR or EoC are sustained. Each new tapered prednisone dose until 20 mg/day is maintained for 2 weeks. Then, the prednisone dose is further tapered by 2.5 mg/day per week. When 10 mg/day is reached, this dose level is maintained until CR on minimal therapy has been achieved. Prednisone can then be further tapered upon clinical judgment by the investigator.

[0294] In case of flare in the period between DC and CR on minimal therapy, the prednisone dose is increased. A flare is defined by the appearance of 3 or more new lesions in a 4-week period that do not heal spontaneously within 1 week, or by the extension of established lesions in a patient who had achieved DC. If the flare occurs after CR and efgartigimod PH20 SC was administered at home by a nurse, the patient will resume weekly on-site visits until he/she achieves CR again. Patients who are not controlled by a dose that is 2 dose levels above the dose at which the flare between DC and CR on minimal therapy is observed and that is at least 0.3 mg/kg/day are considered treatment failures. At visits when at least 1 new lesion is observed or established lesions remain extensive without being defined as a flare, the prednisone dose is maintained or may be increased, according to clinical judgment. If the lesion resolves, tapering of the prednisone dose is pursued as planned.

[0295] Patients who experience treatment failure, or flare after achieving CR on minimal therapy, are allowed to roll over into the OLE trial ARGX-113-1905 (Example 13) earlier than week 30. Patients who do not roll over into trial ARGX-113-1905 will complete an 8-week treatment-free follow-up period. Patients experiencing an SAE related to prednisone may also benefit from an early roll over to the OLE, according to clinical judgment.

Number of Participants:

[0296] A total of 150 patients with PV or PF are randomized as follows:

- A total of 126 patients with PV are randomized in a 2:1 ratio to receive efgartigimod PH20 SC or placebo, respectively
- Up to 24 patients with PF may be randomized in a 2:1 ratio to receive efgartigimod PH20 SC or placebo, respectively

Intervention Groups and Duration:

[0297] The maximum possible total trial duration for each patient is up to 41 weeks:

- Screening period: up to 3 weeks
- Treatment period: 30 weeks from baseline
- Follow-up period: 8 weeks after the last dose of efgartigimod PH20 SC for patients who do not enroll in the OLE trial

[0298] The trial is a prospective, multicenter, randomized, double-blinded, placebo-controlled trial to investigate the efficacy, safety, tolerability, immunogenicity, PK, and PD of efgartigimod PH20 SC in adult patients aged 18 to 80 years with moderate to severe pemphigus vulgaris (PV) or pemphigus foliaceus (PF). This trial demonstrates that efgartigimod PH20 SC with an add-on therapy of low doses of oral prednisone is a highly effective treatment modality for PV and PF, the administration of which leads to early disease remission at minimal prednisone dose.

Prednisone tapering

[0299] Two weeks after DC is achieved after an escalated prednisone dose above 0.5 mg/kg qd, prednisone tapering begins. Prednisone tapering is a procedure that is applied by all practitioners, in order to minimize the side effects of corticosteroids while preventing a flare of

the disease. In this trial:

- Two-week intervals between tapering are defined with a step down of 25% from 1.5 mg to 0.5 mg/kg per day (e.g., from 1.5 to 1.25 mg/kg per day, then 1 mg/kg and 0.75 mg/kg per day). The dosage of prednisone at which CR is achieved (which cannot be lower than the starting dose 0.5 mg/kg per day) is maintained for 2 weeks after CR for observing a sustained CR, or prednisone tapering may be initiated or continued in case of sustained EoC (defined by the time at which no new lesions have developed for a minimum of 2 weeks and approximately 80% of lesions have healed) for at least 4 weeks.
- When CR is observed on daily prednisone of no lower than 0.5 mg/kg for 2 weeks, or when sustained EoC is observed, tapering may be initiated. Each new tapered prednisone dose until 20 mg/day is maintained for 2 weeks. Then, the prednisone dose is further tapered by 2.5 mg/day per week. When 10 mg/day is reached, this dose level is maintained until CR on minimal therapy has been achieved.
- When CR on minimal therapy is achieved, prednisone can be tapered further upon clinical judgment, with the recommendation of a prednisone decrease by 2.5 mg/day every 4 weeks.

[0300] When transient lesions appear, prednisone tapering is temporarily delayed or prednisone may be increased according to clinical judgment. Dose escalation is performed when flare occurs. Taken together, the trial aims to be as close as possible to the real-world clinical practice regarding the prednisone regimen policy, while it allows studying the potential of prednisone sparing when associated with efgartigimod PH20 SC treatment.

Proportion of patients who achieve CR on minimal therapy within 30 weeks as primary endpoint and treatment duration of 30 weeks:

[0301] A trial duration of 30 weeks was selected based on results from the phase 2 trial ARGX-113-1701 in patients with pemphigus for encompassing the observation of the primary endpoint of CR on minimal therapy in the majority of patients belonging to the active group. The time point at which CR on minimal therapy is anticipated to be achieved for most patients varies between 24 and 30 weeks in patients under active treatment and depends on the body weight of the patients. Accounting for this variation between patients, all patients having achieved CR on minimal therapy within the 30 weeks of the trial are considered as responsive to treatment in the primary endpoint.

Tailored regimen of efgartigimod PH20 SC and rollover to the OLE:

[0302] The trial comprises a screening period of up to 3 weeks, a treatment period until CR on

minimal therapy of 30 weeks, and an 8-week follow-up period for patients who do not enroll into the OLE trial ARGX-113-1905 (Example 13). A tailored efgartigimod PH20 SC regimen is proposed due to its rapid effect and the variability of disease activity between patients. Patients are administered the efgartigimod PH20 SC with concomitant low dose prednisone regimen subcutaneously until CR on minimal therapy is achieved. Then, efgartigimod PH20 SC administration is stopped, whereas the concomitant treatment by prednisone is pursued with the option for further tapering to achieve CR off therapy.

Inclusion Criteria

[0303] Participants are eligible to be included in the trial only if all of the following criteria apply:

- Ability to understand the requirements of the trial, to provide written informed consent (including consent for the use and disclosure of research-related health information), willingness and ability to comply with the trial protocol procedures (including required trial visits).
- The patient is male or female, and aged 18 years to 80 years at the time of signing the informed consent form (ICF).
- The patient has a clinical diagnosis of PV (mucosal, cutaneous, mucocutaneous) or PF which has been confirmed by cutaneous histology, positive direct immunofluorescence (IF), and positive indirect IF and/or ELISA.
- The patient meets 1 of the following profiles:

  Newly diagnosed disease with PDAI activity score ≥15 at baseline and naive to treatment.

  Newly diagnosed disease with PDAI activity score ≥15 while receiving a first course of oral prednisone (or equivalent). According to clinical judgment, the patient has shown no significant improvement of PV or PF signs for at least 2 weeks before baseline and is considered fit to start prednisone treatment at 0.5 mg/kg qd at baseline.

[0304] Experiencing flare with PDAI activity score ≥15, a maximum of 4 years since diagnosis, and off prednisone therapy ± a conventional immunosuppressant (eg, azathioprine, cyclophosphamide, methotrexate, mycophenolate mofetil). Conventional immunosuppressants are discontinued before baseline.

[0305] Experiencing flare with PDAI activity score ≥15, a maximum of 4 years since diagnosis, and receiving a tapered dose of oral prednisone (or equivalent), provided that prednisone has been given at stable dose ± a conventional immunosuppressant (e.g., azathioprine, cyclophosphamide, methotrexate, mycophenolate mofetil) for at least 2 weeks and patients are fit to start prednisone treatment at 0.5 mg/kg qd at baseline. Conventional immunosuppressants are discontinued before baseline.

Exclusion Criteria

[0306] Participants are excluded from the trial if any of the following criteria apply:

- Patient has a confirmed diagnosis of paraneoplastic pemphigus, drug-induced pemphigus, pemphigus vegetans, pemphigus erythematosus, or any other non-PV/non-PF autoimmune blistering disease.
- Patients with mild disease severity as defined by PDAI activity score <15 at baseline.
- Patients who show a significant improvement of PV or PF in the period from screening to baseline according to clinical judgment (e.g., the patient has achieved DC or a substantial reduction in PDAI activity score during screening period).
- The patient has been administered therapy(ies) other than oral prednisone or conventional immunosuppressants (e.g., azathioprine, cyclophosphamide, methotrexate, mycophenolate mofetil) within 2 months before the baseline visit and that can affect clinical disease activity. For example, excluded medications are intravenous methylprednisolone, dapsone, sulfasalazine, tetracyclines, nicotinamide, plasmapheresis/ plasma exchange, immunoadsorption, and IVIg.
- Use of any monoclonal antibody (including rituximab or another anti-CD20 biologic) within 6 months before the baseline visit.
- Known hypersensitivity to any of the components of the administered treatments.
- The patient has a known contraindication to oral prednisone.
- The patient has a history of refractory disease, as defined by a failure to respond to first-line and second-line therapies.

Trial Intervention(s) Administered

[0307] A list of trial interventions is presented in **Table 11.** Fixed doses of efgartigimod PH20 SC are administered on body sites spared of any cutaneous pemphigus lesions, the abdomen being used as preferred site. If the abdomen is affected by lesions, optional sites (thighs and the arms) may be chosen. The placebo solution for injection contains the same excipients as the efgartigimod PH20 SC solution for injection but without efgartigimod. Masked vials are provided to preserve the study blind. Additionally, patients receive oral prednisone (or equivalent such as prednisolone) as concomitant therapy.

Table 11. Trial Interventions - ARGX-113-1904

Intervention	Efgartigimod PH20 SC	Placebo	Prednisone (or Equivalent)
Type	Biologic	Other: placebo	Non-IMP
Dose formulation	Efgartigimod 165 mg/mL + 2000	Vehicle + rHuPH20 for SC	Prednisone/prednisolone tablets for oral

Intervention	Efgartigimod PH20 SC	Placebo	Prednisone (or Equivalent)
	U/mL rHuPH20 solution for	administration	administration
	SC injection to be dosed at a fixed dose of 1000 mg per injection		
Unit dose strength(s)	165 mg/mL	Placebo	Prednisone 5 mg, 10 mg, 20 mg, 50 mg
			Prednisolone 5 mg
Dosage level(s)	2 × 1000 mg administered in separate sites on day 1 and day 8 followed by weekly 1000 mg administrations until CR on minimal therapy	2 × SC administrations in separate sites on day 1 and day 8 followed by weekly single administrations until CR on minimal therapy	Refer to Concomitant Pemphigus Therapy
Route of administration	Abdominal SC injection(s); preferred site[a]	Abdominal SC injection(s); preferred site"	Oral administration
Sourcing	Provided by the sponsor to the trial site	Provided by the sponsor to the trial site	Provided by the sponsor to the trial site, or provided locally
Packaging and Labeling	The IMP is provided in glass vials. Each glass vial is labeled as required per country requirement	The IMP is provided in glass vials. Each glass vial is labeled as required per country requirement	Non-IMP is provided in the commercial package, or as magistral preparation upon prescription by the investigator

IMP=investigational medicinal product; SC=subcutaneous(ly)

[a] IMP is administered on body sites spared of any cutaneous pemphigus lesions, the abdomen being used as preferred site. If the abdomen is affected by lesions, optional sites (thighs or arms) may be chosen.

End of Study (EoS)/Early Termination (ET)

[0308] All patients complete week 30/EoS/ET, which is end of study for patients who enroll in the OLE trial ARGX-113-1905.

Treatment-free Follow-Up Period

[0309] Patients who reach week 30, discontinue early, or patients under treatment failure, and who do not roll over to the OLE trial, enter the 8-week treatment-free follow-up period.

Efficacy Assessments

[0310] The efficacy of efgartigimod PH20 SC is assessed at on-site visits by the investigator, assessing the following: PDAI, DC, EoC, CR, CR on minimal therapy, CR off therapy, and flare. The investigator records the daily prednisone dose since the last visit, and treatment failures at on-site visits. For home visits, the investigator calls the patient at least every 2 weeks until CR on minimal therapy is achieved to confirm the patient is still in CR.

[0311] The clinical activity of pemphigus is assessed using the PDAI.

Example 15 - Open Label Phase 3 Extension Clinical Trial with Subcutaneous Administration - ARGX-113-1905

[0312] ARGX-113-1905 is an open-label, long-term extension (OLE) follow-up study, evaluating the safety, tolerability, and efficacy of efgartigimod PH20 SC in adult patients with PV or PF. Trial ARGX-113-1905 provides efgartigimod PH20 SC extended treatment and retreatment options for patients who were randomized to the efgartigimod PH20 SC treatment arm in trial ARGX-113-1904 **(Example 14),** and provides first efgartigimod PH20 SC treatment and retreatment options for patients who were randomized to the placebo treatment arm in trial ARGX-113-1904. The trial is given the acronym ADDRESS+.

[0313] At blinded rollover from trial ARGX-113-1904 into trial ARGX-113-1905 patients may be in different clinical stages. Patients may have achieved CR on minimal therapy, patients may have achieved CR without meeting the minimal therapy criterion, patients may have achieved DC or EoC (the time at which no new lesions have developed for a minimum of 2 weeks and approximately 80% of lesions have healed) without having achieved CR, patients may have experienced treatment failure, or patients may have experienced a flare after having achieved CR on minimal therapy.

[0314] Trial ARGX-113-1905 evaluates the ability to (further) taper prednisone therapy and achieve CR off therapy, the ability to achieve CR and CR on minimal therapy for patients who had not yet achieved CR on minimal therapy, and the ability to treat flare; and assesses patient outcome measures and the safety, PD, PK, and immunogenicity of efgartigimod PH20 SC over the duration of the trial.

Overall Design

[0315] This is a prospective, multicenter, OLE trial on the efficacy, safety, tolerability, immunogenicity, PK, and PD of efgartigimod PH20 SC in adult PV or PF patients, who participated in the antecedent trial ARGX-113-1904. See **Figure 14** for a diagram of the trial design.

[0316] Patients at the EoS visit or early termination visit in trial ARGX-113-1904 are given the option to enroll into trial ARGX-113-1905 after confirmation of eligibility while retaining the blinding of trial ARGX-113-1904. For each patient, the date of the baseline visit in trial ARGX-113-1905 is the same as the date of the EoS visit in trial ARGX-113-1904.

Concomitant Pemphigus Therapy

[0317] At baseline, patients are treated according to their clinical status at EoS of trial ARGX-113-1904, as follows:

- Patients in CR on minimal or off therapy are observed every 4 weeks through on-site visits without efgartigimod PH20 SC treatment. In patients who have achieved CR on minimal therapy (oral prednisone ≤10 mg per day for 8 weeks), prednisone tapering may be pursued upon investigator's judgment until patients achieve CR off therapy.
- Patients who have achieved CR but not CR on minimal therapy receive weekly efgartigimod PH20 SC administrations of 1000 mg while continuing the add-on therapy of oral prednisone until CR on minimal therapy is achieved. Prednisone tapering may be pursued upon investigator's judgment until patients achieve CR off therapy. Patients are treated through on-site visits, or through home visits with an on-site visit once every 4 weeks.
- Patients who have achieved DC but not CR receive weekly on-site administrations of 1000 mg efgartigimod PH20 SC with an add-on therapy of oral prednisone until CR has been achieved, and then continue to receive weekly efgartigimod PH20 SC administrations of 1000 mg until CR on minimal therapy is achieved either on site, or through home visits with an on-site visit once every 4 weeks.
- Patients under certain conditions of treatment failure as defined in trial ARGX-113-1904 (i.e., absence of DC with oral prednisone 1.5 mg/kg/day for a minimum of 3 weeks, or flare between DC and achieving CR on minimal therapy that is not controlled by 2 dose levels above the dose at which the flare is observed and that is at least 0.3 mg/kg/day), or patients with flare after having achieved CR on minimal therapy, may rollover prematurely into trial ARGX-113-1905 and are treated with efgartigimod PH20 SC at a dose of 2000 mg on day 1 and day 8, followed by weekly administrations of 1000 mg with an add-on therapy of oral prednisone until CR on minimal therapy. In the absence of DC, patients receive prednisone at the starting dose of 0.5 mg/kg/day. In case of flare,

patients receive as starting dose the last dose from ARGX-113-1904, with the recommendation of a minimum of 0.3 mg/kg/day when the flare is of mild severity (PDAI activity score <15) and of 0.5 mg/kg/day when the flare is moderate to severe (PDAI activity score ≥15). Dosing of efgartigimod PH20 SC is done during on-site visits or home visits.

- Patients who developed a prednisone-related SAE may prematurely rollover into trial ARGX-113-1905 and may be treated with efgartigimod PH20 SC through on-site or home visits according to the clinical statuses as described above. Concomitant prednisone is considered to fit the clinical statuses as described above as far as compatible with the nature and severity of the SAE, with a lower dose to no concomitant prednisone being considered otherwise.

[0318] At any post-baseline visit before DC is achieved, the prednisone dose is adjusted by incrementing dosage by 1 or 2 steps according to clinical judgment in case of disease progression or insufficient clinical change. Disease progression and insufficient clinical change is defined as follows:

- Disease progression: increase of at least 5 in PDAI activity score compared to baseline score, observed at any post-baseline visit before DC
- Insufficient clinical change: the absence of DC after 3 to 4 weeks of the patient being treated at the starting baseline prednisone dose or after 3 to 4 weeks of any new incremented dose of prednisone

[0319] The prednisone dose escalation rules based on starting dose are as follows:

- Stepwise escalation of daily prednisone dose occurs in 1 or 2 steps according to clinical judgment, with a recommendation of 1 step for moderate progression and 2 steps for severe progression
- Adjustment by incrementing dosage by 1 step in case of insufficient clinical change
- Possible further escalation from the previous step by 1 or 2 steps (e.g., 0.75 mg/kg to 1 or 1.25 mg/kg qd, or from 1 mg/kg to 1.25 or 1.5 mg/kg qd), according to clinical judgment and under the same recommendation as above
- Maximum escalation to 1.5 mg/kg qd for 3 weeks

[0320] If DC is not attained after a minimum of 3 weeks of the patient receiving oral prednisone 1.5 mg/kg qd, then the patient is considered a treatment failure and is withdrawn from the trial.

[0321] For patients who achieve DC with the daily starting dose at baseline, the prednisone dose is maintained at that starting dose qd until CR and for the next 2 weeks. For patients who achieve DC with an escalated prednisone dose, that dose is maintained until 2 weeks after

achieving DC and then is tapered in 25% reductions every 2 weeks until the starting dose has been achieved. This starting dose is maintained until a sustained CR is achieved for 2 weeks, or in case of end of consolidation (EoC) for 4 weeks. Further tapering is performed thereafter, as long as CR or EoC is sustained. Each new tapered prednisone dose until 20 mg/day is maintained for 2 weeks. Then, the prednisone dose is further tapered by 2.5 mg/day per week. When 10 mg/day is reached, this dose level is maintained until CR on minimal therapy has been achieved. The prednisone dose can then be further tapered to reach CR off therapy according to the clinical judgment of the investigator.

Number of Participants:

[0322]

- All participants who were randomized into ARGX-113-1904 (i.e., up to 150 patients with pemphigus vulgaris (PV) or pemphigus foliaceus (PF)) are eligible to rollover to trial ARGX-113-1905

Intervention Groups and Duration:

[0323] Up to 60 weeks for patients who receive efgartigimod PH20 SC administrations up to week 52 and with a follow-up period of 8 weeks after the last efgartigimod PH20 SC administration.

[0324] At visits when at least 1 new lesion is observed or established lesions remain extensive without being defined as a flare (i.e., the appearance of 3 or more new lesions in a 4-week period that do not heal spontaneously within 1 week, or by the extension of established lesions in a patient who had achieved DC), the prednisone dose is maintained or may be increased, according to clinical judgment. If the lesion resolves, tapering of the prednisone dose is pursued as planned.

[0325] In case of flare in the period between DC and CR on minimal therapy, the prednisone dose is increased to achieve DC again. If the flare occurs after CR and IMP was administered at home by a nurse, the patient will resume weekly on-site visits until he/she achieves CR again. Patients who are not controlled by a dose that is 2 dose levels above the dose at which the flare is observed and that is at least 0.3 mg/kg/day, are managed according to clinical judgment, i.e., either receive a further increased prednisone dose or be withdrawn from the trial. Withdrawal of patients with a flare before CR on minimal therapy is defined as treatment failure.

[0326] In case of flare after having achieved CR on minimal therapy (i.e., while being off

efgartigimod treatment), patients are immediately treated/re-treated with a new cycle of efgartigimod PH20 SC. The first day of new efgartigimod treatment will define a new baseline for assessments of outcomes (DC, CR, etc) and prednisone escalation or tapering schedule.

[0327] In the new treatment cycle, patients are administered efgartigimod PH20 SC at a weekly dose of 2000 mg for the first 2 weeks, followed by weekly administrations of 1000 mg until CR on minimal therapy. Oral prednisone is administered at a dose chosen according to clinical judgment, with the recommendation of 0.3 mg/kg qd in the case of mild flare (PDAI activity score < 15) and 0.5 mg/kg qd in the case of moderate to severe flare (PDAI activity score ≥15). The treatment goal of a new treatment cycle is to first achieve DC again and therefore these patients are treated accordingly.

[0328] A new treatment cycle of efgartigimod PH20 SC can be initiated in eligible patients until week 44. In patients requiring a new treatment cycle of efgartigimod PH20 SC between weeks 45 and 49, the initiation is optional and based on clinical judgment and patient consent. A new efgartigimod treatment cycle will not be permitted after week 49 to ensure a minimum of 4 weeks of efgartigimod treatment/cycle.

[0329] Except for oral prednisone, no other systemic therapies (e.g., immunosuppressants, IVIg, dapsone, immunoadsorption, anti-CD20 biologics) are permitted during the trial.

[0330] SC administration is highly preferred over an IV formulation for pemphigus patients in view of the expected need for prolonged weekly administration. Therefore, efgartigimod with rHuPH20 is administered SC weekly until CR on minimal therapy; retreatment may be initiated in case of flare after having achieved CR on minimal therapy. To achieve a fast PD effect and clinical response a dose of 2000 mg is administered in the first 2 weeks, followed by weekly doses of 1000 mg to maintain the PD effect and related clinical response.

Trial Interventions Administered

[0331] A list of trial interventions is presented in **Table 12.** Fixed doses of efgartigimod PH20 SC are administered on body sites spared of any cutaneous pemphigus lesions, the abdomen being used as preferred site. If the abdomen is affected by lesions, optional sites (the thighs and the arms) may be chosen.

Table 12. Trial Interventions - ARGX-113-1905

Intervention	Efgartigimod PH20 SC	Prednisone (or Equivalent)
Type	Biologic	Non-IMP
Dose formulation	Efgartigimod 165 mg/mL + 2000 U/mLrHuPH20 solution for SC injection to be dosed at a fixed dose of 1000 mg per injection	Prednisone/prednisolone tablets for oral administration

Intervention	Efgartigimod PH20 SC	Prednisone (or Equivalent)
Unit dose strength(s)	165 mg/mL	Prednisone 5 mg, 10 mg, 20 mg, 50 mg
		Prednisolone 5 mg
Dosage level(s)	Weekly 1000 mg administrations until CR on minimal therapy, for patients in DC or CR at rollover from ARGX-113-1904.	Refer to Concomitant Pemphigus Therapy
	2 × 1000 mg administered in separate sites on day 1 and day 8 followed by weekly 1000 mg administrations until CR on minimal therapy, for patients under treatment failure in ARGX-113-1904, or patients experiencing flare after CR on minimal therapy.	
Route of administration	Abdominal SC injection(s); preferred site[a]	Oral administration
Use	Investigational drug	Non-IMP or concomitant therapy
IMP	IMP	Non-IMP
Sourcing	Provided by the sponsor to the trial site	Provided by the sponsor to the trial site, or provided locally
Packaging and labeling	The IMP is provided in glass vials. Each glass vial is labeled as required per country requirement	Non-IMP is provided in the commercial package, or as magistral preparation upon prescription by the investigator

CR=complete clinical remission; CR on minimal therapy=complete remission on minimal therapy; DC=disease control; IMP=investigational medicinal product; SC=subcutaneous.

[a] IMP is administered on body sites spared of any cutaneous pemphigus lesions, the abdomen being used as preferred site. If the abdomen is affected by lesions, optional sites (thighs or arms) may be chosen.

[0332] The efficacy of efgartigimod PH20 SC is assessed at on-site and home visits by the investigator, assessing by chronological order the following: PDAI, DC, EoC, CR, CR on minimal therapy, CR off therapy, and flare. The investigator records the average daily prednisone dose since the last visit, and treatment failures at on-site visits. For home visits, the investigator calls the patient every 2 weeks until CR on minimal therapy to confirm the patient is still in CR and to determine the prednisone tapering schedule.

[0333] Additionally, the clinical activity of pemphigus is assessed using the PDAI.

Efficacy Endpoints

[0334]

- Proportion of patients who achieve CR on minimal therapy in patients with PV
- Proportion of patients who achieve CR on minimal prednisone dose in patients with PV and PF
- Time to DC
- Time to complete clinical remission (CR)
- Time to CR on minimal therapy
- Time to CR off therapy
- Time to flare
- Rate of treatment failure
- Rate of flare
- Cumulative prednisone dose over the trial
- PDAI at each visit

Pharmacokinetics

[0335]

- Efgartigimod serum concentrations

Pharmacodynamics

[0336]

- Total IgG and subtype (IgG1, IgG2, IgG3, IgG4) serum levels
- Anti-Dsg-1 and -3 autoantibody serum levels

Immunogenicity

[0337] ADAs to efgartigimod PH20 SC (both to efgartigimod and to rHuPH20)

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US8216805B [0087] [0149]
- US20090252729A1 [0087] [0149]
- US20110081345A1 [0087] [0149]
- WO2019118791A [0096]
- WO2014019727A [0097]
- US8067232B [0105] [0151]
- US8021856B [0106] [0151]
- US20110243966A1 [0138]
- US5624821A [0146] [0147] [0148] [0148]
- US6277375B [0146] [0147] [0148] [0148]
- US6737056B [0146] [0147] [0148]
- WO0158957A [0146] [0147]
- WO0206919A [0146] [0147]
- WO04016750A [0146] [0147]
- WO04029207A [0146] [0147] [0148]
- WO04035752A [0146] [0147]
- WO05040217A [0146] [0147]
- US5885573A [0148]
- US5677425A [0148]
- US6165745A [0148]
- US5869046A [0148]
- US6121022A [0148]
- US5648260A [0148]
- US6528624B [0148]
- US6194551B [0148]

- US6821505B [0148]
- US20040002587A [0148]
- WO9429351A [0148]
- WO9958572A [0148]
- WO0042072A [0148]
- WO02060919A [0148]
- WO04099249A [0148]
- WO04063351A [0148]
- US20040010124A1 [0150]
- US20040047862A1 [0150]
- US20040265321A1 [0150]
- US10316073B [0157] [0159]

Non-patent literature cited in the description

- **CROFF. JOLY P. et al.**Lancet, 2017, vol. 389, 2031-2040 [0002]
- **ULRICHTS et al.**Neonatal Fc receptor antagonist efgartigimod safely and sustainably reduces IgG in humansJ. Clin. Investig., 2018, vol. 128, 104372-4386 [0003]
- **JAMES et al.**Randomized phase 2 study of FcRn antagonist efgartigimod in generalized myasthenia gravisNeurology, 2019, vol. 92, 232661-2673 [0004]
- **BALGHI et al.**Comparing early and late treatments with rituximab in permphigues vulgaris; which one is better?Arch. Dermatol. Res., 2018, vol. 311, 163-69 [0005]
- **EDELMAN, G.M. et al.**Proc. Natl. Acad. USA, 1969, vol. 63, 78-85 [0090]
- **KABAT et al.**Sequences of Proteins of Immunological InterestU.S. Dept. Health and Human Services19910000 [0090]
- **ROUX et al.**J. Immunol., 1998, vol. 161, 4083- [0092]
- **DRICKAMER KTAYLOR ME**Introduction to Glycobiology20060000 [0104]
- **BELLONI-FORTINA A et al.**Clin Dev Immunol., 2009, 187864- [0110]
- **ABASQ C et al.**Arch Dermatol., 2009, vol. 145, 5529-35 [0110]
- **KASPERKIEWICZ**Nat. Re. Dis. Primers, 2017, vol. 3, 17026- [0112]
- **HERTL M et al.**J EurAcad Dermatol Venereol., 2015, vol. 29, 3405-14 [0114]
- **REN Z et al.**J Eur Acad Dermatol Venereol., 2018, vol. 32, 101768-1776 [0115] [0123] [0131]
- **MURRELL et al.**J Am Acad Dermatol., 2008, vol. 58, 1043- [0119]
- **MURRELL et al.**J Am Acad Dermatol., 2018, [0119]
- **HARMAN et al.**Br J Dermatol., 2017, vol. 177, 51170-1201 [0119]
- **JOLY et al.**Lancet, 2017, vol. 389, 100832031-2040 [0119] [0131]
- **HÉBERT et al.**J Invest Dermatol., 2019, vol. 139, 131-37 [0119]
- **ROSENBACH M et al.**J Invest Dermatol., 2009, vol. 129, 102404-10 [0120]
- **RAHBAR Z et al.**JAMA Dermatol., 2014, vol. 150, 3266-72 [0120]

- **SHIMIZU et al.**J Dermatol., 2014, vol. 41, 11969-973 [0121]
- **BOULARD et al.**Br J Dennatol., 2016, vol. 175, 1142-9 [0121]
- **MURRELL et al.**J Am Acad Dermatol., 2020, vol. 82, 3575-585 [0121]
- **SEBARATNAM DF et al.**JAMA Dermatol., 2013, vol. 149, 101186-91 [0122]
- **RATNAM KV et al.**Int J Dermatol., 1990, vol. 29, 5363-7 [0123]
- **CZERNIK A et al.**Arch Dermatol., 2008, vol. 144, 5658-61 [0123]
- **CHAIDEMENOS G et al.**J Eur Acad Dermatol Venereol., 2011, vol. 25, 2206-10 [0123]
- **BEISSERT S et al.**J Invest Dermatol., 2010, vol. 130, 82041-8 [0128]
- **MIMOUNI D et al.**J Eur Acad Dermatol Venereol., 2010, vol. 24, 8947-52 [0128]
- **WANG HH et al.**Acta Derm Venereol., 2015, vol. 95, 8928-32 [0131] [0132]
- **HÉBERT V et al.**J Invest Dermatol., 2019, vol. 139, 131-37 [0132]
- **CHALLA DK et al.**Curr Top Microbiol Immunol., 2014, vol. 382, 249-72 [0134]
- **ROOPENIAN DC et al.**Nat Rev Immunol., 2007, vol. 7, 9715-25 [0134]
- **ROOPENIAN DC et al.**J Immunol., 2003, vol. 170, 73528-33 [0134]
- **WALDMANN TA et al.**J Clin Invest., 1990, vol. 86, 62093-2098 [0134]
- **WANI MA et al.**Proc Natl Acad Sci U S A., 2006, vol. 103, 135084-5089 [0134]
- **GHETIE et al.**Nat. Biotech., 1997, vol. 15, 637-40 [0148]
- **DUNCAN et al.**Nature, 1988, vol. 332, 563-564 [0148]
- **LUND et al.**J. Immunol., 1991, vol. 147, 2657-2662 [0148]
- **LUND et al.**Mol. Immunol., 1992, vol. 29, 53-59 [0148]
- **ALEGRE et al.**Transplantation, 1994, vol. 57, 1537-1543 [0148]
- **HUTCHINS et al.**Proc Natl. Acad Sci USA, 1995, vol. 92, 11980-11984 [0148]
- **JEFFERIS et al.**Immunol Lett., 1995, vol. 44, 111-117 [0148]
- **LUND et al.**FASEB J., 1995, vol. 9, 115-119 [0148]
- **JEFFERIS et al.**Immunol Lett., 1996, vol. 54, 101-104 [0148]
- **LUND et al.**J. Immunol., 1996, vol. 157, 4963-4969 [0148]
- **ARMOUR et al.**Eur J Immunol, 1999, vol. 29, 2613-2624 [0148]
- **IDUSOGIE et al.**J. Immunol., 2000, vol. 164, 4178-4184 [0148]
- **REDDY et al.**J. Immunol., 2000, vol. 164, 1925-1933 [0148]
- **XU et al.**Cell Immunol., 2000, vol. 200, 16-26 [0148]
- **IDUSOGIE et al.**J. Immunol., 2001, vol. 166, 2571-2575 [0148]
- **SHIELDS et al.**J Biol. Chem., 2001, vol. 276, 6591-6604 [0148]
- **JEFFERIS et al.**Immunol Lett., 2002, vol. 82, 57-65 [0148]
- **PRESTA et al.**Biochem Soc Trans., 2002, vol. 30, 487-490 [0148]
- **CLARKSON et al.**New Engl J Med, 2013, vol. 314, 91236-1239 [0150]
- **VACCARO C et al.**Nat Biotechnol., 2005, vol. 23, 101283- [0159]
- **SHPILBERG O et al.**Br J Cancer., 2013, vol. 109, 61556-1561 [0166] [0166]
- **ROSENBACH, M et al.**J Invest Dermatol, 2009, vol. 129, 102404-10 [0214]
- **ULRICHTS, P. et al.**Neonatal Fc receptor antagonist efgartigimod safely and sustainably reduces IgGs in humansJ Clin Invest, 2018, vol. 128, 104372-4386 [0264]
- **HOWARD, J.F., JR. et al.**Randomized phase 2 study of FcRn antagonist efgartigimod in generalized myasthenia gravisNeurology, 2019, vol. 92, 23e2661-e2673 [0264]
- **NEWLAND, A.C. et al.**Phase 2 study of efgartigimod, a novel FcRn antagonist, in adult patients with primary immune thrombocytopeniaAm J Hematol, 2020, vol. 95, 2178-

187 [0264]

- **GRANDO, SA et al.**Int Immunopharmacol, 2020, vol. 82, 106385- [0274]
- **AMAGAI, M et al.**J Am Acad Dermatol, 2009, vol. 60, 4595-603 [0274]
- **LI, N et al.**J Clin Invest, 2005, vol. 115, 123440-50 [0274]
- **CAUZA, K et al.**J Invest Dermatol, 2005, vol. 124, 1132-9 [0274]
- **NGUYEN, VT et al.**J Biol Chem, 2004, vol. 279, 32135-46 [0276]
- **COLLIOU, N et al.**Sci Transl Med, 2013, vol. 5, 175175-30 [0277]

Krav:

1. FcRn-antagonist til anvendelse i behandlingen af pemphigus hos et individ med behov derfor, hvori
   (i) individet har
      (a) nyligt diagnosticeret pemphigus,
      (b) recidiverende pemphigus, eller
      (c) refraktær pemphigus,
   (ii) FcRn-antagonisten består af en varierende Fc-region, hvori nævnte varierende Fc-region består af to Fc-domæner, som danner en homodimer, og hvori aminosyresekvensen af hvert af Fc-domænerne er valgt fra gruppen bestående af SEQ ID NO: 1, SEQ ID NO: 2 og SEQ ID NO: 3, og
   (iii) FcRn-antagonisten administreres i kombination med et kortikosteroid, og kortikosteroidet er i en dosis på ≤ 0,5 mg prednison/kg/dag eller tilsvarende.

2. FcRn-antagonist til anvendelse ifølge krav 1, hvori pemphigus omfatter pemphigus vulgaris (PV), pemphigus foliaceus (PF), eller både PV og PF, og/eller hvori individet har mild, moderat eller svær pemphigus som klassificeret ved Pemphigus Disease Area Index (PDAI).

3. FcRn-antagonist til anvendelse ifølge krav 1 eller krav 2, hvori FcRn-antagonisten administreres én gang om ugen eller hver anden uge, indtil sygdomskontrol eller fuldstændig remission er opnået.

4. FcRn-antagonist til anvendelse ifølge ethvert af kravene 1 – 3, hvori FcRn-antagonisten administreres intravenøst i en dosis på 10 mg/kg til 30 mg/kg én gang om ugen eller hver anden uge,
   valgfrit hvori FcRn-antagonisten administreres intravenøst i en dosis på 10 mg/kg én gang om ugen eller hver anden uge,
   eller valgfrit hvori FcRn-antagonisten administreres intravenøst i en dosis på 25 mg/kg én gang om ugen eller hver anden uge.

5. FcRn-antagonist til anvendelse ifølge ethvert af kravene 1 til 3, hvori FcRn-antagonisten administreres subkutant i en fast dosis på 750 mg til 3000 mg én gang om ugen eller hver anden uge,
   valgfrit hvori FcRn-antagonisten administreres subkutant i en fast dosis på 1000 mg eller 2000 mg én gang om ugen eller hver anden uge,

eller valgfrit hvori FcRn-antagonisten først administreres subkutant i en fast dosis på ca. 1000 mg to gange på den samme dag.

6. FcRn-antagonist til anvendelse ifølge krav 5, hvori FcRn-antagonisten er formuleret sammen med hyaluronidase og administreret subkutant.

7. FcRn-antagonist til anvendelse ifølge ethvert af kravene 1 – 6, yderligere omfattende administration af en nedtrappende dosis af kortikosteroid efter sygdomskontrol eller fuldstændig remission er blevet opnået.

8. FcRn-antagonist til anvendelse ifølge ethvert af kravene 1 – 7, hvori den nedtrappende dosis af kortikosteroid begynder ved ≤ 0,5 mg prednison/kg/dag eller ækvivalent, eller hvori den nedtrappende dosis af kortikosteroid begynder ved en dosis på 10 mg prednison/dag eller ækvivalent.

9. FcRn-antagonist til anvendelse ifølge ethvert af kravene 1 – 8, hvori nedtrapningen af kortikosteroid initieres inden for ca. 1 måned, inden for ca. 3 uger, inden for ca. 2 uger eller inden for ca. 1 uge efter administration af den initiale dosis af FcRn-antagonist.

10. FcRn-antagonist til anvendelse ifølge ethvert af kravene 1 – 9, hvori en reduktion af den initiale eller efterfølgende nedtrapningsdosis kan forekomme hver måned, hver 3. uge, hver 2. uge eller hver uge.

11. FcRn-antagonist til anvendelse ifølge ethvert af kravene 1 – 10, hvori nedtrapningen af kortikosteroider omfatter en reduktion på ca. 0,5 mg prednison/kg/dag eller ækvivalent, ca. 0,25 mg prednison/kg/dag eller ækvivalent, eller ca. 0,1 mg prednison/kg/dag eller ækvivalent.

12. FcRn-antagonist til anvendelse ifølge ethvert af kravene 1 - 11, hvori kortikosteroidet er et systemisk kortikosteroid.

13. FcRn-antagonist til anvendelse ifølge ethvert af kravene 1 - 12, hvori FcRn-antagonisten administreres i en induktionsfase og en konsolideringsfase, hvori
(i) FcRn-antagonisten under induktionsfasen administreres én gang om ugen eller hver anden uge, og et kortikosteroid administreres i en dosis på ≤ 0,5 mg prednison/kg/dag eller ækvivalent indtil sygdommen er under kontrol, og

(ii) FcRn-antagonistdosen under konsolideringsfasen reduceres eller FcRn-antagonistdoseringsintervallet forlænges, og/eller kortikosteroiddosen nedsættes eller kortikosteroiddoseringsintervallet forlænges, til en slut-på-konsolidering-dosis eller et doseringsinterval, som er effektivt til at forhindre nye læsioner i at dukke op.

14. FcRn-antagonist til anvendelse ifølge krav 13, yderligere omfattende en vedligeholdelsesfase, hvori
(iii) slut-på-konsolidering-dosen eller doseringsintervallet for FcRn-antagonisten og/eller prednisonen eller ækvivalent fortsættes under vedligeholdelsesfasen indtil fuldstændig fjernelse af læsioner.

15. FcRn-antagonist til anvendelse ifølge krav 13 eller krav 14, hvori FcRn-antagonisten under induktionsfasen er:
(i) administreret intravenøst i en dosis på 10 mg/kg til 30 mg/kg,
(ii) administreret intravenøst i en dosis på 10 mg/kg,
(iii) administreret intravenøst i en dosis på 25 mg/kg,
(iv) administreret subkutant i en fast dosis på 750 mg til 3000 mg, eller
(v) først administreret subkutant i en fast dosis på ca. 1000 mg to gange dagligt.

16. FcRn-antagonist til anvendelse ifølge ethvert af kravene 1 – 15, hvori individet:
(i) har refraktær pemphigus,
(ii) har nyligt diagnosticeret pemphigus vulgaris med en Pemphigus Disease Area Index (PDAI) score på ≥ 15,
(iii) har recidiverende pemphigus vulgaris,
(iv) har nyligt diagnosticeret pemphigus foliaceus med en Pemphigus Disease Area Index (PDAI) score på ≥ 15,
(v) har recidiverende pemphigus foliaceus,
(vi) er rituximab-refraktær og/eller
(vii) er kortikosteroid-intolerant.

17. FcRn-antagonist til anvendelse ifølge ethvert af kravene 1 – 16, yderligere omfattende administration af et B-celle-depleterende middel til individet, valgfrit hvori det B-celledepleterende middel er rituximab.

DRAWINGS

Drawing



Fig. 1


53 patients for eligibility
18 patients screen failure
1 patient consent withdrawal before first dosing
Safety analysis set
(N = 34)
Cohort 1 (N = 6)
Cohort 2 (N = 5)
Cohort 3 (N = 8)
Cohort 4 (N = 15)
Non-evaluable for efficacy
(N = 3)
1 patient: insufficient exposure to IMP (cohort 1)
1 patient: confounding factor (cohort 1)
1 patient: violation of the inclusion criteria of the study (cohort 3)
Efficacy analysis set
(N = 31)
Cohort 1
(N = 4)
Cohort 2
(N = 5)
Cohort 3
(N = 7)
Cohort 4
(N = 15)

Fig. 2


PDAI activity
IgG4
Dsg-1
Dsg-3
% from baseline
100
50
0
BL
W4
W8
W12
W16
W20
W24
Weekly/biweekly maintenance dosing

Fig. 3



Fig. 4A



Fig. 4B



Fig. 4C



Fig. 4D



Fig. 5A



Fig. 5B



Fig. 5C



Fig. 5D



Fig. 6A



Fig. 6B



Fig. 6C



Fig. 6D



Fig. 6E


Total IgG
% from baseline, mean (SD)
150
100
50
0
0
50
100
150
Study days
Cohort 1
Cohort 2
Cohort 3
Cohort 4

Fig. 7A


PDAI activity
PDAI activity score, mean
25
20
15
10
5
0
0
50
100
150
Study day
Cohort 1
Cohort 2
Cohort 3
Cohort 4

Fig. 7B


150
100
50
0
Total IgG (% T_0 ± SE)
0 7 14 21 28 35 42 49 56
Study day
PV 25 mg/kg
ARGX-113-1701 (n=4)

Fig. 8A


150
100
50
0
Total IgG (% T_0 ± SE)
0 7 14 21 28 35 42 49 56 63 70 77 84 91
Study day
HV 25 mg/kg ARGX-113-1501
PV 25 mg/kg ARGX-113-1701

Fig. 8B


150
100
50
0
Total IgG (% T_0 ± SE)
0 7 14 21 28 35 42 49 56
Study day
PV 10 mg/kg
(n=13)

Fig. 9A


150
100
50
0
Total IgG (% T_0 ± SE)
0 7 14 21 28 35 42 49 56 63 70 77 84 91
Study day
HV 10 mg/kg
PV 10 mg/kg
MG 10 mg/kg
ITP 10 mg/kg

Fig. 9B


Cohort 1 (N=4)
Cohort 2 (N=5)
Cohort 3 (N=7)
Mean Total IgG (mg/L)
20000
15000
10000
5000
0
0
50
100
150
Study Day

Fig. 10A


Mean Total IgG (mg/L))
12000
10000
8000
6000
4000
0
28
56
84
112
140
168
196
224
252
280
308
336
Time (Days)

Fig. 10B


Cohort 1 (N=4)
Cohort 2 (N=5)
Cohort 3 (N=7)
Mean IgG1 (mg/L)
12500
10000
7500
5000
2500
0
0
50
100
150
Study Day

Fig. 11A


Cohort 1 (N=4)
Cohort 2 (N=5)
Cohort 3 (N=7)
Mean IgG2 (mg/L)
10000
8000
6000
4000
2000
0
0
50
100
150
Study Day

Fig. 11B


Cohort 1 (N=4)
Cohort 2 (N=5)
Cohort 3 (N=7)
Mean IgG3 (mg/L)
1500
1250
1000
750
500
250
0
0
50
100
150
Study Day

Fig. 11C


Cohort 1 (N=4)
Cohort 2 (N=5)
Cohort 3 (N=7)
Mean IgG4 (mg/L)
2000
1500
1000
500
0
0
50
100
150
Study Day

Fig. 11D



Fig. 11F



Fig. 11E



Fig. 11H



Fig. 11G

Fig. 12A



Fig. 12B



Fig. 12C



Fig. 12D





Fig. 13

DK/EP 4087875 T3


Main study: 30 weeks
Open-label extension: 60 weeks
Status in 1904
CRmin
CR
DC, no CR
Trt Fail,
Flare after CRmin
Flare after CRmin in 1905
Observation every 4 weeks as long as CRmin or CRoff is sustained
Observation every 4 weeks
Observation every 4 weeks
Observation every 4 weeks
W1 W4 W8 W12 W16 W20 W24 W28 W32 W36 W40 W44 W48 W52 W56 W60
EOT
FU
Efgartigimod PH20 SC weekly
Placebo SC weekly
Efg 2000 mg
Efg 1000 mg
on-site
home

Fig. 14

SEKVENSLISTE

Sekvenslisten er udeladt af skriftet og kan hentes fra det Europæiske Patent Register.

The Sequence Listing was omitted from the document and can be downloaded from the European Patent Register.