

**(19) AUSTRALIAN PATENT OFFICE**

(54) Title  
Novel triazolo pyrimidine compounds

(51)<sup>6</sup> International Patent Classification(s)  
C07D 239/47 61/00  
(2006.01) 20060101ALI2010021  
C07B 61/00 (2006.01) 1BMAU C07D  
C07D 239/56 239/56  
(2006.01) 20060101ALI2010021  
C07D 317/44 1BMAU C07D  
(2006.01) 317/44  
C07D 405/12 20060101ALI2010021  
(2006.01) 1BMAU C07D  
C07D 487/04 405/12  
(2006.01) 20060101ALI2010021  
C07D 239/47 1BMAU C07D  
20060101AFI2010021 487/04  
1BMAU C07B 20060101ALI2010021  
1BMAU

(21) Application No: 2010200461

(22) Application Date: 2010.02.09

(43) Publication Date : 2010.02.25

(43) Publication Journal Date : 2010.02.25

(62) Divisional of:  
2007200776

(71) Applicant(s)  
AstraZeneca AB


(72) Inventor(s)  
Larsson, Ulf; Magnusson, Mattias; Musil, Tibor; Palmgren, Andreas

(74) Agent/Attorney  
Davies Collison Cave, 1 Nicholson Street, Melbourne, VIC, 3000

**ABSTRACT**

The present invention relates to a pyrimidine compound (I) useful as a pharmaceutical intermediate, to a process for preparing said pyrimidine compound, to intermediates used in said process, and to the use of said pyrimidine compound in the preparation of pharmaceuticals.

5



2010200461 09 Feb 2010

*Australian Patents Act 1990 – Regulation 3.2A*

**ORIGINAL COMPLETE SPECIFICATION  
STANDARD PATENT**

Invention Title

"Novel triazolo pyrimidine compounds"


The following statement is a full description of this invention, including the best method of performing it known to us:-

**NOVEL TRIAZOLO PYRIMIDINE COMPOUNDS**


This application is a divisional of Australian Patent Application No. 2007200776, the entire contents of which is incorporated herein by reference.

The present invention relates to a pyrimidine compound useful as a pharmaceutical intermediate, to a process for preparing said pyrimidine compound, to intermediates used in said process, and to the use of said pyrimidine compound in the preparation of pharmaceuticals.


The present invention provides a compound of formula (I):



The present invention also provides a process for preparing a compound of formula (I), comprising reacting a compound of formula (II):




with a salt of a compound of formula (III):



in the presence of a suitable base (such as an alkali metal hydroxide (such as sodium or potassium hydroxide) a tertiary amine (such as a tri(C<sub>1-6</sub> alkyl)amine, for example triethylamine)), a suitable solvent (such as an alcohol, such as an aliphatic alcohol containing from 1 to 6 carbon atoms, for example ethanol), preferably at a temperature in the range 100-150°C and, where necessary (for example when the temperature exceeds the boiling point of the solvent), in a sealed system under autogenic pressure.

A suitable salt of a compound of formula (III) is a salt of a mineral or organic acid. Suitable mineral acids include hydrochloric, hydrobromic, hydroiodic, nitric or sulphuric acid. A suitable organic acid is, for example, an organic achiral acid such as acetic, trifluoroacetic, oxalic or p-toluenesulphonic acid, or an organic chiral acid such as L-tartaric acid, dibenzoyl-L-tartaric acid or di-p-toluoyl-L-tartaric acid.


In another aspect the present invention provides a process for preparing a compound of formula (I) comprising hydrogenating a compound of formula (IV):



wherein Ar is phenyl optionally substituted by halogen, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy; to give a compound of formula (II), and reacting the compound of formula (II) with a compound of formula (III) (as described above) to provide the compound of formula (I).

The hydrogenation is preferably conducted using a heavy metal catalyst (such as platinum on carbon), in a suitable solvent (such as a C<sub>1-6</sub> aliphatic alcohol, for example 2-propanol (iso-propanol)), at a suitable temperature (such as 10-70°C, for example 20-50°C) and at a suitable pressure (such as 1-5 bar, for example about 3 bar).


A compound of formula (IV) can be prepared by chlorinating a compound of formula (VIII):



wherein Ar is as defined above, with a suitable chlorinating agent (such as phosphorus oxychloride) in the presence of a suitable nitrogen containing base (such as triethylamine, especially pyridine) and at a suitable temperature (such as in the range 50°C to the boiling point of phosphorus oxychloride; for example 70 to 90°C). A compound of formula (VIII) can be prepared by routine adaptation of literature methods.


In a further aspect the present invention provides a process, as hereinbefore described, for the preparation of a compound of formula (II).

The compound of formula (I) can be used to prepare the pharmaceutical compound of formula (A):




as described below.

- 5 Thus, a compound of formula (A) can be prepared by deprotecting a compound of formula (V):



for example using a strong mineral acid (such as hydrochloric acid) in a suitable solvent (such as methanol or ethanol).

- 10 A compound of formula (V) can be prepared by coupling a compound of formula (VI) {or a salt thereof (such as a mandelate salt) from which the compound of formula (VI) is generated *in situ*}, with a compound of formula (VII):




for example in the presence of a suitable base (such as a tertiary amine, such as a tri(C<sub>1-6</sub> alkyl)amine, for example triethylamine) and a suitable solvent (for example a polar solvent, such as an alcohol (such as an aliphatic alcohol containing from 1 to 6 carbon atoms, for example ethanol) or a nitrile (such as acetonitrile)) and at a suitable temperature (such as a temperature in the range 10-40°C, for example ambient temperature).

A compound of formula (VII) can be prepared by reacting a compound of formula (I) with an alkali metal nitrite (such as NaNO<sub>2</sub>) or an organic nitrite (for example *iso*-amyl nitrite) in the presence of a suitable acid (such as acetic acid) and a suitable solvent (such as water or a mixture of water and acetic acid) and at a suitable temperature (such as a temperature in the range -10 to 15°C, for example -10 to 10°C).

Thus, in a further aspect the present invention provides the use of the compound of formula (I) in a process for the preparation of the compound of formula (A).

A salt of a compound of formula (III) can be prepared by reacting a compound of formula (III) with the necessary acid in a suitable solvent (such as water, an aliphatic alcohol containing 1 to 4 carbon atoms (for example ethanol) or a simple ester (such as ethyl acetate)) at a suitable temperature (such as from 10 to 60°C, for example 30 to 50°C).


A compound of formula (III) can be prepared by deprotecting a compound of formula (IX):



for example by hydrogenation {such as with a heavy metal catalyst (such as palladium on carbon) in the presence of a solvent (such as an aliphatic alcohol containing 1 to 4 carbon


atoms, for example ethanol) at ambient temperature at a suitable pressure (such as 1 to 3 bar, for example 1.0 to 1.5 bar).

A compound of formula (IX) can be prepared by reducing a compound of formula (X):




wherein R\* is C<sub>1-4</sub> alkyl (preferably ethyl), such as with a suitable borohydride (for example an alkali metal borohydride, such as lithium borohydride), lithium aluminiumhydride or DIBAL-H in a suitable polar solvent (such as tetrahydrofuran).

A compound of formula (X) can be prepared by reacting a compound of formula (XI):



with a suitable compound L-CH<sub>2</sub>CO<sub>2</sub>R\* {wherein R\* is C<sub>1-4</sub> alkyl (especially ethyl); and L is a leaving group, especially halogen (for example bromo)}, in the presence of a suitable polar solvent (such as tetrahydrofuran) and in the presence of a suitable base (such as potassium *tert*-butoxide, sodium hydride or a C<sub>1-6</sub> alkyl lithium species).

A compound of formula (XI) can be prepared by reacting a compound of formula (XII):



with benzyl chloroformate in the presence of a suitable base (such as potassium carbonate) and a suitable solvent (such as a ketone (for example 4-methyl-2-pentanone) or a hydrocarbon (for example toluene)).



In a still further aspect the present invention provides a process for the preparation of a salt of a compound of formula (III) as hereinbefore described.

In further aspects the present invention provides an intermediate compound of formulae (II), (IV), (VII), (VIII), (X) or (XI), or a salt of a compound of formula (III).

The following Examples illustrate the invention.

#### EXAMPLE 1

This Example illustrates the preparation of 2-([(3 $\alpha$ R,4S,6R,6 $\alpha$ S)-6-amino-2,2-dimethyltetrahydro-3 $\alpha$ H-cyclopenta[*d*][1,3]-dioxol-4-yl]oxy)-1-ethanol, L-tartaric acid salt (1:1).

**Step a:** Preparation of [3 $\alpha$ S-(3 $\alpha\alpha$ ,4 $\alpha$ ,6 $\alpha$ ,6 $\alpha\alpha$ )]-[tetrahydro-6-hydroxy-2,2-dimethyl-4H-cyclopenta-1,3-dioxol-4-yl]-carbamic acid, phenylmethyl ester.

Potassium carbonate (39.3g) was added to a suspension of [3 $\alpha$ R-(3 $\alpha\alpha$ ,4 $\alpha$ ,6 $\alpha$ ,6 $\alpha\alpha$ )]-6-amino-tetrahydro-2,2-dimethyl-4H-cyclopenta-1,3-dioxol-4-ol, hydrochloride, (prepared as described in WO 9905142) (27.1g) in 4-methyl-2-pentanone (500ml). Water (150ml) was then added followed by dropwise addition of benzyl chloroformate (23.1g). The reaction mixture was stirred at room temperature for 4 hours before the organic phase was separated. The aqueous phase was extracted with 4-methyl-2-pentanone (2x50ml). The combined organics were concentrated and the residue was purified (SiO<sub>2</sub>, dichloromethane:methanol, 95:5 to 90:10 as eluant) to give the subtitle compound (39.23g).

<sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.32 (5H, m), 5.65 (1H, br s), 5.10 (2H, br s), 4.59 (1H, d), 4.48 (1H, d), 4.27 (1H, m), 4.19 (1H, br m), 2.24 (1H, br s), 1.69 (1H, d), 1.41 (3H, s), 1.26 (3H, s).

**Step b:** Preparation of [3 $\alpha$ S-(3 $\alpha\alpha$ ,4 $\alpha$ ,6 $\alpha$ ,6 $\alpha\alpha$ )]-[2,2-dimethyl-6-(2-hydroxyethoxy)-tetrahydro-4H-cyclopenta-1,3-dioxol-4-yl]-carbamic acid, phenylmethyl ester.

Potassium *tert*-butoxide (3.6g) in tetrahydrofuran (20ml) was added over 5 minutes to a solution of the product from Step (a) (39.23g) in tetrahydrofuran (200ml). After 15 minutes, ethyl bromoacetate (3.7ml) in tetrahydrofuran (10ml) was added dropwise. The mixture was stirred at 0°C for 10 minutes, then further ethyl bromoacetate was added (3.7ml x4). The reaction mixture was stirred at 0°C a further 2 hours.

Lithium borohydride (2.79g) was then added portionwise to the reaction mixture which was then stirred at  $<5^{\circ}\text{C}$  for 16 hours. Glacial acetic acid (23g) was added dropwise to the cold mixture. After stirring for 30 minutes, water (100ml) was added dropwise and the resulting mixture was stirred for 30 minutes. The phases were then separated and the aqueous phase was extracted with ethyl acetate. The combined organics were washed with saturated sodium bicarbonate and brine, dried and concentrated. The residue was purified (SiO<sub>2</sub>, ethyl acetate:hexane, 25:75 to 50:50 as eluant) to give the subtitle compound (38.6g).

MS (APCI) 218 (M+H<sup>+</sup>, 100%).

Step c: Preparation of [3aR-(3a $\alpha$ ,4 $\alpha$ ,6 $\alpha$ ,6a $\alpha$ )]-2-[[6-amino-2,2-dimethyl-tetrahydro-4H-cyclopenta-1,3-dioxol-4-yl]oxy]-ethanol (alternatively named: 2-[[[(3aR,4S,6R,6aS)-6-amino-2,2-dimethyltetrahydro-3aH-cyclopenta[d][1,3]-dioxol-4-yl]oxy]-1-ethanol).

A slurry of 5% palladium on charcoal (4g) in ethanol was added to a solution of the product from Step (b) (39.96g) in ethanol (250ml) and the mixture was hydrogenated at 1.2 bar for 20 hours. The catalyst was filtered off and the filtrate was concentrated to give the subtitle compound (23.65g).

MS (APCI) 160 (M+H<sup>+</sup>, 100%).

Step d: Preparation of [3aR-(3a $\alpha$ ,4 $\alpha$ ,6 $\alpha$ ,6a $\alpha$ )]-2-[[6-amino-2,2-dimethyl-tetrahydro-4H-cyclopenta-1,3-dioxol-4-yl]oxy]-ethanol L tartrate (alternatively named: 2-[[[(3aR,4S,6R,6aS)-6-amino-2,2-dimethyltetrahydro-3aH-cyclopenta[d][1,3]-dioxol-4-yl]oxy]-1-ethanol, L-tartaric acid salt (1:1)).

A stirred solution of the product obtained in Step (c) (545g) in ethanol (3:8l) was heated to 35°C. L-tartaric acid (352g) was added (temperature rise to 45°C) and the mixture was stirred at 40 - 45°C for 1 h. The mixture was cooled to 20°C and the resulting thick slurry stirred for 16 h then filtered. The collected solid was washed with two portions of 2-propanol (300ml, then 500ml), sucked dry then dried *in vacuo* at 40°C to give the product as white crystals (728g).

## EXAMPLE 2

This Example illustrates the preparation of *trans*-(1*R*,2*S*)-2-(3,4-difluorophenyl)-cyclopropylamine, *R*-mandelate salt (alternatively named *trans*-(1*R*,2*S*)-2-(3,4-difluorophenyl)cyclopropanaminium (2*R*)-2-hydroxy-2-phenylethanoate.

5 Step 1: Preparation of (*E*)-3-(3,4-difluorophenyl)-2-propenoic acid.

A stirred mixture of pyridine (15.5kg) and piperidine (0.72kg) were heated to 90°C. Malonic acid (17.6kg) was added, followed by slow addition, over 50 minutes, of 3,4-difluorobenzaldehyde (12.0kg). The reaction mixture was stirred at 90°C for a further 4 hours and 36 minutes. Water (58.5kg) was added and 32 litres of the pyridine/water  
10 mixture then was distilled out of the reactor under reduced pressure. The reaction mixture was acidified to pH 1 with 37% hydrochloric acid (6.4kg) over a 40 minute period, then cooled to 25°C with strong stirring. The solids were collected by filtration, washed twice with 1% hydrochloric acid (34.8 L per wash), once with water (61 L) and then deliquored thoroughly in the filter. The product was then dried under vacuum at 40°C for 24 hours and  
15 40 minutes, affording 13.7kg of the crystalline product.

Step 2: Preparation of (*E*)-3-(3,4-difluorophenyl)-2-propenoyl chloride.

A stirred mixture of (*E*)-3-(3,4-difluorophenyl)-2-propenoic acid (8.2 kg), toluene (7.4kg) and pyridine (0.18kg) was heated to 65°C and then thionyl chloride (7.4kg) was  
20 added over 30 minutes. The reaction was stirred for a further 2h 15 minutes after the addition was complete, then diluted with toluene (8.7kg). Excess thionyl chloride, sulfur dioxide and hydrogen chloride were then distilled out, together with toluene (10 L), under reduced pressure, yielding a solution of the (*E*)-3-(3,4-difluorophenyl)-2-propenoyl chloride (approximately 9kg) in toluene.

25 Step 3: Preparation of (1*R*,2*S*,5*R*)-2-isopropyl-5-methylcyclohexyl (*E*)-3-(3,4-difluorophenyl)-2-propenoate.

A solution of *L*-menthol (7.1kg) in toluene (8.5kg) was added over a 20 minute period to the solution of (*E*)-3-(3,4-difluorophenyl)-2-propenoyl chloride (prepared as in  
30 Step 2) and pyridine (0.18kg, 2.28 mol) stirring at 65°C. The reaction mixture was stirred at 65°C for a further 4 hours and 40 minutes after the addition was complete, then cooled

to 25°C and stirred for a 14 hours. The solution was diluted with toluene (16kg), washed with 5% aqueous sodium chloride (6.4kg), then 6% sodium hydrogen carbonate (6.47kg), then water (6.1kg). The solution was dried azeotropically by distillation of the solvent (20 L) under reduced pressure. Dimethyl sulfoxide (33.9kg) was added and the remaining  
5 toluene was distilled off under reduced pressure, affording 47.3kg of a solution of (1*R*,2*S*,5*R*)-2-isopropyl-5-methylcyclohexyl (*E*)-3-(3,4-difluorophenyl)-2-propenoate (approx. 13.3kg) in dimethyl sulfoxide.

Step 4: Preparation of dimethylsulfoxonium methylide (dimethyl(methylene)oxo- $\lambda^6$ -  
10 sulfane.

Sodium hydroxide powder (1.2kg), prepared by milling sodium hydroxide pellets in a rotary mill through a 1mm metal sieve, and trimethylsulfoxonium iodide (6.2kg) were stirred in dimethyl sulfoxide (25.2kg) under a nitrogen atmosphere at 25°C for 90 min. The solution was used directly in the preparation of (1*R*, 2*S*, 5*R*)-2-isopropyl-5-  
15 methylcyclohexyl *trans*-2-(3,4-difluorophenyl)cyclopropanecarboxylate.

Step 5: Preparation of (1*R*, 2*S*, 5*R*)-2-isopropyl-5-methylcyclohexyl *trans*-2-(3,4-difluorophenyl)cyclopropanecarboxylate.

A solution of (1*R*,2*S*,5*R*)-2-isopropyl-5-methylcyclohexyl (*E*)-3-(3,4-  
20 difluorophenyl)-2-propenoate (approximately 8.6kg) in dimethyl sulfoxide (approximately 27.9kg) was added with stirring over 20 minutes to a mixture of dimethylsulfoxonium methylide (approximately 2.6kg, prepared as described above), sodium iodide (approximately 4.2kg), water (approximately 500g) and sodium hydroxide (approximately 56g) in dimethylsulfoxide(27.7kg) at 25°C. The reaction mixture was stirred for a further  
25 hours and 50 minutes at 25°C, then used directly for the preparation of (1*R*,2*S*,5*R*)-2-isopropyl-5-methylcyclohexyl *trans*-(1*R*,2*R*)-2-(3,4-difluorophenyl)cyclopropanecarboxylate.

Step 6: Preparation of (1*R*,2*S*,5*R*)-2-isopropyl-5-methylcyclohexyl *trans*-(1*R*,2*R*)-2-(3,4-  
30 difluorophenyl)cyclopropanecarboxylate

A crude solution of (1*R*, 2*S*, 5*R*)-2-isopropyl-5-methylcyclohexyl *trans*-2-(3,4-difluorophenyl)cyclopropanecarboxylate produced as described in step 5 was heated with

stirring from 25°C to 50°C over a 1 hour period and the temperature was maintained for a further hour. The mixture was then cooled with stirring from 50°C to 35°C over 4 hours, kept at 35°C for 1 hour, then cooled to 26°C over 4 hours, kept at 26°C for 1 hour, then cooled to 19°C over 3 hours and kept at 19°C for 5 hours and 10 minutes. The product was collected by filtration, affording a crystalline solid (2.7kg) which was shown to contain a mixture of (1*R*,2*S*,5*R*)-2-isopropyl-5-methylcyclohexyl *trans*-(1*R*,2*R*)-2-(3,4-difluorophenyl)cyclopropanecarboxylate (1.99 kg) and (1*R*,2*S*,5*R*)-2-isopropyl-5-methylcyclohexyl *trans*-(1*S*,2*S*)-2-(3,4-difluorophenyl)cyclopropanecarboxylate (85g).

**Step 7:** Preparation of *trans*-(1*R*, 2*R*)-2-(3,4-difluorophenyl)cyclopropanecarboxylic acid. (1*R*,2*S*,5*R*)-2-isopropyl-5-methylcyclohexyl *trans*-(1*R*,2*R*)-2-(3,4-difluorophenyl)cyclopropanecarboxylate (9.6kg, 91.8% diastereomeric excess) was dissolved in ethanol (13.8kg) and heated with stirring to 46°C. 45% aqueous sodium hydroxide (3.1kg) was added over a 20 minute period and the mixture was stirred for a further 2 hours and 27 minutes. Solvent (28 L) was distilled out of the mixture under reduced pressure, then the mixture was cooled to 24°C and diluted with water (29.3kg), after which the liberated menthol was extracted into toluene (3 washes of 3.3kg each). The remaining aqueous material was acidified to pH 2 with 37% hydrochloric acid (3.3 L) and the product was extracted into toluene (8.6kg, then 2 more washes of 4.2kg and 4.3kg). The combined toluene extracts were washed with 1% hydrochloric acid (4.9 L), then diluted with further toluene (4.2kg) and azeotropically dried by distillation of the solvent (25 L) under reduced pressure. A final dilution with toluene (24.2kg) was followed by distillation of the solvent under reduced pressure (10 L) affording a solution containing *trans*-(1*R*, 2*R*)-2-(3,4-difluorophenyl)cyclopropanecarboxylic acid (approximately 3.45kg) suitable for the production of *trans*-(1*R*, 2*R*)-2-(3,4-difluorophenyl)cyclopropanecarbonyl chloride.

**Step 8:** Preparation of *trans*-(1*R*, 2*R*)-2-(3,4-difluorophenyl)cyclopropanecarbonyl chloride.

Pyridine (70ml) was added to a solution of *trans*-(1*R*, 2*R*)-2-(3,4-difluorophenyl)cyclopropanecarboxylic acid (approximately 3.45kg) in toluene (approximately 12 -15kg), prepared as described above and the mixture was then heated to 65°C. Thionyl chloride (2.3kg) was added over a period of 1 hour and the mixture was

stirred at 70°C for 3 hours. Thionyl chloride (0.5kg) was added and the mixture was stirred a further 2 hours at 70°C. A final aliquot of thionyl chloride (0.5kg) was added and the reaction mixture was stirred for 1 hour at 70°C, then cooled to 40°C. Periodic additions of toluene (45kg, 3 additions of 15kg each) were made during distillation of solvent  
5 (approximately 60 L) from the mixture under reduced pressure, then the solution of *trans*-(1*R*, 2*R*)-2-(3,4-difluorophenyl)cyclopropanecarbonyl chloride (approximately 3.8kg) in toluene (approximately 6-9 L) was cooled to 20°C.

Step 9: Preparation of *trans*-(1*R*, 2*R*)-2-(3,4-difluorophenyl)cyclopropanecarbonyl azide.

10 A solution of *trans*-(1*R*, 2*R*)-2-(3,4-difluorophenyl)cyclopropanecarbonyl chloride (approximately 3.8kg) in toluene (approximately 6-9 L), prepared in Step 8, at 1°C was added over a period of 74 minutes to a mixture of sodium azide (1.24kg), tetrabutylammonium bromide (56g) and sodium carbonate (922g) in water (6.2kg), stirring at 1.5°C. The mixture was stirred at 0°C for 1 hour and 55 minutes, then the aqueous layer  
15 was diluted with cold water (3.8kg), stirred briefly, then separated. The toluene layer was washed once more at 0°C with water (3.8kg), then with 20% aqueous sodium chloride (3.8 L), then stored at 3°C for further use.

Step 10: Preparation of *trans*-(1*R*, 2*S*)-2-(3,4-difluorophenyl)cyclopropylamine.

20 A cold solution of *trans*-(1*R*, 2*R*)-2-(3,4-difluorophenyl)cyclopropanecarbonyl azide prepared as described in Step 9 was added over a period of 41 minutes to toluene (6.0kg) stirring at 100°C. The mixture was stirred for a further 55 minutes at 100°C, then cooled to 20°C and added over a period of 2 hours and 15 minutes to hydrochloric acid (3M, 18.2kg) stirring at 80°C. After 65 minutes the solution was diluted with water (34kg)  
25 and cooled to 25°C. The toluene layer was removed and the aqueous layer was basified to pH 12 with 45% aqueous sodium hydroxide (3.8kg) and the product was then extracted into ethyl acetate (31kg) and washed twice with water (13.7kg per wash), affording a solution containing *trans*-(1*R*, 2*S*)-2-(3,4-difluorophenyl)cyclopropylamine (2.6kg, 91.8% enantiomeric excess) in ethyl acetate (29.5 L).

30


Step 11: Preparation of *trans*-(1*R*, 2*S*)-2-(3,4-difluorophenyl)cyclopropanaminium (2*R*)-2-hydroxy-2-phenylethanoate.

*R*-(-)-Mandelic acid (2.26kg) was added to a solution containing *trans*-(1*R*,2*S*)-2-(3,4-difluorophenyl)cyclopropylamine (2.6kg, 91.8% enantiomeric excess), stirring at 17°C in ethyl acetate (45.3 L). The mixture was stirred at 25°C for 3 hours and 8 minutes, then filtered and washed twice with ethyl acetate (13.8kg total) The crystalline product was dried at 40°C under reduced pressure for 23 hours, affording *trans*-(1*R*,2*S*)-2-(3,4-difluorophenyl)-cyclopropanaminium (2*R*)-2-hydroxy-2-phenylethanoate (4.45kg).

### EXAMPLE 3


This Example illustrates the preparation of 4,6-dichloro-2-(propylsulfanyl)-5-pyrimidinamine.

Step 1: 4,6-Dihydroxy-2-(propylsulfanyl)pyrimidine.



Water (670ml) was added to 2-thiobarbituric acid (200 g). The resulting mixture was stirred and sodium hydroxide (126.3g) was added in portions. The mixture was stirred for 40 min. then diluted with water. 1-Methyl-2-pyrrolidinone (400 ml) and 1-iodopropane (140.9ml) were then added. The resulting slurry was stirred at 20°C for 22h. The pH of the mixture was then adjusted to 6.5 by addition of 1M HCl (600ml) over 30 min, then to pH 2.5 by the addition of 6 M HCl (180ml) over a further 30 min. The resulting slurry was stirred for 18h and the product was isolated by filtration and washed successively with water (4 X 100ml), ethanol (200ml) and water (2 X 200ml). The product was dried under reduced pressure overnight at 50°C to yield the title product as a white powder (185g).


Step 2: 4,6-Dihydroxy-5-[(*E*)-2-(4-methylphenyl)diazonyl]-2-(propylsulfanyl)pyrimidine.



Ethanol (25ml), 4,6-dihydroxy-2-(propylsulfanyl)pyrimidine (Step 1; 5g) and water (25 ml) were stirred together at room temperature. Sodium hydroxide (1.02g) was added and a clear solution obtained. The resulting solution was cooled to 0°C and then sodium acetate (9.42g) was added to give Solution A.

5 In a separate vessel, a solution of p-toluidine (3.01g) in water (10 ml) was prepared. To this was added concentrated hydrochloric acid (37% w/w aqueous solution; 8.45 ml). The resulting mixture was cooled to 0°C, and a solution of sodium nitrite (2.16g) in water (10ml) was cooled to 0°C and added dropwise to the toluidine – containing reaction mixture over 30 minutes. The temperature during the addition was kept between 0 and 5°C.  
10 The resulting mixture was cooled to 0°C and added to the cold (0°C) Solution A (the temperature rose to 8°C). The resulting yellow suspension was stirred overnight and the pH of the mixture was adjusted to pH 1 by addition of 6 M HCl. The mixture was filtered and the collected product washed with water (25ml) ethanol (10ml). The product was dried under reduced pressure at 50°C over 24 h to give the product as a yellow solid (6.97g).


15 Step 3: 4,6-Dichloro-5-[(E)-2-(4-methylphenyl)diazenyl]-2-(propylsulfanyl)pyrimidine.



Pyridine (2.58ml) was added to a stirred, heated (70°C) slurry of 4,6-dihydroxy-5-[(E)-2-(4-methylphenyl)diazenyl]-2-(propylsulfanyl)pyrimidine (Step 2, 5g) in toluene  
20 (15ml). Phosphorous oxychloride (18.7ml) was added dropwise to the mixture over 15 min (exotherm to 94°C). The reaction mixture was heated for a further 4.5 h then evaporated. The residue was azeotroped twice with toluene (2 x 30ml). The residue was dissolved in toluene (50 ml) and filtered to remove some solids. The collected solid was washed with toluene and the combined filtrates washed with water (30ml) and saturated aqueous sodium  
25 bicarbonate solution (30ml). Evaporation gave the title product (4.98g) as a red oil that slowly crystallised on standing.



Step 4: Preparation of 4,6-dichloro-2-(propylsulfanyl)-5-pyrimidinamine.



A stirred solution of 4,6-dichloro-5-[(E)-2-(4-methylphenyl)diazanyl]-2-(propylsulfanyl)pyrimidine (Step (3), 1.1kg) in 2-propanol (16.6kg) was hydrogenated for 5  
1h at 40°C/3.2 bar over a platinum on carbon catalyst (0.81kg, 50% w/w Pt/C). The hydrogen gas pressure was released and the reactor flushed with nitrogen. The reaction mixture was filtered. The collected solid was washed with 2-propanol (1.7kg) and the combined filtrates were concentrated under reduced pressure. The residual oil was cooled to 20°C and dissolved in ethyl acetate (5kg) and water (5.5l) was added. The pH of the 10  
stirred mixture was adjusted to pH 2 by the addition of 3M aqueous hydrochloric acid (800ml). The phases were allowed to separate and the aqueous phase was discharged. Water (2.75l) was added to the organic phase and the pH was adjusted to 2 by the addition of a small amount of 3M HCl (45 ml). The aqueous phase was separated and the organic phase concentrated under reduced pressure at 30-50°C, to give 4,6-dichloro-2-  
15 (propylsulfanyl)-5-pyrimidinamine as a reddish viscous oil containing ethyl acetate that was dissolved in ethanol (8.5kg). Solvent (6.5l of ethanol/ethyl acetate) was then removed by distillation at reduced pressure. A further portion of ethanol (4.5kg) was added to the residue and the distillation repeated to remove 6.5 l of solvent. The ethanol solution of the product was used without further purification in the following step.


20

#### EXAMPLE 4

This Example illustrates the preparation of [1S-[1 $\alpha$ ,2 $\alpha$ ,3 $\beta$ (1S\*,2R\*),5 $\beta$ ]]-3-[7-[2-(3,4-difluorophenyl)cyclopropyl-amino]-5-(propylthio)-3H-1,2,3-triazolo[4,5-d]pyrimidin-3-yl]-5-(2-hydroxyethoxy)cyclopentane-1,2-diol (alternatively named (1S,2S,3R,5S)-3-[7-  
25 {{{(1R,2S)-2-(3,4-difluorophenyl)cyclopropyl}amino}-5-(propylsulfanyl)-3H-[1,2,3]triazolo[4,5-d]pyrimidin-3-yl]-5-(2-hydroxyethoxy)-1,2-cyclopentanediol).

Step 1: Preparation of [3aR-(3 $\alpha$ ,4 $\alpha$ ,6 $\alpha$ ,6 $\alpha$ )]-2-[[6-[[5-amino-6-chloro-2-(propylthio)-4-pyrimidinyl]amino]tetrahydro-2,2-dimethyl-3aH-cyclopenta[d][1,3]dioxol-4-yl]oxy]ethanol (alternatively named 2-[[[(3aR,4S,6R,6aS)-6-[[5-amino-6-chloro-2-


(propylsulfanyl)-4-pyrimidinyl]amino)-2,2-dimethyltetrahydro-3a*H*-cyclopenta[*d*][1,3]dioxol-4-yl]oxy]-1-ethanol).



The ethanolic solution of 4,6-dichloro-2-(propylsulfanyl)-5-pyrimidinamine,  
 5 (prepared as in Example 3, Step 4) was added to 2-(((3a*R*,4*S*,6*R*,6a*S*)-6-amino-2,2-dimethyltetrahydro-3a*H*-cyclopenta[*d*][1,3]-dioxol-4-yl]oxy)-1-ethanol, L-tartaric acid salt (1:1) (1.18kg). To the resulting stirred thick slurry was charged triethylamine (0.95kg) maintaining the temperature between 20 and 25°C. The reactor was sealed and the temperature increased to 120-125°C. The reaction mixture was kept within this  
 10 temperature range for 30h, then cooled to 75°C, and the pressure released. The temperature of the mixture was adjusted to 50°C and the solvent distilled off under reduced pressure at 30 to 40°C. Ethyl acetate (4.95kg) and water (5.5l) were added, the pH of the mixture adjusted to pH 5 by the addition of 3M hydrochloric acid (100 ml), and the phases were separated. The organic phase was washed with 15% w/w brine (5.5l), then separated. The  
 15 organic phase was concentrated under reduced pressure (4.8l of solvent removed) to give 2-(((3a*R*,4*S*,6*R*,6a*S*)-6-[[5-amino-6-chloro-2-(propylsulfanyl)-4-pyrimidinyl]amino)-2,2-dimethyltetrahydro-3a*H*-cyclopenta[*d*][1,3]dioxol-4-yl]oxy)-1-ethanol as a brown-red viscous oil containing ethyl acetate. The product is used without further purification in the following step.


20

Step 2: Preparation of [3a*R*-(3aα,4α,6α,6aα)]-2-[[6-[7-chloro-5-(propylthio)-3*H*-[1,2,3]triazolo[4,5-*d*]pyrimidin-3-yl]tetrahydro-2,2-dimethyl-3a*H*-cyclopenta[*d*][1,3]dioxol-4-yl]oxy]ethanol (alternatively named 2-(((3a*R*,4*S*,6*R*,6a*S*)-6-[7-chloro-5-(propylsulfanyl)-3*H*-[1,2,3]triazolo[4,5-*d*]pyrimidin-3-yl]-2,2-dimethyltetrahydro-3a*H*-cyclopenta-*d*][1,3]dioxol-4-yl]oxy)-1-ethanol).  
 25




2-(((3aR,4S,6R,6aS)-6-((5-Amino-6-chloro-2-(propylsulfanyl)-4-pyrimidinyl)amino)-2,2-dimethyltetrahydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)oxy)-1-ethanol, as obtained in Step 1 was dissolved in acetic acid (5.75kg) and water (650ml). The resulting solution was cooled to 2°C (with stirring) and a solution of sodium nitrite (232g) in water (1.25l) was added such that the mixture temperature was held below 7°C. The mixture was then allowed to warm to 7°C then ethyl acetate (8.9kg) was added. Aqueous potassium carbonate solution (4l, 37%w/w) was added. The mixture was separated and the organic phase washed with further aqueous potassium carbonate solution (3.8kg, 21% w/w). The aqueous phase was discarded and the organic phase concentrated under reduced pressure to give the sub-titled compound as a red-brown viscous oil used without further purification in the following step.

Step 3: Preparation of {3aR-[3aα,4α,6α(1R\*,2S\*),6aα]}-2-[6-({7-[2-(3,4-difluorophenyl)cyclopropyl]amino-5-(propylthio)-3H-1,2,3-triazolo[4,5-d]pyrimidin-3-yl)tetrahydro-2,2-dimethyl-4H-cyclopenta-1,3-dioxol-4-yl)oxy]ethanol (alternatively named 2-(((3aR,4S,6R,6aS)-6-[7-(((1R,2S)-2-(3,4-difluorophenyl)-cyclopropyl)amino)-5-(propylsulfanyl)-3H-[1,2,3]triazolo[4,5-d]pyrimidin-3-yl]-2,2-dimethyltetrahydro-3aH-cyclopenta[d][1,3]dioxol-4-yl)oxy)-1-ethanol).



*trans*-(1*R*,2*S*)-2-(3,4-Difluorophenyl)cyclopropanaminium (2*R*)-2-hydroxy-2-phenylethanoate (0.77kg) was charged to a vessel followed by a solution of 2-(((3*aR*,4*S*,6*R*,6*aS*)-6-[7-chloro-5-(propylsulfanyl)-3*H*-[1,2,3]triazolo[4,5-*d*]pyrimidin-3-yl]-2,2-dimethyltetrahydro-3*aH*-cyclopenta-*d*[1,3]dioxol-4-yl)oxy)-1-ethanol (prepared as in Step 2) dissolved in acetonitrile (3.85kg). To the resulting stirred mixture was added triethylamine (0.81 kg) at such a rate that the reaction temperature was maintained between 20-25°C. The reaction mixture was stirred for 13 h then concentrated at reduced pressure at 30°C. To the residue was added ethyl acetate (8.1 kg) and water (4.6l). The pH of the stirred two phase mixture was adjusted to pH 4 by the addition of 3M HCl (450ml). The mixture was then allowed to settle and separate. The aqueous phase was separated and the retained organic phase was washed with 15% w/w aqueous sodium chloride solution (4.15kg), the organic phase was concentrated under reduced pressure at 30-50°C giving the crude title compound as a red oil, that was used directly in the next step.

**Step 4:** Preparation of [1*S*-[1 $\alpha$ ,2 $\alpha$ ,3 $\beta$ (1*S*\*,2*R*\*),5 $\beta$ ]]-3-[7-[2-(3,4-difluorophenyl)-cyclopropylamino]-5-(propylthio)-3*H*-1,2,3-triazolo[4,5-*d*]pyrimidin-3-yl]-5-(2-hydroxyethoxy)cyclopentane-1,2-diol (alternatively named (1*S*,2*S*,3*R*,5*S*)-3-[7-(((1*R*,2*S*)-2-(3,4-Difluorophenyl)cyclopropyl)amino)-5-(propylsulfanyl)-3*H*-[1,2,3]triazolo[4,5-*d*]pyrimidin-3-yl]-5-(2-hydroxyethoxy)-1,2-cyclopentanediol).



Aqueous hydrochloric acid (3 M, 4.8l), was added to a stirred solution of 2-(((3*aR*,4*S*,6*R*,6*aS*)-6-[7-(((1*R*,2*S*)-2-(3,4-difluorophenyl)cyclopropyl)-amino)-5-(propylsulfanyl)-3*H*-[1,2,3]triazolo[4,5-*d*]pyrimidin-3-yl]-2,2-dimethyltetrahydro-3*aH*-cyclopenta-*d*[1,3]dioxol-4-yl)oxy)-1-ethanol (1.931kg) in methanol (13.4kg), maintaining the temperature during the addition between 20 and 25°C. The mixture was then stirred for 24 h at 20°C. Sodium hydroxide (45% w/w aqueous solution; 780ml) was then added to adjust the pH of the mixture to pH 7.2. The methanol was then removed by distillation at

reduced pressure and ethyl acetate (14.3 kg) added. The mixture was heated to 45° and the aqueous layer separated. The organic phase was then washed 15% w/w aqueous sodium chloride solution (7.2kg). Ethyl acetate (10l) was removed by distillation at reduced pressure. Fresh ethyl acetate (7.2kg) was charged and the mixture filtered. The filter was washed with ethyl acetate (1.5kg). The combined filtrates were dried by repeated addition/distillation of ethyl acetate. When the solution was dry, the product content of the ethyl acetate solution was determined by a chromatographic assay technique and found to contain 1016g of product, the concentration of the ethyl acetate was adjusted until a concentration of 5ml ethyl acetate / g of the crude product was reached. The ethyl acetate solution was heated to 47°C and isooctane (2.5ml/g product, 2540ml) was then added over 15 min. The resulting slurry was stirred for 30 min. then more iso-octane (2540ml) was added over 5 mins. The resulting mixture was stirred at 48-50°C for 30 min then cooled to 20°C over 3h. The slurry was stirred at 20°C for 6.5 h then filtered and washed with a mixture consisting of iso-octane (1.25kg) and ethyl acetate (1.6kg). The collected solid was dried in vacuo to give the title compound (920g).

If desired, the crude product can be further purified by employing one of the following three methods.

Recrystallization from ethyl acetate / iso-octane

The crude product is dissolved in ethyl acetate (4.8 ml/g) at 55°C, then filtered to remove particles. The clear solution is taken back to the reactor for recrystallisation and the temperature is set at 50°C. iso-Octane is then added (4.8 ml/g) during 10 min. The slurry is allowed to stand for 30 minutes after which it is cooled to 20°C during 2-3 hours and finally the temperature is kept at 20°C for about 30 minutes. The product is then filtered and washed with iso-octane (2x1.5 ml/g). The product is dried under reduced pressure at 50°C giving pure product (>98% pure by h.p.l.c. analysis).

Slurry with n-butyl acetate

The crude product is suspended in n-butyl acetate 4 ml/g and stirred at room temperature for 10 hours. The slurry is cooled to 0°C during 3-4 hours and kept at 0°C for 1 hour. The product is filtered and washed with 2 ml/g of cold n-butyl acetate (<0°C). The

product is then dried in vacuo at 50°C, giving pure product (>98% pure by h.p.l.c. analysis).

Slurry with iso-propanol


5           The crude product is suspended in iso-propanol 3 ml/g and stirred at 50°C for 72h. The slurry is then cooled to 20°C during 3 hours and the temperature is kept at 20°C for about 30 minutes. The product is then filtered and washed with 1 ml/g of cold iso-propanol (<0°C). Finally, the product is dried under reduced pressure at 50°C giving pure product (>98% pure by h.p.l.c. analysis).

10           Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.


15           The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to  
20           which this specification relates.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:


1. A compound of formula (I):




2. A process for preparing a compound of formula (I), comprising reacting a compound of formula (II):



with a salt of a compound of formula (III):




3. A process for preparing a compound of formula (I) as claimed in claim 2 comprising hydrogenating a compound of formula (IV):




wherein Ar is phenyl optionally substituted by halogen, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy; to give a compound of formula (II), and reacting the compound of formula (II) with a compound of formula (III) to provide the compound of formula (I).

4. A process for preparing a compound of formula (II):




comprising hydrogenating a compound of formula (IV):




wherein Ar is phenyl optionally substituted by halogen, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy.

5. A process for preparing a compound of formula (II) as claimed in claim 2, comprising chlorinating a compound of formula (VIII):



wherein Ar is phenyl optionally substituted by halogen, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy, to provide a compound of formula (IV), and then hydrogenating the compound of formula (IV).


6. The use of a compound of formula (I) to prepare a compound of formula (A):




comprising:




- (a) reacting a compound of formula (I) with an alkali metal nitrite in the presence of a suitable acid to prepare a compound of formula (VII):



- (b) coupling the compound of formula (VII) with a compound of formula (VI):




to provide a compound of formula (V):



and,


- (c) deprotecting a compound of formula (V) to provide the compound of formula (A).

7. A process for preparing a salt of a compound of formula (III):




comprising reacting a compound of formula (III) with the necessary acid.

8. A process for preparing a salt of a compound of formula (III):




comprising the steps:

- (a) preparing a compound of formula (XI):




by reacting a compound of formula (XII):




with benzyl chloroformate;

- (b) preparing a compound of formula (X):




by reacting a compound of formula (XI) with a compound L-CH<sub>2</sub>CO<sub>2</sub>R\*  
{wherein R\* is C<sub>1-4</sub> alkyl; and L is a leaving group};

- (c) preparing a compound of formula (IX):



- by reducing a compound of formula (X);  
(d) preparing a compound of formula (III):



- 5 by deprotecting a compound of formula (IX); and,  
(e) preparing a salt of a compound of formula (III) by reacting the compound of  
formula (III) with the necessary acid.
9. An intermediate compound of formula (II), (IV), (VII), (VIII), (X) or (XI) or a salt  
10 of a compound of formula (III).