



US00PP16390P3

(12) **United States Plant Patent**
Iredell et al.

(10) **Patent No.:** **US PP16,390 P3**

(45) **Date of Patent:** **Mar. 28, 2006**

(54) **BOUGAINVILLEA PLANT NAMED 'NINGILI'**

(50) Latin Name: *Bougainvillea hybrida*
Varietal Denomination: **Ningili**

(75) Inventors: **Peter Iredell**, Moggil Brisbane (AU);
Jan Iredell, Moggil Brisbane (AU)

(73) Assignee: **Bougainvillea Nursery**, Queensland
(AU)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 118 days.

(21) Appl. No.: **10/973,064**

(22) Filed: **Oct. 23, 2004**

(65) **Prior Publication Data**

US 2005/0262607 P1 Nov. 24, 2005

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./256**

(58) **Field of Classification Search** **Plt./256**
See application file for complete search history.

Primary Examiner—Kent Bell
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Bougainvillea* plant named
'Ningili', characterized by its outwardly spreading and
bushy plant habit; vigorous growth habit; dark green-colored
leaves; freely flowering habit; and light red purple-colored
flower bracts.

1 Drawing Sheet

1

Botanical classification/cultivar designation: *Bougainvillea hybrida* cultivar Ningili.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct cultivar of *Bougainvillea* plant, botanically known as *Bougainvillea hybrida*, and hereinafter referred to by the name 'Ningili'.

The new *Bougainvillea* is a naturally-occurring branch mutation of an unnamed selection of *Bougainvillea hybrida*, not patented. The new *Bougainvillea* was discovered and selected by the Inventors from within a population of plants of the unnamed selection in a controlled environment in Brisbane, Queensland, Australia in 1989. The selection of this plant was based on its attractive flower coloration.

Asexual reproduction of the new cultivar by cuttings at Brisbane, Queensland, Australia since 1990, has shown that the unique features of this new *Bougainvillea* are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the cultivar Ningili have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and/or light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Ningili'. These characteristics in combination distinguish 'Ningili' as a new and distinct cultivar:

1. Outwardly spreading and bushy plant habit.
2. Vigorous growth habit.
3. Dark green-colored leaves.
4. Freely flowering habit.
5. Light red purple-colored flower bracts.

2

Plants of the new *Bougainvillea* are most similar to plants of the parent, the unnamed selection. In side-by-side comparisons conducted in Moggil, Queensland, Australia, plants of the new *Bougainvillea* differed primarily from plants of parent selection primarily in flower bract coloration as plants of the parent selection had pink-colored flower bracts.

Plants of the new *Bougainvillea* can be compared to plants of the cultivar Blushing Beauty, not patented. In side-by-side comparisons conducted in Moggil, Queensland, Australia, plants of the new *Bougainvillea* differed primarily from plants of the cultivar Blushing Beauty flower in bract coloration as flower bracts of plants of the cultivar Blushing Beauty were soft pink in color.

Plants of the new *Bougainvillea* can also be compared to plants of the cultivar Donya, not patented. In side-by-side comparisons conducted in Moggil, Queensland, Australia, plants of the new *Bougainvillea* differed primarily from plants of the cultivar Donya in growth habit as plants of the cultivar Donya were larger than plants of the new *Bougainvillea*.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ from the color values cited in the detailed botanical description which accurately describe the colors of the new *Bougainvillea*.

The photograph at the top of the sheet is a side perspective view of typical one-year old plants of 'Ningili' grown in containers in an outdoor nursery.

The photograph in the middle of the sheet is a close-up view of typical developing flowers of 'Ningili'.

The photograph at the bottom of the sheet is a close-up view of typical developing leaves of 'Ningili'.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, except where general terms of ordinary dictionary significance are used. Plants used for the following botanical description were grown in Moggill, Queensland, Australia in a polyethylene-covered greenhouse. Plants used for the description were about one to two years old.

Botanical classification: *Bougainvillea hybrida* cultivar Ningili.

Parentage: Naturally-occurring branch mutation of an unnamed selection of *Bougainvillea hybrida*, not patented.

Propagation:

Type.—By cuttings.

Time to initiate roots.—About one week at 25° C.

Time to develop roots.—About two weeks at 25° C.

Root description.—Thick; white in color.

Plant description:

Form.—Outwardly spreading and bushy plant habit; vigorous growth habit. Freely branching with lateral branches potentially developing at every node.

Plant height, soil level to top of plant plane.—About 1 to 1.5 meters.

Plant width.—About 1 to 1.5 meters.

Stem color.—153A.

Stem texture.—Smooth, glabrous.

Foliage description.—Arrangement: Opposite, simple. Length: About 8.1 cm. Width: About 6.1 mm. Shape: Broadly obovate. Apex: Acuminate. Base: Obtuse. Margin: Entire; slightly undulate. Texture, upper and lower surfaces: Smooth, glabrous. Venation pattern: Pinnate. Color: Developing foliage, upper surface: 137B; towards the center, 144A. Developing foliage, lower surface: 137C. Fully expanded foliage, upper surface: 139A; towards the center, 144A. Fully expanded foliage, lower surface: 137A; towards the center, 144A. Venation, upper surface: 137B. Venation, lower surface: 144A. Petiole length: About 2.2 cm. Petiole diameter: About 1 mm. Petiole texture, upper and lower surfaces: Smooth, glabrous. Petiole color, upper and lower surfaces: 138B.

Flower description:

Flower type and habit.—Single flowers arranged in axillary panicles with showy bracts. Flowers face mostly upright. Flowers not persistent.

Fragrance.—None.

Natural flowering season.—Cyclically flowering year-round in Moggill, Queensland, Australia; flowering periods about six to twelve weeks long.

Quantity.—Freely flowering with about 35 flowers per panicle.

Flower longevity.—About six to ten days.

Flower diameter.—About 4.5 mm.

Flower depth (height).—About 1.4 cm.

Flower buds.—Length: About 1.4 cm. Diameter: About 2 mm. Shape: Elongated. Color: N144C.

Petals.—Quantity per flower: About five; fused tubular corolla. Length: About 1.4 cm. Width: About 3 mm. Shape: Fan-shaped. Apex: Emarginate. Margin: Entire. Texture, upper and lower surfaces: Smooth; glabrous. Color: When opening, upper and lower surfaces: 160D. Fully opened, upper and lower surfaces: 159D.

Sepals.—Quantity per flower: About five; fused tubular corona. Length: About 1.7 cm. Diameter: About 2.5 mm. Shape: Linear. Apex: Acute. Margin: Entire. Color: When opening, upper and lower surfaces: 191B. Fully opened, upper and lower surfaces: 191B.

Flower bracts.—Quantity per flower: Three. Length: About 3.3 cm. Width: About 2.3 cm. Shape: Broadly ovate. Apex: Acuminate. Base: Obtuse. Margin: Entire; undulate. Texture, upper and lower surfaces: Smooth; glabrous; rugose. Color: When opening, upper and lower surfaces: N57D. Fully opened, upper and lower surfaces: N57D.

Peduncles.—Length: About 4.4 cm. Diameter: About 1.25 mm. Texture: Smooth, glabrous. Angle: About 45° from vertical. Strength: Strong. Color: 153B.

Pedicels.—Length: About 1.5 cm. Diameter: About 1 mm. Texture: Smooth, glabrous. Angle: About 45° from vertical. Strength: Strong. Color: N144A.

Reproductive organs.—Androecium: Stamens per flower: About seven. Anther shape: Oblong. Anther length: About 0.75 mm. Anther color: 22C. Pollen color: 22C. Gynoecium: Quantity of pistils per flower: One. Pistil length: About 9 mm. Stigma shape: Linear. Stigma color: Close to 155D. Style length: About 2.5 mm. Style color: N144D. Ovary color: N144D.

Seeds/fruits.—Seed and fruit development has not been observed.

Disease/pest resistance: Under commercial production conditions, plants of the new *Bougainvillea* have not been noted to be resistant to pathogens or pests common to *Bougainvillea*.

Garden performance: Plants of the new *Bougainvillea* have good garden performance as plants of the new *Bougainvillea* have been observed to tolerate wind, rain and temperatures from 3 to 40° C.

It is claimed:

1. A new and distinct cultivar of *Bougainvillea* plant named 'Ningili', as illustrated and described.

* * * * *

