



(72) HESSEL, LASSE, DK

(71) HESSEL, LASSE, DK

(51) Int.Cl.<sup>6</sup> A61K 35/78

(30) 1997/12/08 (60/067,706) US

(54) **COMPOSITIONS ET TECHNIQUES POUR PERTE DE POIDS**

(54) **COMPOSITIONS AND METHODS FOR WEIGHT REDUCTION**

(57) Cette invention concerne des compositions renfermant une combinaison d'extraits d'herbes sélectionnés qui ralentit la vidange gastrique et augmente la vitesse de métabolisme, ce qui favorise la perte de poids chez les patients.

(57) Compositions containing a combination of selected herbal plant extracts that inhibit gastric emptying time and increase metabolic rate are provided which are useful in reducing weight in patients.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b>  <b>A61K 35/78</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/29333</b>  <b>(43) International Publication Date:</b> 17 June 1999 (17.06.99)
<b>(21) International Application Number:</b> PCT/IB98/02106  <b>(22) International Filing Date:</b> 7 December 1998 (07.12.98)  <b>(30) Priority Data:</b> 045,650                      8 December 1997 (08.12.97)                      US  <b>(71)(72) Applicant and Inventor:</b> HESSEL, Lasse [DK/DK]; Rantzausmindevej 111, DK-5700 Svendborg (DK).  <b>(74) Agent:</b> KING, James, B.; Kings Patent Agency Limited, 73 Farringdon Road, London EC1M 3JB (GB).	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.  Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(54) Title:</b> COMPOSITIONS AND METHODS FOR WEIGHT REDUCTION		
<b>(57) Abstract</b>		
<p>Compositions containing a combination of selected herbal plant extracts that inhibit gastric emptying time and increase metabolic rate are provided which are useful in reducing weight in patients.</p>		

WO 99/29333

PCT/IB98/02106

## TITLE

**Compositions and Methods for Weight Reduction****5 Background of the Invention.**

Obesity is a condition that affects millions of Americans. Recent statistics show that when obesity is defined as a 20% excess over desirable weight, 20 – 30% of adult men and 30-40% of adult women are obese. Even mild obesity increases the risk for premature death, diabetes, hypertension, atherosclerosis, gallbladder disease, and even some forms of cancer (Olefsky, J.M. 1994) in: Harrison's Principles of Internal Medicine, pp. 46-452). Therefore, effective methods for weight reduction are constantly being sought.

Although there are many different treatment regimens in use today that can produce a short term or temporary weight loss, most are associated with a rapid increase in weight once treatment is terminated. Caloric restriction is the main goal of most weight reduction treatment regimens. The basic principle is that if intake of food is less than energy expenditure, stored calories will be consumed, mainly in the form of fat. However, once the diet regimen is broken, weight is quickly regained.

20 Other treatment regimens are based on the principle of increasing metabolism. By increasing metabolism, calories are burned thereby decreasing body weight. However, these treatments often have side effects, particularly those involving use of non-prescription and prescription drug products. Further, these treatments also often result in rapid weight increases once treatment is

terminated, unless modification of behavior that led to weight gain is undertaken.

In the case of nutritional supplement regimens, poor taste is often a problem despite addition of taste-improving substances to the product.

Weight reduction has also been attempted using surgical intervention  
5 wherein the size of the stomach is reduced so that a feeling of gastric fullness is produced, resulting in a decrease in appetite and food intake. One such method is placement of a mechanical device such as an inflatable balloon into the stomach. However, such invasive methods are not routinely used.

The use of herbal plant extracts to control weight has also been described.  
10 For example, the herb Guarana (paullinia cupana, P. sorbolis) which contains a high concentration of the active ingredient caffeine has been incorporated into slimming products (Hurel, J.-P. 1993. FR 2 712 191-A1). Caffeine is recognized to have pharmacological activity as a central nervous system stimulant and is a major constituent in many weight-reducing products for its  
15 ability to increase metabolic rate. However, use of such a product alone has only a temporary effect, with weight gain seen immediately upon cessation of treatment. Similarly, Prunet (FR 2 687 548 A1) describes the use of Guarana as a nutritional supplement in short term weight reduction. Finally, Primez (Belgium Patent 100593A7) describes a phyto-active mixture referred to as  
20 Lycopodium which contains Guarana and other plant extracts including Scillia maritime, Ephedra vulgaris, and Betula alba which may be ingested, applied as a cream or lotion, or injected to produce weight loss. It is suggested that administration of Lycopodium slows gastric draining while improving intestinal transit and evacuation of the intestines. However, no clinical data is provided to

support this suggestion.

It has now been found that a combination of selected herbal extracts wherein at least one of the extracts contains caffeine and at least one of the extracts controls gastric emptying is capable of producing weight loss.

5 **Summary of the Invention**

An object of the present invention is to provide a composition comprising a combination of selected herbal extracts that produces weight loss in a patient through inhibition of gastric emptying and an increase in metabolic rate in the patient.

10 Another object of the present invention is to provide a method of reducing weight in patients which comprises administering to the patient a composition comprising a combination of selected herbal extracts that inhibits gastric emptying and increases metabolic rate in the patient.

**Detailed Description of the Invention**

15 The periodic discharge of food from the stomach into the small intestine, also referred to as gastric emptying, is caused by contraction of the muscles in the wall of the stomach. These muscles are innervated by the cranial vagus nerves, which stimulate contraction of the gastric muscles and allow sphincter between the stomach and the duodenum to open. The present invention  
20 relates to a composition comprising a combination of selected herbal extracts wherein at least one of the herbal extracts inhibits gastric emptying. Because nutritional uptake through the mucosal lining of the stomach is extremely low, the extended retention period in the stomach of food resulting from these compositions does not have any discernable effect on the eventual uptake of

nutrients. However, inhibition of gastric emptying results in a decreased appetite thereby decreasing food intake. The compositions of the present invention further comprise at least one herbal extract capable of modifying metabolic rate through the presence of significant concentrations of caffeine.

5 Increasing the metabolic rate of a patient while inhibiting gastric emptying in the patient by administering a composition of the present invention results in weight loss. Active ingredients extracted from the plants are used as a natural nutritional supplement in such a way to control the uptake of nutrients by delaying gastric emptying. At the same time plant extracts are incorporated

10 which are known to promote weight loss by increasing metabolism. The composition of the present invention thus provides a combination of selected herbal extracts which has been shown to be effective in producing weight loss in clinical studies.

Herbal plant extracts that have been assessed and found to be suitable for

15 selection and incorporation into a composition of the present invention for achieving a controlled and durable weight loss include Buchu (*Barosma betulina*, *B. crenulata*, *B. serratifolia*), Vervain (*Verbena officinales*, *V. jamaicensis*, *V. lappulacae*, *V. hesitate*, *V. urticifolia*, *V. Sinuata*), Damiana (*Tumera diffusa* var. *aphrodisiaca*, *T. opifera*, *T. ulmifoliei*), Guarana (*Paullinia cupana*, *P. sorbalis*), Paraguay (*Ilex paraguayensis*, *I. vomitora*, *I. Dahoon*), Kola (*Cola nitida*, *C. Vera*), and Ginseng (*Panax ginseng*, *P. quinquefolius* L.) The

20 active ingredients of each herb are listed in Table 1.

<b>TABLE 1</b> <b>Active Ingredients of Selected Herbs</b>	
<b>Herb</b>	<b>Active Ingredient(s)</b>
<b>Buchu</b>	volatile oil of buchu camphor diosphenol
<b>Vervain</b>	verbanaline (glycoside, adenosine, essential oils, tannin, livertin, emulin
<b>Damiana</b>	ethers, terpenes ( $\alpha$ -pinene, cineol, p-cymol, sesquiterpenes), resin, bitter pineapple, tannin, caoutchouc, albuminoids, starch, arbutin
<b>Guarana</b>	caffeine, other xanthines (tetramethylxanthine, theobromine, theophylline, tannin)
<b>Paraguay</b>	caffeine
<b>Kola</b>	caffeine
<b>Ginseng</b>	triterpenoid saponin

As will be obvious to those of skill in the art, other herbal extracts having  
 5 these active ingredients can also be selected for use in a composition of the  
 present invention. Herbal extracts for combination into a composition of the  
 present invention are obtained in accordance with methods well known and  
 routine to those of skill in the art.

A composition of the present invention comprising a combination of selected  
 10 herbal extracts was administered to patients in a double blind controlled clinical  
 trial. The combination tested included Guarana, Damiana, and Paraguay.  
 Guarana is a dough from the seeds of Paullinia sorbolis, which grows in Brazil  
 and Venezuela. It contains 3-6% caffeine, 5-8.5% tannin, 7.8% resins, 2-3%  
 fat, 0.06% saponin, 5-6% starch, and 1.5% coloring agents. Paraguay is an  
 15 extract of Ilex paraguensis which grows in Brazil, Argentina, and Paraguay. It

contains 1-1.5% caffeine, 4-10% tannin, and 3% resins and fat. Damiana is obtained from the leaves of the plant *Turnera diffusa* var. *aphorodisiaca* from California, Mexico and Brazil, and Bolivia and contains ethereal oils, resins, and tannin. These extracts were obtained as powders. The components were  
5 mixed and prepared as capsules. Each capsule contained 95 mg Guarana, 112 mg Paraguay, and 36 mg Damiana extract. The subjects for the study were 20 otherwise healthy subjects, complaining of light-moderate overweight with a body mass index between 25 and 30 kg/m<sup>2</sup>. None of the subjects were taking any drug or dietary supplement at the time of the study. All were briefed on the  
10 protocol and gave consent to the trial.

In this double blind placebo controlled trial, the subjects were randomized into two groups A and B. Group A was supplied with test capsules and group B with placebo (water coated capsules containing lactose) for 20 days. The participants took three meals a day and were instructed to take 2 capsules with  
15 a large glass of water (250 ml) from 10-15 minutes before each meal. Using a stopwatch, each subject then recorded the time elapsing to perception of gastric fullness. Subjects were also asked to note any side effects. Three days after the end of the first 20 day trial, the procedure was repeated with the test capsules now being given to the group B subjects and the placebo capsules to  
20 group A subjects. All subjects completed the test.

The subjects in Group B, receiving the placebo capsules in the first period reported an average time for perception of fullness of 60 minutes (range 55-65 minutes). In the second period, when taking the test combination product, the average time for perception of fullness in this same group was 36 minutes



(range 33-41 minutes). These average values were statistically different ( $p < 0.01$ ). Therefore, the study showed that treatment with the herbal extract combination produced statistically significant decreases in the time to perception of fullness.

5 The rate of gastric emptying was assessed by scintigraphy. Three volunteer male subjects with no gastrointestinal illness or intake of medicinal drugs took part in the study. They were given a meal consisting of 18 g peas, 100 g dried potatoes, and 200 ml of water containing 16-20 Mbc 113 Indium-DPTA. The subjects were fasting before the test and the meal was consumed  
10 in five minutes. The test was conducted with the subjects in a semi-upright position with a gamma camera placed in front. The time course of radioactivity in the stomach was determined by measuring the radioactivity in an appropriate region of interest every minute over 90 minutes. The test as repeated after each subject had taken three capsules of the drug combination three times daily  
15 and three capsules having been mixed with the food. Results showed that the rate of gastric emptying was significantly decreased in the three subjects after taking the herbal extract combination product. Halving times of gastric emptying were 49, 31, and 32 minutes after taking the test product and 61, 50, and 49 minutes, respectively, after taking the placebo.

20 Ultrasound examination of the stomach was also employed using a 3.5 MHz curved array transducer and an Aloka 630 standard unit employing a modification of the techniques by Holt et al. (Holt, S. et al. 1980. Gut 1:597-601) and Bateman and Whittingham (Bateman, D.N. and T.A. Whittingham. 1982. Gut 23:524-527. Continuous scans were performed switching the

transducer between two alternative projections, one oblique upward view with the transducer positioned under the left curvature allowing the gastric fundus, corpus and antrum to be inspected or a transverse view across the epigastrium with a slight upward direction viewing the antrum pylorus and the duodenal bulb.

5 All examinations were recorded on videotape and still pictures were taken every five minutes. This technique was used in a further double-blind crossover study on 7 healthy normal volunteers with no history of gastrointestinal diseases. Each volunteer had 2 to 8 examinations with the test capsules or placebo capsules (lactose), followed by 20 ml of apple juice and 15  
10 minutes later with 400 ml of apple juice. The projections gave clear visual estimation of the volume of the stomach. Gastric emptying time (GET) was defined as the elapsed time between ingestion of the 400 ml of apple juice and the time when the fundus and corpus of the stomach were completely empty. The results were noted immediately and controlled by playback of the  
15 videotapes. After termination of the study the codes were broken and GET values were compared by an independent analyst. The results showed that there was a considerable variation among the subjects and within the same subject. However, even with this variability, a delaying effect on gastric emptying was associated with administration of the herbal extract combination  
20 product, an effect that was evident in all seven subjects and is shown in Table 2.

TABLE 2 Results of Ultrasound Study: Comparison of Gastric Emptying Time (GET)		
Subject	Placebo GET (minutes)	Herbal Extract GET (minutes)
A	37.0	63.5
B	47.5	70.0
E	45.5	80.0
M	29.0	44.5
H	31.0	46.0
J	34.0	39.0
T	44.0	60.0
Mean Value	31.8	57.6

The effect on body weight of 10 days treatment with the test compound and placebo was recorded in a double blind pilot study of 44 healthy subjects was also determined. None of the subjects took any drugs and none were on a specific diet. The patients were instructed to take three capsules (test compounds or placebo) with a large glass of water 15 minutes before main meals and also to take care to not change their normal food habits. They also were asked to note side effects. The subjects were weighed before and after the period of 10 days. Twenty-two patients with a BMI range 25.1 to 29.5 took the test capsules and 22 patients with a BMI of 24.9 to 29.0 received the placebo. Results showed that of the 22 subjects who received the placebo, there was a mean decrease in body weight of 0.3 kg (SEM = 0.03) while in the 22 subjects who received the herbal extract product, there was a mean decrease of 0.8 kg (SEM = 0.05).

The effect on body weight of 45 days of treatment with the herb combination and placebo was also studied in a double blind randomized crossover study of 47 patients. These subjects were healthy but overweight (BMI range of 25.8 to

30.4). They did not take any medicinal drug or diet before or during the study. All 47 were between the ages of 20 and 60 years of age and gave full consent. In the study, the 24 subjects in group A received the test capsules in the first period. This group showed a mean decrease in body weight of 5.1 kg (SEM = 5 0.5) while taking the herbal extract product. In contrast, the placebo group in the first period had a mean decrease in body weight of only 0.5 kg (SEM = 0.08).

A recording of weight maintenance over 12 months after an initial weight loss was made to assess the long term effectiveness of the treatment. Twenty- 10 two of the subjects from the various studies above who had lost an average of 3.6 kg were invited to take part. Each subject received a month's supply of the test drug each month they returned to the study center for weight measurement.

No side effects were noted in any of the clinical tests. The results of these experiments demonstrate the effectiveness of the herbal extract combination in 15 weight reduction in humans.

What is claimed is:

1. A composition which produces weight loss in a patient comprising a combination of selected herbal extracts wherein said combination comprises at least one herbal extract capable of inhibiting gastric emptying and one herbal  
5 extract which increases metabolic rate in a patient.
  
2. The composition of claim 1 wherein the combination of selected herbal extracts comprises Guarana, Damiana, and Paraguay.
  
- 10 3. A method of reducing weight in a patient comprising administering to a patient a composition of claim 1 so that gastric emptying is inhibited and metabolic rate is increased in the patient.