



(86) Date de dépôt PCT/PCT Filing Date: 2013/01/23
(87) Date publication PCT/PCT Publication Date: 2013/08/01
(85) Entrée phase nationale/National Entry: 2014/07/23
(86) N° demande PCT/PCT Application No.: US 2013/022743
(87) N° publication PCT/PCT Publication No.: 2013/112582
(30) Priorité/Priority: 2012/01/23 (US61/589,601)

(51) Cl.Int./Int.Cl. *A61K 36/48* (2006.01),
A61P 1/00 (2006.01), *A61P 31/04* (2006.01)
(71) Demandeur/Applicant:
SANTALIS PHARMACEUTICALS INC., US
(72) Inventeurs/Inventors:
CLEMENTS, IAN, US;
CASTELLA, PAUL, US;
LEVENSON, COREY, US
(74) Agent: MARKS & CLERK

(54) Titre : HUILE DE BOIS DE SANTAL ET SES UTILISATIONS AFFERENTES AUX INFECTIONS A CLOSTRIDIUM
(54) Title: SANDALWOOD OIL AND ITS USES RELATED TO CLOSTRIDIUM INFECTIONS

(57) **Abrégé/Abstract:**

Provided herein are methods of using therapeutically effective compositions of sandalwood oil to treat, prevent or reduce the recurrence of Clostridium infections.



(51) International Patent Classification:
A61K 36/48 (2006.01) A61P 1/00 (2006.01)
A61P 31/04 (2006.01)

(21) International Application Number:
PCT/US2013/022743

(22) International Filing Date:
23 January 2013 (23.01.2013)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/589,601 23 January 2012 (23.01.2012) US

(71) Applicant: SANTALIS PHARMACEUTICALS INC.
[US/US]; 12621 Silicon Drive, Suite 100, San Antonio,
Texas 78249 (US).

(72) Inventors: CLEMENTS, Ian; 12621 Silicon Drive, Suite
100, San Antonio, Texas 78249 (US). CASTELLA, Paul;
12621 Silicon Drive, Suite 100, San Antonio, Texas 78249
(US). LEVENSON, Corey; 12621 Silicon Drive, Suite
100, San Antonio, Texas 78249 (US).

(74) Agents: PRATT, John S. et al.; 1100 Peachtree Street,
Suite 2800, Atlanta, Georgia 30309 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,

HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))

— as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii))

Published:

— with international search report (Art. 21(3))

— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))

(54) Title: SANDALWOOD OIL AND ITS USES RELATED TO CLOSTRIDIUM INFECTIONS

(57) Abstract: Provided herein are methods of using therapeutically effective compositions of sandalwood oil to treat, prevent or re-
duce the recurrence of Clostridium infections.

WO 2013/112582 A1

**SANDALWOOD OIL AND ITS USES RELATED TO
CLOSTRIDIUM INFECTIONS
RELATED APPLICATIONS**

This application claims the benefit of U.S. Application No. 61/589,601, filed on
5 January 23, 2012, which is hereby incorporated herein in its entirety.

SUMMARY

Provided herein are methods of using therapeutically effective compositions of
sandalwood oil to treat Clostridium infections. More specifically, provided herein is a
method of treating or preventing a Clostridium infection in a subject. The method
10 comprises administering to the subject a composition including a therapeutically effective
amount of sandalwood oil, wherein the subject has a Clostridium infection or is at risk of
developing a Clostridium infection.

Further provided is a method of reducing the recurrence of a Clostridium infection in
a subject. The method comprises administering to the subject a composition including a
15 therapeutically effective amount of sandalwood oil, wherein the subject is at risk for
recurrence of a Clostridium infection.

DESCRIPTION OF DRAWING

Figure 1 is a comparison of *Clostridium difficile* (*C. difficile*) inhibiting activity of
sandalwood oils as compared to other oils. Light gray wells indicate bacterial growth. Dark
20 gray wells indicate no bacterial growth. Figure 1 shows that West Australian sandalwood
oil and East Indian sandalwood oil are effective inhibitors of *C. difficile*.

DETAILED DESCRIPTION

In the present methods, oil from any member of the genus *Santalum* can be used.
For example, and not to be limiting, East Indian sandalwood (*Santalum album*) or West
25 Australian sandalwood (*Santalum spicatum*) can be utilized in the methods set forth herein.
Several other species of the genus also have fragrant wood and are found across India,
Australia, Indonesia, and the Pacific Islands. For example, *Santalum ellipticum*, *S.*
freycinetianum, and *S. paniculatum*, the Hawaiian sandalwoods, can also be used.

By way of example, *Santalum spicatum* (West Australian sandalwood) can be used.
30 Other species produced in Australia that can be utilized in the methods and compositions set
forth herein include, but are not limited to, *S. acuminatum*, *S. lanceolatum*, *S. murrayanum*,
S. obtusifolium and *S. album*. The compositions set forth herein can comprise one or more

sandalwood oils. The oil(s) can be from one or more members of the genus *Santalum*. In the compositions set forth herein, the sandalwood oil can be a sandalwood heartwood oil, a sandalwood nut oil, or a combination of both.

The components of *S. spicatum* and *S. album* species are different. A comparison of the components of steam distilled Australian and Indian sandalwood heartwood oils is presented in Table 1. The components and their percentages can vary with the extraction method.

Table 1: Typical Sandalwood Heartwood Oil Profiles

Compound	<i>S. spicatum</i> %	<i>S. album</i> %
E nerolidol	2.1%	0.1%
Alpha-santalene	nd	0.5%
Cis-alpha-(trans) bergamotene	nd	0.7%
Epi-beta-santalene	nd	1.1%
Beta-santalene	nd	0.3%
Gamma-curcumene	nd	0.2%
Dendrolasin	1.2%	0.2%
Alpha-santalol	17.2%	48.7%
Beta-bisabolol	2.3%	0.5%
Epi -alpha-bisabolol	8%	nd
Z-alpha trans -bergamotol	4.2%	2.4%
Epi beta -santalol	1.2%	5%
Cis -beta-santalol	11.4%	20.4%
E,E, farnesol	6.5%	nd
Cis nuciferol	13.5%	0.6%
Z-beta-curcumen-12-ol	7.9%	0.2%
cis lanceol	2.9%	1.5%

The sandalwood heartwood oil can be prepared by steam distillation, supercritical CO₂ extraction, solvent extraction, hydro-distillation and combinations thereof. The sandalwood heartwood oil can also be double distilled. It is also possible to synthesize one or more of the active ingredients of sandalwood heartwood oil, as identified in Table 1 and thereafter combine individual active ingredients together.

As used herein, a sandalwood heartwood oil can be a sandalwood heartwood oil that conforms with International Organization for Standardization (ISO) specifications for the oil and therefore comprises 20-45% santalols, when derived from *S. spicatum*, and 57-79% santalols when derived from *S. album*. However, the 20-45% santalols from *S. spicatum* and the 57-79% santalols from *S. album* are determined against the pure oil and before such oil is combined with any other solvents, excipients or active ingredients. It is understood that an efficacious preparation of sandalwood heartwood oil may have a concentration of

santalols lower (or higher) than the sandalwood heartwood oil it is prepared from and that the efficacious concentrations may be derived from sandalwood heartwood oils that are outside of the ISO specification prior to formulation. A santalol can be an α -santalol (shown below), a β -santalol (shown below), or any other active isomers or derivatives (such as
 5 esters) thereof.

As used herein, a sandalwood heartwood oil can comprise at least about 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 99% santalols or any percentage in between the percentages set forth herein, when derived from *S. spicatum*. The sandalwood heartwood oil can comprise at least 45%, 50%, 55%, 60%,
 10 65%, 70%, 75%, 80%, 85%, 90%, 95%, 99% santalols or any percentage in between the percentages set forth herein, when derived from *S. album*. As set forth above, the oil can be extracted from cultivated trees or from cell culture of tree cells.

In the methods and compositions set forth herein, the sandalwood heartwood oil can comprise the ingredients in the amounts listed in Table 1 plus or minus about 20%, and
 15 more preferably plus or minus about 10%, 5%, 2%, 1% or any percentage in between the percentages set forth herein.

It is also understood that the activity of sandalwood heartwood oil can be due to one or more components set forth in Table 1 acting either separately or together. Therefore, formulations that increase the concentration of the active component(s) and reduce the
 20 concentration of the inactive component(s) are set forth herein. Synthetic versions of the active components, or their derivatives, may be formulated in conjunction with or to replace the naturally occurring components of sandalwood heartwood oil. Compositions comprising purified or synthetic versions of an active component or derivative thereof can also be utilized in the methods set forth herein. For example, a composition comprising
 25 alpha-santalol (shown below) or beta-santalol (shown below) can be utilized to treat, prevent or reduce the recurrence of a Clostridium infection.



alpha-santalol



beta-santalol

In the methods and compositions set forth herein, sandalwood nut oil can be from the nut of any member of the genus *Santalum*. For example, and not to be limiting, East Indian sandalwood (*Santalum album*) or West Australian sandalwood (*Santalum spicatum*)

nut oil can be utilized in the methods and compositions set forth herein. Nut oil from *Santalum ellipticum*, *S. freycinetianum*, and *S. paniculatum*, the Hawaiian sandalwoods, can also be used. Typical analytical values for sandalwood nut oil are as follows:

Fatty Acid	% concentration
Palmitic (C16.0)	3.5%
Stearic (C18.0)	1.6%
Oleic (C18.1 c-9)	50.7%
Linoleic (C18.2 c-9 12)	1.1%
Linolenic (C18.3 c-9, 12, 15)	4.6%
Gondoic (C20.1 c-11)	3%
Ximenynic	35.8%

The therapeutically effective amount of sandalwood oil utilized in the compositions set forth herein can be, for example, a concentration greater of about 0.2% (w/w) and up to about 100% (w/w). For example, the therapeutically effective amount can be about 0.2%, 0.3%, 0.4%, 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1.0%, 1.1%, 1.2%, 1.3%, 1.4%, 1.5%, 1.6%, 1.7%, 1.8%, 1.9%, 2.0%, 2.1%, 2.2%, 2.3%, 2.4%, 2.5%, 2.6%, 2.7%, 2.8%, 2.9%, 3.0%, 3.1%, 3.2%, 3.3%, 3.4%, 3.5%, 3.6%, 3.7%, 3.8%, 3.9%, 4.0%, 4.1%, 4.2%, 4.3%, 4.4%, 4.5%, 4.6%, 4.7%, 4.8%, 4.9%, 5.0%, 5.1%, 5.2%, 5.3%, 5.4%, 5.5%, 5.6%, 5.7%, 5.8%, 5.9%, 6.0%, 6.1%, 6.2%, 6.3%, 6.4%, 6.5%, 6.6%, 6.7%, 6.8%, 6.9%, 7.0%, 7.1%, 7.2%, 7.3%, 7.4%, 7.5%, 7.6%, 7.7%, 7.8%, 7.9%, 8.0%, 8.1%, 8.2%, 8.3%, 8.4%, 8.5%, 8.6%, 8.7%, 8.8%, 8.9%, 9.0%, 9.1%, 9.2%, 9.3%, 9.4%, 9.5%, 9.6%, 9.7%, 9.8%, 9.9%, 10.0%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100% or any percentage (w/w) in between the percentages set forth herein.

As set forth above, the compositions set forth herein can comprise sandalwood heartwood oil, sandalwood nut oil or a combination of both. Therefore, the sandalwood oil can comprise only sandalwood heartwood oil, wherein the concentration of the sandalwood heartwood oil is from about 0.2% to about 100% (w/w). For example, the concentration of sandalwood heartwood oil can be about 0.2%, 0.3%, 0.4%, 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1.0%, 1.1%, 1.2%, 1.3%, 1.4%, 1.5%, 1.6%, 1.7%, 1.8%, 1.9%, 2.0%, 2.1%, 2.2%, 2.3%, 2.4%, 2.5%, 2.6%, 2.7%, 2.8%, 2.9%, 3.0%, 3.1%, 3.2%, 3.3%, 3.4%, 3.5%, 3.6%, 3.7%, 3.8%, 3.9%, 4.0%, 4.1%, 4.2%, 4.3%, 4.4%, 4.5%, 4.6%, 4.7%, 4.8%, 4.9%, 5.0%, 5.1%, 5.2%, 5.3%, 5.4%, 5.5%, 5.6%, 5.7%, 5.8%, 5.9%, 6.0%, 6.1%, 6.2%, 6.3%, 6.4%, 6.5%, 6.6%, 6.7%, 6.8%, 6.9%, 7.0%, 7.1%, 7.2%, 7.3%, 7.4%, 7.5%, 7.6%, 7.7%, 7.8%, 7.9%, 8.0%, 8.1%, 8.2%, 8.3%, 8.4%, 8.5%, 8.6%, 8.7%, 8.8%, 8.9%, 9.0%, 9.1%, 9.2%, 9.3%, 9.4%, 9.5%, 9.6%, 9.7%, 9.8%, 9.9%, 10.0%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%,

55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100% or any percentage (w/w) in between the percentages set forth herein.

The sandalwood oil can also comprise only sandalwood nut oil, wherein the concentration of the sandalwood nut oil is from about 0.2% to about 100% (w/w). For example, the concentration is sandalwood nut oil can be about 0.3%, 0.4%, 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1.0%, 1.1%, 1.2%, 1.3%, 1.4%, 1.5%, 1.6%, 1.7%, 1.8%, 1.9%, 2.0%, 2.1%, 2.2%, 2.3%, 2.4%, 2.5%, 2.6%, 2.7%, 2.8%, 2.9%, 3.0%, 3.1%, 3.2%, 3.3%, 3.4%, 3.5%, 3.6%, 3.7%, 3.8%, 3.9%, 4.0%, 4.1%, 4.2%, 4.3%, 4.4%, 4.5%, 4.6%, 4.7%, 4.8%, 4.9%, 5.0%, 5.1%, 5.2%, 5.3%, 5.4%, 5.5%, 5.6%, 5.7%, 5.8%, 5.9%, 6.0%, 6.1%, 6.2%, 6.3%, 6.4%, 6.5%, 6.6%, 6.7%, 6.8%, 6.9%, 7.0%, 7.1%, 7.2%, 7.3%, 7.4%, 7.5%, 7.6%, 7.7%, 7.8%, 7.9%, 8.0%, 8.1%, 8.2%, 8.3%, 8.4%, 8.5%, 8.6%, 8.7%, 8.8%, 8.9%, 9.0%, 9.1%, 9.2%, 9.3%, 9.4%, 9.5%, 9.6%, 9.7%, 9.8%, 9.9%, 10.0%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100% or any percentage (w/w) in between the percentages set forth herein.

The sandalwood oil can also comprise a combination of sandalwood heartwood oil and sandalwood nut oil, wherein the concentration of the sandalwood heartwood oil and the concentration of the sandalwood nut oil can each range from about 0.2% to about 99.8%, and the combined concentrations do not exceed 100% (w/w).

Provided herein is method of treating or preventing a Clostridium infection in a subject. The method comprises administering to the subject a composition with a therapeutically effective amount of sandalwood oil, wherein the subject has a Clostridium infection or is at risk of developing a Clostridium infection

Further provided is a method of reducing the recurrence of a Clostridium infection in a subject. The method comprises administering to the subject a composition including a therapeutically effective amount of sandalwood oil, wherein the subject is at risk for recurrence of a Clostridium infection

Clostridium is a genus of rod-shaped Gram-positive bacteria capable of producing endospores. The methods set forth herein can be used to treat the vegetative as well as the spore form of any *Clostridium* infection. Several *Clostridium* species are responsible for disease in humans. These include, but are not limited to, *Clostridium difficile* (*C. difficile*), *Clostridium botulinum* (*C. botulinum*), *Clostridium perfringens* (*C. perfringens*), *Clostridium tetani* (*C. tetani*) and *Clostridium sordelli* (*C. sordelli*).

C. difficile is the most serious cause of antibiotic-associated diarrhea and can lead to pseudomembranous colitis, a severe inflammation of the colon, often resulting from

eradication of the normal gut flora by antibiotics. In a very small percentage of the adult population, *C. difficile* bacteria naturally reside in the gut. Other people accidentally ingest spores of the bacteria while they are patients in a hospital, nursing home, or similar facility. When the bacteria are in a colon in which the normal gut flora has been destroyed, for
 5 example, after a broad-spectrum antibiotic has been used, the gut becomes overrun with *C. difficile*. This overpopulation is harmful because the bacteria release toxins that can cause bloating, diarrhea and abdominal pain.

C. botulinum produces neurotoxins that cause the flaccid muscular paralysis seen in botulism.

10 *C. perfringens* is a common cause of food-borne illness, with poorly prepared meat and poultry the main culprits in harboring the bacterium. The *clostridium perfringens* enterotoxin (CPE) mediating the disease is heat-labile (inactivated at 74°C) and can be detected in contaminated food, if not heated properly, and feces. Incubation time is between 6 and 24 (commonly 10-12) hours after ingestion of contaminated food. Often, meat is well
 15 prepared, but too far in advance of consumption. Since *C. perfringens* forms spores that can withstand cooking temperatures, if let stand for long enough, germination ensues and infective bacterial colonies develop. Symptoms typically include abdominal cramping and diarrhea; vomiting and fever are unusual.

C. tetani usually enters a host through a wound to the skin and then it replicates.
 20 Once an infection is established, *C. tetani* produces two exotoxins, tetanolysin and tetanospasmin. Tetanospasmin released in the wound is absorbed into the circulation and reaches motor nerves all over the body. The toxin acts at several sites within the central nervous system including nerve terminals in the spinal cord, brain, peripheral nerves and autonomic nervous system. The clinical manifestations of tetanus are caused when tetanus
 25 toxin blocks inhibitory impulses, by interfering with the release of neurotransmitters. These inhibitory neurotransmitters inhibit the alpha motor neurons. With diminished inhibition, the resting firing rate of the alpha motor neuron increases, producing rigidity, unopposed muscle contraction and spasm. Characteristic features are risus sardonicus (a rigid smile), trismus (commonly known as "lock-jaw"), and opisthotonus (rigid, arched back). Seizures
 30 may occur, and the autonomic nervous system may also be affected. Hypertension and tachycardia alternating with hypotension and bradycardia may develop.

The Clostridium infection can be in the digestive tract, for example, in the stomach, small intestine, lower intestine or colon of a subject. The infection can also be in the central or peripheral nervous system or autonomic nervous system of a subject. A Clostridium

infection can also be in the urinary tract of a subject. All strains are contemplated herein. A decrease or reduction of infection can occur *in vitro*, *ex vivo* or *in vivo*.

As utilized throughout, the term “infection” encompasses all phases of the Clostridium life cycle including, but not limited to, replication, spore formation, transition
5 from a spore form of the bacteria to a vegetative form of the bacteria, toxin production and receptor-mediated cell entry of toxins.

As used herein, the term subject can be a vertebrate, more specifically a mammal (e.g., a human, horse, pig, rabbit, dog, sheep, goat, non-human primate, cow, cat, guinea pig or rodent), a bird, a reptile or an amphibian. The term does not denote a particular age or
10 sex. Thus, adult and newborn subjects, whether male or female, are intended to be covered. As used herein, patient or subject may be used interchangeably and can refer to a subject with a disease or disorder. The term patient or subject includes human and veterinary subjects.

As used herein the terms treatment, treat, treating or ameliorating refers to a method
15 of reducing one or more signs or symptoms of the disease or condition. Thus in the disclosed method, treatment can refer to a 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% reduction or amelioration in the severity of an established disease or condition, as evidenced by a reduction in one or more signs or symptoms of the disease or condition. For example, and not to be limiting, a method for treating a Clostridium
20 infection is considered to be a treatment if there is a 10% reduction in one or more signs or symptoms of the infection in a subject as compared to control. For example, the method for treating a Clostridium infection is considered to be a treatment if there is a 10% reduction in one or more signs or symptoms of the Clostridium infection in a subject as compared to a control subject that did not receive a composition comprising sandalwood oil described
25 herein. Thus the reduction can be a 10, 20, 30, 40, 50, 60, 70, 80, 90, 100%, or any percent reduction in between 10 and 100 as compared to control levels. It is understood that treatment does not necessarily refer to a cure or complete ablation of the disease, condition, or symptoms of the disease or condition.

A subject can also harbor Clostridium resulting in carriage of the bacteria, which can
30 be asymptomatic. For example, the subject can harbor Clostridium spores that can germinate to become active, vegetative cells. Thus, also provided herein is a method of reducing Clostridium carriage comprising administering to the subject a composition comprising a therapeutically effective amount of sandalwood oil, wherein the subject is a carrier or is at risk of becoming a carrier for Clostridium. In this method, a reduction in

spore formation, a reduction in the number of spores and/or a reduction in the number of spores that germinate to become active, vegetative cells can occur. The reduction can be a 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% reduction as compared to a control.

5 Further provided is a method of reducing the infectivity of Clostridium spores, comprising contacting a population of Clostridium spores with an effective amount of sandalwood oil. In this method, the number of spores and/or the number of spores that germinate to become active, vegetative cells can be reduced. The reduction can be a 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% reduction as compared to a control.

10 The population of spores can be an *in vivo* population or an *in vitro* population.

As utilized herein, by reducing the recurrence of a Clostridium infection is meant a method of preventing, precluding, delaying, averting, obviating, forestalling, stopping, or hindering the onset, incidence or severity of the reappearance of a Clostridium infection in a subject. As utilized herein, by reappearance of a Clostridium infection is meant the
15 reappearance of one or more signs or clinical symptoms of a Clostridium infection after a period devoid of one or more clinical symptoms of a Clostridium infection. The recurrence of a Clostridium infection can be after treatment for a Clostridium infection, for example, after treatment of *C. difficile* with vancomycin, metronidazole and/or fidaxomicin. The recurrence of a Clostridium infection can also be after a remission. A recurrence can occur
20 days, weeks, months or years after treatment or after a remission. For example, the disclosed method is considered to reduce the occurrence of a Clostridium infection if there is a reduction or delay in onset, incidence or severity of the reappearance of a Clostridium infection, or one or more symptoms of a Clostridium infection in a subject at risk for a recurrence of a Clostridium infection compared to control subjects at risk for a recurrence of
25 a Clostridium infection that did not receive a composition comprising sandalwood oil disclosed herein. The disclosed method is also considered to reduce the recurrence of a Clostridium infection if there is a reduction or delay in onset, incidence or severity of the reappearance of a Clostridium infection, or one or more symptoms of a Clostridium infection in a subject at risk for recurrence of a Clostridium infection after receiving a
30 composition comprising sandalwood oil disclosed herein as compared to the subject's progression prior to receiving treatment. Thus, the reduction or delay in onset, incidence or severity of recurrence of a Clostridium infection can be about a 10, 20, 30, 40, 50, 60, 70, 80, 90, 100%, or any amount of reduction in between.

In the methods set forth herein, a composition comprising a therapeutically effective amount of sandalwood oil can selectively inhibit Clostridium infection in the subject. By selective inhibition is meant that inhibition of Clostridium infection occurs while maintaining substantial survival of the natural flora in the digestive tract of the subject.

5 For the administration methods disclosed herein, each method can optionally comprise the step of diagnosing a subject with a Clostridium infection or at risk of developing a Clostridium infection. The method can also include assessing the effectiveness of the sandalwood oil composition and modifying the treatment regimen.

The compositions set forth herein can be provided in a pharmaceutical composition.

10 The compositions include a therapeutically effective amount of the sandalwood oil in combination with a pharmaceutically acceptable carrier and, in addition, may include other medicinal agents, pharmaceutical agents, carriers, or diluents. Depending on the intended mode of administration, the pharmaceutical composition can be in the form of solid, semi-solid or liquid dosage forms, such as, for example, tablets, suppositories, pills, capsules,

15 powders, liquids, or suspensions, preferably in unit dosage form suitable for single administration of a precise dosage. By pharmaceutically acceptable is meant a material that is not biologically or otherwise undesirable, which can be administered to an individual along with sandalwood oil without causing unacceptable biological effects or interacting in a deleterious manner with the other components of the pharmaceutical composition in

20 which it is contained.

The compositions comprising a therapeutically effective amount of sandalwood oil can be concurrently administered with another pharmaceutical agent or agents. The pharmaceutical agent can be formulated with the therapeutically effective amount of sandalwood in a single composition or can be formulated in a separate composition. The

25 compositions comprising a therapeutically effective amount of sandalwood oil can also be administered subsequent to or prior to treatment with another pharmaceutical agent. For example, and not to be limiting, a composition comprising a therapeutically effective amount of sandalwood oil can be administered concurrently with, subsequent to, or prior to administration of vancomycin, metronidazole and/or fidaxomicin in order to treat, prevent

30 or reducing the recurrence of *C. difficile*.

Any appropriate route of administration may be employed, for example, administration can be systemic or local. Systemic administration includes administration via injection or infusion. Other routes of administration, such as intranasal, dermal, aerosol, vaginal, rectal or oral administration are also contemplated. By pharmaceutically acceptable

carrier is meant a material that is not biologically or otherwise undesirable, which can be administered to an individual along with the selected compound without causing unacceptable biological effects or interacting in a deleterious manner with the other components of the pharmaceutical composition in which it is contained.

5 Administration can be carried out using therapeutically effective amounts of the compositions described herein for periods of time effective to treat or reduce recurrence of a Clostridium infection. The effective amount may be determined by one of ordinary skill in the art and includes exemplary dosage amounts for a mammal of from about 0.5 to about 200mg/kg of body weight of active composition per day, which may be administered in a
10 single dose or in the form of individual divided doses, such as from 1 to 4 times per day. Such a dosing schedule can continue for days, weeks or months, depending on the subject and the type of infection. Alternatively, the dosage amount can be from about 0.5 to about 150mg/kg of body weight of active composition per day, about 0.5 to 100mg/kg of body weight of active composition per day, about 0.5 to about 75mg/kg of body weight of active
15 composition per day, about 0.5 to about 50mg/kg of body weight of active composition per day, about 0.5 to about 25mg/kg of body weight of active composition per day, about 1 to about 20mg/kg of body weight of active composition per day, about 1 to about 10mg/kg of body weight of active composition per day, about 20mg/kg of body weight of active composition per day, about 10mg/kg of body weight of active composition per day, or about
20 5mg/kg of body weight of active composition per day.

According to the methods taught herein, the subject is administered an effective amount of sandalwood oil. The terms effective amount and effective dosage are used interchangeably. The term effective amount is defined as any amount necessary to produce a desired physiologic response. Effective amounts and schedules for administering the
25 agent may be determined empirically, and making such determinations is within the skill in the art. The dosage ranges for administration are those large enough to produce the desired effect in which one or more symptoms of the disease or disorder are affected (e.g., reduced or delayed). The dosage should not be so large as to cause substantial adverse side effects, such as unwanted cross-reactions, anaphylactic reactions, and the like. Generally, the
30 dosage will vary with the activity of the specific compound employed, the metabolic stability and length of action of that compound, the species, age, body weight, general health, sex and diet of the subject, the mode and time of administration, rate of excretion, drug combination, and severity of the particular condition and can be determined by one of skill in the art. The dosage can be adjusted by the individual physician in the event of any

contraindications. Dosages can vary, and can be administered in one or more dose administrations daily, for one or several days. Guidance can be found in the literature for appropriate dosages for given classes of pharmaceutical products.

As used herein, the term carrier encompasses any excipient, diluent, filler, salt, buffer, stabilizer, solubilizer, lipid, stabilizer, or other material well known in the art for use in pharmaceutical formulations. The choice of a carrier for use in a composition will depend upon the intended route of administration for the composition. The preparation of pharmaceutically acceptable carriers and formulations containing these materials is described in, *e.g.*, Remington's Pharmaceutical Sciences, 21st Edition, ed. University of the Sciences in Philadelphia, Lippincott, Williams & Wilkins, Philadelphia Pa., 2005. Examples of physiologically acceptable carriers include buffers such as phosphate buffers, citrate buffer, and buffers with other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN[®] (ICI, Inc.; Bridgewater, New Jersey), polyethylene glycol (PEG), and PLURONICS[™] (BASF; Florham Park, NJ). Compositions suitable for parenteral injection may comprise physiologically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (propyleneglycol, polyethyleneglycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

These compositions may also contain adjuvants such as preserving, wetting, emulsifying, and dispensing agents. Prevention of the action of microorganisms can be promoted by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. Isotonic agents, for example, sugars, sodium chloride, and the like may also be included. Prolonged absorption of the injectable

pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

Solid dosage forms for oral administration of the compounds described herein or derivatives thereof include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the compounds described herein or derivatives thereof is admixed with at least one inert customary excipient (or carrier) such as sodium citrate or dicalcium phosphate or (a) fillers or extenders, as for example, starches, lactose, sucrose, glucose, mannitol, and silicic acid, (b) binders, as for example, carboxymethylcellulose, alignates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, (c) humectants, as for example, glycerol, (d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain complex silicates, and sodium carbonate, (e) solution retarders, as for example, paraffin, (f) absorption accelerators, as for example, quaternary ammonium compounds, (g) wetting agents, as for example, cetyl alcohol, and glycerol monostearate, (h) adsorbents, as for example, kaolin and bentonite, and (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, or mixtures thereof. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethyleneglycols, and the like. As set forth in the Examples, sandalwood oil can be formulated with, for example, polyethylene glycol 8000. This formulation can further comprise hydroxypropyl methylcellulose acetate succinate (HPMCAS-LF). For example, a formulation can comprise from about 40% to about 60% sandalwood oil, polyethylene glycol 8000 and HPMCAS-LF, wherein the polyethylene glycol 8000 and the HPMCAS-LF are present in the formulation in about a 2:1 ratio to about a 10:1 ratio. For example, the percentage of sandalwood oil in a formulation can be about 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59 or 60 %. The formulation can further comprise polyethylene glycol 8000 and HPMCAS-LF wherein the ration of polyethylene glycol 8000 to HPMCAS-LF is about a 2:1, 2.5:1, 3:1, 3.5:1, 4:1, 4.5:1, 5:1, 5.5:1, 6:1, 6.5:1, 7:1, 7.5:1, 8:1, 8.5:1, 9:1, 9.5:1 or a 10:1 ratio.

Solid dosage forms such as tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells, such as enteric coatings and others known in the art. They may contain opacifying agents and can also be of such composition that they release the active compound or compounds in a certain part of the intestinal tract in a delayed

manner. Examples of embedding compositions that can be used are polymeric substances and waxes. The active compounds can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration of the compounds described herein or derivatives thereof include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art, such as water or other solvents, solubilizing agents, and emulsifiers, as for example, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propyleneglycol, 1,3-butyleneglycol, dimethylformamide, oils, in particular, cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil, sesame oil, glycerol, tetrahydrofurfuryl alcohol, polyethyleneglycols, and fatty acid esters of sorbitan, or mixtures of these substances, and the like.

Besides such inert diluents, the composition can also include additional agents, such as wetting, emulsifying, suspending, sweetening, flavoring, or perfuming agents.

Ranges may be expressed herein as from about one particular value, and/or to about another particular value. When such a range is expressed, this includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent about, it will be understood that the particular value is disclosed.

Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application.

A number of aspects have been described. Nevertheless, it will be understood that various modifications may be made. Furthermore, when one characteristic or step is described it can be combined with any other characteristic or step herein even if the combination is not explicitly stated. Accordingly, other aspects are within the scope of the claims.

EXAMPLE I

Bacterial Cells and Maintenance

Bacterial *C. difficile* culture strains (BAA-1875, 43594, 43596) were obtained from the American Type Culture Collection (ATCC; Manassas, VA). The *C. difficile* cultures were maintained in a liquid complete growth medium composed of Reinforced Clostridial Medium (RCM) with 0.1% 2-mercaptoethanol (Beta-ME; BME) as a supplement. Cultures

were grown in a 37°C incubator under anaerobic conditions (tightly sealed gas jars with a gas mixture: 80% N₂, 10% CO₂, 10% H₂). For this study, cultures of the indicated organisms were grown for 48 hr in 14 ml snap-cap tubes containing 10 ml complete growth medium. Cultures were still actively growing in log phase and were triturated (in anaerobic gas) until homogeneous cell suspensions were obtained.

Quantified Cell Density

Cell suspensions were diluted 1/10 into RCM in polystyrene cuvettes and OD₆₀₀ readings were measured compared to the RCM blank. For each bacterial strain, a 40 ml suspension of a 0.02 OD₆₀₀ was then made in complete growth medium.

Serial Dilutions and Test Concentrations of Oils

Serial dilutions of test oils were prepared in sterile 15 ml tubes to final 2X concentrations as described in Table 2. Each oil was mixed with 50% glycerol plus 0.1% Tween 80 to a final volume of 1.0 ml at 20X final concentration, then vortexed to create an emulsion. To make a final volume of 10 ml at 2X final concentration, 9 ml of complete growth medium was added.

Each concentration was vortexed before distributing 1 ml of suspension to appropriately pre-labeled 5 ml snap-top culture tubes as per the Sample tube numbers (#1 to #96) shown in Tables 3-5. Control and test concentrations were distributed in duplicates, labeled “a” and “b”.

Table 2. Formulations, Volumes and Methods

Final 2X Concentration (%)	Final Volume RCM + Oil (ml)	Volume RCM (ml)	Transfer Volume (ml)	Volume Oil 20X Final Concentr'n (ul)	Volume Glycerol/Tween 80 (ul)
0	10	9	1	0.000	1000
0.0625	10	9	1	6.25	993.75
0.125	10	9	1	12.5	987.5
0.25	10	9	1	25	975
0.5	10	9	1	50	950
1	10	9	1	100	900
2	10	9	1	200	800
4	10	9	1	400	600
Total	80	72	8	793.75	7206.25

Inoculation with Test Organisms

The experiment was started by inoculating each of the 96 tubes as follows: 1.0 ml of each stock culture was distributed to the appropriate tubes as indicated (Tables 2-4). Tubes were monitored by visualization of individual tubes for growth, and pictures were taken of each individual tube after 4 days of incubation in complete growth medium alone, or in growth medium containing the test oils as described.

Results

Tables 3-5 summarize the results. *C. difficile* growth was seen in all control tubes by visual examination. However, the 3 species had different growth patterns: BAA-1875 grew as a flocculent mass from the bottom of the tube approximately $\frac{3}{4}$ up the liquid column, whereas the others grew as a flocculent mass (strain 43594) or more pelleted mass (strain 43596) on the bottom of the tube. All of the test concentrations of both *S. album* and *S. spicatum* oils inhibited growth in all 3 test species of *C. difficile*. Thus sandalwood oil is an effective antimicrobial against these organisms.

Table 3. Effects of Santalum species oils on growth of *C. difficile* strain BAA-1875.

Sample #	Tube	Species Oil	Final Test Concentration %	Replicate	Relative Growth
1		S. album	Control	a	+++
2			Control	b	+++
3			0.0313	a	0
4			0.0313	b	0
5			0.0625	a	0
6			0.0625	b	0
7			0.125	a	0
8			0.125	b	0
9			0.25	a	0
10			0.25	b	0
11			0.5	a	0
12			0.5	b	0
13			1.0	a	0
14			1.0	b	0
15			2.0	a	0
16			2.0	b	0
17		S. spicatum	Control	a	+++
18			Control	b	+++
19			0.0313	a	0
20			0.0313	b	0
21			0.0625	a	0
22			0.0625	b	0

Sample #	Tube	Species Oil	Final Test Concentration %	Replicate	Relative Growth
23			0.125	a	0
24			0.125	b	0
25			0.25	a	0
26			0.25	b	0
27			0.5	a	0
28			0.5	b	0
29			1.0	a	0
30			1.0	b	0
31			2.0	a	0
32			2.0	b	0

Table 4. Effects of Santalum species oils on growth of *C. difficile* strain 43594.

Sample #	Tube	Species Oil	Final Test Concentration %	Replicate	Relative Growth
33		S. album	Control	a	+++
34			Control	b	+++
35			0.0313	a	0
36			0.0313	b	0
37			0.0625	a	0
38			0.0625	b	0
39			0.125	a	0
40			0.125	b	0
41			0.25	a	0
42			0.25	b	0
43			0.5	a	0
44			0.5	b	0
45			1.0	a	0
46			1.0	b	0
47			2.0	a	0
48			2.0	b	0
49		S. spicatum	Control	a	+++
50			Control	b	+++
51			0.0313	a	0
52			0.0313	b	0
53			0.0625	a	0
54			0.0625	b	0
55			0.125	a	0
56			0.125	b	0
57			0.25	a	0
58			0.25	b	0
59			0.5	a	0
60			0.5	b	0

61	1.0	a	0
62	1.0	b	0
63	2.0	a	0
64	2.0	b	0

Table 5. Effects of Santalum species oils on growth of *C. difficile* strain 43596.

Sample #	Tube	Species Oil	Final Test Concentration %	Replicate	Relative Growth
65		S. album	Control	a	+++
66			Control	b	+++
67			0.0313	a	0
68			0.0313	b	0
69			0.0625	a	0
70			0.0625	b	0
71			0.125	a	0
72			0.125	b	0
73			0.25	a	0
74			0.25	b	0
75			0.5	a	0
76			0.5	b	0
77			1.0	a	0
78			1.0	b	0
79			2.0	a	0
80			2.0	b	0
81		S. spicatum	Control	a	+++
82			Control	b	+++
83			0.0313	a	0
84			0.0313	b	0
85			0.0625	a	0
86			0.0625	b	0
87			0.125	a	0
88			0.125	b	0
89			0.25	a	0
90			0.25	b	0
91			0.5	a	0
92			0.5	b	0
93			1.0	a	0
94			1.0	b	0
95			2.0	a	0
96			2.0	b	0

Comparison of Sandalwood Oil with Other Oils

Test oils were prepared at 2X final concentration. Briefly, each oil was diluted 1:00 in a sterile solution of 50% glycerol and 0.1% Tween 80 (1% final concentration of oil). This suspension was vortexed vigorously for 10 seconds and then further diluted 1:20 with
5 Proteose Fructose Neutral Red (PFNR) growth indicating medium (0.05% final oil concentration). This working stock was then diluted 2X with PFNR three more times to generate the 4 working concentrations at 2X the final concentration. The glycerol:Tween 80 solution was diluted 1:20 with PFNR followed by 3 more 2X dilutions to be inoculated as vehicle controls. All samples were vortexed between dilutions to ensure homogeneity of
10 the suspension. All work was performed within the anaerobic glovebox to ensure exclusion of oxygen in the test samples and growth medium.

100 µl of each suspension was plated per well in duplicate as described in Figure 1. 4 replicate plates were made. Bacterial cultures were maintained in PFNR broth in a 35°C incubator under anaerobic conditions (glovebox 80% N₂, 10% CO₂, 10% H₂). 24 hour
15 cultures of the indicated organisms grown in 14 ml snap-top tubes were pipetted up and down until a homogeneous suspension was obtained. Cell suspensions were diluted 1/10 into PBS in polystyrene cuvettes and OD₆₀₀ readings were measured compared to a PBS blank. 40 ml of a 0.02 OD₆₀₀ suspension was made for each bacterial strain in PFNR. 100
20 ul of 0.02 OD₆₀₀ cell suspension was plated on top of the wells preloaded with test suspensions. The final volume was 200 ul, with a cell concentration 0.01 OD₆₀₀, and each test suspension at 1X concentration. The first column of each plate contained 200ul of PFNR only as a negative control.

Plates were covered and then placed into the 35°C incubator within the glovebox and allowed to incubate overnight. After 14 hours of incubation the plates were removed
25 from the glovebox to photograph and read the absorbance at 450nm (yellow peak absorbance). The plates were replaced in the glovebox and allowed to incubate further. After a total of 38 hours of incubation the plates were removed from the glovebox again to photograph and the absorbance was read at 450nm. Figure 1 shows that both Western Australian sandalwood oil and East Indian sandalwood oil are effective antimicrobial
30 inhibitors of *C. difficile*.

EXAMPLE II

Strain

Clostridium difficile NCTC 13366 Ribotype 27 (vegetative bacteria and spores) was used. Vegetative cells of this strain were grown in Brain-Heart Infusion (BHIS) broth and

agar and 0.1% taurocholate. As explained below, the spore form of this strain was grown in BHIS broth and agar without 0.1% taurocholate.

Test Compound Preparation

200 μ L of sandalwood oil (test compound) were mixed with 100 μ L of DMSO and then combined with 4.7mL of 0.01% Tween 80 in 7H12. The resulting solution was composed of 4% sandalwood oil and 1% DMSO. Once the compound was assayed, the concentrations of sandalwood oil and DMSO were 2% and 1% respectively.

Minimum Inhibitory Concentration (MIC) Determination

The MIC of the test compound was determined according to CLSI guidelines for the microdilution method. Briefly, the test compound dissolved in appropriate broth growth medium was assayed in triplicate in a 2-fold dilution range from 2% to 0.0001%. A fresh working stock of organism was thawed and diluted to a concentration of approximately 5×10^5 CFU/mL and then added to all test and negative control wells and incubated for the appropriate time period and conditions. At the end of the incubation period, the test wells were visually scored for the lowest concentration of test or control compound to inhibit growth of the organism.

Spore Inhibition Assay

Taurocholate is necessary for *C. difficile* spores to transition (germinate) into vegetative bacteria. According to data from the MIC of vegetative *C. difficile*, the compound is inhibitory across the entire concentration range assayed with the organism in its vegetative state. This made it necessary to exclude taurocholate from the compound exposure portion of the assay. In order to test the inhibition of *C. difficile* spores to the test compound, exposure was carried out in BHIS broth without taurocholate to eliminate the possibility of spore germination. After a 24 hour exposure, an aliquot from each test well was transferred to BHIS agar plates containing taurocholate to allow the spores to germinate and then produce colonies. The aliquot was spread across the surface of the plate and then incubated for 48hrs in an anaerobic environment at 37°C. The colonies from each test sample were counted and compared to the number of colonies produced by the negative (untreated) controls. Test samples that resulted in colony counts lower than the negative control samples indicated inhibition of the *C. difficile* spores by the test compound. The number of colonies from the test samples were divided by the number of colonies from the negative control samples and then multiplied by 1.0 to produce a percent inhibition value.

<i>Clostridium difficile</i> NCTC 13366 Spore Inhibition Assay	
Test Compound Concentration	Percent Inhibition
0.5% - 0.0001%	no significant inhibition
1%	≥50%
2%	≥99%

EXAMPLE III

5 Formulations

Challenge Preparation: Challenge preparation was performed inside an anaerobe glove box in IW 305 in order to minimize oxygen exposure. A seed stock of the challenge organism (*C. difficile*) was thawed and added to the appropriate growth medium on Day -1. The organism was incubated anaerobically overnight (18-24 hrs) at 37°C. On the day of
10 challenge (Day 0), the culture was diluted in DPBS to achieve a target concentration of 10⁴ CFU/dose. Syringes with attached gavage needles were loaded with challenge material and placed on ice for transfer to the animal group.

Challenge Formulation Storage, Stability, and Handling: The challenge doses were kept on ice or refrigerated until dosing and considered stable throughout the dosing period.

15 Challenge Dose Formulation Concentration and Homogeneity Analysis: Challenge concentration was determined spectrophotometrically on the day of challenge and confirmed by retrospective plate count.

Polymer Carrier: The polymer carrier used in these studies was a composition comprising polyethylene glycol 8000 (PEG8000) and hydroxypropyl methylcellulose
20 acetate succinate (HPMCAS-LF). HPMCAS-LF was used to make the polymer insoluble at low pH. The formulation comprising a polymer carrier and sandalwood oil comprised about 50% PEG8000, about 41.67% sandalwood oil and about 8.33% HPMCAS-LF.

Test System

25 60 female C57BL/6 mice that weighed approximately 26-30 grams and were approximately 7-11 weeks old on Day 0 were used. The C57BL/6 mouse is susceptible to *C. difficile* and used extensively as a good animal model for the evaluation of antibacterial therapeutics. Treatments were administered via oral gavage and/or intraperitoneally in order to mimic treatment in humans. The challenge was administered via oral gavage as this is
30 both an effective and minimally invasive route of administration that mimics the natural route of infection.

Animals were housed under A/BSL2 containment conditions upon arrival at Southern Research Institute. Mice were group-housed (up to 5/cage) in micro-isolators. Appropriate bedding was used in the bottom of the cages. Analyses of the bedding, supplied by the vendor, was reviewed by Southern Research's veterinarian, or designee, to ensure
5 that no known contaminants were present that could interfere with or affect the outcome of the study.

Animals were housed in an environmentally monitored, well-ventilated room maintained at a temperature of 68-79 °F (approximately 20-26 °C) and a relative humidity of 30%-70%. Fluorescent lighting provided illumination, approximately 12 hours per day.

10 Mice were fed certified rodent chow #2016C (Harlan, Inc.; Madison, WI) during the pre-study and study periods. Feed was provided *ad libitum*. Analyses of the feed, provided by the manufacturer, were reviewed by Southern Research's Veterinarian, or designee, to ensure that no known contaminants were present that would be expected to interfere with or affect the outcome of the study.

15 Water (Birmingham public water supply) was supplied *ad libitum* during the pre-study and study periods. Samples of water from the animal facility were periodically analyzed, and the analysis was reviewed by Southern Research's Veterinarian, or designee, to assure that no known contaminants are present that would be expected to interfere with or affect the outcome of the study.

20 Animals were provided a psychological well-being program for social enrichment as approved by the Institutional Animal Care and Use Committee (IACUC). Animals were provided cage modifications and consumable enrichment for their psychological well-being.

Upon arrival, mice were housed in the AIBSL2 facility. Prior to the start of the study, the animals were observed for general health and acceptability for use in this study.

25 During Week -1, each mouse was uniquely identified by ear punch and an implantable micro-identification device (chip) (IPTT-300 Chip; Biomedic Data Systems Inc.; Seaford, DE) inserted subcutaneously on the back. The microidentification chip also provided body temperature data. Body temperature was taken rectally if chip or transponder was not available.

30 Experimental Design

The Provantis application (Instem Life Sciences Systems, Ltd.; Staffordshire, United Kingdom) was used for the direct on-line capture of most in-life data. Environmental monitoring of animal rooms (i.e., temperature/humidity and light/dark cycles) was performed using the Edstrom Watchdog system (Edstrom Industries, Inc.; Waterford, WI)

or manually. The remainder of the data was collected manually or by the appropriate automated system.

Randomization and Group Assignment: In order to obtain groups that are comparable by body weight, all mice were assigned to their respective treatment groups using a computer-generated randomization procedure. The body weights required for randomization were determined in Week -1 (the lowest body weights were excluded from randomization). After randomization, mice were assigned to one of six groups as indicated below in Table 6:

Table 6

Dose Group	Pre-Challenge Treatments				Clostridium difficile Challenge		Post-Challenge Treatments			No. of Mice
	Control Article	Dose Volume (mL)	Control Article	Dose Volume (mL)	Pathogenic Concentration	Dose Volume (mL)	Test/Control Articles	Dose Volume (mL)	Dose concentration	
1	Antibiotic Cocktail	0.3	Clindamycin	0.4	1.0×10^4	0.2	Untreated	NA	NA	10
2	Antibiotic Cocktail	0.3	Clindamycin	0.4	1.0×10^4	0.2	Vancomycin	0.3	50mg/kg	10
3	Antibiotic Cocktail	0.3	Clindamycin	0.4	1.0×10^4	0.2	Polymer Carrier	0.3	62.5mg/dose	10
4	Antibiotic Cocktail	0.3	Clindamycin	0.4	1.0×10^4	0.2	Sandalwood + Carrier	0.3	62.5mg/dose	10
5	Antibiotic Cocktail	0.3	Clindamycin	0.4	1.0×10^4	0.2	Sandalwood Oil	0.3	25mg/dose	10
6	Antibiotic Cocktail	0.3	Clindamycin	0.4	1.0×10^4	0.2	Sandalwood Oil	0.3	50mg/dose	10

10

Treatment Procedure: All treatment doses were vortexed before administration to animals. On Days -6, -5 and -4, animals were administered the antibiotic cocktail via oral gavage at a volume of 0.3 mL. Treatments were administered $\pm$ 1 hour from the initial day of dosing. On Day -1, animals were administered Clindamycin (10 mg/kg) via intraperitoneal (IP) at a volume of 0.4 mL. Post-challenge on Days 0 -4, animals were administered the appropriate test article as listed in the table above via oral gavage at a volume of 0.3 mL. Treatments were administered $\pm$ 1 hour from the initial day of dosing.

Challenge: On Day 0, mice were challenged via oral gavage with *Clostridium difficile* at a volume of 0.2mL. The bacterial suspension was supplied (and maintained) on ice in pre filled syringes and inverted prior to administration to each group. The challenge of mice was performed in the A/BSL2 laboratory within a primary containment device. All unused challenge material was placed on ice and given to the bacteriology group for subsequent disposal by addition of bleach and subsequent autoclaving.

Fasting: Animals were not fasted during the course of the study.

Clinical Observations

Daily Observations: All mice were observed twice daily throughout the prestudy and study period for signs of mortality and moribundity.

5 Detailed Observations: Detailed observations were recorded daily on Days -6 through Day 11 (or more often as warranted). Clinical signs include, but are not be limited to, hypoactive, hunched posture, poor grooming, diarrhea, squinting, neurological signs, or other abnormalities.

Body Weights: Body weights for all animals were obtained during Week -1 (randomization), then daily on Days -6 through Day 11 (or more often as warranted).

10 Body Temperatures: Body temperatures were obtained daily on Days -6 through Day 11 (or more often as warranted).

Fecal Samples: Fecal samples for each animal were collected on Days 1 -4 (or more often as warranted) for microbiology analysis (Section 11.10). An attempt was made to collect a minimum of 2 fecal pellets from each animal; if unable to collect a sample, a
15 culture swab was used and the sample was placed in DPBS. Collected fecal samples were divided into to two sterile polypropylene vials and stored at -70 °C (± 10 °C). In cases where solid pellets could not be collected, external swabs of the rectum were done.

Microbiology Analysis: To isolate *C. difficile*, fecal pellets or swabs were solubilized in PBS and then cultured onto CDSA. The agar plates were then placed in the
20 anaerobe chamber and incubated for 48-72 hours at 36 °C (± 10 °C). *C. difficile* colonies were identified by colony morphology. Results were recorded as positive (for the presence of *C. difficile* in the stool) or negative.

To determine the presence of *C. difficile* toxin, fecal pellets were qualitatively assayed by enzyme inununoassay. To isolate *C. difficile*, the tissues were manually
25 homogenized, serially diluted in sterile saline, plated in duplicate on *Clostridium difficile* Selective Agar (CDSA) and incubated for colony formation. The plates were counted to determine the number of CFU/tissue.

30 Results

As shown in Table 7, administration of sandalwood oil, alone or with a polymer carrier, resulted in decreased *C. difficile* infection as compared to untreated and placebo controls.

Table 7

Qualitative Fecal Pellet Analysis					
Group Number and Treatment	Animal Number	Day of Pellet Collection			
		Day 1	Day 2	Day 3	Day 4
1 Untreated	1	-	-	-	+
	2	-	-	+	+
	3	NA	NA	NA	NA
	4	-	+	+	-
	5	-	+	+	-
	6	+	+	-	-
	7	-	+	-	-
	8	-	+	-	-
	9	-	+	-	-
	10	-	+	-	-
2 Vancomycin	11	-	-	-	-
	12	-	-	-	-
	13	-	-	-	-
	14	-	-	-	-
	15	-	+	-	-
	16	-	-	-	-

	17	-	-	-	-
	18	-	-	-	-
	19	-	-	-	-
	20	-	-	-	-
3 Polymer Carrier (Placebo)	21	-	+	+	+
	22	+	+	+	+
	23	-	+	+	+
	24	-	+	+	+
	25	-	+	+	+
	26	-	+	+	-
	27	-	-	+	-
	28	-	+	-	-
	29	+	+	+	-
	30	-	-	-	-
4 Sandalwood + Carrier	31	-	-	+	-
	32	-	-	-	-
	33	-	-	-	-
	34	-	-	-	-
	35	-	-	-	-
	36	-	-	+	+
	37	NA	NA	NA	NA

	38	+	+	+	-
	39	-	+	-	-
	40	-	-	+	+
5 Oil (25mg)	41	-	-	+	NA
	42	-	-	+	-
	43	-	-	+	+
	44		-	-	+
	45	-	-	+	+
	46	-	-	+	+
	47	-	-	+	+
	48	-	-	+	+
	49	-	-	-	NA
	50	-	+	+	+
6 Oil (50mg)	51	NA	NA	NA	NA
	52	NA	NA	NA	NA
	53	NA	NA	NA	NA
	54	NA	NA	NA	NA
	55	NA	NA	NA	NA
	56	NA	NA	NA	NA
	57	-	NA	NA	NA
	58	NA	NA	NA	NA
	59	NA	NA	NA	NA
	60	NA	NA	NA	NA

What is claimed is:

1. A method of treating or preventing a Clostridium infection in a subject, said method comprising administering to the subject a composition comprising a therapeutically effective amount of sandalwood oil, wherein the subject has a Clostridium infection or is at risk of developing a Clostridium infection.
2. The method of claim 1, wherein the infection is in the digestive tract of the subject.
3. The method of claim 1 or 2, wherein the Clostridium infection is a *C. perfringens* infection, *C. difficile* infection, *C. tetani* infection or a *C. botulinum* infection.
4. The method of claim 1-3, further comprising selecting a subject with a Clostridium infection or at risk of developing a Clostridium infection.
5. The method of any of claims 1-4, wherein the sandalwood oil is from *Santalum album* or *Santalum spicatum* or a combination thereof.
6. The method of any of claims 1-5, wherein the therapeutically effective amount of sandalwood oil selectively inhibits Clostridium infection in the subject.
7. A method of reducing the recurrence of a Clostridium infection in a subject, comprising administering to the subject a composition comprising a therapeutically effective amount of sandalwood oil, wherein the subject is at risk for recurrence of a Clostridium infection.
8. The method of claim 7, wherein the infection is in the digestive tract of the subject.
9. The method of claim 7 or 8, wherein the Clostridium infection is a *C. difficile* infection.
10. The method of any of claims 7-9, wherein the subject has been treated with vancomycin, metronidazole or fidaxomicin.
11. The method of claim 10, wherein the composition comprising a therapeutically effective amount of sandalwood oil is administered concurrently with or subsequent to treatment with vancomycin, metronidazole or fidaxomicin.

12. The method of claim 11, wherein the composition comprising a therapeutically effective amount of sandalwood oil is administered subsequent to treatment with vancomycin, metronidazole or fidaxomicin.
13. The method of any of claims 1-12, wherein the sandalwood oil is from *Santalum album* or *Santalum spicatum* or a combination thereof.
14. The method of any of claims 1-13, wherein the composition comprises about 0.03% to about 100% sandalwood oil (w/w).
15. The method of claim 14, wherein the composition comprises about 0.03% to about 1% sandalwood oil (w/w).
16. The method of claim 15, wherein the composition comprises about 0.03% to about 0.5% sandalwood oil (w/w).
17. The method of any of claims 1-16, wherein the composition comprising the therapeutically effective amount of sandalwood oil further comprises a pharmaceutical carrier.
18. The method of claim 17 wherein the pharmaceutical carrier is a high molecular weight polyethylene glycol.
19. The method of any of claims 1-18, wherein the composition is administered as a tablet, a capsule, a liquid, or a suppository.
20. A method of reducing Clostridium carriage in a subject comprising administering to the subject a composition comprising a therapeutically effective amount of sandalwood oil, wherein the subject is a carrier or is at risk of becoming a carrier for Clostridium.
21. A method of reducing the infectivity of Clostridium spores, comprising contacting a population of Clostridium spores with an effective amount of sandalwood oil.
22. The method of claim 21, wherein the population is an *in vitro* population.
23. The method of claim 21, wherein the population is an *in vivo* population.

