



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/60, 15/82, 9/88 A01H 5/00	A1	(11) International Publication Number: WO 92/12249 (43) International Publication Date: 23 July 1992 (23.07.92)
(21) International Application Number: PCT/US91/09437 (22) International Filing Date: 17 December 1991 (17.12.91) (30) Priority data: 632,440 26 December 1990 (26.12.90) US (71) Applicant: MONSANTO COMPANY [US/US]; 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US). (72) Inventors: KLEE, Harry, John ; 802 Rotherham Drive, Ballwin, MO 63011 (US). KISHORE, Ganesh, Murthy ; 15354 Grantley Drive, Chesterfield, MO 63017 (US). (74) Agent: BOLDING, James, Clifton; Monsanto Company, 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US).		(81) Designated States: AT (European patent), AU, BB, BE (European patent), BG, BR, CA, CH (European patent), CS, DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), MC (European patent), NL (European patent), NO, PL, RO, SE (European patent), SU . Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: CONTROL OF FRUIT RIPENING AND SENESCENCE IN PLANTS (57) Abstract A method for controlling the ripening of fruits and vegetables as well as a method for controlling senescence of plant tissue is described. The method generally embraces the expression of an ACC metabolizing enzyme in the fruit or other desired plant tissue to inhibit the production of ethylene in the fruit or plant tissue. The use of the ACC metabolizing enzyme ACC deaminase is described in detail. The ripening or senescence process in the fruit or plant tissue is inhibited by the expression of the ACC deaminase gene such that the shelf-life and marketability of the fruit or plant is enhanced. The ACC metabolizing enzyme may be used in combination with other methods for reducing ethylene production in transformed plants to further reduce the production of ethylene in the fruit or plant. DNA constructs containing the ACC deaminase gene are also described.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
DE	Germany	MC	Monaco	TG	Togo
DK	Denmark			US	United States of America

CONTROL OF FRUIT RIPENING AND
SENESCENCE IN PLANTS

5 This is a continuation-in-part of our copending application having U.S. Serial No. 07/632,440 filed on December 26, 1990 entitled "Control of Fruit Ripening and Senescence in Plants."

10 Field of the Invention

 This invention relates in general to plant molecular biology and more particularly to a method for controlling the ripening of fruit and vegetables as well as controlling the effects of senescence in plants and recombinant DNA molecules capable of
15 affecting the desired control.

Background of the Invention

 One of the major problems facing the fruit, vegetable and cut flower industry is the loss of a considerable amount of goods
20 due to spoilage. It is estimated that 12 to 20 percent of the fruit and vegetable products become spoiled from the time they leave the farm until they get to the retail or processing outlets. In the cut flower industry, senescence (the wilting or dying) of the flower before it can be effectively marketed is a significant problem. The
25 spoilage or senescence process observed in fruits, vegetables and cut flowers results in a number of undesirable problems. Chief among these problems is the short harvesting season for the goods and the short shelf life of the goods following the harvest. Furthermore, these spoilage losses ultimately result in a higher
30 cost of the goods to the consumer.

-2-

A primary cause of the spoilage of fruits and vegetables is the natural ripening process of the fruit or vegetable. As the fruit or vegetable becomes more ripe it becomes softer and more easily bruised and susceptible to disease or other spoilage causing agents. It is known that ethylene production in the plant stimulates the fruit ripening process and is the key component in the ripening of fruits and vegetables. Others have attempted to control the ripening of fruits and vegetables in an attempt to extend the shelf life and/or harvesting season of the goods. Many of these attempts have been topical applications of chemicals to the fruit or vegetable itself. These chemical solutions have involved direct applications to the plant in the field or post-harvest applications to the fruit or vegetable itself. Several of these methods are discussed in United States Patent No. 4,957,757 or United States Patent No. 4,851,035. Due to the increasing importance of reducing additional stresses on the environment, a non-chemical means for controlling ripening would be advantageous and beneficial to the industry.

More recently, researchers have used a molecular biology approach to block ethylene synthesis in plants in an attempt to control the ripening of tomatoes. This approach involved transforming a tomato plant with an antisense gene that inhibited the synthesis of ethylene. The antisense gene produces (-) strand RNA that lowers the steady state levels of the (+) strand mRNA encoding a polypeptide involved in the conversion of 1-aminocyclopropane-1-carboxylic acid (ACC) to ethylene by the ethylene forming enzyme ACC oxidase. (Hamilton et al. 1990) While this method exhibits some degree of utility, it would be neither easy nor efficient to apply this technology to other plants, because the antisense gene would probably be species and gene

-3-

specific which would entail obtaining a different antisense gene for each species of plant desired to be transformed.

Thus a need exists in the fruit, vegetable and cut flower industries for a non-chemical method of controlling fruit ripening and senescence in plants that can easily and efficiently be utilized across a wide variety of plant species.

Summary of the Invention

A method for controlling the ripening of fruits and vegetables as well as a method for controlling senescence in cut flowers is presented. In general, the method involves expressing an ACC metabolizing enzyme in the desired plant tissue which lowers the level of ACC in the tissue which thereby reduces the level of ethylene in the desired plant tissue. More particularly, the method comprises transforming plant cells with a chimeric gene comprising a promoter that functions in plant cells to cause the production of an RNA sequence, a structural DNA sequence that causes the production of an RNA sequence that encodes an ACC deaminase enzyme and a 3' non-translated region that functions in plant cells to cause the addition of a stretch of polyadenyl nucleotides to the 3' end of the RNA sequence, with the promoter being heterologous with respect to the structural coding sequence, and then growing the plant to maturity. The expression of the ACC deaminase in the fruit delays the ripening process which provides an extended harvesting season and an extended shelf life for the goods. Likewise, expression of an ACC metabolizing enzyme in floral species suitable for use in the cut flower industry delays senescence of the flowers, thus extending the shelf life and marketability of the flowers.

-4-

In another aspect of the present invention, a recombinant, double stranded DNA molecule comprising a promoter that functions in plant cells to cause the production of an RNA sequence, a structural DNA sequence that encodes an ACC deaminase enzyme and a 3' non-translated region that functions in plant cells to cause the addition of a stretch of polyadenyl nucleotides to the 3' end of the RNA sequence, where the promoter is heterologous with respect to the structural DNA sequence, is also provided that enables one to obtain plants capable of expressing ACC deaminase in order to control ripening and senescence. The expression of the ACC deaminase in the plant cells extends the harvesting season and the shelf life of the goods by reducing the production of ethylene in the plants.

Among the many aims and objects of the present invention, one primary object is to provide a method of controlling ripening and senescence in plants utilizing a molecular biology technique that is efficiently and broadly applicable to many plant species.

Another object of the present invention is to provide a method for extending the harvesting season and shelf life of fruits, vegetables and flowers by controlling the production of ethylene in the plant by lowering the steady state levels of ACC using an ACC metabolizing enzyme, such as ACC deaminase or ACC malonyl transferase, expressed in the plant.

It is a further object of the present invention to reduce the synthesis of ethylene in plants by expressing the enzyme ACC deaminase in the plant.

It is still another object of the present invention to extend the market life of cut flowers by expressing the enzyme ACC

-5-

deaminase in the flower thereby reducing the senescence effects of ethylene synthesis in the flower.

It is a still further object of the present invention to provide transformed plants expressing an enzyme, ACC deaminase, in the plant so as to delay ripening of the fruit of the plant whether the fruit is allowed to ripen on the vine or if picked at an unripe stage of development to be ripened at a later time.

It is also a primary aim of the present invention to provide a fruit-bearing plant capable of expressing ACC deaminase specifically in the fruit of the plant.

Other and further objectives and aims of the invention will be made clear or become apparent from the following description and claims when read in light of the accompanying drawings.

Brief Description of the Drawings

Figure 1 illustrates the contents of the bacterial collection used to screen for ACC deaminase.

Figure 2 shows the nucleotide sequence of the ACC deaminase gene from *Pseudomonas chloroaphis* (isolate 6G5) (SEQ ID NO:1).

Figure 3 illustrates a plasmid map of pMON977.

Figure 4 illustrates a plasmid map of pMON10028.

Figure 5 illustrates a plasmid map of pMON10037.

Figure 6 illustrates a plasmid map of pMON10054.

Figure 7 illustrates a plasmid map of pMON11027.

Figure 8 illustrates a plasmid map of pMON7258.

Figure 9 illustrates a plasmid map of pMON11014.

Figure 10 illustrates a plasmid map of pMON981.

Figure 11 illustrates a plasmid map of pMON11016.

-6-

Figure 12 illustrates a plasmid map of pMON11032.

Figure 13 illustrates a plasmid map of pMON10086.

5 Figure 14 illustrates the nucleotide sequence of the fruit specific promoter E8 with the 5' HindIII and 3'BglII restriction sites underlined (SEQ ID NO:10).

Figure 15 illustrates the nucleotide sequence of the S-adenosyl methionine (SAM) decarboxylase gene (SEQ ID NO:9).

Figure 16 illustrates the nucleotide sequence of the ACC synthase gene (SEQ ID NO:8).

10 Figure 17 illustrates the nucleotide sequence of the ACC deaminase gene isolated from isolate 3F2. (SEQ ID NO:15)

Figure 18 illustrates graphically the relationship between the level of ethylene in control tomato fruit and transgenic tomato fruit expressing ACC deaminase.

15 Figure 19 illustrates a plasmid map of pMON11030.

Figure 20 illustrates the DNA sequence of the chloroplast transit peptide CTP2. (SEQ ID NO:13)


20 Figure 21 illustrates the DNA sequence of the CP4 synthetic 5-enolpyruvyl-3-shikimate phosphate synthase (EPSPS) gene. (SEQ ID NO:14)

Figure 22 illustrates the DNA sequence of a full-length transcript promoter from figwort mosaic virus (SEQ ID NO:17).

Detailed Description of the Preferred Embodiments

25 The metabolic pathway for the production of ethylene in plants is as follows:

-7-



10 In order to inhibit the biosynthesis of ethylene in plant tissues, one possible method would be to metabolize 1-aminocyclopropane-1-carboxylic acid (hereinafter ACC) and remove it from the metabolic pool. While it was unknown whether any ACC metabolizing enzyme would be capable of reducing the level of ACC sufficient to inhibit ethylene
15 biosynthesis, this approach was investigated. A number of enzymes are capable of metabolizing ACC. Examples of ACC metabolizing enzymes are ACC deaminase and ACC malonyl transferase. The ACC deaminase enzyme metabolizes ACC by converting it to α -ketobutyrate and ammonia. Thus, if the enzyme
20 ACC deaminase, or another ACC metabolizing enzyme, having sufficient kinetic capabilities can be expressed at sufficient levels in the plant, the synthesis of ethylene would be inhibited by the removal of ACC from the metabolic pool in the tissues where the ACC metabolizing enzyme is being expressed. A significant
25 aspect of the present invention is to provide a mechanism for delaying the ripening of fruit or senescence in plants by reducing the steady state levels of ACC in the plant tissues which reduces the level of ethylene in the plant tissues. It is preferred that the steady state concentrations of ethylene or ACC in the plant be

-8-

reduced by at least about 70% from normal levels in a non-modified cultivar. Preferably, the ethylene or ACC concentrations are reduced by at least about 90% from normal levels. It is believed that the reduction of the steady state levels of ACC or
5 ethylene in a plant or the fruit of a plant can be achieved by various methods, all of which are considered within the scope of the instant invention.

Regarding the delaying of ripening of fruit, it is preferred that the fruit be delayed from ripening on the vine by 1 to
10 30 days. This delay is to be measured from the onset of ripening and, specifically with respect to tomato, from when the fruit reaches the breaker stage of ripening. Likewise, the fruit is preferably delayed in ripening from 1 to 90 days following detachment from the vine and more preferably between 5 and 30
15 days. With respect to tomato, this delay in ripening is measured from the time of detachment of the fruit from the vine when the fruit is removed at the mature green or breaker stage of ripening. It is to be understood that the delay in ripening after detachment from the vine can be extended beyond the terms described by cold
20 storage or other methods known in the art.

The enzyme ACC deaminase was chosen for further experimentation. ACC deaminase is not known in the art to be produced or expressed naturally in plants. Therefore, in order to pursue a method of inhibiting ethylene synthesis in plants by
25 degrading ACC, an ACC deaminase encoding gene must be identified and then be made capable of being expressed in plants.

ACC deaminase is known to be expressed in certain microorganisms (Honma, M. and Shimomura, T. 1978). In order to isolate an ACC deaminase enzyme, a bacterial screen to isolate
30 bacteria expressing the enzyme can be designed to identify such a

-9-

bacteria or microorganism. Other methods for identifying an ACC deaminase enzyme, such as screening strains of yeast or fungi, would be equally applicable and routine to one of skill in the art. The following is a description of a bacterial screen that
5 identified bacteria expressing an ACC deaminase enzyme.

A collection of bacterial strains (Drahos, D. 1988) was screened for organisms that are capable of degrading ACC. This bacterial collection was composed of 597 microorganisms. The majority of the organisms were fluorescent *Pseudomonas* species
10 with the remaining being microbes typically found in the soil. A description of the bacterial collection is found in Figure 1. The screen was designed to select for microorganisms that would grow in a minimal medium containing ACC at 3.0 mM as the sole source of nitrogen. A sample of each bacteria in the bacterial
15 collection was grown individually in 96-well microliter dishes at 30°C for four days. Each well contained 0.2 ml of DF medium supplemented with ACC. DF medium was made by combining in 1 liter of autoclaved water, 1ml each of Reagent A, Reagent B, Reagent C and 5mg of thiamine HCl. Reagent A is made up of
20 1mg H_3BO_3 , 1mg $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$, 12.5mg $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 8mg $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ and 1.7mg $\text{NaMoO}_3 \cdot 3\text{H}_2\text{O}$ in 100mls of autoclaved water. Reagent B is made up of 0.1g $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ in 100mls of autoclaved water. Reagent C contains 20g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ in 100mls of autoclaved water. To the combined solution, carbon
25 sources glucose, gluconate and citrate are added to final concentrations of 0.1% (w/v) each, inorganic phosphate is added to a final concentration of 1.0mM (w/v) and ACC is added as the sole nitrogen source to a 3.0mM (w/v) final concentration. Finally,

-10-

Yeast Extract (DIFCO) is added to a final concentration of 0.01% (w/v).

Based on this screen, three organisms were identified as being capable of growing on ACC-containing medium. Their
5 ability to grow on ACC-containing minimal medium was confirmed by regrowth in 300 ml liquid cultures of the same medium. The two isolates that grew best on ACC were chosen for further characterization. These two isolates were designated 3F2 and 6G5. Both of these organisms were determined to be
10 *Pseudomonads* as was the organism not chosen for further characterization. Both of the selected organisms were screened for ACC deaminase enzyme activity by an in vitro assay described below. The 6G5 isolate was chosen for further experimentation. The 6G5 bacterium was identified as a *Pseudomonas chloroaphis*
15 strain by gas chromatography analysis of fatty acid methyl esters as described in Miller (1982). From the above screen results, it is apparent that other bacterial strains could be identified which degrade ACC by performing more extensive screens. Thus, other ACC deaminases and those identified in the screen but not
20 utilized for further experimentation are considered to be within the scope of the present invention.

A number of novel organisms capable of degrading ACC have also been isolated from diverse soil samples. These organisms were isolated on the basis of being able to grow on
25 minimal medium with ACC as the sole nitrogen source. Soil samples were collected from St. Charles (Missouri, USA), Sarawak (Malaysia), Iquitos (Peru), San Juan (Puerto Rico) and Mujindi (Tanzania). One gram of each soil sample was suspended into 99 ml of a Dilution buffer bottle (Fisher), shaken
30 well and the soil suspension was diluted 1:100 before plating.

-11-

Final dilution of the soil samples was 10^{-4} . One hundred (100) microliters of the diluted sample was spread on the isolation media in petri-plates (100X15 mm) with a hockey-stick glass rod. The isolation media contains a minimal salt base with K_2HPO_4 (10 g/L), $MgSO_4 \cdot 7H_2O$ (5 g/L), and trace metals: $FeSO_4$ (1 mg/L), $MnCl_2$ (1 mg/L), $CuSO_4$ (1 mg/L), $ZnSO_4$ (1 mg/L), $CaCl_2$ (1 mg/L). The pH of the base was adjusted to 7.0, before autoclaving, with 1N HCl. Noble agar (Difco) was used as the solidifying agent (1.5%). Any of the following three media may be used for isolation of ACC degrading microorganisms; (1) base + glucose (5 g/L) + ACC (0.1 to 1.0 g/L); (2) base + NH_4NO_3 (5 g/L) + ACC (1 g/L); (3) base + ACC (0.1 to 1.0 g/L). ACC, glucose, NH_4NO_3 were dissolved in distilled water, filter-sterilized and added into the autoclaved base media cooling at 50°C. Plates were incubated at 30°C for 1 week.

ACC was added to some of the soil samples obtained from St. Charles to enrich for ACC degrading bacteria in the soil. In these experiments, ACC (250 mg) was added into 50 ml of dilution buffer containing 0.5 g of St. Charles soil in a 250 ml Erlenmeyer flask. The flask was incubated on a rotary shaker (250 rpm, 30°C) for 3 days. The ACC enriched sample was then plated as previously described for non-enriched samples. Bacterial colonies capable of growth in the presence of ACC on plates were then isolated into pure cultures and grown in test tubes (20X150 mm) containing 5 ml of the following medium: KH_2PO_4 (4 g/L), K_2HPO_4 (6.5 g/L), $MgSO_4 \cdot 7H_2O$ (1 g/L), trace metals (same as isolation media), and ACC (0.3 g/L). Glucose (2 g/L) may be added to assist the growth of the bacteria. Bacterial strains which grew in the minimal salt medium with ACC as the sole carbon and nitrogen sources are listed in Table I.

-12-

TABLE I

	<u>Strain</u>	<u>Line #</u>	<u>Source</u>
5	388	B27444	St. Charles (ACC enriched)
	391	B27447	Malaysia
	392	B27448	Peru
	393	B27449	St. Charles
	401	B27457	St. Charles (ACC enriched)
10	T44	B27817	Tanzania
	PR-1	B27813	Puerto Rico

All of these organisms were shown to express ACC deaminase by two criteria. The first was that extracts from all of the organisms were capable of converting ACC to α -ketobutyric acid and the second was that all contained a protein of approximately 37,000 daltons that strongly cross-reacted with an antibody raised against the 6G5 ACC deaminase protein. To further demonstrate the equivalence of these organisms, kinetic parameters were determined for each of the isolated ACC deaminase enzymes.

The K_m for the ACC deaminases isolated from the various soil sources was determined using crude, desalted extracts. Individual strains of bacteria were grown in liquid media containing 4g KH_2PO_4 , 6.5g K_2HPO_4 , 1g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 2g glucose, 1mg FeSO_4 , 1mg MnCl_2 , 1mg ZnSO_4 , 1mg CuSO_4 , 1mg CaCl_2 , and 300mg ACC, all in 1 liter H_2O . Cells were grown for 2 to 3 days at 30°C. Cells were pelleted by centrifugation and resuspended in extraction buffer containing 0.1 M phosphate, pH 7.5, 1 mM EDTA, 0.1% β -mercaptoethanol. The cells were broken with a French Press, 1000 psi, and the cell debris was pelleted by centrifugation. The supernatants were desalted on Sephadex G-

-13-

25 columns pre-equilibrated with extraction buffer, which resulted in a crude, desalted extract. Glycerol was added to the extract (20% v/v) and enzyme solutions were stored at -20°C. ACC deaminase enzyme assays were conducted as described in the

5 Examples to follow. The assay mixture contained 100 μ l of 0.2 M Tris buffer, pH 8.0, 30 μ l of 500mM ACC solution, and enzyme solution to make a final volume of 200 μ l. Reactions were run for 10 minutes at 30°C. The reaction was stopped with 1.8 ml of 2 N HCl. After adding 300 μ l 0.1% 2,4-dinitrophenylhydrazine, the

10 mixture was incubated for 15 minutes at 30°C. The solution was then made basic by adding 2 ml of 2 N NaOH. The optical density of the resulting brownish-red solution was determined at 540 nm with a spectrophotometer.

The kinetic value, K_m , for ACC deaminase was

15 determined against ACC as the enzyme substrate for each of the ACC deaminases isolated. ACC deaminase activity was shown to be linear with respect to enzyme concentration using saturating levels of ACC (50 mM). An estimated K_m was determined for each extracted enzyme with ACC at sub-saturating concentrations.

20 Activity was shown to be linear over time with respect to ACC concentration for the concentrations used to determine the actual K_m values. Actual K_m values were then determined for each extract using ACC concentrations between 0.2X and 2X of the estimated K_m , or ACC concentrations between 1 and 10 mM ACC.

25 K_m values were calculated from double reciprocal plots, plotting the reciprocal of the substrate concentration on the x-axis and the reciprocal of the velocity (α -ketobutyrate formed) on the y-axis. The x-intercept (at y equals 0) is equal to $-1/K_m$. The K_m values for the ACC deaminases extracted from nine different strains were

-14-

determined and were generally within 3-fold of one another (from ~4 to ~12 mM). The K_m data demonstrates that essentially all ACC deaminases are functionally equivalent and can be used in the present invention. The K_m values for ACC deaminases from numerous isolates are listed in Table II.

TABLE II
Kinetic Values for Different Bacterial Isolates

	<u>Strain</u>	<u>K_m [mM ACC]</u>
10	6G5	9.0
	3F2	5.8
	388	8.6
	391	17.4
	392	7.1
15	393	5.9
	401	7.8
	T44	11.8
	PR-1	4.1

20 Once an isolate capable of degrading ACC is selected for further study, the gene encoding the ACC deaminase must be isolated. A general strategy for isolation and purification of the ACC deaminase gene from the selected *Pseudomonas* strain 6G5 is as follows. Isolate 6G5 is an exemplary embodiment for further
25 illustrative embodiments, but other isolates would be useful as well. A cosmid bank of the *Pseudomonas* strain 6G5 is constructed, cloned and introduced into *E. coli*. The clone carrying the ACC deaminase gene is identified by selection on minimal media containing ACC as the sole nitrogen source. The
30 coding region of the ACC deaminase gene is then identified and

-15-

sequenced. Cloning and genetic techniques, unless otherwise indicated, are generally those described by Sambrook et al. (1989). While this strategy was utilized to obtain the ACC deaminase gene from the 6G5 strain, other strategies could be employed with similar success and are considered to be within the scope of the invention. The detailed procedure for isolating the ACC deaminase gene from the 6G5 strain is set forth below.

The cell pellet from a 200 ml L-Broth (Miller 1972) late log phase culture of strain 6G5 was resuspended in 10 ml of Solution I (Birnboim and Doly 1979) in order to obtain chromosomal DNA. Sodium dodecylsulfate (SDS) is added to a final concentration of 1% and the suspension subjected to three freeze-thaw cycles, each consisting of immersion in dry ice for 15 minutes and in water at 70°C for 10 minutes. The lysate is then extracted four times with equal volumes of phenol:chloroform (1:1; phenol saturated with TE buffer at pH8.0) (TE = 10mM Tris; 1.0mM EDTA) and the phases separated by centrifugation (15000g; 10 minutes). The ethanol-precipitable material is pelleted from the supernatant by brief centrifugation (8000g; 5 minutes) following addition of two volumes of ethanol. The pellet is resuspended in 5 mls of TE buffer and dialyzed for 16 hours at 4°C against 2 liters of TE buffer. This preparation yields a 5 ml DNA solution of about 552 µg/ml.

Three 50 µg fractions of the *Pseudomonas* 6G5 DNA are then partially digested with EcoRI to generate fragments greater than 20 Kb. The three 50 µg fractions are digested with 0.125 units, 0.062 units, and 0.032 units, respectively, of EcoRI per µg DNA in a total volume of 1.25 ml each and incubated at 37°C for 30 minutes. The fractions are pooled and extracted once with an equal volume of 1:1 phenol:chloroform saturated with TE buffer at

-16-

pH 7.6 to remove the enzyme. The DNA is precipitated with two volumes of ethanol and pelleted by centrifugation (12000 g, 5 minutes). The dried DNA pellet is resuspended in 500 μ l TE buffer, and layered on top of a sucrose gradient. The 10%-40% sucrose gradient is prepared in seven 5.5 ml layers using 5% sucrose increments in 50 mM Tris pH8.0, 5 mM EDTA, 0.5 mM NaCl. The gradients are centrifuged at 26,000 rpm for 18 hours in a Beckmann SW28 rotor. The tube is punctured on the bottom and 1 ml fractions are collected. From each fraction, 20 μ l aliquots are run on a 1% agarose gel along with lambda DNA HindIII digested size standards. The fractions which contain DNA fragments greater than 20 Kb are combined. In the instant description, seven fractions were combined. The pooled sample is desalted and concentrated over Amicon Centricon-10® columns. The 0.5 ml concentrated sample is rinsed with 2 ml TE buffer, and again concentrated to 0.5 ml. The DNA sample is precipitated with 1 ml ethanol and the dry pellet resuspended in 50 μ l TE buffer. To estimate the DNA yield, 2 μ l of the sample is run on a 1% agarose gel along with 0.8 μ g lambda DNA cut with BstEII as a standard. From the gel, the concentration is estimated at 35 ng/ μ l of the *Pseudomonas* 6G5 DNA partial EcoRI fragments which are greater than 20 Kb.

A cosmid bank is constructed using the vector pMON17016. This vector is a derivative of the phage lambda cos plasmid pHC79 (Hohn and Collins 1980). The pMON17016 plasmid is constructed by introducing the HindIII-BglII fragment from pT7-7 (Tabor and Richardson 1985) containing the gene 10 promoter region from phage T7 into the HindIII-BamHI cut pHC79. The clone interrupts and inactivates the tetracycline resistance gene of pHC79 leaving the ampicillin resistance gene

-17-

intact. The introduced T7 promoter is not required for the function of the cosmid clone. The pMON17016 vector is cut with EcoRI and treated with calf alkaline phosphatase (CAP) in preparation for cloning. The vector and target sequences are
5 ligated as follows. 1.25 μg (25 μl of 50 $\text{ng}/\mu\text{l}$) of the pMON17016 vector DNA (EcoRI/CAP) is combined with 0.63 μg (18 μl of 35 $\text{ng}/\mu\text{l}$) of size fractionated 6G5 EcoRI fragments, and precipitated with two volumes of ethanol. The sample is centrifuged and the dry DNA pellet resuspended in 6 μl H_2O . To this solution, 1 μl of
10 the 10X ligation buffer (250 mM Tris-HCl pH 8.0, 100 mM MgCl_2 , 100 mM Dithiothreitol, 2 mM Spermidine), 2 μl of 100 mM ATP (Adenosine 5'-triphosphate) solution, and 1 μl of 400 unit/ μl T4 DNA ligase (New England Biolabs) is added. The ligation mix is incubated at room temperature (RT) for 6 hours.

15 From the 10 μl of pMON17016/6G5 ligated DNA sample, 3 μl is packaged into lambda phage particles (Stratagene; Gigapack Plus) using the manufacturer's procedure. To establish the cosmid titer, serial dilutions are made and used to infect the host bacteria. A culture of the host MM294 (Talmadge
20 and Gilbert 1980) *E. coli* is grown at 30°C in L-Broth containing 0.2% maltose. A 100 μl sample of MM294 is diluted with 100 μl SM buffer (SM = 50 mM Tris pH7.5, 100 mM NaCl, 8 mM MgSO_4 , 0.01% gelatin) and infected with 10 μl fractions of the packaged cosmid. The sample is incubated at RT for 15 minutes. One ml of
25 L-Broth is added to the sample and incubated at 37°C for 30 minutes. The infected bacteria are then concentrated by centrifugation (4000rpm, 4 minutes.) and plated on L-Broth agar plates containing 100 $\mu\text{g}/\text{ml}$ carbenicillin. The plates are incubated at 37°C overnight. The cosmid titer typically observed is

-18-

estimated at $\sim 8.5 \times 10^5$ clones total from the 3 μ l ligated pMON17016/6G5 DNA, or 2.8×10^6 clones per μ g 6G5 EcoRI DNA.

To select the cosmid clones which contain the ACC deaminase gene, the 6G5 library is then plated on media containing ACC as a sole nitrogen source. The plates contain 1.5% nitrogen free agar, 2 mM MgSO_4 , 0.2% glucose, 0.1 mM CaCl_2 , 1X M9 salts (M9 salts = 6 g $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 3 g KH_2PO_4 , 1.5 g NaCl , per liter), 1 mM Thiamine-HCl, 100 μ g/ml carbenicillin, and 3 mM ACC. The MM294 cells are infected with 35 μ l ($\sim 5.6 \times 10^4$ clones) packaged cosmid as described above, washed two times with 1X M9 salts, and plated on five plates. Growth was evident after a 3 day incubation at 37°C. After a 6 day incubation, approximately 300 cosmids (1 per 200) grew on the minimal media plates containing ACC as a sole nitrogen source. There is no growth evident after 6 days on the control plate which did not contain ACC as a supplemental source of nitrogen.

Several colonies that grew on the minimal media containing ACC are then screened. All the samples in the instant description had different size cosmid inserts and most contained several common EcoRI fragments. The three smallest clones are screened by restriction deletions and subcloning of the common fragments. The activity of the ACC deaminase gene is monitored by plating the clones on minimal media containing ACC as described above. The screens identified a clone containing a ~ 10.6 Kb insert which retained activity. The insert is then subcloned on a BamHI-XbaI fragment into the pUC118 plasmid (Viera and Messing 1987). Subsequent HindIII and SmaI deletions narrowed down the ACC deaminase activity to the 2.4 Kb insert which allowed the clone to grow on minimal media with ACC as the sole

-19-

nitrogen source. The pUC118 plasmid containing the 2.4 Kb insert is designated pMON10027.

Both strands of the 2.4 Kb insert of pMON10027 were then sequenced using the USB Sequenase® DNA sequencing kit following the manufacturer's directions. A 1017 base pair (bp) open reading frame was identified as the coding sequence of the ACC deaminase gene (Figure 2). This sequence is identified as SEQ ID NO:1.

To further demonstrate the equivalence of the ACC deaminase genes from different organisms, the DNA sequence of a second gene was determined. The *Pseudomonas* 3F2 isolate was identified in the initial screen as an organism capable of growth on medium containing ACC as sole nitrogen source as previously described. Conversion of ACC to α -ketobutyric acid *in vitro* (as described for the 6G5 organism) demonstrated that this organism also contained an ACC deaminase enzyme. The polymerase chain reaction (PCR) was used to clone the 3F2 ACC deaminase. Oligodeoxynucleotides for priming off of 3F2 DNA based on the known 6G5 sequence were designed. The sequences of the 5' and 3' oligonucleotides are as follows:

5' oligonucleotide:

CCCGGATCCATGAATCTGAATCGTTTT

(SEQ ID NO:11)

3' oligonucleotide:

CCCGGATCCGCCGTTACGAAACAGGAA

(SEQ ID NO:12)

These oligonucleotides begin with a sequence that incorporates a BamHI site into the PCR product to facilitate

-20-

subsequent cloning. Each is identical to either the 6G5 sequence over the first 18 (5') or last 18 (3') nucleotides, which are underlined. The 3F2 DNA was prepared as previously described for 6G5. The PCR reaction was carried out under conditions that
5 would permit annealing of the oligonucleotides to 3F2 DNA even if some mismatch between the 3F2 and 6G5 sequences existed. The PCR reaction was run for 30 cycles with 15 second extensions for each subsequent cycle. Each cycle consisted of:

	94°C	1 minute
10	40°C	2 minutes
	72°C	3 minutes plus 15 second extensions

The PCR-amplified 3F2 DNA contains the first 18 (5') and last 18 (3') nucleotides of isolate 6G5's ACC deaminase nucleotide sequence incorporated into the oligonucleotides and
15 thus may not correspond to the actual 3F2 gene in the areas of the first and last 18 nucleotides. Therefore, the actual identity of the first and last six amino acids of the 3F2 ACC deaminase may not be the same as the enzyme in the original 3F2 organism. Because a high degree of homology between the 3F2 DNA and the
20 oligonucleotide primers is essential for successful DNA amplification, the 3F2 and 6G5 sequences must be quite similar.

The product of the PCR amplification was cloned into BamHI-cut pBSSK+ (Stratagene) and subjected to dideoxy DNA sequencing as previously described. The sequence of the gene was
25 determined using a series of oligonucleotide primers derived from internal DNA sequences. The sequence of the 3F2 gene and the derived amino acid sequence of the ACC deaminase is shown in Figure 17. The nucleotide sequence is identified as SEQ ID NO:15 and the amino acid sequence is identified as SEQ ID NO:16. A
30 comparison of the derived amino acid sequences of the 6G5 and

-21-

3F2 enzymes indicates that they are highly homologous, having 96% identity and 99% similarity when conservative amino acid substitutions are considered. The sequence conservation, taken together with the kinetic data obtained on these two enzymes
5 clearly indicates the conserved nature of the ACC deaminase in nature.

Once an ACC deaminase gene has been identified and isolated, it must be engineered for plant expression. To introduce the ACC deaminase gene into a plant, a suitable chimeric gene
10 and transformation vector must be constructed. A typical chimeric gene for transformation into a plant will include a promoter region, a heterologous structural DNA coding sequence and a 3' non-translated polyadenylation site. A heterologous structural DNA coding sequence means a structural coding
15 sequence that is not native to the plant being transformed or a structural coding sequence that has been engineered for improved characteristics of its protein product. Heterologous with respect to the promoter means that the coding sequence does not exist in nature in the same gene with the promoter to which it is now
20 attached. Chimeric means a novel non-naturally occurring gene which is comprised of parts of different genes. In preparing the transformation vector, the various DNA fragments may be manipulated as necessary to create the desired vector. This includes using linkers or adaptors as necessary to form suitable
25 restriction sites or to eliminate unwanted restriction sites or other like manipulations which are known to those of ordinary skill in the art.

Promoters which are known or found to cause transcription of the ACC deaminase gene in plant cells can be
30 used in the present invention. Such promoters may be obtained

-22-

from plants, plant pathogenic bacteria or plant viruses and include, but are not necessarily limited to, the 35S and 19S promoters of cauliflower mosaic virus (CaMV35S and CaMV19S), the full-length transcript promoter from the figwort mosaic virus (FMV35S) and promoters isolated from plant genes such as EPSP synthase, ssRUBISCO genes and promoters obtained from T-DNA genes of *Agrobacterium tumefaciens* such as nopaline and mannopine synthases. The particular promoter selected should be capable of causing sufficient expression to result in the production of an effective amount of ACC deaminase to substantially inhibit the production of ethylene. Those skilled in the art will recognize that the amount of ACC deaminase needed to inhibit ethylene production may vary with the type of plant and the tissues within the plant of interest.

Particularly useful promoters for use in the present invention are fruit specific promoters which are expressed during ethylene production in the fruit and the full-length transcript promoter from the figwort mosaic virus (FMV35S). The FMV35S promoter is particularly useful because of its ability to cause uniform and high levels of expression of ACC deaminase in plant tissues. The DNA sequence of a FMV35S promoter is presented in Figure 22 and is identified as SEQ ID NO:17. Examples of fruit specific promoters include the E8, E4, E17 and J49 promoters from tomato (Lincoln, J.E., and Fischer, R.L. 1988), as well as the 2A11 promoter as described in U. S. Patent No. 4,943,674.

The promoters used for expressing the ACC deaminase gene of this invention may be further modified if desired to alter their expression characteristics. For example, the CaMV35S promoter may be ligated to the portion of the ssRUBISCO gene which represses the expression of ssRUBISCO in the absence of

-23-

light, to create a promoter which is active in leaves but not in roots. The resulting chimeric promoter may be used as described herein. As used herein, the phrase "CaMV35S" or "FMV35S" promoter includes variations of these promoters, e.g. promoters
5 derived by means of ligation with operator regions, random or controlled mutagenesis, addition or duplication of enhancer sequences, etc.

The 3' non-translated region contains a polyadenylation signal which functions in plants to cause the addition of
10 polyadenylated nucleotides to the 3' end of the RNA sequence. Examples of suitable 3' regions are the 3' transcribed, non-translated regions containing the polyadenylation signal of the tumor-inducing (Ti) plasmid genes of *Agrobacterium*, such as the nopaline synthase (NOS) gene, and plant genes like the 7s soybean
15 storage protein genes and the pea E9 small subunit of the RuBP carboxylase gene (ssRUBISCO).

The RNA produced by a DNA construct of the present invention also preferably contains a 5' non-translated leader sequence. This sequence can be derived from the promoter
20 selected to express the gene, and can be specifically modified so as to increase translation of the mRNA. The 5' non-translated regions can also be obtained from viral RNA's, from suitable eukaryotic genes, or from a synthetic gene sequence. The present invention is not limited to constructs, as presented in the
25 following examples, wherein the non-translated region is derived from the 5' non-translated sequence that accompanies the promoter sequence. Rather, the non-translated leader sequences can be part of the 5' end of the non-translated region of the native coding sequence for the heterologous coding sequence, or part of

-24-

the promoter sequence, or can be derived from an unrelated promoter or coding sequence as discussed above.

A DNA construct of the present invention can be inserted into the genome of a plant by any suitable method. Suitable plant transformation vectors include those derived from a Ti plasmid of *Agrobacterium tumefaciens*, such as those disclosed by Herrera-Estrella (1983), Bevan (1983), Klee (1985) and U.S. Patent No. 4,940,838. In addition to plant transformation vectors derived from the Ti or root-inducing (Ri) plasmids of *Agrobacterium*, alternative methods can be used to insert the DNA constructs of this invention into plant cells. Such methods may involve, for example, the use of liposomes, electroporation, chemicals that increase free DNA uptake, particle gun technology, and transformation using viruses. Methods for the introduction of vectors into maize, or other monocot cells would include, but are not limited to, the injection method of Neuhaus et al. (1987), the injection method of de la Pena et al. (1987) or the microprojectile methods of Klein et al. (1987) and McCabe et al. (1988).

The construction of vectors capable of being inserted into a plant genome via *Agrobacterium tumefaciens* mediated delivery is known to those of ordinary skill in the art. Typical plant cloning vectors comprise selectable and scoreable marker genes, T-DNA borders, cloning sites, appropriate bacterial genes to facilitate identification of transconjugates, broad host-range replication and mobilization functions and other elements as desired.

If *Agrobacterium* mediated delivery is chosen, once the vector has been introduced into the disarmed *Agrobacterium* strain, the desired plant can then be transformed. Any known

-25-

method of transformation that will work with the desired plant can be utilized.

Plants particularly suitable for use in this invention are tomato, banana, kiwi fruit, avocado, melon, mango, papaya, apple, peach, and other climacteric fruit plants. The present invention should also be suitable for use in the following non-climacteric species: strawberry, lettuce, cabbage, cauliflower, onions, broccoli, cotton, canola and oilseed rape. Other plant species that are affected by the ethylene induced ripening process may also benefit from the teachings of the present invention especially those in which ethylene production is critical to the growth of the plant or the ripening or development of the fruit of the plant. In the flower industry, particularly desirable flower species would be carnations, roses and the like. This list should be interpreted as only illustrative and not limiting in any sense.

In order to obtain constitutive expression of the ACC deaminase gene in plants, the gene was cloned into the transformation vector pMON977. The ACC deaminase gene isolated from the 6G5 isolate was used in the transformation vectors prepared herein. The pMON977 plasmid (Figure 3) contains the following well characterized DNA segments. First, the 0.93 Kb fragment isolated from transposon Tn7 which encodes bacterial spectinomycin/streptomycin resistance (Spc/Str), and is a determinant for selection in *E. coli* and *Agrobacterium tumefaciens* (Fling et al. 1985). This is joined to the chimeric kanamycin resistance gene engineered for plant expression to allow selection of the transformed tissue. The chimeric gene consists of the 0.35 Kb cauliflower mosaic virus 35S promoter (P-35S) (Odell et al. 1985), the 0.83 Kb neomycin phosphotransferase type II gene (NPTII), and the 0.26 Kb 3'-nontranslated region of

-26-

the nopaline synthase gene (NOS 3') (Fraley et al. 1983). The next segment is the 0.75 Kb origin of replication from the RK2 plasmid (ori-V) (Stalker et al. 1981). This is joined to the 3.1 Kb SalI to PvuI fragment from pBR322 which provides the origin of replication for maintenance in *E. coli* (ori-322), and the bom site for the conjugational transfer into the *Agrobacterium tumefaciens* cells. Next is the 0.36 Kb PvuI to BclI fragment from the pTiT37 plasmid, which contains the nopaline-type T-DNA right border region (Fraley et al. 1985). The last segment is the expression cassette consisting of the 0.65 Kb cauliflower mosaic virus (CaMV) 35S promoter enhanced by duplication of the promoter sequence (P-E35S) (Kay et al. 1987), a synthetic multilinker with several unique cloning sites, and the 0.7 Kb 3' nontranslated region of the pea *rbcS-E9* gene (E9 3') (Coruzzi et al. 1984 and Morelli et al. 1985).

Two different size fragments both containing the ACC deaminase gene from pMON10027 were introduced between the E35S promoter and the E9 3' end of pMON977. First, the 1071 bp EcoRV-SacI fragment from pMON10027 was introduced into the StuI-SacI cut pMON977, generating the pMON10028 vector (Figure 4). Second, the 1145 bp EcoRV-EcoRV fragment from pMON10027 was introduced into the StuI cut pMON977, generating the pMON10037 vector (Figure 5).

In order to construct vectors capable of directing expression of ACC deaminase specifically to fruit, a tomato fruit specific transcriptional promoter needed to be isolated. The promoter that was chosen is known to be induced to express at high levels in the presence of ethylene and is also known to be limited to the tomato fruit (Lincoln, J. and Fischer, R. 1988). The DNA sequence of the promoter for this gene, E8, has been published (Deikman et al. 1988). The DNA sequence of the E8

-27-

promoter is designated SEQ ID NO:10 and is illustrated in Figure 14. While this promoter was chosen, other fruit specific promoters would also be useful and their identification and isolation routine to one of ordinary skill in the art. The promoter
5 fragment E8 was isolated using standard polymerase chain reaction techniques. Oligonucleotides complementary to the E8 promoter were synthesized. The DNA sequences of the 5' and 3' oligonucleotides were as follows:

10 5' oligonucleotide:
 GAAGGAAGCT TCACGAAATC GGCCCTTATT C
(SEQ ID NO:2)

 3' oligonucleotide:
15 GGGGCTTTAG ATCTTCTTTT GCACTGTGAA TG
(SEQ ID NO:3).

The 5' oligonucleotide introduced a HindIII site approximately 1040 nucleotides 5' to the start of transcription.
20 The 3' oligonucleotide introduces a BglII site approximately 20 nucleotides beyond the start of transcription. The PCR product is an approximately 1060 nucleotide fragment that can be cloned as a HindIII to BglII fragment. This promoter fragment will confer tissue-specific expression upon any coding sequence placed
25 adjacent to it in an appropriate orientation.

The PCR reaction was performed essentially as recommended by the manufacturer of the GeneAmp kit (Perkin Elmer-Cetus). The reaction mix consisted of the following:

30

-28-

	water	58.5 μ l
	10X buffer	10 μ l
	dNTP mix	16 μ l
	5' primer	75 pM in 3.0 μ l
5	3' primer	75 pM in 3.0 μ l
	tomato DNA	1.24 μ g in 2 μ l
	Ampltaq DNA polymerase	0.5 μ l

- 10 The PCR reaction was run using the following temperature/time combination for 28 cycles:

	94°C	1 minute
	60°C	2 minutes
15	72°C	3 minutes.

Following completion, a PCR product of the correct size was observed. The fragment was purified by extraction with an equal volume of 1:1 phenol:chloroform followed by ethanol precipitation. The PCR fragment was then cut with HindIII and BglII so that it could be ligated to pMON10037 DNA. The PCR fragment was then ligated to pMON10037 DNA that had been cut with the same enzymes to remove the CaMV35S promoter sequence. The resulting plasmid contains the E8 promoter in the same location as the CaMV35S promoter of pMON10037 and was named pMON10054 (Figure 6).

Both of the pMON10028 and pMON10037 vectors can be mobilized into the ABI *Agrobacterium* strain. The ABI strain is the A208 *Agrobacterium tumefaciens* carrying the disarmed pTiC58 plasmid pMP90RK (Koncz and Schell 1986). The Ti

-29-

plasmid does not carry the T-DNA phytohormone genes, and the strain is therefore unable to cause the crown gall disease. Mating of pMON vectors into ABI is done by the triparental conjugation system using the helper plasmid pRK2013 (Ditta et al. 1980).
5 When the plant tissue is incubated with the ABI::pMON conjugate, the vector is transferred to the plant cells by the vir functions encoded by the disarmed pMP90RK Ti plasmid. The vector opens at the T-DNA right border region, and the entire pMON vector sequence is inserted into the host plant
10 chromosome. The Ti plasmid does not transfer to the plant cell but remains in the *Agrobacterium*.

The following examples further demonstrate several preferred embodiments of this invention. Those skilled in the art will recognize numerous equivalents to the specific embodiments
15 described herein. Such equivalents are intended to be within the scope of the claims.

Example 1

Transformed tobacco plants have been generated using
20 the ABI::pMON10028 and the ABI::pMON10037 vectors, to demonstrate the expression of the ACC deaminase gene in plants.

Tobacco cells were transformed using the tobacco leaf disc method. The tobacco leaf disc transformation protocol employed healthy leaf tissue about 1 month old. After a 15-20
25 minute surface sterilization with 10% Clorox plus a surfactant, the tobacco leaves were rinsed 3 times in sterile water. Using a sterile paper punch, leaf discs were punched and placed upside down on MS104 media (MS salts 4.3 g/l, sucrose 30 g/l, B5 vitamins 500X 2 ml/l, NAA 0.1 mg/l, and BA 1.0 mg/l) for a 1 day
30 preculture.

-30-

The discs were then inoculated with an overnight culture of disarmed *Agrobacterium* ABI containing the subject vector that had been diluted 1/5 (i.e. about 0.6 OD). The inoculation was done by placing the discs in centrifuge tubes with the culture. After 30 to 60 seconds, the liquid was drained off and the discs were blotted between sterile filter paper. The discs were then placed upside down on MS104 feeder plates with a filter disc to co-culture.

After 2-3 days of co-culture, the discs were transferred, still upside down, to selection plates with MS104 media. After 2-3 weeks, callus formed, and individual clumps were separated from the leaf discs. Shoots were cleanly cut from the callus when they were large enough to distinguish from stems. The shoots were placed on hormone-free rooting media (MSO: MS salts 4.3 g/l, sucrose 30 g/l, and B5 vitamins 500X 2 ml/l) with selection. Roots formed in 1-2 weeks. Any leaf callus assays were preferably done on rooted shoots while still sterile. Rooted shoots were placed in soil and were kept in a high humidity environment (i.e. plastic containers or bags). The shoots were hardened off by gradually exposing them to ambient humidity conditions.

In order to assay for ACC deaminase in the leaves, tobacco leaf samples were collected and frozen in liquid nitrogen. One gram of tissue was kept frozen under liquid nitrogen and ground to a fine powder. One ml of extraction buffer (100 mM Tris pH7.1, 10 mM EDTA, 35 mM KCl, 20% glycerol, 5 mM DTT, 5 mM L-ascorbate, 1 mM benzamidine, 1 mg/ml BSA) was added to the sample and ground for 45 seconds, then immediately centrifuged (12,000 g, 3 minutes) to remove the leaf debris. To remove small molecules, 250 μ l of the extract was run over a 1 ml Sephadex G-

-31-

50 spin column which was previously equilibrated with the above extraction buffer (less the BSA).

The extracts were assayed for the relative amount of the ACC deaminase enzyme activity in the transformed plant tissue.

- 5 The ACC deaminase enzyme converts the ACC substrate into α -ketobutyrate and ammonia. The α -ketobutyrate was reacted with 2,4-dinitrophenyl-hydrazine hydrochloride to form a hydrazone derivative whose optical density was measured at 520nm following addition of NaOH. The optical density values are a measure of the
- 10 amount of ACC deaminase in the plant extract. The assay reaction mix contained a 50 μ l sample of the tobacco leaf extract, 100 mM Tris pH8.6, and 50 mM ACC in a final volume of 150 μ l. The reaction was incubated at 30°C for 1 minute, and terminated with 50 μ l of 0.56 M HCl. A 0.6 ml aliquot of 0.1% 2,4-
- 15 dinitrophenyl-hydrazine in 2 N HCl was added. The sample was boiled for 2 minutes, cooled to room temperature, and 0.2 ml of 40% NaOH was added. A centrifugation (12,000 g, 5 minutes) removes the precipitate. The optical density of the supernatant was measured at 520nm, which indicated the relative amount of
- 20 the ACC deaminase enzyme being produced in the plants. Non-transformed tobacco plants were used as negative controls.

- Several tobacco leaf extracts were assayed and the ACC deaminase activity was found to range from 0.6 to 7.5mmoles product (α -ketobutyrate acid)/mg total protein/minute. These
- 25 assay results demonstrated that the ACC deaminase was being expressed in the tobacco plant.

Example 2

Transformed tomato plants have been generated using the ABI::pMON10028 and the ABI::pMON10037 vectors, and the expression of the ACC deaminase gene has been demonstrated in these plants.

Tomato plant cells were transformed utilizing the *Agrobacterium* strains described above generally by the method as described in McCormick et al. (1986). In particular, cotyledons were obtained from 7-8 day old seedlings. The seeds were surface sterilized for 20 minutes in 30% Clorox bleach and were germinated in Plantcons boxes on Davis germination media. Davis germination media is comprised of 4.3g/l MS salts, 20g/l sucrose and 10 mls/l Nitsch vitamins, pH5.8. The Nitsch vitamin solution is comprised of 100mg/l myo-inositol, 5mg/l nicotinic acid, 0.5mg/l pyridoxine HCl, 0.5mg/l thiamine HCl, 0.05mg/l folic acid, 0.05mg/l biotin, 2mg/l glycine. The seeds were allowed to germinate for 7-8 days in the growth chamber at 25°C, 40% humidity under cool white lights with an intensity of 80 einsteins m⁻²s⁻¹. The photoperiod was 16 hours of light and 8 hours of dark.

Once germination occurred, the cotyledons were explanted using a #15 feather blade by cutting away the apical meristem and the hypocotyl to create a rectangular explant. These cuts at the short ends of the germinating cotyledon increased the surface area for infection. The explants were bathed in sterile Davis regeneration liquid to prevent desiccation. Davis regeneration media is composed of 1X MS salts, 3% sucrose, 1X Nitsch vitamins, 2.0 mg/l zeatin, pH 5.8. This solution was autoclaved with 0.8% Noble Agar.

The cotyledons were pre-cultured on "feeder plates" composed of media containing no antibiotics. The media is

-33-

composed of 4.3g/l MS salts, 30g/l sucrose, 0.1g/l myo-inositol, 0.2g/l KH_2PO_4 , 1.45mls/l of a 0.9mg/ml solution of thiamine HCl, 0.2mls of a 0.5mg/ml solution of kinetin and 0.1ml of a 0.2mg/ml solution of 2,4 D. This solution was adjusted to pH 6.0 with KOH.

5 These plates were overlaid with 1.5-2.0 mls of tobacco suspension cells (TXD's) and a sterile Whitman filter which was soaked in 2COO5K media. 2COO5K media is composed of 4.3g/l Gibco MS salt mixture, 1ml B5 vitamins (1000X stock), 30g/l sucrose, 2mls/l PCPA from 2mg/ml stock, and 10 μ l/l kinetin from 0.5mg/ml

10 stock. The cotyledons were cultured for 1 day in a growth chamber at 25°C under cool white lights with a light intensity of 40-50 einsteins $\text{m}^{-2}\text{s}^{-1}$ with a continuous light photoperiod.

Cotyledons were then inoculated with a log phase solution of *Agrobacterium* containing the desired transgenic

15 gene. The concentration of the *Agrobacterium* was approximately 5×10^8 cells/ml. The cotyledons were allowed to soak in the bacterial solution for six minutes and were then blotted to remove excess solution on sterile Whatman filter disks and were subsequently replaced to the original feeder plate where they were

20 allowed to co-culture for 2 days. After the two days, cotyledons were transferred to selection plates containing Davis regeneration media with 2mg/l zeatin riboside, 500 μ g/ml carbenicillin, and 100 μ g/ml kanamycin. After 2-3 weeks, cotyledons with callus and/or shoot formation were transferred to fresh Davis

25 regeneration plates containing carbenicillin and kanamycin at the same levels. The experiment was scored for transformants at this time. The callus tissue was subcultured at regular 3 week intervals and any abnormal structures were trimmed so that the developing shoot buds would continue to regenerate. Shoots

30 developed within 3-4 months.

-34-

Once shoots developed, they were excised cleanly from callus tissue and were planted on rooting selection plates. These plates contained 0.5X MSO containing 50 μ g/ml kanamycin and 500 μ g/ml carbenicillin. These shoots formed roots on the selection media within two weeks. If no roots appeared after 2 weeks, shoots were trimmed and replanted on the selection media. Shoot cultures were incubated in percivals at a temperature of 22°C. Shoots with roots were then potted when roots were about 2cm in length. The plants were hardened off in a growth chamber at 21°C with a photoperiod of 18 hours light and 6 hours dark for 2-3 weeks prior to transfer to a greenhouse. In the greenhouse, the plants were grown at a temperature of 26°C during the day and 21°C during the night. The photoperiod was 13 hours light and 11 hours dark and the plants were allowed to mature.

Green tomato fruit and leaf samples were collected and frozen in liquid nitrogen. The samples were extracted and assayed using the procedures described for tobacco. The tomato extraction buffer contained 100 mM Tris pH7.1, 1 mM EDTA, 10% glycerol, 5 mM DTT, 5 mM L-ascorbate, 1 mM benzamidine, 1 mg/ml BSA. The extracts were assayed and the ACC deaminase activity was found to range from 1.6 to 11.2 mmoles of product/mg total protein/minutes reaction for the leaf tissue, and from 3.0 to 25.1 mmoles of product/mg total protein/minutes reaction for the tomato fruit tissue. The results of these assays demonstrated that the ACC deaminase was being expressed constitutively in the tomato plant.

-35-

Example 3

Tomato plants transformed with a chimeric gene encoding ACC deaminase have also been assayed to determine the effect of the expression of ACC deaminase on the ripening of fruit of the tomato plant.

Plasmids pMON10028 and pMON10037 were introduced into tomato (*Lycopersicon esculentum* cv. UC82B) as described in Example 2.

Plants containing the genes were initially identified by resistance to kanamycin. Kanamycin resistant plants were further analyzed by ACC deaminase enzyme assays (as described above) and by routine western blot analysis using antibody prepared against purified ACC deaminase protein. Plants that expressed the ACC deaminase protein were chosen for further analysis.

Tomato plants that were identified as expressing the ACC deaminase gene were examined for inhibition of fruit ripening. R1 progeny of the primary transformants from two lines, designated 5673 and 5854, as well as nontransformed UC82B plants were grown under identical conditions in a greenhouse. Progeny of the transgenic plants were screened for the presence of the NPTII gene, indicating inheritance of the T-DNA. All plants, including the UC82B controls, produced flowers and initiated fruit development simultaneously. Plants were then scored for the day at which fruit entered the breaker stage (the stage when the fruit begins to turn red), indicating initiation of ripening. Plants that had been scored as NPTII positive from both of the transgenic lines showed a significant delay in initiation of ripening. The delay in onset of ripening was approximately one week. Fruits from the transgenic plants as well as UC82B controls were then

-36-

removed from the plants at the breaker stage. Fruits were stored individually in 200 ml beakers at room temperature and allowed to ripen. The fruits from transgenic plants exhibited delays of from two to six weeks in the time it took to reach a fully ripe state.

5 Thus, tomato plants expressing the ACC deaminase gene exhibited delays in both the initiation of ripening and the time that it took to progress through the stages of ripening after the process had been initiated.

10 Example 4

Nicotiana tabacum plants transformed with pMON10028 and pMON10077 as described above have also been assayed to determine the effect of the expression of ACC deaminase in the plant on the life of the tobacco flowers. Tobacco plants expressing
15 the ACC deaminase gene were identified using the same enzyme assay as used for the tomato plants. Enzyme assays were performed on tobacco leaves and flowers. Plants expressing the gene were assayed for the length of time that flowers were retained. Flowers were tagged at the point of anthesis (flower
20 opening) and the time it took to reach a senesced stage was measured. While flowers from control plants showed significant wilting two days after anthesis, flowers from the transgenic plants expressing ACC deaminase were delayed in wilting by a full day.

25

Example 5

The present invention may also be used in combination with other methods known to delay ripening in fruits. One such combination involves use of the ACC deaminase gene in
30 combination with an antisense gene that inhibits ethylene

-37-

production. A plasmid containing ACC deaminase in combination with an antisense gene for the pTOM13 cDNA has been prepared for this purpose (Holdsworth et al. 1987). The gene designated pTOM13 has been previously shown to inhibit ethylene
5 production when placed in an antisense orientation in plants (Hamilton et al. 1980). It has been postulated that this gene encodes an enzyme that converts ACC to ethylene (presumably the enzyme is ACC oxidase) and inhibition of the synthesis of this
10 enzyme with an antisense RNA leads to accumulation of ACC in plant tissue. A cDNA clone corresponding to the pTOM13 gene was isolated from a cDNA library prepared from ripening tomato fruit on the basis of its ability to hybridize to synthetic oligonucleotides prepared from the published pTOM13 sequence.

A cDNA library was purchased from Stratagene (Cat. #
15 936004). This library was prepared from RNA isolated from ripening tomato fruit in the bacteriophage lambda cloning vector lambda-ZAP II. Oligonucleotide probes were prepared from segments of the pTOM13 published sequence as follows:

20 Oligonucleotide 1:

5' GGTGAACCAT GGAATTCCAC ATG 3' (SEQ ID
NO:4)

Oligonucleotide 2:

5' GCAATTGGAT CCCTTTCCAT AGC 3' (SEQ ID
25 NO:5)

Twenty thousand phage were plated on agar-containing plates as recommended by the manufacturer. The *E. coli* strain XL1-Blue, supplied by the manufacturer, was used for phage
30 preparation. Phage plaques were transferred to nitrocellulose

-38-

filters and baked in an 80°C oven for 2 hours. Plates were prehybridized at 65°C for 2 hours in the following solution:

5 6X SSC, 5X Denhardt's solution, 100 µg/ml denatured salmon sperm DNA, 20mM Tris:HCl, pH 7.0, 0.1% SDS, 1.0 mM EDTA.

10 50X Denhardt's Solution = 1.0 % each of Ficoll, polyvinylpyrrolidone, bovine serum albumin (Fraction V; Sigma) in water.

20X SSC = 175 g sodium chloride and 88.2 g sodium citrate per liter of water. pH adjusted to 7.0 with NaOH.

15 After prehybridization, ³²P-labelled oligonucleotides (Sambrook et al. 1989) were added to a final concentration of 500,000 cpm/ml hybridization solution for each oligonucleotide. Hybridization was performed at 50°C for 48 hours. Filters were washed twice in 6X SSC at room temperature for 15 minutes and
20 once at 50°C for 15 minutes. They were then dried and exposed to X-ray film for 48 hours. Plaques corresponding to hybridizing phage were isolated and purified by repeating the above procedure at a density of phage where single plaques could easily be separated from adjacent, non-hybridizing plaques. The pTOM13
25 cDNA insert was rescued in the plasmid vector pBS SK- as described by the manufacturer (Stratagene). This plasmid was designated pMON11023.

30 A vector designed for expression of the pTOM13 cDNA insert in an antisense orientation was then prepared. The cDNA insert with adjacent polylinker was excised from pMON11023 by

-39-

cutting with the restriction endonucleases BamHI and ClaI. The cDNA-containing portion of the plasmid was then cloned into pMON999 which had been cut with BglII and ClaI and treated with calf intestinal alkaline phosphatase. The resulting plasmid, pMON11025, contains the cDNA insert in an antisense orientation with respect to the CaMV35S promoter and a nopaline synthase 3' transcriptional terminator/polyadenylation site. This gene cassette can be excised as a single 2.2 kb NotI fragment. This NotI fragment was excised from pMON11025 and placed into the unique NotI site of pMON10028 to create pMON11027 (Figure 7). This plasmid thus contains an antisense pTOM13 gene and a CaMV35S/ACC deaminase gene. This plasmid was introduced into *Agrobacterium* ABI using triparental mating as described above and used to transform tomato plants.

The resulting transformed plants should significantly inhibit the production of ethylene in the plant. It is expected that the action of the ACC deaminase gene in combination with the pTOM13 antisense gene will virtually eliminate ethylene synthesis and should further delay ripening of the fruit. It is expected that the combination of the ACC deaminase and the pTOM13 antisense gene will exhibit synergistic properties in the reduction of the formation of ethylene in the fruit or plant.

Example 6

An alternate approach to reducing the rate of ethylene production in plant tissue involves overexpression of the gene encoding S-adenosylmethionine (SAM) decarboxylase. This enzyme degrades SAM which is the immediate precursor of ACC. The decarboxylated SAM is then converted to spermidine, a common polyamine. Since polyamines have themselves been

-40-

reported to have anti-senescence properties in plants, it is anticipated that SAM decarboxylase may prevent ripening in two ways 1) the production of spermidine and 2) degradation of a precursor to ethylene.

5 The gene encoding SAM decarboxylase (SEQ ID NO: 9), illustrated in Figure 15, has been cloned and its DNA sequence has been reported (Tabor and Tabor). The gene was cloned using PCR as described above in the protocol for isolation of the E8 promoter. *E. coli* DNA was purified as described above for the
10 isolation of *Pseudomonas* 6G5 genomic DNA. Purified DNA was subjected to PCR as described above using the following oligonucleotides as primers:

5' oligonucleotide:
15 GGAGAAGATA AGATCTATGA AAAAACTGAA
 ACTGC
 (SEQ ID NO:6)

3' oligonucleotide:
20 GCAGAAGTAA ATAGATCTGG CGGAGCC (SEQ ID
 NO:7).

25 The two primers used each introduced a BglII restriction site into the amplified DNA sequence to facilitate subsequent cloning steps. Following amplification, the DNA was cut with BglII and ligated with BglII cut pMON7258 (Figure 8). The resultant plasmid, pMON11014 (Figure 9), contained the SAM decarboxylase gene. The gene was subsequently cloned into plant transformation vectors that would permit expression of the gene
30 under the control of either a constitutive promoter such as the full

-41-

length transcript promoter from FMV or a fruit specific promoter such as the E8 promoter discussed above. The constitutive expression vector was constructed by cloning the pMON11014 BglII fragment containing SAM decarboxylase into BglII cut pMON981 (Figure 10). The resulting plasmid, pMON11016 (Figure 11), contained the gene in the correct orientation for expression in plants. The tissue specific expression vector, pMON11032 (Figure 12), was constructed by insertion of the same BglII fragment from pMON11014 into BamHI cut pMON10086 (Figure 13). Both transformation vectors were then introduced into *Agrobacterium* ABI using triparental mating. The *Agrobacterium* strains containing either pMON11016 or pMON11032 were then used to transform tomato plants as described above.

It is expected that plants expressing the ACC deaminase gene in combination with the SAM decarboxylase gene will inhibit synthesis of ethylene in plants, in a synergistic manner, such that the ripening or senescence process in the resulting plant is controlled to enhance the shelf life of the goods derived from the plant.

Example 7

An ACC metabolizing enzyme such as ACC deaminase may also be used in combination with an antisense ACC synthase gene. The DNA sequence for ACC synthase is known (Van Der Straeten et al. 1990) (SEQ ID NO:8) and is presented in Figure 16. Through routine manipulations, one can isolate a cDNA of the ACC synthase gene from a suitable cDNA library and prepare a vector containing the ACC synthase gene in an antisense orientation. This vector would contain the ACC synthase gene in

-42-

an antisense direction and an ACC metabolizing enzyme such as ACC deaminase in addition with the other DNA fragments necessary for successful plant transformation. Preferably, both the antisense ACC synthase and the ACC deaminase are under the transcriptional control of a fruit specific promoter, such as E8.

The resulting transformed plants should significantly inhibit the production of ethylene in the fruit of the plant transformed. It is expected that the action of the ACC metabolizing enzyme in combination with the ACC synthase antisense gene will virtually eliminate ethylene synthesis and further delay ripening of the fruit. The fruit may be ripened at a desired time by exposure of the fruit to ambient ethylene.

Example 8

This experiment was performed to evaluate the effect of reduction in ethylene levels in a plant when an ACC deaminase is expressed at high levels in the plant. Plant lines 5673 and 5854, as described in Example 3, were examined for ethylene generation in the plants and for phenotypic effects of expression of the ACC deaminase gene in the plant. Ethylene generation assays were performed on young leaf tissue from the plants by enclosing whole leaves or fruit in sealed containers and withdrawing 1.0 ml gas samples after one hour. Ethylene was quantified on a gas chromatograph (Ward et al. 1978) equipped with an alumina column and flame ionization detector. The results of ethylene generation assays are shown in Table 3 below.

-43-

TABLE 3
Ethylene Synthesis (nl/g/h)

<u>Plant</u>	<u>Leaf</u>	<u>Fruit</u>
UC82B	5.15±0.69	11.73±0.86
UB82B-2	5.53±0.37	ND
5673	0.60±0.09	1.43±0.36
5673-2	0.18±0.02	ND
5854	1.14±0.21	ND

(ND = not determined)

10

The ethylene level in plant line 5673 was reduced by 90% in one experiment utilizing young leaf tissue and by 97% in a second experiment. Plant line 5854 showed a reduction of approximately 78%. These data are consistent with the gene expression data in these plant lines. Line 5673 contained approximately 0.5% of the soluble protein as ACC deaminase while plant line 5854 contained approximately 0.05% of the soluble protein as ACC deaminase, as measured by protein gel blot analysis.

20

Protein gel blotting was performed by boiling protein samples for three minutes in the gel-loading buffer (50mM TrisCl, pH 7, 100mM dithiothreitol, 2% SDS, 10% glycerol, 0.1% bromophenol blue) and run on a 4-20% polyacrylamide MINI-PROTEAN II ready gels (BIO-RAD). The protein was transferred to nitrocellulose membrane using a MilliBlot-SDE electroblotting apparatus (Millipore, Bedford, MA) following the manufacturers directions. The membrane was incubated overnight at 4°C in 1% BSA, TBST (10 mM Tris, pH 8, 150 mM NaCl, 0.05% Tween-20). The incubations were performed at room temperature with gentle agitation to hybridize the membrane. The primary ACC

30

-44-

deaminase antibody was bound by incubating the membrane in a 1:1000 dilution of the goat serum in TBST for one hour. This was followed by three 10 minute washes in TBST. The secondary reagent was bound by incubating the membrane with 5 μ C of 125 I-labelled protein G in 20 ml of TBST for 30 minutes. The membrane was washed four times for 10 minutes with 0.1% Triton X-100 and exposed to film. Antibodies were obtained to the ACC deaminase protein by injecting a goat with 1.5 mg of protein and isolating antibodies from the goat pursuant to standard techniques known to those skilled in the art.

Homozygous plants from plant line 5673 were also examined for phenotypic effects. Seed from the transgenic plants germinated normally, and plants were phenotypically indistinguishable from controls. The plants exhibited no delay in the onset of flowering or ripening. They did, however, show significant differences in the progression of ripening. The fruits of transgenic plants exhibited a peak of ethylene synthesis concomitant with control fruit, but at a level of only 10% that of controls. This is illustrated in Figure 18. Ethylene generation by transgenic plants is represented by -•- and ethylene generation by control plants (UC82B) is represented by -■-. The bars represent means $\pm$ standard error at specific time points. The fruit was detached at the breaker stage and ethylene generation measured daily as previously described. The delay in ripening of fruits detached at the breaker stage was also significant. Control fruit passed from breaker to fully red in seven days and exhibited a marked degree of softening after only two weeks. Transgenic tomato fruit reached the fully red stage after 24 days and remained firm for an extended period from the breaker stage. Fruit from transgenic plants remained firm for longer than 40

-45-

days and did not abscise while the control fruit had abscised after 14 days. These data are presented in Table 4.

TABLE 4

<u>Plant</u>	<u>Ripening Stage</u>			
	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>
Transgenic	2.8±0.53	5.3±0.98	11.3±3.1	23.5±3.8
Control	1.4±0.19	2.8±0.26	5.1±0.45	7.0±0.53

The data in Table 4 are expressed as the number of days to reach a particular ripening stage after being detached, with a standard error. Ripening stages were defined as follows: Breaker, first sign of color change; 3, fully orange; 4, orange to red; 4, greater than 50% red; 6, fully red.

Example 9

This example illustrates the expression of the ACC deaminase protein in a flowering plant species. The ACC deaminase gene was transformed into petunia plants. The petunia plants were transformed with a transformation vector that allows for the direct selection of transformed plants on glyphosate. Petunia explants were generally prepared for pre-culture as described for the tobacco plants in Example 1. Leaves from a one month old petunia plant were surface sterilized for fifteen minutes in a solution of 10% Clorox plus surfactant and washed three times with distilled water. The explants were cut in 0.5 cm squares, removing the leaf edges, mid-rib, tip, and petiole end for uniform tissue type. The explants were then placed in a single layer, upside down, on MS104 plates containing 2 mL 4COO5K media to moisten the surface and pre-cultured for 1-2

-46-

days. Explants were inoculated using an overnight culture of *Agrobacterium* containing the plant transformation vector that has been adjusted to a titer of 1.2×10^9 bacteria/mL with 4COO5K media. Explants were placed into a centrifuge tube, the *Agrobacterium* suspension was added and the mixture of bacteria and explants was "vortexed" on maximum setting for 25 seconds to insure even penetration of bacteria. The bacteria were poured off and the explants were blotted between layers of dry sterile filter paper to remove excess bacteria. The blotted explants were placed upside down on MS104 plates to which 2mL 4COO5K media and a filter disk have been placed on top of the agar and co-cultured for two to three days. The explants were transferred to MS104 plates containing carbenicillin 1000mg/l and cefotaxime at 100mg/l for 3 days. The explants were then transferred to a new MS104 media that contains glyphosate at 0.05mM, carbenicillin at 1000mg/l and cefotaxime at 100mg/l for the selection phase. At 4-6 weeks, shoots were cut from callus and placed on MSO and carbenicillin at 500mg/l rooting media. Roots formed in 3-5 days, at which time leaf pieces were taken from rooted plates to confirm glyphosate tolerance and that the material was transformed.

The petunia plants were transformed with plant transformation vector pMON11030. A map of pMON11030 is presented in Figure 19. This plasmid is essentially composed of the previously described bacterial replicon system that enables this plasmid to replicate in *E. coli* and to be introduced into and to replicate in *Agrobacterium*. Referring to Figure 19, this plasmid additionally contains the bacterial spectinomycin-/streptomycin selectable marker gene (Spc/Str), and located between the T-DNA right border and left border is the CTP2-CP4 synthetic 5-enolpyruvyl-3-shikimate phosphate synthase (EPSPS) gene in the

-47-

FMV35S promoter-E9 3' cassette. The CTP2-CP4 synthetic gene permits for selection of transformed cells by their ability to grow in the presence of glyphosate. The CTP2 is a chloroplast transit peptide and its DNA sequence is presented in Figure 20 (SEQ ID NO:13). The DNA sequence of the CP4 EPSPS, a gene capable of conferring resistance to glyphosate, is presented in Figure 21 (SEQ ID NO:14). The ACC deaminase gene from isolate 6G5 was placed between the FMV promoter and a nopaline synthase 3' region as a 2.0 kb NotI fragment into the unique NotI site to create pMON11037.

The presence of the ACC deaminase protein in transformed petunia tissues has been confirmed by immunoblot analysis of leaf discs as described in Example 8. ACC deaminase protein has been detected in leaf tissues in five out of six regenerated petunia plants.

Ethylene levels of transgenic petunia plants transformed with pMON11030 have also been determined in petunia plants expressing ACC deaminase. The level of ethylene in the plant is reduced to about one-half of the ethylene level in a control plant that has not been transformed. The results of ethylene generation assays are presented in Table 5 below.

TABLE 5
ETHYLENE SYNTHESIS (nl/g/h)

Plant Line	Leaf Tissue
35861	0.58
35860	0.53
35862	0.62
Control	1.09

-48-

These data illustrate that transgenic plants expressing the ACC deaminase protein have reduced ethylene levels in leaf tissues. It is expected that such plants will show reduced senescence of flowers and leaves when compared to
5 nontransformed plants.

All publications and patents mentioned in this specification are herein incorporated by reference as if each individual publication or patent was specifically and individually stated to be incorporated by reference.

10 From the foregoing, it will be seen that this invention is one well adapted to attain all the ends and objects hereinabove set forth together with advantages which are obvious and which are inherent to the invention.

It will be understood that certain features and
15 subcombinations are of utility and may be employed without reference to other features and subcombinations. This is contemplated by and is within the scope of the claims.

Since many possible embodiments may be made of the invention without departing from the scope thereof, it is to be
20 understood that all matter herein set forth or shown in the accompanying drawings is to be interpreted as illustrative and not in a limiting sense.

BIBLIOGRAPHY

- 5 Birnboim, H. C. and Doly, J. (1979) A rapid alkaline extraction
procedure for screening recombinant plasmid DNA. Nucl. Acids.
Res. 7:1513-1525.
- 10 Coruzzi, G., Broglie, R., Edwards, C., and Chua, N.H. (1984).
Tissue-specific and light-regulated expression of a pea nuclear
gene encoding the small subunit of ribulose-1,5-bisphosphate
carboxylase. EMBO J 3, 1671-1679.
- 15 Deikman, J. and Fischer, R. (1988). Interaction of a DNA binding
factor with the 5'-flanking region of an ethylene-responsive fruit
ripening gene from tomato. EMBO J. 7, 3315-3320.
- de la Pena, A., Lorz, H. and Schell, J. (1987) Nature 325:274-276.
- 20 Ditta, G., Stanfield, S., Corbin, D., and Helinski, D.R. (1980).
Broad host range DNA cloning system for Gram-Negative
bacteria: construction of a gene bank of *Rhizobium meliloti*. Proc
Natl Acad Sci USA 77, 7347-7351.
- 25 Drahos, D., Barry, G., Hemming, B., Brandt, F., Skipper, H.,
Kline, E., Kluepfel, D., Hughes, T., and Gooden, D., in The
Release of Genetically-Engineered Microorganisms. (1988).
Sussman, M., Collins, C., Skinner, F. and Stewart-Tull, D. eds.
Academic Press, New York.
- 30 Fling, M.E., Kopf, J., and Richards, C. (1985). Nucleotide
sequence of the transposon Tn7 gene encoding an aminoglycoside-

-50-

modifying enzyme, 3'(9)-O-nucleotidyltransferase. Nucleic Acids Research 13 no.19, 7095-7106.

- 5 Fraley, R.T., Rogers, S.G., Horsch, R.B., Sanders, P.R., Flick, J.S., Adams, S.P., Bittner, M.L., Brand, L.A., Fink, C.L., Fry, J.S., Galluppi, G.R., Goldberg, S.B., Hoffmann, N.L., and Woo, S.C. (1983). Expression of bacterial genes in plant cells. Proc Natl Acad Sci USA 80, 4803-4807.
- 10 Fraley, R.T., Rogers, S.G., Horsch, R.B., Eichholtz, D.A., Flick, J.S., Fink, C.L., Hoffmann, N.L., and Sanders, P.R. (1985). The SEV system: a new disarmed Ti plasmid vector system for plant transformation. Bio/Technology 3, 629-635.
- 15 Hamilton, A., Lycett, G. and Grierson, D. (1990). Antisense gene that inhibits synthesis of the hormone ethylene in transgenic plants. Nature 346:284-287.
- 20 Hohn, B. and Collins J. (1980) A small cosmid for efficient cloning of large DNA fragments. Gene 11: 291-298.
- Holdsworth, M. Schuch, W. and Grierson, D. (1987). Nucleotide sequence of an ethylene-related gene from tomato. Nucleic Acids Res. 15:10600
- 25 Honma, M. and Shimomura, T. (1978). Metabolism of 1-Aminocyclopropane-1-carboxylic Acid. Agric, Biol. Chem. 42(10), pp 1825-1831.

-51-

Kay, R., Chan, A., Daly, M., and McPherson, J. (1987). Duplication of the CaMV 35S promoter sequence creates a strong enhancer for plants. *Science* 236, 1299-1302.

5 Klein, T.M., Wold, E.D., Wu, R. and Sanford, J.C. (1987) Nature 327:70-73.

Koncz, C., and Schell, J. (1986). The promoter of TL-DNA gene 5 controls the tissue-specific expression of chimeric genes carried
10 by novel type of Agrobacterium binary vector. *Mol Gen Genet* 204, 383-396.

Lincoln, J. and Fischer, R. (1988). Diverse mechanisms for the regulation of ethylene-inducible gene expression. *Mol Gen Genet*
15 212, 71-75.

McCabe, D.E., et al. (1988) Bio/Technology 6:923.

20 Miller, J. H. (1972). *Experiments in Molecular Genetics*. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York.

Miller, L. T. (1982). Single derivatization method for routine analysis of bacterial whole-cell fatty acid methyl esters, including hydroxy acids. *J. Clinical Microbiology* 16:584-586.

25 Morelli, G., Nagy, F., Fraley, R.T., Rogers, S.G., and Chua, N.H. (1985). A short conserved sequence is involved in the light-inducibility of a gene encoding ribulose 1,5-bisphosphate carboxylase small subunit of pea. *Nature* 315, 200-204.

30

-52-

Neuhaus, G. et al. (1987) Theor. Appl. Genet. 75:30.

5 Odell, J.T., Nagy, F., and Chua, N.H. (1985). Identification of DNA sequences required for activity of the cauliflower mosaic virus 35S promoter. *Nature* 313, 810-812.

10 Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual* - second edition. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.

Stalker, D.M., Thomas, C.M., and Helinski, D.R. (1981). Nucleotide sequence of the region of the origin of replication of the broad host range plasmid RK2. *Mol Gen Genet* 181, 8-12.

15 Tabor, S., and Richardson, C.C. (1985). A bacteriophage T7 RNA polymerase/promoter system for controlled expression of specific genes. *Proc. Natl. Acad. Sci. USA* 82, 1074-1078.

20 Tabor, C. and Tabor H. 1987. The *speEspeD* operon of *E. coli*. *J. Biol. Chem.* 262:16037-16040.

Talmadge, K., and Gilbert, W., "Construction of plasmid vectors with unique PstI cloning sites in the signal sequence coding region" *Gene*, (12) 235-241 (1980).

25 Van Der Straeten, D., Van Wiemeersch, L., Goodman, H. and Van Montagu, M. (1990) *Proc. Natl. Acad. Sci. USA* 87:4859-4863.

30 Vieira, J. and Messing, J., Production of single-stranded plasmid DNA. *Methods. Enzymol.* 153: 3 (1987).

- Ward, T., Wright, M., Roberts, J., Self, R., and Osborne, D. (1978)
Analytical procedures for the assay and identification of ethylene.
In Isolation of plant growth substances, J. Hillman, ed.
5 (Cambridge: Cambridge University Press), pp. 135-151.

-54-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Klee, Harry J.
Kishore, Ganesh M.
- (ii) TITLE OF INVENTION: Control of Fruit Ripening and Senescence
in Plants
- (iii) NUMBER OF SEQUENCES: 17
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Monsanto Co. BB4F
 - (B) STREET: 700 Chesterfield Village Parkway
 - (C) CITY: St. Louis
 - (D) STATE: Missouri
 - (E) COUNTRY: USA
 - (F) ZIP: 63198
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 07/632,440
 - (B) FILING DATE: 26-DEC-1990
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Hoerner Jr., Dennis R.
 - (B) REGISTRATION NUMBER: 30,914
 - (C) REFERENCE/DOCKET NUMBER: 38-21(10538)A
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (314)537-6099
 - (B) TELEFAX: (314)537-6047

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1079 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)

-55-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GATATCCCAT ATCAAGGAGC AGAGTC ATG AAT CTG AAT CGT TTT GAA CGT TAT	53
Met Asn Leu Asn Arg Phe Glu Arg Tyr	
1 5	
CCA TTG ACC TTC GGT CCT TCT CCC ATC ACG CCC TTG AAG CGC CTC AGT	101
Pro Leu Thr Phe Gly Pro Ser Pro Ile Thr Pro Leu Lys Arg Leu Ser	
10 15 20 25	
CAA CAT CTG GGG GGC AAG GTC GAG CTG TAT GCC AAA CGT GAA GAC TGC	149
Gln His Leu Gly Gly Lys Val Glu Leu Tyr Ala Lys Arg Glu Asp Cys	
30 35 40	
AAC AGT GGC CTG GCC TTT GGT GGG AAC AAG ACG CGC AAG CTC GAA TAC	197
Asn Ser Gly Leu Ala Phe Gly Gly Asn Lys Thr Arg Lys Leu Glu Tyr	
45 50 55	
CTC ATT CCC GAA GCG ATC GAG CAA GGC TGC GAT ACG CTG GTT TCC ATC	245
Leu Ile Pro Glu Ala Ile Glu Gln Gly Cys Asp Thr Leu Val Ser Ile	
60 65 70	
GGC GGC ATC CAG TCG AAC CAG ACC CGT CAG GTC GCT GCC GTC GCT GCC	293
Gly Gly Ile Gln Ser Asn Gln Thr Arg Gln Val Ala Ala Val Ala Ala	
75 80 85	
CAC TTG GGC ATG AAG TGC GTG TTG GTG CAG GAA AAC TGG GTG AAC TAT	341
His Leu Gly Met Lys Cys Val Leu Val Gln Glu Asn Trp Val Asn Tyr	
90 95 100 105	
TCC GAC GCG GTG TAT GAC CGC GTA GGC AAC ATC GAG ATG TCG CGG ATC	389
Ser Asp Ala Val Tyr Asp Arg Val Gly Asn Ile Glu Met Ser Arg Ile	
110 115 120	
ATG GGC GCT GAT GTG CGG CTT GAC GCC GCT GGC TTC GAT ATT GGC ATT	437
Met Gly Ala Asp Val Arg Leu Asp Ala Ala Gly Phe Asp Ile Gly Ile	
125 130 135	
CGG CCA AGT TGG GAA AAG GCC ATG AGC GAT GTC GTG GAA CAG GGT GGC	485
Arg Pro Ser Trp Glu Lys Ala Met Ser Asp Val Val Glu Gln Gly Gly	
140 145 150	
AAA CCG TTT CCG ATT CCA GCG GGT TGC TCC GAG CAT CCC TAT GGC GGC	533
Lys Pro Phe Pro Ile Pro Ala Gly Cys Ser Glu His Pro Tyr Gly Gly	
155 160 165	
CTC GGT TTC GTC GGC TTT GCC GAA GAG GTG CGG CAG CAG GAA AAG GAA	581
Leu Gly Phe Val Gly Phe Ala Glu Glu Val Arg Gln Gln Glu Lys Glu	
170 175 180 185	
CTG GGC TTC AAG TTT GAC TAC ATC GTG GTC TGC TCG GTG ACC GGC AGT	629
Leu Gly Phe Lys Phe Asp Tyr Ile Val Val Cys Ser Val Thr Gly Ser	
190 195 200	

-56-

ACG CAG GCG GGC ATG GTT GTT GGT TTC GCG GCT GAC GGT CGT TCG AAG Thr Gln Ala Gly Met Val Val Gly Phe Ala Ala Asp Gly Arg Ser Lys 205 210 215	677
AAT GTG ATT GGT ATC GAT GCT TCG GCC AAG CCG GAA CAG ACC AAG GCA Asn Val Ile Gly Ile Asp Ala Ser Ala Lys Pro Glu Gln Thr Lys Ala 220 225 230	725
CAG ATC CTG CGC ATC GCC CGA CAC ACC GCT GAG TTG GTG GAG TTG GGG Gln Ile Leu Arg Ile Ala Arg His Thr Ala Glu Leu Val Glu Leu Gly 235 240 245	773
CGC GAG ATT ACG GAA GAG GAC GTG GTG CTC GAT ACG CGT TTT GCC TAC Arg Glu Ile Thr Glu Glu Asp Val Val Leu Asp Thr Arg Phe Ala Tyr 250 255 260 265	821
CCG GAA TAT GGC TTG CCC AAC GAA GGC ACA TTG GAA GCC ATC CGA CTG Pro Glu Tyr Gly Leu Pro Asn Glu Gly Thr Leu Glu Ala Ile Arg Leu 270 275 280	869
TGC GGC AGC CTT GAA GGC GTG CTG ACA GAC CCG GTA TAT GAA GGT AAA Cys Gly Ser Leu Glu Gly Val Leu Thr Asp Pro Val Tyr Glu Gly Lys 285 290 295	917
TCG ATG CAC GGC ATG ATT GAA ATG GTC CGT CGT GGT GAA TTC CCC GAA Ser Met His Gly Met Ile Glu Met Val Arg Arg Gly Glu Phe Pro Glu 300 305 310	965
GGT TCC AAA GTG CTT TAC GCA CAC TTG GGT GGG GCG CCG GCG CTG AAC Gly Ser Lys Val Leu Tyr Ala His Leu Gly Gly Ala Pro Ala Leu Asn 315 320 325	1013
GCC TAC AGC TTC CTG TTT CGT AAC GGC TAAGCGTAGA ACTGCTTTTG Ala Tyr Ser Phe Leu Phe Arg Asn Gly 330 335	1060
GAGTCATCTG TGGGAGCTC	1079

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 31 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GAAGGAAGCT TCACGAAATC GGCCCTTATT C

31

(2) INFORMATION FOR SEQ ID NO:3:

-57-

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 32 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGGGCTTTAG ATCTTCTTTT GCACTGTGAA TG

32

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 23 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GGTGAACCAT GGAATCCAC ATG

23

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 23 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GCAATTGGAT CCCTTCCAT AGC

23

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

-58-

GGAGAAGATA AGATCTATGA AAAAAGTGA ACTGC

35

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 27 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GCAGAAGTAA ATAGATCTGG CGGAGCC

27

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1800 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CCAAACACAT AATACTTTTA ATACAATTAG TTATTTATTA GAAGTATTTA AAGTAAAGCA 60

CTTGTGAGTT GTGTACATT TATTAATCTT CATCTCTTA ATTCTCTTCA GTTTTAAATT 120

TCTTCACTTC TAAACTCATT TAGTAAAAAA AAA ATG GGA TTT GAG ATT GCA AAG 174
 Met Gly Phe Glu Ile Ala Lys
 1 5

ACC AAC TCA ATC TTA TCA AAA TTG GCT ACT AAT GAA GAG CAT GGC GAA 222
 Thr Asn Ser Ile Leu Ser Lys Leu Ala Thr Asn Glu Glu His Gly Glu
 10 15 20

AAC TCG CCA TAT TTT GAT GGG TGG AAA GCA TAC GAT AGT GAT CCT TTC 270
 Asn Ser Pro Tyr Phe Asp Gly Trp Lys Ala Tyr Asp Ser Asp Pro Phe
 25 30 35

CAC CCT CTA AAA AAC CCC AAC GGA GTT ATC CAA ATG GGT CTT GCT GAA 318
 His Pro Leu Lys Asn Pro Asn Gly Val Ile Gln Met Gly Leu Ala Glu
 40 45 50 55

AAT CAG CTT TGT TTA GAC TTG ATA GAA GAT TGG ATT AAG AGA AAC CCA 366
 Asn Gln Leu Cys Leu Asp Leu Ile Glu Asp Trp Ile Lys Arg Asn Pro
 60 65 70

-59-

AAA GGT TCA ATT TGT TCT GAA GGA ATC AAA TCA TTC AAG GCC ATT GCC Lys Gly Ser Ile Cys Ser Glu Gly Ile Lys Ser Phe Lys Ala Ile Ala 75 80 85	414
AAC TTT CAA GAT TAT CAT GGC TTG CCT GAA TTC AGA AAA GCG ATT GCG Asn Phe Gln Asp Tyr His Gly Leu Pro Glu Phe Arg Lys Ala Ile Ala 90 95 100	462
AAA TTT ATG GAG AAA ACA AGA GGA GGA AGA GTT AGA TTT GAT CCA GAA Lys Phe Met Glu Lys Thr Arg Gly Gly Arg Val Arg Phe Asp Pro Glu 105 110 115	510
AGA GTT GTT ATG GCT GGT GGT GCC ACT GGA GCT AAT GAG ACA ATT ATA Arg Val Val Met Ala Gly Gly Ala Thr Gly Ala Asn Glu Thr Ile Ile 120 125 130 135	558
TTT TGT TTG GCT GAT CCT GGC GAT GCA TTT TTA GTA CCT TCA CCA TAC Phe Cys Leu Ala Asp Pro Gly Asp Ala Phe Leu Val Pro Ser Pro Tyr 140 145 150	606
TAC CCA GCA TTT AAC AGA GAT TTA AGA TGG AGA ACT GGA GTA CAA CTT Tyr Pro Ala Phe Asn Arg Asp Leu Arg Trp Arg Thr Gly Val Gln Leu 155 160 165	654
ATT CCA ATT CAC TGT GAG AGC TCC AAT AAT TTC AAA ATT ACT TCA AAA Ile Pro Ile His Cys Glu Ser Ser Asn Asn Phe Lys Ile Thr Ser Lys 170 175 180	702
GCA GTA AAA GAA GCA TAT GAA AAT GCA CAA AAA TCA AAC ATC AAA GTA Ala Val Lys Glu Ala Tyr Glu Asn Ala Gln Lys Ser Asn Ile Lys Val 185 190 195	750
AAA GGT TTG ATT TTG ACC AAT CCA TCA AAT CCA TTG GGC ACC ACT TTG Lys Gly Leu Ile Leu Thr Asn Pro Ser Asn Pro Leu Gly Thr Thr Leu 200 205 210 215	798
GAC AAA GAC ACA CTG AAA AGT GTC TTG AGT TTC ACC AAC CAA CAC AAC Asp Lys Asp Thr Leu Lys Ser Val Leu Ser Phe Thr Asn Gln His Asn 220 225 230	846
ATC CAC CTT GTT TGT GAC GAA ATC TAC GCA GCC ACT GTC TTT GAC ACG Ile His Leu Val Cys Asp Glu Ile Tyr Ala Ala Thr Val Phe Asp Thr 235 240 245	894
CCT CAA TTC GTC AGT ATA GCT GAA ATC CTC GAT GAA CAG GAA ATG ACT Pro Gln Phe Val Ser Ile Ala Glu Ile Leu Asp Glu Gln Glu Met Thr 250 255 260	942
TAC TGC AAC AAA GAT TTA GTT CAC ATC GTC TAC AGT CTT TCA AAA GAC Tyr Cys Asn Lys Asp Leu Val His Ile Val Tyr Ser Leu Ser Lys Asp 265 270 275	990
ATG GGG TTA CCA GGA TTT AGA GTC GGA ATC ATA TAT TCT TTT AAC GAC Met Gly Leu Pro Gly Phe Arg Val Gly Ile Ile Tyr Ser Phe Asn Asp 280 285 290 295	1038

-60-

GAT GTC GTT AAT TGT GCT AGA AAA ATG TCG AGT TTC GGT TTA GTA TCT Asp Val Val Asn Cys Ala Arg Lys Met Ser Ser Phe Gly Leu Val Ser 300 305 310	1086
ACA CAA ACG CAA TAT TTT TTA GCG GCA ATG CCA TCG GAC GAA AAA TTC Thr Gln Thr Gln Tyr Phe Leu Ala Ala Met Pro Ser Asp Glu Lys Phe 315 320 325	1134
GTC GAT AAT TTT CTA AGA GAA AGC GCG ATG AGG TTA GGT AAA AGG CAC Val Asp Asn Phe Leu Arg Glu Ser Ala Met Arg Leu Gly Lys Arg His 330 335 340	1182
AAA CAT TTT ACT AAT GGA CTT GAA GTA GTG GGA ATT AAA TGC TTG AAA Lys His Phe Thr Asn Gly Leu Glu Val Val Gly Ile Lys Cys Leu Lys 345 350 355	1230
AAT AAT GCG GGG CTT TTT TGT TGG ATG GAT TTG CGT CCA CTT TTA AGG Asn Asn Ala Gly Leu Phe Cys Trp Met Asp Leu Arg Pro Leu Leu Arg 360 365 370 375	1278
GAA TCG ACT TTC GAT AGC GAA ATG TCG TTA TGG AGA GTT ATT ATA AAC Glu Ser Thr Phe Asp Ser Glu Met Ser Leu Trp Arg Val Ile Ile Asn 380 385 390	1326
GAT GTT AAG CTT AAC GTC TCG CTT GGA TCT TCG TTT GAA TGT CAA GAG Asp Val Lys Leu Asn Val Ser Leu Gly Ser Ser Phe Glu Cys Gln Glu 395 400 405	1374
CCA GGG TGG TTC CGA GTT TGT TTT GCA AAT ATG GAT GAT GGA ACG GTT Pro Gly Trp Phe Arg Val Cys Phe Ala Asn Met Asp Asp Gly Thr Val 410 415 420	1422
GAT ATT GCG CTC GCG AGG ATT CGG AGG TTC GTA GGT GTT GAG AAA AGT Asp Ile Ala Leu Ala Arg Ile Arg Arg Phe Val Gly Val Glu Lys Ser 425 430 435	1470
GGA GAT AAA TCG AGT TCG ATG GAA AAG AAG CAA CAA TGG AAG AAG AAT Gly Asp Lys Ser Ser Ser Met Glu Lys Lys Gln Gln Trp Lys Lys Asn 440 445 450 455	1518
AAT TTG AGA CTT AGT TTT TCG AAA AGA ATG TAT GAT GAA AGT GTT TTG Asn Leu Arg Leu Ser Phe Ser Lys Arg Met Tyr Asp Glu Ser Val Leu 460 465 470	1566
TCA CCA CTT TCG TCA CCT ATT CCT CCC TCA CCA TTA GTT CGT Ser Pro Leu Ser Ser Pro Ile Pro Pro Ser Pro Leu Val Arg 475 480 485	1608
TAAGACTTAA TTAAAAGGGA AGAATTTAAT TTATGTTTTT TTATATTTTG AAAAAAATTT	1668
GTAAGAATAA GATTATAATA GGAAAAGAAA ATAAGTATGT AGGATGAGGA GTATTTTCAG	1728
AAATAGTTGT TAGCGTATGT ATTGACAACCT GGTCTATGTA CTTAGACATC ATAATTTGTC	1788
TTAGCTAATT AA	1800

-61-

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 900 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ACAGCCGTCC TAAGGAGAAG ATAAGATCT ATG AAA AAA CTG AAA CTG CAT GGC	53
Met Lys Lys Leu Lys Leu His Gly	
1 5	
TTT AAT AAT CTG ACC AAA AGT CTG AGT TTT TGT ATT TAC GAT ATC TGC	101
Phe Asn Asn Leu Thr Lys Ser Leu Ser Phe Cys Ile Tyr Asp Ile Cys	
10 15 20	
TAC GCC AAA ACT GCC GAA GAG CGC GAC GGT TAT ATT GCT TAT ATC GAT	149
Tyr Ala Lys Thr Ala Glu Glu Arg Asp Gly Tyr Ile Ala Tyr Ile Asp	
25 30 35 40	
GAA CTC TAT AAT GCC AAC CGT CTG ACC GAA ATC CTG TCA GAA ACC TGT	197
Glu Leu Tyr Asn Ala Asn Arg Leu Thr Glu Ile Leu Ser Glu Thr Cys	
45 50 55	
TCC ATT ATC GGG GCT AAT ATT CTT AAC ATC GCC CGC CAG GAT TAC GAA	245
Ser Ile Ile Gly Ala Asn Ile Leu Asn Ile Ala Arg Gln Asp Tyr Glu	
60 65 70	
CCA CAG GGT GCC AGC GTC ACT ATT CTG GTG AGT GAA GAA CCG GTT GAC	293
Pro Gln Gly Ala Ser Val Thr Ile Leu Val Ser Glu Glu Pro Val Asp	
75 80 85	
CCG AAA CTC ATC GAC AAA ACA GAA CAC CCC GGC CCA CTG CCA GAA ACG	341
Pro Lys Leu Ile Asp Lys Thr Glu His Pro Gly Pro Leu Pro Glu Thr	
90 95 100	
GTC GTT GCC CAT CTT GAT AAA AGT CAT ATT TGC GTA CAT ACC TAC CCG	389
Val Val Ala His Leu Asp Lys Ser His Ile Cys Val His Thr Tyr Pro	
105 110 115 120	
GAA AGT CAT CCT GAA GGC GGT TTA TGT ACC TTC CGC GCC GAT ATT GAA	437
Glu Ser His Pro Glu Gly Gly Leu Cys Thr Phe Arg Ala Asp Ile Glu	
125 130 135	
GTC TCT ACC TGC GGC GTG ATT TCT CCG CTG AAG GCG CTG AAT TAC CTG	485
Val Ser Thr Cys Gly Val Ile Ser Pro Leu Lys Ala Leu Asn Tyr Leu	
140 145 150	
ATC CAC CAG CTT GAG TCC GAT ATC GTA ACC ATT GAT TAT CGC GTG CGC	533
Ile His Gln Leu Glu Ser Asp Ile Val Thr Ile Asp Arg Val Arg	
155 160 165	

-62-

GGT TTT ACC CGC GAC ATT AAC GGT ATG AAG CAC TTT ATC GAC CAT GAG Gly Phe Thr Arg Asp Ile Asn Gly Met Lys His Phe Ile Asp His Glu 170 175 180	581
ATT AAT TCG ATT CAG AAC TTT ATG TCT GAC GAT ATG AAG GCG CTG TAT Ile Asn Ser Ile Gln Asn Phe Met Ser Asp Asp Met Lys Ala Leu Tyr 185 190 195 200	629
GAC ATG GTG GAT GTG AAC GTC TAT CAG GAA AAT ATC TTC CAT ACC AAG Asp Met Val Asp Val Asn Val Tyr Gln Glu Asn Ile Phe His Thr Lys 205 210 215	677
ATG TTG CTT AAA GAG TTC GAC CTT AAG CAC TAC ATG TTC CAC ACC AAA Met Leu Leu Lys Glu Phe Asp Leu Lys His Tyr Met Phe His Thr Lys 220 225 230	725
CCG GAA GAC TTA ACC GAC AGC GAG CGC CAG GAA ATT ACC GCT GCG CTG Pro Glu Asp Leu Thr Asp Ser Glu Arg Gln Glu Ile Thr Ala Ala Leu 235 240 245	773
TGG AAA GAA ATG CGC GAG ATT TAT TAC GGG CGC AAT ATG CCA GCT GTT Trp Lys Glu Met Arg Glu Ile Tyr Tyr Gly Arg Asn Met Pro Ala Val 250 255 260	821
TAACGGCTCT GGCGGAGCTC CCAGGCTCCG CCAGATCTAT TTA CTCTCTGC TGCACGAAAT	881
TGCGGTAAGC CGCCACGAC	900

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1138 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CTAGAAGGAA GCTTCACGAA ATCGGCCCTT ATTCAAAAAT AACTTTTAAA TAATGAATTT	60
TAAATTTTAA GAAATAATAT CCAATGAATA AATGACATGT AGCATTTTAC CTAAATATTT	120
CAACTATTTT AATCCAATAT TAATTGTGTT TATTCCTAAC AATAGAAAGT CTTGTGCAGA	180
CATTTAATCT GACTTTTCCA GTACTAAATA TTAATTTTCT GAAGATTTTC GGGTTTAGTC	240
CACAAGTTTT AGTGAGAAGT TTTGCTCAAA ATTTTAGGTG AGAAGGTTTG ATATTTATCT	300
TTTGTTAAAT TAATTTATCT AGGTGACTAT TATTTATTTA AGTAGAAATT CATATCATTA	360
CTTTTGCCAA CTTGTAGTCA TAATAGGAGT AGGTGTATAT GATGAAGGAA TAAACAAGTT	420

-63-

CAGTGAAGTG ATTAAATATA AATATAATTT AGGTGTACAT CAAATAAAAA CCTTAAAGTT	480
TAGAAAGGCA CCGAATAATT TTGCATAGAA GATATTAGTA AATTTATAAA AATAAAAGAA	540
ATGTAGTTGT CAAGTTGTCT TCTTTTTTTT GGATAAAAAAT AGCAGTTGGC TTATGTCATT	600
CTTTTACAAC CTCCATGCCA CTTGTCCAAT TGTGACACT TAACTAATTA GTTTGATTCA	660
TGTATGAATA CTAAATAATT TTTTAGGACT GACTCAAATA TTTTATATT ATCATAGTAA	720
TATTTATCTA ATTTTATAGGA CCACTTATTA CTAAATAATA AATTAACACT TACTATATTA	780
TTGTTGTGAA ACAACAACGT TTTGGTTGTT ATGATGAAAC GTACACTATA TCAGTATGAA	840
AAATTCAAAA CGATTAGTAT AAATTATATT GAAAATTTGA TATTTTCTA TTCTTAATCA	900
GACGTATTGG GTTTCATATT TAAAAAGGG ACTAACTTA GAAGAGAAGT TTGTTTGAAA	960
CTACTTTTGT CTCTTCTTG TTCCCATTTT TCTCTTAGAT TTCAAAAAGT GAACTACTTT	1020
ATCTCTTCT TTGTTACAT TTTATTTTAT TCTATTATAA ATATGGCATC CTCATATTGA	1080
GATTTTCTA AATTATTCTA ATCATTACCA GTGCAAAAGA AGATCTAAAG CCCTAGAG	1138

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CCCGGATCCA TGAATCTGAA TCGTTTT

27

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CCCGGATCCG CCGTTACGAA ACAGGAA

27

(2) INFORMATION FOR SEQ ID NO:13:

-64-

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 318 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

```

AGATCTATCG ATAAGCTTGA TGTAATTGGA GGAAGATCAA AATTTTCAAT CCCCATTTCTT      60
CGATTGCTTC AATTGAAGTT TCTCCG ATG GCG CAA GTT AGC AGA ATC TGC AAT      113
                               Met Ala Gln Val Ser Arg Ile Cys Asn
                               1               5

GGT GTG CAG AAC CCA TCT CTT ATC TCC AAT CTC TCG AAA TCC AGT CAA      161
Gly Val Gln Asn Pro Ser Leu Ile Ser Asn Leu Ser Lys Ser Ser Gln
10               15               20               25

CGC AAA TCT CCC TTA TCG GTT TCT CTG AAG ACG CAG CAG CAT CCA CGA      209
Arg Lys Ser Pro Leu Ser Val Ser Leu Lys Thr Gln Gln His Pro Arg
30               35               40

GCT TAT CCG ATT TCG TCG TCG TGG GGA TTG AAG AAG AGT GGG ATG ACG      257
Ala Tyr Pro Ile Ser Ser Ser Trp Gly Leu Lys Lys Ser Gly Met Thr
45               50               55

TTA ATT GGC TCT GAG CTT CGT CCT CTT AAG GTC ATG TCT TCT GTT TCC      305
Leu Ile Gly Ser Glu Leu Arg Pro Leu Lys Val Met Ser Ser Val Ser
60               65               70

ACG GCG TGC ATG C      318
Thr Ala Cys Met
75

```

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1377 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

```

GC ATG CTT CAC GGT GCA AGC AGC CGT CCA GCA ACT GCT CGT AAG TCC      47
Met Leu His Gly Ala Ser Ser Arg Pro Ala Thr Ala Arg Lys Ser
1               5               10               15

```

-65-

TCT GGT CTT TCT GGA ACC GTC CGT ATT CCA GGT GAC AAG TCT ATC TCC Ser Gly Leu Ser Gly Thr Val Arg Ile Pro Gly Asp Lys Ser Ile Ser 20 25 30	95
CAC AGG TCC TTC ATG TTT GGA GGT CTC GCT AGC GGT GAA ACT CGT ATC His Arg Ser Phe Met Phe Gly Gly Leu Ala Ser Gly Glu Thr Arg Ile 35 40 45	143
ACC GGT CTT TTG GAA GGT GAA GAT GTT ATC AAC ACT GGT AAG GCT ATG Thr Gly Leu Leu Glu Gly Glu Asp Val Ile Asn Thr Gly Lys Ala Met 50 55 60	191
CAA GCT ATG GGT GCC AGA ATC CGT AAG GAA GGT GAT ACT TGG ATC ATT Gln Ala Met Gly Ala Arg Ile Arg Lys Glu Gly Asp Thr Trp Ile Ile 65 70 75	239
GAT GGT GTT GGT AAC GGT GGA CTC CTT GCT CCT GAG GCT CCT CTC GAT Asp Gly Val Gly Asn Gly Gly Leu Leu Ala Pro Glu Ala Pro Leu Asp 80 85 90 95	287
TTC GGT AAC GCT GCA ACT GGT TGC CGT TTG ACT ATG GGT CTT GTT GGT Phe Gly Asn Ala Ala Thr Gly Cys Arg Leu Thr Met Gly Leu Val Gly 100 105 110	335
GTT TAC GAT TTC GAT AGC ACT TTC ATT GGT GAC GCT TCT CTC ACT AAG Val Tyr Asp Phe Asp Ser Thr Phe Ile Gly Asp Ala Ser Leu Thr Lys 115 120 125	383
CGT CCA ATG GGT CGT GTG TTG AAC CCA CTT CGC GAA ATG GGT GTG CAG Arg Pro Met Gly Arg Val Leu Asn Pro Leu Arg Glu Met Gly Val Gln 130 135 140	431
GTG AAG TCT GAA GAC GGT GAT CGT CTT CCA GTT ACC TTG CGT GGA CCA Val Lys Ser Glu Asp Gly Asp Arg Leu Pro Val Thr Leu Arg Gly Pro 145 150 155	479
AAG ACT CCA ACG CCA ATC ACC TAC AGG GTA CCT ATG GCT TCC GCT CAA Lys Thr Pro Thr Pro Ile Thr Tyr Arg Val Pro Met Ala Ser Ala Gln 160 165 170 175	527
GTG AAG TCC GCT GTT CTG CTT GCT GGT CTC AAC ACC CCA GGT ATC ACC Val Lys Ser Ala Val Leu Leu Ala Gly Leu Asn Thr Pro Gly Ile Thr 180 185 190	575
ACT GTT ATC GAG CCA ATC ATG ACT CGT GAC CAC ACT GAA AAG ATG CTT Thr Val Ile Glu Pro Ile Met Thr Arg Asp His Thr Glu Lys Met Leu 195 200 205	623
CAA GGT TTT GGT GCT AAC CTT ACC GTT GAG ACT GAT GCT GAC GGT GTG Gln Gly Phe Gly Ala Asn Leu Thr Val Glu Thr Asp Ala Asp Gly Val 210 215 220	671
CGT ACC ATC CGT CTT GAA GGT CGT GGT AAG CTC ACC GGT CAA GTG ATT Arg Thr Ile Arg Leu Glu Gly Arg Gly Lys Leu Thr Gly Gln Val Ile 225 230 235	719

-66-

GAT GTT CCA GGT GAT CCA TCC TCT ACT GCT TTC CCA TTG GTT GCT GCC Asp Val Pro Gly Asp Pro Ser Ser Thr Ala Phe Pro Leu Val Ala Ala 240 245 250 255	767
TTG CTT GTT CCA GGT TCC GAC GTC ACC ATC CTT AAC GTT TTG ATG AAC Leu Leu Val Pro Gly Ser Asp Val Thr Ile Leu Asn Val Leu Met Asn 260 265 270	815
CCA ACC CGT ACT GGT CTC ATC TTG ACT CTG CAG GAA ATG GGT GCC GAC Pro Thr Arg Thr Gly Leu Ile Leu Thr Leu Gln Glu Met Gly Ala Asp 275 280 285	863
ATC GAA GTG ATC AAC CCA CGT CTT GCT GGT GGA GAA GAC GTG GCT GAC Ile Glu Val Ile Asn Pro Arg Leu Ala Gly Gly Glu Asp Val Ala Asp 290 295 300	911
TTG CGT GTT CGT TCT TCT ACT TTG AAG GGT GTT ACT GTT CCA GAA GAC Leu Arg Val Arg Ser Ser Thr Leu Lys Gly Val Thr Val Pro Glu Asp 305 310 315	959
CGT GCT CCT TCT ATG ATC GAC GAG TAT CCA ATT CTC GCT GTT GCA GCT Arg Ala Pro Ser Met Ile Asp Glu Tyr Pro Ile Leu Ala Val Ala Ala 320 325 330 335	1007
GCA TTC GCT GAA GGT GCT ACC GTT ATG AAC GGT TTG GAA GAA CTC CGT Ala Phe Ala Glu Gly Ala Thr Val Met Asn Gly Leu Glu Glu Leu Arg 340 345 350	1055
GTT AAG GAA AGC GAC CGT CTT TCT GCT GTC GCA AAC GGT CTC AAG CTC Val Lys Glu Ser Asp Arg Leu Ser Ala Val Ala Asn Gly Leu Lys Leu 355 360 365	1103
AAC GGT GTT GAT TGC GAT GAA GGT GAG ACT TCT CTC GTC GTG CGT GGT Asn Gly Val Asp Cys Asp Glu Gly Glu Thr Ser Leu Val Val Arg Gly 370 375 380	1151
CGT CCT GAC GGT AAG GGT CTC GGT AAC GCT TCT GGA GCA GCT GTC GCT Arg Pro Asp Gly Lys Gly Leu Gly Asn Ala Ser Gly Ala Ala Val Ala 385 390 395	1199
ACC CAC CTC GAT CAC CGT ATC GCT ATG AGC TTC CTC GTT ATG GGT CTC Thr His Leu Asp His Arg Ile Ala Met Ser Phe Leu Val Met Gly Leu 400 405 410 415	1247
GTT TCT GAA AAC CCT GTT ACT GTT GAT GAT GCT ACT ATG ATC GCT ACT Val Ser Glu Asn Pro Val Thr Val Asp Asp Ala Thr Met Ile Ala Thr 420 425 430	1295
AGC TTC CCA GAG TTC ATG GAT TTG ATG GCT GGT CTT GGA GCT AAG ATC Ser Phe Pro Glu Phe Met Asp Leu Met Ala Gly Leu Gly Ala Lys Ile 435 440 445	1343
GAA CTC TCC GAC ACT AAG GCT GCT TGATGAGCTC Glu Leu Ser Asp Thr Lys Ala Ala 450 455	1377

-67-

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1029 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 7..1020

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GGATCC ATG AAT TTG AAT CGT TTT AAA CGT TAT CCG TTG ACC TTC GGT	48
Met Asn Leu Asn Arg Phe Lys Arg Tyr Pro Leu Thr Phe Gly	
1 5 10	
CCT TCT CCC ATC ACG CCC TTG AAG CGC CTC AGT GAA CAC TTG GGT GGC	96
Pro Ser Pro Ile Thr Pro Leu Lys Arg Leu Ser Glu His Leu Gly Gly	
15 20 25 30	
AAG GTC GAG CTG TAT GCC AAG CGT GAA GAC TGC AAC AGT GGC CTG GCC	144
Lys Val Glu Leu Tyr Ala Lys Arg Glu Asp Cys Asn Ser Gly Leu Ala	
35 40 45	
TTC GGC GGG AAC AAA ACG CGC AAG CTC GAA TAT TTG ATT CCC GAA GCG	192
Phe Gly Gly Asn Lys Thr Arg Lys Leu Glu Tyr Leu Ile Pro Glu Ala	
50 55 60	
CTC GAG CAA GGC TGC GAT ACC TTG GTT TCC ATC GGC GGC ATC CAG TCG	240
Leu Glu Gln Gly Cys Asp Thr Leu Val Ser Ile Gly Gly Ile Gln Ser	
65 70 75	
AAC CAG ACC CGC CAG GTG GCC GCC GTT GCC GCT CAC CTG GGC ATG AAG	288
Asn Gln Thr Arg Gln Val Ala Ala Val Ala Ala His Leu Gly Met Lys	
80 85 90	
TCG GTG CTG GTC GAG GAA AAC TGG GTG AAC TAC TCC GAT GCG GTG TAT	336
Ser Val Leu Val Glu Glu Asn Trp Val Asn Tyr Ser Asp Ala Val Tyr	
95 100 105 110	
GAC CGC GTT GGC AAT ATC GAA ATG TCT CGC ATC ATG GGC GCC GAG GTA	384
Asp Arg Val Gly Asn Ile Glu Met Ser Arg Ile Met Gly Ala Glu Val	
115 120 125	
CGA CTG GAC GCC GCC GGG TTC GAT ATC GGC ATT CGG CCC AGC TGG GAG	432
Arg Leu Asp Ala Ala Gly Phe Asp Ile Gly Ile Arg Pro Ser Trp Glu	
130 135 140	

-68-

AAG GCC ATG GAC GAT GTG GTG GCG CGG GGT GGC AAG CCG TTC CCG ATA Lys Ala Met Asp Asp Val Val Ala Arg Gly Gly Lys Pro Phe Pro Ile 145 150 155	480
CCG GCG GGT TGT TCC GAA CAC CCC TAC GGC GGC CTT GGG TTC GTC GGC Pro Ala Gly Cys Ser Glu His Pro Tyr Gly Gly Leu Gly Phe Val Gly 160 165 170	528
TTT GCC GAG GAA GTG CGA GAG CAG GAA AAA CAA CTG GGG TTC ACG TTC Phe Ala Glu Glu Val Arg Glu Gln Glu Lys Gln Leu Gly Phe Thr Phe 175 180 185 190	576
GAC TAC ATC GTG GTC TGC TCT GTG ACC GGC AGT ACC CAG GCC GGC ATG Asp Tyr Ile Val Val Cys Ser Val Thr Gly Ser Thr Gln Ala Gly Met 195 200 205	624
GTC GTC GGT TTC GCC GCG GAC GGC CGT TCG AAG AAC GTT ATC GGC ATT Val Val Gly Phe Ala Ala Asp Gly Arg Ser Lys Asn Val Ile Gly Ile 210 215 220	672
GAT GCC TCG GCC AAG CCG GAG CAA ACC AAG GCA CAG ATC CTG CGT ATC Asp Ala Ser Ala Lys Pro Glu Gln Thr Lys Ala Gln Ile Leu Arg Ile 225 230 235	720
GCC CGG CAC ACC GCA GAG TTG GTG GAA CTG GGC CGT GAG ATC ACC GAA Ala Arg His Thr Ala Glu Leu Val Glu Leu Gly Arg Glu Ile Thr Glu 240 245 250	768
GAC GAC GTG GTG CTC GAT ACA CGT TTT GCC TAC CCG GAA TAC GGT TTG Asp Asp Val Val Leu Asp Thr Arg Phe Ala Tyr Pro Glu Tyr Gly Leu 255 260 265 270	816
CCC AAC GAA GGC ACG CTG GAA GCC ATT CGT TTG TGC GGG AGC CTG GAA Pro Asn Glu Gly Thr Leu Glu Ala Ile Arg Leu Cys Gly Ser Leu Glu 275 280 285	864
GGT GTG CTG ACC GAT CCG GTG TAC GAG GGC AAA TCC ATG CAC GGG ATG Gly Val Leu Thr Asp Pro Val Tyr Glu Gly Lys Ser Met His Gly Met 290 295 300	912
ATT GAA ATG GTC CGC CGT GGC GAG TTC CCC GAA GGC TCC AAA GTG CTG Ile Glu Met Val Arg Arg Gly Glu Phe Pro Glu Gly Ser Lys Val Leu 305 310 315	960
TAT GCG CAC TTG GGT GGG GCG CCT GCG CTG AAT GCC TAC AGC TTC CTG Tyr Ala His Leu Gly Gly Ala Pro Ala Leu Asn Ala Tyr Ser Phe Leu 320 325 330	1008
TTT CGT AAC GGC GGATCCGGG Phe Arg Asn Gly 335	1029

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

-69-

(A) LENGTH: 338 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

```

Met Asn Leu Asn Arg Phe Lys Arg Tyr Pro Leu Thr Phe Gly Pro Ser
 1             5             10             15

Pro Ile Thr Pro Leu Lys Arg Leu Ser Glu His Leu Gly Gly Lys Val
      20             25             30

Glu Leu Tyr Ala Lys Arg Glu Asp Cys Asn Ser Gly Leu Ala Phe Gly
      35             40             45

Gly Asn Lys Thr Arg Lys Leu Glu Tyr Leu Ile Pro Glu Ala Leu Glu
      50             55             60

Gln Gly Cys Asp Thr Leu Val Ser Ile Gly Gly Ile Gln Ser Asn Gln
      65             70             75             80

Thr Arg Gln Val Ala Ala Val Ala Ala His Leu Gly Met Lys Ser Val
      85             90             95

Leu Val Glu Glu Asn Trp Val Asn Tyr Ser Asp Ala Val Tyr Asp Arg
      100            105            110

Val Gly Asn Ile Glu Met Ser Arg Ile Met Gly Ala Glu Val Arg Leu
      115            120            125

Asp Ala Ala Gly Phe Asp Ile Gly Ile Arg Pro Ser Trp Glu Lys Ala
      130            135            140

Met Asp Asp Val Val Ala Arg Gly Gly Lys Pro Phe Pro Ile Pro Ala
      145            150            155            160

Gly Cys Ser Glu His Pro Tyr Gly Gly Leu Gly Phe Val Gly Phe Ala
      165            170            175

Glu Glu Val Arg Glu Gln Glu Lys Gln Leu Gly Phe Thr Phe Asp Tyr
      180            185            190

Ile Val Val Cys Ser Val Thr Gly Ser Thr Gln Ala Gly Met Val Val
      195            200            205

Gly Phe Ala Ala Asp Gly Arg Ser Lys Asn Val Ile Gly Ile Asp Ala
      210            215            220

Ser Ala Lys Pro Glu Gln Thr Lys Ala Gln Ile Leu Arg Ile Ala Arg
      225            230            235            240

His Thr Ala Glu Leu Val Glu Leu Gly Arg Glu Ile Thr Glu Asp Asp
      245            250            255

```

-70-

Val Val Leu Asp Thr Arg Phe Ala Tyr Pro Glu Tyr Gly Leu Pro Asn
 260 265 270

Glu Gly Thr Leu Glu Ala Ile Arg Leu Cys Gly Ser Leu Glu Gly Val
 275 280 285

Leu Thr Asp Pro Val Tyr Glu Gly Lys Ser Met His Gly Met Ile Glu
 290 295 300

Met Val Arg Arg Gly Glu Phe Pro Glu Gly Ser Lys Val Leu Tyr Ala
 305 310 315 320

His Leu Gly Gly Ala Pro Ala Leu Asn Ala Tyr Ser Phe Leu Phe Arg
 325 330 335

Asn Gly

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 597 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

TCATCAAAAT ATTTAGCAGC ATTCCAGATT GGGTCAATC AACAGGTAC GAGCCATATC	60
ACTTTATTCA AATTGGTATC GCCAAAACCA AGAAGGAAC CCCATCCTCA AAGGTTTGTA	120
AGGAAGAATT CTCAGTCCAA AGCCTCAACA AGGTCAGGGT ACAGAGTCTC CAAACCATTA	180
GCCAAAAGCT ACAGGAGATC AATGAAGAAT CTTCAATCAA AGTAACTAC TGTTCCAGCA	240
CATGCATCAT GGTCAAGTAAG TTTCAGAAAA AGACATCCAC CGAAGACTTA AAGTTAGTGG	300
GCATCTTTGA AAGTAATCTT GTCAACATCG AGCAGCTGGC TTGTGGGGAC CAGACAAAAA	360
AGGAATGGTG CAGAATTGTT AGGCGCACCT ACCAAAAGCA TCTTGCCTT TATTGCAAAG	420
ATAAAGCAGA TTCCTTAGT ACAAGTGGGG AACAAAATAA CGTGGAAG AGCTGTCCTG	480
ACAGCCCACT CACTAATGCG TATGACGAAC GCAGTGACGA CCACAAAAGA ATTCCCTCTA	540
TATAAGAAGG CATTCAATCC CATTTGAAGG ATCATCAGAT ACTAACCAAT ATTTCTC	597

-71-

CLAIMS

1. A recombinant, double stranded DNA molecule comprising in sequence:

5 a promoter that functions in plant cells to cause the production of an RNA sequence;

a structural DNA sequence that causes the production of an RNA sequence that encodes a 1-aminocyclopropane-1-carboxylic acid deaminase enzyme; and

10 a 3' non-translated region that functions in plant cells to cause the addition of polyadenylated nucleotides to the 3' end of the RNA sequence, said promoter being heterologous with respect to said structural DNA sequence.

15 2. The DNA molecule of claim 1 wherein said promoter is a plant DNA virus promoter.

3. The DNA molecule of claim 2 wherein said promoter is selected from the group consisting of the CaMV35S promoter and the FMV35S promoter.

4. The DNA molecule of claim 1 wherein said promoter is a fruit specific promoter.

25 5. The DNA molecule of claim 1 wherein said promoter is the E8 promoter from tomato.

6. The DNA molecule of claim 1 wherein said structural DNA sequence is SEQ ID NO:1.

30

-72-

7. The DNA molecule of claim 1 wherein said structural DNA sequence is from a microorganism capable of sustaining growth in media containing ACC as the sole nitrogen source.

5 8. A method for controlling the ripening of the fruit of a fruit-bearing plant, said method comprising the steps of:

 obtaining cells of said fruit-bearing plant;
 transforming said cells of said fruit-bearing plant with a
10 chimeric gene comprising a promoter that functions in plant cells to cause the production of an RNA sequence, a structural DNA sequence that causes the production of an RNA sequence that encodes a 1-aminocyclopropane-1-carboxylic acid deaminase enzyme, and a 3' non-translated region that functions in plant
15 cells to cause the addition of polyadenylated nucleotides to the 3' end of the RNA sequence, said promoter being heterologous with respect to said structural DNA sequence, and wherein said chimeric gene becomes integrated into the genome of said plant cell; regenerating a plant from said transformed plant cell and
20 growing said transformed fruit-bearing plant to produce fruit.

 9. The DNA molecule of claim 8 wherein said promoter is a plant DNA virus promoter.

25 10. The DNA molecule of claim 9 wherein said promoter is selected from the group consisting of the CaMV35S promoter and the FMV35S promoter.

 11. The DNA molecule of claim 8 wherein said promoter
30 is a fruit specific promoter.

-73-

12. The DNA molecule of claim 8 wherein said promoter is the E8 promoter from tomato.

5 13. The DNA molecule of claim 8 wherein said structural DNA sequence is SEQ ID NO:1.

10 14. The DNA molecule of claim 8 wherein said structural DNA sequence is from a microorganism capable of sustaining growth in media containing ACC as the sole nitrogen source.

15 15. A method for delaying the ripening of fruit of a plant comprising expressing 1-aminocyclopropane-1-carboxylic acid deaminase in said plant at a level sufficient to reduce the production of ethylene in said fruit.

20 16. A method for extending the shelf life of a fruit from a fruit-bearing plant, said method comprising the steps of:

obtaining cells of said fruit-bearing plant;
transforming said cells with a chimeric gene comprising a promoter that functions in plant cells to cause the production of an RNA sequence, a structural DNA sequence that
25 causes the production of an RNA sequence that encodes a 1-aminocyclopropane-1-carboxylic acid deaminase enzyme and a 3' non-translated region that functions in plant cells to cause the addition of polyadenylated nucleotides to the 3' end of the RNA sequence, said promoter being heterologous with respect to said

-74-

structural DNA sequence and wherein said chimeric gene becomes integrated into the genome of said plant;

regenerating a plant from said transformed plant cell;

growing said transformed plant until fruit begins to be

5 produced; and

removing said fruit from said plant when said fruit reaches the breaker stage of development.

10 17. The method of claim 16 further comprising the step of:

exposing said fruit to ethylene to achieve the desired ripeness of said fruit.

18. A transformed plant comprising:

15 a promoter that functions in plant cells to cause the production of an RNA sequence, a structural DNA sequence that causes the production of an RNA sequence that encodes a 1-aminocyclopropane-1-carboxylic acid deaminase enzyme, and a 3' non-translated region that functions in plant cells to cause the
20 addition of polyadenylated nucleotides to the 3' end of the RNA sequence, said promoter being heterologous with respect to said structural DNA sequence.

25 19. The transformed plant of claim 18 wherein said plant is selected from the group consisting of tomato, banana, kiwifruit, avocado, melon, strawberry, mango, papaya, apple, peach, cabbage, cauliflower, lettuce, onions, broccoli, cotton, canola, oilseed rape, carnations, and roses.

-75-

20. The transformed plant of claim 19 wherein said plant is tomato.

5 21. The transformed plant of claim 18 wherein said structural DNA sequence is SEQ ID NO:1.

22. A tomato fruit comprising:
a promoter that functions in plant cells to
cause the production of an RNA sequence;
10 a structural DNA sequence that causes the
production of an RNA sequence that encodes a 1-
aminocyclopropane-1-carboxylic acid deaminase
enzyme; and
a 3' non-translated region that functions in
15 plant cells to cause the addition of polyadenylated nucleotides to
the 3' end of the RNA sequence,
said promoter being heterologous with respect to said
structural DNA sequence.

20 23. The tomato fruit of claim 22 wherein said promoter
is the full-length transcript (35S) promoter from figwort mosaic
virus.

24. The tomato fruit of claim 23 wherein said structural
25 DNA sequence is SEQ ID NO:1.

25. A method for reducing the level of ethylene produced
in a plant which comprises expressing a 1-aminocyclopropane-1-
carboxylic acid metabolizing enzyme in said plant at a sufficient

-76-

level to reduce the steady- state 1-aminocyclopropane-1-carboxylic acid concentration by at least about 70%.

26. The method of claim 25 wherein said 1-aminocyclopropane-1-carboxylic acid metabolizing enzyme is 1-aminocyclopropane-1-carboxylic acid deaminase.

27. A method of claim 25 in which the steady-state 1-aminocyclopropane-1-carboxylic acid concentration is reduced by at least about 90%.

28. A method for reducing the level of ethylene produced in a plant which comprises expressing a 1-aminocyclopropane-1-carboxylic acid metabolizing enzyme in said plant at a sufficient level to reduce the steady- state ethylene concentration by at least about 70%.

29. The method of claim 28 wherein said 1-aminocyclopropane-1-carboxylic acid metabolizing enzyme is 1-aminocyclopropane-1-carboxylic acid deaminase.

30. A method of claim 28 in which the steady-state ethylene concentration is reduced by at least about 90%.

1/30

Strain	Number of Isolates Tested
<i>Pseudomonas putida</i> biovar A	58
<i>Pseudomonas putida</i> biovar B	23
<i>Pseudomonas chlororaphis</i>	170
<i>Pseudomonas tolaasii</i>	41
<i>Pseudomonas aureofaciens</i>	28
<i>Pseudomonas corrugata</i>	13
<i>Pseudomonas fragi</i>	18
<i>Pseudomonas marginalis</i>	1
<i>Pseudomonas syringae</i> (multiple pathovars)	93
<i>Pseudomonas fluorescens</i> A	4
<i>Pseudomonas fluorescens</i> B	5
<i>Pseudomonas fluorescens</i> C (Inc.ATCC 10844)	12
<i>Pseudomonas fluorescens</i> G (Inc.ATCC 13524)	14
<i>Pseudomonas coronafaciens</i>	3
<i>Pseudomonas aeruginosa</i> (Inc.ATCC 15526)	9
fluorescent pseudomonads (incomplete identification)	61
<i>Pseudomonas mendocina</i>	1
<i>Pseudomonas stutzeri</i>	1
<i>Pseudomonas alcaligenes</i>	1
<i>Pseudomonas testosteroni</i> (Inc.ATCC 17409, 17510, 11996)	7
<i>Pseudomonas cepacia</i> ATCC 10856	1
<i>Pseudomonas delafieldii</i> ATCC 17505	1
<i>Pseudomonas diminuta</i> ATCC 11568	1
<i>Pseudomonas acidovorans</i>	3
<i>Pseudomonas cruciariae</i> ATCC 13262	1
<i>Pseudomonas methanolica</i> ATCC 21704	1
<i>Pseudomonas pickettii</i> ATCC 27511	1
<i>Pseudomonas vesicularis</i> ATCC 11426	1
<i>Xanthomonas maltophilia</i> ATCC 13637	2
<i>Agrobacterium tumefaciens</i>	3
<i>Erwinia herbicola</i>	1
<i>Enterobacter cloacae</i> ATCC 13047	1
<i>Enterobacter aerogenes</i> ATCC 13048	1
<i>Hafnia alvei</i>	2
non-fluorescent (incomplete identification)	11
<i>Bacillus thuringiensis</i>	1
<i>Bacillus licheniformis</i>	1
<i>Corynebacterium fascians</i>	41

FIG. 1

SUBSTITUTE SHEET

2/30

1 GATATCCCATATCAAGGAGCAGAGTCATGAATCTGAATCGTTTGAACGTTATCCATTGACC
MetAsnLeuAsnArgPheGluArgTyrProLeuThr

63 TTCGGTCCTTCTCCCATCACGCCCTTGAAGCGCCTCAGTCAACATCTGGGGGGCAAGGTCGA
PheGlyProSerProIleThrProLeuLysArgLeuSerGlnHisLeuGlyGlyLysValGI

125 GCTGTATGCCAAACGTGAAGACTGCAACAGTGGCCTGGCCTTTGGTGGGAACAAGACGCGCA
uLeuTyrAlaLysArgGluAspCysAsnSerGlyLeuAlaPheGlyGlyAsnLysThrArgL

187 AGCTCGAATACCTCATTTCCGAAGCGATCGAGCAAGGCTGCGATACGCTGGTTTCCATCGGC
ysLeuGluTyrLeuIleProGluAlaIleGluGlnGlyCysAspThrLeuValSerIleGly

249 GGCAATCCAGTCGAACCAAGCCGTCAGGTCGCTGCCGTCGCTGCCACTTGGGCATGAAGTG
GlyIleGlnSerAsnGlnThrArgGlnValAlaAlaValAlaHisLeuGlyMetLysCy

311 CGTGTGGTGCAGGAAACTGGGTGAACATAATCCGACGCGGTGTATGACCCGCTAGGCAACA
sValLeuValGlnGluAsnTrpValAsnTyrSerAspAlaValTyrAspArgValGlyAsnI

373 TCGAGATGTCGGGATCATGGCGCTGATGTGCGGCTTGACGCCGCTGGCTTCGATATTGGC
leGluMetSerArgIleMetGlyAlaAspValArgLeuAspAlaAlaGlyPheAspIleGly

435 ATTCGGCCAAGTTGGGAAAAGGCCCATGAGCGATGTCGTGGAACAGGGTGGCAACCGTTTCC
IleArgProSerTrpGluLysAlaMetSerAspValValGluGlnGlyGlyLysProphePr

497 GATTCCAGCGGGTTGCTCCGAGCATCCCTATGGCGGCCCTCGGTTTCGTGGGCTTTGCCGAAG
oileProAlaGlyCysSerGluHisProTyrGlyGlyLeuGlyPheValGlyPheAlaGluG

FIG. 2

SUBSTITUTE SHEET

3/30

559 AGGTGCGGCAGCAGGAAAAGGAACTGGGCTTCAAGTTTGACTACATCGTGGTCTGCTCGGTG
luValArgGlnGlnGluLysGluLeuGlyPheLysPheAspTyrIleValValCysSerVal

621 ACCGGCAGTACGCAGGCGGCATGGTTGTTGGTTTCGGGGCTGACGGTCGTTCGAAGAATGT
ThrGlySerThrGlnAlaGlyMetValValGlyPheAlaAlaAspGlyArgSerLysAsnVa

683 GATTGGTATCGATGCTTCGGCCCAAGCCGGAACAGACCAAGGCACAGATCCTGCGCATCGCCC
IleGlyIleAspAlaSerAlaLysProGluGlnThrLysAlaGlnIleLeuArgIleAlaA

745 GACACCCGCTGAGTTGGTGGAGTTGGGGCCGAGATTACGGAAGAGGACGTGGTCTCGAT
rgHisThrAlaGluLeuValGluLeuGlyArgGluIleThrGluGluAspValValLeuAsp

807 ACGCGTTTTCCTACCCGGAAATATGGCTTGCCCCAACGAAGGCACATTGGAAGCCATCCGACT
ThrArgPheAlaTyrProGluTyrGlyLeuProAsnGluGlyThrLeuGluAlaIleArgLe

869 GTGCGGCAGCCTTGAAGCGGTGCTGACAGACCCGGTATATGAAGGTAATCGATGCACGGCA
uCysGlySerLeuGluGlyValLeuThrAspProValTyrGluGlyLysSerMetHisGlyM

931 TGATTGAAATGGTCCGTCGTGGTGAATTCCCCGAAGGTTCCAAAGTGCTTTACGCACACTTG
etIleGluMetValArgArgGlyGluPheProGluGlySerLysValLeuTyrAlaHisLeu

993 GGTGGGGCGCGCTGAACGCCCTACAGCTTCCTGTTTCGTAAACGGCTAAGCGTAGAAGT
GlyGlyAlaProAlaLeuAsnAlaTyrSerPheLeuPheArgAsnGlyEnd

1055 CTTTGGAGTCATCTGTGGAGCTC 1079

FIG. 2 cont.

SUBSTITUTE SHEET

4/30



FIG. 3

5/30




FIG. 4

6/30




FIG. 5

7/30




FIG. 6

8/30




FIG. 7

9/30




FIG. 8

10/30




FIG. 9

11/30




FIG. 10

12/30




FIG. II

13/30




FIG. 12

14/30




FIG. 13

15/30

1 CTAGAAGGAA GCTTCACGAA ATCGGCCCTT ATTCAAAAAT AACTTTTAAA
51 TAATGAATTT TAAATTTTAA GAAATAATAT CCAATGAATA AATGACATGT
101 AGCATTTTAC CTAAATATTT CAACTATTTT AATCCAATAT TAATTTGTTT
151 TATCCCCAAC AATAGAAAGT CTGTGCAGA CATTAAATCT GACTTTTCCA
201 GTAATAATA TTAATTTTCT GAAGATTTTC GGGTTTAGTC CACAAGTTTT
251 AGTGAGAAAGT TTTGCTCAAA ATTTAGGTG AGAAGGTTG ATATTATCT
301 TTTGTTAAAT TAATTTATCT AGGTGACTAT TATTTATTTA AGTAGAAATT
351 CATATCATT CTTTGGCCAA CTTGTAGTCA TAATAGGAGT AGGTGTATAT
401 GATGAAGGAA TAAACAAGTT CAGTGAAGTG ATTAAAAATAA AATAAATTT
451 AGGTGTACAT CAAATAAAAA CCTTAAAGTT TAGAAAGGCA CCGAATAATT
501 TTGCATAGAA GATATTAGTA AATTATATAA AATAAAAGAA ATGTAGTTGT
551 CAAGTTGTCT TCTTTTTTTT GGATAAAAAAT AGCAGTTGGC TTATGTCATT

FIG. 14

16/30

601 CTTTACAAAC CTCATGCCA CTGTCCAAT TGTGACACT TAACTAATTA
651 GTTTGATTCA TGTATGAATA CTAATAAATT TTTTAGGACT GACTCAAATA
701 TTTTATATT ATCATAGTAA TATTATCTA ATTTTAGGA CCACTTATTA
751 CTAAATAATA AATTAAC TACTATATTA TTGTTGTGAA ACAACAACGT
801 TTTGGTTGTT ATGATGAAAC GTACACTATA TCAGTATGAA AAATTCAAAA
851 CGATTAGTAT AAATTATATT GAAAATTGGA TATTTTCTA TTCTTAATCA
901 GACGTATTGG GTTTCATATT TTAAAAAGGG ACTAAACTTA GAAGAGAAAGT
951 TTGTTTGAAA CTACTTTTGT CTCTTTCTTG TTCCCATTTT TCTCTTAGAT
1001 TTCAAAAAGT GAACTACTTT ATCTCTTTCT TTGTTCACAT TTTATTTTAT
1051 TCTATTATAA ATATGGCATC CTCATATTGA GATTTTGA AATTATTCTA
1101 ATCATTCACA GTGCAAAAGA AGATCTAAAG CCTAGAG

FIG. 14 cont.

17/30

ACAGCCGTCCTAAGGAGAAGATAAGATCTATGAAAAAACTGAAACTGCATGGCTTTAATA 60
MetLysLysLeuLysLeuHisGlyPheAsna

ATCTGACCAAAAGTCTGAGTTTGTATTTACGATATCTGCTACGCCAAACTGCCGAAG 120
snLeuThrLysSerLeuSerPheCysIleTyrAspIleCysTyrAlaLysThrAlaGluG

AGCGCGACGGTTATATTGCTTATATCGATGAACCTCTATAATGCCAACCGTCTGACCGAAA 180
luArgAspGlyTyrIleAlaTyrIleAspGluLeuTyrAsnAlaAsnArgLeuThrGluI

TCCTGTCAGAAACCTGTTCATTATCGGGGCTAATATTCTTAACATCGCCCGCCAGGATT 240
leLeuSerGluThrCysSerIleIleGlyAlaAsnIleLeuAsnIleAlaargGlnAspT

ACGAACACACAGGGTGCCAGCGTCACTATTCTGGTGAGTGAAGAACCGGTTGACCCGAAAC 300
yrGluProGlnGlyAlaSerValThrIleLeuValSerGluGluProValAspProLysL

TCATCGACAAACAGAACACCCCGCCCACTGCCAGAAACGGTCGTTGCCCATCTTGATA 360
euIleAspLysThrGluHisProGlyProLeuProGluThrValValAlaHisLeuAspL

AAAGTCATATTGCGGTACATACCTACCCGGAAGTCATCCTGAAGCGGTTTATGTACCT 420
ysSerHisIleCysValHisThrTyrProGluSerHisProGluGlyGlyLeuCysThrp

FIG. 15

18/30

TCCGGCCGATATTGAAGTCTCTACCTGCGCGTGATTTCTCCGCTGAAGCGCTGAATT 480
heArgAlaAspIleGluValSerThrCysGlyValIleSerProLeuLysAlaLeuAsnT
ACCTGATCCACGAGCTTGAGTCCGATATCGTAACCATGATTAATTCGCGTGGCGGTTTA 540
yrLeuIleHisGlnLeuGluSerAspIleValThrIleAspTyrArgValArgGlyPheT
CCCGGACATTAACGGTATGAAGCACTTTATCGACCATGAGATTAATTCGATTCAGAACT 600
hrArgAspIleAsnGlyMetLysHisPheIleAspHisGluIleAsnSerIleGlnAsnP
TTATGTCTGACGATATGAAGCGCTGTATGACATGGTGATGTGAACGCTCTATCAGGAAA 660
heMetSerAspAspMetLysAlaLeuTyrAspMetValAspValAsnValTyrGlnGluA
ATATCTTCCATACCAAGATGTGTGCTTAAAGAGTTCGACCTTAAGCACTACATGTTCCACA 720
snIlePheHisThrLysMetLeuLeuLysGluPheAspLeuLysHisTyrMetPheHist
CCAAACCGGAAGACTTAACCGACAGCGAGCGCCAGGAAATTACCGCTGCGCTGTGGAAG 780
hrLysProGluAspLeuThrAspSerGluArgGlnGluIleThrAlaAlaLeuTrpLysG
AAATGCGCGAGATTTATTACGGGCGCAATATGCCAGCTGTTTAACGGCTCTGGCGGAGCT 840
luMetArgGluIleTyrTyrGlyArgAsnMetProAlaVal*
CCCAGGCTCCGCCAGATCTATTTACTTCTGCTGCACGAAATTGCGGTAAGCCGCCACGAC 900

FIG. 15 cont.

SUBSTITUTE SHEET

19/30

CCAAACACATAACTTTTAAATACAATTAGTTATTATTAGAAAGTATTTAAAGTAAAGCA 60
CTTGTGAGTTGTGTACATTTTATTATTAATCTTCACTCTTCTTAATCTCTTCAGTTTTTAATT 120
TCTTCACTTCTAAACTCATTTAGTAAAAAAAATGGGATTTGAGATTTGCAAAGACCAAC 180
MetGlyPheGluIleAlaLysThrAsn
TCAATCTTATCAAAATTTGGCTACTAATGAAGAGCATGGCGAAAACCTCGCCATATTTTGAT 240
SerIleLeuSerLysLeuAlaThrAsnGluGluHisGlyGluAsnSerProTyrPheAsp
GGTGGAAGCATACGATAGTGATCCTTTCCACCCTCTAAAAAACCCCAACGGAGTTATC 300
GlyTrpLysAlaTyrAspSerAspProPheHisProLeuLysAsnProAsnGlyValIle
CAAATGGGCTCTTGCTGAAAAATCAGCTTTTGTATTAGACTTGATAGAAGATTGGATTAAAGAGA 360
GlnMetGlyLeuAlaGluAsnGlnLeuCysLeuAspLeuIleGluAspTrpIleLysArg
AACCCAAAAGGTTCAATTTGTTCTGAAGGAATCAAAATCATTCAGGCCCATTTGCCAACTTT 420
AsnProLysGlySerIleCysSerGluGlyIleLysSerPheLysAlaIleAlaAsnPhe
CAAGATTATCATGGCTTGCCCTGAATTCAGAAAAGCGATTGCGAAATTTATGGAGAAAACA 480
GlnAspTyrHisGlyLeuProGluPheArgLysAlaIleAlaLysPheMetGluLysThr
AGAGGAGGAAGAGTTAGATTTGATCCAGAAAAGAGTTGTTATGCGTGGTGGTCCACTGGA 540
ArgGlyGlyArgValArgPheAspProGluArgValValMetAlaGlyGlyAlaThrGly

FIG. 16A

20/30

GCTAATGAGACAAATTATTTGTTGGCTGATCCTGGCGATGCATTTTTAGTACCTTCA 600
AlaAsnGluThrIleIlePheCysLeuAlaAspProGlyAspAlaPheLeuValProSer
CCATACTACCCAGCATTTAAACAGAGATTTAAGATGGAGAACTGGAGTACAACCTTATTCCA 660
ProTyrTyrProAlaPheAsnArgAspLeuArgTrpArgThrGlyValGlnLeuIlePro
ATTCACCTGTGAGAGCTCCAAATAATTTCAAAAATTACTTCAAAAGCAGTAAAAGAGCATAT 720
IleHisCysGluSerSerAsnAsnPheLysIleThrSerLysAlaValLysGluAlaTyr
GAAAATGCACAAAAAATCAAAACATCAAAAGTAAAGGTTTGATTTTGACCCAATCCATCAAAT 780
GluAsnAlaGlnLysSerAsnIleLysValLysGlyLeuIleLeuThrAsnProSerAsn
CCATTGGGCACCACTTTGGACAAAGACACACTGAAAAGTGCTCTTGAGTTTCACCAACCAA 840
ProLeuGlyThrThrLeuAspLysAspThrLeuLysSerValLeuSerPheThrAsnGln
CACAAACATCCACCTTGTGTGACGAAATCTACGCAGCCCACTGTCTTTTGACACGCCCTCAA 900
HisAsnIleHisLeuValCysAspGluIleTyrAlaAlaThrValPheAspThrProGln
TTCGTCAGTATAGCTGAAATCCTCGATGAACAGGAAATGACTTACTGCAACAAAGATTTA 960
PheValS IleAlaGluIleLeuAspGLUGlnGluMetThrTyrCysAsnLysAspLeu
GTTACATCGTCTACAGTCTTTCAAAAGACATGGGGTTACCAGGATTTAGAGTCGGAATC 1020
ValHisIleValTyrSerLeuSerLysAspMetGlyLeuProGlyPheArgValGlyIle

FIG. 16B

ATATATTCTTTAACGACGATGTCGTTAAATTGTGCTAGAAAAATGTCGAGTTTCGGTTTA 1080
IleTyrSerPheAsnAspValValAsnCysAlaArgLysMetSerSerPheGlyLeu
GTATCTACACAAACGCAATATTTTtagCGGCAATGCCATCGGACGAAAAATTCGTCGAT 1140
ValSerThrGlnThrGlnTyrPheLeuAlaAlaMetProSerAspGluLysPheValAsp
AATTTCTAAGAGAAAGCGCGATGAGGTTAGGTAAAGGCACAAACATTTTACTAATGGA 1200
AsnPheLeuArgGluSerAlaMetArgLeuGlyLysArgHisLysHisPheThrAsnGly
CTTGAAAGTAGTGGGAATTAAATGCTTGAAAAATAATGCGGGGCTTTTTTGTGGATGGAT 1260
LeuGluValValGlyIleLysCysLeuLysAsnAsnAlaGlyLeuPheCysTrpMetAsp
TTGCGTCCACTTTTAAGGGAATCGACTTTCGATAGCGAAATGTCGTTATGGAGAGTTATT 1320
LeuArgProLeuLeuArgGluSerThrPheAspSerGluMetSerLeuTrpArgValIle
ATAAACGATGTTAAGCTTAACGTCTCGCTTGGAATCTTCGTTTGAATGTCAAGAGCCAGGG 1380
IleAsnAspValLysLeuAsnValSerLeuGlySerSerPheGluCysGlnGluProGly
TGGTTCGGAGTTTGTTTGCAAAATATGGATGATGGAACGGTTGATATTGCGCTCGCGAGG 1440
TrpPheArgValCysPheAlaAsnMetAspAspGlyThrValAspIleAlaLeuAlaArg

21/30

FIG. 16C

SUBSTITUTE SHEET

22/30

ATTCGGAGGTTTCGTAGGTGTTGAGAAAAGTGGAGATAAATCGAGTTCGATGGAAAAGAG 1500
IleArgArgPheValGlyValGluLysSerGlyAspLysSerSerSerMetGluLysLys

CAACAAATGGAAGAAGAAATAATTGAGACTTAGTTTTTCGAAAAGAAATGTATGATGAAAGT 1560
GlnGlnTrpLysLysAsnAsnLeuArgLeuSerPheSerLysArgMetTyrAspGluSer

GTTTGTCAACCACTTTCGTCACCTATTCCCTCCCTCACCATTAGTTCGTTAAGACTTAATT 1620
ValLeuSerProLeuSerSerProIleProProSerProLeuValArg*

AAAAGGGAAGAAATTTATGTTTTTTTATATATTTTGAAAAAAATTTGTAAGAAATAAGA 1680

TTATAATAGGAAAAGAAAATAAGTATGTAGGATGAGGAGTATTTTCAGAAAATAGTTGTTA 1740

GCGTATGTATTGACAACCTGGTCTATGTACTTAGACATCATATAATTTGTCTTAGCTAATTAA 1800

TGAATGCACAAAAGTGAAGTT

FIG. 16D

SUBSTITUTE SHEET

23/30

GGATCC	ATG	AAT	TTG	AAT	CGT	TTT	AAA	CGT	TAT	CCG	TTG	ACC	TTC	GGT	48
	Met	Asn	Leu	Asn	Arg	Phe	Lys	Arg	Tyr	Pro	Leu	Thr	Phe	Gly	
CCT	TCT	CCC	ATC	ACG	TTG	AAG	CGC	CTC	AGT	GAA	CAC	TTG	GGT	GGC	96
Pro	Ser	Pro	Ile	Thr	Pro	Leu	Lys	Arg	Leu	Glu	His	Leu	Gly	Gly	
AAG	GTC	GAG	CTG	TAT	GCC	AAG	CGT	GAA	GAC	AAC	AGT	GGC	CTG	GCC	144
Lys	Val	Glu	Leu	Tyr	Ala	Lys	Arg	Glu	Asp	Cys	Ser	Gly	Leu	Ala	
TTC	GGC	GGG	AAC	AAA	ACG	CGC	AAG	CTC	GAA	TAT	TTG	ATT	CCC	GAA	192
Phe	Gly	Gly	Asn	Lys	Thr	Arg	Lys	Glu	Tyr	Leu	Ile	Pro	Glu	Ala	
CTC	GAG	CAA	GGC	TGC	GAT	ACC	TTG	GTT	TCC	ATC	GGC	ATC	CAG	TCG	240
Leu	Glu	Gln	Gly	Cys	Asp	Thr	Leu	Val	Ser	Ile	Gly	Ile	Gln	Ser	
AAC	CAG	ACC	CGC	CAG	GTG	GCC	GTT	GCC	GCT	CAC	CTG	GGC	ATG	AAG	288
Asn	Gln	Thr	Arg	Gln	Val	Ala	Ala	Val	Ala	His	Leu	Gly	Met	Lys	
TGC	GTG	CTG	GTG	CAG	GAA	AAC	TGG	GTG	AAC	TAC	TCC	GAT	GCG	TAT	336
Cys	Val	Leu	Val	Gln	Glu	Asn	Trp	Val	Asn	Tyr	Ser	Asp	Ala	Val	Tyr
GAC	CGC	GTT	GGC	AAT	ATC	GAA	ATG	TCT	CGC	ATC	ATG	GGC	GCC	GAG	GTA
Asp	Arg	Val	Gly	Asn	Ile	Glu	Met	Ser	Arg	Ile	Met	Gly	Ala	Val	
CGA	CTG	GAC	GCC	GCC	GGG	TTC	GAT	ATC	GGC	ATT	CGG	CCC	AGC	TGG	GAG
Arg	Leu	Asp	Ala	Ala	Gly	Phe	Asp	Ile	Gly	Ile	Arg	Pro	Ser	Trp	Glu
AAG	GCC	ATG	GAC	GAT	GTG	GTG	GCG	CGG	GGT	GGC	AAG	CCG	TTC	CCG	ATA
Lys	Ala	Met	Asp	Asp	Val	Val	Ala	Arg	Gly	Gly	Lys	Pro	Phe	Pro	Ile
CCG	GCG	GGT	TGT	TCC	GAA	CAC	CCC	TAC	GGC	GGC	CTT	GGG	TTC	GTC	GGC
Pro	Ala	Gly	Cys	Ser	Glu	His	Pro	Tyr	Gly	Gly	Leu	Gly	Phe	Val	Gly

FIG. 17

24/30

TTT	GCC	GAG	GAA	GTG	CGA	GAG	CAG	GAA	AAA	CAA	CTG	GGG	TTC	ACG	TTC	576
Phe	Ala	Glu	Glu	Val	Arg	Glu	Gln	Glu	Lys	Gln	Leu	Gly	Phe	Thr	Phe	
GAC	TAC	ATC	GTG	GTC	TGC	TCT	GTG	ACC	GGC	AGT	ACC	CAG	GCC	GGC	ATG	624
Asp	Tyr	Ile	Val	Val	Cys	Ser	Val	Thr	Gly	Ser	Thr	Gln	Ala	Gly	Met	
GTC	GTC	GGT	TTC	GCC	GCG	GAC	GGC	CGT	TCG	AAG	AAC	GTT	ATC	GGC	ATT	672
Val	Val	Gly	Phe	Ala	Ala	Asp	Gly	Arg	Ser	Lys	Asn	Val	Ile	Gly	Ile	
GAT	GCC	TCG	GCC	AAG	CCG	GAG	CAA	ACC	AAG	GCA	CAG	ATC	CTG	CGT	ATC	720
Asp	Ala	Ser	Ala	Lys	Pro	Glu	Gln	Thr	Lys	Ala	Gln	Ile	Leu	Arg	Ile	
GCC	CGG	CAC	ACC	GCA	GAG	TTG	GTG	GAA	CTG	GGC	CGT	GAG	ATC	ACC	GAA	768
Ala	Arg	His	Thr	Ala	Glu	Leu	Val	Glu	Leu	Gly	Arg	Glu	Ile	Thr	Glu	
GAC	GAC	GTG	GTG	CTC	GAT	ACA	CGT	TTT	GCC	TAC	CCG	GAA	TAC	GGT	TTG	816
Asp	Asp	Val	Val	Leu	Asp	Thr	Arg	Phe	Ala	Tyr	Pro	Glu	Tyr	Gly	Leu	
CCC	AAC	GAA	GGC	ACG	CTG	GAA	GCC	ATT	CGT	TTG	TGC	GGG	AGC	CTG	GAA	864
Pro	Asn	Glu	Gly	Thr	Leu	Glu	Ala	Ile	Arg	Leu	Cys	Gly	Ser	Leu	Glu	
GGT	GTG	CTG	ACC	GAT	CCG	GTG	TAC	GAG	GGC	AAA	TCC	ATG	CAC	GGG	ATG	912
Gly	Val	Leu	Thr	Asp	Pro	Val	Tyr	Glu	Gly	Lys	Ser	Met	His	Gly	Met	
ATT	GAA	ATG	GTC	CGC	CGT	GGC	GAG	TTC	CCC	GAA	GGC	TCC	AAA	GTG	CTG	960
Ile	Glu	Met	Val	Arg	Arg	Gly	Glu	Phe	Pro	Glu	Gly	Ser	Lys	Val	Leu	
TAT	GCG	CAC	TTG	GGT	GGG	GCG	CCT	GCG	CTG	AAT	GCC	TAC	AGC	TTC	CTG	1008
Tyr	Ala	His	Leu	Gly	Gly	Ala	Pro	Ala	Leu	Asn	Ala	Tyr	Ser	Phe	Leu	
TTT	CGT	AAC	GGC	GGATCCGGG												1029
Phe	Arg	Asn	Gly													

FIG. 17 cont.

25/30




FIG. 18

26/30




FIG. 19

27/30

AGATCTATCGATAAGCTTGATGTAATTGGAGGAAGATCAAAATTTTCAAT 50
CCCCATTCTTCGATTGCTTCAATTGAAGTTTCTCCGATGGCGCAAGTTAG 100
CAGAACTGCAATGGTGTGCAGAACCCCATCTCTTATCTCCAATCTCTCGA 150
AATCCAGTCAACGCAAAATCTCCCTTATCGGTTTCTCTGAAGACGCAGCAG 200
CATCCACGAGCTTATCCGATTTCGTCGTCGTGGGGATTGAAGAAGAGTGG 250
GATGACGTTAATTGGCTCTGAGCTTCGTCCTCTTAAGGTCATGTCTCTG 300
TTTCCACGGCGTGCAATGC 318

FIG. 20

28/30

GCATGCTTCACGGTGCAAGCAGCCGTCAGCAACTGCTCGTAAGTCCTCT 50
GGTCTTTCTGGAACCGTCCGTATTCCAGGTGACAAGTCTATCTCCACAG 100
GTCTTCATGTTTGGAGGTCTCGCTAGCGGTGAAACTCGTATCACCGGTC 150
TTTGTGAAGGTGAAGATGTTATCAACACTGGTAAGGCTATGCAAGCTATG 200
GGTGCCAGAAATCCGTAAGGAAGGTGATACTTGGATCATTTGATGGTGTGG 250
TAACGGTGGAATCCTTGCTCCTGAGGCTCCTCTCGATTTCGGTAACGCTG 300
CAACTGGTTGCCGTTTGACTATGGGTCTTGTGTGTTTACGATTTCGAT 350
AGCACTTTTCATTGGTGACGCTTCTCTCACTAAGCGTCCAATGGGTCGTGT 400
GTTGAACCCACTTCGCCGAAATGGGTGTGCGAGGTGAAGTCTGAAGACGGTG 450
ATCGTCTTCCAGTTACCTTGCGTGGACCAAGAACTCCAACGCCAATCACC 500
TACAGGGTACCCTATGGCTTCCGCTCAAGTGAAGTCCGCTGTTCTGCTTGC 550
TGGTCTCAACACCCCAAGGTATCACCACTGTATTATCGAGCCAATCATGACTC 600
GTGACCACACTGAAAAGATGCTTCAAGGTTTTTGGTGCTAACCTTACCGTT 650
GAGACTGATGCTGACGGGTGTGCGGTACCATCCGCTCTTGAAGGTCGTGGTAA 700

FIG. 21

29/30

GCTCACC GGTC AAGT GATT GAT GTTCC AGGTG ATCCAT CCTCT ACTGCTT 750
TCCCAT TGGTT GCTGCC TTGCT TTGTTCC AGGTTCCGAC GTCACCAT CCTT 800
AACGTTT GATGA ACCCA ACCCGT ACTGGTCT CATCTT GACTCT GCAGGA 850
AATGGTG CCGACA TCGAAG TGATCA ACCCAC GTCTTG CTGGTG GAGAAG 900
ACGTGGC TGA CTGCGT GTCGTTCT TCTACT TTGAAG GGTGTT ACTGTT 950
CCAGAAG ACCGTG CTCTTCT ATGATCGAC GAGTATCCA ATTCTCGCTGT 1000
TGCAGCT GCATT CGCTGA AGGTGCTACCGT TATGAACGGTTTGG AAGAAC 1050
TCCGTGT AAAGG AAAGC GACCGTCTT TCTGCTGTCGCA AACGGTCTCAAG 1100
CTCAACGGT GTTGA TTGCGATGA AGGTGAGACTTCTCTCGT CGTGCGTGG 1150
TCGTCCCTGACGGT AAGGTCTCGGT AACGCTTCTG GAGCAGCTGTCGCTA 1200
CCCACCTCGATC ACCGTATCGCTATGAGCTT CCTCGTTATGGGTCTCGTT 1250
TCTGAAA ACCCTGTTACTGTGATGATGCTACTATGATCGCTACTAGCTT 1300
CCCAGAGTTCATGGATT TGATGGCTGGTCTTGGAGCTAAGATCGAACTCT 1350
CCGACACTAAGGCTGCTTGATGAGCTC 1377

FIG. 21 cont.

SUBSTITUTE SHEET

30/30

TCATCAAAATATTTAGCAGCATTCAGATTGGGTTCAATCAACAAGGTAC 50
GAGCCATATCACTTTATTCAAATTGGTATCGCCAAAACCAAGAAGGAACT 100
CCCATCCTCAAAGGTTTGTAAAGGAAGAAATTCTCAGTCCAAAGCCTCAACA 150
AGGTCAGGGTACAGAGTCTCCAAACCATTAGCCAAAAGCTACAGGAGATC 200
AATGAAGAATCTTCAATCAAAGTAAACTACTGTTCAGCACATGCATCAT 250
GGTCAGTAAGTTTCAGAAAAAGACATCCACCGAAGACTTAAAGTTAGTGG 300
GCATCTTTGAAAGTAATCTTGTCAACATCGAGCAGCTGGCTTGTGGGGAC 350
CAGACAAAAAGGAATGGTGCAGAATTGTTAGGCGCACCTACCAAAAAGCA 400
TCTTTGCCCTTTATTGCAAAGATAAAGCAGATTCCCTCTAGTACAAGTGGGG 450
AACAAAAATAACGTGGAAAAGAGCTGTCCCTGACAGCCCACTCACTAATGCG 500
TATGACGAACGCAGTGACGACCACAAAAGAATTCCCTCTATATAAGAAGG 550
CATTCAATCCCATTTGAAGGATCATCAGATATAACCAATATTCTC 596

FIG. 22

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

PCT/US 91/09437

International Application No

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C12N15/60;	C12N15/82;	C12N9/88; A01H5/00
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C12N ; A01H	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
P,X	WO,A,9 116 417 (CALGENE) 31 October 1991 see the whole document ---	
P,X	THE PLANT CELL. vol. 3, no. 11, November 1991, ROCKVILLE, MD, USA. pages 1187 - 1193; KLEE, H. J., ET AL.: 'Control of ethylene synthesis by expression of a bacterial enzyme in transgenic tomato plants' see the whole document ---	1-3, 6-10, 13-16, 18-30
A	AGRIC. BIOL. CHEM. vol. 42, no. 10, 1978, pages 1825 - 1831; HONMA, M., ET AL.: 'Metabolism of 1-aminocyclopropane-1-carboxylic acid' see the whole document --- -/--	1,6,7, 13,14
¹⁰ Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
23 APRIL 1992	20. 05. 92	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	MADDOX A.D.	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category °	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	NATURE. vol. 346, 19 July 1990, LONDON GB pages 284 - 287; HAMILTON, A. J., ET AL.: 'Antisense gene that inhibits synthesis of the plant hormone ethylene in transgenic plants' see the whole document ---	8-17

US 9109437
SA 54963

The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 23/04/92

WO-A-9116417	31-10-91	None
--------------	----------	------