



(86) Date de dépôt PCT/PCT Filing Date: 2000/02/28
(87) Date publication PCT/PCT Publication Date: 2000/09/08
(45) Date de délivrance/Issue Date: 2006/01/10
(85) Entrée phase nationale/National Entry: 2001/08/24
(86) N° demande PCT/PCT Application No.: EP 2000/001660
(87) N° publication PCT/PCT Publication No.: 2000/052007
(30) Priorité/Priority: 1999/03/03 (99104306.8) EP


(51) Cl.Int.⁷/Int.Cl.⁷ C07D 417/14, A61K 31/506, A61P 9/00, C07D 401/14

(72) Inventeurs/Inventors:
BREU, VOLKER, DE;
COASSOLO, PHILIPPE, FR;
NEIDHART, WERNER, FR;
ROUX, SEBASTIEN, CH;
WEISS, PETER, CH

(73) Propriétaire/Owner:
F. HOFFMANN-LA ROCHE AG, CH

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : DERIVES 4-HETEROCYCLYSULFONAMIDYL-6-METHOXY-5-(2-METHOXY-PHENOXY)-2-PYRIDYL-PYRIMIDINE, PREPARATIONS ET UTILISATIONS COMME ANTAGONISTES DU RECEPTEUR D'ENDOTHELINE
(54) Title: 4-HETEROCYCLYSULFONAMIDYL-6-METHOXY-5-(2-METHOXY-PHENOXY)-2-PYRIDYL-PYRIMIDINE DERIVATIVES, THEIR PREPARATION AND USE AS ENDOTHELIN RECEPTOR ANTAGONISTS



(I)

(57) Abrégé/Abstract:




The compounds of formula (I) wherein R¹ is a heterocyclic residue selected from pyridyl and thiazolyl, where said heterocyclic residue may optionally be substituted with lower alkyl or lower alkenyl; and pharmaceutically acceptable salts thereof, are inhibitors of endothelin receptors and can therefore be used for the treatment of disorders which are associated with abnormal vascular tone and endothelial dysfunction; and the corresponding intermediates of formulae (III) and (V).



**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 401/14, 417/14, A61K 31/505, A61P 9/00	A1	(11) International Publication Number: WO 00/52007 (43) International Publication Date: 8 September 2000 (08.09.00)
(21) International Application Number: PCT/EP00/01660 (22) International Filing Date: 28 February 2000 (28.02.00) (30) Priority Data: 99104306.8 3 March 1999 (03.03.99) EP (71) Applicant: F. HOFFMANN-LA ROCHE AG [CH/CH]; Gren- zacherstrasse 124, CH-4070 Basle (CH). (72) Inventors: BREU, Volker; Belchenstrasse 7, D-79418 Schlien- gen (DE). COASSOLO, Philippe; 25, rue du Cantal, F-68270 Wittenheim (FR). NEIDHART, Werner; 9, rue du Steinler, F-68220 Hagenthal le Bas (FR). ROUX, Sébastien; Missionsstrasse 21b, CH-4055 Basle (CH). WEISS, Peter; Rosentalstrasse 52, CH-4058 Basle (CH). (74) Agent: WITTE, Hubert; Grenzacherstrasse 124, CH-4070 Basle (CH).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: 4-HETEROCYCLYSULFONAMIDYL-6-METHOXY-5-(2-METHOXY-PHENOXY)-2-PYRIDYL-PYRIMIDINE DERIVATIVES, THEIR PREPARATION AND USE AS ENDOTHELIN RECEPTOR ANTAGONISTS (57) Abstract The compounds of formula (I) wherein R ¹ is a heterocyclic residue selected from pyridyl and thiazolyl, where said heterocyclic residue may optionally be substituted with lower alkyl or lower alkenyl; and pharmaceu- tically acceptable salts thereof, are inhibitors of endothelin receptors and can therefore be used for the treatment of disorders which are associated with abnormal vascular tone and endothelial dysfunction; and the corresponding intermediates of formulae (III) and (V).		



4-HETEROCYCLYSULFONAMIDYL-6-METHOXY-5-(2-METHOXY-PHENOXY)-2-PYRIDYL-PYRIMIDINE
DERIVATIVES, THEIR PREPARATION AND USE AS ENDOTHELIN RECEPTOR ANTAGONISTS

The present invention relates to heterocyclic sulfonamides and their use as medicaments. In particular, the present invention relates to compounds of the formula (I)



wherein

- 5 R^1 is a heterocyclic residue selected from pyridyl and thiazolyl, where said heterocyclic residue may optionally be substituted with lower alkyl or lower alkenyl;

and pharmaceutically acceptable salts thereof.

- 10 The present invention also relates to a pharmaceutical composition comprising a compound of formula (I) and a pharmaceutically acceptable carrier and/or adjuvant.

Furthermore the present invention relates to the use of such compounds for the preparation of medicaments for the treatment or prophylaxis of disorders which are associated with abnormal vascular tone and endothelial dysfunction.

- 15 The present invention also relates to processes for the preparation of the compounds of formula (I).

In addition, the present invention relates to a method for the prophylactic and/or therapeutic treatment of disorders which are associated with abnormal vascular tone and endothelial dysfunction, which method comprises administering a compound of formula
20 (I) to a human being or an animal.

The sulfonamides of the present invention are inhibitors of endothelin receptors. They can accordingly be used for the treatment of disorders which are associated with abnormal vascular tone and endothelial dysfunction. EP 0 713 875 discloses sulfonamide
5 compounds as endothelin receptor inhibitors. However, the compounds of the present invention have a high antagonistic potency in vitro and show unexpectedly high plasma levels following oral administration.

Unless otherwise indicated the following definitions are set forth to illustrate and
10 define the meaning and scope of the various terms used to describe the invention herein.

The term "lower alkyl" refers to a branched or straight chain monovalent saturated aliphatic hydrocarbon radical of one to seven carbon atoms, preferably one to four carbon atoms. This term is further exemplified by such radicals as methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *s*-butyl, *t*-butyl and the like.

15 The term „lower alkenyl“ refers to a lower alkyl group containing one or more double bond(s) in the alkylene chain.

The term "pharmaceutically acceptable salts" embraces salts of the compounds of formula (I) with inorganic or organic acids such as hydrochloric acid, hydrobromic acid, nitric acid, sulphuric acid, phosphoric acid, citric acid, formic acid, maleic acid, acetic
20 acid, succinic acid, tartaric acid, methanesulphonic acid, p-toluenesulphonic acid and the like, which are non-toxic to living organisms. It also includes salts with inorganic or organic bases such as alkali salts like sodium and potassium salts, alkaline earth metal salts like calcium and magnesium salts, N-methyl-D-glutamine salts and salts with amino acids like arginine, lysine and the like.

- 3 -

More particularly, the present invention relates to compounds of the above formula (I), wherein

R^1 is a heterocyclic residue selected from pyridyl and thiazolyl, where said heterocyclic residue may optionally be substituted with lower alkyl or lower alkenyl. The term "lower alkyl" preferably means methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *s*-butyl or *t*-butyl, more preferably methyl, ethyl or isopropyl, still more preferably methyl or isopropyl, most preferably methyl. The term "lower alkenyl" preferably means vinyl, 1-propenyl, allyl, isopropenyl, 1-butenyl, 2-butenyl or 3-butenyl, more preferably vinyl, 1-propenyl, allyl and isopropenyl, most preferably isopropenyl.

Preferred heterocyclic residues in R^1 are 2-pyridyl and 2-thiazolyl, most preferred is 2-pyridyl. In a preferred embodiment, the heterocyclic residues in R^1 are substituted with lower alkyl or lower alkenyl, with lower alkyl being preferred.

Particularly preferred residues R^1 are 5-methyl-pyridine-2-yl, 5-isopropyl-pyridine-2-yl, 5-isopropenyl-pyridine-2-yl and 5-methyl-thiazol-2-yl. More preferred are 5-methyl-pyridine-2-yl, 5-isopropyl-pyridine-2-yl and 5-isopropenyl-pyridine-2-yl. Still more preferred are 5-methyl-pyridine-2-yl and 5-isopropyl-pyridine-2-yl. Most preferred is 5-methyl-pyridine-2-yl.

Particularly preferred compounds of formula (I) are 5-methyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide, 5-isopropyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide, 5-isopropenyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide and 5-methyl-thiazole-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide.

More particularly preferred compounds of formula (I) are 5-methyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide and 5-isopropyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide, most preferred is 5-methyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide.


The compounds of general formula (I) can be prepared in analogy to known processes or processes described below and summarized in the following schemes 1 and 2:

4,6-Dichloro-5-(2-methoxy-phenoxy)-2-(pyridin-4-yl)-pyrimidine (described in EP 0 799 209) can be transformed to the intermediate of general formula (III) - according to scheme 1 - on reaction with an appropriate sulfonamide of general formula (II), wherein R¹ is as defined in claim 1, in a suited solvent such as DMSO or DMF at room temperature or at elevated temperature and in the presence of a suited base such as potassium carbonate.

The corresponding sulfonamides can also be applied in above reaction in form of their pre-formed sodium or potassium salts.

Compounds of formula (III) can be further transformed to compounds of general formula (I) by treatment with sodium methylate in a solvent such as methanol.

Scheme 1



15


The heterocyclic sulfonamides of general formula (II) are either already known in the literature, prepared in a manner analogous to established procedures and/or can be derived from corresponding mercapto derivatives in analogy to a known reaction sequence comprising oxidation with Cl₂ in an acidic aqueous medium, such as diluted aqueous HCl, to yield the corresponding sulfonyl chlorides which can be transformed with liquid ammonia or aqueous ammonium hydroxide to the sulfonamides. The corresponding sodium or potassium salts can be obtained on treatment with sodium or potassium alkoxide in an appropriate solvent such as methanol.

25

Alternatively, compounds of general formula (I) can be prepared – according to scheme 2 - starting from 4-[4,6-dichloro-5-(2-methoxy-phenoxy)-pyrimidin-2-yl]-pyridine-1-oxide (preparation described in EP 0 799 209) in a reaction sequence analogous to that of scheme 1, to give compounds of general formula (V).

- 5 Pyridine-N-Oxide reduction of compounds of general formula (V) can be accomplished with a reagent system such as TiCl_4/NaI in analogy to a method described in Chem. Ber. 123, 647 (1990), or with a trialkylsilyl chloride (for example, *t*-butyldimethylsilyl chloride) in the presence of a suited base, such as triethylamine, and with acetonitrile as a solvent. Other methods for pyridine-N-Oxide deoxygenation are
 10 known per se and comprise, for example, catalytic hydrogenation or reduction with reducing agents such as trivalent phosphorus compounds or metals in acid.

Scheme 2



15

The intermediates of above formula (III) and formula (V), wherein R^1 is as defined above 1, preferably wherein R^1 is 5-methyl-pyridine-2-yl or 5-isopropyl-pyridine-2-yl, and the pharmaceutically acceptable salts thereof are new and are also subject of the present invention.

The inhibitory activity of the compounds of formula (I) on endothelin receptors can be demonstrated using the test procedures described hereinafter:

I. Inhibition of endothelin binding to recombinant human ET_A receptors expressed in
Baculovirus-infected insect cells

A cDNA coding for human ET_A receptors of human placenta was cloned (M. Adachi, Y.-Y. Yang, Y. Furuichi and C. Miyamoto, BBRC 180, 1265-1272) and expressed in the Baculovirus-insect cell system. Baculovirus-infected insect cells from a 23 l fermenter are centrifuged off (3000 x g, 15 minutes, 4°C) 60 hours after the infection, re-suspended in Tris buffer (5 mM, pH 7.4, 1 mM MgCl₂) and again centrifuged. After a further re-suspension and centrifugation the cells are suspended in 800 ml of the same buffer and freeze-dried at -120°C. The cells disintegrate when the suspension in this hypotonic buffer mixture is thawed. After a repeated freeze-drying/thawing cycle the suspension is homogenised and centrifuged (25000 x g, 15 minutes, 4°C). After suspension in Tris buffer (75 mM, pH 7.4, 25 mM MgCl₂, 250 mM sucrose) 1 ml aliquots (protein content about 3.5 mg/ml) are stored at -85°C.

For the binding assay, the freeze-dried membrane preparations are thawed and, after centrifugation at 20°C and 25000 g for 10 minutes, re-suspended in assay buffer (50 mM Tris buffer, pH 7.4, containing 25 mM MnCl₂, 1 mM EDTA and 0.5% bovine serum albumin). 100 µl of this membrane suspension containing 70 µg of protein are incubated with 50 µl of ¹²⁵I-endothelin (specific activity 2200 Ci/mMol) in assay buffer (25000 cpm, final concentration 20 pM) and 100 µl of assay buffer containing varying concentrations of test compound. The incubation is carried out at 20°C for 2 hours or at 4°C for 24 hours. The separation of free and membrane-bound radio- ligands is carried out by filtration over a glass fibre filter. The inhibitory activity of compounds of formula (I) determined in this test procedure is given in Table 1 as the IC₅₀, i.e. as the concentration [nM] which is required to inhibit 50% of the specific binding of ¹²⁵I-endothelin.

Table 1.

Compound of example	1	2	3
IC ₅₀ [nM]	≤ 50	≤ 50	≤ 50

II. Inhibition of endothelin-induced contractions in isolated rat aorta rings

Rings with a length of 5 mm were cut out from the thorax aorta of adult Wistar-Kyoto rats. The endothelium was removed by lightly rubbing the internal surface. Each ring was immersed at 37°C in 10 ml of Krebs-Henseleit solution in an isolated bath while
5 gassing with 95% O₂ and 5% CO₂. The isometric stretching of the rings was measured. The rings were stretched to a pre-tension of 3 g. After incubation for 10 minutes with the test compound or vehicle cumulative dosages of endothelin-1 were added. The activity of the test compound was ascertained by the observed shift to the right of the dosage-activity curve of endothelin-1 in the presence of different concentrations of antagonist. This shift
10 to the right (or "dose ratio", DR) corresponds to the quotient from the EC₅₀ values of endothelin-1 in the presence and in the absence of antagonist, with the EC₅₀ value denoting the endothelin concentration required for a half-maximum contraction.

The corresponding pA₂ value, which is a measure of the activity of the test compound, was calculated using a computer programme according to the following
15 equation from the "dose ratio" DR for each individual dosage-activity curve.

$$pA_2 = \log(DR-1) - \log(\text{antagonist-concentration})$$

The EC₅₀ of endothelin in the absence of test compounds is 0.3 nM.

The pA₂ values obtained with compounds of formula (I) are given in the following Table 3.

20

Table 2.

Compound of example	1	2
pA ₂	≥ 8.0	≥ 8.0

III. Pharmacokinetics of the endothelin receptor antagonists

25 The pharmacokinetics of the newly synthesised endothelin receptor antagonists were assessed in Wistar rats. The test compounds were dissolved in DMSO at a concentration of 5 mg/mL and administered orally by gavage at a dose of 1 mL/kg body weight corresponding to 5 mg/kg body weight. Two rats were administered per test compound. Blood samples were collected from the retro-orbital sinus at 1 and 5 h post
30 dose in one rat, and at 3 and 7 h post dose in the other rat. In addition a terminal 24 h blood sample was collected from both rats by heart puncture. Blood was collected on

EDTA-NaF. Plasma was derived by centrifugation at 2000 g at +4°C for 15 min. Plasma samples were assayed for active drug related material with a bioassay, based on the binding competition of tested compounds and ^{125}I ET-1 on recombinant ET_A receptors.

Quantitation of plasma samples was by comparison to a calibration curve build up from control rat plasma spiked with known concentrations of the test compounds. Selected findings are summarised in the following table:

Table 3.

Test compound of example	Peak concentration in rat plasma (ng/mL)	Area under the plasma concentration time curve (ng.h/mL)
1	$\geq 1'500$	$\geq 10'000$

On the basis of their capability of inhibiting endogenous endothelin binding, the compounds of formula (I) can be used as medicaments for the treatment of disorders which are associated with abnormal vascular tone and endothelial dysfunction.

Therefore, the application field of the compounds of formula (I) could be heart
5 failure (acute and chronic), systemic and pulmonary hypertension, acute ischaemic coronary syndrome, angina pectoris, renal failure (acute and chronic), organ transplant (e.g. liver, heart, kidney), cyclosporin nephrotoxicity, vasospastic disease (subarachnoid haemorrhage but also haemorrhagic and non-haemorrhagic stroke, Raynaud syndrome), peripheral artery occlusive disease, prevention of restenosis after stent or balloon
10 angioplasty, septic shock or multiple organ failure as that occurring in intensive care, asthma, chronic obstructive pulmonary disease, gastric and duodenal ulcer, liver cirrhosis, pancreatitis (acute and chronic), inflammatory bowel disease, fibrosis, atherosclerosis, obesity, glaucoma, prostatic adenoma, migraine, erectile dysfunction, adjunct to cancer therapy as well as other disorders associated with endothelin activities.

15

The compounds of formula (I) can also be administered in combination with antihypertensive drugs, antiarrhythmics, anti angina, antithrombotic and lipid lowering agents as well as antioxidants.

20 It will be appreciated that the compounds of general formula (I) in this invention may be derivatised at functional groups to provide prodrug derivatives which are capable of conversion back to the parent compounds in vivo. Additionally, any physiologically acceptable equivalents of the compounds of general formula (I), which are capable of producing the parent compounds of general formula (I) in vivo, are within the scope of
25 this invention.

As mentioned earlier, medicaments containing a compound of formula (I) are also an object of the present invention, as is a process for the manufacture of such medicaments, which process comprises bringing one or more compounds of formula (I)
30 and, if desired, one or more other therapeutically valuable substances into a galenical administration form.

The pharmaceutical compositions may be administered orally, for example in the form of tablets, coated tablets, dragées, hard or soft gelatine capsules, solutions, emulsions or suspensions. Administration can also be carried out rectally, for example using suppositories; locally or percutaneously, for example using ointments, creams, gels or solutions; or parenterally, e.g. intravenously, intramuscularly, subcutaneously, intrathecally or transdermally, using for example injectable solutions. Furthermore, administration can be carried out sublingually or as ophthalmological preparations or as an aerosol, for example in the form of a spray.

For the preparation of tablets, coated tablets, dragées or hard gelatine capsules the compounds of the present invention may be admixed with pharmaceutically inert, inorganic or organic excipients. Examples of suitable excipients for tablets, dragées or hard gelatine capsules include lactose, maize starch or derivatives thereof, talc or stearic acid or salts thereof.

Suitable excipients for use with soft gelatine capsules include for example vegetable oils, waxes, fats, semi-solid or liquid polyols etc.; according to the nature of the active ingredients it may however be the case that no excipient is needed at all for soft gelatine capsules.

For the preparation of solutions and syrups, excipients which may be used include for example water, polyols, saccharose, invert sugar and glucose.

For injectable solutions, excipients which may be used include for example water, alcohols, polyols, glycerine, and vegetable oils.

For suppositories, and local or percutaneous application, excipients which may be used include for example natural or hardened oils, waxes, fats and semi-solid or liquid polyols.

The pharmaceutical compositions may also contain preserving agents, solubilising agents, stabilising agents, wetting agents, emulsifiers, sweeteners, colorants, odorants, salts for the variation of osmotic pressure, buffers, coating agents or antioxidants. As mentioned earlier, they may also contain other therapeutically valuable agents.

It is a prerequisite that all adjuvants used in the manufacture of the preparations are non-toxic.

Preferred forms of use are intravenous, intramuscular or oral administration, most preferred is oral administration. The dosages in which the compounds of formula (I) are administered in effective amounts depend on the nature of the specific active ingredient, the age and the requirements of the patient and the mode of application. In general, dosages of about 0.01-10 mg/kg body weight per day come into consideration.

The following Examples shall illustrate preferred embodiments of the present invention but are not intended to limit the scope of the invention. Of the abbreviations used therein, MeOH signifies methanol, AcOEt signifies ethyl acetate, DMF signifies dimethylformamide, RT signifies room temperature, HPLC signifies high performance liquid chromatography, TLC signifies thin layer chromatography, ISN signifies Ion Spray Mass Spectrometry – negative mode, EI signifies Electron Impact Mass Spectrometry and M signifies molecular mass.

Example 1

10 a) To a solution of 6.9 g sodium in MeOH (300 ml) were added 14.52 g of 5-methyl-pyridine-2-sulfonic acid [6-chloro-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide at RT and the mixture was refluxed for 5 days until completion of the reaction according to TLC analysis. The reaction mixture was concentrated in vacuo to half its volume upon which the crude reaction product precipitated as a sodium salt. It was
15 filtered off by suction and dried in a high vacuum. The solid was dissolved in water, which was then made acidic by addition of acetic acid. The precipitating free sulfonamide was extracted into Me₂Cl₂. The organic layer was dried over Mg₂SO₄, concentrated on a rotary evaporator, and the crystalline solid that had formed was filtered off. It was then dried in a high vacuum for 12 h at 120 °C to give the desired 5-methyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide as white crystals.
20 Melting point 225-226 °C. ISN mass spectrum, m/e 478.2 (M-1 calculated for C₂₃H₂₁N₅O₅S₁: 478).

C₂₃H₂₁N₅O₅S₁: Calc: C 57.61; H 4.41; N 14.61; S 6.69. Found: C 57.56; H 4.38; N 14.61; S 6.83

25 Preparation of the starting material:

b) 11.3 g of 4,6-dichloro-5-(2-methoxy-phenoxy)-2-pyridin-4-yl)-pyrimidine and 19.66 g of 5-methylpyridyl-2-sulfonamide potassium salt (preparations described in EP 0 799 209) were dissolved in DMF (255 ml) under argon. The solution was stirred for 2 h at 40°C until completion of the reaction according to TLC analysis. The reaction mixture
30 was cooled to RT and the solvent removed in a high vacuum. The residue was suspended in water (850 ml), acetic acid (85 ml) was added and the mixture was stirred for 30 minutes at RT. The solid that precipitated was collected by filtration and dried in a high vacuum at 60 °C for 16 h to give 5-methyl-pyridine-2-sulfonic acid [6-chloro-5-(2-

- 12 -

methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide as yellow crystals. Melting point 177-179 °C. ISN mass spectrum, m/e 482.2 (M-1 calculated for C₂₂H₁₈ClN₅O₅S₁: 482).

Example 2

5 a) In analogy to example 1a), from 5-isopropyl-pyridine-2-sulfonic acid[6-chloro-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide and NaOMe, there was obtained 5-isopropyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide as slightly yellow crystals. Melting point 195-197 °C. ISN mass spectrum, m/e 506.2 (M-1 calculated for C₂₅H₂₅N₅O₅S₁: 506).

10 C₂₃H₂₁N₅O₅S₁: Calc: C 59.16; H 4.96; N 13.80; S 6.32. Found: C 58.99; H 4.90; N 13.83; S 6.37

Preparation of the starting material:

b) In analogy to example 1b), from 4,6-dichloro-5-(2-methoxy-phenoxy)-2-pyridin-4-yl)-pyrimidine and 5-isopropylpyridyl-2-sulfonamide potassium salt
15 (preparation described in EP 799 209) there was obtained 5-isopropyl-pyridine-2-sulfonic acid[6-chloro-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide as a brown crystalline solid.

Example 3

20 a) A solution of 0.156 g of 5-isopropenyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-(1-oxy-pyridin-4-yl)-pyrimidin-4-yl]-amide in acetonitrile (2 ml) was treated at RT with 0.025 ml TiCl₄ followed by 0.135 g of sodium iodide and then stirred at RT for 20 h. Additional 0.1 ml of TiCl₄ were added, the mixture was refluxed for 1 h and then further 0.1 ml of TiCl₄ were added and refluxing was continued
25 for 3 h to complete the reaction according to HPLC analysis. The reaction mixture was cooled to RT, concentrated on rotary evaporator, the residue taken up in AcOEt, which was washed with water, dried over NaSO₄ and finally removed in vacuo. The residue was purified by preparative thin layer chromatography, with AcOEt/MeOH: 9/1 as solvent system, to give the desired 5-isopropenyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-
30 methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide as a light yellow solid. ISN mass spectrum, m/e 504.2 (M-1 calculated for C₂₅H₂₃N₅O₅S₁: 504).

- 13 -

Preparation of the starting material:

b) To a solution of 5-isopropyl-pyridine-2-sulfonamide potassium salt (synthesis described in EP 799 209) in water (10 ml) were added 1.2 g of KMnO_4 at RT and the mixture was then refluxed for 30 minutes. The mixture was cooled to RT, acidified with diluted HCl and the product was extracted into AcOEt. The organic layer was washed with water, dried over Na_2SO_4 and concentrated in vacuo to give (5-(1-hydroxy-1-methyl-ethyl))-pyridine-2-sulfonic acid amide as yellow oil. EI mass spectrum, m/e 216 (M calculated for: $\text{C}_8\text{H}_{12}\text{N}_2\text{O}_3\text{S}$ 216).

c) A solution of 0.1 g of (5-(1-hydroxy-1-methyl-ethyl))-pyridine-2-sulfonic acid amide in $\text{CF}_3\text{CO}_2\text{H}$ (2 ml) was refluxed for 20 h. The solvent was then removed in vacuo to give 5-isopropenyl-pyridine-2-sulfonic acid amide as a white solid which was essentially pure. EI mass spectrum, m/e 198 (M calculated for $\text{C}_8\text{H}_{10}\text{N}_2\text{O}_2\text{S}$: 198).

The corresponding potassium salt was prepared from the sulfonamide on treatment with potassium *t*-butylate in MeOH.

d) In analogy to example 1b), from 4-[4,6-dichloro-5-(2-methoxy-phenoxy)-pyrimidin-2-yl]-pyridine-1-oxide (synthesis described in EP 799 209) and 5-isopropenyl-pyridine-2-sulfonic acid amide potassium salt, there was obtained 5-isopropenyl-pyridine-2-sulfonic acid [6-chloro-5-(2-methoxy-phenoxy)-2-(1-oxy-pyridin-4-yl)-pyrimidin-4-yl]-amide as a light yellow solid. ISN mass spectrum, m/e 524.3 (M-1 calculated for $\text{C}_{24}\text{H}_{20}\text{ClN}_5\text{O}_5\text{S}$: 524).

e) In analogy to example 1a), from 5-isopropenyl-pyridine-2-sulfonic acid [6-chloro-5-(2-methoxy-phenoxy)-2-(1-oxy-pyridin-4-yl)-pyrimidin-4-yl]-amide and sodium methoxide there was obtained 5-isopropenyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-(1-oxy-pyridin-4-yl)-pyrimidin-4-yl]-amide as a light yellow solid. ISN mass spectrum, m/e 520.2 (M-1 calculated for $\text{C}_{25}\text{H}_{23}\text{N}_5\text{O}_6\text{S}$: 520).

30

Example 4

a) A suspension of 0.4 g of 5-methyl-thiazole-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-(1-oxy-pyridin-4-yl)-pyrimidin-4-yl]-amide in acetonitrile (5 ml)

was treated subsequently with 1.12 ml of triethylamine and 1.243 g of *t*-butyldimethylsilyl chloride at RT. The mixture was stirred 5 minutes at RT, its pH adjusted to neutral (by addition of triethylamine), and then heated to reflux for 60 h. The reaction mixture was cooled to RT and concentrated on a rotary evaporator. The residue was taken up in
5 CH₂Cl₂, which was subsequently washed with acetic acid and water. The organic layer was dried over MgSO₄ and the solvent removed in vacuo. The residue was applied to a silica gel column with AcOEt/MeOH: 4/1 as eluent. Combination of the purified fractions and concentration in vacuo gave the desired 5-methyl-thiazole-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide as a beige crystalline solid.
10 ISN mass spectrum, m/e 484.2 (M-1 calculated for C₂₁H₁₉N₅O₅S₁: 484).

Preparation of the starting material:

b) 2.23 g of 5-methylene-thiazolidine-2-thione (preparation described in: Liebigs Ann. Chem., 1985, 58-64) were dissolved in 36% aqueous HCl (150 ml), cooled to -20°C and Cl₂ was bubbled through the solution for 0.5 h while keeping its temperature below
15 -20°C. Ether (400 ml cooled to -15°C) was then added and after stirring for 5 minutes the layers were separated. The organic layer was treated with liquid NH₃ (200 ml) and the mixture allowed warming slowly to RT. The solvent was removed in vacuo to give 5-methyl-thiazole-2-sulfonic acid amide as an off-white solid. EI mass spectrum, m/e 178 (M calculated for C₄H₆N₂O₂S₂: 178).

20 The corresponding potassium salt was prepared from the sulfonamide on treatment with potassium *t*-butylate in MeOH.

c) In analogy to example 1b), from 5-methyl-thiazole-2-sulfonic acid amide potassium salt, and 4-[4,6-dichloro-5-(2-methoxy-phenoxy)-pyrimidin-2-yl]-pyridine-
25 1-oxide there was obtained 5-methyl-thiazole-2-sulfonic acid [6-chloro-5-(2-methoxy-phenoxy)-2-(1-oxy-pyridin-4-yl)-pyrimidin-4-yl]-amide as a light yellow solid. ISN mass spectrum, m/e 504 (M-1 calculated for C₂₀H₁₆ClN₅O₅S₂: 504).

d) In analogy to example 1a), from 5-methyl-thiazole-2-sulfonic acid [6-chloro-5-
30 (2-methoxy-phenoxy)-2-(1-oxy-pyridin-4-yl)-pyrimidin-4-yl]-amide and sodium methoxide there was obtained 5-methyl-thiazole-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-(1-oxy-pyridin-4-yl)-pyrimidin-4-yl]-amide as a light yellow solid. ISN mass spectrum, m/e 500.1 (M-1 calculated for C₂₁H₁₉ClN₅O₆S₂: 500).

- 15 -

Example A

Tablets containing the following ingredients can be produced in a conventional manner:

	<u>Ingredients</u>	<u>mg per tablet</u>
5	Compound of formula (I)	10.0 - 100.0
	Lactose	125.0
	Corn starch	75.0
	Talc	4.0
	Magnesium stearate	1.0
10		

Example B

Capsules containing the following ingredients can be produced in a conventional manner:

	<u>Ingredients</u>	<u>mg per capsule</u>
15	Compound of formula (I)	25.0
	Lactose	150.0
	Corn starch	20.0
	Talc	5.0

20

Example C

Injection solutions can have the following composition:

	Compound of formula (I)	3.0 mg
	Gelatine	150.0 mg
	Phenol	4.7 mg
25	Water for injection	ad 1.0 ml

WO 00/52007

PCT/EP00/01660

- 16 -


Example D

500 mg of compound of formula (I) are suspended in 3.5 ml of Myglyol 812 and 0.08 g of benzyl alcohol. This suspension is filled into a container having a dosage valve. 5.0 g of Freon^{*} 12 are filled into the container under pressure through the valve. The Freon^{*} is dissolved in the Myglyol-benzyl alcohol mixture by shaking. This spray container contains about 100 single doses which can be applied individually.

* trademark

Claims

1. A compound of formula (I)



5 wherein

R^1 is a heterocyclic residue selected from pyridyl and thiazolyl, where said heterocyclic residue may optionally be substituted with lower alkyl or lower alkenyl;

and pharmaceutically acceptable salts thereof.

- 10 2. A compound according to claim 1, wherein R^1 is pyridyl, optionally substituted with lower alkyl or lower alkenyl.
3. A compound according to any of claims 1-2, wherein R^1 is pyridyl, optionally substituted with lower alkyl.
4. A compound of claim 1, selected from the group consisting of:
- 15 5-methyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide,
- 5-isopropyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide,
- 5-isopropenyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide and
- 20 5-methyl-thiazole-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide.

5. A compound of any of claims 1-2, selected from the group consisting of:
- 5-methyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide,
- 5-isopropyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide and
- 5-isopropenyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide.
6. A compound of any of claims 1-3, selected from the group consisting of:
- 5-methyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide and
- 5-isopropyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide.
7. The compound of any of claims 1-3, which is:
- 5-methyl-pyridine-2-sulfonic acid [6-methoxy-5-(2-methoxy-phenoxy)-2-pyridin-4-yl-pyrimidin-4-yl]-amide.
8. A pharmaceutical composition comprising a compound of any one of claims 1-7 and a pharmaceutically acceptable carrier and/or adjuvant.
9. A pharmaceutical composition according to claim 8, additionally comprising any active compounds selected from antihypertensive drugs, antiarrhythmics, anti angina, antithrombotic and lipid lowering agents as well as antioxidants.
10. The use of a compound as defined in any of claims 1-7 for the preparation of medicaments containing a compound as defined in any of claims 1-7 for the treatment or prophylaxis of disorders which are associated with abnormal vascular tone and endothelial dysfunction.
11. A use of a compound as defined in any of claims 1-7 for the treatment or prophylaxis of disorders which are associated with abnormal vascular tone and endothelial dysfunction.

12. A process for the preparation of the compounds described in any one of claims 1-7, which process comprises:

- a) reacting 4,6-dichloro-5-(2-methoxy-phenoxy)-2-(pyridin-4-yl)-pyrimidine with an appropriate sulfonamide of the general formula (II)



5

wherein R^1 is as defined in claim 1,

in a suitable solvent in the presence of a suited base, followed by further treatment with sodium methylate in methanol, or

- b) reacting 4-[4,6-dichloro-5-(2-methoxy-phenoxy)-pyrimidin-2-yl]-pyridine-1-oxide with an appropriate sulfonamide of the general formula (II)

10




wherein R^1 is as defined in claim 1,

in a suitable solvent in the presence of a suited base, followed by further treatment with sodium methylate in methanol and pyridine-N-oxide reduction.

15


13. The compounds of formula (III)



(III)


wherein R^1 is as defined in claim 1.

14. The compounds of formula (V)



wherein R^1 is as defined in claim 1.

15. Compounds according to any of claims 1-7, whenever prepared by the process of
5 claim 12.
16. Compounds according to any of claims 1-7 for the treatment of diseases which are associated with abnormal vascular tone and endothelial dysfunction.



(I)