

(12) DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2004/02/27

(87) Date publication PCT/PCT Publication Date: 2004/09/10

(85) Entrée phase nationale/National Entry: 2005/08/12

(86) N° demande PCT/PCT Application No.: JP 2004/002438

(87) N° publication PCT/PCT Publication No.: 2004/076478

(30) **Priorité/Priority:** 2003/02/28 (2003-052926) JP

(51) Cl.Int.⁷/Int.Cl.⁷ C07K 5/08, A61K 38/55, A61P 25/28, A61P 25/16, A61P 43/00, A61P 25/00, C07K 7/02, C07K 5/02, C07K 7/06



(71) Demandeur/Applicant:
KISO, YOSHIKAKI, JP

(72) Inventeur/Inventor:
KISO, YOSHIKI, JP

(74) **Agent:** KIRBY EADES GALE BAKER

(54) Titre : DERIVES PEPTIDIQUES A ACTIVITE INHIBITRICE DE LA β -SECRETASE

(54) Title: PEPTIDE DERIVATIVES HAVING β -SECRETASE INHIBITORY ACTIVITY




(57) Abrégé/Abstract:


Provided are excellent β -secretase inhibitors, that is, compounds represented by the general formula (1) or prodrugs thereof: (1) wherein A is amino or protected amino; B¹ and B² are each a single bond, alkylene of 1 to 3 carbon atoms, or the like; D is a single bond, -NHCO-, or the like; E is -COOH, a tetrazole ring, or the like; n is an integer of 1 to 3; m is an integer of 1 to 3; G is hydroxyl, a group represented by the general formula (2), or the like: (2) [wherein Z is -NH-, -Asp-Ala-NH-, -Asp-Ala-, -Asp-NH-, or the like; L is a 5- to 10-membered ring optionally containing a heteroatom and/or an unsaturated bond; X is -COOH, a tetrazole ring, or the like; Y is hydrogen, -COOH, or the like; and k is an integer of 1 to 4]; and R¹, R², and R³ are each alkyl of 1 to 6 carbon atoms, or the like.

ABSTRACT

Provided are excellent β -secretase inhibitors, that is, compounds represented by the general formula (1) or prodrugs thereof:



wherein A is amino or protected amino; B¹ and B² are each a single bond, alkylene of 1 to 3 carbon atoms, or the like; D is a single bond, -NHCO, or the like; E is -COOH, tetrazole ring, or the like; n is an integer of 1 to 3; m is an integer of 1 to 3; G is hydroxyl, a group represented by the general formula (2), or the like:



in which Z is -NH, -Asp-Ala-NH-, -Asp-Ala-, -Asp-NH-, or the like; L is a 5-to 10-membered ring optionally containing a heteroatom and/or unsaturated bond; X is -COOH, tetrazole ring, or the like; Y is hydrogen, -COOH, or the like; and k is an integer of 1 to 4; and R¹, R², and R³ are each alkyl of 1 to 6 carbon atoms, or the like.

SPECIFICATION

PEPTIDE DERIVATIVES HAVING β -SECRETASE INHIBITORY ACTIVITY

5 TECHNICAL FIELD

The present invention relates to a peptide derivative having excellent β -secretase inhibitory activity.

BACKGROUND ART

10 Alzheimer's disease (abbreviated as AD) is a neurodegenerative disease characterized by dementia. AD insidiously develops in old age and gradually progresses to dementia. AD is pathologically characterized by large amount of senile plaques and neurofibrillary tangles (abbreviated as NFT) observed in the cerebral cortex and hippocampus, and by cerebral
15 atrophy due to decudation of neuron.

Senile plaque has, as a main component, an amyloid fiber comprising of an extracellularly sedimented amyloid β -peptide (abbreviated as A β). The appearance of AB is extremely specific for AD and aging. Senile plaque is a pathological change which is seen at the earliest stage of
20 the disease's progression. Therefore, an understanding of the mechanism of formation and sedimentation of A β has a very important significance in elaborating the morbid state of AD.

A β includes, two main molecular species; A β 40 that ends at the 40th amino acid residue Val; and A β 42 which is larger by two residues (Ile, Ala). Among the A β secreted from cells, about 90% is A β 40, and 10% is
25 A β 42. However, A β 42 exhibits aggregating property which is far greater than that of A β 40. Aggregated A β 42 entangles with A β 40 to form amyloids. In addition, because A β 40 accumulated during the initial stage of plaque formation, it has been suggested that the accumulation of A β 40
30 was strongly associated with the development of AD.

A β is produced by cleavage of an amyloid precursor protein (abbreviated as APP) by two kinds of proteases. APP is a type I transmembrane protein consisting of about 700 residues, with A β contained in its extracellular region at a transmembrane region. A protease that is involved in the production of A β from APP is called "secretase". A secretase that cuts (cleaves) the amino terminal of A β is a " β secretase"; a secretase that cuts the carboxyl terminal is a " γ secretase"; and a secretase that cuts the carboxyl terminal of the 16th Lys in the interior of A β is an " α secretase". In an amyloidogenic pathway, APP is cut with a β secretase to produce a secretion-type APP β (sAPP β) and a carboxyl terminal fragment called β -stub. A γ secretase cuts this fragment to produce A β and a γ -stub. In a non-amyloidogenic pathway, an α secretase cuts APP to produce secretion-type APP α (sAPP α) and a carboxyl terminal fragment called α -stub. A γ secretase cuts this fragment to produce p3 and γ -stub. These two pathways both exist in the normal state. The majority of A β that is produced as a physiological molecule is secreted extracellularly.

One of the current mainstream strategies is to target secretase as a point of action for preventing or treating AD. Among them, particular attentions are paid for β and γ secretases since direct effects can be readily observed. In recent years, pharmaceutical enterprises and venture companies vigorously progress in the development of secretase inhibitors.

In 1999, it has been reported by some laboratories that β secretase is a β -site APP cleaving enzyme called BACE1 (also referred to as BACE, Asp2, or memapsin2).

Since BACE1 is an aspartic acid protease, based on the substrate transition state concept established by the study on the same class of protease inhibitor such as renin and HIV, various β secretase inhibitors have been reported in recent years. Swedish variant-type APP readily produces A β . A peptide-based inhibitor mimics Swedish APP by forming a

substrate transition state analogue at a β secretase cleaving site. Tung et al. determined the IC_{50} value of 30nM in a 14 residue peptide (P10-P4' statV) in which a hydroxyethylcarbonylisostere (statine) structure was introduced into a substrate cleaving site of β secretase. They reported that other substrate transition state analogues (AHPPA, ACHPA) can also inhibit β secretase (for example, see Nature, 402, 537-540 (1999), J. Med. Chem., 45, 259-262 (2002)). Ghosh et al. searched various compounds containing hydroxyethyleneisostere, and discovered that OM 99-2 had strong inhibitory activity ($K_i=1.6nM$) (for example, see WO 01/00665, J. Am. Chem. Soc. 122, 3522-3523 (2000). Based on X-ray crystal structural analysis of an enzyme-OM99-2 complex (for example, see Science 290, 150-153 (2000)), they synthesized derivatives, and found a smaller inhibitor having equipotent activity (for example, see J. Med. Chem., 44, 2865-2868 (2001)).

However, in considering an acting site of a β -secretase inhibitor, a drug needs to pass through the blood-brain barrier. There are many points to be considered such as a molecular size and hydrophilic/hydrophobic balance. In addition, since substrate specificity greatly differs from previous aspartic acid proteases (for example, see Biochemistry, 40, 10001-10006 (2001)), it is expected that the design of inhibitors is very difficult. Furthermore, in order to be physiologically active, it is necessary for the inhibitor to be able to pass through the blood-brain barrier. For this reason, compounds in which the molecular weight is relatively small, and stability in blood is excellent are desired. Designing compounds with such features are thought to be difficult.

DISCLOSURE OF THE INVENTION


The development of a compound that has excellent β -secretase inhibitory activity and is satisfactory as a medicament is solicited.

The present inventors synthesized an octapeptide in which an HMC

isostere was introduced into a β -secretase cleaving site of Sweden variant APP, and optimized each site of the octapeptide. Thereafter, they examined the minimum molecular size at which BACE1 inhibitory activity could be maintained, studied correlations between structure and activity, and derived the present compound that had excellent activity. Thus the present invention was completed.

BEST MODE FOR CARRYING OUT THE INVENTION

That is, the present invention relates to a compound represented by the general formula (1):



wherein A is amino group or amino group protected with a protecting group which can be degraded in a living body,

B¹ and B² are each a single bond or optionally substituted alkylene of 1 to 3 carbon atoms optionally substituted with R⁴,

in which R⁴ is halogen atom; alkyl of 1 to 6 carbon atoms; alkenyl of 2 to 6 carbon atoms; alkynyl of 2 to 6 carbon atoms; cycloalkyl of 3 to 7 carbon atoms; optionally substituted heterocycle; optionally substituted aralkyl; or optionally substituted aryl, and the substituent thereof is alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or aryl,


D is a single bond, -NHCO- or NHCO-R⁵-,

in which R⁵ is alkylene of 1 to 6 carbon atoms; alkenylene of 2 to 6 carbon atoms; alkynylene of 2 to 6 carbon atoms; cycloalkylene of 3 to 7 carbon atoms; optionally substituted heterocycle; optionally substituted aralkylene; or optionally substituted arylene, and the substituent thereof is halogen atom, alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms,

m is an integer of 1 to 3,

$$-Z-\textcircled{L}-X-(M)_K$$

X is hydrogen atom, -COOH, a group equivalent to COOH, -CONHR⁶, -OCH₂COOR⁶, -OCH₂CONHR⁶, hydroxyl group, -PO₄H, -SO₃H, -CZ¹(Z²)-X', or a group of the formula:



k is an integer of 1 to 4, and,

R¹, R² and R³ are each -COOH, a group equivalent to COOH, alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or alkyl of 1 to 6 carbon

atoms substituted with any of phenyl, phenylthio, alkylthio of 1 to 6 carbon atoms, COOH or heterocycle, provided that the phenyl, the phenylthio and the heterocycle may be substituted with alkyl of 1 to 6 carbon atoms, hydroxyl group, nitro, halogen atom, -SO₃H or -PO₄H or a pharmaceutically acceptable salt thereof, or a prodrug thereof.

The present invention relates to a compound in which, in the general formula (1),

B¹ and B² are each a single bond or alkylene of 1 to 3 carbon atoms optionally substituted with R⁴,

in which R⁴ is halogen atom; alkyl of 1 to 6 carbon atoms; alkenyl of 2 to 6 carbon atoms; alkynyl of 2 to 6 carbon atoms; cycloalkyl of 3 to 7 carbon atoms; optionally substituted heterocycle; optionally substituted aralkyl; or optionally substituted aryl, and the substituent thereof is alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or aryl,

D is a single bond, -NHCO- or -NHCO-R⁵-,


in which R⁵ is alkylene of 1 to 6 carbon atoms; arkenylene of 2 to 6 carbon atoms; alkynylene of 2 to 6 carbon atoms; cycloalkylene of 3 to 7 carbon atoms; optionally substituted heterocycle; optionally substituted aralkylene; or optionally substituted arylene, and the substituent thereof is halogen atom, alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or aryl,

E is -COOH or a group equivalent to COOH, hydrogen atom, hydroxyl group, amino group, halogen atom, or -COOCH₃,

n is an integer of 1 to 3,

m is 1,

G is hydroxyl group, or a group of the formula:




in which Z is -NH-, -Asp-Ala-NH-, -Asp-Ala-, -NH-CZ¹(Z²)-, -Asp-

NH-, -Asp-NH-CZ¹(Z²)-, or -NH-CZ¹(Z²)-CZ¹(Z²),

Ring L is a 5- to 10-membered ring optionally containing a heteroatom and/or unsaturated bond,

Z¹ and Z² are each a hydrogen atom, hydroxyl group, amino group,
5 cyano group, nitro group, alkyl of 1 to 6 carbon atoms, or halogen atom,

X is hydrogen atom, -COOH, a group equivalent to COOH,
-CONHR⁶, -OCH₂COOR⁶, -OCH₂CONHR⁶, hydroxyl group, -PO₄H, -SO₃H,
-CZ¹(Z²)-X', or a group of the formula:



10 in which X' is hydrogen atom, -COOH, a group equivalent to COOH,
-CONHR⁶, -OCH₂COOR⁶, -OCH₂CONHR⁶, hydroxyl group, -PO₄H, or -SO₃H,

Y and Y' are each hydrogen atom, -COOH, a group equivalent to
COOH, -CONHR⁶, -OR⁶, -PO₄H, -SO₃H, halogen atom, or methyl,

R⁶ is alkyl of 1 to 4 carbon atoms optionally substituted with
15 halogen atom, or hydrogen atom,

k is an integer of 1 to 4, and


R¹, R² and R³ are each alkyl of 1 to 6 carbon atoms, alkenyl of 2 to
6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon
atoms, or alkyl of 1 to 6 carbon atoms substituted with any of phenyl,
20 phenylthio, alkylthio of 1 to 6 carbon atoms, or heterocycle, provided that
the phenyl, phenylthio and heterocycle may be substituted with alkyl of 1
to 6 carbon atoms, hydroxyl group, nitro, halogen atom, -SO₃H or -PO₄H,
or a pharmaceutically acceptable salt thereof, or a prodrug thereof.

The present invention relates to a compound in which, in the
25 general formula (1),

B¹ is a single bond or methylene,


B² is a single bond or alkylene of 1 to 3 carbon atoms,

D is a single bond, -NHCO-, a group of the formula:




in which the benzene ring is optionally substituted with halogen atom, amino group, alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms or aryl,

a group of the following formula:




or a group of the following formula:



in which Q is oxygen atom or NH,
E is hydrogen atom, hydroxyl group, amino group, halogen atom, -COOH, -COOCH₃, or tetrazole ring,
n is an integer of 1 to 2,
m is 1,

R¹ is alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or aryl,
R² is alkyl of 1 to 6 carbon atoms, phenyl, naphthyl, isobutyl, 2-carbamoyl ethyl, 2-methylthioethyl, methylthiomethyl, 2-ethylthioethyl, ethylthiomethyl, naphthylmethyl, phenethyl, 4-thiazoylmethyl, 4-imidazolylmethyl, 3-tetrahydrofuranyl, 3-indolylmethyl, cyclohexylmethyl, or benzyl in which the benzene ring may be substituted with halogen atom, hydroxyl group, or alkyloxy group of 1 to 6 carbon atoms, and

G is any one of the group represented by the following formulas:



in which X and X' are each hydrogen atom, -COOH, -CONHR⁶, -OCH₂COOR⁶, -OCH₂CONHR⁶, hydroxy group, -PO₄H, -SO₃H or tetrazole ring,

5 Y and Y' are each hydrogen atom, -COOH, -CONHR⁶, -OR⁶, -PO₄H, -SO₃H, halogen atom, methyl or terazole ring,

R⁶ is alkyl of 1 or 2 carbon atoms optionally substituted with halogen atom, or hydrogen atom,

k is an integer of 1 or 2, and


10 Z¹ and Z² are each hydrogen atom, hydroxyl group, amino group, cyano group, nitro group, alkyl of 1 to 6 carbon atoms, or halogen atom, or a pharmaceutically acceptable salt thereof, or a prodrug thereof.

The present invention relates to a compound in which, in the general formula (1),

15 B¹ is a single bond,

B² is alkylene of 1 or 2 carbon atoms,

D is a single bond, -NHCO-, or a group of the following formula:



E is hydrogen atom, hydroxyl group, halogen atom, —COOH or tetrazole ring,

n is an integer of 1 or 2,


m is 1,

R^1 is isopropyl,

R^2 is isobutyl or cyclohexylmethyl,

R^3 is benzyl, isobutyl, phenylthiomethyl or cyclohexylmethyl, and

G is any one of the group represented by the following formulas:



in which X and X' are each COOH , CONH_2 , hydroxyl group, OCH_2COOH or tetrazole ring,

Y and Y' are each hydrogen atom, COOH , CONH_2 , halogen atom, methyl or a tetrazole ring,


k is an integer of 1 or 2, and

Z¹ and Z² are each hydrogen atom,

or a pharmaceutically acceptable salt thereof, or a prodrug thereof.


The present invention relates to a compound in which, in the
 5 general formula (1),

a partial structure other than G is a group of the formula:




in which R² is isobutyl or cyclohexylmethyl,

R⁷ is a group of the following formula:



and

G is the following formula:



or a pharmaceutically acceptable salt thereof, or a prodrug thereof.

It is preferred that the molecular weight of the compound represented by the general formula (1) is 1100 or less.

5 The present invention also relates to a β -secretase inhibitor containing the above compound as an active ingredient.

The present invention relates to an agent for preventing or treating (i) neurodegenerative disease, (ii) neuropathy in cerebrovascular disease, head trauma·spinal damage, encephalitis sequela, or cerebral palsy, (iii) 10 memory disorder, or (iv) mental disease associated with β -secretase containing the above compound as an active ingredient.

The present invention relates to an agent for preventing or treating (i) neurodegenerative disease, (ii) neuropathy in cerebrovascular disease, head trauma·spinal damage, encephalitis sequela, or cerebral palsy, (iii) 15 memory disorder or (iv) mental disease by promoting secretion of sAPP α , or promoting secretion of sAPP α , and inhibiting production·secretion of a β amyloid protein, or inhibiting production·secretion of a β amyloid protein, containing the aforementioned compound as an active ingredient.

20 The present invention relates to an agent for treating the neurodegenerated disease, Alzheimer's disease or Parkinson's disease,

containing the aforementioned compound as an active ingredient.

The present invention relates to an agent for promoting sAPP α secretion containing the aforementioned compound as an active ingredient.

The present invention relates to an agent for inhibiting production
5 secretion of a β amyloid protein.

The present invention relates to a neurotrophic factor-like acting agent containing the aforementioned compound as an active ingredient.

The present invention relates to an agent for preventing or treating neuropathy or mental disease in head trauma·spinal damage, encephalitis
10 sequela, or cerebral palsy containing the aforementioned compound as an active ingredient.

The present invention relates to the use of a β -secretase inhibitor of the aforementioned compound in the preparation of a sAPP α secretion promoter, a neurotrophic factor-like acting agent, or an agent for
15 preventing or treating neuropathy, memory disorder or mental disease head trauma·spinal damage, encephalitis sequela, or cerebral palsy.

The present invention also relates to a method for treating (i) neurogenerative disease, (ii) neuropathy in cerebrovascular disease, head
20 trauma·spinal damage, encephalitis sequela, or cerebral palsy, (iii) memory disorder, or (iv) mental disease associated with β -secretase, which comprises administering an effective amount of a β -secretase inhibitor containing the aforementioned compound as an active ingredient to a mammal.

As used herein,

25 the “alkyl of 1 to 6 carbon atoms” include straight or branched alkyl of 1 to 6 carbon atoms, specifically, methyl, ethyl, n-propyl, 2-propyl, n-butyl, 2-butyl, isobutyl, 3-methyl-2-propyl, 1,1-dimethylethyl, n-pentyl, and n-hexyl, preferably, methyl, ethyl, n-propyl, and 2-propyl.

The “alkenyl of 2 to 6 carbon atoms” includes vinyl, 2-propenyl, 1-
30 propenyl, 2-butenyl, 1-butenyl, and 2-methyl-1-propenyl.

The “alkynyl of 2 to 6 carbon atoms” includes ethynyl, 2-pronynyl, 2-bunynyl, 3-butynyl, 2-pentynyl, 3-pentynyl, and 4-pentynyl.

The “cycloalkyl of 3 to 7 carbon atoms” includes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cyclopentyl.

5 The “aryl” includes phenyl, and naphthyl, preferably phenyl.

The “heterocycle” includes a heterocycle having an unsaturated bond such as an aromatic heterocycle and a saturated heterocycle. Examples of the aromatic heterocycle include a 5- to 10- membered aromatic ring containing 1 to 3 heteroatoms such as pyridyl, pyrimidyl, 10 pyridazyl, thienyl, furyl, pyrrolyl, imidazolyl, oxazolyl, isooxazolyl, thiazolyl, isothiazolyl, triazolyl, indolyl, benzothiazolyl, quinazolyl, and isoquinazolyl, preferably pyridyl, pyrimidyl, thienyl, pyrrolyl, indolyl, benzothiazolyl, and isoquinazolyl, particularly preferably pyridyl, pyrimidyl, and thienyl. Examples of the saturated heterocycle include piperidinyl, pyrrolydyl, 15 piperazinyl, imidazolyl, morpholinyl, thiomorpholinyl, and tetrafuryl.

The “Ring L” includes a cycloalkyl ring such as cyclohexyl and cyclopentyl; an aromatic ring such as phenyl and naphthyl, an unsaturated heterocycle such as aromatic heterocycle; and a saturated heterocycle. Examples of the aromatic heterocycle include pyridyl, 20 pyrimidyl, pyridazyl, thienyl, furyl, pyrrolyl, imidazolyl, oxazolyl, isooxazolyl, thiazolyl, isothiazolyl, triazolyl, indolyl, benzothiazolyl, quinazolyl, and isoquinazolyl, preferably pyridyl, thienyl, pyrrolyl, indolyl, and benzothiazolyl, particularly preferably pyridyl, and thienyl. Examples of the saturated heterocycle include piperidinyl, pyrrolydyl, piperazinyl, 25 imidazolyl, morpholinyl, thiomorpholinyl, and tetrafuryl.

The “aralkyl” represents alkyl of 1 to 6 carbon atoms substituted with aryl, specifically, benzyl, phenethyl, phenylpropyl, phenylbutyl, and naphthylmethyl, preferably benzyl, and phenethyl.

30 The “a group equivalent to COOH” includes hydroxamic acid, acyl-cyanamide, tetrazole, oxooxadiazole, oxothiadiazole, thioxooxadiazole,

oxooxathiadiazole, mercaptoazole, sulfinylazole, sulfonylazole, isooxazole, isothiazole, hydroxythiadiazole, hydroxyl- γ -pyrone, phosphinic acid, phosphonic acid, phosphonamide, sulfonamide, and acylsulfonamide, preferably tetrazole.

5 The "prodrug" refers to a compound which is hydrolyzed in a living body, and reproduces the compound of the present invention. The prodrug of the present invention includes compounds which are prepared by any procedure for converting into a prodrug known to a person skilled in the art. For example, when the compound of the present invention has
10 a carboxyl group or an amino group, compounds in which those groups are derived into an ester group or an amide group easily hydrolyzable in a living body correspond to a prodrug. When a compound has a carboxyl group, examples of the prodrug include compounds in which the carboxyl group is introduced into alkyl such as methyl, and ethyl, alkyloxyalkyl such as
15 methyloxymethyl, ethyloxymethyl, 2-methyloxyethyl, and 2-methyloxyethyloxymethyl, acyloxymethyl such as pivaloyloxymethyl, acetyloxymethyl, cyclohexylacetyloxymethyl, and 1-methylcyclohexylcarbonyloxymethyl, alkoxycarbonylalkyl such as ethyloxycarbonyloxy-1-ethyl, or cycloalkyloxycarbonylalkyl such as
20 cyclohexyloxycarbonyloxy-1-ethyl. When a compound has an amino group, examples of a prodrug include compounds in which an amino group is introduced into acetamide.


 The compound (1) which is an active ingredient of a medicament of the present invention can be formed in a pharmaceutically acceptable salt
25 thereof. Examples of the pharmaceutically acceptable salt include an acid addition salt and a base addition salt. Examples of the acid addition salt include inorganic acid salts such as hydrochloride, hydrobromide, sulfate, etc. and organic acid salts such as citrate, oxalate, malate, tartarate, fumarate, maleate, etc. and examples of a base addition salt include
30 inorganic base salts such as a sodium salt, a calcium salt, etc. and organic

base salts such as a meglumine salt, a trishydroxymethylaminomethane salt, etc. In addition, the present invention includes a hydrate and a solvate of the compound (1) or a pharmaceutically acceptable salt thereof.


The compound of the present invention or a pharmaceutically acceptable salt thereof can be prepared by the following method or a similar method.

Since the compound of the present invention has a peptidic linkage(s), each amino acid unit can be prepared by the method which is conventionally used in peptide chemistry, for example, the method described in "The Peptides", vol., by Schroder and Luhker, Academic Press, New York, U.S.A (1966), and "Base and Experiment of Peptide Synthesis" by Nobuo Izumiya, et al., Maruzen (1985), and the compound of the present invention, for example, can be prepared using a liquid phase method shown by the following scheme and a solid phase method shown by the following scheme II.

Scheme I (liquid phase method)



Scheme II (Solid phase method)



Further, it can be also prepared by the column or batch method.

The formation of a peptidic linkage can be exemplified by the azide,

acid chloride, acid anhydride, carbodiimide, carbodiimide-active, active ester, carbonylimidazole, redox, or enzyme method, and a method using Woodward's reagent K. Examples of a condensation reaction in a solid phase method include, as a main method, the acid anhydride, carbodiimide, or active ester method among the aforementioned methods.

Furthermore, when a peptide chain is extended by the solid phase method, the C-terminal amino acid is bound to a support such as a resin which is insoluble in the organic solvent used. Herein, a resin in which a functional group is introduced for the purpose of binding an amino acid to a resin, an entity in which a spacer is inserted between a resin and a functional group, and a resin in which a chain referred to as a handle, which is cut at various places depending on the conditions, can also be used depending on the purpose. Examples of such a resin include Wong resin, 2-chlorotriyl resin, halomethyl resin such as chloromethyl resin, oxymethyl resin, and C-terminal amidated resin.

Furthermore, before these condensation reactions are performed, a carboxyl group, an amino group, a hydroxyl group, or an amidino group not involved in the condensation reaction can be protected by conventional means. Conversely, a carboxyl group and an amino group directly involved in the condensation reaction can be activated.

To protect a functional group not involved in the condensation reaction, the functional group can be protected with a protecting group which is usually used in organic chemistry, or a protecting group described, for example, in "Protective Groups in Organic Synthesis", by Green, John Wiley & Sons, Inc. (1981).

A condensation reaction used in peptide synthesis is usually performed in a solvent. Examples of solvents include chloroform, dichloromethane, ethyl acetate, N,N-dimethylformamide, dimethyl sulfoxide, pyridine, dioxane, tetrahydrofuran, N-methylpyrrolidone, water, methanol, and a mixture thereof. The condensation reaction is carried out

at a range of -30 to 50°C as in conventional cases.

Furthermore, the elimination of a protecting group in the preparation of the present invention can be selected according to the kind of protecting group used as far as the protecting group can be eliminated without influencing on peptidic linkage. Examples include acid treatment with hydrogen chloride, hydrogen bromide, anhydrous hydrogen fluoride, methanesulfonic acid, trifluoromethanesulfonic acid, trifluoroacetic acid, or a mixture thereof, alkali treatment with sodium hydroxide, potassium hydroxide, hydrazine, diethylamine, or piperidine, sodium treatment in liquid ammonia, reduction with palladium carbon, and treatment with a silylating agent such as trimethylsilyl triflate, trimethylsilyl bromide, etc. In a deprotecting reaction by the treatment with the acid or the silylating agent mentioned above, it is preferable to add a cation scavenger such as anisole, phenol, cresol, anisole, thioanisole, ethanedithiol, etc. to efficiently perform the deprotecting reaction. A method for cleaving a peptide of the present invention synthesized by the solid phase method, from a solid phase is also performed according to conventional methods.

The prepared peptide of the present invention can be subjected to conventional separation and purification after completion of the aforementioned series of reactions. For example, by extraction, distribution, re-precipitation, recrystallization, column chromatography, etc., the peptide of the present invention can be obtained in a purer form.

The compound of the present invention or a pharmaceutically acceptable salt thereof has asymmetry in some cases, or has a substituent having an asymmetric carbon in some cases, such that there is an optical isomer. The compound of the present invention includes a mixture of the respective or isolated isomers. Examples of a method for obtaining such optical isomer pure include optical resolution.

As an optical resolution, the compound of the present invention or an intermediate thereof may form a salt with an optically active acid (e.g.

monocarboxylic acids such as mandelic acid, N-benzyloxyalanine, lactic acid, etc., dicarboxylic acids such as tartaric acid, o-diisopropylidenetartaric acid, malic acid, etc., sulfonic acids such as camphorsulfonic acid, bromocamphorsulfonic acid, etc.) in an inert solvent (e.g. alcohol solvents such as methanol, ethanol, and 2-propanol, ether solvents such as diethyl ether, ester solvents such as ethyl acetate, aromatic hydrocarbon solvents such as toluene, acetonitrile, and a mixed solvent thereof).

In addition, when the compound of the present invention or an intermediate thereof has an acidic substituent such as a carboxyl group, it may be formed as a salt with optically active amines (e.g. organic amines such as α -phenethylamine, quinine, quinidine, cinchonidine, cinchonine, strychnine, etc.).

Examples of the temperature used to form a salt include a range from room temperature to the boiling point of the solvent. In order to improve optical purity, it is desirable to raise the temperature to near boiling point of the solvent only once. Chemical yield can be improved by cooling before filtering of the precipitated salt, if necessary. It is suitable that an amount of an optically active acid or amine to be used in a range of 0.5 to 2.0 molar equivalents, preferably in a range of around 1 molar equivalent relative to that of the substrate. If necessary, a crystal may be recrystallized in an inert solvent (e.g. alcohol solvents such as methanol, ethanol, and 2-propanol, ether solvents such as diethyl ether, ester solvents such as ethyl acetate, aromatic hydrocarbon solvents such as toluene, and acetonitrile, and a mixed solvent thereof) to obtain an optically active salt having a high purity. If necessary, the resulting salt may be treated with an acid or a base by conventional methods to obtain a free compound.

The treating agent of the present invention can be administered orally or parenterally. When orally administered, the agent can be

administered in an administration form which is conventionally used.

When administered parenterally, the agent can be administered in a form of an agent for topical application, injection, transdermal agent, or nasal agent.

5 The above dosage form is formulated into a preparation with a pharmaceutically acceptable excipient or additive by conventional methods. Examples of a pharmaceutically acceptable excipient or additive include a carrier, binder, flavor, buffer, thickener, colorant, stabilizer, emulsifier, dispersant, suspending agent, and antiseptic.

10 Examples of a pharmaceutically acceptable carrier include magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, low melting point wax, and cacao butter. A capsule can be formulated by putting the present compound together with
15 a pharmaceutically acceptable carrier therein. The treating agent of the present invention can be added by mixing with a pharmaceutically acceptable excipient, or without the excipient into a capsule. A cachet can be prepared by similar methods.

 Examples of a liquid preparation for injection include a solution,
20 suspension, emulsion, etc. such as an aqueous solution, or water-propylene glycol solution. The liquid preparation may also be prepared in the form of a solution of polyethylene glycol or/and propylene glycol, which may contain water. A liquid preparation suitable for oral administration can be prepared by adding water to the present compound and, if
25 necessary, adding a colorant, flavor, stabilizer, sweetener, solubilizer, and thickener. A liquid preparation suitable for oral preparation may be prepared by adding the present compound or a pharmaceutically acceptable salt thereof together with a dispersant to water to increase viscosity. Examples of a thickener include pharmaceutically acceptable
30 natural or synthetic gum, resin, methylcellulose, sodium

carboxymethylcellulose or known suspending agent, etc.

Examples of an agent for topical application include the
aforementioned liquid preparation, and a cream, aerosol, spray, powder,
lotion, and ointment. The above agent for topical application can be
5 prepared by mixing the present compound or a pharmaceutically
acceptable salt thereof with a pharmaceutically acceptable diluent and
carrier that are conventionally used. An ointment or cream is obtained,
for example, by formulating into a preparation by adding a thickener
and/or a gelling agent to an aqueous or oily base. Examples of a base
10 include water, liquid paraffin, and vegetable oil (peanut oil, castor oil).
Examples of a thickener include soft paraffin, aluminum stearate,
cetostearyl alcohol, propylene glycol, polyethylene glycol, lanolin,
hydrogenated lanolin, beeswax, etc.

A lotion can be prepared by adding one or more kinds of
15 pharmaceutically acceptable stabilizer, suspending agent, emulsifier,
diffusing agent, thickener, colorant, and flavor to an aqueous or oily base.

A powder is formulated into a preparation with a base for a
pharmaceutically acceptable powder. Examples of a powder include talc,
lactose, starch, etc. Drops can be formulated into a preparation from an
20 aqueous or non-aqueous base and one or more kinds of pharmaceutically
acceptable diffusing agent, suspending agent, and solubilizer.

The agent for topical application may contain an antiseptic such as
methyl hydroxybenzoate, propyl hydroxybenzoate, chlorocresol, and
benzalkonium chloride, or a bacteriostat.

25 A preparation of a liquid spray, a powder or a drop containing the
present compound or a pharmaceutically salt thereof as an active
ingredient can be administered nasally.

The dose and administration interval are different depending on
symptoms, age, weight, and administration form and, when orally
30 administered, usually a range of about 1 to about 500mg, preferably a

range of about 5 to about 100mg per adult per day which can be administered once or in divided doses. When administered as an injection, a range of about 0.1 to about 300mg, preferably a range of about 1 to about 100mg can be administered once or in divided doses.

5 A process for preparing the present compound will be explained below by Examples.

When the present reaction is performed, the technique of protection and deprotection can be used, if necessary. The technique of protection and deprotection is described in detail in the aforementioned Protective
10 Groups in Organic Synthesis [authored by Green, John Wiley & Sons, Inc. (1981)].

Examples

The meanings of abbreviations used in Examples are as follows:

15 Me: methyl group

Boc: tert-butoxycarbonyl group

DIPCDI: diisopropylcarbodiimide

Fmoc: 9-fluorenylmethoxycarbonyl group

HOBt: N-hydroxybenzotriazole

20 DMF: N,N'-dimethylformamide

Apns: amino acid residue of (2S,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid

DMAP: 4-dimethylaminopyridine

Dap: amino acid residue of L- α,β -diaminopropanoic acid

25 Pns: amino acid residue of (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid

Nst: amino acid residue of (2R,3S)-3-amino-2-hydroxy-5-methylhexanoic acid

Dmt: (R)-5,5-dimethyl-1,3-thiazolidine-4-carboxylic acid

Reference Examples

Upon synthesis of the present compound, first, Fmoc-Nst-OH (compound 6) which was a fundamental block, was synthesized according to the method described in the literatures. Namely, using Z-Leo-OH as a raw material, TMSCN was reacted on N- α Z protected aminoaldehyde derived by conventional methods, and the resulting cyanohydrin was introduced into an α -hydroxy- β -amino acid derivative by acid hydrolysis. After being separated and purified into each isomer by column chromatography, the objective compound 6 was obtained via deprotection and 9-fluorenylmethoxycarbonylation. Fmoc-Pns-OH (compound 7) and Fmoc-Apns were synthesized separately as an HMC isostere similar to Nst. Using H-Pns-OH and H-Apns-OH as starting raw materials, 9-fluorenylmethoxycarbonylation was elaborated using the same manner as that described above to obtain the objective product.

Reference Example 1

Introduction of amino acid into 2-chlorotrityl chloride resin (Clt resin)

A Clt resin, Fmoc amino acid derivative (2.5 eq), DCE (2mL) and DIEA (2.5 eq) were successively added to a reactor under Ar atmosphere, and these were reacted for 2 hours. After completion of the reaction, MeOH (200 μ L) and DIEA (0.5 eq) were added and, the mixture was shaken for 20 minutes. Thereafter, the mixture was thoroughly washed with DMF, CHCl₃ and MeOH, and the resin was dried under vacuum. Using the dry resin, a substitution rate was calculated.

Reference Example 2

Introduction of amino acid into Wang resin

Wang resin was added to a reactor, and the resin was adopted with DMF. Then, an Fmoc amino acid derivative (3 eq), DMF (2mL), DIPCDI (3 eq) and DMAP (0.1 eq) were added to react for 90 minutes. After the reaction, the reaction was thoroughly washed with DMF, CHCl₃ and MeOH, and dried under vacuum. Using the dry resin, a substitution rate was

obtained.

Reference Example 3

Removal of N α -amino group

5 After a protected peptide resin in a reactor was manually shaken with 20% piperidine/DMF for 1 minute, the reaction solution was filtered, 20% piperidine/DMF was newly added thereto, followed by shaking for 20 minutes. After the reaction, the resin was sufficiently washed with DMF and CHCl₃ until the filtrate became neutral.

Reference Example 4

10 Condensation reaction of amino acid

An Fmoc amino acid derivative (2.5 eq), and HOBt·H₂O (2.5 eq) were added to a protected peptide resin in a reactor, followed by dissolution with DMF. Then, DIPCDI (2.5 eq) was added, and the mixture was shaken for 90 minutes. After completion of the reaction, the resin
15 was thoroughly washed with DMF, a part of the resin was taken, and the progress of the reaction was confirmed by Kaiser test. When it was determined that the reaction was not completed, the condensation reaction was repeated.

Reference Example 5

20 Removal of resin and deprotection

A protected peptide resin was reacted with m-cresol (2.5%), thioanisole (2.5%), H₂O (2.5%) and TFA (92.5%) for 90 minutes. After the resin was distilled off from the reaction solution, ether was added to wash the material, and the supernatant was removed by centrifugation. The
25 resulting crude product was dissolved in water, and lyophilized or dried under vacuum.

Reference Example 6

Fmoc-Nst-OH (6)

H-Nst-OH (250mg, 1.6mmol) was dissolved in THF: H₂O (1:2),
30 K₂CO₃ (214mg, 1.6mmol) and Fmoc-OSu (470mg, 1.4mmol) were added

under ice-cooling, and the mixture was stirred at room temperature overnight. 1N HCl was added to the reaction solution to adjust pH to 3 or lower, and the solvent was concentrated under reduced pressure. The residue was dissolved in AcOEt. The solution was washed with 1N HCl and brine, and dried over anhydrous MgSO₄. The solvent was concentrated under reduced pressure, and the residue was crystallized from n-hexane to obtain the objective product as white crystals.

Yield, 580mg (98%); mp, 161-164°C; $[\alpha]^{23}_D$ -22.8° (c = 0.50, DMSO); Rt, 29.27 min; ¹H NMR (DMSO-d₆) δ (ppm); 0.83-0.85 (m, 6H), 1.12-1.37 (m, 2H), 1.42-1.51 (m, 1H), 3.60 (brs, 1H), 3.78 (brs, 1H), 4.19-4.23 (m, 2H), 4.31-4.39(m, 1H), 6.81 (d, 1H, J = 9.3 Hz), 7.31-7.42(m, 4H), 7.64-7.68 (m, 2H), 7.86-7.89 (m, 2H); TOF-MS m/z: 406.29 (M+Na)⁺.

Reference Example 7

Fmoc-Pns-OH (7).

Using H-Pns-OH (2.00 g, 10mmol) as a starting raw material, the title compound was synthesized in the same method as that for Fmoc-Nst-OH.

Yield, 3.76 g (88%); mp, 184-188°C; $[\alpha]^{24}_D$ -57.4° (c = 0.50, dioxane); Rt, 30.32 min; ¹H NMR (DMSO-d₆) δ (ppm); 2.68-2.76 (m, 1H), 2.82-2.88 (m, 1H), 3.93 (d, 1H, J = 2.7 Hz), 4.01-4.17 (m, 4H), 7.10-7.43 (m, 9H), 7.66 (dd, 2H, J = 4.8, 6.9 Hz), 7.88 (dd, 2H, J = 8.4, 16 Hz); TOF-MS m/z: 440.33 (M+Na)⁺.

Reference Example 8

Fmoc-Apns-OH

Using H-Apns-OH (661mg, 3.4mmol) as a starting raw material, the title compound was synthesized according to the same method as that for Fmoc-Nst-OH.

Yield, 1.20 g (85%); mp, 184-186°C; $[\alpha]^{26}_D$ -5.80° (c = 0.50, dioxane); Rt, 29.75 min; ¹H NMR (DMSO-d₆) δ (ppm); 2.71-2.74 (m, 2H), 3.95-4.12 (m, 5H), 7.12-7.47 (m, 9H), 7.63 (t, 2H, J = 6.6 Hz), 7.87 (d, 2H, J = 7.5 Hz);

TOF-MS m/z : 440.43 ($M+Na$)⁺.

Reference Example 9

Fmoc-Dmt-OH

L-Penicillamine (500mg, 3.4mmol) was dissolved in water, and 37%
5 HCHO (750 μ L) was added, and the mixture was stirred at room
temperature. After 1 hour, K₂CO₃ (460mg, 3.4mmol) and Fmoc-OSu
(790mg, 2.3mmol) were added under ice-cooling, and the mixture was
stirred at room temperature overnight. 1N HCl was added to the reaction
10 solution to adjust the pH to 3 or lower, and the solvent was concentrated
under reduced pressure. The residue was dissolved in AcOEt, and the
solution was washed with 1 N HCl, and brine, and dried over anhydrous
Na₂SO₄. The solvent was concentrated under reduced pressure, and the
residue was crystallized from n-hexane to obtain the objective product as
white crystals.

15 Yield, 880mg (69%); mp, 48-52°C; [α]_D²⁸ -37.8° (c = 0.50, DMSO); Rt, 28.06
min; ¹H NMR (DMSO-d₆) δ (ppm); 1.40 (d, 3H, J = 13 Hz), 1.52 (d, 3H, J =
7.2 Hz), 4.12-4.38 (m, 4H), 4.56-4.68 (m, 2H), 7.29-7.44 (m, 4H), 7.61-7.68
(m, 2H), 7.87-7.91 (m, 2H); TOF-MS m/z : 406.47 ($M+Na$)⁺

Example 1

20 H-Glu-Val-Asn-Nst-Asp-Ala-Glu-Phe-OH (8).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as the starting resin,
and compound 6, the title compound was synthesized according to the
aforementioned procedure. The resulting crude product was purified
using preparative HPLC to obtain the objective product as a white powder.

25 Yield, 52%; purity, >98% by analytical HPLC (Rt, 18.68 min); TOF-MS m/z :
967.10 ($M+H$)⁺.

Example 2

H-Glu-Val-Asn-Pns-Asp-Ala-Glu-Phe-OH (9).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin,
30 and compound 7, the title compound was synthesized according to the

aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 73%; purity, >98% by analytical HPLC (Rt, 19.30 min); TOF-MS m/z: 1001.1 (M+H)⁺.

5 Example 3

H-Glu-Val-Asp-Pns-Asp-Ala-Glu-Phe-OH (10).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified
10 using preparative HPLC to obtain the objective product as a white powder. Yield, 64%; purity, >98% by analytical HPLC (Rt, 19.28 min); TOF-MS m/z: 1002.1 (M+H)⁺.

Example 4

H-Glu-Val-Gln-Pns-Asp-Ala-Glu-Phe-OH (11).

15 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 63%; purity, >98% by analytical HPLC (Rt, 18.12 min); TOF-MS m/z:
20 1015.1 (M+H)⁺.

Example 5

H-Glu-Val-Glu-Pns-Asp-Ala-Glu-Phe-OH (12).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the
25 aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 48%; purity, >98% by analytical HPLC (Rt, 19.31 min); TOF-MS m/z: 1016.1 (M+H)⁺.

Example 6

30 H-Glu-Val-Met-Pns-Asp-Ala-Glu-Phe-OH (13).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 51%; purity, >98% by analytical HPLC (Rt, 20.23 min); TOF-MS m/z: 1018.2 (M+H)⁺.

Example 7

H-Glu-Val-Leu-Pns-Asp-Ala-Glu-Phe-OH (14).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 39%; purity, >98% by analytical HPLC (Rt, 18.96 min); TOF-MS m/z: 1000.2 (M+H)⁺.

Example 8

H-Glu-Val-Lys-Pns-Asp-Ala-Glu-Phe-OH (15).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 88%; purity, >98% by analytical HPLC (Rt, 17.04 min); TOF-MS m/z: 1015.0 (M+H)⁺.

Example 9

H-Glu-Leu-Leu-Pns-Asp-Ala-Glu-Phe-OH (16).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 58%; purity, >98% by analytical HPLC (Rt, 19.66 min); TOF-MS m/z: 1014.1 (M+H)⁺.

Example 10

H-Glu-Ile-Leu-Pns-Asp-Ala-Glu-Phe-OH (17).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the
5 aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 68%; purity, >98% by analytical HPLC (Rt, 21.34 min); TOF-MS m/z: 1014.2 (M+H)⁺.

Example 11

10 H-Glu-Phe-Leu-Pns-Asp-Ala-Glu-Phe-OH (18).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the
aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder.
15 Yield, 58%; purity, >98% by analytical HPLC (Rt, 22.34 min); TOF-MS m/z: 1048.2 (M+H)⁺.

Example 12

H-Glu-Met-Leu-Pns-Asp-Ala-Glu-Phe-OH (19).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the
20 aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 60%; purity, >98% by analytical HPLC (Rt, 19.00 min); TOF-MS m/z: 1032.0 (M+H)⁺.

25 Example 13

H-Glu-Val-Leu-Pns-Ala-Ala-Glu-Phe-OH (20).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the
aforementioned procedure. The resulting crude product was purified
30 using preparative HPLC to obtain the objective product as a white powder.

Yield, 43%; purity, >98% by analytical HPLC (Rt, 19.16 min); TOF-MS m/z: 956.11 (M+H)⁺.

Example 14

H-Glu-Val-Leu-Pns-Asn-Ala-Glu-Phe-OH (21).

5 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 44%; purity, >98% by analytical HPLC (Rt, 19.79 min); TOF-MS m/z:
10 999.22 (M+H)⁺.

Example 15

H-Glu-Val-Leu-Pns-Glu-Ala-Glu-Phe-OH (22).

 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the
15 aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 36%; purity, >98% by analytical HPLC (Rt, 21.00 min); TOF-MS m/z:
 1014.0 (M+H)⁺.

Example 16

20 H-Glu-Val-Leu-Pns-Gln-Ala-Glu-Phe-OH (23).

 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the
 aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder.
25 Yield, 63%; purity, >98% by analytical HPLC (Rt, 23.15 min); TOF-MS m/z:
 1013.0 (M+H)⁺.

Example 17

H-Glu-Val-Leu-Pns-Dmt-Ala-Glu-Phe-OH (24).

 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin,
30 and compound 7 and Fmoc-Dmt-OH, the title compound was synthesized

according to the aforementioned procedure. The resulting crude product was purified by preparative HPLC to obtain the objective product as a white powder.

Yield, 21%; purity, >98% by analytical HPLC (Rt, 21.50 min); TOF-MS m/z: 1027.75 (M+H)⁺.

Example 18

H-Glu-Val-Leu-Apns-Asp-Ala-Glu-Phe-OH (25).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and Fmoc-Apns-OH, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 41%; purity, >98% by analytical HPLC (Rt, 18.74 min); TOF-MS m/z: 1000.1 (M+H)⁺.

Example 19

H-Glu-Val-Leu-Apns-Ala-Ala-Glu-Phe-OH (26).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and Fmoc-Apns-OH, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 36%; purity, >98% by analytical HPLC (Rt, 19.34 min); TOF-MS m/z: 956.29 (M+H)⁺.

Example 20

H-Glu-Val-Leu-Apns-Asn-Ala-Glu-Phe-OH (27).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and Fmoc-Apns-OH, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 63%; purity, >98% by analytical HPLC (Rt, 19.16 min); TOF-MS m/z: 999.13 (M+H)⁺.

Example 21

H-Glu-Val-Leu-Apns-Glu-Ala-Glu-Phe-OH (28).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and Fmoc-Apns-OH, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 59%; purity, >98% by analytical HPLC (Rt, 19.56 min); TOF-MS m/z: 1014.1 (M+H)⁺.

Example 22

H-Glu-Val-Leu-Apns-Gln-Ala-Glu-Phe-OH (29).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and Fmoc-Apns-OH, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 63%; purity, >98% by analytical HPLC (Rt, 19.15 min); TOF-MS m/z: 1013.1 (M+H)⁺.

Example 23

H-Glu-Val-Leu-Apns-Dmt-Ala-Glu-Phe-OH (30).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and Fmoc-Apns-OH and Fmoc-Dmt-OH, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 28%; purity, >98% by analytical HPLC (Rt, 20.74 min); TOF-MS m/z: 1028.8 (M+H)⁺.

Example 24

H-Glu-Val-Leu-Pns-Ala-Val-Glu-Phe-OH (31).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder.

Yield, 27%; purity, >98% by analytical HPLC (Rt, 21.75 min); TOF-MS m/z: 984.16 (M+H)⁺.

Example 25

H-Glu-Val-Leu-Pns-Ala-Leu-Glu-Phe-OH (32).

5 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin,
and compound 7, the title compound was synthesized according to the
aforementioned procedure. The resulting crude product was purified
using preparative HPLC to obtain the objective product as a white powder.
Yield, 42%; purity, >98% by analytical HPLC (Rt, 22.54 min); TOF-MS m/z:
10 998.15 (M+H)⁺.

Example 26

H-Glu-Val-Leu-Pns-Ala-Phe-Glu-Phe-OH (33).

 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin,
and compound 7, the title compound was synthesized according to the
15 aforementioned procedure. The resulting crude product was purified
using preparative HPLC to obtain the objective product as a white powder.
Yield, 31%; purity, >98% by analytical HPLC (Rt, 22.90 min); TOF-MS m/z:
1032.0 (M+H)⁺.

Example 27

20 H-Glu-Val-Leu-Pns-Ala-Glu-Glu-Phe-OH (34).

 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin,
and compound 7, the title compound was synthesized according to the
aforementioned procedure. The resulting crude product was purified
using preparative HPLC to obtain the objective product as a white powder.
25 Yield, 58%; purity, >98% by analytical HPLC (Rt, 20.06 min); TOF-MS m/z:
1014.2 (M+H)⁺.

Example 28

H-Glu-Val-Leu-Pns-Asp-Ala-Gln-Phe-OH (35).

 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin,
30 and compound 7, the title compound was synthesized according to the

aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 77%; purity, >98% by analytical HPLC (Rt, 19.47 min); TOF-MS m/z: 999.15 (M+H)⁺.

5 Example 29

H-Glu-Val-Leu-Pns-Asp-Ala-Val-Phe-OH (36).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified
10 using preparative HPLC to obtain the objective product as a white powder. Yield, 27%; purity, >98% by analytical HPLC (Rt, 21.90 min); TOF-MS m/z: 970.18 (M+H)⁺.

Example 30

H-Glu-Val-Leu-Pns-Asp-Ala-Leu-Phe-OH (37).

15 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 47%; purity, >98% by analytical HPLC (Rt, 23.18 min); TOF-MS m/z:
20 984.05 (M+H)⁺.

Example 31

H-Glu-Val-Leu-Pns-Asp-Ala-Phe-Phe-OH (38).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the
25 aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 49%; purity, >98% by analytical HPLC (Rt, 23.11 min); TOF-MS m/z: 1018.2 (M+H)⁺.

Example 32

30 H-Glu-Val-Leu-Pns-Asp-Ala-Glu-OH (39).

Using Fmoc-Glu (OtBu) Wang resin (0.560 mmol/g) as a starting resin and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 37%; purity, >98% by analytical HPLC (Rt, 18.48 min); TOF-MS m/z: 852.87 (M+H)⁺.

Example 33

H-Glu-Val-Leu-Pns-Asp-Ala-OH (40).

Using Fmoc-Ala Wang resin (0.536 mmol/g) as a starting resin and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 64%; purity, >98% by analytical HPLC (Rt, 16.50 min); TOF-MS m/z: 723.82 (M+H)⁺.

Example 34

H-Glu-Val-Leu-Pns-Asp-OH (41).

Using Fmoc-Asp (OtBu) Wang resin (0.586 mmol/g) as a starting resin and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 40%; purity, >98% by analytical HPLC (Rt, 19.62 min); TOF-MS m/z: 652.57 (M+H)⁺.

Example 35

H-Glu-Val-Leu-Pns-OH (42).

Using Fmoc-Pns Clt resin (0.582 mmol/g) as a starting resin, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 39%; purity, >98% by analytical HPLC (Rt, 15.80 min); TOF-MS m/z: 537.64 (M+H)⁺.

Example 36

H-Val-Leu-Pns-Asp-Ala-Glu-Phe-OH (43).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin and compound 7, the title compound was synthesized according to the
5 aforementioned procedure. The resulting crude product was purified
using preparative HPLC to obtain the objective product as a white powder.
Yield, 61%; purity, >98% by analytical HPLC (Rt, 21.15 min); TOF-MS m/z:
871.05 (M+H)⁺.

Example 37

10 H-Leu-Pns-Asp-Ala-Glu-Phe-OH (44).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin and compound 7, the title compound was synthesized according to the
aforementioned procedure. The resulting crude product was purified
using preparative HPLC to obtain the objective product as a white powder.
15 Yield, 49%; purity, >98% by analytical HPLC (Rt, 18.03 min); TOF-MS m/z:
771.66 (M+H)⁺.

Example 38

H-Pns-Asp-Ala-Glu-Phe-OH (45).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin and
20 compound 7, the title compound was synthesized according to the
aforementioned procedure. The resulting crude product was purified
using preparative HPLC to obtain the objective product as a white powder.
Yield, 49%; purity, >98% by analytical HPLC (Rt, 16.12 min); TOF-MS m/z:
658.53 (M+H)⁺.

Example 39

25 H-Gly-Val-Leu-Pns-Asp-Ala-Glu-Phe-OH (46).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin,
and compound 7, the title compound was synthesized according to the
aforementioned procedure. The resulting crude product was purified
30 using preparative HPLC to obtain the objective product as a white powder.

Yield, 78%; purity, >98% by analytical HPLC (Rt, 18.68 min); TOF-MS m/z: 928.11 (M+H)⁺.

Example 40

H-Glu-Gly-Leu-Pns-Asp-Ala-Glu-Phe-OH (47).

5 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 61%; purity, >98% by analytical HPLC (Rt, 18.88 min); TOF-MS m/z: 10 957.97 (M+H)⁺.

Example 41

H-Glu-Val-Gly-Pns-Asp-Ala-Glu-Phe-OH (48).

 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the 15 aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 48%; purity, >98% by analytical HPLC (Rt, 16.90 min); TOF-MS m/z: 943.89 (M+H)⁺.

Example 42

20 H-Glu-Val-Leu-Pns-Gly-Ala-Glu-Phe-OH (49).

 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the 25 aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 63%; purity, >98% by analytical HPLC (Rt, 18.84 min); TOF-MS m/z: 942.05 (M+H)⁺.

Example 43

H-Glu-Val-Leu-Pns-Asp-Gly-Glu-Phe-OH (50).

 Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, 30 and compound 7, the title compound was synthesized according to the

aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 44%; purity, >98% by analytical HPLC (Rt, 18.22 min); TOF-MS m/z: 986.04 (M+H)⁺.

5 Example 44

H-Glu-Val-Leu-Pns-Asp-Ala-Gly-Phe-OH (51).

Using Fmoc-Phe Wang resin (0.520 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 40%; purity, >98% by analytical HPLC (Rt, 19.21 min); TOF-MS m/z: 928.13 (M+H)⁺.

Example 45

H-Glu-Val-Leu-Pns-Asp-Ala-Glu-Gly-OH (52).

15 Using Fmoc-Phe Wang resin (0.462 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 35%; purity, >98% by analytical HPLC (Rt, 15.66 min); TOF-MS m/z: 909.92 (M+H)⁺.

Example 46

Boc-Pns-NH-Bzl (54).

To a solution of Boc-Pns-OH (650mg, 2.2 mmol) in DMF, were added HOBt·H₂O (337mg, 2.2mmol), EDC·HCl (422mg, 2.2mmol) and benzylamine (219μL, 2.0mmol), and the mixture was stirred overnight. The reaction solution was concentrated under reduced pressure. The residue was dissolved in AcOEt and was washed with 10% citric acid, 5% NaHCO₃ and brine, and dried over anhydrous MgSO₄. The solvent was concentrated under reduced pressure to obtain the objective product as a white solid.

Yield, 799mg (104%); mp, 138-140°C; ^1H NMR (DMSO- d_6) δ (ppm); 1.30 (s, 9H), 2.62-2.84 (m, 2H), 3.87 (m, 1H), 3.99 (m, 1H), 4.21-4.34 (m, 2H), 5.81 (d, 1H, $J = 6.0$ Hz), 6.19 (d, 1H, $J = 9.0$ Hz), 7.17-7.29 (m, 10H); HRMS (FAB $^+$) Calcd for $\text{C}_{22}\text{H}_{29}\text{N}_2\text{O}_4$ 385.2127 (M+H) $^+$, found 385.2134.

5 Example 47

Boc-Leu-Pns-NH-Bzl (55).

The compound 54 (385mg, 1.0mmol) of Example 46 was dissolved in 4 N HCl/ dioxane (3.0mL), and the mixture was stirred at room temperature for 1 hour. The reaction mixture was concentrated under reduced pressure, and the residue was precipitated with ether to obtain a hydrochloride. The hydrochloride was dissolved in a DMF solution, and neutralized with TEA (140 μ L, 1.0mmol). Thereto were added HOBt $\cdot$ H $_2$ O (169mg, 1.1mmol), Boc-Leu-OH $\cdot$ H $_2$ O (274mg, 1.1mmol), and EDC $\cdot$ HCl (211mg, 1.1mmol), and the mixture was stirred overnight. The reaction solution was concentrated under reduced pressure, and the residue was dissolved in AcOEt. The solution was washed with 10% citric acid, 5% NaHCO $_3$, and brine, and dried over anhydrous MgSO $_4$. The residue was concentrated under reduced pressure to obtain the objective product as a white solid.

20 Yield, 507mg (102%); mp, 151-154°C; ^1H NMR (DMSO- d_6) δ (ppm); 0.87 and 0.81 (each d, each 3H, $J = 6.6$ Hz), 1.23-1.55 (m, 3H), 1.36 (s, 9H), 2.56-2.63 (m, 1H), 2.83 (dd, 1H, $J = 6.6, 13$ Hz,), 3.85 (m, 1H), 4.16 (dd, 1H, $J = 6.6, 15$ Hz), 4.22 (m, 1H), 4.33 (dd, 1H, $J = 6.6, 15$ Hz), 6.16 (d, 1H, $J = 6.0$ Hz), 6.96 (d, 1H, $J = 9.0$ Hz), 7.20-7.28 (m, 10H), 7.44 (d, 1H, $J = 9.3$ Hz); HRMS (FAB $^+$) Calcd for $\text{C}_{28}\text{H}_{40}\text{N}_3\text{O}_5$ 498.2964 (M+H) $^+$, found 498.2968.

25 Example 48

Boc-Val-Leu-Pns-NH-Bzl (56).

Using compound 55 of Example 47 and Boc-Val-OH, the title compound was synthesized in the same method as that for compound 55.

Yield, 94%; mp, 219-222°C; ¹H NMR (DMSO-d₆) δ (ppm); 0.76 and 0.82 (m, 13H), 1.36 (s, 9H), 1.49-1.61 (m, 1H), 1.81-2.00 (m, 1H), 2.66-2.68 (m, 1H), 2.76-2.83 (m, 1H), 3.72-3.90 (m, 2H), 4.11-4.36 (m, 4H), 6.09 (m, 1H), 6.76 (d, 1H, J = 9.0 Hz), 7.18-7.28 (m, 10H), 7.56 (d, 1H, J = 9.0 Hz), 7.80 (d, 1H, J = 9.0 Hz); HRMS (FAB⁺) Calcd for C₃₃H₄₉N₄O₆ 597.3652 (M+H)⁺, found 597.3655.

Example 49


H-Glu-Val-Leu-Pns-NH-Bzl (57, KMI-205).

Compound 56 (119mg, 0.20mmol) of Example 48 was dissolved in 4 N HCl/dioxane, and the solution was stirred at room temperature for 1 hour. The reaction solution was concentrated under reduced pressure, and the residue was precipitated with ether to obtain a hydrochloride. This hydrochloride was dissolved in a DMF solution and neutralized with TEA (28.0μL, 0.20mmol). Then HOBt·H₂O (33.7mg, 0.22mmol), Z-Glu (OtBu)-OH (74.2mg, 0.22mmol) and EDC·HCl (42.2mg, 0.22mmol) were added thereto, and the mixture was stirred overnight. The reaction solution was concentrated under reduced pressure, the residue was dissolved in AcOEt. The solution was washed with 10% citric acid, 5% NaHCO₃, and brine, and dried over anhydrous MgSO₄. The solvent was concentrated under reduced pressure, and the residue was crystallized from ether to obtain white crystals. The crystals were added to thioanisole (100μL), and the mixture was dissolved in TFA (1.9mL), followed by stirring at room temperature for 3 hours. The solvent was concentrated under reduced pressure, and the residue was crystallized from ether to obtain a crude product. A part thereof was purified with preparative HPLC to obtain the objective product as a white powder.

Yield, 5.5mg (23%); mp, 147-150°C; Rt, 24.68 min; ¹H NMR (DMSO-d₆) δ (ppm); 0.76-0.87 (m, 13H), 1.23-1.49 (m, 3H), 1.81-2.00 (m, 2H), 2.26-2.31 (m, 2H), 2.61-2.83 (m, 2H), 3.81-3.90 (m, 2H), 4.12-4.35 (m, 5H), 6.09 (m, 1H), 7.18-7.26 (m, 10 H), 7.52 (d, 1H, J = 9.0 Hz), 8.10-8.22 (m, 2H), 8.45

(d, 1H, J = 9.0 Hz); HRMS (FAB⁺) Calcd for C₃₃H₄₈N₅O₇ 626.3554 (M+H)⁺, found 626.3547.

Example 50




(58).

To a solution of Boc-Pns-OH (500mg, 1.7mmol) in DMF, were added 5-aminoisophthalic acid dimethyl ester (354mg, 1.7mmol), HOBt·H₂O (285mg, 1.9mmol), and EDC·HCl (357mg, 1.9mmol), and the mixture was stirred overnight. The reaction solution was concentrated under reduced pressure, and the residue was dissolved in AcOEt. The solution was washed with 10% citric acid, 5% NaHCO₃, and brine, and dried over anhydrous Na₂SO₄. The solvent was concentrated under reduced pressure to obtain the objective product as a white solid.

Yield, 786mg (95%); mp, 180-183°C; ¹H NMR (DMSO-d₆) δ (ppm); 1.19 (s, 9H), 2.70-2.89 (m, 2H), 3.88 (s, 6H), 4.01-4.10 (m, 2H), 5.86 (d, 1H, J = 6.0 Hz), 6.42 (d, 1H, J = 9.6 Hz), 7.19-7.29 (m, 5H), 8.16 (s, 1H), 8.57 (m, 2H); HRMS (FAB⁺) Calcd for C₂₅H₃₀N₂O₈Na 509.1900 (M+Na)⁺, found 509.1905.

Example 51



(59).

Compound 58 (786mg, 1.6mmol) of Example 50 was added to anisole (211μL, 1.9mmol), and the mixture was dissolved in 4 N HCl/dioxane (8.0mL). The solution was stirred at room temperature for 1 hour. The reaction solution was concentrated under reduced pressure, and the residue was precipitated with ether to obtain a hydrochloride. The hydrochloride was dissolved in a DMF solution, and the solution was

5

10


15



20

25

Example 53



5

10

NC(CCC(=O)NCC(=O)NCC(=O)OCC(=O)O)C1=CC(=CC=C1C(=O)O)C(=O)O

15

20


25

Yield, 6.6mg (11%); mp, 198-203°C; Rt, 18.74 min; ¹H NMR (DMSO-d₆) δ (ppm); 0.47 (dd, 6H, J = 6.3, 28 Hz), 0.82 (d, 6H, J = 6.9 Hz), 1.11-1.23 (m, 3H), 1.86-1.91 (m, 3H), 2.27 (t, 2H, J = 8.1 Hz), 2.72-2.92 (m, 2H), 3.85 (t, 1H, J = 6.0 Hz), 3.96 (brs, 1H), 4.17 (t, 1H, 7.8 Hz), 4.32-4.34 (m, 2H), 6.28 (brs, 1H), 7.20-7.29 (m, 5H), 7.79 (d, 1H, J = 9.0 Hz), 8.04 (d, 1H, J = 8.1 Hz), 8.14 (s, 1H), 8.39 (d, 1H, J = 8.4 Hz), 8.50 (s, 2H); HRMS (FAB⁺) Calcd for C₃₄H₄₆N₅O₁₁ 700.3194 (M+H)⁺, found 700.3188.

NC(CCC(N)=O)C(CCC(N)=O)C(CCC(N)=O)C(CCC(N)=O)Cc1ccc(cc1)C(=O)O

Using Fmoc-NH-CH₂-C₆H₄-CO-Clt resin (0.492 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder.

Example 56




Using Fmoc-NH-CH₂-C₆H₄-CO-Clt resin (0.606 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder.

Yield, 76%; purity, >98% by analytical HPLC (Rt, 19.44 min); TOF-MS m/z:

670.77 (M+H)⁺.


Example 57



(65).

5 Using Fmoc-NH-C₆H₄-CO-Clt resin (0.612 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 56%; purity, >98% by analytical HPLC (R_t, 19.91 min); TOF-MS m/z: 656.73 (M+H)⁺.

Example 58



(66).

15 Using Fmoc-NH-C₆H₄-CO-Clt resin (0.505 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 45%; purity, >98% by analytical HPLC (R_t, 19.95 min); TOF-MS m/z: 656.74 (M+H)⁺.

Example 59

glt-Val-Leu-Pns-Asp-Ala-Gly-Phe-OH (67).

20 Using Fmoc-Glu (OtBu) Wang resin (0.560 mmol/g) as a starting resin, compound 7 and glutaric anhydride, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder.

Yield, 22%; purity, >98% by analytical HPLC (Rt, 19.94 min); TOF-MS m/z: 959.81 (M+Na)⁺.

Example 60

H-Gln-Val-Leu-Pns-Asp-Ala-Gly-Phe-OH (68).

5 Using Fmoc-Glu (OtBu) Wang resin (0.560 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 47%; purity, >98% by analytical HPLC (Rt, 16.82 min); TOF-MS m/z:
10 851.81 (M+H)⁺.

Example 61

H-Asp-Val-Leu-Pns-Asp-Ala-Gly-Phe-OH (69).

 Using Fmoc-Glu (OtBu) Wang resin (0.560 mmol/g) as a starting resin, and compound 7, the title compound was synthesized according to
15 the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the objective product as a white powder. Yield, 32%; purity, >98% by analytical HPLC (Rt, 17.82 min); TOF-MS m/z:
 834.00 (M+H)⁺.

Example 62

20 H-homoGlu-Val-Leu-Pns-Asp-Ala-Gly-Phe-OH (70)

 Using Fmoc-Glu (OtBu) Wang resin (0.560 mmol/g) as a starting resin, compound 7 and Fmoc-homoGlu (OtBu)-OH²¹, the title compound was synthesized according to the aforementioned procedure. The resulting crude product was purified using preparative HPLC to obtain the
25 objective product as a white powder. Yield, 22%; purity, >98% by analytical HPLC (Rt, 17.67 min); TOF-MS m/z:
 866.94 (M+H)⁺.


Example 63

β-ODap-Val-Leu-Pns-Asp-Ala-Gly-Phe-OH (71)

30 Using Fmoc-Glu (OtBu) Wang resin (0.560 mmol/g) as a starting

resin, compound 7, Boc-Dap (Fmoc)-OH and 2-(tert-butoxy)-2-oxoacetic acid²², the title compound was synthesized according to the
 aforementioned procedure. The resulting crude product was purified
 using preparative HPLC to obtain the objective product as a white powder.
 Yield, 18%; purity, >98% by analytical HPLC (Rt, 17.39 min); TOF-MS m/z:
 881.82 (M+H)⁺.

Example 64




(72).

Using compound 60 of Example 52 and Boc-Dap (Fmoc)-OH, the
 title compound was synthesized in the same method as that for compound
 59 of Example 51.

Yield, 84%; mp, 198-202°C; ¹H NMR (DMSO-d₆) δ (ppm); 0.53 (dd, 6H, J =
 6.6, 26 Hz), 0.77 (t, 6H, J = 6.3 Hz), 1.07-1.20 (m, 2H), 1.33 (brs, 1H), 1.36
 (s, 9H), 1.88-1.99 (m, 1H), 2.70-2.91 (m, 2H), 3.26 (brs, 2H), 3.90 (s, 6H),
 3.97 (d, 1H, J = 4.2 Hz), 4.01-4.07 (m, 1H), 4.18-4.31 (m, 6H), 6.30 (d, 1H,
 J = 5.4 Hz), 6.99 (d, 1H, J = 7.8 Hz), 7.21 (d, 1H, J = 8.4 Hz), 7.25-7.42 (m,
 9H), 7.58 (d, 1H, J = 8.1 Hz), 7.67 (d, 2H, J = 7.2 Hz), 7.74 (d, 1H, J = 8.4
 Hz), 7.88 (d, 2H, J = 7.5 Hz), 7.94 (d, 1H, J = 8.1 Hz), 8.15 (s, 1H), 8.59 (s,
 2H); HRMS (FAB⁺) Calcd for C₅₄H₆₇N₆O₁₃ 1007.4766 (M+H)⁺, found
 1007.4760.

Example 65




(73)

Compound 72 (60.0mg, 0.060mmol) of Example 64 was dissolved in 20% diethylamine/DMF, and the mixture was stirred at room temperature for 20 minutes, and concentrated under reduced pressure. The residue was dissolved in AcOEt, and the solution was washed with 5% NaHCO₃, and brine, and dried over anhydrous Na₂SO₄. The solvent was concentrated under reduced pressure, and the residue was dissolved in DMF again. TEA (16.6μL, 0.12mmol) and chloroformic acid ethyl ester (14.5μL, 0.13mmol) were added thereto, and the mixture was stirred at room temperature overnight. The reaction solution was concentrated under reduced pressure, and the residue was crystallized from ether to obtain the objective product as a white crystal.

Yield, 64.7mg (123%); mp, 143-148°C; ¹H NMR (DMSO-d₆) δ (ppm); 0.53 (dd, 6H, J = 6.3, 24 Hz), 0.75 (t, 6H, J = 6.0 Hz), 1.16-1.36 (m, 15H), 1.87-1.93 (m, 1H), 2.73-2.89 (m, 4H), 3.87 (s, 6H), 3.96-3.97 (m, 1H), 4.11-4.23 (m, 4H), 4.30 (t, 2H, J = 8.1 Hz), 6.36 (d, 1H, J = 5.7 Hz), 7.15 (d, 1H, J = 7.8 Hz), 7.23-7.29 (m, 5H), 7.62 (d, 1H, J = 7.8 Hz), 7.74 (d, 1H, J = 8.7 Hz), 7.94 (d, 1H, J = 8.7 Hz), 8.16 (s, 1H), 8.59 (s, 2H), 8.78 (brs, 1H); HRMS (FAB⁺) Calcd for C₄₃H₆₁N₆O₁₄ 885.4246 (M+H)⁺, found 885.4243.

Example 66



(74)

Compound 73 (34.7mg, 0.039mmol) of Example 65 was dissolved in DMF, and 4N NaOH (98.0μL, 0.39mmol) was added thereto under ice-cooling, followed by stirring at room temperature overnight. The reaction solution was acidified with citric acid, and concentrated under reduced pressure, and the residue was dissolved in AcOEt. The solution was washed with 10% citric acid, and brine, and dried with anhydrous Na₂SO₄. The solvent was concentrated under reduced pressure, and anisole (5.11μL,

0.047mmol) was added to the residue. The mixture was dissolved in 4N HCl/dioxane (3.0mL) again, and the solution was stirred. After 1 hour, the reaction solution was concentrated under reduced pressure, and the residue was crystallized from ether to obtain a crude product. A part thereof was purified by preparative HPLC to obtain the objective product as a white powder.

Yield, 3.7mg (14%); mp, 220-224°C; Rt, 18.55 min; ¹H NMR (DMSO-d₆) δ (ppm); 0.54 (dd, 6H, J = 6.3, 23 Hz), 0.74-0.82 (m, 6H), 1.16-1.23 (m, 3H), 1.90-1.99 (m, 2H), 2.27 (brs, 1H), 2.72-2.89 (m, 2H), 3.96-4.00 (m, 2H), 4.16 (t, 1H, J = 7.2 Hz), 4.30 (t, 2H, J = 7.5 Hz), 7.21 (d, 1H, J = 6.6 Hz), 7.27-7.29 (m, 5H), 7.84 (d, 1H, J = 8.7 Hz), 7.98 (d, 1H, J = 8.4 Hz), 8.06 (d, 1H, J = 8.1 Hz), 8.14 (s, 1H), 8.50 (s, 2H), 8.59 (brs, 1H); HRMS (FAB⁺) Calcd for C₃₄H₄₅N₆O₁₂ 729.3095 (M+H)⁺, found 729.3103.

Example 67

Experiment regarding measurement of BACE1 inhibitory activity

Enzyme used in measurement

As an enzyme having β-secretase activity, recombinant human BACE-1 (rhBACE-1) purchased from R&D systems (Minneapolis, MN) was used. rhBACE-1 is expressed as only an extracellular region (amino acid 1-460 residues) of recombinant human β-secretase, with a His tag at its carboxyl terminal in murine myeloma cell line, NS0. Purified secreted rhBACE1 is a mixture of a pro entity (about 75 kDa) and a mature entity (about 72 kDa). The purity is > 90% (as determined by SDS-PAGE and visualized by silver stain), and enzyme activity is exhibited under the weak acidity (pH 3.5-5.5).

Assessment of enzyme inhibitory activity

Enzyme inhibitory activity of an inhibitor was assessed by quantitating an N-terminal fragment produced after cleaving the substrate with BACE1 by reverse HPLC using (7-methoxycoumarin-4-yl)acetyl-Ser-Glu-Val-Asn-Leu-Asp-Ala-Glu-Phe-Arg-Lys(2,4-dinitrophenyl)-Arg-Arg-NH₂

purchased from Peptide Institute (Osaka, Japan) as a substrate for BACE1. To 40 μ L of 62.5 mM 2-(N-morpholino)ethansulfonic acid (MES)-CH₃COOH-NaOH buffer (pH 5.0), were added 1 μ L of a solution of an inhibitor diluted with DMSO (100 μ M), 4 μ L of rhBACE-1 (6.25 ng/ μ L), and 5 μ L of an aqueous solution (50% DMSO) of 250 μ M substrate (7-methoxycoumarin-4-yl)acetyl-Ser-Glu-Val-Asn-Leu-Asp-Ala-Glu-Phe-Arg-Lys(2,4-dinitrophenyl)-Arg-Arg-NH₂ to initiate an enzyme reaction. After incubation at 37°C for 60 minutes, 20 μ L of 20% trichloroacetic acid was added to stop the reaction. And, inhibition % was calculated by measuring a fluorescent intensity of the N-terminal fragment (Ex, 328nm; Em, 393nm) using reverse phase HPLC (linear gradient of MeCN in aqueous TFA; flow rate, 1.0 mL/min).

BACE1 inhibitory activity

Enzyme inhibitory activity of an inhibitor was assessed by quantitating a N-terminal fragment produced after cleavage of a substrate with BACE1 by reverse phase HPLC using (7-methoxycoumarin-4-yl)acetyl-Ser-Glu-Val-Asn-Leu-Asp-Ala-Glu-Phe-Arg-Lys(2,4-dinitrophenyl)-Arg-Arg-NH₂ purchased from Peptide Institute (Osaka, Japan) as a substrate for BACE1. Recombinant human BACE-1 purchased from R&D systems (Minneapolis, MN) was used. To 40 μ L of 62.5 mM 2-(N-morpholino)ethansulfonic acid (MES)-CH₃COOH-NaOH buffer (pH 5.0), were added 1 μ L of a solution of an inhibitor diluted with DMSO (100 μ M), 4 μ L of BACE1 (6.25 ng/ μ L), and 5 μ L of an aqueous solution (50% DMSO) of 250 μ M substrate (7-methoxycoumarin-4-yl)acetyl-Ser-Glu-Val-Asn-Leu-Asp-Ala-Glu-Phe-Arg-Lys(2,4-dinitrophenyl)-Arg-Arg-NH₂ to initiate an enzyme reaction. After incubation at 37°C for 60 minutes, 20 μ L of 20% trichloroacetic acid was added to stop a reaction. Inhibition % was calculated by measuring a fluorescent intensity of the N-terminal fragment (Ex, 328nm; Em, 393nm) using reverse phase HPLC (linear gradient of MeCN in aqueous TFA; flow rate, 1.0 mL/min).


Results are shown in the following table.

Table 1

Example Number	Activity	Example Number	Activity
127	100	142	89.83
185	100	7	89.65
186	100	76	89.49
187	100	172	89.45
189	100	163	88.31
135	99.77	184	88.13
66	99.75	77	87.92
134	99.56	173	86.06
137	99.47	78	85.85
188	99.46	79	85.32
136	99.43	166	84.88
190	99.35	179	83.67
125	99.07	58	83.67
126	98.93	80	83.24
68	98.83	81	82.89
167	98.77	181	82.52
177	98.50	174	81.83
69	98.36	82	81.76
54	98.32	57	81.46
70	98.31	43	81.23
71	97.89	149	80.13
124	97.28	83	80.13
63	96.66	84	79.08
72	95.01	27	78.43
148	93.76	156	78.28
73	93.01	182	75.72
171	91.80	85	75.23
150	90.80	86	74.98
74	90.52	146	74.74
75	90.42	131	74.08
165	73.69	26	60.23
178	73.43	168	60.03
87	72.25	138	59.62
32	71.60	147	59.46
162	71.54	96	59.25
133	71.49	97	59.11
160	71.33	98	58.41
88	70.42	99	58.04
154	69.89	29	57.47
89	69.77	130	57.22
155	69.70	151	57.00
90	69.04	100	56.93

Example Number	Activity	Example Number	Activity
143	68.96	42	56.72
91	68.90	30	56.57
132	68.08	44	56.30
141	67.49	101	56.12
45	66.79	55	55.98
31	66.56	28	55.88
158	66.10	103	54.60
153	65.79	15	54.52
92	65.37	104	54.42
93	64.84	105	54.04
161	64.74	13	53.71
164	64.12	128	53.57
56	64.05	18	53.38
34	63.71	106	52.95
94	61.80	183	52.15
95	61.66	107	52.14
169	61.57	180	51.43
123	61.38	108	51.32
129	61.04	145	50.49
170	60.53	109	50.03


Example 68



The above compound was prepared in the same manner as in Example 55.

5 Yield 14.1mg (68%), TOF-MS: 685.87


Example 69



The above compound was prepared in the same manner as in Example 55.

Yield 4.0mg (46%), TOF-MS: 685.75


Example 70



5 The above compound was prepared in the same manner as in Example 55.

Yield 3.7mg (37%), TOF-MS: 800.91


Example 71



The above compound was prepared in the same manner as in Example 55.

10 Yield 3.1mg (19%), TOF-MS: 800.84

Example 72




The above compound was prepared in the same manner as in Examples 50-54.

15 Yield 2.5mg (5%), TOF-MS: 803.60

Example 73


56



The above compound was prepared in the same manner as in Example 1.

Yield 4.8mg (14%), TOF-MS: 906.03

Example 74




5

The above compound was prepared in the same manner as in Example 55.

Yield 8.1mg (31%), TOF-MS: 685.84

Example 75




10

The above compound was prepared in the same manner as in Example 1.

Yield 12.5mg (42%), TOF-MS: 1002.04

Example 76


57



The above compound was prepared in the same manner as in Example 55.

Yield 1.0mg (4%), TOF-MS: 771.73

Example 77




5

The above compound was prepared in the same manner as in Example 55.

Yield 5.0mg (58%), TOF-MS: 685.85

Example 78




10

The above compound was prepared in the same manner as in Example 55.

Yield 0.3mg (1%), TOF-MS: 771.92


Example 79



The above compound was prepared in the same manner as in Example 55.

Yield 10.5mg (32%), TOF-MS: 771.53


Example 80



The above compound was prepared in the same manner as in Example 1.

5 Yield 11.9mg (48%), TOF-MS: 1076.02


Example 81



The above compound was prepared in the same manner as in Example 1.

Yield 13.8mg (39%), TOF-MS: 957.01

10 Example 82




The above compound was prepared in the same manner as in Example 1.

Yield 11.6mg (26%), TOF-MS: 1032.15

Example 83


59



The above compound was prepared in the same manner as in Example 55.

Yield 1.0mg (8%), TOF-MS: 691.08

Example 84




5

The above compound was prepared in the same manner as in Example 1.

Yield 10.1mg (24%), TOF-MS: 957.24

Example 85




10

The above compound was prepared in the same manner as in Example 1.

Yield 19.6mg (52%), TOF-MS: 881.70


Example 86



The above compound was prepared in the same manner as in Example 1.

Yield 26.3mg (28%), TOF-MS: 1042.28


Example 87



5 The above compound was prepared in the same manner as in Example 55.

Yield 3.3mg (4%), TOF-MS: 842.83


Example 88



The above compound was prepared in the same manner as in Example 55.

10 Yield 5.4mg (12%), TOF-MS: 763.82


Example 89



The above compound was prepared in the same manner as in Example 55.

Yield 18.0mg (42%), TOF-MS: 670.40


15 Example 90



The above compound was prepared in the same manner as in Example 1.

Yield 18.2mg (52%), TOF-MS: 684.71

Example 91




5

The above compound was prepared in the same manner as in Example 1.

Yield 20.4mg (39%), TOF-MS: 866.82

Example 92




10

The above compound was prepared in the same manner as in Example 55.

Yield 22.7mg (67%), TOF-MS: 842.34


Example 93



The above compound was prepared in the same manner as in Example 55.

Yield 32.0mg (79%), TOF-MS: 842.92


Example 94



5 The above compound was prepared in the same manner as in Example 55.

Yield 7.9mg (19%), TOF-MS: 834.87


Example 95



The above compound was prepared in the same manner as in Example 1.

10 Yield 17.7mg (33%), TOF-MS: 909.82

Example 96




The above compound was prepared in the same manner as in Example 1.

Yield 32.2mg (51%), TOF-MS: 984.18

15 Example 97


63



The above compound was prepared in the same manner as in Example 1.

Yield 22.1mg (48%), TOF-MS: 966.19

Example 98




5

The above compound was prepared in the same manner as in Example 1.

Yield 19.9mg (60%), TOF-MS: 838.94

Example 99




10

The above compound was prepared in the same manner as in Example 1.

Yield 18.0mg (34%), TOF-MS: 1041.14


Example 100



The above compound was prepared in the same manner as in Example 1.

Yield 23.9mg (65%), TOF-MS: 652.65


Example 101



The above compound was prepared in the same manner as in Example 1.

5 Yield 9.3mg (24%), TOF-MS: 909.12


Example 102



The above compound was prepared in the same manner as in Example 55.

Yield 3.2mg (32%), TOF-MS: 656.79


10 Example 103



The above compound was prepared in the same manner as in Example 55.

Yield 17.0mg (32%), TOF-MS: 785.79


Example 104



The above compound was prepared in the same manner as in Example 55.

Yield 11.0mg (30%), TOF-MS: 785.24

Example 105




5

The above compound was prepared in the same manner as in Example 1.

Yield 13.3mg (40%), TOF-MS: 618.89

Example 106




10

The above compound was prepared in the same manner as in Example 1.

Yield 11.2mg (21%), TOF-MS: 1027.75


Example 107



The above compound was prepared in the same manner as in Example 1.

Yield 32.2mg (75%), TOF-MS: 998.60


Example 108



5 The above compound was prepared in the same manner as in Example 55.

Yield 22.3mg (45%), TOF-MS: 856.94


Example 109



The above compound was prepared in the same manner as in Example 55.

10 Yield 2.8mg (10%), TOF-MS: 834.89

Example 110




The above compound was prepared in the same manner as in Example 55.

Yield 0.1mg (10%), TOF-MS: 834.97

15 Example 111


67



The above compound was prepared in the same manner as in Example 1.

Yield 11.0mg (29%), TOF-MS: 965.99

Example 112




5

The above compound was prepared in the same manner as in Example 55.

Yield 24.4mg (47%), TOF-MS: 848.85

Example 113




10

The above compound was prepared in the same manner as in Example 1.


Yield 26.9mg (55%), TOF-MS: 851.91

Example 114



Yield 7.8mg (23%), TOF-MS: 1004.07


Example 115



5 The above compound was prepared in the same manner as in Example 1.

Yield 12.4mg (18%), TOF-MS: 880.88


Example 116



The above compound was prepared in the same manner as in Example 1.

10 Yield 8.8mg (16%), TOF-MS: 982.36


Example 117



The above compound was prepared in the same manner as in Example 1.

Yield 13.4mg (25%), TOF-MS: 1014.38


Example 118



5 The above compound was prepared in the same manner as in Examples 50-54.

Yield 2.6mg (25%), TOF-MS: 812.96


Example 119



10 The above compound was prepared in the same manner as in Example 1.

Yield 27.7mg (77%), TOF-MS: 808.56

Example 120




The above compound was prepared in the same manner as in Example 1.

15 Yield 12.1mg (20%), TOF-MS: 1000.11

Example 121


70



The above compound was prepared in the same manner as in Examples 46-49.

Yield 3.9mg (34%), TOF-MS: 640.87


5 Example 122



The above compound was prepared in the same manner as in Examples 46-49.

Yield 21.2mg (49%), TOF-MS: 707.78

10 Example 123




The above compound was prepared in the same manner as in Example 1.

Yield 7.3mg (72%), TOF-MS: 986.12

Example 124

71




The above compound was prepared in the same manner as in Examples 46-49.

Yield 5.7mg (57%), TOF-MS: 708.81

5


Example 125



The above compound was prepared in the same manner as in Example 55.

Yield 0.6mg (7.4%), TOF-MS: 709.73

Example 126




10

The above compound was prepared in the same manner as in Example 55.

Yield 4.1mg (25%), TOF-MS: 691.73

Example 127


72



The above compound was prepared in the same manner as in Examples 55-54.

Yield 4.1mg (19%), TOF-MS: 753.92


5 Example 128



The above compound was prepared in the same manner as in Example 55.

Yield 15.0mg (14%), TOF-MS: 670.86

Example 129




10

The above compound was prepared in the same manner as in Examples 46 to 49.

Yield 30.0mg (20%), TOF-MS: 672.97

Example 130


73



The above compound was prepared in the same manner as in Examples 46 to 49.

Yield 3.2mg (6.4%), TOF-MS: 628.66


5 Example 131



The above compound was prepared in the same manner as in Example 55.

Yield 8.6mg (46%), TOF-MS: 660.79

Example 132




10

The above compound was prepared in the same manner as in Example 55.

Yield 5.8mg (29%), TOF-MS: 674.83

Example 133


74



The above compound was prepared in the same manner as in Example 1.

Yield 4.2mg (13%), TOF-MS: 945.88

Example 134




5

The above compound was prepared in the same manner as in Examples 50-54.

Yield 4.6mg (46%), TOF-MS: 769.88

Example 135




10

The above compound was prepared in the same manner as in Examples 50-54.

Yield 5.0mg (50%), TOF-MS: 793.89

Example 136


75



The above compound was prepared in the same manner as in Example 55.

Yield 6.2mg (35%), TOF-MS: 769.79

Example 137




5

The above compound was prepared in the same manner as in Example 55.

Yield 2.1mg (9%), TOF-MS: 749.93

Example 138




10

The above compound was prepared in the same manner as in Examples 46-49.

Yield 4.4mg (44%), TOF-MS: 738.92

Example 139


76



The above compound was prepared in the same manner as in Example 55.

Yield 7.7mg (57%), TOF-MS: 672.78

Example 140




5

The above compound was prepared in the same manner as in Examples 46-49.

Yield 2.5mg (9.3%), TOF-MS: 642.77

Example 141




10

The above compound was prepared in the same manner as in Example 55.

Yield 6.1mg (16%), TOF-MS: 656.75

Example 142


77



The above compound was prepared in the same manner as in Example 55.

Yield 6.5mg (43%), TOF-MS: 751.96

Example 143




5

The above compound was prepared in the same manner as in Example 55.

Yield 3.5mg (17%), TOF-MS: 706.85

Example 144




10

The above compound was prepared in the same manner as in Example 55.

Yield 5.6mg (42%), TOF-MS: 766.81

Example 145


78



The above compound was prepared in the same manner as in Example 55.

Yield 4.8mg (26%), TOF-MS: 747.84

Example 146




5

The above compound was prepared in the same manner as in Example 55.

Yield 3.1mg (16%), TOF-MS: 733.87

Example 147




10

The above compound was prepared in the same manner as in Example 55.

Yield 4.5mg (29%), TOF-MS: 748.00

Example 148


79



The above compound was prepared in the same manner as in Example 55.

Yield 3.3mg (19%), TOF-MS: 749.88

Example 149




5

The above compound was prepared in the same manner as in Examples 46-49.

Yield 10.0mg (6.5%), TOF-MS: 698.67

Example 150




10

The above compound was prepared in the same manner as in Example 55.

Yield 7.3mg (58%), TOF-MS: 733.93

Example 151


80



The above compound was prepared in the same manner as in Example 55.

Yield 7.3mg (53%), TOF-MS: 670.62

Example 152




5

The above compound was prepared in the same manner as in Example 55.

Yield 6.3mg (48%), TOF-MS: 690.64

Example 153




10

The above compound was prepared in the same manner as in Example 55.

Yield 6.2mg (44%), TOF-MS: 656.78


Example 154



The above compound was prepared in the same manner as in Example 55.

Yield 5.3mg (53%), TOF-MS: 733.93


Example 155



5 The above compound was prepared in the same manner as in Example 55.

Yield 2.2mg (4.6%), TOF-MS: 662.86


Example 156



10 The above compound was prepared in the same manner as in Examples 46-49.

Yield 35mg (23%), TOF-MS: 682.81


Example 157



The above compound was prepared in the same manner as in Example 55.

15 Yield 3.0mg (5.5%), TOF-MS: 662.91


Example 158



The above compound was prepared in the same manner as in Examples 46-49.

Yield 40mg (23%), TOF-MS: 782.89


5 Example 159



The above compound was prepared in the same manner as in Example 55.

Yield 4.9mg (40%), TOF-MS: 706.87

Example 160




The above compound was prepared in the same manner as in Example 55.

Yield 3.5mg (32%), TOF-MS: 729.87

Example 161


83



The above compound was prepared in the same manner as in Example 55.

Yield 6.2mg (52%), TOF-MS: 642.87

Example 162




5

The above compound was prepared in the same manner as in Examples 46-49.

Yield 50.0mg (33%), TOF-MS: 694.87

Example 163




10

The above compound was prepared in the same manner as in Examples 50-54.

Yield 10.0mg (6.0%), TOF-MS: 756.58

Example 164


84



The above compound was prepared in the same manner as in Example 55.

Yield 4.5mg (48%), TOF-MS: 676.62

Example 165




5

The above compound was prepared in the same manner as in Example 55.

Yield 6.4mg (62%), TOF-MS: 740.75

Example 166




10

The above compound was prepared in the same manner as in Example 55.

Yield 6.6mg (53%), TOF-MS: 704.87

Example 167


85



The above compound was prepared in the same manner as in Example 55.

Yield 3.7mg (44%), TOF-MS: 877.78

Example 168




5

The above compound was prepared in the same manner as in Example 55.

Yield 1.3mg (8%), TOF-MS: 725.02

Example 169




10

The above compound was prepared in the same manner as in Example 55.

Yield 6.0mg (58%), TOF-MS: 690.72

Example 170


86



The above compound was prepared in the same manner as in Example 55.

Yield 5.2mg (34%), TOF-MS: 758.80

Example 171




5

The above compound was prepared in the same manner as in Examples 46-49.

Yield 3.1mg (13%), TOF-MS: 775.89

Example 172




10

The above compound was prepared in the same manner as in Example 55.

Yield 4.1mg (44%), TOF-MS: 749.82

Example 173


87



The above compound was prepared in the same manner as in Example 55.

Yield 4.7mg (47%), TOF-MS: 749.85

Example 174




5

The above compound was prepared in the same manner as in Example 55.

Yield 5.5mg (56%), TOF-MS: 707.76

Example 175




10

The above compound was prepared in the same manner as in Example 55.

Yield 4.1mg (39%), TOF-MS: 628.66

Example 176


88



The above compound was prepared in the same manner as in Example 55.

Yield 4.8mg (47%), TOF-MS: 720.91

Example 177




5

The above compound was prepared in the same manner as in Example 55.

Yield 0.77mg (2.1%), TOF-MS: 808.99

Example 178




10

The above compound was prepared in the same manner as in Examples 50-54.

Yield 0.4mg (1.4%), TOF-MS: 714.83

Example 179

89




The above compound was prepared in the same manner as in Examples 50-54.

Yield 0.03mg (0.2%), TOF-MS: 699.71

5

Example 180




The above compound was prepared in the same manner as in Examples 50-54.

Yield 30.0mg (19%), TOF-MS: 726.86

10

Example 181




The above compound was prepared in the same manner as in Example 55.

Yield 4.4mg (14%), TOF-MS: 790.92

Example 182

90




The above compound was prepared in the same manner as in Examples 46-49.

Yield 5.0mg (5.0%), TOF-MS: 755.65

5


Example 183



The above compound was prepared in the same manner as in Example 55.

Yield 4.2mg (12%), TOF-MS: 775.97

Example 184




10

The above compound was prepared in the same manner as in Examples 50-54.

Yield 15mg (8.1%), TOF-MS: 844.72

Example 185


91



The above compound was prepared in the same manner as in Example 55.

Yield 4.5mg (36%), TOF-MS: 733.98

Example 186




5

The above compound was prepared in the same manner as in Example 55.

Yield 4.9mg (43%), TOF-MS: 777.82

Example 187




10

The above compound was prepared in the same manner as in Example 55.

Yield 7.0mg (55%), TOF-MS: 793.87

Example 188


92



The above compound was prepared in the same manner as in Example 55.

Yield 5.1mg (40%), TOF-MS: 813.88

Example 189




5

The above compound was prepared in the same manner as in Example 55.

Yield 6.9mg (54%), TOF-MS: 833.98

Example 190




10

The above compound was prepared in the same manner as in Example 55.

Yield 5.0mg (38%), TOF-MS: 773.94


Example 191



The above compound was prepared in the same manner as in Example 55.

Yield 6.4mg (47%), TOF-MS: 658.66

Example 192



5

The above compound was prepared in the same manner as in Example 55.

Yield 7.1mg (32%), TOF-MS: 514.44

1/38

SEQUENCE LISTING

<110> KISO, Yoshiaki

<120> PEPTIDE DERIVATIVES HAVING β SECRETASE INHIBITORY ACTIVITY

<130> 664382

<150> JP 2003-052926

<151> 2003-02-28

<160> 95

<210> 1

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-5-methylhexanoic acid.

<400> 1

Glu Val Asn Xaa Asp Ala Glu Phe

1

5

<210> 2

<211> 8

<212> PRT

<213> Artificial Sequence

2/38

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 2

Glu Val Asn Xaa Asp Ala Glu Phe

1

5

<210> 3

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 3

Glu Val Asp Xaa Asp Ala Glu Phe

1

5

<210> 4

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 4

Glu Val Gln Xaa Asp Ala Glu Phe

1

5

<210> 5

3/38

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 5

Glu Val Glu Xaa Asp Ala Glu Phe

1

5

<210> 6

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 6

Glu Val Met Xaa Asp Ala Glu Phe

1

5

<210> 7

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 7

4/38

Glu Val Leu Xaa Asp Ala Glu Phe

1

5

<210> 8

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 8

Glu Val Lys Xaa Asp Ala Glu Phe

1

5

<210> 9

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 9

Glu Leu Leu Xaa Asp Ala Glu Phe

1

5

<210> 10

<211> 8

<212> PRT

<213> Artificial Sequence

5/38

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 10

Glu Ile Leu Xaa Asp Ala Glu Phe

1

5

<210> 11

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 11

Glu Phe Leu Xaa Asp Ala Glu Phe

1

5

<210> 12

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 12

Glu Met Leu Xaa Asp Ala Glu Phe

1

5

6/38

<210> 13

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 13

Glu Val Leu Xaa Ala Ala Glu Phe

1

5

<210> 14

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 14

Glu Val Leu Xaa Asn Ala Glu Phe

1

5

<210> 15

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

7/38

<400> 15

Glu Val Leu Xaa Glu Ala Glu Phe

1

5

<210> 16

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 16

Glu Val Leu Xaa Gln Ala Glu Phe

1

5

<210> 17

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 5 is (R)-5,5-dimethyl-1,3-thiazolidine-4-carboxylic acid.

<400> 17

Glu Val Leu Xaa Xaa Ala Glu Phe

1

5

<210> 18

8/38

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2S,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 18

Glu Val Leu Xaa Asp Ala Glu Phe

1

5

<210> 19

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2S,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 19

Glu Val Leu Xaa Ala Ala Glu Phe

1

5

<210> 20

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2S,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 20

9/38

Glu Val Leu Xaa Asn Ala Glu Phe

1

5

<210> 21

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2S, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 21

Glu Val Leu Xaa Glu Ala Glu Phe

1

5

<210> 22

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2S, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 22

Glu Val Leu Xaa Gln Ala Glu Phe

1

5

<210> 23

<211> 8

<212> PRT

<213> Artificial Sequence

10/38

<220> Xaa at position 4 is (2S, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 5 is (R)-5,5-dimethyl-1,3-thiazolidine-4-carboxylic acid.

<400> 23

Glu Val Leu Xaa Xaa Ala Glu Phe

1

5

<210> 24

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 24

Glu Val Leu Xaa Ala Val Glu Phe

1

5

<210> 25

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 25

Glu Val Leu Xaa Ala Leu Glu Phe

11/38

1 5

<210> 26

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 26

Glu Val Leu Xaa Ala Phe Glu Phe

1 5

<210> 27

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 27

Glu Val Leu Xaa Ala Glu Glu Phe

1 5

<210> 28

<211> 8

<212> PRT

<213> Artificial Sequence

12/38

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 28

Glu Val Leu Xaa Asp Ala Gln Phe

1

5

<210> 29

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 29

Glu Val Leu Xaa Asp Ala Val Phe

1

5

<210> 30

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 30

Glu Val Leu Xaa Asp Ala Leu Phe

1

5

<210> 31

13/38

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 31

Glu Val Leu Xaa Asp Ala Phe Phe

1

5

<210> 32

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 32

Glu Val Leu Xaa Asp Ala Glu

1

5

<210> 33

<211> 6

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 33

14/38

Glu Val Leu Xaa Asp Ala

1 5

<210> 34

<211> 5

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 34

Glu Val Leu Xaa Asp

1 5

<210> 35

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 35

Val Leu Xaa Asp Ala Glu Phe

1 5

<210> 36

<211> 6

<212> PRT

<213> Artificial Sequence

15/38

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 36

Leu Xaa Asp Ala Glu Phe

1 5

<210> 37

<211> 5

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 37

Xaa Asp Ala Glu Phe

1 5

<210> 38

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 38

Gly Val Leu Xaa Asp Ala Glu Phe

1 5

16/38

<210> 39

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 39

Glu Gly Leu Xaa Asp Ala Glu Phe

1

5

<210> 40

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 40

Glu Val Gly Xaa Asp Ala Glu Phe

1

5

<210> 41

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

17/38

<400> 41

Glu Val Leu Xaa Gly Ala Glu Phe

1

5

<210> 42

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 42

Glu Val Leu Xaa Asp Gly Glu Phe

1

5

<210> 43

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 43

Glu Val Leu Xaa Asp Ala Gly Phe

1

5

<210> 44

<211> 8

<212> PRT

18/38

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 44

Glu Val Leu Xaa Asp Ala Glu Gly

1

5

<210> 45

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 1 is glt Val. Xaa at position 3 is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 45

Xaa Leu Xaa Asp Ala Gly Phe

1

5

<210> 46

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 46

Gln Val Leu Xaa Asp Ala Gly Phe

19/38

1

5

<210> 47

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 47

Asp Val Leu Xaa Asp Ala Gly Phe

1

5

<210> 48

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 1 is homo Glu. Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 48

Xaa Val Leu Xaa Asp Ala Gly Phe

1

5

<210> 49

<211> 7

<212> PRT

<213> Artificial Sequence

20/38

<220> Xaa at position 1 is β -ODap Val. Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 49

Xaa Leu Xaa Asp Ala Gly Phe

1

5

<210> 50

<211> 13

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 1 is (7-methoxycoumarin-4-yl)acetyl Ser. Xaa at position 12 is (2,4-dinitrophenyl)Arg.

<400> 50

Xaa Glu Val Asn Leu Asp Ala Glu Phe Arg Lys Xaa Arg

1

5

10

<210> 51

<211> 6

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 1 is β -(tetrazole-5-carbonyl)-Dap Val. Xaa at position 3 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 51

21/38

Xaa Leu Xaa Asp Ala Glu

1

5

<210> 52

<211> 6

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 1 is β -(3,5-dicarboxybenzoyl)-Dap Val. Xaa at position 3 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 52

Xaa Leu Xaa Asp Ala Glu

1

5

<210> 53

<211> 6

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 6 is 4-aminobenzoic acid.

<400> 53

Glu Val Leu Xaa Asp Xaa

1

5

<210> 54

<211> 6

22/38

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 6 is 2-aminobenzoic acid.

<400> 54

Glu Val Leu Xaa Asp Xaa

1

5

<210> 55

<211> 6

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 6 is 3-aminobenzoic acid.

<400> 55

Glu Val Leu Xaa Asp Xaa

1

5

<210> 56

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

23/38

<400> 56

Glu Val Leu Xaa Asp Phe Glu Phe

1

5

<210> 57

<211> 6

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 1 is β -(4-carboxybenzoyl)-Dap Val. Xaa at position 3 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 57

Xaa Leu Xaa Asp Ala Glu

1

5

<210> 58

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3R)-3-amino-2-hydroxy-4-phenylthiobutanoic acid.

<400> 58

Glu Val Leu Xaa Asp Ala Glu Phe

1

5

<210> 59

<211> 6

24/38

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 1 is β -(3-carboxybenzoyl)-Dap Val. Xaa at position 3 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 59

Xaa Leu Xaa Asp Ala Glu

1

5

<210> 60

<211> 6

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 1 is α -ODap Val. Xaa at position 3 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 60

Xaa Leu Xaa Asp Ala Glu

1

5

<210> 61

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 6 is tert-Leu.

25/38

<400> 61

Glu Val Leu Xaa Asp Xaa Glu Phe

1

5

<210> 62

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 7 is 2-aminobenzoic acid.

<400> 62

Glu Val Leu Xaa Asp Ala Xaa

1

5

<210> 63

<211> 6

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 6 is 2-piperidinecarboxylic acid.

<400> 63

Glu Val Leu Xaa Asp Xaa

1

5

26/38

<210> 64

<211> 5

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3R)-3-amino-2-hydroxy-4-phenylthiobutanoic acid.

<400> 64

Glu Val Leu Xaa Asp

1

5

<210> 65

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 6 is 2-aminoisobutyric acid.

<400> 65

Glu Val Leu Xaa Asp Xaa Glu

1

5

<210> 66

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic

27/38

acid. Xaa at position 7 is 3-aminobenzoic acid.

<400> 66

Glu Val Leu Xaa Asp Ala Xaa

1

5

<210> 67

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 7 is 4-aminobenzoic acid.

<400> 67

Glu Val Leu Xaa Asp Ala Xaa

1

5

<210> 68

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 7 is 2-piperidinecarboxylic acid.

<400> 68

Glu Val Leu Xaa Asp Ala Xaa

1

5

28/38

<210> 69

<211> 6

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 1 is β -ethoxyoxalyl-Dap Val. Xaa at position 3 is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 69

Xaa Leu Xaa Asp Ala Glu

1 5

<210> 70

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 70

Glu Val Leu Xaa Val Ala Glu Phe

1 5

<210> 71

<211> 8

<212> PRT

<213> Artificial Sequence

29/38

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-5-methylhexanoic acid.

<400> 71

Glu Val Leu Xaa Asp Ala Glu Phe

1

5

<210> 72

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 72

Glu Val Leu Xaa Asp Ala Glu

1

5

<210> 73

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 3 is (S)-2-amino-3-(4-thiazole)propanoic acid.

Xaa at position 4 is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 73

Glu Val Xaa Xaa Asp Ala Glu Phe

1

5

30/38

<210> 74

<211> 5

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 74

Glu Val Leu Xaa Asp

1 5

<210> 75

<211> 6

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 1 is 5-oxalyl-(S)-2,5-diaminopentanoic acid Val.

Xaa at position 3 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 75

Xaa Leu Xaa Asp Ala Glu

1 5

<210> 76

<211> 6

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic

31/38

acid. Xaa at position 6 is 4-aminomethylbenzoic acid.

<400> 76

Glu Val Leu Xaa Asp Xaa

1

5

<210> 77

<211> 5

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-5-methylhexanoic acid.

<400> 77

Glu Val Leu Xaa Asp

1

5

<210> 78

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 5 is (R)-5,5-dimethyl-1,3-thiazolidine-4-carboxylic acid.

<400> 78

Glu Val Leu Xaa Xaa Ala Glu Phe

1

5

32/38

<210> 79

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 79

Glu Val Leu Xaa Leu Ala Glu Phe

1

5

<210> 80

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 7 is 4-aminomethylbenzoic acid.

<400> 80

Glu Val Leu Xaa Asp Ala Xaa

1

5

<210> 81

<211> 7

<212> PRT

<213> Artificial Sequence

33/38

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 7 is 3-piperidinecarboxylic acid.

<400> 81

Glu Val Leu Xaa Asp Ala Xaa

1

5

<210> 82

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 7 is 4-piperidinecarboxylic acid.

<400> 82

Glu Val Leu Xaa Asp Ala Xaa

1

5

<210> 83

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2S,3S)-3-amino-2-hydroxy-5-methylhexanoic acid.

<400> 83

Glu Val Leu Xaa Asp Ala Glu Phe

34/38

1

5

<210> 84

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 7 is 4-aminocyclohexanecarboxylic acid.

<400> 84

Glu Val Leu Xaa Asp Ala Xaa

1

5

<210> 85

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 85

Glu Val Leu Xaa Asp Ala Glu

1

5

<210> 86

<211> 8

<212> PRT

<213> Artificial Sequence

35/38

<220> Xaa is (2R, 3R)-3-amino-2-hydroxy-4-phenylthiobutanoic acid.

<400> 86

Glu Ala Leu Xaa Asp Ala Glu Phe

1

5

<210> 87

<211> 7

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 87

Glu Val Leu Xaa Asp Ala Arg

1

5

<210> 88

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R, 3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 88

Glu Val Leu Xaa Pro Ala Glu Phe

1

5

36/38

<210> 89

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 3 is (2S)-2-amino-2-(3-tetrahydrofuranyl)acetic acid. Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 89

Glu Val Xaa Xaa Asp Ala Glu Phe

1

5

<210> 90

<211> 6

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 90

Glu Val Leu Xaa Asp Ala

1

5

<210> 91

<211> 7

<212> PRT

<213> Artificial Sequence

37/38

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 7 is 4-aminobutanoic acid.

<400> 91

Glu Val Leu Xaa Asp Ala Xaa

1

5

<210> 92

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 4 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid. Xaa at position 5 is (R)-1,3-thiazolidine-4-carboxylic acid.

<400> 92

Glu Val Leu Xaa Xaa Ala Glu Phe

1

5

<210> 93

<211> 5

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 93

Glu Val Leu Xaa Asp

1

5

38/38

<210> 94

<211> 8

<212> PRT

<213> Artificial Sequence

<220> Xaa is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 94

Glu Val Val Xaa Asp Ala Glu Phe

1

5

<210> 95

<211> 6

<212> PRT

<213> Artificial Sequence

<220> Xaa at position 1 is β -(3,4-dihydroxybenzoyl)Dap Val. Xaa at position 3 is (2R,3S)-3-amino-2-hydroxy-4-phenylbutanoic acid.

<400> 95


Xaa Val Xaa Asp Ala Glu

1

5

CLAIMS

1. A compound represented by the general formula (1):



5 wherein A is amino group, or amino group protected with a protecting group which can be degraded in a living body,

B¹ and B² are each a single bond or alkylene of 1 to 3 carbon atoms optionally substituted with R⁴,

10 in which R⁴ is halogen atom; alkyl of 1 to 6 carbon atoms; alkenyl of 2 to 6 carbon atoms; alkynyl of 2 to 6 carbon atoms; cycloalkyl of 3 to 7 carbon atoms; optionally substituted heterocycle; optionally substituted aralkyl; or optionally substituted aryl, and the substituent thereof is alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or aryl,

15 D is a single bond, -NHCO- or NHCO-R⁵-,


20 in which R⁵ is alkylene of 1 to 6 carbon atoms; alkenylene of 2 to 6 carbon atoms; alkynylene of 2 to 6 carbon atoms; cycloalkylene of 3 to 7 carbon atoms; optionally substituted heterocycle; optionally substituted aralkylene; or optionally substituted arylene, and the substituent thereof is halogen atom, alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or aryl,

E is -COOH, a group equivalent to COOH, hydrogen atom, hydroxy group, amino group, halogen atom, or -COOCH₃,

n is an integer of 1 to 3,

25 m is an integer of 1 to 3,

G is hydroxy group, amino acid residue, a peptide constructed of 4 to 2 amino acids, or a group represented by the formula




in which Z is -NH-, -Asp-Ala-NH-, -Asp-Ala-, -NH-CZ¹(Z²)-, -Asp-NH-, -Asp-NH-CZ¹(Z²)-, or -NH-CZ¹(Z²)-CZ¹(Z²)-,

Ring L is a 5- to 10-membered ring optionally containing a heteroatom and/or unsaturated bond,

Z¹ and Z² are each hydrogen atom, hydroxy group, amino group, cyano group, nitro group, alkyl of 1 to 6 carbon atoms, or halogen atom,

X is hydrogen atom, -COOH, a group equivalent to COOH, -CONHR⁶, -OCH₂COOR⁶, -OCH₂CONHR⁶, hydroxy group, -PO₄H, -SO₃H, -CZ¹(Z²)-X' or a group represented by the formula:



in which X' is hydrogen atom, -COOH, a group equivalent to COOH, -CONHR⁶, -OCH₂COOR⁶, -OCH₂CONHR⁶, hydroxy group, -PO₄H, or -SO₃H,

Y and Y' are each hydrogen atom, -COOH, a group equivalent to COOH, -CONHR⁶, -OR⁶, -PO₄H, -SO₃H, halogen atom, or methyl,

in which R⁶ is an alkyl of 1 to 4 carbon atoms optionally substituted with halogen atom, or hydrogen atom, and

k is an integer of 1 to 4, and

R¹, R² and R³ are each -COOH, a group equivalent to COOH, alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or alkyl of 1 to 6 carbon atoms substituted with any of phenyl, phenylthio, alkylthio of 1 to 6 carbon atoms, -COOH or heterocycle, provided that the phenyl, phenylthio and heterocycle may be substituted with an alkyl of 1 to 6 carbon atoms, hydroxy group, nitro, a halogen atom, -SO₃H or -PO₄H or a pharmaceutically acceptable salt thereof, or a prodrug thereof.

2. The compound of claim 1, wherein in the general formula (1), B¹ and B² are each a single bond, or alkylene of 1 to 3 carbon

atoms optionally substituted with R⁴,

in which R⁴ is halogen atom; alkyl of 1 to 6 carbon atoms; alkenyl of 2 to 6 carbon atoms; alkynyl of 2 to 6 carbon atoms; cycloalkyl of 3 to 7 carbon atoms; optionally substituted heterocycle; optionally substituted aralkyl; or optionally substituted aryl, and the substituent thereof is alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or aryl,

D is a single bond, -NHCO- or -NHCO-R⁵-,


in which R⁵ is alkylene of 1 to 6 carbon atoms; alkenylene of 2 to 6 carbon atoms; alkynylene of 2 to 6 carbon atoms; cycloalkylene of 3 to 7 carbon atoms; optionally substituted heterocycle; optionally substituted aralkylene; or optionally substituted arylene, and the substituent thereof is halogen atom, alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or aryl,

E is -COOH or a group equivalent to COOH, hydrogen atom, hydroxy group, amino group, halogen atom, or -COOCH₃,

n is an integer of 1 to 3,

m is 1,

G is hydroxy group, or a group represented by the formula:




in which Z is -NH-, -Asp-Ala-NH-, -Asp-Ala-, -NH-CZ¹(Z²)-, -Asp-NH-, -Asp-NH-CZ¹(Z²)-, or -NH-CZ¹(Z²)-CZ¹(Z²)-,

Ring L is a 5- to 10-membered ring optionally containing a heteroatom and/or unsaturated bond,

Z¹ and Z² are each hydrogen atom, hydroxy group, amino group, cyano group, nitro group, alkyl of 1 to 6 carbon atoms, or halogen atom,

X is hydrogen atom, -COOH, a group equivalent to COOH, -CONHR⁶, -OCH₂COOR⁶, -OCH₂CONHR⁶, hydroxy group, -PO₄H, -SO₃H, -CZ¹(Z²)-X' or a group represented by the formula:



where X' is hydrogen atom, -COOH, a group equivalent to COOH, -CONHR⁶, -OCH₂COOR⁶, -OCH₂CONHR⁶, hydroxy group, -PO₄H, or -SO₃H,

Y and Y' are each hydrogen atom, a group equivalent to COOH, -CONHR⁶, -OR⁶, -PO₄H, -SO₃H, halogen atom, or methyl,

in which R⁶ is alkyl of 1 to 4 carbon atoms optionally substituted with halogen atom, or hydrogen atom,

k is an integer of 1 to 4, and


R¹, R² and R³ are each alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or alkyl of 1 to 6 carbon atoms substituted with any of phenyl, phenylthio, alkylthio of 1 to 6 carbon atoms, or heterocycle, provided that the phenyl, phenylthio and heterocycle may be substituted with alkyl of 1 to 6 carbon atoms, hydroxy group, nitro, halogen atom, -SO₃H or -PO₄H, or a pharmaceutically acceptable salt thereof, or a prodrug thereof.

3. The compound of claim 1, wherein in the general formula (1),


B¹ is a single bond or methylene,

B² is a single bond or alkylene of 1 to 3 carbon atoms,


D is a single bond, -NHCO-, a group represented by the formula:



in which the benzene ring may be substituted with halogen atom, amino group, alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or aryl, a group represented by the following formula:



a group represented by the following formula:



in which Q is oxygen atom or NH,

E is hydrogen atom, hydroxy group, amino group, halogen atom, -COOH, -COOCH₃ or tetrazole ring,


5 n is an integer of 1 or 2,

m is 1,

R¹ is alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, alkynyl of 2 to 6 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or aryl,

10 R² is alkyl of 1 to 6 carbon atoms, phenyl, naphthyl, isobutyl, 2-carbamoyl ethyl, 2-methylthioethyl, methylthiomethyl, 2-ethylthioethyl, ethylthiomethyl, naphthylmethyl, phenethyl, 4-thiazolylmethyl, 4-imidazolylmethyl, 3-tetrahydrofuranyl, 3-indolylmethyl, cyclohexylmethyl, or benzyl in which the benzene ring may be substituted with halogen atom, hydroxy group, or alkyloxy group of 1 to 6 carbon atoms, and

15 G is any one of the group represented by the following formulas:



in which X and X' are each hydrogen atom, -COOH, -CONHR⁶, -OCH₂COOR⁶, -OCH₂CONHR⁶, hydroxy group, -PO₄H, -SO₃H or tetrazole ring,

Y and Y' are each hydrogen atom, -COOH, -CONHR⁶, -OR⁶, -PO₄H, -SO₃H, halogen atom, methyl or tetrazole ring,

R⁶ is alkyl of 1 to 2 carbon atoms optionally substituted with halogen atom, or hydrogen atom,

5 k is an integer of 1 or 2, and


Z¹ and Z² are each hydrogen, hydroxy group, amino group, cyano group, nitro group, alkyl of 1 to 6 carbon atoms, or halogen atom, or a pharmaceutically acceptable salt thereof, or a prodrug thereof.

4. The compound of claim 1, wherein in the general formula (1),

10 B¹ is a single bond,

B² is alkylene of 1 to 2 carbon atoms,

D is a single bond, -NHCO-, or a group represented by the following formula:



15 E is hydrogen atom, hydroxy group, halogen atom, -COOH or tetrazole ring,

n is an integer of 1 or 2,


m is 1,

R¹ is isopropyl,

20 R² is isobutyl or cyclohexylmethyl,

R³ is benzyl, isobutyl, phenylthiomethyl or cyclohexylmethyl, and

G is any one of the group represented by the following formulas:



in which X and X' are each -COOH, -CONH₂, hydroxy group, -OCH₂COOH or tetrazole ring,


Y and Y' are each hydrogen atom, COOH, CONH₂, halogen atom, methyl or tetrazole ring,

k is an integer of 1 or 2, and

Z¹ and Z² are each hydrogen atom,


or a pharmaceutically acceptable salt thereof, or a prodrug thereof.

5. The compound of claim 1, wherein in the general formula (1), a partial structure other than G is a group represented by the formula:




R² is isobutyl or cyclohexylmethyl,

R⁷ is a group represented by the following formula:



and

G is a group represented by the following formula:



- 5 or a pharmaceutically acceptable salt thereof, or a prodrug thereof.
6. The compound of any one of claims 1 to 5, which has a molecular weight of 1100 or lower.
7. A β -secretase inhibitor comprising the compound of any one of claims 1 to 5 as an active ingredient.
- 10 8. An agent for preventing or treating (i) neurodegenerative disease, (ii) neuropathy in cerebrovascular disease, head trauma·spinal damage, encephalitis sequela, or cerebral palsy, (iii) memory disorder, or (iv) mental disease associated with β -secretase, comprising the compound of any one of claims 1 to 5 as an active ingredient.
- 15 9. An agent for preventing or treating (i) neurodegenerative disease, (ii) neuropathy in cerebrovascular disease, head trauma·spinal damage, encephalitis sequela, or cerebral palsy, (iii) memory disorder, or (iv) mental

disease by promoting secretion of sAPP α , or promoting secretion of sAPP α , and inhibiting production·secretion of β -amyloid protein, or inhibiting production·secretion of β -amyloid protein, comprising the compound of any one of claims 1 to 5 as an active ingredient.

5 10. The preventing or treating agent of claim 7 or 8, wherein the neurodegenerative disease is Alzheimer's disease or Parkinson's disease.

11. An agent for promoting secretion of sAPP α , comprising the compound of any one of claims 1 to 5 as an active ingredient.

10 12. An agent for inhibiting production·secretion of β -amyloid protein, comprising the compound of any one of claims 1 to 5 as an active ingredient.

13. A neurotrophic factor-like acting agent, comprising the compound of any one of claims 1 to 5 as an active ingredient.

15 14. An agent for preventing or treating neuropathy or mental disease in head trauma·spinal damage, encephalitis sequela, or cerebral palsy, comprising the compound of any one of claims 1 to 5 as an active ingredient.

20 15. Use of the compound of any one of claims 1 to 4 for preparing an agent for promoting secretion of sAPP α , a neurotrophic factor-like acting agent, or an agent for preventing or treating neuropathy, memory disorder or mental disease in head trauma·spinal damage, encephalitis sequela, or cerebral palsy.

25 16. A method for treating (i) neurodegenerative disease, (ii) neuropathy in cerebrovascular disease, head trauma·spinal damage, encephalitis sequela, or cerebral palsy, (iii) memory disorder or (iv) mental disease associated with β -secretase, which comprises administering an effective amount of the compound of any one of claims 1 to 5 to a mammal.

