



(51) International Patent Classification:

C07D 235/18 (2006.01) C07D 263/57 (2006.01)
 C07D 277/68 (2006.01) A01N 43/52 (2006.01)
 C07D 471/04 (2006.01) A01N 43/76 (2006.01)
 C07D 498/04 (2006.01) A01N 43/78 (2006.01)
 C07D 513/04 (2006.01) A01N 43/90 (2006.01)

(21) International Application Number:

PCT/JP2011/080557

(22) International Filing Date:

22 December 2011 (22.12.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

2010-287412 24 December 2010 (24.12.2010) JP
 2011-166768 29 July 2011 (29.07.2011) JP
 2011-263374 1 December 2011 (01.12.2011) JP

(71) Applicant (for all designated States except US): SUMITOMO CHEMICAL COMPANY, LIMITED [JP/JP];

27-1, Shinkawa 2-chome, Chuo-ku, Tokyo, 1048260 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): TAKYO, Hayato [JP/JP]; c/o SUMITOMO CHEMICAL COMPANY, LIMITED, 27-1, Shinkawa 2-chome, Chuo-ku, Tokyo, 1048260 (JP). TAKAHASHI, Masaki [JP/JP]; c/o SUMITOMO CHEMICAL COMPANY, LIMITED, 2-1, Takatsukasa 4-chome, Takarazuka-shi, Hyogo, 6658555 (JP). TANABE, Takamasa [JP/JP]; c/o SUMITOMO CHEMICAL COMPANY, LIMITED, 2-1, Takatsukasa 4-chome, Takarazuka-shi, Hyogo, 6658555 (JP). NOKURA, Yoshihiko [JP/JP]; c/o SUMITOMO CHEMICAL COMPANY, LIMITED, 2-1, Takatsukasa 4-chome, Takarazuka-shi, Hy-

ogo, 6658555 (JP). ITO, Mai [JP/JP]; c/o SUMITOMO CHEMICAL COMPANY, LIMITED, 2-1, Takatsukasa 4-chome, Takarazuka-shi, Hyogo, 6658555 (JP). IWATA, Atsushi [JP/JP]; c/o SUMITOMO CHEMICAL COMPANY, LIMITED, 2-1, Takatsukasa 4-chome, Takarazuka-shi, Hyogo, 6658555 (JP).

(74) Agents: TANAKA, Mitsuo et al.; AOYAMA & PARTNERS, IMP Building, 3-7, Shiromi 1-chome, Chuo-ku, Osaka-shi, Osaka, 5400001 (JP).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: FUSED HETEROCYCLIC COMPOUND AND USE FOR PEST CONTROL THEREOF



(57) Abstract: A fused heterocyclic compound the formula (1): wherein A¹ represents -NR⁸-, and the like; A² represents a nitrogen atom, and the like; A³ represents a nitrogen atom, and the like; R¹ represents a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from Group X, and the like; R², R³, R⁴, and R⁵ are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, and the like; R⁶ and R⁷ are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from Group X, and the like; R⁸ represents a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from Group W, and the like; n represents 0, 1 or 2. The compound has an excellent activity of controlling pests.

DESCRIPTION

FUSED HETEROCYCLIC COMPOUND AND USE FOR PEST CONTROL
THEREOF

5 Technical Field

[0001]

The present invention relates to a certain fused heterocyclic compound and use for pest control thereof.

10 Background Art

[0002]

For controlling pests, various compounds have heretofore been developed and used practically.

Certain fused heterocyclic compounds are known (see,
15 Patent Literature 1).

Citation List

[0003]

Patent Literature

20 Patent Literature 1: JP-A-01-501473

Summary of Invention

Technical Problem

[0004]

25 An object of the present invention is to provide a

compound having an excellent activity of controlling pests and a method of controlling pests using the compound.

Solution to Problem


5 [0005]

The present inventors have studied so as to resolve the above problem and found that a fused heterocyclic compound of the following formula (1) has an excellent activity of controlling pests, thus leading to the present invention.

[0006]

That is, the present invention provides:

[1] A fused heterocyclic compound of the formula (1):



15 wherein:

A¹ represents -NR⁸-, an oxygen atom, or a sulfur atom;

A² represents a nitrogen atom or =CR⁹-;

A³ represents a nitrogen atom or =CR¹⁰-;

R¹ represents a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from Group X, or a C3-C6 alicyclic hydrocarbon group optionally having one or more atoms or groups selected from Group Y;

R^2 , R^3 , R^4 , and R^5 are same or different and independently represent a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally having one or more atoms or groups selected from
5 Group Z, a 5-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z, a 6-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z, $-OR^{11}$, $-S(O)_mR^{11}$, $-NR^{11}R^{12}$, $-CO_2R^{11}$, $-C(O)R^{11}$, a cyano group, a nitro group, a
10 halogen atom, $-SF_5$, or a hydrogen atom, provided that at least two of R^2 , R^3 , R^4 , and R^5 represent a hydrogen atom;

R^6 and R^7 are same or different and independently represent a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from Group X, a phenyl
15 group optionally having one or more atoms or groups selected from Group Z, a 5-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z, a 6-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z, $-OR^{11}$, $-S(O)_mR^{11}$, $-S(O)_2NR^{11}R^{12}$, $-NR^{11}R^{12}$, $-NR^{11}CO_2R^{12}$, $-NR^{11}C(O)R^{12}$, $-CO_2R^{11}$, $-C(O)R^{11}$, a cyano group, a nitro group, a halogen
20 atom, $-SF_5$, or a hydrogen atom;

R^8 represents a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from
25 Group W, $-CO_2R^{11}$, $-C(O)R^{11}$, a C3-C6 alicyclic hydrocarbon

group optionally having one or more atoms or groups selected from Group Y, or a hydrogen atom;

R^9 and R^{10} are same or different and independently represent a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, $-NR^{11}R^{12}$, $-CO_2R^{11}$, $-C(O)R^{11}$, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

R^{11} and R^{12} are same or different and independently represent a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a hydrogen atom; and

m represents 0, 1, or 2; and n represents 0, 1, or 2;

(except in cases as follows: both R^6 and R^7 are a hydrogen atom; in $-S(O)_mR^{11}$, when m is 1 or 2, R^{11} is a hydrogen atom; and when A^1 represents $-NH-$ or $-N(CH_3)-$, A^2 and A^3 represent $=CH-$, R^1 represents a methyl group, R^2 , R^3 , R^4 , and R^5 represent a hydrogen atom, and n is 0, 1, or 2, R^6 and R^7 are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from Group X, a phenyl group optionally having one or more atoms or groups selected from Group Z, a 5-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z, a 6-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z, $-OR^{11}$, $-S(O)_mR^{11}$, $-S(O)_2NR^{11}R^{12}$, $-NR^{11}R^{12}$, $-NR^{11}-CO_2R^{12}$, $-NR^{11}C(O)R^{12}$, $-CO_2H$, -

C(O)R¹¹, a cyano group, a nitro group, a bromine atom, an iodine atom, -SF₅ or a hydrogen atom)

the Group X consists of a C1-C6 alkoxy group optionally having one or more halogen atoms, a C2-C6 alkenyloxy group optionally having one or more halogen atoms, a C2-C6 alkynyloxy group optionally having one or more halogen atoms, a C1-C6 alkylsulfanyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfinyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfonyl group optionally having one or more halogen atoms, a C2-C6 alkylcarbonyl group optionally having one or more halogen atoms, a C2-C6 alkoxycarbonyl group optionally having one or more halogen atoms, a C3-C6 cycloalkyl group optionally having one or more halogen atoms, a cyano group, a hydroxy group, and a halogen atom;

the Group Y consists of a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 alkoxy group optionally having one or more halogen atoms, a C2-C6 alkenyloxy group optionally having one or more halogen atoms, a C2-C6 alkynyloxy group optionally having one or more halogen atoms, a hydroxy group, and a halogen atom;

the Group Z consists of a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 alkoxy group optionally having one or more halogen atoms, a

C1-C6 alkylsulfanyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfinyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfonyl group optionally having one or more halogen atoms, a C2-C6
5 alkylcarbonyl group optionally having one or more halogen atoms, a C2-C6 alkoxy carbonyl group optionally having one or more halogen atoms, a C1-C6 alkylamino group optionally having one or more halogen atoms, a C2-C8 dialkylamino group optionally having one or more halogen atoms, a
10 halogen atom, a cyano group, and a nitro group; and

the Group W consists of a C1-C6 alkoxy group optionally having one or more halogen atoms, a C2-C6 alkenyloxy group optionally having one or more halogen atoms, a C2-C6 alkynyloxy group optionally having one or
15 more halogen atoms, a C1-C6 alkylsulfanyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfinyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfonyl group optionally having one or more halogen atoms, a C2-C6 alkylcarbonyl group optionally having one or
20 more halogen atoms, a C2-C6 alkoxy carbonyl group optionally having one or more halogen atoms, a C3-C6 cycloalkyl group optionally having one or more halogen atoms, a halogen atom, a cyano group, and a hydroxy group (hereinafter referred to as the present compound);

25 [2] The fused heterocyclic compound according to the above

[1] wherein R^6 and R^7 are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, $-NR^{11}R^{12}$, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

[3] The fused heterocyclic compound according to the above [1] or [2] wherein R^6 and R^7 are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, a halogen atom, or a hydrogen atom;

[4] The fused heterocyclic compound according to any one of the above [1]-[3] wherein A^2 is $=CH-$;

[5] The fused heterocyclic compound according to any one of the above [1]-[4] wherein A^3 is a nitrogen atom or $=CR^{10}-$, and R^{10} is a halogen atom or a hydrogen atom;

[6] The fused heterocyclic compound according to any one of the above [1]-[4] wherein A^3 is a nitrogen atom;

[7] The fused heterocyclic compound according to any one of the above [1]-[6] wherein R^2 , R^4 , and R^5 are same or different and are independently a hydrogen atom or a halogen atom, and R^3 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, a halogen atom, or a hydrogen atom;

[8] The fused heterocyclic compound according to any one of the above [1]-[7] wherein A^1 is $-NR^8-$, and R^8 is a C1-C6


chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a cyclopropyl group;

5 [9] The fused heterocyclic compound according to any one of the above [1]-[7] wherein A^1 is $-NR^8-$, and R^8 is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group;

10 [10] The fused heterocyclic compound according to any one of the above [1]-[7] wherein A^1 is an oxygen atom;

[11] The fused heterocyclic compound according to any one of the above [1]-[7] wherein A^1 is a sulfur atom;

[12] The fused heterocyclic compound according to the above [1] which is represented by the formula (1-1):



15

wherein:

A^{1a} represents $-NR^{8a}-$ or a sulfur atom;

A^{3a} represents a nitrogen atom or $=CR^{10a}-$;

20 R^{1a} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

R^{2a} , R^{4a} , and R^{5a} are same or different and independently represent a C1-C6 chain hydrocarbon group

optionally having one or more halogen atoms, a halogen atom, or a hydrogen atom;

R^{3a} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11a}$, a
5 halogen atom, or a hydrogen atom;

R^{6a} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11a}$, $-S(O)_mR^{11a}$, a bromine atom, or an iodine atom;

R^{8a} represents a C1-C6 chain hydrocarbon group
10 optionally having one or more halogen atoms or a hydrogen atom;

R^{10a} represents a halogen atom or a hydrogen atom;

R^{11a} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms; and

15 m represents 0, 1, or 2, and n represents 0, 1, or 2;

[13] The fused heterocyclic compound according to the above

[12] wherein A^{1a} is $-NR^{8a}-$ or a sulfur atom, R^{8a} is a methyl group, A^{3a} is a nitrogen atom, R^{1a} is an ethyl group, R^{2a} , R^{4a} , and R^{5a} are same or different and are independently a

20 halogen atom or a hydrogen atom, R^{3a} is a trifluoromethyl group, a halogen atom, or a hydrogen atom, and R^{6a} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a
25 C1-C3 alkylsulfinyl group having one or more fluorine atoms,

or a C1-C3 alkylsulfonyl group having one or more fluorine atoms;

[14] A pest controlling composition which comprises the fused heterocyclic compound according to any one of the above [1]-[13] and an inert carrier; and

[15] A method of controlling pests which comprises applying an effective amount of the fused heterocyclic compound according to any one of the above [1]-[13] to pests or habitats of pests.

10

Effect of Invention

[0007]

The present compound has an excellent activity of controlling pests and is useful as an active ingredient of a pest controlling agent.

15

Description of Embodiments

[0008]

Various substituents used in the present specification will be illustrated by way of examples.

20

In the present invention, the "halogen atom" includes a fluorine atom, a chlorine atom, a bromine atom, and an iodine atom.

[0009]

25

In the present compound, examples of the "C1-C6 chain

hydrocarbon group optionally having one or more atoms or groups selected from Group X" include a C1-C6 alkyl group optionally having one or more atoms or groups selected from Group X such as a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, a neopentyl group, a hexyl group, a methoxymethyl group, an ethoxymethyl group, a propyloxymethyl group, an isopropyloxymethyl group, a butyloxymethyl group, a sec-butyloxymethyl group, a tert-butyloxymethyl group, a 2-methoxyethyl group, a 2-ethoxyethyl group, a 2-propyloxyethyl group, a 2-isopropyloxyethyl group, a 2-butyloxyethyl group, a 2-sec-butyloxyethyl group, a 2-tert-butyloxyethyl group, a trifluoromethyl group, a trichloromethyl group, a 2-fluoroethyl group, a 2,2-difluoroethyl group, a 2,2,2-trifluoroethyl group, a pentafluoroethyl group, a methylsulfanylethyl group, an ethylsulfanylethyl group, a methylsulfinylethyl group, and a methylsulfonylethyl group, a methoxycarbonylethyl group, a cyanomethyl group, a cyclopropylmethyl group, and a cyclobutylmethyl group;

a C2-C6 alkenyl group optionally having one or more atoms or groups selected from Group X such as a vinyl group, a 1-propenyl group, a 2-propenyl group, a 1-methylvinyl group, a 2-methyl-1-propenyl group, a 1-butenyl group, a 2-butenyl

group, a 3-butenyl group, a 1-pentenyl group, a 1-hexenyl group, a 1,1-difluoroallyl group, and a pentafluoroallyl group; and

a C2-C6 alkynyl group optionally having one or more atoms or groups selected from Group X such as an ethynyl group, a propargyl group, a 2-butyngyl group, a 3-butyngyl group, a 1-pentyngyl group, a 1-hexyngyl group, and a 4,4,4-trifluoro-2-butyngyl group.

[0010]

In the present compound, examples of the "C3-C6 alicyclic hydrocarbon group optionally having one or more atoms or groups selected from Group Y" include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, a 1-cyclohexenyl group, a 2-cyclohexenyl group, a 3-cyclohexenyl group, a 1-methylcyclohexyl group, a 2-methylcyclohexyl group, a 3-methylcyclohexyl group, a 4-methylcyclohexyl group, a 2-methoxycyclohexyl group, a 3-methoxycyclohexyl group, a 4-methoxycyclohexyl group, a 1-fluorocyclohexyl group, a 2-fluorocyclohexyl group, a 3-fluorocyclohexyl group, and a 4-fluorocyclohexyl group.

[0011]

In the present compound, examples of the "C1-C6 chain hydrocarbon group optionally having one or more halogen atoms" include a C1-C6 alkyl group optionally having one or

more halogen atoms such as a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, a neopentyl group, a hexyl group, a trifluoromethyl group, a trichloromethyl group, a 2-fluoroethyl group, a 2,2-difluoroethyl group, a 2,2,2-trifluoroethyl group, a pentafluoroethyl group, and a heptafluoroisopropyl group;

a C2-C6 alkenyl group optionally having one or more halogen atoms such as a vinyl group, a 1-propenyl group, a 2-propenyl group, a 1-methylvinyl group, a 2-methyl-1-propenyl group, a 1-butenyl group, a 2-butenyl group, a 3-butenyl group, a 1-pentenyl group, a 1-hexenyl group, a 1,1-difluoroallyl group, and a pentafluoroallyl group; and a C2-C6 alkynyl group optionally having one or more halogen atoms such as an ethynyl group, a propargyl group, a 2-butyne group, a 3-butyne group, a 1-pentyne group, a 1-hexyne group, and a 4,4,4-trifluoro-2-butyne group.

[0012]

In the present compound, examples of the "phenyl group optionally having one or more atoms or groups selected from Group Z" include a phenyl group, a 2-fluorophenyl group, a 3-fluorophenyl group, a 4-fluorophenyl group, a 2,3-difluorophenyl group, a 2,4-difluorophenyl group, a 2,5-difluorophenyl group, a 2,6-difluorophenyl group, a 3,4-

difluorophenyl group, a 3,5-difluorophenyl group, a
2,3,4,5,6-pentafluorophenyl group, a 2-chlorophenyl group,
a 3-chlorophenyl group, a 4-chlorophenyl group, a 2-
bromophenyl group, a 3-bromophenyl group, a 4-bromophenyl
5 group, a 2-iodophenyl group, a 3-iodophenyl group, a 4-
iodophenyl group, a 2-trifluoromethylphenyl group, a 3-
trifluoromethylphenyl group, a 4-trifluoromethylphenyl
group, a 2-trifluoromethoxyphenyl group, a 3-
trifluoromethoxyphenyl group, a 4-trifluoromethoxyphenyl
10 group, a 2-trifluoromethylsulfanylphenyl group, a 3-
trifluoromethylsulfanylphenyl group, a 4-
trifluoromethylsulfanylphenyl group, a 4-
methoxycarbonylphenyl group, a 4-nitrophenyl group, a 4-
cyanophenyl group, a 4-methylaminophenyl group, a 4-
15 dimethylaminophenyl group, a 4-methylsulfinylphenyl group,
a 4-methylsulfonylphenyl group, a 4-acetylphenyl group, and
4-methoxycarbonylphenyl group.

[0013]

In the present compound, the "heterocyclic group" in
20 the "5-membered heterocyclic group optionally having one or
more atoms or groups selected from Group Z" represents a
heterocyclic residue. Examples of "5-membered
heterocyclic group optionally having one or more atoms or
groups selected from Group Z" include a 5-membered
25 saturated heterocyclic group optionally having one or more

atoms or groups selected from Group Z such as a pyrrolidin-1-yl group and a tetrahydrofuran-2-yl group; and
a 5-membered unsaturated heterocyclic group optionally having one or more atoms or groups selected from Group Z
5 such as a 2-pyrrolyl group, a 2-furyl group, a 3-furyl group, a 5-pyrazolyl group, a 4-pyrazolyl group, a 1-pyrrolyl group, a 1-methyl-2-pyrrolyl group, a 5-bromo-2-furyl group, a 5-nitro-2-furyl group, a 2-methyl-3-furyl group, a 2,5-dimethyl-3-furyl group, a 2,4-dimethyl-3-furyl group, a 5-methyl-2-thienyl group, a 3-methyl-2-thienyl group, a 1-methyl-3-trifluoromethyl-5-pyrazolyl group, a 5-chloro-1,3-dimethyl-4-pyrazolyl group, a pyrazol-1-yl group, a 3-chloro-pyrazol-1-yl group, a 3-bromopyrazol-1-yl group, a 4-chloropyrazol-1-yl group, a 4-bromopyrazol-1-yl group,
10 a imidazol-1-yl group, a pyrrol-1-yl group, a 1,2,4-triazol-1-yl group, a 3-chloro-1,2,4-triazol-1-yl group, a 1,2,3,4-tetrazol-1-yl group, a 1,2,3,5-tetrazol-1-yl group, a 2-thienyl group, a 3-thienyl group, a 3-trifluoromethyl-1,2,4-triazol-1-yl group, and a 4-trifluoromethylpyrazol-1-yl group.
20 yl group.

[0014]

In the present compound, the "heterocyclic group" in the "6-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z" represents a
25 heterocyclic residue. Examples of the "6-membered

heterocyclic group optionally having one or more atoms or groups selected from Group Z" include a 6-membered saturated heterocyclic group optionally having one or more atoms or groups selected from Group Z such as a piperidyl group, a morpholinyl group, and a thiomorpholinyl group; and

a 6-membered aromatic heterocyclic group optionally having one or more atoms or groups selected from Group Z such as a pyrazinyl group, a 2-pyrimidyl group, a 4-pyrimidyl group, a 5-pyrimidyl group, a 2-pyridyl group, a 3-pyridyl group, a 4-pyridyl group, a 3-fluoro-2-pyridyl group, a 4-fluoro-2-pyridyl group, a 5-fluoro-2-pyridyl group, a 6-fluoro-2-pyridyl group, a 2-pyrimidyl group, and a 5-trifluoromethylpyridin-2-yl group.

[0015]

In the present compound, the "C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from Group W" include a C1-C6 alkyl group optionally having one or more atoms or groups selected from Group W such as a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, a neopentyl group, a hexyl group, a trifluoromethyl group, a trichloromethyl group, a 2-fluoroethyl group, a 2,2-difluoroethyl group, a 2,2,2-trifluoroethyl group, a

pentafluoroethyl group, a methoxymethyl group, an ethoxymethyl group, a propyloxymethyl group, an isopropyloxymethyl group, a butyloxymethyl group, a sec-butyloxymethyl group, an isobutyloxymethyl group, a tert-butyloxymethyl group, a methoxyethyl group, an ethoxyethyl group, a propyloxyethyl group, an isopropyloxyethyl group, a butyloxyethyl group, a sec-butyloxyethyl group, an isobutyloxyethyl group, a tert-butyloxyethyl group, a methylsulfanylethyl group, an ethylsulfanylethyl group, a methylsulfinylethyl group, a methylsulfonylethyl group, a methoxycarbonylmethyl group, a methoxycarbonylethyl group, a cyclopropylmethyl group, and a cyclohexylmethyl group; a C2-C6 alkenyl group optionally having one or more atoms or groups selected from Group W such as a vinyl group, a 1-propenyl group, a 2-propenyl group, a 1-methylvinyl group, a 2-methyl-1-propenyl group, a 1-butenyl group, a 2-butenyl group, a 3-butenyl group, a 1-pentenyl group, a 1-hexenyl group, a 1,1-difluoroallyl group, and a pentafluoroallyl group; and

a C2-C6 alkynyl group optionally having one or more atoms or groups selected from Group W such as an ethynyl group, a propargyl group, a 2-butylnyl group, a 3-butylnyl group, a 1-pentylnyl group, a 1-hexynyl group, and a 4,4,4-trifluoro-2-butylnyl group.

In the present compound, examples of the "C1-C6 alkylsulfanyl group optionally having one or more halogen atoms" include a methylsulfanyl group, an ethylsulfanyl group, a propylsulfanyl group, an isopropylsulfanyl group, a butylsulfanyl group, a pentylsulfanyl group, a hexylsulfanyl group, a trifluoromethylsulfanyl group, a 2,2,2-trifluoroethylsulfanyl group, and a pentafluoroethylsulfanyl group.

[0017]

In the present compound, examples of the "C1-C6 alkylsulfinyl group optionally having one or more halogen atoms" include a methylsulfinyl group, an ethylsulfinyl group, a propylsulfinyl group, an isopropylsulfinyl group, a butylsulfinyl group, a pentylsulfinyl group, a hexylsulfinyl group, a trifluoromethylsulfinyl group, a 2,2,2-trifluoroethylsulfinyl group, and a pentafluoroethylsulfinyl group.

[0018]

In the present compound, examples of the "C1-C6 alkylsulfonyl group optionally having one or more halogen atoms" include a methylsulfonyl group, an ethylsulfonyl group, a propylsulfonyl group, an isopropylsulfonyl group, a butylsulfonyl group, a pentylsulfonyl group, a hexylsulfonyl group, a trifluoromethylsulfonyl group, a 2,2,2-trifluoroethylsulfonyl group, and a

pentafluoroethylsulfonyl group.

[0019]

In the present compound, examples of the "C1-C6 alkoxy group optionally having one or more halogen atoms" include
5 a methoxy group, a trifluoromethoxy group, an ethoxy group, a 2,2,2-trifluoroethoxy group, a propyloxy group, an isopropyloxy group, a butyloxy group, an isobutyloxy group, a sec-butyloxy group, a tert-butyloxy group, a pentyloxy group, and a hexyloxy group.

10 [0020]

In the present compound, examples of the "C2-C6 alkenyloxy group optionally having one or more halogen atoms" include a 2-propenyloxy group, a 2-methyl-2-propenyloxy group, a 2-butenyloxy group, a 3-butenyloxy
15 group, a 2-pentenyloxy group, a 2-hexenyloxy group, a 3,3-difluoroallyloxy group, and a 3,3-dichloroallyloxy group.

[0021]

In the present compound, examples of the "C2-C6 alkynyloxy group optionally having one or more halogen
20 atoms" include a propargyloxy group, a 2-butyloxy group, a 3-butyloxy group, a 2-pentyloxy group, a 2-hexyloxy group, and a 4,4,4-trifluoro-2-butyloxy group.

[0022]

In the present compound, examples of the "C2-C6
25 alkylcarbonyl group optionally having one or more halogen

atoms" include an acetyl group, a propionyl group, a propylcarbonyl group, a butylcarbonyl group, a pentylcarbonyl group, and a trifluoroacetyl group.

[0023]

5 In the present compound, examples of the "C2-C6 alkoxy carbonyl group optionally having one or more halogen atoms" include a methoxycarbonyl group, an ethoxycarbonyl group, a propyloxycarbonyl group, a butyloxycarbonyl group, a pentyloxycarbonyl group, a tert-butyloxycarbonyl group, and a 2,2,2-trifluoroethylcarbonyl group.

[0024]

In the present compound, examples of the "C1-C6 alkylamino group optionally having one or more halogen atoms" include a methylamino group, an ethylamino group, a 2,2,2-trifluoroethylamino group, a propylamino group, an isopropylamino group, and a butylamino group.

[0025]

In the present compound, examples of the "C2-C8 dialkylamino group optionally having one or more halogen atoms" include a dimethylamino group, a diethylamino group, a bis(2,2,2-trifluoroethyl)amino group, and a dipropylamino group.

[0026]

In the present compound, examples of the "C3-C6 cycloalkyl group optionally having one or more halogen

atoms" include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, and a cyclohexyl group.

[0027]

In the present compound, examples of the "C1-C3 alkyl group having one or more a fluorine atoms" include a fluoromethyl group, a difluoromethyl group, a trifluoromethyl group, a trichloromethyl group, a 2,2,2-trifluoroethyl group, a pentafluoroethyl group, a heptafluoropropyl group, and a heptafluoroisopropyl group.

[0028]

In the present compound, examples of the "C1-C3 alkoxy group having one or more a fluorine atoms" include a trifluoromethoxy group, a 2,2,2-trifluoroethoxy group, and a pentafluoroethoxy group.

[0029]

In the present compound, examples of the "C1-C3 alkylsulfanyl group having one or more a fluorine atoms" include a trifluoromethylsulfanyl group, a 2,2,2-trifluoroethylsulfanyl group, a pentafluoroethylsulfanyl group, a heptafluoropropylsulfanyl group, and a heptafluoroisopropylsulfanyl group.

[0030]

In the present compound, examples of the "C1-C3 alkylsulfinyl group having one or more a fluorine atoms" include a trifluoromethylsulfinyl group, a 2,2,2-

trifluoroethylsulfinyl group, a pentafluoroethylsulfinyl group, a heptafluoropropylsulfinyl group, and a heptafluoroisopropylsulfinyl group.

[0031]

5 In the present compound, examples of the "C1-C3 alkylsulfonyl group having one or more a fluorine atoms" include a trifluoromethylsulfonyl group, a 2,2,2-trifluoroethylsulfonyl group, a pentafluoroethylsulfonyl group, a heptafluoropropylsulfonyl group, and a
10 heptafluoroisopropylsulfonyl group.

[0032]

 In the present compound, examples of the "C1-C3 alkyl group" include a methyl group, an ethyl group, a propyl group, and an isopropyl group.

15 [0033]

 In the present compound, examples of the "C1-C3 alkyl group optionally having one or more halogen atoms" include a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a
20 pentafluoroethyl group, a 2,2,2-trifluoroethyl group, a 2,2,2-trichloroethyl group, a heptafluoropropyl group, and a heptafluoroisopropyl group.

[0034]

 Examples of the present compound include the following
25 pyrimidine compounds.

A compound of the formula (1) wherein A^1 is $-NR^8-$, an oxygen atom, or a sulfur atom, and R^8 is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group having one C2-C6 alkoxycarbonyl group, $-CO_2R^{11}$, or a cyclopropyl group;

A compound of the formula (1) wherein A^1 is $-NR^8-$, an oxygen atom, or a sulfur atom, and R^8 is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a cyclopropyl group;

A compound of the formula (1) wherein A^1 is $-NR^8-$, an oxygen atom, or a sulfur atom, and R^8 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a cyclopropyl group;

A compound of the formula (1) wherein A^1 is $-NR^8-$, an oxygen atom, or a sulfur atom, and R^8 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a methoxymethyl group, an ethoxymethyl group, or a cyclopropyl group;

A compound of the formula (1) wherein A^1 is $-NR^8-$, an oxygen atom, or a sulfur atom, and R^8 is a methyl group;

A compound of the formula (1) wherein A^1 is $-NR^8-$ or

an oxygen atom, and R^8 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a cyclopropyl group;

A compound of the formula (1) wherein A^1 is $-NR^8-$, and
5 R^8 is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group having one C2-C6 alkoxycarbonyl group, $-CO_2R^{11}$, or a cyclopropyl group;

10 A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a cyclopropyl group;

15 A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a methoxymethyl group, an ethoxymethyl group, or a cyclopropyl group;

A compound of the formula (1) wherein A^1 is $-NR^8-$, and
20 R^8 is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group;

A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is a methyl group;

A compound of the formula (1) wherein A^1 is an oxygen
25 atom;

A compound of the formula (1) wherein A^1 is a sulfur atom;

[0035]

5 A compound of the formula (1) wherein A^2 is a nitrogen atom;

A compound of the formula (1) wherein A^2 is a nitrogen atom or $=CR^9-$, and R^9 is a C1-C3 chain hydrocarbon group, a halogen atom, or a hydrogen atom;

10 A compound of the formula (1) wherein A^2 is a nitrogen atom or $=CR^9-$, and R^9 is a halogen atom or a hydrogen atom;

A compound of the formula (1) wherein A^2 is a nitrogen atom or $=CR^9-$, and R^9 is a hydrogen atom;

15 A compound of the formula (1) wherein A^2 is $=CR^9-$, and R^9 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, $-NR^{11}R^{12}$, $-CO_2R^{11}$, $-C(O)R^{11}$, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

20 A compound of the formula (1) wherein A^2 is $=CR^9-$, and R^9 is a C1-C3 chain hydrocarbon group, a halogen atom, or a hydrogen atom;

A compound of the formula (1) wherein A^2 is $=CR^9-$, and R^9 is a halogen atom or a hydrogen atom;

A compound of the formula (1) wherein A^2 is $=CR^9-$, and R^9 is a hydrogen atom;

25 [0036]

A compound of the formula (1) wherein A^3 is a nitrogen atom;

A compound of the formula (1) wherein A^3 is a nitrogen atom or $=CR^{10}-$, and R^{10} is a C1-C3 chain hydrocarbon group,
5 a halogen atom, or a hydrogen atom;

A compound of the formula (1) wherein A^3 is a nitrogen atom or $=CR^{10}-$, and R^{10} is a halogen atom or a hydrogen atom;

A compound of the formula (1) wherein A^3 is a nitrogen atom or $=CR^{10}-$, and R^{10} is a hydrogen atom;
10

A compound of the formula (1) wherein A^3 is $=CR^{10}-$, and R^{10} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, $-NR^{11}R^{12}$, $-CO_2R^{11}$, $-C(O)R^{11}$, a cyano group, a nitro group, a halogen atom, or
15 a hydrogen atom;

A compound of the formula (1) wherein A^3 is $=CR^{10}-$, and R^{10} is a C1-C3 chain hydrocarbon group, a halogen atom, or a hydrogen atom;

A compound of the formula (1) wherein A^3 is $=CR^{10}-$, and
20 R^{10} is a halogen atom or a hydrogen atom;

A compound of the formula (1) wherein A^3 is $=CR^{10}-$, and R^{10} is a halogen atom;

A compound of the formula (1) wherein A^3 is $=CR^{10}-$, and R^{10} is a hydrogen atom;

25 [0037]

A compound of the formula (1) wherein A^1 is $-NR^8-$, an oxygen atom, or a sulfur atom, R^8 is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group having one C2-C6 alkoxycarbonyl group, $-CO_2R^{11}$, or a cyclopropyl group, A^2 is a nitrogen atom or $=CR^9-$, R^9 is a halogen atom or a hydrogen atom, A^3 is a nitrogen atom or $=CR^{10}-$, and R^{10} is a halogen atom or a hydrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, an oxygen atom, or a sulfur atom, R^8 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a methoxymethyl group, an ethoxymethyl group, or a cyclopropyl group, A^2 is a nitrogen atom or $=CR^9-$, R^9 is a halogen atom or a hydrogen atom, A^3 is a nitrogen atom or $=CR^{10}-$, and R^{10} is a halogen atom or a hydrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, an oxygen atom, or a sulfur atom, R^8 is a methyl group, A^2 is a nitrogen atom or $=CR^9-$, R^9 is a halogen atom or a hydrogen atom, A^3 is a nitrogen atom or $=CR^{10}-$, and R^{10} is a halogen atom or a hydrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, an oxygen atom, or a sulfur atom, R^8 is a methyl group, A^2 is $=CR^9-$, R^9 is a hydrogen atom, and A^3 is a nitrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, A^2 is $=CR^9-$, R^9 is a hydrogen atom, and A^3 is a nitrogen atom;

5 A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a methyl group, A^2 is $=CR^9-$, R^9 is a hydrogen atom, and A^3 is a nitrogen atom;

10 A compound of the formula (1) wherein A^1 is an oxygen atom, A^2 is $=CR^9-$, R^9 is a hydrogen atom, and A^3 is a nitrogen atom;

A compound of the formula (1) wherein A^1 is a sulfur atom, A^2 is $=CR^9-$, R^9 is a hydrogen atom, and A^3 is a nitrogen atom;

[0038]

15 A compound of the formula (1) wherein R^1 is a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from Group X;

20 A compound of the formula (1) wherein R^1 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (1) wherein R^1 is a C3-C6 alicyclic hydrocarbon group optionally having one or more atoms or groups selected from Group Y;

25 A compound of the formula (1) wherein R^1 is a methyl group, an ethyl group, a propyl group, an isopropyl group,

a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, and a 2,2,2-trifluoroethyl group;

A compound of the formula (1) wherein R¹ is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, and a 2,2,2-trifluoroethyl group, and R², R³, R⁴, and R⁵ are a hydrogen atom;

A compound of the formula (1) wherein R¹ is a methyl group, an ethyl group, or a propyl group;

A compound of the formula (1) wherein R¹ is an ethyl group;

[0039]

A compound of the formula (1) wherein R², R³, R⁴, and R⁵ are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, -OR¹¹, -NR¹¹R¹², a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

A compound of the formula (1) wherein R², R³, R⁴, and R⁵ are same or different and are independently a methyl group, an ethyl group, a trifluoromethyl group, or a hydrogen atom;

A compound of the formula (1) wherein R², R³, R⁴, and R⁵ are same or different and are independently a methoxy

group, an ethoxy group, a trifluoromethoxy group, or a hydrogen atom;

A compound of the formula (1) wherein R^2 , R^3 , R^4 , and R^5 are same or different and are independently an amino group, a methylamino group, a dimethylamino group, or a hydrogen atom;

A compound of the formula (1) wherein R^2 , R^3 , R^4 , and R^5 are same or different and are independently a cyano group or a hydrogen atom;

10 A compound of the formula (1) wherein R^2 , R^3 , R^4 , and R^5 are same or different and are independently a nitro group or a hydrogen atom;

A compound of the formula (1) wherein R^2 , R^3 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom;

15 A compound of the formula (1) wherein R^2 , R^3 , R^4 , and R^5 are same or different and are independently a fluorine atom, a chlorine atom, a bromine atom, an iodine atom, or a hydrogen atom;

20 A compound of the formula (1) wherein R^2 , R^3 , R^4 , and R^5 are same or different and are independently a fluorine atom, a chlorine atom, a bromine atom, a methyl group, a trifluoromethoxy group, a pentafluoroethyl group, a methoxy group, a cyano group, a nitro group, or a hydrogen atom;

25 A compound of the formula (1) wherein R^2 , R^4 , and R^5

are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 alkylsulfanyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfinyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfonyl group optionally having one or more halogen atoms, a C2-C6 alkylcarbonyl group optionally having one or more halogen atoms, a C2-C6 alkoxy carbonyl group optionally having one or more halogen atoms, a C1-C6 alkylamino group optionally having one or more halogen atoms, C2-C8 dialkylamino group optionally having one or more halogen atoms, an aldehyde group, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a halogen atom;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a methyl group, a trifluoromethyl group, a trifluoromethoxy group, a fluorine atom, a chlorine atom, a bromine atom, an iodine atom, or a

hydrogen atom;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a trifluoromethyl group;

5 A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

10 A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a methyl group, a trifluoromethyl group, a trifluoromethoxy group, a chlorine atom, a bromine atom, an iodine atom, or a hydrogen atom;

15 A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a methyl group, a trifluoromethyl group, a trifluoromethoxy group, a fluorine atom, a chlorine atom, a bromine atom, or an iodine atom;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a trifluoromethyl group;

A compound of the formula (1) wherein R^2 , R^3 , R^4 , and R^5 are a hydrogen atom;

20 A compound of the formula (1) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a methyl group, a trifluoromethyl group, a trifluoromethoxy group, a chlorine atom, a bromine atom, an iodine atom, or
25 a hydrogen atom;

A compound of the formula (1) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a trifluoromethyl group;

5 A compound of the formula (1) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a methyl group, a trifluoromethyl group, a trifluoromethoxy group, a chlorine atom, a bromine atom, an iodine atom, or a hydrogen atom;

10 A compound of the formula (1) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a trifluoromethyl group;

[0040]

A compound of the formula (1) wherein R^6 and R^7 are
15 same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, $-NR^{11}R^{12}$, $-CO_2R^{11}$, $-C(O)R^{11}$, a cyano group, a nitro group, a halogen atom, or a hydrogen atom, and R^{11} and R^{12} are same or different and are independently
20 a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a hydrogen atom;

A compound of the formula (1) wherein R^6 and R^7 are
same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen
25 atoms, $-OR^{11}$, $-S(O)_mR^{11}$, a halogen atom, or a hydrogen atom,

and R^{11} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a hydrogen atom;

A compound of the formula (1) wherein R^6 and R^7 are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally having one or more halogen atoms, an aldehyde group, a cyano group, a halogen atom, or a hydrogen atom;

A compound of the formula (1) wherein R^6 and R^7 are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, an aldehyde group, a cyano group, a halogen atom, or a hydrogen atom;

A compound of the formula (1) wherein R^6 and R^7 are same or different and are independently a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, an iodine atom, or a hydrogen atom;

A compound of the formula (1) wherein R^6 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, or a halogen atom, and R^7 is a hydrogen atom;

A compound of the formula (1) wherein R^6 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, and R^7 is a hydrogen atom;

5 A compound of the formula (1) wherein R^6 is a C1-C6 chain hydrocarbon group having one or more halogen atoms, and R^7 is a hydrogen atom;

A compound of the formula (1) wherein R^6 is a C1-C6 chain hydrocarbon group having one or more a fluorine atoms, and R^7 is a hydrogen atom;

10 A compound of the formula (1) wherein R^6 is $-OR^{11}$, and R^7 is a hydrogen atom;

A compound of the formula (1) wherein R^6 is $-S(O)_mR^{11}$, and R^7 is a hydrogen atom;

15 A compound of the formula (1) wherein R^6 is a C1-C6 chain hydrocarbon group having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, or a halogen atom, and R^7 is a hydrogen atom;

20 A compound of the formula (1) wherein R^6 is a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom, and R^7 is a hydrogen atom;

25 A compound of the formula (1) wherein R^6 is a hydrogen

atom, and R⁷ is a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom;

A compound of the formula (1) wherein R⁶ is a bromine atom, and R⁷ is a hydrogen atom;

10 A compound of the formula (1) wherein R⁶ is an iodine atom, and R⁷ is a hydrogen atom;

A compound of the formula (1) wherein R⁶ is a trifluoromethyl group, and R⁷ is a hydrogen atom;

A compound of the formula (1) wherein R⁶ is a pentafluoroethyl group, and R⁷ is a hydrogen atom;

15 A compound of the formula (1) wherein R⁶ is a trifluoromethoxy group, and R⁷ is a hydrogen atom;

A compound of the formula (1) wherein R⁶ is a trifluoromethylsulfanyl group, and R⁷ is a hydrogen atom;

20 A compound of the formula (1) wherein R⁶ is a trifluoromethylsulfinyl group, and R⁷ is a hydrogen atom;

A compound of the formula (1) wherein R⁶ is a trifluoromethylsulfonyl group, and R⁷ is a hydrogen atom;

A compound of the formula (1) wherein R⁶ is a heptafluoroisopropyl group, and R⁷ is a hydrogen atom;

25 [0041]

A compound of the formula (1) wherein n is 0;

A compound of the formula (1) wherein n is 1;

A compound of the formula (1) wherein n is 2;

[0042]

5 A compound of the formula (1) wherein m is 0;

A compound of the formula (1) wherein m is 1;

A compound of the formula (1) wherein m is 2;

[0043]

10 A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group, A^2 is $=CR^9-$, R^9 is a hydrogen atom, A^3 is a nitrogen atom or $=CR^{10}-$, and R^{10} is a halogen atom or a hydrogen atom;

15 A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group, A^2 is $=CR^9-$, R^9 is a hydrogen atom, A^3 is a nitrogen atom or $=CR^{10}-$, R^{10} is a halogen atom or a hydrogen atom, and R^1 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-
20 butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group;

25 A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group, A^2 is $=CR^9-$, R^9 is a hydrogen atom,

A³ is a nitrogen atom or =CR¹⁰-, R¹⁰ is a halogen atom or a hydrogen atom, R¹ is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, and R², R³, R⁴, and R⁵ are a hydrogen atom;

[0044]

A compound of the formula (1) wherein A¹ is -NR⁸-, R⁸ is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group, and R¹ is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group;

A compound of the formula (1) wherein A¹ is -NR⁸-, R⁸ is a methyl group, an ethyl group, a methoxymethyl group or an ethoxymethyl group, R¹ is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, and R², R³, R⁴, and R⁵ are a hydrogen atom;

[0045]

A compound of the formula (1) wherein A¹ is -NR⁸-, R⁸

is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group, R¹ is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, R², R³, R⁴, and R⁵ are a hydrogen atom, R⁶ is a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom, and R⁷ is a hydrogen atom;

A compound of the formula (1) wherein A¹ is -NR⁸-, R⁸ is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group, R¹ is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, R², R³, R⁴, and R⁵ are a hydrogen atom, and R⁷ is a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom;

[0046]

A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group, A^2 is $=CR^9-$, R^9 is a hydrogen atom, 5 A^3 is a nitrogen atom or $=CR^{10}-$, R^{10} is a halogen atom or a hydrogen atom, R^1 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, or a 2,2,2-10 trifluoroethyl group, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl 15 group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom, and R^7 is a hydrogen atom;

A compound of the formula (1) wherein A^1 is a sulfur atom, A^2 is $=CR^9-$, R^9 is a hydrogen atom, A^3 is a nitrogen atom or $=CR^{10}-$, R^{10} is a halogen atom or a hydrogen atom, R^1 20 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a trifluoromethyl 25 group, a difluoromethyl group, a fluoromethyl group, a

pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom, and R^7 is a hydrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group, A^2 is $=CR^9-$, R^9 is a hydrogen atom, A^3 is a nitrogen atom or $=CR^{10}-$, R^{10} is a halogen atom or a hydrogen atom, R^1 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom, R^7 is a hydrogen atom, and n is 0;

A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group, A^2 is $=CR^9-$, R^9 is a hydrogen atom, A^3 is a nitrogen atom or $=CR^{10}-$, R^{10} is a halogen atom or a hydrogen atom, R^1 is a methyl group, an ethyl group, a

propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom, R^7 is a hydrogen atom, and n is 1;

A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group, A^2 is $=CR^9-$, R^9 is a hydrogen atom, A^3 is a nitrogen atom or $=CR^{10}-$, R^{10} is a halogen atom or a hydrogen atom, R^1 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom, R^7 is a hydrogen atom, and n is 2;

A compound of the formula (1) wherein A^1 is $-NR^8-$, an oxygen atom, or a sulfur atom, A^2 is a nitrogen atom or $=CR^9-$, A^3 is a nitrogen atom or $=CR^{10}-$, R^1 is a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from Group X, R^2 , R^3 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, R^6 and R^7 are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally having one or more halogen atoms, an aldehyde group, a cyano group, a halogen atom, or a hydrogen atom, R^8 is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a cyclopropyl group, R^9 and R^{10} are same or different and are independently a halogen atom or a hydrogen atom, and R^{11} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (1) wherein A^1 is $-NR^8-$;

A compound of the formula (1) wherein A^1 is $-NR^8-$, an oxygen atom, or a sulfur atom, and R^8 is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group having one C2-C6 alkoxycarbonyl

group, $-\text{CO}_2\text{R}^{11}$, a cyclopropyl group, or a hydrogen atom;

A compound of the formula (1) wherein A^1 is $-\text{NR}^8-$, an oxygen atom, or a sulfur atom, and R^8 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (1) wherein A^1 is $-\text{NR}^8-$, an oxygen atom, or a sulfur atom, and R^8 is a C1-C6 chain hydrocarbon group;

A compound of the formula (1) wherein A^1 is $-\text{NR}^8-$, an oxygen atom, or a sulfur atom, and R^8 is a C1-C6 alkyl group;

A compound of the formula (1) wherein A^1 is $-\text{NR}^8-$, an oxygen atom, or a sulfur atom, and R^8 is a C1-C3 alkyl group;

A compound of the formula (1) wherein A^1 is $-\text{NR}^8-$, an oxygen atom, or a sulfur atom, and R^8 is a hydrogen atom;

A compound of the formula (1) wherein A^1 is $-\text{NR}^8-$, and R^8 is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a cyclopropyl group, or a hydrogen atom;

A compound of the formula (1) wherein A^1 is $-\text{NR}^8-$, and R^8 is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, or a C1-C6 chain hydrocarbon group optionally having one or

more halogen atoms;

A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

5 A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is a C1-C6 chain hydrocarbon group;

A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is a C1-C6 alkyl group;

10 A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is a C1-C3 alkyl group;

A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is a methyl group, an ethyl group, a propyl group, or an isopropyl group;

15 A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a methoxymethyl group, an ethoxymethyl group, a cyclopropyl group, or a hydrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is an ethyl group;

20 A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is a propyl group;

A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is an isopropyl group;

25 A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is a methyl group or a hydrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is a hydrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, and R^8 is a methoxymethyl group, an ethoxymethyl group, or a
5 hydrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$ or a sulfur atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$ or a sulfur atom, and R^8 is a methyl group, an ethyl group, a
10 propyl group, or an isopropyl group;

A compound of the formula (1) wherein A^1 is $-NR^8-$ or a sulfur atom, and R^8 is a methoxymethyl group, an ethoxymethyl group, or a hydrogen atom;

A compound of the formula (1) wherein A^2 is $=CR^9-$, and
15 A^3 is $=CR^{10}-$;

A compound of the formula (1) wherein A^2 is $=CR^9-$, A^3 is $=CR^{10}-$, and R^9 and R^{10} are a hydrogen atom;

A compound of the formula (1) wherein A^2 is a nitrogen atom, and A^3 is $=CR^{10}-$;

20 A compound of the formula (1) wherein A^2 is $=CR^9-$, and A^3 is a nitrogen atom;

A compound of the formula (1) wherein A^2 is $=CR^9-$, R^9 is a hydrogen atom, and A^3 is a nitrogen atom;

25 A compound of the formula (1) wherein A^2 is a nitrogen atom, and A^3 is a nitrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, A^2 is $=CR^9-$, and A^3 is $=CR^{10}-$;

A compound of the formula (1) wherein A^1 is $-NR^8-$, A^2 is a nitrogen atom, and A^3 is $=CR^{10}-$;

5 A compound of the formula (1) wherein A^1 is $-NR^8-$, A^2 is $=CR^9-$, and A^3 is a nitrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, A^2 is a nitrogen atom, and A^3 is a nitrogen atom;

10 A compound of the formula (1) wherein A^1 is an oxygen atom, A^2 is $=CR^9-$, and A^3 is $=CR^{10}-$;

A compound of the formula (1) wherein A^1 is an oxygen atom, A^2 is a nitrogen atom, and A^3 is $=CR^{10}-$;

A compound of the formula (1) wherein A^1 is an oxygen atom, A^2 is $=CR^9-$, and A^3 is a nitrogen atom;

15 A compound of the formula (1) wherein A^1 is an oxygen atom, A^2 is a nitrogen atom, and A^3 is a nitrogen atom;

A compound of the formula (1) wherein A^1 is a sulfur atom, A^2 is $=CR^9-$, and A^3 is $=CR^{10}-$;

20 A compound of the formula (1) wherein A^1 is a sulfur atom, A^2 is a nitrogen atom, and A^3 is $=CR^{10}-$;

A compound of the formula (1) wherein A^1 is a sulfur atom, A^2 is $=CR^9-$, and A^3 is a nitrogen atom;

A compound of the formula (1) wherein A^1 is a sulfur atom, A^2 is a nitrogen atom, and A^3 is a nitrogen atom;

25 A compound of the formula (1) wherein A^1 is $-NR^8-$ or a

sulfur atom, R^8 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a methoxymethyl group, an ethoxymethyl group, a cyclopropyl group, or a hydrogen atom, A^2 is $=CR^9-$, R^9 is a hydrogen atom, and A^3 is a nitrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$ or a sulfur atom, R^8 is a methyl group, A^2 is $=CR^9-$, R^9 is a hydrogen atom, and A^3 is a nitrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a methoxymethyl group, an ethoxymethyl group, a cyclopropyl group, or a hydrogen atom, A^2 is $=CR^9-$, R^9 is a hydrogen atom, and A^3 is a nitrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a methyl group, A^2 is $=CR^9-$, R^9 is a hydrogen atom, and A^3 is a nitrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a methyl group, A^2 is $=CR^9-$, A^3 is $=CR^{10}-$, and R^9 and R^{10} are a hydrogen atom;

A compound of the formula (1) wherein A^1 is an oxygen atom, A^2 is $=CR^9-$, A^3 is $=CR^{10}-$, and R^9 and R^{10} are a hydrogen atom;

A compound of the formula (1) wherein A^1 is a sulfur atom, A^2 is $=CR^9-$, A^3 is $=CR^{10}-$, and R^9 and R^{10} are a hydrogen atom;

A compound of the formula (1) wherein R^1 is a C1-C6 chain hydrocarbon group optionally having one or more fluorine atoms;

5 A compound of the formula (1) wherein R^1 is a C1-C6 chain hydrocarbon group;

A compound of the formula (1) wherein R^1 is a C1-C6 alkyl group;

A compound of the formula (1) wherein R^1 is a C1-C3 alkyl group;

10 A compound of the formula (1) wherein R^1 is a C3-C6 alicyclic hydrocarbon group;

A compound of the formula (1) wherein R^1 is a cyclopropyl group, a cyclobutyl group, or a cyclopentyl group;

15 A compound of the formula (1) wherein R^1 is a methyl group;

A compound of the formula (1) wherein R^1 is a propyl group;

20 A compound of the formula (1) wherein R^1 is an isopropyl group;

A compound of the formula (1) wherein R^1 is a trifluoromethyl group;

A compound of the formula (1) wherein R^1 is a 2,2,2-trifluoroethyl group;

25 A compound of the formula (1) wherein R^2 , R^3 , R^4 , and

R⁵ are same or different and are independently a phenyl group optionally having one or more atoms or groups selected from Group Z, a 5-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z, a 6-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z, or a hydrogen atom;

A compound of the formula (1) wherein R², R³, R⁴, and R⁵ are same or different and are independently a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, or a hydrogen atom;

A compound of the formula (1) wherein R², R³, R⁴, and R⁵ are same or different and are independently a 2-pyridyl group, a 2-pyrimidyl group, a 3-chloro-2-pyridyl group, a 3-chloro-5-trifluoromethyl-2-pyridyl group, or a hydrogen atom;

A compound of the formula (1) wherein R², R³, R⁴, and R⁵ are same or different and are independently -SF₅ or a hydrogen atom;

A compound of the formula (1) wherein R^2 , R^3 , R^4 , and R^5 are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a hydrogen atom;

5 A compound of the formula (1) wherein R^2 , R^3 , R^4 , and R^5 are same or different and are independently a C1-C3 alkyl group optionally having one or more halogen atoms or a hydrogen atom;

10 A compound of the formula (1) wherein R^2 , R^3 , R^4 , and R^5 are same or different and are independently $-OR^{11}$ or a hydrogen atom;

A compound of the formula (1) wherein R^2 , R^3 , R^4 , and R^5 are same or different and are independently $-NR^{11}R^{12}$ or a hydrogen atom;

15 A compound of the formula (1) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, R^3 is a phenyl group optionally having one or more atoms or groups selected from Group Z, a 5-membered heterocyclic group optionally having one or more
20 atoms or groups selected from Group Z, or a 6-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z;

25 A compound of the formula (1) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, R^3 is a phenyl group optionally having

one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, or a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a methyl group;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a trifluoromethoxy group;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a fluorine atom;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a chlorine atom;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a bromine atom;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is an iodine atom;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a C1-C6 alkyl group;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a C1-C6 alkyl group
5 optionally having one or more fluorine atoms;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a C1-C3 alkyl group;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a C1-C3 alkyl group
10 optionally having one or more fluorine atoms;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a 5-membered heterocyclic group optionally having one or more atoms or groups selected from Z group, or a 6-membered heterocyclic group
15 optionally having one or more atoms or groups selected from Group Z;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a 6-membered heterocyclic group optionally having one or more atoms or groups
20 selected from Group Z;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a methyl group;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a trifluoromethoxy group;

25 A compound of the formula (1) wherein R^2 , R^4 , and R^5

are a hydrogen atom, and R^3 is a fluorine atom;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a chlorine atom;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a bromine atom;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is an iodine atom;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, or a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a 2-pyridyl group, a 2-pyrimidyl group, a 3-chloro-2-pyridyl group, or a 3-chloro-5-trifluoromethyl-2-pyridyl group;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a 2-pyridyl group;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a 2-pyrimidyl group;

A compound of the formula (1) wherein R^2 , R^4 , and R^5

are a hydrogen atom, and R^3 is $-SF_5$;

A compound of the formula (1) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a 2-pyridyl group, a 2-pyrimidyl group, a 3-chloro-2-pyridyl group, a 3-chloro-5-trifluoromethyl-2-pyridyl group, a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom;

A compound of the formula (1) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a methyl group;

A compound of the formula (1) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a trifluoromethoxy group;

A compound of the formula (1) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a chlorine atom;

A compound of the formula (1) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a bromine atom;

A compound of the formula (1) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is an iodine atom;

A compound of the formula (1) wherein R^6 is a C1-C3

alkyl group having one or more fluorine atoms, and R⁷ is a hydrogen atom;

A compound of the formula (1) wherein R⁶ is a C1-C3 alkoxy group having one or more fluorine atoms, and R⁷ is a
5 hydrogen atom;

A compound of the formula (1) wherein R⁶ is a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms or a C1-C3 alkylsulfonyl group having one or more fluorine
10 atoms, and R⁷ is a hydrogen atom;

A compound of the formula (1) wherein R⁶ is a halogen atom, and R⁷ is a hydrogen atom;

A compound of the formula (1) wherein R⁶ is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3
15 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or a halogen atom;

20 A compound of the formula (1) wherein R⁶ is -SF₅, and R⁷ is a hydrogen atom;

A compound of the formula (1) wherein A¹ is -NR⁸- or a sulfur atom, R⁸ is a C1-C6 alkyl group, A² is =CR⁹-, A³ is =CR¹⁰-, R⁹ and R¹⁰ are a hydrogen atom, R¹ is a C1-C3 alkyl
25 group optionally having one or more fluorine atoms, R², R⁴,

and R^5 are a hydrogen atom, R^3 is a C1-C3 alkyl group optionally having one or more fluorine atoms or a halogen atom, R^6 is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or a halogen atom, and R^7 is a hydrogen atom;

10 A compound of the formula (1) wherein A^1 is $-NR^8-$ or a sulfur atom, R^8 is a C1-C6 alkyl group, A^2 is $=CR^9-$, R^9 is a hydrogen atom, A^3 is a nitrogen atom, R^1 is a C1-C3 alkyl group optionally having one or more fluorine atoms, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a C1-C3 alkyl group optionally having one or more fluorine atoms or a halogen atom, R^6 is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or a halogen atom, and R^7 is a hydrogen atom;

A compound of the formula (1) wherein A^1 is a sulfur atom, A^2 is $=CR^9-$, R^9 is a hydrogen atom, A^3 is a nitrogen atom or $=CR^{10}-$, R^{10} is a halogen atom or a hydrogen atom, R^1

25

is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom, R^7 is a hydrogen atom, and n is 0;

A compound of the formula (1) wherein A^1 is a sulfur atom, A^2 is $=CR^9-$, R^9 is a hydrogen atom, A^3 is a nitrogen atom or $=CR^{10}-$, R^{10} is a halogen atom or a hydrogen atom, R^1 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom, R^7 is a hydrogen atom, and n is 1;

A compound of the formula (1) wherein A^1 is a sulfur atom, A^2 is $=CR^9-$, R^9 is a hydrogen atom, A^3 is a nitrogen atom or $=CR^{10}-$, R^{10} is a halogen atom or a hydrogen atom, R^1 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl group, a 2,2,2-trifluoroethyl group, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom, R^7 is a hydrogen atom, and n is 2;

A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a methyl group, an ethyl group, a propyl group, or an isopropyl group, A^2 is $=CR^9-$, A^3 is $=CR^{10}-$, R^9 and R^{10} are a hydrogen atom, R^1 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom, R^6 is a trifluoromethyl group, a

pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom or an iodine atom, and R^7 is a hydrogen atom;

A compound of the formula (1) wherein A^1 is a sulfur atom, A^2 is $=CR^9-$, A^3 is $=CR^{10}-$, R^9 and R^{10} are a hydrogen atom, R^1 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom or a bromine atom, R^6 is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom, and R^7 is a hydrogen atom;

A compound of the formula (1) wherein A^1 is $-NR^8-$, R^8 is a methyl group, an ethyl group, a propyl group, or an isopropyl group, A^2 is $=CR^9-$, R^9 is a hydrogen atom, A^3 is a nitrogen atom, R^1 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group,


or a 2,2,2-trifluoroethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom, R^6 is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom, and R^7 is a hydrogen atom;

A compound of the formula (1) wherein A^1 is a sulfur atom, A^2 is $=CR^9-$, R^9 is a hydrogen atom, A^3 is a nitrogen atom, R^1 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom or a bromine atom, R^6 is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a

bromine atom, or an iodine atom, and R^7 is a hydrogen atom;

[0047]

A compound of the formula (1-1)



5 wherein;

A^{1a} represents $-NR^{8a}-$ or a sulfur atom;

A^{3a} represents a nitrogen atom or $=CR^{10a}-$;

R^{1a} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

10 R^{2a} , R^{4a} , and R^{5a} are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a halogen atom, or a hydrogen atom;

R^{3a} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11a}$, a halogen atom, or a hydrogen atom;

R^{6a} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11a}$, $-S(O)_mR^{11a}$, a bromine atom, or an iodine atom;

20 R^{8a} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a hydrogen atom;

R^{10a} represents a halogen atom or a hydrogen atom;

R^{11a} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, m represents 0, 1, or 2, and n represents 0, 1 or 2;

5 A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$ or a sulfur atom, R^{8a} is a methyl group, A^{3a} is a nitrogen atom, R^{1a} is an ethyl group, R^{2a} , R^{4a} , and R^{5a} are a halogen atom or a hydrogen atom, R^{3a} is a trifluoromethyl group, a halogen atom, or a hydrogen atom, and R^{6a} is a C1-C3 chain
10 hydrocarbon group having one or more a fluorine atoms, a C1-C3 alkoxy group having one or more a fluorine atoms, a C1-C3 alkylsulfanyl group having one or more a fluorine atoms, a C1-C3 alkylsulfinyl group having one or more a fluorine atoms, or a C1-C3 alkylsulfonyl group having one
15 or more a fluorine atoms;

 A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$ or a sulfur atom, R^{8a} is a methyl group, A^{3a} is a nitrogen atom, R^{1a} is an ethyl group, R^{2a} , R^{4a} , and R^{5a} are a hydrogen atom, R^{3a} is a chlorine atom, a bromine atom, a
20 trifluoromethyl group, or a hydrogen atom, and R^{6a} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

25 A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$

or a sulfur atom, and R^{8a} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$, and R^{8a} is a C1-C6 chain hydrocarbon group optionally
5 having one or more halogen atoms;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$, and R^{8a} is a methyl group;

A compound of the formula (1-1) wherein A^{1a} is a sulfur atom;

10 [0048]

A compound of the formula (1-1) wherein A^{3a} is a nitrogen atom or $=CR^{10a}-$, and R^{10a} is a halogen atom or a hydrogen atom;

A compound of the formula (1-1) wherein A^{3a} is $=CR^{10a}-$,
15 and R^{10a} is a halogen atom or a hydrogen atom;

A compound of the formula (1-1) wherein A^{3a} is a nitrogen atom;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$ or a sulfur atom, R^{8a} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, A^{3a} is a
20 nitrogen atom or $=CR^{10a}-$, and R^{10a} is a hydrogen atom;

[0049]

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$ or a sulfur atom, R^{8a} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, and A^{3a} is a
25

nitrogen atom;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$ or a sulfur atom, R^{8a} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, A^{3a} is $=CR^{10a}-$,
5 and R^{10a} is a hydrogen atom;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$ or a sulfur atom, R^{8a} is a methyl group, and A^{3a} is a nitrogen atom;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$,
10 R^{8a} is a methyl group, and A^{3a} is a nitrogen atom;

A compound of the formula (1-1) wherein A^{1a} is a sulfur atom, and A^{3a} is a nitrogen atom;

[0050]

A compound of the formula (1-1) wherein R^{1a} is a C1-C6
15 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (1-1) wherein R^{1a} is an ethyl group;

A compound of the formula (1-1) wherein R^{2a} , R^{4a} , and
20 R^{5a} are same or different and are independently a fluorine atom, a chlorine atom, or a hydrogen atom, and R^{3a} is a fluorine atom, a chlorine atom, a bromine atom, a methyl group, a trifluoromethyl group, or a hydrogen atom;

A compound of the formula (1-1) wherein R^{2a} , R^{4a} , and
25 R^{5a} are a hydrogen atom, and R^{3a} is a fluorine atom, a

chlorine atom, a bromine atom, a methyl group, or a trifluoromethyl group;

A compound of the formula (1-1) wherein R^{6a} is a trifluoromethyl group, a pentafluoroethyl group, a
5 heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

A compound of the formula (1-1) wherein R^{6a} is a bromine atom;

10 A compound of the formula (1-1) wherein R^{6a} is an iodine atom;

A compound of the formula (1-1) wherein R^{6a} is a trifluoromethyl group;

15 A compound of the formula (1-1) wherein R^{6a} is a pentafluoroethyl group;

A compound of the formula (1-1) wherein R^{6a} is a heptafluoroisopropyl group;

A compound of the formula (1-1) wherein R^{6a} is a trifluoromethoxy group;

20 A compound of the formula (1-1) wherein R^{6a} is a trifluoromethylsulfanyl group;

A compound of the formula (1-1) wherein R^{6a} is a trifluoromethylsulfinyl group;

25 A compound of the formula (1-1) wherein R^{6a} is a trifluoromethylsulfonyl group;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$,
and R^{8a} is a C1-C3 alkyl group;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$,
and R^{8a} is a methyl group, an ethyl group, a propyl group,
5 or an isopropyl group;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$,
and R^{8a} is an ethyl group;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$,
and R^{8a} is a propyl group;

10 A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$,
and R^{8a} is an isopropyl group;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$,
and R^{8a} is a methyl group or a hydrogen atom;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$,
15 and R^{8a} is a hydrogen atom;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$,
and R^{8a} is a methoxymethyl group, an ethoxymethyl group, or
a hydrogen atom;

A compound of the formula (1-1) wherein A^{3a} is $=CR^{10a}-$,
20 and R^{10a} is a hydrogen atom;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$,
and A^{3a} is $=CR^{10a}-$;

A compound of the formula (1-1) wherein A^{1a} is a
sulfur atom, and A^{3a} is $=CR^{10a}-$;

25 A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$,

and A^{3a} is a nitrogen atom;

A compound of the formula (1-1) wherein A^{1a} is -NR^{8a}-,
R^{8a} is a hydrogen atom group, and A^{3a} is a nitrogen atom;

A compound of the formula (1-1) wherein R^{1a} is a C1-C3
5 alkyl group;

A compound of the formula (1-1) wherein R^{1a} is a
methyl group, an ethyl group, a propyl group, an isopropyl
group, a butyl group, a sec-butyl group, an isobutyl group,
a tert-butyl group, a cyclopropyl group, a trifluoromethyl
10 group, or a 2,2,2-trifluoroethyl group;

A compound of the formula (1-1) wherein R^{1a} is a
methyl group;

A compound of the formula (1-1) wherein R^{1a} is a
propyl group;

15 A compound of the formula (1-1) wherein R^{2a}, R^{3a}, R^{4a},
and R^{5a} are a hydrogen atom;

A compound of the formula (1-1) wherein R^{6a} is a
halogen atom;


A compound of the formula (1-1) wherein A^{1a} is -NR^{8a}-
20 or a sulfur atom, R^{8a} is a C1-C6 alkyl group, A^{3a} is =CR^{10a}-,
R^{10a} is a hydrogen atom, R^{1a} is a C1-C3 alkyl group
optionally having one or more fluorine atoms, R^{2a}, R^{4a}, and
R^{5a} are a hydrogen atom, R^{3a} is a C1-C3 alkyl group
optionally having one or more fluorine atoms or a halogen
25 atom, and R^{6a} is a C1-C3 alkyl group having one or more

fluorine atoms, a C1-C3 alkoxy group having one or more
 fluorine atoms, a C1-C3 alkylsulfanyl group having one or
 more fluorine atoms, a C1-C3 alkylsulfinyl group having one
 or more fluorine atoms, a C1-C3 alkylsulfonyl group having
 5 one or more fluorine atoms, or a halogen atom;

A compound of the formula (1-1) wherein A^{1a} is $-NR^{8a}-$
 or a sulfur atom, R^{8a} is a C1-C6 alkyl group, A^{3a} is a
 nitrogen atom, R^{1a} is a C1-C3 alkyl group optionally having
 one or more fluorine atoms, R^{2a} , R^{4a} , and R^{5a} are a hydrogen
 10 atom, R^{3a} is a C1-C3 alkyl group optionally having one or
 more fluorine atoms or a halogen atom, and R^{6a} is a C1-C3
 alkyl group having one or more fluorine atoms, a C1-C3
 alkoxy group having one or more fluorine atoms, a C1-C3
 alkylsulfanyl group having one or more fluorine atoms, a
 15 C1-C3 alkylsulfinyl group having one or more fluorine atoms,
 or a C1-C3 alkylsulfonyl group having one or more fluorine
 atoms, or a halogen atom;

[0051]

A compound of the formula (1-0):



wherein:

A^{1j} represents $-NR^{8j}-$, an oxygen atom, or a sulfur
 atom;

A^{2j} represents a nitrogen atom or =CR^{9j}-;

A^{3j} represents a nitrogen atom or =CR^{10j}-;

R^{1j} represents a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from Group X^j, or a C3-C6 alicyclic hydrocarbon group optionally having one or more atoms or groups selected from Group Y^j;

R^{2j}, R^{3j}, R^{4j}, and R^{5j} are same or different and independently represent a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, -OR^{11j}, -NR^{11j} R^{12j}, a cyano group, a nitro group, a halogen atom, or a hydrogen atom: provided that at least two of R^{2j}, R^{3j}, R^{4j}, and R^{5j} represent a hydrogen atom;

R^{6j} and R^{7j} are same or different and independently represent a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms, -OR^{11j}, -S(O)_mR^{11j}, -NR^{11j} R^{12j}, -CO₂R^{11j}, -C(O)R^{11j}, an aldehyde group, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

R^{8j} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group having one -OR^{11j}, or a cyclopropyl group;

R^{9j} and R^{10j} are same or different and independently

represent a C1-C3 chain hydrocarbon group, a halogen atom,
or a hydrogen atom;

R^{11j} and R^{12j} are same or different and independently represent a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a hydrogen atom; and


m represents 0, 1, or 2, and n represents 0, 1, or 2;

(except in cases as follows: both R^{6j} and R^{7j} are a hydrogen atom; in $-S(O)_m R^{11j}$, when m is 1 or 2, R^{11j} represents a hydrogen atom; and when A^{1j} represents a methylimino group, A^{2j} and A^{3j} represent a methine group, R^{1j} represents a methyl group, R^{2j} , R^{3j} , R^{4j} , and R^{5j} represent a hydrogen atom, and n represents 0 or 1, both R^{6j} and R^{7j} are a chlorine atom and one of R^{6j} and R^{7j} is a chlorine atom and the other is a hydrogen atom)

the Group X^j consists of -OR^{11j} and a halogen atom; and
the Group Y^j consists of a C1-C6 chain hydrocarbon
group, a -OR^{11j}, and a halogen atom.

[0052]

A compound of the formula (Z):



wherein, R^{1j} , R^{2j} , R^{3j} , R^{4j} , R^{5j} , R^{6j} , A^{1j} , A^{3j} , and n are as defined in the formula (1-0);

A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-, an

oxygen atom, or a sulfur atom, and R^{8j} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

5 A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$ or a sulfur atom, and R^{8j} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$ or a sulfur atom;

A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$;

10 A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$, and R^{8j} is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a cyclopropyl group, or a hydrogen atom;

15 A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$, and R^{8j} is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms or a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

20 A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$, and R^{8j} is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a cyclopropyl group;

25 A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$,

and R^{8j} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-, and R^{8j} is a C1-C6 chain hydrocarbon group;

5 A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-, and R^{8j} is a C1-C6 alkyl group;

A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-, and R^{8j} is a C1-C3 alkyl group;

10 A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-, and R^{8j} is a methyl group, an ethyl group, a propyl group, or an isopropyl group;

A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-, and R^{8j} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a methoxymethyl group, an ethoxymethyl group, a cyclopropyl group, or a hydrogen atom;

15 A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-, and R^{8j} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a methoxymethyl group, an ethoxymethyl group, or a cyclopropyl group;

20 A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-, and R^{8j} is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group;

A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-, and R^{8j} is a methyl group;

25 A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-,

and R^{8j} is an ethyl group;

A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-,
and R^{8j} is a propyl group;

A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-,
5 and R^{8j} is an isopropyl group;

A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-,
and R^{8j} is a methyl group or a hydrogen atom;

A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-,
and R^{8j} is a hydrogen atom;

10 A compound of the formula (Z) wherein A^{1j} is -NR^{8j}-,
and R^{8j} is a methoxymethyl group, an ethoxymethyl group, or
a hydrogen atom;

A compound of the formula (Z) wherein A^{1j} is an oxygen
atom;

15 A compound of the formula (Z) wherein A^{1j} is a sulfur
atom;

A compound of the formula (Z) wherein A^{1j} is -NR^{8j}- or
a sulfur atom;

A compound of the formula (Z) wherein A^{1j} is -NR^{8j}- or
20 a sulfur atom, and R^{8j} is a methyl group, an ethyl group, a
propyl group, or an isopropyl group;

A compound of the formula (Z) wherein A^{1j} is -NR^{8j}- or
a sulfur atom, and R^{8j} is a methoxymethyl group, an
ethoxymethyl group, or a hydrogen atom;

25 A compound of the formula (Z) wherein A^{3j} is a

nitrogen atom or $=CR^{10j}-$, and R^{10j} is a halogen atom or a hydrogen atom;

A compound of the formula (Z) wherein A^{3j} is $=CR^{10j}-$, and R^{10j} is a halogen atom or a hydrogen atom;

5 A compound of the formula (Z) wherein A^{3j} is $=CR^{10j}-$, and R^{10j} is a hydrogen atom;

A compound of the formula (Z) wherein A^{3j} is a nitrogen atom;

10 A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$, and A^{3j} is $=CR^{10j}-$;

A compound of the formula (Z) wherein A^{1j} is a sulfur atom, and A^{3j} is $=CR^{10j}-$;

A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$, and A^{3j} is a nitrogen atom;

15 A compound of the formula (Z) wherein A^{1j} is a sulfur atom, and A^{3j} is a nitrogen atom;

20 A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$ or a sulfur atom, R^{8j} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, A^{3j} is a nitrogen atom or $=CR^{10j}-$, and R^{10j} is a hydrogen atom;

A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$ or a sulfur atom, R^{8j} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, and A^{3j} is a nitrogen atom;

25 A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$ or

a sulfur atom, R^{8j} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, A^{3j} is $=CR^{10j}-$, and R^{10j} is a hydrogen atom;

A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$ or
5 a sulfur atom, R^{8j} is a methyl group, and A^{3j} is a nitrogen atom;

A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$, R^{8j} is a methyl group, and A^{3j} is a nitrogen atom;

A compound of the formula (Z) wherein R^{1j} is a C1-C6
10 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (Z) wherein R^{1j} is a C1-C6 chain hydrocarbon group optionally having one or more fluorine atoms;

15 A compound of the formula (Z) wherein R^{1j} is a C1-C6 chain hydrocarbon group;

A compound of the formula (Z) wherein R^{1j} is a C1-C6 alkyl group;

A compound of the formula (Z) wherein R^{1j} is a C1-C3
20 alkyl group;

A compound of the formula (Z) wherein R^{1j} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, a sec-butyl group, an isobutyl group, a tert-butyl group, a cyclopropyl group, a trifluoromethyl
25 group, or a 2,2,2-trifluoroethyl group;

A compound of the formula (Z) wherein R^{1j} is a methyl group;

A compound of the formula (Z) wherein R^{1j} is an ethyl group;

5 A compound of the formula (Z) wherein R^{1j} is a propyl group;

A compound of the formula (Z) wherein R^{1j} is an isopropyl group;

10 A compound of the formula (Z) wherein R^{1j} is a trifluoromethyl group;

A compound of the formula (Z) wherein R^{1j} is a 2,2,2-trifluoroethyl group;

15 A compound of the formula (Z) wherein R^{1j} is a C3-C6 alicyclic hydrocarbon group optionally having one or more atoms or groups selected from Group Y^j ;

A compound of the formula (Z) wherein R^{1j} is a C3-C6 alicyclic hydrocarbon group;

20 A compound of the formula (Z) wherein R^{1j} is a cyclopropyl group, a cyclobutyl group, or a cyclopentyl group;

A compound of the formula (Z) wherein R^{2j} , R^{4j} , and R^{5j} are same or different and are independently a fluorine atom, a chlorine atom, or a hydrogen atom, and R^{3j} is a fluorine atom, a chlorine atom, a bromine atom, a trifluoromethyl group, or a hydrogen atom;

25

A compound of the formula (Z) wherein R^{2j} , R^{4j} , and R^{5j} are a hydrogen atom, R^{3j} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

5 A compound of the formula (Z) wherein R^{2j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{3j} is a C1-C6 alkyl group;

A compound of the formula (Z) wherein R^{2j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{3j} is a optionally having one or more fluorine atoms C1-C6 alkyl group;

10 A compound of the formula (Z) wherein R^{2j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{3j} is a C1-C3 alkyl group;

A compound of the formula (Z) wherein R^{2j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{3j} is a C1-C3 alkyl group optionally having one or more fluorine atoms;

15 A compound of the formula (Z) wherein R^{2j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{3j} is a chlorine atom, a bromine atom, a trifluoromethyl group, or a hydrogen atom;

A compound of the formula (Z) wherein R^{2j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{3j} is a fluorine atom;

20 A compound of the formula (Z) wherein R^{2j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{3j} is a chlorine atom;

A compound of the formula (Z) wherein R^{2j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{3j} is a bromine atom;

A compound of the formula (Z) wherein R^{2j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{3j} is a methyl group;

25 A compound of the formula (Z) wherein R^{2j} , R^{4j} , and R^{5j}

are a hydrogen atom, and R^{3j} is a trifluoromethyl group;

A compound of the formula (Z) wherein R^{2j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{3j} is a trifluoromethoxy group;

A compound of the formula (Z) wherein R^{2j} , R^{3j} , R^{4j} , and
5 R^{5j} are a hydrogen atom;

A compound of the formula (Z) wherein R^{6j} is a halogen atom;

A compound of the formula (Z) wherein R^{6j} is a trifluoromethyl group, a pentafluoroethyl group, a
10 heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

A compound of the formula (Z) wherein R^{6j} is a bromine atom;

15 A compound of the formula (Z) wherein R^{6j} is an iodine atom;

A compound of the formula (Z) wherein R^{6j} is a trifluoromethyl group;

20 A compound of the formula (Z) wherein R^{6j} is a pentafluoroethyl group;

A compound of the formula (Z) wherein R^{6j} is a heptafluoroisopropyl group;

A compound of the formula (Z) wherein R^{6j} is a trifluoromethoxy group;

25 A compound of the formula (Z) wherein R^{6j} is a

trifluoromethylsulfanyl group;

A compound of the formula (Z) wherein R^{6j} is a trifluoromethylsulfinyl group;

5 A compound of the formula (Z) wherein R^{6j} is a trifluoromethylsulfonyl group;

A compound of the formula (Z) wherein R^{2j} , R^{3j} , R^{4j} , and R^{5j} are same or different and are independently a fluorine atom, a chlorine atom, or a hydrogen atom, and R^{6j} is a trifluoromethyl group or a pentafluoroethyl group;

10 A compound of the formula (Z) wherein R^{2j} , R^{3j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{6j} is a trifluoromethyl group;

A compound of the formula (Z) wherein R^{2j} , R^{3j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{6j} is a pentafluoroethyl group or a heptafluoroisopropyl group;

15 A compound of the formula (Z) wherein R^{2j} , R^{3j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{6j} is a trifluoromethoxy group;

A compound of the formula (Z) wherein R^{2j} , R^{3j} , R^{4j} , and R^{5j} are a hydrogen atom, and R^{6j} is a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

20

A compound of the formula (Z) wherein A^{1j} is $-NR^{8j}-$ or a sulfur atom, R^{8j} is a C1-C6 alkyl group, A^{3j} is $=CR^{10j}-$, R^{10j} is a hydrogen atom, R^{1j} is a C1-C3 alkyl group optionally having one or more fluorine atoms, R^{2j} , R^{4j} , and

25

R^{5j} are a hydrogen atom, R^{3j} is a C1-C3 alkyl group optionally having one or more fluorine atoms or a halogen atom, and R^{6j} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more
5 fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or a halogen atom; and

A compound of the formula (Z) wherein A^{1j} is -NR^{8j} - or
10 a sulfur atom, R^{8j} is a C1-C6 alkyl group, A^{3j} is a nitrogen atom, R^{1j} is a C1-C3 alkyl group optionally having one or more fluorine atoms, R^{2j}, R^{4j}, and R^{5j} are a hydrogen atom, R^{3j} is a C1-C3 alkyl group optionally having one or more fluorine atoms or a halogen atom, and R^{6j} is a C1-C3 alkyl
15 group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine
20 atoms, or a halogen atom.


[0053]

The present compound can be produced, for example, by the following Production methods A-F.

[0054]

25 (Production method A)

The present compound (1) can be produced by reacting Compound (1F) with Compound (5). Alternatively, the present compound (1) can be produced by reacting Compound (F) with Compound (5) to give Compound (B), followed by cyclizing Compound (B):




wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^1 , A^2 , A^3 , and n are as defined above.

[0055]

10 (Production method B)

The present compound (1) can be produced by reacting Compound (1F) with Compound (M3). Alternatively, the present compound (1) can be produced by reacting Compound (F) with Compound (5) to give Compound (B), followed by cyclizing Compound (B):

15 cyclizing Compound (B):




wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^1 , A^2 , A^3 and n are as defined above.

[0056]

5 (Production method C)

The present compound (1) can be produced by reacting Compound (F) with Compound (3). Alternatively, the present compound (1) can be produced by reacting Compound (1F) with Compound (3) to give Compound (1J), followed by cyclizing Compound (1J):




wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^1 , A^2 , A^3 , and n are as

defined above.

[0057]

(Production method D)

Compound (12) of the formula (1) wherein A¹ is a
5 sulfur atom can be produced by reacting Compound (K) with a
sulfating agent:




wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, A², A³, and n are as
defined above.

10 [0058]

(Production method E)

Compound (23) of the formula (1) wherein n is 0 can be
produced, for example, by the following method:




15 wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, A¹, A², A³, and n are as
defined above, and V² represents a fluorine atom or a

chlorine atom.

[0059]

(Production method F)

Compound (23) of the formula (1) wherein n is 0 can be
5 produced, for example, by the following method:



wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, A¹, A², A³, n, and V² are
as defined above.

[0060]

10 The production method of the present compound is
illustrated below.


[0061]

The production methods A-F of the present compound are
further illustrated bellow. In addition, the production
15 methods which the present compound is converted to other
present compound are illustrated. The present compound can
be produced, for example, by the following Production
methods 1-26.

[0062]

20 (Production method 1) (Step (C-1))

Compound (4) of the formula (1) wherein A¹ is -NR⁸-
can be produced by reacting Compound (2) with Compound (3):



wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , A^2 , A^3 , and n are as defined above.

The reaction is usually performed in the presence of a
 5 base, acid, sulfite, or disulfite.

The reaction is usually performed in the presence of a solvent.

Examples of the base to be used in the reaction include hydrogen carbonates such as sodium hydrogen
 10 carbonate and potassium hydrogen carbonate; carbonates such as sodium carbonate and potassium carbonate; and mixtures thereof.

Examples of the acid to be used in the reaction include sulfonic acids such as p-toluenesulfonic acid; and
 15 carboxylic acids such as acetic acid.

Examples of the sulfite to be used in the reaction include sodium sulfite and potassium sulfite.

Examples of the disulfite to be used in the reaction include sodium disulfite and potassium disulfite.

20 Examples of the solvent to be used in the reaction include ethers such as tetrahydrofuran (hereinafter referred to as THF), ethylene glycol dimethyl ether, tert-

butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as N,N-dimethylformamide (hereinafter referred to as DMF) and N-methylpyrrolidone (hereinafter referred to as NMP); sulfoxides such as dimethylsulfoxide (hereinafter referred to as DMSO); nitrogen-containing aromatic compounds such as pyridine and quinoline; and mixtures thereof.

An oxidizing agent can be added to the reaction as necessary.

Examples of the oxidizing agent to be used in the reaction include an oxygen, copper (II) oxide, and 2,3-dichloro-5,6-dicyano-p-benzoquinon.

The amount of Compound (3) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (2). The amount of the base to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (2). The amount of the acid to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (2). The amount of the sulfite to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (2). The amount of the disulfite to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (2). The amount of the

oxidizing agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (2).


The reaction temperature of the reaction is usually within a range of 30 to 200 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (4) can be isolated by post-treatments, for example, pouring water into the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (4) can be further purified by chromatography, recrystallization, and the like.

[0063]

(Production method 2) (Step A-1)

Compound (4) of the formula (1) wherein A¹ is -NR⁸- can be produced by reacting Compound (2) with Compound (5) in the presence of a dehydration-condensing agent:



wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, A², A³, and n are as defined above.

The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, tert-butyl methyl ether, ethylene glycol dimethyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; nitrogen-containing aromatic compounds such as pyridine and quinoline; and mixtures thereof.

Examples of the dehydration-condensing agent to be used in the reaction include carbodiimides such as 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (hereinafter referred to as WSC), and 1,3-dicyclohexylcarbodiimide; and (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (hereinafter referred to as BOP reagent).

A catalyst can be added to the reaction as necessary.

Examples of the catalyst to be used in the reaction include 1-hydroxybenzotriazole (hereinafter referred to as HOBT).

The amount of Compound (5) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (2). The amount of the dehydration-condensing agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of

Compound (2). The amount of the catalyst to be used in the reaction is usually 0.01 to 0.1 moles based on 1 mole of Compound (2).

The reaction temperature of the reaction is usually
5 within a range of 30 to 200 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (4) can be isolated by post-treatments, for example, pouring water into the reaction mixture and then extracting the mixture
10 with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (4) can be further purified by chromatography, recrystallization, and the like.

Alternatively, Compound (4) can be produced according
15 to the above method using Compound (M3) instead of Compound (5).

When Compound (M3) is used, the reaction is usually performed without a dehydration-condensing agent.

A base can be added to the reaction as necessary.


20 Examples of the base to be used in the reaction include carbonates of alkali metal such as sodium carbonate and potassium carbonate; tertiary amines such as triethylamine and diisopropylethylamine; and nitrogen-containing aromatic compounds such as pyridine and 4-
25 dimethylaminopyridine.

The amount of Compound (M3) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (2). The amount of the base to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (2).

5 [0064]

(Production method 3) (Step (A-3), Step (B-3))

Compound (4) of the formula (1) wherein A^1 is $-NR^8-$ can be produced by dehydration-condensing Compound (6):



10 wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , A^2 , A^3 , and n are as defined above.

The reaction is usually performed in the presence of a solvent.

15 Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; 20 alcohols such as methanol, ethanol, propanol, butanol, and

pentanol; and mixtures thereof.

In the reaction, an acid or a dehydrating agent can be used as necessary. Examples of the acid to be used in the reaction include sulfonic acids such as p-toluenesulfonic acid, and carboxylic acids such as acetic acid. Examples of the dehydrating agent to be used in the reaction include phosphorous oxychloride, acetic anhydride, and trifluoroacetic anhydride.

The amount of the acid or the dehydrating agent to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (6).


The reaction temperature of the reaction is usually within a range of 30 to 200 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (4) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (4) can be further purified by chromatography, recrystallization, and the like.

[0065]

(Production method 4) (Step (A-3), Step (B-3))

Compound (4) of the formula (1) wherein A^1 is $-NR^8-$ can be produced by reacting Compound (6) in the presence of a base:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , A^2 , A^3 , and n are as defined above.

The reaction is usually performed in the presence of a
5 solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic
10 hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; alcohols such as methanol, ethanol, propanol, butanol,
15 tert-butanol, and pentanol; and mixtures thereof.

Examples of the base to be used in the reaction include tripotassium phosphate.

The amount of the base to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (6).

20 The reaction temperature of the reaction is usually within a range of 30 to 200 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (4) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated


5 Compound (4) can be further purified by chromatography, recrystallization, and the like.

[0066]

(Production method 5)

Compound (4) of the formula (1) wherein A¹ is -NR⁸- can be produced by reacting Compound (32) with Compound

10 (33) in the presence of a base.



wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, A², A³, and n are as defined above, and L represents a leaving group such as a

15 chlorine atom, a bromine atom, an iodine atom, a trifluoromethylsulfonyloxy group and a methylsulfonyloxy group.

The reaction is usually performed in the presence of a solvent.

20 Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aromatic

hydrocarbons such as toluene and xylene; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; and mixtures thereof.

Examples of the base to be used in the reaction
5 include hydrides of alkali metal or alkali earth metal such as sodium hydride, potassium hydride, and calcium hydride; inorganic bases such as sodium carbonate and potassium carbonate; and organic bases such as triethylamine.

The amount of Compound (33) to be used in the reaction
10 is usually 1 to 5 moles based on 1 mole of Compound (32).

The amount of the base to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (32).

The reaction temperature of the reaction is usually within a range of 0 to 100 °C. The reaction time of the
15 reaction is usually within a range of 0.1 to 24 hours.


After the completion of the reaction, Compound (4) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated
20 Compound (4) can be further purified by chromatography, recrystallization, and the like.

[0067]

(Production method 6) (Step (A-1))

Compound (8) of the formula (1) wherein A¹ is an
25 oxygen atom can be produced by reacting Compound (7) with

Compound (5) in the presence of an acid:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^2 , A^3 , and n are as defined above.

5 The reaction is usually performed in the presence or absence of a solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic
 10 hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene and dichlorobenzene; and mixtures thereof.

Examples of the acid to be used in the reaction
 15 include polyphosphoric acid and trimethylsilyl polyphosphate.

When polyphosphoric acid is used as an acid, though the reaction is usually performed in the absence of a solvent, the reaction may be performed in the presence of a
 20 solvent.

The amount of Compound (5) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (7).

The amount of the acid to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (7).

The reaction temperature of the reaction is usually within a range of 50 to 200 °C. The reaction time of the
5 reaction is usually within a range of 0.5 to 24 hours.

After the completion of the reaction, Compound (8) can be isolated by post-treatments, for example, pouring the reaction mixture into water and then extracting the mixture with an organic solvent, followed by drying and
10 concentrating the organic layer. The isolated Compound (8) can be further purified by chromatography, recrystallization, and the like.

Alternatively, Compound (8) can be produced according to the above method using Compound (M3) instead of Compound
15 (5).

When Compound (M3) is used, the reaction is usually performed without a dehydration-condensing agent.

A base can be added to the reaction as necessary.

Examples of the base to be used in the reaction
20 include carbonates of alkali metal such as sodium carbonate and potassium carbonate; tertiary amines such as triethylamine and diisopropylethylamine; and nitrogen-containing aromatic compounds such as pyridine and 4-dimethylaminopyridine.


25 The amount of Compound (M3) to be used in the reaction

is usually 1 to 3 moles based on 1 mole of Compound (7).
The amount of the base to be used in the reaction is
usually 1 to 10 moles based on 1 mole of Compound (7).

[0068]

5 (Production method 7) (Step (C-3))

Compound (8) of the formula (1) wherein A¹ is an
oxygen atom can be produced by oxidizing Compound (9):



wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, A², A³, and n are as
10 defined above.

The reaction is usually performed in the presence of a
solvent.

Examples of the solvent to be used in the reaction
include ethers such as THF, ethylene glycol dimethyl ether,
15 tert-butyl methyl ether, and 1,4-dioxane; aliphatic
hydrocarbons such as hexane, heptane, and octane; aromatic
hydrocarbons such as toluene and xylene; halogenated
hydrocarbons such as dichloromethane, chloroform, and
chlorobenzene; esters such as ethyl acetate and butyl
20 acetate; alcohols such as methanol and ethanol; nitriles
such as acetonitrile; acid amides such as DMF and NMP;
sulfoxides such as DMSO; acetic acid; and mixtures thereof.

Examples of the oxidizing agent to be used in the reaction include a metal oxidizing agent such as lead (IV) acetate and lead (IV) oxide, and a hypervalent iodine compound such as iodobenzene diacetate.

5 The amount of the oxidizing agent to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (9).


10 The reaction temperature of the reaction is usually within a range of 0 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (8) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated
15 Compound (8) can be further purified by chromatography, recrystallization, and the like.

[0069]

(Production method 8) (Step (A-3), Step (B-3))

20 Compound (8) of the formula (1) wherein A¹ is an oxygen atom can be produced by reacting Compound (10) in the presence of a dehydration-condensing agent:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^2 , A^3 , and n are as defined above.

The reaction is usually performed in the presence of a solvent.

5 Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as dichloromethane, chloroform, carbon
10 tetrachloride, and chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; and mixtures thereof. Among of them, carbon tetrachloride can be used as a dehydration-condensing agent.

 Examples of the dehydration-condensing agent to be
15 used in the reaction include a mixture of triphenylphosphine, a base and carbon tetrachloride or carbon tetrabromide, and a mixture of triphenylphosphine and azodiester such as diethyl azodicarboxylate ester.

 Examples of the base to be used in the reaction
20 include tertiary amines such as triethylamine and diisopropylethylamine.

 The amount of the dehydration-condensing agent to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (10). The amount of the base to be used
25 in the reaction is usually 1 to 5 moles based on 1 mole of

Compound (10).


The reaction temperature of the reaction is usually within a range of -30 to 100 °C. The reaction time of the reaction is usually within a range of 0.5 to 24 hours.

5 After the completion of the reaction, Compound (8) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (8) can be further purified by chromatography,
10 recrystallization, and the like.

[0070]

(Production method 9) (Step (A-3), Step (B-3))

Compound (8) of the formula (1) wherein A¹ is an oxygen atom can be produced by reacting Compound (10) in
15 the presence of an acid:



wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, A², A³, and n are as defined above.

The reaction is usually performed in the presence of a
20 solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether,

tert-butyl methyl ether, and 1,4-dioxane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as dichloromethane, chloroform, and chlorobenzene; and mixtures thereof.

5 Examples of the acid include sulfonic acids such as p-toluenesulfonic acid and polyphosphoric acid.

 The amount of the acid to be used in the reaction is usually 0.1 to 3 moles based on 1 mole of Compound (10).


10 The reaction temperature of the reaction is usually within a range of 50 to 200 °C. The reaction time of the reaction is usually within a range of 1 to 24 hours.

 After the completion of the reaction, Compound (8) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by
15 drying and concentrating the organic layer. The isolated Compound (8) can be further purified by chromatography, recrystallization, and the like.

[0071]

(Production method 10) (Step (C-1))

20 Compound (12) of the formula (1) wherein A¹ is a sulfur atom can be produced by reacting Compound (11) with Compound (3):



wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, A², A³, and n are as defined above.

The reaction is usually performed in the presence of a
 5 base, acid, sulfite, or disulfite.

Examples of the base to be used in the reaction include hydrogen carbonates such as sodium hydrogen carbonate and potassium hydrogen carbonate; carbonates such as sodium carbonate and potassium carbonate; and mixtures
 10 thereof.

Examples of the acid to be used in the reaction include sulfonic acids such as p-toluenesulfonic acid; and carboxylic acids such as acetic acid.

Examples of the sulfite to be used in the reaction
 15 include sodium sulfite and potassium sulfite.

Examples of the disulfite to be used in the reaction include sodium disulfite and potassium disulfite.

The reaction is usually performed in the presence of a solvent.

20 Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic

hydrocarbons such as hexane, heptane, and octane; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate nitriles such as acetonitrile; acid amides such as DMF, and NMP; sulfoxides
5 such as DMSO; aromatic hydrocarbons such as toluene, xylene, nitrobenzene; and mixtures thereof.

An oxidizing agent can be added to the reaction as necessary.

10 Examples of the oxidizing agent to be used in the reaction include an oxygen, copper (II) oxide, and 2,3-dichloro-5,6-dicyano-p-benzoquinon.

The amount of Compound (3) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (11). The amount of the acid to be used in the reaction is
15 usually 1 to 5 moles based on 1 mole of Compound (11). The amount of the sulfite to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (11). The amount of the disulfite to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (11). The amount of
20 the oxidizing agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (11). The reaction temperature of the reaction is usually within a range of 50 to 200 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.


25 After the completion of the reaction, Compound (12)

can be isolated by post-treatments, for example, adding the reaction mixture to water and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (12) can be further purified by chromatography, recrystallization, and the like.

[0072]

(Production method 11) (Step (C-1))

Compound (12) of the formula (1) wherein A¹ is a sulfur atom can be produced by reacting hydrochloride of Compound (11) with Compound (3):



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^2 , A^3 , and n are as defined above.

The reaction is performed in the presence of a base and usually in the presence of a solvent.

Examples of the base to be used in the reaction include tertiary amines such as diisopropylethylamine and triethylamine.

Examples of the solvent to be used in the reaction include sulfoxides such as DMSO; aromatic hydrocarbons such as nitrobenzene; and mixtures thereof.

The amount of Compound (3) to be used in the reaction is usually 0.5 to 3 moles based on 1 mole of Compound (11). The amount of the base to be used in the reaction is usually 1 to 2 moles based on 1 mole of Compound (11).


5 The reaction temperature of the reaction is usually within a range of 50 to 200 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (12) can be isolated by post-treatments, for example, adding the
 10 reaction mixture to water and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (12) can be further purified by chromatography, recrystallization, and the like.

15 [0073]

(Production method 12)

Compound (14) of the formula (1) wherein R⁶ is a cyano group or a C1-C6 alkyl group can be produced by reacting
 Compound (13) with a cyanating agent or di(C1-C6 alkyl)
 20 zinc in the presence of a palladium compound:



wherein, R¹, R², R³, R⁴, R⁵, R⁷, A¹, A², and A³ are as defined

above, V^1 represents a leaving group such as a bromine atom and an iodine atom, and R^{6z} represents a cyano group or a C1-C6 alkyl group.

5 The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic
10 hydrocarbons such as toluene and xylene; alcohols such as methanol and ethanol; acid amides such as DMF and NMP; sulfoxides such as DMSO; and mixtures thereof.

Examples of the cyanating agent to be used in the reaction include zinc cyanide. Examples of di(C1-C6
15 alkyl)zinc to be used in the reaction include dimethyl zinc, diethyl zinc, and diisopropyl zinc.

Examples of the palladium compound to be used in the reaction include tetrakis(triphenylphosphine)palladium.

The amount of the cyanating agent or di(C1-C6 alkyl)
20 zinc to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (13). The amount of the palladium compound to be used in the reaction is usually 0.01 to 0.5 moles based on 1 mole of Compound (13).

The reaction temperature of the reaction is usually
25 within a range of 50 to 200 °C. The reaction time of the


reaction is usually within a range of 0.5 to 24 hours.

After the completion of the reaction, Compound (14) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (14) can be further purified by chromatography, recrystallization, and the like.

[0074]

(Production method 13)

Compound (16) of the formula (1) wherein R^6 is $C(O)R^{6y}$ can be produced by reacting Compound (15) with a organometallic agent, then a carbonylating agent:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^7 , A^1 , A^2 , and A^3 are as defined above, R^{6y} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a hydrogen atom, or $-OR^{11}$, and R^{11} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a hydrogen atom.

The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether,

tert-butyl methyl ether, and 1,4-dioxane.

Examples of the organometallic agent to be used in the reaction include alkylolithiums such as n-butyllithium and tert-butyllithium, organic magnesium halides such as
5 isopropylmagnesium chloride.

Examples of the carbonylating agent to be used in the reaction include DMF, 1-formylpiperidine, 4-formylmorpholine, ethyl formate, ethyl acetate, methyl trifluoroacetate, methyl chloroformate, ethyl chloroformate,
10 and carbon dioxide.

The amount of the organometallic agent to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (15). The amount of the carbonylating agent to be used in the reaction is usually 1 to 10 moles based
15 on 1 mole of Compound (15).


The reaction temperature of the reaction is usually within a range of -100 to 0 °C. The reaction time of the reaction is usually within a range of 0.5 to 24 hours.

After the completion of the reaction, Compound (16)
20 can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (16) can be further purified by chromatography, recrystallization, and the like.

25 [0075]

(Production method 14)

Compound (18) of the formula (1) wherein R⁶ is a C1-C3 perfluoroalkyl group can be produced by reacting Compound (13) with Compound (17) in the presence of copper iodide:



wherein, R¹, R², R³, R⁴, R⁵, R⁷, A¹, A², A³, and V¹ are as defined above, and R^f represents a C1-C3 perfluoroalkyl group.

10 The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include aromatic hydrocarbons such as toluene and xylene; acid amides such as DMF and NMP; and mixtures thereof.

15 The amount of Compound (17) to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (13). The amount of copper iodide to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (13).

20 The reaction temperature of the reaction is usually within a range of 50 to 200 °C. The reaction time of the reaction is usually within a range of 0.5 to 24 hours.


After the completion of the reaction, Compound (18) can be isolated by post-treatments, for example, extracting

the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (18) can be further purified by chromatography, recrystallization, and the like.

5 [0076]

(Production method 15)

Compound (20) of the formula (1) wherein R⁶ is a phenyl group optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more
 10 halogen atoms, or a 6-membered heterocyclic group optionally having one or more halogen atoms can be produced by reacting Compound (13) with boronic acid compound (19) in the presence of a palladium compound:



15 wherein, R¹, R², R³, R⁴, R⁵, R⁷, A¹, A², A³, and V¹ are as defined above, and R^{Ar} represents a phenyl group optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms, or a 6-membered heterocyclic group optionally having one or more
 20 halogen atoms.

The reaction is usually performed in the presence of a solvent.

The reaction can be performed in the presence of a base and /or a phase transfer catalyst as necessary.

Examples of the base to be used in the reaction include inorganic salts such as sodium acetate, potassium acetate, potassium carbonate, tripotassium phosphate, and sodium hydrogen carbonate.

Examples of the phase transfer catalyst to be used in the reaction include quaternary ammonium salts such as tetrabutylammonium bromide and benzyltriethylammonium bromide.

The reaction can be performed in the presence of a ligand as necessary.

Examples of the ligand to be used in the reaction include 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; alcohols such as methanol and ethanol; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; acid amides such as DMF and NMP; water; and mixtures thereof.

Examples of the palladium compound to be used in the reaction include palladium acetate, tetrakis(triphenylphosphine) palladium, {1,1'-bis(diphenylphosphino)ferrocene}dichloropalladium-methylene

chloride complex, and bis(triphenylphosphine)palladium dichloride, and tris(dibenzylideneacetone)dipalladium (0).

The amount of boronic acid compound (19) to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (13). The amount of the palladium compound to be used in the reaction is usually 0.01 to 0.5 moles based on 1 mole of Compound (13).


The reaction temperature of the reaction is usually within a range of 50 to 200 °C. The reaction time of the reaction is usually within a range of 0.5 to 24 hours.

After the completion of the reaction, Compound (20) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (20) can be further purified by chromatography, recrystallization, and the like.

[0077]

(Production method 16) (Step (E-4))

Compound (23) of the formula (1) wherein n is 0 can be produced by reacting Compound (21) with Compound (22) in the presence of a base:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^1 , A^2 , A^3 , and V^2 are as defined above.

The reaction is usually performed in the presence of a solvent.

5 Examples of the solvent to be used in the reaction include water; ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aromatic hydrocarbons such as toluene and xylene; alcohols such as methanol and ethanol; nitriles such as acetonitrile; acid
10 amides such as DMF and NMP; sulfoxides such as DMSO; and mixtures thereof.

Examples of the base to be used in the reaction include alkali metal hydrides such as sodium hydride.

The amount of Compound (22) to be used in the reaction
15 is usually 1 to 10 moles based on 1 mole of Compound (21). The amount of the base to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (21).

The reaction temperature of the reaction is usually within a range of 0 to 150 °C. The reaction time of the
20 reaction is usually within a range of 0.5 to 24 hours.


After the completion of the reaction, Compound (23) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated
25 Compound (23) can be further purified by chromatography,

recrystallization, and the like.

[0078]

(Production method 17) (Step (F-2))

Compound (23) of the formula (1) wherein n is 0 can be
 5 produced by reacting Compound (24) with Compound (25) in
 the presence of a base:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^1 , A^2 , A^3 , and L are as defined above.

10 The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aromatic
 15 hydrocarbons such as toluene and xylene; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; and mixtures thereof.

Examples of the base to be used in the reaction include hydrides of alkali metal or alkali earth metal such
 20 as sodium hydride, potassium hydride, and calcium hydride; inorganic bases such as sodium carbonate and potassium carbonate; and organic bases such as triethylamine.

The amount of the base to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (24). The amount of Compound (25) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (24).


5 The reaction temperature of the reaction is usually within a range of 0 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (23) can be isolated by post-treatments, for example, extracting
10 the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (23) can be further purified by chromatography, recrystallization, and the like.

[0079]

15 (Production method 18)

Compound (26) of the formula (1) wherein n is 1 can be produced by oxidizing Compound (23):



wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, A¹, A², and A³ are as
20 defined above.

The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include halogenated hydrocarbons such as dichloromethane and chloroform; alcohols such as methanol and ethanol; acetic acid; water; and mixtures thereof.

5 Examples of the oxidizing agent to be used in the reaction include sodium periodate or m-chloroperbenzoic acid.

10 The amount of the oxidizing agent to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (23). Preferably, the amount of the oxidizing agent to be used in the reaction is 1 to 1.2 moles based on 1 mole of Compound (23).


15 The reaction temperature of the reaction is usually within a range of -20 to 80 °C. The reaction time of the reaction is usually within a range of 0.1 to 12 hours.

20 After the completion of the reaction, Compound (26) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by washing the organic layer with an aqueous solution of a reducing agent (e.g. sodium sulfite, thiosodium sulfate) and an aqueous solution of a base (e.g. sodium hydrogen carbonate) as necessary, followed by drying and concentrating the organic layer. The isolated Compound (26) can be further purified by chromatography, 25 recrystallization, and the like.

[0080]

(Production method 19)

Compound (27) of the formula (1) wherein n is 2 can be produced by reacting Compound (23) in the presence of an oxidizing agent:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^1 , A^2 , and A^3 are as defined above.

The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include halogenated hydrocarbons such as dichloromethane and chloroform; alcohols such as methanol and ethanol; acetic acid; water; and mixtures thereof.

Examples of the oxidizing agent to be used in the reaction include m-chloroperbenzoic acid or an aqueous solution of hydrogen peroxide.

The amount of the oxidizing agent to be used in the reaction is usually 2 to 5 moles based on 1 mole of Compound (23). Preferably, the amount of the oxidizing agent to be used in the reaction is 2 to 3 moles based on 1 mole of Compound (23).


The reaction temperature of the reaction is usually within a range of -20 to 120 °C. The reaction time of the reaction is usually within a range of 0.1 to 12 hours.

After the completion of the reaction, Compound (27) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by washing the organic layer with an aqueous solution of a reducing agent (e.g. sodium sulfite, thiosodium sulfate) and an aqueous solution of a base (e.g. sodium hydrogen carbonate) as necessary, followed by drying and concentrating the organic layer. The isolated Compound (27) can be further purified by chromatography, recrystallization, and the like.

[0081]

(Production method 20)

Compound (27) of the formula (1) wherein n is 2 can be produced by reacting Compound (26) in the presence of an oxidizing agent:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^1 , A^2 , and A^3 are as defined above.

The reaction is usually performed in the presence of a

solvent.

Examples of the solvent to be used in the reaction include halogenated hydrocarbons such as dichloromethane and chloroform; alcohols such as methanol and ethanol;
5 acetic acid; water; and mixtures thereof.

Examples of the oxidizing agent to be used in the reaction include m-chloroperbenzoic acid or an aqueous solution of hydrogen peroxide.

The amount of the oxidizing agent to be used in the
10 reaction is usually 1 to 4 moles based on 1 mole of Compound (26). Preferably, the amount of the oxidizing agent to be used in the reaction is 1 to 2 moles based on 1 mole of Compound (26).

The reaction temperature of the reaction is usually
15 within a range of -20 to 120 °C. The reaction time of the reaction is usually within a range of 0.1 to 12 hours.


After the completion of the reaction, Compound (27) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by
20 washing the organic layer with an aqueous solution of a reducing agent (e.g. sodium sulfite, thiosodium sulfate) and an aqueous solution of a base (e.g. sodium hydrogen carbonate) as necessary, followed by drying and concentrating the organic layer. The isolated Compound
25 (27) can be further purified by chromatography,

recrystallization, and the like.

[0082]

(Production method 21) (Step (A-3), Step (B-3))

Compound (12) of the formula (1) wherein A¹ is a
5 sulfur atom can be produced by reacting Compound (35) in
the presence of an acid:



wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, A², A³, and n are as
defined above.

10 The reaction is usually performed in the presence of a
solvent.

Examples of the solvent to be used in the reaction
include ethers such as THF, ethylene glycol dimethyl ether,
tert-butyl methyl ether, and 1,4-dioxane; aromatic
15 hydrocarbons such as toluene and xylene; halogenated
hydrocarbons such as dichloromethane, chloroform, and
chlorobenzene; and mixtures thereof.

Examples of the acid include sulfonic acids such as p-
toluenesulfonic acid, and polyphosphoric acid.

20 The amount of the acid to be used in the reaction is
usually 0.1 to 3 moles based on 1 mole of Compound (35).


The reaction temperature of the reaction is usually within a range of 50 to 200 °C. The reaction time of the reaction is usually within a range of 1 to 24 hours.

After the completion of the reaction, Compound (12) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (12) can be further purified by chromatography, recrystallization, and the like.

[0083]

(Production method 22)

Compound (34-a) of the formula (1) wherein R^6 is SH, Compound (34-b) which is a disulfide form of Compound (34-a), Compound (36) of the formula (1) wherein R^6 is SR^{6x} , and Compound (37) of the formula (1) wherein R^6 is $S(O)_mR^{6x}$ can be produced, for example, by the following method:



wherein, $R^1, R^2, R^3, R^4, R^5, R^{10}, R^7, A^1, A^2, A^3, n, m, V^1$, and L are as defined above.

Step (22-1)

Compound (34-a) and / or Compound (34-b) can be produced by reacting Compound (13) with a thiolating agent in the presence of a catalyst.

5 The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include aromatic hydrocarbons such as toluene and xylene; acid amides such as DMF and NMP; sulfoxides such as DMSO;
10 and mixtures thereof.

Examples of the thiolating agent to be used in the reaction include sodium sulfide, sodium sulfide nonahydrate, and thiourea.

Examples of the catalyst to be used in the reaction
15 include copper (I) chloride, copper (I) bromide, and copper (I) iodide.

The reaction can be performed in the presence of a base as necessary.

Examples of the base to be used in the reaction
20 include potassium carbonate, cesium carbonate, tripotassium phosphate, and triethylamine.

The amount of the thiolating agent to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (13). The amount of the catalyst to be used in
25 the reaction is usually 0.1 to 5 moles based on 1 mole of

Compound (13).

The reaction temperature of the reaction is usually within a range of 50 to 200 °C. The reaction time of the reaction is usually within a range of 0.5 to 24 hours.

5 After the completion of the reaction, Compound (34-a) and / or Compound (34-b) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (34-a) and / or
10 Compound (34-b) can be further purified by chromatography, recrystallization, and the like.

[0084]

Step (22-2)

Compound (36) can be produced by reacting Compound
15 (34-a) and / or Compound (34-b) with Compound (38-1) in the presence of a base.

The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction
20 include alcohols such as methanol and ethanol; ethers such as 1,4-dioxane, diethyl ether, THF, and tert-butyl methyl ether; halogenated hydrocarbons such as dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, and chlorobenzene; hydrocarbons such as toluene, benzene, and
25 xylene; esters such as ethyl acetate and butyl acetate;

nitriles such as acetonitrile; aprotic polar solvents such as DMF, NMP, 1,3-dimethyl-2-imidazolidinone, and dimethylsulfoxide; nitrogen-containing aromatic compounds such as pyridine and quinoline; water; and mixtures thereof.

5 Examples of the base to be used in the reaction include nitrogen-containing heterocyclic compounds such as pyridine, picoline, 2,6-lutidine, DBU, and 1,5-diazabicyclo[4.3.0]-5-nonen; tertiary amines such as triethylamine and N-ethyldiisopropylamine; and inorganic
10 bases such as tripotassium phosphate, potassium carbonate, sodium hydride, sodium hydroxide, and potassium hydroxide.

 The amount of Compound (38-1) to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (34-a) and / or Compound (34-b). The amount of
15 the base to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (34-a) and / or Compound (34-b).

 The reaction temperature of the reaction is usually within a range of 0 to 120 °C. The reaction time of the reaction is usually within a range of 0.5 to 24 hours.

20 After the completion of the reaction, Compound (36) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (36) can be further purified by chromatography,
25 recrystallization, and the like.

[0085]

Step (22-3)

Compound (37) wherein m is 1 or 2 can be produced by oxidizing Compound (36).

5 The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include halogenated hydrocarbons such as dichloromethane and chloroform; alcohols such as methanol and ethanol;
10 acetic acid; water; and mixtures thereof.

Examples of the oxidizing agent to be used in the reaction include m-chloroperbenzoic acid and an aqueous solution of hydrogen peroxide.

The reaction can be performed in the presence of a
15 catalyst as necessary,

Examples of the catalyst to be used in the reaction include sodium tungstate.

The amount of the oxidizing agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of
20 Compound (36).

The reaction temperature of the reaction is usually within a range of -20 to 120 °C. The reaction time of the reaction is usually within a range of 0.1 to 12 hours.


After the completion of the reaction, Compound (37)
25 can be isolated by post-treatments, for example, extracting

the reaction mixture with an organic solvent, followed by washing the organic layer with an aqueous solution of a reducing agent (e.g. sodium sulfite, thiosodium sulfate) and an aqueous solution of a base (e.g. sodium hydrogen carbonate) as necessary, followed by drying and concentrating the organic layer. The isolated Compound (37) can be further purified by chromatography, recrystallization, and the like.

[0086]

10 (Production method 23)

Compound (39) of the formula (1) wherein R^6 is NH_2 , Compound (40) of the formula (1) wherein R^5 is NHR^{10} , and Compound (41) of the formula (1) wherein R^5 is $NR^{10}R^{11}$ can be produced, for example, by the following method:



15

wherein, $R^1, R^2, R^3, R^4, R^5, R^{10}, R^{11}, R^7, A^1, A^2, A^3, n, V^1$, and L are as defined above.

Step (23-1)

Compound (39) can be produced by aminating Compound

(13).

The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction
5 include water; alcohols such as methanol and ethanol;
ethers such as 1,4-dioxane, diethyl ether, and THF; esters
such as ethyl acetate and butyl acetate; halogenated
hydrocarbons such as dichloromethane, chloroform, carbon
tetrachloride, and 1,2-dichloroethane; nitriles such as
10 acetonitrile; aprotic polar solvents such as DMF, NMP, and
dimethylsulfoxide; nitrogen-containing aromatic compounds
such as pyridine and quinoline; and mixtures thereof.

Examples of the aminating agent to be used in the
reaction include ammonia, ammonia water, and lithium amide.

15 Examples of the copper compound to be used in the
reaction include copper, copper (I) iodide, copper (I)
oxide, copper (II) oxide, copper (II) acetylacetonate, copper
(II) acetate, and copper (II) sulfate.

A ligand can be added to the reaction as necessary.

20 Examples of the ligand to be used in the reaction
include acetylacetonate, salen, and phenanthroline.

A base can be added to the reaction as necessary.

Examples of the base to be used in the reaction
include nitrogen-containing heterocyclic compounds such as
25 pyridine, picoline, 2,6-lutidine, DBU, and 1,5-

diazabicyclo[4.3.0]-5-nonen; tertiary amines such as triethylamine and N-ethyldiisopropylamine; and inorganic bases such as tripotassium phosphate, potassium carbonate, cesium carbonate, and sodium hydroxide.

5 The amount of the aminating agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (13). The amount of the copper compound to be used in the reaction is usually 0.02 to 0.5 moles based on 1 mole of Compound (13). The amount of the ligand to be
10 used in the reaction is 0.02 to 2 moles based on 1 mole of Compound (13) as necessary. The amount of the base to be used in the reaction is 1 to 5 moles based on 1 mole of Compound (13) as necessary.

 The reaction temperature of the reaction is usually
15 within a range of 30 to 200 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

 After the completion of the reaction, Compound (39) can be isolated by pouring the reaction mixture into water, extracting with an organic solvent, and concentrating the
20 organic layer; pouring the reaction mixture into water, and collecting the generated solids by filtration; or collecting the solids generated in the reaction mixture by filtration. The isolated Compound (39) can be further purified by recrystallization, chromatography, and the like.

25 [0087]

Step (23-2)

Compound (40) can be produced by reacting Compound (39) with Compound (38-1) in the presence of a base.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aromatic hydrocarbons such as toluene and xylene; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; and mixtures thereof.

Examples of the base to be used in the reaction include hydrides of alkali metal or alkali earth metal such as sodium hydride, potassium hydride, and calcium hydride; inorganic bases such as sodium carbonate and potassium carbonate; and organic bases such as triethylamine.

The amount of Compound (38-1) to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (40). The amount of the base to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (40).

The reaction temperature of the reaction is usually within a range of 0 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (40) can be isolated by pouring the reaction mixture into water, extracting with an organic solvent, and concentrating the

organic layer; pouring the reaction mixture into water, and collecting the generated solids by filtration; or collecting the solids generated in the reaction mixture by filtration. The isolated Compound (40) can be further purified by recrystallization, chromatography, and the like.

[0088]

Step (23-3)

Compound (41) can be produced by reacting Compound (40) with Compound (38-2) in the presence of a base.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aromatic hydrocarbons such as toluene and xylene; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; and mixtures thereof.

Examples of the base to be used in the reaction include hydrides of alkali metal or alkali earth metal such as sodium hydride, potassium hydride, and calcium hydride; inorganic bases such as sodium carbonate and potassium carbonate; and organic bases such as triethylamine.


The amount of Compound (38-2) to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (40). The amount of the base to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (40).

The reaction temperature of the reaction is usually within a range of 0 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (41) can be isolated by pouring the reaction mixture into water, extracting with an organic solvent, and concentrating the organic layer; pouring the reaction mixture into water, and collecting the generated solids by filtration; or collecting the solids generated in the reaction mixture by filtration. The isolated Compound (41) can be further purified by recrystallization, chromatography, and the like. [0089]

(Production method 24)

Compound (43) of the formula (1) wherein R⁶ is OR¹¹ can be produced, for example, by the following method:



wherein, R¹, R², R³, R⁴, R⁵, R⁷, R¹¹, A¹, A², A³, V¹, and n are as defined above.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aromatic hydrocarbons such as toluene and xylene; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides

such as DMSO; and mixtures thereof.

Examples of the base to be used in the reaction include hydrides of alkali metal or alkali earth metal such as sodium hydride, potassium hydride, and calcium hydride; 5 inorganic bases such as sodium carbonate, potassium carbonate, and cesium carbonate; and organic bases such as triethylamine.

A copper compound can be added to the reaction as necessary.

10 Examples of the copper compound to be used in the reaction include copper, copper (I) iodide, copper (I) bromide, and copper (I) chloride.

The amount of Compound (42) to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (13).

15 The amount of the base to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (13).

The reaction temperature of the reaction is usually within a range of 20 to 250 °C. The reaction time of the reaction is usually within a range of 0.1 to 48 hours.


20 After the completion of the reaction, Compound (43) can be isolated by pouring the reaction mixture into water, extracting with an organic solvent, and concentrating the organic layer; pouring the reaction mixture into water, and collecting the generated solids by filtration; or 25 collecting the solids generated in the reaction mixture by

filtration. The isolated Compound (43) can be further purified by recrystallization, chromatography, and the like.

[0090]

(Production method 25) (Step (A-1))

5 Compound (12) of the formula (1) wherein A^1 is a sulfur atom can be produced by reacting Compound (11) with Compound (5):




10 wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^2 , A^3 , and n are as defined above.

Compound (12) can be produced according to the method described in Production method 2 using Compound (11) instead of Compound (2).

[0091]

15 (Production method 26) (Production method D)

Compound (12) of the formula (1) wherein A^1 is a sulfur atom can be produced by reacting Compound (1K) with a sulfating agent:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^2 , A^3 , and n are as defined above.

The reaction is performed in the presence or absence of a solvent.

5 Examples of the solvent to be used in the reaction include ethers such as 1,4-dioxane, diethyl ether, tetrahydrofuran, tert-butyl methyl ether, and Diglyme; halogenated hydrocarbons such as dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, and
10 chlorobenzene; hydrocarbons such as toluene, benzene, and xylene; nitrile such as acetonitrile; pyridines such as pyridine, picoline, and lutidine; and mixtures thereof.

 Examples of the sulfating agent to be used in the reaction include phosphorus pentasulfide and Lawesson's
15 reagent (2,4-bis-(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane 2,4-disulfide).

 The amount of the sulfating agent to be used in the reaction is usually one or more moles based on 1 mole of Compound (1K).

20 The reaction temperature of the reaction is usually within a range of 0 to 200 °C. The reaction time of the reaction is usually within a range of 1 to 24 hours.


 After the completion of the reaction, Compound (12) can be isolated by pouring the reaction mixture into water,
25 extracting with an organic solvent, and concentrating the

organic layer; pouring the reaction mixture into water, and collecting the generated solids by filtration; or collecting the solids generated in the reaction mixture by filtration. The isolated Compound (12) can be further purified by recrystallization, chromatography, and the like.
 5 [0092]

The intermediates of the present invention can be produced, for example, by the following method.

(Production method of intermediate 1)

10 Compound (2) can be produced by the following method:



wherein, R⁶, R⁷, R⁸, A², and A³ are as defined above.

(Step 1)

Compound (M2) can be produced by reacting Compound (M1) in the presence of a nitrating agent.
 15

The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include halogenated hydrocarbons such as dichloromethane and chloroform, acetic acid, concentrated sulfuric acid, concentrated nitric acid, water and mixtures thereof.
 20

Examples of the nitrating agent to be used in the reaction include concentrated nitric acid.

The amount of the nitrating agent to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (M1).

5 The reaction temperature of the reaction is usually within a range of -10 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (M2) can be isolated by post-treatments, for example, pouring the reaction mixture into water and then extracting the mixture with an organic solvent, followed by drying and
10 concentrating the organic layer. The isolated Compound (M2) can be further purified by chromatography, recrystallization, and the like.

[0093]

15 (Step 2)

Compound (2) can be produced by reacting Compound (M2) with hydrogen in the presence of a hydrogenating catalyst.

The reaction is usually performed under 1 to 100 atms of hydrogen in the presence of a catalyst.

20 Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; esters such as ethyl acetate and butyl acetate; alcohols such as methanol and ethanol; water; and mixtures thereof.

25 Examples of the hydrogenating catalyst to be used in

the reaction include transition metal compounds such as palladium-carbon, palladium hydroxide, Raney[®] nickel, and platinum oxide.

5 The amount of hydrogen to be used in the reaction is usually 3 moles based on 1 mole of Compound (M1). The amount of the hydrogenating catalyst to be used in the reaction is usually 0.001 to 0.5 moles based on 1 mole of Compound (M1).

10 An acid (or base, and the like) can be added to the reaction as necessary.


The reaction temperature of the reaction is usually within a range of -20 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

15 After the completion of the reaction, Compound (2) can be isolated by post-treatments, for example, filtrating the reaction mixture and then extracting the mixture with an organic solvent as necessary, followed by drying and concentrating the organic layer. The isolated Compound (2) can be further purified by chromatography, 20 recrystallization, and the like.

[0094]

(Production method of intermediate 2) (Step (A-2))

25 Compound (6) can be produced by reacting Compound (2) with Compound (5) in the presence of a dehydration-condensing agent:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , A^2 , A^3 , and n are as defined above.

The reaction is usually performed in the presence of a
5 solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic
10 hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; nitrogen-containing aromatic compounds such as pyridine and
15 quinoline; and mixtures thereof.

Examples of the dehydration-condensing agent to be used in the reaction include carbodiimides such as WSC and 1,3-dicyclohexylcarbodiimide, and BOP reagent.

The amount of Compound (5) to be used in the reaction
20 is usually 1 to 3 moles based on 1 mole of Compound (2). The amount of the dehydration-condensing agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of

Compound (2).


The reaction temperature of the reaction is usually within a range of 0 to 140 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

5 After the completion of the reaction, Compound (6) can be isolated by post-treatments, for example, pouring water into the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (6)
10 can be further purified by chromatography, recrystallization, and the like.

[0095]

(Production method of intermediate 3) (Step (B-2))

Compound (6) can be produced by reacting Compound (2)
15 with Compound (M3) in the presence of a base:



wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, A², A³, and n are as defined above.

The reaction is usually performed in the presence of a
20 solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether,

tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; and mixtures thereof.

Examples of the base include alkali metal carbonates such as sodium carbonate and potassium carbonate; tertiary amines such as triethylamine and diisopropylethylamine; and nitrogen-containing aromatic compounds such as pyridine and 4-dimethylaminopyridine.

The amount of Compound (M3) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (2). The amount of the base to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (2).

The reaction temperature of the reaction is usually within a range of -20 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.


After the completion of the reaction, Compound (6) can be isolated by post-treatments, for example, pouring water into the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (6) can be further purified by chromatography,

recrystallization, and the like.

[0096]

(Production method of intermediate 4)

Compound (7) can be produced by the following method:



wherein, R^6 , R^7 , A^2 , and A^3 are as defined above.

(Step 1)

Compound (M5) can be produced by reacting Compound (M4) in the presence of a nitrating agent.

10 The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include halogenated hydrocarbons such as chloroform, acetic acid, concentrated sulfuric acid, concentrated nitric acid, 15 water, and mixtures thereof.

Examples of the nitrating agent to be used in the reaction include concentrated nitric acid.

The amount of the nitrating agent to be used in the reaction is usually 1 to 3 moles based on 1 mole of 20 Compound (M4).

The reaction temperature of the reaction is usually within a range of -10 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (M5) can be isolated by post-treatments, for example, pouring the reaction mixture into water and then extracting the mixture with an organic solvent, followed by drying and
5 concentrating the organic layer. The isolated Compound (M5) can be further purified by chromatography, recrystallization, and the like.

[0097]

(Step 2)

10 Compound (7) can be produced by reacting Compound (M5) with hydrogen in the presence of a hydrogenating catalyst.

The reaction is usually performed under 1 to 100 atms of hydrogen in the presence of a catalyst.

Examples of the solvent to be used in the reaction
15 include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; esters such as ethyl acetate and butyl acetate; alcohols such as methanol and ethanol; water; and mixtures thereof.

Examples of the hydrogenating catalyst to be used in
20 the reaction include transition metal compounds such as palladium-carbon, palladium hydroxide, Raney[®] nickel, and platinum oxide.

The amount of hydrogen to be used in the reaction is usually 3 moles based on 1 mole of Compound (M5). The
25 amount of the hydrogenating catalyst to be used in the

reaction is usually 0.001 to 0.5 moles based on 1 mole of Compound (M5).

An acid (or base, and the like) can be added to the reaction as necessary.


5 The reaction temperature of the reaction is usually within a range of -20 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (7) can be isolated by post-treatments, for example, filtrating the reaction mixture and then extracting the mixture with an organic solvent as necessary, followed by drying and concentrating the organic layer. The isolated Compound (7) can be further purified by chromatography, recrystallization, and the like.

15 [0098]

(Production method of intermediate 5) (Step (C-2))

Compound (9) can be produced by reacting Compound (7) with Compound (3):



20 wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, A², A³, and n are as defined above.

The reaction is usually performed in the presence of a

solvent.

Examples of the solvent to be used in the reaction include alcohols such as methanol and ethanol; ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl
5 ether, and 1,4-dioxane; aromatic hydrocarbons such as toluene and xylene; and mixtures thereof.

The amount of Compound (3) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (7).

10 An acid or base can be added to the reaction as necessary.


The reaction temperature of the reaction is usually within a range of 0 to 150 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

15 After the completion of the reaction, Compound (9) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (9) can be further purified by chromatography, recrystallization, and the like.

20 [0099]

(Production method of intermediate 6) (Step (A-2))

Compound (10) can be produced by reacting Compound (7) with Compound (5) in the presence of a dehydration-condensing agent:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^2 , A^3 , and n are as defined above.

The reaction is usually performed in the presence of a
5 solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; nitrogen-containing aromatic compounds such as pyridine and quinoline; and mixtures thereof.

Examples of the dehydration-condensing agent to be used in the reaction include carbodiimides such as WSC and 1,3-dicyclohexylcarbodiimide, and BOP reagent.

The amount of Compound (5) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (7). The amount of the dehydration-condensing agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of

Compound (7).


The reaction temperature of the reaction is usually within a range of 0 to 140 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

5 After the completion of the reaction, Compound (10) can be isolated by post-treatments, for example, pouring water into the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound
10 (10) can be further purified by chromatography, recrystallization, and the like.

[0100]

(Production method of intermediate 7) (Step (B-2))

Compound (10) can be produced by reacting Compound (7)
15 with Compound (M3) in the presence of a base:



wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, A², A³, and n are as defined above.

The reaction is usually performed in the presence of a
20 solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether,

tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; and mixtures thereof.

Examples of the base to be used in the reaction include alkali metal carbonates such as sodium carbonate and potassium carbonate; tertiary amines such as triethylamine and diisopropylethylamine; and nitrogen-containing aromatic compounds such as pyridine and 4-dimethylaminopyridine.

The amount of Compound (M3) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (7). The amount of the base to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (7).

The reaction temperature of the reaction is usually within a range of -20 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.


After the completion of the reaction, Compound (10) can be isolated by post-treatments, for example, pouring water into the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound

(10) can be further purified by chromatography, recrystallization, and the like.

[0101]

(Production method of intermediate 8)

5 Compound (11) can be produced by the following method:



wherein, R^6 , R^7 , A^2 , and A^3 are as defined above.

(Step 1)

Compound (M7) can be produced by reacting Compound
10 (M6) with thiourea in the presence of a base.

The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction
include alcohols such as methanol and ethanol; water; and
15 mixtures thereof.

Examples of the base to be used in the reaction
include alkali metal hydroxides such as sodium hydroxide
and potassium hydroxide.

The amount of thiourea to be used in the reaction is
20 usually 0.5 to 3 moles based on 1 mole of Compound (M6).
The amount of the base to be used in the reaction is
usually 1 to 10 moles based on 1 mole of Compound (M6).

The reaction temperature of the reaction is usually

within a range of 0 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (M7) can be isolated by post-treatments, for example, pouring an acid into the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (M7) can be further purified by chromatography, recrystallization, and the like.

10 [0102]

(Step 2)

Compound (11) can be produced by reducing Compound (M7).

The reduction reaction can be performed, for example, in the presence of a reducing agent such as iron powder and zinc powder; acids such as hydrochloric acid and acetic acid; and water.

The reaction can be usually performed in the presence of a solvent.

20 Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; esters such as ethyl acetate and butyl acetate; alcohols such as methanol and ethanol; acid amides such as DMF and NMP; and mixtures thereof.

25

The amount of the reducing agent to be used in the reaction is usually 3 to 10 moles based on 1 mole of Compound (M7).


The reaction temperature of the reaction is usually within a range of 0 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (11) can be isolated by post-treatments, for example, adding water to the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (11) can be further purified by chromatography, recrystallization, and the like.

[0103]

(Production method of intermediate 9)

Compound (28) of Compound (21) wherein A¹ is -NR⁸- can be produced by reacting Compound (2) with Compound (M8) in the presence of a base:



wherein, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, A², A³, and V² are as defined above.

The reaction is usually performed in the presence of a solvent.

Examples of the base to be used in the reaction include hydrogen carbonates such as sodium hydrogen carbonate and potassium hydrogen carbonate; carbonates such as sodium carbonate and potassium carbonate; sulfites such as sodium sulfite and potassium sulfite; and mixtures thereof.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; nitrogen-containing aromatic compounds such as pyridine and quinoline; and mixtures thereof.

The amount of Compound (M8) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (2). The amount of the base to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (2).

The reaction temperature of the reaction is usually within a range of 30 to 200 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.


After the completion of the reaction, Compound (28) can be isolated by post-treatments, for example, pouring

water into the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (28) can be further purified by chromatography, recrystallization, and the like.

[0104]

(Production method of intermediate 10) (Step (E-1))

Compound (28) of Compound (21) wherein A^1 is $-NR^8-$ can be produced by reacting Compound (2) with Compound (M9) in the presence of a dehydration-condensing agent:



wherein, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , A^2 , A^3 , and V^2 are as defined above.

The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, tert-butyl methyl ether, ethylene glycol dimethyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO;

nitrogen-containing aromatic compounds such as pyridine and quinoline; and mixtures thereof.

Examples of the dehydration-condensing agent to be used in the reaction include carbodiimides such as WSC and
5 1,3-dicyclohexylcarbodiimide, and BOP reagent.

A catalyst can be added to the reaction as necessary.

Examples of the catalyst to be used in the reaction include 1-hydroxybenzotriazole (hereinafter referred to as HOBt).

10 The amount of Compound (M9) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (2). The amount of the dehydration-condensing agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (2). The amount of the catalyst to be used in the
15 reaction is usually 0.01 to 0.1 moles based on 1 mole of Compound (2).

The reaction temperature of the reaction is usually within a range of 30 to 200 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.


20 After the completion of the reaction, Compound (28) can be isolated by post-treatments, for example, pouring water into the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound
25 (28) can be further purified by chromatography,

recrystallization, and the like.

[0105]

(Production method of intermediate 11) (Step (E-3))

Compound (28) of Compound (21) wherein A^1 is $-NR^8-$ can
5 be produced by dehydration-condensing Compound (M10):



wherein, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , A^2 , A^3 , and V^2 are as defined above.

The reaction is usually performed in the presence of a
10 solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic
15 hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; alcohols such as methanol, ethanol, propanol, butanol, and
20 pentanol; and mixtures thereof.

In the reaction, an acid or a dehydrating agent can be used as necessary. Examples of the acid to be used in the

reaction include sulfonic acids such as p-toluenesulfonic acid, and carboxylic acids such as acetic acid. Examples of the dehydrating agent to be used in the reaction include phosphorous oxychloride, acetic anhydride, and
5 trifluoroacetic anhydride.

The amount of the acid or dehydrating agent to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (M10).


The reaction temperature of the reaction is usually
10 within a range of 30 to 200 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (28) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by
15 drying and concentrating the organic layer. The isolated Compound (28) can be further purified by chromatography, recrystallization, and the like.

[0106]

(Production method of intermediate 12) (Step (E-3))

20 Compound (28) of Compound (21) wherein A¹ is -NR⁸- can be produced by reacting Compound (M10) in the presence of a base:



wherein, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , A^2 , A^3 , and V^2 are as defined above.

The reaction is usually performed in the presence of a
5 solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic
10 hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; alcohols such as methanol, ethanol, propanol, butanol,
15 tert-butanol, and pentanol; and mixtures thereof.

Examples of the base to be used in the reaction include tripotassium phosphate.

The amount of the base to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (M10).

20 The reaction temperature of the reaction is usually within a range of 30 to 200 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (28) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated


5 Compound (28) can be further purified by chromatography, recrystallization, and the like.

[0107]

(Production method of intermediate 13) (Step (E-2))

Compound (M10) can be produced by reacting Compound

10 (2) with Compound (M9) in the presence of a dehydration-condensing agent:



wherein, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , A^2 , A^3 , and V^2 are as defined above.

15 The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic

20 hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl

acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; nitrogen-containing aromatic compounds such as pyridine and quinoline; and mixtures thereof.

5 Examples of the dehydration-condensing agent to be used in the reaction include carbodiimides such as WSC and 1,3-dicyclohexylcarbodiimide, and BOP reagent.

 The amount of Compound (M9) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (2).
10 The amount of the dehydration-condensing agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (2).

 The reaction temperature of the reaction is usually within a range of 0 to 140 °C. The reaction time of the
15 reaction is usually within a range of 0.1 to 24 hours.


 After the completion of the reaction, Compound (M10) can be isolated by post-treatments, for example, pouring water into the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and
20 concentrating the organic layer. The isolated Compound (M10) can be further purified by chromatography, recrystallization, and the like.

[0108]

(Production method of intermediate 14)

25 Compound (M10) can be produced by reacting Compound

(2) with Compound (M12) in the presence of a base:



wherein, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , A^2 , A^3 , and V^2 are as defined above.

5 The reaction is usually performed in the presence of a
solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; and mixtures thereof.

Examples of the base include alkali metal carbonates such as sodium carbonate and potassium carbonate; tertiary amines such as triethylamine and diisopropylethylamine; and nitrogen-containing aromatic compounds such as pyridine and 4-dimethylaminopyridine.

The amount of Compound (M12) to be used in the reaction is usually 1 to 3 moles based on 1 mole of

Compound (2). The amount of the base to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (2).


The reaction temperature of the reaction is usually within a range of -20 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (M10) can be isolated by post-treatments, for example, pouring water into the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (M10) can be further purified by chromatography, recrystallization, and the like.

[0109]

(Production method of intermediate 15)

Compound (29) of Compound (3) wherein n is 0 can be produced by the following method:



wherein, R¹, R², R³, R⁴, R⁵, and V² are as defined above.

(Step I15-1)

Compound (M14) can be produced by reacting Compound

(M13) with Compound (22) in the presence of a base.

The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aromatic hydrocarbons such as toluene and xylene; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; and mixtures thereof.

Examples of the base to be used in the reaction include alkali metal hydrides such as sodium hydride.

The amount of Compound (22) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (M13). The amount of the base to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (M13).

The reaction temperature of the reaction is usually within a range of 0 to 150 °C. The reaction time of the reaction is usually within a range of 0.5 to 24 hours.

After the completion of the reaction, Compound (M14) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (M14) can be further purified by chromatography, recrystallization, and the like.

[0110]

(Step I15-2)

Compound (29) can be produced by reducing Compound (M14).

5 The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aromatic hydrocarbons such as toluene and xylene; halogenated
10 hydrocarbons such as dichloromethane and chloroform; and mixtures thereof.

Examples of the reducing agent to be used in the reaction include diisobutylaluminium hydrate.

The amount of the reducing agent to be used in the
15 reaction is usually 1 to 2 moles based on 1 mole of Compound (M14).


The reaction temperature of the reaction is usually within a range of -20 to 100 °C. The reaction time of the reaction is usually within a range of 0.5 to 24 hours.

20 After the completion of the reaction, Compound (29) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (29) can be further purified by chromatography,
25 recrystallization, and the like.

[0111]

(Production method of intermediate 16)

Compound (29) of Compound (3) wherein n is 0 can be produced by reacting Compound (M8) with Compound (22) in the presence of a base:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , and V^2 are as defined above.

The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aromatic hydrocarbons such as toluene and xylene; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; and mixtures thereof.

Examples of the base to be used in the reaction include alkali metal hydrides such as sodium hydride.

The amount of Compound (22) to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (M15).

The amount of the base to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (M8).

The reaction temperature of the reaction is usually


within a range of 0 to 150 °C. The reaction time of the reaction is usually within a range of 0.5 to 24 hours.

After the completion of the reaction, Compound (29) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (29) can be further purified by chromatography, recrystallization, and the like.

[0112]

(Production method of intermediate 17)

Compound (30) of Compound (5) wherein n is 0 can be produced by hydrolyzing Compound (M14) in the presence of a base, and Compound (M23) of Compound (5) wherein n is 1 or 2 can be produced by the following method:



wherein, R^1 , R^2 , R^3 , R^4 , and R^5 are as defined above, and r

represents 1 or 2.

[0113]

(Step I17-1)

5 The reaction is usually performed in the presence of a solvent.

 Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; alcohols such as methanol and ethanol; water; and mixtures thereof.

10 Examples of the base to be used in the reaction include alkali metal hydroxides such as sodium hydroxide and potassium hydroxide.

 The amount of the base to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (M14).

15 The reaction temperature of the reaction is usually within a range of 0 to 120 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

 After the completion of the reaction, Compound (30) can be isolated by post-treatments, for example, acidifying
20 the reaction mixture and then extracting with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (30) can be further purified by chromatography, recrystallization, and the like.

[0114]

25 (Step I17-2)

The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include halogenated hydrocarbons such as dichloromethane and chloroform; alcohols such as methanol and ethanol; 5 acetic acid; water; and mixtures thereof.

Examples of the oxidizing agent to be used in the reaction include m-chloroperbenzoic acid and an aqueous solution of hydrogen peroxide.

10 The reaction can be performed in the presence of a catalyst as necessary.

Examples of the catalyst to be used in the reaction include sodium tungstate.

The amount of the oxidizing agent to be used in the 15 reaction is usually 1 to 5 moles based on 1 mole of Compound (M14).

The reaction temperature of the reaction is usually within a range of -20 to 120 °C. The reaction time of the reaction is usually within a range of 0.1 to 12 hours.

20 After the completion of the reaction, Compound (M22) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by washing the organic layer with an aqueous solution of a reducing agent (e.g. sodium sulfite, thiosodium sulfate) 25 and an aqueous solution of a base (e.g. sodium hydrogen

carbonate) as necessary, followed by drying and concentrating the organic layer. The isolated Compound (M22) can be further purified by chromatography, recrystallization, and the like.

5 [0115]

(Step I17-3)

The reaction is usually performed in the presence of a solvent.

10 Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; alcohols such as methanol and ethanol; water; and mixtures thereof.

15 Examples of the base to be used in the reaction include alkali metal hydroxides such as sodium hydroxide and potassium hydroxide.

The amount of the base to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (M22).

20 The reaction temperature of the reaction is usually within a range of 0 to 120 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

25 After the completion of the reaction, Compound (M23) can be isolated by post-treatments, for example, acidifying the reaction mixture and then extracting with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (M23) can be further purified

by chromatography, recrystallization, and the like.

[0116]

(Step I17-4)

5 The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include halogenated hydrocarbons such as dichloromethane and chloroform; alcohols such as methanol and ethanol; acetic acid; water; and mixtures thereof.

10 Examples of the oxidizing agent to be used in the reaction include m-chloroperbenzoic acid and an aqueous solution of hydrogen peroxide.

The reaction can be performed in the presence of a catalyst as necessary.

15 Examples of the catalyst to be used in the reaction include sodium tungstate.

The amount of the oxidizing agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (30).

20 The reaction temperature of the reaction is usually within a range of -20 to 120 °C. The reaction time of the reaction is usually within a range of 0.1 to 12 hours.


25 After the completion of the reaction, Compound (M23) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by

washing the organic layer with an aqueous solution of a reducing agent (e.g. sodium sulfite, thiosodium sulfate) and an aqueous solution of a base (e.g. sodium hydrogen carbonate) as necessary, followed by drying and concentrating the organic layer. The isolated Compound (M23) can be further purified by chromatography, recrystallization, and the like.

[0117]

(Production method of intermediate 18)

Compound (30) of Compound (5) wherein n is 0 can be produced by hydrolyzing Compound (M14) in the presence of an acid:



wherein, R^1 , R^2 , R^3 , R^4 , and R^5 are as defined above.

In the reaction, usually, an aqueous solution of the acid is used as a solvent.

Examples of the acid to be used in the reaction include mineral acids such as hydrochloric acid, nitric acid, phosphoric acid, and sulfuric acid, and carboxylic acids such as acetic acid and trifluoroacetic acid.

The reaction temperature of the reaction is usually within a range of 0 to 100 °C. The reaction time of the


reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (30) can be isolated by post-treatments, for example, extracting the reaction mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated
 5 Compound (30) can be further purified by chromatography, recrystallization, and the like.

[0118]

(Production method of intermediate 19)

10 Compound (M3) can be produced by chlorinating Compound (5) in the presence of a chlorinating agent:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , and n are as defined above.

The reaction is usually performed in the presence of a
 15 solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aromatic hydrocarbons such as toluene and xylene; halogenated
 20 hydrocarbons such as dichloromethane and chloroform; and mixtures thereof.

Examples of the chlorinating agent to be used in the

reaction include thionyl chloride and oxalyl dichloride.

The amount of the chlorinating agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (5).


5 The reaction temperature of the reaction is usually within a range of 0 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (M3) can be isolated by evaporating the solvent.

10 [0119]

(Production method of intermediate 20)

Compound (M21) of Compound (2) wherein A³ is a nitrogen atom can be produced, for example, by the following method:



wherein, R⁶, R⁷, R⁸, and A² are as defined above, and X⁹ represents a chlorine atom, a bromine atom, or an iodine atom.

(Step I20-1)

20 Compound (M19) can be produced by reacting Compound (M17) with Compound (M18).

The reaction is performed in the presence or absence of a solvent.

Examples of the solvent to be used in the reaction include water, alcohols such as methanol and ethanol; ethers such as 1,4-dioxane, diethyl ether, and THF; esters such as ethyl acetate and butyl acetate; halogenated hydrocarbons such as dichloromethane, chloroform, carbon tetrachloride, and 1,2-dichloroethane; nitriles such as acetonitrile; aprotic polar solvents such as DMF, NMP, and dimethylsulfoxide; nitrogen-containing aromatic compounds such as pyridine and quinoline; and mixtures thereof.

10 A base can be added to the reaction as necessary.

Examples of the base to be used in the reaction include nitrogen-containing heterocyclic compounds such as pyridine, picoline, 2,6-lutidine, diazabicycloundecene (hereinafter referred to as DBU), and 1,5-diazabicyclo[4.3.0]-5-nonen; tertiary amines such as triethylamine and N-ethyldiisopropylamine; and inorganic bases such as potassium carbonate, cesium carbonate, and sodium hydroxide.

20 The amount of Compound (M18) to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (M17).

The reaction temperature of the reaction is usually within a range of 0 to 200 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

25 After the completion of the reaction, Compound (M19)

can be isolated by pouring the reaction mixture into water, extracting with an organic solvent, and concentrating the organic layer; pouring the reaction mixture into water, and collecting the generated solids by filtration; or
5 collecting the solids generated in the reaction mixture by filtration. The isolated Compound (M19) can be further purified by recrystallization, chromatography, and the like.
[0120]

(Step I20-2)

10 Compound (M20) can be produced by reacting Compound (M19) with a halogenating agent.

The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction
15 include water; acetic acid; ethers such as 1,4-dioxane, diethyl ether, and THF; esters such as ethyl acetate and butyl acetate; halogenated hydrocarbons such as dichloromethane, chloroform, carbon tetrachloride, and 1,2-dichloroethane; nitriles such as acetonitrile; aprotic
20 polar solvents such as DMF and NMP; and mixtures thereof.

Examples of the halogenating agent to be used in the reaction include a chlorinating agent such as N-chlorosuccinimide and chlorine; a brominating agent such as N-bromosuccinimide and bromine; and an iodinating agent
25 such as N-iodosuccinimide and iodine.

The amount of the halogenating agent to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (M19).

5 The reaction temperature of the reaction is usually within a range of -10 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (M20) can be isolated by pouring the reaction mixture into water, extracting with an organic solvent, and concentrating the organic layer; pouring the reaction mixture into water, and
10 collecting the generated solids by filtration; or collecting the solids generated in the reaction mixture by filtration. The isolated Compound (M20) can be further purified by recrystallization, chromatography, and the like.

15 [0121]

(Step I20-3)

Compound (M21) can be produced by reacting Compound (M20) with an aminating agent in the presence of a copper compound.

20 The reaction is usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include water; alcohols such as methanol and ethanol; ethers such as 1,4-dioxane, diethyl ether, and THF; esters
25 such as ethyl acetate and butyl acetate; halogenated

hydrocarbons such as dichloromethane, chloroform, carbon tetrachloride, and 1,2-dichloroethane; nitriles such as acetonitrile; aprotic polar solvents such as DMF, NMP, and dimethylsulfoxide; nitrogen-containing aromatic compounds
5 such as pyridine and quinoline; and mixtures thereof.

Examples of the aminating agent to be used in the reaction include ammonia, ammonia water, and lithium amide.

Examples of the copper compound to be used in the reaction include copper, copper (I) iodide, copper (I)
10 oxide, copper (II) oxide, copper (II) acetylacetonate, copper (II) acetate, and copper (II) sulfate.

A ligand can be added to the reaction as necessary.

Examples of the ligand to be used in the reaction include acetylacetonate, salen, and phenanthroline.

15 A base can be added to the reaction as necessary.

Examples of the base to be used in the reaction include nitrogen-containing heterocyclic compounds such as pyridine, picoline, 2,6-lutidine, DBU, and 1,5-diazabicyclo[4.3.0]-5-nonen; tertiary amines such as
20 triethylamine and N-ethyldiisopropylamine; and inorganic bases such as tripotassium phosphate, potassium carbonate, cesium carbonate, and sodium hydroxide.

The amount of the aminating agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of
25 Compound (M20). The amount of the copper compound to be


used in the reaction is usually 0.02 to 0.5 moles based on 1 mole of Compound (M20). The amount of the ligand to be used in the reaction is 0.02 to 2 moles based on 1 mole of Compound (M20) as necessary. The amount of the base to be
5 used in the reaction is 1 to 5 moles based on 1 mole of Compound (M20) as necessary.

The reaction temperature of the reaction is usually within a range of 30 to 200 °C. The reaction time of the reaction is usually within a range of 0.1 to 48 hours.

10 After the completion of the reaction, Compound (M21) can be isolated by pouring the reaction mixture into water, extracting with an organic solvent, and concentrating the organic layer; pouring the reaction mixture into water, and collecting the generated solids by filtration; or
15 collecting the solids generated in the reaction mixture by filtration. The isolated Compound (M21) can be further purified by recrystallization, chromatography, and the like.
[0122]

(Production method of intermediate 21) (Step (A-2))

20 Compound (35) can be produced by reacting Compound (11) with Compound (5) in the presence of a dehydration-condensing agent:



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^2 , A^3 , and n are as defined above.

The reaction is usually performed in the presence of a
5 solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic
10 hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; nitrogen-containing aromatic compounds such as pyridine and
15 quinoline; and mixtures thereof.

Examples of the dehydration-condensing agent to be used in the reaction include carbodiimides such as WSC and 1,3-dicyclohexylcarbodiimide, and BOP reagent.

The amount of Compound (5) to be used in the reaction
20 is usually 1 to 3 moles based on 1 mole of Compound (11). The amount of the dehydration-condensing agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of

Compound (11).


The reaction temperature of the reaction is usually within a range of 0 to 140 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

5 After the completion of the reaction, Compound (35) can be isolated by post-treatments, for example, pouring water into the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound
10 (35) can be further purified by chromatography, recrystallization, and the like.

[0123]

(Production method of intermediate 22) (Step (B-2))

Compound (35) can be produced by reacting Compound
15 (11) with Compound (M3) in the presence of a base:



wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, A², A³, and n are as defined above.

The reaction is usually performed in the presence of a
20 solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether,

tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; halogenated hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; and mixtures thereof.

Examples of the base include alkali metal carbonates such as sodium carbonate and potassium carbonate; tertiary amines such as triethylamine and diisopropylethylamine; and nitrogen-containing aromatic compounds such as pyridine and 4-dimethylaminopyridine.

The amount of Compound (M3) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (11). The amount of the base to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (11).

The reaction temperature of the reaction is usually within a range of -20 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.


After the completion of the reaction, Compound (35) can be isolated by post-treatments, for example, pouring water into the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (35) can be further purified by chromatography,

recrystallization, and the like.

[0124]

(Production method of intermediate 23) (Step (E-1))

Compound (M24) of Compound (21) wherein A¹ is an
 5 oxygen atom can be produced by reacting Compound (7) with
 Compound (M9):




wherein, R², R³, R⁴, R⁵, R⁶, R⁷, A², A³, n, and V² are as
 defined above.

10 Compound (M24) can be produced according to the method
 described in (Production method 6) using Compound (M9)
 instead of Compound (5).

[0125]

(Production method of intermediate 24) (Step (E-2))

15 Compound (M25) of Compound (1E) wherein A¹ is an
 oxygen atom can be produced by reacting Compound (7) with
 Compound (M9). Additionally, Compound (M26) of Compound
 (1E) wherein A¹ is a sulfur atom can be produced by
 reacting Compound (11) with Compound (M9):



wherein, R², R³, R⁴, R⁵, R⁶, R⁷, A², A³, n, and V² are as defined above.


Compound (M25) can be produced according to the method described in Production method of intermediate 6 using Compound (M9) instead of Compound (5).

Compound (M26) can be produced according to the method described in Production method of intermediate 21 using Compound (11) instead of Compound (5).

[0126]

(Production method of intermediate 25) (Step (E-3))

Compound (M27) of Compound (21) wherein A¹ is an oxygen atom can be produced by cyclizing Compound (M25). Additionally, Compound (M28) of Compound (21) wherein A¹ is a sulfur atom can be produced by cyclizing Compound (M26):



wherein, $R^2, R^3, R^4, R^5, R^6, R^7, A^2, A^3, n$, and V^2 are as defined above.


Compound (M27) can be produced according to the method described in Production method 8 or Production method 9 using Compound (M25) instead of Compound (10).

Compound (M28) can be produced according to the method described in Production method 21 using Compound (M26) instead of Compound (35).

[0127]

(Production method of intermediate 26) (Step (F-1))

Compound (24) can be produced by reacting Compound (21) with sodium sulfide, sodium hydrogen sulfide, or hydrogen sulfide:



wherein, $R^2, R^3, R^4, R^5, R^6, R^7, A^1, A^2, A^3, n$, and V^2 are as

defined above.


Compound (24) can be produced according to the method described in Production method 16 using sodium sulfide, sodium hydrogen sulfide, or hydrogen sulfide instead of
5 Compound (22).

When sodium sulfide or sodium hydrogen sulfide is used, a reaction is usually performed without a base.

[0128]

(Production method of intermediate 27) (Production method
10 D)

Compound (1K) can be produced by reacting Compound (M30) with Compound (5):



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^2 , A^3 , and n are as
15 defined above.

[0129]

(Step I27-1)

Compound (M30) can be produced by reducing Compound (M29).

The reduction reaction can be performed in the presence of a reducing agent such as iron powder and zinc powder; acids such as hydrochloric acid and acetic acid; and water.

5 The reaction can be usually performed in the presence of a solvent.

Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; esters such as
10 ethyl acetate and butyl acetate; alcohols such as methanol and ethanol; acid amides such as DMF and NMP; and mixtures thereof.

The amount of the reducing agent to be used in the reaction is usually 3 to 10 moles based on 1 mole of
15 Compound (M29).

The reaction temperature of the reaction is usually within a range of 0 to 100 °C. The reaction time of the reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (M30)
20 can be isolated by post-treatments, for example, adding water to the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (M30) can be further purified by chromatography,
25 recrystallization, and the like.

[0130]

(Step I27-2)

The reaction is usually performed in the presence or absence of a solvent.

5 Examples of the solvent to be used in the reaction include ethers such as THF, ethylene glycol dimethyl ether, tert-butyl methyl ether, and 1,4-dioxane; aliphatic hydrocarbons such as hexane, heptane, and octane; aromatic hydrocarbons such as toluene and xylene; halogenated
10 hydrocarbons such as chlorobenzene; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile; acid amides such as DMF and NMP; sulfoxides such as DMSO; nitrogen-containing aromatic compounds such as pyridine and quinoline; and mixtures thereof.

15 Examples of the dehydration-condensing agent to be used in the reaction include carbodiimides such as WSC and 1,3-dicyclohexylcarbodiimide, and BOP reagent.

 The amount of Compound (5) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (M30).
20 The amount of the dehydration-condensing agent to be used in the reaction is usually 1 to 5 moles based on 1 mole of Compound (M30).

 The reaction temperature of the reaction is usually within a range of 0 to 140 °C. The reaction time of the
25 reaction is usually within a range of 0.1 to 24 hours.

After the completion of the reaction, Compound (1K) can be isolated by post-treatments, for example, pouring water into the reaction mixture and then extracting the mixture with an organic solvent, followed by drying and concentrating the organic layer. The isolated Compound (1K) can be further purified by chromatography, recrystallization, and the like.

Alternatively, Compound (1K) can be produced according to the above method using Compound (M3) instead of Compound (5).

When Compound (M3) is used, the reaction is usually performed without a dehydration-condensing agent

A base can be added to the reaction as necessary.

Examples of the base include carbonates of alkali metal such as sodium carbonate and potassium carbonate; tertiary amines such as triethylamine and diisopropylethylamine; and nitrogen-containing aromatic compounds such as pyridine and 4-dimethylaminopyridine.


The amount of Compound (M3) to be used in the reaction is usually 1 to 3 moles based on 1 mole of Compound (M30). The amount of the base to be used in the reaction is usually 1 to 10 moles based on 1 mole of Compound (M30).

[0131]

(Production method of intermediate 28)

Compound (F) can be produced by reacting Compound (G)

with an aminating agent:



wherein, X^9 , R^6 , R^7 , A^1 , A^2 , and A^3 are as defined the above.


Compound (F) can be produced according to the method described in Step (I20-3) of Production method of intermediate 20 using Compound (G) instead of Compound (M20).

[0132]

Examples of Compound (B) include the following compounds:

[0133]


An amide compound of the formula (6):



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , A^2 , A^3 , and n are as defined above (referred to as Compound (6));

[0134]


An amide compound of the formula (10):



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^2 , A^3 , and n are as defined above (referred to as Compound (10));

[0135]


5 An amide compound of the formula (35):



wherein, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , A^2 , A^3 , and n are as defined above (referred to as Compound (35));

[0136]

10 An amide compound of the formula (1B):



wherein:

A^{1b} represents $-NR^{8b}-$, an oxygen atom or a sulfur atom;

A^{2b} represents a nitrogen atom or $=CR^{9b}-$;

15 A^{3b} represents a nitrogen atom or $=CR^{10b}-$;

R^{1b} represents a C1-C6 chain hydrocarbon group

optionally having one or more halogen atoms or a C3-C6 alicyclic hydrocarbon group optionally having one or more halogen atoms;

R^{2b} , R^{3b} , R^{4b} , and R^{5b} are same or different and
5 represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more
10 halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, $-OR^{11b}$, $-S(O)_mR^{11b}$, $-SF_5$, a cyano group, a nitro group, a halogen
15 atom, or a hydrogen atom: provided that at least two of R^{2b} , R^{3b} , R^{4b} , and R^{5b} represent a hydrogen atom;

R^{6b} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl
20 groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl
25 groups optionally having one or more halogen atoms, $-OR^{11b}$,

-S(O)_mR^{11b}, -SF₅, a cyano group, a nitro group, or a halogen atom;

R^{7b} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, -OR^{11b}, -S(O)_mR^{11b}, -SF₅, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

R^{8b} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group, a C3-C6 alicyclic hydrocarbon group optionally having one or more halogen atoms, or a hydrogen atom;

R^{9b} and R^{10b} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, -OR^{11b}, -S(O)_mR^{11b}, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

R^{11b} and R^{12b} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a hydrogen atom; and

m represents 0, 1, or 2, and n represents 0, 1, or 2;

provided that: in $-S(O)_mR^{11b}$, when m is 1 or 2, R^{11b} is not a hydrogen atom; and when A^{1b} is an oxygen atom, and A^{2b} is a methine group, any one of R^{2b} , R^{3b} , R^{4b} , and R^{5b} are a
5 C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl
10 groups optionally having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, $-OR^{11b}$, $-S(O)_mR^{11b}$, a cyano group, a nitro group, or a halogen atom (hereinafter, referred to as
15 Compound (1B));

[0137]

And examples of Compound (1B) include the following compounds:

20 An amide compound of the formula (1B) wherein A^{1b} is -
 $NR^{8b}-$;

An amide compound of the formula (1B) wherein A^{1b} is -
 $NR^{8b}-$, and R^{8b} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

25 An amide compound of the formula (1B) wherein A^{1b} is -
 $NR^{8b}-$, and R^{8b} is a C1-C6 chain hydrocarbon group;

An amide compound of the formula (1B) wherein A^{1b} is -
 NR^{8b} -, and R^{8b} is a C1-C6 alkyl group;

An amide compound of the formula (1B) wherein A^{1b} is -
 NR^{8b} -, and R^{8b} is a C1-C3 alkyl group;

5 An amide compound of the formula (1B) wherein A^{1b} is -
 NR^{8b} -, and R^{8b} is a methyl group, an ethyl group, a propyl
group, or an isopropyl group;

An amide compound of the formula (1B) wherein A^{1b} is -
 NR^{8b} -, and R^{8b} is a methyl group;

10 An amide compound of the formula (1B) wherein A^{1b} is -
 NR^{8b} -, and R^{8b} is an ethyl group;

An amide compound of the formula (1B) wherein A^{1b} is -
 NR^{8b} -, and R^{8b} is a propyl group;

15 An amide compound of the formula (1B) wherein A^{1b} is -
 NR^{8b} -, and R^{8b} is an isopropyl group;

An amide compound of the formula (1B) wherein A^{1b} is -
 NR^{8b} -, and R^{8b} is a C1-C6 chain hydrocarbon group having one
C1-C6 alkoxy group;

20 An amide compound of the formula (1B) wherein A^{1b} is -
 NR^{8b} -, and R^{8b} is a methoxymethyl group or an ethoxymethyl
group;

An amide compound of the formula (1B) wherein A^{1b} is -
 NR^{8b} -, and R^{8b} is a C3-C6 alicyclic hydrocarbon group
optionally having one or more halogen atoms;

25 An amide compound of the formula (1B) wherein A^{1b} is -

NR^{8b}-, and R^{8b} is a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, or a cyclohexyl group;

An amide compound of the formula (1B) wherein A^{1b} is - NR^{8b}-, and R^{8b} is a hydrogen atom;

5 An amide compound of the formula (1B) wherein A^{1b} is an oxygen atom;

An amide compound of the formula (1B) wherein A^{1b} is a sulfur atom;

10 An amide compound of the formula (1B) wherein A^{1b} is - NR^{8b}- or a sulfur atom;

An amide compound of the formula (1B) wherein A^{1b} is - NR^{8b}- or a sulfur atom, and R^{8b} is a methyl group, an ethyl group, a propyl group, or an isopropyl group;

15 An amide compound of the formula (1B) wherein A^{1b} is - NR^{8b}- or a sulfur atom, and R^{8b} is a methoxymethyl group, an ethoxymethyl group, or a hydrogen atom;

An amide compound of the formula (1B) wherein A^{2b} is =CR^{9b}-, and A^{3b} is =CR^{10b}-;

20 An amide compound of the formula (1B) wherein A^{2b} is =CR^{9b}-, A^{3b} is =CR^{10b}-, and R^{9b} and R^{10b} are same or different and are independently a halogen atom or a hydrogen atom;

An amide compound of the formula (1B) wherein A^{2b} is =CR^{9b}-, A^{3b} is =CR^{10b}-, and R^{9b} and R^{10b} are a hydrogen atom;

25 An amide compound of the formula (1B) wherein A^{2b} is a nitrogen atom, and A^{3b} is =CR^{10b}-;

An amide compound of the formula (1B) wherein A^{2b} is $=CR^{9b}-$, and A^{3b} is a nitrogen atom;

An amide compound of the formula (1B) wherein A^{2b} is $=CR^{9b}-$, R^{9b} is a halogen atom or a hydrogen atom, and A^{3b} is
5 a nitrogen atom;

An amide compound of the formula (1B) wherein A^{2b} is $=CR^{9b}-$, R^{9b} is a hydrogen atom, and A^{3b} is a nitrogen atom;

An amide compound of the formula (1B) wherein A^{2b} is a nitrogen atom, and A^{3b} is a nitrogen atom;

10 An amide compound of the formula (1B) wherein A^{1b} is -
 $NR^{8b}-$, A^{2b} is $=CR^{9b}-$, and A^{3b} is $=CR^{10b}-$;

An amide compound of the formula (1B) wherein A^{1b} is -
 $NR^{8b}-$, A^{2b} is $=CR^{9b}-$, A^{3b} is $=CR^{10b}-$, and R^{9b} and R^{10b} are a
hydrogen atom;

15 An amide compound of the formula (1B) wherein A^{1b} is
an oxygen atom, A^{2b} is $=CR^{9b}-$, and A^{3b} is $=CR^{10b}-$;

An amide compound of the formula (1B) wherein A^{1b} is a
sulfur atom, A^{2b} is $=CR^{9b}-$, and A^{3b} is $=CR^{10b}-$;

20 An amide compound of the formula (1B) wherein A^{1b} is a
sulfur atom, A^{2b} is $=CR^{9b}-$, A^{3b} is $=CR^{10b}-$, and R^{9b} and R^{10b}
are a hydrogen atom;

An amide compound of the formula (1B) wherein A^{1b} is -
 $NR^{8b}-$, A^{2b} is $=CR^{9b}-$, and A^{3b} is a nitrogen atom;

25 An amide compound of the formula (1B) wherein A^{1b} is -
 $NR^{8b}-$, A^{2b} is $=CR^{9b}-$, R^{9b} is a hydrogen atom, and A^{3b} is a

nitrogen atom;

An amide compound of the formula (1B) wherein A^{1b} is an oxygen atom, A^{2b} is $=CR^{9b}-$, and A^{3b} is a nitrogen atom;

5 An amide compound of the formula (1B) wherein A^{1b} is a sulfur atom, A^{2b} is $=CR^{9b}-$, and A^{3b} is a nitrogen atom;

An amide compound of the formula (1B) wherein A^{1b} is a sulfur atom, A^{2b} is $=CR^{9b}-$, R^{9b} is a hydrogen atom, and A^{3b} is a nitrogen atom;

10 An amide compound of the formula (1B) wherein A^{1b} is -
 $NR^{8b}-$ or a sulfur atom, R^{8b} is a methyl group, an ethyl group, a propyl group, or an isopropyl group, A^{2b} is $=CR^{9b}-$, A^{3b} is $=CR^{10b}-$, and R^{9b} and R^{10b} are a hydrogen atom;

15 An amide compound of the formula (1B) wherein A^{1b} is -
 $NR^{8b}-$ or a sulfur atom, R^{8b} is a methyl group, A^{2b} is $=CR^{9b}-$, A^{3b} is $=CR^{10b}-$, and R^{9b} and R^{10b} are a hydrogen atom;

An amide compound of the formula (1B) wherein A^{1b} is -
 $NR^{8b}-$ or a sulfur atom, R^{8b} is a methyl group, an ethyl group, a propyl group or an isopropyl group, A^{2b} is $=CR^{9b}-$, R^{9b} is a hydrogen atom, and A^{3b} is a nitrogen atom;

20 An amide compound of the formula (1B) wherein A^{1b} is -
 $NR^{8b}-$ or a sulfur atom, R^{8b} is a methyl group, A^{2b} is $=CR^{9b}-$, R^{9b} is a hydrogen atom, and A^{3b} is a nitrogen atom;

25 An amide compound of the formula (1B) wherein R^{1b} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

An amide compound of the formula (1B) wherein R^{1b} is a C1-C6 chain hydrocarbon group optionally having one or more fluorine atoms;

5 An amide compound of the formula (1B) wherein R^{1b} is a C1-C6 chain hydrocarbon group;

An amide compound of the formula (1B) wherein R^{1b} is a C1-C6 alkyl group;

10 An amide compound of the formula (1B) wherein R^{1b} is a C1-C6 alkyl group optionally having one or more fluorine atoms;

An amide compound of the formula (1B) wherein R^{1b} is a C1-C3 alkyl group;

15 An amide compound of the formula (1B) wherein R^{1b} is a C1-C3 alkyl group optionally having one or more fluorine atoms;

An amide compound of the formula (1B) wherein R^{1b} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group;

20 An amide compound of the formula (1B) wherein R^{1b} is a methyl group;

An amide compound of the formula (1B) wherein R^{1b} is an ethyl group;

25 An amide compound of the formula (1B) wherein R^{1b} is a propyl group;

An amide compound of the formula (1B) wherein R^{1b} is an isopropyl group;

An amide compound of the formula (1B) wherein R^{1b} is a trifluoromethyl group;

5 An amide compound of the formula (1B) wherein R^{1b} is a 2,2,2-trifluoroethyl group;

An amide compound of the formula (1B) wherein R^{1b} is a C3-C6 alicyclic hydrocarbon group optionally having one or more halogen atoms;

10 An amide compound of the formula (1B) wherein R^{1b} is a C3-C6 alicyclic hydrocarbon group;

An amide compound of the formula (1B) wherein R^{1b} is a cyclopropyl group, a cyclobutyl group, or a cyclopentyl group;

15 An amide compound of the formula (1B) wherein R^{2b}, R^{3b}, R^{4b}, and R^{5b} are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a hydrogen atom;

20 An amide compound of the formula (1B) wherein R^{2b}, R^{3b}, R^{4b}, and R^{5b} are same or different and are independently a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally
25 having one or more halogen atoms, a 6-membered heterocyclic

group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, or a hydrogen atom;

5 An amide compound of the formula (1B) wherein R^{2b} , R^{3b} , R^{4b} , and R^{5b} are same or different and are independently -OR^{11b} or a hydrogen atom;

An amide compound of the formula (1B) wherein R^{2b} , R^{3b} , R^{4b} , and R^{5b} are same or different and are independently -S(O)_mR^{11b} or a hydrogen atom;

10 An amide compound of the formula (1B) wherein R^{2b} , R^{3b} , R^{4b} , and R^{5b} are same or different and are independently a cyano group, a nitro group, or a hydrogen atom;

15 An amide compound of the formula (1B) wherein R^{2b} , R^{3b} , R^{4b} , and R^{5b} are same or different and are independently a halogen atom or a hydrogen atom;

An amide compound of the formula (1B) wherein R^{2b} , R^{3b} , R^{4b} , and R^{5b} are a hydrogen atom;

20 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a 2-pyridyl group, a 2-pyrimidyl group, a 3-chloro-2-pyridyl group, or a 3-chloro-5-trifluoromethyl-2-pyridyl group;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a 2-pyridyl group;

25 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a 2-pyrimidyl group;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a 3-chloro-2-pyridyl group;

5 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a 3-chloro-5-trifluoromethyl-2-pyridyl group;

10 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a C1-C6 alkyl group;

15 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a C1-C6 alkyl group optionally having one or more fluorine atoms;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a C1-C3 alkyl group;

20 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a C1-C3 alkyl group optionally having one or more fluorine atoms;

25 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, or a heptafluoroisopropyl group;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a methyl group;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is an ethyl group;

5 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a propyl group;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is an isopropyl group;

10 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a trifluoromethyl group;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a pentafluoroethyl group;

15 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a heptafluoropropyl group;

20 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a heptafluoroisopropyl group;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a halogen atom;

25 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a fluorine atom, a chlorine atom, or a bromine atom;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a fluorine atom;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a chlorine atom;

5 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a bromine atom;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is $-SF_5$;

10 An amide compound of the formula (1B) wherein R^{2b} , R^{3b} , R^{4b} , and R^{5b} are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a halogen atom, or a hydrogen atom;

15 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms;

20 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a halogen atom;

25 An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a C1-C3 alkyl group optionally having one or more fluorine atoms or a halogen atom;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a 2-pyridyl group, a 2-pyrimidyl group, a 3-chloro-2-pyridyl group, a 3-chloro-5-trifluoromethyl-2-pyridyl group, a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom;

An amide compound of the formula (1B) wherein R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, R^{3b} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom;

An amide compound of the formula (1B) wherein R^{1b} is a C1-C6 alkyl group, R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

An amide compound of the formula (1B) wherein R^{1b} is a C1-C6 alkyl group, R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, and R^{3b} is a halogen atom;

An amide compound of the formula (1B) wherein R^{6b} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

An amide compound of the formula (1B) wherein R^{6b} is a

C1-C6 chain hydrocarbon group;

An amide compound of the formula (1B) wherein R^{6b} is a C1-C6 chain hydrocarbon group having one or more fluorine atoms;

5 An amide compound of the formula (1B) wherein R^{6b} is a C1-C3 alkyl group having one or more fluorine atoms;

An amide compound of the formula (1B) wherein R^{6b} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, or a heptafluoroisopropyl group;

10 An amide compound of the formula (1B) wherein R^{6b} is a trifluoromethyl group;

An amide compound of the formula (1B) wherein R^{6b} is a pentafluoroethyl group;

15 An amide compound of the formula (1B) wherein R^{6b} is a heptafluoropropyl group;

An amide compound of the formula (1B) wherein R^{6b} is a heptafluoroisopropyl group;

20 An amide compound of the formula (1B) wherein R^{6b} is a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, or a 6-membered heterocyclic group optionally having one or more halogen atoms or a C1-C3 alkyl groups optionally having one or more

25

halogen atoms;

An amide compound of the formula (1B) wherein R^{6b} is -
OR^{11b};

5 An amide compound of the formula (1B) wherein R^{6b} is a
C1-C6 alkoxy group having one or more fluorine atoms;

An amide compound of the formula (1B) wherein R^{6b} is a
C1-C3 alkoxy group having one or more fluorine atoms;

An amide compound of the formula (1B) wherein R^{6b} is a
trifluoromethoxy group;

10 An amide compound of the formula (1B) wherein R^{6b} is -
S(O)_mR^{11b};

An amide compound of the formula (1B) wherein R^{6b} is a
C1-C6 alkylsulfanyl group having one or more fluorine atoms,
a C1-C6 alkylsulfinyl group having one or more fluorine
15 atoms, or a C1-C6 alkylsulfonyl group having one or more
fluorine atoms;

An amide compound of the formula (1B) wherein R^{6b} is a
C1-C3 alkylsulfanyl group having one or more fluorine atoms,
a C1-C3 alkylsulfinyl group having one or more fluorine
20 atoms, or a C1-C3 alkylsulfonyl group having one or more
fluorine atoms;

An amide compound of the formula (1B) wherein R^{6b} is a
trifluoromethylsulfanyl group, a trifluoromethylsulfinyl
group, or a trifluoromethylsulfonyl group;

25 An amide compound of the formula (1B) wherein R^{6b} is a

trifluoromethylsulfanyl group;

An amide compound of the formula (1B) wherein R^{6b} is a trifluoromethylsulfinyl group;

5 An amide compound of the formula (1B) wherein R^{6b} is a trifluoromethylsulfonyl group;

An amide compound of the formula (1B) wherein R^{6b} is a cyano group or a nitro group;

An amide compound of the formula (1B) wherein R^{6b} is a halogen atom;

10 An amide compound of the formula (1B) wherein R^{6b} is a chlorine atom, a bromine atom, or an iodine atom;

An amide compound of the formula (1B) wherein R^{6b} is a chlorine atom;

15 An amide compound of the formula (1B) wherein R^{6b} is a bromine atom;

An amide compound of the formula (1B) wherein R^{6b} is an iodine atom;

An amide compound of the formula (1B) wherein R^{6b} is -SF₅;

20 An amide compound of the formula (1B) wherein R^{6b} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms,
25 a C1-C3 alkylsulfonyl group having one or more fluorine

atoms, or a halogen atom;

An amide compound of the formula (1B) wherein R^{6b} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a
5 trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom;

An amide compound of the formula (1B) wherein R^{7b} is a C1-C6 chain hydrocarbon group optionally having one or more
10 halogen atoms;

An amide compound of the formula (1B) wherein R^{7b} is a C1-C6 chain hydrocarbon group;

An amide compound of the formula (1B) wherein R^{7b} is a C1-C6 chain hydrocarbon group having one or more fluorine
15 atoms;

An amide compound of the formula (1B) wherein R^{7b} is a C1-C3 alkyl group having one or more fluorine atoms;

An amide compound of the formula (1B) wherein R^{7b} is a trifluoromethyl group, a pentafluoroethyl group, a
20 heptafluoropropyl group, or a heptafluoroisopropyl group;

An amide compound of the formula (1B) wherein R^{7b} is a trifluoromethyl group;

An amide compound of the formula (1B) wherein R^{7b} is a pentafluoroethyl group;

25 An amide compound of the formula (1B) wherein R^{7b} is a

heptafluoropropyl group;

An amide compound of the formula (1B) wherein R^{7b} is a heptafluoroisopropyl group;

5 An amide compound of the formula (1B) wherein R^{7b} is a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, or a 6-membered
10 heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms;

An amide compound of the formula (1B) wherein R^{7b} is -
OR^{11b};

15 An amide compound of the formula (1B) wherein R^{7b} is a C1-C6 alkoxy group having one ore more fluorine atoms;

An amide compound of the formula (1B) wherein R^{7b} is a C1-C3 alkoxy group having one or more fluorine atoms;

20 An amide compound of the formula (1B) wherein R^{7b} is a trifluoromethoxy group;

An amide compound of the formula (1B) wherein R^{7b} is -
S(O)_mR^{11b};

25 An amide compound of the formula (1B) wherein R^{7b} is a C1-C6 alkylsulfanyl group having one or more fluorine atoms, a C1-C6 alkylsulfinyl group having one or more fluorine

atoms, or a C1-C6 alkylsulfonyl group having one or more fluorine atoms;

An amide compound of the formula (1B) wherein R^{7b} is a C1-C3 alkylsulfanyl group having one or more fluorine atoms,
5 a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms;

An amide compound of the formula (1B) wherein R^{7b} is a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl
10 group, or a trifluoromethylsulfonyl group;

An amide compound of the formula (1B) wherein R^{7b} is a trifluoromethylsulfanyl group;

An amide compound of the formula (1B) wherein R^{7b} is a trifluoromethylsulfinyl group;

15 An amide compound of the formula (1B) wherein R^{7b} is a trifluoromethylsulfonyl group;

An amide compound of the formula (1B) wherein R^{7b} is a cyano group or a nitro group;

20 An amide compound of the formula (1B) wherein R^{7b} is a halogen atom;

An amide compound of the formula (1B) wherein R^{7b} is a chlorine atom, a bromine atom, or an iodine atom;

An amide compound of the formula (1B) wherein R^{7b} is a chlorine atom;

25 An amide compound of the formula (1B) wherein R^{7b} is a

bromine atom;

An amide compound of the formula (1B) wherein R^{7b} is an iodine atom;

5 An amide compound of the formula (1B) wherein R^{7b} is a hydrogen atom;

An amide compound of the formula (1B) wherein R^{7b} is -SF₅;

10 An amide compound of the formula (1B) wherein R^{7b} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or a halogen atom;

15 An amide compound of the formula (1B) wherein R^{7b} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom;

20 An amide compound of the formula (1B) wherein R^{6b} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms,

25

a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or a halogen atom, and R^{7b} is a hydrogen atom;

An amide compound of the formula (1B) wherein R^{6b} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom, and R^{7b} is a hydrogen atom;

10 An amide compound of the formula (1B) wherein m is 0;

An amide compound of the formula (1B) wherein m is 1;

An amide compound of the formula (1B) wherein m is 2;

An amide compound of the formula (1B) wherein n is 0;

An amide compound of the formula (1B) wherein n is 1;

15 An amide compound of the formula (1B) wherein n is 2;

An amide compound of the formula (1B) wherein A^{1b} is - NR^{8b} - or a sulfur atom, R^{8b} is a C1-C6 alkyl group, A^{2b} is $=CR^{9b}$ -, A^{3b} is $=CR^{10b}$ -, R^{9b} and R^{10b} are a hydrogen atom, R^{1b} is a C1-C3 alkyl group optionally having one or more fluorine atoms, R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, R^{3b} is a C1-C3 alkyl group optionally having one or more fluorine atoms or a halogen atom, R^{6b} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group

20

25

having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or a halogen atom, and R^{7b} is a hydrogen atom;

An amide compound of the formula (1B) wherein A^{1b} is -
5 NR^{8b}- or a sulfur atom, R^{8b} is a C1-C6 alkyl group, A^{2b} is =CR^{9b}-, R^{9b} is a hydrogen atom, A^{3b} is a nitrogen atom, R^{1b} is a C1-C3 alkyl group optionally having one or more fluorine atoms, R^{2b}, R^{4b}, and R^{5b} are a hydrogen atom, R^{3b} is a C1-C3 alkyl group optionally having one or more fluorine
10 atoms or a halogen atom, R^{6b} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl
15 group having one or more fluorine atoms, or a halogen atom, and R^{7b} is a hydrogen atom;

An amide compound of the formula (1B) wherein A^{1b} is -
NR^{8b}-, R^{8b} is a methyl group, an ethyl group, a propyl group, or an isopropyl group, A^{2b} is =CR^{9b}-, A^{3b} is =CR^{10b}-, R^{9b} and
20 R^{10b} are a hydrogen atom, R^{1b} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group or a 2,2,2-trifluoroethyl group, R^{2b}, R^{4b}, and R^{5b} are a hydrogen atom, R^{3b} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a
25 trifluoromethyl group, a pentafluoroethyl group, a

heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom, R^{6b} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom, and R^{7b} is a hydrogen atom;

An amide compound of the formula (1B) wherein A^{1b} is a sulfur atom, A^{2b} is $=CR^{9b}-$, A^{3b} is $=CR^{10b}-$, R^{9b} and R^{10b} are a hydrogen atom, R^{1b} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group or a 2,2,2-trifluoroethyl group, R^{2b} , R^{4b} , and R^{5b} are a hydrogen atom, R^{3b} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom or a bromine atom, R^{6b} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom, and R^{7b} is a hydrogen atom;

An amide compound of the formula (1B) wherein A^{1b} is -

NR^{8b}-, R^{8b} is a methyl group, an ethyl group, a propyl group, or an isopropyl group, A^{2b} is =CR^{9b}-, R^{9b} is a hydrogen atom, A^{3b} is a nitrogen atom, R^{1b} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, or a 2,2,2-trifluoroethyl group, R^{2b},
5 R^{4b}, and R^{5b} are a hydrogen atom, R^{3b} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a
10 fluorine atom, a chlorine atom or a bromine atom, R^{6b} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl
15 group, a chlorine atom, a bromine atom, or an iodine atom, and R^{7b} is a hydrogen atom;

An amide compound of the formula (1B) wherein A^{1b} is a sulfur atom, A^{2b} is =CR^{9b}-, R^{9b} is a hydrogen atom, A^{3b} is a nitrogen atom, R^{1b} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group,
20 or a 2,2,2-trifluoroethyl group, R^{2b}, R^{4b}, and R^{5b} are a hydrogen atom, R^{3b} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a
25 fluorine atom, a chlorine

atom or a bromine atom, R^{6b} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom, and R^{7b} is a hydrogen atom;

[0138]

And examples of Compound (6) include the following compounds:

An amide compound of the formula (6) wherein A^2 is $=CH-$;

An amide compound of the formula (6) wherein A^2 is a nitrogen atom;

An amide compound of the formula (6) wherein A^3 is $=CH-$;

An amide compound of the formula (6) wherein A^3 is a nitrogen atom or $=CR^{10}-$, and R^{10} is a halogen atom or a hydrogen atom;

An amide compound of the formula (6) wherein A^3 is a nitrogen atom;

An amide compound of the formula (6) wherein A^2 is $=CH-$, A^3 is a nitrogen atom or $=CR^{10}-$, and R^{10} is a halogen atom or a hydrogen atom;

An amide compound of the formula (6) wherein A^2 is

=CH-, and A³ is a nitrogen atom;

An amide compound of the formula (6) wherein A² is =CH-, A³ is =CR¹⁰-, and R¹⁰ is a halogen atom or a hydrogen atom;

5 An amide compound of the formula (6) wherein R¹ is a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from Group X;

10 An amide compound of the formula (6) wherein R¹ is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

 An amide compound of the formula (6) wherein R¹ is a C1-C6 chain hydrocarbon group;

 An amide compound of the formula (6) wherein R¹ is a C1-C3 alkyl group;

15 An amide compound of the formula (6) wherein R¹ is a methyl group;

 An amide compound of the formula (6) wherein R¹ is an ethyl group;

20 An amide compound of the formula (6) wherein R¹ is a propyl group;

 An amide compound of the formula (6) wherein R¹ is an isopropyl group;

25 An amide compound of the formula (6) wherein R², R³, R⁴, and R⁵ are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more

halogen atoms, $-OR^{11}$, $-NR^{11}R^{12}$, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

An amide compound of the formula (6) wherein R^2 , R^4 , and R^5 are same or different and are independently a
5 hydrogen atom or a halogen atom, R^3 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, a halogen atom, or a hydrogen atom;

An amide compound of the formula (6) wherein R^2 , R^4 , and R^5 are same or different and are independently a
10 halogen atom or a hydrogen atom, and R^3 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a halogen atom, or a hydrogen atom;

An amide compound of the formula (6) wherein R^2 , R^4 , and R^5 are same or different and are independently a
15 halogen atom or a hydrogen atom, and R^3 is a methyl group, a trifluoromethyl group, a trifluoromethoxy group, a chlorine atom, a bromine atom, an iodine atom, or a hydrogen atom;

An amide compound of the formula (6) wherein R^2 , R^4 ,
20 and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a methyl group;

An amide compound of the formula (6) wherein R^2 , R^4 , and R^5 are same or different and are independently a
25 halogen atom or a hydrogen atom, and R^3 is a trifluoromethyl group;

An amide compound of the formula (6) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a trifluoromethoxy group;

5 An amide compound of the formula (6) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a chlorine atom;

10 An amide compound of the formula (6) wherein R^2 , R^4 , and R^5 are same or different and are independently a halogen atom or a hydrogen atom, and R^3 is a bromine atom;

An amide compound of the formula (6) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

15 An amide compound of the formula (6) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a methyl group, a trifluoromethyl group, a trifluoromethoxy group, a chlorine atom, a bromine atom, an iodine atom, or a hydrogen atom;

20 An amide compound of the formula (6) wherein R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a methyl group, a trifluoromethyl group, a trifluoromethoxy group, a chlorine atom, a bromine atom, or an iodine atom;

An amide compound of the formula (6) wherein R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a methyl group;

25 An amide compound of the formula (6) wherein R^2 , R^4 ,

and R⁵ are a hydrogen atom, and R³ is a trifluoromethyl group;

5 An amide compound of the formula (6) wherein R², R⁴, and R⁵ are a hydrogen atom, and R³ is a trifluoromethoxy group;

An amide compound of the formula (6) wherein R², R⁴, and R⁵ are a hydrogen atom, and R³ is a chlorine atom;

An amide compound of the formula (6) wherein R², R⁴, and R⁵ are a hydrogen atom, and R³ is a bromine atom;

10 An amide compound of the formula (6) wherein R², R³, R⁴, and R⁵ are a hydrogen atom;

An amide compound of the formula (6) wherein R¹ is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, R², R⁴, and R⁵ are same or different and are
15 independently a hydrogen atom or a halogen atom, and R³ is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, -OR¹¹, a halogen atom, or a hydrogen atom;

20 An amide compound of the formula (6) wherein R¹ is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, R², R⁴, and R⁵ are same or different and are independently a halogen atom or a hydrogen atom, and R³ is a methyl group, a trifluoromethyl group, a trifluoromethoxy group, a chlorine atom, a bromine atom, an iodine atom, or
25 a hydrogen atom;

An amide compound of the formula (6) wherein R^1 is a methyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

5 An amide compound of the formula (6) wherein R^1 is a methyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a methyl group or a hydrogen atom;

10 An amide compound of the formula (6) wherein R^1 is a methyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a trifluoromethyl group or a hydrogen atom;

 An amide compound of the formula (6) wherein R^1 is a methyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a chlorine atom or a hydrogen atom;

15 An amide compound of the formula (6) wherein R^1 is a methyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a bromine atom or a hydrogen atom;

20 An amide compound of the formula (6) wherein R^1 is a methyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a methyl group, a trifluoromethyl group, a trifluoromethoxy group, a chlorine atom, a bromine atom, or an iodine atom;

 An amide compound of the formula (6) wherein R^1 is a methyl group, and R^2 , R^3 , R^4 , and R^5 are a hydrogen atom;

25 An amide compound of the formula (6) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a C1-C6 chain hydrocarbon group optionally having one or more

halogen atoms;

An amide compound of the formula (6) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a methyl group or a hydrogen atom;

5 An amide compound of the formula (6) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a trifluoromethyl group or a hydrogen atom;

10 An amide compound of the formula (6) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a chlorine atom or a hydrogen atom;

An amide compound of the formula (6) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a bromine atom or a hydrogen atom;

15 An amide compound of the formula (6) wherein R^1 is an ethyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a methyl group, a trifluoromethyl group, a trifluoromethoxy group, a chlorine atom, a bromine atom, or an iodine atom;

An amide compound of the formula (6) wherein R^1 is an ethyl group, and R^2 , R^3 , R^4 , and R^5 are a hydrogen atom;

20 An amide compound of the formula (6) wherein R^1 is a propyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

25 An amide compound of the formula (6) wherein R^1 is a propyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is

a methyl group or a hydrogen atom;

An amide compound of the formula (6) wherein R^1 is a propyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a trifluoromethyl group or a hydrogen atom;

5 An amide compound of the formula (6) wherein R^1 is a propyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a chlorine atom or a hydrogen atom;

10 An amide compound of the formula (6) wherein R^1 is a propyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a bromine atom or a hydrogen atom;

An amide compound of the formula (6) wherein R^1 is a propyl group, R^2 , R^4 , and R^5 are a hydrogen atom, and R^3 is a methyl group, a trifluoromethyl group, a trifluoromethoxy group, a chlorine atom, a bromine atom, or an iodine atom;

15 An amide compound of the formula (6) wherein R^1 is a propyl group, and R^2 , R^3 , R^4 , and R^5 are a hydrogen atom;

20 An amide compound of the formula (6) wherein R^6 and R^7 are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, $-NR^{11}R^{12}$, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

25 An amide compound of the formula (6) wherein R^6 and R^7 are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, a halogen atom, or a hydrogen atom;

An amide compound of the formula (6) wherein R^6 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, or a halogen atom, and R^7 is a hydrogen atom;

5 An amide compound of the formula (6) wherein R^6 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, and R^7 is a hydrogen atom;

10 An amide compound of the formula (6) wherein R^6 is a C1-C6 chain hydrocarbon group having one or more halogen atoms, and R^7 is a hydrogen atom;

 An amide compound of the formula (6) wherein R^6 is a C1-C6 chain hydrocarbon group having one or more fluorine atoms, and R^7 is a hydrogen atom;

15 An amide compound of the formula (6) wherein R^6 is $-OR^{11}$, and R^7 is a hydrogen atom;

 An amide compound of the formula (6) wherein R^6 is $-S(O)_mR^{11}$, and R^7 is a hydrogen atom;

20 An amide compound of the formula (6) wherein R^6 is a C1-C6 chain hydrocarbon group having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, or a halogen atom, and R^7 is a hydrogen atom;

25 An amide compound of the formula (6) wherein R^6 is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl

group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom, and R⁷ is a hydrogen atom;

An amide compound of the formula (6) wherein R⁸ is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a cyclopropyl group;

An amide compound of the formula (6) wherein R⁸ is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group;

An amide compound of the formula (6) wherein R⁸ is a methyl group;

An amide compound of the formula (6) wherein A² is =CH-, A³ is a nitrogen atom or =CR¹⁰-, R¹⁰ is a hydrogen atom, R⁸ is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, R¹ is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, R², R⁴, and R⁵ are same or different and are independently a hydrogen atom or a halogen atom, R³ is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, -OR¹¹, a halogen atom, or a hydrogen atom, and R⁶ and R⁷ are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, -OR¹¹, -S(O)_mR¹¹, a halogen atom, or a hydrogen atom;


An amide compound of the formula (6) wherein A^2 is $=CH-$, A^3 is a nitrogen atom, R^8 is a methyl group or a hydrogen atom, R^1 is a methyl group, an ethyl group or propyl group, R^2 , R^4 , and R^5 are same or different and are
5 independently a halogen atom or a hydrogen atom, R^3 is a trifluoromethyl group, a halogen atom or a hydrogen atom, R^6 is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms,
10 a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms, and R^7 is a hydrogen atom;

An amide compound of the formula (6) wherein A^2 is $=CH-$, A^3 is a nitrogen atom or $=CR^{10}-$, R^{10} is a hydrogen
15 atom, R^8 is a methyl group or a hydrogen atom, R^1 is a methyl group, an ethyl group, or propyl group, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a methyl group, a trifluoromethyl group, a trifluoromethoxy group, a chlorine atom, a bromine atom, or an iodine atom, R^6 is a
20 trifluoromethyl group, a pentafluoroethyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a bromine atom, or an iodine atom, and R^7 is a hydrogen atom.

Examples of Compound (C) include the following compounds:

[0140]

An amide compound of the formula (M10):




5

wherein, V², R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, A² and A³ are as defined above (referred to as Compound (M10));

[0141]

An amide compound of the formula (M25):




10

wherein, V², R², R³, R⁴, R⁵, R⁶, R⁷, A² and A³ are as defined above (referred to as Compound (M25));

[0142]

An amide compound of the formula (M26):




15

wherein, V², R², R³, R⁴, R⁵, R⁶, R⁷, A² and A³ are as defined

above (referred to as Compound (M26));

[0143]

A compound of the formula (1C):



5 wherein,

X^c represents a fluorine atom or a chlorine atom;

A^{1c} represents $-NR^{8c}-$, an oxygen atom, or a sulfur atom;

A^{2c} represents a nitrogen atom or $=CR^{9c}-$;

10 A^{3c} represents a nitrogen atom or $=CR^{10c}-$;

R^{2c} , R^{3c} , and R^{5c} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, $-OR^{11c}$, $-S(O)_mR^{11c}$, $-SF_5$, a cyano group, a halogen atom, or a hydrogen atom: provided that at least one of R^{2c} , R^{3c} , and

15

20

R^{5c} represent a hydrogen atom;

R^{4c} represents a hydrogen atom;

R^{6c} represents a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or -SF₅;

R^{7c} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, -OR^{11c}, -S(O)_mR^{11c}, -SF₅, a cyano group, a halogen atom, or a hydrogen atom;

R^{8c} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group, a C3-C6 alicyclic hydrocarbon group optionally having one or more halogen atoms, or a hydrogen atom;

R^{9c} and R^{10c} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally

having one or more halogen atoms, $-OR^{11c}$, $-S(O)_mR^{11c}$, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

R^{11c} and R^{12c} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally

5 having one or more halogen atoms or a hydrogen atom; and

m represents 0, 1, or 2;

provided that: in $-S(O)_mR^{11c}$, when m is 1 or 2, R^{11c} is not a hydrogen atom; and when A^{1c} is $-NR^{8c}-$, and A^{2c} and A^{3c} are a methine group, R^{8c} is a hydrogen atom or a methyl group (hereinafter, referred to as Compound (1C));

[0144]

And examples of Compound (1C) include the following compounds:

15 An amide compound of the formula (1C) wherein X^c is a fluorine atom;

An amide compound of the formula (1C) wherein X^c is a chlorine atom;

An amide compound of the formula (1C) wherein A^{1c} is $-NR^{8c}-$;

20 An amide compound of the formula (1C) wherein A^{1c} is $-NR^{8c}-$, and R^{8c} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

An amide compound of the formula (1C) wherein A^{1c} is $-NR^{8c}-$, and R^{8c} is a C1-C6 chain hydrocarbon group;

25 An amide compound of the formula (1C) wherein A^{1c} is $-$

NR^{8c}-, and R^{8c} is a C1-C6 alkyl group;

An amide compound of the formula (1C) wherein A^{1c} is - NR^{8c}-, and R^{8c} is a C1-C3 alkyl group;

5 An amide compound of the formula (1C) wherein A^{1c} is - NR^{8c}-, and R^{8c} is a methyl group, an ethyl group, a propyl group, or an isopropyl group;

An amide compound of the formula (1C) wherein A^{1c} is - NR^{8c}-, and R^{8c} is a methyl group;

10 An amide compound of the formula (1C) wherein A^{1c} is - NR^{8c}-, and R^{8c} is an ethyl group;

An amide compound of the formula (1C) wherein A^{1c} is - NR^{8c}-, and R^{8c} is a propyl group;

An amide compound of the formula (1C) wherein A^{1c} is - NR^{8c}-, and R^{8c} is an isopropyl group;

15 An amide compound of the formula (1C) wherein A^{1c} is - NR^{8c}-, and R^{8c} is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group;

20 An amide compound of the formula (1C) wherein A^{1c} is - NR^{8c}-, and R^{8c} is a methoxymethyl group or an ethoxymethyl group;

An amide compound of the formula (1C) wherein A^{1c} is - NR^{8c}-, and R^{8c} is a C3-C6 alicyclic hydrocarbon group optionally having one or more halogen atoms;

25 An amide compound of the formula (1C) wherein A^{1c} is - NR^{8c}-, and R^{8c} is a cyclopropyl group, a cyclobutyl group, a

cyclopentyl group, or a cyclohexyl group;

An amide compound of the formula (1C) wherein A^{1c} is -
NR^{8c}-, and R^{8c} is a hydrogen atom;

5 An amide compound of the formula (1C) wherein A^{1c} is
an oxygen atom;

An amide compound of the formula (1C) wherein A^{1c} is a
sulfur atom;

An amide compound of the formula (1C) wherein A^{1c} is -
NR^{8c}- or a sulfur atom;

10 An amide compound of the formula (1C) wherein A^{1c} is -
NR^{8c}- or a sulfur atom, and R^{8c} is a methyl group, an ethyl
group, a propyl group, or an isopropyl group;

An amide compound of the formula (1C) wherein A^{1c} is -
NR^{8c}- or a sulfur atom, and R^{8c} is a methoxymethyl group, an
15 ethoxymethyl group, or a hydrogen atom;

An amide compound of the formula (1C) wherein A^{2c} is
=CR^{9c}-, and A^{3c} is =CR^{10c}-;

An amide compound of the formula (1C) wherein A^{2c} is
=CR^{9c}-, A^{3c} is =CR^{10c}-, and R^{9c} and R^{10c} are same or different
20 and are independently a halogen atom or a hydrogen atom;

An amide compound of the formula (1C) wherein A^{2c} is
=CR^{9c}-, A^{3c} is =CR^{10c}-, and R^{9c} and R^{10c} are a hydrogen atom;

An amide compound of the formula (1C) wherein A^{2c} is a
nitrogen atom, and A^{3c} is =CR^{10c}-;

25 An amide compound of the formula (1C) wherein A^{2c} is

$=\text{CR}^{9c}-$, and A^{3c} is a nitrogen atom;

An amide compound of the formula (1C) wherein A^{2c} is $=\text{CR}^{9c}-$, R^{9c} is a halogen atom or a hydrogen atom, and A^{3c} is a nitrogen atom;

5 An amide compound of the formula (1C) wherein A^{2c} is $=\text{CR}^{9c}-$, R^{9c} is a hydrogen atom, and A^{3c} is a nitrogen atom;

An amide compound of the formula (1C) wherein A^{2c} is a nitrogen atom, and A^{3c} is a nitrogen atom;

10 An amide compound of the formula (1C) wherein A^{1c} is -
 $\text{NR}^{8c}-$, A^{2c} is $=\text{CR}^{9c}-$, and A^{3c} is $=\text{CR}^{10c}-$;

An amide compound of the formula (1C) wherein A^{1c} is -
 $\text{NR}^{8c}-$, A^{2c} is $=\text{CR}^{9c}-$, A^{3c} is $=\text{CR}^{10c}-$, and R^{9c} and R^{10c} are a
hydrogen atom;

15 An amide compound of the formula (1C) wherein A^{1c} is
an oxygen atom, A^{2c} is $=\text{CR}^{9c}-$, and A^{3c} is $=\text{CR}^{10c}-$;

An amide compound of the formula (1C) wherein A^{1c} is a
sulfur atom, A^{2c} is $=\text{CR}^{9c}-$, and A^{3c} is $=\text{CR}^{10c}-$;

20 An amide compound of the formula (1C) wherein A^{1c} is a
sulfur atom, A^{2c} is $=\text{CR}^{9c}-$, A^{3c} is $=\text{CR}^{10c}-$, and R^{9c} and R^{10c}
are a hydrogen atom;

An amide compound of the formula (1C) wherein A^{1c} is -
 $\text{NR}^{8c}-$, A^{2c} is $=\text{CR}^{9c}-$, and A^{3c} is a nitrogen atom;

25 An amide compound of the formula (1C) wherein A^{1c} is -
 $\text{NR}^{8c}-$, A^{2c} is $=\text{CR}^{9c}-$, R^{9c} is a hydrogen atom, and A^{3c} is a
nitrogen atom;

An amide compound of the formula (1C) wherein A^{1c} is an oxygen atom, A^{2c} is $=CR^{9c}-$, and A^{3c} is a nitrogen atom;

An amide compound of the formula (1C) wherein A^{1c} is a sulfur atom, A^{2c} is $=CR^{9c}-$, and A^{3c} is a nitrogen atom;

5 An amide compound of the formula (1C) wherein A^{1c} is a sulfur atom, A^{2c} is $=CR^{9c}-$, R^{9c} is a hydrogen atom, and A^{3c} is a nitrogen atom;

10 An amide compound of the formula (1C) wherein A^{1c} is -
 $NR^{8c}-$ or a sulfur atom, R^{8c} is a methyl group, an ethyl group, a propyl group or an isopropyl group, A^{2c} is $=CR^{9c}-$,
 A^{3c} is $=CR^{10c}-$, and R^{9c} and R^{10c} are a hydrogen atom;

An amide compound of the formula (1C) wherein A^{1c} is -
 $NR^{8c}-$ or a sulfur atom, R^{8c} is a methyl group, A^{2c} is $=CR^{9c}-$,
 A^{3c} is $=CR^{10c}-$, and R^{9c} and R^{10c} are a hydrogen atom;

15 An amide compound of the formula (1C) wherein A^{1c} is -
 $NR^{8c}-$ or a sulfur atom, R^{8c} is a methyl group, an ethyl group, a propyl group or an isopropyl group, A^{2c} is $=CR^{9c}-$,
 R^{9c} is a hydrogen atom, and A^{3c} is a nitrogen atom;

20 An amide compound of the formula (1C) wherein A^{1c} is -
 $NR^{8c}-$ or a sulfur atom, R^{8c} is a methyl group, A^{2c} is $=CR^{9c}-$,
 R^{9c} is a hydrogen atom, and A^{3c} is a nitrogen atom;

25 An amide compound of the formula (1C) wherein R^{2c} , R^{3c} , and R^{5c} are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a hydrogen atom;

An amide compound of the formula (1C) wherein R^{2c} , R^{3c} , and R^{5c} are same or different and are independently a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, or a hydrogen atom;

An amide compound of the formula (1C) wherein R^{2c} , R^{3c} , and R^{5c} are same or different and are independently $-OR^{11c}$ or a hydrogen atom;

An amide compound of the formula (1C) wherein R^{2c} , R^{3c} , and R^{5c} are same or different and are independently $-S(O)_mR^{11c}$ or a hydrogen atom;

An amide compound of the formula (1C) wherein R^{2c} , R^{3c} , and R^{5c} are same or different and are independently a cyano group, a nitro group, or a hydrogen atom;

An amide compound of the formula (1C) wherein R^{2c} , R^{3c} , and R^{5c} are same or different and are independently a halogen atom or a hydrogen atom;

An amide compound of the formula (1C) wherein R^{2c} , R^{3c} , and R^{5c} is a hydrogen atom;

An amide compound of the formula (1C) wherein R^{2c} and

R^{5c} are a hydrogen atom, and R^{3c} is a 2-pyridyl group, a 2-pyrimidyl group, a 3-chloro-2-pyridyl group, or a 3-chloro-5-trifluoromethyl-2-pyridyl group;

5 An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a 2-pyridyl group;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a 2-pyrimidyl group;

10 An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a 3-chloro-2-pyridyl group;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a 3-chloro-5-trifluoromethyl-2-pyridyl group;

15 An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a C1-C6 alkyl group;

20 An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a C1-C6 alkyl group optionally having one or more fluorine atoms;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a C1-C3 alkyl group;

25 An amide compound of the formula (1C) wherein R^{2c} and

R^{5c} are a hydrogen atom, and R^{3c} is a C1-C3 alkyl group optionally having one or more fluorine atoms;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, or a heptafluoroisopropyl group;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a methyl group;

10 An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is an ethyl group;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a propyl group;

15 An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is an isopropyl group;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a trifluoromethyl group;

20 An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a pentafluoroethyl group;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a heptafluoropropyl group;

25 An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a heptafluoroisopropyl

group;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a halogen atom;

5 An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a fluorine atom, a chlorine atom, or a bromine atom;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a fluorine atom;

10 An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a chlorine atom;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a bromine atom;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is $-SF_5$;

15 An amide compound of the formula (1C) wherein R^{2c} , R^{3c} , and R^{5c} are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a halogen atom, or a hydrogen atom;

20 An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms;

25 An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a C1-C6 chain

hydrocarbon group optionally having one or more halogen atoms or a halogen atom;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a C1-C3 alkyl group optionally having one or more fluorine atoms or a halogen atom;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a 2-pyridyl group, a 2-pyrimidyl group, a 3-chloro-2-pyridyl group, a 3-chloro-5-trifluoromethyl-2-pyridyl group, a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom;

An amide compound of the formula (1C) wherein R^{2c} and R^{5c} are a hydrogen atom, and R^{3c} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom;

An amide compound of the formula (1C) wherein R^{6c} is a C1-C3 alkyl group having one or more fluorine atoms;

An amide compound of the formula (1C) wherein R^{6c} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, or a heptafluoroisopropyl group;

An amide compound of the formula (1C) wherein R^{6c} is a trifluoromethyl group;

An amide compound of the formula (1C) wherein R^{6c} is a pentafluoroethyl group;

5 An amide compound of the formula (1C) wherein R^{6c} is a heptafluoropropyl group;

An amide compound of the formula (1C) wherein R^{6c} is a heptafluoroisopropyl group;

10 An amide compound of the formula (1C) wherein R^{6c} is a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms;

15 An amide compound of the formula (1C) wherein R^{6c} is a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

An amide compound of the formula (1C) wherein R^{6c} is a trifluoromethylsulfanyl group;

20 An amide compound of the formula (1C) wherein R^{6c} is a trifluoromethylsulfinyl group;

An amide compound of the formula (1C) wherein R^{6c} is a trifluoromethylsulfonyl group;

An amide compound of the formula (1C) wherein R^{6c} is -
 SF_5 ;

25 An amide compound of the formula (1C) wherein R^{6c} is a

C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms;

An amide compound of the formula (1C) wherein R^{6c} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

An amide compound of the formula (1C) wherein R^{7c} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

An amide compound of the formula (1C) wherein R^{7c} is a C1-C6 chain hydrocarbon group;

An amide compound of the formula (1C) wherein R^{7c} is a C1-C6 chain hydrocarbon group having one or more fluorine atoms;

An amide compound of the formula (1C) wherein R^{7c} is a C1-C3 alkyl group having one or more fluorine atoms;

An amide compound of the formula (1C) wherein R^{7c} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, or a heptafluoroisopropyl group;

An amide compound of the formula (1C) wherein R^{7c} is a

trifluoromethyl group;

An amide compound of the formula (1C) wherein R^{7c} is a pentafluoroethyl group;

5 An amide compound of the formula (1C) wherein R^{7c} is a heptafluoropropyl group;

An amide compound of the formula (1C) wherein R^{7c} is a heptafluoroisopropyl group;

10 An amide compound of the formula (1C) wherein R^{7c} is a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, or a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms;

An amide compound of the formula (1C) wherein R^{7c} is -
OR^{11c};

20 An amide compound of the formula (1C) wherein R^{7c} is a C1-C6 alkoxy group having one or more fluorine atoms;

An amide compound of the formula (1C) wherein R^{7c} is a C1-C3 alkoxy group having one or more fluorine atoms;

An amide compound of the formula (1C) wherein R^{7c} is a trifluoromethoxy group;

25 An amide compound of the formula (1C) wherein R^{7c} is -

$S(O)_mR^{11c}$;

An amide compound of the formula (1C) wherein R^{7c} is a C1-C6 alkylsulfanyl group having one or more fluorine atoms, a C1-C6 alkylsulfinyl group having one or more fluorine atoms, or a C1-C6 alkylsulfonyl group having one or more fluorine atoms;

An amide compound of the formula (1C) wherein R^{7c} is a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms;

An amide compound of the formula (1C) wherein R^{7c} is a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

An amide compound of the formula (1C) wherein R^{7c} is a trifluoromethylsulfanyl group;

An amide compound of the formula (1C) wherein R^{7c} is a trifluoromethylsulfinyl group;

An amide compound of the formula (1C) wherein R^{7c} is a trifluoromethylsulfonyl group;

An amide compound of the formula (1C) wherein R^{7c} is a cyano group or a nitro group;

An amide compound of the formula (1C) wherein R^{7c} is a halogen atom;

An amide compound of the formula (1C) wherein R^{7c} is a

chlorine atom, a bromine atom, or an iodine atom;

An amide compound of the formula (1C) wherein R^{7c} is a chlorine atom;

5 An amide compound of the formula (1C) wherein R^{7c} is a bromine atom;

An amide compound of the formula (1C) wherein R^{7c} is an iodine atom;

An amide compound of the formula (1C) wherein R^{7c} is a hydrogen atom;

10 An amide compound of the formula (1C) wherein R^{7c} is -SF₅;

An amide compound of the formula (1C) wherein R^{7c} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or a halogen atom;

15

An amide compound of the formula (1C) wherein R^{7c} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom;

20

25 An amide compound of the formula (1C) wherein R^{6c} is a

C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms, and R^{7c} is a hydrogen atom;

An amide compound of the formula (1C) wherein R^{6c} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, and R^{7c} is a hydrogen atom;

An amide compound of the formula (1C) wherein m is 0;

An amide compound of the formula (1C) wherein m is 1;

15 An amide compound of the formula (1C) wherein m is 2;

An amide compound of the formula (1C) wherein A^{1c} is -NR^{8c}- or a sulfur atom, R^{8c} is a C1-C6 alkyl group, A^{2c} is =CR^{9c}-, A^{3c} is =CR^{10c}-, R^{9c} and R^{10c} are a hydrogen atom, R^{2c} and R^{5c} are a hydrogen atom, R^{3c} is a C1-C3 alkyl group optionally having one or more fluorine atoms or a halogen atom, R^{6c} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms, and R^{7c} is a hydrogen

20

25

atom;

An amide compound of the formula (1C) wherein A^{1c} is -
NR^{8c}- or a sulfur atom, R^{8c} is a C1-C6 alkyl group, A^{2c} is
=CR^{9c}-, R^{9c} is a hydrogen atom, A^{3c} is a nitrogen atom, R^{2c}
5 and R^{5c} are a hydrogen atom, R^{3c} is a C1-C3 alkyl group
optionally having one or more fluorine atoms or a halogen
atom, R^{6c} is a C1-C3 alkyl group having one or more
fluorine atoms, a C1-C3 alkylsulfanyl group having one or
more fluorine atoms, a C1-C3 alkylsulfinyl group having one
10 or more fluorine atoms, or a C1-C3 alkylsulfonyl group
having one or more fluorine atoms, and R^{7c} is a hydrogen
atom;

An amide compound of the formula (1C) wherein A^{1c} is -
NR^{8c}-, R^{8c} is a methyl group, an ethyl group, a propyl group,
15 or an isopropyl group, A^{2c} is =CR^{9c}-, A^{3c} is =CR^{10c}-, R^{9c} and
R^{10c} are a hydrogen atom, R^{2c} and R^{5c} are a hydrogen atom, R^{3c}
is a methyl group, an ethyl group, a propyl group, an
isopropyl group, a trifluoromethyl group, a
pentafluoroethyl group, a heptafluoropropyl group, a
20 heptafluoroisopropyl group, a fluorine atom, a chlorine
atom, or a bromine atom, R^{6c} is a trifluoromethyl group, a
pentafluoroethyl group, a heptafluoropropyl group, a
heptafluoroisopropyl group, a trifluoromethylsulfanyl group,
a trifluoromethylsulfinyl group, or a
25 trifluoromethylsulfonyl group, and R^{7c} is a hydrogen atom;

An amide compound of the formula (1C) wherein A^{1c} is a sulfur atom, A^{2c} is $=CR^{9c}-$, A^{3c} is $=CR^{10c}-$, R^{9c} and R^{10c} are a hydrogen atom, R^{2c} and R^{5c} are a hydrogen atom, R^{3c} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom, R^{6c} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, and R^{7c} is a hydrogen atom;


An amide compound of the formula wherein A^{1c} is $-NR^{8c}-$, R^{8c} is a methyl group, an ethyl group, a propyl group, or an isopropyl group, A^{2c} is $=CR^{9c}-$, R^{9c} is a hydrogen atom, A^{3c} is a nitrogen atom, R^{2c} and R^{5c} are a hydrogen atom, R^{3c} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom, R^{6c} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, and R^{7c} is a hydrogen atom;

An amide compound of the formula (1C) wherein A^{1c} is a sulfur atom, A^{2c} is $=CR^{9c}-$, R^{9c} is a hydrogen atom, A^{3c} is a nitrogen atom, R^{2c} and R^{5c} are a hydrogen atom, R^{3c} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom or a bromine atom, R^{6c} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, and R^{7c} is a hydrogen atom.

[0145]

Examples of Compound (24) include the following compounds:

A compound of the formula (1D):



wherein:

A^{1d} represents $-NR^{8d}-$, an oxygen atom, or a sulfur atom;

A^{2d} represents a nitrogen atom or $=CR^{9d}-$;

A^{3d} represents a nitrogen atom or $=CR^{10d}-$;

R^{2d} , R^{3d} , R^{4d} , and R^{5d} are same or different and

represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, -OR^{11d},
10 -S(O)_mR^{11d}, -SF₅, a cyano group, a nitro group, a halogen atom, or a hydrogen atom: provided that at least two of R^{2d}, R^{3d}, R^{4d}, and R^{5d} represent a hydrogen atom;

R^{6d} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, -OR^{11d},
15 -S(O)_mR^{11d}, -SF₅, a cyano group, a nitro group, or a halogen atom;

R^{7d} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group
25

optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, $-OR^{11d}$, $-S(O)_mR^{11d}$, $-SF_5$, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

R^{8d} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group, a C3-C6 alicyclic hydrocarbon group optionally having one or more halogen atoms, or a hydrogen atom;

R^{9d} and R^{10d} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

R^{11d} and R^{12d} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a hydrogen atom; and

m represents 0, 1, or 2:

provided that in $-S(O)_mR^{11d}$, R^{11d} is not a hydrogen atom (hereinafter, referred to as Compound (1D));

[0146]

And examples of Compound (1D) include the following compounds:

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$,
5 and R^{8d} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$,
and R^{8d} is a C1-C6 chain hydrocarbon group;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$,
10 and R^{8d} is a C1-C6 alkyl group;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$,
and R^{8d} is a C1-C3 alkyl group;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$,
and R^{8d} is a methyl group, an ethyl group, a propyl group,
15 or an isopropyl group;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$,
and R^{8d} is a methyl group;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$,
and R^{8d} is an ethyl group;

20 A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$,
and R^{8d} is a propyl group;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$,
and R^{8d} is an isopropyl group;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$,
25 and R^{8d} is a C1-C6 chain hydrocarbon group having one C1-C6

alkoxy group;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$, and R^{8d} is a methoxymethyl group or an ethoxymethyl group;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$,
5 and R^{8d} is a C3-C6 alicyclic hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$, and R^{8d} is a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, or a cyclohexyl group;

10 A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$, and R^{8d} is a hydrogen atom;

A compound of the formula (1D) wherein A^{1d} is an oxygen atom;

A compound of the formula (1D) wherein A^{1d} is a sulfur
15 atom;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$ or a sulfur atom;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$ or a sulfur atom, and R^{8d} is a methyl group, an ethyl group, a
20 propyl group, or an isopropyl group;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$ or a sulfur atom, and R^{8d} is a methoxymethyl group, an ethoxymethyl group, or a hydrogen atom;

A compound of the formula (1D) wherein A^{2d} is $=CR^{9d}-$,
25 and A^{3d} is $=CR^{10d}-$;

A compound of the formula (1D) wherein A^{2d} is $=CR^{9d}-$, A^{3d} is $=CR^{10d}-$, and R^{9d} and R^{10d} are same or different and are independently a halogen atom or a hydrogen atom;

A compound of the formula (1D) wherein A^{2d} is $=CR^{9d}-$,
5 A^{3d} is $=CR^{10d}-$, and R^{9d} and R^{10d} are a hydrogen atom;

A compound of the formula (1D) wherein A^{2d} is a nitrogen atom, and A^{3d} is $=CR^{10d}-$;

A compound of the formula (1D) wherein A^{2d} is $=CR^{9d}-$, and A^{3d} is a nitrogen atom;

10 A compound of the formula (1D) wherein A^{2d} is $=CR^{9d}-$, R^{9d} is a halogen atom or a hydrogen atom, and A^{3d} is a nitrogen atom;

A compound of the formula (1D) wherein A^{2d} is $=CR^{9d}-$, R^{9d} is a hydrogen atom, and A^{3d} is a nitrogen atom;

15 A compound of the formula (1D) wherein A^{2d} is a nitrogen atom, and A^{3d} is a nitrogen atom;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$, A^{2d} is $=CR^{9d}-$, and A^{3d} is $=CR^{10d}-$;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$,
20 A^{2d} is $=CR^{9d}-$, A^{3d} is $=CR^{10d}-$, and R^{9d} and R^{10d} are a hydrogen atom;

A compound of the formula (1D) wherein A^{1d} is an oxygen atom, A^{2d} is $=CR^{9d}-$, and A^{3d} is $=CR^{10d}-$;

A compound of the formula (1D) wherein A^{1d} is a sulfur
25 atom, A^{2d} is $=CR^{9d}-$, and A^{3d} is $=CR^{10d}-$;

A compound of the formula (1D) wherein A^{1d} is a sulfur atom, A^{2d} is $=CR^{9d}-$, A^{3d} is $=CR^{10d}-$, and R^{9d} and R^{10d} are a hydrogen atom;

5 A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$, A^{2d} is $=CR^{9d}-$, and A^{3d} is a nitrogen atom;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$, A^{2d} is $=CR^{9d}-$, R^{9d} is a hydrogen atom, and A^{3d} is a nitrogen atom;

10 A compound of the formula (1D) wherein A^{1d} is an oxygen atom, A^{2d} is $=CR^{9d}-$, and A^{3d} is a nitrogen atom;

A compound of the formula (1D) wherein A^{1d} is a sulfur atom, A^{2d} is $=CR^{9d}-$, and A^{3d} is a nitrogen atom;

15 A compound of the formula (1D) wherein A^{1d} is a sulfur atom, A^{2d} is $=CR^{9d}-$, R^{9d} is a hydrogen atom, and A^{3d} is a nitrogen atom;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$ or a sulfur atom, R^{8d} is a methyl group, an ethyl group, a propyl group or an isopropyl group, A^{2d} is $=CR^{9d}-$, A^{3d} is $=CR^{10d}-$, and R^{9d} and R^{10d} are a hydrogen atom;

20 A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$ or a sulfur atom, R^{8d} is a methyl group, A^{2d} is $=CR^{9d}-$, A^{3d} is $=CR^{10d}-$, and R^{9d} and R^{10d} are a hydrogen atom;

25 A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$ or a sulfur atom, R^{8d} is a methyl group, an ethyl group, a propyl group, or an isopropyl group, A^{2d} is $=CR^{9d}-$, R^{9d} is a

hydrogen atom, and A^{3d} is a nitrogen atom;

A compound of the formula (1D) wherein A^{1d} is -NR^{8d}- or a sulfur atom, R^{8d} is a methyl group, A^{2d} is =CR^{9d}-, R^{9d} is a hydrogen atom, and A^{3d} is a nitrogen atom;

5 A compound of the formula (1D) wherein R^{2d}, R^{3d}, R^{4d}, and R^{5d} are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a hydrogen atom;

10 A compound of the formula (1D) wherein R^{2d}, R^{3d}, R^{4d}, and R^{5d} are same or different and are independently a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally
15 having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, or a hydrogen atom;

20 A compound of the formula (1D) wherein R^{2d}, R^{3d}, R^{4d}, and R^{5d} are same or different and are independently -OR^{11d} or a hydrogen atom;

A compound of the formula (1D) wherein R^{2d}, R^{3d}, R^{4d}, and R^{5d} are same or different and are independently -S(O)_mR^{11d} or a hydrogen atom;

25 A compound of the formula (1D) wherein R^{2d}, R^{3d}, R^{4d},

and R^{5d} are same or different and are independently a cyano group, a nitro group, or a hydrogen atom;

A compound of the formula (1D) wherein R^{2d}, R^{3d}, R^{4d}, and R^{5d} are same or different and are independently a
5 halogen atom or a hydrogen atom;

A compound of the formula (1D) wherein R^{2d}, R^{3d}, R^{4d}, and R^{5d} are a hydrogen atom;

A compound of the formula (1D) wherein R^{2d}, R^{4d}, and R^{5d} are a hydrogen atom, and R^{3d} is a 2-pyridyl group, a 2-pyrimidyl group, a 3-chloro-2-pyridyl group, or a 3-chloro-
10 5-trifluoromethyl-2-pyridyl group;

A compound of the formula (1D) wherein R^{2d}, R^{4d}, and R^{5d} are a hydrogen atom, and R^{3d} is a 2-pyridyl group;

A compound of the formula (1D) wherein R^{2d}, R^{4d}, and R^{5d}
15 are a hydrogen atom, and R^{3d} is a 2-pyrimidyl group;

A compound of the formula (1D) wherein R^{2d}, R^{4d}, and R^{5d} are a hydrogen atom, and R^{3d} is a 3-chloro-2-pyridyl group;

A compound of the formula (1D) wherein R^{2d}, R^{4d}, and R^{5d} are a hydrogen atom, and R^{3d} is a 3-chloro-5-trifluoromethyl-2-pyridyl group;
20

A compound of the formula (1D) wherein R^{2d}, R^{4d}, and R^{5d} are a hydrogen atom, and R^{3d} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (1D) wherein R^{2d}, R^{4d}, and R^{5d}
25 are a hydrogen atom, and R^{3d} is a C1-C6 alkyl group;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a C1-C6 alkyl group optionally having one or more fluorine atoms;

5 A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a C1-C3 alkyl group;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a C1-C3 alkyl group optionally having one or more fluorine atoms;

10 A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, or a heptafluoroisopropyl group;

15 A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a methyl group;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is an ethyl group;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a propyl group;

20 A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is an isopropyl group;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a trifluoromethyl group;

25 A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a pentafluoroethyl group;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a heptafluoropropyl group;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a heptafluoroisopropyl group;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a halogen atom;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a fluorine atom, a chlorine atom or a bromine atom;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a fluorine atom;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a chlorine atom;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a bromine atom;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is $-SF_5$;

A compound of the formula (1D) wherein R^{2d} , R^{3d} , R^{4d} , and R^{5d} are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a halogen atom, or a hydrogen atom;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3

alkyl groups optionally having one or more halogen atoms;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a
5 halogen atom;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a C1-C3 alkyl group optionally having one or more fluorine atoms or a halogen atom;

10 A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a 2-pyridyl group, a 2-pyrimidyl group, a 3-chloro-2-pyridyl group, a 3-chloro-5-trifluoromethyl-2-pyridyl group, a methyl group, an ethyl group, a propyl group, an isopropyl group, a
15 trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom;

A compound of the formula (1D) wherein R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, and R^{3d} is a methyl group, an ethyl
20 group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom;

A compound of the formula (1D) wherein R^{6d} is a C1-C6
25 chain hydrocarbon group optionally having one or more

halogen atoms;

A compound of the formula (1D) wherein R^{6d} is a C1-C6 chain hydrocarbon group;

5 A compound of the formula (1D) wherein R^{6d} is a C1-C6 chain hydrocarbon group having one or more fluorine atoms;

A compound of the formula (1D) wherein R^{6d} is a C1-C3 alkyl group having one or more fluorine atoms;

10 A compound of the formula (1D) wherein R^{6d} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, or a heptafluoroisopropyl group;

A compound of the formula (1D) wherein R^{6d} is a trifluoromethyl group;

A compound of the formula (1D) wherein R^{6d} is a pentafluoroethyl group;

15 A compound of the formula (1D) wherein R^{6d} is a heptafluoropropyl group;

A compound of the formula (1D) wherein R^{6d} is a heptafluoroisopropyl group;

20 A compound of the formula (1D) wherein R^{6d} is a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, or a 6-membered heterocyclic group
25 optionally having one or more halogen atoms or C1-C3 alkyl

groups optionally having one or more halogen atoms;

A compound of the formula (1D) wherein R^{6d} is $-OR^{11d}$;

A compound of the formula (1D) wherein R^{6d} is a C1-C6 alkoxy group having one or more fluorine atoms;

5 A compound of the formula (1D) wherein R^{6d} is a C1-C3 alkoxy group having one or more fluorine atoms;

A compound of the formula (1D) wherein R^{6d} is a trifluoromethoxy group;

10 A compound of the formula (1D) wherein R^{6d} is $-S(O)_mR^{11d}$;

15 A compound of the formula (1D) wherein R^{6d} is a C1-C6 alkylsulfanyl group having one or more fluorine atoms, a C1-C6 alkylsulfinyl group having one or more fluorine atoms, or a C1-C6 alkylsulfonyl group having one or more fluorine atoms;

20 A compound of the formula (1D) wherein R^{6d} is a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms;

A compound of the formula (1D) wherein R^{6d} is a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

25 A compound of the formula (1D) wherein R^{6d} is a trifluoromethylsulfanyl group;

A compound of the formula (1D) wherein R^{6d} is a trifluoromethylsulfinyl group;

A compound of the formula (1D) wherein R^{6d} is a trifluoromethylsulfonyl group;

5 A compound of the formula (1D) wherein R^{6d} is a cyano group or a nitro group;

A compound of the formula (1D) wherein R^{6d} is a halogen atom;

10 A compound of the formula (1D) wherein R^{6d} is a chlorine atom, a bromine atom, or an iodine atom;

A compound of the formula (1D) wherein R^{6d} is a chlorine atom;

A compound of the formula (1D) wherein R^{6d} is a bromine atom;

15 A compound of the formula (1D) wherein R^{6d} is an iodine atom;

A compound of the formula (1D) wherein R^{6d} is $-SF_5$;

20 A compound of the formula (1D) wherein R^{6d} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or a halogen atom;

25 A compound of the formula (1D) wherein R^{6d} is a

trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom;

A compound of the formula (1D) wherein R^{7d} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (1D) wherein R^{7d} is a C1-C6 chain hydrocarbon group;

A compound of the formula (1D) wherein R^{7d} is a C1-C6 chain hydrocarbon group having one or more fluorine atoms;

A compound of the formula (1D) wherein R^{7d} is a C1-C3 alkyl group having one or more fluorine atoms;

A compound of the formula (1D) wherein R^{7d} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, or a heptafluoroisopropyl group;

A compound of the formula (1D) wherein R^{7d} is a trifluoromethyl group;

A compound of the formula (1D) wherein R^{7d} is a pentafluoroethyl group;

A compound of the formula (1D) wherein R^{7d} is a heptafluoropropyl group;

A compound of the formula (1D) wherein R^{7d} is a heptafluoroisopropyl group;

A compound of the formula (1D) wherein R^{7d} is a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, or a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms;

A compound of the formula (1D) wherein R^{7d} is $-OR^{11d}$;

10 A compound of the formula (1D) wherein R^{7d} is a C1-C6 alkoxy group having one or more fluorine atoms;

A compound of the formula (1D) wherein R^{7d} is a C1-C3 alkoxy group having one or more fluorine atoms;

15 A compound of the formula (1D) wherein R^{7d} is a trifluoromethoxy group;

A compound of the formula (1D) wherein R^{7d} is $-S(O)_mR^{11d}$;

20 A compound of the formula (1D) wherein R^{7d} is a C1-C6 alkylsulfanyl group having one or more fluorine atoms, a C1-C6 alkylsulfinyl group having one or more fluorine atoms, or a C1-C6 alkylsulfonyl group having one or more fluorine atoms;

25 A compound of the formula (1D) wherein R^{7d} is a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms,

or a C1-C3 alkylsulfonyl group having one or more fluorine atoms;

A compound of the formula (1D) wherein R^{7d} is a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

A compound of the formula (1D) wherein R^{7d} is a trifluoromethylsulfanyl group;

A compound of the formula (1D) wherein R^{7d} is a trifluoromethylsulfinyl group;

A compound of the formula (1D) wherein R^{7d} is a trifluoromethylsulfonyl group;

A compound of the formula (1D) wherein R^{7d} is a cyano group or a nitro group;

A compound of the formula (1D) wherein R^{7d} is a halogen atom;

A compound of the formula (1D) wherein R^{7d} is a chlorine atom, a bromine atom, or an iodine atom;

A compound of the formula (1D) wherein R^{7d} is a chlorine atom;

A compound of the formula (1D) wherein R^{7d} is a bromine atom;

A compound of the formula (1D) wherein R^{7d} is an iodine atom;

A compound of the formula (1D) wherein R^{7d} is a hydrogen atom;

A compound of the formula (1D) wherein R^{7d} is $-SF_5$;

A compound of the formula (1D) wherein R^{7d} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or a halogen atom;

A compound of the formula (1D) wherein R^{7d} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom;

A compound of the formula (1D) wherein R^{6d} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or a halogen atom, and R^{7d} is a hydrogen atom;

A compound of the formula (1D) wherein R^{6d} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a

trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom, and R^{7d} is a hydrogen atom;

A compound of the formula (1D) wherein m is 0;

5 A compound of the formula (1D) wherein m is 1;

A compound of the formula (1D) wherein m is 2;

A compound of the formula (1D) wherein A^{1d} is -NR^{8d}- or a sulfur atom, R^{8d} is a C1-C6 alkyl group, A^{2d} is =CR^{9d}-, A^{3d} is =CR^{10d}-, R^{9d} and R^{10d} are a hydrogen atom, R^{2d}, R^{4d}, and R^{5d}
10 are a hydrogen atom, R^{3d} is a C1-C3 alkyl group optionally having one or more fluorine atoms or a halogen atom, R^{6d} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a
15 C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or a halogen atom, and R^{7d} is a hydrogen atom;

A compound of the formula (1D) wherein A^{1d} is -NR^{8d}- or a sulfur atom, R^{8d} is a C1-C6 alkyl group, A^{2d} is =CR^{9d}-, R^{9d}
20 is a hydrogen atom, A^{3d} is a nitrogen atom, R^{2d}, R^{4d}, and R^{5d} are a hydrogen atom, R^{3d} is a C1-C3 alkyl group optionally having one or more fluorine atoms or a halogen atom, R^{6d} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-
25 C3 alkylsulfanyl group having one or more fluorine atoms, a

C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or a halogen atom, and R^{7d} is a hydrogen atom;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$,
5 R^{8d} is a methyl group, an ethyl group, a propyl group, or an isopropyl group, A^{2d} is $=CR^{9d}-$, A^{3d} is $=CR^{10d}-$, R^{9d} and R^{10d} are a hydrogen atom, R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, R^{3d} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a
10 pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom, R^{6d} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a
15 trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom, and R^{7d} is a hydrogen atom;

A compound of the formula (1D) wherein A^{1d} is a sulfur
20 atom, A^{2d} is $=CR^{9d}-$, A^{3d} is $=CR^{10d}-$, R^{9d} and R^{10d} are a hydrogen atom, R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, R^{3d} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a
pentafluoroethyl group, a heptafluoropropyl group, a
25 heptafluoroisopropyl group, a fluorine atom, a chlorine

atom, or a bromine atom, R^{6d} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom, and R^{7d} is a hydrogen atom;

A compound of the formula (1D) wherein A^{1d} is $-NR^{8d}-$, R^{8d} is a methyl group, an ethyl group, a propyl group, or an isopropyl group, A^{2d} is $=CR^{9d}-$, R^{9d} is a hydrogen atom, A^{3d} is a nitrogen atom, R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, R^{3d} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom, R^{6d} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom, and R^{7d} is a hydrogen atom;


A compound of the formula (1D) wherein A^{1d} is a sulfur atom, A^{2d} is $=CR^{9d}-$, R^{9d} is a hydrogen atom, A^{3d} is a nitrogen atom, R^{2d} , R^{4d} , and R^{5d} are a hydrogen atom, R^{3d} is

a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom, R^{6d} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a chlorine atom, a bromine atom, or an iodine atom, and R^{7d} is a hydrogen atom.

[0147]

Examples of Compound (21) include the following compounds:

A compound of the formula (1E):



wherein:

X^e represents a fluorine atom or a chlorine atom;

A^{1e} represents -NR^{8e}-, an oxygen atom or a sulfur atom;

A^{2e} represents a nitrogen atom or =CR^{9e}-;

A^{3e} represents a nitrogen atom or =CR^{10e}-;

R^{2e}, R^{3e}, and R^{5e} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally

having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more
5 halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, -OR^{11e}, -S(O)_mR^{11e}, -SF₅, a cyano group, a halogen atom, or a hydrogen atom: provided that at least one of R^{2e}, R^{3e}, and
10 R^{5e} represent a hydrogen atom;

R^{4e} represents a hydrogen atom;

R^{6e} represents a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one
15 or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, or -SF₅;

R^{7e} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, -OR^{11e}, -S(O)_mR^{11e}, -SF₅, a cyano group, a halogen atom, or a
20 hydrogen atom;

R^{8e} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group, a C3-C6 alicyclic hydrocarbon group optionally having one or more
25 halogen atoms, or a hydrogen atom;

R^{9e} and R^{10e} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11e}$, $-S(O)_mR^{11e}$, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

5 R^{11e} and R^{12e} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a hydrogen atom; and

m represents 0, 1, or 2:

provided that; in $-S(O)_mR^{11e}$, when m is 1 or 2, R^{11e} is
10 not a hydrogen atom; when A^{1e} is an oxygen atom, and A^{2e} and A^{3e} are a methine group, one of R^{2e} , R^{3e} , and R^{5e} are a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one
15 or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one
20 or more halogen atoms, $-OR^{11e}$, $-S(O)_mR^{11e}$, a cyano group, a nitro group, or a halogen atom, and two others are a hydrogen atom; and when A^{1e} is $-NR^{8e}-$, and A^{2e} is a methine group, R^{8e} is a methyl group (hereinafter, referred to as Compound (1E));

25 [0148]

Examples of Compound (1E) include the following compounds:

A compound of the formula (1E) wherein X^e is a fluorine atom;

5 A compound of the formula (1E) wherein X^e is a chlorine atom;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
and R^{8e} is a C1-C6 chain hydrocarbon group optionally
10 having one or more halogen atoms;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
and R^{8e} is a C1-C6 chain hydrocarbon group;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
and R^{8e} is a C1-C6 alkyl group;

15 A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
and R^{8e} is a C1-C3 alkyl group;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
and R^{8e} is a methyl group, an ethyl group, a propyl group,
or an isopropyl group;

20 A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
and R^{8e} is a methyl group;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
and R^{8e} is an ethyl group;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
25 and R^{8e} is a propyl group;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
and R^{8e} is an isopropyl group;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
and R^{8e} is a C1-C6 chain hydrocarbon group having one C1-C6
5 alkoxy group;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
and R^{8e} is a methoxymethyl group or an ethoxymethyl group;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
and R^{8e} is a C3-C6 alicyclic hydrocarbon group optionally
10 having one or more halogen atoms;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
and R^{8e} is a cyclopropyl group, a cyclobutyl group, a
cyclopentyl group, or a cyclohexyl group;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$,
15 and R^{8e} is a hydrogen atom;

A compound of the formula (1E) wherein A^{1e} is an
oxygen atom;

A compound of the formula (1E) wherein A^{1e} is a sulfur
atom;

20 A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$ or
a sulfur atom;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$ or
a sulfur atom, and R^{8e} is a methyl group, an ethyl group, a
propyl group, or an isopropyl group;

25 A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$ or

a sulfur atom, and R^{8e} is a methoxymethyl group, an ethoxymethyl group, or a hydrogen atom;

A compound of the formula (1E) wherein A^{2e} is $=CR^{9e}-$, and A^{3e} is $=CR^{10e}-$;

5 A compound of the formula (1E) wherein A^{2e} is $=CR^{9e}-$, A^{3e} is $=CR^{10e}-$, and R^{9e} and R^{10e} are same or different and are independently a halogen atom or a hydrogen atom;

A compound of the formula (1E) wherein A^{2e} is $=CR^{9e}-$, A^{3e} is $=CR^{10e}-$, and R^{9e} and R^{10e} are a hydrogen atom;

10 A compound of the formula (1E) wherein A^{2e} is a nitrogen atom, and A^{3e} is $=CR^{10e}-$;

A compound of the formula (1E) wherein A^{2e} is $=CR^{9e}-$, and A^{3e} is a nitrogen atom;

A compound of the formula (1E) wherein A^{2e} is $=CR^{9e}-$, R^{9e} is a halogen atom or a hydrogen atom, and A^{3e} is a nitrogen atom;

A compound of the formula (1E) wherein A^{2e} is $=CR^{9e}-$, R^{9e} is a hydrogen atom, and A^{3e} is a nitrogen atom;

A compound of the formula (1E) wherein A^{2e} is a nitrogen atom, and A^{3e} is a nitrogen atom;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$, A^{2e} is $=CR^{9e}-$, and A^{3e} is $=CR^{10e}-$;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$, A^{2e} is $=CR^{9e}-$, A^{3e} is $=CR^{10e}-$, and R^{9e} and R^{10e} are a hydrogen

25 atom;

A compound of the formula (1E) wherein A^{1e} is an oxygen atom, A^{2e} is $=CR^{9e}-$, and A^{3e} is $=CR^{10e}-$;

A compound of the formula (1E) wherein A^{1e} is a sulfur atom, A^{2e} is $=CR^{9e}-$, and A^{3e} is $=CR^{10e}-$;

5 A compound of the formula (1E) wherein A^{1e} is a sulfur atom, A^{2e} is $=CR^{9e}-$, A^{3e} is $=CR^{10e}-$, and R^{9e} and R^{10e} are a hydrogen atom;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$, A^{2e} is $=CR^{9e}-$, and A^{3e} is a nitrogen atom;

10 A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$, A^{2e} is $=CR^{9e}-$, R^{9e} is a hydrogen atom, and A^{3e} is a nitrogen atom;

A compound of the formula (1E) wherein A^{1e} is an oxygen atom, A^{2e} is $=CR^{9e}-$, and A^{3e} is a nitrogen atom;

15 A compound of the formula (1E) wherein A^{1e} is a sulfur atom, A^{2e} is $=CR^{9e}-$, and A^{3e} is a nitrogen atom;

A compound of the formula (1E) wherein A^{1e} is a sulfur atom, A^{2e} is $=CR^{9e}-$, R^{9e} is a hydrogen atom, and A^{3e} is a nitrogen atom;

20 A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$ or a sulfur atom, R^{8e} is a methyl group, an ethyl group, a propyl group, or an isopropyl group, A^{2e} is $=CR^{9e}-$, A^{3e} is $=CR^{10e}-$, and R^{9e} and R^{10e} are a hydrogen atom;

25 A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$ or a sulfur atom, R^{8e} is a methyl group, A^{2e} is $=CR^{9e}-$, A^{3e} is

=CR^{10e}-, and R^{9e} and R^{10e} are a hydrogen atom;

A compound of the formula (1E) wherein A^{1e} is -NR^{8e}- or a sulfur atom, R^{8e} is a methyl group, an ethyl group, a propyl group, or an isopropyl group, A^{2e} is =CR^{9e}-, R^{9e} is a
5 hydrogen atom, and A^{3e} is a nitrogen atom;

A compound of the formula (1E) wherein A^{1e} is -NR^{8e}- or a sulfur atom, R^{8e} is a methyl group, A^{2e} is =CR^{9e}-, R^{9e} is a hydrogen atom, and A^{3e} is a nitrogen atom;

A compound of the formula (1E) wherein R^{2e}, R^{3e}, and R^{5e}
10 are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a hydrogen atom;

A compound of the formula (1E) wherein R^{2e}, R^{3e}, and R^{5e}
are same or different and are independently a phenyl group
15 optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 5-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, a 6-membered heterocyclic group
20 optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms, or a hydrogen atom;

A compound of the formula (1E) wherein R^{2e}, R^{3e}, and R^{5e}
are same or different and are independently -OR^{11e} or a
25 hydrogen atom;

A compound of the formula (1E) wherein R^{2e} , R^{3e} , and R^{5e} are same or different and are independently $-S(O)_mR^{11e}$ or a hydrogen atom;

5 A compound of the formula (1E) wherein R^{2e} , R^{3e} , and R^{5e} are same or different and are independently a cyano group, a nitro group, or a hydrogen atom;

A compound of the formula (1E) wherein R^{2e} , R^{3e} , and R^{5e} are same or different and are independently a halogen atom or a hydrogen atom;

10 A compound of the formula (1E) wherein R^{2e} , R^{3e} , and R^{5e} are a hydrogen atom;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, R^{3e} is a 2-pyridyl group, a 2-pyrimidyl group, a 3-chloro-2-pyridyl group, or a 3-chloro-5-trifluoromethyl-2-pyridyl group;

15

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a 2-pyridyl group;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a 2-pyrimidyl group;

20 A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a 3-chloro-2-pyridyl group;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a 3-chloro-5-trifluoromethyl-2-pyridyl group;

25 A compound of the formula (1E) wherein R^{2e} and R^{5e} are

a hydrogen atom, and R^{3e} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a C1-C6 alkyl group;

5 A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a C1-C6 alkyl group optionally having one or more fluorine atoms;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a C1-C3 alkyl group;

10 A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a C1-C3 alkyl group optionally having one or more fluorine atoms;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a methyl group, an ethyl group,
15 a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, or a heptafluoroisopropyl group;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a methyl group;

20 A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is an ethyl group;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a propyl group;

25 A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is an isopropyl group;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a trifluoromethyl group;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a pentafluoroethyl group;

5 A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a heptafluoropropyl group;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a heptafluoroisopropyl group;

10 A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a halogen atom;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a fluorine atom, a chlorine atom, or a bromine atom;

15 A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a fluorine atom;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a chlorine atom;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a bromine atom;

20 A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is $-SF_5$;

25 A compound of the formula (1E) wherein R^{2e} , R^{3e} , and R^{5e} are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a halogen atom, or a hydrogen atom;

A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a 6-membered heterocyclic group optionally having one or more halogen atoms or C1-C3 alkyl groups optionally having one or more halogen atoms;

5 A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a halogen atom;

10 A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a C1-C3 alkyl group optionally having one or more fluorine atoms, or a halogen atom;

15 A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a 2-pyridyl group, a 2-pyrimidyl group, a 3-chloro-2-pyridyl group, a 3-chloro-5-trifluoromethyl-2-pyridyl group, a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom;

20 A compound of the formula (1E) wherein R^{2e} and R^{5e} are a hydrogen atom, and R^{3e} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine
25 atom, or a bromine atom;

A compound of the formula (1E) wherein R^{6e} is a C1-C3 alkyl group having one or more fluorine atoms;

A compound of the formula (1E) wherein R^{6e} is a trifluoromethyl group, a pentafluoroethyl group, a
5 heptafluoropropyl group, or a heptafluoroisopropyl group;

A compound of the formula (1E) wherein R^{6e} is a trifluoromethyl group;

A compound of the formula (1E) wherein R^{6e} is a pentafluoroethyl group;

10 A compound of the formula (1E) wherein R^{6e} is a heptafluoropropyl group;

A compound of the formula (1E) wherein R^{6e} is a heptafluoroisopropyl group;

15 A compound of the formula (1E) wherein R^{6e} is a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms;

20 A compound of the formula (1E) wherein R^{6e} is a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

A compound of the formula (1E) wherein R^{6e} is a trifluoromethylsulfanyl group;

25 A compound of the formula (1E) wherein R^{6e} is a trifluoromethylsulfinyl group;

A compound of the formula (1E) wherein R^{6e} is a trifluoromethylsulfonyl group;

A compound of the formula (1E) wherein R^{6e} is -SF₅;

A compound of the formula (1E) wherein R^{6e} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms;

10 A compound of the formula (1E) wherein R^{6e} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

15 A compound of the formula (1E) wherein R^{7e} is a C1-C3 alkyl group having one or more fluorine atoms;

A compound of the formula (1E) wherein R^{7e} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, or a heptafluoroisopropyl group;

20 A compound of the formula (1E) wherein R^{7e} is a trifluoromethyl group;

A compound of the formula (1E) wherein R^{7e} is a pentafluoroethyl group;

25 A compound of the formula (1E) wherein R^{7e} is a heptafluoropropyl group;

A compound of the formula (1E) wherein R^{7e} is a heptafluoroisopropyl group;

A compound of the formula (1E) wherein R^{7e} is a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a
5 C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms;

A compound of the formula (1E) wherein R^{7e} is a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl
10 group, or a trifluoromethylsulfonyl group;

A compound of the formula (1E) wherein R^{7e} is a trifluoromethylsulfanyl group;

A compound of the formula (1E) wherein R^{7e} is a trifluoromethylsulfinyl group;

15 A compound of the formula (1E) wherein R^{7e} is a trifluoromethylsulfonyl group;

A compound of the formula (1E) wherein R^{7e} is -SF₅;

A compound of the formula (1E) wherein R^{7e} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3
20 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms;

A compound of the formula (1E) wherein R^{7e} is a
25 trifluoromethyl group, a pentafluoroethyl group, a

heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

A compound of the formula (1E) wherein R^{6e} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms, and R^{7e} is a hydrogen atom;

A compound of the formula (1E) wherein R^{6e} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, and R^{7e} is a hydrogen atom;

A compound of the formula (1E) wherein m is 0;

A compound of the formula (1E) wherein m is 1;

A compound of the formula (1E) wherein m is 2;

A compound of the formula (1E) wherein A^{1e} is -NR^{8e}- or a sulfur atom, R^{8e} is a C1-C6 alkyl group, A^{2e} is =CR^{9e}-, A^{3e} is =CR^{10e}-, R^{9e} and R^{10e} are a hydrogen atom, R^{2e} and R^{5e} are a hydrogen atom, R^{3e} is a C1-C3 alkyl group optionally having one or more fluorine atoms or a halogen atom, R^{6e} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3

alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms, and R^{7e} is a hydrogen atom;

5 A compound of the formula (1E) wherein A^{1e} is -NR^{8e}- or a sulfur atom, R^{8e} is a C1-C6 alkyl group, A^{2e} is =CR^{9e}-, R^{9e} is a hydrogen atom, A^{3e} is a nitrogen atom, R^{2e} and R^{5e} are a hydrogen atom, R^{3e} is a C1-C3 alkyl group optionally having one or more fluorine atoms or a halogen atom, R^{6e} is a C1-
10 C3 alkyl group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms, and R^{7e} is a hydrogen atom;

15 A compound of the formula (1E) wherein A^{1e} is -NR^{8e}-, R^{8e} is a methyl group, an ethyl group, a propyl group, or an isopropyl group, A^{2e} is =CR^{9e}-, A^{3e} is =CR^{10e}-, R^{9e} and R^{10e} are a hydrogen atom, R^{2e} and R^{5e} are a hydrogen atom, R^{3e} is a methyl group, an ethyl group, a propyl group, an
20 isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom, R^{6e} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a
25 heptafluoroisopropyl group, a trifluoromethylsulfanyl group,

a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, and R^{7e} is a hydrogen atom;

A compound of the formula (1E) wherein A^{1e} is a sulfur atom, A^{2e} is $=CR^{9e}-$, A^{3e} is $=CR^{10e}-$, R^{9e} and R^{10e} are a hydrogen atom, R^{2e} and R^{5e} are a hydrogen atom, R^{3e} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom, R^{6e} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethylsulfonyl group, or a trifluoromethylsulfinyl group, and R^{7e} is a hydrogen atom;

A compound of the formula (1E) wherein A^{1e} is $-NR^{8e}-$, R^{8e} is a methyl group, an ethyl group, a propyl group, or an isopropyl group, A^{2e} is $=CR^{9e}-$, R^{9e} is a hydrogen atom, A^{3e} is a nitrogen atom, R^{2e} and R^{5e} are a hydrogen atom, R^{3e} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom, R^{6e} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethylsulfonyl group,

a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, and R^{7e} is a hydrogen atom;


A compound of the formula (1E) wherein A^{1e} is a sulfur atom, A^{2e} is $=CR^{9e}-$, R^{9e} is a hydrogen atom, A^{3e} is a nitrogen atom, R^{2e} , R^{4e} and R^{5e} are a hydrogen atom, R^{3e} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a fluorine atom, a chlorine atom, or a bromine atom, R^{6e} is a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethylsulfonyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, and R^{7e} is a hydrogen atom.

[0149]

Examples of Compound (F) include the following compounds:

[0150]


A compound of the formula (M21):



wherein, R^6 , R^7 , R^8 and A^2 are as defined above (referred to as Compound (M21));

[0151]

A compound of the formula (1F):



wherein:

A^{2f} represents a nitrogen atom or $=CR^{9f}-$;

A^{3f} represents a nitrogen atom or $=CR^{10f}-$;

5 R^{6f} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11f}$, $-S(O)_mR^{11f}$, $-SF_5$, or a halogen atom;

R^{7f} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11f}$, $-S(O)_mR^{11f}$, $-SF_5$, hydrogen atom, or a halogen atom;

10

R^{8f} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a hydrogen atom;

R^{9f} and R^{10f} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a halogen atom, or a hydrogen atom;

15

R^{11f} and R^{12f} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a hydrogen atom;

20

m represents 0, 1, or 2:

provided that in $-S(O)_mR^{11f}$, when m is 1 or 2, R^{11f} is not a hydrogen atom (hereinafter, referred to as Compound

(1F));

[0152]

And examples of Compound (1F) include the following compounds:

5 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$,
and A^{3f} is $=CR^{10f}-$;

 A compound of the formula (1F) wherein A^{2f} is a
nitrogen atom, and A^{3f} is $=CR^{10f}-$;

10 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$,
and A^{3f} is a nitrogen atom;

 A compound of the formula (1F) wherein A^{2f} is a
nitrogen atom, and A^{3f} is a nitrogen atom;

 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$,
and A^{3f} is $=CR^{10f}-$, R^{9f} and R^{10f} are a hydrogen atom;

15 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$,
 R^{9f} is a hydrogen atom, and A^{3f} is a nitrogen atom;

 A compound of the formula (1F) wherein R^{6f} is a C2-C3
alkyl group having one or more fluorine atoms, a C1-C3
alkylsulfanyl group having one or more fluorine atoms, a
20 C1-C3 alkylsulfinyl group having one or more fluorine atoms,
or a C1-C3 alkylsulfonyl group having one or more fluorine
atoms;

 A compound of the formula (1F) wherein R^{6f} is a C2-C3
alkyl group having one or more fluorine atoms;

25 A compound of the formula (1F) wherein R^{6f} is a C1-C3

alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms;

5 A compound of the formula (1F) wherein R^{6f} is a pentafluoroethyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, or an iodine atom;

10 A compound of the formula (1F) wherein R^{6f} is a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

 A compound of the formula (1F) wherein R^{6f} is a pentafluoroethyl group;

15 A compound of the formula (1F) wherein R^{6f} is a trifluoromethylsulfanyl group;

 A compound of the formula (1F) wherein R^{6f} is a trifluoromethylsulfinyl group;

 A compound of the formula (1F) wherein R^{6f} is a trifluoromethylsulfonyl group;

20 A compound of the formula (1F) wherein R^{6f} is an iodine atom;

 A compound of the formula (1F) wherein R^{6f} is $-SF_5$;

 A compound of the formula (1F) wherein R^{7f} is a hydrogen atom;

25 A compound of the formula (1F) wherein R^{8f} is a C1-C3

alkyl group or a hydrogen atom;

A compound of the formula (1F) wherein R^{8f} is a C1-C3 alkyl group;

5 A compound of the formula (1F) wherein R^{8f} is a methyl group;

A compound of the formula (1F) wherein R^{8f} is an ethyl group;

A compound of the formula (1F) wherein R^{8f} is a propyl group;

10 A compound of the formula (1F) wherein R^{8f} is a hydrogen atom;

A compound of the formula (1F) wherein R^{8f} is a methyl group or a hydrogen atom;

15 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, R^{9f} is a hydrogen atom, A^{3f} is a nitrogen atom, R^{6f} is a C2-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine
20 atoms, R^{7f} is a hydrogen atom, and R^{8f} is a C1-C3 alkyl group or a hydrogen atom;

25 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, R^{9f} is a hydrogen atom, A^{3f} is a nitrogen atom, R^{6f} is a C2-C3 alkyl group having one or more fluorine atoms, R^{7f} is a hydrogen atom, and R^{8f} is a C1-C3 alkyl group or a hydrogen

atom;

A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, R^{9f} is a hydrogen atom, A^{3f} is a nitrogen atom, R^{6f} is a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms, R^{7f} is a hydrogen atom, and R^{8f} is a C1-C3 alkyl group or a hydrogen atom;

A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, R^{9f} is a hydrogen atom, A^{3f} is a nitrogen atom, R^{6f} is a pentafluoroethyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group or an iodine atom, R^{7f} is a hydrogen atom, and R^{8f} is a C1-C3 alkyl group or a hydrogen atom: provided that when R^{6f} is an iodine atom, R^{8f} is a C1-C3 alkyl group;

A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, R^{9f} is a hydrogen atom, A^{3f} is a nitrogen atom, R^{6f} is a pentafluoroethyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, R^{7f} is a hydrogen atom, and R^{8f} is a methyl group or a hydrogen atom;

A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, R^{9f} is a hydrogen atom, A^{3f} is a nitrogen atom, R^{6f} is a pentafluoroethyl group, R^{7f} is a hydrogen atom, and R^{8f} is a methyl group;

A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, R^{9f} is a hydrogen atom, A^{3f} is a nitrogen atom, R^{6f} is a trifluoromethylsulfanyl group, R^{7f} is a hydrogen atom, and R^{8f} is a methyl group;

5 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, R^{9f} is a hydrogen atom, A^{3f} is a nitrogen atom, R^{6f} is a trifluoromethylsulfonyl group, R^{7f} is a hydrogen atom, and R^{8f} is a methyl group;

10 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, R^{9f} is a hydrogen atom, A^{3f} is a nitrogen atom, R^{6f} is an iodine atom, R^{7f} is a hydrogen atom, and R^{8f} is a C1-C3 alkyl group;

15 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, R^{9f} is a hydrogen atom, A^{3f} is a nitrogen atom, R^{6f} is an iodine atom, R^{7f} is a hydrogen atom, and R^{8f} is a methyl group;

20 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, A^{3f} is $=CR^{10f}-$, R^{9f} and R^{10f} are a hydrogen atom, R^{6f} is a C2-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms, R^{7f} is a hydrogen atom, and R^{8f} is a C1-C3 alkyl group or a hydrogen atom;

25 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$,

A^{3f} is $=CR^{10f}-$, R^{9f} and R^{10f} are a hydrogen atom, R^{6f} is a C2-C3 alkyl group having one or more fluorine atoms, R^{7f} is a hydrogen atom, and R^{8f} is a C1-C3 alkyl group or a hydrogen atom;

5 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, A^{3f} is $=CR^{10f}-$, R^{9f} and R^{10f} are a hydrogen atom, R^{6f} is a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine
10 atoms, R^{7f} is a hydrogen atom, and R^{8f} is a C1-C3 alkyl group or a hydrogen atom;

 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, A^{3f} is $=CR^{10f}-$, R^{9f} and R^{10f} are a hydrogen atom, R^{6f} is a pentafluoroethyl group, a trifluoromethylsulfanyl group, a
15 trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, R^{7f} is a hydrogen atom, and R^{8f} is a methyl group or a hydrogen atom;

 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, A^{3f} is $=CR^{10f}-$, R^{9f} and R^{10f} are a hydrogen atom, R^{6f} is a
20 pentafluoroethyl group, R^{7f} is a hydrogen atom, and R^{8f} is a methyl group;

 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, A^{3f} is $=CR^{10f}-$, R^{9f} and R^{10f} are a hydrogen atom, R^{6f} is a trifluoromethylsulfanyl group, R^{7f} is a hydrogen atom, and
25 R^{8f} is a methyl group;

A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, A^{3f} is $=CR^{10f}-$, R^{9f} and R^{10f} are a hydrogen atom, R^{6f} is a trifluoromethylsulfonyl group, R^{7f} is a hydrogen atom, and R^{8f} is a methyl group;

5 A compound of the formula (1F) wherein A^{2f} is $=CR^{9f}-$, A^{3f} is $=CR^{10f}-$, R^{9f} and R^{10f} are a hydrogen atom, R^{6f} is $-SF_5$, R^{7f} is a hydrogen atom, and R^{8f} is a methyl group;

[0153]

10 And examples of Compound (M21) include the following compounds:

A compound of the formula (M21) wherein A^2 is $=CH-$;

A compound of the formula (M21) wherein A^2 is nitrogen;

15 A compound of the formula (M21) wherein R^8 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or hydrogen;

A compound of the formula (M21) wherein R^8 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

20 A compound of the formula (M21) wherein R^8 is a C1-C3 alkyl group;

A compound of the formula (M21) wherein R^8 is a methyl group, an ethyl group, a propyl group, or hydrogen;

25 A compound of the formula (M21) wherein R^8 is a methyl group;

A compound of the formula (M21) wherein R⁸ is an ethyl group;

A compound of the formula (M21) wherein R⁸ is a propyl group;

5 A compound of the formula (M21) wherein R⁸ is hydrogen;

A compound of the formula (M21) wherein R⁶ is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 alkylsulfanyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfinyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfonyl group optionally having one or more halogen atoms, a bromine atom, or an iodine atom;

15 A compound of the formula (M21) wherein R⁶ is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 alkylsulfanyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfinyl group optionally having one or more halogen atoms, or a C1-C6 alkylsulfonyl group optionally having one or more halogen atoms;

25 A compound of the formula (M21) wherein R⁶ is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (M21) wherein R⁶ is a C1-C6 alkoxy group optionally having one or more halogen atoms;

A compound of the formula (M21) wherein R⁶ is a C1-C6 alkylsulfanyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfinyl group optionally having one or more halogen atoms, or a C1-C6 alkylsulfonyl group optionally having one or more halogen atoms;

A compound of the formula (M21) wherein R⁶ is a bromine atom or an iodine atom;

A compound of the formula (M21) wherein R⁶ is a trifluoromethyl group, a pentafluoroethyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

A compound of the formula (M21) wherein R⁶ is a pentafluoroethyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

A compound of the formula (M21) wherein R⁶ is a pentafluoroethyl group;

A compound of the formula (M21) wherein R⁶ is a trifluoromethylsulfanyl group;

A compound of the formula (M21) wherein R⁶ is a trifluoromethylsulfinyl group;

A compound of the formula (M21) wherein R⁶ is a

trifluoromethylsulfonyl group;

A compound of the formula (M21) wherein R^6 is a trifluoromethyl group, a pentafluoroethyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, and R^7 is hydrogen;

A compound of the formula (M21) wherein R^6 is a pentafluoroethyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, and R^7 is hydrogen;

A compound of the formula (M21) wherein R^6 is a pentafluoroethyl group, and R^7 is hydrogen;

A compound of the formula (M21) wherein R^6 is a trifluoromethylsulfanyl group, and R^7 is hydrogen;

A compound of the formula (M21) wherein R^6 is a trifluoromethylsulfinyl group, and R^7 is hydrogen;

A compound of the formula (M21) wherein R^6 is a trifluoromethylsulfonyl group, and R^7 is hydrogen;

A compound of the formula (M21) wherein A^2 is $=CH-$, R^6 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, R^7 is hydrogen, and R^8 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (M21) wherein A^2 is $=CH-$, R^6

is a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a trifluoromethoxy group, or a trifluoromethylsulfanyl group, R⁷ is hydrogen, and R⁸ is a methyl group, an ethyl group, a propyl group, or an isopropyl group;

A compound of the formula (M21) wherein A² is =CH-, R⁶ is a trifluoromethyl group or pentafluoroethyl group, R⁷ is hydrogen, and R⁸ is a methyl group;

A compound of the formula (M21) wherein A² is =CH-, R⁶ is a trifluoromethyl group, a pentafluoroethyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, R⁷ is hydrogen, and R⁸ is a methyl group;

A compound of the formula (M21) wherein A² is =CH-, R⁶ is a pentafluoroethyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, R⁷ is hydrogen, and R⁸ is a methyl group;


A compound of the formula (M21) wherein A² is =CH-, R⁶ is a pentafluoroethyl group, R⁷ is hydrogen, and R⁸ is a methyl group;

A compound of the formula (M21) wherein A² is =CH-, R⁶ is a trifluoromethylsulfonyl group, R⁷ is hydrogen, and R⁸ is a methyl group.

Examples of Compound (G) include the following compounds:

[0155]


A compound of the formula (M20):



wherein, X^g , R^6 , R^7 , R^8 and A^2 are as defined above (referred to as Compound (M20));

[0156]

A compound of the formula (1G):



wherein:

X^g represents a chlorine atom, a bromine atom, or an iodine atom;

A^{2g} represents a nitrogen atom or $=CR^{9g}-$;

15 A^{3g} represents a nitrogen atom or $=CR^{10g}-$;

R^{6g} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11g}$, $-S(O)_mR^{11g}$, $-SF_5$, or a halogen atom;

20 R^{7g} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11g}$, $-S(O)_mR^{11g}$, $-SF_5$, or a halogen atom;

R^{8g} represents a C1-C6 chain hydrocarbon group

optionally having one or more halogen atoms or a hydrogen atom;

R^{9g} and R^{10g} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a halogen atom or a hydrogen atom;

R^{11g} and R^{12g} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a hydrogen atom;

m represents 0, 1, or 2:

provided that in $-S(O)_mR^{11g}$, when m is 1 or 2, R^{11g} is not a hydrogen atom (hereinafter, referred to as Compound (1G));

[0157]

And examples of Compound (1G) include the following compounds:

A compound of the formula (1G) wherein X^g is a chlorine atom;

A compound of the formula (1G) wherein X^g is a bromine atom;

A compound of the formula (1G) wherein X^g is an iodine atom;

A compound of the formula (1G) wherein A^{2g} is $=CR^{9g}-$, and A^{3g} is $=CR^{10g}-$;

A compound of the formula (1G) wherein A^{2g} is a

nitrogen atom, and A^{3g} is =CR^{10g}-;

A compound of the formula (1G) wherein A^{2g} is =CR^{9g}-, and A^{3g} is a nitrogen atom;

5 A compound of the formula (1G) wherein A^{2g} is a nitrogen atom, and A^{3g} is a nitrogen atom;

A compound of the formula (1G) wherein A^{2g} is =CR^{9g}-, A^{3g} is =CR^{10g}-, and R^{9g} and R^{10g} are a hydrogen atom;

A compound of the formula (1G) wherein A^{2g} is =CR^{9g}-, R^{9g} is a hydrogen atom, and A^{3g} is a nitrogen atom;

10 A compound of the formula (1G) wherein R^{6g} is a C2-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine
15 atoms;

A compound of the formula (1G) wherein R^{6g} is a C2-C3 alkyl group having one or more fluorine atoms;

A compound of the formula (1G) wherein R^{6g} is a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a
20 C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms;

A compound of the formula (1G) wherein R^{6g} is a pentafluoroethyl group, a trifluoromethylsulfanyl group, a
25 trifluoromethylsulfinyl group, a trifluoromethylsulfonyl

group, or an iodine atom;

A compound of the formula (1G) wherein R^{6g} is a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

5 A compound of the formula (1G) wherein R^{6g} is a pentafluoroethyl group;

A compound of the formula (1G) wherein R^{6g} is a trifluoromethylsulfanyl group;

10 A compound of the formula (1G) wherein R^{6g} is a trifluoromethylsulfinyl group;

A compound of the formula (1G) wherein R^{6g} is a trifluoromethylsulfonyl group;

A compound of the formula (1G) wherein R^{6g} is an iodine atom;

15 A compound of the formula (1G) wherein R^{6g} is $-SF_5$;

A compound of the formula (1G) wherein R^{7g} is a hydrogen atom;

A compound of the formula (1G) wherein R^{8g} is a C1-C3 alkyl group or a hydrogen atom;

20 A compound of the formula (1G) wherein R^{8g} is a C1-C3 alkyl group;

A compound of the formula (1G) wherein R^{8g} is a methyl group;

25 A compound of the formula (1G) wherein R^{8g} is an ethyl group;

A compound of the formula (1G) wherein R^{8g} is a propyl group;

A compound of the formula (1G) wherein R^{8g} is a hydrogen atom;

5 A compound of the formula (1G) wherein R^{8g} is a methyl group or a hydrogen atom;

A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, R^{9g} is a hydrogen atom, A^{3g} is a nitrogen atom, R^{6g} is a C2-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms, R^{7g} is a hydrogen atom, and R^{8g} is a C1-C3 alkyl group or a hydrogen atom;

10

15

A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, R^{9g} is a hydrogen atom, A^{3g} is a nitrogen atom, R^{6g} is a C2-C3 alkyl group having one or more fluorine atoms, R^{7g} is a hydrogen atom, and R^{8g} is a C1-C3 alkyl group or a hydrogen atom;

20

A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, R^{9g} is a hydrogen atom, A^{3g} is a nitrogen atom, R^{6g} is a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms

25

or a C1-C3 alkylsulfonyl group having one or more fluorine atoms, R^{7g} is a hydrogen atom, and R^{8g} is a C1-C3 alkyl group or a hydrogen atom;

A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, R^{9g} is a hydrogen atom, A^{3g} is a nitrogen atom, R^{6g} is a pentafluoroethyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, or an iodine atom, R^{7g} is a hydrogen atom, and R^{8g} is a C1-C3 alkyl group or a hydrogen atom: provided that when R^{6g} is an iodine atom, R^{8g} is a C1-C3 alkyl group;

A compound of the formula (1G) wherein A^{2g} is $=CR^{9g}-$, R^{9g} is a hydrogen atom, A^{3g} is a nitrogen atom, R^{6g} is a pentafluoroethyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, R^{7g} is a hydrogen atom, and R^{8g} is a methyl group or a hydrogen atom;

A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, R^{9g} is a hydrogen atom, A^{3g} is a nitrogen atom, R^{6g} is a pentafluoroethyl group, R^{7g} is a hydrogen atom, and R^{8g} is a methyl group;

A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, R^{9g} is a hydrogen atom, A^{3g} is a nitrogen atom, R^{6g} is a

trifluoromethylsulfanyl group, R^{7g} is a hydrogen atom, and R^{8g} is a methyl group;

A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, R^{9g} is a hydrogen atom, A^{3g} is a nitrogen atom, R^{6g} is a trifluoromethylsulfonyl group, R^{7g} is a hydrogen atom, and R^{8g} is a methyl group;

A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, R^{9g} is a hydrogen atom, A^{3g} is a nitrogen atom, R^{6g} is an iodine atom, R^{7g} is a hydrogen atom, and R^{8g} is a C1-C3 alkyl group;

A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, R^{9g} is a hydrogen atom, A^{3g} is a nitrogen atom, R^{6g} is an iodine atom, R^{7g} is a hydrogen atom, and R^{8g} is a methyl group;

A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, A^{3g} is $=CR^{10g}-$, R^{9g} and R^{10g} are a hydrogen atom, R^{6g} is a C2-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, a C1-C3 alkylsulfonyl group having one or more fluorine atoms, R^{7g} is a hydrogen atom, and R^{8g} is a C1-C3 alkyl group or a hydrogen atom;

A compound of the formula (1G) wherein X^g is a

chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, A^{3g} is $=CR^{10g}-$, R^{9g} and R^{10g} are a hydrogen atom, R^{6g} is a C2-C3 alkyl group having one or more fluorine atoms, R^{7g} is a hydrogen atom, and R^{8g} is a C1-C3 alkyl group or a hydrogen atom;

5 A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, A^{3g} is $=CR^{10g}-$, R^{9g} and R^{10g} are a hydrogen atom, R^{6g} is a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms,
10 or a C1-C3 alkylsulfonyl group having one or more fluorine atoms, R^{7g} is a hydrogen atom, and R^{8g} is a C1-C3 alkyl group or a hydrogen atom;

 A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, A^{3g} is $=CR^{10g}-$,
15 R^{9g} and R^{10g} are a hydrogen atom, R^{6g} is a pentafluoroethyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, R^{7g} is a hydrogen atom, and R^{8g} is a methyl group or a hydrogen atom;

20 A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, A^{3g} is $=CR^{10g}-$, R^{9g} and R^{10g} are a hydrogen atom, R^{6g} is a pentafluoroethyl group, R^{7g} is a hydrogen atom, and R^{8g} is a methyl group;

 A compound of the formula (1G) wherein X^g is a
25 chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, A^{3g} is $=CR^{10g}-$,

R^{9g} and R^{10g} are a hydrogen atom, R^{6g} is a trifluoromethylsulfanyl group, R^{7g} is a hydrogen atom, and R^{8g} is a methyl group;

5 A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, A^{3g} is $=CR^{10g}-$, R^{9g} and R^{10g} are a hydrogen atom, R^{6g} is a trifluoromethylsulfonyl group, R^{7g} is a hydrogen atom, and R^{8g} is a methyl group;

10 A compound of the formula (1G) wherein X^g is a chlorine atom or a bromine atom, A^{2g} is $=CR^{9g}-$, A^{3g} is $=CR^{10g}-$, R^{9g} and R^{10g} are a hydrogen atom, R^{6g} is $-SF_5$, R^{7g} is a hydrogen atom, and R^{8g} is a methyl group;
[0158]

15 And examples of Compound (M20) include the following compounds:

A compound of the formula (M20) wherein X^g is a chlorine atom;

A compound of the formula (M20) wherein X^g is a bromine atom;

20 A compound of the formula (M20) wherein X^g is an iodine atom;

A compound of the formula (M20) wherein A^2 is $=CH-$;

A compound of the formula (M20) wherein A^2 is nitrogen;

25 A compound of the formula (M20) wherein R^8 is a C1-C6

chain hydrocarbon group optionally having one or more halogen atoms or hydrogen;

A compound of the formula (M20) wherein R^8 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (M20) wherein R^8 is a C1-C3 alkyl group;

A compound of the formula (M20) wherein R^8 is a methyl group, an ethyl group, a propyl group, or hydrogen;

A compound of the formula (M20) wherein R^8 is a methyl group;

A compound of the formula (M20) wherein R^8 is an ethyl group;

A compound of the formula (M20) wherein R^8 is a propyl group;

A compound of the formula (M20) wherein R^8 is hydrogen;

A compound of the formula (M20) wherein R^6 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 alkylsulfanyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfinyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfonyl group optionally having one or more halogen atoms, a bromine atom, or an iodine atom;

A compound of the formula (M20) wherein R⁶ is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 alkylsulfanyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfinyl group optionally having one or more halogen atoms, or a C1-C6 alkylsulfonyl group optionally having one or more halogen atoms;

A compound of the formula (M20) wherein R⁶ is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;

A compound of the formula (M20) wherein R⁶ is a C1-C6 alkoxy group optionally having one or more halogen atoms;

A compound of the formula (M20) wherein R⁶ is a C1-C6 alkylsulfanyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfinyl group optionally having one or more halogen atoms, or a C1-C6 alkylsulfonyl group optionally having one or more halogen atoms;

A compound of the formula (M20) wherein R⁶ is a bromine atom or an iodine atom;

A compound of the formula (M20) wherein R⁶ is a trifluoromethyl group, a pentafluoroethyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

A compound of the formula (M20) wherein R^6 is a pentafluoroethyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

5 A compound of the formula (M20) wherein R^6 is a pentafluoroethyl group;

A compound of the formula (M20) wherein R^6 is a trifluoromethylsulfanyl group;

10 A compound of the formula (M20) wherein R^6 is a trifluoromethylsulfinyl group;

A compound of the formula (M20) wherein R^6 is a trifluoromethylsulfonyl group;

15 A compound of the formula (M20) wherein R^6 is a trifluoromethyl group, a pentafluoroethyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, and R^7 is hydrogen;

20 A compound of the formula (M20) wherein R^6 is a pentafluoroethyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, and R^7 is hydrogen;

A compound of the formula (M20) wherein R^6 is a pentafluoroethyl group, and R^7 is hydrogen;

25 A compound of the formula (M20) wherein R^6 is a

trifluoromethylsulfanyl group, and R^7 is hydrogen;

A compound of the formula (M20) wherein R^6 is a trifluoromethylsulfinyl group, and R^7 is hydrogen;

A compound of the formula (M20) wherein R^6 is a
5 trifluoromethylsulfonyl group, and R^7 is hydrogen;

A compound of the formula (M20) wherein A^2 is $=CH-$, R^6 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, R^7 is hydrogen, and R^8 is a C1-C6 chain hydrocarbon group optionally having one or more halogen
10 atoms;

A compound of the formula (M20) wherein X^9 is a bromine atom, A^2 is $=CH-$, R^6 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, R^7 is hydrogen, and R^8 is a C1-C6 chain hydrocarbon group
15 optionally having one or more halogen atoms;

A compound of the formula (M20) wherein A^2 is $=CH-$, R^6 is a trifluoromethyl group, a difluoromethyl group, a fluoromethyl group, a pentafluoroethyl group, a trifluoromethoxy group, or a trifluoromethylsulfanyl group,
20 R^7 is hydrogen, and R^8 is a methyl group, an ethyl group, a propyl group, or an isopropyl group;

A compound of the formula (M20) wherein A^2 is $=CH-$, R^6 is a trifluoromethyl group or a pentafluoroethyl group, R^7 is hydrogen, and R^8 is a methyl group;

25 A compound of the formula (M20) wherein A^2 is $=CH-$, R^6

is a trifluoromethyl group, a pentafluoroethyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, R^7 is hydrogen, and R^8 is a methyl group;


5 A compound of the formula (M20) wherein A^2 is $=CH-$, R^6 is a pentafluoroethyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, R^7 is hydrogen, and R^8 is a methyl group;

10 A compound of the formula (M20) wherein X^9 is a bromine atom, A^2 is $=CH-$, R^6 is a pentafluoroethyl group, a trifluoromethoxy group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group, R^7 is hydrogen, and R^8 is a methyl group.

15 [0159]

Examples of Compound (3), Compound (5), and Compound (M3) include the following compounds:

A compound of the formula (1H):



20 wherein:

R^{1h} represents a C1-C6 alkyl group optionally having one or more halogen atoms;

R^{13} represents a hydrogen atom, hydroxy group, or a chlorine atom;

R^{2h} , R^{3h} , R^{4h} , and R^{5h} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11h}$, $-S(O)_mR^{11h}$, $-SF_5$, a halogen atom, or a hydrogen atom;

R^{11h} and R^{12h} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a hydrogen atom; and

n represents 0, 1, or 2 (hereinafter, referred to as Compound (1H));

[0160]

And examples of Compound (1H) include the following compounds:

A compound of the formula (1H) wherein R^{1h} is a C1-C3 alkyl group optionally having one or more halogen atoms;

A compound of the formula (1H) wherein R^{1h} is a C1-C3 alkyl group;

A compound of the formula (1H) wherein R^{1h} is a methyl group;

A compound of the formula (1H) wherein R^{1h} is an ethyl group;

A compound of the formula (1H) wherein R^{1h} is a propyl group;

A compound of the formula (1H) wherein R^{13} is a

hydrogen atom;

A compound of the formula (1H) wherein R^{13} is hydroxy group;

5 A compound of the formula (1H) wherein R^{13} is a chlorine atom;

A compound of the formula (1H) wherein R^{13} is a hydrogen atom or hydroxy group;

A compound of the formula (1H) wherein R^{2h} , R^{3h} , R^{4h} , and R^{5h} are a hydrogen atom;

10 A compound of the formula (1H) wherein R^{2h} , R^{3h} , R^{4h} , and R^{5h} are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a hydrogen atom;

15 A compound of the formula (1H) wherein R^{2h} , R^{3h} , R^{4h} , and R^{5h} are same or different and are independently $-OR^{11h}$ or a hydrogen atom;

A compound of the formula (1H) wherein R^{2h} , R^{3h} , R^{4h} , and R^{5h} are same or different and are independently $-S(O)_mR^{11h}$ or a hydrogen atom;

20 A compound of the formula (1H) wherein R^{2h} , R^{3h} , R^{4h} , and R^{5h} are same or different and are independently a halogen atom or a hydrogen atom;

25 A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11h}$, -

$S(O)_m R^{11h}$ or a halogen atom;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a C1-C3 alkyl group;

5 A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a C1-C3 alkyl group optionally having one or more fluorine atoms;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a C1-C3 alkyl group having one or more fluorine atoms;

10 A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, or a heptafluoroisopropyl group;

15 A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a methyl group;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is an ethyl group;

20 A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a propyl group;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is an isopropyl group;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a trifluoromethyl group;

25 A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h}

are a hydrogen atom, and R^{3h} is a pentafluoroethyl group;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a heptafluoropropyl group;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h}
5 are a hydrogen atom, and R^{3h} is a heptafluoroisopropyl group;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is $-OR^{11h}$;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h}
10 are a hydrogen atom, and R^{3h} is a C1-C3 alkoxy group optionally having one or more fluorine atoms;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a trifluoromethoxy group;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h}
15 are a hydrogen atom, and R^{3h} is $-S(O)_mR^{11h}$;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a C1-C3 alkylsulfanyl group optionally having one or more fluorine atoms, a C1-C3 alkylsulfinyl group optionally having one or more fluorine
20 atoms, or a C1-C3 alkylsulfonyl group optionally having one or more fluorine atoms;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group or a
25 trifluoromethylsulfonyl group;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a trifluoromethylsulfanyl group;

5 A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a trifluoromethylsulfinyl group;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a trifluoromethylsulfonyl group;

10 A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a halogen atom;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a fluorine atom, a chlorine atom, or a bromine atom;

15 A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a fluorine atom;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a chlorine atom;

20 A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a bromine atom;

A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is $-SF_5$;

25 A compound of the formula (1H) wherein R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a

trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a fluorine atom, a chlorine atom, a bromine atom, or $-SF_5$;

A compound of the formula (1H) wherein n is 0;

A compound of the formula (1H) wherein n is 1;

A compound of the formula (1H) wherein n is 2;

A compound of the formula (1H) wherein R^{1h} is a methyl group, R^{2h} , R^{4h} , and R^{5h} are same or different and are independently a halogen atom or a hydrogen atom, and R^{3h} a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11h}$, $-S(O)_mR^{11h}$, a halogen atom, or a hydrogen atom;

A compound of the formula (1H) wherein R^{1h} is a methyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a C1-C3 alkyl group optionally having one or more fluorine atoms;

A compound of the formula (1H) wherein R^{1h} is a methyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a C1-C3 alkoxy group optionally having one or more fluorine atoms;

A compound of the formula (1H) wherein R^{1h} is a methyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a C1-C3 alkylsulfanyl group optionally having one or more

fluorine atoms, a C1-C3 alkylsulfinyl group optionally having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group optionally having one or more fluorine atoms;

5 A compound of the formula (1H) wherein R^{1h} is a methyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a halogen atom;

A compound of the formula (1H) wherein R^{1h} is a methyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a methyl group;

10 A compound of the formula (1H) wherein R^{1h} is a methyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is an ethyl group;

A compound of the formula (1H) wherein R^{1h} is a methyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a
15 trifluoromethyl group;

A compound of the formula (1H) wherein R^{1h} is a methyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a trifluoromethoxy group;

20 A compound of the formula (1H) wherein R^{1h} is a methyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a fluorine atom;

A compound of the formula (1H) wherein R^{1h} is a methyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a chlorine atom;

25 A compound of the formula (1H) wherein R^{1h} is a methyl

group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a bromine atom;

A compound of the formula (1H) wherein R^{1h} is a methyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is $-SF_5$;

5 A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are same or different and are independently a halogen atom or a hydrogen atom, and R^{3h} a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11h}$, $-S(O)_mR^{11h}$, a halogen atom, or a
10 hydrogen atom;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a C1-C3 alkyl group optionally having one or more fluorine atoms;

15 A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a C1-C3 alkoxy group optionally having one or more fluorine atoms;

A compound of the formula (1H) wherein R^{1h} is an ethyl
20 group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a C1-C3 alkylsulfanyl group optionally having one or more fluorine atoms, a C1-C3 alkylsulfinyl group optionally having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group optionally having one or more fluorine atoms;

25 A compound of the formula (1H) wherein R^{1h} is an ethyl

group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a halogen atom;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a methyl group;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, R^{3h} is a methyl group, and R^{13} is hydroxy group;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, R^{3h} is a methyl group, and R^{13} is a chlorine atom;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, R^{3h} is a methyl group, and R^{13} is a hydrogen atom;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is an ethyl group;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a trifluoromethyl group;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, R^{3h} is a trifluoromethyl group, and R^{13} is hydroxy group;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, R^{3h} is a

trifluoromethyl group, and R^{13} is a chlorine atom;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, R^{3h} is a trifluoromethyl group, and R^{13} is a hydrogen atom;

5 A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a trifluoromethoxy group;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a
10 fluorine atom;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a chlorine atom;

A compound of the formula (1H) wherein R^{1h} is an ethyl
15 group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, R^{3h} is a chlorine atom, and R^{13} is hydroxy group;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, R^{3h} is a chlorine atom, and R^{13} is a chlorine atom;

20 A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, R^{3h} is a chlorine atom, and R^{13} is a hydrogen atom;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a
25 bromine atom;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, R^{3h} is a bromine atom, and R^{13} is hydroxy group;

5 A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, R^{3h} is a bromine atom, and R^{13} is a chlorine atom;

A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, R^{3h} is a bromine atom, and R^{13} is a hydrogen atom;

10 A compound of the formula (1H) wherein R^{1h} is an ethyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is $-SF_5$;

A compound of the formula (1H) wherein R^{1h} is a propyl group, R^{2h} , R^{4h} , and R^{5h} are same or different and are independently a halogen atom or a hydrogen atom, and R^{3h} a
15 Cl-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11h}$, $-S(O)_mR^{11h}$, a halogen atom, or a hydrogen atom;

A compound of the formula (1H) wherein R^{1h} is a propyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a
20 Cl-C3 alkyl group optionally having one or more fluorine atoms;

A compound of the formula (1H) wherein R^{1h} is a propyl group, and R^{3h} is a Cl-C3 alkoxy group optionally having one or more fluorine atoms;

25 A compound of the formula (1H) wherein R^{1h} is a propyl

group, and R^{3h} is a C1-C3 alkylsulfanyl group optionally having one or more fluorine atoms, a C1-C3 alkylsulfinyl group optionally having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group optionally having one or more
5 fluorine atoms;

A compound of the formula (1H) wherein R^{1h} is a propyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a halogen atom;

A compound of the formula (1H) wherein R^{1h} is a propyl
10 group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a methyl group;

A compound of the formula (1H) wherein R^{1h} is a propyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is an ethyl group;

15 A compound of the formula (1H) wherein R^{1h} is a propyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a trifluoromethyl group;

A compound of the formula (1H) wherein R^{1h} is a propyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a
20 trifluoromethoxy group;

A compound of the formula (1H) wherein R^{1h} is a propyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a fluorine atom;

A compound of the formula (1H) wherein R^{1h} is a propyl
25 group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a


chlorine atom;

A compound of the formula (1H) wherein R^{1h} is a propyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is a bromine atom;

- 5 A compound of the formula (1H) wherein R^{1h} is a propyl group, R^{2h} , R^{4h} , and R^{5h} are a hydrogen atom, and R^{3h} is $-SF_5$.
[0161]

Examples of Compound (M22) include the following compounds:

- 10 A compound of the formula (1I):



wherein:

R^{1i} represents a C1-C6 alkyl group optionally having one or more halogen atoms;

- 15 R^{2i} , R^{3i} , R^{4i} , and R^{5i} are same or different and represent independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11i}$, $-S(O)_mR^{11i}$, $-SF_5$, a halogen atom, or a hydrogen atom;

- R^{11i} and R^{12i} are same or different and represent
20 independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a hydrogen atom; and

r represents 1 or 2 (hereinafter, referred to as Compound (1I));

[0162]

And examples of Compound (1I) include the following compounds:

A compound of the formula (1I) wherein R^{1i} is a C1-C3 alkyl group optionally having one or more halogen atoms;

A compound of the formula (1I) wherein R^{1i} is a C1-C3 alkyl group;

A compound of the formula (1I) wherein R^{1i} is a methyl group;

A compound of the formula (1I) wherein R^{1i} is an ethyl group;

A compound of the formula (1I) wherein R^{1i} is a propyl group;

A compound of the formula (1I) wherein R^{2i} , R^{3i} , R^{4i} , and R^{5i} are a hydrogen atom;

A compound of the formula (1I) wherein R^{2i} , R^{3i} , R^{4i} , and R^{5i} are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms or a hydrogen atom;

A compound of the formula (1I) wherein R^{2i} , R^{3i} , R^{4i} , and R^{5i} are same or different and are independently $-OR^{11i}$ or a hydrogen atom;

A compound of the formula (1I) wherein R^{2i} , R^{3i} , R^{4i} , and R^{5i} are same or different and are independently $-S(O)_mR^{11i}$ or a hydrogen atom;

A compound of the formula (1I) wherein R^{2i} , R^{3i} , R^{4i} , and R^{5i} are same or different and are independently a halogen atom or a hydrogen atom;

5 A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11i}$, $-S(O)_mR^{11i}$, or a halogen atom;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a C1-C3 alkyl group;

10 A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a C1-C3 alkyl group optionally having one or more fluorine atoms;

15 A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a C1-C3 alkyl group having one or more fluorine atoms;

20 A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, or a heptafluoroisopropyl group;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a methyl group;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is an ethyl group;

25 A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i}

are a hydrogen atom, and R^{3i} is a propyl group;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is an isopropyl group;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a trifluoromethyl group;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a pentafluoroethyl group;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a heptafluoropropyl group;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a heptafluoroisopropyl group;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is $-OR^{11i}$;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a C1-C3 alkoxy group optionally having one or more fluorine atoms;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a trifluoromethoxy group;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is $-S(O)_mR^{11i}$;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a C1-C3 alkylsulfanyl group optionally having one or more fluorine atoms, a C1-C3 alkylsulfinyl group optionally having one or more fluorine

atoms, or a C1-C3 alkylsulfonyl group optionally having one or more fluorine atoms;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, or a trifluoromethylsulfonyl group;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a trifluoromethylsulfanyl group;

10 A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a trifluoromethylsulfinyl group;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a trifluoromethylsulfonyl group;

15 A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a halogen atom;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a fluorine atom, a chlorine atom, or a bromine atom;

20 A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a fluorine atom;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a chlorine atom;

25 A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i}

are a hydrogen atom, and R^{3i} is a bromine atom;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is $-SF_5$;

A compound of the formula (1I) wherein R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a methyl group, an ethyl group, a propyl group, an isopropyl group, a trifluoromethyl group, a pentafluoroethyl group, a heptafluoropropyl group, a heptafluoroisopropyl group, a trifluoromethylsulfanyl group, a trifluoromethylsulfinyl group, a trifluoromethylsulfonyl group, a fluorine atom, a chlorine atom, a bromine atom, or $-SF_5$;

A compound of the formula (1I) wherein r is 1;

A compound of the formula (1I) wherein r is 2;

A compound of the formula (1I) wherein R^{1i} is a methyl group, R^{2i} , R^{4i} , and R^{5i} are same or different and are independently a halogen atom or a hydrogen atom, and R^{3i} a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11i}$, $-S(O)_mR^{11i}$, a halogen atom, or a hydrogen atom;

A compound of the formula (1I) wherein R^{1i} is a methyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a C1-C3 alkyl group optionally having one or more fluorine atoms;

A compound of the formula (1I) wherein R^{1i} is a methyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a

C1-C3 alkoxy group optionally having one or more fluorine atoms;

A compound of the formula (1I) wherein R^{1i} is a methyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a
5 C1-C3 alkylsulfanyl group optionally having one or more fluorine atoms, a C1-C3 alkylsulfinyl group optionally having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group optionally having one or more fluorine atoms;

A compound of the formula (1I) wherein R^{1i} is a methyl
10 group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a halogen atom;

A compound of the formula (1I) wherein R^{1i} is a methyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a methyl group;

15 A compound of the formula (1I) wherein R^{1i} is a methyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is an ethyl group;

A compound of the formula (1I) wherein R^{1i} is a methyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a
20 trifluoromethyl group;

A compound of the formula (1I) wherein R^{1i} is a methyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a
(trifluoromethoxy group;

A compound of the formula (1I) wherein R^{1i} is a methyl
25 group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a

fluorine atom;

A compound of the formula (1I) wherein R^{1i} is a methyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a chlorine atom;

5 A compound of the formula (1I) wherein R^{1i} is a methyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a bromine atom;

A compound of the formula (1I) wherein R^{1i} is a methyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is $-SF_5$;

10 A compound of the formula (1I) wherein R^{1i} is an ethyl group, R^{2i} , R^{4i} , and R^{5i} are same or different and are independently a halogen atom or a hydrogen atom, and R^{3i} a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11i}$, $-S(O)_mR^{11i}$, a halogen atom, or a
15 hydrogen atom;

A compound of the formula (1I) wherein R^{1i} is an ethyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a C1-C3 alkyl group optionally having one or more fluorine atoms;

20 A compound of the formula (1I) wherein R^{1i} is an ethyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a C1-C3 alkoxy group optionally having one or more fluorine atoms;

25 A compound of the formula (1I) wherein R^{1i} is an ethyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a

C1-C3 alkylsulfanyl group optionally having one or more fluorine atoms, a C1-C3 alkylsulfinyl group optionally having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group optionally having one or more fluorine atoms;

5 A compound of the formula (1I) wherein R^{1i} is an ethyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a halogen atom;

10 A compound of the formula (1I) wherein R^{1i} is an ethyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a methyl group;

 A compound of the formula (1I) wherein R^{1i} is an ethyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is an ethyl group;

15 A compound of the formula (1I) wherein R^{1i} is an ethyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a trifluoromethyl group;

 A compound of the formula (1I) wherein R^{1i} is an ethyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a trifluoromethoxy group;

20 A compound of the formula (1I) wherein R^{1i} is an ethyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a fluorine atom;

25 A compound of the formula (1I) wherein R^{1i} is an ethyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a chlorine atom;

A compound of the formula (1I) wherein R^{1i} is an ethyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a bromine atom;

5 A compound of the formula (1I) wherein R^{1i} is an ethyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is $-SF_5$;

A compound of the formula (1I) wherein R^{1i} is a propyl group, R^{2i} , R^{4i} , and R^{5i} are same or different and are independently a halogen atom or a hydrogen atom, and R^{3i} a C1-C6 chain hydrocarbon group optionally having one or more
10 halogen atoms, $-OR^{11i}$, $-S(O)_mR^{11i}$, a halogen atom, or a hydrogen atom;

A compound of the formula (1I) wherein R^{1i} is a propyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a C1-C3 alkyl group optionally having one or more fluorine
15 atoms;

A compound of the formula (1I) wherein R^{1i} is a propyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a C1-C3 alkoxy group optionally having one or more fluorine atoms;

20 A compound of the formula (1I) wherein R^{1i} is a propyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a C1-C3 alkylsulfanyl group optionally having one or more fluorine atoms, a C1-C3 alkylsulfinyl group optionally having one or more fluorine atoms, or a C1-C3 alkylsulfonyl
25 group optionally having one or more fluorine atoms;

A compound of the formula (1I) wherein R^{1i} is a propyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a halogen atom;

5 A compound of the formula (1I) wherein R^{1i} is a propyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a methyl group;

A compound of the formula (1I) wherein R^{1i} is a propyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is an ethyl group;

10 A compound of the formula (1I) wherein R^{1i} is a propyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a trifluoromethyl group;

A compound of the formula (1I) wherein R^{1i} is a propyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a
15 trifluoromethoxy group;

A compound of the formula (1I) wherein R^{1i} is a propyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a fluorine atom;

A compound of the formula (1I) wherein R^{1i} is a propyl
20 group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a chlorine atom;

A compound of the formula (1I) wherein R^{1i} is a propyl group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is a bromine atom;


25 A compound of the formula (1I) wherein R^{1i} is a propyl

group, R^{2i} , R^{4i} , and R^{5i} are a hydrogen atom, and R^{3i} is $-SF_5$.

[0163]

Examples of the present compound are shown bellow.

A compound of the formula (A):



wherein R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19:

[0164]

Table 1

R ¹	A ¹	A ³	n
Me	-NMe-	=N-	0
Me	-NMe-	=N-	1
Me	-NMe-	=N-	2
Et	-NMe-	=N-	0
Et	-NMe-	=N-	1
Et	-NMe-	=N-	2
Pr	-NMe-	=N-	0
Pr	-NMe-	=N-	1
Pr	-NMe-	=N-	2
iPr	-NMe-	=N-	0
iPr	-NMe-	=N-	1
iPr	-NMe-	=N-	2
tBu	-NMe-	=N-	0
tBu	-NMe-	=N-	1
tBu	-NMe-	=N-	2
CF ₃	-NMe-	=N-	0
CF ₃	-NMe-	=N-	1
CF ₃	-NMe-	=N-	2
CH ₂ CF ₃	-NMe-	=N-	0
CH ₂ CF ₃	-NMe-	=N-	1
CH ₂ CF ₃	-NMe-	=N-	2
CH=CH ₂	-NMe-	=N-	0
CH=CH ₂	-NMe-	=N-	1
CH=CH ₂	-NMe-	=N-	2

[0165]

Table 2

R ¹	A ¹	A ³	n
CH ₂ CH=CH ₂	-NMe-	=N-	0
CH ₂ CH=CH ₂	-NMe-	=N-	1
CH ₂ CH=CH ₂	-NMe-	=N-	2
C≡CH	-NMe-	=N-	0
C≡CH	-NMe-	=N-	1
C≡CH	-NMe-	=N-	2
CH ₂ C≡CH	-NMe-	=N-	0
CH ₂ C≡CH	-NMe-	=N-	1
CH ₂ C≡CH	-NMe-	=N-	2
CyPr	-NMe-	=N-	0
CyPr	-NMe-	=N-	1
CyPr	-NMe-	=N-	2
Me	-NMe-	=CH-	0
Me	-NMe-	=CH-	1
Me	-NMe-	=CH-	2
Et	-NMe-	=CH-	0
Et	-NMe-	=CH-	1
Et	-NMe-	=CH-	2
Pr	-NMe-	=CH-	0
Pr	-NMe-	=CH-	1
Pr	-NMe-	=CH-	2
iPr	-NMe-	=CH-	0
iPr	-NMe-	=CH-	1
iPr	-NMe-	=CH-	2

[0166]

Table 3

R ¹	A ¹	A ³	n
t B u	-NMe-	=CH-	0
t B u	-NMe-	=CH-	1
t B u	-NMe-	=CH-	2
CF ₃	-NMe-	=CH-	0
CF ₃	-NMe-	=CH-	1
CF ₃	-NMe-	=CH-	2
CH ₂ CF ₃	-NMe-	=CH-	0
CH ₂ CF ₃	-NMe-	=CH-	1
CH ₂ CF ₃	-NMe-	=CH-	2
CH=CH ₂	-NMe-	=CH-	0
CH=CH ₂	-NMe-	=CH-	1
CH=CH ₂	-NMe-	=CH-	2
CH ₂ CH=CH ₂	-NMe-	=CH-	0
CH ₂ CH=CH ₂	-NMe-	=CH-	1
CH ₂ CH=CH ₂	-NMe-	=CH-	2
C≡CH	-NMe-	=CH-	0
C≡CH	-NMe-	=CH-	1
C≡CH	-NMe-	=CH-	2
CH ₂ C≡CH	-NMe-	=CH-	0
CH ₂ C≡CH	-NMe-	=CH-	1
CH ₂ C≡CH	-NMe-	=CH-	2
Cy P r	-NMe-	=CH-	0
Cy P r	-NMe-	=CH-	1
Cy P r	-NMe-	=CH-	2

[0167]

Table 4

R ¹	A ¹	A ³	n
Me	-NMe-	=CB r-	0
Me	-NMe-	=CB r-	1
Me	-NMe-	=CB r-	2
Et	-NMe-	=CB r-	0
Et	-NMe-	=CB r-	1
Et	-NMe-	=CB r-	2
Pr	-NMe-	=CB r-	0
Pr	-NMe-	=CB r-	1
Pr	-NMe-	=CB r-	2
iPr	-NMe-	=CB r-	0
iPr	-NMe-	=CB r-	1
iPr	-NMe-	=CB r-	2
tBu	-NMe-	=CB r-	0
tBu	-NMe-	=CB r-	1
tBu	-NMe-	=CB r-	2
CF ₃	-NMe-	=CB r-	0
CF ₃	-NMe-	=CB r-	1
CF ₃	-NMe-	=CB r-	2
CH ₂ CF ₃	-NMe-	=CB r-	0
CH ₂ CF ₃	-NMe-	=CB r-	1
CH ₂ CF ₃	-NMe-	=CB r-	2
CH=CH ₂	-NMe-	=CB r-	0
CH=CH ₂	-NMe-	=CB r-	1
CH=CH ₂	-NMe-	=CB r-	2

[0168]

Table 5

R ¹	A ¹	A ³	n
CH ₂ CH=CH ₂	-NMe-	=CB r-	0
CH ₂ CH=CH ₂	-NMe-	=CB r-	1
CH ₂ CH=CH ₂	-NMe-	=CB r-	2
C≡CH	-NMe-	=CB r-	0
C≡CH	-NMe-	=CB r-	1
C≡CH	-NMe-	=CB r-	2
CH ₂ C≡CH	-NMe-	=CB r-	0
CH ₂ C≡CH	-NMe-	=CB r-	1
CH ₂ C≡CH	-NMe-	=CB r-	2
CyPr	-NMe-	=CB r-	0
CyPr	-NMe-	=CB r-	1
CyPr	-NMe-	=CB r-	2
Me	-N(CH ₂ OMe)-	=N-	0
Me	-N(CH ₂ OMe)-	=N-	1
Me	-N(CH ₂ OMe)-	=N-	2
Et	-N(CH ₂ OMe)-	=N-	0
Et	-N(CH ₂ OMe)-	=N-	1
Et	-N(CH ₂ OMe)-	=N-	2
Pr	-N(CH ₂ OMe)-	=N-	0
Pr	-N(CH ₂ OMe)-	=N-	1
Pr	-N(CH ₂ OMe)-	=N-	2
iPr	-N(CH ₂ OMe)-	=N-	0
iPr	-N(CH ₂ OMe)-	=N-	1
iPr	-N(CH ₂ OMe)-	=N-	2

[0169]

Table 6

R ¹	A ¹	A ³	n
t B u	-N (CH ₂ OMe) -	=N-	0
t B u	-N (CH ₂ OMe) -	=N-	1
t B u	-N (CH ₂ OMe) -	=N-	2
CF ₃	-N (CH ₂ OMe) -	=N-	0
CF ₃	-N (CH ₂ OMe) -	=N-	1
CF ₃	-N (CH ₂ OMe) -	=N-	2
CH ₂ CF ₃	-N (CH ₂ OMe) -	=N-	0
CH ₂ CF ₃	-N (CH ₂ OMe) -	=N-	1
CH ₂ CF ₃	-N (CH ₂ OMe) -	=N-	2
CH=CH ₂	-N (CH ₂ OMe) -	=N-	0
CH=CH ₂	-N (CH ₂ OMe) -	=N-	1
CH=CH ₂	-N (CH ₂ OMe) -	=N-	2
CH ₂ CH=CH ₂	-N (CH ₂ OMe) -	=N-	0
CH ₂ CH=CH ₂	-N (CH ₂ OMe) -	=N-	1
CH ₂ CH=CH ₂	-N (CH ₂ OMe) -	=N-	2
C≡CH	-N (CH ₂ OMe) -	=N-	0
C≡CH	-N (CH ₂ OMe) -	=N-	1
C≡CH	-N (CH ₂ OMe) -	=N-	2
CH ₂ C≡CH	-N (CH ₂ OMe) -	=N-	0
CH ₂ C≡CH	-N (CH ₂ OMe) -	=N-	1
CH ₂ C≡CH	-N (CH ₂ OMe) -	=N-	2
C y P r	-N (CH ₂ OMe) -	=N-	0
C y P r	-N (CH ₂ OMe) -	=N-	1
C y P r	-N (CH ₂ OMe) -	=N-	2

[0170]

Table 7

R ¹	A ¹	A ³	n
Me	-N (CH ₂ OMe) -	=CH-	0
Me	-N (CH ₂ OMe) -	=CH-	1
Me	-N (CH ₂ OMe) -	=CH-	2
Et	-N (CH ₂ OMe) -	=CH-	0
Et	-N (CH ₂ OMe) -	=CH-	1
Et	-N (CH ₂ OMe) -	=CH-	2
Pr	-N (CH ₂ OMe) -	=CH-	0
Pr	-N (CH ₂ OMe) -	=CH-	1
Pr	-N (CH ₂ OMe) -	=CH-	2
iPr	-N (CH ₂ OMe) -	=CH-	0
iPr	-N (CH ₂ OMe) -	=CH-	1
iPr	-N (CH ₂ OMe) -	=CH-	2
tBu	-N (CH ₂ OMe) -	=CH-	0
tBu	-N (CH ₂ OMe) -	=CH-	1
tBu	-N (CH ₂ OMe) -	=CH-	2
CF ₃	-N (CH ₂ OMe) -	=CH-	0
CF ₃	-N (CH ₂ OMe) -	=CH-	1
CF ₃	-N (CH ₂ OMe) -	=CH-	2
CH ₂ CF ₃	-N (CH ₂ OMe) -	=CH-	0
CH ₂ CF ₃	-N (CH ₂ OMe) -	=CH-	1
CH ₂ CF ₃	-N (CH ₂ OMe) -	=CH-	2
CH=CH ₂	-N (CH ₂ OMe) -	=CH-	0
CH=CH ₂	-N (CH ₂ OMe) -	=CH-	1
CH=CH ₂	-N (CH ₂ OMe) -	=CH-	2

[0171]

Table 8

R ¹	A ¹	A ³	n
CH ₂ CH=CH ₂	-N(CH ₂ OMe)-	=CH-	0
CH ₂ CH=CH ₂	-N(CH ₂ OMe)-	=CH-	1
CH ₂ CH=CH ₂	-N(CH ₂ OMe)-	=CH-	2
C≡CH	-N(CH ₂ OMe)-	=CH-	0
C≡CH	-N(CH ₂ OMe)-	=CH-	1
C≡CH	-N(CH ₂ OMe)-	=CH-	2
CH ₂ C≡CH	-N(CH ₂ OMe)-	=CH-	0
CH ₂ C≡CH	-N(CH ₂ OMe)-	=CH-	1
CH ₂ C≡CH	-N(CH ₂ OMe)-	=CH-	2
CyPr	-N(CH ₂ OMe)-	=CH-	0
CyPr	-N(CH ₂ OMe)-	=CH-	1
CyPr	-N(CH ₂ OMe)-	=CH-	2
Me	-N(CH ₂ OMe)-	=CBr-	0
Me	-N(CH ₂ OMe)-	=CBr-	1
Me	-N(CH ₂ OMe)-	=CBr-	2
Et	-N(CH ₂ OMe)-	=CBr-	0
Et	-N(CH ₂ OMe)-	=CBr-	1
Et	-N(CH ₂ OMe)-	=CBr-	2
Pr	-N(CH ₂ OMe)-	=CBr-	0
Pr	-N(CH ₂ OMe)-	=CBr-	1
Pr	-N(CH ₂ OMe)-	=CBr-	2
iPr	-N(CH ₂ OMe)-	=CBr-	0
iPr	-N(CH ₂ OMe)-	=CBr-	1
iPr	-N(CH ₂ OMe)-	=CBr-	2

[0172]

Table 9

R ¹	A ¹	A ³	n
t B u	-N (CH ₂ OMe) -	=C B r -	0
t B u	-N (CH ₂ OMe) -	=C B r -	1
t B u	-N (CH ₂ OMe) -	=C B r -	2
C F ₃	-N (CH ₂ OMe) -	=C B r -	0
C F ₃	-N (CH ₂ OMe) -	=C B r -	1
C F ₃	-N (CH ₂ OMe) -	=C B r -	2
CH ₂ C F ₃	-N (CH ₂ OMe) -	=C B r -	0
CH ₂ C F ₃	-N (CH ₂ OMe) -	=C B r -	1
CH ₂ C F ₃	-N (CH ₂ OMe) -	=C B r -	2
CH=CH ₂	-N (CH ₂ OMe) -	=C B r -	0
CH=CH ₂	-N (CH ₂ OMe) -	=C B r -	1
CH=CH ₂	-N (CH ₂ OMe) -	=C B r -	2
CH ₂ CH=CH ₂	-N (CH ₂ OMe) -	=C B r -	0
CH ₂ CH=CH ₂	-N (CH ₂ OMe) -	=C B r -	1
CH ₂ CH=CH ₂	-N (CH ₂ OMe) -	=C B r -	2
C≡CH	-N (CH ₂ OMe) -	=C B r -	0
C≡CH	-N (CH ₂ OMe) -	=C B r -	1
C≡CH	-N (CH ₂ OMe) -	=C B r -	2
CH ₂ C≡CH	-N (CH ₂ OMe) -	=C B r -	0
CH ₂ C≡CH	-N (CH ₂ OMe) -	=C B r -	1
CH ₂ C≡CH	-N (CH ₂ OMe) -	=C B r -	2
C y P r	-N (CH ₂ OMe) -	=C B r -	0
C y P r	-N (CH ₂ OMe) -	=C B r -	1
C y P r	-N (CH ₂ OMe) -	=C B r -	2

[0173]

Table 10

R ¹	A ¹	A ³	n
Me	O	=N-	0
Me	O	=N-	1
Me	O	=N-	2
Et	O	=N-	0
Et	O	=N-	1
Et	O	=N-	2
Pr	O	=N-	0
Pr	O	=N-	1
Pr	O	=N-	2
iPr	O	=N-	0
iPr	O	=N-	1
iPr	O	=N-	2
tBu	O	=N-	0
tBu	O	=N-	1
tBu	O	=N-	2
CF ₃	O	=N-	0
CF ₃	O	=N-	1
CF ₃	O	=N-	2
CH ₂ CF ₃	O	=N-	0
CH ₂ CF ₃	O	=N-	1
CH ₂ CF ₃	O	=N-	2
CH=CH ₂	O	=N-	0
CH=CH ₂	O	=N-	1
CH=CH ₂	O	=N-	2

[0174]

Table 11

R ¹	A ¹	A ³	n
CH ₂ CH=CH ₂	O	=N-	0
CH ₂ CH=CH ₂	O	=N-	1
CH ₂ CH=CH ₂	O	=N-	2
C≡CH	O	=N-	0
C≡CH	O	=N-	1
C≡CH	O	=N-	2
CH ₂ C≡CH	O	=N-	0
CH ₂ C≡CH	O	=N-	1
CH ₂ C≡CH	O	=N-	2
CyPr	O	=N-	0
CyPr	O	=N-	1
CyPr	O	=N-	2
Me	O	=CH-	0
Me	O	=CH-	1
Me	O	=CH-	2
Et	O	=CH-	0
Et	O	=CH-	1
Et	O	=CH-	2
Pr	O	=CH-	0
Pr	O	=CH-	1
Pr	O	=CH-	2
iPr	O	=CH-	0
iPr	O	=CH-	1
iPr	O	=CH-	2

[0175]

Table 12

R ¹	A ¹	A ³	n
t B u	O	=CH-	0
t B u	O	=CH-	1
t B u	O	=CH-	2
CF ₃	O	=CH-	0
CF ₃	O	=CH-	1
CF ₃	O	=CH-	2
CH ₂ CF ₃	O	=CH-	0
CH ₂ CF ₃	O	=CH-	1
CH ₂ CF ₃	O	=CH-	2
CH=CH ₂	O	=CH-	0
CH=CH ₂	O	=CH-	1
CH=CH ₂	O	=CH-	2
CH ₂ CH=CH ₂	O	=CH-	0
CH ₂ CH=CH ₂	O	=CH-	1
CH ₂ CH=CH ₂	O	=CH-	2
C≡CH	O	=CH-	0
C≡CH	O	=CH-	1
C≡CH	O	=CH-	2
CH ₂ C≡CH	O	=CH-	0
CH ₂ C≡CH	O	=CH-	1
CH ₂ C≡CH	O	=CH-	2
C y P r	O	=CH-	0
C y P r	O	=CH-	1
C y P r	O	=CH-	2

[0176]

Table 13

R ¹	A ¹	A ³	n
Me	O	=CBr-	0
Me	O	=CBr-	1
Me	O	=CBr-	2
Et	O	=CBr-	0
Et	O	=CBr-	1
Et	O	=CBr-	2
Pr	O	=CBr-	0
Pr	O	=CBr-	1
Pr	O	=CBr-	2
iPr	O	=CBr-	0
iPr	O	=CBr-	1
iPr	O	=CBr-	2
tBu	O	=CBr-	0
tBu	O	=CBr-	1
tBu	O	=CBr-	2
CF ₃	O	=CBr-	0
CF ₃	O	=CBr-	1
CF ₃	O	=CBr-	2
CH ₂ CF ₃	O	=CBr-	0
CH ₂ CF ₃	O	=CBr-	1
CH ₂ CF ₃	O	=CBr-	2
CH=CH ₂	O	=CBr-	0
CH=CH ₂	O	=CBr-	1
CH=CH ₂	O	=CBr-	2

[0177]

Table 14

R^1	A^1	A^3	n
$CH_2CH=CH_2$	O	$=CB r -$	0
$CH_2CH=CH_2$	O	$=CB r -$	1
$CH_2CH=CH_2$	O	$=CB r -$	2
$C \equiv CH$	O	$=CB r -$	0
$C \equiv CH$	O	$=CB r -$	1
$C \equiv CH$	O	$=CB r -$	2
$CH_2C \equiv CH$	O	$=CB r -$	0
$CH_2C \equiv CH$	O	$=CB r -$	1
$CH_2C \equiv CH$	O	$=CB r -$	2
CyPr	O	$=CB r -$	0
CyPr	O	$=CB r -$	1
CyPr	O	$=CB r -$	2

[0178]

Table 15

R ¹	A ¹	A ³	n
Me	S	=N-	0
Me	S	=N-	1
Me	S	=N-	2
Et	S	=N-	0
Et	S	=N-	1
Et	S	=N-	2
Pr	S	=N-	0
Pr	S	=N-	1
Pr	S	=N-	2
iPr	S	=N-	0
iPr	S	=N-	1
iPr	S	=N-	2
tBu	S	=N-	0
tBu	S	=N-	1
tBu	S	=N-	2
CF ₃	S	=N-	0
CF ₃	S	=N-	1
CF ₃	S	=N-	2
CH ₂ CF ₃	S	=N-	0
CH ₂ CF ₃	S	=N-	1
CH ₂ CF ₃	S	=N-	2
CH=CH ₂	S	=N-	0
CH=CH ₂	S	=N-	1
CH=CH ₂	S	=N-	2

[0179]

Table 16

R^1	A^1	A^3	n
$CH_2CH=CH_2$	S	$=N-$	0
$CH_2CH=CH_2$	S	$=N-$	1
$CH_2CH=CH_2$	S	$=N-$	2
$C\equiv CH$	S	$=N-$	0
$C\equiv CH$	S	$=N-$	1
$C\equiv CH$	S	$=N-$	2
$CH_2C\equiv CH$	S	$=N-$	0
$CH_2C\equiv CH$	S	$=N-$	1
$CH_2C\equiv CH$	S	$=N-$	2
CyPr	S	$=N-$	0
CyPr	S	$=N-$	1
CyPr	S	$=N-$	2
Me	S	$=CH-$	0
Me	S	$=CH-$	1
Me	S	$=CH-$	2
Et	S	$=CH-$	0
Et	S	$=CH-$	1
Et	S	$=CH-$	2
Pr	S	$=CH-$	0
Pr	S	$=CH-$	1
Pr	S	$=CH-$	2
iPr	S	$=CH-$	0
iPr	S	$=CH-$	1
iPr	S	$=CH-$	2

[0180]

Table 17

R^1	A^1	A^3	n
tBu	S	=CH-	0
tBu	S	=CH-	1
tBu	S	=CH-	2
CF ₃	S	=CH-	0
CF ₃	S	=CH-	1
CF ₃	S	=CH-	2
CH ₂ CF ₃	S	=CH-	0
CH ₂ CF ₃	S	=CH-	1
CH ₂ CF ₃	S	=CH-	2
CH=CH ₂	S	=CH-	0
CH=CH ₂	S	=CH-	1
CH=CH ₂	S	=CH-	2
CH ₂ CH=CH ₂	S	=CH-	0
CH ₂ CH=CH ₂	S	=CH-	1
CH ₂ CH=CH ₂	S	=CH-	2
C≡CH	S	=CH-	0
C≡CH	S	=CH-	1
C≡CH	S	=CH-	2
CH ₂ C≡CH	S	=CH-	0
CH ₂ C≡CH	S	=CH-	1
CH ₂ C≡CH	S	=CH-	2
CyPr	S	=CH-	0
CyPr	S	=CH-	1
CyPr	S	=CH-	2

[0181]

Table 18

R ¹	A ¹	A ³	n
Me	S	=C B r -	0
Me	S	=C B r -	1
Me	S	=C B r -	2
E t	S	=C B r -	0
E t	S	=C B r -	1
E t	S	=C B r -	2
P r	S	=C B r -	0
P r	S	=C B r -	1
P r	S	=C B r -	2
i P r	S	=C B r -	0
i P r	S	=C B r -	1
i P r	S	=C B r -	2
t B u	S	=C B r -	0
t B u	S	=C B r -	1
t B u	S	=C B r -	2
CF ₃	S	=C B r -	0
CF ₃	S	=C B r -	1
CF ₃	S	=C B r -	2
CH ₂ CF ₃	S	=C B r -	0
CH ₂ CF ₃	S	=C B r -	1
CH ₂ CF ₃	S	=C B r -	2
CH=CH ₂	S	=C B r -	0
CH=CH ₂	S	=C B r -	1
CH=CH ₂	S	=C B r -	2

[0182]

Table 19

R ¹	A ¹	A ³	n
CH ₂ CH=CH ₂	S	=CB r -	0
CH ₂ CH=CH ₂	S	=CB r -	1
CH ₂ CH=CH ₂	S	=CB r -	2
C≡CH	S	=CB r -	0
C≡CH	S	=CB r -	1
C≡CH	S	=CB r -	2
CH ₂ C≡CH	S	=CB r -	0
CH ₂ C≡CH	S	=CB r -	1
CH ₂ C≡CH	S	=CB r -	2
CyPr	S	=CB r -	0
CyPr	S	=CB r -	1
CyPr	S	=CB r -	2

[0183]

In the above Tables 1 to 19, Me represents a methyl group, Et represents an ethyl group, Pr represents a propyl group, iPr represents an isopropyl group, tBu represents a tert-butyl group, and CyPr represents a cyclopropyl group.

[0184]

A compound of the formula (A) wherein R² is a fluorine atom, R³, R⁴, and R⁵ are a hydrogen atom, and R¹, A¹, A³, and n are a combination shown in Tables 1-19.

[0185]

A compound of the formula (A) wherein R², R⁴, and R⁵ are a hydrogen atom, R³ is a fluorine atom, and R¹, A¹, A³, and n are a combination shown in Tables 1-19.

[0186]

A compound of the formula (A) wherein R², R³, and R⁵

are a hydrogen atom, R^4 is a fluorine atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0187]

A compound of the formula (A) wherein R^2 , R^3 , and R^4 are a hydrogen atom, R^5 is a fluorine atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0188]

A compound of the formula (A) wherein R^2 is a chlorine atom, R^3 , R^4 , and R^5 are a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0189]

A compound of the formula (A) wherein R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a chlorine atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

15 [0190]

A compound of the formula (A) wherein R^2 , R^3 , and R^5 are a hydrogen atom, R^4 is a chlorine atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0191]

20 A compound of the formula (A) wherein R^2 , R^3 , and R^4 are a hydrogen atom, R^5 is a chlorine atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0192]

25 A compound of the formula (A) wherein R^2 is a methyl group, R^3 , R^4 , and R^5 are a hydrogen atom, and R^1 , A^1 , A^3 ,

and n are a combination shown in Tables 1-19.

[0193]

A compound of the formula (A) wherein R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a methyl group, and R^1 , A^1 , A^3 ,
5 and n are a combination shown in Tables 1-19.

[0194]

A compound of the formula (A) wherein R^2 , R^3 , and R^5 are a hydrogen atom, R^4 is a methyl group, and R^1 , A^1 , A^3 ,
and n are a combination shown in Tables 1-19.

10 [0195]

A compound of the formula (A) wherein R^2 , R^3 , and R^4 are a hydrogen atom, R^5 is a methyl group, and R^1 , A^1 , A^3 ,
and n are a combination shown in Tables 1-19.

[0196]

15 A compound of the formula (A) wherein R^2 is a trifluoromethyl group, R^3 , R^4 , and R^5 are a hydrogen atom,
and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0197]

20 A compound of the formula (A) wherein R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a trifluoromethyl group, and R^1 ,
 A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0198]

25 A compound of the formula (A) wherein R^2 , R^3 , and R^5 are a hydrogen atom, R^4 is a trifluoromethyl group, and R^1 ,
 A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0199]

A compound of the formula (A) wherein R^2 , R^3 , and R^4 are a hydrogen atom, R^5 is a trifluoromethyl group, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

5 [0200]

A compound of the formula (A) wherein R^2 is a bromine atom, R^3 , R^4 , and R^5 are a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0201]

10 A compound of the formula (A) wherein R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a bromine atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0202]

15 A compound of the formula (A) wherein R^2 , R^3 , and R^5 are a hydrogen atom, R^4 is a bromine atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0203]

20 A compound of the formula (A) wherein R^2 , R^3 and R^4 are a hydrogen atom, R^5 is a bromine atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0204]

A compound of the formula (A) wherein R^2 is a pentafluoroethyl group, R^3 , R^4 , and R^5 are a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

25 [0205]

A compound of the formula (A) wherein R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a pentafluoroethyl group, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0206]

5 A compound of the formula (A) wherein R^2 , R^3 , and R^5 are a hydrogen atom, R^4 is a pentafluoroethyl group, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0207]

10 A compound of the formula (A) wherein R^2 , R^3 and R^4 are a hydrogen atom, R^5 is a pentafluoroethyl group, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0208]

15 A compound of the formula (A) wherein R^2 is a trifluoromethoxy group, R^3 , R^4 , and R^5 are a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0209]

A compound of the formula (A) wherein R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a trifluoromethoxy group, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

20 [0210]

A compound of the formula (A) wherein R^2 , R^3 , and R^5 are a hydrogen atom, R^4 is a trifluoromethoxy group, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0211]

25 A compound of the formula (A) wherein R^2 , R^3 , and R^4

are a hydrogen atom, R^5 is a trifluoromethoxy group, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0212]

A compound of the formula (A) wherein R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a 2-pyridyl group, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0213]

A compound of the formula (A) wherein R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a 3-chloro-2-pyridyl group, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0214]

A compound of the formula (A) wherein R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a 2-pyrimidyl group, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0215]

A compound of the formula (A) wherein R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is $-SF_5$, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0216]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a hydrogen atom, R^7 is a trifluoromethyl group, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0217]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 ,

R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a pentafluoroethyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0218]

5 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a heptafluoroisopropyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0219]

10 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a fluorine atom, R^6 is a pentafluoroethyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0220]

15 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a chlorine atom, R^6 is a pentafluoroethyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0221]

20 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a bromine atom, R^6 is a pentafluoroethyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0222]

25 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 ,

R⁴, and R⁵ are a hydrogen atom, R³ is a methyl group, R⁶ is a pentafluoroethyl group, R⁷ is a hydrogen atom, and R¹, A¹, A³, and n are a combination shown in Tables 1-19.

[0223]

5 A compound of the formula (A) wherein A² is =CH-, R², R⁴, and R⁵ are a hydrogen atom, R³ is a trifluoromethyl group, R⁶ is a pentafluoroethyl group, R⁷ is a hydrogen atom, and R¹, A¹, A³, and n are a combination shown in Tables 1-19.

10 [0224]

 A compound of the formula (A) wherein A² is =CH-, R², R⁴, and R⁵ are a hydrogen atom, R³ is a pentafluoroethyl group, R⁶ is a pentafluoroethyl group, R⁷ is a hydrogen atom, and R¹, A¹, A³, and n are a combination shown in
15 Tables 1-19.

[0225]

 A compound of the formula (A) wherein A² is =CH-, R², R⁴, and R⁵ are a hydrogen atom, R³ is a trifluoromethoxy group, R⁶ is a pentafluoroethyl group, R⁷ is a hydrogen
20 atom, and R¹, A¹, A³, and n are a combination shown in Tables 1-19.

[0226]

 A compound of the formula (A) wherein A² is =CH-, R², R³, R⁴, and R⁵ are a hydrogen atom, R⁶ is a difluoromethyl
25 group, R⁷ is a hydrogen atom, and R¹, A¹, A³, and n are a

combination shown in Tables 1-19.

[0227]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a trifluoromethoxy group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a
5 combination shown in Tables 1-19.

[0228]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a fluorine atom, R^6 is
10 a trifluoromethoxy group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0229]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a chlorine atom, R^6 is
15 a trifluoromethoxy group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0230]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a bromine atom, R^6 is
20 a trifluoromethoxy group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0231]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a methyl group, R^6 is
25 a trifluoromethoxy group, R^7 is a hydrogen atom, and R^1 , A^1 ,

A³, and n are a combination shown in Tables 1-19.

[0232]

A compound of the formula (A) wherein A² is =CH-, R², R⁴, and R⁵ are a hydrogen atom, R³ is a trifluoromethyl group, R⁶ is a trifluoromethoxy group, R⁷ is a hydrogen atom, and R¹, A¹, A³, and n are a combination shown in Tables 1-19.

[0233]

A compound of the formula (A) wherein A² is =CH-, R², R⁴, and R⁵ are a hydrogen atom, R³ is a pentafluoroethyl group, R⁶ is a trifluoromethoxy group, R⁷ is a hydrogen atom, and R¹, A¹, A³, and n are a combination shown in Tables 1-19.

[0234]

A compound of the formula (A) wherein A² is =CH-, R², R⁴, and R⁵ are a hydrogen atom, R³ is a trifluoromethoxy group, R⁶ is a trifluoromethoxy group, R⁷ is a hydrogen atom, and R¹, A¹, A³, and n are a combination shown in Tables 1-19.

[0235]

A compound of the formula (A) wherein A² is =CH-, R², R³, R⁴, and R⁵ are a hydrogen atom, R⁶ is a trifluoromethylsulfanyl group, R⁷ is a hydrogen atom, and R¹, A¹, A³, and n are a combination shown in Tables 1-19.

[0236]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a fluorine atom, R^6 is a trifluoromethylsulfanyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

5 [0237]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a chlorine atom, R^6 is a trifluoromethylsulfanyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

10 [0238]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a bromine atom, R^6 is a trifluoromethylsulfanyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

15 [0239]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a methyl group, R^6 is a trifluoromethylsulfanyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

20 [0240]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a trifluoromethyl group, R^6 is a trifluoromethylsulfanyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown
25 in Tables 1-19.

[0241]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a pentafluoroethyl group, R^6 is a trifluoromethylsulfanyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0242]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a trifluoromethoxy group, R^6 is a trifluoromethylsulfanyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0243]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a trifluoromethylsulfinyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0244]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a fluorine atom, R^6 is a trifluoromethylsulfinyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0245]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a chlorine atom, R^6 is

a trifluoromethylsulfinyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0246]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 ,
5 R^4 , and R^5 are a hydrogen atom, R^3 is a bromine atom, R^6 is
a trifluoromethylsulfinyl group, R^7 is a hydrogen atom, and
 R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0247]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 ,
10 R^4 , and R^5 are a hydrogen atom, R^3 is a methyl group, R^6 is
a trifluoromethylsulfinyl group, R^7 is a hydrogen atom, and
 R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0248]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 ,
15 R^4 , and R^5 are a hydrogen atom, R^3 is a trifluoromethyl
group, R^6 is a trifluoromethylsulfinyl group, R^7 is a
hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown
in Tables 1-19.

[0249]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 ,
20 R^4 , and R^5 are a hydrogen atom, R^3 is a pentafluoroethyl
group, R^6 is a trifluoromethylsulfinyl group, R^7 is a
hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown
in Tables 1-19.

25 [0250]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a trifluoromethoxy group, R^6 is a trifluoromethylsulfinyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown
5 in Tables 1-19.

[0251]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a trifluoromethylsulfonyl group, R^7 is a hydrogen atom, and
10 R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0252]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a fluorine atom, R^6 is a trifluoromethylsulfonyl group, R^7 is a hydrogen atom, and
15 R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0253]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a chlorine atom, R^6 is a trifluoromethylsulfonyl group, R^7 is a hydrogen atom, and
20 R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0254]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a bromine atom, R^6 is a trifluoromethylsulfonyl group, R^7 is a hydrogen atom, and
25 R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0255]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a methyl group, R^6 is a trifluoromethylsulfonyl group, R^7 is a hydrogen atom, and
5 R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0256]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a trifluoromethyl group, R^6 is a trifluoromethylsulfonyl group, R^7 is a
10 hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0257]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a pentafluoroethyl
15 group, R^6 is a trifluoromethylsulfonyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0258]

A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a trifluoromethoxy
20 group, R^6 is a trifluoromethylsulfonyl group, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0259]

25 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 ,

R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is a bromine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0260]

- 5 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a fluorine atom, R^6 is a bromine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0261]

- 10 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a chlorine atom, R^6 is a bromine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0262]

- 15 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a bromine atom, R^6 is a bromine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0263]

- 20 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a methyl group, R^6 is a bromine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0264]

- 25 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 ,

R^4 , and R^5 are a hydrogen atom, R^3 is a trifluoromethyl group, R^6 is a bromine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0265]

5 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a pentafluoroethyl group, R^6 is a bromine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0266]

10 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a trifluoromethoxy group, R^6 is a bromine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0267]

15 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is an iodine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0268]

20 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a fluorine atom, R^6 is an iodine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0269]

25 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 ,

R^4 , and R^5 are a hydrogen atom, R^3 is a chlorine atom, R^6 is an iodine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0270]

5 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a bromine atom, R^6 is an iodine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0271]

10 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a methyl group, R^6 is an iodine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0272]

15 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a trifluoromethyl group, R^6 is an iodine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0273]

20 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a pentafluoroethyl group, R^6 is an iodine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0274]

25 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 ,

R^4 , and R^5 are a hydrogen atom, R^3 is a trifluoromethoxy group, R^6 is an iodine atom, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0275]

5 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^3 , R^4 , and R^5 are a hydrogen atom, R^6 is $-SF_5$, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0276]

10 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a fluorine atom, R^6 is $-SF_5$, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0277]

15 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a chlorine atom, R^6 is $-SF_5$, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0278]

20 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 , R^4 , and R^5 are a hydrogen atom, R^3 is a bromine atom, R^6 is $-SF_5$, R^7 is a hydrogen atom, and R^1 , A^1 , A^3 , and n are a combination shown in Tables 1-19.

[0279]

25 A compound of the formula (A) wherein A^2 is $=CH-$, R^2 ,

R⁴, and R⁵ are a hydrogen atom, R³ is a methyl group, R⁶ is -SF₅, R⁷ is a hydrogen atom, and R¹, A¹, A³, and n are a combination shown in Tables 1-19.

[0280]

5 A compound of the formula (A) wherein A² is =CH-, R², R⁴, and R⁵ are a hydrogen atom, R³ is a trifluoromethyl group, R⁶ is -SF₅, R⁷ is a hydrogen atom, and R¹, A¹, A³, and n are a combination shown in Tables 1-19.

[0281]

10 A compound of the formula (A) wherein A² is =CH-, R², R⁴, and R⁵ are a hydrogen atom, R³ is a pentafluoroethyl group, R⁶ is -SF₅, R⁷ is a hydrogen atom, and R¹, A¹, A³, and n are a combination shown in Tables 1-19.

[0282]

15 A compound of the formula (A) wherein A² is =CH-, R², R⁴, and R⁵ are a hydrogen atom, R³ is a trifluoromethoxy group, R⁶ is -SF₅, R⁷ is a hydrogen atom, and R¹, A¹, A³, and n are a combination shown in Tables 1-19.

[0283]

20 Examples of pests against which the present compound has an activity include noxious arthropods such as noxious insects and noxious acarines, and nematodes. In particular, examples of the pests include the following.

[0284]

25 Hemiptera: Delphacidae such as *Laodelphax striatellus*,

Nilaparvata lugens, and Sogatella furcifera;
Deltocephalidae such as Nephotettix cincticeps, Nephotettix
virescens, and Empoasca onukii; Aphididae such as Aphis
gossypii, Myzus persicae, Brevicoryne brassicae, Aphis
5 spiraecola, Macrosiphum euphorbiae, Aulacorthum solani,
Rhopalosiphum padi, Toxoptera citricidus, and Hyalopterus
pruni; Pentatomidae such as Nezara antennata, Riptortus
clavetus, Leptocorisa chinensis, Eysarcoris parvus, and
Halyomorpha mista; Aleyrodidae such as Trialeurodes
10 vaporariorum, Bemisia tabaci, Dialeurodes citri, and
Aleurocanthus spiniferus; Coccidae such as Aonidiella
aurantii, Comstockaspis perniciosus, Unaspis citri,
Ceroplastes rubens, Icerya purchasi, Planococcus kraunhiae,
Pseudococcus longispinis, and Pseudaulacaspis pentagona;
15 Tingidae; Cimices such as Cimex lectularius; and Psyllidae.
[0285]

Lepidoptera: Pyralidae such as Chilo suppressalis,
Tryporyza incertulas, Cnaphalocrocis medinalis, Notarcha
derogata, Plodia interpunctella, Ostrinia furnacalis,
20 Hellula undalis, and Pediasia teterrellus; Noctuidae such
as Spodoptera litura, Spodoptera exigua, Pseudaletia
separata, Mamestra brassicae, Agrotis ipsilon, Plusia
nigrisigna, Thoricoplusia spp., Heliothis spp., and
Helicoverpa spp.; Pieridae such as Pieris rapae;
25 Tortricidae such as Adoxophyes spp., Grapholita molesta,

Leguminivora glycinivorella, Matsumuraeses azukivora,
Adoxophyes orana fasciata, Adoxophyes honmai., Homona
magnanima, Archips fuscocupreanus, and Cydia pomonella;
Gracillariidae such as Caloptilia theivora and
5 Phyllonorycter ringoneella; Carposinidae such as Carposina
niponensis; Lyonetiidae such as Lyonetia spp.; Lymantriidae
such as Lymantria spp. and Euproctis spp; Yponomeutidae
such as Plutella xylostella; Gelechiidae such as
Pectinophora gossypiella and Phthorimaea operculella;
10 Arctiidae such as Hyphantria cunea; and Tineidae such as
Tinea translucens and Tineola bisselliella.

[0286]

Thysanoptera: Thripidae such as Frankliniella
occidentalis, Thrips palmi, Scirtothrips dorsalis, Thrips
15 tabaci, and Frankliniella intonsa.

[0287]

Diptera: Culices such as Culex pipiens pallens, Culex
tritaeniorhynchus, and Culex quinquefasciatus; Aedes spp.
such as Aedes aegypti and Aedes albopictus; Anopheles spp.
20 such as Anopheles sinensis; Chironomidae; Muscidae such as
Musca domestica and Muscina stabulans; Calliphoridae;
Sarcophagidae; Fanniidae; Anthomyiidae such as Delia
platura and Delia antiqua; Agromyzidae such as Agromyza
oryzae, Hydrellia griseola, Liriomyza sativae, Liriomyza
25 trifolii, and Chromatomyia horticola; Chloropidae such as

Chlorops oryzae; Tephritidae such as Dacus cucurbitae and
Ceratitis capitata; Drosophilidae; Phoridae such as
Megaselia spiracularis; Psychodidae such as Clogmia
albipunctata; Simuliidae; Tabanidae such as Tabanus
5 trigonus; and stable flies.

[0288]

Coleoptera: Diabrotica spp. such as Diabrotica
virgifera virgifera and Diabrotica undecimpunctata howardi;
Scarabaeidae such as Anomala cuprea, Anomala rufocuprea,
10 and Popillia japonica; weevils such as Sitophilus zeamais,
Lissorhoptrus oryzophilus, Callosobruchus chienensis,
Echinocnemus squameus, Anthonomus grandis, and Sphenophorus
venatus; Tenebrionidae such as Tenebrio molitor and
Tribolium castaneum; Chrysomelidae such as Oulema oryzae,
15 Aulacophora femoralis, Phyllotreta striolata, and
Leptinotarsa decemlineata; Dermestidae such as Anthrenus
verbasci and Dermestes maculatus; Anobiidae such as
Lasioderma serricorne; Epilachna such as Epilachna
vigintioctopunctata; Scolytidae such as Lyctus brunneus and
20 Tomicus piniperda; Bostrychidae; Ptinidae; Cerambycidae
such as Anoplophora malasiaca; Agriotes spp., and Paederus
fuscipes.

[0289]

Orthoptera: Locusta migratoria, Gryllotalpa africana,
25 Oxya yezoensis, Oxya japonica, and Gryllidae.

[0290]

Hymenoptera: Formicidae such as *Monomorium pharaonis*,
Formica fusca japonica, *Ochetellus glaber*, *Pristomyrmex*
pungens, *Pheidole noda*, *Acromyrmex* spp., and *Solenopsis*
5 spp.; Vespidae; Betylidae; and Tenthredinidae such as
Athalia rosae and *Athalia japonica*.

[0291]

Nematoda: *Aphelenchoides besseyi*, *Nothotylenchus acris*,
Meloidogyne incognita, *Meloidogyne hapla*, *Meloidogyne*
10 *javanica*, *Heterodera glycines*, *Globodera rostochiensis*,
Pratylenchus coffeae, and *Pratylenchus neglectus*.

[0292]

Blattodea: *Blattella germanica*, *Periplaneta fuliginosa*,
Periplaneta americana, *Periplaneta brunnea*, and *Blatta*
15 *orientalis*.

[0293]

Acarina: Tetranychidae such as *Tetranychus urticae*,
Tetranychus kanzawai, *Panonychus citri*, *Panonychus ulmi*,
and *Oligonychus* spp.; Eriophyidae such as *Aculops pelekassi*,
20 *Phyllocoptruta citri*, *Aculops lycopersici*, *Calacarus*
carinatus, *Acaphylla theavagrans*, *Eriophyes chibaensis*, and
Aculus schlechtendali; Tarsonemidae such as
Polyphagotarsonemus latus; Tenuipalpidae such as
Brevipalpus phoenicis; Tuckerellidae; Ixodidae such as
25 *Haemaphysalis longicornis*, *Haemaphysalis flava*, *Dermacentor*

taiwanicus, Ixodes ovatus, Ixodes persulcatus, Ixodes
scapularis, Boophilus microplus, and Rhipicephalus
sanguineus; Acaridae such as Tyrophagus putrescentiae and
Tyrophagus similis; Pyroglyphidae such as Dermatophagoides
5 farinae and Dermatophagoides pteromyssus; Cheyletidae such
as Cheyletus eruditus, Cheyletus malaccensis, and Cheyletus
moorei; Dermanyssidae such as Ornithonyssus bacoti,
Ornithonyssus sylvarum, and Dermanyssus gallinae;
Trombiculidae such as Leptotrombidium akamushi; and Araneae
10 such as Chiracanthium japonicum and Latrodectus hasseltii.
[0294]

The pest controlling agent of the present invention
contains the present compound and an inert carrier.
Generally, the pest controlling agent of the present
15 invention is a formulation such as an emulsion, an oil
solution, a powder, a granule, a wettable powder, a
flowable formulation, a microcapsule, an aerosol, a smoking
agent, a poison bait, and a resin formulation which are
obtained by mixing the present compound and an inert
20 carrier such as a solid carrier, a liquid carrier and a
gaseous carrier, and further adding a surfactant and other
adjuvant for formulation, if necessary.

The pest controlling agent of the present invention
usually contains the present compound in an amount of
25 0.01 % to 95 % by weight.

[0295]

Examples of the solid carrier to be used for formulation include a fine power and a granule of clays (such as kaolin clay, diatomite, bentonite, Fubasami clay, and acid clay), synthetic hydrated silicon oxide, talc, ceramic, other inorganic minerals (such as sericite, quartz, sulfur, activated carbon, calcium carbonate, and hydrated silica), and chemical fertilizers (such as ammonium sulfate, ammonium phosphate, ammonium nitrate, urea, and ammonium chloride) as well as synthetic resins (such as polypropylene, polyacrylonitrile, polyester resins such as methyl polymethacrylate and polyethylene terephthalate, nylon resins such as nylon-6, nylon-11, and nylon-66, polyamide resins, polyvinyl chloride, polyvinylidene chloride, and vinyl chloride-propylene copolymer).

[0296]

Examples of the liquid carrier include water, alcohols (such as methanol, ethanol, isopropyl alcohol, butanol, hexanol, benzyl alcohol, ethylene glycol, propylene glycol, and phenoxyethanol), ketones (such as acetone, methyl ethyl ketone, and cyclohexanone), aromatic hydrocarbons (such as toluene, xylene, ethylbenzene, dodecylbenzene, phenylxylylethane, and methylnaphthalene), aliphatic hydrocarbons (such as hexane, cyclohexane, kerosine, and light oil), esters (such as ethyl acetate, butyl acetate,

isopropyl mylistate, ethyl oleate, diisopropyl adipate, diisobutyl adipate, and propyleneglycol monomethyl ether acetate), nitriles (such as acetonitrile and isobutyronitrile), ethers (such as diisopropyl ether, 1,4-dioxane, ethylene glycol dimethyl ether, diethylene glycol dimethyl ether, diethylene glycol monomethyl ether, propylene glycol monomethyl ether, dipropylene glycol monomethyl ether, and 3-methoxy-3-methyl-1-butanol), acid amides (such as N,N-dimethylformamide and N,N-dimethylacetamide), halogenated hydrocarbons (such as dichloromethane, trichloroethane, and tetrachlorocarbon), sulfoxides (such as dimethylsulfoxide), propylene carbonate, and vegetable oils (such as soy bean oil and cotton seed oil).

[0297]

Examples of the gaseous carrier include fluorocarbons, butane gas, liquefied petroleum gas (LPG), dimethyl ether, and carbon dioxide.

[0298]

Examples of the surfactant include nonionic surfactant such as polyoxyethylene alkyl ether, polyoxyethylene alkylaryl ether, and polyethyleneglycol fatty acid ester; and anionic surfactant such as alkylsulfonic acid salts, alkylbenzenesulfonic acid salts, and alkylsulfuric acid salts.

[0299]

Examples of the other adjuvant for formulation include binders, dispersants, colorants and stabilizers, and particularly for example, casein, gelatin, polysaccharides (such as starch, gum arabic, cellulose derivatives, and alginic acid), lignin derivatives, synthetic water-soluble polymers (such as polyvinyl alcohol, polyvinylpyrrolidone, and polyacrylic acid), PAP (acidic isopropyl phosphate), BHT (2,6-di-t-butyl-4-methylphenol), and BHA (a mixture of 2-t-butyl-4-methoxyphenol and 3-t-butyl-4-methoxyphenol).

10 [0300]

The method for controlling pests of the present invention is applying an effective amount of the present compound to pests directly and/or habitats of pests (such as plant, soil, indoor, and in-body of animals). The present compound is usually used as the pest controlling agent of the present invention in the method for controlling pests of the present invention.

[0301]

When the pest controlling agent of the present invention is used for a control of pests in agriculture, the application amount is usually 1 to 10,000 g as the present compound per 10,000 m². When the pest controlling agent of the present invention is a formulation of emulsions, wettable powders or flowables, they are usually applied after a dilution with water to have an active

ingredient concentration of 0.01 to 10,000 ppm. When the pest controlling agent of the present invention is a formulation of granules or powders, they are usually applied as such.

5 [0302]

The formulations and the dilute aqueous solutions of the formulation may be sprayed directly to the plant to be protected from pests, or may be applied to the soil to control the pests living in a soil.

10 [0303]

Furthermore, the resin formulations of sheets or strip form can be applied by a method such as winding around plants, stretching in the vicinity of plants, and laying on the soil surface at the plant bottom.

15 [0304]

When the pest controlling agent of the present invention is used for a control of pests in indoor, the application amount is usually 0.01 to 1,000 mg as the present compound per 1 m² in case of application for plane surface, and 0.01 to 500 mg as the present compound per 1 m³ in case of application for space. When the pest controlling agent of the present invention is a formulation of emulsions, wettable powders or flowables, they are usually applied after a dilution with water to have an active ingredient concentration of 0.1 to 1,000 ppm. When

20

25

the pest controlling agent of the present invention is a formulation of oil solutions, aerosols, smoking agents and poison baits, they are usually applied as such.

[0305]

5 The pest controlling agent of the present invention could be used in farmlands on which "crops" shown below are cultivated.

 Agricultural crops: corn, rice, wheat, barley, rye, oat, sorghum, cotton, soybean, peanut, sarrazin, sugar beet,
10 rapeseed, sunflower, sugar cane, and tobacco;

 Vegetables: Solanaceae vegetables (such as eggplant, tomato, green pepper, hot pepper, and potato), Cucurbitaceae vegetables (such as cucumber, pumpkin, zucchini, watermelon, and melon), Cruciferae vegetables
15 (such as Japanese radish, turnip, horseradish, kohlrabi, Chinese cabbage, cabbage, brown mustard, broccoli, and cauliflower), Compositae vegetables (such as burdock, garland chrysanthemum, artichoke, and lettuce), Liliaceae vegetables (such as Welsh onion, onion, garlic, and
20 asparagus), Umbelliferae vegetables (such as carrot, parsley, celery, and parsnip), Chenopodiaceae vegetables (such as spinach, and Swiss chard), Labiatae vegetables (such as Japanese basil, mint, and basil), strawberry, sweat potato, yam, and aroid;

25 Fruit trees: pomaceous fruits (such as apple, common

pear, Japanese pear, Chinese quince, and quince), stone
fleshy fruits (such as peach, plum, nectarine, Japanese
plum, cherry, apricot, and prune), citrus plants (such as
Satsuma mandarin, orange, lemon, lime, and grapefruit),
5 nuts (such as chestnut, walnut, hazel nut, almond,
pistachio, cashew nut, and macadamia nut), berry fruits
(such as blueberry, cranberry, blackberry, and raspberry),
grape, persimmon, olive, loquat, banana, coffee, date,
coconut palm, and oil palm;

10 Trees other fruit trees: tea, mulberry, flowering
trees (such as azalea, japonica, hydrangea, sasanqua,
lillicium anisatum, cherry tree, tulip poplar, crepe myrtle,
and orange osmanthus), street trees (such as ash tree,
birch, dogwood, eucalyptus, ginkgo, lilac, maple tree, oak,
15 poplar, cercis, Chinese sweet gum, plane tree, zelkova,
Japanese arborvitae, fir tree, Japanese hemlock, needle
juniper, pine, spruce, yew, elm, and horse-chestnut), sweet
viburnum, Podocarpus macrophyllus, Japanese cedar, Japanese
cypress, croton, spindle tree, and Chinese hawthorn.

20 Lawn: zoysia (such as Japanese lawn grass and
mascarene grass), Bermuda grass (such as Cynodon dactylon),
bent grass (such as creeping bent grass, Agrostis
stolonifera, and Agrostis tenuis), bluegrass (such as
Kentucky bluegrass and rough bluegrass), fescue (such as
25 tall fescue, chewing fescue, and creeping fescue), ryegrass

(such as darnel and perennial ryegrass), cocksfoot, and timothy grass;

Others: flowers (such as rose, carnation, chrysanthemum, Eustoma grandiflorum Shinners, gypsophila, gerbera, pot marigold, salvia, petunia, verbena, tulip, aster, gentian, lily, pansy, cyclamen, orchid, lily of the valley, lavender, stock, ornamental kale, primula, poinsettia, gladiolus, cattleya, daisy, cymbidium, and begonia), biofuel plants (such as Jatropha, curcas, safflower, Camelina alyssum, switchgrass, miscanthus, reed canary grass, Arundo donax, kenaf, cassava, willow, and algae), and foliage plant.

[0306]

The "crops" include genetically modified crops.

[0307]

The pest controlling agent of the present invention can be used as a mixture with or together with other insecticides, acaricides, nematocides, fungicides, plant growth regulators, herbicides, and synergists. Examples of active ingredients of the insecticide, the acaricide, the nematocide, the fungicide, the herbicide, and the synergist are shown below.

[0308]

Active ingredients of insecticides:

(1) Organic phosphorus compounds:

Acephate, Aluminium phosphide, butathiofos, cadusafos, chlorethoxyfos, chlorfenvinphos, chlorpyrifos, chlorpyrifos-methyl, cyanophos: CYAP, DCIP (diazinon, dichlorodiisopropyl ether), dichlofenthion: ECP, 5 dichlorvos: DDVP, dimethoate, dimethylvinphos, disulfoton, EPN, ethion, ethoprophos, etrimfos, fenthion: MPP, fenitrothion: MEP, fosthiazate, formothion, Hydrogen phosphide, isofenphos, isoxathion, malathion, mesulfenfos, methidathion: DMTP, monocrotophos, naled: BRP, oxydeprofos: 10 ESP, parathion, phosalone, phosmet: PMP, pirimiphos-methyl, pyridafenthion, quinalphos, phenthoate: PAP, profenofos, propaphos, prothiofos, pyraclorfos, salithion, sulprofos, tebupirimfos, temephos, tetrachlorvinphos, terbufos, thiometon, trichlorphon: DEP, vamidothion, phorate, and 15 cadusafos.

(2) Carbamate compounds:

Alanycarb, bendiocarb, benfuracarb, BPMC, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb, furathiocarb, 20 isoproc carb: MIPC, metolcarb, methomyl, methiocarb, NAC, oxamyl, pirimicarb, propoxur: PHC, XMC, thiodicarb, xylylcarb, and aldicarb.

(3) Pyrethroid compounds:

Acrinathrin, allethrin, benfluthrin, beta-cyfluthrin, 25 bifenthrin, cycloprothrin, cyfluthrin, cyhalothrin,

cypermethrin, deltamethrin, esfenvalerate, ethofenprox,
fenpropathrin, fenvalerate, flucythrinate, flufenoprox,
flumethrin, fluvalinate, halfenprox, imiprothrin,
permethrin, prallethrin, pyrethrins, resmethrin, sigma-
5 cypermethrin, silafluofen, tefluthrin, tralomethrin,
transfluthrin, tetramethrin, phenothrin, cyphenothrin,
alpha-cypermethrin, zeta-cypermethrin, lambda-cyhalothrin,
gamma-cyhalothrin, furamethrin, tau-fluvalinate,
metofluthrin, profluthrin, dimefluthrin, 2,3,5,6-
10 tetrafluoro-4-(methoxymethyl)benzyl (EZ)-(1RS,3RS;1RS,3SR)-
2,2-dimethyl-3-prop-1-enylcyclopropanecarboxylate, 2,3,5,6-
tetrafluoro-4-methylbenzyl (EZ)-(1RS,3RS;1RS,3SR)-2,2-
dimethyl-3-prop-1-enylcyclopropanecarboxylate, and 2,3,5,6-
tetrafluoro-4-(methoxymethyl)benzyl (1RS,3RS;1RS,3SR)-2,2-
15 dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate.

(4) Nereistoxin compounds:

Cartap, bensultap, thiocyclam, monosultap, and
bisultap.

(5) Neonicotinoid compounds:

20 Imidacloprid, nitenpyram, acetamiprid, thiamethoxam,
thiacloprid, dinotefuran, and clothianidin.

(6) Benzoylurea compounds:

Chlorfluazuron, bistrifluron, diafenthiuron,
diflubenzuron, fluazuron, flucycloxuron, flufenoxuron,
25 hexaflumuron, lufenuron, novaluron, noviflumuron,

teflubenzuron, triflumuron, and triazuron.

(7) Phenylpyrazole compounds:

Acetoprole, ethiprole, fipronil, vaniliprole, pyriprole, and pyrafluprole.

5 (8) Bt toxins:

Live spores derived from and crystal toxins produced from *Bacillus thuringiensis* and a mixture thereof;

(9) Hydrazine compounds:

10 Chromafenozide, halofenozide, methoxyfenozide, and tebufenozide.


(10) Organic chlorine compounds:

Aldrin, dieldrin, dieldrin, endosulfan, and methoxychlor.

(11) Other insecticidal active ingredients:

15 Mmachine oil, nicotine-sulfate; avermectin-B, bromopropylate, buprofezin, chlorphenapyr, cyantraniliprole, cyromazine, D-D (1,3-Dichloropropene), emamectin-benzoate, fenazaquin, flupyrzofos, hydroprone, methoprene, indoxacarb, metoxadiazole, milbemycin-A, pymetrozine, 20 pyridalyl, pyriproxyfen, spinosad, sulfluramid, tolfenpyrad, triazamate, flubendiamide, lepimectin, arsenic acid, benclothiaz, calcium cyanamide, calcium polysulfide, chlordane, DDT, DSP, flufenimer, flonicamid, flurimfen, formetanate, metam-ammonium, metam-sodium, Methyl bromide, 25 Potassium oleate, prothifenebut, spiromesifen, sulfoxaflor,

sulfur, metaflumizone, spirotetramat, pyrifluquinazone, spinetoram, chlorantraniliprole, tralopyril, cyantraniliprole, a compound of the following formula (K):




5 wherein:

R^{100} is chlorine, bromine, or a trifluoromethyl group,

R^{200} is chlorine, bromine, or a methyl group, and

R^{300} is chlorine, bromine, or a cyano group,

and a compound of the following formula (L):



10

wherein:

R^{1000} is chlorine, bromine, or iodine.

[0309]

Active ingredients of acardides:

15 Acequinocyl, amitraz, benzoximate, bifenaate, bromopropylate, chinomethionat, chlorobenzilate, CPCBS (chlorfenson), clofentezine, cyflumetofen, Kelthane

(dicofol), etoxazole, fenbutatin oxide, fenothiocarb, fenpyroximate, fluacrypyrim, fluproxyfen, hexythiazox, propargite: BPPS, polynactins, pyridaben, Pyrimidifen, tebufenpyrad, tetradifon, spirodiclofen, spiromesifen, 5 spirotetramat, amidoflumet, and cyenopyrafen.

[0310]

Active ingredients of nematocides:

DCIP, fosthiazate, levamisole hydrochloride (levamisole), methyisothiocyanate, morantel tartarate, and 10 imicyafos.

[0311]

Active ingredients of fungicides:

Azole fungicidal compounds such as propiconazole, prothioconazole, triadimenol, prochloraz, penconazole, 15 tebuconazole, flusilazole, diniconazole, bromuconazole, epoxiconazole, difenoconazole, cyproconazole, metconazole, triflumizole, tetraconazole, myclobutanil, fenbuconazole, hexaconazole, fluquinconazole, triticonazole, bitertanol, imazalil, and flutriafol;

20 Cyclic amine fungicidal compounds such as fenpropimorph, tridemorph, and fenpropidin;

Benzimidazole fungicidal compounds such as carbendazim, benomyl, thiabendazole, and thiophanate-methyl;

Procymidone; cyprodinil; pyrimethanil; diethofencarb; 25 thiuram; fluazinam; mancozeb; iprodione; vinclozolin;

chlorothalonil; captan; mepanipyrim; fenpiclonil;
fludioxonil; dichlofluanid; folpet; kresoxim-methyl;
azoxystrobin; trifloxystrobin; fluoxastrobin;
picoxystrobin; pyraclostrobin; dimoxystrobin; pyribencarb;
5 spiroxamine; quinoxyfen; fenhexamid; famoxadone;
fenamidone; zoxamide; ethaboxam; amisulbrom; iprovalicarb;
benthiavalicarb; cyazofamid; mandipropamid; boscalid;
penthioopyrad; metrafenone; fluopiran; bixafen;
cyflufenamid; proquinazid; isotianil, and tiadinil.

10 [0312]

Active ingredients of herbicides:

(1) Phenoxyfatty acid herbicidal compounds:

2,4-PA, MCP, MCPB, phenothiol, mecoprop, fluroxypyr,
triclopyr, clomeprop, and naproanilide.

15 (2) Benzoic acid herbicidal compounds:

2,3,6-TBA, dicamba, clopyralid, picloram, aminopyralid,
quinclorac, and quinmerac.

(3) Urea herbicidal compounds:

20 Diuron, linuron, chlortoluron, isoproturon,
fluometuron, isouron, tebuthiuron, methabenzthiazuron,
cumyluron, daimuron, and methyl-daimuron.

(4) Triazine herbicidal compounds:

25 Atrazine, ametryn, cyanazine, simazine, propazine,
simetryn, dimethametryn, prometryn, metribuzin, triaziflam,
and indaziflam.

(5) Bipyridinium herbicidal compounds:

Paraquat and diquat.

(6) Hydroxybenzonitrile herbicidal compounds:

Bromoxynil and ioxynil.

5 (7) Dinitroaniline herbicidal compounds:

Pendimethalin, prodiamine, and trifluralin.

(8) Organic phosphorus herbicidal compounds:

Amiprofos-methyl, butamifos, bensulide, piperophos,
anilofos, glyphosate, glufosinate, glufosinate-P, and
10 bialaphos.

(9) Carbamate herbicidal compounds:

Di-allate, tri-allate, EPTC, butylate, benthocarb,
esprocarb, molinate, dimepiperate, swep, chlorpropham,
phenmedipham, phenisopham, pyributicarb, and asulam.

15 (10) Acid amide herbicidal compounds:

Propanil, propyzamide, bromobutide, and etobenzanid.

(11) Chloroacetanilide herbicidal compounds:

Acetochlor, alachlor, butachlor, dimethenamid,
propachlor, metazachlor, metolachlor, pretilachlor,
20 thenylchlor, and pethoxamid.

(12) Diphenylether herbicidal compounds:

Acifluorfen-sodium, bifenox, oxyfluorfen, lactofen,
fomesafen, chlomethoxynil, and aclonifen.

(13) Cyclic imide herbicidal compounds:

25 Oxadiazon, cinidon-ethyl, carfentrazone-ethyl,

surfentrazone, flumiclorac-pentyl, flumioxazin, pyraflufen-ethyl, oxadiargyl, pentoxazone, fluthiacet-methyl, butafenacil, benzfendizone, bencarbazine, and saflufenacil.

(14) Pyrazole herbicidal compounds:

5 Benzofenap, pyrazolate, pyrazoxyfen, topramezone, and pyrasulfotole.

(15) Triketone herbicidal compounds:

 Isoxaflutole, benzobicyclon, sulcotrione, mesotrione, tembotrione, and tefuryltrione.

10 (16) Aryloxyphenoxypropionic acid herbicidal compounds:

 Clodinafop-propargyl, cyhalofop-butyl, diclofop-methyl, fenoxaprop-ethyl, fluazifop-butyl, haloxyfop-methyl, quizalofop-ethyl, and metamifop.

(17) Trioneoxime herbicidal compounds:

15 Alloxydim-sodium, sethoxydim, butroxydim, clethodim, cloproxydim, cycloxydim, tepraloxym, tralkoxydim, and profoxydim.

(18) Sulfonylurea herbicidal compounds:

 Chlorsulfuron, sulfometuron-methyl, metsulfuron-methyl,
20 chlorimuron-ethyl, tribenuron-methyl, triasulfuron,
 bensulfuron-methyl, thifensulfuron-methyl, pyrazosulfuron-ethyl, primisulfuron-methyl, nicosulfuron, amidosulfuron, cinosulfuron, imazosulfuron, rimsulfuron, halosulfuron-methyl, prosulfuron, ethametsulfuron-methyl,
25 triflusulfuron-methyl, flazasulfuron, cyclosulfamuron,

flupyrsulfuron, sulfosulfuron, azimsulfuron, ethoxysulfuron, oxasulfuron, iodosulfuron-methyl-sodium, foramsulfuron, mesosulfuron-methyl, trifloxysulfuron, tritosulfuron, orthosulfamuron, flucetosulfuron, and propyrisulfuron.

5 (19) Imidazolinone herbicidal compounds:

Imazamethabenz-methyl, imazamethapyr, imazamox, imazapyr, imazaquin, and imazethapyr.

(20) Sulfonamide herbicidal compounds:

10 Flumetsulam, metosulam, diclosulam, florasulam, cloransulam-methyl, penoxsulam, and pyroxsulam.

(21) Pyrimidinyloxybenzoic acid herbicidal compounds:

Pyrithiobac-sodium, bispyribac-sodium, pyriminobac-methyl, pyribenzoxim, pyriftalid, and pyrimisulfan.

(22) Other herbicidal compounds:

15 Bentazon, bromacil, terbacil, chlorthiamid, isoxaben, dinoseb, amitrole, cinmethylin, tridiphan, dalapon, diflufenzopyr-sodium, dithiopyr, thiazopyr, flucarbazone-sodium, propoxycarbazine-sodium, mefenacet, flufenacet, fentrazamide, cafenstrole, indanofan, oxaziclonofone,
20 benfuresate, ACN, pyridate, chloridazon, norflurazon, flurtamone, diflufenican, picolinafen, beflubutamid, clomazone, amicarbazone, pinoxaden, pyracidonil, pyroxasulfone, thienicarbazone-methyl, aminocyclopyrachlor, ipfencarbazone, and methiozolin.

25 [0313]

Active ingredients of synergists:

Piperonyl butoxide, sesamex, sulfoxide, N-(2-ethylhexyl)-8,9,10-trinorborn-5-ene-2,3-dicarboxyimide (MGK 264), N-decylimidazole, WARF-antiresistant, TBPT, TPP, IBP,
5 PSCP, methyl iodide (CH_3I), t-phenylbutenone, diethylmaleate, DMC, FDMC, ETP, and ETN.

Examples

[0314]

10 The present invention is described in more detail by Production Examples, Formulation Examples, and Test Examples, but the present invention is not limited to these Examples.

[0315]

15 First, Production Examples of the present compound are shown below.


[0316]

Production Example 1

20 A mixture of N²-methyl-5-trifluoromethylpyridin-2,3-diamine (1.34 g), 2-methylsulfanylbenzaldehyde (1.28 g), sodium hydrogen sulfite (2.19 g), and DMF (14 ml) was stirred with heating at 80 °C for 1 hour, and then heated to 120 °C, and stirred with heating for further 3 hours. Into the reaction mixture cooled to room temperature,
25 saturated aqueous ammonium chloride solution was poured,

and extracted with ethyl acetate 3 times. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give
5 0.74 g of 3-methyl-2-(2-methylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 1).

Present Compound 1



10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.72-8.71 (1H,m), 8.33-8.32 (1H,m), 7.58-7.53 (1H,m), 7.45-7.42 (2H,m), 7.36-7.31 (1H,m), 3.79 (3H,s), 2.44 (3H,s)


[0317]

Production Example 2

15 To a mixture of 3-methyl-2-(2-methylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.32 g), methanol (4 ml) and water (6 ml), sodium periodate (0.42 g) was added under ice-cooling. The mixture was heated to room temperature, and stirred for 1 hour, and then heated to 40
20 $^{\circ}\text{C}$, and stirred with heating for further 1 hour. Into the reaction mixture cooled to room temperature saturated aqueous sodium hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution were poured, and

extracted with chloroform 2 times. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.32 g of 2-(2-methylsulfinylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 2).

Present Compound 2



¹H-NMR (CDCl₃) δ: 8.76-8.74 (1H,m), 8.38-8.35 (1H,m), 8.33-8.32 (1H,m), 7.89-7.84 (1H,m), 7.73-7.69 (1H,m), 7.62-7.59 (1H,m), 3.92 (3H,s), 3.03 (3H,s)


[0318]

Production Example 3

To a mixture of 3-methyl-2-(2-methylsulfonylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.30 g) and chloroform (9 ml), 69 % of 3-chloroperbenzoic acid (0.70 g) was added under ice-cooling, and then heated to room temperature, and stirred for 2 hours. Then, into the mixture, saturated aqueous sodium hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution were poured, and extracted with chloroform 2 times. The combined organic layer was dried over magnesium sulfate,

and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.32 g of 2-(2-methylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 3).

Present Compound 3




$^1\text{H-NMR}$ (CDCl_3) δ : 8.76-8.74 (1H,m), 8.31-8.27 (2H,m), 7.88-7.80 (2H,m), 7.59-7.55 (1H,m), 3.72 (3H,s), 3.29 (3H,s)

[0319]

Production Example 4-1

A mixture of N^2 -methyl-5-trifluoromethylpyridin-2,3-diamine (1.0 g), 2-ethylsulfonylbenzaldehyde (0.96 g), sodium hydrogen sulfite (1.80 g), sodium hydrogen sulfite (1.80 g) and DMF (10 ml) was stirred with heating at 160 °C for 5 hours. The reaction mixture was ice-cooled, and water was added thereto. The precipitated crystal was collected by filtration, and washed with water, then hexane. The obtained crystal was dried under reduced pressure to give 1.09 g of 2-(2-ethylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 4).

Present Compound 4



$^1\text{H-NMR}$ (CDCl_3) δ : 8.72-8.70 (1H,m), 8.33-8.31 (1H,m), 7.56-7.49 (2H,m), 7.47-7.43 (1H,m), 7.39-7.34 (1H,m), 3.77 (3H,s), 2.87 (2H,q), 1.24 (3H,t)

5 [0320]

Alternatively, Present Compound 4 can be synthesized by the following method.

[0321]

Production Example 4-2


10 To a mixture of N-(2-methylamino-5-trifluoromethylpyridine-3-yl)-2-ethylsulfanyl-benzamide (6.0 g) and xylene (170 ml), p-toluenesulfonic acid monohydrate (5.47 g) was added, and stirred under reflux for 9 hours. After the reaction mixture was allowed to
15 stand to cool to room temperature, the mixture was concentrated under reduced pressure, and then saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced
20 pressure. The resulting residue was subjected to silica gel column chromatography to give 5.17 g of Present Compound 4.

[0322]

Production Example 5

A mixture of 2-(2-ethylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.25 g), sodium periodate (0.24 g), methanol (6 ml), water (2 ml), and THF (0.8 ml) was stirred at room temperature for 20 minutes, and then heated to 50 °C, and stirred with heating for further 1.5 hours. To the ice-cooled reaction mixture, water was added. The precipitated crystal was collected by filtration. The obtained crystal was dissolved in ethyl acetate, and washed with saturated aqueous sodium thiosulfate solution, saturated aqueous sodium hydrogen carbonate solution, then saturated brine, sequentially. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was washed with hexane to give 0.20 g of 2-(2-ethylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 5).

Present Compound 5



¹H-NMR (CDCl₃) δ: 8.76-8.74 (1H,m), 8.32-8.30 (1H,m), 8.27-8.24 (1H,m), 7.86-7.81 (1H,m), 7.72-7.68 (1H,m), 7.62-7.59 (1H,m), 3.89 (3H,s), 3.42-3.31 (1H,m), 3.02-2.92 (1H,m),


1.31 (3H,t)

[0323]

Production Example 6

To a mixture of 2-(2-ethylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.25 g) and chloroform (3 ml), 3-chloroperbenzoic acid (purity: not less than 65 %, 0.43 g) was added under ice-cooling, and then heated to room temperature, and stirred for 1 hour. To the reaction mixture, chloroform was added under ice-cooling, and then saturated aqueous sodium thiosulfate solution was poured, and extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium hydrogen carbonate solution, then saturated brine, dried over sodium sulfate, and concentrated under reduced pressure. The obtained crystal was washed with hexane, then methyl-tert-butyl ether to give 0.27 g of 2-(2-ethylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 6).

Present Compound 6



¹H-NMR (CDCl₃) δ: 8.75-8.73 (1H,m), 8.29-8.27 (1H,m), 8.25-8.21 (1H,m), 7.87-7.79 (2H,m), 7.58-7.55 (1H,m), 3.72


(3H,s), 3.42 (2H,q), 1.26 (3H,t)

[0324]

Production Example 7

A mixture of N²-methyl-5-trifluoromethylpyridin-2,3-diamine (0.96 g), 2-propylsulfanylbzoic acid (1.08 g), WSC (0.78 g), and pyridine (10 ml) was stirred with heating at 120 °C for 1.5 hours. To the reaction mixture, HOBT (0.74 g) was added, and stirred with heating for 1.5 hours. WSC (0.38 g) was added thereto, and stirred with heating for further 2 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.23 g of 3-methyl-2-(2-propylsulfanyphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 7).

Present Compound 7




¹H-NMR (CDCl₃) δ: 8.72-8.71 (1H,m), 8.33-8.31 (1H,m), 7.53-7.51 (2H,m), 7.46-7.43 (1H,m), 7.38-7.33 (1H,m), 3.77 (3H,s), 2.81 (2H,t), 1.65-1.54 (2H,m), 0.93 (3H,t)

[0325]

Production Example 8

A mixture of 3-methyl-2-(2-propylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.22 g), sodium periodate (0.27 g), methanol (2 ml), and water (4 ml) was stirred with heating at 40 °C for 1.5 hours. To the reaction mixture cooled to room temperature, saturated aqueous sodium hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution were added, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.17 g of 3-methyl-2-(2-propylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 8).

Present Compound 8




¹H-NMR (CDCl₃) δ: 8.77-8.74 (1H,m), 8.31-8.27 (2H,m), 7.86-7.80 (1H,m), 7.72-7.66 (1H,m), 7.63-7.59 (1H,m), 3.90 (3H,s), 3.44-3.36 (1H,m), 2.94-2.85 (1H,m), 2.02-1.87 (1H,m), 1.85-1.72 (1H,m), 1.06 (3H,t)

[0326]

Production Example 9

To a mixture of 2-(2-propylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.22 g) and chloroform (3 ml), 3-chloroperbenzoic acid (purity: not less than 65 %, 0.33 g) was added under ice-cooling, and then heated to room temperature, and stirred for 1.5 hours. Into the reaction mixture, saturated aqueous sodium hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution were poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.24 g of 2-(2-ethylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 9).

Present Compound 9




$^1\text{H-NMR}$ (CDCl_3) δ : 8.75-8.73 (1H,m), 8.29-8.27 (1H,m), 8.24-8.21 (1H,m), 7.86-7.78 (2H,m), 7.58-7.54 (1H,m), 3.71 (3H,s), 3.41-3.33 (2H,m), 1.76-1.65 (2H,m), 1.00 (3H,t)

[0327]

Production Example 10

A mixture of N²-methyl-5-trifluoromethylpyridin-2,3-diamine (0.96 g), 2-allylsulfanylbenzoic acid (1.46 g), WSC (1.06 g), and xylene (10 ml) was stirred with heating at 145 °C for 3 hours. After the mixture was evaporated to remove xylene, acetic acid (10 ml) was added thereto, and stirred with heating at 120 °C for further 1 hour. To the reaction mixture cooled to room temperature, 5 mol/l of aqueous sodium hydroxide solution was added to neutralize, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.83 g of 2-(2-allylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 10).

Present Compound 10




¹H-NMR (CDCl₃) δ: 8.73-8.71 (1H,m), 8.34-8.32 (1H,m), 7.55-7.49 (2H,m), 7.48-7.45 (1H,m), 7.42-7.36 (1H,m), 5.81-5.70 (1H,m), 5.15-5.04 (2H,m), 3.77 (3H,s), 3.49-3.45 (2H,m)
[0328]

Production Example 11

The procedure was performed according to the method described in Production Example 9 using 2-(2-allylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine instead of 3-methyl-2-(2-propylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 2-(2-allylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 11).

Present Compound 11




¹H-NMR (CDCl₃) δ: 8.76-8.74 (1H,m), 8.31-8.29 (1H,m), 8.17-8.14 (1H,m), 7.86-7.76 (2H,m), 7.58-7.55 (1H,m), 5.81-5.69 (1H,m), 5.37-5.28 (2H,m), 4.21-4.14 (2H,m), 3.72 (3H,s)
[0329]

Production Example 12

A mixture of N²-methyl-5-trifluoromethylpyridin-2,3-diamine (0.96 g), 2-isopropylsulfanylbenzoic acid (1.47 g), WSC (1.06 g), and pyridine (10 ml) was stirred with heating at 120 °C for 2 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure.

A mixture of the resulting residue and acetic acid (10 ml) was stirred with heating at 118 °C for 1.5 hours, and then acetic anhydride (10 ml) was poured thereinto, and stirred with heating at 118 °C for 1.5 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with tert-butyl methyl ether. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 1.05 g of 2-(2-isopropylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 12).

Present Compound 12



¹H-NMR (CDCl₃) δ: 8.72-8.71 (1H,m), 8.32-8.30 (1H,m), 7.61-7.58 (1H,m), 7.56-7.51 (1H,m), 7.49-7.46 (1H,m), 7.43-7.38 (1H,m), 3.75 (3H,s), 3.38-3.27 (1H,m), 1.18 (6H,d)


[0330]

Production Example 13

The procedure was performed according to the method described in Production Example 8 using 2-(2-isopropylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine instead of 3-methyl-2-(2-

propylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 2-(2-isopropylsulfinylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 13).

5 Present Compound 13




¹H-NMR (CDCl₃) δ: 8.76-8.75 (1H,m), 8.31-8.30 (1H,m), 8.18-8.15 (1H,m), 7.84-7.80 (1H,m), 7.72-7.67 (1H,m), 7.62-7.59 (1H,m), 3.87 (3H,s), 3.50-3.39 (1H,m), 1.33 (3H,d), 1.09 (3H,d)

[0331]

Production Example 14

The procedure was performed according to the method described in Production Example 9 using 2-(2-isopropylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine instead of 3-methyl-2-(2-propylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 2-(2-isopropylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 14).

Present Compound 14




$^1\text{H-NMR}$ (CDCl_3) δ : 8.75-8.74 (1H,m), 8.28-8.27 (1H,m), 8.22-8.19 (1H,m), 7.86-7.78 (2H,m), 7.57-7.54 (1H,m), 3.85-3.74 (1H,m), 3.71 (3H,s), 1.24 (6H,d)

5 [0332]

Production Example 15

To a mixture of tert-butylthiol (0.67 g) and DMF (12 ml), sodium hydride (60 % in oil) was added under ice-cooling. The mixture was stirred under ice-cooling for 10 minutes, and then 2-(2-fluorophenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (1.09 g) was added. The mixture was stirred at room temperature for 1 hour, and then stirred with heating at 60 °C for 1.5 hours, then at 80 °C for 2 hours. Into the reaction mixture cooled to room temperature, saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 1.19 g of 2-(2-tert-butylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 15).

Present Compound 15




$^1\text{H-NMR}$ (CDCl_3) δ : 8.73-8.70 (1H,m), 8.31-8.29 (1H,m), 7.78-7.75 (1H,m), 7.62-7.52 (3H,m), 3.72 (3H,s), 1.12 (9H,s)

[0333]

5 Production Examples 16 and 17


To a mixture of 2-(2-tert-butylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.87 g) and chloroform (12 ml), 3-chloroperbenzoic acid (purity: not less than 65 %) (0.95 g) was added under ice-cooling, heated to room temperature, and stirred for 2 hours. Into the reaction mixture, saturated aqueous sodium hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution were poured, and extracted with chloroform. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.33 g of 2-(2-tert-butylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 16) and 0.52 g of 2-(2-tert-butylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 17).

Present Compound 16



$^1\text{H-NMR}$ (CDCl_3) δ : 8.76-8.74 (1H,m), 8.33-8.31 (1H,m), 8.19-8.15 (1H,m), 7.85-7.80 (1H,m), 7.75-7.70 (1H,m), 7.63-7.59 (1H,m), 3.84 (3H,s), 1.05 (9H,s)

5 Present Compound 17



$^1\text{H-NMR}$ (CDCl_3) δ : 8.73-8.71 (1H,m), 8.28-8.26 (1H,m), 8.17-8.12 (1H,m), 7.86-7.80 (2H,m), 7.57-7.53 (1H,m), 3.66 (3H,s), 1.31 (9H,s)

10 [0334]

Production Example 18


A mixture of N^2 -methyl-5-trifluoromethylpyridin-2,3-diamine (0.86 g), 2-trifluoromethylsulfonylbenzoic acid (1.00 g), WSC (0.76 g), and pyridine (9 ml) was stirred with heating at 120 °C for 3.5 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure.

20 A mixture of the resulting residue, tripotassium phosphate (2.87 g), and 1-pentanol (9 ml) was stirred with

heating at 135 °C for 3.5 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure.

5 The resulting residue was subjected to silica gel column chromatography to give 0.98 g of 3-methyl-6-trifluoromethyl-2-(2-trifluoromethylsulfanylphenyl)-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 18).

10 Present Compound 18



¹H-NMR (CDCl₃) δ: 8.76-8.74 (1H,m), 8.35-8.33 (1H,m), 7.98-7.93 (1H,m), 7.71-7.59 (3H,m), 3.77 (3H,s)

[0335]


15 Production Example 19

To a mixture of 3-methyl-6-trifluoromethyl-2-(2-trifluoromethylsulfanylphenyl)-3H-imidazo[4,5-b]pyridine (0.80 g) and chloroform (7 ml), 3-chloroperbenzoic acid (purity: not less than 65 %) (0.49 g) was added under ice-cooling. The mixture was heated to room temperature, stirred for 20 minutes, and then heated to 62 °C, and stirred under reflux for 2 hours. Into the reaction mixture cooled to room temperature, saturated aqueous sodium

20

hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution were poured, and extracted with chloroform. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.44 g of 3-methyl-6-trifluoromethyl-2-(2-trifluoromethylsulfinylphenyl)-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 19).

Present Compound 19



$^1\text{H-NMR}$ (CDCl_3) δ : 8.77-8.74 (1H,m), 8.46-8.41 (1H,m), 8.36-8.33 (1H,m), 7.92-7.83 (3H,m), 4.04 (3H,s)


[0336]

Production Example 20

A mixture of 3-methyl-6-trifluoromethyl-2-(2-trifluoromethylsulfonylphenyl)-3H-imidazo[4,5-b]pyridine (0.25 g) and acetic acid (7 ml) was heated to 80 °C, and 30 % of an aqueous solution of hydrogen peroxide (3 ml) was added dropwise thereto. The mixture was heated to 100 °C, and then stirred with heating for 5 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer

was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 76 mg of 3-methyl-6-trifluoromethyl-2-(2-trifluoromethylsulfonylphenyl)-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 20).

Present Compound 20



$^1\text{H-NMR}$ (CDCl_3) δ : 8.76-8.74 (1H,m), 8.35-8.32 (2H,m), 8.05-8.01 (1H,m), 7.97-7.93 (1H,m), 7.72-7.70 (1H,m), 3.71 (3H,s)

[0337]


Production Example 21

To a mixture of N^2 -ethyl-5-trifluoromethylpyridin-2,3-diamine (700 mg), 2-ethylsulfanylbenzoic acid (690 mg), and pyridine (20 ml), WSC (720 mg) was added at room temperature, and heated to 95 °C, and then stirred with heating for 10 hours. Into the reaction mixture cooled to room temperature, saturated aqueous sodium carbonate solution was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure.

The resulting residue was dissolved in xylene (20 ml),

and p-toluenesulfonic acid monohydrate (1.6 g) was added thereto. The mixture was heated to 170 °C, and stirred with heating for 9.5 hours. Into the reaction mixture cooled to room temperature, saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 295 mg of 3-ethyl-2-(2-ethylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 21).

Present Compound 21



¹H-NMR (CDCl₃) δ: 8.70 (1H,s), 8.32 (1H,s), 7.60-7.30 (4H,m), 4.25 (2H,q), 2.89 (2H,q), 1.35-1.30 (3H,m), 1.27-1.21 (3H,m)


[0338]

Production Examples 22 and 23

The procedure was performed according to the method described in Production Examples 16 and 17 using 3-ethyl-2-(2-ethylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give


140 mg of 2-(2-ethylsulfinylphenyl)-3-ethyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 22) and 60 mg of 2-(2-ethylsulfonylphenyl)-3-ethyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 23).

Present Compound 22



¹H-NMR (CDCl₃) δ: 8.75 (1H,d), 8.31 (1H,d), 8.24 (1H,dd),
10 7.84 (1H,dt), 7.70 (1H,dt), 7.58 (1H,dd), 4.35 (2H,q),
3.43-3.30 (1H,m), 3.06-2.94 (1H,m), 1.41 (3H,t), 1.30
(3H,t)

Present Compound 23



15 ¹H-NMR (CDCl₃) δ: 8.71 (1H,d), 8.32 (1H,d), 7.56-7.47
(2H,m), 7.43 (1H,dd), 7.39-7.31 (1H,m), 4.25 (2H,q), 2.89
(2H,q), 1.32 (3H,t), 1.25 (3H,t)


[0339]

Production Example 24

20 A mixture of 2-ethylsulfonyl-N-(2-propylamino-5-trifluoromethyl-pyridin-3-yl)-benzamide (710 mg), p-

toluenesulfonic acid monohydrate (1.1 g), and xylene (10 ml) was stirred with heating at 160 °C for 10 hours. Into the reaction mixture cooled to room temperature, saturated aqueous sodium hydrogen carbonate solution was poured, and
5 extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 522 mg of 2-(2-ethylsulfanylphenyl)-3-propyl-6-trifluoromethyl-3H-
10 imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 24).

Present Compound 24



¹H-NMR (CDCl₃) δ: 8.70 (1H,d), 8.31 (1H,d), 7.56-7.32
15 (4H,m), 4.18 (2H,t), 2.89 (2H,q), 1.78-1.73 (2H,m), 1.28-1.21 (3H,m), 0.79 (3H,t)


[0340]

Production Examples 25 and 26

The procedure was performed according to the method
20 described in Production Examples 16 and 17 using 2-(2-ethylsulfanylphenyl)-3-propyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-


butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 100 mg of 2-(2-ethylsulfinylphenyl)-3-propyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present
5 Compound 25) and 42 mg of 2-(2-ethylsulfonylphenyl)-3-propyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 26).

Present Compound 25



10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.74 (1H,d), 8.31 (1H,d), 8.24 (1H,dd), 7.84 (1H,td), 7.70 (1H,td), 7.58 (1H,dd), 4.33-4.24 (2H,m), 3.40-3.35 (1H,m), 3.03-2.98 (1H,m), 1.84-1.71 (2H,m), 1.29 (3H,t), 0.81 (3H,t)

Present Compound 26



15


$^1\text{H-NMR}$ (CDCl_3) δ : 8.65 (1H,s), 8.21-8.13 (2H,m), 7.79-7.70 (2H,m), 7.52-7.49 (1H,m), 4.12-3.92 (2H,m), 3.50-3.30 (2H,m), 1.81-1.70 (2H,m), 1.22-1.16 (3H,m), 0.84-0.76 (3H,m)

[0341]

Production Example 27

The procedure was performed according to the method described in Production Example 21 using N²-isopropyl-5-trifluoromethyl-pyridin-2,3-diamine instead of N²-ethyl-5-trifluoromethyl-pyridin-2,3-diamine to give 130 mg of 2-(2-ethylsulfanylphenyl)-3-isopropyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 27).

10 Present Compound 27



¹H-NMR (CDCl₃) δ: 8.68 (1H,d), 8.28 (1H,d), 7.55-7.47 (2H,m), 7.39-7.31 (2H,m), 4.40-4.33 (1H,m), 2.91 (2H,q), 1.77-1.66 (6H,m), 1.25 (3H,t)


15 [0342]

Production Examples 28 and 29

The procedure was performed according to the method described in Production Examples 16 and 17 using 2-(2-ethylsulfanylphenyl)-3-isopropyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 60 mg of 2-(2-


ethylsulfinylphenyl)-3-isopropyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 28) and 55 mg of 2-(2-ethylsulfonylphenyl)-3-isopropyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine
5 (hereinafter referred to as Present Compound 29).

Present Compound 28



¹H-NMR (CDCl₃) δ: 8.73 (1H,d), 8.28 (1H,d), 8.22 (1H,dd),
7.83 (1H,td), 7.69 (1H,td), 7.50 (1H,dd), 4.60-4.48 (1H,m),
10 3.38-3.26 (1H,m), 3.05-2.95 (1H,m), 1.77-1.71 (6H,m), 1.27
(3H,t)

Present Compound 29



¹H-NMR (CDCl₃) δ: 8.71 (1H,d), 8.24-8.22 (2H,m), 7.85-7.78
15 (2H,m), 7.55-7.52 (1H,m), 4.33-4.26 (1H,m), 3.72-3.62
(1H,m), 3.44-3.34 (1H,m), 1.77-1.69 (6H,m), 1.28 (3H,t)


[0343]

Production Example 30

The procedure was performed according to the method

described in Production Example 21 using N²-cyclopropyl-5-trifluoromethyl-pyridin-2,3-diamine instead of N²-ethyl-5-trifluoromethyl-pyridin-2,3-diamine to give 280 mg of 3-cyclopropyl-2-(2-ethylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 30).

Present Compound 30




¹H-NMR (CDCl₃) δ: 8.74 (1H,s), 8.31 (1H,s), 7.53-7.47 (3H,m), 7.39-7.31 (1H,m), 3.55-3.49 (1H,m), 2.90 (2H,q), 1.25 (3H,t), 1.01-0.94 (2H,m), 0.93-0.86 (2H,m)
[0344]

Production Examples 31 and 32

The procedure was performed according to the method described in Production Examples 16 and 17 using 3-cyclopropyl-2-(2-ethylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 114 mg of 3-cyclopropyl-2-(2-ethylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 31) and 109 mg of 3-cyclopropyl-2-(2-ethylsulfonylphenyl)-6-


trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 32).

Present Compound 31



5 $^1\text{H-NMR}$ (CDCl_3) δ : 8.77 (1H,d), 8.29-8.26 (2H,m), 7.85-7.79 (2H,m), 7.72-7.67 (1H,m), 3.57-3.51 (1H,m), 3.49-3.39 (1H,m), 3.09-2.95 (1H,m), 1.34 (3H,t), 1.29-1.16 (1H,m), 1.09-0.92 (2H,m), 0.80-0.65 (1H,m)

Present Compound 32



10

$^1\text{H-NMR}$ (CDCl_3) δ : 8.76 (1H,s), 8.27-8.22 (2H,m), 7.87-7.78 (2H,m), 7.65 (1H,dd), 3.60 (2H,brs), 3.39-3.33 (1H,m), 1.29 (3H,t), 1.11-1.11 (2H,m), 0.93 (2H,brs).


[0345]

15 Production Examples 33 and 34

To a mixture of 2-(2-ethylsulfonylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (270 mg) and DMF (10 ml), 60 % of sodium hydride (in oil) (80 mg) and chloromethyl ethyl ether (118 μl) were sequentially added


under ice-cooling. The mixture was heated to room temperature, and stirred for 5 hours. Into the reaction mixture, water was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 180 mg of 3-ethoxymethyl-2-(2-ethylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 33) and 70 mg of 1-ethoxymethyl-2-(2-ethylsulfanylphenyl)-6-trifluoromethyl-1H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 34)).

Present Compound 33



$^1\text{H-NMR}$ (CDCl_3) δ : 8.73 (1H,d), 8.35 (1H,d), 7.58-7.47 (3H,m), 7.35 (1H,td), 5.60 (2H,s), 3.50 (2H,q), 2.89 (2H,q), 1.25 (3H,t), 1.10 (3H,t)

Present Compound 34




¹H-NMR (CDCl₃) δ: 8.90 (1H,d), 8.19 (1H,d), 7.56-7.50 (3H,m), 7.40-7.36 (1H,m), 5.45 (2H,s), 3.25 (2H,q), 2.85 (2H,q), 1.22 (3H,t), 1.04 (3H,t)
[0346]

5 Production Examples 35 and 36


The procedure was performed according to the method described in Production Examples 16 and 17 using 3-ethoxymethyl-2-(2-ethylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 86 mg of 2-(2-ethylsulfanylphenyl)-3-ethoxymethyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 35) and 96 mg of 2-(2-ethylsulfonylphenyl)-3-ethoxymethyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 36).

Present Compound 35



¹H-NMR (CDCl₃) δ: 8.75 (1H,d), 8.32 (1H,d), 8.29 (1H,dd), 8.04 (1H,dd), 7.82 (1H,td), 7.70 (1H,td), 5.83 (1H,d), 5.47 (1H,d), 3.80-3.70 (2H,m), 3.61-3.49 (1H,m), 3.10-3.01 (1H,m), 1.40 (3H,t), 1.25 (3H,t)

Present Compound 36




$^1\text{H-NMR}$ (CDCl_3) δ : 8.75 (1H,d), 8.29 (1H,d), 8.21 (1H,dd),
7.86-7.78 (2H,m), 7.72 (1H,dd), 5.51 (2H,brs), 3.64 (2H,q),
5 3.49 (2H,q), 1.28 (3H,t), 1.17 (3H,t)

[0347]

Production Example 37

To a mixture of 2-(2-ethylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (98 mg) and DMF
10 (6 ml), 60 % of sodium hydride (in oil) (29 mg) and chloromethyl methyl ether (25 μl) were sequentially added under ice-cooling. The mixture was heated room temperature, and stirred for 4 hours. Then, water was poured thereinto, and extracted with ethyl acetate. The organic layer was
15 dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 72 mg of 3-methoxymethyl-2-(2-ethylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 37).

20 Present Compound 37




$^1\text{H-NMR}$ (CDCl_3) δ : 8.74 (1H,d), 8.36 (1H,d), 7.58-7.47 (3H,m), 7.35 (1H,td), 5.55 (2H,s), 3.31 (3H,s), 2.90 (2H,q), 1.25 (3H,t)

5 [0348]

Production Example 38

The procedure was performed according to the method described in Production Example 37 using 2-(2-ethylsulfonylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-ethylsulfonylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 79 mg of 2-(2-ethylsulfonylphenyl)-3-methoxymethyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 38).

15 Present Compound 38




$^1\text{H-NMR}$ (CDCl_3) δ : 8.76 (1H,d), 8.31 (1H,d), 8.21 (1H,dd), 7.87-7.78 (2H,m), 7.72 (1H,dd), 5.48 (2H,brs), 3.50 (2H,q), 3.43 (3H,s), 1.28 (3H,t)

[0349]

Production Example 39

A mixture of 2-ethylsulfanyl-N-(5-methyl-2-methylaminopyridin-3-yl)-benzamide (1.40 g), tripotassium
5 phosphate (1.97 g), and tert-butanol (10 ml) was stirred
under reflux at 82 °C for 2 hours. Into the reaction
mixture cooled to room temperature, water was poured, and
extracted with ethyl acetate. The organic layer was dried
over magnesium sulfate, and concentrated under reduced
10 pressure. The resulting residue was subjected to silica gel
column chromatography to give 0.27 g of 2-(2-
ethylsulfanylphenyl)-3,6-dimethyl-3H-imidazo[4,5-b]pyridine
(hereinafter referred to as Present Compound 39).

Present Compound 39



¹H-NMR (CDCl₃) δ: 8.28-8.26 (1H,m), 7.89-7.88 (1H,m), 7.50-
7.43 (3H,m), 7.35-7.30 (1H,m), 3.72 (3H,s), 2.85 (2H,q),
2.51 (3H,s), 1.22 (3H,t)


[0350]

20 Production Examples 40 and 41

The procedure was performed according to the method
described in Production Examples 16 and 17 using 2-(2-
ethylsulfanylphenyl)-3,6-dimethyl-3H-imidazo[4,5-b]pyridine


instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 2-(2-ethylsulfinylphenyl)-3,6-dimethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 40) and 2-(2-ethylsulfonylphenyl)-3,6-dimethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 41).

Present Compound 40



¹H-NMR (CDCl₃) δ: 8.32-8.31 (1H,m), 8.24-8.21 (1H,m), 7.88-7.86 (1H,m), 7.81-7.77 (1H,m), 7.69-7.64 (1H,m), 7.60-7.57 (1H,m), 3.83 (3H,s), 3.38-3.28 (1H,m), 2.97-2.87 (1H,m), 2.52 (3H,s), 1.27 (3H,t)

Present Compound 41



¹H-NMR (CDCl₃) δ: 8.31-8.29 (1H,m), 8.23-8.20 (1H,m), 7.84-7.75 (3H,m), 7.57-7.54 (1H,m), 3.66 (3H,s), 3.45 (2H,q), 2.51 (3H,s), 1.24 (3H,t)


[0351]

Production Example 42

A mixture of 5-bromo-N²-methylpyridin-2,3-diamine (0.70 g), 2-ethylsulfanylbenzoic acid (0.66 g), WSC (0.80

g), HOBt (23 mg), and pyridine (20 ml) was stirred under reflux at 120 °C for 30 minutes. After the reaction mixture was allowed to stand overnight, the mixture was stirred under reflux at 120 °C for 9.5 hours again. Into the
5 reaction mixture cooled to room temperature, water was poured under ice-cooling, and extracted with ethyl acetate. The organic layer was washed with water and saturated brine, dried over sodium sulfate, and concentrated under reduced pressure. A mixture of the resulting residue and acetic
10 anhydride (7 ml) was stirred under reflux at 140 °C for 1 hour. Aqueous sodium hydroxide solution was added to the reaction mixture cooled to room temperature to neutralize, and extracted with ethyl acetate. The organic layer was washed with water and saturated brine, dried over sodium
15 sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.60 g of 6-bromo-2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 42).

20 Present Compound 42



¹H-NMR (CDCl₃) δ: 8.47 (1H,d), 8.22 (1H,d), 7.54-7.48 (2H,m), 7.45-7.42 (1H,m), 7.37-7.32 (1H,m), 3.71 (3H,s),


2.86 (2H,q), 1.23 (3H,t)

[0352]

Production Example 43

A mixture of 6-bromo-2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (0.20 g), sodium periodate (0.18 g), methanol (3 ml), water (1 ml), and THF (0.4 ml) was stirred at room temperature for 3.5 hours. Water was poured thereinto under ice-cooling, and then the precipitated solid was collected by filtration. The obtained solid was washed with water and hexane, and dried to give 0.19 g of 6-bromo-2-(2-ethylsulfinylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 43).

Present Compound 43



¹H-NMR (CDCl₃) δ: 8.52-8.51 (1H,m), 8.26-8.22 (1H,m), 8.21-8.20 (1H,m), 7.84-7.78 (1H,m), 7.71-7.65 (1H,m), 7.60-7.57 (1H,m), 3.83 (3H,s), 3.39-3.28 (1H,m), 2.99-2.88 (1H,m), 1.29 (3H,t)


[0353]

Production Example 44

To a mixture of 6-bromo-2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (0.20 g) and chloroform (3

ml), 3-chloroperbenzoic acid (purity: not less than 65 %) (0.34 g) was added under ice-cooling, and then heated to room temperature, and stirred for 3.5 hours. Chloroform and saturated aqueous sodium thiosulfate solution were poured
5 thereinto under ice-cooling, and stirred, and then, water was poured, and extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium hydrogen carbonate solution and saturated brine, dried over sodium sulfate, and concentrated under reduced pressure. The
10 resulting solid was washed with hexane, and dried to give 0.20 g of 6-bromo-2-(2-ethylsulfonylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 44).

Present Compound 44




¹H-NMR (CDCl₃) δ: 8.50 (1H,d), 8.23-8.20 (1H,m), 8.17 (1H,d), 7.86-7.77 (2H,m), 7.57-7.54 (1H,m), 3.66 (3H,s), 3.42 (2H,q), 1.24 (3H,t)

[0354]

Production Example 45

The procedure was performed according to the method described in Production Example 21 using 5-iodo-N²-methylpyridin-2,3-diamine instead of N²-ethyl-5-trifluoromethyl-


pyridin-2,3-diamine to give 435 mg of 2-(2-ethylsulfanylphenyl)-6-iodo-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 45).
Present Compound 45



¹H-NMR (CDCl₃) δ: 8.60 (1H,d), 8.39 (1H,d), 7.54-7.41 (3H,m), 7.38-7.31 (1H,m), 3.71 (3H,s), 2.85 (2H,q), 1.23 (3H,t)
[0355]


10 Production Examples 46 and 47

The procedure was performed according to the method described in Production Examples 16 and 17 using 2-(2-ethylsulfanylphenyl)-6-iodo-3-methyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 136 mg of 2-(2-ethylsulfanylphenyl)-6-iodo-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 46) and 107 mg of 2-(2-ethylsulfonylphenyl)-6-iodo-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 47).
Present Compound 46



¹H-NMR (CDCl₃) δ: 8.63 (1H,d), 8.38 (1H,d), 8.24 (1H,dd), 7.81 (1H,td), 7.68 (1H,td), 7.59 (1H,dd), 3.83 (3H,s), 3.39-3.29 (1H,m), 2.99-2.88 (1H,m), 1.29 (3H,t)

5 Present Compound 47



¹H-NMR (CDCl₃) δ: 8.63 (1H,d), 8.37 (1H,d), 8.21 (1H,dd), 7.85-7.77 (2H,m), 7.56 (1H,dd), 3.65 (3H,s), 3.41 (2H,q), 1.24 (3H,t)

10 [0356]

Production Example 48


A mixture of zinc cyanide (0.47 g), tetrakis(triphenylphosphine)palladium(0.12 g), 6-bromo-2-(2-ethylsulfonylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (0.70 g), and N-methylpyrrolidone (hereinafter referred to as NMP) (4 ml) was stirred at 90 °C for 2 hours. Into the reaction mixture cooled to room temperature, saturated aqueous ammonium chloride solution was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel

15

20

column chromatography to give 0.59 g of 6-cyano-2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 48).

Present Compound 48




$^1\text{H-NMR}$ (CDCl_3) δ : 8.70 (1H,d), 8.35 (1H,d), 7.49-7.58 (2H,m), 7.42-7.47 (1H,m), 7.34-7.40 (1H,m), 3.77 (3H,s), 2.88 (2H,q), 1.24 (3H,t)

[0357]

10 Production Example 49

To a mixture of 6-cyano-2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (0.40 g) and chloroform (5 ml), 70 % of 3-chloroperbenzoic acid (0.35 g) was added under ice-cooling, and stirred at room temperature for 2 hours. To the reaction mixture, saturated aqueous sodium hydrogen carbonate solution was added, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure to give 0.40 g of 6-cyano-2-(2-ethylsulfinylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 49).

Present Compound 49




$^1\text{H-NMR}$ (CDCl_3) δ : 8.74 (1H,d), 8.35 (1H,d), 8.26 (1H,d),
7.84 (1H,t), 7.71 (1H,t), 7.56-7.64 (1H,m), 3.89 (3H,s),
3.28-3.41 (1H,m), 2.91-3.04 (1H,m), 1.26-1.35 (3H,m)

5 [0358]

Production Example 50

To a mixture of 6-cyano-2-(2-ethylsulfinylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (0.25 g) and chloroform (5 ml), 70 % of 3-chloroperbenzoic acid (0.35 g) was added,
10 and stirred at 50 °C for 2 hours. To the reaction mixture, saturated aqueous sodium hydrogen carbonate solution and sodium sulfite were added, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue
15 was subjected to silica gel column chromatography to give 0.27 g of 6-cyano-2-(2-ethylsulfinylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 50).

Present Compound 50



20

$^1\text{H-NMR}$ (CDCl_3) δ : 8.73 (1H,d), 8.32 (1H,d), 8.20-8.26


(1H,m), 7.80-7.89 (2H,m), 7.54-7.60 (1H,m), 3.71 (3H,s),
3.33-3.45 (2H,m), 1.25 (3H,t)

[0359]

Production Example 51

5 To a mixture of 6-bromo-2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (2 g) and THF (60 ml), n-butyllithium (1.6 M in hexane) (4.3 ml) was added dropwise with cooling in dry-ice-acetone bath. The mixture was stirred for 30 minutes, and then DMF (2.2 ml) was poured.
10 The mixture was heated to room temperature, and then saturated aqueous ammonium chloride solution was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel
15 column chromatography to give 189 mg of 2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridin-6-carbaldehyde (hereinafter referred to as Present Compound 51).

Present Compound 51




20 ¹H-NMR (CDCl₃) δ: 10.20 (1H,s), 8.96 (1H,d), 8.55 (1H,d),
7.56-7.24 (4H,m), 3.79 (3H,s), 2.90-2.83 (2H,m), 1.25-1.21
(3H,m)

[0360]

Production Example 52

To a mixture of 2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridin-6-carbaldehyde (122 mg) and
5 chloroform (3 ml), bis(2-methoxyethyl)aminosulfur
trifluoride (700 μ l) was added under ice-cooling. The
mixture was heated to room temperature, and stirred for 3
hours. Into the reaction mixture, saturated aqueous sodium
hydrogen carbonate solution was poured, and extracted with
10 chloroform and ethyl acetate, sequentially. The organic
layer was dried over sodium sulfate, and concentrated under
reduced pressure. The resulting residue was subjected to
silica gel column chromatography to give 49 mg of 6-
difluoromethyl-2-(2-ethylsulfanylphenyl)-3-methyl-3H-
15 imidazo[4,5-b]pyridine.

Present Compound 52



$^1\text{H-NMR}$ (CDCl_3) δ : 8.59 (1H,s), 8.23 (1H,s), 7.56-7.43
(3H,m), 7.38-7.33 (1H,m), 6.88 (1H,t), 3.77 (3H,s), 2.87
20 (2H,q), 1.23 (3H,t)


[0361]

Production Examples 53 and 54

The procedure was performed according to the method


described in Production Examples 16 and 17 using 6-difluoromethyl-2-(2-ethylsulfonylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 110 mg of 6-difluoromethyl-2-(2-ethylsulfonylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 53) and 101 mg of 6-difluoromethyl-2-(2-ethylsulfonylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 54).

Present Compound 53



$^1\text{H-NMR}$ (CDCl_3) δ : 8.62 (1H,s), 8.26-8.23 (2H,m), 7.83 (1H,td), 7.70 (1H,td), 7.62 (1H,dd), 6.90 (1H,t), 3.89 (3H,s), 3.42-3.32 (1H,m), 3.02-2.92 (1H,m), 1.30 (3H,t)

Present Compound 54




$^1\text{H-NMR}$ (CDCl_3) δ : 8.61 (1H,s), 8.24-8.19 (2H,m), 7.87-7.78 (2H,m), 7.58 (1H,dd), 6.89 (1H,t), 3.71 (3H,s), 3.43 (2H,q), 1.25 (3H,t)

[0362]

Production Example 55

A mixture of 6-bromo-2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (500 mg), NMP (24 ml), xylene (10 ml), copper iodide (1.1 g), and sodium pentafluoropropionate (1.1 g) was heated to 170 °C, and stirred with heating for 3 days. Into the reaction mixture cooled to room temperature, saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with tert-butyl methyl ether. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 43 mg of 2-(2-ethylsulfanylphenyl)-3-methyl-6-pentafluoroethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 55).

15 Present Compound 55



¹H-NMR (CDCl₃) δ: 8.66 (1H,d), 8.30 (1H,d), 7.56-7.49 (2H,m), 7.47-7.43 (1H,m), 7.38-7.34 (1H,m), 3.78 (3H,s), 2.88 (2H,q), 1.24 (3H,t)


20 [0363]

Production Example 56

A mixture of 6-bromo-2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (0.35 g), tripotassium

phosphate (0.42 g), phenylboronic acid (0.13 g), 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (0.03 g), tris(dibenzylideneacetone)dipalladium(0) (0.01 g), and 1,4-dioxane (3 ml) was stirred with heating at 70 °C for 2 hours, and then heated to 90 °C, and stirred with heating for further 2 hours. Into the reaction mixture cooled to room temperature, saturated aqueous ammonium chloride solution was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.37 g of 2-(2-ethylsulfanylphenyl)-3-methyl-6-phenyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 56).

Present Compound 56



¹H-NMR (CDCl₃) δ: 8.67 (1H,d), 8.26 (1H,d), 7.63-7.69 (2H,m), 7.45-7.54 (5H,m), 7.38-7.43 (1H,m), 7.32-7.37 (1H,m), 3.77 (3H,s), 2.88 (2H,q), 1.25 (3H,t)


[0364]

Production Example 57

The procedure was performed according to the method described in Production Example 3 using 2-(2-

ethylsulfanylphenyl)-3-methyl-6-phenyl-3H-imidazo[4,5-b]pyridine instead of 3-methyl-2-(2-methylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 0.04 g of 2-(2-ethylsulfonylphenyl)-3-methyl-6-phenyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 57).

Present Compound 57



¹H-NMR (CDCl₃) δ: 8.70 (1H,d), 8.22-8.25 (1H,m), 8.20 (1H,d), 7.77-7.86 (2H,m), 7.64-7.68 (2H,m), 7.57-7.60 (1H,m), 7.48-7.54 (2H,m), 7.38-7.44 (1H,m), 3.72 (3H,s), 3.40-3.53 (2H,m), 1.23-1.29 (3H,m)


[0365]

Production Example 58

To a mixture of 6-bromo-2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (350 mg) and 1,4-dioxane (5 ml), 2-fluorophenylboronic acid (150 mg), tripotassium phosphate (600 mg), 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (25 mg), and tris(dibenzylideneacetone)palladium(0) (14 mg) were sequentially added, heated to 80 °C, and stirred with heating for 8 hours. The reaction mixture was cooled to room temperature, and then water was poured into the

reaction mixture, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give
5 349 mg of 2-(2-ethylsulfanylphenyl)-6-(2-fluorophenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 58).

Present Compound 58



10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.63-8.61 (1H,m), 8.27-8.25 (1H,m), 7.54-7.46 (4H,m), 7.41-7.18 (4H,m), 3.78 (3H,s), 2.88 (2H,q), 1.28-1.22 (3H,m)


[0366]

Production Examples 59 and 60

15 The procedure was performed according to the method described in Production Examples 16 and 17 using 2-(2-ethylsulfanylphenyl)-6-(2-fluorophenyl)-3-methyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 150 mg of 2-(2-ethylsulfanylphenyl)-6-(2-fluorophenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present
20 Compound 59) and 140 mg of 2-(2-ethylsulfonylphenyl)-6-(2-


fluorophenyl)-3-methyl-3H-imidazo[4,5-b]pyridine
(hereinafter referred to as Present Compound 60).

Present Compound 59



5 $^1\text{H-NMR}$ (CDCl_3) δ : 8.67-8.65 (1H,m), 8.27-8.23 (2H,m), 7.81 (1H,td), 7.69 (1H,t), 7.62 (1H,dd), 7.53 (1H,td), 7.43-7.37 (1H,m), 7.31-7.21 (2H,m), 3.89 (3H,s), 3.42-3.32 (1H,m), 3.02-2.92 (1H,m), 1.29 (3H,t)

Present Compound 60



10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.64 (1H,d), 8.24-8.20 (2H,m), 7.84-7.76 (2H,m), 7.60-7.57 (1H,m), 7.55-7.49 (1H,m), 7.40-7.33 (1H,m), 7.29-7.18 (2H,m), 3.72 (3H,s), 3.48 (2H,q), 1.28-1.22 (3H,m)

15 [0367]


Production Example 61

To a mixture of 6-bromo-2-(2-ethylsulfonylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (350 mg) and 1,4-dioxane (5 ml), 3-fluorophenylboronic acid (150 mg), sodium carbonate (320 mg), and dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II) (22 mg) were

20

sequentially added. The mixture was heated to 80 °C, and stirred with heating for 8 hours. Into the mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 240 mg of 2-(2-ethylsulfanylphenyl)-6-(3-fluorophenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 61).

10 Present Compound 61



¹H-NMR (CDCl₃) δ: 8.65 (1H,d), 8.29 (1H,d), 7.55-7.41 (5H,m), 7.38-7.32 (2H,m), 7.15-7.05 (1H,m), 3.77 (3H,s), 2.86 (2H,q), 1.24 (3H,t)

15 [0368]


Production Examples 62 and 63

The procedure was performed according to the method described in Production Examples 16 and 17 using 2-(2-ethylsulfanylphenyl)-6-(3-fluorophenyl)-3-methyl-3H-

20 imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 120 mg of 2-(2-ethylsulfanylphenyl)-6-(3-fluorophenyl)-3-methyl-3H-


imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 62) and 34 mg of 2-(2-ethylsulfonylphenyl)-6-(3-fluorophenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 63).

5 Present Compound 62



$^1\text{H-NMR}$ (CDCl_3) δ : 8.69 (1H,d), 8.26 (1H,dd), 8.22 (1H,d), 7.82 (1H,td), 7.69 (1H,td), 7.62 (1H,dd), 7.51-7.42 (2H,m), 7.39-7.34 (1H,m), 7.15-7.08 (1H,m), 3.89 (3H,s), 3.43-3.33 (1H,m), 3.02-2.92 (1H,m), 1.30 (3H,t)

10 Present Compound 63



$^1\text{H-NMR}$ (CDCl_3) δ : 8.60 (1H,d), 8.16 (1H,dd), 8.11 (1H,d), 7.78-7.71 (2H,m), 7.51 (1H,dd), 7.42-7.34 (2H,m), 7.28 (1H,dd), 7.06-6.99 (1H,m), 3.65 (3H,s), 3.40 (2H,q), 1.19 (3H,t)


[0369]

Production Example 64

The procedure was performed according to the method described in Production Example 61 using 4-fluorophenylboronic acid instead of 3-fluorophenylboronic

acid to give 140 mg of 2-(2-ethylsulfanylphenyl)-6-(4-fluorophenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 64).

Present Compound 64




¹H-NMR (CDCl₃) δ: 8.61 (1H,d), 8.22 (1H,d), 7.63-7.57 (2H,m), 7.54-7.45 (3H,m), 7.38-7.33 (1H,m), 7.23-7.16 (2H,m), 3.77 (3H,s), 2.88 (2H,q), 1.24 (3H,t)
[0370]

10 Production Example 65

The procedure was performed according to the method described in Production Example 3 using 2-(2-ethylsulfanylphenyl)-6-(4-fluorophenyl)-3-methyl-3H-imidazo[4,5-b]pyridine instead of 3-methyl-2-(2-methylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 58 mg of 2-(2-ethylsulfonylphenyl)-6-(4-fluorophenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 65).

Present Compound 65



¹H-NMR (CDCl₃) δ: 8.64 (1H,d), 8.24-8.22 (1H,m), 8.15


(1H,d), 7.86-7.78 (2H,m), 7.62-7.58 (3H,m), 7.22-7.18 (2H,m), 3.72 (3H,s), 3.47 (2H,q), 1.26 (3H,t)

[0371]

Production Example 66

5 The procedure was performed according to the method described in Production Example 21 using N³-methyl-5-trifluoromethyl-pyridin-2,3-diamine instead of N²-ethyl-5-trifluoromethylpyridin-2,3-diamine to give 67 mg of 2-(2-ethylsulfanylphenyl)-1-methyl-6-trifluoromethyl-1H-
10 imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 66).

Present Compound 66



¹H-NMR (CDCl₃) δ: 8.87 (1H,d), 7.99 (1H,d), 7.54-7.48 (3H,m), 7.41-7.34 (1H,m), 3.73 (3H,s), 2.84 (2H,q), 1.21 (3H,t)


[0372]

Production Example 67

20 The procedure was performed according to the method described in Production Example 1 using N¹-methyl-4-trifluoromethylbenzene-1,2-diamine instead of N²-methyl-5-trifluoromethylpyridin-2,3-diamine to give 0.28 g of 1-methyl-2-(2-methylsulfanylphenyl)-5-trifluoromethyl-1H-

benzimidazole (hereinafter referred to as Present Compound 67).

Present Compound 67




5 $^1\text{H-NMR}$ (CDCl_3) δ : 8.12 (1H,brs), 7.62-7.58 (1H,m), 7.55-7.48 (2H,m), 7.45-7.39 (2H,m), 7.34-7.29 (1H,m), 3.70 (3H,s), 2.41 (3H,s)

[0373]

Production Example 68

10 A mixture of 2-ethylsulfanyl-N-(2-methylamino-5-trifluoromethylphenyl)-benzamide (1.64 g), p-toluenesulfonic acid monohydrate (1.76 g), and xylene (50 ml) was stirred under reflux at 150 °C for 1 hour. Into the reaction mixture cooled to room temperature, saturated
15 aqueous sodium hydrogen carbonate solution was poured, and extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium hydrogen carbonate solution and saturated brine, dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue
20 was subjected to silica gel column chromatography to give 1.40 g of 2-(2-ethylsulfanylphenyl)-1-methyl-5-trifluoromethyl-1H-benzimidazole (hereinafter referred to as Present Compound 68).

Present Compound 68




¹H-NMR (CDCl₃) δ: 8.12-8.10 (1H,m), 7.61-7.58 (1H,m), 7.53-7.44 (4H,m), 7.38-7.32 (1H,m), 3.69 (3H,s), 2.84 (2H,q),
5 1.22 (3H,t)

[0374]

Production Example 69

The procedure was performed according to the method described in Production Example 2 using 2-(2-ethylsulfanyphenyl)-1-methyl-5-trifluoromethyl-1H-benzimidazole instead of 3-methyl-2-(2-methylsulfanyphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 0.30 g of 2-(2-ethylsulfanyphenyl)-1-methyl-5-trifluoromethyl-1H-benzimidazole (hereinafter referred to as Present Compound 69).

Present Compound 69




¹H-NMR (CDCl₃) δ: 8.24-8.20 (1H,m), 8.10-8.07 (1H,m), 7.83-7.78 (1H,m), 7.70-7.62 (2H,m), 7.57-7.51 (2H,m), 3.79 (3H,s), 3.36-3.26 (1H,m), 2.98-2.88 (1H,m), 1.26 (3H,t)

[0375]

Production Example 70

The procedure was performed according to the method described in Production Example 3 using 2-(2-ethylsulfanylphenyl)-1-methyl-5-trifluoromethyl-1H-benzimidazole instead of 3-methyl-2-(2-methylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 0.24 g of 2-(2-ethylsulfonylphenyl)-1-methyl-5-trifluoromethyl-1H-benzimidazole (hereinafter referred to as Present Compound 70).

Present Compound 70



¹H-NMR (CDCl₃) δ: 8.24-8.20 (1H,m), 8.07-8.05 (1H,m), 7.84-7.77 (2H,m), 7.64-7.61 (1H,m), 7.57-7.54 (1H,m), 7.53-7.49 (1H,m), 3.63 (3H,s), 3.46-3.34 (2H,m), 1.23 (3H,t)


[0376]

Production Example 71

To a mixture of 3-chloro-N²-methyl-5-trifluoromethylbenzene-1,2-diamine (1.12 g) and THF (5 ml), 2-ethylsulfanylbenzoic acid hydrochloride (1.00 g) was added under ice-cooling, heated to room temperature, and stirred for 1 hour. Sodium hydrogen carbonate (0.46 g) was added thereto, and then the reaction mixture was allowed to stand overnight. The reaction mixture was heated to 66 °C,

and stirred under reflux for 1.5 hours. After the mixture was cooled to room temperature, 2-ethylsulfanylbenzoic acid hydrochloride (0.5 g) was added at room temperature. After the reaction mixture was allowed to stand at room temperature overnight, sodium hydrogen carbonate (1.00 g) was added, and stirred. The reaction mixture was ice-cooled, and then water was added under ice-cooling, and extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium hydrogen carbonate solution and saturated brine, dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.48 g of 7-chloro-2-(2-ethylsulfanylphenyl)-1-methyl-5-trifluoromethyl-1H-benzimidazole (hereinafter referred to as Present Compound 71).

Present Compound 71



$^1\text{H-NMR}$ (CDCl_3) δ : 7.99-7.98 (1H,m), 7.54-7.48 (3H,m), 7.44-7.41 (1H,m), 7.37-7.32 (1H,m), 3.94 (3H,s), 2.86 (2H,q), 1.24 (3H,t)


[0377]

Production Example 72

The procedure was performed according to the method

described in Production Example 2 using 7-chloro-2-(2-ethylsulfanylphenyl)-1-methyl-5-trifluoromethyl-1H-benzimidazole instead of 3-methyl-2-(2-methylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 0.13 g of 7-chloro-2-(2-ethylsulfanylphenyl)-1-methyl-5-trifluoromethyl-1H-benzimidazole (hereinafter referred to as Present Compound 72).

Present Compound 72




¹H-NMR (CDCl₃) δ: 8.23-8.20 (1H,m), 7.97-7.95 (1H,m), 7.85-7.80 (1H,m), 7.71-7.66 (1H,m), 7.58-7.51 (2H,m), 4.04 (3H,s), 3.32-3.21 (1H,m), 2.96-2.87 (1H,m), 1.25 (3H,t)
[0378]

Production Example 73

The procedure was performed according to the method described in Production Example 3 using 7-chloro-2-(2-ethylsulfanylphenyl)-1-methyl-5-trifluoromethyl-1H-benzimidazole instead of 3-methyl-2-(2-methylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 0.16 g of 7-chloro-2-(2-ethylsulfanylphenyl)-1-methyl-5-trifluoromethyl-1H-benzimidazole (hereinafter referred to as Present Compound

73).

Present Compound 73




¹H-NMR (CDCl₃) δ: 8.24-8.20 (1H,m), 7.94-7.92 (1H,m), 7.86-
5 7.78 (2H,m), 7.58-7.52 (2H,m), 3.90 (3H,s), 3.60-3.16
(2H,m), 1.24 (3H,t)

[0379]

Production Example 74

A mixture of 3-bromo-N²-methyl-5-
10 trifluoromethylbenzene-1,2-diamine (2.22 g), 2-
ethylsulfonylbenzoic acid (1.80 g), WSC (1.40 g), and
pyridine (10 ml) was stirred under reflux at 115 °C for 6
hours. Into the reaction mixture cooled to room temperature,
water was poured, and extracted with ethyl acetate. The
15 organic layer was dried over magnesium sulfate, and
concentrated under reduced pressure. The resulting residue
was subjected to silica gel column chromatography to give
1.48 g of 7-bromo-2-(2-ethylsulfonylphenyl)-1-methyl-5-
trifluoromethyl-1H-benzimidazole (hereinafter referred to
20 as Present Compound 74).

Present Compound 74




$^1\text{H-NMR}$ (CDCl_3) δ : 8.03-8.02 (1H,m), 7.71-7.70 (1H,m), 7.54-7.47 (2H,m), 7.44-7.41 (1H,m), 7.37-7.32 (1H,m), 3.95 (3H,s), 2.86 (2H,q), 1.24 (3H,t)

5 [0380]

Production Examples 75 and 76

The procedure was performed according to the method described in Production Examples 16 and 17 using 7-bromo-2-(2-ethylsulfanylphenyl)-1-methyl-5-trifluoromethyl-1H-benzimidazole instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 7-bromo-2-(2-ethylsulfinylphenyl)-1-methyl-5-trifluoromethyl-1H-benzimidazole (hereinafter referred to as Present Compound 75) and 7-bromo-2-(2-ethylsulfonylphenyl)-1-methyl-5-trifluoromethyl-1H-benzimidazole (hereinafter referred to as Present Compound 76).


Present Compound 75



20 $^1\text{H-NMR}$ (CDCl_3) δ : 8.24-8.20 (1H,m), 8.03-7.99 (1H,m), 7.86-7.80 (1H,m), 7.77-7.75 (1H,m), 7.72-7.66 (1H,m), 7.56-7.52

(1H,m), 4.05 (3H,s), 3.33-3.23 (1H,m), 2.98-2.88 (1H,m),
1.26 (3H,t)

Present Compound 76




5 $^1\text{H-NMR}$ (CDCl_3) δ : 8.24-8.20 (1H,m), 7.99-7.96 (1H,m), 7.86-
7.78 (2H,m), 7.75-7.73 (1H,m), 7.57-7.53 (1H,m), 3.90
(3H,s), 3.58-3.19 (2H,m), 1.25 (3H,t)
[0381]

Production Example 77

10 To a mixture of N-(2-hydroxy-5-trifluoromethylphenyl)-
2-methylsulfanylbenzamide (0.33 g), triphenylphosphine
(0.34 g), and THF (7 ml), diethyl azodicarboxylate (40 % in
toluene) (0.57 g) was added dropwise, and stirred at room
temperature for 1.5 hours. Triphenylphosphine (79 mg) and
15 diethyl azodicarboxylate (40 % in toluene, 0.13 g) were
added to the mixture, and stirred for further 1 hour. The
mixture was heated to 50 °C, and stirred for 1.5 hours. The
reaction mixture was cooled to room temperature, and then
the mixture was allowed to stand at room temperature
20 overnight. The reaction mixture was subjected to silica gel
column chromatography to give 0.28 g of 2-(2-
methylsulfanyphenyl)-5-trifluoromethylbenzoxazole
(hereinafter referred to as Present Compound 77).

Present Compound 77



¹H-NMR (CDCl₃) δ: 8.22-8.19 (1H,m), 8.17-8.15 (1H,m), 7.71-7.67 (1H,m), 7.66-7.63 (1H,m), 7.55-7.50 (1H,m), 7.41-7.37 (1H,m), 7.32-7.27 (1H,m), 2.57 (3H,s)

[0382]


Production Example 78

A mixture of 2-amino-4-trifluoromethyl-phenol (0.97 g), 2-ethylsulfanylbenzoic acid (1.10 g), WSC (1.27 g), HOBt (37 mg), and pyridine (5 ml) was stirred under reflux at 115 °C for 2.5 hours. After the mixture was allowed to stand overnight, the mixture was stirred under reflux at 115 °C for 6 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was washed with water, saturated aqueous sodium hydrogen carbonate solution, and saturated brine, sequentially, dried over sodium sulfate, and concentrated under reduced pressure.

A mixture of the resulting residue, p-toluenesulfonic acid monohydrate (2.09 g), and xylene (50 ml) was stirred under reflux at 153 °C for 2 hours. Into the reaction mixture cooled to room temperature, saturated aqueous sodium hydrogen carbonate solution was poured, and

extracted with ethyl acetate. The organic layer was washed
with saturated aqueous sodium hydrogen carbonate solution,
water, 10 % aqueous citric acid solution, water, saturated
aqueous sodium hydrogen carbonate solution, and saturated
5 brine, sequentially, dried over sodium sulfate, and
concentrated under reduced pressure. The resulting residue
was subjected to silica gel column chromatography to give
0.98 g of 2-(2-ethylsulfanylphenyl)-5-
trifluoromethylbenzoxazole (hereinafter referred to as
10 Present Compound 78).

Present Compound 78



¹H-NMR (CDCl₃) δ: 8.19-8.15 (2H,m), 7.71-7.67 (1H,m), 7.66-
7.63 (1H,m), 7.52-7.47 (1H,m), 7.45-7.42 (1H,m), 7.31-7.27
15 (1H,m), 3.06 (2H,q), 1.44 (3H,t)


[0383]

Production Example 79

The procedure was performed according to the method
described in Production Example 2 using 2-(2-
20 ethylsulfanylphenyl)-5-trifluoromethylbenzoxazole instead
of 3-methyl-2-(2-methylsulfanylphenyl)-6-trifluoromethyl-
3H-imidazo[4,5-b]pyridine to give 0.27 g of 2-(2-
ethylsulfanylphenyl)-5-trifluoromethylbenzoxazole

(hereinafter referred to as Present Compound 79).

Present Compound 79




¹H-NMR (CDCl₃) δ: 8.35-8.30 (2H,m), 8.12-8.10 (1H,m), 7.85-
5 7.79 (1H,m), 7.75-7.66 (3H,m), 3.48-3.38 (1H,m), 3.00-2.90
(1H,m), 1.41 (3H,t)

[0384]

Production Example 80

The procedure was performed according to the method
10 described in Production Example 3 using 2-(2-ethylsulfonylphenyl)-5-trifluoromethylbenzoxazole instead
of 3-methyl-2-(2-methylsulfonylphenyl)-6-trifluoromethyl-
3H-imidazo[4,5-b]pyridine to give 0.24 g of 2-(2-ethylsulfonylphenyl)-5-trifluoromethylbenzoxazole
15 (hereinafter referred to as Present Compound 80).

Present Compound 80




¹H-NMR (CDCl₃) δ: 8.28-8.24 (1H,m), 8.11-8.09 (1H,m), 7.99-
7.95 (1H,m), 7.85-7.77 (2H,m), 7.72-7.70 (2H,m), 3.82
20 (2H,q), 1.40 (3H,t)

[0385]

Production Example 81

A mixture of 2-ethylsulfanyphenyl-N-(2-hydroxy-5-trifluoromethylpyridin-3-yl)-benzamide (0.92 g) and phosphorous oxychloride (5 ml) was heated to 120 °C, and stirred under reflux. The reaction mixture cooled to room temperature was allowed to stand overnight. The mixture was heated to 120 °C again, and stirred under reflux for 2 hours. Into the reaction mixture cooled to room temperature, water was poured, and then the precipitated solid was collected by filtration, washed with water and hexane, and dried to give 0.53 g of 2-(2-ethylsulfanyphenyl)-6-trifluoromethyloxazolo[5,4-b]pyridine (hereinafter referred to as Present Compound 81).

Present Compound 81



¹H-NMR (CDCl₃) δ: 8.67 (1H,s), 8.40 (1H,s), 8.25 (1H,d), 7.53 (1H,t), 7.45 (1H,d), 7.32 (1H,t), 3.08 (2H,q), 1.45 (3H,t)


[0386]

20 Production Example 82

The procedure was performed according to the method described in Production Example 2 using 2-(2-ethylsulfanyphenyl)-6-trifluoromethyloxazolo[5,4-

b]pyridine instead of 3-methyl-2-(2-methylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 0.17 g of 2-(2-ethylsulfonylphenyl)-6-trifluoromethyloxazolo[5,4-b]pyridine (hereinafter referred to as Present Compound 82).

5 Present Compound 82



¹H-NMR (CDCl₃) δ: 8.73-8.72 (1H,m), 8.41-8.38 (2H,m), 8.36-8.33 (1H,m), 7.90-7.84 (1H,m), 7.74-7.69 (1H,m), 3.45-3.35 (1H,m), 3.00-2.90 (1H,m), 1.40 (3H,t)


10 [0387]

Production Example 83

The procedure was performed according to the method described in Production Example 3 using 2-(2-

15 b]pyridine instead of 3-methyl-2-(2-methylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 0.19 g of 2-(2-ethylsulfonylphenyl)-6-trifluoromethyloxazolo[5,4-b]pyridine (hereinafter referred to as Present Compound 83).

Present Compound 83



20

¹H-NMR (CDCl₃) δ: 8.75-8.73 (1H,m), 8.40-8.37 (1H,m), 8.29-


8.26 (1H,m), 8.05-8.02 (1H,m), 7.89-7.81 (2H,m), 3.81 (2H,q), 1.43 (3H,t)

[0388]

Production Example 84

5 A mixture of 2-amino-4-trifluoromethylbenzenethiol hydrochloride (0.50 g), 2-methylsulfanylbenzaldehyde (0.33 g), diisopropylethylamine (0.28 g), and DMSO (4 ml) was heated to 170-180 °C, and stirred with heating for 3 hours. Into the reaction mixture cooled to room temperature, water
10 was poured, and extracted with ethyl acetate. The organic layer was washed with water and saturated brine, dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.42 g of 2-(2-
15 methylsulfanylphenyl)-5-trifluoromethylbenzothiazole (hereinafter referred to as Present Compound 84).

Present Compound 84




¹H-NMR (CDCl₃) δ: 8.42-8.40 (1H,m), 8.05-8.02 (1H,m), 7.92-
20 7.89 (1H,m), 7.66-7.63 (1H,m), 7.50-7.42 (2H,m), 7.32-7.27 (1H,m), 2.51 (3H,s)

[0389]

Production Example 85

The procedure was performed according to the method described in Production Example 71 using 2-amino-4-trifluoromethylbenzenethiol hydrochloride instead of 3-chloro-N²-methyl-5-trifluoromethylbenzene-1,2-diamine to
5 give 0.50 g of 2-(2-ethylsulfanylphenyl)-5-trifluoromethylbenzothiazole (hereinafter referred to as Present Compound 85).

Present Compound 85




10 ¹H-NMR (CDCl₃) δ: 8.41-8.39 (1H,m), 8.06-8.00 (2H,m), 7.66-7.62 (1H,m), 7.55-7.51 (1H,m), 7.48-7.42 (1H,m), 7.37-7.32 (1H,m), 2.96 (2H,q), 1.33 (3H,t)

[0390]

Production Example 86

15 The procedure was performed according to the method described in Production Example 2 using 2-(2-ethylsulfanylphenyl)-5-trifluoromethylbenzothiazole instead of 3-methyl-2-(2-methylsulfanylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 0.25 g of 2-(2-
20 ethylsulfanylphenyl)-5-trifluoromethylbenzothiazole (hereinafter referred to as Present Compound 86).

Present Compound 86




$^1\text{H-NMR}$ (CDCl_3) δ : 8.36-8.32 (1H,m), 8.32-8.30 (1H,m), 8.08-8.05 (1H,m), 7.97-7.94 (1H,m), 7.80-7.75 (1H,m), 7.71-7.68 (1H,m), 7.66-7.61 (1H,m), 3.56-3.45 (1H,m), 3.02-2.93 (1H,m), 1.46 (3H,t)

[0391]

Production Example 87

The procedure was performed according to the method described in Production Example 3 using 2-(2-ethylsulfonylphenyl)-5-trifluoromethylbenzothiazole instead of 3-methyl-2-(2-methylsulfonylphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 0.30 g of 2-(2-ethylsulfonylphenyl)-5-trifluoromethylbenzothiazole (hereinafter referred to as Present Compound 87).

Present Compound 87



$^1\text{H-NMR}$ (CDCl_3) δ : 8.33-8.31 (1H,m), 8.26-8.23 (1H,m), 8.10-8.06 (1H,m), 7.81-7.69 (4H,m), 3.75 (2H,q), 1.36 (3H,t)

[0392]


Production Example 88

A mixture of N^2 -methyl-5-trifluoromethylpyridin-2,3-

diamine (1.14 g), 3-chloro-2-ethylsulfanylbenzoic acid (1.56 g), WSC (1.02 g), and pyridine (12 ml) was stirred under reflux at 115 °C for 3 hours. Into the reaction mixture cooled to room temperature, water was poured, and
5 extracted with ethyl acetate. The organic layer was washed with water and saturated brine, dried over magnesium sulfate, and concentrated under reduced pressure.

To the resulting residue, tripotassium phosphate (3.82 g) and pyridine (12 ml) were added, and stirred under
10 reflux at 115 °C for 2.5 hours. The reaction mixture was cooled to room temperature, and then the solvent was evaporated. Tripotassium phosphate (3.82 g) and DMSO (12 ml) were added to the resulting residue, and stirred with heating at 150 °C for 1 hour. The mixture was heated to 165
15 °C, and stirred with heating for further 2 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was washed with water and saturated brine, dried over magnesium sulfate, and concentrated under reduced pressure.
20 The resulting residue was subjected to silica gel column chromatography to give 0.96 g of 2-(3-chloro-2-ethylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 88).

25 Present Compound 88




$^1\text{H-NMR}$ (CDCl_3) δ : 8.74-8.72 (1H,m), 8.32-8.30 (1H,m), 7.71 (1H,dd), 7.48-7.42 (2H,m), 3.73 (3H,s), 2.73 (2H,q), 1.05 (3H,t)

5 [0393]

Production Example 89

To a mixture of 2-(3-chloro-2-ethylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.72 g) and chloroform (10 ml), 3-chloroperbenzoic acid (purity: 69-75 %, 0.66 g) was added under ice-cooling, and then heated to room temperature, and stirred for 20 minutes. Into the reaction mixture, saturated aqueous sodium hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution were poured, and extracted with chloroform 2 times. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.67 g of 2-(3-chloro-2-ethylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 89).

Present Compound 89




$^1\text{H-NMR}$ (CDCl_3) δ : 8.69 (1H,d), 8.25 (1H,d), 7.64 (1H,dd), 7.58 (1H,t), 7.39 (1H,dd), 3.71 (3H,s), 3.60-3.49 (1H,m), 3.33-3.23 (1H,m), 1.33 (3H,t)

5 [0394]

Production Example 90

A mixture of 2-(3-chloro-2-ethylsulfinylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.48 g) and acetic acid (12 ml) was stirred with heating at 80 °C, and then 30 % of an aqueous solution of hydrogen peroxide (6 ml) was added dropwise. The mixture was heated to 100 °C, and stirred with heating for 3 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium hydrogen carbonate solution, dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.21 g of 2-(3-chloro-2-ethylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 90).

Present Compound 90




$^1\text{H-NMR}$ (CDCl_3) δ : 8.73-8.71 (1H,m), 8.25 (1H,d), 7.85-7.81 (1H,m), 7.71 (1H,t), 7.46-7.43 (1H,m), 3.75 (3H,s), 3.59-3.41 (2H,m), 1.33 (3H,t)

5 [0395]

Production Examples 91 and 92

The procedure was performed according to the method described in Production Examples 16 and 17 using 2-(2-ethylsulfanyl-phenyl)-3-methyl-6-pentafluoroethyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 27 mg of 2-(2-ethylsulfinyl-phenyl)-3-methyl-6-pentafluoroethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 112) and 31 mg of 2-(2-ethylsulfonylphenyl)-3-methyl-6-pentafluoroethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 113).


Present Compound 112



20 $^1\text{H-NMR}$ (CDCl_3) δ : 8.70 (1H,d), 8.29 (1H,d), 8.27 (1H,d),

7.84 (1H,t), 7.71 (1H,t), 7.60 (1H,d), 3.90 (3H,s), 3.43-3.33 (1H,m), 3.04-2.94 (1H,m), 1.31 (3H,t)

Present Compound 113



5 $^1\text{H-NMR}$ (CDCl_3) δ : 8.69 (1H,s), 8.25 (1H,s), 8.23-8.22 (1H,m), 7.88-7.79 (2H,m), 7.59-7.52 (1H,m), 3.72 (3H,s), 3.43 (2H,q), 1.26 (3H,t)

[0396]


Production Example 93

10 A mixture of N²-methyl-5-trifluoromethylpyridin-2,3-diamine (1.14 g), 2-ethylsulfonyl-4-fluorobenzoic acid (1.44 g), WSC (1.02 g), and pyridine (12 ml) was stirred with heating at 120 °C for 1.5 hours. Into the reaction mixture cooled to room temperature, water was poured, and
15 extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure.

To the resulting residue, p-toluenesulfonic acid (3.42 g), xylene (10 ml) and NMP (2 ml) were added, and stirred
20 under reflux at 150 °C for 4.5 hours with removing water with Dean-Stark apparatus. Into the reaction mixture cooled to room temperature, water was poured, and extracted with

ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 1.05 g of 2-(2-ethylsulfanyl-4-fluorophenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 116).

Present Compound 116




¹H-NMR (CDCl₃) δ: 8.70-8.68 (1H,m), 8.31-8.28 (1H,m), 7.43-7.38 (1H,m), 7.17-7.13 (1H,m), 7.04-6.98 (1H,m), 3.75 (3H,s), 2.89 (2H,q), 1.26 (3H,t)
[0397]

Production Examples 94 and 95

To a mixture of 2-(2-ethylsulfanyl-4-fluorophenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.85 g) and chloroform (12 ml), 3-chloroperbenzoic acid (purity: not less than 65 %, 0.81) was added under ice-cooling, heated to room temperature, and stirred for 30 minutes. Into the reaction mixture, saturated aqueous sodium hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution were poured, and extracted with


chloroform. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.33 g of 2-(2-ethylsulfinyl-4-fluorophenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 117) and 0.52 g of 2-(2-ethylsulfonyl-4-fluorophenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 118).

10 Present Compound 117



$^1\text{H-NMR}$ (CDCl_3) δ : 8.76-8.75 (1H,m), 8.31-8.30 (1H,m), 8.01-7.98 (1H,m), 7.65-7.61 (1H,m), 7.41-7.36 (1H,m), 3.90 (3H,s), 3.47-3.37 (1H,m), 3.04-2.94 (1H,m), 1.33 (3H,t)

15 Present Compound 118




$^1\text{H-NMR}$ (CDCl_3) δ : 8.76-8.74 (1H,m), 8.29-8.27 (1H,m), 7.97-7.94 (1H,m), 7.60-7.51 (2H,m), 3.72 (3H,s), 3.44 (2H,q), 1.28 (3H,t)

[0398]

Production Example 96

To a pressure-resistant reaction container, 6-bromo-2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (1.74 g), copper (II) acetylacetonate (0.26 g), acetylacetonate (0.50 g), cesium carbonate (3.25 g), NMP (9 ml), and 28 % of aqueous ammonia (4 ml) were added, and stirred at 120 °C for 6 hours. The mixture was cooled to room temperature, and then saturated aqueous ammonium chloride solution was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 1.20 g of 6-amino-2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 125).

Present Compound 125




¹H-NMR (CDCl₃) δ: 7.99 (1H,d), 7.40-7.49 (4H,m), 7.28-7.34 (1H,m), 3.61-3.70 (5H,m), 2.85 (2H,q), 1.22 (3H,t)

[0399]

Production Example 97

A mixture of 2-(2-ethylsulfanyl-phenyl)-6-iodo-3-methyl-3H-imidazo[4,5-b]pyridine (311 mg), copper iodide (1.5 g), sodium heptafluorobutyrate (1.8 g), NMP (5 ml), and xylene (25 ml) was stirred with heating at 150 °C for 12 hours. Into the reaction mixture cooled to room temperature, saturated aqueous sodium hydrogen carbonate solution and 28 % aqueous ammonia were poured, and extracted with tert-butyl methyl ether. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 118 mg of 2-(2-ethylsulfanyl-phenyl)-6-heptafluoropropyl-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 128).

Present Compound 128



¹H-NMR (CDCl₃) δ: 8.65 (1H,d), 8.29 (1H,d), 7.56-7.51 (2H,m), 7.48-7.43 (1H,m), 7.38-7.34 (1H,m), 3.78 (3H,s), 2.89 (2H,q), 1.25 (3H,t)


[0400]

Production Examples 98 and 99

The procedure was performed according to the method


described in Production Examples 16 and 17 using 2-(2-ethylsulfanyl-phenyl)-6-heptafluoropropyl-3-methyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 2-(2-ethylsulfanyl-phenyl)-6-heptafluoropropyl-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 129) and 2-(2-ethylsulfonylphenyl)-6-heptafluoropropyl-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 130).

Present Compound 129



$^1\text{H-NMR}$ (CDCl_3) δ : 8.68 (1H,d), 8.29-8.24 (2H,m), 7.87-7.81 (1H,m), 7.74-7.68 (1H,m), 7.61 (1H,dd), 3.91 (3H,s), 3.43-3.32 (1H,m), 3.05-2.94 (1H,m), 1.31 (3H,t)

Present Compound 130



$^1\text{H-NMR}$ (CDCl_3) δ : 8.67 (1H,d), 8.26-8.22 (2H,m), 7.87-7.81 (2H,m), 7.59-7.55 (1H,m), 3.73 (3H,s), 3.43 (2H,q), 1.26


(3H,t).

[0401]

Production Example 100

To a mixture of 2-(4-chloro-2-fluorophenyl)-3-methyl-
5 6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.83 g) and
DMF (5 ml), a suspension of sodium ethylmercaptide (80 %, 0.26 g) was added dropwise in DMF under ice-cooling, and stirred for 1 hour under ice-cooling. Into the reaction mixture, saturated aqueous sodium hydrogen carbonate
10 solution was poured under ice-cooling, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography. 0.20 g of the resulting product was
15 subjected to recycle preparation liquid chromatography to give 0.12 g of 2-(2-ethylsulfanyl-4-chlorophenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 133).

Present Compound 133



20

¹H-NMR (CDCl₃) δ: 8.73-8.71 (1H,m), 8.33-8.31 (1H,m), 7.44 (1H,d), 7.38 (1H,d), 7.34-7.30 (1H,m), 3.77 (3H,s), 2.91


(2H,q), 1.27 (3H,t)

[0402]

Production Examples 101 and 102


The procedure was performed according to the method
5 described in Production Examples 16 and 17 using 2-(2-
ethylsulfanyl-4-chlorophenyl)-3-methyl-6-trifluoromethyl-
3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-
butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-
imidazo[4,5-b]pyridine to give 0.34 g of 2-(2-
10 ethylsulfanyl-4-chlorophenyl)-3-methyl-6-trifluoromethyl-
3H-imidazo[4,5-b]pyridine (hereinafter referred to as
Present Compound 134) and 0.17 g of 2-(2-ethylsulfanyl-4-
chlorophenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-
b]pyridine (hereinafter referred to as Present Compound
15 135).

Present Compound 134



¹H-NMR (CDCl₃) δ: 8.76-8.75 (1H,m), 8.32-8.30 (1H,m), 8.25
(1H,d), 7.68-7.65 (1H,m), 7.56 (1H,d), 3.91 (3H,s), 3.49-
20 3.39 (1H,m), 3.05-2.95 (1H,m), 1.35 (3H,t)

Present Compound 135




$^1\text{H-NMR}$ (CDCl_3) δ : 8.76-8.74 (1H,m), 8.29-8.27 (1H,m), 8.23-8.21 (1H,m), 7.83-7.80 (1H,m), 7.51 (1H,d), 3.72 (3H,s), 3.45 (2H,q), 1.28 (3H,t)

5 [0403]

Production Example 103

To a mixture of 2-(2-ethylsulfonyl-phenyl)-6-iodo-3-methyl-3H-imidazo[4,5-b]pyridine (430 mg), copper iodide (229 mg), and DMF (3 ml), sodium sulfide nonahydrate (721 mg) was added at room temperature. The mixture was stirred with heating at 110 °C for 6 hours. Into the reaction mixture cooled to room temperature, saturated aqueous ammonium chloride water solution was poured, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 85 mg of 2-(2-ethylsulfonyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridin-6-thiol (hereinafter referred to as Present Compound 145):


20 Present Compound 145



$^1\text{H-NMR}$ (CDCl_3) δ : 8.41 (1H,d), 8.11 (1H,d), 7.53-7.47 (2H,m), 7.46-7.42 (1H,m), 7.37-7.31 (1H,m), 3.71 (3H,s), 3.56 (1H,s), 2.85 (2H,q), 1.22 (3H,t); and

5 and 98 mg of a compound of the following formula (hereinafter referred to as Compound (145A)).

Compound (145A)



$^1\text{H-NMR}$ (CDCl_3) δ : 8.48 (2H,d), 8.25 (2H,d), 7.53-7.47 (4H,m), 7.47-7.43 (2H,m), 7.36-7.32 (2H,m), 3.73 (6H,s), 2.86 (4H,q), 1.23 (6H,t)


[0404]

Production Example 104

To a mixture of 6-bromo-2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (348 mg), [1,1-bis(diphenylphosphino)ferrocene]palladium(II) dichloride dichloromethane adduct (41 mg), and THF (5 mL), diisopropyl zinc (1 M in toluene, 2 mL) was added at room temperature. The mixture was stirred at room temperature for 4 hours,

washed with water, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 290 mg of 2-(2-ethylsulfanyl-phenyl)-6-isopropyl-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 147).

Present Compound 147



¹H-NMR (CDCl₃) δ: 8.35 (1H,s), 7.97 (1H,s), 7.51-7.42 (3H,m), 7.32-7.26 (1H,m), 3.73 (3H,s), 3.15-3.08 (1H,m), 2.83 (2H,q), 1.35 (6H,d), 1.20 (3H,t).

[0405]


Production Example 105

To a mixture of copper cyanide (514 mg) and THF (20 ml), tert-butyilmagnesium chloride (1 M in THF) was added with cooling at -78 °C, and stirred at the temperature for 2 hours. To the mixture, a mixture of 6-bromo-2-(2-ethylsulfanyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (1 g) and THF (10 ml) was added dropwise with cooling at -78 °C.

The reaction mixture was stirred for 16 hours with

warming to room temperature gradually, and then saturated aqueous ammonium chloride solution was poured, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to 111 mg of 6-tert-butyl-2-(2-ethylsulfanyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 150).

Present Compound 150



¹H-NMR (CDCl₃) δ: 8.53 (1H,d), 8.11 (1H,d), 7.51-7.46 (2H,m), 7.45-7.40 (1H,m), 7.36-7.29 (1H,m), 3.72 (3H,s), 2.86 (2H,q), 1.45 (9H,s), 1.24 (3H,t)


[0406]

Production Example 106

A mixture of N²-methyl-6-trifluoromethylpyridin-2,3-diamine (0.96 g), 2-ethylsulfanylbenzoic acid (1.01 g), WSC (1.06 g), and pyridine (5 ml) was stirred under reflux for 3 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. To the resulting

residue, a mixture of tripotassium phosphate (2.65 g) and 1-propanol (10 ml) was added, and stirred under reflux for 4 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.98 g of 2-(2-ethylsulfanylphenyl)-3-methyl-5-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 151).

Present Compound 151



$^1\text{H-NMR}$ (CDCl_3) δ : 8.20 (1H,d), 7.67 (1H,d), 7.56-7.48 (2H,m), 7.47-7.42 (1H,m), 7.40-7.33 (1H,m), 3.78 (3H,s), 2.87 (2H,q), 1.24 (3H,t)


[0407]

Production Examples 107 and 108

The procedure was performed according to the method described in Production Examples 16 and 17 using 2-(2-ethylsulfanylphenyl)-3-methyl-5-trifluoromethyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-


imidazo[4,5-b]pyridine to give 0.23 g of 2-(2-ethylsulfinylphenyl)-3-methyl-5-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 159) and 0.42 g of 2-(2-ethylsulfonylphenyl)-3-methyl-5-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 160).

Present Compound 159



$^1\text{H-NMR}$ (CDCl_3) δ : 8.26 (1H,dd), 8.19 (1H,dd), 7.86-7.80 (1H,m), 7.74-7.67 (2H,m), 7.60 (1H,dd), 3.90 (3H,s), 3.41-3.27 (1H,m), 3.02-2.89 (1H,m), 1.31-1.26 (3H,m)

Present Compound 160



$^1\text{H-NMR}$ (CDCl_3) δ : 8.25-8.21 (1H,m), 8.15 (1H,d), 7.88-7.78 (2H,m), 7.69 (1H,d), 7.59-7.54 (1H,m), 3.73 (3H,s), 3.43 (2H,q), 1.25 (3H,t)


[0408]

Production Example 109

A mixture of Compound (145A) (3.3 g) and DMF (60m) was

cooled to -50°C , and bubbled with trifluoroiodomethane (17 g). The mixture was kept at -50°C , and tetrakis(dimethylamino)ethylene (3.8 ml) was added dropwise. The reaction mixture was stirred for 5 hours with warming gradually. The mixture was diluted with water, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 3.2 g of 2-(2-ethylsulfanyl-phenyl)-3-methyl-6-trifluoromethylsulfanyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 161).

Present Compound 161



$^1\text{H-NMR}$ (CDCl_3) δ : 8.66 (1H,d), 8.39 (1H,d), 7.57-7.49 (2H,m), 7.48-7.42 (1H,m), 7.39-7.33 (1H,m), 3.76 (3H,s), 2.88 (2H,q), 1.24 (3H,t)


[0409]

Production Example 110-a

To a mixture of 6-bromo-2-(2-ethylsulfanyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (1 g) and THF (20 ml), n-butyllithium (1.5 M in hexane, 2.1 ml) was added dropwise

at -78 °C. The mixture was stirred at -78 °C for 30 minutes,
and then methyl trifluoroacetate (577 µl) was added
dropwise. The mixture was heated to room temperature, and
then saturated aqueous ammonium chloride solution was
5 poured, and extracted with ethyl acetate. The combined
organic layer was dried over sodium sulfate, and
concentrated under reduced pressure to give 778 mg of crude
1-[2-(2-ethylsulfanyl-phenyl)-3-methyl-3H-imidazo[4,5-
b]pyridin-6-yl]-2,2,2-trifluoro-ethanone (hereinafter
10 referred to as Present Compound 91).

Present Compound 91



[0410]


Production Example 110-b

15 To a mixture of the crude 1-[2-(2-ethylsulfanyl-
phenyl)-3-methyl-3H-imidazo[4,5-b]pyridin-6-yl]-2,2,2-
trifluoro-ethanone from Production Example 110-a (778 mg),
methanol (5 ml), and THF (20 ml), sodium borohydride (109
mg) was added under ice-cooling. The mixture was stirred at
20 room temperature for 12 hours, and then 1N aqueous sodium
hydroxide solution was poured, and extracted with ethyl
acetate. The combined organic layer was dried over sodium

sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 170 mg of 1-[2-(2-ethylsulfanyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridin-6-yl]-2,2,2-trifluoro-ethanol

5 (hereinafter referred to as Present Compound 265).

Present Compound 265



¹H-NMR (CDCl₃) δ: 8.56 (1H,s), 8.25 (1H,s), 7.58-7.35 (4H,m), 5.24 (1H,brs), 3.77 (3H,s), 2.90 (2H,q), 1.27-1.19 (3H,m)


[0411]

Production Example 110-c

To a mixture of 1-[2-(2-ethylsulfanyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridin-6-yl]-2,2,2-trifluoro-ethanol (98 mg), toluene(10 ml), and pyridine (50 μl), thionyl chloride (1 ml) was added. The mixture was stirred with heating at 50 °C for 5 hours, and then saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 66 mg of 6-(1-chloro-2,2,2-

trifluoro-ethyl)-2-(2-ethylsulfanyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 266).

Present Compound 266



$^1\text{H-NMR}$ (CDCl_3) δ : 8.52 (1H,d), 8.29 (1H,d), 7.52-7.49 (2H,m), 7.44 (1H,dd), 7.37-7.33 (1H,m), 5.36 (1H,q), 3.75 (3H,s), 2.88 (2H,q), 1.24 (3H,t)

[0412]


10 Production Example 110-d

To a mixture of 6-(1-chloro-2,2,2-trifluoro-ethyl)-2-(2-ethylsulfanyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (61 mg) and methanol (5 ml), sodium borohydride (20 mg) was added under ice-cooling. The mixture was stirred at room temperature for 3 hours, and then 1N aqueous sodium hydroxide solution was poured, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 40 mg of 2-(2-ethylsulfanyl-phenyl)-3-methyl-6-(2,2,2-trifluoro-ethyl)-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 165).

15

20

Present Compound 165



$^1\text{H-NMR}$ (CDCl_3) δ : 8.53 (1H,s), 8.45-8.37 (1H,m), 7.78-7.31
(4H,m), 3.84 (3H,s), 3.60-3.50 (2H,m), 2.91-2.82 (2H,m),
5 1.23-1.16 (3H,m)


[0413]

Production Example 111

A mixture of N^3 -methyl-6-trifluoromethylpyridin-2,3-diamine (0.38 g), 2-ethylsulfanylbenzoic acid (0.40 g), WSC
10 (0.42 g), and pyridine (3 ml) was stirred under reflux for 3 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. To the resulting
15 residue, tripotassium phosphate (1.06 g) and 1-propanol (4 ml) were added, and stirred under reflux for 4 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under
20 reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.33 g of 2-(2-ethylsulfanylmethyl)-1-methyl-5-trifluoromethyl-1H-

imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 166).

Present Compound 166




5 $^1\text{H-NMR}$ (CDCl_3) δ : 7.85 (1H,d), 7.68 (1H,d), 7.58-7.47 (3H,m), 7.43-7.33 (1H,m), 3.71 (3H,s), 2.83 (2H,q), 1.20 (3H,t)

[0414]

Production Examples 112 and 113


10 The procedure was performed according to the method described in Production Examples 16 and 17 using 2-(2-ethylsulfanylphenyl)-1-methyl-5-trifluoromethyl-1H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 0.23 g of 2-(2-ethylsulfanylphenyl)-1-methyl-5-trifluoromethyl-1H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 167) and 0.42 g of 2-(2-ethylsulfonylphenyl)-1-methyl-5-trifluoromethyl-1H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 168).

Present Compound 167



¹H-NMR (CDCl₃) δ: 8.25 (1H,dd), 7.91 (1H,d), 7.83 (1H,td), 7.74 (1H,d), 7.69 (1H,td), 7.56 (1H,dd), 3.83 (3H,s), 3.53-3.37 (1H,m), 3.18-3.05 (1H,m), 1.30 (3H,t)

5 Present Compound 168



¹H-NMR (CDCl₃) δ: 8.26-8.19 (1H,m), 7.90-7.86 (1H,m), 7.85-7.80 (2H,m), 7.72 (1H,d), 7.59-7.53 (1H,m), 3.66 (3H,s), 3.60-3.30 (2H,m), 1.24 (3H,t)

10 [0415]

Production Example 114


To a mixture of ethylmercaptide (80 %, 0.35 g) and DMF (9 ml), a solution of 2-(2-fluoro-4-trifluoromethylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (1.0 g) in DMF was added dropwise under ice-cooling, heated to room temperature, and stirred at room temperature for 30 minutes. To the reaction mixture, saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with ethyl acetate. The organic layer was dried over magnesium

15

sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 1.10 g of 2-(2-ethylsulfanyl-4-trifluoromethylphenyl)-3-methyl-6-trifluoromethyl-3H-

5 imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 174).

Present Compound 174



¹H-NMR (CDCl₃) δ: 8.75-8.73 (1H,m), 8.35-8.33 (1H,m), 7.70-
10 7.68 (1H,m), 7.62-7.56 (2H,m), 3.79 (3H,s), 2.95 (2H,q),
1.28 (3H,t)


[0416]

Production Examples 115 and 116

The procedure was performed according to the method
15 described in Production Examples 16 and 17 using 2-(2-ethylsulfanyl-4-trifluoromethylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 0.51 g of 2-(2-ethylsulfanyl-4-trifluoromethylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter
20 referred to as Present Compound 175) and 0.26 g of 2-(2-


ethylsulfonyl-4-trifluoromethylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 176).

Present Compound 175



¹H-NMR (CDCl₃) δ: 8.79-8.78 (1H,m), 8.57-8.55 (1H,m), 8.35-8.34 (1H,m), 7.97-7.94 (1H,m), 7.77 (1H,d), 3.94 (3H,s), 3.53-3.43 (1H,m), 3.07-2.98 (1H,m), 1.36 (3H,t)

Present Compound 176



¹H-NMR (CDCl₃) δ: 8.78-8.76 (1H,m), 8.51-8.49 (1H,m), 8.31-8.30 (1H,m), 8.12-8.09 (1H,m), 7.74 (1H,d), 3.74 (3H,s), 3.48 (2H,q), 1.29 (3H,t)

[0417]


15 Production Example 117

A mixture of 3-amino-5-(trifluoromethyl)pyridin-2-thiol 0.56g, 2-ethylsulfonylbenzoic acid (0.52 g), WSC (0.80 g), HOBT (39 mg), and pyridine (6 ml) was stirred at 60 °C for 2 hours. After the reaction mixture was allowed to

stand to cool, water was poured thereinto, and extracted with ethyl acetate. The organic layer was washed with water, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure.

5 A mixture of the resulting residue, p-toluenesulfonic acid monohydrate (0.65 g), and N-methylpyrrolidinone (5 ml) was stirred with heating at 150 °C for 2 hours. After the reaction mixture was allowed to cool, water was poured thereinto, and extracted with ethyl acetate. The organic
10 layer was washed with water, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.38 g of 2-(2-ethylsulfanylphenyl)-6-(trifluoromethyl)thiazolo[5,4-b]pyridine (hereinafter
15 referred to as Present Compound 183).

Present Compound 183



¹H-NMR (CDCl₃) δ: 8.86 (1H,d), 8.57 (1H,d), 8.03 (1H,dd),
7.55 (1H,dd), 7.48 (1H,td), 7.36 (1H,td), 2.98 (2H,q), 1.34
20 (3H,t).


[0418]

Production Examples 118 and 119

The procedure was performed according to the method


described in Production Examples 16 and 17 using 2-(2-ethylsulfanylphenyl)-6-(trifluoromethyl)thiazolo[5,4-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give
5 0.13 g of 2-(2-ethylsulfanylphenyl)-6-(trifluoromethyl)thiazolo[5,4-b]pyridine (hereinafter referred to as Present Compound 184) and 0.14 g of 2-(2-ethylsulfonylphenyl)-6-(trifluoromethyl)thiazolo[5,4-b]pyridine (hereinafter referred to as Present Compound
10 185).

Present Compound 184



¹H-NMR (CDCl₃) δ: 8.90 (1H,d), 8.49 (1H,d), 8.37 (1H,dd),
7.99 (1H,dd), 7.81 (1H,td), 7.67 (1H,td), 3.52-3.42 (1H,m),
15 3.01-2.92 (1H,m), 1.45 (3H,t).

Present Compound 185




¹H-NMR (CDCl₃) δ: 8.92 (1H,d), 8.52 (1H,d), 8.25 (1H,dd),
7.84-7.71 (3H,m), 3.73 (2H,q), 1.37 (3H,t).

[0419]

Production Example 120-1

A mixture of 2-(2-fluorophenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (7.83 g), sodium sulfide (6.20 g), and NMP (27 ml) was stirred with heating at 150 °C for 2 hours. Into the reaction mixture cooled to room temperature, water and 12N hydrochloric acid were poured, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 1.74 g of a compound of the following formula:




(hereinafter referred to as Compound (186A)).

¹H-NMR (CDCl₃) δ: 8.73-8.71 (2H,m), 8.31-8.29 (2H,m), 7.77-7.73 (2H,m), 7.47-7.37 (6H,m), 3.77 (6H,s)

[0420]


Production Example 120-2

To a solution of sodium hydride (60 % in oil, 0.12 g) in DMF, Compound (186A) (0.80 g) of the following formula:



was added under ice-cooling, and stirred for 10 minutes under ice-cooling. To the mixture, 2,2,2-trifluoroethyl iodide (0.65 g) was added under ice-cooling, heated to room temperature, and stirred for 15 minutes. Then, the mixture was heated to 60 °C, and stirred with heating for 1 hour. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.23 g of 3-methyl-2-[2-(2,2,2-trifluoroethylsulfanylphenyl)]-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 186).

Present Compound 186




¹H-NMR (CDCl₃) δ: 8.75-8.73 (1H,m), 8.34-8.32 (1H,m), 7.77-7.74 (1H,m), 7.61-7.48 (3H,m), 3.75 (3H,s), 3.37 (2H,q)

[0421]

Production Examples 121 and 122


The procedure was performed according to the method described in Production Examples 16 and 17 using 3-methyl-
5 2-[2-(2,2,2-trifluoroethylsulfanylphenyl)]-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 73 mg of 3-methyl-2-[2-(2,2,2-trifluoroethylsulfanylphenyl)]-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present
10 Compound 187) and 68 mg of 3-methyl-2-[2-(2,2,2-trifluoroethylsulfanylphenyl)]-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 188).

15 Present Compound 187



¹H-NMR (CDCl₃) δ: 8.77-8.75 (1H,m), 8.47-8.44 (1H,m), 8.29-8.26 (1H,m), 7.91-7.87 (1H,m), 7.80-7.74 (2H,m), 4.85-4.73 (1H,m), 4.03 (3H,s), 3.55-3.43 (1H,m)

20 Present Compound 187




$^1\text{H-NMR}$ (CDCl_3) δ : 8.77-8.75 (1H,m), 8.35-8.31 (2H,m), 7.95-7.84 (2H,m), 7.63-7.60 (1H,m), 4.61-4.47 (2H,m), 3.75 (3H,s)

5 [0422]

Production Examples 123 and 124


The procedure was performed according to the method described in Production Examples 16 and 17 using 2-(2-ethylsulfonylphenyl)-3-methyl-6-(trifluoromethylsulfonyl)-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfonylphenyl)-3-methyl-6-(trifluoromethylsulfonyl)-3H-imidazo[4,5-b]pyridine to give 2-(2-ethylsulfonylphenyl)-6-(trifluoromethylsulfonyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 209) and 2-(2-ethylsulfinylphenyl)-6-(trifluoromethylsulfonyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 210).

Present Compound 209



¹H-NMR (CDCl₃) δ: 8.68 (1H,d), 8.36 (1H,d), 8.21 (1H,dd),
7.87-7.77 (2H,m), 7.59 (1H,dd), 3.71 (3H,s), 3.44 (2H,q),
1.24 (3H,t)

Present Compound 210



¹H-NMR (CDCl₃) δ: 8.69 (1H,d), 8.38 (1H,d), 8.25 (1H,dd),
7.86-7.80 (1H,m), 7.72-7.67 (1H,m), 7.60 (1H,dd), 3.88
(3H,s), 3.43-3.31 (1H,m), 3.03-2.92 (1H,m), 1.31 (3H,t)


[0423]

10 Production Examples 125 and 126

The procedure was performed according to the method described in Production Examples 16 and 17 using 2-(2-ethylsulfonylphenyl)-6-trifluoromethylsulfonyl-3-methyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-tert-butylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine to give 2-(2-ethylsulfonylphenyl)-6-trifluoromethylsulfonyl-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 214) and 2-(2-ethylsulfonylphenyl)-6-trifluoromethylsulfonyl-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 215).


Present Compound 214

500



$^1\text{H-NMR}$ (CDCl_3) δ : 8.77 (1H,d), 8.55 (1H,d), 8.24 (1H,dd), 7.90-7.83 (2H,m), 7.61 (1H,dd), 3.75 (3H,s), 3.43 (2H,q), 1.26 (3H,t)

5 Present Compound 215



$^1\text{H-NMR}$ (CDCl_3) δ : 9.05 (1H,d), 8.65 (1H,d), 8.26-8.23 (1H,m), 7.90-7.85 (2H,m), 7.61-7.57 (1H,m), 3.77 (3H,s), 3.41 (2H,q), 1.27 (3H,t)

10 [0424]


Production Example 127

To a mixture of 2-[2-fluoro-4-(trifluoromethyl)phenyl]-6-(trifluoromethyl)thiazolo[5,4-b]pyridine (0.18 g) and DMF (2 ml), sodium thiomethoxide (63 mg) was added under ice-cooling, and stirred at room temperature for 4 hours. Into the reaction mixture, water was poured, and extracted with ethyl acetate. The organic layer was washed with water, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The

15

resulting residue was subjected to silica gel column chromatography to give 0.11 g of 2-[2-ethylsulfanyl-4-(trifluoromethyl)phenyl]-6-(trifluoromethyl)thiazolo[5,4-b]pyridine (hereinafter referred to as Present Compound 267).

Present Compound 267



$^1\text{H-NMR}$ (CDCl_3) δ : 8.90 (1H,d), 8.61 (1H,d), 8.14 (1H,d), 7.75 (1H,s), 7.58 (1H,d), 3.04 (2H,q), 1.38 (3H,t).


[0425]

Production Example 128

To a mixture of 2-[2-ethylsulfanyl-4-(trifluoromethyl)phenyl]-6-(trifluoromethyl)thiazolo[5,4-b]pyridine (0.11 g) and chloroform (3 ml), 3-chloroperbenzoic acid (purity: not less than 65 %) (0.13 g) was added, and stirred at room temperature for 12 hours. The reaction mixture was diluted with chloroform, washed with 10 % aqueous sodium thiosulfate solution and saturated aqueous sodium hydrogen carbonate solution, sequentially, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.11 g of 2-[2-ethylsulfonyl-4-(trifluoromethyl)phenyl]-6-

(trifluoromethyl)thiazolo[5,4-b]pyridine (hereinafter referred to as Present Compound 251).

Present Compound 251




5 $^1\text{H-NMR}$ (CDCl_3) δ : 8.95 (1H,s), 8.55 (1H,s), 8.52 (1H,s), 8.07 (1H,d), 7.88 (1H,d), 3.77 (2H,q), 1.40 (3H,t).

[0426]

Production Example 129

To a mixture of 2-(2-ethylsulfonyl-phenyl)-3-methyl-
10 3H-imidazo[4,5-b]pyridin-6-thiol (535 mg), iodomethane (166 μl), and ethanol (5 ml), potassium hydroxide (200 mg) was added at room temperature, and stirred for 5 hours. To the reaction mixture, saturated aqueous ammonium chloride solution was added, and extracted with ethyl acetate. The
15 combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 515 mg of 2-(2-ethylsulfonyl-phenyl)-3-methyl-6-methylsulfonyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as
20 Present Compound 253).

Present Compound 253




$^1\text{H-NMR}$ (CDCl_3) δ : 8.46 (1H,d), 8.10 (1H,d), 7.52-7.42 (3H,m), 7.37-7.26 (1H,m), 3.73 (3H,s), 2.86 (2H,q), 2.54 (3H,s), 1.22 (3H,t)

5 [0427]

Production Example 130

To a mixture of 2-(2-ethylsulfanylphenyl)-3-methyl-6-methylsulfanyl-3H-imidazo[4,5-b]pyridine (363 mg) and chloroform (5 ml), 69-75 % of 3-chloroperbenzoic acid (1.13 g) was added under ice-cooling. The mixture was heated to room temperature, and stirred for 5 hours, and then saturated aqueous sodium hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution were poured, and extracted with chloroform. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 356 mg of 2-(2-ethylsulfonylphenyl)-6-methylsulfonyl-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Present Compound 254).

Present Compound 254



$^1\text{H-NMR}$ (CDCl_3) δ : 9.02 (1H,d), 8.58 (1H,d), 8.23 (1H,dd), 7.90-7.81 (2H,m), 7.59 (1H,dd), 3.74 (3H,s), 3.42 (2H,q), 3.19 (3H,s), 1.26 (3H,t).

5 [0428]


Production Example 131-1

A mixture of 6-bromo-2-(2-ethylsulfanyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (1.02 mg), copper iodide (280 mg), 1,10-phenanthroline (268 mg), benzyl alcohol (456 μL), cesium carbonate (1.9 g), NMP (4 mL), and xylene (20 mL) was stirred with heating at 150 $^\circ\text{C}$ for 8 hours. After the mixture was allowed to stand to cool to room temperature, into the mixture, 28 % of aqueous ammonia and saturated aqueous sodium hydrogen carbonate solution were poured, extracted with tert-butyl methyl ether. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 402 mg of 6-benzyloxy-2-(2-ethylsulfanyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter referred to as Compound 268A).

10

15

20




$^1\text{H-NMR}$ (CDCl_3) δ : 8.27 (1H,d), 7.67 (1H,d), 7.45-7.33 (9H,m), 5.16 (2H,s), 3.70 (3H,s), 2.85 (2H,q), 1.22 (3H,t).
[0429]

5 Production Example 131-2

To a mixture of 6-benzyloxy-2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine 262mg, and chloroform (3 mL), boron tribromide (1M in dichloromethane) (698 μL) was added under ice-cooling, and stirred for 3
10 hours. Into the mixture, saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with chloroform then ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel
15 column chromatography to give 168 mg of 2-(2-ethylsulfanylphenyl)-3-methyl-3H-imidazo[4,5-b]pyridine-6-ol (hereinafter, Present Compound 268).

Present Compound 268




¹H-NMR (DMSO-D₆) δ: 9.58 (1H, brs), 8.02 (1H, d), 7.60-7.54 (2H, m), 7.47 (1H, d), 7.41 (1H, d), 7.38-7.33 (1H, m), 3.54 (3H, s), 2.94 (2H, q), 1.16 (3H, t).

[0430]

5 Production Example 132

To a mixture of 2-(2-ethylsulfanyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridine-6-ol (579 mg), chloroform (20 mL), and acetonitrile (6 mL), 69 % of 3-chloroperbenzoic acid (1.05 g) was added under ice-cooling. The mixture was
10 heated to room temperature, stirred for 3 hours, and filtered. The resulting filtered substance was washed with chloroform to give 566 mg of 2-(2-ethylsulfonyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridine-6-ol (hereinafter, Present Compound 269).

15 Present Compound 269




[0431]

Production Example 133-1

To a mixture of 2-(2-ethylsulfonyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridine-6-ol (566 mg) and DMF (5 mL),
20 60 % of sodium hydride (in oil, 129 mg) was added, and stirred with heating at 50 °C for 1 hour. After the mixture

was allowed to stand to cool to room temperature, to the mixture, carbon disulfide (1.08 mL) was added, and stirred at room temperature (2.5 hours). To a mixture, iodomethane (446 μ L) was added, and stirred at room temperature for 1.5 hours. Into the mixture, saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 681 mg of S-methyl O-[2-(2-ethylsulfonyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridine-6-yl]dithiocarboxylate (hereinafter referred to as Compound 270A).

Compound 270A



$^1\text{H-NMR}$ (CDCl_3) δ : 8.30-8.25 (1H,m), 8.24-8.18 (1H,m), 7.85-7.76 (3H,m), 7.58-7.53 (1H,m), 3.69 (3H,brs), 3.44 (2H,q), 2.71 (3H,brs), 1.26-1.22 (3H,m).


[0432]

Production Example 133-2

To a mixture of S-methyl O-[2-(2-ethylsulfonyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridine-6-yl]dithiocarboxylate (371 mg) and chloroform (3 mL), HF-


pyridine complex (1.25 mL) and 1,3-dibromo-5,5-dimethylhydantoin (1.58 g) were added with cooling at -55 °C. The mixture was stirred for 7 hours with heating to room temperature gradually, and then saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with chloroform. The combined organic layer was washed with aqueous sodium thiosulfate solution, dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to reversed-phase silica gel column chromatography to give 29 mg of 6-(bromo-difluoromethoxy)-2-(2-ethylsulfonyl-phenyl)-3-methyl-3H-imidazo[4,5-b]pyridine (hereinafter, Present Compound 270) and 76 mg of 2-(2-ethylsulfonyl-phenyl)-3-methyl-6-trifluoromethoxy-3H-imidazo[4,5-b]pyridine (hereinafter, Present Compound 271).

Present Compound 270



¹H-NMR (CDCl₃) δ: 8.43 (1H,d), 8.26-8.21 (1H,m), 7.97-7.97 (1H,m), 7.87-7.80 (2H,m), 7.56 (1H,dd), 3.70 (3H,s), 3.45 (2H,q), 1.26 (3H,t).

Present Compound 271



$^1\text{H-NMR}$ (CDCl_3) δ : 8.42 (1H,d), 8.22 (1H,dd), 7.94-7.94 (1H,m), 7.87-7.79 (2H,m), 7.56 (1H,dd), 3.70 (3H,s), 3.44 (2H,q), 1.25 (3H,t).

5 [0433]

Production Example 134


The procedure was performed according to the method described in Production Example 103 using 2-(2-ethylsulfonyl-4-trifluoromethylphenyl)-6-iodo-3-methyl-3H-imidazo[4,5-b]pyridine instead of 2-(2-ethylsulfonyl-phenyl)-6-iodo-3-methyl-3H-imidazo[4,5-b]pyridine to give Compound 301A.

A mixture of Compound 301A (0.94 g) and DMF (13 ml) was cooled to $-50\text{ }^\circ\text{C}$, and bubbled with excessive amounts of CF_3I gas to dissolve the compound in DMF. To the mixture, tetrakis(dimethylamino)ethylenediamine (1.2 ml) was added dropwise at a rate that the internal temperature did not exceed $-40\text{ }^\circ\text{C}$. Then, the mixture was heated to $-10\text{ }^\circ\text{C}$ over 1 hour, and stirred at $-10\text{ }^\circ\text{C}$ for 1 hour. Into the reaction mixture, water was poured, heated to room temperature, and extracted with ethyl acetate. The combined organic layer was dried over magnesium sulfate, and concentrated under

reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.68 g of 2-(2-ethylsulfanyl-4-trifluoromethylphenyl)-3-methyl-6-trifluoromethylsulfanyl-3H-imidazo[4,5-b]pyridine

5 (hereinafter, Present Compound 301).

Present Compound 301



$^1\text{H-NMR}$ (CDCl_3) δ : 8.66 (1H,d) 8.39 (1H,d), 7.67-7.64 (1H,m), 7.59-7.52 (2H,m), 3.75 (3H,s), 2.94 (2H,q), 1.27 (3H,t).

10 [0434]

Production Example 135


To a mixture of 2-(2-ethylsulfanyl-4-trifluoromethylphenyl)-3-methyl-6-trifluoromethylsulfanyl-3H-imidazo[4,5-b]pyridine and chloroform (5 ml), 69 % of 3-chloroperbenzoic acid (1.05 g) was added under ice-cooling, and then heated to room temperature, and stirred for 1.5 hours. Then, saturated aqueous sodium hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution was poured, and extracted with chloroform. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give

15

20

0.20 g of 2-(2-ethylsulfonyl-4-trifluoromethylphenyl)-3-methyl-6-trifluoromethylsulfinyl-3H-imidazo[4,5-b]pyridine (hereinafter, Present Compound 306).

Present Compound 306



¹H-NMR (CDCl₃) δ: 8.71-8.70 (1H,m), 8.50-8.49 (1H,m), 8.38-8.36 (1H,m), 8.12-8.08 (1H,m), 7.74-7.71 (1H,m), 3.72 (3H,s), 3.49 (2H,q), 1.29 (3H,t).


[0435]

10 Production Example 136

A mixture of 2-(2-ethylsulfonyl-4-trifluoromethylphenyl)-3-methyl-6-trifluoromethylsulfinyl-3H-imidazo[4,5-b]pyridine (0.26 g), sodium tungstate dihydrate (36 mg), 30 % of aqueous hydrogen peroxide solution (1 ml) and acetonitrile (5 ml) was stirred under reflux for 4.5 hours. The mixture was cooled to room temperature. Into the reaction mixture, water was poured, and extracted with ethyl acetate. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.24 g of 2-(2-ethylsulfonyl-4-trifluoromethylphenyl)-3-methyl-6-

trifluoromethylsulfonyl-3H-imidazo[4,5-b]pyridine
(hereinafter, Present Compound 302).

Present Compound 302



5 $^1\text{H-NMR}$ (CDCl_3) δ : 9.08-9.07 (1H,m), 8.68-8.66 (1H,m), 8.52-8.50 (1H,m), 8.16-8.12 (1H,m), 7.76-7.73 (1H,m), 3.78 (3H,s), 3.46 (2H,q), 1.30 (3H,t)


[0436]

Production Example 137-1

10 To a mixture of 4-(1,2,2,2-tetrafluoro-1-trifluoromethyl-ethyl)-benzene-1,2-diamine (552 mg), 2-ethylsulfanylbenzoic acid (401 mg), and pyridine (27 mL), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (422 mg) and 1-hydroxybenzotriazole (27 mg) were added at
15 room temperature. The reaction mixture was stirred at room temperature for 5 hours, diluted with water, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was dissolved in a mixed
20 solution of DMF (7.5 mL) and toluene (30 mL), and then p-toluenesulfonic acid (837 mg) was added at room temperature. The mixture was stirred with heating at 130 °C for 8 hours,

and allowed to stand to cool to room temperature. Into the reaction mixture, saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 97 mg of 2-(2-ethylsulfanyl-phenyl)-5-(1,2,2,2-tetrafluoro-1-trifluoromethyl-ethyl)-1H-benzimidazole (hereinafter, Present Compound 346).

10 Present Compound 346



¹H-NMR (CDCl₃) δ: 12.08-11.87 (1H,m), 8.31 (1H,s), 8.12-7.44 (4H,m), 7.42-7.30 (2H,m), 2.86 (2H,q), 1.22 (3H,t).


[0437]

15 Production Example 137-2

To a mixture of 2-(2-ethylsulfanyl-phenyl)-5-(1,2,2,2-tetrafluoro-1-trifluoromethyl-ethyl)-1H-benzimidazole (97 mg) and DMF (10 mL), sodium hydride (11 mg) and iodomethane (16 μL) were added under ice-cooling. The mixture was heated to room temperature, and stirred for 5 hours. The reaction mixture was diluted with water, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure.


The residue was subjected to silica gel column chromatography to give 35 mg of 2-(2-ethylsulfanyl-phenyl)-1-methyl-6-(1,2,2,2-tetrafluoro-1-trifluoromethyl-ethyl)-1H-benzimidazole (hereinafter, Present Compound 304) and 80
5 mg of 2-(2-ethylsulfanyl-phenyl)-1-methyl-5-(1,2,2,2-tetrafluoro-1-trifluoromethyl-ethyl)-1H-benzimidazole (hereinafter, Present Compound 305).

Present Compound 304



10 $^1\text{H-NMR}$ (CDCl_3) δ : 7.93 (1H,d), 7.68 (1H,s), 7.55-7.44 (4H,m), 7.37-7.29 (1H,m), 3.71 (3H,s), 2.87 (2H,q), 1.24 (3H,t).

Present Compound 305




15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.12 (1H,s), 7.61-7.41 (5H,m), 7.36-7.30 (1H,m), 3.68 (3H,s), 2.86 (2H,q), 1.23 (3H,t).

[0438]

Production Example 138-1

To a mixture of 2-nitro-4-(pentafluorosulphur)-

phenylamine (2.0 g) and DMF (15 mL), sodium hydride (313 mg) and iodomethane (447 μ L) were added under ice-cooling. The mixture was heated to room temperature, and stirred for 5 hours. The reaction mixture was diluted with water, and then the precipitated powder was collected by filtration. The resulting powder was washed with hexane to give 2.0 g of methyl-(2-nitro-4-(pentafluorosulphur)-phenyl)-amine. methyl-[2-nitro-4-(pentafluorosulphur)-phenyl]-amine




$^1\text{H-NMR}$ (CDCl_3) δ : 8.60 (1H,d), 8.29 (1H,brs), 7.78 (1H,dd), 6.89 (1H,d), 3.10 (3H,d).

[0439]

Production Example 138-2

To a mixture of methyl-[2-nitro-4-(pentafluorosulphur)-phenyl]-amine (2.0 g), acetic acid (1.9 mL), ethanol (80 mL), and water (20 mL), iron powder (1.5 g) was added with stirring with heating at 80 $^{\circ}\text{C}$. The reaction mixture was stirred with heating at 80 $^{\circ}\text{C}$ for 6 hours, and filtered through Celite[®]. The resulting filtrate was concentrated under reduced pressure, and diluted with 1N aqueous sodium hydroxide solution. The mixed solution was extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under

reduced pressure. The residue was subjected to silica gel column chromatography to give 1.6 g of N¹-methyl-4-(pentafluorosulphur)-benzene-1,2-diamine (Compound (1F)-35).
N¹-methyl-4-(pentafluorosulphur)-benzene-1,2-diamine



5

[0440]

Production Example 138-3

To a mixture of N¹-methyl-4-(pentafluorosulphur)-benzene-1,2-diamine (575 mg), 2-fluoro-4-trifluoromethyl-benzaldehyde (348 μ L), and DMF (5 mL), sodium hydrogen sulfite (290 mg) was added at room temperature. The mixture was heated to 100 °C, and stirred with heating for 10 hours. After the mixture was allowed to stand to cool to room temperature, the mixture was diluted with water, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was dissolved in DMF, and then sodium ethanethiolate (175 mg) was added at room temperature thereto. The mixture was stirred at room temperature for 8 hours, and then into the mixture, saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and


10

15

20

concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 537 mg of 2-(2-ethylsulfanyl-4-trifluoromethyl-phenyl)-1-methyl-5-(pentafluorosulphur)-1H-benzimidazole (hereinafter, Present Compound 314).

Present Compound 314



$^1\text{H-NMR}$ (CDCl_3) δ : 8.28 (1H,d), 7.79 (1H,dd), 7.67 (1H,s), 7.57 (2H,s), 7.46 (1H,d), 3.70 (3H,s), 2.93 (2H,q), 1.27 (3H,t).


[0441]

Production Example 139

To a mixture of 2-(2-ethylsulfanyl-4-trifluoromethyl-phenyl)-1-methyl-5-(pentafluorosulphur)-1H-benzimidazole (370 mg) and chloroform (10 mL), 69-75 % of 3-chloroperbenzoic acid (335 mg) was added under ice-cooling. The mixture was heated to room temperature, and stirred for 0.5 hours. Into the mixture, saturated aqueous sodium hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution were poured, and extracted with chloroform. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure.


The residue was subjected to silica gel column chromatography to give 115 mg of 2-(2-ethylsulfinyl-4-trifluoromethyl-phenyl)-1-methyl-5-(pentafluorosulphur)-1H-benzimidazole(hereinafter, Present Compound 315) and 244 mg of 2-(2-ethylsulfonyl-4-trifluoromethyl-phenyl)-1-methyl-5-(pentafluorosulphur)-1H-benzimidazole (hereinafter, Present Compound 316).

Present Compound 315



¹H-NMR (CDCl₃) δ: 8.54-8.49 (1H,m), 8.26-8.21 (1H,m), 7.95-7.89 (1H,m), 7.85-7.80 (1H,m), 7.74-7.68 (1H,m), 7.52-7.47 (1H,m), 3.82 (3H,s), 3.48-3.36 (1H,m), 3.06-2.94 (1H,m), 1.35-1.27 (3H,m).

Present Compound 316



¹H-NMR (CDCl₃) δ: 8.50-8.41 (1H,m), 8.24-8.16 (1H,m), 8.09-8.02 (1H,m), 7.83-7.65 (2H,m), 7.50-7.39 (1H,m), 3.64 (3H,s), 3.49-3.37 (2H,m), 1.30-1.24 (3H,m).

[0442]


Production Example 140

A mixture of 2-ethylsulfanylbenzoic acid (1.0 g), oxalyl chloride (0.7 ml), DMF (1 drop) and chloroform (4 ml) was stirred at room temperature for 1 hour, and concentrated under reduced pressure.

A mixture of the resulting residue, 2-amino-3-chloro-5-trifluoromethylpyridine (1.08 g) and N-ethyldiisopropylamine (0.85 g) was stirred at 140 °C for 2 hours. After the reaction mixture was allowed to stand to cool, into the mixture, water was poured, and extracted with ethyl acetate. The organic layer was washed with water, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure.

A mixture of the resulting residue, Lawesson's reagent (2.1 g), and chlorobenzene (10 ml) was refluxed for 3 hours. After the reaction mixture was allowed to stand to cool, the reaction mixture was concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.14 g of 2-(2-ethylsulfanylphenyl)-6-(trifluoromethyl)thiazolo[4,5-b]pyridine (hereinafter, Present Compound 328).

Present Compound 328




¹H-NMR (CDCl₃) δ: 9.02 (1H,d), 8.55 (1H,d), 8.35 (1H,dd), 7.60 (1H,dd), 7.49 (1H,td), 7.40 (1H,td), 2.99 (2H,q), 1.33 (3H,t).

[0443]

5 Production Example 141

To a mixture of 2-(2-ethylsulfanylphenyl)-6-(trifluoromethyl)thiazolo[4,5-b]pyridine (0.14 g) and chloroform (3 ml), 3-chloroperbenzoic acid (purity: not less than 65 %, 0.21 g) was added, and stirred at room temperature for 12 hours. The reaction mixture was diluted with chloroform, washed with 10 % of aqueous sodium thiosulfate solution then saturated aqueous sodium hydrogen carbonate solution, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 57 mg of 2-(2-ethylsulfonylphenyl)-6-(trifluoromethyl)thiazolo[4,5-b]pyridine (hereinafter, Present Compound 329).

Present Compound 329




¹H-NMR (CDCl₃) δ: 9.04 (1H,d), 8.62 (1H,d), 8.26 (1H,dd), 7.84-7.77 (2H,m), 7.71 (1H,dd), 3.81 (2H,q), 1.36 (3H,t).

[0444]

Production Example 142

To a mixture of sodium ethyl mercaptide (80 %, 0.63 g) and DMF (10 ml), a solution of 2-(4-bromo-2-fluorophenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (2.08 g) in DMF was added dropwise under ice-cooling, then heated to room temperature, and stirred for 30 minutes. Into the reaction mixture, saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with ethyl acetate. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 1.57 g of 2-(4-bromo-2-ethylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter, Present Compound 238).

Present Compound 238




[0445]

Production Example 143


To a mixture of 2-(4-bromo-2-ethylsulfanylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.40 g) and chloroform (5 ml), 3-chloroperbenzoic acid (purity: not less than 65 %, 0.29 g) was added under ice-cooling, then

heated to room temperature, and stirred for 2 hours. Into the reaction mixture, saturated aqueous sodium hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution were poured, and extracted with chloroform. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.26g of 2-(4-bromo-2-ethylsulfinylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine

(hereinafter, Present Compound 239) and 0.17 g of 2-(4-bromo-2-ethylsulfonylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter, Present Compound 240). Present Compound 239



Present Compound 240




[0446]

Production Example 144

A mixture of 2-(4-bromo-2-ethylsulfonylphenyl)-3-

methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.20 g),
2-tributylstannylpyrimidine (0.17 g),
tetrakis(triphenylphosphine)palladium (27 mg) and toluene
(5 ml) was heated at reflux under nitrogen atmosphere for
5 5.5 hours. After the mixture was cooled to room temperature,
to the mixture, 2-tributylstannylpyrimidine (0.17 g) and
tetrakis(triphenylphosphine)palladium (27 mg) were added,
and stirred under reflux for further 8 hours. After the
mixture was cooled to room temperature, water was poured
10 thereinto, and extracted with ethyl acetate. The combined
organic layer was dried over magnesium sulfate, and
concentrated under reduced pressure. The resulting residue
was subjected to silica gel column chromatography to give
0.20 g of 2-[2-ethylsulfonyl-4-(pyrimidine-2-yl)-phenyl]-3-
15 methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine
(hereinafter, Present Compound 333).

Present Compound 333




¹H-NMR (CDCl₃) δ: 8.77-8.74 (1H,m), 8.38-8.36 (1H,m), 8.30-
20 8.27 (1H,m), 8.00-7.95 (1H,m), 7.63-7.55 (1H,m), 7.43
(1H,d), 7.41-7.30 (2H,m), 3.72 (3H,s), 3.44 (2H,q), 1.28
(3H,t).

[0447]

Production Example 145-1

To a mixture of 2-chloro-1,3-dinitro-5-trifluoromethyl-benzene (20 g) and DMF (30 mL), 40 % of aqueous methylamine solution (15 mL) was added dropwise under ice-cooling. The mixture was stirred for 6 hours under ice-cooling, and then the reaction mixture was added to iced water. The precipitated powder was collected by filtration, and washed with water to give 20 g of (2,6-dinitro-4-trifluoromethyl-phenyl)-methyl-amine.

(2,6-dinitro-4-trifluoromethyl-phenyl)-methyl-amine




[0448]

Production Example 145-2

To a mixture of (2,6-dinitro-4-trifluoromethyl-phenyl)-methyl-amine (5.0 g), acetic acid (11 mL), ethanol (100 mL), and ice (20 g), iron powder (6.4 g) was added under ice-cooling. The reaction mixture was stirred at 0 °C for 3.5 hours, and filtered through Celite®. The resulting filtrate was concentrated under reduced pressure, and diluted with 1N aqueous sodium hydroxide solution. The solution was extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and

concentrated under reduced pressure to give 3.1 g of N'-methyl-5-trifluoromethyl-benzene-1,2,3-triamine.

N'-methyl-5-trifluoromethyl-benzene-1,2,3-triamine




5 [0449]

Production Example 145-3

To a mixture of N'-methyl-5-trifluoromethyl-benzene-1,2,3-triamine (1.3 g), 2-ethylsulfanylbobenzoic acid (1.3 g), and pyridine (12 mL), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (1.4 g) and
10 1-hydroxybenzotriazole (86 mg) were added at room temperature. The mixture was stirred at room temperature for 5 hours, then heated to 110 °C, and stirred with heating for 8 hours. After the reaction mixture was allowed
15 to cool to room temperature, the reaction mixture was diluted with water, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 339
20 mg of 2-(2-ethylsulfanyl-phenyl)-3-methyl-6-trifluoromethyl-3H-benzimidazole-4-ylamine (hereinafter, Present Compound 336).

Present Compound 336




$^1\text{H-NMR}$ (CDCl_3) δ : 7.55 (1H, s), 7.50–7.36 (3H, m), 7.33–7.28 (1H, m), 6.77 (1H, d), 4.04 (2H, brs), 3.91 (3H, s), 2.83 (2H, q), 1.21 (3H, t).

5 [0450]

Production Example 146

To a mixture of sodium ethyl mercaptide (80 %, 0.33 g) and DMF (4 ml), a solution of 2-(2-fluoro-4-(pentafluorosulphur)phenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (1.19 g) in DMF was added dropwise under ice-cooling, then heated to room temperature, and stirred at room temperature for 2 hours. Into the reaction mixture, saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with ethyl acetate. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 1.21 g of 2-(2-ethylsulfanyl-4-(pentafluorosulphur)phenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter, Present Compound 341).
Present Compound 341

10
15
20



$^1\text{H-NMR}$ (CDCl_3) δ : 8.75-8.74 (1H,m), 8.36-8.34 (1H,m), 7.81 (1H,d), 7.73 (1H,dd), 7.58-7.54 (1H,m), 3.80 (3H,s), 2.95 (2H,q), 1.29 (3H,t).


5 [0451]

Production Example 147

To a mixture of 2-(2-ethylsulfonyl-4-(pentafluorosulphur)phenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (0.90 g) and chloroform (6 ml), 3-chloroperbenzoic acid (purity: not less than 65 %, 0.68 g)
10 was added under ice-cooling, then heated to room temperature, and stirred for 1 hour. Into the reaction mixture, saturated aqueous sodium hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution
15 were poured, and extracted with chloroform. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.45 g of 2-(2-ethylsulfonyl-4-(pentafluorosulphur)phenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine
20 (hereinafter, Present Compound 342) and 0.44 g of 2-(2-ethylsulfonyl-4-(pentafluorosulphur)phenyl)-3-methyl-6-


trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter, Present Compound 343).

Present Compound 342



- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 8.80-8.78 (1H,m), 8.65 (1H,d), 8.35-8.34 (1H,m), 8.07 (1H,dd), 7.78-7.75 (1H,m), 3.95 (3H,s), 3.55-3.45 (1H,m), 3.10-3.00 (1H,m), 1.36 (3H,t).

Present Compound 343



- 10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.78-8.76 (1H,m), 8.61 (1H,d), 8.32-8.30 (1H,m), 8.22 (1H,dd), 7.74-7.70 (1H,m), 3.75 (3H,s), 3.50 (2H,q), 1.30 (3H,t).


[0452]

Production Example 148-1

- 15 To a mixture of 5-trifluoromethyl-pyridine-2-ylamine (65 g) and chloroform (100 mL), N-bromosuccinimide (71 g) was added in 5 parts under ice-cooling. The mixture was heated to room temperature, and stirred for 1 hour, then heated to 80 °C, and stirred with heating for 30 minutes.

After the mixture was allowed to cool to room temperature, into the mixture, saturated aqueous sodium thiosulfate solution and saturated aqueous sodium hydrogen carbonate solution were poured, and extracted with chloroform. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 96 g of 3-bromo-5-trifluoromethyl-pyridine-2-ylamine.

3-bromo-5-trifluoromethyl-pyridine-2-ylamine



$^1\text{H-NMR}$ (CDCl_3) δ : 8.27 (1H,d), 7.86 (1H,d), 5.38 (2H,brs).


[0453]

Production Example 148-2

To an autoclave reactor, 3-bromo-5-trifluoromethyl-pyridine-2-ylamine (40 g), acetylacetonate copper(II) (2.2 g), acetylacetonate (6.6 g), cesium carbonate (59 g), and NMP (105 mL) were added, and then 28 % of aqueous ammonia solution (25 mL) was added under ice-cooling. After the reactor was sealed, the mixture heated to 110 °C, and stirred with heating for 12 hours. The mixture was ice-cooled to room temperature, then diluted with water, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel

column chromatography to give 15g 5-trifluoromethyl-pyridine-2,3-diamine (Compound (1F)-22).

5-trifluoromethyl-pyridine-2,3-diamine




5 $^1\text{H-NMR}$ (CDCl_3) δ : 7.93 (1H,d), 7.04 (1H,d), 4.71 (2H,brs), 3.46 (2H,brs).

[0454]

Production Example 148-3

To a mixture of 5-trifluoromethyl-pyridine-2,3-diamine
10 (8.6 g), 2-formyl-5-trifluoromethylphenylethylsulfide (11 g), and DMF (67 mL), sodium hydrogen sulfite (6.1 g) was added at room temperature. After the mixture was stirred with heating at 100 °C for 3 hours, to the mixture, copper (II) chloride dihydrate (1 g) was added, and stirred at 100
15 °C for further 1 hour. After the reaction mixture was allowed to cool to room temperature, the reaction mixture was added to water, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was
20 subjected to silica gel column chromatography to give a yellow powdered solid. The solid was washed with hot hexane to give 12 g of 2-(2-ethylsulfanyl-4-trifluoromethyl-phenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter, Present Compound 344).

Present Compound 344




$^1\text{H-NMR}$ (CDCl_3) δ : 12.79 (1H,brs), 8.72 (1H,brs), 8.49-8.34 (2H,m), 7.79 (1H,s), 7.64 (1H,d), 3.00 (2H,q), 1.31 (3H,t).

5 [0455]

Production Example 149

To a mixture of 2-(2-ethylsulfanyl-4-trifluoromethyl-phenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (12 g) and chloroform (111 mL), 69-75 % of 3-chloroperbenzoic acid
10 (8.0 g) was added under ice-cooling. The mixture was heated to room temperature, and stirred for 0.5 hours. Into the mixture, saturated aqueous sodium hydrogen carbonate solution and saturated aqueous sodium thiosulfate solution were poured, and extracted with chloroform. The combined
15 organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 9.1 g of 2-(2-ethylsulfonyl-4-trifluoromethyl-phenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter,
20 Present Compound 345).

Present Compound 345



¹H-NMR (DMSO-D₆) δ: 14.15 (1H, brs), 8.83 (1H, s), 8.58 (1H, s), 8.41 (1H, d), 8.37 (1H, s), 8.19 (1H, d), 3.97 (2H, q), 1.23 (3H, t).


5 [0456]

Production Example 150

A mixture of N-(2-amino-5-trifluoromethylpyridine-3-yl)-2-ethylsulfanyl-benzamide (200 mg), tert-butylalcohol (1 ml), and THF (9 ml) was stirred with heating at 80 °C, then 60 % of sodium hydride (in oil, 56 mg) was added thereto. The mixture was stirred with heating at 80 °C for 2 hours, and then 60 % of sodium hydride (in oil, 56 mg) was added. The mixture was further stirred with heating at the same temperature for 2 hours, and then to the mixture, 60 % of sodium hydride (in oil, 56 mg) was added, and stirred with heating at the same temperature for 2 hours. The reaction mixture was cooled to room temperature, and evaporated to remove the solvent, and then to the mixture, saturated ammonium chloride solution was added, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 132 mg of 2-(2-

ethylsulfanyphenyl)-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (hereinafter, Present Compound 351).

Present Compound 351




¹H-NMR (CDCl₃) δ: 8.67 (1H,d), 8.51-8.48 (1H,m), 8.33 (1H,d), 7.66-7.61 (1H,m), 7.51-7.46 (2H,m), 2.93 (2H,q), 1.27 (3H,t)

[0457]

The above present compounds and present compounds produced by a similar production method to those of the above present compounds are shown in the following tables.

[0458]

The compound of the formula (1-2):



wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, A¹, A², A³, and n represent a combination shown in Tables 20-34.

[0459]

Table 20

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
1	Me	H	H	H	H	CF ₃	H	NMe	CH	N	0
2	Me	H	H	H	H	CF ₃	H	NMe	CH	N	1

3	Me	H	H	H	H	CF ₃	H	NMe	CH	N	2
4	Et	H	H	H	H	CF ₃	H	NMe	CH	N	0
5	Et	H	H	H	H	CF ₃	H	NMe	CH	N	1
6	Et	H	H	H	H	CF ₃	H	NMe	CH	N	2
7	Pr	H	H	H	H	CF ₃	H	NMe	CH	N	0
8	Pr	H	H	H	H	CF ₃	H	NMe	CH	N	1
9	Pr	H	H	H	H	CF ₃	H	NMe	CH	N	2
10	CH ₂ =CHCH ₂	H	H	H	H	CF ₃	H	NMe	CH	N	0
11	CH ₂ =CHCH ₂	H	H	H	H	CF ₃	H	NMe	CH	N	2
12	iPr	H	H	H	H	CF ₃	H	NMe	CH	N	0
13	iPr	H	H	H	H	CF ₃	H	NMe	CH	N	1
14	iPr	H	H	H	H	CF ₃	H	NMe	CH	N	2
15	tBu	H	H	H	H	CF ₃	H	NMe	CH	N	0
16	tBu	H	H	H	H	CF ₃	H	NMe	CH	N	1
17	tBu	H	H	H	H	CF ₃	H	NMe	CH	N	2
18	CF ₃	H	H	H	H	CF ₃	H	NMe	CH	N	0
19	CF ₃	H	H	H	H	CF ₃	H	NMe	CH	N	1
20	CF ₃	H	H	H	H	CF ₃	H	NMe	CH	N	2
21	Et	H	H	H	H	CF ₃	H	NEt	CH	N	0
22	Et	H	H	H	H	CF ₃	H	NEt	CH	N	1
23	Et	H	H	H	H	CF ₃	H	NEt	CH	N	2
24	Et	H	H	H	H	CF ₃	H	NPr	CH	N	0
25	Et	H	H	H	H	CF ₃	H	NPr	CH	N	1

[0460]

Table 21

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
26	Et	H	H	H	H	CF ₃	H	NPr	CH	N	2
27	Et	H	H	H	H	CF ₃	H	NiPr	CH	N	0
28	Et	H	H	H	H	CF ₃	H	NiPr	CH	N	1
29	Et	H	H	H	H	CF ₃	H	NiPr	CH	N	2
30	Et	H	H	H	H	CF ₃	H	NCycPr	CH	N	0
31	Et	H	H	H	H	CF ₃	H	NCycPr	CH	N	1
32	Et	H	H	H	H	CF ₃	H	NCycPr	CH	N	2
33	Et	H	H	H	H	CF ₃	H	NCH ₂ OEt	CH	N	0
34	Et	H	H	H	H	H	CF ₃	NCH ₂ OEt	N	CH	0
35	Et	H	H	H	H	CF ₃	H	NCH ₂ OEt	CH	N	1
36	Et	H	H	H	H	CF ₃	H	NCH ₂ OEt	CH	N	2
37	Et	H	H	H	H	CF ₃	H	NCH ₂ OMe	CH	N	0
38	Et	H	H	H	H	CF ₃	H	NCH ₂ OMe	CH	N	2
39	Et	H	H	H	H	Me	H	NMe	CH	N	0
40	Et	H	H	H	H	Me	H	NMe	CH	N	1
41	Et	H	H	H	H	Me	H	NMe	CH	N	2
42	Et	H	H	H	H	Br	H	NMe	CH	N	0
43	Et	H	H	H	H	Br	H	NMe	CH	N	1
44	Et	H	H	H	H	Br	H	NMe	CH	N	2
45	Et	H	H	H	H	I	H	NMe	CH	N	0
46	Et	H	H	H	H	I	H	NMe	CH	N	1
47	Et	H	H	H	H	I	H	NMe	CH	N	2

48	Et	H	H	H	H	CN	H	NMe	CH	N	0
49	Et	H	H	H	H	CN	H	NMe	CH	N	1
50	Et	H	H	H	H	CN	H	NMe	CH	N	2

[0461]

Table 22

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
51	Et	H	H	H	H	CHO	H	NMe	CH	N	0
52	Et	H	H	H	H	CF ₂ H	H	NMe	CH	N	0
53	Et	H	H	H	H	CF ₂ H	H	NMe	CH	N	1
54	Et	H	H	H	H	CF ₂ H	H	NMe	CH	N	2
55	Et	H	H	H	H	C ₂ F ₅	H	NMe	CH	N	0
56	Et	H	H	H	H	Ph	H	NMe	CH	N	0
57	Et	H	H	H	H	Ph	H	NMe	CH	N	2
58	Et	H	H	H	H	2-F-Ph	H	NMe	CH	N	0
59	Et	H	H	H	H	2-F-Ph	H	NMe	CH	N	1
60	Et	H	H	H	H	2-F-Ph	H	NMe	CH	N	2
61	Et	H	H	H	H	3-F-Ph	H	NMe	CH	N	0
62	Et	H	H	H	H	3-F-Ph	H	NMe	CH	N	1
63	Et	H	H	H	H	3-F-Ph	H	NMe	CH	N	2
64	Et	H	H	H	H	4-F-Ph	H	NMe	CH	N	0
65	Et	H	H	H	H	4-F-Ph	H	NMe	CH	N	2
66	Et	H	H	H	H	H	CF ₃	NMe	N	CH	0
67	Me	H	H	H	H	CF ₃	H	NMe	CH	CH	0
68	Et	H	H	H	H	CF ₃	H	NMe	CH	CH	0
69	Et	H	H	H	H	CF ₃	H	NMe	CH	CH	1
70	Et	H	H	H	H	CF ₃	H	NMe	CH	CH	2
71	Et	H	H	H	H	CF ₃	H	NMe	CH	CCl	0
72	Et	H	H	H	H	CF ₃	H	NMe	CH	CCl	1
73	Et	H	H	H	H	CF ₃	H	NMe	CH	CCl	2
74	Et	H	H	H	H	CF ₃	H	NMe	CH	CBr	0
75	Et	H	H	H	H	CF ₃	H	NMe	CH	CBr	1

[0462]

Table 23

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
76	Et	H	H	H	H	CF ₃	H	NMe	CH	CBr	2
77	Me	H	H	H	H	CF ₃	H	O	CH	CH	0
78	Et	H	H	H	H	CF ₃	H	O	CH	CH	0
79	Et	H	H	H	H	CF ₃	H	O	CH	CH	1
80	Et	H	H	H	H	CF ₃	H	O	CH	CH	2
81	Et	H	H	H	H	CF ₃	H	O	CH	N	0
82	Et	H	H	H	H	CF ₃	H	O	CH	N	1
83	Et	H	H	H	H	CF ₃	H	O	CH	N	2
84	Me	H	H	H	H	CF ₃	H	S	CH	CH	0
85	Et	H	H	H	H	CF ₃	H	S	CH	CH	0
86	Et	H	H	H	H	CF ₃	H	S	CH	CH	1

87	Et	H	H	H	H	CF ₃	H	S	CH	CH	2
88	Et	Cl	H	H	H	CF ₃	H	NMe	CH	N	0
89	Et	Cl	H	H	H	CF ₃	H	NMe	CH	N	1
90	Et	Cl	H	H	H	CF ₃	H	NMe	CH	N	2
91	Et	H	H	H	H	C(O)CF ₃	H	NMe	CH	N	0
92	Me	H	H	H	SMe	CF ₃	H	NMe	CH	N	0
93	Me	H	H	H	SOMe	CF ₃	H	NMe	CH	N	1
94	Me	H	H	H	SO ₂ Me	CF ₃	H	NMe	CH	N	2
95	Et	H	H	H	SEt	CF ₃	H	NMe	CH	N	0
96	Et	H	H	H	SOEt	CF ₃	H	NMe	CH	N	1
97	Et	H	H	H	SO ₂ Et	CF ₃	H	NMe	CH	N	2
98	Et	H	H	H	H	Cl	H	NMe	CH	N	0
99	Et	H	H	H	H	Cl	H	NMe	CH	N	1
100	Et	H	H	H	H	Cl	H	NMe	CH	N	2

[0463]

Table 24

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
101	Et	H	H	H	H	Br	H	NMe	CCHO	N	0
102	Et	H	H	SEt	H	CF ₃	H	NMe	CH	N	0
103	Et	H	H	H	H	CF ₃	H	NCH ₂ OEt	CH	C H	0
104	Et	H	H	H	H	H	CF ₃	NCH ₂ OEt	N	C H	0
105	Et	H	H	H	H	CF ₃	H	NCH ₂ CO ₂ Me	CH	N	0
106	Et	H	H	H	H	CF ₃	H	NCH ₂ CO ₂ Et	CH	N	0
107	Et	H	H	H	H	CF ₃	H	N(CH ₂) ₂ OMe	CH	N	0
108	Et	H	H	H	H	CF ₃	H	NCH ₂ SMe	CH	N	0
109	Et	H	H	H	H	CF ₃	H	N(CH ₂) ₂ SM	CH	N	0
110	Et	H	H	H	H	CF ₃	H	NBu	CH	N	0
111	Et	H	H	H	H	CF ₃	H	NCO ₂ tBu	CH	N	0
112	Et	H	H	H	H	C ₂ F ₅	H	NMe	CH	N	1
113	Et	H	H	H	H	C ₂ F ₅	H	NMe	CH	N	2
114	Et	H	H	H	H	CH(OH)CF ₃	H	NMe	CH	N	0
115	Et	H	H	H	H	CHF ₂ CF ₃	H	NMe	CH	N	0
116	Et	H	F	H	H	CF ₃	H	NMe	CH	N	0
117	Et	H	F	H	H	CF ₃	H	NMe	CH	N	1
118	Et	H	F	H	H	CF ₃	H	NMe	CH	N	2
119	Et	OMe	H	H	H	CF ₃	H	NMe	CH	N	0
120	Et	OMe	H	H	H	CF ₃	H	NMe	CH	N	1
121	Et	H	OMe	H	H	CF ₃	H	NMe	CH	N	0
122	Et	H	OMe	H	H	CF ₃	H	NMe	CH	N	1
123	Et	H	OMe	H	H	CF ₃	H	NMe	CH	N	2
124	Et	H	OH	H	H	CF ₃	H	NMe	CH	N	0
125	Et	H	H	H	H	NH ₂	H	NMe	CH	N	0

[0464]

Table 25

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
126	Et	H	H	H	H	CHFCE ₃	H	NMe	CH	N	1
127	Et	H	H	H	H	CHFCE ₃	H	NMe	CH	N	2
128	Et	H	H	H	H	C ₃ F ₇	H	NMe	CH	N	0
129	Et	H	H	H	H	C ₃ F ₇	H	NMe	CH	N	1
130	Et	H	H	H	H	C ₃ F ₇	H	NMe	CH	N	2
131	Et	Cl	H	H	H	C ₂ F ₅	H	NMe	CH	N	1
132	Et	Cl	H	H	H	C ₂ F ₅	H	NMe	CH	N	2
133	Et	H	Cl	H	H	CF ₃	H	NMe	CH	N	0
134	Et	H	Cl	H	H	CF ₃	H	NMe	CH	N	1
135	Et	H	Cl	H	H	CF ₃	H	NMe	CH	N	2
136	Et	H	H	Cl	H	CF ₃	H	NMe	CH	N	0
137	Et	H	H	Cl	H	CF ₃	H	NMe	CH	N	1
138	Et	H	H	Cl	H	CF ₃	H	NMe	CH	N	2
139	Et	H	H	H	Cl	CF ₃	H	NMe	CH	N	0
140	Et	H	H	H	Cl	CF ₃	H	NMe	CH	N	1
141	Et	H	H	H	Cl	CF ₃	H	NMe	CH	N	2
142	Et	H	H	OMe	H	CF ₃	H	NMe	CH	N	0
143	Et	H	H	OMe	H	CF ₃	H	NMe	CH	N	1
144	Et	H	H	OMe	H	CF ₃	H	NMe	CH	N	2
145	Et	H	H	H	H	SH	H	NMe	CH	N	0
146	Et	H	H	H	H	Et	H	NMe	CH	N	0
147	Et	H	H	H	H	iPr	H	NMe	CH	N	0
148	Et	H	H	H	H	NHEt	H	NMe	CH	N	0
149	Et	H	H	H	H	NEt ₂	H	NMe	CH	N	0
150	Et	H	H	H	H	tBu	H	NMe	CH	N	0

[0465]

Table 26

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
151	Et	H	H	H	H	H	CF ₃	NMe	CH	N	0
152	Et	H	H	H	OMe	CF ₃	H	NMe	CH	N	0
153	Et	H	H	H	OMe	CF ₃	H	NMe	CH	N	1
154	Et	H	H	H	OMe	CF ₃	H	NMe	CH	N	2
155	Et	H	H	H	OH	CF ₃	H	NMe	CH	N	0
156	Et	F	H	H	H	CF ₃	H	NMe	CH	N	0
157	Et	F	H	H	H	CF ₃	H	NMe	CH	N	1
158	Et	F	H	H	H	CF ₃	H	NMe	CH	N	2
159	Et	H	H	H	H	H	CF ₃	NMe	CH	N	1
160	Et	H	H	H	H	H	CF ₃	NMe	CH	N	2
161	Et	H	H	H	H	SCF ₃	H	NMe	CH	N	0
162	Et	H	H	H	H	NMe ₂	H	NMe	CH	N	0
163	Et	H	H	H	H	pyrrolidin-1-yl	H	NMe	CH	N	0
164	Et	H	H	H	H	NHCOMe	H	NMe	CH	N	0
165	Et	H	H	H	H	CH ₂ CF ₃	H	NMe	CH	N	0
166	Et	H	H	H	H	CF ₃	H	NMe	N	CH	0

167	Et	H	H	H	H	CF ₃	H	NMe	N	CH	1
168	Et	H	H	H	H	CF ₃	H	NMe	N	CH	2
169	Et	H	H	H	F	CF ₃	H	NMe	CH	N	0
170	Et	H	H	H	F	CF ₃	H	NMe	CH	N	1
171	Et	H	H	H	F	CF ₃	H	NMe	CH	N	2
172	Et	H	H	H	H	N(Me) COMe	H	NMe	CH	N	0
173	Et	H	H	H	H	NH2	H	NMe	CH	N	1
174	Et	H	CF ₃	H	H	CF ₃	H	NMe	CH	N	0
175	Et	H	CF ₃	H	H	CF ₃	H	NMe	CH	N	1

[0466]

Table 27

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
176	Et	H	CF ₃	H	H	CF ₃	H	NMe	CH	N	2
177	Et	H	H	H	H	NHCOCF ₃	H	NMe	CH	N	0
178	Et	H	H	H	H	NHCOCF ₃	H	NMe	CH	N	1
179	Et	H	H	H	H	NHCOCF ₃	H	NMe	CH	N	2
180	Et	H	H	H	H	2-CF ₃ -Ph	H	NMe	CH	N	0
181	Et	H	H	H	H	3-CF ₃ -Ph	H	NMe	CH	N	0
182	Et	H	H	H	H	4-CF ₃ -Ph	H	NMe	CH	N	0
183	Et	H	H	H	H	CF ₃	H	S	CH	N	0
184	Et	H	H	H	H	CF ₃	H	S	CH	N	1
185	Et	H	H	H	H	CF ₃	H	S	CH	N	2
186	CH ₂ CF ₃	H	H	H	H	CF ₃	H	NMe	CH	N	0
187	CH ₂ CF ₃	H	H	H	H	CF ₃	H	NMe	CH	N	1
188	CH ₂ CF ₃	H	H	H	H	CF ₃	H	NMe	CH	N	2
189	Et	Me	H	H	H	CF ₃	H	NMe	CH	N	0
190	Et	Me	H	H	H	CF ₃	H	NMe	CH	N	1
191	Et	Me	H	H	H	CF ₃	H	NMe	CH	N	2
192	Et	H	Me	H	H	CF ₃	H	NMe	CH	N	0
193	Et	H	Me	H	H	CF ₃	H	NMe	CH	N	1
194	Et	H	Me	H	H	CF ₃	H	NMe	CH	N	2
195	Et	H	H	H	H	2-CF ₃ -Ph	H	NMe	CH	N	1
196	Et	H	H	H	H	2-CF ₃ -Ph	H	NMe	CH	N	2
197	Et	H	H	H	H	3-CF ₃ -Ph	H	NMe	CH	N	1
198	Et	H	H	H	H	3-CF ₃ -Ph	H	NMe	CH	N	2
199	Et	H	H	H	H	4-CF ₃ -Ph	H	NMe	CH	N	1
200	Et	H	H	H	H	4-CF ₃ -Ph	H	NMe	CH	N	2

[0467]

Table 28

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
201	Et	H	H	CF ₃	H	CF ₃	H	NMe	CH	N	0
202	Et	H	H	CF ₃	H	CF ₃	H	NMe	CH	N	1
203	Et	H	H	CF ₃	H	CF ₃	H	NMe	CH	N	2
204	Et	H	H	H	H	2-Cl-Ph	H	NMe	CH	N	0
205	Et	H	H	H	H	3-Cl-Ph	H	NMe	CH	N	0
206	Et	H	H	H	H	4-Cl-Ph	H	NMe	CH	N	0

207	Et	H	H	H	H	6-Cl-3-Py	H	NMe	CH	N	0
208	Et	H	H	H	H	5-F-3-Py	H	NMe	CH	N	0
209	Et	H	H	H	H	SCF ₃	H	NMe	CH	N	2
210	Et	H	H	H	H	SCF ₃	H	NMe	CH	N	1
211	Et	H	H	H	H	3-Py	H	NMe	CH	N	0
212	Et	H	H	H	H	4-Py	H	NMe	CH	N	0
213	Et	H	H	H	H	4-Cl-Pyrazole	H	NMe	CH	N	0
214	Et	H	H	H	H	SOCF ₃	H	NMe	CH	N	2
215	Et	H	H	H	H	SO ₂ CF ₃	H	NMe	CH	N	2
216	Et	H	H	H	H	2-Cl-Ph	H	NMe	CH	N	1
217	Et	H	H	H	H	2-Cl-Ph	H	NMe	CH	N	2
218	Et	H	H	H	H	3-Cl-Ph	H	NMe	CH	N	1
219	Et	H	H	H	H	3-Cl-Ph	H	NMe	CH	N	2
220	Et	H	H	H	H	4-Cl-Ph	H	NMe	CH	N	2
221	Et	H	H	H	H	4-Py	H	NMe	CH	N	1
222	Et	H	H	H	H	4-Py	H	NMe	CH	N	2
223	Et	H	H	H	H	6-Cl-3-Py	H	NMe	CH	N	2
224	Et	H	H	H	H	5-F-3-Py	H	NMe	CH	N	1
225	Et	H	H	H	H	5-F-3-Py	H	NMe	CH	N	2

[0468]

Table 29

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
226	Et	H	H	H	H	4-Cl-Pyrazole	H	NMe	CH	N	2
227	Et	H	H	H	H	3-Cl-triazole	H	NMe	CH	N	0
228	Et	H	H	H	H	4-CF ₃ -imidazole	H	NMe	CH	N	0
229	Et	H	H	H	H	2-NO ₂ -Ph	H	NMe	CH	N	0
230	Et	H	H	H	H	3-NO ₂ -Ph	H	NMe	CH	N	0
231	Et	H	H	H	H	2-CN-Ph	H	NMe	CH	N	0
232	Et	H	H	H	H	3-CN-Ph	H	NMe	CH	N	0
233	Et	H	H	H	H	4-CN-Ph	H	NMe	CH	N	0
234	Et	H	H	H	H	3-CF ₃ -triazole	H	NMe	CH	N	0
235	Et	H	H	H	H	3-CF ₃ -5-Me-triazole	H	NMe	CH	N	0
236	Et	H	H	H	H	3-Cl-triazole	H	NMe	CH	N	2
237	Et	H	H	H	H	4-CF ₃ -imidazole	H	NMe	CH	N	1
238	Et	H	Br	H	H	CF ₃	H	NMe	CH	N	0
239	Et	H	Br	H	H	CF ₃	H	NMe	CH	N	1
240	Et	H	Br	H	H	CF ₃	H	NMe	CH	N	2
241	Et	H	CN	H	H	CF ₃	H	NMe	CH	N	0
242	Et	H	CN	H	H	CF ₃	H	NMe	CH	N	1
243	Et	H	CN	H	H	CF ₃	H	NMe	CH	N	2

246	Et	H	C ₂ F ₅	H	H	CF ₃	H	NMe	CH	N	0
247	Et	H	C ₂ F ₅	H	H	CF ₃	H	NMe	CH	N	1
248	Et	H	C ₂ F ₅	H	H	CF ₃	H	NMe	CH	N	2
249	Et	H	CHO	H	H	CF ₃	H	NMe	CH	N	0
250	Et	H	Ph	H	H	CF ₃	H	NMe	CH	N	0

[0469]

Table 30

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
251	Et	H	CF ₃	H	H	CF ₃	H	S	CH	N	2
253	Et	H	H	H	H	SMe	H	NMe	CH	N	0
254	Et	H	H	H	H	SO ₂ Me	H	NMe	CH	N	2
255	Et	H	H	H	H	SEt	H	NMe	CH	N	0
256	Et	H	H	H	H	SO ₂ Et	H	NMe	CH	N	2
257	Et	H	H	H	H	SiPr	H	NMe	CH	N	0
258	Et	H	H	H	H	SO ₂ iPr	H	NMe	CH	N	2
259	Et	H	H	H	H	SCH ₂ CF ₃	H	NMe	CH	N	0
260	Et	H	H	H	H	SO ₂ CH ₂ CF ₃	H	NMe	CH	N	2
261	Et	H	H	H	H	SCH ₂ CH=CH ₂	H	NMe	CH	N	0
262	Et	H	H	H	H	SCF ₂ CF ₃	H	NMe	CH	N	0
263	Et	H	H	H	H	SCF ₂ CF ₂ CF ₃	H	NMe	CH	N	0
264	Et	H	H	H	H	SCF(CF ₃) ₂	H	NMe	CH	N	0
265	Et	H	H	H	H	CH(OH)CF ₃	H	NMe	CH	N	0
266	Et	H	H	H	H	CH(Cl)CF ₃	H	NMe	CH	N	0
267	Et	H	CF ₃	H	H	CF ₃	H	S	CH	N	0
268	Et	H	H	H	H	OH	H	NMe	CH	N	0
269	Et	H	H	H	H	OH	H	NMe	CH	N	2
270	Et	H	H	H	H	OCF ₂ Br	H	NMe	CH	N	2
271	Et	H	H	H	H	OCF ₃	H	NMe	CH	N	2
272	Et	H	H	H	H	SCF ₂ CF ₃	H	NMe	CH	N	1
273	Et	H	H	H	H	SCF ₂ CF ₃	H	NMe	CH	N	2
274	Et	H	H	H	H	SCF ₂ CF ₂ CF ₃	H	NMe	CH	N	1
275	Et	H	H	H	H	SCF ₂ CF ₂ CF ₃	H	NMe	CH	N	2

[0470]

Table 31

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
276	Et	H	H	H	H	SC(CH ₃) ₃	H	NMe	CH	N	0
277	Et	H	H	H	H	SO ₂ C(CH ₃) ₃	H	NMe	CH	N	2
278	Et	H	CF ₃	H	H	Br	H	NMe	CH	N	0
279	Et	H	CF ₃	H	H	Br	H	NMe	CH	N	1
280	Et	H	CF ₃	H	H	Br	H	NMe	CH	N	2
281	Et	H	H	H	H	SCH=C=CH ₂	H	NMe	CH	N	0
282	Et	H	H	H	H	SO ₂ CH=C=CH ₂	H	NMe	CH	N	2
283	Et	H	H	H	H	SO ₂ CH ₂ CH=CH ₂	H	NMe	CH	N	2
284	Et	H	H	H	H	NHSO ₂ CH ₃	H	NMe	CH	N	2
285	Et	H	NO ₂	H	H	CF ₃	H	NMe	CH	N	0

286	Et	H	NO ₂	H	H	CF ₃	H	NMe	CH	N	1
287	Et	H	NO ₂	H	H	CF ₃	H	NMe	CH	N	2
289	Et	H	H	H	H	NHSO ₂ CF ₃	H	NMe	CH	N	2
290	Et	H	H	H	H	NCH ₃ SO ₂ CF ₃	H	NMe	CH	N	2
291	Et	H	H	H	H	NCH ₃ SO ₂ CH ₃	H	NMe	CH	N	2
292	Et	H	CF ₃	H	H	CF ₂ CF ₃	H	NMe	CH	N	0
293	Et	H	Cl	H	H	CF ₃	H	S	CH	N	0
294	Et	H	Cl	H	H	CF ₃	H	S	CH	N	2
295	Et	H	H	H	H	C(OH)(CF ₃) ₂	H	NMe	CH	N	0
296	Et	H	H	H	H	C(Cl)(CF ₃) ₂	H	NMe	CH	N	0
297	Et	H	H	H	H	C(Cl)(CF ₃) ₂	H	NMe	CH	N	1
298	Et	H	H	H	H	C(Cl)(CF ₃) ₂	H	NMe	CH	N	2
299	Et	H	CF ₃	H	H	CF ₂ CF ₃	H	NMe	CH	N	2
300	Et	H	CF ₃	H	H	I	H	NMe	CH	N	0

[0471]

Table 32

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
301	Et	H	CF ₃	H	H	SCF ₃	H	NMe	CH	N	0
302	Et	H	CF ₃	H	H	SO ₂ CF ₃	H	NMe	CH	N	2
303	Et	H	H	H	H	I	H	NMe	CH	CH	0
304	Et	H	H	H	H	H	CF(CF ₃) ₂	NMe	CH	CH	0
305	Et	H	H	H	H	CF(CF ₃) ₂	H	NMe	CH	CH	0
306	Et	H	CF ₃	H	H	SCF ₃	H	NMe	CH	N	2
307	Et	H	CF ₃	H	H	I	H	NMe	CH	N	2
308	Et	H	H	H	H	CF ₂ CF ₃	H	NMe	CH	CH	0
309	Et	H	H	H	H	CF ₂ CF ₃	H	NMe	CH	CH	1
310	Et	H	H	H	H	CF ₂ CF ₃	H	NMe	CH	CH	2
311	Et	H	H	H	H	SF ₅	H	NMe	CH	CH	0
312	Et	H	H	H	H	SF ₅	H	NMe	CH	CH	1
313	Et	H	H	H	H	SF ₅	H	NMe	CH	CH	2
314	Et	H	CF ₃	H	H	SF ₅	H	NMe	CH	CH	0
315	Et	H	CF ₃	H	H	SF ₅	H	NMe	CH	CH	1
316	Et	H	CF ₃	H	H	SF ₅	H	NMe	CH	CH	2
317	Et	H	Me	H	H	CF ₂ CF ₃	H	NMe	CH	N	0
318	Et	H	Me	H	H	CF ₂ CF ₃	H	NMe	CH	N	1
319	Et	H	Me	H	H	CF ₂ CF ₃	H	NMe	CH	N	2
320	Et	H	H	H	H	I	H	S	CH	N	0
321	Et	H	CF ₃	H	H	I	H	S	CH	N	0
322	Et	H	H	H	H	CF ₂ CF ₃	H	S	CH	N	0
323	Et	H	CF ₃	H	H	CF ₂ CF ₃	H	S	CH	N	0
324	Et	H	H	H	H	CF ₂ CF ₃	H	S	CH	N	2
325	Et	H	CF ₃	H	H	CF ₂ CF ₃	H	S	CH	N	2

[0472]

Table 33

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
326	Et	H	H	H	Cl	CF ₃	H	S	CH	N	0
327	Et	H	H	H	Cl	CF ₃	H	S	CH	N	2
328	Et	H	H	H	H	H	CF ₃	S	N	CH	0
329	Et	H	H	H	H	H	CF ₃	S	N	CH	2
330	Et	H	CH=CH ₂	H	H	CF ₃	H	NMe	CH	N	2
331	Et	H	Et	H	H	CF ₃	H	NMe	CH	N	2
332	Et	H	CF ₂ CF ₂ C F ₃	H	H	CF ₃	H	NMe	CH	N	2
333	Et	H	2- pyrimi dyl	H	H	CF ₃	H	NMe	CH	N	2
334	Et	H	H	H	H	SO ₂ N Me ₂	H	NMe	CH	N	1
335	Et	H	H	H	H	SO ₂ N Me ₂	H	NMe	CH	N	2
336	Et	H	H	H	H	CF ₃	H	NMe	CH	CNH ₂	0
337	Et	H	H	H	H	CF ₃	H	NMe	CH	CNHCO CH ₃	0
338	Et	H	H	H	H	CF ₃	H	NMe	CH	CNMe ₂	0
339	Et	H	CF ₃	H	H	CF ₃	H	NMe	CH	CNH ₂	0
340	Et	H	CF ₃	H	H	CF ₃	H	NMe	CH	CNMe ₂	0
341	Et	H	SF ₅	H	H	CF ₃	H	NMe	CH	N	0
342	Et	H	SF ₅	H	H	CF ₃	H	NMe	CH	N	1
343	Et	H	SF ₅	H	H	CF ₃	H	NMe	CH	N	2
344	Et	H	CF ₃	H	H	CF ₃	H	NH	CH	N	0
345	Et	H	CF ₃	H	H	CF ₃	H	NH	CH	N	2
346	Et	H	H	H	H	CF(C F ₃) ₂	H	NH	CH	CH	0
347	Et	H	H	H	H	Br	H	NMe	CCF ₂ H	N	0
348	Et	H	H	H	H	Br	H	NMe	CCF ₂ H	N	1
349	Et	H	H	H	H	Br	H	NMe	CCF ₂ H	N	2
350	Et	H	H	H	H	Br	H	NMe	CCH(C H ₃) ₂	N	0

[0473]

Table 34

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
351	Et	H	H	H	H	CF ₃	H	NH	CH	N	0
352	Et	H	H	H	H	CF ₃	H	NH	CH	N	1
353	Et	H	H	H	H	CF ₃	H	NH	CH	N	2
354	Et	H	H	H	H	CF ₃	H	NH	CH	CH	0
355	Et	H	CF ₃	H	H	CF ₃	H	NEt	CH	N	2
356	Et	H	CF ₃	H	H	CF ₃	H	NCH ₂ CH=CH ₂	CH	N	2
357	Et	H	CF ₃	H	H	CF ₃	H	NCH ₂ CCH	CH	N	2

358	Et	H	CF ₃	H	H	CF ₃	H	NCH ₂ CN	CH	N	2
359	Et	H	CF ₃	H	H	H	CF ₃	NCH ₂ CN	N	CH	2
360	Et	H	CF ₃	H	H	CF ₃	H	NCH ₂ OEt	CH	N	2
361	Et	H	CF ₃	H	H	H	CF ₃	NCH ₂ OEt	N	CH	2
362	Et	H	CF ₃	H	H	CF ₃	H	NCH ₂ SMe	CH	N	2
363	Et	H	CF ₃	H	H	CF ₃	H	NPr	CH	N	2
364	Et	H	CF ₃	H	H	CF ₃	H	N(CH ₂) ₃ CH ₃	CH	N	2
365	Et	H	CF ₃	H	H	CF ₃	H	NCH ₂ CO ₂ Me	CH	N	2
366	Et	H	CF ₃	H	H	H	CF ₃	NCH ₂ CO ₂ Me	N	CH	2
367	Et	H	CF ₃	H	H	CF ₃	H	NCH ₂ CH=CCl ₂	CH	N	2
368	Et	H	CF ₃	H	H	CF ₃	H	NCO ₂ tBu	CH	N	2
369	Et	H	CF ₃	H	H	CF ₃	H	NMe	CH	CH	0
370	Et	H	CF ₃	H	H	CF ₃	H	NMe	CH	CH	1
371	Et	H	CF ₃	H	H	CF ₃	H	NMe	CH	CH	2
372	Et	H	CF ₃	H	H	CF ₃	H	NCO ₂ Me	CH	N	2
373	Et	H	CF ₃	H	H	CF ₃	H	NCOMe	CH	N	2
374	Et	H	OCF ₃	H	H	CF ₃	H	NMe	CH	N	0
375	Et	H	OCF ₃	H	H	CF ₃	H	NMe	CH	N	1

[0474]

Table 35

Present Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	A ¹	A ²	A ³	n
376	Et	H	OCF ₃	H	H	CF ₃	H	NMe	CH	N	2
377	Et	H	CF ₂ CF ₂ C F ₂ CF ₃	H	H	CF ₃	H	NMe	CH	N	2
378	Et	H	2- pyridyl	H	H	CF ₃	H	NMe	CH	N	2
379	Et	H	NH ₂	H	H	CF ₃	H	NMe	CH	N	2
380	Et	H	NHCOCF ₃	H	H	CF ₃	H	NMe	CH	N	2
381	Et	H	iPr	H	H	CF ₃	H	NMe	CH	N	2
382	Et	H	CHO	H	H	CF ₃	H	NMe	CH	N	2
383	CH ₂ CH ₂ C H ₂ CH ₃	H	H	H	H	CF ₃	H	NMe	CH	N	0
384	CH ₂ CO ₂ C H ₃	H	H	H	H	CF ₃	H	NMe	CH	N	0
385	CH ₂ CH=C Cl ₂	H	H	H	H	CF ₃	H	NMe	CH	N	0
386	CH ₂ CCCH 3	H	H	H	H	CF ₃	H	NMe	CH	N	0
387	CH ₂ CN	H	H	H	H	CF ₃	H	NMe	CH	N	0
388	CH ₂ C(CH ₃) ₃	H	H	H	H	CF ₃	H	NMe	CH	N	0
389	CH ₂ CH ₂ C N	H	H	H	H	CF ₃	H	NMe	CH	N	0
390	CH ₂ CyBu	H	H	H	H	CF ₃	H	NMe	CH	N	0
391	CF ₂ Br	H	H	H	H	CF ₃	H	NMe	CH	N	0
392	Et	H	CF ₂ H	H	H	CF ₃	H	NMe	CH	N	2
393	Et	H	CH ₂ OH	H	H	CF ₃	H	NMe	CH	N	2

394	$(\text{CH}_2)_3\text{C}$ H_3	H	H	H	H	CF_3	H	NMe	CH	N	2
395	$\text{CH}_2\text{CO}_2\text{C}$ H_3	H	H	H	H	CF_3	H	NMe	CH	N	2
396	$\text{CH}_2\text{CH}=\text{C}$ Cl_2	H	H	H	H	CF_3	H	NMe	CH	N	2
397	CH_2CCCH $_3$	H	H	H	H	CF_3	H	NMe	CH	N	2
398	CH_2CN	H	H	H	H	CF_3	H	NMe	CH	N	2
399	$\text{CH}_2\text{C}(\text{CH})$ $_3)_3$	H	H	H	H	CF_3	H	NMe	CH	N	2
400	$\text{CH}_2\text{CH}_2\text{C}$ N	H	H	H	H	CF_3	H	NMe	CH	N	2

[0475]

Table 36

Present Compound	R^1	R^2	R^3	R^4	R^5	R^6	R^7	A^1	A^2	A^3	n
401	CH_2CyBu	H	H	H	H	CF_3	H	NMe	CH	N	2
402	CF_2Br	H	H	H	H	CF_3	H	NMe	CH	N	2
403	Et	H	CH_2F	H	H	CF_3	H	NMe	CH	N	2
404	$\text{CH}=\text{CH}_2$	H	H	H	H	CF_3	H	NMe	CH	N	0
405	$\text{CH}=\text{CH}_2$	H	H	H	H	CF_3	H	NMe	CH	N	1
406	$\text{CH}=\text{CH}_2$	H	H	H	H	CF_3	H	NMe	CH	N	2
407	Et	H	H	H	H	H	CF_3	S	CH	N	0
408	Et	H	H	H	H	H	CF_3	S	CH	N	2
409	Et	H	OCF_3	H	H	CF_2CF_3	H	NMe	CH	N	0
410	Et	H	OCF_3	H	H	CF_2CF_3	H	NMe	CH	N	1
411	Et	H	OCF_3	H	H	CF_2CF_3	H	NMe	CH	N	2
412	Et	H	CF_3	H	H	CF_3	H	NMe	CH	CCH_3	0
413	Et	H	CF_3	H	H	CF_3	H	NMe	CH	CCH_3	1
414	Et	H	CF_3	H	H	CF_3	H	NMe	CH	CCH_3	2
415	Et	H	CF_3	H	H	CF_3	H	NMe	CH	CF	0
416	Et	H	CF_3	H	H	CF_3	H	NMe	CH	CF	1
417	Et	H	CF_3	H	H	CF_3	H	NMe	CH	CF	2
418	Et	H	CF_3	H	H	CF_2CF_3	H	NMe	CH	CH	0
419	Et	H	CF_3	H	H	CF_2CF_3	H	NMe	CH	CH	1
420	Et	H	CF_3	H	H	CF_2CF_3	H	NMe	CH	CH	2
421	CH_2CyPr	H	H	H	H	CF_3	H	NMe	CH	N	0
422	CH_2CyPr	H	H	H	H	CF_3	H	NMe	CH	N	1
423	CH_2CyPr	H	H	H	H	CF_3	H	NMe	CH	N	2
424	Et	H	CF_3	H	H	CF_3	H	O	CH	CH	0
425	Et	H	CF_3	H	H	CF_3	H	O	CH	CH	2

[0476]

Table 37

Present Compound	R^1	R^2	R^3	R^4	R^5	R^6	R^7	A^1	A^2	A^3	n
426	Et	H	CF_3	H	H	CF_3	H	NMe	CH	CBr	0

427	Et	H	CF ₃	H	H	CF ₃	H	NMe	CH	CSCH ₂ CH ₃	0

[0477]

In the above Tables 20-37, Me represents a methyl group, Et represents an ethyl group, Pr represents a propyl group, iPr represents an isopropyl group, tBu represents a tert-butyl group, CyPr represents a cyclopropyl group, CyBu represents a cyclobutyl group, 2-F-Ph represents a 2-fluorophenyl group, 3-F-Ph represents a 3-fluorophenyl group, 4-F-Ph represents a 4-fluorophenyl group, 2-CF₃-Ph represents a 2-trifluoromethylphenyl group, 3-CF₃-Ph represents a 3-trifluoromethylphenyl group, 4-CF₃-Ph represents a 4-trifluoromethylphenyl group, 2-Cl-Ph represents a 2-chlorophenyl group, 3-Cl-Ph represents a 3-chlorophenyl group, 4-Cl-Ph represents a 4-chlorophenyl group, 2-NO₂-Ph represents a 2-nitrophenyl group, 3-NO₂-Ph represents a 3-nitrophenyl group, 2-CN-Ph represents a 2-cyanophenyl group, 3-CN-Ph represents a 3-cyanophenyl group, 4-CN-Ph represents a 4-cyanophenyl group, 3-Py represents a pyridin-3-yl group, 4-Py represents a pyridin-4-yl group, 6-Cl-3-Py represents a 6-chloropyridin-3-yl group, 5-F-3-Py represents a 5-fluoropyridin-3-yl group, 4-Cl-pyrazole represents a 4-chloropyrazol-1-yl group, 3-Cl-triazole represents a 3-chloro-(1H-1,2,4-triazol)-1-yl group, 3-CF₃-triazole represents a 3-trifluoromethyl-(1H-1,2,4-triazol)-

1-yl group, 3-CF₃-5-Me-triazole represents a 3-trifluoromethyl-5-methyl-(1H-1,2,4-triazol)-1-yl group, and 4-CF₃-imidazole represents a 4-trifluoromethylimidazol-1-yl group.

5 [0478]

The ¹H-NMR data of the present compounds listed in Tables 20-37 are shown below.

[0479]

Present Compound 92

10 ¹H-NMR (CDCl₃) δ: 8.73-8.71 (1H,m), 8.36-8.35 (1H,m), 7.49 (1H,t), 7.17 (2H,d), 3.70 (3H,s), 2.41 (6H,s).

Present Compound 94

¹H-NMR (CDCl₃) δ: 8.80-8.78 (1H,m), 8.60 (2H,d), 8.30-8.28 (1H,m), 8.08 (1H,t), 3.70 (3H,s), 3.09 (6H,s).

15 Present Compound 95

¹H-NMR (CDCl₃) δ: 8.73-8.71 (1H,m), 8.35-8.34 (1H,m), 7.47-7.43 (1H,m), 7.28-7.25 (2H,m), 3.69 (3H,s), 2.97-2.79 (4H,m), 1.24 (6H,t).

Present Compound 96

20 ¹H-NMR (CDCl₃) δ: 8.85-8.75 (1H,m), 8.38-7.96 (4H,m), 3.81-3.71 (3H,m), 3.15-2.65 (4H,m), 1.22-1.03 (6H,m).

Present Compound 97

¹H-NMR (CDCl₃) δ: 8.79 (1H,d), 8.54 (2H,d), 8.27 (1H,d), 8.06 (1H,t), 3.71 (3H,s), 3.32-3.07 (4H,m), 1.19 (6H,t).

25 Present Compound 98

^1H -NMR (CDCl_3) δ : 8.39 (1H,d), 8.07 (1H,d), 7.54-7.48 (2H,m), 7.45-7.43 (1H,m), 7.37-7.32 (1H,m), 3.72 (3H,s), 2.86 (2H,q), 1.23 (3H,t).

Present Compound 99

5 ^1H -NMR (CDCl_3) δ : 8.43 (1H,d), 8.26-8.22 (1H,m), 8.06 (1H,d), 7.84-7.79 (1H,m), 7.71-7.65 (1H,m), 7.60-7.57 (1H,m), 3.84 (3H,s), 3.39-3.29 (1H,m), 2.99-2.89 (1H,m), 1.29 (3H,t).

Present Compound 100

10 ^1H -NMR (CDCl_3) δ : 8.42 (1H,d), 8.23-8.20 (1H,m), 8.02 (1H,d), 7.86-7.77 (2H,m), 7.57-7.54 (1H,m), 3.67 (3H,s), 3.42 (2H,q), 1.25 (3H,t).

Present Compound 101

15 ^1H -NMR (CDCl_3) δ : 10.83 (1H,s), 8.63 (1H,s), 7.55-7.43 (3H,m), 7.38-7.32 (1H,m), 3.76 (3H,s), 2.89 (2H,q), 1.24 (3H,t).

Present Compound 102

20 ^1H -NMR (CDCl_3) δ : 8.73-8.71 (1H,m), 8.33-8.31 (1H,m), 7.47-7.42 (2H,m), 7.39-7.38 (1H,m), 3.77 (3H,s), 2.97 (2H,q), 2.82 (2H,q), 1.34 (3H,t), 1.21 (3H,t).

Present Compound 103

^1H -NMR (CDCl_3) δ : 8.12 (1H,s), 7.69 (1H,d), 7.62-7.57 (1H,m), 7.53-7.45 (3H,m), 7.36-7.31 (1H,m), 5.43 (2H,s), 3.26 (2H,q), 2.84 (2H,q), 1.22 (3H,t), 1.04 (3H,t).

25 Present Compound 104

^1H -NMR (CDCl_3) δ : 7.94-7.88 (2H,m), 7.61-7.56 (1H,m), 7.54-7.44 (3H,m), 7.37-7.30 (1H,m), 5.44 (2H,s), 3.27 (2H,q), 2.84 (2H,q), 1.22 (3H,t), 1.05 (3H,t).

Present Compound 107

5 ^1H -NMR (CDCl_3) δ : 8.72-8.69 (1H,m), 8.33-8.30 (1H,m), 7.55-7.45 (3H,m), 7.37-7.32 (1H,m), 4.40 (2H,t), 3.65 (2H,t), 3.14 (3H,s), 2.87 (2H,q), 1.24 (3H,t).

Present Compound 108

10 ^1H -NMR (CDCl_3) δ : 8.65 (1H,d), 8.26 (1H,d), 7.50-7.39 (3H,m), 7.33-7.26 (1H,m), 5.23 (2H,s), 2.81 (2H,q), 1.98 (3H,s), 1.17 (3H,t).

Present Compound 109

15 ^1H -NMR (CDCl_3) δ : 8.70 (1H,d), 8.32 (1H,d), 7.56-7.46 (3H,m), 7.39-7.33 (1H,m), 4.40 (2H,t), 2.92-2.80 (4H,m), 1.85 (3H,s), 1.24 (3H,t).

Present Compound 110

20 ^1H -NMR (CDCl_3) δ : 8.71 (1H,d), 8.31 (1H,d), 7.55-7.47 (2H,m), 7.45-7.41 (1H,m), 7.38-7.32 (1H,m), 4.21 (2H,t), 2.89 (2H,q), 1.75-1.63 (2H,m), 1.28-1.11 (5H,m), 0.78 (3H,t).

Present Compound 111

^1H -NMR (CDCl_3) δ : 8.84-8.81 (1H,m), 8.65-8.61 (1H,m), 7.48-7.37 (3H,m), 7.35-7.27 (1H,m), 2.74-2.64 (2H,m), 1.26 (9H,s), 1.10-1.04 (3H,m).

25 Present Compound 114

^1H -NMR (CDCl_3) δ : 8.59-8.52 (1H,m), 8.31-8.17 (1H,m), 7.61-7.32 (4H,m), 5.24 (1H,brs), 3.77 (3H,s), 2.97-2.82 (2H,m), 1.31-1.15 (3H,m).

Present Compound 115

5 ^1H -NMR (CDCl_3) δ : 8.51 (1H,s), 8.22 (1H,s), 7.55-7.50 (2H,m), 7.47-7.43 (1H,m), 7.38-7.32 (1H,m), 5.81 (1H,dq), 3.76 (3H,s), 2.88 (2H,q), 1.24 (3H,t).

Present Compound 119

10 ^1H -NMR (CDCl_3) δ : 8.71-8.70 (1H,m), 8.31-8.29 (1H,m), 7.49-7.45 (1H,m), 7.14-7.09 (2H,m), 4.00 (3H,s), 3.71 (3H,s), 2.74 (2H,q), 1.04 (3H,t).

[0480]

Present Compound 121

15 ^1H -NMR (CDCl_3) δ : 8.68-8.66 (1H,m), 8.28-8.27 (1H,m), 7.35 (1H,d), 6.99 (1H,d), 6.86-6.83 (1H,m), 3.88 (3H,s), 3.74 (3H,s), 2.84 (2H,q), 1.22 (3H,t).

Present Compound 122

20 ^1H -NMR (CDCl_3) δ : 8.73-8.72 (1H,m), 8.29-8.27 (1H,m), 7.77 (1H,d), 7.53 (1H,d), 7.19-7.15 (1H,m), 4.00 (3H,s), 3.89 (3H,s), 3.46-3.36 (1H,m), 3.00-2.90 (1H,m), 1.33 (3H,t).

Present Compound 123

^1H -NMR (CDCl_3) δ : 8.73-8.72 (1H,m), 8.27-8.25 (1H,m), 7.71 (1H,d), 7.46 (1H,d), 7.32-7.28 (1H,m), 3.99 (3H,s), 3.70 (3H,s), 3.43 (2H,q), 1.26 (3H,t).

25 Present Compound 124

¹H-NMR (CDCl₃) δ: 9.12 (1H,brs), 8.72-8.70 (1H,m), 8.33-8.30 (1H,m), 7.15 (1H,d), 6.88 (1H,d), 6.63-6.60 (1H,m), 3.77 (3H,s), 2.77 (2H,q), 1.19 (3H,t).

Present Compound 126

5 ¹H-NMR (CDCl₃) δ: 8.55 (1H,d), 8.26 (1H,d), 8.21 (1H,d), 7.83 (1H,t), 7.69 (1H,t), 7.60 (1H,d), 5.84 (1H,dq), 3.88 (3H,s), 3.42-3.30 (1H,m), 3.03-2.91 (1H,m), 1.33-1.25 (3H,m).

Present Compound 127

10 ¹H-NMR (CDCl₃) δ: 8.53 (1H,s), 8.26-8.16 (2H,m), 7.88-7.78 (2H,m), 7.59-7.53 (1H,m), 5.82 (1H,dq), 3.70 (3H,s), 3.44 (2H,q), 1.26 (3H,t).

Present Compound 131

15 ¹H-NMR (CDCl₃) δ: 8.68 (1H,d), 8.29 (1H,d), 7.74-7.68 (1H,m), 7.49-7.40 (2H,m), 3.73 (3H,s), 2.74 (2H,q), 1.06 (3H,t).

Present Compound 132

20 ¹H-NMR (CDCl₃) δ: 8.64 (1H,s), 8.24 (1H,s), 7.65 (1H,d), 7.58 (1H,t), 7.39 (1H,d), 3.72 (3H,s), 3.63-3.47 (1H,m), 3.37-3.22 (1H,m), 1.38-1.30 (3H,m).

Present Compound 136

¹H-NMR (CDCl₃) δ: 8.74-8.72 (1H,m), 8.33-8.32 (1H,m), 7.52-7.49 (1H,m), 7.47-7.43 (2H,m), 3.79 (3H,s), 2.85 (2H,q), 1.23 (3H,t).

25 Present Compound 137

¹H-NMR (CDCl₃) δ: 8.78-8.76 (1H,m), 8.33-8.31 (1H,m), 8.20 (1H,d), 7.81-7.78 (1H,m), 7.60 (1H,d), 3.93 (3H,s), 3.42-3.32 (1H,m), 3.02-2.92 (1H,m), 1.31 (3H,t).

Present Compound 138

5 ¹H-NMR (CDCl₃) δ: 8.76-8.75 (1H,m), 8.29-8.28 (1H,m), 8.16 (1H,d), 7.80-7.77 (1H,m), 7.57 (1H,d), 3.75 (3H,s), 3.40 (2H,q), 1.26 (3H,t).

Present Compound 139

10 ¹H-NMR (CDCl₃) δ: 8.75-8.73 (1H,m), 8.37-8.35 (1H,m), 7.48-7.44 (1H,m), 7.38-7.34 (2H,m), 3.72 (3H,s), 3.00-2.83 (2H,m), 1.26 (3H,t).

Present Compound 140

15 ¹H-NMR (CDCl₃) δ: 8.80-8.78 (1H,m), 8.35-8.34 (1H,m), 8.11-8.08 (1H,m), 7.79 (1H,t), 7.73-7.71 (1H,m), 3.77 (3H,s), 3.37-3.27 (1H,m), 3.09-2.99 (1H,m), 1.25 (3H,t).

Present Compound 141

¹H-NMR (CDCl₃) δ: 8.77-8.76 (1H,m), 8.32-8.30 (1H,m), 8.16-8.13 (1H,m), 7.89-7.87 (1H,m), 7.77 (1H,t), 3.73 (3H,s), 3.61-3.49 (1H,m), 3.34-3.24 (1H,m), 1.26 (3H,t).

20 Present Compound 142

¹H-NMR (CDCl₃) δ: 8.72-8.71 (1H,m), 8.32-8.30 (1H,m), 7.54 (1H,d), 7.10-7.06 (1H,m), 7.03 (1H,d), 3.85 (3H,s), 3.77 (3H,s), 2.68 (2H,q), 1.12 (3H,t).

Present Compound 143

25 ¹H-NMR (CDCl₃) δ: 8.76-8.74 (1H,m), 8.31-8.30 (1H,m), 8.14

(1H,d), 7.34-7.30 (1H,m), 7.08 (1H,d), 3.93 (3H,s), 3.89 (3H,s), 3.29-3.19 (1H,m), 2.97-2.86 (1H,m), 1.26 (3H,t).

Present Compound 144

¹H-NMR (CDCl₃) δ: 8.74-8.73 (1H,m), 8.28-8.27 (1H,m), 8.13 (1H,d), 7.25-7.21 (1H,m), 7.03 (1H,d), 3.94 (3H,s), 3.74 (3H,s), 3.34 (2H,q), 1.24 (3H,t).

Present Compound 146

¹H-NMR (CDCl₃) δ: 8.30 (1H,s), 7.93 (1H,s), 7.49-7.42 (3H,m), 7.33-7.27 (1H,m), 3.72 (3H,s), 2.89-2.78 (4H,m), 1.33 (3H,t), 1.22 (3H,t).

Present Compound 148

¹H-NMR (CDCl₃) δ: 7.94 (1H,d), 7.48-7.39 (3H,m), 7.34-7.25 (2H,m), 3.67 (3H,s), 3.26-3.16 (2H,m), 2.89-2.80 (2H,m), 1.34-1.27 (3H,m), 1.26-1.19 (3H,m).

Present Compound 149

¹H-NMR (CDCl₃) δ: 8.09 (1H,d), 7.50-7.41 (4H,m), 7.34-7.27 (1H,m), 3.68 (3H,s), 3.38 (4H,q), 2.86 (2H,q), 1.23 (3H,t), 1.17 (6H,t).

[0481]

Present Compound 152

¹H-NMR (CDCl₃) δ: 8.71-8.69 (1H,m), 8.33-8.31 (1H,m), 7.47 (1H,t), 7.06 (1H,d), 6.86 (1H,d), 3.76 (3H,s), 3.69 (3H,s), 2.98-2.82 (2H,m), 1.25 (3H,t).

Present Compound 153

¹H-NMR (CDCl₃) δ: 8.75-8.73 (1H,m), 8.30-8.29 (1H,m), 7.79-

7.76 (2H,m), 7.22-7.18 (1H,m), 3.86 (3H,s), 3.73 (3H,s),
3.42-3.30 (1H,m), 3.09-2.97 (1H,m), 1.29 (3H,t).

Present Compound 154

¹H-NMR (CDCl₃) δ: 8.73-8.71 (1H,m), 8.27-8.25 (1H,m), 7.82-
5 7.73 (2H,m), 7.37-7.34 (1H,m), 3.83 (3H,s), 3.68 (3H,s),
3.67-3.56 (1H,m), 3.36-3.26 (1H,m), 1.26 (3H,t).

Present Compound 155

¹H-NMR (CDCl₃) δ: 8.73-8.72 (1H,m), 8.29-8.27 (1H,m), 7.93
(1H,brs), 7.36 (1H,t), 7.09-7.06 (1H,m), 6.95-6.92 (1H,m),
10 3.85 (3H,s), 2.87-2.78 (2H,m), 1.21 (3H,t).

Present Compound 156

¹H-NMR (CDCl₃) δ: 8.74-8.72 (1H,m), 8.33-8.31 (1H,m), 7.51-
7.45 (1H,m), 7.36-7.31 (2H,m), 3.75 (3H,s), 2.78 (2H,q),
1.10 (3H,t).

15 Present Compound 157

¹H-NMR (CDCl₃) δ: 8.73-8.71 (1H,m), 8.29-8.27 (1H,m), 7.70-
7.64 (1H,m), 7.44-7.38 (1H,m), 7.33 (1H,d), 3.81 (3H,s),
3.53-3.43 (1H,m), 3.42-3.31 (1H,m), 1.33 (3H,t).

Present Compound 158

20 ¹H-NMR (CDCl₃) δ: 8.73-8.72 (1H,m), 8.28-8.26 (1H,m), 7.84-
7.78 (1H,m), 7.55-7.49 (1H,m), 7.38-7.35 (1H,m), 3.77
(3H,s), 3.50-3.34 (2H,m), 1.34 (3H,t).

Present Compound 162

¹H-NMR (CDCl₃) δ: 8.14 (1H,d), 7.51-7.40 (4H,m), 7.35-7.27
25 (1H,m), 3.69 (3H,s), 3.00 (6H,s), 2.90-2.81 (2H,m), 1.26-

1.19 (3H,m).

Present Compound 163

¹H-NMR (CDCl₃) δ: 7.92 (1H,d), 7.48-7.43 (3H,m), 7.34-7.28
(1H,m), 7.26 (1H,d), 3.68 (3H,s), 3.40-3.32 (4H,m), 2.84
5 (2H,q), 2.08-2.05 (4H,m), 1.22 (3H,t).

Present Compound 164

¹H-NMR (CDCl₃) δ: 9.72 (1H,s), 8.41 (1H,d), 8.23 (1H,d),
7.37-7.28 (3H,m), 7.18-7.13 (1H,m), 3.58 (3H,s), 2.74
(2H,q), 2.02 (3H,s), 1.09 (3H,t).

10 Present Compound 169

¹H-NMR (CDCl₃) δ: 8.74-8.72 (1H,m), 8.37-8.35 (1H,m), 7.54-
7.48 (1H,m), 7.27-7.24 (1H,m), 7.09-7.03 (1H,m), 3.77
(3H,d), 2.93 (2H,q), 1.28 (3H,t).

Present Compound 170

15 ¹H-NMR (CDCl₃) δ: 8.79-8.76 (1H,m), 8.35-8.32 (1H,m), 8.03
(1H,d), 7.87-7.80 (1H,m), 7.43 (1H,t), 3.84 (3H,d), 3.51-
3.40 (1H,m), 3.13-3.02 (1H,m), 1.33 (3H,t).

Present Compound 171

20 ¹H-NMR (CDCl₃) δ: 8.77-8.75 (1H,m), 8.32-8.30 (1H,m), 8.07-
8.04 (1H,m), 7.86-7.80 (1H,m), 7.62-7.57 (1H,m), 3.76
(3H,s), 3.69-3.58 (1H,m), 3.41-3.31 (1H,m), 1.29 (3H,t).

Present Compound 172

25 ¹H-NMR (CDCl₃) δ: 8.32 (1H,d), 7.94 (1H,d), 7.57-7.48
(2H,m), 7.47-7.43 (1H,m), 7.39-7.33 (1H,m), 3.77 (3H,s),
3.36 (3H,s), 2.91 (2H,q), 1.93 (3H,s), 1.27 (3H,t).

Present Compound 173

$^1\text{H-NMR}$ (CDCl_3) δ : 8.21 (1H,dd), 8.03 (1H,d), 7.77 (1H,td), 7.65 (1H,td), 7.57 (1H,dd), 7.37 (1H,d), 3.78 (3H,s), 3.36-3.24 (1H,m), 2.96-2.83 (1H,m), 1.26 (3H,t).

5 Present Compound 177

$^1\text{H-NMR}$ (CDCl_3) δ : 8.55 (1H,d), 8.39 (1H,d), 8.21 (1H,brs), 7.55-7.42 (3H,m), 7.38-7.31 (1H,m), 3.74 (3H,s), 2.86 (2H,q), 1.23 (3H,t).

Present Compound 178

10 $^1\text{H-NMR}$ (DMSO-D_6) δ : 11.57 (1H,brs), 8.68-8.64 (1H,m), 8.45-8.42 (1H,m), 8.10 (1H,d), 7.90 (2H,dd), 7.78 (1H,t), 3.83 (3H,s), 3.41-3.30 (1H,m), 2.94-2.81 (1H,m), 1.16 (3H,t).

Present Compound 179

15 $^1\text{H-NMR}$ (CDCl_3) δ : 9.18 (1H,brs), 8.52-8.48 (1H,m), 8.38-8.34 (1H,m), 8.25-8.17 (1H,m), 7.87-7.76 (2H,m), 7.59-7.52 (1H,m), 3.65 (3H,s), 3.47 (2H,q), 1.26 (3H,t).

Present Compound 180

20 $^1\text{H-NMR}$ (CDCl_3) δ : 8.40 (1H,d), 8.06 (1H,d), 7.82 (1H,d), 7.66-7.59 (1H,m), 7.58-7.46 (4H,m), 7.44-7.39 (1H,m), 7.38-7.31 (1H,m), 3.79 (3H,s), 2.90 (2H,q), 1.26 (3H,t).

[0482]

Present Compound 181

25 $^1\text{H-NMR}$ (CDCl_3) δ : 8.65 (1H,d), 8.26 (1H,d), 7.89 (1H,s), 7.83 (1H,d), 7.69-7.60 (2H,m), 7.53-7.46 (3H,m), 7.38-7.33 (1H,m), 3.78 (3H,s), 2.88 (2H,q), 1.25 (3H,t).

Present Compound 182

$^1\text{H-NMR}$ (CDCl_3) δ : 8.67 (1H,d), 8.27 (1H,d), 7.78-7.75 (4H,m), 7.56-7.44 (3H,m), 7.39-7.32 (1H,m), 3.78 (3H,s), 2.89 (2H,q), 1.25 (3H,t).

5 Present Compound 189

$^1\text{H-NMR}$ (CDCl_3) δ : 8.72-8.70 (1H,m), 8.30-8.28 (1H,m), 7.50-7.47 (1H,m), 7.40 (1H,t), 7.35-7.32 (1H,m), 3.71 (3H,s), 2.64 (3H,s), 2.53 (2H,q), 1.00 (3H,t).

Present Compound 190

10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.71-8.70 (1H,m), 8.26-8.24 (1H,m), 7.53 (1H,t), 7.47-7.44 (1H,m), 7.31-7.28 (1H,m), 3.73 (3H,s), 3.54-3.43 (1H,m), 3.27-3.17 (1H,m), 2.66 (3H,s), 1.29 (3H,t).

Present Compound 191

15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.72-8.70 (1H,m), 8.24-8.22 (1H,m), 7.65 (1H,t), 7.60-7.56 (1H,m), 7.34-7.31 (1H,m), 3.73 (3H,s), 3.52-3.32 (2H,m), 2.85 (3H,s), 1.32 (3H,t).

Present Compound 192

20 $^1\text{H-NMR}$ (CDCl_3) δ : 8.71-8.69 (1H,m), 8.31-8.30 (1H,m), 7.35-7.31 (2H,m), 7.19-7.15 (1H,m), 3.76 (3H,s), 2.85 (2H,q), 2.46 (3H,s), 1.22 (3H,t).

Present Compound 193

25 $^1\text{H-NMR}$ (CDCl_3) δ : 8.74-8.73 (1H,m), 8.30-8.28 (1H,m), 8.06-8.05 (1H,m), 7.49-7.48 (2H,m), 3.88 (3H,s), 3.42-3.32 (1H,m), 3.00-2.90 (1H,m), 2.58 (3H,s), 1.32 (3H,t).

Present Compound 194

^1H -NMR (CDCl_3) δ : 8.74-8.72 (1H,m), 8.27-8.26 (1H,m), 8.03-8.02 (1H,m), 7.64-7.61 (1H,m), 7.44 (1H,d), 3.70 (3H,s), 3.41 (2H,q), 2.59 (3H,s), 1.26 (3H,t).

5 Present Compound 195

^1H -NMR (CDCl_3) δ : 8.45 (1H,d), 8.26 (1H,d), 8.05 (1H,d), 7.85-7.78 (2H,m), 7.73-7.61 (3H,m), 7.56 (1H,t), 7.44 (1H,d), 3.91 (3H,s), 3.46-3.32 (1H,m), 3.03-2.91 (1H,m), 1.29 (3H,t).

10 Present Compound 196

^1H -NMR (CDCl_3) δ : 8.43 (1H,d), 8.28-8.20 (1H,m), 8.06-8.00 (1H,m), 7.90-7.75 (3H,m), 7.67-7.59 (2H,m), 7.54 (1H,t), 7.44 (1H,d), 3.74 (3H,s), 3.49 (2H,q), 1.25 (3H,t).

Present Compound 197

15 ^1H -NMR (CDCl_3) δ : 8.70 (1H,d), 8.29-8.23 (2H,m), 7.92-7.78 (3H,m), 7.74-7.60 (4H,m), 3.90 (3H,s), 3.47-3.33 (1H,m), 3.06-2.93 (1H,m), 1.32 (3H,t).

Present Compound 198

20 ^1H -NMR (CDCl_3) δ : 8.69 (1H,d), 8.26-8.19 (2H,m), 7.92-7.89 (1H,m), 7.87-7.76 (3H,m), 7.69-7.57 (3H,m), 3.73 (3H,s), 3.50 (2H,q), 1.27 (3H,t).

Present Compound 199

25 ^1H -NMR (CDCl_3) δ : 8.71 (1H,d), 8.29-8.23 (2H,m), 7.85-7.79 (1H,m), 7.78-7.75 (4H,m), 7.70 (1H,td), 7.63 (1H,dd), 3.90 (3H,s), 3.44-3.32 (1H,m), 3.05-2.92 (1H,m), 1.31 (3H,t).

Present Compound 200

$^1\text{H-NMR}$ (CDCl_3) δ : 8.70 (1H,d), 8.27-8.19 (2H,m), 7.87-7.78 (2H,m), 7.77-7.75 (4H,m), 7.62-7.57 (1H,m), 3.73 (3H,s), 3.48 (2H,d), 1.26 (3H,t).

5 Present Compound 201

$^1\text{H-NMR}$ (CDCl_3) δ : 8.75-8.74 (1H,m), 8.35-8.34 (1H,m), 7.78-7.74 (1H,m), 7.71-7.69 (1H,m), 7.54 (1H,d), 3.80 (3H,s), 2.98 (2H,q), 1.31 (3H,t).

Present Compound 202

10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.79-8.78 (1H,m), 8.44-8.41 (1H,m), 8.35-8.33 (1H,m), 8.10-8.06 (1H,m), 7.89-7.87 (1H,m), 3.94 (3H,s), 3.50-3.40 (1H,m), 3.07-2.97 (1H,m), 1.34 (3H,t).

Present Compound 203

15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.78-8.77 (1H,m), 8.40-8.37 (1H,m), 8.31-8.30 (1H,m), 8.10-8.06 (1H,m), 7.86-7.84 (1H,m), 3.75 (3H,s), 3.45 (2H,q), 1.28 (3H,t).

Present Compound 204

$^1\text{H-NMR}$ (CDCl_3) δ : 8.53 (1H,d), 8.19 (1H,d), 7.56-7.26 (8H,m), 3.79 (3H,s), 2.88 (2H,q), 1.24 (3H,t).

20 Present Compound 205

$^1\text{H-NMR}$ (CDCl_3) δ : 8.63 (1H,d), 8.23 (1H,d), 7.65-7.29 (8H,m), 3.77 (3H,s), 2.87 (2H,q), 1.23 (3H,t).

Present Compound 206

25 $^1\text{H-NMR}$ (CDCl_3) δ : 8.62 (1H,d), 8.22 (1H,d), 7.59-7.53 (2H,m), 7.51-7.40 (5H,m), 7.36-7.29 (1H,m), 3.77 (3H,s),

2.87 (2H,q), 1.23 (3H,t).

Present Compound 207

¹H-NMR (CDCl₃) δ: 8.68 (1H,d), 8.62 (1H,d), 8.24 (1H,d),
7.93 (1H,dd), 7.56-7.42 (4H,m), 7.40-7.34 (1H,m), 3.79
5 (3H,s), 2.89 (2H,q), 1.25 (3H,t).

Present Compound 208

¹H-NMR (CDCl₃) δ: 8.78-8.74 (1H,m), 8.67-8.64 (1H,m), 8.52
(1H,d), 8.27 (1H,d), 7.73-7.65 (1H,m), 7.57-7.46 (3H,m),
7.41-7.34 (1H,m), 3.79 (3H,s), 2.89 (2H,q), 1.25 (3H,t).

10 [0483]

Present Compound 211

¹H-NMR (CDCl₃) δ: 8.93 (1H,d), 8.69-8.62 (2H,m), 8.27
(1H,d), 8.00-7.92 (1H,m), 7.55-7.31 (5H,m), 3.79 (3H,s),
2.89 (2H,q), 1.25 (3H,t).

15 Present Compound 212

¹H-NMR (CDCl₃) δ: 8.75-8.69 (3H,m), 8.33 (1H,d), 7.63-7.57
(2H,m), 7.56-7.44 (3H,m), 7.40-7.32 (1H,m), 3.79 (3H,s),
2.89 (2H,q), 1.25 (3H,t).

Present Compound 213

20 ¹H-NMR (CDCl₃) δ: 8.79 (1H,d), 8.28 (1H,d), 7.97 (1H,s),
7.70 (1H,s), 7.55-7.43 (3H,m), 7.40-7.30 (1H,m), 3.77
(3H,s), 2.87 (2H,q), 1.23 (3H,t).

Present Compound 216

25 ¹H-NMR (CDCl₃) δ: 8.56 (1H,d), 8.26 (1H,dd), 8.16 (1H,d),
7.84-7.78 (1H,m), 7.69 (1H,td), 7.62 (1H,dd), 7.56-7.52

(1H,m), 7.46-7.43 (1H,m), 7.42-7.33 (2H,m), 3.90 (3H,s),
3.44-3.31 (1H,m), 3.03-2.91 (1H,m), 1.30 (3H,t).

Present Compound 217

¹H-NMR (CDCl₃) δ: 8.55 (1H,d), 8.24-8.20 (1H,m), 8.14
5 (1H,d), 7.85-7.76 (2H,m), 7.59 (1H,dd), 7.53 (1H,dd), 7.44
(1H,dd), 7.40-7.29 (2H,m), 3.73 (3H,s), 3.49 (2H,q), 1.25
(3H,t).

Present Compound 218

¹H-NMR (CDCl₃) δ: 8.68-8.64 (1H,m), 8.28-8.23 (1H,m), 8.22-
10 8.19 (1H,m), 7.84-7.78 (1H,m), 7.71-7.66 (1H,m), 7.65-7.59
(2H,m), 7.56-7.49 (1H,m), 7.46-7.41 (1H,m), 7.40-7.34
(1H,m), 3.89 (3H,s), 3.45-3.33 (1H,m), 3.04-2.92 (1H,m),
1.31 (3H,t).

Present Compound 219

¹H-NMR (CDCl₃) δ: 8.65 (1H,d), 8.22 (1H,dd), 8.18 (1H,d),
15 7.86-7.76 (2H,m), 7.64-7.62 (1H,m), 7.59 (1H,dd), 7.52
(1H,dt), 7.42 (1H,t), 7.36 (1H,dt), 3.72 (3H,s), 3.48
(2H,q), 1.26 (3H,t).

Present Compound 220

¹H-NMR (CDCl₃) δ: 8.66-8.62 (1H,m), 8.24-8.20 (1H,m), 8.19-
20 8.17 (1H,m), 7.86-7.75 (2H,m), 7.62-7.54 (3H,m), 7.48-7.42
(2H,m), 3.71 (3H,s), 3.56-3.41 (2H,m), 1.25 (3H,t).

Present Compound 221

¹H-NMR (CDCl₃) δ: 8.76 (1H,d), 8.74 (2H,dd), 8.30 (1H,d),
25 8.26 (1H,dd), 7.83 (1H,td), 7.71 (1H,td), 7.65-7.59 (3H,m),

3.91 (3H,s), 3.44-3.32 (1H,m), 3.05-2.93 (1H,m), 1.31 (3H,t).

Present Compound 222

¹H-NMR (CDCl₃) δ: 9.07-8.56 (3H,m), 8.28 (1H,d), 8.25-8.20 (1H,m), 7.89-7.78 (2H,m), 7.66-7.56 (3H,m), 3.73 (3H,s), 3.47 (2H,q), 1.26 (3H,t).

Present Compound 223

¹H-NMR (CDCl₃) δ: 8.68 (1H,d), 8.64 (1H,d), 8.26-8.21 (1H,m), 8.18 (1H,d), 7.92 (1H,dd), 7.89-7.78 (2H,m), 7.60 (1H,dd), 7.47 (1H,d), 3.73 (3H,s), 3.52-3.41 (2H,m), 1.26 (3H,t).

Present Compound 224

¹H-NMR (CDCl₃) δ: 8.77-8.74 (1H,m), 8.69 (1H,d), 8.53 (1H,d), 8.29-8.23 (2H,m), 7.87-7.80 (1H,m), 7.74-7.67 (2H,m), 7.66-7.61 (1H,m), 3.91 (3H,s), 3.45-3.31 (1H,m), 3.05-2.93 (1H,m), 1.31 (3H,t).

Present Compound 225

¹H-NMR (CDCl₃) δ: 8.77-8.74 (1H,m), 8.67 (1H,d), 8.53 (1H,d), 8.26-8.23 (1H,m), 8.22-8.20 (1H,m), 7.89-7.79 (2H,m), 7.71-7.66 (1H,m), 7.64-7.59 (1H,m), 3.74 (3H,s), 3.48 (2H,q), 1.27 (3H,t).

Present Compound 226

¹H-NMR (DMSO-D₆) δ: 8.96 (1H,d), 8.89 (1H,s), 8.54 (1H,d), 8.14 (1H,dd), 8.02-7.87 (4H,m), 3.60 (3H,s), 3.55 (2H,q), 1.12 (3H,t).

Present Compound 227

$^1\text{H-NMR}$ (CDCl_3) δ : 8.74 (1H,d), 8.48 (1H,s), 8.28 (1H,d), 7.57-7.50 (2H,m), 7.49-7.45 (1H,m), 7.40-7.34 (1H,m), 3.79 (3H,s), 2.88 (2H,q), 1.24 (3H,t).

5 Present Compound 228

$^1\text{H-NMR}$ (CDCl_3) δ : 8.51 (1H,d), 8.10 (1H,d), 7.89 (1H,s), 7.66-7.63 (1H,m), 7.58-7.43 (3H,m), 7.41-7.33 (1H,m), 3.80 (3H,s), 2.90 (2H,q), 1.25 (3H,t).

Present Compound 229

10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.38 (1H,d), 8.05 (1H,d), 7.98 (1H,dd), 7.70 (1H,td), 7.60-7.43 (5H,m), 7.39-7.31 (1H,m), 3.78 (3H,s), 2.89 (2H,q), 1.26 (3H,t).

Present Compound 230

15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.69 (1H,d), 8.54-8.51 (1H,m), 8.31-8.29 (1H,m), 8.27 (1H,dd), 7.98 (1H,d), 7.69 (1H,t), 7.57-7.46 (3H,m), 7.40-7.34 (1H,m), 3.79 (3H,s), 2.89 (2H,q), 1.26 (3H,t).

Present Compound 231

20 $^1\text{H-NMR}$ (CDCl_3) δ : 8.62 (1H,d), 8.25 (1H,d), 7.83 (1H,dd), 7.71 (1H,td), 7.59 (1H,dd), 7.55-7.46 (4H,m), 7.39-7.32 (1H,m), 3.79 (3H,s), 2.89 (2H,q), 1.25 (3H,t).

Present Compound 232

25 $^1\text{H-NMR}$ (CDCl_3) δ : 8.63 (1H,d), 8.24 (1H,d), 7.93 (1H,s), 7.89 (1H,d), 7.74-7.58 (2H,m), 7.56-7.45 (3H,m), 7.40-7.33 (1H,m), 3.79 (3H,s), 2.89 (2H,q), 1.25 (3H,t).

Present Compound 233

$^1\text{H-NMR}$ (CDCl_3) δ : 8.66 (1H,d), 8.27 (1H,d), 7.83-7.73 (4H,m), 7.56-7.45 (3H,m), 7.40-7.32 (1H,m), 3.79 (3H,s), 2.89 (2H,q), 1.25 (3H,t).

5 Present Compound 234

$^1\text{H-NMR}$ (CDCl_3) δ : 8.78 (1H,d), 8.68 (1H,s), 8.34 (1H,d), 7.54-7.26 (4H,m), 3.80 (3H,s), 2.93-2.82 (2H,m), 1.28-1.19 (3H,m).

Present Compound 235

10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.55 (1H,d), 8.17 (1H,d), 7.57-7.24 (4H,m), 3.81 (3H,s), 2.93-2.86 (2H,m), 2.60 (3H,s), 1.28-1.23 (3H,m).

Present Compound 236

15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.75 (1H,d), 8.49 (1H,s), 8.27 (1H,d), 8.25-8.22 (1H,m), 7.89-7.78 (2H,m), 7.63-7.57 (1H,m), 3.73 (3H,s), 3.43 (2H,q), 1.26 (3H,t).

Present Compound 237

20 $^1\text{H-NMR}$ (CDCl_3) δ : 8.57 (1H,d), 8.27 (1H,d), 8.11 (1H,d), 7.92 (1H,s), 7.88-7.82 (1H,m), 7.75-7.68 (2H,m), 7.65 (1H,d), 3.93 (3H,s), 3.46-3.32 (1H,m), 3.06-2.94 (1H,m), 1.32 (3H,t).

Present Compound 238

25 $^1\text{H-NMR}$ (CDCl_3) δ : 8.73-8.71 (1H,m), 8.33-8.32 (1H,m), 7.59 (1H,d), 7.50-7.47 (1H,m), 7.30 (1H,d), 3.77 (3H,s), 2.91 (2H,q), 1.27 (3H,t).

Present Compound 239

$^1\text{H-NMR}$ (CDCl_3) δ : 8.77-8.75 (1H,m), 8.39 (1H,d), 8.32-8.31 (1H,m), 7.84-7.81 (1H,m), 7.49 (1H,d), 3.91 (3H,s), 3.50-3.40 (1H,m), 3.06-2.96 (1H,m), 1.35 (3H,t).

5 Present Compound 240

$^1\text{H-NMR}$ (CDCl_3) δ : 8.77-8.75 (1H,m), 8.37 (1H,d), 8.31-8.29 (1H,m), 7.99-7.96 (1H,m), 7.44 (1H,d), 3.72 (3H,s), 3.44 (2H,q), 1.28 (3H,t).

[0484]

10 Present Compound 241

$^1\text{H-NMR}$ (CDCl_3) δ : 8.76-8.74 (1H,m), 8.35-8.34 (1H,m), 7.70-7.69 (1H,m), 7.63-7.60 (1H,m), 7.57-7.54 (1H,m), 3.79 (3H,s), 2.95 (2H,q), 1.30 (3H,t).

Present Compound 242

15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.80-8.78 (1H,m), 8.59-8.57 (1H,m), 8.36-8.34 (1H,m), 7.99-7.96 (1H,m), 7.78 (1H,d), 3.96 (3H,s), 3.57-3.46 (1H,m), 3.10-3.00 (1H,m), 1.37 (3H,t).

Present Compound 243

20 $^1\text{H-NMR}$ (CDCl_3) δ : 8.79-8.77 (1H,m), 8.53-8.51 (1H,m), 8.32-8.30 (1H,m), 8.14-8.10 (1H,m), 7.74-7.71 (1H,m), 3.74 (3H,s), 3.49 (2H,q), 1.29 (3H,t).

Present Compound 246

25 $^1\text{H-NMR}$ (CDCl_3) δ : 8.75-8.74 (1H,m), 8.35-8.34 (1H,m), 7.66-7.65 (1H,m), 7.61-7.55 (2H,m), 3.80 (3H,s), 2.95 (2H,q), 1.28 (3H,t).

Present Compound 247

$^1\text{H-NMR}$ (CDCl_3) δ : 8.80-8.78 (1H,m), 8.54-8.53 (1H,m), 8.35-8.34 (1H,m), 7.95-7.91 (1H,m), 7.79 (1H,d), 3.95 (3H,s), 3.52-3.41 (1H,m), 3.09-2.99 (1H,m), 1.33 (3H,t).

5 Present Compound 248

$^1\text{H-NMR}$ (CDCl_3) δ : 8.78-8.76 (1H,m), 8.47-8.46 (1H,m), 8.31-8.30 (1H,m), 8.10-8.07 (1H,m), 7.76 (1H,d), 3.76 (3H,s), 3.48 (2H,q), 1.28 (3H,t).

Present Compound 249

10 $^1\text{H-NMR}$ (CDCl_3) δ : 10.12 (1H,s), 8.75-8.74 (1H,m), 8.36-8.34 (1H,m), 7.97-7.96 (1H,m), 7.83-7.81 (1H,m), 7.64 (1H,d), 3.80 (3H,s), 3.00 (2H,q), 1.30 (3H,t).

Present Compound 250

15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.74-8.71 (1H,m), 8.35-8.33 (1H,m), 7.72-7.64 (3H,m), 7.59-7.42 (5H,m), 3.83 (3H,s), 2.92 (2H,q), 1.27 (3H,t).

Present Compound 255

20 $^1\text{H-NMR}$ (CDCl_3) δ : 8.51 (1H,d), 8.19 (1H,d), 7.53-7.42 (3H,m), 7.38-7.29 (1H,m), 3.73 (3H,s), 2.97-2.83 (4H,m), 1.33-1.19 (6H,m).

Present Compound 256

$^1\text{H-NMR}$ (CDCl_3) δ : 8.96 (1H,d), 8.54 (1H,d), 8.28-8.18 (1H,m), 7.91-7.80 (2H,m), 7.63-7.55 (1H,m), 3.74 (3H,s), 3.43 (2H,q), 3.24 (2H,q), 1.38 (3H,t), 1.26 (3H,t).

25 Present Compound 257

^1H -NMR (CDCl_3) δ : 8.53 (1H,d), 8.23 (1H,d), 7.54-7.42 (3H,m), 7.38-7.29 (1H,m), 3.74 (3H,s), 3.33-3.22 (1H,m), 2.87 (2H,q), 1.30 (6H,d), 1.24 (3H,t).

Present Compound 258

5 ^1H -NMR (CDCl_3) δ : 8.92 (1H,d), 8.51 (1H,d), 8.22 (1H,dd), 7.92-7.81 (2H,m), 7.62 (1H,dd), 3.75 (3H,s), 3.43 (2H,q), 3.36-3.26 (1H,m), 1.38 (6H,d), 1.26 (3H,t).

Present Compound 259

10 ^1H -NMR (CDCl_3) δ : 8.64-8.58 (1H,m), 8.33-8.29 (1H,m), 7.53-7.30 (4H,m), 3.75-3.73 (3H,m), 3.45-3.38 (2H,m), 2.89-2.82 (2H,m), 1.27-1.21 (3H,m).

Present Compound 260

15 ^1H -NMR (CDCl_3) δ : 9.01 (1H,d), 8.59 (1H,d), 8.25-8.21 (1H,m), 7.91-7.82 (2H,m), 7.63-7.58 (1H,m), 4.07 (2H,q), 3.75 (3H,s), 3.41 (2H,q), 1.26 (3H,t).

Present Compound 261

^1H -NMR (CDCl_3) δ : 8.48 (1H,d), 8.18 (1H,d), 7.53-7.41 (3H,m), 7.38-7.28 (1H,m), 5.95-5.81 (1H,m), 5.06-4.92 (2H,m), 3.73 (3H,s), 3.51 (2H,d), 2.86 (2H,q), 1.23 (3H,t).

20 Present Compound 262

^1H -NMR (CDCl_3) δ : 8.65-8.60 (1H,m), 8.40-8.35 (1H,m), 7.55-7.48 (2H,m), 7.47-7.41 (1H,m), 7.37-7.31 (1H,m), 3.76 (3H,s), 2.92-2.83 (2H,m), 1.27-1.18 (3H,m).

Present Compound 263

25 ^1H -NMR (CDCl_3) δ : 8.64 (1H,d), 8.39 (1H,d), 7.56-7.49

(2H,m), 7.47-7.43 (1H,m), 7.38-7.32 (1H,m), 3.76 (3H,s),
2.89 (2H,q), 1.24 (3H,t).

Present Compound 264

¹H-NMR (CDCl₃) δ: 8.63 (1H,d), 8.37 (1H,d), 7.56-7.42
5 (3H,m), 7.38-7.32 (1H,m), 3.76 (3H,s), 2.88 (2H,q), 1.24
(3H,t).

Present Compound 272

¹H-NMR (CDCl₃) δ: 8.67 (1H,d), 8.37 (1H,d), 8.26 (1H,dd),
7.83 (1H,td), 7.70 (1H,td), 7.61 (1H,dd), 3.88 (3H,s),
10 3.40-3.35 (1H,m), 3.01-2.95 (1H,m), 1.31 (3H,t).

Present Compound 273

¹H-NMR (CDCl₃) δ: 8.66 (1H,d), 8.34 (1H,d), 8.24-8.22
(1H,m), 7.89-7.78 (2H,m), 7.59-7.56 (1H,m), 3.71 (3H,s),
3.43 (2H,q), 1.25 (3H,t).

15 Present Compound 274

¹H-NMR (CDCl₃) δ: 8.68 (1H,d), 8.37 (1H,d), 8.26 (1H,dd),
7.83 (1H,td), 7.70 (1H,td), 7.61 (1H,dd), 3.89 (3H,s),
3.43-3.32 (1H,m), 3.04-2.93 (1H,m), 1.31 (3H,t).

Present Compound 275

20 ¹H-NMR (CDCl₃) δ: 8.67 (1H,d), 8.35 (1H,d), 8.23 (1H,dd),
7.89-7.77 (2H,m), 7.57 (1H,dd), 3.71 (3H,s), 3.43 (2H,q),
1.25 (3H,t).

Present Compound 276

25 ¹H-NMR (CDCl₃) δ: 8.56 (1H,d), 8.27 (1H,d), 7.53-7.41
(3H,m), 7.39-7.25 (1H,m), 3.74 (3H,s), 2.89 (2H,q), 1.33

(9H,s), 1.24 (3H,t).

Present Compound 277

¹H-NMR (CDCl₃) δ: 8.92 (1H,d), 8.52 (1H,d), 8.25-8.20
(1H,m), 7.89-7.84 (2H,m), 7.63-7.58 (1H,m), 3.75 (3H,s),
5 3.41 (2H,q), 1.43 (9H,s), 1.26 (3H,t).

Present Compound 278

¹H-NMR (CDCl₃) δ: 8.50 (1H,d), 8.24 (1H,d), 7.68-7.66
(1H,m), 7.60-7.54 (2H,m), 3.73 (3H,s), 2.94 (2H,q), 1.28
(3H,t).

10 Present Compound 279

¹H-NMR (CDCl₃) δ: 8.56-8.54 (2H,m), 8.23 (1H,d), 7.95-7.91
(1H,m), 7.77-7.74 (1H,m), 3.88 (3H,s), 3.50-3.40 (1H,m),
3.04-2.94 (1H,m), 1.34 (3H,t).

Present Compound 280

15 ¹H-NMR (CDCl₃) δ: 8.53 (1H,d), 8.49-8.48 (1H,m), 8.19
(1H,d), 8.10-8.07 (1H,m), 7.74-7.71 (1H,m), 3.68 (3H,s),
3.48 (2H,q), 1.28 (3H,t).

Present Compound 281

20 ¹H-NMR (CDCl₃) δ: 8.50 (1H,d), 8.20 (1H,d), 7.51-7.48
(2H,m), 7.46-7.43 (1H,m), 7.37-7.31 (1H,m), 5.96 (1H,t),
4.97 (2H,d), 3.73 (3H,s), 2.90-2.83 (2H,m), 1.23 (3H,t).

Present Compound 282

25 ¹H-NMR (CDCl₃) δ: 8.99 (1H,d), 8.54 (1H,d), 8.26-8.20
(1H,m), 7.88-7.80 (2H,m), 7.58-7.11 (1H,m), 6.36 (1H,t),
5.53 (2H,d), 3.76-3.68 (3H,m), 3.46-3.35 (2H,m), 1.30-1.21

(3H,m).

Present Compound 283

¹H-NMR (CDCl₃) δ: 8.93 (1H,d), 8.51 (1H,d), 8.25-8.22
(1H,m), 7.88-7.81 (2H,m), 7.59-7.57 (1H,m), 5.95-5.84
5 (1H,m), 5.42 (1H,d), 5.23 (1H,dd), 3.93 (2H,d), 3.73 (3H,s),
3.42 (2H,q), 1.26 (3H,t).

Present Compound 284

¹H-NMR (CDCl₃) δ: 8.26-8.15 (1H,m), 8.05-7.97 (1H,m), 7.87-
7.74 (2H,m), 7.62-7.55 (1H,m), 7.40-7.32 (1H,m), 3.70-3.57
10 (6H,m), 3.54-3.42 (2H,m), 1.30-1.19 (3H,m).

Present Compound 285

¹H-NMR (CDCl₃) δ: 8.77-8.75 (1H,m), 8.37-8.35 (1H,m), 8.28-
8.27 (1H,m), 8.18-8.14 (1H,m), 7.65-7.62 (1H,m), 3.81
(3H,s), 3.04 (2H,q), 1.34 (3H,t).

15 Present Compound 286

¹H-NMR (CDCl₃) δ: 9.13-9.11 (1H,m), 8.81-8.79 (1H,m), 8.54-
8.51 (1H,m), 8.37-8.34 (1H,m), 7.87 (1H,d), 3.98 (3H,s),
3.60-3.49 (1H,m), 3.13-3.02 (1H,m), 1.40 (3H,t).

Present Compound 287

20 ¹H-NMR (CDCl₃) δ: 9.08-9.05 (1H,m), 8.80-8.77 (1H,m), 8.68
(1H,dd), 8.34-8.31 (1H,m), 7.81 (1H,d), 3.75 (3H,s), 3.52
(2H,q), 1.32 (3H,t).

Present Compound 289

¹H-NMR (CDCl₃) δ: 8.19 (1H,dd), 8.01 (1H,d), 7.83-7.73
25 (2H,m), 7.55 (1H,dd), 7.35 (1H,d), 3.61 (3H,s), 3.46 (2H,q),

1.23 (3H,t).

Present Compound 290

¹H-NMR (CDCl₃) δ: 8.51 (1H,d), 8.21 (1H,dd), 8.07 (1H,d),
7.90-7.81 (2H,m), 7.59 (1H,dd), 3.69 (3H,s), 3.50-3.42
5 (5H,m), 1.25 (3H,t).

Present Compound 291

¹H-NMR (CDCl₃) δ: 8.21 (1H,dd), 8.16 (1H,d), 7.83-7.72
(2H,m), 7.55 (1H,dd), 7.40 (1H,d), 3.63 (3H,s), 3.48 (2H,q),
3.01 (6H,s), 1.24 (3H,t).

10 Present Compound 292

¹H-NMR (CDCl₃) δ: 8.69-8.68 (1H,m), 8.33-8.31 (1H,m), 7.70-
7.68 (1H,m), 7.62-7.55 (2H,m), 3.79 (3H,s), 2.97 (2H,q),
1.29 (3H,t).

Present Compound 293

15 ¹H-NMR (CDCl₃) δ: 8.86 (1H,d), 8.56 (1H,d), 7.96 (1H,d),
7.48 (1H,d), 7.32 (1H,dd), 3.02 (2H,t), 1.37 (3H,d).

Present Compound 294

¹H-NMR (CDCl₃) δ: 8.93 (1H,d), 8.52 (1H,d), 8.25 (1H,d),
7.78 (1H,dd), 7.67 (1H,d), 3.76 (2H,q), 1.40 (3H,t).

20 Present Compound 295

¹H-NMR (CDCl₃) δ: 8.74 (1H,s), 8.59 (1H,d), 7.56-7.49
(2H,m), 7.41 (1H,dd), 7.37-7.31 (1H,m), 6.84 (1H,brs), 3.75
(3H,s), 2.88 (2H,q), 1.23 (3H,t).

Present Compound 296

25 ¹H-NMR (CDCl₃) δ: 8.89 (1H,d), 8.56 (1H,d), 7.56-7.42

(3H,m), 7.39-7.33 (1H,m), 3.77 (3H,s), 2.89 (2H,q), 1.25 (3H,t).

Present Compound 297

¹H-NMR (CDCl₃) δ: 8.93 (1H,d), 8.53 (1H,d), 8.26 (1H,dd),
5 7.84 (1H,td), 7.71 (1H,td), 7.60 (1H,dd), 3.89 (3H,s),
3.42-3.32 (1H,m), 3.05-2.96 (1H,m), 1.31 (3H,t).

Present Compound 298

¹H-NMR (CDCl₃) δ: 8.91 (1H,d), 8.50 (1H,d), 8.23 (1H,dd),
7.87-7.80 (2H,m), 7.56 (1H,dd), 3.71 (3H,s), 3.45 (2H,q),
10 1.26 (3H,t).

Present Compound 299

¹H-NMR (CDCl₃) δ: 8.72-8.70 (1H,m), 8.51-8.49 (1H,m), 8.29-
8.27 (1H,m), 8.13-8.09 (1H,m), 7.74-7.71 (1H,m), 3.74
(3H,s), 3.49 (2H,q), 1.29 (3H,t).

15 Present Compound 300

¹H-NMR (CDCl₃) δ: 8.60 (1H,d), 8.39 (1H,d), 7.65-7.64
(1H,m), 7.58-7.51 (2H,m), 3.70 (3H,s), 2.91 (2H,q), 1.25
(3H,t).

Present Compound 303

20 ¹H-NMR (CDCl₃) δ: 8.09 (1H,s), 7.52 (1H,d), 7.46-7.35
(3H,m), 7.25 (1H,t), 7.12 (1H,d), 3.56 (3H,s), 2.78 (2H,q),
1.17 (3H,t).

Present Compound 307

¹H-NMR (CDCl₃) δ: 8.65 (1H,d), 8.49-8.47 (1H,m), 8.37
25 (1H,d), 8.10-8.06 (1H,m), 7.74-7.71 (1H,m), 3.67 (3H,s),

3.47 (2H,q), 1.28 (3H,t).

Present Compound 308

¹H-NMR (CDCl₃) δ: 8.09 (1H,s), 7.59-7.42 (5H,m), 7.37-7.31 (1H,m), 3.69 (3H,s), 2.85 (2H,q), 1.23 (3H,t).

5 Present Compound 309

¹H-NMR (CDCl₃) δ: 8.22 (1H,d), 8.07 (1H,s), 7.80 (1H,t), 7.67 (1H,t), 7.62-7.52 (3H,m), 3.79 (3H,s), 3.37-3.26 (1H,m), 3.01-2.89 (1H,m), 1.27 (3H,t).

Present Compound 310

10 ¹H-NMR (CDCl₃) δ: 8.20-8.18 (1H,m), 8.06 (1H,s), 7.82-7.74 (2H,m), 7.59-7.52 (3H,m), 3.61 (3H,s), 3.43 (2H,brs), 1.22 (3H,t).

Present Compound 311

15 ¹H-NMR (CDCl₃) δ: 7.83 (1H,d), 7.69 (1H,d), 7.60 (1H,dd), 7.54-7.49 (1H,m), 7.48-7.42 (1H,m), 7.41-7.35 (1H,m), 6.69 (1H,d), 3.00 (2H,q), 2.93 (3H,s), 1.32 (3H,t).

Present Compound 312

20 ¹H-NMR (CDCl₃) δ: 8.25-8.19 (2H,m), 7.83-7.78 (2H,m), 7.68 (1H,t), 7.55 (1H,d), 7.47 (1H,d), 3.79 (3H,s), 3.36-3.26 (1H,m), 3.00-2.90 (1H,m), 1.27 (3H,t).

Present Compound 313

¹H-NMR (CDCl₃) δ: 8.23-8.20 (2H,m), 7.85-7.76 (3H,m), 7.57-7.53 (1H,m), 7.46 (1H,d), 3.62 (3H,s), 3.39 (2H,brs), 1.24 (3H,t).

25 Present Compound 317

^1H -NMR (CDCl_3) δ : 8.65-8.64 (1H,m), 8.29-8.28 (1H,m), 7.35-7.31 (2H,m), 7.18-7.15 (1H,m), 3.76 (3H,s), 2.86 (2H,q), 2.46 (3H,s), 1.23 (3H,t).

Present Compound 318

5 ^1H -NMR (CDCl_3) δ : 8.69-8.68 (1H,m), 8.28-8.26 (1H,m), 8.07-8.05 (1H,m), 7.50-7.48 (2H,m), 3.89 (3H,s), 3.44-3.33 (1H,m), 3.01-2.92 (1H,m), 2.58 (3H,s), 1.32 (3H,t).

Present Compound 319

10 ^1H -NMR (CDCl_3) δ : 8.68-8.66 (1H,m), 8.25-8.23 (1H,m), 8.04-8.02 (1H,m), 7.65-7.61 (1H,m), 7.45-7.42 (1H,m), 3.71 (3H,s), 3.47-3.38 (2H,m), 2.59 (3H,s), 1.26 (3H,t).

Present Compound 320

15 ^1H -NMR (CDCl_3) δ : 8.77 (1H,d), 8.68 (1H,d), 8.00-7.94 (1H,m), 7.55-7.51 (1H,m), 7.48-7.43 (1H,m), 7.36-7.31 (1H,m), 2.97 (2H,q), 1.33 (3H,t).

Present Compound 321

^1H -NMR (CDCl_3) δ : 8.81 (1H,d), 8.71 (1H,d), 8.08 (1H,d), 7.72 (1H,s), 7.56 (1H,d), 3.03 (2H,q), 1.37 (3H,t).

Present Compound 322

20 ^1H -NMR (CDCl_3) δ : 8.81 (1H,d), 8.55 (1H,d), 8.07-8.00 (1H,m), 7.59-7.53 (1H,m), 7.52-7.45 (1H,m), 7.41-7.33 (1H,m), 2.99 (2H,q), 1.35 (3H,t).

Present Compound 323

25 ^1H -NMR (CDCl_3) δ : 8.85 (1H,d), 8.59 (1H,d), 8.15 (1H,d), 7.75 (1H,s), 7.59 (1H,d), 3.05 (2H,q), 1.38 (3H,t).

Present Compound 324

$^1\text{H-NMR}$ (CDCl_3) δ : 8.87 (1H,d), 8.50 (1H,d), 8.28-8.22 (1H,m), 7.85-7.76 (2H,m), 7.74-7.70 (1H,m), 3.74 (2H,q), 1.37 (3H,t).

5 Present Compound 325

$^1\text{H-NMR}$ (CDCl_3) δ : 8.90 (1H,s), 8.57-8.49 (2H,m), 8.07 (1H,d), 7.88 (1H,d), 3.77 (2H,q), 1.40 (3H,t).

Present Compound 326

10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.93 (1H,d), 8.61 (1H,d), 7.45-7.34 (3H,m), 2.91 (2H,q), 1.27 (3H,t).

Present Compound 327

$^1\text{H-NMR}$ (CDCl_3) δ : 8.95 (1H,d), 8.55 (1H,d), 8.14 (1H,dd), 7.86 (1H,dd), 7.73 (1H,t), 3.40 (2H,q), 1.28 (3H,t).

Present Compound 330

15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.74-8.73 (1H,m), 8.28-8.27 (1H,m), 8.22 (1H,d), 7.82 (1H,dd), 7.51 (1H,d), 6.86 (1H,dd), 6.03 (1H,d), 5.58 (1H,d), 3.72 (3H,s), 3.44 (2H,q), 1.27 (3H,t).

Present Compound 331

20 $^1\text{H-NMR}$ (CDCl_3) δ : 8.74-8.72 (1H,m), 8.28-8.26 (1H,m), 8.05-8.03 (1H,m), 7.67-7.64 (1H,m), 7.46 (1H,d), 3.71 (3H,s), 3.41 (2H,q), 2.88 (2H,q), 1.37 (3H,t), 1.26 (3H,t).

Present Compound 332

25 $^1\text{H-NMR}$ (CDCl_3) δ : 8.78-8.76 (1H,m), 8.46-8.44 (1H,m), 8.31-8.30 (1H,m), 8.09-8.05 (1H,m), 7.76 (1H,d), 3.75 (3H,s), 3.48 (2H,q), 1.27 (3H,t).

Present Compound 334

¹H-NMR (CDCl₃) δ: 8.89 (1H,d), 8.46 (1H,d), 8.27 (1H,dd),
7.85 (1H,td), 7.71 (1H,td), 7.62 (1H,dd), 3.92 (3H,s),
3.45-3.34 (1H,m), 3.04-2.94 (1H,m), 2.18 (6H,s), 1.32
5 (3H,t).

Present Compound 335

¹H-NMR (CDCl₃) δ: 8.87 (1H,d), 8.44 (1H,d), 8.26-8.19
(1H,m), 7.93-7.81 (2H,m), 7.64-7.60 (1H,m), 3.74 (3H,s),
3.43 (2H,q), 2.80 (6H,s), 1.26 (3H,t).

10 Present Compound 337

¹H-NMR (CDCl₃) δ: 9.26 (1H,s), 7.80 (1H,s), 7.43-7.36
(1H,m), 7.34-7.30 (1H,m), 7.27-7.21 (2H,m), 7.17 (1H,td),
3.66 (3H,s), 2.77 (2H,q), 2.12 (3H,s), 1.15 (3H,t).

Present Compound 338

15 ¹H-NMR (CDCl₃) δ: 7.82-7.80 (1H,m), 7.50-7.43 (3H,m), 7.37-
7.30 (1H,m), 7.19 (1H,s), 3.93 (3H,s), 2.87-2.82 (8H,m),
1.22 (3H,t).

Present Compound 339

20 ¹H-NMR (CDCl₃) δ: 7.65 (1H,s), 7.60-7.53 (3H,m), 6.82
(1H,s), 4.01 (2H,brs), 3.94 (3H,s), 2.92 (2H,q), 1.27
(3H,t).

Present Compound 340

25 ¹H-NMR (CDCl₃) δ: 7.82 (1H,s), 7.67 (1H,s), 7.61-7.53
(2H,m), 7.23 (1H,s), 3.96 (3H,s), 2.93 (2H,q), 2.86 (6H,s),
1.28 (3H,t).

Present Compound 347

$^1\text{H-NMR}$ (CDCl_3) δ : 8.58 (1H,s), 7.56-7.29 (5H,m), 3.73 (3H,s), 2.88 (2H,q), 1.24 (3H,t).

Present Compound 348

5 $^1\text{H-NMR}$ (CDCl_3) δ : 8.61 (1H,s), 8.28 (1H,dd), 7.81 (1H,td), 7.68 (1H,td), 7.62 (1H,dd), 7.27 (1H,t), 3.91 (3H,s), 3.68-3.58 (1H,m), 3.12-3.02 (1H,m), 1.37 (3H,t).

Present Compound 349

10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.59 (1H,s), 8.21 (1H,dd), 7.86-7.77 (2H,m), 7.60 (1H,dd), 7.27 (1H,t), 3.71 (3H,s), 3.54 (2H,q), 1.27 (3H,t).

Present Compound 350

15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.46 (1H,s), 7.51-7.32 (4H,m), 3.86-3.71 (1H,m), 3.68 (3H,s), 2.86 (2H,q), 1.61 (3H,s), 1.59 (3H,s), 1.22 (3H,t).

Present Compound 352

$^1\text{H-NMR}$ (CDCl_3) δ : 8.69 (1H,s), 8.33 (1H,s), 8.25 (1H,dd), 8.21 (1H,dd), 7.73-7.58 (2H,m), 3.47-3.36 (1H,m), 3.13-3.01 (1H,m), 1.57-0.71 (3H,m).

20 Present Compound 355

$^1\text{H-NMR}$ (CDCl_3) δ : 8.76 (1H,d), 8.51 (1H,d), 8.29 (1H,d), 8.10 (1H,dd), 7.74 (1H,d), 4.22 (2H,q), 3.55 (2H,q), 1.42 (3H,t), 1.30 (3H,t).

Present Compound 356

25 $^1\text{H-NMR}$ (CDCl_3) δ : 8.77 (1H,d), 8.48 (1H,s), 8.31 (1H,d),

8.05 (1H,d), 7.73 (1H,d), 6.02-5.90 (1H,m), 5.19 (1H,d),
4.96 (1H,d), 4.79 (2H,s), 3.51 (2H,q), 1.29 (3H,t).

Present Compound 357

¹H-NMR (CDCl₃) δ: 8.81 (1H,d), 8.49 (1H,s), 8.33 (1H,d),
5 8.11 (1H,d), 7.94 (1H,d), 4.95 (2H,brs), 3.46 (2H,q), 2.34-
2.31 (1H,m), 1.29 (3H,t).

Present Compound 358

¹H-NMR (CDCl₃) δ: 8.83 (1H,d), 8.50 (1H,d), 8.37 (1H,d),
8.18 (1H,dd), 7.93 (1H,d), 5.10 (2H,brs), 3.35 (2H,q), 1.27
10 (3H,t).

Present Compound 359

¹H-NMR (CDCl₃) δ: 8.98 (1H,d), 8.50 (1H,d), 8.21-8.17
(2H,m), 7.89 (1H,d), 4.97 (1H,brs), 4.85 (1H,brs), 3.47
(1H,brs), 3.34 (1H,brs), 1.28-1.20 (3H,m).

15 Present Compound 360

¹H-NMR (CDCl₃) δ: 8.77 (1H,d), 8.47 (1H,d), 8.32 (1H,d),
8.08 (1H,dd), 7.90 (1H,d), 5.53 (2H,s), 3.67 (2H,q), 3.58
(2H,q), 1.31 (3H,t), 1.18 (3H,t).

Present Compound 361

20 ¹H-NMR (CDCl₃) δ: 8.91 (1H,d), 8.48 (1H,d), 8.21 (1H,d),
8.09 (1H,dd), 7.81 (1H,d), 5.36 (2H,s), 3.60-3.49 (4H,m),
1.28 (3H,t), 1.19 (3H,t).

Present Compound 362

¹H-NMR (CDCl₃) δ: 8.76 (1H,d), 8.47 (1H,d), 8.31 (1H,d),
25 8.10 (1H,dd), 7.98 (1H,d), 5.23 (2H,brs), 3.45 (2H,q), 2.26

(3H,s), 1.28 (3H,t).

Present Compound 363

¹H-NMR (CDCl₃) δ: 8.75 (1H,d), 8.50 (1H,d), 8.29 (1H,d),
8.09 (1H,dd), 7.74 (1H,d), 4.12 (2H,q), 3.53 (2H,q), 1.89-
5 1.79 (2H,m), 1.29 (3H,t), 0.90 (3H,t).

Present Compound 364

¹H-NMR (CDCl₃) δ: 8.75 (1H,d), 8.51 (1H,d), 8.29 (1H,d),
8.09 (1H,dd), 7.74 (1H,d), 4.17-4.10 (2H,m), 3.53 (2H,q),
1.82-1.75 (2H,m), 1.34-1.24 (5H,m), 0.86 (3H,t).

10 Present Compound 365

¹H-NMR (CDCl₃) δ: 8.75 (1H,d), 8.46 (1H,d), 8.33(1H,d),
8.05 (1H,dd), 7.87 (1H,d), 4.91 (2H,brs), 3.77 (3H,s), 3.41
(2H,q), 1.27 (3H,t).

Present Compound 366

15 ¹H-NMR (CDCl₃) δ: 8.92 (1H,d), 8.47 (1H,d), 8.07 (1H,dd),
7.96 (1H,d), 7.83 (1H,d), 4.82 (1H,brs), 4.69 (1H,brs),
3.81 (3H,s), 3.54 (1H,brs), 3.40 (1H,brs), 1.26 (3H,t).

Present Compound 367

20 ¹H-NMR (CDCl₃) δ: 8.77 (1H,d), 8.50 (1H,d), 8.32 (1H,d),
8.11 (1H,dd), 7.73 (1H,d), 6.12 (1H,t), 4.92 (2H,d), 3.48
(2H,q), 1.29 (3H,t).

Present Compound 368

25 ¹H-NMR (CDCl₃) δ: 8.76 (1H,d), 8.42 (1H,d), 8.21 (1H,d),
7.99 (1H,dd), 7.67 (1H,d), 3.60-3.50 (1H,m), 3.40-3.30
(1H,m), 1.70 (9H,s), 1.28 (3H,t).

Present Compound 369

¹H-NMR (CDCl₃) δ: 8.13 (1H,s), 7.67 (1H,s), 7.62 (1H,d), 7.59-7.57 (2H,m), 7.52 (1H,d), 3.70 (3H,s), 2.92 (2H,q), 1.27 (3H,t).

5 Present Compound 370

¹H-NMR (CDCl₃) δ: 8.53 (1H,s), 8.10 (1H,s), 7.93 (1H,d), 7.75 (1H,d), 7.65 (1H,d), 7.57 (1H,d), 3.85 (3H,s), 3.52-3.41 (1H,m), 3.05-2.95 (1H,m), 1.32 (3H,t).

Present Compound 371

10 ¹H-NMR (CDCl₃) δ: 8.48 (1H,s), 8.10-8.05 (2H,m), 7.74 (1H,d), 7.62 (1H,d), 7.53 (1H,d), 3.63 (3H,s), 3.47 (2H,q), 1.25 (3H,t).

Present Compound 372

15 ¹H-NMR (CDCl₃) δ: 8.77 (1H,d), 8.50 (1H,d), 8.30 (1H,d), 8.10 (1H,dd), 7.73 (1H,d), 3.74 (3H,s), 3.48 (2H,q), 1.29 (3H,t).

Present Compound 373

20 ¹H-NMR (CDCl₃) δ: 8.83 (1H,d), 8.37-8.34 (2H,m), 8.04 (1H,dd), 7.72 (1H,d), 3.30-3.11 (2H,brm), 3.04 (3H,s), 1.24 (3H,t).

Present Compound 374

¹H-NMR (CDCl₃) δ: 8.74-8.72 (1H,m), 8.33-8.32 (1H,m), 7.48 (1H,d), 7.29-7.27 (1H,m), 7.21-7.17 (1H,m), 3.79 (3H,s), 2.92 (2H,q), 1.29 (3H,t).

25 Present Compound 375

$^1\text{H-NMR}$ (CDCl_3) δ : 8.78-8.76 (1H,m), 8.33-8.31 (1H,m), 8.14-8.12 (1H,m), 7.68 (1H,d), 7.54-7.50 (1H,m), 3.93 (3H,s), 3.49-3.39 (1H,m), 3.06-2.96 (1H,m), 1.33 (3H,t).

Present Compound 376

5 $^1\text{H-NMR}$ (CDCl_3) δ : 8.77-8.75 (1H,m), 8.30-8.28 (1H,m), 8.09-8.07 (1H,m), 7.70-7.66 (1H,m), 7.63 (1H,d), 3.74 (3H,s), 3.46 (2H,q), 1.28 (3H,t).

Present Compound 377

10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.78-8.77 (1H,m), 8.47-8.45 (1H,m), 8.32-8.30 (1H,m), 8.09-8.06 (1H,m), 7.76 (1H,d), 3.75 (3H,s), 3.48 (2H,q), 1.27 (3H,t).

Present Compound 378

15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.83-8.74 (3H,m), 8.54 (1H,dd), 8.31-8.29 (1H,m), 7.95-7.86 (2H,m), 7.68 (1H,d), 7.41-7.37 (1H,m), 3.75 (3H,s), 3.49 (2H,q), 1.30 (3H,t).

Present Compound 379

$^1\text{H-NMR}$ (CDCl_3) δ : 8.72-8.70 (1H,m), 8.27-8.25 (1H,m), 7.43 (1H,d), 7.29 (1H,d), 6.99 (1H,dd), 4.28 (2H,brs), 3.70 (3H,s), 3.40 (2H,q), 1.26 (3H,t).

20 Present Compound 380

$^1\text{H-NMR}$ (DMSO-D_6) δ : 11.94 (1H,brs), 8.86-8.84 (1H,m), 8.62-8.60 (1H,m), 8.53 (1H,d), 8.28 (1H,dd), 7.95 (1H,d), 3.64 (3H,s), 3.56 (2H,q), 1.14 (3H,t).

Present Compound 381

25 $^1\text{H-NMR}$ (CDCl_3) δ : 8.74-8.72 (1H,m), 8.28-8.25 (1H,m), 8.07-

8.05 (1H,m), 7.69-7.66 (1H,m), 7.47 (1H,d), 3.72 (3H,s),
3.41 (2H,q), 3.18-3.10 (1H,m), 1.37 (6H,d), 1.26 (3H,t).

Present Compound 382

¹H-NMR (CDCl₃) δ: 10.23 (1H,s), 8.78-8.76 (1H,m), 8.71
5 (1H,d), 8.35 (1H,dd), 8.31 (1H,d), 7.77 (1H,d), 3.74 (3H,s),
3.49 (2H,q), 1.30 (3H,t).

Present Compound 383

¹H-NMR (CDCl₃) δ: 8.72-8.70 (1H,m), 8.32-8.31 (1H,m), 7.55-
7.49 (2H,m), 7.46-7.43 (1H,m), 7.38-7.33 (1H,m), 3.77
10 (3H,s), 2.84 (2H,t), 1.58-1.50 (2H,m), 1.39-1.29 (2H,m),
0.85 (3H,t).

Present Compound 384

¹H-NMR (CDCl₃) δ: 8.73-8.72 (1H,m), 8.33-8.31 (1H,m), 7.66-
7.63 (1H,m), 7.58-7.53 (1H,m), 7.49-7.41 (2H,m), 3.79
15 (3H,s), 3.66 (3H,s), 3.60 (2H,s).

Present Compound 385

¹H-NMR (CDCl₃) δ: 8.74-8.72 (1H,m), 8.34-8.32 (1H,m), 7.61-
7.55 (2H,m), 7.52-7.44 (2H,m), 5.85 (1H,t), 3.77 (3H,s),
3.54 (2H,d).

20 Present Compound 386

¹H-NMR (CDCl₃) δ: 8.72-8.71 (1H,m), 8.32-8.31 (1H,m), 7.68-
7.65 (1H,m), 7.59-7.54 (1H,m), 7.47 (1H,dd), 7.44-7.39
(1H,m), 3.78 (3H,s), 3.55 (2H,q), 1.75 (3H,t).

Present Compound 387

25 ¹H-NMR (CDCl₃) δ: 8.75-8.73 (1H,m), 8.33-8.31 (1H,m), 7.87-

7.84 (1H,m), 7.70-7.64 (1H,m), 7.63-7.58 (1H,m), 7.56
(1H,dd), 3.78 (3H,s), 3.61 (2H,s).

Present Compound 388

¹H-NMR (CDCl₃) δ: 8.72-8.71 (1H,m), 8.33-8.31 (1H,m), 7.57-
5 7.48 (2H,m), 7.45-7.42 (1H,m), 7.37-7.32 (1H,m), 3.78
(3H,s), 2.82 (2H,s), 0.91 (9H,s).

Present Compound 389

¹H-NMR (CDCl₃) δ: 8.75-8.73 (1H,m), 8.33-8.31 (1H,m), 7.71-
7.68 (1H,m), 7.63-7.58 (1H,m), 7.55-7.51 (2H,m), 3.76
10 (3H,s), 2.97 (2H,t), 2.52 (2H,t).

Present Compound 390

¹H-NMR (CDCl₃) δ: 8.72-8.70 (1H,m), 8.33-8.31 (1H,m), 7.53-
7.49 (2H,m), 7.46-7.42 (1H,m), 7.39-7.33 (1H,m), 3.76
(3H,s), 2.90 (2H,d), 2.50-2.37 (1H,m), 2.06-1.96 (2H,m),
15 1.87-1.71 (2H,m), 1.67-1.56 (2H,m).

Present Compound 391

¹H-NMR (CDCl₃) δ: 8.76-8.74 (1H,m), 8.35-8.34 (1H,m), 7.98-
7.94 (1H,m), 7.71-7.67 (2H,m), 7.63-7.60 (1H,m), 3.78
(3H,s).

20 Present Compound 392

¹H-NMR (CDCl₃) δ: 8.77-8.75 (1H,m), 8.38-8.36 (1H,m), 8.31-
8.29 (1H,m), 8.02-7.99 (1H,m), 7.69 (1H,d), 6.85 (1H,t),
3.73 (3H,s), 3.54-3.33 (2H,m), 1.28 (3H,t).

Present Compound 393

25 ¹H-NMR (CDCl₃) δ: 8.75-8.73 (1H,m), 8.29-8.27 (1H,m), 8.21-

8.19 (1H,m), 7.88-7.84 (1H,m), 7.56 (1H,d), 4.97-4.93
(2H,m), 3.71 (3H,s), 3.43 (2H,q), 1.26 (3H,t).

Present Compound 394

¹H-NMR (CDCl₃) δ: 8.75-8.73 (1H, m), 8.28-8.26 (1H, m),
5 8.25-8.22 (1H, m), 7.87-7.79 (2H, m), 7.58-7.55 (1H, m),
3.72 (3H, s), 3.43-3.37 (2H, m), 1.70-1.60 (2H, m), 1.45-
1.35 (2H, m), 0.90 (3H, t).

Present Compound 395

¹H-NMR (CDCl₃) δ: 8.76-8.74 (1H, m), 8.33-8.29 (2H, m),
10 7.91-7.81 (2H, m), 7.60-7.57 (1H, m), 4.68 (2H, br s), 3.74
(3H, s), 3.72 (3H, s).

Present Compound 396

¹H-NMR (CDCl₃) δ: 8.75-8.74 (1H, m), 8.29-8.27 (1H, m),
8.22-8.20 (1H, m), 7.92-7.82 (2H, m), 7.61-7.58 (1H, m),
15 5.92 (1H, t), 4.40 (2H, d), 3.73 (3H, s).

Present Compound 397

¹H-NMR (CDCl₃) δ: 8.75-8.73 (1H, m), 8.32-8.28 (2H, m),
7.89-7.80 (2H, m), 7.60-7.57 (1H, m), 4.35 (2H, br s), 3.73
(3H, s), 1.78 (3H, t).

20 Present Compound 398

¹H-NMR (CDCl₃) δ: 8.78-8.76 (1H, m), 8.43-8.40 (1H, m),
8.30-8.29 (1H, m), 8.00-7.95 (1H, m), 7.94-7.89 (1H, m),
7.68-7.65 (1H, m), 4.83 (2H, br s), 3.78 (3H, s).

Present Compound 399

25 ¹H-NMR (CDCl₃) δ: 8.75-8.74 (1H, m), 8.27-8.24 (2H, m),

7.83-7.80 (2H, m), 7.55-7.52 (1H, m), 3.71 (3H, s), 3.45
(2H, br s), 1.06 (9H, s).

Present Compound 400

¹H-NMR (CDCl₃) δ: 8.77-8.75 (1H, m), 8.33-8.32 (1H, m),
5 8.29-8.26 (1H, m), 7.95-7.90 (1H, m), 7.89-7.84 (1H, m),
7.64-7.61 (1H, m), 3.87 (2H, t), 3.73 (3H, s), 2.78 (2H, t).

Present Compound 401

¹H-NMR (CDCl₃) δ: 8.75-8.73 (1H, m), 8.32-8.30 (1H, m),
8.22-8.19 (1H, m), 7.86-7.78 (2H, m), 7.57-7.54 (1H, m),
10 3.70 (3H, s), 3.51 (2H, d), 2.75-2.63 (1H, m), 2.06-1.96
(2H, m), 1.94-1.70 (4H, m).

Present Compound 402

¹H-NMR (CDCl₃) δ: 8.75-8.73 (1H, m), 8.33-8.30 (2H, m),
8.03-7.98 (1H, m), 7.95-7.90 (1H, m), 7.70-7.68 (1H, m),
15 3.70 (3H, s).

Present Compound 403

¹H-NMR (CDCl₃) δ: 8.76-8.74 (1H, m), 8.29-8.28 (1H, m),
8.21-8.19 (1H, m), 7.87-7.84 (1H, m), 7.63-7.60 (1H, m),
5.61 (2H, d), 3.72 (3H, s), 3.54-3.32 (2H, m), 1.27 (3H, t).

20 Present Compound 404

¹H-NMR (CDCl₃) δ: 8.72-8.71 (1H, m), 8.33-8.31 (1H, m),
7.61-7.50 (3H, m), 7.47-7.42 (1H, m), 6.42 (1H, dd), 5.37
(1H, d), 5.35 (1H, d), 3.79 (3H, s).

Present Compound 405

25 ¹H-NMR (CDCl₃) δ: 8.76-8.75 (1H, m), 8.35-8.33 (1H, m),

8.17-8.14 (1H, m), 7.82-7.77 (1H, m), 7.69-7.61 (2H, m),
7.42 (1H, dd), 6.17 (1H, d), 5.86 (1H, d), 3.97 (3H, s).

Present Compound 406

¹H-NMR (CDCl₃) δ: 8.75-8.74 (1H, m), 8.30-8.29 (1H, m),
5 8.25-8.22 (1H, m), 7.84-7.80 (2H, m), 7.57-7.54 (1H, m),
7.10 (1H, dd), 6.31 (1H, d), 6.03 (1H, d), 3.74 (3H, s).

Present Compound 407

¹H-NMR (CDCl₃) δ: 8.47 (1H, d), 8.04 (1H, dd), 7.84 (1H, d),
7.56 (1H, dd), 7.48 (1H, td), 7.37 (1H, td), 2.98 (2H, q),
10 1.34 (3H, t).

Present Compound 408

¹H-NMR (CDCl₃) δ: 8.44 (1H, d), 8.25 (1H, dd), 7.89 (1H, d),
7.84-7.71 (3H, m), 3.72 (2H, q), 1.37 (3H, t).

Present Compound 409

¹H-NMR (CDCl₃) δ: 8.68-8.66 (1H, m), 8.31-8.30 (1H, m),
15 7.48 (1H, d), 7.29-7.26 (1H, m), 7.21-7.17 (1H, m), 3.80
(3H, s), 2.93 (2H, q), 1.29 (3H, t).

Present Compound 410

¹H-NMR (CDCl₃) δ: 8.72-8.71 (1H, m), 8.30-8.29 (1H, m),
20 8.14-8.12 (1H, m), 7.68 (1H, d), 7.55-7.51 (1H, m), 3.93
(3H, s), 3.50-3.40 (1H, m), 3.08-2.98 (1H, m), 1.33 (3H, t).

Present Compound 411

¹H-NMR (CDCl₃) δ: 8.71-8.69 (1H, m), 8.27-8.26 (1H, m),
8.09-8.07 (1H, m), 7.70-7.66 (1H, m), 7.62 (1H, d), 3.74
25 (3H, s), 3.47 (2H, q), 1.28 (3H, t).

Present Compound 412

^1H -NMR (DMSO- D_6) δ : 7.88 (1H, s), 7.81 (1H, s), 7.73 (1H, d), 7.69 (1H, d), 7.39 (1H, s), 3.84 (3H, s), 3.09 (2H, q), 2.84 (3H, s), 1.19 (3H, t).

5 Present Compound 413

^1H -NMR (CDCl_3) δ : 8.50 (1H, s), 7.94-7.89 (2H, m), 7.69 (1H, d), 7.35 (1H, s), 3.99 (3H, s), 3.42-3.32 (1H, m), 3.01-2.90 (1H, m), 2.85 (3H, s), 1.28 (3H, t).

Present Compound 414

10 ^1H -NMR (CDCl_3) δ : 8.48 (1H, d), 8.07 (1H, dd), 7.88 (1H, s), 7.72 (1H, d), 7.32 (1H, s), 3.82 (3H, s), 3.45 (2H, br s), 2.82 (3H, s), 1.26 (3H, t).

Present Compound 415

15 ^1H -NMR (DMSO- D_6) δ : 8.13 (1H, d), 7.96 (1H, d), 7.82 (1H, s), 7.77-7.71 (2H, m), 3.65 (3H, s), 3.09 (2H, q), 1.18 (3H, t).

Present Compound 416

20 ^1H -NMR (CDCl_3) δ : 8.52 (1H, d), 8.06 (1H, d), 7.93 (1H, dd), 7.72 (1H, d), 7.28 (1H, d), 3.80 (3H, s), 3.49-3.39 (1H, m), 3.05-2.95 (1H, m), 1.33 (3H, t).

Present Compound 417

^1H -NMR (CDCl_3) δ : 8.48 (1H, d), 8.09 (1H, dd), 8.02 (1H, d), 7.74 (1H, d), 7.25 (1H, d), 3.60 (3H, s), 3.45 (2H, q), 1.26 (3H, t).

25 Present Compound 418

^1H -NMR (CDCl_3) δ : 8.11 (1H, s), 7.67 (1H, s), 7.61-7.56 (3H, m), 7.53 (1H, d), 3.70 (3H, s), 2.93 (2H, q), 1.28 (3H, t).

Present Compound 419

5 ^1H -NMR (CDCl_3) δ : 8.53 (1H, d), 8.09 (1H, s), 7.93 (1H, dd), 7.74 (1H, d), 7.64 (1H, d), 7.59 (1H, d), 3.85 (3H, s), 3.51-3.40 (1H, m), 3.06-2.97 (1H, m), 1.32 (3H, t).

Present Compound 420

10 ^1H -NMR (CDCl_3) δ : 8.49 (1H, d), 8.10-8.04 (2H, m), 7.73 (1H, d), 7.60 (1H, d), 7.55 (1H, d), 3.64 (3H, s), 3.47 (2H, q), 1.25 (3H, t).

Present Compound 421

15 ^1H -NMR (CDCl_3) δ : 8.72-8.71 (1H, m), 8.32-8.31 (1H, m), 7.57-7.49 (2H, m), 7.47-7.44 (1H, m), 7.39-7.34 (1H, m), 3.78 (3H, s), 2.78 (2H, d), 0.99-0.89 (1H, m), 0.54-0.48 (2H, m), 0.18-0.14 (2H, m).

Present Compound 422

20 ^1H -NMR (CDCl_3) δ : 8.76-8.75 (1H, m), 8.34-8.31 (1H, m), 8.28-8.26 (1H, m), 7.85-7.80 (1H, m), 7.71-7.67 (1H, m), 7.61-7.58 (1H, m), 3.91 (3H, s), 3.27 (1H, dd), 3.04 (1H, dd), 1.33-1.22 (1H, m), 0.71-0.62 (2H, m), 0.43-0.31 (2H, m).

Present Compound 423

25 ^1H -NMR (CDCl_3) δ : 8.75-8.73 (1H, m), 8.31-8.26 (2H, m), 7.88-7.79 (2H, m), 7.57-7.54 (1H, m), 3.72 (3H, s), 3.34 (2H, d), 1.05-0.94 (1H, m), 0.57-0.50 (2H, m), 0.26-0.20

(2H, m).

Present Compound 424

¹H-NMR (CDCl₃) δ: 8.30 (1H, d), 8.20 (1H, s), 7.78-7.67 (2H, m), 7.63 (1H, s), 7.52 (1H, d), 3.11 (2H, q), 1.47 (3H, t).

5 Present Compound 425

¹H-NMR (CDCl₃) δ: 8.53 (1H, s), 8.18-8.12 (2H, m), 8.08 (1H, d), 7.77-7.72 (2H, m), 3.91 (2H, q), 1.44 (3H, t).

Present Compound 426

10 ¹H-NMR (CDCl₃) δ: 8.12 (1H, s), 7.66 (1H, s), 7.62 (1H, dd), 7.57 (1H, br s), 7.52 (1H, d), 3.70 (3H, s), 2.92 (2H, q), 1.26 (3H, t).

Present Compound 427

15 ¹H-NMR (CDCl₃) δ: 7.95 (1H, d), 7.64 (1H, s), 7.57-7.51 (3H, m), 4.03 (3H, s), 3.04 (2H, q), 2.93 (2H, q), 1.37 (3H, t), 1.29-1.20 (3H, m).

[0485]


The productions of the intermediates of the present compound are shown in the following Reference Production Examples.

20 [0486]

Reference Production Example 1

25 A mixture of N²-methyl-5-trifluoromethylpyridin-2,3-diamine (1.91 g), 2-fluorobenzaldehyde (1.86 g), sodium hydrogen sulfite (1.56 g), and DMF (10 ml) was stirred with heating under reflux at 153 °C for 3 hours. Into the


reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.89 g of 2-(2-fluorophenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine (Compound (1E)-F1).



¹H-NMR (CDCl₃) δ: 8.73-8.72 (1H,m), 8.33-8.32 (1H,m), 7.76-7.72 (1H,m), 7.64-7.58 (1H,m), 7.41-7.36 (1H,m), 7.32-7.27 (1H,m), 3.89 (3H,d)
[0487]

Reference Production Example 2-a

To a mixture of 2-chloro-5-trifluoromethylpyridine (18.2) and N-methylpyrrolidone (100 ml), 40 % of aqueous methylamine solution (23.3 g) was added, and stirred at room temperature for 3 hours. Water was poured thereinto, and extracted with ethyl acetate 3 times. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 17.3 g of N-methyl-(5-trifluoromethyl-pyridin-2-yl)-amine.




$^1\text{H-NMR}$ (CDCl_3) δ : 8.34 (s, 1H), 7.60 (dd, 1H), 6.40 (d, 1H), 4.89 (brs, 1H), 2.97 (d, 3H)

[0488]

5 Reference Production Example 2-b

To a mixture of N-methyl-(5-trifluoromethyl-pyridin-2-yl)-amine (17.3 g) and sulfuric acid (100 ml), nitric acid (12.3 g) was added dropwise, and stirred with heating at 80 °C for 2 hours. The mixture was cooled to room temperature, and then the reaction mixture was poured into water. The precipitate was collected by filtration, washed with water, and dried under reduced pressure to give 18.8 g of N-methyl-(3-nitro-5-trifluoromethyl-pyridin-2-yl)-amine.




15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.66-8.64 (m, 2H), 8.43 (brs, 1H), 3.23 (d, 3H)

[0489]

Reference Production Example 2-c

A mixture of N-methyl-(3-nitro-5-trifluoromethyl-pyridin-2-yl)-amine (6.93 g), 5 % of palladium-carbon (0.69 g), and ethanol was stirred under about 1 atm of hydrogen at room temperature for 10 hours. The reaction mixture was filtered through Celite®, and then the filtrate was concentrated under reduced pressure to give 5.02 g of N²-

methyl-5-trifluoromethylpyridin-2,3-diamine (Compound (1F)-
1).



¹H-NMR (CDCl₃) δ: 8.04 (s, 1H), 6.98 (d, 1H), 4.56 (brs, 1H),
5 3.25 (brs, 2H), 3.06 (d, 3H)


[0490]

The compound of Reference Production Example 2-c can
also be produced by the following production method.

[0491]

10 Reference Production Example 2-d

To a mixture of N-methyl-(5-trifluoromethyl-pyridin-2-
yl)-amine (3.48 g) and DMF (20 ml), N-bromosuccinimide
(4.27 g) was added, and stirred at room temperature for 1
hour. Into the reaction mixture, saturated aqueous ammonium
15 chloride solution was poured, and extracted with ethyl
acetate. The organic layer was dried over sodium sulfate,
and concentrated under reduced pressure. The resulting
residue was subjected to silica gel column chromatography
to give 4.8 g of N-methyl-(3-bromo-5-trifluoromethyl-
20 pyridin-2-yl)-amine.



¹H-NMR (CDCl₃) δ: 3.08 (3H, d), 5.40 (1H, brs), 7.78 (1H, d),
8.34 (1H, s).

[0492]


Reference Production Example 2-e

To a pressure-resistant reaction container, N-methyl-(3-bromo-5-trifluoromethyl-pyridin-2-yl)-amine (0.51 g),
5 copper (II) acetylacetonate (0.11 g), acetylacetonate (0.20 g), cesium carbonate (1.30 g), NMP (2 ml), and 28 % of aqueous ammonia (1 ml) were added, and stirred at 120 °C for 7 hours, then at 130 °C for 3 hours. The mixture was allowed to cool to room temperature, and then saturated aqueous
10 ammonium chloride solution was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.28 g of N²-methyl-5-
15 trifluoromethylpyridin-2,3-diamine.

[0493]

Reference Production Example 3-a

A mixture of 2-chloro-5-trifluoromethylpyridine (12 g), 70 % of aqueous ethylamine solution (16 ml), and NMP (7 ml)
20 was stirred with heating at 70 °C for 1 day. Into the reaction mixture cooled to room temperature, water was poured, and then the precipitated solid was collected by filtration. The obtained solid was washed with water, and dried to give 6.3 g of N-ethyl-(5-trifluoromethylpyridin-2-
25 yl)-amine.




$^1\text{H-NMR}$ (CDCl_3) δ : 8.25 (1H,s), 7.51 (1H,d), 6.31 (1H,d), 4.84 (1H,brs), 3.32-3.25 (2H,m), 1.27-1.11 (3H,m)

[0494]

5 Reference Production Example 3-b

A mixture of N-methyl-(5-trifluoromethylpyridin-2-yl)-amine (1.0 g) and sulfuric acid (7 ml) was stirred with heating at 95 °C, and then a mixture of fuming nitric acid (0.45 ml) and sulfuric acid (3 ml) were added thereto, and
10 stirred with heating for 1 hour. The reaction mixture cooling to room temperature was poured into ice, and then saturated aqueous sodium hydrogen carbonate solution was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under
15 reduced pressure. The resulting residue was subjected to silica gel chromatography to give 600 mg of N-ethyl-(3-nitro-5-trifluoromethylpyridin-2-yl)-amine.




[0495]

20 Reference Production Example 3-c

A mixture of iron powder (700 mg), acetic acid (60 μl),

ethanol (5 ml), and water (3 ml) was stirred with heating at 70 °C, and then a mixture of N-ethyl-(3-nitro-5-trifluoromethylpyridin-2-yl)-amine (490 mg) and ethanol (10 ml) was added dropwise. The mixture was stirred with heating at 70 °C for 3 hours. The reaction mixture cooled to room temperature was filtered through Celite®, and the resulting filtrate was concentrated under reduced pressure. The resulting residue was diluted with saturated aqueous sodium hydrogen carbonate solution, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 0.29 g of N²-ethyl-5-trifluoromethylpyridin-2,3-diamine (Compound (1F)-8).




¹H-NMR (CDCl₃) δ: 8.01 (1H,d), 6.97 (1H,d), 3.51 (2H,q), 1.27 (3H,t)

[0496]

Reference Production Example 4-a

The procedure was performed according to the method described in Reference Production Example 3-a using n-propylamine instead of 70 % of aqueous ethylamine solution to give 10 g of N-propyl-(5-trifluoromethylpyridin-2-yl)-

amine.




[0497]

Reference Production Example 4-b

5 A mixture of N-propyl-(5-trifluoromethylpyridin-2-yl)-
amine (3.0 g) and sulfuric acid (20 ml) was stirred with
heating at 95 °C, and then a mixture of fuming nitric acid
(1.3 ml) and sulfuric acid (10 ml) was added dropwise, and
stirred with heating for 1 hour. The reaction mixture was
10 poured into ice, and then saturated aqueous sodium hydrogen
carbonate solution was poured, and extracted with ethyl
acetate. The organic layer was dried over sodium sulfate,
and concentrated under reduced pressure.


 To a mixture of the resulting residue, ethanol (12 ml)
15 and water (8 ml), iron powder (2.5 g) and acetic acid (0.5
ml) were added, heated to 70 °C, and stirred with heating
for 2 hours. The reaction mixture cooled to room
temperature was filtered through Celite®, and then the
resulting filtrate was concentrated under reduced pressure.
20 The resulting residue was subjected to silica gel column
chromatography to give 1.5 g of N²-propyl-5-
trifluoromethylpyridin-2,3-diamine.



[0498]

Reference Production Example 4-c

A mixture of N-propyl-5-trifluoromethylpyridin-2,3-
5 diamine (770 mg), 2-ethylsulfanyl-benzoic acid (700 mg),
WSC (740 mg), and pyridine (20 ml) was stirred with heating
at 95 °C for 7 hours. Into the reaction mixture cooled to
room temperature, saturated aqueous sodium carbonate
solution was poured, and extracted with ethyl acetate. The
10 organic layer was dried over sodium sulfate, and
concentrated under reduced pressure. The resulting residue
was subjected to silica gel column chromatography to give
422 mg of 2-ethylsulfanyl-N-(2-propylamino-5-
trifluoromethylpyridin-3-yl)-benzamide (Compound (1B)-513).




15

[0499]

Reference Production Example 5-a

The procedure was performed according to the method
described in Reference Production Example 3-b using 5-
20 trifluoromethylpyridin-2-ol instead of N-methyl-(5-
trifluoromethylpyridin-2-yl)-amine to give 3.3 g of 3-

nitro-5-trifluoromethylpyridin-2-ol.




$^1\text{H-NMR}$ (DMSO- d_6) δ : 13.54 (1H, brs), 8.67 (1H, d), 8.46 (1H, d)

[0500]

5 Reference Production Example 5-b

To a mixture of 3-nitro-5-trifluoromethylpyridin-2-ol (3.2 g) and DMF (1 drop), phosphorous oxychloride (8.2 ml) was added at room temperature. The mixture was heated to 105 °C, and stirred with heating for 2 days. The reaction
10 mixture was cooled to room temperature, and then the reaction mixture was poured into ice, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure to give 2.5 g of 2-chloro-3-nitro-5-trifluoromethylpyridine.



15


[0501]

Reference Production Example 5-c

To a mixture of 2-chloro-3-nitro-5-trifluoromethylpyridine (1.5 g) and ethanol (15 ml),
20 isopropylamine (1.8 ml) was added under ice-cooling, heated to room temperature, and stirred with heating for 3 hours. Into the reaction mixture, 2 mol/l of aqueous solution of

hydrochloric acid was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure.


To the resulting residue, ethanol (30 ml), water (20
5 ml), ammonium chloride (0.7 g), iron powder (445 mg), and
ammonium chloride (0.7 g) were sequentially added at room
temperature. The mixture was heated to 70 °C, and stirred
with heating for 1 day. Into the reaction mixture cooled to
room temperature, aqueous sodium hydroxide solution was
10 poured, and extracted with ethyl acetate. The organic layer
was dried over sodium sulfate, and concentrated under
reduced pressure to give 970 mg of N²-isopropyl-5-
trifluoromethylpyridin-2,3-diamine (Compound (1F)-15).



15 [0502]

Reference Production Example 6


The procedure was performed according to the method
described in Reference Production Example 5-c using
cyclopropylamine instead of isopropylamine to give 750 mg
20 of N²-cyclopropyl-5-trifluoromethylpyridin-2,3-diamine.



[0503]

Reference Production Example 7

The procedure was performed according to the method described in Reference Production Example 3-c using 5-trifluoromethyl-pyridin-2,3-diamine instead of N²-propyl-5-trifluoromethylpyridin-2,3-diamine to give 4.7 g of N-(2-amino-5-trifluoromethylpyridin-3-yl)-2-ethylsulfanylbenzamide (Compound (1B)-512).




¹H-NMR (DMSO-D₆) δ: 9.75 (1H,s), 8.17 (1H,s), 8.09 (1H,s), 7.64 (1H,d), 7.53-7.45 (2H,m), 7.34-7.28 (1H,m), 6.71 (2H,s), 2.97 (2H,q), 1.22 (3H,t)

[0504]

Reference Production Example 8

A mixture of 5,N²-dimethylpyridin-2,3-diamine (1.13 g), 2-ethylsulfanylbenzoic acid (1.79 g), WSC (1.38 g), and pyridine (20 ml) was stirred with heating under reflux at 120 °C for 2 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 1.58 g of 2-ethylsulfanylphenyl-N-

(5-methyl-2-methylaminopyridin-3-yl)-benzamide (Compound (1B)-514).



¹H-NMR (CDCl₃) δ: 8.18 (1H,brs), 7.97-7.93 (1H,m), 7.85-7.80 (1H,m), 7.54-7.42 (3H,m), 7.39-7.33 (1H,m), 3.03-2.93 (5H,m), 1.32 (3H,t)


[0505]

Reference Production Example 9

To a mixture of 5-iodo-3-nitropyridin-2-ylamine (1.0 g) and DMF (20 ml), 60 % of sodium hydride (in oil, 180 mg) and methyl iodide (240 μl) were sequentially added under ice-cooling. The mixture was heated to room temperature, and stirred for 1 hour. Into the reaction mixture, water was poured, and then the precipitated solid was collected by filtration, and dried.

The obtained solid was dissolved in THF. The solution was added dropwise to a mixture of iron powder (633 mg), acetic acid (1 ml), ethanol (30 ml), and water (20 ml) with heating at 70 °C. The mixture was stirred with heating at 70 °C for 3.5 hours. The reaction mixture cooled to room temperature was filtered through Celite®. The resulting filtrate was concentrated under reduced pressure. The resulting residue was subjected to silica gel column


chromatography to give 980 mg of 5-iodo-N²-methylpyridin-2,3-diamine (Compound (1F)-6).



[0506]

5 Reference Production Example 10-a

To a mixture of 5-trifluoromethylpyridin-2,3-diamine (1.2 g) and pyridine (20 ml), methyl chloroformate (0.63 ml) was added under ice-cooling. The mixture was heated to room temperature, stirred for 8 hours, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 628 mg of methyl (2-amino-5-trifluoromethylpyridin-3-yl)-carbamate.




¹H-NMR (CDCl₃) δ: 8.65 (1H,brs), 8.18-7.99 (2H,m), 5.75 (2H,brs), 3.79 (3H,s)

[0507]

Reference Production Example 10-b

To a mixture of methyl (2-amino-5-trifluoromethylpyridin-3-yl)-carbamate (100 mg) and THF (15 ml), lithium aluminum hydride (26 mg) was added under ice-cooling. The mixture was stirred for 15 minutes under ice-


cooling, heated to 90 °C, and stirred with heating for 1.5 hours. Into the reaction mixture cooled to room temperature, aqueous sodium hydroxide solution was poured, and extracted with tert-butyl methyl ether. The organic layer was dried
5 over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 77 mg of N³-methyl-5-trifluoromethylpyridin-2,3-diamine.



10 [0508]

Reference Production Example 11-a

To a mixture of methyl-(2-nitro-4-trifluoromethylphenyl)-amine (4.40 g) and DMF (20 ml), N-chlorosuccinimide (3.67 g) was added under ice-cooling, and
15 stirred for 3 hours. The mixture was heated to room temperature, and allowed to stand. Water was poured thereinto under ice-cooling, and then the precipitated solid was collected by filtration. The obtained solid was washed with water, and dried to give 3.96 g of (2-chloro-6-nitro-4-trifluoromethylphenyl)-methylaniline.
20




¹H-NMR (CDCl₃) δ: 8.19 (1H,d), 7.68 (1H,d), 7.17 (1H,brs),
3.19 (3H,d)

[0509]

Reference Production Example 11-b

5 To a mixture of (2-chloro-6-nitro-4-trifluoromethylphenyl)-methylamine (3.70 g) and ethanol (60 ml), Raney[®] nickel (slurry in water, 0.5 ml) was added, and stirred under about 1 atm of hydrogen at room temperature for 1.5 hours. The reaction mixture was filtered. The
10 filtrate was concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 2.95 g of 3-chloro-N²-methyl-5-trifluoromethylbenzene-1,2-diamine.




15 ¹H-NMR (CDCl₃) δ: 7.02 (1H,d), 6.82 (1H,d), 4.09 (2H,brs),
3.49 (1H,brs), 2.74 (3H,d)

[0510]

Reference Production Example 12

20 A mixture of 2-amino-4-trifluoromethylphenol (1.00 g), 2-methylsulfanylbobenzoic acid (0.95 g), WSC (1.41 g), and pyridine (10 ml) was stirred with heating at 80 °C for 2.5 hours. Into the reaction mixture cooled to room temperature, 1 mol/l of hydrochloric acid was poured, and extracted with

ethyl acetate. The organic layer was washed with saturated aqueous sodium hydrogen carbonate solution, water, and saturated brine, sequentially, dried over sodium sulfate, and concentrated under reduced pressure. The resulting
5 residue was subjected to silica gel column chromatography to give 0.33 g of N-(2-hydroxy-5-trifluoromethylphenyl)-2-methylsulfanylbenzamide (Compound (1B)-515).




¹H-NMR(DMSO-D₆) δ: 9.50 (1H,brs), 8.24 (1H,s), 7.61 (1H,d),
10 7.50 (1H,t), 7.42 (1H,d), 7.37 (1H,dd), 7.27 (1H,t), 7.06
(1H,d), 2.45 (3H,s)

[0511]

Reference Production Example 13

To a mixture of 3-amino-5-trifluoromethylpyridin-2-ol
15 (0.75 g) and THF (5 ml), 2-ethylsulfanylbenzoic acid
hydrochloride (0.93 g) was added under ice-cooling, heated
to room temperature, and stirred for 2 hours. After the
mixture was allowed to stand overnight, the mixture was
heated to 50 °C, and stirred with heating for 6 hours. To
20 the mixture, sodium hydrogen carbonate (0.35 g) was added,
and stirred with heating for further 2.5 hours. After the
mixture was allowed to stand, water was poured thereinto,
and the precipitated solid was collected by filtration. The

obtained solid was washed with water, and dried to give 1.37 g of 2-ethylsulfanyl-N-(2-hydroxy-5-trifluoromethylpyridin-3-yl)-benzamide (Compound (1B)-506).




5 $^1\text{H-NMR}$ (CDCl_3) δ : 11.68 (1H,brs), 9.35 (1H,brs), 8.87 (1H,d), 7.75-7.71 (1H,m), 7.49-7.43 (3H,m), 7.33-7.29 (1H,m), 2.99 (2H,q), 1.34 (3H,t)
[0512]

Reference Production Example 14

10 To a mixture of 3-chloroanthranilic acid (3.43 g) and 1 mol/l of hydrochloric acid (20 ml), sodium nitrite (1.38 g) was added portionwise at room temperature, and stirred for 10 minutes. The reaction mixture was added dropwise to a mixture of potassium butylxanthogenate (5.65 g) and water
15 (20 ml) at room temperature, and stirred for 1 hour. 3N aqueous sodium hydroxide solution (20 ml) and diethylsulfuric acid (2.9 ml) were added thereto, heated to 100 °C, and stirred under reflux for 1 hour. To the reaction mixture cooled to room temperature, 12 mol/l of
20 hydrochloric acid was added dropwise under ice-cooling until the reaction mixture reached pH 1-2. The precipitated solid was collected by filtration, and dried under reduced pressure.

To the obtained solid, 3N aqueous sodium hydroxide solution (10 ml), ethanol (10 ml), and ethyl iodide (6.86 g) were added, heated to 80 °C, and stirred under reflux for 2 hours. To the reaction mixture cooled to room temperature, 12 mol/l of hydrochloric acid was added dropwise under ice-cooling until the reaction mixture reached pH 1-2, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 2.79 g of 3-chloro-2-ethylsulfanylbenzoic acid.




¹H-NMR (CDCl₃) δ: 8.10-8.05 (1H,m), 7.68 (1H,dd), 7.43 (1H,t), 3.00 (2H,q), 1.26 (3H,t)

[0513]

Reference Production Example 15

To a pressure-resistant reaction container, 3-amino-2-chloro-5-trifluoromethylpyridine (5 g), copper (II) acetylacetonate (1.31 g), acetylacetonate (2.50 g), cesium carbonate (16.25 g), NMP (20 ml), and 40 % of aqueous methylamine solution (15 ml) were added, and stirred at 100 °C for 3 hours, at 120 °C for 3 hours, then at 140 °C for 5 hours. After the mixture was allowed to cool to room

temperature, saturated aqueous ammonium chloride solution was poured thereinto, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 2.09 g of N²-methyl-6-trifluoromethylpyridin-2,3-diamine.




¹H-NMR (CDCl₃) δ: 3.04 (3H,s), 3.41 (2H,brs), 4.22 (1H,brs), 6.84 (1H,d), 6.91 (1H,d).

[0514]

Reference Production Example 16-a

To a mixture of 3-amino-2-chloro-6-trifluoromethylpyridine (3.93 g) and chloroform (30 ml), pivaloyl chloride (2.65 g) and triethylamine (4.04 g) were added under ice-cooling, and stirred at room temperature for 2 hours, then at 60 °C for 2 hours. Into the reaction mixture cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 6.59 g of N-(2-chloro-6-trifluoromethylpyridin-3-yl)-2,2-dimethylpropionamide.



$^1\text{H-NMR}$ (CDCl_3) δ : 1.36 (9H,s), 7.65 (1H,d), 8.17 (1H,brs), 8.97 (1H,d).

[0515]

5 Reference Production Example 16-b

To a mixture of N-(2-chloro-6-trifluoromethylpyridin-3-yl)-2,2-dimethylpropionamide (6.59 g) and THF (30 ml), sodium hydride (60 % in oily, 1 g) was added under ice-cooling, and stirred at room temperature for 30 minutes.

10 The reaction mixture was ice-cooled, and then methyl iodide (10 g) was added, and stirred at room temperature for 4 hours. Into the reaction mixture, water was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced

15 pressure. To the resulting residue, acetonitrile (40 ml), concentrated hydrochloric acid (40 ml), and isopropanol (40 ml) were added, and stirred at room temperature for 2 hours, at 60 °C for 2 hours, then at 80 °C for 2 hours. Into the reaction mixture cooled to room temperature, saturated

20 aqueous sodium hydrogen carbonate solution was poured, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel

column chromatography to give 2.60 g of (2-chloro-6-trifluoromethylpyridin-3-yl)-methylaniline.




¹H-NMR (CDCl₃) δ: 7.49 (1H,d), 6.89 (1H,d), 4.82 (1H,brs),
5 2.96 (3H,d).

[0516]

Reference Production Example 16-c

To a pressure-resistant reaction container, (2-chloro-6-trifluoromethylpyridin-3-yl)-methylaniline (2.60 g), copper
10 (II) acetylacetonate (0.64 g), acetylacetonate (1.23 g), cesium carbonate (7.99 g), NMP (10 ml), and 28 % of aqueous ammonia (7 ml) were added, and stirred at 140 °C for 8 hours. After the mixture was allowed to cool to room temperature, saturated aqueous ammonium chloride solution
15 was poured thereinto, and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 2.09 g of N³-methyl-6-trifluoromethylpyridin-2,3-diamine.




¹H-NMR (CDCl₃) δ: 2.89 (3H,d), 3.56 (1H,brs), 4.29 (2H,brs),
20 6.77 (1H,d), 7.13 (1H,d).

[0517]

Reference Production Example 17

A mixture of 3-amino-5-(trifluoromethyl)pyridin-2-thiol (0.20 g), 2-ethylsulfanylbenzaldehyde (0.20 g), sodium hydrogen sulfite (0.13 g), copper (II) chloride (0.41 g), and DMF (2.5 ml) was stirred at 120 °C for 8 hours. After the reaction mixture was allowed to cool, water was added thereto, and ethyl acetate was poured. The mixture was filtered, and the resulting filtrate was extracted with ethyl acetate. The organic layer was washed with water, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 0.18 g of 2-[2-fluoro-4-(trifluoromethyl)phenyl]-6-(trifluoromethyl)thiazolo[5,4-b]pyridine ((1E)-F57).




¹H-NMR (CDCl₃) δ: 7.58 (1H,d), 7.64 (1H,d), 8.57-8.63 (2H,m), 8.92 (1H,s).

[0518]

20 Reference Production Example 18

A mixture of N²-methyl-5-trifluoromethylpyridin-2,3-diamine (5.37 g), 2-fluoro-4-trifluorobenzaldehyde (6.92 g), sodium hydrogen sulfite (3.75 g), and DMF (30 ml) was stirred at 150 °C for 3 hours. Into the reaction mixture

cooled to room temperature, water was poured, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 7.83 g of 2-(2-fluoro-4-trifluoromethylphenyl)-3-methyl-6-trifluoromethyl-3H-imidazo[4,5-b]pyridine ((1E)-F6).



$^1\text{H-NMR}$ (CDCl_3) δ : 8.75 (1H,d), 8.35 (1H,d), 7.91 (1H,t), 7.67 (1H,d), 7.59 (1H,d), 3.92 (3H,d).


[0519]

Reference Production Example 19-a

To a mixture of 2-chloro-5-iodopyridine (7.2 g), NMP (75 mL), and xylene (75 mL), pentafluorosodium propionate (28 g) and copper iodide (14 g) were added at room temperature, heated to 150 °C, and stirred with heating for 5.5 hours. After the mixture was cooled to 80 °C, to the mixture, 40 % of aqueous methylamine solution (180 mL) was added in parts 4 every 2 hours, and stirred with heating for 8.5 hours. The mixture was ice-cooled to 0 °C, and then 28 % of aqueous ammonia and saturated aqueous sodium hydrogen carbonate solution were poured thereinto, and extracted with MTBE. The combined organic layer was dried

over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 4.6 g of methyl-(5-pentafluoroethyl-pyridine-2-yl)-amine.

5 methyl-(5-pentafluoroethyl-pyridine-2-yl)-amine




¹H-NMR (CDCl₃) δ: 8.28 (1H,d), 7.56 (1H,dd), 6.41 (1H,d), 5.74 (1H,brs), 2.95 (3H,d).

[0520]

10 Reference Production Example 19-b

To a mixture of methyl-(5-pentafluoroethyl-pyridine-2-yl)-amine (4.6 g) and acetonitrile (70 mL), N-bromosuccinimide (5.0 g) was added under ice-cooling, and stirred at room temperature for 8 hours. Into a mixture, saturated aqueous sodium thiosulfate solution and aqueous sodium hydrogen carbonate solution were poured, and extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 6.3 g of (3-bromo-5-pentafluoroethyl-pyridine-2-yl)-methyl-amine.

(3-bromo-5-pentafluoroethyl-pyridine-2-yl)-methyl-amine




¹H-NMR (CDCl₃) δ: 8.29 (1H,d), 7.74 (1H,d), 5.45 (1H,brs), 3.09 (3H,d).

[0521]

Reference Production Example 19-c

5 To an autoclave reactor, (3-bromo-5-pentafluoroethyl-pyridine-2-yl)-methyl-amine (2.0 g), acetylacetone copper(II) (86 mg), acetylacetone (263 mg), cesium carbonate (3.2 g), and NMP (13 mL) were added, and then 28 % of aqueous ammonia (5 mL) was added under ice-cooling. 10 After the reactor was sealed, the mixture was heated to 110 °C, and stirred with heating for 15 hours. After the mixture was cooled to room temperature, the reaction mixture was diluted with water, and extracted with MTBE. The combined organic layer was dried over sodium sulfate, 15 and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 623 mg of N²-methyl-5-pentafluoroethyl-2,3-diamine (Compound (1F)-2).

N²-methyl-5-pentafluoroethyl-2,3-diamine




¹H-NMR (CDCl₃) δ: 7.99 (1H,d), 6.94 (1H,d), 4.64 (1H,brs), 3.30 (2H,brs), 3.06 (3H,d).

[0522]

Reference Production Example 20-a

A mixture of 2-chloro-5-iodopyridine (23.9 g), thiobenzoic acid (14 ml), copper iodide (1.90 g), 1,10-phenanthroline (3.60 g), diisopropylethylamine (35 ml), and toluene (200 ml) was stirred at 110 °C for 4 hours. After the reaction mixture was cooled to room temperature, water was poured thereinto. The insoluble material was filtered through Celite®, and then the filtered material was washed with ethyl acetate, and the filtrate was extracted with ethyl acetate. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 21.2 g of S-(6-chloropyridine-3-yl) thiobenzoate.

S-(6-chloropyridine-3-yl) thiobenzoate




¹H-NMR (CDCl₃) δ: 8.43-8.42 (1H,m), 8.01-7.98 (2H,m), 7.79-7.76 (1H,m), 7.66-7.61 (1H,m), 7.52-7.47 (2H,m), 7.44-7.41 (1H,m).

[0523]

Reference Production Example 20-b

A mixture of S-(6-chloropyridine-3-yl) thiobenzoate (21.2 g), potassium carbonate (17.6 g), and methanol (170 ml) was stirred at room temperature for 2 hours. The reaction mixture was filtered, and then the filtered

material was washed with methanol, and filtrate was concentrated under reduced pressure. To the resulting crude, 1N aqueous sodium hydroxide solution (170 ml) was added, and then an aqueous solution of potassium ferricyanide
5 (56.0 g) was added dropwise, and stirred at room temperature for 2 hours. The reaction mixture was extracted with methyl-tert-butyl ether. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The residue was subjected to silica gel
10 column chromatography to give 11.5 g of a compound of the following formula:




¹H-NMR (CDCl₃) δ: 8.41 (2H,d), 7.74 (2H,dd), 7.29 (2H,d).

[0524]

15 Reference Production Example 20-c


A mixture of a compound (11.5 g) of the following formula:



DMF (133 ml) was cooled to -50 °C, and bubbled with
20 excessive amounts of CF₃I gas to dissolve the compound in DMF. To the mixture, tetrakis(dimethylamino)ethylenediamine (37.0 ml) was added dropwise at a rate that the internal temperature did not exceed -40 °C. Then, the mixture was

heated to -10 °C over 1 hour, and stirred at -10 °C for 2 hour. Into the reaction mixture, water was poured, heated to room temperature, and extracted with ethyl acetate. The combined organic layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 7.25 g of 2-chloro-5-trifluoromethylsulfanylpuridine.

2-chloro-5-trifluoromethylsulfanylpuridine



¹H-NMR (CDCl₃) δ: 8.62 (1H,d), 7.93 (1H,dd), 7.43 (1H,d).


[0525]

Reference Production Example 20-d

To a mixture of 2-chloro-5-trifluoromethylsulfanylpuridine (1.71 g) and NMP (16 ml), 40 % of aqueous methylamine solution (1.86 g) was added dropwise, and then the mixture was heated to 60 °C, and stirred with heating for 2 hours. After the mixture was cooled to room temperature, potassium carbonate (1.66 g) was added thereto, and 40 % of aqueous methylamine solution (1.86 g) was added dropwise. The mixture was heated to 60 °C, and stirred with heating for further 2 hours. After the mixture was cooled to room temperature, water was poured, and extracted with ethyl acetate. The combined organic

layer was dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 1.52 g of methyl-(5-trifluoromethylsulfanylpuridine-2-yl)-amine.

5 methyl-(5-trifluoromethylsulfanylpuridine-2-yl)-amine




¹H-NMR (CDCl₃) δ: 8.28 (1H,d), 7.63 (1H,dd), 6.41-6.38 (1H,m), 4.90 (1H,brs), 2.96 (3H,d,).

[0526]

10 Reference Production Example 20-e

To a mixture of methyl-(5-trifluoromethylsulfanylpuridine-2-yl)-amine (1.52 g) and chloroform (24 mL), N-bromosuccinimide (1.43 g) was added, and stirred at room temperature for 1 hour. Into the
15 reaction mixture, water was poured, and extracted with chloroform. The combined organic layer was dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 1.96 g of (3-bromo-5-
20 trifluoromethylsulfanylpuridine-2-yl)-methylamine.
(3-bromo-5-trifluoromethylsulfanylpuridine-2-yl)-methylamine




¹H-NMR (CDCl₃) δ: 8.28 (1H,d), 7.83 (1H,d), 5.40 (1H,brs), 3.07 (3H,d).

[0527]

Reference Production Example 20-f

5 To a pressure-resistant reaction container, (3-bromo-5-trifluoromethylsulfanylp^{yl}pyridine-2-yl)-methylamine (1.96 g), acetylaceton^e copper (II) (89 mg), acetylaceton^e (0.27 g), cesium carbonat^e (2.34 g), and NMP (7 ml), 28 % of aqueous ammoni^a (5 ml) was added, and stirred at 110 °C for
10 8.5 hours. After the mixture was cooled to room temperature, into the reaction mixture, water was poured. The insoluble material was filtered through Celite[®], and then the filtered material was washed with ethyl acetate, and filtrate was extracted with ethyl acetate. The combined
15 organic layer was dried over sodium sulfat^e, and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography to give 1.18 g of N²-methyl-5-trifluoromethylsulfanylp^{yl}pyridine-2,3-diamine (Compound (1F)-3).

20 N²-methyl-5-trifluoromethylsulfanylp^{yl}pyridine-2,3-diamine




¹H-NMR (CDCl₃) δ: 8.01-7.99 (1H,m), 7.07-7.05 (1H,m), 4.60 (1H,brs), 3.22 (2H,brs), 3.05 (3H,d).

[0528]

Reference Production Example 21-a

To a mixture of 2-ethylsulfanylbenzoic acid (9.11 g), chloroform (100 ml), and oxalyl chloride (9.72 g), DMF (2 drops) was added, and stirred at room temperature for 6
5 hours. The reaction mixture was concentrated under reduced pressure to give 10.03 g of 2-ethylsulfanylbenzoic acid chloride (Compound (1H)-241).




¹H-NMR (CDCl₃) δ: 8.31 (1H,dd), 7.57-7.50 (1H,m), 7.42-7.32
10 (1H,m), 7.28-7.21 (1H,m), 3.00 (2H,q), 1.41 (3H,t).
[0529]

Reference Production Example 21-b

To a mixture of N²-methyl-5-trifluoromethylpyridine-2,3-diamine (3.82 g) and THF (40 ml), 2-
15 ethylsulfanylbenzoic acid chloride (4.42 g) was added, stirred at 50 °C for 2 hours, then stirred under reflux for 2 hours. After the mixture was allowed to cool to room temperature, sodium hydrogen carbonate (1.85 g) was added thereto, and stirred at 50 °C for 2 hours. After the
20 mixture was allowed to cool to room temperature, water was poured thereinto, and then the precipitated solid was collected by the filtration. The resulting solid was washed with water and hexane, and dried to give 7.15 g of N-(2-

methylamino-5-trifluoromethylpyridine-3-yl)-2-ethylsulfanyl-benzamide (Compound ((1B)-19)).




¹H-NMR (CDCl₃) δ: 8.36 (1H,s), 8.26 (1H,s), 7.85 (1H,d),
 5 7.76 (1H,s), 7.53 (1H,d), 7.47 (1H,t), 7.39 (1H,t), 5.55
 (1H,brs), 3.08 (3H,d), 3.01 (2H,q), 1.32 (3H,t).

[0530]

The above intermediate, Compounds (1B)-(1E) are shown
 in the following tables.

10 [0531]


A compound of the formula (1B):



wherein, A^{2b} represents =CH-, R^{2b}, R^{4b}, R^{5b}, R^{7b} represents a
 hydrogen atom, and R^{1b}, R^{3b}, R^{6b}, A^{1b}, A^{3b}, and n represent a
 15 combination of R^{1x}, R^{3x}, R^{6x}, A^{1x}, A^{3x}, and n, respectively,
 shown in Tables 38-58 (Compounds ((1B)-1)-((1B)-504)).

[0532]

A compound of the formula (1C):



wherein, X^c represents a fluorine atom, A^{2c} represents $=CH-$, R^{2c} , R^{4c} , R^{5c} , R^{7c} represents a hydrogen atom, and R^{3c} , R^{6c} , A^{1c} , and A^{3c} represent a combination of R^{3x} , R^{6x} , A^{1x} , and A^{3x} , respectively, shown in Tables 38-52 (Compound (1C)-F1 to Compound (1C)-F9, Compound (1C)-F37 to Compound (1C)-F45, Compound (1C)-F55 to Compound (1C)-F63, Compound (1C)-F73 to Compound (1C)-F81, Compound (1C)-F109 to Compound (1C)-F117, Compound (1C)-F127 to Compound (1C)-F135, Compound (1C)-F145 to Compound (1C)-F153, Compound (1C)-F181 to Compound (1C)-F189, Compound (1C)-F199 to Compound (1C)-F207, Compound (1C)-F217 to Compound (1C)-F225, Compound (1C)-F253 to Compound (1C)-F261, Compound (1C)-F271 to Compound (1C)-F279, Compound (1C)-F289 to Compound (1C)-F297, Compound (1C)-F325 to Compound (1C)-F333, and Compound (1C)-F343 to Compound (1C)-F351).


[0533]

A compound of the formula (1C) wherein X^c is a chlorine atom, A^{2c} is $=CH-$, R^{2c} , R^{4c} , R^{5c} , R^{7c} is a hydrogen atom, R^{3c} , R^{6c} , A^{1c} and A^{3c} , represent a combination of R^{3x} , R^{6x} , A^{1x} and A^{3x} , respectively, shown in Tables 38-52 (Compound (1C)-C1 to Compound (1C)-C9, Compound (1C)-C37 to

Compound (1C)-C45, Compound (1C)-C55 to Compound (1C)-C63,
 Compound (1C)-C73 to Compound (1C)-C81, Compound (1C)-C109
 to Compound (1C)-C117, Compound (1C)-C127 to Compound (1C)-
 C135, Compound (1C)-C145 to Compound (1C)-C153, Compound
 5 (1C)-C181 to Compound (1C)-C189, Compound (1C)-C199 to
 Compound (1C)-C207, Compound (1C)-C217 to Compound (1C)-
 C225, Compound (1C)-C253 to Compound (1C)-C261, Compound
 (1C)-C271 to Compound (1C)-C279, Compound (1C)-C289 to
 Compound (1C)-C297, Compound (1C)-C325 to Compound (1C)-
 10 C333, and Compound (1C)-C343 to Compound (1C)-C351).

[0534]

A compound of the formula (1D):




wherein, A^{2d} represents $=CH-$, R^{2d} , R^{4d} , R^{5d} , R^{7d} represents a
 15 hydrogen atom, and R^{3d} , R^{6d} , A^{1d} and A^{3d} , represent a
 combination of R^{3x} , R^{6x} , A^{1x} and A^{3x} , respectively, shown in
 Tables 38-58 (Compound (1D)-1 to Compound (1D)-9, Compound
 (1D)-37 to Compound (1D)-45, Compound (1D)-55 to Compound
 (1D)-63, Compound (1D)-73 to Compound (1D)-81, Compound
 20 (1D)-109 to Compound (1D)-117, Compound (1D)-127 to
 Compound (1D)-135, Compound (1D)-145 to Compound (1D)-153,
 Compound (1D)-181 to Compound (1D)-189, Compound (1D)-199
 to Compound (1D)-207, Compound (1D)-217 to Compound (1D)-

225, Compound (1D)-253 to Compound (1D)-261, Compound (1D)-
 271 to Compound (1D)-279, Compound (1D)-289 to Compound
 (1D)-297, Compound (1D)-325 to Compound (1D)-333, Compound
 (1D)-343 to Compound (1D)-351, Compound (1D)-361 to
 5 Compound (1D)-369, Compound (1D)-397 to Compound (1D)-405,
 Compound (1D)-415 to Compound (1D)-423, Compound (1D)-433
 to Compound (1D)-441, Compound (1D)-469 to Compound (1D)-
 477, and Compound (1D)-487 to Compound (1D)-495).

[0535]

10 A compound of the formula (1E):



wherein, X^e represents a fluorine atom, A^{2e} represents $=CH-$,
 R^{2e} , R^{4e} , R^{5e} , R^{7e} represents a hydrogen atom, R^{3e} , R^{6e} , A^{1e} ,
 and A^{3e} represent a combination of R^{3x} , R^{6x} , A^{1x} , and A^{3x} ,
 15 respectively, shown in Tables 38-52 (Compound (1E)-F1 to
 Compound (1E)-F9, Compound (1E)-F37 to Compound (1E)-F45,
 Compound (1E)-F55 to Compound (1E)-F63, Compound (1E)-F73
 to Compound (1E)-F81, Compound (1E)-F109 to Compound (1E)-
 F117, Compound (1E)-F127 to Compound (1E)-F135, Compound
 20 (1E)-F145 to Compound (1E)-F153, Compound (1E)-F181 to
 Compound (1E)-F189, Compound (1E)-F199 to Compound (1E)-
 F207, Compound (1E)-F217 to Compound (1E)-F225, Compound
 (1E)-F253 to Compound (1E)-F261, Compound (1E)-F271 to

Compound (1E)-F279, Compound (1E)-F289 to Compound (1E)-F297, Compound (1E)-F325 to Compound (1E)-F333, and Compound (1E)-F343 to Compound (1E)-F351).

[0536]

5 A compound of the formula (1E) wherein, X^e represents a chlorine atom, A^{2e} represents $=CH-$, R^{2e} , R^{4e} , R^{5e} , R^{7e} represents a hydrogen atom, R^{3e} , R^{6e} , A^{1e} , and A^{3e} represent a combination of R^{3x} , R^{6x} , A^{1x} , and A^{3x} , respectively, shown in Tables 38-52 (Compound (1E)-C1 to Compound (1E)-C9,
10 Compound (1E)-C37 to Compound (1E)-C45, Compound (1E)-C55 to Compound (1E)-C63, Compound (1E)-C73 to Compound (1E)-C81, Compound (1E)-C109 to Compound (1E)-C117, Compound (1E)-C127 to Compound (1E)-C135, Compound (1E)-C145 to Compound (1E)-C153, Compound (1E)-C181 to Compound (1E)-
15 C189, Compound (1E)-C199 to Compound (1E)-C207, Compound (1E)-C217 to Compound (1E)-C225, Compound (1E)-C253 to Compound (1E)-C261, Compound (1E)-C271 to Compound (1E)-C279, Compound (1E)-C289 to Compound (1E)-C297, Compound (1E)-C325 to Compound (1E)-C333, and Compound (1E)-C343 to
20 Compound (1E)-C351).

[0537]

Table 38

Intermediate	R^{1x}	R^{3x}	R^{6x}	A^{1x}	A^{3x}	n
1	Me	H	CF ₃	NMe	N	0
2	Me	F	CF ₃	NMe	N	0
3	Me	Cl	CF ₃	NMe	N	0

4	Me	Br	CF ₃	NMe	N	0
5	Me	Me	CF ₃	NMe	N	0
6	Me	CF ₃	CF ₃	NMe	N	0
7	Me	OCF ₃	CF ₃	NMe	N	0
8	Me	C ₂ F ₅	CF ₃	NMe	N	0
9	Me	SF ₅	CF ₃	NMe	N	0
10	Me	H	CF ₃	NMe	N	2
11	Me	F	CF ₃	NMe	N	2
12	Me	Cl	CF ₃	NMe	N	2
13	Me	Br	CF ₃	NMe	N	2
14	Me	Me	CF ₃	NMe	N	2
15	Me	CF ₃	CF ₃	NMe	N	2
16	Me	OCF ₃	CF ₃	NMe	N	2
17	Me	C ₂ F ₅	CF ₃	NMe	N	2
18	Me	SF ₅	CF ₃	NMe	N	2
19	Et	H	CF ₃	NMe	N	0
20	Et	F	CF ₃	NMe	N	0
21	Et	Cl	CF ₃	NMe	N	0
22	Et	Br	CF ₃	NMe	N	0
23	Et	Me	CF ₃	NMe	N	0
24	Et	CF ₃	CF ₃	NMe	N	0
25	Et	OCF ₃	CF ₃	NMe	N	0

[0538]

Table 39

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
26	Et	C ₂ F ₅	CF ₃	NMe	N	0
27	Et	SF ₅	CF ₃	NMe	N	0
28	Et	H	CF ₃	NMe	N	2
29	Et	F	CF ₃	NMe	N	2
30	Et	Cl	CF ₃	NMe	N	2
31	Et	Br	CF ₃	NMe	N	2
32	Et	Me	CF ₃	NMe	N	2
33	Et	CF ₃	CF ₃	NMe	N	2
34	Et	OCF ₃	CF ₃	NMe	N	2
35	Et	C ₂ F ₅	CF ₃	NMe	N	2
36	Et	SF ₅	CF ₃	NMe	N	2
37	Et	H	CF ₃	NMe	CH	0
38	Et	F	CF ₃	NMe	CH	0
39	Et	Cl	CF ₃	NMe	CH	0
40	Et	Br	CF ₃	NMe	CH	0

41	Et	Me	CF ₃	NMe	CH	0
42	Et	CF ₃	CF ₃	NMe	CH	0
43	Et	OCF ₃	CF ₃	NMe	CH	0
44	Et	C ₂ F ₅	CF ₃	NMe	CH	0
45	Et	SF ₅	CF ₃	NMe	CH	0
46	Et	H	CF ₃	NMe	CH	2
47	Et	F	CF ₃	NMe	CH	2
48	Et	Cl	CF ₃	NMe	CH	2
49	Et	Br	CF ₃	NMe	CH	2
50	Et	Me	CF ₃	NMe	CH	2

[0539]

Table 40

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
51	Et	CF ₃	CF ₃	NMe	CH	2
52	Et	OCF ₃	CF ₃	NMe	CH	2
53	Et	C ₂ F ₅	CF ₃	NMe	CH	2
54	Et	SF ₅	CF ₃	NMe	CH	2
55	Et	H	CF ₃	S	N	0
56	Et	F	CF ₃	S	N	0
57	Et	Cl	CF ₃	S	N	0
58	Et	Br	CF ₃	S	N	0
59	Et	Me	CF ₃	S	N	0
60	Et	CF ₃	CF ₃	S	N	0
61	Et	OCF ₃	CF ₃	S	N	0
62	Et	C ₂ F ₅	CF ₃	S	N	0
63	Et	SF ₅	CF ₃	S	N	0
64	Et	H	CF ₃	S	N	2
65	Et	F	CF ₃	S	N	2
66	Et	Cl	CF ₃	S	N	2
67	Et	Br	CF ₃	S	N	2
68	Et	Me	CF ₃	S	N	2
69	Et	CF ₃	CF ₃	S	N	2
70	Et	OCF ₃	CF ₃	S	N	2
71	Et	C ₂ F ₅	CF ₃	S	N	2
72	Et	SF ₅	CF ₃	S	N	2
73	Me	H	CF ₂ CF ₃	NMe	N	0
74	Me	F	CF ₂ CF ₃	NMe	N	0
75	Me	Cl	CF ₂ CF ₃	NMe	N	0

[0540]

Table 41

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
76	Me	Br	CF ₂ CF ₃	NMe	N	0
77	Me	Me	CF ₂ CF ₃	NMe	N	0
78	Me	CF ₃	CF ₂ CF ₃	NMe	N	0
79	Me	OCF ₃	CF ₂ CF ₃	NMe	N	0
80	Me	C ₂ F ₅	CF ₂ CF ₃	NMe	N	0
81	Me	SF ₅	CF ₂ CF ₃	NMe	N	0
82	Me	H	CF ₂ CF ₃	NMe	N	2
83	Me	F	CF ₂ CF ₃	NMe	N	2
84	Me	Cl	CF ₂ CF ₃	NMe	N	2
85	Me	Br	CF ₂ CF ₃	NMe	N	2
86	Me	Me	CF ₂ CF ₃	NMe	N	2
87	Me	CF ₃	CF ₂ CF ₃	NMe	N	2
88	Me	OCF ₃	CF ₂ CF ₃	NMe	N	2
89	Me	C ₂ F ₅	CF ₂ CF ₃	NMe	N	2
90	Me	SF ₅	CF ₂ CF ₃	NMe	N	2
91	Et	H	CF ₂ CF ₃	NMe	N	0
92	Et	F	CF ₂ CF ₃	NMe	N	0
93	Et	Cl	CF ₂ CF ₃	NMe	N	0
94	Et	Br	CF ₂ CF ₃	NMe	N	0
95	Et	Me	CF ₂ CF ₃	NMe	N	0
96	Et	CF ₃	CF ₂ CF ₃	NMe	N	0
97	Et	OCF ₃	CF ₂ CF ₃	NMe	N	0
98	Et	C ₂ F ₅	CF ₂ CF ₃	NMe	N	0
99	Et	SF ₅	CF ₂ CF ₃	NMe	N	0
100	Et	H	CF ₂ CF ₃	NMe	N	2

[0541]

Table 42

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
101	Et	F	CF ₂ CF ₃	NMe	N	2
102	Et	Cl	CF ₂ CF ₃	NMe	N	2
103	Et	Br	CF ₂ CF ₃	NMe	N	2
104	Et	Me	CF ₂ CF ₃	NMe	N	2
105	Et	CF ₃	CF ₂ CF ₃	NMe	N	2
106	Et	OCF ₃	CF ₂ CF ₃	NMe	N	2
107	Et	C ₂ F ₅	CF ₂ CF ₃	NMe	N	2
108	Et	SF ₅	CF ₂ CF ₃	NMe	N	2
109	Et	H	CF ₂ CF ₃	NMe	CH	0

110	Et	F	CF ₂ CF ₃	NMe	CH	0
111	Et	Cl	CF ₂ CF ₃	NMe	CH	0
112	Et	Br	CF ₂ CF ₃	NMe	CH	0
113	Et	Me	CF ₂ CF ₃	NMe	CH	0
114	Et	CF ₃	CF ₂ CF ₃	NMe	CH	0
115	Et	OCF ₃	CF ₂ CF ₃	NMe	CH	0
116	Et	C ₂ F ₅	CF ₂ CF ₃	NMe	CH	0
117	Et	SF ₅	CF ₂ CF ₃	NMe	CH	0
118	Et	H	CF ₂ CF ₃	NMe	CH	2
119	Et	F	CF ₂ CF ₃	NMe	CH	2
120	Et	Cl	CF ₂ CF ₃	NMe	CH	2
121	Et	Br	CF ₂ CF ₃	NMe	CH	2
122	Et	Me	CF ₂ CF ₃	NMe	CH	2
123	Et	CF ₃	CF ₂ CF ₃	NMe	CH	2
124	Et	OCF ₃	CF ₂ CF ₃	NMe	CH	2
125	Et	C ₂ F ₅	CF ₂ CF ₃	NMe	CH	2

[0542]

Table 43

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
126	Et	SF ₅	CF ₂ CF ₃	NMe	CH	2
127	Et	H	CF ₂ CF ₃	S	N	0
128	Et	F	CF ₂ CF ₃	S	N	0
129	Et	Cl	CF ₂ CF ₃	S	N	0
130	Et	Br	CF ₂ CF ₃	S	N	0
131	Et	Me	CF ₂ CF ₃	S	N	0
132	Et	CF ₃	CF ₂ CF ₃	S	N	0
133	Et	OCF ₃	CF ₂ CF ₃	S	N	0
134	Et	C ₂ F ₅	CF ₂ CF ₃	S	N	0
135	Et	SF ₅	CF ₂ CF ₃	S	N	0
136	Et	H	CF ₂ CF ₃	S	N	2
137	Et	F	CF ₂ CF ₃	S	N	2
138	Et	Cl	CF ₂ CF ₃	S	N	2
139	Et	Br	CF ₂ CF ₃	S	N	2
140	Et	Me	CF ₂ CF ₃	S	N	2
141	Et	CF ₃	CF ₂ CF ₃	S	N	2
142	Et	OCF ₃	CF ₂ CF ₃	S	N	2
143	Et	C ₂ F ₅	CF ₂ CF ₃	S	N	2
144	Et	SF ₅	CF ₂ CF ₃	S	N	2
145	Me	H	SCF ₃	NMe	N	0
146	Me	F	SCF ₃	NMe	N	0

147	Me	Cl	SCF ₃	NMe	N	0
148	Me	Br	SCF ₃	NMe	N	0
149	Me	Me	SCF ₃	NMe	N	0
150	Me	CF ₃	SCF ₃	NMe	N	0

[0543]

Table 44

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
151	Me	OCF ₃	SCF ₃	NMe	N	0
152	Me	C ₂ F ₅	SCF ₃	NMe	N	0
153	Me	SF ₅	SCF ₃	NMe	N	0
154	Me	H	SCF ₃	NMe	N	2
155	Me	F	SCF ₃	NMe	N	2
156	Me	Cl	SCF ₃	NMe	N	2
157	Me	Br	SCF ₃	NMe	N	2
158	Me	Me	SCF ₃	NMe	N	2
159	Me	CF ₃	SCF ₃	NMe	N	2
160	Me	OCF ₃	SCF ₃	NMe	N	2
161	Me	C ₂ F ₅	SCF ₃	NMe	N	2
162	Me	SF ₅	SCF ₃	NMe	N	2
163	Et	H	SCF ₃	NMe	N	0
164	Et	F	SCF ₃	NMe	N	0
165	Et	Cl	SCF ₃	NMe	N	0
166	Et	Br	SCF ₃	NMe	N	0
167	Et	Me	SCF ₃	NMe	N	0
168	Et	CF ₃	SCF ₃	NMe	N	0
169	Et	OCF ₃	SCF ₃	NMe	N	0
170	Et	C ₂ F ₅	SCF ₃	NMe	N	0
171	Et	SF ₅	SCF ₃	NMe	N	0
172	Et	H	SCF ₃	NMe	N	2
173	Et	F	SCF ₃	NMe	N	2
174	Et	Cl	SCF ₃	NMe	N	2
175	Et	Br	SCF ₃	NMe	N	2

[0544]

Table 45

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
176	Et	Me	SCF ₃	NMe	N	2
177	Et	CF ₃	SCF ₃	NMe	N	2
178	Et	OCF ₃	SCF ₃	NMe	N	2

179	Et	C ₂ F ₅	SCF ₃	NMe	N	2
180	Et	SF ₅	SCF ₃	NMe	N	2
181	Et	H	SCF ₃	NMe	CH	0
182	Et	F	SCF ₃	NMe	CH	0
183	Et	Cl	SCF ₃	NMe	CH	0
184	Et	Br	SCF ₃	NMe	CH	0
185	Et	Me	SCF ₃	NMe	CH	0
186	Et	CF ₃	SCF ₃	NMe	CH	0
187	Et	OCF ₃	SCF ₃	NMe	CH	0
188	Et	C ₂ F ₅	SCF ₃	NMe	CH	0
189	Et	SF ₅	SCF ₃	NMe	CH	0
190	Et	H	SCF ₃	NMe	CH	2
191	Et	F	SCF ₃	NMe	CH	2
192	Et	Cl	SCF ₃	NMe	CH	2
193	Et	Br	SCF ₃	NMe	CH	2
194	Et	Me	SCF ₃	NMe	CH	2
195	Et	CF ₃	SCF ₃	NMe	CH	2
196	Et	OCF ₃	SCF ₃	NMe	CH	2
197	Et	C ₂ F ₅	SCF ₃	NMe	CH	2
198	Et	SF ₅	SCF ₃	NMe	CH	2
199	Et	H	SCF ₃	S	N	0
200	Et	F	SCF ₃	S	N	0

[0545]

Table 46

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
201	Et	Cl	SCF ₃	S	N	0
202	Et	Br	SCF ₃	S	N	0
203	Et	Me	SCF ₃	S	N	0
204	Et	CF ₃	SCF ₃	S	N	0
205	Et	OCF ₃	SCF ₃	S	N	0
206	Et	C ₂ F ₅	SCF ₃	S	N	0
207	Et	SF ₅	SCF ₃	S	N	0
208	Et	H	SCF ₃	S	N	2
209	Et	F	SCF ₃	S	N	2
210	Et	Cl	SCF ₃	S	N	2
211	Et	Br	SCF ₃	S	N	2
212	Et	Me	SCF ₃	S	N	2
213	Et	CF ₃	SCF ₃	S	N	2
214	Et	OCF ₃	SCF ₃	S	N	2
215	Et	C ₂ F ₅	SCF ₃	S	N	2

216	Et	SF ₅	SCF ₃	S	N	2
217	Me	H	SOCF ₃	NMe	N	0
218	Me	F	SOCF ₃	NMe	N	0
219	Me	Cl	SOCF ₃	NMe	N	0
220	Me	Br	SOCF ₃	NMe	N	0
221	Me	Me	SOCF ₃	NMe	N	0
222	Me	CF ₃	SOCF ₃	NMe	N	0
223	Me	OCF ₃	SOCF ₃	NMe	N	0
224	Me	C ₂ F ₅	SOCF ₃	NMe	N	0
225	Me	SF ₅	SOCF ₃	NMe	N	0

[0546]

Table 47

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
226	Me	H	SOCF ₃	NMe	N	2
227	Me	F	SOCF ₃	NMe	N	2
228	Me	Cl	SOCF ₃	NMe	N	2
229	Me	Br	SOCF ₃	NMe	N	2
230	Me	Me	SOCF ₃	NMe	N	2
231	Me	CF ₃	SOCF ₃	NMe	N	2
232	Me	OCF ₃	SOCF ₃	NMe	N	2
233	Me	C ₂ F ₅	SOCF ₃	NMe	N	2
234	Me	SF ₅	SOCF ₃	NMe	N	2
235	Et	H	SOCF ₃	NMe	N	0
236	Et	F	SOCF ₃	NMe	N	0
237	Et	Cl	SOCF ₃	NMe	N	0
238	Et	Br	SOCF ₃	NMe	N	0
239	Et	Me	SOCF ₃	NMe	N	0
240	Et	CF ₃	SOCF ₃	NMe	N	0
241	Et	OCF ₃	SOCF ₃	NMe	N	0
242	Et	C ₂ F ₅	SOCF ₃	NMe	N	0
243	Et	SF ₅	SOCF ₃	NMe	N	0
244	Et	H	SOCF ₃	NMe	N	2
245	Et	F	SOCF ₃	NMe	N	2
246	Et	Cl	SOCF ₃	NMe	N	2
247	Et	Br	SOCF ₃	NMe	N	2
248	Et	Me	SOCF ₃	NMe	N	2
249	Et	CF ₃	SOCF ₃	NMe	N	2
250	Et	OCF ₃	SOCF ₃	NMe	N	2

[0547]

Table 48

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
251	Et	C ₂ F ₅	SOCF ₃	NMe	N	2
252	Et	SF ₅	SOCF ₃	NMe	N	2
253	Et	H	SOCF ₃	NMe	CH	0
254	Et	F	SOCF ₃	NMe	CH	0
255	Et	Cl	SOCF ₃	NMe	CH	0
256	Et	Br	SOCF ₃	NMe	CH	0
257	Et	Me	SOCF ₃	NMe	CH	0
258	Et	CF ₃	SOCF ₃	NMe	CH	0
259	Et	OCF ₃	SOCF ₃	NMe	CH	0
260	Et	C ₂ F ₅	SOCF ₃	NMe	CH	0
261	Et	SF ₅	SOCF ₃	NMe	CH	0
262	Et	H	SOCF ₃	NMe	CH	2
263	Et	F	SOCF ₃	NMe	CH	2
264	Et	Cl	SOCF ₃	NMe	CH	2
265	Et	Br	SOCF ₃	NMe	CH	2
266	Et	Me	SOCF ₃	NMe	CH	2
267	Et	CF ₃	SOCF ₃	NMe	CH	2
268	Et	OCF ₃	SOCF ₃	NMe	CH	2
269	Et	C ₂ F ₅	SOCF ₃	NMe	CH	2
270	Et	SF ₅	SOCF ₃	NMe	CH	2
271	Et	H	SOCF ₃	S	N	0
272	Et	F	SOCF ₃	S	N	0
273	Et	Cl	SOCF ₃	S	N	0
274	Et	Br	SOCF ₃	S	N	0
275	Et	Me	SOCF ₃	S	N	0

[0548]

Table 49

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
276	Et	CF ₃	SOCF ₃	S	N	0
277	Et	OCF ₃	SOCF ₃	S	N	0
278	Et	C ₂ F ₅	SOCF ₃	S	N	0
279	Et	SF ₅	SOCF ₃	S	N	0
280	Et	H	SOCF ₃	S	N	2
281	Et	F	SOCF ₃	S	N	2
282	Et	Cl	SOCF ₃	S	N	2
283	Et	Br	SOCF ₃	S	N	2
284	Et	Me	SOCF ₃	S	N	2

285	Et	CF ₃	SOCF ₃	S	N	2
286	Et	OCF ₃	SOCF ₃	S	N	2
287	Et	C ₂ F ₅	SOCF ₃	S	N	2
288	Et	SF ₅	SOCF ₃	S	N	2
289	Me	H	SO ₂ CF ₃	NMe	N	0
290	Me	F	SO ₂ CF ₃	NMe	N	0
291	Me	Cl	SO ₂ CF ₃	NMe	N	0
292	Me	Br	SO ₂ CF ₃	NMe	N	0
293	Me	Me	SO ₂ CF ₃	NMe	N	0
294	Me	CF ₃	SO ₂ CF ₃	NMe	N	0
295	Me	OCF ₃	SO ₂ CF ₃	NMe	N	0
296	Me	C ₂ F ₅	SO ₂ CF ₃	NMe	N	0
297	Me	SF ₅	SO ₂ CF ₃	NMe	N	0
298	Me	H	SO ₂ CF ₃	NMe	N	2
299	Me	F	SO ₂ CF ₃	NMe	N	2
300	Me	Cl	SO ₂ CF ₃	NMe	N	2

[0549]

Table 50

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
301	Me	Br	SO ₂ CF ₃	NMe	N	2
302	Me	Me	SO ₂ CF ₃	NMe	N	2
303	Me	CF ₃	SO ₂ CF ₃	NMe	N	2
304	Me	OCF ₃	SO ₂ CF ₃	NMe	N	2
305	Me	C ₂ F ₅	SO ₂ CF ₃	NMe	N	2
306	Me	SF ₅	SO ₂ CF ₃	NMe	N	2
307	Et	H	SO ₂ CF ₃	NMe	N	0
308	Et	F	SO ₂ CF ₃	NMe	N	0
309	Et	Cl	SO ₂ CF ₃	NMe	N	0
310	Et	Br	SO ₂ CF ₃	NMe	N	0
311	Et	Me	SO ₂ CF ₃	NMe	N	0
312	Et	CF ₃	SO ₂ CF ₃	NMe	N	0
313	Et	OCF ₃	SO ₂ CF ₃	NMe	N	0
314	Et	C ₂ F ₅	SO ₂ CF ₃	NMe	N	0
315	Et	SF ₅	SO ₂ CF ₃	NMe	N	0
316	Et	H	SO ₂ CF ₃	NMe	N	2
317	Et	F	SO ₂ CF ₃	NMe	N	2
318	Et	Cl	SO ₂ CF ₃	NMe	N	2
319	Et	Br	SO ₂ CF ₃	NMe	N	2
320	Et	Me	SO ₂ CF ₃	NMe	N	2
321	Et	CF ₃	SO ₂ CF ₃	NMe	N	2

322	Et	OCF ₃	SO ₂ CF ₃	NMe	N	2
323	Et	C ₂ F ₅	SO ₂ CF ₃	NMe	N	2
324	Et	SF ₅	SO ₂ CF ₃	NMe	N	2
325	Et	H	SO ₂ CF ₃	NMe	CH	0

[0550]

Table 51

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
326	Et	F	SO ₂ CF ₃	NMe	CH	0
327	Et	Cl	SO ₂ CF ₃	NMe	CH	0
328	Et	Br	SO ₂ CF ₃	NMe	CH	0
329	Et	Me	SO ₂ CF ₃	NMe	CH	0
330	Et	CF ₃	SO ₂ CF ₃	NMe	CH	0
331	Et	OCF ₃	SO ₂ CF ₃	NMe	CH	0
332	Et	C ₂ F ₅	SO ₂ CF ₃	NMe	CH	0
333	Et	SF ₅	SO ₂ CF ₃	NMe	CH	0
334	Et	H	SO ₂ CF ₃	NMe	CH	2
335	Et	F	SO ₂ CF ₃	NMe	CH	2
336	Et	Cl	SO ₂ CF ₃	NMe	CH	2
337	Et	Br	SO ₂ CF ₃	NMe	CH	2
338	Et	Me	SO ₂ CF ₃	NMe	CH	2
339	Et	CF ₃	SO ₂ CF ₃	NMe	CH	2
340	Et	OCF ₃	SO ₂ CF ₃	NMe	CH	2
341	Et	C ₂ F ₅	SO ₂ CF ₃	NMe	CH	2
342	Et	SF ₅	SO ₂ CF ₃	NMe	CH	2
343	Et	H	SO ₂ CF ₃	S	N	0
344	Et	F	SO ₂ CF ₃	S	N	0
345	Et	Cl	SO ₂ CF ₃	S	N	0
346	Et	Br	SO ₂ CF ₃	S	N	0
347	Et	Me	SO ₂ CF ₃	S	N	0
348	Et	CF ₃	SO ₂ CF ₃	S	N	0
349	Et	OCF ₃	SO ₂ CF ₃	S	N	0
350	Et	C ₂ F ₅	SO ₂ CF ₃	S	N	0

[0551]

Table 52

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
351	Et	SF ₅	SO ₂ CF ₃	S	N	0
352	Et	H	SO ₂ CF ₃	S	N	2
353	Et	F	SO ₂ CF ₃	S	N	2

354	Et	Cl	SO ₂ CF ₃	S	N	2
355	Et	Br	SO ₂ CF ₃	S	N	2
356	Et	Me	SO ₂ CF ₃	S	N	2
357	Et	CF ₃	SO ₂ CF ₃	S	N	2
358	Et	OCF ₃	SO ₂ CF ₃	S	N	2
359	Et	C ₂ F ₅	SO ₂ CF ₃	S	N	2
360	Et	SF ₅	SO ₂ CF ₃	S	N	2

[0552]

Table 53

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
361	Me	H	Br	NMe	N	0
362	Me	F	Br	NMe	N	0
363	Me	Cl	Br	NMe	N	0
364	Me	Br	Br	NMe	N	0
365	Me	Me	Br	NMe	N	0
366	Me	CF ₃	Br	NMe	N	0
367	Me	OCF ₃	Br	NMe	N	0
368	Me	C ₂ F ₅	Br	NMe	N	0
369	Me	SF ₅	Br	NMe	N	0
370	Me	H	Br	NMe	N	2
371	Me	F	Br	NMe	N	2
372	Me	Cl	Br	NMe	N	2
373	Me	Br	Br	NMe	N	2
374	Me	Me	Br	NMe	N	2
375	Me	CF ₃	Br	NMe	N	2
376	Me	OCF ₃	Br	NMe	N	2
377	Me	C ₂ F ₅	Br	NMe	N	2
378	Me	SF ₅	Br	NMe	N	2
379	Et	H	Br	NMe	N	0
380	Et	F	Br	NMe	N	0
381	Et	Cl	Br	NMe	N	0
382	Et	Br	Br	NMe	N	0
383	Et	Me	Br	NMe	N	0
384	Et	CF ₃	Br	NMe	N	0
385	Et	OCF ₃	Br	NMe	N	0

[0553]

Table 54

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n

386	Et	C ₂ F ₅	Br	NMe	N	0
387	Et	SF ₅	Br	NMe	N	0
388	Et	H	Br	NMe	N	2
389	Et	F	Br	NMe	N	2
390	Et	Cl	Br	NMe	N	2
391	Et	Br	Br	NMe	N	2
392	Et	Me	Br	NMe	N	2
393	Et	CF ₃	Br	NMe	N	2
394	Et	OCF ₃	Br	NMe	N	2
395	Et	C ₂ F ₅	Br	NMe	N	2
396	Et	SF ₅	Br	NMe	N	2
397	Et	H	Br	NMe	CH	0
398	Et	F	Br	NMe	CH	0
399	Et	Cl	Br	NMe	CH	0
400	Et	Br	Br	NMe	CH	0
401	Et	Me	Br	NMe	CH	0
402	Et	CF ₃	Br	NMe	CH	0
403	Et	OCF ₃	Br	NMe	CH	0
404	Et	C ₂ F ₅	Br	NMe	CH	0
405	Et	SF ₅	Br	NMe	CH	0
406	Et	H	Br	NMe	CH	2
407	Et	F	Br	NMe	CH	2
408	Et	Cl	Br	NMe	CH	2
409	Et	Br	Br	NMe	CH	2
410	Et	Me	Br	NMe	CH	2

[0554]

Table 55

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
411	Et	CF ₃	Br	NMe	CH	2
412	Et	OCF ₃	Br	NMe	CH	2
413	Et	C ₂ F ₅	Br	NMe	CH	2
414	Et	SF ₅	Br	NMe	CH	2
415	Et	H	Br	S	N	0
416	Et	F	Br	S	N	0
417	Et	Cl	Br	S	N	0
418	Et	Br	Br	S	N	0
419	Et	Me	Br	S	N	0
420	Et	CF ₃	Br	S	N	0
421	Et	OCF ₃	Br	S	N	0
422	Et	C ₂ F ₅	Br	S	N	0

423	Et	SF ₅	Br	S	N	0
424	Et	H	Br	S	N	2
425	Et	F	Br	S	N	2
426	Et	Cl	Br	S	N	2
427	Et	Br	Br	S	N	2
428	Et	Me	Br	S	N	2
429	Et	CF ₃	Br	S	N	2
430	Et	OCF ₃	Br	S	N	2
431	Et	C ₂ F ₅	Br	S	N	2
432	Et	SF ₅	Br	S	N	2
433	Me	H	I	NMe	N	0
434	Me	F	I	NMe	N	0
435	Me	Cl	I	NMe	N	0

[0555]

Table 56

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
436	Me	Br	I	NMe	N	0
437	Me	Me	I	NMe	N	0
438	Me	CF ₃	I	NMe	N	0
439	Me	OCF ₃	I	NMe	N	0
440	Me	C ₂ F ₅	I	NMe	N	0
441	Me	SF ₅	I	NMe	N	0
442	Me	H	I	NMe	N	2
443	Me	F	I	NMe	N	2
444	Me	Cl	I	NMe	N	2
445	Me	Br	I	NMe	N	2
446	Me	Me	I	NMe	N	2
447	Me	CF ₃	I	NMe	N	2
448	Me	OCF ₃	I	NMe	N	2
449	Me	C ₂ F ₅	I	NMe	N	2
450	Me	SF ₅	I	NMe	N	2
451	Et	H	I	NMe	N	0
452	Et	F	I	NMe	N	0
453	Et	Cl	I	NMe	N	0
454	Et	Br	I	NMe	N	0
455	Et	Me	I	NMe	N	0
456	Et	CF ₃	I	NMe	N	0
457	Et	OCF ₃	I	NMe	N	0
458	Et	C ₂ F ₅	I	NMe	N	0
459	Et	SF ₅	I	NMe	N	0

460	Et	H	I	NMe	N	2

[0556]

Table 57

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
461	Et	F	I	NMe	N	2
462	Et	Cl	I	NMe	N	2
463	Et	Br	I	NMe	N	2
464	Et	Me	I	NMe	N	2
465	Et	CF ₃	I	NMe	N	2
466	Et	OCF ₃	I	NMe	N	2
467	Et	C ₂ F ₅	I	NMe	N	2
468	Et	SF ₅	I	NMe	N	2
469	Et	H	I	NMe	CH	0
470	Et	F	I	NMe	CH	0
471	Et	Cl	I	NMe	CH	0
472	Et	Br	I	NMe	CH	0
473	Et	Me	I	NMe	CH	0
474	Et	CF ₃	I	NMe	CH	0
475	Et	OCF ₃	I	NMe	CH	0
476	Et	C ₂ F ₅	I	NMe	CH	0
477	Et	SF ₅	I	NMe	CH	0
478	Et	H	I	NMe	CH	2
479	Et	F	I	NMe	CH	2
480	Et	Cl	I	NMe	CH	2
481	Et	Br	I	NMe	CH	2
482	Et	Me	I	NMe	CH	2
483	Et	CF ₃	I	NMe	CH	2
484	Et	OCF ₃	I	NMe	CH	2
485	Et	C ₂ F ₅	I	NMe	CH	2

[0557]

Table 58

Intermediate	R ^{1x}	R ^{3x}	R ^{6x}	A ^{1x}	A ^{3x}	n
486	Et	SF ₅	I	NMe	CH	2
487	Et	H	I	S	N	0
488	Et	F	I	S	N	0
489	Et	Cl	I	S	N	0
490	Et	Br	I	S	N	0
491	Et	Me	I	S	N	0
492	Et	CF ₃	I	S	N	0
493	Et	OCF ₃	I	S	N	0

494	Et	C ₂ F ₅	I	S	N	0
495	Et	SF ₅	I	S	N	0
496	Et	H	I	S	N	2
497	Et	F	I	S	N	2
498	Et	Cl	I	S	N	2
499	Et	Br	I	S	N	2
500	Et	Me	I	S	N	2
501	Et	CF ₃	I	S	N	2
502	Et	OCF ₃	I	S	N	2
503	Et	C ₂ F ₅	I	S	N	2
504	Et	SF ₅	I	S	N	2


[0558]

In the above Tables 38-58, Me represents a methyl group, and Et represents an ethyl group.

[0559]

5 The ¹H-NMR data of Compound (1B) are shown below.

Compound (1B)-505




¹H-NMR (CDCl₃) δ: 8.76 (1H,s), 8.33 (1H,s), 8.04 (1H,s),
 7.88-7.76 (2H,m), 7.78-7.60 (2H,m), 6.30 (1H,brs), 3.30-
 10 3.20 (1H,m), 3.10 (3H,d), 3.10-2.98 (1H,m), 1.26 (3H,t).

Compound (1B)-28


¹H-NMR (CDCl₃) δ: 8.39 (1H,s), 8.08 (1H,d), 7.85-7.55
 (4H,m), 7.26-7.20 (1H,m), 6.29 (1H,brs), 3.51 (2H,q), 3.05
 (3H,d), 1.33 (3H,t).

15 Compound (1B)-379




$^1\text{H-NMR}$ (CDCl_3) δ : 8.28 (1H, brs), 8.14 (1H, d), 7.90-7.75 (2H, m), 7.58-7.35 (3H, m), 5.06 (1H, brs), 3.20-2.96 (5H, m), 1.32 (3H, q).

5 Compound (1B)-507



$^1\text{H-NMR}$ (CDCl_3) δ : 8.39-8.38 (1H, m), 7.80 (1H, brs), 7.70-7.68 (1H, m), 7.67-7.61 (2H, m), 7.44-7.39 (1H, m), 5.80 (1H, brs), 3.10-3.07 (3H, m), 3.00 (2H, q), 1.26 (3H, t).

10 Compound (1B)-508



$^1\text{H-NMR}$ (CDCl_3) δ : 8.41-8.38 (1H, m), 8.29-8.26 (1H, m), 8.16-8.12 (1H, m), 7.93-7.89 (1H, m), 7.86 (1H, brs), 7.76-7.73 (1H, m), 5.39-5.33 (1H, brm), 3.14 (2H, q), 3.08 (3H, d), 1.42 (3H, t).


15

Compound (1B)-24

$^1\text{H-NMR}$ (CDCl_3) δ : 8.39 (1H,brs), 7.96 (1H,brs), 7.92 (1H,d), 7.77-7.71 (2H,m), 7.62 (1H,d), 5.42 (1H,brs), 3.12-3.03 (5H,m), 1.37 (3H,t).

Compound (1B)-509

5




$^1\text{H-NMR}$ (CDCl_3) δ : 8.38 (1H,brs), 7.89 (1H,d), 7.80 (1H,d), 7.57-7.51 (1H,m), 7.50-7.44 (1H,m), 7.43-7.36 (1H,m), 7.05-6.99 (1H,m), 5.22 (1H,brs), 3.06 (3H,s), 3.00 (2H,q), 1.32 (3H,t).

10 Compound (1B)-451

$^1\text{H-NMR}$ (CDCl_3) δ : 8.26 (1H,brs), 7.87 (1H,s), 7.84 (1H,d), 7.52 (1H,d), 7.49-7.42 (1H,m), 7.41-7.35 (1H,m), 7.20-7.15 (1H,m), 5.12 (1H,brs), 3.02-2.97 (5H,m), 1.32 (3H,t).

15 Compound (1B)-511



$^1\text{H-NMR}$ (CDCl_3) δ : 8.30 (1H,s), 8.10 (1H,brs), 7.76-7.76 (2H,m), 7.50 (1H,d), 7.45 (1H,t), 7.36 (1H,t), 5.41 (1H,d), 4.37-4.27 (1H,m), 2.99 (2H,q), 1.30-1.28 (9H,m).


Compound (1B)-37

$^1\text{H-NMR}$ (CDCl_3) δ : 8.18 (1H,brs), 7.84 (1H,d), 7.60-7.30 (5H,m), 6.77 (1H,d), 4.85 (1H,brs), 3.01 (2H,q), 2.93 (3H,d), 1.32 (3H,t).

5 Compound (1B)-33

$^1\text{H-NMR}$ (Acetone- d_6) δ : 9.56 (1H,brs), 8.36-8.35 (1H,m), 8.30-8.23 (3H,m), 7.87-7.85 (1H,m), 6.44 (1H,brs), 3.65 (2H,q), 3.03 (3H,d), 1.31 (3H,t).

Compound (1B)-516




10

$^1\text{H-NMR}$ (CDCl_3) δ : 9.35 (1H,s), 8.92 (1H,s), 8.03 (1H,d), 7.74 (1H,s), 7.62 (1H,d), 7.57 (1H,s), 7.44 (1H,d), 7.14 (1H,d), 3.05 (2H,q), 1.36 (3H,t).

[0560]

15 The $^1\text{H-NMR}$ data of Compound (1C) are shown below.

Compound (1C)-F505



$^1\text{H-NMR}$ (CDCl_3) δ : 8.40 (1H,s), 7.73-7.71 (1H,m), 7.29-7.24 (3H,m), 4.99 (1H,brs), 3.08 (3H,d).

20 Compound (1C)-F3

$^1\text{H-NMR}$ (CDCl_3) δ : 8.40 (1H,s), 8.14 (1H,dd), 7.93 (1H,d),

7.72 (1H,s), 7.36 (1H,d), 7.32-7.27 (1H,m), 4.99 (1H,brs),
3.08 (3H,d).

Compound (1C)-F4

¹H-NMR (CDCl₃) δ: 8.39 (1H,s), 8.06 (1H,dd), 7.93 (1H,d),
5 7.72 (1H,s), 7.51 (1H,d), 7.44 (1H,d), 4.99 (1H,brs), 3.07
(3H,d).

Compound (1C)-F5

¹H-NMR (CDCl₃) δ: 8.38 (1H,s), 8.15-7.94 (2H,m), 7.73 (1H,s),
7.15 (1H,d), 7.03 (1H,d), 5.06 (1H,brs), 3.07 (3H,d), 2.45
10 (3H,s).

Compound (1C)-F6

¹H-NMR (CDCl₃) δ: 8.41 (1H,brs), 8.33 (1H,t), 7.99 (1H,d),
7.75 (1H,d), 7.64 (1H,d), 7.53 (1H,d), 4.97 (1H,brs), 3.08
(3H,d).

15 Compound (1C)-F37

¹H-NMR (CDCl₃) δ: 8.25-8.06 (2H,m), 7.64-7.53 (2H,m), 7.47
(1H,d), 7.34 (1H,t), 7.24-7.15 (1H,m), 6.80 (1H,d), 4.38
(1H,brs), 2.93 (3H,d).

Compound (1C)-C37

20 ¹H-NMR (CDCl₃) δ: 7.85 (1H,d), 7.64 (1H,brs), 7.56 (1H,s),
7.52-7.38 (4H,m), 6.80 (1H,d), 4.49 (1H,brs), 2.93 (3H,s).

Compound (1C)-F42

¹H-NMR (CDCl₃) δ: 8.33 (1H,t), 8.10 (1H,d), 7.66-7.57
(2H,m), 7.55-7.45 (2H,m), 6.82 (1H,d), 4.29 (1H,brs), 2.93
25 (3H,d).

Compound (1C)-F145

$^1\text{H-NMR}$ (CDCl_3) δ : 8.35 (1H,s), 8.19 (1H,t), 8.02 (1H,d), 7.76 (1H,s), 7.64-7.54 (1H,m), 7.36 (1H,t), 7.25-7.20 (1H,m), 5.06 (1H,brs), 3.07 (3H,d).

5 Compound (1C)-C145

$^1\text{H-NMR}$ (CDCl_3) δ : 8.35 (1H,brs), 7.86 (1H,d), 7.77-7.71 (1H,m), 7.63-7.39 (4H,m), 5.19 (1H,brs), 3.08 (3H,d).

Compound (1C)-F150

10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.40-8.28 (2H,m), 7.98 (1H,d), 7.77 (1H,s), 7.64 (1H,d), 7.53 (1H,d), 4.98 (1H,brs), 3.08 (3H,d).

Compound (1C)-F73

15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.34 (1H,brs), 8.24-8.15 (1H,m), 8.05 (1H,d), 7.68 (1H,s), 7.65-7.54 (1H,m), 7.40-7.32 (1H,m), 7.25-7.18 (1H,m), 5.09 (1H,brs), 3.09 (3H,d).

Compound (1C)-C73

$^1\text{H-NMR}$ (CDCl_3) δ : 8.35 (1H,s), 7.87 (1H,d), 7.65 (1H,s), 7.62 (1H,brs), 7.54-7.40 (3H,m), 5.21 (1H,brs), 3.09 (3H,d).

Compound (1C)-F78

20 $^1\text{H-NMR}$ (CDCl_3) δ : 8.40-8.29 (2H,m), 8.01 (1H,d), 7.69 (1H,s), 7.64 (1H,d), 7.53 (1H,d), 5.00 (1H,brs), 3.09 (3H,d).

Compound (1C)-F1


25 $^1\text{H-NMR}$ (CDCl_3) δ : 8.39 (1H,s), 8.19 (1H,t), 8.03 (1H,d), 7.74 (1H,s), 7.66-7.55 (1H,m), 7.36 (1H,t), 7.25-7.19 (1H,m),

5.05 (1H,brs), 3.08 (3H,d).

Compound (1C)-C1

$^1\text{H-NMR}$ (CDCl_3) δ : 8.40 (1H,s), 7.86 (1H,d), 7.72 (1H,s),
7.59 (1H,brs), 7.54-7.40 (3H,m), 5.15 (1H,brs), 3.08 (3H,d).

5 Compound (1C)-F506



$^1\text{H-NMR}$ (DMSO-D_6) δ : 9.59 (1H,d), 8.50 (1H,s), 7.91 (1H,t),
7.70-7.60 (1H,m), 7.45-7.33 (4H,m), 7.07 (1H,d).

[0561]

10 The $^1\text{H-NMR}$ data of Compound (1D) are shown below.


Compound (1D)-1

$^1\text{H-NMR}$ (CDCl_3) δ : 8.75-8.72 (1H,m), 8.35-8.33 (1H,m), 7.55-
7.51 (1H,m), 7.47-7.41 (2H,m), 7.37-7.32 (1H,m), 4.12
(1H,brs), 3.84 (3H,s).

15 [0562]

The $^1\text{H-NMR}$ data of Compound (1E) are shown below.

Compound (1E)-F505



$^1\text{H-NMR}$ (CDCl_3) δ : 8.29 (1H,d), 7.90 (1H,t), 7.81 (1H,dd),
20 7.66 (1H,d), 7.57 (1H,d), 7.48 (1H,d), 3.82 (3H,d).

Compound (1E)-F9

$^1\text{H-NMR}$ (CDCl_3) δ : 8.77-8.75 (1H,m), 8.37-8.35 (1H,m), 7.93-7.87 (1H,m), 7.80 (1H,dd), 7.74 (1H,dd), 3.92 (3H,d).

Compound (1E)-F7

$^1\text{H-NMR}$ (CDCl_3) δ : 8.75-8.73 (1H,m), 8.34-8.32 (1H,m), 7.81
5 (1H,t), 7.29-7.25 (1H,m), 7.22-7.18 (1H,m), 3.91 (3H,d).

Compound (1E)-F42

$^1\text{H-NMR}$ (CDCl_3) δ : 8.13 (1H,s), 7.91 (1H,t), 7.67-7.61 (2H,m), 7.58-7.51 (2H,m), 3.82 (3H,d).


Compound (1E)-C1

10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.73 (1H,dd), 8.33 (1H,dd), 7.61-7.51 (3H,m), 7.51-7.45 (1H,m), 3.81 (3H,s)

Compound (1E)-F1


$^1\text{H-NMR}$ (CDCl_3) δ : 8.72 (1H,d), 8.32 (1H,d), 7.79-7.65 (1H,m), 7.63-7.58 (1H,m), 7.42-7.36 (1H,m), 7.33-7.25
15 (1H,m), 3.89 (3H,d)

Compound (1E)-F506




$^1\text{H-NMR}$ (CDCl_3) δ : 8.76-8.74 (1H,m), 8.37-8.35 (1H,m), 7.63-7.55 (1H,m), 7.15 (2H,t), 3.86 (3H,s).

20 Compound (1E)-F507




$^1\text{H-NMR}$ (CDCl_3) δ : 8.76-8.73 (1H,m), 8.35-8.33 (1H,m), 7.53-7.40 (2H,m), 7.37-7.30 (1H,m), 3.92 (3H,d).

Compound (1E)-F508



5 $^1\text{H-NMR}$ (CDCl_3) δ : 8.73-8.71 (1H,m), 8.33-8.31 (1H,m), 7.33-7.18 (3H,m), 3.99 (3H,s), 3.89 (3H,d).

Compound (1E)-F509




10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.70-8.69 (1H,m), 8.30-8.28 (1H,m), 7.66 (1H,t), 6.91 (1H,dd), 6.81 (1H,dd), 3.91 (3H,s), 3.88 (3H,d).

Compound (1E)-F3


$^1\text{H-NMR}$ (CDCl_3) δ : 8.74-8.72 (1H,m), 8.33-8.31 (1H,m), 7.70 (1H,t), 7.39 (1H,dd), 7.34 (1H,dd), 3.89 (3H,d).

15 Compound (1E)-C505




$^1\text{H-NMR}$ (CDCl_3) δ : 8.76-8.74 (1H,m), 8.38-8.36 (1H,m), 7.58-7.51 (1H,m), 7.43 (1H,d), 7.23 (1H,t), 3.80 (3H,s).

Compound (1E)-F510




$^1\text{H-NMR}$ (CDCl_3) δ : 8.72-8.70 (1H,m), 8.33-8.31 (1H,m), 7.56-7.49 (1H,m), 6.93-6.86 (2H,m), 3.84 (3H,s), 3.77 (3H,s).

Compound (1E)-F511




$^1\text{H-NMR}$ (CDCl_3) δ : 10.78 (1H,brs), 8.76-8.74 (1H,m), 8.33-8.31 (1H,m), 7.46-7.39 (1H,m), 6.98 (1H,d), 6.84-6.78 (1H,m), 3.96 (3H,d).

Compound (1E)-C506




$^1\text{H-NMR}$ (CDCl_3) δ : 8.79-8.77 (1H,m), 8.48 (1H,d), 8.38-8.36 (1H,m), 8.34 (1H,dd), 7.82 (1H,d), 3.84 (3H,s).

Compound (1E)-C507



$^1\text{H-NMR}$ (CDCl_3) δ : 8.73-8.71 (1H,m), 8.33-8.31 (1H,m), 7.48-7.43 (2H,m), 7.32 (1H,dd), 3.81 (3H,s), 3.06 (2H,q), 1.41 (3H,t).

Compound (1E)-F512



$^1\text{H-NMR}$ (CDCl_3) δ : 8.73-8.70 (1H,m), 8.32-8.30 (1H,m), 7.55-7.49 (1H,m), 7.47-7.41 (1H,m), 7.31-7.22 (1H,m), 3.88 (3H,d), 2.40 (3H,d).


Compound (1E)-F5

$^1\text{H-NMR}$ (CDCl_3) δ : 8.71-8.69 (1H,m), 8.31-8.30 (1H,m), 7.61 (1H,t), 7.18 (1H,d), 7.10 (1H,d), 3.88 (3H,d), 2.48 (3H,s).

Compound (1E)-F4


$^1\text{H-NMR}$ (CDCl_3) δ : 8.74-8.72 (1H,m), 8.33-8.32 (1H,m), 7.63 (1H,t), 7.56-7.48 (2H,m), 3.88 (3H,d).

Compound (1E)-F513




$^1\text{H-NMR}$ (CDCl_3) δ : 8.70-8.68 (1H,m), 8.35-8.33 (1H,m), 7.64-7.55 (1H,m), 7.18-7.12 (2H,m), 3.86 (3H,s).

Compound (1E)-F514




$^1\text{H-NMR}$ (CDCl_3) δ : 8.77-8.75 (1H,m), 8.40-8.38 (1H,m), 6.96-6.90 (2H,m), 6.75 (1H,t), 3.86-3.85 (3H,m).

Compound (1E)-F515




$^1\text{H-NMR}$ (CDCl_3) δ : 8.76-8.74 (1H,m), 8.37-8.35 (1H,m), 7.38-7.33 (2H,m), 3.85 (3H,s).

5 Compound (1E)-F516




$^1\text{H-NMR}$ (CDCl_3) δ : 8.74-8.73 (1H,m), 8.35-8.33 (1H,m), 7.81 (1H,t), 7.69-7.65 (2H,m), 7.63-7.60 (1H,m), 7.55-7.49 (3H,m), 7.48-7.43 (1H,m), 3.94 (3H,d).

10 Compound (1E)-F517



$^1\text{H-NMR}$ (CDCl_3) δ : 8.79-8.77 (1H,m), 8.40-8.38 (1H,m), 7.47-7.42 (2H,m), 3.87 (3H,s).


Compound (1E)-F518



15


$^1\text{H-NMR}$ (CDCl_3) δ : 7.90 (1H,t), 7.63 (1H,dd), 7.59 (1H,d), 7.54 (1H,d), 6.84 (1H,d), 4.05 (3H,d), 4.02 (2H,brs).

Compound (1E) -C508



$^1\text{H-NMR}$ (CDCl_3) δ : 8.74-8.72 (1H,m), 8.35-8.33 (1H,m), 7.50-7.47 (1H,m), 7.41-7.34 (2H,m), 3.79 (3H,s), 2.50 (3H,s).

5 Compound (1E) -F519



$^1\text{H-NMR}$ (CDCl_3) δ : 8.76-8.75 (1H,m), 8.36-8.34 (1H,m), 7.99-7.94 (1H,m), 7.91-7.86 (1H,m), 7.54-7.49 (1H,m), 3.92 (3H,d).

10 Compound (1E) -F78

$^1\text{H-NMR}$ (CDCl_3) δ : 8.71-8.69 (1H,m), 8.34-8.33 (1H,m), 7.93-7.88 (1H,m), 7.69-7.66 (1H,m), 7.61-7.57 (1H,m), 3.92 (3H,d).

Compound (1E) -F77

15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.65 (1H,s), 8.29 (1H,s), 7.64-7.58 (1H,m), 7.20-7.17 (1H,m), 7.13-7.08 (1H,m), 3.90-3.87 (3H,m), 2.49 (3H,s).

Compound (1E) -F149

20 $^1\text{H-NMR}$ (CDCl_3) δ : 8.66-8.64 (1H,m), 8.39-8.37 (1H,m), 7.60 (1H,t), 7.20-7.16 (1H,m), 7.12-7.08 (1H,m), 3.86 (3H,d), 2.48 (3H,s).

Compound (1E)-F73

$^1\text{H-NMR}$ (CDCl_3) δ : 8.67 (1H,d), 8.30 (1H,d), 7.74 (1H,td),
7.66-7.57 (1H,m), 7.39 (1H,td), 7.34-7.27 (1H,m), 3.90
(3H,d).

5 Compound (1E)-C73

$^1\text{H-NMR}$ (CDCl_3) δ : 8.68 (1H,d), 8.31 (1H,d), 7.62-7.53
(3H,m), 7.48 (1H,td), 3.81 (3H,s).

Compound (1E)-F79

10 $^1\text{H-NMR}$ (CDCl_3) δ : 8.70-8.68 (1H, m), 8.33-8.30 (1H, m),
7.81 (1H, t), 7.30-7.25 (1H, m), 7.22-7.18 (1H, m), 3.91
(3H, d).

[0563]

The above intermediate, Compounds (1F)-(1G) are shown
in the following tables.

15 [0564]


A compound of the formula (1F):



wherein A^{2f} represents =CH-, R^{7f} represents a hydrogen atom,
and R^{6f}, R^{8f}, and A^{3f} represent a combination of R^{6x}, R^{8x},
20 and A^{3x}, respectively, shown in Tables 59-61 (Compounds
((1F)-1)-((1F)-56)).

[0565]

A compound of the formula (1G):



wherein X^g represents a chlorine atom, A^{2g} represents $=CH-$, R^{7g} represents a hydrogen atom, and R^{6g} , R^{8g} , and A^{3g} represent a combination of R^{6x} , R^{8x} , and A^{3x} , respectively, shown in Tables 59-61 (Compounds ((1G)-C1)-((1G)-C56)).

A compound of the formula (1G) wherein X^g represents a bromine atom, A^{2g} represents $=CH-$, R^{7g} represents a hydrogen atom, and R^{6g} , R^{8g} , and A^{3g} represent a combination of R^{6x} , R^{8x} , and A^{3x} , respectively, shown in Tables 59-61 (Compounds ((1G)-B1)-((1G)-B56)).

[0566]

Table 59

Intermediate	R^{6x}	R^{8x}	A^{3x}
1	CF_3	CH_3	N
2	CF_2CF_3	CH_3	N
3	SCF_3	CH_3	N
4	$SOCF_3$	CH_3	N
5	SO_2CF_3	CH_3	N
6	I	CH_3	N
7	SF_5	CH_3	N
8	CF_3	CH_2CH_3	N
9	CF_2CF_3	CH_2CH_3	N
10	SCF_3	CH_2CH_3	N
11	$SOCF_3$	CH_2CH_3	N
12	SO_2CF_3	CH_2CH_3	N
13	I	CH_2CH_3	N
14	SF_5	CH_2CH_3	N
15	CF_3	$CH(CH_3)_2$	N
16	CF_2CF_3	$CH(CH_3)_2$	N
17	SCF_3	$CH(CH_3)_2$	N
18	$SOCF_3$	$CH(CH_3)_2$	N

19	SO ₂ CF ₃	CH(CH ₃) ₂	N
20	I	CH(CH ₃) ₂	N
21	SF ₅	CH(CH ₃) ₂	N
22	CF ₃	H	N
23	CF ₂ CF ₃	H	N
24	SCF ₃	H	N
25	SOCF ₃	H	N

[0567]

Table 60

Intermediate	R ^{6x}	R ^{8x}	A ^{3x}
26	SO ₂ CF ₃	H	N
27	I	H	N
28	SF ₅	H	N
29	CF ₃	CH ₃	CH
30	CF ₂ CF ₃	CH ₃	CH
31	SCF ₃	CH ₃	CH
32	SOCF ₃	CH ₃	CH
33	SO ₂ CF ₃	CH ₃	CH
34	I	CH ₃	CH
35	SF ₅	CH ₃	CH
36	CF ₃	CH ₂ CH ₃	CH
37	CF ₂ CF ₃	CH ₂ CH ₃	CH
38	SCF ₃	CH ₂ CH ₃	CH
39	SOCF ₃	CH ₂ CH ₃	CH
40	SO ₂ CF ₃	CH ₂ CH ₃	CH
41	I	CH ₂ CH ₃	CH
42	SF ₅	CH ₂ CH ₃	CH
43	CF ₃	CH(CH ₃) ₂	CH
44	CF ₂ CF ₃	CH(CH ₃) ₂	CH
45	SCF ₃	CH(CH ₃) ₂	CH
46	SOCF ₃	CH(CH ₃) ₂	CH
47	SO ₂ CF ₃	CH(CH ₃) ₂	CH
48	I	CH(CH ₃) ₂	CH
49	SF ₅	CH(CH ₃) ₂	CH
50	CF ₃	H	CH

[0568]

Table 61

Intermediate	R ^{6x}	R ^{8x}	A ^{3x}
51	CF ₂ CF ₃	H	CH
52	SCF ₃	H	CH
53	SOCF ₃	H	CH

655


54	SO ₂ CF ₃	H	CH
55	I	H	CH
56	SF ₅	H	CH

[0569]

The above intermediate, Compounds (1H)-(1I) are shown in the following tables.

[0570]

5 A compound of the formula (1H):




wherein, R^{1h}, R^{2h}, R^{3h}, R^{4h}, R^{5h}, R^{13h}, and n represent a combination of R^{1x}, R^{2x}, R^{3x}, R^{4x}, R^{5x}, R^{13x}, and n, respectively, shown in Tables 62-75 (Compounds ((1H)-1)-((1H)-288).

10 ((1H)-288).

[0571]

A compound of the formula (1I):



wherein, R¹ⁱ, R²ⁱ, R³ⁱ, R⁴ⁱ, R⁵ⁱ, and r represent a combination of R^{1x}, R^{2x}, R^{3x}, R^{4x}, R^{5x}, and n, respectively, shown in Tables 63, 64, 66, and 67 (Compounds ((1I)-17)-((1I)-48) and Compounds ((1I)-65)-((1I)-96).

15 ((1I)-48) and Compounds ((1I)-65)-((1I)-96).

[0572]

Table 62

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n
1	Me	H	H	H	H	H	0
2	Me	H	F	H	H	H	0
3	Me	H	Cl	H	H	H	0
4	Me	H	Br	H	H	H	0
5	Me	H	Me	H	H	H	0
6	Me	H	Et	H	H	H	0
7	Me	H	Pr	H	H	H	0
8	Me	H	CF ₃	H	H	H	0
9	Me	H	C ₂ F ₅	H	H	H	0
10	Me	H	SF ₅	H	H	H	0
11	Me	H	OCH ₃	H	H	H	0
12	Me	H	OC ₂ H ₅	H	H	H	0
13	Me	H	OCF ₃	H	H	H	0
14	Me	H	SCF ₃	H	H	H	0
15	Me	H	SOCF ₃	H	H	H	0
16	Me	H	SO ₂ CF ₃	H	H	H	0

[0573]

Table 63

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n
17	Me	H	H	H	H	H	1
18	Me	H	F	H	H	H	1
19	Me	H	Cl	H	H	H	1
20	Me	H	Br	H	H	H	1
21	Me	H	Me	H	H	H	1
22	Me	H	Et	H	H	H	1
23	Me	H	Pr	H	H	H	1
24	Me	H	CF ₃	H	H	H	1
25	Me	H	C ₂ F ₅	H	H	H	1

[0574]

Table 64

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n
26	Me	H	SF ₅	H	H	H	1
27	Me	H	OCH ₃	H	H	H	1
28	Me	H	OC ₂ H ₅	H	H	H	1
29	Me	H	OCF ₃	H	H	H	1
30	Me	H	SCF ₃	H	H	H	1
31	Me	H	SOCF ₃	H	H	H	1

32	Me	H	SO ₂ CF ₃	H	H	H	1
33	Me	H	H	H	H	H	2
34	Me	H	F	H	H	H	2
35	Me	H	Cl	H	H	H	2
36	Me	H	Br	H	H	H	2
37	Me	H	Me	H	H	H	2
38	Me	H	Et	H	H	H	2
39	Me	H	Pr	H	H	H	2
40	Me	H	CF ₃	H	H	H	2
41	Me	H	C ₂ F ₅	H	H	H	2
42	Me	H	SF ₅	H	H	H	2
43	Me	H	OCH ₃	H	H	H	2
44	Me	H	OC ₂ H ₅	H	H	H	2
45	Me	H	OCF ₃	H	H	H	2
46	Me	H	SCF ₃	H	H	H	2
47	Me	H	SOCF ₃	H	H	H	2
48	Me	H	SO ₂ CF ₃	H	H	H	2

[0575]

Table 65

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n
49	Et	H	H	H	H	H	0
50	Et	H	F	H	H	H	0
51	Et	H	Cl	H	H	H	0
52	Et	H	Br	H	H	H	0
53	Et	H	Me	H	H	H	0
54	Et	H	Et	H	H	H	0
55	Et	H	Pr	H	H	H	0
56	Et	H	CF ₃	H	H	H	0
57	Et	H	C ₂ F ₅	H	H	H	0
58	Et	H	SF ₅	H	H	H	0
59	Et	H	OCH ₃	H	H	H	0
60	Et	H	OC ₂ H ₅	H	H	H	0
61	Et	H	OCF ₃	H	H	H	0
62	Et	H	SCF ₃	H	H	H	0
63	Et	H	SOCF ₃	H	H	H	0
64	Et	H	SO ₂ CF ₃	H	H	H	0

[0576]

Table 66

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n
65	Et	H	H	H	H	H	1
66	Et	H	F	H	H	H	1
67	Et	H	Cl	H	H	H	1
68	Et	H	Br	H	H	H	1
69	Et	H	Me	H	H	H	1
70	Et	H	Et	H	H	H	1
71	Et	H	Pr	H	H	H	1
72	Et	H	CF ₃	H	H	H	1
73	Et	H	C ₂ F ₅	H	H	H	1
74	Et	H	SF ₅	H	H	H	1
75	Et	H	OCH ₃	H	H	H	1
76	Et	H	OC ₂ H ₅	H	H	H	1
77	Et	H	OCF ₃	H	H	H	1
78	Et	H	SCF ₃	H	H	H	1
79	Et	H	SOCF ₃	H	H	H	1
80	Et	H	SO ₂ CF ₃	H	H	H	1
81	Et	H	H	H	H	H	2
82	Et	H	F	H	H	H	2
83	Et	H	Cl	H	H	H	2
84	Et	H	Br	H	H	H	2
85	Et	H	Me	H	H	H	2
86	Et	H	Et	H	H	H	2
87	Et	H	Pr	H	H	H	2
88	Et	H	CF ₃	H	H	H	2
89	Et	H	C ₂ F ₅	H	H	H	2

[0577]

Table 67

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n
90	Et	H	SF ₅	H	H	H	2
91	Et	H	OCH ₃	H	H	H	2
92	Et	H	OC ₂ H ₅	H	H	H	2
93	Et	H	OCF ₃	H	H	H	2
94	Et	H	SCF ₃	H	H	H	2
95	Et	H	SOCF ₃	H	H	H	2
96	Et	H	SO ₂ CF ₃	H	H	H	2

[0578]

Table 68

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n
97	Me	H	H	H	H	OH	0
98	Me	H	F	H	H	OH	0
99	Me	H	Cl	H	H	OH	0
100	Me	H	Br	H	H	OH	0
101	Me	H	Me	H	H	OH	0
102	Me	H	Et	H	H	OH	0
103	Me	H	Pr	H	H	OH	0
104	Me	H	CF ₃	H	H	OH	0
105	Me	H	C ₂ F ₅	H	H	OH	0
106	Me	H	SF ₅	H	H	OH	0
107	Me	H	OCH ₃	H	H	OH	0
108	Me	H	OC ₂ H ₅	H	H	OH	0
109	Me	H	OCF ₃	H	H	OH	0
110	Me	H	SCF ₃	H	H	OH	0
111	Me	H	SOCF ₃	H	H	OH	0
112	Me	H	SO ₂ CF ₃	H	H	OH	0
113	Me	H	H	H	H	OH	1
114	Me	H	F	H	H	OH	1
115	Me	H	Cl	H	H	OH	1
116	Me	H	Br	H	H	OH	1
117	Me	H	Me	H	H	OH	1
118	Me	H	Et	H	H	OH	1
119	Me	H	Pr	H	H	OH	1
120	Me	H	CF ₃	H	H	OH	1

[0579]

Table 69

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n
121	Me	H	C ₂ F ₅	H	H	OH	1
122	Me	H	SF ₅	H	H	OH	1
123	Me	H	OCH ₃	H	H	OH	1
124	Me	H	OC ₂ H ₅	H	H	OH	1
125	Me	H	OCF ₃	H	H	OH	1
126	Me	H	SCF ₃	H	H	OH	1
127	Me	H	SOCF ₃	H	H	OH	1
128	Me	H	SO ₂ CF ₃	H	H	OH	1
129	Me	H	H	H	H	OH	2
130	Me	H	F	H	H	OH	2
131	Me	H	Cl	H	H	OH	2
132	Me	H	Br	H	H	OH	2

133	Me	H	Me	H	H	OH	2
134	Me	H	Et	H	H	OH	2
135	Me	H	Pr	H	H	OH	2
136	Me	H	CF ₃	H	H	OH	2
137	Me	H	C ₂ F ₅	H	H	OH	2
138	Me	H	SF ₅	H	H	OH	2
139	Me	H	OCH ₃	H	H	OH	2
140	Me	H	OC ₂ H ₅	H	H	OH	2
141	Me	H	OCF ₃	H	H	OH	2
142	Me	H	SCF ₃	H	H	OH	2
143	Me	H	SOCF ₃	H	H	OH	2
144	Me	H	SO ₂ CF ₃	H	H	OH	2
145	Et	H	H	H	H	OH	0

[0580]

Table 70

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n
146	Et	H	F	H	H	OH	0
147	Et	H	Cl	H	H	OH	0
148	Et	H	Br	H	H	OH	0
149	Et	H	Me	H	H	OH	0
150	Et	H	Et	H	H	OH	0
151	Et	H	Pr	H	H	OH	0
152	Et	H	CF ₃	H	H	OH	0
153	Et	H	C ₂ F ₅	H	H	OH	0
154	Et	H	SF ₅	H	H	OH	0
155	Et	H	OCH ₃	H	H	OH	0
156	Et	H	OC ₂ H ₅	H	H	OH	0
157	Et	H	OCF ₃	H	H	OH	0
158	Et	H	SCF ₃	H	H	OH	0
159	Et	H	SOCF ₃	H	H	OH	0
160	Et	H	SO ₂ CF ₃	H	H	OH	0
161	Et	H	H	H	H	OH	1
162	Et	H	F	H	H	OH	1
163	Et	H	Cl	H	H	OH	1
164	Et	H	Br	H	H	OH	1
165	Et	H	Me	H	H	OH	1
166	Et	H	Et	H	H	OH	1
167	Et	H	Pr	H	H	OH	1
168	Et	H	CF ₃	H	H	OH	1
169	Et	H	C ₂ F ₅	H	H	OH	1

170	Et	H	SF ₅	H	H	OH	1

[0581]

Table 71

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n
171	Et	H	OCH ₃	H	H	OH	1
172	Et	H	OC ₂ H ₅	H	H	OH	1
173	Et	H	OCF ₃	H	H	OH	1
174	Et	H	SCF ₃	H	H	OH	1
175	Et	H	SOCF ₃	H	H	OH	1
176	Et	H	SO ₂ CF ₃	H	H	OH	1
177	Et	H	H	H	H	OH	2
178	Et	H	F	H	H	OH	2
179	Et	H	Cl	H	H	OH	2
180	Et	H	Br	H	H	OH	2
181	Et	H	Me	H	H	OH	2
182	Et	H	Et	H	H	OH	2
183	Et	H	Pr	H	H	OH	2
184	Et	H	CF ₃	H	H	OH	2
185	Et	H	C ₂ F ₅	H	H	OH	2
186	Et	H	SF ₅	H	H	OH	2
187	Et	H	OCH ₃	H	H	OH	2
188	Et	H	OC ₂ H ₅	H	H	OH	2
189	Et	H	OCF ₃	H	H	OH	2
190	Et	H	SCF ₃	H	H	OH	2
191	Et	H	SOCF ₃	H	H	OH	2
192	Et	H	SO ₂ CF ₃	H	H	OH	2
193	Me	H	H	H	H	Cl	0
194	Me	H	F	H	H	Cl	0
195	Me	H	Cl	H	H	Cl	0

[0582]

Table 72

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n
196	Me	H	Br	H	H	Cl	0
197	Me	H	Me	H	H	Cl	0
198	Me	H	Et	H	H	Cl	0
199	Me	H	Pr	H	H	Cl	0
200	Me	H	CF ₃	H	H	Cl	0
201	Me	H	C ₂ F ₅	H	H	Cl	0

202	Me	H	SF ₅	H	H	Cl	0
203	Me	H	OCH ₃	H	H	Cl	0
204	Me	H	OC ₂ H ₅	H	H	Cl	0
205	Me	H	OCF ₃	H	H	Cl	0
206	Me	H	SCF ₃	H	H	Cl	0
207	Me	H	SOCF ₃	H	H	Cl	0
208	Me	H	SO ₂ CF ₃	H	H	Cl	0
209	Me	H	H	H	H	Cl	1
210	Me	H	F	H	H	Cl	1
211	Me	H	Cl	H	H	Cl	1
212	Me	H	Br	H	H	Cl	1
213	Me	H	Me	H	H	Cl	1
214	Me	H	Et	H	H	Cl	1
215	Me	H	Pr	H	H	Cl	1
216	Me	H	CF ₃	H	H	Cl	1
217	Me	H	C ₂ F ₅	H	H	Cl	1
218	Me	H	SF ₅	H	H	Cl	1
219	Me	H	OCH ₃	H	H	Cl	1
220	Me	H	OC ₂ H ₅	H	H	Cl	1

[0583]

Table 73

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n
221	Me	H	OCF ₃	H	H	Cl	1
222	Me	H	SCF ₃	H	H	Cl	1
223	Me	H	SOCF ₃	H	H	Cl	1
224	Me	H	SO ₂ CF ₃	H	H	Cl	1
225	Me	H	H	H	H	Cl	2
226	Me	H	F	H	H	Cl	2
227	Me	H	Cl	H	H	Cl	2
228	Me	H	Br	H	H	Cl	2
229	Me	H	Me	H	H	Cl	2
230	Me	H	Et	H	H	Cl	2
231	Me	H	Pr	H	H	Cl	2
232	Me	H	CF ₃	H	H	Cl	2
233	Me	H	C ₂ F ₅	H	H	Cl	2
234	Me	H	SF ₅	H	H	Cl	2
235	Me	H	OCH ₃	H	H	Cl	2
236	Me	H	OC ₂ H ₅	H	H	Cl	2
237	Me	H	OCF ₃	H	H	Cl	2
238	Me	H	SCF ₃	H	H	Cl	2

239	Me	H	SOCl ₂	H	H	Cl	2
240	Me	H	SO ₂ Cl ₂	H	H	Cl	2
241	Et	H	H	H	H	Cl	0
242	Et	H	F	H	H	Cl	0
243	Et	H	Cl	H	H	Cl	0
244	Et	H	Br	H	H	Cl	0
245	Et	H	Me	H	H	Cl	0

[0584]

Table 74

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n
246	Et	H	Et	H	H	Cl	0
247	Et	H	Pr	H	H	Cl	0
248	Et	H	CF ₃	H	H	Cl	0
249	Et	H	C ₂ F ₅	H	H	Cl	0
250	Et	H	SF ₅	H	H	Cl	0
251	Et	H	OCH ₃	H	H	Cl	0
252	Et	H	OC ₂ H ₅	H	H	Cl	0
253	Et	H	OCF ₃	H	H	Cl	0
254	Et	H	SCF ₃	H	H	Cl	0
255	Et	H	SOCl ₂	H	H	Cl	0
256	Et	H	SO ₂ Cl ₂	H	H	Cl	0
257	Et	H	H	H	H	Cl	1
258	Et	H	F	H	H	Cl	1
259	Et	H	Cl	H	H	Cl	1
260	Et	H	Br	H	H	Cl	1
261	Et	H	Me	H	H	Cl	1
262	Et	H	Et	H	H	Cl	1
263	Et	H	Pr	H	H	Cl	1
264	Et	H	CF ₃	H	H	Cl	1
265	Et	H	C ₂ F ₅	H	H	Cl	1
266	Et	H	SF ₅	H	H	Cl	1
267	Et	H	OCH ₃	H	H	Cl	1
268	Et	H	OC ₂ H ₅	H	H	Cl	1
269	Et	H	OCF ₃	H	H	Cl	1
270	Et	H	SCF ₃	H	H	Cl	1

[0585]

Table 75

Intermediate	R ^{1x}	R ^{2x}	R ^{3x}	R ^{4x}	R ^{5x}	R ^{13x}	n

271	Et	H	SOCF ₃	H	H	Cl	1
272	Et	H	SO ₂ CF ₃	H	H	Cl	1
273	Et	H	H	H	H	Cl	2
274	Et	H	F	H	H	Cl	2
275	Et	H	Cl	H	H	Cl	2
276	Et	H	Br	H	H	Cl	2
277	Et	H	Me	H	H	Cl	2
278	Et	H	Et	H	H	Cl	2
279	Et	H	Pr	H	H	Cl	2
280	Et	H	CF ₃	H	H	Cl	2
281	Et	H	C ₂ F ₅	H	H	Cl	2
282	Et	H	SF ₅	H	H	Cl	2
283	Et	H	OCH ₃	H	H	Cl	2
284	Et	H	OC ₂ H ₅	H	H	Cl	2
285	Et	H	OCF ₃	H	H	Cl	2
286	Et	H	SCF ₃	H	H	Cl	2
287	Et	H	SOCF ₃	H	H	Cl	2
288	Et	H	SO ₂ CF ₃	H	H	Cl	2

[0586]

In the above Tables 62-75, Me represents a methyl group, and Et represents an ethyl group.

[0587]

5 The ¹H-NMR data of Compound (1H) are shown below.

Compound (1H)-184

¹H-NMR (CDCl₃) δ: 9.23 (1H,brs), 8.37 (1H,brs), 8.03-7.98 (2H,m), 3.63 (2H,q), 1.39 (3H,t)

Compound (1H)-280

10 ¹H-NMR (CDCl₃) δ: 8.33 (1H,brs), 8.09-7.98 (2H,m), 3.43 (2H,q), 1.38 (3H,t)

Compound (1H)-177

¹H-NMR (CDCl₃) δ: 8.12 (1H,dd), 7.87 (1H,dd), 7.77-7.68 (2H,m), 3.59 (2H,q), 1.35 (3H,t).

Compound (1H)-248

$^1\text{H-NMR}$ (CDCl_3) δ : 8.22 (1H,d), 7.55 (1H,s), 7.43 (1H,d),
3.02 (2H,q), 1.43 (3H,t).

[0588]

5 The $^1\text{H-NMR}$ data of Compound (1I) are shown below.

Compound (1I)-88

$^1\text{H-NMR}$ (CDCl_3) δ : 8.43-8.42 (1H,m), 8.10-8.03 (2H,m), 3.45
(2H,q), 1.38 (3H,t)

[0589]

10 Formulation Example 1

Any one of Present Compounds 1-427 (10 parts) is
dissolved in a mixture of xylene (35 parts) and N,N-
dimethylformamide (35 parts), and to the mixture is added
polyoxyethylene styryl phenyl ether (14 parts) and calcium
15 dodecylbenzenesulfonate (6 parts), and stirred to give
emulsions of each compound.

[0590]

Formulation Example 2

Sodium lauryl sulfate (4 parts), calcium lignin
20 sulfonate (2 parts), synthetic hydrated silicone oxide
powder (20 parts) and diatomite (54 parts) are mixed, then
to the mixture is added any one of Present Compounds 1-427
(20 parts), and mixed to give wettable powders of each
compound.

25 [0591]

Formulation Example 3

To any one of Present Compounds 1-427 (2 parts) is added synthetic hydrated silicone oxide powder (1 part), calcium lignin sulfonate (2 parts), bentonite (30 parts), and kaolin clay (65 parts), and mixed. Then, to the mixture is added an appropriate amount of water, further stirred, granulated with a granulator, and draft-dried to give granules of each compound.

[0592]

10 Formulation Example 4

Any one of Present Compounds 1-427 (1 part) is dissolved in an appropriate amount of acetone. To the mixture is added synthetic hydrated silicone oxide powder (5 parts), PAP (0.3 parts), and Fubasami clay (93.7 parts), and well stirred. Then, acetone is removed by evaporation to give powders of each compound.

[0593]

Formulation Example 5

A mixture (ratio by weight = 1:1) of polyoxyethylene alkyl ether sulfate ammonium salt and white carbon (35 parts), any one of Present Compounds 1-427 (10 parts), and water (55 parts) are mixed, pulverized by a wet grinding method to give formulations of each compounds.

[0594]

25 Formulation Example 6

Any one of Present Compounds 1-427 (0.1 parts) is dissolved in xylene (5 parts) and trichloroethane (5 parts), and mixed with deodorized kerosine (89.9 parts) to give oil solutions of each compounds.

5 [0595]

Formulation Example 7

Any one of Present Compounds 1-427 (10 mg) is dissolved in acetone (0.5 ml). The mixture is added to animal powdered solid feed (powdered solid feed for
10 breeding, CE-2, from CLEA Japan, Inc.), (5 g) and mixed uniformly. Then, acetone is removed by evaporation to give poison baits of each compound.

[0596]

Formulation Example 8

15 Any one of Present Compounds 1-427 (0.1 parts) and Neothiosol (Chuo Kasei Co. Ltd.) (49.9 parts) are charged into an aerosol container. After an aerosol valve is attached to the container, dimethyl ether (25 parts) and LPG (25 parts) are charged into the container. The
20 container is vibrated, and attaching an actuator to give oily aerosols of each compound.

[0597]

Formulation Example 9

Any one of Present Compounds 1-427 (0.6 parts), BHT
25 (2,6-di-tert-butyl-4-methylphenol) (0.01 parts), xylene (5

parts), deodorized kerosine (3.39 parts), and an emulsifier (Atmos 300 (a registered trade name for Atmos Chemical Ltd.)) (1 part) are mixed and dissolved. The mixture and distilled water (50 parts) are charged into an aerosol container, and attaching a valve. Then, propellant (LPG) (40 parts) is pressure-charged into the container through the valve to give aqueous aerosols of each compound.

[0598]

The effects of the present compounds to control pests are shown in Test Examples.

[0599]

Test Example 1

The test spray solutions were prepared by diluting the formulations of each of Present Compounds 1-7, 10-14, 18-23, 33-38, 46, 53, 55, 66, 68-70, 72, 75, 81-83, 88-90, 98-99, 102-104, 112-113, 115-118, 121-123, 126-142, 144, 151, 156-159, 161, 167-171, 174-176, 183, 185-186, 188, 192-194, 209-210, 214-215, 235, 238-243, 246-256, 259, 262, 264, 270-273, 279-280, 285, 291-292, 294, 297-299, 301-302, 306-313, 316-319, 324-325, 327, 330-331, 333, 343, 345, 353, 357-361, 371 and 374-379 obtained in Formulation Example 5 with water so as to give 500 ppm of an active ingredient concentration.

On the other hand, on a cucumber seedling (the first true leaf stage) planted in a plastic cup was inoculated

with about 30 *Aphis gossypii*, and leaving it for a day. Twenty ml of each of the above test spray solutions was sprayed on this seedling.

Six days after spraying, the number of the surviving
5 *Aphis gossypii* parasitized on the leaves of the cucumber was examined, and a control value was calculated according to the following equation:

$$\text{Control value (\%)} = \{1 - (\text{Cb} \times \text{Tai}) / (\text{Cai} \times \text{Tb})\} \times 100$$

10 wherein symbols represent as follows:

Cb: the number of insects in a non-treated section before treatment,

Cai: the number of insects in a non-treated section in observation,

15 Tb: the number of insects in a treated-section before treatment,

Tai: the number of insects in a treated-section in observation;

wherein the non-treated section represents a section
20 where the test spray solution prepared by diluting the formulation without the present compound in Formulation Example 5 with the same amount of water as in the treated-section was sprayed.

25 As a result, in the treated-section of the test spray

solution of each of Present Compounds 1-7, 10-14, 18-23, 33-38, 46, 53, 55, 66, 68-70, 72, 75, 81-83, 88-90, 98-99, 102-104, 112-113, 115-118, 121-123, 126-142, 144, 151, 156-159, 161, 167-171, 174-176, 183, 185-186, 188, 192-194, 209-210, 214-215, 235, 238-243, 246-256, 259, 262, 264, 270-273, 279-280, 285, 291-292, 294, 297-299, 301-302, 306-313, 316-319, 324-325, 327, 330-331, 333, 343, 345, 353, 357-361, 371 and 374-379, the control value of 90 % or more was shown.

[0600]

Test Example 2

The test diluted solutions were prepared by diluting the formulations of each of Present Compounds 1-6, 13, 14, 16, 21-23, 33, 34, 36, 43, 46, 48-50, 53, 55, 66, 69, 70, 82, 83, 88, 89, 98-100, 116-118, 122-123, 133-135, 137, 139-140, 142-143, 156-159, 167, 169, 171, 175-176, 192-194, 224, 237-239, 241-243, 247, 254, 256, 280, 299, 306, 317-319, 330-331, 333, 353, 376 and 379 obtained in Formulation Example 5 with water so as to give 500 ppm of an active ingredient concentration.

A root part of a cucumber seedling (the first true leaf stage) from which soil had been washed off was immersed in 5 ml of each of the diluted solutions, and one day after treatment, on the cucumber leaf surface was inoculated with 30 *Aphis gossypii* (whole stage). After

further seven days, the number of insect of living *Aphis gossypii* parasitized on the leaves of the cucumber was examined, and a control value was calculated according to the following equation:

5

$$\text{Control value (\%)} = \{1 - (\text{Cb} \times \text{Tai}) / (\text{Cai} \times \text{Tb})\} \times 100$$

wherein symbols represent as follows:

Cb: the number of insects in a non-treated section before treatment,

10

Cai: the number of insects in a non-treated section in observation,

Tb: the number of insects in a treated-section before treatment,

15

Tai: the number of insects in a treated-section in observation;

20

wherein the non-treated section represents a section where the test diluted solution prepared by diluting the formulation without the present compound in Formulation Example 5 with the same amount of water as in the treated-section was used.

25

As a result, in the treated-section of the test diluted solution of each of Present Compounds 1-6, 13, 14, 16, 21-23, 33, 34, 36, 43, 46, 48-50, 53, 55, 66, 69, 70, 82, 83, 88, 89, 98-100, 116-118, 122-123, 133-135, 137,

139-140, 142-143, 156-159, 167, 169, 171, 175-176, 192-194, 224, 237-239, 241-243, 247, 247, 254, 256, 280, 299, 306, 317-319, 330-331, 333, 353, 376 and 379, the control value of 90 % or more was shown.

5 [0601]

Test Example 3

The test diluted solutions were prepared by diluting the formulations of each of Present Compounds 2-6, 35-38, 50, 55, 89, 90, 112-113, 116-118, 123, 130, 132, 134-135, 10 157-158, 169-171, 175-176, 194, 209-210, 214-215, 239-240, 242-243, 248, 254, 270-273, 280, 299, 302, 306, 309, 318-319, 330-331, 333, 345, 359, 361 and 375-376 obtained in Formulation Example 5 with water so as to give 500 ppm of an active ingredient concentration.

15 On the other hand, a cucumber seedling (the second true leaf stage) planted in a plastic cup was drenched at its foot with 5 ml of each of the diluted solutions, and kept in a greenhouse of 25 °C for 7 days. On the cucumber leaf surface was inoculated with 30 *Aphis gossypii* (whole 20 stage), and further kept in the greenhouse for 6 days, then the number of insect of living *Aphis gossypii* parasitized on the leaves of this cucumber was examined, and a control value was calculated according to the following equation:

25 Control value (%) = $\{1 - (C_b \times T_{ai}) / (C_{ai} \times T_b)\} \times 100$

wherein symbols represent as follows:

Cb: the number of insects in a non-treated section before treatment,

5 Cai: the number of insects in a non-treated section in observation,

Tb: the number of insects in a treated-section before treatment,

Tai: the number of insects in a treated-section in observation;

10 wherein the non-treated section represents a section where the test diluted solution prepared by diluting the formulation without the present compound in Formulation Example 5 with the same amount of water as in the treated-section was used.

15 As a result, in the treated-section of the test diluted solution of each of Present Compounds 2-6, 35-38, 50, 55, 89, 90, 112-113, 116-118, 123, 130, 132, 134-135, 157-158, 169-171, 175-176, 194, 209-210, 214-215, 239-240, 20 242-243, 248, 254, 270-273, 280, 299, 302, 306, 309, 318-319, 330-331, 333, 345, 359, 361 and 375-376, the control value of 90 % or more was shown.

[0602]

Test Example 4

25 The test spray solutions were prepared by diluting the

formulations of each of Present Compounds 1, 3-7, 12, 14, 21, 33-38, 46, 55, 68, 69, 83, 84, 88-90, 102, 112-113, 116-118, 128, 130-133, 136-139, 142-143, 152, 156-158, 161, 169-171, 174-176, 183-186, 188, 201, 209-210, 214-215, 238, 246, 248, 251, 262, 292, 298-299, 301-302, 306, 308-309, 314, 324-325, 327, 343, 353, 360, 374 and 376 obtained in Formulation Example 5 with water so as to give 500 ppm of an active ingredient concentration.

On the other hand, a rice seedling (the second leaf stage) planted in a polyethylene cup was sprayed with 10 ml of each of the above test spray solutions. After air-drying, 20 third-fourth instar larvae of *Nilaparvata lugens* were released, and kept in the greenhouse of 25 °C. After 6 days, the number of insect of living *Nilaparvata lugens* parasitized on the rice was examined, and a control value was calculated according to the following equation:

$$\text{Control value (\%)} = \{1 - (\text{Cb} \times \text{Tai}) / (\text{Cai} \times \text{Tb})\} \times 100$$

wherein symbols represent as follows:

Cb: the number of insects in a non-treated section before treatment,

Cai: the number of insects in a non-treated section in observation,

Tb: the number of insects in a treated-section before treatment,

Tai: the number of insects in a treated-section in observation;

wherein the non-treated section represents a section where the test spray solution prepared by diluting the formulation without the present compound in Formulation Example 5 with the same amount of water as in the treated-section was sprayed.

As a result, in the treated-section of the test spray solution of each of Present Compounds 1, 3-7, 12, 14, 21, 33-38, 46, 55, 68, 69, 83, 84, 88-90, 102, 112-113, 116-118, 128, 130-133, 136-139, 142-143, 152, 156-158, 161, 169-171, 174-176, 183-186, 188, 201, 209-210, 214-215, 238, 246, 248, 251, 262, 292, 298-299, 301-302, 306, 308-309, 314, 324-325, 327, 343, 353, 360, 374 and 376, the control value of 90 % or more was shown.

[0603]

Test Example 5

The test diluted solutions were prepared by diluting the formulations of each of Present Compounds 4-6, 33-36, 38, 55, 89, 90, 112-113, 117-118, 123, 130, 132, 137-138, 143, 156-158, 169-171, 174-176, 185, 214, 248, 251, 272, 292, 294, 297-299, 301-302, 306, 324, 327, 343, 345, 353, 357, 359-361 and 374-376 obtained in Formulation Example 5 with water so as to give 500 ppm of an active ingredient

concentration.

On the other hand, a rice seedling (2 weeks after sowing, the second leaf stage) planted in a plastic cup was drenched at its foot with 5 ml of each of the diluted solutions, and kept in a greenhouse of 25 °C for 7 days. Twenty third-fourth instar larvae of *Nilaparvata lugens* were released, and further kept in the greenhouse for 6 days, then the number of insect of living *Nilaparvata lugens* parasitized on the rice was examined, and a control value was calculated according to the following equation:

$$\text{Control value (\%)} = \{1 - (\text{Cb} \times \text{Tai}) / (\text{Cai} \times \text{Tb})\} \times 100$$

wherein symbols represent as follows:

Cb: the number of insects in a non-treated section before treatment,

Cai: the number of insects in a non-treated section in observation,

Tb: the number of insects in a treated-section before treatment,

Tai: the number of insects in a treated-section in observation;

wherein the non-treated section represents a section where the test diluted solution prepared by diluting the formulation without the present compound in Formulation Example 5 with the same amount of water as in the treated-

section was used.

As a result, in the treated-section of the test diluted solution of each of Present Compounds 4-6, 33-36, 38, 55, 89, 90, 112-113, 117-118, 123, 130, 132, 137-138, 143, 156-158, 169-171, 174-176, 185, 214, 248, 251, 272, 292, 294, 297-299, 301-302, 306, 324, 327, 343, 345, 353, 357, 359-361 and 374-376, the control value of 90 % or more was shown.

10 [0604]

Test Example 6

The test spray solutions were prepared by diluting the formulations of each of Present Compounds 4-6, 21, 22, 49, 50, 55, 75, 88, 112-113, 116-118, 128-132, 161, 169-171, 174-176, 188, 209-210, 214-215, 238-240, 247-248, 255, 259, 262, 270-273, 280, 291-292, 296-299, 301-302, 306-311, 317, 319, 331, 333, 343, 345, 357, 360, 374 and 376 obtained in Formulation Example 5 with water so as to give 500 ppm of an active ingredient concentration.

20 On the other hand, Bemisia tabaci adult was released on a tomato seedling (the third true leaf stage) planted in a polyethylene cup, and made to lay eggs for about 72 hours. The tomato seedling was kept in a greenhouse for 8 days. When instar larvae hatched from the eggs, the above test
25 spray solution was sprayed in the amount of 20 ml/cup. The

cup was kept in a greenhouse at 25 °C. After the keeping for 7 days, the number of surviving instar larvae on the tomato leaves was examined, and a control value was calculated according to the following equation:

5

$$\text{Control value (\%)} = \{1 - (\text{Cb} \times \text{Tai}) / (\text{Cai} \times \text{Tb})\} \times 100$$

wherein symbols represent as follows:

Cb: the number of insects in a non-treated section before treatment,

10

Cai: the number of insects in a non-treated section in observation,

Tb: the number of insects in a treated-section before treatment,

15

Tai: the number of insects in a treated-section in observation;

20

wherein the non-treated section represents a section where the test spray solution prepared by diluting the formulation without the present compound in Formulation Example 5 with the same amount of water as in the treated-section was sprayed.

25

As a result, in the treated-section of the test spray solution of each of Present Compounds 4-6, 21, 22, 49, 50, 55, 75, 88, 112-113, 116-118, 128-132, 161, 169-171, 174-176, 188, 209-210, 214-215, 238-240, 247-248, 255, 259, 262,

270-273, 280, 291-292, 296-299, 301-302, 306-311, 317, 319, 331, 333, 343, 345, 357, 360, 374 and 376, the control value of 90 % or more was shown.

[0605]

5 Test Example 7

The test spray solutions were prepared by diluting the formulations of each of Present Compounds 1-14, 16-23, 25, 28-36, 42-47, 49, 50, 53-55, 59, 64, 66, 68-76, 80, 81-83, 85-90, 97-98, 100, 116-118, 123, 133-142, 144, 151-154, 10 156-160, 169-171, 174-176, 181, 183-188, 190-191, 193-194, 197-198, 201-203, 209-210, 214-215, 230, 238-243, 246-248, 262-264, 278-281, 285-287, 292, 294, 299-303, 306-307, 311, 313-316, 318-319, 327, 330-333, 335, 341-343, 345, 348-349, 353, 358, 360, 365, 369-371 and 374-378 obtained in 15 Formulation Example 5 with water so as to give 500 ppm of an active ingredient concentration.

On the other hand, Cabbage (the third leaf stage) planted in a polyethylene cup was sprayed with 20 mL/cup of each of the test spray solution. After the test solution 20 was dried, the aerial part was cut off, and then placed in a 50 mL volume cup. Five second instar larvae of *Plutella xylostella* were released into the cup, and the cup was sealed with a lid. After the cup was kept at 25 °C for 5 days, the number of living insects was counted. A death 25 rate was calculated according to the following equation:

Death rate (%) = (Number of dead insects/Number of tested insects) x 100

5 As a result, in the treated-section of the test spray solution of each of Present Compounds 1-14, 16-23, 25, 28-36, 42-47, 49, 50, 53-55, 59, 64, 66, 68-76, 80-83, 85-90, 97-98, 100, 116-118, 123, 133-142, 144, 151-154, 156-160, 169-171, 174-176, 181, 183-188, 190-191, 193-194, 197-198, 10 201-203, 209-210, 214-215, 230, 238-243, 246-248, 262-264, 278-281, 285-287, 292, 294, 299-303, 306-307, 311, 313-316, 318-319, 327, 330-333, 335, 341-343, 345, 348-349, 353, 358, 360, 365, 369-371 and 374-378, the death rate of 80 % or more was shown.

15 [0606]

Test Example 8

 The test spray solutions were prepared by diluting the formulations of each of Present Compounds 2-9, 11, 13, 14, 17-20, 22, 23, 28, 29, 31, 35-37, 43-47, 54-55, 68-83, 86-90, 20 106, 112-113, 117-118, 128-129, 131-138, 140-142, 157-158, 161, 164, 169-171, 174-176, 183-185, 187-188, 191, 194, 201-203, 209-210, 214-215, 230, 238-240, 242-243, 246-248, 251, 262-263, 270-275, 278-281, 285-287, 290, 292, 294, 296-302, 305-316, 318-319, 324-325, 327, 329-333, 335, 341-25 343, 345, 349, 355-357, 360, 369-371 and 374-378 obtained

in Formulation Example 5 with water so as to give 500 ppm of an active ingredient concentration.

On the other hand, an apple tree was planted in a plastic cup, and grown until the seventh-eighth leaf was spread. The apple tree was sprayed with 20 mL/cup of each of the test spray solution. After the test solution was dried, 60 first-instar *Adoxophyes orana fasciata* were released, and the cup was covered with a plastic cup upside-down which the bottom was cut off and a filter paper was put thereon. After 7 days, the number of living insects was counted, and a death rate was calculated according to the following equation:

$$\text{Death rate (\%)} = (\text{Number of dead insects} / \text{Number of tested insects}) \times 100$$

As a result, in the treated-section of the test spray solution of each of Present Compounds 2-9, 11, 13, 14, 17-20, 22, 23, 28, 29, 31, 35-37, 43-47, 54-55, 68-83, 86-90, 106, 112-113, 117-118, 128-129, 131-138, 140-142, 157-158, 161, 164, 169-171, 174-176, 183-185, 187-188, 191, 194, 201-203, 209-210, 214-215, 230, 238-240, 242-243, 246-248, 251, 262-263, 270-275, 278-281, 285-287, 290, 292, 294, 296-302, 305-316, 318-319, 324-325, 327, 329-333, 335, 341-343, 345, 349, 355-357, 360, 369-371 and 374-378, the death

rate of 90 % or more was shown.


Industrial Applicability

[0607]

- 5 The present compound has an activity of controlling pests and is useful as an active ingredient of a pest controlling agent.

CLAIMS

1. A fused heterocyclic compound of the formula (1):



wherein:

- 5 A¹ represents -NR⁸-, an oxygen atom, or a sulfur atom;
 A² represents a nitrogen atom or =CR⁹-;
 A³ represents a nitrogen atom or =CR¹⁰-;
 R¹ represents a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from
 10 Group X, or a C3-C6 alicyclic hydrocarbon group optionally having one or more atoms or groups selected from Group Y;
 R², R³, R⁴, and R⁵ are same or different and independently represent a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a phenyl group
 15 optionally having one or more atoms or groups selected from Group Z, a 5-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z, a 6-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z, -OR¹¹, -S(O)_mR¹¹, -
 20 NR¹¹R¹², -CO₂R¹¹, -C(O)R¹¹, a cyano group, a nitro group, a halogen atom, -SF₅, or a hydrogen atom, provided that at least two of R², R³, R⁴, and R⁵ represent a hydrogen atom;

R^6 and R^7 are same or different and independently represent a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from Group X, a phenyl group optionally having one or more atoms or groups selected from Group Z, a 5-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z, a 6-membered heterocyclic group optionally having one or more atoms or groups selected from Group Z, $-OR^{11}$, $-S(O)_mR^{11}$, $-S(O)_2NR^{11}R^{12}$, $-NR^{11}R^{12}$, $-NR^{11}-CO_2R^{12}$, $-NR^{11}C(O)R^{12}$, $-CO_2R^{11}$, $-C(O)R^{11}$, a cyano group, a nitro group, a halogen atom, $-SF_5$, or a hydrogen atom;

R^8 represents a C1-C6 chain hydrocarbon group optionally having one or more atoms or groups selected from Group W, $-CO_2R^{11}$, $-C(O)R^{11}$, a C3-C6 alicyclic hydrocarbon group optionally having one or more atoms or groups selected from Group Y, or a hydrogen atom;

R^9 and R^{10} are same or different and independently represent a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, $-NR^{11}R^{12}$, $-CO_2R^{11}$, $-C(O)R^{11}$, a cyano group, a nitro group, a halogen atom, or a hydrogen atom;

R^{11} and R^{12} are same or different and independently represent a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a hydrogen atom; and

m represents 0, 1, or 2; and n represents 0, 1, or 2;

(except in cases as follows: both R^6 and R^7 are a hydrogen atom; in $-S(O)_mR^{11}$, when m is 1 or 2, R^{11} is a hydrogen atom; and when A^1 represents a methylimino group, A^2 and A^3 represent a methine group, R^1 represents a methyl group, R^2 , R^3 , R^4 , and R^5 represent a hydrogen atom, and n is 0 or 1, both R^6 and R^7 are a chlorine atom and one of R^6 and R^7 is a chlorine atom and the other is a hydrogen atom)

the Group X consists of a C1-C6 alkoxy group optionally having one or more halogen atoms, a C2-C6 alkenyloxy group optionally having one or more halogen atoms, a C2-C6 alkynyloxy group optionally having one or more halogen atoms, a C1-C6 alkylsulfanyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfinyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfonyl group optionally having one or more halogen atoms, a C2-C6 alkylcarbonyl group optionally having one or more halogen atoms, a C2-C6 alkoxycarbonyl group optionally having one or more halogen atoms, a C3-C6 cycloalkyl group optionally having one or more halogen atoms, a cyano group, a hydroxy group, and a halogen atom;

the Group Y consists of a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 alkoxy group optionally having one or more halogen atoms, a C2-C6 alkenyloxy group optionally having one or more halogen atoms, a C2-C6 alkynyloxy group optionally having

one or more halogen atoms, a hydroxy group, and a halogen atom;

the Group Z consists of a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 alkylsulfanyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfinyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfonyl group optionally having one or more halogen atoms, a C2-C6 alkylcarbonyl group optionally having one or more halogen atoms, a C2-C6 alkoxy carbonyl group optionally having one or more halogen atoms, a C1-C6 alkylamino group optionally having one or more halogen atoms, a C2-C8 dialkylamino group optionally having one or more halogen atoms, a halogen atom, a cyano group, and a nitro group; and

the Group W consists of a C1-C6 alkoxy group optionally having one or more halogen atoms, a C2-C6 alkenyloxy group optionally having one or more halogen atoms, a C2-C6 alkynyloxy group optionally having one or more halogen atoms, a C1-C6 alkylsulfanyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfinyl group optionally having one or more halogen atoms, a C1-C6 alkylsulfonyl group optionally having one or more halogen atoms, a C2-C6 alkylcarbonyl group, a C2-C6 alkoxy carbonyl group, a C3-C6 cycloalkyl group, a halogen atom, a cyano

group, and a hydroxy group.

2. The fused heterocyclic compound according to claim 1 wherein R^6 and R^7 are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, $-NR^{11}R^{12}$, a cyano group, a nitro group, a halogen atom, or a hydrogen atom.

3. The fused heterocyclic compound according to claim 1 or claim 2 wherein R^6 and R^7 are same or different and are independently a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, $-S(O)_mR^{11}$, a halogen atom, or a hydrogen atom.

4. The fused heterocyclic compound according to any one of claims 1-3 wherein A^2 is $=CH-$.

5. The fused heterocyclic compound according to any one of claims 1-4 wherein A^3 is a nitrogen atom or $=CR^{10}-$, and R^{10} is a halogen atom or a hydrogen atom.

6. The fused heterocyclic compound according to any one of claims 1-4 wherein A^3 is a nitrogen atom.

7. The fused heterocyclic compound according to any one of claims 1-6 wherein R^2 , R^4 , and R^5 are same or different and are independently a hydrogen atom or a halogen atom, and R^3 is a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, $-OR^{11}$, a halogen atom, or a hydrogen atom.


8. The fused heterocyclic compound according to any one of claims 1-7 wherein A^1 is $-NR^8-$, and R^8 is a C1-C6 chain hydrocarbon group having one C1-C6 alkoxy group optionally having one or more halogen atoms, a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, or a cyclopropyl group.

9. The fused heterocyclic compound according to any one of claims 1-7 wherein A^1 is $-NR^8-$, and R^8 is a methyl group, an ethyl group, a methoxymethyl group, or an ethoxymethyl group.

10. The fused heterocyclic compound according to any one of claims 1-7 wherein A^1 is an oxygen atom.

11. The fused heterocyclic compound according to any one of claims 1-7 wherein A^1 is a sulfur atom.

12. The fused heterocyclic compound according to claim 1 which is represented by the formula (1-1):



wherein:

- 5 A^{1a} represents -NR^{8a}- or a sulfur atom;
- A^{3a} represents a nitrogen atom or =CR^{10a}-;
- R^{1a} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms;
- R^{2a}, R^{4a}, and R^{5a} are same or different and
- 10 independently represent a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, a halogen atom, or a hydrogen atom;
- R^{3a} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, -OR^{11a}, a
- 15 halogen atom, or a hydrogen atom;
- R^{6a} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms, -OR^{11a}, -S(O)_mR^{11a}, a bromine atom, or an iodine atom;
- R^{8a} represents a C1-C6 chain hydrocarbon group
- 20 optionally having one or more halogen atoms or a hydrogen atom;
- R^{10a} represents a halogen atom or a hydrogen atom;

R^{11a} represents a C1-C6 chain hydrocarbon group optionally having one or more halogen atoms; and

m represents 0, 1, or 2, and n represents 0, 1, or 2;

provided that in $-S(O)_mR^{11}$, when m is 1 or 2, R^{11} is not a hydrogen atom.

13. The fused heterocyclic compound according to claim 12 wherein A^{1a} is $-NR^{8a}-$ or a sulfur atom, R^{8a} is a methyl group, A^{3a} is a nitrogen atom, R^{1a} is an ethyl group, R^{2a} , R^{4a} , and R^{5a} are same or different and are independently a halogen atom or a hydrogen atom, R^{3a} is a trifluoromethyl group, a halogen atom, or a hydrogen atom, and R^{6a} is a C1-C3 alkyl group having one or more fluorine atoms, a C1-C3 alkoxy group having one or more fluorine atoms, a C1-C3 alkylsulfanyl group having one or more fluorine atoms, a C1-C3 alkylsulfinyl group having one or more fluorine atoms, or a C1-C3 alkylsulfonyl group having one or more fluorine atoms.

14. A pest controlling composition which comprises the fused heterocyclic compound according to any one of claims 1-13 and an inert carrier.

15. A method of controlling pests which comprises applying an effective amount of the fused heterocyclic compound

according to any one of claims 1-13 to pests or habitats of
pests.

INTERNATIONAL SEARCH REPORT

International application No

PCT/JP2011/080557

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D235/18 C07D277/68 C07D471/04 C07D498/04 C07D513/04
C07D263/57 A01N43/52 A01N43/76 A01N43/78 A01N43/90

ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 262 845 A1 (FISONS PLC [GB]) 6 April 1988 (1988-04-06) cited in the application Ex 8 a-e, Ex 9a i-v, Ex 9b i-iv, Ex 9c i-v, Ex 9d i,ii, Ex 9e i-iv; page 19 - page 21 -----	1-5,7-9
X	GB 1 445 824 A (THOMAE GMBH DR K) 11 August 1976 (1976-08-11) page 12 - page 14; examples 63, 64, 73, 74 ----- -/-	1-7

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

27 February 2012

Date of mailing of the international search report

06/03/2012

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Sotoca Usina, E

INTERNATIONAL SEARCH REPORT

International application No

PCT/JP2011/080557

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SLUKA J ET AL: "2-Phenylbenzimidazoles as Potential Anthelmintics", COLLECTION OF CZECHOSLOVAK CHEMICAL COMMUNICATIONS, INSTITUTE OF ORGANIC CHEMISTRY & BIOCHEMISTRY, PRAGUE; CZ, vol. 41, 1 January 1976 (1976-01-01), pages 3628-3631, XP009156807, ISSN: 0010-0765 page 3629; examples XXXV-XXXIX, LXXV-LXXVIII Not active against worms or coccidia in animals.; page 3630, paragraph 2 -----	1-4,7
X	US 2004/198768 A1 (PARK CHOO HEA YOUNG [KR] ET AL) 7 October 2004 (2004-10-07) page 6; example 15; table 6 -----	1-4,7,11
X	US 4 038 396 A (SHEN TSUNG-YING ET AL) 26 July 1977 (1977-07-26) Compounds in example 65; column 20; example 65 -----	1-3,5,7,10
X	COATES W J ET AL: "CYCLIC NUCLEOTIDE PHOSPHODIESTERASE INHIBITION BY IMIDAZOPYRIDINES: ANALOGUES OF SULMAZOLE AND ISOMALZOLE AS INHIBITORS OF THE CGMP SPECIFIC PHOSPHODIESTERASE", JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, US, vol. 36, no. 10, 1 May 1993 (1993-05-01), pages 1387-1392, XP002924367, ISSN: 0022-2623, DOI: 10.1021/JM00062A011 page 1389; table I; compound 6 -----	1-3,5
X	JP 2008 308448 A (SANKYO AGRO KK) 25 December 2008 (2008-12-25) page 8, paragraph 0001 - page 9, paragraph 0004 page 49 - page 52; compounds 13 1-50; 14 1-50 -----	1-15
X	GB 1 175 013 A (CIBA LTD [CH]) 23 December 1969 (1969-12-23) the whole document -----	1-5,7,8,12-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/JP2011/080557

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0262845	A1	06-04-1988	AU 604771 B2 03-01-1991
		AU 8024487 A 21-04-1988	
		DK 287688 A 01-07-1988	
		EP 0262845 A1 06-04-1988	
		EP 0283504 A1 28-09-1988	
		FI 882394 A 20-05-1988	
		IE 872588 L 27-03-1988	
		JP H01501473 A 25-05-1989	
		NZ 221934 A 28-05-1990	
		PT 85793 A 01-10-1987	
		US 4900751 A 13-02-1990	
		WO 8802367 A1 07-04-1988	

GB 1445824	A	11-08-1976	AT 332873 B 25-10-1976
		AU 6512974 A 07-08-1975	
		BG 23902 A3 10-11-1977	
		CA 1041502 A1 31-10-1978	
		CH 605939 A5 13-10-1978	
		CS 200169 B2 29-08-1980	
		DD 108989 A5 12-10-1974	
		DK 140760 B 12-11-1979	
		ES 422450 A1 01-05-1976	
		FI 58126 B 29-08-1980	
		FR 2215968 A1 30-08-1974	
		GB 1445824 A 11-08-1976	
		HK 4080 A 08-02-1980	
		HU 170909 B 28-09-1977	
		IE 39066 B1 02-08-1978	
		IL 44127 A 31-05-1977	
		JP 1156096 C 15-07-1983	
		JP 49102693 A 27-09-1974	
		JP 57048556 B 16-10-1982	
		NL 7401254 A 06-08-1974	
		NO 139386 B 20-11-1978	
		PH 14988 A 12-03-1982	
		PL 93127 B1 30-05-1977	
		RO 79057 A1 25-06-1982	
		RO 84276 A1 23-05-1984	
		SE 411451 B 27-12-1979	
		SU 563917 A3 30-06-1977	
		SU 634673 A3 25-11-1978	
		YU 17674 A 18-06-1982	
		YU 110480 A 18-06-1982	

US 2004198768	A1	07-10-2004	US 2004198768 A1 07-10-2004
			US 2007066577 A1 22-03-2007

US 4038396	A	26-07-1977	NONE

JP 2008308448	A	25-12-2008	NONE

GB 1175013	A	23-12-1969	AT 283813 B 25-08-1970
		BE 716252 A 09-12-1968	
		CH 475709 A 15-04-1969	
		DE 1770517 B1 09-03-1972	
		FR 1583887 A 05-12-1969	
		GB 1175013 A 23-12-1969	
		NL 6808053 A 09-12-1968	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/JP2011/080557

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
SE361475 B05-11-1973			