

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization

International Bureau

(43) International Publication Date
26 October 2017 (26.10.2017)



(10) International Publication Number
WO 2017/182427 A1

(51) International Patent Classification:

C07K 16/28 (2006.01) A61P 35/00 (2006.01)

(21) International Application Number:

PCT/EP2017/059108

(22) International Filing Date:

18 April 2017 (18.04.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/324,451 19 April 2016 (19.04.2016) US

(71) Applicant: AMGEN RESEARCH (MUNICH) GMBH
[DE/DE]; Staffelseestrasse 2, 81477 Munich (DE).

(72) Inventors: ZUGMAIER, Gerhard; c/o Amgen Research (Munich) GmbH, Staffelseestrasse 2, 81477 Munich (DE). KUFER, Peter; c/o Amgen Research (Munich) GmbH, Staffelseestrasse 2, 81477 Munich (DE). KISCHEL, Roman; c/o Amgen Research (Munich) GmbH, Staffelseestrasse 2, 81477 Munich (DE). SUBKLEWE, Marion; c/o Klinikum der Universitaet Muenchen, Med. Klinik III, Marchioninistrasse 15, 81377 Munich (DE). KRUPKA, Christina; c/o Klinikum der Universitaet Muenchen, Med. Klinik III, Marchioninistrasse 15, 81377 Munich (DE).

(74) Agent: WEINZIERL, Gerhard et al.; Schiweck Weinzierl Koch, Landsberger Strasse 98, 80339 Munich (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,

EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: ADMINISTRATION OF A BISPECIFIC CONSTRUCT BINDING TO CD33 AND CD3 FOR USE IN A METHOD FOR THE TREATMENT OF MYELOID LEUKEMIA

(57) Abstract: The present invention provides a bispecific construct comprising a first binding domain specifically binding to CD33 and a second binding domain specifically binding to CD3 for use in a method for the treatment of myeloid leukemia, wherein the construct is administered for a maximal period of 14 days followed by a period of at least 14 days without administration of the construct. Moreover, the invention provides a method for the treatment of myeloid leukemia comprising the administration of a therapeutically efficient amount of such bispecific construct and the use of such bispecific construct for the preparation of a pharmaceutical composition for the treatment of myeloid leukemia.



WO 2017/182427 A1

**Administration of a bispecific construct binding to CD33 and CD3 for use in a method
for the treatment of myeloid leukemia**

5

Field of the Invention

The present invention relates to a bispecific construct comprising a first binding domain specifically binding to CD33 and a second binding domain specifically binding to CD3 for use
10 in a method for the treatment of myeloid leukemia, wherein the construct is administered for a maximal period of 14 days followed by a period of at least 14 days without administration of the construct. Moreover, the invention relates to method for the treatment of myeloid leukemia comprising the administration of a therapeutically efficient amount of such bispecific
15 construct and the use of such bispecific construct for the preparation of a pharmaceutical composition for the treatment of myeloid leukemia.

Background of the Invention

Bispecific molecules such as BiTE[®] (bispecific T cell engager) antibody constructs are recombinant protein constructs made from two flexibly linked antibody derived binding
20 domains. One binding domain of BiTE[®] antibody constructs is specific for a selected tumor-associated surface antigen on target cells; the second binding domain is specific for CD3, a subunit of the T cell receptor complex on T cells. By their particular design BiTE[®] antibody constructs are uniquely suited to transiently connect T cells with target cells and, at the same time, potently activate the inherent cytolytic potential of T cells against target cells. The first
25 generation of BiTE[®] antibody constructs (see WO 99/54440 and WO 2005/040220) developed into the clinic as AMG 103 (blinatumomab) and AMG 110 (solitomab). These BiTE[®] antibody constructs are administered via continuous intravenous infusion. For example, blinatumomab is administered in B acute lymphoblastic leukemia as 4-week infusing with a lower initial dose in the 1st week and a higher dose in the remaining treatment for the 1st
30 cycle and in all other cycles from start. Before starting a second cycle, there is a treatment-free period of two weeks. A similar administration schema has been used for solitomab which was administered as continuous intravenous infusion over at least 28 days with increasing doses and also a treatment-free period of two weeks between two cycles.

An important further development of the first generation of BiTE[®] antibody constructs was the
35 provision of bispecific antibody constructs binding to a context independent epitope at the N-terminus of the CD3 ϵ chain of human and *Callithrix jacchus*, *Saguinus oedipus* or *Saimiri sciureus* (WO 2008/119567). Also in this application bispecific antibodies additionally binding

to CD33 are disclosed. CD33 is a sialic-acid-dependent cytoadhesion molecule known as a myeloid differentiation antigen found inter alia on AML blasts in most patients and, perhaps, leukemic stem cells in some. Therefore, CD33 has been identified as a promising marker for myeloid leukemia and a target molecule in the treatment of such diseases.

5 Mylotarg[®] (gemtuzumab ozogamcin), a cytotoxic antibiotic linked to a recombinant monoclonal antibody directed against the CD33 antigen present on leukemic myeloblasts, had been approved in the United States for patients with AML through accelerated approval. However, following the drug's failure to demonstrate clinical benefit in the confirmatory trial, and an increased risk of venoocclusive disease observed in the postmarketing setting, the
10 drug was withdrawn voluntarily by the manufacturer from the United States market. Frequently reported toxicities observed with gemtuzumab ozogamcin included neutropenia and thrombocytopenia, and less frequently reported toxicities included events related to acute infusion-related reactions (anaphylaxis), hepatotoxicity, and veno-occlusive disease.

15 **Definitions**

It must be noted that as used herein, the singular forms "a", "an", and "the", include plural references unless the context clearly indicates otherwise. Thus, for example, reference to "a reagent" includes one or more of such different reagents and reference to "the method" includes reference to equivalent steps and methods known to those of ordinary skill in the art
20 that could be modified or substituted for the methods described herein.

Unless otherwise indicated, the term "at least" preceding a series of elements is to be understood to refer to every element in the series. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the
25 specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the present invention.

The term "and/or" wherever used herein includes the meaning of "and", "or" and "all or any other combination of the elements connected by said term".

30

The term "about" or "approximately" as used herein means within $\pm 20\%$, preferably within $\pm 15\%$, more preferably within $\pm 10\%$, and most preferably within $\pm 5\%$ of a given value or range.

35 Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but

not the exclusion of any other integer or step or group of integer or step. When used herein the term "comprising" can be substituted with the term "containing" or "including" or sometimes when used herein with the term "having".

- 5 When used herein "consisting of" excludes any element, step, or ingredient not specified in the claim element. When used herein, "consisting essentially of" does not exclude materials or steps that do not materially affect the basic and novel characteristics of the claim.

10 In each instance herein any of the terms "comprising", "consisting essentially of" and "consisting of" may be replaced with either of the other two terms.

Detailed Description of the Invention

15 Blinatumomab is given in the treatment of acute lymphoblastic leukemia by continuous intravenous infusion for 4 weeks (5 $\mu\text{g}/\text{m}^2/\text{day}$ for the first week and 15 $\mu\text{g}/\text{m}^2/\text{day}$ thereafter), followed by two treatment-free weeks (one cycle) for up to five cycles. Doing so, the treatment eliminates the compartment of CD19⁺ cells, which is a compartment limited to the B cell lineage.

20 In animal studies testing a CD33 specific BiTE[®], consistent with the proposed mode of action for such molecule, transient myelosuppression was observed, including reductions in circulating neutrophils, platelets, and red cell mass. The decrease in leukocytes, along with the expected increase in activated animal studies, consistent with the proposed mode of action, resulted in transient myelosuppression including reductions in circulating neutrophils, platelets, and red cell mass. The decrease in leukocytes, along with the expected increase in activated T-lymphocytes and increase in cytokine levels were observed in all dose groups.

25 Febrile neutropenia and neutropenia are common events observed in patients with hematological malignancies and prior combination chemotherapies.

Bleeding is a common and potentially serious complication of the treatment of AML, most often secondary to thrombocytopenia. Among bleeding complications, of particular importance is the disseminated intravascular coagulation (DIC) syndrome, due to the massive intravascular activation of blood coagulation with consumption of clotting factors and platelets, leading to severe hemorrhages. In adult patients with AML, 1% of lethal bleedings on day of admission have been observed, all in the presence of hyperleukocytosis or acute promyelocytic leukemia (APL). Recent data in patients with AML show a rate of hemorrhagic death of 9.9%. In addition, in this AML patient population, there may be a strong correlation

30 between unresolved infection in the pancytopenic patient and terminal hemorrhage.

It is also well accepted that immunocompromised patients are susceptible to both common community-acquired and opportunistic infections. Infections are a major cause of morbidity

and mortality in cancer patients and although certain cancers are intrinsically associated with immune compromise, the risk of infection is principally related to the intensity and duration of cytotoxic and immunosuppressive therapy.

- 5 In view of the above the problem underlying the present invention is that in acute myeloid leukemia the situation is different compared to acute lymphoblastic leukemia. The myeloid compartment includes a broader spectrum of cell lineages that are necessary for survival. Therefore, it is not possible to simply transfer the administration scheme of blinatumomab in ALL to a treatment of AML using an AML specific bispecific antibody construct. For an
- 10 efficient treatment of AML using a CD33⁺ cell eliminating therapy approach, the treatment needs to be long enough to be efficacious and short enough to be minimize toxicity on those cell types in the myeloid compartment that are essential for survival. In addition the dose needs to be sufficient for efficacy as well. It has been shown that duration of treatment and higher dose can improve efficacy in ex vivo studies.
- 15 This problem was solved e.g. by providing a bispecific construct comprising a first binding domain specifically binding to CD33 and a second binding domain specifically binding to CD3 (CD33/CD3) for use in a method for the treatment of myeloid leukemia, wherein the construct is administered for a maximal period of 14 days followed by a period of at least 14 days without administration of the construct.
- 20 Using an administration schedule in line with the present invention, preferably applying a step dosing with increasing dose levels, it is possible to efficiently eliminate myeloid leukemic cells during the up to 14 days of CD33/CD3 bispecific construct administration period, while still allowing the patient to recover the myeloic compartment in the period of at least 14 days without administration of the construct. At the same time, the step dosing preferably
- 25 eliminates the risk of severe immunologic side effects such as a cytokine release syndrome. As apparent from **figure 1** the expression of CD33 on the surface of myeloid cells comprising the *common myeloid progenitor* cells, *Myeloblasts*, *Monocytes* has been demonstrated in the literature by flow cytometry. Moreover, CD33 expression on the surface of Macrophages has been demonstrated via immunohistochemistry. Those CD33⁺ cell populations in the myeloid
- 30 compartment are eliminated under treatment of a patient with bispecific constructs described herein. Due to the fact that some of those cell populations are itself the progenitor cells for other cell populations in the myeloid compartment the hematopoiesis of all cell types downstream of the *common myeloid progenitor* cells is affected which results in pancytopenia.
- 35 For a successful treatment of a myeloid leukemia a significant exposure of a patient (i.e. a certain length of exposure) with the bispecific constructs described herein is required to

induce T cell activation/proliferation and cytotoxic activity of those T cells. However, based on the above described observations the longer the administration period of the bispecific constructs lasts, the longer pancytopenia is to be expected. This in mind, the solution to the problem underlying this invention is to balance the length of exposure and the dose of the bispecific constructs which enable the effective elimination of the leukemic cells with an off treatment period during which the myeloid compartment of a patient is allowed to recover. This is reflected by the above described administration scheme.

The end of the period of administration is understood to be reached, when the serum level of the active compound, e.g. the bispecific compound drops under a defined threshold. An example for such threshold is a serum level below an EC_{90} value, preferably below an EC_{50} value, more preferably below an EC_{10} value. Such EC values can be defined in a cytotoxic assay using $CD33^+$ target cells and human PBL as effector cells in line with the assays.

In case of a bispecific single chain antibody construct such as AMG 330 (see SEQ ID NO: 104), which is known to have a short serum half-life (based on the PK parameters shown in appended Example 2, the half-life of AMG 330 in mice is 6.5 to 8.7h, while the predicted half-life of AMG 330 in human is about 2 hours), the serum level would fall below the above discussed threshold value within short time after stopping a continuous iv administration, i.e. almost immediately after the end of the administration phase. In case of a half-life extended bispecific single chain antibody construct the end of the administration phase must be planned in order to ensure a tapering below the threshold in line with the treatment schemata of the invention.

An assay for the determination of a specific EC_x value of a bispecific construct suitable for the present invention is described in the examples herein below.

The term "bispecific construct" refers to a molecule having a structure suitable for the specific binding of two individual target structures. In the context of the present invention such bispecific constructs specifically bind to CD33 on the cell surface of target cells and CD3 on the cell surface of T cells. In a preferred embodiment of a bispecific construct at least one, more preferably both binding domains of the bispecific construct are is/are based on the structure and/or function of an antibody. Such constructs may be designated as "bispecific antibody constructs" in line with the present invention.

The term "antibody construct" refers to a molecule in which the structure and/or function is/are based on the structure and/or function of an antibody, e.g. of a full-length or whole immunoglobulin molecule. An antibody construct is hence capable of binding to its specific target or antigen. Furthermore, an antibody construct according to the invention comprises the minimum structural requirements of an antibody which allow for the target binding. This

minimum requirement may e.g. be defined by the presence of at least the three light chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VL region) and/or the three heavy chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VH region). The antibodies on which the constructs according to the invention are based include for example monoclonal, recombinant, chimeric, deimmunized, humanized and human antibodies.

Within the definition of "antibody constructs" according to the invention are full-length or whole antibodies also including camelid antibodies and other immunoglobulin antibodies generated by biotechnological or protein engineering methods or processes. These full-length antibodies may be for example monoclonal, recombinant, chimeric, deimmunized, humanized and human antibodies. Also within the definition of "antibody constructs" are fragments of full-length antibodies, such as VH, VHH, VL, (s)dAb, Fv, Fd, Fab, Fab', F(ab')₂ or "r IgG" ("half antibody"). Antibody constructs according to the invention may also be modified fragments of antibodies, also called antibody variants, such as scFv, di-scFv or bi(s)-scFv, scFv-Fc, scFv-zipper, scFab, Fab₂, Fab₃, diabodies, single chain diabodies, tandem diabodies (Tandab's), tandem di-scFv, tandem tri-scFv, "minibodies" exemplified by a structure which is as follows: (VH-VL-CH₃)₂, (scFv-CH₃)₂, ((scFv)₂-CH₃ + CH₃), ((scFv)₂-CH₃) or (scFv-CH₃-scFv)₂, multibodies such as triabodies or tetrabodies, and single domain antibodies such as nanobodies or single variable domain antibodies comprising merely one variable domain, which might be VHH, VH or VL, that specifically bind an antigen or epitope independently of other V regions or domains.

A binding domain may typically comprise an antibody light chain variable region (VL) and an antibody heavy chain variable region (VH); however, it does not have to comprise both. Fd fragments, for example, have two VH regions and often retain some antigen-binding function of the intact antigen-binding domain. Additional examples for the format of antibody fragments, antibody variants or binding domains include (1) a Fab fragment, a monovalent fragment having the VL, VH, CL and CH1 domains; (2) a F(ab')₂ fragment, a bivalent fragment having two Fab fragments linked by a disulfide bridge at the hinge region; (3) an Fd fragment having the two VH and CH1 domains; (4) an Fv fragment having the VL and VH domains of a single arm of an antibody, (5) a dAb fragment (Ward et al., (1989) Nature 341 :544-546), which has a VH domain; (6) an isolated complementarity determining region (CDR), and (7) a single chain Fv (scFv), the latter being preferred (for example, derived from an scFv-library). Examples for embodiments of antibody constructs according to the invention are e.g. described in WO 00/006605, WO 2005/040220, WO 2008/119567, WO 2010/037838, WO 2013/026837, WO 2013/026833, US 2014/0308285, US 2014/0302037, WO 2014/144722, WO 2014/151910, and WO 2015/048272.

Furthermore, the definition of the term "antibody constructs" includes monovalent, bivalent and polyvalent/ multivalent constructs and, thus, monospecific constructs, specifically binding to only one antigenic structure, as well as bispecific and polyspecific / multispecific constructs, which specifically bind more than one antigenic structure, e.g. two, three or more, through distinct binding domains. Moreover, the definition of the term "antibody constructs" includes molecules consisting of only one polypeptide chain as well as molecules consisting of more than one polypeptide chain, which chains can be either identical (homodimers, homotrimers or homo oligomers) or different (heterodimer, heterotrimer or heterooligomer).
5
10 Examples for the above identified antibodies and variants or derivatives thereof are described *inter alia* in Harlow and Lane, Antibodies a laboratory manual, CSHL Press (1988) and Using Antibodies: a laboratory manual, CSHL Press (1999), Kontermann and Dübel, Antibody Engineering, Springer, 2nd ed. 2010 and Little, Recombinant Antibodies for Immunotherapy, Cambridge University Press 2009.

15
The antibody constructs of the present invention are preferably "*in vitro* generated antibody constructs". This term refers to an antibody construct according to the above definition where all or part of the variable region (e.g., at least one CDR) is generated in a non-immune cell selection, e.g., an *in vitro* phage display, protein chip or any other method in which candidate
20 sequences can be tested for their ability to bind to an antigen. This term thus preferably excludes sequences generated solely by genomic rearrangement in an immune cell in an animal. A "recombinant antibody" is an antibody made through the use of recombinant DNA technology or genetic engineering.

25 An embodiment of the bispecific antibody construct of the present invention is a "single chain antibody constructs". Those single chain antibody constructs include only above described embodiments of antibody constructs, which consist of a single peptide chain.

The term "monoclonal antibody" (mAb) or monoclonal antibody construct as used herein
30 refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations and/or post-translation modifications (e.g., isomerizations, amidations) that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site or determinant on the antigen, in
35 contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (or epitopes). In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma

culture, hence uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method.

5

For the preparation of monoclonal antibodies, any technique providing antibodies produced by continuous cell line cultures can be used. For example, monoclonal antibodies to be used may be made by the hybridoma method first described by Koehler *et al.*, Nature, 256: 495 (1975), or may be made by recombinant DNA methods (see, e.g., U.S. Patent
10 No. 4,816,567). Examples for further techniques to produce human monoclonal antibodies include the trioma technique, the human B-cell hybridoma technique (Kozbor, Immunology Today 4 (1983), 72) and the EBV-hybridoma technique (Cole *et al.*, Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc. (1985), 77-96).

15 Hybridomas can then be screened using standard methods, such as enzyme-linked immunosorbent assay (ELISA) and surface plasmon resonance (BIAcore™) analysis, to identify one or more hybridomas that produce an antibody that specifically binds with a specified antigen. Any form of the relevant antigen may be used as the immunogen, e.g., recombinant antigen, naturally occurring forms, any variants or fragments thereof, as well as
20 an antigenic peptide thereof. Surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of phage antibodies which bind to an epitope of a target antigen, such as the target cell surface antigen CD33 or CD3 epsilon (Schier, Human Antibodies Hybridomas 7 (1996), 97-105; Malmborg, J. Immunol. Methods 183 (1995), 7-13).

25

Another exemplary method of making monoclonal antibodies includes screening protein expression libraries, e.g., phage display or ribosome display libraries. Phage display is described, for example, in Ladner *et al.*, U.S. Patent No. 5,223,409; Smith (1985) Science 228:1315-1317, Clackson *et al.*, Nature, 352: 624-628 (1991) and Marks *et al.*, J. Mol. Biol.,
30 222: 581-597 (1991).

In addition to the use of display libraries, the relevant antigen can be used to immunize a non-human animal, e.g., a rodent (such as a mouse, hamster, rabbit or rat). In one embodiment, the non-human animal includes at least a part of a human immunoglobulin
35 gene. For example, it is possible to engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig (immunoglobulin) loci. Using the hybridoma technology, antigen-specific monoclonal antibodies derived from the genes with the desired

specificity may be produced and selected. See, e.g., XENOMOUSE™, Green *et al.* (1994) Nature Genetics 7:13-21, US 2003-0070185, WO 96/34096, and WO96/33735.

5 A monoclonal antibody can also be obtained from a non-human animal, and then modified, e.g., humanized, deimmunized, rendered chimeric etc., using recombinant DNA techniques known in the art. Examples of modified antibody constructs include humanized variants of non-human antibodies, "affinity matured" antibodies (see, e.g. Hawkins *et al.* J. Mol. Biol. 254, 889-896 (1992) and Lowman *et al.*, Biochemistry 30, 10832- 10837 (1991)) and antibody mutants with altered effector function(s) (see, e.g., US Patent 5,648,260, 10 Kontermann and Dübel (2010), *loc. cit.* and Little (2009), *loc. cit.*).

In immunology, affinity maturation is the process by which B cells produce antibodies with increased affinity for antigen during the course of an immune response. With repeated 15 exposures to the same antigen, a host will produce antibodies of successively greater affinities. Like the natural prototype, the *in vitro* affinity maturation is based on the principles of mutation and selection. The *in vitro* affinity maturation has successfully been used to optimize antibodies, antibody constructs, and antibody fragments. Random mutations inside the CDRs are introduced using radiation, chemical mutagens or error-prone PCR. In addition, the genetic diversity can be increased by chain shuffling. Two or three rounds of 20 mutation and selection using display methods like phage display usually results in antibody fragments with affinities in the low nanomolar range.

A preferred type of an amino acid substitutional variation of the antibody constructs involves substituting one or more hypervariable region residues of a parent antibody (e. g. a 25 humanized or human antibody). Generally, the resulting variant(s) selected for further development will have improved biological properties relative to the parent antibody from which they are generated. A convenient way for generating such substitutional variants involves affinity maturation using phage display. Briefly, several hypervariable region sites (e. g. 6-7 sites) are mutated to generate all possible amino acid substitutions at each site. The 30 antibody variants thus generated are displayed in a monovalent fashion from filamentous phage particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed variants are then screened for their biological activity (e. g. binding affinity) as herein disclosed. In order to identify candidate hypervariable region sites for modification, alanine scanning mutagenesis can be performed to identify hypervariable region residues 35 contributing significantly to antigen binding. Alternatively, or additionally, it may be beneficial to analyze a crystal structure of the antigen-antibody complex to identify contact points between the binding domain and, e.g., human the target cell surface antigen CD33. Such

contact residues and neighboring residues are candidates for substitution according to the techniques elaborated herein. Once such variants are generated, the panel of variants is subjected to screening as described herein and antibodies with superior properties in one or more relevant assays may be selected for further development.

5

The monoclonal antibodies and antibody constructs of the present invention specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is/are identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816, 567; Morrison et al., Proc. Natl. Acad. Sci. USA, 81: 6851-6855 (1984)). Chimeric antibodies of interest herein include "primitized" antibodies comprising variable domain antigen-binding sequences derived from a non-human primate (e.g., Old World Monkey, Ape etc.) and human constant region sequences. A variety of approaches for making chimeric antibodies have been described. See e.g., Morrison *et al.*, Proc. Natl. Acad. ScL U.S.A. 81:6851 , 1985; Takeda *et al.*, Nature 314:452, 1985, Cabilly *et al.*, U.S. Patent No. 4,816,567; Boss *et al.*, U.S. Patent No. 4,816,397; Tanaguchi *et al.*, EP 0171496; EP 0173494; and GB 2177096.

An antibody, antibody construct or antibody fragment may also be modified by specific deletion of human T cell epitopes (a method called "deimmunization") by the methods disclosed in WO 98/52976 and WO 00/34317. Briefly, the heavy and light chain variable domains of an antibody can be analyzed for peptides that bind to MHC class II; these peptides represent potential T cell epitopes (as defined in WO 98/52976 and WO 00/34317). For detection of potential T cell epitopes, a computer modeling approach termed "peptide threading" can be applied, and in addition a database of human MHC class II binding peptides can be searched for motifs present in the VH and VL sequences, as described in WO 98/52976 and WO 00/34317. These motifs bind to any of the 18 major MHC class II DR allotypes, and thus constitute potential T cell epitopes. Potential T cell epitopes detected can be eliminated by substituting small numbers of amino acid residues in the variable domains, or preferably, by single amino acid substitutions. Typically, conservative substitutions are made. Often, but not exclusively, an amino acid common to a position in human germline antibody sequences may be used. Human germline sequences are disclosed e.g. in Tomlinson, *et al.* (1992) J. Mol. Biol. 227:776-798; Cook, G.P. *et al.* (1995) Immunol. Today Vol. 16 (5): 237-242; and Tomlinson et al. (1995) EMBO J. 14: 14:4628-4638. The V BASE

directory provides a comprehensive directory of human immunoglobulin variable region sequences (compiled by Tomlinson, LA. et al. MRC Centre for Protein Engineering, Cambridge, UK). These sequences can be used as a source of human sequence, e.g., for framework regions and CDRs. Consensus human framework regions can also be used, for example as described in US Patent No. 6,300,064.

"Humanized" antibodies, antibody constructs or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) are antibodies or immunoglobulins of mostly human sequences, which contain (a) minimal sequence(s) derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region (also CDR) of the recipient are replaced by residues from a hypervariable region of a non-human (e.g., rodent) species (donor antibody) such as mouse, rat, hamster or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, "humanized antibodies" as used herein may also comprise residues which are found neither in the recipient antibody nor the donor antibody. These modifications are made to further refine and optimize antibody performance. The humanized antibody may also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, Nature, 321: 522-525 (1986); Reichmann *et al.*, Nature, 332: 323-329 (1988); and Presta, Curr. Op. Struct. Biol., 2: 593-596 (1992).

Humanized antibodies or fragments thereof can be generated by replacing sequences of the Fv variable domain that are not directly involved in antigen binding with equivalent sequences from human Fv variable domains. Exemplary methods for generating humanized antibodies or fragments thereof are provided by Morrison (1985) Science 229:1202-1207; by Oi *et al.* (1986) BioTechniques 4:214; and by US 5,585,089; US 5,693,761; US 5,693,762; US 5,859,205; and US 6,407,213. Those methods include isolating, manipulating, and expressing the nucleic acid sequences that encode all or part of immunoglobulin Fv variable domains from at least one of a heavy or light chain. Such nucleic acids may be obtained from a hybridoma producing an antibody against a predetermined target, as described above, as well as from other sources. The recombinant DNA encoding the humanized antibody molecule can then be cloned into an appropriate expression vector.

Humanized antibodies may also be produced using transgenic animals such as mice that express human heavy and light chain genes, but are incapable of expressing the

endogenous mouse immunoglobulin heavy and light chain genes. Winter describes an exemplary CDR grafting method that may be used to prepare the humanized antibodies described herein (U.S. Patent No. 5,225,539). All of the CDRs of a particular human antibody may be replaced with at least a portion of a non-human CDR, or only some of the CDRs may be replaced with non-human CDRs. It is only necessary to replace the number of CDRs required for binding of the humanized antibody to a predetermined antigen.

A humanized antibody can be optimized by the introduction of conservative substitutions, consensus sequence substitutions, germline substitutions and/or back mutations. Such altered immunoglobulin molecules can be made by any of several techniques known in the art, (e.g., Teng *et al.*, Proc. Natl. Acad. Sci. U.S.A., 80: 7308-7312, 1983; Kozbor *et al.*, Immunology Today, 4: 7279, 1983; Olsson *et al.*, Meth. Enzymol., 92: 3-16, 1982, and EP 239 400.

The term "human antibody", "human antibody construct" and "human binding domain" includes antibodies, antibody constructs and binding domains having antibody regions such as variable and constant regions or domains which correspond substantially to human germline immunoglobulin sequences known in the art, including, for example, those described by Kabat *et al.* (1991) (*loc. cit.*). The human antibodies, antibody constructs or binding domains of the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*), for example in the CDRs, and in particular, in CDR3. The human antibodies, antibody constructs or binding domains can have at least one, two, three, four, five, or more positions replaced with an amino acid residue that is not encoded by the human germline immunoglobulin sequence. The definition of human antibodies, antibody constructs and binding domains as used herein also contemplates fully human antibodies, which include only non-artificially and/or genetically altered human sequences of antibodies as those can be derived by using technologies or systems such as the Xenomouse.

In some embodiments, the antibody constructs of the invention are "isolated" or "substantially pure" antibody constructs. "Isolated" or "substantially pure" when used to describe the antibody construct disclosed herein means an antibody construct that has been identified, separated and/or recovered from a component of its production environment. Preferably, the antibody construct is free or substantially free of association with all other components from its production environment. Contaminant components of its production environment, such as that resulting from recombinant transfected cells, are materials that would typically interfere

with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. The antibody constructs may e.g. constitute at least about 5%, or at least about 50% by weight of the total protein in a given sample. It is understood that the isolated protein may constitute from 5% to 99.9% by weight of the total protein content, depending on the circumstances. The polypeptide may be made at a significantly higher concentration through the use of an inducible promoter or high expression promoter, such that it is made at increased concentration levels. The definition includes the production of an antibody construct in a wide variety of organisms and/or host cells that are known in the art. In preferred embodiments, the antibody construct will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Ordinarily, however, an isolated antibody construct will be prepared by at least one purification step.

The term "binding domain" characterizes in connection with the present invention a domain which (specifically) binds to / interacts with / recognizes a given target epitope or a given target site on the target molecules (antigens) and CD3, respectively. The structure and function of the first binding domain (recognizing the target cell surface antigen CD33), and preferably also the structure and/or function of the second binding domain (CD3), is/are based on the structure and/or function of an antibody, e.g. of a full-length or whole immunoglobulin molecule. According to the invention, the first binding domain is characterized by the presence of three light chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VL region) and three heavy chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VH region). The second binding domain preferably also comprises the minimum structural requirements of an antibody which allow for the target binding. More preferably, the second binding domain comprises at least three light chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VL region) and/or three heavy chain CDRs (i.e. CDR1, CDR2 and CDR3 of the VH region). It is envisaged that the first and/or second binding domain is produced by or obtainable by phage-display or library screening methods rather than by grafting CDR sequences from a pre-existing (monoclonal) antibody into a scaffold.

According to the present invention, binding domains are preferably in the form of polypeptides. Such polypeptides may include proteinaceous parts and non-proteinaceous parts (e.g. chemical linkers or chemical cross-linking agents such as glutaraldehyde). Proteins (including fragments thereof, preferably biologically active fragments, and peptides, usually having less than 30 amino acids) comprise two or more amino acids coupled to each

other via a covalent peptide bond (resulting in a chain of amino acids). The term "polypeptide" as used herein describes a group of molecules, which usually consist of more than 30 amino acids. Polypeptides may further form multimers such as dimers, trimers and higher oligomers, i.e. consisting of more than one polypeptide molecule. Polypeptide molecules forming such dimers, trimers etc. may be identical or non-identical. The corresponding higher order structures of such multimers are, consequently, termed homo- or heterodimers, homo- or heterotrimers etc. An example for a heteromultimer is an antibody molecule, which, in its naturally occurring form, consists of two identical light polypeptide chains and two identical heavy polypeptide chains. The terms "peptide", "polypeptide" and "protein" also refer to naturally modified peptides / polypeptides / proteins wherein the modification is effected e.g. by post-translational modifications like glycosylation, acetylation, phosphorylation and the like. A "peptide", "polypeptide" or "protein" when referred to herein may also be chemically modified such as pegylated. Such modifications are well known in the art and described herein below.

Antibodies and antibody constructs comprising at least one human binding domain avoid some of the problems associated with antibodies or antibody constructs that possess non-human such as rodent (e.g. murine, rat, hamster or rabbit) variable and/or constant regions. The presence of such rodent derived proteins can lead to the rapid clearance of the antibodies or antibody constructs or can lead to the generation of an immune response against the antibody or antibody construct by a patient. In order to avoid the use of rodent derived antibodies or antibody constructs, human or fully human antibodies / antibody constructs can be generated through the introduction of human antibody function into a rodent so that the rodent produces fully human antibodies.

The ability to clone and reconstruct megabase-sized human loci in YACs and to introduce them into the mouse germline provides a powerful approach to elucidating the functional components of very large or crudely mapped loci as well as generating useful models of human disease. Furthermore, the use of such technology for substitution of mouse loci with their human equivalents could provide unique insights into the expression and regulation of human gene products during development, their communication with other systems, and their involvement in disease induction and progression.

An important practical application of such a strategy is the "humanization" of the mouse humoral immune system. Introduction of human immunoglobulin (Ig) loci into mice in which the endogenous Ig genes have been inactivated offers the opportunity to study the mechanisms underlying programmed expression and assembly of antibodies as well as their

role in B-cell development. Furthermore, such a strategy could provide an ideal source for production of fully human monoclonal antibodies (mAbs) – an important milestone towards fulfilling the promise of antibody therapy in human disease. Fully human antibodies or antibody constructs are expected to minimize the immunogenic and allergic responses intrinsic to mouse or mouse-derivatized mAbs and thus to increase the efficacy and safety of the administered antibodies / antibody constructs. The use of fully human antibodies or antibody constructs can be expected to provide a substantial advantage in the treatment of chronic and recurring human diseases, such as inflammation, autoimmunity, and cancer, which require repeated compound administrations.

10

One approach towards this goal was to engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig loci in anticipation that such mice would produce a large repertoire of human antibodies in the absence of mouse antibodies. Large human Ig fragments would preserve the large variable gene diversity as well as the proper regulation of antibody production and expression. By exploiting the mouse machinery for antibody diversification and selection and the lack of immunological tolerance to human proteins, the reproduced human antibody repertoire in these mouse strains should yield high affinity antibodies against any antigen of interest, including human antigens. Using the hybridoma technology, antigen-specific human mAbs with the desired specificity could be readily produced and selected. This general strategy was demonstrated in connection with the generation of the first XenoMouse mouse strains (see Green et al. Nature Genetics 7:13-21 (1994)). The XenoMouse strains were engineered with yeast artificial chromosomes (YACs) containing 245 kb and 190 kb-sized germline configuration fragments of the human heavy chain locus and kappa light chain locus, respectively, which contained core variable and constant region sequences. The human Ig containing YACs proved to be compatible with the mouse system for both rearrangement and expression of antibodies and were capable of substituting for the inactivated mouse Ig genes. This was demonstrated by their ability to induce B cell development, to produce an adult-like human repertoire of fully human antibodies, and to generate antigen-specific human mAbs. These results also suggested that introduction of larger portions of the human Ig loci containing greater numbers of V genes, additional regulatory elements, and human Ig constant regions might recapitulate substantially the full repertoire that is characteristic of the human humoral response to infection and immunization. The work of Green et al. was recently extended to the introduction of greater than approximately 80% of the human antibody repertoire through introduction of megabase sized, germline configuration YAC fragments of the human heavy chain loci and kappa light chain loci, respectively. See Mendez *et al.* Nature Genetics 15:146-156 (1997) and U.S. patent application Ser. No. 08/759,620.

35

The production of the XenoMouse mice is further discussed and delineated in U.S. patent applications Ser. No. 07/466,008, Ser. No. 07/610,515, Ser. No. 07/919,297, Ser. No. 07/922,649, Ser. No. 08/031,801, Ser. No. 08/112,848, Ser. No. 08/234,145, 5 Ser. No. 08/376,279, Ser. No. 08/430,938, Ser. No. 08/464,584, Ser. No. 08/464,582, Ser. No. 08/463,191, Ser. No. 08/462,837, Ser. No. 08/486,853, Ser. No. 08/486,857, Ser. No. 08/486,859, Ser. No. 08/462,513, Ser. No. 08/724,752, and Ser. No. 08/759,620; and U.S. Pat. Nos. 6,162,963, 6,150,584, 6,114,598, 6,075,181, and 5,939,598 and Japanese Patent Nos. 3 068 180 B2, 3 068 506 B2, and 3 068 507 B2. See also Mendez *et al.* Nature Genetics 15:146-156 (1997) and Green and Jakobovits J. Exp. Med. 188:483-495 10 (1998), EP 0 463 151 B1, WO 94/02602, WO 96/34096, WO 98/24893, WO 00/76310, and WO 03/47336.

In an alternative approach, others, including GenPharm International, Inc., have utilized a 15 "minilocus" approach. In the minilocus approach, an exogenous Ig locus is mimicked through the inclusion of pieces (individual genes) from the Ig locus. Thus, one or more VH genes, one or more DH genes, one or more JH genes, a mu constant region, and a second constant region (preferably a gamma constant region) are formed into a construct for insertion into an animal. This approach is described in U.S. Pat. No. 5,545,807 to Surani *et al.* and 20 U.S. Pat. Nos. 5,545,806, 5,625,825, 5,625,126, 5,633,425, 5,661,016, 5,770,429, 5,789,650, 5,814,318, 5,877,397, 5,874,299, and 6,255,458 each to Lonberg and Kay, U.S. Pat. Nos. 5,591,669 and 6,023,010 to Krimpenfort and Berns, U.S. Pat. Nos. 5,612,205, 5,721,367, and 5,789,215 to Berns *et al.*, and U.S. Pat. No. 5,643,763 to Choi and Dunn, and GenPharm International U.S. patent application Ser. No. 07/574,748, Ser. No. 07/575,962, 25 Ser. No. 07/810,279, Ser. No. 07/853,408, Ser. No. 07/904,068, Ser. No. 07/990,860, Ser. No. 08/053,131, Ser. No. 08/096,762, Ser. No. 08/155,301, Ser. No. 08/161,739, Ser. No. 08/165,699, Ser. No. 08/209,741. See also EP 0 546 073 B1, WO 92/03918, WO 92/22645, WO 92/22647, WO 92/22670, WO 93/12227, WO 94/00569, WO 94/25585, WO 96/14436, WO 97/13852, and WO 98/24884 and U.S. Pat. No. 5,981,175. See further 30 Taylor *et al.* (1992), Chen *et al.* (1993), Tuailon *et al.* (1993), Choi *et al.* (1993), Lonberg *et al.* (1994), Taylor *et al.* (1994), and Tuailon *et al.* (1995), Fishwild *et al.* (1996).

Kirin has also demonstrated the generation of human antibodies from mice in which, through microcell fusion, large pieces of chromosomes, or entire chromosomes, have been 35 introduced. See European Patent Application Nos. 773 288 and 843 961. Xenerex Biosciences is developing a technology for the potential generation of human antibodies. In this technology, SCID mice are reconstituted with human lymphatic cells, e.g., B and/or

T cells. Mice are then immunized with an antigen and can generate an immune response against the antigen. See U.S. Pat. Nos. 5,476,996; 5,698,767; and 5,958,765.

Human anti-mouse antibody (HAMA) responses have led the industry to prepare chimeric or otherwise humanized antibodies. It is however expected that certain human anti-chimeric antibody (HACA) responses will be observed, particularly in chronic or multi-dose utilizations of the antibody. Thus, it would be desirable to provide antibody constructs comprising a fully human binding domain against the target cell surface antigen and a fully human binding domain against CD3 in order to vitiate concerns and/or effects of HAMA or HACA response.

The terms “(specifically) binds to”, “(specifically) recognizes”, “is (specifically) directed to”, and “(specifically) reacts with” mean in accordance with this invention that a binding domain interacts or specifically interacts with one or more, preferably at least two, more preferably at least three and most preferably at least four amino acids of an epitope located on the target protein or antigen (the target cell surface antigen CD33 / CD3).

The term “epitope” refers to the site on an antigen to which a binding domain, such as an antibody or immunoglobulin or derivative or fragment of an antibody or of an immunoglobulin, specifically binds. An “epitope” is antigenic and thus the term epitope is sometimes also referred to herein as “antigenic structure” or “antigenic determinant”. Thus, the binding domain is an “antigen interaction site”. Said binding/interaction is also understood to define a “specific recognition”. The term “epitope” is understood in connection with this application as describing the complete antigenic structure, whereas the term “part of the epitope” may be used to describe one or more subgroups of the specific epitope of a given binding domain.

“Epitopes” can be formed both by contiguous amino acids or non-contiguous amino acids juxtaposed by tertiary folding of a protein. A “linear epitope” is an epitope where an amino acid primary sequence comprises the recognized epitope. A linear epitope typically includes at least 3 or at least 4, and more usually, at least 5 or at least 6 or at least 7, for example, about 8 to about 10 amino acids in a unique sequence.

A “conformational epitope”, in contrast to a linear epitope, is an epitope wherein the primary sequence of the amino acids comprising the epitope is not the sole defining component of the epitope recognized (e.g., an epitope wherein the primary sequence of amino acids is not necessarily recognized by the binding domain). Typically a conformational epitope comprises an increased number of amino acids relative to a linear epitope. With regard to recognition of conformational epitopes, the binding domain recognizes a three-dimensional structure of the

antigen, preferably a peptide or protein or fragment thereof (in the context of the present invention, the antigen for one of the binding domains is comprised within the target cell surface antigen CD33). For example, when a protein molecule folds to form a three-dimensional structure, certain amino acids and/or the polypeptide backbone forming the conformational epitope become juxtaposed enabling the antibody to recognize the epitope. Methods of determining the conformation of epitopes include, but are not limited to, x-ray crystallography, two-dimensional nuclear magnetic resonance (2D-NMR) spectroscopy and site-directed spin labelling and electron paramagnetic resonance (EPR) spectroscopy.

The interaction between the binding domain and the epitope or epitope cluster implies that a binding domain exhibits appreciable affinity for the epitope or epitope cluster on a particular protein or antigen (here: the target cell surface antigen CD33 and CD3, respectively) and, generally, does not exhibit significant reactivity with proteins or antigens other than the target cell surface antigen CD33 or CD3. "Appreciable affinity" includes binding with an affinity of about 10^{-6} M (KD) or stronger. Preferably, binding is considered specific when the binding affinity is about 10^{-12} to 10^{-8} M, 10^{-12} to 10^{-9} M, 10^{-12} to 10^{-10} M, 10^{-11} to 10^{-8} M, preferably of about 10^{-11} to 10^{-9} M. Whether a binding domain specifically reacts with or binds to a target can be tested readily by, *inter alia*, comparing the reaction of said binding domain with a target protein or antigen with the reaction of said binding domain with proteins or antigens other than the target cell surface antigen CD33 or CD3. Preferably, a binding domain of the invention does not essentially or substantially bind to proteins or antigens other than the target cell surface antigen CD33 or CD3 (*i.e.*, the first binding domain is not capable of binding to proteins other than the target cell surface antigen CD33 and the second binding domain is not capable of binding to proteins other than CD3).

The term "does not essentially / substantially bind" or "is not capable of binding" means that a binding domain of the present invention does not bind a protein or antigen other than the target cell surface antigen CD33 or CD3, *i.e.*, does not show reactivity of more than 30%, preferably not more than 20%, more preferably not more than 10%, particularly preferably not more than 9%, 8%, 7%, 6% or 5% with proteins or antigens other than the target cell surface antigen CD33 or CD3, whereby binding to the target cell surface antigen CD33 or CD3, respectively, is set to be 100%.

Specific binding is believed to be effected by specific motifs in the amino acid sequence of the binding domain and the antigen. Thus, binding is achieved as a result of their primary, secondary and/or tertiary structure as well as the result of secondary modifications of said structures. The specific interaction of the antigen-interaction-site with its specific antigen may

result in a simple binding of said site to the antigen. Moreover, the specific interaction of the antigen-interaction-site with its specific antigen may alternatively or additionally result in the initiation of a signal, e.g. due to the induction of a change of the conformation of the antigen, an oligomerization of the antigen, etc.

5

The term "variable" refers to the portions of the antibody or immunoglobulin domains that exhibit variability in their sequence and that are involved in determining the specificity and binding affinity of a particular antibody (i.e., the "variable domain(s)"). The pairing of a variable heavy chain (VH) and a variable light chain (VL) together forms a single antigen-binding site.

10

Variability is not evenly distributed throughout the variable domains of antibodies; it is concentrated in sub-domains of each of the heavy and light chain variable regions. These sub-domains are called "hypervariable regions" or "complementarity determining regions" (CDRs). The more conserved (i.e., non-hypervariable) portions of the variable domains are called the "framework" regions (FRM) and provide a scaffold for the six CDRs in three dimensional space to form an antigen-binding surface. The variable domains of naturally occurring heavy and light chains each comprise four FRM regions (FR1, FR2, FR3, and FR4), largely adopting a β -sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases forming part of, the β -sheet structure. The hypervariable regions in each chain are held together in close proximity by the FRM and, with the hypervariable regions from the other chain, contribute to the formation of the antigen-binding site (see Kabat *et al.*, *loc. cit.*).

15

20

25

The terms "CDR", and its plural "CDRs", refer to the complementarity determining region of which three make up the binding character of a light chain variable region (CDR-L1, CDR-L2 and CDR-L3) and three make up the binding character of a heavy chain variable region (CDR-H1, CDR-H2 and CDR-H3). CDRs contain most of the residues responsible for specific interactions of the antibody with the antigen and hence contribute to the functional activity of an antibody molecule: they are the main determinants of antigen specificity.

30

The exact definitional CDR boundaries and lengths are subject to different classification and numbering systems. CDRs may therefore be referred to by Kabat, Chothia, contact or any other boundary definitions, including the numbering system described herein. Despite differing boundaries, each of these systems has some degree of overlap in what constitutes the so called "hypervariable regions" within the variable sequences. CDR definitions according to these systems may therefore differ in length and boundary areas with respect to

35

the adjacent framework region. See for example Kabat (an approach based on cross-species sequence variability), Chothia (an approach based on crystallographic studies of antigen-antibody complexes), and/or MacCallum (Kabat *et al.*, *loc. cit.*; Chothia *et al.*, J. Mol. Biol, 1987, 196: 901-917; and MacCallum *et al.*, J. Mol. Biol, 1996, 262: 732). Still another
5 standard for characterizing the antigen binding site is the AbM definition used by Oxford Molecular's AbM antibody modeling software. See, *e.g.*, Protein Sequence and Structure Analysis of Antibody Variable Domains. In: Antibody Engineering Lab Manual (Ed.: Duebel, S. and Kontermann, R., Springer-Verlag, Heidelberg). To the extent that two residue
10 identification techniques define regions of overlapping, but not identical regions, they can be combined to define a hybrid CDR. However, the numbering in accordance with the so-called Kabat system is preferred.

Typically, CDRs form a loop structure that can be classified as a canonical structure. The term "canonical structure" refers to the main chain conformation that is adopted by the
15 antigen binding (CDR) loops. From comparative structural studies, it has been found that five of the six antigen binding loops have only a limited repertoire of available conformations. Each canonical structure can be characterized by the torsion angles of the polypeptide backbone. Correspondent loops between antibodies may, therefore, have very similar three
20 dimensional structures, despite high amino acid sequence variability in most parts of the loops (Chothia and Lesk, J. Mol. Biol., 1987, 196: 901; Chothia *et al.*, Nature, 1989, 342: 877; Martin and Thornton, J. Mol. Biol, 1996, 263: 800). Furthermore, there is a relationship between the adopted loop structure and the amino acid sequences surrounding it. The conformation of a particular canonical class is determined by the length of the loop and the amino acid residues residing at key positions within the loop, as well as within the conserved
25 framework (*i.e.*, outside of the loop). Assignment to a particular canonical class can therefore be made based on the presence of these key amino acid residues.

The term "canonical structure" may also include considerations as to the linear sequence of the antibody, for example, as catalogued by Kabat (Kabat *et al.*, *loc. cit.*). The Kabat
30 numbering scheme (system) is a widely adopted standard for numbering the amino acid residues of an antibody variable domain in a consistent manner and is the preferred scheme applied in the present invention as also mentioned elsewhere herein. Additional structural considerations can also be used to determine the canonical structure of an antibody. For example, those differences not fully reflected by Kabat numbering can be described by the
35 numbering system of Chothia *et al.* and/or revealed by other techniques, for example, crystallography and two- or three-dimensional computational modeling. Accordingly, a given antibody sequence may be placed into a canonical class which allows for, among other

things, identifying appropriate chassis sequences (e.g., based on a desire to include a variety of canonical structures in a library). Kabat numbering of antibody amino acid sequences and structural considerations as described by Chothia *et al.*, loc. cit. and their implications for construing canonical aspects of antibody structure, are described in the literature. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known in the art. For a review of the antibody structure, see Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, eds. Harlow *et al.*, 1988.

The CDR3 of the light chain and, particularly, the CDR3 of the heavy chain may constitute the most important determinants in antigen binding within the light and heavy chain variable regions. In some antibody constructs, the heavy chain CDR3 appears to constitute the major area of contact between the antigen and the antibody. *In vitro* selection schemes in which CDR3 alone is varied can be used to vary the binding properties of an antibody or determine which residues contribute to the binding of an antigen. Hence, CDR3 is typically the greatest source of molecular diversity within the antibody-binding site. H3, for example, can be as short as two amino acid residues or greater than 26 amino acids.

In a classical full-length antibody or immunoglobulin, each light (L) chain is linked to a heavy (H) chain by one covalent disulfide bond, while the two H chains are linked to each other by one or more disulfide bonds depending on the H chain isotype. The CH domain most proximal to VH is usually designated as CH1. The constant ("C") domains are not directly involved in antigen binding, but exhibit various effector functions, such as antibody-dependent, cell-mediated cytotoxicity and complement activation. The Fc region of an antibody is comprised within the heavy chain constant domains and is for example able to interact with cell surface located Fc receptors.

The sequence of antibody genes after assembly and somatic mutation is highly varied, and these varied genes are estimated to encode 10^{10} different antibody molecules (Immunoglobulin Genes, 2nd ed., eds. Jonio *et al.*, Academic Press, San Diego, CA, 1995). Accordingly, the immune system provides a repertoire of immunoglobulins. The term "repertoire" refers to at least one nucleotide sequence derived wholly or partially from at least one sequence encoding at least one immunoglobulin. The sequence(s) may be generated by rearrangement *in vivo* of the V, D, and J segments of heavy chains, and the V and J segments of light chains. Alternatively, the sequence(s) can be generated from a cell in response to which rearrangement occurs, e.g., *in vitro* stimulation. Alternatively, part or all of the sequence(s) may be obtained by DNA splicing, nucleotide synthesis, mutagenesis, and other methods, see, e.g., U.S. Patent 5,565,332. A repertoire may include only one

sequence or may include a plurality of sequences, including ones in a genetically diverse collection.

5 The term "bispecific" as used herein refers to a construct which is "at least bispecific", i.e., it comprises at least a first binding domain and a second binding domain, wherein the first binding domain binds to one antigen or target, and the second binding domain binds to another antigen or target (here: CD3). Accordingly, bispecific constructs according to the invention comprise specificities for at least two different antigens or targets. The term "bispecific construct" of the invention also encompasses multispecific constructs such as trispecific constructs, the latter ones including three binding domains, or constructs having more than three (e.g. four, five...) specificities. In case the construct used in connection with this invention is an antibody construct, these encompassed corresponding constructs are multispecific antibody constructs such as trispecific antibody constructs, the latter ones including three binding domains, or constructs having more than three (e.g. four, five...) specificities.

15 Given that the antibody constructs according to the invention are (at least) bispecific, they do not occur naturally and they are markedly different from naturally occurring products. A "bispecific" antibody construct or immunoglobulin is hence an artificial hybrid antibody or immunoglobulin having at least two distinct binding sites with different specificities. Bispecific antibodies can be produced by a variety of methods including fusion of hybridomas or linking of Fab' fragments. See, e.g., Songsvilai & Lachmann, Clin. Exp. Immunol. 79:315-321 (1990).

25 The at least two binding domains and the variable domains of the antibody construct of the present invention may or may not comprise peptide linkers (spacer peptides). The term "peptide linker" defines in accordance with the present invention an amino acid sequence by which the amino acid sequences of one (variable and/or binding) domain and another (variable and/or binding) domain of the antibody construct of the invention are linked with each other. An essential technical feature of such peptide linker is that said peptide linker does not comprise any polymerization activity. Among the suitable peptide linkers are those described in U.S. Patents 4,751,180 and 4,935,233 or WO 88/09344. The peptide linkers can also be used to attach other domains or modules or regions (such as half-life extending domains) to the antibody construct of the invention.

35 In the event that a linker is used, this linker is preferably of a length and sequence sufficient to ensure that each of the first and second domains can, independently from one another,

retain their differential binding specificities. For peptide linkers which connect the at least two binding domains in the antibody construct of the invention (or two variable domains), those peptide linkers are preferred which comprise only a few number of amino acid residues, e.g. 12 amino acid residues or less. Thus, peptide linker of 12, 11, 10, 9, 8, 7, 6 or 5 amino acid residues are preferred. An envisaged peptide linker with less than 5 amino acids comprises 4, 3, 2 or one amino acid(s) wherein Gly-rich linkers are preferred. A particularly preferred "single" amino acid in context of said "peptide linker" is Gly. Accordingly, said peptide linker may consist of the single amino acid Gly. Another preferred embodiment of a peptide linker is characterized by the amino acid sequence Gly-Gly-Gly-Gly-Ser, i.e. Gly₄Ser, or polymers thereof, i.e. (Gly₄Ser)_x, where x is an integer of 1 or greater. The characteristics of said peptide linker, which comprise the absence of the promotion of secondary structures are known in the art and are described e.g. in Dall'Acqua et al. (Biochem. (1998) 37, 9266-9273), Cheadle et al. (Mol Immunol (1992) 29, 21-30) and Raag and Whitlow (FASEB (1995) 9(1), 73-80). Peptide linkers which also do not promote any secondary structures are preferred.

The linkage of said domains to each other can be provided by, e.g. genetic engineering, as described in the examples. Methods for preparing fused and operatively linked bispecific single chain constructs and expressing them in mammalian cells or bacteria are well-known in the art (e.g. WO 99/54440 or Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 2001).

Bispecific single chain molecules are known in the art and are described in WO 99/54440, Mack, J. Immunol. (1997), 158, 3965-3970, Mack, PNAS, (1995), 92, 7021-7025, Kufer, Cancer Immunol. Immunother., (1997), 45, 193-197, Löffler, Blood, (2000), 95, 6, 2098-2103, Brühl, Immunol., (2001), 166, 2420-2426, Kipriyanov, J. Mol. Biol., (1999), 293, 41-56.

Techniques described for the production of single chain antibodies (see, *inter alia*, US Patent 4,946,778, Kontermann and Dübel (2010), *loc. cit.* and Little (2009), *loc. cit.*) can be adapted to produce single chain antibody constructs specifically recognizing (an) elected target(s).

Bivalent (also called divalent) or bispecific single-chain variable fragments (bi-scFvs or di-scFvs having the format (scFv)₂) can be engineered by linking two scFv molecules. In case these two scFv molecules have the same binding specificity, the resulting (scFv)₂ molecule will preferably be called bivalent (*i.e.* it has two valences for the same target epitope). In case the two scFv molecules have different binding specificities, the resulting (scFv)₂ molecule will preferably be called bispecific. The linking can be done by producing a single peptide chain with two VH regions and two VL regions, yielding tandem scFvs (see e.g. Kufer P. *et al.*, (2004) Trends in Biotechnology 22(5):238-244). Another possibility is the creation of scFv molecules with linker peptides that are too short for the two variable regions to fold together

(e.g. about five amino acids), forcing the scFvs to dimerize. This type is known as diabodies (see e.g. Hollinger, Philipp *et al.*, (July 1993) Proceedings of the National Academy of Sciences of the United States of America 90 (14): 6444-8.).

5 Single domain antibodies comprise merely one (monomeric) antibody variable domain which is able to bind selectively to a specific antigen, independently of other V regions or domains. The first single domain antibodies were engineered from heavy chain antibodies found in camelids, and these are called V_{HH} fragments. Cartilaginous fishes also have heavy chain antibodies (IgNAR) from which single domain antibodies called V_{NAR} fragments can be
10 obtained. An alternative approach is to split the dimeric variable domains from common immunoglobulins e.g. from humans or rodents into monomers, hence obtaining VH or VL as a single domain Ab. Although most research into single domain antibodies is currently based on heavy chain variable domains, nanobodies derived from light chains have also been shown to bind specifically to target epitopes. Examples of single domain antibodies are
15 called sdAb, nanobodies or single variable domain antibodies.

A (single domain mAb)₂ is hence a monoclonal antibody construct composed of (at least) two single domain monoclonal antibodies, which are individually selected from the group comprising VH, VL, VHH and V_{NAR} . The linker is preferably in the form of a peptide linker.
20 Similarly, an "scFv-single domain mAb" is a monoclonal antibody construct composed of at least one single domain antibody as described above and one scFv molecule as described above. Again, the linker is preferably in the form of a peptide linker.

It is also envisaged that the antibody construct of the invention has, in addition to its function
25 to bind to the target antigen CD33 and CD3, a further function. In this format, the antibody construct is a trifunctional or multifunctional antibody construct by targeting target cells through binding to the target antigen, mediating cytotoxic T cell activity through CD3 binding and providing a further function such as a label (fluorescent etc.), a therapeutic agent such as a toxin or radionuclide, etc.

30 Covalent modifications of the antibody constructs are also included within the scope of this invention, and are generally, but not always, done post-translationally. For example, several types of covalent modifications of the antibody construct are introduced into the molecule by reacting specific amino acid residues of the antibody construct with an organic derivatizing
35 agent that is capable of reacting with selected side chains or the N- or C-terminal residues.

Cysteiny l residues most commonly are reacted with α -haloacetates (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteiny l residues also are derivatized by reaction with bromotrifluoroacetone, α -bromo- β -(5-imidozoyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole.

Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1 M sodium cacodylate at pH 6.0. Lysinyl and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysinyl residues. Other suitable reagents for derivatizing alpha-amino-containing residues include imidoesters such as methyl picolinimate; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4-pentanedione; and transaminase-catalyzed reaction with glyoxylate.

Arginyl residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pKa of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

The specific modification of tyrosyl residues may be made, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Tyrosyl residues are iodinated using ^{125}I or ^{131}I to prepare labeled proteins for use in radioimmunoassay, the chloramine T method described above being suitable.

Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides ($\text{R}'\text{—N=C=N—R}'$), where R and R' are optionally different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginyl and glutaminyl residues by reaction with ammonium ions.

Derivatization with bifunctional agents is useful for crosslinking the antibody constructs of the present invention to a water-insoluble support matrix or surface for use in a variety of methods. Commonly used crosslinking agents include, *e.g.*, 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), and bifunctional maleimides such as bis-N-maleimido-1,8-octane. Derivatizing agents such as methyl-3-[(p-azidophenyl)dithio]propioimide yield photoactivatable intermediates that are capable of forming crosslinks in the presence of light. Alternatively, reactive water-insoluble matrices such as cyanogen bromide-activated carbohydrates and the reactive substrates described in U.S. Pat. Nos. 3,969,287; 3,691,016; 4,195,128; 4,247,642; 4,229,537; and 4,330,440 are employed for protein immobilization.

Glutamyl and asparagyl residues are frequently deamidated to the corresponding glutamyl and aspartyl residues, respectively. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains (T. E. Creighton, *Proteins: Structure and Molecular Properties*, W. H. Freeman & Co., San Francisco, 1983, pp. 79-86), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the antibody constructs included within the scope of this invention comprises altering the glycosylation pattern of the protein. As is known in the art, glycosylation patterns can depend on both the sequence of the protein (*e.g.*, the presence or absence of particular glycosylation amino acid residues, discussed below), or the host cell or organism in which the protein is produced. Particular expression systems are discussed below.

Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose,

or xylose, to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

5 Addition of glycosylation sites to the antibody construct is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tri-peptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the starting sequence (for O-linked glycosylation sites). For ease, the amino acid sequence of an antibody construct is preferably altered through changes at the DNA level, particularly by
10 mutating the DNA encoding the polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

Another means of increasing the number of carbohydrate moieties on the antibody construct is by chemical or enzymatic coupling of glycosides to the protein. These procedures are
15 advantageous in that they do not require production of the protein in a host cell that has glycosylation capabilities for N- and O-linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as those of phenylalanine,
20 tyrosine, or tryptophan, or (f) the amide group of glutamine. These methods are described in WO 87/05330, and in Aplin and Wriston, 1981, CRC Crit. Rev. Biochem., pp. 259-306.

Removal of carbohydrate moieties present on the starting antibody construct may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of
25 the protein to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the polypeptide intact. Chemical deglycosylation is described by Hakimuddin *et al.*, 1987, *Arch. Biochem. Biophys.* 259:52 and by Edge *et al.*, 1981, *Anal. Biochem.* 118:131. Enzymatic cleavage of carbohydrate
30 moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura *et al.*, 1987, *Meth. Enzymol.* 138:350. Glycosylation at potential glycosylation sites may be prevented by the use of the compound tunicamycin as described by Duskin *et al.*, 1982, *J. Biol. Chem.* 257:3105. Tunicamycin blocks the formation of protein-N-glycoside linkages.

35 Other modifications of the antibody construct are contemplated herein. For example, another type of covalent modification of the antibody construct comprises linking the antibody

construct to various non-proteinaceous polymers, including, but not limited to, various polyols such as polyethylene glycol, polypropylene glycol, polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol, in the manner set forth in U.S. Pat. Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337. In addition, as is known in the art, amino acid substitutions may be made in various positions within the antibody construct, e.g. in order to facilitate the addition of polymers such as PEG.

In some embodiments, the covalent modification of the antibody constructs of the invention comprises the addition of one or more labels. The labelling group may be coupled to the antibody construct *via* spacer arms of various lengths to reduce potential steric hindrance. Various methods for labelling proteins are known in the art and can be used in performing the present invention. The term "label" or "labelling group" refers to any detectable label. In general, labels fall into a variety of classes, depending on the assay in which they are to be detected – the following examples include, but are not limited to:

- a) isotopic labels, which may be radioactive or heavy isotopes, such as radioisotopes or radionuclides (e.g., ³H, ¹⁴C, ¹⁵N, ³⁵S, ⁸⁹Zr, ⁹⁰Y, ⁹⁹Tc, ¹¹¹In, ¹²⁵I, ¹³¹I)
- b) magnetic labels (e.g., magnetic particles)
- c) redox active moieties
- d) optical dye (including, but not limited to, chromophores, phosphors and fluorophores) such as fluorescent groups (e.g., FITC, rhodamine, lanthanide phosphors), chemiluminescent groups, and fluorophores which can be either "small molecule" fluoeres or proteinaceous fluoeres
- e) enzymatic groups (e.g. horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase)
- f) biotinylated groups
- g) predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags, etc.)

By "fluorescent label" is meant any molecule that may be detected *via* its inherent fluorescent properties. Suitable fluorescent labels include, but are not limited to, fluorescein, rhodamine, tetramethylrhodamine, eosin, erythrosin, coumarin, methyl-coumarins, pyrene, Malacite green, stilbene, Lucifer Yellow, Cascade BlueJ, Texas Red, IAEDANS, EDANS, BODIPY FL, LC Red 640, Cy 5, Cy 5.5, LC Red 705, Oregon green, the Alexa-Fluor dyes (Alexa Fluor 350, Alexa Fluor 430, Alexa Fluor 488, Alexa Fluor 546, Alexa Fluor 568, Alexa Fluor 594, Alexa Fluor 633, Alexa Fluor 660, Alexa Fluor 680), Cascade Blue, Cascade Yellow and R-

phycoerythrin (PE) (Molecular Probes, Eugene, OR), FITC, Rhodamine, and Texas Red (Pierce, Rockford, IL), Cy5, Cy5.5, Cy7 (Amersham Life Science, Pittsburgh, PA). Suitable optical dyes, including fluorophores, are described in Molecular Probes Handbook by Richard P. Haugland.

5

Suitable proteinaceous fluorescent labels also include, but are not limited to, green fluorescent protein, including a Renilla, Ptilosarcus, or Aequorea species of GFP (Chalfie *et al.*, 1994, *Science* 263:802-805), EGFP (Clontech Laboratories, Inc., Genbank Accession Number U55762), blue fluorescent protein (BFP, Quantum Biotechnologies, Inc. 1801 de
10 Maisonneuve Blvd. West, 8th Floor, Montreal, Quebec, Canada H3H 1J9; Stauber, 1998, *Biotechniques* 24:462-471; Heim *et al.*, 1996, *Curr. Biol.* 6:178-182), enhanced yellow fluorescent protein (EYFP, Clontech Laboratories, Inc.), luciferase (Ichiki *et al.*, 1993, *J. Immunol.* 150:5408-5417), β galactosidase (Nolan *et al.*, 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:2603-2607) and Renilla (WO92/15673, WO95/07463, WO98/14605, WO98/26277,
15 WO99/49019, U.S. Patent Nos. 5292658, 5418155, 5683888, 5741668, 5777079, 5804387, 5874304, 5876995, 5925558).

Leucine zipper domains are peptides that promote oligomerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins
20 (Landschulz *et al.*, 1988, *Science* 240:1759), and have since been found in a variety of different proteins. Among the known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble oligomeric proteins are described in PCT application WO 94/10308, and the leucine zipper derived from lung surfactant protein D (SPD) described in Hoppe *et al.*,
25 1994, *FEBS Letters* 344:191. The use of a modified leucine zipper that allows for stable trimerization of a heterologous protein fused thereto is described in Fanslow *et al.*, 1994, *Semin. Immunol.* 6:267-78. In one approach, recombinant fusion proteins comprising the target antigen antibody fragment or derivative fused to a leucine zipper peptide are expressed in suitable host cells, and the soluble oligomeric target antigen antibody fragments
30 or derivatives that form are recovered from the culture supernatant.

The antibody construct of the invention may also comprise additional domains, which are e.g. helpful in the isolation of the molecule or relate to an adapted pharmacokinetic profile of the molecule. Domains helpful for the isolation of an antibody construct may be selected from
35 peptide motives or secondarily introduced moieties, which can be captured in an isolation method, e.g. an isolation column. Non-limiting embodiments of such additional domains comprise peptide motives known as Myc-tag, HAT-tag, HA-tag, TAP-tag, GST-tag, chitin

binding domain (CBD-tag), maltose binding protein (MBP-tag), Flag-tag, Strep-tag and variants thereof (e.g. StreptII-tag) and His-tag. All herein disclosed antibody constructs characterized by the identified CDRs are preferred to comprise a His-tag domain, which is generally known as a repeat of consecutive His residues in the amino acid sequence of a molecule, preferably of six His residues.

T cells or T lymphocytes are a type of lymphocyte (itself a type of white blood cell) that play a central role in cell-mediated immunity. There are several subsets of T cells, each with a distinct function. T cells can be distinguished from other lymphocytes, such as B cells and NK cells, by the presence of a T cell receptor (TCR) on the cell surface. The TCR is responsible for recognizing antigens bound to major histocompatibility complex (MHC) molecules and is composed of two different protein chains. In 95% of the T cells, the TCR consists of an alpha (α) and beta (β) chain. When the TCR engages with antigenic peptide and MHC (peptide / MHC complex), the T lymphocyte is activated through a series of biochemical events mediated by associated enzymes, co-receptors, specialized adaptor molecules, and activated or released transcription factors

The CD3 receptor complex is a protein complex and is composed of four chains. In mammals, the complex contains a CD3 γ (gamma) chain, a CD3 δ (delta) chain, and two CD3 ϵ (epsilon) chains. These chains associate with the T cell receptor (TCR) and the so-called ζ (zeta) chain to form the T cell receptor CD3 complex and to generate an activation signal in T lymphocytes. The CD3 γ (gamma), CD3 δ (delta), and CD3 ϵ (epsilon) chains are highly related cell-surface proteins of the immunoglobulin superfamily containing a single extracellular immunoglobulin domain. The intracellular tails of the CD3 molecules contain a single conserved motif known as an immunoreceptor tyrosine-based activation motif or ITAM for short, which is essential for the signaling capacity of the TCR. The CD3 epsilon molecule is a polypeptide which in humans is encoded by the *CD3E* gene which resides on chromosome 11. The sequence of a preferred human CD3 epsilon extracellular domain is shown in SEQ ID NO: 1, and the most preferred CD3 binding epitope corresponding to amino acid residues 1-27 of the human CD3 epsilon extracellular domain is represented in SEQ ID NO: 2.

The redirected lysis of target cells via the recruitment of T cells by a multispecific, at least bispecific, antibody construct involves cytolytic synapse formation and delivery of perforin and granzymes. The engaged T cells are capable of serial target cell lysis, and are not affected by immune escape mechanisms interfering with peptide antigen processing and presentation, or clonal T cell differentiation; see, for example, WO 2007/042261.

Cytotoxicity mediated by bispecific constructs can be measured in various ways. Effector cells can be e.g. stimulated enriched (human) CD8 positive T cells or unstimulated (human) peripheral blood mononuclear cells (PBMC). If the target cells are of macaque origin or express or are transfected with macaque target cell antigen, the effector cells should also be of macaque origin such as a macaque T cell line, e.g. 4119LnPx. The target cells should express (at least the extracellular domain of) target cell antigen, e.g. human or macaque target cell antigen. Target cells can be a cell line (such as CHO) which is stably or transiently transfected with target cell antigen, e.g. human or macaque target cell antigen. Alternatively, the target cells can be a target cell antigen positive natural expresser cell line, such as a human cancer cell line. Usually EC₅₀ values are expected to be lower with target cell lines expressing higher levels of target cell antigen on the cell surface. The effector to target cell (E:T) ratio is usually about 10:1, but can also vary. Cytotoxic activity of bispecific constructs can be measured in a ⁵¹chromium release assay (incubation time of about 18 hours) or in a FACS-based cytotoxicity assay (incubation time of about 48 hours). Modifications of the assay incubation time (cytotoxic reaction) are also possible. Other methods of measuring cytotoxicity are well-known to the skilled person and comprise MTT or MTS assays, ATP-based assays including bioluminescent assays, the sulforhodamine B (SRB) assay, WST assay, clonogenic assay and the ECIS technology.

The cytotoxic activity mediated by bispecific constructs of the present invention is preferably measured in a cell-based cytotoxicity assay. It is represented by the EC₅₀ value, which corresponds to the half maximal effective concentration (concentration of the antibody construct which induces a cytotoxic response halfway between the baseline and maximum). Preferably, the EC₅₀ value of the bispecific constructs is ≤20.000 pg/ml, more preferably ≤5000 pg/ml, even more preferably ≤1000 pg/ml, even more preferably ≤500 pg/ml, even more preferably ≤350 pg/ml, even more preferably ≤250 pg/ml, even more preferably ≤100 pg/ml, even more preferably ≤50 pg/ml, even more preferably ≤10 pg/ml, and most preferably ≤5 pg/ml.

Any of the above given EC₅₀ values can be combined with any one of the indicated scenarios of a cell-based cytotoxicity assay, e.g. in line with the methods described in the appended example. For example, when (human) CD8 positive T cells or a macaque T cell line are used as effector cells, the EC₅₀ value of the bispecific construct of the invention (e.g. a target cell antigen/CD3 bispecific construct) is preferably ≤1000 pg/ml, more preferably ≤500 pg/ml, even more preferably ≤250 pg/ml, even more preferably ≤100 pg/ml, even more preferably ≤50 pg/ml, even more preferably ≤10 pg/ml, and most preferably ≤5 pg/ml. If in this assay the

target cells are (human or macaque) cells transfected with the target antigen (e.g. the target cell antigen CD33), such as CHO cells, the EC₅₀ value of the bispecific construct is preferably ≤150 pg/ml, more preferably ≤100 pg/ml, even more preferably ≤50 pg/ml, even more preferably ≤30 pg/ml, even more preferably ≤10 pg/ml, and most preferably ≤5 pg/ml. If the target cells are a positive natural expresser cell line (e.g. of target cell antigen), then the EC₅₀ value is preferably ≤350 pg/ml, more preferably ≤250 pg/ml, even more preferably ≤200 pg/ml, even more preferably ≤100 pg/ml, even more preferably ≤150 pg/ml, even more preferably ≤100 pg/ml, and most preferably ≤50 pg/ml, or lower. When (human) PBMCs are used as effector cells, the EC₅₀ value of the bispecific construct is preferably ≤1000 pg/ml, more preferably ≤750 pg/ml, more preferably ≤500 pg/ml, even more preferably ≤350 pg/ml, even more preferably ≤250 pg/ml, even more preferably ≤100 pg/ml, and most preferably ≤50 pg/ml, or lower.

Preferably, the bispecific constructs of the present invention do not induce / mediate lysis or do not essentially induce / mediate lysis of target cell antigen negative cells such as CHO cells. The term “do not induce lysis”, “do not essentially induce lysis”, “do not mediate lysis” or “do not essentially mediate lysis” means that an antibody constructs of the present invention does not induce or mediate lysis of more than 30%, preferably not more than 20%, more preferably not more than 10%, particularly preferably not more than 9%, 8%, 7%, 6% or 5% of target cell antigen negative cells, whereby lysis of a target cell antigen positive cell line is set to be 100%. This usually applies for concentrations of the antibody construct of up to 500 nM. The skilled person knows how to measure cell lysis without further ado. Moreover, the present specification teaches specific instructions how to measure cell lysis.

Preferably, the bispecific construct for the use according to the invention is administered according to a schedule comprising the following steps:

- (a) administration of a first dose of the bispecific construct, followed by
- (b) administration of a second dose of the bispecific construct, wherein said second dose exceeds said first dose, optionally followed by
- (a) administration of a third dose of the bispecific construct, wherein said optional third dose exceeds said second dose.

In line with the above it is further preferred that the period of administration of the first dose is up to seven days. This period of administration of the first dose may be used during the initial phase/first cycle of administration of the bispecific construct e.g to reduce the tumor load in a patient (tumor debulking) while avoiding conditions such as cytokine storm and/or cytokine

release syndrome which one might expect in case a higher dose is used during the period of administration of the first dose.

While in one embodiment of the invention the period of administration of the first dose is up to seven days, it is also within this preferred embodiment that this first dose is administered for a period of six days, five days, four days, three days, two days or one day. In the case that the tumor load or general condition of the individual patient does require the administration of the limited dose of the bispecific construct in the first limited dose step, this first dose step is understood as a run-in phase/adaptation phase which should avoid or limit side effects resulting from the first contact of the patient with the bispecific construct. A preferred range for a dose in such run-in phase/adaptation phase may be in a range of 1 to 50 $\mu\text{g}/\text{d}$, preferably in a range of 3 to 30 $\mu\text{g}/\text{d}$, further preferably in a range of 4 to 20 $\mu\text{g}/\text{d}$ and even more preferably in a range of 5 to 15 $\mu\text{g}/\text{d}$ for a canonical BiTE[®] such as AMG 330, which is a 54 kDa single chain polypeptide. In a very preferred embodiment, the bispecific construct according to the present invention is administered at a dose of 10 $\mu\text{g}/\text{d}$. In the case of a half-life extended bispecific antibody construct, respective equimolar doses can be easily determined. However, in case the tumor load or general condition of the individual patient does not require such administration of the limited dose of the bispecific construct in the first limited dose step, already the first does may be in the range.

Preferred ranges for a second dose of the bispecific construct are e.g. for a canonical BiTE[®] such as AMG 330 in the range of 10 $\mu\text{g}/\text{d}$ to 10 mg/d, more preferably in the range of 25 $\mu\text{g}/\text{d}$ to 1 mg/d and even more preferably in the range of 30 $\mu\text{g}/\text{d}$ to 500 $\mu\text{g}/\text{d}$. In a very preferred embodiment, the second dose is 30 $\mu\text{g}/\text{d}$ or 60 $\mu\text{g}/\text{d}$. In line with the above, the preferred ranges for the third dose of the bispecific construct exceed the respective dose of the second dose. The third dose is typically in the range of 60 $\mu\text{g}/\text{d}$ to 500 $\mu\text{g}/\text{d}$ and preferably eradicates residual target cells which may have evaded treatment equivalent to the second dose according to the present invention.

In one embodiment of the bispecific construct for the use of the invention, the complete period of administration of the dose steps subsequent to the first dose is in the range of eight to thirteen days. In line with the above, the complete period of administration in the different dose steps does not exceed 14 days. Accordingly, the period of administration of the bispecific construct of the second and optional third dose may be preferably 13, 12, 11, 10, 9, or 8 days, most preferably 10 days.

It was surprisingly found that when a step dosing is applied, e.g. if a first dose is given for four days and a second dose is given for 10 days, wherein the first dose is 5 to 15 $\mu\text{g}/\text{d}$,

preferably 10 µg/d, and wherein the second dose is 30/d µg to 500 µg/d, preferably 30 µg/d or 60 µg/d, then immunologic side effects such as undesired cytokine release, e.g. a cytokine release syndrome, may be effectively prevented. In contrast, if a dose equivalent to the second dose is given without a prior lower dose equitant to the first dose of the present invention, then side effects, such as undesired cytokine release, e.g. a cytokine release syndrome, may occur.

It is also preferred for the present invention that the period of administration of the second dose is four to seven days, preferably four days, and the period of administration of the third dose is four to seven days. Accordingly, it is particularly preferred that the period of administration of the second dose is 7, 6, 5, or 4 days and the period of administration of the optional third dose is 7, 6, 5, or 4 days.

Also in line with the present invention the treatment of the myeloid leukemia preferably comprises two or more cycles of treatment which each comprises the maximal period of 14 days of construct administration followed by the period of at least 14 days without administration of the construct. In case the treatment in line with the invention comprises two or more cycles, it is further preferred in one embodiment that only the first cycle of the treatment comprises the administration according to step (a), whereas the following cycles start with the dose according to step (b).

Also in line with the present invention it is preferred for the bispecific construct used in the treatment of myeloid leukemia that the first binding domain of the bispecific construct comprises groups of six CDRs selected from the group consisting of SEQ ID NOs: 10 to 12 and 14 to 16, 22 to 24 and 26 to 28, 34 to 36 and 38 to 40, 46 to 48 and 50 to 52, 58 to 60 and 62 to 64, 70 to 72 and 74 to 76, 82 to 84 and 86 to 88, 94 to 96 and 98 to 100.

Moreover, in line with the present invention it is preferred for the bispecific construct used in the treatment of myeloid leukemia that the second binding domain of the bispecific construct comprises groups of six CDRs selected from the group consisting of SEQ ID NOs: 9 to 14, 27 to 32, 45 to 50, 63 to 68, 81 to 86, 99 to 104, 117 to 122, 135 to 140, 153 to 158 and 171 to 176 of WO 2008/119567.

As well as the second binding domain the first (or any further) binding domain(s) of the antibody construct of the invention is/are preferably cross-species specific for members of the mammalian order of primates. Cross-species specific CD3 binding domains are, for example, described in WO 2008/119567. According to one embodiment, the first and second

binding domain, in addition to binding to human CD33 target cell antigen and human CD3, respectively, will also bind to the CD33 target cell antigen / CD3 of primates including (but not limited to) new world primates (such as *Callithrix jacchus*, *Saguinus Oedipus* or *Saimiri sciureus*), old world primates (such baboons and macaques), gibbons, and non-human
5 *homininae*. *Callithrix jacchus* and *Saguinus oedipus* are both new world primate belonging to the family of *Callitrichidae*, while *Saimiri sciureus* is a new world primate belonging to the family of *Cebidae*.

In a preferred embodiment of the invention the bispecific construct is a bispecific antibody
10 construct. In line with the definitions provided herein above, this embodiment relates to bispecific constructs, which are antibody constructs. In a preferred embodiment of the invention the bispecific antibody construct is a single chain construct. Such bispecific single chain antibody construct may comprise in line with the invention an amino acid sequence selected from the group consisting of SEQ ID NOs: 18, 19, 20, 30, 31, 32, 42, 43, 44, 54, 55,
15 56, 66, 67, 68, 78, 79, 80, 90, 91, 92, 102, 103, 104, 105, 106, 107 and 108.

Amino acid sequence modifications of the bispecific constructs described herein are also contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the bispecific construct. Amino acid sequence variants of the
20 bispecific constructs are prepared by introducing appropriate nucleotide changes into the bispecific constructs nucleic acid, or by peptide synthesis. All of the below described amino acid sequence modifications should result in a bispecific construct which still retains the desired biological activity (binding to target cell antigen and to CD3) of the unmodified parental molecule.

25 The term "amino acid" or "amino acid residue" typically refers to an amino acid having its art recognized definition such as an amino acid selected from the group consisting of: alanine (Ala or A); arginine (Arg or R); asparagine (Asn or N); aspartic acid (Asp or D); cysteine (Cys or C); glutamine (Gln or Q); glutamic acid (Glu or E); glycine (Gly or G); histidine (His or H);
30 isoleucine (He or I); leucine (Leu or L); lysine (Lys or K); methionine (Met or M); phenylalanine (Phe or F); pro line (Pro or P); serine (Ser or S); threonine (Thr or T); tryptophan (Trp or W); tyrosine (Tyr or Y); and valine (Val or V), although modified, synthetic, or rare amino acids may be used as desired. Generally, amino acids can be grouped as having a nonpolar side chain (e.g., Ala, Cys, He, Leu, Met, Phe, Pro, Val); a negatively
35 charged side chain (e.g., Asp, Glu); a positively charged sidechain (e.g., Arg, His, Lys); or an uncharged polar side chain (e.g., Asn, Cys, Gln, Gly, His, Met, Phe, Ser, Thr, Trp, and Tyr).

Amino acid modifications include, for example, deletions from, and/or insertions into, and/or substitutions of, residues within the amino acid sequences of the bispecific constructs. Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid changes also may alter post-translational processes of the bispecific constructs, such as changing the number or position of glycosylation sites.

For example, 1, 2, 3, 4, 5, or 6 amino acids may be inserted or deleted in each of the CDRs (of course, dependent on their length), while 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or 25 amino acids may be inserted or deleted in each of the FRs. Preferably, amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 residues to polypeptides containing a hundred or more residues, as well as intra-sequence insertions of single or multiple amino acid residues. An insertional variant of the bispecific construct of the invention includes the fusion to the N-terminus or to the C-terminus of the bispecific construct to an enzyme or a fusion to a polypeptide which increases the serum half-life of the bispecific construct.

An increased half-life is generally useful in *in vivo* applications of immunoglobulins, especially antibodies and most especially antibody fragments of small size. Although such antibody constructs based on antibody fragments (Fvs, disulphide bonded Fvs, Fabs, scFvs, dAbs) are able to rapidly reach most parts of the body, those antibody constructs are likely to suffer from rapid clearance from the body. Strategies described in the art for extending the half-life of antibody constructs such as single-chain diabodies include the conjugation of polyethylene glycol chains (PEGylation), the fusion to the IgG Fc region or to an albumin or albumin-binding domain.

Serum albumin is a protein physiologically produced by the liver; it occurs dissolved in blood plasma and is the most abundant blood protein in mammals. Albumin is essential for maintaining the oncotic pressure needed for proper distribution of body fluids between blood vessels and body tissues. It also acts as a plasma carrier by non-specifically binding several hydrophobic steroid hormones and as a transport protein for hemin and fatty acids. The term "serum albumin" respectively the human variant thereof ("human albumin") defines in the context of the invented proteins either the parental human serum albumin protein (sequence as described in SEQ ID NO: 109) or any variant (e.g. such as albumin protein as depicted in SEQ ID NOs: 110-138) or fragment thereof preferably expressed as genetic fusion proteins and by chemical crosslinking etc. at least with one therapeutic protein. Variants comprising single or multiple mutations or fragments of albumin provide improved properties such as affinities to FcRn receptor and extended plasma half-life compared to its parent or reference. Variants of human albumin are described e.g. in WO 2014/072481. In line with the invention

the serum albumin may be linked to the antibody construct via a peptide linker. It is preferred that the peptide linker has the amino acid sequence (GGGGS)_n (SEQ ID NO: 13)_n wherein “n” is an integer in the range of 1 to 5. Further preferred is that “n” is an integer in the range of 1 to 3, and most preferably “n” is 1 or 2.

5

The sites of greatest interest for substitutional mutagenesis include the CDRs of the heavy and/or light chain, in particular the hypervariable regions, but FR alterations in the heavy and/or light chain are also contemplated. The substitutions are preferably conservative substitutions as described herein. Preferably, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids may be substituted in a CDR, while 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or 25 amino acids may be substituted in the framework regions (FRs), depending on the length of the CDR or FR. For example, if a CDR sequence encompasses 6 amino acids, it is envisaged that one, two or three of these amino acids are substituted. Similarly, if a CDR sequence encompasses 15 amino acids it is envisaged that one, two, three, four, five or six of these amino acids are substituted.

15

A useful method for identification of certain residues or regions of the bispecific constructs that are preferred locations for mutagenesis is called “alanine scanning mutagenesis” as described by Cunningham and Wells in *Science*, 244: 1081-1085 (1989). Here, a residue or group of target residues within the bispecific construct is/are identified (e.g. charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with the epitope.

20

Those amino acid locations demonstrating functional sensitivity to the substitutions then are refined by introducing further or other variants at, or for, the sites of substitution. Thus, while the site or region for introducing an amino acid sequence variation is predetermined, the nature of the mutation *per se* needs not to be predetermined. For example, to analyze or optimize the performance of a mutation at a given site, alanine scanning or random mutagenesis may be conducted at a target codon or region, and the expressed bispecific construct variants are screened for the optimal combination of desired activity. Techniques for making substitution mutations at predetermined sites in the DNA having a known sequence are well known, for example, M13 primer mutagenesis and PCR mutagenesis. Screening of the mutants is done using assays of target antigen binding activities.

30

35

Generally, if amino acids are substituted in one or more or all of the CDRs of the heavy and/or light chain, it is preferred that the then-obtained “substituted” sequence is at least

60%, more preferably 65%, even more preferably 70%, particularly preferably 75%, more particularly preferably 80% identical to the "original" CDR sequence. This means that it is dependent of the length of the CDR to which degree it is identical to the "substituted" sequence. For example, a CDR having 5 amino acids is preferably 80% identical to its substituted sequence in order to have at least one amino acid substituted. Accordingly, the CDRs of the bispecific antibody construct may have different degrees of identity to their substituted sequences, e.g., CDRL1 may have 80%, while CDRL3 may have 90%.

Preferred substitutions (or replacements) are conservative substitutions. However, any substitution (including non-conservative substitution or one or more from the "exemplary substitutions" listed in Table 1, below) is envisaged as long as the bispecific construct retains its capability to bind to target cell antigen via the first binding domain and to CD3 epsilon via the second binding domain and/or its CDRs have an identity to the then substituted sequence (at least 60%, more preferably 65%, even more preferably 70%, particularly preferably 75%, more particularly preferably 80% identical to the "original" CDR sequence).

Conservative substitutions are shown in Table 1 under the heading of "preferred substitutions". If such substitutions result in a change in biological activity, then more substantial changes, denominated "exemplary substitutions" in Table A, or as further described below in reference to amino acid classes, may be introduced and the products screened for a desired characteristic.

Table A: Amino acid substitutions

Original	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val, leu, ile	val
Arg (R)	lys, gln, asn	lys
Asn (N)	gln, his, asp, lys, arg	gln
Asp (D)	glu, asn	glu
Cys (C)	ser, ala	ser
Gln (Q)	asn, glu	asn
Glu (E)	asp, gln	asp
Gly (G)	Ala	ala
His (H)	asn, gln, lys, arg	arg
Ile (I)	leu, val, met, ala, phe	leu
Leu (L)	norleucine, ile, val, met, ala	ile
Lys (K)	arg, gln, asn	arg

Met (M)	leu, phe, ile	leu
Phe (F)	leu, val, ile, ala, tyr	tyr
Pro (P)	Ala	ala
Ser (S)	Thr	thr
Thr (T)	Ser	ser
Trp (W)	tyr, phe	tyr
Tyr (Y)	trp, phe, thr, ser	phe
Val (V)	ile, leu, met, phe, ala	leu

Substantial modifications in the biological properties of the bispecific construct of the present invention are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties: (1) hydrophobic: norleucine, met, ala, val, leu, ile; (2) neutral hydrophilic: cys, ser, thr, asn, gln; (3) acidic: asp, glu; (4) basic: his, lys, arg; (5) residues that influence chain orientation: gly, pro; and (6) aromatic : trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Any cysteine residue not involved in maintaining the proper conformation of the bispecific construct may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the antibody to improve its stability (particularly where the antibody is an antibody fragment such as an Fv fragment).

For amino acid sequences, sequence identity and/or similarity is determined by using standard techniques known in the art, including, but not limited to, the local sequence identity algorithm of Smith and Waterman, 1981, *Adv. Appl. Math.* 2:482, the sequence identity alignment algorithm of Needleman and Wunsch, 1970, *J. Mol. Biol.* 48:443, the search for similarity method of Pearson and Lipman, 1988, *Proc. Nat. Acad. Sci. U.S.A.* 85:2444, computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.), the Best Fit sequence program described by Devereux *et al.*, 1984, *Nucl. Acid Res.* 12:387-395, preferably using the default settings, or by inspection. Preferably, percent identity is calculated by FastDB based upon the following parameters: mismatch penalty of 1; gap penalty of 1; gap size penalty of 0.33; and joining penalty of 30, "Current

Methods in Sequence Comparison and Analysis," Macromolecule Sequencing and Synthesis, Selected Methods and Applications, pp 127-149 (1988), Alan R. Liss, Inc.

5 An example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments. It can also plot a tree showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, 1987, *J. Mol. Evol.* 35:351-360; the method is similar to that described by Higgins and Sharp, 1989, *CABIOS* 5:151-153. Useful PILEUP parameters including a default gap weight of 3.00, a default gap
10 length weight of 0.10, and weighted end gaps.

Another example of a useful algorithm is the BLAST algorithm, described in: Altschul *et al.*, 1990, *J. Mol. Biol.* 215:403-410; Altschul *et al.*, 1997, *Nucleic Acids Res.* 25:3389-3402; and Karin *et al.*, 1993, *Proc. Natl. Acad. Sci. U.S.A.* 90:5873-5787. A particularly useful BLAST
15 program is the WU-BLAST-2 program which was obtained from Altschul *et al.*, 1996, *Methods in Enzymology* 266:460-480. WU-BLAST-2 uses several search parameters, most of which are set to the default values. The adjustable parameters are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=11. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending
20 upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched; however, the values may be adjusted to increase sensitivity.

An additional useful algorithm is gapped BLAST as reported by Altschul *et al.*, 1993, *Nucl.*
25 *Acids Res.* 25:3389-3402. Gapped BLAST uses BLOSUM-62 substitution scores; threshold T parameter set to 9; the two-hit method to trigger ungapped extensions, charges gap lengths of k a cost of 10+k; Xu set to 16, and Xg set to 40 for database search stage and to 67 for the output stage of the algorithms. Gapped alignments are triggered by a score corresponding to about 22 bits.

30 Generally, the amino acid homology, similarity, or identity between individual variant CDRs are at least 60% to the sequences depicted herein, and more typically with preferably increasing homologies or identities of at least 65% or 70%, more preferably at least 75% or 80%, even more preferably at least 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%,
35 99%, and almost 100%. In a similar manner, "percent (%) nucleic acid sequence identity" with respect to the nucleic acid sequence of the binding proteins identified herein is defined as the percentage of nucleotide residues in a candidate sequence that are identical with the

nucleotide residues in the coding sequence of the bispecific construct. A specific method utilizes the BLASTN module of WU-BLAST-2 set to the default parameters, with overlap span and overlap fraction set to 1 and 0.125, respectively.

5 Generally, the nucleic acid sequence homology, similarity, or identity between the nucleotide sequences encoding individual variant CDRs and the nucleotide sequences depicted herein are at least 60%, and more typically with preferably increasing homologies or identities of at least 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%, and almost 100%. Thus, a "variant CDR" is
10 one with the specified homology, similarity, or identity to the parent CDR of the invention, and shares biological function, including, but not limited to, at least 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% of the specificity and/or activity of the parent CDR.

15 In one embodiment the bispecific construct for the use in accordance with this invention is administered in combination with one or more epigenetic factors selected from the group consisting of histone deacetylase (HDAC) inhibitors, DNA methyltransferase (DNMT) I inhibitors, hydroxyurea, Granulocyte-Colony Stimulating Factor (G-CSF), histone demethylase inhibitors and ATRA (All Trans-retinoic acid) and wherein:

- 20 (a) the one or more epigenetic factors are administered prior to the administration of the bispecific construct;
- (b) the one or more epigenetic factors are administered subsequent to the administration of the bispecific construct; or
- (c) the one or more epigenetic factors and the bispecific construct are administered
25 simultaneously.

The term "epigenetic factor" in connection with the present invention defines a compound which is capable of changing the gene expression or cellular phenotype of a cell population upon administration. It is understood that such change refers to one or more functional relevant modifications to the genome without involving a change in the nucleic acid
30 sequence. Examples of such modifications are DNA methylation and histone modification, which are both important for the regulation of gene expression without altering the underlying DNA sequence.

Details for a treatment of myeloid leukemia comprising the administration of the bispecific construct in combination with one or more of the above described epigenetic factors have
35 been provided in PCT/EP2014/069575.

In one embodiment of the invention it is preferred that the one or more epigenetic factors are administered up to seven days prior to the administration of the bispecific construct.

Also in one embodiment of the invention it is preferred that the epigenetic factor is hydroxyurea.

5

It is preferred for the present invention that the myeloid leukemia is selected from the group consisting of acute myeloblastic leukemia, chronic neutrophilic leukemia, myeloid dendritic cell leukemia, accelerated phase chronic myelogenous leukemia, acute myelomonocytic leukemia, juvenile myelomonocytic leukemia, chronic myelomonocytic leukemia, acute
10 basophilic leukemia, acute eosinophilic leukemia, chronic eosinophilic leukemia, acute megakaryoblastic leukemia, essential thrombocytosis, acute erythroid leukemia, polycythemia vera, myelodysplastic syndrome, acute panmyeloic leukemia, myeloid sarcoma, and acute biphenotypic leukaemia. More preferably, the myeloid leukemia is an acute myeloid leukemia (AML). The definition of AML inter alia comprises acute myeloblastic
15 leukemia, acute myeloid dendritic cell leukemia, acute myelomonocytic leukemia, acute basophilic leukemia, acute eosinophilic leukemia, acute megakaryoblastic leukemia, acute erythroid leukemia, and acute panmyeloic leukemia

The bispecific construct described in connection with this invention may be formulated for an
20 appropriate administration to a subject in the need thereof in form of a pharmaceutical composition.

Formulations described herein are useful as pharmaceutical compositions in the treatment, amelioration and/or prevention of the pathological medical condition as described herein in a patient in need thereof. The term "treatment" refers to both therapeutic treatment and
25 prophylactic or preventative measures. Treatment includes the application or administration of the formulation to the body, an isolated tissue, or cell from a patient who has a disease/disorder, a symptom of a disease/disorder, or a predisposition toward a disease/disorder, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect the disease, the symptom of the disease, or the predisposition toward the
30 disease.

The term "disease" refers to any condition that would benefit from treatment with the bispecific construct or the pharmaceutical composition described herein. This includes chronic and acute disorders or diseases including those pathological conditions that
35 predispose the mammal to the disease in question.

The terms "subject in need" or those "in need of treatment" includes those already with the disorder, as well as those in which the disorder is to be prevented. The subject in need or "patient" includes human and other mammalian subjects that receive either prophylactic or therapeutic treatment.

5

The bispecific construct of the invention will generally be designed for specific routes and methods of administration, for specific dosages and frequencies of administration, for specific treatments of specific diseases, with ranges of bio-availability and persistence, among other things. The materials of the composition are preferably formulated in concentrations that are acceptable for the site of administration.

10

Formulations and compositions thus may be designed in accordance with the invention for delivery by any suitable route of administration. In the context of the present invention, the routes of administration include, but are not limited to

15

- topical routes (such as epicutaneous, inhalational, nasal, ophthalmic, auricular / aural, vaginal, mucosal);
- enteral routes (such as oral, gastrointestinal, sublingual, sublabial, buccal, rectal); and
- parenteral routes (such as intravenous, intraarterial, intraosseous, intramuscular, intracerebral, intracerebroventricular, epidural, intrathecal, subcutaneous, intraperitoneal, extra-amniotic, intraarticular, intracardiac, intradermal, intralesional, intrauterine, intravesical, intravitreal, transdermal, intranasal, transmucosal, intrasynovial, intraluminal).

20

25

The pharmaceutical compositions and the bispecific construct described in connection with the invention are particularly useful for parenteral administration, e.g., subcutaneous or intravenous delivery, for example by injection such as bolus injection, or by infusion such as continuous infusion. Pharmaceutical compositions may be administered using a medical device. Examples of medical devices for administering pharmaceutical compositions are described in U.S. Patent Nos. 4,475,196; 4,439,196; 4,447,224; 4,447, 233; 4,486,194; 4,487,603; 4,596,556; 4,790,824; 4,941,880; 5,064,413; 5,312,335; 5,312,335; 5,383,851; and 5,399,163.

30

35

In particular, the present invention provides for an uninterrupted administration of the suitable composition. As a non-limiting example, uninterrupted or substantially uninterrupted, i.e. continuous administration may be realized by a small pump system worn by the patient for metering the influx of therapeutic agent into the body of the patient. The pharmaceutical

composition comprising the bispecific construct described in connection with the invention can be administered by using said pump systems. Such pump systems are generally known in the art, and commonly rely on periodic exchange of cartridges containing the therapeutic agent to be infused. When exchanging the cartridge in such a pump system, a temporary
5 interruption of the otherwise uninterrupted flow of therapeutic agent into the body of the patient may ensue. In such a case, the phase of administration prior to cartridge replacement and the phase of administration following cartridge replacement would still be considered within the meaning of the pharmaceutical means and methods of the invention together make up one "uninterrupted administration" of such therapeutic agent.

10

The continuous or uninterrupted administration of the bispecific construct described in connection with the invention may be intravenous or subcutaneous by way of a fluid delivery device or small pump system including a fluid driving mechanism for driving fluid out of a reservoir and an actuating mechanism for actuating the driving mechanism. Pump systems
15 for subcutaneous administration may include a needle or a cannula for penetrating the skin of a patient and delivering the suitable composition into the patient's body. Said pump systems may be directly fixed or attached to the skin of the patient independently of a vein, artery or blood vessel, thereby allowing a direct contact between the pump system and the skin of the patient. The pump system can be attached to the skin of the patient for 24 hours
20 up to several days. The pump system may be of small size with a reservoir for small volumes. As a non-limiting example, the volume of the reservoir for the suitable pharmaceutical composition to be administered can be between 0.1 and 50 ml.

25

The continuous administration may also be transdermal by way of a patch worn on the skin and replaced at intervals. One of skill in the art is aware of patch systems for drug delivery suitable for this purpose. It is of note that transdermal administration is especially amenable to uninterrupted administration, as exchange of a first exhausted patch can advantageously be accomplished simultaneously with the placement of a new, second patch, for example on the surface of the skin immediately adjacent to the first exhausted patch and immediately
30 prior to removal of the first exhausted patch. Issues of flow interruption or power cell failure do not arise.

35

If the pharmaceutical composition has been lyophilized, the lyophilized material is first reconstituted in an appropriate liquid prior to administration. The lyophilized material may be reconstituted in, e.g., bacteriostatic water for injection (BWHI), physiological saline, phosphate buffered saline (PBS), or the same formulation the protein had been in prior to lyophilization.

40

The compositions of the present invention can be administered to the subject at a suitable dose which can be determined e.g. by dose escalating studies by administration of increasing doses of the bispecific construct described in connection with the invention exhibiting cross-species specificity described herein to non-chimpanzee primates, for instance macaques. As set forth above, the bispecific construct described in connection with the invention exhibiting cross-species specificity described herein can be advantageously used in identical form in preclinical testing in non-chimpanzee primates and as drug in humans. The dosage regimen will be determined by the attending physician and clinical factors. As is well known in the medical arts, dosages for any one patient depend upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently.

The term "effective dose" or "effective dosage" is defined as an amount sufficient to achieve or at least partially achieve the desired effect. The term "therapeutically effective dose" is defined as an amount sufficient to cure or at least partially arrest the disease and its complications in a patient already suffering from the disease. Amounts or doses effective for this use will depend on the condition to be treated (the indication), the delivered bispecific construct, the therapeutic context and objectives, the severity of the disease, prior therapy, the patient's clinical history and response to the therapeutic agent, the route of administration, the size (body weight, body surface or organ size) and/or condition (the age and general health) of the patient, and the general state of the patient's own immune system. The proper dose can be adjusted according to the judgment of the attending physician such that it can be administered to the patient once or over a series of administrations, and in order to obtain the optimal therapeutic effect.

A typical dosage may range from about 0.1 µg/kg to up to about 30 mg/kg or more, depending on the factors mentioned above. In specific embodiments, the dosage may range from 1.0 µg/kg up to about 20 mg/kg, optionally from 10 µg/kg up to about 10 mg/kg or from 100 µg/kg up to about 5 mg/kg.

A therapeutic effective amount of a bispecific construct described in connection with the invention preferably results in a decrease in severity of disease symptoms, an increase in frequency or duration of disease symptom-free periods or a prevention of impairment or disability due to the disease affliction. For treating target cell antigen-expressing tumors, a therapeutically effective amount of the bispecific construct described in connection with the

invention, e.g. an anti-target cell antigen/anti-CD3 antibody construct, preferably inhibits cell growth or tumor growth by at least about 20%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, or at least about 90% relative to untreated patients. The ability of a compound to inhibit tumor growth may be evaluated in an animal model predictive of efficacy in human tumors.

The pharmaceutical composition can be administered as a sole therapeutic or in combination with additional therapies such as anti-cancer therapies as needed, e.g. other proteinaceous and non-proteinaceous drugs. These drugs may be administered simultaneously with the composition comprising the bispecific construct described in connection with the invention as defined herein or separately before or after administration of said bispecific construct in timely defined intervals and doses.

Further, the present inventors observed that rare side effects, such as immunologic side effects could be prevented or alleviated by means of a glucocorticoid (pre) and/or (co)therapy.

Accordingly, the present invention establishes that glucocorticoids such as dexamethasone mitigate or even prevent adverse effects which might occur in the course of a treatment with CD33/CD3 specific bispecific constructs according to the present invention.

Glucocorticoids (GCs) are still the most widely used immunosuppressive agents for the treatment of inflammatory disorders and autoimmune diseases. Glucocorticoids (GC) are a class of steroid hormones that bind to the glucocorticoid receptor (GR), which is present in almost every vertebrate animal cell, including humans. These compounds are potent anti-inflammatory agents, regardless of the inflammation's cause. Glucocorticoids suppress, inter alia, the cell-mediated immunity by inhibiting genes that code for the cytokines IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-8 and IFN- γ .

Cortisone which belongs to the group of GCs is an important therapeutic drug which is used to fight many ailments ranging from Addison's disease to rheumatoid arthritis. Ever since the discovery of its anti-rheumatic properties, which led to its acclaim as a wonder drug, many derivatives of cortisone with enhanced properties to better fight a specific ailment have been produced. Cortisone belongs to a group of steroids known as corticosteroids. These steroids are produced by the adrenal cortex, which is the outer part of the adrenal glands, near the kidneys. The corticosteroids are divided into two main groups: the glucocorticoids (GCs), which control fat, protein, calcium and carbohydrate metabolism, and the mineralocorticoids controlling sodium and potassium levels. Cortisone belongs to the former group, i.e. to the

GCs. Cortisone and its many derivatives are used for a variety of diseases. Cortisone also helped to make organ transplants a reality due to its ability to minimize the defence reaction of the body towards foreign proteins present in the implanted organ and thus damage the functionality of the implanted organ. However, despite clinical use during more than 50
5 years, the specific anti-inflammatory effects of GC on different cellular compartments of the immune system are not yet clear. GCs affect nearly every cell of the immune system, and there is growing evidence for cell type-specific mechanisms.

In one specific embodiment, the present invention relates to a glucocorticoid (GC) for use in
10 the amelioration, treatment or prophylaxis of adverse effects caused by a CD33/CD3 bispecific construct. As outlined above, these unwanted adverse effects may be prevented by a step dosing as described herein. However, for mere precaution, glucocorticoid(s) for use in the amelioration, treatment or prophylaxis of (immunological) adverse effects in a patient may be provided wherein said patient is subject to therapy with a CD33/CD3 bispecific
15 antibody construct. Accordingly, in one further aspect the present invention relates to a glucocorticoid (GC) for use in a method in the amelioration, treatment or prophylaxis of immunological adverse effects caused by a CD33/CD3 bispecific antibody construct according to the present invention.

20 Also, the present invention relates to a method of amelioration, treatment or prophylaxis of immunological adverse effects caused by a CD33/CD3 bispecific antibody construct, said method comprising administering to a patient in need thereof a glucocorticoid (GC). The GC is preferably administered in an amount which is sufficient to ameliorate, treat or prevent said immunological adverse effects caused by a CD33/CD3 bispecific antibody construct.

25 The term "glucocorticoid" means compounds that bind, preferably specifically, to the glucocorticoid receptor. Said term includes compound(s) selected from the group consisting of cortisone, cortisol (hydrocortisone), cloprednol, prednisone, prednisolone, methylprednisolone, deflazacort, flucortolone, triamcinolone, dexamethasone,
30 beatamethasone, cortivazol, paramethasone, and/or fluticasone, including pharmaceutically acceptable derivatives thereof. In the context of the embodiments of the present invention, the mentioned compounds may be used alone or in combination. Dexamethasone is preferred. The present invention is however not limited to the above mentioned specific GCs. It is envisaged that all substances which already are or will be classified as a "glucocorticoid",
35 may be employed in the context of the present invention as well. Such future glucocorticoids include compounds which specifically bind to and activate the glucocorticoid receptor. The term "specifically binds to the GC receptor" means in accordance with the present invention

that the GC (or a compound which is assumed to act like a GC) associates with (e.g., interacts with) the GC receptor (also known as NR3C1) to a statistically significant degree as compared to association with proteins/receptors generally (i.e., non-specific binding). When the GC receptor binds to glucocorticoids, its primary mechanism of action is the regulation of gene transcription. In the absence of GC, the glucocorticoid receptor (GR) resides in the cytosol complexed with a variety of proteins including heat shock protein 90 (hsp90), the heat shock protein 70 (hsp70) and the protein FKBP52 (FK506-binding protein 52). The binding of the GC to the glucocorticoid receptor (GR) results in release of the heat shock proteins. It is thus envisaged that a future GC, or a pharmaceutically acceptable derivative or salt of a GC is preferably able to bind to the GC receptor and to release the above mentioned heat shock proteins. The activated GR complex up-regulates the expression of anti-inflammatory proteins in the nucleus or represses the expression of pro-inflammatory proteins in the cytosol (by preventing the translocation of other transcription factors from the cytosol into the nucleus).

15

In a preferred embodiment, said GC is selected from the most clinical used and relevant GCs like dexamethasone, fluticasonepropionate, prednisolone, methylprednisolone, betamethasone, triamcinolonacetone or combinations thereof.

20 In an even more preferred embodiment, said GC is dexamethasone.

Dexamethasone has the highest glucocorticoid potency of the most commonly used steroids and also has the longest half-life (see Table below). But a person skilled in the field can select one of the other known glucocorticoids, some of which are disclosed herein, and select an appropriate effective dose to ameliorate or prevent immunological adverse events that may result from the treatment of a patient in need thereof.

25

Agent	Approx. equiv. dose (mg)	Relative anti-inflammatory (glucocorticoid) potency	Relative mineralocorticoid (Na⁺ retaining) potency	Biologic half-life (hrs)
Cortisone	25	0.8	0.8	8-12
Hydrocortisone	20	1	1	8-12
Prednisone	5	4	0.8	18-36
Prednisolone	5	4	0.8	18-36
Methylprednisolone	5	5	0.5	18-36
Dexamethasone	0.75	25	0	36-54

Dexamethasone also possesses a beneficial effect in malignant central nervous system (CNS) disease (e.g. CNS lymphoma or brain metastases) – possibly due to specific penetration to the CNS. It is also preferentially (over other steroids) used to treat brain edema. Although corticosteroids decrease capillary permeability in the tumor itself, it has been found in animal models that dexamethasone may act differently and decrease edema by effects on bulk flow away from the tumor (Molnar, Lapin, & Goothuis, 1995, Neurooncol. 1995;25(1):19-28.

For the clinical trials in connection with the application of a CD33/CD3 bispecific antibody construct, the present inventors had to develop a treatment regime which was efficient and would be well tolerated by most of the patients. To this end, the present inventors applied a step-wise application of a CD33/CD3 bispecific antibody construct as outlined herein. Thereby, adverse effects could be reduced in number, ameliorated and even prevented. The appropriate dosage can be selected by the clinician on the basis of efficacy, tolerability and safety with a minimum of adverse effects in the patient.

The dose of the GC that is to be used in accordance with the embodiments of the present invention is not limited, i.e. it will depend on the circumstances of the individual patient. GC can be administered intravenously or orally. Preferred dosages of the GC include, however, between 1 to 6 mg (dexamethasone equivalent) at the lower end of dosing to 40 mg (dexamethasone equivalent). Said dosage can be administered all at once or subdivided into smaller dosages. Preferred is a subdivide dose wherein one dose of GC is given prior to the infusion of the first and/or second dose according to the step dosing as described herein, and the other dose of GC is given prior to the administration of the second or third dose according to the step dosing as described herein. Hence, GC is preferably two times dosed per treatment cycle. Even more preferably, GC is administered one 24 or 8 h or 4 h or 1 h before the beginning of a treatment cycle or the beginning of the administration of the next higher dose within said treatment cycle. In this regard, 1 h is most preferred. The dose is 1 to 40 mg each, preferably 5 to 20 mg, most preferably 8 mg each. “d” denotes one day. Further dosage regimens are derivable from the appended examples. All dosages given in this paragraph refer to dexamethasone equivalents.

The term “effective and non-toxic dose” as used herein refers to a tolerable dose of a bispecific construct which is high enough to cause depletion of pathologic cells, tumor elimination, tumor shrinkage or stabilization of disease without or essentially without major toxic effects. Such effective and non-toxic doses may be determined e.g. by dose escalation

studies described in the art and should be below the dose inducing severe adverse side events (dose limiting toxicity, DLT).

5 The term "toxicity" as used herein refers to the toxic effects of a drug manifested in adverse events or severe adverse events. These side events might refer to a lack of tolerability of the drug in general and/or a lack of local tolerance after administration. Toxicity could also include teratogenic or carcinogenic effects caused by the drug.

10 The term "safety", "*in vivo* safety" or "tolerability" as used herein defines the administration of a drug without inducing severe adverse events directly after administration (local tolerance) and during a longer period of application of the drug. "Safety", "*in vivo* safety" or "tolerability" can be evaluated e.g. at regular intervals during the treatment and follow-up period. Measurements include clinical evaluation, e.g. organ manifestations, and screening of laboratory abnormalities. Clinical evaluation may be carried out and deviations to normal
15 findings recorded/coded according to NCI-CTC and/or MedDRA standards. Organ manifestations may include criteria such as allergy/immunology, blood/bone marrow, cardiac arrhythmia, coagulation and the like, as set forth e.g. in the Common Terminology Criteria for adverse events v3.0 (CTCAE). Laboratory parameters which may be tested include for instance hematology, clinical chemistry, coagulation profile and urine analysis and
20 examination of other body fluids such as serum, plasma, lymphoid or spinal fluid, liquor and the like. Safety can thus be assessed e.g. by physical examination, imaging techniques (i.e. ultrasound, x-ray, CT scans, Magnetic Resonance Imaging (MRI), other measures with technical devices (i.e. electrocardiogram), vital signs, by measuring laboratory parameters and recording adverse events. For example, adverse events in non-chimpanzee primates in
25 the uses and methods according to the invention may be examined by histopathological and/or histochemical methods.

The above terms are also referred to e.g. in the Preclinical safety evaluation of biotechnology-derived pharmaceuticals S6; ICH Harmonised Tripartite Guideline; ICH
30 Steering Committee meeting on July 16, 1997.

The invention also provides a method for the treatment of myeloid leukemia comprising the administration of a therapeutically efficient amount of a bispecific construct comprising a first binding domain specifically binding to CD33 and a second binding domain specifically
35 binding to CD3 to a subject in need thereof, the method comprising the step of administering to said subject the construct for a maximal period of 14 days followed by a step without administration of the construct for a period of at least 14 days.

It is preferred for the method of the invention that the bispecific construct is administered according to a schedule comprising the following steps:

- (a) administration of a first dose of the bispecific construct, followed by
- (b) administration of a second dose of the bispecific construct, wherein said second dose exceeds said first dose, optionally followed by
- (c) administration of a third dose of the bispecific construct, wherein said optional third dose exceeds said second dose.

In a preferred embodiment of the method of the invention the period of administration of the first dose is up to seven days.

Also in a preferred embodiment of the method of the invention the complete period of administration of the dose steps subsequent to the first dose is in the range of eight to thirteen days.

In line with one embodiment of the method of the invention the complete period of administration of the dose steps subsequent to the first dose is in the range of eight to thirteen days.

For a preferred embodiment of the method of the invention it is envisaged that the period of administration of the second dose is four to seven days and the period of administration of the third dose is four to seven days.

The method of the invention may preferably comprise two or more cycles of treatment which each comprises the maximal period of 14 days of bispecific construct administration followed by the period of at least 14 days without administration of the bispecific construct.

In a preferred embodiment of the method of the invention only the first cycle of the treatment comprises the administration according to step (a), whereas the following cycles start with the dose according to step (b).

It is preferred for the method of the invention that the first binding domain of the bispecific construct comprises groups of six CDRs selected from the group consisting of SEQ ID NOs: 10 to 12 and 14 to 16, 22 to 24 and 26 to 28, 34 to 36 and 38 to 40, 46 to 48 and 50 to 52, 58 to 60 and 62 to 64, 70 to 72 and 74 to 76, 82 to 84 and 86 to 88, 94 to 96 and 98 to 100.

Also in line with a preferred embodiment of the method of the invention the second binding domain of the bispecific construct comprises groups of six CDRs selected from the group consisting of SEQ ID NOs: 9 to 14, 27 to 32, 45 to 50, 63 to 68, 81 to 86, 99 to 104, 117 to 122, 135 to 140, 153 to 158 and 171 to 176 of WO 2008/119567.

In a preferred embodiment of the method of the invention the bispecific construct is a bispecific antibody construct.

Moreover, it is preferred for the method of the invention that the bispecific antibody construct is a single chain construct comprising an amino acid sequence selected from the group

consisting of SEQ ID NOs: 18, 19, 20, 30, 31, 32, 42, 43, 44, 54, 55, 56, 66, 67, 68, 78, 79, 80, 90, 91, 92, 102, 103, 104, 105, 106, 107 and 108.

In one embodiment of the method of the invention the bispecific construct is administered in combination with one or more epigenetic factors selected from the group consisting of
5 histone deacetylase (HDAC) inhibitors, DNA methyltransferase (DNMT) I inhibitors, hydroxyurea, Granulocyte-Colony Stimulating Factor (G-CSF), histone demethylase inhibitors and ATRA (All Trans-retinoic acid) and wherein:

- (a) the one or more epigenetic factors are administered prior to the administration of the bispecific construct;
- 10 (b) the one or more epigenetic factors are administered subsequent to the administration of the bispecific construct; or
- (c) the one or more epigenetic factors and the bispecific construct are administered simultaneously.

It is preferred for the method of the invention that the one or more epigenetic factors are
15 administered up to seven days prior to the administration of the bispecific construct.

For one embodiment of the method of the invention it is preferred that the epigenetic factor is hydroxyurea

As described herein above, in line with the present invention the myeloid leukemia is selected from the group consisting of acute myeloblastic leukemia, chronic neutrophilic
20 leukemia, myeloid dendritic cell leukemia, accelerated phase chronic myelogenous leukemia, acute myelomonocytic leukemia, juvenile myelomonocytic leukemia, chronic myelomonocytic leukemia, acute basophilic leukemia, acute eosinophilic leukemia, chronic eosinophilic leukemia, acute megakaryoblastic leukemia, essential thrombocytosis, acute erythroid leukemia, polycythemia vera, myelodysplastic syndrome, acute panmyeloic leukemia,
25 myeloid sarcoma, and acute biphenotypic leukaemia. It is preferred that the myeloid leukemia is an acute myeloid leukemia (AML).

Also in one embodiment the invention provides a use of a bispecific construct comprising a first binding domain specifically binding to CD33 and a second binding domain specifically
30 binding to CD3 for the preparation of a pharmaceutical composition for the treatment of myeloid leukemia, wherein the construct is to be administered for a maximal period of 14 days followed by a period of at least 14 days without administration of the construct.

It is preferred of the use of the invention that the bispecific construct is to be administered according to a schedule comprising the following steps:

- 35 (a) administration of a first dose of the bispecific construct, followed by
- (b) administration of a second dose of the bispecific construct, wherein said second dose exceeds said first dose, optionally followed by

(a) administration of a third dose of the bispecific construct, wherein said optional third dose exceeds said second dose.

In a preferred embodiment of the use of the invention the period of administration of the first dose is up to seven days.

5 Also in a preferred embodiment of the use of the invention the complete period of administration of the dose steps subsequent to the first dose is in the range of eight to thirteen days.

In line with one embodiment of the use of the invention the complete period of administration of the dose steps subsequent to the first dose is in the range of eight to thirteen days.

10 For a preferred embodiment of the method of the invention it is envisaged that the period of administration of the second dose is four to seven days and the period of administration of the third dose is four to seven days.

In one preferred embodiment of the invention the treatment of the myeloid leukemia comprise two or more cycles of treatment which each comprises the maximal period of 14
15 days of bispecific construct administration followed by the period of at least 14 days without administration of the bispecific construct.

In a preferred embodiment of the use of the invention only the first cycle of the treatment comprises the administration according to step (a), whereas the following cycles start with the dose according to step (b).

20 It is preferred for the use of the invention that the first binding domain of the bispecific construct comprises groups of six CDRs selected from the group consisting of SEQ ID NOs: 10 to 12 and 14 to 16, 22 to 24 and 26 to 28, 34 to 36 and 38 to 40, 46 to 48 and 50 to 52, 58 to 60 and 62 to 64, 70 to 72 and 74 to 76, 82 to 84 and 86 to 88, 94 to 96 and 98 to 100.

Also in line with a preferred embodiment of the use of the invention the second binding
25 domain of the bispecific construct comprises groups of six CDRs selected from the group consisting of SEQ ID NOs: 9 to 14, 27 to 32, 45 to 50, 63 to 68, 81 to 86, 99 to 104, 117 to 122, 135 to 140, 153 to 158 and 171 to 176 of WO 2008/119567.

In a preferred embodiment of the use of the invention the bispecific construct is a bispecific antibody construct.

30 Moreover, it is preferred for the use of the invention that the bispecific antibody construct is a single chain construct comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 18, 19, 20, 30, 31, 32, 42, 43, 44, 54, 55, 56, 66, 67, 68, 78, 79, 80, 90, 91, 92, 102, 103, 104, 105, 106, 107 and 108.

In one embodiment of the use of the invention the bispecific construct is administered in
35 combination with one or more epigenetic factors selected from the group consisting of histone deacetylase (HDAC) inhibitors, DNA methyltransferase (DNMT) I inhibitors,

hydroxyurea, Granulocyte-Colony Stimulating Factor (G-CSF), histone demethylase inhibitors and ATRA (All Trans-retinoic acid) and wherein:

- (a) the one or more epigenetic factors are administered prior to the administration of the bispecific construct;
- 5 (b) the one or more epigenetic factors are administered subsequent to the administration of the bispecific construct; or
- (c) the one or more epigenetic factors and the bispecific construct are administered simultaneously.

It is preferred for the use of the invention that the one or more epigenetic factors are administered up to seven days prior to the administration of the bispecific construct.

For one embodiment of the use of the invention it is preferred that the epigenetic factor is hydroxyurea

As described herein above, in line with the present invention the myeloid leukemia is selected from the group consisting of acute myeloblastic leukemia, chronic neutrophilic leukemia, myeloid dendritic cell leukemia, accelerated phase chronic myelogenous leukemia, acute myelomonocytic leukemia, juvenile myelomonocytic leukemia, chronic myelomonocytic leukemia, acute basophilic leukemia, acute eosinophilic leukemia, chronic eosinophilic leukemia, acute megakaryoblastic leukemia, essential thrombocytosis, acute erythroid leukemia, polycythemia vera, myelodysplastic syndrome, acute panmyeloic leukemia, myeloid sarcoma, and acute biphenotypic leukaemia. It is preferred that the myeloid leukemia is an acute myeloid leukemia (AML).

Brief description of the drawings

25 Figure 1:

The figure shows the hematopoietic cell types and their origin (progenitor cell types).

Source: Wikipedia

Figure 2:

30 Human CD33⁺ EOL-1 cells or CD33⁻negative Kato III cells were co-cultured with human peripheral blood mononuclear cells (PBMCs) at an effector-to-target (E:T) cell ratio of 5:1 and AMG 330 for 48 h. Cell lysis was monitored by flow cytometric determination of PI uptake (A). T cell activation was analyzed by determination of CD69 surface expression (B). Each data point represents the mean of triplicate measurements. Error bars represent the standard error of the mean.

Figure 3:

HEK293-huCD33, HEK293-cyCD33 or HEK293 cells were co-cultured with cynomolgus monkey PBMCs (E:T cell ratio 5:1) and increasing concentrations of AMG 330 for 48 h. Target cell lysis was monitored by flow cytometric determination of PI uptake (A). Surface expression of CD69 on T cells was detected by flow cytometry using antigen-specific
5 fluorochrome-conjugated monoclonal antibodies (B). Each data point represents the mean of duplicates (A) or triplicate wells (B). Error bars represent standard errors of the mean.

It should be understood that the inventions herein are not limited to particular methodology, protocols, or reagents, as such can vary. The discussion and examples provided herein are
10 presented for the purpose of describing particular embodiments only and are not intended to limit the scope of the present invention, which is defined solely by the claims.

All publications and patents cited throughout the text of this specification (including all patents, patent applications, scientific publications, manufacturer's specifications,
15 instructions, etc.), whether supra or infra, are hereby incorporated by reference in their entirety. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention. To the extent the material incorporated by reference contradicts or is inconsistent with this specification, the specification will supersede any such material.

20

Examples:

The following examples are provided for the purpose of illustrating specific embodiments or features of the present invention. These examples should not be construed as to limit the
25 scope of this invention. The examples are included for purposes of illustration, and the present invention is limited only by the claims.

Example 1:

AMG 330-mediated Redirected Lysis and T Cell Activation

30 The activity of AMG 330 requires the simultaneous binding to both target cells and T cells. The pharmacological effect of AMG 330 is mediated by specific redirection of previously primed cytotoxic CD8⁺ or CD4⁺ T lymphocytes to kill CD33⁺ cells (Laszlo et al, 2014 Blood, 123 (4): 554-561; Krupka et al, 2014 Blood, 123 (3): 356-365; Friedrich et al, 2014 Mol Cancer Ther, 13 (6): 1549-1557; Aigner et al, 2013 Leukemia, 27 (5): 1107-1115). AMG 330
35 is a potent molecule showing half-maximal lysis of AML cell lines by human effector cell in vitro over a range of 24 to 200 pg/mL (0.4 to 3.7 pM) (Friedrich et al, 2014).

In vitro and ex vivo experiments demonstrated that AMG 330 can also recruit and activate cynomolgus monkey T cells; EC₅₀ values for redirected lysis of CD33⁺ tumor cell lines are slightly higher compared with human (79 to 254 pg/mL [1.5 to 4.7 pM]). A maximal 3-fold higher potency was observed for human compared with cynomolgus monkey effector cells.

5 The binding and bioactivity data validate the cynomolgus monkey as a relevant species for toxicity assessment.

AMG 330 induced activation of human as well as of cynomolgus monkey T cells and mediated redirected lysis of CD33⁺ cells in a dose-dependent manner.

10 The specificity of AMG 330 activity was verified using CD33⁺ human EOL-1 cells or CD33⁻ negative human Kato III cells. AMG 330 selectively mediated redirected lysis of EOL-1 cells by human T cells, while viability of Kato III remained unaltered, even when exposed to high AMG 330 concentrations (Figure 2A). In addition, AMG 330 increased the surface expression of the activation marker CD69 on T cells only in the presence of CD33⁺ cells, while no significant increase in CD69⁺ T cells was observed in co-cultures with CD33⁻ Kato III cells (Figure 2B).

15 AMG 330 selectively mediated redirected lysis of HEK293 cells stably expressing either human (huCD33) or cynomolgus monkey CD33 (cyCD33) by cynomolgus monkey effector cells, while viability of target-negative cells was not affected (Figure 3A). Cynomolgus monkey T cell activation (CD69) by AMG 330 was highly specific and was not observed in the absence of CD33-expressing cells (Figure 3B).

20 AMG 330 activity was investigated using a panel of CD33⁺ human AML tumor cell lines expressing human CD33. PBMCs or isolated CD3⁺ T cells of healthy human donors were co-cultivated with CD33-expressing target cells in the presence of increasing AMG 330 concentrations and specific cell lysis was determined by flow cytometric analyses. AMG 330-mediated half-maximal lysis (EC₅₀) of AML cell lines by human effector cells in vitro ranged from 24 to 200 pg/mL (0.4 to 3.7 pM) (Table 1).

Table 1. EC_x Values of AMG 330-mediated Redirected Lysis by Human T Cells

<i>Cell line</i>	<i>EC50 [pg/mL]</i>		<i>EC90 [pg/mL]</i>		<i>Amplitude [%]</i>		<i>N</i>
	<i>Mean</i>	<i>± SEM</i>	<i>Mean</i>	<i>± SEM</i>	<i>Mean</i>	<i>± SEM</i>	
KG-1	23.6	± 4.1	62.8	± 10.2	66.4	± 1.9	16
HL-60	25.1	± 4.1	64.9	± 11.8	58.4	± 4.3	11
THP-1	29.3	± 6.3	150.3	± 45.4	58.5	± 5.6	4
MOLM-13	32.5	± 9.1	91.2	± 43.5	61.5	± 5.6	9
U937	34.1	± 10.9	56.5	± 10.9	71.8	± 5.1	4
KASUMI-1	60.4	± 10.3	250.4	± 43.9	55.4	± 3.9	4

SKM-1	63.4	± 31.1	163.5	± 82.0	62.5	± 9.3	5
NB-4	123.6	± 44.0	335.4	± 151.3	70.9	± 2.5	5
M-07e	174.8	± 62.7	261.2	± 71.4	77.6	± 0.4	4
EOL-1	180.7	± 44.6	1605.2	± 447.8	78.1	± 3.7	6
NOMO-1	199.2	± 36.9	813.4	± 231.5	57.6	± 2.2	5

Differences between cell lines are most likely due to variation in the amount of CD33 cell surface expression. A quantitative correlation between target expression and EC50 values was observed for AMG 330 cytotoxicity, with increasing levels of CD33 expression resulting in lower EC50 values (Laszlo et al, 2014, loc cit.).

Example 2:

Pharmacokinetics of AMG 330

The PK of AMG 330 was characterized after a single IV bolus injection of 300 µg/kg or 900 µg/kg, a single SC bolus injection of 900 µg/kg, and a single IP bolus injection of 900 µg/kg to BALB/c mice. The non-compartmental PK parameters of AMG 330 after single dose IV, SC, and IP administration are shown in Table 2. AMG 330 serum concentrations were quantifiable up to 24 hours post IV administration. The half-life, clearance, and volume of distribution were similar in both dose groups, indicating linear pharmacokinetics of AMG 330. Area under the concentration-time curve from time 0 to infinity (AUC_{inf}) and C_{max} increased in a dose-linear manner from 300 µg/kg to 900 µg/kg. After SC administration, a C_{max} of 210.3 ng/mL was reached 2 hours post dose and AMG 330 was quantifiable in serum up to 36 hours. Compared with AUC_{inf} from the 900 µg/kg IV bolus PK, the absolute SC bioavailability was 22%. The C_{max} of 2201 ng/mL was achieved 2 hours after IP administration and AMG 330 was quantifiable for up to 36 hours. The absolute IP bioavailability was 99%, based on comparison to the AUC_{inf} after a 900 µg/kg IV bolus. The apparent elimination half-life was similar for all routes of administration and ranged between 6.5 - 8.7 hours (Table 2).

Table 2. Mean Pharmacokinetic Parameters after Single Dose IV, SC, or IP Administration to BALB/c Mice

Dose (µg/kg)	Route	t_{max} (hr)	C_{max} (ng/mL)	$t_{1/2}$ (hr)	AUC_{inf} (ng*hr/mL)	Vd/F (mL/kg)	CL/F (mL/hr/kg)	F (%)
300	IV	Na	5614	6.5	4468	201.7	67.2	NA
900	IV	Na	16092	7.6	15788	236.4	57.0	na
900	SC	2	210.3	8.7	3536	3184.7	254.5	22
900	IP	2	2201.3	7.5	15557	621.8	57.9	99

t_{max} = time of maximum concentration; C_{max} = maximum concentration; $t_{1/2}$ = terminal half-life; AUC_{inf} = area under the concentration-time curve from time 0 to infinity; Vd = volume of distribution; Cl = clearance; F = bioavailability; na = not applicable

Example 3:**Predicted AMG 330 Concentrations in Humans**

The human PK of AMG 330 was predicted based on previous experience with other BiTE[®] molecules in development. The clinical PK of blinatumomab, AMG 211, and AMG 110 were similar (Table 3), with half-lives ranging from 2.1 – 4.4 hours based on noncompartmental analyses of the clinical concentration-time data. The PK of AMG 330 is expected to be consistent with those data, as the structure and molecular weight of these compounds are similar and they have a similar mechanism of action; all of these BiTE[®] molecules have one arm that targets CD3⁺ T cells, with the other arm targeting CD19, CEA, or EpCAM for blinatumomab, AMG 211, and AMG 110, respectively. Because AMG 330 is intended to be administered via cIV infusion to treat patients with acute myeloid leukemia (AML), a liquid tumor indication, and blinatumomab is administered via cIV infusion to treat ALL, also a liquid tumor indication, the human PK parameters of AMG 330 were derived from blinatumomab. CL and Vd for AMG 330 were assumed to be 2920 mL/hr and 4500 mL, respectively, with the PK parameter estimates obtained from non-compartmental analysis of the PK data from 4 studies in patients with non-Hodgkin's lymphoma (NHL), MRD⁺ ALL, and relapsed/refractory (R/R) ALL receiving cIV infusion of blinatumomab. Although blinatumomab and AMG 330 target different cell receptors (CD19 on B-cells vs CD33 on hematopoietic cells), the prior clinical experience with other BiTE[®] molecules does not suggest any target mediated drug disposition (TMDD) in the dosing ranges tested. Thus, the assumption of linear PK for AMG 330 is consistent with observed linear PK of other clinically tested BiTE[®] molecules.

Based on these PK parameters, the predicted half-life of AMG 330 is 1 hour; half-life in cynomolgus monkeys after cIV administration was similarly short and ranged from 1.6 – 2.7 hours, although half-life was longer after SC administration and ranged from 4 – 7 hours. Furthermore, blinatumomab PK parameters were also used to predict the PK of AMG 110, which is being clinically investigated for the treatment of metastatic solid tumors. The predicted half-life of AMG 110 was within approximately 2-fold of the observed half-life. The clearance of AMG 110 was approximately 4-fold lower than expected based on blinatumomab PK, but still within the range of clearances observed for blinatumomab across the 4 clinical studies from which the PK parameters were derived; mean clearance values ranged from 1.81 - 3.36 L/hr with variability ranging from 32% - 103% (as measured by the coefficient of variation).

35

Table 3. Mean Clinical PK Parameters of Blinatumomab, AMG 110, and AMG 211 Based on Non-Compartmental Analyses

Compound	Cl (L/hr)	V _z (L)	t _{1/2,z} (hr)	Reference
Blinatumomab	2.92.	4.5	2.1	MT103-104, MT103-202, MT103-206, and MT103-211 clinical study reports
AMG 110	0.65	4.0	4.4	AMG 110 Investigator's Brochure
AMG 211	2.3	8.0	2.9	AMG 211 Investigator's Brochure

Predicted human exposure (as measured by C_{ss}) and exposure margins after cIV infusion of AMG 330 at the proposed FIH doses are presented in Table 4. The exposure margins are based on the average C_{ss} in male and female cynomolgus monkeys of 8.36 ng/mL at the HNSTD of 10 µg/kg/day cIV and are adjusted for potency differences between human and monkey effector cells; AMG 330 was 3-fold more potent in human than monkey effector cells

Table 4. Predicted Human Exposures and Exposure Margins at Steady-State Relative to the HNSTD after cIV Infusion of AMG 330

AMG 330 Dose (µg/day)	Predicted Human C _{ss} (ng/mL)	Predicted Exposure Margins ^{a,b,c} (HNSTD = 10 µg/kg/day)
0.5	0.0071	391
2.5	0.036	78
10.0	0.14	20
30.0	0.43	6.7
60.0	0.86	3.3
120.0	1.7	1.6
240.0	3.4	0.8
480.0	6.8	0.4
720.0	10.3	0.3
960.0	13.7	0.2

^aAverage C_{ss} in male and female monkeys at HNSTD divided by predicted human C_{ss}

^bAverage C_{ss} was 8.36 ng/mL at 10 µg/kg/day.

^cCorrected for 3-fold greater potency in human vs monkey effector cells

15

SEQ ID NO.	DESIGNATION	SOURCE	SEQUENCE
1	Human CD3ε extracellular domain	human	QDGNEEMGGITQTPYKVSISGTTVILTCPQYPGSEILWQHNDKNIIGDEDDKNIIGDEDDKNIIGDEDDHLSLKEFFSELE QSGYVVCYPRGSKPEDANFYLYLRARVCENCMEMD
2	Human CD3ε 1-27	human	QDGNEEMGGITQTPYKVSISGTTVILT
3	<i>Callithrix jacchus</i> CD3ε extracellular domain	<i>Callithrix jacchus</i>	QDGNEEMGDTTQNPKVSIISGTTVTLTTPRYDGHEIKWLNVSNQKKEGHEHLLLEDFSEMEQSGYYACL KETPAEEASHLYLYLKARVCENCVEVD
4	<i>Callithrix jacchus</i> CD3ε 1-27	<i>Callithrix jacchus</i>	QDGNEEMGDTTQNPKVSIISGTTVTLT
5	<i>Saguinus oedipus</i> CD3ε extracellular domain	<i>Saguinus oedipus</i>	QDGNEEMGDTTQNPKVSIISGTTVTLTTPRYDGHEIKWLNVSNQKKEGHEHLLLEDFSEMEQSGYYACL KETPAEEASHLYLYLKARVCENCVEVD
6	<i>Saguinus oedipus</i> CD3ε 1-27	<i>Saguinus oedipus</i>	QDGNEEMGDTTQNPKVSIISGTTVTLT
7	<i>Saimiri sciureus</i> CD3ε extracellular domain	<i>Saimiri sciureus</i>	QDGNEEIGDTTQNPKVSIISGTTVTLTTPRYDGQEIKWLVNDQNKKEGHEHLLLEDFSEMEQSGYYACL KETPTEEASHLYLYLKARVCENCVEVD
8	<i>Saimiri sciureus</i> CD3ε 1-27	<i>Saimiri sciureus</i>	QDGNEEIGDTTQNPKVSIISGTTVTLT
9	CD33 VH of AH3	artificial	QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWRQAPGQGLEWMGWINTYTGEPTAYDDFKGRVTMSDSTSTS TAYLEINSLRSDDTAIYYCARWSWSDGYVYFDYWGQGTVTVTVSS
10	CD33 HCDR1 of AH3	artificial	NYGMN
11	CD33 HCDR2 of AH3	artificial	WINTYTGEPTAYDDFKG
12	CD33 HCDR3 of AH3	artificial	WSWSDGYVYFDY
13	CD33 VL of AH3	artificial	DIVMTQSPDSTITVSLIGERTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGSGTDF TLTIDSLQPEDSATYYCQQSAHFPIITFGQGTREIK
14	CD33 LCDR1 of AH3	artificial	KSSQSVLDSKKNKNSLA
15	CD33 LCDR2 of AH3	artificial	WASTRES
16	CD33 LCDR3 of AH3	artificial	QQSAHFPIIT
17	CD33 CD33 HL of AH3	artificial	QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWRQAPGQGLEWMGWINTYTGEPTAYDDFKGRVTMSDSTSTS TAYLEINSLRSDDTAIYYCARWSWSDGYVYFDYWGQGTVTVTVSSGGGSGGGGGSDIVMTQSPDSTITVSLIGE RTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSLQPEDSATYYC

18	CD33 AH3 HL x H2C HL	artificial	QQSAHFPIITFGQTRLEIK QVQLVQSGAEVKKPQGSVKVSCKASGYTFITNYGMNWVRAQPGQGLEWMMGWINTYTGEPTIYADDFKGRVTMSSDSTSTS TAYLEINSLRSDDTAIYYCARWSWSDGYVYFDYWGQGTITVTVSSGGGSGGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGGTDFTLTIDSLQPEDSATIYC QQSAHFPIITFGQTRLEIKSGGGSEVQLVEGGGLVQPGGSLKLSAAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLIKTEDTAVYYCVRHGNFGNSIYSWAYWGQGTITVTVSSGGGG SGGGSGGGGQTVVTEPEPSLTVSPGGTITLTCGSSTGAVTSGYYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSG SLGGKAALTLISGVQPEDEAEYICVILWYSNRWVFGGGTKLITVL
19	CD33 AH3 HL x F12Q HL	artificial	QVQLVQSGAEVKKPQGSVKVSCKASGYTFITNYGMNWVRAQPGQGLEWMMGWINTYTGEPTIYADDFKGRVTMSSDSTSTS TAYLEINSLRSDDTAIYYCARWSWSDGYVYFDYWGQGTITVTVSSGGGSGGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGGTDFTLTIDSLQPEDSATIYC QQSAHFPIITFGQTRLEIKSGGGSEVQLVEGGGLVQPGGSLKLSAAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLIKTEDTAVYYCVRHGNFGNSIYSWAYWGQGTITVTVSSGGGG SGGGSGGGGQTVVTEPEPSLTVSPGGTITLTCGSSTGAVTSGYYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSG SLGGKAALTLISGVQPEDEAEYICVILWYSNRWVFGGGTKLITVL
20	CD33 AH3 HL x I2C HL	artificial	QVQLVQSGAEVKKPQGSVKVSCKASGYTFITNYGMNWVRAQPGQGLEWMMGWINTYTGEPTIYADDFKGRVTMSSDSTSTS TAYLEINSLRSDDTAIYYCARWSWSDGYVYFDYWGQGTITVTVSSGGGSGGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGGTDFTLTIDSLQPEDSATIYC QQSAHFPIITFGQTRLEIKSGGGSEVQLVEGGGLVQPGGSLKLSAAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLIKTEDTAVYYCVRHGNFGNSIYSWAYWGQGTITVTVSSGGGG SGGGSGGGGQTVVTEPEPSLTVSPGGTITLTCGSSTGAVTSGYYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSG SLGGKAALTLISGVQPEDEAEYICVILWYSNRWVFGGGTKLITVL
21	CD33 VH of AF5	artificial	QVQLVQSGAEVKKPGASVKVSCKASGYTFITNYGMNWVRAQPGQGLEWMMGWINTYTGEPTIYADDFKGRVTMTSDTSTS TAYLELHNLRSDDTAVYYCARWSWSDGYVYFDYWGQGTITVTVSS
22	CD33 HCDR1 of AF5	artificial	NYGMN
23	CD33 HCDR2 of AF5	artificial	WINTYTGEPTIYADDFKFG
24	CD33 HCDR3 of AF5	artificial	WSWSDGYVYFDY
25	CD33 VL of AF5	artificial	DIVMTQSPDSLTVSLGERTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGGTDF TLTIDSLQPEDSATIYCQSAHFPIITFGQTRLEIK
26	CD33 LCDR1 of AF5	artificial	KSSQSVLDSKKNKNSLA
27	CD33 LCDR2 of AF5	artificial	WASTRES
28	CD33 LCDR3 of AF5	artificial	QQSAHFPIIT
29	CD33HL of AF5	artificial	QVQLVQSGAEVKKPGASVKVSCKASGYTFITNYGMNWVRAQPGQGLEWMMGWINTYTGEPTIYADDFKGRVTMTSDTSTS

30	CD33 AF5 HL x H2C HL	artificial	<p>TAYLELHNLRSDDTAVYYCARMSWSDGYVYFDYWGQGTTVTVSSGGGGGGGGGGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSKKNKNSLAWYQOKPGQPPKLLLSWASTRESGIPDRFSGSGGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGTREIK</p> <p>QVQLVQSGAEVKKPKGASVKVCKASGYTFITNYGMNWVKQAPGQGLKWMGWINTYTGETTYADDFKGRVTMTSDTSTS TAYLELHNLRSDDTAVYYCARMSWSDGYVYFDYWGQGTTVTVSSGGGGGGGGGGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSKKNKNSLAWYQOKPGQPPKLLLSWASTRESGIPDRFSGSGGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGTREIKSGGGGSEVQLVEVSGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLIKTEDEAVYYCVRHGNFGNSYISWAYWGQGTLLVTVSSGGGG SGGGGGGGGQTVVITQEPSTLTVSPGGTVTLTCGSSTGAVTSGYYPNWVQOKPGQAPRGLIGGTFKFLAPGTPARFSG SLLGGKAALTLISGVQPEDEAEYCVLWYSNRWVFGGGTKLITVL</p>
31	CD33 AF5 HL x F12Q HL	artificial	<p>QVQLVQSGAEVKKPKGASVKVCKASGYTFITNYGMNWVKQAPGQGLKWMGWINTYTGETTYADDFKGRVTMTSDTSTS TAYLELHNLRSDDTAVYYCARMSWSDGYVYFDYWGQGTTVTVSSGGGGGGGGGGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSKKNKNSLAWYQOKPGQPPKLLLSWASTRESGIPDRFSGSGGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGTREIKSGGGGSEVQLVEVSGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLIKTEDEAVYYCVRHGNFGNSYISWAYWGQGTLLVTVSSGGGG SGGGGGGGGQTVVITQEPSTLTVSPGGTVTLTCGSSTGAVTSGYYPNWVQOKPGQAPRGLIGGTFKFLAPGTPARFSG SLLGGKAALTLISGVQPEDEAEYCVLWYSNRWVFGGGTKLITVL</p>
32	CD33 AF5 HL x 12C HL	artificial	<p>QVQLVQSGAEVKKPKGASVKVCKASGYTFITNYGMNWVKQAPGQGLKWMGWINTYTGETTYADDFKGRVTMTSDTSTS TAYLELHNLRSDDTAVYYCARMSWSDGYVYFDYWGQGTTVTVSSGGGGGGGGGGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSKKNKNSLAWYQOKPGQPPKLLLSWASTRESGIPDRFSGSGGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGTREIKSGGGGSEVQLVEVSGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLIKTEDEAVYYCVRHGNFGNSYISWAYWGQGTLLVTVSSGGGG SGGGGGGGGQTVVITQEPSTLTVSPGGTVTLTCGSSTGAVTSGYYPNWVQOKPGQAPRGLIGGTFKFLAPGTPARFSG SLLGGKAALTLISGVQPEDEAEYCVLWYSNRWVFGGGTKLITVL</p>
33	CD33 VH of AC8	artificial	<p>QVQLVQSGAEVKKPGEISVKVCKASGYTFITNYGMNWVKQAPGQGLKWMGWINTYTGETTYADDFKGRVTMTSDTSTS TAYMEIRNLRNDDTAVYYCARMSWSDGYVYFDYWGQGTTVTVSS</p>
34	CD33 HCDR1 of AC8	artificial	<p>WINTYTGETTYADDFKKG NYGMN</p>
35	CD33 HCDR2 of AC8	artificial	<p>WINTYTGETTYADDFKKG</p>
36	CD33 HCDR3 of AC8	artificial	<p>WSWSDGYVYFDY</p>
37	CD33 VL of AC8	artificial	<p>DIVMTQSPDSLTVSLGERTTINCKSSQSVLDSKKNKNSLAWYQOKPGQPPKLLLSWASTRESGIPDRFSGSGGTDF TLTIDSLQPEDSATYYCQSAHFPIITFGQGTREIK</p>
38	CD33 LCDR1 of AC8	artificial	<p>KSSQSVLDSKKNKNSLA</p>
39	CD33 LCDR2 of AC8	artificial	<p>WASTRES</p>

40	CD33 LCDR3 of AC8	artificial	QQSAHFPIIT QVQLVQSGAEVKKPGEISVKVSCKASGYTFITNYGMNWKQAPGQGLKMMGWINTYTGEPYADDFKGRVTMTTDTSTST TAYMEIRNLRNDDTAVYYCARWSWSDGYVYFDYWGGQGTITVTVSSGGGGSGGGSDIVMTQSPDLSLTVSLGE RRTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIK
41	CD33 HL of AC8	artificial	QVQLVQSGAEVKKPGEISVKVSCKASGYTFITNYGMNWKQAPGQGLKMMGWINTYTGEPYADDFKGRVTMTTDTSTST TAYMEIRNLRNDDTAVYYCARWSWSDGYVYFDYWGGQGTITVTVSSGGGGSGGGSDIVMTQSPDLSLTVSLGE RRTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNILKTEDTAVYYCVRHGNFGNSYISWAYWGQGLTVTVSSGGGG SGGGSGGGGQTVVVTQEPSTLTVSPGGTTLTTCGSSTGAVTSGYYPNWWQQKPGQAPRGLIGGTKFLAPGTPARFSG SLGGKAALTLISGVQPEDEAEYYCALWYSNRWVFGGTKLITVL
42	CD33 AC8 HL x H2C HL	artificial	QVQLVQSGAEVKKPGEISVKVSCKASGYTFITNYGMNWKQAPGQGLKMMGWINTYTGEPYADDFKGRVTMTTDTSTST TAYMEIRNLRNDDTAVYYCARWSWSDGYVYFDYWGGQGTITVTVSSGGGGSGGGSDIVMTQSPDLSLTVSLGE RRTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNILKTEDTAVYYCVRHGNFGNSYISWAYWGQGLTVTVSSGGGG SGGGSGGGGQTVVVTQEPSTLTVSPGGTTLTTCGSSTGAVTSGYYPNWWQQKPGQAPRGLIGGTKFLAPGTPARFSG SLGGKAALTLISGVQPEDEAEYYCALWYSNRWVFGGTKLITVL
43	CD33 AC8 HL x F12Q HL	artificial	QVQLVQSGAEVKKPGEISVKVSCKASGYTFITNYGMNWKQAPGQGLKMMGWINTYTGEPYADDFKGRVTMTTDTSTST TAYMEIRNLRNDDTAVYYCARWSWSDGYVYFDYWGGQGTITVTVSSGGGGSGGGSDIVMTQSPDLSLTVSLGE RRTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNILKTEDTAVYYCVRHGNFGNSYISWAYWGQGLTVTVSSGGGG SGGGSGGGGQTVVVTQEPSTLTVSPGGTTLTTCGSSTGAVTSGYYPNWWQQKPGQAPRGLIGGTKFLAPGTPARFSG SLGGKAALTLISGVQPEDEAEYYCVLWYSNRWVFGGTKLITVL
44	CD33 AC8 HL x I2C HL	artificial	QVQLVQSGAEVKKPGEISVKVSCKASGYTFITNYGMNWKQAPGQGLKMMGWINTYTGEPYADDFKGRVTMTTDTSTST TAYMEIRNLRNDDTAVYYCARWSWSDGYVYFDYWGGQGTITVTVSSGGGGSGGGSDIVMTQSPDLSLTVSLGE RRTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNILKTEDTAVYYCVRHGNFGNSYISWAYWGQGLTVTVSSGGGG SGGGSGGGGQTVVVTQEPSTLTVSPGGTTLTTCGSSTGAVTSGYYPNWWQQKPGQAPRGLIGGTKFLAPGTPARFSG SLGGKAALTLISGVQPEDEAEYYCVLWYSNRWVFGGTKLITVL
45	CD33 VH of AH11	artificial	QVQLVQSGAEVKKPGEISVKVSCKASGYTFITNYGMNWKQAPGQGLKMMGWINTYTGEPYADDFKGRVTMTTDTSTST TAYMEIRNLRNDDTAVYYCARWSWSDGYVYFDYWGGQGTITVTVSSGGGGSGGGSDIVMTQSPDLSLTVSLGE RRTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNILKTEDTAVYYCVRHGNFGNSYISWAYWGQGLTVTVSSGGGG SGGGSGGGGQTVVVTQEPSTLTVSPGGTTLTTCGSSTGAVTSGYYPNWWQQKPGQAPRGLIGGTKFLAPGTPARFSG SLGGKAALTLISGVQPEDEAEYYCVLWYSNRWVFGGTKLITVL
46	CD33 HCDR1 of AH11	artificial	NYGMN
47	CD33 HCDR2 of AH11	artificial	WINTYTGEPYADDFKKG
48	HCDR3	artificial	WSWSDGYVYFDY
49	CD33 VL of AH11	artificial	DIVMTQSPDLSLTVSLGERTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSGTDF TLTIDSLQPEDSATYYCQSAHFPIITFGQGRLEIK
50	CD33 LCDR1 of AH11	artificial	KSSQSVLDSKKNKNSLA

51	CD33 LCDR2 of AH11	artificial	WASTRES	QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWVKQAPGQGLKWMGWINTYTGEPYADDKFKGRVTMTSDTSTS TAYMEISLRSDDTAVYYCARWSDGYYVYFDYWGGQTTVTVSSGGGGSGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIK
52	CD33LCDR3 of AH11	artificial	QQSAHFPIIT	QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWVKQAPGQGLKWMGWINTYTGEPYADDKFKGRVTMTSDTSTS TAYMEISLRSDDTAVYYCARWSDGYYVYFDYWGGQTTVTVSSGGGGSGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIK
53	CD33 HL of AH11	artificial		QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWVKQAPGQGLKWMGWINTYTGEPYADDKFKGRVTMTSDTSTS TAYMEISLRSDDTAVYYCARWSDGYYVYFDYWGGQTTVTVSSGGGGSGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIK
54	CD33 AH11 HL x H2C HL	artificial		QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWVKQAPGQGLKWMGWINTYTGEPYADDKFKGRVTMTSDTSTS TAYMEISLRSDDTAVYYCARWSDGYYVYFDYWGGQTTVTVSSGGGGSGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNILKTEDTAVYYCVRHGNFGNSYISWAYWGQGLTVTVSSGGGG SGGGSGGGSTVVTQEPSTLTVSPGGTTLTCGSSTGAVTSGYYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSG SLGGKAALTLISGVQPEDEAEYCVLWYSNRWVFGGKTLITVL
55	CD33 AH11 HL x F12Q HL	artificial		QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWVKQAPGQGLKWMGWINTYTGEPYADDKFKGRVTMTSDTSTS TAYMEISLRSDDTAVYYCARWSDGYYVYFDYWGGQTTVTVSSGGGGSGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNILKTEDTAVYYCVRHGNFGNSYISWAYWGQGLTVTVSSGGGG SGGGSGGGSTVVTQEPSTLTVSPGGTTLTCGSSTGAVTSGYYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSG SLGGKAALTLISGVQPEDEAEYCVLWYSNRWVFGGKTLITVL
56	CD33 AH11 HL x I2C HL	artificial		QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWVKQAPGQGLKWMGWINTYTGEPYADDKFKGRVTMTSDTSTS TAYMEISLRSDDTAVYYCARWSDGYYVYFDYWGGQTTVTVSSGGGGSGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNILKTEDTAVYYCVRHGNFGNSYISWAYWGQGLTVTVSSGGGG SGGGSGGGSTVVTQEPSTLTVSPGGTTLTCGSSTGAVTSGYYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSG SLGGKAALTLISGVQPEDEAEYCVLWYSNRWVFGGKTLITVL
57	CD33 VH of B3	artificial		QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWVKQAPGQGLKWMGWINTYTGEPYADDKFKGRVTMTSDTSTS TAYMELRNLSDDTAVYYCARWSDGYYVYFDYWGGQTTVTVSS NYGMN
58	CD33 HCDR1 of B3	artificial		WINTYTGEPYADKFKQG
59	CD33 HCDR2 of B3	artificial		WSWSDGYVYVFDY
60	CD33 HCDR3 of B3	artificial		
61	CD33 VL of B3	artificial		DIVMTQSPDSMTVSLGERTTINCKSSQSVLDSSTNKNLSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSTDF TLTIDSLQPEDSATYYCQQSAHFPIITFGQGRLEIK

62	CD33 LCDR1 of B3	artificial	KSSQSVLDSSTNKNSLA
63	CD33 LCDR2 of B3	artificial	WASTRE
64	CD33LCDR3 of B3	artificial	QOQSAHFPIIT
65	CD33 HL of B3	artificial	QVQLVQSGAEVKKPQGSVKVSKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGETNYADKFKQGRVFTTSDTSTS TAYMELRNLKSDDTAVYYCARMSWSDGYVYFDYWGGQTTVTVSSGGGGGGGGGGSDIVMTQSPDMSMTVSLGE RTTINCKSSQSVLDSSTNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSLQPEDSATYYC QOQSAHFPIITFGQTRLDIKSGGGSEVQLVESSGGLVQPGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLIKTEDTAVYYCVRHGNFGNSYISWAYWGQGLVTVSSGGGG SGGGSGGGGQTVVTTQPEPSLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSG SLLGGKAALTLISGVQPEDEAEYCVLWYSNRWVFGGTTKLTIVL
66	CD33 B3 HL x H2C HL	artificial	QVQLVQSGAEVKKPQGSVKVSKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGETNYADKFKQGRVFTTSDTSTS TAYMELRNLKSDDTAVYYCARMSWSDGYVYFDYWGGQTTVTVSSGGGGGGGGGGSDIVMTQSPDMSMTVSLGE RTTINCKSSQSVLDSSTNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSLQPEDSATYYC QOQSAHFPIITFGQTRLDIKSGGGSEVQLVESSGGLVQPGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNLIKTEDTAVYYCVRHGNFGNSYISWAYWGQGLVTVSSGGGG SGGGSGGGGQTVVTTQPEPSLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSG SLLGGKAALTLISGVQPEDEAEYCVLWYSNRWVFGGTTKLTIVL
67	CD33 B3 HL x F12Q HL	artificial	QVQLVQSGAEVKKPQGSVKVSKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGETNYADKFKQGRVFTTSDTSTS TAYMELRNLKSDDTAVYYCARMSWSDGYVYFDYWGGQTTVTVSSGGGGGGGGGGSDIVMTQSPDMSMTVSLGE RTTINCKSSQSVLDSSTNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSLQPEDSATYYC QOQSAHFPIITFGQTRLDIKSGGGSEVQLVESSGGLVQPGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNLIKTEDTAVYYCVRHGNFGNSYISWAYWGQGLVTVSSGGGG SGGGSGGGGQTVVTTQPEPSLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSG SLLGGKAALTLISGVQPEDEAEYCVLWYSNRWVFGGTTKLTIVL
68	CD33 B3 HL x I2C HL	artificial	QVQLVQSGAEVKKPQGSVKVSKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGETNYADKFKQGRVFTTSDTSTS TAYMELRNLKSDDTAVYYCARMSWSDGYVYFDYWGGQTTVTVSSGGGGGGGGGGSDIVMTQSPDMSMTVSLGE RTTINCKSSQSVLDSSTNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSLQPEDSATYYC QOQSAHFPIITFGQTRLDIKSGGGSEVQLVESSGGLVQPGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLIKTEDTAVYYCVRHGNFGNSYISWAYWGQGLVTVSSGGGG SGGGSGGGGQTVVTTQPEPSLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSG SLLGGKAALTLISGVQPEDEAEYCVLWYSNRWVFGGTTKLTIVL
69	CD33 VH of F2	artificial	QVQLVQSGAEVKKPQGSVKVSKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGETNYADKFKQGRVFTTSDTSTS TAYMELRNLKSDDTAVYYCARMSWSDGYVYFDYWGGQTTVTVSSGGGGGGGGGGSDIVMTQSPDMSMTVSLGE RTTINCKSSQSVLDSSTNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSLQPEDSATYYC QOQSAHFPIITFGQTRLDIKSGGGSEVQLVESSGGLVQPGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLIKTEDTAVYYCVRHGNFGNSYISWAYWGQGLVTVSSGGGG SGGGSGGGGQTVVTTQPEPSLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSG SLLGGKAALTLISGVQPEDEAEYCVLWYSNRWVFGGTTKLTIVL
70	CD33 HCDR1 of F2	artificial	NYGMN
71	CD33 HCDR2 of F2	artificial	WINTYTGETNYADKFKQ
72	CD33 HCDR3 of F2	artificial	WSWSDGYVYFDY

73	CD33 VL of F2	artificial	DIVMTQSPDLSVSLGERTTINCKSSQSVLDSSTNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGSGTDF TLTIDSLQPEDSATYYCQSAHFPIITFGQGRLEIK
74	CD33 LCDR1 of F2	artificial	KSSQSVLDSSTNKNSLA
75	CD33 LCDR2 of F2	artificial	WASTRES
76	CD33 LCDR3 of F2	artificial	QQSAHFPIIT
77	CD33 HL of F2	artificial	QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGETNYADKFKQGRVFTTSDTSTS TAYMELRNLKSDDTAVYYCARWSWSDGYVYFDYWGGTITVTVSSGGSGGGGGSDIVMTQSPDLSVSLGE RRTINCKSSQSVLDSSTNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIK
78	CD33 F2 HL x H2C HL	artificial	QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGETNYADKFKQGRVFTTSDTSTS TAYMELRNLKSDDTAVYYCARWSWSDGYVYFDYWGGTITVTVSSGGSGGGGGSDIVMTQSPDLSVSLGE RRTINCKSSQSVLDSSTNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVESSGGLVQPGSLKLSCAASGTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLLKTEDTAVYYCVRHGNFGNSYISWAYWGQGLVTVSSGGGG SGGGSGGGGQTVVITQEPSTVSPGGTTLTCGSSTGAVTSGYYPNWVQQKPGQAPRGLIGGKFLAPGTPARFSG SLLGKKAALTLISGVQPEDEAEYCVLWYSNRWVFGGKTLTVL
79	CD33 F2 HL x F12Q HL	artificial	QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGETNYADKFKQGRVFTTSDTSTS TAYMELRNLKSDDTAVYYCARWSWSDGYVYFDYWGGTITVTVSSGGSGGGGGSDIVMTQSPDLSVSLGE RRTINCKSSQSVLDSSTNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVESSGGLVQPGSLKLSCAASGTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNLLKTEDTAVYYCVRHGNFGNSYISWAYWGQGLVTVSSGGGG SGGGSGGGGQTVVITQEPSTVSPGGTTLTCGSSTGAVTSGYYPNWVQQKPGQAPRGLIGGKFLAPGTPARFSG SLLGKKAALTLISGVQPEDEAEYCVLWYSNRWVFGGKTLTVL
80	CD33 F2 HL x I2C HL	artificial	QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGETNYADKFKQGRVFTTSDTSTS TAYMELRNLKSDDTAVYYCARWSWSDGYVYFDYWGGTITVTVSSGGSGGGGGSDIVMTQSPDLSVSLGE RRTINCKSSQSVLDSSTNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVESSGGLVQPGSLKLSCAASGTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLLKTEDTAVYYCVRHGNFGNSYISWAYWGQGLVTVSSGGGG SGGGSGGGGQTVVITQEPSTVSPGGTTLTCGSSTGAVTSGYYPNWVQQKPGQAPRGLIGGKFLAPGTPARFSG SLLGKKAALTLISGVQPEDEAEYCVLWYSNRWVFGGKTLTVL
81	CD33 VH of B10	artificial	QVQLVQSGAEVKKPGEISVKVSKKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGETNYADKFKQGRVFTTSDTSTS TAYMELRNLKSDDTAVYYCARWSWSDGYVYFDYWGGTITVTVSSGGSGGGGGSDIVMTQSPDLSVSLGE RRTINCKSSQSVLDSSTNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGGTDFTLTIDSLQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVESSGGLVQPGSLKLSCAASGTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLLKTEDTAVYYCVRHGNFGNSYISWAYWGQGLVTVSSGGGG SGGGSGGGGQTVVITQEPSTVSPGGTTLTCGSSTGAVTSGYYPNWVQQKPGQAPRGLIGGKFLAPGTPARFSG SLLGKKAALTLISGVQPEDEAEYCVLWYSNRWVFGGKTLTVL
82	CD33 HCDR1 of B10	Artificial	NYGMN
83	CD33HCDR2 of B10	artificial	WINTYTGETTYADKFKQ

84	CD33 HCDR3 of B10	artificial	WSWSDGYVYVFDY
85	CD33 VL of B10	artificial	DIVMTQSPDLSLTVSLGERTTINCKSSQSVLDSNNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGSGTDF TLTIDGLQPEDSATYCYQQSAHFPIITFGQGRLEIK
86	CD33 LCDR1 of B10	artificial	KSSQSVLDSNNKNSLA
87	CD33 LCDR2 of B10	artificial	WASTRES
88	CD33 LCDR3 of B10	artificial	QQSAHFPIIT
89	CD33 HL of B10	artificial	QVQLVQSGAEVKKPGEISVKVSKKASGYTFTNYGMNWVKQAPGQGLEWMGWINTYTGEPYADKFKQGRVTMTTDTST TAYMEIRNLRSDDTAVYYCARWSDGYVYFDYWGQGTFTVTVSSGGSGGGGGSDIVMTQSPDLSLTVSLGE RRTINCKSSQSVLDSNNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDGLQPEDSATYCY QQSAHFPIITFGQGRLEIK
90	CD33 B10 HL x H2C HL	artificial	QVQLVQSGAEVKKPGEISVKVSKKASGYTFTNYGMNWVKQAPGQGLEWMGWINTYTGEPYADKFKQGRVTMTTDTST TAYMEIRNLRSDDTAVYYCARWSDGYVYFDYWGQGTFTVTVSSGGSGGGGGSDIVMTQSPDLSLTVSLGE RRTINCKSSQSVLDSNNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDGLQPEDSATYCY QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNILKTEDTAVYYCVRHGTFGNSYISYWAYWGQGTFLVTVSSGGGG SGGGSGGGGQTVVITQEPSTLTVSPGGTTLTCGSSTGAVTSGYYPNWWQQKPGQAPRGLIGGTFKFLAPGTPARFSG SLLGKKAALTLISGVQPEDEAEYCVLWYSNRWVFGGTKLITVL
91	CD33 B10 HL x F12Q HL	artificial	QVQLVQSGAEVKKPGEISVKVSKKASGYTFTNYGMNWVKQAPGQGLEWMGWINTYTGEPYADKFKQGRVTMTTDTST TAYMEIRNLRSDDTAVYYCARWSDGYVYFDYWGQGTFTVTVSSGGSGGGGGSDIVMTQSPDLSLTVSLGE RRTINCKSSQSVLDSNNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDGLQPEDSATYCY QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNILKTEDTAVYYCVRHGTFGNSYISYWAYWGQGTFLVTVSSGGGG SGGGSGGGGQTVVITQEPSTLTVSPGGTTLTCGSSTGAVTSGYYPNWWQQKPGQAPRGLIGGTFKFLAPGTPARFSG SLLGKKAALTLISGVQPEDEAEYCVLWYSNRWVFGGTKLITVL
92	CD33 B10 HL x I2C HL	artificial	QVQLVQSGAEVKKPGEISVKVSKKASGYTFTNYGMNWVKQAPGQGLEWMGWINTYTGEPYADKFKQGRVTMTTDTST TAYMEIRNLRSDDTAVYYCARWSDGYVYFDYWGQGTFTVTVSSGGSGGGGGSDIVMTQSPDLSLTVSLGE RRTINCKSSQSVLDSNNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDGLQPEDSATYCY QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNILKTEDTAVYYCVRHGTFGNSYISYWAYWGQGTFLVTVSSGGGG SGGGSGGGGQTVVITQEPSTLTVSPGGTTLTCGSSTGAVTSGYYPNWWQQKPGQAPRGLIGGTFKFLAPGTPARFSG SLLGKKAALTLISGVQPEDEAEYCVLWYSNRWVFGGTKLITVL
93	CD33 VH of E11	artificial	QVQLVQSGAEVKKPGEISVKVSKKASGYTFTNYGMNWVKQAPGQGLEWMGWINTYTGEPYADKFKQGRVTMTTDTST TAYMEIRNLGGDDTAVYYCARWSDGYVYFDYWGQGTSTVTVSS
94	CD33 HCDR1 of E11	artificial	NYGMN

95	CD33 HCDR2 of E11	artificial	WINTYTGEPITYADKFKQG
96	CD33 HCDR3 of E11	artificial	WSWSDGYVYFYDY
97	CD33 VL of E11	artificial	DIVMTQSPDSLTVSLGERTTINCKSSQSVLDSSTNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGSGTDF TLTIDSPQPEDSATYYCQSAHFPIITFGQGRLEIK
98	CD33 LCDR1 of E11	artificial	KSSQSVLDSSTNKNSLA
99	CD33 LCDR2 of E11	artificial	WASTRES
100	CD33 LCDR3 of E11	artificial	QQSAHFPIIT
101	CD33 HL of E11	artificial	QVQLVQSGAEVKKPGEISVKVSKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGEPITYADKFKQGRVTMTTDTSTS TAYMEIRNLGGDDTAVYYCARWSWDGYVYFYDYWGQGTSTVTVSSGGSGGGGGSDIVMTQSPDSLTVSLGE RRTINCKSSQSVLDSSTNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSPQPEDSATYYC QQSAHFPIITFGQGRLEIK
102	CD33 E11 HL x H2C HL	artificial	QVQLVQSGAEVKKPGEISVKVSKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGEPITYADKFKQGRVTMTTDTSTS TAYMEIRNLGGDDTAVYYCARWSWDGYVYFYDYWGQGTSTVTVSSGGSGGGGGSDIVMTQSPDSLTVSLGE RRTINCKSSQSVLDSSTNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSPQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLIKTEDTAVYYCVRHGNFNSIYSWAYWGQGTIVTVSSGGGG SGGGSGGGGQTVVTTQEPSTLTVSPGGTVTLTCGSSTGAVTSGYYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSG SLLGKKAALTLISGVQPEDEAEYCVLWYSNRWVFGGTKLITVL
103	CD33 E11 HL x F12Q HL	artificial	QVQLVQSGAEVKKPGEISVKVSKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGEPITYADKFKQGRVTMTTDTSTS TAYMEIRNLGGDDTAVYYCARWSWDGYVYFYDYWGQGTSTVTVSSGGSGGGGGSDIVMTQSPDSLTVSLGE RRTINCKSSQSVLDSSTNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSPQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKGRFTISRDDSKNTAYLQMNLIKTEDTAVYYCVRHGNFNSIYSWAYWGQGTIVTVSSGGGG SGGGSGGGGQTVVTTQEPSTLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSG SLLGKKAALTLISGVQPEDEAEYCVLWYSNRWVFGGTKLITVL
104	CD33 E11 HL x I2C HL	artificial	QVQLVQSGAEVKKPGEISVKVSKASGYTFITNYGMNWKQAPGQGLEWMGWINTYTGEPITYADKFKQGRVTMTTDTSTS TAYMEIRNLGGDDTAVYYCARWSWDGYVYFYDYWGQGTSTVTVSSGGSGGGGGSDIVMTQSPDSLTVSLGE RRTINCKSSQSVLDSSTNKNSLAWYQQKPGQPKLLLSWASTRESGIPDRFSGSGSGTDFTLTIDSPQPEDSATYYC QQSAHFPIITFGQGRLEIKSGGGSEVQLVEISGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLIKTEDTAVYYCVRHGNFNSIYSWAYWGQGTIVTVSSGGGG SGGGSGGGGQTVVTTQEPSTLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSG SLLGKKAALTLISGVQPEDEAEYCVLWYSNRWVFGGTKLITVL
105	CD33 UD H2C HL x AF5 HL	artificial	EVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSD KNTAYLQMNLIKTEDTAVYYCVRHGNFNSIYSWAYWGQGTIVTVSSGGSGGGGGQTVVTTQEPSTLTVSP

				<p>GGTDTLTCGSSGTAVTSGYYPNWQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAAALTLVSGVQPEDEAEYYCA LWYSNRWVFGGGLTLVLSGGGSSQVQLVQSGAEVKKPGASVKVCKASGYTFNYGMNWKQAPGQGLKMMGWINT YTGEPTYADDFKGRVTMTSDTSTAYLELHNLRSDDTAVYYCARWSWSDGYVYFDYWGQGTIVTVSSGGGSSGGG SGGGSDIVMTQSPDSLTVSLGERTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSG SGSGTDFTLTIDSLQPEDSATYCCQSAHFPIITFGQGRTRLEIK</p>
106	CD33 UD F12Q HL x AF5 HL	artificial		<p>EVQLVDSGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNIYATYADSVKGRFTISRDDSD KNTAYLQMNLLKTEDTAVYYCVRHGNFGNSYVSWWAYWGQGTIVTVSSGGGSGGGGSGGGGQTVVTVQEPSTLTVSP GGTDTLTCGSSGTAVTSGYYPNWQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAAALTLVSGVQPEDEAEYYCV LWYSNRWVFGGGLTLVLSGGGSSQVQLVQSGAEVKKPGASVKVCKASGYTFNYGMNWKQAPGQGLKMMGWINT YTGEPTYADDFKGRVTMTSDTSTAYLELHNLRSDDTAVYYCARWSWSDGYVYFDYWGQGTIVTVSSGGGSSGGG SGGGSDIVMTQSPDSLTVSLGERTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSG SGSGTDFTLTIDSLQPEDSATYCCQSAHFPIITFGQGRTRLEIK</p>
107	CD33 UD I2C HL x AF5 HL	artificial		<p>EVQLVDSGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNIYATYADSVKDRFTISRDDSD KNTAYLQMNLLKTEDTAVYYCVRHGNFGNSYVSWWAYWGQGTIVTVSSGGGSGGGGSGGGGQTVVTVQEPSTLTVSP GGTDTLTCGSSGTAVTSGYYPNWQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAAALTLVSGVQPEDEAEYYCV LWYSNRWVFGGGLTLVLSGGGSSQVQLVQSGAEVKKPGASVKVCKASGYTFNYGMNWKQAPGQGLKMMGWINT YTGEPTYADDFKGRVTMTSDTSTAYLELHNLRSDDTAVYYCARWSWSDGYVYFDYWGQGTIVTVSSGGGSSGGG SGGGSDIVMTQSPDSLTVSLGERTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSG SGSGTDFTLTIDSLQPEDSATYCCQSAHFPIITFGQGRTRLEIK</p>
108	CD33 E11 HL x I2C HL H6	artificial		<p>QVQLVQSGAEVKKPGEYSKVSCKASGYTFNYGMNWKQAPGQGLWMMGWINTYTGEPTYADKDFQGRVTMTTDTSTST TAYMEIRNLGGDDTAVYYCARWSWSDGYVYFDYWGQGTIVTVSSGGGSGGGGSGGGGSDIVMTQSPDSLTVSLGE RTTINCKSSQSVLDSSTNKNLSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGSGGSDTFLITIDSPQEDSATYCY QQSAHFPIITFGQGRTRLEIKSGGGSEVQLVQSGAEVKKPGASVKVCKASGYTFNYGMNWKQAPGKGLEWVARIR SKYNNIYATYADSVKDRFTISRDDSKNTAYLQMNLLKTEDTAVYYCVRHGNFGNSYVSWWAYWGQGTIVTVSSGGG SGGGSDIVMTQSPDSLTVSLGERTTINCKSSQSVLDSKKNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSG SLLGGKAAALTLVSGVQPEDEAEYYCVLWYSNRWVFGGGLTLVLSHHHHHH</p>
109	HALB	human		<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQCCPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLQHKDDNPNLPRLVREVDVMTAFHDNEETFLLKKYLYEIAARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAAACLLPKLDELDEGKASAKQRLKASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKYICENQDSISSKLECECEKPLLEKSHCIAEVENDEMPADLP SLAAD FVE SKDVCKNYAEAKDVFLGMFLYEVARRHPDYVSVLLLRLLAKTYETLEKCCAAAADPHECYAKVDFEFKPLVEEPQ NLIKONCELFELGELYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPKAEAKRMPCAEYDLSVVLNQLCV LHEKTPVSDRVTKCTESLVNRRPCFSALEVDETYVPKEFNAETFFHADICTLSEKERQIKKQTTALVELLVKHKPKA TKEQLKAVMDDFAAFVEKCKADDKETCFAEEGKLVAAASQAALGL</p>
110	HALB7	artificial		<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQCCPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLQHKDDNPNLPRLVREVDVMTAFHDNEETFLLKKYLYEIAARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAAACLLPKLDELDEGKASAKQRLKASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKYICENQDSISSKLECECEKPLLEKSHCIAEVENDEMPADLP SLAAD FVE SKDVCKNYAEAKDVFLGMFLYEVARRHPDYVSVLLLRLLAKTYETLEKCCAAAADPHECYAKVDFEFKPLVEEPQ</p>

111	HALB098	artificial	<p>NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTESLVNRRPCFSALEVDETYVPKEFNAGTTFHADICTLSEKERQIKKQTTALVELVKKHPKA TKEQLKAAMDFAAFVEKCKKADDDKTECFAEEGPKLVAASQAALGL</p> <p>DAHKSEVAHRFKDLGEEENFKALVLI AFAQYLQCCPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQE PERNECF LQHKDDNPNI PRLV RPEVDVMTAFHDNEETFLKKYLYEIARRHPYFYAPEL LFFAKRYKAAFTCCQAADKAAACLLPKLDEL RDEGKASSAKQRLK CASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKY ICENQDS ISSKKECCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVE SKDVCKNYAEAKDVFLGMFLYEYARRHPDYSVLLRLAKTYETLEKCCAAAADPHECYAKVDFEFKPLVEEPQ NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTESLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQTTALVELVKKHPKA TKEQLKAAMDFAAFVEKCKKADDDKTECFAEEGPKLVAASQAALGL</p>
112	HALB114	artificial	<p>DAHKSEVAHRFKDLGEEENFKALVLI AFAQYLQCCPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQE PERNECF LQHKDDNPNI PRLV RPEVDVMTAFHDNEETFLKKYLYEIARRHPYFYAPEL LFFAKRYKAAFTCCQAADKAAACLLPKLDEL RDEGKASSAKQRLK CASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKY ICENQDS ISSKKECCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVE SKDVCKNYAEAKDVFLGMFLYEYARRHPDYSVLLRLAKTYETLEKCCAAAADPHECYAKVDFEFKPLVEEPQ NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTESLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQTTALVELVKKHPKA TKEQLKAAMDFAAFVEKCKKADDDKTECFAEEGPKLVAASQAALGL</p>
113	HALB254	artificial	<p>DAHKSEVAHRFKDLGEEENFKALVLI AFAQYLQCCPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQE PERNECF LQHKDDNPNI PRLV RPEVDVMTAFHDNEETFLKKYLYEIARRHPYFYAPEL LFFAKRYKAAFTCCQAADKAAACLLPKLDEL RDEGKASSAKQRLK CASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKY ICENQDS ISSKKECCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVE SKDVCKNYAEAKDVFLGMFLYEYARRHPDYSVLLRLAKTYETLEKCCAAAADPHECYAKVDFEFKPLVEEPQ NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTESLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQTTALVELVKKHPKA TKEQLKAAMDFAAFVEKCKKADDDKTECFAEEGPKLVAASQAALGL</p>
114	HALB253	artificial	<p>DAHKSEVAHRFKDLGEEENFKALVLI AFAQYLQCCPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQE PERNECF LQHKDDNPNI PRLV RPEVDVMTAFHDNEETFLKKYLYEIARRHPYFYAPEL LFFAKRYKAAFTCCQAADKAAACLLPKLDEL RDEGKASSAKQRLK CASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKY ICENQDS ISSKKECCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVE SKDVCKNYAEAKDVFLGMFLYEYARRHPDYSVLLRLAKTYETLEKCCAAAADPHECYAKVDFEFKPLVEEPQ NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTESLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQTTALVELVKKHPKA TKEQLKAAMDFAAFVEKCKKADDDKTECFAEEGPKLVAASQAALGL</p>
115	HALB131	artificial	<p>DAHKSEVAHRFKDLGEEENFKALVLI AFAQYLQCCPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQE PERNECF LQHKDDNPNI PRLV RPEVDVMTAFHDNEETFLKKYLYEIARRHPYFYAPEL LFFAKRYKAAFTCCQAADKAAACLLPKLDEL RDEGKASSAKQRLK CASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKY ICENQDS ISSKKECCEKPLLEKSHCIAEVENDEMPADLPSLAAD</p>

116	HALB135	artificial	<p>FVE SKDVC KNYAE AKDVFLGMFLIYEYARRHPDY SVVLLIIRLAKTYETILEKCCAAADPHECYAKVDFEFKPLVEEPQ NLI KQNC ELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTE SLVNRPCFSALEVDVDETYVPKEFNAETTFHADICTLSEKERQIKKQTALVELVKKHKPKA TKEQLKAVMDDFAAFVEKCKKADDDKTECFAEEGPHLVAASQAALGL</p> <p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQCCPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRILVRPEVDVMTAFHDNEETFLKKYLYEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAACLIPKLDLDRDEGKASAKQRLKASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKYI CENQDSI SSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVE SKDVC KNYAE AKDVFLGMFLIYEYARRHPDY SVVLLIIRLAKTYETILEKCCAAADPHECYAKVDFEFKPLVEEPQ NLI KQNC ELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTE SLVNRPCFSALEVDVDETYVPKEFNAETTFHADICTLSEKERQIKKQTALVELVKKHKPKA TKEQLKAVMDDFAAFVEKCKKADDDKTECFAEEGPHLVAASQAALGL</p>
117	HALB133	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQCCPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRILVRPEVDVMTAFHDNEETFLKKYLYEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAACLIPKLDLDRDEGKASAKQRLKASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKYI CENQDSI SSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVE SKDVC KNYAE AKDVFLGMFLIYEYARRHPDY SVVLLIIRLAKTYETILEKCCAAADPHECYAKVDFEFKPLVEEPQ NLI KQNC ELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTE SLVNRPCFSALEVDVDETYVPKEFNAETTFHADICTLSEKERQIKKQTALVELVKKHKPKA TKEQLKAVMDDFAAFVEKCKKADDDKTECFAEEGPHLVAASQAALGL</p>
118	HALB234	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQCCPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRILVRPEVDVMTAFHDNEETFLKKYLYEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAACLIPKLDLDRDEGKASAKQRLKASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKYI CENQDSI SSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVE SKDVC KNYAE AKDVFLGMFLIYEYARRHPDY SVVLLIIRLAKTYETILEKCCAAADPHECYAKVDFEFKPLVEEPQ NLI KQNC ELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTE SLVNRPCFSALEVDVDETYVPKEFNAETTFHADICTLSEKERQIKKQTALVELVKKHKPKA TKEQLKAVMDDFAAFVEKCKKADDDKTECFAEEGPHLVAASQAALGL</p>
119	HALB C34S	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQCCPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRILVRPEVDVMTAFHDNEETFLKKYLYEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAACLIPKLDLDRDEGKASAKQRLKASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKYI CENQDSI SSKLKECCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVE SKDVC KNYAE AKDVFLGMFLIYEYARRHPDY SVVLLIIRLAKTYETILEKCCAAADPHECYAKVDFEFKPLVEEPQ NLI KQNC ELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTE SLVNRPCFSALEVDVDETYVPKEFNAETTFHADICTLSEKERQIKKQTALVELVKKHKPKA TKEQLKAVMDDFAAFVEKCKKADDDKTECFAEEGPHLVAASQAALGL</p>
120	HALB7 C34S	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQCCPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRILVRPEVDVMTAFHDNEETFLKKYLYEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAACLIPKLDLDRDEGKASAKQRLKASLQKFGERAFKAWAVARLSQRFPKAEFAEV</p>

					<p>SKLVTDLTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVESKDVCKNYAEAKDVFLGMFLYAYARRHPDYVWLLLRLLAKTYETLEKCCAAADPHCEYAKVDFEFKPLVEEPQ NLIKQNCLELFEQIGEYKFNQNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTESLVNRPCFSALEVDETYVPKEFNAGTTFHADICTLSEKERQIKKQTTALVELVKKHKPKA TKEQLKAVMDKFAAFVEKCKKADDDKTECFAEEGPKLVAAASQAALGL</p>			
121	HALB098 C34S	artificial			<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQOSPFEHDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRLVPEVDVMTAFHDNEETFLLKLYLIEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAAACLLPKLDELIRDEGKASSAKQRLKCASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVESKDVCKNYAEAKDVFLGMFLYAYARRHPDYVWLLLRLLAKTYETLEKCCAAADPHCEYAKVDFEFKPLVEEPQ NLIKQNCLELFEQIGEYKFNQNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTESLVNRPCFSALEVDETYVPKEFNAGTTFHADICTLSEKERQIKKQTTALVELVKKHKPKA TKEQLKAVMDKFAAFVEKCKKADDDKTECFAEEGPKLVAAASQAALGL</p>			
122	HALB114 C34S	artificial			<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQOSPFEHDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRLVPEVDVMTAFHDNEETFLLKLYLIEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAAACLLPKLDELIRDEGKASSAKQRLKCASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVESKDVCKNYAEAKDVFLGMFLYAYARRHPDYVWLLLRLLAKTYETLEKCCAAADPHCEYAKVDFEFKPLVEEPQ NLIKQNCLELFEQIGEYKFNQNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTESLVNRPCFSALEVDETYVPKEFNAGTTFHADICTLSEKERQIKKQTTALVELVKKHKPKA TKEQLKAVMDKFAAFVEKCKKADDDKTECFAEEGPKLVAAASQAALGL</p>			
123	HALB254 C34S	artificial			<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQOSPFEHDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRLVPEVDVMTAFHDNEETFLLKLYLIEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAAACLLPKLDELIRDEGKASSAKQRLKCASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVESKDVCKNYAEAKDVFLGMFLYAYARRHPDYVWLLLRLLAKTYETLEKCCAAADPHCEYAKVDFEFKPLVEEPQ NLIKQNCLELFEQIGEYKFNQNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTESLVNRPCFSALEVDETYVPKEFNAGTTFHADICTLSEKERQIKKQTTALVELVKKHKPKA TKEQLKAVMDKFAAFVEKCKKADDDKTECFAEEGPKLVAAASQAALGL</p>			
124	HALB253 C34S	artificial			<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQOSPFEHDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRLVPEVDVMTAFHDNEETFLLKLYLIEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAAACLLPKLDELIRDEGKASSAKQRLKCASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVESKDVCKNYAEAKDVFLGMFLYAYARRHPDYVWLLLRLLAKTYETLEKCCAAADPHCEYAKVDFEFKPLVEEPQ NLIKQNCLELFEQIGEYKFNQNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTESLVNRPCFSALEVDETYVPKEFNAGTTFHADICTLSEKERQIKKQTTALVELVKKHKPKA TKEQLKAVMDKFAAFVEKCKKADDDKTECFAEEGPKLVAAASQAALGL</p>			
125	HALB131 C34S	artificial			<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQOSPFEHDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNLPRLVPEVDVMTAFHDNEETFLLKLYLIEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAAACLLPKLDELIRDEGKASSAKQRLKCASLQKFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVESKDVCKNYAEAKDVFLGMFLYAYARRHPDYVWLLLRLLAKTYETLEKCCAAADPHCEYAKVDFEFKPLVEEPQ NLIKQNCLELFEQIGEYKFNQNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVVLNQLCV LHEKTPVSDRVTKCCTESLVNRPCFSALEVDETYVPKEFNAGTTFHADICTLSEKERQIKKQTTALVELVKKHKPKA TKEQLKAVMDKFAAFVEKCKKADDDKTECFAEEGPKLVAAASQAALGL</p>			

126	HALB135 C34S	artificial	<p>LFFAKRYKAAFTTECCQAADKAAACLLPKLDELIRDEGKASAKQRLKASLQKFFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADDDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVESKDVCKNYAEAKDVFLGMFLYERARRHPDYSVWLLLRRLAKTYETTLKCCAAAADPHECYAKVDFEFKPLVEEPQ NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVWLNQLCV LHEKTPVSDRVTKCCTESLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQTALVELLVKHKPKA TKEQLKAVMDKFAAFVEKCKKADDKETCFAAEEGPHLVAASQAALGL</p> <p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLQHKDDNPNIPLRVRPEVDVMTAFHDNEETFLKKLYLIEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAAACLLPKLDELIRDEGKASAKQRLKASLQKFFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADDDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVESKDVCKNYAEAKDVFLGMFLYERARRHPDYSVWLLLRRLAKTYETTLKCCAAAADPHECYAKVDFEFKPLVEEPQ NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVWLNQLCV LHEKTPVSDRVTKCCTESLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQTALVELLVKHKPKA TKEQLKAVMDKFAAFVEKCKKADDKETCFAAEEGPHLVAASQAALGL</p>
127	HALB133 C34S	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLQHKDDNPNIPLRVRPEVDVMTAFHDNEETFLKKLYLIEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAAACLLPKLDELIRDEGKASAKQRLKASLQKFFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADDDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVESKDVCKNYAEAKDVFLGMFLYERARRHPDYSVWLLLRRLAKTYETTLKCCAAAADPHECYAKVDFEFKPLVEEPQ NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVWLNQLCV LHEKTPVSDRVTKCCTESLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQTALVELLVKHKPKA TKEQLKAVMDKFAAFVEKCKKADDKETCFAAEEGPHLVAASQAALGL</p>
128	HALB234 C34S	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLQHKDDNPNIPLRVRPEVDVMTAFHDNEETFLKKLYLIEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAAACLLPKLDELIRDEGKASAKQRLKASLQKFFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADDDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVESKDVCKNYAEAKDVFLGMFLYERARRHPDYSVWLLLRRLAKTYETTLKCCAAAADPHECYAKVDFEFKPLVEEPQ NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVWLNQLCV LHEKTPVSDRVTKCCTESLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQTALVELLVKHKPKA TKEQLKAVMDKFAAFVEKCKKADDKETCFAAEEGPHLVAASQAALGL</p>
129	HALB C34A	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQQSPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLQHKDDNPNIPLRVRPEVDVMTAFHDNEETFLKKLYLIEIARRHPYFYAPEL LFFAKRYKAAFTTECCQAADKAAACLLPKLDELIRDEGKASAKQRLKASLQKFFGERAFKAWAVARLSQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADDDRADLAKYICENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLPSLAAD FVESKDVCKNYAEAKDVFLGMFLYERARRHPDYSVWLLLRRLAKTYETTLKCCAAAADPHECYAKVDFEFKPLVEEPQ NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVWLNQLCV LHEKTPVSDRVTKCCTESLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQTALVELLVKHKPKA TKEQLKAVMDKFAAFVEKCKKADDKETCFAAEEGPHLVAASQAALGL</p>

130	HALB7 C34A	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI.AFAQYLQOAPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNI.PRLVPEVDVMTAFHDNEETFLLKKLYE.IARRHPYFYAPEL LFFAKRYKAAAFTECCQAADKAAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFKAWAVARL.SQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKYI.CENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLP.SLAAD FVE.SKDVCKNYAEAKDVFLGMFLYEYARRHPDYSVVLLRLAKTYETLEKCCAAAADPHECYAKVDFEFKPLVEEPO NLI.KQNCELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPKAEKRMPCAEIDYLSVVLNQLCV LHEKTPVSDRVTKCCTE.SLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQ.TALVELLVKHKPKA TKEQLKAAVMDDDFAAFVEKCCADDKETE.CFAEEGPKLVAASQAALGL</p>
131	HALB098 C34A	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI.AFAQYLQOAPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNI.PRLVPEVDVMTAFHDNEETFLLKKLYE.IARRHPYFYAPEL LFFAKRYKAAAFTECCQAADKAAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFKAWAVARL.SQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKYI.CENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLP.SLAAD FVE.SKDVCKNYAEAKDVFLGMFLYEYARRHPDYSVVLLRLAKTYETLEKCCAAAADPHECYAKVDFEFKPLVEEPO NLI.KQNCELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPKAEKRMPCAEIDYLSVVLNQLCV LHEKTPVSDRVTKCCTE.SLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQ.TALVELLVKHKPKA TKEQLKAAVMDDDFAAFVEKCCADDKETE.CFAEEGPKLVAASQAALGL</p>
132	HALB114 C34A	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI.AFAQYLQOAPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNI.PRLVPEVDVMTAFHDNEETFLLKKLYE.IARRHPYFYAPEL LFFAKRYKAAAFTECCQAADKAAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFKAWAVARL.SQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKYI.CENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLP.SLAAD FVE.SKDVCKNYAEAKDVFLGMFLYEYARRHPDYSVVLLRLAKTYETLEKCCAAAADPHECYAKVDFEFKPLVEEPO NLI.KQNCELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPKAEKRMPCAEIDYLSVVLNQLCV LHEKTPVSDRVTKCCTE.SLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQ.TALVELLVKHKPKA TKEQLKAAVMDDDFAAFVEKCCADDKETE.CFAEEGPKLVAASQAALGL</p>
133	HALB254 C34A	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI.AFAQYLQOAPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNI.PRLVPEVDVMTAFHDNEETFLLKKLYE.IARRHPYFYAPEL LFFAKRYKAAAFTECCQAADKAAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFKAWAVARL.SQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKYI.CENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLP.SLAAD FVE.SKDVCKNYAEAKDVFLGMFLYEYARRHPDYSVVLLRLAKTYETLEKCCAAAADPHECYAKVDFEFKPLVEEPO NLI.KQNCELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPKAEKRMPCAEIDYLSVVLNQLCV LHEKTPVSDRVTKCCTE.SLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQ.TALVELLVKHKPKA TKEQLKAAVMDDDFAAFVEKCCADDKETE.CFAEEGPKLVAASQAALGL</p>
134	HALB253 C34A	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI.AFAQYLQOAPFEDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV ATLRETYGEMADCCAKQEPERNECFLOHKDDNPNI.PRLVPEVDVMTAFHDNEETFLLKKLYE.IARRHPYFYAPEL LFFAKRYKAAAFTECCQAADKAAACLLPKLDELDRDEGKASSAKQRLKCASLQKFGERAFKAWAVARL.SQRFPKAEFAEV SKLVTDLTKVHTECCHGDLLECADRADLAKYI.CENQDSISSKLEKCEKPLLEKSHCIAEVENDEMPADLP.SLAAD FVE.SKDVCKNYAEAKDVFLGMFLYEYARRHPDYSVVLLRLAKTYETLEKCCAAAADPHECYAKVDFEFKPLVEEPO NLI.KQNCELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPKAEKRMPCAEIDYLSVVLNQLCV LHEKTPVSDRVTKCCTE.SLVNRRPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQ.TALVELLVKHKPKA TKEQLKAAVMDDDFAAFVEKCCADDKETE.CFAEEGPKLVAASQAALGL</p>

135	HALB131 C34A	artificial	<p>TKEQLKAVMDKFAAFVEKCCKADDKETEFAEEGPKLVAAASQAALGL</p> <p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQQAPEFDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV</p> <p>ATLRETYGEMADCCAKQEPERNECFLQHKDDNPNI PRLVVRPEVDVMCTAFHDNEETFLLKKYLYEIAARRHPYFYAPEL</p> <p>LFFAKRYKAAAFTECCQAADKAAACLLPKLDELDRDEGKASSAKQRLKASLQKFGERAFKAWAVARLSQRFPKAEFAEV</p> <p>SKLVTDLTKVHTECCHGDLLECADRADLAKY ICENQDS ISSKKECCEKPLLEKSHCIAEVENDEMPADLPSLAAD</p> <p>FVESKDVCKNYAAEAKDVFLGMFLYEYARRHPDYSVWLLRLAKTYETLEKCCAAAADPHCEYAKVDFEFKPLVEEPPQ</p> <p>NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVWLNQLCV</p> <p>LHEKTPVSDRVTKCCTESLVNRPPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQ TALVELLVKHKPKA</p> <p>TKEQLKAVMDKFAAFVEKCCKADDKETEFAEEGPKLVAAASQAALGL</p>
136	HALB135 C34A	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQQAPEFDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV</p> <p>ATLRETYGEMADCCAKQEPERNECFLQHKDDNPNI PRLVVRPEVDVMCTAFHDNEETFLLKKYLYEIAARRHPYFYAPEL</p> <p>LFFAKRYKAAAFTECCQAADKAAACLLPKLDELDRDEGKASSAKQRLKASLQKFGERAFKAWAVARLSQRFPKAEFAEV</p> <p>SKLVTDLTKVHTECCHGDLLECADRADLAKY ICENQDS ISSKKECCEKPLLEKSHCIAEVENDEMPADLPSLAAD</p> <p>FVESKDVCKNYAAEAKDVFLGMFLYEYARRHPDYSVWLLRLAKTYETLEKCCAAAADPHCEYAKVDFEFKPLVEEPPQ</p> <p>NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVWLNQLCV</p> <p>LHEKTPVSDRVTKCCTESLVNRPPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQ TALVELLVKHKPKA</p> <p>TKEQLKAVMDKFAAFVEKCCKADDKETEFAEEGPKLVAAASQAALGL</p>
137	HALB133 C34A	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQQAPEFDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV</p> <p>ATLRETYGEMADCCAKQEPERNECFLQHKDDNPNI PRLVVRPEVDVMCTAFHDNEETFLLKKYLYEIAARRHPYFYAPEL</p> <p>LFFAKRYKAAAFTECCQAADKAAACLLPKLDELDRDEGKASSAKQRLKASLQKFGERAFKAWAVARLSQRFPKAEFAEV</p> <p>SKLVTDLTKVHTECCHGDLLECADRADLAKY ICENQDS ISSKKECCEKPLLEKSHCIAEVENDEMPADLPSLAAD</p> <p>FVESKDVCKNYAAEAKDVFLGMFLYEYARRHPDYSVWLLRLAKTYETLEKCCAAAADPHCEYAKVDFEFKPLVEEPPQ</p> <p>NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVWLNQLCV</p> <p>LHEKTPVSDRVTKCCTESLVNRPPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQ TALVELLVKHKPKA</p> <p>TKEQLKAVMDKFAAFVEKCCKADDKETEFAEEGPKLVAAASQAALGL</p>
138	HALB234 C34A	artificial	<p>DAHKSEVAHRFKDLGEENFKALVLI AFAQYLQQAPEFDHVKLVNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTV</p> <p>ATLRETYGEMADCCAKQEPERNECFLQHKDDNPNI PRLVVRPEVDVMCTAFHDNEETFLLKKYLYEIAARRHPYFYAPEL</p> <p>LFFAKRYKAAAFTECCQAADKAAACLLPKLDELDRDEGKASSAKQRLKASLQKFGERAFKAWAVARLSQRFPKAEFAEV</p> <p>SKLVTDLTKVHTECCHGDLLECADRADLAKY ICENQDS ISSKKECCEKPLLEKSHCIAEVENDEMPADLPSLAAD</p> <p>FVESKDVCKNYAAEAKDVFLGMFLYEYARRHPDYSVWLLRLAKTYETLEKCCAAAADPHCEYAKVDFEFKPLVEEPPQ</p> <p>NLIKQNCLELFEQLGEYKFNALLVRYTKKVPQVSTPTLVEVSRNLGKVGSKCKHPEAKRMPCAEDYLSVWLNQLCV</p> <p>LHEKTPVSDRVTKCCTESLVNRPPCFSALEVDETYVPKEFNAETTFHADICTLSEKERQIKKQ TALVELLVKHKPKA</p> <p>TKEQLKAVMDKFAAFVEKCCKADDKETEFAEEGPKLVAAASQAALGL</p>
139	Peptide linker	artificial	GGGG
140	Peptide linker	artificial	GGGGS
141	Peptide linker	artificial	GGGGQ

142	Peptide linker	artificial	PGGGS
143	Peptide linker	artificial	FGGDGS
144	Peptide linker	artificial	SGGGGS
145	Peptide linker	artificial	GGGSGGGGS
146	Peptide linker	artificial	GGGSGGGGS
147	Peptide linker	artificial	GGGSGGGSGGGGS
148	SEQ ID NO: 9 of WO-2008/119567	artificial	GSSTGAVTSGYYPN
149	SEQ ID NO: 10 of WO-2008/119567	artificial	GTKFLAP
150	SEQ ID NO: 11 of WO-2008/119567	artificial	ALWYSNRWV
151	SEQ ID NO: 12 of WO-2008/119567	artificial	IYAMN
152	SEQ ID NO: 13 of WO-2008/119567	artificial	RIRSKYNNYATYYADSVKS
153	SEQ ID NO: 14 of WO-2008/119567	artificial	HGNFGNSYVSFFAY
154	SEQ ID NO: 27 of WO-2008/119567	artificial	GSSTGAVTSGYYPN
155	SEQ ID NO: 28 of WO-2008/119567	artificial	GTKFLAP
156	SEQ ID NO: 29 of WO-2008/119567	artificial	ALWYSNRWV
157	SEQ ID NO: 30 of WO-2008/119567	artificial	KYAMN
158	SEQ ID NO: 31 of WO-2008/119567	artificial	RIRSKYNNYATYYADSVKD
159	SEQ ID NO: 32 of WO-2008/119567	artificial	HGNFGNSYISYWAY
160	SEQ ID NO: 45 of WO-2008/119567	artificial	GSSTGAVTSGYYPN

161	SEQ ID NO: 46 of WO-2008/119567	artificial	GTKFLAP
162	SEQ ID NO: 47 of WO-2008/119567	artificial	ALWYSNRWV
163	SEQ ID NO: 48 of WO-2008/119567	artificial	SYAMN
164	SEQ ID NO: 49 of WO-2008/119567	artificial	RIRSKYNNYATYYADSVKKG
165	SEQ ID NO: 50 of WO-2008/119567	artificial	HGNFGNSYLSFWAY
166	SEQ ID NO: 63 of WO-2008/119567	artificial	GSSTGAVTSGYYPN
167	SEQ ID NO: 64 of WO-2008/119567	artificial	GTKFLAP
168	SEQ ID NO: 65 of WO-2008/119567	artificial	ALWYSNRWV
169	SEQ ID NO: 66 of WO-2008/119567	artificial	RYAMN
170	SEQ ID NO: 67 of WO-2008/119567	artificial	RIRSKYNNYATYYADSVKKG
171	SEQ ID NO: 68 of WO-2008/119567	artificial	HGNFGNSYLSYFAY
172	SEQ ID NO: 81 of WO-2008/119567	artificial	RSSTGAVTSGYYPN
173	SEQ ID NO: 82 of WO-2008/119567	artificial	ATDMRPS
174	SEQ ID NO: 83 of WO-2008/119567	artificial	ALWYSNRWV
175	SEQ ID NO: 84 of WO-2008/119567	artificial	VYAMN
176	SEQ ID NO: 85 of WO-2008/119567	artificial	RIRSKYNNYATYYADSVKKG
177	SEQ ID NO: 86 of WO-2008/119567	artificial	HGNFGNSYLSFWAY
178	SEQ ID NO: 99 of WO-2008/119567	artificial	GSSTGAVTSGYYPN
179	SEQ ID NO: 100 of WO-2008/119567	artificial	GTKFLAP

180	SEQ ID NO: 101 of WO-2008/119567	artificial	ALWYSNRWV
181	SEQ ID NO: 102 of WO-2008/119567	artificial	KYAMN
182	SEQ ID NO: 103 of WO-2008/119567	artificial	RIRSKYNNYATYYADSVKS
183	SEQ ID NO: 104 of WO-2008/119567	artificial	HGNFGNSYTSYYAY
184	SEQ ID NO: 117 of WO-2008/119567	artificial	RSSTGAVTSGYYPN
185	SEQ ID NO: 118 of WO-2008/119567	artificial	ATDMRPS
186	SEQ ID NO: 119 of WO-2008/119567	artificial	ALWYSNRWV
187	SEQ ID NO: 120 of WO-2008/119567	artificial	GYAMN
188	SEQ ID NO: 121 of WO-2008/119567	artificial	RIRSKYNNYATYYADSVKE
189	SEQ ID NO: 122 of WO-2008/119567	artificial	HRNFGNSYLSWFAY
190	SEQ ID NO: 135 of WO-2008/119567	artificial	GSSTGAVTSGYYPN
191	SEQ ID NO: 136 of WO-2008/119567	artificial	GTKFLAP
192	SEQ ID NO: 137 of WO-2008/119567	artificial	ALWYSNRWV
193	SEQ ID NO: 138 of WO-2008/119567	artificial	VYAMN
194	SEQ ID NO: 139 of WO-2008/119567	artificial	RIRSKYNNYATYYADSVKK
195	SEQ ID NO: 140 of WO-2008/119567	artificial	HGNFGNSYISWWAY
196	SEQ ID NO: 153 of WO-2008/119567	artificial	GSSTGAVTSGNYYPN
197	SEQ ID NO: 154 of WO-2008/119567	artificial	GTKFLAP
198	SEQ ID NO: 155 of WO-2008/119567	artificial	VLIWYSNRWV

199	SEQ ID NO: 156 of WO-2008/119567	artificial	SYAMN
200	SEQ ID NO: 157 of WO-2008/119567	artificial	RIRSKYNNYATYYADSVKG
201	SEQ ID NO: 158 of WO-2008/119567	artificial	HGNFGNSYVSWWAY
202	SEQ ID NO: 171 of WO-2008/119567	artificial	GSSTGAVTSGNYPN
203	SEQ ID NO: 172 of WO-2008/119567	artificial	GTKFLAP
204	SEQ ID NO: 173 of WO-2008/119567	artificial	VLWYSNRWV
205	SEQ ID NO: 174 of WO-2008/119567	artificial	KYAMN
206	SEQ ID NO: 175 of WO-2008/119567	artificial	RIRSKYNNYATYYADSVKD
207	SEQ ID NO: 176 of WO-2008/119567	artificial	HGNFGNSYISYWAY

Claims

1. A bispecific construct comprising a first binding domain specifically binding to CD33 and a second binding domain specifically binding to CD3 for use in a method for the treatment of myeloid leukemia, wherein the construct is administered for a maximal period of 14 days followed by a period of at least 14 days without administration of the construct.
2. The bispecific construct for the use according to claim 1, wherein the construct is administered according to a schedule comprising the following steps:
 - (a) administration of a first dose of the bispecific construct, followed by
 - (b) administration of a second dose of the bispecific construct, wherein said second dose exceeds said first dose, optionally followed by
 - (a) administration of a third dose of the bispecific construct, wherein said optional third dose exceeds said second dose.
3. The bispecific construct for the use according to claim 2, wherein the period of administration of the first dose is up to seven days.
4. The bispecific construct for the use according to claim 2 or 3, wherein the complete period of administration of the dose steps subsequent to the first dose is in the range of eight to thirteen days.
5. The bispecific construct for the use according to any one of claims 2 to 4, wherein the period of administration of the second dose is four to seven days and the period of administration of the third dose is four to seven days.
6. The bispecific construct for the use according to any one of claims 2 to 5, wherein the treatment of the myeloid leukemia comprises two or more cycles of treatment which each comprises the maximal period of 14 days of construct administration followed by the period of at least 14 days without administration of the construct.
7. The bispecific construct for the use according to claim 6, wherein only the first cycle of the treatment comprises the administration according to step (a), whereas the following cycles start with the dose according to step (b).

8. The bispecific construct for the use according to any one of claim 1 to 7, wherein the first binding domain of the bispecific construct comprises groups of six CDRs selected from the group consisting of SEQ ID NOs: 10 to 12 and 14 to 16, 22 to 24 and 26 to 28, 34 to 36 and 38 to 40, 46 to 48 and 50 to 52, 58 to 60 and 62 to 64, 70 to 72 and 74 to 76, 82 to 84 and 86 to 88, 94 to 96 and 98 to 100.
9. The bispecific construct for the use according to any one of claim 1 to 8, wherein the second binding domain of the bispecific construct comprises groups of six CDRs selected from the group consisting of SEQ ID NOs: 148-153, 154-159, 160-165, 166-171, 172-177, 178-183, 184-189, 190-195, 196-201 and 202-207.
10. The bispecific construct for the use according to any one of claims 1 to 9, wherein the bispecific construct is a bispecific antibody construct.
11. The bispecific construct for the use according to claim 10, wherein the bispecific antibody construct is a single chain construct comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 18, 19, 20, 30, 31, 32, 42, 43, 44, 54, 55, 56, 66, 67, 68, 78, 79, 80, 90, 91, 92, 102, 103, 104, 105, 106, 107 and 108.
12. The bispecific construct for the use according to any one of the preceding claims, wherein the bispecific construct is administered in combination with one or more epigenetic factors selected from the group consisting of histone deacetylase (HDAC) inhibitors, DNA methyltransferase (DNMT) I inhibitors, hydroxyurea, Granulocyte-Colony Stimulating Factor (G-CSF), histone demethylase inhibitors and ATRA (All Trans-retinoic acid) and wherein:
 - (a) the one or more epigenetic factors are administered prior to the administration of the bispecific construct;
 - (b) the one or more epigenetic factors are administered subsequent to the administration of the bispecific construct; or
 - (c) the one or more epigenetic factors and the bispecific construct are administered simultaneously.
13. The bispecific construct for the use according to claim 12, wherein the one or more epigenetic factors are administered up to seven days prior to the administration of the bispecific construct.

14. The bispecific construct for the use according to claim 13, wherein the epigenetic factor is hydroxyurea.
15. The bispecific construct for the use according to any one of the preceding claims, wherein the myeloid leukemia is selected from the group consisting of acute myeloblastic leukemia, chronic neutrophilic leukemia, myeloid dendritic cell leukemia, accelerated phase chronic myelogenous leukemia, acute myelomonocytic leukemia, juvenile myelomonocytic leukemia, chronic myelomonocytic leukemia, acute basophilic leukemia, acute eosinophilic leukemia, chronic eosinophilic leukemia, acute megakaryoblastic leukemia, essential thrombocytosis, acute erythroid leukemia, polycythemia vera, myelodysplastic syndrome, acute panmyeloic leukemia, myeloid sarcoma, and acute biphenotypic leukaemia
16. A method for the treatment of myeloid leukemia comprising the administration of a therapeutically efficient amount of a bispecific construct comprising a first binding domain specifically binding to CD33 and a second binding domain specifically binding to CD3 to a subject in need thereof, the method comprising the step of administering to said subject the construct for a maximal period of 14 days followed by a step without administration of the construct for a period of at least 14 days.
17. The method according to claim 16, wherein the bispecific construct is administered according to a schedule comprising the following steps:
 - (a) administration of a first dose of the bispecific construct, followed by
 - (b) administration of a second dose of the bispecific construct, wherein said second dose exceeds said first dose, optionally followed by
 - (c) administration of a third dose of the bispecific construct, wherein said optional third dose exceeds said second dose.
18. The method according to claim 17, wherein the period of administration of the first dose is up to seven days.
19. The method according claim 17 or 18, wherein the complete period of administration of the dose steps subsequent to the first dose is in the range of eight to thirteen days.
20. The method according to any one of claims 17 to 19, wherein the period of administration of the second dose is four to seven days and the period of administration of the third dose is four to seven days.

21. The method according to any one of claims 17 to 20, comprising two or more cycles of treatment which each comprises the maximal period of 14 days of bispecific construct administration followed by the period of at least 14 days without administration of the bispecific construct.
22. The method according to claim 21, wherein only the first cycle of the treatment comprises the administration according to step (a), whereas the following cycles start with the dose according to step (b).
23. The method according to any one of claims 16 to 22, wherein the first binding domain of the bispecific construct comprises groups of six CDRs selected from the group consisting of SEQ ID NOs: 10 to 12 and 14 to 16, 22 to 24 and 26 to 28, 34 to 36 and 38 to 40, 46 to 48 and 50 to 52, 58 to 60 and 62 to 64, 70 to 72 and 74 to 76, 82 to 84 and 86 to 88, 94 to 96 and 98 to 100.
24. The method according to any one of claim 16 to 23, wherein the second binding domain of the bispecific construct comprises groups of six CDRs selected from the group consisting of SEQ ID NOs: 148-153, 154-159, 160-165, 166-171, 172-177, 178-183, 184- 189, 190-195, 196-201 and 202-207.
25. The method according to any one of claims 16 to 24, wherein the bispecific construct is a bispecific antibody construct.
26. The method according to claim 25, wherein the bispecific antibody construct is a single chain construct comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 18, 19, 20, 30, 31, 32, 42, 43, 44, 54, 55, 56, 66, 67, 68, 78, 79, 80, 90, 91, 92, 102, 103, 104, 105, 106, 107 and 108.
27. The method according to any one of claims 16 to 26, wherein the bispecific construct is administered in combination with one or more epigenetic factors selected from the group consisting of histone deacetylase (HDAC) inhibitors, DNA methyltransferase (DNMT) I inhibitors, hydroxyurea, Granulocyte-Colony Stimulating Factor (G-CSF), histone demethylase inhibitors and ATRA (All Trans-retinoic acid) and wherein:
 - (a) the one or more epigenetic factors are administered prior to the administration of the bispecific construct;

- (b) the one or more epigenetic factors are administered subsequent to the administration of the bispecific construct; or
 - (c) the one or more epigenetic factors and the bispecific construct are administered simultaneously.
28. The method according to claim 27, wherein the one or more epigenetic factors are administered up to seven days prior to the administration of the bispecific construct.
29. The method according to claim 28, wherein the epigenetic factor is hydroxyurea.
30. The method according to any one of claims 16 to 29, wherein the myeloid leukemia is selected from the group consisting of acute myeloblastic leukemia, chronic neutrophilic leukemia, myeloid dendritic cell leukemia, accelerated phase chronic myelogenous leukemia, acute myelomonocytic leukemia, juvenile myelomonocytic leukemia, chronic myelomonocytic leukemia, acute basophilic leukemia, acute eosinophilic leukemia, chronic eosinophilic leukemia, acute megakaryoblastic leukemia, essential thrombocytosis, acute erythroid leukemia, polycythemia vera, myelodysplastic syndrome, acute panmyeloic leukemia, myeloid sarcoma, and acute biphenotypic leukaemia
31. Use of a bispecific construct comprising a first binding domain specifically binding to CD33 and a second binding domain specifically binding to CD3 for the preparation of a pharmaceutical composition for the treatment of myeloid leukemia, wherein the construct is to be administered for a maximal period of 14 days followed by a period of at least 14 days without administration of the construct.
32. The use according to claim 31, wherein the bispecific construct is to be administered according to a schedule comprising the following steps:
- (a) administration of a first dose of the bispecific construct, followed by
 - (b) administration of a second dose of the bispecific construct, wherein said second dose exceeds said first dose, optionally followed by
 - (a) administration of a third dose of the bispecific construct, wherein said optional third dose exceeds said second dose.
33. The use according to claim 32, wherein the period of administration of the first dose is up to seven days.

34. The use according to claim 32 or 33, wherein the complete period of administration of the dose steps subsequent to the first dose is in the range of eight to thirteen days.
35. The use according to any one of claims 32 to 34, wherein the period of administration of the second dose is four to seven days and the period of administration of the third dose is four to seven days.
36. The use according to any one of claims 32 to 35, wherein the treatment of the myeloid leukemia comprises two or more cycles of treatment which each comprises the maximal period of 14 days of bispecific construct administration followed by the period of at least 14 days without administration of the bispecific construct.
37. The use according to claim 36, wherein only the first cycle of the treatment comprises the administration according to step (a), whereas the following cycles start with the dose according to step (b).
38. The use according to any one of claim 31 to 37, wherein the first binding domain of the bispecific construct comprises groups of six CDRs selected from the group consisting of SEQ ID NOs: 10 to 12 and 14 to 16, 22 to 24 and 26 to 28, 34 to 36 and 38 to 40, 46 to 48 and 50 to 52, 58 to 60 and 62 to 64, 70 to 72 and 74 to 76, 82 to 84 and 86 to 88, 94 to 96 and 98 to 100.
39. The use according to any one of claim 31 to 38, wherein the second binding domain of the bispecific construct comprises groups of six CDRs selected from the group consisting of SEQ ID NOs: 148-153, 154-159, 160-165, 166-171, 172-177, 178-183, 184- 189, 190-195, 196-201 and 202-207.
40. The use according to any one of claims 31 to 39, wherein the bispecific construct is a bispecific antibody construct.
41. The use according to claim 40, wherein the bispecific antibody construct is a single chain construct comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 18, 19, 20, 30, 31, 32, 42, 43, 44, 54, 55, 56, 66, 67, 68, 78, 79, 80, 90, 91, 92, 102, 103, 104, 105, 106, 107 and 108.
42. The use according to any one of claims 31 to 41, wherein the bispecific construct is administered in combination with one or more epigenetic factors selected from the

group consisting of histone deacetylase (HDAC) inhibitors, DNA methyltransferase (DNMT) I inhibitors, hydroxyurea, Granulocyte-Colony Stimulating Factor (G-CSF), histone demethylase inhibitors and ATRA (All Trans-retinoic acid) and wherein:

- (a) the one or more epigenetic factors are administered prior to the administration of the bispecific construct;
 - (b) the one or more epigenetic factors are administered subsequent to the administration of the bispecific construct; or
 - (c) the one or more epigenetic factors and the bispecific construct are administered simultaneously.
43. The use according to claim 42, wherein the one or more epigenetic factors are administered up to seven days prior to the administration of the bispecific construct.
44. The use according to claim 43, wherein the epigenetic factor is hydroxyurea.
45. The use according to any one of claims 31 to 44, wherein the myeloid leukemia is selected from the group consisting of acute myeloblastic leukemia, chronic neutrophilic leukemia, myeloid dendritic cell leukemia, accelerated phase chronic myelogenous leukemia, acute myelomonocytic leukemia, juvenile myelomonocytic leukemia, chronic myelomonocytic leukemia, acute basophilic leukemia, acute eosinophilic leukemia, chronic eosinophilic leukemia, acute megakaryoblastic leukemia, essential thrombocytosis, acute erythroid leukemia, polycythemia vera, myelodysplastic syndrome, acute panmyeloic leukemia, myeloid sarcoma, and acute biphenotypic leukaemia




Figure 1

Figure 2





Figure 3



INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2017/059108

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07K16/28 A61P35/00
ADD.
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
C07K
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2016/004108 A2 (AMPHIVENA THERAPEUTICS INC [US]) 7 January 2016 (2016-01-07) examples	1-45
X	Amgen: "A Phase 1 Study of AMG 330 in Subjects With Relapsed/Refractory Acute Myeloid Leukemia", ClinicalTrials.gov, 17 June 2015 (2015-06-17), XP55382070, Retrieved from the Internet: URL:https://clinicaltrials.gov/ct2/show/NC T02520427?term=amg+330&rank=1#wrapper [retrieved on 2017-06-15] the whole document	1-45
	----- -/--	

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>
---	---

Date of the actual completion of the international search 19 June 2017	Date of mailing of the international search report 26/06/2017
---	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Covone-van Hees, M
--	--

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2017/059108

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2015/036583 A2 (AMGEN INC [US]; AMGEN RES MUNICH GMBH [DE]) 19 March 2015 (2015-03-19) examples	1-45
X,P	----- WO 2016/196230 A1 (AMPHIVENA THERAPEUTICS INC [US]) 8 December 2016 (2016-12-08) examples	1-45
T	----- KIMBERLY H. HARRINGTON ET AL: "The Broad Anti-AML Activity of the CD33/CD3 BiTE Antibody Construct, AMG 330, Is Impacted by Disease Stage and Risk", PLOS ONE, vol. 10, no. 8, 25 August 2015 (2015-08-25), page e0135945, XP55370197, DOI: 10.1371/journal.pone.0135945 the whole document -----	1-45

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/EP2017/059108

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
WO 2016004108	A2	07-01-2016	AU 2015284200 A1	02-02-2017
			CA 2953992 A1	07-01-2016
			CN 106794266 A	31-05-2017
			EP 3164159 A2	10-05-2017
			KR 20170041697 A	17-04-2017
			SG 11201610973Y A	27-01-2017
			TW 201613973 A	16-04-2016
			US 9212225 B1	15-12-2015
			US 2016194409 A1	07-07-2016
			WO 2016004108 A2	07-01-2016
WO 2015036583	A2	19-03-2015	AR 097648 A1	06-04-2016
			AU 2014320246 A1	18-02-2016
			CA 2923354 A1	19-03-2015
			CN 105764505 A	13-07-2016
			EA 201690485 A1	31-08-2016
			EP 3043794 A2	20-07-2016
			JP 2016536341 A	24-11-2016
			KR 20160058120 A	24-05-2016
			PH 12016500434 A1	16-05-2016
			SG 11201600671Q A	26-02-2016
			TW 201605473 A	16-02-2016
			US 2016317657 A1	03-11-2016
			WO 2015036583 A2	19-03-2015
WO 2016196230	A1	08-12-2016	TW 201706301 A	16-02-2017
			WO 2016196230 A1	08-12-2016