



US 20150238545A1

(19) **United States**(12) **Patent Application Publication**
BORODY(10) **Pub. No.: US 2015/0238545 A1**(43) **Pub. Date: Aug. 27, 2015**(54) **PROBIOTIC RECOLONISATION THERAPY***A23L 1/30* (2006.01)*A61K 35/747* (2006.01)(71) Applicant: **Thomas Julius BORODY**, Five Dock
(AU)(52) **U.S. Cl.**CPC *A61K 35/741* (2013.01); *A61K 35/747*
(2013.01); *A61K 35/38* (2013.01); *A23L 1/3014*
(2013.01); *A23V 2002/00* (2013.01)(72) Inventor: **Thomas Julius BORODY**, Five Dock
(AU)(21) Appl. No.: **14/710,481**(22) Filed: **May 12, 2015**(57) **ABSTRACT****Related U.S. Application Data**(60) Continuation of application No. 14/270,034, filed on
May 5, 2014, now Pat. No. 9,050,358, Continuation of
application No. 13/910,579, filed on Jun. 5, 2013, now
Pat. No. 9,040,036, which is a division of application
No. 10/332,986, filed on Aug. 4, 2003, now Pat. No.
8,460,648, filed as application No. PCT/AU01/00907
on Jul. 25, 2001.(30) **Foreign Application Priority Data**

Jul. 25, 2000 (AU) PQ8997

Publication Classification(51) **Int. Cl.***A61K 35/741* (2006.01)*A61K 35/38* (2006.01)

The present invention relates to pharmaceutical compositions suitable for the treatment of chronic diseases associated with the presence of abnormal or an abnormal distribution of microflora in the gastrointestinal tract of a mammalian host, which compositions comprise viable non-pathogenic or attenuated pathogenic *Clostridia*. The compositions further comprise one or more additional viable non-pathogenic or attenuated pathogenic microorganisms selected from the group consisting of *Bacteroides*, Eubacteria, Fusobacteria, Propionibacteria, Lactobacilli, anaerobic cocci, *Ruminococcus*, *E. coli*, *Gemmiger*, *Desulfomonas*, *Peptostreptococcus*, and fungi. The present invention also provides pharmaceutical compositions suitable for the treatment of the same chronic diseases comprising viable non-pathogenic or attenuated pathogenic *Escherichia coli*, at least one strain of viable non-pathogenic or attenuated pathogenic *Bacteroides* and at least one strain of viable non-pathogenic or attenuated pathogenic microorganism.

PROBIOTIC RECOLONISATION THERAPY

TECHNICAL FIELD

[0001] The present invention relates to pharmaceutical compositions suitable for the treatment of diseases in mammals, in particular to the treatment of chronic disorders associated with the presence of abnormal or an abnormal distribution of microflora in the gastrointestinal tract. The invention also relates to methods of treating such diseases.

BACKGROUND ART

[0002] There are large numbers of patients suffering from gastro-intestinal symptoms referable to the lower small bowel and large bowel which to date have eluded explanation. These disorders include irritable bowel syndrome (IBS) or spastic colon, idiopathic ulcerative colitis, mucous colitis, collagenous colitis, Crohn's disease, inflammatory bowel disease in general, microscopic colitis, antibiotic-associated colitis, idiopathic or simple constipation, diverticular disease, and AIDS enteropathy. Pathophysiology of these disorders eludes logical explanation in spite of decades of research and millions of dollars of research funds. A common underlying factor shared by all these disorders observed by the present inventor is their onset or aggravation following some extraneous invading infection eg travelers diarrhoea. In all the disorders, a specific causal infection generally cannot be demonstrated due to our inability to detect infecting agents whose cultural characteristics are unknown to medical science.

[0003] Circumstantial evidence which suggests that these disorders are "infection-related" includes:

[0004] (a) onset following a gastro-intestinal infection which failed to completely resolve;

[0005] (b) transient improvement with use of certain antibiotics, but recurrence upon cessation of antibiotics;

[0006] (c) transient improvement following orthostatic lavage prior to colonoscopy and;

[0007] (d) transient symptom improvement with use of "colonic" irrigation.

[0008] It is impractical to use long-term antibiotic therapy (with its associated complications) in such patients since cure is not obtained with its use. Furthermore, chronic gut infections with recognised, specific pathogens such as *Clostridium difficile*, *Yersinia enterocolitica* or *Campylobacter jejuni/coli* are generally not eradicated with antibiotics. Some previous attempts have been made to alter the enteric microflora in order to eradicate such chronic infections. These measures nevertheless indicate that alteration of bacterial flora may effect dramatic clinical improvement in conditions characterised by chronic, resistant enterocolitic infection. However there remain many chronic disorders of uncertain aetiology or causation, which are resistant to cure by current therapeutic techniques.

[0009] The use of probiotics in the human population has been largely confined to the inclusion in various foods of live organism of Lactobacilli and Bifidobacteria and less frequently *Streptococcus faecalis* or several strains of *Escherichia coli*. These organisms are thought to promote health via immune stimulation and reconstitution of what is presumed to be normal flora. Such usage stems back to the beliefs generated by Mechnikov in the early 1900s. The use of probiotics to treat established infections in the gastrointestinal tract has been lesser but a growing part of the use of probiot-

ics. Fungal agents such as *Saccharomyces boulardii* have been used to treat, albeit inefficiently, *Clostridium difficile* infection and *Lactobacillus* GG has also been used for this purpose (Floch M. Probiotics and Dietary Fibre. *J Clin Gastroenterol* 1998; 27(2):99-100). Various patents have claimed the use of probiotics for narrow disease conditions including treatment of *Clostridium difficile* with a combination of Vancomycin and butyric acid bacteria (U.S. Pat. No. 5,266,315), diarrhoea prevention using *Lactobacillus* (U.S. Pat. No. 5,837,238) or *Bifidobacterium* (U.S. Pat. No. 5,902,743), *Lactobacillus acidophilus* to inhibit *cryptosporidium* (U.S. Pat. No. 5,858,356) and mixtures of Lactobacilli and Bifidobacteria in infants to prevent diarrhoea. *Enterococcus faecium* has been claimed to be useful in alleviating symptoms of Irritable Bowel Syndrome in humans (U.S. Pat. No. 5,902,578) (U.S. Pat. No. 5,728,380) but this has not recognised *Clostridium* as the underlying agent in this condition. *Clostridium butyricum* as a single agent has been claimed to be a biological intestinal antiseptic for treatment of bacterial food poisonings (U.S. Pat. No. 4,892,731), but its use in chronic disease treatment was not contemplated.

[0010] Previous attempts to alter the enteric microflora of a patient have prescribed the removal of at least a part of the host's existing enteric microflora, for instance by lavage, prior to substitution with predetermined desired microflora. This procedure, which was the preferred embodiment of WO90/01335 has the distinct disadvantages of complicating the treatment and of causing further discomfort to the patient. This patent also advocated the use of dried, reconstituted faeces or a synthetic mixture comprising *Bacteroides* sp. and *Escherichia coli*. It has now been surprisingly found that lavage or other methods of removal of at least a part of the host's existing enteric microflora can be omitted provided a non-pathogenic *Clostridium* sp. is included within the probiotic replacement mixture. Such a replacement mixture has the dual ability of displacing pathogenic bacteria, frequently Clostridial in nature and also establishing a normal environment in which commensal bacteria can establish. Such a treatment permits long-term recovery both from gastrointestinal disorders and from systemic afflictions not hitherto considered to be caused by harmful enteric flora. These are also called 'para-infective' phenomena and can include rheumatological, neurological, regressive, hepatic, and dermatological conditions among others.

[0011] Autism is a regressive disorder of childhood, affecting boys four times more often than girls. It has been observed that the onset of autism is often preceded by broad spectrum antibiotic use eg for recurrent ear infections. Antibiotic therapy is non-discriminatory in its action and apart from treating the ear infection the microflora of the healthy gastrointestinal tract can be severely disrupted by such treatment. This creates an environment where vulnerability to opportunistic microorganism colonisation is heightened.

[0012] *Clostridium tetani* is a widely distributed, spore forming anaerobe. Toxigenic strains of *Clostridium tetani* produce the extremely potent tetanus neurotoxin which is known to enter the central nervous system from the intestinal tract via the vagus nerve (Hensel B et al. Naunyn-Schmiedeberg's Arch Pharmacol 1973; 276:395). Bolte (Med Hypotheses 1998; 51:133) has hypothesised that opportunistic infection by *Clostridium tetani* may be responsible for the behavioural and medical symptoms present in a sub-group of individuals diagnosed with autism. Others have also raised

the possibility of clostridia in general as a cause of disease (Borriello S P. Clin Infect Dis 1995; Suppl 2:5242).

[0013] Sandler et al. (Fourth Int. Symp. Brain-Gut Interactions. 1998; 10: 363) report a trial in which children with delayed onset autism were treated with vancomycin over an 8 week period. All children in the trial had had antecedent broad-spectrum antibiotic exposure, followed by chronic persistent diarrhoea and then onset of autistic features. Although significant post-treatment improvement was noted, all children eventually regressed towards baseline.

[0014] It is on the background of these known facts and later the results of trials of treatment, that the present invention was formulated. In brief, it was noted that autistic children (as well as related syndromes) who were referred for treatment of refractory ‘irritable bowel syndrome’ (IBS) viz diarrhoea, flatulence, constipation, distension, abdominal pains etc—responded to treatment of their IBS when treated with a novel mix of probiotics. However, not only did their IBS improve dramatically but also their autistic features progressively regressed. Even after the initial 2-6 weeks of treatment eye contact was re-established, repetitive movements were much reduced, and word power (observed vocabulary) expanded—initially 20 words and ultimately >600 words at 12 months (estimated), creating ability of the autistic children to form long sentences. Continuing improvement was observed to occur over 12 months of treatment. These observations (to a lesser but definite degree at this stage of observations) also applied to those with Rett syndrome and children with Attention Deficit/Hyperactivity Disorder (ADHD), Attention Deficit Disorder (ADD), and autism variant Asbergers syndrome. The observations strongly suggest that the treatment of presumed enteric infection/s (eg Clostridial) in these conditions not only improves the IBS present but also the attendant neurological ‘para-infective’ phenomena called collectively autism, Asbergers, Rett syndrome, ADD or ADHD.

[0015] The inclusion within this specification of reference to published documents is not to be taken to be an admission that any one or more of those documents, nor the disclosure of any one or more of those documents, is part of the common general knowledge.

Objects of the Invention

[0016] It is thus an object of the present invention to provide novel pharmaceutical compositions suitable for the treatment of various disease states related to the presence of ‘abnormal’ microflora in the gastrointestinal tract. It is a further object of the invention to propose the use of these pharmaceutical compositions in various disease states which have not previously been considered to owe their causation to the presence of abnormal flora in the gastrointestinal tract.

Disclosure of the Invention

[0017] The present invention recognises chronic infection/infestation as the underlying pathological process in a wide range of chronic disorders such as irritable bowel syndrome, particularly when characterised by chronic abdominal pain, bloating, or excessive flatulence, together with chronic diarrhoea or alternating constipation/diarrhoea, and also in spastic colon, mucous colitis, collagenous colitis, ulcerative colitis, Crohn’s colitis, microscopic colitis, idiopathic

inflammatory bowel disease, antibiotic-associated colitis, idiopathic or simple constipation, diverticular disease and AIDS enteropathy.

[0018] The invention has also been found to relate to other gastrointestinal disorders of unexplained aetiology such as polyposis coli and colonic polyps, which may well be influenced by the local bowel microflora.

[0019] In addition the present invention also provides a method of treatment of chronic gastrointestinal infections with specific microorganisms such as *Clostridium difficile*, *Yersinia* spp, *Campylobacter* spp, *Aeromonas* spp, *Escherichia coli*, *Cryptosporidium* spp, *Amoebae*, *Blastocystis hominis*, *Giardia* and even chronic viral infections, and of small bowel bacterial overgrowth.

[0020] The present invention furthermore, recognises the close association between the intestine and liver disease, and the intestine and migraines and chronic fatigue syndrome, and possibly other neurological syndromes such as, multiple sclerosis, amyotrophic lateral sclerosis, myasthenia gravis, Parkinson’s disease, Alzheimer’s disease, Chronic Inflammatory Demyelinating Polyneuropathy (CIDP), Guillain Barre Syndrome, and other degenerative disorders. Hence, it is proposed that a considerable proportion of currently unexplained diseases of the liver and nervous system of unknown aetiology may be explicable by the chronic growth of pathogens within the small/large intestine and the subsequent passage of antigenic material, pathogenic toxins or biological response modifiers (BRMs) into the portal system (liver damage) or systemic circulation with antibody formation (neurological conditions). Specifically, such hepato/biliary system disorders as primary biliary cirrhosis, primary sclerosing cholangitis, fatty liver of unknown aetiology, or cryptogenic cirrhosis, may be secondary to chronic pathogen carrier state in the intestine.

[0021] The links between the intestine and joint disease are also recognised. Joint diseases such as rheumatoid arthritis, the non-rheumatoid arthritides including, ankylosing spondylitis, and Reiter’s syndrome, may also be causally related to a chronic intestinal carrier state, as may other syndromes with an immune mediated component such as glomerulonephritis, haemolytic uraemic syndrome, juvenile diabetes mellitus, Behcet’s syndrome, coeliac disease and dermatitis herpetiformis. Similarly, syndromes with an immune complex mediated component, such as scleroderma, systemic lupus erythematosus, mixed cryoglobulinaemia, polyarteritis, familial Mediterranean fever, amyloidosis, and the various presentations of such syndromes, together with such “idiopathic” states as chronic urticaria, may be manifestations of variations of immune regulated responses to related bowel-origin pathogens chronically shedding their antigen (s), toxins or biological response modifiers into the circulation. Other chronic conditions such as acne, and chronic idiopathic pseudo-obstructive syndrome, may well be influenced by similar mechanisms.

[0022] For many of these syndromes present therapy offers only palliation of symptoms and/or the induction of remission of the disease process but not cure. The present inventor therefore recognised the need to find a curative therapy for these wide ranging disease processes associated with considerable morbidity.

[0023] By judicious selection of the microorganisms of the invention it has been surprisingly found by the present inventor that lasting recolonisation of the gut microflora does not require pretreatment to remove a portion of the host’s existing

enteric microflora. Thus, by incorporation of *Clostridia* spp. in the therapy, it has been surprisingly found that the prior art requirement for removal of at least a portion of the existing enteric microflora before administration of the substitute microflora is rendered unnecessary. Without the addition specifically of *Clostridia* species, the use of probiotic mixtures, eg such as those of *bacteroides* and *Escherichia coli* failed to have the necessary impact on the above-mentioned clinical disorders for the treatment to be clinically useful.

[0024] It required a prior purging of the gut of its presumably infected and abnormal bowel flora, re colonisation with *bacteroides* and *Escherichia coli*—the main components of lower intestinal tract, and ongoing feeding of patients with such bacteria until colonisation was established. The use of *Clostridia* appears to be the mainstay of this new therapy and the *Clostridia* appear to have power of themselves to remove offending bacterial species which may be responsible for the underlying condition (presumably pathogenic clostridia—yet to be identified scientifically). Hence, the combination of non-pathogenic clostridia together with the crucial major colonic bacterial components of *bacteroides* and *Escherichia coli* can now be used as oral therapy to crowd out/destroy/replace and recolonise the dysbiotic flora of patients with various gastrointestinal conditions which are caused by abnormal bowel flora. In fact, such a therapy becoming available has permitted or allowed greater understanding of the pathogenesis of many other conditions which hitherto were thought to be caused by degenerative, inflammatory, or auto immune mechanisms.

[0025] Thus according to a first embodiment of the invention there is provided a pharmaceutical composition useful for the treatment and/or prophylaxis of chronic disorders associated with the presence in the gastrointestinal tract of a mammalian host of abnormal or an abnormal distribution of microflora, which composition comprises viable non-pathogenic or attenuated pathogenic *Clostridia*.

[0026] Typically the composition includes *Clostridia* selected from the group consisting of *Clostridium absonum*, *Clostridium argentinense*, *Clostridium baratii*, *Clostridium bifermentans*, *Clostridium botulinum*, *Clostridium butyricum*, *Clostridium cadaveris*, *Clostridium camis*, *Clostridium celatum*, *Clostridium chauvoei*, *Clostridium clostridioforme*, *Clostridium cochlearium*, *Clostridium difficile*, *Clostridium fallax*, *Clostridium felsineum*, *Clostridium ghonii*, *Clostridium glycolicum*, *Clostridium haemolyticum*, *Clostridium hastiforme*, *Clostridium histolyticum*, *Clostridium indolis*, *Clostridium innocuum*, *Clostridium irregulare*, *Clostridium limosum*, *Clostridium malenominatum*, *Clostridium novyi*, *Clostridium oroticum*, *Clostridium paraputrificum*, *Clostridium perfringens*, *Clostridium piliforme*, *Clostridium putrefaciens*, *Clostridium putrificum*, *Clostridium ramosum*, *Clostridium sardiniense*, *Clostridium sartagoforme*, *Clostridium scindens*, *Clostridium septicum*, *Clostridium sordellii*, *Clostridium sphenoides*, *Clostridium spiroforme*, *Clostridium sporogenes*, *Clostridium subterminale*, *Clostridium symbiosum*, *Clostridium tedium*, *Clostridium tetani*, *Clostridium welchii*, *Clostridium villosum*.

[0027] In a preferred form the composition further comprises one or more additional viable non-pathogenic or attenuated pathogenic microorganisms selected from the group consisting of *Bacteroides*, *Eubacteria*, *Fusobacteria*, *Propionibacteria*, *Lactobacilli*, *anaerobic cocci*, *Ruminococcus*, *Escherichia coli*, *Gemmiger*, *Desulfomonas*, *Peptostrep-*

tococcus, species and, more specifically, bacteria selected from Table 1. Preferably fungi are also present such as *Monilia*.

[0028] In a preferred form the composition comprises *Clostridia*, *Bacteroides*, *Peptostreptococcus*, *Escherichia coli*, *Bifidobacterium*, and *Lactobacillus*.

[0029] In a more preferred form the composition comprises *Clostridium innocuum*, *Clostridium bifermentans*, *Clostridium butyricum*, *Bacteroides fragilis*, *Bacteroides thetaiotaomicron*, *Bacteroides uniformis*, one or more strains of *Escherichia coli*, and one or more strains of *Lactobacillus*.

[0030] Alternatively, in a preferred form the composition comprises *Clostridium bifermentans*, *Clostridium innocuum*, and *Clostridium butyricum* in combination one or more strains of *Escherichia coli*, one or more strains of *bacteroides* and *Peptostreptococcus productus*.

[0031] According to a second embodiment of the invention there is provided a pharmaceutical composition useful for the treatment and/or prophylaxis of chronic disorders associated with the presence in the gastrointestinal tract of a mammalian host of abnormal or an abnormal distribution of microflora, which composition comprises viable non-pathogenic or attenuated pathogenic *Escherichia coli*, at least one strain of viable non-pathogenic or attenuated pathogenic *Bacteroides*, and at least one other viable non-pathogenic or attenuated pathogenic microorganism.

[0032] In a preferred form the other viable non-pathogenic or attenuated pathogenic microorganism is selected from the group consisting of *Clostridia*, *Peptostreptococcus*, *Bifidobacterium*, and *Lactobacillus*.

[0033] Typically the composition of the first or second embodiments of the invention is derived from disease screened fresh homologous faeces, equivalent freeze-dried and reconstituted faeces or a “synthetic” faecal composition. The fresh homologous faeces does not include an antibiotic resistant population.

[0034] Typically, the composition of the first or second embodiments of the invention is a synthetic faecal composition.

[0035] In a preferred form the synthetic faecal composition comprises a preparation of viable flora which preferably in proportional content, resembles normal healthy human faecal flora which does not include antibiotic resistant populations. Suitable microorganisms may be selected from the following: *Bacteroides*, *Eubacteria*, *Fusobacteria*, *Propionibacteria*, *Lactobacilli*, *anaerobic cocci*, *Ruminococcus*, *Escherichia coli*, *Gemmiger*, *Clostridium*, *Desulfomonas*, *Peptostreptococcus*, *Bifidobacterium*, species and, more specifically, bacteria selected from Table 1. Preferably fungi are also present such as *Monilia*.

[0036] In a preferred form the composition of the first or second embodiments of the invention comprises a liquid culture.

[0037] Preferably, the composition of the first or the second embodiments of the present invention is lyophilised, pulverised and powdered. It may then be infused, dissolved such as in saline, as an enema.

[0038] Alternatively the powder may be encapsulated as enteric-coated capsules for oral administration. These capsules may take the form of enteric-coated microcapsules. As a powder it can preferably be provided in a palatable form for reconstitution for drinking or for reconstitution as a food additive. The composition can be provided as a powder for sale in combination with a food or drink. Typically, the food

or drink is a dairy-based product or a soy-based product. The invention therefore also includes a food or food supplement containing a composition according to the first or second embodiment. In a preferred form the food or food supplement contains enteric-coated microcapsules of the composition of the invention. In a preferred form the food is yogurt.

[0039] The powder may be reconstituted also to be infused via naso-duodenal infusion.

[0040] The composition can be combined with other adjuvants such as antacids to dampen bacterial inactivation in the stomach., eg Mylanta, Mucaine, Gastrogel. Acid secretion in the stomach could also be pharmacologically suppressed using H2-antagonists or proton pump inhibitors. Typically, the H2-antagonist is ranitidine. Typically the proton pump inhibitor is omeprazole.

[0041] The composition of the first or second embodiments of the invention is therefore preferably in the form of:

[0042] an enema composition which can be reconstituted with an appropriate diluent, or

[0043] enteric-coated capsules, or

[0044] enteric-coated microcapsules, or

[0045] powder for reconstitution with an appropriate diluent for naso-enteric infusion or colonoscopic infusion, or

[0046] powder for reconstitution with appropriate diluent, flavouring and gastric acid suppression agent for oral ingestion, or

[0047] powder for reconstitution with food or drink, or

[0048] food or food supplement comprising enteric-coated microcapsules of the composition, powder, jelly, or liquid.

[0049] According to a third embodiment of the invention there is provided a method for the treatment and/or prophylaxis of a chronic disorder associated with the presence in the gastrointestinal tract of a mammalian host of abnormal or an abnormal distribution of microflora, which method comprises administering an effective amount of a composition according to the first or second embodiment of the invention.

[0050] In its preferred form the treatment should effect a cure of the symptoms of such disorders. The change of flora is preferably as "near-complete" as possible and the flora is replaced by viable organisms which will crowd out any remaining, original flora.

[0051] The method of the present invention is applicable to animals in general, in particular humans and economically significant domestic animals.

[0052] In the case of humans, the present invention encompasses methods of treatment of chronic disorders associated with the presence of abnormal enteric microflora. Such disorders include but are not limited to those conditions in the following categories:

[0053] gastro-intestinal disorders including irritable bowel syndrome or spastic colon, functional bowel disease (FBD), including constipation predominant FBD, pain predominant FBD, upper abdominal FBD, non-ulcer dyspepsia (NUD), gastro-oesophageal reflux, inflammatory bowel disease including Crohn's disease, ulcerative colitis, indeterminate colitis, collagenous colitis, microscopic colitis, chronic *Clostridium difficile* infection, pseudomembranous colitis, mucous colitis, antibiotic associated colitis, idiopathic or simple constipation, diverticular disease, AIDS enteropathy, small

bowel bacterial overgrowth, coeliac disease, polyposis coli, colonic polyps, chronic idiopathic pseudo obstructive syndrome;

[0054] chronic gut infections with specific pathogens including bacteria, viruses, fungi and protozoa;

[0055] viral gastrointestinal disorders, including viral gastroenteritis, Norwalk viral gastroenteritis, rotavirus gastroenteritis, AIDS related gastroenteritis;

[0056] liver disorders such as primary biliary cirrhosis, primary sclerosing cholangitis, fatty liver or cryptogenic cirrhosis;

[0057] rheumatic disorders such as rheumatoid arthritis, non-rheumatoid arthritides, non rheumatoid factor positive arthritis, ankylosing spondylitis, Lyme disease, and Reiter's syndrome;

[0058] immune mediated disorders such as glomerulonephritis, haemolytic uraemic syndrome, juvenile diabetes mellitus, mixed cryoglobulinaemia, polyarteritis, familial Mediterranean fever, amyloidosis, *scleroderma*, systemic lupus erythematosus, and Behcets syndrome;

[0059] autoimmune disorders including systemic lupus, idiopathic thrombocytopenic purpura, Sjogren's syndrome, haemolytic uremic syndrome or *scleroderma*;

[0060] neurological syndromes such as chronic fatigue syndrome, migraine, multiple sclerosis, amyotrophic lateral sclerosis, myasthenia gravis, Guillain-Barre syndrome, Parkinson's disease, Alzheimer's disease, Chronic Inflammatory Demyelinating Polyneuropathy, and other degenerative disorders;

[0061] psychiatric disorders including chronic depression, schizophrenia, psychotic disorders, manic depressive illness;

[0062] regressive disorders including Aspergers syndrome, Rett syndrome, attention deficit hyperactivity disorder (ADHD), and attention deficit disorder (ADD);

[0063] the regressive disorder, autism;

[0064] sudden infant death syndrome (SIDS), anorexia nervosa;

[0065] dermatological conditions such as, chronic urticaria, acne, dermatitis herpetiformis and vasculitic disorders.

[0066] The above disorders are all characterised by their response to treatment with the method of the present invention.

[0067] Typically the change in enteric flora comprises introduction of an array of predetermined flora into the gastro-intestinal system, and thus in a preferred form the method of treatment comprises substantially completely displacing pathogenic enteric flora in patients requiring such treatment.

[0068] Furthermore, in some of these disorders a short course of antibiotics prior to probiotic treatment may be preferred to rid tissue-invasive pathogens originating in the bowel lumen. For example, in Crohn's disease, anti-tuberculosis therapy may be required for six to twelve weeks before the bowel is cleared out and the flora content exchanged for a predetermined flora.

[0069] Typically the antibiotic is an anti-Clostridial antibiotic such as vancomycin, rifampicin, and nitroimidazole or chloramphenicol. Typically the nitroimidazole is metronidazole.

[0070] In a preferred form of the invention, the method of treatment or prophylaxis further includes administration of at

least one acid suppressant prior to administering, or in co-administration with, the composition of the invention.

[0071] In a preferred form of the invention the method of treatment or prophylaxis further includes nasogastric and/or nasoduodenal washout prior to administering said composition.

[0072] The introduction of the composition into the gastrointestinal system can be effected by enema or per-colonoscopy, via intubation of the small bowel using for example a large bore catheter equipped with distal balloon to effect rapid passage down the jejunum, or via the oral route with enteric-coated capsules, including enteric-coated microcapsules, or via the oral route with a supplemented food or drink.

[0073] In a preferred form the supplemented food or drink is a dairy-based or soy-based product. Typically the supplemented food product is yogurt.

[0074] According to the method of the invention each dose of the composition is in the range of about 10^3 cells to about 10^{13} cells. Preferably each dose is in the range of about 10^5 cells to about 10^{11} cells. More preferably each dose is in the range of about 10^9 cells to about 10^{11} cells. In a preferred form of the invention an initial treatment regimen consisting of about 10^{10} cells per dose is administered about 3 to 6 times per day for a period sufficient to stabilise the gut flora. According to the method of the invention the treatment regimen may then comprise a maintenance dose of about 10^{10} cells per day.

[0075] Furthermore the present invention also relates to the treatment of animals, in particular to the treatment of gastrointestinal disorders in economically important domestic animals, such as cattle, sheep, horses, pigs, goats etc. The method of the present invention has been found to be especially useful in the treatment of the various forms of necrotising enterocolitis which can be a major problem in animal stocks.

[0076] Obviously in the treatment of animals the appropriate composition of microflora will vary according to the species being treated and the constituent normal flora known to inhabit the gut. Thus the composition according to the invention would comprise, a preparation of viable flora which preferably in proportional content, resembles the normal

healthy faecal flora of the species involved. The compositions may be prepared in any of the forms already described and administered accordingly.

Best Method of Performing the Invention

[0077] In the practice of the invention a synthetic faecal composition of predetermined flora in the form of a liquid or dry powdered culture of *Clostridia*, *Bacteroides*, *Peptostreptococcus*, *Escherichia coli*, *Bifidobacterium*, and *Lactobacillus*, which composition does not include antibiotic resistant populations, is prepared as a liquid culture.

[0078] Typically the method of the invention is applicable to a patient suffering from a chronic disorder associated with the presence of abnormal microflora in the gastrointestinal tract such as irritable bowel syndrome.

[0079] In the practice of the invention a composition of predetermined flora in the form of a liquid culture of *Clostridia*, *Bacteroides*, *Peptostreptococcus*, *Escherichia coli*, *Bifidobacterium*, and *Lactobacillus* is ingested by the patient in an amount sufficient to replace and recolonise the dysbiotic flora of the gastrointestinal tract, and reverse the disease process. Alternatively fresh homologous faeces obtained from a disease screened donor are liquefied and mixed with unprocessed bran. The mixture is then homogenised anaerobically under CO_2 cover and infused into the patient per colonoscopy.

[0080] Cure or remission of symptoms is then monitored subjectively and by assessment of stool frequency or other appropriate criteria.

[0081] Using liquid cultures of *Clostridia*, *Bacteroides*, *Peptostreptococcus*, *Escherichia coli*, *Bifidobacterium*, and *Lactobacillus* the inventor has achieved total reversal of colitis, irritable bowel syndrome and constipation.

[0082] As indicated in the method of treatment aspect of the invention, a preparatory course of appropriate antibiotics may be used. For example, Septrin for chronic yersiniosis, Metronidazole for ulcerative colitis, anti-TB therapy in Crohn's disease, or Vancomycin in chronic *Clostridium difficile* infections.

TABLE 1

% of flora ^b	Organism(s)
11.8(0.90)	<i>Bacteroides fragilis</i> ss. <i>Vulgatus</i>
9.9(0.83)	<i>Eubacterium aerofaciens</i>
8.9(0.78)	<i>Bacteroides fragilis</i> ss. <i>Thetaiotaomicron</i>
6.6(0.68)	<i>Peptostreptococcus productus</i> II
6.0(0.64)	<i>Bacteroides fragilis</i> ss. <i>Distasonis</i>
4.4(0.55)	<i>Fusobacterium prausnitzii</i>
3.5(0.49)	<i>Coprococcus eutactus</i>
3.0(0.45)	<i>Eubacterium aerofaciens</i> III
2.8(0.44)	<i>Peptostreptococcus productus</i> I
2.7(0.43)	<i>Ruminococcus bronii</i>
2.6(0.43)	<i>Bifidobacterium adolescentis</i>
2.2(0.39)	<i>Gemmiger formicilis</i> , <i>Bifidobacterium longum</i>
2.1(0.38)	<i>Eubacterium siraeum</i>
1.8(0.35)	<i>Ruminococcus torques</i>
1.7(0.34)	<i>Eubacterium rectale</i> III-H
1.6(0.33)	<i>Eubacterium rectale</i> IV, <i>Eubacterium eligens</i>
1.5(0.32)	<i>Bacteroides eggerthii</i>
1.4(0.31)	<i>Clostridium leptum</i>
1.3(0.29)	<i>Bacteroides fragilis</i> ss. A
1.2(0.29)	<i>Eubacterium bifforme</i>
0.91(0.25)	<i>Bifidobacterium infantis</i>
0.84(0.24)	<i>Eubacterium rectale</i> III-F
0.57(0.20)	<i>Coprococcus comes</i> , <i>Bacteroides capillosus</i>

TABLE 1-continued

% of flora ^b	Organism(s)
0.50(0.18)	<i>Ruminococcus albus</i> , <i>Eubacterium formicigenerans</i> , <i>Eubacterium hallii</i> , <i>Eubacterium ventriosum</i> I, <i>Fusobacterium russii</i>
0.43(0.17)	<i>Ruminococcus obeum</i> , <i>Eubacterium rectale</i> II, <i>Clostridium ramosum</i> I, <i>Lactobacillus leichmanfi</i>
0.36(0.16)	<i>Ruminococcus cailidus</i> , <i>Butyrivibrio crossotus</i>
0.30(0.14)	<i>Acidaminococcus fermentans</i> , <i>Eubacterium ventriosum</i> , <i>Bacteroides fragilis</i> ss. <i>fragilis</i> , <i>Bacteroides AR</i>
0.23(0.12)	<i>Coprococcus catus</i> , <i>Eubacterium hadrum</i> , <i>Eubacterium cylindroides</i> , <i>Eubacterium ruminantium</i> , <i>Eubacterium CH-1</i> , <i>Staphylococcus epidermidis</i>
0.17(0.10)	<i>Peptostreptococcus BL</i> , <i>Eubacterium limosum</i> , <i>Bacteroides praeacutus</i> , <i>Bacteroides L</i> , <i>Fusobacterium mortiferum</i> I, <i>Fusobacterium naviforme</i> , <i>Clostridium innocuum</i> , <i>Clostridium ramosum</i> , <i>Propionibacterium acnes</i> , <i>Ruminococcus flavefaciens</i>
0.10(0.08)	<i>Ruminococcus AT</i> , <i>Peptococcus AU-1</i> , <i>Eubacterium AG</i> , -AK, -AL, -AL-1, -AN; <i>Bacteroides fragilis</i> ss. <i>ovatus</i> , -ss. d, -ss. f; <i>Bacteroides L-1</i> , L-5; <i>Fusobacterium nucleatum</i> , <i>Fusobacterium mortiferum</i> , <i>Escherichia coli</i> , <i>Streptococcus morbillorum</i>
0.05(0.05)	<i>Peptococcus magnus</i> , <i>Peptococcus G</i> , -AU-2; <i>Streptococcus intermedius</i> , <i>Ruminococcus lactaris</i> , <i>Ruminococcus CO Gemmiger X</i> , <i>Coprococcus BH</i> , -CC; <i>Eubacterium tenue</i> , <i>Eubacterium ramulus</i> , <i>Eubacterium AE</i> , -AG-H, -AG-M, -AJ, -BN-1; <i>Bacteroides clostridiformis</i> ss. <i>clostridiformis</i> , <i>Bacteroides coagulans</i> , <i>Bacteroides orails</i> , <i>Bacteroides ruminicola</i> ss. <i>brevis</i> , -ss. <i>ruminicola</i> , <i>Bacteroides splanchnicus</i> , <i>Desulfomonas pigra</i> , <i>Bacteroides L-4</i> , -N-i; <i>Fusobacterium H</i> , <i>Lactobacillus G</i> , <i>Succinivibrio A</i>

^bThe percentage of the faecal population (the standard deviation of the estimate is given in parentheses).

[0083] The invention will now be further described with reference to the following non-limiting examples.

EXAMPLES

Formulations

[0084] The probiotic therapeutic agents may be prepared in liquid culture anaerobically or aerobically (depending on bacterium cultured) in pure form. Alternatively the probiotics may be cultured on solid media and scraped into a liquid carrier. The resulting product may be spray-dried into a powder form and encapsulated or combined with excipients to be delivered in sachets.

[0085] Combinations of *Clostridia*, *Escherichia coli*, *Bacteroides*, and *Peptostreptococcus* with or without *Lactobacilli*, *Bifidobacteria* and *Eubacteria* may be used in varying disorders.

Example No 1

43 Year Old Female

[0086] Patient with long standing constipation not responsive to high-dose fibre usage together with prokinetics and standard anti-constipation treatments, was treated with increasing doses of orally administered bacterial mix (mixture composition included *Clostridium innocuum*, *bifermantans*, *butyricum*, together with *Bacteroides fragilis*, *thetaiotaomicron* and *uniformis*. Three strains of *Escherichia coli* were also included, as was *Lactobacillus*). This was ingested twice daily in the first two weeks and then daily thereafter. The patient was not given any pre-treatment purgative nor any antibiotics. However, she did take Ranitidine (an acid suppressant) three hours prior to ingestion of the bacterial mix. Two weeks after commencing the treatment the patient's constipation—which would prevent her from defecating for up to four days—reversed to increased frequency with reduction of bloating. Initially, gas production increased and there was burbulence and gurgling in the abdomen but after four weeks of treatment the patient was defecating on a daily basis with no sensation of incomplete emptying and an almost total

absence of bloating. Following the treatment she remained virtually normal, defecating on a daily basis with 3 month follow up.

Example No 2

4½ Year Old Male

[0087] Patient with 3 year history of diagnosis of autism associated with Irritable Bowel Syndrome characterised by constipation alternating with diarrhoea and flatulence, with foul motions, was treated with oral administration of bacterial mix consisting of *Clostridium bifermantans*, *Clostridium innocuum*, and *Clostridium butyricum* in combination with three strains of *Escherichia coli*, three strains of *bacteroides* and *Peptostreptococcus productus*. These were ingested following acid suppression with Ranitidine and were at first taken 3 times daily, reducing to twice daily and then once daily maintenance for eight weeks. The patient's autistic symptoms were reversed quite dramatically with word power increasing from 20 to 200 words (counted by teacher at special 'autistic' school), he began to sleep through the night, and his IBS-type symptoms reverted to near-normality with less constipation, less diarrhoea and less foul flatulence. He developed eye contact, was able to speak sentences up to six words constructed to commands and he began to look, to the untrained eye, as a relatively normal child by about week 10.

Example 3

Male Child, 5½ Years Old

[0088] Male child, 5½ years of age with autism symptoms dating back to age of around 15 months—but diagnosed significantly later. The patient presented initially with gastrointestinal symptoms in association with classical autism—for treatment of the bowel symptoms. Although stool test did not indicate any specific pathogen the bowel symptoms resembled those of a chronic infection or adult Irritable Bowel Syndrome (IBS), ie intermittent diarrhoea, constipation, cramping, colicky pain, inability to sleep at night, occasional explosive diarrhoea and incontinence. The patient was

treated with orthostatic lavage using sodium pico-sulfate followed by water to produce voluminous diarrhoea and to flush out the enteric contents. He was then given 125 mg Vancomycin three times daily orally followed by oral re-colonisation with bacteria at a concentration of 10^9 through to 10^{10} , suspended in yoghurt—of strains which included *bacteroides*, *Escherichia coli*, and non pathogenic *Clostridia*—including *Clostridium innocuum*, *bifementans* and *ramosum*. The response was quite noticeable, in the reversal of the abnormal stool function towards normality. The patient was also able to sleep through the night without any explosive diarrhoea and produced formed stools within five days of commencing the bacterial therapy. While the bacteriotherapy was continued the bowel symptoms were well controlled. Within 3-4 weeks of missing out the treatment for a week or two some of the symptoms would begin to recur. This suggested that the abnormal bacterial flora was suppressed rather than being cured with this treatment in this patient. The unexpected finding however, was a noticeable and marked reversal of symptoms of autism. Whereas previously repetitive movements were present with lack of eye contact, eye contact returned fairly rapidly together with cessation of repetitive movement and progressive increase of word power from around 20 words to around 600 words by the sixth month of treatment. The therapy continues now for more than 12 months with sustained reversal of autism and IBS symptoms.

Example 4

Male Child, 7 Years Old

[0089] A seven year old male patient was referred for treatment initially of bowel problems. He had developed autism between age 1 and 2 years characterised by lack of eye contact, repetitive movements, poorly developed cognitive abilities, vocabulary of fewer than 20 words. The marked bowel symptoms were characterised by either constipation or large voluminous motions, sometimes diarrhoea and explosive stools. Stool examination was negative.

[0090] The patient was given a pre-treatment of Vancomycin 125 mg twice daily and at one week he was given an orthostatic lavage consisting of picosulfate preparation which flushed out his bowel. He was then given twice daily oral bacteriotherapy consisting of cultures containing living probiotics. These included several *bacteroides* species, *Escherichia coli* and non-pathogenic *Clostridia* such as *Clostridium butyricum*, *Clostridium bifementans* and *Clostridium innocuum*. *Within two weeks the bowel symptoms reversed to normal defecation with soft, formed stool—once or twice per day. Constipation disappeared, eye contact returned over the next six weeks and vocabulary and word use quite dramatically improved, to everyone's surprise. When followed for eight months over 600 words could be counted in the vocabulary with sentences of up to eight words being constructed where previously this was not possible. Some abstract thinking was noted by teachers at the special autism school. Parents in particular noted reduced aggression, greater co-operation, and general increasing ability to develop a more normal relationship with the child. Repetitive action also disappeared.*

Example 5

Male Child, 6 Years Old

[0091] A male patient aged 6 was referred to the clinic for treatment of chronic diarrhoea and at times incontinence. The

child had been autistic since the age of one year and three months. The diagnosis however was delayed. He had slow cognitive development and very limited vocabulary. There was virtually absent eye contact and at times violent and explosive behaviour. The greatest problem with management was that of control of defecation as the child developed a fascination with the stools which would then be spread over furniture and walls. This brought severe pressure upon the family with respect to difficulty with management. Stool test was collected and again was negative for any pathogen. The patient was given Vancomycin 250 mg twice daily for 10 days after which a polyethylene glycol orthostatic lavage achieved a large volume flush of the bowel. He was then given twice daily oral bacteriotherapy in a neutral yogurt as a carrier. Within one week the bowel function returned to virtual normality. However, the behavioural changes were just as rapid in reversing again characterised by fairly rapid reduction in aggressiveness and uncontrollable behaviour, sleeping through the night, increased eye contact, and progressively increased word power. The behaviour of spreading stools also disappeared, more as a behavioural change than learnt phenomenon. The patient was continued on medications for over a year and progressively improved in all parameters—at times fluctuating in severity.

1.-182. (canceled)

183. A method for treating autism in a subject in need thereof, the method comprising administering to the subject an amount of a pharmaceutical composition effective for treating the autism,

wherein the pharmaceutical composition is selected from the group consisting of:

- a. a disease screened fresh homologous feces;
- b. a freeze-dried, or a freeze-dried and reconstituted feces;
- c. a reconstituted feces, reconstituted using cultured viable non-pathogenic or attenuated microorganisms;
- d. a synthetic fecal composition of predetermined flora;
- e. a synthetic or reconstituted fecal composition comprising a preparation of viable flora in proportional content that resembles a normal healthy human fecal flora, which does not include antibiotic resistant populations;
- f. a composition comprising viable, non-pathogenic colonic bacterial cells selected from the group consisting of a *Clostridia*, a *Collinsella*, a *Bacteroides*, a *Fusobacteria*, a *Propionibacteria*, a *Lactobacilli*, an anaerobic cocci, a *Ruminococcus*, an *E. coli*, a *Gemmiger*, a *Desulfomonas*, a *Peptostreptococcus*, a *Bifidobacteria* and any combination thereof;
- g. a composition comprising viable, non-pathogenic colonic bacterial components of fecal flora, wherein said bacterial components comprise *Clostridium bifementans*, *Clostridium innocuum*, *Clostridium butyricum*, *Escherichia coli*, *Bacteroides* and *Peptostreptococcus productus*;
- h. a composition comprising viable, non-pathogenic colonic bacterial components of fecal flora, wherein said bacterial components comprise a *Bacteroides*, an *Escherichia coli*, and a non-pathogenic *Clostridia*,
- i. a composition comprising a plurality of viable non-pathogenic *Clostridia* spores and a plurality of viable non-pathogenic *Collinsella* cells,
- j. a composition comprising a plurality of viable non-pathogenic *Clostridia*,

k. a composition comprising a plurality of viable non-pathogenic *Clostridia* and a plurality of viable non-pathogenic *Collinsella*,

l. a composition comprising viable non-pathogenic *Clostridia* spores, a viable non-pathogenic *Bacteroides*, and a viable non-pathogenic *Escherichia coli*.

184. The method of claim **083**, wherein said viable non-pathogenic *Clostridia* are from a species selected from the group consisting of *Clostridium absonum*, *Clostridium argentinense*, *Clostridium baratii*, *Clostridium bifermentans*, *Clostridium botulinum*, *Clostridium butyricum*, *Clostridium cadaveris*, *Clostridium camis*, *Clostridium celatum*, *Clostridium chauvoei*, *Clostridium clostridioforme*, *Clostridium cochlearium*, *Clostridium difficile*, *Clostridium fallax*, *Clostridium felsineum*, *Clostridium ghonii*, *Clostridium glycolicum*, *Clostridium haemolyticum*, *Clostridium hastiforme*, *Clostridium histolyticum*, *Clostridium indolis*, *Clostridium innocuum*, *Clostridium irregulare*, *Clostridium limosum*, *Clostridium malenominatum*, *Clostridium novyi*, *Clostridium oroticum*, *Clostridium paraputrificum*, *Clostridium perfringens*, *Clostridium piliforme*, *Clostridium putrefaciens*, *Clostridium putrificum*, *Clostridium ramosum*, *Clostridium sardiniense*, *Clostridium sartagoforme*, *Clostridium scindens*, *Clostridium septicum*, *Clostridium sordellii*, *Clostridium sphenoides*, *Clostridium spiroforme*, *Clostridium sporogenes*, *Clostridium subterminale*, *Clostridium symbiosum*, *Clostridium tertium*, *Clostridium tetani*, *Clostridium welchii*, and *Clostridium villosum*.

185. The method of claim **183**, wherein said viable non-pathogenic *Clostridia* are from a species selected from the group consisting of *Clostridium bifermentans*, *Clostridium innocuum*, and *Clostridium butyricum*.

186. The method of claim **183**, wherein said viable non-pathogenic *Collinsella* comprise *Collinsella aerofaciens*.

187. The method of claim **183**, wherein said pharmaceutical composition further comprises an acid suppressant, an antacid, an H2 antagonist, a proton pump inhibitor or a combination thereof.

188. The method of claim **183**, wherein said pharmaceutical composition is formulated as an enteric coated capsule, an enteric coated microcapsule, a powder suitable for reconstitution, a naso-duodenal infusion, or for delivery in the form of an enema or a colonoscopic infusion.

189. The method of claim **1830**, wherein said pharmaceutical composition is added to a food, a food additive, a dairy-based product, a soy-based product or a derivative thereof, a jelly, or a yogurt.

190. The method of claim **183**, wherein no existing enteric microflora is removed from said subject prior to administration of said pharmaceutical composition.

191. The method of claim **183**, wherein some existing enteric microflora is removed from said subject prior to administration of said pharmaceutical composition.

192. The method of claim **183**, wherein the subject is pretreated with an antibiotic prior to administration of said pharmaceutical composition.

193. The method of claim **183**, wherein the subject is pretreated with an acid suppressant prior to administration of the pharmaceutical composition.

194. The method of claim **183**, wherein the pharmaceutical composition is administered three times daily (tid).

195. The method of claim **183**, wherein the subject is pretreated with an orthostatic lavage prior to administration of the pharmaceutical composition.

196. The method of claim **183**, wherein the subject is pretreated with an orthostatic lavage and antibiotic treatment prior to administration of the pharmaceutical composition.

197. The method of claim **183**, wherein the subject is pretreated with an orthostatic lavage and antibiotic treatment prior to administration of the pharmaceutical composition, and the pharmaceutical composition comprises a plurality of bacteria comprising a *Bacteroides*, an *Escherichia coli*, and a non-pathogenic *Clostridia*,

and the pharmaceutical composition at a concentration of 10^9 through to 10^{10} .

198. The method of claim **183**, wherein the subject is pretreated with an antibiotic prior to administration of the pharmaceutical composition,

and after the pre-treatment with the antibiotic the subject is pretreated with an orthostatic lavage prior to administration of the pharmaceutical composition.

199. The method of claim **183**, wherein the pharmaceutical composition comprises between about 10^3 and about 10^{13} viable bacterial cells.

200. The method of claim **199**, wherein the pharmaceutical composition comprises between about 10^5 and about 10^{11} viable bacterial cells.

201. The method claim **200**, wherein the pharmaceutical composition comprises between about 10^9 and about 10^{11} viable bacterial cells.

202. The method of claim **201**, wherein the pharmaceutical composition comprises about 10^{10} viable bacterial cells.

203. The method of claim **183**, wherein the autism is associated with Irritable Bowel Syndrome or gastrointestinal symptoms.

* * * * *