Office de la Propriété Intellectuelle du Canada
Un organisme d'Industrie Canada

Canadian Intellectual Property Office
An agency of Industry Canada

CA 2604115 C 2014/11/04

(11)(21) 2 604 115

(12) BREVET CANADIEN
CANADIAN PATENT

(13) C

(86) Date de dépôt PCT/PCT Filing Date: 2006/04/11
(87) Date publication PCT/PCT Publication Date: 2006/10/19
(45) Date de délivrance/Issue Date: 2014/11/04
(85) Entrée phase nationale/National Entry: 2007/10/09
(86) N° demande PCT/PCT Application No.: US 2006/014352
(87) N° publication PCT/PCT Publication No.: 2006/110917
(30) Priorité/Priority: 2005/04/11 (US60/670,896)

(51) Cl.Int./Int.Cl. *C07D 491/20* (2006.01), *A61K 31/407* (2006.01), *A61P 29/00* (2006.01), *C07D 491/10* (2006.01)

(72) Inventeurs/Inventors:
CHAFEEV, MIKHAIL, CA;
CHOWDHURY, SULTAN, CA;
FRASER, ROBERT, CA;
FU, JIANMIN, CA;
KAMBOJ, RAJENDER K., CA;
HOU, DUANJIE, CA;
...

(73) Propriétaire/Owner:
XENON PHARMACEUTICALS INC., CA

(74) Agent: AITKEN KLEE LLP

(54) Titre : COMPOSES DE SPIRO-OXINDOLE ET UTILISATIONS ASSOCIEES EN TANT QU'AGENTS THERAPEUTIQUES

(54) Title: SPIRO-OXINDOLE COMPOUNDS AND THEIR USES AS THERAPEUTIC AGENTS



(I)

(57) Abrégé/Abstract:
This invention is directed to spiro-oxindole compounds of formula (I): wherein k, j, Q, R^1, R^{2a}, R^{2b}, R^{2c}, R^{2d}, R^{3a}, R^{3b}, R^{3c}, and R^{3d} are as defined herein, as a stereoisomer, enantiomer, tautomer thereof or mixtures thereof; or a pharmaceutically acceptable salt, solvate or prodrug thereof, which are useful for the treatment and/or prevention of sodium channel-mediated diseases or conditions, such as pain. Pharmaceutical compositions comprising the compounds and methods of preparing and using the compounds are also disclosed.

CA 2604115 C 2014/11/04

(11)(21) 2 604 115

(13) C

(72) **Inventeurs(suite)/Inventors(continued)**: LIU, SHIFENG, CA; BAGHERZADEH, MEHRAN SEID, CA; SVIRIDOV, SERGUEI V., CA; SUN, SHAOYI, CA; SUN, JIANYU, CA; CHAKKA, NAGASREE, CA; HSIEH, TOM, CA; RAINA, VANDNA, CA

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
19 October 2006 (19.10.2006)

PCT

(10) International Publication Number
WO 2006/110917 A3

(51) International Patent Classification:
C07D 498/10 (2006.01) *A61P 29/00* (2006.01)
A61K 31/404 (2006.01)

(21) International Application Number:
PCT/US2006/014352

(22) International Filing Date: 11 April 2006 (11.04.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/670,896 11 April 2005 (11.04.2005) US

(71) Applicant *(for all designated States except US)*: **XENON PHARMACEUTICALS INC.** [CA/CA]; 3650 Gilmore Way, Burnaby, British Columbia V5G 4W8 (CA).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: **CHAFEEV, Mikhail** [CA/CA]; 1403-4788 Hazel Street, Burnaby, British Columbia V5H 4V9 (CA). **CHOWDHURY, Sultan** [CA/CA]; 97 Orleans Drive, Scarborough, Ontario M1C 4Y3 (CA). **FRASER, Robert** [CA/CA]; 1398 East 15th Street, North Vancouver, British Columbia V7J 1K8 (CA). **FU, Jianmin** [US/CA]; 3170 Quintette Crescent, Coquitlam, British Columbia V3E 3B6 (CA). **KAMBOJ, Rajender** [CA/CA]; 5809 Sperling Avenue, Burnaby, British Columbia V5E 2T6 (CA). **HOU, Duanjie** [CA/CA]; #46-7088 17th Avenue, Burnaby, British Columbia V3N 1K5 (CA). **LIU, Shifeng** [CA/CA]; 15-2615 Fortress Drive, Port Coquitlam, British Columbia V3C 6E8 (CA). **BAGHERZADEH, Mehran Seid** [CA/CA]; 615 East Osborne Road, North Vancouver, British Columbia V7N 1M8 (CA). **SVIRIDOV, Serguei** [CA/CA]; 1904-5885 Olive Avenue, Burnaby, British Columbia V5H 4N8 (CA). **SUN, Shaoyi** [CA/CA]; #5-1200 Brunette Avenue, Coquitlam, British Columbia V3K 1G3 (CA). **SUN, Jianyu** [CN/CA]; 23-8700 Bennett Road, Richmond, British Columbia V6Y 1N7 (CA). **CHAKKA, Nagasree** [CA/CA]; 7480 Vista Crescent, Burnaby, British Columbia V5E 2E1 (CA). **HSIEH, Tom** [CA/CA]; 8392 Hollis Place, Burnaby, British Columbia V5J 4Z8 (CA). **RAINA, Vandna** [IN/CA]; 6654 Linden Avenue, Burnaby, British Columbia V5E 3G4 (CA).

(74) Agents: **ROTH, Carol, J.** et al.; Seed Intellectual Property Law Group PLLC, Suite 6300, 701 Fifth Avenue, Seattle, Washington 98104-7092 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— *with international search report*

(88) Date of publication of the international search report:
4 January 2007

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SPIRO-OXINDOLE COMPOUNDS AND THEIR USES AS THERAPEUTIC AGENTS



(I)

(57) Abstract: This invention is directed to spiro-oxindole compounds of formula (I): wherein k, j, Q, R^1, R^{2a}, R^{2b}, R^{2c}, R^{2d}, R^{3a}, R^{3b}, R^{3c}, and R^{3d} are as defined herein, as a stereoisomer, enantiomer, tautomer thereof or mixtures thereof; or a pharmaceutically acceptable salt, solvate or prodrug thereof, which are useful for the treatment and/or prevention of sodium channel-mediated diseases or conditions, such as pain. Pharmaceutical compositions comprising the compounds and methods of preparing and using the compounds are also disclosed.

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 340

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 340

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHIER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

SPIRO-OXINDOLE COMPOUNDS AND THEIR USES AS THERAPEUTIC AGENTS

FIELD OF THE INVENTION

The present invention is directed to spiro-oxindole compounds. In particular, this invention is directed to spiro-oxindole compounds that are sodium channel blockers and are therefore useful in treating sodium channel-mediated diseases or conditions, such as pain, as well as other diseases and conditions associated with the mediation of sodium channels.

BACKGROUND OF THE INVENTION

Voltage-gated sodium channels, transmembrane proteins that initiate action potentials in nerve, muscle and other electrically excitable cells, are a necessary component of normal sensation, emotions, thoughts and movements (Catterall, W.A., *Nature* (2001), Vol. 409, pp. 988-990). These channels consist of a highly processed alpha subunit that is associated with auxiliary beta subunits. The pore-forming alpha subunit is sufficient for channel function, but the kinetics and voltage dependence of channel gating are in part modified by the beta subunits (Goldin et al., *Neuron* (2000), Vol. 28, pp. 365–368). Each alpha-subunit contains four homologous domains, I to IV, each with six predicted transmembrane segments. The alpha-subunit of the sodium channel, forming the ion-conducting pore and containing the voltage sensors regulating sodium ion conduction has a relative molecular mass of 260,000. Electrophysiological recording, biochemical purification, and molecular cloning have identified ten different sodium channel alpha subunits and four beta subunits (Yu, F.H., et al., *Sci. STKE* (2004), 253; and Yu, F.H., et al., *Neurosci.* (2003), 20:7577-85).

The hallmarks of sodium channels include rapid activation and inactivation when the voltage across the plasma membrane of an excitable cell is depolarized (voltage-dependent gating), and efficient and selective conduction of sodium ions through conducting pores intrinsic to the structure of the protein (Sato, C., *et al., Nature* (2001), 409:1047–1051). At negative or hyperpolarized membrane potentials, sodium channels are closed. Following membrane depolarization, sodium channels open rapidly and then inactivate. Channels only conduct currents in the open state and, once inactivated, have to return to the resting state, favoured by membrane hyperpolarization, before they can reopen. Different sodium channel subtypes vary in the voltage range over which they activate and inactivate as well as their activation and inactivation kinetics.

The sodium channel family of proteins has been extensively studied and shown to be involved in a number of vital body functions. Research in this area has identified variants of the alpha subunits that result in major changes in channel function and activities, which can ultimately lead to major pathophysiological conditions. Implicit with function, this family of proteins are considered prime points of therapeutic intervention. $Na_V1.1$ and $Na_V1.2$ are highly expressed in the brain (Raymond, C.K., et al., *J. Biol. Chem.* (2004), 279(44):46234-41) and are vital to normal brain function. In humans, mutations in $Na_V1.1$ and $Na_V1.2$ result in severe epileptic states and in some cases mental decline (Rhodes, T.H., et al., *Proc. Natl. Acad. Sci.* USA (2004),101(30):11147-52; Kamiya, K., et al., *J. Biol. Chem.* (2004), 24(11) :2690-8; Pereira, S., et al., *Neurology* (2004), 63(1):191-2). As such both channels have been considered as validated targets for the treatment of epilepsy (see PCT Published Patent Publication No. WO 01/38564).

$Na_V1.3$ is broadly expressed throughout the body (Raymond, C.K., et al., *op. cit.*). It has been demonstrated to have its expression upregulated in the dorsal horn sensory neurons of rats after nervous system injury (Hains, B.D., et al., *J. Neurosci.* (2003), 23(26):8881-92). Many experts in the field have considered $Na_V1.3$ as a suitable target for pain therapeutics (Lai, J., et al., *Curr. Opin. Neurobiol.* (2003), (3):291-72003; Wood, J.N., et al., *J. Neurobiol.* (2004), 61(1):55-71; Chung, J.M., et al., *Novartis Found Symp.* (2004), 261:19-27; discussion 27-31, 47-54).

$Na_V1.4$ expression is essentially limited to muscle (Raymond, C.K., et al., *op. cit.*). Mutations in this gene have been shown to have profound effects on muscle function including paralysis, (Tamaoka A., *Intern. Med.* (2003), (9):769-70). Thus, this channel can be considered a target for the treatment of abnormal muscle contractility, spasm or paralysis.

The cardiac sodium channel, $Na_V1.5$, is expressed mainly in the heart ventricles and atria (Raymond, C.K., et al., *op. cit.*), and can be found in the sinovial node, ventricular node and possibly Purkinje cells. The rapid upstroke of the cardiac action potential and the rapid impulse conduction through cardiac tissue is due to the opening of $Na_V1.5$. As such, $Na_V1.5$ is central to the genesis of cardiac arrhythmias. Mutations in human $Na_V1.5$ result in multiple arrhythmic syndromes, including, for example, long QT3 (LQT3), Brugada syndrome (BS), an inherited cardiac conduction defect, sudden unexpected nocturnal death syndrome (SUNDS) and sudden infant death syndrome (SIDS) (Liu, H. et al., *Am. J. Pharmacogenomics* (2003), 3(3):173-9). Sodium channel blocker therapy has been used extensively in treating cardiac arrhythmias. The first

antiarrhythmic drug, quinidine, discovered in 1914, is classified as a sodium channel blocker.

$Na_V1.6$ encodes an abundant, widely distributed voltage-gated sodium channel found throughout the central and peripheral nervous systems, clustered in the nodes of Ranvier of neural axons (Caldwell, J.H., et al., *Proc. Natl. Acad. Sci.* USA (2000), 97(10): 5616-20). Although no mutations in humans have been detected, $Na_V1.6$ is thought to play a role in the manifestation of the symptoms associated with multiple sclerosis and has been considered as a target for the treatment of this disease (Craner, M.J., et al., *Proc. Natl. Acad. Sci.* USA (2004), 101(21):8168-73).

$Na_V1.7$ was first cloned from the pheochromocytoma PC12 cell line (Toledo-Aral, J. J., et al., *Proc. Natl.Acad. Sci.* USA (1997), 94:1527–1532). Its presence at high levels in the growth cones of small-diameter neurons suggested that it could play a role in the transmission of nociceptive information. Although this has been challenged by experts in the field as $Na_V1.7$ is also expressed in neuroendocrine cells associated with the autonomic system (Klugbauer, N., et al., *EMBO J.* (1995), 14(6):1084-90) and as such has been implicated in autonomic processes. The implicit role in autonomic functions was demonstrated with the generation of $Na_V1.7$ null mutants; deleting $Na_V1.7$ in all sensory and sympathetic neurons resulted in a lethal perinatal phenotype. (Nassar, et al., *Proc. Natl. Acad. Sci.* USA (2004), 101(34):12706-11.). In contrast, by deleting the $Na_V1.7$ expression in a subset of sensory neurons that are predominantly nociceptive, a role in pain mechanisms, was demonstrated (Nassar, et al., *op. cit.*). Further support for $Na_V1.7$ blockers active in a subset of neurons is supported by the finding that two human heritable pain conditions, primary erythermalgia and familial rectal pain, have been shown to map to $Na_V1.7$ (Yang, Y., et al., *J. Med. Genet.* (2004), 41(3):171-4).

The expression of $Na_V1.8$ is essentially restricted to the DRG (Raymond, C.K., et al., *op. cit.*). There are no identified human mutations for $Na_V1.8$. However, $Na_V1.8$-null mutant mice were viable, fertile and normal in appearance. A pronounced analgesia to noxious mechanical stimuli, small deficits in noxious thermoreception and delayed development of inflammatory hyperalgesia suggested to the researchers that $Na_V1.8$ plays a major role in pain signalling (Akopian, A. N., et al., *Nat. Neurosci.* (1999), 2(6): 541-8). Blocking of this channel is widely accepted as a potential treatment for pain (Lai, J, et al., *op. cit.*; Wood, J.N., et al., *op. cit.*; Chung, J.M., et al., *op. cit.*). PCT Published Patent Application No. WO03/037274A2 describes pyrazole-amides and sulfonamides for the treatment of central or peripheral nervous system

conditions, particularly pain and chronic pain by blocking sodium channels associated with the onset or recurrance of the indicated conditions. PCT Published Patent Application No. WO03/037890A2 describes piperidines for the treatment of central or peripheral nervous system conditions, particularly pain and chronic pain by blocking sodium channels associated with the onset or recurrence of the indicated conditions. The compounds, compositions and methods of these inventions are of particular use for treating neuropathic or inflammatory pain by the inhibition of ion flux through a channel that includes a PN3 ($Na_V1.8$) subunit.

The tetrodotoxin insensitive, peripheral sodium channel $Na_V1.9$, disclosed by Dib-Hajj, S.D., et al. (see Dib-Hajj, S.D., et al., *Proc. Natl. Acad. Sci.* USA (1998), 95(15):8963-8) was shown to reside solely in the dorsal root ganglia. It has been demonstrated that $Na_V1.9$ underlies neurotrophin (BDNF)-evoked depolarization and excitation, and is the only member of the voltage gated sodium channel superfamily to be shown to be ligand mediated (Blum, R., Kafitz, K.W., Konnerth, A., *Nature* (2002), 419 (6908):687-93). The limited pattern of expression of this channel has made it a candidate target for the treatment of pain (Lai, J, et al., *op. cit.*; Wood, J.N., et al., *op. cit.*; Chung, J.M. et al., *op. cit.*).

NaX is a putative sodium channel, which has not been shown to be voltage gated. In addition to expression in the lung, heart, dorsal root ganglia, and Schwann cells of the peripheral nervous system, NaX is found in neurons and ependymal cells in restricted areas of the CNS, particularly in the circumventricular organs, which are involved in body-fluid homeostasis (Watanabe, E., et al., *J. Neurosci.* (2000), 20(20):7743-51). NaX-null mice showed abnormal intakes of hypertonic saline under both water- and salt-depleted conditions. These findings suggest that the NaX plays an important role in the central sensing of body-fluid sodium level and regulation of salt intake behaviour. Its pattern of expression and function suggest it as a target for the treatment of cystic fibrosis and other related salt regulating maladies.

Studies with the sodium channel blocker tetrodotoxin (TTX) used to lower neuron activity in certain regions of the brain, indicate its potential use in the treatment of addiction. Drug-paired stimuli elicit drug craving and relapse in addicts and drug-seeking behavior in rats. The functional integrity of the basolateral amygdala (BLA) is necessary for reinstatement of cocaine-seeking behaviour elicited by cocaine-conditioned stimuli, but not by cocaine itself. BLA plays a similar role in reinstatement of heroin-seeking behavior. TTX-induced inactivation of the BLA on conditioned and heroin-primed reinstatement of extinguished heroin-seeking behaviour in a rat model

(Fuchs, R.A. and See, R.E., *Psychopharmacology* (2002) 160(4):425-33).

This closely related family of proteins has long been recognised as targets for therapeutic intervention. Sodium channels are targeted by a diverse array of pharmacological agents. These include neurotoxins, antiarrhythmics, anticonvulsants and local anesthetics (Clare, J.J., et al., *Drug Discovery Today* (2000) 5:506-520). All of the current pharmacological agents that act on sodium channels have receptor sites on the alpha subunits. At least six distinct receptor sites for neurotoxins and one receptor site for local anesthetics and related drugs have been identified (Cestèle, S. et al., *Biochimie* (2000), Vol. 82, pp. 883-892).

The small molecule sodium channel blockers or the local anesthetics and related antiepileptic and antiarrhythmic drugs, interact with overlapping receptor sites located in the inner cavity of the pore of the sodium channel (Catterall, W.A., *Neuron* (2000), 26:13-25). Amino acid residues in the S6 segments from at least three of the four domains contribute to this complex drug receptor site, with the IVS6 segment playing the dominant role. These regions are highly conserved and as such most sodium channel blockers known to date interact with similar potency with all channel subtypes. Nevertheless, it has been possible to produce sodium channel blockers with therapeutic selectivity and a sufficient therapeutic window for the treatment of epilepsy (e.g. lamotrignine, phenytoin and carbamazepine) and certain cardiac arrhythmias (e.g. lignocaine, tocainide and mexiletine). However, the potency and therapeutic index of these blockers is not optimal and have limited the usefulness of these compounds in a variety of therapeutic areas where a sodium channel blocker would be ideally suited.

<u>Management of Acute and Chronic Pain</u>

Drug therapy is the mainstay of management for acute and chronic pain in all age groups, including neonates, infants and children. The pain drugs are classified by the American Pain Society into three main categories: 1) non-opioid analgesics-acetaminophen, and non-steroidal anti-inflammatory drugs (NSAIDs), including salicylates (e.g. aspirin*), 2) opioid analgesics and 3) co-analgesics.

Non-opioid analgesics such as acetaminophen and NSAIDs are useful for acute and chronic pain due to a variety of causes including surgery, trauma, arthritis and cancer. NSAIDs are indicated for pain involving inflammation because acetaminophen lacks anti-inflammatory activity. Opioids also lack anti-inflammatory activity. All NSAIDs inhibit the enzyme cyclooxygenase (COX), thereby inhibiting prostaglandin synthesis and reducing the inflammatory pain response. There are at least two COX isoforms, COX-1 and COX-2. Common non-selective COX inhibitors

*Trademark

include, ibuprofen and naproxen. Inhibition of COX-1, which is found in platelets, GI tract, kidneys and most other human tissues, is thought to be associated with adverse effects such as gastrointestinal bleeding. The development of selective COX-2 NSAIDs, such as Celecoxib, Valdecoxib and Rofecoxib, have the benefits of non-selective NSAIDs with reduced adverse effect profiles in the gut and kidney. However, evidence now suggests that chronic use of certain selective COX-2 inhibitors can result in an increased risk of stroke occurrence.

The use of opioid analgesics is recommended by the American Pain Society to be initiated based on a pain-directed history and physical that includes repeated pain assessment. Due to the broad adverse effect profiles associated with opiate use, therapy should include a diagnosis, integrated interdisciplinary treatment plan and appropriate ongoing patient monitoring. It is further recommended that opioids be added to non-opioids to manage acute pain and cancer related pain that does not respond to non-opioids alone. Opioid analgesics act as agonists to specific receptors of the mu and kappa types in the central and peripheral nervous system. Depending on the opioid and its formulation or mode of administration it can be of shorter or longer duration. All opioid analgesics have a risk of causing respiratory depression, liver failure, addiction and dependency, and as such are not ideal for long-term or chronic pain management.

A number of other classes of drugs may enhance the effects of opioids or NSAIDSs, have independent analgesic activity in certain situations, or counteract the side effects of analgesics. Regardless of which of these actions the drug has, they are collectively termed "coanalgesics". Tricyclic antidepressants, antiepileptic drugs, local anaesthetics, glucocorticoids, skeletal muscle relaxants, anti-spasmodil agents, antihistamines, benzodiazepines, caffeine, topical agents (e.g. capsaicin), dextroamphetamine and phenothizines are all used in the clinic as adjuvant therapies or individually in the treatment of pain. The antiepeileptic drugs in particular have enjoyed some success in treating pain conditions. For instance, Gabapentin, which has an unconfirmed therapeutic target, is indicated for neuropathic pain. Other clinical trials are attempting to establish that central neuropathic pain may respond to ion channel blockers such as blockers of calcium, sodium and/or NMDA (N-methyl-D-aspartate) channels. Currently in development are low affinity NMDA channel blocking agents for the treatment of neuropathic pain. The literature provides substantial pre-clinical electrophysiological evidence in support of the use of NMDA antagonists in the treatment of neuropathic pain. Such agents also may find use in the control of pain

after tolerance to opioid analgesia occurs, particularly in cancer patients.

Systemic analgesics such as NSAIDs and opioids are to be distinguished from therapeutic agents which are useful only as local analgesics/anaesthetics. Well known local analgesics such as lidocaine and xylocaine are non-selective ion channel blockers which can be fatal when administered systemically. A good description of non-selective sodium channel blockers is found in Madge, D. et al., *J. Med. Chem.* (2001), 44(2):115-37.

Several sodium channel modulators are known for use as anticonvulsants or antidepressants, such as carbamazepine, amitriptyline, lamotrigine and riluzole, all of which target brain tetradotoxin- sensitive (TTX-S) sodium channels. Such TTX-S agents suffer from dose- limiting side effects, including dizziness, ataxia and somnolence, primarily due to action at TTX-S channels in the brain.

Sodium Channels Role in Pain

Sodium channels play a diverse set of roles in maintaining normal and pathological states, including the long recognized role that voltage gated sodium channels play in the generation of abnormal neuronal activity and neuropathic or pathological pain (Chung, J.M. et al.). Damage to peripheral nerves following trauma or disease can result in changes to sodium channel activity and the development of abnormal afferent activity including ectopic discharges from axotomised afferents and spontaneous activity of sensitized intact nociceptors. These changes can produce long-lasting abnormal hypersensitivity to normally innocuous stimuli, or allodynia. Examples of neuropathic pain include, but are not limited to, post-herpetic neuralgia, trigeminal neuralgia, diabetic neuropathy, chronic lower back pain, phantom limb pain, and pain resulting from cancer and chemotherapy, chronic pelvic pain, complex regional pain syndrome and related neuralgias.

There has been some degree of success in treating neuropathic pain symptoms by using medications, such as gabapentin, and more recently pregabalin, as short-term, first-line treatments. However, pharmacotherapy for neuropathic pain has generally had limited success with little response to commonly used pain reducing drugs, such as NSAIDS and opiates. Consequently, there is still a considerable need to explore novel treatment modalities.

There remains a limited number of potent effective sodium channel blockers with a minimum of adverse events in the clinic. There is also an unmet medical need to treat neuropathic pain and other sodium channel associated pathological states effectively and without adverse side effects. The present invention provides

compounds, methods of use and compositions that include these compounds to meet these critical needs.

SUMMARY OF THE INVENTION

The present invention is directed to spiro-oxindole compounds that are useful for the treatment and/or prevention of sodium channel-mediated diseases or conditions, such as pain. The compounds of the present invention are also useful for the treatment of other sodium channel-mediated diseases or conditions, including, but not limited to central nervous conditions such as epilepsy, anxiety, depression and bipolar disease; cardiovascular conditions such as arrhythmias, atrial fibrillation and ventricular fibrillation; neuromuscular conditions such as restless leg syndrome, essential tremour and muscle paralysis or tetanus; neuroprotection against stroke, glaucoma, neural trauma and multiple sclerosis; and channelopathies such as erythromyalgia and familial rectal pain syndrome.

Accordingly, in one aspect, the invention provides compounds of formula (I):



(I)

wherein:

j and k are each independently 0, 1, 2 or 3;

Q is $-C(R^{1a})H-$, $-C(O)-$, $-O-$, $-S(O)_m-$ (where m is 0, 1 or 2), $-CF_2-$, $-C(O)O-$, $-C(O)N(R^5)-$ or $-N(R^5)C(O)-$;

R^{1a} is hydrogen or $-OR^5$;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

or R^1 is aralkyl substituted by $-C(O)N(R^6)R^7$ where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is hydrogen, alkyl, haloalkyl, $-R^9-CN$, $-R^9-OR^5$, $-R^9-N(R^4)R^5$, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl;

or R^6 and R^7, together with the nitrogen to which they are attached, form a heterocyclyl or heteroaryl;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroaryl groups for R^6 and R^7 may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, $-R^8$-CN, $-R^8$-OR^5, heterocyclyl and heteroaryl;

or R^1 is aralkyl optionally substituted by one or more substituents selected from the group consisting of $-R^8$-OR^5, $-C(O)OR^5$, halo, haloalkyl, alkyl, nitro, cyano, aryl, aralkyl, heterocyclyl and heteroaryl;

or R^1 is $-R^9$-$N(R^{10})R^{11}$, $-R^9$-$N(R^{12})C(O)R^{11}$ or $-R^9$-$N(R^{10})C(O)N(R^{10})R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9$-$OC(O)R^5$, $-R^9$-$C(O)OR^5$, $-R^9$-$C(O)N(R^4)R^5$, $-R^9$-$C(O)R^5$, $-R^9$-$N(R^4)R^5$, $-R^9$-OR^5, or $-R^9$-CN;

R^{12} is hydrogen, alkyl, aryl, arakyl or $-C(O)R^5$;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, $-R^8$-CN, $-R^8$-OR^5, $-R^8$-$C(O)R^5$, heterocyclyl and heteroaryl;

or R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroaryl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8$-OR^5, $-R^8$-$C(O)OR^5$, $-R^8$-$N(R^4)R^5$, $-R^8$-$C(O)N(R^4)R^5$, $-R^8$-$N(R^5)C(O)R^4$, $-R^8$-$S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8$-CN, or $-R^8$-NO_2;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8$-CN, $-R^8$-NO_2, $-R^8$-OR^5, $-R^8$-$N(R^4)R^5$, $-N{=}C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8$-$C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8$-$C(O)OR^4$, $-C(S)OR^4$, $-R^8$-$C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$,

$-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2; and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$; $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain;

as a stereoisomer, enantiomer, tautomer thereof or mixtures thereof;

or a pharmaceutically acceptable salt, solvate or prodrug thereof.

In another aspect, the invention provides methods for the treatment of pain in a mammal, preferably a human, wherein the methods comprise administering to the mammal in need thereof a therapeutically effective amount of a compound of the invention as set forth above.

In another aspect, the present invention provides a method for treating or lessening the severity of a disease, condition, or disorder where activation or hyperactivity of one or more of $Na_V1.1$, $Na_V1.2$, $Na_V1.3$, $Na_V1.4$, $Na_V1.5$, $Na_V1.6$, $Na_V1.7$, $Na_V1.8$, or $Na_V1.9$ is implicated in the disease state.In another aspect, the invention provides methods of treating a range of sodium channel-mediated diseases or conditions, for example, pain associated with HIV, HIV treatment induced neuropathy, trigeminal neuralgia, post-herpetic neuralgia, eudynia, heat sensitivity, tosarcoidosis, irritable bowel syndrome, Crohns disease, pain associated with multiple sclerosis (MS), amyotrophic lateral sclerosis (ALS), diabetic neuropathy, peripheral neuropathy, arthritic, rheumatoid arthritis, osteoarthritis, atherosclerosis, paroxysmal dystonia, myasthenia syndromes, myotonia, malignant hyperthermia, cystic fibrosis, pseudoaldosteronism, rhabdomyolysis, hypothyroidism, bipolar depression, anxiety, schizophrenia, sodium channel toxin related illnesses, familial erythermalgia, primary

erythermalgia, familial rectal pain, cancer, epilepsy, partial and general tonic seizures, restless leg syndrome, arrhythmias, fibromyalgia, neuroprotection under ischaemic conditions caused by stroke, glaucoma or neural trauma, tachy-arrhythmias, atrial fibrillation and ventricular fibrillation.

In another aspect, the invention provides methods of treating a range of sodium channel-mediated disease or condition through inhibition of ion flux through a voltage-dependent sodium channel in a mammal, preferably a human, wherein the methods comprise administering to the mammal in need thereof a therapeutically effective amount of a compound of the invention as set forth above.

In another aspect, the invention provides pharmaceutical compositions comprising the compounds of the invention, as set forth above, and pharmaceutically acceptable excipients. In one embodiment, the present invention relates to a pharmaceutical composition comprising a compound of the invention in a pharmaceutically acceptable carrier and in an amount effective to treat diseases or conditions related to pain when administered to an animal, preferably a mammal, most preferably a human.

In another aspect, the invention provides pharmaceutical therapy in combination with one or more other compounds of the invention or one or more other accepted therapies or as any combination thereof to increase the potency of an existing or future drug therapy or to decrease the adverse events associated with the accepted therapy. In one embodiment, the present invention relates to a pharmaceutical composition combining compounds of the present invention with established or future therapies for the indications listed in the invention.

DETAILED DESCRIPTION OF THE INVENTION

DEFINITIONS

Certain chemical groups named herein are preceded by a shorthand notation indicating the total number of carbon atoms that are to be found in the indicated chemical group. For example; C_7-C_{12}alkyl describes an alkyl group, as defined below, having a total of 7 to 12 carbon atoms, and C_4-C_{12}cycloalkylalkyl describes a cycloalkylalkyl group, as defined below, having a total of 4 to 12 carbon atoms. The total number of carbons in the shorthand notation does not include carbons that may exist in substituents of the group described. For example, the following terms have the meaning indicated:

"C_1-C_{10}alkyl" refers to an alkyl radical as defined below containing one to ten carbon atoms. The C_1-C_{10}alkyl readical may be optionally substituted as defined below for an alkyl group.

"C_2-C_{12}alkynyl" refers to an alknyl radical as defined below containing two to twelve carbon atoms. The C_2-C_{12}alknyl radical may be optionally substituted as defined below for an alkenyl group.

"C_1-C_{12}alkoxy" refers to an alkoxy radical as defined below containing one to twelve carbon atoms. The alkyl part of the C_1-C_{12}alkoxy radical may be optionally substituted as defined below for an alkyl group.

"C_2-C_{12}alkoxyalkyl" refers to an alkoxyalkyl radical as defined below containing two to twelve carbon atoms. Each alkyl part of the C_2-C_{12}alkoxyalkyl radical may be optionally substituted as defined below for an alkyl group.

"C_7-C_{12}aralkyl" refers to an aralkyl group as defined below containing seven to twelve carbon atoms. The aryl part of the C_7-C_{12}aralkyl radical may be optionally substituted as described below for an aryl group. The alkyl part of the C_7-C_{12}aralkyl radical may be optionally substituted as defined below for an alkyl group.

"C_7-C_{12}aralkenyl" refers to an aralkenyl group as defined below containing seven to twelve carbon atoms. The aryl part of the C_7-C_{12}aralkenyl radical may be optionally substituted as described below for an aryl group. The alkenyl part of the C_7-C_{12}aralkenyl radical may be optionally substituted as defined below for an alkenyl group.

"C_3-C_{12}cycloalkyl" refers to a cycloalkyl radical as defined below having three to twelve carbon atoms. The C_3-C_{12}cycloalkyl radical may be optionally substituted as defined below for a cycloalkyl group.

"C_4-C_{12}cycloalkylalkyl" refers to a cycloalkylalkyl radical as defined below having four to twelve carbon atoms. The C_4-C_{12}cycloalkylalkyl radical may be optionally substituted as defined below for a cycloalkylalkyl group.

In addition to the foregoing, as used in the specification and appended claims, unless specified to the contrary, the following terms have the meaning indicated:

"Amino" refers to the $-NH_2$ radical.

"Cyano" refers to the -CN radical.

"Hydroxyl" refers to the -OH radical.

"Imino" refers to the =NH substituent.

"Nitro" refers to the $-NO_2$ radical.

"Oxo" refers to the =O substituent.

"Thioxo" refers to the =S substituent.

"Trifluoromethyl" refers to the $-CF_3$ radical.

"Alkyl" refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, containing no unsaturation, having from one to twelve carbon atoms, preferably one to eight carbon atoms or one to six carbon atoms, and which is attached to the rest of the molecule by a single bond, *e.g.*, methyl, ethyl, *n*-propyl, 1-methylethyl (*iso*-propyl), *n*-butyl, *n*-pentyl, 1,1-dimethylethyl (*t*-butyl), 3-methylhexyl, 2-methylhexyl, and the like. Unless stated otherwise specifically in the specification, an alkyl group may be optionally substituted by one of the following groups: alkyl, alkenyl, halo, haloalkenyl, cyano, nitro, aryl, cycloalkyl, heterocyclyl, heteroaryl, oxo, trimethylsilanyl, $-OR^{14}$, $-OC(O)-R^{14}$, $-N(R^{14})_2$, $-C(O)R^{14}$, $-C(O)OR^{14}$, $-C(O)N(R^{14})_2$, $-N(R^{14})C(O)OR^{16}$, $-N(R^{14})C(O)R^{16}$, $-N(R^{14})S(O)_tR^{16}$ (where t is 1 to 2), $-S(O)_tOR^{16}$ (where t is 1 to 2), $-S(O)_tR^{16}$ (where t is 0 to 2), and $-S(O)_tN(R^{14})_2$ (where t is 1 to 2) where each R^{14} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl (optionally substituted with one or more halo groups), aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; and each R^{16} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, and where each of the above substituents is unsubstituted unless otherwise indicated.

"Alkenyl" refers to a straight or branched hydrocarbon chain radical group consisting solely of carbon and hydrogen atoms, containing at least one double bond, having from two to twelve carbon atoms, preferably one to eight carbon atoms and which is attached to the rest of the molecule by a single bond, *e.g.*, ethenyl, prop-1-enyl, but-1-enyl, pent-1-enyl, penta-1,4-dienyl, and the like. Unless stated otherwise specifically in the specification, an alkenyl group may be optionally substituted by one of the following groups: alkyl, alkenyl, halo, haloalkenyl, cyano, nitro, aryl, cycloalkyl, heterocyclyl, heteroaryl, oxo, trimethylsilanyl, $-OR^{14}$, $-OC(O)-R^{14}$, $-N(R^{14})_2$, $-C(O)R^{14}$, $-C(O)OR^{14}$, $-C(O)N(R^{14})_2$, $-N(R^{14})C(O)OR^{16}$, $-N(R^{14})C(O)R^{16}$, $-N(R^{14})S(O)_tR^{16}$ (where t is 1 to 2), $-S(O)_tOR^{16}$ (where t is 1 to 2), $-S(O)_tR^{16}$ (where t is 0 to 2), and $-S(O)_tN(R^{14})_2$ (where t is 1 to 2) where each R^{14} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl (optionally substituted with one or more halo groups), aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; and each R^{16} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, and where each of the above substituents is unsubstituted unless otherwise indicated.

"Alkylene" or "alkylene chain" refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, containing no unsaturation and having from one to twelve carbon atoms, *e.g.*, methylene, ethylene, propylene, *n*-butylene, and the like. The alkylene chain is attached to the rest of the molecule through a single bond and to the radical group through a single bond. The points of attachment of the alkylene chain to the rest of the molecule and to the radical group can be through one carbon or any two carbons within the chain. Unless stated otherwise specifically in the specification, an alkylene chain may be optionally substituted by one of the following groups: alkyl, alkenyl, halo, haloalkenyl, cyano, nitro, aryl, cycloalkyl, heterocyclyl, heteroaryl, oxo, trimethylsilanyl, $-OR^{14}$, $-OC(O)-R^{14}$, $-N(R^{14})_2$, $-C(O)R^{14}$, $-C(O)OR^{14}$, $-C(O)N(R^{14})_2$, $-N(R^{14})C(O)OR^{16}$, $-N(R^{14})C(O)R^{16}$, $-N(R^{14})S(O)_tR^{16}$ (where t is 1 to 2), $-S(O)_tOR^{16}$ (where t is 1 to 2), $-S(O)_tR^{16}$ (where t is 0 to 2), and $-S(O)_tN(R^{14})_2$ (where t is 1 to 2) where each R^{14} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl (optionally substituted with one or more halo groups), aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; and each R^{16} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, and where each of the above substituents is unsubstituted unless otherwise indicated.

"Alkenylene" or "alkenylene chain" refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, containing at least one double bond and having from two to twelve carbon atoms, *e.g.*, ethenylene, propenylene, *n*-butenylene, and the like. The alkenylene chain is attached to the rest of the molecule through a single bond and to the radical group through a double bond or a single bond. The points of attachment of the alkenylene chain to the rest of the molecule and to the radical group can be through one carbon or any two carbons within the chain. Unless stated otherwise specifically in the specification, an alkenylene chain may be optionally substituted by one of the following groups: alkyl, alkenyl, halo, haloalkenyl, cyano, nitro, aryl, cycloalkyl, heterocyclyl, heteroaryl, oxo, trimethylsilanyl, $-OR^{14}$, $-OC(O)-R^{14}$, $-N(R^{14})_2$, $-C(O)R^{14}$, $-C(O)OR^{14}$, $-C(O)N(R^{14})_2$, $-N(R^{14})C(O)OR^{16}$, $-N(R^{14})C(O)R^{16}$, $-N(R^{14})S(O)_tR^{16}$ (where t is 1 to 2), $-S(O)_tOR^{16}$ (where t is 1 to 2), $-S(O)_tR^{16}$ (where t is 0 to 2), and $-S(O)_tN(R^{14})_2$ (where t is 1 to 2) where each R^{14} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl (optionally substituted with one or more halo groups), aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; and each

R^{16} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, and where each of the above substituents is unsubstituted unless otherwise indicated.

"Alkynylene" or "alkynylene chain" refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, containing at least one triple bond and having from two to twelve carbon atoms, *e.g.*, propynylene, *n*-butynylene, and the like. The alkynylene chain is attached to the rest of the molecule through a single bond and to the radical group through a double bond or a single bond. The points of attachment of the alkynylene chain to the rest of the molecule and to the radical group can be through one carbon or any two carbons within the chain. Unless stated otherwise specifically in the specification, an alkynylene chain may be optionally substituted by one of the following groups: alkyl, alkenyl, halo, haloalkenyl, cyano, nitro, aryl, cycloalkyl, heterocyclyl, heteroaryl, oxo, trimethylsilanyl, $-OR^{14}$, $-OC(O)-R^{14}$, $-N(R^{14})_2$, $-C(O)R^{14}$, $-C(O)OR^{14}$, $-C(O)N(R^{14})_2$, $-N(R^{14})C(O)OR^{16}$, $-N(R^{14})C(O)R^{16}$, $-N(R^{14})S(O)_tR^{16}$ (where t is 1 to 2), $-S(O)_tOR^{16}$ (where t is 1 to 2), $-S(O)_tR^{16}$ (where t is 0 to 2), and $-S(O)_tN(R^{14})_2$ (where t is 1 to 2) where each R^{14} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl (optionally substituted with one or more halo groups), aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; and each R^{16} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, and where each of the above substituents is unsubstituted unless otherwise indicated.

"Alkynyl" refers to a straight or branched hydrocarbon chain radical group consisting solely of carbon and hydrogen atoms, containing at least one triple bond, having from two to twelve carbon atoms, preferably one to eight carbon atoms and which is attached to the rest of the molecule by a single bond, e.g., ethynyl, propynyl, butynyl, pentynyl, hexynyl, and the like. Unless stated otherwise specifically in the specification, an alkynyl group may be optionally substituted by one of the following groups: alkyl, alkenyl, halo, haloalkyl, haloalkenyl, cyano, nitro, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-OR^{14}$, $-OC(O)-R^{14}$, $-N(R^{14})_2$, $-C(O)R^{14}$, $-C(O)OR^{14}$, $-C(O)N(R^{14})_2$, $-N(R^{14})C(O)OR^{16}$, $-N(R^{14})C(O)R^{16}$, $-N(R^{14})S(O)_tR^{16}$ (where t is 1 to 2), $-S(O)_tOR^{16}$ (where t is 1 to 2), $-S(O)_tR^{16}$ (where t is 0 to 2), and $-S(O)_tN(R^{14})_2$ (where t is 1 to 2) where each R^{14} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; and each R^{16} is alkyl,

haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, and where each of the above substituents is unsubstituted.

"Alkoxy" refers to a radical of the formula $-OR_a$ where R_a is an alkyl radical as defined above containing one to twelve carbon atoms. The alkyl part of the alkoxy radical may be optionally substituted as defined above for an alkyl radical.

"Alkoxyalkyl" refers to a radical of the formula $-R_a-O-R_a$ where each R_a is independently an alkyl radical as defined above. The oxygen atom may be bonded to any carbon in either alkyl radical. Each alkyl part of the alkoxyalkyl radical may be optionally substituted as defined above for an alkyl group.

"Aryl" refers to aromatic monocyclic or multicyclic hydrocarbon ring system consisting only of hydrogen and carbon and containing from 6 to 18 carbon atoms, where the ring system may be partially saturated. Aryl groups include, but are not limited to groups such as fluorenyl, phenyl and naphthyl. Unless stated otherwise specifically in the specification, the term "aryl" or the prefix "ar-" (such as in "aralkyl") is meant to include aryl radicals optionally substituted by one or more substituents independently selected from the group consisting of alkyl, akenyl, halo, haloalkyl, haloalkenyl, cyano, nitro, aryl, heteroaryl, heteroarylalkyl, $-R^{15}-OR^{14}$, $-R^{15}-OC(O)-R^{14}$, $-R^{15}-N(R^{14})_2$, $-R^{15}-C(O)R^{14}$, $-R^{15}-C(O)OR^{14}$, $-R^{15}-C(O)N(R^{14})_2$, $-R^{15}-N(R^{14})C(O)OR^{16}$, $-R^{15}-N(R^{14})C(O)R^{16}$, $-R^{15}-N(R^{14})S(O)_tR^{16}$ (where t is 1 to 2), $-R^{15}-S(O)_tOR^{16}$ (where t is 1 to 2), $-R^{15}-S(O)_tR^{16}$ (where t is 0 to 2), and $-R^{15}-S(O)_tN(R^{14})_2$ (where t is 1 to 2) where each R^{14} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; each R^{15} is independently a direct bond or a straight or branched alkylene or alkenylene chain; and each R^{16} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, and where each of the above substituents is unsubstituted.

"Aralkyl" refers to a radical of the formula $-R_aR_b$ where R_a is an alkyl radical as defined above and R_b is one or more aryl radicals as defined above, *e.g.*, benzyl, diphenylmethyl and the like. The aryl radical(s) may be optionally substituted as described above.

"Aryloxy" refers to a radical of the formula $-OR_b$ where R_b is an aryl group as defined above. The aryl part of the aryloxy radical may be optionally substituted as defined above.

"Aralkenyl" refers to a radical of the formula $-R_cR_b$ where R_c is an alkenyl radical

as defined above and R_b is one or more aryl radicals as defined above, which may be optionally substituted as described above. The aryl part of the aralkenyl radical may be optionally substituted as described above for an aryl group. The alkenyl part of the aralkenyl radical may be optionally substituted as defined above for an alkenyl group.

"Aralkyloxy" refers to a radical of the formula $-OR_b$ where R_b is an aralkyl group as defined above. The aralkyl part of the aralkyloxy radical may be optionally substituted as defined above.

"Cycloalkyl" refers to a stable non-aromatic monocyclic or polycyclic hydrocarbon radical consisting solely of carbon and hydrogen atoms, which may include fused or bridged ring systems, having from three to fifteen carbon atoms, preferably having from three to ten carbon atoms, and which is saturated or unsaturated and attached to the rest of the molecule by a single bond. Monocyclic radicals include, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptly, and cyclooctyl. Polycyclic radicals include, for example, adamantyl, norbornyl, decalinyl, 7,7-dimethyl-bicyclo[2.2.1]heptanyl, and the like. Unless otherwise stated specifically in the specification, the term "cycloalkyl" is meant to include cycloalkyl radicals which are optionally substituted by one or more substituents independently selected from the group consisting of alkyl, alkenyl, halo, haloalkyl, haloalkenyl, cyano, nitro, oxo, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^{15}-OR^{14}$, $-R^{15}-OC(O)-R^{14}$, $-R^{15}-N(R^{14})_2$, $-R^{15}-C(O)R^{14}$, $-R^{15}-C(O)OR^{14}$, $-R^{15}-C(O)N(R^{14})_2$, $-R^{15}-N(R^{14})C(O)OR^{16}$, $-R^{15}-N(R^{14})C(O)R^{16}$, $-R^{15}-N(R^{14})S(O)_tR^{16}$ (where t is 1 to 2), $-R^{15}-S(O)_tOR^{16}$ (where t is 1 to 2), $-R^{15}-S(O)_tR^{16}$ (where t is 0 to 2), and $-R^{15}-S(O)_tN(R^{14})_2$ (where t is 1 to 2) where each R^{14} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; each R^{15} is independently a direct bond or a straight or branched alkylene or alkenylene chain; and each R^{16} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, and where each of the above substituents is unsubstituted.

"Cycloalkylalkyl" refers to a radical of the formula $-R_aR_d$ where R_a is an alkyl radical as defined above and R_d is a cycloalkyl radical as defined above. The alkyl radical and the cycloalkyl radical may be optionally substituted as defined above.

"Halo" refers to bromo, chloro, fluoro or iodo.

"Haloalkyl" refers to an alkyl radical, as defined above, that is substituted by one or more halo radicals, as defined above, *e.g.*, trifluoromethyl, difluoromethyl,

trichloromethyl, 2,2,2-trifluoroethyl, 1-fluoromethyl-2-fluoroethyl, 3-bromo-2-fluoropropyl, 1-bromomethyl-2-bromoethyl, and the like. The alkyl part of the haloalkyl radical may be optionally substituted as defined above for an alkyl group.

"Heterocyclyl" refers to a stable 3- to 18-membered non-aromatic ring radical which consists of two to twelve carbon atoms and from one to six heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. Unless stated otherwise specifically in the specification, the heterocyclyl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heterocyclyl radical may be optionally oxidized; the nitrogen atom may be optionally quaternized; and the heterocyclyl radical may be partially or fully saturated. Examples of such heterocyclyl radicals include, but are not limited to, dioxolanyl, thienyl[1,3]dithianyl, decahydroisoquinolyl, imidazolinyl, imidazolidinyl, isothiazolidinyl, isoxazolidinyl, morpholinyl, octahydroindolyl, octahydroisoindolyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, oxazolidinyl, piperidinyl, piperazinyl, 4-piperidonyl, pyrrolidinyl, pyrazolidinyl, thiazolidinyl, tetrahydrofuryl, trithianyl, tetrahydropyranyl, thiomorpholinyl, thiamorpholinyl, 1-oxo-thiomorpholinyl, and 1,1-dioxo-thiomorpholinyl. Unless stated otherwise specifically in the specification, the term "heterocyclyl" is meant to include heterocyclyl radicals as defined above which are optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, halo, haloalkyl, haloalkenyl, cyano, oxo, thioxo, nitro, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^{15}-OR^{14}$, $-R^{15}-OC(O)-R^{14}$, $-R^{15}-N(R^{14})_2$, $-R^{15}-C(O)R^{14}$, $-R^{15}-C(O)OR^{14}$, $-R^{15}-C(O)N(R^{14})_2$, $-R^{15}-N(R^{14})C(O)OR^{16}$, $-R^{15}-N(R^{14})C(O)R^{16}$, $-R^{15}-N(R^{14})S(O)_tR^{16}$ (where t is 1 to 2), $-R^{15}-S(O)_tOR^{16}$ (where t is 1 to 2), $-R^{15}-S(O)_tR^{16}$ (where t is 0 to 2), and $-R^{15}-S(O)_tN(R^{14})_2$ (where t is 1 to 2) where each R^{14} is independently hydrogen, alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; each R^{15} is independently a direct bond or a straight or branched alkylene or alkenylene chain; and each R^{16} is alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, and where each of the above substituents is unsubstituted.

"Heterocyclylalkyl" refers to a radical of the formula $-R_aR_e$ where R_a is an alkyl radical as defined above and R_e is a heterocyclyl radical as defined above, and if the heterocyclyl is a nitrogen-containing heterocyclyl, the heterocyclyl may be attached to the alkyl radical at the nitrogen atom. The alkyl part of the heterocyclylalkyl radical

may be optionally substituted as defined above for an alkyl group. The heterocyclyl part of the heterocyclylalkyl radical may be optionally substituted as defined above for a heterocyclyl group.

"Heteroaryl" refers to a 5- to 18-membered aromatic ring radical which consists of one to seventeen carbon atoms and from one to ten heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. For purposes of this invention, the heteroaryl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heteroaryl radical may be optionally oxidized; the nitrogen atom may be optionally quaternized. Examples include, but are not limited to, azepinyl, acridinyl, benzimidazolyl, benzthiazolyl, benzindolyl, benzodioxolyl, benzofuranyl, benzooxazolyl, benzothiazolyl, benzothiadiazolyl, benzo[*b*][1,4]dioxepinyl, 1,4-benzodioxanyl, benzonaphthofuranyl, benzoxazolyl, benzodioxolyl, benzodioxinyl, benzopyranyl, benzopyranonyl, benzofuranyl, benzofuranonyl, benzothienyl (benzothiophenyl), benzotriazolyl, benzo[4,6]imidazo[1,2-a]pyridinyl, carbazolyl, cinnolinyl, dibenzofuranyl, dibenzothiophenyl, furanyl, furanonyl, isothiazolyl, imidazolyl, indazolyl, indolyl, indazolyl, isoindolyl, indolinyl, isoindolinyl, isoquinolyl, indolizinyl, isoxazolyl, naphthyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, oxiranyl, 1-phenyl-1*H*-pyrrolyl, phenazinyl, phenothiazinyl, phenoxazinyl, phthalazinyl, pteridinyl, purinyl, pyrrolyl, pyrazolyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, quinazolinyl, quinoxalinyl, quinolinyl, quinuclidinyl, isoquinolinyl, tetrahydroquinolinyl, thiazolyl, thiadiazolyl, triazolyl, tetrazolyl, triazinyl, and thiophenyl (i.e. thienyl). Unless stated otherwise specifically in the specification, the term "heteroaryl" is meant to include heteroaryl radicals as defined above which are optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkoxy, halo, haloalkyl, haloalkenyl, cyano, oxo, thioxo, nitro, oxo, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^{15}-OR^{14}$, $-R^{15}-OC(O)-R^{14}$, $-R^{15}-N(R^{14})_2$, $-R^{15}-C(O)R^{14}$, $-R^{15}-C(O)OR^{14}$, $-R^{15}-C(O)N(R^{14})_2$, $-R^{15}-N(R^{14})C(O)OR^{16}$, $-R^{15}-N(R^{14})C(O)R^{16}$, $-R^{15}-N(R^{14})S(O)_tR^{16}$ (where t is 1 to 2), $-R^{15}-S(O)_tOR^{16}$ (where t is 1 to 2), $-R^{15}-S(O)_tR^{16}$ (where t is 0 to 2), and $-R^{15}-S(O)_tN(R^{14})_2$ (where t is 1 to 2) where each R^{14} is independently hydrogen, alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; each R^{15} is independently a direct bond or a straight or branched alkylene or alkenylene chain; and each R^{16} is alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, and where each of the

above substituents is unsubstituted.

"Heteroarylalkyl" refers to a radical of the formula $-R_aR_f$ where R_a is an alkyl radical as defined above and R_f is a heteroaryl radical as defined above. The heteroaryl part of the heteroarylalkyl radical may be optionally substituted as defined above for a heteroaryl group. The alkyl part of the heteroarylalkyl radical may be optionally substituted as defined above for an alkyl group.

"Heteroarylalkenyl" refers to a radical of the formula $-R_bR_f$ where R_b is an alkenyl radical as defined above and R_f is a heteroaryl radical as defined above. The heteroaryl part of the heteroarylalkenyl radical may be optionally substituted as defined above for a heteroaryl group. The alkenyl part of the heteroarylalkenyl radical may be optionally substituted as defined above for an alkenyl group.

"Trihaloalkyl" refers to an alkyl radical, as defined above, that is substituted by three halo radicals, as defined above, *e.g.*, trifluoromethyl. The alkyl part of the trihaloalkyl radical may be optionally substituted as defined above for an alkyl group.

"Trihaloalkoxy" refers to a radical of the formula $-OR_g$ where R_g is a trihaloalkyl group as defined above. The trihaloalkyl part of the trihaloalkoxy group may be optionally substituted as defined above for a trihaloalkyl group.

"Analgesia" refers to an absence of pain in response to a stimulus that would normally be painful.

"Allodynia" refers to a condition in which a normally innocuous sensation, such as pressure or light touch, is perceived as being extremely painful.

"Prodrugs" is meant to indicate a compound that may be converted under physiological conditions or by solvolysis to a biologically active compound of the invention. Thus, the term "prodrug" refers to a metabolic precursor of a compound of the invention that is pharmaceutically acceptable. A prodrug may be inactive when administered to a subject in need thereof, but is converted *in vivo* to an active compound of the invention. Prodrugs are typically rapidly transformed *in vivo* to yield the parent compound of the invention, for example, by hydrolysis in blood. The prodrug compound often offers advantages of solubility, tissue compatibility or delayed release in a mammalian organism (see, Bundgard, H., Design of Prodrugs (1985), pp. 7-9, 21-24 (Elsevier, Amsterdam)).A discussion of prodrugs is provided in Higuchi, T., *et al.*, "Pro-drugs as Novel Delivery Systems," A.C.S. Symposium Series, Vol. 14, and in Bioreversible Carriers in Drug Design, Ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

The term "prodrug" is also meant to include any covalently bonded carriers, which release the active compound of the invention *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound of the invention may be prepared by modifying functional groups present in the compound of the invention in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound of the invention. Prodrugs include compounds of the invention wherein a hydroxy, amino or mercapto group is bonded to any group that, when the prodrug of the compound of the invention is administered to a mammalian subject, cleaves to form a free hydroxy, free amino or free mercapto group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol or amide derivatives of amine functional groups in the compounds of the invention and the like.

The invention disclosed herein is also meant to encompass all pharmaceutically acceptable compounds of formula (I) being isotopically-labelled by having one or more atoms replaced by an atom having a different atomic mass or mass number. Examples of isotopes that can be incorporated into the disclosed compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine, chlorine, and iodine, such as ^{2}H, ^{3}H, ^{11}C, ^{13}C, ^{14}C, ^{13}N, ^{15}N, ^{15}O, ^{17}O, ^{18}O, ^{31}P, ^{32}P, ^{35}S, ^{18}F, ^{36}Cl, ^{123}I, and ^{125}I, respectively. These radiolabelled compounds could be useful to help determine or measure the effectiveness of the compounds, by characterizing, for example, the site or mode of action on the sodium channels, or binding affinity to pharmacologically important site of action on the sodium channels. Certain isotopically-labelled compounds of formula (I), for example, those incorporating a radioactive isotope, are useful in drug and/or substrate tissue distribution studies. The radioactive isotopes tritium, *i.e.* ^{3}H, and carbon-14, *i.e.* ^{14}C, are particularly useful for this purpose in view of their ease of incorporation and ready means of detection.

Substitution with heavier isotopes such as deuterium, *i.e.* ^{2}H, may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life or reduced dosage requirements, and hence may be preferred in some circumstances.

Substitution with positron emitting isotopes, such as ^{11}C, ^{18}F, ^{15}O and ^{13}N, can be useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy. Isotopically-labeled compounds of formula (I) can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the Examples and Preparations as set out below using

an appropriate isotopically-labeled reagent in place of the non-labeled reagent previously employed.

The invention disclosed herein is also meant to encompass the *in vivo* metabolic products of the disclosed compounds. Such products may result from, for example, the oxidation, reducation, hydrolysis, amidation, esterification, and the like of the administered compound, primarily due to enzymatic processes. Accordingly, the invention includes compounds produced by a process comprising contacting a compound of this invention with a mammal for a period of time sufficient to yield a metabolic product thereof. Such products are typically are identified by administering a radiolabelled compound of the invention in a detectable dose to an animal, such as rat, mouse, guinea pig, monkey, or to human, allowing sufficient time for metabolism to occur, and isolating its coversion products from the urine, blood or other biological samples.

"Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

"Mammal" includes humans and both domestic animals such as laboratory animals and household pets, (e.g. cats, dogs, swine, cattle, sheep, goats, horses, rabbits), and non-domestic animals such as wildelife and the like.

"Optional" or "optionally" means that the subsequently described event of circumstances may or may not occur, and that the description includes instances where said event or circumstance occurs and instances in which it does not. For example, "optionally substituted aryl" means that the aryl radical may or may not be substituted and that the description includes both substituted aryl radicals and aryl radicals having no substitution.

"Pharmaceutically acceptable carrier, diluent or excipient" includes without limitation any adjuvant, carrier, excipient, glidant, sweetening agent, diluent, preservative, dye/colorant, flavor enhancer, surfactant, wetting agent, dispersing agent, suspending agent, stabilizer, isotonic agent, solvent, or emulsifier which has been approved by the United States Food and Drug Administration as being acceptable for use in humans or domestic animals.

"Pharmaceutically acceptable salt" includes both acid and base addition salts.

"Pharmaceutically acceptable acid addition salt" refers to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which are formed with inorganic acids such

as, but are not limited to, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as, but not limited to, acetic acid, 2,2-dichloroacetic acid, adipic acid, alginic acid, ascorbic acid, aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, camphoric acid, camphor-10-sulfonic acid, capric acid, caproic acid, caprylic acid, carbonic acid, cinnamic acid, citric acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glucoheptonic acid, gluconic acid, glucuronic acid, glutamic acid, glutaric acid, 2-oxo-glutaric acid, glycerophosphoric acid, glycolic acid, hippuric acid, isobutyric acid, lactic acid, lactobionic acid, lauric acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, mucic acid, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, propionic acid, pyroglutamic acid, pyruvic acid, salicylic acid, 4-aminosalicylic acid, sebacic acid, stearic acid, succinic acid, tartaric acid, thiocyanic acid, *p*-toluenesulfonic acid, trifluoroacetic acid, undecylenic acid, and the like.

"Pharmaceutically acceptable base addition salt" refers to those salts which retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise undesirable. These salts are prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Preferred inorganic salts are the ammonium, sodium, potassium, calcium, and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as ammonia, isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, diethanolamine, ethanolamine, deanol, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, benethamine, benzathine, ethylenediamine, glucosamine, methylglucamine, theobromine, triethanolamine, tromethamine, purines, piperazine, piperidine, *N*-ethylpiperidine, polyamine resins and the like. Particularly preferred organic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline and caffeine.

Often crystallizations produce a solvate of the compound of the invention. As

used herein, the term "solvate" refers to an aggregate that comprises one or more molecules of a compound of the invention with one or more molecules of solvent. The solvent may be water, in which case the solvate may be a hydrate. Alternatively, the solvent may be an organic solvent. Thus, the compounds of the present invention may exist as a hydrate, including a monohydrate, dihydrate, hemihydrate, sesquihydrate, trihydrate, tetrahydrate and the like, as well as the corresponding solvated forms. The compound of the invention may be true solvates, while in other cases, the compound of the invention may merely retain adventitious water or be a mixture of water plus some adventitious solvent.

A "pharmaceutical composition" refers to a formulation of a compound of the invention and a medium generally accepted in the art for the delivery of the biologically active compound to mammals, e.g., humans. Such a medium includes all pharmaceutically acceptable carriers, diluents or excipients therefor.

"Therapeutically effective amount" refers to that amount of a compound of the invention which, when administered to a mammal, preferably a human, is sufficient to effect treatment, as defined below, of a sodium channel-mediated disease or condition in the mammal, preferably a human. The amount of a compound of the invention which constitutes a "therapeutically effective amount" will vary depending on the compound, the condition and its severity, the manner of administration, and the age of the mammal to be treated, but can be determined routinely by one of ordinary skill in the art having regard to his own knowledge and to this disclosure.

"Treating" or "treatment" as used herein covers the treatment of the disease or condition of interest in a mammal, preferably a human, having the disease or condition of interest, and includes:

(i) preventing the disease or condition from occurring in a mammal, in particular, when such mammal is predisposed to the condition but has not yet been diagnosed as having it;

(ii) inhibiting the disease or condition, *i.e.*, arresting its development;

(iii) relieving the disease or condition, *i.e.*, causing regression of the disease or condition; or

(iv) relieving the symptoms resulting from the disease or condition, *i.e.*, relieving pain without addressing the underlying disease or condition.As used herein, the terms "disease" and "condition" may be used interchangeably or may be different in that the particular malady or condition may not have a known causative agent (so that etiology has not yet been worked out) and it is therefore not yet recognized as a

disease but only as an undesirable condition or syndrome, wherein a more or less specific set of symptoms have been identified by clinicians.

The compounds of the invention, or their pharmaceutically acceptable salts may contain one or more asymmetric centres and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (*R*)- or (*S*)- or, as (D)- or (L)- for amino acids. The present invention is meant to include all such possible isomers, as well as their racemic and optically pure forms. Optically active (+) and (-), (*R*)- and (*S*)-, or (D)- and (L)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques, for example, chromatography and fractional crystallisation. Conventional techniques for the preparation/isolation of individual enantiomers include chiral synthesis from a suitable optically pure precursor or resolution of the racemate (or the racemate of a salt or derivative) using, for example, chiral high pressure liquid chromatography (HPLC). When the compounds described herein contain olefinic double bonds or other centres of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both E and Z geometric isomers. Likewise, all tautomeric forms are also intended to be included.

A "stereoisomer" refers to a compound made up of the same atoms bonded by the same bonds but having different three-dimensional structures, which are not interchangeable. The present invention contemplates various stereoisomers and mixtures thereof and includes "enantiomers", which refers to two stereoisomers whose molecules are nonsuperimposeable mirror images of one another.

A "tautomer" refers to a proton shift from one atom of a molecule to another atom of the same molecule. The present invention includes tautomers of any said compounds.

Also within the scope of the invention are intermediate compounds of formula (I) and all polymorphs of the aforementioned species and crystal habits thereof.

The chemical naming protocol and structure diagrams used herein are a modified form of the I.U.P.A.C. nomenclature system, using the ACD/Name Version 9.07 software program, wherein the compounds of the invention are named herein as derivatives of the central core structure, *i.e.*, the 2-oxindole structure. For complex chemical names employed herein, a substituent group is named before the group to which it attaches. For example, cyclopropylethyl comprises an ethyl backbone with cyclopropyl substituent. In chemical structure diagrams, all bonds are identified, except for some carbon atoms, which are assumed to be bonded to sufficient hydrogen

atoms to complete the valency.

Thus, for example, a compound of formula (I) wherein j is 0, k is 1, Q is -O-, R^1 is pentyl, R^{2a} is 3,5-dichlorophenyl, R^{2b}, R^{2c} and R^{2d} are each hydrogen, R^{3a} and R^{3d} are each hydrogen, and R^{3b} and R^{3c}, together with the carbon ring atoms to which they are attached, form a fused dioxolyl ring; *i.e.*, a compound of the following formula:



is named herein as 4'-(3,5-dichlorophenyl)-1'-pentylspiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one.

EMBODIMENTS OF THE INVENTION

Of the various aspects of the invention set forth above in the Summary of the Invention, certain embodiments are preferred.

One embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$,

$-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2; and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$; $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached,

may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen,halo or alkyl;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached,

form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3c} are as defined above;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkylalkyl, $-R^8-OR^5$, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen,halo or alkyl;

R^{3a} and R^{3d} are both hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the Invention wherein:

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkylalkyl, $-R^8-OR^5$, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^8-CN$,

$-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen,halo or alkyl;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the Invention wherein:

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkylalkyl, $-R^8-OR^5$, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen,halo or alkyl;

R^{3a} and R^{3d} are both hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused optionally substituted heterocyclyl ring or a fused optionally substituted cycloalkyl ring;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the Invention wherein:

R^1 is aryl, heteroaryl or heterocyclyl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen; and

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is hydrogen, alkyl, $-R^8-C(O)OR^5$ or $-R^8-C(O)N(R^4)R^5$;

R^{2a} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a} is optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$,

-R^8-C(O)R^4; -R^8-C(O)OR^4, -R^8-C(O)N(R^4)R^5, -N(R^5)C(O)R^4, and -N(R^5)S(O)$_n$$R^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from hydrogen or halo;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl and R^{3a} and R^{3d} are as defined above;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1 or j is 1 and k is 0;

Q is -O-;

R^1 is hydrogen or alkyl;

R^{2a} is selected from the group consisting of alkyl, haloalkenyl, aryl, aralkyl, aralkenyl, heterocyclyl, heteroaryl, -R^8-C(O)N(R^4)R^5, and -R^8-N(R^4)R^5;

wherein each of the aryl, aralkyl, aralkenyl, heterocyclyl and heteroaryl groups for R^{2a} is optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-N(R^4)R^5, -S(O)$_m$$R^4$, -$R^8$-S(O)$_n$N($R^4$)$R^5$, -$R^8$-C(O)$R^4$; -$R^8$-C(O)O$R^4$, -$R^8$-C(O)N($R^4$)$R^5$, -N($R^5$)C(O)$R^4$, and -N($R^5$)S(O)$_n$$R^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from hydrogen or halo;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring or a fused optionally substituted tetrahydrofuranyl ring, and R^{3a} and R^{3d} are as defined above;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is aralkyl substituted by $-C(O)N(R^6)R^7$ where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is hydrogen, alkyl, haloalkyl, $-R^9-CN$, $-R^9-OR^5$, $-R^9-N(R^4)R^5$, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl;

or R^6 and R^7, together with the nitrogen to which they are attached, form a heterocyclyl or heteroaryl;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroaryl groups for R^6 and R^7 may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, $-R^8-CN$, $-R^8-OR^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$,

$-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2; and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$; $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is aralkyl substituted by $-C(O)N(R^6)R^7$ where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is hydrogen, alkyl, haloalkyl, $-R^9-CN$, $-R^9-OR^5$, $-R^9-N(R^4)R^5$, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl;

or R^6 and R^7, together with the nitrogen to which they are attached, form a heterocyclyl or heteroaryl;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroaryl groups for R^6 and R^7 may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, $-R^8-CN$, $-R^8-OR^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached,

form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is aralkyl substituted by $-C(O)N(R^6)R^7$ where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is hydrogen, alkyl, haloalkyl, $-R^9-CN$, $-R^9-OR^5$ or $-R^9-N(R^4)R^5$;

or R^6 and R^7, together with the nitrogen to which they are attached, form a heterocyclyl or heteroaryl;

wherein each aryl, aralkyl, heterocyclyl and heteroaryl groups for R^6 and R^7 is optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, $-R^8-CN$, $-R^8-OR^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or

branched alkenylene chain or a straight or branched alkynylene chain; and each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is aralkyl substituted by $-C(O)N(R^6)R^7$ where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroaryl groups for R^6 and R^7 may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, $-R^8-CN$, $-R^8-OR^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is aralkyl optionally substituted by one or more substituents selected from the group

consisting of -R^8-OR^5, -C(O)OR^5, halo, haloalkyl, alkyl, nitro, cyano, aryl, aralkyl, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$N{=}C(R^4)R^5$, -$S(O)_mR^4$, -$OS(O)_2CF_3$, -R^8-C(O)R^4; -C(S)R^4, -$C(R^4)_2C(O)R^5$, -R^8-C(O)OR^4, -C(S)OR^4, -R^8-C(O)$N(R^4)R^5$, -C(S)$N(R^4)R^5$, -$N(R^5)C(O)R^4$, -$N(R^5)C(S)R^4$, -$N(R^5)C(O)OR^4$, -$N(R^5)C(S)OR^4$, -$N(R^5)C(O)N(R^4)R^5$, -$N(R^5)C(S)N(R^4)R^5$, -$N(R^5)S(O)_nR^4$, -$N(R^5)S(O)_nN(R^4)R^5$, -R^8-$S(O)_nN(R^4)R^5$,-$N(R^5)C(=NR^5)N(R^4)R^5$, and -$N(R^5)C(=N$-CN$)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2; and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$S(O)_mR^4$, -R^8-$S(O)_nN(R^4)R^5$, -R^8-C(O)R^4; -R^8-C(O)OR^4, -R^8-C(O)$N(R^4)R^5$, -$N(R^5)C(O)R^4$, and -$N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5,

$-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$,

wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is aralkyl optionally substituted by one or more substituents selected from the group consisting of $-R^8-OR^5$, $-C(O)OR^5$, halo, haloalkyl, alkyl, nitro, cyano, aryl, aralkyl, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen, alkyl or halo;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl,

haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$N{=}C(R^4)R^5$, -$S(O)_mR^4$, -$OS(O)_2CF_3$, -R^8-$C(O)R^4$; -$C(S)R^4$, -$C(R^4)_2C(O)R^5$, -R^8-$C(O)OR^4$, -$C(S)OR^4$, -R^8-$C(O)N(R^4)R^5$, -$C(S)N(R^4)R^5$, -$N(R^5)C(O)R^4$, -$N(R^5)C(S)R^4$, -$N(R^5)C(O)OR^4$, -$N(R^5)C(S)OR^4$, -$N(R^5)C(O)N(R^4)R^5$, -$N(R^5)C(S)N(R^4)R^5$, -$N(R^5)S(O)_nR^4$, -$N(R^5)S(O)_nN(R^4)R^5$, -R^8-$S(O)_nN(R^4)R^5$,-$N(R^5)C(=NR^5)N(R^4)R^5$, and -$N(R^5)C(N{=}C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is aralkyl optionally substituted by one or more substituents selected from the group consisting of -R^8-OR^5, -$C(O)OR^5$, halo, haloalkyl, alkyl, nitro, cyano, aryl, aralkyl, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen, alkyl or halo;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, halo, and -R^8-OR^5

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, dioxolyl, tetrahydrofuranyl, and heteroaryl, and R^{3a} and R^d are each hydrogen;

each R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is $-R^9-N(R^{10})R^{11}$, $-R^9-N(R^{12})C(O)R^{11}$ or $-R^9-N(R^{10})C(O)N(R^{10})R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9-OC(O)R^5$, $-R^9-C(O)OR^5$, $-R^9-C(O)N(R^4)R^5$, $-R^9-C(O)R^5$, $-R^9-N(R^4)R^5$, $-R^9-OR^5$, or $-R^9-CN$;

R^{12} is hydrogen, alkyl, aryl, arakyl or $-C(O)R^5$;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, $-R^8-CN$, $-R^8-OR^5$, $-R^8-C(O)R^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$,

wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl,

alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$; $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl,

alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is $-R^9-N(R^{10})R^{11}$, $-R^9-N(R^{12})C(O)R^{11}$ or $-R^9-N(R^{10})C(O)N(R^{10})R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9-OC(O)R^5$, $-R^9-C(O)OR^5$, $-R^9-C(O)N(R^4)R^5$, $-R^9-C(O)R^5$, $-R^9-N(R^4)R^5$, $-R^9-OR^5$, or $-R^9-CN$;

R^{12} is hydrogen, alkyl, aryl, arakyl or $-C(O)R^5$;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, $-R^8-CN$, $-R^8-OR^5$, $-R^8-C(O)R^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is $-R^9-N(R^{10})R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9-OC(O)R^5$, $-R^9-C(O)OR^5$, $-R^9-C(O)N(R^4)R^5$, $-R^9-C(O)R^5$, $-R^9-N(R^4)R^5$, $-R^9-OR^5$, or $-R^9-CN$;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, $-R^8-CN$, $-R^8-OR^5$, $-R^8-C(O)R^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is $-R^9-N(R^{12})C(O)R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9-OC(O)R^5$, $-R^9-C(O)OR^5$, $-R^9-C(O)N(R^4)R^5$, $-R^9-C(O)R^5$, $-R^9-N(R^4)R^5$, $-R^9-OR^5$, or $-R^9-CN$;

R^{12} is hydrogen, alkyl, aryl, arakyl or $-C(O)R^5$;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, $-R^8-CN$, $-R^8-OR^5$, $-R^8-C(O)R^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is $-R^9-N(R^{10})C(O)N(R^{10})R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl,

heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^9-OC(O)R^5, -R^9-C(O)OR^5, -R^9-C(O)N(R^4)R^5, -R^9-C(O)R^5, -R^9-N(R^4)R^5, -R^9-OR^5, or -R^9-CN;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, -R^8-CN, -R^8-OR^5, -R^8-C(O)R^5, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroaryl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-OR^5, -R^8-C(O)OR^5, -R^8-N(R^4)R^5, -R^8-C(O)N(R^4)R^5, -R^8-N(R^5)C(O)R^4, -R^8-S(O)$_m$$R^4$ (where m is 0, 1 or 2), -R^8-CN, or -R^8-NO_2;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl,

heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$N{=}C(R^4)R^5$, -$S(O)_mR^4$, -$OS(O)_2CF_3$, -R^8-$C(O)R^4$; -$C(S)R^4$, -$C(R^4)_2C(O)R^5$, -R^8-$C(O)OR^4$, -$C(S)OR^4$, -R^8-$C(O)N(R^4)R^5$, -$C(S)N(R^4)R^5$, -$N(R^5)C(O)R^4$, -$N(R^5)C(S)R^4$, -$N(R^5)C(O)OR^4$, -$N(R^5)C(S)OR^4$, -$N(R^5)C(O)N(R^4)R^5$, -$N(R^5)C(S)N(R^4)R^5$, -$N(R^5)S(O)_nR^4$, -$N(R^5)S(O)_nN(R^4)R^5$, -R^8-$S(O)_nN(R^4)R^5$, -$N(R^5)C(=NR^5)N(R^4)R^5$, and -$N(R^5)C(=N\text{-}CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2; and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$S(O)_mR^4$, -R^8-$S(O)_nN(R^4)R^5$, -R^8-$C(O)R^4$; -R^8-$C(O)OR^4$, -R^8-$C(O)N(R^4)R^5$, -$N(R^5)C(O)R^4$, and -$N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$N{=}C(R^4)R^5$, -$S(O)_mR^4$, -$OS(O)_2CF_3$, -R^8-$C(O)R^4$; -$C(S)R^4$, -$C(R^4)_2C(O)R^5$, -R^8-$C(O)OR^4$, -$C(S)OR^4$, -R^8-$C(O)N(R^4)R^5$, -$C(S)N(R^4)R^5$, -$N(R^5)C(O)R^4$, -$N(R^5)C(S)R^4$, -$N(R^5)C(O)OR^4$, -$N(R^5)C(S)OR^4$, -$N(R^5)C(O)N(R^4)R^5$, -$N(R^5)C(S)N(R^4)R^5$, -$N(R^5)S(O)_nR^4$, -$N(R^5)S(O)_nN(R^4)R^5$, -R^8-$S(O)_nN(R^4)R^5$,-$N(R^5)C(=NR^5)N(R^4)R^5$, and -$N(R^5)C(N{=}C(R^4)R^5)N(R^4)R^5$,

wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroaryl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-OR^5$, $-R^8-C(O)OR^5$, $-R^8-N(R^4)R^5$, $-R^8-C(O)N(R^4)R^5$, $-R^8-N(R^5)C(O)R^4$, $-R^8-S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8-CN$, or $-R^8-NO_2$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen, halo, alkyl or $-R^8-OR^5$;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from hydrogen, halo, alkyl or $-R^8-OR^5$;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached,

form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are each hydrogen;

each R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroaryl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-OR^5$, $-R^8-C(O)OR^5$, $-R^8-N(R^4)R^5$, $-R^8-C(O)N(R^4)R^5$, $-R^8-N(R^5)C(O)R^4$, $-R^8-S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8-CN$, or $-R^8-NO_2$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen, halo, alkyl or $-R^8-OR^5$;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from hydrogen, halo, alkyl or $-R^8-OR^5$;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from dioxolyl or tetrahydrofuranyl, and R^{3a} and R^{3d} are each hydrogen;

each R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroaryl group is optionally substituted by one or more substituents selected from the

group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-OR^5$, $-R^8-C(O)OR^5$, $-R^8-N(R^4)R^5$, $-R^8-C(O)N(R^4)R^5$, $-R^8-N(R^5)C(O)R^4$, $-R^8-S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8-CN$, or $-R^8-NO_2$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2; and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$; $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

R^{3a} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached,

may form a fused ring selected from dioxolyl or tetrahydrofuranyl, and R^{3a} and R^{3d} are as defined above;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroaryl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-OR^5$, $-R^8-C(O)OR^5$, $-R^8-N(R^4)R^5$, $-R^8-C(O)N(R^4)R^5$, $-R^8-N(R^5)C(O)R^4$, $-R^8-S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8-CN$, or $-R^8-NO_2$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2; and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more

substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$; $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$; $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1 or 2;

Q is -O-;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1 or 2;

Q is $-C(R^{1a})H-$, $-C(O)-$, $-CF_2-$, $-C(O)O-$ or $-N(R^5)C(O)-$;

R^{1a} is hydrogen or $-OR^5$;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

or R^1 is aralkyl substituted by $-C(O)N(R^6)R^7$ where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is hydrogen, alkyl, haloalkyl, $-R^9-CN$, $-R^9-OR^5$, $-R^9-N(R^4)R^5$, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl;

or R^6 and R^7, together with the nitrogen to which they are attached, form a heterocyclyl or heteroaryl;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroaryl groups for R^6 and R^7 may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, $-R^8-CN$, $-R^8-OR^5$, heterocyclyl and heteroaryl;

or R^1 is aralkyl optionally substituted by one or more substituents selected from the group consisting of $-R^8-OR^5$, $-C(O)OR^5$, halo, haloalkyl, alkyl, nitro, cyano, aryl, aralkyl, heterocyclyl and heteroaryl;

or R^1 is $-R^9-N(R^{10})R^{11}$, $-R^9-N(R^{12})C(O)R^{11}$ or $-R^9-N(R^{10})C(O)N(R^{10})R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9-OC(O)R^5$, $-R^9-C(O)OR^5$, $-R^9-C(O)N(R^4)R^5$, $-R^9-C(O)R^5$, $-R^9-N(R^4)R^5$, $-R^9-OR^5$, or $-R^9-CN$;

R^{12} is hydrogen, alkyl, aryl, arakyl or $-C(O)R^5$;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, $-R^8-CN$, $-R^8-OR^5$, $-R^8-C(O)R^5$, heterocyclyl and heteroaryl;

or R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroaryl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-OR^5$, $-R^8-C(O)OR^5$, $-R^8-N(R^4)R^5$, $-R^8-C(O)N(R^4)R^5$, $-R^8-N(R^5)C(O)R^4$, $-R^8-S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8-CN$, or $-R^8-NO_2$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$,

wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl,

alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$; $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl,

alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1 or 2;

Q is $-C(R^{1a})H-$;

R^{1a} is hydrogen or $-OR^5$;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2; and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$,

$-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$; $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5,

together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1 or 2;

Q is $-C(R^{1a})H-$;

R^{1a} is hydrogen or $-OR^5$;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl;

each R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1 or 2;

Q is $-C(R^{1a})H-$;

R^{1a} is hydrogen or $-OR^5$;

R^1 is pentyl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring; and

each R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1 or 2;

Q is -C(O)-, $-CF_2-$, -C(O)O- or $-N(R^5)C(O)-$;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

or R^1 is aralkyl substituted by $-C(O)N(R^6)R^7$ where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is hydrogen, alkyl, haloalkyl, $-R^9-CN$, $-R^9-OR^5$, $-R^9-N(R^4)R^5$, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl;

or R^6 and R^7, together with the nitrogen to which they are attached, form a heterocyclyl or heteroaryl;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroaryl groups for R^6 and R^7 may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, $-R^8-CN$, $-R^8-OR^5$, heterocyclyl and heteroaryl;

or R^1 is aralkyl optionally substituted by one or more substituents selected from the group consisting of $-R^8-OR^5$, $-C(O)OR^5$, halo, haloalkyl, alkyl, nitro, cyano, aryl, aralkyl, heterocyclyl and heteroaryl;

or R^1 is $-R^9-N(R^{10})R^{11}$, $-R^9-N(R^{12})C(O)R^{11}$ or $-R^9-N(R^{10})C(O)N(R^{10})R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9-OC(O)R^5$, $-R^9-C(O)OR^5$, $-R^9-C(O)N(R^4)R^5$, $-R^9-C(O)R^5$, $-R^9-N(R^4)R^5$, $-R^9-OR^5$, or $-R^9-CN$;

R^{12} is hydrogen, alkyl, aryl, arakyl or $-C(O)R^5$;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, $-R^8-CN$, $-R^8-OR^5$, $-R^8-C(O)R^5$, heterocyclyl and heteroaryl;

or R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroaryl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-OR^5$, $-R^8-C(O)OR^5$, $-R^8-N(R^4)R^5$, $-R^8-C(O)N(R^4)R^5$, $-R^8-N(R^5)C(O)R^4$, $-R^8-S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8-CN$, or $-R^8-NO_2$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2; and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$; $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached,

may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$; $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$,$-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set

forth above in the Summary of the invention wherein:

j is 0 and k is 1 or 2;

Q is -C(O)-, -CF_2-, -C(O)O- or -N(R^5)C(O)-;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, -R^8-C(O)R^5, -R^8-C(O)OR^5, -R^8-C(O)N(R^4)R^5, -S(O)$_2$-R^5, -R^9-S(O)$_m$-R^5 (where m is 0, 1 or 2), -R^8-OR^5, -R^8-CN, -R^9-P(O)(OR^5)$_2$, or -R^9-O-R^9-OR^5;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl;

each R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

Another embodiment of the invention are the compounds of formula (I) as set forth above in the Summary of the invention wherein:

j is 0 and k is 1 or 2;

Q is -C(O)-, -CF_2-, -C(O)O- or -N(R^5)C(O)-;

R^1 is pentyl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring; and

each R^5 is independently selected from group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl.

Specific embodiments of the compounds of formula (I) are described in more detail below in the Preparation of the Compounds of the Invention.

UTILITY AND TESTING OF THE COMPOUNDS OF THE INVENTION

The present invention relates to compounds, pharmaceutical compositions and methods of using the compounds and pharmaceutical compositions for the treatment of sodium channel-mediated diseases, preferably diseases related to pain, central nervous conditions such as epilepsy, anxiety, depression and bipolar disease; cardiovascular conditions such as arrhythmias, atrial fibrillation and ventricular fibrillation; neuromuscular conditions such as restless leg syndrome and muscle paralysis or tetanus; neuroprotection against stroke, neural trauma and multiple sclerosis; and channelopathies such as erythromyalgia and familial rectal pain syndrome, by administering to a patient in need of such treatment an effective amount of a sodium channel blocker modulating, especially inhibiting, agent.

In general, the present invention provides a method for treating a patient for, or protecting a patient from developing, a sodium channel-mediated disease, especially pain, comprising administering to an animal, such as a mammal, especially a human patient in need thereof, a therapeutically effective amount of a compound of the invention or a pharmaceutical composition comprising a compound of the invention wherein the compound modulates the activity of one or more voltage-dependent sodium channels.

The general value of the compounds of the invention in mediating, especially inhibiting, the sodium channel ion flux can be determined using the assays described below in the Biological Assays section. Alternatively, the general value of the compounds in treating conditions and diseases may be established in industry standard animal models for demonstrating the efficacy of compounds in treating pain. Animal models of human neuropathic pain conditions have been developed that result in reproducible sensory deficits (allodynia, hyperalgesia, and spontaneous pain) over a sustained period of time that can be evaluated by sensory testing. By establishing the degree of mechanical, chemical, and temperature induced allodynia and hyperalgesia present, several physiopathological conditions observed in humans can be modeled allowing the evaluation of pharmacotherapies.

In rat models of peripheral nerve injury, ectopic activity in the injured nerve corresponds to the behavioural signs of pain. In these models, intravenous application of the sodium channel blocker and local anesthetic lidocaine can suppress the ectopic activity and reverse the tactile allodynia at concentrations that do not affect general behaviour and motor function (Mao, J. and Chen, L.L, *Pain* (2000), 87:7-17). Allimetric

scaling of the doses effective in these rat models, translates into doses similar to those shown to be efficacious in humans (Tanelian, D.L. and Brose, W.G., *Anesthesiology* (1991), 74(5):949-951). Furthermore, Lidoderm®, lidocaine applied in the form of a dermal patch, is currently an FDA approved treatment for post-herpetic neuralgia (Devers, A. and Glaler, B.S., *Clin. J. Pain* (2000), 16(3):205-8).

Sodium channel blockers have clinical uses in addition to pain. Epilepsy and cardiac arrhythmias are often targets of sodium channel blockers. Recent evidence from animal models suggest that sodium channel blockers may also be useful for neuroprotection under ischaemic conditions caused by stroke or neural trauma and in patients with multiple sclerosis (MS) (Clare, J.J. et al., *op. cit.* and Anger, T. et al., *op. cit.*).

The compounds of the invention modulate, preferably inhibit, ion flux through a voltage-dependent sodium channel in a mammal, especially in a human. Any such modulation, whether it be partial or complete inhibition or prevention of ion flux, is sometimes referred to herein as "blocking" and corresponding compounds as "blockers". In general, the compounds of the invention modulates the activity of a sodium channel downwards, inhibits the voltage-dependent activity of the sodium channel, and/or reduces or prevents sodium ion flux across a cell membrane by preventing sodium channel activity such as ion flux.

The compounds of the instant invention are sodium channel blockers and are therefore useful for treating diseases and conditions in humans and other organisms, including all those human diseases and conditions which are the result of aberrant voltage-dependent sodium channel biological activity or which may be ameliorated by modulation of voltage-dependent sodium channel biological activity.

As defined herein, a sodium channel-mediated disease or condition refers to a disease or condition which is ameliorated upon modulation of the sodium channel and includes, but is not limited to, pain, central nervous conditions such as epilepsy, anxiety, depression and bipolar disease; cardiovascular conditions such as arrhythmias, atrial fibrillation and ventricular fibrillation; neuromuscular conditions such as restless leg syndrome and muscle paralysis or tetanus; neuroprotection against stroke, neural trauma and multiple sclerosis; and channelopathies such as erythromyalgia and familial rectal pain syndrome.

A sodium channel-mediated disease or condition also includes pain associated with HIV, HIV treatment induced neuropathy, trigeminal neuralgia, glossopharyngeal neuralgia, neuropathy secondary to metastatic infiltration, adiposis dolorosa, thalamic

lesions, hypertension, autoimmune disease, asthma, drug addiction (e.g. opiate, benzodiazepine, amphetamine, cocaine, alcohol, butane inhalation), Alzheimer, dementia, age-related memory impairment, Korsakoff syndrome, restenosis, urinary dysfunction, incontinence, parkinson's disease, cerebrovascular ischemia, neurosis, gastrointestinal disease, sickle cell anemia, transplant rejection, heart failure, myocardial infarction, reperfusion injury, intermittant claudication, angina, convulsion, respiratory disorders, cerebral or myocardial ischemias, long-QT syndrome, Catecholeminergic polymorphic ventricular tachycardia, ophthalmic diseases, spasticity, spastic paraplegia, myopathies, myasthenia gravis, paramyotonia congentia, hyperkalemic periodic paralysis, hypokalemic periodic paralysis, alopecia, anxiety disorders, psychotic disorders, mania, paranoia, seasonal affective disorder, panic disorder, obsessive compulsive disorder (OCD), phobias, autism, Aspergers Syndrome, Retts syndrome, disintegrative disorder, attention deficit disorder, aggressivity, impulse control disorders, thrombosis, pre clampsia, congestive cardiac failure, cardiac arrest, Freidrich's ataxia, Spinocerebellear ataxia, myelopathy, radiculopathy, systemic lupus erythamatosis, granulomatous disease, olivo-ponto-cerebellar atrophy, spinocerebellar ataxia, episodic ataxia, myokymia, progressive pallidal atrophy, progressive supranuclear palsy and spasticity, traumatic brain injury, cerebral oedema, hydrocephalus injury, spinal cord injury, anorexia nervosa, bulimia, Prader-Willi syndrome, obesity, optic neuritis, cataract, retinal haemorrhage, ischaemic retinopathy, retinitis pigmentosa, acute and chronic glaucoma, macular degeneration, retinal artery occlusion, Chorea, Huntington's chorea, cerebral edema, proctitis, post-herpetic neuralgia, eudynia, heat sensitivity, sarcoidosis, irritable bowel syndrome, Tourette syndrome, Lesch-Nyhan Syndrome, Brugado syndrome, Liddle syndrome, Crohns disease, multiple sclerosis and the pain associated with multiple sclerosis (MS), amyotrophic lateral sclerosis (ALS), disseminated sclerosis, diabetic neuropathy, peripheral neuropathy, charcot marie tooth syndrome, arthritic, rheumatoid arthritis, osteoarthritis, chondrocalcinosis, atherosclerosis, paroxysmal dystonia, myasthenia syndromes, myotonia, myotonic dystrophy, muscular dystrophy, malignant hyperthermia, cystic fibrosis, pseudoaldosteronism, rhabdomyolysis, mental handicap, hypothyroidism, bipolar depression, anxiety, schizophrenia, sodium channel toxin related illnesses, familial erythermalgia, primary erythermalgia, rectal pain, cancer, epilepsy, partial and general tonic seizures, febrile seizures, absence seizures (petit mal), myoclonic seizures, atonic seizures, clonic seizures, Lennox Gastaut, West Syndome (infantile spasms), multiresistant seizures, seizure prophylaxis (anti-

epileptogenic), familial Mediterranean fever syndrome, gout, restless leg syndrome, arrhythmias, fibromyalgia, neuroprotection under ischaemic conditions caused by stroke or neural trauma, tachy-arrhythmias, atrial fibrillation and ventricular fibrillation and as a general or local anaesthetic.

As used herein, the term "pain" refers to all categories of pain and is recognized to include, but is not limited to, neuropathic pain, inflammatory pain, nociceptive pain, idiopathic pain, neuralgic pain, orofacial pain, burn pain, burning mouth syndrome, somatic pain, visceral pain, myofacial pain, dental pain, cancer pain, chemotherapy pain, trauma pain, surgical pain, post-surgical pain, childbirth pain, labor pain, reflex sympathetic dystrophy, brachial plexus avulsion, neurogenic bladder, acute pain (e.g. musculoskeletal and post-operative pain), chronic pain, persistent pain, peripherally mediated pain, centrally mediated pain, chronic headache, migraine headache, familial hemiplegic migraine, conditions associated with cephalic pain, sinus headache, tension headache, phantom limb pain, peripheral nerve injury, pain following stroke, thalamic lesions, radiculopathy,HIV pain, post-herpetic pain, non-cardiac chest pain, irritable bowel syndrome and pain associated with bowel disorders and dyspepsia, and combinations thereof.

The compounds identified in the instant specification inhibit the ion flux through a voltage-dependent sodium channel. Preferably, the compounds are state or frequency dependent modifers of the sodium channels, having a low affinity for the rested/closed state and a high affinity for the inactivated state. These compounds are likely to interact with overlapping sites located in the inner cavity of the sodium conducting pore of the channel similar to that described for other state-dependent sodium channel blockers (Cestèle, S., et al., *op. cit.*). These compounds may also be likely to interact with sites outside of the inner cavity and have allosteric effects on sodium ion conduction through the channel pore.

Any of these consequences may ultimately be responsible for the overall therapeutic benefit provided by these compounds.

The present invention readily affords many different means for identification of sodium channel modulating agents that are useful as therapeutic agents. Identification of modulators of sodium channel can be assessed using a variety of *in vitro* and *in vivo* assays, e.g. measuring current, measuring membrane potential, measuring ion flux, (e.g. sodium or guanidinium), measuring sodium concentration, measuring second messengers and transcription levels, and using e.g., voltage-sensitive dyes, radioactive tracers, and patch-clamp electrophysiology.

One such protocol involves the screening of chemical agents for ability to modulate the activity of a sodium channel thereby identifying it as a modulating agent.

A typical assay described in Bean et al., *J. General Physiology* (1983), 83:613-642, and Leuwer, M., et al., *Br. J. Pharmacol* (2004), 141(1):47-54, uses patch-clamp techniques to study the behaviour of channels. Such techniques are known to those skilled in the art, and may be developed, using current technologies, into low or medium throughput assays for evaluating compounds for their ability to modulate sodium channel behaviour.

A competitive binding assay with known sodium channel toxins such as tetrodotoxin, alpha-scorpion toxins, aconitine, BTX and the like, may be suitable for identifying potential therapeutic agents with high selectivity for a particular sodium channel. The use of BTX in such a binding assay is well known and is described in McNeal, E.T., et al., *J. Med. Chem.* (1985), 28(3):381-8; and Creveling, C.R., et al., *Methods in Neuroscience, Vol.8: Neurotoxins* (Conn PM Ed) (1992), pp. 25-37, Academic Press, New York.

These assays can be carried out in cells, or cell or tissue extracts expressing the channel of interest in a natural endogenous setting or in a recombinant setting. The assays that can be used include plate assays which measure Na+ influx through surrogate markers such as ^{14}C-guanidine influx or determine cell depolarization using fluorescent dyes such as the FRET based and other fluorescent assays or a radiolabelled binding assay employing radiolabelled aconitine, BTX, TTX or STX. More direct measurements can be made with manual or automated electrophysiology systems. The guanidine influx assay is explained in more detail below in the Biological Assays section.

Throughput of test compounds is an important consideration in the choice of screening assay to be used. In some strategies, where hundreds of thousands of compounds are to be tested, it is not desirable to use low throughput means. In other cases, however, low throughput is satisfactory to identify important differences between a limited number of compounds. Often it will be necessary to combine assay types to identify specific sodium channel modulating compounds.

Electrophysiological assays using patch clamp techniques is accepted as a gold standard for detailed characterization of sodium channel compound interactions, and as described in Bean et al., *op. cit.* and Leuwer, M., et al., *op. cit.* There is a manual low-throughput screening (LTS) method which can compare 2-10 compounds per day; a recently developed system for automated medium-throughput screening

(MTS) at 20-50 patches (i.e. compounds) per day; and a technology from Molecular Devices Corporation (Sunnyvale, CA) which permits automated high-throughput screening (HTS) at 1000-3000 patches (i.e. compounds) per day.

One automated patch-clamp system utilizes planar electrode technology to accelerate the rate of drug discovery. Planar electrodes are capable of achieving high-resistance, cells-attached seals followed by stable, low-noise whole-cell recordings that are comparable to conventional recordings. A suitable instrument is the PatchXpress 7000A (Axon Instruments Inc, Union City, CA). A variety of cell lines and culture techniques, which include adherent cells as well as cells growing spontaneously in suspension are ranked for seal success rate and stability. Immortalized cells (e.g. HEK and CHO) stably expressing high levels of the relevant sodium ion channel can be adapted into high-density suspension cultures.

Other assays can be selected which allow the investigator to identify compounds which block specific states of the channel, such as the open state, closed state or the resting state, or which block transition from open to closed, closed to resting or resting to open. Those skilled in the art are generally familiar with such assays.

Binding assays are also available, however these are of only limited functional value and information content. Designs include traditional radioactive filter based binding assays or the confocal based fluorescent system available from Evotec OAI group of companies (Hamburg, Germany), both of which are HTS.

Radioactive flux assays can also be used. In this assay, channels are stimulated to open with veratridine or aconitine and held in a stabilized open state with a toxin, and channel blockers are identified by their ability to prevent ion influx. The assay can use radioactive ^{22}[Na] and ^{14}[C] guanidinium ions as tracers. FlashPlate & Cytostar-T plates in living cells avoids separation steps and are suitable for HTS. Scintillation plate technology has also advanced this method to HTS suitability. Because of the functional aspects of the assay, the information content is reasonably good.

Yet another format measures the redistribution of membrane potential using the FLIPR system membrane potential kit (HTS) available from Molecular Dynamics (a division of Amersham Biosciences, Piscataway, NJ). This method is limited to slow membrane potential changes. Some problems may result from the fluorescent background of compounds. Test compounds may also directly influence the fluidity of the cell membrane and lead to an increase in intracellular dye concentrations. Still,

because of the functional aspects of the assay, the information content is reasonably good.

Sodium dyes can be used to measure the rate or amount of sodium ion influx through a channel. This type of assay provides a very high information content regarding potential channel blockers. The assay is functional and would measure Na+ influx directly. CoroNa Red, SBFI and/or sodium green (Molecular Probes, Inc. Eugene OR) can be used to measure Na influx; all are Na responsive dyes. They can be used in combination with the FLIPR instrument. The use of these dyes in a screen has not been previously described in the literature. Calcium dyes may also have potential in this format.

In another assay, FRET based voltage sensors are used to measure the ability of a test compound to directly block Na influx. Commercially available HTS systems include the VIPR™ II FRET system (Aurora Biosciences Corporation, San Diego, CA, a division of Vertex Pharmaceuticals, Inc.) which may be used in conjunction with FRET dyes, also available from Aurora Biosciences. This assay measures sub-second responses to voltage changes. There is no requirement for a modifier of channel function. The assay measures depolarization and hyperpolarizations, and provides ratiometric outputs for quantification. A somewhat less expensive MTS version of this assay employs the FLEXstation™ (Molecular Devices Corporation) in conjunction with FRET dyes from Aurora Biosciences. Other methods of testing the compounds disclosed herein are also readily known and available to those skilled in the art.

These results provide the basis for analysis of the structure-activity relationship (SAR) between test compounds and the sodium channel. Certain substituents on the core structure of the test compound tend to provide more potent inhibitory compounds. SAR analysis is one of the tools those skilled in the art may now employ to identify preferred embodiments of the compounds of the invention for use as therapeutic agents.

Modulating agents so identified are then tested in a variety of *in vivo* models so as to determine if they alleviate pain, especially chronic pain or other conditions such as arrhythmias and epilepsy with minimal adverse events. The assays described below in the Biological Assays Section are useful in assessing the biological activity of the instant compounds.

Typically, a successful therapeutic agent of the present invention will meet some or all of the following criteria. Oral availability should be at or above 20%. Animal model efficacy is less than about 0.1 µg to about 100 mg/Kg body weight and

the target human dose is between 0.1 µg to about 100 mg/Kg body weight, although doses outside of this range may be acceptable ("mg/Kg" means milligrams of compound per kilogram of body mass of the subject to whom it is being administered). The therapeutic index (or ratio of toxic dose to therapeutic dose) should be greater than 100. The potency (as expressed by IC_{50} value) should be less than 10 µM, preferably below 1 µM and most preferably below 50 nM. The IC_{50} ("Inhibitory Concentration – 50%") is a measure of the amount of compound required to achieve 50% inhibition of ion flux through a sodium channel, over a specific time period, in an assay of the invention. Compounds of the present invention in the guanidine influx assay have demonstrated IC-50s ranging from less than a nanomolar to less than 10 micromolar.

In an alternative use of the invention, the compounds of the invention can be used in *in vitro* or *in vivo* studies as exemplary agents for comparative purposes to find other compounds also useful in treatment of, or protection from, the various diseases disclosed herein.

Another aspect of the invention relates to inhibiting $Na_V1.1$, $Na_V1.2$, $Na_V1.3$, $Na_V1.4$, $Na_V1.5$, $Na_V1.6$, $Na_V1.7$, $Na_V1.8$, or $Na_V1.9$ activity in a biological sample or a patient, which method comprises administering to the patient, or contacting said biological sample with a compound of formula I or a composition comprising said compound. The term "biological sample", as used herein, includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

Inhibition of $Na_V1.1$, $Na_V1.2$, $Na_V1.3$, $Na_V1.4$, $Na_V1.5$, $Na_V1.6$, $Na_V1.7$, $Na_V1.8$, or $Na_V1.9$ activity in a biological sample is useful for a variety of purposes that are known to one of skill in the art. Examples of such purposes include, but are not limited to, the study of sodium ion channels in biological and pathological phenomena; and the comparative evaluation of new sodium ion channel inhibitors.

PHARMACEUTICAL COMPOSITIONS OF THE INVENTION AND ADMINISTRATION

The present invention also relates to pharmaceutical composition containing the compounds of the invention disclosed herein. In one embodiment, the present invention relates to a composition comprising compounds of the invention in a pharmaceutically acceptable carrier and in an amount effective to modulate, preferably inhibit, ion flux through a voltage-dependent sodium channel to treat sodium channel

mediated diseases, such as pain, when administered to an animal, preferably a mammal, most preferably a human patient.

The pharmaceutical compositions useful herein also contain a pharmaceutically acceptable carrier, including any suitable diluent or excipient, which includes any pharmaceutical agent that does not itself induce the production of antibodies harmful to the individual receiving the composition, and which may be administered without undue toxicity. Pharmaceutically acceptable carriers include, but are not limited to, liquids, such as water, saline, glycerol and ethanol, and the like. A thorough discussion of pharmaceutically acceptable carriers, diluents, and other excipients is presented in REMINGTON'S PHARMACEUTICAL SCIENCES (Mack Pub. Co., N.J. current edition).

Those skilled in the art know how to determine suitable doses of the compounds for use in treating the diseases and conditions contemplated herein. Therapeutic doses are generally identified through a dose ranging study in humans based on preliminary evidence derived from animal studies. Doses must be sufficient to result in a desired therapeutic benefit without causing unwanted side effects for the patient.

A typical regimen for treatment of sodium channel-mediated disease comprises administration of an effective amount over a period of one or several days, up to and including between one week and about six months, or it may be chronic. It is understood that the dosage of a diagnostic/pharmaceutical compound or composition of the invention administered *in vivo* or *in vitro* will be dependent upon the age, sex, health, and weight of the recipient, severity of the symptons, kind of concurrent treatment, if any, frequency of treatment, the response of the individual, and the nature of the diagnostic/pharmaceutical effect desired. The ranges of effective doses provided herein are not intended to be limiting and represent preferred dose ranges. However, the most preferred dosage will be tailored to the individual subject, as is understood and determinable by one skilled in the relevant arts. (see, e.g., Berkowet al., eds., The Merck Manual, 16th edition, Merck and Co., Rahway, N.J., 1992; Goodmanetna., eds.,Goodman and Cilman's The Pharmacological Basis of Therapeutics, 10th edition, Pergamon Press, Inc., Elmsford, N.Y., (2001); Avery's Drug Treatment: Principles and Practice of Clinical Pharmacology and Therapeutics, 3rd edition, ADIS Press, LTD., Williams and Wilkins, Baltimore, MD. (1987), Ebadi, Pharmacology, Little, Brown and Co., Boston, (1985); Osolci al., eds.,Remington's Pharmaceutical Sciences, 18th edition, Mack Publishing Co., Easton, PA (1990);

Katzung, Basic and Clinical Pharmacology, Appleton and Lange, Norwalk, CT (1992)).

The total dose required for each treatment can be administered by multiple doses or in a single dose over the course of the day, if desired. Generally, treatment is initiated with smaller dosages, which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. The diagnostic pharmaceutical compound or composition can be administered alone or in conjunction with other diagnostics and/or pharmaceuticals directed to the pathology, or directed to other symptoms of the pathology. Effective amounts of a diagnostic pharmaceutical compound or composition of the invention are from about 0.1 μg to about 100 mg/Kg body weight, administered at intervals of 4-72 hours, for a period of 2 hours to 1 year, and/or any range or value therein, such as 0.0001-0.001, 0.001-0.01, 0.01-0.1, 0.1-1.0,1.0-10, 5-10, 10-20, 20-50 and 50-100 mg/Kg, at intervals of 1-4, 4-10, 10-16, 16-24, 24-36, 24-36, 36-48, 48-72 hours, for a period of 1-14, 14-28, or 30-44 days, or 1-24 weeks, or any range or value therein. The recipients of administration of compounds and/or compositions of the invention can be any vertebrate animal, such as mammals. Among mammals, the preferred recipients are mammals of the Orders Primate (including humans, apes and monkeys), Arteriodactyla (including horses, goats, cows, sheep, pigs), Rodenta (including mice, rats, rabbits, and hamsters), and Carnivora (including cats, and dogs). Among birds, the preferred recipients are turkeys, chickens and other members of the same order. The most preferred recipients are humans.

For topical applications, it is preferred to administer an effective amount of a pharmaceutical composition according to the invention to target area, e.g., skin surfaces, mucous membranes, and the like, which are adjacent to peripheral neurons which are to be treated. This amount will generally range from about 0.0001 mg to about 1 g of a compound of the invention per application, depending upon the area to be treated, whether the use is diagnostic, prophylactic or therapeutic, the severity of the symptoms, and the nature of the topical vehicle employed. A preferred topical preparation is an ointment, wherein about 0.001 to about 50 mg of active ingredient is used per cc of ointment base. The pharmaceutical composition can be be formulated as transdermal compositions or transdermal delivery devices ("patches"). Such compositions include, for example, a backing, active compound reservoir, a control membrane, liner and contact adhesive. Such transdermal patches may be used to provide continuous pulsatile, or on demand delivery of the compounds of the present invention as desired.

The composition may be intended for rectal administration, in the form, e.g., of a suppository which will melt in the rectum and release the drug. A typical suppository formulation will generally consist of active ingredient with a binding and/or lubricating agent such as a gelatine or cocoa butter or other low melting vegetable or synthetic wax or fat.

A typical formulation for intramuscular or intrathecal administration will consist of a suspension or solution of active in an oil or solution of active ingredient in an oil, for example arachis oil or seasame oil. A typical formulation for intravenous or intrathecal administration will consist of sterile isotonic aqueous solution containing, for example active ingredient and dextrose or sodium chloride or a mixture of dextrose and sodium chloride.

The compositions of the invention can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art. Controlled release drug delivery systems include osmotic pump systems and dissolutional systems containing polymer-coated reservoirs or drug-polymer matrix formulations. Examples of controlled release systems are given in U.S. Pat. Nos. 3,845,770 and 4,326,525 and in P. J. Kuzma et al, Regional Anesthesia 22 (6): 543-551 (1997).

The compositions of the invention can also be delivered through intra-nasal drug delivery systems for local, systemic, and nose-to-brain medical therapies. Controlled Particle Dispersion (CPD)™ technology, traditional nasal spray bottles, inhalers or nebulizers are known by those skilled in the art to provide effective local and systemic delivery of drugs by targeting the olfactory region and paranasal sinuses.

The invention also relates to an intravaginal shell or core drug delivery device suitable for administration to the human or animal female. The device may be comprised of the active pharmaceutical ingredient in a polymer matrix, surrounded by a sheath, and capable of releasing the compound in a substantially zero order pattern on a daily basis similar to devises used to apply testosterone as desscribed in PCT Patent No. WO 98/50016.

Current methods for ocular delivery include topical administration (eye drops), subconjunctival injections, periocular injections, intravitreal injections, surgical implants and iontophoresis (uses a small electrical current to transport ionized drugs into and through body tissues) Those skilled in the art would combine the best suited excipients with the compound for safe and effective intra-occular administration.

The most suitable route will depend on the nature and severity of the condition being treated. Those skilled in the art are also familiar with determining administration methods (oral, intravenous, inhalation, sub-cutaneous, rectal etc.), dosage forms, suitable pharmaceutical excipients and other matters relevant to the delivery of the compounds to a subject in need thereof.

COMBINATION THERAPY

The compounds of the invention may be usefully combined with one or more other compounds of the invention or one or more other therapeutic agent or as any combination thereof, in the treatment of sodium channel-mediated diseases and conditions. For example, a compound of formula (I) may be administered simultaneously, sequentially or separately in combination with other therapeutic agents, including, but not limited to:

- opiates analgesics, e.g. morphine, heroin, cocaine, oxymorphine, levorphanol, levallorphan, oxycodone, codeine, dihydrocodeine, propoxyphene, nalmefene, fentanyl, hydrocodone, hydromorphone, meripidine, methadone, nalorphine, naloxone, naltrexone, buprenorphine, butorphanol, nalbuphine and pentazocine;
- non-opiate analgesics, e.g. acetomeniphen, salicylates (e.g. aspirin);
- nonsteroidal antiinflammatory drugs (NSAIDs), e.g. ibuprofen, naproxen, fenoprofen, ketoprofen, celecoxib, diclofenac, diflusinal, etodolac, fenbufen, fenoprofen, flufenisal, flurbiprofen, ibuprofen, indomethacin, ketoprofen, ketorolac, meclofenamic acid, mefenamic acid, meloxicam, nabumetone, naproxen, nimesulide, nitroflurbiprofen, olsalazine, oxaprozin, phenylbutazone, piroxicam, sulfasalazine, sulindac, tolmetin and zomepirac;
- anticonvulsants, e.g. carbamazepine, oxcarbazepine, lamotrigine, valproate, topiramate, gabapentin and pregabalin;
- antidepressants such as tricyclic antidepressants, e.g. amitriptyline, clomipramine, despramine, imipramine and nortriptyline;,
- COX-2 selective inhibitors, e.g. celecoxib, rofecoxib, parecoxib, valdecoxib, deracoxib, etoricoxib, and lumiracoxib;
- alpha-adrenergics, e.g. doxazosin, tamsulosin, clonidine, guanfacine, dexmetatomidine, modafinil, and 4-amino-6,7-dimethoxy-2-(5- methane sulfonamido-1,2,3,4-tetrahydroisoquinol-2-yl)-5-(2-pyridyl) quinazoline;
- barbiturate sedatives, e.g. amobarbital, aprobarbital, butabarbital, butabital,

mephobarbital, metharbital, methohexital, pentobarbital, phenobartital, secobarbital, talbutal, theamylal and thiopental;

- tachykinin (NK) antagonist, particularly an NK-3, NK-2 or NK-1 antagonist, e.g. (αR, 9R)-7-[3,5-bis(trifluoromethyl)benzyl)]-8,9,10,11-tetrahydro-9-methyl-5-(4-methylphenyl)-7H-[1,4]diazocino[2,1-g][1,7]-naphthyridine-6-13-dione (TAK-637), 5-[[2R,3S)-2-[(1R)-1-[3,5-bis(trifluoromethylphenyl]ethoxy-3-(4-fluorophenyl)-4-morpholinyl]-methyl]-1,2-dihydro-3H-1,2,4-triazol-3-one (MK-869), aprepitant, lanepitant, dapitant or 3-[[2-methoxy5-(trifluoromethoxy)phenyl]-methylamino]-2-phenylpiperidine (2S,3S);
- coal-tar analgesics, in particular paracetamol;
- serotonin reuptake inhibitors, e.g. paroxetine, sertraline, norfluoxetine (fluoxetine desmethyl metabolite), metabolite demethylsertraline, '3 fluvoxamine, paroxetine, citalopram, citalopram metabolite desmethylcitalopram, escitalopram, d,l-fenfluramine, femoxetine, ifoxetine, cyanodothiepin, litoxetine, dapoxetine, nefazodone, ericlamine, trazodone and fluoxetine;
- noradrenaline (norepinephrine) reuptake inhibitors, e.g. maprotiline, lofepramine, mirtazepine, oxaprotiline, fezolamine, tomoxetine, mianserin, buproprion, buproprion metabolite hydroxybuproprion, nomifensine and viloxazine (Vivalan®)), especially a selective noradrenaline reuptake inhibitor such as reboxetine, in particular (S,S)-reboxetine, and venlafaxine duloxetine neuroleptics sedative/anxiolytics;
- dual serotonin-noradrenaline reuptake inhibitors, such as venlafaxine, venlafaxine metabolite O- desmethylvenlafaxine, clomipramine, clomipramine metabolite desmethylclomipramine, duloxetine, milnacipran and imipramine;
- acetylcholinesterase inhibitors such as donepezil;
- 5-HT3 antagonists such as ondansetron;
- metabotropic glutamate receptor (mGluR) antagonists;
- local anaesthetic such as mexiletine and lidocaine;
- corticosteroid such as dexamethasone;
- antiarrhythimics, e.g. mexiletine and phenytoin;
- muscarinic antagonists, e.g., tolterodine, propiverine, tropsium t chloride,

darifenacin, solifenacin, temiverine and ipratropium;

- cannabinoids;
- vanilloid receptor agonists (e.g. resinferatoxin) or antagonists (e.g. capsazepine);
- sedatives, e.g. glutethimide, meprobamate, methaqualone, and dichloralphenazone;
- anxiolytics such as benzodiazepines,
- antidepressants such as mirtazapine,
- topical agents (e.g. lidocaine, capsacin and resiniferotoxin);
- muscle relaxants such as benzodiazepines, baclofen, carisoprodol, chlorzoxazone, cyclobenzaprine, methocarbamol and orphrenadine;
- anti-histamines or H1 antagonists;
- NMDA receptor antagonists;
- 5-HT receptor agonists/antagonists;
- PDEV inhibitors;
- Tramadol®;
- cholinergic (nicotinc) analgesics;
- alpha-2-delta ligands;
- prostaglandin E2 subtype antagonists;
- leukotriene B4 antagonists;
- 5-lipoxygenase inhibitors; and
- 5-HT3 antagonists.

Sodium channel-mediated diseases and conditions that may be treated and/or prevented using such combinations include but not limited to, pain, central and peripherally mediated, acute, chronic, neuropathic as well as other diseases with associated pain and other central nervous disorders such as epilepsy, anxiety, depression and bipolar disease; or cardiovascular disorders such as arrhythmias, atrial fibrillation and ventricular fibrillation; neuromuscular disorders such as restless leg syndrome and muscle paralysis or tetanus; neuroprotection against stroke, neural trauma and multiple sclerosis; and channelopathies such as erythromyalgia and familial rectal pain syndrome.

As used herein "combination" refers to any mixture or permutation of one or more compounds of the invention and one or more other compounds of the invention or one or more additional therapeutic agent. Unless the context makes clear

otherwise, "combination" may include simultaneous or sequentially delivery of a compound of the invention with one or more therapeutic agents. Unless the context makes clear otherwise, "combination" may include dosage forms of a compound of the invention with another therapeutic agent. Unless the context makes clear otherwise, "combination" may include routes of administration of a compound of the invention with another therapeutic agent. Unless the context makes clear otherwise, "combination" may include formulations of a compound of the invention with another therapeutic agent. Dosage forms, routes of administration and pharmaceutical compositions include, but are not limited to, those described herein.

KITS-OF-PARTS

The present invention also provides kits that contain a pharmaceutical composition which includes one or more compounds of the above formulae. The kit also includes instructions for the use of the pharmaceutical composition for modulating the activity of ion channels, for the treatment of pain, as well as other utilities as disclosed herein. Preferably, a commercial package will contain one or more unit doses of the pharmaceutical composition. For example, such a unit dose may be an amount sufficient for the preparation of an intravenous injection. It will be evident to those of ordinary skill in the art that compounds which are light and/or air sensitive may require special packaging and/or formulation. For example, packaging may be used which is opaque to light, and/or sealed from contact with ambient air, and/or formulated with suitable coatings or excipients.

PREPARATION OF THE COMPOUNDS OF THE INVENTION

The following Reaction Schemes illustrate methods to make compounds of this invention, *i.e.*, compounds of formula (I): :



(I)

wherein k, j, Q, R^1, R^{2a}, R^{2b}, R^{2c}, R^{2d}, R^{3a}, R^{3b}, R^{3c}, and R^{3d} are as defined herein, as a stereoisomer, enantiomer, tautomer thereof or mixtures thereof; or a pharmaceutically

acceptable salt, solvate or prodrug thereof.

It is understood that in the following description, combinations of substituents and/or variables of the depicted formulae are permissible only if such contributions result in stable compounds.

It will also be appreciated by those skilled in the art that in the process described below the functional groups of intermediate compounds may need to be protected by suitable protecting groups. Such functional groups include hydroxy, amino, mercapto and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl or diarylalkylsilyl (*e.g.*, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl or trimethylsilyl), tetrahydropyranyl, benzyl, and the like. Suitable protecting groups for amino, amidino and guanidino include *t*-butoxycarbonyl, benzyloxycarbonyl, and the like. Suitable protecting groups for mercapto include -C(O)-R" (where R" is alkyl, aryl or arylalkyl), *p*-methoxybenzyl, trityl and the like. Suitable protecting groups for carboxylic acid include alkyl, aryl or arylalkyl esters.

Protecting groups may be added or removed in accordance with standard techniques, which are known to one skilled in the art and as described herein.

The use of protecting groups is described in detail in Green, T.W. and P.G.M. Wuts, *Protective Groups in Organic Synthesis* (1999), 3rd Ed., Wiley. The protecting group may also be a polymer resin such as a Wang resin or a 2-chlorotrityl-chloride resin.

It will also be appreciated by those skilled in the art, although such protected derivatives of compounds of this invention may not possess pharmacological activity as such, they may be administered to a mammal and thereafter metabolized in the body to form compounds of the invention which are pharmacologically active. Such derivatives may therefore be described as "prodrugs". All prodrugs of compounds of this invention are included within the scope of the invention.

The following Reaction Schemes illustrate methods to make compounds of this invention. It is understood that one skilled in the art would be able to make these compounds by similar methods or by methods known to one skilled in the art. It is also understood that one skilled in the art would be able to make in a similar manner as described below other compounds of formula (I) not specifically illustrated below by using the appropriate starting components and modifying the parameters of the synthesis as needed. In general, starting components may be obtained from sources such as Sigma Aldrich, Lancaster Synthesis, Inc., Maybridge, Matrix Scientific, TCI, and Fluorochem USA, etc. or synthesized according to sources known to those skilled

in the art (see, e.g., Smith, M.B. and J. March, *Advanced Organic Chemistry: Reactions*, Mechanisms, and Structure, 5th edition (Wiley, December 2000)) or prepared as described herein.

In the following Reaction Schemes, R^1, R^{2a}, R^{2b}, R^{2c}, R^{2d}, R^{3a}, R^{3b}, R^{3c} and R^{3d} are defined as in the Specification unless specifically defined otherwise. X is Cl or Br. R^{11} is an alkyl group.

In general, the compounds of formula (I) of the invention where Q is -O-, j is 0 and k is 1 can be synthesized following the general procedure as described below in REACTION SCHEME 1. As set forth below, an isatin compound of formula (101) is alkylated with the chloro or bromo compound of formula (102) to afford the product of formula (103). The phenol compound of formula (104) is treated with a Grignard reagent of formula (105) at low temperature (0 °C) to form the phenoxymagnesium halide intermediate which reacts with the keto-carbonyl group of the isatin compound of formula (103) in a solvent, such as, but not limited to, methylene chloride or toluene, to afford the oxindole of formula (106). The compound of formula (107) is obtained after the removal of the hydroxyl group at C-3 position of the oxindole by treating the compound of formula (106) with silane such as triethylsilane. The compound of formula (107) can also be achieved by treating the compound of formula (106) with $SOCl_2/NEt_3$ then reduction with Zn dust. Compound of formula (107) is treated with a silyl compound, such as, but not limited to, trimethylsilyl chloride, to generate the silyl ether intermediate which is treated with ytterbium (III) trifluoromethanesulfonate and formaldehyde to afford the compound of formula (108). Alternatively, compound of formula (108) can be obtained by treating the compound of formula (107) with a base, such as, but not limited to, LiOH, iPr_2NH, LDA, and subsequently reacting with formaldehyde. Intramolecular cyclization via Mitsunobu reaction affords the compound of formula (I) of the invention where Q is -O-, j is 0 and k is 1.

REACTION SCHEME 1



Formula (I)

REACTION SCHEME 1.1 below illustrates a schematic synthesis of amide and heterocyclic compounds as compounds of formula (I). When R^1 consists of an ester group, a compound such as a compound of formula (109) (in which A is alkyl or aralkyl) can be converted to the corresponding carboxylic acid compound of formula (110) by treatment of a compound of formula (109) with a base such as, but not limited

to, lithium hydroxide, sodium hydroxide or potassium hydroxide, in a mixed solvent such as, but not limited to, tetrahydrofuran or methanol with water. The acid compound of formula (110) can be converted to a mixed anhydride, by treatment with iso-butyl chloroformate in the presence of a base such as, but not limited to, *N*-methylmorpholine, or to the corresponding acid chloride, by treament with oxalyl chloride in the presence of catalytic amount of *N*,*N*-dimethylformamide in a solvent such as, but not limited to, toluene, dichloromethane or chloroform. The mixed anhydride can react directly with, or the acid chloride can react with, in the presence of a base such as, but not limited to, triethylamine or diisopropyl ethylamine, a primary or secondary amine to form the amide compound of formula (111) as a compound of formula (I). The acid compound of formula (110) can react with an aromatic diamine compound in a solvent such as, but not limited to, toluene to form the benzimidazole compound of formula (111.1) as a compound of formula (I).

REACTION SCHEME 1.1



(109) **(110)** **(111)** **(111.1)**

REACTION SCHEME 1.2 below illustrates a schematic synthesis of amine

compounds as compounds of formula (I). From compound (112), after removal of the protecting group (PG) such as, but not limited to, phthalimido or *tert*-butyloxycarbonyl, either the primary or secondary amino compound of formula (113) can be formed. Reaction of the amino compound of formula (113) with an acyl chloride in the presence of a base such as, but not limited to, triethylamine or diisopropyl ethylamine, in a solvent such as, but not limited to, toluene, dichloromethane or chloroform provides the amide compound of formula (114) as a compound of formula (I). Treatment of amino compound of formula (113) with an isocyanate in the presence of a base such as, but not limited to, triethylamine or diisopropyl ethylamine, in a solvent such as, but not limited to, dichloromethane or chloroform leads to the formation of the urea compound of formula (115) as a compound of formula (I). When the primary or secondary amino compound of formula (113) is treated with an aldehyde or a ketone in the presence of a reducing agent such as, but not limited to, sodium cyanoborohydride or sodium triacetoxyborohydride, in a solvent such as, but not limited to, dichloromethane, a high order functionalized amine (116) is produced as a compound of formula (I).

REACTION SCHEME 1.2



REACTION SCHEME 1.3 below illustrates a schematic synthesis of amine compounds as compounds of formula (I). The alcohol compound of formula (118), upon removal of the protecting group in compound of formula (117), can be oxidized to the aldehyde compound of (119) by using an oxidant such as, but not limited to, pyridinium dichromate or Dess-Martin's reagent. Similarly to the transformation of the compound of formula (113) to the compound of formula (116) as illustrated in REACTION SCHEME 1.2, the amine compound of formula (120) can be obtained as a

compound of formula (I) through the reductive amination of the aldehyde compound of formula (119) with a primary or secondary amine.

REACTION SCHEME 1.3



(117) (118) (119) (120)

REACTION SCHEME 1.4 below illustrates an alternative synthesis of compounds of formula (I) with the introduction of a variety of R^1 groups. Compound of formula (121) where PG is a protecting group such as, but not limited to, diphenylmethyl, can be synthesized through the sequence as shown in REACTION SCHEME 1 above. The protecting group can be removed under a high pressure of hydrogen such as 60 psi to form the oxindole compound of formula (122). The formation of a compound of formula (I) can be achieved by alkylation of the compound of formula (122) with a halide reagent XR^1 (where X is chloro, bromo or iodo) in the presence of a base such as, but not limited to, sodium hydride, sodium bis(trimethylsilyl)amide, and lithium hydroxide, in a solvent such as, but not limited to, *N*,*N*-dimethylformamide, tetrahydrofuran, acetone or acetonitrile. Alternatively, reaction of compound of formula (122) with an alcohol under Mitsunobu reaction conditions in the presence of a phosphine reagent such as, but not limited to,

triphenylphosphine, tributylphosphine or trimethyl phosphine, and azadicarboxylate of diethyl, diisopropyl or di-*tert*-butyl in a solvent such as, but not limited to, tetrahydrofuran, ethyl acetate or dichloromethane, provides the compound of formula (I). Alternatively, treatment of compound of formula (122) with a base such as, but not limited to, sodium hydride or lithium hydroxide, followed by reacting with an acyl chloride or anhydride, or with a sulfonyl chloride reagent, provides the corresponding acyl or sulfonyl (R^1) compound of formula (I) respectively.

REACTION SCHEME 1.4



When R^{2a}, R^{2b}, R^{2c}, R^{2d}, R^{3a}, R^{3b}, R^{3c} or R^{3d} of the compound of formula (I) is a bromo or trifluoromethylsulfonyloxy group, further derivatives can be synthesized as shown in REACTION SCHEME 1.5 and REACTION SCHEME 1.6 below. The triflate compound can be obtained by treating the bromo compound with diborane in the presence of a palladium catalyst followed by sequential oxidation with hydrogen peroxide/sodium hydroxide and reaction with trifluoromethanesulfonyl anhydride. Compounds of formula (123) or (129) (with either a bromo or a trifluoromethylsulfonyloxy group for R^{2a}, R^{2b}, R^{2c}, R^{2d}, R^{3a}, R^{3b}, R^{3c} or R^{3d}) can react

with zinc cyanide or tributyltin cyanide and potassium cyanide in the presence of a palladium catalyst such as, but not limited to, palladium acetate or tris(dibenzylideneacetone)dipalladium(0), and a ligand such as, but not limited to, tri(o-tolyl)phosphine, 1,1'-bis(diphenylphosphino)ferrocene or 2-(di-*tert*-butylphosphino)biphenyl in a solvent such as, but not limited to, *N*,*N*-dimethylformamide or acetonitrile to provide the cyano compounds of formula (124) or formula (130) as compounds of formula (I) (see Marcantonio, K.M., et al, *Org. Lett.* (2004), 6:3723-5 and Yang, C., et al, *Org. Lett.* (2004), 6:2837-40). Reaction of compounds of formula (123) or formula (129) (with either a bromo or a trifluoromethylsulfonyloxy group for R^{2a}, R^{2b}, R^{2c}, R^{2d}, R^{3a}, R^{3b}, R^{3c} or R^{3d}) with a primary or secondary amine in the presence of a palladium catalyst such as, but not limited to, palladium acetate, tetrakis(triphenylphosphine)palladium(0), or tris(dibenzylideneacetone)dipalladium(0), under a pressure of carbon monoxide in a solvent such as, but not limited to, *N*,*N*-dimethylformamide or acetonitrile leads to the formation of the amide compound of formula (125) or formula (131) as compounds of formula formula (I) (See Takahashi, T., et al, *Tetrahedron Lett.* (1999), 40:7843-6 and Schnyder, A., et al, *J. Org. Chem.* (2001), 66:4311-5). Under a typical Ullmann coupling reaction conditions, compounds of formula (123) or formula (129) (with either bromo or trifluoromethylsulfonyloxy group for R^{2a}, R^{2b}, R^{2c}, R^{2d}, R^{3a}, R^{3b}, R^{3c} or R^{3d}) can react with a phenol compound in the presence of a copper reagent such as, but not limited to, copper iodide or copper bromide, a base such as, but not limited to, cesium carbonate or potassium carbonate, an amino acid such as, but not limited to, *N*,*N*-dimethylglycine, in a solvent such as, but not limited to, dimethyl sulfoxide, dioxane or acetonitrile, to form the diaryl ether compounds of formula (126) or formula (132) as compounds of formula (I) (see Sawyer, J.S. *Tetrahedron* (2000), 56:5045-65 and Ma, D., et al, *Org. Lett.* (2003), 5(21):3799-802). Compounds of formula (123) or formula (129) (with either bromo or trifluoromethylsulfonyloxy group for R^{2a}, R^{2b}, R^{2c}, R^{2d}, R^{3a}, R^{3b}, R^{3c} or R^{3d}) can react with an arylboronic acid in the presence of a palladium catalyst such as, but not limited to, palladium acetate, tetrakis(triphenylphosphine)palladium(0), or tris(dibenzylideneacetone)dipalladium(0), with or without a ligand such as, but not limited to, triphenylphosphine, tri(o-tolyl)phosphine, 1,1'-bis(diphenylphosphino)ferrocene or 2-(di-*tert*-butylphosphino)biphenyl, a base such as, but not limited to, sodium carbonate, cesium carbonate, or sodium bicarbonate, in a solvent such as, but not limited to, dimethoxyethane, dioxane, or tetrahedrofuran to provide the coupled product of

formula (127) or formula (133) as compounds of formula (I) (see Kotha, S., et al, *Tetrahedron* (2002), 58:9633 and Miyaura, N., et al, *Chem. Rev.* (1995), 95:2457). Compound of formula (123) or formula (129) (with either a bromo or a trifluoromethylsulfonyloxy group for R^{2a}, R^{2b}, R^{2c}, R^{2d}, R^{3a}, R^{3b}, R^{3c} or R^{3d}) can react with a primary or secondary amine in the presence of a palladium catalyst such as, but not limited to, tetrakis(triphenylphosphine)palladium(0) or tris(dibenzylideneacetone)dipalladium(0), with or without a ligand such as, but not limited to, triphenylphosphine, tri(o-tolyl)phosphine, 1,1'-bis(diphenylphosphino)ferrocene or 2-(di-*tert*-butylphosphino)biphenyl, a base such as, but not limited to, sodium carbonate, cesium carbonate or sodium *tert*-butoxide, in a solvent such as, but not limited to, dioxane or tetrahedrofuran, to provide the amino compound of formula (128) or formula (134) as compounds of formula (I) (see Muci, A. R., et al, *Topics in Current Chemistry* (2002), 219:131).

REACTION SCHEME 1.5



REACTION SCHEME 1.6



(129) (130) (131) (132) (133) (134)

Alternatively, the compound of formula (I) of the invention where Q is -O and k is 1 can be synthesized following the general procedure as described below in REACTION SCHEME 2. As set forth below, a compound of formula (201) is treated

with a lithium reagent of formula (202), such as, but not limited to, *n*-BuLi at low temperature followed by the reaction with keto-carbonyl group of the isatin compound of formula (103) in a solvent, such as, but not limited to, THF to afford the oxindole of formula (203). The compound of formula (204) is obtained after the removal of the hydroxyl group at C-3 position of the oxindole by treating the compound of formula (203) with silane such as triethylsilane. The compound of formula (204) can also be achieved by treating the compound of formula (203) with $SOCl_2/NEt_3$ then reduction with Zn dust. Compound of formula (204) is treated with a silyl compound, such as, but not limited to, trimethylsilyl chloride to generate the silyl ether intermediate which is treated with ytterbium (III) trifluoromethanesulfonate and formaldehyde to afford the compound of formula (205). Alternatively, a compound of formula (205) can be obtained by treating the compound of formula (204) with a base, such as, but not limited to, LiOH, iPr_2NH, or LDA, and subsequently reacting with formaldehyde. Intramolecular cyclization via Mitsunobu reaction affords the compound of formula (I) of the invention where Q is -O- and k is 1.

REACTION SCHEME 2



Alternatively, the compound of formula (I) of the invention where Q is -O-or -S- and k is 0 can be synthesized following the general procedure as described below in REACTION SCHEME 3 wherein intramolecular cyclization of the compound of formula (203) via Mitsunobu reaction affords the compound of formula (I) of the invention where Q is -O- or -S-and k is 0.

REACTION SCHEME 3



Formula (I)

Alternatively, the compound of formula (I) of the invention where Q is -C(O)-, -(CH_2)- or –(CF_2)- and j is 0 can be synthesized following the general procedure as described below in REACTION SCHEME 4. As set forth below, the Grignard reagent of formula (401) reacts with keto-carbonyl group of the isatin compound of formula (103) in a solvent, such as, but not limited to, methylene chloride or toluene to afford the oxindole of formula (402). The compound of formula (403) is obtained after the removal of the hydroxyl group at C-3 position of the oxindole by treating the compound of formula (402) with silane such as triethylsilane. The compound of formula (403) can also be achieved by treating the compound of formula (402) with $SOCl_2/NEt_3$, followed by reduction with Zn dust. Compound of formula (403) is alkylated at C-3 position of oxindole ring with a compound of formula (404) to afford the compound of formula (405) which is subjected to saponification to generate the carboxylic acid of formula (406). This carboxylic acid is then converted to an acid chloride of formula (407) following procedures known to one skilled in the art. Intramolecular cyclization in the presence of an Lewis acid, such as, but not limited to, tin(IV) chloride, yields the compound of formula (I) of the invention where Q is -C(O)- and j is 0. The removal of the carbonyl group of the compound of formula (I) using a silane, such as triethylsilane, or other reagents known to the one skilled in the art, yields the compound of formula (I) of the invention where Q is -CH_2- and j is 0. Reaction of the carbonyl group of the compound (408) of formula (I) with a fluorinating reagent such as, but not limited to, bis(2-methoxyethyl)aminosulfur trifluoride leads to the formation of di-fluoro compound of formula (I) of the invention where Q is -CF_2- and j is 0. Reduction of the carbonyl group of the compound (408) of formula (I) with a reducing agent such as, but not limited to, sodium borohydride provides the hydroxy compound of formula (I) of the invention where Q is –CH(OH)- and j is 0. Further alkylation of the hydroxy compound of formula (I) by the method known to one skilled in the art gives the alkylated

compound of formula (I) of the invention where Q is –CH(OR^5)- and j is 0.

REACTION SCHEME 4



Formula (I)

REACTION SCHEME 4 CONTINUED



Formula (I)

Formula (I)



Formula (I)

Alternatively, the compound of formula (I) of the invention where Q is -O-, j is 0 and k is 1 can be synthesized following the general procedure as described below in REACTION SCHEME 5. As set forth below, the phenol compound of formula (104) is treated with a Grignard reagent of formula (105) at low temperature (0 °C) to form the phenoxymagnesium halide intermediate which reacts with the keto-carbonyl group of the isatin compound of formula (101) in a solvent, such as, but not limited to, tetrahedrofuran, methylene chloride or toluene, to afford the heterocyclic compound of formula (501). The compound of formula (502) can be obtained after the removal of the hydroxyl group of the heterocyclic compound by treating the compound of formula (501) with a silane such as triethylsilane. The compound of formula (502) can also be achieved by treating the compound of formula (501) with $SOCl_2/NEt_3$ followed by

reduction with Zn dust. Compound of formula (502) is treated with a silyl compound such as, but not limited to, trimethylsilyl chloride, to generate the silyl ether intermediate which is treated with ytterbium (III) trifluoromethanesulfonate and formaldehyde to afford the compound of formula (503). Alternatively, compound of formula (503) can be obtained by treating the compound of formula (502) with a base such as, but not limited to, LiOH, iPr_2NH, or LDA, and by subsequently reacting with formaldehyde. Intramolecular cyclization via Mitsunobu reaction affords the compound of formula (504) which can be alkylated with a chloro or bromo compound of formula (102) to afford the compound of formula (I) of the invention where Q is -O-, j is 0 and k is 1.

REACTION SCHEME 5



Formula (I)

Alternatively, the compound of formula (I) of the invention where Q is -NHC(O)-, j is 0 can be synthesized following the general procedure as described below in REACTION SCHEME 6. As set forth below, treatment of the ketone compound (408) with an azide such as, but not limited to, sodium azide in an acid such as, but not limited to, trifluoroacetic acid provides the Schmidt reaction product of formula (I) of the

invention where Q is –NHC(O)-, j is 0.

REACTION SCHEME 6



(408) **formula (I)**

Alternatively, the compound of formula (I) of the invention where Q is –C(O)O-, j is 0 can be synthesized following the general procedure as described below in REACTION SCHEME 7. As set forth below, treatment of the compound of formula (701), which can be obtained following a similar procedure as for the synthesis of compound of formula (405) as described in REACTION SCHEME 4, with a base such as, but not limited to, lithium hydroxide, sodium hydroxide or potassium hydroxide, in a mixed solvent such as, but not limited to, tetrahydrofuran or methanol with water, leads to the formation of the lactone product of formula (I) of the invention where Q is -C(O)O-, j is 0.

REACTION SCHEME 7



(701) **formula (I)**

In the following Preparations, which are directed to intermediates used in the preparation of the compounds of formula (I), and in the following Examples, which are directed to compounds of formula (I), the compound numbers presented therein do not correspond to the compound numbers in the above REACTION SCHEMES.

PREPARATION 1

Synthesis of 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 4-bromo-1-pentyl-1*H*-indole

To a mixture of sodium hydride (2.54 g, 66.3 mmol, 60% dispersion in mineral oil) in anhydrous *N*,*N*-dimethylformamide (50.0 mL) was added 4-bromoindole (10.0 g, 51.0 mmol) at 0 °C. The reaction mixture was stirred for 0.5 h followed by the addition of 1-bromopentane (9.25 g, 61.2 mmol) at 0 °C. The reaction mixture was stirred at ambient temperature for 6 h and quenched with brine solution (20.0 mL). The reaction mixture was diluted with water (100 mL) and extracted with ether (3 x 200 mL). The combined organic layers was washed with brine (100 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with hexane (100%) to give the title compound (13.3 g, 98%) as a yellow oil: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.30-7.27 (m, 2H), 7.14 (t, 1H), 6.88 (t, 1H), 6.55 (d, 1H), 4.08 (t, 2H), 1.87-1.77 (m, 2H), 1.39-1.22 (m, 4H), 0.89 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 136.3, 129.2, 128.4, 122.2, 122.1, 114.9, 108.7, 101.3, 46.8, 29.9, 29.1, 22.3, 13.9.

B. Syntheisis of 4-bromo-1-pentyl-1*H*-indole-2,3-dione

To a solution of 4-bromo-1-pentyl-1*H*-indole (25.0 g, 93.9 mmol) in anhydrous dimethylsulfoxide (350 mL) was added *N*-bromosuccinimide (50.2 g, 282 mmol) in portions over 30 min. The reaction mixture was heated at 60 °C for 3 h, upon which time the internal temperature increased to 120 °C. After cooling down to ambient temperature, the reaction mixture was poured onto ethyl acetate/water (1/1, 600 mL). The organic layer was separated and the aqueous layer was extracted with ethyl acetate (3 x 500 mL). The combined organic layers was washed with water (3 x 500 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness to yield the title compound (25.7 g, 92%) as a yellow solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.38 (t, 1H), 7.21 (t, 1H), 6.82 (d, 1H), 3.68 (t, 2H), 1.72-1.59 (m, 2H), 1.39-1.25 (m, 4H), 0.86 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 180.9, 157.2, 152.6, 138.4, 128.3, 121.7, 116.3, 108.9, 40.4, 28.9, 26.9, 22.3, 13.9.

C. Synthesis of 4-bromo-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

To a solution of 1,3-benzodioxol-5-ol (12.8 g, 92.9 mmol) in tetrahydrofuran (200 mL) was added isopropylmagnesium chloride solution (50.7 mL, 101 mmol, 2.0 M

in ether) at 0 °C. The reaction mixture was stirred at 0 °C for 0.5 h, upon which time the colorless precipitate was formed. After the solvent was removed under reduced pressure, the residue was dissolved in methylene chloride (100 mL) and added to a solution of 4-bromo-1-pentyl-1*H*-indole-2,3-dione (25.0 g, 84.5 mmol) in dichloromethane (100 mL) via a canula over 10 min at 0 °C. The reaction mixture was stirred at ambient temperature for 16 h, quenched with saturated ammonium chloride solution (100 mL) and the organic layer was separated. The aqueous layer was extracted with dichloromethane (100 mL). The combined organic layers was washed with brine, dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate-hexane to give the title compound (34.9 g, 97%) as a brown gummy material: ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.95 (s, 1H), 7.29-7.21 (m, 2H), 6.88-6.81 (m, 1H), 6.55, (s, 1H), 6.14 (s, 1H), 5.86 (dd, 2H), 4.24 (s, 1H), 3.70-3.52 (m, 2H), 1.69-1.55 (m, 2H), 1.31-1.24 (m, 4H), 0.83 (t, 3H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 177.6, 152.6, 149.1, 144.8, 141.2, 131.7, 127.7, 127.6, 121.0, 113.8, 108.3, 106.7, 101.7, 101.4, 80.5 40.5, 28.8, 26.7 22.2, 13.9.

D. Synthesis of 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

To a solution of 4-bromo-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one (34.9 g, 80.4 mmol) in dichloromethane (100 mL) was added trifluoroacetic acid (18.7 g, 161 mmol) and triethylsilane (18.3 g, 161 mmol). The brown solution was stirred at ambient temperature for 3 h and concentrated *in vacuo* to dryness. The residue was diluted with dichloromethane (200 mL), washed with saturated ammonium chloride solution (50.0 mL), brine (3 x 50.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was crystallized from ether to give the title compound (16.5 g, 49%) as a brown solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.29-7.21 (m, 2H), 7.14 (dd, 1H), 6.58 (s, 1H), 6.10 (s, 1H), 5.85 (dd, 2H), 5.01 (s, 1H), 3.75-3.55 (m, 2H), 1.69-1.56 (m, 2H), 1.35-1.21 (m, 4H), 0.86 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.9, 150.9, 147.6, 145.4, 141.6, 130.3 127.1 126.8, 120.8, 113.3 108.0, 106.7, 101.5, 101.2, 59.9, 48.6, 40.7, 28.9, 26.9, 22.3 13.9; MS (ES+) *m/z* 418.3 (M + 1), 420.3 (M + 1).

E. Synthesis of 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

To a solution of 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one (7.50 g, 17.9 mmol) in dry dichloromethane (150 mL) was added triethylamine (10.9 g, 108 mmol) and chlorotrimethylsilane (7.80 g, 71.8 mmol) at 0 °C. The reaction mixture was stirred at 0 °C for 2 h and diluted with dichloromethane (100 mL). The mixture was washed with water (3 x 50.0 mL), dried over anhydrous magnesium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was dissolved in THF (150 mL) followed by the additions of formaldehyde solution (4.90 mL, 179 mmol, 37 wt% in water) and ytterbium (III) trifluoromethanesulfonate (1.11 g, 1.79 mmol). The resulting mixture was stirred at ambient temperature for 36 h. After the solvent was removed under reduced pressure, the residue was diluted with dichloromethane (200 mL), washed with saturated sodium bicarbonate (50.0 mL), saturated ammonium chloride (50.0 mL) and water (100 mL). The organic layer was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness to yield the title compound (6.32 g, 79%) as a fluffy solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.28 (s, 1H), 7.10 (t, 1H), 7.00 (dd, 1H), 6.89 (dd, 1H), 6.83 (s, 1H), 6.27 (s, 1H), 6.85 (dd, 2H), 4.52-4.41 (m, 2H), 3.90 (dd, 1H), 3.70-3.65 (m, 2H), 1.68-1.57 (m, 2H), 1.36-1.29 (m, 4H), 0.83 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.1, 150.3, 147.2, 147.2, 140.5, 129.6, 129.2, 125.6, 118.4, 114.8, 109.2, 106.9, 101.0, 98.2, 62.6, 57.6, 39.9, 28.9, 26.7, 22.2, 13.5 .

PREPARATION 2

Synthesis of 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione

To a suspension of sodium hydride (1.61 g, 41.9 mmol, 60% dispersion in mineral oil) in anhydrous *N*,*N*-dimethylformamide (25.0 mL) was added isatin (6.17 g, 41.9 mmol) at 0 °C. The reaction mixture was stirred for 0.5 h followed by the addition of (2-bromoethyl)cyclopropane (Maercker, A., et al, *Justus Liebigs Ann. Chem.* (1972), 759:132-157) (9.25 g, 61.2 mmol). The resulting mixture was stirred at ambient temperature for 16 h and quenched with water (50.0 mL). The mixture was extracted with ethyl acetate (3 x 100 mL). The combined organic layers was washed with water (3 x 50.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness to yield the title compound (6.50 g, 90%) as a

viscous gum: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.57-7.51 (m, 2H), 7.05 (t, 1H), 6.88 (d, 1H), 3.79-3.74 (m, 2H), 1.59-1.52 (m, 2H), 0.70-0.61 (m, 1H), 0.44-0.38 (m, 2H), 0.05-0.02 (m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 183.7, 158.2, 151.2, 138.4, 125.4, 123.6, 117.5, 110.3, 40.3, 32.2, 8.6, 4.3.

B. Synthesis of 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one

To a solution of 1,3-benzodioxol-5-ol (1.25 g, 9.06 mmol) in THF (20.0 mL) was added dropwise a solution of isopropylmagnesium chloride solution (4.53 mL, 9.06 mmol, 2.0 M in THF) at 0 °C over 5 min. The reaction mixture was stirred for 0.5 h upon which time colorless precipitate was formed. After the solvent was removed under reduced pressure, the residue was dissolved in dichloromethane (20.0 mL) and cooled to 0 °C. A solution of 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione (1.77 g, 8.23 mmol) in dichloromethane (20.0 mL) was added to the above solution at 0 °C. The resulting mixture was stirred at ambient temperature for 16 h and quenched with saturated ammonium chloride solution (30.0 mL). The organic layer was separated and washed with water (3 x 25.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was crystallized from ethyl acetate and ether to give the title compound (2.22 g, 76%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.52 (s, 1H), 7.46 (d, 1H), 7.37 (dt, 1H), 7.18 (dt, 1H), 6.90 (d, 1H), 6.56 (s, 1H), 6.23 (s, 1H), 5.84 (dd, 2H), 4.55 (s, 1H), 3.87-3.63 (m, 2H), 1.64-1.44 (m, 2H), 0.68-0.55 (m, 1H), 0.41-0.27 (m, 2H), -0.02-(-0.07) (m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 179.1, 152.4, 148.8, 142.7, 141.3, 130.3, 129.1, 126.3, 123.7, 117.3, 109.5, 106.9, 101.9, 101.4, 79.3, 40.6, 32.2, 8.6, 4.3, 4.2; MS (ES+) *m/z* 337.6 (M - 17).

C. Synthesis of 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one

To a solution of 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one (2.22 g, 6.27 mmol) in dichloromethane (30.0 mL) was added trifluoroacetic acid (2.12 g, 18.8 mmol) and triethylsilane (2.14 g, 18.8 mmol). The brown solution was stirred at ambient temperature for 0.5 h and concentrated *in vacuo* to dryness. The residue was diluted with dichloromethane (100 mL), washed with water (3 x 50.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate/hexane (20/80) to give the title compound

(1.69 g, 80%) as a brown solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.21-9.10 (br, 1H), 7.38-7.30 (m, 2H), 7.16 (t, 1H), 6.96 (d, 1H), 6.63 (s, 1H), 6.33 (s, 1H), 5.84 (dd, 2H), 5.01 (s, 1H), 3.87-3.72 (m, 2H), 1.66-1.46 (m, 2H), 0.69-0.59 (m, 1H), 0.43-0.30 (m, 2H), 0.09-0.06 (m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.8, 151.3, 147.6, 144.1, 141.5, 128.7, 126.2, 123.1, 115.2, 109.5, 109.4, 106.5, 101.5, 101.2, 47.4, 40.5, 32.2, 8.6, 4.3, 4.2; MS (ES+) *m/z* 338.3 (M + 1).

D. Synthesis of 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, making variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (53%): R_f = 0.28 (EtOAc/Hexanes, 1/1).

PREPARATION 3

Synthesis of ethyl [3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

A. Synthesis of ethyl (2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace (2-bromoethyl)cyclopropane with ethyl bromoacetate, the title compound was obtained (79%) as a light yellow powder: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.64-7.54 (m, 2H), 7.16-7.11 (m, 1H), 6.77 (d, 1H), 4.47 (s, 2H), 4.22 (q, 2H), 1.26 (t, 3H); MS(ES+) *m/z* 256.2 (M + 23).

B. Synthesis of ethyl [3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with ethyl (2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate, the title compound was obtained (95%): ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.08 (s, 1H), 7.21-7.13 (m, 2H), 6.93-6.86 (m, 3H), 6.57 (s, 1H), 6.19 (s, 1H), 5.88 (m, 2H), 4.47 (m, 2H), 4.13 (q, 2H), 1.19 (t, 3H); MS (ES-) *m/z* 370.2 (M - 1).

C. Synthesis of ethyl [3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1D, and making non-critical variations to replace 4-bromo-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-

pentyl-1,3-dihydro-2*H*-indol-2-one with ethyl [3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained (84%) as a white powder: ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.37 (s, 1H), 7.19 (m, 1H), 7.01-6.90 (m, 3H), 6.43 (s, 2H), 5.84 (m, 2H), 4.86 (s, 1H), 4.56 (s, 2H), 4.13 (q, 2H), 1.18 (t, 3H); MS (ES+) *m/z* 378.2 (M + 23).

D. Synthesis of ethyl [3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with ethyl [3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained as a white powder: ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.03 (s, 1H), 7.17-6.85 (m, 5H), 6.22 (s, 1H), 5.83 (s, 2H), 5.04 (t, 1H), 4.56-4.08 (m, 5H), 3.69 (m, 1H), 1.18 (t, 3H); MS (ES+) *m/z* 408.1 (M + 23).

PREPARATION 4

Synthesis of methyl 3-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate

A. Synthesis of methyl 3-[(2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)methyl]benzoate

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace (2-bromoethyl)cyclopropane with methyl 3-(bromomethyl)benzoate, the title compound was obtained (84%) as a orange solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.99-7.95 (m, 2H), 7.60 (d, 1H), 7.53-7.47 (m, 2H), 7.43 (d, 1H), 7.09 (t, 1H), 6.43 (d, 1H), 4.95 (s, 2H), 3.89 (s, 3H).

B. Synthesis of methyl 3-{[3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with methyl 3-[(2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)methyl]benzoate, the title compound was obtained (96%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.65 (s, 1H), 7.92 (s, 1H), 7.85 (d, 1H), 7.41-7.38 (m, 1H), 7.32-7.24 (m, 2H), 7.19-7.13 (m, 1H), 7.04-6.9 (m, 1H), 6.63 (d, 1H), 6.44 (s, 1H), 6.39 (s, 1H), 5.79 (s, 2H), 5.05 (s, 1H), 4.83 (dd, 2H), 3.80 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.7, 167.0, 151.0, 148.5, 142.1, 141.1, 135.7, 131.6, 130.5, 130.1, 129.1, 129.0, 128.4, 125.5, 123.9, 116.7, 109.7, 106.5, 101.3, 100.5, 78.6, 60.6, 52.4, 43.6; MS (ES+) *m/z* 456.1 (M + 23).

C. Synthesis of methyl 3-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with methyl 3-{[3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate, the title compound was obtained (98%): MS (ES+) *m/z* 418.2 (M + 1).

D. Synthesis of methyl 3-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with methyl 3-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate, the title compound was obtained (81%) as a white powder: MS (ES+) *m/z* 470.3 (M + 23), 448.3 (M + 1).

PREPARATION 5

Synthesis of methyl 4-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate

A. Synthesis of methyl 4-[(2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)methyl]benzoate

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace (2-bromoethyl)cyclopropane with methyl 4-(bromomethyl)benzoate, the title compound was obtained (84%) as an orange solid: ^{1}H-NMR (300 MHz, $CDCl_3$) δ 8.00 (d, 2H), 7.61 (d, 1H), 7.46 (t, 1H), 7.38 (d, 2H), 7.09 (t, 1H), 6.69 (d, 1H), 4.96 (s, 2H), 3.88 (s, 3H); MS (ES+) *m/z* 296.1 (M + 1).

B. Synthesis of methyl 4-{[3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with methyl 4-[(2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)methyl]benzoate, the title compound was obtained (79%): MS (ES+) *m/z* 416.1 (M -17).

C. Synthesis of methyl 4-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate

Following the procedure as described in PREPARATION 2C, and making the variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with methyl 4-{[3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-

5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate, the title compound was obtained (98%) as a solid: MS (ES+) *m/z* 418.1 (M +1).

D. Synthesis of methyl 4-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with methyl 4-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate, the title compound was obtained (81%): MS (ES+) *m/z* 448.1 (M + 1).

PREPARATION 6

Synthesis of 2-{3-[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]propyl}-1*H*-isoindole-1,3(2*H*)-dione

A. Synthesis of 1-[3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propyl]-1*H*-indole-2,3-dione

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace (2-bromoethyl)cyclopropane with 2-(3-bromopropyl)-1*H*-isoindole-1,3(2*H*)-dione, the title compound was obtained (92%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.80-7.79 (m, 4H), 7.61-7.56 (m, 1H), 7.49-7.46 (m, 1H), 7.18-7.16 (m, 1H), 7.07-7.05 (m, 1H), 3.72-3.60 (m, 4H), 1.97-1.92 (m, 2H).

B. Synthesis of 2-{3-[3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]propyl}-1*H*-isoindole-1,3(2*H*)-dione

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-[3-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propyl]-1*H*-indole-2,3-dione, the title compound was obtained (96%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.86-7.78 (m, 4H), 7.21-7.13 (m, 2H), 7.00-6.97 (m, 1H), 6.87-6.85 (m, 2H), 6.15 (s, 1H), 5.86-5.84 (m, 2H), 3.69-3.65 (m, 4H), 2.46-2.45 (m, 1H), 1.94-1.87 (m, 2H); MS (ES+) *m/z* 473.4 (M - 17).

C. Synthesis of 2-{3-[3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]propyl}-1*H*-isoindole-1,3(2*H*)-dione

Following the procedure as described in PREPARATION 1D, and making non-critical variations to replace 4-bromo-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 2-{3-[3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]propyl}-1*H*-isoindole-1,3(2*H*)-dione, the title compound was obtained (94%): ^{1}H NMR (300 MHz, $CDCl_3$,) δ 7.81-7.78 (m, 2H), 7.70-

7.67 (m, 2H), 7.32-7.27 (m, 2H), 7.12-7.07 (m, 1H), 6.90-6.87 (m, 1H), 6.54 (s, 1H), 6.45 (s, 1H), 5.86 (dd, 2H), 4.82 (s, 1H), 3.96-3.66 (m, 4H), 2.17-2.04 (m, 2H); MS (ES+) *m/z* 457.0 (M + 1).

D. Synthesis of 2-{3-[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]propyl}-1*H*-isoindole-1,3(2*H*)-dione

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 2-{3-[3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]propyl}-1*H*-isoindole-1,3(2*H*)-dione, the title compound was obtained (94%) as a foam solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.20 (s, 1H), 7.81-7.79 (m, 2H), 7.68-7.61 (m, 2H), 7.35-7.25 (m, 2H), 7.16-7.14 (m, 1H), 6.90 (d, 1H), 6.80 (s, 1H), 6.48 (s, 1H), 5.86 (dd, 2H), 4.64 (d, 1H), 3.67-4.13 (m, 5H), 2.18-2.05- (m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 180.6, 168.6, 151.2, 147.8, 143.2, 141.2, 134.2, 134.2, 131.9, 130.0, 128.7, 125.1, 123.2, 113.9, 108.7, 108.3, 101.3, 100.6, 64.9, 58.0, 37.6, 36.1, 26.5; MS (ES+) *m/z* 487.3 (M + 1).

PREPARATION 7

Synthesis of 2-{2-[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]ethyl}-1*H*-isoindole-1,3(2*H*)-dione

A. Synthesis of 1-[2-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-ethyl]-1*H*-indole-2,3-dione

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace (2-bromoethyl)cyclopropane with 2-(2-bromoethyl)-1*H*-isoindole-1,3(2*H*)-dione, the title compound was obtained (75%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.85-7.78 (m, 4H), 7.65 (td, 1H), 7.55 (dd, 1H), 7.25(d, 1H), 7.12 (t, 1H), 4.00-3.80 (m, 4H); MS (ES+) *m/z* 321 (M + 1), 343 (M + 23).

B. Synthesis of 2-{2-[3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]ethyl}-1*H*-isoindole-1,3(2*H*)-dione

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-[2-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-ethyl]-1*H*-indole-2,3-dione, the title compound was obtained (99%): ^{1}H NMR (300 MHz, CD_3OD) δ 7.85-7.68 (m, 4H), 7.29 (td, 1H), 7.18-6.96 (m, 3H), 6.88 (s, 1H), 6.16 (s, 1H), 5.85 (s, 1H), 5.82 (s, 1H), 3.81-4.01 (m, 4H); MS (ES+) *m/z* 441 (M - 17), 458 (M + 23).

C. Synthesis of 2-{2-[3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]ethyl}-1*H*-isoindole-1,3(2*H*)-dione

Following the procedure as described in PREPARATION 1D, and making non-critical variations to replace 4-bromo-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 2-{2-[3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]ethyl}-1*H*-isoindole-1,3(2*H*)-dione, the title compound was obtained (90%) as a white solid: ^{1}H NMR (300 MHz, CD_3OD) δ 10.15-10.05 (br, 1H), 8.66-8.58 (m, 4H), 8.07-7.70 (m, 4H), 7.12 (s, 1H), 7.18 (s, 1H), 6.70 (s, 1H), 6.69 (s, 1H), 5.50 (s, 1H), 4.91-4.56 (m, 4H); MS (ES+) *m/z* 443 (M + 1).

D. Synthesis of 2-{2-[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]ethyl}-1*H*-isoindole-1,3(2*H*)-dione

Following the procedure as described in PREPARATION 1E, making variation to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 2-{2-[3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]ethyl}-1*H*-isoindole-1,3(2*H*)-dione, the title compound was obtained (56%): ^{1}H NMR (300 MHz, CD_3OD) δ 9.97 (s, 1H), 8.72-8.62 (m, 4H), 8.07-7.67 (m, 5H), 7.01 (s, 1H), 6.71 (s, 1H), 6.70 (s, 1H), 5.79 (t, 1H), 4.88-4.50 (m, 6H); MS (ES+) *m/z* 455 (M - 17), 473 (M + 1), 495 (M + 23).

PREPARATION 8

Synthesis of 1-(diphenylmethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 1-(diphenylmethyl)-1*H*-indole-2,3-dione

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace (2-bromoethyl)cyclopropane with 1,1'-(bromomethylene)dibenzene, the title compound was obtained (68%) as an orange solid: MS (ES+) *m/z* 336.4 (M + 23).

B. Synthesis of 1-(diphenylmethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-(diphenylmethyl)-1*H*-indole-2,3-dione, the title compound was obtained (99%) as an off-white powder: MS (ES+) *m/z* 474.5 (M + 23).

C. Synthesis of 1-(diphenylmethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making the variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 1-(diphenylmethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (84%) as an off-white solid: MS (ES+) *m/z* 458.4 (M + 23).

D. Synthesis of 1-(diphenylmethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations using 1-(diphenylmethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (56%): MS (ES+) *m/z* 488.3 (M + 23).

PREPARATION 9

Synthesis of 1-[3-(benzyloxy)propyl]-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 1-[3-(benzyloxy)propyl]-1*H*-indole-2,3-dione

Following the procedure as described in PREPARATION 1A, and making non-critical variations to replace 4-bromoindole with isatin, and 1-bromopentane with benzyl 3-bromopropyl ether, the title compound was obtained (95%) as a pale yellow syrup: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.57-6.92 (m, 9H), 4.50 (s, 2H), 3.84 (t, 2H), 3.54 (t, 2H), 2.03-1.94 (m, 2H); MS (ES^+) *m/z* 296.3 (M + 1), 318.3 (M + 23).

B. Synthesis of 1-[3-(benzyloxy)propyl]-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1C, and making non-critical variations to replace 4-bromo-1-pentyl-1*H*-indole-2,3-dione with 1-[3-(benzyloxy)propyl]-1*H*-indole-2,3-dione, the title compound was obtained (70%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.42 (s, 1H), 7.32-7.16 (m, 8H), 6.96 (d,), 6.61 (s, 1H), 6.23 (s, 1H), 5.86-5.83 (m, 2H), 4.44 (s, 2H), 3.88-3.73(m, 2H), 3.46 (t, 2H), 2.06-1.85 (m, 2H); MS (ES+) *m/z* 416.3 (M - 17), 456.3 (M + 23).

C. Synthesis of 1-[3-(benzyloxy)propyl]-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1D, and making non-

critical variations to replace 4-bromo-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 1-[3-(benzyloxy)propyl]-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (92%): ^{1}H NMR (300 MHz, $CDCl_3$) δ7.42-6.95 (m, 9H), 6.56 (s, 1H), 6.24 (s, 1H), 5.86 (ABq, 1H), 5.81(ABq, 1H), 4.99 (s, 1H), 4.42 (s, 2H), 3.91-3.76 (m, 2H), 3.46 (t, 2H), 2.03-1.93 (m, 2H); MS (ES+) *m/z* 418.3 (M + 1).

D. Synthesis of 1-[3-(benzyloxy)propyl]-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 1-[3-(benzyloxy)propyl]-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (93%): MS (ES+) *m/z* 448.2 (M + 1).

PREPARATION 10

Synthesis of methyl 2-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate

A. Synthesis of methyl 2-[(2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)methyl]benzoate

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace (2-bromoethyl)cyclopropane with methyl 2-(bromomethyl)benzoate, the title compound was obtained (68%) as a yellow solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.05 (dd, 1H), 7.64 (dd, 1H), 7.50-7.31 (m, 3H), 7.22 (d, 1H), 7.10 (t, 1H), 6.72 (d, 1H), 5.41 (s, 2H), 3.95 (s, 3H).

B. Synthesis of methyl 2-{[3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with methyl 2-[(2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)methyl]benzoate, the title compound was obtained (97%) as a colorless solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.29 (s, 1H), 7.97 (dd, 1H), 7.53-7.36 (m, 3H), 7.28 (s, 1H), 7.10 (td, 1H), 6.96-6.83 (m, 2H), 6.59 (d, 2H), 6.25 (s, 1H), 5.95-5.86 (m, 2H), 5.31-5.07 (m, 2H), 3.88 (s, 3H); MS (ES+) *m/z* 456.1 (M + 23).

C. Synthesis of methyl 2-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with methyl 2-{[3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate, the title compound was obtained (100%) as a white solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.32 (s, 1H), 7.94 (dd, 1H), 7.50-7.34 (m, 2H), 7.26 (d, 1H), 7.08 (t, 1H), 7.00-6.86 (m, 2H), 6.76 (s, 1H), 6.64 (d, 1H), 6.38 (s, 1H), 5.93-5.86 (m, 2H), 5.34-5.12 (m, 2H), 4.83 (s, 1H), 3.87 (s, 3H); MS (ES+) *m/z* 418.2 (M + 1).

D. Synthesis of methyl 2-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate

A solution of methyl 2-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate (17.1 g, 40.0 mmol) and paraformaldehyde (10.3 g, 330 mmol) in THF (500 mL) was degassed by bubbling through argon for 2 hours. To this solution was added lithium diisopropylamide solution (45.1 mL, 2 M solution, 90.0 mmol) slowly at -78 °C. The mixture was stirred at ambient temperature overnight and quenched with saturated ammonium chloride solution. The mixtrue was concentrated *in vacuo* to remove THF followed by the addition of ethyl acetate (500 mL). The organic layer was washed with water, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was recrystallized from ethyl acetate/hexanes to give the title compound (13.7 g, 75%): ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.20 (s, 1H), 7.95 (dd, 1H), 7.53-7.33 (m, 3H), 7.08-6.82 (m, 4H), 6.53 (d, 1H), 6.25 (s, 1H), 5.93-5.86 (m, 2H), 5.31-5.07 (m, 3H), 4.26-4.17 (m, 1H), 4.00-3.92 (m, 1H), 3.88 (s, 3H); MS (ES+) *m/z* 448.3 (M + 1).

PREPARATION 11

Synthesis of [3-(1,3-benzodioxol-5-yl)-2-oxo-1-pentyl-2,3-dihydro-1*H*-indol-3-yl]acetic acid

A. Synthesis of 1-pentyl-1*H*-indole-2,3-dione

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace (2-bromoethyl)cyclopropane with 1-bromopentane, the title compound was obtained (85%) as an orange solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.60-7.52 (m, 2H), 7.08 (td, 1H), 6.87 (d, 1H), 3.69 (t, 2H), 1.74-1.61 (m, 2H), 1.40-1.28 (m, 4H), 0.88 (t, 3H).

B. Synthesis of 3-(1,3-benzodioxol-5-yl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one

To a solution of 1-pentyl-1*H*-indole-2,3-dione (2.80 g, 12.8 mmol) in THF (50.0 mL) was added 3,4-(methylenedioxy)phenylmagnesium bromide (14.0 mL, 1 M THF solution, 14.0 mmol) slowly at -78 °C. The mixture was stirred at 0 °C for 1 h and quenched with ammonium chloride solution. The mixture was poured into water (150 mL) and extracted with ethyl acetate (200 mL). The organic layer was washed with water, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to flash column chromatography to afford the title compound (3.10 g, 71%) as an orange oil: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.34-7.23 (m, 2H), 7.05 (t, 1H), 6.91-6.85 (m, 2H), 6.83-6.78 (m, 1H), 6.71 (d, 1H), 5.92-5.89 (m, 2H), 3.82-3.55 (m, 2H), 3.40 (br, 1H), 1.76-1.61 (m, 2H), 1.39-1.28 (m, 4H), 0.87 (t, 3H).

C. Synthesis of 3-(1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 3-(1,3-benzodioxol-5-yl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (90%) as an oil: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.30 (td, 1H), 7.14 (d, 1H), 7.03 (td, 1H), 6.89 (d, 1H), 6.75 (d, 1H), 6.67 (dd, 1H), 6.57 (d, 1H), 5.90 (s, 2H), 4.50 (s, 1H), 3.81-3.62 (m, 2H), 1.76-1.62 (m, 2H), 1.41-1.28 (m, 4H), 0.88 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.8, 148.1, 147.2, 143.5, 130.0, 129.4, 128.4, 125.1, 122.9, 122.0, 108.7, 108.6, 101.1, 51.9, 40.4, 29.0, 27.1, 22.3, 14.0.

D. Synthesis of methyl [3-(1,3-benzodioxol-5-yl)-2-oxo-1-pentyl-2,3-dihydro-1*H*-indol-3-yl]acetate

A solution of 3-(1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one (1.00 g, 3.10 mmol) and methyl bromoacetate (0.44 mL, 4.60 mmol) in THF (20.0 mL) was degassed by bubbling through argon for one hour. Sodium hydride (0.19 g, 4.60 mmol) was added at 0 °C. The mixture was stirred at 0 °C for 1 h and quenched with ammonium chloride solution. The mixture was poured into water (150 mL), and extracted with ethyl acetate (200 mL). The organic layer was washed with water, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to flash column chromatography to afford the title compound (0.94 g, 76%) as a colorless oil: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.30 (td, 1H), 7.25 (dd, 1H), 7.06 (td, 1H), 6.89 (d, 1H), 6.81 (d, 1H), 6.74-6.65 (m, 2H), 5.90-5.87 (m,

2H), 3.71-3.64 (m, 2H), 3.45 (d, 1H), 3.41 (s, 3H), 3.18 (d, 1H), 1.74-1.60 (m, 2H), 1.39-1.22 (m, 4H), 0.85 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$,) δ 177.8, 170.0, 147.9, 147.0, 143.9, 133.1, 131.3, 128.6, 124.6, 122.3, 119.9, 108.7, 108.1, 107.4, 101.2, 52.8, 51.6, 41.8, 40.4, 29.0, 26.8, 22.3, 14.0; MS (ES+) *m/z* 418.1 (M + 23), 396.1 (M + 1).

E. Synthesis of [3-(1,3-benzodioxol-5-yl)-2-oxo-1-pentyl-2,3-dihydro-1*H*-indol-3-yl]acetic acid

To a solution of methyl [3-(1,3-benzodioxol-5-yl)-2-oxo-1-pentyl-2,3-dihydro-1*H*-indol-3-yl]acetate (5.90 g, 15.0 mmol) in THF/water (2/1 v/v, 120 mL) was added lithium hydroxide monohydrate (1.26 g, 28.0 mmol). The mixture was stirred at ambient temperature overnight. Most THF was removed under vacuum and water (150 mL) was added. The solution was extracted with ethyl acetate/hexanes (1/3 v/v, 50.0 mL). The water layer was acidified with 1 N HCl solution until the pH value reached 2 and extracted with ethyl acetate (200 mL). The organic layer was washed with water, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness to afford the title compound (5.00 g, 88%) as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.29 (td, 1H), 7.21 (dd, 1H), 7.05 (td, 1H), 6.87 (d, 1H), 6.76 (d, 1H), 6.72-6.64 (m, 2H), 5.90-5.86 (m, 2H), 3.65 (t, 2H), 3.43 (d, 1H), 3.11 (d, 1H), 1.70-1.55 (m, 2H), 1.36-1.22 (m, 4H), 0.85 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.2, 174.0, 148.0, 147.1, 143.4, 132.6, 131.4, 128.7, 124.4, 122.7, 119.8, 108.9, 108.2, 107.2, 101.2, 52.6, 41.5, 40.4, 29.0, 26.6, 22.3, 14.0; MS (ES+) *m/z* 404.0 (M + 23), 382.0 (M + 1).

PREPARATION 12

Synthesis of 3-[3-(1,3-benzodioxol-5-yl)-2-oxo-1-pentyl-2,3-dihydro-1*H*-indol-3-yl]propanoic acid

A. Synthesis of methyl 3-[3-(1,3-benzodioxol-5-yl)-2-oxo-1-pentyl-2,3-dihydro-1*H*-indol-3-yl]propanoate

Following the procedure as described in PREPARATION 11D, and making non-critical variations to replace methyl bromoacetate with methyl 3-bromopropionate, the title compound was obtained (76%) as a colorless oil: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.28 (td, 1H), 7.17 (dd, 1H), 7.06 (td, 1H), 6.89 (d, 1H), 6.84 (d, 1H), 6.77 (dd, 1H), 6.68 (d, 1H), 5.89-5.84 (m, 2H), 3.67 (t, 2H), 3.53 (s, 3H), 2.69-2.56 (m, 1H), 2.54-2.41 (m, 1H), 2.21-2.08 (m, 1H), 1.99-1.86 (m, 1H), 1.72-1.59 (m, 2H), 1.38-1.24 (m, 4H), 0.85 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.8, 173.1, 147.9, 146.9, 143.2, 133.5,

131.6, 128.5, 124.9, 122.6, 120.1, 108.7, 108.1, 107.6, 101.1, 55.2, 51.6, 40.2, 32.4, 29.5, 29.1, 27.1, 22.3, 14.0; MS (ES+) *m/z* 410.1 (M + 1), 432.0 (M + 23).

B. Synthesis of 3-[3-(1,3-benzodioxol-5-yl)-2-oxo-1-pentyl-2,3-dihydro-1*H*-indol-3-yl]propanoic acid

Following the procedure as described in PREPARATION 11E, and making non-critical variations to replace methyl [3-(1,3-benzodioxol-5-yl)-2-oxo-1-pentyl-2,3-dihydro-1*H*-indol-3-yl]acetate with methyl 3-[3-(1,3-benzodioxol-5-yl)-2-oxo-1-pentyl-2,3-dihydro-1*H*-indol-3-yl]propanoate, the title compound was obtained (92%) as a colorless solid: MS (ES-) *m/z* 394.2 (M - 1).

PREPARATION 13

Synthesis of 3-(4,5-difluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 3-(4,5-difluoro-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-pentyl-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 3, 4-difluorophenol, the title compound was obtained (31%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.69-9.65 (br, 1H), 7.51-7.41 (m, 2H), 7.26-7.21 (m, 1H), 6.99-6.57 (m, 3H), 4.18-4.14 (br, 1H), 3.78-3.58 (m, 2H), 1.76-1.62 (m, 2H), 1.40-1.28 (m, 4H), 0.87 (t, 3H); MS (ES+) *m/z* 330 (M - 17), 370 (M + 23).

B. Synthesis of 3-(4,5-difluoro-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 3-(4,5-difluoro-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (98%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.46-7.19 (m, 3H), 7.03-6.68 (m, 3H), 5.03 (s, 1H), 3.76-3.67 (m, 2H), 1.76-1.62 (m, 2H), 1.40-1.28 (m, 4H), 0.87 (t, 3H); MS (ES+) *m/z* 332 (M + 1).

C. Synthesis of 3-(4,5-difluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-

dihydro-2*H*-indol-2-one with 3-(4,5-difluoro-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (96%): MS (ES+) *m/z* 344 (M - 17), 384 (M + 23).

PREPARATION 14

Synthesis of 3-(5-fluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 3-(5-fluoro-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1C, and making non-critical variations to replace 4-bromo-1-pentyl-1*H*-indole-2,3-dione with 1-pentyl-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 4-fluorophenol, the title compound was obtained (53%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.42-9.14 (br, 1H), 7.53-6.86 (m, 6H), 6.56-6.48 (m, 1H), 4.58-4.28 (br, 1H), 3.79-3.58 (m, 2H), 1.77-1.61 (m, 2H), 1.41-1.24 (m, 4H), 0.87 (t, 3H); MS (ES+) *m/z* 312 (M - 17), 352 (M + 23).

B. Synthesis of 3-(5-fluoro-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

To a solution of 3-(5-fluoro-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one (2.42 g, 7.35 mmol) in dichloromethane (10.0 mL) were added trifluoroacetic acid (1.00 mL) and triethylsilane (1.00 mL) at ambient temperature. The reaction mixture was stirred at 40 °C for 15 hrs and concentrated *in vacuo* to dryness. The residue was triturated with ether to give the title compound (2.10 g, 91%) as a solid: MS (ES+) *m/z* 314 (M + 1).

C. Synthesis of 3-(5-fluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

To a solution of 3-(5-fluoro-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one (2.10 g, 6.70 mmol) in THF (20.0 mL) were added paraformaldehyde (1.76 g, 58.8 mmol) and lithium diisopropylamide (7.35 mL, 2.0 M in THF, 14.7 mmol) at 0 °C. The reaction mixture was stirred at 0 °C for 2 hrs followed by the addition of ammonium chloride solution (10.0 mL) and ethyl acetate (100 mL). The organic layer was washed with water and brine, dried over Na_2SO_4 and filtered. The filtrate was concentrated *in vacuo* to dryness to give the title compound: ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.55-9.10 (br, 1H), 7.53-6.86 (m, 6H), 6.57-6.49 (m, 1H), 4.74-4.30 (br, 1H), 4.18-4.07 (m, 2H), 3.79-3.60 (m, 2H), 1.77-1.61 (m, 2H), 1.41-1.24 (m, 4H), 0.87 (t, 3H); MS (ES+) *m/z* 326 (M - 17), 366 (M + 23).

PREPARATION 15

Synthesis of 3-(5-bromo-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 3-(5-bromo-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1C, and making non-critical variations to replace 4-bromo-1-pentyl-1*H*-indole-2,3-dione with 1-pentyl-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 4-bromophenol, the title compound was obtained (41%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.46-9.25 (br, 1H), 7.51-6.80 (m, 7H), 4.73-4.51(br, 1H), 3.79-3.56 (m, 2H), 1.76-1.60 (m, 2H), 1.41-1.22 (m, 4H), 0.87 (t, 3H); MS (ES+) *m/z* 377 (M - 17), 379 (M - 17), 412 (M + 23), 414 (M + 23).

B. Synthesis of 3-(5-bromo-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

To a solution of 3-(5-bromo-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one (2.22 g, 5.64 mmol) in dichloromethane (10.0 mL) were added trifluoroacetic acid (1.00 mL) and triethylsilane (1.00 mL) at ambient temperature. The reaction mixture was stirred at 50 °C for 15 hrs and concentrated *in vacuo* to dryness to give the title compound: MS (ES+) *m/z* 374 (M + 1), 376 (M + 1).

C. Synthesis of 3-(5-bromo-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 14C, and making non-critical variations to replace 3-(5-fluoro-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(5-bromo-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained: MS (ES+) *m/z* 386 (M - 17), 388 (M - 17), 426 (M + 23), 428 (M + 23).

PREPARATION 16

Synthesis of 3-(5-chloro-4-fluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 3-(5-chloro-4-fluoro-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1C, and making non-critical variations to replace 4-bromo-1-pentyl-1*H*-indole-2,3-dione with 1-pentyl-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 4-chloro-3-fluorophenol, the title compound was obtained (33%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.80 (s, 1H), 7.52-7.41 (m, 2H), 7.23 (t, 1H), 6.96 (d, 1H), 6.84 (d, 1H), 6.80 (d, 1H), 4.15 (s, 1H), 3.79-3.58

(m, 2H), 1.76-1.62 (m, 2H), 1.40-1.28 (m, 4H), 0.87 (t, 3H); MS (ES+) *m/z* 346 (M - 17), 386 (M + 23).

B. Synthesis of 3-(5-chloro-4-fluoro-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 15B, and making non-critical variations to replace 3-(5-bromo-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(5-chloro-4-fluoro-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (99%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 10.0-9.70 (br, 1H), 7.45-7.18 (m, 3H), 6.98 (d, 1H), 6.90-6.82 (m, 2H), 5.01 (s, 1H), 3.75-3.66 (m, 2H), 1.76-1.62 (m, 2H), 1.40-1.28 (m, 4H), 0.87 (t, 3H); MS (ES+) *m/z* 348 (M + 1).

C. Synthesis of 3-(5-chloro-4-fluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(5-chloro-4-fluoro-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (46%): MS (ES+) *m/z* 360 (M - 17), 400 (M + 23).

PREPARATION 17

Synthesis of 3-(4-chloro-5-fluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 3-(4-chloro-5-fluoro-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1C, and making non-critical variations to replace 4-bromo-1-pentyl-1*H*-indole-2,3-dione with 1-pentyl-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 3-chloro-4-fluorophenol, the title compound was obtained (14%): MS (ES+) *m/z* 346 (M - 17), 386 (M + 23).

B. Synthesis 3-(4-chloro-5-fluoro-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 15B, and making non-critical variations to replace 3-(5-bromo-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(4-chloro-5-fluoro-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained: MS (ES+) *m/z* 348 (M + 1).

C. Synthesis of 3-(4-chloro-5-fluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(4-chloro-5-fluoro-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (50% for two steps): MS (ES+) *m/z* 360 (M - 17), 400 (M + 23).

PREPARATION 18

Synthesis of 3-(4,5-dichloro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 3-(4,5-dichloro-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1C, and making non-critical variations to replace 4-bromo-1-pentyl-1*H*-indole-2,3-dione with 1-pentyl-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 3,4-dichlorophenol, the title compound was obtained (26%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.60 (s, 1H), 7.50-7.40 (m, 2H), 7.22 (td, 1H), 7.11 (s, 1H), 6.95 (d, 1H), 6.86 (s, 1H), 4.31-4.12 (br, 1H), 3.79-3.59 (m, 2H), 1.76-1.62 (m, 2H), 1.40-1.27 (m, 4H), 0.88 (t, 3H); MS (ES+) *m/z* 363 (M - 17), 403 (M + 23).

B. Synthesis of 3-(4,5-dichloro-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 15B, and making non-critical variations to replace 3-(5-bromo-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(4,5-dichloro-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (86%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 10.0-9.50 (br, 1H), 7.42 (t, 1H), 7.32 (d, 1H), 7.22 (td, 1H), 7.09 (s, 1H), 6.95 (d, 1H), 6.93 (s, 1H), 5.04(s, 1H), 3.77-3.68 (m, 2H), 1.77-1.62 (m, 2H), 1.40-1.27 (m, 4H), 0.88 (t, 3H); MS (ES+) *m/z* 348 (M + 1).

C. Synthesis of 3-(4,5-dichloro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(4,5-dichloro-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained: MS (ES+) *m/z* 376 (M - 17), 416 (M +

23).

PREPARATION 19

Synthesis of 3-(hydroxymethyl)-3-[2-hydroxy-5-(trifluoromethyl)phenyl]-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 3-hydroxy-3-[2-hydroxy-5-(trifluoromethyl)phenyl]-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1C, and making non-critical variations to replace 4-bromo-1-pentyl-1*H*-indole-2,3-dione with 1-pentyl-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with α, α, α-trifluorocresol, the title compound was obtained (46%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.75 (s, 1H), 7.50-7.39 (m, 3H), 7.21 (td, 1H), 7.10-7.02 (m, 2H), 6.96 (d, 1H), 4.26 (s, 1H), 3.82-3.59 (m, 2H), 1.77-1.63 (m, 2H), 1.40-1.27 (m, 4H), 0.88 (t, 3H); MS (ES+) *m/z* 362 (M - 17), 402 (M + 23).

B. Synthesis of 3-[2-hydroxy-5-(trifluoromethyl)phenyl]-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 15B, and making non-critical variations to replace 3-(5-bromo-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-hydroxy-3-[2-hydroxy-5-(trifluoromethyl)phenyl]-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (78%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.20-8.00 (br, 1H), 7.43-7.14 (m, 5H), 7.02 (d, 1H), 6.95 (d, 1H), 5.11(s, 1H), 3.82-3.72 (m, 2H), 1.79-1.66 (m, 2H), 1.40-1.27 (m, 4H), 0.88 (t, 3H); MS (ES+) *m/z* 364 (M + 1).

C. Synthesis of 3-(hydroxymethyl)-3-[2-hydroxy-5-(trifluoromethyl)phenyl]-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-[2-hydroxy-5-(trifluoromethyl)phenyl]-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained: MS (ES+) *m/z* 376 (M - 17), 416 (M + 23).

PREPARATION 20

Synthesis of 3-(2-hydroxy-4-methoxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 3-(5-bromo-2-hydroxy-4-methoxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1C, and making non-critical variations to replace 4-bromo-1-pentyl-1*H*-indole-2,3-dione with 1-pentyl-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 4-bromo-3-methoxyphenol, the title compound was obtained (48%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.85 (s, 1H), 7.52-7.38 (m, 2H), 7.22 (td, 1H), 6.94 (d, 1H), 6.89 (s, 1H), 6.63 (s, 1H), 4.13-4.03 (br, 1H), 3.86 (s, 3H), 3.80-3.57 (m, 2H), 1.75-1.63 (m, 2H), 1.40-1.25 (m, 4H), 0.88 (t, 3H); MS (ES+) *m/z* 402 (M - 17), 404 (M - 17), 442 (M + 23), 444 (M + 23).

B. Synthesis of 3-(2-hydroxy-4-methoxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 15B, and making non-critical variations to replace 3-(5-bromo-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(5-bromo-2-hydroxy-4-methoxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (83%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.78-9.20 (br, 1H), 7.43-7.31 (m, 2H), 7.19 (t, 1H), 6.97 (d, 1H), 6.79 (d, 1H), 6.70-6.64 (m, 1H), 6.38 (dd, 1H), 5.02 (s, 1H), 3.77 (s, 3H), 3.70 (t, 2H), 1.75-1.63 (m, 2H), 1.40-1.25 (m, 4H), 0.87 (t, 3H); MS (ES+) *m/z* 326 (M + 1).

C. Synthesis of 3-(2-hydroxy-4-methoxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(2-hydroxy-4-methoxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (41%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 10.79 (s, 1H), 7.51-7.37 (m, 2H), 7.26 (td, 1H), 6.99 (d, 1H), 6.95 (d, 1H), 6.59 (d, 1H), 6.34 (dd, 1H), 4.67 (d, 1H), 4.14 (d, 1H), 3.76 (s, 3H), 3.78-3.69 (m, 2H), 1.75-1.63 (m, 2H), 1.40-1.25 (m, 4H), 0.87 (t, 3H); MS (ES+) *m/z* 338 (M - 17), 378 (M + 23).

PREPARATION 21

Synthesis of ethyl [3-(6-hydroxy-2,3-dihydro-1*H*-inden-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

A. Synthesis of ethyl [3-hydroxy-3-(6-hydroxy-2,3-dihydro-1*H*-inden-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1C, and making non-critical variations to replace 4-bromo-1-pentyl-1*H*-indole-2,3-dione with ethyl (2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate, and 1,3-benzodioxol-5-ol with 5-indanol, the title compound was obtained (84%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.76 (s, 1H), 7.55 (d, 1H), 7.38 (td, 1H), 7.20 (t, 1H), 6.9 (s, 1H), 6.80 (d, 1H), 6.65 (s, 1H), 4.45 (ABq, 2H), 4.32-4.25 (br, 1H), 4.20 (q, 2H), 2.83 (t, 2H), 2.74-2.65 (m, 2H), 2.06-1.94 (m, 2H), 1.27 (t, 3H); MS (ES+) *m/z* 350 (M - 17), 390 (M + 23).

B. Synthesis of ethyl [3-(6-hydroxy-2,3-dihydro-1*H*-inden-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 15B, and making non-critical variations to replace 3-(5-bromo-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one with ethyl [3-hydroxy-3-(6-hydroxy-2,3-dihydro-1*H*-inden-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.50-7.90 (br, 1H), 7.40-7.32 (m, 2H), 7.38 (td, 1H), 6.94 (s, 1H), 6.84 (d, 1H), 6.75 (s, 1H), 5.16 (s, 1H), 4.48 (ABq, 2H), 4.21 (q, 2H), 2.85 (t, 2H), 2.81-2.61 (m, 2H), 2.09-1.92 (m, 2H), 1.25 (t, 3H); MS (ES+) *m/z* 352 (M + 1).

C. Synthesis of ethyl [3-(6-hydroxy-2,3-dihydro-1*H*-inden-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with ethyl [3-(6-hydroxy-2,3-dihydro-1*H*-inden-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained: MS (ES+) *m/z* 364 (M - 17), 404 (M + 23).

PREPARATION 22

Synthesis of ethyl [3-(hydroxymethyl)-3-(3-hydroxy-5,6,7,8-tetrahydronaphthalen-2-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

A. Synthesis of ethyl [3-hydroxy-3-(3-hydroxy-5,6,7,8-tetrahydronaphthalen-2-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1C, and making non-

critical variations to replace 4-bromo-1-pentyl-1*H*-indole-2,3-dione with ethyl (2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate, and 1,3-benzodioxol-5-ol with 5, 6, 7, 8-tetrahydronapthalen-2-ol, the title compound was obtained (81%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.61 (s, 1H), 7.54 (dd, 1H), 7.38 (td, 1H), 7.20 (t, 1H), 6.80 (d, 1H), 6.76 (s, 1H), 6.50 (s, 1H), 4.45 (ABq, 2H), 4.21 (q, 2H), 4.18-4.14 (br, 1H), 2.73-2.47 (m, 4H), 1.77-1.63 (m, 4H), 1.24 (t, 3H); MS (ES+) *m/z* 364 (M - 17), 404 (M + 23)

B. Synthesis of ethyl [3-(3-hydroxy-5,6,7,8-tetrahydronaphthalen-2-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 15B, and making non-critical variations to replace 3-(5-bromo-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one with ethyl [3-hydroxy-3-(3-hydroxy-5,6,7,8-tetrahydronaphthalen-2-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.42-7.32 (m, 2H), 7.20 (t, 1H), 6.84 (d, 1H), 6.78 (s, 1H), 6.61 (s, 1H), 5.12 (s, 1H), 4.47 (ABq, 2H), 4.21 (q, 2H), 2.76-2.44 (m, 4H), 1.78-1.64 (m, 4H), 1.24 (t, 3H); MS (ES+) *m/z* 366 (M + 1).

C. Synthesis of ethyl [3-(hydroxymethyl)-3-(3-hydroxy-5,6,7,8-tetrahydronaphthalen-2-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with ethyl [3-(3-hydroxy-5,6,7,8-tetrahydronaphthalen-2-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained: MS (ES+) *m/z* 378 (M-17), 418 (M + 23).

PREPARATION 23

Synthesis of ethyl [4-bromo-3-(4,5-difluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

A. Synthesis of ethyl (4-bromo-2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace isatin with 4-bromoisatin and (2-bromoethyl)cyclopropane with ethyl bromoacetate, the title compound was obtained as a yellow solid (68%):^{1}H NMR (300 MHz, $CDCl_3$) δ 7.39 (t, 1H), 7.27(dd, 1H), 6.71 (dd, 1H), 4.47 (s, 2H), 4.23 (q, 2H), 1.27 (t, 3H); MS (ES+) *m/z* 312 (M + 1), 314 (M + 1), 334 (M + 23), 336 (M + 23).

B. Synthesis of ethyl [4-bromo-3-(4,5-difluoro-2-hydroxyphenyl)-3-hydroxy-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with ethyl (4-bromo-2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate and 1,3-benzodioxol-5-ol with 3,4-difluorophenol, the title compound was obtained as a white solid (42%); MS (ES+) *m/z* 424 (M - 17), 426 (M - 17), 464 (M + 23), 466 (M + 23).

C. Synthesis of ethyl [4-bromo-3-(4,5-difluoro-2-hydroxyphenyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

A mixture of ethyl [4-bromo-3-(4,5-difluoro-2-hydroxyphenyl)-3-hydroxy-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate (0.90 g, 2.00 mmol), triethylsilane (2.00 mL, 12.2 mmol) and trifluroacetic acid (0.94 mL, 12.2 mmol) was heated at 90 °C for two days. After cooling down to ambient temperature, the mixture was diluted with ethyl acetate (200 mL), washed with water, dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography (ethyl acetate/hexanes, 1/3) to give the title compound (0.37 g, 43%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.35-7.22 (m, 3H), 6.82-6.71 (m, 2H), 6.52 (t, 1H), 5.10 (s, 1H), 4.45 (s, 2H), 4.21 (q, 2H), 1.23 (t, 3H); MS (ES+) *m/z* 426.4 (M + 1), 428.4 (M + 1).

D. Synthesis of ethyl [4-bromo-3-(4,5-difluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with ethyl [4-bromo-3-(4,5-difluoro-2-hydroxyphenyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained (83%): MS (ES+) *m/z* 456.3 (M + 1), 458.3 (M + 1).

PREPARATION 24

Synthesis of ethyl [4-bromo-3-(6-hydroxy-2,2-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

A. Synthesis of 6-(benzyloxy)-2,2-dimethylbenzofuran-3(2*H*)-one

To a solution of 6-(benzyloxy)benzofuran-3(2*H*)-one (Adams, J. L., et al., *J. Med. Chem.* (1996), 39(26):5035-46) (1.60 g, 6.67 mmol) in DMF (50.0 mL) were added sodium hydride (0.59 g, 14.7 mmol) and iodomethane (1.46 mL, 23.3 mmol) at 0 °C. The reaction mixture was stirred at ambient temperature for 16 h and quenched

with saturated ammonium chloride (50.0 mL). The aqueous mixture was extracted with ethyl acetate (3 x 50.0 mL). The combined organic layers was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/10) to give the title compound (0.85 g, 47%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.55 (d, 1H), 7.44-7.30 (m, 5H), 6.69 (dd, 1H), 6.54 (d, 1H), 5.10 (s, 2H), 1.43 (s, 6H); MS (ES+) *m/z* 269.5 (M + 1).

B. Synthesis of 2,2-dimethyl-2,3-dihydrobenzofuran-6-ol

To a solution of 6-(benzyloxy)-2,2-dimethylbenzofuran-3(2*H*)-one (0.85 g, 3.20 mmol) in methanol (100 mL) was added palladium hydroxide (0.22 g, 20 wt. % loading, 0.32 mmol). The resulting mixture was hydrogenated for 16 hours under 60 psi of hydrogen. The reaction mixture was filtered through celite*, washed with methanol. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/5) to give the title compound (0.46 g, 88%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 6.92 (d, 1H), 6.30-6.21 (m, 2H), 4.77 (s, 1H), 2.90 (s, 2H), 1.44 (s, 6H).

C. Synthesis of ethyl [4-bromo-3-hydroxy-3-(6-hydroxy-2,2-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with ethyl (4-bromo-2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate and 1,3-benzodioxol-5-ol with 2,2-dimethyl-2,3-dihydrobenzofuran-6-ol, the title compound was obtained: MS (ES+) *m/z* 498.5 (M + 23), 500.5 (M + 23).

D. Synthesis of ethyl [4-bromo-3-(6-hydroxy-2,2-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

A mixture of ethyl [4-bromo-3-hydroxy-3-(6-hydroxy-2,2-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate (1.32 g, 2.80 mmol), triethylsilane (2.00 mL, 12.2 mmol) and trifluoroacetic acid (0.94 mL, 12.2 mmol) in dichloromethane (50.0 mL) was stirred at 35 °C for 3 hours. The mixture was diluted with dichloromethane (100 mL), washed with water, dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/3) to give the title compound (1.04 g, 81%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.32-7.15 (m, 2H), 6.74 (d, 1H), 6.50-6.36 (br, 2H), 5.04 (s, 1H), 4.51-4.34 (m, 2H), 4.25-4.14 (m, 2H), 2.92-2.69 (m,

*Trademark

2H), 1.43 (s, 3H), 1.37 (s, 3H), 1.23 (t, 3H); MS (ES+) *m/z* 460.5 (M + 1), 462.5 (M + 1).

E. Synthesis of ethyl [4-bromo-3-(6-hydroxy-2,2-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with ethyl [4-bromo-3-(6-hydroxy-2,2-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained (25%): MS (ES+) *m/z* 490.5 (M + 1), 492.5 (M + 1).

PREPARATION 25

Synthesis of 3,3-dimethyl-2,3-dihydro-1-benzofuran-6-ol

A. Synthesis of 4-(benzyloxy)-1-bromo-2-(2-methylallyloxy)benzene

To a solution of 5-(benzyloxy)-2-bromophenol (Simas, A.B.C., et al, *Synthesis*, (1999):1017-21) (8.15 g, 29.3 m mol) in DMF (150 mL) was added potassium carbonate (4.46 g, 32.2 mmol) slowly at 0 °C. The mixture was stirred at ambient temperature for half an hour, followed by the addition of 3-bromo-2-methylpropene (3.35 mL, 32.2 mmol) during half an hour at 0 °C. The mixture was stirred at ambient temperature overnight and quenched with saturated ammonium chloride (50 mL). The aqueous mixture was extracted with ethyl acetate (3 x 200 mL). The combined organic layers was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo*. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/20) to give the title compound (10.0 g, 94%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.43-7.29 (m, 5H), 6.53 (d, 1H), 6.45 (dd, 1H), 5.15-4.94 (m, 4H), 4.43 (s, 2H), 1.82 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 159.1, 155.6, 140.1, 136.5, 133.1, 128.6, 128.1, 127.5, 112.9, 107.2, 103.3, 102.0, 72.4, 70.3, 19.3.

B. Synthesis of 6-(benzyloxy)-3,3-dimethyl-2,3-dihydrobenzofuran

To a solution of 4-(benzyloxy)-1-bromo-2-(2-methylallyloxy)benzene (5.00 g, 15.1 mmol) in benzene (400 mL) was added tributyltin hydride (7.42 mL, 27.2 mmol) and benzoyl peroxide (0.70 g, 2.90 mmol) at 0 °C. The resulting mixture was refluxed at 100 °C overnight. After cooling down to ambient temperature, the mixture was washed with water, dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/20) to give the title compound (3.48 g, 91%): MS (ES+) *m/z* 255.6 (M + 1).

C. Synthesis of 3,3-dimethyl-2,3-dihydro-1-benzofuran-6-ol

To a solution of 6-(benzyloxy)-3,3-dimethyl-2,3-dihydrobenzofuran (3.48 g, 13.7 mmol) in methanol (200 mL) was added Pd/C (1.45 g) and the mixture was hydrogenated under 40 psi of hydrogen overnight. The reaction mixture was filtered through celite, washed with methanol. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/7) to give the title compound (1.66 g, 74%): MS (ES+) *m/z* 165.4 (M + 1).

PREPARATION 26

Synthesis of 1-(diphenylmethyl)-3-(5-hydroxy-2,3-dihydro-1-benzofuran-6-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 1-(diphenylmethyl)-3-hydroxy-3-(5-hydroxy-2,3-dihydro-1-benzofuran-6-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1,3-benzodioxol-5-ol with 2,3-dihydrobenzofuran-5-ol, (Alabaster, R.J., et al.; *Synthesis* (1988), 12:950-2) and 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-(diphenylmethyl)-1*H*-indole-2,3-dione, the title compound was obtained: MS (ES+) *m/z* 472.2 (M + 23).

B. Synthesis of 1-(diphenylmethyl)-3-(5-hydroxy-2,3-dihydro-1-benzofuran-6-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 1-(diphenylmethyl)-3-hydroxy-3-(5-hydroxy-2,3-dihydro-1- benzofuran-6-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained: MS (ES+) *m/z* 434.4 (M + 1).

C. Synthesis of 1-(diphenylmethyl)-3-(5-hydroxy-2,3-dihydro-1-benzofuran-6-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

To a solution of 1-(diphenylmethyl)-3-(5-hydroxy-2,3-dihydro-1-benzofuran-6-yl)-1,3-dihydro-2*H*-indol-2-one (1.01 g, 2.30 mmol) in THF (50.0 mL) was added paraformaldehyde (1.00 g, 30.0 mmol). Argon was bubbled through the reaction mixture for one hour followed by the addition of diisopropylamine (1.00 g, 10.0 mmol) at 0 °C. The reaction mixture was stirred at ambient temperature for 20 hours and diluted with ethyl acetate (100 mL). The resulting mixture was washed with water (2 x 50.0 mL), dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness to give 0.67 g (65%) of the title compound: MS (ES+) *m/z* 486.4 (M

+ 23).

PREPARATION 27

Synthesis of 3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-4-methoxy-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1.3-dihydro-2*H*-indol-2-one

A. Synthesis of ethyl 2-(2-(*tert*-butoxycarbonylamino)-6-methoxyphenyl)-2-oxoacetate

To a solution of *tert*-butyl 3-methoxyphenylcarbamate (25.6 g, 0.11 mol) in THF (300 mL) was added n-BuLi (0.25 mol, 1.6 M solution in pentane) at -78 °C. The resulted solution was stirred at 0 °C for 3 hours and re-cooled to -78 °C followed by the addition of diethyl oxalate (20.1 g, 0.14 mol). The mixture was stirred at -78 °C for 45 min and at ambient temperature for one hour, and quenched with 1 N HCl. The mixture was extracted with ether. The organic solution was dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography to give 3.70 g (27% based on recovered starting material) of the title compound: MS (ES+) *m/z* 324.3 (M + 1).

B. Synthesis of 4-methoxy-1*H*-indole-2,3-dione

A mixture of ethyl 2-(2-(tert-butoxycarbonylamino)-6-methoxyphenyl)-2-oxoacetate (3.70 g, 110 mmol) and 10% H_2SO_4 (100 mL) was heated at 100 °C for 10 hours. After cooling down to ambient temperature, the reaction mixture was extracted with ether (3 x 100 mL). The combined ether solution was washed with water (2 x 50.0 mL), dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography to give 0.37 g (19%) of the title compound: MS (ES+) *m/z* 200.1 (M + 23).

C. Synthesis of 4-methoxy-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1*H*-indole-2,3-dione

Following the procedure as described in PREPARATION 1A, and making non-critical variations to replace 4-bromoindole with 4-methoxy-1*H*-indole-2,3-dione, and 1-bromopentane with 2-(bromomethyl)-5-(trifloromethyl)furan, the title compound was obtained (26%): MS (ES+) *m/z* 348.2 (M + 23).

D. Synthesis of 3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-4-methoxy-1-{[5-(trifluoromethyl)-2-furil]methyl}-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 4-methoxy-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1*H*-indole-2,3-dione, the title compound was

obtained (56%): MS (ES+) *m/z* 486.4 (M + 23).

E. Synthesis of 3-(6-hydroxy-1,3-benzodioxol-5-yl)-4-methoxy-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-4-methoxy-1-{[5-(trifluoromethyl)-2-furil]methyl}-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (86%): MS (ES+) *m/z* 448.4 (M + 1).

F. Synthesis of 3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-4-methoxy-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1.3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 26C, and making non-critical variations to replace 1-(diphenylmethyl)-3-(5-hydroxy-2,3-dihydro-1-benzofuran-6-yl)-1,3-dihydro-2*H*-indol-2-one with 3-(6-hydroxy-1,3-benzodioxol-5-yl)-4-methoxy-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (64%): MS (ES+) *m/z* 500.4 (M + 23).

PREPARATION 28

Synthesis of 4,7-dichloro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 4,7-dichloro-1-pentyl-1*H*-indole-2,3-dione

To a mixture of sodium hydride (0.17 g, 6.94 mmol, 60% dispersion in mineral oil) in anhydrous *N*,*N*-dimethylformamide (5.00 mL) was added a solution of 4,7-dichloro-1*H*-indole-2,3-dione (1.00 g, 4.60 mmol) in *N*,*N*-dimethylformamide (5.00 mL) at 0 ^{0}C. The brown reaction mixture was stirred for 0.5 h followed by the addition of a solution of 1-bromopentane (0.84 g, 5.55 mmol) in anhydrous *N*,*N*-dimethylformamide (5.00 mL). The reaction mixture was stirred at ambient temperature for 16 h and poured into wet ethyl ether (30.0 mL). After the organic layer was separated, it was washed with water (2 x 20.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The gummy residue was dried under vacuum and the solid was triturated with ether to give the title compound (0.98 g, 98%): MS (ES+) *m/z* 286.2 (M + 1).

B. Synthesis of 4,7-dichloro-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 4,7-

dichloro-1-pentyl-1*H*-indole-2,3-dione, the title compound was obtained (68%) as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.61 (s, br, 1H), 7.26 (t, 1H), 7.03 (d, 1H), 6.52 (s, 1H), 6.12 (s, 1H), 5.86 (dd, 2H), 4.21 (s, br, 1H), 4.01-3.96 (m, 2H), 1.73-1.58 (m, 2H), 1.34-1.21 (m, 4H), 0.84 (t, 3H); MS (ES+) *m/z* 408.2 (M - 17).

C. Synthesis of 4,7-dichloro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 4,7-dichloro-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (72%) as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.27-7.23 (m, 1H), 7.03 (d, 1H), 6.55 (s, 1H), 6.04 (s, 1H), 5.84 (dd, 2H), 5.03 (s, 1H), 4.09-3.99 (m, 2H), 1.72-1.62 (m, 2H), 1.33-1.24 (m, 4H), 0.86 (t, 3H); MS (ES+) *m/z* 409.2 (M + 1).

D. Synthesis of 4,7-dichloro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 4,7-dichloro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (94%) as a gummy solid: MS (ES+) *m/z* 439.3 (M + 1).

PREPARATION 29

Synthesis of ethyl 2-(4-chloro-3-(6-hydroxy-2,3-dihydrobenzofuran-5-yl)-3-(hydroxymethyl)-2-oxoindolin-1-yl)acetate

A. Synthesis of ethyl (4-chloro-2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace isatin with 4-chloro-1*H*-indole-2,3-dione, and (2-bromoethyl)cyclopropane with ethyl bromoacetate, the title compound was obtained (95%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.48 (t, 1H), 7.08 (d, 1H), 6.67 (d, 1H), 4.47 (s, 2H), 4.23 (q, 2H), 1.27 (t, 3H); MS (ES+) *m/z* 268.6 (M + 1).

B. Synthesis of ethyl [4-chloro-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with ethyl (4-chloro-2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate, the title compound was obtained

(75%) as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.70 (br, 1H), 7.31 (t, 1H), 7.12 (d, 1H), 6.68 (d, 1H), 6.46 (d, 2H), 4.53-4.46 (m, 2H), 4.18 (q, 2H), 3.08-2.88 (m, 2H), 1.23 (t, 3H); MS (ES+) *m/z* 387.8 (M - 17).

C. Synthesis ethyl [4-chloro-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with ethyl [4-chloro-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained (75%) as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.33-7.27 (m, 2H), 7.12 (d, 1H), 6.71 (d, 1H), 6.50-6.48 (m, 1H), 5.10 (s, 1H), 4.54-4.42 (m, 4H), 4.19 (q, 2H), 3.11-2.90 (m, 2H), 1.23 (t, 3H); MS (ES+) *m/z* 388.8 (M + 1).

D. Synthesis of ethyl [4-chloro-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with ethyl [4-chloro-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained (99%) as a gummy solid: MS (ES+) *m/z* 418.7 (M + 1).

PREPARATION 30

Synthesis of 3-hydroxy-3-[6-(hydroxymethyl)-1,3-benzodioxol-5-yl]-1-pentyl-1,3-dihydro-2*H*-indol-2-one

To a solution of (6-bromo-1,3-benzodioxol-5-yl)methanol (Mann, J., et al, *J. Chem. Soc. Perkin Trans. 1* (1984):2081-8) (1.27 g, 5.50 mmol) in THF (45.0 mL) was added n-BuLi (5.00 mL, 2.0 M, 10.0 mmol) dropwise at -75 °C. The reaction mixture was stirred at -75 °C for 45 min followed by the addition of a solution of 1-pentyl-1*H*-indole-2,3-dione (1.00 g, 4.60 mmol) in THF (20.0 mL) at -75 °C. The resulting mixture was stirred at ambient temperature for 12 hrs and quenched with ammonium chloride solution (5.00 mL). More ethyl acetate and water were added and separated. The organic layer was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with 50% EtOAc:Hexanes to yield the title compound (0.29 g, 25%) as a solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.38-7.24 (m, 2H), 7.11 (t, 1H), 6.91 (d, 1H), 6.81(s, 1H), 6.43 (s, 1H), 5.90-5.87 (m, 2H), 4.77 (dd, 2H), 3.75-3.56 (m, 2H), 1.75-1.58 (m, 2H), 1.26-1.35 (m, 2H), 0.89-0.83 (m, 3H); ^{13}C NMR (75 MHz,

$CDCl_3$) δ 177.8, 147.4, 147.2, 142.8, 133.5, 132.2, 131.1, 130.1, 125.3, 123.8, 111.4, 109.2, 108.1, 101.5, 79.5, 64.7, 40.4, 29.0, 26.8, 22.3, 13.9; MS (ES+) *m/z* 352.1 (M - 17).

PREPARATION 31

Synthesis of ethyl [1-hexyl-3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-3-yl]acetate

A. Synthesis of 1-hexyl-1*H*-indole-2,3-dione

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace (2-bromoethyl)cyclopropane with n-bromohexane, the title compound was obtained (90%) as a viscous gum: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.58-7.51 (m, 2H), 7.08 (t, 1H), 6.87 (d, 1H), 3.68 (t, 2H), 1.71-1.62 (m, 2H), 1.41-1.22 (m, 6H), 0.85 (t, 3H).

B. Synthesis of 1-hexyl-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-hexyl-1*H*-indole-2,3-dione, the title compound was obtained (53%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.44 (br, 1H), 7.47-7.44 (m, 1H), 7.40-7.34 (m, 1H), 7.17 (t, 1H), 6.89 (d, 1H), 6.55 (s, 1H), 6.21 (s, 1H), 5.84-5.82 (m, 2H), 4.58 (br, 1H), 3.71-3.56 (m, 2H), 1.67-1.62 (m, 2H), 1.32-1.21 (m, 6H), 0.84-0.80 (m, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 179.0, 152.3, 148.8, 142.5, 141.3, 130.3, 129.2, 126.1, 123.7, 117.2, 109.5, 106.8, 101.9, 101.4, 79.2, 40.4, 31.3, 27.1, 26.4, 22.4, 13.9; MS (ES+) *m/z* 352.5 (M - 17).

C. Synthesis of 1-hexyl-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 1-hexyl-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (98%) as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.38-7.13 (m, 3H), 6.94 (d, 1H), 6.60 (s, 1H), 6.32 (s, 1H), 5.84 (dd, 2H), 5.02 (s, 1H), 3.74-3.63 (m, 2H), 1.70-1.61 (m, 2H), 1.37-1.19 (m, 6H), 0.83 (t, 3H); MS (ES+) *m/z* 354.2 (M + 1).

D. Synthesis of ethyl [1-hexyl-3-(6-hydroxy-1,3-benzodioxol-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-3-yl]acetate

To a solution of diisopropylamine (1.14 g, 11.0 mmol) in THF (10.0 mL) was added n-butyl lithium (7.00 mL, 11.0 mmol, 1.6 M solution in hexane) at -75 °C. The resulting mixture was stirred at -75 °C for half an hour and added slowly to a solution of 1-hexyl-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one in THF (20.0 mL) at -75 °C. After stirring at -75 °C for another half an hour, ethyl bromoacetate was added. The mixture was stirred at ambient temperature for 18 hrs and quenched with saturated ammonim chloride solution. The organic solvent was removed *in vacuo* and the aqueous residue was diluted with ethyl acetate (100 mL). The orgnic layer was washed with saturated ammonium chloride (25.0 mL), brine (50.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with 40% EtOAc/Hexanes to yield the title compound (0.19 g, 8%) as an oil: MS (ES+) *m/z* 440.5 (M + 1).

PREPARATION 32

Synthesis of 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 4-bromo-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 4-bromoisatin, the title compound was obtained (95%) as a beige solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.40 (s, 1H), 9.09 (s, 1H), 7.22 (s, 1H), 7.04 (t, 1H), 6.90 (d, 1H), 6.75 (d, 1H), 6.43 (br, 1H), 6.21 (s, 1H), 5.88 (d, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 178.0, 148.7, 147.0, 145.8, 139.5, 131.3, 130.8, 125.4, 118.8, 118.4, 109.4, 108.9, 101.0, 97.4, 76.6; MS (ES+) *m/z* 366.4 (M + 1), 364.5 (M + 1).

B. Synthesis of 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1D, and making non-critical variations to replace 4-bromo-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 4-bromo-3hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (95%) as a cream solid: MS (ES+) *m/z* 348.5 (M + 1), 346.3 (M + 1).

C. Synthesis of 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 31D, and making non-critical variations to replace 1-hexyl-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one, and ethyl bromoacetate with *para*-formaldehyde, the title compound was obtained (70%) as a colorless solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.00 (br, 1H), 7.13-6.95 (m, 3H), 6.84 (d, 1H), 6.16 (d, 1H), 5.90-5.84 (m, 2H), 5.16-4.83 (m, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 177.8, 150.4, 147.1, 146.8, 139.8, 130.2, 129.3, 125.8, 117.7, 115.8, 109.3, 107.9, 101.2, 97.6, 63.5, 57.4.

PREPARATION 33

Synthesis of 4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 4-bromo-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 4-bromoisatin, and 1,3-benzodioxol-5-ol with 2,3-dihydrobenzofuran-6-ol, the title compound was obtained (78%) as a colorless solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.36 (s, 1H), 9.15 (s, 1H), 7.49 (1H), 7.04 (t, 1H), 6.89 (d, 1H), 6.74 (d, 1H), 6.35 (br, 1H), 5.90 (s, 1H), 4.45 (t, 2H), 3.05 (t, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 178.4, 160.2, 154.0, 145.7, 131.6, 130.7, 125.5, 125.4, 118.9, 117.7, 116.1, 108.8, 96.8, 76.9, 71.8, 29.1; MS (ES-) *m/z* 344.4 (M - 17), 360.4 (M - 1).

B. Synthesis of4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1D, and making non-critical variations to replace 4-bromo-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 4-bromo-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (62%) as a solid: MS (ES+) *m/z* 346.5 (M + 1), 348.5 (M + 1).

C. Synthesis of 4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 31D, and making non-critical variations to replace 1-hexyl-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-

indol-2-one with 4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one, and ethyl bromoacetate with *para*-formaldehyde, the title compound was obtained that was used directly for further reaction.

PREPARATION 34

Synthesis of 4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1-(pyridin-2-ylmethyl)-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 4-bromo-1-(pyridin-2-ylmethyl)-1*H*-indole-2,3-dione

To a solution of 4-bromoisatin (8.94 g, 39.5 mmol) in anhydrous *N*,*N*-dimethylformamide (100 mL) was added sodium hydride (3.34 g, 86.9 mmol, 60% dispersion in mineral oil) in portions at 0 °C. The brown reaction mixture was stirred for 30 min followed by the addition of a solution of 2-(bromomethyl)pyridine hydrobromide (10.0 g, 39.5 mmol) neutralized with sodium hydride (1.52 g, 39.5 mmol, 60% dispersion in mineral oil) in *N*,*N*-dimethylformamide at 0 °C. The reaction mixture was stirred for 16 h and quenched with water (100 mL). The reaction mixture was extracted with diethyl ether (3 x 100 mL) and the aqueous layer was extracted with ethyl acetate (3 x 100 mL). The combined organic layers was washed with water (5 x 200 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was triturated with ether to afford the title compound (10.6 g, 85%) as a brown solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.53 (d, 1H), 7.67 (t, 1H), 7.30 (t, 2H), 7.25-7.19 (m, 2H), 6.94 (d, 1H), 5.04 (s, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 180.5, 157.3, 154.2, 152.3, 149.5, 138.4, 137.5, 128.6, 123.3, 122.3, 121.5, 116.4, 110.3, 45.8.

B. Synthesis of 4-bromo-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1-(pyridin-2-ylmethyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 4-bromo-1-(pyridin-2-ylmethyl)-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 2,3-dihydrobenzofuran-6-ol, the title compound was obtained (91%) as a colorless solid: mp >225 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.29 (s, 1H), 8.54 (d, 1H), 7.70 (dt, 1H), 7.61 (br, 1H), 7.32-7.26 (m, 2H), 7.07 (d, 1H), 7.00 (d, 1H), 6.72 (d, 1H), 6.60 (br, 1H), 6.02 (s, 1H), 4.91 (ABq, 2H), 4.47 (t, 2H), 3.06 (d, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 176.9, 160.4, 156.3, 153.8, 149.6, 146.1, 137.5, 130.9, 130.8, 126.5, 125.8, 123.1, 121.5, 118.8, 117.3, 116.4, 108.3, 96.7, 76.6, 71.9, 45.7, 29.1; MS (ES+) *m/z* 455.4 (M + 1), 437.4 (M - 17).

C. Synthesis of 4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1-(pyridin-2-ylmethyl)-1,3-dihydro-2*H*-indol-2-one

To a solution of 4-bromo-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1-(pyridin-2-ylmethyl)-1,3-dihydro-2*H*-indol-2-one (1.12 g, 2.48 mmol) in anhydrous dichloromethane (25.0 mL) was added triethylamine (1.40 mL, 9.91 mmol) and $SOCl_2$ (0.40 mL, 4.96 mmol) at 0 °C. The reaction mixture was stirred at 0 °C for 2 h and quenched with water (30.0 mL). The organic layer was separated, washed with water (3 x 30.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness to give a gummy material. The residue was dissolved in acetic acid/tetrahydrofuran (3.00 mL/22.0 mL) followed by the addition of zinc dust (0.81 g, 12.4 mmol) in one portion. The reaction mixture was stirred at ambient temperature for 16 h. After the solid was filtered, the solvent was removed *in vacuo*. The residue was dissolved in ethyl acetate (100 mL), washed with water (3 x 30.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness to give the title compound (1.50 g, 77%) as a gummy material: MS (ES+) *m/z* 437.3 (M + 1).

D. Synthesis of 4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1-(pyridin-2-ylmethyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1-(pyridin-2-ylmethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (34%): MS (ES+) *m/z* 468.4 (M + 1).

PREPARATION 35

Synthesis of 5-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 5-fluoro-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1*H*-indole-2,3-dione

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace isatin with 5-fluoroisatin, and (2-bromoethyl)cyclopropane with 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (59%) as a red solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.54-7.50 (m, 1H), 7.47-7.44 (m, 1H), 7.20 (dd, 1H), 7.14-7.13 (m, 1H), 6.75 (d, 1H), 4.99 (s, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 182.4 (d), 160.7, 158.5 (d), 157.5, 153.0 (d), 146.5 (d), 139.9 (q), 124.3, 119.3 (d), 114.5 (d), 112.7 (d), 112.0 (d), 110.5, 36.8.

B. Synthesis of 5-fluoro-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 5-fluoro-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1*H*-indole-2,3-dione, the title compound was obtained (66%) as a pale yellow solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.15 (s, 1H), 7.21 (s,1H), 7.15 (dd, 1H), 7.08-6.95 (m, 2H), 6.74 (s, 1H), 6.54 (s, 1H), 6.22 (d, 1H), 5.90 (d, 2H), 4.96 (s, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 176.7, 160.57, 157.4, 154.0, 148.6, 147.4, 140.1 (m), 139.6 (m), 134.7 (d, $^2J_{CF}$ = 29.4 Hz), 121.3, 119.5, 117.7, 115.1 (d, $^1J_{CF}$ = 92.1 Hz), 114.5, 111.8 (d, $^1J_{CF}$ = 97.5 Hz), 109.7, 109.6, 107.2, 101.3, 97.8, 75.1, 36.9; MS (ES+) *m/z* 450.3 (M + 1)

C. Synthesis of 5-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1D, and making non-critical variations to replace 4-bromo-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 5-fluoro-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (72%) as a pale yellow solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.31 (s, 1H), 7.13 (dd, 1H), 7.02 (dd, 2H), 6.82 (d, 1H), 6.59 (d, 2H), 6.39 (s, 1H), 5.87 (d, 2H), 5.07-4.96 (m, 2H), 4.84 (s, 1H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 176.1, 160.5, 157.4, 153.9, 150.5, 147.5, 140.2, 139.6, 139.1, 132.3 (d, $^2J_{CF}$ = 33.3 Hz), 115.3, 114.5 (m), 114.2, 113.9, 111.9 (d, $^1J_{CF}$ = 98.7 Hz), 109.9, 109.7 (d, $^2J_{CF}$ = 32.7 Hz), 101.3, 98.3, 48.5, 36.8; MS (ES+) *m/z* 436.2 (M + 1).

D. Synthesis of 5-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one

A mixture of 5-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one (3.64 g, 8.41 mmol), *para*-formaldehyde (2.52 g, 84.1 mmol) and lithium hydroxide monohydrate (1.06 g, 25.2 mmol) in tetrahydrofuran (84.0 mL) and water (10.0 mL) was stirred at 0 °C for 4 h. After the solvent was removed *in vacuo*, the residue was dissolved in ethyl acetate (100 mL), washed with 10% aqueous HCl (3 x 25.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate/hexanes (50%) to give the title compound (0.65 g, 59%) as a colorless solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ

9.11 (s, 1H), 7.12 (d, 1H), 6.99-6.87 (m, 3H), 6.80 (dd, 1H), 6.48 (d, 1H), 6.23 (s, 1H), 5.89 (d, 2H), 5.09 (br, 1H), 4.97 (ABq, 2H), 4.01(ABq, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 176.1, 160.5; MS (ES+) *m/z* 466.2 (M + 1), 448.2 (M - 17).

PREPARATION 36

Synthesis of *tert*-butyl-(2-chloromethyl-5-trifluoromethylthiophen-3-yloxy)dimethylsilane

A. Synthesis of methyl 3-*tert*-butyl dimethylsilanyloxy-5-trifluoromethyl-2-thiophenecarboxylate

To a solution of methyl 3-hydroxy-5-trifluoromethyl-2-thiophenecarboxylate (Karp, G.M., et al, *Synthesis* (2000), 8:1078-1080) (19.4 g, 85.8 mmol) in *N,N*-dimethylformamide (50.0 mL) was added imidazole (8.77 g, 129 mmol) followed by *tert*-butyl dimethylsilyl chloride (19.4 g, 129 mmol) at 0 °C. The reaction mixture was stirred at ambient temperature overnight. More imidazole (7.50 g) and *tert*-butyl dimethylsilyl chloride (10.5 g) were added. The reaction mixture was stirred for another 4 h and quenched with water (100 mL). The reaction mixture was extracted with ether (3 x 500 mL). The combined organic layers was washed with water (3 x 500 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate/hexanes (1/9) to give the titlew compound (26.5 g, 90%) as a yellow oil: ^{1}H NMR (300 MHz, $CDCl_3$) δ 6.91 (s, 1H), 3.82 (s, 3H), 0.21 (s, 6H), 0.06 (s, 9H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 161.5, 155.5, 133.7, 133.2, 123.6 (q, $^1J_{CF}$ = 14.4 Hz), 119.8, 51.9, 25.4, 18.2, -4.6.

B. Synthesis of [3-(*tert*-butyldimethylsilanyloxy)-5-trifluoromethylthiophen-2-yl]methanol

To a mixture of lithium aluminum hydride (1.67 g, 43.9 mmol) in anhydrous ether (75.0 mL) was added a solution of methyl 3-*tert*-butyl dimethylsilanyloxy-5-trifluoromethyl-2-thiophenecarboxylate (10.0 g, 29.3 mmol) in anhydrous ether (25.0 mL) at 0 °C. The reaction mixture was stirred at 0 °C for 30 min and quenched by the slow addition of water (50.0 mL). After the aqueous layer was separated, the organic layer was washed with saturated ammonium chloride (3 x 20.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate/hexanes (1/9) to give the title compound (6.95 g, 76%) as a yellow oil: ^{1}H NMR (300 MHz, $CDCl_3$) δ 6.89 (s, 1H), 4.68 (s, 2H), 2.11 (br, 1H), 0.96 (s, 9H), 0.88 (s, 6H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 148.0, 127.0, 124.0, 122.4 (q, $^1J_{CF}$ = 14.7 Hz), 120.5,

56.1, 25.5, 18.1, -4.6.

C. Synthesis of *tert*-butyl-(2-chloromethyl-5-trifluoromethylthiophen-3-yloxy)dimethylsilane

To a solution of [3-(*tert*-butyldimethylsilanyloxy)-5-trifluoromethylthiophen-2-yl]methanol in anhydrous dichloromethane (100 mL) was added triethylamine (4.05 g, 40.0 mmol) followed by thionyl chloride (2.38 g, 20.0 mmol) at 0 °C. The reaction mixture was stirred for 30 min and quenched with water (50.0 mL). After separation, the organic layer was washed with water (3 x 50.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with hexane to afford the title compound (2.31 g, 70%) as a yellow oil, which was directly used.

PREPARATION 37

Synthesis of 1-(diphenylmethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-5-methyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 1-(diphenylmethyl)-5-methyl-1*H*-indole-2,3-dione

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace isatin with 5-methylisatin, and (2-bromoethyl)cyclopropane with 1,1'-(bromomethylene)dibenzene, the title compound was obtained (74%) as a bright orange solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.42-7.26 (m, 11H), 7.09 (d, 1H), 6.95 (s, 1H), 6.37 (d, 1H), 2.24 (s, 3H).

B. Synthesis of 1-(diphenylmethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-5-methyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-(diphenylmethyl)-5-methyl-1*H*-indole-2,3-dione, the title compound was obtained (92%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.23 (br, 1H), 7.40-7.15 (m, 11H), 6.90-6.85 (m, 2H), 6.57 (s, 1H), 6.33 (d, 1H), 6.31 (s, 1H), 5.87 (s, 2H), 4.46 (br s, 1H), 2.28 (s, 3H); MS (ES+) *m/z* 448.4 (M - 17).

C. Synthesis of 1-(diphenylmethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-5-methyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 1-(diphenylmethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-5-methyl-1,3-dihydro-2*H*-indol-2-one, the title compound

was obtained (84%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.37-7.25 (m, 9H), 7.22-7.17 (m, 2H), 7.10 (s, 1H), 6.91 (s, 1H), 6.86 (d, 1H), 6.63 (s, 1H), 6.40 (s, 1H), 6.38 (d, 1H), 5.88 (ABq, 2H), 5.07 (s, 1H), 2.23 (s, 3H); MS (ES+) *m/z* 450.3 (M + 1).

D. Synthesis of 1-(diphenylmethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-5-methyl-1,3-dihydro-2*H*-indol-2-one

To a solution of 1-(diphenylmethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-5-methyl-1,3-dihydro-2*H*-indol-2-one (1.61 g, 3.60 mmol) and *para*-formaldehyde (0.43 g, 14.6 mmol) in dichloromethane (60.0 mL) was added diisopropylamine (7.20 mmol). After stirring at ambient temperature for 3 h, the reaction was quenched with saturated aqueous ammonium chloride (60.0 mL). The organic layer was separated and washed with water (3 x 100 mL), dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate/hexanes (20-60%) to afford the title compound (1.07 g, 63%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 10.09 (br, 1H), 7.37-7.16 (m, 12H), 6.99 (s, 1H), 6.87 (d, 1H), 6.62 (s, 1H) 6.54 (s, 1H), 6.37 (d, 1H), 5.87 (d, 2H), 4.45 (ABq, 2H), 2.33 (s, 3H); MS (ES+) *m/z* 480.4 (M + 1).

PREPARATION 38

Synthesis of 3-(hydroxymethyl)-3-(5-hydroxy-2-methyl-1,3-benzothiazol-6-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 3-hydroxy-3-(5-hydroxy-2-methyl-1,3-benzothiazol-6-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-pentyl-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 2-methyl-1,3-benzothiazol-5-ol, the title compound was obtained (81%) as a colorless solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.90 (br, 1H), 9.05 (br, 1H), 7.78 (d, 1H), 7.25 (dd, 1H), 7.10-6.95 (m, 2H), 6.90-6.80 (m, 2H), 3.81-3.58 (m, 2H), 2.75 (br, 3H), 1.80-1.60 (m, 2H), 1.50-1.31 (m, 4H), 0.90 (t, 3H); MS (ES+) *m/z* 383.4 (M + 1).

B. Synthesis of 3-(5-hydroxy-2-methyl-1,3-benzothiazol-6-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A suspension of 3-hydroxy-3-(5-hydroxy-2-methyl-1,3-benzothiazol-6-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one (0.50 g, 1.31 mmol) in hydroiodic acid (10.0 mL) was refluxed for 1.5 days. The reaction mixture was concentrated *in vacuo* to dryness.

The residue was used directly in next step.

C. Synthesis of 3-(hydroxymethyl)-3-(5-hydroxy-2-methyl-1,3-benzothiazol-6-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(5-hydroxy-2-methyl-1,3-benzothiazol-6-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained: MS (ES+) *m/z* 367.5 (M + 1).

PREPARATION 39

Synthesis of (5-chloro-1,3,4-thiadiazol-2-yl)methanol

To a a solution of ethyl 5-chloro-1,3,4-thiadiazole-2-carboxylate (0.51 g, 2.60 mmol) in anhydrous methanol (5.00 mL) was added sodium borohydride (0.30 g, 7.99 mmol) at 0 °C. The reaction mixture was stirred at ambient temperature for 16 h, diluted with acetic acid (3.00 mL) and extracted with ethyl acetate (2 x 150 mL). The combined organics was washed with aqueous saturated sodium bicarbonate (3 x 25.0 mL) and aqueous saturated sodium chloride (2 x 25.0 mL). The organic layer was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness to give the title compound (0.30 g, 75%) as a light yellow semi-solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 5.04 (s, 2H), 2.80 (br, 1H); MS (ES+) 151.1 (M + 1), 153.1 (M + 1).

PREPARATION 40

Synthesis of 1-(diphenylmethyl)-3-(6-hydroxy-3,3-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 1-(diphenylmethyl)-3-hydroxy-3-(6-hydroxy-3,3-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-(diphenylmethyl)-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 3,3-dimethyl-2,3-dihydro-1-benzofuran-6-ol, the title compound was obtained: MS (ES+) *m/z* 478.5 (M + 1).

B. Synthesis of 1-(diphenylmethyl)-3-(6-hydroxy-3,3-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and makingnon-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-

benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 1-(diphenylmethyl)-3-hydroxy-3-(6-hydroxy-3,3-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (73% for two steps): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.38-7.20 (m, 12H), 7.11-7.04 (m, 2H), 6.97 (s, 1H), 6.58 (s, 1H), 6.57-6.51 (m, 1H), 6.50 (s, 1H), 5.08 (s, 1H), 4.19 (s, 2H), 1.25 (s, 3H), 1.18 (s, 3H); MS (ES+) *m/z* 426.6 (M + 1).

C. Synthesis of 1-(diphenylmethyl)-3-(6-hydroxy-3,3-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 26C, and making non-critical variations to replace 1-(diphenylmethyl)-3-(5-hydroxy-2,3-dihydro-1-benzofuran-6-yl)-1,3-dihydro-2*H*-indol-2-one with 1-(diphenylmethyl)-3-(6-hydroxy-3,3-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained: MS (ES+) *m/z* 492.5 (M + 1)

PREPARATION 41

Synthesis of 7-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 7-fluoro-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 7-fluoro-1*H*-indole-2,3-dione, the title compound was obtained (80%): ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.66 (s, 1H), 9.11 (s, 1H), 7.18 (s, 1H), 7.07-6.98 (m, 1H), 6.83-6.74 (m, 1H), 6.66 (d, 1H), 6.48 (s, 1H), 6.18 (s, 1H), 5.92-5.85 (m, 2H); MS (ES+) *m/z* 304.5 (M + 1).

B. Synthesis of 7-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 7-fluoro-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (100%): ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.84 (s, 1H), 9.22 (s, 1H), 7.01 (t, 1H), 6.87-6.78 (m, 1H), 6.71 (d, 1H), 6.62 (s, 1H), 6.35 (s, 1H), 5.90-5.85 (m, 2H), 4.67 (s, 1H); MS (ES+) *m/z* 288.5 (M + 1).

PREPARATION 42

Synthesis of ethyl [4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

A. Synthesis of ethyl [4-bromo-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with ethyl (4-bromo-2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate, and 1,3-benzodioxol-5-ol with 2,3-dihydrobenzofuran-6-ol, the title compound was obtained (68%) as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.66 (br, 1H), 7.31-7.19 (m, 3H), 6.73 (dd, 1H), 6.49-6.45 (m, 1H), 5.09-4.36 (m, 4H), 4.20 (q, 2H), 3.14-2.90 (m, 2H), 1.23 (t, 3H); MS (ES+) *m/z* 432.2 (M - 17).

B. Synthesis of ethyl [4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with ethyl [4-bromo-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained (81%) as a white solid: ^{1}H NMR ($CDCl_3$, 300 MHz) δ 7.31-7.19 (m, 3H), 6.75 (d, 1H), 6.50-6.45 (m, 1H), 5.08 (s, 1H), 5.09-4.36 (m, 4H), 4.20 (q, 2H), 3.14-2.90 (m, 2H), 1.23 (t, 3H); MS (ES+) *m/z* 433.3 (M + 1).

C. Synthesis of ethyl [4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with ethyl [4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained (99%): MS (ES+) *m/z* 463.2 (M + 1).

PREPARATION 43

Synthesis of ethyl [5-chloro-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

A. Synthesis of ethyl (5-chloro-2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace isatin with 5-chloro-1*H*-indole-2,3-dione, and (2-

bromoethyl)cyclopropane with ethyl 2-bromoacetate, the title compound was obtained (98%) as solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.60 (d, 1H), 7.54 (dd, 1H), 6.74 (d, 1H), 4.46 (s, 2H), 4.23 (q, 2H), 1.27 (t, 3H); MS (ES+) *m/z* 268.6 (M + 1).

B. Synthesis of ethyl [5-chloro-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1C, and making non-critical variations to replace 4-bromo-1-pentyl-1*H*-indole-2,3-dione with ethyl (5-chloro-2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate, and 1,3-benzodioxol-5-ol with 2,3-dihydrobenzofuran-6-ol, the title compound was obtained (85%) as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.70 (br, 1H), 7.31-7.24 (m, 2H), 6.92 (d, 1H), 6.68 (s, 1H), 6.46 (s, 1H), 4.53-4.46 (m, 2H), 5.09-4.40 (d, 2H), 4.18 (q, 2H), 3.08-2.88 (m, 2H), 1.23 (t, 3H); MS (ES+) *m/z* 387.8 (M - 17).

C. Synthesis of ethyl [5-chloro-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1D, and making non-critical variations to replace 4-bromo-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with ethyl [5-chloro-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained (94%) as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.30-7.24 (m, 2H), 6.72 (d, 1H), 6.66 (s, 1H), 6.39 (s, 1H), 5.05 (s, 1H), 4.53-4.46 (m, 4H), 4.21 (q, 2H), 3.14-2.94 (m, 2H), 1.25 (t, 3H); MS (ES+) *m/z* 388.8 (M + 1).

D. Synthesis of ethyl [5-chloro-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with ethyl [5-chloro-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained (99%): MS (ES+) *m/z* 418.7 (M + 1).

PREPARATION 44

Synthesis of methyl [3-(4-chloro-2-hydroxyphenyl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

A. Synthesis of methyl (2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace (2-bromoethyl)cyclopropane with methyl bromoacetate, the

title compound was obtained (72%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.64-7.53 (m, 2H), 7.14 (t, 1H), 6.77 (d, 1H), 4.48 (s, 2H), 3.76 (s, 3H); MS (ES+) *m/z* 220.4 (M + 1).

B. Synthesis of methyl [3-(4-chloro-2-hydroxyphenyl)-3-hydroxy-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1C, and making non-critical variations to replace 4-bromo-1-pentyl-1*H*-indole-2,3-dione with methyl (2,3-dioxo-2,3-dihydro-1*H*-indol-1-yl)acetate, and 1,3-benzodioxol-5-ol with 3-chlorophenol, the title compound was obtained (29%) as a yellow solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.10 (s, 1H), 7.48 (d, 1H), 7.38 (t, 1H), 7.19 (t, 1H), 7.01 (br, 1H), 6.80-6.64 (m, 3H), 5.28 (br s, 1H), 4.51 (d, 1H), 4.44 (d, 1H), 3.75 (s, 3H); MS (ES+) *m/z* 370.5 (M + 23), 372.4 (M + 23).

C. Synthesis of methyl [3-(4-chloro-2-hydroxyphenyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with methyl [3-(4-chloro-2-hydroxyphenyl)-3-hydroxy-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained (83%) as a semi-solid; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.36 (t, 1H), 7.29 (bd, 1H), 7.18 (t, 1H), 6.95 (br, 1H), 6.86-6.78 (m, 3H), 5.13 (br, 1H), 4.55 (d, 1H), 4.45 (d, 1H), 3.75 (s, 3H); MS (ES+) *m/z* 332.5 (M + 1), 334.5 (M + 1).

D. Synthesis of methyl [3-(4-chloro-2-hydroxyphenyl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with methyl [3-(4-chloro-2-hydroxyphenyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained: MS (ES+) *m/z* 362.5 (M + 1) 364.5 (M + 1).

PREPARATION 45

Synthesis of ethyl [3-(4,5-difluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

A. Synthesis of ethyl [3-(4,5-difluoro-2-hydroxyphenyl)-3-hydroxy-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 2B, and non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with ethyl (2,3-dioxo-

2,3-dihydro-1*H*-indol-1-yl)acetate, the title compound was obtained as a brown oil: MS (ES+) *m/z* 364.3 (M + 1), 348.5 (M - 17).

B. Synthesis of ethyl [3-(4,5-difluoro-2-hydroxyphenyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with ethyl [3-(4,5-difluoro-2-hydroxyphenyl)-3-hydroxy-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained (83%) as a light yellow oil: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.39 (t, 1H), 7.34 (d, 1H), 7.26-7.22 (m, 1H), 6.92-6.82 (m, 2H), 6.73 (dd, 1H), 5.11 (br, 1H), 4.50 (d, 1H), 4.43 (d, 1H), 4.21 (q, 2H), 1.23 (t, 3H); MS (ES+) *m/z* 448.5 (M + 1).

C. Synthesis of ethyl [3-(4,5-difluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with methyl [3-(4-chloro-2-hydroxyphenyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained: MS (ES+) *m/z* 378.3 (M + 1), 361.3 (M - 17).

PREPARATION 46

Synthesis of 3-(4-bromo-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 3-(4-bromo-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1C, and making non-critical variations to replace 4-bromo-1-pentyl-1*H*-indole-2,3-dione with 1-pentyl-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 3-bromophenol, the title compound was obtained (48%) as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.66 (br, 1H), 7.50-7.38 (m, 2H), 7.24-7.16 (m, 2H), 6.98-6.86 (m, 2H), 6.64 (d, 1H), 4.15 (br, 1H), 3.80-3.55 (m, 2H), 1.75-1.62 (m, 2H), 1.40-1.34 (m, 4H), 0.89 (t, 3H); MS (ES+) *m/z* 391.4 (M + 1), 393.4 (M + 1).

B. Synthesis of 3-(4-bromo-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1D, and making non-critical variations to replace 4-bromo-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(4-bromo-2-hydroxyphenyl)-3-hydroxy-1-

pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (91%) as a white powder: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.40 (t, 1H), 7.31 (d, 1H) 7.24-7.23 (m, 2H), 7.01-6.91 (m, 2H), 6.74 (d, 1H), 5.05 (br, 1H), 3.80-3.65 (m, 2H), 1.75-1.63 (m, 2H), 1.38-1.29 (m, 4H), 0.88 (t, 3H); MS (ES+) *m/z* 374.4 (M + 1), 376.4 (M + 1).

C. Synthesis of 3-(4-bromo-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(4-bromo-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained: R_f = 0.5 (EtOAc/Hexanes, ¼).

PREPARATION 47

Synthesis of 3-(5-bromo-2-hydroxyphenyl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 3-(5-bromo-2-hydroxyphenyl)-3-hydroxy-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with isatin, 1,3-benzodioxol-5-ol with 4-bromophenol, the title compound was obtained (71%) as a yellowish solid: MS (ES+) *m/z* 319.4 (M + 1), 321.4 (M + 1).

B. Synthesis of 3-(5-bromo-2-hydroxyphenyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1D, and making non-critical variations to replace 4-bromo-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(5-bromo-2-hydroxyphenyl)-3-hydroxy-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (98%) as a white powder: MS (ES+) *m/z* 306.2 (M + 1), 304.2 (M+1).

C. Synthesis of 3-(5-bromo-2-hydroxyphenyl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 35D, and making non-critical variations to replace 5-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one with 3-(5-bromo-2-hydroxyphenyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained: MS (ES+) *m/z* 334.2 (M + 1), 336.2 (M + 1).

PREPARATION 48

Synthesis of 1-(diphenylmethyl)-3-(hydroxymethyl)-3-[2-hydroxy-4-(trifluoromethoxy)phenyl]-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 1-(diphenylmethyl)-3-hydroxy-3-[2-hydroxy-4-(trifluoromethoxy)phenyl]-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-(diphenylmethyl)-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 3-(trifluoromethoxy)phenol, the title compound was obtained (75%): MS (ES+) *m/z* 514.5 (M + 23).

B. Synthesis of 1-(diphenylmethyl)-3-[2-hydroxy-4-(trifluoromethoxy)phenyl]-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 1-(diphenylmethyl)-3-hydroxy-3-[2-hydroxy-4-(trifluoromethoxy)phenyl]-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (82%): MS (ES+) *m/z* 498.4 (M + 23).

C. Synthesis of 1-(diphenylmethyl)-3-(hydroxymethyl)-3-[2-hydroxy-4-(trifluoromethoxy)phenyl]-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 1-(diphenylmethyl)-3-[2-hydroxy-4-(trifluoromethoxy)phenyl]-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained: MS (ES+) *m/z* 488 (M - 17), 528 (M + 23).

PREPARATION 49

Synthesis of 1-(diphenylmethyl)-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 1-(diphenylmethyl)-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1,3-benzodioxol-5-ol with 2,3-dihydrobenzofuran-6-ol (Foster et al., *J. Chem. Soc.* 1948:2254-2258) and 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-(diphenylmethyl)-1*H*-indole-2,3-dione, the title compound was obtained (68%) as a white solid: MS (ES+) *m/z* 450.4 (M + 1).

B. Synthesis of 1-(diphenylmethyl)-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-criticial variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 1-(diphenylmethyl)-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (67%) as a white solid: MS (ES+) *m/z* 434.3 (M + 1).

C. Synthesis of 1-(diphenylmethyl)-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-(3-hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 26C, and making non-critical variations to replace 1-(diphenylmethyl)-3-(5-hydroxy-2,3-dihydro-1-benzofuran-6-yl)-1,3-dihydro-2*H*-indol-2-one with 1-(diphenylmethyl)-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (45%) as a white solid: MS (ES+) *m/z* 464.5 (M + 1).

PREPARATION 50

Synthesis of 4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 4-bromo-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace 1,3-benzodioxol-5-ol with 2,3-dihydrobenzofuran-6-ol and 1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 4-bromo-1*H*-indole-2,3-dione, the title compound was obtained (78%) as a white solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.36 (s, 1H), 9.15 (s, 1H), 7.49 (1H), 7.04 (t, 1H), 6.89 (d, 1H), 6.74 (d, 1H), 6.35 (br, 1H), 5.90 (s, 1H), 4.45 (t, 2H), 3.05 (t, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 178.4, 160.2, 154.0, 145.7, 131.6, 130.7, 125.5, 125.4, 118.9, 117.7, 116.1, 108.8, 96.8, 76.9, 71.8, 29.1; MS (ES-) *m/z* 344.4 (M - 17), 360.4 (M - 1).

B. Synthesis of 4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 4-bromo-3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (62%) as a white solid: MS (ES+) *m/z* 346.5 (M + 1), 348.5 (M + 1).

C. Synthesis of 4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 14C, and making non-critical variations to replace 3-(5-fluoro-2-hydroxyphenyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (16%): R_f = 0.21 (EtOAc/Hexanes, 7/3).

PREPARATION 51

Synthesis of 7-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 7-fluoro-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1*H*-indole-2,3-dione

Following the procedure as described in PREPARATION 1A, and making tnon-critical variations to replace 4-bromoindole with 7-fluoroisatin, and 1-bromopentane with 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (34%): MS (ES+) *m/z* 336.2 (M + 23).

B. Synthesis of 7-fluoro-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 7-fluoro-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1*H*-indole-2,3-dione, the title compound was obtained (75%): MS (ES+) *m/z* 474.3 (M + 23).

C. Synthesis of 7-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 7-fluoro-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (65%): MS (ES+) *m/z* 436.4 (M + 1).

D. Synthesis of 7-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 26C, and making non-critical variations to replace 1-(diphenylmethyl)-3-(5-hydroxy-2,3-dihydro-1-benzofuran-6-yl)-1,3-dihydro-2H-indol-2-one with 7-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (67%): MS (ES+) *m/z* 488.4 (M + 23).

PREPARATION 52

Synthesis of 3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-pentyl-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 2,3-dihydrobenzofuran-6-ol, the title compound was obtained (90%) as a white powder: MS (ES+) *m/z* 376.3 (M + 23).

B. Synthesis of 3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 3-hydroxy-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (76%): MS (ES+) *m/z* 338.3 (M + 1).

C. Synthesis of 3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (46%): MS (ES+) *m/z* 368.3 (M + 1), 380.4 (M + 23).

PREPARATION 53

Synthesis of 3-(5-bromo-2-hydroxyphenyl)-1-(diphenylmethyl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 3-(5-bromo-2-hydroxyphenyl)-1-(diphenylmethyl)-3-hydroxy-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-(diphenylmethyl)-1*H*-indole-2,3-dione, and 1,3-benzodioxol-5-ol with 4-bromophenol, the title compound was obtained (90%) as an orange solid: MS (ES+) *m/z* 486.2 (M + 1), 488.2 (M + 1).

B. Synthesis of 3-(5-bromo-2-hydroxyphenyl)-1-(diphenylmethyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 15B, and making non-critical variations to replace 3-(5-bromo-2-hydroxyphenyl)-3-hydroxy-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(5-bromo-2-hydroxyphenyl)-1-(diphenylmethyl)-3-hydroxy-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (99%) as a white powder: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.39-7.20 (m, 11H), 7.11-7.06 (m, 4H), 6.82 (d, 1H), 6.57-6.51 (m, 1H), 5.04 (s, 1H); MS (ES+) *m/z* 471.2 (M + 1), 473.2 (M + 1).

C. Synthesis of 3-(5-bromo-2-hydroxyphenyl)-1-(diphenylmethyl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 26C, and making non-critical variations to replace 1-(diphenylmethyl)-3-(5-hydroxy-2,3-dihydro-1-benzofuran-6-yl)-1,3-dihydro-2*H*-indol-2-one with 3-(5-bromo-2-hydroxyphenyl)-1-(diphenylmethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained: MS (ES+) *m/z* 500.4 (M + 1), 502.4 (M + 1).

PREPARATION 54

Synthesis of 3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

A. Synthesis of 1-pentyl-1*H*-indole-2,3-dione

Following the procedure as described in PREPARATION 2A, and making non-critical variations to replace (2-bromoethyl)cyclopropane with 1-bromopentane, the title compound was obtained (72%) as a red solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.90 (d, 1H), 7.53-7.45 (m, 1H), 7.03-6.97 (m, 1H), 6.82 (d, 1H), 3.64-3.57 (m, 2H), 1.68-1.52 (m, 2H), 1.34-1.21 (m, 4H), 0.79 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 183.6, 158.1, 151.0, 138.4, 125.3, 123.5, 117.5, 110.2, 40.2, 28.9, 26.9, 22.2, 13.9.

B. Synthesis of 3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2B, and making non-critical variations to replace1-(2-cyclopropylethyl)-1*H*-indole-2,3-dione with 1-pentyl-1*H*-indole-2,3-dione, the title compound was obtained (47%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.41 (s, 1H), 7.46 (dd, 1H), 7.37 (dt, 1H), 7.16 (dt, 1H), 6.89 (d, 1H), 6.53 (s, 1H), 6.22 (s, 1H), 5.83 (dd, 2H), 4.70 (br, 1H), 3.73-3.54 (m, 2H), 1.69-1.60 (m, 2H), 1.34-1.26 (m, 4H), 0.85 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.9, 152.2, 148.8, 142.5, 141.3, 130.3, 129.3, 126.1, 123.8, 117.1, 109.5, 106.8, 101.8,

101.4, 79.3, 40.4, 28.9, 26.8, 22.3, 13.9; MS (ES+1) *m/z* 355.5 (M + 1).

C. Synthesis of 3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 2C, and making non-critical variations to replace 1-(2-cyclopropylethyl)-3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one with 3-hydroxy-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (81%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.67 (br, 1H), 7.39-7.29 (m, 2H), 7.18-7.13 (m, 1H), 6.94 (d, 1H), 6.62 (s, 1H), 6.32 (s, 1H), 5.84 (dd, 2H), 5.01 (s, 1H), 3.71-3.63 (m, 2H), 1.71-1.61 (m, 2H), 1.35-1.27 (m, 4H), 0.86 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.8, 151.3, 147.6, 143.9, 141.53, 128.7, 126.4, 126.2, 123.1, 115.3, 109.4, 106.5, 101.5, 101.2, 47.4, 40.5, 28.9, 26.9, 22.3, 13.9; MS (ES+) *m/z* 340 (M + 1).

D. Synthesis of 3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one

Following the procedure as described in PREPARATION 1E, and making non-critical variations to replace 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one with 3-(6-hydroxy-1,3-benzodioxol-5-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (67%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 10.85-10.63 (br, 1H), 7.48-7.35 (m, 2H), 7.28-7.19 (m, 1H), 6.96 (d, 1H), 6.52 (d, 2H), 5.82 (dd, 2H), 4.63 (d, 1H), 4.11 (d, 1H), 3.70 (d, 2H), 2.04-1.74 (br, 1H), 1.65-162 (m, 2H), 1.31-1.24 (m, 4H), 0.84 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 180.3, 152.6, 148.1, 143.2, 141.3, 129.2, 129.1, 126.2, 123.3, 112.4, 109.6, 108.2, 101.9, 101.3, 64.6, 59.8, 40.6, 28.9, 26.9, 22.2, 13.9; MS (ES+) *m/z* 370.1 (M + 1).

EXAMPLE 1

Synthesis of 1'-(2-cyclopropylethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a solution of 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-

(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one (0.92 g, 2.51 mmol) in anhydrous THF (20.0 mL) was added triphenylphosphine (0.82 g, 3.13 mmol) and diethyl azodicarboxylate (0.55, 3.13 mmol) at –78 °C. The brown reaction solution was stirred at ambient temperature for 16 h, and quenched with saturated ammonium chloride (50.0 mL). The organic solvent was removed under reduced pressure and the aqueous mixture was extracted with ethyl acetate (3 x 50.0 mL). The combined organic layers were dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The brown residue was subjected to column chromatography eluting with ethyl acetate/hexane (5% to 20%, gradient) to give the title compound (0.63 g, 72%) which was crystallized from ether to afford a colorless solid: mp 125-127 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.30-7.25 (m, 1H), 7.14 (d, 1H), 7.02 (t, 1H), 6.89 (d, 1H), 6.49 (s, 1H), 6.11 (s, 1H), 5.84 (m, 2H), 4.76 (m, 2H), 3.93-3.74 (m, 2H), 1.65-1.57 (m, 2H), 0.76-0.56 (m, 1H), 0.48-0.41 (m, 2H), 0.08-0.03 (m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.4, 155.9, 148.8, 142.6, 142.3, 132.4, 128.8, 124.0, 123.1, 119.5, 108.6, 103.1, 101.5, 93.6, 80.6, 58.2, 40.5, 32.5, 30.8, 8.7, 4.4; MS (ES+) *m/z* 350.3 (M + 1).

EXAMPLE 1.1

Synthesis of 1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in Example 1, and making non-critical variations using 3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (80%) as a white solid: mp 85-87 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.28 (t, 1H), 7.15 (d, 1H), 7.02 (t, 1H), 6.89 (d, 1H), 6.49 (s, 1H), 6.11 (s, 1H), 5.84 (dd, 2H), 4.77 (ABq, 2H), 3.85-3.62 (m, 2H), 1.76-1.66 (m, 2H), 1.40-1.33 (m, 4H), 0.89 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.3, 155.9, 148.8, 142.4, 142.3, 132.5, 128.9, 123.9, 119.6, 108.6, 103.0, 101.5, 93.6, 80.5, 58.2, 40.4, 29.0, 27.1, 22.3, 14.0; MS (ES+) *m/z* 352 (M + 1).

EXAMPLE 1.2

Synthesis of 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1, and making non-critical variations using 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.16 (d, 1H), 7.15 (s, 1H), 6.84 (dd, 1H), 6.45 (s, 1H), 6.06 (s, 1H), 5.86 (dd, 2H), 4.90 (ABq, 2H), 3.83-3.60 (m, 2H), 1.74-1.64 (m, 2H), 1.39-1.28 (m, 4H), 0.89 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.1, 157.2, 149.1, 144.6, 142.0, 130.3, 130.1, 127.0, 120.0, 116.5, 107.6, 102.5, 101.5, 93.3, 77.3, 59.6, 40.6, 29.0, 27.0, 22.3, 14.0; MS (ES+) *m/z* 430 (M + 1).

EXAMPLE 1.3

Synthesis of ethyl (2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetate



Following the procedure as described in EXAMPLE 1, and making non-critical variations using ethyl [3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained as a white powder in 90% yield: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.31-7.26 (m, 1H), 7.17-7.00 (m, 3H), 6.67 (s, 1H), 6.18 (s, 1H), 5.90-5.89 (m, 2H), 4.76-4.66 (m, 2H), 4.59 (s, 2H), 4.13 (q, 2H), 1.17 (t, 3H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 177.3, 168.3, 155.6, 148.8, 142.4, 142.1, 132.0, 129.3, 124.1, 123.7, 120.4, 109.6, 103.3, 101.9, 93.8, 79.8, 61.8, 57.8, 41.8, 14.5; MS (ES+) *m/z* 390.2 (M + 23).

EXAMPLE 1.4

Synthesis of methyl 2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoate



Following the procedure as described in EXAMPLE 1, and making non-critical variations using methyl 2-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained as a white powder in 74% yield: mp 166-167 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.05 (m, 1H), 7.44 (m, 1H), 7.34 (t, 1H), 7.22-7.10 (m, 3H), 7.03 (m, 1H), 6.70 (d, 1H), 6.52 (s, 1H), 6.21 (s, 1H), 5.90-5.84 (m, 2H), 5.52-5.33 (m, 2H), 4.99 (d, 1H), 4.72 (d, 1H), 3.95 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.8, 167.5, 156.0, 148.9, 142.4, 142.3, 137.4, 132.8, 132.1, 131.4, 129.0, 128.6, 127.4, 126.5, 123.9, 123.6, 119.4, 109.5, 103.1, 101.5, 93.7, 80.7, 58.4, 52.3, 42.4; MS (ES+) *m/z* 430.3 (M + 1), 452.3 (M + 23).

EXAMPLE 1.5

Synthesis of methyl 3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoate



Following the procedure as described in EXAMPLE 1, and making non-critical variations using methyl 3-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained as a white powder in 73% yield; ^{1}H NMR (300 MHz, $CDCl_3$) δ

7.97-7.95 (m, 2H), 7.53-7.50 (m, 1H), 7.45-7.40 (m, 1H), 7.21-7.15 (m, 2H), 7.04-6.99 (m, 1H), 6.73-6.71 (m, 1H), 6.52 (s, 1H), 6.20 (s, 1H), 5.86 (s, 1H), 5.18 (d, 1H), 4.72 (d, 1H), 4.80 (d, 1H), 4.69 (d, 1H), 3.89 (s, 1H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.7, 166.6, 156.0, 149.0, 142.4, 141.7, 136.1, 132.2, 131.7, 130.9, 129.2, 128.1, 124.0, 123.7, 119.4, 109.2, 103.1, 101.6, 93.7, 80.5, 64.3, 58.3, 52.3, 43.7; MS (ES+) *m/z* 430 (M + 1).

EXAMPLE 1.6

Synthesis of methyl 4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoate



Following the procedure as described in EXAMPLE 1, and making non-critical variations using methyl 4-{[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]methyl}benzoate to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained as a white powder in 87% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.01 (d, 2H), 7.38 (d, 2H), 7.18 (t, 2H), 7.02 (t, 1H), 6.72 (d, 1H), 6.52 (s, 1H), 6.12 (s, 1H), 5.86 (m, 2H), 5.11 (d, 1H), 4.96 (d, 1H), 4.86 (d, 1H), 4.69 (d, 1H), 3.89 (s, 1H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.7, 166.6, 156.0, 149.0, 142.4, 141.8, 140.8, 132.1, 130.3, 129.8, 129.0 127.3, 124.1, 123.7, 119.3, 109.2, 103.0, 102.0, 93.7, 80.5, 58.3, 52.2, 43.9; MS (ES+) *m/z* 430.1 (M + 1).

EXAMPLE 1.7

Synthesis of 1'-(diphenylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1, and making non-critical variations using 1-(diphenylmethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained as a white powder in 26% yield: MS (ES+) *m/z* 462.3 (M + 1).

EXAMPLE 1.8

Synthesis of spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a solution of 1'-(diphenylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (2.10 g, 4.70 mmol) in EtOAc (100 mL) and acetic acid (0.10 mL) was added palladium on carbon (1.00 g). The reaction mixture was hydrogenated under 60 psi of hydrogen at ambient temperature for 5 days and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography to yield the title compound (0.87 g, 66%) as a white powder: mp 252 °C (dec.); ^{1}H NMR (300 MHz, $CDCl_3$) δ 10.55 (s, 1H), 7.25-6.84 (m, 4H), 6.64 (s, 1H), 6.22 (s, 1H), 5.88 (s, 2H), 4.76-4.57 (dd, 2H); MS (ES+) *m/z* 282.2 (M + 1).

EXAMPLE 1.9

Synthesis of 2-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1*H*-isoindole-1,3(2*H*)-dione



Following the procedure as described in EXAMPLE 1, and making non-critical variations using 2-{3-[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]propyl}-1*H*-isoindole-1,3(2*H*)-dione to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-

2*H*-indol-2-one, the title compound was obtained in 45% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.83-7.80 (m, 2H), 7.70-7.68 (m, 2H), 7.28-7.26 (m, 1H), 7.15 (d, 1H), 7.05-7.00 (m, 1H), 6.86 (d, 1H), 6.48 (s, 1H), 6.23 (s, 1H), 5.85-5.83 (m, 2H), 4.91 (d, 1H), 4.65 (d, 1H), 3.94-3.68- (m, 4H), 2.15-2.10- (m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.4, 168.2, 155.9, 148.8, 142.4, 141.9, 134.0, 132.5, 132.0, 128.9, 124.1, 123.4, 123.3, 119.4, 108.4, 103.2, 101.5, 93.6, 80.4, 58.2, 38.0, 35.6, 26.8; MS (ES+) *m/z* 469.3 (M + 1), 491.3 (M + 23).

EXAMPLE 1.10

Synthesis of 2-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-1*H*-isoindole-1,3(2*H*)-dione



Following the procedure as described in EXAMPLE 1, and making non-critical variations using 2-{2-[3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]ethyl}-1*H*-isoindole-1,3(2*H*)-dione to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 61% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.81-7.73 (m, 2H), 7.71-7.62 (m, 2H), 7.18-7.08 (m, 2H), 6.98 (t, 1H), 6.87 (d, 1H), 6.43 (s, 1H), 6.29 (s, 1H), 5.91-5.81 (ABq, 2H), 4.79 (d, 1H), 4.58 (d, 1H), 4.18-3.92 (m, 4H), 3.06 (t, 2H), 1.59-1.35 (br, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178, 168.2, 156.1, 148.7, 142.2, 141.9, 134.1, 132.4, 131.8, 128.7, 124.1, 123.4, 123.3, 119.0, 107.8, 103.7, 101.4, 93.4, 80.9, 58.1, 39.0, 35.6; MS (ES+) *m/z* 455 (M + 1), 477 (M + 23).

EXAMPLE 1.11

Synthesis of 1'-[3-(Benzyloxy)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1, and making non-critical variations using 1-[3-(benzyloxy)propyl]-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 98% yield as a pale yellow syrup: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.38-6.95 (m, 9H), 6.49 (s, 1H), 6.08 (s, 1H), 5.83 (dd, 2H), 5.86 (ABq, 1H), 4.58 (ABq, 1H), 3.96-3.79 (m, 2H), 3.53 (t, 2H), 2.06-2.00 (m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.4, 155.9, 148.8, 142.5, 142.2, 138.1, 132.3, 128.9, 127.9, 127.6, 123.9, 123.1, 119.5, 108.7, 103.0, 101.4, 93.6, 80.4, 73.1, 67.4, 58.1, 37.7, 27.9; MS (ES+) *m/z* 430.3 (M + 1).

EXAMPLE 1.12

Synthesis of 5,6-difluoro-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1, and making non-critical variations using 3-(4,5-difluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 71% yield: mp 48-50 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.33 (td, 1H), 7.18-7.12 (m, 2H), 6.93 (d, 1H), 6.77 (dd, 1H), 6.51 (dd, 1H), 4.96 (d, 1H), 4.71 (d, 1H), 3.87-3.64 (m, 2H), 1.82-1.65 (m, 2H), 1.46-1.28 (m, 4H), 0.92 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.6, 156.8, 152.9, 149.8, 144.2, 142.6, 131.8, 129.4, 124.1, 123.5, 111.7, 109.0, 100.2,

80.9, 57.9, 40.6, 29.1, 27.2, 22.5, 14.1; MS (ES+) *m/z* 344 (M + 1).

EXAMPLE 1.13

Synthesis of 5-fluoro-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1, and making non-critical variations using 3-(5-fluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 3% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.37-7.28 (m, 1H), 7.18-7.11 (m, 1H), 7.09-7.01 (m, 1H), 6.98-6.82 (m, 3H), 6.45-6.37 (m, 1H), 4.95 (d, 1H), 4.69 (d, 1H), 3.89-3.63 (m, 2H), 1.81-1.65 (m, 2H), 1.48-1.28 (m, 4H), 0.92 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.7, 159.4, 156.6, 132.2, 130.1, 129.3, 124.1, 123.4, 116.3, 110.8, 110.5, 108.9, 80.4, 58.4, 40.6, 29.2, 27.3, 22.5, 14.1; MS (ES+) *m/z* 326 (M + 1).

EXAMPLE 1.14

Synthesis of 5-bromo-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1, and making non-critical variations using 3-(5-bromo-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 4% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.38-7.23 (m, 2H), 7.17-7.01 (m, 2H), 6.93 (d, 1H), 6.84 (d, 1H), 6.79 (d, 1H), 4.95 (d, 1H), 4.69 (d, 1H), 3.89-3.64 (m, 2H), 1.81-1.65 (m,

2H), 1.48-1.28 (m, 4H), 0.92 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.7, 160.1, 142.6, 132.7, 132.1, 131.4, 129.3, 126.5, 124.1, 123.5, 113.1, 112.2, 108.9, 80.3, 58.0, 40.6, 29.2, 27.3, 22.5, 14.2; MS (ES+) *m/z* 386 (M + 1), 388 (M + 23).

EXAMPLE 1.15

Synthesis of 5-chloro-6-fluoro-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1, and making non-critical variations using 3-(5-chloro-4-fluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 80% yield; mp 74-76 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.38-7.27 (m, 1H), 7.18-7.02 (m, 2H), 6.94 (d, 1H), 6.77 (d, 1H), 6.69 (d, 1H), 4.98 (d, 1H), 4.72 (d, 1H), 3.87-3.64 (m, 2H), 1.82-1.65 (m, 2H), 1.47-1.28 (m, 4H), 0.92 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.5, 160.6, 157.4, 142.6, 133.9, 131.8, 129.5, 124.6, 124.1, 123.5, 113.1, 109.0, 100.0, 81.2, 57.5, 40.6, 29.2, 27.2, 22.5, 14.1; MS (ES+) *m/z* 360 (M + 1).

EXAMPLE 1.16

Synthesis of 6-methoxy-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1, and making non-critical variations using 3-(2-hydroxy-4-methoxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-

5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 99% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.30 (td, 1H), 7.14 (dd, 1H), 7.03 (t, 1H), 6.91 (d, 1H), 6.58 (d, 1H), 6.52 (d, 1H), 6.36 (dd, 1H), 4.93 (d, 1H), 4.69 (d, 1H), 3.91-3.63 (m, 2H), 3.77 (s, 3H), 1.81-1.65 (m, 2H), 1.46-1.29 (m, 4H), 0.91 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.4, 162.1, 161.5, 142.5, 132.9, 128.8, 123.9, 123.5, 123.1, 121.0, 108.5, 107.5, 96.6, 80.5, 57.6, 55.6, 40.3, 29.0, 27.1, 22.3, 14.0; MS (ES+) *m/z* 338 (M + 1).

EXAMPLE 1.17

Synthesis of 6-chloro-5-fluoro-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1, and making non-critical variations using 3-(4-chloro-5-fluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 44% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.34 (td, 1H), 7.14 (dd, 1H), 7.06 (td, 1H), 6.98 (d, 1H), 6.93 (d, 1H), 6.50 (d, 1H), 4.96 (d, 1H), 4.70 (d, 1H), 3.87-3.63 (m, 2H), 1.81-1.65 (m, 2H), 1.47-1.29 (m, 4H), 0.91 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.2, 156,9, 154.8, 151.6, 142.5, 131.5, 129.4, 128.6, 123.7, 121.7, 121.9, 111.2, 108.9, 80.6, 57.9, 40.5, 29.0, 27.1, 22.3, 14.0; MS (ES+) *m/z* 360 (M + 1).

EXAMPLE 1.18

Synthesis of 1'-pentyl-5-(trifluoromethyl)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1, and making non-critical variations using 3-(hydroxymethyl)-3-[2-hydroxy-5-(trifluoromethyl)phenyl]-1-pentyl-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 27% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.48 (dd, 1H), 7.35 (td, 1H), 7.16-6.90 (m, 5H), 5.02 (d, 1H), 4.76 (d, 1H), 3.91-3.65 (m, 2H), 1.82-1.67 (m, 2H), 1.47-1.29 (m, 4H), 0.91 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.5, 163.3, 142.6, 131.8, 129.8, 129.3, 127.7, 124.1, 124.0, 123.9, 123.6, 121.0, 110.6, 108.9, 80.5, 57.6, 40.5, 29.0, 27.1, 22.3, 13.9; MS (ES+) *m/z* 376 (M + 1).

EXAMPLE 1.19

Synthesis of 5,6-dichloro-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1, and making non-criticial variations using 3-(4,5-dichloro-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 43% yield. ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.35 (td, 1H), 7.17-7.03 (m, 3H), 6.94 (d, 1H), 6.76 (s, 1H), 4.98 (d, 1H), 4.72 (d, 1H), 3.88-3.65 (m, 2H), 1.82-1.67 (m, 2H), 1.47-1.29 (m, 4H), 0.92 (t, 3H). ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.2, 159.9, 142.5, 133.3, 131.5, 129.4, 124.6, 124.5, 123.9, 123.4, 112.4, 108.9, 80.8, 57.5, 40.5, 29.0, 27.1, 22.3, 14.0; MS (ES+) *m/z* 376 (M + 1), 378 (M + 1).

EXAMPLE 1.20

Synthesis of 1'-(diphenylmethyl)-5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE1, and making non-critical variations using 1-(diphenylmethyl)-3-(6-hydroxy-3,3-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained: mp 190-192 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.40-7.26 (m, 10H), 7.19-7.15 (m, 1H), 7.07-6.93 (m, 3H), 6.55-6.51 (m, 1H), 6.38 (s, 1H), 6.20 (s, 1H), 4.98 (d, 1H), 4.71 (d, 1H), 4.17 (s, 2H), 1.17 (s, 3H), 1.14 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.1, 161.1, 161.0, 141.8, 137.9, 137.2, 132.8, 130.0, 128.6, 128.5, 128.4, 128.2, 128.0, 127.8, 123.9, 123.1, 120.8, 116.1, 112.1, 93.4, 85.4, 80.4, 58.7, 57.4, 41.3, 27.7, 27.6; MS (ES+) *m/z* 474.5 (M +1).

EXAMPLE 1.21

Synthesis of 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one



To a solution of 1'-(diphenylmethyl)-5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one (0.23 g, 0.49 mmol) in methanol (50.0 mL) was added palladium on carbon (0.10 g). The mixture was hydrogenated under 120 psi of hydrogen at ambient temperature overnight. The reaction mixture was filtered through celite, washed with methanol. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/2) to give the title compound (0.10 g, 68%): mp 95-100 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.74 (s,

1H), 7.28-7.20 (m, 1H), 7.15 (d, 1H), 7.03 (t, 1H), 6.95 (d, 1H), 6.43 (s, 1H), 6.40 (s, 1H), 4.94 (d, 1H), 4.66 (d, 1H), 4.19 (s, 2H), 1.20 (s, 3H), 1.16 (3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 180.4, 161.3, 161.0, 140.3, 133.0, 130.1, 128.8, 124.2, 123.4, 120.0, 116.6, 110.1, 93.4, 85.5, 80.6, 58.3, 41.4, 27.7, 27.6; MS (ES+) *m/z* 308.6 (M + 1).

EXAMPLE 1.22

Synthesis of 4',7'-dichloro-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a solution of 4,7-dichloro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one (0.69 g, 1.57 mmol) in anhydrous tetrahydrofuran (15.0 mL) was added triphenylphosphine (0.54 g, 2.04 mmol) followed by slow addition of diisopropyl azodicarboxylate (0.41 g, 2.04 mmol) at 0 °C. The brown reaction mixture was stirred at ambient temperature for 16 h and quenched with ammonium chloride solution (2.00 mL). The organic solvent was removed *in vacuo*. The residue was dissolved in ethyl acetate (20.0 mL), washed with 10% aqueous HCl solution (10.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate/hexane (35%) to give a solid, which was crystallized from ethyl acetate/ether to give the title compound (0.13 g, 20%) as a colorless solid: mp 106-108 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.24-7.16 (m, 1H), 6.81 (d, 1H), 6.44 (s, 1H), 6.07 (s, 1H), 5.86 (dd, 2H), 4.87 (dd, 2H), 4.12-4.07 (m, 2H), 1.76-1.66 (m, 2H) 1.36-1.31 (m, 4H), 0.89 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.5, 157. 0, 149.2, 142.1, 140.0, 132.3, 131.0, 130.2, 124.6, 116.0, 113.8, 102.3, 101.5, 93.3, 77.2, 58.5, 42.1, 29.5, 28.7, 22.3, 14.0; MS (ES+) *m/z* 420.4 (M + 1).

EXAMPLE 1.23

Synthesis of 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1.22, and making non-critical variations using 4-bromo-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one to replace 4,7-dichloro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 71% yield as a colorless solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.79 (s, 1H), 7.19-7.08 (m, 2H), 6.90 (dd, 1H), 6.58 (s, 1H), 6.25 (s, 1H), 5.90 (d, 2H), 4.74 (ABq, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 178.5, 157.0, 148.8, 144.5, 141.9, 131.2, 130.6, 126.1, 119.2, 117.5, 109.8, 103.3, 101.8, 93.3, 77.6, 59.7; MS (ES-) *m/z* 360.4 (M - 1), 358.4 (M – 1).

EXAMPLE 1.24

Synthesis of 4'-bromo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one



To a solution of 4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one (1.80 g, 4.80 mmol) in anhydrous ethyl acetate (50 mL) was added tributylphosphine (1.26 g, 1.54 mL, 6.24 mmol) at 0 °C under nitrogen. A solution of di-*tert*-butyl azodicarboxylate (1.44 g, 6.24 mmol) in anhydrous ethyl acetate (15.0 mL) was added over 10 min. The reaction solution was stirred for 2 h, and then quenched with saturated ammonium chloride solution (30.0 mL). After the aqueous layer was separated, the organic layer was washed with 10% aqueous HCl solution (2 x 25.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate-hexane (70%) to obtain a solid which was triturated with diethyl ether to give the title compound (0.64 g, 37%) as a

colorless solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.77 (s, 1H), 7.18-7.13 (m, 1H), 7.08 (d, 1H), 6.90 (d, 1H), 6.47 (s, 1H), 6.30 (s, 1H), 4.80 (ABq, 2H), 4.46 (t, 2H), 2.92 (t, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 178.7, 162.2, 161.7, 144.5, 131.1, 131.0, 126.1, 119.8, 119.2, 119.1, 118.3, 109.7, 92.4, 77.6, 72.5, 59.2, 28.8; MS (ES-) *m/z* 358.4 (M - 1), 356.3 (M - 1).

EXAMPLE 1.25

Synthesis of 4'-bromo-1'-(pyridin-2-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one



A mixture of 4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1-(pyridin-2-ylmethyl)-1,3-dihydro-2*H*-indol-2-one (1.81 g, 3.88 mmol), triphenylphosphine (2.04 g, 7.77 mmol) and diisopropyl azodicarboxylate (1.57 g, 7.77 mmol) in anhydrous dioxane (60 mL) was heated at reflux for 16 h. After cooling down to ambient temperature, the solvent was removed *in vacuo*. The gummy residue was diluted with ethyl acetate (50.0 mL), washed with water (3 x 25.0 mL), brine (3 x 25.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate/hexane (80%) to give the title compound (0.64 g, 37%) as colorless solid: mp >200 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.47 (d, 1H), 7.77 (dt, 1H), 7.37 (d, 1H), 7.27 (dt, 1H), 7.19-7.13 (m, 2H), 6.94 (dd, 1H), 6.61 (s, 1H), 6.33 (s, 1H), 5.08 (d, 1H), 5.03 (d, 1H), 4.93 (d, 1H), 4.74 (d, 1H), 4.48 (t, *J* = 8.6 Hz, 2H), 2.96 (t, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 177.4, 162.2, 161.8, 155.3, 149.8, 145.3, 137.6, 131.0, 130.5, 126.8, 123.3, 122.2, 119.9, 119.5, 119.0, 118.1, 109.3, 92.4, 77.5, 72.5, 58.8, 45.3, 28.8; MS (ES+) *m/z* 451.3 (M + 1).

EXAMPLE 1.26

Synthesis of 5'-fluoro-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a solution of 5-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one (3.34 g, 7.18 mmol) in anhydrous tetrahydrofuran (80.0 mL) was added tributylphosphine (2.18 g, 2.70 mL, 10.8 mmol) under nitrogen. A solution of di-*tert*-butyl azodicarboxylate (2.49 g, 10.8 mmol) in anhydrous tetrahydrofuran (25.0 mL) was added over 10 min. The reaction solution was stirred for 1 h, and quenched with saturated ammonium chloride (30.0 mL). After the solvent was removed under reduced pressure, the gummy material was extracted with ethyl acetate (3 x 75.0 mL). The organic layer was washed with 10% aqueous HCl solution (2 x 25.0 mL), saturated aqueous sodium hydrogen carbonate (3 x 25.0 mL), brine (3 x 25.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate/hexane (30%) to afforded the title compound (1.10 g, 34%) as a colorless solid: mp 139-141 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.02-6.96 (m, 1H), 6.93-6.89 (m, 2H), 6.74-6.73 (m, 1H), 6.50 (s, 1H), 6.38 (d, 1H), 6.09 (s, 1H), 5.87 (dd, 2H), 4.95 (ABq, 2H), 4.78 (Abq, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.9, 161.5, 158.3, 155.9, 151.7, 149.2, 142.6, 137.1, 137.1, 133.7, 118.6, 115.6, 115.3, 112.7, 112.4, 112.0, 109.7, 109.6, 109.4, 102.8, 101.7, 93.8, 80.1, 58.6, 37.1; MS (ES+) *m/z* 448.2 (M + 1).

EXAMPLE 1.27

Synthesis of 1'-(diphenylmethyl)-5'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a solution of 1-(diphenylmethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-5-methyl-1,3-dihydro-2*H*-indol-2-one (1.31 g, 2.72 mmol) in ethyl acetate (50.0 mL) was added tributylphosphine (0.82 g, 4.07 mmol). A solution of di-*tert*-butyl azodicarboxylate (0.94 g, 4.07 mmol) in ethyl acetate (45.0 mL) was added to the above reaction mixture over a period of 5 minutes. After stirring for 10 minutes under N_2, the reaction was quenched with saturated aqueous ammonium chloride (60.0 mL). The organic layer was separated and washed with 1.0 N hydrochloric acid solution (3 x100 mL), dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate/hexane (10-50%) to afford the title compound (0.98 g, 78% yield): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.40-7.25 (m, 10H), 7.02, (s, 1H), 6.96 (s, 1H), 6.79 (d, 1H), 6.50 (s, 1H), 6.36 (d, 1H), 6.08 (s, 1H), 5.86 (d, 2H), 4.82 (ABq, 2H), 2.20 (s, 3H).

EXAMPLE 1.28

Synthesis of 5'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



A stainless steel hydrogenating vessel was successively charged with 1'-(diphenylmethyl)-5'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.90 g, 1.95 mmol), glacial acetic acid (50.0 mL) and palladium hydroxide (0.10 g, 1.35 mmol, 20 wt % on carbon). The vessel was flushed with nitrogen, sealed then heated to 60 ^{o}C and placed under 120 Psi of H_2. After 4 days of stirring, the reaction mixture was diluted with ethyl acetate and passed through a bed of celite. The filtrate

was washed with water (6 x 100 mL), dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate/hexane (40-50%) to afford the title compound (0.25 g, 43%): mp 269-271 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.45 (s, 1H), 7.00 (d, 1H), 6.87 (s, 1H), 6.76 (d, 1H), 6.63 (s, 1H), 6.21 (s, 1H), 5.87 (d, 2H), 4.64 (ABq, 2H), 2.17 (s, 3H); MS (ES+) *m/z* 296.28 (M + 1).

EXAMPLE 1.29

Synthesis of 5'-methyl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a suspension of sodium hydride (0.03 g, 0.63 mmol, 60% dispersion in mineral oil) in *N*, *N*-dimethylformamide (5.00 mL) was slowly added a solution of 5'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.10 g, 0.33 mmol) in *N*,*N*-dimethylformamide (5.00 mL) at 0 °C. After stirring for 15 minutes at 0 °C, a solution of 2-(bromomethyl)-5-(trifluoromethyl)furan (0.11 g, 0.49 mmol) in *N*,*N*-dimethylformamide (40.0 mL) was added. The resulting mixture was stirred at ambient temperature for 4 h and quenched with water (20.0 mL). The mixture was extracted with ethyl acetate (3 x 25.0 mL). The combined organic layers was washed with water (50.0 mL) and brine (2 x 25.0 mL), dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo*. The residue was subjected to column chromatography eluting with ethyl acetate/hexane (15-50%) to afford the title compound (0.11 g, 77% yield): mp 96-98 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.09 (d, 1H), 7.00 (s, 1H), 6.87 (d, 1H), 6.74 (d, 1H), 6.52 (s, 1H), 6.38 (d, 1H), 6.11 (s, 1H), 5.88 (d, 2H), 4.96 (ABq, 2H), 4.80 (ABq, 2H), 2.29 (s, 3H); MS (ES+) *m/z* 444.2 (M + 1).

EXAMPLE 1.30

Synthesis of 6-bromo-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1.22, and making non-critical variations using 3-(4-bromo-2-hydroxyphenyl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one to replace 4,7-dichloro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 82% yield as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.33 (td, 1H), 7.15-7.14 (m, 2H), 7.04 (dd, 1H), 6.96-6.90 (m, 2H), 6.56 (d, 1H), 4.95 (d, 1H), 4.69 (d, 1H), 3.89-3.64 (m, 2H), 1.80-1.68 (m, 2H), 1.43-1.34 (m, 4H), 0.92 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.6, 161.6, 142.5, 132.1, 129.1, 128.4, 124.4, 123.9, 123.3, 122.8, 114.1, 108.8, 80.4, 57.6, 40.4, 29.0, 27.1, 22.3,14.0; MS (ES+) *m/z* 386.3 (M + 1), 388.3 (M + 1).

EXAMPLE 1.31

Synthesis of 5-bromo-1'-(diphenylmethyl)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in Example 1, and making non-critical variations using 3-(5-bromo-2-hydroxyphenyl)-1-(diphenylmethyl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (72%) as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.43-7.25 (m, 11H), 7.14-6.93 (m, 4H), 6.83 (d, 1H), 6.71 (d, 1H), 6.52 (d, 1H), 5.0 (d, 1H), 4.73 (d, 1H); MS

(ES+) *m/z* 484.4 (M + 1), 482.4 (M + 1).

EXAMPLE 1.32

Synthesis of 2-methyl-1'-pentylspiro[furo[2,3-*f*][1,3]benzothiazole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1.22, and making non-critical variations using 3-(hydroxymethyl)-3-(5-hydroxy-2-methyl-1,3-benzothiazol-6-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one to replace 4,7-dichloro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (50%) as a white solid: mp 105-107 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.59 (d, 1H), 7.28 (dt, 1H), 7.02-6.92 (m, 2H), 5.02 (d, 1H), 4.77 (d, 1H), 4.01 (m, 1H), 3.64 (m, 1H), 2.54 (s, 3H), 1.92-1.71 (m, 2H), 1.54-1.34 (m, 4H), 0.92 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.7, 169.2, 160.2, 149.2, 142.7, 138.3, 132.7, 129.0, 128.6, 123.5, 122.7, 122.1, 120.2, 108.6, 108.3, 80.1, 58.1, 40.7, 29.1, 27.0, 22.5, 20.2, 14.1; MS (ES+) *m/z* 379.5 (M + 1).

EXAMPLE 1.33

Synthesis of 5-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE1.22, and making non-

critical variations using 3-(hydroxymethyl)-3-(5-hydroxy-2-methyl-1,3-benzothiazol-6-yl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one to replace 4,7-dichloro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (30%) as a white solid: mp 143-145 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.35-7.27 (m, 2H), 7.17-6.98 (m, 3H), 6.84 (d, 1H), 6.78-6.73 (m, 2H), 6.40 (d, 1H), 5.07-4.87 (m, 3H), 4.69 (d, 1H); MS (ES+) *m/z* 464.2 (M + 1), 466.2 (M + 1).

EXAMPLE 1.34

Synthesis of 5-bromospiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1, and making non-critical variations using 3-(5-bromo-2-hydroxyphenyl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (25%) as a white solid: mp 225-228 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 10.65 (s, 1H), 7.35 (dd, 1H), 7.24 (dt, 1H), 7.11 (d, 1H), 6.99-6.88 (m, 3H), 6.83 (d, 1H), 4.81 (d, 1H), 4.69 (d, 1H); MS (ES+) *m/z* 316.1 (M + 1), 318.1 (M + 1).

EXAMPLE 1.35

Synthesis of 1'-(diphenylmethyl)-6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



To a solution of 1-(diphenylmethyl)-3-(hydroxymethyl)-3-[2-hydroxy-4-(trifluoromethoxy)phenyl]-1,3-dihydro-2*H*-indol-2-one (17.3 mmol) in anhydrous THF (200 mL) was added triphenylphosphine (6.34 g, 24.2 mmol) followed by diethyl

azodicarboxylate (4.39 mL, 24.2 mmol) at 0 °C. The reaction mixture was stirred at ambient temperature for 16 h, quenched with saturated ammonium chloride (40.0 mL). The aqueous mixture was extracted with ethyl acetate (3 x 150 mL). The combined organic layers was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo*. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/4) to give the title compound (6.00 g, 71%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.45-6.51 (m, 18H), 5.08 (d, 1H), 4.81 (d, 1H); MS (ES+) *m/z* 488 (M + 1).

EXAMPLE 1.36

Synthesis of 6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



To a suspension of 1'-(diphenylmethyl)-6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one (6.00 g, 12.3 mmol) in methanol (100 mL) and acetic acid (1.00 mL) was added 10 % palladium on carbon (0.65 g, 0.62 mmol), and the mixture was hydrogenated at ambient temperature under 130 psi of hydrogen for 5 days. The reaction mixture was filtered over celite and the filtrate was concentrated *in vacuo* to dryness. The residue was subjected to flash chromatography eluting with 30% ethyl acetate in hexane to give the title compound (2.95 g, 75%) as a white solid: mp 180-182 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.19 (s, 1H), 7.29-6.92 (m., 4H), 6.86-6.64 (m, 3H), 5.03 (d, 1H), 4.75 (d, 1H); ^{13}C NMR(75 MHz, $CDCl_3$) δ 179.9, 161.9, 150.6, 132.3, 129.4, 127.4, 125.6, 124.3, 124.2, 123.8, 120.5, 114.1, 110.7, 104.3, 80.8, 58.2; MS (ES+) *m/z* 322 (M + 1).

EXAMPLE 1.37

Synthesis of ethyl (4'-bromo-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)acetate



Following the procedure as described in EXAMPLE 1.22, and making non-critical variations using ethyl [4-bromo-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate to replace 4,7-dichloro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (41%) as a colorless solid: MS (ES+) *m/z* 445.5 (M + 1).

EXAMPLE 1.38

Synthesis of ethyl (4'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetate



Following the procedure as described in EXAMPLE 1.22, and making non-critical variations using ethyl [4-chloro-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate to replace 4,7-dichloro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 63% yield as a colorless solid: MS (ES+) *m/z* 400.8 (M + 1).

EXAMPLE 1.39

Synthesis of ethyl (4'-bromo-6,6-dimethyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetate



Following the procedure as described in EXAMPLE 1, and making non-critical variations using ethyl [4-bromo-3-(6-hydroxy-2,2-dimethyl-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate to replace 1-(2-

cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 52% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.22-7.10 (m, 2H), 6.70 (d, 1H), 6.48 (s, 1H), 6.30 (s, 1H), 5.0 (d, 1H); 4.86 (d, 1H), 4.63 (d, 1H), 4.35 (d, 1H), 4.28-4.18 (m, 2H), 2.79 (s, 2H), 1.43 (s, 3H), 1.39 (s, 3H), 1.28 (t, 3H); MS (ES+) *m/z* 472.5 (M + 1), 474.5 (M + 1).

EXAMPLE 1.40

Synthesis of ethyl (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2' *H*)-yl)acetate



Following the procedure as described in EXAMPLE 1, and making non-critical variations using ethyl [4-bromo-3-(4,5-difluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 81% yield: MS (ES+) *m/z* 438.4 (M + 1), 440.4 (M + 1).

EXAMPLE 1.41

Synthesis of ethyl (5'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)acetate



Following the procedure as described in EXAMPLE 1.22, and making non-critical variations using ethyl [5-chloro-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate to replace 4,7-dichloro-3-(6-

hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 90% yield as a colorless solid: MS (ES+) *m/z* 400.8 (M + 1).

EXAMPLE 1.42

Synthesis of 7'-fluorospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



A solution of 7-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-1,3-dihydro-2*H*-indol-2-one (2.00 g, 7.00 mmol) and paraformaldehyde (2.10 g, 61.0 mmol) in THF (50 mL) was degassed by bubbling through argon for one hour, followed by the slow addition of lithium diisopropylamide (48.8 mL, freshly made 0.50 M solution, 25 mmol) at -78 °C. The mixture was stirred at ambient temperature overnight and quenched with saturated ammonium chloride (50.0 mL). The mixture was extracted with ethyl acetate (3 x 100 mL). The combined organic layers was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was dissolved in anhydrous ethyl acetate (50 mL) followed by the addition of tributylphosphine (2.10 mL, 8.00 mmol) and di-*tert*-butyl azodicarboxylate (1.90 g, 8.00 mmol) at 0 °C. The reaction mixture was stirred at ambient temperature for 16 h and quenched with saturated ammonium chloride (30.0 mL). The organic layer was separated, dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was dissolved in methanol (70.0 mL) followed by the addition of saturated sodium bicarbonate solution (30.0 mL). The resulted mixture was refluxed at 100 °C for one hour. After cooling down to ambient temperature, the mixture was extracted with ethyl acetate (3 x 50.0 mL). The combined organic layers was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/5) to give the title compound (0.27 g, 17%): ^{1}H NMR (300 MHz, DMSO-d_6) δ 11.06 (s, 1H), 7.17-7.08 (m, 1H), 7.00-6.88 (m, 2H), 6.64 (s, 1H), 6.33 (s, 1H), 5.92-5.85 (m, 2H), 4.74 (d, 1H), 4.62 (d, 1H).

EXAMPLE 1.43

Synthesis of methyl (6-chloro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetate



Following the procedure as described in EXAMPLE1.22, and making non-critical variations using methyl [3-(4-chloro-2-hydroxyphenyl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate to replace 4,7-dichloro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 74% yield as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.29 (dt, 1H), 7.14 (dd, 1H), 7.06 (t, 1H), 6.95 (d, 1H), 6.81-6.74 (m, 3H), 5.03 (d, 1H), 4.74 (d, 1H), 4.65 (d, 1H), 4.44 (d, 1H), 3.75 (s, 3H); MS (ES+) *m/z* 344.5 (M + 1), 346.5 (M + 1).

EXAMPLE 1.44

Synthesis of ethyl (5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetate



Following the procedure as described in EXAMPLE 1.22, and making non-critical variations using ethyl [3-(4,5-difluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate to replace 4,7-dichloro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-pentyl-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained in 46% yield as a light yellow oil: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.31 (dt, 1H), 7.16 (dd, 1H), 7.08 (dt, 1H), 6.81-6.71 (m, 2H), 6.67 (dd, 1H), 4.98 (d, 1H), 4.74 (d, 1H), 4.64 (d, 1H), 4.37 (d, 4.24 (q, 7.1 Hz), 1.28 (t, 3H); MS (ES+) *m/z* 360.5 (M + 1).

EXAMPLE 1.45

Synthesis of ethyl (2'-oxo-6,7-dihydro-5*H*-spiro[indeno[5,6-*b*]furan-3,3'-indol]-1'(2'*H*)-yl)acetate



To a solution of ethyl [3-(6-hydroxy-2,3-dihydro-1*H*-inden-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate (4.20 mmol) in anhydrous THF (60.0 mL) was added triphenylphosphine (1.43 g, 5.46 mmol) and diethyl azodicarboxylate (0.95 g, 5.46 mmol) at 0 °C. The reaction mixture was stirred at ambient temperature for 16 h and quenched with saturated ammonium chloride (20.0 mL). The mixture was extracted with ethyl acetate (3 x 50.0 mL). The combined organic layers was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/4) to give the title compound (0.25 g, 16 % in three steps): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.29 (td, 1H), 7.22-7.17 (m, 1H), 7.07 (t, 1H), 6.81 (s, 1H), 6.78 (d, 1H), 6.65 (s, 1H), 4.95 (d, 1H), 4.71 (d, 1H), 4.64 (d, 1H), 4.42 (d, 1H), 4.24 (q, 2H), 2.84 (t, 2H), 2.73-2.65 (m, 2H), 2.10-1.95 (m, 2H), 1.29 (t, 3H); MS (ES+) *m/z* 364 (M + 1).

EXAMPLE 1.46

Synthesis of ethyl (2-oxo-5',6',7',8'-tetrahydrospiro[indole-3,3'-naphtho[2,3-*b*]furan]-1(2*H*)-yl)acetate



Following the procedure as described in EXAMPLE 1.45, and making non-critical variations using ethyl [3-(hydroxymethyl)-3-(3-hydroxy-5,6,7,8-

tetrahydronaphthalen-2-yl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate to replace ethyl [3-(6-hydroxy-2,3-dihydro-1*H*-inden-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained (24 % in three steps): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.29 (td, 1H), 7.19 (d, 1H), 7.07 (t, 1H), 6.79 (d, 1H), 6.66 (s, 1H), 6.51 (s, 1H),), 4.91 (d, 1H), 4.67 (d, 1H), 4.52 (ABq, 2H), 4.24 (q, 2H), 2.77-2.51 (m, 4H), 1.77-1.64 (m, 4H), 1.29 (t, 3H); MS (ES+) *m/z* 378 (M + 1).

EXAMPLE 1.47

Synthesis of ethyl (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetate



Following the procedure as described in EXAMPLE 1.45, and making non-critical variations using ethyl [4-bromo-3-(4,5-difluoro-2-hydroxyphenyl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate to replace ethyl [3-(6-hydroxy-2,3-dihydro-1*H*-inden-5-yl)-3-(hydroxymethyl)-2-oxo-2,3-dihydro-1*H*-indol-1-yl]acetate, the title compound was obtained (41%): mp 133-134 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.26-7.15 (m, 3 H), 6.78-6.58 (m, 2H), 5.08 (d, 1H), 4.91 (d, 1H), 4.63 (d, 1H), 4.35 (d, 1H), 4.24 (q, 2H), 1.29 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.5, 166.8, 157.1, 143.7, 130.7, 129.0, 127.8, 120.0, 111.8, 111.6, 107.5, 99.8, 99.5, 62.2, 59.1, 41.7, 14.1; MS (ES+) *m/z* 438 (M + 1), 440 (M + 1), 460 (M + 23), 462 (M + 23).

EXAMPLE 1.48

Synthesis of 1'-(diphenylmethyl)-6,7-dihydrospiro[benzo[1,2-*b*:4,5-*b*']difuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 1, and making non-critical

variations using 1-(diphenylmethyl)-3-(5-hydroxy-2,3-dihydro-1-benzofuran-6-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (43%): MS (ES+) *m/z* 446.4 (M + 1).

EXAMPLE 1.49

Synthesis of 6,7-dihydrospiro[benzo[1,2-*b*:4,5-*b*']difuran-3,3'-indol]-2'(1'*H*)-one



A mixture of 1'-(diphenylmethyl)-6,7-dihydrospiro[benzo[1,2-*b*:4,5-*b*']difuran-3,3'-indol]-2'(1'*H*)-one (0.29 g, 0.65 mmol) and palladium hydroxide (0.10 g, 20% on activated carbon) in acetic acid (20.0 mL) was hydrogenated at 60 °C under normal pressure of hydrogen for 20 hours. The reaction mixture was filtered through celite and washed with acetone (50.0 mL). The filtrate was concentrated *in vacuo* to dryness to give the title compound (0.13 g, 69%): MS (ES+) *m/z* 280.2 (M + 1).

EXAMPLE 1.50

Synthesis of 1'-(diphenylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 1, and making non-critical variations using 1-(diphenylmethyl)-3-(6-hydroxy-2,3-dihydro-1-benzofuran-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (51%) as a white solid: MS (ES+) *m/z* 446.3 (M + 1).

EXAMPLE 1.51

Synthesis of 5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 1.49, and making non-critical variations using 1'-(diphenylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one to replace 1'-(diphenylmethyl)-6,7-dihydrospiro[benzo[1,2-*b*:4,5-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, the title compound was obtained (68%): mp 208-210 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.53 (s, 1H), 7.39-6.76 (m, 4H), 6.45 (s, 1H), 6.35 (s, 1H), 4.68 (ABq, 2H), 4.45 (t, 2H), 2.92 (t, 2H); MS (ES+) *m/z* 280.2 (M + 1).

EXAMPLE 1.52

Synthesis of ethyl (2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)acetate



A mixture of 5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]2'(1'*H*)-one (0.28 g, 1.00 mmol), ethyl 2-bromoacetate (0.17 g, 1.00 mmol) and cesium carbonate (0.98 g, 3.00 mmol) in acetone (20.0 mL) was stirred at reflux for 5 hours. After cooling down to ambient temperature, the mixture was filtered. The filtrate was evaporated under reduced pressure and the residue was subjected to column chromatography to give the title compound (0.23 g, 63%) as a white solid: MS (ES+) *m/z* 366.4 (M + 1).

EXAMPLE 1.53

Synthesis of 4'-methoxy-1'{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 1, and making non-critical variations using 3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-4-methoxy-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1.3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (33%) as a white solid: mp 149-153 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.32-7.24 (m, 1H), 6.82-6.62 (m, 3H), 6.46 (s, 1H), 6.40 (d, 1H), 6.08 (s, 1H), 5.87 (ABq, 2H), 4.92 (ABq, 2H), 4.82 (ABq, 2H), 3.70 (s, 3H); MS (ES+) *m/z* 460.3 (M + 1).

EXAMPLE 1.54

Synthesis of 7'-fluoro-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 1, and making non-critical variations using 7-fluoro-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1-{[5-(trifluoromethyl)-2-furyl]methyl}-1,3-dihydro-2*H*-indol-2-one to replace 1-(2-cyclopropylethyl)-3-(6-hydroxy-1,3-benzodioxol-5-yl)-3-(hydroxymethyl)-1,3-dihydro-2*H*-indol-2-one, the title compound was obtained (32%) as a white solid: mp 116-118 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.36-6.88 (m, 4H), 6.67 (s, 1H), 6.62 (d, 1H), 6.19 (s, 1H), 5.90 (d, 2H), 5.07 (q, 2H), 4.75 (dd, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ176.9,

155.8, 154.1, 149.1, 148.8, 145.6, 142.4, 140.9, 140.3, 139.2, 139.2, 135.2, 135.2, 128.5, 128.4, 124.9, 128.4, 124.9, 124.8, 124.7, 121.2, 120.4, 120.4, 119.8, 117.6, 117.2, 117.0, 114.5, 114.5, 109.5, 103.2, 102.0, 93.8, 80.1, 58.3, 58.2, 39.0, 38.9; MS (ES+) *m/z* 448.3 (M + 1).

EXAMPLE 2

Synthesis of (2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetic acid



To a suspension of ethyl (2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetate (10.5 g, 24.5 mmol) in THF (200 mL) and water (100 mL) was added lithium hydroxide monohydrate (3.98 g, 95.0 mmol) at 0 °C. The reaction mixture was stirred at 0 °C for 30 min and at ambient temperature for 17 h. The mixture was neutralized with of 4 M HCl (15.0 mL). The residue obtained upon removing the solvent was acidified by the addition of 4 M HCl (6.2 mL) to pH 3. The solid was filtered, washed with water and hexane, and dried under the reduced pressure to give the title compound (8.48 g, 87%) as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.32-7.06 (m, 4H), 6.79 (d, 1H), 6.49 (s, 1H), 6.23 (s, 1H), 5.84 (m, 2H), 4.92 (m, 1H), 4.69-4.63 (m, 2H), 4.45 (m, 1H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.8, 171.8, 155.8, 149.0, 142.4, 141.2, 132.0, 129.0, 124.1, 124.0, 119.2, 108.3, 103.4, 101.5, 93.5, 80.2, 58.2, 41.1; MS (ES-) *m/z* 338.2 (M-1).

EXAMPLE 2.1

Synthesis of (2'-oxo-6,7-dihydro-5*H*-spiro[indeno[5,6-*b*]furan-3,3'-indol]-1'(2'*H*)-yl)acetic acid



Following the procedure as described in EXAMPLE 2, and making non-critical variations using ethyl (2'-oxo-6,7-dihydro-5*H*-spiro[indeno[5,6-*b*]furan-3,3'-indol]-1'(2'*H*)-yl)acetate to replace ethyl (2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetate, the title compound was obtained in 74% yield; MS (ES-) *m/z* 354 (M – 1).

EXAMPLE 2.2

Synthesis of (4'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid



Following the procedure as described in EXAMPLE 2, and making non-critical variations using ethyl (4'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetate to replace ethyl (2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetate, the title compound was obtained in 92% yield as a colorless solid: mp 228-229 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.26-7.21 (m, 1H), 7.03 (dd, 1H), 6.71 (dd, 1H), 6.52 (s, 1H), 6.36 (s, 1H), 4.93 (dd, 2H), 4.69-4.63 (m, 1H), 4.54-4.51 (m, 3H), 2.95 (t, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.8, 170.9, 162.1, 162.1, 143.1, 131.7, 130.0, 128.6, 124.7, 119.6, 118.7, 117.0, 106.7, 92.8, 77.2, 72.3, 58.1, 41.2, 28.9; MS (ES-) *m/z* 370.4 (M-1).

EXAMPLE 2.3

Synthesis of (4'-bromo-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid



Following the procedure as described in EXAMPLE 2, and making non-critical

variations using ethyl (4'-bromo-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetate to replace ethyl (2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetate, the title compound was obtained (98%) as a colorless solid: MS (ES-) *m/z* 415.2 (M-1).

EXAMPLE 2.4

Synthesis of (5'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid



Following the procedure as described in EXAMPLE 2, and making non-critical variations using ethyl (5'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetate to replace ethyl (2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetate, the title compound was obtained in 98% yield as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.27-7.25 (m, 1H), 7.16 (d, 1H), 6.72 (d, 1H), 6.54 (s, 1H), 6.39 (s, 1H), 4.93 (dd, 2H), 4.69-4.63 (m, 1H), 4.54-4.51 (m, 3H), 2.95 (t, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.8, 170.9, 162.1, 162.1, 143.1, 131.7, 130.0, 128.6, 124.7, 119.6, 118.7, 117.0, 106.7, 92.8, 77.2, 72.3, 58.1, 41.2, 28.9; MS (ES-) *m/z* 370.4 (M - 1).

EXAMPLE 2.5

Synthesis of (2-oxo-5',6',7',8'-tetrahydrospiro[indole-3,3'-naphtho[2,3-*b*]furan]-1(2*H*)-yl)acetic acid



Following the procedure as described in Example 2, and making non-critical variations using ethyl (2-oxo-5',6',7',8'-tetrahydrospiro[indole-3,3'-naphtho[2,3-*b*]furan]-

1(2*H*)-yl)acetate to replace ethyl (2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetate, the title compound was obtained (99%): MS (ES-) *m/z* 348 (M – 1).

EXAMPLE 2.6

Synthesis of (2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid



A mixture of ethyl (2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetate (0.23 g, 0.63 mmol) and LiOH (0.10 g, 4.20 mmol) in MeOH/H_2O (1/1, 20.0 mL) was stirred at ambient temperature for 20 hours. The mixture was acidified with 0.1 M HCl until pH 2-3. The solid was filtered off and dried to give the title compound (0.15 g, 70%): MS (ES-) *m/z* 336.3 (M-1).

EXAMPLE 3

Synthesis of *N*-(4-chlorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide



A. Preparation of stock solution of isobutyl (2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetyl carbonate

To a solution of (2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetic acid (0.30 g, 0.88 mmol) in dichloromethane (12.5 mL) was added *N*-methylmorpholine (0.09 g, 0.88 mmol) and *iso*-butyl chloroformate (0.12 g, 0.88 mmol) dropwise at 0 °C. The mixture was stirred at 0 °C for 1.5 h and at ambient temperature for 3 h. This mixture was used as a mixed anhydride for the next step amide

formation.

B. Synthesis of *N*-(4-chlorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide

To the above mixed anhydride stock solution (2.50 mL, 0.18 mmol) was added a solution of 4-chlorobenzylamine in dichloromethane (0.35 mL, 0.50 M, 0.18 mmol) at ambient temperature. The reaction mixture was stirred at ambient temperature for 23 h, washed with saturated aqueous sodium carbonate and water. After removal of the solvent, diethyl ether was added and the precipitate was collected by filtration to give the title compound (0.04 g, 46%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.39-6.97 (m, 8H), 6.49 (s, 1H), 6.29 (br, 1H), 6.01 (s, 1H), 5.85 (m, 2H), 4.87 (m, 1H), 4.65 (m, 1H), 4.53-4.29 (m, 4H); MS (ES+), *m/z* 485.2 (M + 23).

EXAMPLE 3.1

The compounds listed in the following table were synthesized using similar conditions as described in Example 3. The compound numbers listed below do not correspond to the compound numbers provided in the general Reaction Schemes above.

Compound Number	Name	MS (*m/z*, M + 1)
1	*N*-(3-fluorophenyl)-2-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'H)-yl)acetamide	433.2
2	*N*-butyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'H)-yl)acetamide	395.2
3	1'-(2-oxo-2-piperidin-1-ylethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	407.2
4	*N*-butyl-*N*-methyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	409.3
5	2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-phenylacetamide	415.2
6	*N*-(4-fluorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'H)-yl)acetamide	433.3
7	*N*-(3-fluorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	447.3
8	*N*-(3-chlorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	449.2

Compound Number	Name	MS (*m/z*, M + 1)
9	*N*-(2-fluorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	433.3
10	*N*-(2-ethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	443.2
11	*N*-(4-ethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	443.3
12	*N*-(3-methylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	429.3
13	*N*-(2,3-dimethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	443.3
14	*N*-(3,5-dimethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	443.4
15	2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-pentylacetamide	409.4
16	2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-propylacetamide	381.3
17	*N*-isopropyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	381.4
18	*N*-(3-methylbutyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	409.3
19	*N*-isobutyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	395.3
20	*N*-hexyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	423.4
21	*N*-cyclohexyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	421.4
22	*N*-cyclopentyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	407.3
23	*N*-heptyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	437.5
24	*N*-(2-chlorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	463.2
25	*N*-(2,6-dimethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	443.3

Compound Number	Name	MS (*m/z*, M + 1)
26	*N*-(2-methoxyphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	445.3
27	*N*-[(5-methyl-2-furyl)methyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	433.2
28	*N*-ethyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	367.4
29	*N*-methyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	353.3
30	*N*-(2-fluorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	447.2
31	*N*-[2-(3-methoxyphenyl)ethyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	473.3
32	*N*-(2-ethoxyethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	411.3
33	*N*-(4-methoxybenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	459.3
34	*N*-(2,4-dimethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	443.3
35	*N*-(3-isopropoxypropyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	439.4
36	*N*-(2-furylmethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	419.3
37	*N*-(cyclohexylmethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	435.3
38	*N*-(3-fluoro-2-methylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	447.3
39	*N*-(4-methoxyphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	445.3
40	*N*-cyclobutyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	393.4
41	*N*-(2,5-difluorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	451.3
42	*N*-benzyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	429.3

Compound Number	Name	MS (*m/z*, M + 1)
43	*N*-(cyclopropylmethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	393.2
44	*N*-butyl-*N*-ethyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	423.3
45	*N*-octyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	451.2
46	*N*-(3,3-dimethylbutyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	423.2
47	*N*-(4-chloro-2-methylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	463.1
48	*N*-(3-methoxyphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	445.0
49	*N*-(2-fluoro-4-methylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	447.0
50	*N*-(3,4-dimethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	443.1
51	*N*-(3-chlorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	463.1
52	*N*-(3-methoxybenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	459.1
53	*N*-(3,4-difluorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	451.1
54	*N*-(3-methylbenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	443.1
55	*N*-(2-methoxybenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	459.2
56	*N*-(4-isopropylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	457.2
57	*N*-(2,3-difluorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	451.1
58	2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-(tetrahydrofuran-2-ylmethyl)acetamide	423.1
59	*N*-[2-(4-methylphenyl)ethyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	457.1

Compound Number	Name	MS (*m/z*, M + 1)
60	*N*-[2-(3-chlorophenyl)ethyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	477.1
61	*N*-(4-cyanophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	440.1
62	*N*-(2,3-dihydro-1H-inden-1-yl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	455.1
63	*N*-(2-methoxyethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	397.1
64	*N*-[2-(4-methoxyphenyl)ethyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	473.1
65	*N*-(2-cyanoethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	392.1
66	*N*-(2,4-dichlorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	497.0
67	*N*-(3,5-difluorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	465.0
68	*N*-(2,4-difluorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	465.0
69	*N*-(2-methylbenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	443.1
70	*N*-(3,4-difluorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	465.1
71	*N*-(2,5-difluorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	465.1
72	2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N,N*-dipropylacetamide	423.2
73	*N,N*-dibutyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	451.2
74	*N*-(2,6-difluorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	465.1
75	*N*-[2-(methylthio)phenyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	461.1
76	*N*-(2-isopropylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	457.2

Compound Number	Name	MS (*m/z*, M + 1)
77	*N*-(4-bromophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	493.0
78	*N*-(4-chlorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	449.0
79	*N*-(2,4-dichlorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	483.0
80	*N*-(3,5-dichlorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	483.1
81	*N,N*-diethyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	395.1
82	*N*-methyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-phenylacetamide	429.1
83	*N*-(4-hydroxybutyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	411.1
84	*N*-allyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	379.1
85	*N*-(2-fluoro-5-methylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	447.1
86	*N*-(1,3-benzodioxol-5-ylmethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	473.1
87	*N*-cyclopropyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	379.1
88	*N*-(2-cyclopropylethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	407.2
89	*N*-(3,4-dichlorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	497.1
90	*N*-(2,3-dichlorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	497.1
91	*N*-(2,5-dimethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	443.1
92	*N*-(3,4-dichlorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	483.3
93	*N,N*-dimethyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	367.2

Compound Number	Name	MS (*m/z*, M + 1)
94	*N*-methyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-(2-phenylethyl)acetamide	457.2
95	2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-(2-phenylpropyl)acetamide	457.2
96	*N*-[(1*R*)-1-cyclohexylethyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	449.2
97	*N*-[(1S)-1-cyclohexylethyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	449.2
98	2-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-(2-piperidin-1-ylethyl)acetamide	451.4
99	2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-[3-(trifluoromethyl)phenyl]acetamide	483.1
100	*N*-(3-cyanophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	440.1
101	1'-(2-morpholin-4-yl-2-oxoethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	409.1
102	2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-(2-phenylethyl)acetamide	443.1
103	*N*-(4-bromo-2-chlorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	527.0
104	*N*-(2-biphenyl-4-ylethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide	519.2

EXAMPLE 3.2

Synthesis of *N*-(2-fluorophenyl)-2-(2'-oxo-6,7-dihydro-5*H*-spiro[indeno[5,6-*b*]furan-3,3'-indol]-1'(2'*H*)-yl)acetamide



To a solution of 2'-oxo-6,7-dihydro-5*H*-spiro[indeno[5,6-*b*]furan-3,3'-indol]-

1'(2'*H*)-yl)acetic acid (0.18 g, 0.54 mmol) in chloroform (5.00 mL) was added oxalyl chloride (0.09 mL, 1.07 mmol) with one drop of DMF. The mixture was refluxed for 2 hours, and evaporated under reduced vaccum to dryness. To the above residue were added Et_3N (0.66 mL, 4.72 mmol), 2-fluoroaniline (0.10 mL, 1.00 mmol) and THF (5.00 mL). The reaction mixture was stirred at ambient temperature overnight, evaporated to dryness. The residue was subjected to column chromatography (25% ethyl acetate in hexane) to yield the title compound (0.04 g, 17%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.26 (t, 1H), 8.05-7.90 (br, 1H), 7.33 (td, 1H), 7.26-6.96 (m, 6H), 6.83 (s, 1H), 6.61 (s, 1H), 4.98 (d, 1H), 4.73 (d, 1H), 4.71 (d, 1H), 4.52 (d, 1H), 2.85 (t, 2H), 2.69 (t, 2H), 2.12-1.94 (m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.6, 165.0, 159.8, 146.9, 141.5, 137.5, 132.5, 129.2, 126.3, 125.2, 124.8, 124.4, 124.3, 121.9, 118.9, 115.2, 114.9, 109.0, 106.8, 79.9, 58.2, 45.3, 33.2, 32.0, 26.1; MS (ES+) *m/z* 429 (M + 1), 451 (M + 23).

EXAMPLE 3.3

Synthesis of *N*-(2-fluorophenyl)-2-(2-oxo-5',6',7',8'-tetrahydrospiro[indole-3,3'-naphtho[2,3-*b*]furan]-1(2*H*)-yl)acetamide



Following the procedure as described in EXAMPLE 3.2, and making non-critical variations using (2-oxo-5',6',7',8'-tetrahydrospiro[indole-3,3'-naphtho[2,3-*b*]furan]-1(2*H*)-yl)acetic acid to replace 2'-oxo-6,7-dihydro-5*H*-spiro[indeno[5,6-*b*]furan-3,3'-indol]-1'(2'*H*)-yl)acetic acid, the title compound was obtained in 5% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.27 (t, 1H), 8.04-7.90 (br, 1H), 7.33 (td, 1H), 7.26-6.97 (m, 6H), 6.69 (s, 1H), 6.48 (s, 1H), 4.94 (d, 1H), 4.69 (d, 1H), 4.71 (d, 1H), 4.52 (d, 1H), 2.81-2.45 (m, 4H), 1.82-1.60 (m, 4H); MS (ES+) *m/z* 443 (M + 1), 465 (M + 23).

EXAMPLE 3.4

Synthesis of 2-(4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide



A. Synthesis of (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid

Following the procedure as described in EXAMPLE 2, and making non-critical variations using ethyl (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetate to replace ethyl (2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetate, the title compound was obtained in 100% yield. The product was used directly in the next step.

B. Synthesis of 2-(4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide

To a solution of (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid (0.24 g, 0.59 mmol) and oxalyl chloride (0.15 mL, 1.76 mmol) in toluene (7.00 mL) was added one drop of DMF and the resulted mixture was stirred at ambient temperature overnight. The mixture was concentrated under vacuum. The residue was dissolved in dichloromethane (5.00 mL) and 2-fluoroaniline (0.18 mL, 1.89 mmol) was added at ambient temperature. The mixture was stirred at ambient temperature for one hour. More dichloromethane (100 mL) was added. The organic layer was washed with water, dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo*. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/5) to give the title compound (0.23 g, 76%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.20 (t, 1H), 7.80 (br, 1H), 7.27-7.17 (m, 2H), 7.16-7.03 (m, 3H), 6.97-6.88 (m, 1H), 6.78-6.60 (m, 2H), 5.08 (d, 1H), 4.93 (d, 1H), 4.68 (d, 1H), 4.49 (d, 1H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.1, 163.9, 157.8, 143.6, 131.0, 128.8, 128.1, 124.7, 121.8, 120.0, 115.0, 111.6, 108.1, 99.8, 77.5, 59.2, 44.7; MS (ES+) *m/z* 503.4 (M + 1), 505.4 (M + 1).

EXAMPLE 3.5

Synthesis of 2-(4'-bromo-6,6-dimethyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide



A. Synthesis of (4'-bromo-6,6-dimethyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid

Following the procedure as described in EXAMPLE 2, and making non-critical variations using ethyl (4'-bromo-6,6-dimethyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetate to replace ethyl (2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetate, the title compound was obtained. The product was used directly in the next step.

B. Synthesis of 2-(4'-bromo-6,6-dimethyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide

Following the procedure as described in EXAMPLE 3.4B, and making non-critical variations using (4'-bromo-6,6-dimethyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid to replace (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2' *H*)-yl)acetic acid, the title compound was obtained (61% for two steps): MS (ES+) *m/z* 537.4 (M + 1), 539.4 (M + 1).

EXAMPLE 3.6

Synthesis of 2-(4'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide



Following the procedure as described in EXAMPLE 3.4B, and making non-critical variations using (4'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid to replace (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2' *H*)-yl)acetic acid, the title compound was obtained (69%) as a colorless solid: mp 243-245 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.19 (t, 1H), 7.96 (s, 1H), 7.21-7.27 (m, 1H), 7.10-7.02 (m, 4H), 6.88 (d, 1H), 6.55 (s, 1H), 6.35 (s, 1H), 4.96 (dd, 2H), 4.70 (d, 1H), 4.57-4.53 (m, 3H), 2.97 (t, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.2, 164.4, 162.2, 162.1, 143.2, 131.6, 130.2, 128.5, 125.2, 125.1, 124.9, 124.6, 121.9, 119.7, 118.6, 116.8, 115.1, 114.8, 107.3, 92.9, 77.2, 72.4, 58.2, 44.9, 28.9; MS (ES+) *m/z* 465.5 (M + 1).

EXAMPLE 3.7

Synthesis of 2-(5'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide



Following the procedure as described in EXAMPLE 3.4B, and making non-critical variations using (5'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid to replace (4'-bromo-5,6-difluoro-2'-oxospiro[1-

benzofuran-3,3'-indol]-1'(2' *H*)-yl)acetic acid, the title compound was obtained (91%) as a colorless solid: mp 229-230 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.21 (t, 1H), 7.88 (s, 1H), 7.28-7.25 (m, 1H), 7.18 (d, 1H), 7.13-7.04 (m, 3H), 6.90 (d, 1H), 6.57 (s, 1H), 6.40 (s, 1H), 4.95 (d, 1H), 4.70-4.66 (m, 2H), 4.56-4.43 (m, 3H), 2.99 (t, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.8, 170.9, 162.1, 162.1, 143.1, 131.7, 130.0, 128.6, 124.7, 119.6, 118.7, 117.0, 106.7, 92.8, 77.2, 72.3, 58.1, 41.2, 28.9; MS (ES+) *m/z* 465.4 (M + 1).

EXAMPLE 3.8

Synthesis of 2-(6-chloro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide



Following the procedure as described in EXAMPLE 3.4B, and making non-critical variations using (6-chloro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid to replace (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2' *H*)-yl)acetic acid, the title compound was obtained (10%) as a white solid: mp 70-75 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.21 (t, 1H), 7.99 (br, 1H), 7.32 (dt, 1H), 7.19-6.93 (m, 7H), 6.80 (dd, 1H), 6.73 (d, 1H), 5.01 (d, 1H), 4.75 (d, 1H), 4.69 (d, 1H), 4.50 (d, 1H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.7, 164.6, 161.4, 141.4, 135.5, 131.6, 129.4, 127.3, 125.2, 125.0, 124.7, 124.6, 124.4, 124.3, 124.1, 121.9, 121.8, 115.1, 114.8, 111.3, 109.0, 80.3, 57.6, 44.9; MS (ES+) *m/z* 423.4 (M + 1).

EXAMPLE 3.9

Synthesis of 2-(5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide



Following the procedure as described in EXAMPLE 3.4B, and making non-critical variations using (5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid to replace (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid, the title compound was obtained (35%) as a white solid: mp 97-100 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.20-8.10 (m, 2H), 7.31 (dt, 1H), 7.19-7.00 (m, 5H), 6.95 (d, 1H), 6.77-6.40 (m, 2H), 5.02 (d, 1H), 4.74 (d, 1H), 4.69 (d, 1H), 4.51 (d, 1H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.4, 164.3, 141.4, 131.2, 129.7, 125.6, 125.5, 125.3, 125.2, 125.1, 124.7, 124.4, 124.1, 121.8, 115.1, 114.9, 112.1, 111.8, 109.1, 100.3, 100.0, 80.7, 57.9; MS (ES+) *m/z* 425.5 (M + 1).

EXAMPLE 3.10

Synthesis of 2-(5-bromo-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide



Following the procedure as described in EXAMPLE 3.4B, and making non-critical variations using (5-bromo-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid to replace (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid, the title compound was obtained (94%) as a light yellow solid: mp 100-

103 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.26 (dd, 1H), 7.92 (br, 1H), 7.39-7.31 (m, 2H), 7.22-7.01 (m, 6H), 6.91 (d, 1H), 6.87 (d, 1H), 5.02 (d, 1H), 4.76 (d, 1H), 4.67 (d, 1H), 4.57 (d, 1H); MS (ES+) *m/z* 467.3 (M + 1).

EXAMPLE 3.11

Synthesis of 2-(4'-fluoro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide



Following the procedure as described in EXAMPLE 3.4B, and making non-critical variations using (4'-fluoro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid to replace (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid, the title compound was obtained: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.20 (t, 1H), 7.91 (s, 1H), 7.29 (dt, 1H), 7.13-7.04 (m, 3H), 6.81-6.75 (m, 2H), 6.61 (s, 1H), 6.39 (s, 1H), 4.95-4.87 (m, 2H), 4.70 (d, 1H), 4.55-4.44 (m, 3H), 2.98 (t, 2H); MS (ES+) *m/z* 449.5 (M + 1)

EXAMPLE 3.12

Synthesis of 2-(4'-bromo-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide



Following the procedure as described in EXAMPLE 3.4B, and making non-critical variations using (4'-bromo-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-

3,3'-indol]-1'(2'*H*)-yl)acetic acid to replace (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid, the title compound was obtained (75%) as a colorless solid: mp 245-246 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.20 (t, 1H), 7.90 (s, 1H), 7.20-7.03 (m, 5H), 6.92 (dd, 1H), 6.53 (s, 1H), 6.36 (s, 1H), 5.05 (d, 1H), 4.90 (d, 1H), 4.69 (d, 1H), 4.55-4.43 (m, 3H), 2.97 (t, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.3, 164.3, 162.4, 162.2, 143.4, 130.4, 130.0, 128.0, 125.5, 125.2, 124.7, 121.9, 120.0, 119.7, 118.6, 116.7, 115.1, 114.8, 107.8, 92.8, 77.2, 72.4, 59.1, 44.9, 28.9; MS (ES+) *m/z* 509 (M + 1), 511 (M + 1).

EXAMPLE 3.13

Synthesis of *N*-(2-fluorophenyl)-2-(2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)acetamide



Following the procedure as described in EXAMPLE 3.4B, and making non-critical variations using (2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid to replace (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid, the title compound was obtained (68%) as a white solid: mp 210-212 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.18 (s, 1H), 8.17-7.67 (m, 1H), 7.55-6.90 (m, 7H), 6.54 (s, 1H), 6.38 (s, 1H), 4.68 (m, 4H), 4.46 (t, 2H), 2.93 (t, 2H); MS (ES+) *m/z* 431.4 (M + 1).

EXAMPLE 3.14

Synthesis of 2-(4'-fluoro-7'-methyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide



Following the procedure as described in EXAMPLE 3.4B and making non-critical variations using (4'-fluoro-7'-methyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid to replace (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid, the title compound was obtained (21%) as a white solid: mp 250-255 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.23 (s, 1H), 7.99-7.69 (m, 1H), 7.36-7.01 (m, 4H), 6.74 (t, 1H), 6.63 (s, 1H), 6.42 (s, 1H), 5.90 (d, 2H), 4.88 (t, 2H), 4.76 (ABq, 2H), 3.30 (s, 3H); MS (ES+) *m/z* 463.4 (M + 1).

EXAMPLE 4

Synthesis of 4'-[6-(dimethylamino)pyridin-3-yl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To an oven-dried flask was charged with [6-(dimethylamino)pyridin-3-yl]boronic acid (37.0 mg, 0.17 mmol), $Pd(PPh_3)_4$ (13.5 mg, 0.012 mmol) followed by flashing with nitrogen. To the flask was added a solution of 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (50.0 mg, 0.12 mmol) in anhydrous dioxane (2.00 mL) followed by the addition of 2.0 M Na_2CO_3 (0.24 mL). The reaction mixture was heated at reflux for 48 h. After cooling down to ambient temperature, the solvent

was removed under reduced pressure. The residue was diluted with ethyl acetate (2.00 mL), washed with saturated ammonium chloride (2.00 mL), and concentrated *in vacuo* to dryness. The residue was subjected to column chromatography to yield the title compounds: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.77 (d, 1H), 7.32 (t, 1H), 6.87 (dd, 2H), 6.71 (dd, 1H), 6.23 (d, 1H), 6.20 (s, 1H), 5.88 (d, 2H), 4.56 (ABq, 2H), 3.89-3.80 (m, 1H), 3.69-3.59 (m, 1H), 3.05 (s, 6H), 1.78-1.69 (m, 2H), 1.39-1.35 (m, 4H), 0.90 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.9, 158.3, 156.1, 148.8, 147.3, 142.9, 142.0, 137.8, 137.2, 129.9, 128.9, 125.6, 122.0, 121.0, 107.6, 104.3, 102.5, 101.5, 93.6, 77.8, 58.5, 40.5, 38.2, 29.1, 27.2, 22.4, 14.0; MS (ES+, *m/z*) 472.0 (M + 1).

EXAMPLE 4.1

The compounds listed in the following table were synthesized using similar conditions as described in Example 4. As previously noted, the compound numbers listed below do not correspond to the compound numbers provided in the general Reaction Schemes above.

Compound Number	Name	MS (*m/z*, M + 1)
105	4'-(3,5-dimethoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	488.4
106	4'-(4-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	447.3
107	4'-(3,5-dichlorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	496.3
108	4'-[4-(dimethylamino)phenyl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	471.3
109	1'-pentyl-4'-(3,4,5-trimethoxyphenyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	518.3
110	4-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)benzonitrile	453.3
111	4'-dibenzo[b,d]furan-4-yl-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	518.3
112	4'-(1-benzyl-1*H*-pyrazol-4-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	508.4

Compound Number	Name	MS (*m/z*, M + 1)
113	4'-(2-methoxypyrimidin-5-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'H)-one	460.3
114	4'-(2,4-dimethoxypyrimidin-5-yl)-1'-pentylspiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	490.4
115	4-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)benzamide	471.3
116	4'-{4-[(dimethylamino)methyl]phenyl}-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	484.6
117	4'-(1-benzofuran-2-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	467.5
118	4'-(6-methoxypyridin-3-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.5
119	*N,N*-dimethyl-4-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)benzamide	498.6
120	4'-dibenzo[*b,d*]thien-4-yl-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	533.6
121	3-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)benzonitrile	452.5
122	1'-pentyl-4'-pyridin-3-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	428.5
123	4'-(3-fluoro-4-methoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	475.5
124	1'-pentyl-4'-[2-(trifluoromethoxy)phenyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	511.5
125	4'-[3,5-bis(trifluoromethyl)phenyl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	563.5
126	1'-pentyl-4'-[4-(trifluoromethyl)pyridin-3-yl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	496.5

Compound Number	Name	MS (*m/z*, M + 1)
127	4'-(2-fluoro-5-methoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	475.5
128	4'-(4-ethoxy-3-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	489.5
129	4'-(1-benzothien-2-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	483.6
130	4'-isobutyl-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	407.5
131	1'-pentyl-4'-[4-(trifluoromethoxy)phenyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	511.5
132	4'-(5-fluoro-2-methoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	475.5
133	4'-(1,3-benzodioxol-5-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	471.5
134	1'-pentyl-4'-phenylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	427.5
135	1'-pentyl-4'-[2-(trifluoromethyl)phenyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	495.5
136	4'-(4-chlorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	461.9
137	4'-(2,3-dihydro-1,4-benzodioxin-6-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	485.5
138	1'-pentyl-4'-quinolin-3-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	478.5
139	4'-(3,5-difluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	463.5
140	4'-isoquinolin-4-yl-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	478.5
141	4'-(6-methoxypyridin-2-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.5

Compound Number	Name	MS (*m/z*, M + 1)
142	4'-(1*H*-indol-5-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	466.5
143	*N*-[2-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)phenyl]acetamide	484.5
144	4'-(4-fluoro-2-methylphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	459.5
145	1'-pentyl-4'-quinolin-6-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	478.5
146	*N*-[4-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)phenyl]methanesulfonamide	520.6
147	4'-(5-chloro-2-methoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	492.0
148	1'-pentyl-4'-[3-(trifluoromethoxy)phenyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	511.5
149	1'-pentyl-4'-(4-phenoxyphenyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	519.6
150	4'-(2,4-dimethoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	487.5
151	4'-(3-furyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	417.5
152	4'-(3,4-dimethoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	487.5
153	*N*-[4-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)phenyl]acetamide	484.5
154	1'-pentyl-4'-[(*E*)-2-phenylvinyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	453.5
155	4'-(4-methoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	457.5

Compound Number	Name	MS (*m/z*, M + 1)
156	4'-(6-fluoropyridin-3-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	446.5
157	4'-(3-chloro-4-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	479.9
158	4'-(3-chlorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	461.9
159	4'-(1-benzothien-3-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	483.6
160	1'-pentyl-4'-(2-phenoxyphenyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	519.6
161	4'-(4-isopropoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	485.6
162	4'-[(*E*)-2-(4-fluorophenyl)vinyl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	471.5
163	4'-(6-fluoropyridin-2-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	446.5
164	1'-pentyl-4'-[1-(phenylsulfonyl)-1*H*-indol-3-yl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	606.7
165	4'-(3-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	445.5
166	4'-(3-acetylphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	469.5
167	4'-(2-furyl)-1'-pentylspiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'H)-one	417.5
168	4'-(4-methylphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	441.5
169	4'-(1-methyl-1H-pyrrol-3-yl)-1'-pentylspiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'H)-one	430.5
170	4'-(2,5-difluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	463.5

Compound Number	Name	MS (*m/z*, M + 1)
171	4'-(2-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	445.5
172	4'-(2-chlorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'H)-one	461.9
173	4'-(2,4-difluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	463.5
174	4'-(4-morpholin-4-ylphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	512.6
175	tert-butyl 5-methoxy-3-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)-1*H*-indole-1-carboxylate	596.7
176	1'-pentyl-4'-pyrimidin-5-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	429.5
177	tert-butyl 4-[2-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)phenyl]piperazine-1-carboxylate	611.7
178	4'-(2-methoxypyridin-3-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.5
179	4'-(5-methoxypyridin-3-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.5
180	4'-(4-butoxy-3-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	517.6
181	1'-pentyl-4'-pyridin-4-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	428.5
182	1'-pentyl-4'-phenoxathiin-4-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	549.6
183	4'-[(1*Z*)-3-chloroprop-1-en-1-yl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	425.9
184	1'-pentyl-4'-(3-thienyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	433.5
185	4'-(2,3-dimethoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	487.5

Compound Number	Name	MS (*m/z*, M + 1)
186	4'-(4-butylphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	483.6
187	4'-(3-fluoro-5-methoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	475.5
188	4'-[3-fluoro-4-(pentyloxy)phenyl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	531.6
189	4'-(2-butoxy-5-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	517.6
190	4'-(3-butoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	499.6
191	4'-(4-butoxyphenyl)-1'-pentylspiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	499.6
192	4'-(4-isobutoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	499.6
193	4'-{2-chloro-4-[(3,5-dimethoxybenzyl)oxy]phenyl}-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	628.1
194	4'-[4-(benzyloxy)-3-chlorophenyl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	568.1
195	4'-(1-methyl-1*H*-indol-5-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	480.6
196	4'-(4-methoxypyridin-3-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.5
197	4'-[(6-methoxypyridin-3-yl)amino]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	473.5

EXAMPLE 4.2

Synthesis of 2-(5,6-difluoro-2'-oxo-4'-pyrimidin-5-ylspiro[1-benzofuran-3,3'-indol]-1'(2' *H*)-yl)-*N*-(2-fluorophenyl)acetamide



To a solution of 2-(4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2' *H*)-yl)-*N*-(2-fluorophenyl)acetamide (0.15 g, 0.30 mmol) in anhydrous 1,4-dioxane (5.00 mL) was added $Pd(PPh_3)_4$ (0.03 g, 0.03 mmol) and stirred at ambient temperature for 10 min. Pyrimidine-5-boronic acid (0.06 g, 0.45 mmol) and sodium carbonate (0.90 mL of 2 M solution, 1.80 mmol) were added. The reaction mixture was reluxed at 120 °C for 16 h, diluted with ethyl acetate (50.0 mL). The organic layer was washed with water, dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo*. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/1) to give the title compound (0.13 g, 84%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.13 (s, 1H), 8.29-8.10 (m, 3H), 7.62 (s, 1H), 7.44 (t, 1H), 7.16-7.03 (m, 4H), 6.91 (d, 1H), 6.85-6.76 (m, 1H), 6.46-6.37 (m, 1H), 4.85-4.73 (m, 2H), 4.61-4.47 (m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.1, 163.9, 157.7, 156.4, 155.7, 141.9, 132.9, 130.0, 126.0, 124.7, 121.9, 115.0, 111.6, 109.8, 100.2, 79.4, 57.9, 44.7; MS (ES+) *m/z* 503.5 (M + 1).

EXAMPLE 4.3

Synthesis of 2-(6,6-dimethyl-2'-oxo-4'-pyrimidin-5-yl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide



Following the procedure as described in EXAMPLE 4.2, making variation using 2-(4'-bromo-6,6-dimethyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide to replace 2-(4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide, the title compound was obtained (95%): mp > 250 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.10 (s, 1H), 8.24 (t, 1H), 8.15 (s, 2H), 8.00 (s, 1H), 7.40 (t, 1H), 7.16-7.03 (m, 4H), 6.88 (d, 1H), 6.61 (s, 1H), 5.99 (s, 1H), 4.84-4.73 (m, 2H), 4.54 (d, 1H), 4.44 (d, 1H), 2.79 (s, 2H), 1.45 (s, 3H), 1.39 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.4, 164.5, 161.1, 157.4, 155.8, 154.1, 150.8, 141.7, 132.9, 132.2, 131.3, 129.4, 125.9, 125.5, 125.2, 124.7, 121.9, 120.8, 119.3, 118.8, 115.0, 109.5, 93.5, 88.5, 78.9, 58.0, 45.0, 42.0, 28.0, 27.9; MS (ES+) *m/z* 537.5 (M + 1).

EXAMPLE 4.4

Synthesis of 4'-(3-furyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4, making variations using 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 3-furanboronic acid to replace [6-(dimethylamino)pyridin-3-yl]boronic acid, the title compound was

obtained (66%) as a colorless solid: mp 270-272 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.32 (t, 1H), 7.25 (d, 2H), 6.97 (d, 1H), 6.91 (d, 1H), 6.83 (s, 1H), 6.44 (s, 1H), 6.30 (s, 1H), 6.04 (d, 1H), 5.89 (dd, 2H), 4.68 (ABq, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 180.4, 156.2, 149.2, 142.9, 142.1, 141.2, 140.4, 131.3, 129.2, 128.5, 125.3, 122.6, 120.6, 111.0, 109.5, 102.9, 101.6, 94.0, 77.2, 59.0; MS (ES+) *m/z* 348.4 (M + 1).

EXAMPLE 4.5

Synthesis of 4'-dibenzo[*b*,*d*]furan-4-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4, and making non-critical variations using 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and dibenzo[*b*,*d*]furan-4-ylboronic acid to replace [6-(dimethylamino)pyridin-3-yl]boronic acid, the title compound was obtained (10%) as a colorless solid: mp >230 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.77 (s, 1H), 8.04 (d, 1H), 7.97 (dd, 1H), 7.43-7.42 (m, 3H), 7.34-7.29 (m, 3H), 7.16 (t, 1H), 6.98 (d, 1H), 6.89 (d, 1H), 6.25 (s, 1H), 5.69 (d, 2H), 4.41 (ABq, 2H); MS (ES+) *m/z* 448.5 (M + 1).

EXAMPLE 4.6

Synthesis of 4'-(6-methoxypyridin-3-yl)-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4, and making non-critical variations using 4'-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and (6-methoxypyridin-3-yl)boronic acid to replace [6-(dimethylamino)pyridin-3-yl]boronic acid, the title compound was obtained (51%) as a colorless solid: mp 174-176 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.73 (d, 1H), 7.69 (d, 1H), 7.34 (t, 1H), 7.03 (d, 1H), 6.89-6.86 (m, 2H), 6.65 (d, 1H), 6.61 (d, 1H), 6.56 (d, 1H), 6.17 (d, 2H), 5.87 (d, 2H), 4.99 (ABq, 2H), 4.56 (ABq, 2H), 3.90 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.7, 163.5, 158.4, 156.1, 151.8, 151.8, 149.2, 147.8, 146.1, 142.3, 141.6, 138.8, 136.9, 132.6, 130.0, 129.1, 127.8, 127.2, 126.2, 119.9, 112.7, 112.7, 110.9, 109.6, 109.5, 108.5, 102.3, 101.6, 93.6, 78.2, 58.5, 53.6, 37.0; MS (ES+) *m/z* 537.4 (M + 1).

EXAMPLE 4.7

Synthesis of 4'-[6-(dimethylamino)pyridin-3-yl]-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4, and making non-critical variations using 4'-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, the title compound was obtained (37%) as a colorless solid: mp 174-176 °C; MS (ES+) *m/z* 550.4 (M + 1).

EXAMPLE 4.8

Synthesis of 4'-pyrimidin-5-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4, and making non-critical variations using 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and pyrimidin-5-ylboronic acid to replace [6-(dimethylamino)pyridin-3-yl]boronic acid, the title compound was obtained (32%) as a colorless solid: mp 185-187 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.85 (s, 1H), 9.02 (s, 1H), 8.19 (s, 2H), 7.33 (t, 1H), 7.00 (d, 1H), 6.83 (d, 1H), 6.37 (s, 1H), 6.19 (s, 1H), 5.89 (d, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 178.7, 157.5, 156.0, 155.6, 148.8, 142.3, 133.1, 132.6, 131.8, 129.5, 124.5, 120.3, 110.9, 103.2, 101.9, 93.3, 79.5, 66.8, 58.5; MS (ES+) *m/z* 360.4 (M + 1).

EXAMPLE 4.9

Synthesis of 4'-(3-furyl)-1'-(pyridin-2-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4, and making non-critical variations using 4'-bromo-1'-(pyridin-2-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 3-furanboronic acid to replace [6-(dimethylamino)pyridin-3-yl]boronic acid, the title compound was obtained (75%) as a colorless solid: mp 195-197 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.58 (d, 1H), 7.68 (t, 1H), 7.32-7.21 (m, 4H), 6.97 (d, 1H), 6.92 (d, 1H), 6.77 (s, 1H), 6.60 (s, 1H), 6.35 (s, 1H), 5.99 (s, 1H), 5.12 (ABq, 2H), 4.71 (ABq, 2H), 4.57 (t, 2H), 3.03 (t, 2H); MS (ES+) *m/z*

437.4 (M + 1).

EXAMPLE 4.10

Synthesis of 1'-(pyridin-2-ylmethyl)-4'-pyrimidin-5-yl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4, and making non-critical variations using 4'-bromo-1'-(pyridin-2-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and pyrimidin-5-ylboronic acid to replace [6-(dimethylamino)pyridin-3-yl]boronic acid, the title compound was obtained (16%) as a colorless solid: mp >200 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.09 (s, 1H), 8.59 (d, 1H), 8.14 (s, 2H), 7.74 (t, 1H), 7.67 (d, 1H), 7.63 (d, 1H), 7.52 (d, 1H), 7.47-7.42 (m, 1H), 7.37 (d, 1H), 7.30 (d, 1H), 7.09 (d, 1H), 6.82 (d, 1H), 6.62 (s, 1H), 6.07 (s, 1H), 5.18 (ABq, 2H), 4.62 (ABq, 2H), 4.62-4.48 (m, 2H), 3.02 (t, 2H); MS (ES+) *m/z* 449.5 (M + 1).

EXAMPLE 4.11

Synthesis of 4'-pyridin-3-yl-1'-(pyridin-2-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4, and making non-critical variations using 4'-bromo-1'-(pyridin-2-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-pentylspiro[furo[2,3-

f][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and pyridin-3-ylboronic acid to replace [6-(dimethylamino)pyridin-3-yl]boronic acid, the title compound was obtained (9%) as a colorless solid: mp >200 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.52 (d, 1H), 8.42 (d, 1H), 7.96 (s, 1H), 7.79 (t, 1H), 7.40 (d, 1H), 7.29 (t, 2H), 7.18-7.08 (m, 2H), 7.00 (d, 1H), 6.81 (d, 1H), 6.72 (s, 1H), 5.98 (s, 1H), 5.08 (ABq, 2H), 4.56-4.40 (m, 4H), 3.10-2.90 (m, 2H); MS (ES+) *m/z* 448.5 (M + 1).

EXAMPLE 4.12

Synthesis of 4'-(3-furyl)-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4, and making non-critical variations using 4'-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 3-furylboronic acid to replace [6-(dimethylamino)pyridin-3-yl]boronic acid, the title compound was obtained (27%) as a colorless solid: mp 167-169 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.34-7.29 (m, 2H), 7.01 (dd, 1H), 6.95 (dd, 1H), 6.77 (dd, 1H), 6.74 (dd, 1H), 6.51 (s, 1H), 6.41 (d, 1H), 6.34 (s, 1H), 6.00 (dd, 1H), 4.97 (ABq, 2H), 4.67 (ABq, 2H), 4.56 (t, 2H), 3.01 (t, 2H); MS (ES+) *m/z* 494.4 (M + 1).

EXAMPLE 4.13

Synthesis of 4'-quinolin-3-yl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4, and making non-critical variations using 4'-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, quinolin-3-ylboronic acid to replace [6-(dimethylamino)pyridin-3-yl]boronic acid, the title compound was obtained (50%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.52 (d, 1H), 8.09 (d, 1H), 7.74-7.68 (m, 1H), 7.56-7.50 (m, 1H), 7.42-7.39 (m, 2H), 7.32 (s, 1H), 7.09 (d, 1H), 7.01 (d, 1H), 6.78-6.77 (m, 1H), 6.45 (d, 1H), 6.26 (s, 1H), 5.94 (d, 1H), 5.91 (s, 1H), 5.89 (d, 1H), 5.03 (ABq, 2H), 4.52 (ABq, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.4, 156.0, 151.8, 150.0, 149.3, 147.0, 142.4, 141.8, 136.8, 135.8, 131.1, 130.2, 129.9, 129.3, 129.0, 128.8, 128.0, 127.1, 127.0, 126.4, 126.0, 120. 4, 112.7, 109.5, 108.8, 102.5, 101.7, 93.7, 78.3, 58.5, 37.1; MS (ES+) *m/z* 557.4 (M + 1).

EXAMPLE 4.14

Synthesis of 4'-pyrimidin-5-yl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one



A mixture of 4'-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one (0.11 g, 0.21 mmol),

pyrimidine-5-boronic acid (0.04 g, 0.33 mmol), tetrakis(triphenylphosphine)palladium(0) (0.03 g, 0.02 mmol), 2.00 M sodium carbonate (1.00 mL) and 1,2-dimethoxyethane (10.0 mL) was heated at reflux for 16 h under nitrogen. After the organic solvent was evaporated *in vacuo*, the black residue was extracted with ethyl acetate (3 x 35.0 mL). The combined organic layers was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo*. The residue was subjected to column chromatography eluting with ethyl acetate:hexane (35%) to afford the title compound (0.03 g, 26%): mp 263-266 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.10 (s, 1H), 8.14 (s, 2H), 7.39 (t, 1H), 7.10, (d, 1H), 6.87 (d, 1H), 6.76 (s, 1H), 6.51 (s, 1H), 6.46 (s, 1H), 6.07 (s, 1H), 5.03 (ABq, 2H), 4.62-4.48 (m, 2H), 4.58 (ABq, 2H), 3.01 (t, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.3, 162.5, 161.0, 157.4, 155.9, 151.6, 142.2, 141.7, 132.8, 132.3, 131.3, 129.3, 125.7, 120.6, 120.2, 118.3, 117.0, 112.7, 109.7, 109.4, 93.5, 78.9, 72.5, 57.7, 37.1, 28.9; MS (ES+) *m/z* 506.5 (M + 1).

EXAMPLE 4.15

Synthesis of *tert*-butyl 4-[(2'-oxo-4'-pyrimidin-5-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]piperidine-1-carboxylate



Following the procedure as described in EXAMPLE 4.14, and making non-critical variations using *tert*-butyl 4-[(4'-bromo-2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]piperidine-1-carboxylate to replace 4'-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, the title compound was obtained (91%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.13 (s, 1H), 8.23 (s, 2H), 8.24-7.37 (m, 5H), 7.01 (d, 1H), 6.85 (d, 1H), 6.14 (dd2H), 5.91 (d, 2H), 4.55 (ABq, 2H), 4.15 (d, 2H), 3.84-3.58 (m, 3H), 2.69 (t, 2H), 1.44 (s, 9H); MS (ES+) *m/z* 557.5 (M + 1).

EXAMPLE 4.16

Synthesis of 1'-methyl-4'-pyrimidin-5-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4.14, and making non-critical variations using 4'-bromo-1'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one, the title compound was obtained (22%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.10 (s, 1H), 8.18 (s, 2H), 7.41 (t, 1H), 7.00 (d, 1H), 6.85 (d, 1H), 6.20 (s, 1H), 6.12 (s, 1H), 5.87 (d, 2H), 4.54 (ABq, 2H), 3.32 (s, 1H); MS (ES+) *m/z* 374.5 (M + 1).

EXAMPLE 4.17

Synthesis of 4'-(3-furyl)-1'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4.14, and making non-critical variations using 4'-bromo-1'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one, and 3-furanboronic acid to replace pyrimidine-5-boronic acid, the title compound was obtained (81%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.40-7.28 (m, 2H), 7.05-6.95 (m, 1H), 6.90-6.78 (m, 2H), 6.45-6.38 (m, 1H), 6.23-6.16 (m, 1H), 6.07-5.97 (m, 1H), 5.97-5.80 (m, 2H), 4.75-4.50 (m, 2H), 3.30-3.22 (m, 3H); MS (ES+) *m/z* 362.4 (M + 1).

EXAMPLE 4.18

Synthesis of 4'-(6-fluoropyridin-3-yl)-1'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4.14, and making non-critical variations using 4'-bromo-1'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one, and (6-fluoropyridin-3-yl)boronic acid to replace pyrimidine-5-boronic acid, the title compound was obtained (100%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.75-7.65 (m, 1H), 7.42-7.30 (m, 1H), 7.19-7.05 (m, 1H), 7.00-6.90 (m, 1H), 6.90-6.80 (m, 1H), 6.78-6.64 (m, 1H), 6.24-6.12 (m, 2H), 5.92-5.79 (m, 2H), 4.74-4.63 (m, 1H), 4.40-4.29 (m, 1H), 3.34-3.26 (m, 3H); MS (ES+) *m/z* 391.4 (M + 1).

EXAMPLE 4.19

Synthesis of 1'-(2-cyclopropylethyl)-4'-quinolin-3-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 4, and making non-critical variations using 4'-bromo-1'-(2-cyclopropylethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-

indol]-2'(1'*H*)-one, and quinolin-3-ylboronic acid to replace [6-(dimethylamino)pyridin-3-yl]boronic acid, the title compound was obtained: MS (ES+) *m/z* 477.5 (M + 1)

EXAMPLE 4.20

Synthesis of *N*-(2-fluorophenyl)-2-(2'-oxo-4'-pyrimidin-5-yl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetamide



Following the procedure as described in EXAMPLE 4, and making non-critical variations using 2-(4'-bromo-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide to replace 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and pyrimidine-5-boronic acid to replace [6-(dimethylamino)pyridin-3-yl]boronic acid, the title compound was obtained (53%) as a colorless solid: mp 229-230 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.10 (s, 1H), 8.26-8.15 (m, 3H), 7.98 (s, 1H), 7.41 (t, 1H), 7.15-7.05 (m, 4H), 6.89 (d, 1H), 6.68 (s, 1H), 6.06 (s, 1H), 4.81-4.76 (m, 2H), 4.59-4.42 (m, 4H), 3.00 (t, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.2, 164.4, 162.5, 161.0, 157.0, 155.8, 141.8, 132.6, 131.1, 129.4, 125.8, 125.5, 125.4, 125.2, 125.1, 124.7, 121.9, 120.6, 119.9, 118.7, 115.1, 114.8, 109.5, 93.4, 79.0, 72.4, 57.8, 44.9, 28.9; MS (ES+) *m/z* 509.5 (M + 1).

EXAMPLE 5

Synthesis of 4'-[(6-methoxypyridin-3-yl)amino]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To an oven-dried 2-neck 25 mL round bottom flask equipped with a condenser was charged with 4'-bromo-1'-pentylspiro-(6,7-dihydrofuro-[2,3-*f*][1,3]benzodioxole-7,3'-indole)-2'-(1'*H*)-one (50.5 mg, 0.12 mmol), 5-amino-2-methoxypyridine (22.3 mg, 0.18 mmol), $Pd_2(dba)_3$ (10 mole%), BINAP (10 mole%) and sodium methoxide (12.9 mg, 0.24 mmol). The flask was flushed with nitrogen for 5 min followed by the addition of degassed toluene (5.00 mL). The reaction mixture was heated at reflux for 16 h. After cooling to ambient temperature, the reaction mixture was diluted with ethyl acetate (20.0 mL) and washed with saturated ammonium chloride (10.0 mL), brine (10.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The brown residue was subjected to column chromatography eluting with ethyl acetate-hexane (20% to 50%) to yield the title compound (30.0 mg) in 54% yield: MS (ES+), *m/z* 474.3 (M + 1).

EXAMPLE 5.1

The compounds listed in the following table were synthesized using similar conditions as described in EXAMPLE 5. As previously noted, the compound numbers listed below do not correspond to the compound numbers provided in the general Reaction Schemes above.

Compound Number	Name	MS (*m/z*, M + 1)
198	4'-[(3,5-difluorophenyl)amino]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	479.1
199	4'-[(4,6-dimethylpyridin-2-yl)amino]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	473.3 (M+2)

Compound Number	Name	MS (*m/z*, M + 1)
200	4'-[(4-methyl-1,3-thiazol-2-yl)amino]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	464.1

EXAMPLE 5.2

Synthesis of 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



A mixture of 4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.05 g, 0.12 mmol), 3-(trifluoromethyl)aniline (0.03 g, 0.17 mmol), $Pd_2(dba)_3$ (0.02 g, 0.01 mmol), xanthphos (0.007 g, 0.01 mmol), and sodium *tert*-butoxide (0.02 g, 0.17 mmol) in toluene (5.00 mL) was heated at 110 °C for 4 days. After cooling down to ambient temperature, the mixture was extracted with ethyl acetate. The organic phase was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate/hexane to give the title compound (0.06 g, 71%) as a solid: MS (ES+) *m/z* 511.5 (M + 1).

EXAMPLE 5.3

The compounds listed in the following table were synthesized using similar conditions as described in EXAMPLE 5.2. As previously noted, the compound numbers listed below do not correspond to the compound numbers provided in the general Reaction Schemes above.

Compound Number	Name	MS (*m/z*, M + 1)
201	4'-morpholino-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	437.5

Compound Number	Name	MS (*m/z*, M + 1)
202	4'-(4-methylpiperazin-1-yl)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	450.5
203	1'-pentyl-4'-(pyrimidin-4-ylamino)-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	445.5
204	1'-pentyl-4'-(pyridin-3-ylamino)-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	444.5
205	4'-(4-chloro-2-(trifluoromethyl)phenylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	545.5
206	1'-pentyl-4'-(pyrimidin-2-ylamino)-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	444.5
207	4'-(benzo[*d*][1,3]dioxol-5-ylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	487.5
208	4'-(3-fluorophenylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	461.5
209	4'-(naphthalen-2-ylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	493.5
210	4'-(2-methoxyphenylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	473.2
211	4'-(4-methylthiazol-2-ylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	464.1
212	4'-(4,6-dimethylpyridin-2-ylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	473.3
213	4'-(3,5-difluorophenylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	479.1
214	4'-(6-methoxypyridin-3-ylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one	474.3

EXAMPLE 6

Synthesis of 2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoic acid



To a solution of methyl 2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoate (7.56 g, 17.6 mmol) in a mixture of THF/water (2/1 v/v, 180 mL) was added lithium hydroxide monohydrate (1.48 g, 35.2 mmol). The resulting mixture was stirred at ambient temperature overnight and concentrated *in vacuo* followed by the addition of water (150 mL). The mixture was extracted with of ethyl acetate/hexanes, 1/3 v/v, 50.0 mL). The water layer was acidified with1 N HCl solution until the pH value reached 2. The precipitate was filtered and dried to give the title compound (7.30 g, 100%) as a white solid: mp > 250 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.25 (s, 1H), 7.95 (dd, 1H), 7.49 (dt, 1H), 7.37 (t, 1H), 7.24-7.16 (m, 2H), 7.11-6.98 (m, 2H), 6.80 (d, 1H), 6.68 (s, 1H), 6.36 (s, 1H), 5.91 (s, 2H), 5.37-5.19 (m, 2H), 4.88-4.68 (m, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 177.6, 168.8, 156.0, 148.8, 143.0, 142.3, 137.6, 133.1, 132.1, 131.5, 129.9, 129.4, 127.7, 126.5, 124.2, 123.6, 120.1, 109.8, 103.8, 101.9, 93.8, 80.5, 58.0, 42.6.

EXAMPLE 7

Synthesis of *N*-[2-(4-chlorophenyl)ethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide



A. Preparation of stock solution of 2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoyl chloride

A solution of 2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoic acid (0.21 g, 0.50 mmol), oxalyl chloride (0.09 mL, 1.00 mmol) and one drop of DMF in toluene (10.0 mL) was stirred at ambient temperature overnight. The mixture was concentrated under vacuum to afford a solid, which was dissolved in dichloromethane (5.00 mL) to form an acid chloride stock solution (0.10 mmol/mL) for use.

B. Synthesis of *N*-[2-(4-chlorophenyl)ethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide

The acid chloride stock solution obtained above (1.00 mL, 0.10 mmol) was added to a mixture of 2-(4-chlorophenyl)ethylamine (0.02 g, 0.13 mmol), triethylamine (0.14 mL, 1.00 mmol) in dichloromethane (1.00 mL). The resulting mixture was stirred at ambient temperature overnight and diluted with dichloromethane (5.00 mL). The mixture was washed with 1 N HCl, saturated sodium bicarbonate solution, dried over sodium sulfate and filtered. The filtrate was concentrated under vacuum to dryness to give the title compound as a white powder: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.70-7.13 (m, 10H), 7.10-6.96 (m, 2H), 6.65-6.54 (br, 1H), 6.50 (s, 1H), 6.11 (s, 1H), 5.89-5.82 (m, 2H), 5.09-4.88 (m, 3H), 4.68 (d, 1H), 3.79-3.66 (m, 2H), 2.93 (t, 2H); MS (ES+), *m/z* 553.3 (M + 1), 575.3 (M + 23).

EXAMPLE 7.1

The compounds listed in the following table were synthesized using similar conditions as described in EXAMPLE 7. As previously noted, the compound numbers listed below do not correspond to the compound numbers provided in the general Reaction Schemes above.

Compound Number	Name	MS (*m/z*, M + 1)
215	*N*-(3-methylbutyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	485.3
216	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-phenylbenzamide	491.2

Compound Number	Name	MS (*m/z*, M + 1)
217	*N,N*-diisopropyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	499.4
218	*N*-(3-fluorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	509.3
219	*N*-(4-chlorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	539.3
220	*N*-butyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	471.3
221	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pentylbenzamide	485.3
222	*N*-hexyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	499.4
223	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-propylbenzamide	457.3
224	*N*-isopropyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	457.3
225	*N*-cyclohexyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	497.3
226	*N*-cyclopentyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	483.3
227	*N*-heptyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	513.4
228	*N*-(4-fluorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	509.1
229	*N*-(3-fluorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	523.1
230	*N*-(3-chlorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	525.0

Compound Number	Name	MS (*m/z*, M + 1)
231	*N*-(2-fluorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	509.1
232	*N*-(2-ethylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.1
233	*N*-(4-ethylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.2
234	*N*-(3-methylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	505.2
235	*N*-(2,3-dimethylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.2
236	*N*-(3,5-dimethylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.1
237	1'-[2-(piperidin-1-ylcarbonyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	483.1
238	*N*-isobutyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	471.2
239	*N*-(2-chlorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	539.1
240	*N*-(2,6-dimethylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.1
241	*N*-(2-methoxyphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	521.1
242	*N*-(3-methoxypropyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	487.2

Compound Number	Name	MS (*m/z*, M + 1)
243	*N*-[2-(4-methylphenyl)ethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	533.2
244	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(tetrahydrofuran-2-ylmethyl)benzamide	499.2
245	*N,N*-dibenzyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	595.2
246	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(3-phenylpropyl)benzamide	533.2
247	*N*-[2-(3-chlorophenyl)ethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	553.1
248	*N*-[2-(4-fluorophenyl)ethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	537.1
249	*N*-(4-fluorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	523.1
250	*N*-(3-ethoxypropyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	501.1
251	*N*-hexyl-*N*-methyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	513.2
252	*N*-(3-isopropoxypropyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	515.2
253	*N*-(4-methoxybenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	535.2
254	*N*-(cyclopropylmethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	469.1

Compound Number	Name	MS (*m/z*, M + 1)
255	*N*-(2-ethoxyethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	487.2
256	*N*-(cyclohexylmethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	511.2
257	*N*-(2-furylmethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	495.0
258	*N*-(2,4-dimethylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.2
259	*N*-(4-cyanophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	516.2
260	*N*-(3,5-dichlorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	559.0
261	*N*-(3-fluoro-2-methylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	523.1
262	*N*-(4-methoxyphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	521.1
263	*N*-(5-chloro-2-methylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	539.1
264	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[3-(trifluoromethyl)phenyl]benzamide	559.1
265	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(pyridin-4-ylmethyl)benzamide	506.1
266	*N*-cyclobutyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	469.1

Compound Number	Name	MS (*m/z*, M + 1)
267	*N*-methyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	429.1
268	*N*-ethyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	443.1
269	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2,2,2-trifluoroethyl)benzamide	497.1
270	*N*-(2,2-diphenylethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	595.2
271	*N*-[2-(diethylamino)ethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	514.2
272	*N*-(3,3-dimethylbutyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	499.2
273	*N*-(2-ethylbutyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	499.2
274	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-pyrrolidin-1-ylethyl)benzamide	512.4
275	*N*-[(1-ethylpyrrolidin-2-yl)methyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	526.4
276	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-piperidin-1-ylethyl)benzamide	526.5
277	*N*-(2-morpholin-4-ylethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	528.4
278	*N*-[(1*S*)-1-cyclohexylethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	525.5
279	*N*-(2-fluoro-5-methylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	523.4

Compound Number	Name	MS (*m/z*, M + 1)
280	*N*-(2,4-difluorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	527.3
281	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(4-propylphenyl)benzamide	533.4
282	*N*-(3,3-diphenylpropyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	609.4
283	*N*-(2-methoxyethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	473.4
284	*N*-(2,5-difluorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	541.3
285	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-thienylmethyl)benzamide	511.4
286	*N*-[4-chloro-2-(trifluoromethyl)phenyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	593.3
287	*N*-[2-(4-methoxyphenyl)ethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	549.4
288	*N*-(3,5-dichlorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	573.2
289	*N*-(3-chlorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	539.3
290	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(2-thienyl)ethyl]benzamide	525.4
291	*N*-(2,3-dihydro-1*H*-inden-1-yl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	531.4

Compound Number	Name	MS (*m/z*, M + 1)
292	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[4-(trifluoromethyl)benzyl]benzamide	573.3
293	*N*-[4-fluoro-2-(trifluoromethyl)phenyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	577.5
294	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(pyridin-3-ylmethyl)benzamide	506.4
295	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(trifluoromethyl)benzyl]benzamide	573.3
296	*N*-(3-methylpyridin-2-yl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	506.4
297	*N*-(1-benzylpiperidin-4-yl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	588.4
298	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(trifluoromethyl)phenyl]benzamide	559.4
299	*N*-[(1*R*)-1-cyclohexylethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	525.5
300	*N*-(2-cyanoethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	468.3
301	*N*-(6-methoxypyridin-3-yl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	522.4
302	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-1,3-thiazol-2-ylbenzamide	498.3
303	2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-1,3,4-thiadiazol-2-ylbenzamide	499.3

Compound Number	Name	MS (*m/z*, M + 1)
304	*N*-(4,6-dimethylpyridin-2-yl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	520.3
305	*N*-(2,3-dihydro-1*H*-inden-5-yl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	531.4
306	1'-{2-[(4-pyrimidin-2-ylpiperazin-1-yl)carbonyl]benzyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	562.4
307	1'-(2-{[4-(1,3-benzodioxol-5-ylmethyl)piperazin-1-yl]carbonyl}benzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	618.4
308	*N*-2-adamantyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	549.5
309	*N*-1-adamantyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	549.5
310	*N*-1-naphthyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	541.4
311	*N*-(3,5-difluorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	527.3
312	1'-[2-(morpholin-4-ylcarbonyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	485.4
313	*N*-[3-(dimethylamino)propyl]-*N*-methyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	514.4

EXAMPLE 8

Synthesis of 3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoic acid



Following the procedure described in EXAMPLE 6, and making non-critical variations using methyl 3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoate to replace methyl 2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoate, the title compound was obtained (100%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 13.03 (s, 1H), 7.86-7.80 (m, 2H), 7.59-7.57 (m, 1H), 7.48-7.44 (m, 1H), 7.25-7.16 (m, 2H), 7.03-6.95 (m, 2H), 6.68 (s, 1H), 6.18 (s, 1H), 5.90 (s, 2H), 5.05 (ABq, 2H), 4.75 (ABq, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.4, 167.5, 155.9, 148.9, 142.5, 142.3, 137.3, 132.2, 131.7, 129.6, 129.4, 128.8, 128.1, 124.3, 123.7, 120.1, 109.9, 103.3, 101.9, 93.9, 80.3, 58.0, 43.2; MS (ES+) *m/z* 416.2 (M + 1).

EXAMPLE 9

Synthesis of *N*-[2-(3-chlorophenyl)ethyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide



A. Preparation of stock solution of 3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoyl chloride

To a stirred slurry of 3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoic acid (2.08 g, 5.00 mmol) in dry chloroform (50.0 mL) was added oxalyl chloride (0.95 g, 7.5 mmol) at ambient temperature followed by 1 drop of DMF. The mixture was stirred at ambient temperature overnight and evaporated to dryness *in vacuo*. The residue was dissolved in dry dichlormethane (60.0 mL) to form

an acid chloride stock solution for use.

B. Synthesis of *N*-[2-(3-Chlorophenyl)ethyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide

To a solution of 2-(3-chlorophenyl)ethylamine (0.02 mL, 0.24 mmol) in dry dichloromethane (2.00 mL) and triethylamine (0.05 mL, 0.32 mmol) was added the acid chloride stock solution (2.0 mL, 0.081 M in dichloromethane) obtained above at ambient temperature. The mixture was stirred for 2 h, washed with 15% HCl solution and water. The organic layer was separated, dried over Na_2SO_4 and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was dissolved in ethyl acetate, and the product was precipitated by the addition of hexane. The white solid was filtered and collected to yield the title compound (0.06 g) in 65% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.97-7.95 (m, 2H), 7.53-7.50 (m, 1H), 7.45-7.40 (m, 1H), 7.21-7.15 (m, 2H), 7.04-6.99 (m, 1H), 6.73-6.71 (m, 1H), 6.52 (s, 1H), 6.20 (s, 1H), 5.86 (s, 1H), 5.18 (d, 1H), 4.72 (d, 1H), 4.80 (d, 1H), 4.69 (d, 1H), 3.89 (s, 1H); MS (ES+) *m/z* 554.0 (M + 1).

EXAMPLE 9.1

The compounds listed in the following table were synthesized using the similar procedure as described in EXAMPLE 9. As previously noted, the compound numbers listed below do not correspond to the compound numbers provided in the general Reaction Schemes above.

Compound Number	Name	MS (*m/z*, M + 1)
314	*N*-(3-methylbutyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	485.5
315	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-phenylbenzamide	491.5
316	*N,N*-diisopropyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	499.6
317	*N*-(3-fluorophenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	509.5

Compound Number	Name	MS (*m/z*, M + 1)
318	*N*-(4-chlorobenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	540.0
319	*N*-butyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	471.5
320	*N*-(3-fluorobenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	523.5
321	*N*-(3-chlorophenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	526.0
322	*N*-(2-fluorophenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	509.5
323	*N*-(2-ethylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.6
324	*N*-(4-ethylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.6
325	*N*-(3-methylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	505.5
326	*N*-(2,3-dimethylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.6
327	*N*-(3,5-dimethylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.6
328	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pentylbenzamide	485.5
329	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-propylbenzamide	457.5

Compound Number	Name	MS (*m/z*, M + 1)
330	*N*-isopropyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	457.5
331	1'-[3-(piperidin-1-ylcarbonyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	483.5
332	*N*-isobutyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	471.5
333	*N*-hexyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	499.6
334	*N*-cyclohexyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	497.6
335	*N*-cyclopentyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	483.5
336	*N*-heptyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	513.6
337	*N*-(2-methoxybenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	535.6
338	*N*-(2-methoxyphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	521.5
339	*N*-cyclopropyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	455.5
340	*N*-(3-methoxypropyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	487.5
341	*N*-(2,4-dimethylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.6
342	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(tetrahydrofuran-2-ylmethyl)benzamide	499.5

Compound Number	Name	MS (*m/z*, M + 1)
343	*N,N*-dibenzyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	595.7
344	*N*-[2-(diethylamino)ethyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	514.6
345	*N*-methyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	429.4
346	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[3-(trifluoromethyl)phenyl]benzamide	559.5
347	*N*-ethyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	443.5
348	*N*-(3-ethoxypropyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	501.5
349	*N*-(4-methoxybenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	535.6
350	*N*-(3,5-dichlorophenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	560.4
351	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pyridin-3-ylbenzamide	492.5
352	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(pyridin-4-ylmethyl)benzamide	506.5
353	*N*-(2-furylmethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	495.5
354	*N*-(3-fluoro-2-methylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	523.5

Compound Number	Name	MS (*m/z*, M + 1)
355	*N*-hexyl-*N*-methyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	513.6
356	*N*-(3-isopropoxypropyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	515.6
357	*N*-(2-ethoxyethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	487.5
358	*N*-(cyclopropylmethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	469.5
359	*N*-(4-methoxyphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	521.5
360	*N*-cyclobutyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	469.5
361	*N*-[2-(4-fluorophenyl)ethyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	537.5
362	*N*-(cyclohexylmethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	511.6
363	*N*-[2-(4-methylphenyl)ethyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	533.6
364	*N*-(2-ethylbutyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	499.6
365	*N*-benzyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	505.5
366	*N*-(2-methoxyethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	473.5

Compound Number	Name	MS (*m/z*, M + 1)
367	1'-[3-(morpholin-4-ylcarbonyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	485.5
368	*N*-(1-benzylpiperidin-4-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	588.7
369	*N*-[2-(4-methoxyphenyl)ethyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	549.6
370	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-piperidin-1-ylethyl)benzamide	526.6
371	*N*-(1-cyclohexylethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	525.6
372	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-thienylmethyl)benzamide	511.6
373	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(4-propylphenyl)benzamide	533.6
374	*N*-(2,4-difluorobenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	541.5
375	*N*-(3,5-difluorophenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	527.5
376	*N*-(2,3-dihydro-1*H*-inden-5-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	531.6
377	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[4-(trifluoromethyl)benzyl]benzamide	573.5
378	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(2-thienyl)ethyl]benzamide	525.6

Compound Number	Name	MS (*m/z*, M + 1)
379	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(pyridin-3-ylmethyl)benzamide	506.5
380	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(trifluoromethyl)benzyl]benzamide	573.5
381	*N*-[2-(4-chlorophenyl)ethyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	554.0
382	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-pyrrolidin-1-ylethyl)benzamide	512.6
383	*N*-(3-methylpyridin-2-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	506.5
384	*N*-1,3-benzodioxol-5-yl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	535.5
385	*N*-(2-morpholin-4-ylethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	528.6
386	1'-{3-[(4-pyrimidin-2-ylpiperazin-1-yl)carbonyl]benzyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	562.6
387	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-1,3-thiazol-2-ylbenzamide	498.5
388	*N*-(6-methoxypyridin-3-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	522.5
389	*N*-(3,5-dichlorobenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	574.4
390	*N*-1-naphthyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	541.6

Compound Number	Name	MS (*m/z*, M + 1)
391	*N*-(4,6-dimethylpyridin-2-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	520.5
392	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pyrimidin-4-ylbenzamide	493.5
393	*N*-(5-methyl-1,3-thiazol-2-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	512.5
394	*N*-(4-methylbenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.6
395	*N*-[3-(1*H*-imidazol-1-yl)propyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	523.6
396	1'-{3-[(4-methylpiperazin-1-yl)carbonyl]benzyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	498.5
397	*N*-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	539.6
398	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-1,3,4-thiadiazol-2-ylbenzamide	499.5
399	*N*-(3,3-dimethylbutyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	499.6
400	*N*-(4-morpholin-4-ylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	576.6
401	*N*-[(1-ethylpyrrolidin-2-yl)methyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	526.6
402	*N*-(2-cyanoethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	

Compound Number	Name	MS (*m/z*, M + 1)
403	*N*-(2,2-diphenylethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	595.4

EXAMPLE 10

Synthesis of 1'-(4-fluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a solution of spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.16 g, 0.57 mmol) in ethylmethylketone (5.00 mL) was added Cs_2CO_3 (0.40 g, 1.20 mmol). The reaction mixture was stirred at ambient temperature for 15 min followed by the addition of 4-fluorobenzyl bromide (0.20 g, 1.0 mmol). The reaction mixture was refluxed for 4 h. After the completion of the reaction, the mixture was filtered and the solvent was removed under reduced pressure. The residue was recrystallized from EtOAc/Hexane to yield the title compound (0.111 g) as a white solid in 50% yield: MS (ES+) *m/z* 390.3 (M + 1).

EXAMPLE 10.1

The compounds listed in the following table were synthesized using the similar procedure as described in EXAMPLE 10. As previously noted, the compound numbers listed below do not correspond to the compound numbers provided in the general Reaction Schemes above.

Compound Number	Name	MS (*m/z*, M + 1)
404	1'-{[1-(2,6-difluorobenzyl)-1*H*-1,2,3-triazol-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	489.3
405	1'-prop-2-yn-1-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	320.3

Compound Number	Name	MS (*m/z*, M + 1)
406	1'-benzylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	372.2
407	1'-{[6-(trifluoromethyl)pyridin-3-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	441.2
408	1'-(3,5-difluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	408.3
409	1'-(3-nitrobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	417.2
410	1'-[(6-chloropyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	407.2
411	1'-(tetrahydro-2*H*-pyran-2-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	380.2
412	1'-(3-fluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	390.3
413	1'-(tetrahydrofuran-2-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	365.2
414	3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzonitrile	397.2
415	1'-(2,3-dihydro-1,4-benzodioxin-2-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	430.3
416	1'-(2-ethoxyethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	354.2
417	1'-[(2E)-pent-2-en-1-yl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	350.2
418	1'-hex-5-en-1-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	364.3
419	1'-(cyclobutylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	350.2
420	1'-pent-2-yn-1-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	348.2
421	1'-(5-chloropentyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	386.2

Compound Number	Name	MS (*m/z*, M + 1)
422	1'-[4-(1*H*-pyrazol-1-yl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	438.3
423	1'-[(7-methoxy-2-oxo-2*H*-1,4-benzoxazin-3-yl)methyl]spiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	471.3
424	1'-(4-fluorobutyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	356.2
425	1'-(5-methylhexyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	380.3
426	1'-[(3*Z*)-4-methylhex-3-en-1-yl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	378.2
427	1'-[(5-methyl-2-phenyl-2*H*-1,2,3-triazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	453.2
428	1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	430.2
429	1'-(3,4-dihydro-2*H*-1,5-benzodioxepin-7-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	444.3
430	1'-(biphenyl-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	448.2
431	4'-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]biphenyl-2-carbonitrile	473.3
432	1'-(2-bromoethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	389.1
433	1'-(1*H*-1,2,3-triazol-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	363.2
434	1'-(biphenyl-2-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	448.3
435	1'-(diphenylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	448.3
436	5-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)pentanenitrile	363.2
437	1'-[2-(2-methoxyethoxy)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	384.2

Compound Number	Name	MS (*m/z*, M + 1)
438	1'-(cyclopropylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	336.2
439	1'-(4,4,4-trifluorobutyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	392.2
440	ethyl 1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-1*H*-1,2,3-triazole-5-carboxylate	449.3
441	ethyl 1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-1*H*-1,2,3-triazole-4-carboxylate	449.3
442	diethyl [2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]phosphonate	446.1
443	1'-(1,3-thiazol-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	379.3
444	1'-[(5-chloro-1-benzothien-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	462.1
445	1'-(pyridin-2-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	373.1
446	1'-(pyridin-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	373.1
447	1'-(pyridin-3-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	373.1
448	1'-[2-(1*H*-pyrrol-1-yl)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	375.2
449	1'-{[4-chloro-2-(trifluoromethyl)quinolin-6-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	525.1
450	1'-(4-fluoro-3-methylbenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	404.3
451	1'-(5-fluoro-2-methylbenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	404.3
452	1'-[(2-methyl-1,3-thiazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	393.2

Compound Number	Name	MS (*m/z*, M + 1)
453	1'-(1,3-benzothiazol-2-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	329.3
454	1'-(2,5-difluorobenzyl)spiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'H)-one	408.2
455	1'-[4-(1*H*-pyrrol-1-yl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	437.3
456	1'-[3-(1*H*-pyrrol-1-yl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	437.3
457	1'-(2,1,3-benzothiadiazol-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	430.3
458	1'-(2,1,3-benzothiadiazol-5-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	430.3
459	1'-[(1-methyl-1*H*-1,2,3-benzotriazol-5-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	427.2
460	1'-[(4-chlorophenoxy)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	422.2
461	1'-[2-fluoro-3-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.1
462	1'-[2-fluoro-6-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.1
463	1'-[3-fluoro-4-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.1
464	1'-[4-fluoro-3-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.1
465	1'-[2-fluoro-5-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.1
466	1'-[4-fluoro-2-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.1
467	1'-[5-fluoro-2-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.1
468	1'-[2-fluoro-4-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.1

Compound Number	Name	MS (*m/z*, M + 1)
469	1'-[3-(1H-pyrrol-1-yl)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	389.2
470	1'-[(2,2,3,3-tetrafluorocyclobutyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	422.1
471	1'-[(3-methyl-5-phenylisoxazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	453.3
472	1'-[2-(diethylamino)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	381.2
473	1'-(2,3-difluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	408.1
474	1'-[(1-bromo-2-naphthyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	501.2
475	1'-[(7-methoxy-2-oxo-2*H*-chromen-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	470.3
476	1'-[(benzyloxy)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	402.3
477	1'-[2-(2,5,5-trimethyl-1,3-dioxan-2-yl)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	438.2
478	1'-[(6-fluoro-4*H*-1,3-benzodioxin-8-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	448.2
479	1'-allylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	322.3
480	1'-(1-naphthylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	423.3
481	1'-[3-fluoro-5-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	458.1
482	1'-(2,4-difluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	408.1
483	1'-(2,6-difluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	408.3

Compound Number	Name	MS (*m/z*, M + 1)
484	1'-[(5-phenyl-1,3-oxazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	439.2
485	1'-(3,5,5-trimethylhexyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	408.3
486	1'-(2-ethylbutyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	366.3
487	1'-(4-methylpentyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	366.3
488	1'-(3-methoxybenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	402.3
489	1'-(3-methylbutyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	352.2
490	1'-(3-methylbut-2-en-1-yl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	350.2
491	1'-pent-4-en-1-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	350.2
492	4-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)butanenitrile	349.1
493	1'-[4-(1*H*-1,2,4-triazol-1-yl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	439.2
494	1'-(1,3-benzodioxol-5-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	416.2
495	1'-[4,4-bis(4-fluorophenyl)butyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	526.2
496	1'-[(2-methylcyclopropyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	350.4
497	1'-(3-cyclopropylpropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	364.3
498	1'-hexylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	366.3
499	1'-[(2-cyclopropyl-6-hydroxypyrimidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	430.3

EXAMPLE 10.2

Synthesis of 1'-(piperidin-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 10, and making non-critical variations using *tert*-butyl 4-(bromomethyl)peridine-1-carboxylate to replace 4-fluorobenzyl bromide, the title compound was obtained in 67% yield as a white solid upon acidification of the intermediate with 33% HBr: MS (ES+) *m/z* 379.3 (M + 1).

EXAMPLE 10.3

Synthesis of 1'-[(1-methylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



To a solution of 1'-(piperidin-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.19 g, 0.50 mmol) in dichloroethane (5.00 mL) was added formaldehyde (0.10 mL, 33% solution, 0.03 g, 1.10 mmol) and sodium triacetoxyborohydride (0.30 g, 1.40 mmol). After stirring at ambient temperature for 20 hours, the reaction mixture was diluted with of dichloromethane (20.0 mL) and washed with water (2 x 20.0 mL). The organic layer was dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography and the product was dissolved in dichloromethane (5.00 mL) and excess of HCl in ether was added. The precipitate was filtered to give the title compound in 20% yield: MS (ES+) *m/z* 393.3 (M + 1).

EXAMPLE 10.4

Synthesis of 1'-[(1-ethylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in EXAMPLE 10.3, and making non-critical variations using acetaldehyde to replace formalin, the title compound was obtained in 20% yield as a white solid: MS (ES+) *m/z* 407.3 (M + 1).

EXAMPLE 10.5

Synthesis of 1'-[(1-cyclohexylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in EXAMPLE 10.3, and making non-critical variations using cyclohexanone to replace formalin, the title compound was obtained 24% yield as a white solid: MS (ES+) *m/z* 461.5 (M + 1).

EXAMPLE 10.6

Synthesis of 1'-{[1-cyclopropylmethyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in EXAMPLE 10.3, and making non-critical variations using cyclopropanecarbaldehyde to replace formalin, the title compound was obtained in 14% yield as a white solid: MS (ES+) *m/z* 433.5 (M + 1).

EXAMPLE 10.7

Synthesis of 1'-[(1-cyclopentylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in EXAMPLE 10.3, and making non-critical variations using cyclopentanone to replace formalin, the title compound was obtained in 37% yield as a white solid: MS (ES+) *m/z* 447.3 (M + 1).

EXAMPLE 10.8

Synthesis of 1'-{[1-(pyridine-3-ylmethyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in EXAMPLE 10.3, and making non-critical variations using nicotinaldehyde to replace formalin, the title compound was obtained in 11% yield as a white solid: MS (ES+) *m/z* 470.4 (M + 1).

EXAMPLE 10.9

Synthesis of 1'-{[1-(3-methylbutyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in EXAMPLE 10.3, and making non-critical variations using 3-methylbutanal to replace formalin, the title compound was obtained in 15% yield as a white solid: MS (ES+) *m/z* 449.5 (M + 1).

EXAMPLE 10.10

Synthesis of 1'-{[1-(1-ethylpropyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in EXAMPLE 10.3, and making non-critical variations using pentan-3-one to replace formalin, the title compound was obtained in 17% yield as a white solid: MS (ES+) *m/z* 449.4 (M + 1).

EXAMPLE 10.11

Synthesis of 1'-[(1-cyclobutylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in EXAMPLE 10.3, and making non-critical variations using cyclobutanone to replace formalin, the title compound was obtained in 31% yield as a white solid: MS (ES+) *m/z* 433.4 (M + 1).

EXAMPLE 10.12

Synthesis of 1'-[(1-isopropylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in Example 10.3, and making non-critical variations using acetone to replace formalin, the title compound was obtained in 31% yield as a white solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.36 (s, 1H), 7.79-6.87 (m, 4H), 6.82-6.48 (m, 1H), 6.38-6.15 (m, 1H), 5.89 (s, 2H), 4.67 (ABq, 2H), 4.12 (s, 1H), 3.79-0.60 (m, 16H); MS (ES+) *m/z* 421.4 (M + 1).

EXAMPLE 10.13

Synthesis of 1'-{[1-(pyridin-2-ylmethyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in EXAMPLE 10.3, and making non-critical variations using picolinaldehyde to replace formalin, the title compound was obtained in 15% yield as a white solid: MS (ES+) *m/z* 470.4 (M + 1).

EXAMPLE 10.14

Synthesis of 1'-{[1-(2-thienylmethyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in EXAMPLE 10.3, and making non-critical variations using thiophene-2-carbaldehyde to replace formalin, the title compound was obtained in 21% yield as a white solid: MS (ES+) *m/z* 475.3 (M + 1).

EXAMPLE 10.15

Synthesis of 1'-({1-[3-(methylthio)propyl]piperidin-4-yl}methyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in EXAMPLE 10.3, and making non-critical variations using 3-(methylthio)propanal to replace formalin, the title compound was obtained in 7% yield as a white solid: MS (ES+) *m/z* 467.5 (M + 1).

EXAMPLE 10.16

Synthesis of 1'-{[1-(tetrahydro-2*H*-pyran-4-yl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in EXAMPLE 10.3, and making non-critical variations using tetrahydro-4*H*-pyran-4-one to replace formalin, the title compound was obtained in 33% yield as a white solid: MS (ES+) *m/z* 463.4 (M + 1).

EXAMPLE 10.17

Synthesis of 1'-{[1-(3,3-dimethylbutyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



· HCl

Following the procedure as described in EXAMPLE 10.3, and making non-critical variations using 3,3-dimethylbutanal to replace formalin, the title compound was obtained in 19% yield as a white solid: MS (ES+) *m/z* 463.5 (M + 1).

EXAMPLE 10.18

Synthesis of *tert*-butyl 4-[(5,5-dimethyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)methyl]piperidine-1-carboxylate



Following the procedure as described in EXAMPLE 10, and making non-critical variations using 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one to replace spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and *tert*-butyl 4-(tosyloxymethyl)piperidine-1-carboxylate to replace 4-fluorobenzyl bromide, the title compound was obtained in 70% yield: mp 65-75 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.30 (td, 1H), 7.18 (d, 1H), 7.05 (t, 1H), 6.89 (d, 1H), 6.38 (s, 1H), 6.28 (s, 1H), 4.88 (d, 1H), 4.64 (d, 1H), 4.18 (s, 2H), 4.17-4.01 (br, 2H), 3.74-3.53 (m, 2H), 2.74-2.59 (m, 2H), 2.11-1.92 (m, 1H), 1.70-1.59 (m, 2H), 1.43 (s, 9H), 1.37-1.19 (m, 2H), 1.17 (s, 3H), 1.14 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.0, 161.3, 161.0, 154.7, 142.8, 132.6, 129.9, 128.7, 124.2, 123.3, 120.3, 116.2, 108.5, 93.4, 85.4, 80.6, 79.5, 57.7, 45.7, 41.3, 35.0, 30.0, 28.4, 27.8, 27.5; MS (ES+) *m/z* 527.5 (M + 23).

EXAMPLE 10.19

Synthesis of 5,5-dimethyl-1'-(piperidin-4-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one hydrochloride



To a stirred solution of *tert*-butyl 4-[(5,5-dimethyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)methyl]piperidine-1-carboxylate (80 mg, 0.16 m mol) in 10.0 mL dichloromethane was added hydrobromic

acid (0.50 mL of hydrobromic acid ≥33% in glacial acetic acid, 1.60 mmol) slowly at 0 °C. The mixture was stirred at ambient temperature for one hour and concentrated *in vacuo* to dryness. The residue was treated with 10.0 mL of 2 N sodium hydroxide solution and extracted with dichloromethane (3 x 30.0 mL). The combined organic layers was washed with brine (100 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography (2% methanol in ethyl acetate) to give 5,5-dimethyl-1'-(piperidin-4-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one (0.03 g, 46%), which was treated with 2.0 M HCl in diethyl ether to give the title compound: ^{1}H NMR (300 MHz, CD_3OD) δ 7.41 (td, 1H), 7.25-7.12 (m, 3H), 6.43 (s, 1H), 6.37 (s, 1H), 4.87 (d, 1H), 4.72 (d, 1H), 4.23 (s, 2H), 3.80 (d, 2H), 3.51-3.38 (m, 2H), 3.11-2.94 (m, 2H), 2.39-2.19 (m, 1H), 2.06-1.94 (m, 2H), 1.67-1.49 (m, 2H), 1.22 (s, 3H), 1.19 (s, 3H); ^{13}C NMR (75 MHz, CD_3OD) δ 181.4, 163.9, 163.4, 144.7, 134.7, 132.3, 131.0, 125.9, 125.8, 122.6, 118.4, 111.3, 94.9, 87.4, 82.6, 60.2, 46.7, 45.7, 45.6, 43.3, 34.6, 28.8, 28.7, 28.6; MS (ES+) *m/z* 405.4 (M + 1).

EXAMPLE 10.20

Synthesis of 7'-fluoro-1'-[(1-isopropylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



HCl

A. Following the procedure as described in EXAMPLES 10.18 and 10.19, and making non-critical variations using 7'-fluorospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one, 7'-fluoro-1'-(piperidin-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one was obtained and used in the next step.

B. To a stirred solution of 7'-fluoro-1'-(piperidin-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (120 mg, 0.28 mmol) and triethylamine (3.9 µL, 0.028 mmol) in 5.00 mL dichloromethane was added acetone (4.1 µL, 0.56 mmol) followed with sodium triacetoxyborohydride (124 mg, 0.56 mmol) at ambient

temperature. The mixture was stirred at ambient temperature overnight, quenched with water (10.0 mL). The mixture was extracted with dichloromethane (3 x 30.0 mL). The combined organic layers was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo*. The residue was subjected to column chromatography (2% methanol in ethyl acetate/hexane) to give 7'-fluoro-1'-[(1-isopropylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (85 mg, 69%) as a white solid, which was treated with 2.0 M HCl in diethyl ether to give the title compound: mp 157-160 °C; ^{1}H NMR (300 MHz, CD_3OD) δ 7.22-7.05 (m, 2H), 7.05-6.98 (m, 1H), 6.52 (s, 1H), 6.21 (s, 1H), 5.86 (s, 2H), 4.83 (d, 1H), 4.68 (d, 1H), 3.95-3.77 (m, 2H), 3.55-3.42 (m, 3H), 3.12-2.96 (m, 2H), 2.30-2.10 (m, 1H), 2.10-1.97 (m, 2H), 1.76-1.52 (m, 2H), 1.34 (d, 6H); ^{13}C NMR (75 MHz, CD_3OD) δ 179.7, 157.6, 150.6, 150.4, 147.1, 143.9, 136.6, 125.8, 121.2, 120.3, 118.0, 103.9, 103.0, 94.3, 81.8, 60.1, 59.7, 47.6, 35.33, 35.30, 28.6, 28.5, 16.98, 16.96; MS (ES+) *m/z* 439.27 (M + 1).

EXAMPLE 10.21

Synthesis of 5,5-dimethyl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one



To a solution of 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one (0.09 g, 0.29 mmol) in 2-butanone (10.0 mL) was added 2-bromomethyl-5-(triflоromethyl)furan (0.08 g, 0.35 mmol) followed by cesium carbonate (0.19 g, 0.58 mmol) at 0 °C. The mixture was stirred at ambient temperature overnight. The reaction mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/5) to give the title compound (0.06 g, 45%): mp 155-160 °C; ^{1}H NMR (300 MHz, $CDCl_3$,) δ 7.29 (t, 1H), 7.19 (d, 1H), 7.07 (t, 1H), 6.97 (d, 1H), 6.73 (t, 1H), 6.42-6.37 (m, 2H), 6.30 (s, 1H), 5.08 (d, 1H), 4.94-4.84 (m, 2H), 4.65 (d, 1H), 4.18 (s, 2H), 1.19 (s, 3H), 1.14 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.5, 161.2, 161.0, 152.0, 141.4, 132.5, 130.1, 128.8, 124.2, 123.8, 120.1, 116.4, 112.6, 109.3, 108.7, 93.4, 85.5, 80.6, 57.7, 41.4,

36.9, 27.6, 27.5; MS (ES+) *m/z* 456.5 (M + 1).

EXAMPLE 10.22

Synthesis of 5,5-dimethyl-1'-(pyridin-3-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one hydrochloride



To a solution of 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one (0.08 g, 0.26 mmol) in DMF (10 mL) was added sodium hydride (0.03 g, 0.78 mmol) slowly at 0 °C. After 30 min, 3-(bromomethyl)-pyridine hydrobromide (0.10 g, 0.39 mmol) was added. The mixture was stirred at ambient temperature overnight, quenched with saturated ammonium chloride (10.0 mL). The mixture was extracted with ethyl acetate (3 x 20.0 mL). The combined organic layers was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo*. The residue was subjected to column chromatography (ethyl acetate/hexane, 2/1) to give 5,5-dimethyl-1'-(pyridin-3-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one as a white solid (0.05 g, 48%), which was treated with 2.0 M HCl in diethyl ether to give the title compound: mp 124-126 °C; ^{1}H NMR (300 MHz, CD_3OD) δ 8.96 (br, 1H), 8.82 (br, 1H), 8.62 (d, 1H), 7.3 (t, 1H), 7.32 (td, 1H), 7.23-7.17 (m, 1H), 7.16-7.08 (m, 2H), 6.42 (s, 1H), 6.32 (s, 1H), 5.35-5.14 (m, 2H), 4.93-4.84 (m, 1H), 4.74 (d, 1H), 4.18 (s, 2H), 1.18 (s, 3H), 1.14 (s, 3H); ^{13}C NMR (75 MHz, CD_3OD) δ 178.9, 161.6, 161.1, 145.4, 141.3, 141.1, 132.5, 130.1, 128.8, 124.0, 123.9, 120.1, 116.2, 108.8, 92.6, 85.1, 80.2, 57.8, 41.0, 40.5, 26.5, 26.4; MS (ES+) *m/z* 399.5 (M + 1).

EXAMPLE 10.23

Synthesis of 5,5-dimethyl-1'-(pyridin-2-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one hydrochloride



Following the procedure as described in EXAMPLE 10.22, and making non-critical variations using 2-(bromomethyl)-pyridine hydrobromide to replace 3-(bromomethyl)-pyridine hydrobromide, the title compound was obtained (45%): mp 145-147 °C; ^{1}H NMR (300 MHz, CD_3OD) δ 8.86 (dd, 1H), 8.56 (td, 1H), 8.05-7.97 (m, 2H), 7.37 (td, 1H), 7.31-7.25 (m, 1H), 7.24-7.16 (m, 1H), 7.11 (d, 1H), 6.55 (s, 1H), 6.37 (s, 1H), 5.52 (d, 1H), 5.38 (d, 1H), 4.97 (d, 1H), 4.79 (d, 1H), 4.23 (s, 2H), 1.24 (s, 3H), 1.20 (s, 3H); ^{13}C NMR (75 MHz, CD_3OD) δ 177.4, 160.0, 159.6, 150.3, 144.7, 141.0, 139.7, 131.0, 128.6, 127.2, 124.3, 123.7, 122.6, 122.4, 118.5, 114.8, 107.2, 91.0, 83.6, 78.7, 56.3, 40.1, 39.4, 24.9, 24.8; MS (ES+) *m/z* 399.5 (M + 1).

EXAMPLE 10.24

Synthesis of 1'-[(6-methylpyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



Following the procedure as described in EXAMPLE 10.21, and making non-critical variations using (6-methylpyridin-3-yl)methyl 4-methylbenzenesulfonate to replace 2-bromomethyl-5-(trifloromethyl)furan, and spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one, 1'-[(6-methylpyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one was obtained (56%), which was treated with

2.0 M HCl in diethyl ether to give the title compound: ^{1}H NMR (300 MHz, CD_3OD) δ 8.77 (s, 1H), 8.46 (dd, 1H), 7.91 (d, 1H), 7.33 (t, 1H), 7.24-7.09 (m, 3H), 6.52 (s, 1H), 6.14 (s, 1H), 5.86 (s, 2H), 5.18 (s, 2H), 4.93-4.85 (m, 1H), 4.71 (d, 1H), 2.77 (s, 3H); ^{13}C NMR (75 MHz, CD_3OD) δ 180.0, 157.6, 155.1, 150.6, 146.6, 143.8, 142.7, 141.2, 135.7, 133.5, 130.3, 129.5, 125.3, 125.2, 120.5, 110.3, 103.8, 103.0, 94.3, 81.5, 59.7, 41.5, 19.5; MS (ES+) *m/z* 387.4 (M + 1).

EXAMPLE 10.25

Synthesis of 1'-[(6-methoxypyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 10.21, and making non-critical variations using (6-methoxypyridin-3-yl)methyl 4-methylbenzenesulfonate to replace 2-bromomethyl-5-(trifloromethyl)furan, and spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, the title compound was obtained (45%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.19 (d, 1H), 7.59 (dd, 1H), 7.26-7.12 (m, 2H), 7.02 (t, 1H), 6.83 (d, 1H), 6.74 (d, 1H), 6.50 (s, 1H), 6.07 (s, 1H), 5.89-5.82 (m, 2H), 5.00-4.90 (m, 2H), 4.76 (d, 1H), 4.65 (d, 1H), 3.93 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.6, 163.7, 155.9, 149.0, 145.3, 142.4, 141.6, 138.9, 132.2, 129.0, 124.4, 124.1, 123.7, 119.2, 111.7, 109.0, 102.9, 101.5, 93.7, 80.4, 58.2, 53.9, 41.1; MS (ES+) *m/z* 403.2 (M + 1).

EXAMPLE 10.26

Synthesis of 1'-[(6-chloropyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using 2-chloro-5-(chloromethyl)pyridine to replace 2-bromomethyl-5-(trifloromethyl)furan, and spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, the title compound was obtained (69%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.42 (d, 1H), 7.63 (dd, 1H), 7.34-7.14 (m, 3H), 7.05 (t, 1H), 6.77 (d, 1H), 6.51 (s, 1H), 6.06 (s, 1H), 5.89-5.84 (m, 2H), 5.07-4.78 (m, 3H), 4.66 (d, 1H); MS (ES+) *m/z* 407.3 (M + 1).

EXAMPLE 10.27

Synthesis of 1'-{[6-(dimethylamino)pyridin-3-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



To a seal tube was added 1'-((6-chloropyridin-3-yl)methyl)-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one (0.10 g, 0.25 mmol) and dimethylamine (2.00 mL of 2 M THF solution, 4.00 mmol). The mixture was stirred at 130 °C overnight. After cooling down to ambient temperature, the mixture was concentrated *in vacuo*. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/2) to give 1'-{[6-(dimethylamino)pyridin-3-yl]methyl}spiro[furo[2,3-

f][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one as a white solid (50 mg, 48%), which was treated with 2.0 M HCl in diethyl ether to give the title compound: mp 146-150 °C; ^{1}H NMR (300 MHz, CD_3OD) δ 8.03 (d, 1H), 7.95 (dd, 1H), 7.36 (td, 1H), 7.25-7.12 (m, 4H), 6.57 (s, 1H), 6.09 (s, 1H), 5.90 (s, 2H), 5.07-4.87 (m, 3H), 4.72 (d, 1H), 3.27 (s, 6H); ^{13}C NMR (75 MHz, CD_3OD) δ 179.9, 157.6, 154.6, 150.5, 143.8, 143.4, 142.9, 137.7, 133.5, 130.2, 125.1, 121.8, 120.6, 113.2, 110.4, 103.7, 103.0, 94.3, 81.4, 59.7, 41.1, 39.5; MS (ES+) *m/z* 416.5 (M + 1).

EXAMPLE 10.28

Synthesis of 1'-[(6-morpholin-4-ylpyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



HCl

Following the procedure as described in EXAMPLE 10.27, and making non-critical variations using morpholine to replace dimethylamine solution, the title compound was obtained (52%): mp 185-200 °C; ^{1}H NMR (300 MHz, CD_3OD) δ 8.13-8.04 (m, 2H), 7.45 (d, 1H), 7.37 (t, 1H), 7.26-7.14 (m, 3H), 6.56 (s, 1H), 6.10 (s, 1H), 5.89 (s, 2H), 5.10-4.87 (m, 3H), 4.72 (d, 1H), 3.91-3.84 (m, 4H), 3.73-3.67 (m, 4H); ^{13}C NMR (75 MHz, CD_3OD) δ 180.0, 157.6, 153.7, 150.5, 145.1, 143.8, 142.8, 136.3, 133.5, 130.3, 125.2, 125.1, 123.6, 120.6, 114.6, 110.4, 103.7, 103.0, 94.3, 81.5, 66.7, 59.7, 47.2, 40.9; MS (ES+) *m/z* 458.5 (M + 1).

EXAMPLE 10.29

Synthesis of 1'-[(6-pyrrolidin-1-ylpyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



HCl

Following the procedure as described in EXAMPLE 10.27, and making non-critical variations using pyrrolidine to replace dimethylamine solution, the title compound was obtained (45%): mp 160-165 °C; ^{1}H NMR (300 MHz, CD_3OD) δ 7.99-7.91 (m, 2H), 7.33 (td, 1H), 7.22-7.06 (m, 4H), 6.53 (s, 1H), 6.05 (s, 1H), 5.86 (s, 2H), 5.04-4.82 (m, 3H), 4.68 (d, 1H), 3.57 (t, 4H), 2.13 (t, 4H); ^{13}C NMR (75 MHz, CD_3OD) δ 177.0, 154.6, 148.0, 147.6, 141.1, 140.8, 139.9, 132.5, 130.5, 127.3, 122.2, 118.8, 117.6, 112.0, 107.4, 100.7, 100.0, 91.3, 78.5, 56.7, 38.0, 23.2; MS (ES+) *m/z* 442.2 (M + 1).

EXAMPLE 10.30

Synthesis of 1'-(2-chloro-4-fluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and 1-(bromomethyl)-2-chloro-4-fluorobenzene to replace 2-(bromomethyl)-5-

(trifluoromethyl)furan, the title compound was obtained (54%) as a white solid: mp 174-175 °C; MS (ES+) *m/z* 424.2 (M + 1).

EXAMPLE 10.31

Synthesis of 1'-[(2-methylcyclopropyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and 1-(bromomethyl)-2-methylcyclopropane to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (37%) as a white solid: MS (ES+) *m/z* 350.3 (M + 1).

EXAMPLE 10.32

Synthesis of 1'-(3-cyclopropylpropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and (3-bromopropyl)cyclopropane to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (51%) as a white solid: mp 111-113 °C; MS (ES+) *m/z* 364.3 (M + 1).

EXAMPLE 10.33

Synthesis of 1'-butylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one, and 1-bromobutane to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (62%) as a white solid: mp 119-120 °C; MS (ES+) *m/z* 338.3 (M + 1).

EXAMPLE 10.34

Synthesis of 1'-[(5-methylisoxazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one, and 4-(bromomethyl)-5-methylisoxazole to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (25%) as a white solid: mp 159-161 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.39-6.91 (m, 4H), 6.50 (s, 1H), 6.11 (s, 1H), 5.94 (d, 1H), 5.85 (ABq, 2H), 4.95 (ABq, 2H), 4.78 (ABq, 2H), 2.37 (s, 3H); MS (ES+) *m/z* 377.3 (M + 1).

EXAMPLE 10.35

Synthesis of 1'-(tetrahydro-2*H*-pyran-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and 4-(bromomethyl)tetrahydro-2*H*-pyran to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (25%) as a white solid: mp 142-144 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.34-6.85 (m, 4H), 6.50 (s, 1H), 6.08 (s, 1H), 5.85 (ABq, 2H), 4.76 (ABq, 2H), 4.18-3.86 (m, 2H), 3.63 (ddd, 2H), 3.34 (t, 2H), 2.38-1.92 (m, 1H), 1.70-1.36 (m, 4H); MS (ES+) *m/z* 380.3 (M + 1).

EXAMPLE 10.36

Synthesis of 1'[2-(trifluoromethoxy)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and 1-(bromomethyl)-2-(trifluoromethoxy)benzene to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (77%) as a white solid: mp 130-135 °C; MS (ES+) *m/z* 456.3 (M + 1).

EXAMPLE 10.37

Synthesis of 1'[3-(trifluoromethoxy)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and 1-(bromomethyl)-3-(trifluoromethoxy)benzene to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (65%) as a white solid: mp 88-91 °C; MS (ES+) *m/z* 456.3 (M + 1).

EXAMPLE 10.38

Synthesis of 1'[4-(trifluoromethoxy)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and 1-(bromomethyl)-4-(trifluoromethoxy)benzene to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (50%) as a white solid: mp 99-101 °C; MS (ES+) *m/z* 456.3 (M + 1).

EXAMPLE 10.39

Synthesis of 1'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and iodomethane to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (72%) as a white solid: mp 142-144 °C; MS (ES+) *m/z* 296.2 (M + 1).

EXAMPLE 10.40

Synthesis of 1'-Propylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and 1-bromopropane to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (64%) as a white solid: mp 158-160 °C; MS (ES+) *m/z* 324.4 (M + 1).

EXAMPLE 10.41

Synthesis of 1'-(2,1,3-benzoxadiazol-5-ylmethyl)spiro[furo2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and 5-(bromomethyl)benzo[c][1,2,5]oxadiazole to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (17%) as a white solid: mp 163-165 °C; MS (ES+) *m/z* 414.4 (M + 1).

EXAMPLE 10.42

Synthesis of 1'[(1-methyl-1*H*-benzotriazol-6-yl)methyl]spiro[furo2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and 6-(bromomethyl)-1-methyl-1*H*-benzotriazole to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (17%) as a white solid: mp 230-235 °C; MS (ES+) *m/z* 427.3 (M + 1).

EXAMPLE 10.43

Synthesis of *tert*-butyl 4-[(2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)methyl]piperidine 1-carboxylate



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using 5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and *tert*-butyl 4-(bromomethyl)peridine-1-carboxylate to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (58%) as a white solid: mp 96-98 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.84-6.85 (m, 4H), 6.40 (s, 1H), 6.37 (s, 1H), 4.68 (ABq, 2H), 4.46 (t, 2H), 4.06-3.73 (m, 2H), 3.68-3.45 (m, 2H), 2.92 (t, 2H), 2.63 (s, 2H), 2.04-1.82 (m, 1H), 1.76-0.66 (m, 13H); MS (ES+) *m/z* 477.4 (M + 1).

EXAMPLE 10.44

Synthesis of 1'-(2,3-difluorobenzyl)-5,6-dihydrospyro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using 5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and 1-(bromomethyl)-2,3-difluorobenzene to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (67%) as a white solid: mp

156-158 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.50-6.85 (m, 7H), 6.43 (s, 1H), 6.39 (s, 1H), 5.01 (q, 2H), 4.75 (dd, 2H), 4.46 (t, 2H), 2.92 (t, 2H); MS (ES+) *m/z* 406.2 (M + 1).

EXAMPLE 10.45

Synthesis of 1'-(pyridin-2-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one hydrochloride



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using 5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and 2-(bromomethyl)pyridine hydrobromide to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, 1'-(pyridin-2-ylmethyl)-5,6-dihydrospyro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]2'(1'*H*)-one was obtained (27%) as a white solid, which was treated in CH_2Cl_2 with excess of HCl in ether to give the tilte compound: mp 208-210 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.78-8.53 (m, 1H), 8.05 (t, 1H), 7.64-7.47 (m, 2H), 7.30-6.92 (m, 4H), 6.59 (s, 1H), 6.38 (s, 1H), 5.24-5.06 (m, 2H), 4.78 (ABq, 2H), 4.46 (t, 2H), 2.94 (t, 2H); MS (ES+) *m/z* 371.4 (M + 1).

EXAMPLE 10.46

Synthesis of 1'-(4-methoxybenzyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical

variations using 5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, and 1-(chloromethyl)-4-methoxybenzene to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (56%) as a white solid: mp 120-121 °C; MS (ES+) *m/z* 400.2 (M + 1).

EXAMPLE 10.47

Synthesis of 4'-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a solution of 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.48 g, 1.33 mmol) in *N*,*N*-dimethylformamide (5.00 mL) was added sodium hydride (0.08 g, 1.98 mmol, 60% dispersion in mineral oil) in one portion at 0 °C. The reaction mixture was stirred for 0.5 h followed by the addition of a solution of 2-(bromomethyl)-5-trifluoromethyl)furan in *N*,*N*-dimethylformamide (1.00 mL). The reaction mixture was stirred at ambient temperature for 16 h and quenched by slow addition of water (5.00 mL). The reaction mixture was extracted with ethyl acetate (3 x 20.0 mL), washed with brine, dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate:hexane (35%) to afford the title compound (0.46 g, 69%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.22-7.14 (m, 2H), 6.94 (dd, 1H), 6.73 (d, 1H), 6.46 (s, 1H), 6.39 (d, 1H), 6.04 (s, 1H) 5.86 (dd, 2H), 4.94 (ABq, 2H), 4.92 (ABq, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.0, 157.2, 151.6, 151.5, 149.3, 143.4, 142.2, 130.5, 127.8, 129.6, 120.1, 116.0, 112.7, 109.5, 107.9, 102.5, 101.6, 93.3, 77.1, 59.6, 37.1; MS (ES+) *m/z* 508.2 (M + 2).

EXAMPLE 10.48

Synthesis of 4'-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using 4'-bromo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, the title compound was obtained (76%) as a colorless solid: mp 182-184 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.21-7.11 (m, 2H), 6.92 (dd, 1H), 6.74 (d1H), 6.41 (d, 1H), 6.38 (s, 1H), 6.37 (s, 1H), 5.10 (d, 1H), 5.02 (d, 1H), 4.87 (d, 1H), 4.81 (d, 1H), 4.53 (t, 2H), 2.98 (t, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.3, 162.4, 162.2, 151.5, 143.4, 130.2, 127.7, 120.5, 120.0, 119.7, 118.4, 117.0, 112.7, 112.6, 109.5, 107.8, 92.9, 77.1, 72.4, 59.1, 37.0, 28.9; MS (ES+) *m/z* 506.3 (M + 1).

EXAMPLE 10.49

Synthesis of 4'-bromo-1'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using iodomethane to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (79%) as a colorless solid: mp 155-157 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.19 (d, 1H), 7.17 (s, 1H), 6.84 (dd, 1H), 6.46 (s, 1H), 6.08 (s, 1H), 5.86 (dd, 2H), 4.90 (ABq, 2H), 3.25 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.3, 157.2, 149.2, 145.1, 142.0, 130.4, 129.9, 127.3, 119.9, 116.3, 107.3, 102.7, 101.5, 93.3, 77.3, 59.7, 26.9; MS (ES+) *m/z* 376.4 (M + 2).

EXAMPLE 10.50

Synthesis of *tert*-butyl 4-[(4'-bromo-2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]piperidine-1-carboxylate



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using *tert*-butyl 4-({[(4-methylphenyl)sulfonyl]oxy}methyl)piperidine-1-carboxylate to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (43%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.17 (d, 2H), 6.83 (t, 1H), 6.46 (s, 1H), 6.04 (s, 1H), 5.87 (d, 2H), 4.89 (ABq, 2H), 4.11 (d, 2H), 3.73-3.42 (m, 3H), 2.66 (t, 2H), 2.03-1.90 (m, 1H), 1.43 (s, 9H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.6, 157.3, 154.7, 149.2, 144.8, 142.1, 130.3, 129.8, 127.3, 120.2, 116.3, 107.6, 102.4, 101.6, 93.4, 79.6, 77.2, 59.6, 46.1, 43.4, 34.9, 28.4; MS (ES+) *m/z* 581.4 (M + 23), 579.4 (M + 23), 503.3 (M - 57), 501.3 (M - 57).

EXAMPLE 10.51

Synthesis of 1'-[(3,5-dimethylisoxazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 4-(chloromethyl)-3,5-dimethylisoxazole to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (35%) as a colorless solid: mp 165-167 °C; ^{1}H NMR (300

MHz, $CDCl_3$) δ 7.24 (t, 1H), 7.16 (d, 1H), 7.05 (t, 1H), 6.72 (d, 1H), 6.50 (s, 1H), 6.05 (s, 1H), 5.85 (d, 2H), 4.75 (ABq, 2H), 4.67 (ABq, 2H), 2.46 (s, 3H), 2.22 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.5, 167.1, 159.1, 156.1, 149.1, 142.4, 141.7, 131.8, 129.0, 124.3, 123.8, 118.9, 108.8, 108.6, 102.9, 101.6, 93.8, 80.6, 58.2, 33.3, 11.5, 10.7; MS (ES+) *m/z* 391.3 (M + 1).

EXAMPLE 10.52

Synthesis of 1'-(2-furylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 2-chloromethylfuran to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound (40%) as a colorless solid: mp 110-112 °C; ^1H NMR (300 MHz, $CDCl_3$) δ 7.35-7.33 (m, 1H), 7.29-7.23 (m, 1H), 7.15 (d, 1H), 7.06-7.00 (m, 2H), 6.50 (s, 1H), 6.34-6.31 (m, 2H), 6.10 (s, 1H), 5.85 (dd, 2H), 4.92 (ABq, 2H), 4.79 (ABq, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.1, 155.9, 149.0, 148.9, 142.6, 142.3, 141.8, 132.2, 128.9, 123.8, 123.5, 119.5, 110.6, 109.3, 108.7, 103.1, 101.5, 93.6, 80.4, 58.2, 37.1; MS (ES+) *m/z* 362.5 (M + 1).

EXAMPLE 10.53

Synthesis of ethyl 5-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)pentanoate



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and ethyl 5-bromovalerate to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (62%) as a gummy material: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.28 (t, 1H), 7.14 (d, 1H), 7.03 (t, 1H), 6.88 (d, 1H), 6.48 (s, 1H), 6.13 (s, 1H), 5.84 (d, 2H), 4.76 (ABq, 2H), 4.07 (q, 2H), 3.87-3.65 (m, 2H), 2.35 (t, 2H), 1.80-1.64 (m, 4H), 1.20 (t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.4, 173.1, 155.9, 148.8, 142.3, 142.2, 132.5, 128.9, 124.04, 123.3, 119.5, 108.6, 103.0, 101.5, 93.6, 80.5, 60.4, 58.2, 39.9, 33.7, 26.8, 22.2, 14.2; MS (ES+) *m/z* 432.09 (M + 23).

EXAMPLE 10.54

Synthesis of ethyl 4-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)butanoate



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-

bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and ethyl 4-bromobutyrate to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (80%) as a gummy material: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.29 (t, 1H), 7.14 (d, 1H), 7.04 (d, 1H), 6.99 (d, 1H), 6.48 (s, 1H), 6.18 (s, 1H), 5.84 (d, 2H), 4.76 (ABq, 2H), 4.11 (q, 2H), 3.88-3.71 (m, 2H), 2.40 (t, 2H), 2.03 (t, 2H), 1.21(t, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.4, 172.8, 156.0, 148.9, 142.4, 142.2, 132.4, 129.0, 124.0, 123.3, 119.4, 108.7, 103.0, 101.5, 93.6, 80.5, 60.7, 58.2, 39.6, 31.2, 22.6, 14.3; MS (ES+) *m/z* 418.08 (M + 23), 396.1 (M + 1).

EXAMPLE 10.55

Synthesis of 1'-(1,2,4-oxadiazol-3-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 3-(chloromethyl)-1,2,4-oxadiazole to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (36%)) as a colorless solid: mp 160-162 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.39 (dt, 1H), 7.20-7.13 (m, 3H), 7.05 (d, 1H), 6.50 (s, 1H), 6.12 (s, 1H), 5.86 (dd, 2H), 4.78 (ABq, 2H), 4.68 (s, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.7, 156.0, 149.3, 142.5, 139.6, 131.6, 129.4, 124.8, 124.5, 118.4, 113.6, 108.7, 103.0, 101.7, 93.7, 80.3, 58.2, 29.7, 28.0; MS (ES+) *m/z* 365.2 (M + 1).

EXAMPLE 10.56

Synthesis of 1'-{[5-(3-chlorophenyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 2-(chloromethyl)-5-[3-chlorophenyl]furan to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (22%) as a colorless solid: mp 205-207 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.55 (t, 1H), 7.46 (dt, 1H), 7.28 (d, 2H), 7.21-7.14 (m, 2H), 7.09-7.04 (m, 2H), 6.59 (d, 1H), 6.50 (s, 1H), 6.40 (d, 1H), 6.10 (s, 1H), 5.84 (dd, 2H), 4.98 (ABq, 2H), 4.80 (ABq, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.2, 155.9, 152.5, 149.2, 148.9, 142.4, 141.7, 134.8, 132.1, 132.0, 130.1, 128.9, 127.5, 124.0, 123.7, 123.6, 121.7, 119.4, 110.8, 109.2, 106.9, 103.0, 101.5, 93.6, 80.4, 58.2, 37.3; MS (ES+) *m/z* 472.2 (M + 1).

EXAMPLE 10.57

Synthesis of 1'-(3-chloropropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 1-bromo-3-chloropropane to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (22%) as a colorless solid: mp 144-146 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.31 (dt, 1H), 7.14-7.12 (m, 2H), 7.01(t, 1H), 6.65 (s, 1H), 6.23 (s, 1H), 5.89 (s, 2H),

4.68 (ABq, 2H), 3.85-3.79 (m, 2H), 3.67 (t, 2H), 2.06 (t, 1H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 177.2, 155.8, 148.7, 142.9, 142.2, 132.4, 129.3, 124.1, 123.3, 120.3, 109.3, 103.6, 101.7, 80.3, 57.8, 43.4, 30.6; MS(ES+) *m/z* 358.2 (M + 1).

EXAMPLE 10.58

Synthesis of 1'-[(2-isopropyl-1,3-oxazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 4-chloromethyl-2-isopropyloxazole to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (22%) as a colorless solid: mp 118-120 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.97 (s, 1H), 7.24 (t, 1H), 7.13 (d, 1H), 7.05 (d, 1H), 6.98 (t, 1H), 6.65 (s, 1H), 6.26 (s, 1H), 5.88 (d, 2H), 4.85 (d, 1H), 4.77 (d, 1H), 4.71-4.66 (m, 2H), 3.04-2.95 (m, 1H), 1.18 (dd, 6H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 176.9, 169.1, 155.6, 148.7, 142.4, 142.2, 136.8, 135.2, 132.4, 129.2, 123.9, 123.4, 120.5, 109.9, 103.6, 101.9, 93.7, 79.9, 57.9, 36.2, 28.1, 20.7, 20.6; MS (ES+) *m/z* 405.2 (M + 1).

EXAMPLE 10.59

Synthesis of 1'-[(1-methyl-1*H*-benzimidazol-2-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-

bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 2-(bromomethyl)-1-methyl-1*H*-benzimidozole to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (22%) as a colorless solid: mp >250 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.71 (d, 1H), 7.65 (d, 1H), 7.54 (t, 1H), 7.39 (t, 1H), 7.30-7.24 (m, 1H), 7.16 (d, 1H), 7.05 (d, 2H), 6.63 (s, 1H), 6.49 (s, 1H), 6.44 (d, 1H), 6.11 (s, 1H), 5.83 (d, 2H), 4.99 (ABq, 2H), 4.80 (ABq, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.2, 155.9, 150.6, 149.6, 148.9, 142.3, 141.7, 132.1, 131.8, 129.8, 128.9, 127.9, 126.8, 126.7, 126.6, 123.9, 123.6, 119.4, 111.0, 110.9, 110.6, 109.3, 103.1, 101.5, 93.6, 80.4, 58.2, 37.3; MS (ES+) *m/z* 506.3 (M + 1).

EXAMPLE 10.60

Synthesis of 1'-[(2-oxo-1,3-benzothiazol-3(2*H*)-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3''-indol]-2''(1''*H*)-one to replace 4''-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3''-indol]-2''(1''*H*)-one, and 3-(bromomethyl)-benzo[*d*]thiazol-2(3*H*)-one to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (31%) as a colorless solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.67 (d, 1H), 7.41 (d, 1H), 7.35-7.29 (m, 3H), 7.23-7.14 (m, 2H), 7.05 (t, 1H), 6.68 (s, 1H), 5.94-5.85 (m, 5H), 4.69 (td, 1H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 178.1, 170.6, 155.9, 148.9, 142.2, 141.0, 136.1, 131.8, 129.6, 127.3, 124.5, 124.4, 124.3, 123.7, 121.5, 119.8, 112.1, 110.0, 103.4, 101.9, 93.8, 80.3, 58.3, 47.9; MS (ES+) *m/z* 467.2 (M + 23).

EXAMPLE 10.61

Synthesis of 1'-[(5-chloro-2-thienyl)methyl]-5'-fluorospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using 5'-fluorospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 5-chloro-2-(chloromethyl)thiophene to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (76%) as a colorless solid: mp 142-144 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.22-7.17 (m, 1H), 7.14-7.13 (m, 1H), 7.12-7.10 (m, 2H), 6.96 (d, 1H), 6.68 (s, 1H), 6.13 (s, 1H), 5.91 (d, 2H), 5.02 (ABq, 2H), 4.73 (ABq, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 176.9, 161.0, 157.8, 156.0, 149.0, 142.2, 138.3, 138.2 (d, $^4J_{CF}$ = 7.0 Hz) 133.6 (d, $^3J_{CF}$ = 33 Hz), 128.3, 127.8, 127.1, 119.5, 115.6 (d, $^1J_{CF}$ = 93 Hz), 112.5 (d, $^1J_{CF}$ = 100 Hz), 110.8 (d, $^3J_{CF}$ = 32 Hz), 103.2, 102.0, 93.9, 79.8, 58.2 (d, $^4J_{CF}$ = 7.0 Hz), 39.0; MS (ES+) *m/z* 430.1 (M + 1).

EXAMPLE 10.62

Synthesis of 1'-[(5-chloro-2-furyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To an ice-cooled solution of (5-chloro-2-furyl)methanol (2.03 g, 15.3 mmol) in anhydrous dichloromethane (50.0 mL) was added triethylamine (4.64 g, 45.9 mmol) followed by thionyl chloride (3.64 g, 30.6 mmol). The reaction mixture was stirred at 0 °C for 30 min and quenched with saturated ammonium chloride (25.0 mL). After the aqueous layer was separated, the organic layer was washed with 10% aqueous HCl

(20.0 mL), brine (20 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to drynessto give 5-chloro-2-chloromethylfuran as a yellow oil. A solution of this oil in anhydrous *N,N*-dimethylformamide (3.00 mL) was added directly without any further purification to a mixture of spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.84 g, 3.00 mmol) and sodium hydroxide (0.48 g, 12.0 mmol) in anhydrous *N,N*-dimethylformamide (9.00 mL). The reaction mixture was heated at 70 °C for 16 h, cooled to ambient temperature followed by the addition of saturated ammonium chloride (5.0). *N,N*-Dimethylformamide was removed under high vacuum. The residue was diluted with ethyl acetate (100 mL), washed with 10% aqueous HCl (25.0 mL), brine (25.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The brown residue was subjected to column chromatography eluting with ethyl acetate/hexane (35%) to afford the title compound (0.74 g, 62%) as a colorless solid: mp 148-150 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.29 (t, 1H), 7.15-7.12 (m, 2H), 7.01 (d, 1H), 6.67 (s, 1H), 6.60 (d, 1H), 6.39 (d, 1H), 6.10 (s, 1H), 5.89 (d, 2H), 4.89 (ABq, 2H), 4.72 (ABq, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 176.9 155.8 149.7, 148.8, 142.2, 142.1, 134.8, 132.1, 129.3 124.1, 123.7 120.2, 112.0, 109.9, 108.2 103.2, 101.9, 93.8, 80.0, 57.9 37.0; MS (ES+) *m/z* 396 (M + 1).

EXAMPLE 10.63

Synthesis of 1'-[(4-hydroxy-1,2,2,6,6-pentamethylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.62, and making non-critical variations using 5,5,6,7,7-pentamethyl-1-oxa-6-azaspiro[2.5]octane to replace (5-chloro-2-furyl)methanol, the title compound was obtained (70%) as a colorless solid: mp 210-214 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.29 (d, 1H), 7.24 (t, 1H), 7.10 (d, 1H), 7.01 (t, 1H), 6.66 (s, 1H), 6.45 (s, 1H), 5.90 (d, 2H), 5.20 (br, 1H), 4.70 (ABq, 2H), 3.57 (q, 2H), 3.30 (s, 3H), 2.01-1.83 (m, 4H), 1.45 (s, 6H), 1.34 (s, 6H); ^{13}C NMR (75

MHz, DMSO-d_6) δ 178.5, 156.0, 148.7, 144.5, 142.2, 132.3, 128.9, 123.8, 123.2, 120.3, 110.9, 104.1, 101.9, 93.7, 80.9, 71.6, 65.3, 57.8, 52.6, 30.2, 28.7, 22.1; MS (ES+) *m/z* 465.4 (M + 1).

EXAMPLE 10.64

Synthesis of 1'-{[5-(2-chlorophenyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.62, and making non-critical variations using [5-(2-chlorophenyl)-2-furyl]methanol to replace (5-chloro-2-furyl)methanol, the title compound was obtained (48%) as a colorless solid: mp 148-150 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.74 (d, 1H), 7.39 (d, 1H), 7.29 (d, 1H), 7.24 (d, 1H), 7.19-7.15 (m, 2H), 7.08-7.04 (m, 3H), 6.51 (s, 1H), 6.45 (d, 1H), 6.12 (s, 1H), 6.84 (s, 2H), 4.99 (ABq, 2H), 4.78 (ABq, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.2, 155.9, 150.3, 149.0, 148.6, 142.4, 141.8, 132.1, 130.8, 130.0, 128.9, 128.7, 128.2, 127.7, 126.9, 123.9, 123.6, 119.4, 111.8, 110.7, 109.2, 103.1, 101.5, 93.6, 80.4, 58.2, 37.3; MS (ES+) *m/z* 472.2 (M + 1).

EXAMPLE 10.65

Synthesis of 1'-[(5-methyl-2-furyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.62, and making non-critical variations using (5-methyl-2-furyl)methanol to replace (5-chloro-2-furyl)methanol, the title compound was obtained (70%) as a colorless solid: mp 117-119 °C; ^{1}H NMR (300

MHz, DMSO-d_6) δ 7.26 (t, 1H), 7.12 (t, 2H), 6.99 (t, 1H), 6.67 (s, 1H), 6.32 (d, 1H), 6.07 (s, 1H), 5.97 (d, 1H), 5.89 (d, 2H), 4.84 (ABq, 2H), 4.72 (ABq, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 176.8, 155.7, 151.9, 148.8, 147.7, 142.3, 142.2, 132.1, 129.2, 124.0, 123.5, 120.4, 110.1, 110.0, 107.0, 103.2, 101.9, 93.8, 79.9, 57.9, 37.2, 13.7; MS (ES+) *m/z* 376 (M + 1).

EXAMPLE 10.66

Synthesis of 1'-[(5-bromo-2-furyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.62, and making non-critical variations using (5-bromo-2-furyl)methanol to replace (5-chloro-2-furyl)methanol, the title compound was obtained (76%) as colorless solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.29 (s, 1H), 7.69 (dt, 1H), 7.32-7.26 (m, 2H), 7.04 (d, 1H), 6.99 (d, 1H), 6.71 (d, 1H), 6.02 (s, 1H), 4.91 (ABq, 2H), 4.47 (t, 2H), 3.08 (t, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 176.9, 160.4, 156.3, 153.8, 149.7, 146.1, 137.5, 130.9, 130.8, 126.5, 125.8, 123.1, 121.5, 118.8, 116.4, 108.3, 96.7, 76.6, 71.9, 45.7, 29.1; MS (ES+) *m/z* 440.1 (M + 1), 442.1 (M + 1).

EXAMPLE 10.67

Synthesis of 1'-[(5-chloro-2-thienyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.62, and making non-critical variations using (5-chloro-2-thienyl)methanol to replace (5-chloro-2-furyl)methanol, the

title compound was obtained (77%) as a colorless solid: mp 145-146 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.28 (t, 1H), 7.20-7.14 (m, 2H), 7.10 (d, 1H), 7.01 (t, 1H), 6.95 (d, 1H), 6.67 (s, 1H), 6.09 (s, 1H), 5.89 (d, 2H), 5.02 (ABq, 2H), 4.71 (ABq, 2H); MS (ES+) *m/z* 411.9 (M + 1).

EXAMPLE 10.68

Synthesis of 1'-((5-(2-(trifluoromethyl)phenyl)furan-2-yl)methyl)-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one



Following the procedure described in EXAMPLE 10.62, and making non-critical variations using 2-(hydroxymethyl)-5-(trifluoromethyl)thiophene-3-ol to replace (5-chloro-2-furyl)methanol, the title compound was obtained (48%) as a colorless solid: mp 225-227 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.4 (s, 1H), 7.29 (dt, 1H), 7.16-7.10 (m, 3H), 7.01 (dt, 1H), 6.68 (s, 1H), 6.09 (s, 1H), 5.89 (d, 2H), 4.94 (ABq, 2H), 4.70 (ABq, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 176.9, 155.8, 152.3, 148.9, 142.2, 141.9, 132.1, 129.4, 125.9, 125.4, 124.5, 124.2, 123.8, 122.6, 120.9, 120.0, 116.5, 109.5, 103.3, 101.9, 93.8, 80.2, 57.9, 34.9; MS (ES+) *m/z* 460.38 (M - 1).

EXAMPLE 10.69

Synthesis of 1'-{[5-(2-trifluoromethylphenyl)-2-furyl]methyl}-4*H*-spiro[furo[2,3-*g*][1,3]benzodioxine-8,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.62, and making non-critical variations using {5-[2-(trifluoromethyl)phenyl]-2-furyl}methanol to replace (5-chloro-2-furyl)methanol, the title compound was obtained (28%) as a colorless solid: mp 124-

126 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.71 (d, 1H), 7.65 (d, 1H), 7.54 (t, 1H), 7.39 (t, 1H), 7.30-7.24 (m, 1H), 7.16 (d, 1H), 7.05 (d, 2H), 6.63 (s, 1H), 6.49 (s, 1H), 6.44 (d, 1H), 6.11 (s, 1H), 5.83 (d, 2H), 4.99 (ABq, 2H), 4.80 (ABq, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.2, 155.9, 150.6, 149.6, 148.9, 142.3, 141.7, 132.1, 131.8, 129.8, 128.9, 127.9, 126.8, 126.7, 126.6, 123.9, 123.6, 119.4, 111.0, 110.9, 110.6, 109.3, 103.1, 101.5, 93.6, 80.4, 58.2, 37.3; MS (ES+) *m/z* 506.27 (M + 1).

EXAMPLE 10.70

Synthesis of 1'[(2-chloro-1,3-thiazol-5-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a solution of (2-chloro-1,3-thiazol-5-yl)methanol (0.30 g, 2.00 mmol) in anhydrous CH_2Cl_2 (20.0 mL) was added thionyl chloride (0.50 g, 4.20 mmol) followed by triethylamine (0.40 g, 4.00 mmol) at 0 °C. After stirring at 0 °C for one hour and ambient temperature for one hour, the reaction mixture was diluted with CH_2Cl_2 (50.0 mL) and extracted with water (2 x 20 mL). The organic phase was dried over Na_2SO_4 and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was dissolved in methyl-ethyl ketone (10.0 mL) followed by the additions of spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.36 g, 2.00 mmol) and cesium carbonate (1.95 g, 6.00 mmol). The reaction mixture was heated at 70 °C overnight, cooled, filtered and the filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography to yield the title compound (0.032 g, 3.4%) as a colorless solid: mp 195-198 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.81 (s, 1H), 7.39-6.93 (m, 5H), 6.66 (s, 1H), 6.15-6.12 (m, 1H), 5.89 (d, 2H), 5.10 (s, 2H), 4.70 (dd, 2H); MS (ES+) *m/z* 413.1 (M + 1).

EXAMPLE 10.71

Synthesis of 1'-{[5-(trifluoromethyl)-2-furyl]methyl}-6,7-dihydrospiro[benzo[1,2-*b*:4,5-*b*']difuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure as described in EXAMPLE 10.21, and making non-critical variations using 6,7-dihydrospiro[benzo[1,2-*b*:4,5-*b*']difuran-3,3'-indol]-2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, the title compound was obtained (19%) as a white solid: mp 174-177 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.33-6.95 (m, 5H), 6.84 (s, 1H), 6.71 (d, 1H), 5.88 (s, 1H), 5.03 (ABq, 2H), 4.70 (ABq, 2H), 4.46-4.31 (m, 2H), 3.07 (t, 2H); MS (ES+) *m/z* 428.0 (M + 1).

EXAMPLE 10.72

Synthesis of 1'{[5-(trifluoromethyl)-2-furyl]methyl}5,6-dihydrospyro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.21, and making non-critical variations using 5,6-dihydrospyro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]2'(1'*H*)-one to replace 5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one, the title compound was obtained (71%) as a white solid: mp 173-176 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.50-6.90 (m, 5H), 6.73 (d, 1H), 6.38 (s, 1H), 6.32 (s, 1H), 5.04 (ABq, 2H), 4.75 (ABq, 2H), 4.55-4.36 (m, 2H), 2.88 (t, 2H); ^{13}C NMR (75 MHz, DMSO-

d_6) δ 177.2, 161.7, 161.0, 153.5, 142.0, 140.3, 139.7, 139.2, 132.6, 129.1, 124.1, 123.8, 121.2, 121.0, 120.3, 119.1, 117.7, 114.6, 114.5, 110.4, 109.6, 93.0, 80.0, 72.5, 57.3, 36.8, 28.7; MS (ES+) *m/z* 428.2 (M + 1).

EXAMPLE 10.73

Synthesis of 1'-{[5-(trifluoromethyl)-2-thienyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a solution of 1'-{[3-hydroxy-5-(trifluoromethyl)-2-thienyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.75 g, 1.62 mmol) in anhydrous dichloromethane (12.0 mL) was added triethylamine (0.49 g, 0.70 mL, 4.85 mmol) and trifluoromethanesulfonic anhydride (0.91 g, 0.50 mL, 3.24 mmol) at 0 °C under nitrogen. The reaction mixture was stirred for 30 min and quenched with saturated ammonium chloride (15.0 mL). After the aqueous layer was separated, the organic layer was washed with 10% HCl (10.0 mL), brine (10.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography to provide brown gummy material as the triflate. A mixture of this triflate (15.3 mmol), tetrakis(triphenylphosphine)palladium(0) (0.19 g, 0.17 mmol), triethylamine (1.66 g, 2.30 mL, 16.5 mmol), and formic acid (0.76 mg, 0.60 mL, 16.5 mmol) in anhydrous dioxane (24 mL) was heated at reflux for 16 h. After the reaction mixture was cooled down to ambient temperature, the solvent was removed under reduced pressure. The black residue was diluted with ethyl acetate (50.0 mL), washed with 10% HCl (20.0 mL), saturated ammonium chloride (20.0 mL), brine (20.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate:hexane (35%) to afford the title compound (0.65 g, 89%) as a colorless solid: mp 127-130 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.58-7.56 (m, 1H), 7.32-7.27 (m, 2H), 7.22 (s, 1H), 7.18 (d, 1H), 7.16 (s, 1H), 7.09 (dt, 1H), 6.68 (s, 1H), 6.10 (s, 1H), 5.89 (d, 2H), 5.17 (ABq, 2H), 4.72 (ABq, 2H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 177.0, 155.9, 148.9, 144.7,

142.2, 141.8, 132.1, 130.6, 130.5, 129.3, 128.7, 128.0, 124.3, 123.9, 120.0, 109.8, 103.3, 102.0, 93.9, 80.2, 57.8, 38.7; MS (ES+) *m/z* 446.1 (M + 1).

EXAMPLE 10.74

Synthesis of 1'-{[3-methoxy-5-(trifluoromethyl)-2-thienyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



A mixture of 1'-{[3-hydroxy-5-(trifluoromethyl)-2-thienyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.18 g, 0.39 mmol), NaOH (0.08 g, 1.96 mmol) and iodomethane (0.17 g, 1.18 mmol) in anhydrous *N*, *N*-dimethylformamide (2.00 mL) was stirred at ambient temperature for 16 h. The reaction was quenched by addition of saturated ammonium chloride (10.0 mL) and extracted with ethyl acetate (3 x 50.0 mL). The combined organic layers was washed with water (3 x 20.0 mL), brine (20.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The colorless solid was triturated with ether to give the title compound (0.15 g, 81%): mp 178-180 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.71 (s, 1H), 7.30 (dt, 1H), 7.15 (d, 1H), 7.07-7.02 (m, 2H), 6.68 (s, 1H), 6.08 (s, 1H), 5.89 (d, 2H), 4.95 (ABq, 2H), 4.70 (ABq, 2H), 3.90 (s, 3H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 177.0, 155.8, 154.9, 148.9, 142.2, 141.8, 132.1, 129.4, 125.7, 124.3, 123.8, 120.4, 120.3, 120.0, 119.5, 109.4, 103.2, 102.0, 93.9, 80.1, 59.9, 57.9, 34.9; MS (ES+) *m/z* 476.3 (M + 1).

EXAMPLE 10.75

Synthesis of 4'-methyl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



A mixture of 4'-bromo-1'-((5-(trifluoromethyl)furan-2-yl)methyl)-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one (0.51 g, 1.00 mmol), lithium chloride (0.09 mg, 2.00 mmol), $Pd_2(dba)_3$ (0.09 mg, 10 mole%) was flushed with nitrogen. To the above mixture was added anhydrous 1-methyl-2-pyrrolidinone (5.00 mL) and tetramethyltin (0.27 mg, 0.20 mL, 1.50 mmol). The reaction mixture was heated at 60 °C for 16 h and quenched with saturated ammonium chloride (10.0 mL). The reaction mixture was extracted with ethyl acetate (3 x 10.0 mL). The combined organic layers was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate:hexane (20%) to afford the title compound (0.07 g, 16%) as a colorless solid: mp 117-119 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.18 (t, 1H), 6.82 (t, 2H), 6.72 (d, 1H), 6.47 (s, 1H), 6.37 (d, 1H), 6.09 (s, 1H), 5.86 (d, 2H), 4.95 (ABq, 2H), 4.83 (ABq, 2H), 2.03 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.6, 156.2, 152.0, 149.1, 142.3, 141.3, 135.6, 129.4, 128.9, 126.0, 120.6, 117.2, 112.7, 112.6, 109.2, 106.5, 102.9, 101.6, 93.3, 78.4, 58.3, 37.0, 17.1; MS (ES+) *m/z* 444.1 (M + 1).

EXAMPLE 10.76

Synthesis of 5'-methyl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making *non-critical*

variations using 5'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, the title compound was obtained (77%): mp 96-98 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.09 (d, 1H), 7.00 (s, 1H), 6.87 (d, 1H), 6.74 (d, 1H), 6.52 (s, 1H), 6.38 (d, 1H), 6.11 (s, 1H), 5.88 (d, 2H), 4.96 (ABq, 2H), 4.80 (ABq, 2H), 2.29 (s, 3H); MS (ES+) *m/z* 444.2 (M + 1).

EXAMPLE 10.77

Synthesis of 1'-({5-[4-(trifluoromethyl)phenyl]-1,2,4-oxadiazol-3-yl}methyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 3-chloromethyl-5-[4-(trifluoromethyl)phenyl]-1,2,4-oxadiazole to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (44%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.22 (d, 2H), 7.76 (d, 2H), 7.30-7.18 (m, 2H), 7.06 (t, 1H), 6.91 (d, 1H), 6.51 (s, 1H), 6.40 (s, 1H), 5.88 (s, 2H), 5.17 (ABq, 2H), 4.86 (ABq, 2H); MS (ES+) *m/z* 508.1 (M + 1).

EXAMPLE 10.78

Synthesis of 1'-(2-thienylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.62, and making non-critical variations using 2-thiophenemethanol to replace (5-chloro-2-furyl)methanol, the title

compound was obtained (37%) as a colorless solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.26-7.20 (m, 2H), 7.18-7.13 (m, 1H), 7.10-6.98 (m, 2H), 6.97-6.90 (m, 2H), 6.50 (s, 1H), 6.12 (s, 1H), 5.85 (d, 2H), 5.10 (ABq, 2H), 4.79 (ABq, 2H); MS (ES+) *m/z* 378.19 (M + 1).

EXAMPLE 10.79

Synthesis of 5-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]thiophene-2-carbonitrile



A mixture of 1'-[(5-bromo-2-thienyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.23 g, 0.49 mmol), zinc cyanide (0.07 g, 0.59 mmol), tris(dibenzylideneacetone)dipalladium(0) (0.10 g, 0.11 mmol), 1,1'-bis(diphenylphosphino)ferrocene (0.06 g, 0.11 mmol), *N*,*N*-dimethylformamide (6.00 mL) and a catalytic amount of water (2 drops) was heated at 120 °C for 24 hours. After cooling down to ambient temperature, the organic solvent was evaporated *in vacuo*. The residue was extracted with ethyl acetate (5 x 15.0 mL) and the combined organic solution was passed through a bed of celite. The filtrate was successively washed with saturated aqueous ammonium chloride (25.0 mL), water (2 x 35.0 mL) and brine (40.0 mL). The organic layer was dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluted with 20-35% ethyl acetate in hexanes to afford solid which was further purified by preparative thin layer chromatography, eluted with 20% ethyl acetate in hexanes to afford the title compound (0.09 g, 44%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.50 (d, 1H), 7.32-7.16 (m, 2H), 7.12-7.04 (m, 2H), 6.86 (d, 1H), 6.51 (s, 1H), 6.08 (s, 1H), 5.87 (d, 2H), 5.10 (ABq, 2H), 4.78 (ABq, 2H); MS (ES+) *m/z* 403.0 (M + 1).

EXAMPLE 10.80

Synthesis of 5-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-2-furonitrile



Following the procedure described in EXAMPLE 10.79, and making non-critical variations using 1'-[(5-bromo-2-furyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 1'-[(5-bromo-2-thienyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, the title compound was obtained (44%) as a colorless solid: mp 167-169 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.34-7.27 (m, 1H), 7.21-7.17 (m, 1H), 7.12-7.08 (m, 1H), 7.07-7.03 (m, 1H), 6.95 (d, 1H), 6.51 (s, 1H), 6.44 (d, 1H), 6.08 (s, 1H), 5.86 (q, 2H), 4.96 (ABq, 2H), 4.78 (ABq, 2H); MS (ES+) *m/z* 387.2 (M + 1).

EXAMPLE 10.81

Synthesis of 1'-{[5-(methylsulfonyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



A mixture of 1'-[(5-bromo-2-furyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.70 g, 1.59 mmol), sodium methanesulfinate (85%, 0.23 g, 1.91 mmol), copper (I) iodide (0.04 g, 0.22 mmol), L-proline (0.04 g, 0.35 mmol) and dimethyl sulfoxide (4.00 mL) was heated at 100 °C. After 3 days, the reaction mixture was cooled down to ambient temperature, quenched with water (50.0 mL) and extracted with ethyl acetate (3 x 40.0 mL). The combined organic layers was washed with brine (2 x 50.0 mL), dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography eluting with ethyl acetate:hexane (30-50%) to afford the title

compound (0.50 g, 71%): mp 177-179 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.29 (t, 1H), 7.19 (d, 1H), 7.12-7.04 (m, 2H), 6.94 (d, 1H), 6.50 (s, 1H), 6.42 (d, 1H), 6.11 (s, 1H), 5.86 (s, 2H), 5.00 (ABq, 2H), 4.79 (ABq, 2H), 3.11 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.3, 155.9, 154.5, 149.1, 149.1, 142.4, 141.1, 131.9, 129.1, 124.2, 124.1, 119.0, 118.4, 109.9, 108.7, 102.9, 101.6, 93.7, 80.3, 58.2, 43.4, 37.2; MS (ES+) *m/z* 440.0 (M + 1).

EXAMPLE 10.82

Synthesis of 1'-[(6-oxo-1,6-dihydropyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a mixture of 1'-[(6-methoxypyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.23 g, 0.57 mmol), sodium iodide (0.28 g, 1.87 mmol), water (2 drops) in anhydrous acetonitrile (5.00 mL) was added chlorotrimethylsilane (0.19 g, 1.78 mmol) at 0 °C. The reaction mixture was stirred at ambient temperature for 16 h and quenched with sodium bisulfite (0.20 g). The reaction mixture was diluted with ethyl acetate (100 mL) and washed with brine (2 x 25.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was triturated with ether (2 x 10.0 mL) to give the title compound (0.16 g, 72%) as la ight yellow solid: mp 247-250 °C; ^{1}H NMR (300 MHz, DMSO-d_6,) δ 11.50 (br, 1H), 7.50 (bd, 1H), 7.36-7.24 (m, 2H), 7.17-7.12 (m, 2H), 7.01 (dt, 1H), 6.67 (s, 1H), 6.28 (d, 1H), 6.09 (s, 1H), 5.91- 5.88 (m, 2H), 4.78 (d, 1H), 4.67- 4.62 (m, 3H); MS (ES+) *m/z* 389.15 (M + 1).

EXAMPLE 10.83

Synthesis of 1'-[(1-methyl-6-oxo-1, 6-dihydropyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in PREPARATION 1A, and making non-critical variations using 1'-[(6-oxo-1,6-dihydropyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4-bromoindole, and methyl iodide to replace 1-bromopentane, the title compound was obtained (78%): mp 115-118 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.39-7.23 (m, 3H), 7.18 (d, 1H), 7.06 (t, 1H), 6.87 (d, 1H), 6.57-6.48 (m, 2H), 6.02 (s, 1H), 5.87-5.83 (m, 2H), 4.90 (d, 1H), 4.75-4.52 (m, 3H), 3.51 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.8, 162.5, 155.9, 149.1, 142.4, 141.6, 139.7, 137.6, 132.1, 129.1, 124.4, 123.9, 121.3, 119.1, 114.0, 108.6, 102.8, 101.6, 93.7, 80.3, 58.2, 40.8, 38.0; MS (ES+) *m/z* 403.3 (M + 1).

EXAMPLE 10.84

Synthesis of 5-bromo-1'-[(5-chloro-2-thienyl)methyl]spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10, and making non-critical variations using 5-bromospiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one to replace

spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 2-chloro-5-(chloromethyl)thiophene to replace 4-fluorobenzyl bromide, the title compound was obtained (95%) as a white solid: mp 140-142 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.33-7.26 (m, 2H), 7.16-7.02 (m, 2H), 6.94 (d, 1H), 6.97-6.75 (m, 4H), 5.07-4.91 (m, 3H), 4.68 (d, 1H); MS (ES+) *m/z* 446.7 (M + 1), 448.7 (M + 1).

EXAMPLE 10.85

Synthesis of 1'-[(5-chloro-1,3,4-thiadiazol-2-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a solution of spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.56 g, 1.99 mmol) and (5-chloro-1,3,4-thiadiazol-2-yl)methanol (0.30 g, 1.99 mmol) in anhydrous tetrahydrofuran (12.0 mL) was added tributylphosphine (0.60 g, 2.99 mmol) at 0 °C. The reaction mixture was stirred for 15 min followed by the addition of *N*,*N*,*N*',*N*'-tetramethylazodicarboxamide (0.51 g, 2.99 mmol). The reaction mixture was stirred at ambient temperature overnight, quenched with aqueous ammonium chloride (10.0 mL) and diluted with ethyl acetate (350 mL). The organic layer was washed with aqueous saturated sodium chloride (2 x 25.0 mL), dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to column chromatography (ethyl acetate/hexane, 1/2) to give the title compound (0.20 g, 24%) as a yellowish solid: mp 194-197 °C; ^{1}H NMR (300 MHz, DMSO-d_6,) δ 7.30 (dt, 1H), 7.20-7.12 (m, 2H), 7.05 (dt, 1H), 6.67 (s, 1H), 6.28 (s, 1H), 5.90 (s, 2H), 5.43 (d, 1H), 5.34 (d, 1H), 4.78 (d, 1H), 4.67 (d, 1H); ^{13}C NMR (75 MHz, DMSO-d_6) δ 177.2, 168.3, 155.8, 155.4, 148.9, 142.2, 141.9, 132.2, 129.4, 124.2, 124.0, 120.1, 109.8, 103.7, 101.9, 93.8, 80.1, 67.5, 57.9, 25.6; MS (ES+) *m/z* 414.2 (M + 1), 416.2 (M + 1).

EXAMPLE 10.86

Synthesis of 1'-[(1-pyridin-2-ylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



A mixture of 1'-(piperidin-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrobromide(0.20 g, 0.44 mmol), 2-bromopyridine (0.16 mL, 0.65 mmol), tetrabutyl ammonium iodide (0.05 g) and DBU (0.16 mL, 1.09 mmol) in DMF (5.00 mL) was heated at 120 °C for 15 hrs. After cooling down to ambient temperature, water (30.0 mL) was added. The above mixture was extracted twice with ethyl acetate (50.0 mL), the combined organic phase was dried over Na_2SO_4 and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to flash chromatography eluting with 30% ethyl acetate in hexane to give a white solid (0.05 g, 27%): mp 95-97 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.17 (d, 1H), 7.47 (td, 1H), 7.31 (t, 1H), 7.18 (d, 1H), 7.06 (t, 1H), 6.92 (d, 1H), 6.66 (d, 1H), 6.59 (dd, 1H), 6.52 (s, 1H), 6.12 (s, 1H), 5.91-5.84 (m, 2H), 4.91 (d, 1H), 4.66 (d, 1H), 4.42-4.27 (m, 2H), 3.82-3.53 (m, 2H), 2.85 (t, 2H), 2.22-2.05 (m, 1H), 1.85-1.70 (m, 2H), 1.53-1.35 (m, 2H); ^{13}C NMR(75 MHz, $CDCl_3$) δ 177.9, 158.8, 156.1, 149.0, 147.4, 142.8, 142.5, 138.0, 132.4, 129.1, 124.2, 123.5, 119.5, 112.9, 108.8, 107.7, 103.1, 101.7, 93.8, 80.7, 58.3, 46.0, 45.5, 45.4, 35.2, 29.8, 29.7; MS (ES+) *m/z* 456 (M + 1).

EXAMPLE 10.87

Synthesis of 1'-[(1-phenyl-2-ylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



A mixture of 1'-(piperidin-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-

indol]-2'(1'*H*)-one hydrobromide (0.20 g, 0.44 mmol), 2-bromobenzene(0.07 mL, 0.65 mmol), $Pd_2(dba)_3$ (0.03 g, 0.03 mmol), BINAP (0.06 g, 0.10 mmol) and $NaOBu^t$ (0.13 g, 1.30 mmol) in toluene was heated at 100 °C for 15 hrs under nitrogen. After cooling down to ambient temperature, water (30.0 mL) was added. The above mixture was extracted twice with ethyl acetate (50.0 mL), the combined organic phase was dried over Na_2SO_4 and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to flash chromatography eluting 30 % ethyl acetate in hexane to give a white solid (0.10 g, 48%): mp 76-78 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.37-7.16 (m, 4H), 7.12-6.80 (m, 5H), 6.53 (s, 1H), 6.14 (s, 1H), 5.87 (dd, 2H), 4.92 (d, 1H), 4.66 (d, 1H), 3.87-3.55 (m, 4H), 2.72 (t, 2H), 2.12-1.94 (m, 1H), 1.89-1.73 (m, 2H), 1.71-1.45 (m, 2H); ^{13}C NMR(75 MHz, $CDCl_3$) δ 177.9, 156.1, 149.0, 142.9, 142.5, 132.4, 129.2, 129.1, 124.2, 123.4, 119.9, 119.5, 116.9, 108.8, 103.1, 101.6, 93.8, 80.8, 60.5, 58.3, 49.8, 46.1, 34.8, 30.1, 30.0; MS (ES+) *m/z* 455 (M + 1).

EXAMPLE 10.88

Synthesis of 1'-(pyridin-2-ylmethyl)-6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one hydrochloride



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using 6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 2-(bromomethyl)pyridine hydrobromide to replace 2-(bromomethyl)-2-(trifluoromethyl)furan, 1'-(pyridin-2-ylmethyl)-6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one was obtained, which was treated with 4.0 M HCl in dioxane to give the title compound (39%): mp 150-152 °C; ^{1}H NMR (300 MHz, CD_3OD) δ 8.89-8.78 (m, 1 H), 8.62-8.47 (m, 1H), 8.07-7.00 (m, 2H), 7.42-6.70 (m, 7H), 5.52-5.31 (m, 2H), 5.05 (d, 1H), 4.89 (d, 1H); ^{13}C NMR(75 MHz, CD_3OD) δ 179.4, 163.5, 153.1, 152.7, 148.2, 143.8, 142.8, 133.1, 130.6, 129.3, 127.4, 126.8, 125.7, 125.6, 125.4, 114.9, 110.4, 104.9, 82.0, 58.9, 43.0; MS (ES+) *m/z* 413 (M + 1).

EXAMPLE 10.89

Synthesis of 1'-(pyridin-3-ylmethyl)-6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one hydrochloride



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using 6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 3-(bromomethyl)pyridine hydrobromide to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, 1'-(pyridin-3-ylmethyl)-6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one was obtained, which was treated with 4.0 M HCl in dioxane to give the title compound (70%) as a white solid: mp 151-153 °C; ^{1}H NMR (300 MHz, CD_3OD) δ 9.07-8.61 (m, 3 H), 8.19-8.04 (m, 1H), 7.42-6.71 (m, 7H), 5.28 (s, 2H), 5.05 (d, 1H), 4.86 (d, 1H); ^{13}C NMR (75 MHz, CD_3OD) δ 179.43, 163.5, 151.8, 147.0, 142.9, 142.4, 142.2, 133.0, 130.6, 129.3, 128.9, 125.5, 125.4, 125.3, 114.9, 110.5, 104.9, 82.0, 58.9, 42.0; MS (ES+) *m/z* 413 (M + 1).

EXAMPLE 10.90

Synthesis of 6-(trifluoromethoxy)-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using 6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, the title compound was obtained (82%) as a white solid: mp 78-80 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.33 (td, 1H), 7.21-6.98 (m, 3H), 6.86-6.73 (m, 2H), 6.67 (s, 2H), 6.42 (d, 1H),

5.09 (d, 1H), 5.04 (d, 1H), 4.88 (d, 1H), 4.77 (d, 1H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.7, 161.7, 151.9, 150.6, 141.5, 131.7, 129.5, 127.5, 124.2, 124.2, 124.0, 114.1, 112.8, 112.8, 109.6, 109.2, 104.3, 80.7, 57.6, 37.1; MS (ES+) *m/z* 470 (M + 1).

EXAMPLE 10.91

Synthesis of 1'-(4-methoxybenzyl)-6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using 6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 4-methoxybenzyl chloride to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (91%) as a white solid: mp 82-84 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.32-6.80 (m, 9H), 6.68 (s, 2H), 5.06 (d, 1H), 5.03 (d, 1H), 4.80 (d, 1H), 4.77 (d, 1H), 3.80 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 171.1, 161.8, 159.4, 150.5, 142.3, 132.0, 129.2, 128.9, 127.7, 124.1, 124.0, 123.7, 122.2, 118.8, 114.4, 114.1, 109.7, 104.3, 80.9, 57.6, 55.4, 43.9; MS (ES+) *m/z* 442 (M + 1).

EXAMPLE 10.92

Synthesis of 1'-(cyclohexylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and bromomethyl

cyclohexane to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (74%) as a white solid: mp 153-154 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.30 (td, 1H), 7.16 (d, 1H), 7.04 (t, 1H), 6.90 (d, 1H), 6.51 (s, 1H), 6.14 (s, 1H), 5.90-5.84 (m, 2H), 4.91 (d, 1H), 4.65 (d, 1H), 3.72-3.44 (m, 2H), 1.94-1.60 (m, 6H), 1.32-0.99 (m, 5H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.8, 156.1, 148.9, 143.0, 142.4, 132.5, 128.9, 124.0, 123.2, 119.7, 109.0, 103.2, 101.6, 93.7, 80.8, 58.3, 46.8, 36.3, 31.1, 31.0, 26.4, 25.9, 25.8; MS (ES+) *m/z* 378 (M + 1), 400 (M + 23).

EXAMPLE 10.93

Synthesis of 1'-(methylsulfonyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



Following the procedure described in EXAMPLE 10.47, and making non-critical variations using spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and methanesulfonyl chloride to replace 2-(bromomethyl)-5-(trifluoromethyl)furan, the title compound was obtained (51%) as a white solid: mp 215-217 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.83 (d, 1H), 7.42-7.31 (m, 1H), 7.25-7.17 (m, 2H), 6.52 (s, 1H), 6.20 (s, 1H), 5.93-5.87 (m, 2H), 4.98 (d, 1H), 4.68 (d, 1H), 3.46 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.9, 155.8, 149.5, 142.7, 138.1, 130.5, 129.7, 125.9, 124.3, 118.5, 113.8, 102.9, 101.8, 93.8, 80.6, 58.8, 41.8; MS (ES+) *m/z* 360 (M + 1), 382 (M + 23).

EXAMPLE 10.94

Synthesis of 1'-(2-piperidin-1-ylethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



A mixture of 1'-(2-aminoethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.20 g, 0.62 mmol), 1,5-dibromopentane (0.08 mL, 0.62 mmol) and triethyl amine (0.17 mL, 1.23 mmol) in THF (10.0 mL) was refluxed for 15 hrs and concentrated *in vacuo* to dryness. The residue was subjected to flash chromatography eluting with 10% methanol in ethyl acetate to give 1'-(2-piperidin-1-ylethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, which was treated with 4.0 M HCl in dioxane to give the title compound (28%): mp >240 °C; ^{1}H NMR (300 MHz, CD_3OD) δ 7.40 (t, 1H), 7.27-7.11 (m, 3H), 6.51 (s, 1H), 6.17 (s, 1H), 5.86 (s, 2H), 4.91 (d, 1H), 4.71 (d, 1H), 4.40-4.13 (m, 2H), 3.95-3.84 (m, 1H), 3.66-3.37 (m, 3H), 3.14-2.96 (m, 2H), 2.06-1.45 (m, 6H); ^{13}C NMR (75 MHz, CD_3OD) δ 180.4, 157.5, 150.5, 143.8, 142.5, 133.9, 130.3, 125.3, 125.1, 120.6, 110.2, 103.9, 102.9, 94.2, 81.4, 59.7, 55.4, 54.9, 53.9, 36.4, 24.2, 22.6; MS (ES+) *m/z* 393 (M + 1).

EXAMPLE 10.95

Synthesis of 1'-[2-(pyridin-2-ylamino)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one and 1'-[2-(dipyridin-2-ylamino)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



and



Following the procedure described in EXAMPLE 10.87, and making non-critical variations using 1'-(2-aminoethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one to replace 1'-(piperidin-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, and 2-bromopyridine to replace 2-bromobenzene, 1'-(2-(pyridin-2-ylamino)ethyl)-spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one was obtained as the first fraction from the chromatography as a white solid (5%): mp 61-63 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.05 (d, 1H), 7.50-6.97 (m, 5H), 6.57 (dd, 1H), 6.50 (s, 1H), 6.38 (d, 1H), 6.03 (s, 1H), 5.85 (s, 1H), 5.84 (s, 1H), 4.84 (d, 1H), 4.79 (t, 1H), 4.60 (d, 1H), 4.15-3.94 (m, 2H), 3.81-3.64 (m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.4, 158.2, 156.0, 149.0, 148.0, 142.4, 142.38, 137.4, 132.3, 129.1, 124.0, 123.5, 119.5, 113.3, 109.0,

108.3, 103.2, 101.6, 93.7, 80.5, 58.3, 40.1, 39.9; MS (ES+) *m/z* 402 (M + 1). 1'-(2-(dipyridin-2-ylamino)ethyl)-spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one was obtained as the second fraction from the chromatography (31%): mp 165-167 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.31 (dd, 2H), 7.48 (d, 2H), 7.23 (t, 1H), 7.09-6.83 (m, 7H), 6.47 (s, 1H), 5.95 (s, 1H), 5.88-5.81(m, 2H), 4.73 (d, 1H), 4.67-4.49(m, 2H), 4.46 (d, 1H), 4.20 (t, 2H); ^{13}C NMR(75 MHz, $CDCl_3$) δ 177.6, 157.0, 156.0, 148.8, 148.4, 143.1, 142.3, 137.4, 132.3, 128.9, 123.6, 123.1, 119.6, 117.5, 114.5, 109.3, 103.4, 101.6, 93.6, 80.7, 58.2, 45.9, 39.5; MS (ES+) *m/z* 479 (M + 1).

EXAMPLE 10.96

Synthesis of *tert*-butyl 4-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]piperidine-1-carboxylate



Following the procedure as described in EXAMPLE 10, and making non-critical variations using *tert*-butyl 4-(2-{[(4-methylphenyl)sulfonyl]oxy}ethyl)piperidine-1-carboxylate to replace 4-fluorobenzyl bromide, the title compound was obtained in 95% yield: mp 173-175 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.31 (t, 1H), 7.17 (d, 1H), 7.06 (t, 1H), 6.88 (d, 1H), 6.51 (s, 1H), 6.10 (s, 1H), 5.90-5.84 (m, 2H), 4.90 (d, 1H), 4.65 (d, 1H), 4.0-3.64 (m, 4H), 2.75-2.58 (m, 2H), 1.85-1.09 (m, 16H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.5, 156.0, 155.0, 149.0, 142.4, 142.2, 132.6, 129.0, 124.2, 123.4, 119.5, 108.6, 103.0, 101.6, 93.8, 80.5, 79.5, 58.3, 38.0, 34.0, 33.9, 32.1, 31.9, 28.6; MS (ES+) *m/z* 515 (M + 23), 393 (M - 100).

EXAMPLE 10.97

Synthesis of 1'-(2-piperidin-4-ylethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



To a solution of *tert*-butyl 4-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]piperidine-1-carboxylate (0.94 g, 1.91 mmol) in dioxane (5.00 mL) was added 4.0 M HCl in dioxane (2.00 mL, 8.00 mmol). The mixture was stirred at ambient temperature for 30 min followed by the addition of anhydrous ether (40.0 mL). The precipitated white solid was filtered, washed with ether and dried to give the title compound (0.75 g, 91%): ^{1}H NMR (300 MHz, CD_3OD) δ 7.37 (t, 1H), 7.20-7.07 (m, 3H), 6.52 (s, 1H), 6.10 (s, 1H), 5.86 (s, 2H), 4.83 (d, 1H), 4.67 (d, 1H), 3.97-3.75 (m, 2H), 3.45-3.33 (m, 2H), 3.01-2.85 (m, 2H), 2.15-2.01 (m, 2H), 1.82-1.37 (m, 5H); ^{13}C NMR(75 MHz, CD_3OD) δ 179.8, 157.6, 150.4, 143.7, 143.4, 133.6, 130.3, 124.9, 124.8, 120.8, 110.3, 103.7, 102.9, 94.3, 81.4, 59.8, 45.2, 38.5, 34.4, 32.6, 29.8, 29.7; MS (ES+) *m/z* 393 (M + 1).

EXAMPLE 10.98

Synthesis of 1'-[2-(1-cyclopentylpiperidin-4-yl)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



To a solution of cyclopentanone (0.04 mL, 0.45 mmol) and triethyl amine (0.12

mL, 0.84 mmol) in dichloroethane (5.00 mL) was added 1'-(2-piperidin-4-ylethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride (0.12 g, 0.28 mmol) and sodium triacetoxyborohydride (0.10 g, 0.45 mmol). The reaction mixture was stirred for 16 hrs and concentrated *in vacuo* to dryness. The residue was subjected to flash chromatography eluting with ethyl acetate /methanol/ammonium hydroxide (15/1/0.1) to give 1'-[2-(1-cyclopentylpiperidin-4-yl)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one as a white solid, which was treated with 4.0 M HCl in dioxane to give the title compound (0.05 g, 32% yield): mp 153-155 °C; ^{1}H NMR (300 MHz, CD_3OD) δ 7.37 (td, 1H), 7.21-7.08 (m, 3H), 6.53 (s, 1H), 6.11 (s, 1H), 5.88 (s, 1H), 5.87 (s, 1H), 4.83 (d, 1H), 4.69 (d, 1H), 3.98-3.75 (m, 2H), 3.68-3.38 (m, 3H), 3.01-2.83 (m, 2H), 2.25-2.08 (m, 4H), 1.92-1.37 (m, 11H); ^{13}C NMR (75 MHz, CD_3OD) δ 179.8, 157.6, 150.4, 143.8, 143.4, 133.6, 130.3, 124.9, 124.8, 120.8, 110.3, 103.7, 103.0, 94.3, 81.4, 69.1, 59.8, 53.2, 38.6, 34.3, 32.5, 30.6, 30.5, 29.4, 24.7; MS (ES+) *m/z* 461 (M + 1).

EXAMPLE 10.99

Synthesis of 1'-[2-(1-isopropylpiperidin-4-yl)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



Following the procedure as described in EXAMPLE 10.98, and making non-critical variations using acetone to replace cyclopentanone, the title compound was obtained (42%) as a white solid: mp 155-156 °C; ^{1}H NMR (300 MHz, CD_3OD) δ 7.37 (t, 1H), 7.21-7.08 (m, 3H), 6.53 (s, 1H), 6.11 (s, 1H), 5.87 (s, 2H), 4.84 (d, 1H), 4.69 (d, 1H), 3.98-3.75 (m, 2H), 3.58-3.38 (m, 3H), 3.05-2.85 (m, 2H), 2.23-2.09 (m, 2H), 1.82-1.44 (m, 5H), 1.35 (d, 6H); ^{13}C NMR (75 MHz, CD_3OD) δ 179.8, 157.6, 150.5, 143.8, 143.4, 133.6, 130.3, 125.0, 124.8, 120.8, 110.3, 103.7, 103.0, 94.3, 81.4, 59.8, 59.6, 38.6, 34.2, 32.7, 30.6, 30.5, 24.2, 16.9. 15.4; MS (ES+) *m/z* 435 (M + 1).

EXAMPLE 10.100

Synthesis of 1'-[2-(1-cyclobutylpiperidin-4-yl)ethyl]spiro[furo[2,3 *f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



Following the procedure as described in EXAMPLE 10.98, and making non-critical variations using cyclobutanone to replace cyclopentanone, the title compound was obtained (81%) as a white solid: mp 158-160 °C; ^{1}H NMR (300 MHz, CD_3OD) δ 7.37 (t, 1H), 7.21-7.05 (m, 3H), 6.52 (s, 1H), 6.10 (s, 1H), 5.86 (s, 2H), 4.83 (d, 1H), 4.67 (d, 1H), 3.98-3.39 (m, 5H), 2.85-2.59 (m, 2H), 2.43-1.42 (m, 13H); ^{13}C NMR (75 MHz, CD_3OD) δ 179.8, 157.6, 150.4, 143.7, 143.4, 133.6, 130.3, 124.9, 124.8, 120.8, 110.4, 103.8, 103.0, 94.2, 81.5, 60.5, 59.8, 50.8, 38.6, 34.4, 32.5, 30.1, 26.8, 14.4; MS (ES+) *m/z* 447 (M + 1).

EXAMPLE 10.101

Synthesis of 1'-{2-[1-(tetrahydro-2*H*-pyran-4-yl)piperidin-4-yl]ethyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride



Following the procedure as described in EXAMPLE 10.98, and making non-critical variations using tetrahydro-4*H*-pyran-4-one to replace cyclopentanone, the title

compound was obtained (45%) as a white solid: mp 168-170 °C; ^{1}H NMR (300 MHz, CD_3OD) δ 7.37 (t, 1H), 7.21-7.06 (m, 3H), 6.52 (s, 1H), 6.11 (s, 1H), 5.86 (s, 2H), 4.83 (d, 1H), 4.67 (d, 1H), 4.12-3.31 (m, 9H), 3.05-2.85 (m, 2H), 2.25-1.45 (m, 11H); ^{13}C NMR (75 MHz, CD_3OD) δ 179.8, 157.6, 150.4, 143.7, 143.4, 133.6, 130.3, 125.0, 124.8, 120.8, 110.3, 103.8, 102.9, 94.3, 81.4, 67.2, 64.1, 59.8, 50.7, 38.6, 34.2, 32,7, 30.6, 30.5, 28.7; MS (ES+) *m/z* 477 (M + 1).

EXAMPLE 11

Synthesis of 4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoic acid



Following the procedure described in EXAMPLE 6, and making non-critical variations using methyl 4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoate to replace methyl 2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoate, the title compound was obtained (100%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 12.96 (s, 1H), 7.90 (d, 2H), 7.43 (d, 2H), 7.22 (t, 1H), 7.17 (d, 1H), 7.00 (t, 1H), 6.94 (d, 1H), 6.68 (s, 1H), 6.21 (s, 1H), 5.90 (s, 2H), 4.98 (s, 2H), 4.76 (ABq, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.4, 167.5, 156.0, 148.9, 142.6, 142.3, 141.8, 132.1, 130.5, 130.3, 129.3, 127.7, 124.2, 123.7, 120.1, 109.9, 103.5, 101.9, 93.8, 80.4, 58.0, 43.4.

EXAMPLE 12

Synthesis of *N*-(3-fluorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide



A. Preparation of stock solution of 4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoyl chloride

To a stirred slurry of 4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoic acid (2.08 g, 5.00 mmol) in dry chloroform (50.0 mL) was added oxalyl chloride (0.95 g, 7.50 mmol) at ambient temperature followed by 1 drop of DMF. The mixture was stirred at ambient temperature for 2 h and evaporated to dryness *in vacuo*. The residue was dissolved in dry dichlormethane (60.0 mL) to form an acid chloride stock solution for use.

B. Synthesis of *N*-(3-fluorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide

To a solution of 3-fluorophenylamine (0.02 mL, 0.24 mmol) in dry dichloromethane (2.00 mL) and triethylamine (0.05 mL, 0.32 mmol) was added the acid chloride stock solution (2.0 mL, 0.081 M in dichloromethane) obtained above at ambient temperature. The mixture was stirred for 2 h, washed with 15% HCl solution and water. The organic layer was separated, dried over Na_2SO_4 and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was dissolved in ethyl acetate, and the product was precipitated by the addition of hexane. The white solid was filtered and collected to yield the title compound (0.06 g) in 70% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.30 (s, 1H), 7.83 (d, 2H), 7.58 (ddd, 1H), 7.38 (d, 2H), 7.27-7.23 (m, 2H), 7.21-7.16 (m, 2H), 7.04 (dt, 1H), 6.85-6.78 (m, 1H), 6.74 (d, 1H), 6.46 (s, 1H), 6.10 (s, 1H), 5.77 (d, 1H), 5.68 (d, 1H), 4.97 (ABq, 2H), 4.76 (ABq, 2H); MS (ES+), *m/z* 509.1 (M + 1).

EXAMPLE 12.1

The compounds listed in the following table were prepared using the similar procedure as described in EXAMPLE 12. As previously noted, the compound numbers listed below do not correspond to the compound numbers provided in the general Reaction Schemes above.

Compound Number	Name	MS (*m/z*, M + 1)
500	*N*-butyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	471.2
501	*N*-(3-fluorobenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	523.1

Compound Number	Name	MS (*m/z*, M + 1)
502	1'-[4-(piperidin-1-ylcarbonyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	483.2
503	*N,N*-diisopropyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	499.1
504	*N*-(4-chlorobenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	540.1
505	*N*-(3-chlorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	526.1
506	*N*-(2-fluorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	509.2
507	*N*-(2-ethylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.2
508	*N*-(4-ethylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'H)-yl)methyl]benzamide	519.2
509	*N*-(4-methylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	505.2
510	*N*-(3,5-dimethylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.2
511	*N*-(2,3-dimethylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.2
512	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pentylbenzamide	485.1
513	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-propylbenzamide	457.1

Compound Number	Name	MS (*m/z*, M + 1)
514	*N*-isopropyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	457.1
515	*N*-isobutyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	471.1
516	*N*-hexyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	499.2
517	*N*-cyclohexyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	497.2
518	*N*-cyclopentyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	483.2
519	*N*-heptyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	513.2
520	*N*-(2-methoxybenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	535.2
521	*N*-(2,6-dimethylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.2
522	*N*-(2-methoxyphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	521.2
523	*N*-cyclopropyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	455.1
524	*N*-(3-methoxypropyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	487.2
525	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-phenylbenzamide	491.2

Compound Number	Name	MS (*m/z*, M + 1)
526	*N*-(2,4-dimethylphenyl)-4-[(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'H)-yl)methyl]benzamide	519.2
527	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(tetrahydrofuran-2-ylmethyl)benzamide	499.2
528	*N,N*-dibenzyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	595.2
529	*N*-[2-(diethylamino)ethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	514.2
530	*N*-methyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	429.2
531	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[3-(trifluoromethyl)phenyl]benzamide	559.1
532	*N*-ethyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	443.1
533	*N*-(3-ethoxypropyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	501.2
534	*N*-(4-methoxybenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	535.2
535	*N*-(3,5-dichlorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	560.2
536	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pyridin-3-ylbenzamide	492.2
537	*N*-(4-cyanophenyl)-4-[(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	516.2

Compound Number	Name	MS (*m/z*, M + 1)
538	*N*-(4-methylpentyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	499.2
539	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2,2,2-trifluoroethyl)benzamide	497.2
540	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(pyridin-4-ylmethyl)benzamide	506.1
541	*N*-[2-(3-chlorophenyl)ethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	554
542	*N*-(2-furylmethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	495.1
543	*N*-(3-fluoro-2-methylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	523.1
544	*N*-hexyl-*N*-methyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	513.1
545	*N*-(3-isopropoxypropyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	515.2
546	*N*-(2-ethoxyethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	487.2
547	*N*-(cyclopropylmethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'H)-yl)methyl]benzamide	469.1
548	*N*-(4-methoxyphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	521.1

Compound Number	Name	MS (*m/z*, M + 1)
549	*N*-cyclobutyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	469.1
550	*N*-(2,2-diphenylethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	595.2
551	*N*-[2-(4-fluorophenyl)ethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	537.2
552	*N*-(cyclohexylmethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	511.2
553	*N*-(2-fluoro-4-methylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	523.2
554	*N*-[2-(4-methylphenyl)ethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	533.2
555	*N*-(2-ethylbutyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	499.2
556	*N*-benzyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	505.1
557	*N*-(2-methoxyethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	473.1
558	1'-[4-(morpholin-4-ylcarbonyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	485.1
559	*N*-(1-benzylpiperidin-4-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	588.2
560	*N*-[2-(4-methoxyphenyl)ethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	549.2

Compound Number	Name	MS (*m/z*, M + 1)
561	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(trifluoromethyl)phenyl]benzamide	559.1
562	*N*-[4-chloro-2-(trifluoromethyl)phenyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	594.1
563	*N*-[4-fluoro-2-(trifluoromethyl)phenyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	577.1
564	*N*-(2-cyanoethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	468.1
565	*N*-[(1S)-1-cyclohexylethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	525.2
566	*N*-[(1R)-1-cyclohexylethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	525.2
567	*N*-(2,4-difluorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	527.1
568	*N*-(2,3-dihydro-1*H*-inden-1-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	531.2
569	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-thienylmethyl)benzamide	511.2
570	*N*-[(1-ethylpyrrolidin-2-yl)methyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	526.2
571	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(4-propylphenyl)benzamide	533.2
572	*N*-(2,5-difluorobenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	541.1

Compound Number	Name	MS (*m/z*, M + 1)
573	*N*-(2,3-dihydro-1*H*-inden-5-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	531.2
574	*N*-(2,5-difluorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	527.1
575	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[4-(trifluoromethyl)benzyl]benzamide	573.1
576	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(2-thienyl)ethyl]benzamide	525.1
577	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(pyridin-3-ylmethyl)benzamide	506.2
578	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(trifluoromethyl)benzyl]benzamide	573.1
579	*N*-[2-(4-chlorophenyl)ethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	554
580	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-pyrrolidin-1-ylethyl)benzamide	512.2
581	*N*-(3-methylpyridin-2-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	506.1
582	*N*-[3-(dimethylamino)propyl]-*N*-methyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	514.2
583	*N*-1,3-benzodioxol-5-yl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	535.1
584	*N*-(2-morpholin-4-ylethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	528.2

Compound Number	Name	MS (*m/z*, M + 1)
585	1'-{4-[(4-pyrimidin-2-ylpiperazin-1-yl)carbonyl]benzyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	562.2
586	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-1,3-thiazol-2-ylbenzamide	498.1
587	*N*-(6-methoxypyridin-3-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	522.1
588	*N*-(3,5-dichlorobenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	574.1
589	*N*-1-naphthyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	541.2
590	1'-(4-{[4-(1,3-benzodioxol-5-ylmethyl)piperazin-1-yl]carbonyl}benzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	618.2
591	*N*-(4,6-dimethylpyridin-2-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	520.2
592	4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pyrimidin-4-ylbenzamide	493.1
593	*N*-(5-methyl-1,3-thiazol-2-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	512.2
594	*N*-(2-cyano-6-fluorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	534.1
595	*N*-(4-methylbenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	519.1
596	*N*-[3-(1*H*-imidazol-1-yl)propyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	523.1

Compound Number	Name	MS (*m/z*, M + 1)
597	*N*-(4-morpholin-4-ylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	576.2
598	1'-{4-[(4-methylpiperazin-1-yl)carbonyl]benzyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	498.2
599	*N*-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide	539.2

EXAMPLE 13

Synthesis of 1'-(3-hydroxypropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



A suspension of 1'-[3-(benzyloxy)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (6.27 g, 14.5 mmol) and 10% Pd/C (0.5 g) in MeOH (150 mL) was hydrogenated under the normal pressure of hydrogen overnight and filtered through a pad of celite. The filtrate was concentrated *in vacuo* to dryness. The residue was crystallized from ether to yield the title compound (4.82 g) as a white solid in 98% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.38-6.93 (m, 4H), 6.49 (s, 1H), 6.10 (s, 1H), 4.87 (m, 1H), 4.63 (m, 1H), 4.01-3.81 (m, 2H), 3.62 (t, 2H), 2.89 (br, 1H), 1.99-1.91(m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.8, 156.0, 148.9, 142.4, 142.1, 132.3, 129.0, 128.9, 124.1, 123.9, 119.0, 108.6, 103.0, 101.5, 93.6, 80.4, 58.3, 37.8, 29.8; MS (ES+) *m/z* 340.2 (M + 1).

EXAMPLE 14

Synthesis of 3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propanal

To a solution of 1'-(3-hydroxypropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (4.82 g, 14.2 mmol) in dichloromethane (150 mL) was added 1,1,1-triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1*H*)-one (7.00 g, 16.7 mmol) at 0 °C. The resulting mixture was stirred at 0 °C for 4 h, diluted with ethyl acetate, washed sequentially with 10% $Na_2S_2O_3$ solution, saturated $NaHCO_3$ and brine, dried over anhydrous Na_2SO_4 and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to flash chromatography and the product was recrystallized from ethyl acetate/hexanes to afford the title compound (3.86 g) in 80% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 9.83 (s, 1H), 7.33-6.92 (m, 4H), 6.48 (s, 1H), 6.08 (s, 1H), 4.86 (m, 1H), 4.61(m, 1H), 4.15-3.98 (m, 2H), 2.97-2.84 (m, 2H); MS (ES+, *m/z*) 338.1 (M + 1).

EXAMPLE 15

Synthesis of 1'-{3-[(cyclopropylmethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a solution of 3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propanal (0.07 g, 0.20 mmol) in THF (5.00 mL) was added (aminomethyl)cyclopropane (0.30 mmol) and MP-triacetoxyborohydride (0.26 g, 0.60 mmol). After overnight shaking, the polymer-bound 4-phenyloxybenzaldehyde (0.25 g, 0.18 mmol) was added. After another overnight shaking, the mixture was diluted with ether (10.0 mL) and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was recrystallized to give the title compound (0.05 g) in 62% yield as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.32-6.92 (m, 4H), 6.49 (s, 1H), 6.12 (s, 1H), 5.83 (m, 2H), 4.86 (m, 1H), 4.64 (m, 1H), 3.97-3.77(m, 2H), 2.87-2.80 (m, 2H), 2.66-2.56 (m, 2H), 1.02-0.94 (m, 1H), 0.56-0.47 (m, 2H), 0.25-0.18 (m, 2H); MS (ES+) *m/z* 393.3

(M + 1).

EXAMPLE 15.1

The compounds listed in the following table were prepared using the similar procedure as described in EXAMPLE 15. As previously noted, the compound numbers listed below do not correspond to the compound numbers provided in the general Reaction Schemes above.

Compound Number	Name	MS (*m/z*, M + 1)
600	1'-{3-[(4-fluorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	447.0
601	1'-{3-[(4-chlorophenyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	449.2
602	1'-[3-(pentylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	409.1
603	1'-{3-[(2-ethoxyethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	411.1
604	1'-{3-[(3-methoxypropyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	411.1
605	1'-{3-[(3-methylbutyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	409.6
606	1'-{3-[(3-ethoxypropyl)amino]propyl}spiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	425.6
607	1'-{3-[(2,2-dimethylpropyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	409.1
608	3-{[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]amino}propanenitrile	392.4
609	1'-{3-[(2,2,2-trifluoroethyl)amino]propyl}spiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	421.1
610	1'-[3-(cyclopropylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	379.1
611	1'-[3-(cyclobutylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	393.1
612	1'-{3-[(2-cyclopropylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	407.1
613	1'-[3-(isobutylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	395.1

Compound Number	Name	MS (*m/z*, M + 1)
614	1'-[3-(hexylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	423.5
615	1'-[3-(heptylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	437.1
616	1'-[3-(isopropylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	381.7
617	1'-{3-[(tetrahydrofuran-2-ylmethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	423.1
618	1'-[3-(benzylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	429.1
619	1'-{3-[(2-phenylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	443.1
620	1'-[3-(dibenzylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	519.4
621	1'-[3-(propylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	381.1
622	1'-(3-{[2-(3-fluorophenyl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	461.5
623	1'-{3-[(3-phenylpropyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	457.5
624	1'-{3-[(2,2-diphenylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	519.4
625	1'-(3-{[2-(4-methylphenyl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	457.5
626	1'-(3-{[2-(3-chlorophenyl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	477.3
627	1'-{3-[(2-pyridin-4-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	444.2
628	1'-{3-[(pyridin-4-ylmethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	430.5
629	1'-(3-{[2-(4-fluorophenyl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	461.0

Compound Number	Name	MS (*m/z*, M + 1)
630	1'-{3-[(pyridin-2-ylmethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	430.2
631	1'-(3-{[(1*R*)-1-cyclohexylethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	449.1
632	1'-{3-[(2-furylmethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	419.4
633	1'-{3-[(4-chlorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	463.1
634	1'-{3-[(4-methoxybenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	459.4
635	1'-{3-[(3-isopropoxypropyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	439.1
636	1'-(3-{[2-(2-fluorophenyl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	461.0
637	1'-{3-[(3,3-dimethylbutyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	423.5
638	1'-{3-[(cyclohexylmethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	435.0
639	1'-(3-{[(1*S*)-1-cyclohexylethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	436.5
640	1'-{3-[(2-piperidin-1-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	449.1
641	1'-{3-[(2-pyrrolidin-1-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	450.5
642	1'-{3-[(2-morpholin-4-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	452.4
643	1'-[3-(cyclohexylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	421.5
644	1'-[3-(cyclopentylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	407.3
645	1'-{3-[(2-chlorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	463.2

Compound Number	Name	MS (*m/z*, M + 1)
646	1'-(3-pyrrolidin-1-ylpropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	393.5
647	1'-[3-(dibutylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	451.5
648	1'-(3-piperidin-1-ylpropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	407.0
649	1'-[3-(dipropylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	423.2
650	1'-(3-{[2-(dimethylamino)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	410.5
651	1'-(3-{[2-(diethylamino)ethyl](methyl)amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	452.1
652	1'-(3-{[2-(diisopropylamino)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	466.1
653	1'-[3-(diisopropylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	423.0
654	1'-[3-(methylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	353.0
655	1'-[3-(ethylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	367.0
656	1'-{3-[bis(2-methoxyethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	455.0
657	1'-{3-[(2-fluorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	446.9
658	1'-{3-[(3,5-difluorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	464.9
659	1'-(3-{[3-(dimethylamino)propyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	424.0
660	1'-[3-(diethylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	394.9
661	1'-[3-(octylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	451.5

Compound Number	Name	MS (*m/z*, M + 1)
662	1'-{3-[(1-methylbutyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	409.0
663	1'-{3-[butyl(methyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	409.0
664	1'-{3-[(2-isopropoxyethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	425.0
665	1'-{3-[(2,4-difluorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	464.9
666	1'-{3-[(2-methylbenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	443.0
667	1'-{3-[(3-fluorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	446.9
668	1'-{3-[(2,6-difluorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	464.9
669	1'-{3-[(1,2-dimethylpropyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	409.0
670	1'-(3-{[2-(1-methylpyrrolidin-2-yl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	450.4
671	1'-{3-[(2-pyridin-3-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	444.5
672	1'-{3-[(1-methyl-2-phenylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	457.0
673	1'-(3-{[2-(2-chlorophenyl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	476.9
674	1'-{3-[(2-cyclohexylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	448.1
675	1'-{3-[(2-pyridin-2-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	444.2
676	1'-{3-[(2-biphenyl-4-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	519.6
677	1'-{3-[(3-morpholin-4-ylpropyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	466.2

Compound Number	Name	MS (*m/z*, M + 1)
678	1'-(3-{[(5-methyl-2-furyl)methyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	433.3
679	1'-{3-[(3-methylbenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one	443.4

EXAMPLE 16

Synthesis of 1'-(3-aminopropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



To a solution of 2-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1*H*-isoindole-1,3(2*H*)-dione (3.20 g, 6.80 mmol) in ethanol (70.0 mL) was added hydrazine monohydrate (1.87 g, 37.0 mmol). The mixture was stirred at ambient temperature for 4 h. The solvent was removed under reduced pressure and the residue was re-dissolved in ethyl acetate. The solution was washed with sodium bicarbonate and brine solution, dried over $MgSO_4$ and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was crystallized from hexane to yield the title compound (2.50 g) in 75% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.31-7.24 (m, 1H), 7.16-7.14 (m, 1H), 7.06-7.01 (m, 1H), 6.94-6.91 (m, 1H), 6.48 (s, 1H), 6.10 (s, 1H), 5.82 (m, 2H), 4.90-4.87 (m, 1H), 4.61 (d, 1H), 3.98-3.71 (m, 2H), 2.77-2.73 (m, 2H), 1.97 (br, 2H), 1.84-1.81 (m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.8, 155.9, 148.9, 142.3, 142.2, 132.4, 129.0, 124.0, 123.4, 119.4, 108.7, 103.0, 101.5, 93.6, 80.5, 58.2, 38.8, 37.5, 30.6; MS (ES+) *m/z* 339.3 (M + 1).

EXAMPLE 17

Synthesis of 3-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]thiophene-2-carboxamide



To a solution of 1'-(3-aminopropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.05 g, 0.13 mmol) in dichloromethane (4.00 mL) was added triethylamine (0.03 g, 0.26 mmol) and 3-chlorothiophene-2-carbonyl chloride (0.02 g, 0.12 mmol) at 0 °C. The mixture was stirred for 2 h, washed with 15% HCl solution and water. The organic layer was dried over $MgSO_4$ and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was dissolved in ethyl acetate, and the product was precipitated with the addition of hexane. The white solid was collected by filtration and dried *in vacuo* to yield the title compound (0.04 g) in 67% yield: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.60 (t, 1H), 7.41 (d, 1H), 7.33-7.27- (m, 1H), 7.17 (d, 1H), 7.08-7.03- (m, 1H), 6.93 (t, 1H), 6.49 (s, 1H), 6.11 (s, 1H), 5.95 (m, 2H), 4.90 (d, 1H), 4.65 (d, 1H), 3.96-3.79- (m, 2H), 3.53-3.36- (m, 2H), 2.04-1.93- (m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.3, 160.6, 156.0, 149.0, 142.4, 141.8, 132.5, 129.4, 129.2, 129.1, 124.2, 123.7, 123.6, 119.2, 108.5, 102.9, 101.6, 93.7, 80.5, 58.3, 37.4, 36.5, 27.3; MS (ES+) *m/z* 483 (M + 1).

EXAMPLE 17.1

The compounds listed in the following table were synthesized using the similar procedure as described in EXAMPLE 17. As previously noted, the compound numbers listed below do not correspond to the compound numbers provided in the general Reaction Schemes above.

Compound Number	Name	MS (m/z, M + 1)
680	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]cyclopropanecarboxamide	407.3
681	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]cyclobutanecarboxamide	421.3

Compound Number	Name	MS (m/z, M + 1)
682	2-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]nicotinamide	478.3
683	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]cyclopentanecarboxamide	435.3
684	2,2-dimethyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]propanamide	423.3
685	2-(4-methoxyphenyl)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide	487.3
686	4-tert-butyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	499.2
687	3,3-dimethyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]butanamide	437.3
688	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]biphenyl-4-carboxamide	519.3
689	3-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1-benzofuran-2-carboxamide	497.2
690	2-(benzyloxy)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide	487.3
691	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-furamide	433.3
692	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1,3-benzodioxole-5-carboxamide	487.3
693	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]quinoline-2-carboxamide	494.3
694	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-phenylacetamide	457.4
695	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]piperidine-1-carboxamide	450.3

Compound Number	Name	MS (m/z, M + 1)
696	2-methoxy-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide	411.2
697	4-(dimethylamino)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	486.3
698	4-ethoxy-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	487.3
699	2-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]butanamide	423.3
700	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-phenoxyacetamide	473.2
701	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]quinoxaline-2-carboxamide	495.0
702	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]cyclohexanecarboxamide	449.3
703	4-fluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	461.3
704	2-ethyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]butanamide	437.3
705	2-(4-fluorophenyl)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide	475.3
706	6-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]nicotinamide	478.0
707	2-fluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	461.4
708	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-phenylcyclopropanecarboxamide	482.8
709	4-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	457.7

Compound Number	Name	MS (m/z, M + 1)
710	1-(4-fluorophenyl)-5-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1*H*-pyrazole-4-carboxamide	541.2
711	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1-benzofuran-5-carboxamide	483.1
712	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2,1,3-benzoxadiazole-5-carboxamide	485.4
713	2,4-dichloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	513.1
714	1-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1*H*-1,2,3-benzotriazole-5-carboxamide	498.4
715	5-fluoro-2-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	475.1
716	2-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]isonicotinamide	478.1
717	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2,3-dihydro-1,4-benzodioxine-6-carboxamide	501.1
718	5-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]isoxazole-3-carboxamide	448.2
719	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1-benzofuran-2-carboxamide	483.1
720	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1-benzothiophene-2-carboxamide	499.1
721	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2,3-dihydro-1,4-benzodioxine-2-carboxamide	501.1
722	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-4-(1*H*-pyrazol-1-yl)benzamide	509.1

Compound Number	Name	MS (m/z, M + 1)
723	1,3-dimethyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1*H*-pyrazole-5-carboxamide	461.2
724	4-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-3,4-dihydro-2*H*-1,4-benzoxazine-7-carboxamide	514.2
725	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]quinoxaline-6-carboxamide	495.2
726	*N*-[3-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'H)-yl)propyl]-2,3-dihydro-1-benzofuran-2-carboxamide	485.2
727	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2,3-dihydro-1-benzothiophene-5-carboxamide	499.1
728	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-(trifluoromethoxy)benzamide	526.9
729	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]pentanamide	445.2
730	*N*-[3-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]heptanamide	473.2
731	3-cyclopentyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]propanamide	463.0
732	9-oxo-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-9*H*-fluorene-4-carboxamide	545.0
733	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-4-(trifluoromethyl)benzamide	511.4
734	2,5-difluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	479.3
735	2,5-dimethyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-3-furamide	461.4

Compound Number	Name	MS (m/z, M + 1)
736	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-4-phenoxybutanamide	523.3
737	4-fluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-(trifluoromethyl)benzamide	529.0
738	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-(2-thienyl)acetamide	463.3
739	2-chloro-5-fluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	495.0
740	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-naphthamide	493.0
741	2-(4-chlorophenoxy)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide	507.3
742	2,4-dimethoxy-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	503.4
743	2-nitro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	488.4
744	2-(4-chlorophenyl)-3-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]butanamide	533.4
745	4-amino-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	458.4
746	3,4-dimethoxy-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	503.3
747	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-5*H*-dibenzo[*b,f*]azepine-5-carboxamide	558.4
748	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]adamantane-1-carboxamide	501.5

Compound Number	Name	MS (m/z, M + 1)
749	2-[(2-isopropyl-5-methylcyclohexyl)oxy]-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide	535.5
750	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-3,5-bis(trifluoromethyl)benzamide	579.3
751	2-(2,5-dimethoxyphenyl)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide	517.4
752	2-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	477.3
753	3-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	477.3
754	4-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	477.3
755	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]hexanamide	437.4
756	2,6-difluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	479.4
757	2-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide	415.3
758	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2,5-bis(trifluoromethyl)benzamide	579.4
759	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]pyrrolidine-1-carboxamide	436.4
760	2-bromo-2,2-difluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide	497.3
761	2,3,5-trifluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	497.3
762	5-fluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-(trifluoromethyl)benzamide	529.4

Compound Number	Name	MS (m/z, M + 1)
763	5-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-(trifluoromethyl)benzamide	545.3
764	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]thiophene-2-carboxamide	449.3
765	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]morpholine-4-carboxamide	452.4
766	2-(1-naphthyl)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide	507.4
767	2-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]propanamide	409.4
768	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-*N*-propionylpropanamide	451.4
769	*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-4-pentylbenzamide	513.3
770	4,7,7-trimethyl-3-oxo-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-oxabicyclo[2.2.1]heptane-1-carboxamide	519.4
771	2-bromo-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	523.2
772	3-cyano-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	468.3
773	4-cyano-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide	468.3

EXAMPLE 18

Synthesis of 1'-(2-aminoethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one



337

To a suspension of 2-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-1*H*-isoindole-1,3(2*H*)-dione (20.0 g, 44.0 mmol) in methanol (400 mL) was added hydrazine (8.00 mL). The mixture was stirred at ambient temperature for 48 h and filtered. The filtrate was concentrated *in vacuo* to dryness. The residue was subjected to flash chromatography eluting with ethyl acetate/methanol/ammonia (10/1/0.2) to afford the crude product which was recrystalized from ethyl acetate to yield the title compound (8.0 g) in 56% yield as a white solid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.34 (m, 1H), 7.17 (dd, 1H), 7.06 (dd, 1H), 6.95 (d, 1H), 6.51 (s, 1H), 6.18 (s, 1H), 5.89-5.82- (ABq, 2H), 4.93 (d, 1H), 4.66 (d, 1H), 3.95-3.74- (m, 2H), 3.06 (t, 2H), 1.59-1.35- (br, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 178.0, 155.9, 148.8, 142.3, 132.4, 128.9, 124.0, 123.4, 119.5, 108.6, 103.1, 101.5, 93.6, 80.5, 58.2, 43.4, 39.8; MS (ES+) *m/z* 325 (M + 1), 308 (M - 16).

EXAMPLE 19

Synthesis of 1-(4-fluorophenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea



To a mixture of 1'-(2-aminoethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one (0.15 mmol) and triethylamine (0.01 mmol) in anhydrous dichloromethane was added 1-fluoro-4-isocyanatobenzene (0.14 mmol) at ambient temperature. The mixture was stirred for 16 h, diluted with of dichloromethane (5.00 mL), washed with 10% HCl solution and brine, dried over Na_2SO_4 and filtered. The filtrate was concentrated *in vacuo* to dryness to give the title compound (0.05 g) in 82% yield: ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.54 (s, 1H), 7.35-7.25 (m, 3H), 7.19 (d, 1H), 7.11 (d, 1H), 7.06-6.91 (m, 3H), 6.67 (s, 1H), 6.52 (t, 1H), 5.94-5.84 (ABq, 2H), 4.74 (d, 1H), 4.61 (d, 1H), 3.91-3.69 (m, 2H), 3.49-3.34 (m, 2H); MS (ES+) *m/z* 462 (M + 1), 484 (M + 23).

EXAMPLE 19.1

The compounds listed in the following table were synthesized using the similar

procedure as described in EXAMPLE 19. As previously noted, the compound numbers listed below do not correspond to the compound numbers provided in the general Reaction Schemes above.

Compound Number	Name	MS (*m/z*, M + 1)
774	1-benzyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea	458
775	1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-(4-phenoxyphenyl)urea	536
776	1-butyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea	424
777	1-cyclohexyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea	450
778	1-ethyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea	396
779	1-isopropyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea	410
780	1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-propylurea	410
781	1-*tert*-butyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea	424
782	1-cyclopentyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea	436
783	1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-pentylurea	438
784	1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-phenylurea	444
785	1-(2-furylmethyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea	448
786	1-hexyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea	452
787	ethyl *N*-({[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]amino}carbonyl)glycinate	454
788	1-(3-methylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea	458

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 340

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 340

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHIER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

WHAT IS CLAIMED IS

1. A compound of formula (I):



(I)

wherein:

j and k are each independently 0, 1, 2 or 3;

Q is $-C(R^{1a})H-$, $-O-$, $-S(O)_m-$ (where m is 1 or 2), $-CF_2-$, $-C(O)O-$, $-C(O)N(R^5)-$ or $-N(R^5)C(O)-$;

R^{1a} is $-OR^5$;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

or R^1 is aralkyl substituted by $-C(O)N(R^6)R^7$ where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is hydrogen, alkyl, haloalkyl, $-R^9-CN$, $-R^9-OR^5$, $-R^9-N(R^4)R^5$, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, or heteroarylalkyl;

or R^6 and R^7, together with the nitrogen to which they are attached, may form a heterocyclyl or heteroaryl;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^6 and R^7 may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, $-R^8-CN$, $-R^8-OR^5$, heterocyclyl and heteroaryl;

or R^1 is aralkyl optionally substituted by one or more substituents selected from the group consisting of $-R^8-OR^5$, $-C(O)OR^5$, halo, haloalkyl, alkyl, nitro, cyano, aryl, aralkyl, heterocyclyl and heteroaryl;

or R^1 is $-R^9-N(R^{10})R^{11}$, $-R^9-N(R^{12})C(O)R^{11}$ or $-R^9-N(R^{10})C(O)N(R^{10})R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9-OC(O)R^5$, $-R^9-C(O)OR^5$, $-R^9-C(O)N(R^4)R^5$, $-R^9-C(O)R^5$, $-R^9-N(R^4)R^5$, $-R^9-OR^5$, or $-R^9-CN$; and

R^{12} is hydrogen, alkyl, aryl, aralkyl or $-C(O)R^5$;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, $-R^8-CN$, $-R^8-OR^5$, $-R^8-C(O)R^5$, heterocyclyl and heteroaryl;

or R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroarylalkyl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-OR^5$, $-R^8-C(O)OR^5$, $-R^8-N(R^4)R^5$, $-R^8-C(O)N(R^4)R^5$, $-R^8-N(R^5)C(O)R^4$, $-R^8-S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8-CN$, or $-R^8-NO_2$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy,

cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$S(O)_mR^4$, -R^8-$S(O)_nN(R^4)R^5$, -R^8-$C(O)R^4$, -R^8-$C(O)OR^4$, -R^8-$C(O)N(R^4)R^5$, -$N(R^5)C(O)R^4$, and -$N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$N{=}C(R^4)R^5$, -$S(O)_mR^4$, -$OS(O)_2CF_3$, -R^8-$C(O)R^4$, -$C(S)R^4$, -$C(R^4)_2C(O)R^5$, -R^8-$C(O)OR^4$, -$C(S)OR^4$, -R^8-$C(O)N(R^4)R^5$, -$C(S)N(R^4)R^5$, -$N(R^5)C(O)R^4$, -$N(R^5)C(S)R^4$, -$N(R^5)C(O)OR^4$, -$N(R^5)C(S)OR^4$, -$N(R^5)C(O)N(R^4)R^5$, -$N(R^5)C(S)N(R^4)R^5$, -$N(R^5)S(O)_nR^4$, -$N(R^5)S(O)_nN(R^4)R^5$, -R^8-$S(O)_nN(R^4)R^5$, -$N(R^5)C(=NR^5)N(R^4)R^5$,

and -$N(R^5)C(N{=}C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain;

or a stereoisomer, enantiomer, or tautomer thereof or mixtures thereof;

or a pharmaceutically acceptable salt thereof.

2. The compound of Claim 1 wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, $-R^8$-CN, $-R^8$-NO_2, $-R^8$-OR^5, $-R^8$-$N(R^4)R^5$, $-S(O)_mR^4$, $-R^8$-$S(O)_nN(R^4)R^5$, $-R^8$-$C(O)R^4$, $-R^8$-$C(O)OR^4$, $-R^8$-$C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, $-R^8$-CN, $-R^8$-NO_2, $-R^8$-OR^5, $-R^8$-$N(R^4)R^5$, $-N{=}C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8$-$C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8$-$C(O)OR^4$, $-C(S)OR^4$, $-R^8$-$C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8$-$S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$,

and $-N(R^5)C(N{=}C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as

defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

3. The compound of Claim 2 wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen, halo or alkyl;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S($

$O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3c} are as defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

4. The compound of Claim 3 wherein:

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkylalkyl, $-R^8-OR^5$, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen, halo or alkyl;

R^{3a} and R^{3d} are both hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene

chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

5. The compound of Claim 4 which is:

1'-(2-cyclopropylethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

ethyl (2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetate;

spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(benzyloxy)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4',7'-dichloro-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-bromospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-bromo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one;

5'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

ethyl (4'-bromo-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)acetate;

ethyl (4'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-b:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)acetate;

ethyl (5'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-b:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)acetate;

7'-fluorospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetic acid;

N-(4-chlorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3-*f*luorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-butyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

1'-(2-oxo-2-piperidin-1-ylethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

N-butyl-*N*-methyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-phenylacetamide;

N-(4-*f*luorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3-*f*luorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3-chlorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-*f*luorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-ethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(4-ethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3-methylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2,3-dimethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3,5-dimethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-pentylacetamide;

2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-propylacetamide;

N-isopropyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3-methylbutyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-isobutyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-hexyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-cyclohexyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-cyclopentyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-heptyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-chlorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2,6-dimethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-methoxyphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-[(5-methyl-2-*f*uryl)methyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-ethyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-methyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-*f*luorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-[2-(3-methoxyphenyl)ethyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-ethoxyethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(4-methoxybenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2,4-dimethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3-isopropoxypropyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-*f*urylmethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(cyclohexylmethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3-*f*luoro-2-methylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(4-methoxyphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-cyclobutyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2,5-difluorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-benzyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(cyclopropylmethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-butyl-*N*-ethyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-octyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3,3-dimethylbutyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(4-chloro-2-methylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3-methoxyphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-*f*luoro-4-methylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3,4-dimethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3-chlorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3-methoxybenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3,4-difluorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3-methylbenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-methoxybenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(4-isopropylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2,3-difluorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-(tetrahydrofuran-2-ylmethyl)acetamide;

N-[2-(4-methylphenyl)ethyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-[2-(3-chlorophenyl)ethyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-

yl)acetamide;

N-(4-cyanophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2,3-dihydro-1*H*-inden-1-yl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-methoxyethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-[2-(4-methoxyphenyl)ethyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-cyanoethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2,4-dichlorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3,5-difluorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2,4-difluorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-methylbenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3,4-difluorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2,5-difluorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N,N*-dipropylacetamide;

N,N-dibutyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2,6-difluorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-[2-(methylthio)phenyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-isopropylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(4-bromophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(4-chlorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2,4-dichlorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3,5-dichlorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N,N-diethyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-methyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-phenylacetamide;

N-(4-hydroxybutyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-allyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-*f*luoro-5-methylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-

yl)acetamide;

N-(1,3-benzodioxol-5-ylmethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-cyclopropyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-cyclopropylethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3,4-dichlorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2,3-dichlorobenzyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2,5-dimethylphenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(3,4-dichlorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N,N-dimethyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-methyl-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-(2-phenylethyl)acetamide;

2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-(2-phenylpropyl)acetamide;

N-[(1*R*)-1-cyclohexylethyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-[(1S)-1-cyclohexylethyl]-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-(2-piperidin-1-ylethyl)acetamide;

2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-[3-(trifluoromethyl)phenyl]acetamide;

N-(3-cyanophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

1'-(2-morpholin-4-yl-2-oxoethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)-*N*-(2-phenylethyl)acetamide;

N-(4-bromo-2-chlorophenyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-biphenyl-4-ylethyl)-2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)acetamide;

1'-prop-2-yn-1-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2-ethoxyethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(2E)-pent-2-en-1-yl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-hex-5-en-1-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(cyclobutylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pent-2-yn-1-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(5-chloropentyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(4-fluorobutyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(5-methylhexyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(3*Z*)-4-methylhex-3-en-1-yl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2-bromoethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

5-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)pentanenitrile;

1'-[2-(2-methoxyethoxy)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(cyclopropylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(4,4,4-trifluorobutyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

diethyl [2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]phosphonate;

1'-[(2,2,3,3-tetrafluorocyclobutyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(benzyloxy)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-allylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2-ethylbutyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(4-methylpentyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-methylbut-2-en-1-yl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pent-4-en-1-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)butanenitrile;

1'-[(2-methylcyclopropyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-cyclopropylpropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-hexylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(2-cyclopropyl-6-hydroxypyrimidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(2-methylcyclopropyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-cyclopropylpropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-butylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'[4-(trifluoromethoxy)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-Propylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-bromo-1'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

tert-butyl 4-[(4'-bromo-2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]piperidine-1-carboxylate;

ethyl 5-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)pentanoate;

ethyl 4-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)butanoate;

1'-(3-chloropropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(cyclohexylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(methylsulfonyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-hydroxypropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propanal; or

2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indole]-4'-carbonitrile.

6. The compound of Claim 3 wherein:

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkylalkyl, $-R^8-OR^5$, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen, halo or alkyl;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

7. The compound of Claim 6 which is:

5,6-difluoro-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

5-fluoro-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

5-bromo-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

5-chloro-6-fluoro-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

6-methoxy-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

6-chloro-5-fluoro-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-pentyl-5-(trifluoromethyl)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

5,6-dichloro-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

6-bromo-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

5-bromospiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

ethyl (4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2' *H*)-yl)acetate;

methyl (6-chloro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetate;

ethyl (5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetate;

2-(4'-bromo-5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide;

2-(6-chloro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide;

2-(5,6-difluoro-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide;

2-(5-bromo-2'-oxospiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide;

1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

6-anilino-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

6-morpholin-4-yl-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

6-amino-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-pentyl-6-phenoxyspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-pentyl-6-phenoxyspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-pentyl-6-pyridin-4-ylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

6-(methylsulfonyl)-1'-pentylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-pentyl-6-(phenylsulfonyl)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-pentyl-5-phenoxyspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

5-hydroxyspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

2'-oxo-1',2'-dihydrospiro[1-benzofuran-3,3'-indol]-5-yl trifluoromethanesulfonate;

1'-pentyl-5-pyridin-3-ylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-pentyl-5-pyrimidin-5-ylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-pentyl-5-pyridin-4-ylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

2'-oxo-1'-pentyl-1',2'-dihydrospiro[1-benzofuran-3,3'-indole]-5-carbonitrile;

N-(2-fluorophenyl)-2-(2'-oxo-5-pyridin-3-ylspiro[1-benzofuran-3,3'-indol]-1'(2'*H*)-yl)acetamide;

tert-butyl 3-(2'-oxo-1',2'-dihydrospiro[1-benzofuran-3,3'-indol]-5-yl)piperidine-1-carboxylate; or

5-methoxy-1'-methylspiro [1-benzofuran-3,3'-indol]-2'(1'*H*)-one.

8. The compound of Claim 3 wherein:

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkylalkyl, $-R^8-OR^5$, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen, halo or alkyl;

R^{3a} and R^{3d} are both hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused optionally substituted heterocyclyl ring or a fused optionally substituted cycloalkyl ring;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl,

heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

9. The compound of Claim 8 which is:

5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one;

ethyl (4'-bromo-6,6-dimethyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)acetate;

ethyl (2'-oxo-6,7-dihydro-5*H*-spiro[indeno[5,6-*b*]furan-3,3'-indol]-1'(2'*H*)-yl)acetate;

ethyl (2-oxo-5',6',7',8'-tetrahydrospiro[indole-3,3'-naphtho[2,3-*b*]furan]-1(2*H*)-yl)acetate;

6,7-dihydrospiro[benzo[1,2-b:4,5-*b*']difuran-3,3'-indol]-2'(1'*H*)-one;

5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one;

ethyl (2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetate;

(2'-oxo-6,7-dihydro-5*H*-spiro[indeno[5,6-*b*]furan-3,3'-indol]-1'(2'*H*)-yl)acetic acid;

(4'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-b:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid;

(4'-bromo-2'-oxo-5,6-dihydrospiro[benzo[1,2-b:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid;

(5'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid;

(2-oxo-5',6',7',8'-tetrahydrospiro[indole-3,3'-naphtho[2,3-*b*]furan]-1(2*H*)-yl)acetic acid;

(2'-oxo-5,6-dihydrospiro[benzo[1,2-b:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)acetic acid;

N-(2-fluorophenyl)-2-(2'-oxo-6,7-dihydro-5*H*-spiro[indeno[5,6-*b*]furan-3,3'-indol]-1'(2'*H*)-yl)acetamide;

N-(2-fluorophenyl)-2-(2-oxo-5',6',7',8'-tetrahydrospiro[indole-3,3'-naphtho[2,3-*b*]furan]-1(2*H*)-yl)acetamide;

2-(4'-bromo-6,6-dimethyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide;

2-(4'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-

fluorophenyl)acetamide;

2-(5'-chloro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide;

2-(4'-fluoro-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide;

2-(4'-bromo-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide;

N-(2-fluorophenyl)-2-(2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)acetamide; or

2-(4'-fluoro-7'-methyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide.

10. The compound of Claim 3 wherein:

R^1 is aryl, heteroaryl or heterocyclyl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen; and

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring.

11. The compound of Claim 2 wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is hydrogen, alkyl, $-R^8-C(O)OR^5$ or $-R^8-C(O)N(R^4)R^5$;

R^{2a} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$,

and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is

independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a} is optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$, $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from hydrogen or halo;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl and R^{3a} and R^{3d} are as defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

12. The compound of Claim 11 wherein:

j is 0 and k is 1 or j is 1 and k is 0;

Q is -O-;

R^1 is hydrogen or alkyl;

R^{2a} is selected from the group consisting of alkyl, haloalkenyl, aryl, aralkyl, aralkenyl, heterocyclyl, heteroaryl, $-R^8-C(O)N(R^4)R^5$, and $-R^8-N(R^4)R^5$;

wherein each of the aryl, aralkyl, aralkenyl, heterocyclyl and heteroaryl groups for R^{2a} is optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$, $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from hydrogen or halo;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring or a fused optionally substituted tetrahydrofuranyl ring, and R^{3a} and R^{3d} are as defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

13. The compound of Claim 12 which is:

4'-[6-(dimethylamino)pyridin-3-yl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3,5-dimethoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(4-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3,5-dichlorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-[4-(dimethylamino)phenyl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-(3,4,5-trimethoxyphenyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)benzonitrile;

4'-dibenzo[*b*,*d*]furan-4-yl-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(1-benzyl-1*H*-pyrazol-4-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(2-methoxypyrimidin-5-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(2,4-dimethoxypyrimidin-5-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)benzamide;

4'-{4-[(dimethylamino)methyl]phenyl}-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(1-benzofuran-2-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(6-methoxypyridin-3-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

N,*N*-dimethyl-4-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)benzamide;

4'-dibenzo[*b*,*d*]thien-4-yl-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

3-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)benzonitrile;

1'-pentyl-4'-pyridin-3-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3-fluoro-4-methoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-[2-(trifluoromethoxy)phenyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-[3,5-bis(trifluoromethyl)phenyl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-[4-(trifluoromethyl)pyridin-3-yl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(2-fluoro-5-methoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(4-ethoxy-3-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(1-benzothien-2-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-isobutyl-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-[4-(trifluoromethoxy)phenyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(5-fluoro-2-methoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(1,3-benzodioxol-5-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-phenylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-[2-(trifluoromethyl)phenyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(4-chlorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(2,3-dihydro-1,4-benzodioxin-6-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-quinolin-3-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3,5-difluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-isoquinolin-4-yl-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(6-methoxypyridin-2-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(1*H*-indol-5-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

N-[2-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)phenyl]acetamide;

4'-(4-fluoro-2-methylphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-quinolin-6-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

N-[4-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)phenyl]methanesulfonamide;

4'-(5-chloro-2-methoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-[3-(trifluoromethoxy)phenyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-(4-phenoxyphenyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(2,4-dimethoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3-furyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3,4-dimethoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

N-[4-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)phenyl]acetamide;

1'-pentyl-4'-[(*E*)-2-phenylvinyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(4-methoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(6-fluoropyridin-3-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3-chloro-4-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3-chlorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(1-benzothien-3-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-(2-phenoxyphenyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(4-isopropoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-[(*E*)-2-(4-fluorophenyl)vinyl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(6-fluoropyridin-2-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-[1-(phenylsulfonyl)-1*H*-indol-3-yl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3-acetylphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(2-furyl)-1'-pentylspiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(4-methylphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(1-methyl-1*H*-pyrrol-3-yl)-1'-pentylspiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(2,5-difluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(2-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(2-chlorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(2,4-difluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(4-morpholin-4-ylphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

tert-butyl 5-methoxy-3-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)-1*H*-indole-1-carboxylate;

1'-pentyl-4'-pyrimidin-5-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

tert-butyl 4-[2-(2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-4'-yl)phenyl]piperazine-1-carboxylate;

4'-(2-methoxypyridin-3-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(5-methoxypyridin-3-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(4-butoxy-3-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-pyridin-4-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-phenoxathiin-4-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-[(1*Z*)-3-chloroprop-1-en-1-yl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-pentyl-4'-(3-thienyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(2,3-dimethoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(4-butylphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3-fluoro-5-methoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-[3-fluoro-4-(pentyloxy)phenyl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(2-butoxy-5-fluorophenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3-butoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(4-butoxyphenyl)-1'-pentylspiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(4-isobutoxyphenyl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-{2-chloro-4-[(3,5-dimethoxybenzyl)oxy]phenyl}-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-[4-(benzyloxy)-3-chlorophenyl]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(1-methyl-1*H*-indol-5-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(4-methoxypyridin-3-yl)-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-[(6-methoxypyridin-3-yl)amino]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3-furyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-dibenzo[*b*,*d*]furan-4-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-pyrimidin-5-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-methyl-4'-pyrimidin-5-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3-furyl)-1'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(6-fluoropyridin-3-yl)-1'-methylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2-cyclopropylethyl)-4'-quinolin-3-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-[(6-methoxypyridin-3-yl)amino]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-[(3,5-difluorophenyl)amino]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-[(4,6-dimethylpyridin-2-yl)amino]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-[(4-methyl-1,3-thiazol-2-yl)amino]-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-bromo-1'-pentylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-morpholino-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

4'-(4-methylpiperazin-1-yl)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

1'-pentyl-4'-(pyrimidin-4-ylamino)-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

1'-pentyl-4'-(pyridin-3-ylamino)-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

4'-(4-chloro-2-(trifluoromethyl)phenylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

1'-pentyl-4'-(pyrimidin-2-ylamino)-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

4'-(benzo[*d*][1,3]dioxol-5-ylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

4'-(3-fluorophenylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

4'-(naphthalen-2-ylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

4'-(2-methoxyphenylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

4'-(4-methylthiazol-2-ylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

4'-(4,6-dimethylpyridin-2-ylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

4'-(3,5-difluorophenylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

4'-(6-methoxypyridin-3-ylamino)-1'-pentyl-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

1'-pentyl-7*H*-spiro[furo[3,4-*f*][1,3]benzodioxole-5,3'-indol]-2'(1'*H*)-one;

2'-oxo-1'-pentyl-*N*-pyridin-2-yl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indole]-4'-carboxamide;

N-(3-methoxyphenyl)-2'-oxo-1'-pentyl-1',2'-dihydrospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indole]-4'-carboxamide;

2-(5,6-difluoro-2'-oxo-4'-pyrimidin-5-ylspiro[1-benzofuran-3,3'-indol]-1'(2' *H*)-yl)-*N*-(2-fluorophenyl)acetamide;

2-(6,6-dimethyl-2'-oxo-4'-pyrimidin-5-yl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-1'(2'*H*)-yl)-*N*-(2-fluorophenyl)acetamide; or

N-(2-fluorophenyl)-2-(2'-oxo-4'-pyrimidin-5-yl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-

indol]-1'(2'*H*)-yl)acetamide.

14. The compound of Claim 1 wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is aralkyl substituted by $-C(O)N(R^6)R^7$ where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is hydrogen, alkyl, haloalkyl, $-R^9-CN$, $-R^9-OR^5$, $-R^9-N(R^4)R^5$, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, or heteroarylalkyl;

or R^6 and R^7, together with the nitrogen to which they are attached, may form a heterocyclyl or heteroaryl;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^6 and R^7 may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, $-R^8-CN$, $-R^8-OR^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$S(O)_mR^4$, -R^8-$S(O)_nN(R^4)R^5$, -R^8-$C(O)R^4$, -R^8-$C(O)OR^4$, -R^8-$C(O)N(R^4)R^5$, -$N(R^5)C(O)R^4$, and -$N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$N{=}C(R^4)R^5$, -$S(O)_mR^4$, -$OS(O)_2CF_3$, -R^8-$C(O)R^4$, -$C(S)R^4$, -$C(R^4)_2C(O)R^5$, -R^8-$C(O)OR^4$, -$C(S)OR^4$, -R^8-$C(O)N(R^4)R^5$, -$C(S)N(R^4)R^5$, -$N(R^5)C(O)R^4$, -$N(R^5)C(S)R^4$, -$N(R^5)C(O)OR^4$, -$N(R^5)C(S)OR^4$, -$N(R^5)C(O)N(R^4)R^5$, -$N(R^5)C(S)N(R^4)R^5$, -$N(R^5)S(O)_nR^4$, -$N(R^5)S(O)_nN(R^4)R^5$, -R^8-$S(O)_nN(R^4)R^5$, -$N(R^5)C{=}NR^5)N(R^4)R^5$,

and -$N(R^5)C(N{=}C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as

defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

15. The compound of Claim 14 wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is aralkyl substituted by $-C(O)N(R^6)R^7$ where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is hydrogen, alkyl, haloalkyl, $-R^9-CN$, $-R^9-OR^5$, $-R^9-N(R^4)R^5$, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, or heteroarylalkyl;

or R^6 and R^7, together with the nitrogen to which they are attached, may form a heterocyclyl or heteroaryl;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^6 and R^7 may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, $-R^8-CN$, $-R^8-OR^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl,

heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

16. The compound of Claim 15 wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is aralkyl substituted by $-C(O)N(R^6)R^7$ where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is hydrogen, alkyl, haloalkyl, $-R^9-CN$, $-R^9-OR^5$ or $-R^9-N(R^4)R^5$;

or R^6 and R^7, together with the nitrogen to which they are attached, may form a heterocyclyl or heteroaryl;

wherein each aryl, aralkyl, heterocyclyl and heteroaryl groups for R^6 and R^7 is optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, $-R^8-CN$, $-R^8-OR^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

17. The compound of Claim 16 which is:

N-(3-methylbutyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N,N-diisopropyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-butyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pentylbenzamide;

N-hexyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-propylbenzamide;

N-isopropyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-heptyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

1'-[2-(piperidin-1-ylcarbonyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

N-isobutyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-methoxypropyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-ethoxypropyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-hexyl-*N*-methyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-isopropoxypropyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-ethoxyethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-methyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-ethyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2,2,2-trifluoroethyl)benzamide;

N-[2-(diethylamino)ethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3,3-dimethylbutyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-ethylbutyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-methoxyethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-cyanoethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

1'-{2-[(4-pyrimidin-2-ylpiperazin-1-yl)carbonyl]benzyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2-{[4-(1,3-benzodioxol-5-ylmethyl)piperazin-1-yl]carbonyl}benzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[2-(morpholin-4-ylcarbonyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

N-[3-(dimethylamino)propyl]-*N*-methyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-methylbutyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N,N-diisopropyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-butyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pentylbenzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-propylbenzamide;

N-isopropyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

1'-[3-(piperidin-1-ylcarbonyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

N-isobutyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-hexyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-heptyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-methoxypropyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[2-(diethylamino)ethyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-methyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-ethyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-ethoxypropyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-hexyl-*N*-methyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-isopropoxypropyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-ethoxyethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-ethylbutyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-methoxyethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

1'-[3-(morpholin-4-ylcarbonyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(4-pyrimidin-2-ylpiperazin-1-yl)carbonyl]benzyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(4-methylpiperazin-1-yl)carbonyl]benzyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

N-(3,3-dimethylbutyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-cyanoethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-butyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

1'-[4-(piperidin-1-ylcarbonyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

N,N-diisopropyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pentylbenzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-propylbenzamide;

N-isopropyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-isobutyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-hexyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-heptyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-methoxypropyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[2-(diethylamino)ethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-methyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-ethyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-ethoxypropyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-methylpentyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2,2,2-trifluoroethyl)benzamide;

N-hexyl-*N*-methyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-isopropoxypropyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-ethoxyethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-ethylbutyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-methoxyethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-

yl)methyl]benzamide;

N-(2-cyanoethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide; or

N-[3-(dimethylamino)propyl]-*N*-methyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide.

18. The compound of Claim 15 wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is aralkyl substituted by $-C(O)N(R^6)R^7$ where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, or heteroarylalkyl;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^6 and R^7 may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, $-R^8-CN$, $-R^8-OR^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

19. The compound of Claim 18 which is:

N-[2-(4-chlorophenyl)ethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-phenylbenzamide;

N-(3-fluorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-chlorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-cyclohexyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-cyclopentyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-fluorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-fluorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-chlorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-fluorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-ethylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-ethylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-methylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,3-dimethylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3,5-dimethylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-chlorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,6-dimethylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-methoxyphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[2-(4-methylphenyl)ethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(tetrahydrofuran-2-ylmethyl)benzamide;

N,N-dibenzyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(3-phenylpropyl)benzamide;

N-[2-(3-chlorophenyl)ethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[2-(4-fluorophenyl)ethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-fluorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-methoxybenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(cyclopropylmethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(cyclohexylmethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-furylmethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,4-dimethylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-cyanophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3,5-dichlorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-fluoro-2-methylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-methoxyphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(5-chloro-2-methylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[3-(trifluoromethyl)phenyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(pyridin-4-ylmethyl)benzamide;

N-cyclobutyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,2-diphenylethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-pyrrolidin-1-ylethyl)benzamide;

N-[(1-ethylpyrrolidin-2-yl)methyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-piperidin-1-ylethyl)benzamide;

N-(2-morpholin-4-ylethyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[(1*S*)-1-cyclohexylethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-fluoro-5-methylphenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,4-difluorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(4-propylphenyl)benzamide;

N-(3,3-diphenylpropyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,5-difluorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-thienylmethyl)benzamide;

N-[4-chloro-2-(trifluoromethyl)phenyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[2-(4-methoxyphenyl)ethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3,5-dichlorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-chlorobenzyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(2-thienyl)ethyl]benzamide;

N-(2,3-dihydro-1*H*-inden-1-yl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[4-(trifluoromethyl)benzyl]benzamide;

N-[4-fluoro-2-(trifluoromethyl)phenyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(pyridin-3-ylmethyl)benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(trifluoromethyl)benzyl]benzamide;

N-(3-methylpyridin-2-yl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-

yl)methyl]benzamide;

N-(1-benzylpiperidin-4-yl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(trifluoromethyl)phenyl]benzamide;

N-[(1*R*)-1-cyclohexylethyl]-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(6-methoxypyridin-3-yl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-1,3-thiazol-2-ylbenzamide;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-1,3,4-thiadiazol-2-ylbenzamide;

N-(4,6-dimethylpyridin-2-yl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,3-dihydro-1*H*-inden-5-yl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-2-adamantyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-1-adamantyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-1-naphthyl-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3,5-difluorophenyl)-2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[2-(3-chlorophenyl)ethyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-phenylbenzamide;

N-(3-fluorophenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-chlorobenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-fluorobenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-chlorophenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-fluorophenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-ethylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-ethylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-methylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,3-dimethylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3,5-dimethylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-cyclohexyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-cyclopentyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-methoxybenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-methoxyphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-cyclopropyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,4-dimethylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(tetrahydrofuran-2-ylmethyl)benzamide;

N,N-dibenzyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[3-(trifluoromethyl)phenyl]benzamide;

N-(4-methoxybenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3,5-dichlorophenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pyridin-3-ylbenzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(pyridin-4-ylmethyl)benzamide;

N-(2-furylmethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-fluoro-2-methylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(cyclopropylmethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-methoxyphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-cyclobutyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[2-(4-fluorophenyl)ethyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(cyclohexylmethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[2-(4-methylphenyl)ethyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-benzyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(1-benzylpiperidin-4-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[2-(4-methoxyphenyl)ethyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-

yl)methyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-piperidin-1-ylethyl)benzamide;

N-(1-cyclohexylethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-thienylmethyl)benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(4-propylphenyl)benzamide;

N-(2,4-difluorobenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3,5-difluorophenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,3-dihydro-1*H*-inden-5-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[4-(trifluoromethyl)benzyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(2-thienyl)ethyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(pyridin-3-ylmethyl)benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(trifluoromethyl)benzyl]benzamide;

N-[2-(4-chlorophenyl)ethyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-pyrrolidin-1-ylethyl)benzamide;

N-(3-methylpyridin-2-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-1,3-benzodioxol-5-yl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-morpholin-4-ylethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-1,3-thiazol-2-ylbenzamide;

N-(6-methoxypyridin-3-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3,5-dichlorobenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-1-naphthyl-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4,6-dimethylpyridin-2-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pyrimidin-4-ylbenzamide;

N-(5-methyl-1,3-thiazol-2-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-methylbenzyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[3-(1*H*-imidazol-1-yl)propyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-1,3,4-thiadiazol-2-ylbenzamide;

N-(4-morpholin-4-ylphenyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[(1-ethylpyrrolidin-2-yl)methyl]-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,2-diphenylethyl)-3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-fluorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-fluorobenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-chlorobenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-chlorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-fluorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-ethylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-ethylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-methylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3,5-dimethylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,3-dimethylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-cyclohexyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-cyclopentyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-methoxybenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,6-dimethylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-methoxyphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-

yl)methyl]benzamide;

N-cyclopropyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-phenylbenzamide;

N-(2,4-dimethylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(tetrahydrofuran-2-ylmethyl)benzamide;

N,N-dibenzyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[3-(trifluoromethyl)phenyl]benzamide;

N-(4-methoxybenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3,5-dichlorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pyridin-3-ylbenzamide;

N-(4-cyanophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(pyridin-4-ylmethyl)benzamide;

N-[2-(3-chlorophenyl)ethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-furylmethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3-fluoro-2-methylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(cyclopropylmethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-methoxyphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-cyclobutyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,2-diphenylethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[2-(4-fluorophenyl)ethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(cyclohexylmethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-fluoro-4-methylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[2-(4-methylphenyl)ethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-benzyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-methoxyethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

1'-[4-(morpholin-4-ylcarbonyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

N-(1-benzylpiperidin-4-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[2-(4-methoxyphenyl)ethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(trifluoromethyl)phenyl]benzamide;

N-[4-chloro-2-(trifluoromethyl)phenyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[4-fluoro-2-(trifluoromethyl)phenyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[(1*S*)-1-cyclohexylethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[(1*R*)-1-cyclohexylethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,4-difluorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,3-dihydro-1*H*-inden-1-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*[1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-thienylmethyl)benzamide;

N-[(1-ethylpyrrolidin-2-yl)methyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(4-propylphenyl)benzamide;

N-(2,5-difluorobenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,3-dihydro-1*H*-inden-5-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2,5-difluorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[4-(trifluoromethyl)benzyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(2-thienyl)ethyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(pyridin-3-ylmethyl)benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-[2-(trifluoromethyl)benzyl]benzamide;

N-[2-(4-chlorophenyl)ethyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-(2-pyrrolidin-1-ylethyl)benzamide;

N-(3-methylpyridin-2-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-

yl)methyl]benzamide;

N-1,3-benzodioxol-5-yl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-morpholin-4-ylethyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

1'-{4-[(4-pyrimidin-2-ylpiperazin-1-yl)carbonyl]benzyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-1,3-thiazol-2-ylbenzamide;

N-(6-methoxypyridin-3-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(3,5-dichlorobenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-1-naphthyl-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

1'-(4-{[4-(1,3-benzodioxol-5-ylmethyl)piperazin-1-yl]carbonyl}benzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

N-(4,6-dimethylpyridin-2-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-*N*-pyrimidin-4-ylbenzamide;

N-(5-methyl-1,3-thiazol-2-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(2-cyano-6-fluorophenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-methylbenzyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-[3-(1*H*-imidazol-1-yl)propyl]-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide;

N-(4-morpholin-4-ylphenyl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-

yl)methyl]benzamide;

1'-{4-[(4-methylpiperazin-1-yl)carbonyl]benzyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one; or

N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)-4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzamide.

20. The compound of Claim 1 wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is aralkyl optionally substituted by one or more substituents selected from the group consisting of $-R^8-OR^5$, $-C(O)OR^5$, halo, haloalkyl, alkyl, nitro, cyano, aryl, aralkyl, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$, $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^{8}$-CN, $-R^{8}$-NO_2, $-R^{8}$-OR^{5}, $-R^{8}$-$N(R^{4})R^{5}$, $-N{=}C(R^{4})R^{5}$, $-S(O)_mR^{4}$, $-OS(O)_2CF_3$, $-R^{8}$-$C(O)R^{4}$, $-C(S)R^{4}$, $-C(R^{4})_2C(O)R^{5}$, $-R^{8}$-$C(O)OR^{4}$, $-C(S)OR^{4}$, $-R^{8}$-$C(O)N(R^{4})R^{5}$, $-C(S)N(R^{4})R^{5}$, $-N(R^{5})C(O)R^{4}$, $-N(R^{5})C(S)R^{4}$, $-N(R^{5})C(O)OR^{4}$, $-N(R^{5})C(S)OR^{4}$, $-N(R^{5})C(O)N(R^{4})R^{5}$, $-N(R^{5})C(S)N(R^{4})R^{5}$, $-N(R^{5})S(O)_nR^{4}$, $-N(R^{5})S(O)_nN(R^{4})R^{5}$, $-R^{8}$-$S(O)_nN(R^{4})R^{5}$, $-N(R^{5})C(=NR^{5})N(R^{4})R^{5}$, and $-N(R^{5})C(N{=}C(R^{4})R^{5})N(R^{4})R^{5}$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl,

heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

21. The compound of Claim 20 wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is aralkyl optionally substituted by one or more substituents selected from the group consisting of $-R^8-OR^5$, $-C(O)OR^5$, halo, haloalkyl, alkyl, nitro, cyano, aryl, aralkyl, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen, alkyl or halo;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as

defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

22. The compound of Claim 21 wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is aralkyl optionally substituted by one or more substituents selected from the group consisting of $-R^8-OR^5$, $-C(O)OR^5$, halo, haloalkyl, alkyl, nitro, cyano, aryl, aralkyl, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen, alkyl or halo;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, halo, and $-R^8-OR^5$

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, dioxolyl, tetrahydrofuranyl, and heteroaryl, and R^{3a} and R^{3d} are each hydrogen;

each R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

23. The compound of Claim 22 which is:

methyl 2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoate;

methyl 3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoate;

methyl 4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoate;

1'-(diphenylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(diphenylmethyl)-5'-methylspiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

2-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoic acid;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoic acid;

1'-(4-fluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-benzylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3,5-difluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-nitrobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-fluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

3-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzonitrile;

1'-[4-(1*H*-pyrazol-1-yl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(biphenyl-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]biphenyl-2-carbonitrile;

1'-(biphenyl-2-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(diphenylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(4-fluoro-3-methylbenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(5-fluoro-2-methylbenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2,5-difluorobenzyl)spiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[4-(1*H*-pyrrol-1-yl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(1*H*-pyrrol-1-yl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(4-chlorophenoxy)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[2-fluoro-3-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[2-fluoro-6-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-fluoro-4-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[4-fluoro-3-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[2-fluoro-5-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[4-fluoro-2-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[5-fluoro-2-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[2-fluoro-4-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2,3-difluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(1-bromo-2-naphthyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(1-naphthylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-fluoro-5-(trifluoromethyl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2,4-difluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2,6-difluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-methoxybenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[4-(1*H*-1,2,4-triazol-1-yl)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[4,4-bis(4-fluorophenyl)butyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2-chloro-4-fluorobenzyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[2-(trifluoromethoxy)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(trifluoromethoxy)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'[4-(trifluoromethoxy)benzyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2,3-difluorobenzyl)-5,6-dihydrospyro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]2'(1'*H*)-one;

1'-(4-methoxybenzyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]2'(1'*H*)-one;

4-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]benzoic acid;

5-bromo-1'-(diphenylmethyl)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-(diphenylmethyl)-6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-(4-methoxybenzyl)-6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-(diphenylmethyl)-5-hydroxyspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-(diphenylmethyl)-5-pyridin-3-ylspiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-(diphenylmethyl)-5,5-dimethyl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one;

2-methyl-1'-pentylspiro[furo[2,3-*f*][1,3]benzothiazole-7,3'-indol]-2'(1'*H*)-one;

1'-(diphenylmethyl)-6,7-dihydrospiro[benzo[1,2-*b*:4,5-*b*']difuran-3,3'-indol]-2'(1'*H*)-one; or

1'-(diphenylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one.

24. The compound of Claim 1 wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is $-R^9-N(R^{10})R^{11}$, $-R^9-N(R^{12})C(O)R^{11}$ or $-R^9-N(R^{10})C(O)N(R^{10})R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9-OC(O)R^5$, $-R^9-C(O)OR^5$, $-R^9-C(O)N(R^4)R^5$, $-R^9-C(O)R^5$, $-R^9-N(R^4)R^5$, $-R^9-OR^5$, or $-R^9-CN$; and

R^{12} is hydrogen, alkyl, aryl, aralkyl or $-C(O)R^5$;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, $-R^8-CN$, $-R^8-OR^5$, $-R^8-C(O)R^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$, $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a

fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^{8}$-CN, $-R^{8}$-NO_2, $-R^{8}$-OR^{5}, $-R^{8}$-$N(R^{4})R^{5}$, $-N{=}C(R^{4})R^{5}$, $-S(O)_mR^{4}$, $-OS(O)_2CF_3$, $-R^{8}$-$C(O)R^{4}$, $-C(S)R^{4}$, $-C(R^{4})_2C(O)R^{5}$, $-R^{8}$-$C(O)OR^{4}$, $-C(S)OR^{4}$, $-R^{8}$-$C(O)N(R^{4})R^{5}$, $-C(S)N(R^{4})R^{5}$, $-N(R^{5})C(O)R^{4}$, $-N(R^{5})C(S)R^{4}$, $-N(R^{5})C(O)OR^{4}$, $-N(R^{5})C(S)OR^{4}$, $-N(R^{5})C(O)N(R^{4})R^{5}$, $-N(R^{5})C(S)N(R^{4})R^{5}$, $-N(R^{5})S(O)_nR^{4}$, $-N(R^{5})S(O)_nN(R^{4})R^{5}$, $-R^{8}$-$S(O)_nN(R^{4})R^{5}$, $-N(R^{5})C(=NR^{5})N(R^{4})R^{5}$, and $-N(R^{5})C(N{=}C(R^{4})R^{5})N(R^{4})R^{5}$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

25. The compound of Claim 24 wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is $-R^9-N(R^{10})R^{11}$, $-R^9-N(R^{12})C(O)R^{11}$ or $-R^9-N(R^{10})C(O)N(R^{10})R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9-OC(O)R^5$, $-R^9-C(O)OR^5$, $-R^9-C(O)N(R^4)R^5$, $-R^9-C(O)R^5$, $-R^9-N(R^4)R^5$, $-R^9-OR^5$, or $-R^9-CN$; and

R^{12} is hydrogen, alkyl, aryl, aralkyl or $-C(O)R^5$;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, $-R^8-CN$, $-R^8-OR^5$, $-R^8-C(O)R^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the

nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

26. The compound of Claim 25 wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is $-R^9-N(R^{10})R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl; and

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9-OC(O)R^5$, $-R^9-C(O)OR^5$, $-R^9-C(O)N(R^4)R^5$, $-R^9-C(O)R^5$, $-R^9-N(R^4)R^5$, $-R^9-OR^5$, or $-R^9-CN$;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, $-R^8-CN$, $-R^8-OR^5$, $-R^8-C(O)R^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene

chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

27. The compound of Claim 26 which is:

1'-[2-(diethylamino)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[2-(pyridin-2-ylamino)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[2-(dipyridin-2-ylamino)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(cyclopropylmethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(4-fluorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(4-chlorophenyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(pentylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-ethoxyethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(3-methoxypropyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(3-methylbutyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(3-ethoxypropyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2,2-dimethylpropyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

3-{[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]amino}propanenitrile;

1'-{3-[(2,2,2-trifluoroethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(cyclopropylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(cyclobutylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-cyclopropylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(isobutylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(hexylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(heptylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(isopropylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(tetrahydrofuran-2-ylmethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(benzylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-phenylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(dibenzylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(propylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[2-(3-fluorophenyl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(3-phenylpropyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2,2-diphenylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[2-(4-methylphenyl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[2-(3-chlorophenyl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-pyridin-4-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(pyridin-4-ylmethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[2-(4-fluorophenyl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(pyridin-2-ylmethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[(1*R*)-1-cyclohexylethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-furylmethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(4-chlorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(4-methoxybenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(3-isopropoxypropyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[2-(2-fluorophenyl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(3,3-dimethylbutyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(cyclohexylmethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[(1*S*)-1-cyclohexylethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-piperidin-1-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-pyrrolidin-1-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-

one;

1'-{3-[(2-morpholin-4-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(cyclohexylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(cyclopentylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-chlorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(dibutylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(dipropylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[2-(dimethylamino)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[2-(diethylamino)ethyl](methyl)amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[2-(diisopropylamino)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(diisopropylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(methylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(ethylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[bis(2-methoxyethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-fluorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(3,5-difluorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[3-(dimethylamino)propyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(diethylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(octylamino)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(1-methylbutyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[butyl(methyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-isopropoxyethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2,4-difluorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-methylbenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(3-fluorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2,6-difluorobenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(1,2-dimethylpropyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[2-(1-methylpyrrolidin-2-yl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-pyridin-3-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(1-methyl-2-phenylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[2-(2-chlorophenyl)ethyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-cyclohexylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-pyridin-2-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(2-biphenyl-4-ylethyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(3-morpholin-4-ylpropyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-{[(5-methyl-2-furyl)methyl]amino}propyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{3-[(3-methylbenzyl)amino]propyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-aminopropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one; or

1'-(2-aminoethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one.

28. The compound of Claim 25 wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is $-R^9-N(R^{12})C(O)R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9-OC(O)R^5$, $-R^9-C(O)OR^5$, $-R^9-C(O)N(R^4)R^5$, $-R^9-C(O)R^5$, $-R^9-N(R^4)R^5$, $-R^9-OR^5$, or $-R^9-CN$; and

R^{12} is hydrogen, alkyl, aryl, aralkyl or -C(O)R^5;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, -R^8-CN, -R^8-OR^5, -R^8-C(O)R^5, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

29. The compound of Claim 28 which is:

1'-(3-aminopropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

3-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]thiophene-2-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]cyclopropanecarboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]cyclobutanecarboxamide;

2-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]nicotinamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-

yl)propyl]cyclopentanecarboxamide;

2,2-dimethyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]propanamide;

2-(4-methoxyphenyl)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide;

4-*tert*-butyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

3,3-dimethyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]butanamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]biphenyl-4-carboxamide;

3-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1-benzofuran-2-carboxamide;

2-(benzyloxy)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-furamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1,3-benzodioxole-5-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]quinoline-2-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-phenylacetamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]piperidine-1-carboxamide;

2-methoxy-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide;

4-(dimethylamino)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

4-ethoxy-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

2-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]butanamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-phenoxyacetamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]quinoxaline-2-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]cyclohexanecarboxamide;

4-fluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

2-ethyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]butanamide;

2-(4-fluorophenyl)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide;

6-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]nicotinamide;

2-fluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-phenylcyclopropanecarboxamide;

4-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

1-(4-fluorophenyl)-5-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1*H*-pyrazole-4-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1-benzofuran-5-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2,1,3-benzoxadiazole-5-carboxamide;

2,4-dichloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

1-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1*H*-1,2,3-benzotriazole-5-carboxamide;

5-fluoro-2-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

2-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]isonicotinamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2,3-dihydro-1,4-benzodioxine-6-carboxamide;

5-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]isoxazole-3-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1-benzofuran-2-

carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1-benzothiophene-2-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2,3-dihydro-1,4-benzodioxine-2-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-4-(1*H*-pyrazol-1-yl)benzamide;

1,3-dimethyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1*H*-pyrazole-5-carboxamide;

4-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-3,4-dihydro-2*H*-1,4-benzoxazine-7-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]quinoxaline-6-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2,3-dihydro-1-benzofuran-2-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2,3-dihydro-1-benzothiophene-5-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-(trifluoromethoxy)benzamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]pentanamide;

N-[3-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]heptanamide;

3-cyclopentyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]propanamide;

9-oxo-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-9*H*-fluorene-4-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-4-(trifluoromethyl)benzamide;

2,5-difluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

2,5-dimethyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-3-

furamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-4-phenoxybutanamide;

4-fluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-(trifluoromethyl)benzamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-(2-thienyl)acetamide;

2-chloro-5-fluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-naphthamide;

2-(4-chlorophenoxy)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide;

2,4-dimethoxy-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

2-nitro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

2-(4-chlorophenyl)-3-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]butanamide;

4-amino-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

3,4-dimethoxy-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-5*H*-dibenzo[*b,f*]azepine-5-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]adamantane-1-carboxamide;

2-[(2-isopropyl-5-methylcyclohexyl)oxy]-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-3,5-bis(trifluoromethyl)benzamide;

2-(2,5-dimethoxyphenyl)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide;

2-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

3-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

4-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]hexanamide;

2,6-difluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

2-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2,5-bis(trifluoromethyl)benzamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]pyrrolidine-1-carboxamide;

2-bromo-2,2-difluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide;

2,3,5-trifluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

5-fluoro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-(trifluoromethyl)benzamide;

5-chloro-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-(trifluoromethyl)benzamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]thiophene-2-carboxamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]morpholine-4-carboxamide;

2-(1-naphthyl)-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]acetamide;

2-methyl-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]propanamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-*N*-propionylpropanamide;

N-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-4-pentylbenzamide;

4,7,7-trimethyl-3-oxo-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-2-oxabicyclo[2.2.1]heptane-1-carboxamide;

2-bromo-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

3-cyano-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide;

4-cyano-*N*-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]benzamide; or

N-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-2-(trifluoromethoxy)benzamide.

30. The compound of Claim 25 wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is $-R^9-N(R^{10})C(O)N(R^{10})R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl; and

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9-OC(O)R^5$, $-R^9-C(O)OR^5$, $-R^9-C(O)N(R^4)R^5$, $-R^9-C(O)R^5$, $-R^9-N(R^4)R^5$, $-R^9-OR^5$, or $-R^9-CN$;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, $-R^8-CN$, $-R^8-OR^5$, $-R^8-C(O)R^5$, heterocyclyl and heteroaryl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene

chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

31. The compound of Claim 30 which is:

1-(4-fluorophenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-benzyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-(4-phenoxyphenyl)urea;

1-butyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-cyclohexyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-ethyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-isopropyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-propylurea;

1-*tert*-butyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-cyclopentyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-pentylurea;

1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-phenylurea;

1-(2-furylmethyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-hexyl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

ethyl *N*-({[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]amino}carbonyl)glycinate;

1-(3-methylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(4-methylphenyl)-3-[2-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

ethyl *N*-({[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]amino}carbonyl)-beta-alaninate;

1-(4-cyanophenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

N-({[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]amino}carbonyl)benzamide;

1-[2-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-(2-phenylethyl)urea;

1-(4-methylbenzyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(2-methylbenzyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(4-ethylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(3-methoxyphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(2-fluoro-5-methylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(3-fluoro-4-methylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(4-chlorophenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

2-[({[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]amino}carbonyl)amino]ethyl 2-methylacrylate;

1-[2-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-(1,1,3,3-tetramethylbutyl)urea;

ethyl 4-[({[2-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]amino}carbonyl)amino]butanoate;

1-[4-(cyanomethyl)phenyl]-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(2,3-dihydro-1*H*-inden-5-yl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(3-acetylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(4-acetylphenyl)-3-[2-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(4-isopropylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(2-methoxybenzyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(4-methoxybenzyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(4-methoxy-2-methylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(4-chloro-2-methylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(3-chloro-4-methylphenyl)-3-[2-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(3-chloro-2-methylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(5-chloro-2-methylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(2-chlorobenzyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(1-naphthyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(2-naphthyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(3-chloro-2-fluorophenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-(5,6,7,8-tetrahydronaphthalen-1-yl)urea;

1-(4-*tert*-butylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(4-butylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[2-(4-ethylphenyl)ethyl]-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(2,3-dihydro-1,4-benzodioxin-6-yl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

methyl 4-[({[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]amino}carbonyl)amino]benzoate;

1-(2-ethoxybenzyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(3,4-dimethoxyphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(3,5-dimethoxyphenyl)-3-[2-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(3-chloro-4-methoxyphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[4-(difluoromethoxy)phenyl]-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[2-(difluoromethoxy)phenyl]-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-[3-(trifluoromethyl)phenyl]urea;

1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-[2-(trifluoromethyl)phenyl]urea;

1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-[4-(trifluoromethyl)phenyl]urea;

1-(3,4-dichlorophenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(2,3-dichlorophenyl)-3-[2-(2'-oxospiro[furo[2,3-f][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(3,5-dichlorophenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

ethyl 4-[({[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]amino}carbonyl)amino]benzoate;

ethyl 2-[({[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]amino}carbonyl)amino]benzoate;

1-[2-(1,3-benzodioxol-5-yl)ethyl]-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

methyl 2-methyl-3-[({[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]amino}carbonyl)amino]benzoate;

1-(4-butoxyphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(2-methoxy-4-nitrophenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-biphenyl-2-yl-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[4-methyl-3-(trifluoromethyl)phenyl]-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-

1'(2'*H*)-yl)ethyl]urea;

1-(2,4-dichlorobenzyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-[2-(trifluoromethoxy)phenyl]urea;

1-[4-fluoro-2-(trifluoromethyl)phenyl]-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(5-*tert*-butyl-2-methoxyphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[2-(3,5-dimethoxyphenyl)ethyl]-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(9*H*-fluoren-2-yl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(9*H*-fluoren-9-yl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-(3,4,5-trimethoxyphenyl)urea;

1-(diphenylmethyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-(2-phenoxyphenyl)urea;

1-(2-biphenyl-4-ylethyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-3-(3,4,5-trimethoxybenzyl)urea;

1-(2-nitrophenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(1,3-benzodioxol-5-yl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-[4-(dimethylamino)phenyl]-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(2-fluorobenzyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(4-fluoro-3-methylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(3-fluorobenzyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(cyclohexylmethyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

1-(2-methylphenyl)-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea;

or

1-[4-(6-methyl-1,3-benzothiazol-2-yl)phenyl]-3-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]urea.

32. The compound of Claim 1 wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroarylalkyl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-OR^5$, $-R^8-C(O)OR^5$, $-R^8-N(R^4)R^5$, $-R^8-C(O)N(R^4)R^5$, $-R^8-N(R^5)C(O)R^4$, $-R^8-S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8-CN$, or $-R^8-NO_2$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl,

heteroaryl,

heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$S(O)_mR^4$, -R^8-$S(O)_nN(R^4)R^5$, -R^8-$C(O)R^4$, -R^8-$C(O)OR^4$, -R^8-$C(O)N(R^4)R^5$, -$N(R^5)C(O)R^4$, and -$N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$N{=}C(R^4)R^5$, -$S(O)_mR^4$, -$OS(O)_2CF_3$, -R^8-$C(O)R^4$, -$C(S)R^4$, -$C(R^4)_2C(O)R^5$, -R^8-$C(O)OR^4$, -$C(S)OR^4$, -R^8-$C(O)N(R^4)R^5$, -$C(S)N(R^4)R^5$, -$N(R^5)C(O)R^4$, -$N(R^5)C(S)R^4$, -$N(R^5)C(O)OR^4$, -$N(R^5)C(S)OR^4$, -$N(R^5)C(O)N(R^4)R^5$, -$N(R^5)C(S)N(R^4)R^5$, -$N(R^5)S(O)_nR^4$, -$N(R^5)S(O)_nN(R^4)R^5$, -R^8-$S(O)_nN(R^4)R^5$, -$N(R^5)C(=NR^5)N(R^4)R^5$,

and -$N(R^5)C(N{=}C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a

fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

33. The compound of Claim 32 wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroarylalkyl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-OR^5$, $-R^8-C(O)OR^5$, $-R^8-N(R^4)R^5$, $-R^8-C(O)N(R^4)R^5$, $-R^8-N(R^5)C(O)R^4$, $-R^8-S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8-CN$, or $-R^8-NO_2$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen, halo, alkyl or $-R^8-OR^5$;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from hydrogen, halo, alkyl or $-R^8-OR^5$;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are each hydrogen;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

34. The compound of Claim 33 wherein:

j is 0 and k is 1;

Q is -O-;

R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroarylalkyl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-OR^5$, $-R^8-C(O)OR^5$, $-R^8-N(R^4)R^5$, $-R^8-C(O)N(R^4)R^5$, $-R^8-N(R^5)C(O)R^4$, $-R^8-S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8-CN$, or $-R^8-NO_2$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from hydrogen, halo, alkyl or $-R^8-OR^5$;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from hydrogen, halo, alkyl or $-R^8-OR^5$;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from dioxolyl or tetrahydrofuranyl, and R^{3a} and R^{3d} are each hydrogen;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl; and

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

35. The compound of Claim 34 which is:

2-[3-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)propyl]-1*H*-isoindole-1,3(2*H*)-dione;

2-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-1*H*-isoindole-1,3(2*H*)-dione;

4'-bromo-1'-(pyridin-2-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]-2'(1'*H*)-one;

5'-fluoro-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

5'-methyl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

5-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

4'-Methoxy-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

7'-fluoro-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{[1-(2,6-difluorobenzyl)-1*H*-1,2,3-triazol-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{[6-(trifluoromethyl)pyridin-3-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(6-chloropyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(tetrahydro-2*H*-pyran-2-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(tetrahydrofuran-2-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2,3-dihydro-1,4-benzodioxin-2-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(7-methoxy-2-oxo-2*H*-1,4-benzoxazin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(5-methyl-2-phenyl-2*H*-1,2,3-triazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3,4-dihydro-2*H*-1,5-benzodioxepin-7-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(1*H*-1,2,3-triazol-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

ethyl 1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-1*H*-1,2,3-triazole-5-carboxylate;

ethyl 1-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]-1*H*-1,2,3-triazole-4-carboxylate;

1'-(1,3-thiazol-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(5-chloro-1-benzothien-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(pyridin-2-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(pyridin-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(pyridin-3-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[2-(1*H*-pyrrol-1-yl)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{[4-chloro-2-(trifluoromethyl)quinolin-6-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(2-methyl-1,3-thiazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(1,3-benzothiazol-2-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2,1,3-benzothiadiazol-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2,1,3-benzothiadiazol-5-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(1-methyl-1*H*-1,2,3-benzotriazol-5-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[3-(1*H*-pyrrol-1-yl)propyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(3-methyl-5-phenylisoxazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(7-methoxy-2-oxo-2*H*-chromen-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[2-(2,5,5-trimethyl-1,3-dioxan-2-yl)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(6-fluoro-4*H*-1,3-benzodioxin-8-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(5-phenyl-1,3-oxazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(1,3-benzodioxol-5-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(piperidin-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(1-methylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[(1-ethylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[(1-cyclohexylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-{[1-cyclopropylmethyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[(1-cyclopentylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-{[1-(pyridine-3-ylmethyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-{[1-(3-methylbutyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-{[1-(1-ethylpropyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[(1-cyclobutylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[(1-isopropylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-{[1-(pyridin-2-ylmethyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-{[1-(2-thienylmethyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-({1-[3-(methylthio)propyl]piperidin-4-yl}methyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-{[1-(tetrahydro-2*H*-pyran-4-yl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-{[1-(3,3-dimethylbutyl)piperidin-4-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

7'-fluoro-1'-[(1-isopropylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[(6-methylpyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[(6-methoxypyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(6-chloropyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{[6-(dimethylamino)pyridin-3-yl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[(6-morpholin-4-ylpyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[(6-pyrrolidin-1-ylpyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[(5-methylisoxazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'indol]-2'(1'*H*)-one;

1'-(tetrahydro-2*H*-pyran-4-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2,1,3-benzoxadiazol-5-ylmethyl)spiro[furo2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(1-methyl-1*H*-benzotriazol-6-yl)methyl]spiro[furo2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

tert-butyl 4-[(2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-yl)methyl]piperidine 1-carboxylate;

1'-(pyridin-2-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b*']difuran-3,3'-indol]2'(1'*H*)-one hydrochloride;

4'-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-bromo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one;

1'-[(3,5-dimethylisoxazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2-furylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(1,2,4-oxadiazol-3-ylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{[5-(3-chlorophenyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(2-isopropyl-1,3-oxazol-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(1-methyl-1*H*-benzimidazol-2-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(2-oxo-1,3-benzothiazol-3(2*H*)-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(5-chloro-2-thienyl)methyl]-5'-fluorospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(5-chloro-2-furyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(4-hydroxy-1,2,2,6,6-pentamethylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{[5-(2-chlorophenyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(5-methyl-2-furyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(5-bromo-2-furyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(5-chloro-2-thienyl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{[3-hydroxy-5-(trifluoromethyl)-2-thienyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-((5-(2-(trifluoromethyl)phenyl)furan-2-yl)methyl)-6*H*-spiro[benzofuro[6,5-*d*][1,3]dioxole-7,3'-indolin]-2'-one;

1'[(2-chloro-1,3-thiazol-5-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{[5-(trifluoromethyl)-2-thienyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-{[3-methoxy-5-(trifluoromethyl)-2-thienyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-methyl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

5'-methyl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-({5-[4-(trifluoromethyl)phenyl]-1,2,4-oxadiazol-3-yl}methyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(2-thienylmethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

5-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]thiophene-2-carbonitrile;

5-[(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]-2-furonitrile;

1'-{[5-(methylsulfonyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(6-oxo-1,6-dihydropyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(1-methyl-6-oxo-1, 6-dihydropyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(5-chloro-1,3,4-thiadiazol-2-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(1-pyridin-2-ylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-[(1-phenyl-2-ylpiperidin-4-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(pyridin-2-ylmethyl)-6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-(2-piperidin-1-ylethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

tert-butyl 4-[2-(2'-oxospiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)ethyl]piperidine-1-carboxylate;

1'-(2-piperidin-4-ylethyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[2-(1-cyclopentylpiperidin-4-yl)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[2-(1-isopropylpiperidin-4-yl)ethyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[2-(1-cyclobutylpiperidin-4-yl)ethyl]spiro[furo[2,3 *f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-{2-[1-(tetrahydro-2*H*-pyran-4-yl)piperidin-4-yl]ethyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-(3-pyrrolidin-1-ylpropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

1'-(3-piperidin-1-ylpropyl)spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one; and

1'-[(5-fluoro-1*H*-benzimidazol-2-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one.

36. The compound of Claim 32 wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroarylalkyl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-OR^5$, $-R^8-C(O)OR^5$, $-R^8-N(R^4)R^5$, $-R^8-C(O)N(R^4)R^5$, $-R^8-N(R^5)C(O)R^4$, $-R^8-S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8-CN$, or $-R^8-NO_2$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$N=C(R^4)R^5$, -$S(O)_mR^4$, -$OS(O)_2CF_3$, -R^8-$C(O)R^4$, -$C(S)R^4$, -$C(R^4)_2C(O)R^5$, -R^8-$C(O)OR^4$, -$C(S)OR^4$, -R^8-$C(O)N(R^4)R^5$, -$C(S)N(R^4)R^5$, -$N(R^5)C(O)R^4$, -$N(R^5)C(S)R^4$, -$N(R^5)C(O)OR^4$, -$N(R^5)C(S)OR^4$, -$N(R^5)C(O)N(R^4)R^5$, -$N(R^5)C(S)N(R^4)R^5$, -$N(R^5)S(O)_nR^4$, -$N(R^5)S(O)_nN(R^4)R^5$, -R^8-$S(O)_nN(R^4)R^5$, -$N(R^5)C(=NR^5)N(R^4)R^5$, and -$N(R^5)C(=N\text{-}CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$S(O)_mR^4$, -R^8-$S(O)_nN(R^4)R^5$, -R^8-$C(O)R^4$, -R^8-$C(O)OR^4$, -R^8-$C(O)N(R^4)R^5$, -$N(R^5)C(O)R^4$, and -$N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

R^{3a} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-$N(R^4)R^5$, -$N=C(R^4)R^5$, -$S(O)_mR^4$, -$OS(O)_2CF_3$, -R^8-$C(O)R^4$, -$C(S)R^4$, -$C(R^4)_2C(O)R^5$, -R^8-$C(O)OR^4$, -$C(S)OR^4$, -R^8-$C(O)N(R^4)R^5$, -$C(S)N(R^4)R^5$, -$N(R^5)C(O)R^4$, -$N(R^5)C(S)R^4$, -$N(R^5)C(O)OR^4$, -$N(R^5)C(S)OR^4$, -$N(R^5)C(O)N(R^4)R^5$, -$N(R^5)C(S)N(R^4)R^5$, -$N(R^5)S(O)_nR^4$, -$N(R^5)S(O)_nN(R^4)R^5$, -R^8-$S(O)_nN(R^4)R^5$, -$N(R^5)C(=NR^5)N(R^4)R^5$, and -$N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from dioxolyl or tetrahydrofuranyl, and R^{3a} and R^{3d} are as defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

37. The compound of Claim 36 which is:

4'-(6-methoxypyridin-3-yl)-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-[6-(dimethylamino)pyridin-3-yl]-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-(3-furyl)-1'-(pyridin-2-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one;

1'-(pyridin-2-ylmethyl)-4'-pyrimidin-5-yl-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one;

4'-pyridin-3-yl-1'-(pyridin-2-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one;

4'-(3-furyl)-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one;

4'-quinolin-3-yl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one;

4'-pyrimidin-5-yl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one;

tert-butyl 4-[(2'-oxo-4'-pyrimidin-5-ylspiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-1'(2'*H*)-yl)methyl]piperidine-1-carboxylate;

tert-butyl 4-[(5,5-dimethyl-2'-oxo-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-1'(2'*H*)-

yl)methyl]piperidine-1-carboxylate;

5,5-dimethyl-1'-(piperidin-4-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one hydrochloride;

5,5-dimethyl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one;

5,5-dimethyl-1'-(pyridin-3-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one hydrochloride;

5,5-dimethyl-1'-(pyridin-2-ylmethyl)-5,6-dihydrospiro[benzo[1,2-*b*:5,4-*b'*]difuran-3,3'-indol]-2'(1'*H*)-one hydrochloride;

1'-[(6-methylpyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one hydrochloride; or

1'-[(6-methoxypyridin-3-yl)methyl]spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one.

38. The compound of Claim 32 wherein:

at least one of j and k is 1 and the other is 0 or 1;

Q is -O-;

R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroarylalkyl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-OR^5$, $-R^8-C(O)OR^5$, $-R^8-N(R^4)R^5$, $-R^8-C(O)N(R^4)R^5$, $-R^8-N(R^5)C(O)R^4$, $-R^8-S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8-CN$, or $-R^8-NO_2$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is

independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-S(O)_mR^4$, $-R^8-S(O)_nN(R^4)R^5$, $-R^8-C(O)R^4$, $-R^8-C(O)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$,

and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

39. The compound of Claim 38 which is:

5-bromo-1'-[(5-chloro-2-thienyl)methyl]spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

1'-(pyridin-3-ylmethyl)-6-(trifluoromethoxy)spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one hydrochloride;

6-(trifluoromethoxy)-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one;

2'-oxo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-1',2'-dihydrospiro[1-benzofuran-3,3'-indol]-5-yl trifluoromethanesulfonate;

5-pyridin-3-yl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one hydrochloride;

tert-butyl 3-(2'-oxo-1'-{[5-(trifluoromethyl)-2-furyl]methyl}-1',2'-dihydrospiro[1-benzofuran-3,3'-indol]-5-yl)piperidine-1-carboxylate; or

5-pyridin-4-yl-1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[1-benzofuran-3,3'-indol]-2'(1'*H*)-one hydrochloride.

40. The compound of Claim 2 wherein:

j is 0 and k is 1 or 2;

Q is -O-;

R^1 is hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene

chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

41. A compound of formula (I):



(I)

wherein:

j is 0 and k is 1 or 2;

Q is $-C(R^{1a})H-$, $-C(O)-$, $-CF_2-$, $-C(O)O-$ or $-N(R^5)C(O)-$;

R^{1a} is hydrogen or $-OR^5$;

R^1 is ethyl, *n*-propyl, 1-methylethyl, *n*-butyl, *n*-pentyl, 1,1-dimethylethyl, 3-methylhexyl, 2-methylhexyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

or R^1 is aralkyl substituted by $-C(O)N(R^6)R^7$ where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is hydrogen, alkyl, haloalkyl, $-R^9-CN$, $-R^9-OR^5$, $-R^9-N(R^4)R^5$, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, or heteroarylalkyl;

or R^6 and R^7, together with the nitrogen to which they are attached, may form a heterocyclyl or heteroaryl;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^6 and R^7 may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, $-R^8-CN$, $-R^8-OR^5$, heterocyclyl and heteroaryl;

or R^1 is aralkyl optionally substituted by one or more substituents selected from the group consisting of $-R^8-OR^5$, $-C(O)OR^5$, halo, haloalkyl, alkyl, nitro, cyano, aryl, aralkyl, heterocyclyl

and heteroaryl;

or R^1 is $-R^9-N(R^{10})R^{11}$, $-R^9-N(R^{12})C(O)R^{11}$ or $-R^9-N(R^{10})C(O)N(R^{10})R^{11}$ where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^9-OC(O)R^5$, $-R^9-C(O)OR^5$, $-R^9-C(O)N(R^4)R^5$, $-R^9-C(O)R^5$, $-R^9-N(R^4)R^5$, $-R^9-OR^5$, or $-R^9-CN$; and

R^{12} is hydrogen, alkyl, aryl, aralkyl or $-C(O)R^5$;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, nitro, $-R^8-CN$, $-R^8-OR^5$, $-R^8-C(O)R^5$, heterocyclyl and heteroaryl;

or R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroarylalkyl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-OR^5$, $-R^8-C(O)OR^5$, $-R^8-N(R^4)R^5$, $-R^8-C(O)N(R^4)R^5$, $-R^8-N(R^5)C(O)R^4$, $-R^8-S(O)_mR^4$ (where m is 0, 1 or 2), $-R^8-CN$, or $-R^8-NO_2$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(=N-CN)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl,

heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, $-R^8$-CN, $-R^8$-NO_2, $-R^8$-OR^5, $-R^8$-$N(R^4)R^5$, $-S(O)_mR^4$, $-R^8$-$S(O)_nN(R^4)R^5$, $-R^8$-$C(O)R^4$; $-R^8$-$C(O)OR^4$, $-R^8$-$C(O)N(R^4)R^5$, $-N(R^5)C(O)R^4$, and $-N(R^5)S(O)_nR^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, $-R^8$-CN, $-R^8$-NO_2, $-R^8$-OR^5, $-R^8$-$N(R^4)R^5$, $-N{=}C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8$-$C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8$-$C(O)OR^4$, $-C(S)OR^4$, $-R^8$-$C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8$-$S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$,

and $-N(R^5)C(N{=}C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain;

or a stereoisomer, enantiomer, or tautomer thereof, or mixtures thereof;

or a pharmaceutically acceptable salt thereof.

42. The compound of Claim 41 wherein:

j is 0 and k is 1 or 2;

Q is $-C(R^{1a})H-$;

R^{1a} is hydrogen or $-OR^5$;

R^1 is ethyl, *n*-propyl, 1-methylethyl, *n*-butyl, *n*-pentyl, 1,1-dimethylethyl, 3-methylhexyl, 2-methylhexyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C$

(O)R^4, -C(S)R^4, -C(R^4)$_2$C(O)R^5, -R^8-C(O)OR^4, -C(S)OR^4, -R^8-C(O)N(R^4)R^5, -C(S)N(R^4)R^5, -N(R^5)C(O)R^4, -N(R^5)C(S)R^4, -N(R^5)C(O)OR^4, -N(R^5)C(S)OR^4, -N(R^5)C(O)N(R^4)R^5, -N(R^5)C(S)N(R^4)R^5, -N(R^5)S(O)$_n$$R^4$, -N($R^5$)S(O)$_n$N($R^4$)$R^5$, -$R^8$-S(O)$_n$N($R^4$)$R^5$, -N($R^5$)C(=N$R^5$)N($R^4$)$R^5$,

and -N(R^5)C(=N-CN)N(R^4)R^5, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-N(R^4)R^5, -S(O)$_m$$R^4$, -$R^8$-S(O)$_n$N($R^4$)$R^5$, -$R^8$-C(O)$R^4$, -$R^8$-C(O)O$R^4$, -$R^8$-C(O)N($R^4$)$R^5$, -N($R^5$)C(O)$R^4$, and -N($R^5$)S(O)$_n$$R^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl,

heteroarylalkyl, -R^8-CN-R^8-NO_2, -R^8-OR^5, -R^8-N(R^4)R^5, -N=C(R^4)R^5, -S(O)$_m$$R^4$, -OS(O)$_2$$CF_3$, -$R^8$-C(O)$R^4$, -C(S)$R^4$, -C($R^4$)$_2$C(O)$R^5$, -$R^8$-C(O)O$R^4$, -C(S)O$R^4$, -$R^8$-C(O)N($R^4$)$R^5$, -C(S)N($R^4$)$R^5$, -N($R^5$)C(O)$R^4$, -N($R^5$)C(S)$R^4$, -N($R^5$)C(O)O$R^4$, -N($R^5$)C(S)O$R^4$, -N($R^5$)C(O)N($R^4$)$R^5$, -N($R^5$)C(S)N($R^4$)$R^5$, -N($R^5$)S(

$O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$,

and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

43. The compound of Claim 42 wherein:

j is 0 and k is 1 or 2;

Q is $-C(R^{1a})H-$;

R^{1a} is hydrogen or $-OR^5$;

R^1 is ethyl, *n*-propyl, 1-methylethyl, *n*-butyl, *n*-pentyl, 1,1-dimethylethyl, 3-methylhexyl, 2-methylhexyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, $-R^8-C(O)R^5$, $-R^8-C(O)OR^5$, $-R^8-C(O)N(R^4)R^5$, $-S(O)_2-R^5$, $-R^9-S(O)_m-R^5$ (where m is 0, 1 or 2), $-R^8-OR^5$, $-R^8-CN$, $-R^9-P(O)(OR^5)_2$, or $-R^9-O-R^9-OR^5$;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

44. The compound of Claim 43 wherein:

j is 0 and k is 1 or 2;

Q is $-C(R^{1a})H-$;

R^{1a} is hydrogen or $-OR^5$;

R^1 is pentyl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring; and

each R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl.

45. The compound of Claim 44 which is:

1'-pentyl-6,7-dihydrospiro[indeno[5,6-*d*][1,3]dioxole-5,3'-indol]-2'(1'*H*)-one;

1-pentyl-7',8'-dihydro-6'*H*-spiro[indole-3,5'-naphtho[2,3-*d*][1,3]dioxol]-2(1*H*)-one; or

7-methoxy-1'-pentyl-6,7-dihydrospiro[indeno[5,6-*d*][1,3]dioxole-5,3'-indol]-2'(1'*H*)-one.

46. The compound of Claim 41 wherein:

j is 0 and k is 1 or 2;

Q is -C(O)-, -CF_2-, -C(O)O- or -N(R^5)C(O)-;

R^1 is ethyl, *n*-propyl, 1-methylethyl, *n*-butyl, *n*-pentyl, 1,1-dimethylethyl, 3-methylhexyl, 2-methylhexyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, -R^8-C(O)R^5, -R^8-C(O)OR^5, -R^8-C(O)N(R^4)R^5, -$S(O)_2$-R^5, -R^9-$S(O)_m$-R^5 (where m is 0, 1 or 2), -R^8-OR^5, -R^8-CN, -R^9-P(O)(OR^5)$_2$, or -R^9-O-R^9-OR^5;

or R^1 is aralkyl substituted by -C(O)N(R^6)R^7 where:

R^6 is hydrogen, alkyl, aryl or aralkyl; and

R^7 is hydrogen, alkyl, haloalkyl, -R^9-CN, -R^9-OR^5, -R^9-N(R^4)R^5, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, or heteroarylalkyl;

or R^6 and R^7, together with the nitrogen to which they are attached, may form a heterocyclyl or heteroaryl;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^6 and R^7 may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, halo, haloalkyl, -R^8-CN, -R^8-OR^5, heterocyclyl and heteroaryl;

or R^1 is aralkyl optionally substituted by one or more substituents selected from the group consisting of -R^8-OR^5, -C(O)OR^5, halo, haloalkyl, alkyl, nitro, cyano, aryl, aralkyl, heterocyclyl and heteroaryl;

or R^1 is -R^9-N(R^{10})R^{11}, -R^9-N(R^{12})C(O)R^{11} or -R^9-N(R^{10})C(O)N(R^{10})R^{11} where:

each R^{10} is hydrogen, alkyl, aryl, aralkyl or heteroaryl;

each R^{11} is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^9-OC(O)R^5, -R^9-C(O)OR^5, -R^9-C(O)N(R^4)R^5, -R^9-C(O)R^5, -R^9-N(R^4)R^5, -R^9-OR^5, or -R^9-CN; and

R^{12} is hydrogen, alkyl, aryl, aralkyl or -C(O)R^5;

and wherein each aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{10} and R^{11} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, cycloalkyl,

aryl, aralkyl, halo, haloalkyl, nitro, -R^8-CN, -R^8-OR^5, -R^8-C(O)R^5, heterocyclyl and heteroaryl;

or R^1 is heterocyclylalkyl or heteroarylalkyl where the heterocyclylalkyl or the heteroarylalkyl group is optionally substituted by one or more substituents selected from the group consisting of oxo, alkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-OR^5, -R^8-C(O)OR^5, -R^8-N(R^4)R^5, -R^8-C(O)N(R^4)R^5, -R^8-N(R^5)C(O)R^4, -R^8-S(O)$_m R^4$ (where m is 0, 1 or 2), -R^8-CN, or -R^8-NO_2;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-N(R^4)R^5, -N=C(R^4)R^5, -S(O)$_m R^4$, -$OS(O)_2CF_3$, -R^8-C(O)R^4, -C(S)R^4, -C(R^4)$_2$C(O)R^5, -R^8-C(O)OR^4, -C(S)OR^4, -R^8-C(O)N(R^4)R^5, -C(S)N(R^4)R^5, -N(R^5)C(O)R^4, -N(R^5)C(S)R^4, -N(R^5)C(O)OR^4, -N(R^5)C(S)OR^4, -N(R^5)C(O)N(R^4)R^5, -N(R^5)C(S)N(R^4)R^5, -N(R^5)S(O)$_n R^4$, -N(R^5)S(O)$_n$N(R^4)R^5, -R^8-S(O)$_n$N(R^4)R^5, -N(R^5)C(=NR^5)N(R^4)R^5, and -N(R^5)C(=N-CN)N(R^4)R^5, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

and wherein each of the cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl groups for R^{2a}, R^{2b}, R^{2c} and R^{2d} may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -R^8-CN, -R^8-NO_2, -R^8-OR^5, -R^8-N(R^4)R^5, -S(O)$_m R^4$, -R^8-S(O)$_n$N(R^4)R^5, -R^8-C(O)R^4; -R^8-C(O)OR^4, -R^8-C(O)N(R^4)R^5, -N(R^5)C(O)R^4, and -N(R^5)S(O)$_n R^4$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{2a} and R^{2b}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2c} and R^{2d} are as defined above;

or R^{2b} and R^{2c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2d} are as defined above;

or R^{2c} and R^{2d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, aryl, heterocyclyl and heteroaryl, and R^{2a} and R^{2b} are as defined above;

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, halo, haloalkyl, haloalkenyl, haloalkoxy, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^8-CN$, $-R^8-NO_2$, $-R^8-OR^5$, $-R^8-N(R^4)R^5$, $-N=C(R^4)R^5$, $-S(O)_mR^4$, $-OS(O)_2CF_3$, $-R^8-C(O)R^4$, $-C(S)R^4$, $-C(R^4)_2C(O)R^5$, $-R^8-C(O)OR^4$, $-C(S)OR^4$, $-R^8-C(O)N(R^4)R^5$, $-C(S)N(R^4)R^5$, $-N(R^5)C(O)R^4$, $-N(R^5)C(S)R^4$, $-N(R^5)C(O)OR^4$, $-N(R^5)C(S)OR^4$, $-N(R^5)C(O)N(R^4)R^5$, $-N(R^5)C(S)N(R^4)R^5$, $-N(R^5)S(O)_nR^4$, $-N(R^5)S(O)_nN(R^4)R^5$, $-R^8-S(O)_nN(R^4)R^5$, $-N(R^5)C(=NR^5)N(R^4)R^5$, and $-N(R^5)C(N=C(R^4)R^5)N(R^4)R^5$, wherein each m is independently 0, 1, or 2 and each n is independently 1 or 2;

or R^{3a} and R^{3b}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3c} and R^{3d} are as defined above;

or R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3d} are as defined above;

or R^{3c} and R^{3d}, together with the carbon ring atoms to which they are directly attached, may form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl, and R^{3a} and R^{3b} are as defined above;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl;

or when R^4 and R^5 are each attached to the same nitrogen atom, then R^4 and R^5, together with the nitrogen atom to which they are attached, may form a heterocyclyl or heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

47. The compound of Claim 46 wherein:

j is 0 and k is 1 or 2;

Q is -C(O)-, -CF_2-, -C(O)O- or -N(R^5)C(O)-;

R^1 is ethyl, *n*-propyl, 1-methylethyl, *n*-butyl, *n*-pentyl, 1,1-dimethylethyl, 3-methylhexyl, 2-methylhexyl, alkenyl, alkynyl, haloalkyl, aryl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, -R^8-C(O)R^5, -R^8-C(O)OR^5, -R^8-C(O)N(R^4)R^5, -$S(O)_2$-R^5, -R^9-$S(O)_m$-R^5 (where m is 0, 1 or 2), -R^8-OR^5, -R^8-CN, -R^9-P(O)(OR^5)$_2$, or -R^9-O-R^9-OR^5;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused ring selected from cycloalkyl, heterocyclyl, aryl or heteroaryl;

each R^4 and R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl;

each R^8 is a direct bond or a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain; and

each R^9 is a straight or branched alkylene chain, a straight or branched alkenylene chain or a straight or branched alkynylene chain.

48. The compound of Claim 47 wherein:

j is 0 and k is 1 or 2;

Q is -C(O)-, -CF_2-, -C(O)O- or -N(R^5)C(O)-;

R^1 is pentyl;

R^{2a}, R^{2b}, R^{2c} and R^{2d} are each hydrogen;

R^{3a} and R^{3d} are each hydrogen;

R^{3b} and R^{3c}, together with the carbon ring atoms to which they are directly attached, form a fused dioxolyl ring; and

each R^5 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl and heteroaryl.

49. The compound of Claim 48 which is:

1'-pentylspiro[indeno[5,6-*d*][1,3]dioxole-5,3'-indole]-2',7(1'*H*,6*H*)-dione;

1-pentyl-6'*H*-spiro[indole-3,5'-naphtho[2,3-*d*][1,3]dioxole]-2,8'(1*H*,7'*H*)-dione;

8',8'-difluoro-1-pentyl-7',8'-dihydro-6'*H*-spiro[indole-3,5'-naphtho[2,3-*d*][1,3]dioxol]-2(1*H*)-one;

1'-pentyl-6,7-dihydro-5*H*-spiro[1,3-dioxolo[4,5-*g*]isoquinoline-8,3'-indole]-2',5(1'*H*)-dione; and

1'-hexylspiro[1,3-dioxolo[4,5-*g*]chromene-8,3'-indole]-2',6(1'*H*,7*H*)-dione.

50. Use of a compound according to any one of Claims 1-49 or a stereoisomer, enantiomer, or tautomer thereof or mixtures thereof or a pharmaceutically acceptable salt thereof for treating, preventing or ameliorating a disease or a condition in a mammal selected from the group consisting of pain, depression, cardiovascular diseases, respiratory diseases, and psychiatric diseases, and combinations thereof.

51. The use of Claim 50, wherein the disease or condition is pain selected from the group consisting of neuropathic pain, inflammatory pain, visceral pain, cancer pain, chemotherapy pain, trauma pain, surgical pain, dental pain, post-surgical pain, childbirth pain, labor pain, chronic pain, persistent pain, peripherally mediated pain, centrally mediated pain, chronic headache, migraine headache, sinus headache, tension headache, phantom limb pain, and peripheral nerve injury, or combinations thereof.

52. The use of Claim 50, wherein the disease or condition is pain associated with a disease or condition selected from the group consisting of HIV, HIV treatment induced neuropathy, heat sensitivity, tosarcoidosis, irritable bowel syndrome, Crohns disease, multiple sclerosis, amyotrophic lateral sclerosis, diabetic neuropathy, peripheral neuropathy, rheumatoid arthritis, osteoarthritis, atherosclerosis, paroxysmal dystonia, myasthenia syndromes, myotonia, malignant

hyperthermia, cystic fibrosis, pseudoaldosteronism, rhabdomyolysis, hypothyroidism, bipolar depression, anxiety, schizophrenia, sodium channel toxin related illnesses, neurogenic bladder, ulcerative colitis, cancer, epilepsy, partial and general tonic seizures, restless leg syndrome, arrhythmias, schaemic conditions caused by stroke or neural trauma, tachy-arrhythmias, atrial fibrillation and ventricular fibrillation.

53. The use of Claim 50 wherein the disease or condition is pain is selected from trigeminal neuralgia, post-herpetic neuralgia, eudynia, familial erythromelalgia, primary erythromelalgia, familial rectal pain or fibromyalgia.

54. The use of Claim 50 wherein the disease or condition is trigeminal neuralgia.

55. The use of Claim 50 wherein the disease or condition is post-herpetic neuralgia.

56. The use of Claim 50 wherein the disease or condition is familial erythromelalgia.

57. The use of Claim 50 wherein the disease or condition is primary erythromelalgia.

58. The use of Claim 50 wherein the disease or condition is familial rectal pain.

59. The use of Claim 50 wherein the disease or condition is fibromyalgia.

60. A pharmaceutical composition comprising a pharmaceutically acceptable excipient and a compound of any one of Claims 1-41; or a stereoisomer, enantiomer or tautomer thereof or mixtures thereof; or a pharmaceutically acceptable salt thereof.

61. 1'-{[5-(Trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, or a pharmaceutically acceptable salt thereof.

62. A pharmaceutical composition comprising a pharmaceutically acceptable excipient and 1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-

one, or a pharmaceutically acceptable salt thereof.

63. Use of 1'-{[5-(trifluoromethyl)-2-furyl]methyl}spiro[furo[2,3-*f*][1,3]benzodioxole-7,3'-indol]-2'(1'*H*)-one, or a pharmaceutically acceptable salt thereof, for treating pain in a mammal.

64. The use of Claim 63, wherein the pain is neuropathic pain, inflammatory pain, visceral pain, cancer pain, chemotherapy pain, trauma pain, surgical pain, dental pain, post-surgical pain, childbirth pain, labor pain, chronic pain, persistent pain, peripherally mediated pain, centrally mediated pain, chronic headache, migraine headache, sinus headache, tension headache, phantom limb pain, or peripheral nerve injury, or combinations thereof.

65. The use of Claim 63, wherein the pain is associated with a disease or condition which is HIV, HIV treatment induced neuropathy, heat sensitivity, tosarcoidosis, irritable bowel syndrome, Crohns disease, multiple sclerosis, amyotrophic lateral sclerosis, diabetic neuropathy, peripheral neuropathy, rheumatoid arthritis, osteoarthritis, atherosclerosis, paroxysmal dystonia, myasthenia syndromes, myotonia, malignant hyperthermia, cystic fibrosis, pseudoaldosteronism, rhabdomyolysis, hypothyroidism, bipolar depression, anxiety, schizophrenia, sodium channel toxin related illnesses, neurogenic bladder, ulcerative colitis, cancer, epilepsy, partial and general tonic seizures, restless leg syndrome, arrhythmias, ischaemic conditions caused by stroke or neural trauma, tachy-arrhythmias, atrial fibrillation or ventricular fibrillation.

66. The use of Claim 63 wherein the pain is trigeminal neuralgia, post-herpetic neuralgia, eudynia, familial erythromelalgia, primary erythromelalgia, familial rectal pain or fibromyalgia.



(I)