



(86) Date de dépôt PCT/PCT Filing Date: 2004/12/24
(87) Date publication PCT/PCT Publication Date: 2005/07/14
(45) Date de délivrance/Issue Date: 2013/10/01
(85) Entrée phase nationale/National Entry: 2006/06/21
(86) N° demande PCT/PCT Application No.: JP 2004/019750
(87) N° publication PCT/PCT Publication No.: 2005/063981
(30) Priorité/Priority: 2003/12/25 (JP2003-431408)

(51) Cl.Int./Int.Cl. *C12N 15/09* (2006.01),
A61K 39/395 (2006.01), *A61P 37/00* (2006.01),
C07K 16/28 (2006.01), *C12P 21/08* (2006.01)

(72) Inventeurs/Inventors:
TAKAHASHI, NOBUAKI, JP;
MIURA, TORU, JP;
KITAGAWA, YOSHINORI, JP;
HIRANO, AKI, JP

(73) Propriétaire/Owner:
KYOWA HAKKO KIRIN CO., LTD., JP

(74) Agent: SMART & BIGGAR

(54) Titre : MUTANTS D'UN ANTICORPS ANTI-CD40
(54) Title: ANTI-CD40 ANTIBODY MUTANTS

(57) **Abrégé/Abstract:**

A mutant of a potentially therapeutic anti-CD40 antibody is provided which mutant has reduced ADCC and CDC activities designed to be optimized as a pharmaceutical agent. A mutant of an agonistic anti-CD40 antibody, comprising mutation and/or substitution of at least one amino acid in the constant region to reduce the ADCC and/or CDC activities therein, and a mutant of an antagonistic anti-CD40 antibody, comprising at least one mutation or substitution in the constant region to reduce the ADCC and/or CDC activities therein, both mutants having at least a hinge region derived from a human IgG2.



ABSTRACT

A mutant of a potentially therapeutic anti-CD40 antibody is provided which mutant has reduced ADCC and CDC activities designed to be optimized as a pharmaceutical agent. A mutant of an agonistic anti-CD40 antibody, comprising mutation and/or substitution of at least one amino acid in the constant region to reduce the ADCC and/or CDC activities therein, and a mutant of an antagonistic anti-CD40 antibody, comprising at least one mutation or substitution in the constant region to reduce the ADCC and/or CDC activities therein, both mutants having at least a hinge region derived from a human IgG2.

DESCRIPTION

ANTI-CD40 ANTIBODY MUTANTS

Technical Field

The present invention relates to an anti-CD40 antibody which recognizes CD40 which is a type of cell membrane molecules associated with immunity. Further, the present invention relates to an antibody with a mutation in the constant region of the human antibody or with a subclass having its portion substituted in order to decrease an ADCC and/or CDC activity, while keeping an agonistic or antagonistic activity.

Background Art

1. CD40

CD40 is an antigen having a molecular weight of 50 kDa which is present on the surface of cell membrane, and expressed in B cells, dendritic cells (DCs), some types of cancer cells, and thymic epithelial cells. CD40 is known to play an important role in proliferation and differentiation of B cells and DCs. CD40 was identified as an antigen expressed on the surface of human B cells (E. A. Clark et al., Proc. Natl. Acad. Sci. USA 83: 4494, 1986; and I. Stamenkovic et al., EMBO J. 8: 1403, 1989) and has been considered as a member of the TNF receptor family which includes low-affinity NGF receptors, TNF receptors, CD27, OX40 and CD30. Ligands (CD40Ls) to human and murine CD40s have been recently cloned and found to be membrane proteins type II and expressed in activated CD4+T cells. CD40L has been also found to introduce strong signals for activation into human or murine B cells.

In dendritic cells, CD40 has been observed to be more highly expressed than in B cells, and it has become clear that CD40 plays an important role in dendritic cells. Binding of CD40 to CD40L activates antigen presenting cells (APCs), that is, expresses costimulator molecules such as CD80 (B7-1) and CD86 (B7-2) or enhances production of IL-2 (Caux, C., et al.: Activation of human dendritic cells through CD40 cross-linking. J. Exp. Med., 180: 1263, 1994; and Shu, U., et al.: Activated T cells induce interleukin-12 production by

monocyte via CD40 - CD40 ligand interaction. Eur. J. Immunol., 25: 1125, 1995). Dendritic cells have a strong antigen-presenting capacity and a strong capacity to activate helper T (Th) cells. Dendritic cells are also believed to control differentiation of naive Th cells into Th1 or Th2 cells. When peripheral blood monocytes, which are myeloid dendritic cells, are cultured in the presence of GM-CSF and IL-4, and matured by CD40L, the resulting matured dendritic cells (DC1) can produce IL-12 *in vitro*, and stimulate and activate allogeneic naive Th cells to induce IFN γ -producing T cells (i.e., to promote their differentiation into Th1). This action is inhibited by anti-IL-12 antibody and hence may be effected via IL-12. On the other hand, when plasmacytoid T cells, which are present in lymphoid T regions and peripheral blood, are cultured in the presence of IL-3 and CD40 ligand, the resulting lymphoid dendritic cells (DC2) are shown to be unable to produce IL-12, and stimulate and activate allogeneic naive Th cells to induce IL-4-producing T cells, which indicates promotion of their differentiation into Th2. It is believed that Th1 cells are involved in activation of cellular immunity, while Th2 cells are associated with enhancement of humoral immunity as well as restriction of cellular immunity. When cytotoxic T cells (CTL) are activated with the help of Th1 cells, they may eliminate pathogens (a number of types of virus, listeria, tuberculosis bacteria, toxoplasma protozoa, etc.) growing in the cytoplasm and tumor cells.

Monoclonal anti-CD40 antibodies, which recognize CD40 expressed on the membrane surface, have been demonstrated to have different biological activities to B cells. Monoclonal anti-CD40 antibodies are generally classified into agonistic or antagonistic antibodies against the interaction between CD40 and CD40L.

2. Agonistic Antibodies

Agonistic antibodies are known to activate B cells. For instance, the anti-CD40 antibodies are reported to induce cell adhesion (Barrett et al., J. Immunol. 146: 1722, 1991; and Gordon et al., J. Immunol. 140: 1425, 1998), increase cell size (Gordon et al., J. Immunol. 140: 1425, 1998; and Valle et al., Eur. J. Immunol. 19: 1463, 1989), induce cell division of B cells activated only by an anti-IgM antibody, anti-CD20 antibody or phorbol ester (Clark and Ledbetter, Proc. Natl. Acad. Sci. USA 83: 4494, 1986; Gordon et al., LEUCOCYTE TYPING III. A. J. McMichael ed. Oxford University Press. Oxford. p.426; and Paulie et al., J.

Immunol. 142: 590, 1989), induce cell division of B cells in the presence of IL4 (Valle et al., Eur. J. Immunol. 19: 1463, 1989; and Gordon et al., Eur. J. Immunol. 17: 1535, 1987), induce expression of IgE by cultured cells stimulated with IL-4 and deprived of T cells (Jabara et al., J. Exp. Med. 172: 1861, 1990; and Gascan et al., J. Immunol. 147: 8, 1991), induce expression of IgG and IgM by those cultured cells (Gascan et al., J. Immunol. 147: 8, 1991), secrete soluble CD23/FcεRII from cells via IL-4 (Gordon and Guy, Immunol. Today 8: 39, 1987; and Cairns et al., Eur. J. Immunol. 18: 349, 1988), enhance expression of soluble CD23/FcεRII on the cells via IL4 (Challa, A., Allergy, 54: 576, 1999), and promote IL-6 production (Clark and Shu, J. Immunol. 145: 1400, 1990). Furthermore, it is reported that addition of IL-4 and an anti-CD40 antibody to human primary culture B cells in the presence of CDw32 + adhesive cells led to establishment of cloned B cells derived therefrom (Bancherauet et al., Science 241: 70, 1991), and apoptosis of germinal center cells was inhibited through CD40 irrespective of whether its antigen receptor was active or inactive (Liu et al., Nature 342: 929, 1989). As described above, CD40 has been identified as antigen expressed on the surface of human B cells, and consequently, most of the isolated antibodies have been evaluated, as an index, mainly using their induction potency for proliferation and/or differentiation of human B cells, or their induction activity for cell death of cancer cells (Katira, A. et al., LEUKOCYTE TYPING V. S. F. Schlossossmann, et. al. eds. p.547. Oxford University Press. Oxford; W. C. Flansow et. al., LEUKOCYTE TYPING V. S. F. Schlossossmann, et. al. eds. p.555. Oxford University Press. Oxford; and J. D. Pound et. al., International Immunology, 11: 11, 1999).

The anti-CD40 antibody has been demonstrated to mature DC (Z. H. Zhou et. al., Hybridoma, 18: 471, 1999). Furthermore, the role of CD4 T cells in priming antigen-specific CD8 T cells was reported to be in activation of DC via CD40 - CD40L signaling, and the anti-CD40 monoclonal antibody (mAb) has been found to be able to replace CD40 helper T cells in activation of dendritic cells (DC) (Shoenberger, S. P., et. al.: T-cell help for cytotoxic T lymphocytes is mediated by CD40 - CD40L interactions. Nature, 480, 1998). Also, administration of an anti-CD40 antibody in mice has been found to be able to protect the animal body from CD40-expressing tumor cells as well as CD40-non-expressing tumor cells

(French, R. R., et. al.: CD40 antibody evokes a cytotoxic T-cell response that eradicates lymphoma and bypasses T-cell help. *Nature Medicine*, 5, 1999).

Agonistic anti-CD40 antibodies are expected to be effective for treatment of infectious diseases, due to bacteria, virus, etc., cancer and others, based on their functions described above. Anti-CD40 antibodies with superior agonistic activities are described in WO 02/088186. The representative examples of those agonistic antibodies are KM341-1-19 and 2105 antibodies. The hybridoma KM341-1-19 producing the KM341-1-19 antibody and the hybridoma 2105 producing the 2105 antibody were submitted on 27th, September, 2001 and 17th, April, 2002, respectively, for international deposit under the Budapest Treaty, to International Patent Organisms Depositary, National Institute of Advanced Industrial Science and Technology (central 6, 1-1, Higashi 1, Tsukuba, Ibaraki, Japan). Their accession numbers are FERM BP-7759 (KM341-1-19) and FERM BP-8024 (2105).

3. Antagonistic Antibodies

Taking it in consideration, on the other hand, that CD40 plays an important role in immunologic responses, as aforementioned, it is expected that inhibition of binding of CD40 to its ligands would lead to development of therapeutic agents for immune suppression in organ transplantation and autoimmune diseases. Sawada, Hase and others have reported that the peripheral blood of patients suffering from Crohn's disease has a higher percentage of monocytes highly expressing CD40. However, such antibodies have not been well known yet as inhibit binding of CD40 to its ligands. Those inhibitory antibodies would be useful in functional analysis of CD40 and treatment of diseases requiring activation of CD40. Inhibitory antibodies to CD40 ligands are also suggested to be effective against diseases involving binding of CD40 to the CD40 ligands. However, CD40L was reported to be expressed in activated platelets (V. Henn et al., *Nature* 391: 591, 1998), and if an anti-CD40L antibody is used as a therapeutic agent, thrombus formation may occur reportedly (T. Kawai et al., *Nat. Med.* 6: 114, 2000). From this point of view, antibodies to CD40 are expected to be safer rather than anti-CD40L antibodies as therapeutic antibody agent to inhibit binding of CD40 to its ligands. Anti-CD40 antibodies would be required to inhibit binding of CD40L to CD40 and still not activate CD40 in themselves.

72813-258

Such antagonistic anti-CD40 antibodies may be used for treatment of autoimmune diseases and suppression of immunologic rejections in transplantation of organs, bone marrow, etc., in view of their functions described above. Anti-CD40 antibodies with superior antagonistic activities are described in WO 02/088186. The representative example of those antagonistic antibodies is 4D11 antibody. The hybridoma 4D11 producing the 4D11 antibody was submitted on 27th, September, 2001 for international deposit under the Budapest Treaty, to International Patent Organisms Depositary, National Institute of Advanced Industrial Science and Technology (central 6, 1-1, Higashi 1, Tsukuba, Ibaraki, Japan). The accession number is FERM BP-7758.

Patent Document 1 WO 02/088186

Disclosure of the Invention

The object of the present invention is to create mutants from the potentially therapeutic anti-CD40 antibodies disclosed in WO 02/088186, which mutants are designed optimally as pharmaceutical agent.

As a result of extensive and intensive research, the present inventors have successfully created novel mutants of the agonistic or antagonistic antibodies, which mutants may have a higher therapeutic effect against diseases than known anti-CD40 antibodies, and completed the present invention based thereon. The basic idea on modification of the anti-CD40 antibodies according to the present invention will be described in detail below.

Specific aspects of the invention relate to:

- a pharmaceutical composition for use in prevention or treatment of transplant rejection, autoimmune diseases, allergy or blood clotting factor VIII inhibition, comprising a monoclonal antibody as an active ingredient, wherein the monoclonal antibody consists of two heavy chains, each represented by an amino acid sequence ranging from glutamine at position 27 to lysine at position 474 in SEQ ID NO: 140, and two light chains, each represented by an amino acid sequence ranging from alanine at position 23 to cysteine at position 235 in SEQ ID NO: 142;

72813-258

- use of a monoclonal antibody for production of a pharmaceutical composition for prevention or treatment of transplant rejection, autoimmune diseases, allergy or blood clotting factor VIII inhibition, wherein the monoclonal antibody consists of two heavy chains, each represented by an amino acid sequence ranging from glutamine at position 27 to lysine at position 474 in SEQ ID NO: 140, and two light chains, each represented by an amino acid sequence ranging from alanine at position 23 to cysteine at position 235 in SEQ ID NO: 142; and

- use of a monoclonal antibody for prevention or treatment of transplant rejection, autoimmune diseases, allergy or blood clotting factor VIII inhibition, wherein the monoclonal antibody consists of two heavy chains, each represented by an amino acid sequence ranging from glutamine at position 27 to lysine at position 474 in SEQ ID NO: 140, and two light chains, each represented by an amino acid sequence ranging from alanine at position 23 to cysteine at position 235 in SEQ ID NO: 142.

The present specification shall encompass the description in the specification and/or drawings of JP Patent Publication (Kokai) No. 2003-431408 which is the basis for the priority of the present application.

Biological materials have been deposited with the National Institute of Advanced Industrial Science and Technology as the international patent organism depository and accorded accession numbers FERM BP-7758, FERM BP-7759 and FERM-8024.

Brief Description of the Drawings

Figure 1A-1 shows binding site peptides prepared based on the CD40 sequence, to which the anti-CD40 agonistic antibodies bind;

Figure 1A-2 shows binding site peptides prepared based on the CD40 sequence, to which the anti-CD40 agonistic antibodies bind (a continuation to Figure 1A-1);

Figure 1B-1 shows binding site peptides prepared based on the CD40 sequence, to which the anti-CD40 antagonistic antibodies bind;

Figure 1B-2 shows binding site peptides prepared based on the CD40 sequence, to which the anti-CD40 antagonistic antibodies bind (a continuation to Figure 1B-1);

Figure 2A illustrates binding of the anti-CD40 antibodies to the CD40 mutant;

Figure 2B illustrates binding of the anti-CD40 antibodies to the CD40 mutant;

Figure 2C illustrates binding of the anti-CD40 antibodies to the CD40 mutant;

Figure 3A shows diagrams indicating that the KM341-1-19 antibody having a P331S mutation is as active as the original KM341-1-19 antibody with respect to binding to Ramos cells;

Figure 3B shows diagrams indicating that the KM341-1-19 antibody having a P331S mutation is as active as the original KM341-1-19 antibody with respect to enhancement of CD95 expression of Ramos cells;

Figure 4A shows a diagram indicating that the KM341-1-19 antibody having a P331S mutation has a lower CDC activity via the rabbit complement;

Figure 4B shows a diagram indicating that the G2/4 antibody has a lower complement activity when the human complement is used;

Figure 5A-1 shows diagrams indicating that conversion of the subclass of the 2105 antibody from IgG2 into different subclasses has no effect on its binding to Ramos cells;

Figure 5A-2 shows diagrams indicating that conversion of the subclass of the KM341-1-19 antibody from IgG2 into different subclasses has no effect on its binding to Ramos cells;

Figure 5B-1 shows diagrams indicating that conversion of the subclass of the 2105 antibody from IgG2 into different subclasses lowers an activity to enhance CD95 expression of Ramos cells;

Figure 5B-2 shows diagrams indicating that conversion of the subclass of the KM341-1-19 antibody from IgG2 into different subclasses lowers an activity to enhance CD95 expression of Ramos cells;

Figure 6A-1 shows diagrams indicating that the binding capacity of the KM341-1-19 antibodies to Ramos cells is independent of the varying structure of the hinge region;

Figure 6A-2 shows diagrams indicating that the binding capacity of the 2105 antibodies to Ramos cells is independent of the varying structure of the hinge region;

Figure 6B-1 shows diagrams indicating that the upper and middle hinges of the hinge region are important for the activity of the KM341-1-19 antibodies to enhance CD95 expression of Ramos cells;

Figure 6B-2 shows diagrams indicating that the upper and middle hinges of the hinge region are important for the activity of the 2105 antibodies to enhance CD95 expression of Ramos cells;

Figure 7A shows diagrams indicating that conversion of the subclass of the F72 antibody to IgG2 has no effect on its binding to Ramos cells;

Figure 7B shows diagrams indicating that conversion of the subclass of the F72 antibody to IgG2 raises an activity to enhance CD95 expression of Ramos cells;

Figure 8A shows diagrams indicating that conversion of the subclass of the 4D11 antibody from IgG1 to IgG4 has no effect on its binding to Ramos cells;

Figure 8B shows diagrams indicating that conversion of the subclass of the 4D11 antibody from IgG1 to IgG4 inhibits enhancement by the CD40Ligand of CD95 expression of Ramos cells, at the same extent as otherwise;

Figure 9 shows a diagram indicating that conversion of the subclass of the 4D11 antibody from IgG1 to IgG4 or IgG4PE lowers the ADCC activity;

Figure 10 shows a diagram indicating that conversion of the subclass of the 4D11 antibody from IgG1 to IgG4P lowers the CDC activity;

Figure 11 illustrates a variation in number of B cells in the blood (B220-positive cells among the peripheral blood lymphocytes) over time after 4D11G1, 4D11G4P or 4D11G4PE was administered into human CD40-transgenic mice;

Figure 12A illustrates a higher expression of CD23 of splenic B cells (CD23-positive cells among the splenic B cells) after each anti-CD40 antibody was administered into human CD40-transgenic mice;

Figure 12B illustrates a higher expression of CD86 of splenic B cells (CD86-positive cells among the splenic B cells) after each anti-CD40 antibody was administered into human CD40-transgenic mice;

Figure 12C illustrates a higher expression of CD95 of splenic B cells (CD95-positive cells among the splenic B cells) after each anti-CD40 antibody was administered into human CD40-transgenic mice;

Figure 13A illustrates the suppressive activity of the antigen-specific antibody (IgG1) production by 4D11 and 281-1-10 in human CD40-transgenic mice;

Figure 13B illustrates the suppressive activity of the antigen-specific antibody (IgM) production by 4D11 and 281-1-10 in human CD40-transgenic mice;

Figure 14A illustrates the numbers of B cells in the blood (B220-positive cells among the peripheral blood lymphocytes) during the suppression assay of the antigen-specific antibody producing activity;

Figure 14B illustrates the numbers of splenic B cells (B220-positive cells among the splenic lymphocytes) during the suppression assay of the antigen-specific antibody producing activity;

Figure 15 illustrates a variation in number of B cells in the blood (B220-positive cells among the peripheral blood lymphocytes) over time after 4D11G4P or 4D11G4PE was administered at a dose of 30 mg/kg into cynomolgus monkeys;

Figure 16 illustrates blood IL-12 levels during the assay shown in Figure 15;

Figure 17 shows the suppressive effect of 4D11G4PE on the simian DTH (delayed-type hypersensitivity in male cynomolgus monkeys);

Figure 18 shows the titers of the anti tetanus toxin IgG during the assay with the results shown in Figure 17;

Figure 19 shows the titers of the anti tetanus toxin IgM during the assay with the results shown in Figure 17;

Figure 20A illustrates the respective influences of 4D11G4PE and 5C8 (anti-CD40Ligand antibody) on platelet aggregation;

Figure 20B shows the respective influences of 4D11G4PE and 5C8 (anti-CD40Ligand antibody) on platelet aggregation;

Figure 21 illustrates a variation in oligomer content of 4D11G4P, 4D11G4PE, 4D11G2Ser or 4D11G4/2/4 over time after it was incubated at pH 2.7 and 37°C;

Figure 22 illustrates suppression of rejection of skin grafts by the anti-CD40 antagonistic antibody;

Figure 23 illustrates the volume change of the tumor over time from cell implantation, in a case where 341G2Ser was administered to tumor bearing mice with Ramos cells implanted therein;

Figure 24 illustrates the volume change of the tumor over time from cell implantation, in a case where 341G2Ser was administered to tumor bearing mice with T24 cells implanted therein;

Figure 25 illustrates the volume change of the tumor over time from cell implantation, in a case where 341G2Ser was administered to tumor bearing mice with Hs 766T cells implanted therein; and

Figure 26 illustrates the volume change of the tumor over time from cell implantation, in a case where 341G2Ser was administered to tumor bearing mice with Capan-2 cells implanted therein.

Best Mode for Carrying Out the Invention

1. Modification of Agonistic Antibodies

Antibodies are essentially molecules that function to protect living bodies against foreign bodies, such as microorganisms and viruses, and cancer, and hence they can kill and eliminate such cells binding to themselves. The lethal activity is composed of two different activities, called Antibody-Dependent Cellular Cytotoxicity (abbreviated as ADCC hereinafter) and Complement-Dependent Cytotoxicity (abbreviated as CDC hereinafter).

ADCC refers to a type of cytotoxicity induced by activation of macrophages, NK cells, neutrophil cells, etc., which recognize target cells by binding to the constant region of the antibody via Fc receptors expressed on their surface. In contrast, CDC refers to a type of

cytotoxicity induced by activation of a complement system which occurs through binding of an antibody to an antigen. These activities are known to vary depending on a subclass of the antibody, which has been found to be due to a structural difference in the constant region of antibodies (Charles A. Janeway et al., Immunology, 1997, Current Biology Ltd./Garland Publishing Inc.).

Anti-CD40 agonistic antibodies will be more preferable as therapeutic agent if they have not the activities of ADCC and/or CDC which may induce cell death of CD40-expressing cells, in terms of mechanism of immunoactive action. If CD40-expressing cells are injured by ADCC and/or CDC activities, immunosuppression may occur rather than desired immunoactivation, resulting in exacerbation of the disease. In addition, patients suffering from infectious diseases may have higher ADCC and/or CDC activities. Therefore, when such antibodies are applied to infectious diseases, it is necessary to evaluate them for safety more carefully, for example, using more active rabbit complements than those present in healthy human serum or peripheral blood which could not be effective to detect the above activities in this situation. Accordingly, mutants and recombinants were created which had no activity of ADCC or CDC and examined for their activity.

Since ADCC and/or CDC activities are known to vary depending on a subclass of the antibody of interest, conversion of the subclass may reduce ADCC and/or CDC activities. For the human IgG subclasses, for example, IgG4 is generally known to be a subclass with low activities of both ADCC and CDC, and it is reported that IgG2 is CDC active but poorly active in ADCC, while IgG1 is highly active in both ADCC and CDC (Charles A. Janeway et al., Immunology, 1997, Current Biology Ltd./Garland Publishing Inc.). Selection of a particular subclass by taking advantage of the above characteristics may create a less cytotoxic antibody from the original antibody. A combination of a specific subclass of antibody with such a point mutation as described below may create an antibody with a desired activity. Further, reduction in ADCC and/or CDC activities of an antibody is reported to be attained by incorporation of a mutation into its constant region. For instance, L235, D265, D270, K322, P331, and P329 (each alphabetical letter denotes an amino acid by the single-letter notation, and each number denotes an EU index proposed by Kabat et al. (Kabat et. al., Sequences of

proteins of Immunological Interest, 1991 Fifth edition); such symbols will be used hereinafter.) may play an important role in complement activation by human IgG, and substitution of one of those sites by another amino acid may reduce the CDC activity. Esohe E. Idusogie et. al. J. Immunol. 2000, 164:4178-4184, Yuanyuan Xu et. al. J. Biol. Chem. 1994, 269:3469-3474, Brekke, O. H. et. al. Eur. J. Immunol. 1994, 24:2542, Morgan, A., et. al., Immunology 1995, 86:319, Lund, J., et. al., J. Immunol., 1996, 157:4963, Tao, M. H., et. al., J. Exp. Med. 1993, 178:661). Specifically, substitution of D270, K322, P329, or P331 by A may reduce the CDC activity. Substitution of P331 by S or G may also induce the same thing.

It is believed that Glu233-Ser239, Gly316-Lys338, Lys274-Arg301, Tyr407-Arg416, Asn297, Glu318, Leu234-Ser239, Asp265-Glu269, Asn297-Thr299 and Ala327-Ile332 take a part in binding of IgG to FcR (Duncan, A. R., Woof, J. M., Partridge, L. J., Burton, D. R., and Winter, G. (1988) Nature 332, 563-564, Gessner, J. E., Heiken, H., Tamm, A., and Schmidt, R. E. (1998) Ann. Hematol. 76, 231-248, Gavin, A., Hulett, M, and Hogarth, P. M. (1998) in The Immunoglobulin Receptors and Their Physiological and Pathological Roles in Immunity (van de Winkel, J. G. J., and Hogarth, P. M., eds), pp. 11-35, Kluwer Academic Publishers Group, Dordrecht, The Netherlands, Sautes, C. (1997) in Cell-mediated Effects of Immunoglobulins (Fridman, W. H., and Sautes, C., eds), pp. 29-66, R. G. Landes Co., Austin, TX, Da'ron, M. (1997) Annu. Rev. Immunol. 15, 203-234, Canfield, S. M., and Morrison, S. L. (1991) J. Exp. Med. 173, 1483-1491, Chappel, M. S., Isenman, D. E., Everett, M., Xu, Y.-Y., Dorrington, K. J., and Klein, M. H. (1991) Proc. Natl. Acad. Sci. U.S.A. 88, 9036-9040, Woof, J. M., Partridge, L. J., Jefferis, R., and Burton, D. R. (1986) Mol. Immunol. 23, 319-330, Wines, B. D., Powell, M. S., Parren, P. W. H. I., Barnes, N., and Hogarth, P. M. (2000) J. Immunol. 164, 5313-5318), and thus incorporation of a mutation into one of these regions may reduce the ADCC activity. Specifically, substitution of L235 by E or G237 by A can reduce binding of IgG to FcR.

The antibody according to the present invention has at least one mutation of amino acids to reduce the ADCC and/or CDC activities, preferably 1-20, 1-17, 1-16, 1-15, 1-14, 1-13, 1-12, 1-11, 1-10, 1-9, 1-8, 1-7, 1-6, 1-5, 1-4, 1-3, or 1 or 2 mutations.

The present invention has revealed that in some anti-CD40 antibodies, the hinge region of IgG2 is important in expression of their strong agonistic activities. Replacement of the variable region or the constant region except the hinge region by a counterpart of any different subclass, or incorporation of a point mutation thereinto is expected to not only modulate the ADCC and/or CDC activities, but increase the productivity of the antibody, its stability during purification and storage, and its blood kinetics.

To produce an antibody drug, the stability of the antibody during purification and storage is very important. Since the antibodies so far developed belong mainly to the IgG1 subclass, conversion of the variable region or the constant region except the hinge region to a sequence derived from the IgG1 subclass will be also effective to improve the physical properties of the anti-CD40 agonistic antibodies described above.

The present invention provides mutants of agonistic anti-CD40 antibodies and others as follows:

- [1] A heavy chain of a monoclonal antibody having an agonistic activity, which binds to CD40, wherein the heavy chain comprises an upper hinge and a middle hinge derived from a human IgG2, and a constant region with at least one amino acid deleted or substituted, or with at least one amino acid added thereto, said deletion, substitution or addition being capable of increasing or decreasing ADCC and/or CDC.
- [2] The heavy chain according to [1], wherein the constant region is derived from a human IgG.
- [3] The heavy chain according to [2], wherein the human IgG is a human IgG1.
- [4] The heavy chain according to [2], wherein the human IgG is a human IgG2.
- [5] The heavy chain according to [2], wherein the human IgG is a human IgG3.
- [6] The heavy chain according to [2], wherein the human IgG is a human IgG4.
- [7] The heavy chain according to any of [3] to [5], wherein said substitution of amino acids in the constant region is substitution of proline with serine at position 331 which is indicated by the EU index as in Kabat et al.
- [8] A monoclonal antibody comprising the heavy chain according to any of [1] to [7].

[9] The heavy chain according to any of [1] to [7], wherein the heavy chain comprises a variable region from a heavy chain of a monoclonal antibody produced by the hybridoma KM341-1-19 (Accession No. FERM BP-7759).

[10] A monoclonal antibody consisting of the heavy chain according to [9] and a light chain comprising a variable region from a light chain of a monoclonal antibody produced by the hybridoma KM341-1-19 (Accession No. FERM BP-7759).

[11] The heavy chain according to any of [1] to [7], wherein the heavy chain comprises a variable region of the polypeptide represented by SEQ ID NO: 38.

[12] A monoclonal antibody consisting of the heavy chain according to [11] and a light chain of a monoclonal antibody, wherein the light chain comprises a variable region of the polypeptide represented by SEQ ID NO: 40.

[13] The heavy chain according to [1], wherein the heavy chain consists of a remaining portion provided by removing the signal sequence from the polypeptide represented by SEQ ID NO: 132.

[14] A monoclonal antibody consisting of the heavy chain according to [13] and a light chain of a monoclonal antibody, wherein the light chain consists of a remaining portion provided by removing the signal sequence from the polypeptide represented by SEQ ID NO: 134.

[15] The heavy chain according to [1], wherein the heavy chain is produced by a host comprising an expression vector having the polynucleotide represented by SEQ ID NO: 131.

[16] The monoclonal antibody according to [8], wherein the monoclonal antibody is produced by a host comprising an expression vector having the polynucleotide represented by SEQ ID NO: 131 and the polynucleotide represented by SEQ ID NO: 133.

[17] The heavy chain according to any of [1] to [7], wherein the heavy chain comprises a variable region from a heavy chain of a monoclonal antibody produced by the hybridoma 2105 (Accession No. FERM BP-8024).

[18] A monoclonal antibody consisting of the heavy chain according to [17] and a light chain comprising a variable region from a light chain of a monoclonal antibody produced by the hybridoma 2105 (Accession No. FERM BP-8024).

[19] The heavy chain according to any of [1] to [7], wherein the heavy chain comprises a variable region of the polypeptide represented by SEQ ID NO: 42.

[20] A monoclonal antibody consisting of the heavy chain according to [19] and a light chain of a monoclonal antibody, wherein the light chain comprises a variable region of the polypeptide represented by SEQ ID NO: 44.

[21] The heavy chain according to [1], wherein the heavy chain consists of a remaining portion provided by removing the signal sequence from the polypeptide represented by SEQ ID NO: 136.

[22] A monoclonal antibody consisting of the heavy chain according to [21] and a light chain of a monoclonal antibody, wherein the light chain consists of a remaining portion provided by removing the signal sequence from the polypeptide represented by SEQ ID NO: 138.

[23] The heavy chain according to [1], wherein the heavy chain is produced by a host comprising an expression vector having the polynucleotide represented by SEQ ID NO: 135.

[24] The monoclonal antibody according to [8], wherein the monoclonal antibody is produced by a host comprising an expression vector having the polynucleotide represented by SEQ ID NO: 135 and the polynucleotide represented by SEQ ID NO: 137.

[25] A polynucleotide represented by SEQ ID NO: 131.

[26] A polynucleotide represented by SEQ ID NO: 133.

[27] An expression vector having the polynucleotide according to [25].

[28] An expression vector having the polynucleotide according to [26].

[29] An expression vector having the polynucleotides according to [25] and [26].

[30] A host comprising the expression vector according to [27].

[31] A host comprising the expression vector according to [28].

[32] A host comprising the expression vector according to [29].

[33] A process of producing a heavy chain of a monoclonal antibody, comprising the steps of: culturing the host according to [30] in a culture medium; and obtaining a heavy chain of a monoclonal antibody from the culture and/or the host.

- [34] A process of producing a monoclonal antibody, comprising the steps of: culturing the host according to [32] in a culture medium; and obtaining a monoclonal antibody from the culture and/or the host.
- [35] A polynucleotide represented by SEQ ID NO: 135.
- [36] A polynucleotide represented by SEQ ID NO: 137.
- [37] An expression vector having the polynucleotide according to [35].
- [38] An expression vector having the polynucleotide according to [36].
- [39] An expression vector having the polynucleotides according to [35] and [36].
- [40] A host comprising the expression vector according to [37].
- [41] A host comprising the expression vector according to [38].
- [42] A host comprising the expression vector according to [39].
- [43] A process of producing a heavy chain of a monoclonal antibody, comprising the steps of: culturing the host according to [40] in a culture medium; and obtaining a heavy chain of a monoclonal antibody from the culture and/or the host.
- [44] A process of producing a a monoclonal antibody, comprising the steps of: culturing the host according to [42] in a culture medium; and obtaining a monoclonal antibody from the culture and/or the host.
- [45] A process of producing a heavy chain of a monoclonal antibody having an agonistic activity capable of binding to CD40, comprising the step of substituting the upper hinge and the middle hinge of an antibody, which is not either an upper hinge or a middle hinge derived from a human IgG2, with an upper hinge and a middle hinge derived from a human IgG2, respectively.
- [46] A process of producing a heavy chain of a monoclonal antibody comprising a variable region, and an upper hinge and a middle hinge derived from a human IgG2, comprising the step of identifying a polypeptide forming the variable region, which is from a heavy chain of a monoclonal antibody capable of binding to CD40.
- [47] A process of producing a monoclonal antibody having an agonistic activity capable of binding to CD40, comprising the step of substituting the upper hinge and the middle hinge of

an antibody, which is not either an upper hinge or a middle hinge derived from a human IgG2, with an upper hinge and a middle hinge derived from a human IgG2, respectively.

[48] A process of producing a monoclonal antibody comprising a variable region, and an upper hinge and a middle hinge derived from a human IgG2, comprising the step of identifying a polypeptide forming the variable region, which is from a heavy chain of a monoclonal antibody capable of binding to CD40.

[49] A pharmaceutical composition comprising the monoclonal antibody according to [8], [10], [12], [14], [16], [18], [20], [22] or [24] as an active ingredient.

[50] The pharmaceutical composition according to [49] used for prevention or treatment of a malignant tumor, a pathogen or an autoimmune disease.

[51] A method of prevention or treatment of a malignant tumor, a pathogen or an autoimmune disease, comprising administration of the pharmaceutical composition according to [49] into a mammal.

[52] Use of the monoclonal antibody according to [8], [10], [12], [14], [16], [18], [20], [22] or [24] for production of a pharmaceutical composition used for prevention or treatment of a malignant tumor, a pathogen or an autoimmune disease.

[89] A polynucleotide provided by removing the portion encoding the signal sequence from the polynucleotide represented by SEQ ID NO: 131.

[90] A polynucleotide provided by removing the portion encoding the signal sequence from the polynucleotide represented by SEQ ID NO: 133.

[91] A polynucleotide provided by removing the portion encoding the signal sequence from the polynucleotide represented by SEQ ID NO: 135.

[92] A polynucleotide provided by removing the portion encoding the signal sequence from the polynucleotide represented by SEQ ID NO: 137.

The present invention provides an antibody produced by modification of an agonistic anti-CD40 antibody belonging to the human IgG2, wherein the modified antibody is a mutant having the constant region, exclusive of the upper and middle hinges, substituted with a sequence derived from a different subclass. The subclass is preferably IgG1. The present invention provides an antibody produced by modification of an agonistic anti-CD40 antibody

belonging to the human IgG2, wherein the modified antibody is a mutant having the constant region, exclusive of the hinge region, substituted with a sequence derived from a different subclass. The subclass is preferably IgG1.

Herein, reduction in ADCC and CDC activities means reduction in those activities as compared with the corresponding activities of an anti-CD40 monoclonal antibody other than the mutants described above, for example, as compared with the corresponding activities of a monoclonal antibody produced by the hybridoma KM341-1-19 (Accession No. FERM BP-7759) or 2105 (Accession No. FERM BP-8024). The ADCC and CDC activities may be assayed by any known method, for example, the method described in the Examples herein. The sequences of variable regions in the heavy and light chains of a monoclonal antibody will be presented below which is produced by the hybridoma KM341-1-19 (Accession No. FERM BP-7759) or 2105 (Accession No. FERM BP-8024).

DNA encoding variable regions in the heavy and light chains of the KM341-1-19 antibody and the amino acid sequences of the heavy and light chains will be presented below.

In the heavy chain nucleotide sequence (SEQ ID NO: 37) of the KM341-1-19 antibody, the signal sequence is initiated with adenine (A) at position 50. The boundary between the signal sequence and the variable region is located between "adenine" ([A]) at position 109 and cytosine (C) at position 110, and the boundary between the variable region and the constant region is located between adenine (A) at position 493 and guanine (G) at position 494 (the gene prediction software (Signal P ver.2) was used).

In the heavy chain amino acid sequence (SEQ ID NO: 38) of the KM341-1-19 antibody, the boundary between the signal sequence and the variable region is located between serine (S) at position 20 and glutamine (Q) at position 21, and the boundary between the variable region and the constant region is located between serine (S) at position 148 and alanine (A) at position 149.

Accordingly, the variable region in the heavy chain of the KM341-1-19 antibody has a nucleotide sequence ranging from cytosine (C) at position 110 to adenine (A) at position 493, as seen in SEQ ID NO: 37. Further, the variable region in the heavy chain of the KM341-1-

19 antibody has an amino acid sequence ranging from glutamine (Q) at position 21 to serine (S) at position 148, as seen in SEQ ID NO: 38.

In the light chain nucleotide sequence (SEQ ID NO: 39) of the KM341-1-19 antibody, the signal sequence is initiated with adenine (A) at position 29. The boundary between the signal sequence and the variable region is located between "adenine" ([A]) at position 88 and guanine (G) at position 89, and the boundary between the variable region and the constant region is located between adenine (A) at position 400 and "cytosine" ([C]) at position 401 (the gene prediction software (Signal P ver.2) was used).

In the light chain amino acid sequence (SEQ ID NO: 40) of the KM341-1-19 antibody, the boundary between the signal sequence and the variable region is located between glycine (G) at position 20 and glutamic acid (E) at position 21, and the boundary between the variable region and the constant region is located between lysine (K) at position 124 and "arginine" ([R]) at position 125.

Accordingly, the variable region in the light chain of the KM341-1-19 antibody has a nucleotide sequence ranging from guanine (G) at position 89 to adenine (A) at position 400, as seen in SEQ ID NO: 39. Further, the variable region in the light chain of the KM341-1-19 antibody has an amino acid sequence ranging from glutamic acid (E) at position 21 to lysine (K) at position 124, as seen in SEQ ID NO: 40.

The heavy chain nucleotide sequence (SEQ ID NO: 37) of the KM341-1-19 antibody:

```
GTCGACGCTGAATTCTGGCTGACCAGGGCAGCCACCAGAGCTCCAGACAATGTCTGTCTCCTTCCTCATCTT
CCTGCCCCGTGCTGGGCCTCCCATGGGGTGTCTGTACAGGTCCAAGTGCAGCAGTCAGGTCCAGGACTGGT
GAAGCCCTCGCAGACCCTCTCACTCACCTGTGCCATCTCCGGGGACAGTGTCTCTAGCAACAGTGCTACTTG
GAACTGGATCAGGCAGTCCCCATCGAGAGACCTTGAGTGGCTGGGAAGGACATACTACAGGTCCAAGTGGTA
TCGTGATTATGTAGGATCTGTGAAAAGTCGAATAATCATCAACCCAGACACATCCAACAACAGTTCTCCCT
GCAGCTGAACTCTGTGACTCCCGAGGACACGGCTATATATTACTGTACAAGAGCACAGTGGCTGGGAGGGGA
TTACCCCTACTACTACAGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCTTCAGCCTCCACCAA
GGGCCCATCGGTCTTCCCCCTGGCGCCCTGCTCCAGGAGCACCTCCGAGAGCACAGCGGCCCTGGGCTGCCT
GGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGTGGAAGTCAAGGCGCTCTGACCAGCGGCGTGACAC
CTTCCCAGCTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAACTT
```

CGGCACCCAGACCTACACCTGCAACGTAGATCACAAGCCCAGCAACACCAAGGTGGACAAGACAGTTGAGCG
CAAATGTTGTGTCGAGTGCCACCGTGCCAGCACCACCTGTGGCAGGACCGTCAGTCTTCCTCTTCCCCC
AAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACGTGCGTGGTGGTGGACGTGAGCCACGA
AGACCCCGAGGTCCAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCACGGGA
GGAGCAGTTCAACAGCACGTTCCGTGTGGTCAGCGTCCTCACCGTTGTGCACCAGGACTGGCTGAACGGCAA
GGAGTACAAGTGCAAGGTCTCCAACAAAGGCCTCCCAGCCCCATCGAGAAAACCATCTCCAAAACCAAAGG
GCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTCAGCCT
GACCTGCCTGGTCAAAGGCTTCTACCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAA
CAACTACAAGACCACACCTCCCATGCTGGACTCAGACGGCTCCTTCTTCCTCTACAGCAAGCTCACCGTGGA
CAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACAC
GCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATGAGGATCC

The heavy chain amino acid sequence (SEQ ID NO: 38) of the KM341-1-19 antibody:

MSVSFLIFLPVLGLPWGVLSQVQLQQSGPGLVKPSQTLSTCAISGDSVSSNSATWNWIRQSPSRDLEWLGR
TYYSKWKYRDYVGSVKSRRIINPDTSNNQFSLQLNSVTPEDTAIYYCTRAQWLGGDYPYYYSMDVWGQGTTV
TVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSV
VTVPSSNFGTQTYTCNVDPKPSNTKVDKTVERKCCVECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTC
VVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVVSFLTIVVHQDWLNGKEYKCKVSNKGLPAPIE
KTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSGSFF
LYSKLTVDKSRWQQGNVFCSSVMHEALHNHYTQKSLSLSPGK

The light chain nucleotide sequence (SEQ ID NO: 39) of the KM341-1-19 antibody:

ACTGCTCAGTTAGGACCCAGAGGGAACCATGGAAGCCCCAGCTCAGCTTCTCTTCCTCCTGCTACTCTGGCT
CCCAGATAACCACCGGAGAAATTGTGTTGACACAGTCTCCAGCCACCCTGTCTTTGTCTCCAGGGGAAAGAGC
CACCCTCTCCTGCAGGGCCAGTCAGAGTGTTAGCAGCTACTTAGCCTGGTACCAACAGAAACCTGGCCAGGC
TCCCAGGCTCCTCATCTATGATGCATCCAACAGGGCCACTGGCATCCCAGCCAGGTTTCAGTGGCAGTGGGTC
TGGGACAGACTTCACTCTCACCATCAGCAGCCTAGAGCCTGAAGATTTTGCAGTTTATTACTGTCAGCAGCG
TAGCAACACTTTCGGCCCTGGGACCAAAGTGGATATCAAACGTACG

The light chain amino acid sequence (SEQ ID NO: 40) of the KM341-1-19 antibody:

MEAPAQLLFLLLLWLPD TTGEIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDAS
NRATGIPARFSGSGSGTDFTLTIS SLEPEDFAVYYCQQR SNTFGPGTKVDIKRT

DNA encoding variable regions in the heavy and light chains of the 2105 antibody and the amino acid sequences of the heavy and light chains will be presented below.

In the heavy chain nucleotide sequence (SEQ ID NO: 41) of the 2105 antibody, the signal sequence is initiated with adenine (A) at position 70. The boundary between the signal sequence and the variable region is located between "thymine" ([T]) at position 126 and guanine (G) at position 127, and the boundary between the variable region and the constant region is located between adenine (A) at position 495 and guanine (G) at position 496 (the gene prediction software (Signal P ver.2) was used).

In the heavy chain amino acid sequence (SEQ ID NO: 42) of the 2105 antibody, the boundary between the signal sequence and the variable region is located between cysteine (C) at position 19 and glutamic acid (E) at position 20, and the boundary between the variable region and the constant region is located between serine (S) at position 142 and alanine (A) at position 143.

Accordingly, the variable region in the heavy chain of the 2105 antibody has a nucleotide sequence ranging from guanine (G) at position 127 to adenine (A) at position 495, as seen in SEQ ID NO: 41. Further, the variable region in the heavy chain of the 2105 antibody has an amino acid sequence ranging from glutamic acid (E) at position 20 to serine (S) at position 142, as seen in SEQ ID NO: 42.

In the light chain nucleotide sequence (SEQ ID NO: 43) of the 2105 antibody, the signal sequence is initiated with adenine (A) at position 28. The boundary between the signal sequence and the variable region is located between "adenine" ([A]) at position 87 and guanine (G) at position 88, and the boundary between the variable region and the constant region is located between adenine (A) at position 405 and "cytosine" ([C]) at position 406 (the gene prediction software (Signal P ver.2) was used).

In the light chain amino acid sequence (SEQ ID NO: 44) of the 2105 antibody, the boundary between the signal sequence and the variable region is located between glycine (G) at position 20 and glutamic acid (E) at position 21, and the boundary between the variable region and the constant region is located between lysine (K) at position 126 and "arginine" ([R]) at position 127.

Accordingly, the variable region in the light chain of the 2105 antibody has a nucleotide sequence ranging from guanine (G) at position 88 to adenine (A) at position 405, as seen in SEQ ID NO: 43. Further, the variable region in the light chain of the 2105 antibody has an amino acid sequence ranging from glutamic acid (E) at position 21 to lysine (K) at position 126, as seen in SEQ ID NO: 44.

The heavy chain nucleotide sequence (SEQ ID NO: 41) of the 2105 antibody:

CTGAACACAGACCCGTCGACTCCCAGGTGTTTCCATTCA GTGATCAGCACTGAACACAGAGGACTCACCATG
GAGTTGGGACTGAGCTGGATTTTCCTTTTGGCTATTTTAAAAGGTGTCCAGTGTGAAGTGCAGCTGGTGGAG
TCTGGGGGAGGCTTGGTACAGCCTGGCAGGTCCCTGAGACTCTCCTGTGCAGCCTCTGGATTACCTTTGAT
GATTATGCCATGCACTGGGTCCGGCAAGCTCCAGGGAAGGGCCTGGAGTGGGTCTCAGGTATTAGTTGGAAT
AGTGGTAGCTTGGTGCATGCGGACTCTGTGAAGGGCCGATTACCATCTCCAGAGACAACGCCAAGAACTCC
CTGTATCTGCAAATGAACAGTCTGAGAGCTGAGGACACGGCCTTGTATTACTGTGCAAGAGATAGGCTATTT
CGGGGAGTTAGGTACTACGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCCTCAGCTAGCACC
AAGG

The heavy chain amino acid sequence (SEQ ID NO: 42) of the 2105 antibody:

MELGLSWIFLLAILKGVQCEVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSGISW
NSGSLVHADSVKGRFTISRDNANKNSLYLQMNSLRAEDTALYYCARDRLFGRVRYYGMDVWGQGTITVTVSSAS
TK

The light chain nucleotide sequence (SEQ ID NO: 43) of the 2105 antibody:

CTGCTCAGTTAGGACCCAGAGGGAACCATGGAAGCCCCAGCTCAGCTTCTCTTCCTCCTGCTACTCTGGCTC
CCAGATACCACCGGAGAAATTGTGTTGACACAGTCTCCAGCCACCCTGTCTTTGTCTCCAGGGGAAAGAGCC

ACCCTCTCCTGCAGGGCCAGTCAGAGTGTTAGCAGCTACTTAGCCTGGTACCAACAGAAACCTGGCCAGGCT
 CCCAGGCTCCTCATCTATGATGCATCCAACAGGGCCACTGGCATCCCAGCCAGGTTCAAGTGGCAGTGGGTCT
 GGGACAGACTTCACTCTCACCATCAGCAGCCTAGAGCCTGAAGATTTTGCAGTTTATTACTGTCAGCAGCGT
 AGCCACTGGCTCACTTTCGGCGGGGGGACCAAGGTGGAGATCAAACGTACGGTG

The light chain amino acid sequence (SEQ ID NO: 44) of the 2105 antibody:

MEAPAQLLFLLLLWLPDITGEIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDAS
 NRATGIPARFSGSGSGTDFTLTISLEPEDFAVYYCQQRSHWLTFGGGTKVEIKRTV

In the heavy chain nucleotide sequence (SEQ ID NO: 131) of the 341G2Ser, the boundary between the signal sequence and the variable region is located between "adenine" ([A]) at position 60 and cytosine (C) at position 61, and the boundary between the variable region and the constant region is located between adenine (A) at position 444 and guanine (G) at position 445 (the gene prediction software (Signal P ver.2) was used).

In the heavy chain amino acid sequence (SEQ ID NO: 132) of the 341G2Ser, the boundary between the signal sequence and the variable region is located between serine (S) at position 20 and glutamine (Q) at position 21, and the boundary between the variable region and the constant region is located between serine (S) at position 148 and alanine (A) at position 149.

Accordingly, the variable region in the heavy chain of the 341G2Ser has a nucleotide sequence ranging from cytosine (C) at position 61 to adenine (A) at position 444, as seen in SEQ ID NO: 131. Further, the variable region in the heavy chain of the 341G2Ser has an amino acid sequence ranging from glutamine (Q) at position 21 to serine (S) at position 148, as seen in SEQ ID NO: 132.

The entire heavy chain nucleotide sequence of the 341G2Ser (SEQ ID NO: 131):

ATGTCTGTCTCCTTCCTCATCTTCCTGCCCCGTGCTGGGCCTCCCATGGGGTGTCTGTACAGGTCCAACTG
 CAGCAGTCAGGTCCAGGACTGGTGAAGCCCTCGCAGACCCTCTCACTCACCTGTGCCATCTCCGGGGACAGT
 GTCTCTAGCAACAGTGCTACTTGGAAGTGGATCAGGCAGTCCCCATCGAGAGACCTTGAGTGGCTGGGAAGG

ACATACTACAGGTCCAAGTGGTATCGTGATTATGTAGGATCTGTGAAAAGTCGAATAATCATCAACCCAGAC
 ACATCCAACAACCAGTTCTCCCTGCAGCTGAACTCTGTGACTCCCGAGGACACGGCTATATATTACTGTACA
 AGAGCACAGTGGCTGGGAGGGGATTACCCCTACTACTACAGTATGGACGTCTGGGGCCAAGGGACCACGGTC
 ACCGTCTCCTCAGCTAGCACCAAGGGCCCATCGGTCTTCCCCCTGGCGCCCTGCTCCAGGAGCACCTCCGAG
 AGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCGTGGAACCTCAGGC
 GCTCTGACCAGCGGCGTGCACACCTTCCCAGCTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTG
 GTGACCGTGCCCTCCAGCAACTTCGGCACCCAGACCTACACCTGCAACGTAGATCACAAGCCCAGCAACACC
 AAGGTGGACAAGACAGTTGAGCGCAAATGTTGTGTCGAGTGCCACCGTGCCAGCACCACCTGTGGCAGGA
 CCGTCAGTCTTCCTCTTCCCCCAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACGTGC
 GTGGTGGTGGACGTGAGCCACGAAGACCCCGAGGTCCAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCAT
 AATGCCAAGACAAAGCCACGGGAGGAGCAGTTCAACAGCACGTTCCGTGTGGTCAGCGTCCTCACCGTTGTG
 CACCAGGACTGGCTGAACGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGGCCTCCCAGCCTCCATCGAG
 AAAACCATCTCCAAAACCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAG
 ATGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTACCCCAGCGACATCGCCGTGGAGTGG
 GAGAGCAATGGGCAGCCGGAGAACAACCTACAAGACCACACCTCCCATGCTGGACTCCGACGGCTCCTTCTTC
 CTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCAT
 GAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATGA

The entire heavy chain amino acid sequence of the 341G2Ser (SEQ ID NO: 132):

MSVSFLIFLPVLGLPWGVLSQVQLQQSGPGLVKPSQTLSTCAISGDSVSSNSATWNWIRQSPSRDLEWLGR
 TYYRSKWYRDYVGSVKSRRIINPDTSNNQFSLQLNSVTPEDTAIYYCTRAQWLGGDYPYYYSMDVWGQGTTV
 TVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSV
 VTPSSNFGTQTYTCNVDHKPSNTKVDKTVKCCVECPKPPVAGPSVFLFPPKPKDTLMISRTPEVTC
 VVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQDWLNGKEYKCKVSNKGLPASIE
 KTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFF
 LYSKLTVDKSRWQQGNVFCSSVMHEALHNHYTQKSLSLSPGK

In the light chain nucleotide sequence (SEQ ID NO: 133) of the 341G2Ser, the boundary between the signal sequence and the variable region is located between "adenine"

([A]) at position 60 and guanine (G) at position 61, and the boundary between the variable region and the constant region is located between adenine (A) at position 372 and "cytosine" ([C]) at position 373 (the gene prediction software (Signal P ver.2) was used).

In the light chain amino acid sequence (SEQ ID NO: 134) of the 341G2Ser, the boundary between the signal sequence and the variable region is located between glycine (G) at position 20 and glutamic acid (E) at position 21, and the boundary between the variable region and the constant region is located between lysine (K) at position 124 and "arginine" ([R]) at position 125. Accordingly, the variable region in the light chain of the 341G2Ser has a nucleotide sequence ranging from guanine (G) at position 61 to adenine (A) at position 372, as seen in SEQ ID NO: 133. Further, the variable region in the light chain of the 341G2Ser has an amino acid sequence ranging from glutamic acid (E) at position 21 to lysine (K) at position 124, as seen in SEQ ID NO: 134.

The entire light chain nucleotide sequence of the 341G2Ser (SEQ ID NO: 133):

ATGGAAGCCCCAGCTCAGCTTCTCTTCCTCCTGCTACTCTGGCTCCCAGATACCACCGGAGAAATTGTGTTG
ACACAGTCTCCAGCCACCCTGTCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCCTGCAGGGCCAGTCAGAGT
GTTAGCAGCTACTTAGCCTGGTACCAACAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATGATGCATCC
AACAGGGCCACTGGCATCCCAGCCAGGTTTCAGTGGCAGTGGGTCTGGGACAGACTTCACTCTCACCATCAGC
AGCCTAGAGCCTGAAGATTTTGCAGTTTATTACTGTCAGCAGCGTAGCAACACTTTCGGCCCTGGGACCAAA
GTGGATATCAAACGTACGGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCT
GGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAGGTGGAT
AACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTC
AGCAGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCATCAG
GGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTTGA

The entire light chain amino acid sequence of the 341G2Ser (SEQ ID NO: 134):

MEAPAQLLFLLLLWLPDITGEIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDAS
NRATGIPARFSGSGSGTDFTLTISILEPEDFAVYYCQQRNTFGPGTKVDIKRTVAAPSVFIFPPSDEQLKS
GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYLSSTLTLSKADYEEKHKVYACEVTHQ

GLSSPVTKSFNRGEC

In the heavy chain nucleotide sequence (SEQ ID NO: 135) of the 2105G2Ser, the boundary between the signal sequence and the variable region is located between "thymine" ([T]) at position 57 and guanine (G) at position 58, and the boundary between the variable region and the constant region is located between adenine (A) at position 426 and guanine (G) at position 427 (the gene prediction software (Signal P ver.2) was used).

In the heavy chain amino acid sequence (SEQ ID NO: 136) of the 2105G2Ser, the boundary between the signal sequence and the variable region is located between cysteine (C) at position 19 and glutamic acid (E) at position 20, and the boundary between the variable region and the constant region is located between serine (S) at position 142 and alanine (A) at position 143.

Accordingly, the variable region in the heavy chain of the 2105G2Ser has a nucleotide sequence ranging from guanine (G) at position 58 to adenine (A) at position 426, as seen in SEQ ID NO: 135. Further, the variable region in the heavy chain of the 2105G2Ser has an amino acid sequence ranging from glutamic acid (E) at position 20 to serine (S) at position 142, as seen in SEQ ID NO: 136.

The entire heavy chain nucleotide sequence of the 2105G2Ser (SEQ ID NO: 135):

ATGGAGTTGGGACTGAGCTGGATTTTCCTTTTGGCTATTTTAAAAGGTGTCCAGTGTGAAGTGCAGCTGGTG
GAGTCTGGGGGAGGCTTGGTACAGCCTGGCAGGTCCCTGAGACTCTCCTGTGCAGCCTCTGGATTACCTTT
GATGATTATGCCATGCACTGGGTCCGGCAAGCTCCAGGGAAGGGCCTGGAGTGGGTCTCAGGTATTAGTIGG
AATAGTGGTAGCTTGGTGCATGCGGACTCTGTGAAGGGCCGATTCACCATCTCCAGAGACAACGCCAAGAAC
TCCCTGTATCTGCAAATGAACAGTCTGAGAGCTGAGGACACGGCCTTGTATTACTGTGCAAGAGATAGGCTA
TTTCGGGGAGTTAGGTACTACGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCCTCAGCTAGC
ACCAAGGGCCCATCGGTCTTCCCCCTGGCGCCCTGCTCCAGGAGCACCTCCGAGAGCACAGCGGCCCTGGGC
TGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCGTGGAACCTCAGGCGCTCTGACCAGCGGCGTG
CACACCTTCCCAGCTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGC
AACTTCGGCACCCAGACCTACACCTGCAACGTAGATCACAAGCCCAGCAACACCAAGGTGGACAAGACAGTT

GAGCGCAAATGTTGTGTCGAGTGCCCAACCGTGCCAGCACCACTGTGGCAGGACCGTCAGTCTTCCTCTTC
 CCCCCAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACGTGCGTGGTGGTGGACGTGAGC
 CACGAAGACCCCGAGGTCCAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCA
 CGGGAGGAGCAGTTCAACAGCACGTTCCGTGTGGTCAGCGTCCTCACCGTTGTGCACCAGGACTGGCTGAAC
 GGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGGCCTCCCAGCCTCCATCGAGAAAACCATCTCCAAAACC
 AAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTC
 AGCCTGACCTGCCTGGTCAAAGGCTTCTACCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCG
 GAGAACAACACTACAAGACCACACCTCCCATGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACC
 GTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCAC
 TACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATGA

The entire heavy chain amino acid sequence of the 2105G2Ser (SEQ ID NO: 136):

MELGLSWIFLLAILKGVQCEVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSGISW
 NSGSLVHADSVKGRFTISRDNANKNSLYLQMNSLRAEDTALYYCARDRLFRGVRYYGMDVWGQGTTVTVSSAS
 TKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSS
 NFGTQTYTCNVDHKPSNTKVDKTVERKCCVECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVS
 HEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPASIEKTISK
 KGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSGDSFFLYSKLT
 VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

In the light chain nucleotide sequence (SEQ ID NO: 137) of the 2105G2Ser, the boundary between the signal sequence and the variable region is located between "adenine" ([A]) at position 60 and guanine (G) at position 61, and the boundary between the variable region and the constant region is located between adenine (A) at position 378 and "cytosine" ([C]) at position 379 (the gene prediction software (Signal P ver.2) was used).

In the light chain amino acid sequence (SEQ ID NO: 138) of the 2105G2Ser, the boundary between the signal sequence and the variable region is located between glycine (G) at position 20 and glutamic acid (E) at position 21, and the boundary between the variable region and the constant region is located between lysine (K) at position 126 and "arginine"

([R]) at position 127. Accordingly, the variable region in the light chain of the 2105G2Ser has a nucleotide sequence ranging from guanine (G) at position 61 to adenine (A) at position 378, as seen in SEQ ID NO: 137. Further, the variable region in the light chain of the 2105G2Ser has an amino acid sequence ranging from glutamic acid (E) at position 21 to lysine (K) at position 126, as seen in SEQ ID NO: 138.

The entire light chain nucleotide sequence of the 2105G2Ser (SEQ ID NO: 137):

ATGGAAGCCCCAGCTCAGCTTCTCTTCCTCCTGCTACTCTGGCTCCCAGATACCACCGGAGAAATTGTGTTG
ACACAGTCTCCAGCCACCCTGTCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCCTGCAGGGCCAGTCAGAGT
GTTAGCAGCTACTTAGCCTGGTACCAACAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATGATGCATCC
AACAGGGCCACTGGCATCCCAGCCAGGTTCAGTGGCAGTGGGTCTGGGACAGACTTCACTCTCACCATCAGC
AGCCTAGAGCCTGAAGATTTTGCAGTTTATTACTGTCAGCAGCGTAGCCACTGGCTCACTTTCGGCGGGGGG
ACCAAGGTGGAGATCAAACGTACGGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTG
AAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAG
GTGGATAACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTAC
AGCCTCAGCAGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACC
CATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTTGA

The entire light chain amino acid sequence of the 2105G2Ser (SEQ ID NO: 138):

MEAPAQLLFLLLLWLPDTTGEIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDAS
NRATGIPARFSGSGSGTDFTLTISLEPEDFAVYYCQQRSHWLTFGGGTKVEIKRTVAAPSVFIFPPSDEQL
KSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSSLTLSKADYEEKHKVYACEVT
HQGLSSPVTKSFNRGEC

2. Modification of Antagonistic Antibodies

Anti-CD40 antagonistic antibodies will be more preferable as therapeutic agent as well as the agonistic antibodies, if they have not the activities of ADCC and/or CDC, in terms of mechanism of action. Furthermore, it is important that anti-CD40 antagonistic antibodies have no activity to induce signals by their *in vivo* crosslinking via Fc receptors, even if the ADCC activity cannot be detected. In other words, it is necessary to confirm that they are

not activate the immunity, and such active antibodies may be desired as pharmaceutical agent. Anti-CD40 antagonistic antibodies are promising as therapeutic agent for treating autoimmune diseases or suppressing rejection in organ transplantation. If they induce an agonistic activity due to some effect after they are administered to patients, however weak it may be, the symptoms may worsen in contrast to the desired therapeutic effect. Thus, an antibody without any agonistic activity is more preferable as pharmaceutical agent. In the present invention, incorporation of a point mutation L235E (means substitution of L at position 235 with E; similar symbols will be used hereinafter) into IgG4 has been demonstrated to be effective for *in vivo* reduction in the agonistic activity, in the animal test using monkeys. Although IgG4 is a subclass with low activities of ADCC and CDC, it is reported that when it was attempted to express IgG4 as recombinant protein in cells like CHO, its half-molecules were secreted due to a poor S-S bonding between the heavy chains (Rob C. Aalberse et al., Immunology, 105, 9-19, 2002). To overcome this problem, incorporation of a mutation into the constant region of antibodies is reported to successfully promote the formation of the S-S bonding. Therefore, this type of mutation was also evaluated for its usefulness. Specifically, the mutation of substituting S at position 228 with P was incorporated (S. Angal et al., Molecular Immunology, vol.30, no.1, 105-108, 1993).

For antagonistic antibodies as well as agonistic antibodies, the stability of the antibody during purification and storage is very important. There may be some methods to create such antibodies as are physically better while keeping the antagonistic activity. The antibody pharmaceuticals so far offered commercially belong mostly to the IgG1 subclass, and they are not reported to be problematic in pharmaceutical formulation. Based on these facts, it may be advantageous from the viewpoint of physical properties to derive the constant region of antibodies from IgG1. In case of anti-CD40 antibodies, however, they are desirably lower in ADCC and CDC activities. As a consequence, antibodies having an IgG1-type constant region modified with some point mutations may be desired. The mutations described above are useful to create such antibodies. The IgG1-type constant region may become lower in ADCC and CDC activities by incorporating point mutation P331G therein. It is also observed that incorporation of point mutation L235E into IgG4 eliminates a slight agonistic

activity *in vivo* to make it pharmaceutically more active, but makes it physically less stable at a low pH. Thus, substitution of L235 with an amino acid other than E may make it physically more functional. As to the 4D11 antibody, it is very similar to the 2B11 antibody with respect to the structure of its variable region. The 2B11 antibody has a lower antagonistic activity, but has a higher stability at a low pH, compared with the 4D11 antibody. If some amino acids derived from the constant region of 2B11 are incorporated into the 4D11 antibody, based on the above properties, 4D11 may become more stable. Specifically, point mutation L38V, P58R, G62W, I79M, K81Y, H87Y, S98A, K109R, V120M or T124A in the heavy chain, or N75S in the light chain, or a combination thereof may be effective for that purpose. Specifically, a mutant created by substituting L at position 38 in the variable region of the heavy chain of the 4D11 antibody with V (abbreviated as L38V; similar symbols will be used hereinafter), a P58R mutant, a G62W mutant, a I79M mutant, a K81Y mutant, a H87Y mutant, a S98A mutant, a K109R mutant, a V120M mutant or a T124A mutant, or a N75S mutant in the light chain, or a combination thereof may be provided for that purpose.

The antibody according to the present invention has at least one mutation of amino acids to reduce the ADCC and/or CDC activities, preferably 1-15, 1-13, 1-12, 1-11, 1-10, 1-9, 1-8, 1-7, 1-6, 1-5, 1-4, 1-3, or 1 or 2 mutations.

The present invention provides mutants of antagonistic anti-CD40 antibodies and others as follows:

[53] A heavy chain of a monoclonal antibody having an antagonistic activity capable of binding to CD40, wherein the heavy chain comprises a constant region with at least one amino acid deleted or substituted, or with at least one amino acid added thereto, said deletion, substitution or addition being capable of increasing or decreasing ADCC and/or CDC.

[54] The heavy chain according to [53], wherein the constant region is derived from a human IgG.

[55] The heavy chain according to [54], wherein the human IgG is a human IgG1.

[56] The heavy chain according to [54], wherein the human IgG is a human IgG2.

[57] The heavy chain according to [54], wherein the human IgG is a human IgG3.

[58] The heavy chain according to [54], wherein the human IgG is a human IgG4.

[59] The heavy chain according to any of [55], [57] or [58], wherein said substitution of amino acids in the constant region is substitution of leucine with glutamic acid at position 235 which is indicated by the EU index as in Kabat et al.

[60] A heavy chain according to any of [53] to [58], wherein the heavy chain comprises a constant region with at least one amino acid deleted or substituted, or with at least one amino acid added thereto, said deletion, substitution or addition being capable of promoting the formation of the S-S bond between the heavy chains.

[61] The antibody heavy chain according to [60], wherein said substitution of amino acids in the constant region is substitution of serine with proline at position 228 which is indicated by the EU index as in Kabat et al.

[62] A monoclonal antibody comprising the heavy chain according to any of [53] to [61].

[63] The heavy chain according to any of [53] to [61], wherein the heavy chain comprises a variable region from a heavy chain of a monoclonal antibody produced by the hybridoma 4D11 (Accession No. FERM BP-7758).

[64] A monoclonal antibody comprising the heavy chain according to [63] and a light chain comprising a variable region from a light chain of a monoclonal antibody produced by the hybridoma 4D11 (Accession No. FERM BP-7758).

[65] The heavy chain according to any of [53] to [61], wherein the heavy chain comprises a variable region of the polypeptide represented by SEQ ID NO: 46.

[66] A monoclonal antibody consisting of the heavy chain according to [65] and a light chain of a monoclonal antibody, wherein the light chain comprises a variable region of the polypeptide represented by SEQ ID NO: 48.

[67] The heavy chain according to [53], wherein the heavy chain consists of a remaining portion provided by removing the signal sequence from the polypeptide represented by SEQ ID NO: 140.

[68] A monoclonal antibody consisting of the heavy chain according to [67] and a light chain of a monoclonal antibody, wherein the light chain consists of a remaining portion provided by removing the signal sequence from the polypeptide represented by SEQ ID NO: 142.

- [69] The heavy chain according to [53], wherein the heavy chain is produced by a host comprising an expression vector having the polynucleotide represented by SEQ ID NO: 139.
- [70] The monoclonal antibody according to [62], wherein the monoclonal antibody is produced by a host comprising an expression vector having the polynucleotide represented by SEQ ID NO: 139 and the polynucleotide represented by SEQ ID NO: 141.
- [71] A polynucleotide represented by SEQ ID NO: 139.
- [72] A polynucleotide represented by SEQ ID NO: 141.
- [73] An expression vector having the polynucleotide according to [71].
- [74] An expression vector having the polynucleotide according to [72].
- [75] An expression vector having the polynucleotides according to [71] and [72].
- [76] A host comprising the expression vector according to [73].
- [77] A host comprising the expression vector according to [74].
- [78] A host comprising the expression vector according to [75].
- [79] A process of producing a heavy chain of a monoclonal antibody, comprising the steps of: culturing the host according to [76] in a culture medium; and obtaining a heavy chain of a monoclonal antibody from the culture and/or the host.
- [80] A process of producing a monoclonal antibody, comprising the steps of: culturing the host according to [78] in a culture medium; and obtaining a monoclonal antibody from the culture and/or the host.
- [81] A pharmaceutical composition comprising the monoclonal antibody according to [62], [64], [66], [68] or [70] as an active ingredient.
- [82] The pharmaceutical composition according to [81] used for prevention or treatment of transplant rejection, an autoimmune disease, allergy or blood clotting factor VIII inhibition.
- [83] A method of prevention or treatment of transplant rejection, an autoimmune disease, allergy or blood clotting factor VIII inhibition, which comprises administering the pharmaceutical composition according to [81] into a mammal.
- [84] Use of the monoclonal antibody according to [62], [64], [66], [68] or [70] for production of a pharmaceutical composition used for prevention or treatment of transplant rejection, an autoimmune disease, allergy or blood clotting factor VIII inhibition.

[85] A method of producing a heavy chain of a monoclonal antibody having an antagonistic activity capable of binding to CD40, wherein the agonistic activity is lowered, comprising a step of making deletion or substitution of at least one amino acid, or addition of at least one amino acid in a constant region of a heavy chain of a human antibody.

[86] The method according to [85], wherein the constant region is from a human IgG.

[87] The method according to [86], wherein the human IgG is a human IgG4.

[88] The method according to any of [85] to [87], wherein said substitution of amino acids in the constant region is substitution of leucine with glutamic acid at position 235 which is indicated by the EU index as in Kabat et al.

[93] A polynucleotide provided by removing the portion encoding the signal sequence from the polynucleotide represented by SEQ ID NO: 139.

[94] A polynucleotide provided by removing the portion encoding the signal sequence from the polynucleotide represented by SEQ ID NO: 141.

Additionally, the present invention provides the materials below.

A mutant of an antagonistic anti-CD40 antibody, comprising at least one substitution selected from the group consisting of substitution of L with V at position 38, substitution of P with R at position 58, substitution of G with W at position 62, substitution of I with M at position 79, substitution of K with Y at position 81, substitution of H with Y at position 87, substitution of S with A at position 98, substitution of K with R at position 109, substitution of V with M at position 120 and substitution of T with A at position 124, which substitutions are all carried out in a variable region of a heavy chain of a monoclonal antibody produced by the hybridoma 4D11 (Accession No. FERM BP-7758), and a mutant of an antagonistic anti-CD40 antibody comprising substitution of N with S at position 75 in a variable region of a light chain of the 4D11 antibody.

Herein, reduction in ADCC and CDC activities means reduction in those activities as compared with the corresponding activities of an anti-CD40 monoclonal antibody other than the mutants described above, for example, as compared with the corresponding activities of a monoclonal antibody produced by the hybridoma 4D11 (Accession No. FERM BP-7758). The ADCC and CDC activities may be assayed by any known method, for example, the

method described in the Examples herein. The sequences of variable regions in the heavy and light chains of a monoclonal antibody will be presented below which is produced by the hybridoma 4D11 (Accession No. FERM BP-7758).

DNA encoding variable regions in the heavy and light chains of the 4D11 antibody and the amino acid sequences of the heavy and light chains will be presented below, respectively.

In the heavy chain nucleotide sequence (SEQ ID NO: 45) of the 4D11 antibody, the boundary between the signal sequence and the variable region is located between "cytosine" ([C]) at position 93 and cytosine (C) at position 94, and the boundary between the variable region and the constant region is located between adenine (A) at position 456 and guanine (G) at position 457 (the gene prediction software (Signal P ver.2) was used).

In the heavy chain amino acid sequence (SEQ ID NO: 46) of the 4D11 antibody, the boundary between the signal sequence and the variable region is located between serine (S) at position 26 and glutamine (Q) at position 27, and the boundary between the variable region and the constant region is located between serine (S) at position 147 and alanine (A) at position 148.

Accordingly, the variable region in the heavy chain of the 4D11 antibody has a nucleotide sequence ranging from cytosine (C) at position 94 to adenine (A) at position 456, as seen in SEQ ID NO: 45. Further, the variable region in the heavy chain of the 4D11 antibody has an amino acid sequence ranging from glutamine (Q) at position 27 to serine (S) at position 147, as seen in SEQ ID NO: 46.

In the light chain nucleotide sequence (SEQ ID NO: 47) of the 4D11 antibody, the boundary between the signal sequence and the variable region is located between "thymine" ([T]) at position 124 and guanine (G) at position 125, and the boundary between the variable region and the constant region is located between adenine (A) at position 442 and "cytosine" ([C]) at position 443 (the gene prediction software (Signal P ver.2) was used).

In the light chain amino acid sequence (SEQ ID NO: 48) of the 4D11 antibody, the boundary between the signal sequence and the variable region is located between cytosine (C) at position 22 and alanine (A) at position 23, and the boundary between the variable region

and the constant region is located between lysine (K) at position 128 and "arginine" ([R]) at position 129.

Accordingly, the variable region in the light chain of the 4D11 antibody has a nucleotide sequence ranging from guanine (G) at position 125 to adenine (A) at position 442, as seen in SEQ ID NO: 47. Further, the variable region in the light chain of the 4D11 antibody has an amino acid sequence ranging from alanine (A) at position 23 to lysine (K) at position 128, as seen in SEQ ID NO: 48.

The heavy chain nucleotide sequence (SEQ ID NO: 45) of the 4D11 antibody:

ATATGTCGACGAGTCATGGATCTCATGTGCAAGAAAATGAAGCACCTGTGGTTCTTCCTCCTGCTGGTGGCG
GCTCCCAGATGGGTCCTGTCCCAGCTGCAGCTGCAGGAGTCGGGCCCAGGACTACTGAAGCCTTCGGAGACC
CTGTCCCTCACCTGCACTGTCTCTGGCGGCTCCATCAGCAGTCCTGGTTACTACGGGGGCTGGATCCGCCAG
CCCCCAGGGAAGGGGCTGGAGTGGATTGGGAGTATCTATAAAAGTGGGAGCACCTACCACAACCCGTCCCTC
AAGAGTCGAGTCACCATATCCGTAGACACGTCCAAGAACCAGTTCTCCCTGAAGCTGAGCTCTGTGACCGCC
GCAGACACGGCTGTGTATTACTGTACGAGACCTGTAGTACGATATTTTGGGTGGTTCGACCCCTGGGGCCAG
GGAACCCTGGTCACCGTCTCCTCAGCTAGC

The heavy chain amino acid sequence (SEQ ID NO: 46) of the 4D11 antibody:

MDLMCKKMKHLWFFLLLVAAPRWVLSQLQLQESGPGLLPSETLSLTCTVSGGSISSPGYYGGWIRQPPGKG
LEWIGSIYKSGSTYHNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCTRPVVRVYFGWFDPWGQGLVT
VSSAS

The light chain nucleotide sequence (SEQ ID NO: 47) of the 4D11 antibody:

AGATCTTAAGCAAGTGTAACAACCTCAGAGTACGCGGGGAGACCCACTCAGGACACAGCATGGACATGAGGGT
CCCCGCTCAGCTCCTGGGGCTTCTGCTGCTCTGGCTCCCAGGTGCCAGATGTGCCATCCAGTTGACCCAGTC
TCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGCCGGGCAAGTCAGGGCATTAGCAG
TGCTTTAGCCTGGTATCAGCAGAAACCAGGGAAAGCTCCTAAGCTCCTGATCTATGATGCCTCCAATTTGGA
AAGTGGGGTCCCATCAAGGTTCAAGCGGCAGTGGATCTGGGACAGATTTCCTCTCACCATCAGCAGCCTGCA
GCCTGAAGATTTTGCAACTTATTACTGTCAACAGTTTAATAGTTACCCGACGTTTCGGCCAAGGGACCAAGGT

GGAAATCAAACGTACG

The light chain amino acid sequence (SEQ ID NO: 48) of the 4D11 antibody:

MDMRVPAQLLGLLLLWLPGARCAIQLTQSPSSLSASVGDRVTITCRASQGISSALAWYQQKPGKAPKLLIYD
ASNLESGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQFNSYPTFGQGTKVEIKRT

In the heavy chain nucleotide sequence (SEQ ID NO: 139) of the 4D11 antibody G4PE, the boundary between the signal sequence and the variable region is located between "cytosine" ([C]) at position 78 and cytosine (C) at position 79, and the boundary between the variable region and the constant region is located between adenine (A) at position 441 and guanine (G) at position 442 (the gene prediction software (Signal P ver.2) was used).

In the heavy chain amino acid sequence (SEQ ID NO: 140) of the 4D11 antibody, the boundary between the signal sequence and the variable region is located between serine (S) at position 26 and glutamine (Q) at position 27, and the boundary between the variable region and the constant region is located between serine (S) at position 147 and alanine (A) at position 148.

Accordingly, the variable region in the heavy chain of the 4D11 antibody has a nucleotide sequence ranging from cytosine (C) at position 79 to adenine (A) at position 441, as seen in SEQ ID NO: 139. Further, the variable region in the heavy chain of the 4D11 antibody has an amino acid sequence ranging from glutamine (Q) at position 27 to serine (S) at position 147, as seen in SEQ ID NO: 140.

The entire heavy chain nucleotide sequence (SEQ ID NO: 139) of the 4D11G4PE:

ATGGATCTCATGTGCAAGAAAATGAAGCACCTGTGGTTCTTCCTCCTGCTGGTGGCGGCTCCCAGATGGGTC
CTGTCCCAGCTGCAGCTGCAGGAGTCGGGCCCAGGACTACTGAAGCCTTCGGAGACCCTGTCCCTCACCTGC
ACTGTCTCTGGCGGCTCCATCAGCAGTCCTGGTTACTACGGGGGCTGGATCCGCCAGCCCCAGGGAAGGGG
CTGGAGTGGATTGGGAGTATCTATAAAAGTGGGAGCACCTACCACAACCCGTCCCTCAAGAGTCGAGTCACC
ATATCCGTAGACACGTCCAAGAACCAGTTCTCCCTGAAGCTGAGCTCTGTGACCGCCGCAGACACGGCTGTG
TATTACTGTACGAGACCTGTAGTACGATATTTTGGGTGGTTCGACCCCTGGGGCCAGGGAACCCTGGTCACC

GTCTCCTCAGCTAGCACCAAGGGGCCATCCGTCTTCCCCCTGGCGCCCTGCTCCAGGAGCACCTCCGAGAGC
ACAGCCGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCGTGGAACTCAGGCGCC
CTGACCAGCGGCGTGACACCTTCCCGGCTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTG
ACCGTGCCCTCCAGCAGCTTGGGCACGAAGACCTACACCTGCAACGTAGATCACAAGCCCAGCAACACCAAG
GTGGACAAGAGAGTTGAGTCCAAATATGGTCCCCCATGCCCACCATGCCCAGCACCTGAGTTCGAGGGGGGA
CCATCAGTCTTCCTGTTCCCCCAAAACCCAAGGACACTCTCATGATCTCCCGGACCCCTGAGGTCACGTGC
GTGGTGGTGGACGTGAGCCAGGAAGACCCCGAGGTCCAGTTCAACTGGTACGTGGATGGCGTGGAGGTGCAT
AATGCCAAGACAAAGCCGCGGGAGGAGCAGTTCAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTG
CACCAGGACTGGCTGAACGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGGCCTCCCGTCCTCCATCGAG
AAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAGCCACAGGTGTACACCCTGCCCCCATCCCAGGAGGAG
ATGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTACCCCAGCGACATCGCCGTGGAGTGG
GAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTC
CTCTACAGCAGGCTAACCGTGGACAAGAGCAGGTGGCAGGAGGGGAATGTCTTCTCATGCTCCGTGATGCAT
GAGGCTCTGCACAACCACTACACACAGAAGAGCCTCTCCCTGTCTCTGGGTAAATGA

The entire heavy chain amino acid sequence (SEQ ID NO: 140) of the 4D11G4PE:

MDLMCKKMKHLWFFLLLVAAPRWVLSQLQLQESGPGLLKPSETLSLTCTVSGGSISSPGYYGGWIRQPPGKG
LEWIGSIYKSGSTYHNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCTRPPVVR YFGWFD PWGQGTLVT
VSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVV
TVPSSSLGTKTYTCNVDHKPSNTKVDKRVESKYGPCCPPCPAPEFEGGPSVFLFPPKPKDTLMISRTPEVTC
VVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE
KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFF
LYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSLSLGK

In the light chain nucleotide sequence (SEQ ID NO: 141) of the 4D11G4PE, the boundary between the signal sequence and the variable region is located between "thymine" ([T]) at position 66 and guanine (G) at position 67, and the boundary between the variable region and the constant region is located between adenine (A) at position 384 and "cytosine" ([C]) at position 385 (the gene prediction software (Signal P ver.2) was used).

72813-258

In the light chain amino acid sequence (SEQ ID NO: 142) of the 4D11G4PE, the boundary between the signal sequence and the variable region is located between cytosine (C) at position 22 and alanine (A) at position 23, and the boundary between the variable region and the constant region is located between lysine (K) at position 128 and "arginine" ([R]) at position 129.

Accordingly, the variable region in the light chain of the 4D11G4PE has a nucleotide sequence ranging from guanine (G) at position 67 to adenine (A) at position 384, as seen in SEQ ID NO: 141. Further, the variable region in the light chain of the 4D11 antibody has an amino acid sequence ranging from alanine (A) at position 23 to lysine (K) at position 128, as seen in SEQ ID NO: 142.

The entire light chain nucleotide sequence (SEQ ID NO: 141) of the 4D11G4PE:

ATGGACATGAGGGTCCCCGCTCAGCTCCTGGGGCTTCTGCTGCTCTGGCTCCCAGGTGCCAGATGTGCCATC
CAGTTGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGCCGGGCAAGT
CAGGGCATTAGCAGTGCTTTAGCCTGGTATCAGCAGAAACCAGGGAAAGCTCCTAAGCTCCTGATCTATGAT
GCCTCCAATTTGGAAAGTGGGGTCCCATCAAGGTTACGCGGCAGTGGATCTGGGACAGATTTCACTCTCACC
ATCAGCAGCCTGCAGCCTGAAGATTTTGCAACTTATTACTGTCAACAGTTTAATAGTTACCCGACGTTCCGGC
CAAGGGACCAAGGTGGAAATCAAACGTACGGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAG
CAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAG
TGGAAGGTGGATAACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGC
ACCTACAGCCTCAGCAGCACCTGACGGTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAA
GTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTTGA

The entire light chain amino acid sequence (SEQ ID NO: 142) of the 4D11G4PE:

MDMRVPAQLLGLLLLWLPGARCAIQLTQSPSSLASVGDRVTTICRASQGISSALAWYQQKPGKAPKLLIYD
ASNLESGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQFNSYPTFGQGTKVEIKRTVAAPSVFIFPPSDE
QLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTLSSTLTLSKADYEEKHKVYACE
VTHQGLSSPVTKSFNRGEC

3. Definition

The terms used herein will be defined below.

"CD40" described in the present invention refers to a polypeptide having the amino acid sequence described in E. A. Clark et al., Proc. Natl. Acad. Sci. USA 83: 4494, 1986, or I. Stamenkovic et al., EMBO J. 8: 1403, 1989, and particularly, an antigenic polypeptide expressed on the surface of B cells, DC, macrophages, endothelial cells, epithelial cells or tumor cells derived from them.

"An anti-CD40 antibody" refers to any monoclonal antibody to a cell-expressed CD40, a full-length CD40 or a partial-length CD40.

In addition, "an antibody" of the invention is derived from genes (collectively called antibody genes) encoding a heavy chain variable region and a heavy chain constant region, as well as a light chain variable region and a light chain constant region which together constitute an immunoglobulin. Human immunoglobulins are grouped into 5 different classes consisting of IgG, IgA, IgM, IgD, and IgE. Further, IgG is composed of 4 different subclasses, IgG1, IgG2, IgG3 and IgG4, while IgA is composed of 2 different subclasses, IgA1 and IgA2. IgG1, IgG2, IgG3 and IgG4 are located in 14q32, 33 of the human chromosomes. The fundamental structure of immunoglobulin is composed of two homologous L chains (light chains) and two homologous H chains (heavy chains). The class and subclass of an immunoglobulin is determined by its H chains. The antibody according to the present invention may comprise any class, any subclass or any isotype of immunoglobulin. "A functional fragment" of the inventive antibody refers to a portion (partial fragment) of the antibody defined above that is active singly or multiply on an antigen to the antibody, including, for example, F(ab')₂, Fab', Fab, Fv, disulfide-stabilized FV, single-chain FV (scFV) and a multimer thereof (D. J. King, Applications and Engineering of Monoclonal Antibodies, 1998, T. J. International Ltd.).

Up to now, it has been known that IgG1 includes J00228, Z17370 and Y14737, IgG2 includes J00230, AJ250170, AF449616, AF449617, AF449618, Z49802 and Z49801, IgG3 includes M12958, K01313, X16110, X99549, AJ390236, AJ390237, AJ390238, AJ390241, AJ390242, AJ390246, AJ390247, AJ390252, AJ390244, AJ390254, AJ390260, AJ390262,

AJ390272, AJ390276 and AJ390279, and IgG4 includes K01316, AJ001563 and AJ001564 (the symbols listed above indicates accession numbers of the genes).

In the present invention, CH1, hinge, CH2 and CH3 each denote a portion of the heavy-chain constant region of any antibody, and are based on the EU index as in Kabat et al. (Kabat et al., Sequences of proteins of immunological interest, 1991 Fifth edition). By definition, CH1 ranges from 118 to 215 by the EU index, hinge ranges from 216 to 230 by the EU index, CH2 ranges from 231 to 340 by the EU index, and CH3 ranges from 341 to 446 by the EU index.

The "human antibody" of the present invention means an antibody which is an expression product of a human-derived antibody gene.

"Agonistic" refers to an action of enhancing binding of a ligand to CD40 expressed on the surface of such cells as B cells, tumor cells or dendritic cells, or an action of providing the CD40-expressing cells with at least one effect which the CD40 ligand makes on the CD40-expressing cells. "An agonistic antibody" refers to an antibody having such an agonistic action. An example of the effects provided for the CD40-expressing cells is to promote the expression of CD95.

"Antagonistic" refers to an action of inhibiting binding of the ligand to CD40 expressed on the surface of such cells as B cells, tumor cells or dendritic cells, or an action of neutralizing at least one effect which the CD40 ligand makes on the CD40-expressing cells. "An antagonistic antibody" refers to an antibody having such an antagonistic action. An example of the effects provided for the CD40-expressing cells is to suppress the proliferation of B cells or the production of antibodies.

The present application clearly presents antibodies, or heavy chain or light chain variable regions thereof by amino acid sequences. The present invention also encompasses the amino acid sequences with at least one amino acid deleted or substituted, or added thereto, preferably 1-10, 1-9, 1-8, 1-7, 1-6, 1-5, 1-4, 1-3, or 1 or 2 amino acids.

The present application clearly presents genes which encode antibodies, or heavy chain or light chain variable regions thereof by nucleotide sequences. The present invention also

encompasses the nucleotide sequences with at least one nucleotide deleted or substituted, or added thereto, preferably 1-10, 1-9, 1-8, 1-7, 1-6, 1-5, 1-4, 1-3, or 1 or 2 amino acids.

The anti-CD40 antibody according to the present invention can be provided by incorporating an antibody gene into an expression vector, transfecting the vector into a suitable host cell, harvesting the antibody from the cultured cells or the supernatant, and purifying it.

The vector may be a phage or a plasmid which can replicate in the host cell by itself or can be integrated into the chromosome of the host cell. The plasmid DNA may be derived from *Escherichia coli*, *Bacillus subtilis* or a yeast, while the phage DNA may be from λ phage.

The host cell for transformation is not particularly limited if it can express the target gene. Examples of the host cell may include bacteria (*Escherichia coli*, *Bacillus subtilis*, etc.), yeasts, animal cells (COS cell, CHO cell, etc.) and insect cells.

There are known modes of transferring the gene into the host cells, including any mode, such as mediation by calcium ion, electroporation, spheroplast fusion, mediation by lithium acetate, calcium phosphate transfection or lipofection. In order to transfer the gene into an animal, as described later, the modes include microinjection; electroporation or lipofection for ES cells; and nuclear transplantation.

In the present invention, "culture" refers to (a) culture supernatant, (b) cultured cells, cultured biomass or disrupted matter thereof, or (c) secretion of transformant. To culture the transformant, a medium suitable for the host is used and static culture, roller bottle culture or something else may be employed.

After the culture, if the desired antibody protein is produced within the biomass or cells, the antibody is harvested by disrupting the biomass or cells. If the desired antibody is produced out of the biomass or cells, the culture solution is used as it is or after it is separated from the biomass or cells by centrifugation or other means. Thereafter, a biochemical process utilizing any chromatography, which is appropriate for separation/purification of proteins, is employed alone or optionally in combination with another to separate/isolate the desired antibody from the culture.

Furthermore, the technology of creating a transgenic animal may be used to produce a transgenic animal that is a host animal having the gene integrated into an endogenous gene,

such as a transgenic bovine, a transgenic goat, a transgenic sheep or a transgenic pig (Wright, G., et al., (1991) Bio/Technology 9, 830-834) and a large amount of a monoclonal antibody derived from the antibody gene can be obtained from the milk secreted from the transgenic animal.. The culture of a hybridoma *in vitro* can be done by using a known nutrient medium or any nutrient medium derivatively prepared from known basic media as used to grow, maintain and store the hybridoma and to produce a monoclonal antibody in the supernatant, depending on the properties of the cultured hybridoma, the purpose of the study and the culture method.

4. Antibody Properties

(1) Agonistic Antibodies

The mutant of the agonistic antibody according to the present invention may activate the immune system without injuring immunocompetent cells, since it has an ADCC and/or CDC activity equal to or lower than the original antibody, while keeping an agonistic activity. It is thus expected that the mutant exhibits the immunoactivating action which is equal to or higher than the original antibody and the cytotoxicity to CD40-expressing cells which is equal to or lower than the original antibody.

(2) Antagonistic Antibodies

The mutant of the antagonistic anti-CD40 antibody according to the present invention has the reduced ADCC and/or CDC activity compared to the unmodified antibody, while keeping a suppressive activity against immunoactivating signals induced by CD40L. It is also expected to decrease the activity of signal induction *in vivo* which is considered to occur via Fc receptors.

5. Pharmaceutical Compositions

A pharmaceutical composition containing a formulation of the purified antibody according to the present invention is also within the scope of the present invention. The pharmaceutical composition may preferably contain a physiologically acceptable diluent or carrier in addition to the antibody, and may be a mixture thereof with a different antibody or a different drug such as an antibiotic agent. The suitable carrier may include, but not limited to, physiological saline, phosphate buffered physiological saline, phosphate buffered

physiological saline glucose solution and buffered physiological saline. Alternatively, the antibody may be freeze-dried for storage and when it is used, reconstituted in an aqueous buffer as described above. The pharmaceutical composition may be administered via the oral route, or the parenteral route, such as intravenous, intramuscular, subcutaneous or intraperitoneal injection or dosing.

A single effective dose, which is a combination of the antibody of the present antibody with a suitable diluent and a physiologically acceptable carrier, is from 0.0001 mg to 100 mg per kg of body weight, and it may be taken at a time interval of from 2 days to 8 weeks.

When the pharmaceutical composition of the present antibody is an agonistic antibody, it is used as: immunostimulant (antiviral or anti-infective agent) for pathogens including, for example, hepatitis A, B, C, D or E virus, HIV, influenza virus, simple herpes virus, cytomegalovirus, EB virus, papiloma virus, chlamydia, mycoplasma, toxoplasma, malaria, trypanosome and tubercle bacillus; antitumor agent for malignant tumors having cancer cells with CD40 expressed, including, for example, pancreatic cancer, bladder cancer, lymphoma (e.g., Hodgkin's lymphoma), leukemia, malignant melanoma, pancreatic cancer, lung cancer, ovarian cancer, bladder cancer, breast cancer, colon cancer, prostatic cancer, and head and neck cancer; and therapeutic agent for autoimmune diseases such as rheumatism. The pharmaceutical composition may be used for a combination of the above diseases. It may be also used in combination as adjuvant for a cancer-specific peptide. When the pharmaceutical composition is an antagonistic antibody, on the other hand, it is useful as: immunosuppressant in organ transplantation (preventive or therapeutic agent for rejection in transplantation of pancreatic islets, kidney or something else, or GVHD), therapeutic agent for autoimmune diseases (e.g., rheumatism, psoriasis, chronic ulcerative colitis, Crohn's disease, systemic lupus erythematosus, multiple sclerosis, myasthenia, scleroderma, antiphospholipid antibodies syndrome, autoimmune hepatitis, idiopathic thrombocytopenic purpura, Behcet's syndrome, arteriosclerosis, nephritis and respiratory distress syndrome), therapeutic agent for allergy (e.g., asthma), and therapeutic agent for blood clotting factor VIII inhibition. The pharmaceutical composition may be used for a combination of the above diseases.

6. Epitopes

The binding epitopes of CD40 were determined for the KM341-1-19 and 2105 antibodies having a superior agonistic activity, and for the 4D11 antibody having a superior antagonistic activity, respectively (Example 2). The present invention provides antibodies having an agonistic or antagonistic activity which have a different variable region sequence from those of the above antibodies but recognize the same epitope as one of the above antibodies. These antibodies can be obtained in such a procedure as described below.

When it is intended to acquire an agonistic anti-CD40 antibody recognizing the same epitope as the KM341-1-19 antibody, for example, mice or the likes are immunized with CD40 to provide monoclonal antibodies, from which some monoclonal antibodies competing with the KM341-1-19 antibody to bind to CD40 are screened according to the standard procedure. From the screened antibodies, an antibody having the same pattern of binding to the peptide as the KM341-1-19 antibody is selected according to the method described in Example 2.

The present invention will be described in more detail below with reference to examples. However, the present invention is not limited to embodiments described in the examples.

Example 1: Expression and purification of antibody and antigen proteins

A vector plasmid containing a variable region of an antibody was transfected into CHO cells (ATCC), and antibody-expressing cells were selected by G418 to prepare a stable expression cell line.

A mutant antigen was expressed by transiently introducing a vector into HEK cells (ATCC).

An anti-CD40 antibody was purified from the above culture supernatant by the following method. The culture supernatant containing an anti-CD40 antibody was affinity purified in a Hyper D Protein A column (manufactured by NGK Insulators, Ltd.) or in case of mouse IgG1 purification, a Protein G column (Amersham Pharmacia Biotech) according to the attached instruction using PBS(-) as an adsorption buffer and a 0.1 M sodium citrate buffer (pH 3) as an elution buffer. The eluted fraction was adjusted to about pH 7.2 by addition of a 1 M Tris-HCl (pH 8.0) or Na₂HPO₄ solution. The prepared antibody solution was substituted

72813-258

with PBS(-) using a dialysis membrane (10,000 cuts, manufactured by Spectrum Laboratories, Inc.) or an SP column (Amersham Pharmacia Biotech), and filtered and sterilized using a membrane filter MILLEXTM-GV with a pore diameter of 0.22 μ m (manufactured by Millipore Corp.). The concentration of the purified antibody was calculated by measurement of the absorbance at 280 nm, taking 1 mg/ml as 1.450D.

Example 2: Determination of epitopes

A 13-mer peptide covering amino acid 175 (SEQ ID NO: 1) in an extracellular region of CD40 was shifted by two amino acids each to synthesize 82 peptides in total (SEQ ID NOS: 49 to 130) as spots from the C-terminal on a cellulose membrane and acetylate the N-terminal thereof (Jerini AG, Germany). The reaction thereafter was carried out based on a conventional Western analysis (see Reineke, U. et al. (2001), "Epitope mapping with synthetic peptides prepared by SPOT synthesis." Antibody Engineering (Springer Lab Manual) Eds.: Kontermann/Dubel, 433-459, for example). In the analysis, coloring intensity of each spot was quantified using LumiImagerTM (Boehringer-Mannheim Corp.) (Figures 1A-1, A-2, B-1 and B-2).

The results confirmed that an 4D11 antibody strongly recognizes the 20th to 24th and 41st peptides, a 2105 antibody strongly recognizes the 12th to 23rd and 64th peptides, a KM341-1-19 antibody strongly recognizes the 41st and 42nd peptides, KM643-4-11 strongly recognizes the 43rd peptide, F72 strongly recognizes the 75th peptide, 110 strongly recognizes the 64th peptide, F4-465 strongly recognizes the 34th, 35th, 54th, 55th, 65th, 66th and 75 peptides, KM281-1-10 strongly recognizes the 21st, 24th, 64th and 75th peptides, 2B11 (novel antibody) strongly recognizes the 21st, 24th and 64th peptides, and F76 (novel antibody) strongly recognizes the 21st, 35th, 51st and 52nd peptides.

In order to confirm the binding site of the anti-CD40 antibody, a CD40-FC fusion protein having a mutation introduced therein was prepared, and the binding ability thereto was examined by ELISA. Since the anti-CD40 antibody does not cross-react with mouse B cells, five CD40Fc fusion proteins were prepared by partially converting the amino acid sequence into that of mouse CD40. Binding of the antibody to the antigens was examined.

The method for preparing the mutant CD40-FC fusion proteins is shown below. The mutation site was prepared by introducing a mouse CD40 sequence into a part to which the antibody strongly binds of the peptide sequence. CD40mut1 converted EFTE at a site corresponding to the 15th peptide into ALEK, CD40mut2 converted LDT at a site corresponding to the 21st peptide into SAQ, CD40mut3 converted TH at a site corresponding to the 24th peptide into IR, CD40mut4 converted EEGW at a site corresponding to the 42nd peptide into KEGQ, and CD40mut5 converted VSSA at a site corresponding to the 64th peptide into QSSL. The mutants were prepared according to a gene engineering technique (Figures 2A, B and C). The analysis results confirmed that the 2105 antibody has extremely reduced binding ability to CD40mut1. The results also confirmed that the 4D11 antibody and 2B11 have reduced binding ability to CD40mut2.

Example 3: Binding activity of anti-CD40 agonistic antibody to Ramos cells

A Ramos cell line was suspended in a PBS staining buffer (SB) containing 0.1% NaN₃ and 2% FCS at a concentration of 2×10^6 /ml. The cell suspension (100 μ l/well) was dispensed to a 96-well round-bottom plate (manufactured by Becton, Dickinson and Company). Each hybridoma culture supernatant (50 μ l) was added, and incubated at an ice temperature for 30 minutes. A human IgG1 antibody to human serum albumin as a negative control was adjusted to a concentration of 2 μ g/ml in a hybridoma culture medium, added in an amount of 50 μ l, and then incubated at an ice temperature for 15 minutes. After washing the plate with SB, 50 μ l of a 250-fold diluted R-PE fluorescently labeled anti-human antibody (manufactured by Southern Biotechnology Associates, Inc.) was added, and incubated at an ice temperature for 15 minutes. After washing the plate with SB twice, it was suspended in 300 to 500 μ l of a FACS buffer, and fluorescence intensity of each cell was measured using FACS (FACSort, FACScan, manufactured by Becton, Dickinson and Company).

Example 4: Evaluation of agonistic activity of anti-CD40 agonistic antibody to Ramos cells

5.0×10^5 cells/ml of a Ramos cell suspension was seeded on a 96-well plate at 100 μ l/well. A hybridoma culture supernatant or purified antibody was diluted to 20 μ g/ml in a

72813-258

medium, and the dilution was added to the 96-well plate at a concentration of 100 μ l/well. After culturing overnight, cells were harvested, and R-PE labeled anti-CD95 antibody (Pharmingen NJ) was used for the cells. Analysis was carried out using FACScan or FACSsort (Becton, Dickinson and Company).

Example 5: Inhibition of CD95 expression by anti-CD40 antagonistic antibody in Ramos cells

1.0×10^6 cells/ml of a Ramos cell suspension was seeded on a 96-well plate at 50 μ l/well. A hybridoma culture supernatant or purified antibody was adjusted to 2 μ g/ml in a medium, and the medium was added to the 96-well plate at 100 μ l/well. 4 μ g/ml of a soluble CD40 ligand (Alexis Corporation) and 4 μ g/ml of an anti-FLAGTM antibody (M2, Sigma) were added to a medium, and the medium was added to a 96-well plate at 50 μ l/well. After culturing overnight, cells were harvested, and an R-PE labeled anti-CD95 antibody (Pharmingen NJ) was used for the cells. Analysis was carried out using FACS.

Example 6: Measurement of CDC activity in anti-CD40 antibody

In the CDC assay, 2,000 Cr⁵¹-labeled target cells and a human serum-derived complement (manufactured by Sigma Co.) or rabbit serum-derived complement (Cedarlane Laboratories Limited, Ontario, Canada) at a final concentration of 5% were cultured in a round-bottom 96-well plate in a total volume of 200 μ L together with the antibody at various concentrations at 37°C in the presence of 5% CO₂ for two hours.

After culturing, the plate was centrifuged to cause the cells to precipitate, and then 50 μ L of the supernatant was transferred to a 96-well plate including a powder scintillator (LumaplateTM-96: manufactured by Packard Instrument Co., Inc.) and dried at 55°C for 1.5 hours. After confirming that the plate was dried, it was covered with a special cover (TopSealTM-A: 96-well microplates: manufactured by Packard Instrument Co., Inc.), and the γ -ray dose was measured with a scintillation counter (TopCount: manufactured by Packard Instrument Co., Inc.).

Example 7: Measurement of ADCC activity of anti-CD40 antibody

As antibody-mediated cytotoxicity, cytotoxicity to target cells in the presence of cells having killer activity such as NK cells or neutrophils and an antibody (Antibody-Dependent Cellular Cytotoxicity, hereinafter ADCC), and cytotoxicity to target cells in the presence of a complement and an antibody (Complement-Dependent Cytotoxicity, hereinafter CDC) were measured. hIgG was used as a control.

The measurement method is simply described as follows. Radioactive chromium (Cr^{51}) was incorporated into the cytoplasm of the target cells, and the amount of Cr^{51} released in the culture solution by cell death was measured as a γ -ray dose.

Specifically, 10^5 target cells of a Burkitt's lymphoma cell line Raji (ATCC CCL-86) were suspended in 15 μL of fetal calf serum (FCS). 50 μL (37 MBq/mL) of Cr^{51} -labeled sodium chromate (manufactured by PerkinElmer, Inc.: hereinafter referred to as Cr^{51}) was added to the suspension, and the cells were cultured at 37°C for one hour. Next, 10 mL of a medium was added, and the medium was discarded by centrifugation. This operation was repeated three times to remove Cr^{51} not incorporated in the cells.

In the ADCC assay, 2,000 Cr^{51} -labeled target cells and 200,000 healthy human peripheral blood mononuclear leukocytes obtained by the method described in Example 6 were cultured in a round-bottom 96-well plate (manufactured by Falcon) in a total volume of 200 μL together with the antibody at various concentrations at 37°C in the presence of 5% CO_2 for four hours.

After culturing, the plate was centrifuged to cause the cells to precipitate, and then 50 μL of the supernatant was transferred to a 96-well plate including a powder scintillator (LumaplateTM-96: manufactured by Packard Instrument Co., Inc.) and dried at 55°C for 1.5 hours. After confirming that the plate was dried, the plate was covered with a special cover (TopSealTM-A: 96-well microplates: manufactured by Packard Instrument Co., Inc.), and the γ -ray dose was measured with a scintillation counter (TopCount: manufactured by Packard Instrument Co., Inc.).

Example 8: Preparation and activity evaluation of anti-CD40 agonistic antibody P331S mutant

Gene cloning of anti-CD40 agonistic antibodies KM341-1-19 and 2105 is described in WO 02/099186. It is reported that CDC activity is reduced by converting Pro at position 331 in the IgG2 constant region into Ser. To reduce CDC activity of the KM341-1-19 antibody and the 2105 antibody, a P331S mutation was introduced into the IgG2 constant region thereof.

The human IgG1 constant region of an antibody-expressing vector N5KG1-Val Lark (IDEC Pharmaceuticals: hereinafter abbreviated to N5KG1) was substituted with human IgG2 to prepare N5KG2, and the Pro at position 331 of IgG2 was converted into Ser to prepare a mutation. cDNA cloning of the IgG2 constant region was carried out by harvesting KM341-1-19 hybridoma by centrifugation, adding TRIZOL (Gibco BRL), and extracting total RNA according to the instruction. The antibody cDNA variable region was cloned using a SMART RACE cDNA amplification kit of Clontech Laboratories, Inc. according to the attached instruction. 1st strand cDNA was prepared using 5 µg of total RNA as a template. PCR was carried out with tnIgG3Nhe: atatGCTAGCACCAAGGGCCCATCGGTCTTCCCCCTGGC (SEQ ID NO: 2)G and tnIgG2Bam: atatggatccTCATTTACCCGGAGACAGGGAGAGGCTC (SEQ ID: 3) as primer sequences using a ZtaqPCR kit (Takara) in 30 cycles each consisting of reaction at 98°C for 1 second, at 55°C for 30 seconds and at 72°C for 1 minute to amplify the gene. After the reaction, the amplified product was purified by a QIAGEN PCR purification kit, digested with NheI and BamHI, and incorporated into N5KG1 to confirm the sequence. This vector was defined as N5KG2.

N5KG2Ser (with Pro at position 331 converted into Ser) was prepared as follows. Reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG2 as a template and primers IgG3Nhe: atatGCTAGCACCAAGGGCCCATCGGTCTTCCCCCTGGCG (SEQ ID NO: 4) and G2Ser2: GTTTTCTCGATGGAGGCTGGGAGGCC (SEQ ID NO: 5). At the same time, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG2 as a template and primers IgG2Bam: atatggatccTCATTTACCCGGAGACAGGGAGAGGCTC (SEQ ID NO: 6) and G2Ser1: GGCCTCCCAGCCTCCATCGAGAAAAC (SEQ ID NO: 7). The amplified DNA

fragments were purified using a PCR purification kit, and the same amounts of the two purified DNA fragments were mixed. Thereafter, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out five times. Primers IgG3Nhe and IgG2Bam were added to the mixture, and the same reaction was carried out 15 times. The amplified DNA fragment was cleaved with NheI and BamHI, and substituted with the IgG1 constant region of the N5KG1 vector (N5KG2Ser). The fragment containing the sequence of the antibody variable region digested with BglII and NheI was incorporated into the N5KG2Ser vector.

The antibody expressed and purified by the above method was evaluated in terms of binding ability to Ramos cells (Figure 3A) and agonistic activity (Figure 3B). The fluctuation in activity due to introduction of the P331S variation was not observed.

Example 9: Measurement of CDC activity of anti-CD40 agonistic antibody 331ser mutant

CDC activity was measured by the above method. A rabbit serum-derived complement was used, and Ramos cells were used as target cells. The results confirmed that, in a KM341-1-19 antibody at an antibody concentration of 1 µg/ml, IgG2ser exhibited CDC activity significantly reduced as compared with IgG2 (Figure 4A). On the other hand, when a human supplement was used, no change was observed (Figure 4B).

Example 10: Preparation and activity measurement of agonist anti-CD40 antibody having constant region converted

Among the anti-CD antibodies described in WO 02/088186, two antibodies exhibiting strongest agonistic activity (KM341-1-19 antibody and 2105 antibody) belong to IgG2 subclass. In order to examine whether or not the IgG2 subclass is important for activation of CD40, recombinant proteins having an antibody constant region converted into IgG1, IgG3 and IgG4, respectively, were prepared, and measured in terms of binding ability to an antigen and CD95 expression enhancing activity in Ramos cells according to Examples 4 and 6. IgG1 was expressed using N5KG1, and IgG2 and IgG3 were respectively expressed using expression vectors N5KG2 and N5KG3 obtained by substituting the N5KG1 constant region

with IgG2 and IgG3, respectively. cDNA cloning of the IgG3 constant region was carried out according to the IgG2 cloning method partially modified, using an IgG3-specific primer. IgG4 was expressed using N5KG4PE (IDEC Pharmaceuticals).

The antibody protein was expressed according to Example 1. Binding activity to Ramos cells expressing human CD40 of the KM341-1-19 antibody and the 2105 antibody was not affected by converting IgG2 into IgG1, IgG3 or IgG4 (Figures 5A-1 and 5A-2). However, these antibodies were found to have CD95 expression enhancing activity in Ramos cells reduced by 10% or more (Figures 5B-1 and 5B-2). This shows that not only the structure of the variable region defining the binding region of the antibody but also the structure of the constant region of the antibody are important for the strong agonistic activity of the 2105 antibody and the KM341-1-19 antibody. Thus, in order to examine which region in the IgG2 constant region is important for agonistic activity, a domain swap mutant in which the IgG2 structure is mixed with the IgG4 structure was prepared to measure its activity. As described below, a domain swap mutant is prepared by substitution of a hinge region. In this case, the "hinge region" includes the upper hinge (from Kabat EU code 216), middle hinge (from Kabat EU code 226) and lower hinge (Kabat EU code 231), as described Ole H Brekke et. al., Immunology Today 1995, 16, 85-90. Four domain swap mutants IgG2/4 (CH1 and hinge region: IgG2, other regions: IgG4), IgG4/2/4 (hinge region: IgG2, other regions: IgG4), IgG2/4/4 (CH1: IgG2, other regions: IgG4) and IgG4/2/2 (CH1: IgG4, other regions: IgG2) were prepared respectively for the KM341-1-19 antibody and the 2105 antibody.

A vector N5KG2/4 for expressing the IgG2/4 antibody was prepared using a Ztaq PCR kit (Takara). Reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG2 as a template and primers IgG3Bam: atatgatccTCATTTACCCGGAGACAGGGAGAGGC (SEQ ID NO: 8) and 24Chi4: AGGGGTCCGGGAGATCATGAGAGTGTCCTT (SEQ ID NO: 9). At the same time, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG4 (IDEC Pharmaceuticals) as a template and primers 24Chi3: AAGGACACTCTCATGATCTCCCGGACCCCT (SEQ ID NO: 10) and linkH2: tgatcatacgtagatatcacggc (SEQ ID NO: 11). The amplified DNA fragments were purified

using a PCR purification kit, and the same amounts of the two purified DNA fragments were mixed. Thereafter, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out five times. Primers IgG3Bam and linkH2: tgatcatagtagatatacagggc (SEQ ID NO: 12) were added to the mixture, and the same reaction was carried out 15 times. The amplified DNA fragment was cleaved with NheI and BamHI, and substituted with the IgG1 constant region of the N5KG1 vector.

A vector N5KG4/2/4 for expressing IgG4/2/4 was prepared as follows. Reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG4 as a template and primers linkH: gggtacgtcctcacattcagtgatcag (SEQ ID NO: 13), G2Hin3: TTTGCGCTCAACTGTCTTGTCCACCTTGGTGTGCTGGG (SEQ ID NO: 14), linkH2: tgatcatagtagatatacagggc (SEQ ID NO: 15) and G2Hin4: ACAGTTGAGCGCAAATGTTGTGTCGAGTGCCACCG (SEQ ID NO: 16). The amplified DNA fragments were purified with a PCR purification kit, and the same amounts of the two purified DNA fragments were mixed. Thereafter, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out five times using the mixture as a template. Primers linkH and linkH2 were added to the mixture, and the same reaction was carried out 15 times. The amplified DNA fragment was cleaved with NheI and BamHI, and substituted with the IgG1 constant region of the N5KG1 vector.

A vector N5KG2/4/4 for expressing IgG2/4/4 was prepared as follows. Reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG2 as a template and primers linkH: gggtacgtcctcacattcagtgatcag (SEQ ID NO: 17) and G4CH1-2: GGTGTTGCTGGGCTTGTGATCTACGTTGCAG (SEQ ID NO: 18). At the same time, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG4 as a template and primers G4CH1-1: CTGCAACGTAGATCACAAGCCCAGCAACACC (SEQ ID NO: 19) and linkH2: tgatcatagtagatatacagggc (SEQ ID NO: 20). The amplified DNA fragments were purified using a PCR purification kit, and the same amounts of the two purified DNA fragments were mixed. Thereafter, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out five times. Primers linkH and linkH2 were added to the mixture, and

the same reaction was carried out 15 times. The amplified DNA fragment was cleaved with *NheI* and *BamHI*, and substituted with the IgG1 constant region of the N5KG1 vector.

A vector N5KG4/2/2 for expressing IgG4/2/2 was prepared as follows. Reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out using N5KG4 as a template and primers linkH: gggtacgtcctcacattcagtgatcag (SEQ ID NO: 21) and G4CH1-2: GGTGTTGCTGGGCTTGTGATCTACGTTGCAG (SEQ ID NO: 22). At the same time, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG2 as a template and primers G4CH1-1: CTGCAACGTAGATCACAAGCCCAGCAACACC (SEQ ID NO: 23) and linkH2: tgatcatagtagatcacggc (SEQ ID NO: 24). The amplified DNA fragments were purified using a PCR purification kit, and the same amounts of the two purified DNA fragments were mixed. Thereafter, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out five times. Primers linkH and linkH2 were added to the mixture, and the same reaction was carried out 15 times. The amplified DNA fragment was cleaved with *NheI* and *BamHI*, and substituted with the IgG1 constant region of the N5KG1 vector.

Binding activity of the respective four domain swap mutants of the KM341-1-19 antibody and the 2105 antibody was examined. As a result, no difference between them and the original IgG2 was observed in terms of binding ability (Figures 6A-1 and 6A-2). However, only IgG2/4/4 of both the KM341-1-19 antibody and the 2105 antibody exhibited significantly reduced agonistic activity (Figures 6B-1 and 6B-2). The results confirmed that the hinge region of IgG2 is important for agonistic activity.

Further, it was examined which sequence is important in the hinge region. The hinge region is divided into three sites, specifically, upper hinge, middle hinge and lower hinge (Ole H Brekke et al. Immunology Today 1995, 16, 85-90). IgG2-specific sequences in these regions were respectively substituted with IgG4-specific sequences. Antibodies obtained by introducing a mutation into the upper hinge (from Kabat EU code 216), middle hinge (from Kabat EU code 226) and lower hinge (from Kabat EU code 231) were respectively defined as IgG2UH4, IgG2MH4 and IgG2LH4. Their respective expression vectors were defined as

N5KG2UH4, N5KG2MH4 and N5KG2LH4. "Hinge" is defined as EU indices 216 to 230 according to Kabat et al., Sequences of proteins of immunological interest, 1991 Fifth edition.

N5KG2UH4 was prepared as follows. Reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG2 as a template and primers linkH: gggtacgtcctcacattcagtgatcag (SEQ ID NO: 25) and UH4-2: CACAACATTTggaCTCAACTcTCTTGTCCACC (SEQ ID NO: 26). At the same time, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG2 as a template and primers UH4-1: GGTGGACAAGAgAGTTGAGtccAAATGTTGTG (SEQ ID NO: 27) and linkH2: tgatcatagtagatatacggc (SEQ ID NO: 28). The amplified DNA fragments were purified using a PCR purification kit, and the same amounts of the two purified DNA fragments were mixed. Thereafter, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out five times. Primers linkH and linkH2 were added to the mixture, and the same reaction was carried out 15 times. The amplified DNA fragment was cleaved with NheI and BamHI, and substituted with the IgG1 constant region of the N5KG1 vector.

N5KG2MH4 was prepared as follows. Reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG2 as a template and primers linkH: gggtacgtcctcacattcagtgatcag (SEQ ID NO: 29) and UM4-2: GGCACGGTGGGCAtgggggaccataTTTGCGCTC (SEQ ID NO: 30). At the same time, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG2 as a template and primers UM4-1: GAGCGCAAAtatggtcccccaTGCCCACCGTGCC (SEQ ID NO: 31) and linkH2: tgatcatagtagatatacggc (SEQ ID NO: 32). The amplified DNA fragments were purified using a PCR purification kit, and the same amounts of the two purified DNA fragments were mixed. Thereafter, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out five times. Primers linkH and linkH2 were added to the mixture, and the same reaction was carried out 15 times. The amplified DNA fragment was cleaved with NheI and BamHI, and substituted with the IgG1 constant region of the N5KG1 vector.

N5KG2LH4 was prepared as follows. Reaction at 98°C for 4 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG2 as a template and primers linkH: gggtacgtcctcacattcagtgatcag (SEQ ID NO: 33) and UL4-2: GAAGACTGACGGTCCccccaggaactcTGGTGCTGGGCA (SEQ ID NO: 34). At the same time, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out 15 times using N5KG2 as a template and primers UL4-1: TGCCCAGCACCAgagttcctggggGGACCGTCAGTCTTC (SEQ ID NO: 35) and linkH2: tgatcatagctagatatcacggc (SEQ ID NO: 36). The amplified DNA fragments were purified using a PCR purification kit, and the same amounts of the two purified DNA fragments were mixed. Thereafter, reaction at 98°C for 1 second, at 60°C for 30 seconds and at 72°C for 30 seconds was carried out five times. Primers linkH and linkH2 were added to the mixture, and the same reaction was carried out 15 times. The amplified DNA fragment was cleaved with NheI and BamHI, and substituted with the IgG1 constant region of the N5KG1 vector.

The three respective domain swap mutants of the KM341-1-19 antibody and the 2105 antibody were examined to have the same binding activity to an antigen (Figures 6A-1 and 6A-2). However, IgG2UH4 and IgG2MH4 exhibited significantly reduced agonistic activity to Ramos cells (Figures 6B-1 and 6B-2). It was found from the above that the structures of upper hinge and middle hinge in the hinge region are important for IgG2 subclass-dependent agonistic activity of the anti-CD40 antibodies KM341-1-19 and 2105.

Since IgG2 subclass was found to be important for agonistic activity, antibodies of subclass other than IgG2 were converted to those of IgG2 subclass to examine whether or not the agonistic activity was enhanced. In the examination on several clones, agonistic activity of F76 could be enhanced by converting IgG1 subclass to IgG2 subclass (Figures 7A and B).

Example 11: Preparation of anti-CD40 antagonist antibody mutants

A DNA fragment containing a heavy chain and a light chain of a 4D11 antibody gene described in WO 02/088186, whose original subclass is IgG1, was digested with BglII and NheI, purified, and then integrated into N5KG4PE, N5KG4P and N5KG4 vectors (IDEC Pharmaceuticals). N5KG4PE contains point mutations S228P and L235E in the IgG4

constant region, and N5KG4P contains a point mutation S228P in the IgG4 constant region. The antibody protein was expressed and purified according to the above method. The antibody was purified according to the above method using binding to Ramos cells as an index. Change in binding activity of IgG1, IgG4, IgG4P and IgG4PE to Ramos cells was not observed (Figure 8A). Antagonistic activity of IgG1 was compared with those of various IgG4 mutants according to the above method to find that antagonistic activity of IgG1 does not differ from those of the IgG4 mutants (Figure 8B).

Example 12: Evaluation of ADCC activity and CDC activity of anti-CD40 antagonist antibody mutants

ADCC activity and CDC activity of anti-CD40 mutant antibodies were evaluated according to the above method.

When using human MNC as effector cells and CD40-expressing Daudi cells as target cells, two mutants IgG4 and IgG4PE were respectively observed to have ADCC activity significantly reduced as compared with IgG1 as the original subclass of the 4D11 antibody (Figure 9).

CDC activity of IgG1 was compared with that of IgG4P using Daudi cells as target cells. IgG4P was found to have CDC activity significantly reduced as compared with IgG1 (Figure 10).

Example 13: Effect of anti-CD40 antagonistic antibody on B cells

100 µg each of IgG1, IgG4P and IgG4PE of the 4D11 antibody was administered to the tail vein of mice having a genetic background whereby they were homozygotes for mouse endogenous disrupted CD40 and harboring a transgene of a human CD40 gene (Yasui. et al. Int. Immunol. 2002 Vol 14: 319). 24 hours after the administration, blood was collected from the orbital venous plexus. After hemolysis with 0.16 mol/L of ammonium chloride, an FITC-labeled anti-B220 antibody was added to the hemolysate, and it was analyzed using FACS. The results are shown in Figure 11. In the figure, the longitudinal axis indicates the ratio of B cells in the total lymphocytes. IgG1 reduced the ratio of B cells most, IgG4P

reduced the ratio to a lesser extent, and IgG4PE reduced the ratio to a much lesser extent. 24 hours after the administration, the spleen was removed and crushed with a slide glass to prepare a cell suspension. After hemolysis of the cell suspension, a PE-labeled anti-B220 antibody and an FITC-labeled anti-CD23, CD86 or CD95 antibody were used for the hemolysate, and it was analyzed using FACS. The results are shown in Figures 12A, B and C. In the figures, the longitudinal axis indicates the ratio of B cells expressing each surface marker in the total lymphocytes. 4D11G1 was found to achieve the same level of increase in expression of each marker as in a commercially available mouse anti-human CD40 agonistic antibody 5C3 (Pharmingen). IgG4PE achieved a smaller increase in expression of each activation surface marker as compared with IgG1 and IgG4P.

Example 14: Effect of inhibiting production of antigen-specific antibody and change in the number of B cells caused by anti-CD40 antagonistic antibody

100 µg (based on NP-CGG) of a complex of 4-hydroxy-3-nitrophenylacetyl-chicken γ-globulin conjugates (NP-CGG: distributed by Professor Hitoshi KIKUTANI, Research Institute for Microbial Diseases, Osaka University) and alum (aluminum hydroxide gel) was intraperitoneally administered to mice having a genetic background whereby they were homozygotes for mouse endogenous disrupted CD40 and harboring a transgene of a human CD40 gene (Yasui et al. Int. Immunol. 2002 Vol 14: 319) to sensitize the mice. Immediately before the antigen sensitization, 50 or 100 µg of each antibody was administered to the tail vein. 100 µg of an anti-human albumin human IgG4PE antibody was administered as a negative control. 7 and 14 days after the sensitization, blood was collected from the orbital venous plexus. The amounts of NP-specific IgG1 and IgM antibodies in the serum were measured by the ELISA method. The ELISA method was carried out as follows. 50 µl/well of NP-bound bovine serum albumin (NP-BSA: 2.5 µg/ml) was added to each well of a 96-well microplate for ELISA (Maxisorp, manufactured by Nunc A/S) and incubated at 4°C to cause NP-BSA to be adsorbed thereon. Next, the supernatant was discarded, and a blocking reagent (SuperBlock, manufactured by Pierce Biotechnology, Inc.) was added to each well and incubated at room temperature to carry out blocking. Then, each well was washed with a

72813-258

phosphate buffer (PBS-T) containing 0.1% TweenTM 20 three times. Next, each serum diluted with PBS-T containing 10% Block Ace (50 μ l/well) was added to each well, and incubated and reacted at 37°C for two hours. The microplate was washed with PBS-T three times. Then, a 1,000-fold dilution of a goat anti-mouse IgG1 antibody or IgM antibody labeled with alkaline phosphatase (Cosmo Bio, 1070-04 or 1020-04) with PBS-T containing 10% Block Ace (50 μ g/well) was added to each well, and incubated at 37°C for two hours. Next, the microplate was washed with PBS-T, and then a coloring substrate solution (50 μ l/well, Sigma 104, phosphatase substrate) was added to each well. The absorbance at a wavelength of 405 nm was measured using a microplate reader. The results are shown in Figures 13A and B. In the figures, the longitudinal axis indicates values obtained by converting a 10,000-fold dilution (in the case of IgG1) or 100-fold dilution (in the case of the IgM antibody) of serum collected from C57BL/6 mice, to which NP-CGG was injected twice, and pooled into one unit. The 4D11 antibody and the IgG4P or IgG4PE antibody of 281 inhibited production of NP-specific IgG1 and IgM antibodies equally strongly.

The change in the number of B cells in the peripheral blood and spleen in the mice used for examining the effect of inhibiting antibody production was measured according to the same method as in Example 1. The results are shown in Figures 14A and B. The 4D11 antibody and the IgG4P antibody of 281 reduced the ratio of B cells in peripheral blood significantly as compared with the IgG4PE antibody. Administration of 100 μ g of the IgG4PE antibody did not change the ratio of B cells in the spleen removed 14 days after the antigen sensitization. However, administration of IgG4P changed or tended to change the ratio.

Example 15: Effect of anti-CD40 antagonistic antibody on cynomolgus monkeys

30 mg/kg of IgG4P or IgG4PE of a 4D11 antibody was administered to the forearm cephalic vein of cynomolgus monkeys, and blood was collected from the femoral vein after a certain period of time. In the subset analysis of peripheral blood lymphocytes, an FITC-labeled anti-CD3 antibody, PE-labeled anti-CD20 antibody and APC-labeled anti-CD45 antibody were used for each cell suspension, and the ratio of positive cells was measured using

FACS to calculate the ratio of CD45 positive cells. The results are shown in Figure 15. In the figure, the longitudinal axis indicates the ratio of CD20 positive cells at each time to CD20 positive cells before antibody administration. 1 to 7 days after the antibody administration, CD20 positive cells were reduced by about 40% in individuals to which the IgG4P antibody was administered. However, 4 days after the administration, CD20 positive cells were reduced by only about 20% in individuals to which the IgG4PE antibody was administered.

The IL12 concentration in serum was measured by the ELISA method. Blood collected from the femoral vein was allowed to stand at room temperature for 20 to 60 minutes, and then centrifuged at 3,000 rpm at room temperature for 15 minutes. The IL12 concentration in the resulting serum was measured using a monkey IL12 ELISA kit (BioSource International Inc.). The results are shown in Figure 16. No increase in IL12 production by the IgG4PE antibody was observed at any blood collection point. However, maximum IL12 production by the IgG4P antibody was observed on the 4th day.

Example 16: Effect of anti-CD40 antagonistic antibody on cynomolgus monkey delayed hypersensitivity model

Nine male cynomolgus monkeys were intradermally and intramuscularly sensitized with Tetanus toxoid (TTx) (10 Lf/ml; Denka Seiken Co., Ltd.) to induce delayed hypersensitivity to TTx. At the same time, 10 minutes before the start of sensitization, 0.1 and 10 mg/kg of a 4D11G4PE antibody was intravenously administered to each three animals three times (once a week) to examine the effect of 4D11G4PE on delayed hypersensitivity. Under anesthesia by intramuscular administration of ketamine, sensitization was carried out by intradermal administration of TTx to the back (50 μ L/site $\times$ 12 sites) and intramuscular administration of TTx to the femur (0.6 mL/body), and challenge was carried out by intradermal administration of TTx to the thorax (10 μ L/site, 0 to 10 Lf/ml for each three sites) 21 days after the sensitization. 24 and 48 hours after the elicitation, skin reaction at the administration sites was observed and evaluated according to the Draize skin irritation score. The results of determining the TTx concentration in each three sites were respectively a mean value. The results are shown in Figure 17. Administration of the 4D11G4PE antibody

apparently inhibited the delayed hypersensitivity reaction observed 24 and 48 hours after the administration.

The effect of TTx on TTx-specific IgG and IgM antibody titers was examined. Blood collected from the femoral vein over time was allowed to stand at room temperature for 20 to 60 minutes, and then centrifuged at 3,000 rpm at room temperature for 15 minutes. The antibody titer in the resulting serum was measured using the ELISA method. The ELISA method was carried out as follows. 100 µl/well of TTx (0.5 Lf/ml) was added to each well of a 96-well microplate for ELISA (Maxisorp, manufactured by Nunc A/S) and incubated at 4°C to cause TTx to be adsorbed thereon. Next, the supernatant was discarded, and a blocking reagent (phosphate buffer containing 0.5% BSA) was added to each well and incubated at room temperature to carry out blocking. Then, each well was washed with a phosphate buffer (PBS-T) containing 0.05% Tween 20 three times. Next, each serum diluted with PBS-T containing 0.5% BSA (100 to 819,200-fold dilution, dilution magnification: 2; 100 µl/well) was added to each well, and incubated and reacted at room temperature for two hours. The microplate was washed with PBS-T three times. Then, a 3,000-fold dilution of a goat anti-monkey IgG antibody or IgM antibody labeled with peroxidase (Nordic Immunology) with PBS-T containing 0.5% BSA (100 µg/well) was added to each well, and incubated at room temperature for one hour. Next, the microplate was washed with PBS-T, and then a coloring substrate solution (100 µl/well, o-phenylenediamine hydrochloride + aqueous hydrogen peroxide) was added to each well. The absorbance at a wavelength of 492 nm was measured using a microplate reader. The anti-TTx antibody titer was defined as a maximum dilution magnification to make the absorbance 0.1 and more. The antibody titer was 0 when the absorbance did not reach 0.1 even at 100-fold dilution. The results are shown in Figures 18 and 19. Administration of 1 mg/kg of 4D11G4PE suppressed the TTx-specific IgG and IgM antibody titers to about 1/10. When 10 mg/kg of 4D11G4PE was administered, the antibody titers were below the detection sensitivity at any blood collection point.

Example 17: Effect of anti-CD40 antagonistic antibody on platelet thrombus formation

Blood collected from a healthy human was divided into four aliquots (each 6 ml). Control human IgG4PE, control mouse IgG2a, human anti-human CD40 IgG4PE (4D11) and mouse anti-human CD154 IgG2a (5C8) were respectively added to the fractions so that each fraction had a blood concentration of 100 µg/ml. A flat perfusion chamber (GlycoTech Corp.) and a collagen-coated Petri dish were assembled according to the attached instruction. The blood treated with various antibodies was caused to flow into the chamber at a rate that can apply a shear stress of 1,500/s to the blood for seven minutes. Thereafter, a 4% paraformaldehyde phosphate buffer was caused to flow into the chamber at a rate that can apply a shear stress of 1,500/s to the buffer for 10 minutes. The platelet aggregate formed on the Petri dish was fixed, stained with a platelet-specific PE-labeled CD41a antibody, and observed with a fluorescence microscope. The results are shown in Figures 20A and B. The blood treated with human anti-human CD40 IgG4PE (4D11) formed a platelet aggregate on the collagen-coated Petri dish, as the blood treated with the control antibodies did. However, the blood treated with mouse anti-human CD154 IgG2a did not form a platelet aggregate.

Example 18: Evaluation of stability of anti-CD40 antagonistic antibody

Constant region-modified antibodies of the 4D11 antibody were compared and examined in terms of stability. In the evaluation method, culture supernatants obtained by respectively transiently expressing G4P, G4PE, G2Ser and G4/2/4 in HEK293 cells were charged with a Protein A column (Amersham Pharmacia Biotech), eluted with a 0.1 M citrate buffer (pH 2.7), and then incubated at 37°C for 1 minute and 10 minutes. Thereafter, they are neutralized with a 50 nM phosphate buffer (pH 7.0). The oligomer content in the resulting antibody solutions was measured using a gel filtration column (Tosoh Corp.). As a result, it was found that the oligomer content increases in proportion with the incubation time, and G4/2/4 produces an oligomer easiest, G4PE second easiest, G2Ser third easiest, and G4P fourth easiest (Figure 21).

Example 19: Effect of inhibiting skin graft rejection by anti-CD40 antagonistic antibody

A graft collected from the tail of DBA/2 mice was grafted into the side dorsal thorax of C57BL/6 background mice having a genetic background whereby they were homozygotes for mouse endogenous disrupted CD40 and harboring a transgene of a human CD40 gene, and the graft was fixed with a plaster for seven days. 100 µg of a test substance 4D11G4PE or a vehicle was administered to the tail vein 0, 2, 4, 7, 9, 11 and 14 days after the skin graft, respectively. To inhibit graft rejection by NK cells, 100 µg of an anti-asialo GML antibody was intraperitoneally administered to all mice 3 days before the operation and 1, 4 and 9 days after the operation. The results are shown in Figure 22. The delay in graft rejection was observed to be significant in the 4D11G4PE-administered group as compared with the vehicle-administered group.

Example 20: Analysis of CD40 expression in human tumor cell lines

Expression of CD40 in a Burkitt's lymphoma cell line Ramos, bladder cancer cell line T24 (ATCC, HTB-4), pancreatic cancer cell line Hs 766T (ATCC, HTB-134) and Capan-2 (ATCC, HTB-80) was confirmed by FACS analysis using 341G2Ser.

T24, Hs 766T and Capan-2 were digested with trypsin and harvested, and Ramos was harvested as is. The cell lines were washed with PBS, and then re-suspended in a staining buffer containing 1 µg/ml of 341G2Ser. The staining buffer was prepared by adding 0.05 mM EDTA, 0.05% sodium azide and 5% immobilized bovine serum to PBS. After incubation at 4°C for 15 minutes, the cells were washed with the staining buffer twice, and re-suspended in a 1:250 dilution of PE-bound goat anti-human IgG (γ) (Southern Biotechnology Associates, Inc) with the staining buffer. After incubation at 4°C for 15 minutes, the cells were washed with the staining buffer twice, and analyzed with FACSCalibur (manufactured by BD Biosciences). The same amount of a human anti-2,4-dinitrophenol (DNP) antibody was used as a negative control. The analysis was carried out using Cellquest (manufactured by BD Biosciences) as data analysis software to calculate the mean fluorescence intensity.

As a result, Ramos, T24, Hs766T and Capan-2 had a mean fluorescence intensity obviously higher than that of the negative control when stained with 341G2Ser, and thus expression of CD40 was confirmed.

Example 21: Effect of anti-CD40 agonistic antibody on human tumor cell lines

2.5×10^3 Ramos cells, 2.5×10^2 T24 cells, 5×10^3 Hs766T cells and 5×10^3 Capan-2 cells were respectively suspended in a medium to make the total volume of 100 μ L in a flat bottom 96-well plate (manufactured by Falcon). Ramos and Hs766T, Capan-2 and T24 were cultured for 66 hours, 90 hours and 114 hours, respectively, together with 341G2Ser at a concentration of 1 ng/ml to 1,000 ng/ml at 37°C in the presence of 5% CO₂. 10 μ L (3.7 MBq/mL) of ³H-labeled thymidine (manufactured by Amersham Biosciences) was added and cultured at 37°C in the presence of 5% CO₂ for six hours. Ramos cells were harvested on Printed Filtermat A (manufactured by PerkinElmer, Inc.) using a 96 micro cell harvester (manufactured by Skatron Instruments, Inc.), and covered with a sample bag (manufactured by PerkinElmer, Inc.). 12 mL of Betaplate Scint (manufactured by PerkinElmer, Inc.) was added, and the β -ray dose was measured with a liquid scintillation counter (Pharmacia 1205 Betaplate: manufactured by Pharmacia Corp.). Hs 766T cells, T24 cells and Capan-2 cells were respectively harvested on Unifilter (manufactured by PerkinElmer, Inc.) using a harvester (manufactured by PerkinElmer, Inc.). A special seal was attached to the back of each filter, and 20 μ L/well of MicroScint 20 (manufactured by PerkinElmer, Inc.) was added thereto. The β -ray dose was measured with a scintillation counter (TopCount: manufactured by Packard Instrument Co., Inc.). Data were expressed as cell survival rates (%) obtained by dividing a mean of triplicate measurements obtained in three independent tests by a value of non-treatment control.

As a result, the cell survival rates were reduced in all cell lines depending on the 341G2Ser concentration (Table 1). When adding 100 ng/ml of 341G2Ser, the Ramos cell survival rate was 58%, the T24 cell survival rate was 22%, the Hs 766T cell survival rate was 15%, and the Capan-2 cell survival rate was 77%. 341G2Ser was found to have activity of inhibiting growth of Ramos cells, T24 cells, Hs 766T cells and Capan-2 cells.

Table 1

Cell survival
rate

Cell line	341G2Ser concentration (ng/ml)			
	1	10	100	1000
Ramos	98.49%	81.68%	57.77%	55.26%
T24	97.94%	50.72%	21.97%	25.35%
Hs 766T	34.67%	21.50%	14.67%	15.18%
Capan-2	100.94%	85.34%	76.76%	72.89%

Example 22: Effect of anti-CD40 agonistic antibody on mouse cancer-bearing model

(1) Ramos cells

Six-week-old female Balb/c nude mice (purchased from CLEA Japan, Inc.) were irradiated with 3Gy radiation, and 2×10^7 cells/mouse of Ramos cells were subcutaneously grafted into the back thereof. 16 days after the graft, the size of tumors that took there was measured. Cancer-bearing mice having a tumor size of 50 to 170 mm³ were classified into groups each consisting of five mice. 100 µg/mouse of 341G2Ser (a solution in 200 µl of PBS containing 1% nude mouse serum) was intravenously administered to the cancer-bearing mice once on the 16th day, and the tumor size was measured until the 47th day. A human anti-human serum albumin (HAS) antibody was used as a negative control.

(2) T24 cells

A T24 cell mass that had undergone subcutaneous passage in the back of nude mice three times were removed, and subcutaneously grafted into the back of six-week-old female Balb/c nude mice (purchased from CLEA Japan, Inc.). The tumor cell mass to be grafted is appropriately about 3 mm square. 10 days after the graft, the size of engrafted tumors was measured. Cancer-bearing mice having a tumor size of 80 to 200 mm³ were classified into groups each consisting of five mice. 100 µg/mouse of 341G2Ser (a solution in 200 µl of PBS containing 1% nude mouse serum) was intravenously administered to the cancer-bearing mice once on the 10th day, and the tumor size was measured until the 29th day. The same amount of a human anti-DNP antibody was used as a negative control.

(3) Hs 766T cells

7×10^5 cells/mouse of Hs 766T cells were subcutaneously grafted into the back of eight-week-old female Balb/c nude mice (purchased from CLEA Japan, Inc.). 16 days after the graft, the size of engrafted tumors was measured. Cancer-bearing mice having a tumor size of 50 to 140 mm³ were classified into groups each consisting of five mice. 100 µg/mouse of 341G2Ser (a solution in 200 µl of PBS containing 1% nude mouse serum) was intravenously administered to the cancer-bearing mice once on the 16th day, and the tumor size was measured until the 32nd day. The same amount of a human anti-DNP antibody was used as a negative control.

(4) Capan-2 cells

2×10^6 cells/mouse of Capan-2 cells were subcutaneously grafted into the back of six-week-old female Balb/c nude mice (purchased from CLEA Japan, Inc.). 13 days after the graft, the size of engrafted tumors was measured. Cancer-bearing mice having a tumor size of 30 to 130 mm³ were classified into groups each consisting of five mice. 10 or 100 µg/mouse of 341G2Ser (a solution in 200 µl of PBS containing 1% nude mouse serum) was intravenously administered to the cancer-bearing mice twice a week from the 13th day, and the tumor size was measured until the 34th day. A human polyclonal antibody (hIgG) (manufactured by Sigma Co.) was used as a negative control.

The tumor growth inhibition ratio (TGIR) was calculated from the following formula.

$$100 - \left[\frac{\text{mean tumor volume of 341G2Ser-administered group on the last measurement day} - \text{mean tumor volume of 341G2Ser-administered group on the day of start of antibody administration}}{\text{mean tumor volume of negative control-administered group on the last measurement day} - \text{mean tumor volume of negative control-administered group on the day of start of antibody administration}} \right] \times 100$$

As a result, TGIR exceeded 100% in the T24, Hs766T and Capan-2 cancer-bearing mice, and a decrease in the tumor volume was observed in the mice. On the other hand, TGIR was 73.4% in the Ramos cancer-bearing mice, and an increase in the tumor volume was considerably suppressed in the mice (Table 2). Figures 23 to 26 respectively show the

72813-258

change in the tumor volume of cancer-bearing mice to which Ramos cells, T24 cells, Hs 766T cells and Capan-2 cells were respectively engrafted.

Table 2

Tumor growth
inhibition ratio

Cell line	Amount of 341G2Ser administered	
	10 µg/head	100 µg/head
Ramos	-	73.40%
T24	-	109.05%
Hs 766T	-	108.55%
Capan-2	103.49%	119.20%

The inhibition ratio in each cell line is a value on the last measurement day.

Industrial Applicability

As shown in the Examples, the anti-CD40 antibody of the present invention having a constant region into which a mutation is introduced and an anti-CD40 antibody in which a part of the structure of the subclass is substituted with that of another subclass have reduced ADCC activity and CDC activity, while maintaining its activity. Accordingly, the antibody of the present invention has the reduced cytotoxicity to CD40-expressing cells when administered to a subject as a therapeutic antibody, and thus can be used with safety.

Sequence Listing Free Text

SEQ ID NOS: 2 to 36: Synthetic DNAs

SEQ ID NOS: 49 to 130: Synthetic peptides

SEQUENCE LISTING IN ELECTRONIC FORM

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 72813-258 Seq 05-11-09 v1.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

The sequences in the sequence listing in electronic form are reproduced in the following table.

SEQUENCE TABLE

<110> KIRIN BEER KABUSIKI KAISHA

<120> A mutant of anti CD40 antibody

<130> PH-2356-PCT

<140>

<141>

<150> JP 2003-431408

<151> 2003-12-25

<160> 142

<170> PatentIn Ver. 2.1

<210> 1

<211> 175

<212> PRT

<213> Homo sapiens

<400> 1

Glu	Pro	Pro	Thr	Ala	Cys	Arg	Glu	Lys	Gln	Tyr	Leu	Ile	Asn	Ser	Gln	1	5	10	15
Cys	Cys	Ser	Leu	Cys	Gln	Pro	Gly	Gln	Lys	Leu	Val	Ser	Asp	Cys	Thr	20	25	30	
Glu	Phe	Thr	Glu	Thr	Glu	Cys	Leu	Pro	Cys	Gly	Glu	Ser	Glu	Phe	Leu	35	40	45	
Asp	Thr	Trp	Asn	Arg	Glu	Thr	His	Cys	His	Gln	His	Lys	Tyr	Cys	Asp	50	55	60	
Pro	Asn	Leu	Gly	Leu	Arg	Val	Gln	Gln	Lys	Gly	Thr	Ser	Glu	Thr	Asp	65	70	75	80
Thr	Ile	Cys	Thr	Cys	Glu	Glu	Gly	Trp	His	Cys	Thr	Ser	Glu	Ala	Cys	85	90	95	
Glu	Ser	Cys	Val	Leu	His	Arg	Ser	Cys	Ser	Pro	Gly	Phe	Gly	Val	Lys	100	105	110	
Gln	Ile	Ala	Thr	Gly	Val	Ser	Asp	Thr	Ile	Cys	Glu	Pro	Cys	Pro	Val	115	120	125	
Gly	Phe	Phe	Ser	Asn	Val	Ser	Ser	Ala	Phe	Glu	Lys	Cys	His	Pro	Trp	130	135	140	
Thr	Ser	Cys	Glu	Thr	Lys	Asp	Leu	Val	Val	Gln	Gln	Ala	Gly	Thr	Asn	145	150	155	160
Lys	Thr	Asp	Val	Val	Cys	Gly	Pro	Gln	Asp	Arg	Leu	Arg	Ala	Leu		165	170	175	

<210> 2
 <211> 39
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 2
 atatgctagc accaagggcc catcgggtctt ccccttggc 39

<210> 3
 <211> 38
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 3
 atatggatcc tcattttaccc ggagacaggg agaggctc 38

<210> 4
 <211> 40
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 4
 atatgctagc accaagggcc catcgggtctt ccccttggcg 40

<210> 5
 <211> 26
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 5
 gttttctcga tggaggctgg gaggcc 26

<210> 6
 <211> 38
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 6
 atatggatcc tcattttaccc ggagacaggg agaggctc 38

<210> 7
 <211> 26
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 7
 ggccctcccag cctccatcga gaaaac 26

<210> 8
 <211> 36
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 8
 atatggatcc tcatttaccc ggagacaggg agaggg 36

<210> 9
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 9
 aggggtccgg gagatcatga gagtgtcctt 30

<210> 10
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 10
 aaggacactc tcatgatctc ccggaccctt 30

<210> 11
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 11
 tgatcatacg tagatatcac ggc 23

<210> 12
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic DNA

<400> 12
tgatcatacg tagatatcac ggc 23

<210> 13
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic DNA

<400> 13
gggtacgtcc tcacattcag tgatcag 27

<210> 14
<211> 39
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic DNA

<400> 14
tttgcgctca actgtcttgt ccaccttggt gttgctggg 39

<210> 15
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic DNA

<400> 15
tgatcatacg tagatatcac ggc 23

<210> 16
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic DNA

<400> 16
acagttgagc gcaaattgtg tgtcgagtgcc ccaccg 36

<210> 17
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic DNA

<400> 17
gggtacgtcc tcacattcag tgatcag 27

<210> 18
 <211> 31
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 18
 ggtgttgctg ggcttgtgat ctacgttgca g 31

<210> 19
 <211> 31
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 19
 ctgcaacgta gatcacaagc ccagcaacac c 31

<210> 20
 <211> 23
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 20
 tgatcatacg tagatatcac ggc 23

<210> 21
 <211> 27
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 21
 gggtagctcc tcacattcag tgatcag 27

<210> 22
 <211> 31
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 22
 ggtgttgctg ggcttgtgat ctacgttgca g 31

<210> 23
 <211> 31
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 23
 ctgcaacgta gatcacaagc ccagcaacac c 31

 <210> 24
 <211> 23
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 24
 tgatcatatcg tagatatcac ggc 23

 <210> 25
 <211> 27
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 25
 gggtagctcc tcacattcag tgatcag 27

 <210> 26
 <211> 32
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 26
 cacaacattt ggactcaact ctcttgcca cc 32

 <210> 27
 <211> 32
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

 <400> 27
 ggtggacaag agagttgagt ccaaattgttg tg 32

 <210> 28
 <211> 23
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic DNA

<400> 28
 tgatcatacg tagatatcac ggc 23

<210> 29
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic DNA

<400> 29
 gggtacgtcc tcacattcag tgatcag 27

<210> 30
 <211> 34
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic DNA

<400> 30
 ggcacggtgg gcatggggga ccatatttgc gctc 34

<210> 31
 <211> 34
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic DNA

<400> 31
 gagcgcaa at atggtccccc atgcccaccg tgcc 34

<210> 32
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic DNA

<400> 32
 tgatcatacg tagatatcac ggc 23

<210> 33
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic DNA

<400> 33
 gggtacgtcc tcacattcag tgatcag 27

<210> 34
 <211> 39
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic DNA

<400> 34

gaagactgac ggtcccccca ggaactctgg tgctgggca 39

<210> 35
 <211> 39
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic DNA

<400> 35

tgcccagcac cagagttcct ggggggaccg tcagtcttc 39

<210> 36
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic DNA

<400> 36

tgatcatacg tagatatcac ggc 23

<210> 37
 <211> 1480
 <212> DNA
 <213> Homo sapiens

<400> 37

```

gtcgacgctg aattctggct gaccagggca gccaccagag ctccagacaa tgtctgtctc 60
cttcctcatc ttccctgccg tgctgggcct cccatggggg gtccctgtcac aggtccaact 120
gcagcagtca ggtccaggac tgggtgaagcc ctccgagacc ctctcactca cctgtgccat 180
ctccgggggac agtgtctcta gcaacagtgc tacttggaac tggatcaggc agtccccatc 240
gagagacctt gagtggctgg gaaggacata ctacagggtcc aagtgggtatc gtgattatgt 300
aggatctgtg aaaagtcgaa taatcatcaa cccagacaca tccaacaacc agttctccct 360
gcagctgaac tctgtgactc ccgaggacac ggctatatat tactgtacaa gagcacagtg 420
gctgggaggg gattacccct actactacag tatggacgtc tggggccaag ggaccacggt 480
caccgtctct tcagcctcca ccaagggccc atcggctctc ccctgggcgc cctgctccag 540
gagcacctcc gagagcacag cggccctggg ctgcctggtc aaggactact tccccgaacc 600
ggtgacggtg tcgtggaact caggcgtctc gaccagcggc gtgcacacct tcccagctgt 660
cctacagtcc tcaggactct actccctcag cagcgtgggtg accgtgccct ccagcaactt 720
cggcacccag acctacacct gcaacgtaga tcacaagccc agcaacacca aggtggacaa 780
gacagttgag cgcaaattgt gtgtcgagtg cccaccgtgc ccagcaccac ctgtggcagg 840
accgtcagtc ttccctcttc ccccaaaacc caaggacacc ctcatgatct cccggacccc 900
tgaggtcacg tgcgtgggtg tggacgtgag ccacgaagac cccgaggtcc agttcaactg 960
gtacgtggac ggcgtggagg tgcataatgc caagacaaag ccacgggagg agcagttcaa 1020
cagcacgttc cgtgtggtca gcgtcctcac cgttgtgcac caggactggc tgaacggcaa 1080
ggagtacaag tgcaaggtct ccaacaaagg cctcccagcc cccatcgaga aaaccatctc 1140
caaaaccaaa gggcagcccc gagaaccaca ggtgtacacc ctgcccccat cccgggagga 1200
gatgaccaag aaccaggtca gcctgacctg cctgggtcaa ggcttctacc ccagcgacat 1260
cgccgtggag tgggagagca atgggcagcc ggagaacaac tacaagacca cacctcccat 1320

```

```

gctggactca gacggctcct tcttcctcta cagcaagctc accgtggaca agagcaggtg 1380
gcagcagggg aacgtcttct catgctccgt gatgcatgag gctctgcaca accactacac 1440
gcagaagagc ctctccctgt ctccgggtaa atgaggatcc 1480

```

```

<210> 38
<211> 474
<212> PRT
<213> Homo sapiens

```

```

<400> 38
Met Ser Val Ser Phe Leu Ile Phe Leu Pro Val Leu Gly Leu Pro Trp
 1      5      10      15
Gly Val Leu Ser Gln Val Gln Leu Gln Gln Ser Gly Pro Gly Leu Val
      20      25      30
Lys Pro Ser Gln Thr Leu Ser Leu Thr Cys Ala Ile Ser Gly Asp Ser
      35      40      45
Val Ser Ser Asn Ser Ala Thr Trp Asn Trp Ile Arg Gln Ser Pro Ser
      50      55      60
Arg Asp Leu Glu Trp Leu Gly Arg Thr Tyr Tyr Arg Ser Lys Trp Tyr
      65      70      75      80
Arg Asp Tyr Val Gly Ser Val Lys Ser Arg Ile Ile Ile Asn Pro Asp
      85      90      95
Thr Ser Asn Asn Gln Phe Ser Leu Gln Leu Asn Ser Val Thr Pro Glu
      100     105     110
Asp Thr Ala Ile Tyr Tyr Cys Thr Arg Ala Gln Trp Leu Gly Gly Asp
      115     120     125
Tyr Pro Tyr Tyr Tyr Ser Met Asp Val Trp Gly Gln Gly Thr Thr Val
      130     135     140
Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala
      145     150     155     160
Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu
      165     170     175
Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly
      180     185     190
Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser
      195     200     205
Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Asn Phe
      210     215     220
Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr
      225     230     235     240
Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys Val Glu Cys Pro Pro
      245     250     255
Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro
      260     265     270
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
      275     280     285
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp
      290     295     300
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
      305     310     315     320
Glu Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val
      325     330     335
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
      340     345     350
Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly
      355     360     365
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
      370     375     380
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
      385     390     395     400
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
      405     410     415

```

```

Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe
      420                      425                      430
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
      435                      440                      445
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
      450                      455                      460
Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
465                      470

```

```

<210> 39
<211> 406
<212> DNA
<213> Homo sapiens

```

```

<400> 39
actgctcagt taggacccag aggggaaccat ggaagcccca gctcagcttc tcttcctcct 60
gctactctgg ctcccagata ccaccggaga aattgtgttg acacagtctc cagccaccct 120
gtctttgtct ccaggggaaa gagccaccct ctctgcaggg gccagtcaga gtgttagcag 180
ctacttagcc tggtagcaac agaaacctgg ccaggctccc aggctcctca tctatgatgc 240
atccaacagg gccactggca tcccagccag gttcagtggc agtgggtctg ggacagactt 300
cactctcacc atcagcagcc tagagcctga agattttgca gtttattact gtcagcagcg 360
tagcaacact ttcggccctg ggaccaaagt ggatatcaaa cgtacg 406

```

```

<210> 40
<211> 126
<212> PRT
<213> Homo sapiens

```

```

<400> 40
Met Glu Ala Pro Ala Gln Leu Leu Phe Leu Leu Leu Leu Trp Leu Pro
  1          5          10          15
Asp Thr Thr Gly Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser
      20          25          30
Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser
      35          40          45
Val Ser Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro
      50          55          60
Arg Leu Leu Ile Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala
      65          70          75          80
Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
      85          90          95
Ser Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser
      100          105          110
Asn Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys Arg Thr
      115          120          125

```

```

<210> 41
<211> 508
<212> DNA
<213> Homo sapiens

```

```

<400> 41
ctgaacacag acccgctcgac tcccaggtgt ttccattcag tgatcagcac tgaacacaga 60
ggactcacca tggagttggg actgagctgg attttccttt tggctatattt aaaaggtgtc 120
cagtgtgaag tgcagctggg ggagtctggg ggaggcttgg tacagcctgg caggtccttg 180
agactctcct gtgcagcctc tggattcacc tttgatgatt atgcatgca ctgggtccgg 240
caagctccag ggaagggcct ggagtgggtc tcagggtatta gttggaatag tggtagcttg 300
gtgcatgcgg actctgtgaa gggccgattc accatctcca gagacaacgc caagaactcc 360
ctgtatctgc aaatgaacag tctgagagct gaggacacgg ccttggtatta ctgtgcaaga 420
gataggctat ttcggggagt taggtactac ggtatggacg tctggggcca agggaccacg 480
gtcaccgtct cctcagctag caccaagg 508

```

<210> 42
 <211> 146
 <212> PRT
 <213> Homo sapiens

<400> 42
 Met Glu Leu Gly Leu Ser Trp Ile Phe Leu Leu Ala Ile Leu Lys Gly
 1 5 10 15
 Val Gln Cys Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln
 20 25 30
 Pro Gly Arg Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Asp Asp Tyr Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Val Ser Gly Ile Ser Trp Asn Ser Gly Ser Leu Val His Ala
 65 70 75 80
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95
 Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Leu
 100 105 110
 Tyr Tyr Cys Ala Arg Asp Arg Leu Phe Arg Gly Val Arg Tyr Tyr Gly
 115 120 125
 Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser
 130 135 140
 Thr Lys
 145

<210> 43
 <211> 414
 <212> DNA
 <213> Homo sapiens

<400> 43
 ctgctcagtt aggacccaga gggaaccatg gaagccccag ctcagcttct cttcctcctg 60
 ctactctggc tcccagatac caccggagaa attgtgttga cacagtctcc agccaccctg 120
 tctttgtctc caggggaaag agccaccctc tcctgcaggg ccagtcagag tgtagcagc 180
 tacttagcct ggtaccaaca gaaacctggc caggctccca ggctcctcat ctatgatgca 240
 tccaacaggg ccactggcat cccagccagg ttcagtggca gtgggtctgg gacagacttc 300
 actctcacca tcagcagcct agagcctgaa gattttgcag tttattactg tcagcagcgt 360
 agccactggc tcactttcgg cgggggggacc aaggtggaga tcaaacgtac ggtg 414

<210> 44
 <211> 129
 <212> PRT
 <213> Homo sapiens

<400> 44
 Met Glu Ala Pro Ala Gln Leu Leu Phe Leu Leu Leu Leu Trp Leu Pro
 1 5 10 15
 Asp Thr Thr Gly Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser
 20 25 30
 Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser
 35 40 45
 Val Ser Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro
 50 55 60
 Arg Leu Leu Ile Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala
 65 70 75 80
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
 85 90 95
 Ser Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser
 100 105 110

His Trp Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr
 115 120 125
 Val

<210> 45
 <211> 462
 <212> DNA
 <213> Homo sapiens

<400> 45
 atatgtcgac gagtcacgga tctcatgtgc aagaaaatga agcacctgtg gttcttcctc 60
 ctgctggtgg cggctcccag atgggtcctg tcccagctgc agctgcagga gtcgggcca 120
 ggactactga agccttcgga gaccctgtcc ctcacctgca ctgtctctgg cggctccatc 180
 agcagtcctg gttactacgg gggctggatc cgccagcccc caggaagg gctggagtgg 240
 attgggagta tctataaaag tgggagcacc taccacaacc cgccctcaa gagtcgagtc 300
 accatatccg tagacacgtc caagaaccag ttctccctga agctgagctc tgtgaccgcc 360
 gcagacacgg ctgtgtatta ctgtacgaga cctgtagtac gatattttgg gtggttcgac 420
 ccctggggcc aggaaccct ggtcacccgc tcctcagcta gc 462

<210> 46
 <211> 149
 <212> PRT
 <213> Homo sapiens

<400> 46
 Met Asp Leu Met Cys Lys Lys Met Lys His Leu Trp Phe Phe Leu Leu
 1 5 10 15
 Leu Val Ala Ala Pro Arg Trp Val Leu Ser Gln Leu Gln Leu Gln Glu
 20 25 30
 Ser Gly Pro Gly Leu Leu Lys Pro Ser Glu Thr Leu Ser Leu Thr Cys
 35 40 45
 Thr Val Ser Gly Gly Ser Ile Ser Ser Pro Gly Tyr Tyr Gly Gly Trp
 50 55 60
 Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile Gly Ser Ile Tyr
 65 70 75 80
 Lys Ser Gly Ser Thr Tyr His Asn Pro Ser Leu Lys Ser Arg Val Thr
 85 90 95
 Ile Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu Lys Leu Ser Ser
 100 105 110
 Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Thr Arg Pro Val Val
 115 120 125
 Arg Tyr Phe Gly Trp Phe Asp Pro Trp Gly Gln Gly Thr Leu Val Thr
 130 135 140
 Val Ser Ser Ala Ser
 145

<210> 47
 <211> 448
 <212> DNA
 <213> Homo sapiens

<400> 47
 agatcttaag caagtgtaac aactcagagt acgcggggag acccactcag gacacagcat 60
 ggacatgagg gtccccgctc agctcctggg gcttctgctg ctctggctcc caggtgccag 120
 atgtgccatc cagttgaccc agtctccatc ctccctgtct gcactctgtag gagacagagt 180
 caccatcact tgccgggcaa gtcagggcat tagcagtgtt ttagcctggg atcagcagaa 240
 accagggaaa gctcctaagc tcctgatcta tgatgcctcc aatttggaag gtgggggtccc 300
 atcaagggttc agcggcagtg gatctgggac agatttcact ctaccatca gcagcctgca 360
 gcctgaagat tttgcaactt attactgtca acagtttaat agttaccgga cgttcggcca 420
 agggaccaag gtggaaatca aacgtacg 448

<210> 48
 <211> 130
 <212> PRT
 <213> Homo sapiens

<400> 48
 Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu Trp
 1 5 10 15
 Leu Pro Gly Ala Arg Cys Ala Ile Gln Leu Thr Gln Ser Pro Ser Ser
 20 25 30
 Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
 35 40 45
 Gln Gly Ile Ser Ser Ala Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys
 50 55 60
 Ala Pro Lys Leu Leu Ile Tyr Asp Ala Ser Asn Leu Glu Ser Gly Val
 65 70 75 80
 Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 85 90 95
 Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
 100 105 110
 Phe Asn Ser Tyr Pro Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 115 120 125
 Arg Thr
 130

<210> 49
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 49
 Glu Pro Pro Thr Ala Cys Arg Glu Lys Gln Tyr Leu Ile
 1 5 10

<210> 50
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 50
 Pro Thr Ala Cys Arg Glu Lys Gln Tyr Leu Ile Asn Ser
 1 5 10

<210> 51
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 51

Ala Cys Arg Glu Lys Gln Tyr Leu Ile Asn Ser Gln Cys
1 5 10

<210> 52

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic
peptide

<400> 52

Arg Glu Lys Gln Tyr Leu Ile Asn Ser Gln Cys Cys Ser
1 5 10

<210> 53

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic
peptide

<400> 53

Lys Gln Tyr Leu Ile Asn Ser Gln Cys Cys Ser Leu Cys
1 5 10

<210> 54

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic
peptide

<400> 54

Tyr Leu Ile Asn Ser Gln Cys Cys Ser Leu Cys Gln Pro
1 5 10

<210> 55

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic
peptide

<400> 55

Ile Asn Ser Gln Cys Cys Ser Leu Cys Gln Pro Gly Gln
1 5 10

<210> 56

<211> 13

<212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 56

Ser	Gln	Cys	Cys	Ser	Leu	Cys	Gln	Pro	Gly	Gln	Lys	Leu
1				5					10			

<210> 57

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 57

Cys	Cys	Ser	Leu	Cys	Gln	Pro	Gly	Gln	Lys	Leu	Val	Ser
1				5				10				

<210> 58

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 58

Ser	Leu	Cys	Gln	Pro	Gly	Gln	Lys	Leu	Val	Ser	Asp	Cys
1				5				10				

<210> 59

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 59

Cys	Gln	Pro	Gly	Gln	Lys	Leu	Val	Ser	Asp	Cys	Thr	Glu
1				5				10				

<210> 60

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 60

Pro	Gly	Gln	Lys	Leu	Val	Ser	Asp	Cys	Thr	Glu	Phe	Thr
1				5					10			

<210> 61

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 61

Gln	Lys	Leu	Val	Ser	Asp	Cys	Thr	Glu	Phe	Thr	Glu	Thr
1				5				10				

<210> 62

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 62

Leu	Val	Ser	Asp	Cys	Thr	Glu	Phe	Thr	Glu	Thr	Glu	Cys
1				5				10				

<210> 63

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 63

Ser	Asp	Cys	Thr	Glu	Phe	Thr	Glu	Thr	Glu	Cys	Leu	Pro
1				5				10				

<210> 64

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 64

Cys	Thr	Glu	Phe	Thr	Glu	Thr	Glu	Cys	Leu	Pro	Cys	Gly
1				5				10				

<210> 65

<211> 13

<212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 65

Glu Phe Thr Glu Thr Glu Cys Leu Pro Cys Gly Glu Ser
 1 5 10

<210> 66

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 66

Thr Glu Thr Glu Cys Leu Pro Cys Gly Glu Ser Glu Phe
 1 5 10

<210> 67

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 67

Thr Glu Cys Leu Pro Cys Gly Glu Ser Glu Phe Leu Asp
 1 5 10

<210> 68

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 68

Cys Leu Pro Cys Gly Glu Ser Glu Phe Leu Asp Thr Trp
 1 5 10

<210> 69

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 69
 Pro Cys Gly Glu Ser Glu Phe Leu Asp Thr Trp Asn Arg
 1 5 10

<210> 70
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 70
 Gly Glu Ser Glu Phe Leu Asp Thr Trp Asn Arg Glu Thr
 1 5 10

<210> 71
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 71
 Ser Glu Phe Leu Asp Thr Trp Asn Arg Glu Thr His Cys
 1 5 10

<210> 72
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 72
 Phe Leu Asp Thr Trp Asn Arg Glu Thr His Cys His Gln
 1 5 10

<210> 73
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 73
 Asp Thr Trp Asn Arg Glu Thr His Cys His Gln His Lys
 1 5 10

<210> 74
 <211> 13

<212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 74

Trp	Asn	Arg	Glu	Thr	His	Cys	His	Gln	His	Lys	Tyr	Cys
1					5				10			

<210> 75

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 75

Arg	Glu	Thr	His	Cys	His	Gln	His	Lys	Tyr	Cys	Asp	Pro
1				5				10				

<210> 76

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 76

Thr	His	Cys	His	Gln	His	Lys	Tyr	Cys	Asp	Pro	Asn	Leu
1				5				10				

<210> 77

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 77

Cys	His	Gln	His	Lys	Tyr	Cys	Asp	Pro	Asn	Leu	Gly	Leu
1				5				10				

<210> 78

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 78
 Gln His Lys Tyr Cys Asp Pro Asn Leu Gly Leu Arg Val
 1 5 10

<210> 79
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 79
 Lys Tyr Cys Asp Pro Asn Leu Gly Leu Arg Val Gln Gln
 1 5 10

<210> 80
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 80
 Cys Asp Pro Asn Leu Gly Leu Arg Val Gln Gln Lys Gly
 1 5 10

<210> 81
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 81
 Pro Asn Leu Gly Leu Arg Val Gln Gln Lys Gly Thr Ser
 1 5 10

<210> 82
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 82
 Leu Gly Leu Arg Val Gln Gln Lys Gly Thr Ser Glu Thr
 1 5 10

<210> 83
 <211> 13

<212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 83

Leu Arg Val Gln Gln Lys Gly Thr Ser Glu Thr Asp Thr
 1 5 10

<210> 84

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 84

Val Gln Gln Lys Gly Thr Ser Glu Thr Asp Thr Ile Cys
 1 5 10

<210> 85

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 85

Gln Lys Gly Thr Ser Glu Thr Asp Thr Ile Cys Thr Cys
 1 5 10

<210> 86

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 86

Gly Thr Ser Glu Thr Asp Thr Ile Cys Thr Cys Glu Glu
 1 5 10

<210> 87

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 87

Ser Glu Thr Asp Thr Ile Cys Thr Cys Glu Glu Gly Trp
 1 5 10

<210> 88

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 88

Thr Asp Thr Ile Cys Thr Cys Glu Glu Gly Trp His Cys
 1 5 10

<210> 89

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 89

Thr Ile Cys Thr Cys Glu Glu Gly Trp His Cys Thr Ser
 1 5 10

<210> 90

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 90

Cys Thr Cys Glu Glu Gly Trp His Cys Thr Ser Glu Ala
 1 5 10

<210> 91

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 91

Cys Glu Glu Gly Trp His Cys Thr Ser Glu Ala Cys Glu
 1 5 10

<210> 92

<211> 13

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 92
 Glu Gly Trp His Cys Thr Ser Glu Ala Cys Glu Ser Cys
 1 5 10

<210> 93
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 93
 Trp His Cys Thr Ser Glu Ala Cys Glu Ser Cys Val Leu
 1 5 10

<210> 94
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 94
 Cys Thr Ser Glu Ala Cys Glu Ser Cys Val Leu His Arg
 1 5 10

<210> 95
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 95
 Ser Glu Ala Cys Glu Ser Cys Val Leu His Arg Ser Cys
 1 5 10

<210> 96
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 96
 Ala Cys Glu Ser Cys Val Leu His Arg Ser Cys Ser Pro
 1 5 10

<210> 97
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 97
 Glu Ser Cys Val Leu His Arg Ser Cys Ser Pro Gly Phe
 1 5 10

<210> 98
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 98
 Cys Val Leu His Arg Ser Cys Ser Pro Gly Phe Gly Val
 1 5 10

<210> 99
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 99
 Leu His Arg Ser Cys Ser Pro Gly Phe Gly Val Lys Gln
 1 5 10

<210> 100
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 100
 Arg Ser Cys Ser Pro Gly Phe Gly Val Lys Gln Ile Ala
 1 5 10

<210> 101
 <211> 13

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 101
 Cys Ser Pro Gly Phe Gly Val Lys Gln Ile Ala Thr Gly
 1 5 10

<210> 102
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 102
 Pro Gly Phe Gly Val Lys Gln Ile Ala Thr Gly Val Ser
 1 5 10

<210> 103
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 103
 Phe Gly Val Lys Gln Ile Ala Thr Gly Val Ser Asp Thr
 1 5 10

<210> 104
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 104
 Val Lys Gln Ile Ala Thr Gly Val Ser Asp Thr Ile Cys
 1 5 10

<210> 105
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 105

Gln Ile Ala Thr Gly Val Ser Asp Thr Ile Cys Glu Pro
 1 5 10

<210> 106

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 106

Ala Thr Gly Val Ser Asp Thr Ile Cys Glu Pro Cys Pro
 1 5 10

<210> 107

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 107

Gly Val Ser Asp Thr Ile Cys Glu Pro Cys Pro Val Gly
 1 5 10

<210> 108

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 108

Ser Asp Thr Ile Cys Glu Pro Cys Pro Val Gly Phe Phe
 1 5 10

<210> 109

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 109

Thr Ile Cys Glu Pro Cys Pro Val Gly Phe Phe Ser Asn
 1 5 10

<210> 110

<211> 13

<212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 110

Cys	Glu	Pro	Cys	Pro	Val	Gly	Phe	Phe	Ser	Asn	Val	Ser
1				5					10			

<210> 111

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 111

Pro	Cys	Pro	Val	Gly	Phe	Phe	Ser	Asn	Val	Ser	Ser	Ala
1				5					10			

<210> 112

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 112

Pro	Val	Gly	Phe	Phe	Ser	Asn	Val	Ser	Ser	Ala	Phe	Glu
1				5					10			

<210> 113

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 113

Gly	Phe	Phe	Ser	Asn	Val	Ser	Ser	Ala	Phe	Glu	Lys	Cys
1				5					10			

<210> 114

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 114
 Phe Ser Asn Val Ser Ser Ala Phe Glu Lys Cys His Pro
 1 5 10

<210> 115
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 115
 Asn Val Ser Ser Ala Phe Glu Lys Cys His Pro Trp Thr
 1 5 10

<210> 116
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 116
 Ser Ser Ala Phe Glu Lys Cys His Pro Trp Thr Ser Cys
 1 5 10

<210> 117
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 117
 Ala Phe Glu Lys Cys His Pro Trp Thr Ser Cys Glu Thr
 1 5 10

<210> 118
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 118
 Glu Lys Cys His Pro Trp Thr Ser Cys Glu Thr Lys Asp
 1 5 10

<210> 119
 <211> 13

<212> PRT
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic peptide

 <400> 119
 Cys His Pro Trp Thr Ser Cys Glu Thr Lys Asp Leu Val
 1 5 10

 <210> 120
 <211> 13
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic peptide

 <400> 120
 Pro Trp Thr Ser Cys Glu Thr Lys Asp Leu Val Val Gln
 1 5 10

 <210> 121
 <211> 13
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic peptide

 <400> 121
 Thr Ser Cys Glu Thr Lys Asp Leu Val Val Gln Gln Ala
 1 5 10

 <210> 122
 <211> 13
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic peptide

 <400> 122
 Cys Glu Thr Lys Asp Leu Val Val Gln Gln Ala Gly Thr
 1 5 10

 <210> 123
 <211> 13
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:Synthetic peptide

<400> 123

Thr Lys Asp Leu Val Val Gln Gln Ala Gly Thr Asn Lys
 1 5 10

<210> 124

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 124

Asp Leu Val Val Gln Gln Ala Gly Thr Asn Lys Thr Asp
 1 5 10

<210> 125

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 125

Val Val Gln Gln Ala Gly Thr Asn Lys Thr Asp Val Val
 1 5 10

<210> 126

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 126

Gln Gln Ala Gly Thr Asn Lys Thr Asp Val Val Cys Gly
 1 5 10

<210> 127

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic peptide

<400> 127

Ala Gly Thr Asn Lys Thr Asp Val Val Cys Gly Pro Gln
 1 5 10

<210> 128

<211> 13

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 128
 Thr Asn Lys Thr Asp Val Val Cys Gly Pro Gln Asp Arg
 1 5 10

<210> 129
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 129
 Lys Thr Asp Val Val Cys Gly Pro Gln Asp Arg Leu Arg
 1 5 10

<210> 130
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 peptide

<400> 130
 Asp Val Val Cys Gly Pro Gln Asp Arg Leu Arg Ala Leu
 1 5 10

<210> 131
 <211> 1425
 <212> DNA
 <213> Homo sapiens

<400> 131
 atgtctgtct ccttcctcat cttcctgccc gtgctgggcc tcccatgggg tgtcctgtca 60
 caggtccaac tgcagcagtc aggtccagga ctggtgaagc cctcgcagac cctctcactc 120
 acctgtgcca tctccgggga cagtgtctct agcaacagtg ctacttgga ctggatcagg 180
 cagtcccat cgagagacct tgagtggctg ggaaggacat actacaggtc caagtgggat 240
 cgtgattatg taggatctgt gaaaagtcga ataatcatca acccagacac atccaacaac 300
 cagttctccc tgcagctgaa ctctgtgact cccgaggaca cggctatata ttactgtaca 360
 agagcacagt ggctgggagg ggattacccc tactactaca gtatggacgt ctggggccaa 420
 gggaccacgg tcaccgtctc ctcagctagc accaagggcc catcgggtct cccctggcg 480
 ccctgctcca ggagcacctc cgagagcaca gcggccctgg gctgcctggt caaggactac 540
 ttccccgaac cggtgacggt gtcgtggaac tcaggcgctc tgaccagcgg cgtgcacacc 600
 ttcccagctg tcctacagtc ctcaggactc tactccctca gcagcgtggt gaccgtgccc 660
 tccagcaact tcggcaccca gacctacacc tgcaacgtag atcacaagcc cagcaacacc 720
 aaggtggaca agacagttga gcgcaaagt tgtgtcgagt gccaccgtg cccagcacca 780
 cctgtggcag gaccgtcagt cttcctcttc ccccaaaaac ccaaggacac cctcatgatc 840
 tcccggaccc ctgaggtcac gtgctgtggt gtggacgtga gccacgaaga ccccgaggtc 900
 cagttcaact ggtacgtgga cggcgtggag gtgcataatg ccaagacaaa gccacgggag 960
 gagcagttca acagcacgtt ccgtgtggtc agcgtcctca ccgttgtgca ccaggactgg 1020
 ctgaacggca aggagtacaa gtgcaaggtc tccaacaaag gcctcccagc ctccatcgag 1080

```

aaaaccatct ccaaaaccaa agggcagccc cgagaaccac aggtgtacac cctgccccca 1140
tcccgggagg agatgaccaa gaaccaggtc agcctgacct gcctgggtcaa aggcttctac 1200
cccagcgaca tcgccgtgga gtgggagagc aatgggcagc cggagaacaa ctacaagacc 1260
acacctccca tgctggactc cgacggctcc ttcttctctt acagcaagct caccgtggac 1320
aagagcaggt ggcagcaggg gaacgtcttc tcatgctccg tgatgcatga ggctctgcac 1380
aaccactaca cgcagaagag cctctccctg tctccgggta aatga 1425

```

<210> 132
 <211> 474
 <212> PRT
 <213> Homo sapiens

<400> 132

Met	Ser	Val	Ser	Phe	Leu	Ile	Phe	Leu	Pro	Val	Leu	Gly	Leu	Pro	Trp
1				5					10					15	
Gly	Val	Leu	Ser	Gln	Val	Gln	Leu	Gln	Gln	Ser	Gly	Pro	Gly	Leu	Val
			20					25					30		
Lys	Pro	Ser	Gln	Thr	Leu	Ser	Leu	Thr	Cys	Ala	Ile	Ser	Gly	Asp	Ser
		35					40					45			
Val	Ser	Ser	Asn	Ser	Ala	Thr	Trp	Asn	Trp	Ile	Arg	Gln	Ser	Pro	Ser
	50					55					60				
Arg	Asp	Leu	Glu	Trp	Leu	Gly	Arg	Thr	Tyr	Tyr	Arg	Ser	Lys	Trp	Tyr
65					70				75						80
Arg	Asp	Tyr	Val	Gly	Ser	Val	Lys	Ser	Arg	Ile	Ile	Ile	Asn	Pro	Asp
			85					90					95		
Thr	Ser	Asn	Asn	Gln	Phe	Ser	Leu	Gln	Leu	Asn	Ser	Val	Thr	Pro	Glu
		100						105					110		
Asp	Thr	Ala	Ile	Tyr	Tyr	Cys	Thr	Arg	Ala	Gln	Trp	Leu	Gly	Gly	Asp
	115						120					125			
Tyr	Pro	Tyr	Tyr	Tyr	Ser	Met	Asp	Val	Trp	Gly	Gln	Gly	Thr	Thr	Val
	130					135					140				
Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala
145					150				155						160
Pro	Cys	Ser	Arg	Ser	Thr	Ser	Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu
			165					170						175	
Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly
		180						185					190		
Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser
	195						200					205			
Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Asn	Phe
	210					215					220				
Gly	Thr	Gln	Thr	Tyr	Thr	Cys	Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr
225				230					235						240
Lys	Val	Asp	Lys	Thr	Val	Glu	Arg	Lys	Cys	Cys	Val	Glu	Cys	Pro	Pro
			245					250						255	
Cys	Pro	Ala	Pro	Pro	Val	Ala	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro
		260					265						270		
Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys
		275					280					285			
Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp
	290					295						300			
Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu
305					310				315						320
Glu	Gln	Phe	Asn	Ser	Thr	Phe	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Val
			325					330						335	
His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn
		340						345					350		
Lys	Gly	Leu	Pro	Ala	Ser	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Thr	Lys	Gly
	355						360					365			
Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu
	370					375					380				
Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr
385					390					395					400

```

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
      405                      410                      415
Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe
      420                      425                      430
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
      435                      440                      445
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
      450                      455                      460
Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
465                      470

```

<210> 133
 <211> 696
 <212> DNA
 <213> Homo sapiens

```

<400> 133
atggaagccc cagctcagct tctcttcctc ctgctactct ggctcccaga taccaccgga 60
gaaattgtgt tgacacagtc tccagccacc ctgtctttgt ctccagggga aagagccacc 120
ctctcctgca gggccagtc gagtgtagc agctacttag cctgggtacca acagaaacct 180
ggccaggctc ccaggctcct catctatgat gcatccaaca gggccactgg catcccagcc 240
aggttcagtg gcagtgggtc tgggacagac ttcactctca ccatcagcag cctagagcct 300
gaagattttg cagtttatta ctgtcagcag cgtagcaaca ctttcggccc tgggaccaa 360
gtggatatca aacgtacggg ggctgcacca tctgtcttca tcttcccgcc atctgatgag 420
cagttgaaat ctggaactgc ctctgttggt tgctgtctga ataacttcta tcccagagag 480
gccaaagtac agtggaaggt ggataacgcc ctccaatcgg gtaactccca ggagagtgtc 540
acagagcagg acagcaagga cagcacctac agcctcagca gcaccctgac gctgagcaaa 600
gcagactacg agaaacacaa agtctacgcc tgcgaagtca cccatcaggg cctgagctcg 660
cccgtcacia agagcttcaa caggggagag tgttga 696

```

<210> 134
 <211> 231
 <212> PRT
 <213> Homo sapiens

```

<400> 134
Met Glu Ala Pro Ala Gln Leu Leu Phe Leu Leu Leu Leu Trp Leu Pro
  1                      5                      10                      15
Asp Thr Thr Gly Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser
      20                      25                      30
Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser
      35                      40                      45
Val Ser Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro
      50                      55                      60
Arg Leu Leu Ile Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala
      65                      70                      75                      80
Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
      85                      90                      95
Ser Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser
      100                      105                      110
Asn Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys Arg Thr Val Ala
      115                      120                      125
Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser
      130                      135                      140
Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu
      145                      150                      155                      160
Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser
      165                      170                      175
Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu
      180                      185                      190
Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val
      195                      200                      205

```

Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys
 210 215 220
 Ser Phe Asn Arg Gly Glu Cys
 225 230

<210> 135
 <211> 1407
 <212> DNA
 <213> Homo sapiens

<400> 135
 atggagttgg gactgagctg gatttttcctt ttggctatatt taaaagggtgt ccagtgtgaa 60
 gtgcagctgg tggagtctgg gggaggcttg gtacagcctg gcaggccctt gagactctcc 120
 tgtgcagcct ctggattcac ctttgatgat tatgccatgc actgggtccg gcaagctcca 180
 gggaagggcc tggagtgggt ctcagggtatt agttggaata gtggtagctt ggtgcatgcg 240
 gactctgtga agggccgatt caccatctcc agagacaacg ccaagaactc cctgtatctg 300
 caaatgaaca gtctgagagc tgaggacacg gccttgatatt actgtgcaag agataggcta 360
 tttcgggggag ttaggtacta cggatatggac gtctggggcc aagggaccac ggtcaccgtc 420
 tcctcagcta gcaccaaggg cccatcgggtc ttccccctgg cgccctgctc caggagcacc 480
 tccgagagca cagcggccct gggctgcctg gtcaaggact acttccccga accggtgacg 540
 gtgtcgtgga actcaggcgc tctgaccagc ggcgtgcaca ccttcccagc tgtcctacag 600
 tcctcaggac tctactccct cagcagcgtg gtgaccgtgc cctccagcaa cttcggcacc 660
 cagacctaca cctgcaacgt agatcacaaag cccagcaaca ccaagggtgga caagacagtt 720
 gagcgcaaat gttgtgtcga gtgcccaccg tgcccagcac cacctgtggc aggaccgtca 780
 gtcttctctt tccccccaaa acccaaggac accctcatga tctcccggac ccctgagggtc 840
 acgtgcgtgg tgggtggacgt gagccacgaa gaccccgagg tccagttcaa ctggtacgtg 900
 gacggcgtgg aggtgcataa tgccaagaca aagccacggg aggagcagtt caacagcacg 960
 ttccgtgtgg tcagcgtcct caccgttgtg caccaggact ggctgaacgg caaggagtac 1020
 aagtgcagg tctccaacaa aggcctccca gcctccatcg agaaaaccat ctccaaaacc 1080
 aaagggcagc cccgagaacc acagggtgtac accctgcccc catcccggga ggagatgacc 1140
 aagaaccagg tcagcctgac ctgcctgggtc aaaggcttct accccagcga catcgccgtg 1200
 gagtggggaga gcaatgggca gccggagaa aactacaaga ccacacctcc catgctggac 1260
 tccgacggct ccttcttctt ctacagcaag ctcaccgtgg acaagagcag gtggcagcag 1320
 gggaacgtct tctcatgctc cgtgatgcat gaggtctctgc acaaccacta cacgcagaag 1380
 agcctctccc tgtctccggg taaatga 1407

<210> 136
 <211> 468
 <212> PRT
 <213> Homo sapiens

<400> 136
 Met Glu Leu Gly Leu Ser Trp Ile Phe Leu Leu Ala Ile Leu Lys Gly
 1 5 10 15
 Val Gln Cys Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln
 20 25 30
 Pro Gly Arg Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Asp Asp Tyr Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Val Ser Gly Ile Ser Trp Asn Ser Gly Ser Leu Val His Ala
 65 70 75 80
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95
 Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Leu
 100 105 110
 Tyr Tyr Cys Ala Arg Asp Arg Leu Phe Arg Gly Val Arg Tyr Tyr Gly
 115 120 125
 Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser
 130 135 140
 Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr
 145 150 155 160

Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro
 165 170 175
 Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val
 180 185 190
 His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser
 195 200 205
 Ser Val Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr
 210 215 220
 Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val
 225 230 235 240
 Glu Arg Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val
 245 250 255
 Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu
 260 265 270
 Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser
 275 280 285
 His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu
 290 295 300
 Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr
 305 310 315 320
 Phe Arg Val Val Ser Val Leu Thr Val Val His Gln Asp Trp Leu Asn
 325 330 335
 Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Ser
 340 345 350
 Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln
 355 360 365
 Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val
 370 375 380
 Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val
 385 390 395 400
 Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro
 405 410 415
 Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr
 420 425 430
 Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val
 435 440 445
 Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
 450 455 460
 Ser Pro Gly Lys
 465

<210> 137
 <211> 702
 <212> DNA
 <213> Homo sapiens

<400> 137
 atggaagccc cagctcagct tctcttctc ctgctactct ggctcccaga taccaccgga 60
 gaaattgtgt tgacacagtc tccagccacc ctgtctttgt ctccagggga aagagccacc 120
 ctctcctgca gggccagtca gagggttagc agctacttag cctggtacca acagaaacct 180
 ggccaggctc ccaggctcct catctatgat gcatccaaca gggccactgg catcccagcc 240
 aggttcagtg gcagtggtgc tgggacagac ttcactctca ccatcagcag cctagagcct 300
 gaagattttg cagtttatta ctgtcagcag cgtagccact ggctcacttt cggcgggggg 360
 accaaggtgg agatcaaacg tacggtggct gcaccatctg tcttcatctt cccgccatct 420
 gatgagcagt tgaaatctgg aactgcctct gttgtgtgcc tgctgaataa cttctatccc 480
 agagaggcca aagtacagtg gaaggtggat aacgccctcc aatcgggtaa ctcccaggag 540
 agtgtcacag agcaggacag caaggacagc acctacagcc tcagcagcac cctgacgctg 600
 agcaaagcag actacgagaa acacaaagtc tacgcctgcg aagtcaccca tcagggcctg 660
 agctcgcccc tcacaaagag cttcaacagg ggagagtgtt ga 702

<210> 138
 <211> 233

<212> PRT
 <213> Homo sapiens

<400> 138

Met	Glu	Ala	Pro	Ala	Gln	Leu	Leu	Phe	Leu	Leu	Leu	Leu	Trp	Leu	Pro
1				5				10						15	
Asp	Thr	Thr	Gly	Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser
			20					25					30		
Leu	Ser	Pro	Gly	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser
		35					40					45			
Val	Ser	Ser	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro
	50					55					60				
Arg	Leu	Leu	Ile	Tyr	Asp	Ala	Ser	Asn	Arg	Ala	Thr	Gly	Ile	Pro	Ala
65					70				75					80	
Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser
			85					90						95	
Ser	Leu	Glu	Pro	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Arg	Ser
			100					105					110		
His	Trp	Leu	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Val	Glu	Ile	Lys	Arg	Thr
	115						120					125			
Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	Leu
	130					135					140				
Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	Pro
145					150					155					160
Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln	Ser	Gly
			165					170						175	
Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr	Tyr
			180					185					190		
Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys	His
	195						200					205			
Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	Val
	210					215					220				
Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys							
225					230										

<210> 139
 <211> 1425
 <212> DNA
 <213> Homo sapiens

<400> 139

atggatctca	tgtgcaagaa	aatgaagcac	ctgtggttct	tcctcctgct	ggtggcggct	60
cccagatggg	tcctgtccca	gctgcagctg	caggagtcgg	gcccaggact	actgaagcct	120
tcggagaccc	tgtccctcac	ctgcactgtc	tctggcggct	ccatcagcag	tcctgggttac	180
tacgggggct	ggatccgcca	gccccaggg	aaggggctgg	agtggattgg	gagtatctat	240
aaaagtggga	gcacctacca	caaccgtcc	ctcaagagtc	gagtcaccat	atccgtagac	300
acgtccaaga	accagttctc	cctgaagctg	agctctgtga	ccgccgcaga	cacggctgtg	360
tattactgta	cgagacctgt	agtagatat	tttgggtggt	tcgacctctg	gggccaggga	420
accctgggtca	ccgtctcctc	agctagcacc	aagggggccat	ccgtcttccc	cctggcgccc	480
tgctccagga	gcacctccga	gagcacagcc	gccctgggct	gcctgggtcaa	ggactacttc	540
cccgaaccgg	tgacgggtgc	gtggaactca	ggcgccctga	ccagcggcgt	gcacaccttc	600
ccggctgtcc	tacagtcttc	aggactctac	tcctcagca	gcgtgggtgac	cgtgccctcc	660
agcagcttgg	gcacgaagac	ctacacctgc	aacgtagatc	acaagcccag	caacaccaag	720
gtggacaaga	gagttgagtc	caaatatggt	cccccatgcc	caccatgccc	agcacctgag	780
ttcgaggggg	gaccatcagt	cttcctgttc	ccccaaaac	ccaaggacac	tctcatgatc	840
tcccggaccc	ctgagggtcac	gtgcgtgggtg	gtggacgtga	gccaggaaga	ccccgaggtc	900
cagttcaact	ggtacgtgga	tggcgtggag	gtgcataatg	ccaagacaaa	gccgcgggag	960
gagcagttca	acagcacgta	ccgtgtgggtc	agcgtcctca	ccgtcctgca	ccaggactgg	1020
ctgaacggca	aggagtacaa	gtgcaaggtc	tccaacaaag	gcctcccgtc	ctccatcgag	1080
aaaaccatct	ccaaagccaa	agggcagccc	cgagagccac	aggtgtacac	cctgccccca	1140
tcccaggagg	agatgaccaa	gaaccaggtc	agcctgacct	gcctgggtcaa	aggcttctac	1200
cccagcgaca	tcgccgtgga	gtgggagagc	aatgggcagc	cggagaacaa	ctacaagacc	1260
acgcctcccc	tgctggactc	cgacggctcc	ttcttcctct	acagcaggct	aaccgtggac	1320

aagagcaggt ggcaggaggg gaatgtcttc tcatgctccg tgatgcatga ggctctgcac 1380
aaccactaca cacagaagag cctctccctg tctctgggta aatga 1425

<210> 140
<211> 474
<212> PRT
<213> Homo sapiens

<400> 140
Met Asp Leu Met Cys Lys Lys Met Lys His Leu Trp Phe Phe Leu Leu
1 5 10 15
Leu Val Ala Ala Pro Arg Trp Val Leu Ser Gln Leu Gln Leu Gln Glu
20 25 30
Ser Gly Pro Gly Leu Leu Lys Pro Ser Glu Thr Leu Ser Leu Thr Cys
35 40 45
Thr Val Ser Gly Gly Ser Ile Ser Ser Pro Gly Tyr Tyr Gly Gly Trp
50 55 60
Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile Gly Ser Ile Tyr
65 70 75 80
Lys Ser Gly Ser Thr Tyr His Asn Pro Ser Leu Lys Ser Arg Val Thr
85 90 95
Ile Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu Lys Leu Ser Ser
100 105 110
Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Thr Arg Pro Val Val
115 120 125
Arg Tyr Phe Gly Trp Phe Asp Pro Trp Gly Gln Gly Thr Leu Val Thr
130 135 140
Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro
145 150 155 160
Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val
165 170 175
Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala
180 185 190
Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly
195 200 205
Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly
210 215 220
Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys
225 230 235 240
Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys
245 250 255
Pro Ala Pro Glu Phe Glu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
260 265 270
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
275 280 285
Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp
290 295 300
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
305 310 315 320
Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
325 330 335
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
340 345 350
Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
355 360 365
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu
370 375 380
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
385 390 395 400
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
405 410 415
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
420 425 430

Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn
 435 440 445
 Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
 450 455 460
 Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys
 465 470

<210> 141
 <211> 708
 <212> DNA
 <213> Homo sapiens

<400> 141
 atggacatga gggccccgc tcagctcctg gggcttctgc tgctctggct cccaggtgcc 60
 agatgtgcca tccagttgac ccagtctcca tcctccctgt ctgcatctgt aggagacaga 120
 gtcaccatca cttgccgggc aagtcagggc attagcagtg ctttagcctg gtatcagcag 180
 aaaccagggg aagctcctaa gctcctgac tatgatgcct ccaatttgga aagtgggggc 240
 ccatcaaggt tcagcggcag tggatctggg acagatttca ctctcaccat cagcagcctg 300
 cagcctgaag attttgcaac ttattactgt caacagttta atagttaccc gacgttcggc 360
 caagggacca aggtggaaat caaacgtacg gtggctgcac catctgtctt catcttcccg 420
 ccatctgatg agcagttgaa atctggaact gcctctgttg tgtgcctgct gaataacttc 480
 tatcccagag aggccaaagt acagtggaag gtggataacg ccctccaatc gggtaactcc 540
 caggagagtg tcacagagca ggacagcaag gacagcacct acagcctcag cagcaccctg 600
 acgctgagca aagcagacta cgagaaacac aaagtctacg cctgcgaagt caccatcag 660
 ggcctgagct cgcccgtcac aaagagcttc aacaggggag agtggtga 708

<210> 142
 <211> 235
 <212> PRT
 <213> Homo sapiens

<400> 142
 Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu Trp
 1 5 10 15
 Leu Pro Gly Ala Arg Cys Ala Ile Gln Leu Thr Gln Ser Pro Ser Ser
 20 25 30
 Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
 35 40 45
 Gln Gly Ile Ser Ser Ala Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys
 50 55 60
 Ala Pro Lys Leu Leu Ile Tyr Asp Ala Ser Asn Leu Glu Ser Gly Val
 65 70 75 80
 Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 85 90 95
 Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
 100 105 110
 Phe Asn Ser Tyr Pro Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 115 120 125
 Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
 130 135 140
 Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
 145 150 155 160
 Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
 165 170 175
 Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
 180 185 190
 Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
 195 200 205
 Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
 210 215 220
 Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230 235

72813-258

CLAIMS:

1. A pharmaceutical composition for use in prevention or treatment of transplant rejection, autoimmune diseases, allergy or blood clotting factor VIII inhibition, comprising a monoclonal antibody as an active ingredient, wherein the monoclonal antibody consists of two heavy chains, each represented by an amino acid sequence ranging from glutamine at position 27 to lysine at position 474 in SEQ ID NO: 140, and two light chains, each represented by an amino acid sequence ranging from alanine at position 23 to cysteine at position 235 in SEQ ID NO: 142.
2. Use of a monoclonal antibody for production of a pharmaceutical composition for prevention or treatment of transplant rejection, autoimmune diseases, allergy or blood clotting factor VIII inhibition, wherein the monoclonal antibody consists of two heavy chains, each represented by an amino acid sequence ranging from glutamine at position 27 to lysine at position 474 in SEQ ID NO: 140, and two light chains, each represented by an amino acid sequence ranging from alanine at position 23 to cysteine at position 235 in SEQ ID NO: 142.
3. Use of a monoclonal antibody for prevention or treatment of transplant rejection, autoimmune diseases, allergy or blood clotting factor VIII inhibition, wherein the monoclonal antibody consists of two heavy chains, each represented by an amino acid sequence ranging from glutamine at position 27 to lysine at position 474 in SEQ ID NO: 140, and two light chains, each represented by an amino acid sequence ranging from alanine at position 23 to cysteine at position 235 in SEQ ID NO: 142.

Application number: numéro de demande: JP04/19750

Figures: _____

Pages: 8, 9, 12 to 23, 39

DRW-TP

Unscannable items
received with this application
(Request original documents in File Prep. Section on the 10th Floor)

Documents reçus avec cette demande ne pouvant être balayés
(Commander les documents originaux dans la section de préparation des dossiers au
10ième étage)

Fig. 1A-1

Sequence	2105	KM643-4-11	F2-103	110	KM341-1-19
1. EPPTACREKQYLI	16353	13220	31427	13403	14108
2. PTACREKQYLINS	17358	13220	39202	13575	14194
3. ACREKQYLINSQC	20267	13212	35911	13681	14013
4. REKQYLINSQCCS	17096	13305	43762	13685	13952
5. KQYLINSQCCSLC	23249	13292	36839	13894	13779
6. YLINSQCCSLCQP	16140	13470	44975	13964	13909
7. INSQCCSLCQPGQ	15918	13390	54811	13652	13751
8. SQCCSLCQPGQKL	18804	13271	60441	13872	13819
9. CCSLCQPGQKLVS	16534	13436	56601	13821	13851
10. SLCQPGQKLVSDC	18155	13314	54718	13705	13965
11. CQPGQKLVSDCTE	17614	13460	52073	13699	13781
12. PGQKLVSDCTEFT	91270	13334	58177	13464	13725
13. QKLVSDCTEFTET	96926	13465	49003	14916	13875
14. LVSDCTEFTETEC	141877	13634	62580	13394	13783
15. SDCTEFTETECLP	148692	13213	42916	13571	13696
16. CTEFTETECLPCG	145926	13591	38170	13220	13824
17. EFTETECLPCGES	136350	13593	56597	13434	13638
18. TETECLPCGESEF	77805	13298	33201	13398	13723
19. TECLPCGESEFLD	50648	13544	29781	13488	13685
20. CLPCGESEFLDTW	79246	13736	26075	13599	13807
21. PCGESEFLDTWNR	93482	13626	44337	16142	13830
22. GESEFLDTWNRET	68932	14815	45811	13777	13832
23. SEFLDTWNRETHC	105899	13777	40462	14023	13898
24. FLDTWNRETHCHQ	55803	15498	79990	40210	13937
25. DTWNRETHCHQHK	19237	13367	34525	13995	13737
26. WNRETHCHQHKEYC	22963	13843	32912	15601	13595
27. RETHCHQHKEYCDP	16522	13282	41600	13644	13595
28. THCHQHKEYCDPNL	25181	13211	32264	13567	13621
29. CHQHKEYCDPNLGL	16455	13300	34480	13684	13671
30. QHKEYCDPNLGLRV	16627	13541	39057	13461	13670
31. KYCDPNLGLRVQQ	16429	13491	46485	13452	13779
32. CDPNLGLRVQQKG	16877	13454	90926	13293	13799
33. PNLGLRVQQKGTS	18117	13611	123330	13358	13917
34. LGLRVQQKGTSSET	22898	13512	237353	13381	15032
35. LRVQQKGTSSETDT	24787	13599	250401	13364	13774
36. VQQKGTSSETDTIC	28492	13372	35303	13124	13686
37. QKGTSSETDTICTC	27724	13490	35334	13298	13569
38. GTSETDTICTCEE	18834	13340	29905	13289	13525
39. SETDTICTCEEGW	16914	13340	24871	13315	13486
40. TDTICTCEEGWHC	17866	13568	24475	13481	13631

Fig. 1A-2

Sequence	2105	KM643-4-11	F2-103	110	KM341-1-19
41. TICTCEEGWHCTS	18122	13767	24169	13564	68670
42. CTCEEGWHCTSEA	17485	18955	28190	13736	75214
43. CEEGWHCTSEACE	16859	41843	32056	13525	16033
44. EGWHCTSEACESC	17735	13381	35518	13771	13867
45. WHCTSEACESCVL	64890	13444	38190	13708	13687
46. CTSEACESCVLHR	16788	13446	39497	13859	13636
47. SEACESCVLHRSC	16421	13555	40206	13482	13584
48. ACESCVLHRSCSP	15795	13183	36735	13503	13575
49. ESCVLHRSCSPGF	47038	13504	41251	14500	13637
50. CVLHRSCSPGFGV	17469	13279	45795	13484	13573
51. LHRSCSPGFGVKQ	16635	13382	88644	13187	13567
52. RSCSPGFGVKQIA	16642	13339	114128	13368	13623
53. CSPGFGVKQIATG	17457	13485	110624	13323	13703
54. PGFGVKQIATGVS	18818	13577	253876	13234	13769
55. FGVKQIATGVSDT	20521	13429	697242	13178	13903
56. VKQIATGVSDTIC	20855	13510	91544	13176	13681
57. QIATGVSDTICEP	19014	13523	40193	13154	13575
58. ATGVSDTICEPCP	17139	13526	38012	13127	13599
59. GVSDTICEPCPVG	16515	13508	31118	13376	13571
60. SDTICEPCPVGFF	42762	13590	25542	13392	13534
61. TICEPCPVGFFSN	25039	13366	27921	13524	13783
62. CEPCPVGFFSNVS	18141	13429	31377	14364	13717
63. PCPVGFFSNVSSA	20806	13343	51411	19243	13979
64. PVGFFSNVSSAFE	95541	14553	128669	158203	14523
65. GFFSNVSSAFEKC	67908	13506	94016	14323	13831
66. FSNVSSAFEKCHP	22379	13350	124184	14057	13699
67. NVSSAFEKCHPWT	18703	13298	44966	13518	13676
68. SSAFEKCHPWTSC	19809	13459	37784	13433	13553
69. AFEKCHPWTSCET	16212	13387	38013	13279	13683
70. EKCHPWTSCETKD	15198	13423	39248	13315	13494
71. CHPWTSCETKDLV	15754	13383	40627	13228	13673
72. PWTSCETKDLVVQ	15609	13426	46662	13247	13613
73. TSCETKDLVVQQA	16454	13377	46750	13228	13670
74. CETKDLVVQQAGT	16875	13544	42504	13185	13700
75. TKDLVVQQAGTNK	17357	14026	83860	13184	13674
76. DLVVQQAGTNKTD	17735	13434	34853	13154	13776
77. VVQQAGTNKTDVV	17176	13805	51573	13166	13651
78. QQAGTNKTDVVCG	15794	13261	27337	13103	13398
79. AGTNKTDVVCGPQ	15793	13448	25989	13304	13658
80. TNKTDVVCGPQDR	15114	13481	24807	13199	13767
81. KTDVVCGPQDRLR	15782	13328	35951	13548	14112
82. DVVCGPQDRLRAL	16644	13255	33393	13756	14104

Fig. 1B-1

Sequence	F4-465	281-1-10	2B11	F72	F76	4D11
1. EPPTACREKQYLI	20561	4194	14158	3965	3348	14138
2. PTACREKQYLINS	19249	4141	14276	3906	3420	14387
3. ACREKQYLINSQC	20418	4221	14276	3884	3300	14044
4. REKQYLINSQCCS	20278	4080	14214	3849	3546	13956
5. KQYLINSQCCSLC	19642	4169	14082	3806	3378	14443
6. YLINSQCCSLCQP	19658	4184	14231	4082	3464	14432
7. INSQCCSLCQPGQ	18482	4603	13927	3986	3157	13783
8. SQCCSLCQPGQKL	19075	4326	13748	3934	3675	13471
9. CCSLCQPGQKLVS	20282	4349	13578	4046	3336	13431
10. SLCQPGQKLVSDC	20175	4841	13522	3948	3222	13387
11. CQPGQKLVSDCTE	22520	4576	13441	4115	3566	13339
12. PGQKLVSDCTEFT	23688	4499	13618	4012	3525	13290
13. QKLVSDCTEFTET	22029	4771	13528	4294	3539	13450
14. LVSDCTEFTETEC	23692	5212	13439	4177	3601	14898
15. SDCTEFTETECLP	25950	4864	13359	4095	3362	13277
16. CTEFTETECLPCG	20943	4906	13496	4121	3582	13874
17. EFTETECLPCGES	37123	4748	13570	3960	3402	16583
18. TETECLPCGESEF	23400	4807	13444	3811	3407	13808
19. TECLPCGESEFLD	21427	4649	13352	3970	3288	13566
20. CLPCGESEFLDTW	18021	4794	19449	3940	5926	47209
21. PCGESEFLDTWNR	26480	7656	86887	5478	16438	165895
22. GESEFLDTWNRET	25229	4906	24124	4337	4362	32083
23. SEFLDTWNRETHC	23274	5979	30344	4138	4083	72724
24. FLDTWNRETHCHQ	28280	19544	120439	4636	6401	336496
25. DTWNRETHCHQHK	20114	4107	14285	4029	3459	14028
26. WNRETHCHQHKEYC	21110	4714	16070	4111	3819	25715
27. RETHCHQHKEYCDP	19452	4075	13942	4000	3252	13812
28. THCHQHKEYCDPNL	18412	4221	14203	3912	3459	14295
29. CHQHKEYCDPNLGL	18890	4288	16566	4025	3473	13886
30. QHKEYCDPNLGLRV	19296	4313	13580	4132	3383	13355
31. KYCDPNLGLRVQQ	20430	4460	13469	4254	3651	13357
32. CDPNLGLRVQQKG	28700	4570	13568	5394	4575	13336
33. PNLGLRVQQKGTS	39822	5229	13590	24278	6996	13404
34. LGLRVQQKGTSSET	52908	4681	13651	7042	4112	13400
35. LRVQQKGTSSETDT	56985	4376	13514	4839	16679	13557
36. VQQKGTSSETDTIC	21684	4531	13423	3960	3258	13355
37. QKGTSSETDTICTC	21838	4509	13507	3975	3355	13365
38. GTSETDTICTCEE	21896	4074	13348	3856	3298	13453
39. SETDTICTCEEGBW	18789	4082	13377	3687	3297	13606
40. TDTICTCEEGBWHC	17729	4187	13485	3756	3638	19679

Fig. 1B-2

Sequence	F4-465	281-1-10	2B11	F72	F76	4D11
41. TICTCEEGWHCTS	22932	4207	14559	4151	3992	46128
42. CTCEEGWHCTSEA	22823	4167	14428	4077	3394	14309
43. CEEGWHCTSEACE	25545	4071	14192	4118	3375	14162
44. EGWHCTSEACESC	23461	4272	14290	4171	3393	14029
45. WHCTSEACESCVL	20999	4198	14136	4318	3465	14065
46. CTSEACESCVLHR	20553	4273	14301	4152	3396	14337
47. SEACESCVLHRSC	20792	4234	13900	4122	3439	13877
48. ACESCVLHRSCSP	21816	4179	13741	3934	3525	13539
49. ESCVLHRSCSPGF	20599	4725	13758	4141	3514	13658
50. CVLHRSCSPGFGV	19851	4393	13533	4127	3526	13320
51. LHRSCSPGFGVKQ	23160	4317	13486	4851	18376	13355
52. RSCSPGFGVKQIA	24797	4495	13728	5568	15564	13375
53. CSPGFGVKQIATG	30890	4962	13400	8711	3383	13340
54. PGFGVKQIATGVS	82448	5027	13415	11404	4102	13378
55. FGVKQIATGVSDT	126309	4541	13423	4896	3515	13474
56. VKQIATGVSDTIC	32389	4361	13454	4040	3278	13305
57. QIATGVSDTICEP	28554	4308	13325	3807	3318	13350
58. ATGVSDTICEPCP	22484	4176	13463	3868	3241	13399
59. GVSDTICEPCPVG	21417	4277	13329	3584	3259	13475
60. SDTICEPCPVGFF	20192	4360	13337	3725	3358	14425
61. TICEPCPVGFFSN	25475	4390	14453	4101	3446	15502
62. CEPCPVGFFSNVS	26387	4419	14370	4434	3538	14441
63. PCPVGFFSNVSSA	40241	4468	14425	4363	3409	14664
64. PVGFFSNVSSAFE	27432	11225	85432	4772	5735	49493
65. GFFSNVSSAFEKC	77408	4483	14462	4173	3489	14521
66. FSNVSSAFEKCHP	57923	4088	14050	4230	3296	13989
67. NVSSAFEKCHPWT	23154	4031	13752	4025	3343	13619
68. SSAFEKCHPWTSC	20994	4170	13512	4006	3419	13466
69. AFEKCHPWTSCET	21390	4498	13629	4028	3413	13518
70. EKCHPWTSCETKD	19639	4186	13571	3911	3403	16038
71. CHPWTSCETKDLV	20967	4216	13608	4177	3489	13283
72. PWTSCETKDLVVQ	23271	4300	13420	4459	3316	13507
73. TSCETKDLVVQQA	26353	4690	13457	4411	3496	13462
74. CETKDLVVQQAGT	26955	4517	13389	4600	3403	13433
75. TKDLVVQQAGTNK	63164	6695	13475	60382	7609	13271
76. DLVVQQAGTNKTD	25820	4567	13537	4057	3585	13328
77. VVQQAGTNKTDVV	28426	4587	13372	4179	3418	13309
78. QQAGTNKTDVVCG	19474	4111	13455	3709	3809	13332
79. AGTNKTDVVCGPQ	18988	4186	13517	3868	3231	13436
80. TNKTDVVCGPQDR	18178	4662	13423	3654	4897	13350
81. KTDVVCGPQDRLR	27431	4208	13879	4242	4238	13779
82. DVVCGPQDRLRAL	28872	4127	14128	4179	3459	13855

Fig. 2A

CD40FC




Fig. 2B

CD40mut 1




Fig. 2C

CD40mut2




Fig. 4A




Fig. 4B




Fig. 9




Fig. 10




Fig. 11




Fig. 12A




Fig. 12B





Fig. 12C






Fig. 13A

Fig. 13B




Fig. 14A




Fig. 14B




Fig. 15




Fig. 16




Fig. 17

Treatment				Average score*	
Group (mg/kg)	Tetanus toxoid		Abdominal intradermal challenge	24 hrs**	48 hrs**
	Dorsal intradermal	Sensitization Intramuscular			
Control	10 µL/site		0 Lf/mL	0.00	0.00
	x	0.6 mL/animal	1 Lf/mL	1.67	1.00
	12 sites		3 Lf/mL	3.00	2.67
			10 Lf/mL	3.33	3.00
4D11 1 (mg/kg)	10 µL/site		0 Lf/mL	0.00	0.00
	x	0.6 mL/animal	1 Lf/mL	0.67	0.67
	12 sites		3 Lf/mL	1.00	0.67
			10 Lf/mL	1.00	1.00
4D11 10 (mg/kg)	10 µL/site		0 Lf/mL	0.00	0.00
	x	0.6 mL/animal	1 Lf/mL	0.00	0.00
	12 sites		3 Lf/mL	0.67	0.33
			10 Lf/mL	1.00	0.67
Remarks: * Response score					
formation of erythema and/or eschar			formation of edema		
no erythema			no edema		
very slight erythema (barely perceptible)			very slight edema (barely perceptible)		
well-defined erythema			slight edema (edges of area well defined by definite raising)		
moderate to severe erythema			moderate edema (raised about 1 mm)		
from severe erythema to slight crust (deep injury)			severe edema (raised 1 mm or more and extending beyond area of exposure)		
** time after challenge					
			: 0		
			: 1		
			: 2		
			: 3		
			: 4		

Fig. 18

Group	Animal No.	pre	(Maximum dilution rate for positive effect)							
			Days after the 1st dose							
			4	7	11	14	18	21	23	
Control	1	0	200	0	12800	102400	204800	204800	204800	
	2	0	0	0	6400	25600	51200	51200	51200	
	3	0	0	0	6400	12800	25600	25600	25600	
	Mean	0	67	0	8533	46933	93867	93867	93867	
	S.D.	0	115	0	3695	48460	96920	96920	96920	
1 (mg/kg)	4	0	0	0	200	1600	12800	12800	12800	
	5	0	0	0	200	400	6400	6400	6400	
	6	0	0	0	400	3200	6400	6400	6400	
	Mean	0	0	0	267	1733	8533	8533	8533	
	S.D.	0	0	0	115	1405	3695	3695	3695	
10 (mg/kg)	7	0	0	0	0	0	0	0	0	
	8	0	0	0	0	0	0	0	0	
	9	0	0	0	0	0	0	0	0	
	Mean	0	0	0	0	0	0	0	0	
	S.D.	0	0	0	0	0	0	0	0	

Fig. 19

Group	Animal No.	pre	Days after the 1st dose								(Maximum dilution rate for positive effect)			
			4	7	11	14	18	21	23					
Control	1	0	0	0	3200	6400	6400	6400	6400	6400	6400	6400	6400	
	2	0	0	0	1600	1600	1600	1600	1600	1600	1600	800	800	
	3	0	0	0	800	1600	1600	1600	1600	1600	1600	1600	1600	
	Mean	0	0	0	1867	3200	3200	3200	3200	3200	3200	2933	2933	
	S.D.	0	0	0	1222	2771	2771	2771	2771	2771	2771	3029	3029	
1 (mg/kg)	4	0	0	0	0	0	0	400	0	0	0	0	0	
	5	0	0	0	0	0	0	0	0	0	0	0	0	
	6	0	0	0	0	0	800	800	800	800	800	800	800	
	Mean	0	0	0	0	0	267	400	400	267	267	267	267	
	S.D.	0	0	0	0	0	462	400	400	462	462	462	462	
10 (mg/kg)	7	0	0	0	0	0	0	0	0	0	0	0	0	
	8	0	0	0	0	0	0	0	0	0	0	0	0	
	9	0	0	0	0	0	0	0	0	0	0	0	0	
	Mean	0	0	0	0	0	0	0	0	0	0	0	0	
	S.D.	0	0	0	0	0	0	0	0	0	0	0	0	

Fig. 20B




Fig. 21




Fig. 22




Fig. 23




Fig. 24




Fig. 25




Fig. 26

Capan-2

