

(19) **DANMARK**

(10) **DK/EP 3145947 T3**



(12)

Oversættelse af
europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 07 K 14/195 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2025-03-31**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2025-02-12**
- (86) Europæisk ansøgning nr.: **15753495.9**
- (86) Europæisk indleveringsdag: **2015-05-21**
- (87) Den europæiske ansøgnings publiceringsdag: **2017-03-29**
- (86) International ansøgning nr.: **IL2015050540**
- (87) Internationalt publikationsnr.: **WO2015177800**
- (30) Prioritet: **2014-05-22 US 201462001736 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Yeda Research and Development Co., Ltd., at the Weizmann Institute of Science , P.O. Box 95, 7610002 Rehovot, Israel**
- (72) Opfinder: **MILO, Ron, 11 Arazim Street, 4445630 Kfar-Saba, Israel**
ANTONOVSKY, Niv, c/o Yeda Research and Development Co. Ltd., At the Weizmann Institute of Science, P.O. Box 95, 7610002 Rehovot, Israel
NOOR, Elad, c/o Yeda Research and Development Co. Ltd., at the Weizmann Institute of Science, P.O. Box 95, 7610002 Rehovot, Israel
BAR-EVEN, Arren, c/o Yeda Research and Development Co. Ltd., at the Weizmann Institute of Science, P.O. Box 95, 7610002 Rehovot, Israel
ZOHAR, Yehudit, c/o Yeda Research and Development Co. Ltd., at the Weizmann Institute of Science, P.O. Box 95, 7610002 Rehovot, Israel
ZELCBUCH, Lior, C/o Yeda Research and Development Co. Ltd., At the Weizmann Institute of Science, P.O. Box 95, 7610002 Rehovot, Israel
GLEIZER, Shmuel, c/o Yeda Research and Development Co. Ltd., at the Weizmann Institute of Science, P.O. Box 95, 7610002 Rehovot, Israel
AMRAM, Shira, c/o Yeda Research and Development Co. Ltd., at the Weizmann Institute of Science, P.O. Box 95, 7610002 Rehovot, Israel
- (74) Fuldmægtig i Danmark: **Dennemeyer & Associates S.A, P.O. Box 700425, DE-81304 Munich, Tyskland**
- (54) Benævnelse: **REKOMBINANTE MIKROORGANISMER, SOM ER I STAND TIL FIKSERING AF KULSTOF**
- (56) Fremdragne publikationer:
WO-A1-2009/036095
V?CTOR GUADALUPE-MEDINA ET AL: "Carbon dioxide fixation by Calvin-Cycle enzymes improves ethanol yield in yeast", BIOTECHNOLOGY FOR BIOFUELS, vol. 6, no. 1, 1 January 2013 (2013-01-01), GB, pages 125, XP055405759, ISSN: 1754-6834, DOI: 10.1186/1754-6834-6-125

Fortsættes ...

DESCRIPTION

Description

[0001] The present disclosure, in some aspects thereof, relates to recombinant non-autotrophic microorganisms that are capable of carbon fixation.

[0002] Carbon fixation plays an essential role in ecosystems by providing a continuous flux of organic carbon into the biosphere. From a human perspective, the process of CO₂ assimilation dominates humanity's usage of land and water. Importantly, under human cultivation, where water, light and nutrients are abundant, carbon fixation often limits plant growth. Therefore, increasing the rate of carbon fixation is of major importance in the path towards agricultural and energetic sustainability.

[0003] Carbon fixation in plants, algae, cyanobacteria and many other bacterial lineages is achieved by the Calvin-Benson-Bassham (CBB) Cycle. The productivity of the CBB cycle is limited, under many conditions, by the relatively slow rate and lack of substrate specificity of the carboxylating enzyme RuBisCO. Previous attempts to utilize classic molecular biology tools to improve the rate and specificity of RuBisCO have achieved only limited success. Moreover, several lines of evidence indicate that in spite of its shortcomings, RuBisCO is already naturally optimized given the natural tradeoff between the enzyme's rate and specificity. Other enzymes of the CBB also were shown to constrain carbon fixation in certain circumstances. For example, tobacco plants overexpressing sedoheptulose-1,7-bisphosphatase were characterized by an increased photosynthetic rate and a 30% enhancement in biomass yield. Therefore, improving the rate of the CBB cycle seems to be quite a complex task that has many possible leads. Such a challenge can benefit significantly from the utilization of novel selection systems that can go far beyond the sequence divergence presented in plants.

[0004] WO 2011/099006 teaches plants and bacterial cells which are genetically modified so as to express enzymes of alternative (RuBisCO-independent) pathways in order to promote carbon fixation. Autotrophic *E. Coli* cells which express phosphoribulokinase and Ribulose-Bisphosphate Carboxylase are also disclosed.

[0005] WO 2009036095 identifies pathways and mechanisms to confer photoautotrophic properties to a heterotrophic organism. The resultant engineered cell or organism will uniquely enable efficient conversion of carbon dioxide and light into biomass and carbon-based products of interest.

SUMMARY OF THE INVENTION

[0006] The present invention is set out in the appended set of claims.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

[0007] Some aspects of the disclosure are herein described, by way of example only, with reference to the accompanying drawings and images. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of aspects of the disclosure. In this regard, the description taken with the drawings makes apparent to those skilled in the art how aspects of the disclosure may be practiced.

[0008] In the drawings:

FIG. 1 is a diagram illustrating how decoupling energy production and carbon fixation in *E. Coli* achieves semiautotrophic growth. In native autotrophs, the CBB cycle enables the transformation of inorganic carbon and chemical energy into biomass precursors. Two recombinant enzymes are needed to complete the carbon fixation cycle in *E. Coli*: RuBisCO, the carboxylating enzyme, and the kinase *prk*. The remaining reactions required for the reduction and substrate regeneration phases of the cycle are endogenous to the metabolic network of the host, as part of the gluconeogenic and the pentose phosphate pathways. The deletion of the phosphoglycerate mutase genes (*gpmA* and *gpmM*) disrupts carbon flow in the glycolytic/gluconeogenic backbone and generates two disconnected sub-networks: (1) a carbon fixation module containing upper glycolysis, the pentose phosphate pathway and the two foreign CBB enzymes; (2) an energy-module, containing lower

glycolysis and the TCA cycle supplying reducing power and ATP. In a scenario where an organic carbon source (e.g., pyruvate) is utilized by the energy module to supply the energetic demands of the cycle, the cellular building blocks derived from CBB-module metabolites (e.g., serine, pentose phosphates etc.) are synthesized from inorganic carbon using the non-native CBB pathway. The remainder of the biomass building blocks (those emanating from the energy module metabolites, e.g., valine, threonine) as well as the energy requirements of the cell, are supplied via the catabolism of the organic carbon source. In such a semiautotrophic growth mode, CO₂ and energy carriers are the sole inputs for the production of biomass precursors using the CBB cycle.

FIGs. 2A-E illustrate how chemostat evolution leads to a semiautotrophic phenotype. (a) Ancestor strain (left), containing *gpmA* and *gpmM* deletions was evolved in a xylose limited chemostat supplied with an excess of pyruvate and CO₂. Additional deletions of *pfkA* and *pfkB* and *zwf* results in RuBisCO dependent (but not semiautotrophic) catabolism of xylose in the initial heterotrophic growth (Figure 10). Propagation in a chemostat ensures that the xylose limited growth will result in a strong selective pressure towards achieving flux through the carbon fixation cycle. When mutations arise that create a fully functioning CBB cycle, they enable CO₂ to be the sole carbon input for the required biomass precursors. Xylose dependency is alleviated, allowing the semiautotrophic strain (right) to take over the population. (b) Around day 50 (arrow) an increase in OD and a decrease in pyruvate concentration (blue) was observed, indicating a takeover by an evolved clone. (c) In contrast to the ancestor strain, clones isolated from day 50 onwards were able to form colonies on agar plates supplemented with only pyruvate and requiring high CO₂ atmosphere. Evolved clones grew on pyruvate and CO₂ in liquid minimal media with a doubling time of ≈6 hours. (d) Isotopic labeling analysis, in which evolved clones were grown with isotopically labeled ¹³CO₂ and non-labeled pyruvate as an energy source, showed that biomass components synthesized from CBB module metabolites are almost fully labeled while biomass components originating from the energy module show low levels of labeling, as expected due to glycolytic cutoff, if the semiautotrophic growth mode is achieved. This indicates that the evolved strain is able to synthesize CBB module biomass precursors from CO₂ using the non-native CBB cycle, while the biomass precursors originating from the energy module are synthesized from the supplemented pyruvate. Note that threonine (Thr) is produced via an anaplerotic reaction that incorporates labeled inorganic carbon and thus the partial labeling observed was expected. (e) Isotopic carbon composition of total cell biomass was measured using an elemental mass analyzer. In the evolved semiautotrophic strain grown in labeled ¹³CO₂ atmosphere, over one third of cellular carbon originated from fixed inorganic carbon; this result is in line with the expected fraction of biomass precursors originating from the CBB module in the literature and corresponds to the fraction of biomass precursors produced from xylose in the ancestor strain under heterotrophic conditions (Figure 8). Labeling of carbon atoms in the ancestor strain grown with non-labeled xylose is due to the RuBisCO dependent catabolism of pentoses (Figure 10) and labeled carbon assimilation via anaplerotic reactions.

FIGs. 3A-B illustrate the dynamics of sequence evolution towards a semiautotrophic phenotype. (A) Temporal trajectories of mutations present in the evolving chemostat population. Single mutations and the average of mutation cohorts are plotted with a bold line; members of each cohort are dashed with matching color. The time at which semiautotrophic clones were first isolated from chemostat culture is marked with a dotted black line. Mutations that attained a frequency of at least 20% of the population are shown. Cell density and carbon source concentrations throughout the experiment are detailed in Figure 11B (B) A lineage tree of mutations in the population and a Muller diagram depicting the population dynamics as inferred from the temporal trajectories of mutations appearing in (A). Four rapid sweeps by mutations appear in the first 30 days (blue sectors) followed by a bifurcation into two subpopulation, the green and the purple lineage, each of which led to a semiautotrophic phenotype. Mutations marked with a star appeared in a subpopulation but were overtaken by clonal interference. Detailed description of each mutation is given in Tables 2-6 in the Examples section below, including few large chromosomal events that were identified in isolated clones but could not be quantified accurately by sequencing a whole chemostat heterogeneous population and do not appear in the figure.

FIGs. 4A-B illustrate that endogenous Pentose Phosphate Pathway (PPP) enzymes of *E. coli* can regenerate pentoses in a nonnative Calvin cycle. (A) Pentose regeneration in the Calvin cycle takes in most photosynthetic organisms by utilizing an aldolase and a designated Sedoheptulose-bisphosphate phosphatase (SBPase) to dephosphorylate Sedoheptulose-bisphosphate (SBP). Alternatively, pentose regeneration can occur using a bifunctional FBPase (fbp*) as was found to be the case in cyanobacteria. (B) While the *E. coli* genome does not contain bona fide SBPase, RuBP regeneration can be accomplished in SBP-independent manner using the native enzymes of the PPP.

FIGs. 5A-B illustrate the loss of carbon fixation in a host expressing RuBisCO and *prk* on a background of an unperturbed metabolic network. (A) Schematic representation of two possible flux modes for pentose catabolism in recombinant *E. coli* expressing RuBisCO and *prk*: native route via the pentosephosphate pathway (purple), and a RuBisCO dependent bypass from Ru5P to 3PG. While flux through the RuBisCO dependent branch is stoichiometrically feasible, it is not essential for growth due to the presence of the native route. (B) *In vivo* inorganic carbon assimilation was measured by incubating cells in M9 minimal media supplemented with unlabeled xylose (0.2%) and isotopically labeled ¹³CO₂. Assimilation of CO₂ into

biomass was quantified using mass spectrometry of amino acid isotopomers from protein hydrolysate sampled during growth. Upon expression of recombinant CBB enzymes, almost a sixth of the carbons composing the intracellular 3PG pool are labeled (indicated by the labeling of serine in blue and alanine in red), indicating significant flux through the RuBisCO dependent branch. However, after ~100 generations in exponential growth a loss of carbon fixation activity is observed, presumably due to counter selection and takeover of clones in which recombinant expression has been diminished. Labeling in wildtype (WT) sample (without the RuBisCOprk encoding plasmid, dashed) beyond the expected natural fraction of ^{13}C in the supplied xylose (gray) is due to the anaplerotic reactions of the TCA cycle.

FIGs. 6A-F illustrate the systematic identification of metabolic configurations in which pathway activity is coupled to cellular growth. (A) Metabolic toy model containing three endogenous reactions (V1-V3) and one target recombinant reaction (V_{target}). Flux through the target reaction is not essential for biomass generation on either of two considered carbon sources. (B) While frameworks for identifying gene deletion leading to the overproduction of a prespecified metabolite (such as OptKnock) ensure maximal flux through the target pathway at the point of maximal biomass production, maximal biomass production may not be the prevailing growth condition. Therefore, these frameworks do not ensure a global coupling between target flux and biomass production and therefore they do not make the target reaction essential for growth. The present inventors performed an exhaustive search for combinations of reaction knockouts in which V_{target} is essential for growth when metabolizing a specific carbon source. (C) For example, eliminating reactions V1 and V3 results in making V_{target} essential for any growth on either of the carbon sources. While metabolizing carbon source A, the selection slope, defined as V_{target}/V_B at the origin, is 0.5. (D) By changing the carbon source the selection slope is increased, now requiring 1 unit of target flux per biomass flux, and thus a tighter coupling. (E) Phenotypic phase space depiction of the feasible fluxes for each metabolic configuration. While the unmodified network allows biomass to be generated independently from the target flux (gray line), by eliminating reactions V1 and V3 the flux space is reduced only to flux modes in which biomass generation is coupled to the target flux; either with weak or strong coupling slope (purple and cyan lines, respectively) (F) If several alternatives lead to coupling, choosing knockouts combination that yields variable selection slopes on different carbon sources (red circle) is preferred, thus giving the experimenter options to increase the selection stringency in steps.

FIGs. 7A-B illustrate a systematic approach for identifying metabolic configurations, in which carbon fixation activity is essential for cell growth. (A) Each block (3x3 yellow grid) refers to a combination of two metabolic reactions to be knocked out within central carbon metabolism of *E. Coli*. The present inventors analyze the possible growth for each combination of reactions double knockout under 9 different carbon sources. The level of target flux dependence on biomass production is shown using a color code. Feasible fluxes are calculated for the perturbed metabolic network (Methods and Figure 6) using each of the nine different carbon sources as exchange inputs (in addition to CO_2). Metabolic configurations (a specific combination of knockout mutations and carbon source), in which carbon fixation activity is essential for cell growth are highlighted in green. The selection slope is defined as the minimal carbon fixation flux, required per unit of biomass produced. Higher slope (bright green), refers to a higher carbon fixation activity required for cell growth. Grey cells refer to configurations in which carbon fixation is not essential for growth, while black cells are configurations where growth is not possible. (B) The flux dependency phase spaces of four metabolic configurations (red boxed cells). A mutant which lacks the enzymatic activities of phosphofructokinase (*pfkA*, *pfkB*) and glucose6phosphate dehydrogenase (*zwf*) is predicted to grow on pyruvate (top, left) but without a need to use the target RuBisCO flux. In contrast, it will require the expression of RuBisCO and *prk* to metabolize pentoses (e.g. xylose, topright). An interesting scenario is the combined knockout of glucose6phosphate dehydrogenase (*zwf*) and phosphoglycerate mutase (*gpmA*, *gpmM*). In this mutant, expression of CBB enzymes allows growth on carbon sources which enter central carbon metabolism through lower glycolysis (e.g. pyruvate, bottom left). The metabolic cutoff prevents gluconeogenic carbon flow which supplies biomass precursors included in upper glycolysis and the pentose phosphate pathway. However, pyruvate can be metabolized in the TCA cycle to generate ATP and reducing power, hence supplying the energy to allow carbon fixation using the CBB cycle thus achieving a growth mode we term semiautotrophic growth. The RuBisCO dependent assimilation of CO_2 into biomass, supplies the demand for those biomass precursors which cannot be synthesized from organic carbon in a $\Delta gpm \Delta zwf$ mutant in order to allow growth. The present inventors aimed to experimentally test this metabolic scenario. Uptake rate was set to 10 mmol gCDW⁻¹h⁻¹ for xylose, and 16.7 for pyruvate.

FIG. 8 is a schematic representation of the main sinks for biomass precursor metabolites in central carbon metabolism. 12 precursor metabolites in glycolysis and the TCA cycle serve as metabolic branchpoints, in which flux branches out from central carbon metabolism to build the components of the cell's biomass. We detail the percentage (by mass) contributed to the total biomass derived from each of these key metabolites (as calculated based on F. C. Neidhardt et al., Physiology of the bacterial cell, pp. 142, 1990). This serves as a basis for predicting the expected labeled fraction in the semiautotrophic strain. In mutant *E. coli* lacking the activities of glucose phosphomutase (*gpmA*, *gpmM*) and glucose6phosphate dehydrogenase (*zwf*) central carbon metabolism is divided into two subnetworks disconnected in terms of carbon flow between them. Due to the cutoff, heterotrophic growth requires a minimum of two carbon sources: one feeding lower EMP pathway and the TCA cycle (energy module, blue), while the other feeding upper EMP pathway and the pentose phosphate pathway (CBB module, green). The relative percentage of the biomass emanating from precursor metabolites in the CBB module sums to approximately 30% of

total biomass in the cell (or almost equivalently, carbon biomass), while the metabolites derived from the energy module contribute the remaining 70%. Involved cells showing a semiautotrophic phenotype, CO₂ is the sole carbon input into the CBB module, hence about 30% of the cell's carbon is expected to be supplied from RuBisCO dependent assimilation of inorganic carbon.

FIGs. 9A-B illustrate carbon source combinations effecting growth and the condition for semiautotrophic growth. (A) Wild type *E. Coli* cells, with an unperturbed metabolic network, can grow on minimal media supplied with a single carbon source. In contrast, *E. Coli* mutant lacking glucose phosphomutase (*gpmA*, *gpmM*) and glucose6phosphate dehydrogenase (*zwf*) activities requires a minimum of two carbon sources, each feeding into one of the two disconnected sub networks formed by the metabolic cut off. While two carbon sources are supplied (e.g., xylose and pyruvate), lack of either one would limit growth, even in excess of the other. As shown in Figure 9B, per given concentration of supplied pyruvate (feeding the energy module), maximal biomass formation is initially proportional linear to the amount of supplied xylose. However, an excess of xylose beyond the point in which the ratio of carbon sources matches the consumption uptake ratio, does not contribute to the formation of additional biomass since growth becomes limited by the availability of pyruvate.

FIGs. 10A-C illustrate a computational prediction and experimental validation that phosphofructokinase (*pfkA*, *pfkB*) mutant is RuBisCO-dependent when growing on pentoses. (A) We systematically explored metabolic perturbations which couple RuBisCO-dependent carbon fixation to cellular growth even while an organic carbon source (e.g., xylose) is feeding the CBB module (as occurs during the initial phase of chemostat evolution). The computational analysis (Figure 7) predicts that a mutant lacking the activities of glucose6phosphate dehydrogenase (*zwf*) and phosphofructokinase (*pfkA* and *pfkB*) will require the activity of the CBB recombinant enzymes for growth on pentose sugars, but not on trioses. This result stems from the stoichiometry of the pentose phosphate pathway which converts three pentose sugars into two molecules of fructose6P (F6P) and a glyceraldehyde 3phosphate triose. Downstream glycolysis of F6P requires *pfk* activity to generate fructose1,6P and feed the glycolytic pathway. Therefore, *pfk* mutant is predicted not to grow on pentose sugars as a single carbon source. Recombinant expression of RuBisCO and *prk* creates a *pfk* independent route from pentoses to trioses through the RuBisCO-dependent carboxylation of ribulose biphosphate to 3phosphoglycerate, hence rescuing growth. In contrast to pentose metabolism, trioses feeding into glycolysis (e.g., pyruvate, glycerol) do not require *pfk* activity for catabolism nor for the gluconeogenic biosynthesis of hexose phosphates which takes place via fructose 1,6bisphosphatase (*fbp*). Uptake rate was set to 10 mmol gCDW⁻¹h⁻¹ for xylose, and 16.7 for pyruvate and glycerol. (B) To experimentally validate the computational predictions, we constructed $\Delta zwf \Delta pfk$ mutants and tested the effect of RuBisCO and *prk* expression while growing either on pentoses or trioses. As predicted, the RuBisCO-dependent bypass is essential for growth on pentoses. On permissive carbon sources such as glycerol or pyruvate, growth is independent from RuBisCO expression. (C) The ancestor strain ($\Delta gpmAM \Delta zwf \Delta pfkAB::rbcLprk$) is dependent on RuBisCO activity even while xylose is feeding the CBB module, as occurs during the initial state of chemostat evolution. Due to the glycolytic cutoff imposed by the lack of *gpm* activity (Figure 1) the ancestor strain requires a second carbon source feeding into the energy module, in addition to xylose.

FIGs. 11A-D illustrate mass isotopomer distributions shows fixation of unlabeled CO₂ when cells are grown on fully labeled pyruvate. (A-C) Clones displaying semiautotrophic phenotype (i.e., capable of growing on pyruvate and CO₂) were isolated from three independent chemostat experiments. Cells were cultured on M9 minimal media containing no organic carbon except uniformly labeled ¹³C pyruvate (5 g/L), in gas controlled environment with 0.1 atm nonlabeled CO₂ (Methods). Mass spectrometry analysis of amino acids derived from precursor metabolites originating either from the CBB module (serine, histidine) or the energy module (valine, threonine) shows the expected labeling pattern from semiautotrophic growth mode: trace labeling in serine and histidine synthesized from unlabeled CO₂, in contrast to full labeling in valine and threonine which are synthesized from organic carbon. (D) In wild type sample all amino acids are fully labeled since they are directly synthesized from the labeled pyruvate. Figure 11A: Isolated from first chemostat experiment, day 50; Figure 11B: Isolated from second chemostat experiment, day 130; Figure 11C: Isolated from third chemostat experiment, day 70; Figure 11C: Wild-type control.

FIG. 12 illustrates a mass spectrometry analysis of intracellular metabolites pool validates CBB cycle dependent biosynthesis of biomass precursors from CO₂. Clones isolated from three independent chemostat experiments were cultured in minimal media supplemented only with uniformly labeled ¹³C pyruvate and non labeled CO₂. Intracellular metabolites were extracted from samples taken during exponential growth, and analysed using mass spectrometry (Methods). Hexose phosphates (i.e., glucose6phosphate, fructose6phosphate) resulting from the assimilation of CO₂ in the CBB module were not labeled, while phosphoenolpyruvate, resulting from pyruvate metabolism in the energy module, was fully labeled. This result is in line with the expected labeling pattern from a semiautotrophic growth mode, in which CBB cycle intermediate are originated solely from the assimilation of inorganic carbon.

FIGs. 13A-C illustrate evolution dynamics of *E. Coli* towards semiautotrophic growth in three independent chemostat experiments. (A) Ancestor strain ($\Delta pfk \Delta gpm \Delta zwf::rbcLprk$) was cultured in xylose limited chemostat with a dilution rate of 0.08

h1 (Methods). Xylose concentration in the feed media was kept constant at 100 mg/L (purple dashed line). Pyruvate, the organic substrate feeding the energy module, was provided in excess and its concentration in the feed media was 5 g/L. The residual concentration of these carbon sources, and the cell density inside the chemostat were measured by routine sampling. Once initial steadystate was reached, residual xylose concentration fell below detection limit, while cell density remained relatively constant for the next 50 days. From this point onwards, cell density increased while the residual pyruvate levels decreased. In addition, culture samples from day 50 onwards were able to grow on liquid M9 minimal media, as well as on agar plates, in elevated CO₂ conditions (pCO₂= 0.2) while supplemented solely with pyruvate. The number of colonies formed increased from day 50, reaching saturation at approximately day 70. (B) In the second experiment, the ancestor strain was propagated in the same conditions excluding xylose feed regime. Here xylose concentration was decreased in the feed media whenever residual pyruvate concentration in the culture fell below detection limit, avoiding the possibility that pyruvate becomes the limiting nutrient instead of xylose. Clones with semiautotrophic phenotype were first isolated on day 130. Xylose was completely omitted from the feed media from day 150 onwards, validating the ability of cells to grow in the complete absence of an organic carbon source feeding into the CBB module. Every two weeks culture samples were collected, and later subjected for wholegenome wholepopulation sequencing. (C) For the third experiment, the ancestor strain was further modified by knocking out a component of mismatch repair system (*mutS*) known to drastically increase the mutation rate. In addition, the dilution rate was set to 0.035 h⁻¹. Xylose concentration in the feed was set to 25 mg/L and reduced whenever pyruvate concentration in the culture approached the detection limit as described before. Clones displaying semiautotrophic phenotype were isolated from day 55 onwards.

DESCRIPTION OF SPECIFIC ASPECTS OF THE DISCLOSURE

[0009] The present disclosure, in some aspects thereof, relates to recombinant non-autotrophic microorganisms that are capable of carbon fixation.

[0010] Before explaining at least one aspect of the disclosure in detail, it is to be understood that the disclosure is not necessarily limited in its application to the details set forth in the following description or exemplified by the Examples. The disclosure is capable of other aspects or of being practiced or carried out in various ways.

[0011] Carbon fixation is the process by which carbon dioxide is incorporated into organic compounds. In the process of transforming sunlight into biological fuel, plants absorb carbon dioxide using over 70 % of the fresh water utilized by humanity and the majority of cultivatable land resources on earth. These figures point to the central place that carbon fixation by plants plays in our global ecological footprint.

[0012] Carbon fixation in plants and algae is achieved by the Calvin-Benson Cycle (CBBC). The productivity of the Calvin-Benson cycle is limited, under many conditions, by the slow rate and lack of substrate specificity of the carboxylating enzyme Rubisco. As carbon fixation has been shown to limit growth rate in many studies, the present inventors sought to develop alternative pathways that can support carbon fixation in organisms whose genome can be easily manipulated such as *E. Coli*, with the aim to using such bacteria as a platform to experimentally investigate different aspects of the carbon fixation pathways. For example, the microorganism can be used as a screen for identifying engineered enzymes for enhanced carbon fixation properties. Once identified, such enzymes may be expressed in naturally occurring carbon fixing organisms (such as plants) so as to improve carbon fixation.

[0013] To accomplish that, the present inventors expressed the two enzymes of the Calvin-Benson cycle which are absent in wild-type *E. Coli*: phosphoribulokinase (PRK) and ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO). Upon expressing the two necessary genes, the present inventors were able to detect PRK- and RuBisCO-dependent assimilation of inorganic ¹³CO₂ (see Figures 3A-C.) However, this activity was found to be unstable; after less than 50 generations of continuous cultivation at heterotrophic conditions, cells lost all newly gained carbon fixation activity (Figure 3D). This could be attributed to the protein load imposed by the overexpression of the foreign genes, whose activities do not contribute bacterial fitness when the endogenous metabolic network remains intact.

[0014] The present inventors therefore systematically explored all gene deletions that result in the activities of PRK and RuBisCO being essential even for a heterotrophically grown bacterium using a constraint-based modeling framework (Figure 4). From the in silico results, the present inventors established that by combining several gene transfer events, and gene deletions, it is theoretically possible to produce an *E. Coli* strain which obtains its energy from metabolizing an organic compound, while building a portion of its cellular building blocks from carbon fixation.

[0015] The present inventors implemented this strategy and engineered a number of different *E. Coli* strains which expressed two disconnected sub-networks of central metabolism - (1) the CBBC-module, containing the upper glycolysis, the pentose phosphate pathway and the two foreign enzymes; (2) the energy-module, containing the lower glycolysis and the TCA cycle. However, these strains failed to grow semiautotrophically at elevated CO₂, when supplied only with a substrate of the energy-module.

[0016] Thus, for example the present inventor showed that the $\Delta\text{pfkA}\Delta\text{pfkB}\Delta\text{gpmA}\Delta\text{gpmM}\Delta\text{zwf}:\text{PRK-RuBisCO}$ strain was able to grow only in the presence of two carbon sources; one which enters the CBBC-module and the second which enters the energy-module, e.g., xylose and pyruvate, respectively.

[0017] The present inventors hypothesized that the while all the necessary enzymatic components for carbon fixation were present and active, the organism failed to grow semiautotrophically (in the absence of xylose) because a fine tuning of flux distribution is required to support a well-balanced activity of an autocatalytic cycle (a cycle whose product is an intermediate of the cycle), such as the CBBC.

[0018] Hence, the present inventors harnessed the natural selection of the *E. Coli* to optimize and balance pathway activity towards establishing semiautotrophic growth. They cultivated the cells in an environment in which the substrate of the energy-module, as well as CO₂, are in excess, while the substrate of the CBBC-module is limiting. This, they showed, imposed a strong selective pressure on the cells to become dependent on the carbon fixation activity of the CBBC: Semiautotrophs which no longer required the organic substrate of the CBBC-module took over the population.

[0019] Thus according to a first aspect of the present disclosure there is provided a microorganism which is genetically modified so that it produces a first essential biomass precursor by metabolizing CO₂ using a recombinant carbon fixation enzyme, wherein the microorganism produces a second biomass precursor by metabolizing an organic carbon source and not by metabolizing CO₂, wherein the microorganism does not use the organic carbon source for producing the first essential biomass precursor.

[0020] The microorganism of this aspect of the present disclosure generates ATP and a reducing power (e.g. NADH) via metabolism of the organic carbon source.

[0021] As used herein, the term "microorganism" refers to any organism of microscopic size which, in its native state, is not capable of biosynthesizing metabolites by utilizing CO₂ solely as a carbon source.

[0022] Thus, for example the scope of the present disclosure does not include cyanobacteria or *Rhodospirillum Rubrum*, which are capable of carbon fixation in their native state. Non-limiting examples of microorganisms include both prokaryotic and eukaryotic microorganisms, such as bacteria, archae, protozoan, fungi, molds, yeasts, algae etc. The microorganism may be aerobic or anaerobic.

[0023] The organisms can be fermentative organisms. Exemplary microorganisms include, for example, *Clostridium* (e.g., *C. acetobutylicum*, *C. Beijerinckii*, *C. saccharoperbutylacetonicum*, *C. saccharobutylicum*, *C. aurantibutyricum*, *C. tetanomorphum*), *Zymomonas*, *Escherichia* (e.g., *E. Coli*), *Salmonella*, *Rhodococcus*, *Pseudomonas*, *Bacillus*, *Lactobacillus*, *Enterococcus*, *Alcaligenes*, *Klebsiella*, *Paenibacillus*, *Arthrobacter*, *Corynebacterium*, *Brevibacterium*, *Pichia*, *Candida*, *Hansenula*, *Zymomonas* and *Saccharomyces*, e.g., *Saccharomyces cerevisiae*, *Saccharomyces carlsbergensis*, *Kluyveromyces lactis*, *Saccharomyces lactis*.

[0024] The bacteria may be those which are useful in the food industry. For example lactic Acid Bacteria (LAB) play an essential role in the preservation, taste and texture of cheese, yogurt, sausage, sauerkraut and a large variety of traditional indigenous fermented foods.

[0025] Bacteria may be gram positive or gram negative. Examples of bacteria which are contemplated by the present disclosure include, but are not limited to *Agrobacterium*, *Alicyclobacillus*, *Anabaena*, *Anacystis*, *Arthrobacter*, *Azobacter*, *Bacillus*, *Brevibacterium*, *Chromatium*, *Clostridium*, *Corynebacterium*, *Enterobacter*, *Erwinia*, *Escherichia*, *Lactobacillus*, *Lactococcus*, *Mesorhizobium*, *Methylobacterium*, *Microbacterium*, *Phormidium*, *Pseudomonas*, *Rhodobacter*, *Rhodopseudomonas*, *Rhodospirillum*, *Rhodococcus*, *Salmonella*, *Scenedesmun*, *Serratia*, *Shigella*, *Staphylococcus*, *Streptomyces*, *Synnecoccus*, and *Zymomonas*.

[0026] Examples of fungi contemplated by the present disclosure include, but are not limited to *Aspergillus*, *Candida*, *Chlamydomonas*, *Chrysosporium*, *Cryptococcus*, *Fusarium*, *Kluyveromyces*, *Neotyphodium*, *Neurospora*, *Penicillium* (e.g. *P.*

chrysogenum), *Pichia*, *Saccharomyces*, *Trichoderma* and *Xanthophyllomyces*.

[0027] The term "carbon fixation" as used herein refers to a process through which inorganic carbon (e.g. gaseous carbon dioxide or bicarbonate) is assimilated into an organic backbone. Carbon fixation results in the transfer from an inorganic carbon pool to an organic cellular carbon pool.

[0028] The term "enzyme" as used herein refers to a "catalytically functional biomolecule," which includes both whole native (or native-size) molecules, naturally occurring enzymes and derivatives (e.g. genetic modifications) thereof.

[0029] As used herein, the phrase "carbon fixation enzyme" refers to an enzyme which is part of a carbon fixation pathway or cycle.

[0030] As used herein the phrase "carbon fixation pathway" refers to a set of molecules (e.g. enzymes, electron donors, co-factors etc.) that together enable carbon fixation. The pathway may be linear or cyclical.

[0031] An exemplary carbon fixation pathway is the Calvin-Benson-Bassham (CBB) Cycle.

[0032] This cycle can be divided into three phases:

Phase 1: Carbon fixation. The enzyme Rubisco (ribulose biphosphate carboxylase) catalyses the carboxylation of ribulose-1,5-bisphosphate in a two step reaction. Ribulose-1,5-bisphosphate must first be phosphorylated by the enzyme Phosphoribulose kinase. The outcome of this carboxylation are two molecules of 3-Phosphoglycerate.

Phase 2: Reduction. 3-Phosphoglycerate is then phosphorylated with the aid of the enzyme phosphoglycerate kinase to yield 1,3-Bisphosphoglycerate. Next 1,3-Bisphosphoglycerate is reduced by NADPH to yield NADP⁺ and Glyceraldehyde-3-phosphate with the aid of Glyceraldehyde-3-phosphate dehydrogenase. One of every six Glyceraldehyde-3-phosphate molecules is exported into the cytoplasm to be use in the synthesis of Glucose and other metabolic processes.

Phase 3: Regeneration of ribulose. Glyceraldehyde-3-phosphate is then reversibly converted to Dihydroxyacetone phosphate by Triose phosphate isomerase. Next Dihydroxyacetone is converted into fructose-6-phosphate (F-6-P) by Aldolase and Fructose biphosphatase. Aldolase condenses the two DHAP molecules to form Fructose-1,6-bisphosphate. Because of its high (-)delta G the transformation of Fructose-1,6-bisphosphate to Fructose-6-phosphate is thought to be the rate limiting step of the CBB cycle. F-6-P can then be converted into glucose via two enzymatic steps with the help of Phosphoglucoisomerase and glucose-6-Phosphatase. Dihydroxyacetone can also go on to condense with Erythrose-4-phosphate to form Sedoheptulose-1,7-bisphosphate(SBP). This reaction is also catalyzed by Aldolase. SBP is then de-phosphorolated by Sedoheptulose biphosphatase to yield Sedoheptulose-7-phosphate (S7P). After several rearrangement reactions utilizing Transketolase and Transaldolase enzymes, Xylulose-5-Phosphate (X5P) and Ribose-5-phosphate (R5P) are synthesized. Lastly X5P and R5P are isomerised by Phosphopentose epimerase and Phosphopentose isomerase to yield Ribulose-5-phosphate which can then be put back into the cycle.

[0033] According to particular aspects, the microorganism is genetically modified to express at least one enzyme of a carbon fixation pathway (e.g. Calvin-Benson-Bassham (CBB) Cycle) such that the carbon fixation pathway is active in the microorganism (i.e. utilizes carbon dioxide through the non-native carbon fixation pathway) for the production of an essential biomass precursor. As such the recombinant enzymes of this aspect of the present disclosure are expressed such that they are positioned relative to one another such that they are able to function to cause carbon fixation. According to this aspect the enzymes which take part in the carbon fixation pathways are present in the same component of the cell such that they are able to cooperate with one another in order to fulfill their role in the carbon fixation pathway.

[0034] The exact carbon fixation enzymes to be expressed in a particular microorganism will vary according to the enzymes which are natively expressed in that microorganism.

[0035] Thus, for example, out of the 12 reactions of the CBBC, only two are catalyzed by enzymes which are absent in *E. Coli*: phosphoribulokinase (PRK; EC 2.7.1.19) and ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO). The remaining reactions are catalyzed by the native gluconeogenic and pentose phosphate pathway's enzymes. Notably, while *E. Coli* does not contain sedoheptulose-bisphosphate aldolase and sedoheptulose-bisphosphatase, these can be bypassed by the native transketolase enzyme. Thus, for *E. Coli*, the present disclosure contemplates expressing both PRK and RuBisCO.

[0036] The enzymes may be of any origin.

[0037] Thus an enzyme of the present disclosure also refers to homologs and other modifications including additions or deletions of specific amino acids to the sequence (e.g., polypeptides which are at least 50 %, at least 55 %, at least 60 %, at least 65 %, at least 70 %, at least 75 %, at least 80 %, at least 85 %, at least 87 %, at least 89 %, at least 91 %, at least 93 %, at least 95 % or more say 100 % homologous to the native amino acid sequence of the enzyme, as determined using BlastP software of the National Center of Biotechnology Information (NCBI) using default parameters). The homolog may also refer to an ortholog, a deletion, insertion, or substitution variant, including an amino acid substitution, thereof and biologically active polypeptide fragments thereof.

[0038] Thus RuBisCo may for example be a *Rhodospirillum rubrum* RuBisCo being encoded by a sequence as set forth in SEQ ID NO: 1 or having an amino acid sequence as set forth in SEQ ID NO: 4. The amino acid sequence may be at least 90 % identical, at least 91 % identical, at least 92 % identical, at least 93 % identical, at least 94 % identical, at least 95 % identical, at least 96 % identical, at least 97 % identical, at least 98 % identical, at least 99 % identical to SEQ ID NO: 4.

[0039] The PRK enzyme may be *Synechococcus* PRK being encoded by a sequence as set forth in SEQ ID NO: 2 or having an amino acid sequence as set forth in SEQ ID NO: 5. The amino acid sequence may be at least 90 % identical, at least 91 % identical, at least 92 % identical, at least 93 % identical, at least 94 % identical, at least 95 % identical, at least 96 % identical, at least 97 % identical, at least 98 % identical, at least 99 % identical to SEQ ID NO: 5. Alternatively, the RuBisCo and PRK may be expressed using an operon of the proteobacteria *Ralstonia eutropha*, which contains all the Calvin-Benson Cycle genes in tandem.

[0040] According to a particular aspect, the RuBisCo is a plant-derived RuBisCo.

[0041] The present inventors further contemplate expressing a recombinant carbonate dehydratase in the microorganisms of the present disclosure. By doing so, the microorganism becomes capable of utilizing inorganic carbon which is present in the bicarbonate form in order to generate the first biomass precursor.

[0042] The carbonate dehydratase may be a *Rhodospirillum rubrum* carbonate dehydratase being encoded by a sequence as set forth in SEQ ID NO: 3 or having an amino acid sequence as set forth in SEQ ID NO: 6. The amino acid sequence may be at least 90 % identical, at least 91 % identical, at least 92 % identical, at least 93 % identical, at least 94 % identical, at least 95 % identical, at least 96 % identical, at least 97 % identical, at least 98 % identical, at least 99 % identical to SEQ ID NO: 6.

[0043] To express the enzymes of the present disclosure using recombinant technology, a polynucleotide encoding the enzymes is ligated into a nucleic acid expression vector, which comprises the polynucleotide sequence under the transcriptional control of a cis-regulatory sequence (e.g., promoter sequence) suitable for directing constitutive, tissue specific or inducible transcription of the polypeptides of the present disclosure in the host cells.

[0044] The polynucleotide may further comprise ribosome binding sites to differentially control the expression level of the genes. Thus, for example for *rbcL*, the ribosome binding site rbs-C may be used, for *prkA* the ribosome binding site rbs-E may be used and for *CA* the ribosome binding site rbs-C may be used.

[0045] Thus, the present disclosure contemplates isolated polynucleotides encoding the enzymes of the present disclosure.

[0046] The phrase "an isolated polynucleotide" refers to a single or double stranded nucleic acid sequence which is isolated and provided in the form of an RNA sequence, a complementary polynucleotide sequence (cDNA), a genomic polynucleotide sequence and/or a composite polynucleotide sequences (e.g., a combination of the above).

[0047] As used herein the phrase "complementary polynucleotide sequence" refers to a sequence, which results from reverse transcription of messenger RNA using a reverse transcriptase or any other RNA dependent DNA polymerase. Such a sequence can be subsequently amplified *in vivo* or *in vitro* using a DNA dependent DNA polymerase.

[0048] As used herein the phrase "genomic polynucleotide sequence" refers to a sequence derived (isolated) from a chromosome and thus it represents a contiguous portion of a chromosome.

[0049] As used herein the phrase "composite polynucleotide sequence" refers to a sequence, which is at least partially complementary and at least partially genomic. A composite sequence can include some exon sequences required to encode the polypeptide of the present disclosure, as well as some intronic sequences interposing therebetween. The intronic sequences can be of any source, including of other genes, and typically will include conserved splicing signal sequences. Such intronic sequences may further include cis acting expression regulatory elements.

[0050] Nucleic acid sequences encoding the enzymes and enzyme activating proteins of some aspects of the disclosure may be optimized for expression for a particular microorganism. Examples of such sequence modifications include, but are not limited to, an altered G/C content to more closely approach that typically found in the microorganism species of interest, and the removal of codons atypically found in the microorganism species commonly referred to as codon optimization.

[0051] The phrase "codon optimization" refers to the selection of appropriate DNA nucleotides for use within a structural gene or fragment thereof that approaches codon usage within the microorganism of interest. Therefore, an optimized gene or nucleic acid sequence refers to a gene in which the nucleotide sequence of a native or naturally occurring gene has been modified in order to utilize statistically-preferred or statistically-favored codons within the microorganism. The nucleotide sequence typically is examined at the DNA level and the coding region optimized for expression in the microorganism species determined using any suitable procedure, for example as described in Sardana et al. (1996, *Plant Cell Reports* 15:677-681). In this method, the standard deviation of codon usage, a measure of codon usage bias, may be calculated by first finding the squared proportional deviation of usage of each codon of the native gene relative to that of highly expressed genes, followed by a calculation of the average squared deviation. The formula used is: $1 \text{ SDCU} = n = 1 \text{ N} [(X_n - Y_n) / Y_n]^2 / N$, where X_n refers to the frequency of usage of codon n in highly expressed genes, where Y_n to the frequency of usage of codon n in the gene of interest and N refers to the total number of codons in the gene of interest.

[0052] One method of optimizing the nucleic acid sequence in accordance with the preferred codon usage for a particular cell type is based on the direct use, without performing any extra statistical calculations, of codon optimization tables such as those provided on-line at the Codon Usage Database through the NIAS (National Institute of Agrobiological Sciences) DNA bank in Japan (wwwdotkazusadotordotjp/codon/). The Codon Usage Database contains codon usage tables for a number of different species, with each codon usage table having been statistically determined based on the data present in Genbank.

[0053] By using the above tables to determine the most preferred or most favored codons for each amino acid in a particular species (for example, *E. Coli*), a naturally-occurring nucleotide sequence encoding a protein of interest can be codon optimized for that particular species. This is effected by replacing codons that may have a low statistical incidence in the particular species genome with corresponding codons, in regard to an amino acid, that are statistically more favored. However, one or more less-favored codons may be selected to delete existing restriction sites, to create new ones at potentially useful junctions (5' and 3' ends to add signal peptide or termination cassettes, internal sites that might be used to cut and splice segments together to produce a correct full-length sequence), or to eliminate nucleotide sequences that may negatively effect mRNA stability or expression.

[0054] The naturally-occurring encoding nucleotide sequence may already, in advance of any modification, contain a number of codons that correspond to a statistically-favored codon in a particular species. Therefore, codon optimization of the native nucleotide sequence may comprise determining which codons, within the native nucleotide sequence, are not statistically-favored with regards to a particular plant, and modifying these codons in accordance with a codon usage table of the particular species to produce a codon optimized derivative. A modified nucleotide sequence may be fully or partially optimized for microorganism codon usage provided that the protein encoded by the modified nucleotide sequence is produced at a level higher than the protein encoded by the corresponding naturally occurring or native gene. Construction of synthetic genes by altering the codon usage is described in for example PCT Patent Application No. 93/07278.

[0055] As mentioned hereinabove, polynucleotide sequences of the present disclosure are inserted into expression vectors (i.e., a nucleic acid construct) to enable expression of the recombinant polypeptide. The expression vector of the present disclosure includes additional sequences which render this vector suitable for replication and integration in prokaryotes, eukaryotes, or preferably both (e.g., shuttle vectors). Typical cloning vectors contain transcription and translation initiation sequences (e.g., promoters, enhances) and transcription and translation terminators (e.g., polyadenylation signals).

[0056] Various methods can be used to introduce the expression vector of the present disclosure into the host cell system. Such methods are generally described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Springs Harbor Laboratory, New York (1989, 1992), in Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley and Sons, Baltimore, Md. (1989), Chang et al., *Somatic Gene Therapy*, CRC Press, Ann Arbor, Mich. (1995), Vega et al., *Gene Targeting*, CRC Press, Ann Arbor Mich. (1995), *Vectors: A Survey of Molecular Cloning Vectors and Their Uses*, Butterworths, Boston Mass. (1988) and Gilboa et al. [*Biotechniques* 4 (6): 504-512, 1986] and include, for example, stable or transient transfection, lipofection, electroporation and infection with recombinant viral vectors. In addition, see U.S. Pat. Nos. 5,464,764 and 5,487,992 for positive-negative selection methods.

[0057] Exemplary bacterial based expression systems are disclosed in Baneyx et al., *Current Opinion in Biotechnology*, 1999; 10, 411-421 and Macrides et al, *Microbiol Rev* 1996, 60: 512-538.

[0058] Contemplated promoters for expression in bacteria include the l-arabinose inducible araBAD promoter (PBAD), the lac promoter, the l-rhamnose inducible rhaP BAD promoter, the T7 RNA polymerase promoter, the trc and tac promoter, the lambda phage promoter pL, and the anhydrotetracycline-inducible tetA promoter/operator.

[0059] The microorganisms may be transformed stably or transiently with the nucleic acid constructs of the present disclosure. In stable transformation, the nucleic acid molecule of the present disclosure is integrated into the microorganism genome and as such it represents a stable and inherited trait. In transient transformation, the nucleic acid molecule is expressed by the cell transformed but it is not integrated into the genome and as such it represents a transient trait.

[0060] Knock-in methods for expressing a gene in a microorganism are also contemplated.

[0061] It will be appreciated that the microorganisms of this aspect of the present disclosure are not fully autotrophic since they rely on an organic carbon source to produce energy (i.e. ATP). Examples of organic carbon sources include, but are not limited to acetate, pyruvate, malate, succinate, fumarate, oxaloacetate, xylose and citrate. Selection of a particular carbon source will depend on the particular microorganism and the genetic modifications thereof as further described herein below.

[0062] As mentioned, the microorganisms of this aspect of the present disclosure utilize carbon dioxide, which is assimilated via the non-native enzymes, to generate at least one biomass precursor, while utilizing an organic carbon source to generate other (i.e. non-identical) biomass precursors. The "at least one biomass precursor" may be derived from organic carbon sources as well, so long as it is not the same organic carbon source used to generate the "other biomass precursors". By way of an example, the microorganism is modified such that it utilizes carbon dioxide which is assimilated through non-native Calvin cycle enzymes to generate biomass precursors such as 5-phospho-D-ribose α -1-pyrophosphate (PRPP), glucose-6P, fructose-6P, erythrose-4P, 3-phosphoglycerate, glycerol-3-phosphate and aromatic amino acids (referred to herein as group 1 biomass precursors). Other biomass precursors (e.g. non-aromatic amino acids, also referred to herein as group 2 biomass precursors), are derived from an organic carbon source (e.g. pyruvate). Additional examples of biomass precursors are provided in Figure 8, the group 1 biomass precursors being colored in green and the group 2 biomass precursors being colored in blue. The group 1 biomass precursors may also be generated from an organic carbon source, so long as it is not the same organic source used to generate the group 2 biomass precursors (e.g. not pyruvate).

[0063] According to a particular aspect, group 1 biomass precursors are derived from carbon dioxide and xylose and group 2 biomass precursors are derived from pyruvate.

[0064] According to a yet another aspect, group 1 biomass precursors are derived from carbon dioxide and glycerol and group 2 biomass precursors are derived from pyruvate.

[0065] According to still another aspect, the organic carbon source from which group 2 biomass precursors are generated does not comprise a hexose or pentose sugar.

[0066] In order to prevent the microorganism from utilizing the organic carbon source which is used for energy production (and group 2 biomass precursors) for the generation of group 1 biomass precursors, the microorganism is typically further genetically modified.

[0067] Determining which additional genetic modifications should be made to a particular microorganism can be carried out by and identifying sets of knock outs, gene additions and media composition, which ensure that the flux through the carbon fixation pathway is essential for cell growth. This may be effected computationally - e.g. using a constraint-based modeling framework, as described in the Examples section below.

[0068] Thus, for example, for *E. Coli*, the present inventors have determined that according to one aspect, in order to prevent utilization of the organic carbon source which is used for energy production and the generation of group 2 biomass precursors for the generation of group 1 biomass precursors, down-regulation of expression or activity of phosphofructokinase (pfk) or ribose-5-phosphate isomerase (rpi; EC 5.3.1.6) may be effected. Preferably down-regulation of both pfkA (EC 2.7.1.11) and pfkB (EC 2.7.1.105) is effected.

[0069] Additionally, or alternatively down-regulation of expression or activity of phosphoglycerate mutase (gpm; EC 5.4.2.11) may be carried out to prevent utilization of the organic carbon source which is used for energy production and the generation of group 2 biomass precursors for the generation of group 1 biomass precursors. Preferably downregulation of both gpmA and gpmB is effected.

[0070] At least one member of the Entner-Doudoroff (ED) pathway and/or the oxidative pentose P pathway may optionally be knocked out (or down-regulated) to enforce Rubisco dependence via PFK. Thus, for example the genes 6-phosphogluconate dehydratase (edd) and ketohydroxyglutarate aldolase (eda) may be knocked out to prevent utilization of the ED pathway and glucose 6-phosphate-1-dehydrogenase (zwf) may be knocked out to prevent utilization of the of oxidative P pathway).

[0071] Furthermore downregulation of the operon AceBAK can be effected so as to prevent utilization of acetate for the generation of group 1 biomass precursors. Thus, knock-out of malate synthase A (aceB), isocitrate lyase (aceA) and/or isocitrate dehydrogenase kinase/phosphatase (aceK) is also contemplated.

[0072] Down-regulation of expression or activity of genes in the microorganism may be effected using any method known in the art.

[0073] Methods of deleting or downregulating genes from the chromosome of microorganisms are known to those of skill in the art and include homologous recombination, knock out techniques, RNAi etc.

[0074] For bacteria, methods such as P1 transduction from already existing knockout strains (KEIO collection) or via lambda-phage assisted recombination (Pkd46 system) may be used to knock-out specific genes.

[0075] Other methods which include down-regulating genes in microorganisms using CRISPR arrays are also contemplated. These methods are described for example in WO 2012164565.

[0076] For *E. Coli*, the following strains summarized in Table 1 are contemplated:

Table 1

Knockouts							
KO7	KO6	KO5	KO4	KO3	KO2	KO1	strain
		glcB	aceBAK	pfkB	pfkA	zwf	#1
pck	pps	glcB	aceBAK	pfkB	pfkA	zwf	#2
	gpmB	gpmA	aceBAK (optional)	pfkB	pfkA	zwf	#3
		eno	aceBAK	pfkB	pfkA	zwf	#4

[0077] It is appreciated that the strains described in Table 1 (which have been genetically modified to express phosphoribulokinase (prk) and Ribulose-Bisphosphate Carboxylase (RuBisCo), and optionally carbonate dehydratase may be further modified. This may be effected by performing random mutagenesis on the strains (e.g. as described in the Examples section herein below) and/or by deleting a component of mismatch repair system genes (mutS). Such strains may then be propagated under particular conditions so as to harness the natural selection of the *E. Coli* to optimize and balance pathway activity, as further described herein below.

[0078] Alternatively, randomly mutated strains of *E. Coli* may be used as the starting material on which the deletions and modifications described herein above are performed.

[0079] As mentioned, in order to propagate the microorganisms of this aspect of the present disclosure, they are cultured in a medium which comprise both an organic carbon source to feed the TCA cycle (for the production of ATP and a reducing power (e.g. NADH) and a non-organic carbon source (e.g. carbon dioxide) to supply the building blocks for biomass precursors originating from upper glycolysis and Pentose Phosphate intermediates - group 1 biomass precursors.

[0080] The organic carbon source is selected depending on the particular strain of the microorganism and the genetic modifications carried out. Thus, for strain 1 in Table 1 herein above, the *E. Coli* may be cultured in a medium comprising acetate. For Strains 2, 3 or 4, the *E. Coli* may be cultured in a medium comprising pyruvate, malate, succinate, fumarate, oxaloacetate or citrate. According to one aspect, the culture medium comprises a single organic carbon source (e.g. pyruvate).

[0081] As mentioned, the microorganisms may rely solely on carbon dioxide to supply the building blocks for biomass precursors originating from upper glycolysis and Pentose Phosphate intermediates (group 1 biomass precursors) or may rely on a second organic carbon source (pentose or hexose sugar) together with the carbon dioxide to produce those biomass precursors. The former state is referred to herein as a "semi-autotrophic state".

[0082] Contemplated pentose or hexose sugars include for example glycerol or xylose.

[0083] Thus according to another aspect, the culture medium comprises two organic carbon sources (e.g. pyruvate and glycerol; or pyruvate and xylose).

[0084] The present inventors have found that in order to generate microorganisms that rely solely on carbon dioxide to supply the building blocks for biomass precursors originating from upper glycolysis and Pentose Phosphate intermediates it is preferable to initially culture the microorganism in a culture which comprises an organic carbon source and a hexose or pentose sugar, wherein the ratio of organic carbon source: hexose or pentose sugar is at least about 10:1. The gas atmosphere may be manipulated such that the carbon dioxide is provided at saturating levels (e.g. at a minimum of 20 %).

[0085] Preferably, the microorganisms are grown for at least one day, at least two days, at least three days, at least one week, at least one month, at least three months following genetic modification in order for the generation of new strains which are further adapted for semi-autotrophic growth.

[0086] The amount of hexose or pentose sugar may be gradually reduced such that eventually, the microorganism relies solely on the carbon dioxide to supply the building blocks for the generation of biomass precursors originating from upper glycolysis and Pentose Phosphate intermediates (group 1).

[0087] Typically, the medium also comprises an electron acceptor such as nitrate, sulfate or oxygen at low levels (1-5 %). The medium may also comprise appropriate salts, minerals, metals and other nutrients, such as vitamins. Cells of the present disclosure can be cultured in conventional fermentation bioreactors, shake flasks, test tubes, microtiter dishes and petri plates. Culturing can be carried out at a temperature, pH and oxygen content appropriate for a recombinant cell. Such culturing conditions are within the expertise of one of ordinary skill in the art.

[0088] According to one aspect, the microorganisms are cultured on a solid surface - e.g. agarose plates.

[0089] The microorganisms may be immobilized on to a solid surface - e.g. filters and the like.

[0090] According to another aspect, the microorganisms are cultured in a bioreactor - e.g. a chemostat to which fresh medium is continuously added, while culture liquid is continuously removed to keep the culture volume constant. By harnessing the natural selection of the *E. Coli* to optimize and balance pathway activity towards establishing semiautotrophic growth, as described herein, the present inventors have generated populations of *E. Coli* with additional mutations. Thus, the present inventors propose deleting (or down-regulating) any one of the genes or intergenic sequences summarized in Tables 2-6 as well as those described herein above in order to generate additional microorganisms of this aspect of the present disclosure.

[0091] Semiautotrophic microorganisms may be further cultured for a suitable length of time, wherein the amount of the first organic carbon source is slowly reduced so that eventually autotrophic microorganisms are produced (i.e. microorganisms that do not require an organic carbon source to generate energy (e.g. ATP)). Typically a source of energy and reducing power is also required for the sustenance of the microorganism. The energy/reducing power source should not provide direct carbon for the growth of the host (otherwise the carbon fixation cycle becomes redundant) i.e. the energy reducing power source should not enable the microorganism to generate the group 1 precursor without being dependent on inorganic carbon dioxide fixation.

[0092] Thus, according to another aspect of the present disclosure there is provided a genetically modified *E. Coli* which expresses a recombinant phosphoribulokinase (prk) and Ribulose-Bisphosphate Carboxylase (RuBisCo) and has deletions in the genes *zwf*, *pfkA* and *pfkB*, the *E. Coli* being an autotroph.

[0093] The two best candidates for providing *E. Coli* with reducing power (and energy) are formate and phosphite. The soluble enzyme NAD⁺-dependent formate dehydrogenase irreversibly oxidizes formate ($E^{\circ}=-430\text{mV}$) and reduces NAD⁺ - formate cannot be directly assimilated by *E. Coli*. NAD:phosphite oxidoreductase irreversibly oxidizes phosphite to phosphate ($E^{\circ}=-650\text{mV}$) and reduces NAD⁺.

[0094] According to a particular aspect, formate may be used as the source of energy/reducing power as further described in US Application No. 61/913,940.

[0095] The formate which is used may come from any source - e.g., sodium formate, potassium formate, formic acid or formic acid anhydride etc.

[0096] Alternatively, and/or additionally, the formate may be generated using electricity. CO₂ can be directly reduced at the cathode (the electrons are derived from water splitting at the anode, for example) to generate formate at relatively high

efficiency.

[0097] In order to generate the formate for use by the microorganism, the microorganism is placed in a bioreactor in a fluid (e.g., water). The cathode may optionally be placed inside the bioreactor in contact with the microorganism. Alternatively, the cathode may be placed in a separate container to the bioreactor and the formate may be channeled to the chamber comprising the microorganism. The fluid may contain other elements required by the microorganism for growth including for example salts, minerals, metals and other nutrients, such as vitamins.

[0098] Examples of such bioreactors and further methods are provided in Li et al. *Science*, 2012, Vol 335, page 1596, Rabaey et al, *Current Opinion in Biotechnology*, 2011, 22: 371-377; Lovley et al., *Current Opinion in Biotechnology*, 2011, 22: 441-448; Lovley D.R., *Environmental microbiology reports*, 2011, 3(1), 27-35; Nevin et al., *Microbiology*, May/June 2010 Volume 1 Issue 2; Rabaey et al., *Applied and Industrial Microbiology*, *Nature Reviews*, October 2010, Volume 8, page 706-716.

[0099] The electrodes may be fabricated from such conductive polymers and metallic materials including indium tin oxide (ITO), graphite, platinum and silver.

[0100] According to one aspect, the microorganism is one that produces an industrially important product - e.g., a biofuel or a chemical (e.g. astaxanthin). Alternatively, or additionally, the microorganism expresses enzymes such that it is capable of producing an industrially important product - e.g., a biofuel. It will be appreciated that the precise choice of enzymes are selected according to the particular microorganism being used. Alternatively, or additionally, the microorganism expresses an industrially important product - e.g., a recombinant protein. Additional industrial important products include antibiotics or other pharmaceutical, solvents, pigments, food additives, monomers for the plastic industry and industrially valuable polymers.

[0101] Biofuels include for example, an alcohol (e.g., methanol, ethanol, propanol, isobutanol, and n-butanol etc.), a hydrocarbon (e.g., an alkane such as methane, ethane, propane, butane, an alkene such as ethylene, propylene, isoprenes, an alkyne such as acetylene etc.) hydrogen, a biodiesel (long-chain alkyl (methyl, propyl or ethyl) esters), an aldehyde or ketones (e.g. acetone, formaldehyde, 1-propanal, etc.). The biofuel can be a solid, a liquid or a gas.

[0102] Industrially useful microorganisms include the production of ethanol by *Saccharomyces* and the production of butanol by *Clostridium*.

[0103] Other industrially useful microorganisms include those in the production of plastic monomers and solvents.

[0104] The recombinant protein may be any protein - e.g., a human protein used for medicinal purposes. Examples of such proteins include an antibody, insulin, interferon, growth hormone, erythropoietin, growth hormone, follicle stimulating hormone, factor VIII, low density lipoprotein receptor (LDLR) alpha galactosidase A and glucocerebrosidase.

[0105] As mentioned, in order to express recombinant proteins in the microorganism, polynucleotide sequences encoding same are inserted into expression vectors as described herein above.

[0106] It will be appreciated that other than containing the necessary elements for the transcription and translation of the inserted coding sequence (encoding the industrially useful polypeptide), the expression construct for expression of the industrially useful polypeptide can also include sequences engineered to optimize stability, production, purification, yield or activity of the expressed polypeptide.

[0107] Depending on the vector and host system used for production, resultant polypeptides of the present disclosure may either remain within the recombinant cell, secreted into the fermentation medium, secreted into a space between two cellular membranes, such as the periplasmic space in *E. Coli*; or retained on the outer surface of a cell or viral membrane.

[0108] Following a predetermined time in culture, recovery of the recombinant polypeptide is effected.

[0109] The phrase "recovering the recombinant polypeptide" used herein refers to collecting the whole fermentation medium containing the polypeptide and need not imply additional steps of separation or purification.

[0110] Thus, polypeptides of the present disclosure can be purified using a variety of standard protein purification techniques, such as, but not limited to, affinity chromatography, ion exchange chromatography, filtration, electrophoresis, hydrophobic interaction chromatography, gel filtration chromatography, reverse phase chromatography, concanavalin A chromatography, chromatofocusing and differential solubilization.

[0111] To facilitate recovery, the expressed coding sequence can be engineered to encode the polypeptide of the present disclosure and fused cleavable moiety. Such a fusion protein can be designed so that the polypeptide can be readily isolated by affinity chromatography; e.g., by immobilization on a column specific for the cleavable moiety. Where a cleavage site is engineered between the polypeptide and the cleavable moiety, the polypeptide can be released from the chromatographic column by treatment with an appropriate enzyme or agent that specifically cleaves the fusion protein at this site (e.g. 26, 27).

[0112] Recovery of biofuels may be recovered according to methods known in the art. Alcohols such as ethanol, methanol, and/or butanol may be recovered from liquid material by molecular sieves, distillation, and/or other separation techniques. For example, ethanol can be concentrated by fractional distillation to about 90% or about 95% by weight. There are several methods available to further purify ethanol beyond the limits of distillation, and these include drying (e.g., with calcium oxide or rocksalt), the addition of small quantities of benzene or cyclohexane, molecular sieve, membrane, or by pressure reduction.

[0113] Product gas, for example, as produced by anaerobic metabolism or photosynthesis, may be processed to separate the methane and/or hydrogen components. Methane, hydrogen, or biogas may be drawn off from the system as pipeline gas.

[0114] In accordance with the disclosure, methane and/or hydrogen may be recovered as a biofuel product. Methane may be recovered and/or purified from biogas by known methods and systems which are commercially available, including membrane systems known for separating gases on the basis of different permeabilities. See, for example, U.S. Patent No. 6,601,543. Alternatively, various methods of adsorption may be used for separating methane and hydrogen.

[0115] Other ways of collecting biofuel products including centrifugation, temperature fractionalization, chromatographic methods and electrophoretic methods.

[0116] In certain aspects, the biofuel recovery/purification components may be integrated into the microorganism culturing system (e.g. bioreactor), for example, by connecting the respective device or apparatus to the gas or liquid effluents from the bioreactors. The purified biofuels and bioenergy products may be stored in a separate container(s).

[0117] Another use of the microorganisms of the present disclosure is to use them as a screen so as to identify enzymes and/or pathways capable of altering the efficiency of carbon fixation in a plant. Thus, the present disclosure contemplates expressing in the microorganism carbon fixation enzymes of alternate sources (e.g. plants, fungi etc.) in order to determine whether they have enhanced activity with respect to carbon fixation. According to a particular enzyme, the assay may be used to compare different homologs or variations of a particular enzyme.

[0118] Measuring the efficiency of carbon fixation may be performed by analyzing the amount of inorganic carbon assimilation. This may be measured *in-vivo* by incubating the cells in the presence of isotopic labeled $^{13}\text{CO}_2$. The assimilation of inorganic carbon into biomass may be detected via LC-MS analysis of metabolic extracts.

[0119] Once a particular enzyme has been identified (e.g. a naturally occurring enzyme, or a mutant enzyme which was generated by mutagenesis, as described herein or appeared via a natural selection in an evolutionary process, as described herein) which shows enhanced properties with respect to carbon fixation in the microorganisms described herein, the present inventors contemplate expressing those enzymes in autotrophic organisms such as plants or algae. It is envisaged that such organisms genetically modified to express these heterologous enzymes will also show enhanced properties with respect to carbon fixation.

[0120] There are various methods of introducing foreign genes into both monocotyledonous and dicotyledonous plants (Potrykus, I., *Annu. Rev. Plant. Physiol., Plant. Mol. Biol.* (1991) 42:205-225; Shimamoto et al., *Nature* (1989) 338:274-276).

[0121] The principle methods of causing stable integration of exogenous DNA into plant genomic DNA include two main approaches:

1. (i) Agrobacterium-mediated gene transfer: Klee et al. (1987) *Annu. Rev. Plant Physiol.* 38:467-486; Klee and Rogers in *Cell Culture and Somatic Cell Genetics of Plants*, Vol. 6, Molecular Biology of Plant Nuclear Genes, eds. Schell, J., and Vasil, L. K., Academic Publishers, San Diego, Calif. (1989) p. 2-25; Gatenby, in *Plant Biotechnology*, eds. Kung, S. and Amtzen, C. J., Butterworth Publishers, Boston, Mass. (1989) p. 93-112.
2. (ii) direct DNA uptake: Paszkowski et al., in *Cell Culture and Somatic Cell Genetics of Plants*, Vol. 6, Molecular Biology of Plant Nuclear Genes eds. Schell, J., and Vasil, L. K., Academic Publishers, San Diego, Calif. (1989) p. 52-68; including methods for direct uptake of DNA into protoplasts, Toriyama, K. et al. (1988) *Bio/Technology* 6:1072-1074. DNA uptake induced by brief electric shock of plant cells: Zhang et al. *Plant Cell Rep.* (1988) 7:379-384. Fromm et al. *Nature* (1986) 319:791-793. DNA injection into plant cells or tissues by particle bombardment, Klein et al. *Bio/Technology* (1988) 6:559-

563; McCabe et al. *Bio/Technology* (1988) 6:923-926; Sanford, *Physiol. Plant.* (1990) 79:206-209; by the use of micropipette systems: Neuhaus et al., *Theor. Appl. Genet.* (1987) 75:30-36; Neuhaus and Spangenberg, *Physiol. Plant.* (1990) 79:213-217; glass fibers or silicon carbide whisker transformation of cell cultures, embryos or callus tissue, U.S. Pat. No. 5,464,765 or by the direct incubation of DNA with germinating pollen, DeWet et al. in *Experimental Manipulation of Ovule Tissue*, eds. Chapman, G. P. and Mantell, S. H. and Daniels, W. Longman, London, (1985) p. 197-209; and Ohta, *Proc. Natl. Acad. Sci. USA* (1986) 83:715-719.

[0122] The *Agrobacterium* system includes the use of plasmid vectors that contain defined DNA segments that integrate into the plant genomic DNA. Methods of inoculation of the plant tissue vary depending upon the plant species and the *Agrobacterium* delivery system. A widely used approach is the leaf disc procedure which can be performed with any tissue explant that provides a good source for initiation of whole plant differentiation. Horsch et al. in *Plant Molecular Biology Manual A5*, Kluwer Academic Publishers, Dordrecht (1988) p. 1-9. A supplementary approach employs the *Agrobacterium* delivery system in combination with vacuum infiltration. The *Agrobacterium* system is especially viable in the creation of transgenic dicotyledenous plants.

[0123] There are various methods of direct DNA transfer into plant cells. In electroporation, the protoplasts are briefly exposed to a strong electric field. In microinjection, the DNA is mechanically injected directly into the cells using very small micropipettes. In microparticle bombardment, the DNA is adsorbed on microprojectiles such as magnesium sulfate crystals or tungsten particles, and the microprojectiles are physically accelerated into cells or plant tissues.

[0124] Following stable transformation plant propagation is exercised. The most common method of plant propagation is by seed. Regeneration by seed propagation, however, has the deficiency that due to heterozygosity there is a lack of uniformity in the crop, since seeds are produced by plants according to the genetic variances governed by Mendelian rules. Basically, each seed is genetically different and each will grow with its own specific traits. Therefore, it is preferred that the transformed plant be produced such that the regenerated plant has the identical traits and characteristics of the parent transgenic plant. Therefore, it is preferred that the transformed plant be regenerated by micropropagation which provides a rapid, consistent reproduction of the transformed plants.

[0125] Micropropagation is a process of growing new generation plants from a single piece of tissue that has been excised from a selected parent plant or cultivar. This process permits the mass reproduction of plants having the preferred tissue expressing the fusion protein. The new generation plants which are produced are genetically identical to, and have all of the characteristics of, the original plant. Micropropagation allows mass production of quality plant material in a short period of time and offers a rapid multiplication of selected cultivars in the preservation of the characteristics of the original transgenic or transformed plant. The advantages of cloning plants are the speed of plant multiplication and the quality and uniformity of plants produced.

[0126] Micropropagation is a multi-stage procedure that requires alteration of culture medium or growth conditions between stages. Thus, the micropropagation process involves four basic stages: Stage one, initial tissue culturing; stage two, tissue culture multiplication; stage three, differentiation and plant formation; and stage four, greenhouse culturing and hardening. During stage one, initial tissue culturing, the tissue culture is established and certified contaminant-free. During stage two, the initial tissue culture is multiplied until a sufficient number of tissue samples are produced to meet production goals. During stage three, the tissue samples grown in stage two are divided and grown into individual plantlets. At stage four, the transformed plantlets are transferred to a greenhouse for hardening where the plants' tolerance to light is gradually increased so that it can be grown in the natural environment.

[0127] Although stable transformation is presently preferred, transient transformation of leaf cells, meristematic cells or the whole plant is also envisaged by the present disclosure.

[0128] Transient transformation can be effected by any of the direct DNA transfer methods described above or by viral infection using modified plant viruses.

[0129] As used herein the term "about" refers to $\pm 10\%$.

[0130] The terms "comprises", "comprising", "includes", "including", "having" and their conjugates mean "including but not limited to".

[0131] The term "consisting of" means "including and limited to".

[0132] The term "consisting essentially of" means that the composition, method or structure may include additional ingredients, steps and/or parts, but only if the additional ingredients, steps and/or parts do not materially alter the basic and novel characteristics of the claimed composition, method or structure.

[0133] Whenever a numerical range is indicated herein, it is meant to include any cited numeral (fractional or integral) within the indicated range. The phrases "ranging/ranges between" a first indicate number and a second indicate number and "ranging/ranges from" a first indicate number "to" a second indicate number are used herein interchangeably and are meant to include the first and second indicated numbers and all the fractional and integral numerals therebetween.

[0134] As used herein the term "method" refers to manners, means, techniques and procedures for accomplishing a given task including, but not limited to, those manners, means, techniques and procedures either known to, or readily developed from known manners, means, techniques and procedures by practitioners of the chemical, pharmacological, biological, biochemical and medical arts.

[0135] It is appreciated that certain features of the disclosure, which are, for clarity, described in the context of separate aspects, may also be provided in combination in a single aspect. Conversely, various features of the disclosure, which are, for brevity, described in the context of a single aspect, may also be provided separately or in any suitable subcombination or as suitable in any other described aspect of the disclosure. Certain features described in the context of various aspects are not to be considered essential features of those aspects, unless the aspect is inoperative without those elements.

[0136] Various aspects and aspects of the present disclosure as delineated hereinabove and as claimed in the claims section below find experimental support in the following examples.

EXAMPLES

[0137] Reference is now made to the following examples, which together with the above descriptions illustrate some aspects of the disclosure in a non limiting fashion.

[0138] Generally, the nomenclature used herein and the laboratory procedures utilized in the present disclosure include molecular, biochemical, microbiological and recombinant DNA techniques. Such techniques are thoroughly explained in the literature. See, for example, "Molecular Cloning: A laboratory Manual" Sambrook et al., (1989); "Current Protocols in Molecular Biology" Volumes I-III Ausubel, R. M., ed. (1994); Ausubel et al., "Current Protocols in Molecular Biology", John Wiley and Sons, Baltimore, Maryland (1989); Perbal, "A Practical Guide to Molecular Cloning", John Wiley & Sons, New York (1988); Watson et al., "Recombinant DNA", Scientific American Books, New York; Birren et al. (eds) "Genome Analysis: A Laboratory Manual Series", Vols. 1-4, Cold Spring Harbor Laboratory Press, New York (1998); methodologies as set forth in U.S. Pat. Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659 and 5,272,057; "Cell Biology: A Laboratory Handbook", Volumes I-III Cellis, J. E., ed. (1994); "Culture of Animal Cells - A Manual of Basic Technique" by Freshney, Wiley-Liss, N. Y. (1994), Third Edition; "Current Protocols in Immunology" Volumes I-III Coligan J. E., ed. (1994); Stites et al. (eds), "Basic and Clinical Immunology" (8th Edition), Appleton & Lange, Norwalk, CT (1994); Mishell and Shiigi (eds), "Selected Methods in Cellular Immunology", W. H. Freeman and Co., New York (1980); available immunoassays are extensively described in the patent and scientific literature, see, for example, U.S. Pat. Nos. 3,791,932; 3,839,153; 3,850,752; 3,850,578; 3,853,987; 3,867,517; 3,879,262; 3,901,654; 3,935,074; 3,984,533; 3,996,345; 4,034,074; 4,098,876; 4,879,219; 5,011,771 and 5,281,521; "Oligonucleotide Synthesis" Gait, M. J., ed. (1984); "Nucleic Acid Hybridization" Hames, B. D., and Higgins S. J., eds. (1985); "Transcription and Translation" Hames, B. D., and Higgins S. J., eds. (1984); "Animal Cell Culture" Freshney, R. I., ed. (1986); "Immobilized Cells and Enzymes" IRL Press, (1986); "A Practical Guide to Molecular Cloning" Perbal, B., (1984) and "Methods in Enzymology" Vol. 1-317, Academic Press; "PCR Protocols: A Guide To Methods And Applications", Academic Press, San Diego, CA (1990); Marshak et al., "Strategies for Protein Purification and Characterization - A Laboratory Course Manual" CSHL Press (1996). Other general references are provided throughout this document. The procedures therein are believed to be well known in the art and are provided for the convenience of the reader.

GENERAL MATERIALS AND METHODS

[0139] *Strains*: Strains in which the metabolic genes were knocked-out were derived from Keio collection²⁴, *E. Coli* strain BW25113, referred in the text as "wild-type". The genome sequence of the parent used for constructing the ancestor strain, differed from *E. Coli* strain BW25113 in four loci: *ptsI* (D464N), *fabR* (V42G), *butB* (G162A), *fluA* (frameshift). These mutations

were acquired during early handling of the strain and were discovered during the course of resequencing the evolved clones. The strain inoculated in the chemostat for the evolutionary experiment, referred in the text as "ancestor", contained further genomic modifications: we deleted phosphoglycerate mutase genes (*gpmA*, *gpmM*), phosphofructokinase genes (*pfkA*, *pfkB*), 6-phosphate-1-dehydrogenase (*zwf*). In addition, we further deleted the *aceBAK* operon encoding for the enzymes of the glyoxylate shunt (thus ensuring that a bypass using the tartronate semialdehyde pathway cannot be used to grow on pyruvate as a sole organic carbon source in *Δgpm* mutant). Detailed information regarding the genotype of clones showing semiautotrophic phenotype which were isolated from chemostat experiments, referred in the text as "evolved isolated clones", appears in Tables 2-6, herein below.

[0140] Genomic modifications: Multiple gene knockouts were generally obtained by iterative rounds of P1 transductions²⁵ according to the following procedure: an isolate of the bacteria containing the desired gene deletion, derived from the Keio collection, was purified by single-colony isolation. Gene deletion was confirmed by PCR and the clone was used as a donor for the preparation of the P1 transducing lysate using standard protocols. Upon transduction, recipient strain was grown in agar plates containing a suitable carbon source and supplemented with Kanamycin (Km) as a positive selection for transduction. The introduction of the novel gene deletion was confirmed by checking the size of PCR fragments amplified by a set of primers located upstream and downstream of the targeted deletion(s). When the sizes of the deletion and non-deletion (wild-type) alleles were similar, deletions were validated by direct sequencing of the PCR fragments. Due to the size of the transduced DNA fragment (up to 100Kbp), it is unlikely that adjacent genes would be deleted via successive rounds of P1 transduction. In such cases we used a lambda RED-mediated gene replacement using a chloramphenicol (Cm) cassette²⁶. PCP20, a temperature sensitive plasmid encoding the FLP recombinase, was used according to standard procedures²⁶ to eliminate the Km or Cm resistance markers, allowing iterative rounds of deletion. After recombinase expression, all gene deletions and the loss of resistance markers were validated using PCR, and the resulting strain was used as the recipient in the next iteration of gene deletion. Whole genome sequencing was used to validate the genotype at the end of the process.

[0141] Growth conditions: Unless stated otherwise, cells were grown on M9 minimal media supplemented with mentioned carbon sources and chloramphenicol (34 µg/ml). Antibiotics were omitted when validating the ability of evolved semiautotrophic clones to grow solely on pyruvate and CO₂. Agar plates were prepared using ultrapure agarose (Hispanagar, Spain). Liquid cultures used for measuring growth curves, were cultured in 96-well plates and incubated at 37°C in a gas controlled shaking incubator (Infinite M200 - Tecan, USA).

[0142] Recombinant expression of RuBisCO and *prk*: For recombinant expression of the CBB cycle components, we constructed a synthetic operon encoding the His-tagged type-II RuBisCO from *Rhodospirillum rubrum* ATCC 11170 (*rbcL*), His-tagged phosphoribulokinase (*prkA*) from *Synechococcus elongatus* PCC 7942 and carbonate dehydratase (CA, Rru_A2056) from *Rhodospirillum rubrum* (CA). To differentially control the expression level of *rbcL*, *prkA* and CA we placed synthetic ribosome binding sites with varying translation efficiency (rbs-C, rbs-E and rbs-C, respectively) upstream to the open reading frame, as previously described²⁷. The synthetic operon was cloned into a pZ vector (Expressys, Germany) with the backbone containing a PLtetO-1 promoter and a p15A medium copy origin of replication. Construct assembly was based on "no-background" cloning methodology previously described²⁷, with Cm as selective marker. In control experiments, recombinant CBB genes were replaced with an mCherry reporter.

[0143] Computational analysis of RuBisCO dependent strains: To identify candidate mutants in which cell growth is coupled to the flux through non-native CBB enzymes, we implemented an algorithm based on the principles of flux balance analysis (FBA)²⁸. We started by considering all possible combinations of enzymatic reaction knockouts in central metabolism, and filtered out all those combinations that allow growth without any flux in RuBisCO. For those that cannot produce any biomass without RuBisCO, we calculated a slope which is defined as the biomass production rate achieved by allowing a unit of flux in RuBisCO. All other constraints were the same as in standard FBA, where the rate of biomass production is maximized. In more technical terms, the slope is the shadow price²⁹ of the upper bound on the RuBisCO flux, when that upper bound is set to 0. We implemented our algorithm using the COBRA for Python toolbox³⁰ and using the *E. Coli* core model (gcrdotucsddotedu/Downloads/EcoliCore), augmented with the two reactions corresponding to RuBisCO and phosphoribulokinase. It is possible to create a bi-level optimization MILP problem to find the knockout with the highest slope, using the same mathematical principles used in OptKnock^{31,32} and RobustKnock^{33,34}. However, since we only used the core *E. Coli* model, and had to consider about 20 single knockouts, we chose to explicitly calculate the slope of all possible combinations of one to three knockouts. The total runtime was less than 3 hours on a single Intel™ Core-i7 CPU.

[0144] Random Mutagenesis: A single colony of the ancestor strain was inoculated into 10 mL of M9 media with 5 g/L pyruvate, 0.2 g/L xylose, 2.5 g/L sodium bicarbonate and 34 mg/L chloramphenicol. The culture was incubated overnight in a

shaking incubator at 37°C. Subsequently, the cells were harvested and subjected to random mutagenesis using the Aquamutant Mutagenesis Kit (MultiTarget Pharmaceuticals, USA) according to manufacturer instructions. Cells from several mutagenesis intensities regimes (from none to very strong mutagenesis) were pooled and inoculated into the chemostat.

[0145] Chemostat Evolution: We performed a chemostat based laboratory evolution in several independent experiments. In the first and second experiments, bacterial cells following mutagenesis were inoculated into a Bioflo 110 chemostat (New Brunswick Scientific, USA) at working volume of 0.7 L and a dilution rate of 0.08 h⁻¹ (equivalent to a doubling rate of 8.75 hours). Feed input was M9 minimal media supplemented with 5 g/L sodium pyruvate, 0.1 g/L xylose and 30 mg/L chloramphenicol at 37°C. The culture was sparged at 0.5 L min⁻¹ with a gas mixture containing 25% CO₂, 5% O₂ and 70% N₂. Agitation speed was set to 500 rpm. Biomass density was measured daily and samples for glycerol stocks and the quantification of residual substrate concentrations were collected once a week. To rule out contaminations, culture samples were plated on positive (supplemented with pyruvate and xylose) and negative (glucose) control agar plates. Genetic markers of the ancestor strain were routinely validated in the colonies formed on pyruvate and xylose. No colonies were formed on glucose containing media during the period of the experiment. During the first experiment, the chemostat had a technical malfunction on day 20 and the experiment was restarted using a glycerol stock sample taken on day 18. To avoid the possibility that pyruvate becomes the limiting nutrient instead of xylose, in the second and third experiments, xylose concentrations were lowered in the feed media whenever a drop in residual pyruvate was detected in the culture. The third experiment was performed using a DASBox Bacterial Fermentation system (DASGIP - Eppendorf, Germany). The bacteria were grown in a working volume of 100 ml at a dilution rate of 0.035 h⁻¹ (equivalent to 20 hours doubling time). Feed input was the same as described above except that the initial xylose concentration was 25 mg/L. The gassing (0.1 L min⁻¹) mixture throughout the entire experiment contained 20% CO₂, 10% O₂ and 70% N₂. The OD, DO, pH and temperature were monitored on-line. Several attempts to perform the evolutionary process resulted in culture contamination after one to 10 weeks and are not reported.

[0146] Pyruvate and xylose measurements: Reported results of the residual concentrations of pyruvate and xylose in the chemostat culture were quantified enzymatically with K-PYRUV and K-XYLOSE assay kits (Megazyme, Ireland) respectively. In addition, pyruvate concentrations were validated using an Agilent 1200 series high-performance liquid chromatography system equipped with an anion exchange Bio-Rad HPX-87H column (Bio-Rad, USA). The column was eluted with 5 mM sulfuric acid at a flow rate of 0.6 mL/min at 45°C.

[0147] Isotopic analysis of hydrolyzed amino acids: Mass isotopologues distributions (MID) can supply information regarding the topology and the fluxes in a metabolic network. By measuring the MID of specific amino acids, it is possible to infer the labeling pattern in precursors metabolites upstream to it in the biosynthesis pathway. For intermediates directly converted to amino acids, hydrolyzed protein samples hold several advantages such as increased throughput and sample stability. For MID analysis, cells were cultured in M9 minimal media, either in the presence of a fully labeled organic carbon source and non-labeled CO₂, or in a reciprocal set-up with isotopically labeled ¹³CO₂ (Cambridge Isotope Laboratories, USA) and a non-labeled organic carbon source. For experiments in which gaseous ¹³CO₂ was used, culture tubes were placed in a transparent air-tight container and flushed with 5 volumes of isotopically labeled gas mixture (10% ¹³CO₂ 10% O₂ and 80% N₂). Cells were incubated in a shaking incubator at 37 °C and harvested during exponential growth. Protein biomass was hydrolyzed with 6N hydrochloric acid using standard protocols. Hydrolyzed amino acids were separated using ultra performance liquid chromatography (Acquity - Waters, USA) on a C-8 column (Zorbax Eclipse XBD - Agilent, USA) at a flow rate of 0.6 mL/min and eluted off the column using a hydrophobicity gradient. Buffers used were: A) H₂O + 0.1% formic acid and B) acetonitrile + 0.1% formic acid with the following gradient: 100% of A (0-3 min), 100% A to 100% B (3-9 min), 100% B (9-13 min), 100% B to 100% A (13-14 min), 100% A (14-20 min). Overall run time was 20 minutes. The UPLC was coupled online to a triple quadrupole mass spectrometer (TQS -Waters, USA). Data was acquired using MassLynx v4.1 (Waters, USA). Optimization of ionization parameters and determination of retention times was performed by direct infusion of amino acid commercial standards (Sigma-Aldrich, USA). Argon was used as the collision gas with a flow rate of 0.22 mL/min. Cone voltage was 25V, the capillary was set to 3 kV, source temperature was 150°C, desolvation temperature was 500°C, desolvation gas flow was 700 L/min, source offset 50, cone gas flow was 250 L/min and collision energy was 14eV. MID's were detected using multiple reaction monitoring (MRM) with the known molecular masses and the neutral loss of carbonyl carbon as a daughter ion (either 47 or 46 m/z, labeled and unlabeled respectively). Data analysis was performed using TargetLynx (Waters, USA). Total carbon labeling was calculated according to the formula:

$$\text{labeling fraction} = \frac{\sum_{i=0}^n m_i * i}{\sum_{i=0}^n m_i * n}$$

where m_i is total ion current as measured for the m₀..m_n masses of each compound.

[0148] Isotopic carbon ratio in whole biomass samples: Samples were grown in 500 mL glass flasks in M9 minimal media supplemented with 5 g/L sodium pyruvate. Ancestor strain samples were supplemented with 0.2 g/L xylose in addition to pyruvate. Following inoculation flasks were sealed with rubber septa and flushed with 5 volumes of a gas mixture containing isotopically labeled $^{13}\text{C}_2\text{O}_2$. Cells were harvested during exponential growth, washed in M9 minimal media and lyophilized. Between 0.2 and 0.4 mg of each dry biomass sample were weighed into tin capsules and their $^{13}\text{C}/^{12}\text{C}$ ratio was determined using an elemental analyzer linked to a Micromass (Manchester, UK) Optima IRMS.

[0149] Whole-genome sequencing: DNA was extracted from sampled cultures using DNeasy Blood & Tissue kit (QIAGEN, Germany). Samples were prepared for sequencing as previously described³⁵, with the following modifications: 0.6 μg of DNA was sheared using the Covaris E220x system (Covaris Inc., USA) to fragments of ≈ 200 bp, followed by ends repair, ligation to adapters, 8-cycles of PCR amplification with Kapa HiFi polymerase (Kapa Biosystems, USA). Samples were sequenced using an Illumina HiSeq 2500 (Illumina, USA) yielding 100 bp paired-end reads. Minimum of 2.5 million reads was obtained per sample, with a mean of 3.5 million reads per sample.

[0150] Analysis of Sequence Data: A reference genome which is based on *E. Coli* strain BW25113³⁶ (CP009273) and recombinant CBB enzymes encoding the plasmid as a second contig, were prepared. Reads for each sample were mapped independently using BWA³⁷ (version 0.7.5) with the standard parameters of the reference genome. A 95% unique mapping rate was observed. The mapped reads yielded a mean coverage depth of $\times 150$ per sample and above 99% of the genome was covered with more than $\times 50$. A GATK pipeline³⁸ was used to perform joint variant calling with all samples. The pipeline included the following steps: (a) Indel realignment using all samples jointly. (b) Variant calling using the Unified Genotyper, assuming a ploidy of 10 for each sample. (c) Standard filtering of SNPs and INDELS to mark suspected artefacts according to GATK Best Practices recommendations. This pipeline infers SNPs and INDELS, assuming a population of cells at a resolution of 10% in the allele frequency. To detect events that are related to the evolutionary process, we compared the frequencies of the alternative alleles between the ancestor and each of the evolved strains. To achieve that, we performed a Fisher exact test on the reference and alternative allele counts between each pair. Allele frequencies at each time point (as shown in Figure 3A) were estimated using allele counts for the alternative and reference alleles. Fisher test P-values were calculated for all SNP and INDELS. We note, that such analysis does not capture certain types of mutations, such as copy number variation, inversions, or large insertions or deletions. For the detection of large deletions (>100 bp) as well as large sequences ($>1000\text{bp}$) with high copy number variation, the coverage for each base in each sample was calculated and regions differing by more than three standard deviations of the average coverage in sample were annotated as high copy variation. Regions with a depth below 10 reads were marked as suspected deletions. Loci with poorly determined alternative alleles of isolated clones were revalidated by PCR amplification and Sanger sequencing. All mutations were validated manually using Geneious³⁹ version 8.0.5 (Biomatters, New Zealand).

[0151] Muller Diagrams: Muller diagrams were automatically generated by a custom program in python. The algorithm receives two inputs: a hierarchical lineage tree (manually inferred from the mutations frequencies and temporal distribution appearing in Figure 3A to achieve maximal parsimony), and the fraction of every mutation out of the population at each time point. A strain is therefore characterized by the set of mutations that it possesses, but that its ancestor strain does not. Given these two inputs the algorithm first, for every mutation at every time point, rounds it down to 0 if its fraction is smaller than a predefined threshold of 0.005 and given that at neither of the adjacent measurement points the fraction of that mutation is not higher. Next, the algorithm checks, for every mutation, that at every measurement point its fraction out of the population is at least as high as the sum of the fractions of the mutations of its direct descendant strains, where the test is performed recursively. In case this condition fails, the algorithm rounds up the fraction of the mutation to satisfy the constraint. Finally, the algorithm sets, and possibly adds time points to indicate the times at which every mutation arises. If mutation A first appeared in a measurement taken at time t_i , then A's initiation time is set to t_{i-1} . For every descendant strain of A, if it also first appeared at time t_{j-1} , then its initiation time is set to

$$\frac{t_i + t_{j-1}}{2}$$

.This procedure then continues recursively for the descendants of each strain whose initiation time was set. At this point the input is being used to generate the muller diagram in the following manner: at every time point, the [0,1] interval is divided to intervals representing the fractions the different mutations occupy out of the population, such that if mutation M has a frequency f_M at the given time point, it will be assigned an interval $[a, b]$ such that $b - a = f_M$. The division is done recursively, starting with the wild type (the ancestor of all the strains) being assigned the entire [0,1] interval. Then, given that mutation M occupies the interval $[a, b]$, and that there are n direct descendant strains to the strain containing M , the unique mutations of these descendent strains are given evenly spaced intervals in $[a, b]$, representing their frequencies, with the space between the intervals of two adjacent descendants being the frequency of M , f_M , minus the sum of the frequencies of all of its direct descendants, divided by $n + 1$ (to account of the spaces needed between the first and last descendants) and the boundaries of

the interval assigned to M , namely, a and b . For example, if N and L are the only descendants of M and at time point t their fractions are 0.2, 0.4 and 0.75 respectively, and M is assigned the interval $[0.2, 0.95]$ then N and L will be assigned the intervals $[0.25, 0.45]$ and $[0.5, 0.9]$ respectively, so that their intervals are spaced by 0.05 from the boundaries of M 's interval and between each other. The recursion then proceeds to the descendants of M until the entire strain hierarchy is covered. Once the intervals are determined for all time points, the algorithm generates the diagram by traversing the strains hierarchy tree. For every strain, identified by its unique mutation, the algorithm draws two lines, one connecting the lower bounds of the intervals of its mutation in time, and the other connecting the upper bounds. The area between these two lines is filled with the color assigned to that mutation. The algorithm then recursively continues to the direct descendants to overlay their area etc. Figure 3B has been graphically rendered based on the algorithm output in order to clarify locations where mutations occurred and to clearly position adjacent sectors.

RESULTS

[0152] The Calvin-Benson-Bassham (CBB) cycle¹ is the gateway to the organic world, being the main pathway for turning inorganic carbon (CO_2) into biomass and for storing energy in the biosphere. Heterotrophic organisms are dependent on such supply of organic carbon fixed by autotrophs. How difficult is it to evolve from one trophic mode of growth to another? Specifically, can an obligatory heterotrophic organism be evolved to synthesize biomass directly from CO_2 ? A positive answer will affect our understanding of metabolic plasticity and stimulate exciting avenues towards agricultural productivity^{2,3} as well as sustainable production of chemicals⁴. Here the present inventors demonstrate how a combination of rational metabolic rewiring and laboratory evolution under selective environmental conditions leads to the emergence of a fully functional CBB cycle in *E. Coli*. They rewired the metabolic network of an *E. Coli* host by introducing two enzymes (RuBisCO and *prk*) and severing the glycolysis pathway to decouple carbon-fixation from energy production. After several months under selective conditions in a chemostat, the modified bacteria evolved to semiautotrophic growth, in which carbon fixation via the non-native CBB cycle solely provides all carbons for major biomass building blocks (e.g., serine, histidine, pentose phosphates). Reducing power, energy and the rest of the biomass precursors are obtained by metabolizing a supplied organic compound (e.g., pyruvate). Using whole-genome whole-population sequencing, the evolutionary dynamics until the emergence of the semiautotrophic phenotype are described. The success in evolving a non-native carbon fixation pathway in an obligatory heterotrophic host provides a striking demonstration of the capacity for rapid trophic-mode evolution in metabolism with relevance to biotechnology⁵.

[0153] The activity of the CBB cycle accounts for >99% of CO_2 fixation in the biosphere⁶ and dominates agricultural productivity. Out of the 12 reactions of the CBB cycle, only two enzymatic activities are missing in *E. Coli* in order to complete a full cycle: phosphoribulokinase (*prk*) and ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO). *E. Coli*'s native gluconeogenesis and pentose phosphate pathway enzymes can catalyze all other reactions (Figure 1 and 4). Therefore, gene transfer of RuBisCO and *prk* can equip this heterotrophic host with all the enzymatic machinery required for CBB functionality. Previous efforts have shown functional expression of these enzymes⁷⁻¹⁰. However, full CBB operation in which only CO_2 and reducing energy are used as the cycle inputs has so far remained an elusive major challenge¹¹.

[0154] To explore whether the acquisition of only two non-native enzymes is sufficient to establish carbon fixation activity, the present inventors introduced a synthetic operon containing both RuBisCO and *prk* into an *E. Coli* host (*rbcL* from *Rhodospirillum rubrum* and *prkA* from *Synechococcus PCC6301*). Upon expression they were able to detect *prk*- and RuBisCO-dependent assimilation of inorganic $^{13}\text{CO}_2$ into biomass during heterotrophic growth on various carbon sources (Methods). However, this activity was found to be unstable and after less than 50 generations of continuous cultivation the cells lost all newly gained carbon fixation activity (for example, due to toxic effects of CBB cycle intermediates^{12,13} or protein burden, Figure 5).

[0155] Next, they examined whether additional metabolic modifications could ensure a stable fitness advantage of CBB cycle components that will sustain their activity. To identify gene deletions that make RuBisCO-dependent carbon fixation essential for *E. Coli* growth, they computationally analyzed the metabolic effect of RuBisCO and *prk* expression on various combinations of gene deletions (Methods and Figures 6 and 7). Their analysis pointed to a novel scenario in which full CBB cycle activity leads to the synthesis of essential biomass precursors using CO_2 as the sole carbon source, while energy supply relies on the metabolism of organic carbon (Figure 1). This scenario goes beyond previously described RuBisCO-dependent strains^{8,10} by operating a fully functional CBB cycle in which CO_2 , ATP and reducing power are the only inputs, as in autotrophic organisms. This mode of growth, not known to occur in nature, was termed semiautotrophic growth.

[0156] Out of several design alternatives for semiautotrophic growth, the present inventors decided to experimentally test one possible scenario by severing glycolysis through the deletion of phosphoglycerate mutase (*gpmA*, *gpmM*). This deletion effectively separates central metabolism into two disconnected sub-networks (Figure 1): (i) carbon fixing CBB module, containing upper glycolysis, the pentose phosphate pathway and the recombinant CBB enzymes; (ii) energy module, containing lower glycolysis and the TCA cycle, which supplies ATP and reducing power to drive carbon fixation taking place in the first module. Notably, out of the 12 biomass precursor metabolites from which biosynthetic pathways diverge to supply cellular building blocks^{14,15}, six are synthesized by the CBB module, and six by the energy module (Figure 8).

[0157] An organic carbon source feeding the energy module, such as pyruvate, could in principle provide reducing power and ATP for carbon fixation, hence allowing the synthesis of biomass precursors in the CBB module from CO₂. However, the *gpm* double mutant expressing RuBisCO and *prk* failed to grow semiautotrophically at elevated CO₂ when provided with a single organic carbon source (Figures 9 and 10). This strain grew only when a second carbon source feeding the CBB module directly was supplied in addition to pyruvate (e.g., xylose), allowing all biomass precursors to be produced from organic carbon (Figure 2A - left and Figure 10). It was hypothesized that given that all the necessary enzymatic components for carbon fixation are present, the failure to grow semiautotrophically can be solved by subjecting the cells to selective conditions favouring full CBB pathway operation. Natural selection can explore multi-knob fine tuning essential for a shift in growth mode, such as changes in regulation, that cannot be currently predicted by rational design.

[0158] The present inventors chose to harness natural selection to find the necessary metabolic adjustments by using a chemostat-based evolution experiment that continuously maintained selective conditions for the semiautotrophic phenotype¹⁶. The ancestor strain for this evolutionary experiment was the double knockout *gpm* strain expressing RuBisCO and *prk*, coupled with additional deletions of phosphofructokinase (*pfkA*, *pfkB*) and 6-phosphate-1-dehydrogenase (*zwf*) that make RuBisCO essential for xylose catabolism during heterotrophic growth (Figure 2A, 7, 9 and 10). The ancestor strain was subjected to random mutagenesis (Methods) and propagated in a xylose-limited chemostat at a dilution rate of 0.08 h⁻¹ (doubling time ≈9 hours). The CBB-module substrate concentration in the feed media was set to be strongly limiting ([xylose]=0.1 g/L) while the energy-module substrate and CO₂ were supplied in excess ([pyruvate]=5 g/L; pCO₂=0.25 atm). The balance of cell growth and dilution rate ensures that the concentration of xylose in the chemostat is extremely low and decreases even further as cells adapt. The scarcity of xylose imposes a strong and continuous selective pressure on cells to overcome the growth limitation by utilizing the abundant form of inorganic carbon (i.e., CO₂). It was hypothesized that this selection process will favour the fixation of inorganic carbon in the CBB module and enable a semiautotrophic strain to take over the population¹⁷. As expected, following inoculation the chemostat became xylose limited: internal xylose concentration in the chemostat was not detectable (<1 mg/L), while the concentration of pyruvate remained in considerable excess (≈ 2 g/L). Due to Δgpm severing the gluconeogenic pathway, carbon from pyruvate could not enter the carbon fixation module to compensate for the xylose limitation. However, excess pyruvate could serve as a source of energy and reducing power to be utilized by the CBB module, as it evolves to function as a xylose-independent CO₂ fixation cycle.

[0159] During the first 40 days of growth (≈100 chemostat generations), no significant change in cell density and nutrient concentrations was observed in the effluent media (Figure 2B). However, over the following 20 days a gradual increase in cell density was noted, accompanied by a steady decrease in pyruvate concentration. Finally, around day 60 (≈150 chemostat generations, Figure 2B) the concentration of pyruvate dropped to an undetectable level (<1 mg/L). Importantly, in contrast to the ancestor strain, culture samples from day 50 onward (Figure 2B arrow) formed colonies on minimal agar plates and grew in M9 liquid media (doubling time of ≈6 hours) supplied with only pyruvate and elevated CO₂ (Figure 2C). In ambient CO₂, no growth was detected on either liquid media or agar plates. The evolution of clones able to grow solely on CO₂ and pyruvate in two other independent chemostat experiments were observed (appearing on day 60 and 130, Figure 11).

[0160] Isotopic labeling experiments were conducted to test if the source of carbon for the synthesis of CBB-module biomass precursors (in the absence of xylose) results from carbon fixation or by utilizing pyruvate through an evolved Δgpm bypass¹⁸. The evolved bacteria were cultivated on non-labeled pyruvate and isotopically labeled ¹³CO₂ (p¹³CO₂ = 0.1 atm). Liquid chromatography-tandem mass spectrometry (LC-MS/MS) was used to analyze the labeling pattern of amino-acids derived from the energy-module versus those that originate from the CBB-module (Methods). As shown in Figure 2D, amino-acids derived from the CBB-module were almost fully labeled (e.g., serine and histidine), while amino-acids derived from the energy-module were mostly unlabeled (e.g., valine and threonine). Small deviations from complete labeling or no labeling are expected due to the release of unlabeled CO₂ from the TCA cycle and usage of labeled inorganic carbon in anaplerotic reactions respectively. The labeling pattern was verified in a reciprocal experiment where the bacteria was cultivated on uniformly labeled ¹³pyruvate and non-labeled CO₂. In this experiment serine and histidine were not labeled, while valine and

threonine were almost fully labeled (Figure 12). In addition, the labeling of intracellular precursor metabolites in the CBB and energy modules were analyzed, and as in the case of amino-acids, the labeling pattern matched the expected outcome from CO₂ fixation by the CBB module (Figure 13). Semiautotrophic phenotypes of clones isolated from all three chemostats were similarly validated using labeling experiments (Figures 12 and 13). Finally, isotopic analysis of biomass content revealed that ≈35% of cellular carbon originated from inorganic CO₂ (Figure 2E), matching the known fraction of biomass produced from CBB module metabolites¹⁵ (Figure 8). Therefore, it may be concluded that inorganic carbon fixation is entirely responsible for producing key biomass precursors.

[0161] To characterize the evolutionary process that led to the emergence of semiautotrophic growth, the experiment was repeated (Figure 11B) and whole-population samples were sequenced across 12 time points at a temporal resolution of ≈2 weeks. In addition, clones displaying the semiautotrophic phenotype were isolated and sequenced. The temporal trajectories of mutations (Figure 3A) were used to construct a Muller diagram^{19,20} describing the clonal dynamics of the evolving population (Figure 3B, Methods). Each sector in the diagram is a genotype, and a nested sector shares the mutations of its ancestor sector. During the first two months, four rapid nested mutational sweeps were observed (Figure 3B, blue shaded sectors Tables 2 and 3) that led to fixation of mutations in the coding sequence of *xyIE* (R160S in the low affinity transporter of D-xylose), *topA* (R168C in DNA topoisomerase), *crp* (K23NK insertion in transcriptional regulator of metabolic genes) and a short deletion in the promoter region of *yjiY* (carbon starvation associated protein, CstA homolog).

[0162] To identify the mutation which enabled the engineered strain to grow based on a fully functional non-native CBB cycle, three distinct clones which were isolated from the chemostat (~200 generations) and were able to grow on pyruvate only in the presence of CO₂ were isolated. The list of genetic mutations identified within the population samples from the chemostat experiment shown in Figure 3 are provided in Table 2. The list of genetic mutations identified in clones isolated from chemostat experiment shown in Figure 3 are provide in Table 3. The list of large chromosomal deletions in semiautotrophic clones isolated from the chemostat experiment shown in Figure 3 are provided in Table 4. Increased copy number of chromosomal sections in semiautotrophic clones isolated from chemostat experiment shown in Figure 3 are provided in Table 5. Mutations identified in population samples from the first chemostat experiment are provided in Table 6.

Table 2

Gene	Name in Fig.3:	Type (effect in protein)	Genomic change	Position relative to ORF start	Found in clones:	Genomic position	Related gene description
<i>xyIE</i>	<i>xyIE</i>	missense (R -> S)	G->T	478	EV1-5	4,231,705	D-xylose transporter
<i>topA</i>	<i>topA</i>	missense (R -> C)	C ->T	502	EV1-5	1,325,806	DNA topoisomerase I, omega subunit
<i>crp</i>	<i>crp</i>	insertion (K -> NK)	+TAA	69	EV1-5	3,479,547	cAMP-activated global transcription factor, mediator of catabolite repression
<i>crp</i>	<i>crp*</i>	insertion (K -> NIHK)	+CATTCATAA SEQ ID NO: 7	69	None	3,479,547	cAMP-activated global transcription factor, mediator of catabolite repression
<i>yjiY_p</i>	<i>yjiY</i>	deletion	-GGT	-125	EV1-5	4,581,222	putative transporter (promoter region)
<i>fliF</i>	<i>fliF</i>	deletion (frameshift)	-C	298	EV4-5	2,007,007	flagellar basal-body MS-ring and collar protein
<i>mlc</i>	<i>mlc+ptsP</i>	insertion (frameshift)	+AGAGAAA SEQ ID NO: 8	263	EV1-3	1,662,559	glucosamine anaerobic growth regulon transcriptional repressor; autorepressor
<i>ptsP</i>	<i>mlc+ptsP</i>	deletion (frameshift)	-CGACT SEQ ID NO: 9	24	EV1-3	2,961,770	fused PTS enzyme: PEP-protein phosphotransferase (enzyme I) /GAF domain containing protein

Gene	Name in Fig.3:	Type (effect in protein)	Genomic change	Position relative to ORF start	Found in clones:	Genomic position	Related gene description
rpoB	rpoB*	insertion (-> D)	+GAC	598	None	4,171,770	RNA polymerase, beta subunit
malk	malk+2	insertion (frameshift)	+CCGCCAGAA CGACGT GG TGTTGGTA SEQ ID NO: 10	1012	EV1-3	4,237,723	fused maltose transport subunit, ATP-binding component of ABC superfamily/regulatory protein
thrA	malk+2	missense (G->S)	G->A	1,207	EV1-3	1,543	Bifunctional aspartokinase/homoserine dehydrogenase 1
yfiH	malk+2	insertion (frameshift)	+C	492	EV1-3	2,727,902	UPF0124 family protein
ptsl	ptsl*	insertion (N->KVNRRGN)	+AGTTAACC GTGGTAA SEQ ID NO: 11	1,401	None	2,528,825	PEP-protein phosphotransferase of PTS system (enzyme I)
nadB_p	nadB*	deletion	-TTA	-76	None	2,703,703	quinolinate synthase, L-aspartate oxidase (B protein) subunit (promoter region)
prs	prs+4	insertion (A->ARVPITA)	+TCGTGTACC AATCACTGC SEQ ID NO: 12	330	EV1-3	1,257,002	Phosphoribosyl pyrophosphate synthase
nadR	prs+4	missense (A->T)	G->A	904	EV1-3	4,618,035	trifunctional protein: nicotinamide mononucleotide adenylyltransferase, ribosylnicotinamide kinase, transcriptional repressor
brnQ	prs+4	deletion (frameshift)	-C	396	EV1-3	415,442	branched-chain amino acid transport system 2 carrier protein; LIV-II transport system for Ile, Leu, and Val
tufA	prs+4	insertion (frameshift)	+A	317	EV1-3	3,464,372	translation elongation factor EF-Tu 1
ycbZ	prs+4	insertion (I->IAN)	+TTGGCA SEQ ID NO: 13	219	EV1-3	1,013,537	putative peptidase
brnQ	brnQ*	deletion (frameshift)	-G	1,073	None	416,119	branched-chain amino acid transport system 2 carrier protein; LIV-II transport system for Ile, Leu, and Val
mlc	mlc'+malk'+prs'+7	insertion (D->DQIN)	+TTATTTGAT SEQ ID NO: 14	32	EV4-5	1,662,790	glucosamine anaerobic growth regulon transcriptional repressor; autorepressor
prs	mlc'+malk'+prs'+7	insertion (T->TDMAI)	+ATTGCCA TATCG SEQ ID NO: 15	561	EV4-5	1,256,771	Phosphoribosyl pyrophosphate synthase
malk	mlc'+malk'+prs'+7	insertion (frameshift)	+GACGTGGT GTTGGTAGAAG SEQ ID NO: 16	1,016	EV4-5	4,237,727	fused maltose transport subunit, ATP-binding component of ABC superfamily/regulatory protein

<i>Gene</i>	<i>Name in Fig.3:</i>	<i>Type (effect in protein)</i>	<i>Genomic change</i>	<i>Position relative to ORF start</i>	<i>Found in clones:</i>	<i>Genomic position</i>	<i>Related gene description</i>
slyA	mlc'+malk'+prs'+7	insertion (frameshift)	+ATTTC	337	EV4-5	1,714,745	global transcriptional regulator
ppsR	mlc'+malk'+prs'+7	insertion (frameshift)	+GT	736	EV4-5	1,782,437	bifunctional regulatory protein: PEP synthase kinase and PEP synthase pyrophosphorylase
rpoC	mlc'+malk'+prs'+7	missense (G->S)	G->A	2,026	EV4-5	4,177,303	RNA polymerase, beta prime subunit
rne	mlc'+malk'+prs'+7	insertion (->RVRK)	+TTTACGC ACGCG SEQ ID NO: 17	1,462	EV4-5	1,138,362	fused ribonucleaseE: endoribonuclease/RNA-binding protein/RNA degradosome binding protein
ycjQ	mlc'+malk'+prs'+7	synonymous	G->A	84	EV4-5	1,369,303	putative Zn-dependent NAD(P)-binding oxidoreductase
hisL_p	mlc'+malk'+prs'+7	insertion	+TAT	-83	EV4-5	2,083,394	his operon leader peptide (promoter region)
psuK P	mlc'+malk'+prs'+7	insertion	+AA	-120	EV4-5	2,252,896	pseudouridine kinase (promoter region)
cbdA P	cbdA	insertion	+GATAA	-62	EV4-5	1,033,134	cytochrome bd-II oxidase, subunit I (promoter region)

Table 3

<i>Gene</i>	<i>Type (protein effect)</i>	<i>Genomic change</i>	<i>Position relative to ORF start</i>	<i>Found in clones:</i>	<i>Genomic position</i>	<i>Related gene description</i>
malEp	insertion	+CACCTA SEQ ID NO: 18	-48	EV1-3	4,236,396	maltose transporter subunit (promoter region)
opgH	nonsense (W->stop)	G->A	2,007	EV4	1,108,325	membrane glycosyltransferase; nutrient-dependent cell size regulator, FtsZ assembly antagonist
pdhR	insertion (R->HSRR)	+ATTCGC GTC SEQ ID NO: 19	575	EV1	119,153	pyruvate dehydrogenase complex repressor; autorepressor
pyrH	Insertion (K->IMGEK)	+TAATGGGTG AAA SEQ ID NO: 20	701	EV1	189,042	uridylate kinase
brnQ	insertion (truncation)	+TAAGA SEQ ID NO: 21	20	EV1	415,066	branched-chain amino acid transport system 2 carrier protein; LIV-II transport system for Ile, Leu, and Val
ybeL	insertion (frameshift)	+AATA	117	EV3	670,590	DUF1451 family protein
dcm	insertion (frameshift)	+TTTTTC SEQ ID NO: 22	558	EV5	2,025,241	DNA cytosine methyltransferase
asmA	insertion (frameshift)	+T	530	EV5	2,134,564	suppressor of OmpF assembly mutants; putative outer membrane protein assembly factor; inner membrane-anchored periplasmic protein

Gene	Type (protein effect)	Genomic change	Position relative to ORF start	Found in clones:	Genomic position	Related gene description
yejM	insertion (frameshift)	+GATGCC GAGCG SEQ ID NO: 23	1,100	EV1	2,278,954	essential inner membrane DUF3413 domain-containing protein; lipid A production and membrane permeability factor
murP	insertion (frameshift)	+CTTTCAAT SEQ ID NO: 24	1,023	EV2	2,541,054	N-acetylmuramic acid permease, EIBC component, PTS system

Table 4

Genes	Length [bp]	Found in clones:	start	end
mrp	210	EV4	2,186,581	2,186,791
opgH, yceK, msyB, mdtG	3,771	EV1-3	1,106,931	1,110,702
cryptic prophage e14 excision	15,139	EV4-5	1,191,728	1,206,867
yhhJ, rbbA, yhiI, yhiJ, yhiL, yhiM, yhiN, pitA, uspB, uspA, dtpB, rsmJ, prlC, rlmJ, gor	21,848	EV2-3	3,618,291	3,640,139

Table 5

Flanking genes	Length [bp]	Found in clones:	start	end
IS150...xylAB xylFGH...rhsA	83,646	EV2-3	3,758,616	3,842,262
rhsA...uhpB	41,651	EV2	3,713,998	3,755,649
yqeH...ygeN	8,928	EV4	2,980,486	2,989,414
tdcR, yhaB, yhaC	1,999	EV4	3,260,790	3,262,789
yqil, glgS	1,601	EV4	3,183,918	3,185,519

Table 6

Gene	Type (effect in protein)	Genomic change	Position relative to ORF start	Detected in chemostat	Found in clones:	Genomic position	description
ydgC	deletion (frameshift) ()	-C	157	Yes	EV6-9	1,676,131	inner membrane protein, GlpM family
topA	missense (R->P)	G->C	104	Yes	EV6-9	1,325,408	DNA topoisomerase I, omega subunit
gatY p	deletion	-T	-56	Yes	EV6-9	2,170,739	D-tagatose 1,6-bisphosphate aldolase 2, catalytic subunit
truB	deletion (frameshift)	-A	593	Yes	EV6-9	3,305,544	tRNA pseudouridine(55) synthase
brnQ	deletion (frameshift)	-AT	12	Yes	EV6-9	415,058	branched-chain amino acid transport system 2 carrier protein; LIV-II transport system for Ile, Leu, and Val
xylE	deletion (frameshift)	-T	137	Yes	EV6-9	4,232,046	D-xylose transporter
yhdZ_reg	deletion	-A	+4 after stop (IG)	Yes	EV6-9	3,416,557	putative amino-acid transporter subunit
malQ	deletion (frameshift)	-G	1269	Yes	EV6-9	3,542,161	4-alpha-glucanotransferase (amylomaltase)
xylF P	deletion	-A	-69	Yes	EV6-9	3,742,422	D-xylose transporter subunit
nanC_Reg	SNP (transversion)	G->T	-594 (IG)	Yes	EV6-9	4,529,912	N-acetylmuramic acid outer membrane channel protein
yjiY_P	deletion	-TAA	-126	Yes	EV6-9	4,581,222	putative inner membrane protein
yciV	deletion (frameshift)	-C	149	Yes	EV6-9	1,317,625	hypothetical protein

<i>Gene</i>	<i>Type (effect in protein)</i>	<i>Genomic change</i>	<i>Position relative to ORF start</i>	<i>Detected in chemostat</i>	<i>Found in clones:</i>	<i>Genomic position</i>	<i>description</i>
ppsR	deletion (frameshift)	-C	184	Yes	EV6-9	1,781,885	bifunctional regulatory protein: PEP synthase kinase and PEP synthase pyrophosphorylase
prs	missense (A->T)	C->T	283	Yes	EV6-9	1,257,049	phosphoribosylpyrophosphate synthase
dcm	deletion (frameshift)	-TGTA	1102	Yes	EV6-9	2,024,694	DNA cytosine methyltransferase
slyA	deletion (-V)	-GTT	322	Yes	EV6-9	1,714,758	DNA-binding transcriptional activator
trmA	deletion (frameshift)	-AT	1067	Yes	EV6-9	4,152,131	tRNA m(5)U54 methyltransferase, SAM-dependent
yjiY_P	deletion	-GGTAA SEQ ID NO: 25	-123	Yes	?	4,581,219	putative inner membrane protein
yciV	deletion (frameshift)	-A	488	Yes	None	1,317,964	hypothetical protein
yehX	deletion (frameshift)	-CA	830	Yes	None	2,210,056	putative transporter subunit: ATP-binding component of ABC superfamily
pepA	deletion (-I)	-ATC	549	Yes	None	4,475,218	multifunctional aminopeptidase A: a cyteinyglycinase, transcription regulator and site-specific recombination factor
aroK	deletion	-A	-355 (IG)	No	EV6-9	3,512,778	shikimate kinase I
ydbA_2	deletion (frameshift)	-AT	552	No	EV7-9	1,465,324	pseudogene, autotransporter homolog; interrupted by IS2 and IS30
lptG_reg				No	?		lipopolysaccharide export ABC permease of the LptBFGC export complex
cyoB	deletion (frameshift)	-A	1006	No	EV6	445,092	cytochrome o ubiquinol oxidase subunit I
xylF_p	deletion	-A	-17	No	EV6	3,724,474	D-xylose transporter subunit
rimI	deletion (-M)	-ATT	52	No	EV6	4,598,053	ribosomal-protein-S 18-alanine N-acetyltransferase
pitA	deletion (frameshift)	-A	957	No	EV6	3,631,958	phosphate transporter, low-affinity; tellurite importer
gltA_reg	deletion	-T	-299 (IG)	No	EV6	750,224	citrate synthase
chbB	deletion (frameshift)	-A	88	No	EV6	1,815,789	N,N'-diacetylchitobiose-specific enzyme IIB component of PTS
opgH	deletion (frameshift)	-TT	1813	No	EV6	1,108,131	membrane glycosyltransferase; nutrient-dependent cell size regulator, FtsZ assembly antagonist
sdaC_reg	deletion	-CAT	-388 (IG)	No	EV6	2,921,202	putative serine transporter
yqiC_p	deletion	-AA	-36	No	EV6	3,178,162	BMFP family putative fusogenic protein
asmA	deletion (frameshift)	-T	1066	No	EV6	2,134,028	putative assembly protein
xylA_reg	deletion	-AT	-139 (IG)	No	EV6	3,724,264	D-xylose isomerase

Gene	Type (effect in protein)	Genomic change	Position relative to ORF start	Detected in chemostat	Found in clones:	Genomic position	description
ybaM	deletion (frameshift)	-T	106	No	EV6, EV8	485,622	hypothetical protein
pyrH	deletion (-G)	-GTG	695	No	EV7-9	189,036	uridylate kinase
putP	deletion (frameshift)	-TC	1304	No	EV7, EV9	1,076,064	proline:sodium symporter
queA	deletion (frameshift)	-TA	177	No	EV8	420,643	S-adenosylmethionine :tRNA ribosyltransferase-isomerase
gadW_p	deletion	-TG	-51	No	EV8	3,658,029	transcriptional activator of gadA and gadBC; repressor of gadX
rpmH reg	insertion	+TCACCC ATG SEQ ID NO: 26	-80 (IG)	No	EV8	3,877,616	50S ribosomal subunit protein L34
sbp	deletion (frameshift)	-T	294	No	EV8	4,099,055	sulfate transporter subunit
hflC	deletion (frameshift) ()	-AT	277	No	EV8	4,393,393	modulator for HflB protease specific for phage lambda cII repressor
ldhA	insertion (I->IHL)	+GGTGAA SEQ ID NO: 27	359	No	EV8	1,436,743	fermentative D-lactate dehydrogenase, NAD-dependent
sodB p	deletion	-A	-19	No	EV8	1,729,616	superoxide dismutase, Fe
yciQ	deletion (frameshift)	-CT	59	No	EV8	1,319,060	putative inner membrane protein
sdiA_reg	deletion	-A	-80 (IG)	No	EV8	1,990,393	quorum-sensing transcriptional activator
adhE	deletion (frameshift)	-T	426	No	EV8	1,293,152	fused acetaldehyde-CoA dehydrogenase/iron-dependent alcohol dehydrogenase/pyruvate-formate lyase deactivase
rseB	deletion (frameshift)	-T	385	No	EV8	2,701,729	anti-sigma E factor, binds RseA
yheS_p	deletion	-CAT	-44	No	EV8	3,474,602	putative transporter subunit of ABC superfamily:ATP-binding component
nadR	deletion (frameshift)	-T	249	No	EV8	4,617,380	trifunctional protein: nicotinamide mononucleotide adenylyltransferase, ribosylnicotinamide kinase, transcriptional repressor
glgB	deletion (frameshift)	-C	396	No	EV9	3,566,467	1,4-alpha-glucan branching enzyme
hrpA	deletion (frameshift)	-T	1764	No	EV9	1,479,081	putative ATP-dependent helicase
pldA	deletion (frameshift)	-GA	253	No	EV9	3,998,474	outer membrane phospholipase A
yaiL	deletion (frameshift)	-CA	447	No	EV9	372,674	DUF2058 family protein
hflX	deletion (frameshift)	-G	541	No	EV9	4,391,029	GTPase, stimulated by 50S subunit binding
vjiL	deletion (frameshift)	-T	282	No	EV9	4,554,225	putative ATPase, activator of (R)-hydroxyglutaryl-CoA dehydratase

<i>Gene</i>	<i>Type (effect in protein)</i>	<i>Genomic change</i>	<i>Position relative to ORF start</i>	<i>Detected in chemostat</i>	<i>Found in clones:</i>	<i>Genomic position</i>	<i>description</i>
thiB	deletion (frameshift)	-TA	463	No	EV9	71,504	thiamin transporter subunit
ybjC	deletion (frameshift)	-T	15	No	EV9	886,383	conserved protein, DUF1418 family
nlpl	deletion (frameshift)	-AT	417	No	EV9	3,301,866	lipoprotein involved in osmotic sensitivity and filamentation
clpA	deletion (frameshift)	-T	415	No	EV9	919,134	ATPase and specificity subunit of ClpA-ClpP ATP-dependent serine protease, chaperone activity
clpA	deletion (frameshift)	-TC	417	No	EV9	919,136	ATPase and specificity subunit of ClpA-ClpP ATP-dependent serine protease, chaperone activity

REFERENCES

[0163]

1. Bassham, J. A. et al. The Path of Carbon in Photosynthesis. XXI. The Cyclic Regeneration of Carbon Dioxide Acceptor. *J. Am. Chem. Soc.* 76, 1760-1770 (1954).
2. Long, S. P., Marshall-Colon, A. & Zhu, X.-G. Meeting the Global Food Demand of the Future by Engineering Crop Photosynthesis and Yield Potential. *Cell* 161, 56-66 (2015).
3. Raines, C. A. The Calvin cycle revisited. *Photosynth. Res.* 75, 1-10 (2003).
4. Gronenberg, L. S., Marcheschi, R. J. & Liao, J. C. Next generation biofuel engineering in prokaryotes. *Curr. Opin. Chem. Biol.* 17, 462-471 (2013).
5. Church, G. M., Elowitz, M. B., Smolke, C. D., Voigt, C. A. & Weiss, R. Realizing the potential of synthetic biology. *Nat. Rev. Mol. Cell Biol.* 15, 289-294 (2014).
6. Raven, J. A., MacIntyre, H., Berman, T., Frank, I. B. & Others. Contributions of anoxygenic and oxygenic phototrophy and chemolithotrophy to carbon and oxygen fluxes in aquatic environments. in *Aquatic Microbial Ecology* 56, 177-192 (2009).
7. Parikh, M. R., Greene, D. N., Woods, K. K. & Matsumura, I. Directed evolution of RuBisCO hypermorphs through genetic selection in engineered *E. coli*. *Protein Eng. Des. Sel.* 19, 113-119 (2006).
8. Mueller-Cajar, O. & Whitney, S. M. Evolving improved *Synechococcus* Rubisco functional expression in *Escherichia coli*. *Biochem. J.* 414, 205 (2008).
9. Somerville, C. R. & Somerville, S. C. Cloning and expression of the *Rhodospirillum rubrum* ribulosebiphosphate carboxylase gene in *E. Coli*. *Mol. Gen. Genet.* 193, 214-219, (1984).
10. Durão, P. et al. Opposing effects of folding and assembly chaperones on evolvability of Rubisco. *Nat. Chem. Biol.* 11, 148-155 (2015).
11. Schada von Borzyskowski, L., Rosenthal, R. G. & Erb, T. J. Evolutionary history and biotechnological future of carboxylases. *J. Biotechnol.* 168, 243-251 (2013).
12. Wang, D., Zhang, Y., Pohlmann, E. L. & Li, J. The poor growth of *Rhodospirillum rubrum* mutants lacking RubisCO is due to the accumulation of ribulose-1, 5-bisphosphate. *Journal of* (2011).
13. Hudson, G. S., Morell, M. K., Arvidsson, Y. & Andrews, T. J. Synthesis of spinach phosphoribulokinase and ribulose 1, 5-bisphosphate in *Escherichia coli*. *Funct. Plant Biol.* 19, 213-221 (1992).
14. Noor, E., Eden, E., Milo, R. & Alon, U. Central carbon metabolism as a minimal biochemical walk between precursors for biomass and energy. *Mol. Cell* 39, 809-820 (2010).
15. Neidhardt, F. C. et al. *Escherichia coli* and *Salmonella typhimurium*. Cellular and molecular biology. Volumes I and II. (American Society for Microbiology, 1987).
16. Sauer, U. Evolutionary engineering of industrially important microbial phenotypes. *Adv. Biochem. Eng. Biotechnol.* 73, 129-169 (2001).
17. Sonderegger, M. & Sauer, U. Evolutionary engineering of *Saccharomyces cerevisiae* for anaerobic growth on xylose. *Appl. Environ. Microbiol.* 69, 1990-1998 (2003).

18. 18. Feuer, R. et al. Model-based analysis of an adaptive evolution experiment with *Escherichia coli* in a pyruvate limited continuous culture with glycerol. *EURASIP J. Bioinform. Syst. Biol.* 2012, 14 (2012).
19. 19. Elena, S. F. & Lenski, R. E. Evolution experiments with microorganisms: the dynamics and genetic bases of adaptation. *Nat. Rev. Genet.* 4, 457-469 (2003).
20. 20. Lang, G. I. et al. Pervasive genetic hitchhiking and clonal interference in forty evolving yeast populations. *Nature* 500, 571-574 (2013).
21. 21. Crozat, E., Philippe, N., Lenski, R. E., Geiselmann, J. & Schneider, D. Long-term experimental evolution in *Escherichia coli*. XII. DNA topology as a key target of selection. *Genetics* 169, 523-532 (2005).
22. 22. Brown, C. J., Todd, K. M. & Rosenzweig, R. F. Multiple duplications of yeast hexose transport genes in response to selection in a glucose-limited environment. *Mol. Biol. Evol.* 15, 931-942 (1998).
23. 23. Desai, M. M. & Fisher, D. S. Beneficial Mutation-Selection Balance and the Effect of Linkage on Positive Selection. *Genetics* 176, 1759-1798 (2007).
24. 24. Baba, T., Ara, T., Hasegawa, M. & Takai, Y. Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol. Syst. Biol.* (2006).
25. 25. Lennox, E. S. Transduction of linked genetic characters of the host by bacteriophage P1. *Virology* 1, 190-206 (1955).
26. 26. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. U. S. A.* 97, 6640-6645 (2000).
27. 27. Zelcbuch, L. et al. Spanning high-dimensional expression space using ribosome-binding site combinatorics. *Nucleic Acids Res.* 41, e98 (2013).
28. 28. Orth, J. D., Thiele, I. & Palsson, B. Ø. What is flux balance analysis? *Nat. Biotechnol.* 28, 245-248 (2010).
29. 29. Reznik, E., Mehta, P. & Segrè, D. Flux imbalance analysis and the sensitivity of cellular growth to changes in metabolite pools. *PLoS Comput. Biol.* 9, e1003195 (2013).
30. 30. Ebrahim, A., Lerman, J. A., Palsson, B. O. & Hyduke, D. R. COBRApy: COstraints-Based Reconstruction and Analysis for Python. *BMC Syst. Biol.* 7, 74 (2013).
31. 31. Burgard, A. P., Pharkya, P. & Maranas, C. D. OptKnock: a bilevel programming framework for identifying gene knockout strategies for microbial strain optimization. *Biotechnol. Bioeng.* 84, 647-657 (2003).
32. 32. Xu, Z., Zheng, P., Sun, J. & Ma, Y. ReactKnock: identifying reaction deletion strategies for microbial strain optimization based on genome-scale metabolic network. *PLoS One* 8, e72150 (2013).
33. 33. Tepper, N. & Shlomi, T. Predicting metabolic engineering knockout strategies for chemical production: accounting for competing pathways. *Bioinformatics* 26, 536-543 (2010).
34. 34. Tepper, N. & Shlomi, T. Computational design of auxotrophy-dependent microbial biosensors for combinatorial metabolic engineering experiments. *PLoS One* 6, e16274 (2011).
35. 35. Blecher-Gonen, R. et al. High-throughput chromatin immunoprecipitation for genome-wide mapping of in vivo protein-DNA interactions and epigenomic states. *Nat. Protoc.* 8, 539-554 (2013).
36. 36. Grenier, F., Matteau, D., Baby, V. & Rodrigue, S. Complete Genome Sequence of *Escherichia coli* BW25113. *Genome Announc.* 2, (2014).
37. 37. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* (2009).
38. 38. McKenna, A. et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* 20, 1297-1303 (2010).
39. 39. Kearse, M. et al. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28, 1647-1649 (2012).

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2011099005A \[0054\]](#)
- [WO2009038095A \[0055\]](#)
- [WO9907278A \[0054\]](#)
- [US5464764A \[0056\]](#)
- [US5487992A \[0056\]](#)
- [WO2012164565A \[0075\]](#)
- [US61913940B \[0094\]](#)
- [US6601543B \[0114\]](#)
- [US5464765A \[0121\]](#)
- [US4666828A \[0138\]](#)
- [US4683202A \[0138\]](#)
- [US4601531A \[0138\]](#)
- [US5192659A \[0138\]](#)
- [US5272057A \[0138\]](#)
- [US3791932A \[0138\]](#)
- [US3839153A \[0138\]](#)
- [US3850752A \[0138\]](#)
- [US3850578A \[0138\]](#)
- [US3853987A \[0138\]](#)
- [US3867517A \[0138\]](#)
- [US3879262A \[0138\]](#)
- [US3901654A \[0138\]](#)
- [US3935074A \[0138\]](#)
- [US3984533A \[0138\]](#)
- [US3996345A \[0138\]](#)
- [US4034074A \[0138\]](#)
- [US4098876A \[0138\]](#)
- [US4879219A \[0138\]](#)
- [US5011771A \[0138\]](#)
- [US5281521A \[0138\]](#)

Non-patent literature cited in the description

- **F. C. NEIDHARDT et al.** Physiology of the bacterial cell, 1990, 142- [0099]
- **SARDANA et al.** Plant Cell Reports, 1996, vol. 15, 677-681 [0051]
- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual Cold Springs Harbor Laboratory 19890000 [0056]
- **AUSUBEL et al.** Current Protocols in Molecular Biology John Wiley and Sons 19890000 [0056] [0138]
- **CHANG et al.** Somatic Gene Therapy CRC Press 19950000 [0056]
- **VEGA et al.** Gene Targeting CRC Press 19950000 [0056]
- **Vectors: A Survey of Molecular Cloning Vectors and Their Uses** Butterworths 19880000 [0056]
- **GILBOA** Biotechniques, 1986, vol. 4, 6504-512 [0056]
- **BANEYX et al.** Current Opinion in Biotechnology, 1999, vol. 10, 411-421 [0057]
- **MACRIDES et al.** Microbiol Rev, 1996, vol. 60, 512-538 [0057]
- **LI et al.** Science, 2012, vol. 335, 1596- [0098]
- **RABAEY et al.** Current Opinion in Biotechnology, 2011, vol. 22, 371-377 [0098]
- **LOVLEY et al.** Current Opinion in Biotechnology, 2011, vol. 22, 441-448 [0098]
- **LOVLEY D.R.** Environmental microbiology reports, 2011, vol. 3, 127-35 [0098]
- **NEVIN et al.** Microbiology, 2010, vol. 1, 2 [0098]
- **RABAEY et al.** Applied and Industrial Microbiology Nature Reviews, 2010, vol. 8, 706-716 [0098]
- **POTRYKUS, I.** Annu. Rev. Plant. Physiol. Plant. Mol. Biol, 1991, vol. 42, 205-225 [0120]
- **SHIMAMOTO et al.** Nature, 1989, vol. 338, 274-276 [0120]
- **KLEE et al.** Annu. Rev. Plant Physiol., 1987, vol. 38, 467-486 [0121]
- **Cell Culture and Somatic Cell Genetics of Plants** **KLEEROGERS** Molecular Biology of Plant Nuclear Genes Academic Publishers 19890000 vol. 6, 2-25 [0121]
- **GATENBY** Plant Biotechnology Butterworth Publishers 19890000 93-112 [0121]

- Cell Culture and Somatic Cell Genetics of Plants **PASZKOWSKI et al.** Molecular Biology of Plant Nuclear Genes Academic Publishers 19890000 vol. 6, 52-68 [\[0121\]](#)
- **TORIYAMA, K. et al.** Bio/Technology, 1988, vol. 6, 1072-1074 [\[0121\]](#)
- **ZHANG et al.** Plant Cell Rep., 1988, vol. 7, 379-384 [\[0121\]](#)
- **FROMM et al.** Nature, 1986, vol. 319, 791-793 [\[0121\]](#)
- **KLEIN et al.** Bio/Technology, 1988, vol. 6, 559-563 [\[0121\]](#)
- **MCCABE et al.** Bio/Technology, 1988, vol. 6, 923-926 [\[0121\]](#)
- **SANFORD** Physiol. Plant, 1990, vol. 79, 206-209 [\[0121\]](#)
- **NEUHAUS et al.** Theor. Appl. Genet, 1987, vol. 75, 30-36 [\[0121\]](#)
- **NEUHAUSSPANGENBERG** Physiol. Plant., 1990, vol. 79, 213-217 [\[0121\]](#)
- **DEWET et al.** Experimental Manipulation of Ovule Tissue, 1985, 197-209 [\[0121\]](#)
- **OHTA** Proc. Natl. Acad. Sci. USA, 1986, vol. 83, 715-719 [\[0121\]](#)
- **HORSCH et al.** Plant Molecular Biology Manual A5 Kluwer Academic Publishers 198800001-9 [\[0122\]](#)
- **SAMBROOK et al.** Molecular Cloning: A laboratory Manual 19890000 [\[0138\]](#)
- Current Protocols in Molecular Biology 19940000 vol. I-III, [\[0138\]](#)
- **PERBALA** Practical Guide to Molecular Cloning John Wiley & Sons 19880000 [\[0138\]](#)
- **WATSON et al.** Recombinant DNA Scientific American Books [\[0138\]](#)
- Genome Analysis: A Laboratory Manual Series Cold Spring Harbor Laboratory Press 19980000 vol. 1-4, [\[0138\]](#)
- Cell Biology: A Laboratory Handbook 19940000 vol. I-III, [\[0138\]](#)
- **FRESHNEY** Culture of Animal Cells - A Manual of Basic Technique Wiley-Liss 19940000 [\[0138\]](#)
- Current Protocols in Immunology 19940000 vol. I-III, [\[0138\]](#)
- Basic and Clinical Immunology Appleton & Lange 19940000 [\[0138\]](#)
- Selected Methods in Cellular Immunology W. H. Freeman and Co. 19800000 [\[0138\]](#)
- Oligonucleotide Synthesis, 1984, [\[0138\]](#)
- Nucleic Acid Hybridization, 1985, [\[0138\]](#)
- Transcription and Translation, 1984, [\[0138\]](#)
- Animal Cell Culture, 1986, [\[0138\]](#)
- Immobilized Cells and Enzymes IRL Press 19860000 [\[0138\]](#)
- **PERBAL, BA** Practical Guide to Molecular Cloning, 1984, [\[0138\]](#)
- Methods in Enzymology Academic Press vol. 1-317, [\[0138\]](#)
- PCR Protocols: A Guide To Methods And Applications Academic Press 19900000 [\[0138\]](#)
- **MARSHAK et al.** Strategies for Protein Purification and Characterization - A Laboratory Course Manual CSHL Press 19960000 [\[0138\]](#)
- **BASSHAM, J. A et al.** The Path of Carbon in Photosynthesis. XXI. The Cyclic Regeneration of Carbon Dioxide Acceptor J. Am. Chem. Soc., 1954, vol. 76, 1760-1770 [\[0163\]](#)
- **LONG, S. PMARSHALL-COLON, AZHU, X.-G** Meeting the Global Food Demand of the Future by Engineering Crop Photosynthesis and Yield Potential. Cell, 2015, vol. 161, 56-66 [\[0163\]](#)
- **RAINES, C. A.** The Calvin cycle revisited. Photosynth. Res, 2003, vol. 75, 1-10 [\[0163\]](#)
- **GRONENBERG, L. S. MARCHESCHI, R. J. LIAO, J. C.** Next generation biofuel engineering in prokaryotes. Curr. Opin. Chem. Biol., 2013, vol. 17, 462-471 [\[0163\]](#)
- **CHURCH, G. M. ELOWITZ, M. B. SMOLKE, C. D. VOIGT, C. A. WEISS, R.** Realizing the potential of synthetic biology. Nat. Rev. Mol. Cell Biol., 2014, vol. 15, 289-294 [\[0163\]](#)
- **RAVEN, J. A. MACINTYRE, H. BERMAN, T. FRANK, I. B.** Contributions of anoxygenic and oxygenic phototrophy and chemolithotrophy to carbon and oxygen fluxes in aquatic environments. Aquatic Microbial Ecology, 2009, vol. 56, 177-192 [\[0163\]](#)
- **PARIKH, M. R. GREENE, D. N. WOODS, K. K. MATSUMURA, I.** Directed evolution of RuBisCO hypermorphs through genetic selection in engineered E. coli. Protein Eng. Des. Sel., 2006, vol. 19, 113-119 [\[0163\]](#)
- **MUELLER-CAJAR, O. WHITNEY, S. M.** Evolving improved Synechococcus Rubisco functional expression in Escherichia coli. Biochem. J, 2008, vol. 414, 205- [\[0163\]](#)
- **SOMERVILLE, C. R. SOMERVILLE, S. C.** Cloning and expression of the Rhodospirillum rubrum ribulose biphosphate carboxylase gene in E. Coli. Mol. Gen. Genet, 1984, vol. 193, 214-219 [\[0163\]](#)
- **DURÃO, P et al.** Opposing effects of folding and assembly chaperones on evolvability of Rubisco. Nat. Chem. Biol, 2015, vol. 11, 148-155 [\[0163\]](#)
- **SCHADA VON BORZYSKOWSKIL., ROSENTHAL R. G. ERB, T. J.** Evolutionary history and biotechnological future of carboxylases. J. Biotechnol, 2013, vol. 168, 243-251 [\[0163\]](#)
- **WANG, D. ZHANG, Y. POHLMANN, E. L. LI, J.** The poor growth of Rhodospirillum rubrum mutants lacking RubisCO is due to the accumulation of ribulose-1, 5-bisphosphate. Journal of, 2011, [\[0163\]](#)
- **HUDSON, G. S. MORELL, M. K. ARVIDSSON, Y. ANDREWS, T. J.** Synthesis of spinach phosphoribulokinase and ribulose 1, 5-bisphosphate in Escherichia coli. Funct. Plant Biol., 1992, vol. 19, 213-221 [\[0163\]](#)

- **NOOR, E. EDEN, E. MILO, R. ALON, U.** Central carbon metabolism as a minimal biochemical walk between precursors for biomass and energy. *Mol. Cell*, 2010, vol. 39, 809-820 [\[0163\]](#)
- **Escherichia coli and Salmonella typhimurium NEIDHARDT, F. C. et al.** Cellular and molecular biology American Society for Microbiology 19870000 vol. I and II, [\[0163\]](#)
- **SAUER, U.** Evolutionary engineering of industrially important microbial phenotypes *Adv. Biochem. Eng. Biotechnol.*, 2001, vol. 73, 129-169 [\[0163\]](#)
- **SONDEREGGER, MSAUER, U.** Evolutionary engineering of *Saccharomyces cerevisiae* for anaerobic growth on xylose. *Appl. Environ. Microbiol.*, 2003, vol. 69, 1990-1998 [\[0163\]](#)
- **FEUER, R et al.** Model-based analysis of an adaptive evolution experiment with *Escherichia coli* in a pyruvate limited continuous culture with glycerol. *EURASIP J. Bioinform. Syst. Biol.* 2012, 2012, vol. 14, [\[0163\]](#)
- **ELENA, S. FLENSKI, R. E.** Evolution experiments with microorganisms: the dynamics and genetic bases of adaptation. *Nat. Rev. Genet.*, 2003, vol. 4, 457-469 [\[0163\]](#)
- **LANG, G. I. et al.** Pervasive genetic hitchhiking and clonal interference in forty evolving yeast populations *Nature*, 2013, vol. 500, 571-574 [\[0163\]](#)
- **CROZAT, E. PHILIPPE, N. LENSKI, R. E. GEISELMANN, J. SCHNEIDER, D.** Long-term experimental evolution in *Escherichia coli*. XII. DNA topology as a key target of selection. *Genetics*, 2005, vol. 169, 523-532 [\[0163\]](#)
- **BROWN, C. J. TODD, K. M. ROSENZWEIG, R. F.** Multiple duplications of yeast hexose transport genes in response to selection in a glucose-limited environment. *Mol. Biol. Evol.*, 1998, vol. 15, 931-942 [\[0163\]](#)
- **DESAI, M. M. FISHER, D. S.** Beneficial Mutation-Selection Balance and the Effect of Linkage on Positive Selection *Genetics*, 2007, vol. 176, 1759-1798 [\[0163\]](#)
- **BABA, T. ARA, T. HASEGAWA, M. TAKAI, Y.** Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol. Syst. Biol.*, 2006, [\[0163\]](#)
- **LENNOX, E. S.** Transduction of linked genetic characters of the host by bacteriophage P1 *Virology*, 1955, vol. 1, 190-206 [\[0163\]](#)
- **DATSENKO, K. A. WANNER, B. L.** One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products *Proc. Natl. Acad. Sci. U. S. A.*, 2000, vol. 97, 6640-6645 [\[0163\]](#)
- **ZELCIBUCH, L. et al.** Spanning high-dimensional expression space using ribosome-binding site combinatorics. *Nucleic Acids Res.*, 2013, vol. 41, e98- [\[0163\]](#)
- **ORTH, J. D. THIELE, I. PALSSON, B. Ø.** What is flux balance analysis *Nat. Biotechnol.*, 2010, vol. 28, 245-248 [\[0163\]](#)
- **REZNIK, EMEHTA, P. SEGRÈ, D.** Flux imbalance analysis and the sensitivity of cellular growth to changes in metabolite pools. *PLoS Comput. Biol.*, 2013, vol. 9, e1003195- [\[0163\]](#)
- **EBRAHIM, A. LERMAN, J. A. PALSSON, B. O. HYDUKE, D. R.** COBRApy: COstraints-Based Reconstruction and Analysis for Python. *BMC Syst. Biol.*, 2013, vol. 7, 74- [\[0163\]](#)
- **BURGARD, A. P. PHARKYA, P. MARANAS, C. D.** OptKnock: a bilevel programming framework for identifying gene knockout strategies for microbial strain optimization. *Biotechnol. Bioeng.*, 2003, vol. 84, 647-657 [\[0163\]](#)
- **XU, Z. ZHENG, P. SUN, J. MA, Y.** ReacKnock: identifying reaction deletion strategies for microbial strain optimization based on genome-scale metabolic network. *PLoS One*, 2013, vol. 8, e72150- [\[0163\]](#)
- **TEPPER, N. SHLOMI, T.** Predicting metabolic engineering knockout strategies for chemical production: accounting for competing pathways. *Bioinformatics*, 2010, vol. 26, 536-543 [\[0163\]](#)
- **TEPPER, N. SHLOMI, T.** Computational design of auxotrophy-dependent microbial biosensors for combinatorial metabolic engineering experiments. *PLoS One*, 2011, vol. 6, e16274- [\[0163\]](#)
- **BLECHER-GONEN, R. et al.** High-throughput chromatin immunoprecipitation for genome-wide mapping of in vivo protein-DNA interactions and epigenomic states. *Nat. Protoc.*, 2013, vol. 8, 539-554 [\[0163\]](#)
- **GRENIER, F. MATTEAU, DBABY, V. RODRIGUE, S.** Complete Genome Sequence of *Escherichia coli* BW25113. *Genome Announc.*, 2014, vol. 2, [\[0163\]](#)
- **LI, H. DURBIN, R.** Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*, 2009, [\[0163\]](#)
- **MCKENNA, A. et al.** The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.*, 2010, vol. 20, 1297-1303 [\[0163\]](#)
- **KEARSE, M et al.** Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data *Bioinformatics*, 2012, vol. 28, 1647-1649 [\[0163\]](#)

REKOMBINANTE MIKROORGANISMER, SOM ER I STAND TIL FIKSERING AF KULSTOF

PATENTKRAV

1. Genmodificeret *E. Coli* som udtrykker en rekombinant phosphoribulokinase (prk) og Ribulose-bisphosphat carboxylase (RuBisCo) og har deletioner i generne zwf, pfkA, pfkB, gpmA og gpmB.
2. Genetisk modificeret *E. Coli* ifølge krav 1, som ydermere har en deletion i aceBAK.
3. Genetisk modificeret *E. Coli* ifølge krav 1, som ydermere har deletion i aceBAK og malatsyntase G (glcB).
4. Genetisk modificeret *E. Coli* ifølge krav 1, som ydermere har deletioner i aceBAK, malatsyntase G (glcB), phosphoenolpyruvat syntase (pps) og phosphoenolpyruvat carboxykinase (pck).
5. Genetisk modificeret *E. Coli* ifølge krav 1, som ydermere har deletioner i aceBAK og phosphopyruvat hydratase (eno).
6. Cellekultur som omfatter *E. Coli* ifølge ethvert af kravene 1-5 og et medium som omfatter en organisk kulstofkilde.
7. Fremgangsmåde til formering af *E. Coli* ifølge ethvert af kravene 1-6 som omfatter dyrkning af *E. Coli* i et medie som omfatter en organisk kulstofkilde og en uorganisk kulstofkilde, og derved formering af *E. Coli*.
8. Fremgangsmåde til produktion af et kemikalie som omfatter dyrkning af *E. Coli* ifølge ethvert af kravene 1-5 i et medie som omfatter en organisk kulstofkilde og en uorganisk kulstofkilde og udtagning af kemikaliet.

DRAWINGS

Drawing




FIG. 1

FIG. 2A

RuBisCO dependent heterotroph ancestor

evolved semiautotroph (fully functional CBB cycle)




FIG. 2D

LC-MS isotopic analysis




FIG. 2B chemostat evolution dynamics




FIG. 2C CO₂ dependent semiautotrophic growth




FIG. 2E

cellular biomass from ¹³CO₂




FIG. 3A





FIG. 3B






FIG. 4B




FIG. 4A

FIG. 5A




FIG. 5B




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 6D





FIG. 6E





FIG. 6F

FIG. 7B

flux dependency phase space



Δpfk

Δzwf

glu fru gal
rib pyr xpl
2pg acc gly

Δpfk

Δzwf

glu fru gal
rib pyr xpl
2pg acc gly

FIG. 8




FIG. 9A




FIG. 9B




FIG. 10A





FIG. 10B

experimental validation



CBB enzymes are essential even while xylose is feeding the CBB module

FIG. 10C




FIG. 11A




FIG. 11B




FIG. 11C




FIG. 11D




FIG. 12




FIG. 13A

Chemostat evolution dynamics (n=1)




FIG. 13B

Chemostat evolution dynamics (n=2)





FIG. 13C

Chemostat evolution dynamics (n=3)



SEKVENSLISTE

Sekvenslisten er udeladt af skriftet og kan hentes fra det Europæiske Patent Register.

The Sequence Listing was omitted from the document and can be downloaded from the European Patent Register.

