



(51) International Patent Classification:

C11D 1/22 (2006.01) *C11D 3/386* (2006.01)
C11D 1/42 (2006.01) *C11D 17/00* (2006.01)
C11D 3/20 (2006.01) *C11D 17/04* (2006.01)

(21) International Application Number:

PCT/US2016/035372

(22) International Filing Date:

2 June 2016 (02.06.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

15170805.4 5 June 2015 (05.06.2015) EP
 16158410.7 3 March 2016 (03.03.2016) EP

(71) Applicant: **THE PROCTER & GAMBLE COMPANY**
[US/US]; One Procter & Gamble Plaza, Cincinnati, Ohio
45202 (US).(72) Inventors: **SOUTER, Philip, Frank**; Procter & Gamble
Technical Centres Limited, Whitley Road, PO Box: Forest
Hall No. 2, Longbenton Newcastle upon Tyne NE12 9TS
(GB). **SOMERVILLE-ROBERTS, Nigel, Patrick**;
Procter & Gamble Technical Centres Limited, Whitley
Road, PO Box: Forest Hall No. 2, Longbenton Newcastle
upon Tyne NE12 9TS (GB). **BROOKER, Alan, Thomas**;
Procter & Gamble Technical Centres Limited, Whitley
Road, PO Box: Forest Hall No. 2, Longbenton Newcastle
upon Tyne NE12 9TS (GB). **XU, Dan**; Procter & Gamble
Technical Centres Limited, Whitley Road, PO Box: Forest
Hall No. 2, Longbenton Newcastle upon Tyne NE12 9TS
(GB). **GUMMEL, Jeremie, Robert Marcel**; Procter &

Gamble Technical Centres Limited, Whitley Road, PO
Box: Forest Hall No. 2, Longbenton Newcastle upon Tyne
NE12 9TS (GB).

(74) Agent: **KREBS, Jay A.**; c/o The Procter & Gamble Com-
pany, Global Patent Services, One Procter & Gamble
Plaza, C8-229, Cincinnati, Ohio 45202 (US).(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LI, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: COMPACTED LIQUID LAUNDRY DETERGENT COMPOSITION

(57) **Abstract:** The present invention is to a liquid laundry detergent composition comprising: a. a liquid phase; b. between 0.5% and 15% by weight of the liquid detergent composition of a solid enzyme particle, wherein the solid phase is dispersed within the liquid phase and wherein the water-soluble solid phase is defined as the solid obtained when the liquid laundry detergent composition is centrifuged at 1200 G for 10 mins; and wherein the liquid phase comprises between 5% and 40% by weight of the liquid of an alcohol selected from the group comprising ethylene glycol, 1,3 propanediol, 1,2 propanediol, tetramethylene glycol, pentamethylene glycol, hexamethylene glycol, 2,3-butane diol, 1,3 butanediol, diethylene glycol, triethylene glycol, polyethylene glycol, glycerol formal, dipropylene glycol, polypropylene glycol, dipropylene glycol n-butyl ether, and mixtures thereof, preferably the alcohol is selected from the group comprising 1,2 propanediol, dipropylene glycol, polypropylene glycol, 2,3-butane diol, dipropylene glycol n-butyl ether and mixtures thereof; and from 10% to 30% by weight of the composition of a non-amine neutralized linear alkylbenzene sulphonate and wherein the liquid laundry detergent composition comprises less than 10% by weight of the liquid laundry detergent composition of an amine-neutralised anionic surfactant; and wherein the composition comprises between 0.5% and 50% by weight of the composition of water; and wherein the composition comprises less than 5% by weight of the composition of a hydroxyl-containing amine.



COMPACTED LIQUID LAUNDRY DETERGENT COMPOSITION

FIELD OF THE INVENTION

The present invention is the field of liquid laundry detergent compositions and their
5 methods of use.

BACKGROUND OF THE INVENTION

Liquid laundry detergent compositions with low equilibrium relative humidities have the
advantage of being less susceptible to microbial contamination. There is also a trend towards so
10 called compacted liquids that minimise the presence of unnecessary 'filler' liquids such as water.
Such compositions are more environmentally friendly as less unnecessary material needs to be
transported. Therefore, there is a move in the industry to using so called compacted liquids
which minimise the levels of non-active materials such as water. Such liquid laundry detergent
compositions require both the presence of anionic surfactant such as linear alkylbenzene
15 sulphonate and other non-surfactant cleaning and/or care enzymes.

However, such compacted composition can often have high viscosities due to the high
relative concentration of the cleaning materials such as anionic surfactants. Traditionally,
hydroxyl-containing amines have been used in such compositions to ensure consumer acceptable
viscosity of the liquid laundry detergent composition. Also, acceptable viscosity is required to
20 allow processability of the composition during manufacture. The hydroxyl-containing amines
are often used as neutralising agents for the anionic detergent surfactants such as linear
alkylbenzene sulphonate.

However, there is now a desire to reduce the overall level of such hydroxyl-containing
amines.

25 Reduction in the level of the hydroxyl-containing amines of known low relative humidity
laundry detergent compositions can result in high viscosity of the composition which negatively
impacts the ability of the consumer to accurately pour and dose the composition. Also,
processability of the composition is impacted as it is difficult to handle such viscous
compositions during manufacture.

30 Thus, there is a need in the art for low relative humidity liquid laundry detergent
compositions containing lower levels of hydroxyl-containing amine compounds, but which
exhibit consumer acceptable and/or process acceptable viscosities.

It has been surprisingly found that the above problems are overcome by the specific
formulation space of the present invention. The formulation space described below can provide

a liquid composition having a low relative humidity and comprising lower levels of hydroxyl-containing amine compounds but which has acceptable viscosity.

SUMMARY OF THE INVENTION

5 The present invention is also to a liquid laundry detergent composition comprising;

- a. a liquid phase;
- b. between 0.5% and 15% by weight of the liquid detergent composition of a solid enzyme particle,

wherein the solid phase is dispersed within the liquid phase and wherein the water-soluble solid phase is defined as the solid obtained when the liquid laundry detergent composition is centrifuged at 1200 G for 10 mins; and

wherein the liquid phase comprises between 5% and 40% by weight of the liquid of an alcohol selected from the group comprising ethylene glycol, 1,3 propanediol, 1,2 propanediol, tetramethylene glycol, pentamethylene glycol, hexamethylene glycol, 2,3-butane diol, 1,3 butanediol, diethylene glycol, triethylene glycol, polyethylene glycol, glycerol formal, dipropylene glycol, polypropylene glycol, dipropylene glycol n-butyl ether, and mixtures thereof, preferably the alcohol is selected from the group comprising 1,2 propanediol, dipropylene glycol, polypropylene glycol, 2,3- butane diol, dipropylene glycol n-butyl ether and mixtures thereof; and

from 10% to 30% by weight of the composition of a non-amine neutralized linear alkylbenzene sulphonate and wherein the liquid laundry detergent composition comprises less than 10% by weight of the liquid laundry detergent composition of an amine-neutralised anionic surfactant; and

wherein the composition comprises between 0.5% and 50% by weight of the composition of water; and

wherein the composition comprises less than 5% by weight of the composition of a hydroxyl-containing amine.

10 DETAILED DESCRIPTION OF THE INVENTION

Laundry Detergent Composition

The liquid laundry detergent composition of the present invention comprises a liquid phase and between 0.5% and 15% by weight of the composition of a solid enzyme particle. The solid particle is dispersed within the liquid phase. Suitable enzymes are described in more detail
15 below. The solid particle and liquid phases are described in more detail below.

Preferably, the liquid laundry detergent composition has a viscosity of between 300mPa.s and 700mPa.s, more preferably between 350mPa.s and 600mPa.s at a shear rate of 1000s⁻¹. An exemplary method for measuring viscosity is to use a Rheometer DHR1 from TA instruments using a gap of 1000µm at 20°C as according to the manufacturer's instructions.

5 The liquid laundry detergent composition of the present invention is preferably opaque. Without wishing to be bound by theory, consumers prefer opaque compositions as this signals excellent cleaning. By opaque we herein mean the composition has a fresh hunter L value of greater than 70, more preferably greater than 72, more preferably greater than 75. The Hunter colour space is organized as a cube. The L axis runs from top to bottom; the maximum L being
10 100 which is white and the minimum value is zero, which is black. The a and b axes have no specific numerical limits, however positive a is red, negative a is green, positive b is yellow and negative b is blue (see figure 1). Delta values (ΔL , Δa and Δb) can be measured and are associated with a colour change. The total colour difference, ΔE , can also be calculated. The ΔE is a single value that takes into account the differences between the L, a and b of test and
15 comparison samples. The ΔE is calculated as follows;

Using L_1 , a_1 , b_1 and L_2 , a_2 and b_2

$$\Delta E = \sqrt{(L_2 - L_1)^2 + (a_2 - a_1)^2 + (b_2 - b_1)^2}$$

A just noticeable difference (JND) is characterized as a ΔE of greater than 2.3. The JND is the smallest detectable difference possible with the human eye between a starting and
20 secondary level of a particular sensory stimulus.

The measurements of the present invention are taken on a HunterLab colour measurement instrument (Hunter Lab Color Quest XE), set as follows;

Illuminant: D65, Angle of observer: 10°, Mode: reflection

The instrument is used as per the manufacturer's instructions. A sample of 20mL are
25 tested in an optically clear glass cell having a fixed path length of 10mm and dimensions 55mm by 57mm. The measurement type is reflectance measurement RSIN, which measures the diffuse and specular reflectance of the sample at the port. The measurements are made with the specular exclusion port door closed.

Fresh Hunter colour value is a measure of the colour parameters of a fresh sample,
30 immediately after preparation.

The liquid laundry detergent composition of the present invention overall is liquid in nature. That is to say, even though it comprises a solid particle dispersed within a liquid phase, the composition has the nature of a liquid rather than a solid or granular composition. In relation to the laundry detergent composition of the present invention, the term 'liquid' encompasses

forms such as dispersions, gels, pastes and the like. The liquid composition may also include gases in suitably subdivided form. However, the liquid composition excludes forms which are non-liquid overall, such as tablets or granules.

The term 'liquid laundry detergent composition' refers to any laundry detergent composition comprising a liquid capable of wetting and treating fabric e.g., cleaning clothing in a domestic washing machine,

The liquid composition may be formulated into a unit dose article. The unit dose article of the present invention comprises a water-soluble film which fully encloses the liquid composition in at least one compartment. Suitable unit dose articles are described in more detail below.

The liquid laundry detergent composition can be used as a fully formulated consumer product, or may be added to one or more further ingredient to form a fully formulated consumer product. The liquid laundry detergent composition may be a 'pre-treat' composition which is added to a fabric, preferably a fabric stain, ahead of the fabric being added to a wash liquor.

The liquid laundry detergent composition comprises from 10% to 30% by weight of the composition of linear alkylbenzene sulphonate.

The liquid laundry detergent composition comprises less than 10% by weight, or even less than 5% by weight, or even less than 2% by weight of the liquid laundry detergent composition of an amine-neutralised anionic surfactant, wherein the anionic surfactant is preferably selected from the group comprising linear alkylbenzene sulphonate, alkyl sulphate and mixtures thereof.

The liquid laundry detergent composition comprises between 0.5% and 50% by weight of the composition of water and preferably has an equilibrium relative humidity of less than 65% at 20°C.

The composition comprises less than 5% by weight of the composition of a hydroxyl-containing amine. Suitable amines are described in more detail below.

The liquid laundry detergent composition may comprise a structurant. Suitable structurants are described in more detail below.

The liquid laundry detergent composition may comprise a silica.

The liquid laundry detergent composition may comprise a perfume raw material. The perfume raw material is preferably selected from aldehydes, ketones or a mixture thereof.

The liquid laundry detergent composition of the present invention may comprise adjunct ingredients, wherein the adjunct ingredients are present in the solid phase, the liquid phase or both.

Without wishing to be bound by theory, it is believed that the removal of the hydroxyl-containing amine compounds results in a number of detergent ingredients, to come out of solution. This in turn results in increase viscosity of the composition. The present invention carefully balances the form of the ingredients between the solid and liquid forms, so resulting in a composition of acceptable viscosity.

Furthermore, removal of the hydroxyl-containing amine can cause the formulation to phase split (i.e. at least two visibly distinct phases can be seen). The present invention provides the additional benefit of providing a composition having a low relative humidity and lower levels of hydroxyl-containing amine compounds, whilst minimising phase splitting.

Solid enzyme particle

The liquid laundry detergent composition of the present invention comprises a solid enzyme particle, wherein the solid enzyme particle is dispersed in the liquid phase. Preferably the enzyme particle is partially or completely water-soluble.

By 'solid' we herein mean any material that is solid, i.e. not liquid. The solid is in particulate form. The term 'particles' is herein used in its broadest meaning. The particles may have a mean particle size distribution of between 2µm and 50µm.

By 'water-soluble' we herein mean at least 75%, or even at least 85% or even at least 95% of the solid dissolves in water as measured by the method set out here after using a glass-filter with a maximum pore size of 20 microns:

5 grams $\pm$ 0.1 gram of solid is added in a pre-weighed 3L beaker and 2L $\pm$ 5ml of distilled water is added. This is stirred vigorously on a magnetic stirrer, Labline model No. 1250 or equivalent and 15cm magnetic stirrer, set at 600 rpm, for 30 minutes at 35°C. Then, the mixture is filtered through a folded qualitative sintered-glass filter with a pore size as defined above (max. 20 micron). The water is dried off from the collected filtrate by any conventional method, and the weight of the remaining material is determined (which is the dissolved or dispersed fraction). Then, the percentage solubility or dispersability can be calculated.

The water-soluble solid phase consists of any material obtained in the solid fraction when the liquid laundry detergent composition is centrifuged at 1200 G for 10 mins. A preferred method is;

1. Before use, pre heat the centrifuge (Sigma Centrifuge 6-15H, 6-pot rotor) to the desired temperature. When loading centrifuge tubes into the rotor, they should always be placed opposite each other in diametrically opposed positions, number of samples can be tested are 2, 3, 4 and 6.

2. 85ml polycarbonate with screw lids test tubes are used. Each tube was filled with 50g of material and the total mass: tube+lid+testing material measured
3. Place the tubes in the centrifuge rotor so that they are evenly spaced, and fasten the rotor cover firmly. When loading centrifuge tubes into the rotor, they should always be placed opposite each other in diametrically opposed positions, number of samples can be tested are 2, 3, 4 and 6.
4. Set the centrifuge time to 90 minutes. Start the centrifuge (it will gradually increase the speed automatically until 17119 Relative Centrifugal Force (Maximum RCF for this centrifuge is used to maximise the separation rate) is achieved.
5. At the end of the 90 minutes, reweigh each tube to ensure that no material has been lost, as centrifuge tubes can crack after several uses.
6. Different fractions can result at end of the centrifugation and the number of fractions depends on the nature of the sample, solid fraction is the most dense, opaque fraction at the bottom and the relative high viscosity. The bottom fraction can then be obtained by simply removing the top phase(s) from the tube.

The liquid laundry detergent composition may comprise between 0.5% and 10%, or even between 0.5% and 7.5% or even between 0.5% and 5% by weight of the liquid laundry detergent composition of the enzyme particle.

The enzyme may be selected from the group comprising hemicellulases, peroxidases, proteases, cellulases, xylanases, lipases, phospholipases, esterases, cutinases, pectinases, keratanases, reductases, oxidases, phenoloxidases, lipxygenases, ligninases, pullulanases, tannases, pentosanases, malanases, β -glucanases, arabinosidases, hyaluronidase, chondroitinase, laccase, and amylases, or mixtures thereof. A typical combination is a cocktail of conventional applicable enzymes like protease, lipase, cutinase and/or cellulase in conjunction with amylase.

A typical combination is an enzyme cocktail that may comprise, for example, a protease and lipase in conjunction with amylase.

Preferred enzymes could include a protease. Suitable proteases include metalloproteases and serine proteases, including neutral or alkaline microbial serine proteases, such as subtilisins (EC 3.4.21.62). In one aspect, such suitable protease may be of microbial origin. The suitable proteases include chemically or genetically modified mutants of the aforementioned suitable proteases. In one aspect, the suitable protease may be a serine protease, such as an alkaline microbial protease or/and a trypsin-type protease. Examples of suitable neutral or alkaline proteases include:

(a) subtilisins (EC 3.4.21.62), including those derived from *Bacillus*, such as *Bacillus lentus*, *B. alkalophilus*, *B. subtilis*, *B. amyloliquefaciens*, *Bacillus pumilus* and *Bacillus gibsonii*.

(b) trypsin-type or chymotrypsin-type proteases, such as trypsin (*e.g.*, of porcine or bovine origin), including *Fusarium* protease and chymotrypsin proteases derived from
 5 *Cellulomonas*.

(c) metalloproteases, including those derived from *Bacillus amyloliquefaciens*.

(d) subtilisin proteases derived from the *Bacillus* sp TY-145 , NCIMB 40339, especially the variants described with substitutions and/or deletions at positions 171, 173, 175 or 179.

Preferred proteases include those derived from *Bacillus gibsonii*, *Bacillus*

10 *amyloliquefaciens*, *Bacillus* sp. TY-145 or *Bacillus Lentus*.

Suitable commercially available protease enzymes include those sold under the trade names Alcalase®, Savinase®, Primase®, Durazym®, Polarzyme®, Kannase®, Liqueanase®, Liqueanase Ultra®, Savinase Ultra®, Ovozyme®, Neutrase®, Blaze®, Everlase® and Esperase® by Novozymes A/S (Denmark), those sold under the tradename Maxatase®, Maxacal®,

15 Maxapem®, Properase®, Purafect®, Purafect Prime®, Purafect Ox®, FN3®, FN4®, Excellase®, Ultimase®, Purafect OXP® and the Preferenz P® series by DuPont International Biosciences, those sold under the tradename Opticlean® and Optimase® by Solvay Enzymes, those available from BASF, namely BLAP (sequence shown in Figure 29 of US 5,352,604 with the following mutations S99D + S101 R + S103A + V104I + G159S, hereinafter referred to as
 20 BLAP), BLAP R (BLAP with S3T + V4I + V199M + V205I + L217D), BLAP X (BLAP with S3T + V4I + V205I) and BLAP F49 (BLAP with S3T + V4I + A194P + V199M + V205I + L217D) - all from BASF; and KAP (*Bacillus alkalophilus* subtilisin with mutations A230V + S256G + S259N) from Kao.

Suitable alpha-amylases include those of bacterial or fungal origin. Chemically or
 25 genetically modified mutants (variants) are included. A preferred alkaline alpha-amylase is derived from a strain of *Bacillus*, such as *Bacillus licheniformis*, *Bacillus amyloliquefaciens*, *Bacillus stearothermophilus*, *Bacillus subtilis*, or other *Bacillus* sp., such as *Bacillus* sp. NCIB 12289, NCIB 12512, NCIB 12513, DSM 9375, DSM 12368, DSMZ no. 12649, KSM AP1378, KSM K36 or KSM K38. Preferred amylases include:

30 (a) the variants described in WO 94/02597, WO 94/18314, WO96/23874 and WO 97/43424, especially the variants with substitutions in one or more of the following positions versus the enzyme listed as SEQ ID No. 2 in WO 96/23874: 15, 23, 105, 106, 124, 128, 133, 154, 156, 181 , 188, 190, 197, 202, 208, 209, 243, 264, 304, 305, 391, 408, and 444.

(b) the variants described in USP 5,856,164 and WO99/23211, WO 96/23873, WO00/60060 and WO 06/002643, especially the variants with one or more substitutions in the following positions versus the AA560 enzyme listed as SEQ ID No. 12 in WO 06/002643:

26, 30, 33, 82, 37, 106, 118, 128, 133, 149, 150, 160, 178, 182, 186, 193, 203, 214, 231,
 256, 257, 258, 269, 270, 272, 283, 295, 296, 298, 299, 303, 304, 305, 311, 314, 315, 318, 319,
 339, 345, 361, 378, 383, 419, 421, 437, 441, 444, 445, 446, 447, 450, 461, 471, 482, 484,
 preferably that also contain the deletions of D183* and G184*.

(c) variants exhibiting at least 85%, preferably 90% identity with SEQ ID No. 4 in WO06/002643, the wild-type enzyme from *Bacillus* SP722, especially variants with deletions in the 183 and 184 positions and variants described in WO 00/60060, WO11/100410 and WO13/003659, particularly those with one or more substitutions at the following positions versus SEQ ID No. 4 in WO06/002643:

51, 52, 54, 109, 304, 140, 189, 134, 195, 206, 243, 260, 262, 284, 347, 439, 469, 476 and 477.

(d) variants exhibiting at least 95% identity with the wild-type enzyme from *Bacillus* sp.707 (SEQ ID NO:7 in US 6,093, 562), especially those comprising one or more of the following mutations M202, M208, S255, R172, and/or M261. Preferably said amylase comprises one or more of M202L, M202V, M202S, M202T, M202I, M202Q, M202W, S255N and/or R172Q. Particularly preferred are those comprising the M202L or M202T mutations.

(e) variants described in WO 09/149130, preferably those exhibiting at least 90% identity with SEQ ID NO: 1 or SEQ ID NO:2 in WO 09/149130, the wild-type enzyme from *Geobacillus Stearothermophilus* or a truncated version thereof.

(f) variants described in WO10/115021, especially those exhibiting at least 75%, or at least 85% or at least 90% or at least 95% with SEQ ID NO:2 in WO10/115021, the alpha-amylase derived from *Bacillus* sp. TS-23.

Suitable commercially available alpha-amylases include DURAMYL®, LIQUEZYME®, TERMAMYL®, TERMAMYL ULTRA®, NATALASE®, SUPRAMYL®, STAINZYME®, STAINZYME PLUS®, EVEREST®, FUNGAMYL® and BAN® (Novozymes A/S, Bagsvaerd, Denmark), KEMZYM® AT 9000 Biozym Biotech Trading GmbH Wehlistrasse 27b A-1200 Wien Austria, RAPIDASE®, PURASTAR®, ENZYSIZE®, OPTISIZE HT PLUS®, POWERASE® and PURASTAR OXAM®, PREFERENZ® S series, including PREFERENZ S1000 and PREFERENZ S110 (DuPont Industrial Biosciences, Palo Alto, California) and KAM® (Kao, 14-10 Nihonbashi Kayabacho, 1-chome, Chuo-ku Tokyo 103-8210, Japan). In one aspect, suitable amylases include NATALASE®, EVEREST®, PREFERENZ S1000®, STAINZYME® and STAINZYME PLUS® and mixtures thereof.

In one aspect, such enzymes may be selected from the group consisting of: lipases, including “first cycle lipases”. In one aspect, the lipase is a first-wash lipase, preferably a variant of the wild-type lipase from *Thermomyces lanuginosus* comprising one or more of the T231R and N233R mutations. The wild-type sequence is the 269 amino acids (amino acids 23 – 291) of the Swissprot accession number Swiss-Prot O59952 (derived from *Thermomyces lanuginosus* (*Humicola lanuginosa*)). Preferred first cycle lipases include those with mutations at one or more of the following positions: 27, 38, 58, 60, 83, 96, 111, 150, 163, 227, 231, 233, 254, 255 and 256. Further preferred lipase variants have a net charge that is more positive than the wild-type lipase. The net charge can be calculated by assigning a charge of -1 to the anionic groups (D and E) and a net charge of +1 to the cationic groups (R and K) and comparing to the wild-type. Preferred mutations include D27R, G38A, G91A, D96G, D96E, D111A, G163K, G225R, T231R, N233R, D254S, P256T, S58A, V60S, S83T, A150G, L227G, I255A and/or P256K. Preferred variants include those with mutations:

(a) T231R + N233R;

(b) D27R + G38A + D96E + D111A + G163K + T231R + N233R + D254S + P256T

(c) G91A + D96G + G225R + T231R + N233R

Preferred lipases would include those sold under the tradenames Lipex®, Lipoclean®, Calipso® and Lipolex®.

In one aspect, other preferred enzymes include fungal and microbial-derived endoglucanases exhibiting endo-beta-1,4-glucanase activity (E.C. 3.2.1.4), including a bacterial polypeptide endogenous to a member of the genus *Bacillus* which has a sequence of at least 90%, 94%, 97% and even 99% identity to the amino acid sequence SEQ ID NO:2 in US7,141,403B2) and mixtures thereof. Suitable endoglucanases are sold under the tradenames Celluclean®, Carezyme®, Celluzyme®, Carezyme Premium® and Whitezyme® (Novozymes A/S, Bagsvaerd, Denmark).

Other preferred enzymes include pectate lyases sold under the tradenames Pectawash®, Pectaway®, Xpect® and mannanases sold under the tradenames Mannaway® (all from Novozymes A/S, Bagsvaerd, Denmark), and Preferenz F® and Purabrite® (DuPont International Biosciences, Palo Alto, California).

Deoxyribonuclease (DNase): Suitable deoxyribonucleases (DNases) are any enzyme that catalyzes the hydrolytic cleavage of phosphodiester linkages in the DNA backbone, thus degrading DNA. According to the invention, a DNase which is obtainable from a bacterium is preferred; in particular a DNase which is obtainable from a *Bacillus* is preferred; in particular a DNase which is obtainable from *Bacillus subtilis* or *Bacillus licheniformis* is preferred.

Perhydrolases: Suitable perhydrolases are capable of catalyzing a perhydrolysis reaction that results in the production of a peracid from a carboxylic acid ester (acyl) substrate in the presence of a source of peroxygen (e.g., hydrogen peroxide). While many enzymes perform this reaction at low levels, perhydrolases exhibit a high perhydrolysis:hydrolysis ratio, often greater than 1. Suitable perhydrolases may be of plant, bacterial or fungal origin. Chemically modified or protein engineered mutants are included.

Examples of useful perhydrolases include naturally occurring *Mycobacterium* perhydrolase enzymes, or variants thereof. An exemplary enzyme is derived from *Mycobacterium smegmatis*.

Oxidases/peroxidases: Suitable oxidases and peroxidases (or oxidoreductases) include various sugar oxidases, laccases, peroxidases and haloperoxidases.

Suitable peroxidases include those comprised by the enzyme classification EC 1.1.1.17, as set out by the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology (IUBMB), or any fragment derived therefrom, exhibiting peroxidase activity.

Suitable peroxidases include those of plant, bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful peroxidases include peroxidases from *Coprinopsis*, e.g., from *C. cinerea* and variants thereof.

A peroxidase for use in the invention also include a haloperoxidase enzyme, such as chloroperoxidase, bromoperoxidase and compounds exhibiting chloroperoxidase or bromoperoxidase activity. Haloperoxidases are classified according to their specificity for halide ions. Chloroperoxidases (E.C. 1.1.1.10) catalyze formation of hypochlorite from chloride ions.

In an embodiment, the haloperoxidase is a chloroperoxidase. Preferably, the haloperoxidase is a vanadium haloperoxidase, i.e., a vanadate-containing haloperoxidase. In a preferred method of the present invention the vanadate-containing haloperoxidase is combined with a source of chloride ion.

Haloperoxidases have been isolated from many different fungi, in particular from the fungus group dematiaceous hyphomycetes, such as *Caldariomyces*, e.g., *C. fumago*, *Alternaria*, *Curvularia*, e.g., *C. verruculosa* and *C. inaequalis*, *Drechslera*, *Ulocladium* and *Botrytis*.

Haloperoxidases have also been isolated from bacteria such as *Pseudomonas*, e.g., *P. pyrrocinia* and *Streptomyces*, e.g., *S. aureofaciens*.

In an preferred embodiment, the haloperoxidase is derivable from *Curvularia* sp., in particular *Curvularia verruculosa* or *Curvularia inaequalis*, such as *C. inaequalis* CBS 102.42 6;

or *C. verruculosa* CBS 147.63 or *C. verruculosa* CBS 444.70; or from *Drechslera hartlebii*, *Dendryphiella salina*, *Phaeotrichoconis crotalarie*, or *Geniculosporium* sp.

An oxidase according to the invention include, in particular, any laccase enzyme comprised by the enzyme classification EC 1.10.3.2, or any fragment derived therefrom exhibiting laccase activity, or a compound exhibiting a similar activity, such as a catechol oxidase (EC 1.10.3.1), an o-aminophenol oxidase (EC 1.10.3.4), or a bilirubin oxidase (EC 1.3.3.5).

Preferred laccase enzymes are enzymes of microbial origin. The enzymes may be derived from plants, bacteria or fungi (including filamentous fungi and yeasts).

Suitable examples from fungi include a laccase derivable from a strain of *Aspergillus*, *Neurospora*, e.g., *N. crassa*, *Podospora*, *Botrytis*, *Collybia*, *Fomes*, *Lentinus*, *Pleurotus*, *Trametes*, e.g., *T. villosa* and *T. versicolor*, *Rhizoctonia*, e.g., *R. solani*, *Coprinopsis*, e.g., *C. cinerea*, *C. comatus*, *C. friesii*, and *C. plicatilis*, *Psathyrella*, e.g., *P. condelleana*, *Panaeolus*, e.g., *P. papilionaceus*, *Myceliophthora*, e.g., *M. thermophila*, *Schytalidium*, e.g., *S. thermophilum*, *Polyporus*, e.g., *P. pinsitus*, *Phlebia*, e.g., *P. radiata* or *Coriolus*, e.g., *C. irsutus*.

Suitable examples from bacteria include a laccase derivable from a strain of *Bacillus*. A laccase derived from *Coprinopsis* or *Myceliophthora* is preferred; in particular a laccase derived from *Coprinopsis cinerea*; or from *Myceliophthora thermophila*.

Examples of other oxidases include, but are not limited to, amino acid oxidase, glucose oxidase, lactate oxidase, galactose oxidase, polyol oxidase and aldose oxidase. Oxidases and their corresponding substrates may be used as hydrogen peroxide generating enzyme systems, and thus a source of hydrogen peroxide. Several enzymes, such as peroxidases, haloperoxidases and perhydrolases, require a source of hydrogen peroxide. By studying EC 1.1.3., EC 1.2.3., EC 1.4.3., and EC 1.5.3. or similar classes (under the International Union of Biochemistry), other examples of such combinations of oxidases and substrates are easily recognized by one skilled in the art.

The enzyme may also be added in a solid form or as a capsule. Solid forms would include granules that can be made by fluid bed coating such as layered granules. Preferably said microcapsules and granules are coated with a polymer that provides triggered release via an ionic strength trigger such that said granule and/or capsule is stable in product but upon dilution in water will release its enzyme payload. Examples of such polymeric coatings include cellulose derivatives, such as hydroxypropyl methyl cellulose derivatives, particularly hydroxyl propyl methyl cellulose phthalate and cellulose acetate phthalate. A further preferred polymeric coating is polyvinyl alcohol. It is further preferred that any capsules and/or granules are density-matched

to the surrounding liquid matrix to promote stability and prevent settling out of a visible phase. In a further aspect the enzymes can be added as capsules and/or microcapsules derived from interfacial polymerization reaction of a polyamine, preferably a branched polyamine. Said microcapsules can be made by reaction of polyamines, such as those sold under the Lupasol tradename by BASF with an acid chloride.

For the granules preferred particle sizes are from 50 to 1000 μ m, preferable from 50 to 500 μ m, most preferably from 100 – 250 μ m. For the capsules preferred particle sizes are from 1 to 1000 μ m, preferably 5 to 200 μ m, most preferably from 10 to 100 μ m.

Addition of the enzymes in a slurried solid or encapsulated form is particularly advantageous for worker safety as it serves to minimize any aerosolization risk during manufacture as the traditional risks from solid forms (dusting) and liquid forms (fine aerosols, particularly those of < 5 μ m) are mitigated.

Liquid Phase

The liquid laundry detergent composition of the present invention comprises a liquid phase into which the solid enzyme particle is dispersed.

The liquid phase comprises between 5% and 60% by weight of the liquid of an alcohol. The alcohol is described in more detail below.

The liquid phase may comprise a natural or synthetically derived fatty alcohol ethoxylate non-ionic surfactant. Preferred synthetically derived fatty alcohol ethoxylate non-ionic surfactant or those derived from the oxo-synthesis process, or so-called oxo-synthesised non-ionic surfactants. The composition may comprise from 0% to 30% or even from 0.1% to 25% by weight of the composition of fatty alcohol ethoxylate non-ionic surfactant.

The ethoxylated nonionic surfactant may be, e.g., primary and secondary alcohol ethoxylates, especially the C₈-C₂₀ aliphatic alcohols ethoxylated with an average of from 1 to 50 or even 20 moles of ethylene oxide per mole of alcohol, and more especially the C₁₀-C₁₅ primary and secondary aliphatic alcohols ethoxylated with an average of from 1 to 10 moles of ethylene oxide per mole of alcohol.

The ethoxylated alcohol non-ionic surfactant can be, for example, a condensation product of from 3 to 8 mol of ethylene oxide with 1 mol of a primary alcohol having from 9 to 15 carbon atoms.

The non-ionic surfactant may comprise a fatty alcohol ethoxylate of formula R(EO)_n, wherein R represents an alkyl chain between 4 and 30 carbon atoms, (EO) represents one unit of ethylene oxide monomer and n has an average value between 0.5 and 20.

The composition may comprise other non-ionic surfactants, preferably natural or synthetic non-ionic surfactants.

Alcohol

5 The liquid phase comprises between 5% and 40%, or even between 5% and 20% or even between 5% and 15% by weight of the composition of an alcohol, preferably, wherein the alcohol has a molecular weight of between 20 and 400 and an eRH of between 50% and 80%, or even between 52% and 75% at 20°C as measured via the alcohol eRH test.

10 The alcohol eRH test comprises the steps of preparing a solution of 80% alcohol in deionised water, followed by adding this to a calibrated Rotronic Hygrolab meter (in a plastic sample liner of 14mm depth) at room temperature (20°C +/- 1°C) and allowing this to equilibrate for 25 minutes, and finally measuring the eRH recorded. The volume of sample used was sufficient to fill the plastic sample liner.

15 By 'alcohol' we herein mean either a single compound or a mixture of compounds that when taken together collectively each have a molecular weight of between 20 and 400 and an overall eRH of the compound or mixture of between 50% and 80% at 20°C as measured via the alcohol eRH test. Without wishing to be bound by theory, an alcohol is any compound comprising at least one OH unit, preferably polyols and diols, more preferably diols. Preferred diols included glycols.

20 The alcohol may be selected from the group comprising ethylene glycol, 1,3 propanediol, 1,2 propanediol, tetramethylene glycol, pentamethylene glycol, hexamethylene glycol, 2,3-butane diol, 1,3 butanediol, diethylene glycol, triethylene glycol, polyethylene glycol, glycerol formal, dipropylene glycol, polypropylene glycol, dipropylene glycol n-butyl ether, and mixtures thereof.

25 Preferably the alcohol may be selected from the group comprising ethylene glycol, 1,2 propanediol, 2,3-butane diol, 1,3 butanediol, triethylene glycol, polyethylene glycol, glycerol formal, dipropylene glycol, polypropylene glycol, dipropylene glycol n-butyl ether, and mixtures thereof.

30 More preferably the alcohol is selected from the group comprising 1,2 propanediol, dipropylene glycol, polypropylene glycol, 2,3-butane diol, dipropylene glycol n-butyl ether and mixtures thereof.

 Most preferably the alcohol may be selected from the group comprising 1,2 propanediol, dipropylene glycol, polypropylene glycol, dipropylene glycol n-butyl ether and mixtures thereof.

Anionic surfactant

The liquid laundry detergent composition comprises from 10% to 30% by weight of the composition of a non-amine neutralized linear alkylbenzene sulphonate. The linear
5 alkylbenzene sulphonate may be present in the liquid or may be present as a solid, or a mixture thereof. If the linear alkylbenzene sulphonate is present as a solid, preferably it is in the form of a lamellar liquid crystal alkylbenzene sulphonate. By 'lamellar liquid crystal' we herein mean the system being in a state where the surfactant molecules are organised in stacks of bilayers of surfactant in the melted state separated by thin layers of solvent. This structure has both liquid
10 properties in term of flowability as well as solid properties in term of being structured. The structure is characterised by its d-spacing, the sum of the bilayer thickness and the solvent layer between sheets. The repetition and periodicity of this structure yields to sharp x-ray diffraction peaks characteristic of crystal phases.

Non-amine neutralized linear alkylbenzene sulphonates are those in which the linear
15 alkylbenzene sulphonic acid is neutralized to the correspond linear alkylbenzene sulphonate salt using a neutralizing material other than an amine. Non-limiting examples of such neutralizing groups include sodium, potassium, magnesium and mixtures thereof. The non-amine neutralized linear alkylbenzene sulphonate may be a sodium linear alkylbenzene sulphonate, a potassium alkylbenzene sulphonate, a magnesium alkylbenzene sulphonate or a mixture thereof.

20 Exemplary linear alkylbenzene sulphonates are C₁₀-C₁₆ alkyl benzene sulfonic acids, or C₁₁-C₁₄ alkyl benzene sulfonic acids. By 'linear', we herein mean the alkyl group is linear. Alkyl benzene sulfonates are well known in the art. Especially useful are the sodium, potassium and magnesium linear straight chain alkylbenzene sulfonates in which the average number of carbon atoms in the alkyl group is from about 11 to 14.

25 The cleaning technology may comprise an amine neutralized anionic surfactant, preferably an amine neutralized linear alkylbenzene sulphonate, an amine neutralized alkyl sulphate or a mixture thereof. The liquid laundry detergent composition may comprise an amine neutralized anionic surfactant in the solid phase, liquid phase or both.

The liquid laundry detergent composition may comprise an alkyl sulphate anionic
30 surfactant. The liquid laundry detergent composition may comprise from 10% to 30% or even from 15% to 25% by weight of the laundry detergent composition of alkyl sulphate anionic surfactant.

The alkyl sulphate anionic surfactant may be non-amine neutralised, amine neutralised or a mixture thereof, preferably the alkyl sulphate is non-amine neutralised. The non-amine

neutralized alkyl sulphate may be a sodium alkyl sulphate, a potassium alkyl sulphate, a magnesium alkyl sulphate or a mixture thereof.

The alkyl sulphate anionic surfactant may be present in the liquid or may be present as a solid, or a mixture thereof. If the alkyl sulphate is present as a solid, preferably it is in the form of a lamellar liquid crystal alkyl sulphate. The alkyl sulphate anionic surfactant may be alkoxyated or non-alkoxyated or a mixture thereof. The alkyl sulphate anionic surfactant may be a C₁₀-C₂₀ primary, branched-chain and random alkyl sulfates (AS), including predominantly C₁₂ alkyl sulfates. Alternatively, the alkyl sulphate anionic surfactant may be a C₁₀-C₁₈ secondary (2,3) alkyl sulfates. Alternatively, the alkyl sulphate anionic surfactant may be a C₁₀-C₁₈ alkyl alkoxy sulfates (AE_xS) wherein x is from 1-30. Alternatively, the alkyl sulphate anionic surfactant may be a mixture of all the above alkyl sulphate anionic surfactants. Non-limiting examples of suitable cations for the alkyl sulphate anionic surfactant include sodium, potassium, ammonium, amine and mixtures thereof.

Amine

The detergent composition comprises less than 5% by weight of the composition of a hydroxyl-containing amine compound, or even from 0.1% to 5%, or even from 0.1% to 4% by weight of the composition of a hydroxyl-containing amine compound. By 'hydroxyl-containing amine compound' we herein mean a compound comprising an alcohol (OH) group and an amine group. The hydroxyl-containing amine compound may be selected from monoethanolamine, triethanolamine, diisopropanolamine, triisopropanolamine, Monoamino hexanol, 2-[(2-methoxyethyl) methylamino]- ethanol, Propanolamine, N-Methylethanolamine, diethanolamine, Monobutanol amine, Isobutanolamine, Monopentanol amine, 1-Amino-3-(2-methoxyethoxy)- 2-propanol, 2-Methyl-4-(methylamino)- 2-butanol, 6-amino-1-hexanol, Heptaminol, Isoetarine, Norepinephrine, Sphingosine, Phenylpropanolamine and mixtures thereof.

The hydroxyl-containing amine compound may be selected from the group comprising monoethanol amine, triethanolamine and mixtures thereof.

Preferably, the hydroxyl-containing amine compound has a molecular weight of less than 500, or even less than 250.

The detergent composition may comprise other amine containing compounds.

Structurant

The composition of the present invention may comprises less than 2% by weight of the composition of a structurant. If a structurant is present, preferably the composition comprises

from 0.05% to 2%, preferably from 0.1% to 1% by weight of a structurant. The structurant may be selected from non-polymeric or polymeric structurants. The structurant may be a non-polymeric structurant, preferably a crystallisable glyceride. The structurant may be a polymeric structurant, preferably a fibre based polymeric structurant, more preferably a cellulose fibre-based structurant. The structurant may be selected from crystallisable glyceride, cellulose-fibre based structurants, TiO_2 , silica and mixtures thereof.

Suitable structurants are preferably ingredients which impart a sufficient yield stress or low shear viscosity to stabilize the liquid laundry detergent composition independently from, or extrinsic from, any structuring effect of the deterative surfactants of the composition. Preferably, they impart to the laundry detergent composition a high shear viscosity at 20 sec⁻¹ at 21°C of from 1 to 1500 cps and a viscosity at low shear (0.05 sec⁻¹ at 21°C) of greater than 5000 cps. The viscosity is measured using an AR 550 rheometer from TA instruments using a plate steel spindle at 40 mm diameter and a gap size of 500 μm . The high shear viscosity at 20s⁻¹ and low shear viscosity at 0.5s⁻¹ can be obtained from a logarithmic shear rate sweep from 0.1-1 to 25-1 in 3 minutes time at 21°C.

The composition may comprise a non-polymeric crystalline, hydroxyl functional structurant. Such non-polymeric crystalline, hydroxyl functional structurants generally comprise a crystallizable glyceride which can be pre-emulsified to aid dispersion into the final liquid laundry detergent composition. A non-limiting example of such a pre-emulsified external structuring system comprises: (a) crystallizable glyceride(s); (b) anionic surfactant; and (c) water and optionally, non-aminofunctional organic solvents. Each of these components is discussed in detail below.

The structurant may be a polymeric crystalline, hydroxy-functional structurant that comprises a crystallizable glyceride, preferably hydrogenated castor oil or "HCO". HCO as used herein most generally can be any hydrogenated castor oil or derivative thereof, provided that it is capable of crystallizing in the non-polymeric crystalline, hydroxy-functional structurant premix. Castor oils may include glycerides, especially triglycerides, comprising C_{10} to C_{22} alkyl or alkenyl moieties which incorporate a hydroxyl group. Hydrogenation of castor oil, to make HCO, converts the double bonds which may be present in the starting oil as ricinoleyl moieties. As such, the ricinoleyl moieties are converted into saturated hydroxyalkyl moieties, e.g., hydroxystearyl. The HCO herein may be selected from: trihydroxystearin; dihydroxystearin; and mixtures thereof. The HCO may be processed in any suitable starting form, including, but not limited to those selected from solid, molten and mixtures thereof. HCO is typically present at a level of from 2% to 10%, from 3% to 8%, or from 4% to 6% by weight in the external

structuring system. The corresponding percentage of hydrogenated castor oil delivered into a finished laundry detergent product may be below 1.0%, typically from 0.1% to 0.8%. HCO may be present at a level of between 0.01% and 1%, or even between 0.05% and 0.8% by weight of the laundry detergent composition.

5 HCO of use in the present invention includes those that are commercially available. Non-limiting examples of commercially available HCO of use in the present invention include: THIXCIN® from Rheox, Inc. Further examples of useful HCO may be found in U.S. Patent 5,340,390.

10 While the use of hydrogenated castor oil is preferred, any crystallisable glyceride can be used within the scope of the invention. Preferred crystallisable glyceride(s) have a melting point of from 40 °C to 100 °C.

The structurant may comprise a fibre-based structurant. The structurant may comprise a microfibrillated cellulose (MFC), which is a material composed of nanosized cellulose fibrils, typically having a high aspect ratio (ratio of length to cross dimension). Typical lateral
15 dimensions are 1 to 100, or 5 to 20 nanometres, and longitudinal dimension is in a wide range from nanometres to several microns. For improved structuring, the microfibrillated cellulose preferably has an average aspect ratio (l/d) of from 50 to 200,000, more preferably from 100 to 10,000. Microfibrillated cellulose can be derived from any suitable source, including bacterial cellulose, citrus fibers, and vegetables such as sugar beet, chicory root, potato, carrot, and the
20 like.

The structurant may be selected from the group consisting of titanium dioxide, tin dioxide, any forms of modified TiO₂, TiO₂ or stannic oxide, bismuth oxychloride or bismuth oxychloride coated TiO₂, silica coated TiO₂ or metal oxide coated TiO₂ and mixtures thereof. Modified TiO₂ may comprise carbon modified TiO₂, metallic doped TiO₂ or mixtures thereof.
25 Metallic doped TiO₂ may be selected from platinum doped TiO₂, Rhodium doped TiO₂.

The structurant may comprise silica. Those skilled in the art will know suitable silica materials to use. The silica may comprise fumed silica.

Water and equilibrium relative humidity

30 The liquid laundry detergent composition comprises between 0.5% and 15% by weight of the composition of water. The liquid laundry detergent composition may comprise between 0.5% and 12%, or even between 0.5% and 10% by weight of the composition of water.

The liquid laundry detergent composition may have an equilibrium relative humidity of less than 65% at 20°C.

A preferred method for measuring the eRH of the composition is via the composition eRH test. The composition eRH test comprises the steps of adding a sample of the composition to a calibrated Rotronic Hygrolab meter (in a plastic sample liner of 14mm depth) at room temperature (20°C +/- 1°C) and allowing this to equilibrate for 25 minutes, and finally measuring the eRH recorded. The volume of sample used was sufficient to fill the plastic sample liner.

Adjunct ingredients

The liquid laundry detergent composition may comprise an adjunct ingredient. The adjunct ingredient may be selected from the group comprising bleach, bleach catalyst, dye, hueing dye, cleaning polymers including alkoxylated polyamines and polyethyleneimines, soil release polymer, surfactant, solvent, dye transfer inhibitors, chelant, enzyme, perfume, encapsulated perfume, polycarboxylates, structurant and mixtures thereof.

Water-soluble pouch

The liquid laundry detergent composition may be present in a water-soluble unit dose article wherein the composition comprises between 0.5% and 15%, preferably between 0.5% and 12%, more preferably between 0.5% and 10% by weight of the composition of water. In such an embodiment, the water-soluble unit dose article comprises at least one water-soluble film shaped such that the unit-dose article comprises at least one internal compartment surrounded by the water-soluble film. The at least one compartment comprises the liquid laundry detergent composition. The water-soluble film is sealed such that the liquid laundry detergent composition does not leak out of the compartment during storage. However, upon addition of the water-soluble unit dose article to water, the water-soluble film dissolves and releases the contents of the internal compartment into the wash liquor.

The compartment should be understood as meaning a closed internal space within the unit dose article, which holds the composition. Preferably, the unit dose article comprises a water-soluble film. The unit dose article is manufactured such that the water-soluble film completely surrounds the composition and in doing so defines the compartment in which the composition resides. The unit dose article may comprise two films. A first film may be shaped to comprise an open compartment into which the composition is added. A second film is then laid over the first film in such an orientation as to close the opening of the compartment. The first and second films are then sealed together along a seal region. The film is described in more detail below.

The unit dose article may comprise more than one compartment, even at least two compartments, or even at least three compartments. The compartments may be arranged in superposed orientation, i.e. one positioned on top of the other. Alternatively, the compartments may be positioned in a side-by-side orientation, i.e. one orientated next to the other. The compartments may even be orientated in a 'tyre and rim' arrangement, i.e. a first compartment is positioned next to a second compartment, but the first compartment at least partially surrounds the second compartment, but does not completely enclose the second compartment. Alternatively one compartment may be completely enclosed within another compartment.

The film of the present invention is soluble or dispersible in water. The water-soluble film preferably has a thickness of from 20 to 150 micron, preferably 35 to 125 micron, even more preferably 50 to 110 micron, most preferably about 76 micron.

Preferably, the film has a water-solubility of at least 50%, preferably at least 75% or even at least 95%, as measured by the method set out here after using a glass-filter with a maximum pore size of 20 microns:

5 grams $\pm$ 0.1 gram of film material is added in a pre-weighed 3L beaker and 2L $\pm$ 5ml of distilled water is added. This is stirred vigorously on a magnetic stirrer, Labline model No. 1250 or equivalent and 5 cm magnetic stirrer, set at 600 rpm, for 30 minutes at 30°C. Then, the mixture is filtered through a folded qualitative sintered-glass filter with a pore size as defined above (max. 20 micron). The water is dried off from the collected filtrate by any conventional method, and the weight of the remaining material is determined (which is the dissolved or dispersed fraction). Then, the percentage solubility or dispersability can be calculated.

Preferred film materials are preferably polymeric materials. The film material can, for example, be obtained by casting, blow-moulding, extrusion or blown extrusion of the polymeric material, as known in the art.

Preferred polymers, copolymers or derivatives thereof suitable for use as pouch material are selected from polyvinyl alcohols, polyvinyl pyrrolidone, polyalkylene oxides, acrylamide, acrylic acid, cellulose, cellulose ethers, cellulose esters, cellulose amides, polyvinyl acetates, polycarboxylic acids and salts, polyaminoacids or peptides, polyamides, polyacrylamide, copolymers of maleic/acrylic acids, polysaccharides including starch and gelatine, natural gums such as xanthum and carragum. More preferred polymers are selected from polyacrylates and water-soluble acrylate copolymers, methylcellulose, carboxymethylcellulose sodium, dextrin, ethylcellulose, hydroxyethyl cellulose, hydroxypropyl methylcellulose, maltodextrin, polymethacrylates, and most preferably selected from polyvinyl alcohols, polyvinyl alcohol copolymers and hydroxypropyl methyl cellulose (HPMC), and combinations thereof. Preferably,

the level of polymer in the pouch material, for example a PVA polymer, is at least 60%. The polymer can have any weight average molecular weight, preferably from about 1000 to 1,000,000, more preferably from about 10,000 to 300,000 yet more preferably from about 20,000 to 150,000.

5 Mixtures of polymers can also be used as the pouch material.

Preferred films exhibit good dissolution in cold water, meaning unheated distilled water. Preferably such films exhibit good dissolution at temperatures of 24°C, even more preferably at 10°C. By good dissolution it is meant that the film exhibits water-solubility of at least 50%, preferably at least 75% or even at least 95%, as measured by the method set out here after using
10 a glass-filter with a maximum pore size of 20 microns, described above.

Preferred films are those supplied by Monosol under the trade references M8630, M8900, M8779, M8310.

The film may be opaque, transparent or translucent.

The area of print may be achieved using standard techniques, such as flexographic
15 printing or inkjet printing.

The film may comprise an aversive agent, for example a bittering agent. Suitable bittering agents include, but are not limited to, naringin, sucrose octaacetate, quinine hydrochloride, denatonium benzoate, or mixtures thereof. Any suitable level of aversive agent may be used in the film. Suitable levels include, but are not limited to, 1 to 5000ppm, or even
20 100 to 2500ppm, or even 250 to 2000rpm.

Method of making

The liquid laundry detergent composition of the present invention may be made using any suitable manufacturing techniques known in the art. Those skilled in the art would know
25 appropriate methods and equipment to make the composition according to the present invention.

A preferred process comprises the step of adding the solid phase wherein the solid phase comprises particles wherein the particles have a mean particle size distribution of less than 500µm.

30 Method of use

The composition or unit dose article of the present invention can be added to a wash liquor to which laundry is already present, or to which laundry is added. It may be used in an washing machine operation and added directly to the drum or to the dispenser drawer. The washing machine may be an automatic or semi-automatic washing machine. It may be used in

combination with other laundry detergent compositions such as fabric softeners or stain removers. It may be used as pre-treat composition on a stain prior to being added to a wash liquor.

The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and a functionally equivalent range surrounding that value. For example, a dimension disclosed as “40 mm” is intended to mean “about 40 mm.”

EXAMPLES

The viscosity of various compositions were compared. The following compositions were prepared;

Table 1

	Compositions (wt%)		
	A	B	C
water	7.16	7.16	7.44
Dipropylene glycol	14.66	14.66	31.19
1,2-propanediol		10.00	
Dipropylene glycol n-butyl ether	9.80	9.80	
Glycerol	15.00	5.00	5.00
Linear alkylbenzene sulphonate neutralized with monoethanolamine			23.59
Linear alkylbenzene sulphonate neutralized with sodium carbonate	23.59	23.59	
Ethoxylated polyethyleneimine	2.16	2.16	2.16
Alkyl sulphate with an average degree of ethoxylation of 3, neutralized with monoethanolamine			11.00
Alkyl sulphate with an average degree of ethoxylation of 3, neutralized with sodium carbonate	11.00	11.00	
HEDP	1.81	1.81	1.81
Amphiphilic graft copolymer	2.72	2.72	2.72
Brightener 49	0.24	0.24	0.24
Soil release polymer commercially available from Clariant as SRA-300	0.32	0.32	0.32
Carboxymethyl cellulose	1.07	1.07	1.07
Siloxane polymeric suds suppressor	0.13	0.13	0.13
Perfume	2.68	2.68	2.68
protease	0.10	0.10	0.10
TiO ₂	0.50	0.50	0.50
palm kernel fatty acid	3.26	3.26	3.26

22

Guerbet alcohol non-ionic surfactant commercially available from BASF as Lutensol XL100	0.56	0.56	0.56
minors	2.36	2.36	2.77

The compositions were made by preparing a 1L beaker having an IKA Eurostar 200 mixer with 10cm impeller. This was operated at 250rpm. To the beaker with the rotating impeller, the solvent materials were added, followed by the surfactant materials. Once these had dispersed, the polymers and salts were added. The pH of the composition was adjusted using NaOH to approximately 8 (measured using a Sartorius PT-10 pH meter). Remaining ingredients were then added and mixed. All materials were weighed out using a Mettler Toledo PB3002-S balance.

Composition C comprised 6.25% by weight of composition C of monoethanolamine.

Compositions A and B comprised no monoethanolamine.

The viscosity of the compositions were then measured using a Rheometer DHR1 from TA instruments using a gap of 1000 μ m at 20°C. Samples were equilibrated for 1 min at 0.05s⁻¹ followed by a measured flow curve from 0.05s⁻¹ to 1200s⁻¹ over 10 mins. Results for 0.05s⁻¹ and 1000s⁻¹ are shown in Table 2.

Table 2

	0.05s ⁻¹	1000s ⁻¹
	mPa.s	mPa.s
A	1560	870
B	1112	413
C	1310	315

Shear at 0.05s⁻¹ corresponds to that experienced by the composition during pouring of the composition by the consumer. Shear at 1000s⁻¹ corresponds to that experienced by the composition during manufacture.

Composition C which comprises 6.25% monoethanolamine shows an acceptable viscosity profile at low and high shear corresponding to consumer pouring shear and process dosing shear.

However, when the monoethanolamine is removed in composition A (and correspondingly the surfactants are neutralized with sodium carbonate), there is an increase in viscosity to unacceptable levels.

Composition B corresponds to the present invention in which the monoethanolamine has been removed and the surfactants neutralized with sodium carbonate, but also 1,2-propandiol has been added. The viscosity returns to acceptable levels.

CLAIMS

What is claimed is:

1. A liquid laundry detergent composition comprising;
 - a. a liquid phase;
 - b. between 0.5% and 15% by weight of the liquid detergent composition of a solid enzyme particle,wherein the solid phase is dispersed within the liquid phase and wherein the water-soluble solid phase is defined as the solid obtained when the liquid laundry detergent composition is centrifuged at 1200 G for 10 mins; and
wherein the liquid phase comprises between 5% and 40% by weight of the liquid of an alcohol selected from the group comprising ethylene glycol, 1,3 propanediol, 1,2 propanediol, tetramethylene glycol, pentamethylene glycol, hexamethylene glycol, 2,3-butane diol, 1,3 butanediol, diethylene glycol, triethylene glycol, polyethylene glycol, glycerol formal, dipropylene glycol, polypropylene glycol, dipropylene glycol n-butyl ether, and mixtures thereof, preferably the alcohol is selected from the group comprising 1,2 propanediol, dipropylene glycol, polypropylene glycol, 2,3- butane diol, dipropylene glycol n-butyl ether and mixtures thereof; and
from 10% to 30% by weight of the composition of a non-amine neutralized linear alkylbenzene sulphonate and wherein the liquid laundry detergent composition comprises less than 10% by weight of the liquid laundry detergent composition of an amine-neutralised anionic surfactant; and
wherein the composition comprises between 0.5% and 50% by weight of the composition of water; and
wherein the composition comprises less than 5% by weight of the composition of a hydroxyl-containing amine.
2. The detergent composition according to claim 1, wherein the enzyme is selected from the group comprising hemicellulases, peroxidases, proteases, cellulases, xylanases, lipases, phospholipases, esterases, cutinases, pectinases, keratanases, reductases, oxidases, phenoloxidases, lipxygenases, ligninases, pullulanases, tannases, pentosanases, malanases, β -glucanases, arabinosidases, hyaluronidase, chondroitinase, laccase, and amylases, or mixtures thereof. .

3. The detergent composition according to any preceding claims, wherein the composition comprises between 0.5% and 10% or even between 0.5% and 5% by weight of the composition of the solid enzyme particle.
4. The detergent composition according to any preceding claims, wherein the liquid phase comprises between 5% and 20%, or even between 5% and 15% by weight of the composition of the alcohol.
5. The detergent composition according to any preceding claims, wherein the solvent is selected from the group comprising ethylene glycol, 1,3 propanediol, 1,2 propanediol, tetramethylene glycol, pentamethylene glycol, hexamethylene glycol, 2,3-butane diol, 1,3 butanediol, diethylene glycol, triethylene glycol, polyethylene glycol, glycerol formal dipropylene glycol, polypropylene glycol, dipropylene glycol n-butyl ether, and mixtures thereof, preferably the solvent is selected from the group comprising 1,2 propanediol, dipropylene glycol, polypropylene glycol, 2,3- butane diol, dipropylene glycol n-butyl ether and mixtures thereof.
6. The detergent composition according to any preceding claims, wherein the liquid phase comprises alkyl sulphate anionic surfactant, a natural or synthetically derived fatty alcohol ethoxylate non-ionic surfactant or a mixture thereof.
7. The detergent composition according to any preceding claims comprising from 0% to 5%, or even from 0.1% to 5%, or even from 0.1% to 4% by weight of the composition of a hydroxyl-containing amine, preferably wherein the hydroxyl-containing amine is selected from the group comprising monoethanol amine, triethanolamine and mixtures thereof.
8. The detergent composition according to any preceding claims comprising a structurant.
9. The detergent composition according to any preceding claims, wherein the solid particles have a mean particle size distribution of between 2 μ m and 50 μ m.
10. The detergent composition according to any preceding claims comprising a perfume raw material, wherein the perfume raw material is preferably selected from aldehydes, ketones or a mixture thereof.

11. The detergent composition according to any preceding claims comprising an adjunct ingredient, wherein the adjunct ingredient is selected from the group comprising bleach, bleach catalyst, dye, hueing dye, cleaning polymers including alkoxyated polyamines and polyethyleneimines, soil release polymer, surfactant, solvent, dye transfer inhibitors, chelant, enzyme, perfume, encapsulated perfume, polycarboxylates, structurant and mixtures thereof.
12. A water-soluble unit dose article comprising a water-soluble film and a detergent composition according to any preceding claims wherein the composition comprises between 0.5% and 15% by weight of the composition of water.
13. The unit dose article according to claim 12 wherein the unit dose article comprises at least two compartments, or even at least three or even at least four compartments.
14. A process of making a composition according to any preceding claims, comprising the step of adding the solid enzyme particle wherein the solid particles have a mean particle size distribution of less than 500 μ m.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/035372

A. CLASSIFICATION OF SUBJECT MATTER

INV. C11D1/22 C11D1/42 C11D3/20 C11D3/386 C11D17/00
C11D17/04

ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98/00518 A1 (PROCTER & GAMBLE) 8 January 1998 (1998-01-08) claims; tables I,II -----	1-14
X	WO 98/00514 A1 (PROCTER & GAMBLE) 8 January 1998 (1998-01-08) table II -----	1-14
X	WO 99/00471 A1 (PROCTER & GAMBLE) 7 January 1999 (1999-01-07) table I -----	1-14



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

28 July 2016

Date of mailing of the international search report

05/08/2016

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Hillebrecht, Dieter

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2016/035372

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9800518	A1	08-01-1998	AR 008054 A1 09-12-1999
			BR 9710068 A 10-08-1999
			CA 2258993 A1 08-01-1998
			EP 0907716 A1 14-04-1999
			JP H11514030 A 30-11-1999
			WO 9800518 A1 08-01-1998

WO 9800514	A1	08-01-1998	AR 008624 A1 09-02-2000
			AT 212373 T 15-02-2002
			AU 3495497 A 21-01-1998
			BR 9710066 A 10-08-1999
			CA 2258702 A1 08-01-1998
			CN 1228116 A 08-09-1999
			CZ 9804183 A3 14-07-1999
			DE 69710069 D1 14-03-2002
			DE 69710069 T2 17-10-2002
			EP 0907713 A1 14-04-1999
			JP 3233944 B2 04-12-2001
			JP H11514026 A 30-11-1999
			MA 24234 A1 31-12-1997
			US 6576602 B1 10-06-2003
			WO 9800514 A1 08-01-1998
			ZA 9705707 B 26-01-1998

WO 9900471	A1	07-01-1999	AR 016099 A1 20-06-2001
			BR 9810633 A 03-10-2000
			CA 2295142 A1 07-01-1999
			EP 0991746 A1 12-04-2000
			JP 2002512654 A 23-04-2002
			WO 9900471 A1 07-01-1999
