(19) ORGANISATION AFRICAINE DE LA PROPRIETE INTELLECTUELLE

(51) Inter. Cl.[8] A01N 41/06 (2018.01); A61K 31/18 (2018.01)

(11) N° 20932

FASCICULE DE BREVET D'INVENTION

(21) Numéro de dépôt : 1201700130
PCT/US2015/054316

(22) Date de dépôt : 06/10/2015

(30) Priorité(s) :
US n° 62/060,182 du 06/10/2014;
US n° 62/114,767 du 11/02/2015;
US n° 62/153,120 du 27/04/2015

(24) Délivré le : 05/05/2023

(45) Publié le : 24.07.2023

(54) Titre : **Modulators of cystic fibrosis transmembrane conductance regulator.**

(73) Titulaire(s) :


Vertex Pharmaceuticals Incorporated,
50 Northern Avenue, 15th Floor,
BOSTON 02210,
Massachusetts (US)

(72) Inventeur(s) :

ANDERSON, Corey (US);
BEAR, Brian Richard (US);
HADIDA-RUAH, Sara Sabina (US);
TERMIN, Andreas P. (US);
MILLER, Mark Thomas (US);
ARUMUGAM, Vijay-alaksmi (US);
BINCH, Hayley Marie (US);
CLEMENS, Jeremy J. (US);
CLEVELAND, Thomas (US);
CONROY, Erica (US);
COON, Timothy Richard (US);
FRIEMAN, Bryan A. (US);
GROOTENHUIS, Peter Diederik Jan (US);
GROSS, Raymond Stanley (US);
HARIPADA, Khatuya (US);
JOSHI, Pramod Virupax (US);
KRENITSKY, Paul John (US);
LIN, Chun-Chieh (US);
MARELIUS, Gulin Erdogan (US);
MELILLO, Vito (US);
MCCARTNEY, Jason (US);
NICHOLLS, Georgia McGaughey (US);
PIERRE, Fabrice Jean Denis (US);
SILINA, Alina (US);
UY, Johnny (US);
ZHOU, Jinglan (US)

(57) Abrégé :

The present invention features a compound of formula (I) :



I,

(I) or a pharmaceutically acceptable salt thereof, where R1, R2, R3, W, X, Y, Z, n, o, p, and q are defined herein, for the treatment of CFTR mediated diseases, such as cystic fibrosis. The present invention also features pharmaceutical compositions, method of treating, and kits thereof.

Formula (I)

(74) Mandataire : **SPOOR & FISHER Inc. NGWAFOR & PARTNERS Sarl, The House of Gideon, Golf/Bastos Quarters, Opposite The American Embassy, Entrance-Saint John Paul II Boulevard, P.O. Box 8211, YAOUNDE (CM).**

Modulators of Cystic Fibrosis Transmembrane Conductance Regulator

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of priority under 35 U.S.C. § 119 to United States provisional patent application serial numbers 62/060,182, filed October 6, 2014; 62/114,767, filed February 11, 2015; and 62/153,120, filed April 27, 2015, the entire contents of all applications are incorporated herein by reference.

TECHNICAL FIELD OF THE INVENTION

The present invention features modulators of Cystic Fibrosis Transmembrane Conductance Regulator (CFTR), pharmaceutical compositions, methods of treatment, and kits thereof.

BACKGROUND OF THE INVENTION

Cystic fibrosis (CF) is a recessive genetic disease that affects approximately 30,000 children and adults in the United States and approximately 30,000 children and adults in Europe. Despite progress in the treatment of CF, there is no cure.

In patients with CF, mutations in CFTR endogenously expressed in respiratory epithelia leads to reduced apical anion secretion causing an imbalance in ion and fluid transport. The resulting decrease in anion transport contributes to enhanced mucus accumulation in the lung and the accompanying microbial infections that ultimately cause death in CF patients. In addition to respiratory disease, CF patients typically suffer from gastrointestinal problems and pancreatic insufficiency that, if left untreated, results in death. In addition, the majority of males with cystic fibrosis are infertile and fertility is decreased among females with cystic fibrosis. In contrast to the severe effects of two copies of the CF associated gene, individuals with a single copy of the CF associated gene exhibit increased resistance to cholera and to dehydration resulting from diarrhea – perhaps explaining the relatively high frequency of the CF gene within the population.

Sequence analysis of the CFTR gene of CF chromosomes has revealed a variety of disease causing mutations (Cutting, G. R. et al. (1990) Nature 346:366-369; Dean, M. et al.


(1990) Cell 61:863:870; and Kerem, B-S. et al. (1989) Science 245:1073-1080; Kerem, B-S et al. (1990) Proc. Natl. Acad. Sci. USA 87:8447-8451). To date, greater than 1000 disease causing mutations in the CF gene have been identified (http://cftr2.org). The most prevalent mutation is a deletion of phenylalanine at position 508 of the CFTR amino acid sequence, and is commonly referred to as F508del. This mutation occurs in approximately 70% of the cases of cystic fibrosis and is associated with a severe disease.

The deletion of residue 508 in F508del prevents the nascent protein from folding correctly. This results in the inability of the mutant protein to exit the ER, and traffic to the plasma membrane. As a result, the number of channels present in the membrane is far less than observed in cells expressing wild-type CFTR. In addition to impaired trafficking, the mutation results in defective channel gating. Together, the reduced number of channels in the membrane and the defective gating lead to reduced anion transport across epithelia leading to defective ion and fluid transport. (Quinton, P. M. (1990), FASEB J. 4: 2709-2727). Studies have shown, however, that the reduced numbers of F508del in the membrane are functional, albeit less than wild-type CFTR. (Dalemans et al. (1991), Nature Lond. 354: 526-528; Denning et al., supra; Pasyk and Foskett (1995), J. Cell. Biochem. 270: 12347-50). In addition to F508del, other disease causing mutations in CFTR that result in defective trafficking, synthesis, and/or channel gating could be up- or down-regulated to alter anion secretion and modify disease progression and/or severity.

Accordingly, there is a need for novel treatments of CFTR mediated diseases.

SUMMARY OF THE INVENTION

The invention features a compound of formula I:



I,

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring A is a C6-C10 aryl ring; C3-C10 cycloalkyl ring; or a C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring B is a C3-C10 cycloalkyl ring; a C6-C10 aryl ring; or a C4-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C6-C10 aryl ring; a C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, NR, O, or S; or a C3-C10 cycloalkyl ring;

W is O, NR, or S;

X is O or NR;

Y is independently CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

Z is NR or CHR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy;C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are optionally and independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; OH; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH;CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;

o is 0, 1, 2, 3, 4, or 5;

p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5;

---- is a single bond or a double bond;

provided that the moieties containing ring B and ring C are substituted at adjacent positions on ring A.

The present invention also features phanmaceutical compositions thereof, which may include additional agents, and methods of treating CFTR mediated diseases, such as cystic fibrosis, comprising administering compounds of formula I to a subject in need thereof. The present invention also features kits comprising compounds of formula I.

BRIEF DESCRIPTION OF DRAWINGS

Figure 1 discloses a list of CFTR mutations by name, protein name, and legacy name that in one aspect of the invention a patient may possess and be treatable by the compounds and compositions of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

As used herein, "CFTR" stands for cystic fibrosis transmembrane conductance regulator.

As used herein, "mutations" can refer to mutations in the *CFTR* gene or the CFTR protein. A "*CFTR* mutation" refers to a mutation in the *CFTR* gene, and a "CFTR mutation" refers to a mutation in the CFTR protein. A genetic defect or mutation, or a change in the nucleotides in a gene in general results in a mutation in the CFTR protein translated from that gene.

As used herein, a "F508del mutaion" or "F508del" is a specific mutation within the CFTR protein. The mutation is a deletion of the three nucleotides that comprise the codon for amino acid phenylalanine at position 508, resulting in CFTR protein that lacks this phenylalanine residue.

The term "*CFTR* gating mutation" as used herein means a *CFTR* mutation that results in the production of a CFTR protein for which the predominant defect is a low channel open probability compared to normal CFTR (Van Goor, F., Hadida S. and Grootenhuis P., "Pharmacological Rescue of Mutant CFTR function for the Treatment of Cystic Fibrosis", Top. Med. Chem. 3: 91-120 (2008)). Gating mutations include, but are not limited to, G551D, G178R, S549N, S549R, G551S, G970R, G1244E, S1251N, S1255P, and G1349D.

As used herein, a patient who is "homozygous" for a particular mutation, e.g. F508del, has the same mutation on each allele.

As used herein, a patient who is "heterozygous" for a particular mutation, e.g. F508del, has this mutation on one allele, and a different mutation on the other allele.

As used herein, the term "modulator" refers to a compound that increases the activity of a biological compound such as a protein. For example, a CFTR modulator is a compound that increases the activity of CFTR. The increase in activity resulting from a CFTR modulator may be through a corrector mechanism or a potentiator mechanism as described below.

As used herein, the term "CFTR corrector" refers to a compound that increases the amount of functional CFTR protein to the cell surface, resulting in enhanced ion transport.

As used herein, the term "CFTR potentiator" refers to a compound that increases the channel activity of CFTR protein located at the cell surface, resulting in enhanced ion transport.

As used herein, the term "inducing," as in inducing CFTR activity, refers to increasing CFTR activity, whether by the corrector, potentiator, or other mechanism.

As used herein, the term "active pharmaceutical ingredient" or "API" refers to a biologically active compound.

A "patient," "subject" or "individual" are used interchangeably and refer to either a human or non-human animal. The term includes mammals such as humans.

The terms "effective dose" or "effective amount" are used interchangeably herein and refer to that amount that produces the desired effect for which it is administered (e.g., improvement in CF or a symptom of CF or lessening the severity of CF or a symptom of CF). The exact amount will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques (see, e.g., Lloyd (1999) The Art, Science and Technology of Pharmaceutical Compounding).

As used herein, the terms "treatment," "treating," and the like generally mean the improvement of CF or its symptoms or lessening the severity of CF or its symptoms in a subject. "Treatment," as used herein, includes, but is not limited to, the following: increased growth of the subject, increased weight gain, reduction of mucus in the lungs, improved pancreatic and/or liver function, reduced cases of chest infections, and/or reduced instances of coughing or shortness of breath. Improvements in or lessening the severity of any of these conditions can be readily assessed according to standard methods and techniques known in the art.

As used herein, the term "in combination with" when referring to two or more compounds or agents means that the order of administration includes the compounds or agents being administered prior to, concurrent with, or subsequent to each other to the patient.

As used herein, the phrase "optionally substituted" is used interchangeably with the phrase "substituted or unsubstituted."

As described herein, compounds within the compositions of the invention can optionally be substituted with one or more substituents, such as are illustrated generally above, or as exemplified by particular classes, subclasses, and species of the invention (such as the

compounds listed in Table 1). As described herein in formulae I-Id-ii , the variables R_1-R_4 in formulae I-Id-ii encompass specific groups, such as, for example, alkyl, alkenyl, alkynyl, alkoxy, heteroaryl, heterocyclic, cycloalkyl, and aryl, etc. Unless otherwise noted, each of the specific groups for the variables R_1-R_4 can be optionally substituted with one or more group selected from halo, phospho, OH, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, fluoroalkyl, alkyl, alkenyl, alkynyl, nitro, CN, hydroxyl, and (C1-C9alkylene)-E wherein up to 4 CH_2 units are independently replaced with O, S, SO_2, SO, CO, NH, *N*-alkyl, *N*-alkenyl, or *N*-alkynyl, and E is H, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, alkoxy,CN, or CF_3, further wherein each of the aryl, cycloalkyl, heterocycloalkyl, and heteroaryl is optionally substituted with one or more group selected from halo, alkyl, amino, CN, alkenyl, alkynyl, and alkoxy.

As one of ordinary skill in the art will recognize, combinations of substituents envisioned by this invention are those combinations that result in the formation of stable or chemically feasible compounds. The term "stable", as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and preferably their recovery, purification, and use for one or more of the purposes disclosed herein. In some embodiments, a stable compound or chemically feasible compound is one that is not substantially altered when kept at a temperature of 40°C or less, in the absence of moisture or other chemically reactive conditions, for at least a week. When two alkoxy groups are bound to the same atom or adjacent atoms, the two alkoxy groups can form a ring together with the atom(s) to which they are bound.

As disclosed herein, substituents or variables (such as R and R_1-R_4 in formula I-Id-ii) can be selected from more than one specific group. To the extent that one specific group for a variable may include or overlap with another specific group for the same variable, the narrower specific group is provisoed out from the broader specific group. In other words, double inclusion cannot exist.

In general, the term "substituted," whether preceded by the term "optionally" or not, refers to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. Specific substituents are described above in the definitions and below in the description of compounds and examples thereof. Unless otherwise indicated, an optionally substituted group can have a substituent at each substitutable position of the group, and when

more than one position in any given structure can be substituted with more than one substituent selected from a specified group, the substituent can be either the same or different at every position. As one of ordinary skill in the art will recognize, combinations of substituents envisioned by this invention are those combinations that result in the formation of stable or chemically feasible compounds.

The phrase "up to", as used herein, refers to zero or any integer number that is equal or less than the number following the phrase. For example, "up to 3" means any one of 0, 1, 2, and 3.

The terms "about" and "approximately", when used in connection with doses, amounts, or weight percent of ingredients of a composition or a dosage form, mean a dose, amount, or weight percent that is recognized by one of ordinary skill in the art to provide a pharmacological effect equivalent to that obtained from the specified dose, amount, or weight percent. Specifically the term "about" or "approximately" means an acceptable error for a particular value as determined by one of ordinary skill in the art, which depends in part on how the value is measured or determined. In certain embodiments, the term "about" or "approximately" means within 1, 2, 3, or 4 standard deviations. In certain embodiments, the term "about" or "approximately" means within 30%, 25%, 20%, 15%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.1%, or 0.05% of a given value or range.

The term "adjacent" as used herein refers to positions on the ring wherein the two ring atoms are bonded to each other. Two ring atoms with an intervening ring atom are not considered adjacent even when that intervening atom does not allow substitution due to valency.

The term "aliphatic", "aliphatic group," as used herein, means a straight-chain (i.e., unbranched) or branched hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation. Unless otherwise specified, aliphatic groups contain 1-20 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-10 aliphatic carbon atoms. In other embodiments, aliphatic groups contain 1-8 aliphatic carbon atoms. In still other embodiments, aliphatic groups contain 1-6 aliphatic carbon atoms, and in yet other embodiments aliphatic groups contain 1-4 aliphatic carbon atoms. Suitable aliphatic groups include, but are not limited to, linear or branched, substituted or unsubstituted alkyl, alkenyl, alkynyl groups.

The term "spiro" as used herein, means a two-ring system wherein both rings share

only one common atom.

The term "cycloaliphatic" or "cycloalkyl" mean a monocyclic, bicyclic (fused or spiro), tricyclic (fused or spiro), or propellane hydrocarbon that has a single point of attachment to the rest of the molecule, and that is completely saturated or contains one or more units of unsaturation, but none of the individual rings in the monocyclic, bicyclic, or tricyclic hydrocarbon is aromatic. The single point of attachment can be on the saturated or unsaturated carbon. In some embodiments, "cycloaliphatic" or "cycloalkyl" refers to a monocyclic C_3-C_8 hydrocarbon or bicyclic C_8-C_{12} hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but none of the individual ring in the monocyclic C_3-C_8 hydrocarbon or fused bicyclic C_8-C_{12} hydrocarbon is aromatic, and that has a single point of attachment to the rest of the molecule wherein any individual ring in said bicyclic ring system has 3-7 members.

As used herein, an "alkyl" group refers to a saturated hydrocarbon group containing 1-20 (e.g., 1-6 or 1-12) carbon atoms. An alkyl group can be straight or branched. Examples of alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, n-heptyl, or 2-ethylhexyl.

As used herein, an "alkenyl" group refers to a hydrocarbon group that contains 2-20 (e.g., 2-12, 2-6, or 2-4) carbon atoms and at least one double bond. Like an alkyl group, an alkenyl group can be straight or branched. The point of attachement can be on a saturated carbon or unsaturated carbon. Examples of an alkenyl group include, but are not limited to allyl, isoprenyl, 2-butenyl, and 2-hexenyl.

As used herein, an "alkynyl" group refers to a hydrocarbon group that contains 2-20 (e.g., 2-12, 2-6, or 2-4) carbon atoms and has at least one triple bond. An alkynyl group can be straight or branched. The point of attachement can be on a saturated carbon or unsaturated carbon. Examples of an alkynyl group include, but are not limited to, propargyl and butynyl.

As used herein, an "alkoxy" group refers to -O-alkyl, O-alkenyl, or O-alknyl, wherein alkyl, alkenyl, and alkynyl are as defined above.

As used herein, "fluoroalkyl" or "fluoroalkoxy" refers to alkyl or alkoxy wherein one or more hydrogen is substituted with a fluoro.

As used herein, an "amino" refers to NH_2 which is optionally substituted with one or

two groups independently selected from alkyl, cycloalkyl, and heterocycloalkyl.

The term "electron withdrawing group", as used herein means an atom or a group that is electronegative relative to hydrogen. See, e.g., "Advanced Organic Chemistry: Reactions, Mechanisms, and Structure," Jerry March, 4th Ed., John Wiley & Sons (1992), e.g., pp. 14-16, 18-19, etc. Exemplary such substituents include halo such as Cl, Br, or F, CN, COOH, CF_3, etc.

Unless otherwise specified, the term "heterocycle", "heterocyclyl", "heterocycloaliphatic", "heterocycloalkyl" or "heterocyclic" as used herein means monocyclic, bicyclic (fused or spiro), tricyclic (fused or spiro), or propellane ring systems in which one or more ring atoms in one or more ring members is an independently selected heteroatom and none of the individual rings in the system is aromatic. Heterocyclic rings can be saturated or can contain one or more unsaturated bonds. In some embodiments, the "heterocycle", "heterocyclyl", "heterocycloaliphatic", "heterocycloalkyl"or "heterocyclic" group has three to fourteen ring members in which one or more ring members is a heteroatom independently selected from oxygen, sulfur, nitrogen, or phosphorus, and each ring in the ring system contains 3 to 7 ring members. The point of attachment can be on the carbon or heteroatom.

The term "heteroatom" means oxygen, sulfur, nitrogen, phosphorus, or silicon (including, any oxidized form of nitrogen, sulfur, phosphorus, or silicon; the quaternized form of any basic nitrogen or; a substitutable nitrogen of a heterocyclic ring, for example N (as in 3,4-dihydro-2*H*-pyrrolyl), NH (as in pyrrolidinyl) or NR^+ (as in *N*-substituted pyrrolidinyl)).

The term "unsaturated", as used herein, means that a moiety has one or more units of unsaturation but is not aromatic.

The term "alkoxy", or "thioalkyl", as used herein, refers to an alkyl group, as previously defined, attached to the principal carbon chain through an oxygen ("alkoxy") or sulfur ("thioalkyl") atom.

The term "aryl" used alone or as part of a larger moiety as in "aralkyl", "aralkoxy", or "aryloxyalkyl", refers to monocyclic, bicyclic (fused), and tricyclic (fused or spiro) hydrocarbon ring systems having a total of five to fourteen ring carbon atoms, wherein at least one ring in the system is aromatic and wherein each ring in the system contains 3 to 7 ring carbon atoms. The term "aryl" may be used interchangeably with the term "aryl ring".

The term "heteroaryl", used alone or as part of a larger moiety as in "heteroaralkyl" or "heteroarylalkoxy", refers to monocyclic, bicyclic (fused), and tricyclic (fused or spiro) ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic, at least one ring in the system contains one or more heteroatoms, and wherein each ring in the system contains 3 to 7 ring members. The term "heteroaryl" may be used interchangeably with the term "heteroaryl ring" or the term "heteroaromatic".


The term "alkylene" refers to a straight or branched hydrocarbon chain that may be fully saturated or have one or more units of unsaturation and has two points of attachment to the rest of the molecule.

The term "prodrug," as used herein, represents a compound that is transformed *in vivo* into a compound according to any one of the formulae listed herein. Such a transformation can be affected, for example, by hydrolysis in blood or enzymatic transformation of the prodrug form to the parent form in blood or tissue. Prodrugs of the compounds of the invention may be, for example, amides. Amides that may be utilized as prodrugs in the present invention are phenyl amides, aliphatic (C_1-C_{24}) amids, acyloxymethyl amides, ureas, carbamates, and amino acid amides. For example, a compound of the invention that contains an NH group may be acylated at this position in its prodrug form. Other prodrug forms include esters, such as, for example phenyl esters, aliphatic (C_1-C_{24}) esters, acyloxymethyl esters, carbonates, carbamates, and amino acid esters. A thorough discussion of prodrugs is provided in T. Higuchi and V. Stella, *Pro-drugs as Novel Delivery Systems*, Vol. 14 of the A.C.S. Symposium Series, Edward B. Roche, ed., *Bioreversible Carriers in Drug Design*, American Pharmaceutical Association and Pergamon Press, 1987, and Judkins *et al., Synthetic Communications* 26(23):4351-4367, 1996, each of which is incorporated in its entirety herein by reference. In some embodiments, the present invention features a prodrug of any one of the formulas or compounds listed herein.

The term "isosteres" or "bioisosteres," as used herein, refers to compounds resulting from the exchange of an atom or group of atoms to create a new compound with similar biological properties to the parent compound. The bioisosteric replacement may be physicochemically or topologically based. For example, an isosteric replacement for a carboxylic acid is $CONHSO_2$(alkyl or aryl)) such as $CONHSO_2Me$. A further discussion of isosterism is provided in R. Silverman, *The Organic Chemistry of Drug Design and Drug Action*,

second edition, Elsevier Academic Press, 2004, incorporated in its entirety herein by reference. In some embodiments, the present invention features a isostere of any one of the formulas or compounds listed herein.

In the formulas and drawings, a line transversing a ring and bonded to an R group such as, for example, R_2 in the following formula



means that the R group can be bonded to any carbon, or if applicable, heteroatom such as N, of that ring, including any fused ring, as valency allows.

Within a term definition as, for example, R_1 through R_3 when a CH_2 unit or, interchangeably, a methylene unit may be replaced by O, CO, S, SO, SO_2, or NR; it is meant to include any CH_2 unit, including a CH_2 within a terminal methyl or methylene group. For example, $-CH_2CH_2CH_2SH$ is within the definition of C1-C9 alkylene-R_4 wherein up to four CH_2 units are independently replaced by O, CO, S, SO, SO_2, or NR because the CH_2 unit of the terminal methyl group has been replaced by S. The analogous applies to such definitions as $-CH_2CH_2OH$, $-CH_2CH_2CN$, or $-CH_2CH_2NH_2$.

Unless otherwise stated, structures depicted herein are also meant to include all isomeric (e.g., enantiomeric, diastereomeric, and geometric (or conformational)) forms of the structure; for example, the R and S configurations for each asymmetric center, (Z) and (E) double bond isomers, and (Z) and (E) conformational isomers. Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, and geometric (or conformational) mixtures of the present compounds are within the scope of the invention. Where the enantiomers of a racemic mixture have been separated, but the absolute chemistry has not yet been determined, the compound's structure is depicted with a wavy line.

Unless otherwise stated, all tautomeric forms of the compounds of the invention are within the scope of the invention. Thus, included within the scope of the invention are tautomers of compounds of formulas I to Id-ii.

To the extent that a definition in the present application differs from any definition in an application incorporated by reference, the definition in the present application supercedes.

CFTR is a cAMP/ATP-mediated anion channel that is expressed in a variety of cells types, including absorptive and secretory epithelia cells, where it regulates anion flux across the membrane, as well as the activity of other ion channels and proteins. In epithelia cells, normal functioning of CFTR is critical for the maintenance of electrolyte transport throughout the body, including respiratory and digestive tissue. CFTR is composed of approximately 1480 amino acids that encode a protein made up of a tandem repeat of transmembrane domains, each containing six transmembrane helices and a nucleotide binding domain. The two transmembrane domains are linked by a large, polar, regulatory (R)-domain with multiple phosphorylation sites that regulate channel activity and cellular trafficking.

In patients with cystic fibrosis, mutations in endogenously expressed CFTR lead to reduced apical anion secretion causing an imbalance in ion and fluid transport. The resulting decrease in anion transport contributes to enhanced mucus accumulation in the lung and the accompanying microbial infections that ultimately cause death in CF patients. In addition to respiratory disease, CF patients typically suffer from gastrointestinal problems and pancreatic insufficiency that, if left untreated, results in death. In addition, the majority of males with cystic fibrosis are infertile and fertility is decreased among females with cystic fibrosis. In contrast to the severe effects of two copies of the CF associated gene, individuals with a single copy of the CF associated gene exhibit increased resistance to cholera and to dehydration resulting from diarrhea-perhaps explaining the relatively high frequency of the CF gene within the population.


Sequence analysis of the *CFTR* gene of CF chromosomes has revealed a variety of disease-causing mutations (Cutting, G. R. et al. (1990) Nature 346:366-369; Dean, M. et al. (1990) Cell 61:863:870; and Kerem, B-S. et al. (1989) Science 245:1073-1080; Kerem, B-S et al. (1990) Proc. Natl. Acad. Sci. USA 87:8447-8451). To date, greater than 1000 disease-causing mutations in the CF gene have been identified as reported by the scientific and medical literature. The most prevalent mutation is a deletion of phenylalanine at position 508 of the CFTR amino

acid sequence, and is commonly referred to as F508del. This mutation occurs in approximately 70 percent of the cases of cystic fibrosis and is associated with a severe disease. A more complete list of identified mutations can be found at www.cftr2.org.

These elements work together to achieve directional transport across the epithelium via their selective expression and localization within the cell. Chloride absorption takes place by the coordinated activity of ENaC and CFTR present on the apical membrane and the Na^+-K^+-ATPase pump and Cl- channels expressed on the basolateral surface of the cell. Secondary active transport of chloride from the luminal side leads to the accumulation of intracellular chloride, which can then passively leave the cell via Cl^- channels, resulting in a vectorial transport. Arrangement of $Na^+/2Cl^-/K^+$ co-transporter, Na^+-K^+-ATPase pump and the basolateral membrane K^+ channels on the basolateral surface and CFTR on the luminal side coordinate the secretion of chloride via CFTR on the luminal side. Because water is probably never actively transported itself, its flow across epithelia depends on tiny transepithelial osmotic gradients generated by the bulk flow of sodium and chloride.

Compounds of Formula I

In some embodiments the present invention features a compound of formula I:



I,

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring A is a C6-C10 aryl ring; C3-C10 cycloalkyl ring; or a C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring B is a C3-C10 cycloalkyl ring; a C6-C10 aryl ring; or a C4-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C6-C10 aryl ring ; a C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, NR, O, or S; or a C3-C10 cycloalkyl ring;

W is O, NR, or S;

X is O or NR;

Y is independently CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

Z is NR or CHR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy;C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; OH; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O,

S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl ; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;

o is 0, 1, 2, 3, 4, or 5;

p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5;

---- is a single bond or a double bond;



provided that the moieties containing ring B and ring C are substituted at adjacent positions on ring A.


In another embodiment, the compounds of the invention are of formula I and the attendant definitions, provided that: i) when ring A is indole, n is not zero and rings B and C are aryl unsubstituted by CF_3 or halo; ii) when ring A is pyrazole, n is not zero; iii) when ring A is pyridyl, ring B and ring C moieties are substituted at the 2- and 3- positions of the pyridyl ring, interchangeably; and iv) when ring A is imidazole, the ring B moiety is substituted at the 2-position of the imidazole ring.

In some embodiments, the compounds of the present inventions are in the form of a pharmaceutically acceptable prodrug.

In some embodiments, ring A is a C6-C10 aryl ring, such as a phenyl, indane, 1,2,3,4-tetrahydronaphthalene, or naphthalene. In some emboidments, ring A is a C3-C11 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, such as pyridyl, indole, indoline, isoindoline, pyrazole, pyrimidine, 1,2,3,4-tetrahydroquinoline,


quinoline, 5,6,7,8-tetrahydroquinoline, 1,2,3,4-tetrahydroisoquinoline, pyrrolodine, aza-indole, pyrrole, oxazole, pyrazine, triazole, indazole, 2,3,4,5-tetrahydro-1H-benzo[d]azepine, 1H-benzo[d]imidazole, or imidazole ring.


In some embodiments, ring A is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring A is  ,  , ,


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  , or .

In some embodiments, ring C is a C6-C10 aryl ring, such as a phenyl, indane, or naphthalene. In some embodiments, ring C is a C3-C10 cycloalkyl ring, such as a cyclopropane, cyclobutane, cyclopentane, cyclopentene, cyclohexane, cyclohexene, cycloheptane, or (1s,4s)-bicyclo[2.2.1]heptane. In some embodiments, ring C is a C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are indpependently N, NR, O, or S, such as indole, isoindoline, indoline, pyridyl, pyrrolidine, 2,5-dihydro-1H-pyrrole, morpholine, tetrahydro-2H-pyran, tetrahydrofuran, oxetane, piperidine, piperazine, azepane, azetidine, octahydrocyclopenta[c]pyrrole, octahydrocyclopenta[b]pyrrole, octahydro-1H-indole, octahydro-1H-cyclopenta[b]pyridine, (1s,4s)-7-azabicyclo[2.2.1]heptane, 2-azabicyclo[4.1.0]heptane,

(1S,4R)-2-azabicyclo[2.2.1]heptane, (1R,5S)-8-azabicyclo[3.2.1]octane, 3,6-dihydro-2H-pyran, or (1R,5S)-3-azabicyclo[3.2.1]octane.

[0073] In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, Ring C is 















In some embodiments, X is O. In some embodiments, X is NR. In some embodiments, X is NH.


In In some embodiments, Y is O. In some embodiments, Y is CRR, such as CH_2, CH(C1-C6 alkyl), $CH(CH_3)$, or$CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6 alkynyl), or N(C1-C6 alkyl). In some embodiments, $(Y)_o$ is a bond, O, NH, $(CH_2)_m$, O-$(CH_2)_m$, where m=1-4, $CH(CH_3)$, O-$CH(CH_3)$, O-$CH(CH_2CH_3)$, O-$CH(CH_3)CH_2$, O-$C(CH_3)_2CH_2$, O-$CH_2CH_2CH(CH_3)$, $N(CH_3)$, NH-CH_2, $N(CH_3)$-CH_2, $NCH(CH_3)_2$-CH_2, $N(CH_2CH_2CH_3)$-CH_2, NH-$CH(CH_3)$, $N(CH_3)$-$CH(CH_3)$, or $N(CH_2CCH)$-CH_2CH_2.


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, azetidine, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu,

$CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$, 


















In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 carbon atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are optionally and independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,








, or CO_2H.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,












, or .



In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.




In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.





In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.



In some embodiments, ring A is  , and n is 1. In some embodiments, ring A is  , n is 1, and R_1 is tBu. In some embodiments, ring A is  , n is 1, and R_1 is phenyl. In some embodiments, ring A is  , n is 1, R_1 is phenyl, and Y is O. In some embodiments, ring A is  , n is 1, R_1 is tBu, and Y is O. In some emdodiments, ring A is  , ring B is pyridyl, and ring C is phenyl. In some embodiments, R_1 is phenyl, R_2 is amino, and R_3 is CH_3.







In some embodiments, ring A is  , n is 1, and R_1 is pyridyl. In some embodiments, ring A is  , n is 1, R_1 is pyridyl, and o is 0. In some embodiment, ring A is  , ring B is pyridyl, and ring C is pyrrolidine.





In some embodiments, ring A is  , n is 1, and R_1 is pyrazole. In some embodiments, ring A is  , n is 1, R_1 is pyrazole, and o is 0.



In some embodiments, ring A is  , n is 1, and R_1 is cycloalkyl. In some embodiments, ring A is  , n is 1, R_1 is cycloalkyl, and Y is O. In some emdodiments, ring A is  , ring B is pyridyl, and ring C is cycloalkyl. In some embodiments, R_1 is cycloalkyl, R_2 is alkoxy, and R_3 is CH_3.

In some embodiments, ring A is  , and n is 1. In some embodiments, ring A is  , n is 1, and R_1 is phenyl. In some embodiments, ring A is  , n is 1, and R_1 is tBu. In some embodiments, ring A is  , n is 1, R_1 is phenyl, and Y is O. In some


embodiments, ring A is  , n is 1, R_1 is tBu, and Y is O. In one emdodiment, ring A is  , ring B is pyridyl, and ring C is phenyl.

In some embodiments, ring A is  , and n is 1. In some embodiments, ring A is  , n is 1, and R_1 is phenyl. In some embodiments, ring A is  , n is 1, and R_1 is tBu. In some embodiments, ring A is  , n is 1, R_1 is phenyl, and Y is O. In some embodiments, ring A is  , n is 1, R_1 is tBu, and Y is O. In some embodiments, ring A is  , ring B is pyridyl, and ring C is phenyl.

In some embodiments, ring A is  , and n is 1. In some embodiments, ring A is  , n is 1, and R_1 is phenyl. In some embodiments, ring A is  , n is 1, and R_1 is tBu. In some embodiments, ring A is  , n is 1, R_1

is phenyl, and Y is O. In some embodiments, ring A is  , n is 1, R_1 is tBu, and Y is O. In one emdodiment, ring A is  , ring B is pyridyl, and ring C is phenyl.

In some embodiments, the present invention features a compound of formula I-i:



I-i,

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring A is a C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring B is a C3-C10 heteroaryl ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C6-C10 aryl ring , C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are optionally N, O, or S, or a C3-C10 cycloalkyl ring;

X is O or NR;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

Z is NR or CHR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-

C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy;C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl ; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;

o is 0, 1, 2, 3, 4, or 5;



p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5;



provided that the moieties containing ring B and ring C are substituted at adjacent positions on ring A.


In some embodiments, the compound of formula I-i exists as a pharmaceutically acceptable prodrug.

In some embodiments, ring A is a pyridyl, indole, indoline, isoindoline, pyrazole, pyrimidine, quinoline, 5,6,7,8-tetrahydroquinoline, 1,2,3,4-tetrahydroisoquinoline, pyrrolodine, aza-indole, pyrrole, oxazole, pyrazine, triazole, indazole, or imidazole ring.

In some embodiments, ring A is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring A is 



In some embodiments, ring B is pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, thiazole, pyrrolidinone, or quinoline.


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is 




In some embodiments, ring C is phenyl, indole, cycloalkyl, pyridyl, tetrahydro-2H-pyran, azepane, tetrahydrofuran, pyrrolidine, naphthalene, piperidine, azetidine, or dihydroindene.

In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is






In some embodiments, X is O. In some embodiments, X is NR. In some embodiments, X is NH.


In some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6alkynyl), or N(C1-C6 alkyl).


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, azetidine, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S,


SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,












In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are optionally O, S, N, or NR; and a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,








In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, C6-C10 aryl, or C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,




In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, the compound of formula I is a compound of formula I-ii:



I-ii,

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring A is a C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring B is a C6-C10 aryl ring;

Ring C is a C6-C10 aryl ring , C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are optionally N, O, or S, or a C3-C10 cycloalkyl ring;

X is O or NR;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

Z is NR or CHR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy;C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O,

S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;

o is 0, 1, 2, 3, 4, or 5;



p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5;


provided that the moieties containing ring B and ring C are substituted at adjacent positions on ring A.

In some embodiments, the compound of formula I-ii exists as a pharmaceutically acceptable prodrug.

In some embodiments, ring A is a pyridyl, indole, indoline, isoindoline, pyrazole, pyrimidine, quinoline, 5,6,7,8-tetrahydroquinoline, 1,2,3,4-tetrahydroisoquinoline, pyrrolodine, aza-indole, pyrrole, oxazole, pyrazine, triazole, indazole, or imidazole ring.

In some embodiments, ring A is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring A is  ,  , ,







In some embodiments, X is O. In another embodiment, X is NR. In some embodiments, X is NH.


In some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6 alkynyl), or N(C1-C6 alkyl).


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, azetidine, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,










In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.


In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O,


CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,



, or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, C6-C10 aryl, or C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,  ,




In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, the invention features a compound of formula Ia:



Ia

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C6-C10 aryl ring , C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 cycloalkyl ring;

X is O or NR;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy;C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl ; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH;CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, or 2;






o is 0, 1, 2, 3, 4, or 5;

p is 0, 1, 2, or 3; and



q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ia exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.

In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  ,  ,  ,  ,  , or .

In some embodiments, ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, piperidine, or dihydroindene.


In some embodiments, ring C is is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is  ,  ,








In some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6alkynyl), or N(C1-C6 alkyl).

In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$, 








In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 ring

atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.


In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,



In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3,

CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,




In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, ring B is phenyl. In some embodiments, ring B and ring C are phenyl. In some embodiments, R_1 is tBu. In some embodiments, ring B and ring C are phenyl and R_1 is tBu. In some embodiments, Y is CH_2. In some embodiments, ring B and ring C are phenyl, R_1 is tBu, and Y is CH_2. In some embodiments, ring B, ring C, and R_1 are phenyl and Y is CH_2.

In some embodiments, the invention features a compound of formula Ia-i:



Ia-i

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C3-C10 heteroaryl ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C6-C10 aryl ring, C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 cycloalkyl ring;

X is O or NR;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy;C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl ; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH;CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, or 2;

o is 0, 1, 2, 3, 4, or 5;








p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.

In some embodiments, the compound of formula Ia-i exists as a pharmaceutically acceptable prodrug.

In some embodiments, ring B is pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is ,  ,  ,


 ,  ,  ,  ,  ,  ,  ,




In some embodiments, ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, piperidine, or dihydroindene.

In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is










In some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6alkynyl), or N(C1-C6 alkyl).


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example,R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu,












In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,








In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,




In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, the invention features a compound of formula Ia-ii:



Ia-ii

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring;

Ring C is a C6-C10 aryl ring , C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 cycloalkyl ring;

X is O or NR;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy;C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring

wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, or 2;

o is 0, 1, 2, 3, 4, or 5;


p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ia-ii exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is phenyl or naphthalene.

In some embodiments, ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, piperidine, or dihydroindene.

In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is 










In some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6alkynyl), or N(C1-C6 alkyl).


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,










In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O,


CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,




, or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,






In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, the compound of formula I is a compound of formula Ib:



Ib

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C6-C10 aryl ring, C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 mono- or bicyclic cycloalkyl ring;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH;CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;



o is 0, 1, 2, 3, 4, or 5;

p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ib exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.

In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  , 

In some embodiments, ring C is phenyl, indole, cycloalkyl, pyridyl, naphthalene, pyrrolidine, or dihydroindene.

In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is  ,








In some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6alkynyl), or N(C1-C6 alkyl).


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.


In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,














In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O,

CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  , or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O, 



In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, ring B is pyridyl. In some embodiments, ring B is phenyl. In some embodiments, ring B and ring C are phenyl. In some embodiments, ring B is pyridyl and ring C is phenyl. In some embodiments, Y is CH_2. In some embodiments, Y is O.


In some embodiments, R_1 is aryl. In some embodiments, ring B, ring C, and R_1 are phenyl. In some embodiments, ring B is pyridyl, ring C is phenyl, and R_1 is phenyl. In some embodiments, ring B, ring C, and R_1 are phenyl, and Y is CH_2. In some embodiments, ring B, ring C, and R_1 are phenyl, and Y is O. In some embodiments, ring B is pyridyl, ring C is phenyl, R_1 is phenyl, and Y is CH_2. In some embodiments, ring B is pyridyl, ring C is phenyl, R_1 is phenyl, and Y is O.

In some embodiments, ring C is heterocyclic. In some embodiments, ring C is pyrrolodine. In some embodiments, ring C is piperidine. In some embodiments, ring B is pyridyl , ring C is heterocyclic. In some embodiments, ring B is pyridyl, ring C is heterocycloalkyl, and R_1 is phenyl. In some embodiments, ring B is phenyl, ring C is heterocycloalkyl, and R_1 is phenyl.

In some embodiments, R_1 is heteroaryl. In some embodiments, ring C is pyrrolidine and R_1 is heteroaryl. In some embodiments, ring C is pyrrolidine and R_1 is pyrrozole. In some embodiments, ring B is pyridyl, ring C is pyrrolidine, and R_1 is heteroaryl. In some embodiments, ring B is pyridyl, ring C is heterocycloalkyl, and R_1 is pyridyl..

In some embodiments, R_1 is cycloalkyl. In some embodiments, ring B is pyridyl and R_1 is cycloalkyl.

In some embodiments, the invention features a compound of formula Ib-i:



Ib-i

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C3-C10 heteroaryl ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C6-C10 aryl ring , C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 cycloalkyl ring;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or

heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl ; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;


o is 0, 1, 2, 3, 4, or 5;


p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5.

In some embodiments, the compound of formula Ib-i exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  ,



In some embodiments, ring C is phenyl, indole, cycloalkyl, pyridyl, naphthalene, pyrrolidine, or dihydroindene.

In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is








In some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6alkynyl), or N(C1-C6 alkyl).


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,












In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,








In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,




In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, the invention features a compound of formula Ib-ii:



Ib-ii

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring;

Ring C is a C6-C10 aryl ring, C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 cycloalkyl ring;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl ; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;

o is 0, 1, 2, 3, 4, or 5;


p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ib-ii exists as a pharmaceutically acceptable prodrug.

In some embodiments, ring B is phenyl or napthalene.

In some embodiments, ring C is phenyl, indole, cycloalkyl, pyridyl, naphthalene, pyrrolidine, or dihydroindene.

In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is 








In some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6alkynyl), or N(C1-C6 alkyl).


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,










In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O,


CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,




, or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,






In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, the invention features a compound of formula Ib-iii:



Ib-iii

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, and wherein one nitrogen on Ring C is the point of attachment to the pyridine ring;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;


n is 0, 1, 2 or 3;

p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ib-iii exists as a pharmaceutically acceptable prodrug.

In some embodiments, ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is 


In some embodiments, ring C is indole, piperidine, azepane, azetadine, indoline, isoindoline, or pyrrolidine.


In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is 




In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$, 










In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu,




In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, R_1 is phenyl, pyridine, or pyrazole, and n is 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, n is 1, R_2 is amino or alkyl, and p is 0 or 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, n is 1, R_3 is alkyl, such as methyl, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-iii-1:



Ib-iii-1

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C3-C10 heteroaryl ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, and wherein one nitrogen on Ring C is the point of attachment to the pyridine ring;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy;C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH;CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;


o is 0, 1, 2, 3, 4, or 5;

p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ib-iii-1 exists as a pharmaceutically acceptable prodrug.

In some embodiments, ring B is pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.

In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is 


In some embodiments, ring C is indole, piperidine, azepane, azetadine, indoline, isoindoline, or pyrrolidine.


In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is 





In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,






































In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$,

=O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  , and CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,  ,  ,  ,  ,  .




In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, R_1 is phenyl, pyridine, or pyrazole, and n is 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, n is 1, R_2 is amino or alkyl, and p is 0 or 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, n is 1, R_3 is alkyl, such as methyl, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-iii-2:



Ib-iii-2

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring;

Ring C is a C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, and wherein one nitrogen on Ring C is the point of attachment to the pyridine ring;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl ; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;

o is 0, 1, 2, 3, 4, or 5;





p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5.

In some embodiments, the compound of formula Ib-iii-2 exists as a pharmaceutically acceptable prodrug.

In some embodiments, ring B is phenyl or naphthalene.


In some embodiments, ring C is indole, piperidine, azepane, azetadine, indoline, isoindoline, or pyrrolidine.


In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is  ,  ,  ,  , .


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected


from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,













In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,










In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,




In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, R_1 is phenyl, pyridine, or pyrazole, and n is 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, n is 1, R_2 is amino or alkyl, and p is 0 or 1. In

some embodiments, R_1 is phenyl, pyridine, or pyrazole, n is 1, R_3 is alkyl, such as methyl, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-iv:



Ib-iv

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, and wherein one nitrogen on Ring C is the point of attachment to the pyridine ring;

R_1 is C6-C10 aryl or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy;C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-

C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;



R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;


p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ib-iv exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  , 



In some embodiments, ring C is indole, piperidine, azepane, azetadine, indoline, isoindoline, or pyrrolidine.


In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is 








In some embodiments, R_1 is a phenyl, thiophene, pyridine, or pyrazole.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is












In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,








In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.


In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,



In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, R1 is phenyl, pyridine, or pyrazole. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, R_2 is amino or alkyl, and p is 0 or 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, R_3 is alkyl, such as methyl, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-iv-1:



Ib-iv-1

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C3-C10 heteroaryl ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, and wherein one nitrogen on Ring C is the point of attachment to the pyridine ring;

R_1 is C6-C10 aryl or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;





R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

p is 0, 1, 2, or 3; and




q is 0, 1, 2, 3, 4, or 5.

In some embodiments, ring B is pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.



In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  ,  ,  ,  , or .


In some embodiments, the compound of formula Ib-iv-1 exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring C is indole, piperidine, azepane, azetadine, indoline, isoindoline, or pyrrolidine.


In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is  ,  , or  .

In some embodiments, R_1 is a phenyl, thiophene, pyridine, or pyrazole.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is  ,  .








In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH,


$(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,




, or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,






In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, R_1 is phenyl, pyridine, or pyrazole. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, R_2 is amino or alkyl, and p is 0 or 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, R_3 is alkyl, such as methyl, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-iv-2:



Ib-iv-2

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring;

Ring C is a C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, and wherein one nitrogen on Ring C is the point of attachment to the pyridine ring;

R_1 is C6-C10 aryl or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy;C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;


p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.

In some embodiments, the compound of formula Ib-iv-2 exists as a pharmaceutically acceptable prodrug.

In some embodiments, ring B is phenyl or napthalene.


In some embodiments, ring C is indole, piperidine, azepane, azetadine, indoline, isoindoline, or pyrrolidine.


In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is 





In some embodiments, R_1 is a phenyl, thiophene, pyridine, or pyrazole.


In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is
















In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.


In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$,


$N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$,

$NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  ,  ,  ,

 ,


 ,


 ,


 , or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu,

$=CH_2$, $=O$,  ,


 .



In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, R_1 is phenyl, pyridine, or pyrazole. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, R_2 is amino or alkyl, and p is 0 or 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, R_3 is alkyl, such as methyl, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-v:



Ib-v

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O,

S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;



n is 0, 1, 2 or 3;


p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5.

In some embodiments, the compound of formula Ib-v exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  , 





In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,










In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH,


$(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,




, or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,






In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, R_1 is phenyl, pyridine, or pyrazole, and n is 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, n is 1, R_2 is amino or alkyl, and p is 0 or 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, n is 1, R_3 is alkyl, such as methyl, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-v-1:



Ib-v-1

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C3-C10 heteroaryl ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-

C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;






















n is 0, 1, 2 or 3;

p is 0, 1, 2, or 3; and



















q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ib-v-1 exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  , or  .

In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  .








In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH,


$(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,




, or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, $=O$,






In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, R_1 is phenyl, pyridine, or pyrazole, and n is 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, n is 1, R_2 is amino or alkyl, and p is 0 or 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, n is 1, R_3 is alkyl, such as methyl, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-v-2:



Ib-v-2

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-

C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;

p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ib-v-2 exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is phenyl or napthalene.


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9


alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,












In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,








In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,




In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, R_1 is phenyl, pyridine, or pyrazole, and n is 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, n is 1, R_2 is amino or alkyl, and p is 0 or 1. In

some embodiments, R_1 is phenyl, pyridine, or pyrazole, n is 1, R_3 is alkyl, such as methyl, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-vi:



Ib-vi

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

R_1 is C6-C10 aryl or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;





R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ib-vi exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  ,  ,  ,  ,  , or .

In some embodiments, R_1 is a phenyl, thiophene, pyridine, or pyrazole.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is 










In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,








In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.


In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,



In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, R_1 is phenyl, pyridine, or pyrazole. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, R_2 is amino or alkyl, and p is 0 or 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, R_3 is alkyl, such as methyl, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-vi-1:



Ib-vi-1

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C3-C10 heteroaryl ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

R_1 is C6-C10 aryl or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;






















R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

p is 0, 1, 2, or 3; and










q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ib-vi-1 exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  , or  .

In some embodiments, R_1 is a phenyl, thiophene, pyridine, or pyrazole.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is  ,  ,  ,  ,  ,  ,  ,  ,  .








In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH,


$(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,




, or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,






In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, R_1 is phenyl, pyridine, or pyrazole. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, R_2 is amino or alkyl, and p is 0 or 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, R_3 is alkyl, such as methyl, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-vi-2:



Ib-vi-2

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring;

R_1 is C6-C10 aryl or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O,

S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;



p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ib-vi-2 exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is phenyl or napthalene.

In some embodiments, R_1 is a phenyl, thiophene, pyridine, or pyrazole.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is 









In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH,


$(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,




, or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, $=O$,






In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, R_1 is phenyl, pyridine, or pyrazole. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, R_2 is amino or alkyl, and p is 0 or 1. In some embodiments, R_1 is phenyl, pyridine, or pyrazole, R_3 is alkyl, such as methyl, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-vii:



Ib-vii

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O,

S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_5 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;




p is 0, 1, 2, or 3;
























q is 0, 1, 2, 3, 4, or 5; and

r is 0, 1, 2, 3, 4, or 5.












In some embodiments, the compound of formula Ib-vii exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.

In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  ,  ,  ,

$NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,    ,                      , or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.


In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,           



In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.


In some embodiments, R_5 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_5 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH_2C(CH_3)_3$ $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,



In some embodiments, r is 0. In some embodiments, r is 1. In some embodiments, r is 2. In some embodiments, r is 3. In some embodiments, r is 4. In some embodiments, r is 5.

In some embodiments, R_5 is halo or alkoxy. In some embodiments, R_2 is amino or alkyl, R_5 is halo or alkoxy, r is 1 or 2, and p is 0 or 1. In some embodiments, R_5 is halo or alkoxy, R_3 is alkyl, such as methyl, r is 1 or 2, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-viii:



Ib-viii

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_5 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;



p is 0, 1, 2, or 3;


q is 0, 1, 2, 3, 4, or 5; and

r is 0, 1, 2, 3, 4, or 5.

In some embodiments, the compound of formula Ib-viii exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  , 




In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,








In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,




In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, R_5 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_5 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH_2C(CH_3)_3$ $OCH(CH_3)_2$, CO_2H, CO_2NH_2,


OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu,

$=CH_2$, =O, 

In some embodiments, r is 0. In some embodiments, r is 1. In some embodiments, r is 2. In some embodiments, r is 3. In some embodiments, r is 4. In some embodiments, r is 5.

In some embodiments, R_5 is halo or alkoxy. In some embodiments, R_2 is amino or alkyl, R_5 is halo or alkoxy, r is 1 or 2, and p is 0 or 1. In some embodiments, R_5 is halo or alkoxy, R_3 is alkyl, such as methyl, r is 1 or 2, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ib-ix:



Ib-ix

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_5 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;


p is 0, 1, 2, or 3;

q is 0, 1, 2, 3, 4, or 5; and

r is 0, 1, 2, 3, 4, or 5.

In some embodiments, the compound of formula Ib-ix exists as a pharmaceutically acceptable prodrug.




In some embodiments, ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.







In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is 







In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein up to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.







In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$,




$N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$,

$NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  ,  ,  ,

 ,  ,  ,  ,  ,  ,






 ,  ,  ,  ,  ,  ,







5  ,  ,  ,  ,  ,  ,


 ,  ,  , or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

10 In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu,

$=CH_2$, $=O$,  ,  ,  ,  ,  ,


15  ,  ,  ,  ,  ,  ,  ,



In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.


In some embodiments, R_5 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_5 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH_2C(CH_3)_3$ $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,



In some embodiments, r is 0. In some embodiments, r is 1. In some embodiments, r is 2. In some embodiments, r is 3. In some embodiments, r is 4. In some embodiments, r is 5.

In some embodiments, R_5 is halo or alkoxy. In some embodiments, R_2 is amino or alkyl, R_5 is halo or alkoxy, r is 1 or 2, and p is 0 or 1. In some embodiments, R_5 is halo or alkoxy, R_3 is alkyl, such as methyl, r is 1 or 2, and q is 1, 2, 3, or 4.

In some embodiments, the invention features a compound of formula Ic:



Ic

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C6-C10 aryl ring, C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 cycloalkyl ring;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;


o is 0, 1, 2, 3, 4, or 5;


p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5.

In some embodiments, the compound of formula Ic exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is 



In some embodiments, ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, or dihydroindene.

In some embodiments, ring C a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is







In some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6alkynyl), or N(C1-C6 alkyl).


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-


C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,












In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example , R_2 isCl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,








In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,




In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, ring B is phenyl. In some embodiments, ring C is phenyl. In some embodiments, ring B and ring C are phenyl. In some embodiments, R_1 is halo. In some embodiments, R_1 is F. In some embodiments, R_1 is OCH_3. In some embodiments R_2 is amino. In some embodiments, R_3 is CH_3. In some embodiments, R_2 is amino and R_3 is CH_3. In some embodiments, Y is CH_3. In some embodiments, Y is O. In some embodiments, R_1 is F, R_2 is amino, R_3 is CH_3, and Y is CH_2. In some embodiments, Y is O. In some embodiments, R_1 is F, R_2 is amino, R_3 is CH_3, and Y is O.

In some embodiments, the invention features a compound of formula Ic-i:



Ic-i

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C3-C10 heteroaryl ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C6-C10 aryl ring, C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 cycloalkyl ring;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O,

S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH;CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;


o is 0, 1, 2, 3, 4, or 5;


p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ic-i exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is ,  ,



In some embodiments, ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, or dihydroindene.

In some embodiments, ring C is is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is 








in some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6alkynyl), or N(C1-C6 alkyl).


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,












In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,








In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,




In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In other embodiments, the invention features a compound of formula Ic-ii:



Ic-ii

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring;

Ring C is a C6-C10 aryl ring, C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 cycloalkyl ring;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;

o is 0, 1, 2, 3, 4, or 5;



p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Ic-ii exists as a pharmaceutically acceptable prodrug.

In some embodiments, ring B is phenyl or napthalene.

In some embodiments, ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, or dihydroindene.

In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is  , 








In some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6alkynyl), or N(C1-C6 alkyl).


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,












In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O,

CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  , or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O, 




In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, the invention features a compound of formula Id:



Id

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C6-C10 aryl ring, C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 cycloalkyl ring;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy;C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;



o is 0, 1, 2, 3, 4, or 5;

p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Id exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.

In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  , 

In some embodiments, ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, or dihydroindene.

In some embodiments, ring C is is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is  ,








In some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6alkynyl), or N(C1-C6 alkyl).


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,













In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O,

CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  ,  , , or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O, 




In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, ring B is phenyl. In some embodiments, ring B is pyridyl. In some embodiments, ring C is phenyl. In some embodiments, ring B and ring C are phenyl. In some embodiments, ring B is pyridyl and ring C is phenyl. In some embodiments, Y is CH_2. In some embodiments, Y is O. In some embodiments, R_2 is amino. In some embodiments, R_2 is OH. In some embodiments, R_2 is OCH_3. In some embodiments, R_3 is CH_3. In some embodiments, R_2 is amino and R_3 is CH_3. In some embodiments, ring B is pyridyl, ring C is phenyl, and Y is CH_2. In some embodiments, ring B is pyridyl, ring C is phenyl, and Y is O.

In some embodiments, the invention features a compound of formula Id-i:



Id-i

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C3-C10 heteroaryl ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C6-C10 aryl ring, C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 cycloalkyl ring;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;



o is 0, 1, 2, 3, 4, or 5;


p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5.

In some embodiments, the compound of formula Id-i exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, aza-indole, thiophene, dihydrobenzofuran, or quinoline.


In some embodiments, ring B is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring B is  , 



In some embodiments, ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, or dihydroindene.

In some embodiments, ring C is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is
















In some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6alkynyl), or N(C1-C6 alkyl).


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.

In some embodiments, R_1 is is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3,

tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$, 


















In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O, CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,








In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O,




In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In some embodiments, the invention features a compound of formula Id-ii:



Id-ii

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring;

Ring C is a C6-C10 aryl ring, C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are indepednently N, O, or S, or a C3-C10 cycloalkyl ring;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring

wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;

o is 0, 1, 2, 3, 4, or 5;


p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.


In some embodiments, the compound of formula Id-ii exists as a pharmaceutically acceptable prodrug.


In some embodiments, ring B is phenyl or napthalene.

In some embodiments, ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, or dihydroindene.

In some embodiments, ring C is is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, ring C is 










In some embodiments, Y is O. In some embodiments, Y is CH_2. In some embodiments, Y is CH(C1-C6 alkyl). In some embodiments, Y is $CH(CH_3)$. In some embodiments, Y is $CH(CH_2CH_3)$. In some embodiments, Y is NR, such as NH, N(C1-C6alkynyl), or N(C1-C6 alkyl).


In some embodiments, R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, pyrazole, or thiophene ring, or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, wherein all rings are optionally substituted with one or more groups selected from halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, and C3-C10 heterocycloalkyl.


In some embodiments, R_1 is is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_1 is CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, $OCH_2CH_2OCH_2CH_3$,

















In some embodiments, R_2 is halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C3-C10 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR.

In some embodiments, R_2 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, $N(CH_3)CH_2CH_2CH_3$, $N(CH_3)CH_2CH_2CH_2CH_3$, CH_3, CH_2OH, CH_2CH_3, $CH_2CH_2CH_3$, =O,

CH_3SO_2, CH_3SO_2NH, CF_3CONH, CH_3CONH, $CH_3CON(CH_3)$, tBuOCONH, $(CH_3)_2CHOCONH$, $CH(CH_3)_2$, CHF_2, OCH_3, OCH_2CH_3, $OCH_2CH_2CH_3$, $OCH_2CH_2CH(CH_3)_2$, OCF_3, $OCHF_2$, $OC(CH_3)_3$, OCH_2CH_2tBu, $NHCH(CH_3)(CH_2CH_2CH_3)$, $OCH(CH_3)_2$, $NH(CH_2)_2O(CH_2)_2CH_3$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $N(CH_3)CH_2CH_2CH_2OCH_3$, $NHCH_2CH_2COOH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  , or CO_2H.

In some embodiments, R_3 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C3-C10 heteroaryl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR.

In some embodiments, R_3 is a group represented by the corresponding moieties shown in the compounds of Table 1. For example, R_3 is Cl, I, deuterium, F, CN, CH_3, OH, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2$, CO_2H, CO_2NH_2, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CH_2CONH_2, CO_2CH_3, $-CH_2N(CH_3)_2$, CO_2tBu, tBu, $=CH_2$, =O, 



In some embodiments, o is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, n is 0. In some embodiments, o is 1. In some embodiments, o is 2.

In some embodiments, p is 0. In some embodiments, p is 1. In some embodiments, p is 2.

In another embodiment, the present invention provides compounds of formulas I to Id-ii that contain isotope-labelled forms thereof. An isotope-labelled form of a compound of formulas I to Id-ii is identical to this compound apart from the fact that one or more atoms of the compound have been replaced by an atom or atoms having an atomic mass or mass number which differs from the atomic mass or mass number of the atom which usually occurs in greater natural abundance. Examples of isotopes which are readily commercially available and which can be incorporated into a compound of formulas I to Id-ii by well-known methods include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine and chlorine, for example ^{2}H, ^{3}H, ^{13}C, ^{14}C, ^{15}N, ^{18}O, ^{17}O, ^{31}P, ^{32}P, ^{35}S, ^{18}F and ^{36}Cl, respectively. A compound of formulas I to Id-ii, a prodrug thereof, or a pharmaceutically acceptable salt of either which contains one or more of the above-mentioned isotopes and/or other iso-topes of other atoms is intended to be part of the present invention. An isotope-labelled compound of formulas I to Id-ii can be used in a number of beneficial ways. For example, an isotope-labelled compound of formula I to In-i into which, for example, a radioisotope, such as ^{3}H or ^{14}C, has been incorporated is suitable for medicament and/or substrate tissue distribution assays. These radioisotopes, i.e. tritium (^{3}H) and carbon-14 (^{14}C), are particularly preferred owing to simple preparation and excellent

detectability. Incorporation of heavier isotopes, for example deuterium (^{2}H), into a compound of formulas I to Id-ii has therapeutic advantages owing to the higher metabolic stability of this isotope-labelled compound. Higher metabolic stability translates directly into an increased in vivo half-life or lower dosages, which under most circumstances would represent a preferred embodiment of the present invention. An isotope-labelled compound of formulas I to Id-ii can usually be prepared by carrying out the procedures disclosed in the synthesis schemes and the related description, in the example part and in the preparation part in the present text, replacing a non-isotope-labelled reactant by a readily available isotope-labelled reactant.

Deuterium (^{2}H) can also be incorporated into a compound of formulas I to Id-ii for the purpose in order to manipulate the oxidative metabolism of the compound by way of the primary kinetic isotope effect. The primary kinetic isotope effect is a change of the rate for a chemical reaction that results from exchange of isotopic nuclei, which in turn is caused by the change in ground state energies necessary for covalent bond formation after this isotopic exchange. Exchange of a heavier isotope usually results in a lowering of the ground state energy for a chemical bond and thus causes a reduction in the rate-limiting bond breakage. If the bond breakage occurs in or in the vicinity of a saddle-point region along the coordinate of a multi-product reaction, the product distribution ratios can be altered substantially. For explanation: if deuterium is bonded to a carbon atom at a non-exchangeable position, rate differences of k_M/k_D = 2-7 are typical. If this rate difference is successfully applied to a compound of formulas I to Id-ii that is susceptible to oxidation, the profile of this compound in vivo can be drastically modified and result in improved pharmacokinetic properties. For a further discussion, see S. L. Harbeson and R. D. Tung, *Deuterium In Drug Discovery and Development*, Ann. Rep. Med. Chem. 2011, 46, 403-417, incorporated in its entirety herein by reference.

When discovering and developing therapeutic agents, the person skilled in the art attempts to optimise pharmacokinetic parameters while retaining desirable in vitro properties. It is reasonable to assume that many compounds with poor pharmacokinetic profiles are susceptible to oxidative metabolism. In vitro liver microsomal assays currently available provide valuable information on the course of oxidative metabolism of this type, which in turn permits the rational design of deuterated compounds of formulas I to Id-ii with improved stability through resistance to such oxidative metabolism. Significant improvements in the pharmacokinetic profiles of compounds of formulas I to Id-ii are thereby obtained, and can be expressed quantitatively in













terms of increases in the in vivo half-life ($t_{1/2}$), concentration at maximum therapeutic effect (C_{max}), area under the dose response curve (AUC), and bioavailability; and in terms of reduced clearance, dose and materials costs.













The following is intended to illustrate the above: a compound of formulas I to Id-ii which has multiple potential sites of attack for oxidative metabolism, for example benzylic hydrogen atoms and hydrogen atoms bonded to a nitrogen atom, is prepared as a series of analogues in which various combinations of hydrogen atoms are replaced by deuterium atoms, so that some, most or all of these hydrogen atoms have been replaced by deuterium atoms. Half-life determinations enable favourable and accurate determination of the extent to which the improvement in resistance to oxidative metabolism has improved. In this way, it is determined that the half-life of the parent compound can be extended by up to 100% as the result of deuterium-hydrogen exchange of this type.













Deuterium-hydrogen exchange in a compound of formulas I to Id-ii can also be used to achieve a favorable modification of the metabolite spectrum of the starting compound in order to diminish or eliminate undesired toxic metabolites. For example, if a toxic metabolite arises through oxidative carbon-hydrogen (C-H) bond cleavage, it can reasonably be assumed that the deuterated analogue will greatly diminish or eliminate production of the unwanted metabolite, even if the particular oxidation is not a rate-determining step. Further information on the state of the art with respect to deuterium-hydrogen exchange may be found, for example in Hanzlik et al., J. Org. Chem. 55, 3992-3997, 1990, Reider et al., J. Org. Chem. 52, 3326-3334, 1987, Foster, Adv. Drug Res. 14, 1-40, 1985, Gillette et al, Biochemistry 33(10) 2927-2937, 1994, and Jarman et al. Carcinogenesis 16(4), 683-688, 1993.













In some embodiments, the compound of formula I is selected from Table 1:













Table 1.













1	2	3
		
4	5	6
		
7	8	9
		
10	11	12
		













13	14	15
		
16	17	18
		
19	20	21
		
22	23	24
		













25	26	27
		
28	29	30
		
31	32	33
		
34	35	36
		













37	38	39
		
40	41	42
		
43	44	45
		
46	47	48
		













49	50	51
		
52	53	54
		
55	56	57
		
58	59	60
		













61	62	63
		
64	65	66
		
67	68	69
		
70	71	72
		













73	74	75
		
76	77	78
		
79	80	81
		
82	83	84
		













85	86	87
		
88	89	90
		
91	92	93
		
94	95	96
		













97	98	99
		
100	101	102
		
103	104	105
		
106	107	108
		













109	110	111
		
112	113	114
		
115	116	117
		
118	119	120
		













121	122	123
		
124	125	126
		
127	128	129
		
130	131	132
		













133	134	135
		
136	**137**	**138**
		
139	**140**	**141**
		
142	**143**	**144**
		













145	146	147
		
148	**149**	**150**
		
151	**152**	**153**
		
154	**155**	**156**
		













157	158	159
		
160	161	162
		
163	164	165
		
166	167	168
		













169	170	171
		
172	173	174
		
175	176	177
		
178	179	180
		













181	182	183
		
184	185	186
		
187	188	189
		
190	191	192
		













193	194	195
		
196	197	198
		
199	200	201
		
202	203	204
		













205	206	207
		
208	209	210
		
211	212	213
		
214	215	216
		













217	218	219
		
220	221	222
		
223	224	225
		
226	227	228
		













229	230	231
		
232	**233**	**234**
		
235	**236**	**237**
		
238	**239**	**240**
		













241	242	243
		
244	245	246
		
247	248	249
		
250	251	252
		













253	254	255
		
256	257	258
		
259	260	261
		
262	263	264
		













265	266	267
		
268	269	270
		
271	272	273
		
274	275	276
		













277	278	279
		
280	281	282
		
283	284	285
		
286	287	288
		













289	290	291
		
292	293	294
		
295	296	297
		
298	299	300
		













301	302	303
		
304	305	306
		
307	308	309
		
310	311	312
		













313	314	315
		
316	317	318
		
319	320	321
		
322	323	324
		













325	326	327
		
328	329	330
		
331	332	333
		
334	335	336
		













337	338	339
		
340	341	342
		
343	344	345
		
346	347	348
		













349	350	351
		
352	353	354
		
355	356	357
		
358	359	360
		













361	362	363
		
364	365	366
		
367	368	369
		
370	371	372
		













373	374	375
		
376	377	378
		
379	380	381
		
382	383	384
		













385	386	387
		
388	389	390
		
391	392	393
		
394	395	396
		













397	398	399
		
400	**401**	**402**
		
403	**404**	**405**
		
406	**407**	**408**
		













409	410	411
		
412	413	414
		
415	416	417
		
418	419	420
		













421	422	423
		
424	425	426
		
427	428	429
		
430	431	432
		













433	434	435
		
436	437	438
		
439	440	441
		
442	443	444
		













445	446	447
		
448	449	450
		
451	452	453
		
454	455	456
		













457	458	459
		
460	461	462
		
463	464	465
		
466	467	468
		













469	470	471
		
472	473	474
		
475	476	477
		
478	479	480
		













481	482	483
		
484	485	486
		
487	488	489
		
490	491	492
		













493	494	495
		
496	497	498
		
499	500	501
		
502	503	504
		













505	506	507
		
508	509	510
		
511	512	513
		
514	515	516
		













517	518	519
		
520	521	522
		
523	524	525
		
526	527	528
		













529	530	531
		
532	533	534
		
535	536	537
		
538	539	540
		













541	542	543
		
544	545	546
		
547	548	549
		
550	551	552
		













553	554	555
		
556	557	558
		
559	560	561
		
562	563	564
		













565	566	567
		
568	569	570
		
571	572	573
		
574	575	576
		













577	578	579
		
580	581	582
		
583	584	585
		
586	587	588
		













589	590	591
		
592	593	594
		
595	596	597
		
598	599	600
		













601	602	603
		
604	605	606
		
607	608	609
		
610	611	612
		













613	614	615
		
616	617	618
		
619	620	621
		
622	623	624
		













625	626	627
		
628	629	630
		
631	632	633
		
634	635	636
		













637	638	639
		
640	**641**	**642**
		
643	**644**	**645**
		
646	**647**	**648**
		













649	650	651
		
652	653	654
		
655	656	657
		
658	659	660
		













661	662	663
		
664	665	666
		
667	668	669
		
670	671	672
		













673	674	675
		
676	677	678
		
679	680	681
		
682	683	684
		













685	686	687
		
688	689	690
		
691	692	693
		
694	695	696
		













697	698	699
		
700	701	702
		
703	704	705
		
706	707	708
		













709	710	711
		
712	713	714
		
715	716	717
		
718	719	720
		













721	722	723
		
724	725	726
		
727	728	729
		
730	731	732
		













733	734	735
		
736	737	738
		
739	740	741
		
742	743	744
		













745	746	747
		
748	749	750
		
751	752	753
		
754	755	756
		













757	758	759
		
760	761	762
		
763	764	765
		
766	767	768
		













769	770	771
		
772	773	774
		
775	776	777
		
778	779	780
		













781	782	783
		
784	785	786
		
787	788	789
		
790	791	792
		













793	794	795
		
796	797	798
		
799	800	801
		
802	803	804
		













805	806	807
		
808	809	810
		
811	812	813
		
814	815	816
		













817	818	819
		
820	821	822
		
823	824	825
		
826	827	828
		













829	830	831
		
832	833	834
		
835	836	837
		
838	839	840
		













841	842	843
		
844	845	846
		
847	848	849
		
850	851	852
		













853	854	855
		
856	857	858
		
859	860	861
		
862	863	864
		













865	866	867
		
868	869	870
		
871	872	873
		
874	875	876
		













877	878	879
		
880	881	882
		
883	884	885
		
886	887	888
		













889	890	891
		
892	893	894
		
895	896	897
		
898	899	900
		













901	902	903
		
904	905	906
		
907	908	909
		
910	911	912
		













913	914	915
		
916	917	918
		
919	920	921
		
922	923	924
		













925	926	927
		
928	929	930
		
931	932	933
		
934	935	936
		













937	938	939
		
940	941	942
		
943	944	945
		
946	947	948
		













949	950	951
		
952	953	954
		
955	956	957
		
958	959	960
		













961	962	963
		
964	965	966
		
967	968	969
		
970	971	972
		













973	974	975
		
976	977	978
		
979	980	981
		
982	983	984
		













985	986	987
		
988	989	990
		
991	992	993
		
994	995	996
		













997	998	999
		
1000	1001	1002
		
1003	1004	1005
		
1006	1007	1008
		













1009	1010	1011
		
1012	1013	1014
		
1015	1016	1017
		
1018	1019	1020
		













1021	1022	1023
		
1024	1025	1026
		
1027	1028	1029
		
1030	1031	1032
		













1033	1034	1035
		
1036	**1037**	**1038**
		
1039	**1040**	**1041**
		
1042	**1043**	**1044**
		













1045	1046	1047
		
1048	**1049**	**1050**
		
1051	**1052**	**1053**
		
1054	**1055**	**1056**
		













1057	1058	1059
		
1060	1061	1062
		
1063	1064	1065
		
1066	1067	1068
		













1069	1070	1071
		
1072	1073	1074
		
1075	1076	1077
		
1078	1079	1080
		













1081	1082	1083
		
1084	1085	1086
		
1087	1088	1089
		
1090	1091	1092
		













1093	1094	1095
		
1096	**1097**	**1098**
		
1099	**1100**	**1101**
		
1102	**1103**	**1104**
		













1105	1106	1107
		
1108	1109	1110
		
1111	1112	1113
		
1114	1115	1116
		













1117	1118	1119
		
1120	**1121**	**1122**
		
1123	**1124**	**1125**
		
1126	**1127**	**1128**
		













1129	1130	1131
		
1132	1133	1134
		
1135	1136	1137
		
1138	1139	1140
		













1141	1142	1143
		
1144	**1145**	**1146**
		
1147	**1148**	**1149**
		
1150	**1151**	**1152**
		













1153	1154	1155
		
1156	1157	1158
		
1159	1160	1161
		
1162	1163	1164
		













1165	1166	1167
		
1168	1169	1170
		
1171	1172	1173
		
1174	1175	1176
		













1177	1178	1179
		
1180	**1181**	**1182**
		
1183	**1184**	**1185**
		
1186	**1187**	**1188**
		













1189	1190	1191
		
1192	1193	1194
		
1195	1196	1197
		
1198	1199	1200
		













1201	1202	1203
		
1204	**1205**	**1206**
		
1207	**1208**	**1209**
		
1210	**1211**	**1212**
		













1213	1214	1215
		
1216	1217	1218
		
1219	1220	1221
		
1222	1223	1224
		













1225	1226	1227
		
1228	1229	1230
		
1231	1232	1233
		
1234	1235	1236
		













1237	1238	1239
		
1240	**1241**	**1242**
		
1243	**1244**	**1245**
		
1246	**1247**	**1248**
		













1249	1250	1251
		
1252	1253	1254
		
1255	1256	1257
		
1258	1259	1260
		













1261	1262	1263
		
1264	1265	1266
		
1267	1268	1269
		
1270	1271	1272
		













1273	1274	1275
		
1276	**1277**	**1278**
		
1279	**1280**	**1281**
		
1282	**1283**	**1284**
		













1285	1286	1287
		
1288	1289	1290
		
1291	1292	1293
		
1294	1295	1296
		













1297	1298	1299
		
1300	1301	1302
		
1303	1304	1305
		
1306	1307	1308
		













1309	1310	1311
		
1312	1313	1314
		
1315	1316	1317
		
1318	1319	1320
		













1321	1322	1323
		
1324	1325	1326
		
1327	1328	1329
		
1330	1331	1332
		













1333	1334	1335
		
1336	**1337**	**1338**
		
1339	**1340**	**1341**
		
1342	**1343**	**1344**
		













1345	1346	1347
		
1348	**1349**	**1350**
		
1351	**1352**	**1353**
		
1354	**1355**	**1356**
		













1357	1358	1359
		
1360	**1361**	**1362**
		
1363	**1364**	**1365**
		
1366	**1367**	**1368**
		













1369	1370	1371
		
1372	**1373**	**1374**
		
1375	**1376**	**1377**
		
1378	**1379**	**1380**
		













1381	1382	1383
		
1384	1385	1386
		
1387	1388	1389
		
1390	1391	1392
		













1393	1394	1395
		
1396	1397	1398
		
1399	1400	1401
		
1402	1403	1404
		













1405	1406	1407
		
1408	1409	1410
		
1411	1412	1413
		
1414	1415	1416
		













1417	1418	1419
		
1420	1421	1422
		
1423	1424	1425
		
1426	1427	1428
		













1429	1430	1431
		
1432	1433	1434
		
1435	1436	1437
		
1438	1439	1440
		













1441	1442	1443
		
1444	**1445**	**1446**
		
1447	**1448**	**1449**
		
1450	**1451**	**1452**
		













1453	1454	1455
		
1456	**1457**	**1458**
		
1459	**1460**	**1461**
		
1462	**1463**	**1464**
		













1465	1466	1467
		
1468	**1469**	**1470**
		
1471	**1472**	**1473**
		
1474	**1475**	**1476**
		













1477	1478	1479
		
1480	**1481**	**1482**
		
1483	**1484**	**1485**
		
1486	**1487**	**1488**
		













1489	1490	1491
		
1492	1493	1494
		
1495	1496	1497
		
1498	1499	1500
		













1501	1502	1503
		
1504	1505	1506
		
1507	1508	1509
		
1510	1511	1512
		













1513	1514	1515
		
1516	1517	1518
		
1519	1520	1521
		
1522	1523	1524
		













1525	1526	1527
		
1528	1529	1530
		
1531	1532	1533
		
1534	1535	1536
		













1537	1538	1539
		
1540	**1541**	**1542**
		
1543	**1544**	**1545**
		
1546	**1547**	**1548**
		













1549	1550	1551
		
1552	1553	1554
		
1555	1556	1557
		
1558	1559	1560
		













1561	1562	1563
		
1564	**1565**	**1566**
		
1567	**1568**	**1569**
		
1570	**1571**	**1572**
		













1573	1574	1575
		
1576	1577	1578
		
1579	1580	1581
		
1582	1583	1584
		













1585	1586	1587
		
1588	1589	1590
		
1591	1592	1593
		
1594	1595	1596
		













1597	1598	1599
		
1600	1601	1602
		
1603	1604	1605
		
1606	1607	1608
		













1609	1610	1611
		
1612	**1613**	**1614**
		
1615	**1616**	**1617**
		
1618	**1619**	**1620**
		













1621	1622	1623
		
1624	**1625**	**1626**
		
1627	**1628**	**1629**
		
1630	**1631**	**1632**
		













1633	1634	1635
		
1636	**1637**	**1638**
		
1639	**1640**	**1641**
		
1642	**1643**	**1644**
		













1645	1646	1647
		
1648	**1649**	**1650**
		
1651	**1652**	**1653**
		
1654	**1655**	**1656**
		













1657	1658	1659
		
1660	1661	1662
		
1663	1664	1665
		
1666	1667	1668
		













1669	1670	1671
		
1672	**1673**	**1674**
		
1675	**1676**	**1677**
		
1678	**1679**	**1680**
		













1681	1682	1683
		
1684	**1685**	**1686**
		
1687	**1688**	**1689**
		
1690	**1691**	**1692**
		













1693	1694	1695
		
1696	**1697**	**1698**
		
1699	**1700**	**1701**
		
1702	**1703**	**1704**
		













1705	1706	1707
		
1708	1709	1710
		
1711	1712	1713
		
1714	1715	1716
		













1717	1718	1719
		
1720	**1721**	**1722**
		
1723	**1724**	**1725**
		
1726	**1727**	**1728**
		













1729	1730	1731
		
1732	1733	1734
		
1735	1736	1737
		
1738	1739	1740
		













1741	1742	1743
		
1744	**1745**	**1746**
		
1747	**1748**	**1749**
		
1750	**1751**	**1752**
		













1753	1754	1755
		
1756	1757	1758
		
1759	1760	1761
		
1762	1763	1764
		













1765	1766	1767
		
1768	1769	1770
		
1771	1772	1773
		
1774	1775	1776
		













1777	1778	1779
		
1780	1781	1782
		
1783	1784	1785
		
1786	1787	1788
		













1789	1790	1791
		
1792	1793	1794
		
1795	1796	1797
		
1798	1799	1800
		













1801	1802	1803
		
1804	**1805**	**1806**
		
1807	**1808**	**1809**
		
1810	**1811**	**1812**
		













1813	1814	1815
		
1816	1817	1818
		
1819	1820	1821
		
1822	1823	1824
		













1825	1826	1827
		
1828	1829	1830
		
1831	1832	1833
		
1834	1835	1836
		













1837	1838	1839
		
1840	**1841**	**1842**
		
1843	**1844**	**1845**
		
1846	**1847**	**1848**
		













1849	1850	1851
		
1852	**1853**	**1854**
		
1855	**1856**	**1857**
		
1858	**1859**	**1860**
		













1861	1862	1863
		
1864	1865	1866
		
1867	1868	1869
		
1870	1871	1872
		













1873	1874	1875
		
1876	**1877**	**1878**
		
1879	**1880**	**1881**
		
1882	**1883**	**1884**
		













1885	1886	1887
		
1888	1889	1890
		
1891	1892	1893
		
1894	1895	1896
		













1897	1898	1899
		
1900	1901	1902
		
1903	1904	1905
		
1906	1907	1908
		













1909	1910	1911
		
1912	**1913**	**1914**
		
1915	**1916**	**1917**
		
1918	**1919**	**1920**
		













1921	1922	1923
		
1924	**1925**	**1926**
		
1927	**1928**	**1929**
		
1930	**1931**	**1932**
		













1933	1934	1935
		
1936	**1937**	**1938**
		
1939	**1940**	**1941**
		
1942	**1943**	**1944**
		













1945	1946	1947
		
1948	1949	1950
		
1951	1952	1953
		
1954	1955	1956
		













1957	1958	1959
		
1960	1961	1962
		
1963	1964	1965
		
1966	1967	1968
		













1969	1970	1971
		
1972	**1973**	**1974**
		
1975	**1976**	**1977**
		
1978	**1979**	**1980**
		













1981	1982	1983
		
1984	**1985**	**1986**
		
1987	**1988**	**1989**
		
1990	**1991**	**1992**
		













1993	1994	1995
		
1996	**1997**	**1998**
		
1999	**2000**	**2001**
		
2002	**2003**	**2004**
		













2005	2006	2007
		
2008	**2009**	**2010**
		
2011	**2012**	**2013**
		
2014	**2015**	**2016**
		













2017	2018	2019
		
2020	**2021**	**2022**
		
2023	**2024**	**2025**
		
2026	**2027**	**2028**
		













2029	2030	2031
		
2032	2033	2034
		
2035	2036	2037
		
2038	2039	2040
		













2041	2042	2043
		
2044	**2045**	**2046**
		
2047	**2048**	**2049**
		
2050	**2051**	**2052**
		













2053	2054	2055
		
2056	2057	2058
		
2059	2060	2061
		
2062	2063	2064
		













2065	2066	2067
		
2068	**2069**	**2070**
		
2071	**2072**	**2073**
		
2074	**2075**	**2076**
		













2077	2078	2079
		
2080	2081	2082
		
2083	2084	2085
		
2086	2087	2088
		













2089	2090	2091
		
2092	**2093**	**2094**
		
2095	**2096**	**2097**
		
2098	**2099**	**2100**
		













2101	2102	2103
		
2104	**2105**	**2106**
		
2107	**2108**	**2109**
		
2110	**2111**	**2112**
		













2113	2114	2115
		
2116	**2117**	**2118**
		
2119	**2120**	**2121**
		
2122	**2123**	**2124**
		













2125	2126	2127
		
2128	2129	2130
		
2131	2132	2133
		
2134	2135	2136
		













2137	2138	2139
		
2140	**2141**	**2142**
		
2143	**2144**	**2145**
		
2146	**2147**	**2148**
		













2149	2150	2151
		
2152	2153	2154
		
2155	2156	2157
		
2158	2159	2160
		













2161	2162	2163
		
2164	**2165**	**2166**
		
2167	**2168**	**2169**
		
2170	**2171**	**2172**
		













2173	2174	2175
		
2176	2177	2178
		
2179	2180	2181
		
2182	2183	2184
		













2185	2186	2187
		
2188	2189	2190
		
2191	2192	2193
		
2194	2195	2196
		













2197	2198	2199
		
2200	**2201**	**2202**
		
2203	**2204**	**2205**
		
2206	**2207**	**2208**
		













2209	2210	2211
		
2212	2213	2214
		
2215	2216	2217
		
2218	2219	2220
		













2221	2222	2223
		
2224	**2225**	**2226**
		
2227	**2228**	**2229**
		
2230	**2231**	**2232**
		













2233	2234	2235
		
2236	**2237**	**2238**
		
2239	**2240**	**2241**
		
2242	**2243**	**2244**
		













2245	2246	2247
		
2248	**2249**	**2250**
		
2251	**2252**	**2253**
		
2254	**2255**	**2256**
		













2257	2258	2259
		
2260	**2261**	**2262**
		
2263	**2264**	**2265**
		
2266	**2267**	**2268**
		













2269	2270	2271
		
2272	**2273**	**2274**
		
2275	**2276**	**2277**
		
2278	**2279**	**2280**
		













2281	2282	2283
		
2284	**2285**	**2286**
		
2287	**2288**	**2289**
		
2290	**2291**	**2292**
		













2293	2294	2295
		
2296	**2297**	**2298**
		
2299	**2300**	**2301**
		
2302	**2303**	**2304**
		













2305	2306	2307
		
2308	**2309**	**2310**
		
2311	**2312**	**2313**
		
2314	**2315**	**2316**
		













2317	2318	2319
		
2320	2321	2322
		
2323	2324	2325
		
2326	2327	2328
		













2329	2330	2331
		
2332	**2333**	**2334**
		
2335	**2336**	**2337**
		
2338	**2339**	**2340**
		













2341	2342	2343
		
2344	**2345**	**2346**
		
2347	**2348**	**2349**
		
2350	**2351**	**2352**
		













2353	2354	2355
		
2356	**2357**	**2358**
		
2359	**2360**	**2361**
		
2362	**2363**	**2364**
		













2365	2366	2367
		
2368	**2369**	**2370**
		
2371	**2372**	**2373**
		
2374	**2375**	**2376**
		













2377	2378	2379
		
2380	2381	2382
		
2383	2384	2385
		
2386	2387	2388
		













2389	2390	2391
		
2392	2393	2394
		
2395	2396	2397
		
2398	2399	2400
		













2401	2402	2403
		
2404	**2405**	**2406**
		
2407	**2408**	**2409**
		
2410	**2411**	**2412**
		













2413	2414	2415
		
2416	**2417**	**2418**
		
2419	**2420**	**2421**
		
2422	**2423**	**2424**
		













2425	2426	2427
		
2428	**2429**	**2430**
		
2431	**2432**	**2433**
		
2434	**2435**	**2436**
		













2437	2438	2439
		
2440	2441	2442
		
2443	2444	2445
		
2446	2447	2448
		













2449	2450	2451
		
2452	2453	2454
		
2455	2456	2457
		
2458	2459	2460
		













2461	2462	2463
		
2464	**2465**	**2466**
		
2467	**2468**	**2469**
		
2470	**2471**	**2472**
		













2473	2474	2475
		
2476	**2477**	**2478**
		
2479	**2480**	**2481**
		
2482	**2483**	**2484**
		













2485	2486	2487
		
2488	2489	2490
		
2491	2492	2493
		
2494	2495	2496
		













2497	2498	2499
		
2500	**2501**	**2502**
		
2503	**2504**	**2505**
		
2506	**2507**	**2508**
		













2509	2510	2511
		
2512	2513	2514
		
2515	2516	2517
		
2518	2519	2520
		













2521	2522	2523
		
2524	**2525**	**2526**
		
2527	**2528**	**2529**
		
2530	**2531**	**2532**
		













2533	2534	2535
		
2536	**2537**	**2538**
		
2539	**2540**	**2541**
		
2542	**2543**	**2544**
		













2545	2546	2547
		
2548	2549	2550
		
2551	2552	2553
		
2554	2555	2556
		













2557	2558	2559
		
2560	**2561**	**2562**
		
2563	**2564**	**2565**
		
2566	**2567**	**2568**
		













2569	2570	2571
		
2572	**2573**	**2574**
		
2575	**2576**	**2577**
		
2578	**2579**	**2580**
		













2581	2582	2583
		
2584	**2585**	**2586**
		
2587	**2588**	**2589**
		
2590	**2591**	**2592**
		













2593	2594	2595
		
2596	**2597**	**2598**
		
2599	**2600**	**2601**
		
2602	**2603**	**2604**
		













2605	2606	2607
		
2608	2609	2610
		
2611	2612	2613
		
2614	2615	2616
		













2617	2618	2619
		
2620	2621	2622
		
2623	2624	2625
		
2626	2627	2628
		













2629	2630	2631
		
2632	2633	2634
		
2635	2636	2637
		
2638	2639	2640
		













2641	2642	2643
		
2644	**2645**	**2646**
		
2647	**2648**	**2649**
		
2650	**2651**	**2652**
		













2653	2654	2655
		
2656	2657	2658
		
2659	2660	2661
		
2662	2663	2664
		













2665	2666	2667
		
2668	2669	2670
		
2671	2672	2673
		
2674	2675	2676
		












2677	2678	2679
		
2680	2681	2682
		
2683	2684	2685
		
2686	2687	2688
		

2689	2690	2691
		
2692	2693	2694
		
2695	2696	2697
		
2698	2699	2700
		

2701	2702	2703
		
2704	**2705**	**2706**
		
2707	**2708**	**2709**
		
2710	**2711**	**2712**
		


2713	2714	2715
		
2716	**2717**	**2718**
		
2719	**2720**	**2721**
		
2722	**2723**	**2724**
		

2725	2726	2727
		
2728	**2729**	**2730**
		
2731	**2732**	**2733**
		
2734	**2735**	**2736**
		

2737	2738	2739
		
2740	**2741**	**2742**
		
2743	**2744**	**2745**
		
2746	**2747**	
		

In one embodiment, the present invention features any of the below numerated embodiments.

1. In one embodiment, the present invention features a compound of formula I:



I,

or pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring A is a C6-C10 mono- or bicyclic aryl ring, or C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced by O, S, N, or NR;

Ring B is a C6-C10 aryl ring or C5-C10 heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced by O, S, N, or NR;

Ring C is a C6-C10 mono- or bicyclic aryl ring , C4-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 carbon atoms may be replaced by N, O, or S, or a C3-C10 mono- or bicyclic cycloalkyl ring;

X is O or NR;

Y is CR_2, O, S, SO, SO_2, or NR;

Z is NR or CR_2;

R_1 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or

NR; or C3-C8 cycloalkyl, all four of which may be substituted with halo, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, CN, CO_2H, amino, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C5-C10 mono- or bicyclic heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; or two R_1 along with the atoms to which they are attached form a C5-C8 cyclic or heterocyclic ring wherein up to 2 carbon atoms may be replaced with O, S, or NR;

R_2 is halo, OH, amino, azide, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR, C3-C8 cycloalkyl; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR;

R_3 is halo, CN, CO_2R, C1-C6 alkyl, alkene, alkyne, or fluoroalkyl; C1-C6 alkoxy, cycloalkoxy, or fluoroalkoxy; C6-C10 mono- or bicyclic aryl or C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR, both of which may be substituted with halo, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, CN, CO_2H, amino, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C5-C10 mono- or bicyclic heterocycloalkyl wherein up to 4 carbon atoms may be replaced by O, S, or NR; or or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; or two R_3 may form a $=CH_2$ or =O group;

R_4 is H, CF_3, CO_2R, OH, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C3-C8 cycloalkyl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR; $N(R)_2$, NRCOR, $CON(R)_2$, CN, halo, or SO_2R;

R is H, OH, C1-C6 alkyl, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C3-

C8 cycloalkyl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR;

n is 0, 1, 2 or 3;




o is 0, 1, 2, or 3;


p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5;


provided that the moieties containing ring B and ring C are substituted at adjacent positions on ring A.

2. In another embodiment, the present invention features the compound of embodiment 1, wherein ring A is a pyridyl, indole, indoline, isoindoline, pyrazole, pyrimidine, phenyl, quinoline, 5,6,7,8-tetrahydroquinoline, 1,2,3,4-tetrahydroisoquinoline, pyrrolidine, aza-indole, pyrrole, oxazole, pyrazine, triazole, benzimidazole, indazole, or imidazole ring.


3. In another embodiment, the present invention features the compound of embodiment 1 or 2, wherein ring A is selected from  ,  ,  ,




4. In another embodiment, the present invention features the compound of any of embodiments 1 to 3, wherein ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, thiophene, dihydrobenzofuran, pyrazine, indazole, thiazole, pyridine-4(1H)-one, pyrrolidinone, or quinoline.

5. In another embodiment, the present invention features the compound of any of embodiments 1 to 4, wherein ring B is selected from 

6. In another embodiment, the present invention features the compound of any of embodiments 1 to 5, wherein ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, piperidine, azetidine, or dihydroindene.

7. In another embodiment, the present invention features the compound of any of embodiments 1 to 6, wherein ring C is 



8. In another embodiment, the present invention features the compound of any of embodiments 1 to 7, wherein Y is O.

9. In another embodiment, the present invention features the compound of any of embodiments 1 to 7, wherein Y is CH_2.

10. In another embodiment, the present invention features the compound of any of embodiments 1 to 7, wherein Y is CH(C1-C6 alkyl).

11. In another embodiment, the present invention features the compound of any of embodiments 1 to 7, wherein Y is $CH(CH_3)$.


12. In another embodiment, the present invention features the compound of any of embodiments 1 to 7, wherein Y is $CH(CH_2CH_3)$.


13. In another embodiment, the present invention features the compound of any of embodiments 1 to 7, wherein Y is N(C1-C6 alkyl).


14. In another embodiment, the present invention features the compound of any of embodiments 1 to 7, wherein Y is $N(CH_3)$.


15. In another embodiment, the present invention features the compound of any of embodiments 1 to 14, wherein R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, azetidine, or thiophene ring, wherein all rings may be substituted with halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C5-C10 heteroaryl, C5-C10 heterocycloalkyl, or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

16. In another embodiment, the present invention features the compound of any of embodiments 1 to 15, wherein R_1 is selected from CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$,












17. In another embodiment, the present invention features the compound of any of embodiments 1 to 16, wherein R_2 is selected from halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C5-C10 mono- or bicyclic heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

18. In another embodiment, the present invention features the compound of any of embodiments 1 to 17, wherein R_2 is selected from Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, CH_3, CH_2OH, CH_2CH_3, $CH(CH_3)_2$, CHF_2, OCH_3, OCF_3, $OCHF_2$, $OCH(CH_3)_2$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  , or CO_2H.

19. In another embodiment, the present invention features the compound of any of embodiments 1 to 18, wherein R_3 is selected from halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C5-C10 mono- or bicyclic heteroaryl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR.

20. In another embodiment, the present invention features the compound of any of embodiments 1 to 19, wherein R_3 is selected from Cl, F, CN, CH_3, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CO_2CH_3, tBu, $=CH_2$, =O,  ,



21. In another embodiment, the present invention features the compound of any of embodiments 1 to 20, wherein o is 0.

22. In another embodiment, the present invention features the compound of any of embodiments 1 to 20, wherein o is 1.

23. In another embodiment, the present invention features the compound of any of embodiments 1 to 20, wherein o is 2.

24. In another embodiment, the present invention features the compound of any of embodiments 1 to 23, wherein n is 0.

25. In another embodiment, the present invention features the compound of any of embodiments 1 to 23, wherein n is 1.


26. In another embodiment, the present invention features the compound of any of embodiments 1 to 23, wherein n is 2.

27. In another embodiment, the present invention features the compound of any of embodiments 1 to 26, wherein p is 0.

28. In another embodiment, the present invention features the compound of any of embodiments 1 to 26, wherein p is 1.

29. In another embodiment, the present invention features the compound of any of embodiments 1 to 26, wherein p is 2.

30. In another embodiment, the present invention features the compound of embodiment 1, wherein the compound is of formula Ia:



Ia

or pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C5-C10 heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced by O, S, N, or NR;

Ring C is a C6-C10 mono- or bicyclic aryl ring , C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 carbon atoms may be replaced by N, O, or S, or a C3-C10 mono- or bicyclic cycloalkyl ring;

X is O or NR;

Y is CR_2, O, S, SO, SO_2, or NR;

R_1 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or C3-C8 cycloalkyl, all four of which may be substituted with halo, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, CN, CO_2H, amino, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C5-C10 mono- or bicyclic heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; or two R_1 along with the atoms to which they are attached form a C5-C8 cyclic or heterocyclic ring wherein up to 2 carbon atoms may be replaced with O, S, or NR;

R_2 is halo, OH, amino, azide, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR, C3-C8 cycloalkyl; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR;

R_3 is halo, CN, CO_2R, C1-C6 alkyl, alkene, alkyne, or fluoroalkyl; C1-C6 alkoxy, cycloalkoxy, or fluoroalkoxy; C6-C10 mono- or bicyclic aryl or C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR, both of which may be substituted with halo, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, CN, CO_2H, amino, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C5-C10 mono- or bicyclic heterocycloalkyl wherein up to 4 carbon atoms may be replaced by O, S, or NR; or or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; or two R_3 may form a $=CH_2$ or =O group;

R_4 is H, CF_3, CO_2R, OH, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C3-C8 cycloalkyl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR; $N(R)_2$, NRCOR, $CON(R)_2$, CN, halo, or SO_2R;

R is H, OH, C1-C6 alkyl, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C3-C8 cycloalkyl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR;


n is 0, 1, 2 or 3;

o is 0, 1, 2, or 3;


p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.

31. In another embodiment, the present invention features a compound of embodiment 30, wherein ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, thiophene, dihydrobenzofuran, pyrazine, indazole, thiazole, pyridine-4(1H)-one, pyrrolidinone, or quinoline.

32. In another embodiment, the present invention features a compound of embodiment 30 or 31, wherein ring B is selected from 

33. In another embodiment, the present invention features a compound of any of embodiments 30 to 32, wherein ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, piperidine, or dihydroindene.

34. In another embodiment, the present invention features a compound of any of embodiments 30 to 33, wherein ring C is 



35. In another embodiment, the present invention features a compound of any of embodiments 30 to 34, wherein Y is O.

36. In another embodiment, the present invention features a compound of any of embodiments 30 to 34, wherein Y is CH_2.


37. In another embodiment, the present invention features a compound of any of embodiments 30 to 34, wherein Y is CH(C1-C6 alkyl).


38. The compound of claim 30, wherein Y is $CH(CH_3)$.


39. In another embodiment, the present invention features a compound of any of embodiments 30 to 34, wherein Y is $CH(CH_2CH_3)$.

40. In another embodiment, the present invention features a compound of any of embodiments 30 to 39, wherein R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, or thiophene ring, wherein all rings may be substituted with halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C5-C10 heteroaryl, C5-C10 heterocycloalkyl, or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

41. In another embodiment, the present invention features a compound of any of embodiments 30 to 40, wherein R_1 is selected from CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$,












42. In another embodiment, the present invention features a compound of any of embodiments 30 to 41, wherein R_2 is selected from halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C5-C10 mono- or bicyclic heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

43. In another embodiment, the present invention features a compound of any of embodiments 30 to 42, wherein R_2 is selected from Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, CH_3, CH_2OH, CH_2CH_3, $CH(CH_3)_2$, CHF_2, OCH_3, OCF_3, $OCHF_2$, $OCH(CH_3)_2$, $C(O)CH_3$, CH_2CH_2OH,

CH_2NH_2, $NH(CH_2)_2OH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  ,  ,  , or CO_2H.

44. In another embodiment, the present invention features a compound of any of embodiments 30 to 43, wherein R_3 is selected from halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR.

45. In another embodiment, the present invention features a compound of any of embodiments 30 to 44, wherein R_3 is selected from Cl, F, CN, CH_3, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CO_2CH_3, tBu, $=CH_2$, =O,  ,  .

46. In another embodiment, the present invention features a compound of any of embodiments 30 to 45, wherein o is 0.

47. In another embodiment, the present invention features a compound of any of embodiments 30 to 45, wherein o is 1.

48. In another embodiment, the present invention features a compound of any of embodiments 30 to 47, wherein n is 0.

49. In another embodiment, the present invention features a compound of any of embodiments 30 to 47, wherein n is 1.

50. In another embodiment, the present invention features a compound of any of embodiments 30 to 47, wherein n is 2.

51. In another embodiment, the present invention features a compound of any of embodiments 30 to 50, wherein p is 0.


52. In another embodiment, the present invention features a compound of any of embodiments 30 to 50, wherein p is 1.

53. In another embodiment, the present invention features a compound of any of embodiments 30 to 50, wherein p is 2.

54. In another embodiment, the present invention features a compound of any of embodiments 30 to 53, wherein ring B is phenyl.

55. In another embodiment, the present invention features a compound of any of embodiments 30 to 54, wherein ring B and ring C are phenyl.

56. In another embodiment, the present invention features a compound of embodiment 1, wherein the compound is of formula Ib:



Ib

or pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C5-C10 heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced by O, S, N, or NR;

Ring C is a C6-C10 mono- or bicyclic aryl ring , C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 carbon atoms may be replaced by N, O, or S, or a C3-C10 mono- or bicyclic cycloalkyl ring;

Y is CR_2, O, S, SO, SO_2, or NR;

R_1 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or C3-C8 cycloalkyl, all four of which may be substituted with halo, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, CN, CO_2H, amino, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C5-C10 mono- or bicyclic heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; or two R_1 along with the atoms to which they are attached form a C5-C8 cyclic or heterocyclic ring wherein up to 2 carbon atoms may be replaced with O, S, or NR;

R_2 is halo, OH, amino, azide, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR, C3-C8 cycloalkyl; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR;

R_3 is halo, CN, CO_2R, C1-C6 alkyl, alkene, alkyne, or fluoroalkyl; C1-C6 alkoxy, cycloalkoxy, or fluoroalkoxy; C6-C10 mono- or bicyclic aryl or C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR, both of which may be substituted with halo, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, CN, CO_2H, amino, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C5-C10 mono- or bicyclic heterocycloalkyl wherein up to 4 carbon atoms may be replaced by O, S, or NR; or or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; or two R_3 may form a $=CH_2$ or =O group;

R_4 is H, CF_3, CO_2R, OH, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C3-

C8 cycloalkyl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR; $N(R)_2$, NRCOR, $CON(R)_2$, CN, halo, or SO_2R;

R is H, OH, C1-C6 alkyl, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C3-C8 cycloalkyl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR;




n is 0, 1, 2 or 3;

o is 0, 1, 2, or 3;


p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5.

57. In another embodiment, the present invention features a compound of embodiment 56, wherein ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, thiophene, dihydrobenzofuran, pyrazine, indazole, thiazole, pyridine-4(1H)-one, pyrrolidinone, or quinoline.

58. In another embodiment, the presnt invention features a compound of embodiment 56 or 57, wherein ring B is selected from  ,  ,  , or .

59. In another embodiment, the present invention features a compound of any one of embodiments 56 to 58, wherein ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, or dihydroindene.

60. In another embodiment, the present invention features a compound of any one of embodiments 56 to 59, wherein ring C is  , or .

61. In another embodiment, the present invention features a compound of any one of embodiments 56 to 60, wherein Y is O.

62. In another embodiment, the present invention features a compound of any one of embodiments 56 to 60, wherein Y is CH_2.


63. In another embodiment, the present invention features a compound of any one of embodiments 56 to 60, wherein Y is CH(C1-C6 alkyl).


64. In another embodiment, the present invention features a compound of any one of embodiments 56 to 60, wherein Y is $CH(CH_3)$.


65. In another embodiment, the present invention features a compound of any one of embodiments 56 to 60, wherein Y is $CH(CH_2CH_3)$.


66. In another embodiment, the present invention features a compound of any one of embodiments 56 to 65, wherein R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, or thiophene ring, wherein all rings may be substituted with halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C5-C10 heteroaryl, C5-C10 heterocycloalkyl, or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

67. In another embodiment, the present invention features a compound of any one of embodiments 56 to 66, wherein R_1 is selected from CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu,












68. In another embodiment, the present invention features a compound of any one of embodiments 56 to 67, wherein R_2 is selected from halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C5-C10 mono- or bicyclic heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

69. In another embodiment, the present invention features a compound of any one of embodiments 56 to 68, wherein R_2 is selected from Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, CH_3, CH_2OH, CH_2CH_3, $CH(CH_3)_2$, CHF_2, OCH_3, OCF_3, $OCHF_2$, $OCH(CH_3)_2$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  , or CO_2H.

70. In another embodiment, the present invention features a compound of any one of embodiments 56 to 69, wherein R_3 is selected from halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR.

71. In another embodiment, the present invention features a compound of any one of embodiments 56 to 70, wherein R_3 is selected from Cl, F, CN, CH_3, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CO_2CH_3, tBu, $=CH_2$, =O,





72. In another embodiment, the present invention features a compound of any one of embodiments 56 to 71, wherein o is 0.

73. In another embodiment, the present invention features a compound of any one of embodiments 56 to 71, wherein o is 1.

74. In another embodiment, the present invention features a compound of any one of embodiments 56 to 71, wherein o is 2.

75. In another embodiment, the present invention features a compound of any one of embodiments 56 to 74, wherein n is 0.

76. In another embodiment, the present invention features a compound of any one of embodiments 56 to 74, wherein n is 1.

77. In another embodiment, the present invention features a compound of any one of embodiments 56 to 74, wherein n is 2.

78. In another embodiment, the present invention features a compound of any one of embodiments 56 to 77, wherein p is 0.

79. In another embodiment, the present invention features a compound of any one of embodiments 56 to 77, wherein p is 1.


80. In another embodiment, the present invention features a compound of any one of embodiments 56 to 77, wherein p is 2.

81. In another embodiment, the present invention features a compound of any one of embodiments 56 to 80, wherein ring B is pyridyl.

82. In another embodiment, the present invention features a compound of any one of embodiments 56 to 81, wherein ring B is phenyl.

83. In another embodiment, the present invention features a compound of any one of embodiments 56 to 82, wherein ring B and ring C are phenyl.

84. In another embodiment, the present invention features a compound of embodiment 1, wherein the compound is of formula Ic:



Ic

or pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C5-C10 heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced by O, S, N, or NR;

Ring C is a C6-C10 mono- or bicyclic aryl ring , C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 carbon atoms may be replaced by N, O, or S, or a C3-C10 mono- or bicyclic cycloalkyl ring;

Y is CR_2, O, S, SO, SO_2, or NR;

R_1 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or C3-C8 cycloalkyl, all four of which may be substituted with halo, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, CN, CO_2H, amino, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C5-C10 mono- or bicyclic heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; or two R_1 along

with the atoms to which they are attached form a C5-C8 cyclic or heterocyclic ring wherein up to 2 carbon atoms may be replaced with O, S, or NR;

R_2 is halo, OH, amino, azide, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR, C3-C8 cycloalkyl; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR;

R_3 is halo, CN, CO_2R, C1-C6 alkyl, alkene, alkyne, or fluoroalkyl; C1-C6 alkoxy, cycloalkoxy, or fluoroalkoxy; C6-C10 mono- or bicyclic aryl or C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR, both of which may be substituted with halo, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, CN, CO_2H, amino, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C5-C10 mono- or bicyclic heterocycloalkyl wherein up to 4 carbon atoms may be replaced by O, S, or NR; or or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; or two R_3 may form a $=CH_2$ or =O group;

R_4 is H, CF_3, CO_2R, OH, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C3-C8 cycloalkyl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR; $N(R)_2$, NRCOR, $CON(R)_2$, CN, halo, or SO_2R;

R is H, OH, C1-C6 alkyl, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C3-C8 cycloalkyl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR;


n is 0, 1, 2 or 3;

o is 0, 1, 2, or 3;


p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.

85. In another embodiment, the present invention features a compound of embodiment 84, wherein ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, thiophene, dihydrobenzofuran, pyrazine, indazole, thiazole, pyridine-4(1H)-one, pyrrolidinone, or quinoline.

86. In another embodiment, the present invention features a compound of embodiment 84 or 85, wherein ring B is selected from  .

87. In another embodiment, the present invention features a compound of any one of embodiments 84 to 86, wherein ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, or dihydroindene.

88. In another embodiment, the present invention features a compound of any one of embodiments 84 to 87, wherein ring C is  .



89. In another embodiment, the present invention features a compound of any one of embodiments 84 to 88, wherein Y is O.

90. In another embodiment, the present invention features a compound of any one of embodiments 84 to 88, wherein Y is CH_2.

91. In another embodiment, the present invention features a compound of any one of embodiments 84 to 88, wherein Y is CH(C1-C6 alkyl).


92. In another embodiment, the present invention features a compound of any one of embodiments 84 to 88, wherein Y is $CH(CH_3)$.


93. In another embodiment, the present invention features a compound of any one of embodiments 84 to 88, wherein Y is $CH(CH_2CH_3)$.


94. In another embodiment, the present invention features a compound of any one of embodiments 84 to 93, wherein R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, or thiophene ring, wherein all rings may be substituted with halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C5-C10 heteroaryl, C5-C10 heterocycloalkyl, or

a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

95. In another embodiment, the present invention features a compound of any one of embodiments 84 to 94, wherein R_1 is selected from CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$,










96. In another embodiment, the present invention features a compound of any one of embodiments 84 to 95, wherein R_2 is selected from halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C5-C10 mono- or bicyclic heterocyclic ring wherein

up to 4 carbon atoms may be replaced with O, S, N, or NR; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

97. In another embodiment, the present invention features a compound of any one of embodiments 84 to 96, wherein R_2 is selected from Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, CH_3, CH_2OH, CH_2CH_3, $CH(CH_3)_2$, CHF_2, OCH_3, OCF_3, $OCHF_2$, $OCH(CH_3)_2$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  ,  , or CO_2H.

98. In another embodiment, the present invention features a compound of any one of embodiments 84 to 97, wherein R_3 is selected from halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR.

99. In another embodiment, the present invention features a compound of any one of embodiments 84 to 98, wherein R_3 is selected from Cl, F, CN, CH_3, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CO_2CH_3, tBu, $=CH_2$, =O, 

100. In another embodiment, the present invention features a compound of any one of embodiments 84 to 99, wherein o is 0.

101. In another embodiment, the present invention features a compound of any one of embodiments 84 to 99, wherein o is 1.

102. In another embodiment, the present invention features a compound of any one of embodiments 84 to 99, wherein o is 2.

103. In another embodiment, the present invention features a compound of any one of embodiments 84 to 102, wherein n is 0.

104. In another embodiment, the present invention features a compound of any one of embodiments 84 to 102, wherein n is 1.

105. In another embodiment, the present invention features a compound of any one of embodiments 84 to 102, wherein n is 2.

106. In another embodiment, the present invention features a compound of any one of embodiments 84 to 105, wherein p is 0.

107. In another embodiment, the present invention features a compound of any one of embodiments 84 to 105, wherein p is 1.


108. In another embodiment, the present invention features a compound of any one of embodiments 84 to 105, wherein p is 2.

109. In another embodiment, the present invention features a compound of any one of embodiments 84 to 108, wherein ring B is phenyl.

110. In another embodiment, the present invention features a compound of any one of embodiments 84 to 109, wherein ring C is phenyl.

111. In another embodiment, the present invention features a compound of any one of embodiments 84 to 110, wherein ring B and ring C are phenyl.

112. In another embodiment, the present invention features a compound of embodiment 1, wherein the compound is of formula Id:



Id

or pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C5-C10 heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced by O, S, N, or NR;

Ring C is a C6-C10 mono- or bicyclic aryl ring , C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 carbon atoms may be replaced by N, O, or S, or a C3-C10 mono- or bicyclic cycloalkyl ring;

Y is CR_2, O, S, SO, SO_2, or NR;

R_1 is halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or C3-C8 cycloalkyl, all four of which may be substituted with halo, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, CN, CO_2H, amino, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C5-C10 mono- or bicyclic heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; or two R_1 along with the atoms to which they are attached form a C5-C8 cyclic or heterocyclic ring wherein up to 2 carbon atoms may be replaced with O, S, or NR;

R_2 is halo, OH, amino, azide, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl or heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR, C3-C8 cycloalkyl; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR;

R_3 is halo, CN, CO_2R, C1-C6 alkyl, alkene, alkyne, or fluoroalkyl; C1-C6 alkoxy, cycloalkoxy, or fluoroalkoxy; C6-C10 mono- or bicyclic aryl or C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR, both of which may be substituted with halo, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, CN, CO_2H, amino, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C5-C10 mono- or bicyclic heterocycloalkyl wherein up to 4 carbon atoms may be

replaced by O, S, or NR; or or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR; or two R_3 may form a $=CH_2$ or =O group;

R_4 is H, CF_3, CO_2R, OH, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C3-C8 cycloalkyl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR; $N(R)_2$, NRCOR, $CON(R)_2$, CN, halo, or SO_2R;

R is H, OH, C1-C6 alkyl, C6-C10 mono- or bicyclic aryl, C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR; C3-C8 cycloalkyl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced with O, S, or NR;




n is 0, 1, 2 or 3;

o is 0, 1, 2, or 3;


p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5.

113. In another embodiment, the present invention features a compound of embodiment 112, wherein ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, thiophene, dihydrobenzofuran, pyrazine, indazole, thiazole, pyridine-4(1H)-one, pyrrolidinone, or quinoline.

114. In another embodiment, the present invention features a compound of embodiment 112 or 113, wherein ring B is selected from  ,  ,  , or .

115. In another embodiment, the present invention features a compound of any one of embodiments 112 to 114, wherein ring C is phenyl, indole, cycloalkyl, pyridyl, pyrrolidine, naphthalene, or dihydroindene.

116. In another embodiment, the present invention features a compound of any one of embodiments 112 to 115, wherein ring C is  , or .

117. In another embodiment, the present invention features a compound of any one of embodiments 112 to 116, wherein Y is O.

118. In another embodiment, the present invention features a compound of any one of embodiments 112 to 116, wherein Y is CH_2.


119. In another embodiment, the present invention features a compound of any one of embodiments 112 to 116, wherein Y is CH(C1-C6 alkyl).


120. In another embodiment, the present invention features a compound of any one of embodiments 112 to 116, wherein Y is $CH(CH_3)$.


121. In another embodiment, the present invention features a compound of any one of embodiments 112 to 116, wherein Y is $CH(CH_2CH_3)$.


122. In another embodiment, the present invention features a compound of any one of embodiments 112 to 121, wherein R_1 is halo, CN, C1-C6 alkyl, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, or thiophene ring, wherein all rings may be substituted with halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 fluoroalkyl, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C5-C10 heteroaryl, C5-C10 heterocycloalkyl, or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

123. In another embodiment, the present invention features a compound of any one of embodiments 112 to 122, wherein R_1 is selected from CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu,











124. In another embodiment, the present invention features a compound of any one of embodiments 112 to 123, wherein R_2 is selected from halo, OH, CN, azide, amino, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy or fluoroalkoxy, C5-C10 mono- or bicyclic heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or a (C1-C8 alkyl)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

125. In another embodiment, the present invention features a compound of any one of embodiments 112 to 124, wherein R_2 is selected from Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, CH_3, CH_2OH, CH_2CH_3, $CH(CH_3)_2$, CHF_2, OCH_3, OCF_3, $OCHF_2$, $OCH(CH_3)_2$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  , or CO_2H.

126. In another embodiment, the present invention features a compound of any one of embodiments 112 to 125, wherein R_3 is selected from halo, CN, C1-C6 alkyl or fluoroalkyl, C1-C6 alkoxy, or C5-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR.

127. In another embodiment, the present invention features a compound of any one of embodiments 112 to 126, wherein R_3 is selected from Cl, F, CN, CH_3, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CO_2CH_3, tBu, $=CH_2$, =O,



128. In another embodiment, the present invention features a compound of any one of embodiments 112 to 127, wherein o is 0.

129. In another embodiment, the present invention features a compound of any one of embodiments 112 to 127, wherein o is 1.

130. In another embodiment, the present invention features a compound of any one of embodiments 112 to 127, wherein o is 2.

131. In another embodiment, the present invention features a compound of any one of embodiments 112 to 130, wherein n is 0.

132. In another embodiment, the present invention features a compound of any one of embodiments 112 to 130, wherein n is 1.

133. In another embodiment, the present invention features a compound of any one of embodiments 112 to 130, wherein n is 2.

134. In another embodiment, the present invention features a compound of any one of embodiments 112 to 133, wherein p is 0.

135. In another embodiment, the present invention features a compound of any one of embodiments 112 to 133, wherein p is 1.

136. In another embodiment, the present invention features a compound of any one of embodiments 112 to 133, wherein p is 2.

137. In another embodiment, the present invention features a compound of any one of embodiments 112 to 136, wherein ring B is phenyl.

138. In another embodiment, the present invention features a compound of any one of

embodiments 112 to 137, wherein ring B is pyridyl.

139. In another embodiment, the present invention features a compound of any one of embodiments 112 to 138, wherein ring C is phenyl.

140. In another embodiment, the present invention features a compound of any one of embodiments 112 to 139, wherein ring B and ring C are phenyl.

141. In another embodiment, the present invention features a compound of any one of embodiments 112 to 140, wherein ring B is pyridyl and ring C is phenyl.

142. In another embodiment, the present invention features a compound of embodiment 1, wherein the compound is selected from Table 1.

143. In another embodiment, the present invention features a pharmaceutical composition comprising the compound of any one of embodiments 1 to 142 and a pharmaceutically acceptable carrier.


144. In another embodiment, the present invention features the pharmaceutical composition of embodiment 143, further comprising one or more additional therapeutic agent(s).

145. In another embodiment, the present invention features the pharmaceutical composition of embodiment 144, wherein the additional therapeutic agent is selected from a mucolytic agent, bronchodialator, an antibiotic, an anti-infective agent, a CFTR modulator, or an anti-inflammatory agent.

146. In another embodiment, the present invention features the pharmaceutical composition of embodiment 144, wherein the additional therapeutic agent is a CFTR modulator.

147. In another embodiment, the present invention features the pharmaceutical composition of embodiment 144, wherein the additional therapeutic agent is a CFTR corrector.


148. In another embodiment, the present invention features the pharmaceutical composition of embodiment 144, wherein the additional therapeutic agent is



,


or pharmaceutically acceptable salt thereof.

149. In another embodiment, the present invention features the pharmaceutical composition of embodiment 144, wherein the additional therapeutic agent is



or pharmaceutically acceptable salt thereof.


150. In another embodiment, the present invention features the pharmaceutical composition of embodiment 144, wherein the additional therapeutic agent is



or pharmaceutically acceptable salt thereof.

151. In another embodiment, the present invention features the pharmaceutical composition of embodiment 144, wherein the additional therapeutic agent is a CFTR potentiator.



152. In another embodiment, the present invention features the pharmaceutical composition of embodiment 144, wherein the additional therapeutic agent is



or pharmaceutically acceptable salt thereof.



153. In another embodiment, the present invention features the pharmaceutical composition of embodiment 144, wherein the additional therapeutic agents are a CFTR corrector and a CFTR potentiator.

154. In another embodiment, the present invention features the pharmaceutical composition of embodiment 144, wherein the additional therapeutic agents are


and

, or



pharmaceutically acceptable salts thereof.

155. In another embodiment, the present invention features the pharmaceutical composition of embodiment 144, wherein the additional therapeutic agents are


and

, or

pharmaceutically acceptable salts thereof.

156. In another embodiment, the present invention features the pharmaceutical composition of embodiment 144, wherein the additional therapeutic agents are


and

, or

pharmaceutically acceptable salts thereof.

157. In another embodiment, the present invention features a method of treating cystic fibrosis in a patient comprising administering to the patient an effective amount of the compound of any one of embodiments 1 to 142 or the pharmaceutical composition of any one of embodiments 143 to 156.


158. In another embodiment, the present invention features the method of embodiment 157, further comprising administering to the patient one or more additional therapeutic agent(s) prior to, concurrent with, or subsequent to the compound of any one of embodiments 1 to 137 or the pharmaceutical composition of any one of embodiments 138 to 151.

159. In another embodiment, the present invention features the method of embodiment 158, wherein the additional therapeutic agent is selected from a mucolytic agent, bronchodialator, an antibiotic, an anti-infective agent, a CFTR modulator, or an anti-inflammatory agent.

160. In another embodiment, the present invention features the method of embodiment 158, wherein the additional therapeutic agent is a CFTR modulator.


161. In another embodiment, the present invention features the method of embodiment 158, wherein the additional therapeutic agent is a CFTR corrector.

162. In another embodiment, the present invention features the method of embodiment 158, wherein the additional therapeutic agent is




or a pharmaceutically acceptable salt thereof.

163. In another embodiment, the present invention features the method of embodiment 158, wherein the additional therapeutic agent is



or a pharmaceutically acceptable salt thereof.


164. In another embodiment, the present invention features the method of embodiment 158, wherein the additional therapeutic agent is



or a pharmaceutically acceptable salt thereof.

165. In another embodiment, the present invention features the method of embodiment 158, wherein the additional therapeutic agent is a CFTR potentiator.

166. In another embodiment, the present invention features the method of embodiment 158, wherein the additional therapeutic agent is



or a pharmaceutically acceptable salt thereof.

167. In another embodiment, the present invention features the method of any one of embodiments 157 to 166, wherein the patient is homozygous in the F508del CFTR mutation.

168. In another embodiment, the present invention features the method of any one of embodiments 157 to 166, wherein the patient is heterozygous in the F508del CFTR mutation.

169. In another embodiment, the present invention features a kit comprising the compound of any one of embodiments 1 to 142 or the pharmaceutical composition of any one of embodiments 143 to 156, and instructions for use therof.


170. In another embodiment, the present invention features the kit of embodiment 169, further comprising one or more additional therapeutic agent(s).

171. In another embodiment, the present invention features the kit of embodiment 170, wherein the additional therapeutic agent is selected from a mucolytic agent, bronchodialator, an antibiotic, an anti-infective agent, a CFTR modulator, or an anti-inflammatory agent.

172. In another embodiment, the present invention features the kit of embodiment 170, wherein the additional therapeutic agent is a CFTR modulator.

173. In another embodiment, the present invention features the kit of embodiment 170, wherein the additional therapeutic agent is a CFTR corrector.


174. In another embodiment, the present invention features the kit of embodiment 170, wherein the additional therapeutic agent is



,

or a pharmaceutically acceptable salt thereof.


175. In another embodiment, the present invention features the kit of embodiment 170, wherein the additional therapeutic agent is



,

or a pharmaceutically acceptable salt thereof.

176. In another embodiment, the present invention features the kit of embodiment 170, wherein the additional therapeutic agent is




,

or a pharmaceutically acceptable salt thereof.

177. In another embodiment, the present invention features the kit of embodiment 170, wherein the additional therapeutic agent is a CFTR potentiator.


178. In another embodiment, the present invention features the kit of embodiment 170, wherein the additional therapeutic agent is




,


or a pharmaceutically acceptable salt thereof.


179. In another embodiment, the present invention features the kit of embodiment 170, wherein the additional therapeutic agents are a CFTR corrector and a CFTR potentiator.

180. In another embodiment, the present invention features the kit of embodiment 170, wherein the additional therapeutic agents are  and





, or pharmaceutically acceptable salts thereof.

181. In another embodiment, the present invention features the kit of embodiment 170, wherein the additional therapeutic agents are  and



, or pharmaceutically acceptable salts thereof.

182. In another embodiment, the present invention features the kit of embodiment 170, wherein the additional therapeutic agents are  and  , or pharmaceutically acceptable salts thereof.

183. In another embodiment, the present invention features the kit of any one of embodiments 169 to 182, wherein the compound of any one of embodiments 1 to 142 or the pharmaceutical composition of any one of embodiments 143 to 156 and the one or more additional therapeutic agent(s) are in separate containers.


184. In another embodiment, the present invention features the kit of any one of embodiments 169 to 182, wherein the compound of any one of embodiments 1 to 142 or the pharmaceutical composition of any one of embodiments 143 to 156 and the one or more additional therapeutic agent(s) are in the same container.

185. In another embodiment, the present invention features the kit of embodiment 183 or 184, wherein the container is a bottle, vial, or blister pack, or combination thereof.

Methods of Preparing Compounds of Formula I


Compounds of the invention may be prepared by known methods and as illustrated in Schemes I-XX1. Ring A, B, C and R, R_1, R_2, R_3, R_4, X , Y, Z and n, o, p, q are as defined in the Specification unless noted otherwise below for each individual scheme.

Scheme I




a) R_5 = OH, R_6 = NH_2: HATU, Et_3N or $(i\text{-}Pr)_2NEt$ or K_2CO_3 or Na_2CO_3, DMF or CH_2Cl_2 or NMP; R_5 = OH, R_6 = NH_2: $SOCl_2$, CH_2Cl_2 and DMF (catalytic amount), then Et_3N, CH_2Cl_2; R_5 = NH_2, R_6 = Cl: NaH or Cs_2CO_3, DMF or THF; or pyridine, CH_2Cl_2.

Scheme II



a) NaH or Cs_2CO_3, DMF or THF; or pyridine, CH_2Cl_2.


Scheme III



a) R_7 = CN or CO_2R', R_6 = halogen (Cl or Br), Y = bond, R_8 = boronic acid or ester: $Pd(dppf)Cl_2*CH_2Cl_2$, $NaHCO_3$ or Na_2CO_3, water, DMA or DME; Y = O or N(R'), R_8 = H: NaH or Cs_2CO_3 or KHMDS; DMF or DMSO or dioxane or DMF/THF; Y = O, R_8 = H: $Cu(OTf)_2$,


Cs_2CO_3, toluene; Y = C(R')(R''), R_8 = ZnBr: $Pd(dppf)Cl_2*CH_2Cl_2$ or $Pd(PPh_3)_4$ or $Pd(OAc)_2$ and tBu_3P*HBF_4, THF; Y = C(R')(R''), R_8 = Cl or Br: K_2CO_3 or KOH, DMF or *i*-PrOH. b) KOH or NaOH or LiOH, water, *i*-PrOH or EtOH or MeOH; or HCl in AcOH. c) H_2O_2, NaOH, EtOH; or KOH, EtOH.

Scheme IV




a) R_6 = Br, CN or CO_2R'': $Cu(OAc)_2$, pyridine, DCE; b) R_6 = Br or is absent: *n*-BuLi, THF, CO_2.

Scheme V




a) $Cu(OTf)_2$, Cs_2CO_3, toluene. Y = O, R_8= H.

Scheme VI




a) Cl_2, water, $CHCl_3$; b) NH_3 or NH_4OH, MeOH or CH_2Cl_2 or dioxane.

Scheme VII




a) R_6= halogen (Cl), R_8 = $B(OR'')_2$, R_9 = a protecting group (*tert*-butyl) or when R_8 is absent R_1= moiety containing N: EtOH, toluene, water, Na_2CO_3, $Pd(PPh_3)_4$ or NaH and DMF; b) TFA, CH_2Cl_2; c) CDI, NaH, DMF ; d) C=moiety containing N, Y and R_8 are absent or R_8 = NHR, NH_2 or NRR: K_2CO_3, DMSO or CsF, K_2CO_3, DMSO.

Scheme VIII




a) X= NH: $(CH_3)_3COK$, CH_3CN, HMDS, *N*-chlorosuccinamide ; X = NCH_3or NR: CH_3NH_2 or RNH_2, CH_3CN, *N*-chlorosuccinamide.

Scheme IX




a) R_6= halogen (Cl), Y = bond, R_8= $B(OR'')_2$: $Pd(PPh_3)_4$ or $Pd(dppf)Cl_2*CH_2Cl_2$, Na_2CO_3, DMF, water or dioxane or DMA.

Scheme X




a) CSI, CH_2Cl_2, 45 min, then Et_3N, B=moiety containing N.

Scheme XI




a) R_6 = halogen (Cl), Y = bond or Y = O, R_8 = $B(OR'')_2$: $Pd(PPh_3)_4$, Na_2CO_3, K_2CO_3 or K_3PO_4, DMF; R_6 = halogen (Cl), Y = bond, R_8 = H: CsF, DMSO, then R_1= moiety containing N; R_6 = $B(OR'')_2$, Y = C(R')(R''), R_8 = halogen (Br): $Pd(dppf)Cl_2*CH_2Cl_2$, K_2CO_3, DMF; R_6 = halogen (Cl), R_8 = H or R_8 is absent, R_1= moiety containing N: *trans*-diaminocyclohexane, DMF, CuI; R_6= halogen (Cl), Y = bond, R_8 = $B(OR'')_2$, R_1= moiety containing C3-C8 cycloalkyl optionally substituted: PEPPSI™-IPr catalyst, K_2CO_3, dioxane.

Scheme XII




a) R_6 = halogen (F, Cl), Y = bond or Y = O, $R_2 = NH_2$, p = 1, X= O: NH_4OH; $R_6 = NO_2$, Y = O, $R_2 = NH_2$, p = 1, X = O: CH_3COOH, Zn; R_6 = OMe, Y = C(R')(R''), R_2 is =O, p = 1, X = O: CH_3COOH or dioxane, HBr or HCl; $R_6 = NO_2$, $R_2 = NH_2$, p = 1, X = NH: MeOH, Pd/C; R_6 = aldehyde, X = O, Y = C(R')(R''), R_2 = hydroxyalkyl: $NaBH_4$, MeOH.

Scheme XIII




a) R = alkyl or H, R_6 = halogen (Cl), $R_8 = NH_2$ or OH, Y = O: DMA or NaH and DMF; R = alkyl, R_6= halogen (Cl), R_8= alkyne: DMF, Et_3N, $PdCl_2(PPh_3)_2$, CuI, PPh_3; R = alkyl, R_6= halogen (Cl), Y = O, R_8 = ZnBr: $Pd(dppf)Cl_2*CH_2Cl_2$, then aq. sat. sol. of EDTA disodium salt and sat. aq. sol of NH_4Cl.

Scheme XIV




a) R_6 = halogen (I): $(CH_3)_3CMgCl$, CuCN, THF; b) 3-chlorobenzoperoxoic acid, CH_2Cl_2 c) R_7 = CN: TMSCN, Et_3N, CH_3CN.

Scheme XV




a) R_8 = Cl, Y = C(R')(R''), o = 1, R = alkyl (e.g., methyl): methyl *N*-aminocarbamate, K_2CO_3, DMF; b) R_6 = OH: dimethyl but-2-ynedioate , MeOH, then NaOMe c) Pyridine, toluene, triflic anhydride; d)R_1= boronic ester: Bis(pinacolato)diboron, KOAc, dioxane, Pd(dppf)Cl_2*CH_2Cl_2, DPPF e) Y = C(R')(R''), o = 1, R = alkyl (e.g., methyl): R_8 = boronic acid, K_2CO_3, dioxane, Pd(dppf)Cl_2*CH_2Cl_2.

Scheme XVI




a) R_6= halogen (Cl): CDI, NaH, DMF b) R_8 is absent, Y or o = 0: C=moiety containing N or R_8= $B(OR'')_2$, Y or o = 0: K_2CO_3, DMSO or Na_2CO_3, DMF, water, $Pd(PPh_3)_4$, c) $R_8 = B(OR'')_2$, $Pd(PPh_3)_4$, K_2CO_3, DMF or R_8 is absent: R_1=moiety containing N, n = 1: CsF, DMSO.

Scheme XVII




a) R = alkyl (methyl, ethyl): 2-nitropropane, dioxane, benzyltrimethylammonium hydroxide b) R_3 = gem-dimethyl group, q = 1, C = moiety containing N: $NiCl_2$ $6H_2O$, water, MeOH, $NaBH_4$; c) R_{10} = H or D: $LiAlH_4$, THF or $LiAlD_4$, THF.

Scheme XVIII




a) NBS, DCE; b) R_6 = Cl: $POCl_3$, PCl_5; c) R_6 = Cl: KOH, water,*i*-PrOH; d) CDI and DMF, then a mixture containing the desired sulfonoamide and NaH in DMF; e) C=moiety containing N, Y or o = 0: K_2CO_3, DMSO.

Scheme XIX




a) Z = CH_2, R_{11} = CO_2R, R = alkyl (methyl): NaH, DMF.

Scheme XX



a)Et_3N, $(CH_3)_3COH$, DPPA; b) MTBE, $(CH_3)_2CHMgCl$ in THF, then SO_2; c) CH_3COONa, H_2NOSO_3H.

Scheme XXI



a) DMSO, SMOPS; b) THF/MeOH 1:1, NaOMe in MeOH; c) DMSO, H_2NOSO_3H, CH_3COONa.

Pharmaceutically Acceptable Salts and Prodrugs of Compounds of Formulas I-Id

It will also be appreciated that certain compounds of the compositions of the present invention can exist in free form for treatment, or where appropriate, as a pharmaceutically acceptable derivative or a prodrug thereof. According to the present invention, a pharmaceutically acceptable derivative or a prodrug includes, but is not limited to, pharmaceutically acceptable salts, esters, salts of such esters, or any other adduct or derivative which upon administration to a patient in need is capable of providing, directly or indirectly, a compound as otherwise described herein, or a metabolite or residue thereof.

As used herein, the term "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. A "pharmaceutically acceptable salt" means any non-toxic salt or salt of an ester of a compound of this invention that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof.

Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge, *et al.* describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 1977, *66*, 1-19, incorporated herein by reference. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the

art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4}alkyl)_4$ salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersable products may be obtained by such quaternization. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

Pharmaceutical Compositions

Pharmaceutically Acceptable Vehicle, Adjuvant, or Carrier

In one aspect, the present invention features a pharmaceutical composition comprising a compound of formula I or pharmaceutically acceptable salt or prodrug thereof, and a pharmaceutically acceptable vehicle, adjuvant, or carrier.


As described above, the pharmaceutical compositions of the present invention additionally comprise a pharmaceutically acceptable carrier, adjuvant, or vehicle, which, as used herein, includes any and all solvents, diluents, or other liquid vehicle, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. Remington: *The Science and Practice of Pharmacy*, 21st edition, 2005, ed. D.B. Troy, Lippincott Williams & Wilkins, Philadelphia, and *Encyclopedia of Pharmaceutical Technology*, eds. J. Swarbrick and J. C. Boylan, 1988-1999, Marcel Dekker, New York, the contents of each of which is incorporated

by reference herein, disclose various carriers used in formulating pharmaceutically acceptable compositions and known techniques for the preparation thereof. Except insofar as any conventional carrier medium is incompatible with the compounds of the invention, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutically acceptable composition, its use is contemplated to be within the scope of this invention. Some examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, or potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, wool fat, sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil; sesame oil; olive oil; corn oil and soybean oil; glycols; such a propylene glycol or polyethylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.


Additional Therapeutic Agent(s)

In another embodiment, the pharmaceutical compositions of the present invention further comprise one or more additional therapeutic agent(s). In one embodiment, the additional therapeutic agent is a CFTR modulator. In one embodiment, the additional therapeutic agent is a CFTR corrector. In one embodiment, the additional therapeutic agent is a CFTR potentiator. In another embodiment, the pharmaceutical composition comprises a compound of formula I and one or more of the following additional therapeutic agents.


In another embodiment, the additional therapeutic agent is selected from:



3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof;



(*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or a pharmaceutically acceptable salt thereof; or



4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of formula I, or pharmaceutically acceptable salt thereof and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of Table 1, or pharmaceutically acceptable salt thereof and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of formula I, or pharmaceutically acceptable salt thereof and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of Table 1, or pharmaceutically acceptable salt thereof and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of formula I, or pharmaceutically acceptable salt thereof and 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of Table 1, or pharmaceutically acceptable salt thereof and 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of formula I, or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of Table 1, or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-

dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 233), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 238), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 318), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 228), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 196), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 337), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 368), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-

5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-8,8-dimethyl-2-(2,4,6-trimethylphenoxy)-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 182), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(benzenesulfonyl)-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1356), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-methylcyclohexyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 172), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-

difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 639), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(p-tolyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 300), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1660), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chlorophenyl)-2-(2,4,6-

trimethylphenoxy)pyridine-3-carboxamide (Compound 234), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1975), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-tert-butyl-*N*-[[6-(methylamino)-2-pyridyl]sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 262), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3-isobutoxypyrazol-1-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2191), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-

difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-2-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 207), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 320), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 353), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-

difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 171), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-isopropyl-1-piperidyl)pyridine-3-carboxamide (Compound 114), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 321), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-8,8-dimethyl-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 15), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(4-aminothiazol-2-yl)sulfonyl-6-tert-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 265), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 164), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-tert-butyl-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 214), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-

methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(7-fluoro-1,3-benzodioxol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 345), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 8), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 110), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isopropoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 281), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 351), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(3-aminopyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 197), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(4-chloro-3-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 136), or pharmaceutically acceptable salt

thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 2-(2,5-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 307), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(2-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 279), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(4-chloro-3-propoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 72), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-

methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(3-isobutoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 125), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(benzenesulfonyl)-5-tert-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 45), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(3-aminophenyl)sulfonyl-4-isopropyl-2-[(2,4,6-trimethylphenyl)methyl] isoindoline-1-carboxamide (Compound 299), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-methoxy-3-pyridyl)sulfonyl]-6-(1-methylcyclopropyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 365), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 361), or pharmaceutically acceptable salt thereof; and a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid, or pharmaceutically acceptable salts thereof.

In one embodiment, the above recited pharmaceutical compositions contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of formula I, or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of Table 1, or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[-3-phenyl-1-

piperidyl]pyridine-3-carboxamide (Compound 233), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 238), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 318), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 228), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 196), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 337), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 368), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-8,8-dimethyl-2-(2,4,6-trimethylphenoxy)-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 182), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(benzenesulfonyl)-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1356), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-methylcyclohexyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 172), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 639), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(p-tolyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 300), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-

difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1660), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chlorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 234), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1975), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-tert-butyl-*N*-[[6-(methylamino)-2-pyridyl]sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 262), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3-isobutoxypyrazol-1-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2191), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-2-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 207), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 320), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 353), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 171), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-isopropyl-1-piperidyl)pyridine-3-carboxamide (Compound 114), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 321), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-8,8-dimethyl-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 15), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(4-aminothiazol-2-yl)sulfonyl-6-tert-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 265), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 164), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-tert-butyl-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 214), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(7-fluoro-1,3-benzodioxol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 345), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 8), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 110), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isopropoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 281), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 351), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(3-aminopyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 197), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(4-chloro-3-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 136), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 2-(2,5-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 307), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(2-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 279), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(4-chloro-3-propoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 72), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(3-isobutoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 125), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(benzenesulfonyl)-5-tert-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 45), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(3-aminophenyl)sulfonyl-4-isopropyl-2-[(2,4,6-trimethylphenyl)methyl] isoindoline-1-carboxamide (Compound 299), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-methoxy-3-pyridyl)sulfonyl]-6-(1-methylcyclopropyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 365), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 361), or pharmaceutically acceptable salt thereof; and 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In one embodiment, the above recited pharmaceutical compositions contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of formula I, or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of Table 1, or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 233), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 238), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 318), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 228), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 196), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 337), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 368), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-8,8-dimethyl-2-(2,4,6-trimethylphenoxy)-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 182), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(benzenesulfonyl)-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1356), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-methylcyclohexyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 172), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 639), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(p-tolyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 300), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1660), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chlorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 234), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1975), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-tert-butyl-*N*-[[6-(methylamino)-2-pyridyl]sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 262), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3-isobutoxypyrazol-1-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2191), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-2-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 207), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 320), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 353), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 171), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-isopropyl-1-piperidyl)pyridine-3-carboxamide (Compound 114), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 321), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-8,8-dimethyl-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 15), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(4-aminothiazol-2-yl)sulfonyl-6-tert-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 265), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 164), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-tert-butyl-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 214), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(7-fluoro-1,3-benzodioxol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 345), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 8), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 110), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isopropoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 281), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 351), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(3-aminopyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 197), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(4-chloro-3-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 136), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 2-(2,5-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 307), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(2-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 279), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(4-chloro-3-propoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 72), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(3-isobutoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 125), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(benzenesulfonyl)-5-tert-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 45), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(3-aminophenyl)sulfonyl-4-isopropyl-2-[(2,4,6-trimethylphenyl)methyl] isoindoline-1-carboxamide (Compound 299), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-methoxy-3-pyridyl)sulfonyl]-6-(1-methylcyclopropyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 365), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 361), or pharmaceutically acceptable salt thereof; and (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropane carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the above recited pharmaceutical compositions contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of formula I, or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a compound of Table 1, or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 233), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 238), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 318), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 228), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-

2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 196), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 337), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 368), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-8,8-dimethyl-2-(2,4,6-trimethylphenoxy)-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 182), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(benzenesulfonyl)-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1356), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-methylcyclohexyl)-2-(2,4,6-

trimethylphenoxy)pyridine-3-carboxamide (Compound 172), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 639), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(p-tolyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 300), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1660), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chlorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 234), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1975), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-tert-butyl-*N*-[[6-(methylamino)-2-pyridyl]sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 262), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3-isobutoxypyrazol-1-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2191), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-2-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 207), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 320), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 353), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 171), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-isopropyl-1-piperidyl)pyridine-3-carboxamide (Compound 114), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 321), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-8,8-dimethyl-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 15), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(4-aminothiazol-2-yl)sulfonyl-6-tert-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 265), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 164), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-tert-butyl-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 214), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(7-fluoro-1,3-benzodioxol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 345), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 8), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 110), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isopropoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 281), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 351), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(3-aminopyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 197), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(4-chloro-3-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 136), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 2-(2,5-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 307), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(2-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 279), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(4-chloro-3-propoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 72), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises 6-(3-isobutoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 125), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(benzenesulfonyl)-5-tert-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 45), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.




In another embodiment, the pharmaceutical composition of the present invention comprises *N*-(3-aminophenyl)sulfonyl-4-isopropyl-2-[(2,4,6-trimethylphenyl)methyl] isoindoline-1-carboxamide (Compound 299), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.













In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(2-methoxy-3-pyridyl)sulfonyl]-6-(1-methylcyclopropyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 365), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.




In another embodiment, the pharmaceutical composition of the present invention comprises *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 361), or pharmaceutically acceptable salt thereof; and *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.



In another embodiment, the above recited pharmaceutical compositions contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In another embodiment, the additional therapeutic agent is selected from the following table:

1	2	3
		

4	5	6
		
7	8	9
		
10	11	12
		
13	14	15
		

16	17	18
		


19	20
	

In one embodiment, the additional therapeutic agent is selected from the following table:


Compounds disclosed in US Patent No. 7,407,976 (Col 13, ln 35- col 66, ln 67; Compounds 1-100 in Table 1 at col 67, ln 1-col 127, ln 42)
Compounds disclosed in US Patent No. 7,645,789 (Col 16, ln 52-col 50, ln 22; Compounds 1-322 in Table 1 at col 50, ln 24-col 167, ln 42)
Compounds disclosedin US Patent No. 7,659,268 (Col 16, ln 20-col 70, ln 52; Compounds 1-528 in Table 1 at col 70, ln 53-col 331, ln 34)
Compounds disclosed in US Patent No. 7,671,221 (Col 16, ln 12-col 54, ln 48; Compounds 1-1216 in Table 1 at col 54, ln 49-col 699, ln 27)
Compounds disclosed in US Patent No. 7,691,902 (Col 16, ln 11-col 54, ln 29; Compounds 1-959 in Table 1 at col 54, ln 29-col 683, ln 44)
Compounds disclosed in US Patent No. 7,741,321 (Col 16, ln 25-col 72, ln 17; Compounds 1-422 in Table 1 at col 72, ln 20-col 279, ln 15)
Compounds disclosed in US Patent No. 7,754,739 (Col 16, ln 1-col 22, ln 47; Compounds 1-2 in Table 1 at col 18, ln 26-65)
Compounds disclosed in US Patent No. 7,776,905 (Col 16, ln 23-col 38, ln 40; Compounds 1-306 in Table 1 at col 38, ln 45-col 96, ln 40)

Compounds disclosed in US Patent No. 7,973,169 (Col 9, ln 16-col 40, ln 40; Compounds 1-289 in Table 1 at col. 40, ln 41-col 289, ln 39)
Compounds disclosed in US Patent No. 7,977,322 (Col 6, ln 26-col 37, ln 47; Compounds 1-498 in Table 1 at col 37, ln 50-col 141, ln 40)
Compounds disclosed in US Patent No. 7,999,113 (Col 6, ln 13-col 10, ln 67; Compounds 1-13 in Table 1 at col 11, ln 5-col 13, ln 65)
Compounds disclosed in US Patent No. 8,227,615 (Col 6, ln 10-col 29, ln 66; Compounds 1-78 in Table 1 at col 30, ln 1-col 46, ln 48)
Compounds disclosed in US Patent No. 8,299,099 (Col 6, ln 10-col 34, ln 18; Compounds 1-47 in Table 1 at col 34, ln 20-col 42, ln 35)
Compounds disclosed in US Published Application No. 2006-0052358 (Paragraphs [0034]-[0056]; [0077]-[0240]; Compounds 1-320 in Table 1 at paragraph [0241])
Compounds disclosed in US Published Application No. 2009-0143381 (Paragraphs [0102]-[0263]; Compounds 1-28 in Table 1 at paragraph [0264])
Compounds disclosed in US Published Application No. 2009-0170905 (Paragraphs [0012]-[0013]; [0030]-[0051])
Compounds disclosed in US Published Application No. 2009-0253736 (Paragraphs [0031]-[0162]; Compounds 1-15 in Table 1 at paragraph [0163])
Compounds disclosed in US Published Application No. 2011-0263654 (Paragraphs [0012]-[0013]; [0066]-[0141])
Compounds disclosed in US Published Application No. 2011-0251253 (Paragraphs [0012]-[0013]; [0054]-[0079])
Compounds disclosed in PCT application WO2008141119 (Paragraphs [0100]-[0339]; Compounds 1-117 in Table 1 at paragraph [0340])
Compounds disclosed in US Application No. 11/047,361
Compounds disclosed in US Published Application No. 2013-0116238 (Paragraphs [0028]-[0044]; [0117]-[0128]), or combinations thereof.

In another embodiment, the additional therapeutic agent is selected from



N-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; or



N-(4-(7-azabicyclo[2.21]heptan-7-yl)-2-(trifluoromethyl)phenyl)-4-oxo-5-(trifluoromethyl)-1,4-dihydroquinoline-3-carboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) a compound selected from *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide or *N*-(4-(7-azabicyclo[2.21]heptan-7-yl)-2-(trifluoromethyl)phenyl)-4-oxo-5-(trifluoromethyl)-1,4-dihydroquinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) a compound selected from *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide or *N*-(4-(7-azabicyclo[2.21]heptan-7-yl)-2-(trifluoromethyl)phenyl)-4-oxo-5-(trifluoromethyl)-1,4-dihydroquinoline-3-carboxamide.

In another embodiment, the above recited pharmaceutical compositions contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)

cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; and c) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; and c) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; and c) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; and c) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the above recited pharmaceutical compositions contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In another embodiment, the additional therapeutic agent is selected from the following table:

Compounds disclosed in US Published Application No. 2005-0113423 (Paragraph [00146]; Compounds IA-1-IA-136 and Compounds I-1-I-21 in Tables 1 and 2 at paragraphs [0391]-[0392])
Compounds disclosed in US Published Application No. 2005-0059687 (Paragraphs [00100]-[00101]; Compounds 1-405 in Table 1 at paragraph [0169])
Compounds 1-108 disclosed in US Patent No. 7,598,412 (Col 22, ln 14-col 79, ln 20;Table 1)
Compounds 1-485 disclosed in US Patent No. 7,495,103 (Col 51, ln 1-col 63, ln 43; Table 1)
Compounds 1-718 disclosed in US Patent No. 8,354,427 (Col 51, ln 3-col 71, ln 46; Table 1)
Compounds 1-233 disclosed in US Published Application No. 2007-0105833 (Paragraph [00145]; Table 1)
Compounds 1-26 disclosed in US Patent No. 8,242,149 (Col 46, ln 47-col 57, ln 37; Table 1)
Compounds 1-18 disclosed in US Patent No. 8,314,256 (Col 21, ln 1-col 26, ln 19)
Compounds 1-14 disclosed in US Patent No. 8,399,479 (Col 36, ln 20-col 38, ln 40; Table 1)
Compounds 1-18 disclosed in US Patent No. 8,188,283 (Col 38, ln 43-col 43, ln 36; Table 1)
Compounds 1-16 disclosed in US Published Application No. 2010-0249180 (Paragraph [0173]; Table 1)
Compounds 1-19 disclosed in US Published Application No. 2011-0008259 (Paragraph [0172]; Table 1)
Compounds 1-129 disclosed in US Patent No. 8,367,660 (Col 57, ln 31-col 81, ln 24; Table1)

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) a compound selected from *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide or *N*-(4-(7-azabicyclo[2.2.1]heptan-7-yl)-2-(trifluoromethyl)phenyl)-4-oxo-5-(trifluoromethyl)-1,4-dihydroquinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) a compound selected from *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide or *N*-(4-(7-azabicyclo[2.21]heptan-7-yl)-2-(trifluoromethyl)phenyl)-4-oxo-5-(trifluoromethyl)-1,4-dihydroquinoline-3-carboxamide.

In another embodiment, the above recited pharmaceutical compositions contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) a compound selected from certain flavones and isoflavones, such as genistein, which are capable of stimulating CFTR-mediated chloride transport in epithelial tissues in a cyclic-AMP independent manner (See U.S. Pat. No. 6,329,422, incorporated herein by reference in its entirety); phenylglycine-01 (2-[(2-1H-indol-3-yl-acetyl)-methylamino]-*N*-(4-isopropylphenyl)-2-phenylacetamide); felodipine (Ethyl methyl 4-(2,3-dichlorophenyl)-2,6-dimethyl-1,4-dihydro-3,5-pyridinedicarboxylate); sulfonamide SF-01 (6-(ethylphenylsulfamoyl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid cycloheptylamide); and UCCF-152 (3-[2-(benzyloxy)phenyl]-5-(chloromethyl)isoxazole).

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; and c) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; and c) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; and c) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; and c) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the above recited pharmaceutical compositions contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 233), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)

cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 238), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 318), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 228), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 196), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 337), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 368), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-8,8-dimethyl-2-(2,4,6-trimethylphenoxy)-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 182), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-(benzenesulfonyl)-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1356), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-methylcyclohexyl)-2-(2,4,6-

trimethylphenoxy)pyridine-3-carboxamide (Compound 172), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 639), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-(p-tolyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 300), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1660), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chlorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 234), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)

cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1975), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-tert-butyl-*N*-[[6-(methylamino)-2-pyridyl]sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 262), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3-isobutoxypyrazol-1-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2191), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-2-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 207), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 320), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 353), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 171), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-isopropyl-1-piperidyl)pyridine-3-carboxamide (Compound 114), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-methyl-1-

piperidyl)pyridine-3-carboxamide (Compound 321), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-8,8-dimethyl-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 15), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-(4-aminothiazol-2-yl)sulfonyl-6-tert-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 265), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 164), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-tert-butyl-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 214), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt

thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(7-fluoro-1,3-benzodioxol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 345), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 8), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl] pyrazole-3-carboxamide (Compound 110), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isopropoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 281), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 351), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-(3-aminopyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 197), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-(4-chloro-3-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 136), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 2-(2,5-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 307), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-(2-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-

trimethylphenoxy)pyridine-3-carboxamide (Compound 279), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-(4-chloro-3-propoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 72), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-(3-isobutoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 125), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-(benzenesulfonyl)-5-tert-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 45), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-(3-aminophenyl)sulfonyl-4-isopropyl-2-[(2,4,6-trimethylphenyl)methyl] isoindoline-1-carboxamide (Compound 299), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)

cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(2-methoxy-3-pyridyl)sulfonyl]-6-(1-methylcyclopropyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 365), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 361), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, or pharmaceutically acceptable salt thereof.

In another embodiment, the above recited pharmaceutical compositions contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 233), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 238), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-

(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 318), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 228), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 196), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 337), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 368), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-8,8-dimethyl-2-(2,4,6-trimethylphenoxy)-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 182), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-(benzenesulfonyl)-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1356), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-methylcyclohexyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 172), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 639), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-(p-tolyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 300), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1660), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chlorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 234), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1975), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-tert-butyl-*N*-[[6-(methylamino)-2-pyridyl]sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 262), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3-isobutoxypyrazol-1-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2191), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-2-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 207), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 320), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 353), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 171), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-isopropyl-1-piperidyl)pyridine-3-carboxamide (Compound 114), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 321), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-8,8-dimethyl-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 15), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-(4-aminothiazol-2-yl)sulfonyl-6-tert-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 265), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 164), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-tert-butyl-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-

carboxamide (Compound 214), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(7-fluoro-1,3-benzodioxol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 345), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 8), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl] pyrazole-3-carboxamide (Compound 110), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isopropoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 281), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or

pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 351), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-(3-aminopyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5H-quinoline-3-carboxamide (Compound 197), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-(4-chloro-3-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 136), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 2-(2,5-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 307), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-

(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-(2-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 279), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-(4-chloro-3-propoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 72), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) 6-(3-isobutoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 125), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-(benzenesulfonyl)-5-tert-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 45), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-(3-aminophenyl)sulfonyl-4-isopropyl-2-[(2,4,6-trimethylphenyl)methyl] isoindoline-1-carboxamide (Compound 299), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(2-methoxy-3-pyridyl)sulfonyl]-6-(1-methylcyclopropyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 365), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the pharmaceutical composition of the present invention comprises a) *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 361), or pharmaceutically acceptable salt thereof; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide, or pharmaceutically acceptable salt thereof; and c) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or pharmaceutically acceptable salt thereof.

In another embodiment, the above recited pharmaceutical compositions contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In another embodiment, the present invention features a pharmaceutical composition comprising a) a compound of formula I; b) a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) an additional CFTR corrector.

In another embodiment, the present invention features a pharmaceutical composition comprising a) a compound of Table 1; b) a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) an additional CFTR corrector.

In another embodiment, the above recited pharmaceutical composition contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In another embodiment, the additional CFTR corrector from above is (*N*-(2-(5-chloro-2-methoxy-phenylamino)-4'-methyl-[4,5']bithiazolyl-2'-yl)-benzamide (corr-4a). In another embodiment, the additional CFTR corrector is a compound disclosed in published International Patent Application WO2014081820, incorporated herein in its entirety by reference.

In another embodiment, the present invention features a pharmaceutical composition comprising a) a compound of formula I; b) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide or *N*-(4-(7-azabicyclo[2.2.1]heptan-7-yl)-2-(trifluoromethyl)phenyl)-4-oxo-5-(trifluoromethyl)-1,4-dihydroquinoline-3-carboxamide; and a CFTR corrector selected from (*N*-(2-(5-chloro-2-methoxy-phenylamino)-4'-methyl-[4,5']bithiazolyl-2'-yl)-benzamide (corr-4a), a compound disclosed in published International Patent Application WO2014081820, incorporated herein in its entirety by reference, a compound disclosed in Published International Patent Application No. WO2014086687, or a compound disclosed in US Patent No. 8,247,436 or 8,476,269; US Patent Application No. 13/923,349; or Published International Patent Application No: WO2011113894, WO2013038373, WO2013038378, WO2013038381, WO2013038386, or WO2013038390; all US patents, US patent applications, and published international patent applications are hereby incorporated in their entirety by reference.

In another embodiment, the present invention features a pharmaceutical composition comprising a) a compound of formula I; b) a compound selected from certain flavones and isoflavones, such as genistein, which are capable of stimulating CFTR-mediated chloride transport in epithelial tissues in a cyclic-AMP independent manner (See U.S. Pat. No. 6,329,422, incorporated herein by reference in its entirety); phenylglycine-01 (2-[(2-1H-indol-3-yl-acetyl)-

methylamino]-*N*-(4-isopropylphenyl)-2-phenylacetamide); felodipine (Ethyl methyl 4-(2,3-dichlorophenyl)-2,6-dimethyl-1,4-dihydro-3,5-pyridinedicarboxylate); sulfonamide SF-01 (6-(ethylphenylsulfamoyl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid cycloheptylamide); and UCCF-152 (3-[2-(benzyloxy)phenyl]-5-(chloromethyl)isoxazole); and c) a CFTR corrector selected from (*N*-(2-(5-chloro-2-methoxy-phenylamino)-4'-methyl-[4,5']bithiazolyl-2'-yl)-benzamide (corr-4a), a compound disclosed in published International Patent Application WO2014081820, incorporated herein in its entirety by reference, or a compound disclosed in US Patent No. 8,247,436 or 8,476,269; US Patent Application No. 13/923,349; or Published International Patent Application No: WO20111 13894, WO2013038373, WO2013038378, WO2013038381, WO2013038386, or WO2013038390; all US patents, US patent applications, and published international patent applications are hereby incorporated in their entirety by reference.

In another embodiment, the present invention features the pharmaceutical composition described above comprising a) a compound of formula I; b) a compound selected from 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid, (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide, or 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) an additional CFTR corrector, such as those, for example, specifically listed above; and further comprising d) a CFTR potentiator.

In another embodiment, the CFTR potentiator from above is *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide or *N*-(4-(7-azabicyclo[2.21]heptan-7-yl)-2-(trifluoromethyl)phenyl)-4-oxo-5-(trifluoromethyl)-1,4-dihydroquinoline-3-carboxamide. In another embodiment, the CFTR potentiator from above is *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the CFTR potentiator from above is selected from certain flavones and isoflavones, such as genistein, which are capable of stimulating CFTR-mediated chloride transport in epithelial tissues in a cyclic-AMP independent manner (See U.S. Pat. No. 6,329,422, incorporated herein by reference in its entirety); phenylglycine-01 (2-[(2-1H-indol-3-yl-acetyl)-methylamino]-*N*-(4-isopropylphenyl)-2-phenylacetamide); felodipine (Ethyl methyl 4-

(2,3-dichlorophenyl)-2,6-dimethyl-1,4-dihydro-3,5-pyridinedicarboxylate); sulfonamide SF-01 (6-(ethylphenylsulfamoyl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid cycloheptylamide); and UCCF-152 (3-[2-(benzyloxy)phenyl]-5-(chloromethyl)isoxazole).

In one embodiment, the additional therapeutic agent is selected from a compound disclosed in Published Interanational Patent Application No. WO2014078842 or WO2012158885, or in United States Published Application No. US20140073667; all published international patent applications and published US patent application are hereby incorporated in their entirety by reference.

In one embodiment, the additional therapeutic agent is selected from a compound disclosed in Published International Patent Application No: WO2012078909; WO2012154880; WO2012154888; WO2012154967; WO2013112651; WO2013112699; or WO2012078902; all published international patent applications are hereby incorporated in their entirety by reference.

In one embodiment, the additional therapeutic agent is selected from a compound disclosed in US Patent No. 8,247,436 or 8,476,269; US Patent Application No. 13/923,349; and Published International Patent Application No: WO2011113894, WO2013038373, WO2013038378, WO2013038381, WO2013038386, or WO2013038390; all US patents, US patent applications, and published international patent applications are hereby incorporated in their entirety by reference.

In one embodiment, the additional therapeutic agent is slected from a compound disclosed in WO2014210159 or WO2014152213; all published international patent applications are hereby incorporated in their entirety by reference.

In one embodiment, the additional therapeutic agent is selected from a compound disclosed in Published International Patent Application No.: WO2014210159 or WO2014152213; all published international patent applications are hereby incorporated in their entirety by reference.

In one embodiment, the additional therapeutic agent is selected from a compound disclosed in Published International Patent Application No.: WO2015003083, WO2015007517, WO2015007519, or WO2015007516; all published international patent applications are hereby incorporated in their entirety by reference.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; and c) a compound disclosed in US Patent No. 8,247,436 or 8,476,269; US Patent Application No. 13/923,349; and Published International Patent Application No: WO2011113894, WO2013038373, WO2013038378, WO2013038381, WO2013038386, or WO2013038390; all US patents, US patent applications, and published international patent applications are hereby incorporated in their entirety by reference.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; and c) a compound disclosed in US Patent No. 8,247,436 or 8,476,269; US Patent Application No. 13/923,349; and Published International Patent Application No: WO2011113894, WO2013038373, WO2013038378, WO2013038381, WO2013038386, or WO2013038390; all US patents, US patent applications, and published international patent applications are hereby incorporated in their entirety by reference.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) a compound disclosed in US Patent No. 8,247,436 or 8,476,269; US Patent Application No. 13/923,349; and Published International Patent Application No: WO2011113894, WO2013038373, WO2013038378, WO2013038381, WO2013038386, or WO2013038390; all US patents, US patent applications, and published international patent applications are hereby incorporated in their entirety by reference.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) a compound disclosed in US Patent No. 8,247,436 or 8,476,269; US Patent Application No. 13/923,349; and Published International Patent Application No: WO2011113894, WO2013038373, WO2013038378, WO2013038381, WO2013038386, or WO2013038390; all US patents, US patent applications, and published

international patent applications are hereby incorporated in their entirety by reference; and c) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; c) a compound disclosed in US Patent No. 8,247,436 or 8,476,269; US Patent Application No. 13/923,349; and Published International Patent Application No: WO2011113894, WO2013038373, WO2013038378, WO2013038381, WO2013038386, or WO2013038390; all US patents, US patent applications, and published international patent applications are hereby incorporated in their entirety by reference; and d) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; c) a compound disclosed in US Patent No. 8,247,436 or 8,476,269; US Patent Application No. 13/923,349; and Published International Patent Application No: WO2011113894, WO2013038373, WO2013038378, WO2013038381, WO2013038386, or WO2013038390; all US patents, US patent applications, and published international patent applications are hereby incorporated in their entirety by reference; and d) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; c) a compound disclosed in US Patent No. 8,247,436 or 8,476,269; US Patent Application No. 13/923,349; and Published International Patent Application No: WO2011113894, WO2013038373, WO2013038378, WO2013038381, WO2013038386, or WO2013038390; all US patents, US patent applications, and published international patent applications are hereby incorporated in their entirety by reference; and d) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; and c) a compound disclosed in

Published International Patent Application No: WO2012078909; WO2012154880; WO2012154888; WO2012154967; WO2013112651; WO2013112699; or WO2012078902; all published international patent applications are hereby incorporated in their entirety by reference.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; and c) a compound disclosed in Published International Patent Application No: WO2012078909; WO2012154880; WO2012154888; WO2012154967; WO2013112651; WO2013112699; or WO2012078902; all published international patent applications are hereby incorporated in their entirety by reference.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) a compound disclosed in Published International Patent Application No: WO2012078909; WO2012154880; WO2012154888; WO2012154967; WO2013112651; WO2013112699; or WO2012078902; all published international patent applications are hereby incorporated in their entirety by reference.


In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; c) a compound disclosed in Published International Patent Application No: WO2012078909; WO2012154880; WO2012154888; WO2012154967; WO2013112651; WO2013112699; or WO2012078902; all published international patent applications are hereby incorporated in their entirety by reference; and d) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; c) a compound disclosed in Published International Patent Application No: WO2012078909; WO2012154880; WO2012154888; WO2012154967; WO2013112651; WO2013112699; or WO2012078902; all published international patent


applications are hereby incorporated in their entirety by reference; and d) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; c) a compound disclosed in Published International Patent Application No: WO2012078909; WO2012154880; WO2012154888; WO2012154967; WO2013112651; WO2013112699; or WO2012078902; all published international patent applications are hereby incorporated in their entirety by reference; and d) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.


In another embodiment, the additional therapeutic agent is a compound disclosed in Published International Patent Application No. WO2014099673 hereby incorporated in its entirety by reference, including but not limited to the following compounds:




P-1;




P-2;




P-3;




P-4;




P-5;




P-6;




P-7;




P-8;




P-9;




P-10;




P-11;




P-12;




P-13;



P-14;



P-15; or



P-16.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of fonnula I; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; and c) a compound selected from P-1 to P-16 disclosed above.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; and c) a compound selected from P-1 to P-16 disclosed above.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; and c) a compound selected from P-1 to P-16 disclosed above.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-

dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; and c) a compound selected from P-1 to P-16 disclosed above.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) a compound selected from P-1 to P-16 disclosed above.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and c) a compound selected from P-1 to P-16 disclosed above.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; c) a compound selected from P-1 to P-16 disclosed above; and d) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; c) a compound selected from P-1 to P-16 disclosed above; and d) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.


In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; c) a compound selected from P-1 to P-16 disclosed above; and d) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; c) a compound selected from P-1 to

P-16 disclosed above; and d) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In another embodiment, the above recited pharmaceutical compositions contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In some embodiments, the invention features a pharmaceutical composition comprising a) a compound of formula I; and b) a compound disclosed in United States Patent No. 8,865,902. In some embodiments, the invention features a pharmaceutical composition comprising a) a compound of Table 1; and b) a compound disclosed in United States Patent No. 8,865,902. In another embodiment, the compound disclosed in United States Patent No. 8,865,902 has the following structure:



wherein the compound disclosed in United States Patent No. 8,865,902 is any one of the compounds of the following table:

X^1	X^2	X^3	X^4	X^5	X^6	X^7	Y^1	Y^2	Y^3	Y^4	Y^5	Y^6
D	D	D	D	D	D	D	CD_3	CD_3	CD_3	CD_3	CD_3	CD_3
H	H	H	H	D	H	H	CD_3	CD_3	CD_3	CD_3	CD_3	CD_3
H	H	H	H	D	H	H	CD_3	CD_3	CD_3	CH_3	CH_3	CH_3
H	H	H	H	D	H	H	CH_3	CH_3	CH_3	CD_3	CD_3	CD_3
H	H	H	H	D	H	H	CH_3	CH_3	CH_3	CH_3	CH_3	CH_3
H	H	H	H	H	H	H	CD_3	CD_3	CD_3	CD_3	CD_3	CD_3
H	H	H	H	H	H	H	CD_3	CD_3	CD_3	CH_3	CH_3	CH_3
H	H	H	H	H	H	H	CH_3	CH_3	CH_3	CD_3	CD_3	CD_3

.

In another embodiment, the compound disclosed in United States Patent No. 8,865,902 is any one of the compounds of the following table:

X^1	X^2	X^3	X^4	X^5	X^6	X^7	Y^1	Y^2	Y^3	Y^4	Y^5	Y^6
H	H	H	H	H	D	D	CD_3	CD_3	CD_3	CD_3	CD_3	CD_3
H	H	H	H	D	D	D	CD_3	CD_3	CD_3	CD_3	CD_3	CD_3
H	H	H	H	D	D	D	CD_3	CD_3	CD_3	CH_3	CH_3	CH_3
H	H	H	H	D	D	D	CH_3	CH_3	CH_3	CD_3	CD_3	CD_3
H	H	H	H	D	D	D	CH_3	CH_3	CH_3	CH_3	CH_3	CH_3
H	H	H	H	H	D	D	CD_3	CD_3	CD_3	CH_3	CH_3	CH_3
H	H	H	H	H	D	D	CH_3	CH_3	CH_3	CD_3	CD_3	CD_3
D	D	D	D	D	D	D	CD_3	CD_3	CD_3	CH_3	CH_3	CH_3
D	D	D	D	D	D	D	CH_3	CH_3	CH_3	CD_3	CD_3	CD_3
D	D	D	D	D	H	H	CD_3	CD_3	CD_3	CH_3	CH_3	CH_3
D	D	D	D	D	H	H	CH_3	CH_3	CH_3	CD_3	CD_3	CD_3
H	H	H	H	H	H	D	CD_3	CD_3	CD_3	CD_3	CD_3	CD_3
H	H	H	H	H	H	D	CD_3	CD_3	CD_3	CH_3	CH_3	CH_3
H	H	H	H	H	H	D	CH_3	CH_3	CH_3	CD_3	CD_3	CD_3
H	H	H	H	H	D	H	CD_3	CD_3	CD_3	CD_3	CD_3	CD_3
H	H	H	H	H	D	H	CD_3	CD_3	CD_3	CH_3	CH_3	CH_3
H	H	H	H	H	D	H	CH_3	CH_3	CH_3	CD_3	CD_3	CD_3


.

In another embodiment, the compound disclosed in United States Patent No. 8,865,902 is any one of the compounds of the following table:


X^1	X^2	X^3	X^4	X^5	X^6	X^7	Y^1	Y^2	Y^3	Y^4	Y^5	Y^6
D	D	D	D	D	H	H	CD_3	CD_3	CD_3	CD_3	CD_3	CD_3
D	D	D	D	D	D	D	CH_3	CH_3	CH_3	CH_3	CH_3	CH_3
D	D	D	D	D	H	H	CH_3	CH_3	CH_3	CH_3	CH_3	CH_3
D	D	D	D	H	D	D	CD_3	CD_3	CD_3	CD_3	CD_3	CD_3
D	D	D	D	H	H	H	CD_3	CD_3	CD_3	CD_3	CD_3	CD_3
D	D	D	D	H	D	D	CH_3	CH_3	CH_3	CH_3	CH_3	CH_3
D	D	D	D	H	H	H	CH_3	CH_3	CH_3	CH_3	CH_3	CH_3
H	H	H	H	H	D	D	CH_3	CH_3	CH_3	CH_3	CH_3	CH_3

.


In some embodiments, the invention features a pharmaceutical composition comprising a) a compound of formula I; and b) the compound:




In some embodiments, the invention features a pharmaceutical composition comprising a) a compound of Table 1; and b) the compound:




In some embodiments, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; and c) the compound:




In some embodiments, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; and c) the compound:




In some embodiments, the invention features a pharmaceutical composition comprising a) a compound selected from Compound 233, Compound 238, Compound 318, Compound 228, Compound 196, Compound 337, Compound 368, Compound 182, Compound 1356, Compound 172, Compound 639, Compound 300, Compound 1660, Compound 234, Compound 1975, Compound 262, Compound 2191, Compound 207, Compound 320, Compound 353, Compound 171, Compound 114, Compound 321, Compound 15, Compound 265, Compound 164, Compound 214, Compound 345, Compound 8, Compound 110, Compound 281, Compound 351, Compound 197, Compound 136, Compound 307, Compound 279, Compound 72, Compound 125, Compound 45, Compound 299, Compound 365, or Compound 361; b) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; and c) the compound:




In some embodiments, the invention features a pharmaceutical composition comprising a) a compound of formula I; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; and c) the compound:



In some embodiments, the invention features a pharmaceutical composition comprising a) a compound of Table 1; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; and c) the compound:



In some embodiments, the invention features a pharmaceutical composition comprising a) a compound selected from Compound 233, Compound 238, Compound 318, Compound 228, Compound 196, Compound 337, Compound 368, Compound 182, Compound 1356, Compound 172, Compound 639, Compound 300, Compound 1660, Compound 234, Compound 1975, Compound 262, Compound 2191, Compound 207, Compound 320, Compound 353, Compound 171, Compound 114, Compound 321, Compound 15, Compound 265, Compound 164, Compound 214, Compound 345, Compound 8, Compound 110, Compound 281, Compound 351, Compound 197, Compound 136, Compound 307, Compound 279, Compound 72, Compound 125, Compound 45, Compound 299, Compound 365, or Compound 361; b) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; and c) the compound:



In another embodiment, the above recited pharmaceutical compositions contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In one embodiment, any of the below therapeutic agents can be used alone with a compound of formula I or Id, or in combination with any of the above described pharmaceutical compositions, or as a component in any of the above described pharmaceutical compositions.

In one embodiment, the additional therapeutic agent is selected from a mucolytic agent, bronchodialator, an antibiotic, an anti-infective agent, a CFTR modulator, or an anti-inflammatory agent.

In one embodiment, any of the below therapeutic agents, which predominantly treat the symptoms a CFTR mediated disease, such as cystic fibrosis, rather than its underlying cause, can be used alone with a compound of formula I, or in combination with any of the above described pharmaceutical compositions, or as a component in any of the above described pharmaceutical compositions.

In one embodiment, the additional therapeutic agent is an antibiotic. Exemplary antibiotics useful herein include tobramycin, including tobramycin inhaled powder (TIP), azithromycin, cayston, aztreonam, including the aerosolized form of aztreonam, amikacin, including liposomal formulations thereof, ciprofloxacin, including formulations thereof suitable for administration by inhalation, levoflaxacin, including aerosolized formulations thereof, and combinations of two antibiotics, e.g., fosfomycin and tobramycin.

In another embodiment, the additional therapeutic agent is a mucolyte. Exemplary mucolytes useful herein includes Pulmozyme®.

In another embodiment, the additional therapeutic agent is a bronchodialator. Exemplary bronchodialtors include albuterol, metaprotenerol sulfate, pirbuterol acetate, salmeterol, or tetrabuline sulfate.

In another embodiment, the additional therapeutic agent is effective in restoring lung airway surface liquid. Such agents improve the movement of salt in and out of cells, allowing mucus in the lung airway to be more hydrated and, therefore, cleared more easily. Exemplary such agents include hypertonic saline, denufosol tetrasodium ([[(3S,5R)-5-(4-amino-2-oxopyrimidin-1-yl)-3-hydroxyoxolan-2-yl]methoxy-hydroxyphosphoryl] [[[(2R,3S,4R,5R)-5-(2,4-dioxopyrimidin-1-yl)-3, 4-dihydroxyoxolan-2-yl]methoxy-hydroxyphosphoryl]oxy-hydroxyphosphoryl] hydrogen phosphate), or bronchitol (inhaled formulation of mannitol).

In another embodiment, the additional therapeutic agent is an anti-inflammatory agent, i.e., an agent that can reduce the inflammation in the lungs. Exemplary such agents useful herein include ibuprofen, docosahexanoic acid (DHA), sildenafil, inhaled glutathione, pioglitazone, hydroxychloroquine, or simavastatin.

In another embodiment, the additional therapeutic agent is a compound that augments or induces CFTR activity other than a compound of formula I. Exemplary such agents include ataluren ("PTC124®"; 3-[5-(2-fluorophenyl)-1,2,4-oxadiazol-3-yl]benzoic acid), sinapultide, lancovutide, depelestat (a human recombinant neutrophil elastase inhibitor), and cobiprostone (7-{(2R, 4aR, 5R, 7aR)-2-[(3S)-1,1-difluoro-3-methylpentyl]-2-hydroxy-6-oxooctahydrocyclopenta[b]pyran-5-yl}heptanoic acid).

In another embodiment, the additional therapeutic agent is a nutritional agent. Exemplary nutritional agents include pancrelipase (pancreating enzyme replacement), including Pancrease®, Pancreacarb®, Ultrase®, or Creon®, Liprotomase® (formerly Trizytek®), Aquadeks®, or glutathione inhalation. In one embodiment, the additional nutritional agent is pancrelipase.

In another embodiment, the additional therapeutic agent is a compound selected from gentamicin, curcumin, cyclophosphamide, 4-phenylbutyrate, miglustat, felodipine, nimodipine, Philoxin B, geniestein, Apigenin, cAMP/cGMP augmenters or inducers such as rolipram, sildenafil, milrinone, tadalafil, amrinone, isoproterenol, albuterol, and almeterol, deoxyspergualin, HSP 90 inhibitors, HSP 70 inhibitors, proteosome inhibitors such as epoxomicin, lactacystin, etc.

In other embodiments, the additional therapeutic agent is a compound disclosed in WO 2004028480, WO 2004110352, WO 2005094374, WO 2005120497, or WO 2006101740,

incorporated herein in their entirety by reference. In another embodiment, the additional agent is a benzo[c]quinolizinium derivative that exhibits CFTR inducing or augmenting activity or a benzopyran derivative that exhibits CFTR inducing or augmenting activity. In another embodiment, the additional agent is a compound disclosed in U.S. Pat. No. 7,202,262, U.S. Pat. No. 6,992,096, US20060148864, US20060148863, US20060035943, US20050164973, WO2006110483, WO2006044456, WO2006044682, WO2006044505, WO2006044503, WO2006044502, or WO2004091502, incorporated herein in their entirety by reference. In another embodiment, the additional agent is a compound disclosed in WO2004080972, WO2004111014, WO2005035514, WO2005049018, WO2006099256, WO2006127588, or WO2007044560, incorporated herein in their entirety by reference.

In another embodiment, the additional therapeutic agent is a compound disclosed in WO2014180562 or US20140274933, incorporated herein in their entirety by reference.

In another embodiment, the additional therapeutic agent is selected from the categories ENaC inhibitors, betamimetics, anticholinergics, corticosteroids, PDE4-inhibitors, LTD4-antagonists, EGFR-inhibitors, dopamine agonists, H1-antihistamines, PAF-antagonists, MAP-kinase inhibitors, MPR4-Inhibitors, iNOS-Inhibitors, or SYK-Inhibitors, or double or triple combinations thereof.

In another embodiment, the additional therapeutic agent is an ENaC inhibitor selected from 3-amino-6-(4-fluoro-phenyl)-5-trifluoromethyl-pyridine-2-carboxylic acid (3,3,3-trifluoro-2-hydroxy -2-methyl-propyl)-amide;

5-amino-6′-methyl-3-trifluoromethyl-[2,3]bipyridinyl-6-carboxylic acid (3,3,3-trifluoro-2-hydroxy -2-methyl-propyl)-amide;

3-amino-6-cyclopropyl-*N*-(3,3,3-trifluoro-2-hydroxy-2-methylpropyl)-5-(trifluoromethyl)picolinamide; 3-amino-6-methoxy-*N*-(3,3,3-trifluoro-2-hydroxy-2-(trifluoromethyl)propyl)-5-(trifluoro methyl)picolinamide;

3-amino-6-(4-fluoro-phenyl)-5-trifluoromethyl-pyridine-2-carboxylic acid ((S)-3,3,3-trifluoro-2-hydroxy -2-methyl-propyl)-amide;

3-amino-6-methoxy-5-trifluoromethyl-pyridine-2-carboxylic acid((S-3,3,3-trifluoro-2-hydroxy-2-methyl-propyl)-amide;

3-amino-6-methoxy-5-trifluoromethyl-pyridine-2-carboxylic acid ((R)-3,3,3-trifluoro-2-hydroxy-2-methyl-propyl)-amide;

3-amino-6-(2,4-dichloro-phenyl)-5-trifluoromethyl-pyridine-2-carboxylic acid ((S)-3,3,3-trifluoro-2-hydroxy-2-methyl-propyl)-amide;

3-amino-6-(2,4-dichloro-phenyl)-5-trifluoromethyl-pyridine-2-carboxylic acid ((R)-3,3,3-trifluoro -2-hydroxy-2-methyl-propyl)-amide;

3-amino-6-(4-fluoro-phenyl)-5-trifluoromethyl-pyridine-2-carboxylic acid (2-hydroxy-2-methyl-propyl)-amide;

3-amino-5,6-bis-trifluoromethyl-pyridine-2-carboxylic acid ((S)-3,3,3-trifluoro-2-hydroxy-2-methyl-propyl)-amide;

3-amino-5,6-bis-trifluoromethyl-pyridine-2-carboxylic acid ((R)-3,3,3-trifluoro-2-hydroxy-2-methyl-propyl)-amide;

(S)-3-amino-6-ethoxy-*N*-(3,3,3-trifluoro-2-hydroxy-2-methylpropyl)-5-(trifluoro methyl)picolinamide;

3-amino-6-methoxy-5-trifluoromethyl-pyridine-2-carboxylic acid ((S)-3,3,3-trifluoro -2-hydroxy-2-methyl-propyl)-amide;

3-amino-6-methoxy-5-trifluoromethyl-pyridine-2-carboxylic acid ((R)-3,3,3-trifluoro -2-hydroxy-2-methyl-propyl)-amide;

3-amino-6-(4-fluoro-phenyl)-5-trifluoromethyl-pyridine-2-carboxylic acid (3,3,3-trifluoro-2-hydroxy-2-methyl-propyl)-amide;

3-amino-5,6-bis-trifluoromethyl-pyridine-2-carboxylic acid ((S)-3,3,3-trifluoro -2-hydroxy-2-methyl-propyl)-amide;

3-amino-5,6-bis-trifluoromethyl-pyridine-2-carboxylic acid ((R)-3,3,3-trifluoro -2-hydroxy-2-methyl-propyl)-amide, or pharmaceutically acceptable salts thereof.

In another embodiment, the additional agent is a compound disclosed in United States Patent No. 8,247,436 and International PCT Publication WO 2011113894, incorporated herein in their entirety by reference.

In another embodiment, the additional therapeutic agent is a betamimetic selected from Albuterole, Arformoterole, Bambuterole, Bitolterole, Broxaterole, Carbuterole, Clenbuterole, Fenoterole, Formoterole, Hexoprenaline, Ibuterole, Isoetharine, Isoprenaline, Levosalbutamole, Mabuterole, Meluadrine, Metaproterenole, Milveterol, Orciprenaline, Pirbuterole, Procaterole, Reproterole, Rimiterole, Ritodrine, Salmefamole, Salmeterole, Soterenole, Sulphonterole, Terbutaline, Tiaramide, Tolubuterole, Zinterole, Nolomirole, and

1-(2-chloro-4-hydroxyphenyl)-t-butylaminoethanole;

(-)-2-[7(S)-[2(R)-Hydroxy-2-(4-hydroxyphenyl)-ethylamino]-5,6,7,8-tetrahydro-2-naphthyloxy]-N,*N*-dimethylacetamide hydrochloride monohydrate;

3-(4-{6-[2-Hydroxy-2-(4-hydroxy-3-hydroxymethyl-phenyl)-ethylamino]-hexyl-oxy}-butyl)-benzyl-sulfonamide;

5-[2-(5,6-Diethyl-indan-2-ylamino)-1-hydroxy-ethyl]-8-hydroxy-1H-quinoline-2-one;

4-Hydroxy-7-{2-{[2-{[3-(2-phenylethoxy)propyl]sulphonyl}ethyl]-amino}ethy-l]-2(3H)-benzothiazolone;

1-(2-Fluoro-4-hydroxyphenyl)-2-[4-(1-benzimidazolyl)-2-methyl-2-butylamino]ethanol;

1-[3-(4-Methoxybenzyl-amino)-4-hydroxyphenyl]-2-[4-(1-benzimidazolyl)-2-methyl-2-butylamino]ethanol;

1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-N,*N*-dimethylaminophenyl)-2-methyl-2-propylamino]ethanol;

1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-methoxyphenyl)-2-methyl-2-propylamino]ethanol;

1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-n-butyloxyphenyl)-2-methyl-2-propylamino]ethanol;

1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-{4-[3-(4-methoxyphenyl)-1,2,4-triazol-3-yl]-2-methyl-2-butylamino}ethanol;

5-Hydroxy-8-(1-hydroxy-2-isopropylaminobutyl)-2H-1,4-benzoxazin-3-(4H)-one;

1-(4-Amino-3-chloro-5-trifluormethylphenyl)-2-tert.-butylamino)ethanol;

6-Hydroxy-8-{1-hydroxy-2-[2-(4-methoxy-phenyl)-1,1-dimethyl-ethylamino]-ethyl}-4H-benzo[1,4]oxazin-3-one;

6-Hydroxy-8-{1-hydroxy-2-[2-(4-phenoxy-acetic acid ethylester)-1,1-dimethyl-ethylamino]-ethyl}-4H-benzo[1,4]oxazin-3-one;

6-Hydroxy-8-{1-hydroxy-2-[2-(4-phenoxy-acetic acid)-1,1-dimethyl-ethylamino]-ethyl}-4H-benzo[1,4]oxazin-3-one;

8-{2-[1,1-Dimethyl-2-(2,4,6-trimethylphenyl)-ethylamino]-1-hydroxy-ethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one;

6-Hydroxy-8-{1-hydroxy-2-[2-(4-hydroxy-phenyl)-1,1-dimethyl-ethylamino]-e-thyl}-4H-benzo[1,4]oxazin-3-one;

6-Hydroxy-8-{1-hydroxy-2-[2-(4-isopropyl-phenyl)-1,1dimethyl-ethylamino]-ethyl}-4H-benzo[1,4]oxazin-3-one;

8-{2-[2-(4-Ethyl-phenyl)-1,1-dimethyl-ethylamino]-1-hydroxy-ethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one;

8-{2-[2-(4-Ethoxy-phenyl)-1,1-dimethyl-ethylamino]-1-hydroxy-ethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one;

4-(4-{2-[2-Hydroxy-2-(6-hydroxy-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-8-yl)-ethylamino]-2-methyl-propyl}-phenoxy)-butyric acid;

8-{2-[2-(3,4-Difluor-phenyl)-1,1-dimethyl-ethylamino]-1-hydroxy-ethyl}-6-hydroxy-4H-benzo[1,4]oxazin-3-one;

1-(4-Ethoxy-carbonylamino-3-cyano-5-fluorophenyl)-2-(tert-butylamino)ethanol;

N-[2-Hydroxy-5-(1-hydroxy-2-{2-[4-(2-hydroxy-2-phenyl-ethylamino)-phenyl]-ethylamino}-ethyl)-phenyl]-formamide;

8-Hydroxy-5-(1-hydroxy-2-{2-[4-(6-methoxy-biphenyl-3-ylamino)-phenyl]-ethylamino}-ethyl)-1H-quinolin-2-one;

8-Hydroxy-5-[1-hydroxy-2-(6-phenethylamino-hexylamino)-ethyl]-1H-quinolin-2-one;

5-[2-(2-{4-[4-(2-Amino-2-methyl-propoxy)-phenylamino]-phenyl}-ethylamino)-1-hydroxy-ethyl]-8-hydroxy-1H-quinolin-2-one;

[3-(4-{6-[2-Hydroxy-2-(4-hydroxy-3-hydroxymethyl-phenyl)-ethylamino]hexyl-oxy}-butyl)-5-methyl-phenyl]-urea;

4-(2-{6-[2-(2,6-Dichloro-benzyloxy)-ethoxy]-hexylamino}-1-hydroxy-ethyl)-2-hydroxymethyl-phenole;

3-(4-{6-[2-Hydroxy-2-(4-hydroxy-3-hydroxymethyl-phenyl)-ethylamino]-hexyl-oxy}-butyl)-benzenesulfonamide;

3-(3-{7-[2-Hydroxy-2-(4-hydroxy-3-hydroxymethyl-phenyl)-ethylamino]-hepty-loxy}-propyl)-benzenesulfonamide;

4-(2-{6-[4-(3-Cyclopentanesulfonyl-phenyl)-butoxy]-hexylamino}-1-hydroxy-ethyl)-2-hydroxymethyl-phenole;

N-Adamantan-2-yl-2-(3-{2-[2-hydroxy-2-(4-hydroxy-3-hydroxymethyl-phenyl)-ethylamino]-propyl}-phenyl)-acetamide;

(R,S)-4-(2-{[6-(2,2-Difluoro-4-phenylbutoxy)hexyl]amino}-1-hydroxy-ethyl)-2-(hydroxymethyl)phenol;

(R,S)-4-(2-{[6-(2,2-Difluoro-2-phenylethoxy)hexyl]amino}-1-hydroxy-ethyl)-2-(hydroxymethyl)phenol;

(R,S)-4-(2-{[4,4-Difluoro-6-(4-phenylbutoxy)hexyl]amino}-1-hydroxy-ethyl)-2-(hydroxy-methyl)phenol;

(R,S)-4-(2-{[6-(4,4-Difluoro-4-phenylbutoxy)hexyl]amino}-1-hydroxy-ethyl)-2-(hydroxymethyl)phenol;

(R,S)-5-(2-{[6-(2,2-Difluoro-2-phenylethoxy)hexyl]amino}-1-hydroxy-ethyl)-8-hydroxyquinolin-2(1H)-one;

(R,S)-[2-({6-[2,2-Difluoro-2-(3-methylphenyl)ethoxy]hexyl}amino)-1-hydroxyethyl]-2-(hydroxymethyl)phenol;

4-(1R)-2-{[6-(2,2-Difluoro-2-phenylethoxy)hexyl]amino}-1-hydroxyethyl)-2-(hydroxymethyl)phenol;

(R,S)-2-(Hydroxymethyl)-4-(1-hydroxy-2-{[4,4,5l5-tetrafluoro-6-(3-phenylpropoxy)-hexyl]amino}ethyl)phenol;

(R,S)-[5-(2-{[6-(2,2-Difluoro-2-phenylethoxy)hexyl]amino}-1-hydroxy-ethyl-)-2-hydroxy-phenyl]formamide;

(R,S)-4-[2-({6-[2-(3-Bromophenyl)-2,2-difluoroethoxy]hexyl}amino)-1-hydroxyethyl]-2-(hydroxymethyl)phenol;

(R,S)-*N*-[3-(1,1-Difluoro-2-{[6-({2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)-phenyl]-ethyl}amino)hexyl]oxy}ethyl)phenyl]urea;

3-[3-(1,1-difluoro-2-{[6-({2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]ethyl}-amino)hexyl]oxy}ethyl)phenyl]imidazolidine-2,4-dione;

(R,S)-4-[2-({6-[2,2-difluoro-2-(3-methoxyphenyl)ethoxy]hexyl}amino)-1-hydroxyethyl]-2-(hydroxymethyl)phenol;

5-((1R)-2-{[6-(2,2-difluoro-2-phenylethoxy)hexyl]amino}-1-hydroxyethyl)-8-hydroxyquinolin-2(1H)-one;

4-((1R)-2-{[4,4-Difluoro-6-(4-phenylbutoxy)hexyl]amino}-1-hydroxy-ethyl)-2-(hydroxymethyl)phenol;

(R,S)-4-(2-{[6-(3,3-Difluoro-3-phenylpropoxy)hexyl]amino}-1-hydroxy-ethyl-)-2-(hydroxymethyl)phenol;

(R,S)-(2-{[6-(2,2-Difluoro-2-phenylethoxy)-4,4-difluorohexyl]amino}-1-hydroxyethyl)-2-(hydroxymethyl)phenol;

(R,S)-4-(2-{[6-(2,2-difluoro-3-phenylpropoxy)hexyl]amino}-1-hydroxyethyl)-2-(hydroxy-methyl)phenol;

3-[2-(3-Chloro-phenyl)-ethoxy]-*N*-(2-diethylamino-ethyl)-*N*-{2-[2-(4-hydroxy-2-oxo-2,3-dihydro-benzothiazol-7-yl)-ethylamino]-ethyl}-propionamide;

N-(2-Diethylamino-ethyl)-*N*-{2-[2-(4-hydroxy-2-oxo-2,3-dihydro-benzothiazol-7-yl)-ethylamino]-ethyl}-3-(2-naphthalen-1-yl-ethoxy)-propionamide;

7-[2-(2-{3-[2-(2-Chloro-phenyl)-ethylamino]-propylsulfanyl}-ethylamino)-1-hydroxy-ethyl]-4-hydroxy-3H-benzothiazol-2-one; or

7-[(1R)-2-(2-{3-[2-(2-Chloro-phenyl)-ethylamino]-propylsulfanyl}-ethylamino)-1-hydroxyethyl]-4-hydroxy-3H-benzothiazol-2-one; optionally in racemic form, as enantiomers, diastereomeres or as pharmaceutically acceptable salts, solvates or hydrates. Preferred are salts selected from the group consisting of hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethansulfonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydrooxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulfonate.

In one embodiment, the additional therapeutic agent is an anticholinergic selected from Tiotropium salts, preferably the bromide salt, Oxitropium salts, preferably the bromide salt, Flutropium salts, preferably the bromide salt, Ipratropium salts, preferably the bromide salt, Aclidinium salts, preferably the bromide salt, Glycopyrronium salts, preferably the bromide salt, Trospium salts, preferably the chloride salt, Tolterodin. From the above mentioned salts the pharmaceutically active part is the cation, possible anions are chloride, bromide, iodide, sulfate, phosphate, methansulfonate, nitrate, maleate, acetate, citrate, fumarate, tartrate, oxalate, succinate, benzoate or p-toluenesulfonate. Further examples of preferred anticholinergics are selected from among

2,2-Diphenylpropionic acid tropenole ester-methobromide;

2,2-Diphenylpropionic acid scopine ester-methobromide;

2-Fluor-2,2-Diphenylacetic acid scopine ester-methobromide;

2-Fluor-2,2-Diphenylacetic acid tropenole ester-methobromide;

3,3',4,4'-Tetrafluorbenzil acid tropenole ester-methobromide;

3,3',4,4'-Tetrafluorbenzil acid scopine ester-methobromide;

4,4'-Difluorbenzil acid tropenole ester-methobromide;

4,4'-Difluorbenzil acid scopine ester-methobromide;

3,3'-Difluorbenzil acid tropenole ester-methobromide;

3,3'-Difluorbenzil acid scopine ester-methobromide;

9-Hydroxy-fluorene-9-carbon acid tropenole ester-methobromide;

9-Fluor-fluorene-9-carbon acid tropenole ester-methobromide;

9-Hydroxy-fluorene-9-carbon acid scopine ester-methobromide;

9-Fluor-fluorene-9-carbon acid scopine ester methobromide;

9-Methyl-fluorene-9-carbon acid tropenole estermethobromide;

9-Methyl-fluorene-9-carbon acid scopine estermethobromide;

Benzil acid cyclopropyl tropine ester-methobromide;

2,2-Diphenylpropionic acid cyclopropyl tropine ester-methobromide;

9-Hydroxy-xanthene-9-carbon acid cyclopropyl tropine ester-methobromide;

9-Methyl-fluorene-9-carbon acid cyclopropyl tropine ester-methobromide;

9-Methyl-xanthene-9-carbon acid cyclopropyl tropine ester-methobromide;

9-Hydroxy-fluorene-9-carbon acid cyclopropyl tropine ester-methobromide;

4,4'-Difluorbenzil acid methylester cyclopropyl tropine ester-methobromide;

9-Hydroxy-xanthene-9-carbon acid tropenole ester-methobromide;

9-Hydroxy-xanthene-9-carbon acid scopine ester methobromide;

9-Methyl-xanthene-9-carbon acid tropenole ester-methobromide;

9-Methyl-xanthene-9-carbon acid scopine estermethobromide;

9-Ethyl-xanthene-9-carbon acid tropenole ester methobromide;

9-Difluormethyl-xanthene-9-carbon acid tropenole ester-methobromide; or

9-Hydroxymethyl-xanthene-9-carbon acid scopine ester methobromide.

In one embodiment, the additional therapeutic agent is a corticosteroid selected from Beclomethasone, Betamethasone, Budesonide, Butixocorte, Ciclesonide, Deflazacorte, Dexamethasone, Etiprednole, Flunisolide, Fluticasone, Loteprednole, Mometasone, Prednisolone, Prednisone, Rofleponide, Triamcinolone, Tipredane,

{20R-16alpha,17alpha-[butylidenebis(oxy)]-6alpha,9alpha-difluoro-11beta-hydroxy-17beta-(methylthio)androsta-4-en-3-one};

9-fluoro-11beta,17,21-trihydroxy-16alpha-methylpregna-1,4-diene-3,20-dione 21-cyclohexanecarboxylate 17-cyclopropanecarboxylate;

16,17-butylidenedioxy-6,9-difluoro-11-hydroxy-17-(methylthio)androst-4-en-3-one;

Flunisolide-21-[4'-(nitrooxymethyl)benzoate];

6,9-Difluoro-17-[(2-furanylcarbonyl)oxy]-11-hydroxy-16-methyl-3-oxo-androsta-1,4-dien-17-carbothion acid (S)-fluoromethylester;

6,9-Difluoro-11-hydroxy-16-methyl-3-oxo-17-propionyloxy-androsta-1,4-dien-17-carbothion acid (S)-(2-oxo-tetrahydro-furan-3S-yl)ester; or

6alpha,9alpha-difluoro-11beta-hydroxy-16alpha-methyl-3-oxo-17alpha-(2,2,3,3-tertamethylcyclopropylcarbonyl)oxy-androsta-1,4-diene-17beta-carboxylic acid cyanomethyl ester; optionally in racemic form, as enantiomers, diastereomeres or as pharmaceutically acceptable salts, solvates or hydrates. Examples for preferred salts and derivatives are alkali salts, i.e. sodium or potassium salts, sulfobenzoates, phosphates, isonicotinates, acetates, dichloroacetates, propionates, dihydrogenphosphates, palmitates, pivalates or furoates.

In one embodiment, the additional therapeutic agent is a PDE4-inhibitor selected from Enprofylline, Theophylline, Roflumilaste, Ariflo (Cilomilaste), Tofimilaste, Pumafentrine, Lirimilaste, Apremilaste, Arofylline, Atizorame, Oglemilastum, Tetomilaste;

5-[(*N*-(2,5-dichloro-3-pyridinyl)-carboxamide]-8-methoxy-quinoline;

5-[*N*-(3,5-dichloro-1-oxido-4-pyridinyl)-carboxamide]-8-methoxy-2-(trifluoromethyl)-quinoline;

N-(3,5-dichloropyrid-4-yl)-[1-(4-fluorobenzyl)-5-hydroxy-indole-3-yl]glyoxyl acid amide);

9-[(2-fluorophenyl)methyl]-*N*-methyl-2-(trifluoromethyl)-9H-purine-6-amine;

4-[(2R)-2-[3-(cyclopentyloxy)-4-methoxyphenyl]-2-phenylethyl]-pyridine;

N-[(3R)-3,4,6,7-tetrahydro-9-methyl-4-oxo-1-phenylpyrrolo[3,2,1-jk-][1,4]benzodiazepin-3-yl]-4-Pyridinecarboxamide;

4-[6,7-diethoxy-2,3-bis(hydroxymethyl)-1-naphthalenyl]-1-(2-methoxyethyl)-2(1H)-pyridinone;

2-[4-[6,7-diethoxy-2,3-bis(hydroxymethyl)-1-naphthalenyl]-2-pyridinyl]-4-(3-pyridinyl)-1 (2H)-Phthalazinone;

(3-(3-cyclopenyloxy-4-methoxybenzyl)-6-ethylamino-8-isopropyl-3H-purine;

beta-[3-(cyclopentyloxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isoindole-2-propanamide;

9-ethyl-2-methoxy-7-methyl-5-propyl-imidazo[1,5-a]pyrido[3,2-e]pyrazin-6(-5H)-one;

5-[3-(cyclopentyloxy)-4-methoxyphenyl]-3-[(3-methylphenyl)methyl](3S,5S)-2-piperidinone;

4-[1-[3,4-bis(difluoromethoxy)phenyl]-2-(3-methyl-1-oxido-4-pyridinyl)ethyl]-alpha,alpha-bis(trifluoromethyl)-Benzenemethanol;

N-(3,5-Dichloro-1-oxo-pyridine-4-yl)-4-difluoromethoxy-3-cyclopropylmethoxy-benzamide;

(-)p-[(4aR*,10bS*)-9-Ethoxy-1,2,3,4,4a,10b-hexahydro-8-methoxy-2-methyl-benzo[s][1,6]naphthyridin-6-yl]-N,*N*-diisopropylbenzamide;

(R)-(+)-1-(4-Bromobenzyl)-4-[(3-cyclopentyloxy)-4-methoxyphenyl]-2-pyrrolidone;

3-(Cyclopentyloxy-4-methoxyphenyl)-1-(4-N'-[*N*-2-cyano-S-methyl-isothioureido]-benzyl)-2-pyrrolidone;

cis[4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexan-1-carbon acid];

2-Carbomethoxy-4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)-cyclohexan-1-one;

cis[4-Cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-ol];

(R)-(+)-Ethyl[4-(3-cyclopentyloxy-4-methoxyphenyl)pyrrolidine-2-yliden]acetate;

(S)-(-)-Ethyl[4-(3-cyclopentyloxy-4-methoxyphenyl)pyrrolidine-2-yliden]acetate;

9-Cyclopentyl-5,6-dihydro-7-ethyl-3-(2-thienyl)-9H-pyrazolo[3,4-c]-1,2,4-triazolo[4,3-a]pyridine; or

9-Cyclopentyl-5,6-dihydro-7-ethyl-3-(tert-butyl)-9H-pyrazolo[3,4-c]-1,2,4-triazolo[4,3-a]pyridine; optionally in racemic form, as enantiomers, diastereomeres or as pharmaceutically acceptable salts, solvates or hydrates. Preferred are salts selected from the group consisting of hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethansulfonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydrooxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulfonate.

In one embodiment, the additional therapeutic agent is a LTD4-antagonist selected from Montelukast, Pranlukast, Zafirlukast, Masikulast, L-733321 (see compound 2ab of D. Guay et al, Bioorg. Med. Chem. Lett. 8 (1998) 453-458);

(E)-8-[2-[4-[4-(4-Fluorophenyl)butoxy]phenyl]ethenyl]-2-(1H-tetrazole-5-yl)-4H-1-benzopyran-4-one;

4-[6-Acetyl-3-[3-(4-acetyl-3-hydroxy-2-propylphenylthio)propoxy]-2-propyl-phenoxy]-butyric acid;

1-(((R)-(3-(2-(6,7-Difluoro-2-quinolinyl)ethenyl)phenyl)-3-(2-(2-hydroxy-2-propyl)phenyl)thio)methylcyclopropane-acetic acid;

1-(((1(R)-3(3-(2-(2,3-Dichlorothieno[3,2-b]pyridin-5-yl)-(E)-ethenyl)phenyl)-3-(2-(1-hydroxy-1-methylethyl)phenyl)propyl)thio)methyl)cyclopropane acetic acid; or

[2-[[2-(4-tert-Butyl-2-thiazolyl)-5-benzofuranyl]oxymethyl]phenyl]acetic acid; optionally in racemic form, as enantiomers, diastereomeres or as pharmaceutically acceptable salts, solvates or hydrates. Preferred are salts selected from the group consisting of hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethansulfonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydrooxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulfonate.

Further examples for optionally preferred salts and derivatives are alkali salts, i.e. sodium or potassium salts, sulfobenzoates, phosphates, isonicotinates, acetates, propionates, dihydrogenphosphates, palmitates, pivalates or furoates.

In one embodiment, the additional therapeutic agent is an EGFR-inhibitor selected from Cetuximab, Trastuzumab, Panitumumab Gefitinib, Canertinib, Erlotinib, Mab ICR-62;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{[4-(morpholine-4-yl)-1-oxo-2-butene-1-yl]amino}-7-cyclopropylmethoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{[4-(N,*N*-diethylamino)-1-oxo-2-butene-1-yl]amino}-7-cyclopropylmethoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{[4-(N,*N*-dimethylamino)-1-oxo-2-butene-1-yl]amino}-7-cyclopropylmethoxy-quinazoline;

4-[(R)-(1-Phenyl-ethyl)amino]-6-{[4-(morpholine-4-yl)-1-oxo-2-butene-1-yl-]amino}-7-cyclopentyloxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{[4-((R)-6-methyl-2-oxo-morpholine-4-yl)-1-oxo-2-butene-1-yl]amino}-7-cyclopropylmethoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{[4-((R)-6-methyl-2-oxo-morpholine-4-yl)-1-oxo-2-butene-1-yl]amino}-7-[(S)-(tetrahydrofuran-3-yl)oxy]-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{[4-((R)-2-methoxymethyl-6-oxo-morpholine-4-yl)-1-oxo-2-butene-1-yl]amino}-7-cyclopropylmethoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-[2-((S)-6-methyl-2-oxo-morpholine-4-yl)-ethoxy]-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-({4-[*N*-(2-methoxy-ethyl)-*N*-methyl-amino]-1-oxo-2-butene-1-yl}amino)-7-cyclopropylmethoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{[4-(N,*N*-dimethylamino)-1-oxo-2-butene-1-yl]amino}-7-cyclopentyloxy-quinazoline;

4-[(R)-(1-Phenyl-ethyl)amino]-6-{[4-(N,*N*-bis-(2-methoxy-ethyl)-amino)-1-oxo-2-butene-1-yl]amino}-7-cyclopropylmethoxy-quinazoline;

4-[(R)-(1-Phenyl-ethyl)amino]-6-({4-[*N*-(2-methoxy-ethyl)-*N*-ethyl-amino]-1-oxo-2-butene-1-yl}amino)-7-cyclopropylmethoxy-quinazoline;

4-[(R)-(1-Phenyl-ethyl)amino]-6-({4-[*N*-(2-methoxy-ethyl)-*N*-methyl-amino]-1-oxo-2-butene-1-yl}amino)-7-cyclopropylmethoxy-quinazoline;

4-[(R)-(1-Phenyl-ethyl)amino]-6-({4-[*N*-(tetrahydropyran-4-yl)-*N*-methyl-amino]-1-oxo-2-butene-1-yl}amino)-7-cyclopropylmethoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{[4-(N,*N*-dimethylamino)-1-oxo-2-butene-1-yl]amino}-7-((R)-tetrahydrofuran-3-yloxy)-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{[4-(N,*N*-dimethylamino)-1-oxo-2-butene-1-yl]amino}-7-((S)-tetrahydrofuran-3-yloxy)-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-({4-[*N*-(2-methoxy-ethyl)-*N*-methyl-amino]-1-oxo-2-butene-1-yl}amino)-7-cyclopentyloxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{[4-(*N*-cyclopropyl-*N*-methyl-amino)-1-oxo-2-butene-1-yl]amino}-7-cyclopentyloxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{[4-(N,*N*-dimethylamino)-1-oxo-2-butene-1-yl]amino}-7-[(R)-(tetrahydrofuran-2-yl)methoxy]-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{[4-(N,*N*-dimethylamino)-1-oxo-2-butene-1-yl]amino}-7-[(S)-(tetrahydrofuran-2-yl)methoxy]-quinazoline;

4-[(3-Ethinyl-phenyl)amino]-6,7-bis-(2-methoxy-ethoxy)-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-7-[3-(morpholine-4-yl)-propyloxy]-6-[(vinylcarbonyl)amino]-quinazoline;

4-[(R)-(1-Phenyl-ethyl)amino]-6-(4-hydroxy-phenyl)-7H-pyrrolo[2,3-d]pyrimidine;

3-Cyano-4-[(3-chlor-4-fluorophenyl)amino]-6-{[4-(N,*N*-dimethylamino)-1-oxo-2-butene-1-yl]amino}-7-ethoxy-quinoline;

4-{[3-Chlor-4-(3-fluor-benzyloxy)-phenyl]amino}-6-(5-{[(2-methansulfonyl-ethyl)amino]methyl}-furan-2-yl)quinazoline;

4-[(R)-(1-Phenyl-ethyl)amino]-6-{[4-((R)-6-methyl-2-oxo-morpholine-4-yl)-1-oxo-2-butene-1-yl]amino}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{[4-(morpholine-4-yl)-1-oxo-2-butene-1-yl]amino}-7-[(tetrahydrofuran-2-yl)methoxy]-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-({4-[N,*N*-bis-(2-methoxy-ethyl)-amino]-1-oxo-2-butene-1-yl}amino)-7-[(tetrahydrofuran-2-yl)methoxy]-quinazoline;

4-[(3-Ethinyl-phenyl)amino]-6-{[4-(5,5-dimethyl-2-oxo-morpholine-4-yl)-1-oxo-2-butene-1-yl]amino}-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-[2-(2,2-dimethyl-6-oxo-morpholine-4-yl)-ethoxy]-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-[2-(2,2-dimethyl-6-oxo-morpholine-4-yl)-ethoxy]-7-[(R)-(tetrahydrofuran-2-yl)methoxy]-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-7-[2-(2,2-dimethyl-6-oxo-morpholine-4-yl)-ethoxy]-6-[(S)-(tetrahydrofuran-2-yl)methoxy]-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{2-[4-(2-oxo-morpholine-4-yl)-piperidin-1-yl]-ethoxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-[1-(tert-butyloxycarbonyl)-piperidine-4-yloxy]-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(trans-4-amino-cyclohexane-1-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(trans-4-methansulfonylamino-cyclohexane-1-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(tetrahydropyran-3-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(1-methyl-piperidine-4-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{1-[(morpholine-4-yl)carbonyl]-piperidine-4-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{1-[(methoxymethyl)carbonyl]-piperidine-4-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(piperidine-3-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-[1-(2-acetylamino-ethyl)-piperidine-4-yloxy]-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-ethoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-((S)-tetrahydrofuran-3-yloxy)-7-hydroxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-methoxy-ethoxy)-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{trans-4-[(dimethylamino)sulfonylamino]-cyclohexane-1-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{trans-4-[(morpholine-4-yl)carbonylamino]-cyclohexane-1-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{trans-4-[(morpholine-4-yl)sulfonylamino]-cyclohexane-1-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-acetyl-amino-ethoxy)-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-methane-sulfonylamino-ethoxy)-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{1-[(piperidine-1-yl)carbonyl]-piperidine-4-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(1-aminocarbonylmethyl-piperidine-4-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(cis-4-{*N*-[(tetrahydropyran-4-yl)carbonyl]-*N*-methyl-amino}-cyclohexane-1-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(cis-4-{*N*-[(morpholine-4-yl)carbonyl]-*N*-methyl-amino}-cyclohexane-1-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(cis-4-{*N*-[(morpholine-4-yl)sulfonyl]-*N*-methyl-amino}-cyclohexane-1-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(trans-4-ethansulfonylamino-cyclohexane-1-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(1-methansulfonyl-piperidine-4-yloxy)-7-ethoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(1-methansulfonyl-piperidine-4-yloxy)-7-(2-methoxy-ethoxy)-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-[1-(2-methoxy-acetyl)-piperidine-4-yl-oxy]-7-(2-methoxy-ethoxy)-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(cis-4-acetylamino-cyclohexane-1-yloxy)-7-methoxy-quinazoline;

4-[(3-Ethinyl-phenyl)amino]-6-[1-(tert-butyloxycarbonyl)-piperidine-4-yloxy]-7-methoxy-quinazoline;

4-[(3-Ethinyl-phenyl)amino]-6-(tetrahydropyran-4-yloxy]-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(cis-4-{*N*-[(piperidine-1-yl)carbonyl]-*N*-methyl-amino}-cyclohexane-1-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(cis-4-{*N*-[(4-methyl-piperazine-1-yl)-carbonyl]-*N*-methyl-amino}-cyclohexane-1-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{cis-4-[(morpholine-4-yl)carbonylamino]-cyclohexane-1-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{1-[2-(2-oxopyrrolidine-1-yl)ethyl]-piperidine-4-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{1-[(morpholine-4-yl)carbonyl]-piperidine-4-yloxy}-7-(2-methoxy-ethoxy)-quinazoline;

4-[(3-Ethinyl-phenyl)amino]-6-(1-acetyl-piperidine-4-yloxy)-7-methoxy-quinazoline;

4-[(3-Ethinyl-phenyl)amino]-6-(1-methyl-piperidine-4-yloxy)-7-methoxy-quinazoline;

4-[(3-Ethinyl-phenyl)amino]-6-(1-methansulfonyl-piperidine-4-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(1-methyl-piperidine-4-yloxy)-7-(2-methoxy-ethoxy)-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(1-isopropyloxycarbonyl-piperidine-4-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(cis-4-methylamino-cyclohexane-1-yloxy)-7-methoxy-chinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{cis-4-[*N*-(2-methoxy-acetyl)-*N*-methyl-amino]-cyclohexane-1-yloxy}-7-methoxy-quinazoline;

4-[(3-Ethinyl-phenyl)amino]-6-(piperidine-4-yloxy)-7-methoxy-quinazoline;

4-[(3-Ethinyl-phenyl)amino]-6-[1-(2-methoxy-acetyl)-piperidine-4-yloxy]-7-methoxy-quinazoline;

4-[(3-Ethinyl-phenyl)amino]-6-{1-[(morpholine-4-yl)carbonyl]-piperidine-4-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{1-[cis-2,6-dimethyl-morpholine-4-yl)-carbonyl]-piperidine-4-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{1-[(2-methyl-morpholine-4-yl)carbony-l]-piperidine-4-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{1-[(S,S)-(2-oxa-5-aza-bicyclo[2.2.1]-hept-5-yl)carbonyl]-piperidine-4-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{1-[(*N*-methyl-*N*-2-methoxyethyl-amino)-carbonyl]-piperidine-4-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(1-ethyl-piperidine-4-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{1-[(2-methoxyethyl)carbonyl]-piperidine-4-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-{1-[(3-methoxypropyl-amino)-carbonyl]-piperidine-4-yloxy}-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-[cis-4-(*N*-methansulfonyl-*N*-methyl-amino)-cyclohexane-1-yloxy]-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-[cis-4-(*N*-acetyl-*N*-methyl-amino)-cyclohexane-1-yloxy]-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(trans-4-methylamino-cyclohexane-1-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-[trans-4-(*N*-methansulfonyl-*N*-methyl-amino)-cyclohexane-1-yloxy]-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(trans-4-dimethylamino-cyclohexane-1-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(trans-4-{*N*-[(morpholine-4-yl)carbony-l]-*N*-methyl-amino}-cyclohexane-1-yloxy)-7-methoxy-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-[2-(2,2-dimethyl-6-oxo-morpholine-4-yl)-ethoxy]-7-[(S)-(tetrahydrofuran-2-yl)methoxy]-quinazoline;

4-[(3-Chlor-4-fluorophenyl)amino]-6-(1-methansulfonyl-piperidine-4-yloxy)-7-methoxy-quinazoline; or

4-[(3-Chlor-4-fluorophenyl)amino]-6-(1-cyano-piperidine-4-yloxy)-7-methoxy-quinazoline; optionally in racemic form, as enantiomers, diastereomeres or as pharmaceutically acceptable salts, solvates or hydrates. Preferred are salts selected from the group consisting of hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethansulfonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydrooxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulfonate.

In another embodiment, the additional therapeutic agent is a dopamine antagonist selected from Bromocriptine, Cabergoline, Alpha-Dihydroergocryptine, Lisuride, Pergolide, Pramipexole, Roxindole, Ropinirole, Talipexole, Terguride and Viozane, optionally in racemic form, as enantiomers, diastereomeres or as pharmaceutically acceptable salts, solvates or hydrates. Preferred are salts selected from the group consisting of hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethansulfonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydrooxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulfonate.

In another embodiment, the additional therapeutic agent is an antiallergic agent selected from Epinastine, Cetirizine, Azelastine, Fexofenadine, Levocabastine, Loratadine, Mizolastine, Ketotifene, Emedastine, Dimetindene, Clemastine, Bamipine, Cexchlorpheniramine, Pheniramine, Doxylamine, Chlorphenoxamine, Dimenhydrinate, Diphenhydramine, Promethazine, Ebastine, Olopatadine, Desloratidine and Meclozine, optionally in racemic form, as enantiomers, diastereomeres or as pharmaceutically acceptable salts, solvates or hydrates. Preferred are salts selected from the group consisting of hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethansulfonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydrooxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulfonate.

In one embodiment, the additional therapeutic agent is an MAP kinase inhibitor selected from Bentamapimod, Doramapimod, 5-Carbamoylindole, 6-[(aminocarbonyl)(2,6-difluorophenyl)amino]-2-(2,4-difluorophenyl)-3-pyridine carboxamide, alpha-[2-[[2-(3-pyridinyl)ethyl]amino]-4-pyrimidinyl]-2-benzothiazole acetonitrile, 9,12-Epoxy-1H-diindolo[1,2,3-fg:3',2',1'-kl]pyrrolo[3,4-i][1,6]benzodiazo-cine-10-Carboxylic acid, or 4-[3-(4-chlorophenyl)-5-(1-methyl-4-piperidinyl)-1H-pyrazole-4-yl]-pyrimidine, optionally in racemic form, as enantiomers, diastereomeres or as pharmaceutically acceptable salts, solvates or hydrates. Preferred are salts selected from the group consisting of hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethansulfonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydrooxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulfonate.

In one embodiment, the additional therapeutic agent is an MRP4-Inhibitor selected from *N*-Acetyl-dinitrophenyl-Cysteine, cGMP, Cholate, Diclofenac, Dehydroepiandrosterone 3-glucuronide, Dehydroepiandrosterone 3-sulphate, Dilazep, Dinitrophenyl-S-glutathione, Estradiol 17-beta-glucuronide, Estradiol 3,17-disulphate, Estradiol 3-glucuronide, Estradiol 3-sulphate, Estrone 3-sulphate, Flurbiprofen, Folate, N5-formyl-tetrahydrofolate, Glycocholate, Glycolithocholic acid sulphate, Ibuprofen, Indomethacin, Indoprofen, Ketoprofen, Lithocholic acid sulphate, Methotrexate, (E)-3-[[[3-[2-(7-Chloro-2-quinolinyl)ethenyl]phenyl]-[3-dimethylamino)-3-oxopropyl]thio]methyl]thio]-propanoic acid, alpha-Naphthyl-beta-D-glucuronide, Nitrobenzyl mercaptopurine riboside, Probenecid, Valspodar, Sildenafil, Sulfinpyrazone, Taurochenodeoxycholate, Taurocholate, Taurodeoxycholate, Taurolithocholate, Taurolithocholic acid sulphate, Topotecan, Trequinsin, Zaprinast or Dipyridamol, optionally in racemic form, as enantiomers, diastereomers or as pharmaceutically acceptable salts, solvates or hydrates. Preferred are salts selected from the group consisting of hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethansulfonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydrooxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulfonate.

In one embodiment, the additional therapeutic is an iNOS-Inhibitor selected from S-(2-Aminoethyl)isothio-urea, Aminoguanidine, 2-Aminomethylpyridine, 5,6-dihydro-6-methyl-4H-1,3-thiazine-2-amine (AMT), L-Canavanin, 2-Iminopiperidine, S-Isopropylisothiourea, S-Methylisothiourea, S-Ethylisothiourea, S-Methylthiocitrulline, S-Ethylthiocitrulline, L-NA (N^{ω}-Nitro-L-arginin), L-NAME (N^{ω}-Nitro-L-argininmethylester), L-NMMA (N^{ω} -Monomethyl-L-arginin), L-NIO (N^{ω} -Iminoethyl-L-ornithin), L-NIL (N^{ω} -iminoethyl-lysin),

(S)-6-Acetimidoylamino-2-amino-hexanoic acid (1H-tetrazole-5-yl)-amide;

N-[[3-(aminomethyl)phenyl]methyl]-ethanimidamide;

(S)-4-(2-acetimidoylamino-ethylsulfanyl)-2-amino-buturic acid;

2-[2-(4-Methoxy-pyridine-2-yl)-ethyl]-3H-imidazo[4,5-b]pyridine;

2-((R)-3-amino-1-phenyl-propoxy)-4-chlor-5-fluorbenzonitrile;

2-((1R,3S)-3-amino-4-hydroxy-1-thiazole-5-yl-butylsulfanyl)-6-trifluoromethyl-nicotinonitrile;

2-((1R,3S)-3-amino-4-hydroxy-1-thiazole-5-yl-butylsulfanyl)-4-chlor-benzonitrile;

2-((1R,3S)-3-amino-4-hydroxy-1-thiazole-5-yl-butylsulfanyl)-5-chlor-benzonitrile;

(2S,4R)-2-amino-4-(2-chlor-5-trifluoromethyl-phenylsulfanyl)-4-thiazole-5-yl-butane-1-ol;

2-((1R,3S)-3-amino-4-hydroxy-1-thiazole-5-yl-butylsulfanyl)-5-chlor-nicotinonitrile;

4-((S)-3-amino-4-hydroxy-1-phenyl-butylsulfanyl)-6-methoxy-nicotinonitrile; or substituted 3-phenyl-3,4-dihydro-1-isoquinolinamine as for instance 1S,5S,6R)-7-Chlor-5-methyl-2-aza-bicyclo[4.1.0]hept-2-ene-3-ylamin(4R,5R)-5-Ethyl-4-methyl-thiazolidine-2-ylideneamine, (1S,5S,6R)-7-Chlor-5-methyl-2-aza-bicyclo[4.1.0]hept-2-ene-3-ylamin, (4R,5R)-5-Ethyl-4-methyl-thiazolidine-2-ylideneamine, (4R,5R)-5-Ethyl-4-methyl-selenazolidine-2-ylideneamine, 4-Aminotetrahydrobiopterine, (E)-3-(4-Chlor-phenyl)-*N*-(1-{2-oxo-2-[4-(6-trifluormethyl-pyrimidine-4-yloxy)-piperidine-1-yl]-ethylcarbamoyl}-2-pyridine-2-yl-ethyl)-acrylamide, 3-(2,4-Difluor-phenyl)-6-[2-(4-imidazole-1-ylmethyl-phenoxy)-ethoxy]-2-phenyl-pyridine, 3-{[(Benzo[1,3]dioxol-5-ylmethyl)-carbamoyl]-methyl}-4-(2-imidazole-1-yl-pyrimidine-4-yl)-piperazine-1-carbon acid methylester, or (R)-1-(2-imidazole-1-yl-6-methyl-pyrimidine-4-yl)-pyrrolidine-2-carbon acid (2-benzo[1,3]dioxol-5-yl-ethyl)-amide, optionally in racemic form, as enantiomers, diastereomeres or as pharmaceutically acceptable salts, solvates or hydrates. Preferred are salts selected from the group consisting of hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethansulfonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydrooxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulfonate. Further examples of preferred iNOS-Inhibitors which may be mentioned include antisense-Oligonucleotide, especially those antisense-Oligonucleotide binding iNOS-coding nucleic acids, examples therefore are disclosed in WO 01/52902, incorporated herein by reference in its entirety.

In another embodiment, the additional therapeutic agent is a SYK-inhibitor selected from

2-[(2-aminoethyl)amino]-4-[(3-bromophenyl)amino]-5-pyrimidinecarboxamide;

2-[[7-(3,4-dimethoxyphenyl)imidazo[1,2-c]pyrimidine-5-yl]amino]-3-pyridinecarboxamide;

6-[[5-fluoro-2-[3,4,5-trimethoxyphenyl)amino]-4-pyrimidinyl]amino]-2,2-dimethyl-2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one;

N-[3-bromo-7-(4-methoxyphenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

7-(4-methoxyphenyl)-*N*-methyl-1,6-naphthyridine-5-amine;

N-[7-(4-methoxyphenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(2-thienyl)-1,6-naphthyridine-5-yl-1,3-propanediamine;

N-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-1,2-ethanediamine;

N-[7-(4-methoxyphenyl)-2-(trifluoromethyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(4-methoxyphenyl)-3-phenyl-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-(7-phenyl-1,6-naphthyridine-5-yl)-1,3-propanediamine;

N-[7-(3-fluorophenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(3-chlorophenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[3-(trifluoromethoxy)phenyl]-1,6-naphthyridine-5yl]-1,3-propanediamine;

N-[7-(4-fluorophenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(4-fluorophenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(4-chlorophenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(4'-methyl[1,1'-biphenyl]-4-yl)-1,6-naphthyridine-1,3-propane-diamine;

N-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[4-(diethylamino)phenyl]-1,6-naphthyridine-5-yl]-1,3-propane-diamine;

N-[7-[4-(4-morpholinyl)phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[4-[[2-(dimethylamino)ethyl]methylamino]phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(4-bromophenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(4-methylphenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[4-(methylthio)phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[4-(1-methylethyl)phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

7-[4-(dimethylamino)phenyl]-*N*-methyl-1,6-naphthyridine-5-amine;

7-[4-(dimethylamino)phenyl]-N,*N*-dimethyl-1,6-naphthyridine-5-amine;

N-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-1,4-butanediamine;

N-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-1,5-pentanediamine;

3-[[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]oxy]-1-propanole;

4-[5-(4-aminobutoxy)-1,6-naphthyridine-7-yl]-N,*N*-dimethyl-benzenamine;

4-[[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]amino]-1-butanole;

N-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-*N*-methyl-1,3-propanediamine;

N-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-N'-methyl-1,3-propanediamine;

N-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-N,N'-dimethyl-1,3-propanediamine;

1-amino-3-[[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]amino]-2-propanole;

N-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-2,2-dimethyl-1,3-propanediamine;

7-[4-(dimethylamino)phenyl]-*N*-(3-pyridinylmethyl)-1,6-naphthyridine-5-amine;

N-[(2-aminophenyl)methyl]-7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-amine;

N-[7-[6-(dimethylamino)[1,1'-biphenyl]-3-yl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[3-chloro-4-(diethylamino)phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[4-(dimethylamino)-3-methoxyphenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[4-(diethylamino)phenyl]-3-methyl-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(3'-fluoro[1,1'-biphenyl]-3-yl)-1,6-naphthyridine-5-yl]-1,2-ethanediamine;

N-[7-(4-methoxyphenyl)-1,6-naphthyridine-5-yl]-1,6-naphthyridine-1,3-propanediamine;

N,N'-bis(3-aminopropyl)-7-(4-methoxyphenyl)-2,5-diamine;

N-[7-(4-methoxyphenyl)-2-(phenylmethoxy)-1,6-naphthyridine-5-yl]-1,6-naphthyridine-1,3-propanediamine;

-(3-aminopropyl)-7-(4-methoxyphenyl)-N2-(phenylmethyl)-2,5-diamine;

N-[7-(2-naphthalenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(2'-fluoro[1,1'-biphenyl]-4-yl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(3,4,5-trimethoxyphenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(3,4-dimethylphenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

1-amino-3-[[7-(2-naphthalenyl)-1,6-naphthyridine-5-yl]amino]-2-propanole;

1-amino-3-[[7-(2'-fluoro[1,1'-biphenyl]-4-yl)-1,6-naphthyridine-5-yl]amino]-2-propanole;

1-amino-3-[[7-(4'-methoxy[1,1'-biphenyl]-4-yl)-1,6-naphthyridine-5-yl]amino]-2-propanole;

1-amino-3-[[7-(3,4,5-trimethoxyphenyl)-1,6-naphthyridine-5-yl]amino]-2-propanole;

1-amino-3-[[7-(4-bromophenyl)-1,6-naphthyridine-5-yl]amino]-2-propanole;

N-[7-(4'-methoxy[1,1'-biphenyl]-4-yl)-1,6-naphthyridine-5-yl]-2,2-dimethyl-1,3-propanediamine;

1-[[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]amino]-2-propanole;

2-[[2-[[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]amino]-ethyl]thio]-ethanole;

7-[4-(dimethylamino)phenyl]-*N*-(3-methyl-5-isoxazolyl)-1,6-naphthyridine-5-amine;

7-[4-(dimethylamino)phenyl]-*N*-4-pyrimidinyl-1,6-naphthyridine-5-amine;

N-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-1,3-cyclohexane diamine;

N,*N*-dimethyl-4-[5-(1-piperazinyl)-1,6-naphthyridine-7-yl]-benzenamine;

4-[5-(2-methoxyethoxy)-1,6-naphthyridine-7-yl]-N,*N*-dimethyl-benzeneamine;

1-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-4-piperidinole;

1-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-3-pyrrolidinole;

7-[4-(dimethylamino)phenyl]-*N*-(2-furanylmethyl)-1,6-naphthyridine-5-amine;

7-[4-(dimethylamino)phenyl]-*N*-[3-(1H-imidazole-1-yl)propyl]-1,6-naphthyridine-5-amine;

1-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-4-piperidine carboxamide;

1-[3-[[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]amino]propyl]-2-pyrrolidinone;

N-[3'-[5-[(3-aminopropyl)amino]-1,6-naphthyridine-7-yl][1,1'-biphenyl]-3-yl]-acetamide;

N-[7-(4'-fluoro[1,1'-biphenyl]-4-yl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[4'-[5-[(3-aminopropyl)amino]-1,6-naphthyridine-7-yl][1,1'-biphenyl]-3-yl]-acetamide;

N-[7-[4-(1,3-benzodioxol-5-yl)phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[4-(2-thienyl)phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[4-fluoro-3-(trifluoromethyl)phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[4-(3-pyridinyl)phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(1,3-benzodioxol-5-yl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(6-methoxy-2-naphthalenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

7-[4-(dimethylamino)phenyl]-*N*-(4-pyridinylmethyl)-1,6-naphthyridine-5-amine;

3-[[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]methylamino]-propanenitrile;

7-[4-(dimethylamino)phenyl}-*N*-[1-(phenylmethyl)-4-piperidinyl]-1,6-naphthyridine-5-amine;

N-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-1,2-cyclohexanediamine;

N-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-1,2-cyclohexanediamine, (1R,2S)-rel-;

N-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-1,2-benzene dimethanamine;

N-[7-[4-(diethylamino)phenyl]-1,6-naphthyridine-5-yl]-1,4-butanediamine;

N-[7-[3',5'-bis(trifluoromethyl)[1,1'-biphenyl]-4-yl]-1,6-naphthyr- idine-5-yl]-,3-propanediamine;

N-[7-(3'-methoxy[1,1'-biphenyl]-4-yl)-1,6-naphthyridine-5-yl}-1,3-propane-diamine;

N-[7-(3'-fluoro[1,1'-biphenyl]-4-yl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

4-[[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]oxy}-1-butanole;

N-[7-[4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-1,4-cyclohexanediamine;

7-[4-(dimethylamino)phenyl]-*N*-(2,2,6,6-tetramethyl-4-piperidinyl)-1,6-naphthyridine-5-amine;

N-[7-[3-bromo-4-(dimethylamino)phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(1-methyl-1H-indole-5-yl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[3-(trifluoromethyl)phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[4-(trifluoromethyl)phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-(3-bromo-4-methoxyphenyl)-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N-[7-[4-[[3-(dimethylamino)propyl]methylamino]phenyl]-1,6-naphthyridine-5-yl]-1,4-cyclohexanediamine;

N-[7-[4-[[2-(dimethylamino)ethyl]methylamino]phenyl]-1,6-naphthyridine-5-yl]-1,4-cyclohexanediamine;

N-[7-[4-(dimethylamino)-3-methoxyphenyl]-1,6-naphthyridine-5-yl]-1,4-cyclohexanediamine;

N-[7-[4-(4-morpholinyl)phenyl]-1,6-naphthyridine-5-yl]-1,4-cyclohexanediamine;

N-[7-[3-bromo-4-(4-morpholinyl)phenyl]-1,6-naphthyridine-5-yl]-1,4-cyclohexanediamine;

4-[[7-[4-[[2-(dimethylamino)ethyl]methylamino]phenyl]-1,6-naphthyridine-5-yl]oxy]-cyclohexanole;

N-[7-[3-bromo-4-(4-morpholinyl)phenyl]-1,6-naphthyridine-5-yl]-1,3-propanediamine;

N,*N*-dimethyl-4-[5-(4-methyl-1-piperazinyl)-1,6-naphthyridine-7-yl]-benzenamine;

4-[[7-[4-[[3-(dimethylamino)propyl]methylamino]phenyl]-1,6-naphthyridine-5-yl]oxy]-cyclohexanole;

N-[7-[4-[[2-(dimethylamino)ethyl]methylamino]phenyl]-1,6-naphthyridine-5-yl]-1,4-butanediamine; or

[3-[[5-[(3-aminopropyl)amino]-7-(4-methoxyphenyl)-1,6-naphthyridine-2-yl]-amino]propyl]-carbamic acid-1,1-dimethylethyl ester, optionally in racemic form, as enantiomers, diastereomers or as pharmaceutically acceptable salts, solvates or hydrates. Preferred are salts selected from the group consisting of hydrochloride, hydrobromide, hydroiodide, hydrosulfate, hydrophosphate, hydromethansulfonate, hydronitrate, hydromaleate, hydroacetate, hydrocitrate, hydrofumarate, hydrotartrate, hydrooxalate, hydrosuccinate, hydrobenzoate and hydro-p-toluenesulfonate.

In another embodiment, the additional therapeutic agent is an agent that inhibits the interaction between CAL (also known as CFTR-associated ligand) and mutant CFTR proteins. In one embodiment, the agent that inhibits interaction between CAL and mutant CFTR proteins is a peptide or peptidomimetic (e.g., 6 to 20 residues in length). In another embodiment, the agent that inhibits interaction between CAL and mutant CFTR proteins is a peptide or peptidomimetic disclosed in US Published Patent Application No. US2014/0100155 A1, incorporated herein by reference in its entirety.

In another embodiment, the above recited pharmaceutical compositions contain a pharmaceutically acceptable prodrug of the compound of the present invention.

In one aspect, the pharmaceutical compositions of the invention can be administered to a patient once daily or about every twenty four hours. Alternatively, the pharmaceutical compositions of the invention can be administered to a patient twice daily. Alternatively, the pharmaceutical composition of the invention can be administered about every twelve hours. These pharmaceutical compositions are administered as oral formulations containing about 25 mg, 50 mg, 100 mg, 125 mg, 150 mg, 200 mg, 250 mg, 300 mg, or 400 mg of a compound of Formula I. In this aspect, the pharmaceutical compositions further comprise a filler; a disintegrant; a surfactant; a binder; or a lubricant, or combinations thereof.

It will also be appreciated that the pharmaceutical compositions of the invention, including the pharmaceutical compositions comprising combinations described previously, can be employed in combination therapies; that is, the compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutic agents or medical procedures.

These combinations are useful for treating the diseases described herein including cystic fibrosis. These combinations are also useful in the kits described herein.

The amount of additional therapeutic agent present in or with the compositions of this invention will be no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent. Preferably the amount of additional therapeutic agent in the presently disclosed compositions will range from about 50% to 100% of the amount normally present in a composition comprising that agent as the only therapeutically active agent.

Therapeutic Uses of the Compounds of Formula I-Ij and Pharmaceutical Compositions Thereof

In one aspect, the invention also provides a method of treating, lessening the severity of, or symptomatically treating a disease in a patient, the method comprising administering an effective amount of the pharmaceutical composition of the invention to the patient, preferably a mammal, wherein the disease is selected from cystic fibrosis, asthma, smoke induced COPD, chronic bronchitis, rhinosinusitis, constipation, pancreatitis, pancreatic insufficiency, male

infertility caused by congenital bilateral absence of the vas deferens (CBAVD), mild pulmonary disease, idiopathic pancreatitis, allergic bronchopulmonary aspergillosis (ABPA), liver disease, hereditary emphysema, hereditary hemochromatosis, coagulation-fibrinolysis deficiencies, such as protein C deficiency and Type 1 hereditary angioedema, lipid processing deficiencies (such as familial hypercholesterolemia, Type 1 chylomicronemia, and abetalipoproteinemia), lysosomal storage diseases (such as I-cell disease/pseudo-Hurler), mucopolysaccharidoses, Sandhof/Tay-Sachs, Crigler-Najjar type II, polyendocrinopathy/hyperinsulinemia, Diabetes mellitus, Laron dwarfism, myeloperoxidase deficiency, primary hypoparathyroidism, melanoma, glycanosis CDG type 1, congenital hyperthyroidism, osteogenesis imperfecta, hereditary hypofibrinogenemia, ACT deficiency, Diabetes insipidus (DI), neurohypophyseal DI, nephrogenic DI, Charcot-Marie Tooth syndrome, Pelizaeus-Merzbacher disease, neurodegenerative diseases (such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, and Pick's disease), several polyglutamine neurological disorders (such as Huntington's, spinocerebellar ataxia type I, spinal and bulbar muscular atrophy, dentatorubral pallidoluysian atrophy, and myotonic dystrophy), spongiform encephalopathies (such as hereditary Creutzfeldt-Jakob disease (due to prion protein processing defect)), Fabry disease, Gerstmann-Straussler-Scheinker syndrome, COPD, dry-eye disease, or Sjogren's disease, osteoporosis, osteopenia, bone healing and bone growth (including bone repair, bone regeneration, reducing bone resorption and increasing bone deposition), Gorham's Syndrome, chloride channelopathies such as myotonia congenita (Thomson and Becker forms), Bartter's syndrome type III, Dent's disease, epilepsy, lysosomal storage disease, Angelman syndrome, and Primary Ciliary Dyskinesia (PCD), a term for inherited disorders of the structure and/or function of cilia, including PCD with situs inversus (also known as Kartagener syndrome), PCD without situs inversus and ciliary aplasia.

In another embodiment, the method of treating, lessening the severity of, or symptomatically treating a disease in a patient of the invention comprises administering an effective amount of the pharmaceutical composition of the invention, provided that in formula I and the attendant definitions: i) when ring A is indole, n is not zero and rings B and C are aryl unsubstituted by CF_3 or halo; ii) when ring A is pyrazole, n is not zero; iii) when ring A is pyridyl, ring B and ring C moieties are substituted at the 2- and 3- positions of the pyridyl ring,

interchangeably; and iv) when ring A is imidazole, the ring B moiety is substituted at the 2-position of the imidazole ring.

In one aspect, the invention also provides a method of treating, lessening the severity of, or symptomatically treating a disease in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the disease is selected from generalized epilepsy with ferbrile seizures plus (GEFS+), general epilepsy with ferbile and aferbrile seizures, myotonia, paramyotonia congenital, potassium-aggravated myotonia, hyperkalemic periodic paralysis, LQTS, LQTS/Brugada syndrome, autosomal-dominant LQTS with deafness, autosomal-recessive LQTS, LQTS with dysmorphic features, congenital and acquired LQTS, Timothy syndrome, persistent hyperinsulinemic hypolglycemia of infancy, dilated cardiomyopathy, autosomal-dominant LQTS, Dent disease, Osteopetrosis, Bartter syndrome type III, central core disease, malignant hyperthermia, and catecholaminergic polymorphic tachycardia.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation N1303K, ΔI507, or R560T.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation G551D. In another embodiment, the patient is homozygous in G551D. In another embodiment, the patient is heterozygous in G551D. In another embodiment, the patient is heterozygous in G551D, wherein the other CFTR genetic mutation is any one of F508del, G542X, N1303K, W1282X, R117H, R553X, 1717-1G->A, 621+1G->T, 2789+5G->A, 3849+10kbC->T, R1162X, G85E, 3120+1G->A, ΔI507, 1898+1G->A, 3659delC, R347P, R560T, R334W, A455E, 2184delA, or 711+1G->T. In another embodiment, the patient is heterozygous in G551D, wherein the other CFTR genetic mutation is F508del. In another embodiment, the patient is heterozygous in G551D, wherein the other CFTR genetic mutation is R117H.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation F508del. In another embodiment, the patient is homozygous in F508del. In another embodiment, the patient is heterozygous in F508del. In another embodiment, the patient is heterozygous in F508del, wherein the other CFTR genetic mutation is any one of G551D, G542X, N1303K, W1282X, R117H, R553X, 1717-1G->A, 621+1G->T, 2789+5G->A, 3849+10kbC->T, R1162X, G85E, 3120+1G->A, ΔI507, 1898+1G->A, 3659delC, R347P, R560T, R334W, A455E, 2184delA, or 711+1G->T. In another embodiment, the patient is heterozygous in F508del, wherein the other CFTR genetic mutation is G551D. In another embodiment, the patient is heterozygous in F508del, wherein the other CFTR genetic mutation is R117H.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V, G1069R, R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N, D1152H, 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V and G1069R. In one embodiment of this aspect, the invention provides a method of treating CFTR comprising administering Compound 1 to a patient possessing a human CFTR mutation

selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R and S1251N. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from E193K, F1052V and G1069R. In some embodiments of this aspect, the method produces a greater than 10-fold increase in chloride transport relative to baseline chloride transport.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N and D1152H. In one embodiment of this aspect, the method produces an increase in chloride transport which is greater or equal to 10% above the baseline chloride transport.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 1811+1.6kbA->G, 2789+5G->A, 3272-26A->G and 3849+10kbC->T. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient

comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 2789+5G->A and 3272-26A->G.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V, G1069R, R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N, D1152H, 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G, and a human CFTR mutation selected from F508del, R117H, and G551D.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V and G1069R, and a human CFTR mutation selected from F508del, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R and S1251N, and a human CFTR mutation selected from F508del, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient

possesses the CFTR genetic mutation is selected from E193K, F1052V and G1069R, and a human CFTR mutation selected from F508del, R117H, and G551D. In some embodiments of this aspect, the method produces a greater than 10-fold increase in chloride transport relative to baseline chloride transport.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N and D1152H, and a human CFTR mutation selected from F508del, R117H, and G551D. In one embodiment of this aspect, the method produces an increase in chloride transport which is greater or equal to 10% above the baseline chloride transport.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G, and a human CFTR mutation selected from F508del, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 1811+1.6kbA->G, 2789+5G->A, 3272-26A->G and 3849+10kbC->T, and a human CFTR mutation selected from F508del, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective

amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 2789+5G->A and 3272-26A->G, and a human CFTR mutation selected from F508del, R117H.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V, G1069R, R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N, D1152H, 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G, and a human CFTR mutation selected from F508del, R117H, and G551D.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V and G1069R. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R and S1251N. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from E193K,

F1052V and G1069R. In some embodiments of this aspect, the method produces a greater than 10-fold increase in chloride transport relative to baseline chloride transport.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N and D1152H. In one embodiment of this aspect, the method produces an increase in chloride transport which is greater or equal to 10% above the baseline chloride transport.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 1811+1.6kbA->G, 2789+5G->A, 3272-26A->G and 3849+10kbC->T. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 2789+5G->A and 3272-26A->G.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an

effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V, G1069R, R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N, D1152H, 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G, and a human CFTR mutation selected from F508del, R117H, and G551D, and one or more human CFTR mutations selected from F508del, R117H, and G551D.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R, S1251N, E193K, F1052V and G1069R, and one or more human CFTR mutations selected from F508del, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from G178R, G551S, G970R, G1244E, S1255P, G1349D, S549N, S549R and S1251N, and one or more human CFTR mutations selected from F508del, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from E193K, F1052V and G1069R, and one or more human CFTR mutations selected from F508del, R117H, and G551D. In some embodiments of this aspect, the method produces a greater than 10-fold increase in chloride transport relative to baseline chloride transport.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from R117C, D110H, R347H, R352Q, E56K, P67L, L206W, A455E, D579G, S1235R, S945L, R1070W, F1074L, D110E, D1270N and D1152H, and one or more human CFTR mutations selected from F508del, R117H, and G551D. In one embodiment of this aspect, the method produces an increase in chloride transport which is greater or equal to 10% above the baseline chloride transport.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 621+1G->T, 3120+1G->A, 1898+1G->A, 711+1G->T, 2622+1G->A, 405+1G->A, 406-1G->A, 4005+1G->A, 1812-1G->A, 1525-1G->A, 712-1G->T, 1248+1G->A, 1341+1G->A, 3121-1G->A, 4374+1G->T, 3850-1G->A, 2789+5G->A, 3849+10kbC->T, 3272-26A->G, 711+5G->A, 3120G->A, 1811+1.6kbA->G, 711+3A->G, 1898+3A->G, 1717-8G->A, 1342-2A->C, 405+3A->C, 1716G/A, 1811+1G->C, 1898+5G->T, 3850-3T->G, IVS14b+5G->A, 1898+1G->T, 4005+2T->C and 621+3A->G, and one or more human CFTR mutations selected from F508del, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 1717-1G->A, 1811+1.6kbA->G, 2789+5G->A, 3272-26A->G and 3849+10kbC->T, and one or more human CFTR mutations selected from F508del, R117H, and G551D. In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation is selected from 2789+5G->A and 3272-26A->G, and one or more human CFTR mutations selected from F508del, R117H, and G551D.

In one aspect, the present invention is directed to a method of treating, lessening the severity of, or symptomatically treating cystic fibrosis in a patient comprising administering an effective amount of the pharmaceutical composition or tablet of the invention to the patient, preferably a mammal, wherein the patient possesses the CFTR genetic mutation selected from Figure 1.

In certain embodiments, the composition of the present invention is useful for treating, lessening the severity of, or symptomatically treating cystic fibrosis in patients who exhibit residual CFTR activity in the apical membrane of respiratory and non-respiratory epithelia. The presence of residual CFTR activity at the epithelial surface can be readily detected using methods known in the art, e.g., standard electrophysiological, biochemical, or histochemical techniques. Such methods identify CFTR activity using *in vivo* or *ex vivo* electrophysiological techniques, measurement of sweat or salivary Cl^- concentrations, or *ex vivo* biochemical or histochemical techniques to monitor cell surface density. Using such methods, residual CFTR activity can be readily detected in patients heterozygous or homozygous for a variety of different mutations, including patients homozygous or heterozygous for the most common mutation, F508del, as well as other mutations such as the G551D mutation, or the R117H mutation. In certain embodiments, compositions of the present invention are useful for treating, lessening the severity of, or symptomatically treating cystic fibrosis in patients who exhibit little to no residual CFTR activity. In certain embodiments, compositions of the present invention are useful for treating, lessening the severity of, or symptomatically treating cystic fibrosis in patients who exhibit little to no residual CFTR activity in the apical membrane of respiratory epithelia.

In another embodiment, the compositions of the present invention are useful for treating or lessening the severity of cystic fibrosis in patients who exhibit residual CFTR activity using pharmacological methods. In another embodiment, the compositions of the present invention are useful for treating or lessening the severity of cystic fibrosis in patients who have residual CFTR activity using gene therapy. Such methods increase the amount of CFTR present at the cell surface, thereby inducing a hitherto absent CFTR activity in a patient or augmenting the existing level of residual CFTR activity in a patient.

In one embodiment, the compositions of the present invention are useful for treating or lessening the severity of cystic fibrosis in patients within certain genotypes exhibiting residual

CFTR activity, e.g., Class I mutations (not synthesized), class II mutation (misfolding), class III mutations (impaired regulation or gating), class IV mutations (altered conductance), or class V mutations (reduced synthesis).

In one embodiment, compositions of the present invention are useful for treating, lessening the severity of, or symptomatically treating cystic fibrosis in patients within certain clinical phenotypes, e.g., a moderate to mild clinical phenotype that typically correlates with the amount of residual CFTR activity in the apical membrane of epithelia. Such phenotypes include patients exhibiting pancreatic sufficiency.

In one embodiment, the compositions of the present invention are useful for treating, lessening the severity of, or symptomatically treating patients diagnosed with pancreatic sufficiency, idiopathic pancreatitis and congenital bilateral absence of the vas deferens, or mild lung disease wherein the patient exhibits residual CFTR activity.

In one embodiment, the compositions of the present invention are useful for treating, lessening the severity of, or symptomatically treating patients diagnosed with pancreatic sufficiency, idiopathic pancreatitis and congenital bilateral absence of the vas deferens, or mild lung disease wherein the patient has wild type CFTR.

In addition to cystic fibrosis, modulation of CFTR activity may be beneficial for other diseases not directly caused by mutations in CFTR, such as secretory diseases and other protein folding diseases mediated by CFTR. These include, but are not limited to, chronic obstructive pulmonary disease (COPD), dry eye disease, and Sjögren's Syndrome. COPD is characterized by airflow limitation that is progressive and not fully reversible. The airflow limitation is due to mucus hypersecretion, emphysema, and bronchiolitis. Activators of mutant or wild-type CFTR offer a potential treatment of mucus hypersecretion and impaired mucociliary clearance that is common in COPD. Specifically, increasing anion secretion across CFTR may facilitate fluid transport into the airway surface liquid to hydrate the mucus and optimized periciliary fluid viscosity. This would lead to enhanced mucociliary clearance and a reduction in the symptoms associated with COPD. Dry eye disease is characterized by a decrease in tear aqueous production and abnormal tear film lipid, protein and mucin profiles. There are many causes of dry eye, some of which include age, Lasik eye surgery, arthritis, medications, chemical/thermal burns, allergies, and diseases, such as cystic fibrosis and Sjögrens's syndrome. Increasing anion

secretion via CFTR would enhance fluid transport from the corneal endothelial cells and secretory glands surrounding the eye to increase corneal hydration. This would help to alleviate the symptoms associated with dry eye disease. Sjögrens's syndrome is an autoimmune disease in which the immune system attacks moisture-producing glands throughout the body, including the eye, mouth, skin, respiratory tissue, liver, vagina, and gut. Symptoms, include, dry eye, mouth, and vagina, as well as lung disease. The disease is also associated with rheumatoid arthritis, systemic lupus, systemic sclerosis, and polymypositis/dermatomyositis. Defective protein trafficking is believed to cause the disease, for which treatment options are limited. Augmenters or inducers of CFTR activity may hydrate the various organs afflicted by the disease and help to elevate the associated symptoms.

In one embodiment, the invention relates to a method of augmenting or inducing anion channel activity *in vitro* or *in vivo*, comprising contacting the channel with a composition of the present invention. In another embodiment, the anion channel is a chloride channel or a bicarbonate channel. In another embodiment, the anion channel is a chloride channel.

The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular agent, its mode of administration, and the like. The compounds of the invention are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. The expression "dosage unit form" as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the invention will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed, and like factors well known in the medical arts. The term "patient", as used herein, means an animal, preferably a mammal, and most preferably a human.

In recent years a number of attempts have been made to treat cystic fibrosis using gene therapy. Gene therapy is the insertion, alteration, or removal of genes within an individual's cells and biological tissues to treat disease. It is a technique for correcting defective genes that are responsible for disease development. The most common form of gene therapy involves the insertion of functional genes into an unspecified genomic location in order to replace a mutated gene, but other forms involve directly correcting the mutation or modifying normal gene that enables a viral infection. Although the technology is still in its infancy, it has been used with some success.

Cystic fibrosis is a good candidate for gene therapy as it is primarily caused by mutations in a single gene. A normal copy of the gene could be delivered to patients via topical delivery to the lung, not requiring invasive techniques or surgery. A gene complementation approach would aslo directly target the cause of the disease and could correct many aspects of the complex lung pathology.

Soon after the cloning of the CFTR gene, proof-of-principle was established when the Cl-conductance defect was corrected after delivery of a functional copy of human wild-type CFTR DNA to cells isolated from cystic fibrosis patients. To date, a number of trials for cystic fibrosis gene therapy have been tested in humans. These early studies were concerned mainly with safety issues.

In one embodiment, the invention features a method of treating cystic fibrosis comprising administering to the patient a compound of formula I to Id in conjunction with gene therapy. The gene therapy can be as disclosed in Internation Published Patent Application No. WO2013061091, incorporated herein in its entirety by reference. Administration may be prior, concomitant, or subsequent administration of a compound of formula I – Id, or any of the pharmaceutical compositions of the present invention.

In another embodiment, the gene therapy features administering to the patient 1 mL to less than 10 mLs of a complex of (i) a non-viral CpG dinucleotide-free plasmid comprising nucleic acid encoding a CFTR polypeptide operatively linked to hCEF1 promoter, wherein the plasmid is at a concentration of 2 mg/mL to 3 mg/mL, and (ii) GL67A lipid mixture at a concentration of 10 mg/mL to 20 mg/mL.

In one embodiment, gene therapy is administered to a patient with cystic fibrosis prior, concomitant, or subsequent administration of a pharmaceutical composition comprising (i) a compound of formula I – Id and (ii) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid.

In one embodiment, gene therapy is administered to a patient with cystic fibrosis prior, concomitant, or subsequent administration of a pharmaceutical composition comprising (i) a compound of formula I – Id and (ii) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide.

In one embodiment, gene therapy is administered to a patient with cystic fibrosis prior, concomitant, or subsequent administration of a pharmaceutical composition comprising (i) a compound of formula I – Id and (ii) 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid.

In one embodiment, gene therapy is administered to a patient with cystic fibrosis prior, concomitant, or subsequent administration of a pharmaceutical composition comprising (i) a compound of formula I – Id; (ii) 3-(6-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl) cyclopropanecarboxamido)-3-methylpyridin-2-yl)benzoic acid; and (iii) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

In one embodiment, gene therapy is administered to a patient with cystic fibrosis prior, concomitant, or subsequent administration of a pharmaceutical composition comprising (i) a compound of formula I – Id; (ii) (*R*)-1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-*N*-(1-(2,3-dihydroxypropyl)-6-fluoro-2-(1-hydroxy-2-methylpropan-2-yl)-1H-indol-5-yl)cyclopropanecarboxamide; and (iii) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.


In one embodiment, gene therapy is administered to a patient with cystic fibrosis prior, concomitant, or subsequent administration of a pharmaceutical composition comprising (i) a compound of formula I – Id; (ii) 4-(3-(1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)cyclopropanecarboxamido) isoquinolin-1-yl)benzoic acid; and (iii) *N*-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide.

Kits

In another aspect, the present invention features a kit comprising a compound and/or pharmaceutical composition of the present invention and instructions for use thereof.

In another embodiment, the kits of the present invention further comprise one or more additional therapeutic agent(s). In another embodiment, the additional therapeutic agent is selected from a mucolytic agent, bronchodialator, an antibiotic, an anti-infective agent, a CFTR modulator, or an anti-inflammatory agent. In another embodiment, the additional therapeutic agent is a CFTR modulator. In another embodiment, the additional therapeutic agent is a CFTR corrector.


In another embodiment, the additional therapeutic agent is



,

or a pharmaceutically acceptable salt thereof.


In another embodiment, the additional therapeutic agent is



,

or a pharmaceutically acceptable salt thereof.

In another embodiment, the additional therapeutic agent is




,

or a pharmaceutically acceptable salt thereof.

In another embodiment, the additional therapeutic agent is a CFTR potentiator.

In another embodiment, the additional therapeutic agent is




,


or a pharmaceutically acceptable salt thereof.

In another embodiment, the additional therapeutic agents are a CFTR corrector and a CFTR potentiator.

In another embodiment, the additional therapeutic agents are




and




, or pharmaceutically acceptable salts thereof.

In another embodiment, the additional therapeutic agents are




and




, or pharmaceutically acceptable salts thereof.

In another embodiment, the additional therapeutic agents are



and



, or pharmaceutically acceptable salts thereof.

In another embodiment, the kits of the present invention are drawn to kits wherein the compounds or the pharmaceutical compositions of the present invention and the one or more additional therapeutic agent(s) are in separate containers.

In another embodiment, the kits fo the present invention are drawn to kits wherein the compounds or the pharmaceutical compositions of the present invention and the one or more additional therapeutic agent(s) are in the same container.


In another embodiment, the container is a bottle, vial, or blister pack, or combination thereof.

Anywhere in the present application where a name of a compound may not correctly describe the structure of the compound, the structure supersedes the name and governs.

EXAMPLES


PREPARATION 1: *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenyl)pyridine-3-carboxamide (Compound 26)

Step 1: 6-*tert*-butyl-2-chloro-pyridine-3-carbonitrile




2,2-Dimethylpropanoic acid (50. mL, 870 mmol), 2-chloropyridine-3-carbonitrile (30 g, 220 mmol) and $AgNO_3$ (9.2 g, 54 mmol) were suspended in 10% aq. H_2SO_4 (300 mL). A solution of ammonium peroxydisulfate (91 g, 430 mmol) and water (480 mL) was added to the mixture dropwise through an addition funnel. The mixture was stirred at room temperature overnight. The pH of reaction mixture was adjusted to ~8-9 with 30% NH_4OH (~250 mL) and the mixture was extracted with ethyl acetate (3 x 300 mL). The combined organic layers were washed with brine, dried over sodium sulfate, and concentrated. The residue was subjected to silica gel column chromatography (0-25% ethyl acetate/hexanes) to give 6-*tert*-butyl-2-chloro-pyridine-3-carbonitrile (37 g, 83%) as a clear oil which solidified upon standing. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.83 (d, J = 8.1 Hz, 1H), 7.30 (d, J = 8.1 Hz, 1H), 1.29 (s, 9H). ESI-MS *m/z* calc. 194.1, found 195.0 $(M+1)^+$; Retention time: 1.66 min (3 min run).

Step 2: 6-*tert*-butyl-2-(2,4,6-trimethylphenyl)pyridine-3-carbonitrile




A mixture of 6-*tert*-butyl-2-chloro-pyridine-3-carbonitrile (400. mg, 2.06 mmol), (2,4,6-trimethylphenyl)boronic acid (506 mg, 3.08 mmol), $Pd(dppf)_2Cl_2*CH_2Cl_2$ (75 mg, 0.10 mmol), $NaHCO_3$ (173 mg, 2.06 mmol) and DME (2 mL) was heated at 120 °C in a microwave reactor for 30 min. The mixture was partitioned between CH_2Cl_2 and water. The layers were separated and the aqueous layer was extracted with CH_2Cl_2 (2x). The combined organics were dried over sodium sulfate, filtered and concentrated. The residue was subjected to silica gel column chromatography (0-20% ethyl acetate/hexanes) to give 6-*tert*-butyl-2-(2,4,6-trimethylphenyl)pyridine-3-carbonitrile (225 mg, 39%). ESI-MS *m/z* calc. 278.2, found 279.4 $(M+1)^+$; Retention time: 0.84 min (3 min run).

Step 3: 6-*tert*-butyl-2-(2,4,6-trimethylphenyl)pyridine-3-carboxylic acid



A mixture of 6-*tert*-butyl-2-(2,4,6-trimethylphenyl)pyridine-3-carbonitrile (225 mg, 0.808 mmol), KOH (2.0 mL of 4.0 M, 8.0 mmol) and EtOH (2.3 mL) was heated at 90 °C for 2.5 d. The mixture was acidified with conc. HCl to ~pH 2 before it was concentrated under reduced pressure. The residue was subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give 6-*tert*-butyl-2-(2,4,6-trimethylphenyl)pyridine-3-carboxylic acid (40 mg, 17%) as a white solid. ESI-MS *m/z* calc. 297.2, found 298.4 $(M+1)^+$; Retention time: 1.77 min (3 min run).


Step 4: *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenyl)pyridine-3-carboxamide (Compound 26)



Sodium (3-nitrophenyl)sulfonylazanide (60. mg, 0.27 mmol) was added to a mixture of 6-*tert*-butyl-2-(2,4,6-trimethylphenyl)pyridine-3-carboxylic acid (40. mg, 0.13 mmol), *N*-[(dimethylamino)-1*H*-1,2,3-triazolo-[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (51 mg, 0.13 mmol) and *N,N*-dimethylformamide (800.0 µL) at room temperature. The mixture was allowed to stir at 70 °C overnight before it was filtered and subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05% HCl) to give the nitro intermediate, 6-*tert*-butyl-*N*-(3-nitrophenyl)sulfonyl-2-(2,4,6-trimethylphenyl)pyridine-3-carboxamide. The nitro intermediate was taken up in acetic acid (1 mL) before Zn (8.8 mg, 0.13 mmol) was added. The mixture was stirred at room temperature for 30 min before it was filtered. The filtrate was subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenyl)pyridine-3-carboxamide (Compound 26) (13 mg, 22%). ESI-MS *m/z* calc. 451.2, found 452.5 $(M+1)^+$; Retention time: 1.81 min (3 min run).

PREPARATION 2: *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(4-chlorophenyl)pyridine-3-carboxamide (Compound 223)


Step 1: 6-*tert*-butyl-2-chloro-pyridine-3-carboxamide



To a mixture of 6-*tert*-butyl-2-chloro-pyridine-3-carbonitrile (6.81 g, 35.0 mmol) and NaOH (38.5 mL of 1.0 M, 38.5 mmol) in EtOH (175 mL) was slowly added hydrogen peroxide (19.9 mL of 30% w/v, 175 mmol). The reaction mixture was stirred at room temperature for 45 min. before additional hydrogen peroxide (19.9 mL of 30% w/v, 175 mmol) was added. After 1 h, the mixture was slowly quenched with saturated sodium bisulfite solution, diluted with water, and extracted with ethyl acetate. The combined extracts were washed with water, dried over


sodium sulfate, and evaporated to give 6-*tert*-butyl-2-chloro-pyridine-3-carboxamide (7.21 g, 97%) as a white solid. ESI-MS *m/z* calc. 212.1, found 212.9 $(M+1)^+$; Retention time: 0.46 min (1 min run).

Step 2: 6-*tert*-butyl-2-chloro-*N*-(3-nitrophenyl)sulfonyl-pyridine-3-carboxamide



To a solution of 6-(*tert*-butyl)-2-chloronicotinamide (3.19 g, 15.0 mmol) and 3-nitrobenzenesulfonyl chloride (3.32 g, 15.0 mmol) in THF (100.0 mL) was slowly added NaH (1.80 g, 60%, w/w, 45.0 mmol) in portions. The mixture was allowed to stir for 2h at room temperature before it was quenched with saturated aqueous ammonium chloride solution. The mixture was extracted with ethyl acetate (3x). The combined extracts were washed with water, dried over sodium sulfate, and evaporated. The residue was taken up in diethyl ether. The solids were filtered, washed with diethyl ether, and dried under vacuum. The filtrate was evaporated and the residue was purified by silica gel chromatography (1-11% methanol/CH_2Cl_2). The two batches (solid from ether and product from column) were combined to give 6-(*tert*-butyl)-2-chloro-*N*-((3-nitrophenyl)sulfonyl)nicotinamide (3.56 g, 60%) as light yellow solid. ESI-MS *m/z* calc. 397.1, found 398.0 $(M+1)^+$; Retention time: 0.66 min (1 min run).


Step 3: *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-chloro-pyridine-3-carboxamide



To a solution of 6-(*tert*-butyl)-2-chloro-*N*-((3-nitrophenyl)sulfonyl)nicotinamide (1.40 g, 3.52 mmol) in glacial acetic acid (140 mL) was added zinc (5.76 g, 88.0 mmol). The reaction mixture was allowed to stir at room temperature for 1 h before it was filtered and concentrated under reduced pressure. The crude product was subjected silica gel column chromatography (0-5% methanol/CH_2Cl_2) to provide *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-chloro-pyridine-3-carboxamide (1.12 g, 87%) as an orange sticky foam. ESI-MS *m/z* calc. 367.1, found 368.6

$(M+1)^+$; Retention time: 1.09 min (3 min run).

Step 4: *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(4-chlorophenyl)pyridine-3-carboxamide (Compound 223)



To (4-chlorophenyl)boronic acid (26 mg, 0.16 mmol) was added a solution of *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-chloro-pyridine-3-carboxamide (50. mg, 0.14 mmol) in DMA (0.5 mL). Na_2CO_3 (136 μL, 2.0 M, 0.27 mmol) and dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (5 mg, 0.007 mmol) were added and the vessel was sealed under nitrogen gas and heated under microwave irradiation at 120 °C for 20 minutes. The reaction mixture was cooled to room temperature, filtered and subjected to preparatory-HPLC (10-99% water/CH_3CN with 0.05 mM HCl) to give *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(4-chlorophenyl)pyridine-3-carboxamide (Compound 223) (15 mg, 25%). ESI-MS *m/z* calc. 443.1, found 444.1 $(M+1)^+$; Retention time: 1.81 min (3 min run).

The following compounds were synthesized using the procedures described herein:

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(4-*tert*-butylphenyl)pyridine-3-carboxamide (Compound 252),

N-(3-aminophenyl)sulfonyl-8,8-dimethyl-2-phenyl-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 685),

(3-aminophenyl)sulfonyl-[8,8-dimethyl-2-(p-tolyl)-6,7-dihydro-5*H*-quinoline-3-carbonyl]azanide (Compound 1231),

N-(4-aminophenyl)sulfonyl-6-*tert*-butyl-2-(cyclohexen-1-yl)pyridine-3-carboxamide (Compound 880),

N-(4-aminophenyl)sulfonyl-8,8-dimethyl-2-phenyl-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 1190),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(4-fluorophenyl)pyridine-3-carboxamide (Compound 417),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-indan-5-yl-pyridine-3-carboxamide (Compound 308),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(3,4-dimethylphenyl)pyridine-3-carboxamide (Compound 274),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(3-*tert*-butylphenyl)pyridine-3-carboxamide (Compound 229),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(3,5-dimethylphenyl)pyridine-3-carboxamide (Compound 52),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(4-isopropylphenyl)pyridine-3-carboxamide (Compound 314),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(4-ethylphenyl)pyridine-3-carboxamide (Compound 359),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-[4-(trifluoromethyl)phenyl]pyridine-3-carboxamide (Compound 129),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(3-fluoro-5-methyl-phenyl)pyridine-3-carboxamide (Compound 237),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 301),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(4-fluoro-3-methyl-phenyl)pyridine-3-carboxamide (Compound 177),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(3-fluoro-4-methyl-phenyl)pyridine-3-carboxamide (Compound 349),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2,4-dimethylphenyl)pyridine-3-carboxamide (Compound 5),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(m-tolyl)pyridine-3-carboxamide (Compound 282),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(4-fluorophenyl)pyridine-3-carboxamide (Compound 46),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2-fluoro-5-methyl-phenyl)pyridine-3-carboxamide (Compound 1350) (Commpound 1350),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(3,4-difluorophenyl)pyridine-3-carboxamide (Compound 1217),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(o-tolyl)pyridine-3-carboxamide (Compound 1036),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(4-methoxyphenyl)pyridine-3-carboxamide (Compound 550),


N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2-fluoro-4-methyl-phenyl)pyridine-3-carboxamide (Compound 1331),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-phenyl-pyridine-3-carboxamide (Compound 542), and

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(3,4-difluorophenyl)pyridine-3-carboxamide (Compound 1217).


PREPARATION 3: *N*-(benzenesulfonyl)-6-*tert*-butyl-2-(*p*-tolyl)pyridine-3-carboxamide (Compound 74)

Step 1: 6-*tert*-butyl-2-chloro-pyridine-3-carboxylic acid



A solution of KOH (8.65 g, 154 mmol) in water (25 mL) was added to a solution of 6-*tert*-butyl-2-chloro-pyridine-3-carbonitrile (5.00 g, 25.7 mmol) in isopropanol (25 mL). The reaction mixture was allowed to stir at 90 °C overnight. After cooling to room temperature, the reaction mixture was diluted with ethyl acetate (75 mL) and was extracted with aqueous NaOH (1 N, 3×75 mL). The combined aqueous layers were acidified to pH 3 with the addition of aqueous HCl (1 N, 200 mL) and were extracted with ethyl acetate (2×75 mL). The combined organic layers were dried over sodium sulfate, filtered and concentrated to give 6-*tert*-butyl-2-chloro-pyridine-3-carboxylic acid (5.25 g, 96%) as a yellow solid. ESI-MS *m/z* calc. 213.1, found 214.4 $(M+1)^+$; Retention time: 1.09 min (3 min run). ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.39 (s, 1H), 8.26 (d, J = 8.1 Hz, 1H), 7.36 (d, J = 8.1 Hz, 1H), 1.38 (s, 9H).


Step 2: 6-*tert*-butyl-2-(p-tolyl)pyridine-3-carboxylic acid



6-*tert*-Butyl-2-chloro-pyridine-3-carboxylic acid (1.35 g, 6.32 mmol) and *p*-tolylboronic acid (1.03 g, 7.58 mmol) were suspended in DMA (14 mL). Aqueous sodium carbonate (6.3 mL of 2.0 M, 13 mmol) was added followed by dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (231 mg, 0.312 mmol). The reaction vessel was sealed under nitrogen and heated to 140 °C for 1.5 h in a microwave reactor. The reaction mixture was diluted with ethyl acetate (50 mL) and was washed with aqueous NaOH (1 M, 4×50 mL). The combined aqueous layers were acidified to pH 3 with 1 N HCl and then extracted with ethyl acetate (5×50 mL). The organic layers were combined, dried over sodium sulfate, filtered, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography (0-30% ethyl acetate/hexane) to give 6-*tert*-butyl-2-(*p*-tolyl)pyridine-3-carboxylic acid (740 mg, 43%) as a white solid. ESI-MS *m/z* calc. 269.1, found 270.5 $(M+1)^+$; Retention time: 1.53 min (3 min run).

The following compounds can be synthesized using the procedures described above: 8,8-dimethyl-2-(p-tolyl)-5,6,7,8-tetrahydroquinoline-3-carboxylic acid.

Step 3: *N*-(benzenesulfonyl)-6-*tert*-butyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 74)



To benzenesulfonamide (23 mg, 0.14 mmol) was added 6-*tert*-butyl-2-(*p*-tolyl)pyridine-3-carboxylic acid (35 mg, 0.13 mmol) in *N,N*-dimethylformamide (400 µL). A solution of *N*-[(dimethylamino)-1*H*-1,2,3-triazolo-[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (54 mg, 0.14 mmol) in *N,N*-dimethylformamide (125 µL) was added followed by solid potassium carbonate (54 mg, 0.39 mmol). The reaction mixture was allowed to stir at 80 °C for 1 h. Additional *N*-

[(dimethylamino)-1*H*-1,2,3-triazolo-[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (0.5 eq) was added and the reaction mixture was allowed to stir overnight at 80 °C. The mixture was cooled, filtered and subjected to preparatory-HPLC (10-99% water/acetonitrile with 0.05 mM HCl) to give *N*-(benzenesulfonyl)-6-*tert*-butyl-2-(*p*-tolyl)pyridine-3-carboxamide (Compound 74) (28 mg, 53%). ESI-MS *m/z* calc. 408.2, found 409.3 $(M+1)^+$; Retention time: 1.93 min (3 min run).

The following compounds can be synthesized using the procedures described herein:

6-*tert*-butyl-*N*-(3-ethylphenyl)sulfonyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 50),

6-*tert*-butyl-*N*-(3-methoxyphenyl)sulfonyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 327),

6-*tert*-butyl-*N*-(4-methoxyphenyl)sulfonyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 206),

6-*tert*-butyl-*N*-(4-cyanophenyl)sulfonyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 288),

6-*tert*-butyl-2-(p-tolyl)-*N*-(2-pyridylsulfonyl)pyridine-3-carboxamide (Compound 407),

6-*tert*-butyl-2-(p-tolyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 632),

6-*tert*-butyl-*N*-[(5-methyl-2-pyridyl)sulfonyl]-2-(p-tolyl)pyridine-3-carboxamide (Compound 1355),

6-*tert*-butyl-*N*-(2-cyanophenyl)sulfonyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 623),

6-*tert*-butyl-*N*-(2-methoxyphenyl)sulfonyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 1124),

N-(4-aminophenyl)sulfonyl-8,8-dimethyl-2-(p-tolyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 1307),

6-*tert*-butyl-*N*-(2-hydroxyphenyl)sulfonyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 1175),

N-[(6-amino-2-pyridyl)sulfonyl]-8,8-dimethyl-2-(p-tolyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 444),

8,8-dimethyl-2-(p-tolyl)-*N*-(1H-pyrazol-5-ylsulfonyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 1292),

6-*tert*-butyl-*N*-(4-hydroxyphenyl)sulfonyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 809),

6-*tert*-butyl-*N*-(3-hydroxyphenyl)sulfonyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 1265),

6-*tert*-butyl-*N*-[(6-methoxy-3-pyridyl)sulfonyl]-2-(p-tolyl)pyridine-3-carboxamide (Compound

512),

N-[(6-amino-3-pyridyl)sulfonyl]-8,8-dimethyl-2-(p-tolyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 635),

6-(3-fluoro-5-isopropoxy-phenyl)-2-(p-tolyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 849),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isopropoxy-phenyl)-2-(p-tolyl)pyridine-3-carboxamide (Compound 722),

6-(3-fluoro-5-isopropoxy-phenyl)-*N*-[(2-methoxy-3-pyridyl)sulfonyl]-2-(p-tolyl)pyridine-3-carboxamide (Compound 798),

6-*tert*-butyl-*N*-(1H-pyrazol-4-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 401),

6-(4-fluorophenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 518),

N-[(2-methoxy-3-pyridyl)sulfonyl]-8,8-dimethyl-2-(2,4,6-trimethylphenoxy)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 423),

6-*tert*-butyl-*N*-[[6-(difluoromethyl)-2-pyridyl]sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 501),

N-[(2-methoxy-3-pyridyl)sulfonyl]-2,6-bis(p-tolyl)pyridine-3-carboxamide (Compound 524),


6-(6-methoxy-5-methyl-3-pyridyl)-*N*-[(2-methoxy-3-pyridyl)sulfonyl]-2-(p-tolyl)pyridine-3-carboxamide (Compound 861),

6-*tert*-butyl-*N*-[(6-methoxy-2-pyridyl)sulfonyl]-2-(p-tolyl)pyridine-3-carboxamide (Compound 75), and

N-(4-aminophenyl)sulfonyl-6-*tert*-butyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 297).


PREPARATION 4: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 58)

Step 1: 6-*tert*-butyl-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-2-(*p*-tolyl)pyridine-3-carboxamide



N,N-diisopropyl ethyl amine (104 μL, 0.5940 mmol) was added to a mixture of 6-*tert*-butyl-2-(*p*-tolyl)pyridine-3-carboxylic acid (80. mg, 0.30 mmol), 6-fluoropyridine-2-sulfonamide (58 mg, 0.33 mmol), *N*-[(dimethylamino)-1*H*-1,2,3-triazolo-[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (136 mg, 0.356 mmol) and *N,N*-dimethylformamide (800 μL) at room temperature. The mixture was heated at 70 °C overnight before it was cooled. The mixture was partitioned between ethyl acetate and 1N HCl. The layers were separated and the aqueous layer was extracted with ethyl acetate (3x). The combined organics were washed with brine, dried over sodium sulfate, filtered, and concentrated. The residue was subjected to silica gel column chromatography (0-20% ethyl acetate/hexanes) to give 6-*tert*-butyl-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-2-(*p*-tolyl)pyridine-3-carboxamide (66 mg, 38%) as a yellowish solid. ESI-MS *m/z* calc. 427.1, found 428.4 $(M+1)^+$; Retention time: 0.7 min (1 min run).

Step 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(*p*-tolyl)pyridine-3-carboxamide (Compund 58)



A mixture of 6-*tert*-butyl-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-2-(*p*-tolyl)pyridine-3-carboxamide (55 mg, 0.13 mmol) and NH_4OH (30%, 860 μL) was heated at 150 °C for 30 min in a microwave reactor. The mixture was concentrated and the residue was subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(*p*-tolyl)pyridine-3-carboxamide (Compound 58) (17 mg, 31%). ESI-MS *m/z* calc. 424.2, found 425.4 $(M+1)^+$; Retention time: 1.63 min (3 min run).

The following compounds can be synthesized using the procedures described herein: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(m-tolyl)pyridine-3-carboxamide (Compound

31),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(cyclopenten-1-yl)pyridine-3-carboxamide (Compound 529),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(cyclohexen-1-yl)pyridine-3-carboxamide (Compound 478),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-ethoxyphenyl)-2-(p-tolyl)pyridine-3-carboxamide (Compound 409),


N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(4-fluorophenyl)pyridine-3-carboxamide (Compound 417),

and

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-phenyl-pyridine-3-carboxamide (Compound 334).

PREPARATION 5: 6-phenethyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid


Step 1: methyl 6-chloro-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate



NaH (60%, 2.13 g, 53.3 mmol) was added to a solution of 2,4,6-trimethylphenol (6.60 g, 48.5 mmol) and *N,N*-dimethylformamide (150 mL) at 0 °C. The mixture was stirred for 15 min before a solution of methyl 2,6-dichloronicotinate (10.0 g, 48.5 mmol) in *N,N*-dimethylformamide (20 mL) was added dropwise. The mixture was allowed to stir for 1 h at room temperature before water was added. The mixture was extracted with ethyl acetate (2x). The combined extracts were washed with water and brine, dried over sodium sulfate, filtered and concentrated. The residue was subjected to silica gel column chromatography (0-50% CH_2Cl_2/hexanes) to give methyl 6-chloro-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate


(3.58 g, 24%). ESI-MS *m/z* calc. 305.1, found 306.2 $(M+1)^+$; Retention time: 0.87 min (1 min run).

Step 2: methyl 6-phenethyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate



Bromo-(2-phenylethyl)zinc (3.9 mL of 0.5 M, 2.0 mmol) and dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (36 mg, 0.049 mmol) were stirred at room temperature under an atmosphere of nitrogen for 20 min. Methyl 6-chloro-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (150 mg, 0.49 mmol) was added to the reaction mixture and it was then heated at 150 °C for 10 minutes in a microwave reactor. A saturated aqueous solution of ethylenediaminetetraacetic acid disodium salt (2 mL) and a saturated aqueous solution of ammonium chloride were added to the reaction mixture. After stirring for 30 minutes, the layers were separated and the aqueous layer was extracted with dichloromethane. The combined organic layers were evaporated to dryness and the residue was subjected to preparatory-HPLC (20-80% acetonitrile/water with 5 mM HCl) to yield methyl 6-(2-phenylethyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (56 mg, 31%). ESI-MS *m/z* calc. 375.2, found 376.4 $(M+1)^+$; Retention time: 0.87 min (1 min run).

Step 3: 6-phenethyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid




Methyl 6-(2-phenylethyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (56 mg, 0.15 mmol) was dissolved in methanol (5mL). NaOH (245 µL of 4.0 M, 0.98 mmol) was added

and the mixture was stirred at room temperature overnight. The solvent was removed and the pH of the residue was adjusted to ~3 with 1N HCl. The precipitate was filtered, washed with water and dried to give 6-phenethyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (21 mg, 36%). ESI-MS *m/z* calc. 361.2, found 362.5 $(M+1)^+$; Retention time: 0.77 min (1 min run).

The following compound can be synthesized using the procedures described herein: 6-benzyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid.


PREPARATION 6: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 228)

Step 1: 6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carbonitrile




6-*tert*-Butyl-2-chloro-pyridine-3-carbonitrile (5.00 g, 25.7 mmol), 2,4,6-trimethylphenol (3.50 g, 25.7 mmol), K_2CO_3 (3.91 g, 28.3 mmol) and *N,N*-dimethylformamide (10 mL) were added to a 40 mL vial. The mixture was stirred overnight at 95 °C. The reaction mixture was diluted with H_2O (10 mL) and was extracted with ethyl acetate (3 x 20 mL). The combined organic layers were washed with brine, dried over sodium sulfate and concentrated. The residue was subjected to silica gel column chromatography (0-20% ethyl acetate/hexanes) to give 6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carbonitrile (6.70 g, 84%) as a white solid. ESI-MS *m/z* calc. 294.2, found 295.2 $(M+1)^+$; Retention time: 2.31 min (3 min run). ^1H NMR (400 MHz, DMSO) δ 8.31 (d, *J* = 8.0 Hz, 1H), 7.25 (d, *J* = 7.9 Hz, 1H), 6.96 (s, 2H), 2.27 (s, 3H), 2.00 (s, 6H), 1.07 (s, 9H).

Step 2: 6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid



A mixture of 6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carbonitrile (4.50 g, 15.3 mmol), KOH (5.15 g, 91.7 mmol), EtOH (135 mL) and H_2O (32 mL) was heated at 90 °C overnight. Additional KOH (2.00 g, 35.6 mmol) was added and the mixture was heated at 90 °C overnight. The mixture was cooled to room temperature, concentrated and acidified to pH ~2 with 2N HCl. The solids were collected via filtration and were placed in a vacuum oven at 50 °C overnight to give 6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (4.64 g, 96%). ^{1}H NMR (400 MHz, DMSO) δ 8.14 (d, J = 7.8 Hz, 1H), 7.08 (d, J = 7.8 Hz, 1H), 6.91 (s, 2H), 2.26 (s, 3H), 1.97 (s, 6H), 1.04 (s, 9H).


Step 3: 6-*tert*-butyl-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide



N-ethyl-*N*-isopropylpropan-2-amine (560 µL, 3.19 mmol) was added to a mixture of 6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (500. mg, 1.56 mmol), 6-fluoropyridine-2-sulfonamide (309 mg, 1.76 mmol), *N*-[(dimethylamino)-1*H*-1,2,3-triazolo-[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (728 mg, 1.91 mmol) and *N,N*-dimethylformamide (5 mL) at room temperature. The mixture was heated at 70 °C overnight before it was cooled. The mixture was partitioned between ethyl acetate and 1N HCl. The layers were separated and the aqueous layer was extracted with ethyl acetate (3x). The combined organics were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was subjected to silica gel column chromatography (0-20% ethyl

acetate/hexanes), then a second purification by silica gel column chromatography (0-10% ethyl acetate/hexanes) to give 6-*tert*-butyl-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (230 mg, 23%) as a white solid. ESI-MS *m/z* calc. 471.2, found 472.4 $(M+1)^+$; Retention time: 2.32 min (3 min run).

Step 4: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 228)



6-*tert*-Butyl-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (77 mg, 0.16 mmol) was dissolved in ammonium hydroxide (10 mL of 30 %w/w) and irradiated in the microwave reactor for 1 h at 150 °C. The mixture was evaporated and the crude material was purified by preparatory-HPLC (10-99% acetonitrile/water with 5 μM HCl). The material was further purified by silica gel chromatography (0-100% ethyl acetate/hexanes) to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 228) (27 mg, 35%). ^{1}H NMR (400 MHz, DMSO) δ 12.10 (s, 1H), 8.01 (d, J = 7.8 Hz, 1H), 7.67 - 7.58 (m, 1H), 7.20 (d, J = 7.3 Hz, 1H), 7.11 (d, J = 7.9 Hz, 1H), 6.90 (s, 2H), 6.69 (d, J = 8.6 Hz, 1H), 6.49 (s, 2H), 2.26 (s, 3H), 1.97 (s, 6H), 1.05 (s, 9H). ESI-MS *m/z* calc. 468.2, found 469.3 $(M+1)^+$; Retention time: 2.11 min (3 min run).

The following compounds can be synthesized using the procedures described herein:

6-*tert*-butyl-*N*-(1H-indol-4-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 275),

N-(benzenesulfonyl)-6-*tert*-butyl-2-(2-isopropylphenoxy)pyridine-3-carboxamide (Compound 147),

6-*tert*-butyl-*N*-[[6-(dimethylamino)-2-pyridyl]sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 468),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1,1,2,2,2-pentafluoroethyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 853),

N-(4-aminophenyl)sulfonyl-6-(1-methylcyclopropyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1297),

N-(2-aminothiazol-4-yl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1294),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-methylcyclopropyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1310),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(1-methylcyclopropyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 854),

N-(3-aminophenyl)sulfonyl-6-(1-methylcyclopropyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 826),

6-(3-fluoro-5-isopropoxy-phenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 914),

N-[(6-amino-2-pyridyl)sulfonyl]-6-prop-1-ynyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1172),

6-*tert*-butyl-*N*-(2-methylpyrazol-3-yl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 911),

N-[(6-amino-2-pyridyl)sulfonyl]-5,6-dimethyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 841),

N-[[6-(2-hydroxyethylamino)-2-pyridyl]sulfonyl]-6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 595),

N-[[6-(dimethylamino)-2-pyridyl]sulfonyl]-6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 869),

N-[[6-(2-aminoethylamino)-2-pyridyl]sulfonyl]-6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1027),

N-[[6-(methylamino)-2-pyridyl]sulfonyl]-6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1032),

N-[[6-(2-dimethylaminoethylamino)-2-pyridyl]sulfonyl]-6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1122),

N-[(6-cyano-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 771),

N-[(6-amino-2-pyridyl)sulfonyl]-4-methyl-6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-

carboxamide (Compound 1102),
6-*tert*-butyl-*N*-[[6-(2-hydroxyethylamino)-2-pyridyl]sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 916),
6-*tert*-butyl-*N*-[[6-(2-dimethylaminoethylamino)-2-pyridyl]sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 947),
6-[[6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carbonyl]sulfamoyl]pyridine-2-carboxylic acid (Compound 461),
6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[[6-(hydroxymethyl)-2-pyridyl]sulfonyl]-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 1340),
N-[[6-(2-aminoethylamino)-2-pyridyl]sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1054),
N-[(6-amino-2-pyridyl)sulfonyl]-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1137),
N-(2-azidophenyl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide,
6-methyl-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 670),
N-[(6-amino-2-pyridyl)sulfonyl]-5-phenethyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1253),
N-[(6-amino-2-pyridyl)sulfonyl]-6-[1-(trifluoromethyl)cyclopropyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 621),
N-[(6-amino-2-pyridyl)sulfonyl]-5-benzyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1233),
N-[[6-(isopropylamino)-2-pyridyl]sulfonyl]-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 482),
N-[[6-(2-methoxyethylamino)-2-pyridyl]sulfonyl]-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (compound 372),
N-[(6-amino-2-pyridyl)sulfonyl]-8,8-dimethyl-2-(2,4,6-trimethylphenoxy)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 182),
N-(benzenesulfonyl)-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 341),
6-*tert*-butyl-*N*-[3-(methylamino)phenyl]sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-

carboxamide (Compound 289),
6-*tert*-butyl-*N*-[3-(dimethylamino)phenyl]sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 311),
6-*tert*-butyl-*N*-(2-cyanophenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 78),
N-(benzenesulfonyl)-6-*tert*-butyl-2-(1-methylindol-7-yl)oxy-pyridine-3-carboxamide (Compound 326),
6-*tert*-butyl-*N*-(1-methylindol-4-yl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 146),
N-[(6-amino-5-fluoro-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide,
6-*tert*-butyl-*N*-[(6-methoxy-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 34),
6-*tert*-butyl-*N*-(4-pyridylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 117),
N-[(2-amino-3-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 235),
N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 590),
N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methylphenoxy)pyridine-3-carboxamide (Compound 778),
6-*tert*-butyl-*N*-(2-methoxyphenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 115),
6-*tert*-butyl-*N*-[(2-methoxy-3-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 191),
6-*tert*-butyl-*N*-(2-pyridylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 222),
N-[(2-amino-4-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 283),
6-*tert*-butyl-*N*-[(2-methoxy-4-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 217),

6-*tert*-butyl-*N*-(3-pyridylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 330),

6-*tert*-butyl-*N*-(3-methoxyphenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 56),

N-(benzenesulfonyl)-6-(trifluoromethyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 244),

6-*tert*-butyl-*N*-[(3-methoxy-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 92),

N-(benzenesulfonyl)-2-benzyloxy-6-*tert*-butyl-pyridine-3-carboxamide (Compound 130),

6-*tert*-butyl-2-(2,4-dimethylphenoxy)-*N*-[(2-methoxy-3-pyridyl)sulfonyl]pyridine-3-carboxamide (Compound 89),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4-dimethylphenoxy)pyridine-3-carboxamide (Compound 204),

N-(4-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 11),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(1-phenylethoxy)pyridine-3-carboxamide (Compound 219),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)-5,6,7,8-tetrahydroquinoline-3-carboxamide (Compound 285),

N-[(5-amino-3-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 163),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2-methylphenoxy)pyridine-3-carboxamide (Compound 296),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(4-cyano-2,6-dimethyl-phenoxy)pyridine-3-carboxamide (Compound 111),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2,4-dimethylphenoxy)pyridine-3-carboxamide (Compound 251),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethylcyclopentoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 503),

N-(benzenesulfonyl)-6-*tert*-butyl-2-(3-methoxyphenoxy)pyridine-3-carboxamide (Compound 101),

6-*tert*-butyl-*N*-[(6-oxo-1*H*-pyridin-2-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 305),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2-cyano-4,6-dimethyl-phenoxy)pyridine-3-carboxamide (Compound 221),

N-(3-aminophenyl)sulfonyl-2-(2,6-dimethylphenoxy)-6-(trifluoromethyl)pyridine-3-carboxamide (Compound 243),

N-(3-aminophenyl)sulfonyl-6-(trifluoromethyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 96),

6-*tert*-butyl-*N*-[(4-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 107),

N-(benzenesulfonyl)-6-*tert*-butyl-2-(2-pyridylmethoxy)pyridine-3-carboxamide (Compound 154),

6-*tert*-butyl-*N*-[(2-oxo-1*H*-pyridin-4-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 113),

6-*tert*-butyl-2-(2,4-dimethylphenoxy)-*N*-[(2-hydroxy-3-pyridyl)sulfonyl]pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-(trifluoromethyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 175),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(trifluoromethyl)pyridine-3-carboxamide (Compound 246),

N-[(6-amino-2-pyridyl)sulfonyl]-6-cyclopropyl-4-(trifluoromethyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 54),

N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-phenoxy-pyridine-3-carboxamide (Compound 272),

N-(3-aminophenyl)sulfonyl-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 13),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-methylcyclopentoxy)pyridine-3-carboxamide (Compound 452),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-phenoxy-pyridine-3-carboxamide (Compound 376),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[1-(3-fluorophenyl)ethoxy]pyridine-3-carboxamide (Compound 562),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(3-ethyloxetan-3-yl)methoxy]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 618),

N-[[6-(3-aminopropylamino)-2-pyridyl]sulfonyl]-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 982),

N-[[6-(cyclopropylamino)-2-pyridyl]sulfonyl]-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 594),

6-methyl-*N*-[[6-(methylamino)-2-pyridyl]sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 646),

N-[[6-(2-aminoethylamino)-2-pyridyl]sulfonyl]-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 792),

6-methyl-*N*-[(6-morpholino-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 883),

6-methyl-*N*-[[6-(4-methylpiperazin-1-yl)-2-pyridyl]sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1018),

N-[[6-(dimethylamino)-2-pyridyl]sulfonyl]-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1250),

2-(2,4-dimethylphenoxy)-*N*-[(2-methoxy-3-pyridyl)sulfonyl]-8,8-dimethyl-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 1347),

2-(2,4-dimethylphenoxy)-8,8-dimethyl-*N*-(1H-pyrazol-3-ylsulfonyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 1246),

6-*tert*-butyl-*N*-[(2-methoxy-6-methyl-3-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1153),

8,8-dimethyl-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 715),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(oxetan-2-ylmethoxy)pyridine-3-carboxamide (Compound 640),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(cyclohexoxy)pyridine-3-carboxamide (Compound 649),

2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 671),

6-(3-fluoro-5-isobutoxy-phenyl)-2-phenoxy-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 679),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-tetrahydropyran-3-yloxy-pyridine-3-carboxamide (Compound 706),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methylphenoxy)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 738),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-methylphenoxy)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 758),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2-cyano-4,6-dimethyl-phenoxy)pyridine-3-carboxamide (Compound 767),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2,2-difluorocyclopropyl)methoxy]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 769),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[1-(o-tolyl)ethoxy]pyridine-3-carboxamide (Compound 782),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3-methyloxetan-3-yl)methoxy]pyridine-3-carboxamide (Compound 824),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4-difluorophenoxy)pyridine-3-carboxamide (Compound 829),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[1-(4-fluorophenyl)ethoxy]pyridine-3-carboxamide (Compound 867),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(oxetan-3-yloxy)pyridine-3-carboxamide (Compound 872),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(1-phenylethoxy)pyridine-3-carboxamide (Compound 874),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2,2-dimethylcyclopropyl)methoxy]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 894),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(tetrahydropyran-4-ylmethoxy)pyridine-3-carboxamide (Compound 933),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4,4-difluorocyclohexoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 961),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(1-methyl-2-phenyl-ethoxy)pyridine-3-carboxamide (Compound 967),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,6-dimethylcyclohexoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 978),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2-isopropylphenoxy)pyridine-3-carboxamide (Compound 1031),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(1-phenylpropoxy)pyridine-3-carboxamide (Compound 1045),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-phenoxy-pyridine-3-carboxamide (Compound 1046),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3-fluorooxetan-3-yl)methoxy]pyridine-3-carboxamide (Compound 1053),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(1-phenylpropoxy)pyridine-3-carboxamide (Compound 1068),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(1-methylindol-7-yl)oxy-pyridine-3-carboxamide (Compound 1071),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[[(1R,5S)-3-bicyclo[3.1.0]hexanyl]oxy]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1077),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(4-cyano-2,6-dimethyl-phenoxy)pyridine-3-carboxamide (Compound 1089),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(4-fluoro-2-methyl-phenoxy)pyridine-3-carboxamide (Compound 1104),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methylcyclohexoxy)pyridine-3-carboxamide (Compound 1108),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(cyclobutoxy)pyridine-3-carboxamide (Compound 1154),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(1-cyclopropylethoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1165),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(1R,2R,4S)-norbornan-2-yl]oxy-pyridine-3-carboxamide (Compound 1216),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-norbornan-2-yloxy-pyridine-3-carboxamide (Compound 1227),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(cyclopropylmethoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1229),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(1-cyclobutylethoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1238),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,6-dimethylphenoxy)pyridine-3-carboxamide (Compound 1243),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(cycloheptoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1263),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(1-methyl-2-phenyl-ethoxy)pyridine-3-carboxamide (Compound 1314),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2-methylphenoxy)pyridine-3-carboxamide (Compound 1318),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-tetrahydropyran-4-yloxy-pyridine-3-carboxamide (Compound 1324),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-norbornan-2-yloxy-pyridine-3-carboxamide (Compound 1349),

N-[(6-amino-2-pyridyl)sulfonyl]-7,7-dimethyl-2-(2,4,6-trimethylphenoxy)-6,8-dihydro-5*H*-quinoline-3-carboxamide (Compound 665),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,5-dimethylcyclopentoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 416),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-tetrahydrofuran-3-yloxy-pyridine-3-carboxamide (Compound 433),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-methylcyclohexoxy)pyridine-3-carboxamide (Compound 438),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 985),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1461),

3-[[6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carbonyl]sulfamoyl]benzoic acid (Compound 205),

N-(3-aminophenyl)sulfonyl-6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 231),

N-[(6-amino-2-pyridyl)sulfonyl]-6-phenethyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 268),

N-[(6-amino-2-pyridyl)sulfonyl]-6-benzyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1),

N-(benzenesulfonyl)-2-(2,4,6-trimethylphenoxy)quinoline-3-carboxamide (Compound 215),

N-[(6-amino-2-pyridyl)sulfonyl]-8-methyl-2-(2,4,6-trimethylphenoxy)quinoline-3-carboxamide (Compound 70),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)quinoline-3-carboxamide (Compound 91),

N-(3-aminophenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)quinoline-3-carboxamide (Compound 260),


N-(4-pyridylsulfonyl)-2-(2,4,6-trimethylphenoxy)quinoline-3-carboxamide (Compound 240),

6-*tert*-butyl-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 214),

6-*tert*-butyl-*N*-[[6-(methylamino)-2-pyridyl]sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 262), and

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-8,8-dimethyl-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 15).

PREPARATION 7: 6-*tert*-butyl-*N*-(2-hydroxyphenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 131)



6-*tert*-Butyl-*N*-(2-methoxyphenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 115) (99 mg, 0.21 mmol) was taken up in CH_2Cl_2 (10 mL). The mixture was cooled to -78 °C before BBr_3 (1.0 mL, 1M, 1.0 mmol) was added slowly. The mixture allowed to warm to room temperature overnight. The reaction mixture was diluted with saturated $NaHCO_3$ (10 mL) and was extracted with ethyl acetate (2 x 5 mL). The organic layers were dried over sodium sulfate and concentrated. The residue was subjected to preparatory-HPLC (20-80% acetonitrile/water with 5 mM HCl) to give 6-*tert*-butyl-*N*-(2-hydroxyphenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 131) (63 mg, 66%). ^{1}H NMR (400 MHz, DMSO) δ 11.66 (s, 1H), 11.04 (s, 1H), 8.05 (d, J = 7.8 Hz, 1H), 7.83 (d, J = 7.9 Hz, 1H), 7.52 (s, 1H), 7.16 (d, J = 7.9 Hz, 1H), 7.02 (s, 2H), 6.93 (s, 2H), 2.26 (s, 3H), 1.98 (s, 6H), 1.04 (s, 9H). ESI-MS *m/z* calc. 468.2, found 469.0 $(M+1)^+$; Retention time: 2.16 min (3 min run).


The following compound can be synthesized using the procedures described herein: 6-*tert*-butyl-*N*-(3-hydroxyphenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 47)

and

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(3-hydroxyphenyl)sulfonyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 395).


PREPARATION 8: 6-*tert*-butyl-2-(1-phenylethyl)pyridine-3-carboxylic acid

Step 1: 6-*tert*-butyl-2-(1-phenylethyl)pyridine-3-carbonitrile



A mixture of bromo(1-phenylethyl)zinc (4.1 mL of 0.5 M, 2.1 mmol) and dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (38 mg, 0.051 mmol) was stirred at room temperature under an atmosphere of nitrogen for 20 min. 6-*tert*-Butyl-2-chloro-pyridine-3-carbonitrile (100 mg, 0.51 mmol) was added and the reaction mixture was heated at 150 °C for 10 min in a microwave reactor. The mixture was concentrated and the residue was subjected to silica gel column chromatography (0-20% ethyl acetate/hexanes) to give 6-*tert*-butyl-2-(1-phenylethyl)pyridine-3-carbonitrile (120 mg, 80.%). ESI-MS *m/z* calc. 264.2, found 265.4 $(M+1)^+$; Retention time: 2.28 min (3 min run).

Step 2: 6-*tert*-butyl-2-(1-phenylethyl)pyridine-3-carboxylic acid




A mixture of 6-*tert*-butyl-2-(1-phenylethyl)pyridine-3-carbonitrile (120 mg, 0.45 mmol), KOH (1.1 mL of 4.0 M, 4.4 mmol) and EtOH (1.2 mL) was heated at 90 °C overnight. The mixture was concentrated and conc. HCl was added to the residue. The solids were collected

and washed with water. The solids were placed in a vacuum oven at 60 °C overnight to give a ~ 1:1 mixture of 6-*tert*-butyl-2-(1-phenylethyl)pyridine-3-carboxylic acid [ESI-MS *m/z* calc. 283.2, found 284.3 $(M+1)^+$; Retention time: 0.78 min (1 min run)] and 6-*tert*-butyl-2-(1-phenylethyl)pyridine-3-carboxamide [ESI-MS *m/z* calc. 282.2, found 283.4 $(M+1)^+$; Retention time: 0.69 min (1 min run)].

The following compound can be synthesized using the procedures described herein: 6-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyridine-3-carboxylic acid, *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(1-phenylethyl)pyridine-3-carboxamide (Compound 271), and *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyridine-3-carboxamide (Compound 346).


PREPARATION 9: 6-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]pyridine-2-carboxylic acid

Step 1: 5-bromo-2-*tert*-butylpyridine




tert-Butylmagnesium chloride (60 mL of a 1.0 N solution in THF, 60 mmol) was slowly added to a greenish suspension of CuCN (2.7 g, 30 mmol) in THF (60 mL) cooled to -78 °C and the resulting gray mixture was stirred at -78 °C for 1 h. A solution of 5-bromo-2-iodopyridine (4.3 g, 15 mmol) in THF (10 mL) was then added. The reaction mixture was stirred at -78 °C for 1 h then at 0 °C for 3 h before being allowed to warm up gradually to room temperature overnight. Ammonium hydroxide (80 mL of 20%) was added and the resulting mixture was filtered and washed with ethyl acetate (100 mL). The filtrate was washed with brine (40 mL), dried over $MgSO_4$, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (0-10% ethyl acetate/ heptanes) to afford 5-bromo-2-*tert*-butylpyridine (2.2 g, 68%; contains 20% heptanes) as a colorless oil. ^1H NMR (300 MHz, $CDCl_3$) δ 1.43 (s, 9H), 7.23 (dd, J = 8.5, 0.9 Hz, 1H), 7.71 (dd, J = 8.5, 2.3 Hz, 1H), 8.60 (d, J = 2.3 Hz, 1H). $[M+H]^+$ = 214.0.

Step 2: 5-bromo-2-*tert*-butylpyridine-*N*-oxide




3-Chlorobenzoperoxoic acid (4.40 g of 77% grade, 17.9 mmol) was added to a solution of 5-bromo-2-*tert*-butylpyridine (3.70 g, 13.8 mmol) in dichloromehtane (45 mL). The reaction mixture was stirred at room temperature overnight. $Ca(OH)_2$ (4.1 g, 44 mmol) was added and the resulting mixture was then stirred at room temperature for 30 min before being filtered through a pad of Celite. The filtrate was concentrated under reduced pressure and the residue was purified by silica gel chromatography (0-50% ethyl acetate/heptanes) to afford 5-bromo-2-*tert*-butylpyridine-*N*-oxide (2.7 g, 85%) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.48 (s, 9H), 7.19 (d, J = 8.8 Hz, 1H), 7.31 (dd, J = 8.8, 1.8 Hz, 1H), 8.33 (d, J = 1.8 Hz, 1H). $[M+H]^+$ = 230.0.

Step 3: 6-*tert*-butylpyridine-2-carbonitrile




Trimethylsilanecarbonitrile (8.8 mL, 70 mmol) and Et_3N (7.4 mL, 53 mmol) were added to a solution of 5-bromo-2-*tert*-butylpyridine-*N*-oxide (2.7 g, 12 mmol) in acetonitrile (20 mL) at room temperature. The reaction mixture was then heated at 100 °C overnight. Additional trimethylsilanecarbonitrile (2.4 mL, 19 mmol) and Et_3N (2.1 mL, 15 mmol) were added and the reaction mixture was heated at reflux again overnight. Most of the acetonitrile was removed under reduced pressure and the residue was dissolved in ethyl acetate (100 mL). The organic layer was washed with a saturated aqueous solution of $NaHCO_3$ (20 mL), brine (20 mL), dried over $MgSO_4$, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (0-15% ethyl acetate/heptanes) to afford 3-bromo-6-*tert*-butylpyridine-2-carbonitrile (2.6 g, 93%) as a colorless oil. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.34 (s, 9H), 7.42 (d, J = 8.5 Hz, 1H), 7.89 (d, J = 8.5 Hz, 1H). $[M+H]^+$ = 238.9.

Step 4: 6-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]pyridine-2-carbonitrile



Tetrakis(triphenylphosphine)palladium(0) (530 mg, 0.46 mmol) was added to a solution of 3-bromo-6-*tert*-butylpyridine-2-carbonitrile (1.1 g, 4.6 mmol) in THF (12 mL) in a sealed tube. The mixture was evacuated and backfilled with nitrogen three times, then a stock solution of (2,4,6-trimethylphenyl)methylzinc bromide (9.2 mL of a 1.0 M solution in THF, 9.2 mmol) was added at room temperature. The orange solution was stirred at 60 °C overnight. The reaction mixture was quenched with 5% citric acid (20 mL) and the resulting mixture was extracted with ethyl acetate (2 x 40 mL). The combined organic layers were washed with water (2 x 15 mL), brine (15 mL), dried over $MgSO_4$, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (5-30% ethyl acetate/heptanes) to afford 6-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]pyridine-2-carbonitrile (1.2 g, 89%) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.32 (s, 9H), 2.17 (s, 6H), 2.30 (s, 3H), 4.17 (s, 2H), 6.91 (s, 2H), 6.97 (d, J = 8.5 Hz, 1H), 7.32 (d, J = 8.5 Hz, 1H). $[M+H]^+$ = 293.2.

Step 5: 6-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]pyridine-2-carboxylic acid




6-*tert*-Butyl-3-[(2,4,6-trimethylphenyl)methyl]pyridine-2-carbonitrile (900. mg, 3.08 mmol) was dissolved in acetic acid (8 mL). Concentrated HCl (8 mL) was added and the reaction mixture was stirred at 120 °C for 3 d. Additional acetic acid (2 mL) and concentrated HCl (2 mL) were added and the reaction mixture was stirred at 120 °C for another 24 h. Most of the acetic acid was removed under reduced pressure and the resulting heterogeneous mixture was basified with 25% aqueous NaOH to a pH of about 8-10. An aqueous solution of 5% citric acid

(20 mL) was then added and the aqueous layer was extracted with methyl tert-butyl ether (2 x 50 mL). The combined organic layers were washed with water (2 x 20 mL), brine (20 mL), dried over $MgSO_4$, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (0-50% ethyl acetate/heptanes) to afford 6-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]pyridine-2-carboxylic acid (660 mg, 68.8%) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.36 (s, 9H), 2.12 (s, 6H), 2.31 (s, 3H), 4.57 (s, 2H), 6.91 (s, 2H), 7.07 (d, J = 8.2 Hz, 1H), 7.39 (d, J = 8.2 Hz, 1H), 12.25 (br s, 1H). $[M+H]^+$ = 312.2.

The following compounds can be synthesized using the procedures described herein: *N*-(benzenesulfonyl)-6-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]pyridine-2-carboxamide (Compound 22), and *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]pyridine-2-carboxamide (Compound 313).


PREPARATION 10: 6-*tert*-butyl-3-(2,4,6-trimethylphenoxy)pyridine-2-carboxylic acid

Step 1: 6-*tert*-butyl-3-(2,4,6-trimethylphenoxy)pyridine-2-carbonitrile



A mixture of 3-bromo-6-*tert*-butylpyridine-2-carbonitrile (1.0 g, 4.2 mmol), 2,4,6-trimethylphenol (0.57 g, 4.2 mmol) and Cs_2CO_3 (2.0 g, 6.3 mmol) in DMSO (8 mL) was heated at 100 °C overnight. Once cooled to room temperature, ethyl acetate (70 mL) was added and the organic layer was washed with water (2 x 20 mL), brine (20 mL), dried over $MgSO_4$, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography, (0-30% ethyl acetate/heptanes to afford 6-*tert*-butyl-3-(2,4,6-trimethylphenoxy)pyridine-2-carbonitrile (1.0 g, 81%) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.33 (s, 9H), 2.08 (s, 6H), 2.30 (s, 3H), 6.73 (d, J = 9.1 Hz, 1H), 6.92 (s, 2H), 7.32 (d, J = 8.8 Hz, 1H). $[M+H]^+$ = 295.1.

Step 2: 6-*tert*-butyl-3-(2,4,6-trimethylphenoxy)pyridine-2-carboxylic acid




6-*tert*-Butyl-3-(2,4,6-trimethylphenoxy)pyridine-2-carbonitrile (700 mg, 2.38 mmol) was dissolved in acetic acid (9 mL). Concentrated HCl (9 mL) was added and the reaction mixture was heated at 120 °C for 2 d. Once cooled to room temperature, the reaction mixture was concentrated to dryness under reduced pressure. The residue was suspended in water (20 mL) and 2N NaOH (3 mL) was added to provide a clear aqueous solution. The aqueous layer was washed with methyl tert-butyl ether (2 x 10 mL), acidified by the addition of 5% citric acid (20 mL) and the resulting mixture was extracted with ethyl acetate (2 x 50 mL). The combined organic layers were washed with water (2 x 20 mL), brine (20 mL), dried over $MgSO_4$, filtered and concentrated under reduced pressure. The residue was triturated with heptanes (10 mL) at 0 °C to afford 6-*tert*-butyl-3-(2,4,6-trimethylphenoxy)pyridine-2-carboxylic acid (574 mg, 77%) as an off-white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.37 (s, 9H), 2.08 (s, 6H), 2.30 (s, 3H), 6.89 - 6.92 (m, 3H), 7.42 (d, J = 8.9 Hz, 1H), 11.68 (br s, 1H). $[M+H]^+$ = 314.2.

The following compounds can be synthesized using the procedures described herein: *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-3-(2,4,6-trimethylphenoxy)pyridine-2-carboxamide (Compound 106) and *N*-(benzenesulfonyl)-6-*tert*-butyl-3-(2,4,6-trimethylphenoxy)pyridine-2-carboxamide (Compound 350).


PREPARATION 11: *N*-(3-aminophenyl)sulfonyl-5-cyano-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 124)

Step 1: ethyl 5-cyano-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate




Ethyl 2-chloro-5-cyano-6-methyl-pyridine-3-carboxylate (518 mg, 2.31 mmol), 2,4,6-trimethylphenol (377 mg, 2.77 mmol), and cesium carbonate (1.50 g, 4.61 mmol) were combined in *N,N*-dimethylformamide (5 mL) and heated at 90 °C for 1 h. The reaction mixture was cooled and partitioned between ethyl acetate and 1N HCl. The organics were separated, washed with brine, dried over sodium sulfate and evaporated to give ethyl 5-cyano-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate. ESI-MS *m/z* calc. 324.2, found 325.2 $(M+1)^+$; Retention time: 0.77 min (1 min run).

Step 2: 5-cyano-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid




Crude ethyl 5-cyano-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate was dissolved in methanol (12 mL) before NaOH (12 mL of 1.0 M, 12 mmol) was added. The mixture was heated at 70 °C for 3 h. The mixture was cooled and made acidic by the addition of 1 N HCl. The resulting precipitate was collected by filtration. The solid was further dried to give 5-cyano-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (500 mg, 73% for 2 steps). ESI-MS *m/z* calc. 296.1, found 297.4 $(M+1)^+$; Retention time: 0.63 min (1 min run).

Step 3: 5-cyano-6-methyl-*N*-(3-nitrophenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide



5-Cyano-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (65 mg, 0.22 mmol), [(3-nitrophenyl)sulfonylamino]sodium (98 mg, 0.44 mmol), *N*-[(dimethylamino)-1*H*-1,2,3-triazolo-[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (170 mg, 0.44 mmol) and *N,N*-dimethylformamide (1 mL) were combined and the mixture was stirred at room temperature for 1 h. The reaction mixture was diluted with water (1 mL) and was extracted with ethyl acetate (2 x 5 mL). The organic layers were dried over sodium sulfate, filtered and concentrated. The residue was subjected to preparatory-HPLC (20-80% acetonitrile/water with 5 mM HCl) to give 5-cyano-6-methyl-*N*-(3-nitrophenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide. ESI-MS *m/z* calc. 480.1, found 481.5 $(M+1)^+$; Retention time: 0.73 min (1 min run).


Step 4: *N*-(3-aminophenyl)sulfonyl-5-cyano-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 124)



5-Cyano-6-methyl-*N*-(3-nitrophenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (from step 3) was dissolved in methanol (10 mL) before 10% Pd/C (12 mg, 0.011 mmol) was added. The mixture was stirred under a balloon of hydrogen at room temperature for 2 h. The mixture was filtered and concentrated. The residue was subjected to preparatory-HPLC (25-75% acetonitrile/water with 5 mM HCl) to give *N*-(3-aminophenyl)sulfonyl-5-cyano-6-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 124) (15 mg, 15% for 2 steps). ESI-MS *m/z* calc. 450.1, found 451.0 $(M+1)^+$; Retention time: 1.68 min (3 min run).


PREPARATION 12: *N*-(2-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 322)

Step 1: 6-*tert*-butyl-*N*-(2-nitrophenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide



6-*tert*-Butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (50. mg, 0.16 mmol), 2-nitrobenzenesulfonamide (32 mg, 0.16 mmol), *N*-[(dimethylamino)-1*H*-1,2,3-triazolo-[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (91 mg, 0.24 mmol), ethyldiisopropylamine (56 µL, 0.32 mmol) and NMP (1 mL) were combined. The mixture was heated at 75 °C for 16 h before it was cooled to room temperature. The reaction mixture was diluted with water (1 mL) and extracted with ethyl acetate (2 x 5 mL). The combined organic layers were dried over sodium sulfate, filtered and concentrated. The residue was subjected to preparatory-HPLC (30-90% acetonitrile/water with 5 mM HCl) to give 6-*tert*-butyl-*N*-(2-nitrophenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide. ESI-MS *m/z* calc. 497.2, found 498.4 $(M+1)^+$; Retention time: 0.89 min (1 min run).

Step 2: *N*-(2-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 322)




6-*tert*-Butyl-*N*-(2-nitrophenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (23 mg, 0.046 mmol) was dissolved in EtOH (5 mL) before 10% Pd/C (2 mg, 0.02

mmol) was added. The mixture was stirred under a balloon of hydrogen at room temperature for 2h. The mixture was filtered and the filtrate was concentrated. The residue was subjected to preparatory-HPLC (20-80% acetonitrile/water with 5 mM HCl) to give *N*-(2-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 322) (4.8 mg, 33%). ^{1}H NMR (400 MHz, DMSO) δ 8.06 (s, 1H), 7.74 (s, 1H), 7.36 (s, 1H), 7.20 (s, 1H), 6.99 (s, 2H), 6.87 (s, 1H), 6.72 (s, 1H), 2.34 (s, 3H), 2.04 (s, 6H), 1.33 (s, 2H), 1.13 (s, 9H). ESI-MS *m/z* calc. 467.2, found 468.0 $(M+1)^+$; Retention time: 2.25 min (3 min run).

The following compounds can be synthesized using the procedures described herein: *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2,6-dimethylphenoxy)pyridine-3-carboxamide (Compound 286), *N*-(4-aminophenyl)sulfonyl-6-*tert*-butyl-2-cyclohexyl-pyridine-3-carboxamide (Compound 537), *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 294) and *N*-(3-amino-1-methyl-pyrazol-4-yl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 41).

PREPARATION 13: 6-*tert*-butyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 51)



6-*tert*-butyl-*N*-[(2-methoxy-3-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 191) (45 mg, 0.093 mmol) was dissolved in dioxane (1 mL) before aq HCl (0.5 mL of 4.0 M, 2.0 mmol) was added. The reaction mixture was heated at 90 °C for 1 h before it was cooled and concentrated. The residue was subjected to preparatory-HPLC (20-80% acetonitrile/water with 5 μM HCl) to give 6-*tert*-butyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 51) (19 mg, 20.%). ^{1}H NMR (400 MHz, DMSO) δ 12.57 (s, 1H), 11.83 (s, 1H), 8.27 (dd, *J* = 7.3, 2.1 Hz, 1H), 8.07 (d, *J* = 7.8 Hz, 1H), 7.83 (d, *J* = 6.3 Hz, 1H), 7.16 (d, *J* = 7.9 Hz, 1H), 6.93 (s, 2H), 6.46 (t, *J* =

6.8 Hz, 1H), 2.27 (s, 3H), 1.99 (s, 6H), 1.05 (s, 9H). ESI-MS *m/z* calc. 469.2, found 470.3 $(M+1)^+$; Retention time: 1.95 min (3 min run).

The following compound can be synthesized using the procedures described herein: 6-*tert*-butyl-2-(2,4-dimethylphenoxy)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 55),

7,7-dimethyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)-6,8-dihydro-5*H*-quinoline-3-carboxamide (Compound 1199),

2-(2,4-dimethylphenoxy)-8,8-dimethyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 634),

8,8-dimethyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 899),

6-(1-methylcyclopropyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1226),

6-*tert*-butyl-*N*-[(4-methyl-2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1311),

6-(4-ethoxy-3-fluoro-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 901),

6-(4-methylcyclohexyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 743),


6-(4,4-dimethylcyclohexyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1091),

6-(4,4-difluorocyclohexyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1329), and

6-cyclohexyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1357).


PREPARATION 14: 6-Fluoropyridine-2-sulfonamide

Step 1: 2-(Benzylthio)-6-fluoropyridine



To a solution of BnSH (18.1 g, 146 mmol) in THF (1.12 L) was added 60% NaH (6.60 g, 165 mmol) portionwise over 10 minutes. The reaction mixture was stirred at room temperature for 15 minutes after which time a solution of 2,6-difluoropyridine (13.7 mL, 150. mmol) in THF (100 mL) was added dropwise over 10 minutes. The resulting clear solution was stirred at room temperature for 2 h. The reaction mixture was quenched with H_2O (250 mL) and the resultant mixture was extracted with Et_2O (800 mL). The organic layer was washed with H_2O (400 mL) and brine (250 mL), dried over Na_2SO_4 and concentrated to dryness. The residue was purified by silica gel column chromatography (100% hexanes followed by a gradient of 1-10% ethyl acetate in hexanes) to give 2-(benzylthio)-6-fluoropyridine (32.0 g, 100%). ^{1}H NMR ($CDCl_3$, 250 MHz) δ 7.56 (q, J = 8.0 Hz, 1H), 7.43-7.40 (m, 2H), 7.39-7.21 (m, 3H), 7.03 (dd, J = 7.5, 2.3 Hz, 1H), 6.60 (dd, J = 7.8, 2.5 Hz, 1H), 4.41 (s, 2H).

Step 2: 6-Fluoropyridine-2-sulfonyl chloride



2-(Benzylthio)-6-fluoropyridine (32.0 g, 146 mmol) was dissolved in $CHCl_3$ (500 mL) and H_2O (400 mL) was added. Chlorine was slowly bubbled through the reaction mixture for 2 h. (Note: a 30% aq. NaOH trap (1-1.5L) was used, followed by a sat. aq. $Na_2S_2O_3$ (1-1.5 L) trap to neutralize the excess Cl_2). The reaction mixture became warm during the addition and a yellow suspension formed. The reaction was carefully quenched (exothermic and foaming) with a solution of $Na_2S_2O_5$ (200 g) in H_2O (450 mL). The layers were separated and the aqueous layer was extracted once with $CHCl_3$ (200 mL). The organic layers were combined and washed with H_2O (300 mL), dried over Na_2SO_4 and concentrated to dryness to afford impure 6-fluoropyridine-2-sulfonyl chloride as a brownish oil. This material was used as such in the next step.


Step 3: 6-Fluoropyridine-2-sulfonamide



Crude 6-fluoropyridine-2-sulfonyl chloride (from step 2) was dissolved in CH_2Cl_2 (1 L) and the solution was cooled to -78°C. Ammonia was bubbled through the solution for 25 min. The mixture was stirred for 1 h at room temperature and the precipitated product was removed by filtration and washed with CH_2Cl_2 (2x). The filtrate and washings were combined and concentrated to ~50 mL and filtered; the solid was washed with CH_2Cl_2/hexanes (50/50 mL) and dried to afford 6-fluoropyridine-2-sulfonamide (12.0 g, 46% yield, purity ~90%) as a grey solid. ^{1}H NMR (DMSO, 250 MHz) δ 8.27 (q, *J*=8.0 Hz, 1H), 7.88 (dd, *J*=7.5, 2.1 Hz, 1H), 7.66 (s, 2H), 7.48 (dd, *J*=8.0, 2.4 Hz, 1H).

PREPARATION 15: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-ethoxy-5-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1255)


Step 1: 6-chloro-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid



To a solution of 2,4,6-trimethylphenol (13.6 g, 100 mmol) in *N,N*-dimethylformamide (300 mL) at 0-5 °C, was carefully added NaH solid (60% in mineral oil, 8.80 g, 220 mmol). After the resultant suspension was stirred at the same temperature for 15 min, solid 2,6-dichloro-nicotinic acid (19.2 g, 100 mmol) was added. The reaction mixture was stirred at room temperature for 20 min and then at 80 °C overnight. The mixture was cooled to room temperature before it was poured into a cold $NaHSO_4$ solution (24.0 g in 200 mL water). The resultant suspension was stirred at room temperature for 30 min. The precipitate was collected by filtration, washed with water, followed by Et_2O/hexanes (200/500 mL), and dried at 40 °C under vacuum to give 6-chloro-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid as a brownish solid (19.2 g, 66%). ^{1}H NMR (400 MHz, DMSO) δ 13.38 (s, 1H), 8.28 (d, *J* = 7.9 Hz,


1H), 7.27 (d, J = 7.9 Hz, 1H), 6.94 (d, J = 1.1 Hz, 2H), 2.27 (s, 3H), 1.99 (s, 6H). ESI-MS m/z calc. 291.1, found 292.0 $(M+1)^+$; Retention time: 2.34 min (3 min run).

Step 2: 6-chloro-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide




6-Chloro-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (23.7 g, 81.4 mmol) was added to a mixture of *N*-[(dimethylamino)-1*H*-1,2,3-triazolo-[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (35.0 g, 92.1 mmol), 6-fluoropyridine-2-sulfonamide (16.5 g, 93.8 mmol) and ethyldiisopropylamine (36 mL) in *N*,*N*-dimethylformamide (400 mL) at 5 °C. The resulted mixture was stirred at room temperature overnight before it was quenched with 1 N HCl (150 mL). The mixture was diluted with water (450 mL) and the resultant solids were collected by filtration and washed with water (500 mL), methanol (200 mL) and Et_2O (400 mL). The solid was purified by silica gel column chromatography (0-4% methanol/CH_2Cl_2) to afford 6-chloro-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide as a white solid (22.2 g, 60%). ^{1}H NMR ($CDCl_3$, 250 MHz) δ 8.36 (dd, J=15.7, 7.5 Hz, 1H), 8.14 (d, J=7.5 Hz, 1H), 8.08 (d, J=8.0 Hz, 1H), 7.58 (d, J=8.2 Hz, 1H), 7.29 (d, J=7.8 Hz, 1H), 6.91 (s, 2H), 2.25 (s, 3H), 1.94 (s, 6H).

Step 3: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide



6-Chloro-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (12.0 g, 26.7 mmol) was mixed with NH_4OH (28-30%, 250 mL) and DMSO (200 mL) in a 1L pressure vessel at 0 °C. The container was sealed and heated at 92 °C (oil bath) for 2 d, and at 85 °C for another 2 d. Solvents (water and DMSO) were removed and the residue was suspended in a mixture solvent of CH_2Cl_2/ethyl acetate/methanol (25/25/25 mL). The suspension was sonicated at room temperature for 15 min and was filtered. The solids were washed with CH_2Cl_2/ethyl acetate/methanol (10/10/10 mL) and dried to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide as a white solid (6.8g, 57%). ^{1}H NMR ($CDCl_3$, 250 MHz) δ 8.10 (d, *J*=7.8 Hz, 1H), 7.64 (t, *J*=7.8 Hz, 1H), 7.27 (d, *J*=8.0 Hz, 1H), 7.21 (d, *J*=7.2 Hz, 1H), 6.92 (s, 2H), 6.71 (d, *J*=8.5 Hz, 1H), 6.60 (br, s, 2H), 2.26 (s, 3H), 1.98 (s, 6H).

Step 4: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-ethoxy-5-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1255)



A vial containing a mixture of *N*-((6-aminopyridin-2-yl)sulfonyl)-6-chloro-2-(mesityloxy)-nicotinamide (30. mg, 0.067 mmol), 3-ethoxy-5-fluoro-benzeneboronic acid (25 mg, 0.13 mmol), $Pd(PPh_3)_4$ (5.0 mg), K_3PO_4 (2.0 M, 0.13 mL) and *N*,*N*-dimethylformamide (0.6 mL) was flushed with nitrogen and sealed. The mixture was then heated at 80 °C overnight before it was cooled and diluted with DMSO (0.3 mL). The mixture was subjected to preparatory-HPLC (30-90% acetonitrile/water with 5 μM HCl) to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-ethoxy-5-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1255) (20 mg, 54%). ESI-MS *m/z* calc. 550.2, found 551.5 $(M+1)^+$; Retention time: 3.15 min (5 min run).

The following compounds can be synthesized using the procedures described herein: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(2-*tert*-butylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-

carboxamide (Compound 364),

N-[(6-amino-2-pyridyl)sulfonyl]-6-cyclopropyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1050),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,4-dimethylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 399),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 639),

N-[(6-amino-2-pyridyl)sulfonyl]-4-(3-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)benzamide (Compound 1549),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)-6-vinyl-pyridine-3-carboxamide (Compound 430),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-hydroxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 505),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-methyl-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 513),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-4-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 514),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-methylindazol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 451),

N-[[6-(aminomethyl)-2-pyridyl]sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 871),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[2-methyl-5-(trifluoromethyl)pyrazol-3-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 457),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 463),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-fluoro-5-isobutoxy-phenyl)-3-(2,4,6-trimethylphenoxy)pyridine-2-carboxamide (Compound 1456),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-propylpyrazol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 469),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4,4-dimethylcyclohexyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 538),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4,4-difluorocyclohexyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound783),

N-[(6-amino-2-pyridyl)sulfonyl]-6-ethyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 963),

N-[(6-amino-2-pyridyl)sulfonyl]-6-isopropyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1232),

N-[(6-amino-2-pyridyl)sulfonyl]-6-cyclopentyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1280),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-methoxypropyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 791),

N-[(6-amino-2-pyridyl)sulfonyl]-6-pentyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 862),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-ethoxyethyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 929),

N-[(6-amino-2-pyridyl)sulfonyl]-6-cyclohexyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1131),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[2-(1-piperidyl)pyrimidin-5-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 475),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-methoxy-4-methyl-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 375),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-fluoro-2-methoxy-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 387),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,4-difluorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-

carboxamide (Compound 439),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-hydroxy-3-methyl-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 450),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(morpholinomethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 465),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-methoxy-3-methyl-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 477),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-2-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 396),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 436),

6-(2-fluoro-4-methoxy-phenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 374),

6-[2-methyl-5-(trifluoromethyl)pyrazol-3-yl]-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 391),

6-(4-fluoro-2-methoxy-phenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 405),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-6-(1-isobutylpyrazol-4-yl)pyridine-3-carboxamide (Compound 381),

6-(4-isopropoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 494),

6-(1,3-benzodioxol-4-yl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 525),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-fluorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 93),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-methylindol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 386),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(cyanomethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 388),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-methoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 413),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-methoxypyrimidin-5-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 449),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-methoxy-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 531),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-fluoro-4-(isopropylcarbamoyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 543),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-ethoxy-2-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 599),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,6-dimethoxy-4-methyl-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 629),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(cyanomethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 638),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-3-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 661),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(m-tolyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 672),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 673),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 682),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(1-piperidylmethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 686),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[2-(2,2,2-trifluoroethoxy)-3-pyridyl]-2-(2,4,6-

trimethylphenoxy)pyridine-3-carboxamide (Compound 697),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-fluoro-2-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 700),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,4-difluorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 725),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[2-fluoro-5-(hydroxymethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 727),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tetrahydropyran-4-yl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 748),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-hydroxy-3-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 768),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(dimethylamino)-3-pyridyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 787),

6-[3-(aminomethyl)phenyl]-*N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 801),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,5-difluorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 803),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(morpholinomethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 810),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,5-dimethoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 828),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-fluoro-2-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 830),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-methyl-2-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1272),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-methyl-2-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 730),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-methyl-2-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1290),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-ethoxy-2-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 602),

6-(6-methoxy-5-methyl-3-pyridyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1044),

2-(2,4-dimethylphenoxy)-6-(p-tolyl)-*N*-(1H-pyrazol-3-ylsulfonyl)pyridine-3-carboxamide (Compound 610),

2-(2,4-dimethylphenoxy)-6-(6-methoxy-5-methyl-3-pyridyl)-*N*-(1H-pyrazol-3-ylsulfonyl)pyridine-3-carboxamide (Compound 892),

2-(2,4-dimethylphenoxy)-6-(3-fluoro-5-isopropoxy-phenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)pyridine-3-carboxamide (Compound 1213),

2-(2,4-dimethylphenoxy)-6-(4-ethoxyphenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)pyridine-3-carboxamide (Compound 813),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-propoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 535),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-2-hydroxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 866),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-6-(4-ethoxyphenyl)pyridine-3-carboxamide (Compound 641),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-6-(3-fluoro-5-isopropoxy-phenyl)pyridine-3-carboxamide (Compound 1150),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-6-(p-tolyl)pyridine-3-carboxamide (Compound 1015),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-ethylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1118),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isopropylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-

carboxamide (Compound 1016),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(trifluoromethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 709), (Compound 709)

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-ethylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 900),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(trifluoromethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 548),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-6-(6-methoxy-5-methyl-3-pyridyl)pyridine-3-carboxamide (Compound 533),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-chloro-2-methoxy-4-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1284),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-methoxy-2-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1034),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-ethoxy-2-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1133),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isopropoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1138),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-fluoro-2-methoxy-4-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1277),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-ethoxy-4-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 860),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-isopropylpyrazol-3-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1317),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-hydroxy-4-methyl-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 793),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-propoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 996),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-ethoxy-3-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1268),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-chlorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1221),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,4-dihydro-2*H*-pyran-6-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 937),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-4-methyl-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1000),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(cyclopenten-1-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1330),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-2-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 897), (Compound 897)

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-4-methyl-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 875),

N-[(6-amino-2-pyridyl)sulfonyl]-6-isopropenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1211),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-chloro-2-methoxy-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 993),

6-(1-isobutylpyrazol-4-yl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 577),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1278),

N-(1H-pyrazol-3-ylsulfonyl)-6-(2-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1353),

6-(3-ethoxy-5-fluoro-phenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 585),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2,4,6-

trimethylphenoxy)pyridine-3-carboxamide (Compound 1163),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-methoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1173),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-chlorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1295),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 570),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-fluoro-2-hydroxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 622),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,6-dihydro-2*H*-pyran-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 976),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,5-difluorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1325),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,4-dimethoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 950),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[1-methyl-3-(trifluoromethyl)pyrazol-4-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 812),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-chloro-6-methoxy-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1332),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,6-dimethoxy-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 754),

N-[(6-amino-2-pyridyl)sulfonyl}-6-(2-isopropoxy-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 909),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(trifluoromethyl)-1*H*-pyrazol-4-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 582),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(dimethylaminomethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 962),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-cyano-2-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1111),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-fluoro-6-methoxy-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 612),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-ethoxypyrimidin-5-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1130),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-methoxy-5-methyl-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 705),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-methoxy-2-methyl-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1212),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-ethoxy-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 718),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1,3-benzodioxol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 974),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1,3-benzodioxol-5-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1207),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-benzyloxy-2-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1113),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-fluoro-2-hydroxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1409),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-isopropylpyrazol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 808),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-methoxy-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1115),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-ethylpyrazol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1312),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1,5-dimethylpyrazol-4-yl)-2-(2,4,6-

trimethylphenoxy)pyridine-3-carboxamide (Compound 584),

6-phenyl-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 784),

6-(4-methoxyphenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1033),

N-[(2-methoxy-3-pyridyl)sulfonyl]-6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 939),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1H-indazol-5-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 856),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-cyanophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 886),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-fluoro-3-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 919),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-6-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 920),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-cyano-3-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 923),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,5-dimethylpyrazol-3-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 938),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-4-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 956),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1H-indol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 977),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,5-difluoro-4-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 984),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(p-tolyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1035),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,3-dihydrobenzofuran-5-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1084),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-methylindol-6-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1056),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-methylpyrazol-3-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1057),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(o-tolyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1120),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[2-(cyclopropylamino)pyrimidin-5-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1126),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[2-(dimethylamino)pyrimidin-5-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1139),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-4-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1140),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,4-dimethoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1146),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(1-hydroxy-1-methyl-ethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1147),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-isobutylpyrazol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1177),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-5-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1178),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-ethoxy-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1185),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-hydroxy-2-methyl-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1204),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,5-dimethoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-

3-carboxamide (Compound 1225),

6-[4-(aminomethyl)phenyl]-*N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1228),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,3-difluorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1244),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-hydroxy-4-methoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1251),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-ethoxy-5-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1255),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-pyrrolidin-1-ylpyrimidin-5-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1275),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-methylindazol-5-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1276),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-methylindazol-6-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1299),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-methoxy-5-methyl-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1320),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,6-dimethoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1326),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-isopentylpyrazol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1336),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-ethoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1337),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-hydroxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1344),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(hydroxymethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 786),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-hydroxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 908),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(hydroxymethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1183),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,3-dimethoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 485),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,4-dimethoxypyrimidin-5-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 499),

N-[(6-amino-2-pyridyl)sulfonyl]-6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 203),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-methoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 17),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-ethoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 61),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,6-dimethylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 97),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-ethoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 250),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-furyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 284),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(trifluoromethyl)-3-pyridyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 132),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-cyanophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 216),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,4-dihydro-2*H*-1,5-benzodioxepin-7-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 14),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(methoxymethyl)phenyl]-2-(2,4,6-

trimethylphenoxy)pyridine-3-carboxamide (Compound 66),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylphenoxy)-6-phenyl-pyridine-3-carboxamide (Compound 340),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(methoxymethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 69),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-isopropylpyrimidin-5-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 64),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-methyl-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 513),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(hydroxymethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 786),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-hydroxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 908),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(hydroxymethyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1183),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-ethoxy-2-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 599),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(dimethylcarbamoyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-fluoro-5-(methylcarbamoyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-methylindazol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 451),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,5-dimethoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 828),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,5-dimethoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1225),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-isopentylpyrazol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1336),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-propylpyrazol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 469),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-hydroxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 505),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-isobutylpyrazol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1177),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,4-difluorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 725),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(2-cyanoethylcarbamoyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-fluoro-4-(isopropylcarbamoyl)phenyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 543),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(1-methyl-2-tetrahydropyran-4-yl-ethoxy)pyridine-3-carboxamide (Compound 1361),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(1,1-dimethyl-2-phenyl-ethoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1365),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(1-cyclopentylethoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1374),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(cyclohexylmethoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1395),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methylcyclobutoxy)pyridine-3-carboxamide (Compound 406),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-cyclobutylethoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1418),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-phenethyloxy-pyridine-3-

carboxamide (Compound 1426),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[1-(3-pyridyl)ethoxy]pyridine-3-carboxamide (Compound 1434),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-cyclopentylethoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1440),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(cyclopropoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1446),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(cyclopentylmethoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1450),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(cyclopentoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1451),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,3-difluorocyclobutoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1458),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-tetrahydrofuran-3-yl]oxy-pyridine-3-carboxamide (Compound 1464),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-cyclohexylethoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1465),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(cyclohexoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1477),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(1-benzylcyclobutoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1479),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(tetrahydrofuran-3-ylmethoxy)pyridine-3-carboxamide (Compound 1499),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-cyclopropylcyclobutoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1530),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-benzyloxycyclobutoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1559),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(tetrahydropyran-2-ylmethoxy)pyridine-3-carboxamide (Compound 1570),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2,4-dimethyl-3-pyridyl)oxy]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1574),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-tetrahydropyran-4-ylethoxy)pyridine-3-carboxamide (Compound 1579),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(cyclobutoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1580),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-tetrahydrofuran-3-yl]oxy-pyridine-3-carboxamide (Compound 1592),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-tetrahydrofuran-2-ylethoxy)pyridine-3-carboxamide (Compound 1604),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[2-(4-pyridyl)ethoxy]pyridine-3-carboxamide (compound 1605),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methoxycyclohexoxy)pyridine-3-carboxamide (Compound 1622),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[2-(2-pyridyl)ethoxy]pyridine-3-carboxamide (Compound 1625),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethylcyclopentoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1629),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[1-(2,2-dimethylcyclopropyl)propoxy]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1639),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(1-methylcyclobutoxy)pyridine-3-carboxamide (Compound 1644),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[2-(3-pyridyl)ethoxy]pyridine-3-carboxamide (Compound 1654),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(tetrahydrofuran-2-

ylmethoxy)pyridine-3-carboxamide (Compound 1661),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(1-methylcyclopropyl)methoxy]pyridine-3-carboxamide (Compound 1668),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-tetrahydropyran-2-ylethoxy)pyridine-3-carboxamide (Compound 1680),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-cyclopropylethoxy)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1685),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 8),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,6-dimethylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 97),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-ethoxy-3-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 161),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 164),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 196),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-2-fluoro-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 207),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chlorophenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 234),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 238),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isopropoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 281),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isobutoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 320),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(7-fluoro-1,3-benzodioxol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 345),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 351),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-methylcyclohexyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 172),


6-(p-tolyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 300),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 318), and

6-(4-ethoxyphenyl)-*N*-(1H-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 190).

PREPARATION 16: *N*-(3-aminophenyl)sulfonyl-5-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1321)


Step 1: methyl 5-bromo-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate



2,4,6-Trimethylphenol (2.68 g, 19.7 mmol) and 60% NaH (870 mg, 22 mmol) were combined in an anhydrous mixture of *N,N*-dimethylformamide (24 mL) and THF (2.4 mL). The reaction mixture was stirred at room temperature for 15 minutes under an atmosphere of nitrogen. Methyl 5-bromo-2-chloro-pyridine-3-carboxylate (4.93 g, 19.7 mmol) was added and the reaction mixture was heated at 60 °C for 1 h. The reaction mixture was then allowed to cool to room temperature. The reaction mixture was diluted with water and was extracted with ethyl acetate. The ethyl acetate layer was washed three times with water and then was concentrated under reduced pressure. The residue was subjected to silica gel column chromatography (0-50%


ethyl acetate/hexanes) to yield methyl 5-bromo-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (5.50 g, 73%) as a faint yellow solid. ESI-MS *m/z* calc. 349.0, found 350.3 $(M+1)^+$; Retention time: 2.09 min (3 min run).

Step 2: methyl 5-bromo-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate



Methyl 5-bromo-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (2.60 g, 6.76 mmol) was combined in a mixture of methanol (12.95 mL) and water (17.27 mL) containing sodium hydroxide (810 mg, 20.3 mmol). The resulting suspension was heated at 60 °C for 3 h. The reaction mixture was cooled to room temperature and was made acidic with 6 M hydrochloric acid. The resulting precipitate was filtered, washed with water and dried in a vacuum oven at 60 °C overnight to yield 5-bromo-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (2.24 g, 98%) as a white solid. ESI-MS *m/z* calc. 335.0, found 338.2 $(M+1)^+$; Retention time: 1.75 min (3 min run).


Step 3: methyl 5-bromo-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate



5-Bromo-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (89 mg, 0.26 mmol), phenylboronic acid (39 mg, 0.32 mmol), $Pd(PPh_3)_4$ (31 mg, 0.026 mmol), and K_2CO_3 (530 µL of 2.0 M, 1.1 mmol) were combined in DME (2.5 mL). The reaction mixture was briefly sparged with nitrogen, capped, and heated at 80 °C for 2 h. The reaction mixture was allowed to cool to room temperature and the layers were separated. The organic layer was subjected to


preparatory-HPLC (1-99% acetonitrile/water with 5 μM HCl) to give 5-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (54 mg, 61%) as a white solid. ESI-MS *m/z* calc. 333.1, found 334.4 $(M+1)^+$; Retention time: 0.72 min (1 min run).

Step 4: *N*-(3-nitrophenyl)sulfonyl-5-phenyl-2-(2,4,6-trimethylphenoxy)-pyridine-3-carboxamide



5-Phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (54 mg, 0.16 mmol), (3-nitrophenyl)sulfonylazanide sodium (73 mg, 0.32 mmol), *N*-[(dimethylamino)-1*H*-1,2,3-triazolo-[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (62 mg, 0.16 mmol) and *N,N*-dimethylformamide (1 mL) were combined. The reaction mixture was stirred at room temperature for 72 h. The crude mixture was subjected to preparatory-HPLC (1-99% acetonitrile/water with 5 μM HCl) to give *N*-(3-nitrophenyl)sulfonyl-5-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (51 mg, 61%) as a white solid. ESI-MS *m/z* calc. 517.1, found 518.5 $(M+1)^+$; Retention time: 2.04 min (3 min run).

Step 5: *N*-(3-aminophenyl)sulfonyl-5-phenyl-2-(2,4,6-trimethylphenoxy)-pyridine-3-carboxamide



N-(3-nitrophenyl)sulfonyl-5-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (51 mg, 0.099 mmol) was suspended in acetic acid (2 mL) containing zinc (10 mg, 0.15 mmol). The reaction mixture was heated at 65 °C for 30 min. The hot reaction mixture was

filtered and then allowed to cool to room temperature. The mixture was concentrated and the residue was subjected to preparatory-HPLC (1-99% acetonitrile/water with 5 mM HCl) to give *N*-(3-aminophenyl)sulfonyl-5-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1321) (18 mg, 38%) as a white solid. ESI-MS *m/z* calc. 487.2, found 488.5 $(M+1)^+$; Retention time: 2.74 min (3 min run).

The following compound can be synthesized using the procedures described herein: *N*-[(6-amino-2-pyridyl)sulfonyl]-5-(2-propylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 429),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(2-isopropylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 865),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(2-*tert*-butylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 868),


N-(3-aminophenyl)sulfonyl-6-*tert*-butyl-4-methoxy-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 959),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(o-tolyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 817), and

N-[(6-amino-2-pyridyl)sulfonyl]-5-(2-ethylphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 945).

PREPARATION 17: *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(*N*,2-dimethylanilino)pyridine-3-carboxamide (Compound 193)


Step 1: 6-*tert*-butyl-2-(*N*,2-dimethylanilino)pyridine-3-carbonitrile



A mixture of 6-*tert*-butyl-2-chloro-pyridine-3-carbonitrile (100. mg, 0.514 mmol), *N*,2-dimethylaniline (76 µL, 0.62 mmol), KHMDS (1.2 mL of 0.5 M, 0.62 mmol) and dioxane (1.5 mL) was heated at 100 °C for 20 min. The mixture was partitioned between ethyl acetate


and 1N HCl. The layers were separated and the aqueous layer was extracted with ethyl acetate (3x). The combined organics were washed with brine, dried over magnesium sulfate, filtered and concentrated. The residue was subjected to silica gel column chromatography (0-25% ethyl acetate/hexanes) to give 6-*tert*-butyl-2-(N,2-dimethylanilino)pyridine-3-carbonitrile (40 mg, 26%). ESI-MS *m/z* calc. 279.2, found 280.4 $(M+1)^+$; Retention time: 2.28 min (3 min run).

Step 2: 6-*tert*-butyl-2-(N,2-dimethylanilino)pyridine-3-carboxamide



A mixture of 6-*tert*-butyl-2-(*N*,2-dimethylanilino)pyridine-3-carbonitrile (40 mg, 0.14 mmol) and KOH (340 µL of 4.0 M, 1.4 mmol) in EtOH (1.2 mL) was heated at 90 °C overnight. The mixture was concentrated, acidified with 2N HCl and the solids were collected to give an off-white solid. The solids were placed in a vacuum oven at 55 °C overnight to give 6-*tert*-butyl-2-(N,2-dimethylanilino)pyridine-3-carboxamide (40 mg, 30%) as an off-white solid. ESI-MS *m/z* calc. 297.2, found 298.4 $(M+1)^+$; Retention time: 0.68 min (1 min run).


Step 3: 6-*tert*-butyl-2-(*N*,2-dimethylanilino)-*N*-(3-nitrophenyl)sulfonyl-pyridine-3-carboxamide



Sodium hydride (16 mg, 60%, 0.40 mmol) was added to a mixture of 6-*tert*-butyl-2-(*N*,2-dimethylanilino)pyridine-3-carboxamide (40. mg, 0.13 mmol) and *N*,*N*-dimethylformamide (1 mL). The mixture was allowed to stir for 15 min before 3-nitrobenzenesulfonyl chloride (45 mg, 0.20 mmol) was added. The mixture was allowed to stir for 1H before adding two drops of methsanol. The mixture was filtered and subjected to preparatory-HPLC (10-99%

acetonitrile/water with 0.05 mM HCl) to give 6-*tert*-butyl-2-(*N*,2-dimethylanilino)-*N*-(3-nitrophenyl)sulfonyl-pyridine-3-carboxamide. ESI-MS *m/z* calc. 482.2, found 483.3 $(M+1)^+$; Retention time: 0.82 min (1 min run).

Step 4: *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(*N*,2-dimethylanilino)pyridine-3-carboxamide (Compound 193)




Zinc (9.0 mg, 0.13 mmol) was added to a mixture of 6-*tert*-butyl-2-(*N*,2-dimethylanilino)-*N*-(3-nitrophenyl)sulfonyl-pyridine-3-carboxamide and acetic acid (1 mL) at room temperature. The mixture was allowed to stir for 20 min before it was filtered and subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mMHCl) to give *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(*N*,2-dimethylanilino)pyridine-3-carboxamide (Compound 193) (1.2 mg, 2% for 2 steps). ESI-MS *m/z* calc. 452.2, found 453.3 $(M+1)^+$; Retention time: 1.90 minutes (3 min run).

The following compound can be synthesized using the procedures described herein: *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-2-(*N*-methylanilino)pyridine-3-carboxamide (Compound 85).


PREPARATION 18: *N*-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 110)

Step 1: ethyl 5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylate




To a mixture of ethyl 5-*tert*-butyl-4*H*-pyrazole-3-carboxylate (500. mg, 2.55 mmol) and *N*,*N*-dimethylformamide (5 mL) was added 2-(bromomethyl)-1,3,5-trimethyl-benzene (543 mg, 2.55 mmol) and K_2CO_3 (704 mg, 5.10 mmol). The reaction mixture was heated at 80 °C overnight. Additional 2-(bromomethyl)-1,3,5-trimethyl-benzene (200 mg, 1.02 mmol) and K_2CO_3 (704 mg, 5.10 mmol) were added and the mixture was heated at 80 °C for another 18 h. The mixture was cooled to room temperature before water (20 mL) was added. The mixture was extracted with ethyl acetate (2 x 50 mL). The combined organic layers were dried over sodium sulfate, filtered and concentrated. The residue was subjected to silica gel column chromatography (0-50% ethyl acetate/hexanes) to give ethyl 5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylate (642 mg, 77%). ESI-MS *m/z* calc. 328.2, found 329.4 $(M+1)^+$; Retention time: 0.99 min (1 min run).

Step 2: 5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylic acid




Ethyl 5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylate (5.00 g, 15.2 mmol) and NaOH (4.87 g, 122 mmol) were combined in a THF (20 mL) and water (20 mL). The mixture was heated at 60 °C for 3 h. The reaction mixture was made acidic by the addition of 6 M HCl and was extracted with ethyl acetate. The organics were washed with brine, dried over sodium sulfate and evaporated to give 5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)-methyl]pyrazole-3-carboxylic acid (4.38 g, 96%). ^{1}H NMR (400 MHz, DMSO) δ 13.23 (s, 1H), 6.83 (s, 2H), 6.68 (s, 1H), 5.64 (s, 2H), 2.24 (s, 6H), 2.21 (s, 3H), 1.15 (s, 9H). ESI-MS *m/z* calc. 300.2, found 301.4 $(M+1)^+$; Retention time: 2.21 min (3 min run).

Step 3: 5-*tert*-butyl-*N*-(3-nitrophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl) methyl]pyrazole-3-carboxamide



5-*tert*-Butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylic acid (108 mg, 0.356 mmol), 3-nitrobenzenesulfonamide (109 mg, 0.539 mmol), *N*-[(dimethylamino)-1*H*-1,2,3-triazolo-[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (137 mg, 0.356 mmol), and triethylamine (100 µL, 0.719 mmol) were combined in *N,N*-dimethylformamide (2.3 mL) and heated at 60 °C for 16 h. The reaction mixture was filtered and subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mMHCl) to give 5-*tert*-butyl-*N*-(3-nitrophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)-methyl]pyrazole-3-carboxamide (105 mg, 60.9%). ESI-MS *m/z* calc. 484.2, found 485.3 $(M+1)^+$; Retention time: 2.27 min (3 min run).

4: *N*-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]-pyrazole-3-carboxamide



5-*tert*-Butyl-*N*-(3-nitrophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]-pyrazole-3-carboxamide (105 mg, 0.217 mmol) was dissolved in methanol (9 mL) with Pd/C (38 mg, 0.036 mmol) under a balloon of hydrogen and the mixture was stirred for 30 min. The reaction mixture was filtered, evaporated, and the residue was purified by LC/MS utilizing a gradient of 10-99% acetonitrile in 5 mM aqueous HCl to yield *N*-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 110) (72 mg, 44%). ^{1}H NMR (400 MHz, DMSO) δ 7.26 (t, *J* = 7.9 Hz, 1H), 7.21 (t, *J* = 2.0 Hz, 1H), 7.12 - 7.06 (m, 1H), 7.01 (s, 1H), 6.89 - 6.83 (m, 1H), 6.77 (s, 2H), 5.45 (s, 2H), 2.17 (s, 3H), 2.11 (s, 6H), 1.14 (s, 9H). ESI-

MS *m/z* calc. 454.2, found 455.4 $(M+1)^+$; Retention time: 2.06 min (3 min run).

The following compounds can be synthesized using the procedures described herein:

3-(*tert*-butyl)-1-(1-mesitylethyl)-*N*-(phenylsulfonyl)-1*H*-pyrazole-5-carboxamide (Compound 201),

N-(3-aminophenyl)sulfonyl-3-tert-butyl-1-[(2,4,6-trimethylphenyl)methyl]pyrazole-4-carboxamide,

3-(*tert*-butyl)-1-(2-fluoro-5-(trifluoromethyl)benzyl)-*N*-(phenylsulfonyl)-1*H*-pyrazole-5-carboxamide (Compound 35),

3-(*tert*-butyl)-*N*-((3-(dimethylamino)phenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 270),

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(2,4,6-trichlorobenzyl)-1*H*-pyrazole-5-carboxamide (Compound 303),

3-(*tert*-butyl)-1-(2,6-dimethylbenzyl)-*N*-((1-methyl-1*H*-indol-4-yl)sulfonyl)-1*H*-pyrazole-5-carboxamide (Compound 210),

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(4-(*tert*-butyl)-2,6-dimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 156),

3-(4-(difluoromethyl)phenyl)-1-(2,6-dimethylbenzyl)-*N*-(phenylsulfonyl)-1*H*-pyrazole-5-carboxamide (Compound 119),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-4-methoxy-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 1305),

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(2-chloro-6-(trifluoromethyl)benzyl)-1*H*-pyrazole-5-carboxamide (Compound 189),

3-(*tert*-butyl)-*N*-((3-(dimethylamino)-2-methylphenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 104),

N-((3-amino-2-methylphenyl)sulfonyl)-3-(*tert*-butyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 123),

3-(*tert*-butyl)-*N*-((1-methyl-1*H*-indol-7-yl)sulfonyl)-1-(2-methyl-3-(trifluoromethyl)benzyl)-1*H*-

pyrazole-5-carboxamide (Compound 276),

3-(*tert*-butyl)-*N*-(m-tolylsulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 80),

N-((3-amino-4-fluorophenyl)sulfonyl)-3-(*tert*-butyl)-1-(2,6-dimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 150),

3-(*tert*-butyl)-*N*-(quinolin-5-ylsulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 352),

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(2,3,5,6-tetramethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 317),

3-(*tert*-butyl)-1-(2,6-dimethylbenzyl)-*N*-(thiophen-2-ylsulfonyl)-1*H*-pyrazole-5-carboxamide (Compound 77),

3-(*tert*-butyl)-*N*-(phenylsulfonyl)-1-(2,4,6-trichlorobenzyl)-1*H*-pyrazole-5-carboxamide (Compound 145),

3-(*tert*-butyl)-1-(2,6-dimethylbenzyl)-*N*-((2-methoxy-3-methylphenyl)sulfonyl)-1*H*-pyrazole-5-carboxamide (Compound 59),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[2-(2,4,6-trimethylphenyl)ethyl]pyrazole-3-carboxamide (Compound 7),

3-(*tert*-butyl)-*N*-(phenylsulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide,

3-(*tert*-butyl)-1-(naphthalen-1-ylmethyl)-*N*-(phenylsulfonyl)-1*H*-pyrazole-5-carboxamide (Compound 328),

3-(*tert*-butyl)-*N*-((3-(methylamino)phenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 249),

3-(*tert*-butyl)-*N*-((1-(2-hydroxyethyl)-1*H*-indol-4-yl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 188),

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-4-methyl-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 20),

3-(*tert*-butyl)-1-(2,6-dimethylbenzyl)-*N*-((2-fluorophenyl)sulfonyl)-1*H*-pyrazole-5-carboxamide

(Compound 57),

3-(*tert*-butyl)-*N*-((2-cyanophenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 266),

3-(*tert*-butyl)-*N*-((3-hydroxyphenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 84),

3-(*tert*-butyl)-1-(2,6-dimethylbenzyl)-*N*-(phenylsulfonyl)-1*H*-pyrazole-5-carboxamide (Compound 257),

N-((3-amino-4-methylphenyl)sulfonyl)-3-(*tert*-butyl)-1-(2,6-dimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 312),

1-(3-(1*H*-pyrrol-1-yl)benzyl)-3-(*tert*-butyl)-*N*-(phenylsulfonyl)-1*H*-pyrazole-5-carboxamide (Compound 108),

3-(*tert*-butyl)-1-(5-fluoro-2-(trifluoromethyl)benzyl)-*N*-(phenylsulfonyl)-1*H*-pyrazole-5-carboxamide (Compound 3),

N-((3-acetylphenyl)sulfonyl)-3-(*tert*-butyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 332),

N-((1*H*-indol-4-yl)sulfonyl)-3-(*tert*-butyl)-1-(2-methyl-3-(trifluoromethyl)benzyl)-1*H*-pyrazole-5-carboxamide (Compound 60),

3-(*tert*-butyl)-*N*-((2,6-difluorophenyl)sulfonyl)-1-(2,6-dimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 170),

N-((4-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(2,6-dimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 81),

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(2-fluoro-6-(trifluoromethyl)benzyl)-1*H*-pyrazole-5-carboxamide (Compound 239),

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(1-(o-tolyl)ethyl)-1*H*-pyrazole-5-carboxamide (Compound 342),

N-((6-aminopyridin-2-yl)sulfonyl)-3-(*tert*-butyl)-4-methyl-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 356),

N-((2-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(2,6-dimethylbenzyl)-1*H*-pyrazole-5-carboxamide,

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(1-cyclohexylethyl)-1*H*-pyrazole-5-carboxamide (Compound 287),

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(1-phenylpropyl)-1*H*-pyrazole-5-carboxamide (Compound 165),

3-(*tert*-butyl)-*N*-((4-hydroxyphenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 269),

N-((3-aminophenyl)sulfonyl)-3-(1-methylcyclopropyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 158),

3-(*tert*-butyl)-1-(2-methyl-3-(trifluoromethyl)benzyl)-*N*-(phenylsulfonyl)-1*H*-pyrazole-5-carboxamide (Compound 98),

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(2,6-dimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 185),

N-((3-aminophenyl)sulfonyl)-3-isobutyl-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 167),

3-(*tert*-butyl)-*N*-((2,3-dihydrobenzofuran-7-yl)sulfonyl)-1-(2,6-dimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 211),

N-((6-aminopyridin-2-yl)sulfonyl)-3-(*tert*-butyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 173),

3-(*tert*-butyl)-1-(2-methoxyphenethyl)-*N*-(phenylsulfonyl)-1*H*-pyrazole-5-carboxamide (Compound 157),

N-((1*H*-indazol-4-yl)sulfonyl)-3-(*tert*-butyl)-1-(2,6-dimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 86),

N-(benzenesulfonyl)-5-*tert*-butyl-2-[(1-methylindol-4-yl)methyl]pyrazole-3-carboxamide (Compound 245),

3-(*tert*-butyl)-*N*-((3-methoxypyridin-2-yl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (Compound 44),

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(naphthalen-1-ylmethyl)-1*H*-pyrazole-5-carboxamide (Compound 295),

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-((1-methyl-1*H*-indol-7-yl)methyl)-1*H*-pyrazole-5-carboxamide,

3-(*tert*-butyl)-1-(2,6-dimethylbenzyl)-*N*-((2-(methylamino)pyridin-4-yl)sulfonyl)-1*H*-pyrazole-5-carboxamide,

5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]-*N*-(2-methoxyphenyl)sulfonyl-pyrazole-3-carboxamide (Compound 1551),

1-((1*H*-indol-7-yl)methyl)-*N*-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1*H*-pyrazole-5-carboxamide,

N-[(2-amino-3-pyridyl)sulfonyl]-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 1106),

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-((1-methylcyclohexyl)methyl)-1*H*-pyrazole-5-carboxamide,

N-(4-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 1653),

3-(*tert*-butyl)-1-(4-(*tert*-butyl)-2,6-dimethylbenzyl)-*N*-(phenylsulfonyl)-1*H*-pyrazole-5-carboxamide,

3-(*tert*-butyl)-*N*-((3-ethylphenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide,

3-(*tert*-butyl)-*N*-((3-chlorophenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide,

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-(1-phenylethyl)pyrazole-3-carboxamide (Compound 799),

N-((3-aminophenyl)sulfonyl)-3-isopropyl-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide,

5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]-*N*-[3-(2-methylpyrimidin-4-yl)phenyl]sulfonyl-

pyrazole-3-carboxamide (Compound 1496),

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(2,4-dichlorobenzyl)-1*H*-pyrazole-5-carboxamide,

3-(*tert*-butyl)-*N*-((3-fluorophenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide,

3-(*tert*-butyl)-*N*-((2-methoxyphenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide,

3-(*tert*-butyl)-*N*-((3-cyanophenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide,

N-((4-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide,

N-((2-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide,

3-(*tert*-butyl)-*N*-(pyridin-4-ylsulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide,

3-(*tert*-butyl)-*N*-((3-methoxyphenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide,

N-((3-aminophenyl)sulfonyl)-3-(*tert*-butyl)-1-(2,4-dimethylbenzyl)-1*H*-pyrazole-5-carboxamide,

5-*tert*-butyl-*N*-(1H-indol-4-ylsulfonyl)-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 180),

3-(*tert*-butyl)-1-(5-methyl-2-(trifluoromethyl)benzyl)-*N*-(phenylsulfonyl)-1*H*-pyrazole-5-carboxamide (Compound 304),

N-((3-aminophenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-benzo[*d*]imidazole-2-carboxamide (Compound 71),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(1-methylindol-4-yl)methyl]pyrazole-3-carboxamide (Compound 927),

5-*tert*-butyl-*N*-[(2-methoxy-3-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 759),

5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]-*N*-(3-thienylsulfonyl)pyrazole-3-carboxamide

(Compound 603),

N-(2-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 88),

5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]-*N*-(1H-indol-7-ylsulfonyl)pyrazole-3-carboxamide (Compound 1358),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-(1H-indol-7-ylmethyl)pyrazole-3-carboxamide (Compound 1256),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(1-methylindol-7-yl)methyl]pyrazole-3-carboxamide (Compound 755),

5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]-*N*-[[2-(methylamino)-4-pyridyl]sulfonyl]pyrazole-3-carboxamide (Compound 821),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(1-methylcyclohexyl)methyl]pyrazole-3-carboxamide (Compound 1152),

N-(benzenesulfonyl)-5-*tert*-butyl-2-[(4-*tert*-butyl-2,6-dimethyl-phenyl)methyl]pyrazole-3-carboxamide (Compound 443),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(1R)-1-phenylethyl]pyrazole-3-carboxamide (Compound 540),

N-[(2-amino-3-pyridyl)sulfonyl]-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide,

5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]-*N*-(3-thienylsulfonyl)pyrazole-3-carboxamide (Compound 603),

5-*tert*-butyl-*N*-[(2-methoxy-3-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 759),

5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]-*N*-(2-furylsulfonyl)pyrazole-3-carboxamide (Compound 1468),

N-[4-(aminomethyl)phenyl]sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 1670),

5-*tert*-butyl-*N*-(4-pyridylsulfonyl)-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 1585),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4-dichlorophenyl)methyl]pyrazole-3-carboxamide (Compound 797),

5-*tert*-butyl-*N*-(3-ethylphenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 823),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-(1-phenylethyl)pyrazole-3-carboxamide,

5-*tert*-butyl-*N*-(3-chlorophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 472),

N-(2-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 1167),

N-(3-aminophenyl)sulfonyl-5-isopropyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 566),

N-(1H-pyrazol-3-ylsulfonyl)-3-[(2,4,6-trimethylphenyl)methyl]-1,2,4,5-tetrahydro-3-benzazepine-4-carboxamide (Compound 628),

N-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-3-[(2,4,6-trimethylphenyl)methyl]-1,2,4,5-tetrahydro-3-benzazepine-4-carboxamide (Compound 662),

N-[(6-morpholino-2-pyridyl)sulfonyl]-3-[(2,4,6-trimethylphenyl)methyl]-1,2,4,5-tetrahydro-3-benzazepine-4-carboxamide (Compound 669),

N-[[6-(methylamino)-2-pyridyl]sulfonyl]-3-[(2,4,6-trimethylphenyl)methyl]-1,2,4,5-tetrahydro-3-benzazepine-4-carboxamide (Compound 747),

N-[[6-(2-hydroxyethylamino)-2-pyridyl]sulfonyl]-3-[(2,4,6-trimethylphenyl)methyl]-1,2,4,5-tetrahydro-3-benzazepine-4-carboxamide (Compound 838),

N-[(2-methoxy-3-pyridyl)sulfonyl]-3-[(2,4,6-trimethylphenyl)methyl]-1,2,4,5-tetrahydro-3-benzazepine-4-carboxamide (Compound 876),

N-[(6-methoxy-2-pyridyl)sulfonyl]-3-[(2,4,6-trimethylphenyl)methyl]-1,2,4,5-tetrahydro-3-benzazepine-4-carboxamide (Compound 948),

N-(benzenesulfonyl)-3-[(2,4,6-trimethylphenyl)methyl]-1,2,4,5-tetrahydro-3-benzazepine-4-carboxamide (Compound 1039),


N-[(6-amino-2-pyridyl)sulfonyl]-3-[(2,4,6-trimethylphenyl)methyl]-1,2,4,5-tetrahydro-3-benzazepine-4-carboxamide (Compound 1055),

N-(2-pyridylsulfonyl)-3-[(2,4,6-trimethylphenyl)methyl]-1,2,4,5-tetrahydro-3-benzazepine-4-carboxamide (Compound 1093),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(1-methylindol-4-yl)methyl]pyrazole-3-carboxamide (Compound 927), and

N-(benzenesulfonyl)-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 45).

PREPARATION 19: 5-*tert*-butyl-*N*-[3-(hydroxymethyl)phenyl]sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 39)




A solution of 3-(*tert*-Butyl)-*N*-((3-formylphenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (prepared following procedures described herein, 50 mg, 0.11 mmol) and anhydrous methanol (5 mL) was cooled to 0 °C. The mixture was purged with nitrogen before adding sodium borohydride (7.6 mg, 0.20 mmol). The mixture was allowed to warm to room temperature and was stirred overnight. The reaction mixture was again cooled to 0 °C before adding additional sodium borohydride (7.6 mg, 0.20 mmol). The mixture was warmed to room temperature and was allowed to stir for 4 h. Solvents were removed and the crude residue was taken up in DMSO and subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give 5-*tert*-butyl-*N*-[3-(hydroxymethyl)phenyl]sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 39) (28 mg, 56%). ^{1}H NMR (400 MHz, DMSO) δ 12.51 (s, 1H), 7.99 (s, 1H), 7.87 (d, J = 7.7 Hz, 1H), 7.63 (dt, J = 15.3, 7.7 Hz, 2H), 7.01 (s, 1H), 6.75 (s, 2H), 5.42 (s, 2H), 4.63 (s, 2H), 2.16 (s, 3H), 2.07 (d, J =

6.4 Hz, 6H), 1.14 (s, 9H). ESI-MS *m/z* calc. 469.2, found 470.0 $(M+1)^+$; Retention time: 2.12 min (3 min run).


PREPARATION 20: *N*-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(2-methoxyphenyl)pyrazole-3-carboxamide (Compound 347)

Step 1: methyl *N*-[(2,6-dimethylphenyl)methylamino]carbamate



2-(Chloromethyl)-1,3-dimethyl-benzene (25.0 g, 162 mmol), methyl *N*-aminocarbamate (29.1 g, 323 mmol) and K_2CO_3 (33.5 g, 243 mmol) were combined in *N,N*-dimethylformamide (150 mL) and heated at 80 °C for 2 h. The reaction mixture was concentrated and the crude material was partitioned between ethyl acetate and water. The organics were separated, washed with brine and dried over sodium sulfate. The mixture was filtered and the filtrate was concentrated. The residue was subjected to silica gel column chromatography (0-60% ethyl acetate/hexanes) to give methyl *N*-[(2,6-dimethylphenyl)methylamino]carbamate (18.0 g, 53%). ESI-MS *m/z* calc. 208.3, found 209.2 $(M+1)^+$; Retention time: 0.41 min (1 min run).


Step 2: methyl 2-[(2,6-dimethylphenyl)methyl]-5-hydroxy-pyrazole-3-carboxylate



Methyl *N*-[(2,6-dimethylphenyl)methylamino]carbamate (18.0 g, 86.4 mmol) was suspended in methanol (126 mL) and dimethyl but-2-ynedioate (10.6 mL, 86.4 mmol) was added. The reaction mixture was stirred at reflux for 2 h and was then cooled to room temperature. Sodium methoxide (20.5 mL of 25 %w/v, 95.1 mmol) was added and the reaction


mixture was again heated at reflux. After 1 h, the mixture was cooled to room temperature and was added drop-wise to a flask containing 1M aq. HCl (300 mL) in an ice water bath. The resulting precipitate was collected by vacuum filtration. The solids were dissolved in CH_2Cl_2 and the excess water was separated and the organic layer was dried over sodium sulfate and evaporated to give methyl 2-[(2,6-dimethylphenyl)methyl]-5-hydroxy-pyrazole-3-carboxylate (21.6 g, 96%). ^{1}H NMR (400 MHz, DMSO) δ 10.05 (s, 1H), 7.10 (dd, J = 8.4, 6.5 Hz, 1H), 7.02 (d, J = 7.5 Hz, 2H), 6.06 (s, 1H), 5.57 (s, 2H), 3.85 (s, 3H), 2.24 (s, 6H). ESI-MS *m/z* calc. 260.1, found 261.4 $(M+1)^+$; Retention time: 0.57 min (1 min run).

Step 3: 2-[(2,6-dimethylphenyl)methyl]-5-(trifluoromethylsulfonyloxy)-pyrazole-3-carboxylate




Methyl 2-[(2,6-dimethylphenyl)methyl]-5-hydroxy-pyrazole-3-carboxylate (8.40 g, 32.3 mmol) and pyridine (3.13 mL, 38.7 mmol) were combined in toluene (84.00 mL) and cooled in a salt water / ice bath. Triflic anhydride (5.98 mL, 35.5 mmol) was added drop-wise keeping the temperature below 0 °C. After the addition was complete, the reaction mixture was warmed to room temperature and stirred for 1 h. The reaction mixture was quenched with water (30 mL) and the layers were separated. The aqueous layer was extracted with toluene (1 x 40 mL). The organics were combined and washed with water (40 mL), dried over sodium sulfate, filtered and evaporated to give methyl 2-[(2,6-dimethylphenyl)methyl]-5-(trifluoromethylsulfonyloxy)-pyrazole-3-carboxylate (12.4 g, 98%) ESI-MS *m/z* calc. 392.1, found 393.2 $(M+1)^+$; Retention time: 0.82 min (1 min run).

Step 4: 2-[(2,6-dimethylphenyl)methyl]-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-3-carboxylate



A suspension of methyl 2-[(2,6-dimethylphenyl)methyl]-5-(trifluoromethylsulfonyloxy)pyrazole-3-carboxylate (12.1 g, 30.7 mmol), 4,4,5,5-tetramethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (8.59 g, 33.8 mmol) and KOAc (9.05 g, 92.2 mmol) in dioxane (240 mL) was degassed for 10 min. Dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (1.26 g, 1.54 mmol) and 1,1'-bis(diphenylphosphino)ferrocene (852 mg, 1.54 mmol) were added and the reaction was degassed for an additional 10 min. The reaction mixture was heated at 100 °C for 4 h. The mixture was concentrated to ~100 mL total volume and was diluted with toluene (250 mL) and water (100 mL). The mixture was filtered through a pad of Celite to remove small black particulates. The organics were separated, washed with water (100 mL) and dried over sodium sulfate. The organics were filtered through a plug of silica (100 g) and the silica was washed with another 1 L of toluene. The toluene was evaporated to give methyl 2-[(2,6-dimethylphenyl)methyl]-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-3-carboxylate (10.6 g, 93%). ^{1}H NMR (400 MHz, Chloroform-d) δ 7.27 (s, 1H), 7.09 (dd, J = 8.2, 6.7 Hz, 1H), 7.00 (d, J = 7.4 Hz, 2H), 5.86 (s, 2H), 3.81 (s, 3H), 2.25 (s, 6H), 1.32 (s, 12H).


Step 5: 2-[(2,6-dimethylphenyl)methyl]-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-3-carboxylic acid



Methyl 2-[(2,6-dimethylphenyl)methyl]-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-3-carboxylate (7.00 g, 14.2 mmol) was dissolved in THF (53 mL) and NaOH (43 mL of 1.0 M, 43 mmol). The reaction mixture was heated at 60 °C for 2 h and was concentrated.


The resulting solid was partitioned between ethyl acetate and a 1 N HCl solution. The organics were separated, washed with brine, dried over sodium sulfate and evaporated to give 2-[(2,6-dimethylphenyl)methyl]-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-3-carboxylic acid (5.5 g, 98%). ^{1}H NMR (400 MHz, DMSO) δ 13.44 (s, 1H), 7.13 (dd, J = 8.3, 6.6 Hz, 1H), 7.07 - 7.02 (m, 3H), 5.81 (s, 2H), 2.23 (s, 6H), 1.24 (s, 12H). ESI-MS m/z calc. 356.2, found 275.4 $(M+1)^+$; Retention time: 0.42 min (1 min run). Note: found mass is [-pinacol+H].

Step 6: *N*-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-3-carboxamide



2-[(2,6-Dimethylphenyl)methyl]-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-3-carboxylic acid (4.14 g, 11.6 mmol), *N*-[(dimethylamino)-1*H*-1,2,3-triazolo-[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (4.42 g, 11.6 mmol) and ethyldiisopropylamine (6.07 mL, 34.9 mmol) were combined in CH_2Cl_2 (116 mL). Benzenesulfonamide (2.19 g, 13.9 mmol) was added and the reaction mixture was stirred for 16 h before it was diluted with CH_2Cl_2 and washed with 1N HCl. The organics were further washed with brine, dried over sodium sulfate, filtered and evaporated. The crude material was purified by silica gel chromatography (0-100% ethyl acetate/hexanes) to give *N*-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-3-carboxamide (4.14 g, 72%). ESI-MS m/z calc. 495.2, found 414.4 $(M+1)^+$; Retention time: 0.68 min (1 min run). Note: found mass is minus the pinacol ester.

Step 7: *N*-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(2-methoxyphenyl)pyrazole-3-carboxamide (Compound 347)



N-(Benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-3-carboxamide (50 mg, 0.10 mmol), 1-bromo-2-methoxybenzene (19 mg, 0.10 mmol), dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (4 mg, 0.005 mmol), and 2M aq. K_2CO_3 (0.20 mL, 0.40 mmol) were combined in *N,N*-dimethylformamide (0.6 mL) and heated at 70 °C for 2 h. The reaction mixture was cooled, filtered and subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mMHCl) to give *N*-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(2-methoxyphenyl)pyrazole-3-carboxamide (Compound 347). ESI-MS *m/z* calc. 475.2, found 476.3 $(M+1)^+$; Retention time: 2.08 min (3 min run).

The following compounds can be synthesized using the procedures described herein:

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-[2-(trifluoromethoxy)phenyl]pyrazole-3-carboxamide (Compound 149),

N-(benzenesulfonyl)-5-(4-chloro-3-methoxy-phenyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 37),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(3-isopropoxyphenyl)pyrazole-3-carboxamide (Compound 133),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-[3-(trifluoromethoxy)phenyl]pyrazole-3-carboxamide (Compound 225),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-[4-(trifluoromethyl)phenyl]pyrazole-3-carboxamide (Compound 316),

N-(benzenesulfonyl)-5-[3-(difluoromethyl)phenyl]-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 315),

N-(benzenesulfonyl)-5-(2,4-dimethoxyphenyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-

carboxamide (Compound 135),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(1-methylindol-4-yl)pyrazole-3-carboxamide (Compound 363),

N-(3-aminophenyl)sulfonyl-5-(2,6-dimethylphenyl)-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 209),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(m-tolyl)pyrazole-3-carboxamide (Compound 202),

N-(3-aminophenyl)sulfonyl-5-(2-fluoro-3-methyl-phenyl)-2-[(2,4,6-trimethylphenyl)methyl] pyrazole-3-carboxamide,

N-(3-aminophenyl)sulfonyl-5-phenyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 367),

N-(benzenesulfonyl)-5-[3-(dimethylamino)phenyl]-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 178),

N-(benzenesulfonyl)-5-[4-(diethylamino)phenyl]-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 162),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(4-fluorophenyl)pyrazole-3-carboxamide (Compound 333),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(3-methoxyphenyl)pyrazole-3-carboxamide (Compound 176),

N-(benzenesulfonyl)-5-(4-dimethylaminophenyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 76),

N-(benzenesulfonyl)-5-(2-cyanophenyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 99),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(3-fluoro-4-methoxy-phenyl)pyrazole-3-carboxamide (Compound 43),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(4-methoxyphenyl)pyrazole-3-carboxamide (Compound 49),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-[3-(trifluoromethyl)phenyl]pyrazole-3-carboxamide (Compound 220),

N-(benzenesulfonyl)-5-(3-chlorophenyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 83),

N-(benzenesulfonyl)-5-(2-chlorophenyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 339),

N-(benzenesulfonyl)-5-(2,5-dimethoxyphenyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 100),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-[4-(trifluoromethoxy)phenyl]pyrazole-3-carboxamide (Compound 159),

ethyl 2-[4-[5-(benzenesulfonylcarbamoyl)-1-[(2,6-dimethylphenyl)methyl]pyrazol-3-yl]phenyl]acetate (Compound 21),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-[5-(5-methyl-1,2,4-oxadiazol-3-yl)-2-thienyl]pyrazole-3-carboxamide (Compound 151),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(2-ethoxyphenyl)pyrazole-3-carboxamide (Compound 152),

N-(3-aminophenyl)sulfonyl-5-(2-methoxy-3-methyl-phenyl)-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 208),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(2-fluorophenyl)pyrazole-3-carboxamide (Compound 109),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(2-isopropoxyphenyl)pyrazole-3-carboxamide (Compound 277),

N-(benzenesulfonyl)-5-(4-chlorophenyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 28),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(4-ethoxyphenyl)pyrazole-3-carboxamide (Compound 29),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(3-ethoxyphenyl)pyrazole-3-

carboxamide (Compound 280),

N-(3-aminophenyl)sulfonyl-5-(2-methoxyphenyl)-2-(1-phenylethyl)pyrazole-3-carboxamide (Compound 40),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(5-fluoro-2-methoxy-phenyl)pyrazole-3-carboxamide,

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-[2-(trifluoromethyl)phenyl]pyrazole-3-carboxamide (Compound 369),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(p-tolyl)pyrazole-3-carboxamide (Compound 199),

N-(benzenesulfonyl)-5-[2-(dimethylamino)phenyl]-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 128),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(3-fluorophenyl)pyrazole-3-carboxamide (Compound 166),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(1-methylindol-5-yl)pyrazole-3-carboxamide (Compound 263),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-[6-(1-piperidyl)-3-pyridyl]pyrazole-3-carboxamide (Compound 329),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(4-fluoro-2-methoxy-phenyl)pyrazole-3-carboxamide (Compound 155),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-[3-(morpholinomethyl)phenyl]pyrazole-3-carboxamide (Compound 302),

N-(3-aminophenyl)sulfonyl-5-(2-methoxy-3-pyridyl)-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 122),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(3-thienyl)pyrazole-3-carboxamide (Compound 357),

N-(3-aminophenyl)sulfonyl-5-(4-methoxyphenyl)-2-(1-phenylethyl)pyrazole-3-carboxamide (Compound 1024),

3-[5-(benzenesulfonylcarbamoyl)-1-[(2,6-dimethylphenyl)methyl]pyrazol-3-yl]benzoic acid (Compound 343),

N-(benzenesulfonyl)-5-(3-*tert*-butoxyphenyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 744),

N-(benzenesulfonyl)-5-(4*H*-1,3-benzodioxin-7-yl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 1621),

N-(benzenesulfonyl)-5-[2-(dimethylamino)pyrimidin-5-yl]-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 1678),

N-(benzenesulfonyl)-5-(5-chloro-3-fluoro-2-methoxy-phenyl)-2-[(2,6-dimethylphenyl)methyl] pyrazole-3-carboxamide,

N-(benzenesulfonyl)-5-(5-chloro-3-fluoro-2-methoxy-phenyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 379),

N-(benzenesulfonyl)-5-(3-*tert*-butoxyphenyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 744),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(4-isopropoxyphenyl)pyrazole-3-carboxamide (Compound 1478),

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(4-oxochromen-3-yl)pyrazole-3-carboxamide (Compound 1501),


N-(3-aminophenyl)sulfonyl-5-(4-methoxyphenyl)-2-(1-phenylethyl)pyrazole-3-carboxamide (Compound 1024)

N-(3-aminophenyl)sulfonyl-5-(2-fluoro-3-methyl-phenyl)-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 293), and

N-(benzenesulfonyl)-2-[(2,6-dimethylphenyl)methyl]-5-(5-fluoro-2-methoxy-phenyl)pyrazole-3-carboxamide (Compound 27).


PREPARATION 21: 5-Benzyloxy-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylic acid

Step 1: methyl 3-(benzyloxy)-1-(2,6-dimethylbenzyl)-1*H*-pyrazole-5-carboxylate



To a solution of methyl 5-hydroxy-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylate (250 mg, 0.911 mmol) and K_2CO_3 (315 mg, 2.28 mmol) in *N*,*N*-dimethylformamide (5 mL) was added benzyl bromide (130 µL, 1.09 mmol) and the reaction mixture was stirred at room temperature for 16 h. The mixture was concentrated and the resulting solids were partitioned between ethyl acetate and water. The organics were separated, washed with brine, dried over sodium sulfate and evaporated to give methyl 3-(benzyloxy)-1-(2,6-dimethylbenzyl)-1*H*-pyrazole-5-carboxylate. ESI-MS *m/z* calc. 364.2, found 365.4 $(M+1)^+$; Retention time: 0.90 min (1 min run).

Step 2: 5-benzyloxy-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylic acid



Methyl 3-(benzyloxy)-1-(2,6-dimethylbenzyl)-1*H*-pyrazole-5-carboxylate (from step 1) was dissolved in methanol (3 mL) before 1N NaOH (3 mL) was added. The reaction mixture was heated for 2 h at 60 °C and concentrated. The crude mixture was partitioned between ethyl acetate and 1 N HCl. The organics were separated, washed with brine, dried over sodium sulfate, and evaporated to yield 5-benzyloxy-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylic acid (295 mg, 92% for 2 steps) ESI-MS *m/z* calc. 350.2, found 351.4 $(M+1)^+$; Retention time: 0.79 min (1 min run).

The following compound can be synthesized using the procedures described herein:

3-(1-phenylethoxy)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxylic acid,


N-(3-aminophenyl)sulfonyl-5-benzyloxy-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-

carboxamide (Compound 42), and

N-(3-aminophenyl)sulfonyl-5-(1-phenylethoxy)-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 278).


PREPARATION 22: *N*-(3-aminophenyl)sulfonyl-5-(3,3-dimethylbutyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 323)

Step 1: 5-[(E)-3,3-dimethylbut-1-enyl]-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxylate




Methyl 2-[(2,6-dimethylphenyl)methyl]-5-(trifluoromethylsulfonyloxy)pyrazole-3-carboxylate (500. mg, 1.27 mmol), [(*E*)-3,3-dimethylbut-1-enyl]boronic acid (196 mg, 1.53 mmol), dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane (52 mg, 0.064 mmol), and 2M K_2CO_3 (1.9 mL of 2.0 M, 3.8 mmol) were combined in dioxane (10 mL) and irradiated in a microwave reactor for 30 min at 120 °C. The reaction mixture was diluted with ethyl acetate and was filtered through a pad of Celite. The filtrate was washed with water, brine, dried over sodium sulfate and concentrated. The crude material was purified by silica gel column chromatography (0-60% ethyl acetate/hexanes) to give methyl 5-[(*E*)-3,3-dimethylbut-1-enyl]-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxylate (300. mg, 72%) ESI-MS *m/z* calc. 326.2, found 327.5 $(M+1)^+$; Retention time: 0.94 min (1 min run).

Step 2: 5-[(*E*)-3,3-dimethylbut-1-enyl]-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxylate




Methyl 5-[(*E*)-3,3-dimethylbut-1-enyl]-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxylate (300. mg, 0.919 mmol) was dissolved in THF (3 mL) before adding NaOH (3.0 mL of 1.0 M, 3.0 mmol). The reaction mixture was heated for 2 h at 60 °C and then concentrated. The resulting solid was partitioned between ethyl acetate and an aqueous solution of 1 N HCl. The organics were separated, washed with brine, dried over sodium sulfate and concentrated to give 5-[(*E*)-3,3-dimethylbut-1-enyl]-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxylic acid (198 mg, 69%). ESI-MS *m/z* calc. 312.2, found 313.5 $(M+1)^+$; Retention time: 0.77 min (1 min run).

Step 3: (*E*)-3-(3,3-dimethylbut-1-en-1-yl)-*N*-((3-nitrophenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide



5-[(*E*)-3,3-dimethylbut-1-enyl]-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxylic acid (50. mg, 0.16 mmol), 3-nitrobenzenesulfonamide (32 mg, 0.16 mmol), *N*-[(dimethylamino)-1*H*-1,2,3-triazolo-[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (61 mg, 0.16 mmol), and ethyldiisopropylamine (84 µL, 0.48 mmol) were combined in *N,N*-dimethylformamide (1 mL) and stirred at room temperature for 16 h. The reaction mixture was filtered and subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mMHCl) to give (*E*)-3-(3,3-dimethylbut-1-en-1-yl)-*N*-((3-nitrophenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (23 mg, 29%). ESI-MS *m/z* calc. 496.2, found 497.2 $^+$; Retention time: 0.85 min (1 min run).

Step 4: *N*-(3-aminophenyl)sulfonyl-5-(3,3-dimethylbutyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 323)




A mixture of (*E*)-3-(3,3-dimethylbut-1-en-1-yl)-*N*-((3-nitrophenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide (23 mg, 0.046 mmol), methanol (2 mL) and 10% Pd/C (17 mg, 0.016 mmol) at room temperature was placed under a balloon of hydrogen The mixture was stirred for 30 min before it was filtered. The filtrate was subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give *N*-(3-aminophenyl)sulfonyl-5-(3,3-dimethylbutyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 323) (4.0 mg, 17%). ESI-MS *m/z* calc. 468.2, found 469.0 $(M+1)^+$; Retention time: 2.11 min (3 min run).

The following compounds can be synthesized using the procedures described herein: *N*-(3-aminophenyl)sulfonyl-2-[(2,6-dimethylphenyl)methyl]-5-norboman-2-yl-pyrazole-3-carboxamide (Compound 370), *N*-[(6-amino-2-pyridyl)sulfonyl]-7-ethyl-2-[(2,4,6-trimethylphenyl)methyl]indazole-3-carboxamide (Compound 1086), and *N*-(3-aminophenyl)sulfonyl-5-cyclopentyl-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 105).


PREPARATION 23: 1-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazole-4-carboxylic acid

Step 1: (*E*)-1-*tert*-butyl-2-[2-(2,4,6-trimethylphenyl)ethylidene]hydrazine




Acetic acid (0.2 mL) was added to a mixture of 2-(2,4,6-trimethylphenyl)acetaldehyde (0.80 g, 4.9 mmol) and *t*-butylhydrazine hydrochloride (0.73 g, 4.9 mmol) in methanol (10 mL). The mixture was stirred at room temperature for 24 h. The reaction mixture was concentrated under reduced pressure and the residue was treated with ether (20 mL) to give a precipitate which was collected by filtration. The solid was suspended in ethyl acetate (150 mL), the pH adjusted to >7 by adding saturated $NaHCO_3$, the organic layer washed with brine, dried over Na_2SO_4, filtered and concentrated under reduced pressure to afford (*E*)-1-*tert*-butyl-2-[2-(2,4,6-trimethylphenyl)ethylidene]hydrazine (0.80 g, 70.%) as a yellow oil. ^{1}H NMR (300 MHz, DMSO-d_6) δ 1.22 (s, 9H), 2.18 (s, 3H), 2.23 (s, 6H), 3.68 (d, J = 5.2 Hz, 2H), 6.85 (s, 2H), 8.24 (t, J = 5.2 Hz, 1H), 11.39 (br s, 1H). ESI-MS *m/z* calc. 232.2, found 233.2 $(M+1)^+$; Retention time: 1.75 min (5 min run).

Step 2: ethyl 1-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazole-4-carboxylate



A mixture of (*E*)-1-*tert*-butyl-2-[2-(2,4,6-trimethylphenyl)ethylidene]hydrazine (0.86 g, 3.7 mmol) and ethylpropiolate (0.44 g, 4.5 mmol) in acetonitrile (3 mL) and acetic acid (3 mL) was stirred at room temperature for 2 d. The reaction mixture was concentrated under reduced pressure and the residue was taken up with ethyl acetate (100 mL) before it was washed with saturated $NaHCO_3$, brine, dried over Na_2SO_4, filtered and concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel, eluting with 30% ethyl acetate in heptane to afford ethyl 1-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazole-4-carboxylate (0.40 g, 33%) as a yellow solid. ESI-MS *m/z* calc. 328.2, found 329.2 $(M+1)^+$; Retention time: 2.92 min (5 min run).

Step 4: 1-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazole-4-carboxylic acid



A mixture of ethyl 1-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazole-4-carboxylate (0.34 g, 1.0 mmol) and $LiOH \cdot H_2O$ (0.22 g, 5.2 mmol) in methanol (4 mL) and water (4 mL) was heated at reflux for 3 h. The reaction mixture was concentrated under reduced pressure and the pH of the aqueous layer adjusted to ~3 with 5% citric acid. The aqueous layer was extracted with ethyl acetate (150 mL), washed with brine, dried over Na_2SO_4, filtered and concentrated under reduced pressure. The residue was purified by C_{18} column chromatography, eluting at 80% methanol in water to afford 1-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazole-4-carboxylic acid (0.23 g, 74%) as a yellow solid. ESI-MS *m/z* calc. 300.2, found 301.2 $(M+1)^+$; Retention time: 2.41 min (5 min run).


The following compounds can be synthesized using the procedures described herein:

N-(benzenesulfonyl)-1-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]pyrazole-4-carboxamide (Compound 63) and

N-(3-aminophenyl)sulfonyl-1-*tert*-butyl-3-[(2,4,6-trimethylphenyl)methyl]pyrazole-4-carboxamide (Compound 324).

PREPARATION 24: 1-*tert*-Butyl-4-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazole-3-carboxylic acid

Step 1: 4-bromo-1-*tert*-butyl-1*H*-pyrazole-3-carboxylic acid



Concentrated sulfuric acid (0.24 mL, 4.3 mmol) was added to a suspension of 4-bromo-1*H*-pyrazole-3-carboxylic acid (860 mg, 4.3 mmol) in *t*-BuOH (2.1 mL, 22 mmol). The reaction mixture was stirred at 100 °C for 2 h. Most of the solvent evaporated during the heating


and a white solid remained in the flask. After cooling to room temperature, the pH was adjusted to ~4 by adding saturated aq. $NaHCO_3$. The aqueous layer was extracted with ethyl acetate (2 x 40 mL). The combined organic layers were washed with brine (20 mL), dried over $MgSO_4$, filtered and concentrated under reduced pressure to afford 4-bromo-1-*tert*-butyl-1*H*-pyrazole-3-carboxylic acid (1.0 g, 89%) as a white solid. ^1H NMR (300 MHz, DMSO) δ 1.52 (s, 9H), 8.23 (s, 1H), 12.92 (br s, 1H).

Step 2: methyl 4-bromo-1-*tert*-butyl-1*H*-pyrazole-3-carboxylate



Potassium carbonate (1.1 g, 8.0 mmol) and iodomethane (0.3 mL, 4.8 mmol) were added to a solution of 4-bromo-1-*tert*-butyl-1*H*-pyrazole-3-carboxylic acid (980 mg, 4.0 mmol) in *N,N*-dimethylformamide (6 mL). The reaction mixture was stirred at room temperature overnight. A 5% citric acid solution (20 mL) was added and the aqueous layer was extracted with ethyl acetate (60 mL). The organic layer was washed with water (2 x 15 mL), brine (15 mL), dried over $MgSO_4$, filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (0-30% ethyl acetate/hexanes) to afford methyl 4-bromo-1-*tert*-butyl-1*H*-pyrazole-3-carboxylate (1.1 g, quantitative) as a colorless oil. ^1H NMR (300 MHz, $CDCl_3$) δ 1.57 (s, 9H), 3.94 (s, 3H), 7.61 (s, 1H).


Step 3: methyl 1-*tert*-butyl-4-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazole-3-carboxylate



Palladium acetate (45 mg, 0.20 mmol) and $^tBu_3P \cdot HBF_4$ (120 mg, 0.40 mmol) were added to a solution of methyl 4-bromo-1-*tert*-butyl-1*H*-pyrazole-3-carboxylate (1.1 g, 4.2 mmol)

in tetrahydrofuran (10 mL) in a sealed tube. The reaction vessel was evacuated and backfilled with nitrogen three times. A solution of freshly prepared (2,4,6-trimethylphenyl)methyl zinc bromide (8.0 mL of a 1.0 M solution in THF, 8.0 mmol) was added at room temperature. The orange-red solution was stirred at room temperature overnight. The resulting black reaction mixture was quenched with 5% citric acid solution (20 mL) and extracted with ethyl acetate (2 x 40 mL). The combined organic layers were washed with water (2 x 15 mL), brine (15 mL), dried over $MgSO_4$, filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (5 to 30% ethyl acetate/hexanes) to give methyl 1-*tert*-butyl-4-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazole-3-carboxylate (1.1 g, 87%) as a white solid. ^1H NMR (300 MHz, $CDCl_3$) δ 1.47 (s, 9H), 2.19 (s, 6H), 2.29 (s, 3H), 3.96 (s, 3H), 4.05 (s, 2H), 6.67 (s, 1H), 6.87 (s, 2H).

Step 4: 1-*tert*-butyl-4-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazole-3-carboxylic acid



A mixture of 1-*tert*-butyl-4-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazole-3-carboxylate (1.00 g, 3.18 mmol), $LiOH \cdot H_2O$ (667 mg, 15.5 mmol), THF (10 mL) and water (10 mL) was stirred in a sealed tube at 60 °C for 4 h. The mixture was acidified by adding 5% citric acid (30 mL) and was extracted with ethyl acetate (2 x 30 mL). The combined organic layers were washed with water (2 x 15 mL), brine (15 mL), dried over $MgSO_4$, filtered and concentrated under reduced pressure. The residue was triturated with a mixture of methyl tert-butyl ether (2 mL) and heptanes (2 mL) to afford 1-*tert*-butyl-4-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazole-3-carboxylic (769 mg, 81%) as a white solid. ^1H NMR (300 MHz, DMSO) δ 1.41 (s, 9H), 2.14 (s, 6H), 2.21 (s, 3H), 3.94 (s, 2H), 6.80 (s, 1H), 6.84 (s, 2H), 12.60 (s, 1H).


The following compound can be synthesized using the procedures described herein:

N-(3-aminophenyl)sulfonyl-1-*tert*-butyl-4-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-

carboxamide (Compound 366).

PREPARATION 25: 1-*tert*-butyl-3-(2,4,6-trimethylphenoxy)-1*H*-pyrazole-4-carboxylic acid

Step 1: 1-*tert*-butyl-1*H*-pyrazol-3-ol



tert-Butylhydrazine hydrochloride (3.03 g, 24.3 mmol) was added to pyridine (30 mL) and the mixture was stirred for 1 h at room temperature. The mixture containing insoluble material was cooled to 5 °C then 3,3-diethoxypropanoyl chloride (3.27 g, 24.3 mmol) was added dropwise. The mixture was then was heated at 80 °C for 48 h before it was cooled to room temperature. The mixture was diluted with ethyl acetate and the organic phase was washed with 5% aq. $NaHCO_3$ and brine before it was dried over anhydrous Na_2SO_4. The solids were filtered off and the solvent was removed under reduced pressure. The residue was purified by silica gel column chromatography (5-50% ethyl acetate/heptanes) to give a solid which was triturated in heptanes (1 x 10 mL), filtered and dried to afford 1-*tert*-butyl-1*H*-pyrazol-3-ol (1.38 g, 44%) as an off-white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.54 (s, 9H), 5.57 (d, J = 2.2 Hz, 1H), 7.24 (d, J = 2.2 Hz, 1H), 12.11 (br. s, 1H). ESI-MS *m/z* calc. 140.1, found 141.2 $(M+1)^+$; Retention time: 1.12 min (5 min run).


Step 2: 4-bromo-1-*tert*-butyl-1*H*-pyrazol-3-ol



To 1-*tert*-butyl-1*H*-pyrazol-3-ol (1.30 g, 9.27 mmol) in dichloromethane (20 mL) was added $NaHCO_3$ (0.97 g, 11. mmol). The mixture was cooled to 0 °C and a solution of bromine (1.56 g, 9.74 mmol) in dichloromethane (10 mL) was added dropwise. The mixture was stirred for 60 min at 0-10 °C before it was diluted with ethyl acetate. The organic phase was washed with 5% $NaHCO_3$, 1% $Na_2S_2O_3$, 5% $NaHCO_3$, brine and dried over anhydrous Na_2SO_4. The solid was filtered off and the solvent was removed under reduced pressure to afford 4-bromo-1-*tert*-butyl-1*H*-pyrazol-3-ol (1.87 g, 92% yield) as an off-white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ


1.53 (s, 9H), 7.31 (s, 1H), 11.58 (br. s, 1H). ESI-MS *m/z* calc. 218.0, found 219.1/221.1 $(M+1)^+$; Retention time: 1.52 min (5 min run).

Step 3: 4-bromo-1-*tert*-butyl-3-(2,4,6-trimethylphenoxy)-1*H*-pyrazole



1-2-Dichloroethane (125 mL) was added to flame-dried 4Å molecular sieves (4.81 g) and air was bubbled through the mixture for 15 minutes. 4-Bromo-1-*tert*-butyl-1*H*-pyrazol-3-ol (1.75 g, 7.99 mmol) was added followed by 2,4,6-trimethylphenylboronic acid (3.27 g, 20.0 mmol), pyridine (1.94 mL, 24.0 mmol) and copper (II) acetate (2.18 g, 12.0 mmol). The reaction mixture was stirred under an air atmosphere for 72 h at room temperature before being diluted with ethyl acetate. The organic phase was washed with 10% NH_4OH, brine and dried over anhydrous Na_2SO_4. The solid was filtered off and the solvent was removed under reduced pressure. The residue was purified by silica gel column chromatography (5-75% dichloromethane/heptanes) to afford 4-bromo-1-*tert*-butyl-3-(2,4,6-trimethylphenoxy)-1*H*-pyrazole (1.35 g, 50%) as pale yellow solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ1.41 (s, 9H), 2.15 (s, 6H), 2.27 (s, 3H), 6.83 (s, 2H), 7.34 (s, 1H). ESI-MS *m/z* calc. 336.1, found 337.1/339.1 $(M+1)^+$; Retention time: 2.54 min (5 min run).

Step 4: 1-*tert*-butyl-3-(2,4,6-trimethylphenoxy)-1*H*-pyrazole-4-carboxylic acid




To a solution of 4-bromo-1-*tert*-butyl-3-(2,4,6-trimethylphenoxy)-1*H*-pyrazole (1.00 g, 2.96 mmol) in THF (8 mL) cooled at -78 °C was added dropwise a 2.5 M solution of *n*-buthyllithium in hexanes (1.36 mL, 3.41 mmol). The mixture was stirred for 30 minutes at -78

°C and then CO_2 was bubbled through the solution for 45 min after which a chunk of dry ice was added to the reaction mixture. The reaction mixture was allowed to warm from -78 °C to -60 °C over 3 h. The reaction mixture was quenched by the addition of water (5 mL) at 0 °C before it was diluted with ethyl acetate (50 mL). Citric acid (5%, aq.) was added under stirring to obtain pH of 4. The mixture was extracted with ethyl acetate. The combined organic extracts were washed with 5% aq. citric acid, 3% HCl, H_2O and brine and dried over anhydrous Na_2SO_4. The solid was filtered off and the solvent was removed under reduced pressure. The residue was triturated twice in heptanes/methyl tert-butyl ether (8 mL/0.5 mL) then purified by silica gel column chromatography 0-20% methanol/dichloromethane) to afford 1-*tert*-butyl-3-(2,4,6-trimethylphenoxy)-1*H*-pyrazole-4-carboxylic acid (480 mg, 53% yield) as white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.42 (s, 9H), 2.15 (s, 6H), 2.28 (s, 3H), 6.85 (s, 2H), 7.91 (s, 1H). ESI-MS *m/z* calc. 302.2, found 303.2 $(M+1)^+$; Retention time: 3.94 min (5 min run).

The following compounds can be synthesized using the procedures described herein: *N*-[(6-amino-2-pyridyl)sulfonyl]-1-*tert*-butyl-3-(2,4,6-trimethylphenoxy)pyrazole-4-carboxamide (Compound 256) and *N*-(3-aminophenyl)sulfonyl-1-*tert*-butyl-3-(2,4,6-trimethylphenoxy)pyrazole-4-carboxamide (Compound 325).

PREPARATION 26: *N*-(3-aminophenyl)sulfonyl-6-cyano-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 1082)


Step 1: methyl 6-cyano-1-(2,4,6-trimethylbenzyl)-1*H*-indole-2-carboxylate



Methyl 6-cyano-1*H*-indole-2-carboxylate (467 mg, 2.33 mmol) was dissolved in *N,N*-dimethylformamide (25 mL) and cooled in an ice bath. NaH (60%, 112 mg, 2.80 mmol) was added and the reaction mixture was stirred for 10 min. 2-(Chloromethyl)-1,3,5-trimethyl-benzene (472 mg, 2.80 mmol) was added and the reaction mixture was allowed to warm to room temperature and was stirred for 12 h. The reaction mixture was poured into ice water and was made acidic with the addition of 1 N HCl. The resulting precipitate was collected to give methyl


6-cyano-1-(2,4,6-trimethylbenzyl)-1*H*-indole-2-carboxylate. ESI-MS *m/z* calc. 332.2, found 333.2 $(M+1)^+$; Retention time: 0.81 min (1 min run).

Step 2: 6-cyano-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxylic acid



Methyl 6-cyano-1-(2,4,6-trimethylbenzyl)-1*H*-indole-2-carboxylate (from step 1) was stirred in a mixture of methanol (20 mL), THF (10 mL) and NaOH (9.34 mL of 1.0 M, 9.34 mmol) for 30 min. The reaction mixture was cooled in an ice bath and was acidified with 6 M HCl to a pH of 1. The mixture was extracted with ethyl acetate. The organics were washed with brine, dried over sodium sulfate and evaporated to give 6-cyano-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxylic acid (740 mg, 99% over two steps). ESI-MS *m/z* calc. 318.1, found 319.2 $(M+1)^+$; Retention time: 0.68 min (1 min run).


Step 3: 6-cyano-*N*-((3-nitrophenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-indole-2-carboxamide



6-Cyano-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxylic acid (35 mg, 0.11 mmol), (3-nitrophenyl)sulfonylazanide sodium (37 mg, 0.16 mmol), *N*-[(dimethylamino)-1H-1,2,3-triazolo-[4,5-b]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (42 mg, 0.11 mmol), and K_2CO_3 (46 mg, 0.33 mmol) were combined in *N,N*-dimethylformamide (1 mL). The reaction mixture was heated at 60 °C for 2 h and was concentrated. The crude material was subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give 6-cyano-*N*-((3-nitrophenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-

indole-2-carboxamide. ESI-MS *m/z* calc. 502.1, found 503.2 $(M+1)^+$; Retention time: 0.75 min (1 min run).

Step 4: *N*-(3-aminophenyl)sulfonyl-6-cyano-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 1082)



6-Cyano-*N*-((3-nitrophenyl)sulfonyl)-1-(2,4,6-trimethylbenzyl)-1*H*-indole-2-carboxamide (from step 3) was mixed with methanol (2 mL) and 10% Pd/C (12 mg, 0.011 mmol) under a balloon of hydrogen. After 1 h, the reaction mixture was filtered and the filtrate was concentrated. The residue was subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mMHCl) to give *N*-(3-aminophenyl)sulfonyl-6-cyano-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 1082) (6.8 mg, 13% over two steps) ESI-MS *m/z* calc. 472.2, found 473.2 $(M+1)^+$; Retention time: 1.71 min (3 min run).

The following compounds can be synthesized using the procedures described herein:

N-(benzenesulfonyl)-6-chloro-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 10),

N-(benzenesulfonyl)-5-methoxy-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 241),

N-(benzenesulfonyl)-4-chloro-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 127),

N-(benzenesulfonyl)-4-fluoro-1-(1-phenylethyl)indole-2-carboxamide (Compound 160),

N-(benzenesulfonyl)-6-methyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 12),

N-(benzenesulfonyl)-5-chloro-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 137),

N-(benzenesulfonyl)-4-fluoro-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 87),

N-(benzenesulfonyl)-5-methyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 248),

N-(benzenesulfonyl)-4-methyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 143),

N-(benzenesulfonyl)-7-methyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 354),

N-(benzenesulfonyl)-5-fluoro-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 33),

N-(3-aminophenyl)sulfonyl-4-chloro-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 362),

N-(3-aminophenyl)sulfonyl-4-methoxy-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 120),

N-[(6-amino-2-pyridyl)sulfonyl]-5-ethyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 179),

N-(benzenesulfonyl)-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 264),

N-(benzenesulfonyl)-1-[(2,6-dimethylphenyl)methyl]-4-methoxy-indole-2-carboxamide (Compound 290),

N-(benzenesulfonyl)-4-methoxy-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 348),

N-(3-aminophenyl)sulfonyl-5-chloro-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 344),

N-(3-aminophenyl)sulfonyl-5-methyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 259),

N-(3-aminophenyl)sulfonyl-6-chloro-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 9),

N-(3-aminophenyl)sulfonyl-6-fluoro-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 142),

N-(benzenesulfonyl)-6-fluoro-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 212),

N-(3-aminophenyl)sulfonyl-4-fluoro-1-(1-phenylethyl)indole-2-carboxamide (Compound 186),

N-(3-aminophenyl)sulfonyl-4-methyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 144),

N-(3-aminophenyl)sulfonyl-1-[(2,6-dimethylphenyl)methyl]-4-methoxy-indole-2-carboxamide (Compound 168),

N-(3-aminophenyl)sulfonyl-4-fluoro-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 181),

N-(3-methoxyphenyl)sulfonyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 335),

N-(3-aminophenyl)sulfonyl-7-methyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 194),

N-(3-aminophenyl)sulfonyl-5-methoxy-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 336),

N-(2-methoxyphenyl)sulfonyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 183),

N-[(6-methoxy-2-pyridyl)sulfonyl]-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 253),

N-(3-aminophenyl)sulfonyl-5-fluoro-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 153),

N-(3-aminophenyl)sulfonyl-6-methyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 62),

N-(benzenesulfonyl)-3-methyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 232),

N-(3-aminophenyl)sulfonyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 148),

N-(2-hydroxyphenyl)sulfonyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 112),

N-(benzenesulfonyl)-5,6-dimethoxy-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 198),

N-(4-aminophenyl)sulfonyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 134),

N-(benzenesulfonyl)-1-[(2,6-dimethylphenyl)methyl]indole-2-carboxamide (Compound 2),

N-(3-aminophenyl)sulfonyl-4-cyano-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 306),

N-[(2-methoxy-3-pyridyl)sulfonyl]-5-methyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 267),

4-fluoro-1-(1-phenylethyl)-*N*-(4-pyridylsulfonyl)indole-2-carboxamide (Compound 192),

N-[(2-methoxy-3-pyridyl)sulfonyl]-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 18),

N-(3-aminophenyl)sulfonyl-1-[(2,6-dimethylphenyl)methyl]indole-2-carboxamide (Compound 331),

N-(2-pyridylsulfonyl)-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 247),

N-(2-cyanophenyl)sulfonyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 358),

4-chloro-*N*-(4-pyridylsulfonyl)-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 82),

6-methyl-*N*-(4-pyridylsulfonyl)-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 38),

4-methoxy-*N*-(4-pyridylsulfonyl)-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 36),

4-fluoro-*N*-(4-pyridylsulfonyl)-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 298),

N-(4-pyridylsulfonyl)-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 227),

5-methyl-*N*-(4-pyridylsulfonyl)-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 25),

N-(3-aminophenyl)sulfonyl-3-methyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 118),

N-(3-aminophenyl)sulfonyl-1-[(2,6-dimethylphenyl)methyl]-6-methoxy-indole-2-carboxamide (Compound 116),

N-[(3-methoxy-2-pyridyl)sulfonyl]-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 258),

5-chloro-*N*-(4-pyridylsulfonyl)-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 65),

6-chloro-*N*-(4-pyridylsulfonyl)-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 24),

N-[(6-amino-2-pyridyl)sulfonyl]-1-[(4-cyano-2,6-dimethyl-phenyl)methyl]-5-methyl-indole-2-carboxamide (Compound 67),

N-[(4-amino-2-pyridyl)sulfonyl]-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 23),

N-(3-aminophenyl)sulfonyl-5-cyano-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 291),

N-(3-aminophenyl)sulfonyl-5,6-dimethoxy-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 598),

N-(3-aminophenyl)sulfonyl-6-cyano-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 1082),

N-[(2-amino-4-pyridyl)sulfonyl]-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 627),

N-[(6-amino-2-pyridyl)sulfonyl]-5-isopropyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 404),

N-[(6-amino-2-pyridyl)sulfonyl]-5-methyl-1-(p-tolyl)indole-2-carboxamide (Compound 653),

N-(3-aminophenyl)sulfonyl-5,6-dimethoxy-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 598),

N-(benzenesulfonyl)-1-[(2,6-dimethylphenyl)methyl]pyrrolo[2,3-b]pyridine-2-carboxamide (Compound 1291),

N-[(2-amino-4-pyridyl)sulfonyl]-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 627),

(2R)-*N*-[(6-amino-2-pyridyl)sulfonyl]-1-[(2,4,6-trimethylphenyl)methyl]indoline-2-carboxamide (Compound 462),

(2R)-*N*-(benzenesulfonyl)-1-[(2,4,6-trimethylphenyl)methyl]indoline-2-carboxamide (Compound 918),

N-[(6-amino-2-pyridyl)sulfonyl]-5-propyl-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxamide (Compound 1158),


N-[(6-amino-2-pyridyl)sulfonyl]-5-ethyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[2,3-b]pyridine-2-carboxamide (Compound 816),

N-(3-aminophenyl)sulfonyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[2,3-b]pyridine-2-carboxamide (Compound 184), and

N-(benzenesulfonyl)-1-[(2,6-dimethylphenyl)methyl]pyrrolo[2,3-b]pyridine-2-carboxamide (Compound 1291).


PREPARATION 27: 4-methoxy-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[3,2-*c*]pyridine-2-carboxylic acid

Step 1: methyl 4-chloro-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[3,2-*c*]pyridine-2-carboxylate



NaH (60%, 62 mg, 1.5 mmol) was added to a mixture of methyl 4-chloro-1*H*-pyrrolo[3,2-*c*]pyridine-2-carboxylate (250. mg, 1.19 mmol) and *N,N*-dimethylformamide (5 mL) at 0 °C. The mixture was allowed to stir for 15 min at 0 °C before a solution of 2-(chloromethyl)-1,3,5-trimethyl-benzene (260. mg, 1.54 mmol) in *N,N*-dimethylformamide (1 mL) was added dropwise. The mixture was allowed to stir for 1H at room temperature, then 4h at 60 °C. The mixture was partitioned between 1N HCl and ethyl acetate. The layers were separated and the aqueous layer was extracted with ethyl acetate (3x). The combined organic layers were washed with brine (2x), dried over magnesium sulfate, filtered and concentrated under reduced pressure. The clear residue was subjected to silica gel column chromatography (0-20% ethyl acetate/hexanes) to give methyl 4-chloro-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[3,2-*c*]pyridine-2-carboxylate (370 mg, 91%). ESI-MS *m/z* calc. 342.1, found 343.4 $(M+1)^+$; Retention time: 2.09 min (3 min run).


Step 2: methyl 4-methoxy-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[3,2-*c*]pyridine-2-carboxylate



A mixture of methyl 4-chloro-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[3,2-c]pyridine-2-carboxylate (130. mg, 0.379 mmol) and NaOMe (7.6 mL of 0.50 M, 3.8 mmol) was heated at 110 °C for 30 min in the microwave. The mixture was concentrated to give methyl 4-methoxy-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[3,2-*c*]pyridine-2-carboxylate (56 mg, 15%) as a white solid. ESI-MS *m/z* calc. 338.2, found 339.4 $(M+1)^+$; Retention time: 0.64 min (1 min

run).

Step 3: 4-methoxy-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[3,2-*c*]pyridine-2-carboxylic acid




Lithium hydroxide (160 µL of 2.0 M, 0.32 mmol) was added to a mixture of methyl 4-methoxy-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[3,2-*c*]pyridine-2-carboxylate (50. mg, 0.15 mmol) and EtOH (1 mL) at room temperature. The mixture was allowed to stir at room temperature overnight. The mixture was concentrated and the residue was partitioned between 1N HCl and CH_2Cl_2. The layers were separated and the aqueous layer was extracted with CH_2Cl_2 (3x). The combined organics were washed with brine, dried over magnesium sulfate, filtered and concentrated. The residue was taken up in *N,N*-dimethylformamide and subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give 4-methoxy-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[3,2-*c*]pyridine-2-carboxylic acid (39 mg, 81%) as a white solid. ESI-MS *m/z* calc. 324.2, found 325.4 $(M+1)^+$; Retention time: 0.52 min (1 min run).

The following compound can be synthesized using the procedures described herein: *N*-(3-aminophenyl)sulfonyl-4-methoxy-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[3,2-c]pyridine-2-carboxamide (Compound 943)

and

N-[(6-amino-2-pyridyl)sulfonyl]-4-methoxy-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[2,3-b]pyridine-2-carboxamide (Compound 539)

PREPARATION 28: 2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxylic acid hydrochloride



A suspension of isoindoline-1-carboxylic acid (100 mg, 0.613 mmol) and KOH (103 mg, 1.84 mmol) in isopropanol (600 µL) was stirred until a solution was formed. 2-(Chloromethyl)-1,3,5-trimethyl-benzene (114 mg, 0.674 mmol) was added and the reaction mixture was stirred for 3 h. The mixture was concentrated and the residue was subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give 2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxylic acid hydrochloride (99 mg, 49%). ^{1}H NMR (400 MHz, DMSO) δ 7.39 - 7.33 (m, 1H), 7.34 - 7.25 (m, 3H), 6.87 (s, 2H), 4.97 (s, 1H), 4.34 - 3.94 (m, 4H), 2.40 (s, 6H), 2.23 (s, 3H). ESI-MS *m/z* calc. 295.2, found 296.2 $(M+1)^+$; Retention time: 0.43 min (1 min run).

The following compounds can be synthesized using the procedures described herein:

N-(3-aminophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 90),

N-(4-pyridylsulfonyl)-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 94),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 73),

N-(3-methoxyphenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 16),

N-(benzenesulfonyl)-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 6),

N-(3-hydroxyphenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 254),

N-[(6-methoxy-2-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide

(Compound 174),

N-[(6-oxo-1*H*-pyridin-2-yl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 319),

N-[(2-methoxy-3-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 218),

N-(2-methoxyphenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 79),

N-(2-pyridylsulfonyl)-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 48),

N-[(6-methoxy-3-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 200),

N-(3-aminophenyl)sulfonyl-2-(1-phenylethyl)isoindoline-1-carboxamide (Compound 187),

N-(4-aminophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 121),

N-[(4-amino-2-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 32),

N-(2-aminophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 230),

N-[(6-amino-2-pyridyl)sulfonyl]-5-methyl-2-[(2,4,6-trimethylphenyl)methyl]-3,4-dihydro-1*H*-isoquinoline-3-carboxamide (Compound 95),

N-(3-aminophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]-3,4-dihydro-1*H*-isoquinoline-3-carboxamide (Compound 213),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[1-(2,4,6-trimethylphenyl)ethyl]isoindoline-1-carboxamide (Compound 1186),

N-(benzenesulfonyl)-1-[(2,4,6-trimethylphenyl)methyl]-2,3,4,5-tetrahydro-1-benzazepine-2-carboxamide (Compound 604),

N-[(6-amino-2-pyridyl)sulfonyl]-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-

carboxamide (Compound 895),

N-[(3-hydroxy-2-pyridyl)sulfonyl]-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 377),

N-[(6-amino-2-pyridyl)sulfonyl]-4-ethoxy-2-[(2-fluoro-6-methyl-phenyl)methyl]-1-methyl-isoindoline-1-carboxamide (Compound 1392),

N-(3-amino-4-propyl-phenyl)sulfonyl-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1367),

N-(3-amino-4-isopropyl-phenyl)sulfonyl-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1562),

N-(3-amino-4-ethyl-phenyl)sulfonyl-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1584),

N-[(6-amino-2-pyridyl)sulfonyl]-4-ethoxy-1-methyl-2-(o-tolylmethyl)isoindoline-1-carboxamide (Compound 1408)

4-ethoxy-*N*-indolin-5-ylsulfonyl-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1422)

N-(2-aminophenyl)sulfonyl-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1457)

N-(4-aminophenyl)sulfonyl-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1480)

4-ethoxy-1-methyl-*N*-[[6-[(2,2,2-trifluoroacetyl)amino]-2-pyridyl]sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1616)

N-[(6-acetamido-2-pyridyl)sulfonyl]-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1415)

N-[(6-amino-3-pyridyl)sulfonyl]-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1413)

4-ethoxy-*N*-(2-hydroxyphenyl)sulfonyl-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1509)

4-ethoxy-*N*-(4-hydroxyphenyl)sulfonyl-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1453)

4-ethoxy-1-methyl-*N*-(3-pyridylsulfonyl)-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1669)

N-[(6-amino-5-chloro-2-pyridyl)sulfonyl]-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1596)

4-ethoxy-*N*-(3-methoxyphenyl)sulfonyl-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1470)

N-[(6-amino-2-pyridyl)sulfonyl]-4-ethoxy-2-[(3-fluoro-5-isobutoxy-phenyl)methyl]-1-methyl-isoindoline-1-carboxamide (Compound 1681)

4-ethoxy-1-methyl-*N*-thiazol-5-ylsulfonyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1606)

4-ethoxy-*N*-(1H-indol-6-ylsulfonyl)-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1393)

N-[[6-(difluoromethyl)-2-pyridyl]sulfonyl]-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1597)

4-ethoxy-*N*-[(3-hydroxy-2-pyridyl)sulfonyl]-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1642) (Compound 377)

4-ethoxy-1-methyl-*N*-thiazol-4-ylsulfonyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1542)

4-ethoxy-1-methyl-*N*-(4-pyridylsulfonyl)-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1449)

4-ethoxy-*N*-(1H-indol-4-ylsulfonyl)-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1666)

N-(2-acetamidothiazol-5-yl)sulfonyl-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1555)

4-ethoxy-*N*-(3-fluoro-2-methoxy-phenyl)sulfonyl-1-methyl-2-[(2,4,6-

trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1403)

4-ethoxy-1-methyl-*N*-(2-pyridylsulfonyl)-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1474)

4-ethoxy-1-methyl-*N*-thiazol-2-ylsulfonyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1546)

4-ethoxy-*N*-(3-fluorophenyl)sulfonyl-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1425)

N-[(2-amino-3-pyridyl)sulfonyl]-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1591)

N-[(6-amino-2-pyridyl)sulfonyl]-4-ethoxy-2-[(6-isopropoxy-3-pyridyl)methyl]-1-methyl-isoindoline-1-carboxamide (Compound 1662)

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2,6-dimethylphenyl)methyl]-4-ethoxy-1-methyl-isoindoline-1-carboxamide (Compound 1498)

2-[(2,6-dimethylphenyl)methyl]-4-ethoxy-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-1-methyl-isoindoline-1-carboxamide (Compound 1676)

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(6-chloro-3-pyridyl)methyl]-4-ethoxy-1-methyl-isoindoline-1-carboxamide (Compound 1565)

N-(benzenesulfonyl)-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1539)

N-(3-aminophenyl)sulfonyl-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1454)

N-[(6-amino-2-pyridyl)sulfonyl]-4-ethoxy-2-[(4-isopropylphenyl)methyl]-1-methyl-isoindoline-1-carboxamide (Compound 1576)

4-ethoxy-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-2-[(4-isopropylphenyl)methyl]-1-methyl-isoindoline-1-carboxamide (Compound 1655)

N-[(6-amino-2-pyridyl)sulfonyl]-4-ethoxy-2-[(4-fluorophenyl)methyl]-1-methyl-isoindoline-1-carboxamide (Compound 1433)

4-ethoxy-2-[(4-fluorophenyl)methyl]-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-1-methyl-isoindoline-1-carboxamide (Compound 1583)

N-[(6-amino-2-pyridyl)sulfonyl]-4-ethoxy-2-[(3-fluoro-5-methoxy-phenyl)methyl]-1-methyl-isoindoline-1-carboxamide (Compound 1421)

4-ethoxy-2-[(3-fluoro-5-methoxy-phenyl)methyl]-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-1-methyl-isoindoline-1-carboxamide (Compound 1428)

(1S)-*N*-[(6-amino-2-pyridyl)sulfonyl]-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1372) (Compound 1346)

(1R)-*N*-[(6-amino-2-pyridyl)sulfonyl]-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1582) (Compound 1346)

(1S)-4-ethoxy-1-methyl-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1541) (Compound 1237) (Compound 410)

(1R)-4-ethoxy-1-methyl-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1476) (Compound 1237) (Compound 410)

N-(3-amino-4-chloro-phenyl)sulfonyl-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1587)

N-(3-amino-4-fluoro-phenyl)sulfonyl-4-ethoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1394)

N-(3-aminopyrazin-2-yl)sulfonyl-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 480),

1-methyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1237),

N-(3-aminophenyl)sulfonyl-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1273),

N-(benzenesulfonyl)-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 873),

1-methyl-*N*-(1H-pyrazol-5-ylsulfonyl)-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 378),

N-[(6-amino-2-pyridyl)sulfonyl]-1-methyl-2-(o-tolylmethyl)isoindoline-1-carboxamide (Compound 666),

N-[(2-methoxy-3-pyridyl)sulfonyl]-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 765),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2,6-dimethylphenyl)methyl]-1-methyl-isoindoline-1-carboxamide (Compound 790),

N-[(3-methoxy-2-pyridyl)sulfonyl]-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 825),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2,4-dimethylphenyl)methyl]-1-methyl-isoindoline-1-carboxamide (Compound 1058),

N-[(2-methoxy-3-pyridyl)sulfonyl]-3,3-dimethyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1235),

4-ethoxy-1-methyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1237) (Compound 410),

N-[(6-amino-2-pyridyl)sulfonyl]-7-methoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1075),

N-[(6-amino-2-pyridyl)sulfonyl]-6-methoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 474),

N-[(6-amino-2-pyridyl)sulfonyl]-4-methoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1161),

N-[(6-amino-2-pyridyl)sulfonyl]-5-methoxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 406),

4-ethoxy-*N*-[(2-methoxy-3-pyridyl)sulfonyl]-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 765) (Compound 648),

N-[(6-amino-2-pyridyl)sulfonyl]-4-ethoxy-1-methyl-2-[(2,4,6-

trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1346),

N-(3-amino-4-methyl-phenyl)sulfonyl-3,3-dimethyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 448),

N-(5-amino-2-methyl-phenyl)sulfonyl-3,3-dimethyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 840),

N-[(6-amino-2-pyridyl)sulfonyl]-7-hydroxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 394),

N-[(6-amino-2-pyridyl)sulfonyl]-6-hydroxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 885),

N-[(6-amino-2-pyridyl)sulfonyl]-4-hydroxy-1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1338),

N-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]-1,3,4,5-tetrahydro-2-benzazepine-3-carboxamide (Compound 855),

N-[(2-methoxy-3-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]-1,3,4,5-tetrahydro-2-benzazepine-3-carboxamide (Compound 1342),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]-1,3,4,5-tetrahydro-2-benzazepine-3-carboxamide (Compound 690),

3,3-dimethyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1085),

N-[(6-amino-2-pyridyl)sulfonyl]-3,3-dimethyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 1101),

N-(benzenesulfonyl)-2-[(2,4,6-trimethylphenyl)methyl]-1,3,4,5-tetrahydro-2-benzazepine-1-carboxamide (Compound 674),


N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]-1,3,4,5-tetrahydro-2-benzazepine-1-carboxamide (Compound 696),

N-(3-aminophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]-3,4-dihydro-1*H*-isoquinoline-1-carboxamide (Compound 309), and

(2S,4R)-*N*-[(6-amino-2-pyridyl)sulfonyl]-4-phenyl-1-[(2,4,6-trimethylphenyl)methyl] pyrrolidine-2-carboxamide.


PREPARATION 29: 4-isopropyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 103)

Step 1: 2,6-dibromobenzaldehyde



To a solution of iPr_2NH (21.2 mL, 150 mmol) in THF (180 mL) at 0 °C was slowly added *n*-BuLi (1.6 M, 94 mL, 150 mmol). The reaction mixture was stirred at 0 °C for 30 min and then cooled to -78 °C. 1,3-Dibromobenzene (17.6 g, 74.6 mmol) in THF (80 mL) was slowly added over 20 min. The mixture was stirred at -78 °C for 30 min and *N,N*-dimethylformamide (11.6 mL, 150 mmol) was added. The mixture was stirred for 1 h, then treated with H_2SO_4 (2.5 M, 350 mL), and extracted with ethyl acetate/Et_2O (50/50, 3 x 300 mL). The combined extract were dried over $MgSO_4$ and filtered. The filtrate was concentrated to give 2,6-dibromobenzaldehyde (17.6 g, 89%). ^1H NMR (250 MHz, $CDCl_3$) δ 10.3 (s, 1H), 7.65 (d, *J*=8.0 Hz, 2H), 7.23 (t, *J*= 8.1 Hz, 1H).


Step 2: methyl (2,6-dibromobenzyl)glycinate



To a solution of glycine methyl ester hydrochloride (21.0 g, 167 mmol), 2.6-dibromobenzaldehyde (16.9 g, 64.4 mmol) and triethylamine (21.0 mL, 142 mmol) in dichloroethane (380 mL) were added acetic acid (8.40 mL, 145 mmol) and $NaBH(OAc)_3$ (33.5 g, 158 mmol). The mixture was stirred at room temperature under an argon atmosphere for 20 h. Additional glycine methyl ester hydrochloride (8.10 g, 64.4 mmol), triethylamine (9.52 g, 64.4 mmol), acetic acid (3.73 mL, 64.4 mmol) and $NaBH(OAc)_3$ (13.7 g, 64.4 mmol) were added and the reaction mixture was stirred for another 10 h. The reaction mixture was quenched by adding 1N NaOH, and the product was extracted with dichloromethane. The organic extracts were


washed with brine, dried over sodium sulfate, filtered, and the solvent was removed in vacuo. The residue was purified by a plug filtration through a silica gel pad (0-50% ethyl acetate/hexanes to give methyl methyl (2,6-dibromobenzyl)glycinate (13.3g, 62%) as an oil. ^{1}H NMR ($CDCl_3$, 250 MHz) δ 7.52 (d, J = 8.0 Hz, 2H), 6.98 (t, J= 8.0, 1H), 4.20 (s, 2H), 3.70 (s, 3H), 3.50 (s, 2H).

Step 3: methyl *N*-(2,6-dibromobenzyl)-*N*-(2,4,6-trimethylbenzyl)glycinate



A mixture of methyl (2,6-dibromobenzyl)glycinate (13.3 g, 39.5 mmol), 2,4,6-trimethylbenzyl bromide (7.32 g, 43.4 mmol), K_2CO_3 (12.0 g, 86.8 mmol), and LiI (300 mg) in acetonitrile (300 mL) was stirred at 60 °C for 24 h. The reaction mixture was filtered and the filtrate was concentrated. The residue was purified by silica gel column chromatography (0-20% ethyl acetate/hexanes) to give methyl *N*-(2,6-dibromobenzyl)-*N*-(2,4,6-trimethylbenzyl)glycinate (18.0 g, 97%) as an oil which solidified on standing. ^{1}H NMR ($CDCl_3$, 250 MHz) δ 7.54 (d, J = 7.8 Hz, 2H), 7.00 (t, J = 8.0, 1H), 6.79 (s, 2H), 4.34 (s, 2H), 4.04 (s, 2H), 3.71 (s, 3H), 3.50 (s, 2H), 2.23 (s, 3H), 2.18 (s, 6H).


Step 4: methyl 4-bromo-2-(2,4,6-trimethylbenzyl)isoindoline-1-carboxylate



A mixture of methyl *N*-(2,6-dibromobenzyl)-*N*-(2,4,6-trimethylbenzyl)glycinate (16.0 g, 34.1 mmol), Cs_2CO_3 (33.3 g, 102 mmol) and $Pd(PPh_3)_4$ (2.0 g, 1.7 mmol) in THF (200 mL) was stirred at 110 °C in a sealed vessel for 60 h. Some THF (~50 mL) was removed by flushing with N_2 before Cs_2CO_3 (6.5 g, 20 mmol) and $Pd(PPh_3)_4$ (400 mg, 0.34 mmol) were added. The

reaction mixture was stirred at 110 °C for 24 h. The mixture was poured into water and was extracted with Et_2O. The organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was purified by silica gel column chromatography (5-20% CH_2Cl_2/hexanes) to give methyl 4-bromo-2-(2,4,6-trimethylbenzyl)isoindoline-1-carboxylate (2.9 g, 22%). MS: 389 $[M+1]^+$.


Step 5: 4-bromo-2-(2,4,6-trimethylbenzyl)isoindoline-1-carboxylic acid hydrochloride



Methyl 4-bromo-2-(2,4,6-trimethylbenzyl)isoindoline-1-carboxylate (2.9 g, 7.4 mmol) was combined in a mixture of methanol (45 mL) and water (20 mL) containing sodium hydroxide (900. mg, 22.5 mmol). The resulting suspension was heated at 80 °C for 1 h. The reaction mixture was concentrated to remove the methanol. Water (50 mL) was added and the mixture was extracted with Et_2O. The aqueous phase was adjusted to pH ~2 with 6N HCl and was then extracted with CH_2Cl_2. The extract was dried over Na_2SO_4, filtered and concentrated. The residue was purified by silica gel column chromatography (0-5% methanol/CH_2Cl_2) and then subjected to preparatory-HPLC (10-100% acetonitrile/water with 0.05 mM trifluoroacetic acid) to give the desired product as the trifluoroacetic acid salt. The material was dissolved in methanol (30 mL) and was treated with HCl (1M, 10 mL). The mixture was then concentrated to dryness. The solid was suspended in CH_2Cl_2 (15 mL), sonicated for 10 min and filtered. The solid was washed with Et_2O and hexanes, and dried to give 4-bromo-2-(2,4,6-trimethylbenzyl)isoindoline-1-carboxylic acid hydrochloric acid (1.5 g, 49%). ^1H NMR (DMSO, 250 MHz) δ 7.65 (d, J = 7.8 Hz, 1H), 7.51 (d, J= 7.8, 1H), 7.39 (m, 1H), 6.95 (s, 2H), 5.87 (s, 1H), 4.68 (s, 2H), 4.52 (d, J= 10.8 Hz, 1H), 4.30 (d, J= 10.8 Hz, 1H), 2.46 (s, 6H), 2.26 (s, 3H). MS: 375 $[M+1]^+$.


Step 6: 4-isopropenyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxylic

acid hydrochloride



4-Bromo-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxylic acid hydrochloride (374 mg, 0.914 mmol), 2-isopropenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (269 mg, 1.60 mmol) and cesium carbonate (652 mg, 2.00 mmol) in dioxane (9 mL) and water (900 μL) was degassed with a stream of nitrogen for 1 min before adding dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (73 mg, 0.10 mmol). The reaction mixture was degassed with a stream of nitrogen for 1 min and the mixture was stirred at 120 °C for 2 h. The reaction mixture was filtered and subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give 4-isopropenyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxylic acid hydrochloric acid (180 mg, 48%). ESI-MS *m/z* calc. 335.2, found 336.2 $(M+1)^+$; Retention time: 0.53 min (1 min run).


Step 7: 4-isopropyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxylic acid hydrochloride



A mixture of 4-(prop-1-en-2-yl)-2-(2,4,6-trimethylbenzyl)isoindoline-1-carboxylic acid hydrochloride (43 mg, 0.12 mmol) and 10% palladium on carbon (5 mg, 0.005 mmol) in methanol (3 mL) was stirred under an atmosphere of hydrogen for 5 h. The reaction mixture was filtered and subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give 4-isopropyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxylic acid hydrochloride


(28 mg, 65%) as a colorless solid. ESI-MS *m/z* calc. 337.2, found 338.2 $(M+1)^+$; Retention time: 0.55 min (1 min run).

Step 8: 4-isopropyl-*N*-[(2-methoxy-3-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 218)



A mixture of 4-isopropyl-2-(2,4,6-trimethylbenzyl)isoindoline-1-carboxylic acid hydrochloride (28 mg, 0.075 mmol) and sodium carbonate (24 mg, 0.22 mmol) in *N,N*-dimethylformamide (1 mL) was degassed with a stream of nitrogen for 1 min before adding *N*-[(dimethylamino)-1H-1,2,3-triazolo-[4,5-b]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (34 mg, 0.090 mmol). The mixture was degassed for 1 min and 2-methoxypyridine-3-sulfonamide (21 mg, 0.11 mmol) was added to the reaction mixture. The mixture was degassed for 1 min before it was stirred at room temperature for 17 h. The reaction mixture was filtered and subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give 4-isopropyl-*N*-[(2-methoxy-3-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 218) (16 mg, 42%). ESI-MS *m/z* calc. 507.2, found 508.4 $(M+1)^+$; Retention time: 0.68 min (1 min run).

Step 9: 4-isopropyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 103)



A mixture of 4-isopropyl-*N*-((2-methoxypyridin-3-yl)sulfonyl)-2-(2,4,6-

trimethylbenzyl)isoindoline-1-carboxamide (16 mg, 0.032 mmol) in acetic acid (1 mL) and HBr (46 µL of 33 %w/v, 0.19 mmol) (in acetic acid) was stirred at 50 °C for 1 h. The solvent was evaporated and the residue was subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give 4-isopropyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 103) (12 mg, 69%). ESI-MS *m/z* calc. 493.2, found 494.3 $(M+1)^+$; Retention time: 1.44 min (3 min run).

The following compounds can be synthesized using the procedures described herein: *N*-(3-aminophenyl)sulfonyl-4-ethyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 30),

N-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-4-phenyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 508),

N-[(2-methoxy-3-pyridyl)sulfonyl]-4-phenyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 427),

N-[(6-amino-2-pyridyl)sulfonyl]-4-phenyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 554), and

N-(3-aminophenyl)sulfonyl-4-isopropyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 299).

PREPARATION 30: 2-(*tert*-butyl)-4-(mesityloxy)pyrimidine-5-carboxylic acid


Step 1: ethyl 2-(*tert*-butyl)-4-hydroxypyrimidine-5-carboxylate



In EtOH (40 mL) was placed *t*-butylcarbamidine hydrochloride (3.71 g, 27.2 mmol). A solution of NaOEt (21% in EtOH, 8.80 g, 27.2 mmol) was added and the resulting mixture was stirred 15 min at room temperature. Diethyl ethoxymethylenemalonate (5.87 g, 27.2 mmol) was added and the reaction mixture was stirred at room temperature overnight. The reaction mixture was heated at reflux for 1 h and then the solvent was removed. The residue was dissolved in water (100 mL) and the pH adjusted to 3-4 with $NaHSO_4$ (1M). The solid was


collected by filtration, washed with water (50 mL) and dried. The filtrate and washings were combined and extracted with ethyl acetate (2 x 150 mL). The combined extracts were dried over Na_2SO_4, filtered and concentrated. The residue plus the solid collected during filtration were subjected to column chromatography on silica gel, eluted with CH_2Cl_2/hexanes/methanol (50/50/0 to 45/45/10) to afford ethyl 2-(*tert*-butyl)-4-hydroxypyrimidine-5-carboxylate (2.85 g, 60%). ^1H NMR (250 MHz, $CDCl_3$) δ 8.74 (s, 1H), 4.36 (q, J = 7.3 Hz, 2H), 1.43 (s, 9H), 1.37 (t, J = 7.2 Hz, 3H).

Step 2: ethyl 2-(*tert*-butyl)-4-chloropyrimidine-5-carboxylate



At 0 °C, Et_3N (1.1 mL) was added to $POCl_3$ (32 mL), followed by the addition of ethyl 2-*tert*-butyl-4-hydroxypyrimidine-5-carboxylate (3.70 g, 16.5 mmol). The mixture was stirred at 40 °C for 1 h and the excess $POCl_3$ was removed under vacuum. The residue was carefully poured into ice (400 mL) and was extracted with CH_2Cl_2. The extract was washed with $NaHCO_3$ and water, dried over Na_2SO_4, filtered and concentrated. The crude material was subjected to silica gel column chromatography (10-90% CH_2Cl_2/hexanes) to afford ethyl 2-(*tert*-butyl)-4-chloropyrimidine-5-carboxylate (3.8 g, 95%) as a liquid. $(M+1)^+$ = 243.4.


Step 3: 2-*tert*-butyl-4-(2,4,6-trimethylphenoxy)pyrimidine-5-carboxylate



2,4,6-Trimethylphenol (2.10 g, 15.4 mmol) and sodium hydride (60%, 690 mg, 17.3 mmol) were combined in an anhydrous mixture of *N,N*-dimethylformamide (20 mL) and tetrahydrofuran (2 mL). The reaction mixture was stirred at room temperature for 15 min under an atmosphere of nitrogen. Ethyl 2-*tert*-butyl-4-chloropyrimidine-5-carboxylate (3.80 g, 15.7

mmol) was added and the reaction mixture was heated at 60 °C for 1 h. The reaction mixture was then allowed to cool to room temperature. The reaction mixture was diluted with water and extracted with ethyl acetate. The ethyl acetate layer was washed three times with water, dried over sodium sulfate, filtered and concentrated. The crude material was purified by silica gel column chromatography (0-30% ethyl acetate/hexanes) to yield ethyl 2-*tert*-butyl-4-(2,4,6-trimethylphenoxy)pyrimidine-5-carboxylate (3.1 g, 58%) as a liquid. ESI-MS *m/z* 351 $(M+1)^+$; ^{1}H NMR (250 MHz, $CDCl_3$) δ 9.07 (s, 1H), 6.88 (s, 2H), 4.43 (q, *J* = 7.2 Hz, 2H), 2.30 (s, 3H), 2.21 (s, 6H), 1.40 (t, *J* = 7.2 Hz, 3H), 1.17 (s, 9H).

Step 4: 2-*tert*-butyl-4-(2,4,6-trimethylphenoxy)pyrimidine-5-carboxylic acid



Ethyl 2-*tert*-butyl-4-(2,4,6-trimethylphenoxy)pyrimidine-5-carboxylate (3.1 g, 9.1 mmol) was combined in a mixture of methanol (25 mL) and water (30 mL) containing sodium hydroxide (1.2 g, 30 mmol). The resulting suspension was heated at 60 °C for 3 h. The reaction mixture was cooled to room temperature and was made acidic with 6M hydrochloric acid. The resulting precipitate was filtered, washed with water and dried to yield 2-*tert*-butyl-4-(2,4,6-trimethylphenoxy)pyrimidine-5-carboxylic acid (2.2 g, 77%) as a white solid. ESI-MS *m/z* 315 $(M+1)^+$; ^{1}H NMR (250 MHz, DMSO) δ 9.01 (s, 1H), 6.95 (s, 2H), 2.27 (s, 3H), 1.98 (s, 6H), 1.11 (s, 9H).

The following compound can be synthesized using the procedures described herein: *N*-(3-aminophenyl)sulfonyl-2-*tert*-butyl-4-(2,4,6-trimethylphenoxy)pyrimidine-5-carboxamide (Compound 68).


PREPARATION 31: 4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-imidazole-2-carboxylic acid

Step 1: 4-*tert*-butyl-1*H*-imidazole




A mixture of 1-bromo-3,3-dimethylbutan-2-one (2.00 g, 11.2 mmol) in formamide (10 mL) was heated between 160-180°C in a sealed tube equipped with a safety pressure release mechanism for 5 h. The mixture was cooled to room temperature and was poured into 10% aqueous sodium bicarbonate (30 mL). The solution was extracted with dichloromethane (2 x 30 mL). The combined organic layers were washed with 10% potassium carbonate, brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to provide 4-*tert*-butyl-1*H*-imidazole (1.37 g, 99%) as an oil. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.31 (s, 9H), 6.76 (s, 1H), 7.56 (s, 1H), 9.63 (br.s., 1H).

Step 2: 4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-imidazole



To 4-*tert*-butyl-1*H*-imidazole (1.37 g, 11.0 mmol) and cesium carbonate (7.20 g, 22.0 mmol) in acetonitrile (24 mL) was added 2-(bromomethyl)-1,3,5-trimethyl benzene (1.58 g, 9.38 mmol). The resulting suspension was stirred at room temperature overnight. Heptane (10 mL) was added to the reaction mixture before it was filtered through a pad of silica and washed with a 1:1 mixture of ethyl acetate:heptane. The eluent was concentrated to provide 4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-imidazole (1.47g, 52%) as an oil. ^{1}H NMR (300 MHz, DMSO) δ1.15 (s, 9H), 2.26 (s, 6H), 2.29 (s, 3H), 5.02 (s, 2H), 6.43 (s, 1H), 6.90 (s, 2H), 7.23 (s, 1H).

Step 3: 4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-imidazole-2-carboxylic acid




To a solution of 4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-imidazole (1.40 g, 5.46 mmol) in anhydrous THF (40 mL) was added *n*-BuLi (2.5 M, 2.62 mL, 6.55 mmol) at –78 °C. The reaction mixture was stirred at –78 °C for 1H. A piece of dry ice was added in one portion at –78 °C. The solution was stirred for 20 min at –78 °C, and then the reaction mixture was allowed to reach to room temperature and was stirred under a CO_2 atmosphere for 2 h. The mixture was quenched with water (50 mL) and1N HCl (7 ml) and then stirred for 10 min. The mixture was extracted with ethyl acetate (100 mL). The organic layer was separated from the aqueous phase and then concentrated under reduced pressure. The residue was triturated with CH_2Cl_2, concentrated, and then triturated again with methyl *tert*-butyl ether. The solid was washed with methyl *tert*-butyl ether and dried under reduced pressure to provide 4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-imidazole (588 mg, 34%) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ1.23 (s, 9H), 2.16 (s, 6H), 2.22 (s, 3H), 5.76 (s, 2H), 6.27 (s, 1H), 6.91 (s, 2H).

The following compound can be synthesized using the procedures described herein: *N*-(3-aminophenyl)sulfonyl-4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]imidazole-2-carboxamide (Compound 255).

PREPARATION 32: 4,4-dimethyl-2-[(2,4,6-trimethylphenyl)methyl]-1,3-dihydroisoquinoline-1-carboxylic acid


Step 1: methyl (2-methyl-2-phenylpropyl)carbamate



Methyl carbonochloridate (518 µL, 6.70 mmol) was added to a solution of 2-methyl-2-phenyl-propan-1-amine (1.00 g, 6.70 mmol) and pyridine (813 µL, 10.1 mmol) in CH_2Cl_2 (25 mL) at 0 °C. The mixture was allowed to stir at room temperature for 1 h before it was quenched with 1N HCl. The mixture was washed with 1N HCl (2 x 25 mL), brine, dried over sodium


sulfate, filtered and concentrated to give methyl (2-methyl-2-phenylpropyl)carbamate (911 mg, 66%). ESI-MS *m/z* calc. 207.1, found 208.4 $(M+1)^+$; Retention time: 0.54 min (1 min run).

Step 2: 2-methoxycarbonyl-4,4-dimethyl-1,3-dihydroisoquinoline-1-carboxylic acid




Methyl (2-methyl-2-phenylpropyl)carbamate (911 mg, 4.40 mmol) was suspended in acetic acid (15 mL) and H_2SO_4 (5 mL). The mixture was cooled to 0 °C before oxaldehydic acid hydrate (679 mg, 7.37 mmol) was added. The reaction mixture was allowed to warm to room temperature over 1H and was stirred at room temperature for an additional 2 h. The reaction mixture was poured over ice which and the mixture was extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with brine, dried over sodium sulfate, filtered and concentrated to give 2-(methoxycarbonyl)-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline-1-carboxylic acid (1.1 g, 79%) as a white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.55 (s, 1H), 7.45 - 7.17 (m, 3H), 5.66 (d, J = 35.8 Hz, 1H), 3.90 (d, J = 94.9 Hz, 4H), 3.33 (s, 1H), 1.28 (d, J = 51.5 Hz, 6H). ESI-MS *m/z* calc. 263.1, found 264.0 $(M+1)^+$; Retention time: 1.27 min (3 min run).

Step 3: 4,4-dimethyl-2,3-dihydro-1*H*-isoquinoline-1-carboxylic acid



Iodo(trimethyl)silane (254 µL, 1.78 mmol) was added to 2-methoxycarbonyl-4,4-dimethyl-1,3-dihydroisoquinoline-1-carboxylic acid (412 mg, 1.49 mmol) in chloroform (10 mL). The mixture was heated at 65 °C for 2 h. Additional iodo(trimethyl)silane (254 µL, 1.78 mmol) was added and the reaction mixture was heated at 65 °C for 2 h. The reaction mixture was concentrated and the residue was subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give 4,4-dimethyl-2,3-dihydro-1*H*-isoquinoline-1-carboxylic acid (261 mg, 85%). ESI-MS *m/z* calc. 205.1, found 206.4 $(M+1)^+$; Retention time: 0.27 min (1 min run).


Step 4: 4,4-dimethyl-2-[(2,4,6-trimethylphenyl)methyl]-1,3-dihydroisoquinoline-1-carboxylic acid



A suspension of 4,4-dimethyl-2,3-dihydro-1*H*-isoquinoline-1-carboxylic acid (260 mg, 1.27 mmol) and KOH (213 mg, 3.80 mmol) in isopropanol (1.6 mL) was stirred at room temperature for 20 min. 2-(Chloromethyl)-1,3,5-trimethylbenzene (235 mg, 1.39 mmol) was added and the reaction mixture was stirred for 3 h. The mixture was concentrated and the residue was subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give 4,4-dimethyl-2-(2,4,6-trimethylbenzyl)-1,2,3,4-tetrahydroisoquinoline-1-carboxylic acid. ESI-MS *m/z* calc. 337.2, found 338.0 $(M+1)^+$; Retention time: 1.42 min (3 min run).

The following compound can be synthesized using the procedures described herein: *N*-[(6-amino-2-pyridyl)sulfonyl]-4,4-dimethyl-2-[(2,4,6-trimethylphenyl)methyl]-1,3-dihydroisoquinoline-1-carboxamide (Compound 140).

PREPARATION 33: 4-*tert*-butyl-2-(2,4,6-trimethylphenoxy)benzoic acid



A mixture of 2-bromo-4-*tert*-butyl-benzoic acid (500 mg, 1.95 mmol), 2,4,6-trimethylphenol (397 mg, 2.92 mmol), Cs_2CO_3 (1.27 g, 3.89 mmol), $Cu(OTf)_2$ (49 mg, 0.097 mmol) and toluene (10 mL) was combined in a microwave vial before it was sealed and heated at 125 °C for 4 h. The mixture was cooled to room temperature and was partitioned between ethyl acetate (50 mL) and water (50 mL). The layers were separated and the aqueous layer was made

acid by the addition of 1N HCl. The aqueous layer was extracted with ethyl acetate (2 x 30 mL). The combined organics were dried over magnesium sulfate, filtered and concentrated. The residue was subjected to silica gel column chromatography (0-3% methanol/CH_2Cl_2) to give 4-*tert*-butyl-2-(2,4,6-trimethylphenoxy)benzoic acid (250 mg, 39%) as a white solid. ESI-MS *m/z* calc. 312.2, found 313.4 $(M+1)^+$; Retention time: 2.14 min (3 min run). ^{1}H NMR (400 MHz, DMSO) δ 12.66 (s, 1H), 7.73 (d, *J* = 8.2 Hz, 1H), 7.07 (dd, *J* = 8.2, 1.8 Hz, 1H), 6.99 (s, 2H), 6.35 (d, *J* = 1.7 Hz, 1H), 2.28 (s, 3H), 2.01 (s, 6H), 1.10 (s, 9H).

The following compounds can be synthesized using the procedures described herein: *N*-(3-aminophenyl)sulfonyl-4-*tert*-butyl-2-(2,4,6-trimethylphenoxy)benzamide (Compound 224) and *N*-[(6-amino-2-pyridyl)sulfonyl]-4-*tert*-butyl-2-(2,4,6-trimethylphenoxy)benzamide (Compound 102).


PREPARATION 34: 4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrrole-2-carboxylic acid

Step 1: ethyl 5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylate




To a solution of gallium trichloride (4.00 g, 22.7 mmol) in carbon disulfide (30 mL) at room temperature were added ethyl pyrrole-2-carboxylate (2.87 g, 20.6 mmol) and 2-chloro-2-methylpropane (2.50 mL, 23.0 mmol). The mixture was then heated at reflux for 45 min. The reaction mixture was cooled to room temperature and was poured onto a mixture of ice and 1N HCl. The aqueous layer was extracted with dichloromethane. The organic layer was washed with saturated $NaHCO_3$, dried over Na_2SO_4, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (0-10% ethyl acetate/heptanes) to afford 4-*tert*-butyl-1*H*-pyrrole-2-carboxylate (260 mg, 7%), ethyl 5-*tert*-butyl-1*H*-pyrrole-2-carboxylate (2.1 g, 52%) and ethyl 3,5-di-*tert*-butyl-1*H*-pyrrole-2-carboxylate (352 mg, 7%) each isolated as white solids. Ethyl 4-*tert*-butyl-1*H*-pyrrole-2-carboxylate: ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.25 (s, 9H), 1.35 (t, *J* = 7.1 Hz, 3H), 4.30 (q, *J* = 7.1 Hz, 2H), 6.76 (dd, *J* = 2.7, 1.8 Hz, 1H), 6.85 (dd, *J* = 2.3, 1.9 Hz, 1H), 8.83 (br. s, 1H).

Step 2: ethyl 4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrrole-2-carboxylate



Sodium hydride (24 mg as a 60% dispersion in mineral oil, 0.60 mmol) was added to a solution of ethyl 4-*tert*-butyl-1*H*-pyrrole-2-carboxylate (0.10g, 0.51 mmol) in *N,N*-dimethylformamide (3 mL) at 0 °C. The mixture was allowed to stir at 0 °C for 5 min before it was stirred at room temperature for 30 min. 2,4,6-Trimethylbenzyl bromide (0.12 g, 0.55 mmol) was added portion-wise and the mixture was stirred at room temperature for 20 h. The reaction mixture was quenched at 0 °C with aqueous saturated NH_4Cl (4 mL). The mixture was extracted with ethyl acetate (80 mL) and the organic layer was washed with brine (2 x 30 mL) and water (30 mL), dried over Na_2SO_4, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (0-15% ethyl acetate/heptanes) to afford 4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrrole-2-carboxylate (161 mg, 98%) as a white solid. ^1H NMR (300 MHz, $CDCl_3$) δ 1.12 (s, 9H), 1.38 (t, J = 7.1 Hz, 3H), 2.20 (s, 6H), 2.31 (s, 3H), 4.31 (q, J = 7.1 Hz, 2H), 5.47 (s, 2H), 6.11 (d, J = 2.1 Hz, 1H), 6.91 (s, 2H), 6.92 (d, J = 2.1, 1H).

Step 3: 4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrrole-2-carboxylic acid



To a solution of ethyl 4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrrole-2-carboxylate (97 mg, 0.29 mmol) in methanol (6.3 mL) at room temperature was added a 1.0 N solution of NaOH (5.5 mL) and the mixture was heated at reflux for 4 hours. The mixture was

cooled at room temperature and diluted with water (40 mL). The pH was adjusted to 4-5 using 1.0 N HCl and the aqueous layer was extracted with ethyl acetate (120 mL). The organic layer was washed with water (2 x 40 mL), dried over Na_2SO_4, filtered and concentrated under reduced pressure. The solid was triturated with diethyl ether to provide 4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrrole-2-carboxylic acid (74 mg, 84%) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.12 (s, 9H), 2.21 (s, 6H), 2.32 (s, 3H), 5.47 (s, 2H), 6.18 (d, J = 2.0 Hz, 1H), 6.91 (s, 2H), 7.06 (d, J = 2.0 Hz, 1H).

The following compounds can be synthesized using the procedures described herein: *N*-(benzenesulfonyl)-4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrole-2-carboxamide (Compound 242) and *N*-(3-aminophenyl)sulfonyl-4-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrole-2-carboxamide (Compound 195).

PREPARATION 35: 2-(*tert*-butyl)-5-(2,4,6-trimethylbenzyl)oxazole-4-carboxylic acid

Step 1: ethyl 2-hydroxy-2-pivalamidoacetate




Acetic acid (2.61 g, 44.5 mmol) was added to a suspension of trimethylacetamide (9.00 g, 89.1 mmol) and ethyl glyoxylate (50% in toluene, 27.6 g, 133 mmol) in THF (60 mL). The mixture was heated at reflux for 16 h before it was cooled and concentrated to dryness. The residue was treated with heptanes and diethyl ether (10:1 ratio, 110 mL) to give a precipitate which was removed by filtration. The filtrate was concentrated to afford ethyl 2-hydroxy-2-pivalamidoacetate (16.8 g) as a colorless oil. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.22 (s, 9H), 2.27 - 1.35 (m, 3H), 3.63 - 3.75 (m, 1H), 4.20 - 4.36 (m, 2H), 5.50 - 5.60 (m, 1H), 6.65 (br s, 1H). LC-MS analysis: $(M+1)^+$ = 204.1.

Step 2: ethyl 2-chloro-2-pivalamidoacetate




Thionyl chloride (28.6 g, 242 mmol) was added to a solution of ethyl 2-hydroxy-2-pivalamidoacetate (18.6 g, 80.8 mmol) and *N,N*-dimethylformamide (1 mL) in CH_2Cl_2 (100 mL) at 0 °C. The mixture was stirred at room temperature for 16 h before it was concentrated to dryness to afford ethyl 2-chloro-2-pivalamidoacetate (20 g, 64%) as a yellow oil. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.27 (s, 9H), 1.29 - 1.37 (m, 3H), 4.20 - 4.36 (m, 2H), 6.25 (d, J = 9.8, 1H), 6.85 (br s, 1H).

Step 3: ethyl 5-amino-2-(*tert*-butyl)oxazole-4-carboxylate



Diethylaluminum cyanide (1N in toluene, 76.6 mL, 76.7 mmol) was added to a solution of ethyl 2-chloro-2-(2,2-dimethylpropanamido)acetate (11.3 g, 2.71 mmol) in THF (110 mL) at 0 °C. The mixture was stirred at room temperature for 16 h before sat. aqueous NH_4Cl (4 mL) was added to the reaction mixture. The mixture was filtered through Celite and the filtrate was extracted with ethyl acetate (200 mL). The organic layer was washed with brine, dried over Na_2SO_4, filtrated and concentrated. The residue was purified by silica gel column chromatography (50% ethyl acetate/heptane) to afford ethyl 5-amino-2-*tert*-butyl-1,3-oxazole-4-carboxylate (3.21 g, 30%) as a yellow solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.32 (s, 9H), 1.33 - 1.38 (m, 3H), 4.28 - 4.38 (m, 2H), 5.42 (br s, 2H). LC-MS analysis: $[M+H]^+$ = 213.1.


Step 4: ethyl 5-bromo-2-(*tert*-butyl)oxazole-4-carboxylate



A mixture of ethyl 5-amino-2-*tert*-butyl-1,3-oxazole-4-carboxylate (1.50 g, 7.06 mmol) and $CuBr_2$ (1.59 g, 7.12 mmol) in acetonitrile (25 mL) was stirred at room temperature for 1 h followed by addition of *tert*-butyl nitrite (0.94 g, 9.2 mmol). The mixture was stirred at room temperature for 1 h, and then heated at 65 °C for 2.5 h. Brine was added to the reaction mixture and it was extracted with diethyl ether (200 mL). The organic layer was washed by brine spiked with a few drops of 1N HCl, dried over Na_2SO_4, filtrated and concentrated. The residue


was purified by silica gel column chromatography (15% ethyl acetate/hexanes) to afford ethyl 5-bromo-2-*tert*-butyl-1,3-oxazole-4-carboxylate (1.31 g, 56%) as a yellow oil. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.19 - 1.23 (m, 3H), 1.25 (s, 9H), 4.38 (q, J = 6.8 Hz, 2H). LC-MS analysis: $(M+1)^+$ = 276.0.

Step 5: ethyl 2-(*tert*-butyl)-5-(2,4,6-trimethylbenzyl)oxazole-4-carboxylate



A mixture of ethyl 5-bromo-2-*tert*-butyl-1,3-oxazole-4-carboxylate (1.15 g, 4.18 mmol), $Pd(OAc)_2$ (0.09 g, 0.4 mmol) and ^{t}BuP-HBF_4 (0.24 g, 0.83 mmol) in THF (10 mL) was evacuated and back-filled with N_2 three times. The mixture was cooled to 0 °C and (2,4,6-trimethylbenzyl)zinc(II) bromide (1M in THF, 8.36 mL, 8.36 mmol) was added slowly. The mixture was allowed to warm to room temperature and it was stirred for 16 h. The mixture was concentrated and the residue was subjected to silica gel column chromatography (15% ethyl acetate/heptanes) to afford ethyl 2-*tert*-butyl-5-[(2,4,6-trimethylphenyl)methyl]-1,3-oxazole-4-carboxylate (0.48 g, 31%) as yellow oil. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.27 (s, 9H), 1.41 (t, J = 6.9 Hz, 3H), 2.25 (s, 3H), 2.31 (s, 6H), 4.35 (s, 2H), 4.42 (q, J = 7.2 Hz, 2H), 6.85 (s, 2H). LC-MS analysis: $(M+1)^+$ = 300.2.

Step 6: 2-(*tert*-butyl)-5-(2,4,6-trimethylbenzyl)oxazole-4-carboxylic acid




A mixture of ethyl 2-*tert*-butyl-5-[(2,4,6-trimethylphenyl)methyl]-1,3-oxazole-4-carboxylate (0.48 g, 1.5 mmol) and $LiOH.H_2O$ (0.30 g, 7.3 mmol) in THF (4 mL) and water (4

mL) was stirred at room temperature for 16 h. The mixture was diluted with ethyl acetate (150 mL) the the pH was adjusted ~4 using 5% citric acid. The organic layer was washed with brine, dried over Na_2SO_4, filtrated and concentrated. The residue was triturated with diethyl ether and hexanes (8:1, 9 mL) to give 2-*tert*-butyl-5-[(2,4,6-trimethylphenyl)methyl]-1,3-oxazole-4-carboxylic acid (0.33 g, 75%) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.28 (s, 9H), 2.25 (s, 3H), 2.33 (s, 6H), 4.37 (s, 2H), 6.85 (s, 2H). LC-MS analysis: $(M+1)^+ = 302.2$.

The following compound can be synthesized using the procedures described herein: *N*-(3-aminophenyl)sulfonyl-2-*tert*-butyl-5-[(2,4,6-trimethylphenyl)methyl]oxazole-4-carboxamide (Compound 360).


PREPARATION 36: lithium 3-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-1,2,4-triazole-5-carboxylate

Step 1: 2,2-dimethyl-*N*'-[(1*E*)-2,4,6-trimethylphenyl)methylidene]-propanehydrazide




To a solution of 2,4,6-trimethylbenzaldehyde (2.96 g, 20.0 mmol) in 1,2-dichloroethane (16 mL) at room temperature was added 2,2-dimethylpropionic acid hydrazide (2.32 g, 20.0 mmol) and the mixture was stirred for 2 h at room temperature during which time a white precipitate formed. Diethyl ether (30 mL) was added to facilitate precipitation and stirring was continued for 1 h. Filtration over a fritted funnel, followed by washing of the solids with diethyl ether (30 mL) and drying under high vacuum afforded 2,2-dimethyl-*N*'-[(1*E*)-2,4,6-trimethylphenyl)methylidene]propanehydrazide (4.33 g, 88%) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ1.33 (s, 9H), 2.28 (s, 3H), 2.45 (s, 6H), 6.87 (s, 2H), 8.52 (br. s, 1H), 8.61 (br. s, 1H). $(M+1)^+ = 247.2$.

Step 2: 2,2-dimethyl-*N*'-[(2,4,6-trimethylphenyl)methyl]-propanehydrazide




To a solution of 2,2-dimethyl-*N*'-[(1*E*)-2,4,6-trimethylphenyl)methylidene]-propanehydrazide (4.33 g, 17.6 mmol) in methanol (90 mL) was added 10% palladium on carbon (dry basis, 230 mg, 0.21 mmol). The reaction mixture was purged with hydrogen (3 cycles of vacuum/hydrogen) then stirred at room temperature for 2 h under an atmosphere of hydrogen. The reaction mixture was filtered over Celite, washed with methanol and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (0-35% ethyl acetate/heptanes) to afford 2,2-dimethyl-*N*'-[(2,4,6-trimethylphenyl)methyl]propanehydrazide (3.76 g, 86%) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.20 (s, 9H), 2.26 (s, 3H), 2.43 (s, 6H), 3.97 (s, 2H), 4.59 (br. s, 1H), 6.86 (s, 2H), 7.14 (br. s, 1H). $(M+1)^+ = 249.2$.

Step 3: ethyl 3-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-1,2,4-triazole-5-carboxylate



Triethyloxonium tetrafluoroborate (3.42 g, 18.0 mmol) was added portion-wise to a solution of ethyl thiooxamate (2.28 g, 17.1 mmol) in CH_2Cl_2 (35 mL). The resulting mixture was stirred at room temperature for 20 h. The mixture was then cooled to 0 °C, and a solution of 2,2-dimethyl-*N*'-[(2,4,6-trimethylphenyl)methyl]propanehydrazide (4.48 g, 18.0 mmol) and triethylamine (2.5 mL, 18 mmol) in CH_2Cl_2 (20 mL) was added slowly, followed by an additional portion of triethylamine (1.3 mL, 9.0 mmol). The reaction mixture was then heated at reflux for 20 h. The crude mixture was concentrated under reduced pressure and the resulting yellow oil was purified by silica gel column chromatography (0-40% ethyl acetate/heptanes) to afford a solid. Reverse phase chromatographic separation with mixtures of 40% to 95% CH_3CN in water containing 0.1% formic acid afforded ethyl 3-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-1,2,4-triazole-5-carboxylate (620 mg, 11%) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.29 (s, 9H), 1.43 (t, J = 7.1 Hz, 3H), 2.27 (s, 3H), 2.31 (s, 6H), 4.49 (q, J = 7.1 Hz, 2H), 5.71 (s, 2H), 6.86 (s, 2H). $(M+1)^+ = 330.2$.

Step 4: lithium 3-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-1,2,4-triazole-5-carboxylate




To a flask containing ethyl 3-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-1,2,4-triazole-5-carboxylate (110 mg, 0.33 mmol) and LiOH monohydrate (17 mg, 0.40 mmol) were successively added THF (2 mL) and water (1 mL) followed by one drop of methanol. The mixture was stirred at room temperature for 1 h. The volatiles (mostly THF and methanol) were removed under reduced pressure, then the mixture was frozen in a dry-ice / acetone bath and freeze-dried to afford lithium 3-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-1,2,4-triazole-5-carboxylate (98 mg , 95%) as a white solid. ^{1}H NMR (300 MHz, CD_3OD) δ 1.24 (s, 9H), 2.24 (s, 3H), 2.33 (s, 6H), 5.73 (s, 2H), 6.82 (s, 2H).

The following compound can be synthesized using the procedures described herein: *N*-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]-1,2,4-triazole-3-carboxamide (Compound 261).


PREPARATION 37: 5-(*tert*-butyl)-3-(mesityloxy)pyrazine-2-carboxylic acid

Step 1: 3,3-dimethylbutane-1,2-diamine dihydrochloride




A mixture of 2-amino-3,3-dimethyl-butyronitrile hydrochloride (2.94 g, 20.0 mmol), HCl (sat. in EtOH, 10 mL), PtO_2 (600 mg) and methanol (200 mL) at room temperature was treated with hydrogen (50 psi) for 18 h. The reaction mixture was filtered through a pad of Celite and the filtrate was evaporated to give 3,3-dimethylbutane-1,2-diamine dihydrochloride as a white solid (3.9 g, 79%).

Step 2: ethyl 5-(*tert*-butyl)-3-hydroxypyrazine-2-carboxylate




A mixture of 3,3-Dimethyl-butane-1,2-diamine dihydrochloride (3.9 g, 21 mmol), 2-oxo-malonic acid diethyl ester (3.45 mL, 22.7 mmol), Et_3N (5.8 mL, 41 mmol) and EtOH (400 mL) was heated at reflux for 20 h. All volatiles were removed and the residue was subjected to silica gel column chromatography (0-30% ethyl acetate/hexanes) (0/100 to 30/70) to give ethyl 5-(*tert*-butyl)-3-hydroxypyrazine-2-carboxylate (0.59 g, 13%) as a sticky oil, which solidified on standing. ^{1}H NMR (250 MHz, $CDCl_3$) δ 8.36 (s, 1H), 4.55 (q, J = 7.2 Hz, 2H), 1.49 (t, J = 7.2 Hz, 3H), 1.41 (s, 9H).

Step 3: ethyl 5-(*tert*-butyl)-3-chloropyrazine-2-carboxylate



A mixture of ethyl 5-(*tert*-butyl)-3-hydroxypyrazine-2-carboxylate (590. mg, 2.63 mmol) and $POCl_3$ (10 mL) was heated at reflux for 5 h. The excess $POCl_3$ was removed under reduced pressure and the residue was partitioned between ethyl acetate and aq. $NaHCO_3$ (pH adjusted to ~6 by addition of sodium carbonate). The organic phase was separated, dried over Na_2SO_4, filtered and concentrated. The crude material was purified by silica gel column chromatography (0-30% ethyl acetate/hexanes) to give ethyl 5-(*tert*-butyl)-3-chloropyrazine-2-carboxylate (90 mg, 14%). ^{1}H NMR (250 MHz, $CDCl_3$) δ 8.59 (s, 1H), 4.49 (q, J = 7.2 Hz, 2H), 1.44 (t, J = 7.2 Hz, 3H), 1.40 (s, 9H).

Step 4: 5-*tert*-butyl-3-(2,4,6-trimethylphenoxy)pyrazine-2-carboxylic acid




2,4,6-Trimethylphenol (51 mg, 0.38 mmol) and sodium hydride (18 mg, 0.45 mmol) were combined in anhydrous *N,N*-dimethylformamide (3 mL). The reaction mixture was stirred at room temperature for 15 min under an atmosphere of nitrogen. Ethyl 5-*tert*-butyl-3-chloro-pyrazine-2-carboxylate (90. mg, 0.37 mmol) in tetrahydrofuran (1 mL) was then added and the reaction mixture was heated at 60 °C for 1 h. To this mixture was added water (1 mL) and NaOH (120 mg). The resulted mixture was stirred at 70 °C for 1 h and was cooled to room temperature. The mixture was treated with HCl (1 M) to pH 3 and the mixture was extracted with ethyl acetate (3x). The combined extracts were concentrated to about 1 mL (*N,N*-dimethylformamide present). The residue was subjected to preparatory-HPLC (10-99% acetonitrile/water with 0.05 mM HCl) to give 5-*tert*-butyl-3-(2,4,6-trimethylphenoxy)pyrazine-2-carboxylic acid (55 mg, 47%). ^{1}HNMR (250 MHz, DMSO) δ 8.41 (s, 1H), 6.94 (s, 2H), 2.26 (s, 3H), 1.97 (s, 6H), 1.11 (s, 9H).

The following compound can be synthesized using the procedures described herein: *N*-(3-aminophenyl)sulfonyl-5-*tert*-butyl-3-(2,4,6-trimethylphenoxy)pyrazine-2-carboxamide (Compound 141).

PREPARATION 38: 5-(*tert*-butyl)-3-(mesityloxy)pyrazine-2-carboxylic acid


Step 1: ethyl 6-(*tert*-butyl)-4-chloronicotinate



A solution of ethyl 6-*tert*-butyl-1,4-dihydro-4-oxopyridine-3-carboxylate (1.0 g, 4.5 mmol) and phosphorus oxychloride (10 mL) was heated at 110 °C for 1.5 h. The reaction solution was cooled to room temperature and was concentrated under reduced pressure. The


residue was cooled to 0 °C before water (20 mL) and enough saturated aqueous $NaHCO_3$ were carefully added to bring the mixture to pH ~8. The mixture was extracted with ethyl acetate (3x). The combined organics were dried over $MgSO_4$, filtered, and concentrated under vacuum to give ethyl 6-(*tert*-butyl)-4-chloronicotinate (0.80 g, 73%) as a brown oil. ^1H NMR ($CDCl_3$, 250 MHz) δ 8.99 (s, 1H), 7.40 (s, 1H), 4.41 (m, 2H), 1.36 (m, 12H). LC-MS: $(M+1)^+ = 242.2$.

Step 2: 6-*tert*-butyl-4-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate



To a solution of 2,4,6-trimethylphenol (0.43 g, 3.1 mmol) in *N,N*-dimethylformamide (4 mL) and THF (1 mL) was added slowly 60% NaH (0.15 g, 3.9 mmol). The mixture was stirred at room temperature for 25 min before ethyl ethyl 6-(*tert*-butyl)-4-chloronicotinate (0.74 g, 3.1 mmol) was added. The reaction mixture was stirred at 60 °C for 1 h before it was cooled to room temperature and quenched by adding water. The mixture was extracted with ethyl acetate (3x). The combined organic extracts were concentrated and the residue was purified by silica gel column chromatography (0-20% ethyl acetate/hexanes) to afford ethyl 6-*tert*-butyl-4-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (0.45 g, 43%) as a light yellow oil. LC-MS: $(M+1)^+ = 341.9$.

Step 3: 6-*tert*-butyl-4-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid




A solution of 6-(*tert*-butyl)-4-(mesityloxy)nicotinate (0.74 g, 2.2 mmol) and sodium hydroxide (0.21 g, 5.2 mmol) in EtOH (2 mL) and water (1 mL) was stirred at room temperature

for 3 h. The reaction was concentrated and the residue was acidified with concentrated HCl to about pH 3. The crude material was purified by silica gel column chromatography (ethyl acetate) to give 6-*tert*-butyl-4-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (0.38 g, 55%) as a white solid. ^{1}H NMR ($CDCl_3$, 250 MHz) δ 9.22 (s, 1H), 6.96 (s, 2H), 6.24 (s, 1H), 2.34 (s, 3H), 2.09 (s, 6H), 1.22 (s, 9H). LC-MS: $(M+1)^+ = 314.1$.

The following compound can be synthesized using the procedures described herein: *N*-(3-aminophenyl)sulfonyl-6-*tert*-butyl-4-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 338).

PREPARATION 39: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 368) and *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4*R*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1289)


Step 1: *tert*-butyl 2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylate



tert-Butyl 2,6-dichloropyridine-3-carboxylate (15.0 g, 60.5 mmol) and (3-fluoro-5-isobutoxy-phenyl)boronic acid (13.46 g, 63.48 mmol) were combined and fully dissolved in ethanol (150 mL) and toluene (150 mL). A suspsension of sodium carbonate (19.23 g, 181.4 mmol) in water (30 mL) was added. Tetrakis(triphenylphosphine)palladium (0) (2.096 g, 1.814 mmol) was added under nitrogen. The reaction mixture was allowed to stir at 60 °C for 16 hours. Volatiles were removed under reduced pressure. The remaining solids were partitioned between water (100 mL) and ethyl acetate (100 mL). The organic layer was washed with brine (1× 100 mL), dried over sodium sulfate, filtered and concentrated under reduced pressure. The material was subjected silica gel column chromatography on a 330 gram silica gel column, 0 to 20% ethyl acetate in hexanes gradient. The material was repurified on a 220 gram silica gel column, isocratic 100% hexane for 10 minutes, then a 0 to 5% ethyl acetate in hexanes gradient to yield *tert*-butyl 2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylate (18.87 g, 49.68


mmol, 82.2%) was obtained as a colorless oil. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.24 (d, J = 8.0 Hz, 1H), 8.16 (d, J = 8.1 Hz, 1H), 7.48 (dd, J = 9.4, 2.0 Hz, 2H), 6.99 (dt, J = 10.8, 2.2 Hz, 1H), 3.86 (d, J = 6.5 Hz, 2H), 2.05 (dt, J = 13.3, 6.6 Hz, 1H), 1.57 (d, J = 9.3 Hz, 9H), 1.00 (t, J = 5.5 Hz, 6H). ESI-MS *m/z* calc. 379.13504, found 380.2 $(M+1)^+$; Retention time: 2.57 minutes.

Step 2: 2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylic acid




tert-Butyl 2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylate (18.57 g, 48.89 mmol) was dissolved in dichloromethane (200 mL). Trifluoroacetic acid (60 mL, 780 mmol) was added and the reaction mixture was allowed to stir at room temperature for 1 hour. The reaction mixture was stirred at 40 °C for 2 hours. The reaction mixture was concentrated under reduced pressure and taken up in ethyl acetate (100 mL). It was washed with a saturated aqueous sodium bicarbonate solution (1× 100 mL) and brine (1× 100 mL), dried over sodium sulfate, filtered and concentrated under reduced pressure. The crude product was suspended in ethyl acetate (75 mL) and washed with aqueous HCl (1 N, 1× 75 mL). The organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure. The remaining solid (17.7 g) was stirred as a slurry in dichloromethane (35 mL) at 40 °C for 30 minutes. After cooling to room temperature, the remaining slurry was filtered, and then rinsed with cold dichloromethane to give 2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylic acid (11.35 g, 35.06 mmol, 72%) as a white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 13.76 (s, 1H), 8.31 (d, J = 8.0 Hz, 1H), 8.17 (d, J = 8.1 Hz, 1H), 7.54 - 7.47 (m, 2H), 7.00 (dt, J = 10.8, 2.3 Hz, 1H), 3.87 (d, J = 6.5 Hz, 2H), 2.05 (dt, J = 13.3, 6.6 Hz, 1H), 1.01 (d, J = 6.7 Hz, 6H). ESI-MS *m/z* calc. 323.1, found 324.1 $(M+1)^+$; Retention time: 1.96 minutes.

Step 3: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide



2-Chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylic acid (3.00 g, 9.27 mmol) was dissolved in *N,N*-dimethylformamide (30.00 mL), and 1,1'-carbonyldiimidazole (2.254 g, 13.90 mmol) was added to the solution. The solution was allowed to stir at 65 °C for 1 hour. In a separate flask, sodium hydride (444.8 mg, 11.12 mmol) was added to a solution of 6-aminopyridine-2-sulfonamide (1.926 g, 11.12 mmol) in *N,N*-dimethylformamide (15.00 mL). This mixture was stirred for one hour before being added to the prior reaction mixture. The final reaction mixture was stirred at 65 °C for 15 minutes. Volatiles were removed under reduced pressure. The remaining oil was taken up in ethyl acetate and washed with aqueous HCl (1 N, 1× 75 mL) and brine (3× 75 mL). The organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure. The remaining white solid (4.7 g) was fully dissolved in isopropanol (120 mL) in an 85 °C water bath. The colorless solution was allowed to slowly cool to room temperature with slow stirring over 16 hours. The crystalline solids that had formed were collected by vacuum filtration, and then rinsed with cold isopropanol (50 mL). Upon drying, *N*-[(6-amino-2-pyridyl)sulfonyl]-2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (3.24 g, 6.765 mmol, 73%) was obtained as a white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.78 (s, 1H), 8.15 (d, J = 8.0 Hz, 1H), 8.09 (d, J = 7.9 Hz, 1H), 7.73 - 7.63 (m, 1H), 7.49 (dd, J = 8.6, 1.9 Hz, 2H), 7.21 (d, J = 7.3 Hz, 1H), 6.99 (dt, J = 10.7, 2.2 Hz, 1H), 6.74 (d, J = 8.4 Hz, 1H), 6.64 (s, 2H), 3.86 (d, J = 6.5 Hz, 2H), 2.05 (dp, J = 13.3, 6.5 Hz, 1H), 1.02 (dd, J = 12.7, 6.4 Hz, 6H).

Step 4: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4*R*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1289) and *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide



N-[(6-Amino-2-pyridyl)sulfonyl]-2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (309 mg, 0.645 mmol) was dissolved in dimethylsulfoxide (3.708 mL) and potassium carbonate (445.9 mg, 3.226 mmol) was slowly added, followed by 2,2,4-trimethylpyrrolidine (146.0 mg, 1.290 mmol). The reaction mixture was sealed and heated at 150 °C for 72 hours. The reaction was cooled down, diluted with water (50 mL), extracted 3 times with 50 mL portions of ethyl acetate, washed with brine, dried over sodium sulfate, filtered and evaporated to dryness. The crude material was dissolved in 2 mL of dichloromethane and purified by on silica gel using a gradient of 0 to 80% ethyl acetate in hexanes. The stereoisomers were separated using supercritical fluid chromatography on a ChiralPak AD-H (250 x 4.6 mm), 5μm column using 25% isopropanol with 1.0% diehtylamine in CO_2 at a flow rate of 3.0 mL/min. The separated enationmers were separately concentrated, diluted with ethyl acetate (3 mL) and washed with 1N aqueous hydrochloric acid. The organic layers were dried over sodium sulfate, filtered, and evaporated to dryness to give the pure compounds as pale yellow solids.

The first compound to eluet from the SFC conditions given above gave *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4*R*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1289) (Hydrochloric Acid (1)) ^{1}H NMR (400 MHz, DMSO-d6) δ 12.47 (s, 1H), 7.78 (d, *J* = 8.0 Hz, 1H), 7.69 - 7.57 (m, 1H), 7.56 - 7.46 (m, 1H), 7.41 (dt, *J* = 10.1, 1.8 Hz, 1H), 7.26 (d, *J* = 8.0 Hz, 1H), 7.21 (d, *J* = 7.2 Hz, 1H), 6.89 (dt, *J* = 10.7, 2.3 Hz, 1H), 6.69 (d, *J* = 8.3 Hz, 1H), 3.83 (d, *J* = 6.7 Hz, 2H), 2.61 (dq, *J* = 9.7, 4.9 Hz, 2H), 2.24 (d, *J* = 15.8 Hz, 1H), 2.06 (dq, *J* = 13.3, 6.7 Hz, 1H), 1.93 - 1.82 (m, 1H), 1.61 (s, 3H), 1.59 (s, 3H), 1.48 - 1.33 (m, 1H), 1.32 - 1.20 (m, 2H), 0.99 (d, *J* = 6.6 Hz, 6H), 0.88 (d, *J* = 6.2 Hz, 3H). ESI-MS *m/z* calc. 555.2, found 556.4 $(M+1)^+$; Retention time: 2.76 minutes.

The second compound to eluet from the SFC conditions described above gave *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Hydrochloric Acid (1)) ^{1}H NMR (400 MHz, Chloroform-d) δ 15.49 (s, 1H), 8.49 (d, *J* = 8.2 Hz, 1H), 7.75 - 7.56 (m, 3H), 7.34 (t, *J* = 1.8 Hz, 1H), 7.30 (dt, *J* = 9.4, 1.9 Hz, 1H), 6.75 - 6.66 (m, 2H), 3.95 (s, 1H), 3.78 (d, *J* = 6.5 Hz, 2H), 3.42 (s, 1H), 2.88 - 2.74

(m, 1H), 2.23 (dd, J = 12.5, 8.0 Hz, 1H), 2.17 - 2.08 (m, 1H), 1.98 - 1.87 (m, 1H), 1.55 (s, 3H), 1.39 (s, 3H), 1.31 (d, J = 6.7 Hz, 3H), 1.05 (d, J = 6.7 Hz, 6H). ESI-MS m/z calc. 555.2, found 556.4 $(M+1)^+$; Retention time: 2.77 minutes. Absolute stereochemistry was confirmed by X-ray crystallography.

The following compound can be synthesized using the procedures described herein: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-isopropyl-1-piperidyl)pyridine-3-carboxamide (Compound 114),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 946),

N-[(6-amino-4-methyl-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1472),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-phenylbutoxy)pyridine-3-carboxamide (Compound 1686),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[1-(1-naphthyl)ethylamino]pyridine-3-carboxamide (Compound 1769),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[2-(o-tolyl)azepan-1-yl]pyridine-3-carboxamide (Compound 1772),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[3-(cyclopropylmethyl)-2,2-dimethyl-pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1794),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-benzyl-2,2-dimethyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2275),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[1-[2-(trifluoromethyl)phenyl]ethoxy]pyridine-3-carboxamide (Compound 2501),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[2-(1,3-benzoxazol-2-yl)phenoxy]pyridine-3-carboxamide (Compound 2544),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2-cyclohexylcyclohexyl)amino]pyridine-3-carboxamide (Compound 2599),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-phenyl-1-piperidyl)pyridine-3-carboxamide (Compound 2615),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(6-methoxy-4-methyl-2-pyridyl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1482),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-tetrahydrofuran-3-ylsulfonyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1640),

N-cyclopropylsulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1525),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)-4-methoxy-pyridine-3-carboxamide (Compound 1254),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 171),

N-[(6-Amino-2-pyridyl)sulfonyl]-2-[(2*S*,5*R*)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(3R)-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 233),

2-(2,5-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 307),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 321),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-[(2,4-dimethyl-3-pyridyl)oxy]pyridine-3-carboxamide (Compound 460),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 353),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 355),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(2R)-2-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1354),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3,3,4-trimethyl-1-piperidyl)pyridine-3-carboxamide (Compound 435),

6-(4-ethoxyphenyl)-2-(4-methyl-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 445),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,3R)-2,3-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 441),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[4-(4-pyridyl)butoxy]pyridine-3-carboxamide (Compound 1687),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-4-oxo-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-1,2-dihydropyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-3-oxo-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-1,2-dihydropyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[4-(3-pyridyl)butoxy]pyridine-3-carboxamide (Compound 1690),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[4-(2-pyridyl)butoxy]pyridine-3-carboxamide (Compound 1688),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-pyrazol-1-ylpropoxy)pyridine-3-carboxamide (Compound 1692),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-phenylbutoxy)pyridine-3-carboxamide (Compound 1689),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[3-(2-pyridyl)propoxy]pyridine-3-carboxamide (Compound 1693),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[3-(3-pyridyl)propoxy]pyridine-3-carboxamide (Compound 1691),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-phenylpropoxy)pyridine-3-carboxamide (Compound 1694),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2,2,3-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 464),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1R,4S)-3-azabicyclo[2.2.1]heptan-3-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 476),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[3-(2-pyridyl)-1-piperidyl]pyridine-3-carboxamide (Compound 489),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(trifluoromethyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1556),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2,4,4-tetramethylpyrrolidin-1-yl)-6-(trifluoromethyl)pyridine-3-carboxamide (Compound 1635),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(trifluoromethyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1550),

N-[(6-amino-2-pyridyl)sulfonyl]-6-methyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1420),

N-[(6-amino-2-pyridyl)sulfonyl]-6-methoxy-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1536),

N-[(6-amino-2-pyridyl)sulfonyl]-5-fluoro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1607),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]-6,7-dihydro-5H-cyclopenta[b]pyridine-3-carboxamide (Compound 1386),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropylmethyl(propyl)amino]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 500),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2-methyl-4-phenyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 502),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2-phenylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 509),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3,3,5-trimethyl-1-piperidyl)pyridine-3-carboxamide (Compound 520),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-azabicyclo[2.2.1]heptan-3-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 521),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-propyl-1-piperidyl)pyridine-3-carboxamide (Compound 522),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R,4S)-4-fluoro-2-methyl-pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1362)

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(3S)-2,2,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 557),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[3-(3,4-difluorophenyl)-1-piperidyl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 558),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(1-piperidyl)pyridine-3-carboxamide (Compound 567),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[2-(methoxymethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 574),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6-azaspiro[2.5]octan-6-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 578),

N-[(6-amino-2-pyridyl)sulfonyl]-2-indolin-1-yl-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 588),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2,4,4-tetramethylpyrrolidin-1-yl)-5-(trifluoromethyl)pyridine-3-carboxamide (Compound 1638),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2,4,4-tetramethylpyrrolidin-1-yl)-4-(trifluoromethyl)pyridine-3-carboxamide (Compound 1658),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6,6-dimethyl-3-azabicyclo[3.1.0]hexan-3-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 596),

N-[(6-amino-5-hydroxy-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1563),

N-[(6-amino-5-hydroxy-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1375),

N-[(6-amino-5-benzyloxy-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1588),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide,

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1593),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4R)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1444),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-*tert*-butylpyrrolidin-1-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 597),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,5-dimethyl-3-phenyl-pyrrolidin-1-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 601),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[4-(methoxymethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 606),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-ethyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 619),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2-methyl-3-phenyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 625),

N-[(6-amino-5-benzyloxy-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1429),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-benzylpyrrolidin-1-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 647),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[3-(2,2,2-trifluoroethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 656),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-benzyl-2-methyl-pyrrolidin-1-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 658),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2-methyl-2,3,3a,4,5,6,7,7a-octahydroindol-1-yl)pyridine-3-carboxamide (Compound 675),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-azabicyclo[3.1.1]heptan-3-yl)-6-*tert*-butyl-pyridine-3-carboxamide (Compound 1401),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(6-methoxy-2-pyridyl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1483),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[2-(3-methyldiazirin-3-yl)ethyl-prop-2-ynyl-amino]pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5S)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 687),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylpyrrolidin-1-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 689),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(3-methoxyphenyl)sulfonyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1452),

N-(benzenesulfonyl)-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4R)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1396),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azaspiro[2.5]octan-7-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 703),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2-isopropylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 724),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-oxa-7-azaspiro[3.5]nonan-7-yl)pyridine-3-carboxamide (Compound 729),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2-phenyl-1-piperidyl)pyridine-3-carboxamide (Compound 731),

2-(3,3a,4,5,6,6a-hexahydro-2*H*-cyclopenta[b]pyrrol-1-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 732),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azaspiro[3.5]nonan-7-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 733),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 736),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropylmethyl(methyl)amino]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 740),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(4-phenyl-1-piperidyl)pyridine-3-carboxamide (Compound 742),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-fluoro-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 745),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(3R)-3-methyl-1-piperidyl]pyridine-3-carboxamide (Compound 757),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,3S)-2,3-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 764),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(1-phenylethylamino)pyridine-3-carboxamide (Compound 766),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(3-methoxyphenyl)sulfonyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1171),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(6-oxo-1*H*-pyridin-2-yl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1308),

N-(3-aminophenyl)sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1184),

N-(benzenesulfonyl)-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 620),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(2-pyridylsulfonyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1063),

N-(3-aminophenyl)sulfonyl-2-(4,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1014),

2-(4,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(3-methoxyphenyl)sulfonyl-pyridine-3-carboxamide (Compound 510),

N-(benzenesulfonyl)-2-(4,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 583),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(6-methoxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2430),

2-(4,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(6-oxo-1*H*-pyridin-2-yl)sulfonyl]pyridine-3-carboxamide (Compound 775),

2-(4,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(2-pyridylsulfonyl)pyridine-3-carboxamide (Compound 1069),

2-(4,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(6-methoxy-2-pyridyl)sulfonyl]pyridine-3-carboxamide (Compound 1090),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 811),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-spiro[1H-isobenzofuran-3,4'-piperidine]-1'-yl-pyridine-3-carboxamide (Compound 818),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethylpyrrolidin-1-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 834),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(3S)-3-methyl-1-piperidyl]pyridine-3-carboxamide (Compound 836),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[3-(cyclopropylmethoxy)-1-piperidyl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 858),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(3S)-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 864),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[4-(cyclopropylmethoxy)-1-piperidyl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 884),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4,4-difluoro-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 889),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azabicyclo[2.2.1]heptan-7-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 903),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-fluoro-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 913),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2,2,4,4-tetramethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 915),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1S,4R)-3-azabicyclo[2.2.1]heptan-3-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 921),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2,3,5-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 925),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-cyclopropylpyrrolidin-1-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 928),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(indan-1-ylamino)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 935),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(azepan-1-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 940),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(4R)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 949),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-ethynyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 954),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-ethoxy-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 960),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,3-diethyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 971),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,4-dimethyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 979),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(8-azaspiro[3.4]octan-8-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 987),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(3R)-2,2,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 992),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-benzylpyrrolidin-1-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 994),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-benzyloxy-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1012),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-ethyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1023),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopentyl(methyl)amino]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1025),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(benzylamino)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1026),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2,2,5-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1040),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[4-(trifluoromethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 1043),

N-[(6-amino-2-pyridyl)sulfonyl]-2-isoindolin-2-yl-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1072),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(4-phenoxy-1-piperidyl)pyridine-3-carboxamide (Compound 1080),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(3S)-3-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1081),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-ethylpyrrolidin-1-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1087),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-phenoxypyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1092),

2-(3,3a,4,5,6,6a-hexahydro-1*H*-cyclopenta[c]pyrrol-2-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1107),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(3-phenylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1112),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[3-(methoxymethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 1116),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-cyclopropyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1132),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4,4-dimethyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1134),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(3S,5S)-3,5-dimethyl-1-piperidyl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1136),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(3R,5S)-3,5-dimethyl-1-piperidyl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1145),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 1182),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1198),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-*tert*-butylpyrrolidin-1-yl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1206),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(3R)-3-phenyl-1-piperidyl]pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-ethoxy-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1214),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(2R)-2-(trifluoromethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1230),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1247),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(2S)-2-(trifluoromethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1248),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2-methylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1271),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[3-(trifluoromethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 1274),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,3-dimethyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1283),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(3R)-3-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1322),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-(2,3,3-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1327),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[3-(3-chlorophenyl)-1-piperidyl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1339),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-ethyl-3-methyl-1-piperidyl)-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1341),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azaspiro[2.4]heptan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1333),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(3R,5S)-3,5-dimethyl-1-piperidyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 371),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(1-piperidyl)pyridine-3-carboxamide (Compound 380),

2-(3,3a,4,5,6,6a-hexahydro-2*H*-cyclopenta[b]pyrrol-1-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 383),

methyl (2S)-1-[3-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-pyridyl]pyrrolidine-2-carboxylate (Compound 385),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2-(trifluoromethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 398),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-3-methyl-1-piperidyl]pyridine-3-carboxamide (Compound 408),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(azepan-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 418),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[3-(methoxymethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 419),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azabicyclo[2.2.1]heptan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 420),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 421),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(8-oxa-4-azaspiro[4.4]nonan-4-yl)pyridine-3-carboxamide (Compound 424),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 425),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2-phenylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 447),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,5-dihydropyrrol-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 453),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5S)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 458),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-benzylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 473),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 479),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methyl-4-phenyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 488),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[4-(trifluoromethyl)-3-azabicyclo[3.1.0]hexan-3-yl]pyridine-3-carboxamide (Compound 490),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[2-(1,1-dimethylpropyl)pyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 493),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,5-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 496),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[1-cyclopropylethyl(methyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 516),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6-azaspiro[3.3]heptan-6-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 527),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-ethyl-3-methyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 530),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-2,2,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 547),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 555),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3,3,5-trimethyl-1-piperidyl)pyridine-3-carboxamide (Compound 560),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3,3,4-trimethyl-1-piperidyl)pyridine-3-carboxamide (Compound 575),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1R,5S)-3-azabicyclo[3.2.1]octan-3-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 581),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 587),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-pyrrolidin-1-yl-pyridine-3-carboxamide (Compound 592),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-cyclopropylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 593),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6,6-dimethyl-3-azabicyclo[3.1.0]hexan-3-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 609),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-*tert*-butylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 611),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-*tert*-butylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 613),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-ethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 615),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 651),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-cyclopropyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 655),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-2,2,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 657),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[2-(2-pyridyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 659),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-benzyl-2-methyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 663),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-3-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 678),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-azabicyclo[2.2.1]heptan-3-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 680),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclobutyl(methyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 683),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 684),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 694),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,5-dimethyl-3-phenyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 711),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2-(methoxymethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 719),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 720),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 735),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 751),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-benzylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 772),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[2,2-dimethyl-3-(trifluoromethyl)pyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 780),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 837),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(8-azaspiro[3.4]octan-8-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 845),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4,4-tetramethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 850),

2-(3,3a,4,5,6,6a-hexahydro-1*H*-cyclopenta[c]pyrrol-2-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 857),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-phenylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 859),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropylmethyl(propyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 881),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,3-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 887),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-azabicyclo[3.1.0]hexan-4-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 906),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1S,4R)-3-azabicyclo[2.2.1]heptan-3-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 924),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(9-azaspiro[4.4]nonan-9-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 926),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-ethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 931),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1R,5S)-8-azabicyclo[3.2.1]octan-8-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 934),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azaspiro[2.5]octan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 953),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-benzyl-2-methyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 969),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6-azaspiro[2.5]octan-6-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 973),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 991),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1004),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[methyl-[(1-methylcyclopropyl)methyl]amino]pyridine-3-carboxamide (Compound 1009),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropylmethyl(methyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1010),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,5-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1013),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-3-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1051),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6-azaspiro[2.4]heptan-6-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1061),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2,3,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1065),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-4-oxo-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-1,2-dihydropyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S)-2-*tert*-butylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1097),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-isopropyl-1-piperidyl)pyridine-3-carboxamide (Compound 1098),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropyl(methyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1117),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,3-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1121),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2,3,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1144),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2-(methoxymethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1149),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,3,3-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1156),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropylmethyl(isopropyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1157),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-methylene-1-piperidyl)pyridine-3-carboxamide (Compound 1162),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(5R)-2,2,5-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1166),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R)-2-*tert*-butylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1170),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1R,4S)-3-azabicyclo[2.2.1]heptan-3-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1180),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-3-oxo-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-1,2-dihydropyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-3-methyl-1-piperidyl]pyridine-3-carboxamide (Compound 1193),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[1-(3-chlorophenyl)ethyl-methyl-amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1205),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(3S,5S)-3,5-dimethyl-1-piperidyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1210),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-ethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1218),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-ethyl-5-methyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1222),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(5S)-2,2,5-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1241),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-isopropylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1258),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2-phenylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1260),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(8-azaspiro[3.5]nonan-8-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1264),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,3-dimethylmorpholin-4-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1267),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1279),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4R)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1289),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azaspiro[3.5]nonan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1303),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2-(trifluoromethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1304),

2-(2,3,4,4a,5,6,7,7a-octahydrocyclopenta[b]pyridin-1-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1313),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methyl-3-phenyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1319) *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azaspiro[2.4]heptan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1333),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(3R,5S)-3,5-dimethyl-1-piperidyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 371),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(1-piperidyl)pyridine-3-carboxamide (Compound 380),

2-(3,3a,4,5,6,6a-hexahydro-2*H*-cyclopenta[b]pyrrol-1-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 383),

methyl (2S)-1-[3-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-pyridyl]pyrrolidine-2-carboxylate (Compound 385),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2-(trifluoromethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 398),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-3-methyl-1-piperidyl]pyridine-3-carboxamide (Compound 408),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(azepan-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 418),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[3-(methoxymethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 419),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azabicyclo[2.2.1]heptan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 420),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 421),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(8-oxa-4-azaspiro[4.4]nonan-4-yl)pyridine-3-carboxamide (Compound 424),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 425),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2-phenylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 447),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,5-dihydropyrrol-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 453),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5S)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 458),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-benzylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 473),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 479),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methyl-4-phenyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 488),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[4-(trifluoromethyl)-3-azabicyclo[3.1.0]hexan-3-yl]pyridine-3-carboxamide (Compound 490),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[2-(1,1-dimethylpropyl)pyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 493),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,5-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 496),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[1-cyclopropylethyl(methyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 516),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6-azaspiro[3.3]heptan-6-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 527),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-ethyl-3-methyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 530),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-2,2,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 547),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 555),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3,3,5-trimethyl-1-piperidyl)pyridine-3-carboxamide (Compound 560),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3,3,4-trimethyl-1-piperidyl)pyridine-3-carboxamide (Compound 575),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1R,5S)-3-azabicyclo[3.2.1]octan-3-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 581),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 587),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-pyrrolidin-1-yl-pyridine-3-carboxamide (Compound 592),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-cyclopropylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 593),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6,6-dimethyl-3-azabicyclo[3.1.0]hexan-3-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 609),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-*tert*-butylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 611),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-*tert*-butylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 613),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-ethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 615),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 651),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-cyclopropyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 655),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-2,2,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 657),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[2-(2-pyridyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 659),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-benzyl-2-methyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 663),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-3-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 678),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-azabicyclo[2.2.1]heptan-3-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 680),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclobutyl(methyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 683),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 684),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 694),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,5-dimethyl-3-phenyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 711),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2-(methoxymethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 719),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 720),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 735),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 751),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-benzylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 772),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[2,2-dimethyl-3-(trifluoromethyl)pyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 780),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 837),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(8-azaspiro[3.4]octan-8-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 845),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4,4-tetramethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 850),

2-(3,3a,4,5,6,6a-hexahydro-1*H*-cyclopenta[c]pyrrol-2-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 857),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-phenylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 859),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropylmethyl(propyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 881),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,3-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 887),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-azabicyclo[3.1.0]hexan-4-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 906),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1S,4R)-3-azabicyclo[2.2.1]heptan-3-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 924),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(9-azaspiro[4.4]nonan-9-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 926),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-ethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 931),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1R,5S)-8-azabicyclo[3.2.1]octan-8-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 934),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azaspiro[2.5]octan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 953),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-benzyl-2-methyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 969),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6-azaspiro[2.5]octan-6-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 973),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 991),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1004),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[methyl-[(1-methylcyclopropyl)methyl]amino]pyridine-3-carboxamide (Compound 1009),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropylmethyl(methyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1010),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,5-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1013),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-3-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1051),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6-azaspiro[2.4]heptan-6-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1061),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2,3,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1065),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-4-oxo-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-1,2-dihydropyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S)-2-*tert*-butylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1097),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-isopropyl-1-piperidyl)pyridine-3-carboxamide (Compound 1098),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropyl(methyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1117),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,3-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1121),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2,3,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1144),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2-(methoxymethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1149),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,3,3-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1156),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropylmethyl(isopropyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1157),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-methylene-1-piperidyl)pyridine-3-carboxamide (Compound 1162),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(5R)-2,2,5-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1166),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R)-2-*tert*-butylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1170),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1R,4S)-3-azabicyclo[2.2.1]heptan-3-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1180),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-3-oxo-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-1,2-dihydropyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-3-methyl-1-piperidyl]pyridine-3-carboxamide (Compound 1193),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[1-(3-chlorophenyl)ethyl-methyl-amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1205),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(3S,5S)-3,5-dimethyl-1-piperidyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1210),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-ethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1218),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-ethyl-5-methyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1222),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(5S)-2,2,5-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1241),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-isopropylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1258),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2-phenylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1260),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(8-azaspiro[3.5]nonan-8-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1264),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,3-dimethylmorpholin-4-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1267),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1279),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4R)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1289),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azaspiro[3.5]nonan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1303),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2-(trifluoromethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1304),

2-(2,3,4,4a,5,6,7,7a-octahydrocyclopenta[b]pyridin-1-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1313),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methyl-3-phenyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1319),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-methyl-1-piperidyl)-6-(p-tolyl)pyridine-3-carboxamide (Compound 802),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-methoxy-5-methyl-3-pyridyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 863),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-ethoxyphenyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 910),

6-(3-fluoro-5-isopropoxy-phenyl)-2-(4-methyl-1-piperidyl)-*N*-(1H-pyrazol-3-ylsulfonyl)pyridine-3-carboxamide (Compound 1042),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azaspiro[2.4]heptan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1333),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(3R,5S)-3,5-dimethyl-1-piperidyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 371),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(1-piperidyl)pyridine-3-carboxamide (Compound 380),

2-(3,3a,4,5,6,6a-hexahydro-2*H*-cyclopenta[b]pyrrol-1-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 383),

methyl (2S)-1-[3-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-pyridyl]pyrrolidine-2-carboxylate (Compound 385),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2-(trifluoromethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 398),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-3-methyl-1-piperidyl]pyridine-3-carboxamide (Compound 408),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(azepan-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 418),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[3-(methoxymethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 419),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azabicyclo[2.2.1]heptan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 420),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 421),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(8-oxa-4-azaspiro[4.4]nonan-4-yl)pyridine-3-carboxamide (Compound 424),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 425),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2-phenylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 447),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,5-dihydropyrrol-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 453),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5S)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 458),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-benzylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 473),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 479),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methyl-4-phenyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 488),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[4-(trifluoromethyl)-3-azabicyclo[3.1.0]hexan-3-yl]pyridine-3-carboxamide (Compound 490),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[2-(1,1-dimethylpropyl)pyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 493),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,5-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 496),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[1-cyclopropylethyl(methyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 516),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6-azaspiro[3.3]heptan-6-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 527),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-ethyl-3-methyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 530),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-2,2,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 547),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 555),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3,3,5-trimethyl-1-piperidyl)pyridine-3-carboxamide (Compound 560),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3,3,4-trimethyl-1-piperidyl)pyridine-3-carboxamide (Compound 575),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1R,5S)-3-azabicyclo[3.2.1]octan-3-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 581),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 587),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-pyrrolidin-1-yl-pyridine-3-carboxamide (Compound 592),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-cyclopropylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 593),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6,6-dimethyl-3-azabicyclo[3.1.0]hexan-3-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 609),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-*tert*-butylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 611),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-*tert*-butylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 613),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-ethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 615),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 651),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-cyclopropyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 655),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-2,2,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 657),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[2-(2-pyridyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 659),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-benzyl-2-methyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 663),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-3-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 678),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-azabicyclo[2.2.1]heptan-3-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 680),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclobutyl(methyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 683),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 684),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 694),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,5-dimethyl-3-phenyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 711),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2-(methoxymethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 719),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 720),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 735),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 751),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-benzylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 772),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[2,2-dimethyl-3-(trifluoromethyl)pyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 780),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,3-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 837),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(8-azaspiro[3.4]octan-8-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 845),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4,4-tetramethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 850),

2-(3,3a,4,5,6,6a-hexahydro-1*H*-cyclopenta[c]pyrrol-2-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 857),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-phenylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 859),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropylmethyl(propyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 881),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,3-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 887),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-azabicyclo[3.1.0]hexan-4-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 906),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1S,4R)-3-azabicyclo[2.2.1]heptan-3-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 924),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(9-azaspiro[4.4]nonan-9-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 926),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-ethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 931),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1R,5S)-8-azabicyclo[3.2.1]octan-8-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 934),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azaspiro[2.5]octan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 953),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-benzyl-2-methyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 969),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6-azaspiro[2.5]octan-6-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 973),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 991),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1004),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[methyl-[(1-methylcyclopropyl)methyl]amino]pyridine-3-carboxamide (Compound 1009),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropylmethyl(methyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1010),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,5-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1013),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-3-methylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1051),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6-azaspiro[2.4]heptan-6-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1061),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2,3,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1065),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-4-oxo-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-1,2-dihydropyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S)-2-*tert*-butylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1097),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-isopropyl-1-piperidyl)pyridine-3-carboxamide (Compound 1098),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropyl(methyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1117),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,3-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1121),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2,3,3-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1144),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2-(methoxymethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1149),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,3,3-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1156),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclopropylmethyl(isopropyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1157),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-methylene-1-piperidyl)pyridine-3-carboxamide (Compound 1162),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(5R)-2,2,5-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1166),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R)-2-*tert*-butylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1170),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(1R,4S)-3-azabicyclo[2.2.1]heptan-3-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1180),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-3-oxo-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-1,2-dihydropyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-3-methyl-1-piperidyl]pyridine-3-carboxamide (Compound 1193),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[1-(3-chlorophenyl)ethyl-methyl-amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1205),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(3S,5S)-3,5-dimethyl-1-piperidyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1210),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-ethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1218),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-ethyl-5-methyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1222),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(5S)-2,2,5-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1241),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-isopropylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1258),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2-phenylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1260),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(8-azaspiro[3.5]nonan-8-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1264),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,3-dimethylmorpholin-4-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1267),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1279),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4R)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1289),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(7-azaspiro[3.5]nonan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1303),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2-(trifluoromethyl)pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1304),

2-(2,3,4,4a,5,6,7,7a-octahydrocyclopenta[b]pyridin-1-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1313),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methyl-3-phenyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1319),

2-(2,5-dimethyl-3-phenyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1348),

2-[(3,3-difluorocyclobutyl)methylamino]-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 393),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 415),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)-2-[4-(trifluoromethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 426),

2-(3-ethynyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 431),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methyl-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 442),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-isopropyl-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 498),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1*H*-pyrazol-5-ylsulfonyl)-2-spiro[1*H*-isobenzofuran-3,4'-piperidine]-1'-yl-pyridine-3-carboxamide (Compound 506),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-oxa-8-azaspiro[3.5]nonan-8-yl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 519),

2-(3,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 528),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-isopropyl-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 553),

2-[4-(cyanomethyl)-1-piperidyl]-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 556),

2-(3,3-difluoropyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 565),

2-(2-ethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 591),

2-[(3R,5S)-3,5-dimethyl-1-piperidyl]-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 642),

6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3R)-3-methyl-1-piperidyl]-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 668),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)-2-[3-(2-pyridyl)-1-piperidyl]pyridine-3-carboxamide (Compound 677),

6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2S)-2-phenylpyrrolidin-1-yl]-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 716),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-fluoro-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 728),

2-(azepan-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 734),

6-(3-fluoro-5-isobutoxy-phenyl)-2-indolin-1-yl-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 739),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-methoxy-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 749),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-methyl-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 756),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-methoxy-4-methyl-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 774),

2-(3-ethoxy-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 843),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-fluoro-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 878),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 904),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)-2-[3-(trifluoromethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 907),

6-(3-fluoro-5-isobutoxy-phenyl)-2-[(3S)-3-phenyl-1-piperidyl]-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 912),

2-[(3R,5R)-3,5-dimethyl-1-piperidyl]-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 942),

2-(7-azaspiro[3.5]nonan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 944),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,3,3-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 952),

6-(3-fluoro-5-isobutoxy-phenyl)-2-[4-(methoxymethyl)-1-piperidyl]-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 975),

6-(3-fluoro-5-isobutoxy-phenyl)-2-isoindolin-2-yl-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 981),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,2,3-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 997),

2-(3,3-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1005),

2-(4,4-difluoro-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1006),

6-(3-fluoro-5-isobutoxy-phenyl)-2-[3-(methoxymethyl)-1-piperidyl]-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1028),

2-(4-ethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1030),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1041),

6-(3-fluoro-5-isobutoxy-phenyl)-2-[(2R)-2-phenylpyrrolidin-1-yl]-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1060),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-methyl-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1083),

2-[4-(cyclopropylmethoxy)-1-piperidyl]-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1088),

2-(4,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1110),

2-(2,6-dimethylmorpholin-4-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1119),

2-(6-azaspiro[2.5]octan-6-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1127),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methylpyrrolidin-1-yl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1142),

2-(4-ethoxy-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1155),

2-(2,3-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1169),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(3,3,4-trimethyl-1-piperidyl)pyridine-3-carboxamide (Compound 1174),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)-2-[(4R)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1176),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-methoxy-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1245),

2-(2-benzylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1257),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)-2-[[3-(trifluoromethyl)cyclohexyl]amino]pyridine-3-carboxamide (Compound 1281),

2-(8-azaspiro[3.4]octan-8-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1282),

6-(6-isopropoxy-3-pyridyl)-*N*-(1H-pyrazol-5-ylsulfonyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1352),

2-(3-cyclopropyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1286),

2-(2,2-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1302),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-phenyl-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1315),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isopropoxy-phenyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 600),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isopropoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 972),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isopropoxy-phenyl)pyridine-3-carboxamide (Compound 1123),

N-(4-aminophenyl)sulfonyl-6-*tert*-butyl-2-(1-piperidyl)pyridine-3-carboxamide (Compound 753),

N-(4-aminophenyl)sulfonyl-2-(benzylamino)-6-*tert*-butyl-pyridine-3-carboxamide (Compound 1064),

N-(4-aminophenyl)sulfonyl-6-*tert*-butyl-2-(4-methoxy-1-piperidyl)pyridine-3-carboxamide (Compound 1298),

N-(4-aminophenyl)sulfonyl-6-*tert*-butyl-2-[(4-methyltetrahydropyran-4-yl)methylamino]pyridine-3-carboxamide (Compound 752),

N-(4-aminophenyl)sulfonyl-6-*tert*-butyl-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 1037),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 794),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(3,3-dimethylbutoxy)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(3-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 1008),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-[4-(methoxymethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 986),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(3,5-dimethyl-1-piperidyl)pyridine-3-carboxamide (Compound 617),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-[4-(1-hydroxy-1-methyl-ethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 1038),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(4-methoxy-1-piperidyl)pyridine-3-carboxamide (Compound 1164),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(4-ethoxy-1-piperidyl)pyridine-3-carboxamide (Compound 1200),

2-[(4-methyltetrahydropyran-4-yl)methylamino]-6-(p-tolyl)-*N*-(1H-pyrazol-3-ylsulfonyl)pyridine-3-carboxamide (Compound 879)

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(4-methyltetrahydropyran-4-yl)methylamino]-6-(p-tolyl)pyridine-3-carboxamide (Compound 1196),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 373),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-[(3S)-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 1261),

N-[(2-amino-3-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1316),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(4R)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1240),

N-[(2-amino-3-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1159),

N-[(6-amino-2-pyridyl)sulfonyl]-6-methyl-2-[(3S)-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 532),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-[(3R)-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 626),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1074),

6-(6-isopropoxy-3-pyridyl)-2-[(3R)-3-phenyl-1-piperidyl]-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 741),

6-(6-isopropoxy-3-pyridyl)-2-[(3S)-3-phenyl-1-piperidyl]-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 776),

2-[(3R,5S)-3,5-dimethyl-1-piperidyl]-6-(6-isopropoxy-3-pyridyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 446),

2-[(3R,5R)-3,5-dimethyl-1-piperidyl]-6-(6-isopropoxy-3-pyridyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 990),

6-(6-isopropoxy-3-pyridyl)-2-(4-methyl-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 534),

N-(benzenesulfonyl)-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-azabicyclo[2.2.2]octan-3-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1657),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(3S)-2,2-dideuterio-3,5,5-trimethyl-pyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1400),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-azabicyclo[3.1.1]heptan-3-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1529),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,5,5-trimethylmorpholin-4-yl)pyridine-3-carboxamide (Compound 1548),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(5,7-dimethylindolin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1572),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(4R)-4-ethyl-2,2-dimethyl-pyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1611),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(4S)-4-ethyl-2,2-dimethyl-pyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1612),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,5,5-tetramethylmorpholin-4-yl)pyridine-3-carboxamide (Compound 1675),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,5-diisopropylpyrazol-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1387),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,3-difluoroazetidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2155),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-methylazetidin-1-yl)pyridine-3-carboxamide (Compound 1840),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-fluoroazetidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2128),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(6,6-dimethyl-7-azaspiro[3.4]octan-7-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2589),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(5-azabicyclo[4.1.0]heptan-5-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2376),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(6-methyl-3-azabicyclo[3.1.1]heptan-3-yl)pyridine-3-carboxamide (Compound 2224),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-{[1-(2-hydroxy-2-methyl-propyl)cyclopropyl]methylamino}pyridine-3-carboxamide (Compound 1723),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(6-methyl-8-azaspiro[3.4]octan-8-yl)pyridine-3-carboxamide (Compound 2552),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethylazetidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1928),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2532) (Isomer 1) (Compound 2178) (Isomer 2) (Compound 2062) (Isomer 3) (Compound 1863) (Isomer 4),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,5-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2451) (Isomer 1) (Compound 1993) (Isomer 2) (Compound 1864) (Isomer 3) (Compound 1855) (Isomer 4),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-hydroxy-1-piperidyl)pyridine-3-carboxamide (Compound 2323),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[cyclohexylmethyl(methyl)amino]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2289),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,5-trimethyl-1-piperidyl)pyridine-3-carboxamide (Compound 1995),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-isopropyl-2,2-dimethyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2201),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-BLAHyl-pyridine-3-carboxamide (Compound 1821),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(5-methyl-7-azaspiro[2.4]heptan-7-yl)pyridine-3-carboxamide (Compound 1754),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-tert-butyl-2,2-dimethyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2538),

tert-butyl (2R)-1-[3-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-pyridyl]pyrrolidine-2-carboxylate (Compound 2216),

tert-butyl (2S)-1-[3-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-pyridyl]pyrrolidine-2-carboxylate (Compound 2204),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethyl-1-piperidyl)pyridine-3-carboxamide (Compound 2581),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4,4-difluoro-2,2-dimethyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2631),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(5,5-dimethyl-6-azaspiro[2.4]heptan-6-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1746),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3-methylazetidin-1-yl)pyridine-3-carboxamide (Compound 2117),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[2,2-dimethyl-4-(trifluoromethyl)pyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1788),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[2-methyl-5-(trifluoromethyl)-1-piperidyl]pyridine-3-carboxamide (Compound 1835),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,5-dimethylazepan-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2479),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4,4-dimethylazepan-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2171),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3,3-dimethylazepan-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1901),


N-[(6-amino-2-pyridyl)sulfonyl]-2-(cyclohexylmethylamino)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2375),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(1-piperidyl)pyridine-3-carboxamide (Compound 571),

and

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethylpyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 361).

PREPARATION 40: 2-(Benzenesulfonyl)-3-[5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazol-3-yl]-3-oxo-propanoate (Compound 4)




5-*tert*-Butyl-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxylic acid (100 mg, 0.35 mmol) was dissolved in thionyl chloride (26 μL, 0.35 mmol) and a drop of *N,N*-dimethylformamide was added. The mixture was stirred for 1h and evaporated to give the acid chloride.

A mixture of the acid chloride (40 mg, 0.13 mmol), methyl 2-(benzenesulfonyl)acetate (28 mg, 0.13 mmol) and NaH (60%, 5.3 mg, 0.13 mmol) in *N,N*-dimethylformamide (1 mL) were stirred at room temperature for 15 min. The mixture was filtered and purified by reverse-phase preparatory-HPLC (10-99% CH_3CN/H_2O, HCl modifier) to give 2-(benzenesulfonyl)-3-[5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazol-3-yl]-3-oxo-propanoate (Compound 4). ESI-MS *m/z* calc. 482.6, found 483.4 $(M+1)^+$; Retention time: 2.41 min (3 min run).

PREPARATION 41: 6-(3-isoButoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 125)


Step 1: 6-chloro-*N*-[(2-methoxy-3-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide



6-Chloro-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (5.09 g, 17.5 mmol) was dissolved in thionyl chloride (6.36 mL, 87.3 mmol), CH_2Cl_2 (1 mL) and *N,N*-dimethylformamide (one drop). The reaction mixture was stirred for 30 min before it was concentrated to give the acid chloride.

The acid chloride was dissolved in CH_2Cl_2 (1 mL) and was added to solution of 2-methoxypyridine-3-sulfonamide (3.69 g, 17.5 mmol) in CH_2Cl_2 (1 mL) containing triethylamine (7.30 mL, 52.4 mmol) at 0 °C. The mixture was allowed to stir at room temperature overnight before it was diluted with CH_2Cl_2 and washed with 1N HCl, saturated $NaHCO_3$, and then brine. The organic layer was dried over sodium sulfate and concentrated to give 6-chloro-*N*-[(2-methoxy-3-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (6.20 g, 77%) ESI-MS *m/z* calc. 461.1, found 462.2 $(M+1)^+$; Retention time: 0.74 min (1 min run).


Step 2: 6-chloro-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide



A mixture of 6-chloro-*N*-[(2-methoxy-3-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (500 mg, 1.08 mmol), 1,4-dioxane (10 mL) and HCl (2.71 mL of 4 M, 10.8 mmol) was heated at 90 °C for 3.5 h. The volatiles were removed to give 6-chloro-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-

carboxamide (372 mg, 77%) ESI-MS *m/z* calc. 447.1, found 448.2 $(M+1)^+$; Retention time: 0.63 min (1 min run).

Step 3: 6-(3-isoButoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 125)



A mixture of 6-chloro-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (56 mmol, 0.10 mmol), 3-isobutoxyphenyl boronic acid (39 mg, 0.2 mmol), $Pd(PPh_3)_4$ (7 mg, 0.006 mmol), K_2CO_3 (2 M, 200 μL, 0.4 mmol) and *N,N*-dimethylformamide (1 mL) was flushed with nitrogen and sealed. The mixture was then stirred at 80 °C for 18h. The resultant solid was diluted with water and filtered. Then solid was taken up in ethyl acetate, filtered and concentrated to give 6-(3-isobutoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 125). ESI-MS *m/z* calc. 561.2, found 562.2 $(M+1)^+$; Retention time: 2.23 min (3 min run).

The following compound can be synthesized using the procedures described herein:

6-(4-chloro-3-propoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 72),

2-(2,4-dimethylphenoxy)-6-(4-ethoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 517),

6-(4-chloro-3-isobutoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 136),

6-(2,5-difluorophenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 545),

6-(3,4-difluorophenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 546),

6-(2-fluoro-5-propoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 552),

6-(4-fluorophenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 561),

6-(4,4-dimethylcyclohexen-1-yl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 573),

6-(4-chlorophenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 579),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-indol-4-ylsulfonyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 549),

2-(4,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(3-hydroxyphenyl)sulfonyl-pyridine-3-carboxamide (Compound 1309),

2-(4,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(1H-indol-4-ylsulfonyl)pyridine-3-carboxamide (Compound 1323),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-methoxy-3-pyridyl)pyridine-3-carboxamide (Compound 726),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,5-dimethoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1471),

isopropyl 4-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-(2,4,6-trimethylphenoxy)-2-pyridyl]-3,6-dihydro-2H-pyridine-1-carboxylate (Compound 1366),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-ethylcyclohexen-1-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1370),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-benzyl-3,6-dihydro-2H-pyridin-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1389),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1,2-dimethylprop-1-enyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1390),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(trifluoromethyl)cyclohexen-1-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1402),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4,7,7-trimethyl-3-bicyclo[2.2.1]hept-2-enyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1410),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,3-dihydrofuran-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1412),

ethyl 4-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-(2,4,6-trimethylphenoxy)-2-pyridyl]cyclohex-3-ene-1-carboxylate (Compound 1432),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1,4-dioxaspiro[4.5]dec-8-en-9-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1437),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6,6-dimethyl-2,3-dihydropyran-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1460),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(cyclohepten-1-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1513),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-bicyclo[2.2.1]hept-2-enyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1518),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5,5-dimethyl-3-oxo-cyclohexen-1-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1528),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5,5-dimethyl-2-oxo-cyclohexyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1427),

N-[(6-amino-2-pyridyl)sulfonyl]-6-cycloheptyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1376),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,2-dimethyltetrahydropyran-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1466),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-ethylcyclohexyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1475),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1,2-dimethylpropyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1486),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,3-dimethyl-5-oxo-cyclohexyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1492),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,6-dihydro-2H-thiopyran-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1517),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(trifluoromethyl)cyclohexyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1519),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropylcyclohexen-1-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1630),

6-(4-acetylcyclohexyl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1631),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1,7,7-trimethylnorbornan-2-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1650),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[1-(2-methylpropanoyl)-3,6-dihydro-2H-pyridin-4-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1540),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1,4-dioxaspiro[4.5]dec-8-en-8-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1558),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,3-dimethyl-6-oxo-cyclohexen-1-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1598),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,2,5,5-tetramethyl-3-furyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1609),

6-(4-acetylcyclohexen-1-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1615),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1,2,3,6-tetrahydropyridin-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1643),

6-(1-acetyl-3,6-dihydro-2H-pyridin-4-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1649),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-isopropoxy-5-(trifluoromethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1674),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-ethoxy-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1488),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-ethoxy-5-(trifluoromethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1569),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isopropoxy-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1671),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxy-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1646),

N-(2-azidophenyl)sulfonyl-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide,

N-(2-azidophenyl)sulfonyl-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-cyclohexyl-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 704),

6-(4-fluoro-2-methoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 608),

6-(4-fluoro-3-methoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 624),

6-(3-fluoro-5-isopropoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 645),

6-(3-fluoro-5-methoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 652),

N-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 654),

6-(2-fluoro-5-methoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 762),

6-(2-fluoro-3-isobutoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 804),

6-(3,5-difluorophenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 815),

6-(4-ethylphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 831),

6-(3-ethoxy-5-fluoro-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 835),

6-(2-fluoro-4-methoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 839),

6-(2-chlorophenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 877),

6-(cyclohexen-1-yl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 882),

6-(2,5-dimethoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 888),

6-(3-ethoxy-4-fluoro-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 917),

6-(6-methoxy-5-methyl-3-pyridyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 930),

6-(6-ethoxy-3-pyridyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 932),

6-(1-methylindol-6-yl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 936),

6-(2-chloro-3-propoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 955),

6-(3-butoxy-2-chloro-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 964),

6-(4-methoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 980),

6-(3-chloro-4-propoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 999),

6-(4-isopropylphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1002),

6-[4-(cyclopropylmethoxy)phenyl]-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1007),

6-(3-fluoro-4-propoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1022),

6-(4-methylcyclohexen-1-yl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1052),

6-(4-isobutoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1079),

6-(3-chlorophenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1095),

6-[3-(cyclopropylmethoxy)phenyl]-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1105),

2-(2,4-dimethylphenoxy)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6-(p-tolyl)pyridine-3-carboxamide (Compound 1129),

N-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6-(3-propoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1143),

6-(5-fluoro-2-methoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1151),

6-(4-ethoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1179),

6-(4-butoxy-3-fluoro-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1187),

6-(3-isopropoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1188),

6-(2-chloro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1201),

N-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6-(4-propoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1202),

6-(2,3-difluorophenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1296),

N-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6-(p-tolyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1300),

6-(5-ethoxy-2-fluoro-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1301),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 169),

2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6-(p-tolyl)pyridine-3-carboxamide (Compound 392),

6-(6-isopropoxy-3-pyridyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4R)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 403),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isopropoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 487),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 492),

6-(6-isopropoxy-3-pyridyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 497),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-isobutoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 511),

6-[3-(cyclopropylmethoxy)-5-fluoro-phenyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 541),

6-[3-(2,2-dimethylpropoxy)-5-fluoro-phenyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 572),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(4-isopropylphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 589),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4R)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 605),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6-phenyl-pyridine-3-carboxamide (Compound 660),

6-[3-(cyclopropylmethoxy)-5-fluoro-phenyl]-2-(4-methyl-1-piperidyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 664),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isopentyloxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 688),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(2-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 721),

6-(3-fluoro-5-isopropoxy-phenyl)-2-(4-methyl-1-piperidyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 737),

6-(4-chlorophenyl)-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 760),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 770),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-methoxy-2-pyridyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 781),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(4-isobutoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 891),

6-(4-chloro-3-isobutoxy-phenyl)-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 902),

6-[3-(cyclobutylmethoxy)-5-fluoro-phenyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 922),

6-(4-chloro-3-isobutoxy-phenyl)-2-(4-methyl-1-piperidyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 941),

6-(3-isobutoxyphenyl)-2-(4-methyl-1-piperidyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 958),

6-(4-chloro-3-isopropoxy-phenyl)-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 1001),

6-(4-isobutoxyphenyl)-2-(4-methyl-1-piperidyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 1017),

2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-6-(3-ethoxy-5-fluoro-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 1096),

8,8-dimethyl-2-(4-methyl-1-piperidyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 1195),

6-(4-chloro-3-propoxy-phenyl)-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 1293),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(3S)-3-phenyl-1-piperidyl]pyridine-3-carboxamide (Compound 481),

6-(2-methoxy-5-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2530),

6-(3-ethoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2235),

6-(4-methoxy-2-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2469),

6-(4-ethoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2207),

6-(2-methoxy-3-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1880),

6-isobutyl-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2467),

6-(3-methoxy-4-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2440),

6-(3-isopropoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2549),

6-(4-isopropoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1774),

6-(2-isobutoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2629),

6-(3-fluoro-4-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2219),

6-(3-isobutoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2267),

6-(2-fluoro-6-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2422),

6-(4-isobutoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2153),

6-(2-fluoro-3-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2222),

6-(5-fluoro-2-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1830),

N-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-6-[3-(trifluoromethoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1934),

6-(2-fluoro-5-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2606),

N-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-6-[4-(trifluoromethoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2234),

6-(4-fluoro-2-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2356),

6-(3-fluoro-5-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2527),

N-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-6-[3-(trifluoromethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2144),

6-(2-fluoro-4-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1858),

6-(3-fluoro-2-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1779),

6-(4-fluoro-3-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1780),

6-(2-naphthyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1803),

6-(1-naphthyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2007),

6-(2,4-dimethylphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2037),

N-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-6-(2,3,5-trifluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1837),

N-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-6-(2,4,5-trifluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1981),

6-(2,3-dihydro-1,4-benzodioxin-6-yl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2516),

6-(2-hydroxy-3-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2300),

6-(3-ethoxy-5-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2512),

6-(2,3-dimethylphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2073),

6-(2-hydroxy-5-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2174),

6-(5-fluoro-2-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2067),

6-(2-ethoxy-5-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2410),

6-(4-methoxy-2,6-dimethyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2575),

6-[3-(hydroxymethyl)-5-methoxy-phenyl]-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2247),

6-(2,3-dimethoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1786),

6-(2,5-dimethoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1752),

6-(2-hydroxy-4-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2148),

6-(3,5-dimethoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1878),

6-(2-ethoxy-5-fluoro-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2051),

6-(5-ethoxy-2-fluoro-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2138),

6-(3-ethoxy-2-fluoro-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1802),

6-(2-ethoxy-4-fluoro-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1867),

6-(3,5-dimethylphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2626),

6-(3-ethoxy-5-fluoro-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2278),

6-(3-ethoxy-4-fluoro-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2459),

6-(3,4-dimethylphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2154),

6-(4-methyl-1-naphthyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1811),

6-(2-isopropylphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2635),

6-(4,5-difluoro-2-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1904),

6-(3,5-difluoro-2-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2496),

6-(2,3-difluoro-4-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2567),

6-(3,5-difluoro-4-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1801),

6-(3,4-difluoro-5-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1914),

6-(2,5-difluoro-4-methoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1913),

6-(4-hydroxy-3,5-dimethyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2556),

6-(1H-indol-5-yl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1982),

6-(1H-indol-4-yl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2636),

6-(6-ethoxy-4-methyl-2-pyridyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1745),

6-(3-fluoro-4-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2084),

6-(2-fluoro-4-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2000),

6-(3-fluoro-2-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2491),

6-(2-chlorophenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1784),

6-(3-fluoro-5-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2348),

6-(2-fluoro-5-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1946),

6-(3-chlorophenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1791),

6-(3,4-difluorophenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2284),

6-(4-chlorophenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2028),

6-(2,4-difluorophenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2159),

6-(2,3-difluorophenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2162),

6-(3,5-difluorophenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2652),

6-(2,5-difluorophenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1954),

6-(2-fluorophenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2542),

6-(2-isopropoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2541),

6-(3-cyanophenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2185),

6-(3-fluorophenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1925),

6-(4-fluorophenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1807),

N-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-6-[2-(trifluoromethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1960),

6-(2-methoxy-3-pyridyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2454),

N-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-6-[4-(trifluoromethyl)cyclohexyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2315),

N-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-6-[4-(trifluoromethyl)cyclohexyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1710),

6-(2,5-dimethylphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1990),

6-(o-tolyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2176),

6-(4-fluoro-3-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2456),

6-(5-fluoro-6-methoxy-3-pyridyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2444),

6-(6-methoxy-2-pyridyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2101),

6-(2,6-dimethoxy-3-pyridyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1853),

6-(3-isopropylphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2498),

6-(4-isopropylphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2483),

6-(5-fluoro-2-hydroxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2295),

6-(2-methoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2485),

6-(2,3-dihydrobenzofuran-7-yl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2583),

6-(4-methoxy-3-methyl-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2369),


6-(2-ethoxyphenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1848),

6-(3-fluoro-5-isobutoxy-phenyl)-2-(4-methyl-1-piperidyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 422), and

6-(2-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 279).


PREPARATION 42: *N*-(3-aminopyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 197)

Step 1: 8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5*H*-quinoline-3-carbonitrile




A mixture of 2-chloro-8,8-dimethyl-6,7-dihydro-5*H*-quinoline-3-carbonitrile (1.00 g, 4.51 mmol) and 4-methylpiperidine (4.29 mL, 36.3 mmol) was stirred at 100 °C for 1 h. The reaction mixture was diluted with water and was extracted with ethyl acetate (3x). The combined extracts were washed with water, dried over sodium sulfate, and evaporated. The residue was purified by silica gel chromatography with 0-5% ethyl acetate in hexanes to give 8,8-dimethyl-2-(4-methylpiperidin-1-yl)-5,6,7,8-tetrahydroquinoline-3-carbonitrile (1.00 g, 78%) as a colorless oil. ESI-MS *m/z* calc. 283.2, found 284.3 $(M+1)^+$; Retention time: 0.94 min (1 min run).

Step 2: 8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5*H*-quinoline-3-carboxylic acid



A solution of 8,8-dimethyl-2-(4-methylpiperidin-1-yl)-5,6,7,8-tetrahydroquinoline-3-carbonitrile (0.80 g, 2.8 mmol) and KOH (1.6 g, 28 mmol) in ethanol (5.6 mL) was heated at reflux for 4 d. The reaction mixture was neutralized with 1 M HCl, and the solids were filtered, washed with water, and dried under vacuum to give 8,8-dimethyl-2-(4-methylpiperidin-1-yl)-5,6,7,8-tetrahydroquinoline-3-carboxylic acid (0.71 g, 83%) as a tan solid. ESI-MS *m/z* calc. 302.2, found 303.3 $(M+1)^+$; Retention time: 0.58 min (1 min run).


Step 3: *N*-(3-chloropyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide



A solution of 8,8-dimethyl-2-(4-methylpiperidin-1-yl)-5,6,7,8-tetrahydroquinoline-3-carboxylic acid (0.11 g, 0.36 mmol), *N*-[(dimethylamino)-1H-1,2,3-triazolo-[4,5-b]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (170 mg, 0.44 mmol) and sodium carbonate (77 mg, 0.73 mmol) in *N*,*N*-dimethylformamide (1.8 mL) was stirred for 5 min

before 3-chloropyrazine-2-sulfonamide (110 mg, 0.55 mmol) was added. The reaction mixture was stirred for 20 h, filtered, and subjected to preparatory-HPLC (1-99% CH_3CN/water) to give *N*-(3-chloropyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide (75mg, 43%). ESI-MS *m/z* calc. 477.2, found 478.3 $(M+1)^+$; Retention time: 0.65 min (1 min run).

Step 4: *N*-(3-Aminopyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide



N-(3-Chloropyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide (35 mg, 0.073 mmol) and ammonium hydroxide (1 mL, 26 mmol) was heated in a sealed vial in an microwave reactor at 120 °C for 20 min. The reaction mixture was concentrated, the residue was taken up in DMSO and subjected to preparatory-HPLC (1-99% CH_3CN/water) to give *N*-(3-aminopyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 197) (8.7 mg, 26%). ESI-MS *m/z* calc. 458.2, found 459.3 $(M+1)^+$; Retention time: 1.67 min (3 min run).

The following compound can be synthesized using the procedures described herein:

N-(3-aminopyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(p-tolyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 707),

N-(3-aminopyrazin-2-yl)sulfonyl-6-(4-ethoxyphenyl)-2-(p-tolyl)pyridine-3-carboxamide (Compound 819),

N-(3-aminopyrazin-2-yl)sulfonyl-6-(3-fluoro-5-isopropoxy-phenyl)-2-(p-tolyl)pyridine-3-carboxamide (Compound 1351),

N-(3-aminopyrazin-2-yl)sulfonyl-2,6-bis(p-tolyl)pyridine-3-carboxamide (Compound 820),

N-(3-aminopyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(p-tolyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 707),

N-(3-aminopyrazin-2-yl)sulfonyl-2-(2,4-dimethylphenoxy)-6-(4-ethoxyphenyl)pyridine-3-carboxamide (Compound 434),

N-[(6-amino-3-pyridyl)sulfonyl]-6-*tert*-butyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 702),

N-(6-aminopyrazin-2-yl)sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 586),

N-(3-aminopyrazin-2-yl)sulfonyl-8,8-dimethyl-2-(2,4,6-trimethylphenoxy)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 1223),

N-(3-aminopyrazin-2-yl)sulfonyl-2-(2,4-dimethylphenoxy)-8,8-dimethyl-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 1239),

N-(3-aminopyrazin-2-yl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1208),

N-(3-aminopyrazin-2-yl)sulfonyl-2-(2,4-dimethylphenoxy)-6-(3-fluoro-5-isopropoxy-phenyl)pyridine-3-carboxamide (Compound 559),

N-(3-aminopyrazin-2-yl)sulfonyl-2-(2,4-dimethylphenoxy)-6-(p-tolyl)pyridine-3-carboxamide (Compound 607),

N-(3-aminopyrazin-2-yl)sulfonyl-2-(2,4-dimethylphenoxy)-6-(1-isobutylpyrazol-4-yl)pyridine-3-carboxamide (Compound 713),

N-(3-aminopyrazin-2-yl)sulfonyl-2-(2,4-dimethylphenoxy)-6-(6-methoxy-5-methyl-3-pyridyl)pyridine-3-carboxamide (Compound 893),

N-(3-aminopyrazin-2-yl)sulfonyl-6-(3-fluoro-5-isopropoxy-phenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 989),

N-(3-aminopyrazin-2-yl)sulfonyl-6-(4-ethoxyphenyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1019),

N-(3-aminopyrazin-2-yl)sulfonyl-6-(p-tolyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1047),

N-(3-aminopyrazin-2-yl)sulfonyl-6-(1-isobutylpyrazol-4-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1103),

N-(3-aminopyrazin-2-yl)sulfonyl-6-(6-methoxy-5-methyl-3-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1249),

N-(3-aminopyrazin-2-yl)sulfonyl-2-[(4-methyltetrahydropyran-4-yl)methylamino]-6-(p-tolyl)pyridine-3-carboxamide (Compound 1066),

N-(3-aminopyrazin-2-yl)sulfonyl-6-(4-ethoxyphenyl)-2-[(4-methyltetrahydropyran-4-yl)methylamino]pyridine-3-carboxamide (Compound 1203),

N-(3-aminopyrazin-2-yl)sulfonyl-6-(1-isobutylpyrazol-4-yl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 1288),

N-(3-aminopyrazin-2-yl)sulfonyl-6-(6-methoxy-5-methyl-3-pyridyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 1191),

N-(3-aminopyrazin-2-yl)sulfonyl-2-(4-methyl-1-piperidyl)-6-(p-tolyl)pyridine-3-carboxamide (Compound 1084),

N-(3-aminopyrazin-2-yl)sulfonyl-6-(3-fluoro-5-isopropoxy-phenyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 699),

N-(3-aminopyrazin-2-yl)sulfonyl-6-(4-ethoxyphenyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 507),


8,8-dimethyl-2-(4-methyl-1-piperidyl)-*N*-(1H-pyrazol-5-ylsulfonyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 800),

N-[(6-amino-2-pyridyl)sulfonyl]-8,8-dimethyl-2-(4-methyl-1-piperidyl)-6,7-dihydro-5*H*-quinoline-3-carboxamide (Compound 750), and

N-[(2-amino-3-pyridyl)sulfonyl]-6-*tert*-butyl-2-(p-tolyl)pyridine-3-carboxamide (Compound 636).


PREPARATION 43: *N*-[(3-Aminophenyl)sulfonimidoyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 292)

Step 1: sodium 3-nitrobenzene-1-sulfinate




To a colorless solution of sodium sulfite (6.26 g, 49.6 mmol) in water (50 mL) at 80 °C, were added 3-nitrobenzenesulfonyl chloride (5.00 g, 22.6 mmol) and sodium carbonate (4.30 g, 40.6 mmol). The reaction mixture was stirred at 80 °C for 2h. The reaction mixture was concentrated under reduced pressure. The residue was suspended in ethanol (45 mL) and heated at 80 °C for 1H. The reaction mixture was cooled to room temperature and filtered to remove most of the salts. The filtrate was concentrated under reduced pressure to give sodium 3-nitrobenzene-1-sulfinate (4.20g, 89% yield, containing some salts). ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.68 (t, J = 7.8 Hz, 1H), 8.02 (dt, J = 7.7, 1.4 Hz, 1H), 8.21 – 8.28 (m, 1H), 8.47 – 8.51 (m, 1H). $[M+H]^+$ = 188.0.

Step 2: methyl 3-nitrobenzene-1-sulfinate




Thionyl chloride (3.27 mL, 45.0 mmol) was added to a solution of sodium 3-nitrobenzene-1-sulfinate (3.14 g, 15.0 mmol) in dichloromethane (21 mL) at 0 °C. After 20 minutes at 0 °C, the reaction mixture was allowed to warm to room temperature and stirred for 3h. The reaction mixture was concentrated and dried under vacuum to give a residue (3.08 g, 15 mmol). To the residue was added dichloromethane (5 mL) and methanol (100 mL) at 0 °C. After 10 minutes at this temperature, triethylamine (6.27 mL, 45.0 mmol) was added and the mixture was stirred at room temperature overnight. The mixture was concentrated under reduced pressure. The residue was dissolved in dichloromethane (50 mL), washed with water (25 mL), brine (25 mL), dried over anhydrous sodium sulfate and concentrated to give methyl 3-nitrobenzene-1-sulfinate (1.40 g, 46% yield). ^{1}H NMR (300 MHz, $CDCl_3$) δ 3.52 (s, 3H), 7.71 (t, J = 7.8 Hz, 1H), 7.98 (dt, J = 7.7, 1.4 Hz, 1H), 8.21 – 8.28 (m, 1H), 8.47 – 8.51 (m, 1H). $[M+H]^+$ = 202.0.

Step 3: 6-*tert*-butyl-*N*-(3-nitrobenzenesulfinyl)-2-(2,4,6-trimethylphenoxy)-pyridine-3-carboxamide



To a solution methyl 3-nitrobenzene-1-sulfinate (1.35 g, 6.70 mmol) in dimethylformamide (8.0 mL) was added methyl 3-nitrobenzene-1-sulfinate (697 mg, 2.68 mmol), followed by cesium carbonate (2.18 g, 6.69 mmol). The mixture was stirred overnight at 60 °C. The mixture was diluted with ethyl acetate (60 mL), washed with water (30 mL), brine (30 mL), dried over anhydrous sodium sulfate, filtered, and concentrated. The crude material was purified by chromatography on silica gel, eluting with 10 – 20% ethyl acetate in heptane to give 6-*tert*-butyl-*N*-(3-nitrobenzenesulfinyl)-2-(2,4,6-trimethylphenoxy)-pyridine-3-carboxamide (725 mg, impure). This material was used in the next step without further purification. $[M+H]^+$ = 482.1.


Step 4: 6-*tert*-butyl-*N*-[imino(3-nitrophenyl)oxo- λ^6-sulfanyl]-2-(2,4,6-trimethyl-phenoxy)pyridine-3-carboxamide



To a solution of 6-*tert*-butyl-*N*-(3-nitrobenzenesulfinyl)-2-(2,4,6-trimethylphenoxy)-pyridine-3-carboxamide (724 mg, 1.50 mmol) and potassium *tert*-butoxide (337 mg, 3.00 mmol) in acetonitrile (7.5 mL) was added hexamethyldisilazane (0.94 mL, 4.5 mmol), followed by *N*-chlorosuccinimide (0.60 g, 4.5 mmol). After stirring for 15 min, additional hexamethyldisilazane (0.94 mL, 4.5 mmol), followed by *N*-chlorosuccinimide (0.60 g, 4.5 mmol) were added. The mixture was stirred until TLC indicated completion. The mixture was concentrated and the residue was purified by flash chromatography, eluting with 20 – 50% ethyl acetate in heptane to give 6-*tert*-butyl-*N*-[imino(3-nitrophenyl)oxo- λ^6-sulfanyl]-2-(2,4,6-trimethyl-phenoxy)pyridine-


3-carboxamide (95 mg, 3% over two steps). ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.07 (s, 9H), 2.03 (s, 6H), 2.30 (s, 3H), 6.87 (s, 2H), 6.98 (d, J = 8.0 Hz, 1H), 7.74 (t, J = 8.2 Hz, 1H), 8.30 (d, J = 8.0 Hz, 1H), 8.41 – 8.53 (m, 2H), 8.95 (s, 1H). $[M+H]^+$ = 497.2.

Step 5: *N*-[(3-aminophenyl)sulfonimidoyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide



To a solution of 6-*tert*-butyl-*N*-[imino(3-nitrophenyl)oxo-λ^6-sulfanyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (90 mg, 0.18 mmol) in methanol (2.6 mL) was added 10% Pd/C (19 mg, 0.018 mmol) and the mixture was stirred at room temperature for 3h. The solvent was removed under vacuum and the crude material was purified by flash chromatography eluting with 20 – 50% ethyl acetate / heptane to give *N*-[(3-aminophenyl)sulfonimidoyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (52 mg, 62% yield). ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.01 (s, 9H), 1.92 (s, 6H), 2.22 (s, 3H), 5.57 (br. s, 2H), 6.73 (d, J = 8.1 Hz, 1H), 6.85 (s, 2H), 6.97-7.07 (m, 2H), 7.11-7.20 (m, 2H), 7.56 (br. s, 2H), 8.09 (d, J = 7.8 Hz, 1H). $[M+H]^+$ = 467.2.

Step 6: *N*-[(3-aminophenyl)sulfonimidoyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 19 and Compound 53)



N-[(3-Aminophenyl)sulfonimidoyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (30 mg, 0.064 mmol) was taken up in methanol (1.8 mL) and CH_2Cl_2 (0.4 mL). The solution was subjected to chiral SFC (30 µL injections; column: Chiralpak AS-H (250 x 10 mm),5µm; mobile phase: 25% methanol w/20mM NH_3, 75% CO_2; flow: 10 mL/min) to give *N*-

[(3-aminophenyl)sulfonimidoyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)-pyridine-3-carboxamide [peak 1, enantiomer 1, 97.9% ee, ^{1}H NMR (400 MHz, Chloroform-d) δ 8.27 (d, *J* = 7.9 Hz, 1H), 7.49 - 7.36 (m, 2H), 7.22 (t, *J* = 7.9 Hz, 1H), 6.92 (d, *J* = 7.8 Hz, 1H), 6.87 - 6.75 (m, 3H), 3.89 (s, 2H), 2.29 (s, 3H), 2.02 (s, 6H), 1.06 (s, 9H)] and *N*-[(3-aminophenyl)sulfonimidoyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide [peak 2, enantiomer 2, 96% ee, ^{1}H NMR (400 MHz, Chloroform-d) δ 8.27 (d, *J* = 7.9 Hz, 1H), 7.49 - 7.33 (m, 2H), 7.23 (t, *J* = 7.9 Hz, 1H), 6.92 (d, *J* = 7.9 Hz, 1H), 6.88 - 6.74 (m, 3H), 3.89 (s, 2H), 2.29 (s, 3H), 2.02 (s, 6H), 1.06 (s, 9H)].

The following compounds can be prepared using the procedures described herein:


N-[*S*-(3-aminophenyl)-*N*-methyl-sulfonimidoyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 568),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[1-phenylethyl]pyrazole-3-carboxamide can be separated to give *N*-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(1*S*)-1-phenylethyl]pyrazole-3-carboxamide (Compound 138) and *N*-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(1*R*)-1-phenylethyl]pyrazole-3-carboxamide (Compound 540) using Chiralcel OJ-H (250 x 10mm), 5μm column;

N-(3-aminophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide can be separated to give (1*S*)-*N*-(3-aminophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 126) (Compound 126) and (1*R*)-*N*-(3-aminophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (Compound 310) using Chiralpak AD-H (250 x 10mm), 5μm column.

PREPARATION 44: *N*-[(6-Amino-3-fluoro-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 226)

Step 1: 2-(benzylthio)-3,6-difluoropyridine



To a solution of 2,3,6-trifluoropyridine (532 mg, 4.00 mmol) and phenylmethanethiol (470 μL, 4.00 mmol) in THF (20 mL) was added NaH (60%, 176 mg, 4.40 mmol) in portions.


After 1 h, the mixture was quenched with saturated aqueous ammonium chloride and extracted with ethyl acetate (3x). The combined extracts were washed with water, dried over sodium sulfate, and evaporated. The residue was purified by silica gel chromatography with 0-20% ethyl acetate in hexanes to give 2-(benzylthio)-3,6-difluoropyridine (0.63 g, 66%) as a colorless oil. ESI-MS *m/z* calc. 237.0, found 237.9 $(M+1)^+$; Retention time: 0.72 min (1 min run).

Step 2: 3,6-difluoropyridine-2-sulfonyl chloride




Chlorine was bubbled through a vigorously stirred solution of 2-(benzylthio)-3,6-difluoropyridine (0.27 g, 1.1 mmol) in chloroform and water for min, and the reaction mixture was stirred for 1 h. Chlorine was bubbled through the mixture for another 15 min, and the reaction mixture was stirred for 2 h. The reaction mixture was quenched with saturated aqueous sodium metabisulfite solution and extracted with dichloromethane (3x). The combined extracts were washed with water, dried over sodium sulfate, and evaporated to give crude 3,6-difluoropyridine-2-sulfonyl chloride (50% pure, 0.49 g) as a colorless oil. ESI-MS *m/z* calc. 213.0, found 213.8 $(M+1)^+$; Retention time: 0.47 min (1 min run).

Step 3: 3,6-difluoropyridine-2-sulfonamide




A solution of 5,6-difluoropyridine-2-sulfonyl chloride (0.35 g, 1.6 mmol) in dioxane (4 mL) was cooled in an ice bath before ammonia (6.6 mL of 0.5 M in dioxane, 3.3 mmol) was added. The reaction was stirred for 3 h and the solvent was evaporated. The residue was purified by silica gel chromatography with 0-50% ethyl acetate in hexanes to give 3,6-difluoropyridine-2-sulfonamide (72 mg, 23%) as a colorless solid. ESI-MS *m/z* calc. 194.0, found 194.8 $(M+1)^+$; Retention time: 0.15 min (1 min run).

Step 4: 6-*tert*-butyl-*N*-[(3,6-difluoro-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide



A solution of 6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (107 mg, 0.340 mmol), 3,6-difluoropyridine-2-sulfonamide (66 mg, 0.34 mmol), *N*-[(dimethylamino)-1H-1,2,3-triazolo-[4,5-b]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (194 mg, 0.510 mmol) and sodium carbonate (72 mg, 0.68 mmol) in *N,N*-dimethylformamide (2 mL) was stirred at room temperature for 18 h. The reaction was diluted with water and extracted with ethyl acetate (3x). The combined extracts were washed with water, dried over sodium sulfate, and evaporated. The residue was purified by silica gel chromatography with 0-5% methanol in dichloromethane to give 6-(*tert*-butyl)-*N*-((3,6-difluoropyridin-2-yl)sulfonyl)-2-(mesityloxy)nicotinamide (33 mg, 18%) as a colorless solid. ESI-MS *m/z* calc. 489.2, found 490.2 $(M+1)^+$; Retention time: 0.85 min (1 min run).


Step 5: *N*-[(6-amino-3-fluoro-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide



A mixture of 6-*tert*-butyl-*N*-[(5,6-difluoro-2-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (15 mg, 0.028 mmol) and ammonium hydroxide (1.0 mL of 30 %w/v, 8.6 mmol) was heated in a sealed vessel in a microwave reactor at 130 °C for 30 min and 150 °C for 60 min. The solvent was evaporated and the residue was subjected to preparatory-HPLC (30%–99% acetonitrile / water (5 mM HCl)) to give *N*-[(6-amino-3-fluoro-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (0.7 mg, 5%). ESI-MS *m/z* calc. 486.2, found 487.3 $(M+1)^+$; Retention time: 2.15 min (3 min run).


PREPARATION 45: (2*S*,4*R*)-*N*-[(6-Amino-2-pyridyl)sulfonyl]-4-phenyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolidine-2-carboxamide (Compound 236)

Step 1: (2*S*,4*R*)-4-phenylpyrrolidine-2-carboxylic acid trifluoroacetic acid salt



(2*S*,4*R*)-1-(Boc)-4-Phenylpyrrolidine-2-carboxylic acid (1.01 g, 3.47 mmol) was dissolved in trifluoroacetic acid (20 ml) and was stirred for 1 hour at room temperature. The reaction mixture was concentrated under reduced pressure and the resulting residue was left under high vacuum for 18 h. The residue was triturated in a minimal amount of diethyl ether and placed in an ultrasound bath for 1 hour, filtered and then dried under high vacuum to provide (2*S*,4*R*)-4-phenylpyrrolidine-2-carboxylic acid trifluoroacetic acid salt (999 mg, 99%) as solid. $[M+H]^+$ = 192.2. ^{1}H NMR (300MHz,$CDCl_3$) δ 1.94-2.11 (m, 1H), 2.61-2.75 (m, 1H), 3.09-3.22 (m, 1H), 3.42-3.59 (m, 1H), 3.59-3.73 (m, 1H), 4.35-4.48 (m, 1H), 7.15-7.40 (m, 5H), 8.52 - 10.23 (br. s. 2H).


Step 2: (2*S*,4*R*)-4-phenyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolidine-2-carboxylic acid



(2*S*-4*R*)-4-Phenylpyrrolidine-2-carboxylic acid trifluoroacetic acid salts (999 mg, 3.47 mmol) and potassium hydroxide (1.17g, 20.8 mmol) were dissolved in isopropanol (25 mL) and heated at 40 °C. A solution of 2-(bromomethyl)-1,3,5-trimethylbenzene (849 mg, 3.99 mmol) in iso-propanol (20 mL) was added over 60 min via a syringe. The reaction was stirred for additional 2h at this temperature. The reaction mixture was treated with concentrated HCl until the pH was between 3 and 4. The resulting solid was filtered off and the filtrate was concentrated


under reduced pressure. The resultant solid was dissolved in dichloromethane and was purified on silica gel (0-10% methanol in ethyl acetate). Upon dilution of the residue with dichloromethane, the silica was filtered off and the filtrate was concentrated under reduced pressure to provide (2*S*-4*R*)-4-phenyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolidine-2-carboxylic acid (1.1g, 98%) as a solid. $[M+H]^+ = 324.2$. ^1H NMR (300MHz,$CDCl_3$) δ 2.22 (s, 3H), 2.27-2.34 (m, 1H), 2.38(s, 6H), 2.59-2.94 (m, 1H), 2.96-3.30 (m, 1H) , 3.36 (t, *J*=11.2Hz, 1H), 3.47-3.80 (m, 1H), 4.05 (m, 1H), 4.20 (d, *J*=13.4Hz,1H), 4.35 (d, *J*=13.4Hz,1H), 6.83 (s, 2H), 7.05-7.33(m, 5H).

Step 3: (2*S*,4*R*)-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-4-phenyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolidine-2-carboxamide



(2*S*,4*R*)-4-Phenyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolidine-2-carboxylic acid (75 mg, 0.23 mmol), 6-fluoropyridine-2-sulfonamide (192 mg, 1.09 mmol), *N*-[(dimethylamino)-1H-1,2,3-triazolo-[4,5-b]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (497 mg, 1.31 mmol), and ethyldiisopropylamine (380 µL, 2.18 mmol) were combined in *N*,*N*-dimethylformamide (1.440 mL) and heated at 60°C for 16 h. The reaction mixture was diluted with water (1 mL) and was extracted with ethyl acetate (2 x 5mL). The organic layers were dried over sodium sulfate, filtered and concentrated. The residue was subjected to preparatory-HPLC utilizing a gradient of 30-90% acetonitrile in 5 mM aq HCl to give (2*S*,4*R*)-*N*-[(6-fluoro-2-pyridyl)sulfonyl]-4-phenyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolidine-2-carboxamide. ^1H NMR (400 MHz, DMSO) δ 9.18 (s, 1H), 8.16 (d, *J* = 7.8 Hz, 1H), 7.85 (d, *J* = 9.1 Hz, 1H), 7.30 (s, 7H), 6.88 (s, 2H), 4.40 (s, 2H), 4.24 (s, 1H), 3.81 (s, 1H), 3.33 – 3.21 (m, 2H), 2.76 (s, 1H), 2.37 (s, 6H), 2.21 (s, 3H), 2.00 (s, 1H).

Step 4: (2*S*,4*R*)-*N*-[(6-amino-2-pyridyl)sulfonyl]-4-phenyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolidine-2-carboxamide



(2*S*,4*R*)-*N*-[(6-Fluoro-2-pyridyl)sulfonyl]-4-phenyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolidine-2-carboxamide (from step 3) was suspended in NH_4OH (2 mL of 30 %w/v, 17 mmol). The mixture was heated in a microwave at 150 °C. The reaction mixture was concentrated, redissolved in DMSO and subjected to preparatory-HPLC utilizing a gradient of 1-60% acetonitrile in 5 mM aq HCl to give (2*S*,4*R*)-*N*-[(6-amino-2-pyridyl)sulfonyl]-4-phenyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolidine-2-carboxamide (Compound 236) (7.5 mg, 7% over two steps). ^{1}H NMR (400 MHz, MeOD) δ 7.84 (s, 1H), 7.33 (s, 7H), 6.99 (d, J = 8.7 Hz, 1H), 6.90 (s, 2H), 4.60 (d, J = 35.7 Hz, 2H), 4.41 (s, 1H), 3.98 (s, 1H), 3.76 (s, 1H), 3.67 - 3.55 (m, 1H), 2.90 (dt, J = 12.8, 6.3 Hz, 1H), 2.41 (d, J = 14.1 Hz, 7H), 2.24 (s, 4H). ESI-MS *m/z* calc. 478.2, found 479.0 $(M+1)^+$; Retention time: 1.41 min (3 min run).

The following compounds can be synthesized using the procedures described herein:

(2*R*,4*S*)-*N*-[(6-amino-2-pyridyl)sulfonyl]-4-phenyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolidine-2-carboxamide (Compound 467) and

(2*S*,4*S*)-*N*-[(6-amino-2-pyridyl)sulfonyl]-4-phenyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolidine-2-carboxamide (Compound 870)

PREPARATION 46: *N*-(4-Aminothiazol-2-yl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 265)


Step 1: *tert*-butyl thiazol-4-ylcarbamate



A mixture of thiazole-4-carboxylic acid (8.16 g, 63.2 mmol) and triethylamine (9.7 mL, 70 mmol) in *tert*-butanol (320 mL) was treated with diphenyl phosphoryl azide (15 mL, 70

mmol) and heated gradually in an oil bath to 100 °C and stirred for 17 h. Once cooled, the crude reaction mixture was concentrated under reduced pressure to remove most of the volatiles. The residue was transferred to a 1.0-L separatory funnel with ethyl acetate (300 mL) and the organic layer was washed with water (2 x 200 mL) and brine (2 x 100 mL). The organic layer was then dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified silica gel chromatography, eluting from 0% to 50% ethyl acetate in heptanes. The resultant gummy solid was triturated with heptanes, filtered, washed with heptanes and dried under high vacuum to afford *tert*-butyl thiazol-4-ylcarbamate (3.42 g, 27%) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) δ 1.54 (s, 9 H), 7.30 (br. s, 1 H), 8.33 (br. s, 1 H), 8.58 (d, J = 2.3 Hz, 1 H). $[M\text{-}C_4H_8+H]^+$ = 145.1.


Step 2: *tert*-butyl *N*-(2-sulfamoyl-1,3-thiazol-4-yl)carbamate



A solution of *tert*-butyl thiazol-4-ylcarbamate (1.50 g, 7.49 mmol) in methyl *tert*-butyl ether (35 mL) was cooled in an ice bath and treated slowly with *iso*-propylmagnesium chloride (11 mL of a 2.0 M solution in THF, 22 mmol). A voluminous white solid appeared. After 20 minutes in the ice bath, sulfur dioxide was gently bubbled into the reaction mixture for 5 minutes then the reaction mixture was stirred at room temperature for 1 h. The crude reaction mixture was concentrated under reduced pressure to remove most of the volatiles and the solid obtained was suspended in water (80 mL) and cooled in an ice bath. Sodium acetate (3.40 g, 41.5 mmol) and hydroxylamine-*O*-sulfonic acid (3.39 g. 30.0 mmol) were successively added. The ice-bath was removed and the reaction was left to stir at room temperature for 45 h. The reaction mixture was transferred to a separatory funnel and the aqueous layer was extracted with ethyl acetate (2 x 50 mL). The combined organic layers were washed with 5% aqueous sodium bicarbonate (30 mL), brine (30 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography, eluting from 0% to 75% ethyl acetate in heptanes, to provide 1.2 g of a solid that was contaminated with *iso*-propylsulfonamide. The solid was dissolved in methanol (3 mL) and water (2 mL) was added dropwise with stirring. The precipitate was filtered, washed with water and dried under high


vacuum to afford *tert*-butyl *N*-(2-sulfamoyl-1,3-thiazol-4-yl)carbamate (0.90 g, 43%) as a white solid. ^{1}H NMR (300 MHz, DMSO-d_6) δ 1.46 (s, 9 H), 7.56 (s, 2 H), 9.12 (s, 1 H), 9.24 (s, 1 H). $[M-C_4H_8+H]^+ = 224.0$.

Step 3: *N*-[2-[[6-*tert*-Butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carbonyl]sulfamoyl]thiazol-4-yl]carbamate



K_2CO_3 (45 mg, 0.35 mmol) was added to a mixture of 6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (44 mg, 0.14 mmol), *tert*-butyl *N*-(2-sulfamoylthiazol-4-yl)carbamate (39 mg, 0.14 mmol), *N*-[(dimethylamino)-1H-1,2,3-triazolo-[4,5-b]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (64 mg, 0.17 mmol) and *N*,*N*-dimethylformamide (1 mL) at room temperature. The mixture was stirred at 60 °C for 3 h before it was cooled to room temperature. The mixture was partitioned between ethyl acetate and 1N HCl. The layers were separated and the aqueous layer was extracted with ethyl acetate (2x). The combined organics were washed with brine, dried over magnesium sulfate, filtered and concentrated. The residue was subjected to silica gel column chromatography (0-50% ethyl acetate/hexanes), then subjected to preparatory-HPLC(10-99% acetonitrile/water with 0.05 mM HCl) to give *tert*-butyl *N*-[2-[[6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carbonyl]sulfamoyl]thiazol-4-yl]carbamate as a white solid. ESI-MS *m/z* calc. 574.2, found 575.4 $(M+1)^+$; Retention time: 2.48 min (3 min run).

Step 4: *N*-(4-Aminothiazol-2-yl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide




Trifluoroacetic acid (11 μL, 0.14 mmol) was added to a mixture of *tert*-butyl *N*-[2-[[6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carbonyl]sulfamoyl]thiazol-4-yl]carbamate (from step 3) and CH_2Cl_2 (5 mL) at room temperature. The mixture was allowed to stir for 1H before it was concentrated under reduced pressure. The residue was subjected to preparatory-HPLC(10-99% acetonitrile/water with 0.05 mM HCl) to give *N*-(4-aminothiazol-2-yl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 265). ESI-MS *m/z* calc. 474.1, found 475.5 $(M+1)^+$; Retention time: 2.11 min (3 min run). ^1H NMR (400 MHz, DMSO-d6) δ 11.96 (s, 1H), 9.07 (s, 1H), 8.00 (d, *J* = 7.8 Hz, 1H), 7.13 (d, *J* = 7.8 Hz, 1H), 6.91 (s, 2H), 6.81 (s, 2H), 2.25 (s, 3H), 1.96 (s, 6H), 1.04 (s, 9H).

The following compounds can be prepared using the procedures described herein:

N-(4-aminothiazol-2-yl)sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 842) and *N*-(4-aminothiazol-2-yl)sulfonyl-2-(4,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1168)


PREPARATION 47: *N*-[(6-Amino-2-pyridyl)sulfonyl]-6-(*tert*-butylamino)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 273)

Step 1: methyl 6-(*tert*-butylamino)-2-(mesityloxy)pyridine-3-carboxylate




A mixture of methyl 6-chloro-2-(mesityloxy)pyridine-3-carboxylate (170 mg, 0.58 mmol) and *t*-butylamine (4.0 mL, 38 mmol) in 500 mL dimethylacetamide was heated in a microwave at 150 °C for 4 h. The crude mixture was purified by silica gel column chromatography eluting with a gradient of 0-40% ethyl acetate/hexanes to afford methyl 6-(*tert*-butylamino)-2-(mesityloxy)pyridine-3-carboxylate (76 mg, 38% yield) as an orange solid. LC-MS: $(M+H)^+ = 342.5$.

Step 2: 6-(*tert*-butylamino)-2-(mesityloxy)pyridine-3-carboxylic acid



A mixture of methyl 6-(*tert*-butylamino)-2-(mesityloxy)pyridine-3-carboxylate (76 mg, 0.22 mmol) and sodium hydroxide (14 mg, 0.34 mmol) in methanol (1.0 mL), THF (1.0 mL), and water (0.50 mL) was heated at 60 °C for 7 h. A few drops of concentrated HCl were added, solid formed, everything was rotary evaporated, and dried under vacuum to give crude 6-(*tert*-butylamino)-2-(mesityloxy)pyridine-3-carboxylic acid (98 mg) as a white solid. LC-MS: $(M+H)^+ = 329.3$.


Step 3: 6-(*tert*-butylamino)-2-(mesityloxy)-*N*-((6-nitropyridin-2-yl)sulfonyl)nicotinamide



A mixture of crude 6-(*tert*-butylamino)-2-(mesityloxy)pyridine-3-carboxylic acid (0.22 mmol), 6-nitropyridine-2-sulfonamide (58 mg, 0.29 mmol), *N*-[(dimethylamino)-1H-1,2,3-triazolo-[4,5-b]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide


(0.12 g, 0.33 mmol), ethyldiisopropylamine (0.16 mL, 0.92 mmol) and *N,N*-dimethylformamide (1 mL) was stirred in a capped vial at 70 °C for 6 h and then at room temperature for 18 h. LC-MS at that point showed mostly the activated acid. An additional 70 mg of 6-nitropyridine-2-sulfonamide and 0.080 mL of ethyldiisopropylamine were added. The mixture was stirred at 70 °C for 4 h and the crude was purified twice by silica gel column chromatography eluting with a gradient of 0-20% methanol/dichloromethane to give 6-(*tert*-butylamino)-2-(mesityloxy)-*N*-((6-nitropyridin-2-yl)sulfonyl)nicotinamide (34 mg, 30% yield) as a light yellow solid. LC-MS: $(M+H)^+ = 514.5$.

Step 4: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(*tert*-butylamino)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide



A mixture of 6-(*tert*-butylamino)-2-(mesityloxy)-*N*-((6-nitropyridin-2-yl)sulfonyl)nicotinamide (34 mg, 0.064 mmol) and iron metal (48 mg, 0.86 mmol) in THF (0.7 mL), EtOH (0.35 mL), conc. HCl (0.06 mL) and water (0.06 mL) was heated at 60 °C for 50 min. The mixture was filtered hot and was washed with methanol. The filtrate was rotary evaporated to give 120 mg of crude material which was purified by reverse phase HPLC to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(*tert*-butylamino)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 273) (7.2 mg, 23%). LC-MS: $(M+H)^+ = 484.5$.

PREPARATION 48: 6-(*tert*-butoxy)-2-(mesityloxy)nicotinic acid



To a solution of t-butanol (0.70 mL, 7.3 mmol, 11 eq) in 1.0 mL of *N,N*-dimethylformamide was added portion-wise 60% NaH (320 mg, 8.0 mmol, 12 eq). After bubbling stopped, about 10 min, 6-chloro-2-(mesityloxy)pyridine-3-carboxylic acid 7 (0.20 g, 0.69 mmol, 1.0 eq) was added to the reaction mixture. The mixture was heated in a capped vial at 100 °C for 22 hr. After cooling to room temperature, a couple of drops of concentrated hydrochloric acid was added and the reaction mixture was evaporated to dryness. The crude material was purified by column chromatography using 25 g of silica gel and eluting with a gradient of 0-30% methanol in dichlormethane over 15 minutes. The product-containing fractions were combined and concentrated to afford 6-*tert*-butoxy-2-(mesityloxy)pyridine-3-carboxylic acid (109 mg, 48% yield) as a light yellow solid. LC-MS: (M+H)+ = 330.1.

The following compound can be synthesized using the procedures described herein:

N-[(6-amino-2-pyridyl)sulfonyl]-6-(isopropylamino)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 650),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-methylpyrrolidin-1-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1553),

N-[(6-amino-2-pyridyl)sulfonyl]-6-pyrrolidin-1-yl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 746),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-piperidyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 785),

N-[(6-amino-2-pyridyl)sulfonyl]-6-morpholino-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 966),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(methylamino)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 983),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(ethylamino)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1128),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(dimethylamino)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1306),

N-[(6-amino-2-pyridyl)sulfonyl]-6-ethoxy-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 637),

N-[(6-amino-2-pyridyl)sulfonyl]-6-methoxy-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 806),

N-[(6-amino-2-pyridyl)sulfonyl]-8,8-dimethyl-2-(2,4,6-trimethylphenoxy)-5,7-dihydropyrano[4,3-*b*]pyridine-3-carboxamide (Compound 714),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[isobutyl(methyl)amino]-5-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1416),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1494),

N-[(6-amino-2-pyridyl)sulfonyl]-5-methyl-6-(2-methylpyrrolidin-1-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1516),

N-[(6-amino-2-pyridyl)sulfonyl]-5-fluoro-2,6-bis[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1673),

N-[(6-amino-2-pyridyl)sulfonyl]-6-isopropoxy-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1573),


N-[(6-amino-2-pyridyl)sulfonyl]-6-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-5-methyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1589),

N-[(6-amino-2-pyridyl)sulfonyl]-6-isopropoxy-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 968), and

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butoxy-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1070).


PREPARATION 49: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-cyclopentyl-pyridine-3-carboxamide (Compound 1219)

Step 1: 6-(*tert*-Butyl)-2-(cyclohex-1-en-1-yl)-*N*-((6-fluoropyridin-2-yl)sulfonyl)nicotinamide



6-*tert*-butyl-2-chloro-*N*-[(6-fluoro-2-pyridyl)sulfonyl]pyridine-3-carboxamide (75 mg, 0.20 mmol), 1-cyclohexenylboronic acid (25.41 mg, 0.2017 mmol), sodium carbonate (201.7 µL of 2 M, 0.4034 mmol), dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (7.376 mg, 0.01008 mmol) in dioxane (900.0 µL) were added to a microwave reactor vial. The vial was purged with nitrogen, capped and heated to 120 °C for 30 minutes in a microwave reactor. The reaction mixture was diluted with water and extracted with ethyl acetate (2 x 10 mL). The organic layers were dried over sodium sulfate, concentrated, and purified by HPLC utilizing a gradient of 0-70% acetonitrile in 5 mM aq HCl to give the product (19.7 mg, 23%) ESI-MS *m/z* calc. 417.5, found 418.4 $(M+1)^+$; Retention time: 0.64 min (1 min run).

Step 2: 6-(*tert*-butyl)-2-cyclohexyl-*N*-((6-fluoropyridin-2-yl)sulfonyl)nicotinamide



6-(*tert*-Butyl)-2-(cyclohex-1-en-1-yl)-*N*-((6-fluoropyridin-2-yl)sulfonyl)nicotinamide (19.7 mg, 0.047 mmol) was dissolved in methanol (10 mL) and palladium on carbon (12.64 mg, 0.01188 mmol) was added. The mixture was stirred under a hydrogen balloon at room temperature for 16 hours. The crude material was filtered, concentrated, and purified by HPLC utilizing a gradient of 20-80% acetonitrile in 5 mM aq HCl to give 6-(*tert*-butyl)-2-cyclohexyl-*N*-((6-fluoropyridin-2-yl)sulfonyl)nicotinamide.

The following compound can be synthesized using the procedures described herein: *N*-(4-aminophenyl)sulfonyl-6-*tert*-butyl-2-cyclopentyl-pyridine-3-carboxamide (Compound 1003),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(3,3,5,5-tetramethylcyclopenten-1-yl)pyridine-3-carboxamide (Compound 1502),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[4-(trifluoromethyl)cyclohexen-1-yl]pyridine-3-carboxamide (Compound 1667),


N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,6,6-tetramethyl-3*H*-pyran-4-yl)pyridine-3-carboxamide (Compound 1368),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4,4-dimethylcyclohexen-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 1493),

and

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,5,5-tetramethyl-3-furyl)pyridine-3-carboxamide (Compound 1531).

Step 3: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-cyclohexyl-pyridine-3-carboxamide (Compound 411)




6-(*tert*-butyl)-2-cyclohexyl-*N*-((6-fluoropyridin-2-yl)sulfonyl)nicotinamide was suspended in ammonium hydroxide (1mL of 28 %w/v, 8.56 mmol). The mixture was stirred in a microwave reactor for 0.5 h at 150 °C. The reaction mixture was concentrated, redissolved in DMSO and purified by HPLC utilizing a gradient of 10-65% acetonitrile in 5 mM aq HCl to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(cyclohexen-1-yl)pyridine-3-carboxamide ^{1}H NMR (400 MHz, MeOD) δ 8.56 (d, *J* = 8.4 Hz, 1H), 7.98 (d, *J* = 8.4 Hz, 1H), 7.87 (s, 1H), 7.43 (d, *J* = 7.2 Hz, 1H), 7.03 (d, *J* = 8.7 Hz, 1H), 6.08 (s, 1H), 2.31 (s, 2H), 2.20 (s, 2H), 1.69 (d, *J* = 17.3 Hz, 4H), 1.53 (s, 9H). ESI-MS *m/z* calc. 414.17255, found 415.0 $(M+1)^+$; Retention time: 1.48 minutes.

The following compound can be synthesized using the procedures described herein:

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-cyclopentyl-pyridine-3-carboxamide (Compound


1219).

PREPARATION 50: 6-*tert*-butyl-*N*-[(1-methyl-2-oxo-3-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 667)



6-*tert*-butyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (28 mg, 0.060 mmol) was dissolved in *N,N*-dimethylformamide with potassium carbonate (24.72 mg, 0.1789 mmol) and a solution of iodomethane (8.464 mg, 3.712 µL, 0.05963 mmol) was added dropwise. The solution was stirred for 4 h at room temperature. The reaction mixture was filtered and purified by LC/MS utilizing a gradient of 10-99% acetonitrile in 5 mM aq HCl to yield 6-*tert*-butyl-*N*-[(1-methyl-2-oxo-3-pyridyl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 667) (2 mg, 0.004 mmol, 7%) ESI-MS *m/z* calc. 483.1828, found 484.3 $(M+1)^+$; Retention time: 2.13 minutes.


PREPARATION 51: methyl 2-(mesityloxy)-6-(prop-1-yn-1-yl)nicotinate



Prop-1-yne (25.78 µL, 0.4543 mmol) was bubbled through a solution of *N,N*-dimethylformamide (2mL) and triethylamine (409.6 µL, 2.939 mmol) and then added to a mixture of methyl 6-chloro-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (150. mg, 0.491 mmol), dichloropalladium; triphenylphosphane (34.44 mg, 0.04907 mmol), copper (I) iodide (14.02 mg, 0.07362 mmol) and triphenyl phosphine (17.05 µL, 0.07359 mmol) in deoxygenated *N,N*-dimethylformamide (2 mL) under continous nitorgen flow. The resulting solution was

purged with propyne and capped. The reaction was then stirred at 105 °C for 16 hours. The cooled mixture was diluted with ethyl acetate and passed through a plug of celite, washing with ethyl acetate (10 mL). Water (7.5 mL) was added to the filtrate and the layers were separated. The aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic extracts were washed with a saturated aqueous solution of sodium chloride (10 mL) and dried over sodium sulfate. After filtration the solvent was removed under reduced pressure to give a dark brown residue. The residue was purified by column chromatography on silica gel eluting 0-100 % ethyl acetate in hexanes to yield methyl 6-prop-1-ynyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (102.4 mg, 0.331 mmol, 72.9%) ESI-MS *m/z* calc. 309.1365, found 310.2 $(M+1)^+$; Retention time: 0.76 minutes.

Preparation 52: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(trifluoromethyl)pyrazol-1-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 676)



N-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (30. mg, 0.067 mmol), 5-(trifluoromethyl)-1*H*-pyrazole (27.41 mg, 0.2014 mmol), potassium carbonate (19.49 mg, 0.1410 mmol) and *trans*-diaminocyclohexane (1.534 mg, 1.613 µL, 0.01343 mmol) were combined in *N,N*-dimethylformamide. At this point copper(I) iodide (0.6393 mg, 0.003357 mmol) was added and the reaction was heated at 150 °C in the microwave for 2 hours. The reaction was filtered and purified by LC/MS utilizing a gradient of 10-99% acetonitrile in 5 mM aq HCl to yield *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(trifluoromethyl)pyrazol-1-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 676) (8 mg, 0.01 mmol, 20%) ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.54 (s, 1H), 8.32 (d, J = 8.1 Hz, 1H), 7.85 (dd, J = 2.6, 1.1 Hz, 1H), 7.69 - 7.62 (m, 2H), 7.22 (dd, J = 7.4, 0.8 Hz, 1H), 6.99 (d, J = 2.6 Hz, 1H), 6.97 (s, 2H), 6.72 (d, J = 8.4 Hz, 1H), 6.56 (s, 2H), 2.28 (s, 3H), 2.03 (s, 6H). ESI-MS *m/z* calc. 546.1297, found 547.2 $(M+1)^+$; Retention time: 2.1 minutes.

The following compound can be synthesized using the procedures described herein:


N-[(6-amino-2-pyridyl)sulfonyl]-6-pyrazol-1-yl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 564),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[5-methyl-3-(trifluoromethyl)pyrazol-1-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 633),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(trifluoromethyl)pyrazol-1-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 691), and

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-methylpyrazol-1-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1220).


PREPARATION 53: methyl 2-(mesityloxy)-6-(1-methylcyclopropyl)nicotinate



In a microwave vial, trimethylsulfoxonium iodide (5.2 g, 23 mmol), was suspended in anhydrous DMSO (26 mL) under nitrogen. Sodium hydride 60% in oil (1.0 g, 26 mmol) was added and the resulting solution was left stirring 10 min. at room temperature. Methyl 6-(prop-1-en-2-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (2.6 g, 8.4 mmol) was added dissolved in anhydrous THF (8 mL) and left stirring at room temperature 18 hours. The reaction mixture was partitioned between water (100 mL) and ethyl acetate (3 x 100 mL), separated, then the organic phase was concentrated under reduced pressure and the resulting residue was purified on silica gel using 0 to 10% ethyl acetate in heptane which provided the desired product methyl 6-(1-methylcyclopropyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (1.59 g, 58%) as a oil. ESI-MS *m/z* calc. 325.2, found 326.2 $(M+1)^+$; ^{1}H NMR (300 MHz, $CDCl_3$) ppm: 0.60 (m, 2 H), 0.84 (m, 2 H), 1.36 (s, 3 H), 2.01 (s, 6 H), 2.28 (s, 3 H), 3.92 (s, 3 H), 6.83 (s, 2 H), 6.96 (d, J = 7.9 Hz, 1 H), 8.16 (d, J = 7.9 Hz, 1 H).

PREPARATION 54: ethyl 2-(mesityloxy)-6-(1-(trifluoromethyl) cyclopropyl)nicotinate


Step 1: ethyl 2-(mesityloxy)-6-(3-(trifluoromethyl)-4,5-dihydro-3*H*-pyrazol-3-yl)nicotinate



| Preparation of a solution of diazomethane in ethyl ether: *N*-nitriso-*N*-methylurea (1.87 g, 15.8 mmol) was added to a biphasic mixture of 40% aqueous KOH (6.7 mL) and diethyl ether (20 mL) cooled in an ice-bath. The mixture was stirred in the ice-bath for 20 min (turns yellow), and then cooled in a dry-ice/acetone bath. Once the aqueous layer was frozen, the ether layer was decanted and the yellow solution was kept under nitrogen until ready to use.


The above solution of diazomethane (15.8 mmol) was added dropwise to a solution of ethyl 6-(3,3,3-trifluoroprop-1-en-2-yl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (1.5 g, 3.95 mmol) in ethyl ether (20 mL) at 0 °C. The yellow color of the diazomethane spontaneously disappeared upon addition. The reaction was stirred at 0 °C for 30 min, and TLC showed starting material was completely consumed. Acetic acid (1 mL) was added to quench excess the diazomethane until it turns colorless and bubbles evolution ceased. The mixture was then diluted with ethyl acetate (50 mL), washed with saturated aqueous sodium bicarbonate (20 mL), brine (20 mL), dried over anhydrous sodium sulfate and concentrated to give ethyl 2-(mesityloxy)-6-(3-(trifluoromethyl)-4,5-dihydro-3*H*-pyrazol-3-yl)nicotinate as a pale yellow oil (1.7 g). ^{1}H NMR (300 MHz, $CDCl_3$) ppm, 1.40 (t, J = 6.9 Hz, 3 H), 1.68 – 1.92 (m, 2 H), 1.98 (br s, 6 H), 2.31 (s, 3 H), 3.90 – 4.10 (m, 1 H), 4.43 (q, J = 6.9 Hz, 2 H), 4.64 (ddd, J = 18.1, 9.5, 4.2 Hz, 1 H), 6.87 (s, 2 H), 7.57 (d, J = 7.8 Hz, 1 H), 8.31 (d, J = 7.8 Hz, 1 H). LCMS: $[M+H]^+$=422.2.

Step 2: ethyl 2-(mesityloxy)-6-(1-(trifluoromethyl)cyclopropyl)nicotinate



Ethyl 2-(mesityloxy)-6-(3-(trifluoromethyl)-4,5-dihydro-3*H*-pyrazol-3-yl)nicotinate (1.7 g, crude, 3.9 mmol) was dissolved in *m*-xylene (25 mL). The solution was gently refluxed for 2 hours. TLC shows reaction complete. The solvent was removed and the residue was purified by silica gel chromatography, eluting with a mixture of ethyl acetate and heptanes (0-10%), to afford 1.6 g, crude (quantitative yield).as an unseparable mixture (1:1) of ethyl 6-[1-(trifluoromethyl)cyclopropyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate, and ethyl 6-[-1,1,1-trifluorobut-2-en-2-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (undefined stereochemistry of the olefin). In order to remove the alkene, the mixture of 6-[1-(trifluoromethyl)cyclopropyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate and ethyl 6-[(2Z or 2E)-1,1,1-trifluorobut-2-en-2-yl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (1.5 g, 3.81 mmol, 1:1) was dissolved in *tert*-butanol (5 mL). Water (3 mL) was added followed by citric acid (549 mg, 2.86 mmol) and potassium osmium(VI) oxide dehydrate (1.4 mg). Finally, 4-methylmorpholine *N*-oxide monohydrate (568 mg, 4.2 mmol) was added. The mixture was stirred at room temperature overnight. LCMS shows 50% conversion. Potassium osmium(VI) oxide dehydrate (1.4 mg) was added and the reaction was continued stirred until HPLC showed reaction was complete (approximately 40 hours). The reaction was concentrated to remove most of the *tert*-butanol. The resulting aqueous mixture was diluted with ethyl acetate (60 mL) and washed with water (10 mL), brine (10 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography, eluting with a mixture of ethyl acetate and heptanes (0-10% then 50%), to afford ethyl 6-[1-(trifluoromethyl)cyclopropyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (600 mg, 40%) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 0.96 – 1.05 (m, 2 H), 1.15 – 1.23 (m, 2 H), 1.39 (t, J = 7.2 Hz, 3 H), 2.01 (s, 6 H), 2.29 (s, 3 H), 4.41 (q, J = 7.2 Hz, 2 H), 6.85 (s, 2 H), 7.25 (d, J = 8.2 Hz, 1 H), 8.21 (d, J = 8.2 Hz, 1 H). LCMS: $[M+H]^+$ = 394.0.


PREPARATION 55: ethyl 2-(mesityloxy)-6-(perfluoroethyl)nicotinate



In a sealed tube, 3-(ethoxycarbonyl)-2-(mesityloxy)pyridine 1-oxide (1.85 g, 6.14 mmol) was dissolved in anhydrous tetrahydrofuran (18.5 mL) with flame dried (under high vacuum) cesium fluoride (93 mg, 0.05 mmol) then trimethyl(pentafluoroethyl)silane (3.53 g, 18.4 mmol) was added at 0 °C and the reaction mixture was stirred at room temperature under nitrogen for 5 days. Distilled water (10 mL) and ethyl acetate (50 mL) were added to the crude material. The organic phase was separated, dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The resulting residue was purified on silica gel using 0-10% ethyl acetate in heptane to provide ethyl 6-(pentafluoroethyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (480 mg, 25%) as an oil. $[M + H]^+ = 404.1$ ^1H NMR (300 MHz, $CDCl_3$) ppm: 1.41 (t, $J = 7.1$ Hz, 3 H), 2.03 (s, 6 H), 2.29 (s, 3 H), 4.45 (q, $J = 7.1$ Hz, 2H), 6.87 (s, 2 H), 7.39 (d, $J = 7.7$ Hz, 1 H), 8.36 (d, $J = 7.8$Hz, 1 H).

PREPARATION 56: *tert*-butyl (4-sulfamoylthiazol-2-yl)carbamate


Step 1: lithium (*tert*-butoxycarbonyl)(4-sulfinatothiazol-2-yl)amide



Methyllithium (0.70 mL, 1.6 M in diethyl ether, 1.1 mmol) was added to a solution of *tert*-butyl *N*-(4-bromo-1,3-thiazol-2-yl)carbamate (279 mg, 1.00 mmol) in anhydrous tetrahydrofuran (2 mL) and anhydrous diethyl ether (2 mL) at -78 °C. After 5 min, *tert*-butyl lithium (1.2 mL, 1.7 M in pentane, 2 mmol) was added at -78 °C. A light yellow suspension was formed. After 20 min, sulfur dioxide (gas) was bubbled into the reaction at -78 °C for 5 min. A yellow clear solution was obtained. The reaction was allowed to slowly warm up to room temperature and stirred overnight. The reaction mixture was concentrated to dryness and the


residue was triturated with diethyl ether (10 mL) to afford a light yellow solid (390 mg), which is a mixture of lithium (*tert*-butoxycarbonyl)(4-sulfinatothiazol-2-yl)amide and lithium bromide. $[M+Na]^+ = 287.0$.

Step 2: *tert*-butyl (4-sulfamoylthiazol-2-yl)carbamate



Hydroxylamine-O-sulfonic acid (147 mg, 1.2 mmol) was added to a solution of lithium 2-{[(*tert*-butoxy)carbonyl]amino}-1,3-thiazole-4-sulfinate (330 mg, containing lithium bromide, 0.91 mmol) and sodium acetate (148 mg, 1.8 mmol) in water (4 mL) at 0 °C. The reaction was stirred at room temperature for 2 hours. Copious amounts of solids precipitated out. HPLC showed complete conversion. The mixture was stirred at 0 °C for 5 min. The solid was filtered off, washed with cold water (1 mL) and dried under high vacuum to afford *tert*-butyl *N*-(4-carbamoyl-1,3-thiazol-2-yl)carbamate (130 mg, 51%) as a white solid. ^{1}H NMR (300 MHz, DMSO-d_6) δ 1.49 (s, 9 H), 7.43 (s, 2 H), 7.67 (s, 1 H), 11.84 (br. s, 1 H). $[M+Na]^+ = 302.0$.


PREPARATION 57: *tert*-butyl ((6-sulfamoylpyridin-2-yl)methyl)carbamate



A solution of 6-cyanopyridine-2-sulfonamide (1.00 g, 5.46 mmol) in methanol (25 mL) and ammonia (5 mL of a 7.0 N solution in methanol, 35 mmol) was treated with a spatula of Raney-Nickel (2800, slurry in water). The flask was purged three times with hydrogen then left to stir at room temperature under one atmosphere of hydrogen for 18 hours. The crude reaction mixture was filtered over a pad of celite, washed with methanol, and concentrated under reduced pressure. The residue was taken up in dichloromethane (25 mL) and treated successively with triethylamine (1.52 mL, 10.9 mmol) and di-*tert*-butyl dicarbonate (1.55 g, 7.10 mmol). After about 1 hour, tetrahydrofuran (10 mL) was added in an attempt to help solubilize the reaction mixture. After another hour, the reaction was concentrated under reduced pressure, transferred to


a 250-mL separatory funnel with water (50 mL) and extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with brine (50 mL), dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure. The residue was purified on 40g of silica gel utilizing a gradient from 50% to 100% ethyl acetate in heptanes to afford *tert*-butyl ((6-sulfamoylpyridin-2-yl)methyl)carbamate (217 mg, 14%) as a white solid. ^{1}H NMR (300 MHz, Acetone-d_6) ppm 1.43 (s, 9 H), 4.43 (d, J = 6.0 Hz, 2 H), 6.48 - 6.77 (m, 3 H), 7.58 (d, J = 7.8 Hz, 1 H), 7.83 (d, J = 7.8 Hz, 1 H), 7.97 - 8.10 (m, 1 H). $[M+Na]^+$ = 310.1.

PREPARATION 58: *N*-(2-aminothiazol-5-yl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1148)



In a 4mL vial a solution of *N*-(2-acetamidothiazol-5-yl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (14.6 mg, 0.0269 mmol) and aqueous hydrochloric acid (268.5 µL of 4 M, 1.074 mmol) in dioxane (277.4 µL) was stirred at 90 °C for 1 hour. The reaction mixture was cooled and purified directly by reverse-phase preparative chromatography utilizing a C18 column and a gradient of 30-99 acetonitrile in water containing 5 mM hydrochloric acid to afford *N*-(2-aminothiazol-5-yl)sulfonyl-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1148) (5.9 mg, 0.012 mmol, 45%) as a white solid. ^{1}H NMR (400 MHz, DMSO) δ 12.10 (s, 1H), 8.02 (s, 2H), 7.95 (d, J = 7.8 Hz, 1H), 7.67 (s, 1H), 7.12 (d, J = 7.8 Hz, 1H), 6.90 (s, 2H), 2.25 (s, 3H), 1.97 (s, 7H), 1.04 (s, 10H). ESI-MS m/z calc. 474.1, found 475.2 $(M+1)^+$; Retention time: 2.11 minutes.


PREPARATION 59: 6-(*tert*-butyl)-2-(mesityloxy)-4-methoxy-*N*-((3-nitrophenyl)sulfonyl)nicotinamide



In a 10 mL microwave vessel was added 6-*tert*-butyl-*N*-(3-nitrophenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (85 mg, 0.17 mmol) dissolved in anhydrous methanol (684.1 mg, 864.9 µL, 21.35 mmol) and xylene (850.0 µL) and to it was added diacetoxypalladium (3.835 mg, 0.01708 mmol), (diacetoxyiodo)benzene (165.0 mg, 0.5124 mmol), and 4 Angstrom molecular sieves. The mixture was heated in an oil bath to 100 °C for 18 hours. A second aliquot of diacetoxypalladium (3.835 mg, 0.01708 mmol) was added and the reaction mixture heated in an oil bath to 100 °C for an additional 48 hours. The mixture was allowed to cool and the mixture was filtered and the resulting solution was purified by reverse-phase preparative chromatography utilizing a C18 column and a gradient of 30-99 acetonitrile in water containing 5 mM hydrochloric acid to afford 6-*tert*-butyl-4-methoxy-*N*-(3-nitrophenyl)sulfonyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (20.7 mg, 0.0388 mmol, 23%). ^{1}H NMR (400 MHz, DMSO) δ 12.89 (s, 1H), 8.61 (t, J = 2.0 Hz, 1H), 8.49 (d, J = 8.2 Hz, 1H), 8.36 (d, J = 8.1 Hz, 1H), 7.92 (t, J = 8.1 Hz, 1H), 6.82 (s, 2H), 6.72 (s, 1H), 3.84 (s, 3H), 2.22 (s, 3H), 1.86 (s, 6H), 1.03 (s, 10H). ESI-MS *m/z* calc. 527.1726, found 528.2 $(M+1)^+$; Retention time: 2.2 minutes.

PREPARATION 60: 2-chloro-8,8-dimethyl-7,8-dihydro-5*H*-pyrano[4,3-*b*]pyridine-3-carbonitrile


Step 1: 5-(hydroxymethylene)-3,3-dimethyltetrahydro-4*H*-pyran-4-one



To a solution of 3,3-dimethyltetrahydropyran-4-one (1.8 g, 14 mmol) in ether (50 mL) was added ethyl formate (1.040 g, 1.134 mL, 14.04 mmol). The mixture was cooled to 0 °C. Sodium ethoxide (1.051 g, 15.44 mmol) was added at 0 °C. The mixture was allowed to warm to


room temperature, stirred overngiht and then evaporated to dryness. The crude material was used directly in next step without further purification.

Step 2: 2-hydroxy-8,8-dimethyl-7,8-dihydro-5*H*-pyrano[4,3-*b*]pyridine-3-carbonitrile



To a suspension of 5-(hydroxymethylene)-3,3-dimethyl-tetrahydropyran-4-one (2.2 g, 14 mmol) in pyridine (100 mL) was added 2-cyanoacetamide (1.177 g, 14.00 mmol). The mixture was heated at reflux ovemight. The reaction mixture was evaporated to dryness and the residue was partitioned between ethyl acetate and water. The aqueous layer was extracted thee times with ethyl acetate, and then three times with a 2:1 mixture of dichloromethane and isopropanol. The combined organic layers were washed with brine, dried over magnesium sulfate, filtered, and evaporated to dryness. The residue was washed with methanol to yield the product (205 mg, 1.00 mmol,7.1%). ^{1}H NMR (400 MHz, Chloroform-*d*) δ 7.55 (s, 1H), 4.54 (s, 2H), 3.61 (s, 2H), 3.49 (s, 1H), 1.42 (s, 6H).


Step 3: 2-chloro-8,8-dimethyl-7,8-dihydro-5*H*-pyrano[4,3-*b*]pyridine-3-carbonitrile



A mixture of 2-hydroxy-8,8-dimethyl-7,8-dihydro-5*H*-pyrano[4,3-*b*]pyridine-3-carbonitrile (240. mg, 1.17 mmol) in phosphorous oxychloride (2.5 mL, 26.82 mmol) was heated to 105 °C for 6 hours. The reaction mixture was evaporated to dryness. The residue was added dropwise to water, and then extracted three times with ethyl acetate. The combined organic layers were washed with brine, dried over magnesium sulfate, filtered, and concentrated to dryness. The crude material was purified by silica gel column chromatography (0-10% ethyl acetate in hexanes) to provide 2-chloro-8,8-dimethyl-7,8-dihydro-5*H*-pyrano[4,3-*b*]pyridine-3-

carbonitrile (240. mg, 1.08 mmol, 92.3%) as a yellow solid. ^{1}H NMR (400 MHz, Chloroform-*d*) δ 7.57 (s, 1H), 4.78 (d, J = 0.9 Hz, 2H), 3.71 (s, 2H), 1.33 (s, 6H).


PREPARATION 61: 6-(*tert*-butyl)-2-(mesityloxy)-*N*-(*N*-methyl-3-nitrophenylsulfonimidoyl)nicotinamide



To a solution of *rac*-6-*tert*-butyl-*N*-(3-nitrobenzenesulfinyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (253 mg, 0.525 mmol) in acetonitrile (4.1 mL) was added *N*-chlorosuccinimide (217 mg, 1.63 mmol) at 0 °C. The mixture was then stirred at room temperature until no more starting material was left (about 1 hour). Methylamine (0.79 mL of a 2.0 M solution in THF, 1.6 mmol) was added to the reaction at 0 °C and the mixture was stirred for 1 hour at room temperature and then quenched by the addition of water (3 mL). The mixture was extracted with ethyl acetate (2 x 10 mL), the organic layers were combined, dried over sodium sulfate, filtered, and concentrated under reduced pressure. The residue was purified on 25 g of silica gel utilizing a gradient of 0% to 45% ethyl acetate in heptanes to afford 6-(*tert*-butyl)-2-(mesityloxy)-*N*-(*N*-methyl-3-nitrophenylsulfonimidoyl)nicotinamide (170 mg, 63% yield) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.06 (s, 9 H), 2.02 (s, 6 H), 2.28 (s, 3 H), 2.78 (s, 3 H), 6.83 (s, 2 H), 6.91 (d, J = 7.8 Hz, 1 H), 7.74 (t, J = 8.0 Hz, 1 H), 8.24 (d, J = 8.0 Hz, 1 H), 8.38 (d, J = 8.3 Hz, 1 H), 8.45 (d, J = 7.7 Hz, 1 H), 8.83 – 8.87 (m, 1 H). $[M+H]^+$ = 511.1.


PREPARATION 62: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(5-methoxy-2-methyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 470)

Step 1: *N*-((6-aminopyridin-2-yl)sulfonyl)-2,6-dichloronicotinamide



2,6-dichloro-pyridine-3-carboxylic acid (20.2 g, 105 mmol) was dissolved in *N,N*-dimethylformamide (200 mL) and stirred under nitrogen and CDI (25.5 g, 158 mmol) was added in one portion. The reaction was warmed to 65 °C and heated for 1 h. In a separate vessel, 6-aminopyridine-2-sulfonamide (18.2 g, 105 mmol) was dissolved in *N,N*-dimethylformamide (100 mL) under nitrogen and NaH (4.40 g, 60%, 110 mmol) was added in portions to mitigate gas evolution and the reaction was stirred for 1 h. The two reactions were combined in one portion at 65 °C. The reaction was stirred for 30 min and evaporated to remove most of *N,N*-dimethylformamide. The residue was poured over ice and made acidic (pH=~3) by the addition of 6M HCl. Solid was collected by filtration and washed with water (100 mL) and methanol (30 mL) and combined to the following crude material after dried. The filtrate and washing were combined and extracted with ethyl acetate (3 x 500 mL). The organics were washed once with brine (500 mL), dried over sodium sulfate and evaporated. The crude material was suspended in a mixture solvent of dichloromethane/ethyl acetate/methanol (30/30/30 mL), the suspension sonicated for 30 min, and filtered. The solid washed with ether (2 x 150 mL) and dried to give 19.8 g white solid (54%). The filtrate was further concentrated to afford 6.0 g (16%) of product. LCMS: 347 [M+H]+; ^{1}H NMR (DMSO-d_6, 250 MHz): 8.11 (d, J = 7.5 Hz, 1H), 7.80-7.55 (m, 2H), 7.19 (d, J = 7.2 Hz, 1H), 6.75 (d, J = 8.5 Hz, 1H), 6.71 (s, 2H) ppm


Step 2: *N*-((6-aminopyridin-2-yl)sulfonyl)-6-chloro-2-(2,2,4-trimethylpyrrolidin-1-yl)nicotinamide



N-[(6-amino-2-pyridyl)sulfonyl]-2,6-dichloro-pyridine-3-carboxamide (1.02 g, 2.94 mmol), 2,2,4-trimethylpyrrolidine (Hydrochloric Acid (1)) (1.319 g, 8.814 mmol), and potassium carbonate (2.030 g, 14.69 mmol) were combined in dimethylsulfoxide (8.160 mL) and heated at 100 °C for 16 h. The reaction was diluted with water and the pH was adjusted to 7. The aqueous mixture was extracted with ethyl acetate (3 x 100 mL). The organics were combined, washed with brine, dried over sodium sulfate and evaporated. The crude material was purified by silica gel chromatography eluting with 0-100% ethyl acetate in hexanes to yield *N*-[(6-amino-2-

pyridyl)sulfonyl]-6-chloro-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (700. mg, 1.65 mmol, 56.2%) ESI-MS *m/z* calc. 423.1132, found 424.2 $(M+1)^+$; Retention time: 0.63 minutes.

Step 3:



N-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (42 mg, 0.10 mmol), (5-methoxy-2-methyl-phenyl)boronic acid (25 mg, 0.15 mmol), tetrakis(triphenylphosphine)palladium (0) (11 mg, 0.010 mmol), and 2M aqueous potassium carbonate (0.2 mL, 4 mmol) were combined in *N*,*N*-dimethylformamide (1 mL) and irradiated in the microwave reactor for 20 min at 120 °C. The reaction was filtered and purified by LC/MS utilizing a gradient of 10-99% acetonitrile in 5 mM aq HCl to yield *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(5-methoxy-2-methyl-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 470) as a hydrochloric acid salt (12 mg, 0.022 mmol, 22%). ESI-MS *m/z* calc. 509.21, found 510.5 $(M+1)^+$; Retention time: 1.42 minutes.

The following compound can be synthesized using the procedures described herein:

N-[(6-amino-2-pyridyl)sulfonyl]-6-(o-tolyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 761),

N-(benzenesulfonyl)-6-(3-fluoro-5-hydroxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1538),

N-[(6-amino-2-pyridyl)sulfonyl]-6-cyclopropyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1627),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-hydroxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1636)

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 630),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropylphenyl)-2-[(4R)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 890),

N-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 852),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 946),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,4-dimethylphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 614),

tert-butyl 5-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]-2-pyridyl]-3,6-dihydro-2*H*-pyridine-1-carboxylate (Compound 1371),

N-[(6-amino-2-pyridyl)sulfonyl]-3-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-5-(6-isopropoxy-3-pyridyl)pyrazine-2-carboxamide (Compound 1651),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-fluoro-5-isobutoxy-phenyl)-3-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyrazine-2-carboxamide (Compound 1526),

N-[(6-amino-2-pyridyl)sulfonyl]-6-norboman-2-yl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1378),

N-[(6-amino-2-pyridyl)sulfonyl]-6-isopropenyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1423),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6,6-dimethyl-2,3-dihydropyran-4-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1369),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,4-dihydro-2H-pyran-6-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1438),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,3-dihydrofuran-4-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1455),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1,2-dimethylpropyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1489),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tetrahydropyran-2-yl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1500),

N-[(2-oxo-1H-pyridin-3-yl)sulfonyl]-6-tetrahydropyran-2-yl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1503),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1,2-dimethylprop-1-enyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1510),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,4-dihydro-2H-pyran-5-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1524),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tetrahydrofuran-3-yl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1534),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tetrahydropyran-3-yl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1632),

6-(1-acetyl-4-piperidyl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1634) (Compound 946),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(trifluoromethyl)cyclohexen-1-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1664),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-cyclopropylethyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1677),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(trifluoromethyl)cyclohexyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1672),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-hydroxy-5,5-dimethyl-cyclohexyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1439),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,3-dimethyl-6-oxo-cyclohexen-1-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1462),

N-[(6-amino-2-pyridyl)sulfonyl]-6-isopropyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1505),

N-[(6-amino-2-pyridyl)sulfonyl]-6-ethyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1507),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,6-dihydro-2H-pyran-4-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1594),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tetrahydropyran-4-yl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1595),

N-[(6-amino-2-pyridyl)sulfonyl]-6-isobutyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1599),

tert-butyl 4-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]-2-pyridyl]-3,6-dihydro-2H-pyridine-1-carboxylate (Compound 1624),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-methylprop-1-enyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1683),

tert-butyl 4-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]-2-pyridyl]-2,3-dihydropyrrole-1-carboxylate (Compound 1684),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-fluoro-2-methyl-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 822),

N-[(6-amino-2-pyridyl)sulfonyl]-6-phenyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1334),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-fluorophenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 712),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-ethoxy-5-fluoro-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1099),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(p-tolyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 456),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxyphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 695),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1194),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-propoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 644),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1021),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isobutoxyphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 440),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropylphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 693),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isopropoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 504),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropoxyphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1020),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chlorophenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1100),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropylphenyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 847),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-5-isobutoxy-phenyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 1109),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isobutoxyphenyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 1094),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isobutoxy-phenyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 1266),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-propoxy-phenyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 692),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxyphenyl)-2-(4-methyl-1-piperidyl)pyridine-3-carboxamide (Compound 777),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-phenyl-pyridine-3-carboxamide (Compound 805),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(4-fluorophenyl)pyridine-3-carboxamide (Compound 1029),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(4-methoxyphenyl)pyridine-3-carboxamide (Compound 1215),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-6-(3-ethoxy-5-fluoro-phenyl)pyridine-3-carboxamide (Compound 551),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chlorophenyl)-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 459),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isobutoxy-phenyl)-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1197),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(4-isobutoxyphenyl)pyridine-3-carboxamide (Compound 563),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(4-isopropylphenyl)pyridine-3-carboxamide (Compound 1343),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-isopropoxy-phenyl)-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 788),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-propoxy-phenyl)-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 773),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(2-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 807),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-isobutoxyphenyl)pyridine-3-carboxamide (Compound 957),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-methoxy-2-pyridyl)pyridine-3-carboxamide (Compound 1141),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-6-(p-tolyl)pyridine-3-carboxamide (Compound 1078),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(4-fluoro-3-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2472),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-fluoro-4-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2587),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(2-fluoro-4-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1967),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-fluoro-2-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2199),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(m-tolyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2339),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-fluoro-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1819),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(2-fluoro-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2016),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(2-methoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2666),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-methoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2240),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3,4-difluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2229),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(4-methoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2327),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(2,4-difluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1947),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(2-ethoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2050),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(2,3-difluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2110),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-ethoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2570),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3,5-difluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1716),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(4-ethoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2525),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(2,5-difluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2489),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(2-isopropoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1897),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-isopropoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2156),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(4-isopropoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2120),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-[2-(trifluoromethoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2070),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-phenyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1725),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-[3-(trifluoromethoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2029),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(2-fluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2045),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-[4-(trifluoromethoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1763),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-fluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2414),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-[3-(trifluoromethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2220),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-[4-(trifluoromethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1900),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3,5-dimethylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1839),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3,4-dimethylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2605),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(2,3-dimethylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2225),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(2,5-dimethylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1736),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(5-fluoro-2-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2490),

2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-6-(3-ethoxy-5-fluoro-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 382),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-isobutoxyphenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 710),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(2-fluoro-5-isobutoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 681),

6-(4-chloro-3-propoxy-phenyl)-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 832),

6-(4-chloro-3-isobutoxy-phenyl)-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1234),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-methoxy-2-pyridyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1236),

6-(4-chloro-3-isopropoxy-phenyl)-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 701),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isopropoxy-phenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1285),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(4-isobutoxyphenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 1335),

2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(4-isopropylphenyl)-*N*-(1H-pyrazol-5-ylsulfonyl)pyridine-3-carboxamide (Compound 414),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-5-methyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1269),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-5-methyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 995),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-ethoxyphenyl)-5-methyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 455),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-methoxyphenyl)-5-methyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 833),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-fluorophenyl)-5-methyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 763),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)-5-methyl-pyridine-3-carboxamide (Compound 483),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)-5-methyl-pyridine-3-carboxamide (Compound 486),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-methoxy-2-methyl-phenyl)-5-methyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1287),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)-4-methoxy-pyridine-3-carboxamide (Compound 1254),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 643),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropyl-3-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 896),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-isopropyl-4-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1262),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-isopropylpyrazol-4-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1067),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-isobutylpyrazol-4-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 631),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-isopropylpyrazol-3-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1242),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[2-[isobutyl(methyl)amino]-4-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 898),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-ethoxypyrazin-2-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1252),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-isopropoxypyrazin-2-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 965)

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-isopropoxypyrimidin-5-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 844),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isobutoxy-2-methyl-3-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 616),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-fluoro-5-(2-hydroxy-2-methyl-propoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1424),

N-(benzenesulfonyl)-6-[3-fluoro-5-(3-hydroxy-2-methyl-propoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1515),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-fluoro-5-(3-hydroxy-2-methyl-propoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1567),

N-(benzenesulfonyl)-6-[3-fluoro-5-(2-hydroxy-2-methyl-propoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1581),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(5-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 400),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[5-(cyclopentoxy)-3-pyridyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 495),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-methoxy-3-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 526),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(5-isobutoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1224),,

N-[(6-amino-2-pyridyl)sulfonyl]-6-[(E)-2-cyclopropylvinyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1379),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,3-difluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2291),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-ethoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1822),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,5-difluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2105),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-ethoxy-3-fluoro-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2403),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,4-dimethylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1871),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-fluoro-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2205),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-isopropoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2623),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(5-ethoxy-2-fluoro-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2149),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-ethoxy-2-fluoro-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1847),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-ethoxy-4-fluoro-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1808),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-ethoxy-5-fluoro-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2018),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-ethoxy-4-fluoro-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2665),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-chlorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2183),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-methyl-1-naphthyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2126),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[2-(trifluoromethoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2539),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(trifluoromethoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2245),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[4-(trifluoromethoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2223),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[2-(trifluoromethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2040),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4,5-difluoro-2-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2334),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(trifluoromethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2593),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3,5-difluoro-2-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2006),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(o-tolyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1766),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[4-(trifluoromethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2048),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,3-difluoro-4-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2302),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3,5-difluoro-4-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2457),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3,4-difluoro-5-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1998),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-methoxy-2,3-dimethyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2450),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,4-difluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2447),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-methoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2164),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(1H-indol-6-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2054),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-ethoxy-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2241),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(1-methylindol-6-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2401),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-cyanophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2437),

tert-butyl 2-[5-[(2-amino-3-pyridyl)sulfonylcarbamoyl]-6-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]-2-pyridyl]indole-1-carboxylate (Compound 2657),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-fluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1790),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-ethoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1945),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-fluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1873),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-cyanophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1824),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-fluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2435),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-cyanophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2096),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-hydroxy-4-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1731),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-fluoro-2-hydroxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2446),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(1H-pyrrolo[2,3-b]pyridin-5-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1714),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(1-isopropylpyrazol-4-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2290),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-methoxy-2,6-dimethyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2607),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(1-isobutylpyrazol-4-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2231),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(6-methoxy-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2244),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-methoxy-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1980),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-methoxy-4-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2646),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[6-(dimethylamino)-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2319),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(6-methoxy-2-methyl-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1866),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(hydroxymethyl)-5-methoxy-phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2065),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(6-methoxy-5-methyl-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2564),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-isoquinolyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2418),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(5-fluoro-6-methoxy-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2628),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3,4-difluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1854),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(6-methoxy-2-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2268),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,6-dimethoxy-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2091),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3,5-difluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2272),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,3-dimethylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1713),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-chlorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2367),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3,5-dimethylisoxazol-4-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2215),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-hydroxy-2-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1999),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[4-(difluoromethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2644),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(m-tolyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1829),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(p-tolyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2132),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-fluoro-2-hydroxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2653),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(5-fluoro-2-hydroxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2046),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-fluoro-5-hydroxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2349),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,5-dimethylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1951),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,3-dihydrobenzofuran-7-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2163),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-methoxy-3-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1747),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-hydroxy-3-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2592),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-hydroxy-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2426),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-methoxy-3-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2380),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(hydroxymethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1738),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-methoxy-4-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2379),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(5-fluoro-2-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2660),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-hydroxy-3-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2158),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-methoxy-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2413),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-fluoro-4-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1851),

N-[(2-amino-3-pyridyl)sulfonyl]-6-isobutyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2150),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-fluoro-6-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2033),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-fluoro-3-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2562),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3,4-dimethoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2664),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(5-fluoro-2-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2189),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-fluoro-3-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2052),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-fluoro-5-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2474),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-fluoro-2-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2482),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-fluoro-5-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1869),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-fluoro-4-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1818),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-fluoro-2-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2355),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[4-(hydroxymethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2590),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-fluoro-3-methoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2124),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-fluoro-4-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2345),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-naphthyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2280),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(1-naphthyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1857),

N-[(2-amino-3-pyridyl)sulfonyl]-6-phenyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1958),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-chlorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2066),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3,5-dimethylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2127),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,4,5-trifluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2035),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-fluoro-4-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2613),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2522),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-isopropylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2172),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-methoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2667),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-ethoxy-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1964),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-isopropoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2515),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3,4-dimethylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2017),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-isopropoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1903),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-fluoro-2-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1944),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-isobutoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2448),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-isopropylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1991),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,6-dimethoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1994),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-isopropylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2368),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,3-dimethoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2436),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-hydroxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2257),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,5-dimethoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2195),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-hydroxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1820),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[2-(hydroxymethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1706),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-hydroxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1712),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,4-dimethoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1705),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-fluoro-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2505),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-methoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1962),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3,5-dimethoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1825),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-methoxy-2-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2494),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[2-(hydroxymethyl)-4-methoxy-phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1707),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,4-dimethoxypyrimidin-5-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2008),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-fluoro-3-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1930),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-4-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1922),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-ethoxy-2,5-dimethyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2227),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-fluoro-4-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1910),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-2-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2616),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isopropoxypyrazol-1-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2658),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2638),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,2-difluoro-1,3-benzodioxol-5-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2104),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-hydroxy-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2232),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,4-difluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2372),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,5-difluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2236),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-hydroxy-3-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2203),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-hydroxy-5-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2034),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2566),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-2-hydroxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2386),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-hydroxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2555),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-hydroxy-4-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2043),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,4-difluorophenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2559),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxyphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1969),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(trifluoromethoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1785),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,5-difluoro-4-isopropoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2608),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-ethoxy-4,5-dimethyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1931),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(trifluoromethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2392),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3,5-dimethylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1771),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-methyl-2-propoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2085),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(trifluoromethyl)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2296),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,3-dimethylphenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2287),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-isobutylpyrazol-3-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2114),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(trifluoromethoxy)phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2366),


N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(difluoromethoxy)-3-methyl-phenyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2543),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-fluoro-2-methyl-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2394),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-ethoxy-2-methyl-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 536), and

N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-methoxy-2,5-dimethyl-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1259).

PREPARATION 63: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(cyclopropylmethoxy)-3-pyridyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 789)



V-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-fluoro-3-pyridyl)pyridine-3-carboxamide (50. mg, 0.11 mmol) and cyclopropylmethanol (77 mg, 1.1 mmol) were combined in DMSO (1 mL), cooled down to 0 °C and NaH (12.75 mg, 0.5315 mmol) was added to the reaction mixture. The mixture was heated to 90 °C for 3h. The reaction mixture was filtered and then purified using a reverse phase HPLC-MS method using a Luna C18 column (50 × 21.2 mm, 5 μm particle size) sold by Phenomenex (pn: 00B-4252-P0-AX), and a dual gradient run from 1-99% mobile phase B over 15.0 minutes. Mobile phase A = H_2O (containing 5 mM HCl). Mobile phase B = CH_3CN. Flow rate = 35 mL/min, injection volume = 950 μL, and column temperature = 25 °C. The UV absorption at 254nm is used to collect fractions. The desired fractions were collected, neutralized with 20mL of $NaHCO_3$, extracted with ethyl acetate, washed with brine, dried over sodium sulfate, filtered, and concentrated to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(cyclopropylmethoxy)-3-pyridyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 789) (24 mg, 0.046 mmol, 42%). 1H NMR (400 MHz, DMSO-d_6) δ 8.78 (d, *J* = 2.4 Hz, 1H), 8.28 (dd, *J* = 8.7, 2.5 Hz, 1H), 7.60 - 7.42 (m, 2H), 7.01 (bs, 2H), 6.87 (d, *J* = 8.7 Hz, 1H), 6.51 (d, *J* = 8.6 Hz, 1H), 6.23 (bs, 2H), 4.23 (s, 2H), 4.13 (d, *J* = 7.1 Hz, 2H), 1.86 (bs, 2H), 1.56 (bs, 2H), 1.25 (dddd, *J* = 11.9, 7.3, 5.4, 2.6 Hz, 1H), 1.04 (d, *J* = 6.1 Hz, 6H), 0.59 - 0.49 (m, 2H), 0.36 - 0.26 (m, 2H). ESI-MS *m/z* calc. 522.2049, found 523.3 $(M+1)^+$; Retention time: 1.96 minutes

The following compound can be synthesized using the procedures described herein:

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2,2-dimethylpropoxy)-3-pyridyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1270),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isobutoxy-4-methyl-2-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1656),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-ethoxy-4-methyl-2-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1512),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-4-methyl-2-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1637),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(difluoromethoxy)-3-pyridyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1679),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isobutoxy-2-methyl-3-pyridyl)pyridine-3-carboxamide (Compound 1537),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[2-(2-ethoxyethoxy)ethyl-methyl-amino]-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2684),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[2-ethoxyethyl(methyl)amino]-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2685),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2-ethoxyethylamino)-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2686),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-2-methyl-3-pyridyl)pyridine-3-carboxamide (Compound 1623),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-4-methyl-3-pyridyl)pyridine-3-carboxamide (Compound 1514),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-[6-(isobutylamino)-3-pyridyl]pyridine-3-carboxamide (Compound 1398),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[[(1S)-1,3-dimethylbutyl]amino]-3-pyridyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1463),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-[6-[isobutyl(methyl)amino]-3-pyridyl]pyridine-3-carboxamide (Compound 1506),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-6-[6-(1-ethylpropylamino)-3-pyridyl]pyridine-3-carboxamide (Compound 1397),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[[(1S)-1,2-dimethylpropyl]amino]-3-pyridyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1399),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[isopropyl(methyl)amino]-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2687),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(isopropylamino)-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2688),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[ethyl(methyl)amino]-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2689),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[[(1R)-1,2-dimethylpropyl]amino]-3-pyridyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1442),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[[(1R)-1,3-dimethylbutyl]amino]-3-pyridyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1527),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(4-isobutoxy-2-pyridyl)pyridine-3-carboxamide (Compound 389),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isobutoxy-2-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 397),

methyl 2-[[5-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-2-pyridyl]-2-pyridyl]oxy]propanoate (Compound 723),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isobutoxypyrazin-2-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1523),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[isobutyl(methyl)amino]-3-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 412),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isobutoxy-2-pyridyl)pyridine-3-carboxamide (Compound 432),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(isopropylamino)-3-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 437),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isobutoxy-3-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 471),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isobutoxy-3-pyridyl)pyridine-3-carboxamide (Compound 544),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(isobutylamino)-3-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 569),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(1-ethylpropoxy)-2-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 580),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-2-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 708),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(1,2-dimethylpropoxy)-2-pyridyl]-2-(2,2-dimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 717),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2,2-dimethylpropoxy)pyrazin-2-yl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 779),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(diethylamino)-3-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 796),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-2-methyl-3-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 814),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[isobutyl(methyl)amino]-2-methyl-3-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 827),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(1,3-dimethylbutoxy)-2-pyridyl]-2-(2,2-dimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 846),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopentyloxy-3-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 848),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isobutoxy-4-methyl-3-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 851),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-4-methyl-3-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 905),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(isopropylamino)-2-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 951),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isobutoxypyrazin-2-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 988),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethylpyrrolidin-1-yl)-6-(6-isobutoxy-2-pyridyl)pyridine-3-carboxamide (Compound 1011),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(isobutylamino)-2-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1059),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2,2-dimethylpropoxy)-2-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1062),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(1-ethylpropoxy)-3-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1076),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopentyloxy-2-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1135),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2,2-dimethylpropoxy)-4-methyl-3-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1160),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2,2-dimethylpropoxy)-3-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1181),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2,2-dimethylpropoxy)-2-pyridyl]-2-(2,2-dimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1328),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(diethylamino)-2-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1345),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[2-(1,3-dimethylbutoxy)-4-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 384),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[2-(isobutylamino)-4-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 523),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-propoxy-4-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 491),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-isobutoxy-4-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1192),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-isopropoxy-4-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 402),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-isobutoxy-4-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1618),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2,2-dimethylpropoxy)-3-methyl-2-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1645),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isobutoxy-3-methyl-2-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1659),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2,2-dimethylpropoxy)-2-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1663),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2,2-dimethylpropoxy)-5-methyl-2-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1610),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-ethoxy-5-methyl-2-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1363),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(diethylamino)-3-methyl-2-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1383),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(isobutylamino)-3-methyl-2-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1417),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2,2-dimethylpropoxy)-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1504),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(cyclobutoxy)-3-pyridyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 428),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-[6-(2-hydroxy-1-methyl-ethoxy)-3-pyridyl]pyridine-3-carboxamide (Compound 998),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[(1R)-2-benzyloxy-1-methyl-ethoxy]-3-pyridyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 576),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-ethoxy-5-methyl-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1727),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(diethylamino)-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1915),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[2-[2-(2-azidoethoxy)ethoxy]ethoxy]-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isobutoxy-5-methyl-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1741),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[isobutyl(methyl)amino]-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2279),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[5-methyl-6-[(1R)-1-methylpropoxy]-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2576),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-isopropoxy-5-methyl-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2170),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[2-(2-ethoxyethoxy)ethoxy]-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1715),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[5-methyl-6-[(1S)-1-methylpropoxy]-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1728),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[2-[2-(2-isopropoxyethoxy)ethoxy]ethoxy]-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1787),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[2-(3-but-3-ynyldiazirin-3-yl)ethoxy]-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2-ethoxyethoxy)-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2404),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2-isopropoxyethoxy)-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1973),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2-methoxyethoxy)-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2582),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[2-(2-ethoxyethoxy)-6-methyl-4-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1948),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[2-(2-isopropoxyethoxy)-6-methyl-4-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2079),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[2-(diethylamino)-6-methyl-4-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1739),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-isobutoxy-6-methyl-4-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2010),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[2-[2-(2-ethoxyethoxy)ethoxy]-6-methyl-4-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1812),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-ethoxy-6-methyl-4-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2364),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-isopropoxy-6-methyl-4-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2306),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2-methoxy-6-methyl-4-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1748),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(6-isopropoxy-5-methyl-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1702),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[6-(2-ethoxyethoxy)-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2497),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(6-ethoxy-5-methyl-3-pyridyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2651),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[6-[2-(2-ethoxyethoxy)ethoxy]-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1783),


N-[(2-amino-3-pyridyl)sulfonyl]-6-[6-(2-isopropoxyethoxy)-5-methyl-3-pyridyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1695),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-[6-[(1-methylcyclopropyl)methoxy]-3-pyridyl]pyridine-3-carboxamide (Compound 454), and

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(cyclopropoxy)-3-pyridyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1125).

PREPARATION 64: 1-Methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxylic acid


Step 1: 2-Benzyl 1-(*tert*-butyl) 1-methylisoindoline-1,2-dicarboxylate



A solution of *tert*-butyl 2-[benzyloxycarbonyl-[(2-bromophenyl)methyl]amino]propanoate (2.69 g, 6.00 mmol) in dioxane (30 mL) was purged with nitrogen for 2 minutes. 2-(2-Diphenylphosphanylphenyl)-*N*,*N*-dimethyl-aniline (228.9 mg,


0.6000 mmol) was added under nitrogen followed by the addition of tris(dibenzylideneacetone)dipalladium(0) (274.7 mg, 0.3000 mmol). Lithium *tert*-butoxide (960.7 mg, 12.00 mmol) was added to the reaction mixture. The mixture was heated to 90 °C for 16 hours, diluted with ethyl acetate and filtered. The residue was purified by silica gel column chromatography (0-10% ethyl acetate in hexanes) to provide 2-benzyl 1-(*tert*-butyl) 1-methylisoindoline-1,2-dicarboxylate (1.4 g, 3.8 mmol, 63%). ESI-MS *m/z* calc. 367.17834, found 368.4 $(M+1)^+$; Retention time: 2.75 minutes.

Step 2: *tert*-butyl 1-methylisoindoline-1-carboxylate



A solution of 2-benzyl 1-(*tert*-butyl) 1-methylisoindoline-1,2-dicarboxylate (718 mg, 1.95 mmol) in methanol (50 mL) was purged with nitrogen for 2 minutes. Palladium on carbon (10% w/w, 104.0 mg, 0.09770 mmol) was added and the mixture was placed under an atmosphere of hydrogen for 16 hours. The catalyst was removed via filtration through celite and the filtrate was concentrated to dryness to provide *tert*-butyl 1-methylisoindoline-1-carboxylate (400. mg, 1.71 mmol, 87.7%). The crude material was used directly in next step without further purification.


Step 3: 1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxylic acid



To a solution of *tert*-butyl 1-methylisoindoline-1-carboxylate (1.5 g, 6.4 mmol) in acetonitrile (75.00 mL) was added 2-(chloromethyl)-1,3,5-trimethyl-benzene (1.084 g, 6.429


mmol) and cesium carbonate (4.190 g, 12.86 mmol). The mixture was heated at 90 °C overnight, and then partitioned between ethyl acetate and water. The aqueous layer was extracted with ethyl acetate. The combined organic layers were washed with brine, dried over magnesium sulate, filtered, and concentrated to dryness. The crude material was purified by silica column chromatography utilizing a gradient of 0-5% ethyl acetate in hexanes to provide *tert*-butyl 1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxylate. This material was then dissolved in dioxane (3.2 mL) and hydrochloric acid in dioxane (5 mL of 4 M, 20.00 mmol) was added to the mixture. The mixture was stirred for 16 hours at room temperature. The mixture was then stirred at 70 °C for 1 hour. The reaction mixture was cooled to room temperature, the precipitate was filtered and washed with dioxane to give an off-white solid. 1-methyl-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxylic acid (0.98 g, 3.2 mmol, 49%) ESI-MS *m/z* calc. 309.17288, found 310.5 $(M+1)^+$; Retention time: 1.18 minutes (3 min run).

PREPARATION 65: Ethyl 5-methyl-1-(*p*-tolyl)-1*H*-indole-2-carboxylate



A mixture of ethyl 5-methyl-1*H*-indole-2-carboxylate (2.032 g, 10.00 mmol), 1-iodo-4-methyl-benzene (2.180 g, 10.00 mmol), iodocopper (95.22 mg, 0.5000 mmol), *N*1,*N*2-dimethylcyclohexane-1,2-diamine (284.5 mg, 315.4 µL, 2.000 mmol), and potassium phosphate (4.458 g, 21.00 mmol) in toluene (50.00 mL) was heated to reflux for 22 hours. The cooled reaciton was filtered and evaporated. The green residue was purified by silica gel chromatography with 0-15% ethyl acetate in hexanes to give ethyl 5-methyl-1-(*p*-tolyl)-1*H*-indole-2-carboxylate (1.43 g, 4.875 mmol, 48.74%) as a colorless oil. ESI-MS *m/z* calc. 293.14, found 294.2 $(M+1)^+$; Retention time: 0.83 minutes (1 minute run).


PREPARATION 66: *N*-[(6-amino-2-pyridyl)sulfonyl]-5-bromo-2-[(2*S*,5*R*)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 390)



N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2*S*,5*R*)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (147.4 mg, 0.2887 mmol) in acetic acid (1.5mL) was treated with bromine (69.20 mg, 22.31 µL, 0.4330 mmol) and stirred at room temperature for 1 hour. The resulting yellow suspension was diluted with water (20 mL), quenched to pH 2-3 by addition of saturated aqueous sodium bicarbonate and extracted with dichloromethane. The crude material was purified by chromatography over silica gel (12g) with a linear gradient of 0 to 5% methanol in dichloromethane to give *N*-[(6-amino-2-pyridyl)sulfonyl]-5-bromo-2-[(2*S*,5*R*)-2,5-dimethylpyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 390) (90. mg, 0.14 mmol, 50.%) as a yellow foam. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.80 (s, 1H), 8.52 (dd, J = 2.5, 0.7 Hz, 1H), 8.02 (dd, J = 8.7, 2.5 Hz, 1H), 7.85 (s, 1H), 7.64 (t, J = 7.8 Hz, 1H), 7.23 (d, J = 7.2 Hz, 1H), 6.83 (dd, J = 8.6, 0.7 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 6.51 (s, 2H), 5.32 (p, J = 6.2 Hz, 1H), 3.95 (s, 2H), 1.97 - 1.82 (m, 2H), 1.64 (dd, J = 8.9, 5.3 Hz, 2H), 1.33 (d, J = 6.2 Hz, 6H), 1.04 (d, J = 6.1 Hz, 6H). ESI-MS *m/z* calc. 588.1154, found 589.0 $(M+1)^+$; Retention time: 2.25 minutes.

PREPARATION 67: 3-(*tert*-butyl)-4-methoxy-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxylic acid


Step 1: 3-(*tert*-butyl)-*N*-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide



In a 20 mL sealed flask containing 5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylic acid 1 (500. mg, 1.66 mmol) was added *N*-

methylpyrrolidinone (5.000 mL) and diisopropylamine (537.7 mg, 724.7 µL, 4.160 mmol) followed by a propylphosphonic anhydride solution (2.118 mL of 50 %w/v, 3.328 mmol) and 2,3,5,6-tetrafluoro-4-(trifluoromethyl)aniline (426.5 mg, 1.830 mmol) and the resulting mixture was stirred at 80 °C for 16 hours. The material was diluted with ethyl acetate (10 mL), washed with water and the organic layer was extracted. The organic layer was washed twice with a saturated aqueous solution of sodium bicarbonate, followed by brine. The material was dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. The resulting crude material was adsorbed to celite and purified by silica gel column chromatography ((40 g) using a gradient of 0 to 50% ethyl acetate) to afford 5-*tert*-butyl-*N*-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenyl]-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (748 mg, 1.44 mmol, 86.4%). ESI-MS *m/z* calc. 515.1808, found 516.2 $(M+1)^+$; Retention time: 2.58 minutes. ^{1}H NMR (400 MHz, DMSO) δ 10.92 (s, 1H), 7.01 (s, 1H), 6.83 (s, 2H), 5.64 (s, 2H), 2.27 (s, 6H), 2.21 (s, 3H), 1.20 (s, 9H).


Step 2: 3-(*tert*-butyl)-4-methoxy-*N*-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxamide



5-*tert*-Butyl-*N*-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenyl]-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (200. mg, 0.388 mmol), anhydrous methanol (1.554 g, 1.965 mL, 48.50 mmol), xylene (2.000 mL), diacetoxypalladium (8.711 mg, 0.03880 mmol), (diacetoxyiodo)benzene (374.9 mg, 1.164 mmol), and 4A molecular sieves were combined in a capped 10 mL vial. The mixture was heated in an oil bath to 100 °C for 40 hours. The mixture was allowed to cool was diluted with water, extracted three times with dichloromethane (10 mL), and the combined organics were washed with a saturated aqueous sodium bicarbonate solution and then a saturated aqueous solution of sodium chloride. The organic layer was dried over sodium sulfate and then purified by silica gel column chromatography (24 g) utilizing a gradient of 0 to 50% ethyl acetate in hexanes to afford 5-*tert*-


butyl-4-methoxy-*N*-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenyl]-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (161 mg, 0.295 mmol, 76.1%) as an off-white solid. ESI-MS *m/z* calc. 545.19135, found 546.2 $(M+1)^+$; Retention time: 2.4 minutes. ^{1}H NMR (400 MHz, DMSO) δ 10.76 (s, 1H), 6.83 (s, 2H), 5.39 (s, 2H), 3.79 (s, 3H), 2.28 (s, 6H), 2.20 (s, 3H), 1.25 (s, 9 H).

Step 3: Methyl 3-(*tert*-butyl)-4-methoxy-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxylate



5-*tert*-butyl-4-methoxy-*N*-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenyl]-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (161 mg, 0.295 mmol) was dissolved in anhydrous tetrahydrofuran (3.220 mL) and to the mixture was added (bis(trimethylsilyl)amino)lithium (442.7 µL of 1 M in tetrahydrofuran, 0.4427 mmol) at 0 °C and the resulting mixture was allowed to stir for 10 minutes. Then methyl chloroformate (83.66 mg, 68.41 µL, 0.8853 mmol) was added and the reaction mixture was allowed to stir for one hour, and then concentrated under reduced pressure. The residue was redissolved in anhydrous tetrahydrofuran (3.220 mL) and then the mixture was cooled to 0 °C in an ice water bath. Sodium methoxide (319.0 µL of 25 %w/v in methanol, 1.476 mmol) was added and the reaction mixture was stirred for 10 minutes. The mixture was diluted with water, extracted three times with ethyl acetate (10mL) and the combined organics were washed with brine. The mixture was dried over sodium sulfate, and then purified by reverse-phase preparative chromatography utilizing a gradient from 10 to 99% acetonitrile in water containing 5 mM aqueous hydrochloric acid to yield methyl 3-(*tert*-butyl)-4-methoxy-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxylate (25.8 mg, 0.0749 mmol, 25.4%) as an off-white solid. ESI-MS *m/z* calc. 344.21, found 345.2 $(M+1)^+$; Retention time: 1.92 minutes.


Step 4: 3-(*tert*-butyl)-4-methoxy-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxylic acid



To methyl 3-(*tert*-butyl)-4-methoxy-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxylate (30. mg, 0.087 mmol) in anhydrous tetrahydrofuran (600.0 µL) was added sodium hydroxide (108.9 µL of 4 M, 0.4355 mmol), the mixture was heated to 70 °, C and stirred for 3 hours. The mixture was allowed to cool and then it was purified directly by reverse-phase preparative chromatography utilizing a gradient from 10 to 99% acetonitrile in water containing 5 mM aqueous hydrochloric acid to yield 3-(*tert*-butyl)-4-methoxy-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxylic acid (27.3 mg, 0.0826 mmol, 95%) as an off- white solid. ESI-MS *m/z* calc. 330.19434, found 331.2 $(M+1)^+$; Retention time: 2.08 minutes.

PREPARATION 68: *N*-[amino-(6-amino-2-pyridyl)-oxo-$l^{6}-sulfanylidene]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1602) (Compound 1481) (Compound 515)


Step 1: 2-fluoro-6-[(6-fluoropyridin-2-yl)disulfanyl]pyridine



A mixture of 2-fluoro-6-hydroxypyridine (1.13 g, 10.0 mmol) and phosphorus pentasulfide (4.44 g, 20.0 mmol) in toluene (50 mL) was refluxed for 18 hours. Once cooled, the flask was placed in an ice bath and quenched with water (50 mL) and then basified to a pH of about 8-9 with 25% NaOH. The mixture was filtered over celite to remove some solids that were still present and washed with water (50 mL) and ethyl acetate (200 mL). The biphasic mixture was transferred to a 1.0-L separatory funnel, extracted and the layers were separated. The aqueous layer was extracted again with ethyl acetate (100 mL), and the organic layers were then


combined, washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by silica-gel column chromatography, eluting from 0% to 10% ethyl acetate in heptanes to afford 2-fluoro-6-[(6-fluoropyridin-2-yl)disulfanyl]pyridine (304 mg, 23.7% yield) as a colorless oil. ^{1}H NMR (300 MHz, $CDCl_3$): ppm 6.75 (ddd, J = 8.0, 2.7, 0.5 Hz, 1 H), 7.49 (ddd, J = 7.7, 2.1, 0.6 Hz, 1 H), 7.72 (q, J = 7.9 Hz, 1 H). $[M+H]^+$ = 257.0.

Step 2: 6-*tert*-butyl-*N*-[(6-fluoropyridin-2-yl)sulfinyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide




To a solution of 2-fluoro-6-[(6-fluoropyridin-2-yl)disulfanyl]pyridine (744 mg, 2.90 mmol) and 6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (906 mg, 2.90 mmol) in pyridine (10 mL) and acetonitrile (10 mL) was added bromine (0.172 mL, 3.36 mmol) dropwise at 0 °C. After complete addition, the mixture was stirred at room temperature overnight. Additional bromine (0.087 mL, 1.7 mmol) was added at room temperature and the mixture was stirred for an additional hour. Solvent was then removed under reduced pressure and pyridine was further co-evaporated with toluene (10 mL) two times under reduced pressure. The residue was purified by silica gel column chromatography (24g) column, eluting from 0% to 25% ethyl acetate in heptane to afford 6-*tert*-butyl-*N*-[(6-fluoropyridin-2-yl)sulfinyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (850 mg, 64% yield) as an off-white solid. ^{1}H NMR (300 MHz, $CDCl_3$): ppm 1.08 (s, 9 H), 2.00 (s, 6 H), 2.29 (s, 3 H), 6.86 (s, 2 H), 7.03 – 7.89 (m, 1 H), 7.12 (d, J = 8.0, 1 H), 7.99 – 8.11 (m, 2 H), 8.53 (d, J = 8.0 Hz, 1 H). ^{19}F NMR (282 MHz, $CDCl_3$): ppm -64.8 (s, 1 F). $[M+H]^+$ = 456.1.

Step 3: 6-(*tert*-butyl)-*N*-(6-fluoropyridine-2-sulfonimidoyl)-2-(mesityloxy)nicotinamide



To a solution of 6-*tert*-butyl-*N*-[(6-fluoropyridin-2-yl)sulfinyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (419 mg, 0.920 mmol) in dichloromethane (10 mL) was added *tert*-butyl hypochlorite (94.8 µL, 0.87 mmol) at 0 °C. The mixture was stirred at this temperature for 2 days. The solvent was then removed under reduced pressure at below 21 °C. The crude product was used directly in the next step without further purification. The residue was dissolved in acetonitrile (11 mL) then hexamethyldisilazane (1.09 mL, 5.22 mmol) was added at 0 °C. The mixture was stirred for 1 hour at room temperature then quenched by the addition of water (3 mL). The mixture was extracted with ethyl acetate (2 x 20 mL), dried over sodium sulfate, filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography column (24g), eluting from 0% to 5% methanol in dichloromethane, to afford 6-(*tert*-butyl)-*N*-(6-fluoropyridine-2-sulfonimidoyl)-2-(mesityloxy)nicotinamide (286 mg, 70% yield) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$): ppm 1.08 (s, 9 H), 2.03 (s, 6 H), 2.29 (s, 3 H), 6.84 (s, 2 H), 6.97 (d, J = 8.0 Hz, 1 H), 7.16 (ddd, J = 8.0, 2.6, 0.8 Hz, 1 H), 7.94 – 8.11 (m, 2 H), 8.35 (d, J = 8.0 Hz, 1 H). ^{19}F NMR (282 MHz, $CDCl_3$): ppm -64.1 (s, 1 F). $[M+H]^+$ = 471.1.


Step 4: 6-(*tert*-butyl)-*N*-(6-((2,4-dimethoxybenzyl)amino)pyridine-2-sulfonimidoyl)-2-(mesityloxy)nicotinamide



A mixture of 6-(*tert*-butyl)-*N*-(6-((2,4-dimethoxybenzyl)amino)pyridine-2-sulfonimidoyl)-2-(mesityloxy)nicotinamide (400. mg, 0.850 mmol), 2,4-dimethoxybenzylamine

(0.25 mL, 1.7 mmol) and triethylamine (0.24 mL, 1.7 mmol) in dimethylsulfoxide (10 mL) was stirred at 100 °C overnight. After completion, the mixture was diluted with ethyl acetate (20 mL), washed with brine (2 x 10 mL), dried over sodium sulfate, filtered, and concentrated under reduced pressure. The residue was purified by reverse-phase column chromatography on a 25g column, eluting from 0% to 100% acetonitrile in water containing 0.1% formic acid to afford 6-(*tert*-butyl)-*N*-(6-((2,4-dimethoxybenzyl)amino)pyridine-2-sulfonimidoyl)-2-(mesityloxy)nicotinamide (426 mg, 81.1% yield) as a yellow foam. ^{1}H NMR (300 MHz, $CDCl_3$): ppm 1.07 (s, 9 H), 2.03 (s, 6 H), 2.27 (s, 3 H), 3.77 (s, 3 H), 3.82 (s, 3 H), 4.42 (s, 2 H), 5.18 – 5.35 (br. s., 1 H), 6.36 (dd, J = 8.3, 2.3 Hz, 1 H), 6.45 (d, J = 2.3 Hz, 1 H), 6.52 (d, J = 8.3 Hz, 1 H), 6.81 (s, 2 H), 6.94 (d, J = 7.9 Hz, 1 H), 7.18 (d, J = 8.3 Hz, 1 H), 7.28 (d, J = 7.4 Hz, 1 H), 7.45 (dd, J = 8.5, 7.4 Hz, 1 H), 8.38 (d, J = 7.9 Hz, 1 H). $[M+H]^+$ = 618.2.


Step 5: *N*-(6-aminopyridine-2-sulfonimidoyl)-6-(*tert*-butyl)-2-(mesityloxy)nicotinamide



To a stirred solution of *N*-[amino(6-{[(2,4-dimethoxyphenyl)methyl]amino}pyridin-2-yl)oxo-λ^6-sulfanylidene]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (420 mg, 0.68 mmol) in dichloromethane (50 mL) was added trifluoroacetic acid (12 mL) and the mixture was stirred until LCMS indicated completion. The reaction mixture was diluted with dichloromethane (20 mL) and water (20 mL) and the pH was adjusted to greater than 6 by adding solid $NaHCO_3$. The mixture was extracted with dichloromethane (2 x 20 mL) and the combined organic layers were dried over sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by reverse-phase column chromatography on a 25g column, eluting from 0% to 100% acetonitrile in water containing 0.1% formic acid to afford *N*-(6-aminopyridine-2-sulfonimidoyl)-6-(*tert*-butyl)-2-(mesityloxy)nicotinamide (90. mg, 28% yield) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$ + 5 drops of DMSO-d_6) ppm 0.96 (s, 9 H), 1.90 (s,


6 H), 2.17 (s, 3 H), 5.78 (br. s., 2 H), 6.59 (d, J = 8.2 Hz, 1 H), 6.71 (s, 2 H), 6.82 (d, J = 7.8 Hz, 1 H), 7.19 - 7.31 (m, 3 H), 7.36 - 7.47 (m, 1 H), 8.15 (d, J = 7.8 Hz, 1 H). $[M+H]^+$ = 468.1.

Step 6: *N*-[(*S*)-amino(6-aminopyridin-2-yl)oxo-λ^6-sulfanylidene]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide and *N*-[(*R*)-amino(6-aminopyridin-2-yl)oxo-λ^6-sulfanylidene]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide



N-(6-aminopyridine-2-sulfonimidoyl)-6-(*tert*-butyl)-2-(mesityloxy)nicotinamide was dissolved in a methanol and dimethylsulfoxide mixture (4:1 (v/v)) to an approximate concentration of 16 mg/mL. The solution was subjected to chiral SFC (70 µL injections; column: Chiralpak OD-H (250 x 21.2 mm),5µm; mobile phase: 20% methanol, 80% CO_2; flow: 10.0 mL/min) to give *N*-[(*S*)-amino(6-aminopyridin-2-yl)oxo-λ^6-sulfanylidene]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (peak 1, enantiomer 1, 98% ee), and *N*-[(*R*)-amino(6-aminopyridin-2-yl)oxo-λ^6-sulfanylidene]-6-*tert*-butyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (peak 2, enantiomer 2, 97.3% ee). Absolute stereochemistry is inknown.

PREPARATION 69: (*R*)-1-(2,4,6-trimethylbenzyl)indoline-2-carboxylic acid



A suspension of (*R*)-indoline-2-carboxylic acid (300. mg, 1.84 mmol) and potassium hydroxide (309.5 mg, 5.517 mmol) was stirred in isopropanol (2.010 mL) for 20 minutes. The reaction mixture was stirred for 20 minutes and was treated with 2-(chloromethyl)-1,3,5-trimethyl-benzene (341.2 mg, 2.023 mmol). The reaction mixture was stirred overnight. The

reaction mixture was diluted with ethyl acetate and water. The mixture was extracted with ethyl acetate and the aqueous layer was acidified to pH 5 and extracted again with ethyl acetate. The combined organics were washed with water, brine, dried over sodium sulfate, filtered and concentrated to yield (*R*)-1-(2,4,6-trimethylbenzyl)indoline-2-carboxylic acid (220. mg, 0.745 mmol, 40.5%) as a white solid. ESI-MS *m/z* calc. 295.15723, found 296.5 $(M+1)^+$; Retention time: 1.79 minutes.

PREPARATION 70: *tert*-butyl 2,3,4,5-tetrahydro-1*H*-1-benzazepine-2-carboxylate

Step 1: 3-(2-bromophenyl)propan-1-ol



A solution of 3-(2-bromophenyl)propanoic acid (1.0 g, 4.4 mmol) in anhydrous tetrahydrofuran (5.0 mL) was treated with borane-tetrahydrofuran complex in tetrahydrofuran (1.0 M, 7.0 mL, 7.0 mmol) at 0 °C over a period of 15 minutes. The mixture was stirred at 0 °C for an additional 1 hour. After completion, the residual borane was quenched by the cautious addition of water (1 mL) and the solvent was removed under vacuum. The resulting residue was dissolved in ether (20 mL) and the organic layer was washed with water (20 mL), brine (20 mL), dried over sodium sulfate, filtered, and concentrated under pressure to afford 3-(2-bromophenyl)propan-1-ol as a colorless oil (938 mg, quantitative yield). This compound was used in the following step without further purification.


[001864] Step 2: Synthesis of 1-bromo-2-(3-bromopropyl)benzene



To a solution of 3-(2-bromophenyl)propan-1-ol (938 mg, 4.36 mmol) and triphenylphosphine (2.29 g, 8.72 mmol) in anhydrous dichloromethane (25 mL) was added *N*-bromosuccinimide (1.63 g, 9.16 mmol) at 0 °C and the mixture was stirred at room temperature for 2 hours. After completion, the reaction was quenched by the addition of water (20 mL) and the organic layer was separated and dried over sodium sulfate, filtered and concentrated under pressure. The residue was purified by silica gel chromatography (40 g), eluting from 0% to 10%


ethyl acetate in heptanes to afford 1-bromo-2-(3-bromopropyl)benzene as a yellow oil (830 mg, 68.5% yield). ^{1}H NMR (300 MHz, $CDCl_3$) ppm 2.13 – 2.24 (m, 2 H), 2.90 (t, J = 7.7 Hz, 2 H), 3.43 (t, J = 6.5 Hz, 2 H), 7.04 – 7.12 (m, 1H), 7.20 – 7.28 (m, 2 H), 7.54 (d, J = 7.7 Hz, 1 H). $[M+H]^+$ = 279.1.

Step 3: *tert*-butyl 2-amino-5-(2-bromophenyl)pentanoate




To a suspension of potassium hydroxide (1.27 g, 22.6 mmol) in dimethylsulfoxide (10 mL) cooled in a cold water bath was added *N*-(diphenylmethylene)glycine *tert*-butyl ester (2.23 g, 7.55 mmol). Then, 1-bromo-2-(3-bromopropyl)benzene (938 mg, 4.36 mmol) in dimethyl sulfoxide (10 mL) was added dropwise and the mixture was stirred at room temperature for 1 hour. After completion, the mixture was diluted with ethyl acetate (100 mL) and the solution was filtered. The mixture was washed with water (3 x 50 mL), brine, dried over sodium sulfate, filtered, and concentrated under reduced pressure. After drying under high vacuum, the residue was dissolved in tetrahydrofuran (78 mL). To this solution was added 1 N HCl (75.7 mL, 75.7 mmol) dropwise at 0 °C and the mixture was stirred at this temperature for an additional 2 hours. The resulting mixture was then neutralized by addition of solid sodium bicarbonate and extracted with dichloromethane (100 mL). The organic layer was dried over sodium sulfate, filtered and concentrated under pressure. The residue was purified by silica gel chromatography (80 g) eluting from 0% to 10% ethyl acetate in heptanes to afford *tert*-butyl 2-amino-5-(2-bromophenyl)pentanoate as a yellow oil (2.21 g, 86.4% yield over two steps). ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.45 (s, 9 H), 1.60 – 1.81 (m, 2 H), 2.70 – 2.78 (m, 2 H), 3.33 – 3.39 (m, 2 H), 7.00 – 7.09 (m, 1 H), 7.18 – 7.24 (m, 2 H), 7.51 (d, J = 8.1 Hz, 1 H). $[M+H]^+$ = 328.1.

Step 4: *tert*-butyl 2,3,4,5-tetrahydro-1*H*-1-benzazepine-2-carboxylate



A mixture of *tert*-butyl 2-amino-5-(2-bromophenyl)pentanoate (1.7 g, 5.2 mmol), palladium acetate (116 mg, 0.518 mmol), triphenylphosphine (408 mg, 1.55 mmol), and cesium carbonate (3.4 g, 10. mmol) in toluene (80 mL) was stirred at 110 °C overnight. After completion, the mixture was diluted with ethyl acetate (100 mL) and washed with water (50 mL), brine (50 mL), dried over sodium sulfate, filtered, and concentrated under reduced pressure. The residue was purified by silica gel chromatography (80 g) eluting from 0% to 20% ethyl acetate in heptanes to afford *tert*-butyl 2,3,4,5-tetrahydro-1*H*-1-benzazepine-2-carboxylate as a yellow oil (527 mg, 41.0%). ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.46-1.57 (m, 10 H), 1.66-1.82 (m, 1 H), 1.94-2.08 (m, 1 H), 2.24-2.36 (m, 1 H), 2.65-2.77 (m, 1 H), 2.78-2.91 (m, 1 H), 3.44 (dd, J = 10.7, 2.3 Hz, 1 H), 4.50 (br. s, 1 H), 6.79-6.88 (m,2 H), 7.01-7.11 (m, 2 H). $[M+H]^+$ = 248.2

PREPARATION 71: 5-ethyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[2,3-b]pyridine-2-carboxylate




A solution of methyl 5-bromo-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[2,3-b]pyridine-2-carboxylate (150. mg, 0.387 mmol),1,2-dimethoxyethane (1.35 mL) 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (119.3 mg, 0.7746 mmol), and potassium carbonate (187.4 mg, 1.356 mmol) in 1,2-dimethoxyethane (1.35 mL) and water (0.45 mL) was degassed and put under nitrogen. The reaction mixture was treated with tetrakis(triphenylphosphine)palladium (0) (44.75 mg, 0.03873 mmol) and irradiated in a microwave reactor at 120 °C for 30 minutes. The mixture was diluted with water and extracted three times with ethyl acetate. The combined organics were washed with brine, dried oer sodium sulfate, filtered, and concentrated under reduced pressure. The resulting crude methyl 1-[(2,4,6-trimethylphenyl)methyl]-5-vinyl-pyrrolo[2,3-b]pyridine-2-carboxylate was dissolved in methanol (12.45 mL) and put under a nitrogen atmosphere before being treated with wet 10% palladium on carbon (41.22 mg, 0.03873 mmol). The mixture was evacuated and put under a

hydrogen atmosphere for 16 hours. The reaction mixture was put under a nitrogen atmosphere and filtered through celite to afford the crude product which was purified on 12 g of silica gel utilizing a gradient of 0-40% ethyl acetate in hexanes to yield methyl 5-ethyl-1-[(2,4,6-trimethylphenyl)methyl]pyrrolo[2,3-b]pyridine-2-carboxylate ESI-MS *m/z* calc. 336.18378, found 337.5 $(M+1)^+$; Retention time: 2.24 minutes.


PREPARATION 72:

Step 1: Synthesis of 2-(2-bromophenyl)propan-2-amine




Methylmagnesium bromide (54.9 mL, 164.8 mmol, 3M in diethyl ether) was added to a solution of 2-bromobenzonitrile (10.0 g, 54.9 mmol) in diethyl ether (200 mL) and the reaction mixture was stirred at room temperature under nitrogen. After 30 minutes, titanium isopropoxide (16.3 mL, 54.9 mmol) was added and resulting mixture was refluxed overnight. Once cooled to 0 °C, 2N NaOH (400 mL) was added and resulting mixture was stirred at room temperature for 30 minutes. The solution was diluted with saturated aqueous sodium bicarbonate (400 mL), extracted using methyl *tert*-butyl ether (3 x 200 mL). The organic layers were combined and concentrated under reduced pressure. The residue was dissolved in 1N hydrochloric acid (75 mL) and washed with methyl *tert*-butyl ether (150 mL). The aqueous layer was basified to pH 10-11 using 2N sodium hydroxide and the resulting solution was extracted with methyl *tert*-butyl ether (3 x 200 mL). The organic layers were combined, dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford 2-(2-bromophenyl)propan-2-amine (9.0 g, 77% yield) as orange oil. ^{1}H NMR (300 MHz, $CDCl_3$): ppm 1.66 (s, 6 H), 7.02 – 7.10 (m, 1 H), 7.13 – 7.43 (m, 1 H), 7.55 – 7.61 (m, 2 H). $[M+H]^+$ = 214.0.

Step 2: Synthesis of *tert*-butyl 2-{[2-(2-bromophenyl)propan-2-yl]amino}acetate




tert-Butyl 2-bromoacetate (6.24 mL, 42.2 mmol) was slowly added to a solution of 2-(2-bromophenyl)propan-2-amine (9.04 g, 42.22 mmol) and potassium carbonate (14.6 g, 105.6 mmol) in dry tetrahydrofuran (100 mL) at 0 °C and reaction mixture was stirred at 50-55 °C for 16 hours. Once cooled to room temperature, reaction mixture was diluted with brine (100 mL) and extracted using ethyl acetate (3 x 100 mL). The organic layers were combined, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel chromatography, eluting from 0% to 20% ethyl acetate in heptanes to afford *tert*-butyl 2-{[2-(2-bromophenyl)propan-2-yl]amino}acetate (4.5 g, 32% yield) as a yellow oil. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.43 (s, 9 H), 1.63 (s, 6 H), 2.94 (s, 2 H), 7.09 (dt, J = 7.6, 1.7 Hz, 1 H), 7.22 – 7.31 (m, 1 H), 7.43 (dd, J = 7.9, 1.6 Hz, 1 H), 7.58 (dd, J = 7.9, 1.4 Hz, 1 H). $[M+H]^+$ = 328.1.

Step 3: *tert*-butyl 2-{[(benzyloxy)carbonyl][2-(2-bromophenyl)propan-2-yl]amino}acetate




Diisopropylethylamine (5.48 mL, 31.4 mmol) was added to a solution of *tert*-butyl 2-{[2-(2-bromophenyl)propan-2-yl]amino}acetate (4.48 g, 13.7 mmol) in dichloromethane (50 mL) and solution was cooled to 0 °C. Benzyl chloroformate (3.51 mL, 24.6 mmol) was added dropwise at 0 °C and reaction mixture was stirred at room temperature overnight. The reaction mixture was diluted with dichloromethane (50 mL) and washed with 5% aqueous citric acid (2 x 50 mL) and brine (50 mL). The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue was purified by silica gel chromatography, eluting from 0% to 10% ethyl acetate in heptane to afford *tert*-butyl 2-{[(benzyloxy)carbonyl][2-(2-bromophenyl)propan-2-yl]amino}acetate (4.2 g, 67% yield) as a yellow oil. ^{1}H NMR (300 MHz, $CDCl_3$): ppm 1.41 (s, 6 H), 1.55 (s, 3 H), 1.78 (s, 2 H), 1.84 (s, 4 H), 4.21 (s, 1.3 H), 4.32 (s, 0.7 H), 4.83 (s, 0.7 H), 4.99 (s, 1.3 H), 6.70 – 7.62 (m, 9 H). $[M+H]^+$ = 484.1.

Step 4: 2-benzyl 1-*tert*-butyl 3,3-dimethyl-2,3-dihydro-1*H*-isoindole-1,2-dicarboxylate



Tris(dibenzylideneacetone)dipalladium(0) (87 mg, 0.10 mmol), 2'-(diphenylphosphino)-*N*,*N*'-dimethyl-(1,1'-biphenyl)-2-amine (PhDavePhos) (73 mg, 0.19 mmol) and lithium *tert*-butoxide (0.31 g, 3.81 mmol) were added in order to a solution of *tert*-butyl 2-{[(benzyloxy)carbonyl][2-(2-bromophenyl)propan-2-yl]amino}acetate (0.88 g, 1.90 mmol) in dioxane (10 mL) previously degased and purged with nitrogen. The reaction mixture was stirred at 90 °C overnight. Once cooled at room temperature, the reaction mixture was diluted with ethyl acetate (20 mL), filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography, eluting from 0% to 10% ethyl acetate in heptane to afford 2-benzyl 1-*tert*-butyl 3,3-dimethyl-2,3-dihydro-1*H*-isoindole-1,2-dicarboxylate (0.31 g, 43% yield) as an orange oil. ^{1}H NMR (300 MHz, $CDCl_3$): ppm 1.31 (s, 6H), 1.45 (s, 3H), 1.65 (s, 1H), 1.73 (s, 1H), 1.76 (s, 2H), 1.81 (s, 2H), 5.11-5.47 (m, 3H), 7.12-7.48 (m, 9H). $[M+Na]^+$ = 404.2.

PREPARATION 73: 6-hydroxy-1-methyl-*N*-((6-nitropyridin-2-yl)sulfonyl)-2-(2,4,6-trimethylbenzyl)isoindoline-1-carboxamide



Under an atmosphere of nitrogen, 6-methoxy-1-methyl-*N*-[(6-nitro-2-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (333 mg, 0.635 mmol) was dissolved in dichloromethane (30 mL). The mixture was cooled to -78 °C, and tribromoborane (5.046 g, 1.904 mL of 1 M, 1.904 mmol) was added to the reaction mixture dropwise. The mixture was slowly warmed to room temperature and stirred for 16 hours. The reaction mixture was added to approximately 50 mL of ice. The layers were separated, and the aqueous layer was extracted with dichloromethane (3 x 5 mL). The combined organic layers were dried over sodium sulfate, concentrated, and purified on a column of silica gel utilizing a gradient of 0 to 100% ethyl acetate in hexanes to yield 6-hydroxy-1-methyl-*N*-[(6-nitro-2-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]isoindoline-1-carboxamide (90.7 mg, 0.178 mmol, 28.0%) ESI-MS *m/z* calc. 510.15732, found 511.5 $(M+1)^+$; Retention time: 0.56 minutes


PREPARATION 74: *tert*-butyl 2,3,4,5-tetrahydro-1*H*-benzo[*c*]azepine-3-carboxylate

Step 1: 3, 4-dihydro-1*H*-2-benzopyran-1-one




To a solution of isochroman (15.0 g, 112 mmol) in dichloromethane (1.4 L) stirred at room temperature was added, over 15 minutes, a homogeneous mixture of potassium permanganate (106 g, 672 mmol) and manganese dioxide (58.0 g, 672 mmol), prepared by grinding potassium permanganate and then mixing with manganese dioxide using a pestle and mortar. The suspension was vigorously stirred for 4 days at room temperature, and then filtered through a pad of Celite. The filtrate was concentrated and the residue was purified by silica chromatography, eluting with a mixture of heptanes and ethyl acetate (0 to 20%) to provide 3,4-dihydro-1*H*-2-benzopyran-1-one (8.2 g, 49% yield) as a colorless oil. ^{1}H NMR ($CDCl_3$, 300 MHz) ppm 3.06 (t, J = 6.0 Hz, 2 H), 4.52 (t, J = 6.0 Hz, 2 H), 7.20 – 7.30 (m, 1 H), 7.38 (t, J = 7.6 Hz, 1 H), 7.53 (t, J= 7.6 Hz, 1 H), 8.06 (d, J = 7.6 Hz, 1 H).

Step 2: ethyl 2-(2-bromoethyl)benzoate



Phosphorus tribromide (1.05 mL, 11.0 mmol) and bromine (0.62 mL, 12 mmol) were slowly added to a solution of 3,4-dihydro-1*H*-2-benzopyran-1-one (1.5 g, 10. mmol) in carbon tetrachloride (15 mL) cooled in an ice-water bath. The mixture was then stirred at room temperature for 16 hours. The resulting orange suspension was stirred at 60 °C for 3 hours. The red mixture was then cooled down and ethanol (10 mL) was slowly added at 0 °C (exothermic!). The reaction mixture (orange solution) was stirred for one hour. The reaction mixture was partitioned between dichloromethane (50 mL) and water (20 mL). The organic phase was separated and washed with brine (10 mL), dried over sodium sulfate and concentrated under reduced pressure. The residue was purified by silica chromatography, eluting with a mixture of ethyl acetate and heptanes (0 to 10%) to afford ethyl 2-(2-bromoethyl)benzoate (2.0 g, 78% yield) as a pale yellow oil. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.41 (t, J = 6.9 Hz, 3 H), 3.50 (t, J = 6.9 Hz, 2 H), 3.64 (t, J = 6.9 Hz, 2 H), 4.38 (q, J = 6.9 Hz, 2 H), 7.27 – 7.39 (m, 2 H), 7.45 (t, J = 7.5 Hz, 1 H), 7.96 (d, J=7.5 Hz, 1 H).

Step 3: ethyl 2-(3-amino-4-ethoxy-4-oxobutyl)benzoate




The reaction was run in two separate batches.

N-(Diphenylmethylene)glycine ethyl ester (4.17 g, 15.6 mmol) was added to a suspension of finely powdered potassium hydroxide (2.63 g, 46.8 mmol) in dimethylsulfoxide (20 ml) cooled with an ice-water bath (an orange mixture was obtained). Then ethyl 2-(2-bromoethyl)benzoate (4.0 g, 15.6 mmol) was added dropwise. The reaction was stirred at room temperature for 60 minutes and then diluted with ethyl acetate (150 mL). The solution was decanted into a separation funnel and the solid (KOH) was discarded. The organic phase was

washed with water (3 x 50 mL,) brine, dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The second batch was run in a similar way: ethyl 2-(2-bromoethyl) benzoate (9.3 g, 36.1 mmol), *N*-(diphenylmethylene)glycine ethyl ester (9.65 g, 36.1 mmol), potassium hydroxide (6.08 g, 108.3 mmol) and dimethylsulfoxide (50 mL).

The crude products of the two batches were combined (19 g) and dissolved in tetrahydrofuran (50 mL). The solution was cooled with an ice-water bath and 1 N HCl aqueous solution (60 mL) was added. The mixture was stirred at room temperature for 30 minutes. Water (50 mL) and ethyl acetate (50 mL) were added, the biphasic mixture was extracted and the two phases were separated. The aqueous phase was then basified with 2 N NaOH aqueous solution (30 mL) at 0 °C and the resulting aqueous solution was extracted with ethyl acetate (2 x 50 mL). The organic layers were combined, washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was purified with silica chromatography, eluting with a mixture of methanol and dichloromethane (0-7%) to give ethyl 2-(3-amino-4-ethoxy-4-oxobutyl)benzoate (6.6 g, 46% yield) as a yellow oil. ^{1}H NMR (300 MHz, $CDCl_3$) 1.27 (t, J = 7.1 Hz, 3 H), 1.41 (t, J = 7.1 Hz, 3 H), 1.75 - 1.93 (m, 1 H), 1.97 - 2.14 (m, 1 H), 2.95 - 3.15 (m, 2 H), 3.42 - 3.54 (m, 1 H), 4.17 (q, J = 7.1 Hz, 2 H), 4.35 (q, J = 7.1 Hz, 2 H), 7.20 - 7.30 (m, 2 H), 7.41 (t, J = 7.5 Hz, 1 H), 7.88 (d, J = 7.5 Hz, 1 H). LCMS: $[M+H]^+$ = 280.2.


Step 4: Synthesis of ethyl 1-oxo-2,3,4,5-tetrahydro-1*H*-2-benzazepine-3-carboxylate



Sodium metal (823 mg, 35.8 mmol) was added to anhydrous ethanol (270 mL, dried with 3 Å molecular sieves) in a flask at room temperature. The mixture was stirred at room temperature until all the sodium had disappeared. Then a solution of ethyl 2-(3-amino-4-ethoxy-4-oxobutyl)benzoate (5.0 g, 18 mmol) in anhydrous ethanol (30 mL) was added at room temperature. The reaction was stirred at room temperature for two days and then cooled with an ice-water bath. Thionyl chloride (10 mL) was slowly added (reacts violently) and the reaction was refluxed for 2 hours. The solvent was removed under reduced pressure and the residue was

taken up into ethyl acetate (100 mL), washed with saturated aqueous sodium bicarbonate (30 mL), brine, dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by silica chromatography, eluting with a mixture of ethyl acetate and heptanes (10 to 100%) to afford ethyl 1-oxo-2,3,4,5-tetrahydro-1*H*-2-benzazepine-3-carboxylate (2.9 g, 69% yield) as a yellow oil. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.24 (t, J = 6.7 Hz, 3 H), 2.07 – 2.25 (m, 1 H), 2.26 – 2.43 (m, 1 H), 2.70 – 2.84 (m, 1 H), 2.89 – 3.08 (m, 1 H), 3.80 – 3.95 (m, 1 H), 4.20 (q, J = 6.7 Hz, 2 H), 6.69 (s, 1 H), 7.20 (d, J = 7.2 Hz, 1 H), 7.33 (t, J = 7.2 Hz, 1 H), 7.41 (t, J = 7.2 Hz, 1 H), 7.70 (d, J = 7.2 Hz, 1 H). LCMS: $[M+H]^+$ = 234.1.

Step 5: *tert*-butyl 1-oxo-2,3,4,5-tetrahydro-1*H*-2-benzazepine-3-carboxylate




Butyl lithium (17.4 mL of 2.5 M in hexanes, 43.4 mmol) was added to a solution of anhydrous *tert*-butanol (4.7 mL, 50. mmol, distilled and then dried over potassium carbonate) in anhydrous tetrahydrofuran (40 mL) at -78 °C in a pressure tube under nitrogen. The mixture was stirred at room temperature for 10 min, then a solution of ethyl 1-oxo-2,3,4,5-tetrahydro-1*H*-2-benzazepine-3-carboxylate (2.9 g, 12 mmol) in anhydrous tetrahydrofuran (10 mL) was added and the mixture was stirred at 30 °C overnight. Volatiles were removed under reduced pressure and the residue was taken up into a mixture of water (40 mL) and dichloromethane (100 mL). The two phases were separated and the organic phase was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was purified by silica chromatography, eluting with a mixture of ethyl acetate and heptanes (0 to 30%) to give *tert*-butyl 1-oxo-2,3,4,5-tetrahydro-1*H*-2-benzazepine-3-carboxylate (1.4 g, 43% yield) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.44 (s, 9 H), 1.99 - 2.23 (m, 1 H), 2.32 (tt, J = 12.7, 6.3 Hz, 1 H), 2.77 (dd, J = 13.7, 6.3 Hz, 1 H), 2.89 - 3.11 (m, 1 H), 3.68 - 3.88 (m, 1 H), 6.64 (br. s., 1 H), 7.21 (d, J = 7.3 Hz, 1 H), 7.29 - 7.49 (m, 2 H), 7.71 (d, J = 7.5 Hz, 1 H). $[M+H]^+$ = 262.1.

Step 6: *tert*-butyl 1-methoxy-4,5-dihydro- *H*-2-benzazepine-3-carboxylate



The reaction was run in two identical batches. 4 Å molecular sieves were added to a sealed tube. The tube was flame-dried under vacuum and then flushed with nitrogen. After the tube was cooled to room temperature, *tert*-butyl 1-oxo-2,3,4,5-tetrahydro-1*H*-2-benzazepine-3-carboxylate (522 mg, 2.00 mmol) was added followed by anhydrous dichloromethane (9 mL) and trimethyloxonium tetrafluoroborate (296 mg, 2.00 mmol). The mixture was stirred at 30 °C for 16 hours. The mixture was quenched with a saturated aqueous sodium bicarbonate solution (30 mL). The resulting mixture was extracted with ethyl acetate (2 x 40 mL). The organic phases were combined and washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue of the two batches were combined and purified by silica chromatography, eluting with a mixture of ethyl acetate and heptanes (0 to 30%) to provide *tert*-butyl 1-methoxy-4,5-dihydro-3*H*-2-benzazepine-3-carboxylate (900. mg, 3.27 mmol, 81.7% yield) as a colorless oil. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.43 (s, 9 H), 2.33-2.49 (m, 2 H), 2.58 - 2.70 (m, 2 H), 3.50 (dd, J = 11.0, 7.2 Hz, 1 H), 3.96 (s, 3 H), 7.20 - 7.28 (m, 1 H), 7.30 - 7.40 (m, 2 H), 7.41 - 7.47 (m, 1 H). LCMS: $[M+H]^+$=276.1.

Step 7: *tert*-butyl 2,3,4,5-tetrahydro-1*H*-2-benzazepine-3-carboxylate




Sodium cyanoborohydride (821 mg, 13.1 mmol) was added to a solution of *tert*-butyl 1-methoxy-4,5-dihydro-3*H*-2-benzazepine-3-carboxylate (900. mg, 3.27 mmol) in acetic acid (3 mL) at 0 °C. The mixture was stirred at room temperature for 2 hours. The mixture was diluted with ethyl acetate (40 mL) and cooled with an ice-water bath. An aqueous solution of NaOH (25% w/v) was slowly added to adjust the pH to about 8. The two phases were separated and the aqueous phase was extracted with ethyl acetate (40 mL). The organic phases were combined, washed with brine (10 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was purified by silica chromatography, eluting with a mixture of ethyl

acetate and dichloromethane (0 to 50%) to give *tert*-butyl 2,3,4,5-tetrahydro-1*H*-2-benzazepine-3-carboxylate (610. mg, 2.47 mmol, 75.5% yield) as a yellow oil. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.46 (s, 9 H), 1.53 - 1.66 (m, 1 H), 2.17 - 2.37 (m, 1 H), 2.79 - 2.99 (m, 1 H), 2.99 - 3.18 (m, 1 H), 3.65 (dd, J = 10.6, 2.9 Hz, 1 H), 3.93 (d, J = 15.1 Hz, 1 H), 3.97 (d, J = 15.1 Hz, 1 H), 7.06 - 7.21 (m, 4 H). $[M+H]^+$ = 248.1.


PREPARATION 75: 1-[(4-methoxyphenyl)methyl]-2-oxopyrrolidine-3-sulfonamide

Step 1: 2,4-dibromo-*N*-[(4-methoxyphenyl)methyl]butanamide




To a solution of 2,4-dibromobutanoyl chloride (5.00 g, 18.9 mmol) in dichloromethane (250 mL) at 0 °C was added triethylamine (2.38 g, 23.6 mmol) followed by 4-methoxybenzylamine (2.59 g, 18.9 mmol) and the mixture was stirred at 0 °C for 2 hours. The reaction was then poured at 0 °C into a solution of 5% citric acid (80 mL) at pH = 3-4 and diluted with dichloromethane (100 mL) and water (50 mL). The organic layer was separated and the aqueous player was extracted again with dichloromethane (50 mL). The combined organic layer was dried over Na_2SO_4, filtered and concentrated under reduced pressure to afford crude 2,4-dibromo-*N*-[(4-methoxyphenyl)methyl]butanamide (7.09 g, 103%) as a brown solid. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 2.41 – 2.55 (m, 1 H), 2.64 – 2.84 (m, 1 H), 3.52 – 3.61 (m, 2 H), 3.80 (s, 3 H), 4.36 – 4.44 (m, 2 H), 4.55 (dd, J = 9.0, 4.8 Hz, 1 H), 6.54 (br. s, 1 H), 6.85 – 6.91 (m, 2 H), 7.19 – 7.23 (m, 2 H). $[M+H]^+$ = 366.0.

Step 2: 3-bromo-1-[(4-methoxyphenyl)methyl]pyrrolidin-2-one




To a solution of 2,4-dibromo-*N*-[(4-methoxyphenyl)methyl]butanamide (3.25 g, 8.90 mmol) in tetrahydrofuran (120 mL) and *N,N*-dimethylformamide (12 mL) was added sodium hydride (427 mg of 60% dispersion in mineral oil, 17.8 mmol) portionwise at 0 °C and the mixture was stirred at room temperature for 2.5 hours. The reaction mixture was cooled again to 0 °C and quenched carefully with an aqueous 5% citric acid solution (10 mL). The volatiles were then removed under reduced pressure. The reaction mixture was diluted with water and dichloromethane and the aqueous layer was extracted twice with dichloromethane. The combined organic layers were dried over Na_2SO_4, filtered, concentrated under reduced pressure and co-evaporated with heptanes as a co-solvent (twice) to afford crude 3-bromo-1-[(4-methoxyphenyl)methyl]pyrrolidin-2-one (2.5 g, 99%) as a pale yellow oil. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 2.22 – 2.32 (m, 1 H), 2.47 – 2.60 (m, 1 H), 3.14 – 3.21 (m, 1 H), 3.35 – 3.44 (m, 1 H), 3.80 (s, 3 H), 4.33 – 4.52 (m, 3 H), 6.84 – 6.87 (m, 2 H), 7.15 – 7.20 (m, 2 H). $[M+H]^+$ = 284.0.

Step 3: methyl 3-({1-[(4-methoxyphenyl)methyl]-2-oxopyrrolidin-3-yl}sulfonyl)propanoate



To a solution of 3-bromo-1-[(4-methoxyphenyl)methyl]pyrrolidin-2-one (2.50 g, 8.80 mmol) in dimethylsulfoxide (50 mL) was added sodium 1-methyl 3-sulfinopropanoate (1.53 g, 8.80 mmol) and the reaction was stirred at room temperature for three days. The solution was diluted with water (400 mL) and extracted with dichloromethane (2 x 250 mL). The combined organic layer was washed with water (150 mL), brine (100 mL), dried over Na_2SO_4, filtered and concentrated under reduced pressure to afford crude methyl 3-({1-[(4-methoxyphenyl)methyl]-2-oxopyrrolidin-3-yl}sulfonyl)propanoate (2.5 g, 80.% yield). ^{1}H NMR (300 MHz, $CDCl_3$) ppm 2.31 – 2.45 (m, 1 H), 2.65 – 2.75 (m, 1 H), 2.85 – 3.05 (m, 2 H), 3.26 (dt, J = 9.4, 3.8 Hz, 1 H), 3.41 – 3.50 (m, 1 H), 3.74 (s, 3 H), 3.79 (s, 3 H), 3.77 – 3.92 (m, 2 H), 4.01 (dd, J = 10.2, 4.6 Hz, 1 H), 4.44 (s, 2 H), 6.87 (d, J = 8.6 Hz, 2 H), 7.16 (d, J = 8.5 Hz, 2 H). $[M+H]^+$ = 356.1.

Step 4: sodium 1-[(4-methoxyphenyl)methyl]-2-oxopyrrolidine-3-sulfinate



To a solution of methyl 3-({1-[(4-methoxyphenyl)methyl]-2-oxopyrrolidin-3-yl}sulfonyl)propanoate (2.50 g, 7.03 mmol) in a 1:1 (v/v) mixture of tetrahydrofuran and methanol (40 mL) was added sodium methoxide (3.39 mL of 25 wt. % in methanol, 14.8 mmol) and the reaction was stirred at room temperature for 4 hours. The reaction mixture was then concentrated under reduced pressure to remove the methyl acrylate and kept under reduced pressure to provide crude sodium 1-[(4-methoxyphenyl)methyl]-2-oxopyrrolidine-3-sulfinate that was used directly in the next step without further purification $[M+H]^+$ = 270.0 (sulfinic acid).


Step 5: Synthesis of 1-[(4-methoxyphenyl)methyl]-2-oxopyrrolidine-3-sulfonamide



To a solution of crude sodium 1-[(4-methoxyphenyl)methyl]-2-oxopyrrolidine-3-sulfinate (2.05 g, 7.03mmol) in dimethylsulfoxide (20 ml) was added a solution of hydroxylamine-*O*-sulfonic acid (3.98 g, 35.2 mmol) and sodium acetate (2.19 g, 26.7 mmol) in water (7 mL) at 0 °C. The reaction was stirred for 16 hours at room temperature then diluted with ethyl acetate and water. The aqueous layer was extracted with ethyl acetate (2 x 100 mL). The


combined organic layer was washed with brine (75 mL), dried over Na_2SO_4, filtered and concentrated under reduced pressure. The crude material was purified by silica-gel chromatography using 0% to 10% methanol in dichloromethane (containing 1% NH_4OH) to afford 1-[(4-methoxyphenyl)methyl]-2-oxopyrrolidine-3-sulfonamide (1.3 g, 63% yield) as a pale brown solid. ^{1}H NMR (300MHz, $CDCl_3$) ppm 2.29 – 2.73 (m, 2 H), 3.14 – 3.34 (m, 1 H), 3.34 – 3.51 (m, 1 H), 3.80 (s, 3 H), 3.89 – 4.11 (m, 1 H), 4.30 – 4.61 (m, 2 H), 5.25 (br. s., 2 H), 6.75 – 6.98 (m, 2 H), 7.10 – 7.22 (m, 2 H). $[M+H]^+$ = 285.1.

PREPARATION 76: 6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(2-oxopyrrolidin-3-yl)sulfonyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 466)



6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[1-[(4-methoxyphenyl)methyl]-2-oxo-pyrrolidin-3-yl]sulfonyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (32.7 mg, 0.0490 mmol) was dissolved in a mixture of trifluoroacetic acid (0.9319 mL) and dichloromethane (0.9319 mL). Trifluoromethanesulfonic acid (25.75 mg, 15.18 µL, 0.1716 mmol) was then added and the mixture and stirred at ambient temperature for 20 hours. The reaction mixture was concentrated, then diluted with dimethylsulfoxide and purified by reverese-phase chromatography utilizing a C18 column and a gradient of 1-99% acetonitrile in 5 mM aq HCl to yield 6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(2-oxopyrrolidin-3-yl)sulfonyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 466) ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.27 (d, J = 19.3 Hz, 1H), 7.94 (d, J = 8.0 Hz, 1H), 7.72 (d, J = 8.0 Hz, 1H), 7.51 (s, 1H), 7.43 (d, J = 10.1 Hz, 1H), 7.28 (dd, J = 14.4, 8.0 Hz, 1H), 6.90 (dt, J = 10.6, 2.3 Hz, 1H), 4.61 - 4.44 (m, 1H), 3.84 (d, J = 6.7 Hz, 2H), 3.40 - 2.86 (m, 4H), 2.16 - 1.91 (m, 2H), 1.71 - 1.60 (m, 7H), 1.60 - 1.49 (m, 1H), 1.16 (s, 1H), 1.06 - 0.96 (m, 10H). ESI-MS *m/z* calc. 546.2312, found 547.3 $(M+1)^+$; Retention time: 1.87 minutes


PREPARATION 77: 5-isopropyl-1-(2,4,6-trimethylbenzyl)-1*H*-indole-2-carboxylic acid and 5-propyl-1-(2,4,6-trimethylbenzyl)-1*H*-indole-2-carboxylic acid



Step 1: 5-bromo-1-[(2,4,6-trimethylphenyl)methyl]indole-2-carboxylic acid (0.7000 g, 1.880 mmol) , [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (68.78 mg, 0.09400 mmol), aqueous sodium carbonate (1.880 mL of 2 M, 3.760 mmol), and 2-isopropenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (315.9 mg, 1.880 mmol) in dioxane (5 mL) were added to a microwave reactor vial. The vial was purged with nitrogen, capped and heated to 120 °C for 45 minutes in a microwave reactor. The reaction mixture was diluted with water and extracted with ethyl acetate (2 x 10 mL). The organic layers were dried over sodium sulfate, concentrated, and purified by reverse-phase chromatography utilizing a C18 column and a gradient of 30-90% acetonitrile in 5 mM aq HCl to give a mixture of products which could not be separated under these specific condition. This material was dissolved in methanol (10 mL) and palladium on carbon (20.01 mg, 0.01880 mmol) was added to the reaction mixture The mixture was stirred under an atmosphere of hydrogen for 4 hours. The mixture of crude products was filtered, through a pad of silica gel with 100 mL of 15% methanol in dichloromethane and used for the next step without further manipulation. ESI-MS *m/z* calc. 335.19, found 336.5 $(M+1)^+$; Retention time: 0.84 minutes.

PREPARATION 78: (4*R*)-2,2,4-trimethylpyrrolidine


Step 1: Methyl 2,4-dimethyl-4-nitropentanoate



A 1-L three-necked flask was fitted with a dropping funnel, and a thermometer. A solution of 2-nitropropane (178 g, 2.00 mol) in dioxane (100 mL) and a 40% aqueous solution of benzyltrimethylammonium hydroxide (Triton B, 20.0 mL, 420. mmol) were added to the flask. The flask was heated to 70 °C in an oil bath and methyl methacrylate (200. g, 212.3 mL, 2.00

mol) was added via a dropping funnel over 45 minutes. The temperature rose to about 100 °C during the addition then dropped to ~ 90 °C. The mixture was then heated at 100 °C for 4 hours. The reaction mixture was cooled to room temperature and 1 N hydrochloric acid (80 mL) was added. Water (500 mL) and diethyl ether (1000 mL) were added to the reaction flask. The mixture was poured into a separatory funnel and organic layer was washed with water (2 x 400 mL), brine (400 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a yellow liquid. The product was distilled through a short path distillation apparatus to give a pale yellow liquid (356 g, 94%) (oil bath temperature 150 °C, an oil pump, collecting temperature 100 to 108 °C). ^{1}H NMR ($CDCl_3$, 250 MHz): δ 3.67 (s, 3H), 2.55-2.35 (m, 2H), 2.10 – 1.90 (m, 1H), 1.58 (s, 3H), 1.53 (s, 3H), 1.18 (d, J=6.5 Hz, 3H).


Step 2: (*R*)-3,5,5-trimethylpyrrolidin-2-one and (*S*)-3,5,5-trimethylpyrrolidin-2-one



A solution of methyl 2,4-dimethyl-4-nitro-pentanoate (100. g, 528.5 mmol), nickel(II) chloride hexahydrate (25.12 g, 105.7 mmol), water (28.57 g, 28.57 mL, 1.586 mol) and methanol (1.200 L) was cooled to 0 to 5 °C. Sodium borohydride (49.98 g, 1.321 mol) was slowly added to the reaction mixture while maintaining the reaction temperature below 20 °C. The addition was exothermic with steady and prolonged gas generation. THe reaction mixture was allowed to stand for 14 days. A solution of potassium carbonate (146.1 g, 1.057 mol) in water (1.000 L) was slowly added to the reaction mixture. The reaction mixture was then filtered through a pad of celite. The pad was then washed with three 100 mL portions of methanol. The reaction mixture was then concentrated under reduced pressure to remove the methanol. Methyl *tert*-butyl ether (1 L) was added. The layers were separated and the aqueous layer was extracted twice with 400 mL portions of methyl *tert*-butyl ether. The combined organic layers were dried over sodium sulfate, filtered, and evaporated to dryness. This material was then portioned between methyl *tert*-butyl ether and an aqueous solution of sodium chloride. The organic layer was separated, dried over magnesium sulfate, filtered, and then evaporated to dryness to yield 41.5 g (61.7%) of an off-white solid. The stereoisomers were separated using supercritical fluid

chromatography on a ChiralPak AS-H (250 x 4.6 mm), 5 μm column using 10% methanol in CO_2 at a flow rate of 3.0 mL/min.

Step 3: (4*R*)-2,2,4-trimethylpyrrolidine



A solution of (3*R*)-3,5,5-trimethylpyrrolidin-2-one (20.0 g, 157 mmol) in tetrahydrofuran (200.0 mL) was cooled to 0 to 5 °C. Lithium aluminum hydride (8.006 g, 236.0 mmol) was added in portions over 10 minutes while maintain the temperature of the reaction mixture below 20 °C. The addition was somewhat exothermic and degassing was observed. The suspension was heated to reflux for 9.5 hours. After cooling to room temperature an additional portion of lithium aluminum hydride (1.601 g, 1.746 mL, 47.19 mmol) was added to the reaction mixture. The mixture was then heated to reflux for 2.5 hours. The reaction mixture was cooled to 7 °C and an aqueous solution of sodium sulfate (27.93 mL of 20 % w/v, 39.32 mmol) was slowly added to the reaction mixture. The reaction mixture was then diluted with 200 mL of methyl *tert*-butyl ether and filtered through a pad of celite. The filtrate was dried over sodium sulfate, filtered, and concentrated at ambient pressure to remove the ethers. The crude material was purified by distillation (28 °C/50 torr) to afford 10. g (56%) of a colorless liquid. 1H NMR (400 MHz, Chloroform-*d*) δ 3.11 (dd, J = 10.8, 7.3 Hz, 1H), 2.51 (dd, J = 10.9, 8.1 Hz, 1H), 2.32 – 2.15 (m, 1H), 1.80 (dd, J = 12.4, 8.1 Hz, 1H), 1.65 (s, 1H), 1.19 (s, 3H), 1.17 – 1.08 (m, 4H), 1.01 (d, J = 6.8 Hz, 3H). The absolute stereochemistry was determined by X-ray crystallography of the (2*R*)-2-(2-chlorophenyl)-2-hydroxy-acetic acid salt.

The following compounds can be synthesized using the procedures described herein

(*S*)-4-ethyl-2,2-dimethylpyrrolidine, and

(*R*)-4-ethyl-2,2-dimethylpyrrolidine


The following compounds can be synthesized using the procedures described herein using lithium aluminum dueterohydride:

(*S*)-2,2,4-trimethylpyrrolidine-5,5-d_2 and,

(*R*)-2,2,4-trimethylpyrrolidine-5,5-d_2

Preparation 79: 3-Isobutoxy-1*H*-pyrazole


Step 1: 1-(3-isobutoxy-1*H*-pyrazol-1-yl)ethan-1-one



A 4 mL vial was charged with 1-(3-hydroxypyrazol-1-yl)ethanone (100. mg, 0.793 mmol), potassium carbonate (164.3 mg, 1.189 mmol) and anhydrous DMF (1.000 mL). 1-bromo-2-methyl-propane (119.5 mg, 94.84 µL, 0.8722 mmol) was added, the vial was capped and the mixture was stirred at room temperature for 16 hours. The reaction was stirred at 80°C for one hour. The mixture was diluted with ethyl acetate (50 ml) and water (25 mL) and the two phases were separated. The aqueous phase (pH = 9-10) was further extracted with ethyl acetate (20 mL). The combined extracts were dried over sodium sulfate and the solvents evaporated to give 100 mg of a crude oil. The product was dissolved in dichloromethane and purified by flash chromatography on silica gel (4 g column) using a gradient of ethyl acetate (0 to 50 % over 15 minutes) in hexanes. The pure fractions were combined and the solvents removed under reduced pressure to provide 1-(3-isobutoxypyrazol-1-yl)ethanone (78 mg, 0.43 mmol, 54%) as a colorless oil.

ESI-MS *m/z* calc. 182.10553, found 183.2 $(M+1)^+$; Retention time: 1.34 minutes. ^{1}H NMR (400 MHz, Chloroform-d) δ 8.05 (d, *J* = 3.0 Hz, 1H), 5.96 (d, *J* = 3.0 Hz, 1H), 3.99 (d, *J* = 6.6 Hz, 2H), 2.58 (s, 3H), 2.09 (dp, *J* = 13.4, 6.7 Hz, 1H), 1.01 (d, *J* = 6.7 Hz, 6H).

Step 2: 3-Isobutoxy-1*H*-pyrazole



1-(3-Isobutoxy-1*H*-pyrazol-1-yl)ethan-1-one (72 mg, 0.40 mmol) was treated with methanol (1 mL) and NaOH (65.85 µL of 6 M, 0.3951 mmol) at room temperature for 1 hour. The volatiles were removed under reduced pressure. The residue was taken up in ethyl acetate (25 mL) and brine (20 mL)and two phases were separated. The aqueous phase was further extracted with ethyl acetate (20 mL) and the combined extracts were dried over sodium sulfate. After evaporation of the solvent, 3-isobutoxy-1*H*-pyrazole (55 mg, 0.39 mmol, 99%) was isolated as a colorless viscous oil. ESI-MS *m/z* calc. 140.09496, found 141.2 $(M+1)^+$; Retention time: 0.86 minutes. ^{1}H NMR (400 MHz, Chloroform-d) δ 7.34 (d, J = 2.4 Hz, 1H), 5.71 (d, J = 2.5 Hz, 1H), 3.90 (d, J = 6.6 Hz, 2H), 2.08 (dq, J = 13.3, 6.7 Hz, 1H), 1.00 (d, J = 6.7 Hz, 6H).

The following compounds can be synthesized using the procedures described herein


3-isobutoxy-5-methyl-1*H*-pyrazole,

4-isobutoxy-1*H*-pyrazole, and

3-isobutoxy-4-methyl-1*H*-pyrazole

Preparation 80: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxypyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1620)


Step 1: *tert*-Butyl 2-chloro-6-(3-isobutoxypyrazol-1-yl)pyridine-3-carboxylate



A 4 mL vial was charged under nitrogen with with 3-isobutoxy-1*H*-pyrazole (131.5 mg, 0.9383 mmol) and anhydrous DMF (1 mL). Sodium hydride (37.48 mg, 0.9370 mmol, 60% in mineral oil) was added and the mixture was stirred until gas evolution stopped. tert-Butyl 2,6-dichloropyridine-3-carboxylate (155 mg, 0.625 mmol) was added. The vial was capped and the mixture was stirred at 100 °C for 4 hours. LCMS showed the presence of two mono addition isomers (ratio approx. 60:40) and a major product being a double addition product. The reaction mixture was diluted with ethyl acetate (30 mL) and water (20 mL) and the two phases were separated. The aqueous phase was further extracted with ethyl acetate (25 mL). The combined extracts were dried over sodium sulfate and the solvent removed under reduced pressure to give


247 mg of crude solid. The residue was dissolved in dichloromethane and subjected to flash chromatography on silica gel (40 g column) using a gradient of ethyl acetate (0 to 20% over 20 minutes) in hexanes to yield *tert*-butyl 2-chloro-6-(3-isobutoxypyrazol-1-yl)pyridine-3-carboxylate (58 mg, 0.1649 mmol, 26.39%) (off-white solid). ESI-MS *m/z* calc. 351.13498, found 352.4 $(M+1)^+$; Retention time: 2.25 minutes.

Step 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-chloro-6-(3-isobutoxypyrazol-1-yl)pyridine-3-carboxamide



A 100 mL round bottom flask containing tert-butyl 2-chloro-6-(3-isobutoxypyrazol-1-yl)pyridine-3-carboxylate (58 mg, 0.16 mmol) was treated with dichloromethane (600 µL) and trifluoracetic acid (180 µL, 2.336 mmol). The mixture was stirred at 45°C for 1.5 hours. The volatiles were removed under reduced pressure to give a white solid. Anhydrous DMF (500 µL) was then added to the 100 mL flask containing the acid under an atmosphere of nitrogen and the mixture was stirred at 40°C. 1,1'-Carbonyldiimidazole (32.09 mg, 0.1979 mmol) was added and the mixture was stirred at 40 °C for 1 hour. A separate 4 mL vial under nitrogen was charged with 6-aminopyridine-2-sulfonamide (34.27 mg, 0.1979 mmol) and anhydrous DMF (500 µL). NaH (7.915 mg, 0.1979 mmol, 60% oil dispersion) was added. After gas evolution had stopped, the vial was capped and it was stirred at 40°C for 1 hour. This mixture was added to the first flask and the mixture was stirred at room temperature for 2 hours. The reaction was quenched with ice and water an neutralized with acetic acid (148.6 mg, 140.7 µL, 2.474 mmol, final pH = 4). The resulting white solid was filtered, washed with water and dried under reduced pressure. *N*-[(6-amino-2-pyridyl)sulfonyl]-2-chloro-6-(3-isobutoxypyrazol-1-yl)pyridine-3-carboxamide (57 mg, 0.13 mmol, 77%) was isolated as a white solid. ESI-MS *m/z* calc. 450.0877, found 451.3 $(M+1)^+$; Retention time: 1.54 minutes

Step 3: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxypyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1620)



A 4 mL vial was charged under nitrogen with *N*-[(6-amino-2-pyridyl)sulfonyl]-2-chloro-6-(3-isobutoxypyrazol-1-yl)pyridine-3-carboxamide (57 mg, 0.13 mmol), CsF (96.00 mg, 0.6320 mmol), potassium carbonate (87.35 mg, 0.6320 mmol), (4*S*)-2,2,4-trimethylpyrrolidine (47.64 mg, 0.3792 mmol) and anhydrous dimethylsulfoxide (200 µL). The vial was capped and the contents were stirred at 150°C for 4 hours. The mixture was diluted with water (200 uL), *N*-methylpyrrolidinone (600 uL) and the mixture was filtered. The compound was purified by reverse phase preparative HPLC using a gradient of acetonitrile in water (1 to 99% over 15 minutes) in 5 mM aqueous ammonium formate. The pure fractions were combined, a few drops of brine was added and the organic solvents were removed under reduced pressure. The solid that crashed out was extracted with dichloromethane (3 x 20 mL). The combined extracts were dried over sodium sulfate, filtered, and the solvents were removed under reduced pressure to give 28 mg of material that still contained an impurity. The material was purified a second time by preparative HPLC using a gradient of acetonitrile in water (1 to 99% over 15 minutes) in 5 mM aqueous hydrochloric acid. The product was isolated using the same extraction procedure. After concentration of the dichloromethane, a solid was obtained by adding hexanes. Evaporation of the solvents gave *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxypyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1620) (18 mg, 0.03394 mmol, 26.86%) as an off-white solid. ESI-MS *m/z* calc. 527.23145, found 528.5 $(M+1)^+$; Retention time: 1.77 minutes. ^{1}H NMR (400 MHz, Chloroform-d) δ 13.87 (broad s, 1H), 8.29 (d, J = 8.0 Hz, 1H) 8.21 (d, J = 2.9 Hz, 1H), 7.65 - 7.58 (m, 2H), 7.52 (d, J = 8.5 Hz, 1H), 6.65 (p, J = 3.8 Hz, 1H), 5.95 (d, J = 2.8 Hz, 1H), 4.80 (broad s, 2H), 4.02 (d, J = 6.6 Hz, 2H), 3.52 - 3.41 (m, 1H), 3.15 (dd, J = 10.5, 8.1 Hz, 1H), 2.69 - 2.50 (m, 1H), 2.18 - 2.02 (m, 2H), 1.72 (dd, J = 12.3, 9.7 Hz, 1H), 1.45 (s, 3H), 1.39 (s, 3H), 1.19 (d, J = 6.6 Hz, 3H), 1.02 (d, J = 6.7 Hz, 6H).

The following compounds can be synthesized using the procedures described herein

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxy-5-methyl-pyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1407),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(cyclobutylmethoxy)pyrazol-1-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2694),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(cyclopropylmethoxy)pyrazol-1-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2695),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(2,2-dimethylpropoxy)-4-methyl-pyrazol-1-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2699),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutylpyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1411),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(5-isobutylpyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1431),


N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-isobutoxypyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1443),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-ethoxy-5-methyl-pyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1473), and

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-isobutoxy-4-methyl-pyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1532)

Preparation 81: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-(3,4-dihydro-2*H*-pyran-5-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1487)


Step 1: 5-bromo-6-*tert*-butyl-2-hydroxypyridine-3-carbonitrile



N-Bromosuccinimide (10.3 g, 57.9 mmol) was added to 6-*tert*-butyl-2-hydroxypyridine-3-carbonitrile (6.8 g, 38.6 mmol) in 1,2-dichloroethane (50 mL) and the reaction mixture was refluxed for 3h. Water was added and extracted twice with


dichloromethane (2 x 50 mL). The organic phase was washed with water, dried with anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by flash chromatography (100% dichloromethane to 10% methanol/90% dichloromethane) to provide 5-bromo-6-*tert*-butyl-2-hydroxypyridine-3-carbonitrile in 65% yield (6.4 g) as a white solid. ^{1}H NMR (300 MHz, DMSO-d_6) ppm 1.43 (s, 9H), 8.36 (s, 1H). $[M+H]^+$ = 255.0, 257.0.

Step 2: 5-bromo-6-*tert*-butyl-2-chloropyridine-3-carbonitrile



Phosphorus oxychloride (30.3 mL, 325 mmol) and phosphorus pentachloride (20.3 g, 97.6 mmol) were added to 5-bromo-6-*tert*-butyl-2-hydroxypyridine-3-carbonitrile (8.30 g, 32.5 mmol) and the reaction mixture was refluxed for 2 days. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. Water was added to the residue, extracted 3 times with ethyl acetate (75-mL portions), washed with brine (75 mL), dried with anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (100% heptanes to 50% ethyl acetate/50% heptanes) to provide 5-bromo-6-*tert*-butyl-2-chloroxypyridine-3-carbonitrile in 39% yield (3.5 g) as an off-white solid. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.52 (s, 9H), 8.08 (s, 1H). $[M+H]^+$ = 273.0, 275.0.


Step 3: 5-bromo-6-*tert*-butyl-2-chloropyridine-3-carboxylic acid



Potassium hydroxyde (4.3 g, 77 mmol) in water (15 mL) was added to 5-bromo-6-*tert*-butyl-2-chloroxypyridine-3-carbonitrile in 2-propanol (15 mL) and the reaction mixture was heated at 90 °C for 16 hours. Ethyl acetate (50 mL) was added and extracted 3 times with 1N sodium hydroxide (3 x 50 mL). The aqueous phase was acidified to pH ~4 with 10% hydrochloric acid and the aqueous phase was extracted 3 times with ethyl acetate (3 x 75 mL), and washed with brine (100 mL). The desired compound was present in all organic phases so they were combined, dried with anhydrous sodium sulfate, filtered and concentrated under


reduced pressure. The crude compound was purified by reverse phase chromatography (methanol/water, 0 to 100%) to afford 5-bromo-6-*tert*-butyl-2-chloropyridine-3-carboxylic acid (2.3 g, 61% yield) as an off-white solid. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.53 (s, 9H), 8.46 (s, 1H). $[M+H]^+ = 292.0, 294.0$.

Step 4: *N*-[(6-amino-2-pyridyl)sulfonyl]-5-bromo-6-tert-butyl-2-chloro-pyridine-3-carboxamide



To a stirred solution of 5-bromo-6-*tert*-butyl-2-chloro-pyridine-3-carboxylic acid (500.0 mg, 1.709 mmol) in dry DMF (3.5 mL) in a 50 mL round-bottom flask was added 1,1'-carbonyldiimidazole (346.3 mg, 2.136 mmol) in one portion under nitrogen at ambient temperature. The reaction was warmed to 45 °C and stirred for 1.5 h. In a separate 50 mL round-bottom flask, 6-aminopyridine-2-sulfonamide (369.9 mg, 2.136 mmol) was dissolved in dry DMF (2.5 mL) under nitrogen and sodium hydride (60 % in mineral oil) (85.43 mg, 2.136 mmol) was added in portions to mitigate gas evolution at ambient temerature. The heterogeneous mixture was warmed to 40 °C and stirred for 1 hour. This mixture was added to the stirred activated acid mixture in one portion at 45 °C. Heating was stopped and the reaction was stirred for an additional hour. The reaction mixture from was slowly (over 2 minutes) added to ice-cold water (50 mL) in a 100 mL round-bottom flask, and placed in an ice bath. Then acetic acid (528.5 mg, 500.5 µL, 8.801 mmol) was added and the mixture stirred vigorously for 10 minutes and the solids were filtered and washed with water (3 x 20 mL). The solids were slurried in hexanes (20 mL) and filtered. The solid was further and dried to furnish desired coupled product as white solid. *N*-[(6-amino-2-pyridyl)sulfonyl]-5-bromo-6-tert-butyl-2-chloro-pyridine-3-carboxamide (585 mg, 1.31 mmol, 76.5%) ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.87 (s, 1H), 8.15 (s, 1H), 7.60 (d, J = 1.5 Hz, 1H), 7.54 (t, J = 7.8 Hz, 1H), 7.09 (d, J = 7.3 Hz, 1H), 6.58 (d, J = 8.3 Hz, 1H), 6.35 (s, 2H), 1.46 (s, 9H) ESI-MS *m/z* calc. 445.9815, found 449.2 $(M+1)^+$; Retention time: 1.52 minutes


Step 5: *N*-[(6-amino-2-pyridyl)sulfonyl]-5-bromo-6-tert-butyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1628)



2,2,4-Trimethylpyrrolidine (252.9 mg, 2.234 mmol) and potassium carbonate (617.5 mg, 4.468 mmol) were added to a solution of *N*-[(6-amino-2-pyridyl)sulfonyl]-5-bromo-6-*tert*-butyl-2-chloro-pyridine-3-carboxamide (500.0 mg, 1.117 mmol) in anhydrous DMSO (4.0 mL) in a 20 mL reaction vial. The vial was capped under a nitrogen atmosphere and heated in a pre-heated oil bath at 145 °C for 24 h. The reaction vessel was removed and allowed to cool to ambient temerature. The reaction mixture was poured onto a mixture of ice and water (50 mL) with stirring. The resulting reaction mixture was then adjusted to a pH of about 6.0 with 1 M aqueous HCl. The mixture was vigorously stirred for 15 minutes. The soild was filtered, washed with water (4 x 20 mL), and dried under vacuum to yield the desired product as white soild.

N-[(6-amino-2-pyridyl)sulfonyl]-5-bromo-6-*tert*-butyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1628) (495 mg, 0.944 mmol, 84.5%) ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.49 (s, 1H), 8.52 (s, 1H), 7.70 (s, 1H), 7.49 (s, 1H), 7.07 (d, J = 7.2 Hz, 1H), 6.53 (s, 1H), 6.20 (s, 2H), 3.37 (dd, J = 11.6, 8.2 Hz, 1H), 2.77 (dd, J = 11.6, 8.9 Hz, 1H), 2.16 (dt, J = 12.5, 6.4 Hz, 1H), 1.99 (dd, J = 12.9, 7.6 Hz, 1H), 1.81 (dd, J = 11.8, 5.5 Hz, 1H), 1.51 (s, 3H), 1.48 (s, 3H), 1.44 (s, 9H), 0.87 (d, J = 6.3 Hz, 3H). ESI-MS *m/z* calc. 523.12524, found 526.1 $(M+1)^+$; Retention time: 1.43 minutes

Step 6: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-(3,4-dihydro-2*H*-pyran-5-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1487)



A 4 mL vial was charged with a stir bar, *N*-[(6-amino-2-pyridyl)sulfonyl]-5-bromo-6-tert-butyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1628) (29 mg, 0.055 mmol) and vinylboronic acid 34.85 mg, 0.1659 mmol). Then DMF (0.5 mL) and water (0.1 mL) were added, followed by potassium carbonate (30.57 mg, 0.2212 mmol). Nitrogen was bubbled through the vial for 1 min. Then tetrakis(triphenylphosphine)palladium (0) (6.389 mg, 0.005529 mmol) was added and nitrogen was bubbled through the solution for another 1 min, sealed with a screw cap under nitrogen and heated at 130 °C (external temperature) for 13 h. The reaction mixture was filtered through a syringe filter disc, and purified by reverse phase HPLC utilizing a gradient of 10 to 99 percent acetonitrile in water containing 5 mM HCl to yield *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-(3,4-dihydro-2*H*-pyran-5-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1487) (3.3 mg, 0.0058 mmol, 11%) as the hydrochloric acid salt. ESI-MS *m/z* calc. 527.26, found 528.3 $(M+1)^+$; Retention time: 1.21 minutes

The following compounds can be synthesized using the procedures described herein

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[4-(trifluoromethyl)cyclohexen-1-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1380),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-3,3-dimethylbut-1-enyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1613),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-2-cyclohexylvinyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1484),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-hex-1-enyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1377),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-5-methylhex-1-enyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1682),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-2-cyclopentylvinyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1633),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-2-[4-(trifluoromethyl)phenyl]vinyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1566),

(E)-3-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-2-tert-butyl-6-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]-3-pyridyl]prop-2-enoic acid (Compound 1547),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-bicyclo[2.2.1]hept-2-enyl)-6-tert-butyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-phenyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1364),

N-[(6-amino-2-pyridyl)sulfonyl]-5-phenyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1384),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(4,7,7-trimethyl-3-bicyclo[2.2.1]hept-2-enyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1544),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[(E)-2-cyclopropylvinyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1571),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(1,2-dimethylprop-1-enyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1590),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-2-(4-methoxyphenyl)vinyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-pent-1-enyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-bicyclo[2.2.1]hept-2-enyl)-6-tert-butyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide,

ethyl (E)-3-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-2-tert-butyl-6-(2,2,4-trimethylpyrrolidin-1-yl)-3-pyridyl]prop-2-enoate,

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-2-ethoxyvinyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-2-(2,2,4-trimethylpyrrolidin-1-yl)-5-vinyl-pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-2-cyclopropylvinyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-pent-1-enyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-2-(4-methoxyphenyl)vinyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-2-cyclopropylvinyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-2-(2,2,4-trimethylpyrrolidin-1-yl)-5-vinyl-pyridine-3-carboxamide,

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[(E)-2-ethoxyvinyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide,

ethyl (E)-3-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-2-tert-butyl-6-(2,2,4-trimethylpyrrolidin-1-yl)-3-pyridyl]prop-2-enoate,

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-(2-methylprop-1-enyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1561),

benzyl 4-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-2-tert-butyl-6-(2,2,4-trimethylpyrrolidin-1-yl)-3-pyridyl]-3,6-dihydro-2H-pyridine-1-carboxylate (Compound 1404),

tert-butyl 3-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-2-tert-butyl-6-(2,2,4-trimethylpyrrolidin-1-yl)-3-pyridyl]-8-azabicyclo[3.2.1]oct-3-ene-8-carboxylate (Compound 1388),

5-(4-acetylcyclohexen-1-yl)-*N*-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1461),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-(1,4-dioxaspiro[4.5]dec-8-en-8-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1459),

N-[(6-amino-2-pyridyl)sulfonyl]-5-bromo-6-tert-butyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1628),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-(2,2,6,6-tetramethyl-3H-pyran-4-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1647),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-(4,7,7-trimethyl-3-bicyclo[2.2.1]hept-2-enyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1578),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-(3,4-dihydro-2H-pyran-6-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1485),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-(5,5-dimethyl-3-oxo-cyclohexen-1-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1405),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-(1,4-dioxaspiro[4.5]dec-8-en-9-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1469),

tert-butyl 4-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-2-tert-butyl-6-(2,2,4-trimethylpyrrolidin-1-yl)-3-pyridyl]-3,6-dihydro-2H-pyridine-1-carboxylate (Compound 1522),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-(1-methyl-3,6-dihydro-2H-pyridin-4-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1382),

4-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-2-tert-butyl-6-(2,2,4-trimethylpyrrolidin-1-yl)-3-pyridyl]cyclohex-3-ene-1-carboxylic acid (Compound 1381),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-cyano-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1419),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-(3,6-dihydro-2H-pyran-4-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1495),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-methoxy-5-methyl-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2499),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[(E)-hex-1-enyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1916),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-isobutoxy-5-methyl-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2558),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[2-(trifluoromethyl)phenyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1921),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-chlorophenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2429),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-cyanophenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1849),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(4-methoxyphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1792),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-tert-butyl-5-methyl-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2322),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[4-[1-(hydroxymethyl)cyclopropyl]phenyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2112),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(4-tert-butylphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2463),

ethyl 3-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-(2,2,4-trimethylpyrrolidin-1-yl)-3-pyridyl]-4-methoxy-benzoate (Compound 2134),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(4-chlorophenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2399),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-chloro-5-methoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2545),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(2-methoxyphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1735),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(4-cyanophenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2304),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(4-isopropylphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1809),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(2-isopropoxyphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2283),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[3-(2-methoxyethoxymethyl)phenyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1943),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-isobutoxyphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2089),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(4-isobutylphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2395),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(1H-indazol-4-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2157),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[4-(1-hydroxy-1-methyl-ethyl)phenyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2547),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[(E)-2-[4-(trifluoromethyl)phenyl]vinyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2014),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[3-(cyclopentoxy)phenyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1918),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(6-ethoxy-3-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2384),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(o-tolyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1701),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[4-(1-cyano-1-methyl-ethyl)phenyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2521),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(6-methoxy-3-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1886),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[3-(cyclopentoxy)-5-methyl-phenyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2313),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(4-dimethylaminophenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2288),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(1-isobutylpyrazol-3-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2303),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(4-methoxycyclohexen-1-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1920),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(2-cyanophenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2584),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-chloro-5-methyl-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1963),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[4-(trifluoromethoxy)phenyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2649),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(2-ethoxyphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2557),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3,5-dimethylphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2093),

tert-butyl 4-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-(2,2,4-trimethylpyrrolidin-1-yl)-3-pyridyl]-3,6-dihydro-2H-pyridine-1-carboxylate (Compound 2129),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(4-methylcyclohexen-1-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2361),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-isopentyloxyphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1894),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[3-(trifluoromethyl)phenyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1950),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[(E)-5-methylhex-1-enyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2647),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(6-methoxy-2-pyridyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1861),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[3-isopropoxy-5-(trifluoromethyl)phenyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1815),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-fluoro-5-methoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2431),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(1,1,4,4,7-pentamethyltetralin-6-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1759),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-isopropoxy-5-methyl-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1889),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3,5-dichlorophenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1721),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(2-isobutoxyphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1979),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(5-benzyloxy-2-methyl-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2588),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-isopropoxyphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1804),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3,5-difluorophenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2259),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[4-(hydroxymethyl)phenyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2088),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[4-(trifluoromethyl)cyclohexen-1-yl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2510),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(m-tolyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2184),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(2-chlorophenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2312),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[4-(trifluoromethyl)phenyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2337),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(2-chloro-5-isopropoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2210),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-methoxyphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2031),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-tert-butylphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2180),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[(E)-2-(4-methoxyphenyl)vinyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1726),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-bicyclo[2.2.1]hept-2-enyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1888),

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-ethoxy-5-methyl-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1843),


N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-methyl-4-propoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2095),

N-[(6-amino-2-pyridyl)sulfonyl]-5-[(E)-2-cyclohexylvinyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2063),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-(cyclohexen-1-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1601), and

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-(1-methylpyrazol-4-yl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1441)

Preparation 82: *tert*-butyl *N*-[1-[[6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-3-methyl-3-piperidyl]carbamate (Compound 1520)



A solution of 6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxylic acid (40.45 mg, 0.1000 mmol) and chlorosulfonyl isocyanate (16.98 mg, 10.44 µL, 0.1200 mmol) in dichloromethane (1.000 mL) was stirred for 45 minutes. Triethylamine (15.18 mg, 20.91 µL, 0.1500 mmol) and *tert*-butyl-*N*-(3-methyl-3-piperidyl)carbamate (32.15 mg, 0.1500 mmol) were added, and the raction was stirred for three days. The reaction was purified by silica gel chromatography with 0-5% methanol in dichloromethane to give tert-butyl (1-(*N*-(6-(3-fluoro-5-isobutoxyphenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)nicotinoyl)sulfamoyl)-3-methylpiperidin-3-yl)carbamate (34 mg, 0.050 mmol, 50.%) as a colorless solid.

The following compounds can be synthesized using the procedures described herein

N-[(3R)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1360),

N-(3-acetamidopyrrolidin-1-yl)sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1391),

N-[[3-(dimethylamino)-1-piperidyl]sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1414),

N-[[(3S)-3-amino-1-piperidyl]sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1435),

N-[(3-amino-3-methyl-1-piperidyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1445),

N-[(3-amino-1-piperidyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1447)

N-(3-aminopyrrolidin-1-yl)sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1467),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[[3-(methylamino)-1-piperidyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1508),

N-[(5-amino-3-azabicyclo[3.1.0]hexan-3-yl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1521),

N-[3-[acetyl(methyl)amino]pyrrolidin-1-yl]sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1533),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(3-methoxy-1-piperidyl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1545),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[3-(methylamino)pyrrolidin-1-yl]sulfonyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1554),

N-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1557),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(3-methylsulfonylpyrrolidin-1-yl)sulfonyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1560),

N-[(3-acetamido-1-piperidyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1586),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[[3-(methanesulfonamido)-1-piperidyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1600),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(3-hydroxy-1-piperidyl)sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1608),

N-[[(3R)-3-amino-1-piperidyl]sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1614),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-pyrrolidin-1-ylsulfonyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1619), and

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(3-hydroxypyrrolidin-1-yl)sulfonyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1648),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(3-hydroxy-3-methyl-azetidin-1-yl)sulfonyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2398),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(2-pyridylsulfamoyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide,

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[[(4S)-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl]sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1789),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[[(3R)-3-piperidyl]sulfamoyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2677),

N-[(6-amino-2-pyridyl)sulfamoyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide,

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-(3-pyridylsulfamoyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1448),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(2-methoxy-3-pyridyl)sulfamoyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1359),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[[(1R,4R)-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl]sulfonyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2347),

N-(3,3-difluoropyrrolidin-1-yl)sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1956),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(3,3,3-trifluoro-2-hydroxy-2-methyl-propyl)sulfamoyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide,

N-(3,8-diazabicyclo[3.2.1]octan-3-ylsulfonyl)-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2237),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(3R)-3-hydroxypyrrolidin-1-yl]sulfonyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2427),

N-[(3R,4R)-3,4-dihydroxypyrrolidin-1-yl]sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2534),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[[(3S)-pyrrolidin-3-yl]sulfamoyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1935),

N-[(5-amino-3,3-difluoro-1-piperidyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2209),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(6-methoxy-2-pyridyl)sulfamoyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide,

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[[(3R)-pyrrolidin-3-yl]sulfamoyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide,

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(3S)-3-fluoropyrrolidin-1-yl]sulfonyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2228),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(3R)-3-methoxypyrrolidin-1-yl]sulfonyl-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1893),

N-[(3,3-difluoro-5-hydroxy-1-piperidyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2508),


6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[[(3S)-3-piperidyl]sulfamoyl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide,

N-[(3R)-3-aminopyrrolidin-1-yl]sulfonyl-2-(2-benzyl-2-methyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2325) (Isomer 1, *R* or *S* stereoisomer)

N-[(3R)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-phenylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2477) (Isomer 2, *R* or *S* stereoisomer)


Preparation 83: 1-[3-fluoro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy]-2-methyl-propan-2-ol

Step 1: 1-(3-bromo-5-fluoro-phenoxy)-2-methyl-propan-2-ol



To 3-bromo-5-fluoro-phenol (9.0 g, 47 mmol), and 1-chloro-2-methyl-propan-2-ol (5.116 g, 47.12 mmol), aqueous sodium hydroxide (37.69 g, 94.24 mmol) was added and the reaction mixture was heated at 120 °C for 5 days in a pressure vessel. The reaction was cooled and then extracted three times with ethyl acetate. The combined organic layer was dried over sodium sulfate, filtered, and the solvent was evaporated under reduced pressure. The crude product was purified on 220 g of silica gel utilizing a gradient of 0 to 10% ethyl acetate in dichloromethane. The impure product was repurified on 220 g of gold silica gel utilizing a gradient of 0 to 10% ethyl acetate in dichloromethane to yield 1-(3-bromo-5-fluoro-phenoxy)-2-methyl-propan-2-ol (4.75 g, 18.0 mmol, 38%). ESI-MS *m/z* calc. 262.00046, found 263.0 $(M+1)^+$; Retention time: 1.57 minutes (3 min run).

Step 2: 1-[3-Fluoro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy]-2-methyl-propan-2-ol




To 1-(3-bromo-5-fluoro-phenoxy)-2-methyl-propan-2-ol (5.250 g, 19.95 mmol) and 4,4,5,5-tetramethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (7.598 g, 29.92 mmol) in a flask was added potassium acetate (4.894 g, 49.87 mmol) and [1,1'-Bis(diphenylphosphino)ferrocene]palladium(II) dichloride dichloromethane adduct (729.9 mg, 0.9975 mmol) followed by dioxane (32.58 mL). The suspension was degassed for 10 minutes by sparging with nitrogen and then the reaction was heated in an oil bath at 85 °C for 3 h. The reaction was diluted with ethyl acetate, filtered, and concentrated. The crude product was purified on 330 g of silica gel utilizing a gradient of 0 to 50% ethyl acetate in hexane to yield 1-[3-fluoro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy]-2-methyl-propan-2-ol (5.23

g, 16.86 mmol, 84.52%). The product was not pure and it was used in the next step without further purification. ESI-MS *m/z* calc. 310.17517, found 311.5 $(M+1)^+$; Retention time: 1.895 minutes (3 min run).


Preparation 84: 3-[3-fluoro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy]-2-methyl-propan-1-ol

Step 1: 3-(3-bromo-5-fluoro-phenoxy)-2-methyl-propan-1-ol



To a stirred solution of 3-bromo-5-fluoro-phenol (2.270 g, 11.88 mmol), 2-methylpropane-1,3-diol (1.071 g, 11.88 mmol) and triphenylphosphane (3.428 g, 3.028 mL, 13.07 mmol) in tetrahydrofuran (71.82 mL) at 0 °C was added diisopropyl azodicarboxylate (2.643 g, 2.532 mL, 13.07 mmol). The ice bath was removed and the reaction was stirred at 55 °C for 18 h. The reaction was diluted with ethylacetate and washed twice with sodium bicarbonate and twice with brine. The organic layer was dried over sodium sulfate, filtered and the solvent was evaporated under reduced pressure. The crude product was purified on 220 g of silica gel utilizing a gradient of 0 to 15% ethyl acetate in dichloromethane to yield 3-(3-bromo-5-fluoro-phenoxy)-2-methyl-propan-1-ol (1.73 g, 6.57 mmol, 55.3%) as a colorless liquid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.09 - 7.03 (m, 1H), 7.03 - 6.98 (m, 1H), 6.87 (dt, J = 11.2, 2.3 Hz, 1H), 4.57 (t, J = 5.3 Hz, 1H), 3.97 (dd, J = 9.5, 5.9 Hz, 1H), 3.85 (dd, J = 9.5, 6.3 Hz, 1H), 3.48 - 3.34 (m, 2H), 2.05 - 1.84 (m, 1H), 0.94 (d, J = 6.8 Hz, 3H). ESI-MS *m/z* calc. 262.00046, found 265.0 (M+1)+; Retention time: 1.58 minutes (3 min run).


Step 2: 3-[3-fluoro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy]-2-methyl-propan-1-ol



To 3-(3-bromo-5-fluoro-phenoxy)-2-methyl-propan-1-ol (1.7 g, 6.5 mmol) and 4,4,5,5-tetramethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (2.461 g, 9.692 mmol) in a flask was potassium acetate (1.585 g, 16.15 mmol) and [1,1'-Bis(diphenylphosphino)ferrocene]palladium(II) dichloride dichloromethane adduct (236.3 mg, 0.3230 mmol) followed by dioxane (10.55 mL). The suspension was degassed for 10 min by bubbling nitrogen through the reaction mixture and then the reaction was heated in an oil bath at 85 °C for 2.5 h. The reaction was diluted with ethyl acetate, filtered, and concentrated. The crude product was purified on 220 g of silica gel utilizing a gradient of 0 to 50% ethyl acetate in hexane to yield 3-[3-fluoro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy]-2-methyl-propan-1-ol (1.77 g, 5.71 mmol, 88%) as an orange viscous liquid. The product contained some impurity. It was used in the next step without further purification. ESI-MS *m/z* calc. 310.17517, found 311.2 $(M+1)^+$; Retention time: 1.81 minutes (3 min run).


Preparation 85: 5-(Benzyloxy)-6-nitropyridine-2-sulfonamide

Step 1: 3-Benzyloxy-6-bromo-2-nitro-pyridine




To a solution of 6-bromo-2-nitro-pyridin-3-ol (1.36 g, 6.21 mmol) in *N,N*-dimethylformamide (15 mL) was added potassium carbonate (1.717 g, 12.42 mmol) followed by the addition of bromomethylbenzene (1.115 g, 775.4 µL, 6.520 mmol). The mixture was heated at 60 °C overnight, and partitioned between ethyl acetate and water. The aqueous layer was extracted with ethyl acetate three times. The combined organic layers were washed three times with water, brine, dried over magnesium sulfate, filtered, and concentrated to dryness. The crude material was suspended in hexane, collected via filtration and air dried to provide 3-benzyloxy-6-bromo-2-nitro-pyridine (1.61 g, 5.21 mmol, 83.8%) as a yellow solid. ^{1}H NMR (400 MHz, Chloroform-d) δ 7.60 (d, *J* = 8.7 Hz, 1H), 7.45 - 7.29 (m, 6H), 5.25 (s, 2H).

Step 2: Methyl 3-[(5-benzyloxy-6-nitro-2-pyridyl)sulfonyl]propanoate



A mixture of 3-benzyloxy-6-bromo-2-nitro-pyridine (585 mg, 1.89 mmol) and (3-methoxy-3-oxo-propyl)sulfinyloxysodium (329.5 mg, 1.892 mmol) in dimethylsulfoxide (5 mL) was purged with nitrogen for 2 minutes. Iodocopper (720.7 mg, 3.784 mmol) was added to the reaction mixture. The mixture was heated at 80 °C overnight, partitioned between ethyl acetate and a saturated aqueous solution of ammonium chloride. The resulting precipitate was removed by filtration and washed with ethyl acetate. The aqueous layer was extracted three times with ethyl acetate. The combined organic layers were washed three times with water, brine, dried over magnesium sulfate, filtered and concentrated to dryness. The crude material was purified by column chromatography (40-60% ethyl acetate in hexanes) to provide methyl 3-[(5-benzyloxy-6-nitro-2-pyridyl)sulfonyl]propanoate (210. mg, 0.552 mmol, 29.2%) as a pale yellow solid. ^{1}H NMR (400 MHz, Chloroform-*d*) δ 8.20 (d, *J* = 8.6 Hz, 1H), 7.68 (d, *J* = 8.7 Hz, 1H), 7.50 - 7.31 (m, 5H), 5.35 (s, 2H), 3.75 - 3.63 (m, 5H), 2.87 (t, *J* = 7.5 Hz, 2H).

Step 3: 5-(Benzyloxy)-6-nitropyridine-2-sulfonamide




Tetrahydrofuran (10 mL) was added to a suspension of methyl 3-[(5-benzyloxy-6-nitro-2-pyridyl)sulfonyl]propanoate (1.141 g, 3.000 mmol) in methanol (20 mL). The resulting solution was cooled to 0 °C. Sodium methoxide (6.000 mL of 0.5 M in methanol, 3.000 mmol) was added in a dropwise manner. The mixture was stirred at room temperature for 15 minutes. The mixture was evaporated to dryness to provide a light brown solid. This solid was then suspended in dichloromethane (30 mL) and *N*-chlorosuccinimide (400.6 mg, 3.000 mmol) was added to the reaction mixture. The mixture was stirred at room temperature for 10 minutes. The reaction was recharged with *N*-chlorosuccinimide (80.12 mg, 0.6000 mmol). The mixture turned clear. The mixture was stirred at room temperature for 5 minutes. The resulting solution was

added to ammonia (36 mL of 0.5 M, 18.00 mmol) at 0 °C. The mixture was stirred at room temperature for 30 minutes, and then partitioned between dichloromethane and water. The aqueous layer was extracted three times with dichloromethane. The combined organic layers were washed with brine, dried over magnesium sulfate, filtered, and concentrated to dryness to provide a yellow solid (342 mg). The crude material was used directly in next step without further purification. ESI-MS *m/z* calc. 309.04, found 310.3 $(M+1)^+$; Retention time: 1.10 minutes (3 min run).


Preparation 86: *N*-[(6-amino-2-pyridyl)sulfonyl]-3-fluoro-4-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)benzamide (Compound 1497)

Step 1: *N*-[(6-amino-2-pyridyl)sulfonyl]-4-bromo-2,3-difluoro-benzamide




To 4-bromo-2,3-difluoro-benzoic acid (2.66 g, 11.2 mmol) in *N, N*-dimethylformamide (2.3 mL) was added at room temperature di(imidazol-1-yl)methanone (2.7 g, 17 mmol) and reaction was heated at 65 °C for 1 hour. Separately to 6-aminopyridine-2-sulfonamide (2.4 g, 14.0 mmol) in *N, N*-dimethylformamide (2 mL) was added sodium hydride (561 mg, 14.0 mmol) at 0 °C and stirred for 10 minutes for 1 hour. The reaction was cooled back to 0 °C and the adduct from above was added and the reaction was heated at 65 °C for 1.5 hours. The reaction was cooled to room temperature and diluted with ethyl acetate and washed with a brine solution. The organics were separated, dried and concentrated to give a residue which was purified with silica gel eluting with 0 to 100% ethyl acetate in hexanes to give *N*-[(6-amino-2-pyridyl)sulfonyl]-4-bromo-2,3-difluoro-benzamide (3.8 g, 9.7 mmol, 86%) as product. ESI-MS *m/z* calc 390.94 , found 392.0 $(M+1)^+$; Retention time: 1.1 minutes (3 min run)

Step 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-2,3-difluoro-4-(3-fluoro-5-isobutoxy-phenyl)benzamide



To mixture of *N*-[(6-amino-2-pyridyl)sulfonyl]-4-bromo-2,3-difluoro-benzamide (700. mg, 1.79 mmol), (3-fluoro-5-isobutoxy-phenyl)boronic acid (568 mg, 2.68 mmol), and potassium carbonate (740 mg, 5.4 mmol) in a vial was added [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (130.6 mg, 0.1785 mmol) and *N, N*-dimethylformamide (4.8 mL) and water (1.2 mL) and the reaction mixture was stirred under nitrogen at 100 °C for 16 hours. The reaction mixture was cooled, filtered over pad of celite and the filtrate was diluted with ethyl acetate and a brine solution. The organics were separated, dried over sodium sulfate, and concentrated to give a residue which was purified with silica gel using 5 to 100% ethyl acetate in dichloromethane to give *N*-[(6-amino-2-pyridyl)sulfonyl]-2,3-difluoro-4-(3-fluoro-5-isobutoxy-phenyl)benzamide (174 mg, 0.363 mmol, 20.3%) ^{1}H NMR (400 MHz, Methanol-d_4) δ 7.64 - 7.53 (m, 2H), 7.29 (d, J = 7.4 Hz, 1H), 7.25 - 7.18 (m, 1H), 6.90 (d, J = 2.7 Hz, 1H), 6.88 - 6.81 (m, 1H), 6.73 (dt, J = 10.8, 2.3 Hz, 1H), 6.64 (d, J = 8.3 Hz, 1H), 3.79 (d, J = 6.4 Hz, 2H), 2.07 (hept, J = 6.7 Hz, 1H), 1.04 (d, J = 6.7 Hz, 6H). ESI-MS *m/z* calc 479.11 , found 480.2 $(M+1)^+$; LCMS retention time: 1.8 minutes (3 min run).

Step 3: *N*-[(6-amino-2-pyridyl)sulfonyl]-3-fluoro-4-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)benzamide (Compound 1497)




To mixture of 2,2,4-trimethylpyrrolidine (35.41 mg, 0.3128 mmol) and *N*-[(6-amino-2-pyridyl)sulfonyl]-2,3-difluoro-4-(3-fluoro-5-isobutoxy-phenyl)benzamide (30.00 mg, 0.06257 mmol) was added 1-methylpyrrolidin-2-one (0.3 mL) and the reaction mixture was heated 190

°C for 2 hours. The reaction mixture was cooled and purified by reverse phase HPLC utilizing a gradient of 10 to 99 percent acetonitrile in water containing 5 mM ammonium formate to yield *N*-[(6-amino-2-pyridyl)sulfonyl]-3-fluoro-4-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)benzamide (Compound 1497) (7.6 mg, 0.013 mmol, 4.2%) ^{1}H NMR (400 MHz, Methanol-d_4) δ 8.03 (dd, J = 8.3, 1.2 Hz, 1H), 7.68 (dd, J = 8.5, 7.3 Hz, 1H), 7.59 (dd, J = 8.3, 7.1 Hz, 1H), 7.38 (dd, J = 7.4, 0.8 Hz, 1H), 6.93 – 6.82 (m, 2H), 6.82 – 6.74 (m, 2H), 3.92 (t, J = 9.4 Hz, 1H), 3.79 (d, J = 6.4 Hz, 2H), 2.84 – 2.72 (m, 1H), 2.23 (dd, J = 12.6, 7.3 Hz, 1H), 2.12 – 2.02 (m, 1H), 1.87 (t, J = 11.7 Hz, 1H), 1.38 (s, 3H), 1.24 (d, J = 6.6 Hz, 6H), 1.04 (d, J = 6.7 Hz, 6H). ESI-MS *m/z* calc. 572.23, found 573.1 $(M+1)^+$; LCMS retention time: 1.77 minutes (3 min run).

Preparation 87: *N*-[(6-amino-2-pyridyl)sulfonyl]-5-chloro-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1641)


N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (56.79 mg, 0.1022 mmol) and *N*-chlorosuccinimide (13.65 mg, 0.1022 mmol) in dichloromethane (400 µL) was stirred at reflux overnight. The dichloromethane was evaporated under a stream of nitrogen, and the residue was dissolved in dimethyl sulfoxide and purified by preparative HPLC chromatography utilizing a gradient of 10 to 99 percent acetonitrile in water to give *N*-[(6-amino-2-pyridyl)sulfonyl]-5-chloro-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1641) (18 mg, 0.030 mmol, 29.5%) ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.59 (s, 1H), 7.80 (s, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.27 - 7.18 (m, 1H), 7.10 - 6.99 (m, 2H), 6.93 (dt, J = 11.0, 2.3 Hz, 1H), 6.70 (dd, J = 8.4, 0.8 Hz, 1H), 6.54 (s, 2H), 3.80 (d, J = 6.6 Hz, 2H), 2.65 - 2.53 (m, 2H), 2.22 (dt, J = 11.5, 5.9 Hz, 1H), 2.08 - 1.96 (m, 1H), 1.84 (dd, J = 12. 0 , 5.6 Hz, 1H), 1.57 - 1.33 (m, 7H), 0.97 (d, J = 6.7 Hz, 6H), 0.86 (d, J = 6.3 Hz, 3H). ESI-MS *m/z* calc. 589.19257, found 590.0 $(M+1)^+$; Retention time: 3.06 minutes.

Preparation 88: *N*-[(6-amino-2-pyridyl)sulfonyl]-5-chloro-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1603)



N-[(6-amino-2-pyridyl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 946) (40.0 mg, 0.102 mmol) and *N*-chlorosuccinimide (13.65 mg, 0.1022 mmol) in dichloromethane (400 µL) was stirred at reflux for 3 days. The reaction mixture was directly purified by chromatography over silica gel (4g) with a linear gradient of 0 to 5% methanol in dichloromethane to give *N*-[(6-amino-2-pyridyl)sulfonyl]-5-chloro-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1603) (25 mg, 0.058 mmol, 57%) ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.54 (s, 1H), 8.27 - 8.00 (m, 1H), 7.72 - 7.54 (m, 2H), 7.18 (d, J = 7.2 Hz, 1H), 6.66 (d, J = 8.4 Hz, 1H), 6.50 (s, 2H), 2.55 (s, 2H), 2.17 (dq, J = 13.0, 7.1, 6.7 Hz, 1H), 1.83 (dd, J = 11.8, 5.6 Hz, 1H), 1.47 (d, J = 7.1 Hz, 6H), 1.37 (t, J = 12.1 Hz, 1H), 0.85 (d, J = 6.3 Hz, 3H). ESI-MS *m/z* calc. 423.1132, found 424.0 $(M+1)^+$; Retention time: 1.75 minutes.


Preparation 89: *N*-[(6-acetamido-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1652)



N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (56.78 mg, 0.1022 mmol) in pyridine (161.7 mg, 165.3 µL, 2.044 mmol) was treated with acetic anhydride (104.3 mg, 96.40 µL, 1.022 mmol) and heated to 45 °C for 2.5 hours. The reaction mixture was diluted with a 9:1 (v/v) mixture of dimethyl sulfoxide in water and purified by preparative HPLC chromatography to give *N*-[(6-acetamido-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1652) (22 mg, 0.036 mmol, 35%)


^{1}H NMR (400 MHz, DMSO-d_6) δ 12.68 (s, 1H), 10.87 (s, 1H), 8.35 (d, J = 8.4 Hz, 1H), 8.17 - 8.05 (m, 1H), 7.88 - 7.80 (m, 2H), 7.51 - 7.45 (m, 1H), 7.42 (ddd, J = 10.2, 2.4, 1.3 Hz, 1H), 7.30 (d, J = 8.0 Hz, 1H), 6.90 (dt, J = 10.7, 2.3 Hz, 1H), 3.83 (d, J = 6.6 Hz, 2H), 2.60 (dd, J = 10.2, 7.2 Hz, 1H), 2.52 (d, J = 5.3 Hz, 1H), 2.14 (s, 4H), 2.07 - 2.00 (m, 1H), 1.87 (dd, J = 11.8, 5.6 Hz, 1H), 1.59 (d, J = 6.0 Hz, 6H), 1.41 (t, J = 12.1 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.77 (d, J = 6.3 Hz, 3H). ESI-MS m/z calc. 597.2421, found 598.0 $(M+1)^+$; Retention time: 2.83 minutes.

Preparation 90: *N*-((6-aminopyridin-2-yl)sulfonyl)-6-chloro-2-(3-fluoro-5-isobutoxyphenyl)nicotinamide




A mixture of *N*-((6-aminopyridin-2-yl)sulfonyl)-2,6-dichloronicotinamide (2.6 g, 7.4 mmol) and 2-(3-fluoro-5-isobutoxyphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.3 g, 7.7 mmol) in 12 mL of *N,N*-dimethylformamide and 3 mL of water was heated under nitrogen to 70 °C for 30 minutes and then tetrakis(triphenylphosphine)palladium(0) (0.20 g, 0.17 mmol) and sodium carbonate (4.8 g, 45 mmol) were added to the reaction mixture. The mixture was stirred at 70 °C for 18 hours and then purified by silica gel chromatography eluting with a gradient of 40 to 100% ethyl acetate in hexanes. The product was further purified on reverse phase HPLC and a white solid was isolated as *N*-((6-aminopyridin-2-yl)sulfonyl)-6-chloro-2-(3-fluoro-5-isobutoxyphenyl)nicotinamide (0.37 g, 0.77 mmol, 14%). ^{1}H NMR (DMSO-d_6, 250 MHz): δ 8.03 (d, J = 8.3 Hz, 1H), 7.62 (m, 2H), 7.11 (d, J = 7.3 Hz, 1H), 6.93 (m, 2H), 6.81 (d, J = 9.3 Hz, 1H), 6.73 (d, J = 8.5 Hz, 1H), 3.87 (d, J = 6.5 Hz, 2H), 2.02 (m, 1H), 1.00 (d, J = 6.5 Hz, 6H) ppm. LC-MS: $(M+H)^+$ = 479.1

Preparation 91: *N*-((6-aminopyridin-2-yl)sulfonyl)-6-chloro-6'-isopropoxy-[2,3'-bipyridine]-3-carboxamide



A mixture of the sulfonamide *N*-((6-aminopyridin-2-yl)sulfonyl)-2,6-dichloronicotinamide (2.1 g, 6.0 mmol) and 2-isopropoxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (1.7 g, 6.6 mmol) in 12 mL of *N,N*-dimethylformamide and 3 mL of water was heated under nitrogen at 70 °C for 20 minutes and added tetrakis(triphenylphosphine)palladium(0) (0.15 g, 0.13 mmol) and sodium carbonate (1.9 g, 18 mmol). The mixture was stirred at 70 °C for 19 hours. The reaction mixture was evaporated to dryness, and the crude product was purified by silica gel chromatography utilizing a gradient of 40 to 100% ethyl acetate in hexanes. The product was further purified on reverse phase HPLC and a white solid was isolated as *N*-((6-aminopyridin-2-yl)sulfonyl)-6-chloro-6'-isopropoxy-[2,3'-bipyridine]-3-carboxamide (0.72 g, 1.6 mmol, 27%). ^{1}H NMR (DMSO-d_6, 250 MHz): δ 8.33 (s, 1H), 8.06 (d, *J* = 8.0 Hz, 1H), 7.64 (m, 3H), 7.15 (d, J = 7.3 Hz, 1H), 6.93 (m, 2H), 6.80 (d, *J* = 7.3 Hz, 1H), 6.69 (d, *J* = 8.8 Hz, 1H), 5.27 (m, 1H), 1.32 (d, *J* = 6.3 Hz, 6H) ppm.

Preparation 92: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(6-isopropoxy-3-pyridyl)-6-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1436)



N-((6-aminopyridin-2-yl)sulfonyl)-6-chloro-6'-isopropoxy-[2,3'-bipyridine]-3-carboxamide (0.050 g, 0.11mmol) and cesium floride (0.085 g, 0.56 mmol) were combined in dimethyl sulfoxide (0.5 mL). The reaction mixture was stirred at 150 °C for 16 hours. (4S)-2,2,4-trimethylpyrrolidine (0.0379 g, 0.335 mmol) was added to the reaction mixture and the mixture was stirred at 150 °C for 10 hours. The crude product was purified by reverse phase HPLC utilizing a gradient of 10 to 99% acetonitrile in water to yield *N*-[(6-amino-2-

pyridyl)sulfonyl]-2-(6-isopropoxy-3-pyridyl)-6-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1436) (0.0026 g, 0.0050 mmol, 4.5%). ESI-MS *m/z* calc. 424.2, found 525.3 $(M+1)^+$; Retention time: 1.89 minutes.

The following compounds can be synthesized using the procedures described herein

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-fluoro-5-isobutoxy-phenyl)-6-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1409),

N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-fluoro-5-isobutoxy-phenyl)-6-[(4*R*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1385),


N-[(6-amino-2-pyridyl)sulfonyl]-2-(6-isopropoxy-3-pyridyl)-6-[(4*R*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1491).

Preparation 93: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-cyano-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1564)



[(2-Di-*tert*-butylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)-2-(2'-amino-1,1'-biphenyl)] palladium(II) methanesulfonate (4.186 mg, 0.005270 mmol) , di-*tert*-butyl-[2-(2,4,6-triisopropylphenyl)phenyl]phosphane (2.238 mg, 0.005270 mmol), Potassium hexacyanoferrate(II) trihydrate (22.26 mg, 0.05270 mmol), and *N*-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 852) (44.7 mg, 0.1054 mmol) were added to a vial and the vial was purged with nitrogen. Deoxygenated dioxane (500 µL) and a solution of potassium carbonate (263.6 µL of 0.05 M, 0.01318 mmol) was added to reaction mixture, the head space was purged with nitorgen and the sealed vial was stirred at 100 °C for 1 hour. The crude product was purified by reverse phase HPLC utilizing a gradient of 10 to 99% acetonitrile in water to yield *N*-[(6-amino-2-pyridyl)sulfonyl]-6-cyano-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1564) (1.8 mg, 0.0043 mmol, 4.1%) ESI-MS *m/z* calc. 414.1474, found 415.2 $(M+1)^+$; Retention time: 1.49 minutes.


Preparation 94: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)-4-hydroxy-pyridine-3-carboxamide (Compound 1430)



In a nitrogen purged flask, *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)-4-methoxy-pyridine-3-carboxamide (Compound 1254) (17.5 mg, 0.0306 mmol) was dissolved in dichloromethane (1.577 mL). The mixture was cooled to -78°C, and tribromoborane (811.2 mg, 306.1 µL of 1 M, 0.3061 mmol) was added to the reaction mixture in a dropwise manner. The mixture was slowly warmed to room temperature. The mixture was stirred overnight at ambient temperature. The reaction mixture was added to ice (~ 5 mL). The layers were separated, and the aqueous layer was extracted with dichloromethane. The combined organic layers were dried over sodium sulfate, concentrated, and purified by reverse phase HPLC utilizing a gradient of 1-99% acetonitrile in 5 mM aqueous hydrochloric acid. The compound was further puried on silica gel using dichloromethane as the eluent to yield *N*-[(6-amino-2-pyridyl)sulfonyl]-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)-4-hydroxy-pyridine-3-carboxamide (Compound 1430) (3.2 mg, 0.0057 mmol, 19%) as an off -white solid. ESI-MS *m/z* calc. 557.2108, found 558.3 $(M+1)^+$, Retention time: 1.84 minutes.


Preparation 95: 4-Bromo-2-(mesityloxy)benzoic acid

Step 1: 4-Bromo-2-(mesityloxy)benzonitrile



A mixture of 4-bromo-2-fluoro-benzonitrile (1.2 g, 6.0 mmol), 2,4,6-trimethylphenol (780 mg, 6.1 mmol), potassium carbonate (2.50 g, 18.1 mmol), and dimethylsulfoxide (9.0 mL) were microwave irradiated at 140 °C for 2 hours. Water (20 mL) was added and the resulting mixture was stirred for 5 minutes. The resulting suspension was filtered and the solid was washed with water and dried under vacuum at 50 °C for 16 hours to yield 4-bromo-2-(mesityloxy)benzonitrile (1.8 g, 95%) as a brown solid. ^{1}H NMR ($CDCl_3$, 250 MHz): 7.50 (d, J = 8.2 Hz, 1H), 7.20 (d, J = 8.2 Hz, 1H), 6.94 (s, 2H), 6.61 (s, 1H), 2.33 (s, 3H), 2.10 (s, 6H) ppm.


Step 2: 4-Bromo-2-(mesityloxy)benzoic acid



4-bromo-2-(mesityloxy)benzonitrile (1.8 g, 9.0 mmol) was suspended in ethanol (6 mL) and water (12 mL), containing sodium hydroxide (3.0 g, 75 mmol). The mixture was stirred at 90 °C for 16 hours. Hydrochloric acid (6M, 13 mL) was added to the reaction mixture followed by water (20 mL) and the suspension was stirred for 10 minutes, filtered, and washed with water and hthen hexanes to yield 4-bromo-2-(mesityloxy)benzoic acid as an off-white solid (1.7 g, 5.1 mmol, 57%). ^{1}H NMR (DMSO-d_6, 250 MHz): 7.74 (d, J = 8.2 Hz, 1H), 7.28 (d, J = 8.2 Hz, 1H), 7.01 (s, 2H), 6.41 (s, 1H), 2.28 (s, 3H), 2.02 (s, 6H) ppm.

Preparation 96: 6-Methoxy-4-methylpyridine-2-sulfonamide


Step 1: 2-Benzylsulfanyl-6-chloro-4-methylpyridine



2,6-Dichloro-4-methyl-pyridine (4.74 g, 29.3 mmol) was dissolved in tetrahydrofuran (47.40 mL) with phenylmethanethiol (3.634 g, 3.435 mL, 29.26 mmol) and 2-methylpropan-2-olate (Sodium Ion (1)) (29.26 mL of 2 M, 58.52 mmol) was added to the reaction mixture. The


reaction was stirred for 16 hours and partitioned between water and ethyl acetate. The organics were separated, washed with brine, dried over sodium sulfate and evaporated. The crude material was used without fu rther purification. 2-Benzylsulfanyl-6-chloro-4-methyl-pyridine (6.91 g, 27.7 mmol, 94.5%) ESI-MS *m/z* calc. 249.0379, found 250.0 $(M+1)^+$; Retention time: 0.78 minutes.

Step 2: 6-Chloro-4-methylpyridine-2-sulfonamide




A biphasic mixture of 2-benzylsulfanyl-6-chloro-4-methyl-pyridine (6.87 g, 27.5 mmol) in dichloromethane (41.22 mL), water (13.74 mL), and aqueous hydrochloric acid (2.751 mL of I M, 2.751 mmol) was cooled in a salt water ice bath. (Note: Aqueous hydrochloric acid is added to avoid an otherwise delayed exotherm upon the chlorine addition.). Chlorine (7.800 g, 110.0 mmol) was bubbled through the solution (3 times at 2 minute intervals) over an hour. The reaction was stirred at 0 °C. After 1.5 hours (reaction monitored by UPLC) the reaction was added dropwise to an ice bath cooled solution of ammonium hydroxide (41.32 mL of 28 %w/v, 330.1 mmol) . The reaction was warmed to room temperature and stirred for 15 minutes. The reaction mixture was partitioned between ethyl acetate (60 mL) and water (20 mL). The organics were separated, and the aqueous layer was extracted twice with ethyl acetate (60 mL portions). The organics were combined, washed with brine, dried over sodium sulfate and evaporated. The resulting solid was triturated with dichloromethanr (30 mL) and filtered. The solid was washed with minimal dichloromethane to give 6-chloro-4-methyl-pyridine-2-sulfonamide as an off white solid (3.18 g, 15.4 mmol, 56.0%) ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.78 (dd, J = 1.3, 0.7 Hz, 1H), 7.66 - 7.63 (m, 1H), 7.59 (s, 2H), 2.45 (s, 3H). ESI-MS *m/z* calc. 205.99167, found 206.9 $(M+1)^+$; Retention time: 0.3 minutes .

Step 3: 6-Methoxy-4-methylpyridine-2-sulfonamide



6-Chloro-4-methylpyridine-2-sulfonamide (500. mg, 2.42 mmol) was dissolved in sodium methoxide in methanol (2.537 g, 2.615 mL of 25 % w/v, 12.10 mmol) and heated at 65 °C. After 1 hour, all starting material was gone and two products had formed. The reaction mixture was partitioned between ethyl acetate and a saturated aqueous ammonium chloride solution. The organics were separated, washed with water, brine, dried over sodium sulfate and evaporated to dryness. The crude product was purified by silica gel chromatography eluting with 0 to 100% ethyl acetate in hexanes to give 6-methoxy-4-methylpyridine-2-sulfonamide (126 mg, 0.623 mmol, 25.7%) ESI-MS *m/z* calc. 202.04121, found 203.0 $(M+1)^+$; Retention time: 0.32 minutes.


Preparation 97: 6-Amino-4-methylpyridine-2-sulfonamide



6-Chloro-4-methylpyridine-2-sulfonamide (500. mg, 2.42 mmol) was dissolved in aqueous ammonium hydroxide (3.029 mL of 28 %w/v, 24.20 mmol) and irradiated in a microwave reactor for 1 hour at 150 °C. The reaction was evaporated and the solid was washed with water and methanol to remove excess ammonium hydroxide. The resulting solid was a mixture of product and starting material. The mixture was taken on to the next step without further purification. 6-amino-4-methylpyridine-2-sulfonamide (200. mg, 1.07 mmol, 44.2%) ESI-MS *m/z* calc. 187.04155, found 188.0 $(M+1)^+$; Retention time: 0.21 minutes.

PREPARATION 98: 4,4,5,5-Tetramethyl-2-(3,3,5,5-tetramethylcyclopenten-1-yl)-1,3,2-dioxaborolane

Step 1: 3,3,5-trimethyl-5-trimethylstannyl-cyclohexanone




To a stirring solution of trimethyl(trimethylstannyl)stannane (3.555 g, 2.250 mL, 10.85 mmol) in tetrahydrofuran (50 mL) at -20 °C was added methyllithium (4.964 g, 6.781 mL of 1.6 M, 10.85 mmol) and the mixture was stirred at -20°C for 15minutes. Next the 3,5,5-trimethylcyclohex-2-en-1-one (1.00 g, 7.24 mmol) was slowly added and the reaction mixture was warmed to 0°C, stirred 30 minutes, and then quenched with methanol (695.6 mg, 879.4 µL, 21.71 mmol). The reaction mixture was warmed to room temperature and diluted with water then extracted twice with diethyl ether. The combined organic layers were dried (sodium sulfate), filtered and concentrated to a yellow oil which was purified by silica gel chromatography using 100% hexanes as eluent followed by 100% dichloromethane as eluent to yield 3,3,5-trimethyl-5-trimethylstannyl-cyclohexanone (2.01 g, 6.63 mmol, 91.7%) as a clear oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 2.59 - 2.45 (m, 1H), 2.21 - 2.12 (m, 3H), 1.92 (d, J = 14.2 Hz, 1H), 1.59 (d, J = 14.2 Hz, 1H), 1.28 (s, 3H), 1.07 (s, 3H), 1.00 (s, 3H), 0.13 (d, J = 1.1 Hz, 1H), 0.07 (s, 9H), 0.01 (d, J = 1.1 Hz, 1H).

Step 2: 2,2,4,4-Tetramethylcyclopentanone



To a stirring solution of 3,3,5-trimethyl-5-trimethylstannyl-cyclohexanone (4.17 g, 13.8 mmol) in the dichloromethane (79.23 mL) under a nitrogen atmosphere at 0 °C was added (dropwise) tetrachlorotitanium (2.610 g, 2.183 mL, 13.76 mmol) giving a dark brown color on addition. Upon completion of addition the reaction mixture was a brown suspension which was stirred 20 minutes at 0 °C and then quenched by the addition of a solution of saturated aqueous sodium bicarbonate (Caution: Gas evolution, reaction became biphasic with a purple suspension as the aqueous layer). The resulting mixture was extracted twice with dichloromethane, dried (magnesium sulfate), filtered and carefully concentrated to a clear oil. Purified by silica gel chromatography eluting with 100% giving 2,2,4,4-tetramethylcyclopentanone (1.67 g, 11.9 mmol, 86.6%) as a clear oil.


Step 3: 4,4,5,5-Tetramethyl-2-(3,3,5,5-tetramethylcyclopenten-1-yl)-1,3,2-dioxaborolane



To 2,2,4,4-tetramethylcyclopentanone (1.67 g, 11.9 mmol) was added tetrahydrofuran (16.70 mL) and the reaction was chilled to -78 °C. The reaction was stirred for 10 minutes and lithium bis(trimethylsilyl)amide (14.29 mL of 1 M, 14.29 mmol) was added dropwise and the reaction was allowed to stir at -78 °C for 1 hour. 1,1,1-Trifluoro-*N*-phenyl-*N*-(trifluoromethylsulfonyl)methanesulfonamide (5.105 g, 14.29 mmol) dissolved in tetrahydrofuran (12.5 mL) was added dropwise and the reaction was allowed to warm to room temperature and stir 16 hours. The reaction was quenched with a saturated aqueous solution of sodium chloride and extracted with ethyl acetate. The organic layer was dried over sodium sulfate and evaporated to an orange syrup. The crude orange syrup was added to a mixture of bis(pinacol)diboron (4.535 g, 17.86 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (871.5 mg, 1.191 mmol) and potassium acetate (3.507 g, 35.73 mmol) in dimethylformamide (16.70 mL). The reaction was purged with nitrogen and heated to 65 °C for 16 hours. The reaction was quenched with a saturated aqueous solution of sodium chloride and extracted with ethyl acetate. The organic layer was dried over sodium sulfate and evaporated to dryness. The crude product was purified by silica gel chromatography using a gradient from 100% hexanes to 40% ethyl acetate in hexanes giving 4,4,5,5-tetramethyl-2-(3,3,5,5-tetramethylcyclopenten-1-yl)-1,3,2-dioxaborolane (651 mg, 2.60 mmol, 21.8%) as a white solid.


PREPARATION 99: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxamide (Compound 2492) (Isomer 1) (Compound 2374) (Isomer 2)

Step 1: ethyl 2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylate



2-Chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylic acid (5.077 g, 15.53 mmol) was dissolved in ethyl alcohol (50.26 mL) and thionyl chloride (2.218 g, 1.360 mL, 18.64 mmol) was added dropwise. The resulting solution was slowly heated to 76 °C (reflux) and stirred at 76 °C for 5 hours. The brownish solution was concentrated to remove solvent. The crude residue was redissolved in ethyl acetate and extracted with saturated aqueous sodium bicarbonate. The aqueous phase was extracted twice with ethyl acetate. The combined organic layers were extracted with saturated aqueous brine, dried over sodium sulfate, concentrated and purified by silica gel chromatography using a gradient from 100% hexanes to 100% ethyl acetate giving ethyl 2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylate (3.64 g, 10.3 mmol, 66.6%) as a white solid.


Step 2: Ethyl 6-(3-fluoro-5-isobutoxy-phenyl)-2-(3,3,5,5-tetramethylcyclopenten-1-yl)pyridine-3-carboxylate



A mixture of ethyl 2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylate (841.8 mg, 2.393 mmol), 4,4,5,5-tetramethyl-2-(3,3,5,5-tetramethylcyclopenten-1-yl)-1,3,2-dioxaborolane (598.7 mg, 2.393 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (525.3 mg, 0.7179 mmol), and sodium carbonate (760.9 mg, 7.179 mmol) in dioxane (10.77 mL) and water (1.196 mL) was heated in a sealed vessel at 150 °C for 16 hours. The reaction was filtered and the solids were washed with ethyl acetate. The extract was evaporated and the residue was purified by silica gel chromatography using a gradient from 100% hexanes to 25% ethyl acetate


in hexanes to give ethyl 6-(3-fluoro-5-isobutoxy-phenyl)-2-(3,3,5,5-tetramethylcyclopenten-1-yl)pyridine-3-carboxylate (460.3 mg, 1.047 mmol, 43.75%) as a light yellow solid.

Step 3: 6-(3-Fluoro-5-isobutoxy-phenyl)-2-(3,3,5,5-tetramethylcyclopenten-1-yl)pyridine-3-carboxylic acid



A solution of ethyl 6-(3-fluoro-5-isobutoxy-phenyl)-2-(3,3,5,5-tetramethylcyclopenten-1-yl)pyridine-3-carboxylate (460.3 mg, 1.047 mmol) and sodium hydroxide (2.617 mL of 1 M, 2.617 mmol) in tetrahydrofuran (2.606 mL) and water (2.606 mL) was stirred at 50 °C for 6.5 hours. The reaction was cooled to room temperature and then neutralized with 1M hydrochloric acid and extracted twice with ethyl acetate. The combined extracts were dried over sodium sulfate and evaporated to give 6-(3-fluoro-5-isobutoxy-phenyl)-2-(3,3,5,5-tetramethylcyclopenten-1-yl)pyridine-3-carboxylic acid (389.1 mg, 0.9455 mmol, 90.32%) as yellow needles.


Step 4: 6-(3-Fluoro-5-isobutoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxylic acid



A mixture of 6-(3-fluoro-5-isobutoxy-phenyl)-2-(3,3,5,5-tetramethylcyclopenten-1-yl)pyridine-3-carboxylic acid (389.1 mg, 0.9455 mmol), palladium hydroxide (79.67 mg, 0.5673 mmol) and ammonium formate (1.192 g, 18.91 mmol) in ethanol (10 mL) were combined in a microwave vial and heated to 85 °C for 3hours. The reaction mixture was cooled to room temperature, uncapped, ammonium formate (1.192 g, 18.91 mmol) and palladium hydroxide (79.67 mg, 0.5673 mmol) were added, nitrogen was bubbled through the solution for 2 minutes,

and the vial was capped and returned to the microwave reactor at 85°C for 4 additional hours. The reaction was filtered through celite eluting with methanol and concentrated under reduced pressure. The mixture was then diluted with 1N aqueous hydrochloric acid and washed with twice with ethyl acetate. The organic layers were combined, dried (sodium sulfate), filtered and concentrated to a clear oil with some white solid present. The crude material was purified by silica gel chromatography using a gradient from 100% dichloromethane to 20% methanol in dichloromethane followed by 20% methanol in ethyl acetate giving 6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxylic acid (52.3 mg, 0.126 mmol, 13.4%) as a white solid.

Step 5: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxamide (Compound 2492) (Isomer 1) (Compound 2374) (Isomer 2)



A solution of 6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxylic acid (52.0 mg, 0.126 mmol) and carbonyl diimidazole (30.58 mg, 0.1886 mmol) in dimethylformamide (0.5 mL) was stirred at 45°C for 90 minutes. A separate solution of 6-aminopyridine-2-sulfonamide (43.54 mg, 0.2514 mmol) and sodium hydride (10.06 mg, 0.2514 mmol) in dimethylformamide (0.5 mL) which was stirred at 45 °C for 45 minutes was then added and the resulting mixture was stirred at 45 °C for 2hours. The reaction was filtered and purified using a reverse phase HPLC-MS method using a Luna C18 (2) column (75 x 30 mm, 5 μm particle size) sold by Phenomenex (pn: 00C-4252-U0-AX), and a dual gradient run from 1 to 99% mobile phase B over 15.0 minutes (mobile phase A = H_2O (5 mM HCl), mobile phase B = acetonitrile, flow rate = 50 mL/min, injection volume = 950 μL and column temperature = 25 °C) giving *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxamide (Compound 2492) (Isomer 1) (Compound 2374) (Isomer 2) hydrochloride (51 mg, 0.084 mmol, 67%) as an off-white solid ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.63 (s, 1H), 7.91 (dd, J = 20.4, 8.2 Hz, 2H), 7.66

(t, J = 7.9 Hz, 1H), 7.57 (s, 1H), 7.51 (d, J = 9.8 Hz, 1H), 7.25 (d, J = 7.3 Hz, 1H), 6.93 (d, J = 10.7 Hz, 1H), 6.73 (d, J = 8.4 Hz, 1H), 6.51 (s, 2H), 3.86 (d, J = 6.6 Hz, 2H), 3.58 (dd, J = 12.2, 6.5 Hz, 1H), 2.55 (d, J = 12.3 Hz, 1H), 2.06 (dt, J = 13.2, 6.6 Hz, 1H), 1.63 (dd, J = 12.4, 6.4 Hz, 1H), 1.50 (d, J = 13.0 Hz, 1H), 1.39 (d, J = 12.9 Hz, 1H), 1.15 (s, 3H), 1.08 - 0.93 (m, 9H), 0.80 (s, 3H), 0.69 (s, 3H). ESI-MS *m/z* calc. 568.25195, found 569.3 $(M+1)^+$; Retention time: 2.4 minutes.


PREPARATION 100: 4,4,5,5-tetramethyl-2-[(5*S*)-3,3,5-trimethylcyclohexen-1-yl]-1,3,2-dioxaborolane

Step 1: (5*R*)-3,3,5-Trimethylcyclohexanone



To a round bottom flask purged with nitrogen was added copper (II) acetate (14.53 mg, 0.08000 mmol), [(4*S*)-4-[5-bis(3,5-di*tert*-butyl-4-methoxy-phenyl)phosphanyl-1,3-benzodioxol-4-yl]-4,5,6,7-tetrahydro-1,3-benzodioxol-5-yl]-bis(3,5-di*tert*-butyl-4-methoxy-phenyl)phosphane (47.34 mg, 0.04000 mmol), and toluene (27.64 mL). The reaction was stirred at room temperature for 90 minutes. Polymethylhydrosiloxane (4.8 mL) was added and the reaction was allowed to stir at room temperature for 30 minutes. 3,5,5-Trimethylcyclohex-2-en-1-one (5.528 g, 6.009 mL, 40.00 mmol) was then added dropwise and the reaction was allowed to stir for 90 minutes at room temperature. The reaction was poured into a solution of 3M sodium hydroxide and allowed to stir vigorously for 2 hours. The crude reaction was extracted with ethyl acetate and dried over sodium sulfate. The solvent was removed and the crude product was taken onto the next step without further purification to give (5*R*)-3,3,5-trimethylcyclohexanone (5.1 g, 36 mmol, 91%) ESI-MS *m/z* calc. 140.12012, found 141.2 $(M+1)^+$; Retention time: 1.44 minutes was a pale oil.


Step 2: 4,4,5,5-tetramethyl-2-[(5*S*)-3,3,5-trimethylcyclohexen-1-yl]-1,3,2-dioxaborolane



To crude (5*R*)-3,3,5-trimethylcyclohexanone (2.85 g, 20.3 mmol) was added tetrahydrofuran (28.50 mL) and the reaction was chilled to -78 °C. The reaction was stirred for 10 minutes and lithium bis(trimethylsilyl)amide (24.38 mL of 1 M, 24.38 mmol) was added dropwise and the reaction was allowed to stir at -78 °C for 1 hour. 1,1,1-Trifluoro-*N*-phenyl-*N*-(trifluoromethylsulfonyl)methanesulfonamide (8.710 g, 24.38 mmol) dissolved in tetrahydrofuran (20 mL) was added dropwise and the reaction was allowed to warm to room temperatre and stir for 16 hours. The reaction was quenched with brine and extracted with ethyl acetate. The organic layer was dried over sodium sulfate and evaporated to a yellow oil. The crude vinyl triflate was used in the next step without further purification. The crude vinyl triflate was added to a mixture of bis(pinacol)diboron (7.740 g, 30.48 mmol), bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (1.487 g, 2.032 mmol), potassium acetate (5.983 g, 60.96 mmol) in *N.N*-dimethylformamide (28.50 mL). The reaction was purged with nitrogen and heated to 65 °C for 16 hours. The reaction was quenched with brine and extracted with ethyl acetate. The organic layer was dried over sodium sulfate and evaporated. The crude product was purified by silica gel chromatography utilizing a gradient of 0 to 40% ethyl acetate in hexanes to yield 4,4,5,5-tetramethyl-2-[(5*S*)-3,3,5-trimethylcyclohexen-1-yl]-1,3,2-dioxaborolane (2.2 g, 8.8 mmol, 43%) as a mixture of regioisomers ESI-MS *m/z* calc. 250.2104, found 251.6 $(M+1)^+$; Retention time: 2.17 minutes


PREPARATION 101: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[(1*S*,2*S*)-2-(methoxymethyl)cyclopropyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1575)

Step 1: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[(1*S*,2*S*)-2-(methoxymethyl)cyclopropyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1575) and *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[(1*R*,2*R*)-2-(methoxymethyl)cyclopropyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide



To a microwave vial was added *N*-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 852) trifluoroacetic acid salt (100 mg, 0.186 mmol) , [1,3-Bis(2,6-Diisopropylphenyl)imidazol-2-ylidene](3-chloropyridyl)palladium(II) dichloride (6.3 mg, 0.0093 mmol) and dioxane (600 µL) . The reaction was purged with nitrogen for 2 minutes and *trans* 2-[(1*S*,2*S*)-2-(methoxymethyl)cyclopropyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (68.99 mg, 0.325 mmol) was added followed by an aqueous 2M solution of potassium carbonate (279 µL of 2 M, 0.558 mmol). Then the reaction was irradiated to 120 °C over 16 hours in the microwave reactor. The reaction mixture was allowed to cool, filtered and then purified directly by reverse-phase preparative chromatography utilizing a C18 column and HPLC-MS method 1 to 99 acetonitrile in water containing 5 mM hydrochloric acid to afford two compounds. Compound 1 (2.26 mg, 0.0044 mmol, 2.39%) ESI-MS *m/z* calc. 473.2097, found 474.2 $(M+1)^+$; Retention time: 1.13 minutes. Compound 2 (0.86 mg, 0.0018 mmol, 0.97%) ESI-MS *m/z* calc. 473.2097, found 474.2 $(M+1)^+$; Retention time: 1.26 minutes as a light yellow solid.

Preparation 102: *N*-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-chloro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide




Step 1: 6-chloro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxylic acid

6-chloro-2-fluoro-pyridine-3-carboxylic acid (1.50 g, 8.545 mmol) and (4S)-2,2,4-trimethylpyrrolidine (Hydrochloric Acid) (1.534 g, 10.25 mmol) were combined in DMSO (4.500 mL). Potassium carbonate (2.362 g, 17.09 mmol) was added. Note: Substantial bubbling was observed upon mixing. The reaction mixture was sealed and heated at 90 °C overnight. The


reaction mixture was diluted with EtOAc (50 mL) and washed with aqueous 0.5 N HCl. The organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure to a yellow oil. To remove any residual solvents, the oil was subjected to a stream of nitrogen gas for 1 hour, resulting in a crystalline solid. The product was used in the next step without further purification. ESI-MS *m/z* calc. 268.09787, found 269.2 $(M+1)^+$; Retention time: 1.49 minutes

Step 2: tert-butyl *N*-[(3S)-1-[[6-chloro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carbonyl]sulfamoyl]pyrrolidin-3-yl]carbamate




6-chloro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxylic acid (1.61 g, 5.99 mmol) was dissolved in dichloromethane (8 mL), warmed to 40 °C and *N*-(oxomethylene)sulfamoyl chloride (1.017 g, 625.5 µL, 7.189 mmol) was added. Note: Substantial bubbling was observed during addition. After 3 minutes of stirring at 40 °C, additional dichloromethane (10 mL) was added followed by *tert*-butyl *N*-[(3*S*)-pyrrolidin-3-yl]carbamate (2.80 g, 15.0 mmol). Bubbling was observed. After overnight stirring at 40 °C, the reaction mixture was diluted with EtOAc (75 mL) and washed with water and brine. The organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure. The product was isolated by silica gel column chromatography: 0 to 40% EtOAc/hexane gradient on a 80 gram silica gel column. tert-butyl *N*-[(3S)-1-[[6-chloro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carbonyl]sulfamoyl]pyrrolidin-3-yl]carbamate (965 mg, 1.87 mmol, 31.2%) was obtained as a lightly brown foaming solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 11.92 (s, 1H), 7.62 (d, J = 7.9 Hz, 1H), 7.18 (s, 1H), 6.61 (d, J = 7.9 Hz, 1H), 3.96 (s, 1H), 3.65 - 3.45 (m, 3H), 3.14 (dd, J = 9.7, 5.9 Hz, 1H), 3.03 (t, J = 10.3 Hz, 1H), 2.97 - 2.85 (m, 1H), 2.33 (s, 1H), 2.04 (dd, J = 12.7, 5.7 Hz, 1H), 1.92 (dd, J = 11.4, 5.1 Hz, 1H), 1.78 (dd, J = 12.6, 6.9 Hz, 1H), 1.54 (d, J = 6.1 Hz, 6H), 1.38 (s, 9H), 1.01 (d, J = 6.3 Hz, 3H), 0.83 (s, 1H). ESI-MS m/z calc. 515.1969, found 516.4 $(M+1)^+$; Retention time: 0.77 minutes.

Step 3: *N*-[(3S)-3-aminopyrrolidin-1-yl]sulfonyl-6-chloro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide



tert-butyl *N*-[(3S)-1-[[6-chloro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carbonyl]sulfamoyl]pyrrolidin-3-yl]carbamate (100 mg, 0.1938 mmol) was dissolved in CH2Cl2 (1 mL), and TFA (1 mL, 12.98 mmol) was added. The solution was allowed to stir at room temperature for 30 minutes. Volatiles were removed under reduced pressure. The crude product was used in the next step without further purification.

Preparation 103: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[1-(tert-butylcarbamoyl)-3,6-dihydro-2H-pyridin-5-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2568)




To a stirred solution of *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(1,2,3,6-tetrahydropyridin-5-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Hydrochloric Acid (2)) (27 mg, 0.037mmol) in anhydrous dichloromethane (1.0 mL), triethyl amine (50 µL, 0.3587 mmol) and 2-isocyanato-2-methyl-propane (8.0 mg, 0.081 mmol) were added at 0 °C under nitrogen. The yellow solution was stirred at ambient temerature for 2 hours. The volatiles were removed under reduced pressure and the residue was purified by reverse-phase HPLC to furnish *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[1-(tert-butylcarbamoyl)-3,6-dihydro-2H-pyridin-5-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2568) (Hydrochloric Acid

(1)) (9.2 mg, 0.015 mmol, 41%) ESI-MS m/z calc. 569.27844, found 570.4 $(M+1)^+$; Retention time: 1.3 minutes

The following compounds can be synthesized using the procedures described herein

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[1-(cyclohexylcarbamoyl)-3,6-dihydro-2H-pyridin-4-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2520)

Preparation 104: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[1-(2,2-dimethylbutanoyl)-3,6-dihydro-2H-pyridin-5-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2612)



To a stirred solution of *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(1,2,3,6-tetrahydropyridin-5-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Hydrochloric Acid (2)) (27 mg, 0.037 mmol) in anhydrous dichloromethane (1.0 mL), triethyl amine (50. µL, 0.36 mmol) and 2,2-dimethylbutanoyl chloride (11 mg, 0.082 mmol) were added at 0 °C under nitrogen. The yellow solution was stirred at ambient temerature for 2 hours. The volatiles were removed under reduced pressure and the residue was purified from reverse-phase HPLC to furnish *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[1-(2,2-dimethylbutanoyl)-3,6-dihydro-2H-pyridin-5-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2612) (Hydrochloric Acid (1)) (6.7 mg, 0.011 mmol, 27%) ESI-MS m/z calc. 568.2832, found 569.7 (M+1)+; Retention time: 1.44 minutes

The following compounds can be synthesized using the procedures described herein

N-[(6-amino-2-pyridyl)sulfonyl]-6-[1-(3-chlorobenzoyl)-3,6-dihydro-2H-pyridin-5-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1984),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[1-(1-piperidylsulfonyl)-3,6-dihydro-2H-pyridin-5-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1942),

N-[[6-[(2-phenylacetyl)amino]-2-pyridyl]sulfonyl]-6-[1-(2-phenylacetyl)-3,6-dihydro-2H-pyridin-5-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2243),

isopropyl *N*-[6-[[6-(1,2,3,6-tetrahydropyridin-5-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]carbamate (Compound 2624),

isopropyl 5-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-6-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]-2-pyridyl]-3,6-dihydro-2H-pyridine-1-carboxylate (Compound 2622),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[1-(3,3-dimethylbutanoyl)-3,6-dihydro-2H-pyridin-5-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2650),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[1-(2-cyclohexylacetyl)-3,6-dihydro-2H-pyridin-4-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2596),

isopropyl 4-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-2-tert-butyl-6-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]-3-pyridyl]-3,6-dihydro-2H-pyridine-1-carboxylate (Compound 2027),


N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[1-(2-phenylacetyl)-3,6-dihydro-2H-pyridin-4-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2417),

N-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-[1-(3,3-dimethylbutanoyl)-3,6-dihydro-2H-pyridin-4-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2344),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[1-(2-cyclohexylacetyl)-3,6-dihydro-2H-pyridin-5-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2633) and,

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-isobutylsulfonyl-3,6-dihydro-2H-pyridin-5-yl)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1882)

Preparation 105: *N*-[(6-amino-2-pyridyl)sulfonyl]-5-fluoro-6-[(3R)-tetrahydrofuran-3-yl]oxy-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2281)



To a 4 mL vial was added NaH (10. mg, 0.25 mmol), (3*R*)-tetrahydrofuran-3-ol (27.63 mg, 25.19 µL, 0.3136 mmol), and DMSO (800 µL). The mixture was stirred at ambient temperature for 2 minoutes. *N*-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-5-fluoro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Hydrochloric Acid (1)) (30. mg, 0.063 mmol) was then added to the vial and the reaction was stirred at 150 °C for 2 hours. The reaction mixture was filtered and purified by preparative HPLC (1-99% acetonitrile over 30 minutes with 5 mM HCl modifier) to give *N*-[(6-amino-2-pyridyl)sulfonyl]-5-fluoro-6-[(3R)-tetrahydrofuran-3-yl]oxy-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2281) (Hydrochloric Acid (1)) (9 mg, 0.02 mmol, 30%) ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.29 (s, 1H), 7.66 - 7.57 (m, 2H), 7.17 (d, J = 7.3 Hz, 1H), 6.66 (d, J = 8.3 Hz, 1H), 6.41 (s, 2H), 5.55 - 5.48 (m, 1H), 3.94 - 3.83 (m, 2H), 3.83 - 3.76 (m, 2H), 2.56 (d, J = 10.6 Hz, 1H), 2.44 (t, J = 8.7 Hz, 1H), 2.32 - 2.16 (m, 1H), 2.10 - 2.02 (m, 1H), 1.84 (dd, J = 11.8, 5.6 Hz, 1H), 1.53 (s, 3H), 1.44 (s, 3H), 1.37 (t, J = 12.0 Hz, 1H), 0.85 (d, J = 6.3 Hz, 3H). ESI-MS *m/z* calc. 493.1795, found 494.3 $(M+1)^+$; Retention time: 1.08 minutes

The following compounds can be synthesized using the procedures described herein

N-[(6-amino-2-pyridyl)sulfonyl]-6-[(1R)-1,2-dimethylpropoxy]-5-fluoro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2020),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-cyclohexylethoxy)-5-fluoro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1730),

N-[(6-amino-2-pyridyl)sulfonyl]-5-fluoro-6-[(3S)-tetrahydrofuran-3-yl]oxy-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2056),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-cyclopentylethoxy)-5-fluoro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1966),


N-[(6-amino-2-pyridyl)sulfonyl]-5-fluoro-6-(1-tetrahydropyran-4-ylethoxy)-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1891),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(cyclopentoxy)-5-fluoro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2100),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(2,2-dimethylmorpholin-4-yl)-5-fluoro-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2586), and

N-[(6-amino-2-pyridyl)sulfonyl]-5-fluoro-6-[1-(4-pyridyl)ethoxy]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2212)

Preparation 106: 6-tert-butyl-*N*-[[6-[(4-methoxyphenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2724) (Isomer 1) (Compound 2428) (Isomer 2)



6-(*tert*-butyl)-*N*-((6-chloropyridin-2-yl)sulfonyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)nicotinamide (100. mg, 0.215 mmol) and (4-methoxyphenyl)methanol (59.6 mg, 0.431 mmol) were dissolved in anhydrous anhydrous dimethyl sulfoxide (1 mL), and then cesium carbonate (210 mg, 0.645 mmol) was added to the reaction mixture The resulting mixture was heated at 130 °C for 24 hours and cooled to room temperature. Water (10 mL) and aqueous hydrogen chloride solution (1N) was added until pH value of solution was around 2. Reaction solution was extracted with ethyl acetate, and the extract was washed with brine, dried over sodium sulfate and filtered. Solvents were removed under the reduced pressure. The residue obtained was purified by reverse-phase HPLC utilizing a gradient of 0.1% trifluoroacetic acid in water and 0.1% trifluoroacetic acid in acetonitrile. Factions were lyophilized to afford 6-tert-butyl-*N*-[[6-[(4-methoxyphenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2724) (Isomer 1) (Compound 2428) (Isomer 2) (35.2 mg, 0.0621 mmol, 28.9%). MS (ESI, m/z): $[M+H]^+$ 567.3.

The following compounds can be synthesized using the procedures described herein

N-[[6-[2-(4-fluorophenyl)ethoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1704),

N-[[6-(3-phenylpyrrolidin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1708),

N-[[6-[methyl(propyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1711),

N-[[6-[(4-methoxyphenyl)methyl-methyl-amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1717),

N-[(6-tert-butoxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1718),

N-[[6-(3-methoxy-5-methyl-phenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1719),

N-[[6-(2-cyanophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1720),

N-[[6-(4-isopropylanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1722),

N-[[6-(4-cyclopentylpiperazin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1733),

N-[[6-(pyrrolidin-3-ylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1734),

N-[[6-(4-acetyl-2-methoxy-phenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1744),

N-[[6-(3,5-dimethyl-1-piperidyl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1749),

N-[[6-(1-naphthyloxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1751),

N-[(6-prop-2-ynoxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1753),

N-[[6-(3-phenyl-1-piperidyl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1755),

N-[[6-(4-tert-butylanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1756),

N-[[6-[4-(dimethylamino)anilino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1757),

N-[[6-(3,4-dimethoxyphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1761),

N-[[6-(tetrahydrofuran-2-ylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1764),

N-[[6-(cyclohexylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1767),

N-[[6-[(2,5-difluorophenyl)methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1770),

tert-butyl 4-[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]piperazine-1-carboxylate (Compound 1773),

N-[[6-(1,3-benzothiazol-6-ylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1775),

N-[[6-(3,4,4a,5,6,7,8,8a-octahydro-2H-quinolin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1776),

N-[[6-[(5-phenylisoxazol-3-yl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1777),

N-[[6-[(3-chloro-4-methoxy-phenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1778),

N-[[6-(3-ethynylphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1782),

N-[[6-(2,4,5-trifluoroanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1796),

N-[[6-[(2-fluorophenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1798),

N-[[6-(2-cyclohexylethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1800),

N-[[6-[3-(methoxymethyl)-1-piperidyl]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1805),

N-[[6-(2,4-dichlorophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1806),

N-[[6-(4-fluorophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1813),

N-[[6-(1,3-benzodioxol-5-ylmethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1814),

N-[[6-[4-(2-pyridyl)piperazin-1-yl]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1816),

N-[[6-(1H-indol-5-ylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1823),

N-[[6-(8-quinolylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1826),

N-[[6-[benzyl(methyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1827),

N-[[6-(3-ethoxyphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1828),

N-[[6-(3-chloro-4-fluoro-phenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1831),

N-[[6-(1,3,4,9-tetrahydropyrido[3,4-b]indol-2-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1833),

N-[[6-(cyclopentoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1834),

N-[(6-isoindolin-2-yl-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1838),

N-[[6-(3-chloroanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1841),

N-[[6-[(1R)-1-phenylethoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1842),

N-[[6-[(4-ethyl-2-pyridyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1862),

N-[[6-(thiazol-2-ylmethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1865),

N-[[6-[ethyl(2-methoxyethyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1870),

N-[[6-(2-chloroanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1872),

N-[[6-(3,4-dihydro-2H-pyran-2-ylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1874),

N-[[6-(7-quinolyloxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1875),

N-[[6-[(1-acetylindolin-5-yl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1877),

N-[[6-[3-(*N*-methylanilino)propylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1879),

N-[[6-[(E)-but-2-enoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1881),

N-[[6-(3-methylcyclopentoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1884),

N-[[6-(p-tolylmethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1892),

N-[[6-(2-methylanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1898),

N-[[6-[(2-chloro-4-fluoro-phenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1899),

N-[[6-(4-isopropylphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1902),

N-[[6-(1-phenylethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1905),

N-[[6-(2,3-dichloroanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1906),

N-[[6-(cyclohexoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1908),

N-[[6-(indan-5-ylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1909),

N-[[6-(1-phenylethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1917),

N-[[6-(1-cyclohexylethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1919),

N-[[6-(4-ethoxyphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1924),

N-[[6-(3-tert-butylphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1926),

N-[[6-[(2,4-dichlorophenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1929),

N-[[6-(azepan-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1932),

N-[[6-(3-isoquinolylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1933),

N-[[6-(1-methylcyclopentoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1936),

N-[[6-[4-(1,3-dioxolan-2-ylmethyl)piperazin-1-yl]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1937),

N-[[6-(benzylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1938),

N-[[6-(tetrahydropyran-2-ylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1939),

N-[[6-[(3-methoxyphenyl)methyl-methyl-amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1940),

N-[[6-(2-chloro-5-fluoro-anilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1941),

N-[[6-(3-acetylanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1949),

N-[[6-[[3-(trifluoromethyl)phenyl]methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1953),

N-[(6-tetralin-5-yloxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1955),

N-[[6-[(5-chloro-3-pyridyl)oxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1957),

N-[[6-(2-methoxyanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1959),

N-[[6-[(2-methoxyphenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1965),

N-[[6-[(2,4-dichlorophenyl)methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1968),

N-[[6-[(4-tert-butylcyclohexyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1971),

N-[[6-[(3-methoxyphenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1972),

N-[[6-[4-(trifluoromethoxy)anilino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1977),

N-[[6-[methyl(phenethyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1978),

N-[[6-[(2-chlorophenyl)methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1983),

N-[[6-(4-pentylpiperazin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1985),

N-[[6-(1,3-dihydrobenzo[de]isoquinolin-2-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1986),

N-[[6-(4-phenyl-1-piperidyl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1987),

N-[[6-(3-methylcyclohexoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1988),

N-[[6-(2-quinolyloxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1989),

N-[[6-(3-thienylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1992),

N-[[6-[[(1R)-1-cyclohexylethyl]amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1996),

N-[[6-[(6-methylcyclohex-3-en-1-yl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2001),

N-[[6-(1H-indol-5-yloxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2003),

N-[[6-(2-thienylmethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2004),

N-[[6-[methyl-[2-(3-pyridyl)ethyl]amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2005),

N-[[6-(3,4,4a,5,6,7,8,8a-octahydro-1H-isoquinolin-2-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2011),

N-[[6-[[3-(dimethylamino)phenyl]methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2012),

N-[[6-(1,3-benzodioxol-5-yloxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2013),

N-[[6-(2-fluorophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2015),

N-[[6-(2-cyclohexylpropoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2019),

N-[[6-(4-phenylanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2023),

N-[[6-[(2S)-2-(methoxymethyl)pyrrolidin-1-yl]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2024),

N-[[6-(3,3-dimethylbutoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2025),

N-[[6-(cyclopentylmethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2026),

N-[[6-(3-methyl-1-piperidyl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2030),

N-[[6-(tetrahydropyran-4-ylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2032),

N-[[6-(3,3-dimethylcyclohexoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2038),

N-[[6-(3,4-dimethoxyanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2041),

N-[[6-[cyclopropylmethyl(methyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2044),

N-[[6-[1-(1-naphthyl)ethylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2049),

N-[[6-(cyclopentylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2053),

N-[[6-[2-(trifluoromethyl)pyrrolidin-1-yl]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2055),

N-[[6-(1-methylcycloheptoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2057),

N-[[6-(1,7,7-trimethylnorbornan-2-yl)oxy-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2059),

ethyl 3-[[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]amino]benzoate (Compound 2060),

N-[[6-(3,4-dichlorophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2061),

N-[[6-[(3-chlorophenyl)methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2064),

N-[[6-(2-pyridylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2068),

N-[[6-[2-(4-methoxyphenyl)ethylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2072),

N-[[6-(5-quinolylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2074),

N-[(6-isopentyloxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2075),

N-[[6-(4-cyanophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2076),

N-[[6-[(3S)-1-benzylpyrrolidin-3-yl]oxy-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2078),

N-[[6-[(4-methyl-2-pyridyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2082),

N-[[6-(2-tert-butylphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2083),

N-[[6-[4-(1-piperidyl)anilino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2087),

N-[[6-[(2-methoxyphenyl)methyl-methyl-amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2090),

N-[[6-(2-phenylpyrrolidin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2094),

N-[[6-(3,4-dihydro-1H-isoquinolin-2-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2098),

N-[[6-[(2,4-difluorophenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2099),

N-[[6-(4-fluoroanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2102),

N-[[6-(2,4,6-trifluoroanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2106),

N-[[6-(4-benzyl-1-piperidyl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2107),

N-[[6-(3-isopropylphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2108),

N-[[6-[(1S,2S)-2-methylcyclopentoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2109),

N-[[6-[(1-phenylpyrazol-4-yl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2111),

N-[[6-(2,3-dihydro-1,4-benzodioxin-3-ylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2115),

N-[[6-(3-methylpyrrolidin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2116),

N-[[6-[(4-pyrazol-1-ylphenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2118),

N-[[6-(4-phenylbutylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2119),

ethyl 4-[[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]amino]benzoate (Compound 2122),

N-{[6-(3-benzylpyrrolidin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2123),

N-[[6-(1,1-dimethylbutoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2130),

N-[[6-(tetrahydrofuran-2-ylmethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2131),

N-[[6-(3-methoxypropylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2135),

N-[[6-(1-naphthylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2136),

N-[[6-(2-methyl-2-pyrrol-1-yl-propoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2137),

methyl 6-[[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]amino]pyridine-3-carboxylate (Compound 2140),

N-[[6-(3-tert-butylanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2142),

N-[[6-[3-(trifluoromethoxy)anilino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2143),

N-[[6-(4-chlorophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2145),

N-[[6-[2-(2-thienyl)ethylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2146),

N-[(6-cyclohex-2-en-1-yloxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2147),

N-[[6-(2-naphthylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2151),

N-[[6-(3-methoxy-1-piperidyl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2152),

N-[[6-[(4,6-dimethyl-2-pyridyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2160),

N-[[6-[2-(3-thienyl)ethoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2161),

N-[[6-(3-chloro-2-fluoro-phenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2165),

N-[[6-(3,3-difluoropyrrolidin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2166),

N-[(6-butoxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2167),

N-[[6-[(3-methoxyphenyl)methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2168),

N-[[6-(1,3-benzodioxol-5-ylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2169),

N-[[6-(4-isopropylcyclohexoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2173),

N-[[6-[4-(4-pyridylmethyl)piperazin-1-yl]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2175),

N-[[6-(3-chloro-5-cyano-phenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2177),

N-[[6-(2,5-dimethoxyanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2179),

N-[[6-(2-thienylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2181),

N-[[6-[(4-chloro-2-fluoro-phenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2182),

N-[[6-[(1S)-1-phenylethoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2186),

N-[[6-[(4-isopropylphenyl)methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2187),

N-[[6-(cyclobutylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2188),

N-[[6-(2,3-dihydro-1,4-benzodioxin-6-ylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2190),

N-[[6-[(1-methylindol-5-yl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2193),

N-[[6-(1-methylbutylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2197),

N-[[6-[(2-methyl-1,3-benzothiazol-5-yl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2198),

N-[[6-(isopropylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2200),

N-[[6-(cyclooctylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2202),

N-[[6-(2-phenoxyethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2206),

N-[[6-(4-phenylpiperazin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2208),

N-[(6-decalin-2-yloxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2211),

N-[[6-(4-ethoxyanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2214),

N-[[6-(2,4-dichloroanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2217),

N-[[6-[(2-methylcyclohexyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2218),

N-[[6-[1-(methoxymethyl)propylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2221),

N-[[6-[4-(1-methylbutyl)piperazin-1-yl]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2226),

N-[[6-(2,3-dimethoxyphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2233),

N-[[6-(4-methoxyphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2239),

N-[[6-(1,3-benzodioxol-4-ylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2249),

N-[[6-(cyclohexylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2250),

tert-butyl 2-[methyl-[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]amino]acetate (Compound 2251),

N-[[6-(cycloheptoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2252),

ethyl 4-[[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]amino]piperidine-1-carboxylate (Compound 2253),

N-[[6-(4-pyrazin-2-ylpiperazin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2254),

N-[(6-indolin-1-yl-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2255),

N-[[6-(6-methoxy-3,4-dihydro-1H-isoquinolin-2-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2260),

N-[[6-[[2-(trifluoromethyl)phenyl]methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2261),

N-[[6-[2-(4-chlorophenyl)ethylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2262),

N-[[6-[3-(5-methyl-1,2,4-oxadiazol-3-yl)phenoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2263),

N-[[6-[(3-chlorophenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2264),

N-[[6-(1-naphthylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2270),

N-[[6-[(2-chlorophenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2273),

N-[[6-(5-isoquinolylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2274),

N-[[6-(3,5-difluoroanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2282),

N-[[6-[4-(6-methyl-2-pyridyl)piperazin-1-yl]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2285),

N-[[6-[(4-tert-butylphenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2286),

N-[[6-[4-(trifluoromethyl)anilino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2292),

N-[[6-(3,5-dimethoxyphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2293),

N-[[6-[(6-methyl-2-pyridyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2294),

N-[[6-(3-benzyl-1-piperidyl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2297),

N-[[6-(2-chlorophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2298),

N-[[6-[(3-cyanophenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2299),

N-[[6-(2-ethylbutylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2305),

N-[[6-[butyl(methyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2307),

N-[[6-(2-furylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2308),

N-[[6-(2,6-dimethoxyphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2309),

N-[[6-[(2-methoxyphenyl)methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2310),

N-[[6-(4-aminophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2314),

N-[[6-(3-cyano-4-fluoro-anilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2316),

N-[(6-cyclopent-3-en-1-yloxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2317),

N-[[6-[[(1R)-1,2-dimethylpropyl]amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2318),

N-[[6-(3-methylphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2320),

(3S)-2-[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]-3,4-dihydro-1H-isoquinoline-3-carboxylic acid (Compound 2321),

N-[[6-(2-methylmorpholin-4-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2324),

N-[[6-(4-acetylanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2328),

N-[[6-(2-tetrahydropyran-4-ylethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2329),

N-[[6-(4-methyl-1-piperidyl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2330),

N-[[6-(3-fluorophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2332),

N-[[6-(8-quinolyloxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2333),

tert-butyl *N*-[4-[[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]amino]butyl]carbamate (Compound 2335),

N-[[6-(4-methylcyclohexoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2336),

N-[[6-(1H-indol-4-yloxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2338),

N-[[6-(3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2340),

N-[[6-(3,4-difluoroanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2341),

N-[[6-(cyclopropylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2342),

N-[[6-[2-methoxyethyl(propyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2343),

N-[[6-[isopentyl(methyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2346),

N-[[6-(2,2-difluoroethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2350),

N-[[6-(3-isopropoxyanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2353),

N-[[6-(2-ethoxyethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2354),

N-[[6-(3,5-dimethoxyanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2357),

N-[[6-[2-(2-pyridyl)ethylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2359),

N-[[6-[[6-(trifluoromethyl)-2-pyridyl]methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2365),

N-[[6-(cyclobutoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2371),

N-[[6-[(3-chlorophenyl)methyl-methyl-amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2378),

N-[[6-[3-(trifluoromethyl)anilino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2381),

N-[[6-(3,4-dihydro-2H-quinolin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2382),

N-[[6-[4-(2-dimethylaminoethyl)-1-piperidyl]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2383),

N-[[6-(5-chloroindolin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2385),

N-[[6-(3-acetamidoanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2387),

N-[[6-(1H-indazol-5-ylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2389),

N-[[6-[(2,6-dichlorophenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2390),

N-[[6-(m-tolylmethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2391),

N-[[6-[(3S)-3-fluoropyrrolidin-1-yl]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2393),

N-[[6-(tetrahydrofuran-3-ylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2396),

N-[[6-[2-(trifluoromethyl)anilino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2397),

tert-butyl *N*-methyl-*N*-[2-[[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]amino]ethyl]carbamate (Compound 2400),

N-[[6-[(3,4-dichlorophenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2402),

N-[[6-[(1-benzylpyrrolidin-3-yl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2407),

N-[[6-[1-(2-fluorophenyl)ethoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2408),

N-[[6-[(3-fluorophenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2409),

N-[[6-(2,2-dimethylpropoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2412),

N-[[6-[(4-isopropylphenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2415),

N-[[6-[(6-methoxy-2-pyridyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2420),

N-[[6-[(3,5-difluorophenyl)methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2421),

N-[[6-(4-hexylpiperazin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2423),

N-[[6-[(4-methoxyphenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2428),

N-[(6-methoxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2430),

N-[[6-(3-acetamidophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2433),

N-[[6-[(1-methylcyclopropyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2434),

tert-butyl *N*-[4-[[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]amino]cyclohexyl]carbamate (Compound 2439),

N-[[6-(4-imidazol-1-ylphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2441),

methyl 1-[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]piperidine-4-carboxylate (Compound 2442),

N-[[6-[3-(dimethylamino)phenoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2443),

N-[[6-(3-quinolylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2449),

N-[[6-[2-(4-piperidyl)ethoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2452),

N-[[6-(4-fluoro-1-piperidyl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2453),

N-[[6-(4-methylphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2460),

N-[[6-(3-chlorophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2461),

N-[[6-[(3,5-difluorophenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2462),

N-[(6-morpholino-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2464),

N-[[6-[2-(3-methoxyphenyl)ethylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2465),

N-[[6-[[6-(trifluoromethyl)-3-pyridyl]amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2466),

N-[[6-[(1R)-1-methyl-2-phenyl-ethoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2470),

N-[[6-[[6-(trifluoromethyl)-3-pyridyl]methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2476),

N-[[6-(3-cyanoanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2484),

N-[[6-(2-isopropylphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2486),

ethyl 1-[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]piperidine-4-carboxylate (Compound 2488),

methyl 3-[[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]amino]benzoate (Compound 2493),

tert-butyl 3-[[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]amino]propanoate (Compound 2495),

N-[[6-[2-(p-tolyl)ethylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2502),

N-[[6-(5-chloro-2-methoxy-phenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2503),

N-[[6-[4-(2-pyridylmethyl)piperazin-1-yl]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2504),

methyl 4-[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]piperazine-1-carboxylate (Compound 2506),

N-[[6-(3-furylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2507),

N-[(6-phenoxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2509),

N-[[6-(2,6-difluoroanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2513),

N-[[6-[3-(dimethylamino)anilino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2517),

N-[[6-(7-fluoro-3,4-dihydro-1H-isoquinolin-2-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2518),

N-[[6-(1-oxotetralin-6-yl)oxy-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2523),

N-[[6-[(4-fluorophenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2524),

N-[[6-(3-ethoxyanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2526),

N-[(6-propoxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2528),

N-[[6-(3-butoxypropylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2529),

N-[[6-[4-(diethylamino)anilino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2531),

N-[[6-[(3,4-difluorophenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2533),

N-[[6-(methylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2535),

tert-butyl (2R)-1-[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]pyrrolidine-2-carboxylate (Compound 2536),

tert-butyl *N*-[(3R)-1-[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]pyrrolidin-3-yl]carbamate (Compound 2537),

N-[[6-(4-oxazol-5-ylanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2546),

N-[[6-[3-(3-methylpyrazol-1-yl)propylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2550),

N-[[6-(3-methoxyphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2551),

N-[(6-pyrrolidin-1-yl-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2561),

N-[[6-(4-pyrrol-1-ylphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2563),

N-[[6-[4-(1-ethylpropyl)piperazin-1-yl]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2565),

N-[[6-(3-acetylphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2569),

N-[[6-(3-chloro-4-methoxy-anilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2571),

N-[[6-(3-methylanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2572),

N-[[6-(cyclopropylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2574),

N-[[6-[(4-methoxyphenyl)methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2579),

N-[[6-(5-fluoroisoindolin-2-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2580),

N-[[6-[(4-chlorophenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2585),

N-[[6-(2-chloro-5-fluoro-phenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2591),

N-[[6-(chroman-3-ylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2595),

N-[[6-(2-furylmethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2597),

N-[[6-(5-chloro-2-fluoro-phenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2598),

N-[[6-[(3-pyrrol-1-ylphenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2600),

N-[[6-(4-cyclohexylpiperazin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2601),

N-[[6-(2-propoxyethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2602),

N-[[6-(2-methylpyrrolidin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2609),

N-[[6-[(4-ethynylphenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2610),

N-[[6-(2-naphthyloxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2611),

N-[(6-ethoxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2617),

N-[[6-[[4-(trifluoromethyl)phenyl]methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2618),

N-[(6-benzyloxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2619),

N-[[6-(4-butyl-1,4-diazepan-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2620),

N-[[6-(4-pyrimidin-2-ylpiperazin-1-yl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2621),

N-[[6-[2-(2-fluorophenyl)ethoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2625),

N-[[6-(4H-1,3-benzodioxin-2-ylmethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2627),

N-[[6-(2-cyclopentylethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2630),

N-[[6-[2-(2-chlorophenyl)ethylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2632),

N-[[6-(2-methoxyphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2637),

N-[(6-anilino-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2639),

N-[[6-[(2,3-difluorophenyl)methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2640),

N-[[6-[4-(trifluoromethyl)-1-piperidyl]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2641),

N-[[6-[2-(4-ethoxyphenyl)ethylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2642),

N-[[6-[(2-methyl-8-quinolyl)oxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2643),

N-[[6-[(5-chloro-2-methoxy-phenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2645),

N-[[6-(3-methylbut-2-enoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2648),

N-[[6-(2,4,6-trifluorophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2654),

N-[[6-[(5-chloro-2-pyridyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2656),

N-[[6-(2-cyanoanilino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2659),

N-[[6-[(5-chloro-2-pyridyl)oxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2747),

N-[[6-[2-(4-pyridyl)ethylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2746),

N-[[6-(2-aminoethoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2754),

N-[(6-but-2-ynoxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2753),

N-[[6-[3-methoxypropyl(methyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2745),

N-[[6-(4-methoxy-1-piperidyl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2743),

N-[[6-(1-oxoindan-5-yl)oxy-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2744),

N-[[6-[(6-methyl-2-pyridyl)oxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2742),

N-[[6-(4-acetylphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2741),

N-[[6-[(2-methyl-1,3-benzothiazol-5-yl)oxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2740),

N-[[6-[[(1R)-1,2,2-trimethylpropyl]amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2757),

methyl 4-[[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]oxy]benzoate (Compound 2739),

N-[[6-(2,1,3-benzoxadiazol-5-yloxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2738),

N-[[6-(4-tert-butoxyphenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2751),

N-[[6-[isobutyl(methyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2758),

N-[[6-(diethylamino)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2737),

N-[[6-(3-cyanophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2736),

N-[[6-[isopropyl(methyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2735),

tert-butyl *N*-[4-[[6-[[2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carbonyl]sulfamoyl]-2-pyridyl]amino]phenyl]carbamate (Compound 2755),

N-[[6-[2-(3-fluorophenyl)ethoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2752),

N-[[6-[2-(1-piperidyl)phenoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2750),

N-[[6-(2-methyl-1-piperidyl)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2759),

N-[[6-[cyclopropylmethyl(propyl)amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2734),

N-[[6-[(2,5-dimethoxyphenyl)methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2733),

N-[(6-phenethyloxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2732),

N-[[6-(2,3-dichlorophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2731),

N-[[6-(3-chloro-5-fluoro-phenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2730),

N-[[6-[(2-chlorophenyl)methyl-methyl-amino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2756),

N-[(6-chroman-4-yloxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2749),

N-[[6-[3-(trifluoromethyl)phenoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2729),

N-[[6-[(2-chloro-6-fluoro-phenyl)methoxy]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2728),

6-tert-butyl-*N*-[(6-isopentyloxy-2-pyridyl)sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2727),

N-[[6-(1,3-benzodioxol-5-ylmethoxy)-2-pyridyl]sulfonyl]-6-tert-butyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2725),

N-[[6-[(E)-but-2-enoxy]-2-pyridyl]sulfonyl]-6-tert-butyl-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2726),

6-tert-butyl-*N*-[[6-[(4-methoxyphenyl)methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2723),


N-[[6-[(3,4-dimethoxyphenyl)methylamino]-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2662), and

N-[[6-(3,5-dichlorophenoxy)-2-pyridyl]sulfonyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2663)

PREPARATION 107: *N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6-phenyl-2-[(4*S*)-2,2,4-


trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2445)

Step 1: 2-Methoxypyridine-3-sulfonamide



2-Chloropyridine-3-sulfonyl chloride (5.00 g, 23.6 mmol) was dissolved in ammonia in methanol (16.8 mL of 7 M, 117.9 mmol) and the reaction mixture was stirred for 30 minutes. The reaction mixture was concentrated under reduced pressure and dissolved in sodium methoxide (24.9 mL of 25 %w/v, 115.1 mmol) and heated to 60 °C over 2 days. The reaction mixture was evaporated and the residue was partitioned between ethyl acetate and 1N HCl. The organics were separated, washed with brine, dried over sodium sulfate and concentrated under reduced pressure. The crude material was purified by silica gel column chromatography eluting with 0-100% ethyl acetate in hexanes to give 2-methoxypyridine-3-sulfonamide (3.12 g, 70%) ESI-MS m/z calc. 188.02556, found 189.0 $(M+1)^+$; Retention time: 0.23 minutes


Step 2: 2,6-dichloro-*N*-[(2-methoxy-3-pyridyl)sulfonyl]pyridine-3-carboxamide



A mixture of 2,6-dichloropyridine-3-carboxylic acid (2.00 g, 10.4 mmol), thionyl chloride (7.45 g, 4.56 mL, 62.5 mmol), and DMF (0.1 mL) was stirred at 45 °C for 5 hours. The reaction was concentrated under reduced pressure, dissolved in dichloromethane (5.30 mL) and slowly added to a solution of 2-methoxypyridine-3-sulfonamide (1.96 g, 10.4 mmol) and triethylamine (3.16 g, 4.36 mL, 31.3 mmol) in dichloromethane (5.30 mL) cooled in an ice bath. The reaction mixture was stirred overnight, then diluted with dichloromethane and washed with 1 M HCl (20 mL x 1), water (20 mL), dried over sodium sulfate, and evaporated under reduced pressure. The residue was purified by silica gel column chromatography (0-5 % methanol in dichloromethane). The resulting solid was dissolved in dichloromethane and washed with sodium bicarbonate (~ pH 5), the organic layers were dried over sodium sulfate and


evaporated under reduced pressure to give 2,6-dichloro-*N*-[(2-methoxy-3-pyridyl)sulfonyl]pyridine-3-carboxamide (3.50 g, 93%) ESI-MS m/z calc. 360.9691, found 362.0 $(M+1)^+$; Retention time: 0.46 minutes.

Step 3: 2,6-dichloro-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide



A mixture of 2,6-dichloro-*N*-[(2-methoxy-3-pyridyl)sulfonyl]pyridine-3-carboxamide (0.76 g, 2.10 mmol) and HCl (5.25 mL of 4 M in dioxane, 21.0 mmol) in dioxane (5 mL) was stirred at 90 °C for 90 minutes. The solvent was evaporated under reduced pressure to give 2,6-dichloro-*N*-((2-oxo-1,2-dihydropyridin-3-yl)sulfonyl)nicotinamide (0.72 g, 99%) as a tan solid.


Step 4: 6-chloro-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide



To 2,6-dichloro-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (HCl salt) (4.04 g, 10.5 mmol), (4*S*)-2,2,4-trimethylpyrrolidine (HCl salt) (2.36 g, 15.8 mmol), and K_2CO_3 (7.26 g, 52.5 mmol) was added DMSO (8.86 mL) and the reaction mixture was heated at 120 °C for 24 h. The reaction mixture was diluted with ethyl acetate and the residual potassium carbonate was filtered. The organics were washed with 0.1 M aqueous HCl (150 mL). The pH of aqueous layer was brought to 5 by the addition of aqueous HCl, and then washed with ethyl acetate. The combined organic layers were washed with brine, dried over sodium sulfate and concentrated under reduced pressure to give 6-chloro-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-

2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (4.15 g, 94%).

Step 5: *N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6-phenyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2445)



6-chloro-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (30.0 mg, 0.07 mmol), phenylboronic acid (10.3 mg, 0.08 mmol), [1,3-bis(2,6-di*iso*propylphenyl)-2H-imidazol-2-yl]-dichloro-(3-chloro-1-pyridyl)palladium (4.80 mg, 0.01 mmol), and 2M K_2CO_3 (177 uL, 0.35 mmol) were combined in DMF (1.0 mL) and heated at 120 °C for 16 h. The reaction mixture was filtered and purified by LC/MS utilizing a gradient of 10-99% acetonitrile in 5 mM aq HCl to yield *N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6-phenyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2445). ESI-MS m/z 467.4 $(M+1)^+$; Retention time: 1.13 minutes; 1H NMR (400 MHz, DMSO-d6) δ 12.59 (s, 1H), 12.53 (d, J = 6.3 Hz, 1H), 8.24 (dd, J = 7.2, 2.2 Hz, 1H), 8.08 - 8.02 (m, 2H), 7.83 (dt, J = 6.6, 3.3 Hz, 1H), 7.76 (d, J = 8.0 Hz, 1H), 7.54 - 7.46 (m, 2H), 7.46 - 7.40 (m, 1H), 7.19 (d, J = 8.0 Hz, 1H), 6.45 (t, J = 6.8 Hz, 1H), 2.82 - 2.63 (m, 2H), 2.30 - 2.18 (m , 1H), 1.89 (dd, J = 11.8, 5.5 Hz, 1H), 1.62 (s, 6H), 1.45 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).

The following compounds can be synthesized using the procedures described herein:

6-(*m*-tolyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2424), 6-(2-hydroxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2480), *N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6-[4-(trifluoromethyl)phenyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1799), *N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6-(*p*-tolyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2438),

6-(3-methoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1927), *N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-6-[2-(trifluoromethoxy)phenyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2514),

6-[3-(hydroxymethyl)phenyl]-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1997),

6-(4-methoxyphenyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2080),

N-(benzenesulfonyl)-6-(6-methoxy-2-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2411),

6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-(3-hydroxyphenyl)sulfonyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1885),

N-(benzenesulfonyl)-6-(2-hydroxy-3-methyl-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2133),

6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-(4-hydroxyphenyl)sulfonyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1750),

6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-(2-hydroxyphenyl)sulfonyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1836),

6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-(4-methoxyphenyl)sulfonyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1876),

N-(benzenesulfonyl)-6-phenyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2092),

N-(benzenesulfonyl)-2-[(3S)-2,2-dideuterio-3,5,5-trimethyl-pyrrolidin-1-yl]-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2560),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-ethoxy-3-methyl-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1860),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[1-methyl-3-(trifluoromethyl)pyrazol-4-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1740),


N-[(2-amino-3-pyridyl)sulfonyl]-6-(1*H*-indol-5-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2125),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[2-(dimethylamino)pyrimidin-5-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2021),
N-[(2-amino-3-pyridyl)sulfonyl]-6-(1*H*-indol-4-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2478),
N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-hydroxy-3,5-dimethyl-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1923),
N-[(2-amino-3-pyridyl)sulfonyl]-6-[2-methyl-5-(trifluoromethyl)pyrazol-3-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2416),
N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-ethoxy-6-fluoro-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2103),
N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-ethoxy-5-fluoro-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2071),
N-[(2-amino-3-pyridyl)sulfonyl]-6-(6-methyl-3-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2363),
N-[(2-amino-3-pyridyl)sulfonyl]-6-(1-methylindol-5-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2352),
N-[(2-amino-3-pyridyl)sulfonyl]-6-(1,3,5-trimethylpyrazol-4-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2668),
N-[(2-amino-3-pyridyl)sulfonyl]-6-(2,5-difluoro-4-methoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2388),
N-[(2-amino-3-pyridyl)sulfonyl]-6-(6-hydroxy-2-naphthyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2468),
N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-hydroxy-3-methoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2614),
N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-methoxy-5-methyl-3-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2086),
6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-(3-pyridylsulfonyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2377),
6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-(2-pyridylsulfonyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1832),

6-(3,5-dimethylphenyl)-*N*-[(6-isopentyloxy-2-pyridyl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2707) and 2-[(2*S*,5*R*)-2,5-dimethylpyrrolidin-1-yl]-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(2-oxo-1H-pyridin-3-yl)sulfonyl]pyridine-3-carboxamide (Compound 307).


PREPARATION 108: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3-*iso*butoxypyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2191)

Step 1: 1-(3-hydroxypyrazol-1-yl)ethanone



1*H*-pyrazol-5-ol (1.02 g, 12.1 mmol) and pyridine (5.0 mL, 61.8 mmol) were stirred at 95°C and a solution of acetic anhydride (1.24 g, 1.14 mL, 12.1 mmol) in pyridine (2.0 mL, 24.7 mmol) was added dropwise over a period of 3 minutes. The reaction mixture was then stirred at 95°C for an additional 1.5 h. The solvents were removed under reduced pressure and the solid residue was triturated with diethylether (30 mL), filtered, washed with diethylether and dried under high vacuum to give 1-(3-hydroxypyrazol-1-yl)ethanone (1.34 g, 88%) as a yellow solid. ESI-MS m/z calc. 126.04293, found 127.1 $(M+1)^+$; Retention time: 0.3 minutes; 1H NMR (400 MHz, DMSO-d6) δ 10.92 (s, 1H), 8.12 (d, J = 2.9 Hz, 1H), 6.00 (d, J = 3.0 Hz, 1H), 2.48 (s, 3H).


Step 2: 1-(3-*iso*butoxypyrazol-1-yl)ethanone



To 1-(3-hydroxypyrazol-1-yl)ethanone (1.25 g, 9.91 mmol), potassium carbonate (2.06 g, 14.9 mmol) and anhydrous DMF (10 mL) under nitrogen atmosphere was added 1-bromo-2-methyl-propane (1.49 g, 1.19 mL, 10.9 mmol). The reaction was stirred at 80°C for 2 hours. The reaction mixture was diluted with EtOAc (50 ml) and water (25 mL) and two phases were separated. The aqueous phase (pH = 9-10) was further extracted with EtOAc


(20 mL). The combined extracts were dried over sodium sulfate and the solvents evaporated. The resulting oil was purified by silica gel column chromatography using a gradient of EtOAc (0 to 30 %) in hexanes to give 1-(3-*iso*butoxypyrazol-1-yl)ethanone (1.24 g, 69%) as a colorless oil. ESI-MS m/z calc. 182.10553, found 183.3 $(M+1)^+$; Retention time: 1.35 minutes; 1H NMR (400 MHz, Chloroform-d) δ 8.05 (d, J = 3.0 Hz, 1H), 5.96 (d, J = 3.0 Hz, 1H), 3.99 (d, J = 6.6 Hz, 2H), 2.58 (s, 3H), 2.08 (dq, J = 12.9, 6.4 Hz, 1H), 1.01 (d, J = 6.7 Hz, 6H).

Step 3: 3-*iso*butoxy-1*H*-pyrazole



1-(3-*iso*butoxypyrazol-1-yl)ethanone (1.24 g, 6.82 mmol) was treated with MeOH (20 mL) and NaOH (1.14 mL of 6 M, 6.82 mmol) (6N aqueous) at room temperature for 2 hours. The volatiles were removed under reduced pressure. The residue was taken in EtOAc (25 mL) and brine (20 ml) and two phases were separated. The aqueous phase was further extracted with EtOAc (2 x 25 mL) and the combined extracts were dried over sodium sulfate and concentrated under reduced pressure to give 3-*iso*butoxy-1*H*-pyrazole (955 mg, 99%) as a slightly colored viscous oil. ESI-MS m/z calc. 140.09496, found 141.2 $(M+1)^+$; Retention time: 0.86 minutes; 1H NMR (400 MHz, Chloroform-d) δ 9.25 (very broad s, 1H), 7.35 (d, J = 2.5 Hz, 1H), 5.73 (d, J = 2.5 Hz, 1H), 3.91 (d, J = 6.7 Hz, 2H), 2.15 - 2.02 (m, 1H), 1.06 - 0.98 (m, 6H).


Step 4: *tert*-butyl 2-chloro-6-(3-*iso*butoxypyrazol-1-yl)pyridine-3-carboxylate



NaH (64 mg, 1.6 mmol) (60 % oil suspension) was added to 3-*iso*butoxy-1*H*-pyrazole (224 mg, 1.60 mmol) in anhydrous DMF (2 mL). The reaction mixture was stirred until


gas evolution stopped. *Tert*-butyl 2,6-dichloropyridine-3-carboxylate (330 mg, 1.33 mmol) was added and the reaction mixture was stirred at 100°C under nitrogen for 30 minutes. The reaction mixture was diluted with EtOAc (50 mL) and water + brine (40 mL) and the two phases were separated. The aqueous phase was further extracted with EtOAc (2 x 30 mL). The combined extracts were dried over sodium sulfate and the solvent removed under reduced pressure to give a residue, which was purified by silica gel column chromatography using a gradient of AcOEt (0 to 20%) in hexanes to give *tert*-butyl 2-chloro-6-(3-*iso*butoxypyrazol-1-yl)pyridine-3-carboxylate (235 mg, 50%) as an off-white solid). ESI-MS m/z calc. 351.13498, found 352.2 $(M+1)^+$; Retention time: 2.34 minutes, 1H NMR (400 MHz, Chloroform-d) δ 8.35 (d, J = 2.8 Hz, 1H), 8.18 (d, J = 8.4 Hz, 1H), 7.70 (d, J = 8.4 Hz, 1H), 5.97 (d, J = 2.9 Hz, 1H), 4.02 (d, J = 6.6 Hz, 2H), 2.12 (dp, J = 13.4, 6.7 Hz, 1H), 1.62 (s, 9H), 1.03 (d, J = 6.7 Hz, 6H).

Step 5: 2-chloro-6-(3-*iso*butoxypyrazol-1-yl)pyridine-3-carboxylic acid




Tert-butyl 2-chloro-6-(3-*iso*butoxypyrazol-1-yl)pyridine-3-carboxylate (203 mg, 0.57 mmol), DCM (2 mL) and TFA (615 µL, 7.98 mmol) were stirred at 40-45°C for 3 hours. The volatiles were remove under reduced pressure to give 2-chloro-6-(3-*iso*butoxypyrazol-1-yl)pyridine-3-carboxylic acid (170 mg, 100%) as a white solid. ESI-MS m/z calc. 295.07236, found 296.3 $(M+1)^+$; Retention time: 1.64 minutes; 1H NMR (400 MHz, DMSO-d6) δ 13.58 (s, 1H), 8.41 (d, J = 2.9 Hz, 1H), 8.38 (d, J = 8.4 Hz, 1H), 7.73 (d, J = 8.4 Hz, 1H), 6.20 (d, J = 2.9 Hz, 1H), 4.02 (d, J = 6.6 Hz, 2H), 2.06 (hept, J = 6.6 Hz, 1H), 0.98 (d, J = 6.7 Hz, 6H).

Step 6: *N*-[(2-amino-3-pyridyl)sulfonyl]-2-chloro-6-(3-*iso*butoxypyrazol-1-yl)pyridine-3-carboxamide



DMF (1.2 mL) was added to 2-chloro-6-(3-*iso*butoxypyrazol-1-yl)pyridine-3-carboxylic acid (128 mg, 0.43 mmol) and CDI (87 mg, 0.54 mmol) under an atmosphere of nitrogen. The reaction mixture was stirred at 40°C under nitrogen for 1 hour. 2-aminopyridine-3-sulfonamide (91 mg, 0.53 mmol) was dissolved under nitrogen in anhydrous DMF (1.2 mL) and NaH (21 mg, 0.53 mmol) (60% oil suspension) was added and the reaction mixture was stirred under nitrogen until gas evolution stopped and then stirred at 40°C for 45 minutes. The two reaction mixtures were combined and stirred at room temperature for 1.5 hours. Water and ice was added, followed by acetic acid (400 µL, 7.03 mmol). The resulting solid was filtered, washed with water and dried to give *N*-[(2-amino-3-pyridyl)sulfonyl]-2-chloro-6-(3-*iso*butoxypyrazol-1-yl)pyridine-3-carboxamide (189 mg, 97%) as a white solid. ESI-MS m/z calc. 450.0877, found 451.4 (M+1)+; Retention time: 1.42 minutes.

Step 7: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3-*iso*butoxypyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2191)




N-[(2-amino-3-pyridyl)sulfonyl]-2-chloro-6-(3-*iso*butoxypyrazol-1-yl)pyridine-3-carboxamide (177 mg, 0.39 mmol), (4*S*)-2,2,4-trimethylpyrrolidine (HCl salt) (237 mg, 1.58

mmol), K_2CO_3 (327 mg, 2.37 mmol) and anhydrous DMSO (550 µL) were heated at 150°C for 6 hours under an atmosphere of nitrogen. The reaction mixture was then cooled and a stirred mixture of water (40 mL) and acetic acid (400 µL, 7.034 mmol) was added dropwise. The resulting yellow solid was filtered and washed with water. The solid was dissolved in dichloromethane and the organic phase was dried over sodium sulfate and concentrated under reduced pressure. The resulting solid was purified by silica gel column chromatography using a gradient of MeOH (0 to 10%) in DCM, followed by purification by reverse phase preparative HPLC using a gradient of MeCN in water (1 to 99% over 15 min) and HCl as a modifier to give *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3-*iso*butoxypyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2191) (77 mg, 36%) as a pale yellow solid. ESI-MS m/z calc. 527.23145, found 528.3 $(M+1)^+$; Retention time: 1.55 minutes; 1H NMR (400 MHz, DMSO-d6) δ 12.45 (broad s, 1H), 8.28 - 8.22 (m, 1H), 8.20 (d, J = 2.7 Hz, 1H), 8.06 - 7.99 (m, 1H), 7.84 (d, J = 8.3 Hz, 1H), 6.94 (d, J = 8.3 Hz, 1H), 6.77 (dd, J = 7.7, 4.9 Hz, 1H), 6.69 (br s, 2H), 6.12 (d, J = 2.6 Hz, 1H), 3.98 (d, J = 6.6 Hz, 2H), 2.45 (d, J = 8.6 Hz, 2H), 2.13-2.01 (m, 2H), 1.90 - 1.79 (m, 1H), 1.55 (s, 3H), 1.53 (s, 3H), 1.38 (t, J = 12.1 Hz, 1H), 0.98 (d, J = 6.8 Hz, 6H), 0.74 (d, J = 6.2 Hz, 3H).

PREPARATION 109: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1975)


Step 1: 1-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]ethanone



To a suspension of 1-(3-hydroxypyrazol-1-yl)ethanone (1.21 g, 9.60 mmol) in anhydrous THF (20 mL) was added triphenyl phosphine (2.78 g, 10.6 mmol) and 2,2-dimethylpropan-1-ol (846 mg, 9.60 mmol). The reaction mixture was cooled down in an ice-bath under nitrogen. DIAD (2.1 mL, 10.7 mmol) was added dropwise via a syringe over 3 min. The reaction mixture was then stirred under nitrogen warming to room temperature overnight. The reaction mixture was diluted with ethyl acetate (100 mL) and the combined organic phase was washed with aqueous saturated sodium bicarbonate (2 x 50 mL), brine (50 mL) and dried over


sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography using a gradient of ethyl acetate (0 to 30 %) in hexanes. to provide 1-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]ethanone (967 mg, 51%) as a yellow solid. ESI-MS m/z calc. 196.12119, found 197.3 $(M+1)^+$; Retention time: 1.62 minutes; 1H NMR (400 MHz, Chloroform-d) δ 8.05 (d, J = 3.0 Hz, 1H), 5.97 (d, J = 3.0 Hz, 1H), 3.89 (s, 2H), 2.58 (s, 3H), 1.03 (s, 9H).

Step 2: 3-(2,2-dimethylpropoxy)-1*H*-pyrazole



1-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]ethanone (960 mg, 4.89 mmol), MeOH (15 mL) and NaOH (815 µL of 6 M, 4.89 mmol) (6N aqueous) were stirred at room temperature for 21 hours. The volatiles were removed under reduced pressure. The residue was taken in EtOAc (25 mL) and brine (20 ml) and two phases were separated. The aqueous phase was further extracted with EtOAc (2 x 25 mL) and the combined organic extracts were dried over sodium sulfate and evaporated under reduced pressure to give 3-(2,2-dimethylpropoxy)-1*H*-pyrazole (719 mg, 95%) as an off-white solid. ESI-MS m/z calc. 154.11061, found 155.3 $(M+1)^+$; Retention time: 1.13 minutes; 1H NMR (400 MHz, Chloroform-d) δ 7.36 (d, J = 2.4 Hz, 1H), 5.74 (d, J = 2.5 Hz, 1H), 3.81 (s, 2H), 1.02 (s, 9H).


Step 3: *Tert*-butyl 2-chloro-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]pyridine-3-carboxylate



NaH (95 mg, 2.375 mmol) (60 % oil suspension) (slight exotherm noticed) was added to 3-(2,2-dimethylpropoxy)-1*H*-pyrazole (363 mg, 2.35 mmol) in anhydrous DMF (3 mL)


and the reaction mixture was stirred until gas evolution stopped. *Tert*-butyl 2,6-dichloropyridine-3-carboxylate (530 mg, 2.14 mmol) was added portionwise and the reaction mixture was stirred at 100°C under nitrogen for 1 hour. The reaction mixture was diluted with EtOAc (60 mL) and water + brine (60 mL) and the two phases were separated. The aqueous phase was further extracted with EtOAc (2 x 30 mL). The combined extracts were dried over sodium sulfate and the solvent removed under reduced pressure. The residue was purified by silica gel column chromatography using a gradient of EtOAc (0 to 20% over 40 min) in hexanes to give *tert*-butyl 2-chloro-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]pyridine-3-carboxylate (497 mg, 64%) as a yellow solid. ESI-MS m/z calc. 365.1506, found 366.2 $(M+1)^+$; Retention time: 2.44 minutes.

Step 4: 2-chloro-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]pyridine-3-carboxylic acid




Tert-butyl 2-chloro-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]pyridine-3-carboxylate (497 mg, 1.36 mmol), DCM (5 mL) and TFA (1.5 mL, 19.5 mmol) were stirred at 40°C for 4 hours. The volatiles were removed by evaporation under reduced pressure to give 2-chloro-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]pyridine-3-carboxylic acid (413 mg, 98%) as a white solid. ESI-MS m/z calc. 309.088, found 310.1 $(M+1)^+$; Retention time: 1.8 minutes; 1H NMR (400 MHz, DMSO-d6) δ 13.58 (s, 1H), 8.42 (dd, J = 2.8, 0.8 Hz, 1H), 8.38 (d, J = 8.4 Hz, 1H), 7.74 (d, J = 8.5 Hz, 1H), 6.22 (dd, J = 2.8, 0.8 Hz, 1H), 3.93 (s, 2H), 1.01 (s, 9H).

Step 5: *N*-[(2-amino-3-pyridyl)sulfonyl]-2-chloro-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]pyridine-3-carboxamide



DMF (1 mL) was added to 2-chloro-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]pyridine-3-carboxylic acid (105 mg, 0.34 mmol) and CDI (69 mg, 0.43 mmol) and the reaction mixture was stirred at 40°C under nitrogen for 1 hour. In a separate 4 mL vial, 2-aminopyridine-3-sulfonamide (73 mg, 0.42mmol) was dissolved under nitrogen in anhydrous DMF (1 mL). NaH (22 mg, 0.55 mmol) (60% oil suspension) was added and the reaction mixture was stirred under nitrogen until gas evolution stopped, then stirred at 40°C for one hour. The two reaction mixtures were combined and stirred at room temperature for 4 hours. Water and ice was added, followed by acetic acid (400 µL, 7.034 mmol). The resulting solid was filtered, washed with water and dried to give *N*-[(2-amino-3-pyridyl)sulfonyl]-2-chloro-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]pyridine-3-carboxamide (150 mg, 95%) as a white solid. The product was used for the next step without any further purification. ESI-MS m/z calc. 464.10336, found 465.2 $(M+1)^+$; Retention time: 1.5 minutes.

Step 6: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1975)



N-[(2-amino-3-pyridyl)sulfonyl]-2-chloro-6-[3-(2,2-dimethylpropoxy)pyrazol-1-

yl]pyridine-3-carboxamide (150 mg, 0.32 mmol), (4*S*)-2,2,4-trimethylpyrrolidine (HCl salt) (195 mg, 1.30 mmol), K_2CO_3 (274 mg, 1.98 mmol) and anhydrous DMSO (480 µL) were stirred at 150°C for 7.5 hours under an atmosphere of nitrogen. The reaction mixture was poured into a mixture of water (25 mL) and acetic acid (400 µL, 7.034 mmol)and the resulting yellow solid was separated by filtration. The wet solid was dissolved in DCM, the organic phase was dried over sodium sulfate and the solvents evaporated to give a crude glass, which was purified by silica gel column chromatography using a gradient of MeOH (0 to 10%) in DCM. Evaporation of the solvent gave an off-white solid which was dissolved in MeOH (3 mL) and purified by reverse phase preparative HPLC using a gradient of MeCN in water (1 to 99% over 15 min) and HCl as a modifier (3 x 950 uL injections) to give *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1975) (75 mg, 42%) as a tan solid. ESI-MS m/z calc. 541.24713, found 542.3 $(M+1)^+$; Retention time: 1.65 minutes; 1H NMR (400 MHz, DMSO-d6) δ 12.45 (s, 1H), 8.24 (dd, J = 4.7, 1.9 Hz, 1H), 8.20 (d, J = 2.8 Hz, 1H), 8.03 (dd, J = 7.8, 1.9 Hz, 1H), 7.84 (d, J= 8.3 Hz, 1H), 6.95 (d, J = 8.2 Hz, 1H), 6.77 (dd, J = 7.8, 4.8 Hz, 1H), 6.69 (broad s, 2H), 6.14 (d, J = 2.7 Hz, 1H), 3.89 (s, 2H), 2.45 (d, J = 8.9 Hz, 2H), 2.12 (br s, 1H), 1.85 (dd, J = 11.9, 5.4 Hz, 1H), 1.55 (s, 3H), 1.53 (s, 3H), 1.43 - 1.33 (m, 1H), 1.00 (s, 9H), 0.74 (d, J = 6.2 Hz, 3H).

The following compounds can be synthesized using the procedures described herein:

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-*iso*butylpyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2036),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[4-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2406),

6-(3-*iso*propoxypyrazol-1-yl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2360),

N-(benzenesulfonyl)-6-(4-*iso*butoxypyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1912),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-ethoxy-4-methyl-pyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2248),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(2-ethoxyethoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1742),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[4-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2603),

6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2578),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(4-*iso*butoxypyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1895),

6-(3-*iso*butoxy-4-methyl-pyrazol-1-yl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1743),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-*iso*propoxypyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1844),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-[(2*S*)-2-methylbutoxy]pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2458),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-*sec*-butoxypyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1970),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(3-hydroxy-2-methyl-propoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2192),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-*iso*propoxypyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1709),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-*iso*butoxy-4-methyl-pyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1758),

N-cyclopropylsulfonyl-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2554),


6-(4-*iso*butoxypyrazol-1-yl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1760),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-*iso*butoxy-5-methyl-pyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2238),

N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-[2-(2-ethoxyethoxy)ethoxy]pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2326),
N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-*iso*butoxypyrazol-1-yl)-2-(6-methyl-8-azaspiro[3.4]octan-8-yl)pyridine-3-carboxamide (Compound 2604) (Isomer 1) (Compound 2481) (Isomer 2) (Compound 2194) (Isomer 3) (Compound 2615) (Isomer 4),
N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-[2,3,3,3-tetradeuterio-2-(trideuteriomethyl)propoxy]pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1887),
6-(3-ethoxy-4-methyl-pyrazol-1-yl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1856),
6-(3-*iso*butoxypyrazol-1-yl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1896),
N-[(6-amino-2-pyridyl)sulfonyl]-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1952),
N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-[3,3,3-trideuterio-2,2-bis(trideuteriomethyl)propoxy]pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2703),
N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(1-methylpropoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2704),
N-(benzenesulfonyl)-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1114),
N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-cyclopropylpyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2701),
N-cyclopropylsulfonyl-2-[(2*S*,5*R*)-2,5-dimethylpyrrolidin-1-yl]-6-(3-phenylpyrazol-1-yl)pyridine-3-carboxamide (Compound 2676),
N-cyclopropylsulfonyl-6-(3-phenylpyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2674),
N-[(2-amino-3-pyridyl)sulfonyl]-6-(3-phenylpyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2702),
N-[(2-amino-3-pyridyl)sulfonyl]-2-[(2*S*,5*R*)-2,5-dimethylpyrrolidin-1-yl]-6-(3-*iso*butoxypyrazol-1-yl)pyridine-3-carboxamide (Compound 1732),

N-[(3*R*)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-*iso*butoxypyrazol-1-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2716), *N*-cyclopropylsulfonyl-6-[3-(2-pyridyl)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2718), *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(2,2-dimethylpropoxy)-4-methyl-pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2699), *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(cyclobutylmethoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2694) and *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(cyclopropylmethoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2695).

PREPARATION 110: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3-*iso*butoxypyrazol-1-yl)-2-(6-methyl-8-azaspiro[3.4]octan-8-yl)pyridine-3-carboxamide (Compound 2604) (Isomer 1, *R* or *S* stereoisomer) and *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3-*iso*butoxypyrazol-1-yl)-2-(6-methyl-8-azaspiro[3.4]octan-8-yl)pyridine-3-carboxamide (Compound 2481) (Isomer 2, *R* or *S* stereoisomer)



The stereoisomers were separated using supercritical fluid chromatography on a ChiralPak AD-H (250 x 10 mm), 5μm column using 30% methanol in CO_2 at a flow rate of 10.0 mL/min. The separated enantiomers were separately concentrated, diluted with ethyl acetate (3 mL) and washed with 1N aqueous hydrochloric acid. The organic layers were dried over sodium sulfate, filtered, and evaporated to dryness to give the pure compounds. SFC Peak 1: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3-*iso*butoxypyrazol-1-yl)-2-(6-methyl-8-azaspiro[3.4]octan-8-yl)pyridine-3-carboxamide (Compound 2604) (Compound 2481) ESI-MS *m/z* calc. 539.23145, found 540.2 $(M+1)^+$; Retention time: 1.67 minutes . SFC Peak 2: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3-*iso*butoxypyrazol-1-yl)-2-(6-methyl-8-azaspiro[3.4]octan-8-yl)pyridine-3-carboxamide (Compound 2604) (Isomer 1, *R* or *S* stereoisomer) (Compound 2481) (Isomer 2, *R* or *S* stereoisomer) ESI-MS *m/z* calc. 539.23145, found 540.2 $(M+1)^+$; Retention time: 1.67 minutes.


The following compounds can be synthesized using the procedures described herein:

N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-2-[6-methyl-8-azaspiro[3.4]octan-8-yl]pyridine-3-carboxamide (Compound 2358) (Isomer 1, *R* or *S* stereoisomer), followed by SFC column chromatography using ChiralPak AS-H (250 x 10 mm), 5μm, using as eluent 38% MeOH (no modifier) in CO_2.

N-[(2-amino-3-pyridyl)sulfonyl]-6-[3-[(1-methylpropoxy)pyrazol-1-yl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2594) (Isomer 2, *R* or *S* stereoisomer) using ChiralPak IA (250 x 4.6 mm), 5μm; Mobile phase: 25 % MeOH:EtOH (50:50; 0.1% TFA) in CO_2.


PREPARATION 111: 4-(2,2-dimethylpropoxy)-1*H*-pyrazole

Step 1: *tert*-butyl 4-hydroxypyrazole-1-carboxylate



To *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazole-1-carboxylate (25 g, 85 mmol) in THF (170 mL) at 0 °C was added sodium hydroxide (85 mL of 2 M, 170 mmol) followed by hydrogen peroxide (19.27 mL of 30 %w/v, 170.0 mmol) and the reaction mixture was stirred at 0 °C for 10 min and RT for 45 min. The reaction mixture was cooled to 0 °C and diluted with DCM and 2 M HCl was added till pH 2 was reached. The organics were separated, dried and concentrated under reduced pressure to give *tert*-butyl 4-hydroxypyrazole-1-carboxylate (15 g 100%) as yellow solid. ESI-MS m/z calc 184.19, found 185.0 $(M+1)^+$; LCMS retention time (3 min run) 0.63 min.


Step 2: 4-(2,2-dimethylpropoxy)-1*H*-pyrazole



To *tert*-butyl 4-hydroxypyrazole-1-carboxylate (1000 mg, 5.429 mmol) and Ph_3P (2.136 g, 8.144 mmol) in THF (2.500 mL) was added 2,2-dimethylpropan-1-ol (717.9 mg, 8.144 mmol) at 0 °C. To the reaction mixture was added DIAD (1.647 g, 1.578 mL, 8.144 mmol) in THF (2 mL). The reaction mixture was stirred at 60 °C for 16h , then cooled and diluted with DCM and washed with 6 M HCl solution. The organics were separated, dried and concentrated to give a residue which was purified by silica gel column chromatography using 5-100% EA-hexanes followed by 5-40% EA-DCM to give *tert*-butyl 4-(2,2-dimethylpropoxy)pyrazole-1-carboxylate (435.5 mg) ESI-MS m/z calc 254.16 , found 255.4 $(M+1)^+$; LCMS retention time (3 min run) = 1.82 min; 1H NMR (400 MHz, Chloroform-d) δ 7.57 (d, J = 1.0 Hz, 1H), 7.51 (d, J = 1.0 Hz, 1H), 3.52 (s, 2H), 1.64 (s, 9H), 1.00 (s, 9H). To *tert*-butyl 4-(2,2-dimethylpropoxy)pyrazole-1-carboxylate (435.5 mg, 1.712 mmol) in DCM (0.2 mL) was added 4M HCl-dixoane (2 mL). After 5h at RT, the solvent was removed under reduced pressure and the residue was triturated with diethyl ether to give 4-(2,2-dimethylpropoxy)-1*H*-pyrazole (HCl salt) (210 mg, 20%) ESI-MS m/z calc 154.11 , found 155.4 $(M+1)^+$; LCMS retention time(3 min run) 1.13 min.


PREPARATION 112: 3-*iso*butoxy-4-methyl-1*H*-pyrazole

Step 1: 1-(3-hydroxy-4-methyl-pyrazol-1-yl)ethanone




4-methyl-1*H*-pyrazol-3-ol (1.0 g, 10.2 mmol) and pyridine (4.0 mL, 49.5 mmol) were heated at 95°C. A solution of acetic anhydride (1.0 g, 961.2 µL, 10.2 mmol) in pyridine (1.5 mL, 18.6 mmol) was added dropwise over a period of 2 minutes. The reaction mixture was then stirred at 95°C for an additional 2 hours. The solvents were removed under reduced pressure and the solid residue was triturated with diethylether (30 mL), filtered, washed with diethylether and dried to give 1-(3-hydroxy-4-methyl-pyrazol-1-yl)ethanone (1.2 g, 86%) as a white solid. ESI-MS m/z calc. 140.05858, found 141.1 $(M+1)^+$; Retention time 0.39 minutes; 1H NMR (400 MHz, DMSO-d6) δ 10.99 (s, 1H), 7.92 (q, J = 1.1 Hz, 1H), 2.43 (s, 3H), 1.88 (d, J = 1.2 Hz, 3H).

Step 2: 1-(3-*iso*butoxy-4-methyl-pyrazol-1-yl)ethanone



1-Bromo-2-methyl-propane (658 mg, 522 µL, 4.80 mmol) was added to 1-(3-hydroxy-4-methyl-pyrazol-1-yl)ethanone (612 mg, 4.37 mmol), potassium carbonate (905 mg, 6.55 mmol) and anhydrous DMF (5 mL) under nitrogen atmosphere. The reaction mixture was stirred at 80 °C for 2.5 hours, then diluted with EtOAc (50 ml) and water (25 mL) and two phases were separated. The aqueous phase (pH = 9-10) was further extracted with EtOAc (20 mL). The combined extracts were dried over sodium sulfate and the solvents evaporated to give a crude oil. The oil was purified by silica gel column chromatography using a gradient of AcOEt (0 to 30 %) in hexanes to provide 1-(3-*iso*butoxy-4-methyl-pyrazol-1-yl)ethanone (726 mg, 85%) as a colorless liquid. ESI-MS m/z calc. 196.12119, found 197.3 $(M+1)^+$; Retention time: 1.57 minutes; 1H NMR (400 MHz, Chloroform-d) δ 7.82 (s, 1H), 4.01 (d, J = 6.6 Hz, 2H), 2.53 (s, 3H), 2.10 (sep, J = 6.7 Hz, 1H), 1.95 (d, J = 1.0 Hz, 3H), 1.01 (d, J = 6.7 Hz, 6H).

Step 3: 3-*iso*butoxy-4-methyl-1*H*-pyrazole



1-(3-*iso*butoxy-4-methyl-pyrazol-1-yl)ethanone (726 mg, 3.70 mmol) was treated with MeOH (10 mL) and NaOH (617 µL of 6 M, 3.70 mmol) (6N aqueous) at room temperature for 2 hours. The volatiles were removed under reduced pressure. The residue was taken in EtOAc (25 mL) and brine (20 ml) and two phases were separated. The aqueous phase was further extracted with EtOAc (2 x 25 mL) and the combined extracts were dried over sodium sulfate. After evaporation of the solvent, 3-*iso*butoxy-4-methyl-1*H*-pyrazole (564 mg, 99%) was isolated as a slightly colored viscous oil. ESI-MS m/z calc. 154.11061, found 155.0 $(M+1)^+$; Retention time: 1.17 minutes; 1H NMR (400 MHz, Chloroform-d) δ 8.83 (broad s, 1H), 7.16 - 7.09 (m, 1H), 3.95 (d, J = 6.7 Hz, 2H), 2.09 (m, 1H), 1.95 (d, J = 0.9 Hz, 3H), 1.01 (d, J = 6.7 Hz, 6H).

The following compounds can be prepared following the procedures described

herein:

3-(2-ethoxyethoxy)-1*H*-pyrazole,

(*S*)-3-(2-methylbutoxy)-1*H*-pyrazole,

3-(*sec*-butoxy)-1*H*-pyrazole,

3-((1*H*-pyrazol-5-yl)oxy)-2-methylpropan-1-ol,

5-*iso*propoxy-1*H*-pyrazole,


5-*iso*butoxy-3-methyl-1*H*-pyrazole,

3-(2-(2-ethoxyethoxy)ethoxy)-1*H*-pyrazole and

3-(2-(methyl-d3)propoxy-2,3,3,3-d4)-1*H*-pyrazole

PREPARATION 113: *N*-[(6-amino-2-pyridyl)sulfonyl]-3-fluoro-4-(3-fluoro-5-*iso*butoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]benzamide (Compound 2634)


Step 1: *N*-[(6-amino-2-pyridyl)sulfonyl]-4-bromo-2,3-difluoro-benzamide



To 4-bromo-2,3-difluoro-benzoic acid (2.66 g, 11.2 mmol) in DMF (2.30 mL) was added at RT di(imidazol-1-yl)methanone (2.73 g, 16.8 mmol) and reaction mixture was heated at 65 °C for 1 h. Separately to 6-aminopyridine-2-sulfonamide (2.43 g, 14.0 mmol) in DMF (2 mL) was added sodium hydride (561 mg, 14.0 mmol) at 0 °C.and stirred for 10 min and RT for 1h. The reaction mixture was cooled back to 0 °C and the CDI adduct was added and reaction mixture was heated at 65 °C for 1.5 h, then cooled to RT and diluted with ethyl acetate and washed with brine solution. The organics were separated, dried and concentrated under reduced pressure to give a residue which was purified by silica gel column chromatography eluting with 0-100% ethyl acetate-hexanes to give *N*-[(6-amino-2-pyridyl)sulfonyl]-4-bromo-2,3-difluoro-benzamide (3.80 g, 986%). ESI-MS m/z calc 390.94 , found 394.2 $(M+4)^+$; LCMS


retention time (3 min run) 1.13 min.

Step 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-2,3-difluoro-4-(3-fluoro-5-*iso*butoxy-phenyl)benzamide



To mixture of *N*-[(6-amino-2-pyridyl)sulfonyl]-4-bromo-2,3-difluoro-benzamide (1.3 g, 3.3 mmol) and (3-fluoro-5-*iso*butoxy-phenyl)boronic acid (1.1 g, 5.0 mmol) and K_2CO_3 (1.4 g, 9.9 mmol) was added cyclopentyl(diphenyl)phosphane;dichloropalladium;iron (243 mg, 0.33 mmol) and DMF (8.8 mL) and water (2.2 mL) and the reaction mixture was stirred under nitrogen at 100 °C for 16 h. The reaction mixture was cooled, filtered over pad of Celite and filtrate was diluted with ethyl acetate and brine solution. The organics were separated, dried and concentrated to give a residue which was purified by silica gel column chromatography using 5-100% ethyl acetate-DCM as eluent to give *N*-[(6-amino-2-pyridyl)sulfonyl]-2,3-difluoro-4-(3-fluoro-5-*iso*butoxy-phenyl)benzamide (356 mg, 22%). 1H NMR (400 MHz, Methanol-d4) δ 7.64 - 7.53 (m, 2H), 7.29 (d, J = 7.4 Hz, 1H), 7.25 - 7.18 (m, 1H), 6.90 (d, J = 2.7 Hz, 1H), 6.88 - 6.81 (m, 1H), 6.73 (dt, J = 10.8, 2.3 Hz, 1H), 6.64 (d, J = 8.3 Hz, 1H), 3.79 (d, J = 6.4 Hz, 2H), 2.07 (hept, J = 6.7 Hz, 1H), 1.04 (d, J = 6.7 Hz, 6H); ESI-MS m/z calc 479.11 , found 480.2 $(M+1)^+$; LCMS retention time (3 min run) 1.78 min.


Step 3: *N*-[(6-amino-2-pyridyl)sulfonyl]-3-fluoro-4-(3-fluoro-5-*iso*butoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]benzamide (Compound 2634)



To *N*-[(6-amino-2-pyridyl)sulfonyl]-2,3-difluoro-4-(3-fluoro-5-*iso*butoxy-phenyl)benzamide (356 mg, 0.74 mmol) and K_2CO_3 (277 mg, 2.00 mmol) in NMP (650 µl) was added (4*S*)-2,2,4-trimethylpyrrolidine (HCl salt) (150 mg, 1.00 mmol) and the reaction mixture was stirred at 195 °C for 5 h, then cooled and filtered and diluted with EtOAc and brine solution. The organics were separated, dried and concentrated under reduced pressure to give a residue which was purified using silica gel column chromatography using 0-15% MeOH-DCM to give *N*-[(6-amino-2-pyridyl)sulfonyl]-3-fluoro-4-(3-fluoro-5-*iso*butoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]benzamide (Compound 2634) (42 mg, 2%) 1H NMR (400 MHz, Methanol-d4) δ 7.69 (s, 1H), 7.32 (t, J = 7.9 Hz, 1H), 7.26 (s, 1H), 7.08 (dd, J = 7.3, 0.8 Hz, 1H), 6.62 – 6.51 (m, 2H), 6.44 (dd, J = 19.3, 9.7 Hz, 2H), 3.58 (s, 1H), 3.49 (d, J = 6.4 Hz, 2H), 2.52 (t, J = 0.8 Hz, 1H), 2.45 (s, 1H), 1.90 (s, 1H), 1.82 – 1.69 (m, 2H), 1.53 (s, 1H), 1.04 (s, 3H), 0.92 (s, 6H), 0.74 (d, J = 6.7 Hz, 6H). ESI-MS m/z calc 572.2 , found 573.3 $(M+1)^+$; LCMS retention time (3 min run) 1.91 min.

PREPARATION 114: *N*-[(6-amino-5-hydroxy-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2331)


Step 1: 6-bromo-2-nitro-pyridin-3-ol



A solution of phenyl trimethyl ammonium tribromide (8.45 g, 22.3 mmol) in THF (68 mL) was added dropwise to a stirred solution of 2-nitropyridin-3-ol (3.00 g, 21.4 mmol) and triethylamine (4.33 g, 5.97 mL, 42.8 mmol) in THF (91 mL) at -40 °C. The addition was done over a period of 5 minutes and the reaction mixture was kept at -40 °C for 1h.


1N HCl and aq. $Na_2S_2O_3$ were added and the aq. phase was extracted with dichloromethane. The combined organic phase was washed with brine, dried over $MgSO_4$, filtered and the filtrate evaporated under reduced pressure. The residue was purified by silica gel column chromatography eluted with 30 to 100% EtOAc/Hexanes to afford the title compound as a yellow solid.

Step 2: 3-benzyloxy-6-bromo-2-nitro-pyridine



To a solution of 6-bromo-2-nitro-pyridin-3-ol (5.04 g, 23.0 mmol) in DMF (40 mL) was added potassium carbonate (6.36 g, 46.0 mmol) followed by the addition of bromomethylbenzene (4.13 g, 2.87 mL, 24.2 mmol). The reaction mixture was heated at 60 °C overnight, then partitioned between EtOAc and water. The aqueous layer was extracted with EtOAc (3x). The combined organic layers were washed with water (3x), brine, dried over $MgSO_4$, filtered and concentrated under reduced pressure. The crude material was suspended in hexane, collected via filtration and air dried to provide 3-benzyloxy-6-bromo-2-nitro-pyridine (6.22 g, 87%) as a yellow solid.


Step 3: methyl 3-[(5-benzyloxy-6-nitro-2-pyridyl)sulfonyl]propanoate



A mixture of 3-benzyloxy-6-bromo-2-nitro-pyridine (6.21 g, 20.1 mmol) and (3-methoxy-3-oxo-propyl)sulfinyloxysodium (3.50 g, 20.1 mmol) in DMSO (50 mL) was purged with nitrogen for 2 minutes. Iodocopper (7.66 g, 40.2 mmol) was added. The reaction


mixture was heated at 80 °C for 24 h, then partitioned between EtOAc and sat. NH_4Cl. The resulting precipitate was removed via filtration and washed with EtOAc. The aqueous layer was extracted with EtOAc (3x). The combined organic layers were washed with water (3x), brine, dried over $MgSO_4$, filtered and concentrated under reduced pressure. The crude material was purified by silica gel column chromatography (40-60% EtOAc-Hex) to provide methyl 3-[(5-benzyloxy-6-nitro-2-pyridyl)sulfonyl]propanoate (2.50 g, 33%) as a brown solid.

Step 4: 5-benzyloxy-6-nitro-pyridine-2-sulfinate



THF (1 mL) was added to a suspension of methyl 3-[(5-benzyloxy-6-nitro-2-pyridyl)sulfonyl]propanoate (76 mg, 0.2 mmol) in MeOH (2 mL). Sodium methoxide (400 µL of 0.5 M, 0.2 mmol) (in MeOH) was added dropwise. The reaction mixture was stirred at room temperature for 30 minutes, then evaporated under reduced pressure to provide a light brown solid. The crude material was used directly in next step without further purification.


Step 5: 5-benzyloxy-6-nitro-pyridine-2-sulfonamide



To a solution of amino hydrogen sulfate (3.68 g, 32.5 mmol) and sodium acetate (5.32 g, 65.0 mmol) in water (40 mL) was added 5-benzyloxy-6-nitro-pyridine-2-sulfinate (sodium salt) (2.06 g, 6.5 mmol) in MeOH (25 mL). The reaction mixture was stirred at room temperature overnight, then partitioned between water and EtOAc. The aqueous layer was extracted with EtOAc (3x). The combined organic layers were washed with brine, dried over


$MgSO_4$, filtered and concentrated under reduced pressure to provide 5-benzyloxy-6-nitro-pyridine-2-sulfonamide (1.10 g, 55%) as a crude material. The crude material (light yellow solid) was used directly in next step without further purification.

Step 6: *N*-[(5-benzyloxy-6-nitro-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide




To a mixture of 5-benzyloxy-6-nitro-pyridine-2-sulfonamide (155 mg, 0.50 mmol), 2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxylic acid (248 mg, 0.60 mmol) and HATU (228 mg, 0.60 mmol) in DMF (10 mL) was added potassium carbonate (207 mg, 1.50 mmol). The reaction mixture was heated at 60 °C for 7 h, partitioned between EtOAc and water. The aqueous layer was extracted with EtOAc (3x). The combined organic layers were washed with water (3x), brine, dried over $MgSO_4$, filtered and concentrated under reduced pressure. The crude material was purified by silica gel column chromatography (20-40% EtOAc-Hex) to provide *N*-[(5-benzyloxy-6-nitro-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (220 mg, 63%) as a yellow solid.

Step 7: *N*-[(6-amino-5-benzyloxy-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 1762)



To a solution of *N*-[(5-benzyloxy-6-nitro-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (215 mg, 0.31 mmol) in dioxane (4 mL) and ethanol (2 mL) was added Fe (171 mg, 3.07 mmol) and HCl (511 µL of 6 M, 3.07 mmol). The reaction mixture was heated at 65 °C for 30 min. The iron was removed and the precipitate was collected via filtration. The filtrate was concentrated to dryness, dissolved in DMSO, filtered and purified by preparative HPLC to provide *N*-[(6-amino-5-benzyloxy-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 1762) (25 mg, 12%) 1H NMR (400 MHz, DMSO-d6) δ 12.23 (s, 1H), 8.13 (d, J = 7.8 Hz, 1H), 7.81 (d, J = 7.9 Hz, 1H), 7.57 - 7.46 (m, 2H), 7.45 - 7.23 (m, 5H), 7.23 - 7.04 (m, 5H), 6.83 (dt, J = 10.9, 2.3 Hz, 1H), 6.33 (s, 2H), 5.24 (s, 2H), 3.61 (d, J = 6.6 Hz, 2H), 2.05 (s, 6H), 1.98 (dt, J = 13.4, 6.7 Hz, 1H), 0.96 (d, J = 6.6 Hz, 6H). ESI-MS m/z calc. 670.22614, found 670.0 $(M+1)^+$; Retention time: 3.6 minutes.


Step 8: *N*-[(6-amino-5-hydroxy-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2331)



To a suspension of *N*-[(6-amino-5-benzyloxy-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 1762) (21 mg, 0.03 mmol) in MeOH (20 mL) was added EtOAc (1 mL). The solution was purged with nitrogen for 2 min. Pd (3. mg, 0.003 mmol) was added. The reaction mixture was stirred under an atmosphere of hydrogen at room temperature overnight. The Pd catalyst was removed via filtration. The filtrate was concentrated to dryness. The residue was re-dissolved in DMSO and purified by preparative HPLC to provide *N*-[(6-amino-5-hydroxy-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2331) (5.0 mg, 24%). ESI-MS m/z calc. 580.1792, found 580.0 $(M+1)^+$; Retention time: 3.15 minutes.

PREPARATION 115: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(3-fluoro-5-*iso*butoxy-phenyl)-4-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyrimidine-5-carboxamide (Compound 2121)

Step 1: 2,4-dichloro-*N*-((6-nitropyridin-2-yl)sulfonyl)pyrimidine-5-carboxamide



A solution of 2,4-dichloropyrimidine-5-carboxylic acid (4.56 g, 23.8 mmol) in thionylchloride (30 mL) was refluxed for 1 hour, and evaporated to remove thionylchloride. The residue was dissolved in dichloromethane (50 mL), and 6-nitropyridine-2-sulfonamide (5.14 g, 23.7 mmol) was carefully added followed by trimethylamine (10 mL). The reaction mixture was stirred at room temperature for 30 min and concentrated. The crude material was purified by chromatography on silica gel, eluted with 30-100% of ethyl acetate in hexanes, to give the desired product (9.1 g, 100%).


Step 2: *N*-((6-aminopyridin-2-yl)sulfonyl)-2,4-dichloropyrimidine-5-carboxamide



A mixture of 2,4-dichloro-*N*-((6-nitropyridin-2-yl)sulfonyl)pyrimidine-5-carboxamide (1.7 g, 4.5 mmol, 1.0 eq) and iron (2.6 g, 47 mmol, 10 eq) in HCl (1.9 mL 23 mmol, 5.1 eq), ethanol (6 mL), tetrahydrofuran (6 mL), and water (0.90 mL) was heated at 60 °C for 1 hour. The reaction mixture was filtered through celite and washed with methanol and ethyl


acetate. The filtrate was concentrated under reduced pressure to give the desired compound (1.7 g, 100%) LC-MS: $(M+H)^+ = 348.0$.

Step 3: (*S*)-*N*-((6-aminopyridin-2-yl)sulfonyl)-2-chloro-4-(2,2,4-trimethylpyrrolidin-1-yl)pyrimidine-5-carboxamide



A mixture of *N*-((6-aminopyridin-2-yl)sulfonyl)-2,4-dichloropyrimidine-5-carboxamide (0.60 g, 1.7 mmol, 1.0 eq), (*S*)-2,2,4-trimethylpyrrolidine hydrochloride (0.26 g, 1.8 mmol, 1.0 eq), and potassium carbonate (0.96 g, 7.0 mmol, 4.0 eq) in 4.0 mL of dimethylsulfoxide was heated in a capped vial at 125 °C for 20 hours. The reaction mixture was purified chromatography on silica gel, and eluting with a gradient of 0-80% of methanol in ethyl acetate to give (*S*)-*N*-((6-aminopyridin-2-yl)sulfonyl)-2-chloro-4-(2,2,4-trimethylpyrrolidin-1-yl)pyrimidine-5-carboxamide as a brown solid (0.22 g, 30% yield). ^{1}H NMR (DMSO-d_6, 250 MHz): δ 8.04 (m, 1H), 7.48 (t, J = 7.3 Hz, 1H), 7.04 (d, J = 7.0 Hz, 1H), 6.51 (d, J = 8.3 Hz, 1H), 6.25 (s, br, 2H), 2.99 (m, 1H), 2.84 (m, 1H), 2.14 (m, 1H), 1.84 (m, 1H), 1.50 (s, 3H), 1.47 (s, 3H), 1.38 (m, 1H), 0.85 (d, J = 6.3 Hz, 3H) ppm. LC-MS: $(M+H)^+ = 424.9$.


Step 4: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(3-fluoro-5-*iso*butoxy-phenyl)-4-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyrimidine-5-carboxamide (Compound 2121)



To *N*-[(6-amino-2-pyridyl)sulfonyl]-2-chloro-4-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyrimidine-5-carboxamide (70 mg, 0.1647 mmol), cyclopentyl(diphenyl)phosphane;dichloropalladium;iron (12.05 mg, 0.01647 mmol), and dioxane (840.0 µL) was added (3-fluoro-5-*iso*butoxy-phenyl)boronic acid (69.84 mg, 0.3294 mmol), followed by sodium carbonate (247.0 µL of 2 M, 0.4941 mmol) under an atmosphere of nitrogen. The reaction mixture was irradiated at 120°C in the microwave for 1 hour. The reaction

mixture was filtered and then purified directly by reverse-phase preparative chromatography utilizing a C18 column and HPLC-MS method 10-99 A1B1 (Acetonitrile-Water + 5 mmolar HCl, 30 min method) to afford as a tan solid, *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(3-fluoro-5-*iso*butoxy-phenyl)-4-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyrimidine-5-carboxamide (Compound 2121) (14 mg, 15%) ESI-MS m/z calc. 556.2268, found 557.2 $(M+1)^+$; Retention time: 1.71 minutes. 1H NMR (400 MHz, DMSO) δ 8.53 (s, 1H), 7.73 (s, 3H), 7.70 - 7.63 (m, 1H), 7.59 (d, J = 9.7 Hz, 1H), 7.21 (d, J = 7.2 Hz, 1H), 7.07 (d, J = 10.6 Hz, 1H), 6.73 (d, J = 8.4 Hz, 1H), 3.86 (d, J = 6.6 Hz, 2H), 2.86 - 2.74 (m, 1H), 2.69 (t, J = 10.9 Hz, 1H), 2.34 - 2.18 (m, 1H), 2.13 - 1.99 (m, 1H), 1.95 (dd, J = 11.9, 5.5 Hz, 1H), 1.63 (d, J = 23.9 Hz, 6H), 1.50 (t, J = 12.2 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.90 (d, J = 6.3 Hz, 3H).

PREPARATION 116: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,2-dimethyl-3-tetrahydropyran-4-yl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2425) (Compound 2266) (racemic mixture)



In a 5 mL microwave vial was combined the *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-chloro-pyridine-3-carboxamide (85 mg, 0.2305 mmol) and fluorocesium (175.0 mg, 42.53 µL, 1.152 mmol) in DMSO (510.0 µL) followed by K_2CO_3 (318.6 mg, 2.305 mmol) and 2,2-dimethyl-3-tetrahydropyran-4-yl-pyrrolidine (169.0 mg, 0.9220 mmol). The vial was capped and heated to 150°C in an oil bath for 16 hours. Additional fluorocesium (175.0 mg, 42.53 µL, 1.152 mmol), K_2CO_3 (318.6 mg, 2.305 mmol) and 2,2-dimethyl-3-tetrahydropyran-4-yl-pyrrolidine (169.0 mg, 0.9220 mmol) was added and heated to 165°C for 16 additional hours. The reaction mixture was filtered and then purified directly by reverse-phase preparative chromatography utilizing a C18 column and HPLC-MS method 10-99% acetonitrile-water + 5 mmolar HCl to afford *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,2-dimethyl-3-tetrahydropyran-4-yl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2425) (Compound 2266) (80 mg, 66%)as a light yellow solid, ESI-MS m/z calc. 515.25665, found 516.2 $(M+1)^+$;

Retention time: 1.31 minutes. 1H NMR (400 MHz, DMSO) δ 12.47 (s, 1H), 7.70 - 7.52 (m, 2H), 7.17 (dd, J = 7.3, 0.5 Hz, 1H), 6.78 - 6.59 (m, 2H), 5.14 (s, 2H), 3.84 (d, J = 3.5 Hz, 2H), 3.30 (d, J = 10.7 Hz, 2H), 2.84 (t, J = 10.0 Hz, 1H), 2.68 - 2.55 (m, 1H), 2.07 (s, 4H), 1.87 - 1.73 (m, 1H), 1.65 (dd, J = 30.6, 10.5 Hz, 2H), 1.55 (d, J = 8.6 Hz, 6H), 1.49 (s, 3H), 1.28 (d, J = 6.6 Hz, 10H).

The stereoisomers were separated using supercritical fluid chromatography on a ChiralPak AS-H (250 x 10 mm), 5μm column using 12 % MeOH in CO_2 at a flow rate of 10.0 mL/min. The separated enantiomers were separately concentrated to dryness to give the pure compounds.

The following compounds can be prepared in the manner described herein:

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(3-cyclopropyl-2,2-dimethyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2577) (Isomer 1, *R* or *S* stereoisomer) (Compound 2022) (Isomer 2, *R* or *S* stereoisomer),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-[3-(cyclopropylmethyl)-2,2-dimethyl-pyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2242),


N-[(6-amino-2-pyridyl)sulfonyl]-2-(3-benzyl-2,2-dimethyl-pyrrolidin-1-yl)-6-*tert*-butyl-pyridine-3-carboxamide (Compound 1852),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(3-cyclopropyl-2,2-dimethyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2577) (Isomer 1, *R* or *S* stereoisomer) (Compound 2022) (Isomer 2, *R* or *S* stereoisomer),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,2-dimethyl-3-tetrahydropyran-4-yl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2425) (Isomer 1, *R* or *S* stereoisomer) (Compound 2266) (Isomer 2, *R* or *S* stereoisomer),

6-*tert*-butyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1765)

PREPARATION 117: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-(2-cyclopentyl-1,2-dihydroxyethyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Isomer 1, *R* or *S* stereoisomer) and *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-(2-cyclopentyl-1,2-dihydroxyethyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Isomer 2, *R* or *S* stereoisomer)




To a solution of *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-[(E)-2-cyclopentylvinyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1633) (8 mg, 0.01 mmol) in acetone (1.0 mL) and water (0.5 mL), were added 4-methylmorpholine *N*-oxide (16 mg, 0.14 mmol) and osmium tetroxide (15 µL of 2.5 %w/v, 0.001 mmol), in that order, and the reaction mixture was stirred under air at ambient temperature. After 2 h, the reaction mixture turned to a clear solution, and was concentrated to a light yellow paste. The paste was dissolved in DMSO (1 mL) and purified by reverse-phase HPLC-MS method to furnish the two diastereomeric diols as white solids. Isomer 1: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-(2-cyclopentyl-1,2-dihydroxy-ethyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (HCl salt) (2 mg, 23%) ESI-MS m/z calc. 573.29846, found 574.4 $(M+1)^+$; Retention time: 1.51 minutes; Isomer 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-(2-cyclopentyl-1,2-dihydroxy-ethyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (HCl salt) (4 mg, 47%) ESI-MS m/z calc. 573.29846, found 574.4 $(M+1)^+$; Retention time: 1.55 minutes

The following compounds can be prepared in the manner as described herein: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-(2-cyclopentyl-1,2-dihydroxyethyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2661) (Isomer 1), *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-(1,2-dihydroxy-3,3-dimethylbutyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2069) (Isomer 2), *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-(2-cyclohexyl-1,2-dihydroxyethyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (isomer 1), *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-(2-cyclohexyl-1,2-dihydroxyethyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (isomer 2),


PREPARATION 118: 6-methoxy-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]quinoline-3-carboxamide (Compound 2419)

Step 1: 2-chloro-6-methoxy-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]quinoline-3-carboxamide



2-Chloroquinoline-3-carboxylic acid (62 mg, 0.3 mmol) and CDI (73 mg, 0.5mmol) were combined in DMF (2 mL) and stirred at 40 °C for 2 hours. In a separate vial, NaH (15 mg, 0.4 mmol) was added to 2-methoxypyridine-3-sulfonamide (71 mg, 0.4 mmol) in DMF (1 mL). The solutions were combined and stirred overnight at ambient temperature. The reaction mixture was taken up in EtOAc and washed with HCl (1 N, 1 × 5 mL). The aqueous layer was back washed with EtOAc (2 x 10 mL), the combined organic layers were washed with brine (1× 10 mL) and dried over sodium sulfate, filtered and concentrated under reduced pressure. The solid was dissolved in 1,4-dioxane (2 mL) with 4M HCl (750 uL, 4.0 mmol) and heated to 90 °C for 2 hours. The reaction mixture was cooled, the solvent was removed under reduced pressure to yield a solid, which was triturated with ethyl acetate to give the desired compound (89 mg, 79%). ESI-MS m/z 394.2 $(M+1)^+$, retention time 0.39 minutes.

Step 2: 6-methoxy-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]quinoline-3-carboxamide (Compound 2419) (Compound 2405)



2-Chloro-*N*-methylsulfonyl-quinoline-3-carboxamide (40 mg, 0.1 mmol), potassium carbonate (69 mg, 0.5 mmol), and (4*S*)-2,2,4-trimethylpyrrolidine (HCl salt) (30 mg, 0.2 mmol) was suspended in DMSO (1 mL). The reaction was heated at 120 °C for 2 days, then cooled, diluted with DMSO-methanol, filtered, and then purified via reverse phase HPLC-MS method using a dual gradient run from 1-99% mobile phase B over 15.0 minutes (mobile phase A = H_2O (5 mM TFA), mobile phase B = acetonitrile) to give the desired compound (8 mg, 17%). ESI-MS m/z 471.3 $(M+1)^+$, retention time 1.19 minutes.

The following compounds can be synthesized using the procedures described herein:

6-methyl-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]quinoline-3-carboxamide (Compound 2471),

N-[(6-amino-2-pyridyl)sulfonyl]-7-methyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]quinoline-3-carboxamide (Compound 2519),

N-[(6-amino-2-pyridyl)sulfonyl]-6-methoxy-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]quinoline-3-carboxamide (Compound 2196),

N-[(6-amino-2-pyridyl)sulfonyl]-6-methyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]quinoline-3-carboxamide (Compound 2548),


N-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]quinoline-3-carboxamide (Compound 2405),

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]quinoline-3-carboxamide (Compound 2265) and

N-[(2-methoxy-3-pyridyl)sulfonyl]-7-methyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]quinoline-3-carboxamide (Compound 1859)

PREPARATION 119: *N*-[(6-amino-2-pyridyl)sulfonyl]-5-fluoro-6-*iso*propoxy-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2230)


Step 1: *N*-[(6-amino-2-pyridyl)sulfonyl]-2,6-dichloro-5-fluoro-pyridine-3-carboxamide



2,6-Dichloro-5-fluoro-pyridine-3-carboxylic acid (5.0 g, 23.8 mmol), CDI (6.8 g, 41.9 mmol) and DMF (30 mL) were stirred at 60°C for 50 min. In a separate flask, 6-aminopyridine-2-sulfonamide (5.2 g, 29.8 mmol), NaH (1.2 g, 29.8 mmol), and DMF (25 mL). The reaction mixture was stirred at 60°C for 1 hour. The contents containing 6-aminopyridine-2-sulfonamide was added dropwise to the activated acid at room temperature. The reaction mixture


was stirred for 16 hours at room temperature. The reaction mixture was then placed in an ice bath, diluted with water (300 mL), and adjusted pH to 2. The resulting slurry was vigorously stirred for 30 min. The precipitate was filtered, washed with water (50 ML x 3) and hexanes (50 mL x 3), and dried *in vacuo* to furnish *N*-[(6-amino-2-pyridyl)sulfonyl]-2,6-dichloro-5-fluoro-pyridine-3-carboxamide (6.9 g, 79%).

Step 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-5-fluoro-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide



To a solution of *N*-[(6-amino-2-pyridyl)sulfonyl]-2,6-dichloro-5-fluoro-pyridine-3-carboxamide (1.0 g, 2.7 mmol) in DMSO (8 mL) was added 2,2,4-trimethylpyrrolidine (HCl salt) (820 mg, 5.5 mmol) and K_2CO_3 (2.3 g, 16.4 mmol). The reaction was stirred at 150°C for 16 hours. The reaction mixture was filtered and purified via prep HPLC (1-99% ACN, 15 min, HCl modifier) to afford *N*-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-5-fluoro-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (HCl salt) (200 mg, 30%).

Step 3: *N*-[(6-amino-2-pyridyl)sulfonyl]-5-fluoro-6-*iso*propoxy-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2230)



N-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-5-fluoro-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (HCl salt) (50 mg, 0.10 mmol), *iso*propanol (31mg, 40 µL, 0.52 mmol), NaH (10 mg, 0.24 mmol), and DMSO (1mL). The reaction mixture was stirred at

150°C for 2 hours. The reaction mixture was then filtered and purified via prep HPLC (1-99% ACN, 15min, HCL modifier) to yield *N*-[(6-amino-2-pyridyl)sulfonyl]-5-fluoro-6-*iso*propoxy-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2230) (HCl salt) (13 mg, 23%) 1H NMR (400 MHz, DMSO-d6) δ 12.26 (s, 1H), 7.66 - 7.55 (m, 2H), 7.17 (dd, J = 7.3, 0.7 Hz, 1H), 6.66 (dd, J = 8.4, 0.8 Hz, 1H), 6.41 (s, 2H), 5.26 (p, J = 6.1 Hz, 1H), 2.60 - 2.51 (m, 1H), 2.49 - 2.40 (m, 1H), 2.33 (p, J = 1.8 Hz, 0H), 2.20 (s, 1H), 1.84 (dd, J = 11.8, 5.6 Hz, 1H), 1.51 (s, 3H), 1.45 (s, 3H), 1.43 - 1.30 (m, 4H), 1.29 (d, J = 6.2 Hz, 3H), 0.85 (d, J = 6.3 Hz, 3H). ESI-MS m/z calc. 465.1846, found 466.3 $(M+1)^+$; Retention time: 1.38 minutes.


The following compounds can be synthesized using the procedures described herein:

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1-cyclopropylethoxy)-5-fluoro-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2373) and

N-[(6-amino-2-pyridyl)sulfonyl]-6-(1,2-dimethylpropoxy)-5-fluoro-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1810)

PREPARATION 120: *N*-[(3*R*)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(*p-tolyl*)pyridine-3-carboxamide (Compound 2276)


Step 1: 2-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxylic acid



2,6-Dichloropyridine-3-carboxylic acid (1.000 g, 4.861 mmol) in IPA (30 mL), K_2CO_3 (6.075 mL of 2 M, 12.15 mmol) and dichloropalladium;triphenylphosphane (341.2 mg, 0.4861 mmol) was added. The reaction mixture was heated to 80 °C under an atmosphere of nitrogen and (3-fluoro-5-*iso*butoxy-phenyl)boronic acid (1.082 g, 5.104 mmol) in IPA (30 mL) was added slowly over 1 hour. The reaction mixture was heated for a further 4 h. The reaction mixture was cooled, 6 M HCl (2 mL) was added and then pH adjusted to 2-3 with additional 6 M HCl. The organic phase was collected, dried over sodium sulfate and concentrated under reduced pressure. The residue was purified by revere phase column chromatography using 10-80% ACN in Water (with TFA modifier) to give the desired compound. ESI-MS m/z 324.0 $(M+1)^+$;


Retention time: 1.0 minutes.

Step 2: 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(*p-tolyl*)pyridine-3-carboxylic acid



A mixture of 2-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxylic acid (324 mg, 1.00 mmol), *p-tolyl*boronic acid (272 mg, 2.00 mmol), K_2CO_3 (415 mg, 3.00 mmol), and tetrakis(triphenylphosphine)palladium (0) (116 mg, 0.10 mmol) in DMF (10 mL) was degassed under a stream of nitrogen and stirred at 120 °C for 16 hours. A further aliquot of tetrakis(triphenylphosphine)palladium (0) (116 mg, 0.10 mmol) was added, and the reaction mixture was stirred at 150 °C for 17 hours. The reaction mixture was diluted with water, adjusted to pH 5 with 1 M HCl, and extracted with ethyl acetate. The combined extracts were washed with brine, dried over sodium sulfate, and evaporated under reduced pressure. The residue was purified by silica gel chromatography using 0-5% methanol in dichloromethane as eluent to give 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(*p-tolyl*)pyridine-3-carboxylic acid (314 mg, 83%).


Step 3: *N*-[(3*R*)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(*p-tolyl*)pyridine-3-carboxamide (Compound 2276)



A solution of 6-(3-fluoro-5-*iso*butoxyphenyl)-2-(*p-tolyl*)nicotinic acid (38 mg, 0.1 mmol) and *N*-(oxomethylene)sulfamoyl chloride (17 mg, 10 µL, 0.1 mmol) in

dichloromethane (1 mL) was stirred for an hour, then *tert*-butyl *N*-[(3R)-pyrrolidin-3-yl]carbamate (28 mg, 0.2 mmol) and triethylamine (15 mg, 21 µL, 0.2 mmol) were added. After 17 hours, HCl (200 µL of 4 M, 0.8 mmol) (in dioxane) was added, and the reaction mixture was stirred for two hours.The reaction mixture was filtered and purified using a reverse phase HPLC-MS method using a Luna C18 (2) column (75 x 30 mm, 5 µm particle size) and a dual gradient run from 1-99% mobile phase B over 15.0 minutes (mobile phase A = H_2O (5 mM HCl), mobile phase B = CH_3CN. Flow rate = 50 mL/min, and column temperature = 25 °C) to give the desired compound. ESI-MS m/z 527.4 $(M+1)^+$; Retention time: 1.62 minutes.


PREPARATION 121: *N*-[(2-acetamido-3-pyridyl)sulfonyl]-6-(6-*iso*propoxy-3-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1976)



N-[(2-amino-3-pyridyl)sulfonyl]-6-(6-*iso*propoxy-3-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (80 mg, 0.1525 mmol) in pyridine (800 µL, 9.891 mmol) was treated with acetic anhydride (400 µL, 4.239 mmol) and stirred at room temperature overnight. The product was isolated by prep. HPLC to give *N*-[(2-acetamido-3-pyridyl)sulfonyl]-6-(6-*iso*propoxy-3-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1976) (HCl salt) (20 mg, 21%) 1H NMR (400 MHz, DMSO-d6) δ 8.85 (s, 1H), 8.67 (d, J = 4.7 Hz, 1H), 8.43 (d, J = 7.9 Hz, 1H), 8.30 (d, J = 9.0 Hz, 1H), 7.84 (d, J = 8.0 Hz, 1H), 7.53 - 7.42 (m, 1H), 7.28 (d, J = 8.0 Hz, 1H), 6.87 (d, J = 8.7 Hz, 1H), 5.31 (p, J = 6.3 Hz, 1H), 2.41 (s, 2H), 2.25 (s, 3H), 2.18 (dt, J = 16.2, 7.6 Hz, 1H), 1.88 (dd, J = 12.1, 5.8 Hz, 1H), 1.55 (d, J = 4.4 Hz, 6H), 1.42 (t, J = 12.0 Hz, 1H), 1.31 (d, J = 6.1 Hz, 6H), 0.74 (d, J = 6.3 Hz, 3H). ESI-MS m/z calc. 566.23114, found 567.0 $(M+1)^+$; Retention time: 2.19 minutes.


PREPARATION 122: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-ethoxy-4-methyl-2-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2256)

Step 1: 2-chloro-6-(6-fluoro-4-methyl-2-pyridyl)pyridine-3-carboxylic acid



2,6-Dichloronicotinic acid (661 mg, 3.44 mmol), (6-fluoro-4-methyl-2-pyridyl)boronic acid (560 mg, 3.62 mmol), K_2CO_3 (4.13 mL of 2 M, 8.26 mmol) in IPA (20 mL) was added $Pd(PPh_3)_2Cl_2$ (48.3 mg, 0.07 mmol). The reaction mixture was placed under an atmosphere of nitrogen and heated to 80 °C for 10 h. The reaction mixture was filtered, the solvent was evaporated under reduced pressure and the crude mixture was purified by reverse phase column chromatography using 10-80% acetonitrile in water (TFA modifier) to give the desired compound. ESI-MS m/z 267.0 $(M+1)^+$; Retention time: 0.82 minutes.


Step 2: 2-chloro-6-(6-ethoxy-4-methyl-2-pyridyl)pyridine-3-carboxylic acid



2-chloro-6-(6-fluoro-4-methyl-2-pyridyl)pyridine-3-carboxylic acid (4.2 g, 15 mmol) was dissolved into a solution of sodium ethoxide (25 mL of 21 %w/v, 79 mmol) in ethanol. The reaction mixture was heated to 75 °C and stirred for 3 hours. The reaction mixture was diluted with EtOAc (75 mL) and washed with water (1× 75 mL) with some brine (20 mL). The aqueous layer was extracted with EtOAc (1× 75 mL). The organic layers were combined, dried over sodium sulfate, filtered and concentrated under reduced pressure to give an orange solid. The solid was redissolved in EtOAc (75 mL) and aqueous NaOH solution (1 M, 75 mL). The aqueous layer was isolated and acidified to pH 2, extracted with EtOAc (2× 75 mL). The final organic layers were dried over sodium sulfate, filtered and concentrated under reduced pressure. The obtained solid was purified by silica gel column chromatography column, using 0-10% MeOH/DCM as eluent to give 2-chloro-6-(6-ethoxy-4-methyl-2-pyridyl)pyridine-3-carboxylic acid (1.17 g, 25%) as a white solid. ESI-MS m/z calc. 292.06146, found 293.1 $(M+1)^+$; Retention time: 1.72 minutes.


Step 3: *N*-[(2-amino-3-pyridyl)sulfonyl]-2-chloro-6-(6-ethoxy-4-methyl-2-

pyridyl)pyridine-3-carboxamide



2-chloro-6-(6-ethoxy-4-methyl-2-pyridyl)pyridine-3-carboxylic acid (200 mg, 0.68 mg) was dissolved in DMF (2.5 mL). CDI (123 mg, 0.75 mmol) was added, and the reaction mixture was allowed to stir at 60 °C for 1 hour. In a separate flask, sulfonamide was dissolved in DMF (600 µL), and NaH (34 mg, 0.85 mmol) was added and the reaction mixture was stirred for 45 minutes at room temperature. The activated carboxylic acid mixture was then added to the deprotonated sulfonamide. The final reaction mixture was stirred at 60 °C overnight. After cooling to room temperature, the reaction mixture was diluted with EtOAc (50 mL) and washed with water (1× 50 mL) and brine (2× 75 mL). The organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure to give the desired compound (297 mg, 97%). ESI-MS m/z 448.0 $(M+1)^+$; Retention time: 0.53 minutes.

Step 4: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-ethoxy-4-methyl-2-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2256)




2-chloro-6-(6-ethoxy-4-methyl-2-pyridyl)-*N*-methylsulfonyl-pyridine-3-carboxamide (60 mg, 0.13 mmol) and (4*S*)-2,2,4-trimethylpyrrolidine (free base, 45 mg, 0.4 mmol) were combined in DMSO (100 µL). Potassium carbonate (55 mg, 0.4 mmol) was added and the reaction mixture was heated overnight at 140 °C, then 165 °C for 3 hours. The reaction

mixture was purified using a reverse phase HPLC method using a Luna C18 (2) column (50 × 21.2 mm, 5 µm particle size) and a dual gradient run from 10-99% mobile phase B over 15.0 minutes (mobile phase A = water (de-ionized, no acid modifier), mobile phase B = acetonitrile) to give the desired compound. ESI-MS m/z 525.3 $(M+1)^+$; Retention time: 1.57 minutes.


PREPARATION 123: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[6-(diethylamino)-2-pyridyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1845)

Step 1: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-fluoro-2-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide



A mixture of *N*-[(2-amino-3-pyridyl)sulfonyl]-6-chloro-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (0.12 g, 0.28 mmol), (6-fluoro-2-pyridyl)boronic acid (0.06 g, 0.42 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (0.02 g, 0.02 mmol), and Na_2CO_3 (0.09 g, 0.85 mmol) in DMF (2.2 mL) and water (566 µL) was degassed under a stream of nitrogen and stirred at 100 °C for 20 hours. The reaction mixture was diluted with water and extracted with ethyl acetate. The combined extracts were washed with brine, dried over sodium sulfate, and evaporated under reduced pressure. The residue was purified by silica gel column chromatography using 0-5% methanol in dichloromethane as eluent to give *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-fluoro-2-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (0.11 g, 80%) as yellow solid.

Step 2: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[6-(diethylamino)-2-pyridyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1845)



A solution of *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-fluoro-2-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (30 mg, 0.06 mmol) and diethylamine (1 mL, 9.67 mmol) was stirred in a sealed vessel at 120 °C for 17 hours, then 150 °C for 22 hours. The diethylamine was removed under vacuum, and the residue was purified using a reverse phase HPLC-MS method using a Luna C18 (2) column (75 × 30 mm, 5 µm particle size) and a dual gradient run from 25-75% mobile phase B over 15.0 minutes (mobile phase A = H_2O (5 mM HCl), mobile phase B = CH_3CN) to give the desired compound. ESI-MS m/z 483.5 $(M+1)^+$; Retention time: 0.88 minutes

The following compounds can be synthesized using the procedures described herein

N-[(2-amino-3-pyridyl)sulfonyl]-6-[6-(2,2-dimethylpropylamino)-2-pyridyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2500),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(diethylamino)-5-methyl-2-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 1907),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(*iso*butylamino)-5-methyl-2-pyridyl]-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2362),


N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(diethylamino)-4-methyl-2-pyridyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2002),

6-(6-ethoxy-4-methyl-2-pyridyl)-*N*-(1*H*-pyrazol-3-ylsulfonyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2487),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[2-(2-ethoxyethoxy)ethoxy]-4-methyl-2-pyridyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1696),

N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2-*iso*propoxyethoxy)-4-methyl-2-pyridyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1703),
N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2-ethoxyethoxy)-4-methyl-2-pyridyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1700),
N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[isopropyl(methyl)amino]-5-methyl-3-pyridyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2687),
N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(isopropylamino)-5-methyl-3-pyridyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2688)
N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[ethyl(methyl)amino]-5-methyl-3-pyridyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2689),
N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[2-ethoxyethyl(methyl)amino]-5-methyl-3-pyridyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2685),
N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-[2-(2-ethoxyethoxy)ethyl-methyl-amino]-5-methyl-3-pyridyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2684) and
N-[(6-amino-2-pyridyl)sulfonyl]-6-[6-(2-ethoxyethylamino)-5-methyl-3-pyridyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2686).

PREPARATION 124: *N*-(benzenesulfonyl)-6-(6-ethoxy-4-methyl-2-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 2455)



N-[(2-amino-3-pyridyl)sulfonyl]-2-chloro-6-(6-ethoxy-4-methyl-2-pyridyl)pyridine-3-carboxamide (50 mg, 0.1 mmol) and 2,4,6-trimethyl-phenol (45 mg, 0.3 mmol) were dissolved in DMSO (175 µL) and dioxane (175 µL). Sodium hydride (13mg, 0.3 mmol) was added and the reaction mixture was heated under microwave irradiation at 150 °C for 20 minutes. The reaction mixture was purified using a reverse phase HPLC method using a Luna C18 (2) column (50 × 21.2 mm, 5 µm particle size) and a dual gradient run from 10-99% mobile phase B over 15.0 minutes (mobile phase A = water (de-ionized, no acid modifier),

mobile phase B = acetonitrile) to give *N*-(benzenesulfonyl)-6-(6-ethoxy-4-methyl-2-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 2455). ESI-MS m/z 532.5 $(M+1)^+$; Retention time: 2.52 minutes.

The following compounds can be synthesized using the procedures described herein:

N-[(6-amino-2-pyridyl)sulfonyl]-6-(6-ethoxy-4-methyl-2-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 2511),


N-[(2-amino-3-pyridyl)sulfonyl]-6-(6-ethoxy-4-methyl-2-pyridyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1795),

6-(6-ethoxy-4-methyl-2-pyridyl)-*N*-(1*H*-pyrazol-3-ylsulfonyl)-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 2301) and

6-(6-ethoxy-4-methyl-2-pyridyl)-*N*-[(2-oxo-1*H*-pyridin-3-yl)sulfonyl]-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 1768).


PREPARATION 125: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-tetrahydropyran-2-yl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1697)

Step 1:*N*-[(2-amino-3-pyridyl)sulfonyl]-2,6-dichloro-pyridine-3-carboxamide




2,6-dichloropyridine-3-carboxylic acid (1.5 g, 7.8 mmol) and CDI (1.9 g, 11.7 mmol) in DMF (17 mL) was stirred at 60 °C. In a separate vial, 2-aminopyridine-3-sulfonamide (1.7 g, 9.8 mmol) and NaH (0.4 g, 9.8 mmol) in DMF (8 mL) was stirred at ambient temperature. The two reaction mixtures were combined and stirred overnight at room temperature. The reaction mixture was taken up in EtOAc and washed with HCl (1 N, 1 x 25 mL) and the aqueous layer was washed with EtOAc (4 x 20 mL). The combined organic layers were washed with brine (1× 15 mL) and dried over sodium sulfate, filtered, and concentrated under reduced pressure. The resulting solid that crashed was washed with ethyl acetate and the mother liquor was concentrated, the residue was washed with ethyl acetate and hexanes to obtain *N*-[(2-amino-3-pyridyl)sulfonyl]-2,6-dichloro-pyridine-3-carboxamide (2.5 g, 94%) as a white solid. 1H NMR (400 MHz, DMSO-d6) δ 8.28 - 8.12 (m, 2H), 8.07 (d, J = 8.0 Hz, 1H), 7.62 (d, J = 8.0 Hz, 1H), 6.87 (dd, J = 7.7, 5.4 Hz, 1H).

Step 2: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-chloro-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide



DMSO (17 mL) was added to *N*-[(2-amino-3-pyridyl)sulfonyl]-2,6-dichloro-pyridine-3-carboxamide (2.1 g, 6.0 mmol), (4*S*)-2,2,4-trimethylpyrrolidine (HCl salt) (1.1 g, 7.3 mmol), and potassium carbonate (3.3g, 24 mmol). The reaction mixture was sealed and stirred at 120°C for 2 days. The reaction mixture was allowed to cool and then diluted with ethyl acetate (40 mL) and water (60 mL). To the aqueous layer, aq HCl was added to pH 7 and then washed with ethyl acetate (5 x 50 mL). The combined organic layers were concentrated to half volume then washed with brine and dried over sodium sulfate, filtered and evaporated under reduced pressure to give *N*-[(2-amino-3-pyridyl)sulfonyl]-6-chloro-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (2.5 g, 100%) as a yellow powder. 1H NMR (400 MHz, DMSO-d6) δ 8.26 (dd, J = 4.8, 1.8 Hz, 1H), 8.06 (dd, J = 7.9, 1.8 Hz, 1H), 7.71 (d, J = 7.9 Hz, 1H), 6.79 (dd, J = 7.9, 4.8 Hz, 1H), 6.64 (d, J = 7.9 Hz, 1H), 2.46 - 2.34 (m, 2H), 2.19 - 2.03 (m, 1H), 1.82 (dd, J = 12.1, 5.6 Hz, 1H), 1.47 (d, J = 3.3 Hz, 6H), 1.36 (t, J = 12.2 Hz, 1H), 0.74 (d, J = 6.3 Hz, 3H).


Step 3: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3,4-dihydro-2H-pyran-6-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1699)



N-[(2-amino-3-pyridyl)sulfonyl]-6-chloro-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (32 mg, 0.08 mmol), 2-(3,4-dihydro-2H-pyran-6-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (19 mg, 0.09 mmol), and 1,3-bis(2,6-di*iso*propylphenyl)-4,5-dihydroimidazole; 3-chloropyridine; dichloropalladium (3 mg, 0.005 mmol, 5 mol%) was

dissolved in EtOH (600 μL) and solution of potassium carbonate (150 μL, 2M) was added. The reaction mixture was heated at 120°C for 16 hours under an atmosphere of nitrogen. The reaction mixture was cooled, filtered and purified via LC/MS utilizing a gradient of 25-75% acetonitrile in 5 mM aq HCl. Sodium carbonate was added to collected fractions (pH 7-8) and the solvents were removed to dryness to yield the desired compound. ESI-MS m/z 472.3 $(M+1)^+$; Retention time: 1.18 minutes

Step 4: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-tetrahydropyran-2-yl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2351)



To *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(3,4-dihydro-2H-pyran-6-yl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1699) (37 mg, 0.08 mmol) in EtOH (3 mL), palladium on carbon (8 mg, 0.008 mmol, 10w/w%) was added. The reaction mixture was stirred under an atmosphere of hydrogen 16 hours at ambient temperature. The reaction mixture was filtered and evaporated to dryness, dissolved in DMSO-methanol and purified via reverse phase HPLC-MS method using a dual gradient run from 1-99% mobile phase B over 15.0 minutes. Mobile phase A = H_2O (5 mM TFA) Mobile phase B = acetonitrile to yield *N*-[(2-amino-3-pyridyl)sulfonyl]-6-tetrahydropyran-2-yl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2351). ESI-MS m/z 474.4$(M+1)^+$; Retention time: 1.15 minutes.

The following compounds can be synthesized using the procedures described herein:

N-[(2-amino-3-pyridyl)sulfonyl]-6-(1,2-dimethylpropyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1911)

N-[(2-amino-3-pyridyl)sulfonyl]-6-(5,5-dimethyl-2-oxo-cyclohexyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1793)

N-[(2-amino-3-pyridyl)sulfonyl]-6-norboman-2-yl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2097)

N-[(2-amino-3-pyridyl)sulfonyl]-6-tetrahydropyran-4-yl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1850)

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-cyclopropylethyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2553)

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-hydroxy-5,5-dimethyl-cyclohexyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2047)

N-(benzenesulfonyl)-6-tetrahydropyran-2-yl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1868)

N-(benzenesulfonyl)-6-*iso*butyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2246)

N-(benzenesulfonyl)-6-tetrahydrofuran-3-yl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2277)


N-[(2-amino-3-pyridyl)sulfonyl]-6-tetrahydropyran-3-yl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2311)

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-bicyclo[2.2.1]hept-2-enyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2709)

N-[(2-amino-3-pyridyl)sulfonyl]-6-[4-(trifluoromethyl)cyclohexyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2077) (Isomer 1, cis or trans stereoisomer)

N-[(2-amino-3-pyridyl)sulfonyl]-6-[4-(trifluoromethyl)cyclohexyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2432) (Isomer 2, cis or trans stereoisomer).


PREPARATION 126: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-tetrahydropyran-2-yl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2351) (Isomer 1, *RS* or *SS* stereoisomer) and *N*-[(2-amino-3-pyridyl)sulfonyl]-6-tetrahydropyran-2-yl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2351) (Isomer 2, *RS* or *SS* stereoisomer)



The stereoisomers were separated using supercritical fluid chromatography on a Phenomenex LUX-4 (250 x 10 mm), 5µm column using 48 % EtOH/IPA (50/50) in CO_2 at a flow rate of 10.0 mL/min. The separated enantiomers were separated concentrated to dryness to give the pure compounds. SFC Peak 1: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-tetrahydropyran-2-yl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2351) ESI-MS m/z 474.4(M+1)$^+$; Retention time: 1.15 minutes. SFC Peak 2: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-tetrahydropyran-2-yl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2351) ESI-MS m/z 474.4(M+1)$^+$; Retention time: 1.15 minutes.

The following compounds can be synthesized using the procedures described herein: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[2-tetrahydropyran-2-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2672) (Isomer 1, *RS* or *SS* stereoisomer) and *N*-[(2-amino-3-pyridyl)sulfonyl]-6-[2-tetrahydropyran-2-yl]-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2675) (Isomer 2, *RS* or *SS* stereoisomer).


PREPARATION 127: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(3-cyclopropyl-2,2-dimethyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2577) (Compound 2022) (Isomer 1, *R* or *S* stereoisomer) and *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(3-cyclopropyl-2,2-dimethyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2577) (Compound 2022) (Isomer 2, *R* or *S* stereoisomer)



The stereoisomers were separated using supercritical fluid chromatography on a ChiralPak AD-H (250 x 10 mm), 5µm column using 22% EtOH/MeOH (90:10, 20 mM NH_3) in CO_2 at a flow rate of 10.0 mL/min. The separated enantiomers were separated concentrated, diluted with ethyl acetate (3 mL) and washed with 1N aqueous hydrochloric acid. The organic layers were dried over sodium sulfate, filtered, and evaporated to dryness to give the pure compounds. SFC Peak 1: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(3-cyclopropyl-2,2-dimethyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2577) (Compound 2022) 1H NMR (400 MHz, DMSO) δ 12.35 (s, 1H), 7.59 (d, J = 7.7 Hz, 2H), 7.14 (d, J = 6.7 Hz, 1H), 6.73

- 6.58 (m, 2H), 6.43 (s, 2H), 2.79 (s, 1H), 2.62 (s, 1H), 1.80 - 1.59 (m, 2H), 1.56 (s, 3H), 1.51 (s, 3H), 1.27 (s, 9H), 1.13 - 1.03 (m, 1H), 0.71 - 0.58 (m, 1H), 0.56 - 0.46 (m, 1H), 0.46 - 0.33 (m, 1H), 0.24 (td, J = 9.2, 5.0 Hz, 1H), 0.12 - 0.01 (m, 1H) . SFC Peak 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(3-cyclopropyl-2,2-dimethyl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2577) (Compound 2022) 1H NMR (400 MHz, DMSO) δ 12.35 (s, 1H), 7.58 (d, J = 7.1 Hz, 2H), 7.14 (d, J = 7.1 Hz, 1H), 6.77 - 6.56 (m, 2H), 6.42 (s, 2H), 2.79 (s, 1H), 2.66 (dd, J = 12.3, 10.5 Hz, 1H), 1.69 (d, J = 20.9 Hz, 2H), 1.56 (s, 3H), 1.51 (s, 3H), 1.27 (s, 9H), 1.17 - 1.05 (m, 1H), 0.73 - 0.63 (m, 1H), 0.58 - 0.47 (m, 1H), 0.40 (dd, J = 8.2, 3.9 Hz, 1H), 0.24 (dd, J = 9.0, 4.1 Hz, 1H), 0.07 (dd, J = 9.0, 4.0 Hz, 1H).

PREPARATION 128: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,2-dimethyl-3-tetrahydropyran-4-yl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2425) (Compound 2266) (Isomer 1, *R* or *S* stereoisomer) and *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,2-dimethyl-3-tetrahydropyran-4-yl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2425) (Compound 2266) (Isomer 2, *R* or *S* stereoisomer)




The stereoisomers were separated using supercritical fluid chromatography on a ChiralPak AD-H (250 x 10 mm), 5μm column using 12% MeOH in CO_2 at a flow rate of 10.0 mL/min. The separated enantiomers were separately concentrated, diluted with ethyl acetate (3 mL) and washed with 1N aqueous hydrochloric acid. The organic layers were dried over sodium sulfate, filtered, and evaporated to dryness to give the pure compounds. SFC Peak 1: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,2-dimethyl-3-tetrahydropyran-4-yl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2425) (Compound 2266) ESI-MS *m/z* calc. 515.25665, found 516.2 $(M+1)^+$; Retention time: 1.31 minutes. SFC Peak 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-(2,2-dimethyl-3-tetrahydropyran-4-yl-pyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2425) (Compound 2266) ESI-MS *m/z* calc. 515.25665, found 516.2 $(M+1)^+$; Retention time: 1.31 minutes . 1H NMR (400 MHz, DMSO) δ 12.48 (s, 1H), 7.70 -

7.52 (m, 2H), 7.17 (dd, J = 7.3, 0.5 Hz, 1H), 6.70 (dd, J = 9.7, 8.5 Hz, 2H), 5.19 (s, 2H), 3.84 (td, J = 11.1, 3.7 Hz, 2H), 3.38 - 3.18 (m, 2H), 2.84 (t, J = 10.0 Hz, 1H), 2.68 - 2.55 (m, 1H), 2.07 (s, 4H), 1.81 (t, J = 6.4 Hz, 1H), 1.65 (dd, J = 30.6, 10.5 Hz, 2H), 1.55 (d, J = 8.6 Hz, 6H), 1.49 (s, 3H), 1.28 (d, J = 6.6 Hz, 10H).


PREPARATION 129: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(3,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2532) (Isomer 1) (Compound 2178) (Isomer 2) (Compound 2062) (Isomer 3) (Compound 1863) (Isomer 4)

Step 1: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide




2-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxylic acid (3.91 g, 9.66 mmol) was dissolved in DMF (31mL) was stirred under nitrogen and CDI (2.35 g, 14.5 mmol) was added in one portion. The reaction mixture was warmed to 65 °C and heated for 1 h. Meanwhile in a separate vessel, 6-aminopyridine-2-sulfonamide (2.09 g, 12.1 mmol) was slurried in DMF (15 mL) under nitrogen and NaH (483 mg, 12.1 mmol) added portionwise to mitigate gas evolution and the reaction was stirred for 1 h. The two reactions were combined in one portion at 65 °C. The reaction was stirred for 15 min and evaporated. The reaction mixture was poured over ice and acidified (ph=4) by the addition of 6M HCl. The resulting mixture was extracted with ethyl acetate (1 L). The organics were washed with brine (2 x 500 mL), dried over sodium sulfate and evaporated under reduced pressure. The crude material was purified by silica gel column chromatography eluting with 0-100% ethyl acetate in hexanes to give *N*-[(6-amino-2-pyridyl)sulfonyl]-2-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (4.02 g, 887%) ESI-MS m/z calc. 478.08777, found 479.1 $(M+1)^+$; Retention time: 0.7 minutes.

Step2: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(3,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2532) (Compound 2178) (Compound 2062) (Compound 1863)



N-[(6-amino-2-pyridyl)sulfonyl]-2-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (100 mg, 0.21 mmol) and fluorocesium (156 mg, 38.5 µL, 1.04 mmol) in DMSO (600 µL) followed by K_2CO_3 (289 mg, 2.09 mmol) and 3,4-dimethylpiperidine (118 mg, 1.04 mmol). The reaction mixture was sealed and heated to 150°C overnight, then 165°C for 6 hours. The reaction mixture was filtered and then purified directly by reverse-phase HPLC-MS using10-99% Acetonitrile-Water + 5 mmolar HCl as eluent to afford *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(3,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2532) (Compound 2178) (Compound 2062) (Compound 1863) (86 mg, 74%), as a light yellow solid ESI-MS m/z calc. 555.23157, found 556.2 $(M+1)^+$; Retention time: 2.0 minutes. 1H NMR (400 MHz, DMSO) δ 12.66 (s, 1H), 7.81 (d, J = 7.9 Hz, 1H), 7.65 (t, J = 7.8 Hz, 1H), 7.50 - 7.41 (m, 3H), 7.22 (d, J = 7.3 Hz, 1H), 6.91 (d, J = 10.9 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 6.47 (s, 1H), 3.85 (d, J = 6.5 Hz, 2H), 3.51 (d, J = 8.5 Hz, 1H), 3.40 (d, J = 12.8 Hz, 1H), 3.13 (d, J = 11.8 Hz, 1H), 2.97 (d, J = 6.0 Hz, 1H), 2.06 (d, J = 9.3 Hz, 1H), 1.89 - 1.72 (m, 2H), 1.42 (s, 2H), 1.00 (d, J = 6.7 Hz, 6H), 0.87 (d, J = 6.6 Hz, 3H), 0.77 (d, J = 6.6 Hz, 3H

Step 3: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(3,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2532) (Compound 2178) (Compound 2062) (Compound 1863) (Isomers 1-4, *RR*, *RS*, *SS*, or *SR* stereoisomer)



The stereoisomers were separated using supercritical fluid chromatography on a ChiralCel OZ-H (250 x 10 mm), 5µm column using 25% MeOH:EtOH 50:50 in CO_2 at a flow

rate of 10.0 mL/min. The separated enantiomers were separately concentrated to give the pure compound. SFC Peak 1: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(3,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2532) (Compound 2178) (Compound 2062) (Compound 1863) ESI-MS m/z calc. 555.23157, found 556.2 $(M+1)^+$; Retention time: 2.02 minutes. SFC Peak 2 (*N*-[(6-amino-2-pyridyl)sulfonyl]-2-(3,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2532) (Compound 2178) (Compound 2062) (Compound 1863) ESI-MS m/z calc. 555.23157, found 556.2 $(M+1)^+$; Retention time: 2.01 minutes. SFC Peak 3 *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(3,4-dimethyl-1-piperidyl)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2532) (Compound 2178) (Compound 2062) (Compound 1863) ESI-MS m/z calc. 555.23157, found 556.2 $(M+1)^+$; Retention time: 1.96 minutes. SFC Peak 4 ESI-MS m/z calc. 555.23157, found 556.2 $(M+1)^+$; Retention time: 1.96 minutes. 1H NMR (400 MHz, DMSO) δ 12.66 (s, 1H), 7.79 (d, J = 7.8 Hz, 1H), 7.63 (t, J = 10.7 Hz, 1H), 7.45 (d, J = 11.6 Hz, 3H), 7.20 (d, J = 6.5 Hz, 1H), 6.90 (d, J = 10.7 Hz, 1H), 6.70 (d, J = 7.1 Hz, 1H), 6.44 (s, 2H), 3.84 (d, J = 6.5 Hz, 2H), 3.50 (s, 1H), 3.38 (s, 1H), 3.12 (d, J = 13.0 Hz, 1H), 2.96 (s, 1H), 2.08 - 1.98 (m, 1H), 1.80 (s, 2H), 1.41 - 1.37 (m, 1H), 1.30 - 1.22 (m, 1H), 1.00 (d, J = 6.7 Hz, 7H), 0.86 (d, J = 6.5 Hz, 4H), 0.76 (d, J = 6.6 Hz, 3H).

The following compounds can be synthesized using the procedures described herein:


N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,5-dimethyl-1-piperidyl)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2451) (Compound 1993) (Compound 1864) (Compound 1855) (Isomers 1-4, stereoisomers *RR*, *RS*, *SS*, *SR*), using as SFC ChiralCel OZ-H (250 x 10 mm), 5μm column and eluent 25% MeOH (20 mM NH_3) in CO2

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(6-methyl-8-azaspiro[3.4]octa*N*-8-yl)pyridine-3-carboxamide (Compound 2552) (Isomer 1, *R* or *S* stereoisomer) (Compound 2269) (Isomer 2, *R* or *S* stereoisomer), using as SFC ChiralPak IA (250 x 10 mm), 5μm column and eluent 22% EtOH (0.1% TFA) in CO2

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2-ethyl-2,4-dimethyl-pyrrolidin-1-yl)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2213) (Compound 2058) (Compound 1817) (Compound 1729) (Isomers 1-4, stereoisomers *RR*, *RS*, *SS*, *SR*), using as SFC ChiralCel OZ-H (250 x 10 mm), 5μm column and eluent 22% IPA:EtOH 50:50 in CO_2


N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-4-(trifluoromethyl)pyrrolidin-1-yl)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2271) (Isomer 1, *R* or *S* stereoisomer) using as SFC ChiralCel OZ-H (250 x 10 mm), 5μm column and eluent 25 % MeOH:EtOH (50:50) in CO_2.

PREPARATION 130: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-5-iodo-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2009)



N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (100 mg, 0.18 mmol) was dissolved in CH_2Cl_2 (700 μL) and was treated with 1-iodopyrrolidine-2,5-dione (44 mg, 0.19 mmol). The reaction mixture was allowed to stir at 40 °C for 30 minutes. The volatiles were evaporated under a slow stream of nitrogen gas. The remaining residue was redissolved in DMSO, filtered and purified by reverse-phase HPLC using a Luna C18 (2) column (50 x 21.2 mm, 5 μm particle size) and a dual gradient run from 30-99% mobile phase B over 15.0 minutes (mobile phase A = water (de-ionized, no acid modifier), mobile phase B = acetonitrile) to give the desired compound. ESI-MS m/z calc. 681.1282, found 682.2 $(M+1)^+$; Retention time: 2.26 minutes.


PREPARATION 131: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-methyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2258)



N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (500 mg, 0.90 mmol) in DMF (5 mL) was treated with Cs_2CO_3 (352 mg, 1.1 mmol) and the reaction mixture was stirred for 0.5 h at room temperature, then cooled in an ice bath and treated with MeI (141 mg, 61 µL, 1.0 mmol). The reaction mixture was stirred overnight at room temperature. The suspension was added to water (25ml), stirred at room temperature for 1 h and the suspension was filtered and washed with water. The solid was purified by silica gel column chromatography using as dichloromethane as eluent to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-methyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2258) (315 mg, 61%). ESI-MS m/z 570.0 $(M+1)^+$; Retention time: 3.4 minutes ; 1H NMR (400 MHz, DMSO-d6) δ 7.67 - 7.54 (m, 2H), 7.49 (t, J = 1.8 Hz, 1H), 7.40 (dt, J = 10.2, 1.8 Hz, 1H), 7.23 (d, J = 8.0 Hz, 1H), 7.06 (d, J = 7.3 Hz, 1H), 6.89 (dt, J = 10.7, 2.3 Hz, 1H), 6.70 (d, J = 8.4 Hz, 1H), 6.64 (s, 2H), 3.84 (d, J = 6.7 Hz, 2H), 3.43 (s, 3H), 2.82 (s, 1H), 2.64 (s, 1H), 2.34 - 2.22 (m, 1H), 2.05 (hept, J = 6.6 Hz, 1H), 1.91 (dd, J = 11.9, 5.7 Hz, 1H), 1.62 (d, J = 4.4 Hz, 6H), 1.46 (t, J = 12.1 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.94 (d, J = 6.3 Hz, 3H)


PREPARATION 132: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-[3-fluoro-5-(2-hydroxy-2-methyl-propoxy)phenyl]pyridine-3-carboxamide (Compound 1890)

Step 1: 1-[3-fluoro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy]-2-methyl-propan-2-ol



To a suspension of 1-(3-bromo-5-fluoro-phenoxy)-2-methyl-propan-2-ol (25.0 g, 95 mmol), Bis(pinacolato)diboron (26.5 g, 105 mmol), and KOAc (18.7 g, 190 mmol) in dioxane (375 mL) under an atmosphere of nitrogen was added $Pd(dppf)Cl_2$.DCM (768 mg, 1.0 mmol) and the reaction mixture was heated to 103 °C for 2 hours, cooled and stirred overnight at room temperature. The reaction mixture was filtered through Florisil and the filtrate was concentrated under reduced pressure to afford a yellow oil. The oil was purified by silica gel column chromatography eluted with EtOAc/heptane (10 - 20% EtOAc) to afford 1-[3-fluoro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy]-2-methyl-propan-2-ol (12.6 g (86%) as pale yellow liquid.

Step 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-[3-fluoro-5-(2-hydroxy-2-methyl-propoxy)phenyl]pyridine-3-carboxamide (Compound 1890)



N-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-2-(2,6-dimethylphenoxy)pyridine-3-carboxamide (140 mg, 0.32 mmol), 1-[3-fluoro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy]-2-methyl-propan-2-ol (215 mg, 0.69 mmol), potassium carbonate (647 µL of 2 M, 1.29 mmol), and 1,3-bis(2,6-di*iso*propylphenyl)-4,5-dihydroimidazole; 3-chloropyridine; dichloropalladium (PEPPSI-IPr) (11 mg, 0.02 mmol) were combined in ethanol (3 mL) and the reaction mixture was heated at 80 °C for 21 h. The reaction mixture was cooled to 40 °C, glacial AcOH (0.20 mL) was added, filtered and the filtrate evaporated under reduced pressure. The residue was dissolved in DMSO (3 mL), filtered and purified using a reverse phase HPLC-MS method using a Luna C18 (2) column (75 × 30 mm, 5 µm particle size) and a dual gradient run


from 10-99% mobile phase B over 30.0 minutes (mobile phase A = H_2O (5 mM HCl), mobile phase B = CH_3CN) to give *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-[3-fluoro-5-(2-hydroxy-2-methyl-propoxy)phenyl]pyridine-3-carboxamide (Compound 1890) (HCl salt) (80 mg, 40 %) 1H NMR (400 MHz, DMSO-d6) δ 12.39 (s, 1H), 8.16 (d, J = 7.8 Hz, 1H), 7.84 (d, J = 7.9 Hz, 1H), 7.66 (t, J = 7.9 Hz, 1H), 7.24 (d, J = 7.3 Hz, 1H), 7.21 - 7.14 (m, 4H), 7.14 - 7.07 (m, 1H), 6.84 (dt, J = 10.6, 2.3 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 6.54 (s, 2H), 4.61 (s, 1H), 3.61 (s, 2H), 2.07 (s, 1H), 2.06 (s, 6H), 1.19 (s, 6H). ESI-MS m/z calc. 580.1792, found 581.2 $(M+1)^+$; Retention time: 1.76 minutes.

The following compound can be synthesized using the procedures described herein:

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,6-dimethylphenoxy)-6-[3-fluoro-5-(3-hydroxy-2-methyl-propoxy)phenyl]pyridine-3-carboxamide (Compound 1797)


PREPARATION 133: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,5,5-trimethylcyclopenten-1-yl)pyridine-3-carboxamide (Compound 2081) (Isomer 1, *R* or *S* stereoisomer) (Compound 2039) (Isomer 2, *R* or *S* stereoisomer)

Step 1: Ethyl 2-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxylate




To a solution of 2-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxylic acid (1.3 g, 4.0 mmol) in ethanol (40 mL) was slowly added thionyl chloride (1.0 g, 584 µL, 8.0 mmol) and the reaction mixture was stirred at reflux for six hours. The solvent was evaporated under reduced pressure to give ethyl 2-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxylate (1.3 g, 95%) as a colorless solid.

Step 2: Ethyl 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,5,5-trimethylcyclopenten-1-yl)pyridine-3-carboxylate



A mixture of ethyl 2-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxylate (700 mg, 1.99 mmol), 4,4,5,5-tetramethyl-2-(3,5,5-trimethylcyclopenten-1-yl)-1,3,2-dioxaborolane (517 mg, 2.19 mmol), 1,1'-bis(diphenylphosphino)ferrocene-palladium dichloride (437 mg, 0.60 mmol), and Na_2CO_3 (633 mg, 5.97 mmol) in dioxane (9 mL) and water (1 mL) was heated in a sealed vessel at 150 °C for 16 hours. The reaction mixture was filtered, and the solids were washed with EtOAc. The combined organics were evaporated, and the residue was purified by silica gel column chromatography using a gradient of 100% hexanes to 20% EtOAc in hexanes to afford ethyl 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,5,5-trimethylcyclopenten-1-yl)pyridine-3-carboxylate (97 mg, 11%) as a white solid. ESI-MS m/z calc. 425.23663, found 426.2 $(M+1)^+$; Retention time: 2.4 minutes.


Step 3: 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,5,5-trimethylcyclopenten-1-yl)pyridine-3-carboxylic acid



To a solution of ethyl 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,5,5-trimethylcyclopenten-1-yl)pyridine-3-carboxylate (92 mg, 0.2 mmol) in THF (520 µL) was added NaOH (650 µL of 1 M, 0.6 mmol) and then MeOH (520 µL) and the reaction mixture was heated to 50°C for 4 hrs. The reaction mixture was neutralized with 1 M HCl (3.0 mL) and extracted with EtOAc (10 mL x 2). The combined extracts were dried over Na_2SO_4 and evaporated under reduced pressure to give 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,5,5-trimethylcyclopenten-1-yl)pyridine-3-carboxylic acid (77 mg, 90%) as a light yellow solid. ESI-MS m/z calc. 397.20532, found 398.2 $(M+1)^+$; Retention time: 1.92 minutes; 1H NMR (400


MHz, DMSO) δ 13.11 (s, 1H), 7.98 (s, 2H), 7.56 (s, 1H), 7.49 (d, J = 9.9 Hz, 1H), 6.93 (d, J = 10.7 Hz, 1H), 5.67 (s, 1H), 3.85 (d, J = 6.6 Hz, 2H), 2.87 (dd, J = 14.1, 7.0 Hz, 1H), 2.13 - 1.98 (m, 2H), 1.46 - 1.38 (m, 4H), 1.24 (s, 4H), 1.08 (d, J = 6.9 Hz, 3H), 0.99 (d, J = 6.7 Hz, 6H).

Step 4: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,5,5-trimethylcyclopenten-1-yl)pyridine-3-carboxamide (Compound 2081) (Compound 2039)



6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,5,5-trimethylcyclopenten-1-yl)pyridine-3-carboxylic acid (74 mg, 0.19 mmol) and CDI (45 mg, 0.28 mmol) were mixed in DMF (740 µL) under and atmosphere of nitrogen and the reaction mixture was heated to 45 °C for 45 minutes. In a separate vial, 6-aminopyridine-2-sulfonamide (80.62 mg, 0.4655 mmol) and NaH (18 mg, 0.46 mmol) were mixed slowly in DMF (370.0 µL) and mixture was heated to 45 °C for 45minutes. The 6-aminopyridine-2-sulfonamide reaction mixture was added to the activated acid and the reaction mixture was heated at 50°C for 16h. The reaction mixture was filtered and then purified directly by reverse-phase preparative chromatography utilizing a C18 column and HPLC-MS method 10-99% mobile phase B (mobile phase A = water (de-ionized, no acid modifier), mobile phase B = acetonitrile) to afford *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,5,5-trimethylcyclopenten-1-yl)pyridine-3-carboxamide (Compound 2081) (Compound 2039) (64 mg, 62%) as an off-white solid, ESI-MS m/z calc. 552.22064, found 553.4 $(M+1)^+$; Retention time: 2.25 minutes; 1H NMR (400 MHz, DMSO) δ 12.55 (s, 1H), 7.96 (d, J = 8.2 Hz, 1H), 7.86 (d, J = 8.2 Hz, 1H), 7.68 - 7.60 (m, 1H), 7.56 (s, 1H), 7.49 (d, J = 9.6 Hz, 1H), 7.20 (d, J = 7.3 Hz, 1H), 7.01 - 6.84 (m, 1H), 6.72 (d, J = 8.4 Hz, 1H), 6.55 (s, 2H), 5.62 (d, J = 1.7 Hz, 1H), 3.85 (d, J = 6.6 Hz, 2H), 2.66 (ddd, J = 14.3, 7.4, 1.8 Hz, 1H), 2.03 (dd, J = 13.2, 6.6 Hz, 1H), 1.95 (dd, J = 12.1, 7.5 Hz, 1H), 1.38 - 1.28 (m, 4H), 1.25 (s, 3H), 0.99 (d, J = 6.7 Hz, 6H), 0.93 (d, J = 6.9 Hz, 3H).


Step 5: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,5,5-trimethylcyclopenten-1-yl)pyridine-3-carboxamide (Compound 2081) (Isomer 1, *R* or *S* stereoisomer) and *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,5,5-trimethylcyclopenten-1-yl)pyridine-3-carboxamide (Compound 2039) (Isomer 2, *R* or *S* stereoisomer)



The stereoisomers were separated using supercritical fluid chromatography on a ChiralCel AD-H (250 x 10 mm), 5μm column using 20% EtOH (0.1% TFA) in CO_2 at a flow rate of 10.0 mL/min. The separated enantiomers were separately concentrated to give the pure compound. SFC Peak 1: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,5,5-trimethylcyclopenten-1-yl)pyridine-3-carboxamide (Compound 2081) (Isomer 1, *R* or *S* stereoisomer) ESI-MS m/z calc. 552.22064, found 553.2 $(M+1)^+$; Retention time: 2.26 minutes. SFC Peak 2 *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,5,5-trimethylcyclopenten-1-yl)pyridine-3-carboxamide (Compound 2039) (Isomer 2, *R* or *S* stereoisomer) ESI-MS m/z calc. 552.22064, found 553.2 $(M+1)^+$; Retention time: 2.26 minutes.


PREPARATION 134: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxamide (Compound 2492) (Isomer 1, *R* or *S* stereoisomer) (Compound 2374) (Isomer 2, *R* or *S* stereoisomer)

Step 1: 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxylic acid



A mixture of 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3,3,5,5-tetramethylcyclopenten-1-yl)pyridine-3-carboxylic acid (389 mg, 0.94 mmol) (prepared in a similar manner to that described in preparation 133), palladium hydroxide (80 mg, 0.57 mmol) and ammonium formate (1.19 g, 18.91 mmol) in ethanol (10 mL) were combined in a microwave vial and heated to 85°C for 3h. The reaction mixture was cooled to room temperature and a further aliquot of ammonium formate (1.19 g, 18.91 mmol) and palladium hydroxide (80 mg, 0.57 mmol) was added, and irradiated under microwave conditions at 85°C for 4h. The reaction mixture was filtered over celite, washed with methanol and the filtrate concentrated under reduced pressure, then diluted with 1N HCl and washed with ethyl acetate (2X). The combined organic layers were dried over sodium sulfate, filtered and concentrated under reduced pressure to a clear oil. The oil was purified by silica gel column chromatography using a gradient from 100% dichloromethane to 20% methanol in dichloromethane followed by 20% methanol in ethyl acetate to give 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxylic acid (52 mg, 13%) as a white solid.


Step 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxamide (Compound 2492) (Compound 2374)



A solution of 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(2,2,4,4-

tetramethylcyclopentyl)pyridine-3-carboxylic acid (52 mg, 0.13 mmol) and carbonyl diimidazole (31 mg, 0.19 mmol) in dimethylformamide (0.5 mL) was stirred at 45°C for 90 minutes. A separate solution of 6-aminopyridine-2-sulfonamide (44 mg, 0.25 mmol) and sodium hydride (10 mg, 0.25 mmol) in dimethylformamide (0.5 mL) was stirred at 45 °C for 45 minutes was then added to the activated acid and the resulting mixture was stirred at 45 °C for 2 h. The reaction mixture was filtered and purified using a reverse phase HPLC-MS method using a Luna C18 (2) column (75 x 30 mm, 5 µm particle size) and a dual gradient run from 1-99% mobile phase B over 15.0 minutes (mobile phase A = H_2O (5 mM HCl), mobile phase B = acetonitrile) to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxamide (Compound 2492) (Compound 2374) (HCl salt) (51 mg, 67%) as an off-white solid. 1H NMR (400 MHz, DMSO) δ 12.63 (s, 1H), 7.91 (dd, J = 20.4, 8.2 Hz, 2H), 7.66 (t, J = 7.9 Hz, 1H), 7.57 (s, 1H), 7.51 (d, J = 9.8 Hz, 1H), 7.25 (d, J = 7.3 Hz, 1H), 6.93 (d, J = 10.7 Hz, 1H), 6.73 (d, J = 8.4 Hz, 1H), 6.51 (s, 2H), 3.86 (d, J = 6.6 Hz, 2H), 3.58 (dd, J = 12.2, 6.5 Hz, 1H), 2.55 (d, J = 12.3 Hz, 1H), 2.06 (dt, J = 13.2, 6.6 Hz, 1H), 1.63 (dd, J = 12.4, 6.4 Hz, 1H), 1.50 (d, J = 13.0 Hz, 1H), 1.39 (d, J = 12.9 Hz, 1H), 1.15 (s, 3H), 1.08 - 0.93 (m, 9H), 0.80 (s, 3H), 0.69 (s, 3H). ESI-MS m/z calc. 568.25195, found 569.3 $(M+1)^+$; Retention time: 2.4 minutes.

Step 3: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxamide (Compound 2492) (Isomer 1, *R* or *S* stereoisomer) and *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxamide (Compound 2374) (Isomer 2, *R* or *S* stereoisomer)




The stereoisomers were separated using supercritical fluid chromatography on a

Chiralpak IC (250 x 10mm), 5μm column using 17 % EtOH (DeNat) + 0.1% TFA in CO_2 at a flow rate of 10.0 mL/min. The separated enantiomers were separately concentrated to give the pure compound. SFC Peak 1: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxamide (Compound 2492) (Compound 2374) ESI-MS m/z calc. 568.25195, found 569.3 $(M+1)^+$; Retention time: 2.4 minutes. 1H NMR (400 MHz, MeOD) δ 7.87 (d, J = 8.2 Hz, 1H), 7.76 (d, J = 8.2 Hz, 1H), 7.70 (dd, J = 8.3, 7.4 Hz, 1H), 7.54 (s, 1H), 7.46 - 7.38 (m, 2H), 6.83 (d, J = 8.4 Hz, 1H), 6.75 (dt, J = 10.5, 2.3 Hz, 1H), 3.82 (d, J = 6.5 Hz, 2H), 3.67 (dd, J = 12.3, 6.4 Hz, 1H), 2.66 (t, J = 12.5 Hz, 1H), 2.11 (dt, J = 13.3, 6.7 Hz, 1H), 1.69 (dd, J = 12.6, 6.3 Hz, 1H), 1.49 (dd, J = 38.8, 13.1 Hz, 2H), 1.18 (d, J = 5.6 Hz, 3H), 1.10 - 1.00 (m, 9H), 0.81 (s, 3H), 0.75 (s, 3H). SFC Peak 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(2,2,4,4-tetramethylcyclopentyl)pyridine-3-carboxamide (Compound 2492) (Compound 2374). ESI-MS m/z calc. 568.25195, found 569.3 $(M+1)^+$; Retention time: 2.4 minutes. 1H NMR (400 MHz, MeOD) δ 7.87 (d, J = 8.2 Hz, 1H), 7.76 (d, J = 8.2 Hz, 1H), 7.70 (dd, J = 8.3, 7.4 Hz, 1H), 7.54 (s, 1H), 7.46 - 7.38 (m, 2H), 6.83 (d, J = 8.4 Hz, 1H), 6.75 (dt, J = 10.5, 2.3 Hz, 1H), 3.82 (d, J = 6.5 Hz, 2H), 3.67 (dd, J = 12.3, 6.4 Hz, 1H), 2.66 (t, J = 12.5 Hz, 1H), 2.11 (dt, J = 13.3, 6.7 Hz, 1H), 1.69 (dd, J = 12.6, 6.3 Hz, 1H), 1.49 (dd, J = 38.8, 13.1 Hz, 2H), 1.18 (d, J = 5.6 Hz, 3H), 1.10 - 1.00 (m, 9H), 0.81 (s, 3H), 0.75 (s, 3H).

PREPARATION 135: 6-(4-chloro-3-propoxy-phenyl)-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-*N*-[[6-(pent-4-ynoylamino)-2-pyridyl]sulfonyl]pyridine-3-carboxamide (Compound 2540)


Step 1: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-



N-[(6-amino-2-pyridyl)sulfonyl]-2,6-dichloro-pyridine-3-carboxamide (2.0 g, 5.8 mmol) and (2R,5S)-2,5-dimethylpyrrolidine (0.7 g, 7.2 mmol) in DMSO (10 mL) was treated


with K_2CO_3 (2.4 g, 17 mmol) and heated at 120 °C under stirring for 3h. The reaction mixture was cooled down and carefully added to cold mixture of HCl (2.881 mL of 12 M, 35 mmol) and water (50 mL) forming a white foam. The reaction mixture was allowed to stir at room temperature for 30 min, filtered and the white solid was triturated with IPA (100mL) to give the desired compound.

Step 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-propoxy-phenyl)-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 773)



N-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-2-[(2S,5R)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (62 mg, 0.15 mmol), (4-chloro-3-propoxy-phenyl)boronic acid (48 mg, 0.23 mmol), tetrakis(triphenylphosphine)palladium (0) (10 mg, 0.009 mmol), potassium carbonate (300 µL, 2 M, 0.40 mmol) and DMF (1.7 mL) were heated at 100 °C overnight. The reaction mixture was filtered and purified by reverse phase HPLC utilizing a gradient of 1-99% acetonitrile in 5 mM aq HCl. ESI-MS m/z 544.3 $(M+1)^+$; Retention time: 1.7 minutes; 1H NMR (400 MHz, DMSO-d6) δ 12.98 (s, 1H), 7.83 (d, J = 1.9 Hz, 1H), 7.81 (d, J = 8.0 Hz, 1H), 7.70 - 7.61 (m, 2H), 7.52 (d, J = 8.3 Hz, 1H), 7.37 (d, J = 7.9 Hz, 1H), 7.23 (dd, J = 7.3, 0.8 Hz, 1H), 6.72 (d, 1H), 6.48 (s, 2H), 4.13 (t, J = 6.5 Hz, 2H), 4.09 - 3.97 (m, 2H), 2.05 - 1.91 (m, 2H), 1.81 (h, J = 7.2 Hz, 2H), 1.74 - 1.61 (m, 2H), 1.10 (d, J = 6.2 Hz, 6H), 1.03 (t, J = 7.4 Hz, 3H).


Step 3: 6-(4-chloro-3-propoxy-phenyl)-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]-*N*-[[6-(pent-4-ynoylamino)-2-pyridyl]sulfonyl]pyridine-3-carboxamide (Compound 2540)



N-[(6-amino-2-pyridyl)sulfonyl]-6-(4-chloro-3-propoxy-phenyl)-2-[(2R,5S)-2,5-dimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 773) (10 mg, 0.02 mmol) in DCM (2 mL) at 0 °C was added triethylamine (11 mg, 15 µL, 0.11 mmol) and pent-4-ynoyl chloride (4 mg, 0.04 mmol). The reaction mixture was then stirred at 0 °C for 5 minutes, then at rt for 2 h. The reaction mixture was evaporated under reduced pressure and the crude mixture was purified by reverse phase HPLC using 10-100% ACN in Water (TFA modifier). The salt was neutralized with SPE-CO_3H cartridge, eluted with MeOH/DCM to give the desired compound. ESI-MS m/z 624.3 $(M+1)^+$; Retention time: 0.9 minutes; 1H NMR (300 MHz, $CDCl_3$) δ 8.73 (d, J = 8.2 Hz, 1H), 8.28 (d, J = 8.1 Hz, 1H), 8.04 (t, J = 7.9 Hz, 1H), 7.98 - 7.89 (m, 2H), 7.56 (dd, J = 8.2, 1.9 Hz, 1H), 7.51 (d, J = 2.4 Hz, 2H), 4.21 (d, J = 6.1 Hz, 2H), 4.10 (t, J = 6.4 Hz, 2H), 2.73 (t, J = 6.9 Hz, 2H), 2.64 - 2.47 (m, 4H), 2.11 (d, J = 2.8 Hz, 2H), 2.00 - 1.85 (m, 4H), 1.34 (d, J = 6.6 Hz, 6H), 1.25 (s, 6H), 1.11 (t, J = 7.4 Hz, 5H), 0.93 - 0.79 (m, 3H).

PREPARATION 136: *N*-[4-[3-(4-chlorophenyl)propyl]piperazin-1-yl]sulfonyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2141)


Step 1: 2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxylic acid



A mixture of 2-fluoropyridine-3-carboxylic acid (71 mg, 0.5 mmol), (4*S*)-2,2,4-trimethylpyrrolidine (HCl salt) (150 mg, 1.0 mmol), potassium carbonate (207 mg, 1.5 mmol), and cesium fluoride (152 mg, 1.0 mmol) in DMSO (500 µL) was stirred at 130 °C for six hours. The reaction mixture was diluted with water, adjusted to pH 4 with 1 M HCl, and extracted with ethyl acetate. The combined extracts were washed with brine, dried over sodium sulfate, and


evaporated under reduced pressure. The residue was purified using reverse phase HPLC-MS using a Luna C18 (2) column (75 × 30 mm, 5 μm particle size) and a dual gradient run from 1-70% mobile phase B over 15.0 minutes (mobile phase A = H_2O (5 mM HCl). mobile phase B = CH_3CN) to give the desired compound.

Step 2:benzyl 4-[[2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carbonyl]sulfamoyl]piperazine-1-carboxylate




A solution of (S)-2-(2,2,4-trimethylpyrrolidin-1-yl)nicotinic acid (117 mg, 0.5 mmol) and chlorosulfonyl *iso*cyanate (85 mg, 52 μL, 0.6 mmol) in dichloromethane (3 mL) was stirred for 30 minutes, and was slowly added to a solution of benzyl piperazine-1-carboxylate (110 mg, 96 μL, 0.5 mmol) and pyridine (59 mg, 60 μL, 0.8 mmol) in dichloromethane (2 mL). The reaction mixture was stirred for two hours, evaporated under reduced pressure and purified by silica gel column chromatography with 0-5% methanol in dichloromethane as eluent to give benzyl (S)-4-(*N*-(2-(2,2,4-trimethylpyrrolidin-1-yl)nicotinoyl)sulfamoyl)piperazine-1-carboxylate (35 mg, 13%) as a colorless oil. ESI-MS m/z 516.6 $(M+1)^+$; Retention time: 1.65 minutes.

Step 3: *N*-piperazin-1-ylsulfonyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide



A mixture of benzyl (S)-4-(*N*-(2-(2,2,4-trimethylpyrrolidin-1-yl)nicotinoyl)sulfamoyl)piperazine-1-carboxylate (0.22 g, 0.43 mmol) and palladium on carbon (20 mg, 0.02 mmol) in methanol (2 mL) was stirred under a hydrogen atmosphere for 16 hours. The reaction mixture was filtered and evaporated to give (S)-*N*-(piperazin-1-ylsulfonyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)nicotinamide (0.13 g, 80%) as a colorless solid.

Step 4: *N*-[4-[3-(4-chlorophenyl)propyl]piperazin-1-yl]sulfonyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2141)



A solution of (S)-*N*-(piperazin-1-ylsulfonyl)-2-(2,2,4-trimethylpyrrolidin-1-yl)nicotinamide (30 mg, 0.08 mmol) and 3-(4-chlorophenyl)propanal (14 mg, 0.08 mmol) in 1,2-dichloroethane (800 µL) was stirred for five minutes, and sodium triacetoxyborohydride (33 mg, 0.16 mmol) was added. The reaction mixture was stirred for 16 hours, diluted with methanol, filtered, and purified using a reverse phase HPLC-MS method using a Luna C18 (2) column (75 × 30 mm, 5 µm particle size) and a dual gradient run from 1-99% mobile phase B over 15.0 minutes (mobile phase A = H_2O (5 mM HCl), mobile phase B = CH_3CN) to give the desired compound. ESI-MS m/z 534.6 $(M+1)^+$; Retention time: 1.23 minutes.

The following compounds can be synthesized using the procedures described

herein:

N-[(3*R*)-3-[(4-*tert*-butoxyphenyl)methylamino]pyrrolidin-1-yl]sulfonyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2698),

N-[[3-[(4-*tert*-butoxyphenyl)methylamino]-1-piperidyl]sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2697),


N-[(3*R*)-3-[[4-fluoro-3-(trifluoromethoxy)phenyl]methylamino]pyrrolidin-1-yl]sulfonyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2681),

N-[(3*R*)-3-[3-(4-chlorophenyl)propylamino]pyrrolidin-1-yl]sulfonyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2717) and

N-[(3*R*)-3-[(2,4-dichlorophenyl)methylamino]pyrrolidin-1-yl]sulfonyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2693)


PREPARATION 137: 6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-(3-piperidylsulfonyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2655)

Step 1: 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxylic acid




A mixture of 2-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxylic acid (1.5 g, 3.6 mmol), (4*S*)-2,2,4-trimethylpyrrolidine (HCl salt) (0.8 g, 5.8 mmol), K_2CO_3 (1.6 g, 11.6 mmol), and CsF (1.1 g, 7.5 mmol) in DMSO (5 mL) was stirred at 130 °C for two days. The reaction mixture was diluted with water, adjusted to pH 4 with 1 M HCl, and extracted with ethyl acetate. The combined extracts were washed with brine, dried over sodium sulfate, and evaporated under reduced pressure. The residue was purified by silica gel chromatography with 0-6% methanol in dichloromethane to give 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxylic acid (1.2 g, 82%) as a tan foam. ESI-MS m/z calc. 400.21622, found 401.0 $(M+1)^+$; Retention time: 1.92 minutes (3 min).

Step 2: 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide



A solution of 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxylic acid (120 mg, 0.3 mmol) and CDI (73 mg, 0.5 mmol) in DMF (1.5 mL) was stirred at 45 °C for 30 minutes, and ammonium hydroxide (525 µL of 30 %w/v, 4.5 mmol) was added. The reaction mixture was stirred at 45 °C for three days, diluted with water, and extracted with ethyl acetate. The combined extracts were washed with brine, dried over sodium sulfate, and evaporated under reduced pressure. The residue was purified by silica gel column chromatography with 0-5% methanol in dichloromethane to give 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (82 mg, 68%). ESI-MS m/z 400.5 $(M+1)^+$; Retention time: 1.78 minutes (3 min).

Step 3: 6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-(3-piperidylsulfonyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2655)



A solution of 6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (32 mg, 0.1 mmol) and NaH (5 mg, 0.1 mmol) in DMF (800 µL) was stirred for 30 minutes, and benzyl 3-chlorosulfonylpiperidine-1-carboxylate (51 mg, 0.2 mmol) was added. The reaction mixture was stirred for 16 hours. The reaction was filtered and purified using a reverse phase HPLC-MS method using a Luna C18 (2) column (75 × 30 mm, 5

μm particle size) and a dual gradient run from 1-99% mobile phase B over 15.0 minutes (mobile phase A = H_2O (5 mM HCl), mobile phase B = CH_3CN). The intermediate and palladium (10 mg, 0.01 mmol) (10% on carbon) in MeOH was stirred under a hydrogen atmosphere for 17 hours. The reaction was filtered and purified using a reverse phase HPLC-MS method using a Luna C18 (2) column (75 x 30 mm, 5 μm particle size) and a dual gradient run from 1-99% mobile phase B over 15.0 minutes (mobile phase A = H_2O (5 mM HCl), mobile phase B = CH_3CN) to give the desired compound. ESI-MS m/z 547.6 $(M+1)^+$; Retention time: 1.65 minutes (3 min)

The following compounds can be synthesized using the procedures described herein:

6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-pyrrolidin-3-ylsulfonyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2042),


N-cyclopentylsulfonyl-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2113),

6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[[(3*R*)-3-piperidyl]sulfamoyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2677),

6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-(3-pyridylsulfamoyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1448) and

6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-[(2-methoxy-3-pyridyl)sulfamoyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1359).


PREPARATION 138: *N*-[(6-amino-2-pyridyl)sulfonyl]-5-fluoro-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2473)



(3-fluoro-5-*iso*butoxy-phenyl)boronic acid (5 mg, 0.02 mmol), *N*-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-5-fluoro-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (10 mg, 0.02 mmol), 1,3-bis(2,6-di*iso*propylphenyl)-4,5-dihydroimidazole; 3-chloropyridine; dichloropalladium (3 mg, 0.005 mmol) and K_2CO_3 (3 mg, 0.02 mmol) were combined in *N*-propanol (0.5 mL). The reaction mixture was heated to 90 °C for 16 h under an atmosphere of nitrogen. The reaction mixture was filtered and purified by reverse phase HPLC using 1-99% ACN in water (0.05% HCl modifier) over 15 minutes to give the desired compound as a pale solid. 1H NMR (400 MHz, DMSO-d6) δ 12.62 (s, 1H), 7.73 (d, J = 11.2 Hz, 1H), 7.36 (s, 1H), 7.24 (dd, J = 15.0, 8.6 Hz, 2H), 6.96 (dt, J = 10.9, 2.5 Hz, 1H), 6.70 (d, J = 8.3 Hz, 1H), 6.53 (s, 1H), 3.82 (d, J = 6.6 Hz, 2H), 2.59 (p, J = 10.2 Hz, 2H), 2.22 (s, 1H), 2.07 - 1.98 (m, 1H), 1.87 (dd, J = 12.0, 5.7 Hz, 1H), 1.54 (d, J = 6.6 Hz, 6H), 1.41 (t, J = 12.1 Hz, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.87 (d, J = 6.3 Hz, 3H).1H NMR (400 MHz, DMSO-d6) δ 12.62 (s, 1H), 7.73 (d, J = 11.2 Hz, 1H), 7.36 (s, 1H), 7.24 (dd, J = 15.0, 8.6 Hz, 2H), 6.96 (dt, J = 10.9, 2.5 Hz, 1H), 6.70 (d, J = 8.3 Hz, 1H), 6.53 (s, 1H), 3.82 (d, J = 6.6 Hz, 2H), 2.59 (p, J = 10.2 Hz, 2H), 2.22 (s, 1H), 2.07 - 1.98 (m, 1H), 1.87 (dd, J = 12.0, 5.7 Hz, 1H), 1.54 (d, J = 6.6 Hz, 6H), 1.41 (t, J = 12.1 Hz, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.87 (d, J = 6.3 Hz, 3H).

PREPARATION 139: *N*-[(3*R*)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-phenylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2477)


Step 1: 6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-phenylpyrrolidin-1-yl)pyridine-3-carboxylic acid



A mixture of 2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylic acid (227 mg, 0.7 mmol), 2-phenylpyrrolidine (206 mg, 1.4 mmol), CsF (213 mg, 1.4 mmol), and K_2CO_3 (194 mg, 1.4 mmol) in DMSO (3.5 mL) was stirred at 120 °C for 16 hours. The reaction mixture was diluted with water, adjusted to pH 4 with 1 M HCl, and extracted with ethyl acetate (2 x). The combined extracts were washed with brine, dried over sodium sulfate, and evaporated

under reduced pressure. The residue was purified by silica gel column chromatography with 0-5% methanol in dichloromethane to give 6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-phenylpyrrolidin-1-yl)pyridine-3-carboxylic acid (190 mg, 62%). ESI-MS m/z 435.5 $(M+1)^+$; Retention time: 0.9 minutes (3 min).

Step 2: *N*-[(3*R*)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-phenylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2477)




A solution of 6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-phenylpyrrolidin-1-yl)pyridine-3-carboxylic acid (43 mg, 0.1 mmol) and chlorosulfonyl isocyanate (11 uL, 0.1 mmol) in dichloromethane was stirred for 30 minutes, and *tert*-butyl *N*-[(3*R*)-pyrrolidin-3-yl]carbamate (27 mg, 0.2 mmol) and triethylamine (21 uL, 0.2 mmol) were added. The reaction mixture was stirred for 16 hours, and HCl (4M in dioxane) (200 uL, 0.8 mmol) was added. After stirring for four hours, the sovent was removed under reduced pressure, and the residue was purified using a reverse phase HPLC-MS method using a Luna C18 (2) column (75 x 30 mm, 5 μm particle size) and a dual gradient run from 1-99% mobile phase B over 15.0 minutes (mobile phase A = H_2O (5 mM HCl), mobile phase B = CH_3CN) to give *N*-[(3*R*)-3-aminopyrrolidin-1-yl]sulfonyl-6-(3-fluoro-5-isobutoxy-phenyl)-2-(2-phenylpyrrolidin-1-yl)pyridine-3-carboxamide (Compound 2477) (9 mg, 16%). ESI-MS m/z 582.6 $(M+1)^+$; Retention time: 1.82 minutes (3 min).

The following compound can be synthesized using the procedures described herein:

N-[(3*R*)-3-aminopyrrolidin-1-yl]sulfonyl-2-(2-benzyl-2-methyl-pyrrolidin-1-yl)-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (Compound 2325)


PREPARATION 140: 3-(cyclopropylmethyl)-2,2-dimethyl-pyrrolidine

Step 1: 1-benzyl-3-(cyclopropylmethyl)pyrrolidin-2-one



1-Benzylpyrrolidin-2-one (5.0 g, 28.6 mmol) was dissolved in THF (100 mL) and cooled to -78°C. n-BuLi (2.5M, 13.7 mL, 34.3 mmol) was added dropwise at -78 °C and the reaction mixture was stirred at -78 °C for 30 minutes. (Bromo-methyl)cyclopropane (4.6 g, 34.3 mmol) was added and stirring continued at -78 °C for 30 minutes. The reaction mixture was allowed to reach RT slowly overnight. The mixture was quenched with sat. aq. NH_4Cl and extracted with EtOAc (2×). The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (heptanes/EtOAc 2:1) affording 1-benzyl-3-(cyclopropylmethyl)-pyrrolidin-2-one (3.9 g, 60%) as a colorless oil. 1H NMR ($CDCl_3$, 300 MHz): 0.05 (m, 2H); 0.43 (m, 2H); 0.74 (m,1H); 1.22 (m, 1H); 1.68 (m, 1H); 1.76 (m, 1H); 2.21 (m, 1H); 2.55 (m, 1H); 3.08 (m, 2H); 4.43 (q, 2H); 7.24 (m, 5H).


Step 2: 1-benzyl-3-(cyclopropylmethyl)-2,2-dimethyl-pyrrolidine



$ZrCl_4$ (3.96 g, 17.0 mmol) was added to THF (100 mL) at -10°C and the reaction mixture was stirred for 30 minutes. 1-Benzyl-3-(cyclopropylmethyl)pyrrolidin-2-one (3.95 g, 17.0 mmol) was added as a solution in THF (10 mL). The reaction mixture was stirred for 10 minutes, then MeMgBr (3.0 M, 34 mL, 102 mmol) was added dropwise at 0°C. The reaction mixture was allowed to reach RT overnight. It was quenched at 0°C with 30% aq. NaOH (20 mL) and stirred for 10 min. Water (100 mL) was added and the reaction mixture was extracted with CH_2Cl_2 (2×). The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (heptanes/EtOAc 4:1) affording 1-benzyl-3-(cyclopropylmethyl)-2,2-dimethylpyrrolidine (2.0 g, 48%) as a colorless oil. 1H NMR ($CDCl_3$, 300 MHz): -0.03 (m, 1H);

0.07 (m, 1H); 0.42 (m, 2H); 0.66 (m, 1H); 0.76 (s, 3H); 1.12 (s, 3H); 1.19 (m, 2H); 1.39 (m, 1H); 1.90 (m, 1H); 2.03 (m, 1H); 2.32 (m, 1H); 2.85 (m, 1H); 3.20 (d, 1H); 3.87 (d, 1H); 7.25 (m, 5H).


Step 3: 3-(cyclopropylmethyl)-2,2-dimethyl-pyrrolidine



1-Benzyl-3-(cyclopropylmethyl)-2,2-dimethylpyrrolidine (2.0 g, 8.2 mmol) was dissolved in MeOH (100 mL) and conc. HCl (5 mL) was added. Pd/C (400 mg) was added and the reaction mixture was stirred over the weekend under 5 bars of H_2. The reaction mixture was filtered over Celite and concentrated under reduced pressure affording 3-(cyclopropylmethyl)-2,2-dimethyl-pyrrolidine hydrochloride (1.3 g, 84%) as an off-white solid. 1H NMR (CD_3OD, 300 MHz): 0.07 (m, 1H); 0.13 (m, 1H); 0.47 (m, 2H); 0.72 (m, 1H); 1.17 (s, 3H); 1.19 (m, 1H); 1.34 (m, 1H); 1.43 (s, 3H); 1.83 (m, 1H); 2.01 (m, 1H); 2.38 (m, 1H); 3.26 (m, 2H). ESI-MS m/z calc. 153.15175, found 154.0 $(M+1)^+$.

PREPARATION 141: *N*-(benzenesulfonyl)-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide


Step 1: *N*-(benzenesulfonyl)-2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide



To a solution of the 2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylic acid (10.0 g, 30.89 mmol) in DMF (30.0 mL) at ambient temperature in a round bottom flask was slowly added carbonyldiimidazole (5.510 g, 33.98 mmol) portionwise and the mixture stirred for 100 min. Meanwhile to benzenesulfonamide (6.069 g, 38.61 mmol) in DMF (30.0 mL) (homogenous solution) in another round bottom flask was added NaHMDS in THF (38.61 mL of 1.0 M, 38.61 mmol) portionwise via syringe over 30-45 min and on completion of

addition the mixture was stirred a further 30 min. The mixture containing the activated acid was then added to the mixture containing the deprotonated sulfonamide and the combined mixture was stirred 1 h. The reaction was cooled with a 0°C bath and quenched by addition of 12N aqueous HCl (11.58 mL) in portions over 2-3 minutes resulting in precipitated solids. Transferred the reaction mixture to a separatory funnel and ethyl acetate (100.0 mL) was added. Added 1N aqueous HCl (20.0 mL) giving a pH = 2-3 then separated the layers and washed the organic layer with 5:1 water/saturated aqueous brine (120.0 mL), saturated aqueous brine (1 X 50mL, 1 X 30 mL), dried (sodium sulfate), filtered and concentrated under reduced pressure to a clear light yellow oil that was concentrated from isopropanol several more times resulting in precipitation of a solid. The solid was slurried overnight in isopropanol then filtered and washed the solid with heptane (50 mL) and dried in vacuo giving *N*-(benzenesulfonyl)-2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (10.22 g, 22.08 mmol, 71.47%) as a white solid. ^{1}H NMR (400 MHz, DMSO) δ 12.85 (s, 1H), 8.15 (d, J = 8.0 Hz, 1H), 8.09 (d, J = 8.0 Hz, 1H), 8.02 (dd, J = 5.3, 3.3 Hz, 2H), 7.76 (d, J = 7.4 Hz, 1H), 7.69 (t, J = 7.6 Hz, 2H), 7.51 – 7.43 (m, 2H), 6.99 (dd, J = 10.8, 2.3 Hz, 1H), 3.85 (d, J = 6.5 Hz, 2H), 2.04 (dt, J = 13.3, 6.6 Hz, 1H), 1.00 (d, J = 6.7 Hz, 6H). ESI-MS m/z calc. 462.08163, found 463.19 $(M+1)^+$; Retention time: 2.93 minutes [5 minute method].

Step 2: *N*-(benzenesulfonyl)-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide




To a round bottom flask outfitted with a reflux condenser was added *N*-(benzenesulfonyl)-2-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (10.0 g, 21.60 mmol) and NMP (40 mL) and stirring was commenced. Warmed to 50°C and began portionwise addition of potassium carbonate (5.970 g, 43.20 mmol) followed by (4*S*)-2,2,4-trimethylpyrrolidine (4.890 g, 43.20 mmol) in one portion. After stirring for 10 min, heated the mixture to 125°C for 65 h, then cooled to 10°C and added 1N aqueous HCl (50.0 mL, 50.00 mmol) in portions to give pH 1-2 and a precipitated solid. Added ethyl acetate (100.0 mL) to dissolve solid and diluted the aqueous layer with water (50.0 mL) and stirred for 10 min. The

mixture was transferred to a separatory funnel and layers were allowed to separate. Added aqueous 1N HCl dropwise until all solids were dissolved. Separated the layers and the aqueous layer was back extracted with ethyl acetate (50.00 mL) followed by combination of the organic layers. To the combined organic layers was added water (50.00 mL) giving an emulsion which was clarified by the addition of 1N aqueous HCl (25.00 mL). Separated the layers then the organic layer was washed with saturated aqueous brine (50.00 mL), dried over Na_2SO_4, filtered through celite and rinsed with ethyl acetate (30.00 mL). The filtrate was concentrated under reduced pressure and the residue was purified by silica gel chromatography using a gradient from 100% hexanes to 50% EtOAc giving a light amber oil which was evaporated from isopropanol several times under reduced pressure providing *N*-(benzenesulfonyl)-6-(3-fluoro-5-isobutoxy-phenyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (9.73 g, 18.03 mmol, 83.5%) as a yellow solid. ^{1}H NMR (400 MHz, DMSO-d6) δ 12.57 (s, 1H), 8.16 – 7.88 (m, 2H), 7.82 – 7.57 (m, 4H), 7.47 (t, J = 1.8 Hz, 1H), 7.40 (dt, J = 9.9, 2.0 Hz, 1H), 7.27 (d, J = 8.1 Hz, 1H), 6.89 (dt, J = 10.8, 2.3 Hz, 1H), 3.83 (d, J = 6.6 Hz, 2H), 2.48 – 2.28 (m, 2H), 2.07 (dtt, J = 26.6, 13.4, 6.4 Hz, 2H), 1.83 (dd, J = 11.9, 5.5 Hz, 1H), 1.57 (d, J = 17.3 Hz, 6H), 1.38 (t, J = 12.1 Hz, 1H), 1.04 (d, J = 6.1 Hz, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.66 (d, J = 6.3 Hz, 3H). ESI-MS m/z calc. 539.2254, found 540.0 $(M+1)^+$; Retention time: 3.25 minutes [5 minute method].

PREPARATION 142: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide


Step 1: Ethyl 2-chloro-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxylate



Ethyl 2,6-dichloropyridine-3-carboxylate (7.00 g, 31.81 mmol) and (6-isopropoxy-3-pyridyl)boronic acid (5.04 g, 27.85 mmol) were combined and dissolved in ethanol (50.40 mL) and toluene (50.40 mL). A suspension of sodium carbonate (10.12 g, 95.47 mmol) in water (10.08 mL) was added. Under nitrogen, tetrakis(triphenylphosphine)palladium (1.103 g, 0.9547 mmol) was added. The reaction mixture was allowed to warm to 80°C and stirred for 2 h. The volatiles were removed under reduced pressure. The remaining solids were partitioned


between water (75 mL) and ethyl acetate (75 mL). The organic layer was washed with saturated aqueous sodium chloride solution (1× 75 mL), dried over sodium sulfate, filtered and concentrated under reduced pressure. The material was subjected to silica gel column chromatography using a gradient from 100% hexanes to 5% ethyl acetate in hexanes. Ethyl 2-chloro-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxylate (3.95 g, 12.07 mmol, 43.33%) was obtained as a clear colorless oil. ^{1}H NMR (400 MHz, DMSO) δ 8.91 (s, 1H), 8.37 (dd, J = 8.7, 2.3 Hz, 1H), 8.30 (d, J = 8.1 Hz, 1H), 8.10 (d, J = 8.1 Hz, 1H), 6.89 (d, J = 8.7 Hz, 1H), 5.34 (dt, J = 12.3, 6.1 Hz, 1H), 4.36 (q, J = 7.1 Hz, 2H), 2.51 (d, J = 1.7 Hz, 1H), 1.42 - 1.30 (m, 9H). ESI-MS m/z calc. 320.09277, found 321.2 $(M+1)^+$; Retention time: 0.72 minutes.

Step 2: 2-chloro-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxylic acid



A solution of sodium hydroxide (2.594 g, 64.85 mmol) in water (20.80 mL) was added to a solution of ethyl 2-chloro-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxylate (4.16 g, 12.97 mmol) in isopropanol (20.8 mL) stirring at 90°C. The reaction mixture was stirred at 90°C for 30 min. After cooling to room temperature, the reaction mixture was diluted with ethyl acetate (75 mL) and extracted with saturated aqueous sodium bicarbonate (1× 75 mL) and aqueous NaOH (1 N, 5 × 50 mL). The aqueous layers were combined and acidified to pH = 1 with the addition of aqueous 6 N HCl. The white precipitate that formed was collected by vacuum filtration and rinsed with aqueous 1 N HCl to provide 2-chloro-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxylic acid (4.2446 g, 12.89 mmol, 99.43%) as a white solid. ESI-MS m/z calc. 292.06146, found 293.1 $(M+1)^+$; Retention time: 0.59 minutes.


Step 3: 2-aminopyridine-3-sulfonamide



Ammonium hydroxide (590.3 mL of 28 %w/v, 4.716 mol) was stirred and cooled in an ice bath. 2-chloropyridine-3-sulfonyl chloride (50 g, 235.8 mmol) was dissolved in

dichloromethane (150 mL) and added dropwise to the stirred ammonia solution to maintain a temperature between 5-12°C. On completion of addition, the mixture was stirred 1 h. Stirring was halted and the resulting layers were separated. Concentrated the aqueous layer under vacuum to remove ammonia resulting in a white solid precipitate which was collected by filtration and washed with water. The material was dissolved in ammonium hydroxide (236.1 mL of 28 %w/v, 1.886 mol) and heated in a pressure vessel at 100°C overnight then reaction temperature was increased to 150°C and stirred overnight. The mixture was cooled to room temperature resulting in a precipitate then concentrated under vacuum to remove ammonia resulting in a white solid precipitate which was collected by filtration and washed with methyl *tert*-butyl ether (100 mL) to give 2-aminopyridine-3-sulfonamide (25.73 g, 148.6 mmol, 63.00%) ESI-MS m/z calc. 173.0259, found 174.0 $(M+1)^+$.

Step 4: *N*-[(2-amino-3-pyridyl)sulfonyl]-2-chloro-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide



To a solution of 2-aminopyridine-3-sulfonamide (4.068 g, 23.49 mmol) in DMF (38.50 mL) (homogenous solution) in a round bottom flask was added NaH (939.5 mg, 23.49 mmol) portionwise giving an off-white thick slurry and a slight exotherm from 20°C to 32°C during addition. The mixture was stirred at ambient temperature for 30 min. Meanwhile, to a solution of 2-chloro-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxylic acid (5.5 g, 18.79 mmol) in DMF (38.50 mL) at ambient temperature in another round bottom flask was slowly added carbonyldiimidazole (3.352 g, 20.67 mmol) portionwise and the mixture stirred for 20 min over which time it became a light yellow homogenous solution which was then added to the reaction mixture containing the deprotonated sulfonamide. During the addition the mixture stayed homogenous with a slight exotherm to 31°C. The combined mixture was stirred at ambient temperature affording a dark orange homogenous mixture and after 5 min was slowly added to chilled HCl (93.95 mL of 1 M, 93.95 mmol) (slightly exothermic) giving a pH of ~1. The mixture afforded a precipitate and the slurry was stirred for 60 min at ambient temperature. The mixture was then filtered and the solid was washed three times with 50 mL of water. The solid

was then washed three times with 50 mL of isopropanol giving an off-white solid that was air dried for 1 h. The solid was next slurried in 50 mL of isopropanol and the precipitate was filtered and washed three times with 5 mL of isopropanol. The off-white solid obtained was air dried for 2 h, then in vacuo at 45°C for 48 h giving *N*-[(2-amino-3-pyridyl)sulfonyl]-2-chloro-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (8.0 g, 17.86 mmol, 95.06%) ^{1}H NMR (400 MHz, DMSO-d6) δ 8.87 (d, J = 2.5 Hz, 1H), 8.33 (dd, J = 8.7, 2.6 Hz, 1H), 8.24 (dd, J = 5.1, 1.8 Hz, 1H), 8.13 - 8.00 (m, 3H), 6.96 (s, 2H), 6.88 (d, J = 8.7 Hz, 1H), 6.83 (dd, J = 7.8, 5.0 Hz, 1H), 5.33 (hept, J = 6.3 Hz, 1H), 1.32 (d, J = 6.2 Hz, 6H). ESI-MS m/z calc. 447.0768, found 448.18 $(M+1)^+$; Retention time: 1.55 minutes.

Step 5: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide



A vial equipped with magnetic stir bar was charged with *N*-[(2-amino-3-pyridyl)sulfonyl]-2-chloro-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (147.8 mg, 0.330 mmol), (4*S*)-2,2,4-trimethylpyrrolidine (74.71 mg, 0.660 mmol) and potassium carbonate (228.0 mg, 1.65 mmol) in DMSO (739 µL). The vial was sealed and stirred at 120°C for 16 hours then at 140°C for 30 hours. The reaction mixture was cooled, diluted with minimal DMSO and methanol and filtered. The filtrate was purified directly using a reverse phase HPLC-MS method using a Luna C18 (2) column (75 x 30 mm, 5 µm particle size) sold by Phenomenex (pn: 00C-4252-U0-AX), and a dual gradient run from 1-99% mobile phase B over 15.0 minutes (mobile phase A = H_2O (5 mM HCl), mobile phase B = CH_3CN, flow rate = 50 mL/min, injection volume = 950 µL and column temperature = 25 °C) giving *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-isopropoxy-3-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (25.2 mg, 0.048 mmol, 14.6%). ^{1}H NMR (400 MHz, DMSO) δ 12.49 (s, 1H), 8.84 (d, J = 2.4 Hz, 1H), 8.31 – 8.27 (m, 1H), 8.25 (dd, J = 7.4, 4.4 Hz, 1H), 8.03 (dd, J = 7.9, 1.7 Hz, 1H), 7.75 (d, J = 8.0 Hz, 1H), 7.20 (d, J = 8.0 Hz, 1H), 6.86 (d, J = 8.7 Hz, 1H), 6.77 (dd, J = 7.8, 4.8 Hz, 1H), 6.69 (s, 2H), 5.37 – 5.26 (m, 1H), 2.52 (s, 1H), 2.14 (s, 1H), 1.85 (dd, J = 11.9, 5.5 Hz, 1H), 1.58 (d, J = 10.7 Hz, 6H), 1.39 (t, J = 12.1 Hz, 1H), 1.31 (d, J = 6.2 Hz, 6H), 1.26 (d, J = 13.3 Hz, 1H), 0.86

(t, J = 6.9 Hz, 1H), 0.76 (d, J = 6.3 Hz, 3H). ESI-MS m/z calc. 524.2206, found 525.3 $(M+1)^+$; Retention time: 1.38 minutes.

The following compounds can be synthesized using the procedures described herein:


N-[(2-amino-6-methoxy-3-pyridyl)sulfonyl]-6-(6-*iso*propoxy-3-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2692),

N-[(2-amino-3-pyridyl)sulfonyl]-2-[(4*S*)-3,3-dideuterio-2,2-dimethyl-4-(trideuteriomethyl)pyrrolidin-1-yl]-6-(6-isopropoxy-3-pyridyl)pyridine-3-carboxamide (Compound 1543) and

N-[(6-amino-2-pyridyl)sulfonyl]-2-[(2*R*,5*S*)-2,5-dimethylpyrrolidin-1-yl]-6-[6-[1,2,2,2-tetradeuterio-1-(trideuteriomethyl)ethoxy]-3-pyridyl]pyridine-3-carboxamide (Compound 2714)


PREPARATION 143: 6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-*N*-[[6-(1-piperidyl)-2-pyridyl]sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 484)

Step 1: 6-(1-piperidyl)pyridine-2-sulfonamide




To 6-fluoro-2-pyridinesulfonamide (215 mg, 1.220 mmol) was added piperidine (312 mg, 362 µL, 3.66 mmol) and the reaction mixture was heated at 90 °C for 2.5 hours. The reaction mixture was cooled and resulting solid was dried.1 H NMR (400 MHz, DMSO-d6) δ 7.66 (dd, J = 8.8, 7.2 Hz, 1H), 7.03 (d, J = 7.2 Hz, 1H), 6.98 (d, J = 8.6 Hz, 1H), 2.75 (d, J = 5.5 Hz, 3H), 1.61 (p, J = 3.4, 2.9 Hz, 2H), 1.54 (dd, J = 7.0, 3.9 Hz, 3H), 1.49 (s, 4H). ESI-MS m/z found 242.2 $(M+1)^+$; Retention time: 0.45 minutes.

Step 2: 2-chloro-6-(3-(neopentyloxy)-1*H*-pyrazol-1-yl)-*N*-((6-(piperidin-1-yl)pyridin-2-yl)sulfonyl)nicotinamide



To 2-chloro-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]pyridine-3-carboxylic acid (100 mg, 0.323 mmol) in DMF (0.6 mL) at room temperature was added carbonyl diimidazole (78.5 mg, 0.484 mmol) and the reaction mixture was stirred for 40 minutes. Separately 6-(1-piperidyl)pyridine-2-sulfonamide (93.5 mg, 0.387 mmol) in DMF (0.6 mL) was cooled to 0 °C and treated with sodium hydride (15.5 mg, 0.387 mmol) and stirred for 10 min and then at room temperature for 30 min. The reaction mixtures were then combined and stirred at room temperature for 30 min. The reaction mixture was cooled to 0 °C and acetic acid (194 mg, 184 µL, 3.23 mmol) was added followed by HCL (269 µL of 6 M, 1.61 mmol) and the reaction mixture was diluted with EtOAc (5 mL) and the layers were separated. The organics were dried and concentrated under reduced pressure to give a residue which was purified by silica gel column chromatography and eluted with 0-14% MeOH-DCM to give 2-chloro-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-*N*-[[6-(1-piperidyl)-2-pyridyl]sulfonyl]pyridine-3-carboxamide (158 mg, 82%) ESI-MS m/z found 533.2 $(M+1)^+$; Retention time: 0.89 minutes (1 min); 1H NMR (400 MHz, DMSO-d6) δ 8.44 – 8.38 (m, 1H), 8.05 (d, J = 8.3 Hz, 1H), 7.95 (s, 1H), 7.79 – 7.70 (m, 2H), 7.23 (d, J = 7.2 Hz, 1H), 7.12 (d, J = 8.7 Hz, 1H), 6.22 (t, J = 2.5 Hz, 1H), 3.93 (d, J = 1.2 Hz, 2H), 3.59 (t, J = 5.4 Hz, 3H), 1.66 – 1.58 (m, 2H), 1.53 (td, J = 10.2, 4.8 Hz, 3H), 1.00 (s, 9H).

Step 3: 6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-*N*-[[6-(1-piperidyl)-2-pyridyl]sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 484)




A mixture of 2-chloro-6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-*N*-[[6-(1-piperidyl)-2-pyridyl]sulfonyl]pyridine-3-carboxamide (140 mg, 0.223 mmol), potassium carbonate (185 mg, 1.34 mmol) and (4*S*)-2,2,4-trimethylpyrrolidine (HCl salt) (75 mg, 0.501 mmol) in DMSO (744 µL) was heated at 150 °C for 5 h. The reaction mixture was cooled, diluted with EtOAc and the aqueous and organic layers were separated. The organics were washed with brine, dried, concentrated under reduced pressure and the resulting residue was purified by silica gel column chromatography eluting with 0-15% MeOH-DCM to give 6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-*N*-[[6-(1-piperidyl)-2-pyridyl]sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 484) (38 mg, 28%). ESI-MS m/z calc 609.31, found 610.1 $(M+1)^+$; retention time 2.1 min (3 min run); 1H NMR (400 MHz, Methanol-d4) δ 8.25 (d, J = 2.8 Hz, 1H), 7.88 (d, J = 8.3 Hz, 1H), 7.69 (dd, J = 8.7, 7.2 Hz, 1H), 7.31 (d, J = 7.2 Hz, 1H), 6.98 (dd, J = 8.5, 6.5 Hz, 2H), 5.99 (d, J = 2.8 Hz, 1H), 3.90 (s, 2H), 3.63 (q, J = 6.1, 5.3 Hz, 4H), 2.80 - 2.66 (m, 2H), 2.26 (d, J = 10.4 Hz, 1H), 1.90 (dd, J = 11.9, 5.6 Hz, 1H), 1.68 (q, J = 5.1, 4.7 Hz, 2H), 1.61 (d, J = 5.3 Hz, 10H), 1.47 (t, J = 12.2 Hz, 1H), 1.04 (s, 9H), 0.88 (d, J = 6.3 Hz, 3H).

The following compound can be synthesized using the procedures described herein:

6-[3-(2,2-dimethylpropoxy)pyrazol-1-yl]-*N*-[[2-(1-piperidyl)-3-pyridyl]sulfonyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2670)


PREPARATION 144: *N*-[(6-amino-2-pyridyl)sulfonyl]-3-(3-fluoro-5-isobutoxy-phenyl)-5-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyrazine-2-carboxamide (Compound 2673)

Step 1: 3,5-dichloro-*N*-[(6-nitro-2-pyridyl)sulfonyl]pyrazine-2-carboxamide



To a stirred heterogeneous mixture of 3,5-dichloropyrazine-2-carboxylic acid (2.21 g, 11.5 mmol) in anhydrous CH_2Cl_2 (20 mL), thionyl chloride (7.5 mL, 103 mmol) was added under an atmosphere of nitrogen, followed by anhydrous DMF (30 mg, 0.41 mmol) and the reaction mixture was stirred at room temperature for 1 hour. The reaction mixture was evaporated, dissolved in CH_2Cl_2 (40 mL) and added dropwise to a solution of 6-nitropyridine-2-sulfonamide (2.32 g, 11.5 mmol) with triethylamine (8.5 mL, 61.0 mmol) at 0 °C, stirred overnight (24 h) while allowing to warm to ambient temperature. The reaction mixture was concentrated under reduced pressure. The residue was partitioned between CH_2Cl_2 (10 mL) and ice-water (200 mL) and acidified with 2 M HCl to pH about 4.0. The resulting solid was filtered, the layers separated and the aqueous layer was extract with CH_2Cl_2 (3 x 50 mL). The combined organics were washed sequentially with 1N HCl, saturated sodium chloride solution then dried over sodium sulfate, filtered and evaporated under reduced pressure. The crude material was purified by triturating with dichloromethane (10 mL), filtration and drying to give 3,5-dichloro-*N*-[(6-nitro-2-pyridyl)sulfonyl]pyrazine-2-carboxamide (2.90 g, 67%) as a white solid. ESI-MS m/z calc. 376.93884, found 378.0 $(M+1)^+$; Retention time: 0.6 minutes.


Step 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-3,5-dichloro-pyrazine-2-carboxamide



To a solution of 3,5-dichloro-*N*-[(6-nitro-2-pyridyl)sulfonyl]pyrazine-2-


carboxamide (830 mg, 2.20 mmol) in THF (8 mL) and EtOH (4 mL), iron (Fe) powder (613 mg, 11.0 mmol) was added, followed by HCl (6 N) (1.8 mL of 6.0 M, 10.80 mmol). The reaction mixture was heated at 60 °C for 15 min. After cooling to room temperature, the reaction mixture was filtered and the filtrate was concentrated under reduced pressure. The residue was diluted with EtOAc (50 mL) and washed with NaOH (4 mL, 1 N). The layers were filtered again and the aqueous layer was acidified to pH 2 with the addition of 1 N HCl (aq) and extracted with EtOAc (2 × 50 mL). The organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure. The crude material was purified from by silica gel column chromatography using 0-20 % MeOH in CH_2Cl_2 as eluent to give *N*-[(6-amino-2-pyridyl)sulfonyl]-3,5-dichloro-pyrazine-2-carboxamide (404 mg, 53%) ESI-MS m/z calc. 346.96466, found 348.0 $(M+1)^+$; Retention time: 0.53 minutes.

Step 3: *N*-[(6-amino-2-pyridyl)sulfonyl]-5-chloro-3-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyrazine-2-carboxamide and (*S*)-*N*-((6-aminopyridin-2-yl)sulfonyl)-3-chloro-5-(2,2,4-trimethylpyrrolidin-1-yl)pyrazine-2-carboxamide



To a solution of *N*-[(6-amino-2-pyridyl)sulfonyl]-3,5-dichloro-pyrazine-2-carboxamide (36 mg, 0.10 mmol) in DMSO (142 µL), DIEA (90 µL, 0.52 mmol) and (4*S*)-2,2,4-trimethylpyrrolidine (32 mg, 0.26 mmol) was added. The reaction mixture was stirred for 6 hours at 85 °C. The cooled reaction was purified via reverse phase HPLC-MS method using a dual gradient run from 15-99% mobile phase B over 15.0 minutes (mobile phase A = H_2O (5 mM HCl), mobile phase B = acetonitrile) to yield a mixture of *N*-[(6-amino-2-pyridyl)sulfonyl]-5-chloro-3-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyrazine-2-carboxamide and (*S*)-*N*-((6-aminopyridin-2-yl)sulfonyl)-3-chloro-5-(2,2,4-trimethylpyrrolidin-1-yl)pyrazine-2-carboxamide (28 mg, 64%) ESI-MS m/z calc. 424.10843, found 425.2 $(M+1)^+$.


Step 4: *N*-[(6-amino-2-pyridyl)sulfonyl]-3-(3-fluoro-5-isobutoxy-phenyl)-5-[(*4S*)-2,2,4-trimethylpyrrolidin-1-yl]pyrazine-2-carboxamide (Compound 2673)



To a solution of *N*-[(6-amino-2-pyridyl)sulfonyl]-5-chloro-3-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyrazine-2-carboxamide (*S*)-*N*-((6-aminopyridin-2-yl)sulfonyl)-3-chloro-5-(2,2,4-trimethylpyrrolidin-1-yl)pyrazine-2-carboxamide (60 mg, 0.14 mmol) and (3-fluoro-5-isobutoxy-phenyl)boronic acid (60 mg, 0.28 mmol) in dioxane (3.0 mL) were added H_2O (1.0 mL), K_2CO_3 (78 mg, 0.56 mmol) and bis(2-diphenylphosphanylcyclopenta-2,4-dien-1-yl)iron; dichloropalladium [$Cl_2Pd(dppf).CH_2Cl_2$] (8 mg, 0.011 mmol). Nitrogen was bubbled for 5 min and the reaction mixture was sealed and stirred at 120 °C under microwave radiation for 30 min. The reaction mixture was allowed to cool to ambient temperature, filtered through a and purified *via* reverse phase HPLC-MS (using a Luna C18 (2) column (75 x 30 mm, 5 μm particle size) and a dual gradient run from 30-99% mobile phase B over 30.0 minutes; mobile phase A = H_2O (5 mM HCl), mobile phase B = CH_3CN) to furnish *N*-[(6-amino-2-pyridyl)sulfonyl]-3-(3-fluoro-5-isobutoxy-phenyl)-5-[(*4S*)-2,2,4-trimethylpyrrolidin-1-yl]pyrazine-2-carboxamide (Compound 2673) (5.7 mg, 6%) as off-white solid. 1H NMR (400 MHz, DMSO-d6) δ 11.58 (s, 1H), 7.89 (s, 1H), 7.62 - 7.53 (m, 1H), 7.11 (d, J = 7.3 Hz, 1H), 6.88 - 6.79 (m, 2H), 6.75 - 6.69 (m, 1H), 6.66 (d, J = 8.3 Hz, 1H), 6.52 (s, 2H), 3.83 (t, J = 9.1 Hz, 1H), 3.74 (d, J = 6.5 Hz, 2H), 3.12 (t, J = 10.4 Hz, 1H), 2.47 - 2.35 (m, 1H), 2.02 (dd, J = 12.8, 6.3 Hz, 2H), 1.61 (d, J = 12.2 Hz, 1H), 1.56 (s, 3H), 1.45 (s, 3H), 1.10 (d, J = 6.4 Hz, 3H), 0.96 (d, J = 6.8 Hz, 6H). ESI-MS m/z calc. 556.2268, found 557.4 $(M+1)^+$; Retention time: 2.15 minutes.


PREPARATION 145: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-4-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2679)

Step 1: 4-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylic acid



To 6-bromo-4-chloro-pyridine-3-carboxylic acid (205 mg, 0.865 mmol), (3-fluoro-5-isobutoxy-phenyl)boronic acid (184 mg, 0.865 mmol), and palladium;triphenylphosphane (30.0 mg, 0.0260 mmol) in dioxane (4 mL) was added solution of potassium carbonate (1.73 mL of 2 M, 3.46 mmol) under an atmosphere of nitrogen and the reaction mixture was stirred at 80 °C for 10 h. The reaction mixture was concentrated under reduced pressure, partitioned between water (15 mL) and EtOAc (15 mL). The aqueous layer was acidified with HCl and washed with EtOAc (3 x 15 mL). The combined organic layers was washed with brine (1× 25 mL), dried over sodium sulfate, filtered and concentrated under reduced pressure. The solid obtained was washed with hexanes to give 4-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylic acid (232 mg, 83%) as a tan solid. ESI-MS m/z calc. 323.07245, found 324.1 $(M+1)^+$; Retention time: 0.74 minutes.


Step 2: 4-chloro-6-(3-fluoro-5-isobutoxy-phenyl)-*N*-[(6-nitro-2-pyridyl)sulfonyl]pyridine-3-carboxamide



4-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxylic acid (104 mg, 0.32 mmol) and $SOCl_2$ (190 mg, 116 µL, 1.60 mmol) with a drop of DMF was stirred at ambient temperature for 30 minutes. The volatiles were removed under reduced pressure. In another flask, 6-nitropyridine-2-sulfonamide (65 mg, 0.32 mmol) and TEA (97 mg, 134 µL, 0.96 mmol) in DCM (1.2 mL) was cooled in an ice-water bath. A solution of the acid chloride generated in


DCM was added slowly. The reaction mixture was stirred in ice-water bath and allowed to reach room temperature. The reaction mixture was diluted with DCM, washed with 1N HCl, saturated NaHCO3, and brine. The organics were separated and dried over sodium sulfate and evaporated under reduced pressure to give 4-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-[(6-nitro-2-pyridyl)sulfonyl]pyridine-3-carboxamide (157 mg, 97%) ESI-MS m/z calc. 508.06195, found 509.1 $(M+1)^+$; Retention time: 0.74 minutes.

Step 3: *N*-[(6-amino-2-pyridyl)sulfonyl]-4-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide




4-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)-*N*-[(6-nitro-2-pyridyl)sulfonyl]pyridine-3-carboxamide (123 mg, 0.242 mmol) was dissolved in THF (615 µL) and EtOH (615 µL). Powdered Fe (135 mg, 2.42 mmol) with HCl (80.6 µL of 6 M, 0.483 mmol) was added. The reaction mixture was heated at 65 °C for 10 min. After cooling to room temperature, the reaction mixture was filtered, the filtrate was diluted with EtOAc and washed with NaOH (4 mL, 1 N). The layers were filtered again and aqueous layer was acidified to pH 2 with the addition of aqueous HCl and extracted with EtOAc (1× 8 mL). The organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure to give *N*-[(6-amino-2-pyridyl)sulfonyl]-4-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (61 mg, 53%) as a yellow powder. ESI-MS m/z calc. 478.08777, found 479.1 $(M+1)^+$; Retention time: 0.69 minutes.

Step 4: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-4-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2679)



To a solution of *N*-[(6-amino-2-pyridyl)sulfonyl]-4-chloro-6-(3-fluoro-5-isobutoxy-phenyl)pyridine-3-carboxamide (24 mg, 0.050 mmol) in DMSO (0.1 mL) was added CsF (23 mg, 0.15 mmol) and (4*S*)-2,2,4-trimethylpyrrolidine (27 mg, 0.15 mmol). The reaction mixture was sealed and stirred at 120 °C overnight. The reaction mixture was cooled, diluted with DMSO and purified via HPLC/MS utilizing a gradient of 1-99% acetonitrile in 5 mM aq HCl to yield *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-4-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2679) (5 mg, 19%). ESI-MS m/z calc. 555.23157, found 556.4 $(M+1)^+$; Retention time: 1.47 minutes.

PREPARATION 146: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(2-ethoxyethoxy)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2678)



N-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 852) (25 mg, 0.059 mmol), 2-ethoxyethanol (53 mg, 57 µL, 0.59 mmol), NaH (7.0 mg, 0.29 mmol), and scandium(3+) trifluoromethanesulfonate (2.9 mg, 0.0058 mmol) were dissolved in DMSO (0.5 mL) and heated at 120 °C for 8 h. The reaction mixture was partitioned between ethyl acetate and a 1N aq HCl solution. The organics were separated, washed with brine, dried over sodium sulfate and evaporated under reduced pressure. The crude material was purified by silica gel chromatography eluting with 0-100% ethyl acetate in hexanes to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(2-ethoxyethoxy)-2-[(4*S*)-2,2,4-

trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2678) (7 mg, 25%) ESI-MS *m/z* calc. 477.2046, found 478.4 $(M+1)^+$; Retention time: 1.25 minutes.


The following compounds can be synthesized using the procedures described herein:

N-[(2-amino-3-pyridyl)sulfonyl]-6-methoxy-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2704),

N-[(2-amino-3-pyridyl)sulfonyl]-6-(2-ethoxyethoxy)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2705) and

N-[(2-amino-3-pyridyl)sulfonyl]-6-[2-(2-ethoxyethoxy)ethoxy]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2706).

PREPARATION 147: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(4-iodo-2,6-dimethyl-phenoxy)pyridine-3-carboxamide (Compound 2671)



To *N*-[(6-amino-2-pyridyl)sulfonyl]-2-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (69 mg, 0.14 mmol) and 4-iodo-2,6-dimethyl-phenol (71 mg, 0.29 mmol) in NMP (4 mL) was added NaH (17 mg, 0.43 mmol). The reaction mixture was stirred for 5 min at rt then stirred at 150 °C for 2 h. The reaction mixture was quenched with NH_4Cl soln, the crude mixture purified by reverse phase silica gel column chromatography using 10-100% ACN in Water (TFA modifier) as eluent to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(4-iodo-2,6-dimethyl-phenoxy)pyridine-3-carboxamide (Compound 2671) (8 mg, 7%). 1H NMR (300 MHz, CDCl3) δ 8.47 (d, J = 8.1 Hz, 1H), 7.76 - 7.68 (m, 1H), 7.64 (d, J = 7.3 Hz, 1H), 7.52 (d, J = 8.1 Hz, 1H), 7.47 (s, 2H), 7.04 - 6.95 (m, 3H), 6.76 (d, J = 8.2 Hz, 1H), 6.63 (dt, J = 10.3, 2.2 Hz, 1H), 3.56 (t, J = 7.0 Hz, 2H), 2.13 - 2.06 (m, 6H), 1.01 (t, J = 6.6 Hz, 6H), 0.85 (t, J = 6.7 Hz, 3H), 0.47 (s, 1H). ESI-MS m/z found 691.4 $(M+1)^+$; Retention time: 0.83 minutes


The following compounds can be synthesized using the procedures described

herein:

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-benzoyl-2,6-dimethyl-phenoxy)-6-(6-*iso*propoxy-3-pyridyl)pyridine-3-carboxamide (Compound 2715) and

N-[(6-amino-2-pyridyl)sulfonyl]-2-(4-benzoyl-2,6-dimethyl-phenoxy)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 2669).


PREPARATION 148: *N*-[(6-amino-2-pyridyl)sulfonyl]-2,6-bis(4-benzoyl-2,6-dimethyl-phenoxy)pyridine-3-carboxamide (Compound 970)



To *N*-[(6-amino-2-pyridyl)sulfonyl]-2,6-dichloro-pyridine-3-carboxamide (487 mg, 1.40 mmol) in DMF (4 mL) was added (4-hydroxy-3,5-dimethyl-phenyl)-phenyl-methanone (390 mg, 1.57 mmol) and NaH (140.3 mg, 3.508 mmol). The reaction mixture was stirred at rt for 20 min, then 150 °C for 1 hr. The crude mixture purified by reverse phase silica gel column chromatography using 0-80% ACN in Water (TFA modifier) as eluent, then further purified by silica gel column chromatography eluting with 30-100% EtOAc in heptane to give *N*-[(6-amino-2-pyridyl)sulfonyl]-2,6-bis(4-benzoyl-2,6-dimethyl-phenoxy)pyridine-3-carboxamide (Compound 970) (51 mg, 4%). 1H NMR (300 MHz, DMSO) δ 8.24 (dd, J = 15.6, 8.2 Hz, 1H), 7.80 - 7.66 (m, 1H), 7.66 - 7.39 (m, 10H), 7.31 - 7.22 (m, 4H), 7.11 (d, J = 7.2 Hz, 1H), 6.74 (t, J = 10.3 Hz, 1H), 6.55 (d, J = 7.9 Hz, 1H), 6.27 (s, 2H), 1.92 (d, J = 1.9 Hz, 12H); ESI-MS m/z found 727.5 $(M+1)^+$; Retention time: 0.68 minutes (4 min run).


PREPARATION 149: *N*-[(6-fluoro-2-pyridyl)sulfonyl]-6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 698)

Step 1: Methyl 6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate




A mixture of methyl 2-chloro-6-phenyl-pyridine-3-carboxylate (224 mg, 0.904 mmol), 2,4,6-trimethylphenol (123 mg, 0.904 mmol), cesium carbonate (295 mg, 0.904 mmol) and DMF (3.14 mL) was heated at 80 °C overnight. The reaction mixture was cooled to room temperature before partitioned between ethyl acetate and 1N HCl. The layers were separated and the aqueous layer was extracted with ethyl acetate (2x). The combined organics were washed with brine, dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue was subjected to column chromatography (0-25% ethyl acetate/hexanes) to give methyl 6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (250 mg, 71%). ESI-MS m/z calc. 347.15213, found 348.4 $(M+1)^+$; Retention time: 2.28 minutes (3 min run).

Step 2: 6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid



Methyl 6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylate (562 mg, 1.62 mmol) was dissolved in methanol (8 mL): THF (3 mL) and NaOH (8.09 mL of 1 M, 8.09 mmol) was added. The reaction mixture was heated at 65 °C for 3 h. The reaction mixture was poured over ice and acidified with 6M HCl. The resulting white solid was collected and then dissolved in methanol and evaporated under reduced pressure to give 6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (466 mg, 86%) ESI-MS m/z calc. 333.1365, found 334.2 $(M+1)^+$; Retention time: 0.74 minutes.

Step 3: *N*-[(6-fluoro-2-pyridyl)sulfonyl]-6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 698)



To 6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxylic acid (150 mg, 0.450 mmol), HATU (171 mg, 0.450 mmol), DMF (1.5 mL), and triethylamine (137 mg, 188 µL, 1.35 mmol) was added 6-fluoropyridine-2-sulfonamide (79.3 mg, 0.450 mmol) was added and the reaction mixture was allowed to stir at 65°C for 1 h. The reaction mixture was filtered and purified via HPLC (1%-99%) ACN:H_2O with a 0.1% HCl modifier to give *N*-[(6-fluoro-2-pyridyl)sulfonyl]-6-phenyl-2-(2,4,6-trimethylphenoxy)pyridine-3-carboxamide (Compound 698) (79 mg, 36%) 1H NMR (400 MHz, DMSO-d6) δ 8.37 (q, J = 7.8 Hz, 1H), 8.16 (dd, J = 7.5, 2.1 Hz, 1H), 8.12 (d, J =7.8 Hz, 1H), 7.80 - 7.73 (m, 3H), 7.59 (dd, J = 8.3, 2.2 Hz, 1H), 7.41 (ddt, J = 5.5, 3.8, 2.2 Hz, 3H), 6.94 (s, 2H), 2.28 (s, 3H), 2.01 (s, 6H). ESI-MS m/z calc. 491.1315, found 492.4 $(M+1)^+$; Retention time: 2.19 minutes.

PREPARATION 150: (*S*)-2,2-dimethyl-4-(methyl-d_3)pyrrolidine-3,3-d_2

Step 1: Methyl-d_3 4-methyl-2-(methyl-d_3)-4-nitropentanoate-3,3-d_2




A 500-mL, three-neck round bottom flask equipped with a magnetic stir bar, a nitrogen line and a J-Kem thermocouple with heating mantle was charged with 2-nitropropane (34.3 g, 385 mmol) and d_8-methyl methacrylate (50 g, 462 mmol) was stirred at ambient temperature when 1,8-diazabicyclo[5.4.0]undec-7-ene (1.47 g, 9.63 mmol) was added in one portion. The reaction solution exothermed from 20 to ~40 °C and was allowed to stir without heating or cooling for 16 h. The reaction mixture was heated at 80 °C for 4 h. The reaction mixture was diluted with MTBE (170 mL), washed with 1 M HCl (15 mL), dried ($MgSO_4$), filtered and concentrated under reduced pressure to afford methyl-d_3 4-methyl-2-(methyl-d_3)-4-nitropentanoate-3,3-d_2 as light yellow oil (75 g, 99%), which was used in the next step without further purification.

Step 2: Methyl-d_3 (*S*)-4-methyl-2-(methyl-d_3)-4-nitropentanoate-3,3-d_2




A 5-L, three-neck round bottom flask equipped an overhead mechanical stirrer, a nitrogen line and a J-Kem thermocouple with heating mantle was charged with methyl-d_3 4-methyl-2-(methyl-d_3)-4-nitropentanoate-3,3-d_2 (75 g, 380 mmol) and pH 7.5 Na-phosphate buffer @ 0.8 M (2000 mL) . To this was added lipase from Rhizomucor miehei (sigma L4277, palatase from Novozymes) (37.5 mL) and stirred at 30 °C for 25 h. The reaction mixture was extracted twice with MTBE (1 L each time). The combined organics were washed with $NaHCO_3$ (5 vol) twice, brine (5 vol), dried over sodium sulfate and concentrated *in vacuo* to afford methyl-d_3 (*S*)-4-methyl-2-(methyl-d_3)-4-nitropentanoate-3,3-d_2 as pale yellow oil (32.5 g, 43% yield).

Step 3: (*S*)-5,5-dimethyl-3-(methyl-d_3)pyrrolidin-2-one-4,4-d_2



A high-pressure vessel (Parr shaker bottle, 500 mL) was purged with and maintained under N_2. The vessel was charged sequentially with DI water-rinsed (3x) damp Raney®2800 Ni (6.1 g), methyl-d3 (*S*)-4-methyl-2-(methyl-d3)-4-nitropentanoate-3,3-d2 (32.5 g, 165 mmol) and EtOH (290 mL). The vessel was sealed and evacuated/backfilled with N_2 (3 x). With no stirring, the vessel was then evacuated and backfilled with H_2 (30 psi). The Parr bottle was shaken while heating the reaction mixture at 60 °C, and the H_2 pressure was maintained at 30 psi for 8 hours. The vessel was evacuated/backfilled with N_2 (3 x) and the catalyst/Celite were removed by vacuum filtration (Celite pad; N_2 blanket). The flask/filter-pad was washed with EtOH (3 x 50 mL). The filtrate and washes were combined and concentrated *in vacuo* to afford (*S*)-5,5-dimethyl-3-(methyl-d_3)pyrrolidin-2-one-4,4-d_2 (20 g, 92%) as white solid.


Step 4: (*S*)-2,2-dimethyl-4-(methyl-d_3)pyrrolidine-3,3-d_2



A 1-L, three-neck round bottom flask equipped an overhead mechanical stirrer, a nitrogen line and a J-Kem thermocouple was charged with LAH pellets (7.6 g, 202 mmol) in THF (80 mL). A solution of (*S*)-5,5-dimethyl-3-(methyl-d_3)pyrrolidin-2-one-4,4-d_2 (20 g, 151 mmol) in THF (120 mL) was added to the suspension over 30 minutes while allowing the reaction temperature to rise to ~60 °C. The reaction temperature was increased to near reflux (~68 °C) and maintained for 16 h. The reaction mixture was cooled to below 40 °C and diluted with MTBE (200 mL). The reaction mixture was quenched carefully with drop-wise addition of a saturated aqueous solution of Na_2SO_4 (1 vol) over 2 h. After the addition was completed, the reaction mixture was cooled to RT. The solid was removed by filtration (Celite pad) and washed with EtOAc (4 vol). With external cooling and N_2 blanket, the filtrate and washings were combined and treated with drop-wise addition of anhydrous 4 M HCl in dioxane (1 equiv.) while maintaining the temperature below 20 °C. After the addition was completed (20 min), the resultant suspension was concentrated under reduced pressure at 45 °C. The suspension was backfilled with heptanes (4 vol) twice during concentration. The suspension was cooled to below 30 °C when the solid was collected by filtration under a N_2 blanket. The solid was dried under high vacuum at 45 °C to afford (*S*)-2,2-dimethyl-4-(methyl-d_3)pyrrolidine-3,3-d_2 (17.5 g, 75%) as a white, crystalline solid.

PREPARATION 151: (6-((propan-2-yl-d7)oxy)pyridin-3-yl)boronic acid

Step 1: 5-bromo-2-((propan-2-yl-d7)oxy)pyridine



To NaH (245 g, 10.2 mmol) slurried in DMF (7 mL) was added d7-IPA (610 mg, 9.09 mmol) and the reaction mixture was stirred for 30 min at room temperature. 5-bromo-2-fluoropyridine (1.00g, 5.68 mmol) was added and the reaction mixture was heated to 120 °C for 2.5 h, then an additional aliquot of NaH (40 mg) and d7-IPA (0.07 mL) was added. After a further 2 h of heating, the reaction mixture was allowed to cool to room temperature. Water (21

mL), followed by MTBE (14 mL) was added, and after stirring, the layers were separated. The organic layer was washed with water (14 mL), dried over sodium sulfate, filtered and concentrated to give 5-bromo-2-((propan-2-yl-d7)oxy)pyridine. 1H NMR (400 mHz) δ 8.17 (m, 1H), 7.60 (dd, J= 8.8, 2.6 Hz, 1H), 6.58 (dd, J= 8.8, 0.7 Hz, 1H).


Step 2: (6-((propan-2-yl-d7)oxy)pyridin-3-yl)boronic acid



A mixture of 5-bromo-2-[1,2,2,2-tetradeuterio-1-(trideuteriomethyl)ethoxy]pyridine (1.24g, 5.00 mmol), bis(pinacolato)diboron (1.52 g, 6.00 mmol), KOAc (1.47 g, 15.0 mmol), $Pd(OAc)_2$ (44.9 mg, 0.20 mmol), X-Phos (119 mg, 0.25 mmol) in 2-MeTHF (11 mL) was heated to 80 °C for 2.5 h under an atmosphere of nitrogen. The reaction mixture was allowed to cool, then water (8 mL) was added and the reaction mixture was filtered through Celite pad washing with 2-MeTHF (2 x 5 mL). The layers were separated and the organic layer dried over sodium sulfate, filtered, and concentrated under reduced pressure. The crude material was dissolved in 2-MeTHF (12 mL), 10% aq w/v NaOH was added and the reaction mixture was stirred. The layers were separated, the aqueous layer was acidified to pH 5 with conc HCl and stirred with 2-MeTHF (12 mL), then separated, the organic layer dried over sodium sulfate, filtered and concentrated under reduced pressure to give (6-((propan-2-yl-d7)oxy)pyridin-3-yl)boronic acid.


PREPARATION 152: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[3-fluoro-5-[2,3,3,3-tetradeuterio-2-(trideuteriomethyl)propoxy]phenyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2708)

Step 1: 1-bromo-3-fluoro-5-[2,3,3,3-tetradeuterio-2-(trideuteriomethyl)propoxy]benzene



To 3-bromo-5-fluoro-phenol (6.00 g, 31.41 mmol) in DMF (60 mL) was slowly added NaH (1.38 g, 34.55 mmol) and the reaction mixture was stirred at ambient temperature for 15 min. To the mixture was added NaI (1.18 g, 7.85 mmol), followed 2-(bromomethyl)-1,1,1,2,3,3,3-heptadeuterio-propane (5.05 g, 34.55 mmol). The reaction mixture was stirred at 80 °C for 6 h. The reaction mixture was cooled to ambient temperature and slowly quenched with water (120 mL). The reaction mixture was extracted with EtOAc (240 mL) and the organic phase separated. The organic phase was washed with brine, dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude light yellow oil was purified by silica gel column chromatography eluting with 0-100% EtOAc/hexanes to give 1-bromo-3-fluoro-5-[2,3,3,3-tetradeuterio-2-(trideuteriomethyl)propoxy]benzene (6.8 g, 85%). 1H NMR (400 MHz, Chloroform-d) δ 6.90 - 6.77 (m, 2H), 6.55 (dt, J = 10.7, 2.2 Hz, 1H), 3.75 - 3.61 (m, 2H).


Step 2: 2-[3-fluoro-5-[2,3,3,3-tetradeuterio-2-(trideuteriomethyl)propoxy]phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane



To the 1-bromo-3-fluoro-5-[2,3,3,3-tetradeuterio-2-(trideuteriomethyl)propoxy]benzene (10.8 g, 42.50 mmol) in dioxane (162 mL) was added bis(dipinacolato)diboron (16.2 g, 63.75 mmol) and KOAc (12.5 g, 127.5 mmol) and the reaction mixture was purged with N_2 for 20 min. To the reaction mixture was added $PdCl_2(dppf)_2.CH_2Cl_2$ (1.74 g, 2.13 mmol) and the reaction mixture was heated to 100 °C for 3 h. The reaction mixture was cooled to ambient temperature and filtered through celite and concentrated *in vacuo* affording a dark oil. The material was purified using silica gel column chromatography eluting with 0-100% EtOAc/hexanes to give 2-[3-fluoro-5-[2,3,3,3-tetradeuterio-2-(trideuteriomethyl)propoxy]phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (8.92 g, 70%). 1H NMR (400 MHz, DMSO-d6) δ 6.99 (dd, J = 2.4, 0.8 Hz, 1H), 6.96 - 6.87 (m, 2H), 3.76 (s, 2H), 1.29 (s, 12H).

Step 3: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[3-fluoro-5-[2,3,3,3-tetradeuterio-2-(trideuteriomethyl)propoxy]phenyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-

carboxamide (Compound 2708)




To *N*-[(6-amino-2-pyridyl)sulfonyl]-6-chloro-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 852) (1.4 g, 3.3 mmol) in 1-propanol (14 mL) and water (4.2 mL) was slowly added K_2CO_3 (1.4 g, 9.9 mmol), followed by 1,3-bis(2,6-diisopropylphenyl)imidazolidin-2-ide;3-chloropyridine;palladium(2+);dichloride (45 mg, 0.06 mmol) and the reaction mixture was heated to 80°C. To the reaction mixture was slowly added 2-[3-fluoro-5-[2,3,3,3-tetradeuterio-2-(trideuteriomethyl)propoxy]phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.2 g, 4.1 mmol) in 1-propanol (2.8 mL). The reaction mixture was stirred at 80°C for 19 h. The reaction mixture was cooled to ambient temperature and slowly added to water (80 mL) containing HCl (20 mL of 1 M, 19.8 mmol). The reaction mixture was diluted with EtOAc (100 mL) and the organic phase separated. The organic phase was washed with brine, dried over MgSO4, filtered over celite and concentrated *in vacuo*. The crude product was purified by silica gel column chromatography eluting with 0-10% MeOH/DCM, then by reverse phase column chromatography eluting with 5-100% ACN/Water to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-[3-fluoro-5-[2,3,3,3-tetradeuterio-2-(trideuteriomethyl)propoxy]phenyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2708) (495 mg, 27%). 1H NMR (400 MHz, DMSO-d6) δ 12.46 (s, 1H), 7.78 (d, J = 8.0 Hz, 1H), 7.63 (dd, J = 8.4, 7.3 Hz, 1H), 7.53 - 7.45 (m, 1H), 7.40 (dt, J = 10.0, 2.0 Hz, 1H), 7.26 (d, J = 8.0 Hz, 1H), 7.20 (d, J = 7.3 Hz, 1H), 6.89 (dt, J = 10.7, 2.3 Hz, 1H), 6.68 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 3.83 (s, 2H), 2.76 - 2.54 (m, 3H), 2.22 (s, 1H), 1.88 (dd, J = 11.9, 5.7 Hz, 1H), 1.60 (d, J = 5.7 Hz, 6H), 1.42 (t, J = 12.1 Hz, 1H), 0.88 (d, J = 6.2 Hz, 3H). ESI-MS m/z calc. 562.2755, found 563.36 (M+1)+; Retention time: 2.73 minutes.

The following compound can be synthesized using the procedures described herein:

N-(benzenesulfonyl)-6-[3-fluoro-5-[2,3,3,3-tetradeuterio-2-(trideuteriomethyl)propoxy]phenyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 2713).


PREPARATION 153: 5-*tert*-butyl-*N*-(2-cyanophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 266)

Step 1: Ethyl 3-(*tert*-butyl)-1-(2,4,6-trimethylbenzyl)-1*H*-pyrazole-5-carboxylate



A mixture of ethyl 5-*tert*-butyl-4*H*-pyrazole-3-carboxylate (0.5 g, 2.5 mmol), 2-(bromomethyl)-1,3,5-trimethyl-benzene (0.5 g, 2.5 mmol), and K_2CO_3 (0.7 g, 5.1 mmol) in DMF (5 mL) was heated to 80 °C overnight. A further aliquot of 2-(bromomethyl)-1,3,5-trimethyl-benzene (200 mg) and K_2CO_3 (0.7 g, 5.1 mmol) were added and the temperature was increased to 85 °C. The reaction mixture was heated overnight. The reaction mixture was cooled, water (20 ml) was added and extracted with ethyl acetate (2 x 50 ml). The organic layers were dried over Na_2SO_4, concentrated *in vacuo*, and purified by silica gel column chromatography.


Step 2: 5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylic acid



A solution of ethyl 5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylate (2.0 g, 6.0 mmol) and sodium hydroxide (1.9 g, 48 mmol) in THF (10 mL) and water (10 mL) was stirred at 60 °C for 17 hours. The reaction mixture was cooled to room temperature and acidified with 1 M hydrochloric acid. The white precipitate was filtered, washed with water, and dried under vacuum to give 5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylic acid (1.7 g, 96%) as a white powder. 1H NMR

(400 MHz, DMSO) δ 6.92 - 6.77 (m, 2H), 6.67 (s, 1H), 5.63 (s, 2H), 2.25 (dd, J = 23.8, 13.5 Hz, 9H), 1.13 (d, J = 10.4 Hz, 9H). ESI-MS m/z calc. 300.18378, found 301.0 $(M+1)^+$; Retention time: 2.06 minutes.

Step 3: 5-*tert*-butyl-*N*-(2-cyanophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 266)



5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxylic acid (30mg, 0.1 mmol), 2-cyanobenzenesulfonamide (18 mg, 0.1 mmol), HATU (38 mg, 0.1 mmol) and DIEA (39 mg, 52 µL, 0.3 mmol) were combined in DMF (1.5 mL) and heated at 60 °C for 16 h. The reaction mixture was filtered and purified by HPLC/MS utilizing a gradient of 1-99% acetonitrile in 5 mM aq HCl to yield 5-*tert*-butyl-*N*-(2-cyanophenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 266) (11 mg, 24%) 1H NMR (400 MHz, DMSO-d6) δ 8.22 (dd, J = 7.9, 1.3 Hz, 1H), 8.12 (dd, J = 7.6, 1.4 Hz, 1H), 7.97 (td, J = 7.8, 1.5 Hz, 1H), 7.90 (td, J = 7.6, 1.3 Hz, 1H), 7.00 (s, 1H), 6.74 (s, 2H), 5.43 (s, 2H), 2.17 (s, 3H), 2.06 (s, 6H), 1.16 (s, 9H). ESI-MS m/z calc. 464.1882, found 465.4 $(M+1)^+$; Retention time: 2.29 minutes.

The following compounds can be synthesized using the procedures described herein:

N-(benzenesulfonyl)-5-*tert*-butyl-2-[1-(2,4,6-trimethylphenyl)ethyl]pyrazole-3-carboxamide (Compound 201),

N-(benzenesulfonyl)-5-*tert*-butyl-2-[[2-fluoro-5-(trifluoromethyl)phenyl]methyl]pyrazole-3-carboxamide (Compound 35),

5-*tert*-butyl-*N*-[3-(dimethylamino)phenyl]sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 270),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trichlorophenyl)methyl]pyrazole-3-carboxamide (Compound 303),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(4-*tert*-butyl-2,6-dimethyl-phenyl)methyl]pyrazole-3-carboxamide (Compound 156),

N-(benzenesulfonyl)-5-[4-(difluoromethyl)phenyl]-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 119),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[[2-chloro-6-(trifluoromethyl)phenyl]methyl]pyrazole-3-carboxamide (Compound 189),

5-*tert*-butyl-*N*-[3-(dimethylamino)-2-methyl-phenyl]sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 104),

N-(3-amino-2-methyl-phenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 123),

5-*tert*-butyl-*N*-(1-methylindol-7-yl)sulfonyl-2-[[2-methyl-3-(trifluoromethyl)phenyl]methyl]pyrazole-3-carboxamide (Compound 276),

5-*tert*-butyl-*N*-(m-tolylsulfonyl)-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 80),

5-*tert*-butyl-*N*-(5-quinolylsulfonyl)-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 352),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,3,5,6-tetramethylphenyl)methyl]pyrazole-3-carboxamide (Compound 317),

5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]-*N*-(2-thienylsulfonyl)pyrazole-3-carboxamide (Compound 77),

N-(benzenesulfonyl)-5-*tert*-butyl-2-[(2,4,6-trichlorophenyl)methyl]pyrazole-3-carboxamide (Compound 145),

5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]-*N*-(2-methoxy-3-methyl-phenyl)sulfonyl-pyrazole-3-carboxamide (Compound 59),

N-(benzenesulfonyl)-5-*tert*-butyl-2-(1-naphthylmethyl)pyrazole-3-carboxamide (Compound 328),

5-*tert*-butyl-*N*-[3-(methylamino)phenyl]sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 249),

5-*tert*-butyl-*N*-[1-(2-hydroxyethyl)indol-4-yl]sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 188),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-4-methyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 20),

5-*tert*-butyl-*N*-(3-hydroxyphenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 84),

5-*tert*-butyl-*N*-(1-methylimidazol-4-yl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 1568),

N-(benzenesulfonyl)-5-*tert*-butyl-2-[(3-pyrrol-1-ylphenyl)methyl]pyrazole-3-carboxamide (Compound 108),

5-*tert*-butyl-*N*-(1H-indol-4-ylsulfonyl)-2-[[2-methyl-3-(trifluoromethyl)phenyl]methyl]pyrazole-3-carboxamide (Compound 60),

N-(3-acetylphenyl)sulfonyl-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 332),

N-(benzenesulfonyl)-5-*tert*-butyl-2-[[5-methyl-2-(trifluoromethyl)phenyl]methyl]pyrazole-3-carboxamide (Compound 304),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[[2-fluoro-6-(trifluoromethyl)phenyl]methyl]pyrazole-3-carboxamide (Compound 239),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[1-(o-tolyl)ethyl]pyrazole-3-carboxamide (Compound 342),

N-[(6-amino-2-pyridyl)sulfonyl]-5-*tert*-butyl-4-methyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 356),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-(1-cyclohexylethyl)pyrazole-3-carboxamide (Compound 287),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-(1-phenylpropyl)pyrazole-3-carboxamide (Compound 165),

5-*tert*-butyl-*N*-(4-hydroxyphenyl)sulfonyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 269),

N-(3-aminophenyl)sulfonyl-5-(1-methylcyclopropyl)-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 158),

N-(benzenesulfonyl)-5-*tert*-butyl-2-[[2-methyl-3-(trifluoromethyl)phenyl]methyl]pyrazole-3-carboxamide (Compound 98),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 185),

N-(3-aminophenyl)sulfonyl-5-isobutyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 167),

N-[(6-amino-2-pyridyl)sulfonyl]-5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 173),

N-(benzenesulfonyl)-5-*tert*-butyl-2-[2-(2-methoxyphenyl)ethyl]pyrazole-3-carboxamide (Compound 157),

5-*tert*-butyl-*N*-[(3-methoxy-2-pyridyl)sulfonyl]-2-[(2,4,6-trimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 44),

N-(3-aminophenyl)sulfonyl-5-*tert*-butyl-2-(1-naphthylmethyl)pyrazole-3-carboxamide (Compound 295),

5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]-*N*-(2-fluorophenyl)sulfonyl-pyrazole-3-carboxamide (Compound 57),

N-(4-aminophenyl)sulfonyl-5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 81),

5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]-*N*-(1H-indazol-4-ylsulfonyl)pyrazole-3-carboxamide (Compound 86),

N-(3-amino-4-fluoro-phenyl)sulfonyl-5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 150),

5-*tert*-butyl-*N*-(2,6-difluorophenyl)sulfonyl-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 170),

5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]-*N*-(1-methylindol-4-yl)sulfonyl-pyrazole-3-carboxamide (Compound 210),

5-*tert*-butyl-*N*-(2,3-dihydrobenzofuran-7-ylsulfonyl)-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 211),

N-(benzenesulfonyl)-5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 257),

N-(3-amino-4-methyl-phenyl)sulfonyl-5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxamide (Compound 312) and

N-(3-aminophenyl)sulfonyl-1-[(2,4,6-trimethylphenyl)methyl]benzimidazole-2-carboxamide (Compound 71).

PREPARATION 154: (3*S*)-3,5,5-trimethylpyrrolidin-2-one


Step 1: Methyl-2,4-dimethyl-4-nitro-pentanoate



THF (4.5 L) was added to a 20 L glass reactor and stirred under an atmosphere of nitrogen at room temperature. 2-Nitropropane (1.5 kg) and DBU (1.282 kg, 0.5 equiv.) were then charged to the reactor (immediate yellow color observed) and the jacket temperature was increased to 50 °C. Once the reactor contents were close to 50 °C, methyl methacrylate (1.854 kg, 1.1 equiv.) was added slowly over 100 minutes. The reaction temperature was maintained at or close to 50 °C with reaction completion reached at 21 h. During the course of the reaction, a color change from yellow to deep green was observed. The reaction mixture was concentrated *in vacuo* then transferred back to the reactor and diluted with MTBE (14 L). 2 M HCl (7.5 L) was added and this was stirred for 5 minutes then allowed to settle. Two clear layers were visible – a lower yellow aqueous phase and an upper green organic phase. The aqueous was removed and the

organic was stirred again with 2 M HCl (3 L). After separation, the HCl washes were recombined and stirred with MTBE (3 L) for 5 minutes. The aqueous was removed and all of the organics were combined in the reactor and stirred with water (3 L) for 5 minutes. After separation, the organics were concentrated *in vacuo* to afford a cloudy green mobile oilm which was dried with $MgSO_4$ and filtered to afford racemic methyl-2,4-dimethyl-4-nitro-pentanoate (3.16 kg) as a clear green mobile oil (99% yield). ^{1}H NMR (400 MHz, Chloroform-*d*) δ 3.68 (s, 3H), 2.56 – 2.35 (m, 2H), 2.11 – 2.00 (m, 1H), 1.56 (d, J = 20.2 Hz, 6H), 1.19 (d, J = 6.8 Hz, 3H).


Step 2: Methyl (2*S*)-2,4-dimethyl-4-nitro-pentanoate



A reaction vessel was set up with methyl 2,4-dimethyl-4-nitro-pentanoate (7 kg, 105.7 mmol) and stirred with 70 L of pH 6.5 K phosphate buffer @ 0.1 M.

To this was added 3.5 L lipase (palatase from Novozymes) and stirred at 34 °C for ~43 h. The pH of the reaction was maintained at pH 6.5 by pH Stat dosing K_2CO_3 (aq. Soln). The reaction mixture was extracted with MTBE (2 x 5 vol, 1 x 2 vol). The combined MTBE layers were then washed with 2 M Na_2CO_3 (4 x 2.5 vol), water (3 vol) and brine (3 vol). The product was concentrated down to ~10 L in the reactor by distilling off the MTBE and transferred to a large rotary evaporator for final concentration to afford methyl (2*S*)-2,4-dimethyl-4-nitro-pentanoate as a mobile yellow oil (3072.2 g, 44% yield).

Step 3: (3*S*)-3,5,5-trimethylpyrrolidin-2-one




A 20 L Hastelloy reactor vessel, purged with and maintained under N_2. The vessel was charged sequentially with DI water-rinsed, damp Raney® Ni (2800 grade, 250 g), methyl (2*S*)-2,4-dimethyl-4-nitro-pentanoate (1741g, 9.2 mol) and ethanol (13.9 L, 8 vol). Stirring was commenced at 900 rpm and the reactor flushed with H_2 and maintained at ~2.5 bar. After 5

hours, the reaction mixture was filtered to remove Raney nickel and solid cake rinsed with ethanol (3.5 L, 2 vol). The ethanolic solution of product was concentrated *in vacuo* to reduce to a minimum volume of ethanol (~1.5 volumes), then heptane (2.5 L) was added and the suspension concentrated again to ~1.5 volumes. This was repeated a further 3 times, then the resulting suspension was cooled to 0-5 °C and filtered under suction and washed with heptane (2.5 L). The product was dried on the filter under vacuum for 20 minutes then transferred to drying trays and dried in a vacuum oven at 40 °C overnight (with N2 bleed). This afforded (3*S*)-3,5,5-trimethylpyrrolidin-2-one as a white crystalline solid (2.042 kg, 87 %). ^{1}H NMR (400 MHz, Chloroform-*d*) δ 6.39 (s, 1H), 2.62 (ddq, J = 9.9, 8.6, 7.1 Hz, 1H), 2.17 (ddd, J = 12.4, 8.6, 0.8 Hz, 1H), 1.56 (dd, J = 12.5, 9.9 Hz, 1H), 1.31 (s, 3H), 1.25 (s, 3H), 1.20 (d, J = 7.1 Hz, 3H).

PREPARATION 155: *N*-((2-aminopyridin-3-yl)sulfonyl)-6'-(((*S*)-1-hydroxypropan-2-yl)oxy)-6-((*S*)-2,2,4-trimethylpyrrolidin-1-yl)-[2,3'-bipyridine]-5-carboxamide (Compound 2690)


Step 1: *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-fluoro-2-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide



A mixture of *N*-[(2-amino-3-pyridyl)sulfonyl]-6-chloro-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (3.8 g, 8.96 mmol), (6-fluoro-2-pyridyl)boronic acid (2.00 g, 14.2 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) complex with dichloromethane (1.40 g, 2.32 mmol), and Na_2CO_3 (4.00 g, 37.7 mmol) in dimethylformamide (80 mL) and water (20 mL) was degassed under a stream of nitrogen and stirred at 100 °C for 18 hours. (6-Fluoro-2-pyridyl)boronic acid (0.90 g, 6.4 mmol), [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II) complex with dichloromethane (0.40 g, 0.55 mmol) were added, and the reaction mixture stirred at 100°C for another day. The reaction mixture was filtered through a pad of Celite and the filtrate was diluted with water (300 mL), adjusted to pH ~7 and extracted with ethyl acetate (5x300 mL). The extract was washed with brine (2x200 mL), dried over Na_2SO_4, and concentrated. The residue purified by chromatography on silica gel, eluted with 0-100% ethyl acetate in hexanes, affording yellow


foam product (2.2 g, ~85% purity) and starting material (3) (560 mg).

Step 2: *N*-((2-aminopyridin-3-yl)sulfonyl)-6'-(((*S*)-1-(benzyloxy)propan-2-yl)oxy)-6-((*S*)-2,2,4-trimethyl-pyrrolidin-1-yl)-[2,3'-bipyridine]-5-carboxamide



Sodium hydride (44 mg, 1.1 mmol) (60 wt% in mineral oil) was added to a solution of (*R*)-1-benzyloxy-propan-2-ol (117 mg, 0.7 mmol) in dimethylformamide (2.0 mL). After 5 minutes of stirring at room temperature, *N*-[(2-amino-3-pyridyl)sulfonyl]-6-(6-fluoro-2-pyridyl)-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (108 mg, 0.18 mmol) was added. The reaction mixture was heated under microwave irradiation at 75 °C for 30 minutes. The reaction mixture was quenched with water (10 mL), adjusted pH 3-4, and extracted with ethyl acetate (2x10 mL). The combined extract was washed with brine (2x5 mL), dried over Na_2SO_4 and concentrated under reduced pressure to give *N*-((2-aminopyridin-3-yl)sulfonyl)-6'-(((*S*)-1-(benzyloxy)propan-2-yl)oxy)-6-((*S*)-2,2,4-trimethyl-pyrrolidin-1-yl)-[2,3'-bipyridine]-5-carboxamide. ESI-MS m/z 631 $(M+1)^+$.

Step 3: *N*-((2-aminopyridin-3-yl)sulfonyl)-6'-(((*S*)-1-hydroxypropan-2-yl)oxy)-6-((*S*)-2,2,4-trimethylpyrrolidin-1-yl)-[2,3'-bipyridine]-5-carboxamide (Compound 2690)




N-((2-aminopyridin-3-yl)sulfonyl)-6'-(((*S*)-1-(benzyloxy)propan-2-yl)oxy)-6-((*S*)-2,2,4-trimethyl-pyrrolidin-1-yl)-[2,3'-bipyridine]-5-carboxamide was hydrogenated under 1 atmosphere of hydrogen with Pd/C (10%, 260 mg) in methanol (25 mL) at room temperature. After 4 hours, the catalyst was removed by filtration, evaporated under reduced pressure and purified by silica gel column chromatography (0-10% of methanol in dichloromethane), followed by a further purification by silica gel column chromatography eluted with 0-8% of methanol in a mixture solvent of ethyl acetate and hexanes in a ratio of 50/50 to give *N*-((2-

aminopyridin-3-yl)sulfonyl)-6'-(((*S*)-1-hydroxypropan-2-yl)oxy)-6-((*S*)-2,2,4-trimethylpyrrolidin-1-yl)-[2,3'-bipyridine]-5-carboxamide (Compound 2690) (9 mg). ESI-MS m/z 541 $(M+1)^+$.


PREPARATION 156: Methyl 2-(benzenesulfonyl)-3-[5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazol-3-yl]-3-oxo-propanoate (Compound 4)

Step1: 5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carbonyl chloride



5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carboxylic acid (100 mg, 0.3492 mmol) was dissolved in thionyl chloride (41.54 mg, 25.47 µL, 0.3492 mmol) and a drop of DMF was added. The reaction was stirred for 1h and evaporated to give the acid chloride, which was used without further purification.


Step 2: Methyl 2-(benzenesulfonyl)-3-[5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazol-3-yl]-3-oxo-propanoate (Compound 4)



To 5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazole-3-carbonyl chloride (40 mg, 0.13 mmol), methyl 2-(benzenesulfonyl)acetate (28 mg, 22 uL, 0.13 mmol)and NaH (5 mg, 0.13 mmol) in DMF were stirred at room temperature for 15 minutes. The reactions were filtered and purified by reverse phase LC-MS (10-99% CH_3CN/H_2O, HCl modifier) to give methyl 2-(benzenesulfonyl)-3-[5-*tert*-butyl-2-[(2,6-dimethylphenyl)methyl]pyrazol-3-yl]-3-oxo-propanoate (Compound 4) (18 mg, 28%). ESI-MS m/z 483.4 $(M+1)^+$; Retention time: 2.41 minutes.

PREPARATION 157: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-[(*E*)-2-(4-methoxyphenyl)vinyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1535)

Step 1: 5-Iodo-6-*tert*-butyl-2-hydroxypyridine-3-carbonitrile



N-Iodosuccinimide (31.9 g, 142 mmol) was added to a solution of 6-*tert*-butyl-2-hydroxypyridine-3-carbonitrile (10.0 g, 56.8 mmol) in a mixture of 1,2-dichloroethane (120 mL) and trifluoroacetic acid (40 mL) and the reaction mixture was heated at 50 °C overnight. The solution was concentrated under reduced pressure and ethyl acetate (300 mL) was added. The organic layer was washed with 10% sodium thiosulfate solution (300 mL), water (2 x 200 mL) and brine (150 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by flash chromatography, eluting from 30% EtOAc in heptanes to 60% EtOAc in heptanes, to provide 5-iodo-6-*tert*-butyl-2-hydroxypyridine-3-carbonitrile (15.85 g, 92% yield) as a white solid. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.57 (s, 9 H), 8.17 (s, 1 H). $[M+H]^+ = 303.0$.

Step 2: 5-Iodo-6-*tert*-butyl-2-chloropyridine-3-carbonitrile



Phosphorus oxychloride (67.0 mL, 718 mmol) and phosphorus pentachloride (44.9 g, 216 mmol) were added to 5-iodo-6-*tert*-butyl-2-hydroxypyridine-3-carbonitrile (21.7 g, 71.8 mmol) and the reaction mixture was refluxed overnight. The reaction mixture was cooled to room temperature and concentrated under reduced pressure to remove most of the remaining phosphorus oxychloride. Water (***caution***: exotherm) followed by 50% NaOH solution were added until a pH ~9 was reached and the aqueous layer was extracted with ethyl acetate (3 x 100 mL). The combined organic layers were washed with brine (100 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to provide 5-iodo-6-*tert*-butyl-2-chloroxypyridine-3-carbonitrile (17.35 g, 75% yield) as a yellow solid. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.55 (s, 9 H), 8.38 (s, 1 H). $[M+H]^+ = 320.9$.


Step 3: 5-Iodo-6-*tert*-butyl-2-chloropyridine-3-carboxylic acid



Potassium hydroxide (18.2 g, 325 mmol) in water (60 mL) was added to 5-iodo-6-*tert*-butyl-2-chloroxypyridine-3-carbonitrile (17.35 g, 54.1 mmol) in 2-propanol (60 mL) and the


reaction mixture was heated at 90 °C overnight. Ethyl acetate (50 mL) was added and the organic layer was extracted with 1 N NaOH (3 x 50 mL). The basic aqueous layers were combined and concentrated under reduced pressure to remove most of the 2-propanol. The remaining aqueous layer was acidified to pH 1-2 with concentrated HCl and extracted with ethyl acetate (3 x 100 mL). The organic layers were combined, washed with brine (100 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by reverse-phase chromatography (methanol/water, 0% to 100%) to afford 5-iodo-6-*tert*-butyl-2-chloropyridine-3-carboxylic acid (8.2 g, 45% yield) as an off-white solid. ^{1}H NMR (300 MHz, $CDCl_3$) ppm 1.57 (s, 9 H), 8.77 (s, 1 H). $[M+H]^+$ = 339.9.

Step 4: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-chloro-5-iodo-pyridine-3-carboxamide




To a stirred solution of 6-tert-butyl-2-chloro-5-iodo-pyridine-3-carboxylic acid (1.57 g, 4.62 mmol) in dry DMF (11.0 mL) added CDI (0.94 g, 5.78 mmol) in one portion under nitrogen at ambient temperature. The reaction was warmed to 45 °C and stirred for 1.5 h. In a separate flask, 6-aminopyridine-2-sulfonamide (1.00 g, 5.78 mmol) was dissolved in dry DMF (9.0 mL) under nitrogen and sodium hydride (60 % in mineral oil) (231 mg, 5.78 mmol) was added in portions to mitigate gas evolution at ambient temperature. The heterogeneous mixture was warmed to 40 °C and stirred for 1 hour. This reaction mixture was added to the stirred activated acid mixture in one portion at 45 °C. Heat was removed and the reaction was stirred for an additional hour. The reaction mixture was slowly added to ice-cold water (160 mL) then AcOH (1.43 g, 1.35 mL, 23.81 mmol) was added and the reaction mixture was stirred vigorously for 10 mins and the solids were filtered and washed with water (3 x 20 mL). The solids were slurried in hexanes (50 mL) and filtered. The solid was further dried to furnish *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-chloro-5-iodo-pyridine-3-carboxamide (2.11 g, 92%) as white solid.

Step 5: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-iodo-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide



To a solution of *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-chloro-5-iodo-pyridine-3-carboxamide (2.11 g, 4.27 mmol) in anhydrous DMSO (10 mL), (4S)-2,2,4-trimethylpyrrolidine (1.01 g, 8.96 mmol) and K_2CO_3 (2.95 g, 21.32 mmol) were added in that order. The reaction mixture was heated at 150 °C for 18 h under an atmosphere of nitrogen. The reaction mixture was allowed to cool to ambient temperature, then poured onto ice-water (125 mL) with stirring. The pH was adjusted to about 6.0 to 7.0 with the addition of 1 M HCl, the solid was filtered, washed with water (4 x 20 mL) and dried under vacuum to yield *N*-[(6-amino-2-pyridyl)sulfonyl]-6-tert-butyl-5-iodo-2-[(4S)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (2.15 g, 88%) as off-white solid.

Step 6: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-[(*E*)-2-(4-methoxyphenyl)vinyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1535)



DMF (0.5 mL) and H_2O (0.1 mL) were added to *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-iodo-2-(2,2,4-trimethylpyrrolidin-1-yl)pyridine-3-carboxamide (30 mg, 0.05 mmol) and [(*E*)-

2-(4-methoxyphenyl)vinyl]boronic acid (19 mg, 0.11 mmol), followed by K_2CO_3 (29 mg, 0.21 mmol) under an atmosphere of nitrogen. Then tetrakis(triphenylphosphine)palladium (0) (6 mg, 0.005 mmol) was added, the reaction mixture was sealed under an atmosphere of nitrogen and heated at 130 °C for 13 h. The reaction mixture was allowed to cool to ambient temperature and the product was purified using a reverse phase HPLC-MS method using a Luna C18 (2) column (75 × 30 mm, 5 μm particle size) and a dual gradient run from 10-99% mobile phase B over 30.0 minutes (mobile phase A = H_2O (5 mM HCl), mobile phase B = CH_3CN) to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-[(*E*)-2-(4-methoxyphenyl)vinyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1535) (5 mg, 16%). ESI-MS m/z 578.6 $(M+1)^+$; Retention time: 1.78 minutes; 1H NMR (400 MHz, Methanol-d4) δ 8.26 (s, 1H), 7.68 (dd, J = 8.5, 3.4 Hz 1H), 7.48 (d, J = 8.7 Hz, 2H), 7.43 (d, J = 15.9 Hz 1H), 7.37 (d, J = 7.3 Hz, 1H), 6.94 (d, J = 8.8 Hz, 2H), 6.87 (d, J = 15.9 Hz, 1H), 6.77 (d, J = 8.4 Hz, 1H), 3.82 (s, 3H), 3.48 (dd, J = 10.9, 8.2 Hz, 1H), 3.17 (dd, J = 11.1, 9.3 Hz, 1H), 2.69 - 2.52 (m, 1H), 2.11 (dd, J = 12.4, 6.8 Hz, 1H), 1.74 (t, J = 11.8 Hz, 1H), 1.54 (s, 3H), 1.49 (s, 9H), 1.46 (s, 3H), 1.13 (d, J = 6.5 Hz, 3H).

The following compounds can be synthesized using the procedures described herein:

N-[(6-amino-2-pyridyl)sulfonyl]-5-(3-bicyclo[2.2.1]hept-2-enyl)-6-*tert*-butyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1577),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-[(*E*)-pent-1-enyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1617),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-[(*E*)-2-cyclopropylvinyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1665),


Ethyl (*E*)-3-[5-[(6-amino-2-pyridyl)sulfonylcarbamoyl]-2-*tert*-butyl-6-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]-3-pyridyl]prop-2-enoate (Compound 1490),

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]-5-vinyl-pyridine-3-carboxamide (Compound 1373) and

N-[(6-amino-2-pyridyl)sulfonyl]-6-*tert*-butyl-5-[(*E*)-2-ethoxyvinyl]-2-[(4*S*)-2,2,4-trimethylpyrrolidin-1-yl]pyridine-3-carboxamide (Compound 1626)


PREPARATION 158: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(4-phenylbutoxy)pyridine-3-carboxamide (Compound 1686)

Step 1: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-fluoro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide



N-[(6-amino-2-pyridyl)sulfonyl]-2-chloro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (300 mg, 0.63 mmol) was dissolved in DMSO (1.2 mL). Cesium fluoride (476 mg, 3.13 mmol) was added. The reaction mixture was sealed and stirred overnight at 150 °C. Additional cesium fluoride (100 mg) was added, and the reaction mixture was stirred overnight at 150 °C. After cooling to room temperature, the mixture was diluted with EtOAc (75 mL) and washed with diluted aqueous HCl solution (0.1 N, 1× 75 mL). The organic layer was dried over sodium sulfate, filtered and concentrated under reduced pressure. The product was used in the next step without further purification. ESI-MS m/z 463.2 $(M+1)^+$; Retention time: 1.83 minutes.

Step 2: *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(4-phenylbutoxy)pyridine-3-carboxamide (Compound 1686)



4-phenylbutan-1-ol (23 mg, 0.15 mmol) was dissolved in dioxane (250 µL).

Sodium hydride (6 mg, 0.15 mmol) was added, and the reaction mixture was stirred at room temperature for 5 minutes. A solution of *N*-[(6-amino-2-pyridyl)sulfonyl]-2-fluoro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (35 mg, 0.08 mmol) in DMSO (250 µL) was added. After another 5 minutes of stirring, the reaction mixture was heated under microwave irradiation at 100 °C for 30 minutes, then concentrated under reduced pressure and purified by reverse phase HPLC-MS method using a Luna C18 (2) column (75 × 30 mm, 5 µm particle size) and a dual gradient run from 1-99% mobile phase B over 15.0 minutes, (mobile phase A = water (5 mM HCl), mobile phase B = acetonitrile) to give *N*-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(4-phenylbutoxy)pyridine-3-carboxamide (Compound 1686) (30 mg, 64%). ESI-MS m/z 593.4 $(M+1)^+$; Retention time: 2.47 minutes.

The following compounds can be synthesized using the procedures described herein:

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3-phenylbutoxy)pyridine-3-carboxamide (Compound 1689),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3-phenylpropoxy)pyridine-3-carboxamide (Compound 1694),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[4-(3-pyridyl)butoxy]pyridine-3-carboxamide (Compound 1690),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[4-(4-pyridyl)butoxy]pyridine-3-carboxamide (Compound 1687),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[4-(2-pyridyl)butoxy]pyridine-3-carboxamide (Compound 1688),


N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[3-(2-pyridyl)propoxy]pyridine-3-carboxamide (Compound 1693),

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-[3-(3-pyridyl)propoxy]pyridine-3-carboxamide (Compound 1691) and

N-[(6-amino-2-pyridyl)sulfonyl]-6-(3-fluoro-5-*iso*butoxy-phenyl)-2-(3-pyrazol-1-ylpropoxy)pyridine-3-carboxamide (Compound 1692).

PREPARATION 159: *N*-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-3-oxo-

pyrrolidin-1-yl)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 1189)




Solid potassium carbonate (90 mg, 0.65 mmol) was added to a suspension of 2,2-dimethylpyrrolidin-3-one (49 mg, 0.32 mmol) in DMSO (250 µL), followed by *N*-[(6-amino-2-pyridyl)sulfonyl]-2-fluoro-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (HCl salt) (50 mg, 0.11 mmol). The reaction mixture was stirred at 150 °C for 3 hours. The reaction mixture was purified using a reverse phase HPLC method using a Luna C18 (2) column (50 × 21.2 mm, 5 µm particle size) and a dual gradient run from 10-99% mobile phase B over 15.0 minutes (mobile phase A = water (5 mM HCl), mobile phase B = acetonitrile). ESI-MS m/z 556.4 $(M+1)^+$; Retention time: 2.01 minutes.

The following compound can be synthesized using the procedures described herein:

N-[(6-amino-2-pyridyl)sulfonyl]-2-(2,2-dimethyl-4-oxo-pyrrolidin-1-yl)-6-(3-fluoro-5-*iso*butoxy-phenyl)pyridine-3-carboxamide (Compound 1073).

PREPARATION 160: 2-(benzenesulfonyl)-1-[5-*tert*-butyl-2-[(2,4,6-trimethylphenyl)methyl]pyrazol-3-yl]propan-1-one (Compound 139)



To a solution of 2-(benzenesulfonyl)-1-{3-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazol-5-yl}ethan-1-one (50 mg, 0.11 mmol) in THF (3.0 mL) and

DMF (1.0 mL) was added DBU (52 mg, 0.34 mmol) and the mixture was stirred at room temperature for 30 minutes. Iodomethane (40 mg, 0.28 mmol) was then added and the reaction was stirred overnight at room temperature. Analysis of an aliquot by LCMS indicated that only mono-methylation (no bis-methylation) had occurred. The crude reaction mixture was transferred to a separatory funnel with water and extracted twice with diethyl ether. The organic layers were combined, dried over sodium sulfate, filtered and concentrated under reduced pressure to give the crude product. In a separate experiment, a solution of 2-(benzenesulfonyl)-1-{3-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazol-5-yl}ethan-1-one (50 mg, 0.11 mmol) in THF (3.0 mL) and DMF (1.0 mL) was added DBU (52 mg, 0.34 mmol) and the mixture was heated in an oil bath at 80 °C for 10 minutes. Iodomethane (40 mg, 0.28 mmol) was then added, the vial was capped and the reaction was heated in an oil bath at 80 °C overnight. Analysis of an aliquot by LCMS indicated that only mono-methylation (no bis-methylation) had occurred. The crude reaction mixture was transferred to a separatory funnel with water and extracted twice with diethyl ether. The organic layers were combined, dried over sodium sulfate, filtered and concentrated under reduced pressure to give the crude product. Both crude reaction mixtures were combined together and purified by silica-gel column chromatography, eluting with 5% to 20% EtOAc in heptanes, to afford *rac* 2-(benzenesulfonyl)-1-{3-*tert*-butyl-1-[(2,4,6-trimethylphenyl)methyl]-1*H*-pyrazol-5-ylpropan-1-one (64 mg, 60% yield) as a white solid. ^{1}H NMR (300 MHz, DMSO-d_6) ppm 1.10 (s, 9 H), 1.46 (d, J = 6.8 Hz, 3 H), 2.16 (s, 6 H), 2.21 (s, 3 H), 5.40 - 5.53 (m, 3 H), 6.83 (s, 2 H), 7.15 (s, 1 H), 7.55 - 7.62 (m, 2 H), 7.68 - 7.76 (m, 3 H). ESI-MS m/z 453.2 $(M+1)^+$.

Table 2 below recites the analytical data for the compounds of Table 1.

Table 2.

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1	503.00	1.96	
2	419.00	2.04	
3	484.05	2.00	
4	483.40	2.41	
5	438.40	1.74	
6	435.20	1.50	
7	469.40	2.06	1H NMR (400 MHz, DMSO-d6) δ 12.19 (s, 1H), 7.24 (t, J = 7.9 Hz, 1H), 7.17 (t, J = 2.1 Hz, 1H), 7.09 - 7.03 (m,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			2H), 6.83 (ddd, J = 8.1, 2.3, 1.0 Hz, 1H), 6.72 (s, 2H), 5.68 (s, 2H), 4.42 - 4.35 (m, 2H), 2.84 - 2.76 (m, 2H), 2.15 (s, 3H), 2.08 (s, 6H), 1.23 (s, 9H).
8	561.40	2.30	
9	482.10	1.98	
10	467.10	2.18	
11	468.00	2.14	1H NMR (400 MHz, DMSO) δ 11.65 (s, 1H), 7.92 (s, 1H), 7.63 (d, J = 8.8 Hz, 2H), 7.11 (d, J = 7.8 Hz, 1H), 6.91 (s, 2H), 6.62 (d, J = 11.3 Hz, 2H), 6.17 (s, 2H), 2.26 (s, 3H), 1.95 (s, 6H), 1.04 (s, 9H).
12	447.20	2.14	
13	426.40	1.80	
14	561.14	2.91	
15	481.00	2.15	1H NMR (400 MHz, DMSO) δ 12.05 (s, 1H), 7.69 (s, 1H), 7.62 (s, 1H), 7.19 (d, J = 7.3 Hz, 1H), 7.05 (d, J = 26.9 Hz, 3H), 6.69 (d, J = 8.4 Hz, 1H), 6.50 (s, 2H), 2.69 (s, 2H), 2.29 (s, 3H), 2.04 (d, J = 25.4 Hz, 3H), 1.67 (d, J = 28.9 Hz, 4H), 0.98 (s, 6H).
16	465.30	1.52	
17	519.00	1.93	1H NMR (400 MHz, DMSO) δ 8.10 (d, J = 7.9 Hz, 1H), 7.33 (s, 2H), 7.20 (s, 3H), 7.07 (s, 2H), 6.94 (s, 4H), 6.71 (d, J = 8.3 Hz, 1H), 6.52 (s, 2H).
18	464.00	1.97	1H NMR (400 MHz, DMSO) δ 12.71 (s, 1H), 8.47 (d, J = 6.8 Hz, 1H), 8.28 (d, J = 9.5 Hz, 1H), 7.70 (d, J = 7.9 Hz, 1H), 7.55 (s, 1H), 7.25 (s, 1H), 7.15 (s, 1H), 7.07 (s, 1H), 7.01 (d, J = 8.4 Hz, 1H), 5.62 (s, 2H), 3.92 (s, 3H), 2.18 (s, 3H), 1.84 (s, 6H).
19	467.20	2.83	1H NMR (400 MHz, Chloroform-d) δ 8.27 (d, J = 7.9 Hz, 1H), 7.49 - 7.33 (m, 2H), 7.23 (t, J = 7.9 Hz, 1H), 6.92 (d, J = 7.9 Hz, 1H), 6.88 - 6.74 (m, 3H), 3.89 (s, 2H), 2.29 (s, 3H), 2.02 (s, 6H), 1.06 (s, 9H).
20	469.40	2.04	
21	532.30	2.08	
22	451.00	2.44	
23	449.00	1.54	1H NMR (400 MHz, DMSO) δ 8.02 (s,

Cmpd No	LC/MS M+1	LC/RT min	NMR
			1H), 7.60 (s, 1H), 7.25 (s, 1H), 7.18 (s, 1H), 6.98 (s, 2H), 6.78 (s, 4H), 2.17 (s, 3H), 1.97 (s, 6H).
24	468.20	1.92	
25	448.20	1.91	
26	452.50	1.81	
27	494.07	2.08	
28	480.30	2.20	
29	490.00	2.16	
30	478.34	1.45	
31	425.40	1.64	
32	451.00	0.90	1H NMR (400 MHz, MeOD) δ 7.99 (d, J = 7.0 Hz, 1H), 7.64 (d, J = 6.2 Hz, 1H), 7.44 - 7.26 (m, 4H), 6.92 (s, 2H), 6.86 (d, J = 9.5 Hz, 1H), 5.36 (s, 1H), 4.69 (d, J = 20.6 Hz, 3H), 2.44 (s, 6H), 2.25 (s, 3H).
33	451.20	2.07	
34	484.00	2.30	1H NMR (400 MHz, DMSO) δ 12.52 (s, 1H), 8.04 (s, 1H), 7.98 (d, J = 7.8 Hz, 1H), 7.76 (d, J = 7.3 Hz, 1H), 7.15 (dd, J = 12.9, 8.1 Hz, 2H), 6.90 (s, 2H), 3.85 (s, 3H), 2.25 (s, 3H), 1.94 (s, 6H), 1.05 (s, 9H).
35	484.05	1.97	
36	464.20	1.82	
37	510.00	2.13	
38	448.30	1.90	
39	470.00	2.12	1H NMR (400 MHz, DMSO) δ 12.51 (s, 1H), 7.99 (s, 1H), 7.87 (d, J = 7.7 Hz, 1H), 7.63 (dt, J = 15.3, 7.7 Hz, 2H), 7.01 (s, 1H), 6.75 (s, 2H), 5.42 (s, 2H), 4.63 (s, 2H), 2.16 (s, 3H), 2.07 (d, J = 6.4 Hz, 6H), 1.14 (s, 9H).
40	477.25	1.86	
41	472.28	2.06	
42	505.40	1.92	
43	494.07	2.02	
44	471.30	2.28	
45	440.40	2.25	1H NMR (400 MHz, Chloroform-d) δ 8.33 (s, 1H), 8.17 - 8.11 (m, 2H), 7.72 - 7.65 (m, 1H), 7.58 (dd, J = 8.5, 7.1 Hz, 2H), 6.74 (s, 2H), 6.45 (s, 1H), 5.51 (s, 2H), 2.22 (s, 3H), 2.12 (s, 6H), 1.17 (s, 9H).

Cmpd No.	LC/MS M+1	LC/RT min	NMR
46	428.17	1.77	
47	469.00	2.16	1H NMR (400 MHz, DMSO) δ 12.16 (s, 1H), 10.19 (s, 1H), 7.94 (d, J = 7.8 Hz, 1H), 7.42 (d, J = 4.7 Hz, 3H), 7.14 - 7.04 (m, 2H), 6.89 (s, 2H), 2.25 (s, 3H), 1.93 (s, 6H), 1.04 (s, 9H).
48	436.00	1.20	1H NMR (400 MHz, DMSO) δ 10.14 (s, 1H), 8.59 (s, 1H), 7.97 (s, 2H), 7.56 (d, J = 23.1 Hz, 2H), 7.32 (d, J = 21.5 Hz, 3H), 6.89 (s, 2H), 5.27 (s, 1H), 4.49 (d, J = 13.2 Hz, 1H), 4.31 (s, 3H), 2.35 (s, 6H), 2.23 (s, 3H).
49	476.30	2.05	
50	437.40	2.11	
51	470.00	1.94	1H NMR (400 MHz, DMSO) δ 12.58 (s, 1H), 11.83 (s, 1H), 8.28 (s, 1H), 8.07 (s, 1H), 7.84 (s, 1H), 7.16 (s, 1H), 6.92 (s, 1H), 6.46 (s, 1H), 2.26 (s, 3H), 1.98 (s, 6H), 1.04 (s, 9H).
52	438.22	1.88	
53	467.30	2.84	1H NMR (400 MHz, Chloroform-d) δ 8.27 (d, J = 7.9 Hz, 1H), 7.49 - 7.36 (m, 2H), 7.22 (t, J = 7.9 Hz, 1H), 6.92 (d, J = 7.8 Hz, 1H), 6.87 - 6.75 (m, 3H), 3.89 (s, 2H), 2.29 (s, 3H), 2.02 (s, 6H), 1.06 (s, 9H).
54	521.00	1.81	
55	456.40	1.91	1H NMR (400 MHz, DMSO-d6) δ 12.55 (s, 1H), 12.03 (s, 1H), 8.25 (d, J = 7.5 Hz, 1H), 7.99 (d, J = 7.8 Hz, 1H), 7.83 (s, 1H), 7.16 (d, J = 7.9 Hz, 1H), 7.10 (s, 1H), 7.04 (s, 2H), 6.45 (t, J = 6.8 Hz, 1H), 2.30 (s, 3H), 2.02 (s, 3H), 1.09 (s, 9H).
56	483.00	2.34	1H NMR (400 MHz, DMSO) δ 12.23 (s, 1H), 7.95 (s, 1H), 7.54 (d, J = 36.2 Hz, 3H), 7.30 (s, 1H), 7.12 (s, 1H), 6.90 (s, 2H), 3.80 (s, 3H), 2.25 (s, 3H), 1.92 (s, 6H), 1.04 (s, 9H).
57	444.38	3.14	
58	425.40	1.63	
59	470.30	2.27	
60	519.00	1.98	1H NMR (400 MHz, DMSO) δ 12.51 (s, 1H), 11.67 (s, 1H), 7.74 (s, 1H), 7.66 (s, 1H), 7.56 (s, 1H), 7.43 (s, 1H), 7.28

Cmpd No.	LC/MS M+1	LC/RT min	NMR
			(d, J = 21.0 Hz, 3H), 6.86 (s, 1H), 6.55 (s, 1H), 5.51 (s, 2H), 2.20 (s, 3H), 2.07 (s, 2H), 1.24 (s, 9H).
61	533.22	3.09	
62	462.40	1.91	
63	440.00	2.11	1H NMR (400 MHz, DMSO-d6) δ 11.90 (s, 1H), 8.57 (s, 1H), 8.05 - 7.96 (m, 2H), 7.77 - 7.70 (m, 1H), 7.70 - 7.61 (m, 2H), 6.73 (s, 2H), 3.98 (s, 2H), 2.17 (s, 3H), 2.07 (s, 6H), 1.38 (s, 9H).
64	533.22	2.85	1H NMR (400 MHz, DMSO) δ 12.52 (s, 1H), 9.02 (s, 2H), 8.22 - 8.19 (d, 1H), 7.89 - 7.86 (d, 1H), 7.66 (t, J = 7.91, 1H), 7.25 - 7.22 (d, 1H), 6.96 (s, 2H), 6.74 - 7.61 (d, 1H), 6.59 (s, 2H), 3.22 - 3.08 (m, 1H), 2.2 9 (s, 3H), 2.03 (s, 6H), 1.27 - 1.24 (d, 6H)
65	468.20	1.93	
66	533.22	2.90	
67	474.50	1.70	1H NMR (400 MHz, DMSO-d6) δ 7.61 (dd, J = 8.4, 7.3 Hz, 1H), 7.48 (s, 2H), 7.38 (s, 2H), 7.11 (d, J = 7.3 Hz, 1H), 7.02 (dd, J = 8.9, 1.7 Hz, 1H), 6.88 (d, J = 8.7 Hz, 1H), 6.69 (d, J = 8.4 Hz, 1H), 6.46 (s, 2H),, 3.18 (s, 1H), 2.34 (s, 3H), 1.97 (d, J = 15.9 Hz, 6H).
68	469.40	2.04	1H NMR (400 MHz, DMSO-d6) δ 12.39 (s, 1H), 8.78 (s, 1H), 7.30 - 7.21 (m, 2H), 7.12 (dd, J = 7.4, 1.8 Hz, 1H), 6.93 (s, 2H), 6.86 (dd, J = 7.9, 2.2 Hz, 1H), 2.26 (s, 3H), 1.94 (s, 6H), 1.11 (s, 9H); NH2 not observed.
69	533.22	2.89	
70	477.40	2.03	
71	449.20	1.41	
72	582.20	2.26	
73	451.00	1.07	1H NMR (400 MHz, CD3CN) δ 7.70 (s, 1H), 7.37 (d, J = 26.5 Hz, 4H), 7.18 (d, J = 7.3 Hz, 1H), 6.94 (s, 2H), 6.84 (d, J = 8.6 Hz, 1H), 5.06 (s, 1H), 4.62 (d, J = 14.0 Hz, 1H), 4.45 (s, 2H), 4.37 (d, J = 14.0 Hz, 1H), 2.43 (s, 6H), 2.27 (s, 3H).
74	409.30	1.93	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
75	440.40	1.91	
76	489.40	1.58	
77	432.04	2.04	
78	478.00	2.26	1H NMR (400 MHz, DMSO) δ 8.27 (d, J = 7.9 Hz, 1H), 8.14 (d, J = 7.5 Hz, 1H), 8.00 (s, 2H), 7.91 (s, 1H), 7.14 (d, J = 7.8 Hz, 1H), 6.89 (s, 2H), 2.25 (s, 3H), 1.92 (s, 6H), 1.04 (s, 9H).
79	465.00	1.57	1H NMR (400 MHz, DMSO) δ 7.84 (d, J = 7.9 Hz, 1H), 7.60 (s, 1H), 7.33 (s, 4H), 7.16 (d, J = 8.4 Hz, 1H), 7.07 (s, 1H), 6.87 (s, 2H), 5.14 (s, 1H), 4.25 (d, J = 39.6 Hz, 4H), 3.86 (s, 3H), 2.31 (s, 6H), 2.22 (s, 3H).
80	454.00	2.31	1H NMR (400 MHz, DMSO) δ 7.69 (s, 1H), 7.43 (s, 1H), 7.30 (s, 1H), 7.17 (s, 1H), 7.04 (s, 1H), 6.68 (s, 2H), 5.40 (s, 2H), 2.35 (s, 3H), 2.10 (s, 3H), 2.02 (s, 6H), 1.07 (s, 9H).
81	441.38	2.84	1H NMR (400 MHz, DMSO) δ 12.09 (s, 1H), 7.63 - 7.60 (d, 2H), 7.07 - 6.92 (m, 4H), 6.64 - 6.61 (d, 2H), 6.21 (s, 2H), 5.48 (s, 2H), 2.13 (s, 6H), 1.12 (s, 9H)
82	468.20	1.91	
83	480.30	2.19	
84	456.40	2.06	1H NMR (400 MHz, DMSO-d6) δ 12.47 (s, 1H), 10.23 (s, 1H), 7.50 - 7.39 (m, 3H), 7.12 (dt, J = 8.0, 1.8 Hz, 1H), 7.01 (s, 1H), 6.77 (s, 2H), 5.46 (s, 2H), 2.19 (s, 3H), 2.10 (s, 6H), 1.16 (s, 9H).
85	439.30	1.85	
86	466.30	2.93	1H NMR (400 MHz, DMSO) δ 8.43 (s, 1H), 7.99 - 7.96 (d, 1H), 7.87 - 7.84 (d, 1H), 7.60 (t, J = 7.8, 1H), 7.04 - 6.94 (m, 2H), 6.86 - 6.83 (m, 2H), 5.35 (s, 2H), 1.92 (s, 6H), 1.14 (s, 9H)
87	451.20	2.09	
88	441.08	3.03	
89	470.40	2.27	
90	450.20	1.19	
91	463.00	1.76	1H NMR (400 MHz, DMSO) δ 12.66 (s, 1H), 8.63 (s, 1H), 8.08 (d, J = 7.8 Hz, 1H), 7.67 (d, J = 20.4 Hz, 2H), 7.54 (d,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			J = 14.6 Hz, 2H), 7.23 (d, J = 7.3 Hz, 1H), 6.94 (s, 2H), 6.72 (d, J = 8.4 Hz, 1H), 6.59 (s, 2H), 2.28 (s, 3H), 2.01 (s, 6H).
92	484.00	2.12	1H NMR (400 MHz, DMSO) δ 12.15 (s, 1H), 8.22 (d, J = 4.4 Hz, 1H), 8.04 (d, J = 7.8 Hz, 1H), 7.84 (d, J = 8.5 Hz, 1H), 7.73 (s, 1H), 7.16 (d, J = 7.8 Hz, 1H), 6.92 (s, 2H), 3.95 (s, 3H), 2.26 (s, 3H), 1.99 (s, 6H), 1.06 (s, 9H).
93	507.00	1.98	1H NMR (400 MHz, DMSO) δ 12.29 (s, 1H), 8.14 (d, J = 7.9 Hz, 1H), 7.82 (s, 2H), 7.73 (d, J = 7.9 Hz, 1H), 7.65 (s, 1H), 7.25 (s, 3H), 6.95 (s, 2H), 6.71 (d, J = 8.4 Hz, 1H), 6.53 (s, 2H), 2.28 (s, 3H), 2.03 (s, 6H).
94	436.20	1.06	
95	479.50	1.20	
96	480.30	1.88	1H NMR (400 MHz, DMSO-d6) δ 12.64 (s, 1H), 8.20 (d, J = 7.6 Hz, 1H), 7.63 (d, J = 7.6 Hz, 1H), 7.25 - 7.16 (m, 2H), 7.10 - 7.05 (m, 1H), 6.91 (s, 2H), 6.81 (ddd, J = 8.1, 2.3, 1.0 Hz, 1H), 5.68 (s, 2H), 2.26 (s, 3H), 1.94 (s, 6H).
97	517.00	2.03	1H NMR (400 MHz, DMSO) δ 12.42 (s, 1H), 8.11 (d, J = 6.8 Hz, 1H), 7.64 (s, 1H), 7.16 (dd, J = 53.1, 23.0 Hz, 6H), 6.83 (s, 2H), 6.71 (d, J = 8.5 Hz, 1H), 6.54 (s, 2H), 2.18 (s, 3H), 1.98 (s, 6H), 1.88 (s, 6H).
98	480.09	2.06	1H NMR (400 MHz, DMSO) δ 7.91 (s, 2H), 7.69 (s, 1H), 7.58 (s, 2H), 7.45 (s, 1H), 7.37 (s, 1H), 7.21 (s, 1H), 6.60 (s, 1H), 5.60 (s, 2H), 2.31 (s, 3H), 1.25 (s, 9H).
99	471.30	1.95	
100	506.09	2.02	
101	441.40	2.07	
102	468.40	2.11	1H NMR (400 MHz, DMSO-d6) δ 11.83 (s, 1H), 7.69 - 7.54 (m, 2H), 7.21 (d, J = 7.3 Hz, 1H), 7.12 (d, J = 8.2 Hz, 1H), 7.00 (s, 2H), 6.70 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 6.38 - 6.30 (m, 1H), 2.28 (s, 3H), 2.02 (s, 6H), 1.10 (s,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			9H).
103	494.30	1.44	
104	497.00	2.15	1H NMR (400 MHz, DMSO) δ 12.59 (s, 1H), 7.78 (d, J = 7.4 Hz, 1H), 7.43 (s, 2H), 7.06 (s, 1H), 6.75 (s, 2H), 5.40 (s, 2H), 2.66 (s, 6H), 2.56 (s, 3H), 2.17 (s, 3H), 2.06 (s, 6H), 1.15 (s, 9H).
105	453.00	1.95	1H NMR (400 MHz, DMSO-d6) δ 12.34 (s, 1H), 7.25 (t, J = 7.9 Hz, 1H), 7.21 (t, J = 2.1 Hz, 1H), 7.11 - 7.03 (m, 2H), 7.01 - 6.94 (m, 3H), 6.85 (dd, J = 7.8, 2.2 Hz, 1H), 5.49 (s, 2H), 2.90 (p, J = 7.6 Hz, 1H), 2.13 (s, 6H), 1.92 - 1.80 (m, 2H), 1.68 - 1.44 (m, 6H).
106	468.20	2.07	
107	470.00	1.87	1H NMR (400 MHz, DMSO) δ 12.19 (s, 1H), 11.76 (s, 1H), 8.44 (s, 1H), 8.06 (s, 1H), 7.80 (s, 1H), 7.16 (s, 1H), 6.92 (s, 2H), 6.34 (s, 1H), 2.26 (s, 3H), 1.98 (s, 6H), 1.04 (s, 9H).
108	463.00	2.19	
109	464.30	2.08	
110	455.40	4.15	1H NMR (400 MHz, DMSO-d6) δ 7.26 (t, J = 7.9 Hz, 1H), 7.21 (t, J = 2.0 Hz, 1H), 7.12 - 7.06 (m, 1H), 7.01 (s, 1H), 6.89 - 6.83 (m, 1H), 6.77 (s, 2H), 5.45 (s, 2H), 2.17 (s, 3H), 2.11 (s, 6H), 1.14 (s, 9H).
111	479.40	1.91	1H NMR (400 MHz, CDCl3) δ 10.13 (s, 1H), 8.42 (d, J = 8.0 Hz, 1H), 7.52 (d, J = 8.0 Hz, 2H), 7.45 (s, 2H), 7.34 (t, J = 7.7 Hz, 1H), 7.15 (d, J = 8.1 Hz, 1H), 7.00 (s, 1H), 2.11 (d, J = 8.2 Hz, 7H), 1.05 (s, 9H).
112	449.00	1.92	
113	470.00	1.95	1H NMR (400 MHz, DMSO) δ 12.23 (s, 1H), 8.01 (d, J = 7.4 Hz, 1H), 7.63 (d, J = 6.2 Hz, 1H), 7.12 (d, J = 7.3 Hz, 2H), 6.90 (s, 3H), 6.85 (s, 1H), 6.57 (s, 1H), 2.25 (s, 3H), 1.95 (s, 6H), 1.04 (s, 10H).
114	539.30	1.83	
115	483.00	2.27	1H NMR (400 MHz, DMSO) δ 12.11 (s, 1H), 8.03 (s, 2H), 7.78 (s, 1H), 7.35 (d, J = 8.4 Hz, 1H), 7.30 - 7.19 (m, 2H),

Cmpd. No	LC/MS M+1	LC/RT min	NMR
			7.00 (s, 2H), 3.95 (s, 3H), 2.35 (s, 3H), 2.04 (s, 6H), 1.13 (s, 9H).
116	464.09	1.69	
117	454.00	2.15	1H NMR (400 MHz, DMSO) δ 8.90 (d, J = 5.9 Hz, 2H), 7.99 (d, J = 7.8 Hz, 1H), 7.93 (d, J = 6.0 Hz, 2H), 7.11 (d, J = 7.8 Hz, 1H), 6.89 (s, 2H), 2.25 (s, 3H), 1.92 (s, 6H), 1.04 (s, 9H).
118	462.20	1.85	
119	496.21	2.08	
120	478.20	1.85	
121	450.00	1.36	1H NMR (400 MHz, DMSO) δ 7.55 (d, J = 8.6 Hz, 2H), 7.33 (s, 3H), 7.21 (d, J = 7.1 Hz, 1H), 6.83 (s, 2H), 6.61 (d, J = 8.8 Hz, 2H), 5.21 (s, 1H), 4.40 (s, 2H), 4.22 (s, 2H), 2.30 (s, 6H), 2.22 (s, 3H).
122	506.30	1.92	1H NMR (400 MHz, DMSO) δ 8.14 (dd, J = 4.9, 1.9 Hz, 1H), 7.98 (dd, J = 7.5, 1.9 Hz, 1H), 7.68 (s, 1H), 7.47 - 7.38 (m, 2H), 7.33 (d, J = 7.8 Hz, 1H), 7.13 - 7.01 (m, 2H), 6.80 (s, 3H), 5.60 (s, 2H), 4.01 (s, 3H), 2.18 (s, 3H), 2.14 (d, J = 2.9 Hz, 6H).
123	469.30	2.39	
124	451.00	1.68	1H NMR (400 MHz, DMSO) δ 12.38 (s, 1H), 8.44 (s, 1H), 7.19 (s, 3H), 7.08 (s, 1H), 6.90 (s, 2H), 6.82 (s, 1H), 5.69 (s, 2H), 2.42 (s, 3H), 2.24 (s, 3H), 1.92 (s, 9H).
125	562.20	2.23	
126	450.40	1.20	
127	467.30	2.15	
128	489.00	1.45	
129	478.40	1.90	
130	425.40	2.23	
131	469.00	2.16	1H NMR (400 MHz, DMSO) δ 11.66 (s, 1H), 11.04 (s, 1H), 8.05 (d, J = 7.8 Hz, 1H), 7.83 (d, J = 7.9 Hz, 1H), 7.52 (s, 1H), 7.16 (d, J = 7.9 Hz, 1H), 7.02 (s, 2H), 6.93 (s, 2H), 2.26 (s, 3H), 1.98 (s, 6H), 1.04 (s, 9H).
132	558.14	2.91	
133	504.23	2.17	
134	448.00	1.82	1H NMR (400 MHz, DMSO) δ 7.63 (dd, J = 19.9, 8.3 Hz, 3H), 7.38 (s, 1H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			7.07 (d, J = 25.7 Hz, 2H), 6.91 (d, J = 8.3 Hz, 1H), 6.62 (d, J = 8.8 Hz, 2H), 6.15 (s, 2H), 5.66 (s, 2H), 2.16 (s, 3H), 1.90 (s, 6H).
135	506.09	2.04	
136	596.20	2.37	
137	467.10	2.19	
138	427.50	1.84	
139	453.20	6.23	1H NMR (400 MHz, DMSO) δ 1.10 (s, 9 H), 1.46 (d, J = 6.8 Hz, 3 H), 2.16 (s, 6 H), 2.21 (s, 3 H), 5.40 - 5.53 (m, 3 H), 6.83 (s, 2 H), 7.15 (s, 1 H), 7.55 - 7.62 (m, 2 H), 7.68 - 7.76 (m, 3 H)
140	493.00	1.49	1H NMR (400 MHz, DMSO) δ 12.49 (s, 1H), 7.59 (s, 1H), 7.32 (s, 1H), 7.19 (s, 3H), 7.14 (d, J = 7.2 Hz, 1H), 6.80 (s, 2H), 6.67 (d, J = 8.4 Hz, 1H), 6.43 (s, 1H), 4.64 (s, 1H), 2.95 (d, J = 11.1 Hz, 1H), 2.20 (s, 9H), 2.00 (d, J = 11.2 Hz, 1H), 1.05 (s, 3H), 0.91 (s, 3H).
141	469.40	2.02	
142	466.40	1.87	
143	447.20	2.14	
144	462.40	1.93	
145	502.20	2.34	
146	506.00	2.31	1H NMR (400 MHz, DMSO) δ 12.22 (s, 1H), 7.84 (s, 2H), 7.76 (s, 1H), 7.53 (s, 1H), 7.35 (s, 1H), 7.09 (s, 1H), 6.87 (s, 3H), 3.84 (s, 3H), 2.24 (s, 3H), 1.84 (s, 6H), 1.02 (s, 9H).
147	453.40	2.34	
148	448.90	1.85	1H NMR (400 MHz, DMSO-d6) δ 12.35 (s, 1H), 7.67 (d, J = 7.8 Hz, 1H), 7.46 (s, 1H), 7.25 - 7.18 (m, 2H), 7.13 - 7.09 (m, 1H), 7.08 - 7.02 (m, 2H), 6.91 - 6.87 (m, 1H), 6.84 - 6.80 (m, 1H), 6.74 (s, 2H), 5.68 (s, 2H), 2.17 (s, 3H), 1.92 (s, 6H).
149	530.30	2.18	1H NMR (400 MHz, DMSO-d6) δ 8.07 - 8.01 (m, 2H), 7.81 - 7.74 (m, 1H), 7.73 - 7.65 (m, 3H), 7.49 (s, 1H), 7.48 - 7.39 (m, 3H), 7.10 - 7.03 (m, 1H), 6.99 - 6.95 (m, 2H), 5.63 (s, 2H), 2.13 (s, 6H).
150	459.39	3.00	

Cmpd. No.	LC/MS M+1	LC/RT* min	NMR
151	534.19	2.03	
152	490.00	2.17	
153	466.20	1.88	
154	426.40	1.66	
155	494.07	2.09	
156	497.00	2.23	1H NMR (400 MHz, DMSO) δ 7.25 (s, 1H), 7.21 (s, 1H), 7.09 (s, 1H), 7.01 (s, 1H), 6.97 (s, 2H), 6.86 (s, 1H), 5.46 (s, 2H), 2.16 (s, 6H), 2.08 (s, 1H), 1.23 (s, 9H), 1.15 (s, 9H).
157	442.06	1.81	
158	453.50	2.06	1H NMR (400 MHz, DMSO-d6) δ 7.26 - 7.16 (m, 2H), 7.05 (d, J = 7.8 Hz, 1H), 6.88 - 6.78 (m, 2H), 6.76 (s, 2H), 5.44 (s, 2H), 2.17 (s, 3H), 2.10 (s, 6H), 1.27 (s, 3H), 0.75 - 0.69 (m, 2H), 0.67 - 0.62 (m, 2H).
159	530.30	2.27	
160	423.15	1.93	
161	551.20	2.11	
162	517.39	1.70	
163	469.00	1.94	1H NMR (400 MHz, DMSO) δ 8.27 (s, 1H), 8.16 (s, 1H), 7.97 (d, J = 7.8 Hz, 1H), 7.56 (s, 1H), 7.12 (d, J = 7.8 Hz, 1H), 6.90 (s, 2H), 2.25 (s, 3H), 1.94 (s, 6H), 1.04 (s, 9H).
164	531.20	2.34	1H NMR (400 MHz, DMSO-d6) δ 12.24 (s, 1H), 8.12 (d, J = 7.8 Hz, 1H), 7.73 - 7.60 (m, 4H), 7.31 - 7.19 (m, 3H), 6.95 (s, 2H), 6.71 (d, J = 8.2 Hz, 1H), 6.52 (s, 2H), 2.88 (p, J = 6.9 Hz, 1H), 2.28 (s, 3H), 2.03 (s, 6H), 1.18 (d, J = 6.9 Hz, 6H).
165	441.40	2.04	
166	464.30	2.10	
167	455.00	1.92	1H NMR (400 MHz, DMSO) δ 7.26 (d, J = 12.4 Hz, 2H), 7.13 (s, 1H), 6.93 (s, 1H), 6.90 (s, 1H), 6.77 (s, 2H), 5.44 (s, 2H), 3.18 (s, 1H), 2.30 (s, 2H), 2.18 (s, 3H), 2.07 (s, 6H), 1.75 (s, 1H), 0.83 (d, J = 6.6 Hz, 6H).
168	464.09	1.73	
169	580.30	2.24	
170	462.39	3.09	
171	542.60	1.70	

Cmpd. No	LC/MS M+1	LC/RT min	NMR
172	509.20	2.32	
173	456.50	1.92	
174	466.00	1.41	1H NMR (400 MHz, DMSO) δ 10.10 (s, 1H), 7.83 (s, 1H), 7.60 (s, 2H), 7.31 (d, J = 12.3 Hz, 3H), 7.00 - 6.85 (m, 3H), 5.22 (s, 1H), 4.46 (d, J = 13.2 Hz, 1H), 4.38 - 4.17 (m, 3H), 3.78 (s, 3H), 2.36 (s, 6H), 2.23 (s, 3H).
175	481.00	1.76	1H NMR (400 MHz, DMSO) δ 12.76 (s, 1H), 8.28 (s, 1H), 7.66 (s, 2H), 7.20 (s, 1H), 6.92 (s, 2H), 6.65 (d, J = 46.8 Hz, 3H), 2.25 (s, 3H), 1.97 (s, 6H).
176	476.30	2.05	
177	442.17	1.85	
178	489.50	1.72	
179	477.20	2.01	1H NMR (400 MHz, DMSO-d6) δ 12.39 (s, 1H), 7.63 (t, J = 7.9 Hz, 1H), 7.52 - 7.44 (m, 2H), 7.17 (d, J = 7.3 Hz, 1H), 6.97 (dd, J = 8.8, 1.7 Hz, 1H), 6.74 (d, J = 6.0 Hz, 3H), 6.69 (d, J = 8.4 Hz, 1H), 6.46 (s, 2H), 5.66 (s, 2H), 2.61 (q, J = 7.6 Hz, 2H), 2.18 (s, 3H), 1.95 (s, 6H), 1.17 (t, J = 7.5 Hz, 3H).
180	479.30	2.09	
181	466.40	1.88	
182	495.00	2.18	1H NMR (400 MHz, DMSO) δ 11.98 (s, 1H), 7.74 (s, 1H), 7.63 (s, 1H), 7.21 (d, J = 7.3 Hz, 1H), 6.90 (s, 2H), 6.70 (d, J = 8.4 Hz, 1H), 6.51 (s, 2H), 2.68 (s, 2H), 2.26 (s, 3H), 1.97 (s, 6H), 1.66 (d, J = 27.0 Hz, 4H), 0.92 (s, 6H).
183	463.00	2.02	1H NMR (400 MHz, DMSO) δ 12.45 (s, 1H), 7.90 (s, 1H), 7.67 (s, 2H), 7.57 (s, 1H), 7.13 (d, J = 48.8 Hz, 4H), 6.90 (s, 1H), 6.72 (s, 2H), 5.64 (s, 2H), 3.84 (s, 3H), 2.17 (s, 3H), 1.88 (s, 6H).
184	449.20	1.64	1H NMR (400 MHz, DMSO-d6) δ 8.39 (dd, J = 4.6, 1.6 Hz, 1H), 8.15 (dd, J = 8.0, 1.6 Hz, 1H), 7.30 (s, 1H), 7.28 - 7.21 (m, 2H), 7.18 (dd, J = 8.0, 4.6 Hz, 1H), 7.06 - 7.01 (m, 1H), 6.96 - 6.90 (m, 1H), 6.58 (s, 2H), 5.65 (s, 2H), 2.10 (s, 3H), 1.88 (s, 6H).
185	441.00	1.91	1H NMR (400 MHz, DMSO) δ 7.34 (s, 2H), 7.23 (s, 1H), 7.03 (s, 2H), 7.00 (s,

Cmpd No.	LC/MS M+1	LC/RT min	NMR
			1H), 6.96 (s, 2H), 5.50 (s, 2H), 2.14 (s, 6H), 1.14 (s, 9H).
186	438.17	1.72	
187	422.17	0.93	
188	523.00	2.02	1H NMR (400 MHz, MeOD) δ 7.89 (s, 1H), 7.84 (s, 1H), 7.51 (s, 1H), 7.34 (s, 1H), 6.99 (s, 1H), 6.75 (s, 1H), 6.66 (s, 2H), 5.36 (s, 2H), 4.38 (s, 2H), 3.90 (s, 2H), 2.15 (s, 3H), 1.93 (s, 6H), 1.17 (s, 10H).
189	515.00	1.94	1H NMR (400 MHz, DMSO) δ 7.80 (s, 2H), 7.61 (s, 1H), 7.25 (s, 2H), 7.14 (s, 1H), 7.10 (s, 1H), 6.85 (s, 1H), 5.68 (s, 2H), 2.08 (s, 1H), 1.06 (s, 9H).
190	507.00	1.97	1H NMR (400 MHz, DMSO) δ 13.75 (s, 1H), 12.32 (s, 1H), 8.04 (d, J = 7.9 Hz, 1H), 7.96 (s, 1H), 7.71 (d, J = 8.6 Hz, 2H), 7.64 (s, 1H), 6.94 (s, 4H), 6.86 (s, 1H), 4.03 (s, 2H), 2.28 (s, 3H), 2.02 (s, 6H), 1.31 (s, 3H).
191	484.00	2.28	1H NMR (400 MHz, DMSO) δ 12.43 (s, 1H), 8.49 (d, J = 6.5 Hz, 1H), 8.34 (d, J = 9.3 Hz, 1H), 7.95 (d, J = 7.8 Hz, 1H), 7.27 (s, 1H), 7.14 (d, J = 7.8 Hz, 1H), 6.90 (s, 2H), 3.94 (s, 3H), 2.25 (s, 3H), 1.93 (s, 6H), 1.05 (s, 9H).
192	424.17	1.65	
193	453.30	1.90	
194	462.30	1.88	
195	454.00	2.07	
196	579.30	2.39	1H NMR (400 MHz, DMSO-d6) δ 12.37 (s, 1H), 8.13 (d, J = 7.8 Hz, 1H), 7.81 (d, J = 7.9 Hz, 1H), 7.65 (t, J = 7.8 Hz, 1H), 7.27 - 7.14 (m, 3H), 6.95 (s, 2H), 6.85 (dt, J = 10.7, 2.3 Hz, 1H), 6.71 (d, J = 8.4 Hz, 1H), 6.53 (s, 2H), 3.65 (d, J = 6.6 Hz, 2H), 2.27 (s, 3H), 2.02 (s, 6H), 1.97 (dt, J = 13.3, 6.6 Hz, 1H), 0.97 (d, J = 6.8 Hz, 6H).
197	459.32	1.67	
198	493.20	1.91	
199	460.20	2.12	
200	466.60	1.40	
201	454.14	2.22	
202	460.50	2.23	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
203	489.40	2.82	1H NMR (400 MHz, DMSO-d6) δ 12.30 (s, 1H), 8.14 (d, J = 7.8 Hz, 1H), 7.81 - 7.70 (m, 3H), 7.64 (s, 1H), 7.44 - 7.38 (m, 3H), 7.23 (d, J = 7.3 Hz, 1H), 6.95 (s, 2H), 6.69 (s, 1H), 6.52 (s, 2H), 2.28 (s, 3H), 2.04 (s, 6H).
204	455.40	2.04	1H NMR (400 MHz, DMSO-d6) δ 12.14 (s, 1H), 7.97 (d, J = 7.8 Hz, 1H), 7.63 (t, J = 7.8 Hz, 1H), 7.20 (d, J = 7.3 Hz, 1H), 7.15 (d, J = 7.8 Hz, 1H), 7.09 (s, 1H), 7.08 - 7.00 (m, 2H), 6.70 (d, J = 8.4 Hz, 1H), 6.50 (s, 2H), 2.30 (s, 3H), 2.01 (s, 3H), 1.09 (s, 9H).
205	497.00	2.13	1H NMR (400 MHz, DMSO) δ 8.13 (s, 1H), 7.92 (s, 1H), 7.79 (s, 1H), 7.71 (s, 1H), 7.62 (s, 1H), 7.26 (s, 1H), 6.63 (d, J = 28.8 Hz, 3H), 1.98 (s, 3H), 1.68 (s, 6H), 0.76 (s, 9H).
206	439.30	1.95	
207	541.10	2.20	
208	519.30	2.08	
209	503.30	2.07	1H NMR (400 MHz, DMSO) δ 7.26 (dd, J = 14.6, 6.7 Hz, 2H), 7.12 (dd, J = 14.0, 7.3 Hz, 3H), 7.04 (d, J = 7.5 Hz, 2H), 6.86 (d, J = 8.0 Hz, 1H), 6.77 (s, 2H), 5.74 (s, 1H), 5.56 (s, 2H), 2.17 (s, 3H), 2.08 (d, J = 2.2 Hz, 6H), 1.97 (s, 6H).
210	479.20	3.14	
211	468.39	3.11	1H NMR (400 MHz, DMSO) δ 7.65 - 7.62 (d, 1H), 7.59 - 7.56 (d, 1H), 7.06 - 6.94 (m, 5H), 5.47 (s, 2H), 4.68 (t, J = 8.68, 2H), 3.29 (t, J = 9.01, 2H), 2.13 - 2.09 (m, 6H), 1.16 (s, 9H)
212	451.20	2.08	
213	464.30	1.20	
214	443.50	2.86	
215	447.30	2.07	
216	514.31	2.79	
217	484.00	2.33	1H NMR (400 MHz, DMSO) δ 8.46 (d, J = 5.3 Hz, 1H), 8.00 (d, J = 7.8 Hz, 1H), 7.49 (d, J = 5.3 Hz, 1H), 7.29 (s, 1H), 7.12 (d, J = 7.8 Hz, 1H), 6.90 (s, 2H), 3.91 (s, 3H), 2.25 (s, 4H), 1.93 (s, 6H), 1.05 (s, 9H).

Cmpd. No	LC/MS M+1	LC/RT min	NMR
218	466.00	1.38	1H NMR (400 MHz, DMSO) δ 10.21 (s, 1H), 8.28 (s, 1H), 8.18 (d, J = 9.3 Hz, 1H), 7.58 (d, J = 7.2 Hz, 1H), 7.34 (s, 2H), 7.28 (s, 1H), 7.10 (s, 1H), 6.89 (s, 2H), 5.21 (s, 1H), 4.42 (s, 1H), 4.22 (s, 5H), 3.89 (s, 3H), 2.36 (s, 6H), 2.23 (s, 3H).
219	454.00	2.07	1H NMR (400 MHz, MeOD) δ 8.07 (s, 1H), 7.40 (s, 2H), 7.33 (s, 3H), 7.27 (s, 3H), 7.06 (s, 1H), 6.94 (s, 1H), 6.34 (s, 1H), 1.73 (s, 3H), 1.22 (s, 9H).
220	514.03	2.17	
221	479.40	1.93	
222	454.00	2.19	1H NMR (400 MHz, DMSO) δ 12.54 (s, 1H), 8.73 (s, 1H), 8.18 (s, 2H), 7.96 (s, 1H), 7.74 (s, 1H), 7.13 (s, 1H), 6.90 (s, 2H), 2.25 (s, 3H), 1.95 (s, 6H), 1.05 (s, 9H).
223	444.10	1.81	
224	467.40	2.20	
225	530.20	2.22	
226	487.27	2.15	
227	434.20	1.80	
228	469.00	2.07	1H NMR (400 MHz, DMSO) δ 8.02 (d, J = 7.8 Hz, 1H), 7.64 (s, 1H), 7.21 (d, J = 7.3 Hz, 1H), 7.12 (d, J = 7.8 Hz, 1H), 6.91 (s, 2H), 6.70 (d, J = 8.4 Hz, 1H), 6.51 (s, 2H), 2.26 (s, 3H), 1.97 (s, 6H), 1.05 (s, 9H).
229	466.25	2.06	
230	450.00	1.40	1H NMR (400 MHz, DMSO) δ 10.81 (s, 1H), 7.56 (d, J = 7.8 Hz, 1H), 7.39 (d, J = 7.1 Hz, 1H), 7.23 (d, J = 22.0 Hz, 4H), 6.85 (s, 2H), 6.74 (d, J = 8.2 Hz, 1H), 6.56 (s, 1H), 4.97 (s, 1H), 4.19 (d, J = 35.8 Hz, 5H), 2.31 (s, 6H), 2.22 (s, 3H).
231	488.40	2.08	
232	447.20	2.06	
233	573.30	1.97	
234	523.20	2.15	
235	469.00	2.05	1H NMR (400 MHz, DMSO) δ 8.23 (d, J = 5.8 Hz, 1H), 8.05 (dd, J = 7.9, 1.0 Hz, 1H), 8.00 (d, J = 8.2 Hz, 1H), 7.13 (d, J = 7.8 Hz, 1H), 6.90 (s, 2H), 6.82

Cmpd No.	LC/MS M+1	LC/RT min	NMR
			(s, 1H), 6.75 (s, 1H), 2.25 (s, 3H), 1.95 (s, 6H), 1.04 (s, 10H).
236	479.00	1.41	1H NMR (400 MHz, MeOD) δ 7.84 (s, 1H), 7.33 (s, 7H), 6.99 (d, J = 8.7 Hz, 1H), 6.90 (s, 2H), 4.60 (d, J = 35.7 Hz, 2H), 4.41 (s, 1H), 3.98 (s, 1H), 3.76 (s, 1H), 3.67 - 3.55 (m, 1H), 2.90 (dt, J = 12.8, 6.3 Hz, 1H), 2.41 (d, J = 14.1 Hz, 7H), 2.24 (s, 4H).
237	442.21	1.87	
238	595.20	2.43	1H NMR (400 MHz, DMSO-d6) δ 8.09 (d, J = 7.7 Hz, 1H), 7.61 (d, J = 7.9 Hz, 1H), 7.50 - 7.32 (m, 4H), 7.07 (d, J = 7.4 Hz, 1H), 6.90 (s, 2H), 6.45 (d, J = 8.2 Hz, 1H), 6.09 (s, 2H), 3.60 (d, J = 6.5 Hz, 2H), 2.26 (s, 3H), 2.02 (s, 7H), 1.01 (d, J = 6.7 Hz, 6H).
239	499.00	1.89	1H NMR (400 MHz, DMSO) δ 7.57 (d, J = 19.8 Hz, 3H), 7.28 (s, 2H), 7.16 (d, J = 18.2 Hz, 2H), 6.94 (s, 1H), 5.67 (s, 2H), 1.09 (s, 9H).
240	448.30	1.82	
241	463.20	2.04	
242	439.00	2.28	1H NMR (400 MHz, DMSO-d6) δ 11.87 (s, 1H), 8.03 - 7.97 (m, 2H), 7.75 - 7.69 (m, 1H), 7.68 - 7.62 (m, 2H), 7.24 (d, J = 2.0 Hz, 1H), 6.88 (s, 2H), 6.15 (d, J = 1.9 Hz, 1H), 5.27 (s, 2H), 2.23 (s, 3H), 2.05 (s, 6H), 1.07 (s, 9H).
243	466.00	1.70	1H NMR (400 MHz, DMSO) δ 8.23 (d, J = 7.6 Hz, 1H), 7.65 (d, J = 7.6 Hz, 1H), 7.24 (d, J = 11.1 Hz, 2H), 7.12 (s, 4H), 6.85 (d, J = 9.9 Hz, 1H), 1.98 (s, 6H).
244	465.30	2.06	
245	451.40	1.94	1H NMR (400 MHz, DMSO-d6) δ 12.42 (s, 1H), 7.98 - 7.89 (m, 2H), 7.75 (t, J = 7.4 Hz, 1H), 7.63 (t, J = 7.7 Hz, 2H), 7.27 (d, J = 8.2 Hz, 1H), 7.22 (d, J = 3.2 Hz, 1H), 7.05 (s, 1H), 6.87 (dd, J = 8.2, 7.2 Hz, 1H), 6.44 - 6.36 (m, 2H), 5.72 (s, 2H), 3.73 (s, 3H), 1.23 (s, 9H).
246	467.00	1.64	1H NMR (400 MHz, DMSO) δ 8.28 (d,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			J = 7.6 Hz, 1H), 7.64 (s, 2H), 7.21 (d, J = 7.3 Hz, 1H), 7.10 (s, 4H), 6.71 (d, J = 8.4 Hz, 1H), 6.59 (s, 2H), 2.01 (s, 6H).
247	434.00	1.88	1H NMR (400 MHz, DMSO) δ 8.60 (s, 1H), 8.03 (s, 2H), 7.61 (s, 2H), 7.01 (s, 3H), 6.83 (s, 1H), 6.72 (s, 2H), 5.79 (s, 2H), 2.17 (s, 3H), 1.92 (s, 6H).
248	447.20	2.15	
249	469.40	2.19	
250	533.22	3.09	
251	454.00	2.04	1H NMR (400 MHz, DMSO) δ 12.01 (s, 1H), 7.91 (d, J = 7.8 Hz, 1H), 7.21 (s, 2H), 7.14 (d, J = 7.8 Hz, 1H), 7.07 (d, J = 14.6 Hz, 4H), 6.81 (s, 1H), 5.66 (s, 2H), 2.30 (s, 3H), 1.99 (s, 3H), 1.09 (s, 9H).
252	466.50	2.04	
253	464.00	2.02	1H NMR (400 MHz, DMSO) δ 12.78 (s, 1H), 8.01 (s, 1H), 7.70 (s, 2H), 7.57 (s, 1H), 7.15 (s, 2H), 7.07 (s, 1H), 7.00 (s, 1H), 6.69 (s, 2H), 5.63 (s, 2H), 3.79 (s, 3H), 2.16 (s, 3H), 1.85 (s, 6H).
254	451.30	1.32	1H NMR (400 MHz, DMSO-d6) δ 10.51 (s, 1H), 9.87 (s, 1H), 7.39 - 7.34 (m, 1H), 7.32 - 7.21 (m, 6H), 6.91 (ddd, J = 7.7, 2.5, 1.5 Hz, 1H), 6.85 (s, 2H), 4.98 (s, 1H), 4.33 - 4.08 (m, 4H), 2.33 (s, 6H), 2.22 (s, 3H).
255	455.00	1.49	
256	458.00	1.77	1H NMR (400 MHz, DMSO) δ 11.45 (s, 1H), 8.52 (s, 1H), 7.63 (s, 1H), 7.18 (d, J = 7.3 Hz, 1H), 6.86 (s, 2H), 6.69 (d, J = 8.4 Hz, 1H), 6.47 (s, 2H), 2.22 (s, 3H), 1.99 (s, 6H), 1.35 (s, 9H).
257	426.07	3.16	
258	464.00	1.80	1H NMR (400 MHz, DMSO) δ 12.46 (s, 1H), 8.17 (s, 1H), 7.79 (s, 1H), 7.70 (s, 2H), 7.60 (s, 1H), 7.09 (d, J = 15.2 Hz, 2H), 6.90 (s, 1H), 6.76 (s, 2H), 5.69 (s, 2H), 3.91 (s, 3H), 2.19 (s, 3H), 1.96 (s, 6H).
259	462.20	1.95	
260	462.30	1.87	
261	456.09	1.84	
262	483.40	3.30	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
263	499.04	2.06	
264	433.20	2.06	
265	475.50	2.11	1H NMR (400 MHz, DMSO-d6) δ 11.96 (s, 1H), 9.07 (s, 1H), 8.00 (d, J = 7.8 Hz, 1H), 7.13 (d, J = 7.8 Hz, 1H), 6.91 (s, 2H), 6.81 (s, 2H), 2.25 (s, 3H), 1.96 (s, 6H), 1.04 (s, 9H).
266	465.40	2.29	1H NMR (400 MHz, DMSO-d6) δ 8.22 (dd, J = 7.9, 1.3 Hz, 1H), 8.12 (dd, J = 7.6, 1.4 Hz, 1H), 7.97 (td, J = 7.8, 1.5 Hz, 1H), 7.90 (td, J = 7.6, 1.3 Hz, 1H), 7.00 (s, 1H), 6.74 (s, 2H), 5.43 (s, 2H), 2.17 (s, 3H), 2.06 (s, 6H), 1.16 (s, 9H).
267	478.30	2.09	
268	517.00	2.08	1H NMR (400 MHz, DMSO) δ 12.16 (s, 1H), 7.91 (d, J = 7.6 Hz, 1H), 7.62 (s, 1H), 7.18 (s, 3H), 7.11 (d, J = 7.1 Hz, 1H), 6.96 (dd, J = 17.7, 9.9 Hz, 5H), 6.69 (d, J = 8.4 Hz, 1H), 6.50 (s, 2H), 2.85 (s, 2H), 2.75 (s, 2H), 2.26 (s, 3H), 1.97 (s, 6H).
269	456.40	2.04	1H NMR (400 MHz, DMSO-d6) δ 12.28 (s, 1H), 10.62 (s, 1H), 7.85 - 7.80 (m, 2H), 6.99 - 6.93 (m, 3H), 6.75 (s, 2H), 5.43 (s, 2H), 2.17 (s, 3H), 2.07 (s, 6H), 1.14 (s, 9H).
270	483.40	2.28	1H NMR (400 MHz, DMSO-d6) δ 12.35 (s, 1H), 7.46 - 7.39 (m, 1H), 7.27 - 7.20 (m, 2H), 7.07 - 7.01 (m, 1H), 6.98 (d, J = 1.2 Hz, 1H), 6.75 (s, 2H), 5.44 (s, 2H), 2.98 (s, 6H), 2.17 (s, 3H), 2.08 (s, 6H), 1.14 (s, 9H).
271	438.30	1.95	
272	426.00	1.84	1H NMR (400 MHz, DMSO) δ 7.92 (d, J = 7.8 Hz, 1H), 7.43 (s, 2H), 7.24 (d, J = 37.6 Hz, 7H), 6.93 (s, 1H), 3.17 (s, 2H), 1.12 (s, 9H).
273	484.30	2.95	
274	438.22	1.86	
275	492.00	2.36	1H NMR (400 MHz, DMSO) δ 12.28 (s, 1H), 8.01 (d, J = 7.8 Hz, 1H), 7.70 (d, J = 8.0 Hz, 1H), 7.52 (d, J = 8.3 Hz, 1H), 7.32 (s, 1H), 7.24 (s, 1H), 7.13 (s, 2H), 6.89 (s, 2H), 2.25 (s, 3H), 1.94 (s, 6H), 1.04 (s, 9H).

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
276	533.00	2.07	
277	504.40	2.28	
278	519.30	1.98	
279	580.20	2.25	
280	490.16	2.12	
281	581.30	2.29	
282	424.40	1.74	
283	469.00	1.79	1H NMR (400 MHz, DMSO) δ 8.10 (s, 1H), 7.99 (s, 1H), 7.21 (s, 1H), 7.10 (s, 1H), 7.00 (s, 1H), 6.89 (s, 2H), 2.25 (s, 3H), 1.94 (s, 6H), 1.04 (s, 9H).
284	478.90	2.81	
285	467.00	1.89	
286	454.00	2.02	1H NMR (400 MHz, DMSO) δ 12.04 (s, 1H), 7.96 (d, J = 7.8 Hz, 1H), 7.22 (s, 2H), 7.09 (s, 5H), 6.82 (d, J = 9.3 Hz, 1H), 5.66 (s, 2H), 1.99 (s, 6H), 1.03 (s, 9H).
287	433.30	2.08	
288	434.60	1.86	
289	482.00	2.26	1H NMR (400 MHz, DMSO) δ 12.04 (s, 1H), 7.94 (d, J = 7.8 Hz, 1H), 7.30 (s, 1H), 7.13 (s, 3H), 6.90 (s, 2H), 6.81 (d, J = 10.4 Hz, 1H), 6.28 (s, 1H), 2.68 (s, 3H), 2.25 (s, 3H), 1.94 (s, 6H), 1.04 (s, 9H).
290	449.00	2.03	
291	473.20	1.74	1H NMR (400 MHz, DMSO-d6) δ 8.31 - 8.30 (m, 1H), 7.52 (d, J = 0.8 Hz, 1H), 7.49 (dd, J = 8.8, 1.6 Hz, 1H), 7.23 (t, J = 7.9 Hz, 1H), 7.19 (t, J = 2.0 Hz, 1H), 7.11 (d, J = 8.9 Hz, 1H), 7.05 - 7.01 (m, 1H), 6.84 (ddd, J = 8.0, 2.3, 1.0 Hz, 1H), 6.73 (s, 2H), 5.69 (s, 2H), 2.16 (s, 3H), 1.89 (s, 6H).
292	467.20	5.47	1H NMR (300 MHz, DMSO) δ 1.01 (s, 9H), 1.92 (s, 6H), 2.22 (s, 3H), 5.57 (br. s, 2H), 6.73 (d, J=8.1 Hz, 1H), 6.85 (s, 2H), 6.97-7.07 (m, 2H), 7.11-7.20 (m, 2H), 7.56 (br. s, 2H), 8.09 (d, J=7.8 Hz, 1H)
293	507.30	2.11	
294	468.20	2.16	1H NMR (400 MHz, DMSO-d6) δ 12.00 (s, 1H), 7.94 (d, J = 7.8 Hz, 1H), 7.25 - 7.19 (m, 2H), 7.12 (d, J = 7.8

Cmpd No	LC/MS M+1	LC/RT min	NMR
			Hz, 1H), 7.09 (dd, J = 1.8, 0.9 Hz, 1H), 6.90 (s, 2H), 6.83 (s, 1H), 5.66 (s, 2H), 2.26 (s, 3H), 1.95 (s, 6H), 1.04 (s, 9H).
295	463.40	1.90	
296	440.00	1.94	1H NMR (400 MHz, DMSO) δ 12.07 (s, 1H), 7.93 (d, J = 7.8 Hz, 1H), 7.29 (d, J = 7.5 Hz, 1H), 7.27 - 7.12 (m, 6H), 7.08 (d, J = 7.8 Hz, 1H), 6.83 (d, J = 11.0 Hz, 1H), 5.68 (s, 2H), 2.03 (s, 3H), 1.08 (s, 9H).
297	424.30	1.73	
298	452.20	1.82	
299	492.38	1.55	
300	477.00	1.99	
301	424.40	1.74	
302	545.50	1.66	
303	517.20	2.13	
304	480.22	2.09	
305	470.00	2.05	1H NMR (400 MHz, DMSO) δ 12.29 (s, 1H), 11.71 (s, 1H), 8.01 (d, J = 7.8 Hz, 1H), 7.93 (s, 1H), 7.61 (s, 1H), 7.13 (d, J = 7.8 Hz, 1H), 7.05 - 6.85 (m, 3H), 2.25 (s, 3H), 1.96 (s, 6H), 1.05 (s, 9H).
306	473.29	1.84	
307	543.36	1.74	1H NMR (400 MHz, DMSO) δ 8.25 (dd, J = 7.3, 2.1 Hz, 1H), 7.90 - 7.81 (m, J = 6.1 Hz, 1H), 7.78 (d, J = 7.9 Hz, 1H), 7.50 (s, 1H), 7.45 (d, J = 10.1 Hz, 1H), 7.34 (d, J = 7.9 Hz, 1H), 6.98 - 6.84 (m, J = 10.7 Hz, 1H), 6.54 - 6.40 (m, J = 6.7 Hz, 1H), 4.17 - 4.01 (m, 2H), 3.84 (d, J = 6.5 Hz, 2H), 2.15 - 2.01 (m, 1H), 2.01 - 1.90 (m, J = 22.2 Hz, 2H), 1.80 - 1.61 (m, 2H), 1.13 (d, J = 6.2 Hz, 6H), 0.99 (d, J = 6.7 Hz, 6H).
308	450.40	1.87	
309	464.30	1.26	
310	450.40	1.20	
311	496.00	2.38	1H NMR (400 MHz, DMSO) δ 12.08 (s, 1H), 7.95 (s, 1H), 7.39 (s, 1H), 7.22 (s, 2H), 7.12 (s, 1H), 7.00 (s, 1H), 6.90 (s, 2H), 2.92 (s, 6H), 2.25 (s, 3H), 1.93 (s, 6H), 1.04 (s, 9H).
312	455.18	2.91	
313	466.20	2.22	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
314	452.40	1.95	
315	496.08	2.07	
316	514.09	2.20	
317	469.00	2.09	1H NMR (400 MHz, DMSO) δ 7.26 (s, 1H), 7.21 (s, 1H), 7.09 (d, J = 7.8 Hz, 1H), 7.04 (s, 1H), 6.89 (s, 1H), 6.85 (d, J = 10.3 Hz, 1H), 2.14 (s, 6H), 2.04 (s, 6H), 1.13 (s, 9H).
318	553.00	2.24	1H NMR (400 MHz, DMSO) δ 13.80 (s, 1H), 12.51 (s, 1H), 8.08 (d, J = 7.8 Hz, 1H), 7.98 (s, 1H), 7.82 (d, J = 7.9 Hz, 1H), 7.22 (d, J = 9.8 Hz, 1H), 7.18 (s, 1H), 6.95 (s, 2H), 6.89 (s, 1H), 6.84 (d, J = 12.9 Hz, 1H), 3.65 (d, J = 6.5 Hz, 2H), 2.27 (s, 3H), 2.07 (s, 1H), 2.01 (s, 7H), 0.97 (d, J = 6.7 Hz, 6H).
319	452.00	1.11	1H NMR (400 MHz, DMSO) δ 9.99 (s, 1H), 7.64 (d, J = 28.4 Hz, 3H), 7.33 (d, J = 17.6 Hz, 4H), 6.89 (s, 2H), 6.65 (s, 1H), 5.30 (s, 1H), 4.50 (s, 1H), 4.30 (s, 3H), 2.38 (s, 6H), 2.22 (s, 3H).
320	561.30	2.31	1H NMR (400 MHz, DMSO-d6) δ 12.15 (s, 1H), 8.10 (d, J = 7.9 Hz, 1H), 7.75 - 7.61 (m, 4H), 7.23 (d, J = 7.2 Hz, 1H), 6.99 - 6.88 (m, 4H), 6.71 (d, J = 8.4 Hz, 1H), 6.56 - 6.46 (m, 2H), 3.77 (d, J = 6.6 Hz, 2H), 2.28 (s, 3H), 2.03 (s, 7H), 0.95 (d, J = 6.7 Hz, 6H).
321	511.30	1.58	
322	468.00	2.25	1H NMR (400 MHz, DMSO) δ 8.06 (s, 1H), 7.74 (s, 1H), 7.36 (s, 1H), 7.20 (s, 1H), 6.99 (s, 2H), 6.87 (s, 1H), 6.72 (s, 1H), 2.34 (s, 3H), 2.04 (s, 6H), 1.33 (s, 2H), 1.13 (s, 9H).
323	469.00	2.11	
324	455.00	1.88	
325	457.30	1.87	
326	464.40	2.14	
327	439.30	1.98	
328	448.10	2.17	
329	530.34	1.70	
330	454.00	2.16	1H NMR (400 MHz, DMSO) δ 12.52 (s, 1H), 9.15 (s, 1H), 8.90 (s, 1H), 8.41 (d, J = 8.1 Hz, 1H), 7.99 (d, J = 7.8 Hz, 1H), 7.70 (s, 1H), 7.11 (d, J = 7.8 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 6.89 (s, 2H), 2.25 (s, 3H), 1.91 (s, 6H), 1.04 (s, 9H).
331	434.05	1.71	
332	482.00	2.18	1H NMR (400 MHz, DMSO) δ 8.40 (s, 1H), 8.24 (s, 1H), 8.14 (s, 1H), 7.76 (s, 1H), 6.89 (s, 1H), 6.64 (s, 2H), 5.34 (s, 2H), 2.60 (s, 3H), 2.08 (s, 3H), 1.95 (s, 6H), 1.07 (s, 9H).
333	464.30	3.03	
334	411.30	1.54	
335	463.00	2.07	1H NMR (400 MHz, DMSO) δ 12.56 (s, 1H), 7.68 (d, J = 7.7 Hz, 1H), 7.53 (s, 2H), 7.44 (s, 2H), 7.30 (d, J = 7.7 Hz, 1H), 7.14 (s, 1H), 7.06 (s, 1H), 6.98 (d, J = 8.4 Hz, 1H), 6.68 (s, 2H), 5.64 (s, 2H), 3.84 (s, 3H), 2.14 (s, 3H), 1.87 (s, 6H).
336	478.20	1.83	
337	511.30	1.41	
338	468.40	1.58	
339	480.30	2.11	
340	475.40	2.70	1H NMR (400 MHz, DMSO-d6) δ 12.32 (s, 1H), 8.10 (d, J = 7.9 Hz, 1H), 7.82 (dd, J = 6.8, 2.9 Hz, 2H), 7.75 (d, J = 7.9 Hz, 1H), 7.64 (t, J = 7.8 Hz, 1H), 7.46 - 7.39 (m, 3H), 7.22 (d, J = 7.4 Hz, 1H), 7.14 (t, J = 4.2 Hz, 2H), 7.12 - 7.05 (m, 1H), 6.71 (d, J = 8.3 Hz, 1H), 6.52 (s, 2H), 2.33 (s, 3H), 2.07 (s, 3H).
341	453.00	2.38	
342	441.00	1.95	1H NMR (400 MHz, DMSO) δ 7.19 (s, 1H), 7.13 (s, 1H), 7.06 (d, J = 8.0 Hz, 4H), 6.96 (s, 2H), 6.83 (s, 1H), 6.39 (s, 1H), 5.67 (s, 1H), 2.17 (s, 3H), 1.67 (s, 3H), 1.28 (s, 9H).
343	489.50	1.76	
344	482.10	1.99	
345	551.20	2.06	
346	466.00	2.00	
347	476.30	2.08	
348	463.20	2.06	
349	442.17	1.88	
350	453.00	2.28	
351	579.30	2.33	
352	491.30	2.06	1H NMR (400 MHz, DMSO-d6) δ 9.12

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			- 9.10 (m, 1H), 9.10 - 9.08 (m, 1H), 8.49 (dd, J = 7.4, 1.2 Hz, 1H), 8.44 - 8.40 (m, 1H), 8.02 (dd, J = 8.5, 7.4 Hz, 1H), 7.81 - 7.76 (m, 1H), 6.97 (s, 1H), 6.59 (s, 2H), 5.26 (s, 2H), 2.10 (s, 3H), 1.78 (s, 6H), 1.15 (s, 9H).
353	525.50	1.49	
354	447.20	2.08	1H NMR (400 MHz, DMSO-d6) δ 7.72 - 7.67 (m, 2H), 7.41 - 7.30 (m, 4H), 6.94 (s, 1H), 6.84 (dd, J = 7.8, 7.0 Hz, 1H), 6.76 (dt, J = 7.0, 1.2 Hz, 1H), 6.68 (s, 2H), 6.11 (s, 2H), 2.28 (s, 3H), 2.16 (s, 3H), 1.70 (s, 6H).
355	502.15	1.71	
356	470.60	1.91	
357	452.18	2.00	
358	458.00	192.00	1H NMR (400 MHz, DMSO) δ 8.49 (s, 1H), 8.14 (s, 1H), 7.91 (s, 2H), 7.74 (s, 1H), 7.61 (s, 1H), 7.16 (d, J = 38.6 Hz, 3H), 6.81 (s, 2H), 5.80 (s, 2H), 2.10 (d, J = 38.6 Hz, 9H).
359	438.22	1.93	
360	456.09	1.94	
361	542.40	1.84	1H NMR (400 MHz, DMSO-d6) δ 12.48 (s, 1H), 7.74 (d, J = 7.9 Hz, 1H), 7.63 (t, J = 7.8 Hz, 1H), 7.50 - 7.47 (m, 1H), 7.41 (dt, J = 10.0, 1.8 Hz, 1H), 7.26 (d, J = 7.9 Hz, 1H), 7.20 (d, J = 7.3 Hz, 1H), 6.89 (dt, J = 10.8, 2.3 Hz, 1H), 6.70 (d, J = 8.3 Hz, 1H), 6.46 (s, 2H), 3.84 (d, J = 6.6 Hz, 2H), 2.81 - 2.72 (m, 2H), 2.05 (dt, J = 13.4, 6.7 Hz, 1H), 1.82 - 1.71 (m, 4H), 1.59 (s, 6H), 0.99 (d, J = 6.7 Hz, 6H).
362	482.40	1.97	
363	499.10	2.04	
364	545.00	2.13	1H NMR (400 MHz, DMSO) δ 12.42 (s, 1H), 8.08 (d, J = 7.6 Hz, 1H), 7.65 (s, 1H), 7.44 (d, J = 8.1 Hz, 1H), 7.31 (s, 1H), 7.26 - 7.16 (m, 2H), 7.11 (d, J = 7.6 Hz, 1H), 6.92 (d, J = 9.0 Hz, 1H), 6.81 (s, 2H), 6.72 (d, J = 8.4 Hz, 1H), 6.54 (s, 2H), 2.17 (s, 3H), 1.98 (s, 6H), 0.97 (s, 10H).
365	482.00	2.18	1H NMR (400 MHz, DMSO) δ 12.29 (s,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 8.48 (s, 1H), 8.34 (s, 1H), 7.92 (s, 1H), 7.27 (s, 1H), 7.09 (s, 1H), 6.89 (s, 2H), 3.93 (s, 3H), 2.24 (s, 3H), 1.90 (s, 6H), 1.32 (s, 3H), 0.67 (d, J = 15.3 Hz, 4H).
366	475.20	4.99	1H NMR (300 MHz, DMSO-d6): δ 2.16 (s, 6 H), 2.17 (s, 3 H), 5.55 (s, 2 H), 6.80 (s, 2 H), 6.83 - 6.90 (m, 1 H), 7.08 - 7.15 (m, 1 H), 7.21 - 7.34 (m, 3 H), 7.35 - 7.44 (m, 2 H), 7.54 (s, 1 H), 7.60 - 7.67 (m, 2 H).
367	455.00	2.00	
368	556.40	2.77	1H NMR (400 MHz, Chloroform-d) δ 15.49 (s, 1H), 8.49 (d, J = 8.2 Hz, 1H), 7.75 - 7.56 (m, 3H), 7.34 (t, J = 1.8 Hz, 1H), 7.30 (dt, J = 9.4, 1.9 Hz, 1H), 6.75 - 6.66 (m, 2H), 3.95 (s, 1H), 3.78 (d, J = 6.5 Hz, 2H), 3.42 (s, 1H), 2.88 - 2.74 (m, 1H), 2.23 (dd, J = 12.5, 8.0 Hz, 1H), 2.17 - 2.08 (m, 1H), 1.98 - 1.87 (m, 1H), 1.55 (s, 3H), 1.39 (s, 3H), 1.31 (d, J = 6.7 Hz, 3H), 1.05 (d, J = 6.7 Hz, 6H).
369	514.03	2.07	
370	479.00	2.05	
371	556.40	2.87	1H NMR (400 MHz, Chloroform-d) δ 13.89 (s, 1H), 8.43 (d, J = 8.1 Hz, 1H), 7.72 - 7.59 (m, 3H), 7.39 - 7.32 (m, 2H), 6.75 - 6.65 (m, 2H), 3.78 (d, J = 6.5 Hz, 2H), 3.50 - 3.43 (m, 2H), 2.91 (s, 2H), 2.44 (dq, J = 10.0, 6.3 Hz, 2H), 2.13 (hept, J = 6.6 Hz, 1H), 1.20 (d, J = 6.8 Hz, 6H), 1.06 (d, J = 6.7 Hz, 6H).
372	485.30	1.89	
373	432.70	1.29	
374	511.00	1.89	1H NMR (400 MHz, DMSO) δ 13.81 (s, 1H), 12.45 (s, 1H), 8.03 (s, 2H), 7.66 (s, 1H), 7.36 (s, 1H), 6.91 (s, 5H), 3.84 (s, 3H), 2.24 (s, 3H), 2.01 (s, 6H).
375	533.20	2.20	
376	537.30	3.07	
377	466.00	1.38	1H NMR (400 MHz, DMSO) δ 10.52 (s, 1H), 9.72 (s, 1H), 8.03 (s, 1H), 7.53 (s, 1H), 7.31 (t, J = 20.3 Hz, 6H), 6.91 (s, 2H), 4.41 (d, J = 13.3 Hz, 1H), 4.26 (d,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			J = 15.8 Hz, 2H), 4.14 (d, J = 13.2 Hz, 1H), 2.39 (s, 7H), 2.24 (s, 3H), 1.82 (s, 3H).
378	439.00	1.27	1H NMR (400 MHz, DMSO) δ 10.20 (s, 1H), 7.73 (s, 1H), 7.42 (s, 1H), 7.26 (d, J = 18.8 Hz, 3H), 6.90 (s, 2H), 6.60 (s, 1H), 4.13 (s, 4H), 2.40 (s, 7H), 2.24 (s, 3H), 1.73 (s, 3H).
379	528.30	2.28	
380	528.29	1.90	
381	521.50	1.78	1H NMR (400 MHz, DMSO-d6) δ 12.08 (s, 1H), 8.07 (s, 1H), 8.00 (d, J = 7.8 Hz, 1H), 7.68 (s, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.40 (d, J = 7.9 Hz, 1H), 7.21 (d, J = 7.2 Hz, 1H), 7.11 (d, J = 8.1 Hz, 2H), 7.06 (dd, J = 8.3, 2.1 Hz, 1H), 6.70 (d, J = 8.3 Hz, 1H), 6.65 - 6.34 (m, 2H), 3.90 (d, J = 7.1 Hz, 2H), 2.31 (s, 3H), 2.07 (d, J = 4.0 Hz, 4H), 0.81 (d, J = 6.7 Hz, 6H).
382	488.30	1.47	1H NMR (400 MHz, DMSO-d6) δ 13.77 (s, 1H), 13.34 (s, 1H), 7.97 (s, 1H), 7.80 (d, J = 7.9 Hz, 1H), 7.66 - 7.27 (m, 3H), 7.01 - 6.76 (m, 2H), 4.13 (q, J = 6.9 Hz, 2H), 4.02 (s, 2H), 2.00 (s, 2H), 1.74 - 1.61 (m, 2H), 1.36 (t, J = 6.9 Hz, 3H), 1.07 (d, J = 6.2 Hz, 6H).
383	554.40	3.03	
384	567.40	2.73	1H NMR (400 MHz, DMSO-d6) δ 12.50 (s, 1H), 8.24 (d, J = 5.4 Hz, 1H), 7.75 (s, 1H), 7.58 (s, 1H), 7.51 (dd, J = 5.4, 1.5 Hz, 1H), 7.28 (s, 2H), 7.16 (s, 1H), 6.62 (s, 1H), 6.42 (s, 1H), 5.30 (p, J = 6.2 Hz, 1H), 2.19 (td, J = 12.8, 11.1, 6.8 Hz, 1H), 1.86 (dd, J = 11.8, 5.6 Hz, 1H), 1.78 - 1.62 (m, 2H), 1.61 (s, 3H), 1.58 (s, 3H), 1.45 - 1.35 (m, 2H), 1.27 (d, J = 6.1 Hz, 3H), 1.17 (t, J = 7.1 Hz, 1H), 0.89 (dd, J = 12.8, 6.3 Hz, 9H).
385	572.25	1.99	
386	542.23	2.97	
387	538.20	2.03	
388	528.12	2.75	
389	525.33	1.12	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
390	589.00	2.25	1H NMR (400 MHz, DMSO-d6) δ 12.80 (s, 1H), 8.52 (dd, J = 2.5, 0.7 Hz, 1H), 8.02 (dd, J = 8.7, 2.5 Hz, 1H), 7.85 (s, 1H), 7.64 (t, J = 7.8 Hz, 1H), 7.23 (d, J = 7.2 Hz, 1H), 6.83 (dd, J = 8.6, 0.7 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 6.51 (s, 2H), 5.32 (p, J = 6.2 Hz, 1H), 3.95 (s, 2H), 1.97 - 1.82 (m, 2H), 1.64 (dd, J = 8.9, 5.3 Hz, 2H), 1.33 (d, J = 6.2 Hz, 6H), 1.04 (d, J = 6.1 Hz, 6H).
391	534.00	2.00	1H NMR (400 MHz, DMSO) δ 13.80 (s, 1H), 12.63 (s, 1H), 8.14 (d, J = 7.8 Hz, 1H), 7.97 (s, 1H), 7.71 (d, J = 7.9 Hz, 1H), 7.45 (s, 1H), 6.94 (s, 2H), 6.88 (s, 1H), 3.48 (s, 3H), 2.26 (s, 3H), 1.99 (s, 6H).
392	467.20	1.28	1H NMR (400 MHz, DMSO-d6) δ 13.20 (s, 1H), 12.55 (s, 1H), 8.25 (dd, J = 7.2, 2.2 Hz, 1H), 8.02 - 7.96 (m, 2H), 7.82 (d, J = 7.9 Hz, 2H), 7.30 (dd, J = 8.1, 6.6 Hz, 3H), 6.44 (t, J = 6.8 Hz, 1H), 4.13 - 4.02 (m, 2H), 2.36 (s, 3H), 2.03 - 1.92 (m, 2H), 1.76 - 1.61 (m, 2H), 1.14 (d, J = 6.2 Hz, 6H).
393	538.25	2.14	
394	481.00	1.20	1H NMR (400 MHz, DMSO) δ 9.77 (s, 1H), 7.64 (s, 1H), 7.14 (s, 2H), 6.90 (s, 2H), 6.69 (s, 3H), 4.37 (d, J = 12.8 Hz, 1H), 4.25 (s, 2H), 3.96 (d, J = 12.9 Hz, 1H), 2.39 (s, 6H), 2.24 (s, 3H), 1.85 (s, 3H).
395	556.34	2.90	
396	537.50	1.93	
397	539.50	1.72	
398	582.28	2.12	
399	517.20	2.18	
400	511.32	1.04	
401	443.40	2.08	1H NMR (400 MHz, DMSO-d6) δ 13.72 (s, 1H), 12.02 (s, 1H), 8.51 (s, 1H), 7.96 (d, J = 7.8 Hz, 2H), 7.11 (d, J = 7.8 Hz, 1H), 6.90 (s, 2H), 2.25 (s, 3H), 1.94 (s, 6H), 1.04 (s, 9H).
402	525.30	2.16	1H NMR (400 MHz, DMSO-d6) δ 8.21 (d, J = 5.4 Hz, 1H), 7.71 - 7.57 (m, 1H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			7.55 - 7.43 (m, 2H), 7.29 - 7.23 (m, 1H), 7.17 (s, 1H), 7.00 (s, 1H), 6.54 (s, 1H), 6.32 (s, 2H), 5.27 (p, J = 6.1 Hz, 1H), 2.87 (s, 2H), 2.16 - 2.07 (m, 1H), 1.82 (dd, J = 11.7, 5.6 Hz, 1H), 1.62 (s, 3H), 1.55 (s, 3H), 1.37 (t, J = 12.0 Hz, 1H), 1.31 (d, J = 6.2 Hz, 6H), 0.79 (d, J = 6.3 Hz, 3H).
403	526.20	1.56	
404	491.00	2.06	1H NMR (400 MHz, DMSO) δ 12.38 (s, 1H), 7.63 (s, 1H), 7.49 (d, J = 6.2 Hz, 2H), 7.18 (d, J = 7.3 Hz, 1H), 7.02 (d, J = 10.4 Hz, 1H), 6.75 (d, J = 9.0 Hz, 3H), 6.69 (d, J = 8.4 Hz, 1H), 6.46 (s, 2H), 5.66 (s, 2H), 2.90 (s, 1H), 2.18 (s, 3H), 1.96 (s, 6H), 1.19 (d, J = 6.9 Hz, 6H).
405	511.00	2.00	
406	495.00	1.27	1H NMR (400 MHz, DMSO) δ 9.74 (s, 1H), 7.77 (s, 1H), 7.43 (d, J = 8.5 Hz, 1H), 7.17 (d, J = 7.1 Hz, 1H), 6.91 (s, 6H), 4.39 - 4.21 (m, 3H), 4.08 (d, J = 13.2 Hz, 1H), 3.71 (s, 5H), 3.17 (s, 1H), 2.40 (s, 6H), 2.24 (s, 3H), 1.81 (s, 3H).
407	410.30	1.76	
408	542.40	1.98	
409	489.00	1.67	1H NMR (400 MHz, DMSO) δ 12.66 (s, 1H), 8.14 (d, J = 8.8 Hz, 2H), 7.95 (dd, J = 22.2, 8.2 Hz, 2H), 7.64 (s, 1H), 7.47 (d, J = 8.1 Hz, 2H), 7.15 (d, J = 3.9 Hz, 3H), 7.06 (d, J = 8.9 Hz, 2H), 6.81 (d, J = 8.4 Hz, 1H), 4.11 (d, J = 7.0 Hz, 2H), 2.33 (s, 3H), 1.38 (s, 3H).
410	510.00	1.37	1H NMR (400 MHz, DMSO) δ 12.27 (s, 1H), 10.44 (s, 1H), 8.11 (s, 1H), 7.71 (s, 1H), 7.17 (s, 1H), 6.89 (s, 2H), 6.81 (s, 2H), 6.35 (s, 1H), 4.01 (d, J = 16.8 Hz, 4H), 3.76 (s, 2H), 3.17 (s, 4H), 2.39 (s, 6H), 2.24 (s, 3H), 2.07 (s, 1H), 1.62 (s, 3H), 1.24 (s, 3H).
411	417.00	1.90	1H NMR (400 MHz, MeOD) δ 8.21 (s, 1H), 7.91 (s, 1H), 7.69 (s, 1H), 7.48 (d, J = 7.2 Hz, 1H), 7.07 (d, J = 8.8 Hz, 1H), 3.16 (s, 1H), 1.74 (d, J = 21.2 Hz, 7H), 1.46 (s, 9H), 1.26 (s, 3H).

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
412	552.60	1.08	
413	519.42	2.97	
414	468.30	1.62	
415	516.22	1.63	1H NMR (400 MHz, DMSO) δ 13.77 (s, 1H), 13.32 (s, 1H), 7.97 (s, 1H), 7.80 (d, J = 7.9 Hz, 1H), 7.53 - 7.40 (m, 3H), 6.90 (dt, J = 10.8, 2.3 Hz, 1H), 6.86 (d, J = 2.1 Hz, 1H), 4.01 (d, J = 5.0 Hz, 2H), 3.84 (d, J = 6.6 Hz, 2H), 3.31 (s, 3H), 2.51 (dq, J = 7.3, 1.8 Hz, 1H), 2.07 - 1.94 (m, 4H), 1.67 (d, J = 3.6 Hz, 2H), 1.06 (d, J = 6.2 Hz, 6H), 0.99 (d, J = 6.7 Hz, 7H).
416	557.36	2.41	
417	429.40	1.61	
418	542.40	2.80	
419	572.32	2.03	
420	540.40	2.17	1H NMR (400 MHz, DMSO-d6) δ 12.45 (s, 1H), 7.76 - 7.59 (m, 2H), 7.50 - 7.40 (m, 3H), 7.24 (d, J = 7.1 Hz, 1H), 6.91 (dt, J = 10.7, 2.3 Hz, 1H), 6.72 (d, J = 8.6 Hz, 1H), 6.48 (s, 2H), 4.23 (p, J = 2.3 Hz, 2H), 3.85 (d, J = 6.5 Hz, 2H), 2.12 - 1.97 (m, 1H), 1.68 - 1.57 (m, 4H), 1.47 - 1.36 (m, 4H), 1.00 (d, J = 6.7 Hz, 6H).
421	528.50	1.90	
422	543.36	1.95	
423	510.00	2.38	1H NMR (400 MHz, DMSO) δ 12.31 (s, 1H), 8.47 (s, 1H), 8.33 (d, J = 7.6 Hz, 1H), 7.69 (s, 1H), 7.26 (s, 1H), 6.89 (s, 2H), 3.94 (s, 3H), 2.66 (s, 2H), 2.25 (s, 3H), 1.92 (s, 6H), 1.65 (d, J = 35.1 Hz, 4H), 0.91 (s, 6H).
424	570.40	2.66	
425	542.30	2.54	
426	570.12	1.98	
427	542.00	1.74	1H NMR (400 MHz, MeOD) δ 8.27 (s, 1H), 8.21 (d, J = 9.4 Hz, 1H), 7.40 (s, 9H), 7.05 (s, 1H), 6.89 (s, 2H), 5.12 (s, 1H), 4.52 (s, 4H), 3.81 (s, 3H), 2.39 (s, 6H), 2.25 (s, 4H).
428	523.30	2.01	1H NMR (400 MHz, DMSO-d6) δ 8.77 (d, J = 2.4 Hz, 1H), 8.27 (dd, J = 8.7, 2.5 Hz, 1H), 7.54 (d, J = 7.5 Hz, 1H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			7.46 (t, J = 7.8 Hz, 1H), 7.01 (d, J = 7.0 Hz, 2H), 6.82 (d, J = 8.7 Hz, 1H), 6.49 (d, J = 8.3 Hz, 1H), 6.15 (s, 2H), 5.24 - 5.11 (m, 1H), 4.25 (s, 2H), 2.42 (dddd, J = 10.5, 9.4, 5.2, 2.4 Hz, 2H), 2.07 (qdd, J = 9.8, 7.8, 2.7 Hz, 2H), 1.96 - 1.74 (m, 3H), 1.62 (dddd, J = 29.1, 15.5, 11.3, 8.4 Hz, 3H), 1.06 (d, J = 6.1 Hz, 6H).
429	531.00	2.09	1H NMR (400 MHz, DMSO) δ 12.49 (s, 1H), 8.10 (s, 1H), 7.90 (s, 1H), 7.63 (s, 1H), 7.37 - 7.20 (m, 5H), 6.91 (s, 2H), 6.69 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 2.52 (s, 2H), 2.25 (s, 3H), 2.00 (s, 6H), 1.39 (s, 2H), 0.73 (s, 3H).
430	439.20	1.85	
431	526.13	1.91	
432	525.33	1.70	
433	531.18	1.95	
434	520.50	2.00	1H NMR (400 MHz, DMSO-d6) δ 8.31 (d, J = 2.1 Hz, 1H), 8.03 (d, J = 7.9 Hz, 1H), 7.89 (d, J = 2.2 Hz, 1H), 7.78 - 7.73 (m, 2H), 7.66 (d, J = 8.0 Hz, 1H), 7.16 - 7.04 (m, 4H), 6.97 - 6.91 (m, 2H), 4.05 (q, J = 7.0 Hz, 2H), 2.32 (s, 3H), 2.04 (s, 3H), 1.31 (t, J = 6.9 Hz, 3H).
435	539.30	1.81	
436	547.30	2.20	
437	524.50	0.90	
438	557.26	2.42	
439	525.20	2.04	
440	538.24	1.97	
441	511.40	1.59	1H NMR (400 MHz, DMSO-d6) δ 12.47 (s, 1H), 8.85 (d, J = 2.4 Hz, 1H), 8.31 (dd, J = 8.8, 2.5 Hz, 1H), 7.69 (d, J = 7.9 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.19 (dd, J = 9.7, 7.6 Hz, 2H), 6.84 (d, J = 8.7 Hz, 1H), 6.71 (d, J = 8.3 Hz, 1H), 5.31 (p, J = 6.2 Hz, 1H), 4.41 (t, J = 6.6 Hz, 1H), 3.12 - 2.97 (m, 1H), 2.86 - 2.72 (m, 1H), 2.26 - 2.11 (m, 1H), 1.89 (s, 1H), 1.70 - 1.52 (m, 1H), 1.32 (d, J = 6.1 Hz, 6H), 1.04 (d, J = 6.4 Hz, 3H), 0.95 (d, J = 6.8 Hz, 3H).

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
442	516.21	1.67	
443	482.00	2.41	1H NMR (400 MHz, DMSO) δ 8.01 (s, 2H), 7.70 (d, J = 34.2 Hz, 3H), 6.96 (d, J = 10.2 Hz, 3H), 5.43 (s, 2H), 2.09 (s, 6H), 1.21 (s, 9H), 1.15 (s, 9H).
444	451.40	1.63	
445	470.35	1.60	1H NMR (400 MHz, DMSO) δ 8.11 - 8.02 (m, J = 8.9 Hz, 2H), 7.97 (d, J = 2.4 Hz, 1H), 7.84 (d, J = 8.0 Hz, 1H), 7.46 (d, J = 8.0 Hz, 1H), 7.07 - 6.99 (m, J = 8.9 Hz, 2H), 6.85 (d, J = 2.4 Hz, 1H), 4.09 (q, J = 7.0 Hz, 2H), 3.70 - 3.55 (m, J = 12.7 Hz, 2H), 2.97 - 2.81 (m, J = 11.6 Hz, 2H), 1.72 - 1.62 (m, J = 13.0 Hz, 2H), 1.62 - 1.50 (m, 1H), 1.35 (t, J = 7.0 Hz, 3H), 1.32 - 1.16 (m, 2H), 0.94 (d, J = 6.4 Hz, 3H).
446	499.60	1.76	
447	590.33	2.18	
448	492.00	1.59	1H NMR (400 MHz, DMSO) δ 11.09 (s, 1H), 7.64 (s, 1H), 7.33 (s, 4H), 7.20 (s, 1H), 7.11 (s, 1H), 6.72 (s, 2H), 4.58 (s, 1H), 4.16 (s, 2H), 3.91 (s, 2H), 2.22 (s, 3H), 2.19 (s, 3H), 2.12 (s, 6H), 1.60 (s, 3H), 1.48 (s, 3H).
449	521.22	2.59	
450	519.30	2.01	
451	543.43	2.65	
452	543.29	2.35	
453	512.40	2.02	
454	537.30	2.16	1H NMR (400 MHz, DMSO-d6) δ 8.77 (d, J = 2.5 Hz, 1H), 8.27 (dd, J = 8.6, 2.5 Hz, 1H), 7.60 - 7.42 (m, 2H), 7.00 (t, J = 6.7 Hz, 2H), 6.88 (d, J = 8.6 Hz, 1H), 6.56 - 6.46 (m, 1H), 6.25 (s, 2H), 4.23 (bs, 2H), 4.10 (s, 2H), 1.86 (s, 2H), 1.56 (bs, 2H), 1.17 (bs, 3H), 1.03 (d, J = 6.1 Hz, 6H), 0.58 - 0.49 (m, 2H), 0.43 - 0.34 (m, 2H).
455	524.30	1.51	
456	480.24	1.75	
457	561.14	2.97	
458	542.30	1.92	1H NMR (400 MHz, DMSO-d6) δ 12.59 (s, 1H), 7.75 (d, J = 8.0 Hz, 1H), 7.69 - 7.60 (m, 1H), 7.52 - 7.45 (m,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 7.42 (d, J = 10.0 Hz, 1H), 7.32 (d, J = 7.9 Hz, 1H), 7.23 (d, J = 7.3 Hz, 1H), 6.93 - 6.84 (m, 1H), 6.72 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 4.13 (s, 2H), 3.84 (d, J = 6.5 Hz, 2H), 2.11 - 1.99 (m, 3H), 1.52 - 1.45 (m, 2H), 1.04 - 0.95 (m, 12H).
459	486.20	1.43	
460	456.21	1.17	
461	608.00	2.23	1H NMR (400 MHz, DMSO) δ 13.78 (s, 1H), 13.05 (s, 1H), 8.36 (d, J = 28.4 Hz, 3H), 8.16 (d, J = 7.8 Hz, 1H), 7.82 (d, J = 7.9 Hz, 1H), 7.28 - 7.16 (m, 2H), 6.93 (s, 2H), 6.84 (d, J = 10.7 Hz, 1H), 3.65 (d, J = 6.5 Hz, 2H), 2.26 (s, 3H), 2.02 (d, J = 42.5 Hz, 9H), 0.97 (d, J = 6.7 Hz, 6H).
462	451.50	1.72	
463	507.41	3.02	
464	525.30	1.57	
465	588.50	1.31	1H NMR (400 MHz, DMSO-d6) δ 8.10 (d, J = 7.8 Hz, 1H), 7.67 (s, 1H), 7.61 (d, J = 7.9 Hz, 1H), 7.54 (s, 1H), 7.45 - 7.40 (m, 1H), 7.35 - 7.22 (m, 2H), 7.11 - 7.05 (m, 1H), 6.91 (s, 2H), 6.48 - 6.43 (m, 1H), 6.08 (s, 2H), 3.55 (t, J = 4.6 Hz, 4H), 3.42 (s, 2H), 2.55 (s, 1H), 2.32 - 2.27 (m, 5H), 2.27 (s, 3H), 2.03 (s, 6H).
466	547.30	1.87	1H NMR (400 MHz, DMSO-d6) δ 8.27 (d, J = 19.3 Hz, 1H), 7.94 (d, J = 8.0 Hz, 1H), 7.72 (d, J = 8.0 Hz, 1H), 7.51 (s, 1H), 7.43 (d, J = 10.1 Hz, 1H), 7.28 (dd, J = 14.4, 8.0 Hz, 1H), 6.90 (dt, J = 10.6, 2.3 Hz, 1H), 4.61 - 4.44 (m, 1H), 3.84 (d, J = 6.7 Hz, 2H), 3.40 - 2.86 (m, 4H), 2.16 - 1.91 (m, 2H), 1.71 - 1.60 (m, 7H), 1.60 - 1.49 (m, 1H), 1.16 (s, 1H), 1.06 - 0.96 (m, 10H).
467	479.32	1.20	
468	497.51	3.50	
469	521.52	2.73	
470	510.50	1.42	
471	539.50	1.66	
472	474.00	2.36	1H NMR (400 MHz, DMSO) δ 7.88

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(dd, J = 10.8, 4.8 Hz, 2H), 7.77 (d, J = 8.0 Hz, 1H), 7.63 (t, J = 8.0 Hz, 1H), 6.88 (s, 1H), 6.67 (s, 2H), 5.37 (s, 2H), 3.11 (s, 1H), 2.09 (s, 3H), 2.00 (d, J = 6.5 Hz, 6H), 1.08 (s, 9H).
473	604.40	3.25	
474	495.00	1.18	1H NMR (400 MHz, DMSO) δ 9.96 (s, 1H), 9.61 (s, 1H), 7.90 (s, 1H), 7.44 - 6.61 (m, 7H), 4.39 (s, 3H), 4.16 (s, 2H), 3.74 (s, 3H), 2.41 (s, 5H), 2.24 (s, 3H), 1.97 (d, J = 80.3 Hz, 3H).
475	574.34	3.13	
476	509.60	1.54	1H NMR (400 MHz, DMSO-d6) δ 12.39 (s, 1H), 8.85 (d, J = 2.5 Hz, 1H), 8.31 (dd, J = 8.7, 2.5 Hz, 1H), 7.70 (d, J = 7.9 Hz, 1H), 7.64 (dd, J = 8.4, 7.2 Hz, 1H), 7.19 (dd, J = 14.2, 7.6 Hz, 2H), 6.83 (d, J = 8.7 Hz, 1H), 6.71 (d, J = 8.2 Hz, 1H), 6.45 (s, 1H), 5.30 (h, J = 6.2 Hz, 1H), 4.56 (s, 1H), 2.98 (d, J = 9.7 Hz, 1H), 2.44 (s, 1H), 2.35 - 2.26 (m, 1H), 1.71 - 1.13 (m, 7H) 1.32 (d, J = 96.2 Hz, 6H).
477	533.20	2.06	
478	415.00	1.48	1H NMR (400 MHz, MeOD) δ 8.56 (d, J = 8.4 Hz, 1H), 7.98 (d, J = 8.4 Hz, 1H), 7.87 (s, 1H), 7.43 (d, J = 7.2 Hz, 1H), 7.03 (d, J = 8.7 Hz, 1H), 6.08 (s, 1H), 2.31 (s, 2H), 2.20 (s, 2H), 1.69 (d, J = 17.3 Hz, 4H), 1.53 (s, 9H).
479	528.31	1.99	
480	466.00	1.26	
481	605.40	2.21	
482	469.27	2.05	
483	556.30	1.76	1H NMR (400 MHz, DMSO-d6) δ 13.97 (s, 1H), 7.84 (s, 1H), 7.65 - 7.60 (m, 1H), 7.23 - 7.18 (m, 1H), 6.99 - 6.94 (m, 2H), 6.89 (d, J = 11.3 Hz, 1H), 6.72 - 6.67 (m, 1H), 6.42 (s, 2H), 3.92 - 3.87 (m, 2H), 3.81 (d, J = 6.5 Hz, 2H), 2.30 (s, 3H), 2.09 - 1.98 (m, 1H), 1.65 - 1.60 (m, 2H), 1.07 (d, J = 6.2 Hz, 6H), 0.98 (d, J = 6.7 Hz, 6H).
484	610.40	2.10	1H NMR (400 MHz, Methanol-d4) δ 8.25 (d, J = 2.8 Hz, 1H), 7.88 (d, J = 8.3

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			Hz, 1H), 7.69 (dd, J = 8.7, 7.2 Hz, 1H), 7.31 (d, J = 7.2 Hz, 1H), 6.98 (dd, J = 8.5, 6.5 Hz, 2H), 5.99 (d, J = 2.8 Hz, 1H), 3.90 (s, 2H), 3.63 (q, J = 6.1, 5.3 Hz, 4H), 2.80 - 2.66 (m, 2H), 2.26 (d, J = 10.4 Hz, 1H), 1.90 (dd, J = 11.9, 5.6 Hz, 1H), 1.68 (q, J = 5.1, 4.7 Hz, 2H), 1.61 (d, J = 5.3 Hz, 10H), 1.47 (t, J = 12.2 Hz, 1H), 1.04 (s, 9H), 0.88 (d, J = 6.3 Hz, 3H).
485	549.43	2.85	
486	525.30	1.41	1H NMR (400 MHz, DMSO-d6) δ 14.19 (s, 1H), 8.46 (d, J = 2.5 Hz, 1H), 7.98 (dd, J = 8.6, 2.5 Hz, 1H), 7.88 (s, 1H), 7.62 (t, J = 7.9 Hz, 1H), 7.20 (d, J = 7.2 Hz, 1H), 6.84 (d, J = 8.6 Hz, 1H), 6.69 (d, J = 8.3 Hz, 1H), 6.40 (s, 2H), 5.32 (p, J = 6.2 Hz, 1H), 3.97 - 3.85 (m, 2H), 2.36 (s, 3H), 2.11 - 2.00 (m, 2H), 1.71 - 1.57 (m, 2H), 1.33 (d, J = 6.2 Hz, 6H), 1.08 (d, J = 6.3 Hz, 6H).
487	529.18	1.44	
488	604.40	323.20	
489	574.33	1.33	
490	594.30	3.20	1H NMR (400 MHz, Methanol-d4) δ 8.16 (d, J = 8.1 Hz, 1H), 7.74 - 7.62 (m, 2H), 7.49 (t, J = 1.9 Hz, 1H), 7.45 - 7.36 (m, 2H), 6.79 (ddd, J = 8.5, 4.5, 1.9 Hz, 2H), 5.45 (p, J = 6.1 Hz, 1H), 3.89 - 3.77 (m, 2H), 3.62 (d, J = 9.3 Hz, 1H), 3.16 (dd, J = 9.2, 4.8 Hz, 1H), 2.16 - 1.95 (m, 2H), 1.89 (tt, J = 8.1, 4.3 Hz, 1H), 1.15 (s, 1H), 1.06 (dd, J = 6.8, 2.3 Hz, 6H), 0.85 (td, J = 8.1, 5.8 Hz, 1H).
491	525.30	2.23	1H NMR (400 MHz, DMSO-d6) δ 8.20 (d, J = 5.4 Hz, 1H), 7.61 (d, J = 7.5 Hz, 1H), 7.56 - 7.43 (m, 2H), 7.32 (d, J = 1.4 Hz, 1H), 7.17 (d, J = 7.6 Hz, 1H), 6.99 (d, J = 7.1 Hz, 1H), 6.53 (d, J = 8.3 Hz, 1H), 6.32 (s, 2H), 4.24 (t, J = 6.7 Hz, 2H), 2.89 (dd, J = 15.6, 8.8 Hz, 2H), 2.09 (d, J = 13.2 Hz, 1H), 1.87 - 1.67 (m, 3H), 1.62 (s, 3H), 1.55 (s, 3H), 1.36 (t, J = 11.9 Hz, 1H), 0.97 (t, J = 7.4 Hz, 3H), 0.78 (d, J = 6.3 Hz, 3H).

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
492	557.50	1.89	
493	584.40	3.12	1H NMR (400 MHz, DMSO-d6) δ 12.44 (s, 1H), 7.84 (d, J = 7.9 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.53 (t, J = 1.8 Hz, 1H), 7.45 (dt, J = 10.0, 2.0 Hz, 1H), 7.36 (d, J = 8.0 Hz, 1H), 7.21 (d, J = 7.2 Hz, 1H), 6.88 (dt, J = 10.6, 2.3 Hz, 1H), 6.70 (d, J = 8.2 Hz, 1H), 6.44 (s, 2H), 5.12 (dd, J = 8.5, 3.7 Hz, 1H), 3.83 (qd, J = 9.2, 6.6 Hz, 2H), 2.81 (dt, J = 14.0, 7.2 Hz, 1H), 2.67 (t, J = 9.3 Hz, 1H), 2.10 - 1.97 (m, 1H), 1.87 - 1.63 (m, 4H), 1.26 (dp, J = 21.5, 7.3 Hz, 2H), 0.99 (dd, J = 6.6, 1.3 Hz, 6H), 0.83 (t, J = 7.5 Hz, 3H), 0.74 (d, J = 12.5 Hz, 6H).
494	548.20	2.08	
495	537.34	1.23	
496	556.40	2.54	1H NMR (400 MHz, Methanol-d4) δ 8.60 (d, J = 8.3 Hz, 1H), 8.08 (d, J = 8.3 Hz, 1H), 7.70 (dd, J = 8.6, 7.3 Hz, 1H), 7.50 (t, J = 1.9 Hz, 1H), 7.48 - 7.41 (m, 1H), 7.36 (dd, J = 7.4, 0.8 Hz, 1H), 6.85 (dt, J = 10.5, 2.3 Hz, 1H), 6.80 (dd, J = 8.5, 0.8 Hz, 1H), 4.59 (h, J = 7.2 Hz, 1H), 3.84 (d, J = 6.4 Hz, 2H), 2.61 - 2.47 (m, 1H), 2.30 - 2.19 (m, 2H), 2.17 - 1.92 (m, 2H), 1.53 (s, 3H), 1.37 (s, 3H), 1.33 (d, J = 6.5 Hz, 3H), 1.06 (d, J = 6.7 Hz, 6H).
497	526.50	1.50	
498	544.22	1.94	
499	551.53	2.74	
500	525.25	1.49	
501	504.20	2.33	1H NMR (400 MHz, DMSO) δ 12.83 (s, 1H), 8.44 - 8.28 (m, 2H), 8.04 (d, J = 6.5 Hz, 1H), 7.95 (d, J = 7.8 Hz, 1H), 7.11 (dd, J = 31.1, 23.3 Hz, 2H), 6.89 (d, J = 3.4 Hz, 2H), 2.24 (s, 3H), 1.92 (s, 6H), 1.04 (s, 9H).
502	573.30	1.97	
503	557.26	2.39	
504	558.07	2.02	
505	523.32	2.65	1H NMR (400 MHz, DMSO) δ 12.43 (s, 1H), 10.17 (s, 1H), 8.14 - 8.11 (d,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 7.70 - 7.63 (m, 2H), 7.25 - 7.22 (d, 1H), 7.02 - 6.96 (m, 4H), 6.73 - 7.60 (d, 1H), 6.64 - 6.58 (m, 2H), 2.28 (s, 3H), 2.03 (s, 6H)
506	606.15	2.02	
507	497.31	1.64	1H NMR (400 MHz, DMSO) δ 8.31 (d, J = 2.2 Hz, 1H), 8.12 - 8.04 (m, J = 8.9 Hz, 2H), 7.94 - 7.86 (m, J = 5.1, 2.9 Hz, 2H), 7.51 (d, J = 8.1 Hz, 1H), 7.03 (d, J = 8.9 Hz, 2H), 4.10 (q, J = 7.0 Hz, 2H), 3.67 - 3.61 (m, J = 12.5 Hz, 2H), 3.03 - 2.89 (m, J = 11.6 Hz, 2H), 1.73 - 1.51 (m, J = 22.5, 9.5 Hz, 3H), 1.35 (t, J = 7.0 Hz, 3H), 1.33 - 1.18 (m, 2H), 0.94 (d, J = 6.4 Hz, 3H).
508	528.00	1.45	
509	559.20	1.87	
510	570.44	3.43	1H NMR (400 MHz, DMSO) δ 7.82 - 7.79 (d, 1H), 7.63 - 7.57 (m, 2H), 7.52 - 7.46 (m, 3H), 7.35 - 7.32 (d, 1H), 6.95 - 6.91 (d, 1H), 3.85 (s, 3H), 3.85 - 3.83 (d, 2H), 3.17 (m, 4H), 2.10 - 2.00 (m, 1H), 1.30 (m, 4H), 1.01 - 0.98 (d, 6H), 0.94 (s, 6H).
511	525.40	1.62	
512	440.40	1.93	
513	504.41	1.87	
514	508.31	2.51	
515	468.10	5.33	0.96 (s, 9H), 1.90 (s, 6H), 2.17 (s, 3H), 5.78 (br. s., 2H), 6.59 (d, J=8.2 Hz, 1H), 6.71 (s, 2H), 6.82 (d, J=7.8 Hz, 1H), 7.19-7.31 (m, 3H), 7.36-7.47 (m, 1H),8.15 (d, J=7.8 Hz, 1H).
516	542.30	2.58	
517	520.40	1.83	1H NMR (400 MHz, DMSO-d6) δ 12.56 (s, 1H), 12.06 (s, 1H), 8.26 (d, J = 7.3 Hz, 1H), 8.08 (d, J = 8.0 Hz, 1H), 7.84 (d, J = 5.6 Hz, 1H), 7.79 - 7.73 (m, 2H), 7.69 (d, J = 8.0 Hz, 1H), 7.16 - 7.04 (m, 3H), 6.97 - 6.91 (m, 2H), 6.46 (t, J = 6.7 Hz, 1H), 4.05 (q, J = 6.9 Hz, 2H), 2.32 (s, 3H), 2.07 (s, 3H), 1.31 (t, J = 7.0 Hz, 3H).
518	481.00	1.90	1H NMR (400 MHz, DMSO) δ 13.82 (s, 1H), 12.52 (s, 1H), 8.08 (d, J = 7.8 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 7.99 (s, 1H), 7.81 (s, 2H), 7.74 (d, J = 7.9 Hz, 1H), 7.26 (s, 2H), 6.95 (s, 2H), 6.89 (s, 1H), 2.28 (s, 3H), 2.01 (s, 6H).
519	544.19	1.75	
520	539.30	1.85	
521	509.18	1.34	
522	539.23	1.77	
523	538.30	1.58	1H NMR (400 MHz, DMSO-d6) δ 8.00 (d, J = 5.3 Hz, 1H), 7.72 - 7.57 (m, 1H), 7.54 - 7.36 (m, 1H), 7.13 - 6.91 (m, 4H), 6.60 (s, 1H), 6.49 (s, 1H), 6.15 (s, 1H), 3.06 (t, J = 6.4 Hz, 2H), 3.00 - 2.86 (m, 1H), 2.25 - 2.07 (m, 0H), 1.85 (ddd, J = 16.9, 12.2, 6.0 Hz, 2H), 1.61 (s, 3H), 1.57 (s, 3H), 1.39 (t, J = 11.8 Hz, 1H), 1.26 - 1.12 (m, 2H), 0.91 (d, J = 6.7 Hz, 6H), 0.86 (d, J = 6.2 Hz, 3H).
524	474.00	1.85	1H NMR (400 MHz, DMSO) δ 13.10 (s, 1H), 8.56 (d, J = 4.9 Hz, 1H), 8.25 (d, J = 9.4 Hz, 1H), 8.08 (d, J = 8.2 Hz, 2H), 8.00 - 7.92 (m, 2H), 7.35 (d, J = 4.4 Hz, 4H), 7.24 (s, 1H), 7.04 (d, J = 8.0 Hz, 2H), 4.04 (s, 3H), 2.38 (s, 3H), 2.31 (s, 3H).
525	534.20	1.90	
526	497.50	1.25	
527	540.30	2.99	
528	530.22	1.86	
529	401.00	1.40	1H NMR (400 MHz, MeOD) δ 8.48 (d, J = 8.4 Hz, 1H), 7.99 - 7.82 (m, 2H), 7.44 (d, J = 7.3 Hz, 1H), 7.07 (d, J = 8.0 Hz, 1H), 6.30 (s, 1H), 2.71 (s, 2H), 2.54 (s, 2H), 2.01 (s, 2H).
530	570.50	2.12	1H NMR (400 MHz, DMSO-d6) δ 7.78 (d, J = 7.9 Hz, 1H), 7.65 (dd, J = 8.4, 7.3 Hz, 1H), 7.48 (t, J = 1.8 Hz, 1H), 7.43 (dt, J = 10.0, 1.9 Hz, 1H), 7.38 (d, J = 7.9 Hz, 1H), 7.22 (d, J = 7.1 Hz, 1H), 6.90 (dt, J = 10.6, 2.3 Hz, 1H), 6.71 (d, J = 8.4 Hz, 1H), 6.46 (s, 1H), 3.84 (d, J = 6.6 Hz, 2H), 3.29 - 3.15 (m, 2H), 3.15 - 2.96 (m, 2H), 2.05 (hept, J = 6.7 Hz, 1H), 1.43 (s, 2H), 1.38 - 1.16 (m, 4H), 0.99 (d, J = 6.7

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			Hz, 6H), 0.80 (s, 3H), 0.76 (t, J = 7.4 Hz, 3H).
531	520.32	2.85	
532	452.20	2.05	
533	520.50	1.89	1H NMR (400 MHz, DMSO-d6) δ 8.32 (d, J = 2.3 Hz, 1H), 8.08 (d, J = 7.7 Hz, 1H), 7.92 (dd, J = 2.3, 1.0 Hz, 1H), 7.54 (d, J = 7.7 Hz, 1H), 7.43 (dd, J = 8.2, 7.3 Hz, 1H), 7.10 (d, J = 2.1 Hz, 1H), 7.08 - 7.00 (m, 2H), 6.94 (d, J = 8.1 Hz, 1H), 6.46 (dd, J = 8.3, 0.9 Hz, 1H), 6.12 (s, 2H), 3.87 (s, 3H), 2.31 (s, 3H), 2.12 (s, 3H), 2.04 (s, 3H).
534	485.60	1.63	
535	581.30	2.34	1H NMR (400 MHz, DMSO-d6) δ 12.32 (s, 1H), 8.15 (d, J = 7.8 Hz, 1H), 7.80 (d, J = 7.9 Hz, 1H), 7.65 (t, J = 7.9 Hz, 1H), 7.52 - 7.41 (m, 2H), 7.40 (d, J = 1.7 Hz, 1H), 7.23 (d, J = 7.3 Hz, 1H), 6.95 (s, 2H), 6.71 (d, J = 8.4 Hz, 1H), 6.53 (s, 2H), 3.80 (t, J = 6.4 Hz, 2H), 2.27 (s, 3H), 2.02 (s, 6H), 1.73 (h, J = 7.1 Hz, 2H), 1.00 (t, J = 7.4 Hz, 3H).
536	524.50	1.50	
537	416.00	1.96	1H NMR (400 MHz, MeOD) δ 7.87 (d, J = 8.8 Hz, 3H), 7.49 (s, 1H), 6.93 (d, J = 8.7 Hz, 2H), 2.82 (s, 1H), 1.62 (s, 8H), 1.39 (s, 9H), 1.18 (s, 3H).
538	523.30	2.38	
539	480.19	1.46	
540	427.50	1.84	
541	541.36	1.58	
542	410.40	1.64	
543	592.35	2.77	
544	525.40	1.57	
545	526.10	1.89	
546	526.20	1.92	
547	556.40	2.66	1H NMR (400 MHz, DMSO-d6) δ 7.53 - 7.41 (m, 3H), 7.35 (ddd, J = 10.4, 2.3, 1.2 Hz, 1H), 7.08 (d, J = 7.6 Hz, 1H), 6.98 (d, J = 7.3 Hz, 1H), 6.79 (dt, J = 10.8, 2.3 Hz, 1H), 6.49 (d, J = 8.3 Hz, 1H), 6.14 (s, 2H), 3.82 (d, J = 6.6 Hz, 2H), 3.29 - 3.21 (m, 1H), 2.88 (t, J = 9.5 Hz, 1H), 2.05 (hept, J = 6.6 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 1.80 (dp, J = 13.3, 6.8 Hz, 1H), 1.66 (q, J = 6.5, 5.3 Hz, 1H), 1.59 (s, 3H), 1.48 - 1.36 (m, 1H), 1.33 (s, 3H), 0.99 (d, J = 6.7 Hz, 6H), 0.94 (d, J = 6.8 Hz, 3H).
548	557.10	2.21	1H NMR (400 MHz, DMSO-d6) δ 8.10 (d, J = 7.7 Hz, 1H), 7.90 (d, J = 8.2 Hz, 2H), 7.73 (d, J = 8.1 Hz, 2H), 7.66 (d, J = 7.7 Hz, 1H), 7.42 (t, J = 7.9 Hz, 1H), 7.08 (d, J = 7.3 Hz, 1H), 6.91 (s, 2H), 6.46 (d, J = 8.2 Hz, 1H), 6.06 (s, 2H), 3.42 (s, 13H), 2.27 (s, 3H), 2.03 (s, 6H).
549	579.45	3.38	1H NMR (400 MHz, DMSO-d6) δ 12.74 (s, 1H), 12.19 (s, 1H), 7.77 - 7.69 (m, 2H), 7.56 - 7.53 (d, 1H), 7.48 (s, 1H), 7.43 - 7.39 (d, 1H), 7.34 - 7.23 (m, 3H), 7.12 (t, J = 7.47, 1H), 6.92 - 6.88 (d, 1H), 3.84 - 3.81 (d, 2H), 2.09 - 1.98 (m, 2H), 1.81 - 1.73 (m, 1H), 1.58 (s, 3H), 1.53 (s, 3H), 1.30 (t, J = 12.41, 1H), 1.00 - 0.97 (d, 6H), 0.30 - 0.27 (d, 3H)
550	440.40	1.63	
551	514.30	1.48	
552	566.20	2.15	
553	544.34	2.20	
554	527.00	1.41	
555	556.50	1.95	1H NMR (400 MHz, Methanol-d4) δ 7.93 (d, J = 8.0 Hz, 1H), 7.49 (t, J = 1.8 Hz, 1H), 7.42 - 7.30 (m, 2H), 7.26 (d, J = 8.0 Hz, 1H), 6.79 - 6.66 (m, 2H), 3.81 (d, J = 6.5 Hz, 2H), 2.91 (d, J = 7.5 Hz, 1H), 2.85 (d, J = 10.1 Hz, 1H), 2.37 (s, 1H), 1.64 (s, 6H), 1.58 (d, J = 12.0 Hz, 1H), 1.05 (d, J = 6.7 Hz, 6H), 1.00 (d, J = 6.4 Hz, 3H).
556	541.18	1.73	
557	525.30	1.50	
558	609.20	2.06	
559	552.50	2.11	
560	570.30	3.05	
561	508.20	1.88	
562	583.22	2.25	
563	524.40	1.67	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
564	479.20	0.69	
565	524.17	1.83	
566	441.00	1.94	1H NMR (400 MHz, DMSO) δ 7.28 (d, J = 9.7 Hz, 2H), 7.15 (s, 1H), 7.00 (s, 1H), 6.92 (s, 1H), 6.77 (s, 2H), 5.45 (s, 2H), 2.75 (s, 1H), 2.18 (s, 3H), 2.09 (s, 6H), 1.10 (d, J = 6.9 Hz, 6H).
567	497.26	1.45	
568	481.20	5.67	1H NMR (300 MHz, CDCl3) δ 1.06 (s, 9H), 2.04 (s, 6H), 2.29 (s, 3H), 2.71 (s, 3H), 6.80-6.86 (m, 3H), 6.90 (d, J=7.8 Hz, 1H), 7.23-7.31 (m, 1H), 7.32-7.41 (m, 3H), 8.25 (d, J=7.8 Hz, 1H).
569	538.60	0.99	
570	507.20	2.01	
571	418.50	1.26	
572	557.40	1.86	
573	522.30	2.25	
574	541.29	1.45	
575	570.40	3.00	
576	617.20	1.69	
577	509.00	1.87	
578	523.30	1.65	1H NMR (400 MHz, DMSO) δ 8.90 (d, J = 2.4 Hz, 1H), 8.37 (dd, J = 8.7, 2.4 Hz, 1H), 7.89 (d, J = 8.0 Hz, 1H), 7.73 - 7.63 (m, 1H), 7.50 (d, J = 8.0 Hz, 1H), 7.22 (d, J = 7.3 Hz, 1H), 6.87 (d, J = 8.7 Hz, 1H), 6.73 (d, J = 8.4 Hz, 1H), 5.32 (dt, J = 12.3, 6.1 Hz, 1H), 3.43 - 3.28 (m, 4H), 2.60 - 2.44 (m, 7H), 1.44 (d, J = 4.5 Hz, 4H), 1.32 (d, J = 6.2 Hz, 6H).
579	524.10	2.03	1H NMR (400 MHz, DMSO-d6) δ 11.75 (s, 1H), 8.10 (d, J = 7.7 Hz, 1H), 8.05 - 7.93 (m, 2H), 7.76 - 7.68 (m, 2H), 7.68 - 7.53 (m, 2H), 7.47 - 7.39 (m, 2H), 6.91 (s, 2H), 6.23 (s, 1H), 2.27 (s, 3H), 2.01 (s, 6H).
580	553.50	1.82	
581	554.30	2.69	
582	547.20	1.65	
583	540.43	3.43	1H NMR (400 MHz, DMSO-d6) δ 12.89 (s, 1H), 8.02 - 7.99 (d, 2H), 7.80 - 7.64 (m, 4H), 7.51 - 7.44 (m, 3H), 6.94 - 6.90 (d, 1H), 3.85 - 3.82 (d, 2H), 3.16 (m, 4H), 2.09 - 1.99 (m, 1H), 1.29

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(m, 4H), 1.01 - 0.98 (d, 6H), 0.93 (s, 6H)
584	507.20	1.64	
585	525.00	2.03	1H NMR (400 MHz, DMSO) δ 13.81 (s, 1H), 12.52 (s, 1H), 8.08 (d, J = 7.8 Hz, 1H), 7.98 (s, 1H), 7.81 (d, J = 7.8 Hz, 1H), 7.24 - 7.14 (m, 2H), 6.96 (s, 2H), 6.92 - 6.80 (m, 2H), 5.75 (s, 1H), 3.93 (d, J = 7.0 Hz, 2H), 2.27 (s, 3H), 2.00 (s, 6H), 1.29 (s, 3H).
586	580.00	2.23	1H NMR (400 MHz, DMSO) δ 12.65 (s, 1H), 8.31 (s, 1H), 8.13 (d, J = 7.9 Hz, 2H), 7.81 (d, J = 7.8 Hz, 1H), 7.18 (dd, J = 29.3, 16.8 Hz, 4H), 6.94 (s, 2H), 6.84 (d, J = 10.7 Hz, 1H), 3.65 (d, J = 6.5 Hz, 2H), 2.27 (s, 3H), 2.00 (s, 7H), 0.96 (s, 7H).
587	542.31	2.09	
588	531.20	1.82	
589	495.18	1.46	
590	565.40	3.28	
591	516.27	2.05	
592	514.30	2.78	
593	554.40	2.70	
594	467.27	2.00	
595	533.50	1.90	
596	523.23	1.80	
597	539.30	1.76	1H NMR (400 MHz, DMSO-d6) δ 12.41 (s, 1H), 8.87 (d, J = 2.5 Hz, 1H), 8.33 (dd, J = 8.8, 2.5 Hz, 1H), 7.83 (d, J = 8.0 Hz, 1H), 7.65 (dd, J = 8.4, 7.3 Hz, 1H), 7.27 (d, J = 8.0 Hz, 1H), 7.21 (d, J = 7.2 Hz, 1H), 6.86 (d, J = 8.7 Hz, 1H), 6.71 (d, J = 8.2 Hz, 1H), 6.45 (s, 2H), 5.32 (p, J = 6.2 Hz, 1H), 5.06 - 4.95 (m, 1H), 2.89 - 2.76 (m, 1H), 2.75 - 2.60 (m, 1H), 1.88 - 1.63 (m, 4H), 1.32 (d, J = 6.2 Hz, 6H), 0.82 (s, 9H).
598	508.20	1.68	
599	551.53	3.14	
600	528.37	1.82	1H NMR (400 MHz, DMSO) δ 7.83 (d, J = 7.9 Hz, 1H), 7.65 (dd, J = 8.4, 7.3 Hz, 1H), 7.48 (d, J = 7.9 Hz, 1H), 7.47 - 7.40 (m, J = 11.0, 1.7 Hz, 2H), 7.21 (d, J = 6.8 Hz, 1H), 6.91 (dt, J = 10.9, 2.2

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			Hz, 1H), 6.72 (d, J = 8.0 Hz, 1H), 4.73 (hept, 1H), 3.78 - 3.63 (m, J = 13.0 Hz, 2H), 2.94 - 2.78 (m, J = 11.6 Hz, 2H), 1.69 - 1.59 (m, J = 12.8 Hz, 2H), 1.58 - 1.49 (m, 1H), 1.30 (d, J = 6.0 Hz, 6H), 1.24 - 1.07 (m, 2H), 0.91 (d, J = 6.4 Hz, 3H).
601	587.50	1.78	
602	551.50	2.05	
603	432.04	2.02	
604	463.50	2.09	
605	557.50	1.89	
606	541.36	1.49	
607	490.50	2.01	1H NMR (400 MHz, DMSO-d6) δ 8.33 (d, J = 2.2 Hz, 1H), 8.05 (d, J = 7.9 Hz, 1H), 7.91 (d, J = 2.2 Hz, 1H), 7.71 (dd, J = 8.1, 1.8 Hz, 3H), 7.22 (d, J = 8.1 Hz, 2H), 7.17 - 7.04 (m, 5H), 2.31 (d, J = 5.4 Hz, 6H), 2.04 (s, 3H).
608	538.20	1.88	
609	554.50	2.23	
610	463.50	1.92	1H NMR (400 MHz, DMSO-d6) δ 13.79 (s, 1H), 12.40 (s, 1H), 8.03 (d, J = 7.9 Hz, 1H), 7.98 (s, 1H), 7.74 - 7.68 (m, 3H), 7.22 (d, J = 8.0 Hz, 2H), 7.13 (dd, J = 5.2, 3.0 Hz, 2H), 7.07 (dd, J = 8.3, 2.1 Hz, 1H), 6.86 (d, J = 2.4 Hz, 1H), 2.32 (s, 3H), 2.30 (s, 3H), 2.05 (s, 3H).
611	570.40	3.23	1H NMR (400 MHz, DMSO-d6) δ 12.48 (s, 1H), 7.76 (d, J = 7.6 Hz, 1H), 7.64 (t, J = 7.8 Hz, 1H), 7.51 - 7.41 (m, 2H), 7.29 (d, J = 7.6 Hz, 1H), 7.22 (d, J = 7.0 Hz, 1H), 6.89 (d, J = 10.9 Hz, 1H), 6.69 (dd, J = 8.3, 4.0 Hz, 1H), 6.48 (s, 2H), 3.84 (d, J = 6.2 Hz, 2H), 3.25 - 3.06 (m, 5H), 2.10 - 1.90 (m, 2H), 1.88 - 1.75 (m, 1H), 1.68 - 1.52 (m, 1H), 0.99 (d, J = 7.1 Hz, 6H), 0.88 (s, 9H).
612	538.50	1.90	1H NMR (400 MHz, DMSO-d6) δ 8.30 (d, J = 1.9 Hz, 1H), 8.08 (d, J = 7.7 Hz, 1H), 7.93 (dd, J = 11.9, 2.0 Hz, 1H), 7.60 (d, J = 7.7 Hz, 1H), 7.42 (t, J = 7.8 Hz, 1H), 7.07 (dd, J = 7.3, 0.8 Hz, 1H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			6.93 (s, 2H), 6.44 (d, J = 8.3 Hz, 1H), 6.08 (s, 2H), 3.94 (s, 3H), 2.39 (s, 1H), 2.28 (s, 3H), 2.01 (s, 6H).
613	570.38	2.17	
614	494.30	1.58	
615	556.60	2.02	
616	553.40	2.39	1H NMR (400 MHz, DMSO-d6) δ 7.67 (d, J = 8.4 Hz, 1H), 7.55 (d, J = 7.6 Hz, 1H), 7.47 (t, J = 7.8 Hz, 1H), 6.98 (d, J = 7.3 Hz, 1H), 6.69 (d, J = 8.4 Hz, 1H), 6.55 (d, J = 7.6 Hz, 1H), 6.49 (d, J = 8.3 Hz, 1H), 6.22 (s, 2H), 4.04 (d, J = 6.7 Hz, 2H), 3.03 - 2.84 (m, 2H), 2.42 (s, 3H), 2.04 (tq, J = 12.4, 5.7 Hz, 2H), 1.76 (dd, J = 11.8, 5.7 Hz, 1H), 1.49 (s, 3H), 1.45 (s, 3H), 1.33 (t, J = 12.0 Hz, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.80 (d, J = 6.3 Hz, 3H).
617	446.50	1.57	
618	559.28	2.08	
619	525.30	1.70	
620	540.43	2.73	1H NMR (400 MHz, DMSO-d6) δ 12.60 (s, 1H), 8.03 - 8.00 (d, 2H), 7.75 - 7.34 (m, 4H), 7.48 (s, 1H), 7.43 - 7.39 (d, 1H), 7.29 - 7.26 (d, 1H), 6.92 - 6.89 (d, 1H), 3.84 - 3.81 (d, 2H), 2.51 - 2.29 (m, 2H), 2.11 - 1.99 (m, 2H), 1.87 - 1.80 (m, 1H), 1.59 (s, 3H), 1.55 (s, 3H), 1.38 (t, J = 11.76, 1H), 1.00 - 0.97 (d, 6H), 0.67 - 0.64 (d, 3H)
621	521.30	2.06	1H NMR (400 MHz, DMSO) δ 12.34 (s, 1H), 8.08 (d, J = 7.8 Hz, 1H), 7.63 (dd, J = 8.3, 7.4 Hz, 1H), 7.26 (d, J = 7.1 Hz, 1H), 7.20 (d, J = 6.8 Hz, 1H), 6.90 (s, 2H), 6.70 (d, J = 8.3 Hz, 1H), 6.53 (s, 2H), 2.25 (s, 3H), 1.94 (s, 6H), 1.29 (q, J = 4.5 Hz, 2H), 0.97 (s, 2H).
622	523.20	1.91	
623	434.30	1.79	
624	538.20	1.92	
625	573.32	1.90	
626	494.50	1.85	
627	449.00	1.45	1H NMR (400 MHz, DMSO) δ 8.27 (s, 1H), 8.10 (d, J = 6.5 Hz, 1H), 7.67 (d, J = 7.8 Hz, 1H), 7.39 (d, J = 32.1 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			2H), 7.12 (d, J = 29.1 Hz, 3H), 6.97 (d, J = 6.4 Hz, 1H), 6.68 (s, 2H), 5.71 (s, 2H), 2.16 (s, 3H), 1.89 (s, 6H).
628	453.30	1.25	1H NMR (400 MHz, Methanol-d4) δ 7.78 (d, J = 2.7 Hz, 1H), 7.29 - 7.10 (m, 4H), 6.99 (s, 2H), 6.84 (s, 1H), 4.37 (s, 2H), 4.26 - 4.12 (m, 1H), 3.65 (s, 1H), 3.51 - 3.33 (m, 3H), 3.27 - 3.09 (m, 2H), 2.32 (s, 6H), 2.28 (s, 3H).
629	563.24	2.89	
630	508.00	2.49	1H NMR (400 MHz, DMSO) δ 12.41 (s, 11H), 7.97 (d, J = 8.3 Hz, 23H), 7.77 (d, J = 8.0 Hz, 11H), 7.63 (dd, J = 8.2, 7.5 Hz, 12H), 7.36 (d, J = 8.3 Hz, 23H), 7.18 (dd, J = 16.2, 7.6 Hz, 23H), 6.68 (d, J = 8.3 Hz, 11H), 6.48 (s, 22H), 2.92 (dd, J = 13.8, 6.9 Hz, 11H), 2.61 (d, J = 8.8 Hz, 23H), 2.30 (d, J = 22.9 Hz, 1H), 2.27 - 2.09 (m, 11H), 1.87 (dd, J = 11.8, 5.6 Hz, 11H), 1.61 (d, J = 3.5 Hz, 67H), 1.42 (t, J = 12.1 Hz, 13H), 1.23 (d, J = 6.9 Hz, 70H), 0.88 (d, J = 6.3 Hz, 36H).
631	512.30	1.85	1H NMR (400 MHz, DMSO-d6) δ 8.05 (s, 1H), 7.84 (s, 1H), 7.50 (d, J = 7.6 Hz, 1H), 7.44 (t, J = 7.8 Hz, 1H), 6.97 (d, J = 7.2 Hz, 1H), 6.69 (d, J = 7.6 Hz, 1H), 6.46 (d, J = 8.3 Hz, 1H), 6.15 (d, J = 4.6 Hz, 2H), 3.93 (d, J = 7.1 Hz, 2H), 2.95 (t, J = 10.6 Hz, 1H), 2.85 (dd, J = 10.8, 7.4 Hz, 1H), 2.18 - 2.02 (m, 2H), 1.78 (dd, J = 11.7, 5.7 Hz, 1H), 1.59 (s, 3H), 1.54 (s, 3H), 1.34 (t, J = 12.0 Hz, 1H), 0.85 (d, J = 6.7 Hz, 6H), 0.81 (d, J = 6.3 Hz, 3H).
632	399.40	1.63	1H NMR (400 MHz, DMSO-d6) δ 13.90 (s, 1H), 12.73 (s, 1H), 8.01 (d, J = 2.5 Hz, 1H), 7.82 (d, J = 8.1 Hz, 1H), 7.46 (d, J = 8.1 Hz, 1H), 7.42 - 7.34 (m, 2H), 7.11 (d, J = 7.8 Hz, 2H), 6.79 (d, J = 2.5 Hz, 1H), 2.34 (s, 3H), 1.35 (s, 9H).
633	561.20	0.81	
634	482.00	1.95	1H NMR (400 MHz, DMSO) δ 12.56 (s,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 11.91 (s, 1H), 8.24 (d, J = 9.4 Hz, 1H), 7.83 (s, 1H), 7.74 (s, 1H), 7.09 (s, 1H), 7.01 (s, 2H), 6.45 (s, 1H), 2.67 (s, 2H), 2.29 (s, 3H), 2.02 (s, 3H), 1.70 (s, 2H), 1.61 (s, 2H), 0.97 (s, 6H).
635	451.40	1.29	
636	425.40	1.29	
637	457.30	2.84	
638	528.12	2.71	
639	565.60	2.28	1H NMR (400 MHz, DMSO-d6) δ 12.40 (s, 1H), 8.15 (d, J = 7.8 Hz, 1H), 7.84 (d, J = 7.9 Hz, 1H), 7.66 (dd, J = 8.4, 7.3 Hz, 1H), 7.29 - 7.06 (m, 6H), 6.84 (dt, J = 10.8, 2.3 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 6.54 (s, 2H), 3.61 (d, J = 6.7 Hz, 2H), 2.06 (s, 6H), 1.98 (dt, J = 13.4, 6.7 Hz, 1H), 0.97 (d, J = 6.7 Hz, 6H).
640	531.18	1.97	
641	519.50	1.96	1H NMR (400 MHz, DMSO-d6) δ 12.19 (s, 1H), 8.11 - 8.01 (m, 1H), 7.83 - 7.73 (m, 2H), 7.66 (p, J = 8.3 Hz, 2H), 7.28 - 7.04 (m, 4H), 7.00 - 6.90 (m, 2H), 6.71 (d, J = 8.7 Hz, 1H), 6.51 (s, 2H), 4.05 (q, J = 7.0 Hz, 2H), 2.32 (s, 3H), 2.06 (s, 3H), 1.33 (d, J = 7.0 Hz, 3H).
642	530.23	1.88	
643	467.30	1.16	1H NMR (400 MHz, DMSO-d6) δ 12.63 (s, 1H), 9.00 - 8.86 (m, 2H), 8.42 (d, J = 5.8 Hz, 2H), 7.91 (d, J = 7.9 Hz, 1H), 7.65 (dd, J = 8.4, 7.3 Hz, 1H), 7.54 (dd, J = 8.1, 1.5 Hz, 1H), 7.22 (d, J = 7.3 Hz, 1H), 6.70 (d, J = 8.4 Hz, 1H), 6.51 (s, 1H), 2.71 - 2.53 (m, 2H), 2.23 (dt, J = 12.5, 5.7 Hz, 1H), 1.91 (dd, J = 11.9, 5.5 Hz, 1H), 1.63 (s, 3H), 1.59 (s, 3H), 1.45 (t, J = 12.1 Hz, 1H), 0.88 (d, J = 6.3 Hz, 3H).
644	558.07	2.09	
645	566.30	2.10	1H NMR (400 MHz, DMSO-d6) δ 8.10 (d, J = 7.7 Hz, 1H), 7.98 (dd, J = 7.3, 1.9 Hz, 1H), 7.96 (s, 1H), 7.62 (d, J = 7.7 Hz, 1H), 7.46 (s, 1H), 7.16 (d, J = 10.0 Hz, 1H), 7.10 (d, J = 2.1 Hz, 1H),

Cmpd No.	LC/MS M+1	LC/RT min	NMR
			6.92 (s, 2H), 6.70 (dd, J = 10.8, 2.4 Hz, 1H), 6.20 (s, 1H), 4.39 (p, J = 6.1 Hz, 1H), 2.90 (s, 1H), 2.27 (s, 3H), 2.01 (s, 6H), 1.22 (d, J = 6.0 Hz, 6H).
646	441.26	1.88	
647	573.32	1.90	
648	524.00	1.64	1H NMR (400 MHz, DMSO) δ 10.00 (s, 1H), 8.26 (s, 1H), 8.18 (s, 1H), 7.29 (s, 1H), 7.07 (d, J = 8.6 Hz, 2H), 6.92 (s, 3H), 4.28 - 3.92 (m, 7H), 3.81 (s, 3H), 2.40 (s, 6H), 2.24 (s, 3H), 2.07 (s, 1H), 1.74 (s, 3H), 1.24 (s, 3H).
649	433.25	2.10	
650	470.50	2.71	
651	604.38	2.31	
652	538.20	1.91	
653	421.26	1.73	
654	490.30	1.87	
655	568.40	2.98	
656	579.24	1.81	
657	556.30	2.67	1H NMR (400 MHz, DMSO-d6) δ 7.51 (d, J = 7.6 Hz, 1H), 7.47 - 7.42 (m, 2H), 7.37 - 7.33 (m, 1H), 7.08 (d, J = 7.7 Hz, 1H), 6.99 (d, J = 7.3 Hz, 1H), 6.79 (dt, J = 10.8, 2.3 Hz, 1H), 6.48 (d, J = 8.3 Hz, 1H), 6.11 (s, 2H), 3.82 (d, J = 6.6 Hz, 2H), 3.27 (d, J = 10.2 Hz, 1H), 2.95 - 2.84 (m, 1H), 2.09 - 1.99 (m, 1H), 1.79 (dq, J = 13.2, 6.6 Hz, 1H), 1.68 (t, J = 6.3 Hz, 1H), 1.59 (s, 3H), 1.45 - 1.38 (m, 1H), 1.33 (s, 3H), 0.99 (d, J = 6.7 Hz, 6H), 0.95 (d, J = 6.7 Hz, 3H).
658	587.30	1.91	
659	591.29	1.53	
660	453.23	1.17	
661	537.43	2.87	1H NMR (400 MHz, DMSO-d6) δ 8.16 - 8.13 (d, 1H), 7.65 - 7.46 (m, 2H), 7.22 - 7.10 (m, 3H), 7.04 - 6.91 (m, 3H), 6.69 - 6.66 (d, 1H), 6.51 (s, 2H), 3.85 (s, 3H), 2.25 (s, 3H), 2.02 (s, 6H)
662	480.30	1.25	
663	618.36	2.27	
664	541.33	1.78	
665	495.50	2.06	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
666	437.00	1.04	1H NMR (400 MHz, DMSO) δ 7.83 (s, 1H), 7.61 (s, 2H), 7.26 (dd, J = 41.7, 19.3 Hz, 7H), 6.96 (d, J = 8.7 Hz, 1H), 4.68 (d, J = 12.3 Hz, 1H), 4.46 (d, J = 12.6 Hz, 1H), 4.23 (d, J = 12.9 Hz, 2H), 2.43 (s, 3H), 1.81 (s, 3H).
667	484.30	2.13	
668	516.21	1.81	
669	549.40	1.73	1H NMR (400 MHz, DMSO-d6) δ 7.80 - 7.70 (m, 1H), 7.24 (d, J = 7.0 Hz, 1H), 7.08 (dt, J = 14.8, 6.6 Hz, 5H), 6.85 (s, 2H), 3.95 - 3.72 (m, 3H), 3.47 - 3.34 (m, 7H), 3.24 (dd, J = 14.0, 7.4 Hz, 3H), 3.16 - 3.01 (m, 1H), 2.99 - 2.66 (m, 3H), 2.22 (s, 3H), 2.16 (s, 6H).
670	401.30	1.67	
671	539.20	2.26	
672	503.21	3.11	
673	547.33	2.94	
674	463.30	1.59	
675	551.27	1.53	
676	547.20	2.10	1H NMR (400 MHz, DMSO-d6) δ 12.54 (s, 1H), 8.32 (d, J = 8.1 Hz, 1H), 7.85 (dd, J = 2.6, 1.1 Hz, 1H), 7.69 - 7.62 (m, 2H), 7.22 (dd, J = 7.4, 0.8 Hz, 1H), 6.99 (d, J = 2.6 Hz, 1H), 6.97 (s, 2H), 6.72 (d, J = 8.4 Hz, 1H), 6.56 (s, 2H), 2.28 (s, 3H), 2.03 (s, 6H).
677	579.30	1.65	
678	528.30	2.93	
679	511.20	2.09	
680	540.30	2.86	
681	516.30	1.71	1H NMR (400 MHz, DMSO-d6) δ 13.78 (s, 1H), 13.22 (s, 1H), 7.98 (s, 1H), 7.79 (d, J = 7.9 Hz, 1H), 7.53 (dd, J = 6.4, 3.3 Hz, 1H), 7.28 - 7.17 (m, 2H), 7.03 (dt, J = 9.0, 3.6 Hz, 1H), 6.87 (s, 1H), 4.00 (s, 2H), 3.78 (d, J = 6.6 Hz, 2H), 2.11 - 1.96 (m, 1H), 1.98 (s, 2H), 1.66 (s, 2H), 1.06 (d, J = 6.2 Hz, 6H), 0.98 (d, J = 6.7 Hz, 6H).
682	537.43	3.01	
683	528.30	2.64	
684	604.32	2.30	
685	436.40	1.72	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
686	586.35	2.21	
687	511.30	1.53	
688	557.40	1.85	
689	511.40	1.66	1H NMR (400 MHz, DMSO-d6) δ 12.41 (s, 1H), 8.86 (d, J = 2.5 Hz, 1H), 8.31 (dd, J = 8.7, 2.5 Hz, 1H), 7.83 (d, J = 8.0 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.31 - 7.16 (m, 2H), 6.84 (d, J = 8.7 Hz, 1H), 6.70 (t, J = 8.6 Hz, 1H), 6.48 (s, 2H), 5.31 (p, J = 6.1 Hz, 1H), 4.36 - 4.20 (m, 1H), 2.57 (m, obsucured multiplet), 2.29 - 2.14 (m, 1H), 1.31 (d, J = 6.2 Hz, 6H), 1.27 (d, J = 5.9 Hz, 2H), 1.14 (q, J = 11.5 Hz, 1H), 0.92 (d, J = 6.3 Hz, 2H).
690	479.00	1.25	1H NMR (400 MHz, DMSO) δ 10.18 (s, 1H), 7.68 (s, 1H), 7.30 - 7.09 (m, 4H), 6.90 (d, J = 30.5 Hz, 3H), 6.77 (d, J = 8.4 Hz, 1H), 4.40 (s, 1H), 4.15 (s, 3H), 3.96 (s, 2H), 2.96 (s, 2H), 2.23 (s, 3H), 2.15 (s, 6H).
691	547.20	0.77	
692	544.30	1.91	
693	508.26	1.94	
694	528.50	1.89	
695	538.26	2.00	
696	479.30	1.25	
697	588.15	3.05	
698	492.40	2.19	
699	529.39	1.85	1H NMR (400 MHz, DMSO) δ 8.33 (d, J = 2.2 Hz, 1H), 7.91 (d, J = 2.2 Hz, 1H), 7.86 (d, J = 7.9 Hz, 1H), 7.54 (d, J = 8.0 Hz, 1H), 7.48 - 7.40 (m, J = 8.7, 1.8 Hz, 2H), 6.92 (dt, J = 11.0, 2.3 Hz, 1H), 4.74 (hept, J = 6.0 Hz, 1H), 3.73 - 3.60 (m, J = 12.9 Hz, 2H), 2.98 - 2.82 (m, J = 11.7 Hz, 2H), 1.69 - 1.49 (m, J = 11.6 Hz, 3H), 1.30 (d, J = 6.0 Hz, 6H), 1.27 - 1.11 (m, J = 23.3, 11.6 Hz, 2H), 0.92 (d, J = 6.1 Hz, 3H).
700	537.43	2.95	
701	518.50	1.65	1H NMR (400 MHz, DMSO-d6) δ 13.78 (s, 1H), 13.21 (s, 1H), 7.98 (s, 1H), 7.87 (d, J = 2.0 Hz, 1H), 7.79 (d, J = 8.0 Hz, 1H), 7.64 (dd, J = 8.4, 1.9 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 7.51 (d, J = 8.3 Hz, 1H), 7.39 (d, J = 7.9 Hz, 1H), 6.86 (s, 1H), 4.77 (p, J = 6.1 Hz, 1H), 4.02 (s, 2H), 2.06 - 1.94 (m, 3H), 1.75 - 1.63 (m, 2H), 1.35 (d, J = 6.1 Hz, 6H), 1.07 (d, J = 6.2 Hz, 6H).
702	425.40	1.31	
703	523.23	1.51	
704	458.30	1.47	
705	534.50	1.99	1H NMR (400 MHz, DMSO-d6) δ 8.26 (s, 1H), 8.08 (d, J = 7.7 Hz, 1H), 7.89 (s, 1H), 7.51 (d, J = 7.7 Hz, 1H), 7.42 (t, J = 7.7 Hz, 1H), 7.08 (d, J = 7.3 Hz, 1H), 6.92 (s, 2H), 6.44 (d, J = 8.2 Hz, 1H), 6.05 (s, 2H), 3.87 (s, 3H), 2.90 (s, 1H), 2.28 (s, 3H), 2.12 (s, 3H), 2.02 (s, 6H).
706	545.21	2.10	
707	452.00	1.48	1H NMR (400 MHz, DMSO) δ 12.94 (s, 1H), 8.40 (s, 1H), 8.04 (s, 1H), 7.59 (s, 1H), 7.31 (d, J = 8.1 Hz, 2H), 7.06 (d, J = 14.8 Hz, 4H), 2.81 (s, 2H), 2.31 (s, 3H), 1.77 (s, 4H), 1.29 (s, 6H).
708	525.50	1.60	
709	557.20	2.18	1H NMR (400 MHz, DMSO-d6) δ 12.41 (s, 1H), 8.18 (d, J = 7.8 Hz, 1H), 8.12 (d, J = 7.8 Hz, 1H), 8.04 (s, 1H), 7.89 (d, J = 7.8 Hz, 1H), 7.79 - 7.72 (m, 1H), 7.70 - 7.61 (m, 2H), 7.24 (d, J = 7.0 Hz, 1H), 6.95 (s, 2H), 6.72 (d, J = 8.4 Hz, 1H), 6.54 (s, 2H), 2.27 (s, 3H), 2.02 (s, 6H).
710	498.40	1.67	
711	618.36	2.11	
712	484.20	1.47	
713	522.50	1.83	
714	497.20	2.75	1H NMR (400 MHz, Chloroform-d) δ 10.71 - 10.46 (m, 1H), 8.04 (s, 1H), 7.64 (d, J = 3.6 Hz, 2H), 6.90 (s, 2H), 6.65 (s, 1H), 4.70 (s, 2H), 3.60 (s, 2H), 2.33 (s, 3H), 2.08 (s, 6H), 0.99 (s, 6H).
715	468.00	2.20	1H NMR (400 MHz, DMSO) δ 13.78 (s, 1H), 12.16 (s, 1H), 7.97 (s, 1H), 7.69 (s, 1H), 6.90 (s, 2H), 6.86 (s, 1H), 2.67 (s, 2H), 2.25 (s, 3H), 1.95 (s, 6H), 1.68 (s, 2H), 1.60 (s, 2H), 0.92 (s, 6H).

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
716	564.29	2.14	1H NMR (400 MHz, DMSO) δ 13.81 (s, 1H), 12.70 (s, 1H), 8.01 (s, 1H), 7.74 (d, J = 7.9 Hz, 1H), 7.29 - 7.19 (m, 5H), 7.17 - 7.06 (m, 3H), 6.93 (s, 1H), 6.77 (dt, J = 10.7, 2.3 Hz, 1H), 5.31 (t, J = 7.6 Hz, 1H), 3.72 (ddd, J = 31.5, 9.2, 6.4 Hz, 2H), 3.17 (dd, J = 16.6, 9.7 Hz, 1H), 2.81 (t, J = 7.5 Hz, 1H), 2.38 (dd, J = 10.9, 5.0 Hz, 1H), 2.07 - 1.98 (m, 1H), 1.93 - 1.72 (m, 2H), 1.64 (dd, J = 18.4, 8.3 Hz, 1H), 1.01 (dd, J = 6.7, 2.1 Hz, 6H).
717	539.60	1.71	
718	534.20	2.01	
719	558.30	2.00	1H NMR (400 MHz, DMSO-d6) δ 12.52 (s, 1H), 7.94 (d, J = 7.9 Hz, 1H), 7.78 (d, J = 7.9 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.51 (dt, J = 7.0, 1.8 Hz, 2H), 7.48 - 7.42 (m, 2H), 7.35 (dd, J = 7.9, 5.5 Hz, 2H), 7.21 (d, J = 7.2 Hz, 1H), 6.89 (dt, J = 10.6, 2.3 Hz, 2H), 6.71 (d, J = 8.4 Hz, 1H), 4.59 (dd, J = 7.2, 3.3 Hz, 1H), 4.55 - 4.46 (m, 1H), 3.85 (d, J = 3.2 Hz, 2H), 3.83 (d, J = 3.1 Hz, 2H), 3.81 (dd, J = 9.1, 3.3 Hz, 1H), 3.74 (dd, J = 9.1, 3.3 Hz, 1H), 3.59 - 3.52 (m, 2H), 3.35 (dd, J = 9.1, 7.5 Hz, 2H), 3.32 (s, 3H), 3.27 (s, 3H), 2.94 (dd, J = 9.7, 6.7 Hz, 1H), 2.89 - 2.78 (m, 1H), 2.14 - 1.94 (m, 5H), 1.86 - 1.68 (m, 5H), 1.00 (d, J = 2.3 Hz, 6H), 0.99 (d, J = 2.2 Hz, 6H). Two rotamers in about 1:1 ratio. Thus two sets of peaks.
720	542.34	2.02	
721	543.30	1.66	
722	521.00	1.85	1H NMR (400 MHz, DMSO) δ 12.95 (s, 1H), 8.04 (d, J = 8.3 Hz, 2H), 7.64 (s, 1H), 7.56 (s, 2H), 7.47 (d, J = 7.9 Hz, 2H), 7.17 (d, J = 8.2 Hz, 3H), 6.96 (d, J = 10.8 Hz, 1H), 6.81 (d, J = 8.4 Hz, 1H), 4.76 (s, 1H), 2.33 (s, 3H), 1.32 (s, 6H).
723	555.10	1.25	
724	525.25	1.76	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
725	525.42	3.01	
726	483.20	1.17	1H NMR (400 MHz, DMSO-d6) δ 13.13 (s, 1H), 8.90 (d, J = 2.5 Hz, 1H), 8.36 (dd, J = 8.7, 2.5 Hz, 1H), 7.83 (d, J = 7.9 Hz, 1H), 7.65 (dd, J = 8.4, 7.3 Hz, 1H), 7.36 (d, J = 8.0 Hz, 1H), 7.22 (d, J = 7.1 Hz, 1H), 6.94 (d, J = 8.7 Hz, 1H), 6.76 - 6.69 (m, 1H), 4.06 - 4.00 (m, 2H), 3.92 (s, 3H), 2.05 - 1.93 (m, 2H), 1.73 - 1.60 (m, 2H), 1.09 (d, J = 6.3 Hz, 6H).
727	537.43	2.51	
728	520.21	1.83	
729	539.30	1.68	
730	504.30	1.65	
731	573.25	1.70	
732	523.23	1.69	
733	537.30	1.75	
734	516.21	1.82	
735	542.31	2.00	
736	511.30	1.56	
737	529.32	1.77	1H NMR (400 MHz, DMSO) δ 8.25 (dd, J = 7.3, 2.1 Hz, 1H), 7.91 - 7.80 (m, J = 7.9 Hz, 2H), 7.51 (d, J = 8.0 Hz, 1H), 7.44 (dd, J = 9.9, 1.7 Hz, 2H), 6.91 (dt, J = 10.9, 2.2 Hz, 1H), 6.46 (dd, J = 7.1, 6.5 Hz, 1H), 4.73 (hept, J = 6.0 Hz, 1H), 3.78 - 3.66 (m, J = 12.8 Hz, 2H), 2.95 - 2.80 (m, J = 11.6 Hz, 2H), 1.69 - 1.49 (m, J = 20.3, 9.3 Hz, 3H), 1.30 (d, J = 6.0 Hz, 6H), 1.27 - 1.12 (m, J = 12.0 Hz, 2H), 0.92 (d, J = 6.3 Hz, 3H).
738	525.20	2.16	
739	536.16	1.92	
740	497.21	1.37	
741	547.60	1.97	
742	573.32	1.90	1H NMR (400 MHz, DMSO) δ 8.92 (d, J = 2.4 Hz, 1H), 8.38 (dd, J = 8.7, 2.4 Hz, 1H), 7.89 (d, J = 7.9 Hz, 1H), 7.64 (t, J = 7.8 Hz, 1H), 7.49 (d, J = 7.9 Hz, 1H), 7.26 (tt, J = 12.8, 6.5 Hz, 6H), 6.87 (d, J = 8.7 Hz, 1H), 6.69 (d, J = 8.4 Hz, 1H), 5.33 (dd, J = 12.3, 6.2 Hz, 1H), 3.87 (d, J = 12.4 Hz, 2H), 3.02 (t, J

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			= 11.9 Hz, 2H), 2.75 (s, 1H), 2.59 - 2.46 (m, 3H), 1.93 - 1.63 (m, 4H), 1.32 (d, J = 6.2 Hz, 6H).
743	510.30	2.20	
744	518.12	2.21	
745	515.25	1.65	
746	482.50	2.86	10.67 (s, 1H), 7.92 - 7.89 (d, 1H), 7.63 (t, J = 7.80, 1H), 7.23 - 7.20 (d, 1H), 6.94 (s, 2H), 6.70 - 6.67 (d, 1H), 6.49 (s, 2H), 6.22 - 6.19 (d, 1H), 2.27 (s, 3H), 2.06 (s, 6H), 1.81 (s, 4H)
747	493.20	1.72	
748	497.51	2.65	1H NMR (400 MHz, DMSO) δ 7.95 - 7.92 (d, 1H), 7.46 (t, J = 7.03, 1H), 7.09 - 7.06 (d, 1H), 6.89 - 6.87 (m, 3H), 6.52 - 6.49 (d, 1H), 6.20 (s, 2H), 3.80 - 3.76 (m, 2H), 2.67- 2.59 (m, 1H), 2.25 (s, 3H), 1.96 (s, 6H) , 1.56 - 1.34 (m, 4H)
749	532.20	1.80	
750	458.38	1.59	
751	542.40	1.95	1H NMR (400 MHz, DMSO-d6) δ 12.58 (s, 1H), 7.76 (d, J = 7.9 Hz, 1H), 7.65 (t, J = 7.8 Hz, 1H), 7.48 - 7.45 (m, 1H), 7.42 (dt, J = 10.2, 1.8 Hz, 1H), 7.32 (d, J = 7.9 Hz, 1H), 7.23 (d, J = 7.3 Hz, 1H), 6.89 (dt, J = 10.8, 2.3 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 4.13 (s, 2H), 3.84 (d, J = 6.6 Hz, 2H), 2.12 - 1.98 (m, 3H), 1.49 (dd, J = 9.8, 7.4 Hz, 2H), 0.99 (dd, J = 6.7, 3.9 Hz, 12H).
752	461.40	1.73	
753	417.30	1.75	
754	550.20	2.11	
755	466.50	1.79	
756	516.22	1.82	
757	511.30	1.66	
758	525.20	2.17	
759	471.50	3.24	1H NMR (400 MHz, DMSO-d6) δ 12.77 (s, 1H), 8.50 (dd, J = 4.9, 1.9 Hz, 1H), 8.33 (dd, J = 7.7, 1.9 Hz, 1H), 7.27 (dd, J = 7.7, 4.9 Hz, 1H), 7.06 (s, 1H), 6.74 (s, 2H), 5.40 (s, 2H), 3.96 (s, 3H), 2.17 (s, 3H), 2.05 (s, 6H), 1.16 (s, 9H).

Cmpd No.	LC/MS M+1	LC/RT min	NMR
760	487.11	1.32	
761	480.30	1.47	
762	538.20	1.91	
763	498.30	1.45	
764	511.40	1.64	1H NMR (400 MHz, Ethanol-d6) δ 8.85 (d, J = 2.4 Hz, 1H), 8.31 (dd, J = 8.7, 2.5 Hz, 1H), 7.78 (d, J = 8.0 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.22 (dd, J = 13.7, 7.5 Hz, 2H), 6.85 (d, J = 8.7 Hz, 1H), 6.71 (d, J = 8.3 Hz, 1H), 5.31 (p, J = 6.2 Hz, 2H), 3.70 (dd, J = 8.1, 5.9 Hz, 1H), 2.93 - 2.77 (m, 1H), 2.72 - 2.61 (m, 1H), 1.96 - 1.67 (m, 2H), 1.32 (s, 6H) 1.32 (obsucured m, 1H), 1.26 (d, J = 5.9 Hz, 3H), 1.05 (d, J = 6.3 Hz, 3H).
765	480.00	1.51	1H NMR (400 MHz, DMSO) δ 9.89 (s, 1H), 8.20 (d, J = 24.9 Hz, 2H), 7.55 (s, 1H), 7.29 (d, J = 41.2 Hz, 3H), 7.08 (s, 1H), 6.91 (s, 2H), 4.37 - 4.01 (m, 4H), 3.82 (s, 3H), 2.41 (s, 6H), 2.24 (s, 3H), 1.78 (s, 3H).
766	533.31	1.88	
767	480.30	1.86	
768	535.32	2.59	
769	551.13	2.13	
770	512.20	1.25	
771	589.00	2.38	1H NMR (400 MHz, DMSO) δ 8.49 (d, J = 7.9 Hz, 1H), 8.44 (s, 1H), 8.36 (d, J = 7.5 Hz, 1H), 8.13 (d, J = 7.8 Hz, 1H), 7.84 (d, J = 7.9 Hz, 1H), 7.22 (d, J = 11.5 Hz, 1H), 7.18 (s, 1H), 6.94 (s, 2H), 6.84 (d, J = 10.7 Hz, 1H), 3.65 (d, J = 6.5 Hz, 2H), 2.26 (s, 3H), 2.07 (s, 3H), 1.99 (s, 7H), 0.97 (d, J = 6.7 Hz, 6H).
772	604.50	2.25	
773	544.30	1.70	1H NMR (400 MHz, DMSO-d6) δ 12.98 (s, 1H), 7.83 (d, J = 1.9 Hz, 1H), 7.81 (d, J = 8.0 Hz, 1H), 7.70 - 7.61 (m, 2H), 7.52 (d, J = 8.3 Hz, 1H), 7.37 (d, J = 7.9 Hz, 1H), 7.23 (dd, J = 7.3, 0.8 Hz, 1H), 6.72 (d, 1H), 6.48 (s, 2H), 4.13 (t, J = 6.5 Hz, 2H), 4.09 - 3.97 (m, 2H), 2.05 - 1.91 (m, 2H), 1.81 (h, J = 7.2 Hz, 2H), 1.74 - 1.61 (m, 2H), 1.10

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(d, J = 6.2 Hz, 6H), 1.03 (t, J = 7.4 Hz, 3H).
774	546.21	1.75	
775	557.24	2.72	1H NMR (400 MHz, DMSO-d6) δ 7.95 (t, J = 7.80, 1H), 8.86 - 8.83 (d, 1H), 7.62 - 7.59 (d, 1H), 7.52 - 7.48 (m, 3H), 6.99 - 6.90 (m, 2H), 3.86 - 3.83 (d, 2H), 3.26 (m, 4H), 2.10 - 1.97 (m, 1H), 1.33 (m, 4H), 1.01 - 0.98 (d, 6H), 0.94 (s, 6H)
776	547.60	1.97	1H NMR (400 MHz, DMSO) δ 13.74 (s, 1H), 12.85 (s, 1H), 8.86 (d, J = 2.5 Hz, 1H), 8.32 (dd, J = 8.7, 2.5 Hz, 1H), 7.94 (s, 1H), 7.80 (d, J = 7.9 Hz, 1H), 7.45 (d, J = 7.9 Hz, 1H), 7.38 - 7.30 (m, 4H), 7.27 - 7.20 (m, 1H), 6.84 (d, J = 8.8 Hz, 2H), 5.37 - 5.23 (m, 1H), 3.93 (d, J = 11.4 Hz, 1H), 3.59 (d, J = 12.7 Hz, 1H), 3.02 - 2.83 (m, 3H), 1.91 (d, J = 8.9 Hz, 1H), 1.70 (t, J = 11.4 Hz, 3H), 1.31 (d, J = 6.2 Hz, 6H).
777	524.30	1.89	
778	551.40	2.22	
779	554.60	1.79	
780	610.38	2.24	
781	484.30	1.20	
782	579.27	2.29	
783	531.30	1.98	
784	463.30	1.96	1H NMR (400 MHz, DMSO-d6) δ 13.80 (s, 1H), 12.46 (s, 1H), 8.09 (d, J = 7.8 Hz, 1H), 7.98 (s, 1H), 7.82 - 7.71 (m, 3H), 7.41 (dp, J = 3.5, 1.9 Hz, 3H), 6.95 (s, 2H), 6.88 (s, 1H), 2.28 (s, 3H), 2.03 (s, 6H).
785	496.60	2.96	
786	519.42	2.53	1H NMR (400 MHz, DMSO-d6) δ 12.35 (s, 1H), 8.17 - 8.14 (d, 1H), 7.76 - 7.56 (m, 4H), 7.37 - 7.35 (m, 2H), 7.26 - 7.23 (d, 1H), 6.95 (s, 2H), 6.74 - 7.61 (d, 1H), 6.56 (s, 2H), 4.48 (s, 2H), 2.28 (s, 3H), 2.04 (s, 6H)
787	533.22	2.05	
788	544.30	1.65	
789	523.30	1.96	1H NMR (400 MHz, DMSO-d6) δ 8.78 (d, J = 2.4 Hz, 1H), 8.28 (dd, J = 8.7,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			2.5 Hz, 1H), 7.60 - 7.42 (m, 2H), 7.01 (bs, 2H), 6.87 (d, J = 8.7 Hz, 1H), 6.51 (d, J = 8.6 Hz, 1H), 6.23 (bs, 2H), 4.23 (s, 2H), 4.13 (d, J = 7.1 Hz, 2H), 1.86 (bs, 2H), 1.56 (bs, 2H), 1.25 (dddd, J = 11.9, 7.3, 5.4, 2.6 Hz, 1H), 1.04 (d, J = 6.1 Hz, 6H), 0.59 - 0.49 (m, 2H), 0.36 - 0.26 (m, 2H).
790	451.00	1.10	1H NMR (400 MHz, DMSO) δ 10.04 (s, 1H), 7.74 (s, 1H), 7.49 (s, 1H), 7.42 - 7.07 (m, 7H), 6.82 (s, 1H), 4.36 - 4.09 (m, 5H), 2.44 (s, 5H), 1.82 (s, 3H).
791	485.50	2.71	
792	470.32	1.38	
793	519.20	1.87	
794	432.40	1.50	
795	571.34	3.04	1H NMR (400 MHz, DMSO) δ 12.89 (s, 1H), 8.04 (t, J = 8.04, 1H), 7.83 - 7.75 (m, 2H), 7.49 (s, 1H), 7.44 - 7.40 (d, 1H), 7.32 - 7.29 (d, 1H), 7.18 - 7.15 (d, 1H), 6.93 - 6.89 (d, 1H), 3.91 (s, 3H), 3.85 - 3.82 (d, 2 H), 2.10 (m, 1H), 2.07 - 1.99 (m, 1H), 1.88 - 1.83 (m, 1H), 1.59 (s, 6H), 1.39 (t, J = 12.41, 1H), 1.00 - 0.97 (d, 6H), 0.78 - 0.75 (d, 3H)
796	538.60	0.97	
797	481.00	2.01	1H NMR (400 MHz, DMSO) δ 7.60 (s, 1H), 7.29 (d, J = 8.4 Hz, 1H), 7.21 (d, J = 14.5 Hz, 2H), 7.13 (s, 1H), 6.99 (d, J = 7.1 Hz, 1H), 6.81 (d, J = 10.1 Hz, 1H), 6.43 (d, J = 8.4 Hz, 1H), 5.59 (s, 2H), 1.25 (s, 9H).
798	536.00	201.00	1H NMR (400 MHz, DMSO) δ 13.12 (s, 1H), 8.56 (d, J = 6.0 Hz, 1H), 8.25 (d, J = 7.6 Hz, 1H), 8.07 (d, J = 8.1 Hz, 1H), 7.98 (d, J = 8.1 Hz, 1H), 7.54 (d, J = 10.1 Hz, 2H), 7.37 (d, J = 8.0 Hz, 2H), 7.24 (s, 1H), 7.04 (d, J = 8.0 Hz, 2H), 6.95 (d, J = 13.1 Hz, 1H), 4.75 (s, 1H), 4.03 (s, 3H), 2.32 (s, 3H), 1.30 (d, J = 6.0 Hz, 6H).
799	427.30	1.90	
800	432.36	1.60	
801	518.22	2.05	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
802	466.33	1.61	
803	525.42	2.97	
804	580.20	2.19	
805	452.30	1.24	
806	443.18	2.74	
807	542.30	1.70	
808	521.20	1.81	
809	425.30	1.72	
810	588.15	2.07	
811	525.30	1.57	1H NMR (400 MHz, DMSO-d6) δ 12.42 (s, 1H), 8.82 (d, J = 2.4 Hz, 1H), 8.27 (dd, J = 8.8, 2.6 Hz, 1H), 7.72 (d, J = 7.8 Hz, 1H), 7.58 (s, 1H), 7.16 (d, J = 8.6 Hz, 2H), 6.85 (d, J = 8.7 Hz, 1H), 6.63 (s, 1H), 6.41 (s, 2H), 5.32 (h, J = 6.3 Hz, 1H), 2.83 - 2.56 (m, 2H), 2.27 - 2.13 (m, 1H), 1.86 (dd, J = 11.9, 5.7 Hz, 1H), 1.59 (d, J = 4.8 Hz, 6H), 1.41 (t, J = 12.0 Hz, 1H), 1.31 (d, J = 6.2 Hz, 6H), 0.88 (d, J = 6.3 Hz, 3H).
812	561.50	1.75	
813	493.20	1.96	1H NMR (400 MHz, DMSO-d6) δ 13.78 (s, 1H), 12.33 (s, 1H), 8.01 (d, J = 7.9 Hz, 1H), 7.98 (d, J = 2.4 Hz, 1H), 7.79 - 7.73 (m, 2H), 7.66 (d, J = 7.9 Hz, 1H), 7.13 (dd, J = 5.0, 3.0 Hz, 2H), 7.10 - 7.04 (m, 1H), 6.97 - 6.91 (m, 2H), 6.86 (s, 1H), 4.05 (q, J = 6.9 Hz, 2H), 2.32 (s, 3H), 2.05 (s, 3H), 1.31 (t, J = 7.0 Hz, 3H).
814	539.50	1.44	
815	526.10	1.98	
816	478.50	1.78	1H NMR (400 MHz, DMSO) δ 8.25 (d, J = 2.1 Hz, 1H), 7.95 (d, J = 2.1 Hz, 1H), 7.61 (dd, J = 8.3, 7.4 Hz, 1H), 7.32 (s, 1H), 7.08 (d, J = 6.9 Hz, 1H), 6.70 (d, J = 8.2 Hz, 1H), 6.60 (s, 2H), 5.63 (s, 2H), 4.81 (br s, 2H), 2.68 (q, J = 7.5 Hz, 2H), 2.12 (s, 3H), 1.91 (s, 6H), 1.21 (t, J = 7.6 Hz, 3H).
817	503.00	1.94	1H NMR (400 MHz, DMSO) δ 12.47 (s, 1H), 8.15 (s, 1H), 7.95 (s, 1H), 7.63 (s, 1H), 7.31 (d, J = 11.3 Hz, 4H), 7.22 (d, J = 7.3 Hz, 1H), 6.91 (s, 2H), 6.69 (d, J = 8.4 Hz, 1H), 6.50 (s, 2H), 2.24 (d, J =

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			4.4 Hz, 6H), 2.00 (s, 6H).
818	601.31	1.90	
819	490.00	1.68	1H NMR (400 MHz, DMSO) δ 13.14 (s, 1H), 8.42 (s, 1H), 8.14 (d, J = 8.7 Hz, 2H), 8.06 (s, 1H), 7.96 (s, 2H), 7.41 (d, J = 7.9 Hz, 2H), 7.09 (dd, J = 21.2, 8.3 Hz, 4H), 4.10 (s, 2H), 2.34 (s, 3H), 1.36 (s, 3H).
820	460.00	1.68	1H NMR (400 MHz, DMSO) δ 13.12 (s, 1H), 8.42 (d, J = 2.2 Hz, 1H), 8.07 (dd, J = 12.6, 5.2 Hz, 3H), 7.99 (s, 2H), 7.42 (d, J = 8.1 Hz, 2H), 7.34 (d, J = 8.1 Hz, 2H), 7.10 (t, J = 15.8 Hz, 4H), 2.36 (d, J = 14.8 Hz, 6H).
821	456.10	2.19	1H NMR (400 MHz, DMSO) δ 8.01 - 7.99 (d, 1H), 7.46 (s, 2H), 7.06 - 6.77 (m, 6H), 6.41 (s, 1H), 5.67 (s, 2H), 2.80 (s, 3H), 2.23 (s, 6H), 1.12 (s, 9H)
822	498.50	1.46	
823	468.40	2.39	1H NMR (400 MHz, DMSO-d6) δ 12.47 (s, 1H), 7.85 - 7.80 (m, 2H), 7.63 - 7.54 (m, 2H), 6.99 (s, 1H), 6.74 (s, 2H), 5.43 (s, 2H), 2.75 (q, J = 7.6 Hz, 2H), 2.16 (s, 3H), 2.05 (s, 6H), 1.23 (t, J = 7.6 Hz, 3H), 1.14 (s, 9H).
824	545.28	2.00	
825	480.00	1.33	
826	466.00	2.04	1H NMR (400 MHz, DMSO) δ 11.95 (s, 1H), 7.92 (d, J = 7.9 Hz, 1H), 7.23 (s, 2H), 7.10 (s, 2H), 6.94 - 6.81 (m, 3H), 3.51 (s, 2H), 2.25 (s, 3H), 1.92 (s, 6H), 1.31 (s, 3H), 0.67 (d, J = 14.3 Hz, 4H).
827	566.30	1.11	
828	549.43	2.97	
829	463.17	1.86	
830	537.43	2.98	
831	518.20	2.12	
832	518.20	1.71	
833	510.30	1.41	
834	511.20	1.48	1H NMR (400 MHz, DMSO) δ 12.44 (s, 1H), 8.84 (d, J = 2.4 Hz, 1H), 8.28 (dd, J = 8.8, 2.5 Hz, 1H), 7.72 (d, J = 8.0 Hz, 1H), 7.69 - 7.60 (m, 1H), 7.18 (t, J = 7.2 Hz, 2H), 6.86 (d, J = 8.7 Hz, 1H), 6.70 (d, J = 8.3 Hz, 1H), 5.32 (dt, J =

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			12.1, 6.1 Hz, 1H), 2.76 (s, 2H), 1.76 (s, 4H), 1.56 (d, J = 14.3 Hz, 6H), 1.32 (d, J = 6.2 Hz, 6H).
835	552.20	2.04	1H NMR (400 MHz, DMSO-d6) δ 11.83 (s, 1H), 8.10 (d, J = 7.7 Hz, 1H), 8.01 (dd, J = 7.0, 2.0 Hz, 1H), 7.63 (d, J = 7.7 Hz, 1H), 7.51 (s, 1H), 7.20 - 7.08 (m, 2H), 6.92 (s, 2H), 6.75 (dt, J = 10.7, 2.3 Hz, 1H), 6.27 (s, 1H), 3.91 (q, J = 7.0 Hz, 2H), 2.26 (s, 3H), 2.00 (s, 6H), 1.29 (t, J = 7.0 Hz, 3H).
836	511.30	1.66	
837	556.50	1.90	
838	523.40	1.50	
839	538.20	1.89	
840	492.00	1.29	1H NMR (400 MHz, DMSO) δ 11.11 (s, 1H), 7.64 (s, 1H), 7.31 (s, 4H), 7.19 (s, 1H), 7.10 (s, 1H), 6.72 (s, 2H), 4.58 (s, 1H), 4.15 (d, J = 7.4 Hz, 1H), 3.90 (d, J = 12.5 Hz, 1H), 2.22 (s, 3H), 2.19 (s, 3H), 2.12 (s, 6H), 1.60 (s, 3H), 1.48 (s, 3H).
841	441.40	1.82	
842	562.34	3.06	1H NMR (400 MHz, DMSO) δ 12.44 (s, 1H), 9.10 (s, 1H), 7.72 - 7.69 (d, 1H), 7.49 (s, 1H), 7.44 - 7.40 (d, 1H), 7.30 - 7.27 (d, 1H), 6.92 - 6.88 (d, 1H), 3.85 - 3.82 (d, 2H), 2.67 - 2.59 (m, 2H), 2.20 (m, 1H), 2.10 - 2.00 (m, 1H), 1.92 - 1.85 (m, 1H), 1.64 (s, 3H), 1.59 (s, 3H), 1.45 (t, J = 12.09, 1H), 1.00 - 0.97 (d, 6H), 0.84 - 0.81 (d, 3H)
843	546.20	1.86	
844	526.30	2.00	1H NMR (400 MHz, DMSO-d6) δ 9.13 (s, 2H), 7.65 (s, 1H), 7.53 (s, 1H), 7.09 (d, J = 31.1 Hz, 2H), 6.57 (s, 1H), 6.34 (s, 2H), 5.26 (h, J = 6.2 Hz, 1H), 2.80 (s, 1H), 2.13 (s, 1H), 1.84 (dd, J = 11.7, 5.6 Hz, 1H), 1.59 (s, 3H), 1.55 (s, 3H), 1.34 (d, J = 6.2 Hz, 6H), 0.82 (d, J = 6.2 Hz, 3H).
845	554.50	1.94	
846	553.60	1.81	
847	494.30	1.81	
848	553.50	1.79	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
849	495.00	1.81	1H NMR (400 MHz, DMSO) δ 13.93 (s, 1H), 12.89 (s, 1H), 8.02 (d, J = 24.9 Hz, 3H), 7.57 (s, 2H), 7.45 (d, J = 8.0 Hz, 2H), 7.15 (d, J = 7.9 Hz, 2H), 6.96 (d, J = 12.7 Hz, 1H), 6.81 (s, 1H), 4.76 (s, 1H), 2.36 (s, 3H), 1.32 (s, 6H).
850	570.30	2.05	1H NMR (400 MHz, DMSO) δ 12.52 (s, 1H), 7.75 (d, J = 8.0 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.46 (s, 1H), 7.38 (d, J = 9.4 Hz, 1H), 7.27 (d, J = 8.0 Hz, 1H), 7.21 (d, J = 7.3 Hz, 1H), 6.89 (dt, J = 10.7, 2.3 Hz, 1H), 6.70 (d, J = 8.4 Hz, 1H), 6.51 (s, 2H), 3.83 (d, J = 6.6 Hz, 2H), 2.59 (s, 2H), 2.05 (dd, J = 11.6, 5.0 Hz, 1H), 1.76 (s, 2H), 1.65 (s, 6H), 1.03 - 0.95 (m, 12H).
851	553.60	1.64	
852	424.19	2.24	1H NMR (400 MHz, DMSO-d6) δ 12.47 (s, 1H), 7.70 (d, J = 7.9 Hz, 1H), 7.62 (dd, J = 8.4, 7.3 Hz, 1H), 7.18 (d, J = 7.3 Hz, 1H), 6.68 (d, J = 8.4 Hz, 1H), 6.61 (d, J = 7.9 Hz, 1H), 6.51 (s, 2H), 2.60 - 2.52 (m, 2H), 2.20 (dq, J = 16.9, 6.4 Hz, 1H), 1.85 (dd, J = 12.0, 5.6 Hz, 1H), 1.48 (d, J = 9.8 Hz, 6H), 1.39 (t, J = 12.2 Hz, 1H), 0.86 (d, J = 6.2 Hz, 3H).
853	531.30	1.96	1H NMR (400 MHz, DMSO) δ 12.72 (s, 1H), 8.29 (d, J = 7.6 Hz, 1H), 7.68 - 7.59 (m, 2H), 7.20 (d, J = 7.2 Hz, 1H), 6.90 (s, 2H), 6.71 (d, J = 8.4 Hz, 1H), 6.59 (s, 2H), 2.25 (s, 3H), 1.94 (s, 6H).
854	467.00	1.92	1H NMR (400 MHz, DMSO) δ 12.31 (s, 1H), 8.03 (s, 1H), 7.78 (s, 1H), 7.63 (s, 1H), 7.22 (d, J = 7.3 Hz, 1H), 6.87 (s, 2H), 6.71 (d, J = 8.3 Hz, 1H), 6.50 (s, 2H), 2.24 (s, 3H), 1.94 (s, 6H), 1.37 (s, 3H), 0.89 (s, 2H), 0.75 (s, 2H).
855	480.00	1.32	1H NMR (400 MHz, DMSO) δ 12.60 (s, 1H), 8.20 (d, J = 9.3 Hz, 1H), 7.82 (s, 1H), 7.12 (d, J = 28.2 Hz, 3H), 6.93 (s, 1H), 6.82 (s, 2H), 6.43 (s, 1H), 4.41 (s, 1H), 2.83 (s, 2H), 2.21 (s, 3H), 2.14 (s, 6H), 2.07 (s, 2H).
856	529.02	2.55	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
857	554.50	2.13	
858	567.27	1.72	
859	590.30	2.21	
860	563.30	1.99	
861	505.00	1.80	1H NMR (400 MHz, DMSO) δ 13.12 (s, 1H), 8.81 (s, 1H), 8.56 (d, J = 6.8 Hz, 1H), 8.31 (s, 1H), 8.25 (d, J = 7.6 Hz, 1H), 7.97 (s, 2H), 7.37 (d, J = 8.1 Hz, 2H), 7.25 (s, 1H), 7.04 (d, J = 7.9 Hz, 2H), 4.04 (s, 3H), 3.96 (s, 3H), 2.28 (d, J = 30.8 Hz, 6H).
862	483.40	3.22	
863	497.31	1.53	
864	573.28	1.92	
865	531.00	2.10	1H NMR (400 MHz, DMSO) δ 8.07 (s, 1H), 7.88 (s, 1H), 7.62 (s, 1H), 7.42 (d, J = 17.4 Hz, 2H), 7.21 (s, 3H), 6.91 (s, 2H), 6.68 (s, 1H), 6.47 (s, 2H), 2.90 (s, 1H), 2.25 (s, 3H), 2.00 (s, 6H), 1.12 (d, J = 6.8 Hz, 6H).
866	523.20	1.83	
867	583.22	2.27	
868	545.00	2.13	1H NMR (400 MHz, DMSO) δ 12.52 (s, 1H), 8.02 (s, 1H), 7.83 (s, 1H), 7.71 - 7.55 (m, 2H), 7.37 (s, 1H), 7.23 (d, J = 6.7 Hz, 2H), 7.00 (d, J = 6.1 Hz, 1H), 6.90 (s, 2H), 6.69 (d, J = 11.3 Hz, 1H), 6.48 (s, 1H), 2.24 (s, 3H), 2.00 (s, 6H), 1.12 (s, 9H).
869	517.50	2.19	
870	479.32	1.22	
871	593.00	1.81	1H NMR (400 MHz, DMSO) δ 12.76 (s, 1H), 8.49 (s, 2H), 8.25 - 8.09 (m, 3H), 7.76 (s, 2H), 7.25 - 7.13 (m, 2H), 6.94 (s, 2H), 6.82 (d, J = 10.4 Hz, 1H), 4.25 (s, 2H), 3.64 (d, J = 6.5 Hz, 2H), 2.27 (s, 3H), 2.02 (s, 7H), 0.97 (d, J = 6.7 Hz, 6H).
872	517.18	1.89	
873	449.50	1.57	
874	565.20	2.24	
875	521.20	2.14	
876	494.30	1.53	
877	524.10	2.66	
878	520.21	1.84	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
879	470.50	1.46	1H NMR (400 MHz, DMSO-d6) δ 13.72 (s, 1H), 8.24 (d, J = 8.2 Hz, 1H), 8.03 (d, J = 8.0 Hz, 2H), 7.95 (s, 1H), 7.31 (d, J = 8.0 Hz, 2H), 7.17 (d, J = 8.3 Hz, 1H), 6.85 (d, J = 2.4 Hz, 1H), 3.63 (dt, J = 11.9, 4.6 Hz, 2H), 3.55 (s, 2H), 3.49 (ddd, J = 11.8, 9.1, 3.0 Hz, 2H), 2.36 (s, 3H), 1.45 (ddd, J = 13.4, 9.1, 4.1 Hz, 2H), 1.27 - 1.19 (m, 2H), 0.97 (s, 3H).
880	414.00	1.65	1H NMR (400 MHz, DMSO) δ 11.97 (s, 1H), 7.61 (s, 3H), 7.31 (d, J = 8.1 Hz, 1H), 6.69 (d, J = 8.7 Hz, 2H), 5.77 (s, 10H), 5.57 (s, 1H), 2.33 (s, 2H), 1.79 (s, 2H), 1.57 (s, 2H), 1.48 (s, 2H), 1.28 (s, 9H).
881	556.50	1.87	
882	494.30	2.08	
883	497.31	1.93	
884	567.33	1.72	
885	481.00	1.07	1H NMR (400 MHz, DMSO) δ 9.61 (s, 1H), 7.81 (s, 1H), 7.22 (d, J = 7.2 Hz, 1H), 7.05 (s, 2H), 6.91 (s, 3H), 6.72 (d, J = 10.1 Hz, 1H), 4.39 - 4.01 (m, 4H), 2.39 (s, 6H), 2.24 (s, 3H), 1.82 (s, 3H).
886	514.31	2.83	
887	556.50	2.03	
888	550.30	1.84	
889	533.28	1.82	
890	508.00	2.50	1H NMR (400 MHz, DMSO) δ 12.40 (s, 1H), 7.97 (d, J = 8.4 Hz, 2H), 7.77 (d, J = 8.0 Hz, 1H), 7.63 (dd, J = 8.3, 7.4 Hz, 1H), 7.36 (d, J = 8.3 Hz, 2H), 7.24 - 7.11 (m, 2H), 6.68 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 2.93 (dd, J = 13.7, 6.9 Hz, 1H), 2.61 (d, J = 8.9 Hz, 2H), 2.28 - 2.12 (m, 1H), 1.95 - 1.76 (m, 1H), 1.61 (d, J = 3.2 Hz, 6H), 1.42 (t, J = 12.1 Hz, 1H), 1.23 (d, J = 6.9 Hz, 6H), 0.88 (d, J = 6.3 Hz, 3H).
891	525.23	1.52	
892	494.50	1.85	1H NMR (400 MHz, DMSO-d6) δ 13.20 (s, 1H), 12.91 (s, 1H), 8.36 - 8.29 (m, 1H), 8.06 (dd, J = 21.6, 7.8 Hz, 1H), 7.92 (d, J = 1.9 Hz, 1H), 7.66

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(s, 0.5H), 7.40 (s, 0.5H), 7.54 (d, J = 7.8 Hz, 1H), 7.10 (d, J = 2.0 Hz, 1H), 7.03 (dd, J = 8.1, 2.2 Hz, 1H), 6.95 (d, J = 8.2 Hz, 1H), 6.51 (s, 1H), 3.87 (s, 3H), 2.31 (s, 3H), 2.15 - 2.09 (m, 3H), 2.04 (s, 3H).
893	521.50	1.95	1H NMR (400 MHz, DMSO-d6) δ 8.40 (d, J = 2.4 Hz, 1H), 8.32 (d, J = 2.1 Hz, 1H), 8.07 (d, J = 7.9 Hz, 1H), 7.98 (dd, J = 2.3, 1.0 Hz, 1H), 7.92 - 7.89 (m, 1H), 7.71 (d, J = 7.9 Hz, 1H), 7.18 - 7.05 (m, 4H), 3.89 (s, 3H), 2.33 (s, 3H), 2.13 (s, 3H), 2.04 (s, 3H).
894	543.22	2.35	
895	465.30	1.23	
896	509.30	1.50	1H NMR (400 MHz, DMSO-d6) δ 9.11 (d, J = 2.3 Hz, 1H), 8.23 (dd, J = 8.3, 2.4 Hz, 1H), 7.62 (d, J = 7.4 Hz, 1H), 7.49 (t, J = 6.9 Hz, 1H), 7.36 (d, J = 8.2 Hz, 1H), 7.04 (dd, J = 25.8, 6.0 Hz, 2H), 6.50 (d, J = 7.1 Hz, 1H), 6.23 (s, 2H), 3.05 (p, J = 6.9 Hz, 1H), 2.99 - 2.82 (m, 2H), 2.19 - 2.05 (m, 1H), 1.83 (dd, J = 11.8, 5.6 Hz, 1H), 1.62 (s, 3H), 1.57 (s, 3H), 1.37 (t, J = 12.0 Hz, 1H), 1.26 (d, J = 6.9 Hz, 6H), 0.82 (d, J = 6.2 Hz, 3H).
897	553.20	2.16	
898	552.30	1.68	1H NMR (400 MHz, DMSO-d6) δ 8.10 (d, J = 5.3 Hz, 1H), 7.61 (s, 1H), 7.45 (s, 1H), 7.25 (s, 1H), 7.13 - 6.95 (m, 3H), 6.47 (s, 1H), 3.35 (d, J = 7.5 Hz, 2H), 3.06 (s, 3H), 2.98 (s, 1H), 2.22 - 2.01 (m, 2H), 1.83 (dd, J = 11.8, 5.6 Hz, 1H), 1.63 (s, 3H), 1.58 (s, 3H), 1.38 (t, J = 12.0 Hz, 1H), 0.86 (dd, J = 11.1, 6.7 Hz, 9H).
899	496.00	2.06	1H NMR (400 MHz, DMSO) δ 12.58 (s, 1H), 11.71 (s, 1H), 8.25 (s, 1H), 7.83 (s, 2H), 6.92 (s, 2H), 6.45 (s, 1H), 2.67 (s, 2H), 2.26 (s, 3H), 1.98 (s, 6H), 1.65 (d, J = 27.5 Hz, 4H), 0.92 (s, 6H).
900	517.20	2.25	
901	552.20	1.97	
902	559.08	1.65	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
903	509.40	1.78	
904	530.40	1.97	
905	539.60	1.47	
906	526.30	2.85	
907	570.11	2.03	
908	505.31	2.55	
909	548.30	2.13	
910	496.30	1.60	
911	457.40	2.29	1H NMR (400 MHz, DMSO-d6) δ 7.98 (d, J = 7.8 Hz, 1H), 7.60 (d, J = 2.1 Hz, 1H), 7.12 (d, J = 7.8 Hz, 1H), 6.99 - 6.94 (m, 1H), 6.90 (s, 2H), 4.07 (s, 3H), 2.25 (s, 3H), 1.93 (s, 6H), 1.05 (s, 9H).
912	578.16	2.04	
913	515.25	1.64	
914	539.50	2.11	1H NMR (400 MHz, DMSO-d6) δ 13.80 (s, 1H), 12.51 (s, 1H), 8.08 (d, J = 7.8 Hz, 1H), 7.98 (s, 1H), 7.80 (d, J = 7.8 Hz, 1H), 7.25 - 7.18 (m, 1H), 7.13 (t, J = 1.8 Hz, 1H), 6.96 (s, 2H), 6.88 (s, 1H), 6.79 (dt, J = 10.9, 2.4 Hz, 1H), 4.41 (p, J = 6.0 Hz, 1H), 2.27 (s, 3H), 2.00 (s, 6H), 1.21 (d, J = 6.0 Hz, 6H).
915	539.30	1.71	
916	513.41	3.01	
917	552.30	1.99	
918	435.50	1.96	
919	537.43	2.98	
920	537.43	2.79	
921	509.60	1.54	1H NMR (400 MHz, DMSO-d6) δ 8.85 (d, J = 2.5 Hz, 1H), 8.31 (dd, J = 8.7, 2.5 Hz, 1H), 7.84 (s, 0H), 7.71 (d, J = 7.9 Hz, 1H), 7.65 (dd, J = 8.4 Hz, 1H), 7.19 (dd, J = 15.2, 7.6 Hz, 2H), 6.83 (d, J = 8.7 Hz, 1H), 6.72 (d, J = 8.3 Hz, 1H), 5.31 (hept, J = 6.1 Hz, 1H), 4.56 (s, 1H), 2.98 (d, J = 9.9 Hz, 1H), 2.44 (s, 1H), 2.31 (d, J = 9.8 Hz, 1H), 1.68 - 1.19 (m, 7H), 1.32 (d, J = 6.2 Hz, 6H).
922	555.37	1.78	
923	544.33	2.84	
924	540.30	1.98	1H NMR (400 MHz, DMSO-d6) δ 12.44 (s, 1H), 7.72 (d, J = 7.9 Hz, 1H), 7.65 (dd, J = 8.4, 7.3 Hz, 1H), 7.47 -

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			7.39 (m, 2H), 7.30 - 7.17 (m, 2H), 6.89 (dt, J = 10.8, 2.3 Hz, 1H), 6.72 (dd, J = 8.4, 0.8 Hz, 1H), 6.49 (s, 2H), 4.54 (s, 1H), 3.84 (d, J = 6.5 Hz, 2H), 2.99 (d, J = 9.4 Hz, 1H), 2.44 (s, 1H), 2.37 - 2.27 (m, 1H), 2.09 - 2.00 (m, 1H), 1.70 - 1.48 (m, 4H), 1.45 - 1.35 (m, 1H), 1.32 - 1.21 (m, 1H), 0.99 (d, J = 6.7 Hz, 6H).
925	525.30	1.45	1H NMR (400 MHz, DMSO-d6) δ 13.04 (s, 1H), 8.87 (d, J = 2.5 Hz, 1H), 8.32 (dd, J = 8.7, 2.5 Hz, 1H), 7.81 (d, J = 7.9 Hz, 1H), 7.65 (dd, J = 8.4, 7.3 Hz, 1H), 7.32 (d, J = 8.0 Hz, 1H), 7.22 (dd, J = 7.3, 0.8 Hz, 1H), 6.85 (d, J = 8.7 Hz, 1H), 6.72 (d, J = 8.3 Hz, 1H), 5.30 (h, J = 6.1 Hz, 1H), 4.24 - 3.70 (m, 2H), 2.27 - 2.16 (m, 1H), 2.16 - 2.04 (m, 1H), 1.48 - 1.34 (m, 1H), 1.32 (d, J = 6.2 Hz, 6H), 1.08 (d, J = 6.1 Hz, 3H), 0.97 (d, J = 6.5 Hz, 6H).
926	568.40	2.68	1H NMR (400 MHz, DMSO-d6) δ 12.51 (s, 1H), 7.75 (d, J = 7.9 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.38 (t, J = 1.8 Hz, 1H), 7.35 - 7.29 (m, 1H), 7.20 (dd, J = 7.6, 4.1 Hz, 2H), 6.89 (dt, J = 10.9, 2.3 Hz, 1H), 6.71 (d, J = 8.3 Hz, 1H), 6.47 (s, 1H), 3.83 (d, J = 6.5 Hz, 2H), 2.83 (t, J = 13.4 Hz, 2H), 2.74 (t, J = 6.1 Hz, 2H), 2.04 (dt, J = 13.3, 7.0 Hz, 1H), 1.96 - 1.85 (m, 2H), 1.76 (dq, J = 12.5, 6.2 Hz, 4H), 1.69 - 1.57 (m, 2H), 1.34 - 1.23 (m, 2H), 1.17 - 1.13 (m, 1H), 0.99 (d, J = 6.6 Hz, 6H).
927	466.40	1.75	
928	523.23	1.56	
929	485.50	2.73	
930	535.20	1.97	1H NMR (400 MHz, DMSO-d6) δ 8.25 (d, J = 2.2 Hz, 1H), 8.05 (d, J = 7.6 Hz, 1H), 7.85 (dd, J = 2.4, 1.1 Hz, 1H), 7.72 (dd, J = 4.8, 2.4 Hz, 1H), 7.65 (dd, J = 7.1, 2.5 Hz, 1H), 7.42 (d, J = 7.7 Hz, 1H), 6.91 (s, 2H), 5.88 (dd, J = 7.1, 4.8 Hz, 1H), 3.86 (s, 3H), 2.27 (s, 3H), 2.11 (s, 3H), 2.02 (s, 6H).
931	542.30	3.60	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
932	535.20	1.88	1H NMR (400 MHz, DMSO-d6) δ 11.81 (s, 1H), 8.04 (d, J = 7.7 Hz, 1H), 8.00 - 7.93 (m, 2H), 7.60 (d, J = 7.7 Hz, 1H), 7.52 (s, 1H), 7.19 - 7.02 (m, 3H), 6.89 (s, 2H), 6.21 (s, 1H), 3.80 (s, 3H), 2.74 (s, 2H), 2.24 (s, 3H), 2.01 (s, 6H).
933	559.21	2.09	
934	554.30	1.96	1H NMR (400 MHz, DMSO-d6) δ 12.54 (s, 1H), 7.72 - 7.61 (m, 2H), 7.48 - 7.38 (m, 2H), 7.33 (d, J = 7.8 Hz, 1H), 7.22 (d, J = 7.3 Hz, 1H), 6.89 (dt, J = 10.8, 2.4 Hz, 1H), 6.71 (d, J = 8.4 Hz, 1H), 6.46 (s, 2H), 4.23 (d, J = 3.9 Hz, 2H), 3.84 (d, J = 6.5 Hz, 2H), 2.03 (dt, J = 13.4, 6.6 Hz, 1H), 1.78 (dd, J = 8.3, 4.2 Hz, 2H), 1.69 (q, J = 9.9, 7.2 Hz, 5H), 1.41 (s, 1H), 1.33 (d, J = 8.6 Hz, 2H), 0.99 (d, J = 6.7 Hz, 6H).
935	545.28	1.97	
936	543.20	1.93	
937	495.20	1.97	
938	507.11	2.63	
939	504.30	2.21	
940	511.20	1.62	
941	559.30	2.00	1H NMR (400 MHz, DMSO-d6) δ 13.25 (s, 1H), 12.51 (s, 1H), 8.25 (dd, J = 7.2, 2.1 Hz, 1H), 7.90 - 7.81 (m, 2H), 7.77 (d, J = 2.0 Hz, 1H), 7.69 (dd, J = 8.3, 1.9 Hz, 1H), 7.53 (dd, J = 8.1, 4.6 Hz, 2H), 6.46 (t, J = 6.8 Hz, 1H), 3.96 (d, J = 6.5 Hz, 2H), 3.72 (d, J = 12.7 Hz, 2H), 2.88 (t, J = 12.1 Hz, 2H), 2.17 - 2.02 (m, 1H), 1.70 - 1.47 (m, 3H), 1.30 - 1.13 (m, 2H), 1.03 (d, J = 6.7 Hz, 6H), 0.92 (d, J = 6.2 Hz, 3H).
942	530.22	1.93	
943	479.40	1.94	1H NMR (400 MHz, DMSO-d6) δ 12.36 (s, 1H), 7.67 (d, J = 6.2 Hz, 1H), 7.53 (s, 1H), 7.22 (t, J = 7.9 Hz, 1H), 7.20 - 7.16 (m, 1H), 7.06 - 7.00 (m, 1H), 6.82 (dd, J = 7.8, 2.4 Hz, 1H), 6.76 (s, 2H), 6.41 (d, J = 6.2 Hz, 1H), 5.79 - 5.52 (m, 4H), 3.97 (s, 3H), 2.18

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(s, 3H), 1.93 (s, 6H).
944	542.13	1.92	
945	517.00	2.02	1H NMR (400 MHz, DMSO) δ 12.61 (s, 1H), 8.21 (s, 1H), 8.01 (s, 1H), 7.72 (s, 1H), 7.39 (d, J = 50.3 Hz, 5H), 7.00 (s, 2H), 6.79 (d, J = 8.3 Hz, 1H), 6.59 (s, 2H), 2.65 (s, 2H), 2.34 (s, 3H), 2.10 (s, 6H), 1.13 (s, 3H).
946	390.00	1.15	1H NMR (400 MHz, DMSO) δ 12.41 (s, 1H), 8.22 - 8.03 (m, 1H), 7.74 - 7.54 (m, 2H), 7.18 (d, J = 7.3 Hz, 1H), 6.66 (d, J = 8.3 Hz, 1H), 6.57 (dd, J = 7.5, 4.6 Hz, 1H), 6.46 (s, 2H), 2.57 (d, J = 7.3 Hz, 2H), 2.18 (s, 1H), 1.83 (dd, J = 11.8, 5.6 Hz, 1H), 1.50 (d, J = 9.2 Hz, 6H), 1.38 (t, J = 12.0 Hz, 1H), 0.86 (d, J = 6.3 Hz, 3H).
947	540.43	2.73	
948	494.30	1.56	
949	525.30	1.56	1H NMR (400 MHz, DMSO-d6) δ 12.42 (s, 1H), 8.83 (d, J = 2.5 Hz, 1H), 8.27 (dd, J = 8.8, 2.5 Hz, 1H), 7.74 (d, J = 7.9 Hz, 1H), 7.61 (t, J = 7.6 Hz, 1H), 7.17 (dd, J = 12.3, 7.6 Hz, 2H), 6.85 (d, J = 8.7 Hz, 1H), 6.65 (d, J = 8.4 Hz, 1H), 6.45 (s, 2H), 5.31 (hept, J = 6.3 Hz, 1H), 2.63 (s, 2H), 2.22 (dd, J = 13.1, 7.4 Hz, 1H), 1.87 (dd, J = 11.9, 5.7 Hz, 1H), 1.59 (d, J = 3.4 Hz, 6H), 1.41 (t, J = 12.0 Hz, 1H), 1.31 (d, J = 6.2 Hz, 6H), 0.88 (d, J = 6.2 Hz, 3H).
950	549.20	1.98	
951	524.50	1.09	
952	530.30	2.05	
953	554.40	1.93	
954	521.40	1.81	
955	582.29	2.05	
956	537.43	2.97	1H NMR (400 MHz, DMSO-d6) δ 12.35 (s, 1H), 8.09 - 8.06 (d, 1H), 7.62 - 7.56 (d, 2H), 7.33 (t, J = 7.91, 1H), 7.20 - 7.17 (d, 1H), 7.06 - 7.01 (d, 1H), 6.91 (s, 2H), 6.83 - 6.76 (m, 1H), 6.66 - 6.63 (d, 1H), 6.45 (s, 2 H), 3.84 (s, 3H), 2.25 (s, 3H), 2.02 (s, 6H)
957	524.30	1.67	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
958	525.40	1.82	1H NMR (400 MHz, DMSO-d6) δ 13.42 (s, 1H), 12.50 (s, 1H), 8.25 (dd, J = 7.3, 2.1 Hz, 1H), 7.92 - 7.80 (m, 2H), 7.70 - 7.58 (m, 2H), 7.52 (d, J = 8.0 Hz, 1H), 7.39 (t, J = 8.0 Hz, 1H), 7.07 - 6.99 (m, 1H), 6.46 (dd, J = 7.3, 6.3 Hz, 1H), 3.83 (d, J = 6.5 Hz, 2H), 3.70 (d, J = 12.1 Hz, 2H), 2.96 - 2.85 (m, 2H), 2.13 - 1.99 (m, 2H), 1.69 - 1.60 (m, 2H), 1.33 - 1.18 (m, 2H), 1.00 (d, J = 6.7 Hz, 6H), 0.93 (d, J = 6.2 Hz, 3H).
959	498.30	2.03	1H NMR (400 MHz, DMSO) δ 12.23 (s, 1H), 7.20 (dd, J = 9.7, 6.1 Hz, 2H), 7.07 (d, J = 8.1 Hz, 1H), 6.88 - 6.80 (m, 3H), 6.71 (s, 1H), 3.84 (s, 3H), 2.23 (s, 3H), 1.93 (s, 6H), 1.04 (s, 9H).
960	541.50	1.60	
961	579.23	2.23	
962	546.50	1.27	
963	441.20	1.89	
964	596.34	2.15	
965	526.30	2.13	1H NMR (400 MHz, DMSO-d6) δ 8.90 (d, J = 1.3 Hz, 1H), 8.25 (d, J = 1.3 Hz, 1H), 7.62 (d, J = 7.4 Hz, 1H), 7.48 (t, J = 7.7 Hz, 1H), 7.31 (d, J = 7.7 Hz, 1H), 6.99 (d, J = 7.3 Hz, 1H), 6.50 (d, J = 8.2 Hz, 1H), 6.25 (s, 2H), 5.28 (h, J = 6.2 Hz, 1H), 2.93 (d, J = 23.0 Hz, 2H), 2.07 (s, 1H), 1.83 (dd, J = 11.8, 5.6 Hz, 1H), 1.63 (s, 3H), 1.58 (s, 3H), 1.35 (d, J = 1.6 Hz, 3H), 1.34 (d, J = 1.7 Hz, 3H), 0.80 (d, J = 6.3 Hz, 3H).
966	498.40	2.55	
967	579.20	2.38	
968	471.69	3.25	
969	618.40	3.26	
970	727.50	0.68	1H NMR (300 MHz, DMSO) δ 8.24 (dd, J = 15.6, 8.2 Hz, 1H), 7.80 - 7.66 (m, 1H), 7.66 - 7.39 (m, 10H), 7.31 - 7.22 (m, 4H), 7.11 (d, J = 7.2 Hz, 1H), 6.74 (t, J = 10.3 Hz, 1H), 6.55 (d, J = 7.9 Hz, 1H), 6.27 (s, 2H), 1.92 (d, J = 1.9 Hz, 12H).
971	553.28	1.80	
972	542.30	1.77	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
973	554.50	2.03	
974	533.20	1.99	
975	546.20	1.72	
976	495.30	1.76	1H NMR (400 MHz, DMSO-d6) δ 7.98 (d, J = 7.7 Hz, 1H), 7.41 (dd, J = 8.2, 7.4 Hz, 1H), 7.09 - 7.01 (m, 2H), 6.88 - 6.81 (m, 2H), 6.44 (dd, J = 8.2, 0.9 Hz, 1H), 6.35 (dt, J = 3.2, 1.6 Hz, 1H), 6.07 (s, 2H), 4.12 (q, J = 2.8 Hz, 2H), 3.68 (t, J = 5.5 Hz, 2H), 2.23 (d, J = 5.4 Hz, 5H), 1.98 (s, 6H).
977	528.12	2.75	1H NMR (400 MHz, DMSO-d6) δ 12.23 (s, 1H), 11.19 (s, 1H), 8.17 - 8.14 (d, 1H), 7.79 - 7.76 (d, 1H), 7.65 (t, J = 8.02, 1H) 7.53 - 7.50 (d, 1H), 7.47 - 7.44 (d, 1H), 7.26 - 7.23 (d, 1H), 7.15 - 7.09 (m, 2H), 6.97 (s, 2 H), 6.72 - 6.69 (d, 1H), 6.54 (s, 2H), 6.22 (s, 1H), 2.31 (s, 3H), 2.04 (s, 6H)
978	571.32	2.42	
979	525.25	1.66	
980	520.20	1.85	
981	536.17	1.94	
982	484.29	1.41	
983	442.58	2.86	
984	555.13	2.98	
985	446.00	1.35	
986	462.50	1.32	
987	523.30	1.58	1H NMR (400 MHz, DMSO-d6) δ 12.50 (s, 1H), 8.87 (d, J = 2.5 Hz, 1H), 8.33 (dd, J = 8.8, 2.5 Hz, 1H), 7.77 (d, J = 8.0 Hz, 1H), 7.63 (dd, J = 8.4, 7.3 Hz, 1H), 7.24 (d, J = 7.9 Hz, 1H), 7.19 (d, J = 7.4 Hz, 1H), 6.89 (d, J = 8.7 Hz, 1H), 6.71 (d, J = 8.3 Hz, 1H), 5.32 (p, J = 6.2 Hz, 1H), 3.50 (q, J = 10.2, 9.6 Hz, 2H), 2.69 (t, J = 6.6 Hz, 2H), 2.02 (t, J = 6.5 Hz, 2H), 1.98 - 1.86 (m, 1H), 1.86 - 1.75 (m, 1H), 1.68 (q, J = 7.9, 6.0 Hz, 4H), 1.32 (d, J = 6.2 Hz, 6H)(NH2 under H2O peak).
988	540.50	1.69	
989	566.50	2.20	1H NMR (400 MHz, DMSO-d6) δ 8.32 (d, J = 2.2 Hz, 1H), 8.12 (d, J = 7.9 Hz, 1H), 7.91 (d, J = 2.2 Hz, 1H), 7.81 (d, J

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			= 7.9 Hz, 1H), 7.23 (dt, J = 9.8, 2.0 Hz, 1H), 7.13 (t, J = 1.8 Hz, 3H), 6.95 (s, 2H), 6.79 (dt, J = 10.7, 2.3 Hz, 1H), 4.41 (p, J = 6.1 Hz, 1H), 2.27 (s, 3H), 2.00 (s, 6H), 1.21 (d, J = 6.0 Hz, 6H).
990	499.70	1.83	
991	542.40	1.96	1H NMR (400 MHz, DMSO-d6) δ 12.82 (s, 1H), 7.83 (d, J = 7.9 Hz, 1H), 7.65 (dd, J = 8.4, 7.3 Hz, 1H), 7.49 (d, J = 8.0 Hz, 1H), 7.46 (dd, J = 8.6, 2.2 Hz, 2H), 7.21 (d, J = 7.3 Hz, 1H), 6.92 (dd, J = 10.8, 2.3 Hz, 1H), 6.71 (d, J = 8.4 Hz, 1H), 6.44 (s, 2H), 3.85 (d, J = 6.5 Hz, 2H), 3.71 (d, J = 12.8 Hz, 2H), 2.85 (t, J = 12.1 Hz, 2H), 2.05 (dt, J = 13.3, 6.7 Hz, 1H), 1.62 (d, J = 13.3 Hz, 2H), 1.58 - 1.48 (m, 1H), 1.15 (q, J = 10.8, 10.2 Hz, 2H), 1.00 (d, J = 6.7 Hz, 6H), 0.91 (d, J = 6.3 Hz, 3H).
992	525.40	1.50	
993	554.10	2.16	
994	573.30	1.93	
995	539.30	1.53	
996	547.20	2.25	1H NMR (400 MHz, DMSO-d6) δ 8.05 (s, 1H), 7.64 (s, 2H), 7.44 (s, 2H), 7.07 (s, 1H), 6.90 (s, 4H), 6.45 (s, 1H), 6.07 (s, 2H), 3.93 (s, 2H), 2.27 (s, 2H), 2.02 (s, 5H), 1.71 (s, 2H).
997	530.27	2.03	
998	527.20	1.42	1H NMR (400 MHz, DMSO) δ 13.12 (s, 1H), 8.86 (s, 1H), 8.32 (dd, J = 8.8, 2.3 Hz, 1H), 7.79 (s, 1H), 7.62 (s, 1H), 7.31 (s, 1H), 7.21 (d, J = 6.7 Hz, 1H), 6.86 (d, J = 8.7 Hz, 1H), 6.70 (s, 1H), 6.47 (d, J = 22.7 Hz, 2H), 5.21 (dd, J = 11.2, 5.9 Hz, 1H), 4.82 (t, J = 5.7 Hz, 1H), 4.02 (s, 2H), 3.59 (dt, J = 11.3, 5.7 Hz, 1H), 3.56 - 3.46 (m, 1H), 1.96 (s, 2H), 1.65 (s, 2H), 1.26 (d, J = 6.3 Hz, 3H), 1.08 (t, J = 8.4 Hz, 6H).
999	582.20	2.20	
1000	521.20	2.16	
1001	545.11	1.50	
1002	532.20	2.21	
1003	401.00	1.90	1H NMR (400 MHz, DMSO) δ 12.08 (s,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 7.59 (s, 3H), 7.24 (d, J = 8.1 Hz, 1H), 6.64 (d, J = 1.6 Hz, 2H), 3.02 (s, 1H), 1.68 (s, 6H), 1.48 (s, 2H), 1.28 (s, 9H).
1004	556.35	1.83	
1005	530.23	1.86	
1006	538.19	1.91	
1007	560.20	2.07	
1008	432.50	1.41	
1009	542.30	2.75	
1010	528.40	1.78	
1011	525.60	1.62	
1012	603.34	1.87	
1013	556.40	2.10	1H NMR (400 MHz, DMSO-d6) δ 12.76 (s, 1H), 7.81 (d, J = 7.8 Hz, 1H), 7.64 (t, J = 7.8 Hz, 1H), 7.52 - 7.43 (m, 3H), 7.22 (d, J = 7.2 Hz, 1H), 6.91 (d, J = 10.7 Hz, 1H), 6.69 (d, J = 8.3 Hz, 1H), 6.48 (s, 2H), 3.85 (d, J = 6.6 Hz, 2H), 3.69 (d, J = 12.5 Hz, 2H), 3.36 - 3.34 (m, 2H), 2.56 - 2.53 (m, 1H), 2.42 (t, J = 11.8 Hz, 1H), 2.05 (dt, J = 13.3, 6.7 Hz, 1H), 1.79 - 1.69 (m, 2H), 1.00 (d, J = 6.8 Hz, 6H), 0.83 (d, J = 6.5 Hz, 6H).
1014	555.43	3.03	1H NMR (400 MHz, DMSO-d6) δ 7.78 - 7.75 (d, 1H), 7.50 - 7.47 (m, 3H), 7.30 - 7.22 (m, 2H), 7.10 - 7.07 (d, 1H), 6.95 - 6.85 (m, 2H), 3.86 - 3.83 (d, 2H), 3.20 (m, 4H), 2.10 - 2.00 (m, 1H), 1.33 (m, 4H), 1.01 - 0.98 (d , 6H), 0.95 (s, 6H)
1015	489.50	1.95	1H NMR (400 MHz, DMSO-d6) δ 12.25 (s, 1H), 8.07 (d, J = 7.9 Hz, 1H), 7.71 (d, J = 8.1 Hz, 3H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.25 - 7.19 (m, 3H), 7.16 - 7.11 (m, 2H), 7.07 (dd, J = 8.3, 2.1 Hz, 1H), 6.71 (dd, J = 8.5, 0.8 Hz, 1H), 6.52 (s, 2H), 2.32 (s, 3H), 2.31 (s, 3H), 2.06 (s, 3H).
1016	531.30	2.32	
1017	525.30	1.82	
1018	510.34	1.42	
1019	534.50	2.07	
1020	524.24	1.82	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1021	572.16	2.16	
1022	566.20	2.10	
1023	525.18	1.65	
1024	477.25	1.81	
1025	511.30	1.49	
1026	519.28	1.82	
1027	532.50	1.54	
1028	546.22	1.79	
1029	470.30	1.28	
1030	530.22	1.91	
1031	469.23	2.08	
1032	503.50	2.07	
1033	493.00	1.86	1H NMR (400 MHz, DMSO) δ 13.80 (s, 1H), 12.38 (s, 1H), 8.04 (d, J = 7.9 Hz, 1H), 7.98 (s, 1H), 7.73 (d, J = 8.8 Hz, 2H), 7.66 (s, 1H), 3.76 (s, 3H), 2.28 (s, 3H), 2.02 (s, 6H).
1034	520.20	2.08	
1035	503.21	3.11	
1036	424.20	1.63	
1037	431.40	1.90	
1038	476.50	1.21	1H NMR (400 MHz, DMSO-d6) δ 7.82 (d, J = 8.0 Hz, 1H), 7.65 (dd, J = 8.4, 7.3 Hz, 1H), 7.19 (d, J = 6.9 Hz, 1H), 7.01 (d, J = 7.9 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 3.70 - 3.60 (m, 3H), 2.81 (t, J = 12.0 Hz, 2H), 1.74 (d, J =2 Hz, 2H), 1.41 - 1.30 (m, 2H), 1.28 (s, 9H), 1.04 (s, 6H). The three proton are buried in the water peak, thus the integration is skewed
1039	463.30	1.60	
1040	525.40	1.42	1H NMR (400 MHz, DMSO-d6) δ 8.88 (d, J = 2.5 Hz, 1H), 8.34 (dd, J = 8.8, 2.5 Hz, 1H), 8.24 - 8.14 (m, 1H), 7.84 - 7.71 (m, 1H), 7.65 (t, J = 7.9 Hz, 1H), 7.19 (d, J = 7.3 Hz, 1H), 6.87 (d, J = 8.8 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 5.28 (p, J = 6.1 Hz, 1H), 4.20 - 4.03 (m, 1H), 2.29 - 2.16 (m, 1H), 2.00 - 1.88 (m, 2H), 1.80 - 1.65 (m, 1H), 1.46 (s, 3H), 1.30 (s, 6H), 1.29 (s, 3H), 1.09 (d, J = 6.2 Hz, 3H).
1041	530.30	2.00	1H NMR (400 MHz, DMSO) δ 13.75 (s, 1H), 12.59 (s, 1H), 7.97 (s, 1H), 7.69

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(d, J = 7.9 Hz, 1H), 7.48 (s, 1H), 7.40 (d, J = 9.9 Hz, 1H), 7.26 (d, J = 8.0 Hz, 1H), 6.93 - 6.84 (m, 2H), 3.83 (d, J = 6.6 Hz, 2H), 3.17 (s, 0H), 2.58 (d, J = 9.4 Hz, 2H), 2.22 (s, 1H), 2.05 (dt, J = 13.3, 6.7 Hz, 1H), 1.88 (dd, J = 11.8, 5.5 Hz, 1H), 1.60 (d, J = 12.9 Hz, 6H), 1.44 (t, J = 12.1 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.87 (d, J = 6.3 Hz, 3H).
1042	502.36	1.81	1H NMR (400 MHz, DMSO) δ 7.98 (s, 1H), 7.80 (d, J = 7.9 Hz, 1H), 7.50 (d, J = 7.9 Hz, 1H), 7.47 - 7.40 (m, J = 10.2, 1.5 Hz, 2H), 6.91 (dt, J = 10.9, 2.3 Hz, 1H), 6.88 - 6.84 (m, 1H), 4.82 - 4.66 (m, 1H), 3.72 - 3.59 (m, J = 12.9 Hz, 2H), 2.93 - 2.78 (m, J = 11.9 Hz, 2H), 1.70 - 1.50 (m, J = 20.8, 9.2 Hz, 3H), 1.30 (d, J = 6.0 Hz, 6H), 1.27 - 1.09 (m, 2H), 0.93 (d, J = 6.3 Hz, 3H).
1043	565.27	1.87	
1044	508.50	1.94	1H NMR (400 MHz, DMSO-d6) δ 13.81 (d, J = 6.2 Hz, 1H), 12.50 - 12.35 (m, 1H), 8.33 (d, J = 2.1 Hz, 1H), 8.07 (d, J = 7.9 Hz, 1H), 8.03 - 7.93 (m, 2H), 7.70 (d, J = 7.8 Hz, 1H), 6.96 (s, 2H), 6.92 - 6.84 (m, 1H), 3.88 (s, 3H), 2.29 (s, 3H), 2.13 (s, 3H), 2.01 (s, 6H).
1045	469.26	2.08	
1046	427.19	1.83	
1047	504.50	2.10	
1048	531.42	2.96	
1049	523.20	1.87	
1050	453.30	1.93	
1051	528.30	2.93	
1052	508.30	2.19	
1053	549.17	2.00	
1054	512.51	2.61	
1055	479.30	1.17	1H NMR (500 MHz, DMSO-d6) δ 9.38 (s, 1H), 7.67 (t, J = 8.0 Hz, 1H), 7.18 - 7.07 (m, 4H), 7.04 (d, J = 7.3 Hz, 1H), 6.91 (s, 2H), 6.80 (d, J = 8.5 Hz, 1H), 4.31 (d, J = 13.7 Hz, 1H), 4.16 (dd, J = 17.7, 10.2 Hz, 2H), 3.54 (d, J = 13.0 Hz, 1H), 3.39 - 3.25 (m, 2H), 3.22 - 3.08 (m, 2H), 3.02 (d, J = 16.8 Hz, 1H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			2.22 (d, J = 8.3 Hz, 9H).
1056	542.23	3.03	
1057	493.30	2.57	
1058	450.00	1.17	
1059	538.60	1.19	
1060	564.29	2.14	
1061	540.40	2.98	
1062	553.50	1.82	
1063	541.33	3.03	1H NMR (500 MHz, DMSO-d6) δ 12.90 (s, 1H), 8.80 - 8.78 (d, 1H), 8.19 - 8.17 (d, 2H), 7.81 - 7.73 (m, 2H), 7.49 (s, 1H), 7.43 - 7.39 (d, 1H), 7.31 - 7.28 (d, 1H), 6.92 - 6.88 (d, 1H), 3.84 - 3.81 (d, 2H), 2.51 - 2.36 (m, 1H), 2.21 (m, 1H), 2.10 - 1.99 (m, 1H), 1.89 - 1.82 (m, 1H), 1.58 (s, 6H), 1.34 (t, J = 11.87, 1H), 1.00 - 0.97 (d, 6H), 0.82 - 0.79 (d, 3H)
1064	439.40	1.94	
1065	556.60	1.96	
1066	497.50	1.38	1H NMR (400 MHz, DMSO-d6) δ 8.30 (d, J = 2.2 Hz, 1H), 8.24 (d, J = 8.2 Hz, 1H), 8.02 (d, J = 8.4 Hz, 2H), 8.01 - 7.98 (m, 1H), 7.94 - 7.91 (m, 0H), 7.88 (d, J = 2.2 Hz, 1H), 7.31 (d, J = 8.3 Hz, 2H), 7.19 (d, J = 8.3 Hz, 1H), 7.09 (d, J = 8.1 Hz, 0H), 3.63 (ddd, J = 16.6, 10.6, 5.6 Hz, 2H), 3.54 (s, 2H), 3.53 - 3.46 (m, 2H), 2.37 (s, (3+1.5)H), 1.51 (ddt, J = 13.5, 9.3, 4.7 Hz, 1H), 1.41 (ddd, J = 13.4, 9.2, 4.2 Hz, 2H), 1.33 - 1.25 (m, 1H), 1.24 - 1.16 (m, 2H), 1.02 (s, 1.5H), 0.93 (s, 3H). Rotamers in 2:1 ratio
1067	498.30	1.84	1H NMR (400 MHz, DMSO-d6) δ 7.76 (d, J = 2.2 Hz, 1H), 7.55 (d, J = 7.7 Hz, 1H), 7.47 (dd, J = 8.3, 7.3 Hz, 1H), 7.03 - 6.91 (m, 2H), 6.61 (d, J = 2.3 Hz, 1H), 6.49 (d, J = 8.2 Hz, 1H), 6.25 (s, 2H), 4.52 (hept, J = 6.7 Hz, 1H), 2.92 (t, J = 10.6 Hz, 1H), 2.81 (dd, J = 10.7, 7.4 Hz, 1H), 2.07 (q, J = 6.7, 6.1 Hz, 1H), 1.78 (dd, J = 11.6, 5.6 Hz, 1H), 1.60 (s, 3H), 1.54 (s, 3H), 1.45 (d, J = 6.7 Hz, 6H), 1.34 (t, J = 12.0 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 0.78 (d, J = 6.2 Hz, 3H).
1068	579.24	2.30	
1069	541.33	3.16	1H NMR (400 MHz, DMSO-d6) δ 8.76 - 8.74 (d, 1H), 8.19 (m, 2H), 7.84 - 7.74 (m, 2H), 7.52 - 7.45 (m, 3H), 6.95 - 6.91 (d, 1H), 3.86 - 3.83 (d, 2H), 3.23 (m, 4H), 2.10 - 1.99 (m, 1H), 1.33 (m, 4H), 1.01 - 0.98 (d, 6H), 0.94 (s, 6H)
1070	485.50	3.11	
1071	480.30	1.84	
1072	531.30	1.83	
1073	556.31	2.06	
1074	485.40	1.42	
1075	495.00	1.30	1H NMR (400 MHz, DMSO) δ 10.31 (s, 1H), 9.76 (s, 1H), 7.76 (s, 1H), 7.21 (d, J = 27.9 Hz, 2H), 6.91 (d, J = 51.5 Hz, 6H), 3.99 (d, J = 59.4 Hz, 4H), 3.61 (s, 3H), 2.29 (s, 6H), 2.22 (s, 3H), 1.55 (s, 3H).
1076	553.50	1.76	
1077	541.19	2.27	
1078	466.30	1.37	
1079	562.20	2.25	1H NMR (400 MHz, DMSO) δ 12.60 (s, 1H), 11.92 (s, 1H), 8.27 (s, 1H), 8.14 (d, J = 7.9 Hz, 1H), 7.84 (s, 1H), 7.71 (d, J = 8.8 Hz, 3H), 6.96 (s, 4H), 6.47 (s, 1H), 3.77 (d, J = 6.5 Hz, 2H), 2.29 (s, 3H), 2.01 (d, J = 25.1 Hz, 7H), 0.95 (d, J = 6.7 Hz, 6H).
1080	589.31	1.91	
1081	497.21	1.62	
1082	473.20	1.71	
1083	516.20	2.01	1H NMR (400 MHz, DMSO-d6) δ 13.75 (s, 1H), 12.97 (s, 1H), 8.05 - 7.93 (m, 1H), 7.80 (d, J = 7.9 Hz, 1H), 7.55 - 7.42 (m, 3H), 6.92 (dd, J = 10.8, 2.3 Hz, 1H), 6.85 (t, J = 2.2 Hz, 1H), 3.85 (d, J = 6.5 Hz, 2H), 3.67 (d, J = 12.7 Hz, 2H), 2.89 - 2.79 (m, 2H), 2.05 (dt, J = 13.3, 6.6 Hz, 1H), 1.67 - 1.57 (m, 2H), 1.27 - 1.22 (m, 1H), 1.21 - 1.13 (m, 2H), 1.00 (d, J = 6.7 Hz, 6H), 0.93 (d, J = 6.2 Hz, 3H).
1084	467.31	1.65	
1085	480.00	1.43	1H NMR (400 MHz, DMSO) δ 12.40 (s,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 11.29 (s, 1H), 7.97 (d, J = 7.2 Hz, 1H), 7.74 (s, 1H), 7.26 (d, J = 3.7 Hz, 2H), 7.15 (t, J = 12.4 Hz, 2H), 6.71 (s, 2H), 6.32 (s, 1H), 4.57 (s, 1H), 4.02 (d, J = 12.6 Hz, 1H), 3.83 (d, J = 12.6 Hz, 1H), 2.20 (d, J = 15.0 Hz, 9H), 1.56 (s, 3H), 1.37 (s, 3H).
1086	478.30	1.96	
1087	511.26	1.62	
1088	572.19	1.89	
1089	480.30	1.89	
1090	571.34	3.21	1H NMR (400 MHz, DMSO) δ 8.05 (t, J = 7.91, 1H), 8.88 - 8.85 (d, 1H), 7.78 - 7.75 (d, 1H), 7.54 - 7.47 (m, 3H), 7.21 - 7.18 (d, 1H), 6.95 - 6.91 (d, 1H), 3.86 - 3.83 (m, 5H), 3.23 (m, 4H), 2.10 - 1.99 (m, 1H), 1.31 (m, 4H), 1.01 - 0.98 (d, 6H), 0.94 (s, 6H)
1091	524.30	2.32	
1092	575.31	1.84	
1093	464.30	1.36	
1094	524.40	1.86	1H NMR (400 MHz, DMSO-d6) δ 13.03 (s, 1H), 8.09 - 8.00 (m, 2H), 7.85 (d, J = 8.0 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.43 (d, J = 8.0 Hz, 1H), 7.20 (d, J = 7.3 Hz, 1H), 7.04 (d, J = 8.9 Hz, 2H), 6.70 (d, J = 8.4 Hz, 1H), 6.42 (s, 2H), 3.81 (d, J = 6.5 Hz, 2H), 3.67 (d, J = 12.7 Hz, 2H), 2.88 (t, J = 12.4 Hz, 2H), 2.03 (td, J = 14.0, 13.5, 6.9 Hz, 1H), 1.74 - 1.47 (m, 3H), 1.21 (q, J = 13.7, 12.8 Hz, 2H), 1.00 (d, J = 6.7 Hz, 6H), 0.92 (d, J = 6.4 Hz, 3H).
1095	524.10	2.00	
1096	515.30	1.42	
1097	570.60	2.04	
1098	570.50	2.13	
1099	528.19	1.87	
1100	500.20	1.63	
1101	479.00	1.22	1H NMR (400 MHz, DMSO) δ 7.56 (s, 1H), 7.26 (d, J = 29.4 Hz, 4H), 6.89 (d, J = 7.3 Hz, 1H), 6.80 - 6.68 (m, 3H), 4.69 (s, 1H), 4.16 (d, J = 12.9 Hz, 1H), 2.20 (d, J = 25.7 Hz, 9H), 1.63 (s, 3H), 1.45 (s, 3H).

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1102	503.00	1.86	1H NMR (400 MHz, DMSO) δ 12.66 (s, 1H), 7.75 (s, 2H), 7.65 (s, 1H), 7.57 (s, 1H), 7.46 - 7.36 (m, 3H), 7.24 (d, J = 7.2 Hz, 1H), 6.93 (s, 2H), 6.73 (d, J = 8.4 Hz, 1H), 6.48 (s, 2H), 2.29 (d, J = 13.2 Hz, 6H), 2.05 (s, 6H).
1103	536.50	1.91	
1104	459.18	1.96	
1105	560.50	2.08	
1106	456.50	2.78	
1107	523.20	1.79	
1108	557.25	2.41	
1109	542.30	1.92	
1110	530.20	2.09	
1111	544.50	1.77	
1112	559.32	1.87	
1113	613.50	2.15	
1114	526.20	2.19	1H NMR (400 MHz, Methanol-d4) δ 8.23 (d, J = 2.7 Hz, 1H), 8.08 (dd, J = 7.2, 1.7 Hz, 2H), 7.76 (d, J = 8.3 Hz, 1H), 7.66 (d, J = 7.2 Hz, 1H), 7.59 (dd, J = 8.4, 6.7 Hz, 2H), 6.97 (d, J = 8.3 Hz, 1H), 5.98 (d, J = 2.8 Hz, 1H), 3.89 (s, 2H), 2.65 (d, J = 10.4 Hz, 1H), 2.52 (s, 1H), 2.17 (s, 1H), 1.85 (dd, J = 11.7, 5.7 Hz, 1H), 1.59 (d, J = 12.2 Hz, 6H), 1.45 (t, J = 12.1 Hz, 1H), 1.03 (s, 9H), 0.76 (d, J = 6.3 Hz, 3H).
1115	520.20	1.47	
1116	541.33	1.58	
1117	514.30	2.86	
1118	517.20	2.23	
1119	532.20	1.87	
1120	503.21	3.01	
1121	556.50	1.91	
1122	560.50	1.60	1H NMR (400 MHz, DMSO-d6) δ 12.41 (s, 1H), 10.03 (s, 1H), 8.12 (d, J = 7.9 Hz, 1H), 7.80 - 7.68 (m, 4H), 7.57 (s, 1H), 7.44 - 7.37 (m, 3H), 7.32 (d, J = 7.2 Hz, 1H), 6.95 (s, 2H), 6.84 (d, J = 8.5 Hz, 1H), 3.62 - 3.53 (m, 2H), 3.23 (q, J = 5.7 Hz, 2H), 2.78 (d, J = 4.6 Hz, 6H), 2.28 (s, 3H), 2.01 (s, 6H).
1123	528.30	1.57	1H NMR (400 MHz, DMSO-d6) δ 13.04 (s, 1H), 7.81 (dd, J = 7.8, 1.6 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 7.65 (dd, J = 8.4, 7.3 Hz, 1H), 7.52 - 7.46 (m, 1H), 7.46 - 7.35 (m, 2H), 7.23 (dd, J = 7.3, 0.7 Hz, 1H), 6.88 (dt, J = 10.9, 2.3 Hz, 1H), 6.73 (d, J = 8.4 Hz, 1H), 4.73 (h, J = 6.1 Hz, 1H), 4.08 - 3.95 (m, 2H), 2.06 - 1.91 (m, 2H), 1.73 - 1.61 (m, 2H), 1.31 (d, J = 6.0 Hz, 6H), 1.09 (d, J = 6.2 Hz, 6H).
1124	439.30	1.90	
1125	509.30	1.66	1H NMR (400 MHz, DMSO-d6) δ 8.89 - 8.78 (m, 1H), 8.30 (dd, J = 8.8, 2.5 Hz, 1H), 7.56 (s, 1H), 7.48 (d, J = 8.1 Hz, 1H), 7.03 (s, 2H), 6.92 (d, J = 8.6 Hz, 1H), 6.16 (s, 2H), 4.25 (tt, J = 7.1, 3.7 Hz, 3H), 1.87 (s, 2H), 1.57 (q, J = 8.6, 8.2 Hz, 2H), 1.07 (d, J = 6.1 Hz, 6H), 0.82 - 0.74 (m, 2H), 0.74 - 0.64 (m, 2H).
1126	546.13	2.47	
1127	528.20	1.86	
1128	456.38	2.97	
1129	490.40	1.84	1H NMR (400 MHz, DMSO-d6) δ 12.56 (s, 1H), 12.16 (s, 1H), 8.27 (dd, J = 7.3, 2.1 Hz, 1H), 8.10 (d, J = 7.9 Hz, 1H), 7.84 (s, 1H), 7.75 - 7.68 (m, 3H), 7.22 (d, J = 8.1 Hz, 2H), 7.16 - 7.10 (m, 2H), 7.07 (dd, J = 8.2, 2.1 Hz, 1H), 6.46 (dd, J = 7.3, 6.3 Hz, 1H), 2.34 - 2.29 (m, 6H), 2.07 (s, 3H).
1130	535.20	1.82	
1131	495.40	3.25	
1132	537.50	1.76	
1133	534.20	2.19	
1134	525.33	1.67	1H NMR (400 MHz, DMSO) δ 12.97 (s, 1H), 8.90 (s, 1H), 8.37 (d, J = 8.7 Hz, 1H), 7.86 (d, J = 8.0 Hz, 1H), 7.66 (s, 1H), 7.46 (d, J = 8.0 Hz, 1H), 7.22 (d, J = 7.2 Hz, 1H), 6.86 (d, J = 8.7 Hz, 1H), 6.74 (d, J = 8.4 Hz, 1H), 5.32 (s, 1H), 3.29 (s, 4H), 1.40 (s, 4H), 1.32 (d, J = 6.2 Hz, 6H), 0.96 (s, 6H).
1135	553.50	1.83	
1136	525.40	1.73	
1137	427.27	2.61	
1138	565.20	2.26	1H NMR (400 MHz, DMSO-d6) δ 8.08

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(d, J = 7.7 Hz, 1H), 7.61 (d, J = 7.7 Hz, 1H), 7.42 (dd, J = 8.2, 7.3 Hz, 1H), 7.15 (ddd, J = 10.1, 2.4, 1.3 Hz, 1H), 7.11 - 7.04 (m, 2H), 6.91 (s, 2H), 6.69 (dt, J = 10.8, 2.3 Hz, 1H), 6.45 (dd, J = 8.2, 0.9 Hz, 1H), 6.08 (s, 2H), 4.38 (p, J = 6.0 Hz, 1H), 2.26 (s, 3H), 2.00 (s, 6H), 1.21 (d, J = 6.0 Hz, 6H).
1139	534.12	2.71	
1140	537.43	2.95	
1141	483.20	1.28	
1142	502.14	1.78	
1143	548.20	2.12	
1144	556.60	1.96	
1145	525.33	1.64	
1146	549.43	2.81	1H NMR (400 MHz, DMSO-d6) δ 12.19 (s, 1H), 8.13 - 8.10 (d, 1H), 7.74 - 7.63 (m, 2H), 7.49 - 7.46 (d, 1H), 7.29 - 7.23 (m, 2H), 7.00 - 6.96 (m, 3H), 6.74 - 6.71 (d, 1H), 6.56 (s, 2H), 3.77 (s, 3H), 3.61 (s, 3H), 2.26 (s, 3H), 2.04 (s, 6H)
1147	547.33	2.68	
1148	475.20	2.11	1H NMR (400 MHz, DMSO) δ 12.10 (s, 1H), 8.02 (s, 2H), 7.95 (d, J = 7.8 Hz, 1H), 7.67 (s, 1H), 7.12 (d, J = 7.8 Hz, 1H), 6.90 (s, 2H), 2.25 (s, 3H), 1.97 (s, 7H), 1.04 (s, 10H).
1149	558.30	2.00	
1150	551.50	2.09	1H NMR (400 MHz, DMSO-d6) δ 12.37 (s, 1H), 8.09 (d, J = 7.9 Hz, 1H), 7.81 (dd, J = 7.1, 2.4 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.29 - 7.20 (m, 2H), 7.19 - 7.06 (m, 4H), 6.81 (dt, J = 10.8, 2.3 Hz, 1H), 6.71 (d, J = 8.4 Hz, 1H), 6.58 - 6.46 (m, 2H), 4.45 (h, J = 6.0 Hz, 1H), 2.32 (d, J = 3.2 Hz, 3H), 2.04 (s, 3H), 1.22 (d, J = 5.9 Hz, 6H).
1151	538.20	1.87	
1152	433.00	1.94	1H NMR (400 MHz, DMSO) δ 12.24 (s, 1H), 7.21 (s, 1H), 7.16 (s, 1H), 7.04 (d, J = 8.4 Hz, 1H), 6.97 (s, 1H), 6.81 (d, J = 9.6 Hz, 1H), 5.68 (s, 1H), 4.21 (s, 2H), 1.30 (d, J = 48.6 Hz, 14H), 1.11 - 0.97 (m, 5H), 0.60 (s, 3H).

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1153	498.00	2.34	1H NMR (400 MHz, DMSO) δ 12.24 (s, 1H), 8.26 (s, 1H), 7.96 (d, J = 7.8 Hz, 1H), 7.11 (dd, J = 25.3, 6.4 Hz, 2H), 6.91 (s, 2H), 3.89 (s, 3H), 2.70 (s, 3H), 2.26 (s, 3H), 1.94 (s, 6H), 1.05 (s, 9H).
1154	405.21	1.94	
1155	546.20	1.81	
1156	556.50	2.03	
1157	556.50	1.82	
1158	491.00	2.08	1H NMR (400 MHz, DMSO) δ 12.38 (s, 1H), 7.63 (s, 1H), 7.47 (d, J = 17.9 Hz, 2H), 7.17 (d, J = 7.0 Hz, 1H), 6.96 (d, J = 8.7 Hz, 1H), 6.75 (d, J = 8.4 Hz, 3H), 6.69 (d, J = 8.1 Hz, 1H), 6.47 (s, 2H), 5.66 (s, 2H), 2.56 (s, 2H), 2.18 (s, 3H), 1.92 (s, 6H), 1.56 (s, 2H), 0.87 (s, 3H).
1159	542.20	1.58	1H NMR (400 MHz, DMSO-d6) δ 13.47 (s, 1H), 8.26 (dd, J = 5.1, 1.8 Hz, 1H), 8.16 (dd, J = 7.8, 1.8 Hz, 1H), 7.95 (d, J = 8.0 Hz, 1H), 7.59 (d, J = 8.0 Hz, 1H), 7.56 - 7.46 (m, 2H), 7.04 (s, 1H), 6.93 (dt, J = 10.7, 2.3 Hz, 1H), 6.84 (dd, J = 7.8, 5.1 Hz, 1H), 4.04 (q, J = 6.0 Hz, 2H), 3.85 (d, J = 6.5 Hz, 2H), 2.06 - 1.98 (m, 2H), 1.77 - 1.62 (m, 2H), 1.05 (d, J = 6.3 Hz, 6H), 0.99 (d, J = 6.7 Hz, 6H).
1160	567.60	1.74	
1161	495.00	1.38	1H NMR (400 MHz, MeOD) δ 7.94 (s, 1H), 7.36 (s, 2H), 7.01 (s, 5H), 4.38 (d, J = 100.8 Hz, 4H), 3.81 (s, 3H), 2.48 (s, 6H), 2.30 (s, 3H), 1.96 (s, 3H).
1162	540.30	3.03	1H NMR (400 MHz, DMSO-d6) δ 12.70 (s, 1H), 7.82 (d, J = 7.9 Hz, 1H), 7.66 (dd, J = 8.4, 7.3 Hz, 1H), 7.52 - 7.37 (m, 3H), 7.22 (d, J = 7.1 Hz, 1H), 6.92 (dt, J = 10.8, 2.3 Hz, 1H), 6.73 (d, J = 8.3 Hz, 1H), 6.48 (s, 1H), 4.74 (s, 2H), 3.85 (d, J = 6.5 Hz, 2H), 3.32 (dd, J = 6.9, 4.5 Hz, 4H), 2.21 (t, J = 5.6 Hz, 4H), 2.09 - 1.90 (m, 1H), 1.00 (d, J = 6.7 Hz, 6H).
1163	548.30	2.13	
1164	448.50	1.41	1H NMR (400 MHz, CDCl3) δ 14.47 (s, 1H), 8.43 (d, J = 7.5 Hz, 1H), 7.78 (s,

Cmpd No.	LC/MS M+1	LC/RT min	NMR
			1H), 7.58 - 7.41 (m, 2H), 6.94 (s, 1H), 3.87 (s, 2H), 3.70 (s, 3H), 3.42 (d, J = 6.3 Hz, 3H), 2.48 (s, 2H), 2.26 (d, J = 13.3 Hz, 2H), 1.36 (s, 9H).
1165	529.20	2.31	
1166	556.30	2.49	1H NMR (400 MHz, DMSO-d6) δ 7.64 (d, J = 7.8 Hz, 1H), 7.53 - 7.43 (m, 2H), 7.37 (dt, J = 10.1, 2.0 Hz, 1H), 7.15 (d, J = 7.8 Hz, 1H), 6.99 (d, J = 7.3 Hz, 1H), 6.80 (dt, J = 10.8, 2.3 Hz, 1H), 6.50 (d, J = 8.3 Hz, 1H), 6.19 (s, 2H), 4.01 - 3.86 (m, 1H), 3.82 (d, J = 6.6 Hz, 2H), 2.10 - 1.98 (m, 1H), 1.94 - 1.77 (m, 1H), 1.73 - 1.62 (m, 2H), 1.60 (s, 3H), 1.49 (s, 3H), 1.43 (dd, J = 18.5, 9.9 Hz, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.77 (d, J = 6.0 Hz, 3H).
1167	455.40	2.18	1H NMR (400 MHz, DMSO-d6) δ 7.68 (dd, J = 8.2, 1.6 Hz, 1H), 7.34 (ddd, J = 8.5, 7.0, 1.6 Hz, 1H), 6.95 (s, 1H), 6.83 (d, J = 8.3 Hz, 1H), 6.75 (s, 2H), 6.67 (t, J = 7.6 Hz, 1H), 5.41 (s, 2H), 2.17 (s, 3H), 2.09 (s, 6H), 1.15 (s, 9H).
1168	562.04	3.03	1H NMR (400 MHz, DMSO-d6) δ 9.10 (s, 1H), 7.83 - 7.80 (d, 1H), 7.52 - 7.48 (m, 3H), 6.95 - 6.91 (d, 1H), 3.86 - 3.83 (d, 2H), 3.26 (m, 4H), 2.10 - 2.00 (m, 1H), 1.36 (m, 4H), 1.01 - 0.98 (d, 6H), 0.96 (s, 6H)
1169	530.22	1.76	
1170	570.60	2.04	
1171	570.44	3.29	1H NMR (400 MHz, DMSO-d6) δ 12.58 (s, 1H), 7.77 - 7.74 (d, 1H), 7.61 - 7.59 (m, 2H), 7.49 (s, 2H), 7.44 - 7.40 (d, 1H), 7.33 - 7.28 (m, 2H), 6.93 - 6.89 (d, 1H), 3.85 - 3.82 (m, 5H), 2.47 - 2.37 (m, 2H), 2.11 - 2.03 (m, 2H), 1.89 - 1.82 (m, 1H), 1.61 (s, 3H), 1.57 (s, 3H), 1.41 (t, J = 12.52, 1H), 1.01 - 0.98 (d, 6H), 0.71 - 0.68 (d, 3H)
1172	451.30	1.74	
1173	519.20	1.97	
1174	544.23	1.92	
1175	425.30	1.82	
1176	530.30	1.99	1H NMR (400 MHz, DMSO) δ 13.76 (s,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 12.59 (s, 1H), 7.97 (s, 1H), 7.69 (d, J = 8.0 Hz, 1H), 7.48 (s, 1H), 7.40 (d, J = 9.4 Hz, 1H), 7.26 (d, J = 8.0 Hz, 1H), 6.93 - 6.84 (m, 2H), 3.83 (d, J = 6.6 Hz, 2H), 3.17 (s, 0H), 2.58 (d, J = 9.7 Hz, 2H), 2.21 (s, 1H), 2.05 (dt, J = 13.3, 6.7 Hz, 1H), 1.88 (dd, J = 11.7, 5.6 Hz, 1H), 1.60 (d, J = 12.8 Hz, 6H), 1.44 (t, J = 12.1 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.87 (d, J = 6.3 Hz, 3H).
1177	535.32	2.85	
1178	537.43	3.07	
1179	534.20	1.99	
1180	540.40	1.98	1H NMR (400 MHz, DMSO-d6) δ 12.44 (s, 1H), 7.72 (d, J = 7.9 Hz, 1H), 7.65 (dd, J = 8.4, 7.3 Hz, 1H), 7.47 - 7.40 (m, 2H), 7.23 (dd, J = 16.7, 7.6 Hz, 2H), 6.89 (dt, J = 10.7, 2.3 Hz, 1H), 6.72 (d, J = 8.3 Hz, 1H), 6.49 (s, 2H), 4.54 (s, 1H), 3.84 (d, J = 6.5 Hz, 2H), 2.99 (d, J = 9.9 Hz, 1H), 2.44 (s, 1H), 2.33 (d, J = 9.5 Hz, 1H), 2.10 - 1.98 (m, 1H), 1.69 - 1.48 (m, 4H), 1.44 - 1.37 (m, 1H), 1.33 - 1.22 (m, 1H), 0.99 (d, J = 6.7 Hz, 6H).
1181	553.50	1.77	
1182	511.20	1.47	
1183	519.12	2.47	
1184	555.43	2.90	1H NMR (400 MHz, DMSO-d6) δ 12.41 (s, 1H), 7.71 - 7.68 (d, 1H), 7.49 (s, 1H), 7.43 - 7.39 (d, 1H), 7.29 - 7.24 (m, 3H), 7.12 - 7.09 (d, 1H), 6.93 - 9.85 (m, 2H), 3.85 - 3.82 (d, 2H), 2.14 - 2.00 (m, 1H), 1.90 - 1.83 (m, 1H), 1.61 (s, 3H), 1.57 (s, 3H), 1.41 (t, J = 12.42, 1H), 1.00 - 0.97 (d, 6H), 0.78 - 0.75 (d, 3H)
1185	534.12	2.95	1H NMR (400 MHz, DMSO-d6) δ 8.49 (s, 1H), 8.12 - 8.09 (d, 1H), 8.04 - 8.01 (d, 1H), 7.64 - 7.53 (m, 2H), 7.15 - 7.12 (d, 1H), 6.94 (s, 2H), 6.83 - 6.80 (d, 1H), 6.58 - 6.56 (d, 1H), 6.32 (s, 2H), 3.30 (q, J = 7.25, 2H), 2.28 (s, 3H), 2.02 (s, 6H), 1.30 (t, J = 7.02, 3H)
1186	465.50	3.00	

Cmpd No.	LC/MS M+1	LC/RT min	NMR
1187	580.20	2.21	
1188	548.20	2.08	
1189	556.35	2.01	
1190	436.30	1.73	
1191	498.30	1.56	
1192	539.40	2.42	1H NMR (400 MHz, DMSO-d6) δ 8.20 (d, J = 5.5 Hz, 1H), 7.62 (s, 1H), 7.57 - 7.45 (m, 2H), 7.38 - 7.25 (m, 1H), 7.18 (s, 1H), 7.01 (s, 1H), 6.51 (s, 1H), 6.23 (s, 2H), 4.06 (d, J = 6.6 Hz, 2H), 3.00 - 2.86 (m, 2H), 2.19 - 2.00 (m, 2H), 1.83 (dd, J = 11.7, 5.6 Hz, 1H), 1.62 (s, 3H), 1.56 (s, 3H), 1.37 (t, J = 11.9 Hz, 1H), 0.97 (d, J = 6.7 Hz, 6H), 0.81 (d, J = 6.3 Hz, 3H).
1193	542.40	2.00	
1194	556.19	2.05	
1195	459.36	1.54	1H NMR (400 MHz, DMSO) δ 8.20 (dd, J = 7.2, 2.2 Hz, 1H), 7.76 (dd, J = 6.3, 2.0 Hz, 1H), 7.74 (s, 1H), 6.41 (dd, J = 7.2, 6.4 Hz, 1H), 3.44 - 3.33 (m, J = 12.4 Hz, 2H), 3.05 - 2.91 (m, J = 12.1, 9.9 Hz, 2H), 2.76 - 2.64 (m, J = 5.6 Hz, 2H), 1.79 - 1.66 (m, J = 12.8 Hz, 6H), 1.66 - 1.54 (m, 1H), 1.54 - 1.39 (m, 2H), 1.25 (s, 6H), 0.95 (d, J = 6.3 Hz, 3H).
1196	496.50	1.44	1H NMR (400 MHz, DMSO-d6) δ 8.28 (d, J = 8.2 Hz, 1H), 8.02 (d, J = 8.1 Hz, 2H), 7.67 (dd, J = 8.4, 7.3 Hz, 1H), 7.31 (d, J = 7.9 Hz, 2H), 7.20 (d, J = 7.2 Hz, 1H), 7.17 (d, J = 8.3 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 3.62 (dt, J = 11.7, 4.6 Hz, 2H), 3.54 (s, 2H), 3.48 (ddd, J = 11.9, 9.2, 3.0 Hz, 3H), 2.36 (s, 3H), 1.42 (ddd, J = 13.4, 9.2, 4.3 Hz, 2H), 1.21 (dt, J = 11.3, 2.9 Hz, 2H), 0.94 (s, 3H).
1197	558.30	1.82	1H NMR (400 MHz, DMSO-d6) δ 12.97 (s, 1H), 7.83 (d, J = 2.0 Hz, 1H), 7.80 (d, J = 7.9 Hz, 1H), 7.70 - 7.60 (m, 2H), 7.52 (d, J = 8.3 Hz, 1H), 7.38 (d, J = 8.0 Hz, 1H), 7.23 (d, J = 7.3 Hz, 1H), 6.72 (d, J = 8.3 Hz, 1H), 6.48 (s, 2H), 4.12 - 3.99 (m, 2H), 3.94 (d, J =

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			6.6 Hz, 2H), 2.16 - 2.06 (m, 1H), 2.04 - 1.90 (m, 2H), 1.73 - 1.58 (m, 2H), 1.10 (d, J = 6.2 Hz, 6H), 1.03 (d, J = 6.7 Hz, 7H).
1198	511.30	1.55	1H NMR (400 MHz, DMSO-d6) δ 8.83 (d, J = 2.4 Hz, 1H), 8.30 (dd, J = 8.8, 2.5 Hz, 1H), 7.75 (d, J = 7.9 Hz, 1H), 7.65 (dd, J = 8.4, 7.3 Hz, 1H), 7.23 (dd, J = 7.6, 3.3 Hz, 2H), 6.85 (d, J = 8.7 Hz, 1H), 6.73 (d, J = 8.4 Hz, 1H), 5.30 (p, J = 6.2 Hz, 1H), 4.30 - 3.96 (m, 2H), 2.22 - 1.94 (m, 2H), 1.62 - 1.36 (m, 2H), 1.31 (d, J = 6.2 Hz, 6H), 0.97 (d, J = 6.2 Hz, 6H).
1199	496.50	1.94	1H NMR (400 MHz, DMSO) δ 12.76 - 12.41 (m, 1H), 11.80 (bs, 1H), 8.26 (dd, J = 7.3, 2.1 Hz, 1H), 7.86 - 7.72 (m, 2H), 6.89 (s, 2H), 6.49 - 6.38 (m, J = 6.8 Hz, 1H), 2.78 - 2.60 (m, 3H), 2.33 - 2.13 (m, 5H), 1.97 (s, 6H), 1.56 - 1.31 (m, 2H), 0.89 (s, 6H).
1200	462.60	1.55	
1201	596.34	2.23	
1202	548.20	2.13	
1203	527.50	1.40	1H NMR (400 MHz, DMSO-d6) δ 8.61 (dd, J = 4.4, 1.5 Hz, 0H), 8.35 (dd, J = 8.3, 1.5 Hz, 0H), 8.05 (d, J = 7.7 Hz, 0H), 8.02 (d, J = 8.8 Hz, 2H), 7.47 (d, J = 7.8 Hz, 1H), 7.37 (dd, J = 8.3, 4.4 Hz, 0H), 7.06 (d, J = 7.8 Hz, 1H), 7.03 - 6.96 (m, 2H), 6.48 (t, J = 6.2 Hz, 1H), 4.08 (q, J = 6.9 Hz, 2H), 3.66 (dt, J = 11.6, 4.5 Hz, 2H), 3.55 - 3.50 (m, 2H), 3.49 (d, J = 6.2 Hz, 2H), 1.51 (ddd, J = 13.6, 9.3, 4.2 Hz, 2H), 1.35 (t, J = 6.9 Hz, 3H), 1.29 - 1.22 (m, 2H), 1.02 (s, 3H). Rotamer ratio about 5:1
1204	519.12	2.56	
1205	612.32	2.29	
1206	539.23	1.89	
1207	533.20	1.91	
1208	470.28	2.20	
1209	573.30	1.97	1H NMR (400 MHz, CDCl3) δ 14.58 (s, 1H), 8.87 (s, 1H), 8.57 (s, 1H), 8.23 (s, 1H), 7.80 (s, 2H), 7.38 (dd, J = 64.6,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			37.5 Hz, 7H), 6.89 (d, J = 40.0 Hz, 2H), 5.39 (s, 1H), 4.09 (s, 2H), 3.63 (s, 3H), 2.56 (s, 1H), 2.26 (s, 2H), 1.88 (s, 1H), 1.39 (s, 7H).
1210	556.40	2.99	1H NMR (400 MHz, Chloroform-d) δ 14.73 (s, 1H), 8.49 (d, J = 8.1 Hz, 1H), 7.75 - 7.67 (m, 2H), 7.63 (d, J = 7.3 Hz, 1H), 7.35 (dd, J = 9.0, 2.1 Hz, 2H), 6.76 - 6.68 (m, 2H), 3.79 (d, J = 6.5 Hz, 2H), 3.50 (s, 2H), 2.88 (t, J = 11.7 Hz, 2H), 2.34 (s, 2H), 2.13 (dq, J = 13.4, 6.7 Hz, 1H), 2.00 (t, J = 6.7 Hz, 1H), 1.05 (dd, J = 6.7, 4.7 Hz, 12H), 0.89 (t, J = 12.3 Hz, 1H).
1211	453.30	1.97	
1212	534.20	1.87	
1213	525.50	2.04	1H NMR (400 MHz, DMSO-d6) δ 13.80 (s, 1H), 12.50 (s, 1H), 8.05 (d, J = 7.8 Hz, 1H), 7.98 (s, 1H), 7.81 (d, J = 7.9 Hz, 1H), 7.27 - 7.21 (m, 1H), 7.19 - 7.06 (m, 4H), 6.87 (d, J = 2.3 Hz, 1H), 6.81 (dt, J = 10.8, 2.3 Hz, 1H), 4.46 (h, J = 6.0 Hz, 1H), 2.32 (s, 3H), 2.03 (s, 3H), 1.22 (d, J = 6.0 Hz, 6H).
1214	541.33	1.59	
1215	482.30	1.24	
1216	555.29	2.40	
1217	446.30	1.77	
1218	542.40	2.93	
1219	403.00	1.70	1H NMR (400 MHz, MeOD) δ 8.11 (s, 1H), 7.88 (s, 1H), 7.60 (s, 1H), 7.45 (d, J = 7.0 Hz, 1H), 7.04 (d, J = 8.6 Hz, 1H), 3.47 (s, 1H), 1.91 (s, 4H), 1.77 (s, 2H), 1.64 (s, 2H), 1.43 (s, 9H).
1220	493.20	0.72	
1221	523.10	2.01	
1222	556.40	1.85	
1223	496.00	2.23	1H NMR (400 MHz, DMSO) δ 12.42 (s, 1H), 8.32 (s, 1H), 7.90 (s, 1H), 7.72 (s, 1H), 7.10 (s, 2H), 6.89 (s, 2H), 2.69 (s, 2H), 2.25 (s, 3H), 1.95 (s, 6H), 1.65 (d, J = 36.4 Hz, 4H), 0.92 (s, 6H).
1224	525.36	1.23	
1225	549.43	2.95	
1226	468.00	1.94	1H NMR (400 MHz, DMSO) δ 12.57 (s,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 11.69 (s, 1H), 8.26 (d, J = 7.2 Hz, 1H), 8.04 (d, J = 7.9 Hz, 1H), 7.83 (s, 1H), 7.13 (d, J = 7.9 Hz, 1H), 6.91 (s, 2H), 6.46 (s, 1H), 2.26 (s, 3H), 1.96 (s, 6H), 1.32 (s, 3H), 0.68 (d, J = 12.5 Hz, 4H).
1227	555.26	2.37	
1228	518.22	1.99	
1229	515.10	2.25	
1230	551.20	1.82	
1231	450.30	1.78	
1232	455.30	2.04	
1233	503.00	1.88	1H NMR (400 MHz, DMSO) δ 12.36 (s, 1H), 8.04 (s, 1H), 7.86 (s, 1H), 7.61 (s, 1H), 7.28 (s, 4H), 7.20 (s, 2H), 6.87 (s, 2H), 6.69 (d, J = 8.7 Hz, 1H), 6.48 (s, 2H), 3.93 (s, 2H), 2.23 (s, 3H), 1.94 (s, 6H).
1234	532.20	1.84	
1235	493.00	1.74	1H NMR (400 MHz, MeOD) δ 8.32 (s, 1H), 8.06 (s, 1H), 7.12 (dd, J = 130.1, 74.1 Hz, 7H), 4.30 (d, J = 79.9 Hz, 2H), 3.58 (s, 3H), 2.43 (s, 6H), 2.26 (s, 3H), 1.98 (s, 3H), 1.56 (s, 3H).
1236	457.30	1.26	1H NMR (400 MHz, DMSO-d6) δ 13.17 (s, 1H), 12.90 (s, 1H), 7.86 (d, J = 7.3 Hz, 1H), 7.79 (t, J = 7.7 Hz, 1H), 7.76 - 7.35 (m, 3H), 6.79 (d, J = 8.0 Hz, 1H), 6.51 (s, 1H), 4.29 (s, 3H), 3.96 (d, J = 1.2 Hz, 2H), 1.89 (s, 2H), 1.61 (s, 2H), 1.13 (d, J = 6.1 Hz, 6H).
1237	466.00	1.20	1H NMR (400 MHz, DMSO) δ 12.29 (s, 1H), 10.30 (s, 1H), 8.13 (s, 1H), 7.68 (s, 1H), 7.38 (s, 1H), 7.24 (s, 3H), 6.90 (s, 2H), 6.34 (s, 1H), 4.12 (s, 2H), 3.96 (s, 2H), 3.17 (s, 1H), 2.41 (s, 6H), 2.24 (s, 3H), 1.73 (s, 3H).
1238	543.22	2.35	
1239	482.00	2.13	1H NMR (400 MHz, DMSO) δ 12.54 (s, 1H), 8.31 (s, 1H), 7.89 (s, 1H), 7.68 (s, 1H), 7.04 (d, J = 30.5 Hz, 5H), 2.67 (s, 3H), 2.29 (s, 3H), 1.99 (s, 3H), 1.66 (d, J = 20.8 Hz, 4H), 0.98 (s, 6H).
1240	525.40	1.38	1H NMR (400 MHz, DMSO-d6) δ 8.82 (s, 1H), 8.33 - 8.19 (m, 2H), 8.13 (d, J

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			= 8.1 Hz, 1H), 7.79 (d, J = 8.1 Hz, 1H), 7.25 (d, J = 8.0 Hz, 1H), 6.93 - 6.79 (m, 2H), 5.34 - 5.19 (m, 1H), 2.71 - 2.58 (m, 1H), 2.58 - 2.53 (m, 1H, obscured by solvent peak), 2.26 - 2.12 (m, 1H), 1.88 (dd, J = 12.1, 5.8 Hz, 1H), 1.53 (d, J = 4.5 Hz, 6H), 1.41 (t, J = 12.0 Hz, 1H), 1.29 (d, J = 6.1 Hz, 6H), 0.77 (d, J = 6.2 Hz, 3H).
1241	556.30	2.50	1H NMR (400 MHz, DMSO-d6) δ 7.64 (d, J = 7.6 Hz, 1H), 7.48 (d, J = 6.9 Hz, 2H), 7.41 - 7.30 (m, 1H), 7.15 (d, J = 7.9 Hz, 1H), 6.99 (d, J = 7.4 Hz, 1H), 6.85 - 6.75 (m, 1H), 6.50 (d, J = 8.4 Hz, 1H), 6.20 (s, 2H), 3.99 - 3.90 (m, 1H), 3.82 (d, J = 6.7 Hz, 2H), 2.05 (tt, J = 11.5, 5.9 Hz, 1H), 1.93 - 1.79 (m, 1H), 1.74 - 1.62 (m, 2H), 1.63 - 1.56 (m, 3H), 1.49 (s, 3H), 1.44 (t, J = 8.3 Hz, 1H), 0.98 (d, J = 6.8 Hz, 6H), 0.76 (d, J = 6.1 Hz, 3H).
1242	498.30	1.77	1H NMR (400 MHz, DMSO-d6) δ 7.58 (d, J = 7.6 Hz, 1H), 7.48 - 7.42 (m, 2H), 6.99 (d, J = 7.6 Hz, 1H), 6.67 (d, J = 7.6 Hz, 1H), 6.47 (d, J = 8.3 Hz, 1H), 6.40 (d, J = 1.8 Hz, 1H), 6.15 (s, 2H), 5.30 (p, J = 6.6 Hz, 1H), 3.08 - 2.92 (m, 2H), 2.18 - 2.03 (m, 1H), 1.80 (dd, J = 11.7, 5.6 Hz, 1H), 1.49 (d, J = 8.4 Hz, 6H), 1.38 (d, J = 6.4 Hz, 3H), 1.36 - 1.33 (m, 1H), 1.31 (d, J = 6.6 Hz, 3H), 0.82 (d, J = 6.3 Hz, 3H).
1243	455.23	2.02	
1244	525.42	2.95	
1245	532.20	1.90	
1246	455.00	2.02	1H NMR (400 MHz, DMSO) δ 13.78 (s, 1H), 12.23 (s, 1H), 7.96 (s, 1H), 7.65 (s, 1H), 7.08 (s, 1H), 7.01 (s, 2H), 6.83 (s, 1H), 2.67 (s, 2H), 2.29 (s, 3H), 2.00 (s, 3H), 1.66 (d, J = 22.4 Hz, 4H), 0.98 (s, 6H).
1247	525.30	1.61	1H NMR (400 MHz, DMSO) δ 12.43 (s, 1H), 8.84 (d, J = 2.1 Hz, 1H), 8.28 (dd, J = 8.7, 2.5 Hz, 1H), 7.76 (d, J = 8.0 Hz, 1H), 7.63 (dd, J = 8.4, 7.3 Hz, 1H),

Cmpd No	LC/MS M+1	LC/RT min	NMR
			7.19 (dd, J = 10.1, 7.4 Hz, 2H), 6.86 (d, J = 8.7 Hz, 1H), 6.68 (d, J = 8.4 Hz, 1H), 5.38 - 5.27 (m, 1H), 2.59 (d, J = 9.1 Hz, 2H), 2.22 (s, 1H), 1.87 (dd, J = 11.9, 5.6 Hz, 1H), 1.59 (d, J = 1.7 Hz, 6H), 1.42 (t, J = 12.1 Hz, 1H), 1.31 (d, J = 6.2 Hz, 6H), 0.88 (d, J = 6.3 Hz, 3H).
1248	551.40	1.81	
1249	535.50	2.04	1H NMR (400 MHz, DMSO-d6) δ 8.33 (dd, J = 3.3, 2.2 Hz, 2H), 8.11 (d, J = 7.9 Hz, 1H), 7.96 (dd, J = 2.4, 1.1 Hz, 1H), 7.91 (d, J = 2.2 Hz, 1H), 7.71 (d, J = 7.9 Hz, 1H), 7.20 - 7.05 (m, 2H), 6.96 (s, 2H), 3.88 (s, 3H), 2.28 (s, 3H), 2.12 (s, 3H), 2.01 (s, 6H).
1250	455.30	2.04	
1251	535.32	2.59	
1252	512.30	2.05	1H NMR (400 MHz, DMSO-d6) δ 8.92 (s, 1H), 8.27 (d, J = 5.7 Hz, 1H), 7.67 (bs, 1H), 7.45 (bs, 2H), 7.04 (bs, 1H), 6.50 (bs, 1H), 6.16 (bs, 1H), 4.48 (q, J = 7.0 Hz, 2H), 2.99 (bs, 1H), 2.15 (bs, 1H), 1.85 (dd, J = 10.7, 5.0 Hz, 1H), 1.64 (s, 3H), 1.59 (s, 3H), 1.40 (t, J = 7.0 Hz, 4H), 0.85 (d, J = 6.2 Hz, 3H).
1253	517.00	1.95	1H NMR (400 MHz, DMSO) δ 12.34 (s, 1H), 7.95 (d, J = 2.3 Hz, 1H), 7.87 (d, J = 2.3 Hz, 1H), 7.64 (s, 1H), 7.23 (s, 6H), 6.87 (s, 2H), 6.71 (d, J = 8.4 Hz, 1H), 6.54 (s, 2H), 2.86 (s, 4H), 2.23 (s, 3H), 1.94 (s, 6H).
1254	572.40	1.94	1H NMR (400 MHz, DMSO-d6) δ 12.24 (s, 1H), 7.64 (t, J = 7.8 Hz, 1H), 7.54 - 7.50 (m, 1H), 7.50 - 7.44 (m, 1H), 7.29 - 7.19 (m, 1H), 6.99 (s, 1H), 6.90 - 6.84 (m, 1H), 6.71 (d, J = 8.1 Hz, 1H), 6.51 (s, 2H), 4.10 (s, 2H), 3.88 (s, 3H), 3.82 (d, J = 6.6 Hz, 2H), 2.10 - 2.00 (m, 1H), 1.97 - 1.85 (m, 2H), 1.69 - 1.55 (m, 2H), 1.08 (d, J = 6.2 Hz, 6H), 0.99 (d, J = 6.7 Hz, 6H).
1255	551.53	3.15	
1256	452.40	1.77	
1257	578.15	2.02	
1258	556.40	2.20	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1259	524.60	1.51	
1260	590.33	2.18	
1261	494.50	1.86	
1262	509.30	1.51	1H NMR (400 MHz, DMSO-d6) δ 8.54 (d, J = 5.2 Hz, 1H), 7.97 - 7.82 (m, 1H), 7.74 (dd, J = 5.2, 1.7 Hz, 1H), 7.64 (s, 1H), 7.49 (t, J = 7.6 Hz, 1H), 7.21 (d, J = 7.7 Hz, 1H), 7.03 (s, 1H), 6.57 - 6.43 (m, 1H), 6.24 (s, 2H), 3.06 (h, J = 6.9 Hz, 1H), 2.91 (d, J = 11.8 Hz, 2H), 2.16 - 2.09 (m, 1H), 1.84 (dd, J = 11.8, 5.7 Hz, 1H), 1.64 (s, 3H), 1.58 (s, 3H), 1.38 (t, J = 12.0 Hz, 1H), 1.28 (d, J = 6.8 Hz, 6H), 0.82 (d, J = 6.3 Hz, 3H).
1263	557.22	2.39	
1264	568.90	2.98	
1265	425.30	1.76	
1266	558.30	2.04	
1267	558.30	2.75	
1268	563.30	2.00	
1269	570.30	1.86	1H NMR (400 MHz, DMSO-d6) δ 12.46 (s, 1H), 7.68 - 7.59 (m, 2H), 7.21 (d, J = 7.3 Hz, 1H), 6.96 - 6.82 (m, 3H), 6.68 (d, J = 8.3 Hz, 1H), 6.51 (s, 2H), 3.80 (d, J = 6.5 Hz, 2H), 2.71 - 2.53 (m, 2H), 2.27-2.15 (obscured m, 1H), 2.22 (s, 3H), 2.09 - 1.97 (m, 1H), 1.82 (dd, J = 11.9, 5.7 Hz, 1H), 1.47 (d, J = 11.2 Hz, 6H), 1.38 (t, J = 12.0 Hz, 1H), 0.97 (d, J = 6.7 Hz, 6H), 0.87 (d, J = 6.2 Hz, 3H).
1270	539.37	1.68	1H NMR (400 MHz, Chloroform-d) δ 14.94 (s, 1H), 8.88 (d, J = 2.2 Hz, 1H), 8.82 (d, J = 8.3 Hz, 1H), 8.34 (d, J = 8.9 Hz, 1H), 7.95 (d, J = 8.4 Hz, 1H), 7.76 (dd, J = 8.9, 7.0 Hz, 1H), 7.42 (dd, J = 18.0, 8.0 Hz, 2H), 6.95 (d, J = 8.8 Hz, 1H), 4.27 (s, 2H), 4.10 (s, 2H), 2.55 - 2.47 (m, 2H), 2.38 - 2.32 (m, 2H), 1.44 (d, 6H), 1.33 - 1.22 (m, 2H), 1.07 (d, J = 5.4 Hz, 9H).
1271	497.26	1.52	
1272	504.50	1.47	
1273	464.50	1.25	
1274	565.20	1.98	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1275	560.24	2.67	
1276	543.43	2.75	
1277	538.20	1.94	
1278	490.00	1.57	1H NMR (400 MHz, DMSO) δ 12.54 (s, 1H), 8.67 (s, 1H), 8.21 (d, J = 7.8 Hz, 1H), 8.12 (d, J = 7.8 Hz, 1H), 7.84 (s, 1H), 7.65 (s, 2H), 7.43 (s, 1H), 7.23 (d, J = 7.2 Hz, 1H), 6.97 (s, 2H), 6.72 (d, J = 8.4 Hz, 1H), 6.58 (s, 1H), 2.29 (s, 3H), 2.04 (s, 6H).
1279	556.40	2.92	
1280	481.50	2.07	1H NMR (400 MHz, DMSO-d6) δ 7.95 (d, J = 7.7 Hz, 1H), 7.63 (t, J = 7.8 Hz, 1H), 7.20 (d, J = 7.3 Hz, 1H), 7.00 (d, J = 7.7 Hz, 1H), 6.89 (s, 2H), 6.69 (d, J = 8.4 Hz, 1H), 6.45 (s, 2H), 3.03 - 2.93 (m, 1H), 2.24 (s, 3H), 1.96 (s, 6H), 1.83 - 1.66 (m, 2H), 1.52 - 1.33 (m, 6H).
1281	584.20	2.29	
1282	528.27	1.94	
1283	525.30	1.70	
1284	554.20	1.94	
1285	502.30	157.00	
1286	542.16	1.90	
1287	524.30	1.47	
1288	499.35	1.42	
1289	556.40	2.76	1H NMR (400 MHz, DMSO-d6) δ 12.47 (s, 1H), 7.78 (d, J = 8.0 Hz, 1H), 7.69 - 7.57 (m, 1H), 7.56 - 7.46 (m, 1H), 7.41 (dt, J = 10.1, 1.8 Hz, 1H), 7.26 (d, J = 8.0 Hz, 1H), 7.21 (d, J = 7.2 Hz, 1H), 6.89 (dt, J = 10.7, 2.3 Hz, 1H), 6.69 (d, J = 8.3 Hz, 1H), 3.83 (d, J = 6.7 Hz, 2H), 2.61 (dq, J = 9.7, 4.9 Hz, 2H), 2.24 (d, J = 15.8 Hz, 1H), 2.06 (dq, J = 13.3, 6.7 Hz, 1H), 1.93 - 1.82 (m, 1H), 1.61 (s, 3H), 1.59 (s, 3H), 1.48 - 1.33 (m, 1H), 1.32 - 1.20 (m, 2H), 0.99 (d, J = 6.6 Hz, 6H), 0.88 (d, J = 6.2 Hz, 3H).
1290	504.20	1.70	
1291	420.40	1.77	
1292	425.00	1.60	1H NMR (400 MHz, DMSO) δ 13.87 (s, 1H), 12.65 (s, 1H), 8.00 (s, 1H), 7.54 (s, 1H), 7.34 (d, J = 8.0 Hz, 2H), 7.09

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(d, J = 8.0 Hz, 2H), 6.78 (s, 1H), 2.80 (s, 2H), 2.32 (s, 3H), 1.76 (s, 4H), 1.29 (s, 6H).
1293	545.13	1.54	1H NMR (400 MHz, DMSO-d6) δ 12.98 (s, 1H), 12.57 (s, 1H), 8.26 (dd, J = 7.3, 2.1 Hz, 1H), 7.85 (dd, J = 6.7, 2.1 Hz, 2H), 7.79 (d, J = 7.9 Hz, 1H), 7.66 (dd, J = 8.3, 1.9 Hz, 1H), 7.51 (d, J = 8.3 Hz, 1H), 7.34 (d, J = 7.9 Hz, 1H), 6.45 (t, J = 6.8 Hz, 1H), 4.17 - 4.05 (m, 4H), 2.02 - 1.90 (m, 2H), 1.87 - 1.74 (m, 2H), 1.74 - 1.63 (m, 2H), 1.14 (d, J = 6.2 Hz, 6H), 1.03 (t, J = 7.4 Hz, 3H).
1294	475.20	4.69	1H NMR (300 MHz, CDCl3) δ 1.07 (s, 9H), 2.05 (s, 6H), 2.32 (s, 3H), 5.34 (s, 2H), 6.89 (s, 2H), 7.09 (d, J=8.1 Hz, 1H), 7.68 (s, 1H), 8.39 (d, J=7.9 Hz, 1H), 10.50 (s, 1H).
1295	523.20	2.12	
1296	526.20	1.89	
1297	466.00	2.03	1H NMR (400 MHz, DMSO) δ 11.55 (s, 1H), 7.91 (d, J = 7.8 Hz, 1H), 7.62 (d, J = 8.6 Hz, 2H), 7.07 (d, J = 7.8 Hz, 1H), 6.89 (s, 2H), 6.61 (d, J = 8.6 Hz, 2H), 6.14 (s, 2H), 2.25 (s, 3H), 1.92 (s, 6H), 1.31 (s, 3H), 0.66 (d, J = 16.9 Hz, 4H).
1298	447.40	1.73	
1299	543.13	2.65	
1300	504.20	2.01	
1301	552.20	2.01	
1302	516.25	1.67	
1303	568.35	2.14	
1304	582.28	2.12	1H NMR (400 MHz, DMSO-d6) δ 12.68 (s, 1H), 7.88 (d, J = 7.9 Hz, 1H), 7.66 (dd, J = 8.4, 7.3 Hz, 1H), 7.59 - 7.36 (m, 3H), 7.22 (d, J = 7.2 Hz, 1H), 6.91 (dt, J = 10.7, 2.3 Hz, 1H), 6.72 (d, J = 8.5 Hz, 1H), 6.50 (s, 2H), 5.62 (qd, J = 8.1, 3.9 Hz, 1H), 3.93 - 3.73 (m, 2H), 2.98 (dt, J = 10.0, 6.8 Hz, 1H), 2.84 - 2.73 (m, 1H), 2.21 - 1.79 (m, 5H), 0.99 (d, J = 6.7 Hz, 6H).
1305	485.20	2.18	1H NMR (400 MHz, CDCl3) δ 9.71 (s, 1H), 8.11 - 7.78 (m, 2H), 7.44 (s, 2H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			6.76 (s, 2H), 5.45 (s, 2H), 3.85 (s, 3H), 2.21 (s, 3H), 2.14 (s, 6H), 1.20 (d, J = 8.8 Hz, 10H). 1H NMR (400 MHz, DMSO) ? 7.30 (dd, J = 9.0, 6.8 Hz, 2H), 7.18 (d, J = 7.5 Hz, 1H), 6.94 (d, J = 8.0 Hz, 1H), 6.73 (s, 2H), 5.14 (s, 2H), 4.09 (s, 3H), 3.67 (s, 3H), 2.15 (s, 3H), 2.07 (s, 6H), 1.19 (s, 9H).
1306	456.08	2.91	
1307	450.30	1.79	
1308	557.24	2.64	1H NMR (400 MHz, CDCl3) δ 8.78 - 8.75 (d, 1H), 7.90 - 7.87 (d, 1H), 7.49 - 7.43 (m, 1H), 7.30 (s, 2H), 6.96 - 6.93 (d, 1H), 6.77 - 6.67 (m, 2H), 4.05 (t, J = 9.23, 1H), 3.80 - 3.77 (d, 2H), 3.48 (t, J = 9.45, 1H), 2.9 6 - 2.86 (m, 1H), 2.39 - 2.30 (m, 1H), 2.21 - 2.01 (m, 2H), 1.65 (s, 3H), 1.36 - 1.34 (m, 6H), 1.07 - 1.04 (d, 6H)
1309	556.34	3.03	1H NMR (400 MHz, CDCl3) δ 8.47 - 8.44 (d, 1H), 7.75 (s, 1H), 7.67 - 7.62 (m, 2H), 7.41 - 7.34 (m, 3H), 7.11 - 7.08 (d, 1H), 6.74 - 6.70 (d, 1H), 3.79 - 3.76 (d, 2H), 3.27 (m, 4H), 2.18 - 2.08 (m, 1H), 1.79 (m, 4H), 1.1 3 (s, 6H), 1.07 - 1.04 (d, 6H)
1310	467.00	2.08	1H NMR (400 MHz, DMSO) δ 11.99 (s, 1H), 7.99 (d, J = 7.8 Hz, 1H), 7.63 (s, 1H), 7.20 (d, J = 7.4 Hz, 1H), 7.10 (d, J = 7.1 Hz, 1H), 6.89 (s, 2H), 6.69 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 2.25 (s, 3H), 1.94 (s, 6H), 1.32 (s, 3H), 0.67 (d, J = 17.0 Hz, 4H).
1311	484.00	1.99	1H NMR (400 MHz, DMSO) δ 12.34 (s, 1H), 11.45 (s, 1H), 8.14 (d, J = 7.9 Hz, 1H), 7.64 (d, J = 6.5 Hz, 1H), 7.17 (d, J = 7.9 Hz, 1H), 6.94 (s, 2H), 6.30 (d, J = 6.5 Hz, 1H), 2.60 (s, 3H), 2.27 (s, 3H), 2.00 (s, 6H), 1.04 (s, 9H).1H NMR (400 MHz, DMSO) ? 12.34 (s, 1H), 11.45 (s, 1H), 8.14 (d, J = 7.9 Hz, 1H), 7.64 (d, J = 6.5 Hz, 1H), 7.17 (d, J = 7.9 Hz, 1H), 6.94 (s, 2H), 6.30 (d, J = 6.5 Hz, 1H), 2.60 (s, 3H), 2.27 (s, 3H), 2.00 (s, 6H), 1.04 (s, 9H).

Cmpd No.	LC/MS M+1	LC/RT min	NMR
1312	507.50	1.66	
1313	568.40	2.73	
1314	469.26	2.08	
1315	578.16	2.04	
1316	511.20	1.24	1H NMR (400 MHz, DMSO-d6) δ 13.74 (s, 1H), 8.94 (d, J = 2.5 Hz, 1H), 8.39 (dt, J = 8.7, 2.2 Hz, 1H), 8.30 - 8.18 (m, 2H), 8.05 (d, J = 7.9 Hz, 1H), 7.64 (d, J = 8.0 Hz, 1H), 7.25 (s, 2H), 6.98 - 6.82 (m, 2H), 5.41 - 5.26 (m, 1H), 4.21 - 4.04 (m, 2H), 2.23 - 2.06 (m, 2H), 1.83 - 1.67 (m, 2H), 1.32 (d, J = 6.2 Hz, 6H), 1.08 (d, J = 6.4 Hz, 6H).
1317	521.30	1.88	
1318	441.19	1.93	
1319	604.30	3.24	
1320	533.22	3.07	
1321	488.50	2.74	
1322	497.15	1.62	
1323	579.45	3.40	1H NMR (400 MHz, DMSO-d6) δ 12.20 (s, 1H), 7.86 - 7.83 (d, 1H), 7.73 - 7.70 (d, 1H), 7.56 - 7.46 (m, 4H), 7.33 (t, J = 8.13, 1H), 7.21 - 7.11 (m, 2H), 6.94 - 6.90 (d, 1H), 3.85 - 3.82 (d, 2H), 3.13 (m, 4H), 2.06 - 1.9 8 (m, 1H), 1.15 (m, 4H), 1.00 - 0.97 (d, 6H), 0.81 (s, 6H)
1324	545.21	2.05	
1325	525.50	2.00	
1326	549.43	2.77	
1327	525.30	1.74	
1328	539.60	1.73	
1329	532.20	1.85	
1330	479.30	2.16	
1331	442.21	1.80	
1332	554.10	2.14	
1333	540.31	1.96	
1334	466.23	1.66	
1335	498.40	1.68	
1336	549.43	3.01	
1337	533.22	3.10	1H NMR (400 MHz, DMSO-d6) δ 8.10 - 8.07 (d, 1H), 7.71 - 7.52 (m, 4H), 7.18 - 7.15 (d, 1H), 7.93 - 7.90 (m, 4H), 6.60 - 6.57 (d, 1H), 6.36 (s, 2H), 4.04 (q, J = 6.92, 2H), 2.28 (s, 3H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			2.03 (s, 6H), 1.31 (t, J = 6.92, 3H)
1338	481.00	1.10	1H NMR (400 MHz, DMSO) δ 9.61 (s, 1H), 7.52 (s, 1H), 7.13 (s, 1H), 7.06 (d, J = 7.0 Hz, 1H), 6.90 (s, 2H), 6.86 (d, J = 7.5 Hz, 1H), 6.68 (d, J = 7.9 Hz, 1H), 6.56 (d, J = 8.3 Hz, 1H), 6.23 (s, 2H), 4.14 (d, J = 12.7 Hz, 1H), 4.01 (d, J = 12.7 Hz, 1H), 3.92 (s, 2H), 2.38 (s, 6H), 2.24 (s, 3H), 1.69 (s, 3H).
1339	607.21	2.04	
1340	557.00	1.91	1H NMR (400 MHz, DMSO) δ 13.27 (s, 1H), 8.18 (s, 1H), 8.03 (d, J = 7.7 Hz, 1H), 7.83 (dd, J = 12.8, 7.9 Hz, 2H), 7.54 (d, J = 8.0 Hz, 1H), 7.47 (d, J = 8.3 Hz, 2H), 6.94 (s, 1H), 4.58 (s, 2H), 3.84 (s, 2H), 3.64 (d, J = 12.8 Hz, 2H), 2.87 (t, J = 12.1 Hz, 2H), 2.11 - 1.98 (m, 3H), 1.57 (d, J = 10.1 Hz, 3H), 1.18 (t, J = 10.6 Hz, 2H), 1.00 (d, J = 6.7 Hz, 6H), 0.91 (d, J = 6.0 Hz, 3H).
1341	539.20	1.83	
1342	494.00	1.64	1H NMR (400 MHz, DMSO) δ 8.45 (d, J = 6.7 Hz, 1H), 8.26 (d, J = 9.4 Hz, 1H), 8.21 (d, J = 2.1 Hz, 1H), 8.19 (d, J = 2.1 Hz, 1H), 7.24 (s, 1H), 7.13 (s, 2H), 7.05 (s, 1H), 6.88 (s, 1H), 6.81 (s, 2H), 4.33 (d, J = 15.1 Hz, 1H), 3.90 (s, 3H), 3.73 (s, 2H), 3.58 (d, J = 13.6 Hz, 2H), 2.79 (s, 2H), 2.20 (s, 3H), 2.11 (s, 6H), 2.00 (s, 2H).
1343	494.40	1.60	
1344	505.31	2.55	
1345	538.60	1.23	
1346	509.00	1.44	1H NMR (400 MHz, DMSO) δ 9.91 (s, 1H), 7.73 (s, 1H), 7.31 (s, 1H), 7.15 (d, J = 7.1 Hz, 1H), 7.04 (d, J = 7.6 Hz, 1H), 6.92 (s, 3H), 6.82 (s, 1H), 4.15 (s, 3H), 4.03 (s, 4H), 2.38 (s, 6H), 2.25 (s, 3H), 1.78 (s, 3H), 1.25 (s, 3H).
1347	496.00	2.28	
1348	592.19	1.91	
1349	445.22	2.12	
1350	442.21	1.80	
1351	522.00	1.87	1H NMR (400 MHz, DMSO) δ 13.07 (s, 1H), 8.43 (s, 1H), 8.05 (d, J = 15.1 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			3H), 7.56 (s, 2H), 7.42 (d, J = 8.0 Hz, 2H), 7.05 (dd, J = 67.1, 10.4 Hz, 5H), 4.78 (s, 1H), 2.34 (s, 3H), 1.31 (d, J = 5.9 Hz, 6H).
1352	499.30	1.61	1H NMR (400 MHz, DMSO) δ 14.22 - 13.30 (m, 1H), 12.56 (s, 1H), 8.83 (d, J = 2.3 Hz, 1H), 8.28 (dd, J = 8.7, 2.5 Hz, 1H), 7.96 (d, J = 2.4 Hz, 1H), 7.68 (d, J = 8.0 Hz, 1H), 7.17 (d, J = 8.0 Hz, 1H), 6.86 (dd, J = 5.5, 3.1 Hz, 2H), 5.31 (dt, J = 12.3, 6.2 Hz, 1H), 2.58 (dd, J = 17.5, 7.2 Hz, 2H), 2.21 (s, 1H), 1.88 (dd, J = 11.8, 5.7 Hz, 1H), 1.59 (d, J = 8.8 Hz, 6H), 1.43 (t, J = 12.1 Hz, 1H), 1.31 (d, J = 6.2 Hz, 6H), 0.87 (d, J = 6.3 Hz, 3H).
1353	464.00	1.53	1H NMR (400 MHz, DMSO) δ 13.86 (s, 1H), 12.60 (s, 1H), 8.66 (s, 1H), 8.12 (d, J = 4.6 Hz, 2H), 7.99 (s, 1H), 7.85 (s, 1H), 7.63 (d, J = 7.9 Hz, 1H), 7.42 (s, 1H), 6.96 (s, 2H), 6.89 (s, 1H), 2.29 (s, 3H), 2.03 (s, 6H).
1354	497.21	1.50	
1355	424.30	1.84	
1356	540.00	3.27	1H NMR (400 MHz, DMSO) δ 12.36 (s, 1H), 8.13 (d, J = 7.8 Hz, 1H), 7.80 (d, J = 7.6 Hz, 1H), 7.63 (s, 1H), 7.28 - 7.19 (m, 2H), 7.17 (s, 1H), 6.95 (s, 2H), 6.84 (d, J = 10.8 Hz, 1H), 6.70 (d, J = 6.5 Hz, 1H), 6.51 (s, 2H), 3.65 (d, J = 6.5 Hz, 2H), 2.27 (s, 3H), 2.11 - 1.92 (m, 7H), 0.97 (d, J = 6.7 Hz, 6H).
1357	496.20	2.09	1H NMR (400 MHz, DMSO-d6) δ 12.56 (s, 1H), 11.86 (s, 1H), 8.26 (dd, J = 7.2, 2.1 Hz, 1H), 8.03 (d, J = 7.7 Hz, 1H), 7.83 (s, 1H), 7.04 (d, J = 7.9 Hz, 1H), 6.95 - 6.91 (m, 2H), 6.46 (dd, J = 7.3, 6.3 Hz, 1H), 2.47 - 2.36 (m, 1H), 2.26 (s, 3H), 1.98 (s, 6H), 1.78 - 1.50 (m, 6H), 1.21 (h, J = 10.9, 10.4 Hz, 4H).
1358	465.39	3.36	1H NMR (400 MHz, DMSO-d6) δ 11.0 (s, 1H), 7.98 - 7.95 (d, 1H), 7.75 - 7.72 (d, 1H), 7.56 (s, 1H), 7.24 (t, J = 7.8, 1H), 7.01 - 6.95 (m, 2H), 6.87 - 6.84

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(m, 2H), 6.68 (s, 1H), 5.36 (s, 2H), 1.94 (s, 6H), 1.14 (s, 9H)
1359	586.60	2.24	
1360	548.60	1.60	
1361	587.30	2.22	
1362	546.20	2.01	1H NMR (400 MHz, DMSO) δ 12.63 (s, 1H), 7.77 (d, J = 7.9 Hz, 1H), 7.69 - 7.57 (m, 1H), 7.52 - 7.40 (m, 2H), 7.35 (d, J = 7.9 Hz, 1H), 7.22 (d, J = 7.1 Hz, 1H), 6.90 (d, J = 10.8 Hz, 1H), 6.73 (d, J = 8.3 Hz, 1H), 6.52 (s, 2H), 5.32 (d, J = 54.8 Hz, 1H), 4.64 (t, J = 6.6 Hz, 1H), 3.84 (d, J = 6.6 Hz, 2H), 3.34 - 3.15 (m, 2H), 2.36 - 2.15 (m, 1H), 2.05 (dt, J = 13.2, 5.3 Hz, 1H), 1.99 - 1.83 (m, 1H), 1.28 (d, J = 6.4 Hz, 3H), 1.00 (d, J = 6.7 Hz, 6H).
1363	525.50	1.65	
1364	522.30	1.85	1H NMR (400 MHz, DMSO-d6) δ 12.25 (s, 1H), 7.62 (dd, J = 8.4, 7.3 Hz, 1H), 7.45 - 7.32 (m, 3H), 7.36 - 7.24 (m, 3H), 7.18 (dd, J = 7.3, 0.7 Hz, 1H), 6.67 (dd, J = 8.4, 0.8 Hz, 1H), 6.45 (s, 2H), 2.60 - 2.51 (m, 2H), 2.21 (ddt, J = 17.6, 11.9, 6.4 Hz, 1H), 1.86 (dd, J = 11.8, 5.6 Hz, 1H), 1.58 (d, J = 8.4 Hz, 6H), 1.46 - 1.35 (m, 1H), 1.13 (s, 9H), 0.90 - 0.79 (m, 3H).
1365	593.20	2.40	
1366	580.50	1.89	
1367	550.40	2.69	1H NMR (400 MHz, DMSO-d6) δ 10.55 (s, 1H), 7.26 (s, 1H), 7.20 (t, J = 7.9 Hz, 1H), 7.05 (q, J = 7.9 Hz, 2H), 6.85 (d, J = 8.2 Hz, 3H), 4.01 (dd, J = 15.5, 9.3 Hz, 3H), 3.79 (q, J = 14.0, 13.4 Hz, 4H), 2.45 (t, J = 7.7 Hz, 2H), 2.31 (s, 6H), 2.23 (s, 3H), 1.61 (s, 3H), 1.52 (p, J = 7.4 Hz, 2H), 1.24 (t, J = 6.9 Hz, 3H), 0.91 (t, J = 7.4 Hz, 3H).
1368	583.20	2.10	1H NMR (400 MHz, DMSO) δ 12.61 (s, 1H), 7.99 (d, J = 8.2 Hz, 1H), 7.89 (d, J = 8.2 Hz, 1H), 7.64 (dd, J = 8.3, 7.4 Hz, 1H), 7.58 - 7.49 (m, 2H), 7.21 (d, J = 6.9 Hz, 1H), 6.95 (dt, J = 10.8, 2.3 Hz, 1H), 6.79 - 6.68 (m, 1H), 6.54 (s, 2H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			5.90 (s, 1H), 3.86 (d, J = 6.6 Hz, 2H), 3.32 (s, 1H), 2.42 (s, 2H), 1.22 (s, 6H), 1.15 (s, 6H), 1.00 (d, J = 6.7 Hz, 6H).
1369	500.30	1.27	
1370	521.50	2.27	
1371	571.40	1.57	
1372	509.00	1.34	1H NMR (400 MHz, MeOD) δ 8.10 (s, 1H), 7.52 (s, 1H), 7.25 (s, 2H), 6.97 (s, 3H), 6.87 (d, J = 7.9 Hz, 1H), 6.61 (s, 1H), 4.40 (s, 1H), 4.17 (s, 2H), 4.04 (s, 2H), 2.44 (s, 6H), 2.29 (s, 3H), 1.85 (s, 2H), 1.32 (s, 3H), 1.15 (d, J = 6.1 Hz, 1H).
1373	472.50	1.50	
1374	557.08	2.48	
1375	527.20	1.99	1H NMR (400 MHz, Chloroform-d) δ 8.87 (d, J = 2.5 Hz, 1H), 8.54 (d, J = 8.2 Hz, 1H), 8.22 (dd, J = 8.7, 2.6 Hz, 1H), 7.74 (d, J = 8.3 Hz, 1H), 7.39 (d, J = 7.8 Hz, 1H), 6.78 (dd, J = 8.6, 4.5 Hz, 2H), 5.38 (hept, J = 6.2 Hz, 1H), 3.94 (q, J = 6.4 Hz, 2H), 2.34 (ddt, J = 9.8, 7.1, 3.7 Hz, 2H), 2.11 - 1.89 (m, 2H), 1.38 (d, J = 6.1 Hz, 6H), 1.34 - 1.22 (m, 6H).
1376	509.30	2.27	
1377	528.60	1.89	
1378	484.30	1.57	1H NMR (400 MHz, DMSO-d6) δ 12.31 (s, 1H), 7.61 (d, J = 7.6 Hz, 2H), 7.17 (d, J = 7.3 Hz, 1H), 6.66 (d, J = 8.3 Hz, 1H), 6.58 - 6.37 (m, 3H), 3.23 - 3.11 (m, 1H), 2.62 - 2.53 (m, 2H, obscured by solvent peak), 2.28 (s, 1H), 2.25 - 2.12 (m, 1H), 1.88 - 1.69 (m, 3H), 1.54 (dd, J = 7.9, 4.6 Hz, 7H), 1.47 - 1.03 (m, 7H), 0.87 (d, J = 6.2 Hz, 3H).
1379	456.20	1.38	
1380	594.50	1.84	
1381	570.45	1.21	
1382	541.40	0.80	
1383	552.50	1.18	
1384	466.30	1.45	
1385	556.30	2.20	
1386	430.30	1.14	
1387	595.30	2.32	1H NMR (400 MHz, DMSO) δ 12.41 (s,

Cmpd No.	LC/MS M+1	LC/RT min	NMR
			1H), 8.09 (s, 2H), 7.60 (s, 1H), 7.55 - 7.47 (m, 2H), 7.13 (d, J = 7.5 Hz, 1H), 6.97 (d, J = 10.3 Hz, 1H), 6.69 (d, J = 8.6 Hz, 1H), 6.54 (s, 2H), 6.20 (s, 1H), 3.86 (d, J = 6.5 Hz, 2H), 3.62 (s, 1H), 2.82 - 2.71 (m, 1H), 2.09 - 2.00 (m, 1H), 1.25 (d, J = 6.9 Hz, 6H), 1.16 (d, J = 6.9 Hz, 6H), 1.00 (d, J = 6.7 Hz, 6H).
1388	653.40	1.88	1H NMR (400 MHz, Methanol-d4) δ 7.81 (d, J = 13.5 Hz, 1H), 7.78 - 7.66 (m, 1H), 7.35 (d, J = 7.3 Hz, 1H), 6.88 (dd, J = 8.6, 5.6 Hz, 1H), 5.93 (dd, J = 12.2, 5.1 Hz, 1H), 4.45 (s, 1H), 4.39 (t, J = 6.3 Hz, 1H), 3.77 - 3.51 (m, 1H), 3.27 - 3.16 (m, 1H), 3.00 (dd, J = 20.5, 12.5 Hz, 1H), 2.67 (s, 1H), 2.28 (s, 1H), 2.17 (td, J = 13.3, 7.0 Hz, 1H), 2.04 (s, 2H), 1.92 (s, 1H), 1.85 - 1.69 (m, 1H), 1.52 (d, J = 5.2 Hz, 3H), 1.49 (s, 9H), 1.45 - 1.35 (m, 12H), 1.23 - 1.12 (m, 3H).
1389	584.40	1.34	
1390	481.30	2.02	
1391	590.60	2.05	
1392	499.00	1.23	1H NMR (400 MHz, MeOD) δ 7.91 (s, 1H), 7.43 (d, J = 6.1 Hz, 1H), 7.32 (d, J = 16.0 Hz, 2H), 7.21 - 7.09 (m, 3H), 7.05 (d, J = 8.9 Hz, 1H), 6.95 (d, J = 8.3 Hz, 1H), 4.52 (s, 4H), 4.09 (s, 2H), 2.55 (s, 3H), 1.92 (s, 3H), 1.33 (s, 3H).
1393	532.00	1.84	1H NMR (400 MHz, DMSO) δ 11.63 (s, 1H), 10.77 (s, 1H), 8.04 (s, 1H), 7.64 (s, 2H), 7.45 (d, J = 8.3 Hz, 1H), 7.15 (s, 1H), 6.83 (s, 3H), 6.67 (d, J = 7.5 Hz, 1H), 6.55 (s, 1H), 3.97 (d, J = 7.0 Hz, 3H), 3.73 (s, 1H), 3.57 (s, 3H), 2.23 (d, J = 15.3 Hz, 9H), 2.07 (s, 1H), 1.56 (s, 3H), 1.23 (s, 3H).
1394	526.00	1.67	1H NMR (400 MHz, DMSO) δ 10.45 (s, 1H), 7.34 (d, J = 8.5 Hz, 1H), 7.23 (s, 1H), 7.08 (s, 1H), 6.98 (s, 1H), 6.87 (d, J = 8.4 Hz, 3H), 6.81 (d, J = 7.7 Hz, 1H), 5.52 (s, 1H), 4.08 (d, J = 12.3 Hz, 1H), 4.01 (s, 2H), 3.83 (d, J = 19.7 Hz, 3H), 2.34 (s, 6H), 2.24 (s, 3H), 2.08 (s,

Cmpd No.	LC/MS M+1	LC/RT min	NMR
			1H), 1.64 (s, 3H), 1.25 (s, 3H).
1395	557.36	2.50	
1396	540.00	3.26	1H NMR (400 MHz, DMSO-d6) δ 8.09 - 7.95 (m, 2H), 7.79 - 7.70 (m, 2H), 7.70 - 7.61 (m, 2H), 7.48 (dd, J = 2.3, 1.4 Hz, 1H), 7.40 (ddd, J = 10.2, 2.4, 1.4 Hz, 1H), 7.27 (d, J = 8.0 Hz, 1H), 6.89 (dt, J = 10.7, 2.3 Hz, 1H), 3.92 - 3.67 (m, 3H), 2.43 (t, J = 10.3 Hz, 1H), 2.36 (dd, J = 10.1, 7.0 Hz, 1H), 2.08 (ddt, J = 32.6, 13.3, 6.3 Hz, 2H), 1.83 (dd, J = 11.9, 5.5 Hz, 1H), 1.57 (d, J = 17.5 Hz, 6H), 1.38 (t, J = 12.1 Hz, 1H), 1.04 (d, J = 6.1 Hz, 5H), 0.98 (d, J = 6.7 Hz, 6H), 0.66 (d, J = 6.3 Hz, 3H).
1397	538.30	1.25	
1398	524.30	1.17	
1399	538.30	1.25	
1400	558.34	2.73	1H NMR (400 MHz, DMSO-d6) δ 12.46 (s, 1H), 7.77 (d, J = 7.9 Hz, 1H), 7.62 (t, J = 7.8 Hz, 1H), 7.51 - 7.46 (m, 1H), 7.40 (dt, J = 9.9, 1.9 Hz, 1H), 7.22 (dd, J = 22.1, 7.6 Hz, 2H), 6.89 (dt, J = 10.9, 2.3 Hz, 1H), 6.67 (d, J = 8.4 Hz, 1H), 6.48 (s, 2H), 3.83 (d, J = 6.6 Hz, 2H), 2.26 - 2.13 (m, 1H), 2.05 (dt, J = 13.3, 6.7 Hz, 1H), 1.94 - 1.83 (m, 1H), 1.60 (d, J = 6.8 Hz, 6H), 1.42 (t, J = 12.1 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.88 (d, J = 6.3 Hz, 3H).
1401	430.00	1.49	1H NMR (400 MHz, DMSO) δ 12.40 (s, 1H), 7.69 - 7.49 (m, 2H), 7.24 - 7.07 (m, 1H), 6.82 - 6.61 (m, 2H), 4.84 (s, 2H), 3.46 (s, 4H), 2.42 (t, J = 5.8 Hz, 2H), 2.12 - 2.00 (m, 2H), 1.27 (s, 9H), 1.23 (dd, J = 6.7, 2.7 Hz, 2H).
1402	561.40	2.07	
1403	541.00	1.79	1H NMR (400 MHz, DMSO) δ 7.64 (d, J = 7.9 Hz, 1H), 7.42 (s, 1H), 7.26 (s, 1H), 7.16 (s, 1H), 7.07 (d, J = 7.6 Hz, 1H), 6.92 (s, 3H), 4.24 - 3.99 (m, 6H), 3.84 (s, 3H), 2.40 (s, 6H), 2.25 (s, 4H), 2.07 (s, 1H), 1.75 (s, 3H), 1.24 (s, 3H).
1404	661.40	1.84	1H NMR (400 MHz, Methanol-d4) δ 7.73 (s, 1H), 7.67 (t, J = 8.1 Hz, 1H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			7.43 - 7.27 (m, 6H), 6.77 (d, J = 8.5 Hz, 1H), 5.59 (s, 1H), 5.18 (s, 2H), 4.11 (s, 2H), 3.72 (s, 2H), 3.39 (q, J = 16.4, 13.3 Hz, 1H), 3.12 (t, J = 10.2 Hz, 1H), 2.57 (s, 1H), 2.40 (s, 2H), 2.09 (dd, J = 12.5, 6.8 Hz, 1H), 1.72 (t, J = 11.8 Hz, 1H), 1.53 (s, 3H), 1.44 (s, 3H), 1.39 (s, 9H), 1.11 (d, J = 6.5 Hz, 3H).
1405	568.39	1.26	1H NMR (400 MHz, Methanol-d4) δ 7.70 (dd, J = 8.5, 7.3 Hz, 1H), 7.63 (s, 1H), 7.36 (dd, J = 7.3, 0.8 Hz, 1H), 6.81 (dd, J = 8.6, 0.8 Hz, 1H), 5.93 (d, J = 1.7 Hz, 1H), 3.10 (t, J = 9.1 Hz, 1H), 2.95 (t, J = 10.3 Hz, 1H), 2.71 - 2.57 (m, 2H), 2.46 (dd, J = 11.1, 6.4 Hz, 1H), 2.36 (d, J = 2.1 Hz, 2H), 2.02 (dd, J = 12.2, 6.2 Hz, 1H), 1.62 (t, J = 12.0 Hz, 1H), 1.57 (s, 3H), 1.53 (s, 3H), 1.40 (s, 9H), 1.17 (d, J = 1.3 Hz, 6H), 1.04 (d, J = 6.5 Hz, 3H).
1406	529.20	2.32	1H NMR (400 MHz, DMSO) δ 7.90 - 7.84 (m, 1H), 7.54 (d, J = 7.7 Hz, 1H), 7.51 - 7.40 (m, 3H), 7.07 (dt, J = 7.3, 2.6 Hz, 1H), 6.85 (dt, J = 10.7, 2.2 Hz, 1H), 6.47 (d, J = 8.2 Hz, 1H), 6.13 (s, 2H), 5.07 (dq, J = 119.9, 7.4 Hz, 1H), 3.85 (d, J = 6.6 Hz, 2H), 2.46 - 2.18 (m, 2H), 2.14 - 2.01 (m, 3H), 2.01 - 1.76 (m, 2H), 1.36 - 1.21 (m, 1H), 1.16 (d, J = 6.7 Hz, 2H), 1.05 - 0.91 (m, 7H).
1407	542.50	1.65	1H NMR (400 MHz, DMSO-d6) δ 12.46 (s, 1H), 7.82 (broad s, 1H), 7.59 (br s, 1H), 7.16 (br s, 1H), 6.89 (br s, 1H), 6.64 (br s, 1H), 6.45 (br s, 2H), 5.80 (s, 1H), 3.89 (d, J = 6.6 Hz, 2H), 2.67 (br s, 2H), 2.20 (br s, 1H), 2.02 (dt, J = 13.3, 6.7 Hz, 1H), 1.86 (dd, J = 12.0, 5.6 Hz, 1H), 1.53 (s, 3H), 1.49 (s, 3H), 1.39 (t, J = 12.0 Hz, 1H), 0.95 (d, J = 6.7 Hz, 6H), 0.87 (d, J = 6.3 Hz, 3H).
1408	481.00	1.15	1H NMR (400 MHz, MeOD) δ 7.95 (s, 1H), 7.62 (s, 1H), 7.34 (s, 5H), 7.14 (d, J = 42.4 Hz, 2H), 6.96 (s, 1H), 4.65 (s, 2H), 4.37 (s, 2H), 4.08 (s, 2H), 2.53 (s, 3H), 2.07 (s, 3H), 1.90 (s, 3H), 1.35 (s, 3H).

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1409	556.50	2.19	
1410	547.50	2.33	
1411	512.50	1.78	1H NMR (400 MHz, Chloroform-d) δ 13.85 (br s, 1H), 8.35 (d, J = 8.5 Hz, 1H), 8.31 (d, J = 2.6 Hz, 1H), 7.73 (d, J = 8.5 Hz, 1H), 7.70 - 7.59 (m, 2H), 6.68 (d, J = 7.7 Hz, 1H), 6.28 (d, J = 2.7 Hz, 1H), 3.61 (bra s, 1H), 3.24 (br t, 1H), 2.67 (m, 1H), 2.58 (d, J = 7.1 Hz, 2H), 2.14 (dd, J = 12.4, 7.7 Hz, 1H), 1.97-2.04 (m, 1H), 1.78 (br t, J = 11.0 Hz, 1H), 1.42 (s, 3H), 1.29 - 1.19 (m, 6H), 0.98 (d, J = 6.6 Hz, 6H).
1412	481.40	1.77	1H NMR (400 MHz, DMSO-d6) δ 11.86 (s, 1H), 7.97 (d, J = 7.9 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.22 (d, J = 7.2 Hz, 1H), 7.13 (t, J = 1.8 Hz, 1H), 7.04 (d, J = 7.9 Hz, 1H), 6.90 (s, 2H), 6.70 (d, J = 8.3 Hz, 1H), 6.53 - 6.45 (m, 2H), 4.43 (t, J = 9.6 Hz, 2H), 2.69 - 2.62 (m, 2H), 2.25 (s, 3H), 2.00 (s, 6H).
1413	509.00	1.17	1H NMR (400 MHz, DMSO) δ 9.66 (s, 1H), 8.34 (s, 2H), 8.12 (d, J = 9.2 Hz, 1H), 7.29 (s, 1H), 7.04 - 6.86 (m, 4H), 4.14 (d, J = 84.4 Hz, 7H), 2.40 (s, 6H), 2.25 (s, 3H), 1.77 (s, 3H), 1.25 (s, 3H).
1414	590.60	1.63	
1415	551.00	1.46	1H NMR (400 MHz, MeOD) δ 8.17 (d, J = 8.3 Hz, 1H), 7.92 (s, 1H), 7.77 (d, J = 7.4 Hz, 1H), 7.34 (s, 1H), 7.12 (d, J = 7.4 Hz, 1H), 6.99 (s, 2H), 6.94 (d, J = 8.3 Hz, 1H), 4.49 (s, 2H), 4.35 (s, 2H), 4.06 (s, 2H), 2.42 (s, 6H), 2.29 (s, 3H), 2.11 (s, 3H), 2.01 (s, 3H), 1.32 (s, 3H).
1416	512.30	3.06	
1417	552.60	1.19	
1418	543.20	2.40	
1419	471.50	1.79	
1420	404.30	1.03	1H NMR (400 MHz, DMSO-d6) δ 12.33 (s, 1H), 7.66 - 7.55 (m, 2H), 7.18 (dd, J = 7.3, 0.7 Hz, 1H), 6.67 (dd, J = 8.4, 0.8 Hz, 1H), 6.46 (d, J = 7.8 Hz, 1H), 2.55 - 2.50 (m, 2H, obscured by solvent peak), 2.31 (s,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			3H), 2.21 - 2.13 (m, 1H), 1.82 (dd, J = 11.9, 5.7 Hz, 1H), 1.51 (d, J = 9.4 Hz, 6H), 1.37 (t, J = 12.0 Hz, 1H), 0.86 (d, J = 6.3 Hz, 3H).
1421	515.00	1.27	H NMR (400 MHz, DMSO) δ 10.72 (s, 1H), 7.67 (s, 1H), 7.27 (s, 1H), 7.11 (d, J = 7.3 Hz, 1H), 7.05 (s, 1H), 6.95 (dd, J = 18.9, 8.7 Hz, 2H), 6.84 (dd, J = 19.9, 9.2 Hz, 2H), 6.75 (d, J = 8.4 Hz, 1H), 4.14 (dd, J = 49.8, 18.9 Hz, 7H), 3.80 (s, 3H), 1.53 (s, 3H), 1.28 (s, 3H).
1422	534.00	1.72	1H NMR (400 MHz, MeOD) δ 7.67 - 7.54 (m, 2H), 7.23 (s, 1H), 6.96 (s, 2H), 6.88 (d, J = 8.3 Hz, 1H), 6.75 (d, J = 7.7 Hz, 1H), 6.59 (d, J = 8.3 Hz, 1H), 4.30 (d, J = 12.8 Hz, 1H), 4.07 (d, J = 26.0 Hz, 5H), 3.60 (s, 2H), 3.00 (s, 2H), 2.42 (s, 6H), 2.29 (s, 3H), 2.03 (s, 1H), 1.79 (s, 3H), 1.31 (s, 3H).
1423	430.30	1.27	1H NMR (400 MHz, DMSO-d6) δ 7.73 (d, J = 8.0 Hz, 1H), 7.66 (t, J = 7.8 Hz, 1H), 7.22 (d, J = 7.2 Hz, 1H), 6.92 (d, J = 8.1 Hz, 1H), 6.72 (d, J = 8.3 Hz, 1H), 5.89 (s, 1H), 5.35 (s, 1H), 2.66 - 2.56 (m, 2H), 2.25 - 2.20 (m, 1H), 2.11 (s, 3H), 1.87 (dd, J = 12.1, 5.7 Hz, 1H), 1.53 (d, J = 2.5 Hz, 6H), 1.45 - 1.34 (m, 1H), 0.88 (d, J = 6.3 Hz, 3H).
1424	572.30	1.50	1H NMR (400 MHz, DMSO-d6) δ 12.57 - 12.38 (s, 1H), 7.79 (d, J = 8.0 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.53 - 7.45 (m, 1H), 7.45 - 7.34 (m, 1H), 7.26 (d, J = 8.0 Hz, 1H), 7.21 (d, J = 7.2 Hz, 1H), 6.90 (dt, J = 10.6, 2.3 Hz, 1H), 6.69 (d, J = 8.2 Hz, 1H), 6.64 - 6.10 (s, 1H), 3.80 (s, 2H), 2.67 - 2.56 (m, 2H), 2.33 - 2.15 (m, 1H), 1.94 - 1.82 (m, 1H), 1.61 (s, 3H), 1.60 (s, 3H), 1.50 - 1.34 (m, 1H), 1.21 (s, 6H), 0.88 (d, J = 6.3 Hz, 3H).
1425	511.00	1.71	1H NMR (400 MHz, DMSO) δ 9.77 (s, 1H), 7.67 (d, J = 7.8 Hz, 1H), 7.57 (d, J = 8.9 Hz, 1H), 7.51 (s, 1H), 7.30 (d, J = 23.3 Hz, 2H), 7.02 - 6.90 (m, 4H), 4.28 - 4.12 (m, 4H), 4.07 - 4.01 (m, 2H), 2.39 (s, 6H), 2.25 (s, 3H), 1.76 (s, 3H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1.24 (s, 3H).
1426	565.34	2.30	
1427	537.30	2.11	
1428	518.00	1.43	
1429	662.20	2.07	
1430	558.30	1.84	
1431	512.50	1.56	
1432	565.50	1.94	
1433	485.00	1.12	1H NMR (400 MHz, DMSO) δ 10.35 (s, 1H), 7.74 (s, 1H), 7.60 (s, 2H), 7.28 (s, 4H), 7.16 (s, 1H), 7.03 (s, 1H), 6.93 (s, 1H), 6.83 (s, 1H), 4.32 (s, 3H), 4.15 - 4.02 (m, 5H), 1.63 (s, 3H), 1.27 (t, J = 6.9 Hz, 3H).
1434	566.06	1.77	
1435	562.50	1.62	
1436	525.30	1.89	
1437	551.20	1.81	
1438	472.50	1.31	
1439	516.30	1.40	
1440	557.30	2.44	
1441	526.60	1.33	
1442	538.30	1.25	1H NMR (400 MHz, DMSO-d6) δ 8.49 (d, J = 9.8 Hz, 2H), 7.79 (d, J = 7.9 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.24 (dd, J = 13.1, 7.6 Hz, 3H), 6.72 (d, J = 8.3 Hz, 1H), 4.09 - 3.95 (m, 2H), 3.87 (bs, 1H), 1.95 (p, J = 8.0 Hz, 2H), 1.83 (h, J = 6.8 Hz, 1H), 1.67 (dd, J = 9.4, 5.6 Hz, 2H), 1.18 (d, J = 6.4 Hz, 3H), 1.07 (dd, J = 6.3, 2.3 Hz, 6H), 0.95 (dd, J = 12.0, 6.7 Hz, 6H).
1443	528.40	1.83	1H NMR (400 MHz, Chloroform-d) δ 13.43 (s, 1H), 8.27 (d, J = 8.5 Hz, 1H), 7.95 (d, J = 0.9 Hz, 1H), 7.67 - 7.59 (m, 3H), 7.53 (d, J = 0.9 Hz, 1H), 6.65 (p, J = 3.8 Hz, 1H), 4.66 (s, 2H), 3.72 (d, J = 6.5 Hz, 2H), 3.50 - 3.37 (m, 1H), 3.15 (dd, J = 10.4, 8.1 Hz, 1H), 2.61 (d, J = 6.5 Hz, 1H), 2.16 - 2.02 (m, 2H), 1.73 (dd, J = 12.3, 9.9 Hz, 1H), 1.47 (s, 3H), 1.42 (s, 3H), 1.21 (d, J = 6.6 Hz, 3H), 1.03 (d, J = 6.7 Hz, 6H).
1444	556.30	1.69	1H NMR (400 MHz, DMSO-d6) δ 12.56 (s, 1H), 8.26 (dd, J = 4.8, 1.9 Hz,

Cmpd No.	LC/MS M+1	LC/RT min	NMR
			1H), 8.05 (dd, J = 7.9, 1.9 Hz, 1H), 7.78 (d, J = 8.0 Hz, 1H), 7.52 - 7.46 (m, 1H), 7.46 - 7.37 (m, 1H), 7.30 (d, J = 8.0 Hz, 1H), 6.90 (dt, J = 10.7, 2.3 Hz, 1H), 6.78 (dd, J = 7.9, 4.7 Hz, 1H), 6.72 (s, 2H), 3.84 (d, J = 6.7 Hz, 2H), 3.39 (s, 6H), 2.15 (s, 1H), 2.05 (dt, J = 13.4, 6.7 Hz, 1H), 1.87 (dd, J = 11.9, 5.5 Hz, 1H), 1.59 (d, J = 14.2 Hz, 6H), 1.40 (t, J = 12.1 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.76 (d, J = 6.3 Hz, 3H).
1445	576.60	1.63	
1446	501.20	2.12	
1447	562.60	1.61	
1448	556.60	1.68	
1449	494.00	1.33	1H NMR (400 MHz, DMSO) δ 9.48 (s, 1H), 8.83 (d, J = 6.0 Hz, 2H), 7.99 (d, J = 6.0 Hz, 2H), 7.32 (s, 1H), 7.09 (d, J = 7.7 Hz, 1H), 6.93 (s, 3H), 4.34 (s, 10H), 4.04 (s, 4H), 2.40 (s, 6H), 2.25 (s, 3H), 1.82 (s, 3H), 1.25 (s, 3H).
1450	542.81	2.48	
1451	529.10	2.41	
1452	570.00	2.19	
1453	509.00	1.64	
1454	508.00	1.44	1H NMR (400 MHz, DMSO) δ 10.03 (s, 1H), 7.60 (s, 1H), 7.50 (s, 1H), 7.39 (s, 1H), 7.28 (s, 1H), 7.20 (s, 1H), 6.91 (s, 4H), 4.10 (d, J = 78.6 Hz, 6H), 2.37 (s, 6H), 2.24 (s, 3H), 1.73 (s, 3H), 1.25 (s, 3H).
1455	458.50	1.10	
1456	579.45	3.55	1H NMR (400 MHz, DMSO) δ 8.43 (s, 1H), 7.48 (s, 1H), 7.18-7.15 (d, 1H), 6.98 (s, 2H), 6.89-6.85 (m, 3H), 6.71 (s, 1H), 6.52 (s, 1H), 6.27 (s, 1H), 3.80-3.77 (d, 2H), 2.26 (s, 3H), 2.05 (s, 6H), 0.97-0.94 (d, 3H)
1457	508.00	1.66	
1458	551.20	2.14	
1459	584.60	1.54	
1460	523.20	1.89	
1461	568.60	1.54	
1462	512.30	1.36	
1463	552.30	1.42	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1464	531.08	2.14	
1465	571.30	2.55	
1466	525.30	1.84	1H NMR (400 MHz, DMSO-d6) δ 12.14 (s, 1H), 7.99 (d, J = 7.7 Hz, 1H), 7.63 (dd, J = 8.4, 7.3 Hz, 1H), 7.20 (dd, J = 7.3, 0.7 Hz, 1H), 7.03 (d, J = 7.7 Hz, 1H), 6.91 (s, 2H), 6.70 (dd, J = 8.4, 0.8 Hz, 1H), 6.51 (s, 2H), 3.59 - 3.52 (m, 2H), 2.94 - 2.83 (m, 1H), 2.25 (s, 3H), 1.97 (s, 6H), 1.56 - 1.43 (m, 2H), 1.34 - 1.20 (m, 2H), 1.14 (s, 3H), 1.06 (s, 3H).
1467	548.60	1.59	
1468	416.16	2.91	
1469	584.60	1.57	
1470	523.00	1.73	1H NMR (400 MHz, MeOD) δ 7.46 (s, 2H), 7.37 (s, 1H), 7.27 (s, 1H), 7.09 (d, J = 6.2 Hz, 1H), 6.99 (s, 2H), 6.92 (dd, J = 11.5, 8.0 Hz, 2H), 4.41 (s, 2H), 4.27 (d, J = 13.0 Hz, 2H), 4.04 (s, 2H), 3.80 (s, 3H), 2.45 (s, 6H), 2.30 (s, 3H), 2.03 (s, 1H), 1.90 (s, 3H), 1.32 (s, 3H
1471	526.28	1.08	
1472	570.30	1.88	
1473	514.50	1.40	1H NMR (400 MHz, Chloroform-d) δ 15.64 (s, 1H), 8.45 (d, J = 8.7 Hz, 1H), 7.83 (d, J = 8.7 Hz, 1H), 7.69 - 7.58 (m, 2H), 6.64 (dd, J = 7.0, 2.1 Hz, 1H), 5.71 (d, J = 1.0 Hz, 1H), 4.67 (br s, 2H), 4.28 (q, J = 7.1 Hz, 2H), 3.76 (t, J = 9.3 Hz, 1H), 3.21 (dd, J = 10.1, 7.3 Hz, 1H), 2.68 (s, 3H) overlapped with 2.65-2.72 (m, 1H), 2.21 (dd, J = 12.5, 8.2 Hz, 1H), 1.79 (dd, J = 12.5, 9.0 Hz, 1H), 1.41 (t, J = 7.1 Hz, 3H), 1.32 (s, 3H), 1.26 (s, 6H). Some grease signals detected.
1474	494.00	1.48	1H NMR (400 MHz, DMSO) δ 9.99 (s, 1H), 8.56 (s, 1H), 7.98 (s, 2H), 7.51 (s, 1H), 7.29 (s, 1H), 7.04 (d, J = 7.4 Hz, 1H), 6.91 (s, 3H), 4.21 (s, 1H), 4.13 (s, 2H), 4.01 (s, 3H), 2.38 (s, 6H), 2.24 (s, 3H), 1.75 (s, 3H), 1.25 (s, 3H).
1475	523.20	2.37	
1476	510.00	1.37	1H NMR (400 MHz, MeOD) δ 8.26 (d,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			J = 6.5 Hz, 1H), 8.10 (s, 1H), 7.57 (d, J = 4.9 Hz, 1H), 7.21 (s, 1H), 6.99 (s, 2H), 6.88 (dd, J = 19.2, 7.9 Hz, 2H), 6.44 (s, 1H), 4.54 (d, J = 12.3 Hz, 1H), 4.33 (dd, J = 33.8, 9.1 Hz, 1H), 4.13 (s, 2H), 4.08 - 3.99 (m, 2H), 2.50 (s, 6H), 2.30 (s, 3H), 1.82 (s, 3H), 1.32 (s, 3H).
1477	542.96	2.48	
1478	504.50	2.19	
1479	605.20	2.43	
1480	508.00	1.65	1H NMR (400 MHz, DMSO) δ 10.63 (s, 1H), 7.51 (d, J = 8.7 Hz, 2H), 7.18 (s, 1H), 6.84 (d, J = 11.3 Hz, 3H), 6.64 (d, J = 7.6 Hz, 1H), 6.58 (d, J = 8.7 Hz, 2H), 3.99 (s, 4H), 3.73 (s, 3H), 2.29 (s, 6H), 2.23 (s, 3H), 2.07 (s, 1H), 1.76 (s, 1H), 1.55 (s, 3H), 1.24 (s, 3H).
1481	468.20	2.80	
1482	585.30	2.29	
1483	571.00	3.17	1H NMR (400 MHz, DMSO) δ 12.85 (s, 1H), 8.04 (dd, J = 8.4, 7.4 Hz, 1H), 7.82 (d, J = 8.0 Hz, 1H), 7.77 (dd, J = 7.3, 0.5 Hz, 1H), 7.49 (d, J = 1.7 Hz, 1H), 7.42 (dd, J = 9.8, 1.7 Hz, 1H), 7.30 (d, J = 8.1 Hz, 1H), 7.16 (dd, J = 8.4, 0.5 Hz, 1H), 6.90 (dt, J = 10.7, 2.3 Hz, 1H), 3.91 (s, 3H), 3.83 (d, J = 6.6 Hz, 2H), 2.50 - 2.36 (m, 2H), 2.30 - 2.15 (m, 1H), 2.05 (dt, J = 13.3, 6.7 Hz, 1H), 1.87 (dd, J = 11.8, 5.5 Hz, 1H), 1.59 (d, J = 1.3 Hz, 6H), 1.40 (t, J = 12.1 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.77 (d, J = 6.3 Hz, 3H).
1484	554.60	2.01	
1485	528.34	1.30	1H NMR (400 MHz, Methanol-d4) δ 7.94 (s, 1H), 7.91 (s, 1H), 7.77 - 7.70 (m, 1H), 7.69 - 7.62 (m, 1H), 7.37 (d, J = 2.4 Hz, 1H), 7.34 (s, 1H), 6.85 (d, J = 8.5 Hz, 1H), 6.76 (d, J = 8.4 Hz, 1H), 4.20 - 4.09 (m, 2H), 3.62 (t, J = 6.4 Hz, 2H), 3.37 (t, J = 10.0 Hz, 1H), 3.15 - 3.02 (m, 2H), 2.90 (dt, J = 17.7, 7.3 Hz, 1H), 2.73 (d, J = 7.6 Hz, 2H), 2.56 (s, 1H), 2.19 (td, J = 6.4, 4.0 Hz, 2H), 2.09 (dd, J = 12.5, 6.6 Hz, 1H), 2.03 (s,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			3H), 1.92 (dq, J = 12.4, 6.3 Hz, 3H), 1.77 (ddd, J = 14.3, 8.9, 5.4 Hz, 2H), 1.72 - 1.64 (m, 2H), 1.63 (s, 3H), 1.56 (s, 3H), 1.54 (s, 3H), 1.45 (s, 3H), 1.43 (s, 8H), 1.35 (s, 9H), 1.10 (d, J = 6.5 Hz, 3H), 0.95 (d, J = 6.2 Hz, 3H). (two sets of peaks due to a 1:1 mixture of olefin and hemi-acetal)
1486	483.20	2.14	
1487	528.34	1.21	
1488	524.28	1.21	1H NMR (400 MHz, DMSO-d6) δ 12.41 (s, 1H), 7.77 (d, J = 8.0 Hz, 1H), 7.63 (dd, J = 8.4, 7.3 Hz, 1H), 7.42 (d, J = 2.0 Hz, 2H), 7.19 (t, J = 7.4 Hz, 2H), 6.82 (t, J = 1.9 Hz, 1H), 6.68 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 4.07 (q, J = 6.9 Hz, 2H), 2.63 - 2.55 (m, 2H), 2.34 (s, 3H), 2.23 (td, J = 13.6, 13.1, 6.9 Hz, 1H), 1.87 (dd, J = 11.9, 5.7 Hz, 1H), 1.61 (d, J = 4.4 Hz, 6H), 1.42 (t, J = 12.1 Hz, 1H), 1.34 (t, J = 6.9 Hz, 3H), 0.88 (d, J = 6.2 Hz, 3H).
1489	460.30	1.56	
1490	544.60	1.79	
1491	525.30	1.90	
1492	537.30	1.84	
1493	553.50	2.20	
1494	510.30	2.11	1H NMR (400 MHz, DMSO-d6) δ 10.59 (s, 1H), 7.90 (d, J = 8.7 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.22 (d, J = 7.1 Hz, 1H), 6.94 (s, 2H), 6.69 (d, J = 8.2 Hz, 1H), 6.45 (s, 2H), 6.26 (d, J = 8.8 Hz, 1H), 3.69 (s, 2H), 2.26 (s, 3H), 2.05 (s, 6H), 1.96 - 1.85 (m, 2H), 1.57 (q, J = 9.5, 8.6 Hz, 2H), 0.92 (s, 6H).
1495	528.50	1.43	
1496	518.22	2.99	DMSO(d6) δ 8.86 - 8.84 (d, 1H), 8.80 (s, 1H), 8.54 - 8.51 (d, 1H), 8.17 - 8.14 (d, 1H), 8.01 - 7.99 (d, 1H), 7.84 (t, J = 7.81, 1H), 7.01 - 6.84 (m, 4H), 5.44 (s, 2H), 2.72 (s, 3H), 2.03 (s, 6H), 1.13 (s, 9H)
1497	573.10	1.77	1H NMR (400 MHz, Methanol-d4) δ 8.03 (dd, J = 8.3, 1.2 Hz, 1H), 7.68 (dd, J = 8.5, 7.3 Hz, 1H), 7.59 (dd, J =

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			8.3, 7.1 Hz, 1H), 7.38 (dd, J = 7.4, 0.8 Hz, 1H), 6.93 - 6.82 (m, 2H), 6.82 - 6.74 (m, 2H), 3.92 (t, J = 9.4 Hz, 1H), 3.79 (d, J = 6.4 Hz, 2H), 2.84 - 2.72 (m, 1H), 2.23 (dd, J = 12.6, 7.3 Hz, 1H), 2.12 - 2.02 (m, 1H), 1.87 (t, J = 11.7 Hz, 1H), 1.38 (s, 3H), 1.24 (d, J = 6.6 Hz, 6H), 1.04 (d, J = 6.7 Hz, 6H).
1498	495.00	1.23	
1499	545.03	2.16	1H NMR (400 MHz, DMSO) δ 11.84 (s, 1H), 7.98 (d, J = 7.8 Hz, 1H), 7.74 (d, J = 7.9 Hz, 1H), 7.70 - 7.62 (m, 1H), 7.52 (t, J = 5.6 Hz, 2H), 7.21 (d, J = 7.3 Hz, 1H), 6.96 (dd, J = 10.8, 2.1 Hz, 1H), 6.71 (d, J = 8.4 Hz, 1H), 6.51 (s, 2H), 4.48 (dd, J = 10.4, 6.5 Hz, 1H), 4.38 (dd, J = 10.4, 7.8 Hz, 1H), 3.86 (d, J = 6.5 Hz, 2H), 3.85 - 3.76 (m, 2H), 3.74 - 3.66 (m, 1H), 3.66 - 3.58 (m, 1H), 2.71 (dt, J = 14.5, 7.4 Hz, 1H), 2.04 (qd, J = 13.5, 7.5 Hz, 2H), 1.76 (dt, J = 14.3, 6.8 Hz, 1H), 1.00 (d, J = 6.7 Hz, 6H).
1500	474.30	1.34	
1501	514.24	2.00	
1502	567.40	2.39	1H NMR (400 MHz, MeOD) δ 7.86 (d, J = 8.2 Hz, 1H), 7.78 (d, J = 8.2 Hz, 1H), 7.66 - 7.59 (m, 1H), 7.53 (s, 1H), 7.39 (d, J = 9.5 Hz, 1H), 7.36 (d, J = 7.2 Hz, 1H), 6.78 - 6.72 (m, 2H), 5.64 (s, 1H), 3.82 (d, J = 6.5 Hz, 2H), 2.15 - 2.05 (m, 1H), 1.75 (s, 2H), 1.38 (s, 6H), 1.09 - 1.00 (m, 12H).
1503	475.30	1.27	
1504	553.00	1.76	1H NMR (400 MHz, DMSO) δ 12.43 (s, 1H), 8.83 (s, 1H), 8.31 (d, J = 8.7 Hz, 1H), 7.78 (d, J = 8.0 Hz, 1H), 7.64 (s, 1H), 7.20 (s, 2H), 6.96 (d, J = 8.7 Hz, 1H), 6.70 (d, J = 8.3 Hz, 1H), 4.02 (s, 2H), 2.60 (d, J = 9.6 Hz, 2H), 2.22 (s, 1H), 1.87 (s, 1H), 1.60 (s, 6H), 1.42 (s, 1H), 0.88 (d, J = 6.3 Hz, 3H).
1505	432.30	1.31	
1506	538.30	1.29	1H NMR (400 MHz, DMSO-d6) δ 8.61 - 8.48 (m, 2H), 7.81 (d, J = 7.9 Hz, 1H),

Cmpd. No.	LC/MS1 M+1	LC/RT min	NMR
			7.65 (dd, J = 8.4, 7.3 Hz, 1H), 7.38 (d, J = 9.5 Hz, 1H), 7.32 (d, J = 7.9 Hz, 1H), 7.22 (d, J = 7.4 Hz, 1H), 6.74 (d, J = 8.4 Hz, 1H), 4.01 (h, J = 5.6 Hz, 2H), 3.56 (d, J = 7.5 Hz, 2H), 3.28 (s, 3H), 2.15 - 2.02 (m, 1H), 1.96 (qq, J = 6.6, 3.2 Hz, 2H), 1.68 (tq, J = 8.5, 5.2, 4.7 Hz, 2H), 1.08 (d, J = 6.2 Hz, 6H), 0.93 (d, J = 6.6 Hz, 6H).
1507	418.30	1.17	
1508	576.60	1.61	
1509	509.00	1.70	
1510	458.30	1.48	
1511	446.20	1.49	
1512	525.30	2.31	1H NMR (400 MHz, DMSO-d6) δ 7.64 - 7.58 (m, 2H), 7.51 - 7.44 (m, 1H), 7.42 (d, J = 7.7 Hz, 1H), 6.98 (d, J = 7.3 Hz, 1H), 6.60 (t, J = 1.1 Hz, 1H), 6.49 (d, J = 8.2 Hz, 1H), 6.24 (s, 2H), 4.40 (q, J = 7.0 Hz, 2H), 2.93 (dt, J = 21.0, 10.6 Hz, 2H), 2.31 (s, 3H), 2.09 (bs, 1H), 1.81 (dd, J = 11.6, 5.6 Hz, 1H), 1.64 (s, 3H), 1.58 (s, 3H), 1.35 (t, J = 6.9 Hz, 4H), 0.79 (d, J = 6.3 Hz, 3H).
1513	507.50	2.16	
1514	525.30	1.92	1H NMR (400 MHz, DMSO-d6) δ 8.23 (s, 1H), 7.89 (dd, J = 7.8, 2.1 Hz, 1H), 7.70 - 7.55 (m, 1H), 7.22 (d, J = 7.3 Hz, 1H), 7.04 (dd, J = 8.0, 2.7 Hz, 1H), 6.71 (d, J = 10.0 Hz, 2H), 5.28 (h, J = 6.2 Hz, 1H), 3.94 (q, J = 6.0 Hz, 2H), 2.36 (s, 3H), 1.99 (p, J = 7.4 Hz, 2H), 1.70 - 1.55 (m, 2H), 1.30 (d, J = 6.1 Hz, 6H), 1.07 (d, J = 6.2 Hz, 6H).
1515	556.50	1.83	
1516	510.30	2.11	1H NMR (400 MHz, DMSO-d6) δ 10.56 (s, 1H), 7.72 - 7.57 (m, 2H), 7.21 (d, J = 7.0 Hz, 1H), 6.93 (s, 2H), 6.69 (d, J = 8.4 Hz, 1H), 6.45 (s, 2H), 3.76 - 3.60 (m, 2H), 3.53 (dt, J = 10.1, 6.8 Hz, 1H), 2.28 (s, 3H), 2.26 (s, 3H), 2.08 (s, 3H), 2.01 (s, 3H), 1.82 (ddd, J = 11.4, 6.6, 3.2 Hz, 2H), 1.74 - 1.59 (m, 1H), 1.37 (td, J = 9.9, 4.0 Hz, 1H), 0.62 (d, J = 6.1 Hz, 3H).

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1517	511.20	1.92	
1518	505.20	2.18	
1519	563.20	2.13	
1520	676.60	2.46	
1521	560.50	1.61	
1522	627.44	1.34	
1523	540.00	1.67	1H NMR (400 MHz, DMSO) δ 12.56 (s, 1H), 8.94 (s, 1H), 8.36 (s, 1H), 7.88 (d, J = 7.9 Hz, 1H), 7.64 (s, 1H), 7.54 (d, J = 7.9 Hz, 1H), 7.22 (d, J = 7.3 Hz, 1H), 6.70 (d, J = 8.4 Hz, 1H), 4.23 (d, J = 6.4 Hz, 2H), 2.64 (s, 2H), 2.19 (d, J = 38.0 Hz, 2H), 1.89 (s, 1H), 1.62 (s, 6H), 1.44 (s, 1H), 1.03 (d, J = 6.7 Hz, 6H), 0.89 (d, J = 6.2 Hz, 3H).
1524	472.30	1.23	
1525	504.30	2.22	
1526	557.30	2.22	1H NMR (400 MHz, DMSO-d6) δ 12.56 (s, 1H), 8.51 (s, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.56 - 7.45 (m, 2H), 7.22 (d, J = 7.2 Hz, 1H), 6.98 (dt, J = 10.9, 2.3 Hz, 1H), 6.69 (d, J = 8.4 Hz, 1H), 6.55 (s, 2H), 3.85 (d, J = 6.7 Hz, 2H), 2.72 - 2.63 (m, 1H), 2.59 (t, J = 10.6 Hz, 1H), 2.23 (s, 1H), 2.06 (dt, J = 13.3, 6.6 Hz, 1H), 1.92 (dd, J = 12.0, 5.6 Hz, 1H), 1.62 (s, 3H), 1.55 (s, 3H), 1.47 (t, J = 12.2 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.89 (d, J = 6.3 Hz, 3H).
1527	552.30	1.42	
1528	535.50	1.76	
1529	540.30	2.11	1H NMR (400 MHz, DMSO) δ 12.58 (s, 1H), 7.74 (d, J = 7.9 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.45 (dd, J = 8.7, 2.3 Hz, 2H), 7.33 (d, J = 7.9 Hz, 1H), 7.21 (d, J = 7.0 Hz, 1H), 6.89 (dt, J = 10.8, 2.3 Hz, 1H), 6.70 (d, J = 8.1 Hz, 1H), 6.49 (s, 2H), 3.84 (d, J = 6.5 Hz, 2H), 3.54 (s, 4H), 2.46 (t, J = 5.8 Hz, 2H), 2.06 (ddd, J = 19.9, 9.8, 4.6 Hz, 3H), 1.26 (dd, J = 6.6, 2.6 Hz, 2H), 0.99 (d, J = 6.7 Hz, 6H).
1530	555.20	2.34	
1531	569.50	2.01	
1532	542.60	1.89	1H NMR (400 MHz, Chloroform-d) δ

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			14.55 (br s, 1H), 8.31 (br s, 1H), 7.93 (br s, 1H), 7.57 (br m, 3H), 6.59 (br s, 1H), 4.75 and 5.49 (br m, 2H), 4.04 (d, J = 6.7 Hz, 2H), 3.55 (br s, 1H), 3.17 (br s, 1H), 2.66 (br s, 1H), 2.11 (dq, J = 13.4, 6.7 Hz, 2H), 2.00 (s, 3H), 1.82 (br s, 1H), 1.42-1.36 (br m, 6H), 1.21 (br s, 3H), 1.02 (d, J = 6.7 Hz, 6H).
1533	604.60	2.10	
1534	460.30	1.03	
1535	578.60	1.78	
1536	420.20	1.09	
1537	539.30	2.17	1H NMR (400 MHz, DMSO-d6) δ 7.92 (d, J = 7.9 Hz, 1H), 7.85 (d, J = 8.4 Hz, 1H), 7.66 (dd, J = 8.4, 7.3 Hz, 1H), 7.22 (d, J = 7.0 Hz, 1H), 7.07 (d, J = 7.8 Hz, 1H), 6.76 (dd, J = 17.4, 8.5 Hz, 2H), 4.08 (d, J = 6.6 Hz, 2H), 3.96 (q, J = 5.9 Hz, 2H), 2.08 - 1.93 (m, 3H), 1.73 - 1.56 (m, 2H), 1.08 (d, J = 6.2 Hz, 6H), 0.98 (d, J = 6.7 Hz, 6H).
1538	484.30	1.65	1H NMR (400 MHz, DMSO-d6) δ 12.58 - 12.52 (s, 1H), 10.09 (s, 1H), 8.38 - 7.93 (m, 2H), 7.78 - 7.69 (m, 2H), 7.69 - 7.62 (m, 2H), 7.29 (d, J = 2.0 Hz, 1H), 7.23 (d, J = 10.2 Hz, 1H), 7.14 (d, J = 7.9 Hz, 1H), 6.90 - 6.36 (m, 1H), 2.48 - 2.26 (m, 2H), 2.18 - 2.08 (m, 1H), 1.91 - 1.70 (m, 1H), 1.58 (s, 3H), 1.55 (s, 3H), 1.38 (t, J = 12.1 Hz, 1H), 0.66 (d, J = 6.2 Hz, 3H).
1539	493.00	1.72	1H NMR (400 MHz, MeOD) δ 7.93 (s, 2H), 7.50 (d, J = 32.1 Hz, 3H), 7.26 (s, 1H), 6.98 (s, 2H), 6.91 (s, 2H), 4.31 (d, J = 65.9 Hz, 4H), 4.05 (s, 2H), 2.44 (s, 6H), 2.29 (s, 3H), 2.03 (s, 1H), 1.87 (s, 3H), 1.32 (s, 3H).
1540	564.50	1.69	
1541	510.00	1.37	1H NMR (400 MHz, MeOD) δ 8.26 (d, J = 6.4 Hz, 1H), 8.10 (s, 1H), 7.57 (d, J = 5.1 Hz, 1H), 7.21 (s, 1H), 6.99 (s, 2H), 6.88 (dd, J = 19.4, 7.9 Hz, 2H), 6.44 (s, 1H), 4.54 (d, J = 12.7 Hz, 1H), 4.33 (dd, J = 34.0, 9.6 Hz, 1H), 4.13 (s,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			2H), 4.08 - 3.99 (m, 2H), 2.50 (s, 6H), 2.31 (s, 3H), 1.82 (s, 3H), 1.32 (s, 3H).
1542	500.00	1.45	1H NMR (400 MHz, DMSO) δ 9.88 (s, 1H), 9.06 (s, 1H), 8.24 (s, 1H), 7.28 (s, 1H), 7.07 (d, J = 7.6 Hz, 1H), 6.92 (s, 3H), 4.27 - 4.12 (m, 3H), 4.05 - 3.97 (m, 3H), 2.40 (s, 7H), 2.25 (s, 3H), 1.76 (s, 3H), 1.25 (t, J = 6.9 Hz, 3H).
1543			
1544	524.40	1.89	
1545	577.60	2.29	
1546	500.00	1.49	1H NMR (400 MHz, DMSO) δ 9.50 (s, 1H), 7.86 (d, J = 3.1 Hz, 1H), 7.82 (d, J = 3.1 Hz, 1H), 7.34 (s, 1H), 7.19 (d, J = 7.7 Hz, 1H), 6.95 (d, J = 12.8 Hz, 3H), 4.34 (s, 2H), 4.06 (s, 2H), 2.42 (s, 6H), 2.25 (s, 3H), 1.84 (s, 3H), 1.23 (s, 3H).
1547	516.50	1.47	
1548	572.20	1.99	1H NMR (400 MHz, DMSO) δ 15.07 (s, 1H), 8.28 (d, J = 8.2 Hz, 1H), 8.10 (d, J = 8.2 Hz, 1H), 7.65 (dd, J = 8.4, 7.4 Hz, 1H), 7.59 - 7.50 (m, 2H), 7.31 - 7.19 (m, 1H), 6.99 (dt, J = 10.8, 2.3 Hz, 1H), 6.71 (d, J = 7.9 Hz, 1H), 6.38 (s, 2H), 3.93 (ddd, J = 13.4, 6.0, 2.9 Hz, 1H), 3.87 (d, J = 6.6 Hz, 2H), 3.76 (d, J = 11.4 Hz, 1H), 3.63 (d, J = 11.4 Hz, 1H), 3.39 - 3.27 (m, 1H), 3.02 (dd, J = 12.5, 2.6 Hz, 1H), 2.06 (dq, J = 13.3, 6.6 Hz, 1H), 1.52 (s, 3H), 1.18 (d, J = 6.2 Hz, 3H), 1.07 - 0.95 (m, 9H).
1549	578.55	3.75	7.61-7.58 (d, 1H), 7.46 (t, J=7.15, 1H), 7.21-7.18 (d, 1H), 7.04-7.01 (d, 1H), 6.94 (s, 1H), 6.80-6.68 (m, 2H), 6.51-6.47 (d, 1H),6.32-6.28 (d,3H), 3.76-3.74 (d, 2H), 2.24 (s, 3H), 2.03 (s, 6H), 0.96 -0.93 (d, 6H)
1550	458.30	1.75	1H NMR (400 MHz, DMSO-d6) δ 12.62 (s, 1H), 8.47 (d, J = 2.5 Hz, 1H), 7.89 (d, J = 2.5 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.21 (d, J = 7.2 Hz, 1H), 6.70 (d, J = 8.5 Hz, 1H), 6.52 (s, 2H), 2.65 - 2.52 (m, 2H), 2.22 - 2.17 (m, 1H), 1.92 - 1.82 (m, 1H), 1.52 (s, 6H), 1.42 (t, J = 12.1 Hz, 1H), 0.86 (d, J

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			= 6.3 Hz, 3H).
1551	456.38	3.05	
1552	569.30	2.40	1H NMR (400 MHz, DMSO) δ 12.63 (s, 1H), 7.91 (dd, J = 20.4, 8.2 Hz, 2H), 7.66 (t, J = 7.9 Hz, 1H), 7.57 (s, 1H), 7.51 (d, J = 9.8 Hz, 1H), 7.25 (d, J = 7.3 Hz, 1H), 6.93 (d, J = 10.7 Hz, 1H), 6.73 (d, J = 8.4 Hz, 1H), 6.51 (s, 2H), 3.86 (d, J = 6.6 Hz, 2H), 3.58 (dd, J = 12.2, 6.5 Hz, 1H), 2.55 (d, J = 12.3 Hz, 1H), 2.06 (dt, J = 13.2, 6.6 Hz, 1H), 1.63 (dd, J = 12.4, 6.4 Hz, 1H), 1.50 (d, J = 13.0 Hz, 1H), 1.39 (d, J = 12.9 Hz, 1H), 1.15 (s, 3H), 1.08 - 0.93 (m, 9H), 0.80 (s, 3H), 0.69 (s, 3H).
1553	496.50	1.95	
1554	562.60	1.59	
1555	557.00	1.46	1H NMR (400 MHz, DMSO) δ 12.27 (s, 1H), 9.74 (s, 1H), 7.78 (s, 1H), 7.28 (s, 1H), 7.05 (d, J = 7.7 Hz, 1H), 6.93 (s, 3H), 4.22 (dd, J = 26.1, 10.6 Hz, 4H), 4.03 (d, J = 7.4 Hz, 3H), 2.43 (s, 6H), 2.25 (s, 4H), 2.14 (s, 3H), 1.77 (s, 3H), 1.23 (s, 3H).
1556	458.30	1.72	
1557	548.50	1.60	
1558	551.50	1.75	
1559	621.30	2.35	
1560	611.50	2.06	
1561	500.30	1.73	
1562	550.40	2.60	1H NMR (400 MHz, DMSO-d6) δ 10.48 (s, 1H), 7.30 (s, 1H), 7.26 - 7.07 (m, 3H), 6.86 (d, J = 6.7 Hz, 3H), 6.78 (d, J = 7.6 Hz, 1H), 4.09 - 3.94 (m, 3H), 3.79 (d, J = 16.5 Hz, 4H), 3.00 (p, J = 6.8 Hz, 1H), 2.31 (s, 6H), 2.23 (s, 3H), 1.63 (s, 3H), 1.24 (t, J = 6.9 Hz, 3H), 1.14 (dd, J = 6.8, 3.0 Hz, 6H).
1563	571.20	2.73	1H NMR (400 MHz, Methanol-d4) δ 7.74 (d, J = 7.9 Hz, 1H), 7.46 (t, J = 1.8 Hz, 1H), 7.35 - 7.20 (m, 2H), 7.01 (d, J = 7.7 Hz, 1H), 6.92 (d, J = 7.9 Hz, 1H), 6.65 (dt, J = 10.4, 2.3 Hz, 1H), 3.79 (d, J = 6.5 Hz, 2H), 3.02 - 2.90 (m, 1H), 2.18 (dtd, J = 12.0, 9.2, 7.9, 4.3 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 2.08 (dh, J = 13.4, 6.7 Hz, 1H), 1.83 (dd, J = 11.7, 5.6 Hz, 1H), 1.65 (m, 1H), 1.50 - 1.38 (m, 1H), 1.39 - 1.23 (m, 6H), 1.04 (d, J = 6.7 Hz, 6H),
1564	415.20	1.49	
1565	502.00	1.13	
1566	616.50	1.99	
1567	572.40	1.48	1H NMR (400 MHz, DMSO-d6) δ 12.46 (s, 1H), 7.78 (d, J = 8.0 Hz, 1H), 7.63 (dd, J = 8.4, 7.3 Hz, 1H), 7.57 - 7.45 (m, 1H), 7.45 - 7.34 (m, 1H), 7.26 (d, J = 8.0 Hz, 1H), 7.21 (d, J = 7.3 Hz, 1H), 6.93 - 6.80 (m, 1H), 6.69 (d, J = 8.4 Hz, 1H), 6.46 (s, 1H), 4.03 (dd, J = 9.4, 6.0 Hz, 1H), 3.90 (dd, J = 9.6, 6.3 Hz, 1H), 3.46 - 3.39 (m, 2H), 2.64 - 2.55 (m, 2H), 2.28 - 2.14 (m, 1H), 2.06 - 1.95 (m, 1H), 1.95 - 1.80 (m, 1H), 1.61 (s, 3H), 1.59 (s, 3H), 1.42 (t, J = 12.1 Hz, 1H), 0.97 (d, J = 6.8 Hz, 3H), 0.88 (d, J = 6.3 Hz, 3H).
1568	444.00	1.94	1H NMR (400 MHz, DMSO) δ 8.04 (s, 1H), 7.82 (s, 1H), 7.03 (s, 1H), 6.78 (s, 2H), 5.48 (s, 2H), 3.75 (s, 3H), 2.16 (d, J = 21.1 Hz, 9H), 1.14 (s, 9H).
1569	578.30	2.63	
1570	558.98	2.36	
1571	456.40	1.44	1H NMR (400 MHz, DMSO-d6) δ 12.39 (s, 1H), 8.12 (s, 1H), 7.61 (d, J = 13.8 Hz, 2H), 7.19 (d, J = 7.3 Hz, 1H), 6.67 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 6.36 (d, J = 15.9 Hz, 1H), 5.68 (dd, J = 15.9, 9.0 Hz, 1H), 2.17 (s, 1H), 1.82 (dd, J = 11.9, 5.7 Hz, 1H), 1.55 (dq, J = 8.8, 4.7, 4.2 Hz, 1H), 1.48 (d, J = 5.3 Hz, 6H), 1.36 (t, J = 12.1 Hz, 1H), 0.85 (d, J = 6.3 Hz, 3H), 0.81 - 0.71 (m, 2H), 0.50 (dq, J = 8.8, 5.5, 3.7 Hz, 2H).
1572	590.20	2.24	1H NMR (400 MHz, DMSO) δ 12.63 (s, 1H), 7.98 (d, J = 8.0 Hz, 1H), 7.74 - 7.61 (m, 2H), 7.41 - 7.33 (m, 2H), 7.27 - 7.20 (m, 1H), 6.90 (s, 1H), 6.86 (dt, J = 10.6, 2.3 Hz, 1H), 6.82 (s, 1H), 6.73 (d, J = 8.0 Hz, 1H), 6.52 (s, 2H), 3.70 (d, J = 6.6 Hz, 2H), 3.42 (s, 2H), 3.33

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(s, 2H), 2.25 (s, 3H), 2.08 - 1.96 (m, 1H), 1.91 (s, 3H), 0.98 (d, J = 6.7 Hz, 7H).
1573	448.20	1.32	
1574	566.20	1.43	
1575	474.20	1.13	
1576	509.00	1.41	1H NMR (400 MHz, DMSO) δ 10.47 (s, 1H), 7.58 (s, 1H), 7.46 (d, J = 8.0 Hz, 2H), 7.30 (d, J = 7.9 Hz, 3H), 7.09 (d, J = 7.3 Hz, 1H), 6.93 (dd, J = 19.4, 7.9 Hz, 2H), 6.64 (d, J = 8.2 Hz, 1H), 6.40 (s, 1H), 4.20 (d, J = 12.5 Hz, 1H), 4.05 (d, J = 24.9 Hz, 4H), 2.93 (s, 1H), 2.07 (s, 1H), 1.57 (s, 3H), 1.26 (s, 9H).
1577	538.40	1.91	
1578	579.52	1.50	
1579	573.35	2.17	
1580	515.02	2.35	
1581	556.50	1.83	1H NMR (400 MHz, DMSO-d6) δ 12.62 - 12.53 (s,1H), 8.17 - 7.93 (m, 2H), 7.78 - 7.71 (m, 2H), 7.71 - 7.61 (m, 2H), 7.52 - 7.44 (m, 1H), 7.43 - 7.35 (m, 1H), 7.27 (d, J = 8.0 Hz, 1H), 6.90 (dt, J = 10.6, 2.4 Hz, 1H), 3.80 (s, 2H), 2.48 - 2.26 (m, 2H), 2.26 - 2.10 (m, 1H), 1.84 (dd, J = 11.9, 5.5 Hz, 1H), 1.60 (s, 3H), 1.55 (s, 3H), 1.38 (t, J = 12.0 Hz, 1H), 1.21 (s, 6H), 0.66 (d, J = 6.3 Hz, 3H).
1582	509.00	1.34	1H NMR (400 MHz, MeOD) δ 7.52 (s, 1H), 7.24 (s, 2H), 6.97 (s, 3H), 6.87 (d, J = 7.9 Hz, 1H), 6.61 (s, 1H), 4.39 (s, 1H), 4.16 (s, 2H), 4.03 (s, 2H), 2.44 (s, 6H), 2.29 (s, 3H), 1.85 (s, 3H), 1.32 (s, 4H), 1.15 (d, J = 6.1 Hz, 1H).
1583	488.00	1.31	
1584	536.40	2.49	1H NMR (400 MHz, DMSO-d6) δ 10.41 (s, 1H), 7.19 (dt, J = 30.9, 7.9 Hz, 3H), 6.85 (d, J = 22.6 Hz, 4H), 4.16 - 3.92 (m, 4H), 3.82 (d, J = 14.9 Hz, 4H), 2.33 (s, 6H), 2.23 (s, 3H), 1.64 (s, 3H), 1.24 (t, J = 6.9 Hz, 3H), 1.13 (td, J = 7.5, 2.5 Hz, 3H).
1585	441.50	2.01	1H NMR (400 MHz, DMSO-d6) δ 8.96 - 8.85 (m, 2H), 7.96 - 7.87 (m, 2H),

Cmpd No.	LC/MS M+1	LC/RT min	NMR
			6.94 (s, 1H), 6.73 (s, 2H), 5.44 (s, 2H), 2.16 (s, 3H), 2.05 (s, 6H), 1.15 (s, 9H).
1586	604.60	2.09	
1587	542.00	1.75	1H NMR (400 MHz, DMSO) δ 10.28 (s, 1H), 7.35 (d, J = 2.1 Hz, 1H), 7.29 - 7.22 (m, 2H), 6.96 (dd, J = 8.3, 2.2 Hz, 1H), 6.89 (d, J = 7.2 Hz, 4H), 5.70 (s, 2H), 4.13 (d, J = 12.7 Hz, 1H), 3.97 (dd, J = 33.5, 9.7 Hz, 5H), 2.36 (s, 6H), 2.24 (s, 3H), 2.08 (s, 3H), 1.68 (s, 3H), 1.25 (s, 3H).
1588	617.30	2.64	1H NMR (400 MHz, Chloroform-d) δ 8.87 (d, J = 2.5 Hz, 1H), 8.55 (d, J = 8.3 Hz, 1H), 8.22 (dd, J = 8.7, 2.5 Hz, 1H), 7.68 (dd, J = 8.2, 4.0 Hz, 2H), 7.47 - 7.33 (m, 5H), 7.08 (d, J = 8.1 Hz, 1H), 6.78 (d, J = 8.7 Hz, 1H), 5.39 (hept, J = 6.2 Hz, 1H), 5.12 (s, 2H), 3.96 - 3.68 (m, 2H), 2.29 (ddt, J = 9.6, 6.7, 3.6 Hz, 2H), 2.03 (d, J = 13.7 Hz, 2H), 1.38 (d, J = 6.2 Hz, 6H), 1.27 (dd, J = 6.8, 2.8 Hz, 6H).
1589	524.30	2.17	1H NMR (400 MHz, DMSO-d6) δ 10.54 (s, 1H), 7.70 (s, 1H), 7.67 - 7.59 (m, 1H), 7.21 (d, J = 7.4 Hz, 1H), 6.93 (s, 2H), 6.69 (d, J = 8.6 Hz, 1H), 3.99 (p, J = 6.1, 5.7 Hz, 2H), 2.29 (s, 3H), 2.25 (s, 3H), 2.04 (s, 6H), 1.84 (pt, J = 6.6, 3.0 Hz, 2H), 1.61 - 1.48 (m, 2H), 0.88 (d, J = 6.2 Hz, 6H).
1590	458.20	1.54	
1591	509.00	1.20	1H NMR (400 MHz, MeOD) δ 8.52 (s, 1H), 8.00 (s, 1H), 7.31 (s, 1H), 7.05 (d, J = 27.3 Hz, 4H), 6.93 (d, J = 8.2 Hz, 1H), 4.50 (s, 2H), 4.33 (d, J = 45.1 Hz, 2H), 4.06 (s, 2H), 2.49 (s, 6H), 2.31 (s, 3H), 1.93 (s, 3H), 1.33 (s, 3H).
1592	531.09	2.15	
1593	556.30	1.69	
1594	472.30	1.10	1H NMR (400 MHz, DMSO-d6) δ 12.36 (s, 1H), 7.69 (d, J = 8.0 Hz, 1H), 7.62 (dd, J = 8.4, 7.3 Hz, 1H), 7.22 - 7.15 (m, 1H), 6.81 (d, J = 8.0 Hz, 1H), 6.75 - 6.69 (m, 1H), 6.67 (d, J = 8.5, 0.8 Hz, 1H), 6.47 (s, 2H), 4.26 (q, J =

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			2.8 Hz, 2H), 3.81 (dq, J = 14.5, 5.6 Hz, 2H), 2.59 - 2.54 (m, 2H, obscured by solvent), 2.21 - 2.16 (m, 1H), 1.83 (dd, J = 11.8, 5.6 Hz, 1H), 1.52 (s, 6H), 1.38 (t, J = 12.1 Hz, 1H), 1.22 - 1.11 (m, 1H), 0.86 (d, J = 6.3 Hz, 3H).
1595	474.40	1.12	
1596	543.00	1.48	
1597	544.00	1.64	
1598	535.20	1.92	
1599	446.40	1.44	1H NMR (400 MHz, DMSO-d6) δ 7.47 - 7.32 (m, 2H), 6.99 (d, J = 7.5 Hz, 1H), 6.42 (d, J = 8.1 Hz, 1H), 6.21 (d, J = 7.7 Hz, 1H), 6.01 (s, 2H), 3.04 - 2.90 (m, 2H), 2.44 - 2.28 (m, 2H), 2.17 - 2.00 (m, 2H), 1.76 (dd, J = 11.6, 5.7 Hz, 1H), 1.50 (d, J = 5.4 Hz, 6H), 1.34 (t, J = 5.9 Hz, 1H), 0.92 - 0.79 (m, 9H).
1600	640.60	2.11	
1601	526.60	1.78	1H NMR (400 MHz, Methanol-d4) δ 7.79 (s, 1H), 7.60 (dd, J = 8.4, 7.3 Hz, 1H), 7.33 (d, J = 7.8 Hz, 1H), 6.69 (d, J = 8.1 Hz, 1H), 5.56 (tt, J = 3.6, 1.7 Hz, 1H), 3.60 (dd, J = 11.2, 8.3 Hz, 1H), 3.21 (dd, J = 11.2, 9.0 Hz, 1H), 2.66 (dt, J = 13.8, 8.4 Hz, 1H), 2.30 - 2.22 (m, 2H), 2.20 - 2.15 (m, 2H), 2.15 - 2.10 (m, 1H), 1.83 - 1.75 (m, 3H), 1.75 - 1.67 (m, 2H), 1.52 (s, 3H), 1.42 (s, 9H), 1.40 (s, 3H), 1.15 (d, J = 6.7 Hz, 3H).
1602	468.20	2.80	
1603	424.00	1.75	1H NMR (400 MHz, DMSO-d6) δ 12.54 (s, 1H), 8.27 - 8.00 (m, 1H), 7.72 - 7.54 (m, 2H), 7.18 (d, J = 7.2 Hz, 1H), 6.66 (d, J = 8.4 Hz, 1H), 6.50 (s, 2H), 2.55 (s, 2H), 2.17 (dq, J = 13.0, 7.1, 6.7 Hz, 1H), 1.83 (dd, J = 11.8, 5.6 Hz, 1H), 1.47 (d, J = 7.1 Hz, 6H), 1.37 (t, J = 12.1 Hz, 1H), 0.85 (d, J = 6.3 Hz, 3H).
1604	559.30	2.18	
1605	566.32	1.68	
1606	500.00	1.47	1H NMR (400 MHz, DMSO) δ 9.64 (s, 1H), 8.12 (s, 2H), 7.77 (d, J = 7.3 Hz,

Cmpd No.	LC/MS M+1	LC/RT min	NMR
			1H), 7.30 (s, 1H), 7.10 (d, J = 7.6 Hz, 1H), 6.92 (s, 3H), 4.27 (s, 3H), 4.04 (s, 3H), 2.39 (s, 6H), 2.24 (s, 3H), 1.79 (s, 3H), 1.24 (s, 3H).
1607	408.20	0.98	
1608	563.60	2.13	
1609	537.20	1.97	
1610	567.50	1.91	
1611	570.30	2.02	
1612	570.30	2.02	
1613	528.60	1.85	
1614	562.60	1.61	
1615	535.50	1.75	
1616	605.00	1.67	1H NMR (400 MHz, MeOD) δ 8.16 (d, J = 8.2 Hz, 1H), 8.00 (s, 1H), 7.88 (s, 1H), 7.27 (s, 1H), 7.12 (d, J = 6.8 Hz, 1H), 6.99 (s, 2H), 6.89 (d, J = 8.2 Hz, 1H), 4.47 (s, 2H), 4.26 (d, J = 33.1 Hz, 2H), 4.05 (s, 2H), 2.43 (s, 6H), 2.30 (s, 3H), 1.93 (s, 3H), 1.31 (s, 3H).
1617	514.30	1.89	
1618	539.00	1.66	1H NMR (400 MHz, DMSO) δ 12.54 (s, 1H), 8.27 (d, J = 5.4 Hz, 1H), 7.82 (d, J = 7.9 Hz, 1H), 7.64 (s, 1H), 7.58 (d, J = 6.7 Hz, 1H), 7.41 (s, 1H), 7.34 (d, J = 8.0 Hz, 1H), 7.21 (d, J = 7.2 Hz, 1H), 6.70 (d, J = 8.3 Hz, 1H), 4.09 (d, J = 6.7 Hz, 2H), 2.61 (s, 2H), 2.23 (s, 1H), 2.07 (s, 1H), 1.88 (s, 1H), 1.60 (d, J = 7.4 Hz, 6H), 1.43 (s, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.88 (d, J = 6.3 Hz, 3H).
1619	533.30	2.40	
1620	528.50	1.77	1H NMR (400 MHz, Chloroform-d) δ 13.87 (broad s, 1H), 8.29 (d, J = 8.0 Hz, 1H) 8.21 (d, J = 2.9 Hz, 1H), 7.65 - 7.58 (m, 2H), 7.52 (d, J = 8.5 Hz, 1H), 6.65 (p, J = 3.8 Hz, 1H), 5.95 (d, J = 2.8 Hz, 1H), 4.80 (broad s, 2H), 4.02 (d, J = 6.6 Hz, 2H), 3.52 - 3.41 (m, 1H), 3.15 (dd, J = 10.5, 8.1 Hz, 1H), 2.69 - 2.50 (m, 1H), 2.18 - 2.02 (m, 2H), 1.72 (dd, J = 12.3, 9.7 Hz, 1H), 1.45 (s, 3H), 1.39 (s, 3H), 1.19 (d, J = 6.6 Hz, 3H), 1.02 (d, J = 6.7 Hz, 6H).
1621	504.23	1.97	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1622	573.14	2.40	
1623	525.30	1.89	1H NMR (400 MHz, DMSO-d6) δ 7.90 (d, J = 7.8 Hz, 1H), 7.82 (d, J = 8.5 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.22 (d, J = 7.3 Hz, 1H), 7.05 (d, J = 7.9 Hz, 1H), 6.70 (t, J = 8.6 Hz, 2H), 5.31 (p, J = 6.2 Hz, 1H), 3.95 (q, J = 5.9 Hz, 2H), 2.04 - 1.93 (m, 5H), 1.71 - 1.57 (m, 3H), 1.31 (d, J = 6.2 Hz, 6H), 1.07 (d, J = 6.2 Hz, 6H).
1624	571.40	1.55	
1625	566.32	1.67	
1626	516.60	1.58	
1627	430.30	1.19	
1628	524.40	1.65	
1629	557.03	2.48	
1630	535.30	2.43	1H NMR (400 MHz, DMSO-d6) δ 12.08 (s, 1H), 8.01 (d, J = 7.9 Hz, 1H), 7.63 (dd, J = 8.4, 7.3 Hz, 1H), 7.24 - 7.16 (m, 2H), 6.91 (s, 2H), 6.70 (d, J = 8.3 Hz, 1H), 6.57 - 6.44 (m, 3H), 2.38 - 2.30 (m, 1H), 2.25 (s, 3H), 2.19 - 2.05 (m, 2H), 1.99 (s, 6H), 1.88 - 1.76 (m, 2H), 1.44 (h, J = 6.9 Hz, 1H), 1.25 - 1.09 (m, 2H), 0.85 (dd, J = 6.8, 4.7 Hz, 6H).
1631	537.30	1.80	
1632	474.30	1.18	
1633	540.60	1.91	
1634	515.30	0.95	
1635	472.30	1.79	1H NMR (400 MHz, DMSO-d6) δ 12.70 (s, 1H), 7.87 (d, J = 7.7 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.21 (dd, J = 7.3, 0.7 Hz, 1H), 7.00 (d, J = 7.7 Hz, 1H), 6.70 (dd, J = 8.5, 0.8 Hz, 1H), 6.56 (s, 2H), 2.60 (s, 2H), 1.75 (s, 2H), 1.55 (s, 6H), 0.97 (s, 6H).
1636	500.20	1.27	1H NMR (400 MHz, DMSO-d6) δ 12.45 (s, 1H), 10.50 - 9.90 (s, 1H), 7.78 (d, J = 8.0 Hz, 1H), 7.63 (t, J = 7.9 Hz, 1H), 7.31 (s, 1H), 7.26 - 7.17 (m, 2H), 7.14 (d, J = 8.1 Hz, 1H), 6.69 (d, J = 8.4 Hz, 1H), 6.64 (d, J = 10.5 Hz, 1H), 2.65 - 2.53 (m, 2H), 2.29 - 2.11 (m, 1H), 1.88 (dd, J = 11.9, 5.6 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 1.60 (s, 3H), 1.59 (s, 3H), 1.42 (t, J = 12.0 Hz, 1H), 0.88 (d, J = 6.2 Hz, 3H).
1637	539.40	2.47	1H NMR (400 MHz, DMSO-d6) δ 7.65 - 7.56 (m, 2H), 7.48 (dd, J = 8.2, 7.3 Hz, 1H), 7.39 (d, J = 7.7 Hz, 1H), 6.98 (dd, J = 7.4, 0.8 Hz, 1H), 6.60 - 6.46 (m, 2H), 6.28 (s, 2H), 5.38 (p, J = 6.2 Hz, 1H), 3.00 - 2.83 (m, 2H), 2.30 (s, 3H), 2.09 (dq, J = 11.6, 6.0 Hz, 1H), 1.81 (dd, J = 12.0, 5.7 Hz, 1H), 1.65 (s, 3H), 1.57 (s, 3H), 1.41 - 1.29 (m, 8H), 0.78 (d, J = 6.3 Hz, 3H).
1638	472.30	1.82	
1639	571.36	2.49	
1640	534.30	2.08	1H NMR (400 MHz, DMSO-d6) δ 12.26 (s, 1H), 7.73 (d, J = 8.0 Hz, 1H), 7.53 - 7.48 (m, 1H), 7.43 (dd, J = 9.9, 1.9 Hz, 1H), 7.29 (d, J = 8.0 Hz, 1H), 6.91 (dt, J = 10.7, 2.3 Hz, 1H), 4.47 (s, 1H), 4.18 - 3.97 (m, 2H), 3.92 - 3.82 (m, 3H), 3.77 - 3.68 (m, 1H), 3.09 (q, J = 10.6 Hz, 1H), 2.94 - 2.82 (m, 1H), 2.40 - 2.18 (m, 3H), 2.06 (dt, J = 13.3, 6.7 Hz, 1H), 2.01 - 1.93 (m, 1H), 1.68 (d, J = 1.8 Hz, 3H), 1.63 (s, 3H), 1.56 (td, J = 12.3, 3.7 Hz, 1H), 1.05 (dd, J = 6.4, 3.2 Hz, 3H), 0.99 (d, J = 6.7 Hz, 6H).
1641	590.00	3.06	1H NMR (400 MHz, DMSO-d6) δ 12.59 (s, 1H), 7.80 (s, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.27 - 7.18 (m, 1H), 7.10 - 6.99 (m, 2H), 6.93 (dt, J = 11.0, 2.3 Hz, 1H), 6.70 (dd, J = 8.4, 0.8 Hz, 1H), 6.54 (s, 2H), 3.80 (d, J = 6.6 Hz, 2H), 2.65 - 2.53 (m, 2H), 2.22 (dt, J = 11.5, 5.9 Hz, 1H), 2.08 - 1.96 (m, 1H), 1.84 (dd, J = 12.0, 5.6 Hz, 1H), 1.57 - 1.33 (m, 7H), 0.97 (d, J = 6.7 Hz, 6H), 0.86 (d, J = 6.3 Hz, 3H).
1642	510.00	1.52	1H NMR (400 MHz, DMSO) δ 10.48 (s, 1H), 7.37 (d, J = 4.2 Hz, 1H), 7.35 (s, 1H), 7.32 (d, J = 8.7 Hz, 1H), 7.28 (d, J = 7.9 Hz, 1H), 7.05 (d, J = 7.6 Hz, 1H), 6.92 (d, J = 7.8 Hz, 3H), 4.24 (s, 3H), 4.02 (s, 3H), 2.39 (s, 6H), 2.24 (s, 3H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1.78 (s, 3H), 1.24 (s, 3H).
1643	494.60	1.06	
1644	529.20	2.30	1H NMR (400 MHz, DMSO) δ 7.84 (d, J = 7.7 Hz, 1H), 7.50 (t, J = 5.7 Hz, 1H), 7.47 - 7.36 (m, 3H), 7.06 (dd, J = 7.3, 0.7 Hz, 1H), 6.84 (dt, J = 10.7, 2.2 Hz, 1H), 6.45 (dd, J = 8.2, 0.7 Hz, 1H), 6.07 (s, 2H), 3.84 (d, J = 6.6 Hz, 2H), 2.44 - 2.31 (m, 2H), 2.24 (dd, J = 10.0, 6.9 Hz, 2H), 2.06 (dt, J = 13.3, 6.6 Hz, 1H), 1.83 - 1.71 (m, 2H), 1.68 (s, 3H), 1.01 (t, J = 8.3 Hz, 6H).
1645	567.50	1.78	
1646	552.32	1.39	
1647	584.42	1.36	
1648	549.60	2.10	
1649	536.50	1.49	
1650	549.30	2.44	
1651	512.30	1.86	
1652	598.00	2.83	1H NMR (400 MHz, DMSO-d6) δ 12.68 (s, 1H), 10.87 (s, 1H), 8.35 (d, J = 8.4 Hz, 1H), 8.17 - 8.05 (m, 1H), 7.88 - 7.80 (m, 2H), 7.51 - 7.45 (m, 1H), 7.42 (ddd, J = 10.2, 2.4, 1.3 Hz, 1H), 7.30 (d, J = 8.0 Hz, 1H), 6.90 (dt, J = 10.7, 2.3 Hz, 1H), 3.83 (d, J = 6.6 Hz, 2H), 2.60 (dd, J = 10.2, 7.2 Hz, 1H), 2.52 (d, J = 5.3 Hz, 1H), 2.14 (s, 4H), 2.07 - 2.00 (m, 1H), 1.87 (dd, J = 11.8, 5.6 Hz, 1H), 1.59 (d, J = 6.0 Hz, 6H), 1.41 (t, J = 12.1 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.77 (d, J = 6.3 Hz, 3H).
1653	455.40	2.05	1H NMR (400 MHz, DMSO-d6) δ 12.04 (s, 1H), 7.65 - 7.59 (m, 2H), 6.95 (s, 1H), 6.76 (s, 2H), 6.67 - 6.61 (m, 2H), 6.19 (s, 2H), 5.45 (s, 2H), 2.17 (s, 3H), 2.10 (s, 6H), 1.13 (s, 9H).
1654	566.32	1.68	
1655	512.00	1.60	
1656	553.40	2.62	1H NMR (400 MHz, DMSO-d6) δ 12.47 (s, 1H), 7.83 (d, J = 8.0 Hz, 1H), 7.69 - 7.59 (m, 2H), 7.54 (d, J = 7.9 Hz, 1H), 7.20 (d, J = 7.1 Hz, 1H), 6.78 - 6.62 (m, 2H), 4.15 (dd, J = 6.6, 1.7 Hz, 2H), 2.70 - 2.56 (m, 2H), 2.34 (s, 3H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			2.29 - 2.16 (m, 1H), 2.07 (hept, J = 6.7 Hz, 1H), 1.88 (dd, J = 11.9, 5.6 Hz, 1H), 1.62 (d, J = 5.9 Hz, 6H), 1.43 (t, J = 12.0 Hz, 1H), 1.00 (d, J = 6.7 Hz, 6H), 0.89 (d, J = 6.2 Hz, 3H).
1657	554.20	2.05	1H NMR (400 MHz, DMSO) δ 12.47 (s, 1H), 7.76 (d, J = 7.9 Hz, 1H), 7.65 (t, J = 7.8 Hz, 1H), 7.41 (d, J = 12.0 Hz, 2H), 7.27 (d, J = 7.9 Hz, 1H), 7.22 (d, J = 7.3 Hz, 1H), 6.89 (d, J = 10.8 Hz, 1H), 6.71 (d, J = 8.3 Hz, 1H), 6.46 (s, 2H), 4.29 (s, 1H), 3.84 (d, J = 6.5 Hz, 2H), 2.86 (s, 2H), 2.03 (dd, J = 13.5, 6.8 Hz, 1H), 1.86 - 1.75 (m, 3H), 1.54 (ddd, J = 36.2, 22.1, 10.7 Hz, 6H), 0.99 (d, J = 6.7 Hz, 6H).
1658	472.20	1.71	
1659	553.50	1.69	
1660	525.30	1.38	
1661	545.09	2.22	1H NMR (400 MHz, DMSO) δ 11.70 (s, 1H), 8.03 (d, J = 7.9 Hz, 1H), 7.77 (d, J = 7.9 Hz, 1H), 7.69 - 7.61 (m, 1H), 7.53 (t, J = 5.3 Hz, 2H), 7.22 (d, J = 7.3 Hz, 1H), 7.02 - 6.93 (m, 1H), 6.71 (d, J = 8.4 Hz, 1H), 6.53 (s, 2H), 4.50 (qd, J = 11.1, 4.8 Hz, 2H), 4.26 (d, J = 5.2 Hz, 1H), 3.93 - 3.78 (m, 3H), 3.68 (dd, J = 13.9, 7.0 Hz, 1H), 2.13 - 1.98 (m, 2H), 1.98 - 1.74 (m, 3H), 1.00 (d, J = 6.7 Hz, 6H).
1662	526.00	1.21	1H NMR (400 MHz, MeOD) δ 8.53 (s, 1H), 8.30 (s, 1H), 7.96 (s, 1H), 7.15 (dd, J = 97.7, 47.9 Hz, 7H), 5.31 (s, 1H), 4.50 (d, J = 44.1 Hz, 3H), 4.11 (s, 2H), 1.83 (s, 3H), 1.40 (d, J = 29.2 Hz, 10H).
1663	553.00	1.81	1H NMR (400 MHz, DMSO) δ 12.45 (s, 1H), 7.91 - 7.78 (m, 3H), 7.64 (s, 1H), 7.57 (d, J = 7.9 Hz, 1H), 7.21 (d, J = 7.2 Hz, 1H), 6.88 (d, J = 7.8 Hz, 1H), 6.69 (d, J = 8.3 Hz, 1H), 4.10 (s, 2H), 2.62 (s, 2H), 2.23 (s, 1H), 1.88 (s, 1H), 1.62 (d, J = 6.0 Hz, 6H), 1.43 (s, 1H), 1.04 (s, 9H), 0.89 (d, J = 6.3 Hz, 3H).
1664	538.30	1.62	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1665	512.60	1.67	
1666	532.00	1.72	
1667	593.20	2.15	1H NMR (400 MHz, DMSO) δ 12.60 (s, 1H), 8.00 (d, J = 8.2 Hz, 1H), 7.91 (d, J = 8.1 Hz, 1H), 7.66 (dd, J = 8.4, 7.3 Hz, 1H), 7.54 - 7.47 (m, 2H), 7.26 - 7.16 (m, 1H), 6.95 (dt, J = 10.8, 2.3 Hz, 1H), 6.73 (d, J = 8.0 Hz, 1H), 6.56 (s, 2H), 5.81 (s, 1H), 3.86 (d, J = 6.5 Hz, 2H), 3.35 (s, 1H), 2.73 - 2.60 (m, 1H), 2.47 - 2.37 (m, 1H), 2.25 (dt, J = 9.1, 4.0 Hz, 1H), 2.08 - 1.98 (m, 3H), 1.49 (qd, J = 12.2, 5.5 Hz, 1H), 1.00 (d, J = 6.7 Hz, 6H).
1668	529.00	2.40	
1669	494.00	1.38	
1670	469.40	1.60	
1671	538.29	1.27	
1672	540.30	1.58	
1673	519.30	1.59	1H NMR (400 MHz, DMSO-d6) δ 14.11 (s, 1H), 7.73 - 7.46 (m, 2H), 7.16 (d, J = 7.3 Hz, 1H), 6.69 (dd, J = 8.5, 2.4 Hz, 1H), 3.79 (dt, J = 11.0, 6.2 Hz, 1H), 3.31 (td, J = 10.8, 3.7 Hz, 2H), 2.96 (t, J = 9.6 Hz, 1H), 2.41 - 2.27 (m, 2H), 1.99 (ddd, J = 23.1, 12.1, 6.4 Hz, 2H), 1.58 - 1.50 (m, 2H), 1.49 (s, 3H), 1.46 (s, 3H), 1.34 (s, 3H), 1.27 (s, 3H), 1.04 (dd, J = 6.5, 1.9 Hz, 6H).
1674	592.30	1.42	
1675	586.20	1.95	1H NMR (400 MHz, DMSO) δ 13.79 (s, 1H), 8.16 (d, J = 8.1 Hz, 1H), 8.01 (d, J = 8.2 Hz, 1H), 7.64 (dd, J = 8.4, 7.4 Hz, 1H), 7.58 - 7.46 (m, 2H), 7.24 (d, J = 7.0 Hz, 1H), 6.98 (dt, J = 10.7, 2.3 Hz, 1H), 6.71 (d, J = 8.1 Hz, 1H), 6.44 (s, 2H), 3.87 (d, J = 6.6 Hz, 2H), 3.56 (s, 2H), 3.14 (s, 2H), 2.06 (dq, J = 13.3, 6.6 Hz, 1H), 1.26 (d, J = 19.1 Hz, 12H), 1.00 (d, J = 6.7 Hz, 6H).
1676	498.00	1.43	1H NMR (400 MHz, DMSO) δ 9.60 (d, J = 113.0 Hz, 1H), 8.13 (s, 1H), 7.89 (s, 1H), 7.39 - 7.07 (m, 7H), 6.94 (s, 1H), 4.29 (d, J = 27.5 Hz, 3H), 4.04 (s, 3H), 2.44 (s, 6H), 1.81 (s, 3H), 1.26 (s, 3H).

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1677	458.30	1.49	1H NMR (400 MHz, DMSO) δ 12.32 (s, 1H), 7.80 - 7.55 (m, 2H), 7.18 (d, J = 7.3 Hz, 1H), 6.67 (d, J = 8.2 Hz, 1H), 6.47 (d, J = 7.8 Hz, 1H), 6.34 - 5.83 (m, 2H), 2.75 - 2.58 (m, 2H), 2.53 (d, J = 8.9 Hz, 2H), 2.17 (dd, J = 13.5, 7.1 Hz, 1H), 1.82 (dd, J = 11.9, 5.7 Hz, 1H), 1.59 - 1.49 (m, 8H), 1.39 (dd, J = 26.5, 14.4 Hz, 1H), 0.86 (d, J = 6.3 Hz, 3H), 0.69 (ddd, J = 12.8, 7.9, 4.9 Hz, 1H), 0.43 - 0.32 (m, 2H), 0.07 - 0.00 (m, 2H).
1678	491.50	1.74	
1679	519.20	1.78	
1680	573.30	2.25	
1681	557.00	1.66	1H NMR (400 MHz, DMSO) δ 10.84 (s, 1H), 7.62 (s, 1H), 7.26 (s, 1H), 7.13 - 7.01 (m, 2H), 6.92 (s, 2H), 6.80 (s, 2H), 6.69 (d, J = 8.3 Hz, 1H), 6.43 (s, 1H), 4.08 (d, J = 30.9 Hz, 4H), 3.89 (s, 2H), 3.79 (s, 2H), 2.01 (s, 1H), 1.50 (s, 3H), 1.28 (s, 3H), 0.96 (d, J = 6.6 Hz, 6H).
1682	542.60	1.97	
1683	444.30	1.32	
1684	557.40	1.56	
1685	529.29	2.29	
1686	593.37	2.47	
1687	594.20	1.66	
1688	594.20	1.64	
1689	593.33	2.44	
1690	594.20	1.67	
1691	580.35	1.74	
1692	569.33	2.06	
1693	580.32	1.71	
1694	579.34	2.39	
1695	583.30	1.48	
1696	613.50	1.62	
1697	474.40	1.35	
1698	474.40	1.35	
1699	472.30	1.21	
1700	569.40	1.65	
1701	480.21	1.62	
1702	539.30	1.56	
1703	583.50	1.74	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1704	513.41	2.61	
1705	526.16	1.40	
1706	496.31	1.14	
1707	526.18	1.24	
1708	520.32	2.43	
1709	514.30	1.65	1H NMR (400 MHz, DMSO-d6) δ 12.35 (s, 1H), 8.19 (d, J = 2.7 Hz, 1H), 7.85 (d, J = 8.3 Hz, 1H), 7.61 (t, J = 7.9 Hz, 1H), 7.18 (d, J = 7.2 Hz, 1H), 6.90 (d, J = 8.2 Hz, 1H), 6.66 (d, J = 8.3 Hz, 1H), 6.49 (broad s, 2H), 6.07 (d, J = 2.8 Hz, 1H), 4.86 (hept, J = 6.0 Hz, 1H), 2.57 (br m, 2H), 2.21 (br s, 1H), 1.87 (dd, J = 11.9, 5.7 Hz, 1H), 1.56 (s, 3H), 1.55 (s, 3H), 1.44 - 1.35 (m, 1H), 1.33 (dd, J = 6.1, 3.1 Hz, 6H), 0.87 (d, J = 6.1 Hz, 3H)
1710	541.30	1.64	1H NMR (400 MHz, DMSO-d6) δ 12.47 (d, J = 23.3 Hz, 2H), 8.22 (dd, J = 7.2, 2.1 Hz, 1H), 7.81 (s, 1H), 7.59 (d, J = 7.8 Hz, 1H), 6.44 (dd, J = 13.0, 7.1 Hz, 2H), 2.65 (d, J = 9.1 Hz, 2H), 2.58 (d, J = 12.3 Hz, 1H), 2.35 - 2.12 (m, 2H), 1.97 (d, J = 11.7 Hz, 3H), 1.92 - 1.78 (m, 2H), 1.64 - 1.47 (m, 8H), 1.46 - 1.32 (m, 3H), 0.89 (d, J = 6.2 Hz, 3H).
1711	446.18	2.53	
1712	482.24	0.73	
1713	494.57	1.08	
1714	506.20	1.15	
1715	613.50	1.58	1H NMR (400 MHz, DMSO) δ 8.63 (d, J = 2.4 Hz, 1H), 8.10 (d, J = 2.4 Hz, 1H), 7.57 (d, J = 7.6 Hz, 1H), 7.43 (t, J = 7.7 Hz, 1H), 6.99 (dd, J = 10.5, 7.4 Hz, 2H), 6.45 (d, J = 8.1 Hz, 1H), 6.09 (d, J = 4.6 Hz, 2H), 4.44 (dd, J = 5.7, 3.9 Hz, 2H), 3.77 (dd, J = 5.8, 4.0 Hz, 2H), 3.59 (dd, J = 5.8, 3.8 Hz, 2H), 3.49 (dd, J = 5.9, 3.7 Hz, 2H), 3.42 (q, J = 7.0 Hz, 2H), 3.06 - 2.93 (m, 2H), 2.20 (s, 3H), 2.16 - 2.04 (m, 2H), 1.81 (dd, J = 11.5, 5.6 Hz, 1H), 1.61 (s, 3H), 1.57 (s, 3H), 1.37 (t, J = 11.9 Hz, 1H), 1.08 (t, J = 6.9 Hz, 3H), 0.85 (d, J = 6.3

Cmpd No.	LC/MS M+1	LC/RT min	NMR
			Hz, 4H).
1716	494.31	1.52	
1717	524.52	2.49	
1718	447.08	2.71	
1719	511.31	2.49	1H NMR (400 MHz, DMSO) δ 12.91 (s, 1H), 8.20 - 8.14 (m, 2H), 7.92 - 7.89 (d, 1H), 7.53 - 7.50 (dd, 1H), 7.33 - 7.30 (d, 1H), 6.68 (s, 1H), 6.59 - 6.55 (m, 3H), 3.70(s, 3H), 2.26 (s, 3H), 2.26 - 2.17 (m, 1H), 1.89 - 1.82 (m, 1 H), 1.50 (s, 3H), 1.47 (s, 3H), 1.45 - 1.35 (m, 1H), 0.88 - 0.86 (d, 3H)
1720	492.10	2.12	
1721	534.18	1.98	
1722	508.31	2.64	
1723	586.30	2.00	1H NMR (400 MHz, DMSO) δ 13.01 - 11.30 (m, 1H), 8.27 (d, J = 8.2 Hz, 1H), 7.66 (t, J = 7.8 Hz, 1H), 7.56 - 7.42 (m, 2H), 7.22 (t, J = 8.2 Hz, 2H), 6.91 (d, J = 10.6 Hz, 1H), 6.71 (d, J = 8.3 Hz, 1H), 6.58 (s, 2H), 3.84 (d, J = 6.4 Hz, 2H), 3.65 (s, 2H), 2.13 - 1.99 (m, 1H), 1.37 (s, 2H), 1.13 (s, 6H), 1.00 (d, J = 6.6 Hz, 6H), 0.46 (s, 2H), 0.28 (s, 2H).
1724	556.20	2.02	
1725	458.30	1.36	
1726	522.35	1.65	
1727	525.30	2.35	1H NMR (400 MHz, DMSO) δ 8.63 (s, 3H), 8.09 (s, 3H), 7.61 - 7.53 (m, 4H), 7.48 - 7.37 (m, 4H), 6.99 (dq, J = 13.5, 6.1 Hz, 7H), 6.52 - 6.38 (m, 4H), 6.17 - 6.02 (m, 6H), 4.37 (q, J = 7.1 Hz, 6H), 3.07 - 2.88 (m, 7H), 2.19 (s, 8H), 2.17 - 2.04 (m, 1H), 1.81 (q, J = 8.4, 7.7 Hz, 3H), 1.61 (s, 8H), 1.57 (s, 8H), 1.41 - 1.30 (m, 11H), 0.84 (d, J = 6.0 Hz, 9H).
1728			1H NMR (400 MHz, DMSO) δ 8.61 (d, J = 2.3 Hz, 1H), 8.11 - 8.01 (m, 1H), 7.57 (d, J = 7.7 Hz, 1H), 7.46 (t, J = 7.8 Hz, 1H), 6.97 (t, J = 6.9 Hz, 2H), 6.48 (d, J = 8.3 Hz, 1H), 6.20 (s, 2H), 5.16 (h, J = 6.2 Hz, 1H), 2.98 (t, J = 10.6 Hz, 1H), 2.92 - 2.82 (m, 1H), 2.17 (s, 3H), 2.10 (t, J = 8.0 Hz, 1H), 1.81 (dd, J = 11.4, 5.6 Hz, 1H), 1.75 - 1.64 (m, 2H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1.61 (s, 3H), 1.56 (s, 3H), 1.36 (t, J = 11.9 Hz, 1H), 1.28 (d, J = 6.1 Hz, 3H), 0.93 (t, J = 7.4 Hz, 3H), 0.81 (d, J = 6.3 Hz, 3H).
1729	570.20	2.07	
1730	534.30	1.91	
1731	496.30	1.29	1H NMR (400 MHz, DMSO-d6) δ 11.79 (s, 1H), 8.26 (dd, J = 4.9, 1.8 Hz, 1H), 8.08 (dd, J = 7.8, 1.8 Hz, 1H), 7.81 (d, J = 8.2 Hz, 1H), 7.76 (d, J = 8.3 Hz, 1H), 7.39 (d, J = 8.2 Hz, 1H), 6.80 (dd, J = 7.8, 4.8 Hz, 1H), 6.73 (d, J = 7.4 Hz, 2H), 2.70 - 2.53 (m, 2H), 2.27 (s, 3H), 2.23 - 2.12 (m, 1H), 1.89 (dd, J = 12.1, 5.5 Hz, 1H), 1.51 (d, J = 2.0 Hz, 6H), 1.42 (t, J = 12.1 Hz, 1H), 0.77 (d, J = 6.3 Hz, 3H).
1732	513.00	1.12	1H NMR (400 MHz, DMSO-d6) δ 12.80 (s, 1H), 8.30 (d, J = 2.8 Hz, 1H), 8.26 (dd, J = 4.7, 1.9 Hz, 1H), 8.03 (dd, J = 7.9, 1.9 Hz, 1H), 7.84 (d, J = 8.2 Hz, 1H), 6.97 (d, J = 8.2 Hz, 1H), 6.76 (dd, J = 7.8, 4.8 Hz, 2H), 6.10 (d, J = 2.7 Hz, 1H), 3.98 (d, J = 6.6 Hz, 2H), 3.92 (d, J = 5.4 Hz, 2H), 2.05 (dq, J = 13.4, 6.8 Hz, 1H), 1.94 - 1.84 (m, 2H), 1.65 - 1.55 (m, 2H), 0.97 (dd, J = 7.9, 6.4 Hz, 12H).
1733	527.22	1.46	
1734	474.09	2.57	
1735	496.20	1.53	
1736	486.33	1.49	
1737	568.20	2.00	
1738	496.31	1.14	
1739	552.40	1.70	1H NMR (400 MHz, DMSO) δ 12.57 - 12.26 (m, 2H), 7.69 (s, 0H), 7.57 - 7.36 (m, 3H), 7.19 - 7.02 (m, 7H), 6.93 (s, 3H), 6.74 - 5.97 (m, 4H), 3.52 (q, J = 6.9 Hz, 12H), 2.35 (s, 9H), 2.18 (s, 4H), 1.85 (dd, J = 11.6, 5.7 Hz, 3H), 1.62 (s, 9H), 1.59 (s, 8H), 1.40 (t, J = 12.1 Hz, 3H), 1.13 (t, J = 6.9 Hz, 18H), 0.87 (d, J = 6.2 Hz, 9H), 3.05 - 2.58 (m, 6H).
1740	538.22	1.37	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1741	553.40	2.75	1H NMR (400 MHz, DMSO-d6) δ 8.63 (d, J = 2.2 Hz, 1H), 8.10 (d, J = 2.0 Hz, 1H), 7.61 (s, 1H), 7.49 (s, 1H), 7.02 (s, 2H), 6.51 (d, J = 8.7 Hz, 1H), 6.23 (s, 2H), 4.10 (d, J = 6.6 Hz, 2H), 2.87 (s, 2H), 2.21 (s, 3H), 2.19 - 1.98 (m, 2H), 1.82 (dd, J = 11.7, 5.6 Hz, 1H), 1.61 (s, 3H), 1.57 (s, 3H), 1.37 (t, J = 12.0 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.83 (d, J = 6.4 Hz, 3H).
1742	544.30	1.32	1H NMR (400 MHz, DMSO-d6) δ 12.52 (s, 1H), 8.37 (d, J = 2.8 Hz, 0H), 8.26 (dd, J = 4.8, 1.9 Hz, 1H), 8.20 (d, J = 2.8 Hz, 1H), 8.06 (dd, J = 7.8, 1.8 Hz, 1H), 7.85 (d, J = 8.2 Hz, 1H), 6.94 (d, J = 8.3 Hz, 1H), 6.79 (dd, J = 7.8, 4.9 Hz, 1H), 6.14 (d, J = 2.8 Hz, 1H), 4.35 - 4.28 (m, 2H), 3.74 - 3.67 (m, 2H), 3.50 (q, J = 7.0 Hz, 2H), 2.46 (d, J = 8.8 Hz, 2H), 2.22 - 2.03 (m, 1H), 1.85 (dd, J = 11.9, 5.6 Hz, 1H), 1.55 (s, 3H), 1.53 (s, 3H), 1.44 - 1.33 (m, 1H), 1.13 (t, J = 7.0 Hz, 3H), 0.74 (d, J = 6.3 Hz, 3H).
1743	543.30	1.95	1H NMR (400 MHz, DMSO-d6) δ 12.51 (broad d, 1H), 12.47 (s, 1H), 8.22 (d, J = 7.2 Hz, 1H), 8.00 (s, 1H), 7.81 (br t, J = 6.3 Hz, 1H), 7.76 (d, J = 8.4 Hz, 1H), 6.84 (d, J = 8.2 Hz, 1H), 6.44 (t, J = 6.8 Hz, 1H), 4.01 (d, J = 6.7 Hz, 2H), 2.69 (d, J = 9.2 Hz, 2H), 2.21 (br m, 1H), 2.07 (hep, J = 6.6 Hz, 1H), 1.95 (s, 3H), 1.88 (dd, J = 11.9, 5.6 Hz, 1H), 1.57 (s, 3H), 1.55 (s, 3H), 1.42 (t, J = 12.1 Hz, 1H), 0.98 (d, J = 6.7 Hz, 5H), 0.89 (d, J = 6.3 Hz, 3H).
1744	539.53	2.10	
1745	526.21	1.18	
1746	568.50	1.99	1H NMR (400 MHz, DMSO) δ 12.43 (s, 1H), 7.78 (d, J = 8.0 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.50 (d, J = 1.6 Hz, 1H), 7.43 (dd, J = 9.7, 1.7 Hz, 1H), 7.29 (d, J = 8.0 Hz, 1H), 7.19 (dd, J = 7.3, 0.6 Hz, 1H), 6.90 (dt, J = 10.7, 2.3 Hz, 1H), 6.71 (dd, J = 8.4, 0.6 Hz, 1H), 6.49 (s, 2H), 3.84 (d, J = 6.6 Hz, 2H),

Cmpd. No.	LC/MS M+1	LC/RT, min	NMR
			2.69 (s, 2H), 2.12 - 1.98 (m, 1H), 1.77 (s, 2H), 1.69 (s, 6H), 0.99 (d, J = 6.7 Hz, 6H), 0.49 (d, J = 12.5 Hz, 4H).
1747	510.16	1.47	
1748	511.20	1.34	1H NMR (400 MHz, DMSO) δ 7.59 (d, J = 7.6 Hz, 1H), 7.44 (t, J = 7.8 Hz, 1H), 7.40 (s, 1H), 7.16 (s, 1H), 7.11 (d, J = 7.7 Hz, 1H), 7.00 (d, J = 7.3 Hz, 1H), 6.46 (d, J = 8.2 Hz, 1H), 6.13 (s, 2H), 3.86 (s, 3H), 3.01 (t, J = 9.2 Hz, 2H), 2.44 (s, 3H), 2.12 (s, 1H), 1.82 (dd, J = 11.6, 5.6 Hz, 1H), 1.61 (s, 3H), 1.55 (s, 3H), 1.37 (t, J = 12.0 Hz, 1H), 0.84 (d, J = 6.3 Hz, 3H).
1749	486.40	2.76	
1750	556.00	2.08	1H NMR (400 MHz, DMSO) δ 12.30 (s, 1H), 10.56 (s, 1H), 7.83 (d, J = 8.7 Hz, 2H), 7.67 (d, J = 7.9 Hz, 1H), 7.47 (s, 1H), 7.39 (d, J = 10.0 Hz, 1H), 7.24 (d, J = 8.0 Hz, 1H), 6.94 (d, J = 8.7 Hz, 2H), 6.88 (d, J = 10.7 Hz, 1H), 3.83 (d, J = 6.6 Hz, 2H), 2.45 (d, J = 10.4 Hz, 1H), 2.36 (s, 1H), 2.07 (s, 3H), 1.83 (s, 1H), 1.57 (d, J = 18.9 Hz, 6H), 1.41 (d, J = 12.1 Hz, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.71 (d, J = 6.3 Hz, 3H).
1751	517.32	2.45	
1752	527.30	1.38	
1753	429.07	1.97	
1754	554.20	1.98	1H NMR (400 MHz, DMSO) δ 12.57 (s, 1H), 7.78 (d, J = 8.0 Hz, 1H), 7.64 (dt, J = 12.1, 6.0 Hz, 1H), 7.42 (d, J = 1.7 Hz, 1H), 7.35 (t, J = 9.1 Hz, 2H), 7.22 (d, J = 6.9 Hz, 1H), 6.92 - 6.85 (m, 1H), 6.71 (d, J = 8.1 Hz, 1H), 6.50 (s, 2H), 3.84 (d, J = 6.6 Hz, 2H), 2.94 (dd, J = 10.2, 6.9 Hz, 1H), 2.60 (dd, J = 10.3, 7.3 Hz, 1H), 2.26 (dd, J = 14.1, 7.0 Hz, 1H), 2.00 (ddd, J = 14.6, 11.5, 6.1 Hz, 3H), 1.87 (dd, J = 11.9, 6.4 Hz, 1H), 1.66 (dd, J = 11.8, 8.2 Hz, 1H), 0.99 (t, J = 5.2 Hz, 10H), 0.68 - 0.38 (m, 2H).
1755	534.42	2.82	
1756	522.42	2.92	
1757	509.21	1.36	1H NMR (DMSO(d6)) δ 12.85 (s, 1H),

Cmpd No.	LC/MS M+1	LC/RT min	NMR
			10.07 (s, 1H), 8.16 - 8.14 (dd, 1H), 8.00 - 7.97 (d, 2H), 7.89 (t, J = 7.80, 1H), 7.75 - 7.72 (dd, 1H), 7.69 - 7.66 (d, 2H), 7.54 - 7.51 (d, 1H), 7.17 - 7.14 (d, 1H), 6.60 - 6.57 (dd, 1H), 2.44 - 2.40 (m, 1H), 1.92 - 1.84 (m, 1H), 1.77 - 1.70 (m, 1H), 1.47 (s, 3H), 1.43 (s, 3H), 1.29 (t, J = 11.86, 1H), 0.70 - 0.68 (d, 3H)
1758	542.60	1.63	
1759	590.40	2.09	
1760	529.40	1.68	1H NMR (400 MHz, Methanol-d4) δ 8.38 (dd, J = 7.2, 2.3 Hz, 1H), 7.99 (d, J = 1.1 Hz, 1H), 7.87 (d, J = 8.2 Hz, 1H), 7.71 (s, 1H), 7.67 (d, J = 4.8 Hz, 1H), 7.50 (s, 1H), 7.06 (d, J = 8.0 Hz, 1H), 6.55 - 6.47 (m, 1H), 4.54 (s, 1H), 3.74 - 3.70 (m, 2H), 2.92 (s, 2H), 2.31 (s, 1H), 2.04 (dd, J = 13.2, 6.7 Hz, 1H), 1.92 (dd, J = 11.8, 6.0 Hz, 1H), 1.64 (s, 6H), 1.01 (dd, J = 6.8, 2.0 Hz, 6H), 0.97 (d, J = 6.3 Hz, 3H).
1761	527.22	2.10	
1762	670.00	3.60	1H NMR (400 MHz, DMSO-d6) δ 12.23 (s, 1H), 8.13 (d, J = 7.8 Hz, 1H), 7.81 (d, J = 7.9 Hz, 1H), 7.57 - 7.46 (m, 2H), 7.45 - 7.23 (m, 5H), 7.23 - 7.04 (m, 5H), 6.83 (dt, J = 10.9, 2.3 Hz, 1H), 6.33 (s, 2H), 5.24 (s, 2H), 3.61 (d, J = 6.6 Hz, 2H), 2.05 (s, 6H), 1.98 (dt, J = 13.4, 6.7 Hz, 1H), 0.96 (d, J = 6.6 Hz, 6H).
1763	542.29	1.64	
1764	475.29	1.95	
1765	447.30	1.37	1H NMR (400 MHz, DMSO-d6) δ 12.46 (s, 2H), 8.22 (dd, J = 7.3, 2.3 Hz, 1H), 7.81 (s, 1H), 7.62 (d, J = 7.9 Hz, 1H), 6.61 (d, J = 8.1 Hz, 1H), 6.43 (t, J = 6.8 Hz, 1H), 2.67 (d, J = 9.6 Hz, 2H), 2.22 (s, 1H), 1.86 (d, J = 5.8 Hz, 1H), 1.54 (d, J = 11.1 Hz, 6H), 1.27 (d, J = 5.0 Hz, 10H), 0.89 (d, J = 6.2 Hz, 3H).
1766	480.28	1.36	
1767	487.30	2.90	
1768	549.50	2.14	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1769	447.98	1.58	
1770	516.41	2.28	
1771	494.50	1.71	1H NMR (400 MHz, DMSO-d6) δ 12.41 (s, 1H), 7.77 (d, J = 8.0 Hz, 1H), 7.69 - 7.57 (m, 3H), 7.20 (dd, J = 7.3, 0.7 Hz, 1H), 7.16 (d, J = 8.1 Hz, 1H), 7.09 - 7.04 (m, 1H), 6.68 (dd, J = 8.4, 0.8 Hz, 3H), 2.60 (d, J = 9.0 Hz, 2H), 2.33 (s, 6H), 2.22 (s, 1H), 1.87 (dd, J = 11.8, 5.6 Hz, 1H), 1.60 (d, J = 3.6 Hz, 6H), 1.42 (t, J = 12.1 Hz, 1H), 0.88 (d, J = 6.3 Hz, 3H).
1772	466.29	1.96	
1773	559.34	2.56	
1774	525.50	1.68	1H NMR (400 MHz, DMSO-d6) δ 12.52 (d, J = 8.6 Hz, 2H), 8.24 (d, J = 7.3 Hz, 1H), 7.98 (d, J = 8.4 Hz, 2H), 7.82 (t, J = 5.5 Hz, 1H), 7.71 (d, J = 8.0 Hz, 1H), 7.10 (d, J = 8.1 Hz, 1H), 7.01 (d, J = 8.4 Hz, 2H), 6.44 (t, J = 6.8 Hz, 1H), 4.69 (p, J = 6.0 Hz, 1H), 2.72 (d, J = 9.2 Hz, 2H), 2.23 (s, 1H), 1.88 (dd, J = 11.9, 5.6 Hz, 1H), 1.61 (s, 6H), 1.44 (t, J = 12.2 Hz, 1H), 1.29 (d, J = 6.0 Hz, 6H), 0.91 (d, J = 6.3 Hz, 3H).
1775	523.32	1.90	
1776	512.51	3.00	
1777	548.23	2.54	
1778	545.23	2.58	1H NMR DMSO(d6) δ 12.76 (s, 1H), 8.18 - 8.16 (dd, 1H), 8.06 (t, J = 7.80, 1H), 7.81 - 7.75 (m, 2H), 7.58 - 7.56 (m, 2H), 7.16 - 7.13 (d, 1H), 6.94 - 6.91 (d, 1H), 6.64 - 6.61 (dd, 1H), 5.30 (q, J = 11.76, 2H), 3.64 (s, 3H), 1.89 - 1.81 (m, 2H), 1.75 - 1.69 (m, 1H), 1.49 (s, 3H), 1.43 (s, 3H), 1.24 (t, J = 11.86, 1H), 0.58 - 0.56 (d, 3H)
1779	515.15	1.72	
1780	515.21	1.61	
1781	556.00	1.85	
1782	491.20	2.41	
1783	613.30	1.36	1H NMR (400 MHz, DMSO) δ 8.62 (s, 1H), 8.10 (s, 1H), 7.94 (dd, J = 4.8, 1.8 Hz, 1H), 7.84 (dd, J = 7.5, 1.8 Hz, 1H), 7.55 (d, J = 7.7 Hz, 1H), 6.97 (d, J = 7.7

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			Hz, 1H), 6.55 (dd, J = 7.5, 4.8 Hz, 1H), 6.29 (s, 2H), 4.48 - 4.39 (m, 2H), 3.81 - 3.74 (m, 2H), 3.63 - 3.57 (m, 2H), 3.51 - 3.46 (m, 2H), 3.42 (q, J = 7.0 Hz, 2H), 2.94 (p, J = 10.5 Hz, 2H), 2.20 (s, 3H), 2.08 (d, J = 10.6 Hz, 1H), 1.80 (dd, J = 11.4, 5.7 Hz, 1H), 1.58 (d, J = 12.5 Hz, 6H), 1.35 (t, J = 12.0 Hz, 1H), 1.08 (t, J = 7.0 Hz, 3H), 0.78 (d, J = 6.3 Hz, 3H).
1784	501.30	1.48	
1785	550.30	1.71	1H NMR (400 MHz, DMSO-d6) δ 12.45 (s, 1H), 8.19 - 8.11 (m, 2H), 7.81 (d, J = 8.0 Hz, 1H), 7.63 (t, J = 7.9 Hz, 1H), 7.50 (d, J = 8.3 Hz, 2H), 7.22 (t, J = 7.5 Hz, 2H), 6.69 (d, J = 8.4 Hz, 1H), 6.50 (s, 2H), 2.65 - 2.56 (m, 2H), 2.23 (d, J = 9.1 Hz, 1H), 1.88 (dd, J = 12.0, 5.6 Hz, 1H), 1.60 (d, J = 4.6 Hz, 6H), 1.43 (t, J = 12.0 Hz, 1H), 0.88 (d, J = 6.2 Hz, 3H).
1786	527.36	0.99	
1787	671.60	1.94	1H NMR (400 MHz, DMSO) δ 8.63 (d, J = 2.3 Hz, 1H), 8.10 (d, J = 2.3 Hz, 1H), 7.57 (d, J = 7.7 Hz, 1H), 7.48 - 7.39 (m, 1H), 6.98 (dd, J = 7.5, 3.9 Hz, 2H), 6.47 (d, J = 8.3 Hz, 1H), 6.16 (s, 2H), 4.48 - 4.42 (m, 2H), 3.80 - 3.74 (m, 2H), 3.62 - 3.58 (m, 2H), 3.53 (ddd, J = 7.5, 5.3, 2.7 Hz, 3H), 3.48 (dd, J = 5.7, 3.7 Hz, 2H), 3.45 - 3.42 (m, 2H), 3.04 - 2.88 (m, 2H), 2.20 (s, 3H), 2.11 (dd, J = 11.6, 5.8 Hz, 1H), 1.81 (dd, J = 11.5, 5.7 Hz, 1H), 1.61 (s, 3H), 1.56 (s, 3H), 1.36 (t, J = 11.9 Hz, 1H), 1.04 (d, J = 6.1 Hz, 6H), 0.82 (d, J = 6.3 Hz, 3H).
1788	610.30	2.23	
1789	561.60	2.32	
1790	484.27	0.92	
1791	501.40	1.61	1H NMR (400 MHz, DMSO-d6) δ 12.63 (s, 1H), 12.54 (d, J = 6.0 Hz, 1H), 8.25 (dd, J = 7.3, 2.1 Hz, 1H), 8.08 (t, J = 1.9 Hz, 1H), 8.01 (dt, J = 7.4, 1.7 Hz, 1H), 7.83 (dd, J = 6.7, 4.6 Hz, 1H), 7.76 (d, J = 8.0 Hz, 1H), 7.57 -

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			7.47 (m, 2H), 7.25 (d, J = 8.0 Hz, 1H), 6.45 (t, J = 6.8 Hz, 1H), 2.82 - 2.64 (m, 2H), 2.31 - 2.17 (m, 0H), 1.89 (dd, J = 11.9, 5.6 Hz, 1H), 1.61 (d, J = 1.7 Hz, 6H), 1.45 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).
1792	496.27	1.48	
1793	514.30	1.25	
1794	430.27	1.70	1H NMR (400 MHz, DMSO-d6) δ 8.18 - 8.16 (dd, 1H), 7.73 - 7.70 (dd, 1H), 7.865 (t, J = 7.80, 1H), 7.20 - 7.17 (d, 1H), 6.74 - 6.71 (d, 1H), 6.69 - 6.69 (dd, 1H), 2.94 - 2.84 (m, 21H), 2.72 - 2.65 (m, 1H), 1.99 - 1.92 (m, 1H), 1.82 - 1.75 (m, 1H), 1.67 - 1.54 (m, 1H), 1.46 (s, 3H), 1.26 (s, 3H), 1.21 - 1.06 (m, 2H), 0.74 - 0.65 (m, 1H), 0.51 - 0.34 (m, 2H), 0.09 - 0.03 (m, 2H)
1795	548.20	2.27	
1796	520.32	2.35	
1797	581.20	1.77	1H NMR (400 MHz, DMSO-d6) δ 12.40 (s, 1H), 8.15 (d, J = 7.8 Hz, 1H), 7.83 (d, J = 7.9 Hz, 1H), 7.66 (t, J = 7.8 Hz, 1H), 7.24 (d, J = 7.2 Hz, 1H), 7.21 - 7.13 (m, 4H), 7.13 - 7.07 (m, 1H), 6.83 (dt, J = 10.6, 2.3 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 6.54 (s, 2H), 4.56 (s, 1H), 3.83 (dd, J = 9.4, 6.2 Hz, 1H), 3.69 (dd, J = 9.5, 6.2 Hz, 1H), 3.46 - 3.35 (m, 3H), 2.06 (s, 6H), 1.95 (dp, J = 12.7, 6.3 Hz, 1H), 0.95 (d, J = 6.7 Hz, 3H).
1798	499.31	2.48	1H NMR (400 MHz, DMSO-d6) δ 12.81 (s, 1H), 8.19 - 8.17 (dd, 1H), 8.06 (t, J = 7.91, 1H), 7.81 - 7.73 (m, 3H), 7.53 - 7.50 (dd, 1H), 7.38 - 7.21 (m, 3H), 6.64 - 6.61 (dd, 1H), 5.41 (s, 2H), 2.30 - 2.15 (m, 2H), 2.02 - 1.93 (m, 1H), 1.80 - 1.73 (m, 1H), 1.50 (s, 3H), 1.46 (s, 3H), 1.30 (t, J = 11.86, 1H), 0.65 - 0.63 (d, 3H)
1799	535.30	1.72	1H NMR (400 MHz, DMSO-d6) δ 12.66 (s, 1H), 12.54 (d, J = 6.2 Hz, 1H), 8.29 - 8.19 (m, 3H), 7.91 - 7.74 (m, 4H), 7.27 (d, J = 8.0 Hz, 1H), 6.45

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(t, J = 6.8 Hz, 1H), 2.83 - 2.64 (m, 2H), 2.31 - 2.18 (m, 1H), 1.90 (dd, J = 12.0, 5.5 Hz, 1H), 1.61 (d, J = 3.9 Hz, 6H), 1.46 (t, J = 12.1 Hz, 1H), 0.92 (d, J = 6.3 Hz, 3H).
1800	501.41	3.06	
1801	533.31	1.16	
1802	529.20	1.58	
1803	517.40	1.64	
1804	524.30	1.69	
1805	502.31	2.38	
1806	535.02	2.67	
1807	485.50	1.53	
1808	528.19	1.48	
1809	508.28	1.78	1H NMR (400 MHz, DMSO-d6) δ 12.46 (s, 1H), 8.50 (d, J = 2.5 Hz, 1H), 7.89 (d, J = 2.4 Hz, 1H), 7.63 (t, J = 8.3 Hz, 1H), 7.58 (d, J = 8.0 Hz, 2H), 7.32 (d, J = 8.3 Hz, 2H), 7.22 (d, J = 7.2 Hz, 1H), 6.69 (d, J = 8.3 Hz, 1H), 6.49 (s, 2H), 2.92 (p, J = 6.9 Hz, 1H), 2.56 (d, J = 8.7 Hz, 2H), 2.28 - 2.12 (m, 1H), 1.85 (dd, J = 11.8, 5.6 Hz, 1H), 1.55 (s, 3H), 1.53 (s, 3H), 1.40 (t, J = 12.0 Hz, 1H), 1.24 (d, J = 7.0 Hz, 6H), 0.87 (d, J = 6.2 Hz, 3H).
1810	494.60	1.63	1H NMR (400 MHz, DMSO-d6) δ 12.24 (s, 1H), 7.66 - 7.55 (m, 2H), 7.17 (d, J = 7.2 Hz, 1H), 6.66 (d, J = 8.4 Hz, 1H), 6.40 (s, 2H), 5.02 (td, J = 6.1, 3.1 Hz, 1H), 2.56 (d, J = 10.6 Hz, 1H), 2.20 (dt, J = 11.7, 6.0 Hz, 1H), 1.88 (ddt, J = 30.9, 11.8, 6.0 Hz, 2H), 1.51 (d, J = 2.8 Hz, 3H), 1.45 (d, J = 5.1 Hz, 3H), 1.37 (t, J = 12.0 Hz, 1H), 1.22 (dd, J = 19.8, 6.3 Hz, 3H), 0.97 - 0.82 (m, 9H).
1811	531.35	1.24	1H NMR (400 MHz, DMSO) δ 12.63 (s, 1H), 12.55 (s, 1H), 8.27 (dd, J = 7.3, 2.1 Hz, 1H), 8.09 (d, J = 8.4 Hz, 1H), 7.98 (d, J = 8.5 Hz, 1H), 7.84 (s, 1H), 7.81 (d, J = 7.8 Hz, 1H), 7.58 (t, J = 7.6 Hz, 1H), 7.52 - 7.46 (m, 1H), 7.45 (s, 2H), 6.77 (d, J = 7.8 Hz, 1H), 6.46 (t, J = 6.8 Hz, 1H), 2.84 - 2.73 (m, 2H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			2.70 (s, 3H), 2.24 (s, 1H), 1.81 (dd, J = 11.7, 5.6 Hz, 1H), 1.53 - 1.37 (m, 7H), 0.92 (d, J = 6.3 Hz, 3H).
1812	613.30	1.47	
1813	485.20	2.18	
1814	524.22	1.97	
1815	592.30	2.02	
1816	536.22	1.85	
1817	570.20	2.07	1H NMR (400 MHz, DMSO) δ 12.44 (s, 1H), 7.79 (d, J = 7.9 Hz, 1H), 7.63 (t, J = 7.8 Hz, 1H), 7.47 (s, 1H), 7.38 (d, J = 9.7 Hz, 1H), 7.22 (dd, J = 19.7, 7.6 Hz, 2H), 6.88 (d, J = 10.7 Hz, 1H), 6.68 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 3.82 (d, J = 6.6 Hz, 2H), 2.63 - 2.53 (m, 2H), 2.31 (dt, J = 17.2, 8.5 Hz, 1H), 2.25 - 2.12 (m, 1H), 2.05 (dt, J = 13.3, 6.6 Hz, 1H), 1.86 (td, J = 14.2, 7.1 Hz, 1H), 1.67 (dd, J = 11.7, 6.1 Hz, 1H), 1.61 (s, 3H), 1.52 (t, J = 12.1 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.88 (d, J = 6.2 Hz, 3H), 0.69 (t, J = 7.4 Hz, 3H).
1818	514.21	1.42	
1819	490.29	1.52	
1820	482.24	0.76	1H NMR (400 MHz, DMSO) δ 12.50 (s, 1H), 9.57 (s, 1H), 8.25 (dd, J = 4.8, 1.8 Hz, 1H), 8.05 (dd, J = 7.8, 1.8 Hz, 1H), 7.75 (d, J = 8.0 Hz, 1H), 7.48 - 7.39 (m, 2H), 7.27 (t, J = 8.1 Hz, 1H), 7.13 (d, J = 8.0 Hz, 1H), 6.86 - 6.81 (m, 1H), 6.78 (dd, J = 7.8, 4.8 Hz, 1H), 2.14 (s, 1H), 1.85 (dd, J = 11.6, 5.6 Hz, 1H), 1.59 (d, J = 12.1 Hz, 6H), 1.40 (t, J = 12.2 Hz, 1H), 0.76 (d, J = 6.2 Hz, 3H).
1821	566.40	2.24	1H NMR (400 MHz, DMSO) δ 12.40 (s, 1H), 7.77 (d, J = 7.9 Hz, 1H), 7.63 (t, J = 7.8 Hz, 1H), 7.48 (d, J = 1.4 Hz, 1H), 7.43 (d, J = 10.1 Hz, 1H), 7.26 (d, J = 7.9 Hz, 1H), 7.21 (d, J = 7.3 Hz, 1H), 6.89 (dt, J = 10.7, 2.1 Hz, 1H), 6.69 (d, J = 8.4 Hz, 1H), 6.46 (s, 2H), 4.14 (d, J = 5.7 Hz, 1H), 3.83 (d, J = 6.6 Hz, 2H), 3.66 (t, J = 8.8 Hz, 1H), 3.35 (d, J = 9.9 Hz, 1H), 2.30 (s, 2H), 2.11 (s, 1H), 2.08 - 1.99 (m, 1H), 1.54 (d, J = 10.1 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 1.39 (d, J = 7.9 Hz, 2H), 1.12 (d, J = 10.1 Hz, 1H), 1.04 (s, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.80 (d, J = 10.2 Hz, 1H).
1822	510.53	1.06	
1823	505.31	1.34	
1824	491.30	1.16	
1825	526.13	1.42	
1826	517.32	2.03	
1827	494.20	2.49	
1828	511.31	2.52	1H NMR (400 MHz, DMSO) δ 12.89 (s, 1H), 8.22 - 8.15 (m, 2H), 7.92 - 7.89 (d, 1H), 7.51 - 7.48 (dd, 1H), 7.37 - 7.30 (m, 2H), 6.85 - 6.82 (dd, 1H), 6.74 - 6.70 (m, 2H), 6.60 - 6.57 (dd, 1H), 3.94 (q, J = 7.04, 2H), 2.26 - 2.17 (m, 1H), 1.89 - 1.82 (m, 1H), 1.51 (s, 3H), 1.47 (s, 3H), 1.44 - 1.35 (m, 1H), 1.25 (t, J = 6.92, 3H), 0.88 - 0.86 (d, 3H)
1829	480.28	0.99	
1830	515.20	1.58	
1831	519.12	2.38	
1832	541.30	1.72	1H NMR (400 MHz, Methanol-d4) δ 8.65 (d, J = 4.9 Hz, 1H), 8.31 (s, 1H), 8.19 (d, J = 7.9 Hz, 1H), 8.04 (t, J = 7.4 Hz, 1H), 7.91 (d, J = 8.0 Hz, 1H), 7.60 - 7.54 (m, 1H), 7.47 (d, J = 2.1 Hz, 1H), 7.32 (d, J = 10.4 Hz, 1H), 7.21 (d, J = 7.9 Hz, 1H), 6.72 - 6.65 (m, 1H), 3.80 (d, J = 6.6 Hz, 2H), 3.09 (dd, J = 16.7, 7.0 Hz, 1H), 2.93 (t, J = 10.3 Hz, 1H), 2.36 (s, 1H), 2.16 - 2.01 (m, 1H), 1.92 (d, J = 6.3 Hz, 1H), 1.63 (s, 6H), 1.55 (t, J = 12.0 Hz, 1H), 1.03 (dd, J = 6.7, 1.9 Hz, 6H), 0.96 (d, J = 6.3 Hz, 3H).
1833	545.23	2.34	
1834	459.39	2.54	
1835	610.20	2.22	1H NMR (400 MHz, DMSO) δ 12.53 (s, 1H), 7.81 (d, J = 7.9 Hz, 1H), 7.66 (t, J = 7.8 Hz, 1H), 7.55 - 7.48 (m, 2H), 7.46 (d, J = 10.0 Hz, 1H), 7.23 (d, J = 7.3 Hz, 1H), 6.91 (d, J = 10.7 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 6.46 (s, 2H), 4.19 (d, J = 13.8 Hz, 1H), 3.83 (d, J =

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			6.5 Hz, 2H), 3.62 (s, 1H), 3.04 (t, J = 12.4 Hz, 1H), 2.66 (s, 1H), 2.12 - 1.95 (m, 1H), 1.76 (s, 1H), 1.72 - 1.62 (m, 1H), 1.56 (d, J = 17.7 Hz, 2H), 0.97 (dd, J = 13.9, 6.7 Hz, 10H).
1836	556.00	2.13	1H NMR (400 MHz, DMSO) δ 12.47 (s, 1H), 10.90 (s, 1H), 7.80 (d, J = 7.6 Hz, 1H), 7.72 (d, J = 8.0 Hz, 1H), 7.49 (s, 2H), 7.41 (d, J = 9.9 Hz, 1H), 7.24 (d, J = 7.9 Hz, 1H), 7.00 (d, J = 8.2 Hz, 2H), 6.88 (d, J = 10.5 Hz, 1H), 3.83 (d, J = 6.6 Hz, 2H), 2.70 (s, 1H), 2.60 (s, 1H), 2.22 (s, 1H), 2.05 (s, 1H), 1.87 (s, 1H), 1.59 (s, 6H), 1.41 (s, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.84 (d, J = 6.2 Hz, 3H).
1837	521.30	1.64	
1838	492.40	2.52	
1839	486.33	1.53	
1840	514.40	2.00	1H NMR (400 MHz, DMSO) δ 12.44 (s, 1H), 7.80 (d, J = 8.0 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.50 - 7.47 (m, 1H), 7.47 - 7.40 (m, 1H), 7.35 (d, J = 8.0 Hz, 1H), 7.20 (dd, J = 7.3, 0.6 Hz, 1H), 6.90 (dt, J = 10.8, 2.3 Hz, 1H), 6.69 (dd, J = 8.4, 0.6 Hz, 1H), 6.46 (s, 2H), 4.45 (d, J = 8.0 Hz, 1H), 3.84 (d, J = 6.6 Hz, 2H), 3.61 - 3.53 (m, 1H), 3.32 (d, J = 6.8 Hz, 1H), 2.38 - 2.26 (m, 1H), 2.05 - 1.95 (m, 1H), 1.87 - 1.80 (m, 1H), 1.38 (d, J = 6.2 Hz, 3H), 1.00 - 0.95 (m, 6H).
1841	500.21	2.37	1H NMR (400 MHz, DMSO) δ 12.81 (s, 1H), 9.83 (s, 1H), 8.15 - 8.13 (dd, 1H), 7.93 - 7.86 (m, 2H), 7.75 - 7.72 (m, 2H), 7.55 - 7.52 (d, 1H), 7.26 (t, J = 8.13, 1H), 7.12 - 7.09 (d, 1H), 6.99 - 6.96 (dd, 1H), 6.55 - 6 .52 (dd, 1H), 1.91 (m, 1H), 1.79 - 1.72 (m, 1H), 1.45 (s, 3H), 1.43 (s, 3H), 1.31 (t, J = 12.09, 1H), 0.72 - 0.70 (d, 3H)
1842	495.40	2.60	
1843	524.29	1.74	
1844	514.30	1.39	1H NMR (400 MHz, DMSO-d6) δ 12.44 (s, 1H), 8.24 (dd, J = 4.8, 1.9 Hz, 1H), 8.18 (d, J = 2.8 Hz, 1H), 8.03 (dd,

Cmpd. No	LC/MS M+1	LC/RT min	NMR
			J = 7.8, 2.0 Hz, 1H), 7.84 (d, J = 8.2 Hz, 1H), 6.92 (d, J = 8.3 Hz, 1H), 6.77 (dd, J = 7.8, 4.8 Hz, 1H), 6.68 (broad s, 2H), 6.08 (d, J = 2.8 Hz, 1H), 4.95 - 4.77 (m, 1H), 2.44 (m, 2H partially overlapped with solvent peak), 2.12 (br s, 1H), 1.85 (dd, J = 11.8, 5.6 Hz, 1H), 1.54 (s, 3H), 1.53 (s, 3H), 1.41-1.38 (m, 1H overlapped with following signal), 1.34-1.31 (m, 6H), 0.74 (d, J = 6.2 Hz, 3H).
1845	538.60	1.32	
1846	588.50	1.65	
1847	528.26	1.45	
1848	511.50	1.53	1H NMR (400 MHz, DMSO-d6) δ 12.56 (s, 1H), 12.53 (d, J = 6.3 Hz, 1H), 8.24 (dd, J = 7.2, 2.2 Hz, 1H), 7.86 - 7.79 (m, 1H), 7.73 - 7.65 (m, 2H), 7.40 - 7.31 (m, 1H), 7.13 (dd, J = 11.0, 8.1 Hz, 2H), 7.04 (t, J = 7.5 Hz, 1H), 6.45 (t, J = 6.9 Hz, 1H), 4.10 (q, J = 6.9 Hz, 2H), 2.77 - 2.71 (m, 2H), 2.24 (s, 1H), 1.86 (dd, J = 11.8, 5.6 Hz, 1H), 1.55 (s, 7H), 1.43 (t, J = 12.2 Hz, 1H), 1.34 (t, J = 6.9 Hz, 3H), 0.91 (d, J = 6.3 Hz, 3H).
1849	491.23	1.63	
1850	474.30	0.94	1H NMR (400 MHz, DMSO-d6) δ 12.43 (s, 1H), 8.24 (dd, J = 4.8, 1.9 Hz, 1H), 8.03 (dd, J = 7.9, 2.3 Hz, 1H), 7.61 (d, J = 7.8 Hz, 1H), 6.94-6.58 (m, 2H), 6.77 (dd, J = 7.9, 4.8 Hz, 1H, obscured by previous signal), 6.54 (d, J = 7.8 Hz, 1H), 4.00 - 3.85 (m, 2H), 3.53 - 3.30 (m, 2H), 2.89 - 2.73 (m, 1H), 2.49 - 2.37 (m, 2H), 2.17 - 2.04 (m, 1H), 1.86 - 1.61 (m, 5H), 1.52 (s, 6H), 1.36 (t, J = 12.0 Hz, 1H), 0.74 (d, J = 6.3 Hz, 3H).
1851	514.22	1.39	1H NMR (400 MHz, DMSO-d6) δ 12.52 (s, 1H), 8.24 (dd, J = 4.9, 1.8 Hz, 1H), 8.04 (dd, J = 7.8, 1.8 Hz, 1H), 7.91 - 7.81 (m, 2H), 7.74 (d, J = 8.0 Hz, 1H), 7.29 (t, J = 8.9 Hz, 1H), 7.21 (d, J = 8.0 Hz, 1H), 6.77 (dd, J = 7.8, 4.8 Hz, 1H), 3.89 (s, 3H), 2.53 (d, J =

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			10.6 Hz, 1H), 2.14 (s, 1H), 2.06 (s, 1H), 1.85 (dd, J = 11.9, 5.7 Hz, 1H), 1.58 (d, J = 12.2 Hz, 7H), 1.39 (t, J = 12.1 Hz, 1H), 0.75 (d, J = 6.3 Hz, 3H).
1852	522.20	1.64	1H NMR (400 MHz, DMSO) δ 12.38 (s, 1H), 7.63 (d, J = 8.0 Hz, 1H), 7.61 - 7.53 (m, 1H), 7.34 - 7.26 (m, 2H), 7.21 (dd, J = 7.1, 5.5 Hz, 3H), 7.13 (d, J = 7.3 Hz, 1H), 6.67 (d, J = 8.0 Hz, 1H), 6.61 (d, J = 8.4 Hz, 1H), 6.36 (s, 1H), 2.80 (d, J = 11.1 Hz, 2H), 2.62 (dd, J = 17.8, 11.4 Hz, 1H), 2.54 (s, 1H), 2.37 (dd, J = 27.9, 9.4 Hz, 1H), 2.00 (dd, J = 27.6, 18.9 Hz, 1H), 1.57 (s, 5H), 1.48 (s, 3H), 1.28 (s, 9H).
1853	528.14	1.59	
1854	502.52	1.03	
1855	556.20	1.85	1H NMR (400 MHz, DMSO) δ 8.40 (d, J = 8.2 Hz, 1H), 8.18 (d, J = 8.2 Hz, 1H), 7.82 - 7.54 (m, 3H), 7.52 - 7.41 (m, 1H), 7.25 - 7.16 (m, 1H), 6.96 (dd, J = 42.8, 10.7 Hz, 1H), 6.70 (t, J = 8.3 Hz, 1H), 6.37 (s, 1H), 3.86 (dd, J = 12.1, 6.6 Hz, 2H), 3.58 (d, J = 9.5 Hz, 2H), 3.26 (d, J = 11.5 Hz, 1H), 2.82 (dt, J = 24.3, 11.1 Hz, 1H), 2.06 - 1.99 (m, 3H), 1.93 - 1.74 (m, 1H), 1.73 - 1.48 (m, 2H), 1.41 - 1.27 (m, 2H), 1.01 - 0.81 (m, 14H).
1856	515.40	1.69	
1857	516.23	1.51	
1858	515.17	1.58	
1859	469.40	1.55	1H NMR (400 MHz, DMSO-d6) δ 13.03 (s, 1H), 8.45 (s, 1H), 8.31 (d, J = 7.3 Hz, 1H), 8.15 (s, 1H), 7.68 (d, J = 8.2 Hz, 1H), 7.32 (s, 1H), 7.25 (s, 1H), 7.05 (d, J = 8.2 Hz, 1H), 4.02 (s, 3H), 2.83 - 2.64 (m, 1H), 2.59 - 2.53 (m, 1H, obscured by solvent peak), 2.43 (s, 3H), 2.28 - 2.13 (m, 1H), 1.84 (dd, J = 11.8, 5.8 Hz, 1H), 1.60 (d, J = 3.7 Hz, 6H), 1.37 (t, J = 12.0 Hz, 1H), 0.78 (d, J = 6.2 Hz, 3H).
1860	524.25	1.63	
1861	497.27	1.58	

Cmpd. No.	LC/MS M+1	LC/RT min.	NMR
1862	495.40	1.68	
1863	556.20	2.01	
1864	556.00	1.85	
1865	487.30	1.88	
1866	511.15	1.33	
1867	529.36	1.13	
1868	458.40	1.69	
1869	514.15	1.45	
1870	476.19	2.17	
1871	494.60	1.45	
1872	500.21	2.64	1H NMR (400 MHz, DMSO-d6) δ 12.70 (s, 1H), 8.93 (s, 1H), 8.25 - 8.22 (d, 1H), 8.15 - 8.13 (dd, 1H), 7.88 (t, J = 7.91, 1H), 7.62 - 7.59 (dd, 1H), 7.53 - 7.48 (m, 2H), 7.35 - 7.32 (d, 1H), 7.27 (t, J = 7.91, 1H), 7.08 (t, J = 7.91, 1H), 6.53 - 6.50 (dd, 1H), 2.03 - 1.94 (m, 1H), 1.82 - 1.75 (m, 1H), 1.46 (s, 3H), 1.45 (s, 3H), 1.33 (t, J = 11.91, 1H), 0.79 - 0.77 (d, 3H)
1873	484.27	0.94	
1874	487.30	2.59	
1875	518.22	1.47	1H NMR (400 MHz, DMSO-d6) δ 9.28 - 9.26 (dd, 1H), 9.17 - 9.14 (d, 1H), 8.44 - 8.41 (d, 1H), 8.33 (t, J = 7.80, 1H), 8.23 - 8.22 (d, 1H), 8.14 - 8.12 (dd, 1H), 8.06 - 8.00 (m, 2H), 7.82 - 7.78 (dd, 1H), 6.75 - 6.72 (d, 1H), 7.50 - 7.47 (dd, 1H), 6.53 - 6.50 (dd, 1H), 2.33 - 2.22 (m, 1H), 1.88 - 1.81 (m, 1H), 1.48 (s, 3H), 1.45 (s, 3H), 1.39-1.34 (m, 1H), 0.89 - 0.87 (d, 3H)
1876	570.00	2.23	1H NMR (400 MHz, DMSO) δ 12.39 (s, 1H), 7.94 (d, J = 8.9 Hz, 2H), 7.69 (d, J = 8.0 Hz, 1H), 7.47 (s, 1H), 7.40 (d, J = 10.0 Hz, 1H), 7.26 (d, J = 8.0 Hz, 1H), 7.17 (d, J = 8.9 Hz, 2H), 6.89 (d, J = 10.7 Hz, 1H), 3.83 (d, J = 6.7 Hz, 2H), 3.31 (s, 3H), 2.41 (s, 1H), 2.32 (s, 1H), 2.04 (s, 2H), 1.84 (d, J = 11.9 Hz, 1H), 1.57 (d, J = 18.4 Hz, 6H), 1.38 (s, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.65 (d, J = 6.2 Hz, 3H).
1877	549.43	1.81	
1878	527.20	1.47	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1879	537.43	1.63	
1880	511.14	1.59	
1881	445.28	2.38	
1882	591.60	1.39	1H NMR (400 MHz, DMSO-d6) δ 12.41 (s, 1H), 7.74 - 7.66 (m, 1H), 7.66 - 7.59 (m, 1H), 7.18 (dd, J = 7.2, 2.5 Hz, 1H), 6.88 (d, J = 8.1 Hz, 1H), 6.82 (s, 1H), 6.69 (dd, J = 10.1, 8.0 Hz, 1H), 4.19 (q, J = 18.4, 17.6 Hz, 2H), 3.35 (t, J = 5.7 Hz, 2H), 2.95 (t, J = 6.8 Hz, 2H), 2.54 (d, J = 8.8 Hz, 1H), 2.41 (s, 2H), 2.12 (dq, J = 13.4, 6.7 Hz, 1H), 1.85 (dd, J = 11.7, 5.7 Hz, 1H), 1.53 (d, J = 7.3 Hz, 1H), 1.51 (s, 3H), 1.39 - 1.33 (m, 1H), 1.17 (t, J = 7.3 Hz, 1H), 1.05 (s, 1H), 1.03 (s, 3H), 1.01 (s, 3H), 0.99 - 0.92 (m, 1H), 0.89 (t, J = 3.1 Hz, 1H), 0.86 (d, J = 6.3 Hz, 3H).
1883	562.40	1.54	
1884	473.19	2.73	
1885	556.00	2.04	1H NMR (400 MHz, DMSO) δ 12.48 (s, 1H), 10.18 (s, 1H), 7.71 (d, J = 8.1 Hz, 1H), 7.44 (d, J = 25.5 Hz, 5H), 7.26 (d, J = 8.0 Hz, 1H), 7.07 (d, J = 7.7 Hz, 1H), 6.89 (d, J = 10.6 Hz, 1H), 3.83 (d, J = 6.6 Hz, 2H), 2.39 (s, 1H), 2.08 (d, J = 42.4 Hz, 2H), 1.84 (s, 1H), 1.58 (d, J = 17.2 Hz, 6H), 1.39 (s, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.72 (d, J = 6.2 Hz, 3H).
1886	497.27	1.36	
1887	535.40	1.59	1H NMR (400 MHz, DMSO-d6) δ 12.45 (s, 1H), 8.23 (d, J = 3.7 Hz, 1H), 8.19 (d, J = 2.8 Hz, 1H), 8.02 (dd, J = 7.8, 1.9 Hz, 1H), 7.83 (d, J = 8.3 Hz, 1H), 6.93 (d, J = 8.2 Hz, 1H), 6.76 (dd, J = 7.8, 4.8 Hz, 1H), 6.68 (broad s, 2H), 6.12 (d, J = 2.8 Hz, 1H), 3.97 (s, 2H), 2.46 (d, J = 7.8 Hz, 2H), 2.20 - 2.02 (br m, 1H), 1.85 (dd, J = 11.9, 5.6 Hz, 1H), 1.55 (s, 3H), 1.53 (s, 3H), 1.38 (t, J = 12.2 Hz, 1H), 0.74 (d, J = 6.3 Hz, 3H).
1888	482.30	1.59	1H NMR (400 MHz, DMSO-d6) δ 8.03 (t, J = 2.5 Hz, 1H), 7.47 (dd, J = 4.0, 2.5 Hz, 1H), 7.40 (t, J = 7.7 Hz, 1H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			7.01 (d, J = 7.3 Hz, 1H), 6.42 (d, J = 8.1 Hz, 1H), 6.08 (d, J = 3.1 Hz, 1H), 6.01 (s, 2H), 2.97 (ddd, J = 31.6, 19.5, 11.1 Hz, 3H), 2.09 (d, J = 15.9 Hz, 1H), 1.85 - 1.63 (m, 3H), 1.48 (dt, J = 4.5, 2.5 Hz, 7H), 1.43 - 1.25 (m, 2H), 1.25 - 0.91 (m, 3H), 0.85 (d, J = 6.3 Hz, 3H).
1889	538.34	1.78	1H NMR (400 MHz, DMSO-d6) δ 12.44 (s, 1H), 8.50 (d, J = 2.4 Hz, 1H), 7.89 (d, J = 2.5 Hz, 1H), 7.64 (t, J = 7.9 Hz, 1H), 7.22 (d, J = 7.3 Hz, 1H), 7.00 (d, J = 17.8 Hz, 2H), 6.87 - 6.58 (m, 2H), 6.48 (s, 2H), 4.69 (p, J = 6.1 Hz, 1H), 2.33 (s, 3H), 2.20 (s, 1H), 1.85 (dd, J = 11.9, 5.7 Hz, 1H), 1.54 (d, J = 7.2 Hz, 6H), 1.40 (t, J = 12.0 Hz, 1H), 1.28 (d, J = 6.0 Hz, 6H), 0.86 (d, J = 6.3 Hz, 3H).
1890	581.20	1.76	1H NMR (400 MHz, DMSO-d6) δ 12.39 (s, 1H), 8.16 (d, J = 7.8 Hz, 1H), 7.84 (d, J = 7.9 Hz, 1H), 7.66 (t, J = 7.9 Hz, 1H), 7.24 (d, J = 7.3 Hz, 1H), 7.21 - 7.14 (m, 4H), 7.14 - 7.07 (m, 1H), 6.84 (dt, J = 10.6, 2.3 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 6.54 (s, 2H), 4.61 (s, 1H), 3.61 (s, 2H), 2.07 (s, 1H), 2.06 (s, 6H), 1.19 (s, 6H).
1891	536.50	1.35	1H NMR (400 MHz, DMSO-d6) δ 12.24 (s, 1H), 7.66 - 7.55 (m, 2H), 7.17 (dd, J = 7.3, 0.7 Hz, 1H), 6.66 (dd, J = 8.4, 0.8 Hz, 1H), 6.44 (s, 2H), 5.03 (dt, J = 10.8, 6.2 Hz, 1H), 3.86 (s, 2H), 3.28 (dd, J = 12.8, 10.6 Hz, 2H), 2.63 - 2.51 (m, 1H), 2.43 (t, J = 8.8 Hz, 1H), 2.19 (dq, J = 11.9, 6.2 Hz, 1H), 1.83 (dd, J = 12.0, 5.9 Hz, 2H), 1.72 - 1.51 (m, 2H), 1.51 (s, 3H), 1.47 - 1.19 (m, 9H), 0.85 (d, J = 6.3 Hz, 3H).
1892	494.20	2.42	
1893	563.60	2.26	
1894	552.39	1.98	
1895	528.20	1.52	1H NMR (400 MHz, DMSO-d6) δ 12.48 (s, 1H), 8.24 (d, J = 4.8 Hz, 1H), 8.02 (dd, J = 7.9, 1.9 Hz, 1H), 7.92 (d, J = 1.0 Hz, 1H), 7.83 (d, J = 8.2 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 7.69 (d, J = 0.9 Hz, 1H), 7.06 (d, J = 8.3 Hz, 1H), 6.76 (dd, J = 7.8, 4.8 Hz, 1H), 6.67 (dd, J = 7.7, 4.8 Hz, 1H), 3.74 (d, J = 6.4 Hz, 2H), 2.14 (s, 1H), 2.01 (dt, J = 13.3, 6.6 Hz, 1H), 1.86 (dd, J = 11.9, 5.6 Hz, 1H), 1.56 (d, J = 5.3 Hz, 6H), 1.39 (t, J = 12.1 Hz, 1H), 0.96 (d, J = 6.7 Hz, 6H), 0.75 (d, J = 6.3 Hz, 3H).
1896	529.30	1.80	1H NMR (400 MHz, DMSO-d6) δ 12.52 (s, 2H), 8.23 (d, J = 7.2 Hz, 1H), 8.19 (d, J = 2.5 Hz, 1H), 7.87 - 7.72 (m, 2H), 6.88 (d, J = 8.3 Hz, 1H), 6.44 (broad t, 1H), 6.11 (d, J = 2.6 Hz, 1H), 3.98 (d, J = 6.6 Hz, 2H), 2.72-2.70 (m , 2H), 2.22 (br s, 1H), 2.05 (sep, J = 8.0 1H), 1.88 (dd, J = 11.8, 5.5 Hz, 1H), 1.56 (s, 3H), 1.55 (s, 3H), 1.43 (t, J = 12.1 Hz, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.89 (d, J = 6.3 Hz, 3H).
1897	516.36	1.50	
1898	480.10	2.33	
1899	533.22	2.84	
1900	526.30	1.59	
1901	570.50	2.10	1H NMR (400 MHz, DMSO) δ 12.55 (s, 1H), 7.71 (d, J = 7.9 Hz, 1H), 7.64 (dd, J = 8.3, 7.4 Hz, 1H), 7.50 (d, J = 1.9 Hz, 1H), 7.47 - 7.38 (m, 1H), 7.30 (d, J = 7.9 Hz, 1H), 7.24 - 7.17 (m, 1H), 6.88 (dt, J = 10.7, 2.3 Hz, 1H), 6.73 - 6.67 (m, 1H), 6.48 (s, 2H), 3.84 (d, J = 6.6 Hz, 2H), 3.60 (s, 2H), 3.16 (t, J = 5.9 Hz, 2H), 2.09 - 1.99 (m, 1H), 1.55 (d, J = 4.7 Hz, 2H), 1.45 (s, 2H), 1.35 - 1.25 (m, 2H), 0.99 (d, J = 6.7 Hz, 6H), 0.83 (s, 6H).
1902	509.21	2.59	
1903	524.34	1.50	1H NMR (400 MHz, DMSO) δ 12.51 (s, 1H), 8.25 (dd, J = 4.8, 1.8 Hz, 1H), 8.05 (dd, J = 7.8, 1.7 Hz, 1H), 7.98 (d, J = 8.9 Hz, 2H), 7.74 (d, J = 8.0 Hz, 1H), 7.17 (d, J = 8.1 Hz, 1H), 7.02 (d, J = 8.9 Hz, 2H), 6.78 (dd, J = 7.8, 4.8 Hz, 1H), 4.69 (dt, J = 12.0, 6.0 Hz, 1H), 2.57 - 2.51 (m, 2H), 2.15 (s, 1H), 1.85 (dd, J = 11.8, 5.6 Hz, 1H), 1.60 (s, 3H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1.58 (s, 3H), 1.40 (t, J = 12.1 Hz, 1H), 1.29 (s, 3H), 1.28 (s, 3H), 0.76 (d, J = 6.3 Hz, 3H).
1904	533.31	1.12	
1905	494.20	2.32	1H NMR (400 MHz, DMSO) δ 12.41 (s, 1H), 8.19 - 8.15 (m, 1H), 7.80 - 7.77 (d, 1H), 7.63 - 7.54 (m, 1H), 7.45 - 7.42 (d, 2H), 7.33 - 7.06 (m, 4H), 6.75 - 6.53 (m, 2H), 5.14 - 5.12 (d, 1H), 2.21 (t, J = 10.66, 1H), 1. 88 - 1.81 (m, 1H), 1.72 - 1.65 (m, 1H), 1.52 - 1.43 (m, 9H), 1.22 (t, J = 11.65, 1H), 0.85 - 0.83 (d, 1H), 0.56 - 0.54 (d, 2H)
1906	534.12	2.82	
1907	552.50	1.08	
1908	473.19	2.70	
1909	506.21	2.75	1H NMR (400 MHz, DMSO) δ 12.71 (s, 1H), 9.45 (s, 1H), 8.15 - 8.13 (dd, 1H), 7.83 - 7.70 (m, 3H), 7.47 - 7.44 (dd, 1H), 7.42 - 7.39 (d, 1H), 7.10 - 7.07 (d, 1H), 7.05 - 7.02 (d, 1H), 6.52 - 6.49 (dd, 1H), 2.80 - 2.6 1 (m, 3H), 1.96 - 1.90 (m, 2H), 1.75 - 1.69 (m, 2H), 1.45 (s, 3H), 1.40 (s, 3H), 1.29 (t, J = 11.91, 1H), 0.72 - 0.70 (d, 3H)
1910	498.23	1.63	
1911	460.40	1.34	
1912	512.40	1.82	1H NMR (400 MHz, DMSO) δ 12.49 (s, 1H), 8.02 - 7.97 (m, 2H), 7.92 (d, J = 1.0 Hz, 1H), 7.80 (d, J = 8.3 Hz, 1H), 7.76 - 7.71 (m, 1H), 7.70 - 7.63 (m, 3H), 7.04 (d, J = 8.3 Hz, 1H), 3.73 (d, J = 6.4 Hz, 2H), 2.43 (t, J = 10.4 Hz, 1H), 2.34 - 2.25 (m, 1H), 2.10 (dq, J = 12.0, 6.1 Hz, 1H), 2.01 (dq, J = 13.3, 6.7 Hz, 1H), 1.88 - 1.79 (m, 1H), 1.54 (d, J = 8.1 Hz, 6H), 1.38 (t, J = 12.1 Hz, 1H), 0.96 (d, J = 6.7 Hz, 6H), 0.66 (d, J = 6.3 Hz, 3H).
1913	533.31	1.09	1H NMR (400 MHz, DMSO) δ 12.67 (s, 1H), 12.54 (s, 1H), 8.88 (d, J = 1.7 Hz, 1H), 8.42 (d, J = 2.8 Hz, 1H), 8.25 (dd, J = 7.3, 2.2 Hz, 1H), 8.04 (d, J = 1.7 Hz, 1H), 7.83 (s, 1H), 7.79 (d, J = 7.9 Hz, 1H), 7.34 (d, J = 8.0 Hz, 1H), 6.49 -

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			6.42 (m, 1H), 3.93 (s, 3H), 2.73 (tt, J = 16.6, 8.3 Hz, 2H), 2.25 (d, J = 4.1 Hz, 1H), 1.90 (dd, J = 12.0, 5.5 Hz, 1H), 1.61 (d, J = 2.4 Hz, 6H), 1.46 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).
1914	533.31	1.18	
1915	552.30	1.63	1H NMR (400 MHz, DMSO) δ 8.72 (d, J = 2.5 Hz, 1H), 8.01 (d, J = 2.4 Hz, 1H), 7.56 (d, J = 7.8 Hz, 1H), 7.43 (t, J = 7.7 Hz, 1H), 6.98 (dd, J = 13.6, 7.5 Hz, 2H), 6.45 (d, J = 8.1 Hz, 1H), 6.11 (s, 2H), 3.23 (q, J = 6.8 Hz, 4H), 3.04 - 2.88 (m, 2H), 2.27 (d, J = 5.1 Hz, 3H), 2.12 (qd, J = 12.2, 11.5, 5.2 Hz, 1H), 1.83 - 1.75 (m, 1H), 1.62 (s, 2H), 1.57 (s, 2H), 1.35 (d, J = 11.9 Hz, 1H), 1.05 (t, J = 6.9 Hz, 6H), 0.84 (d, J = 6.3 Hz, 4H).
1916	472.30	1.70	
1917	495.40	2.60	
1918	550.38	1.88	
1919	501.41	3.02	
1920	500.50	1.44	
1921	534.25	1.75	
1922	498.50	1.67	
1923	510.16	1.36	
1924	511.31	2.48	
1925	485.50	1.56	
1926	523.32	2.90	
1927	497.40	1.40	1H NMR (400 MHz, DMSO-d6) δ 12.59 (s, 1H), 12.53 (s, 1H), 8.25 (dd, J = 7.3, 2.2 Hz, 1H), 7.83 (s, 1H), 7.75 (d, J = 7.9 Hz, 1H), 7.66 - 7.57 (m, 2H), 7.40 (t, J = 7.9 Hz, 1H), 7.20 (d, J = 8.0 Hz, 1H), 7.05 - 6.96 (m, 1H), 6.45 (t, J = 6.8 Hz, 1H), 3.82 (s, 3H), 2.79 - 2.65 (m, 2H), 2.30 - 2.17 (m, 1H), 1.89 (dd, J = 11.9, 5.5 Hz, 1H), 1.62 (d, J = 1.7 Hz, 6H), 1.45 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).
1928	528.40	2.25	
1929	549.13	2.85	
1930	498.23	1.65	
1931	538.40	2.65	
1932	472.29	2.49	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1933	517.32	1.70	
1934	551.50	1.82	1H NMR (400 MHz, DMSO-d6) δ 12.64 (s, 1H), 12.56 - 12.50 (m, 1H), 8.28 - 8.21 (m, 1H), 8.10 - 8.02 (m, 2H), 7.83 (t, J = 6.3 Hz, 1H), 7.78 (d, J = 8.0 Hz, 1H), 7.63 (t, J = 8.0 Hz, 1H), 7.43 (d, J = 8.1 Hz, 1H), 7.28 (d, J = 8.0 Hz, 1H), 6.45 (t, J = 6.7 Hz, 1H), 2.81 - 2.65 (m, 2H), 2.25 (s, 1H), 1.90 (dd, J = 12.2, 5.5 Hz, 1H), 1.60 (d, J = 3.6 Hz, 6H), 1.45 (t, J = 12.2 Hz, 1H), 0.91 (d, J = 6.2 Hz, 3H).
1935	548.60	1.72	
1936	473.19	2.85	
1937	545.23	1.42	
1938	480.10	2.21	
1939	489.10	2.19	
1940	524.52	2.48	
1941	518.22	2.63	
1942	618.70	1.47	
1943	554.34	1.46	
1944	498.53	1.05	
1945	510.31	1.42	
1946	499.50	1.60	
1947	494.28	1.44	
1948	569.30	1.49	
1949	508.31	2.00	
1950	534.28	1.80	
1951	494.57	1.10	
1952	542.60	1.99	1H NMR (400 MHz, DMSO-d6) δ 12.35 (s, 1H), 8.20 (d, J = 2.8 Hz, 1H), 7.86 (d, J = 8.5 Hz, 1H), 7.62 (broad s, 1H), 7.18 (d, J = 7.0 Hz, 1H), 6.93 (d, J = 8.2 Hz, 1H), 6.50 (br s, 2H), 6.13 (s, 1H), 3.89 (s, 2H), 2.59 (br s, 2H), 2.20 (br s, 1H), 1.93 - 1.82 (m, 1H), 1.56 (s, 3H), 1.55 (s, 3H), 1.41 (t, J = 12.1 Hz, 1H), 1.00 (s, 9H), 0.88 (d, J = 6.3 Hz, 3H).
1953	549.43	2.72	
1954	503.30	1.52	
1955	521.22	2.49	
1956	569.50	2.27	
1957	502.31	2.10	
1958	466.28	1.33	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1959	496.60	2.30	
1960	535.30	1.53	1H NMR (400 MHz, DMSO-d6) δ 12.63 (s, 1H), 12.55 (d, J = 4.9 Hz, 1H), 8.25 (dd, J = 7.3, 2.1 Hz, 1H), 7.88 - 7.78 (m, 2H), 7.76 - 7.71 (m, 2H), 7.64 (t, J = 7.6 Hz, 1H), 7.45 (d, J = 7.4 Hz, 1H), 6.59 (d, J = 7.8 Hz, 1H), 6.45 (t, J = 6.8 Hz, 1H), 2.78 - 2.65 (m, 2H), 2.28 - 2.14 (m, 1H), 1.82 (dd, J = 11.8, 5.6 Hz, 1H), 1.48 - 1.35 (m, 7H), 0.90 (d, J = 6.3 Hz, 3H).
1961	553.20	2.26	
1962	496.28	0.92	
1963	514.21	1.82	
1964	524.25	1.54	
1965	511.31	2.52	
1966	520.40	1.80	1H NMR (400 MHz, DMSO-d6) δ 12.23 (s, 1H), 7.66 - 7.54 (m, 2H), 7.17 (d, J = 7.3 Hz, 1H), 6.66 (d, J = 8.4 Hz, 1H), 6.44 (s, 1H), 5.09 (td, J = 6.4, 3.8 Hz, 1H), 2.59 - 2.51 (m, 1H), 2.43 (dd, J = 10.7, 5.8 Hz, 1H), 2.15 (dt, J = 16.6, 8.6 Hz, 3H), 1.83 (dd, J = 11.9, 5.5 Hz, 1H), 1.70 (d, J = 11.9 Hz, 2H), 1.52 (d, J = 17.4 Hz, 7H), 1.44 (d, J = 4.6 Hz, 3H), 1.38 (d, J = 12.1 Hz, 1H), 1.28 (d, J = 6.1 Hz, 2H), 1.23 (d, J = 6.2 Hz, 2H), 0.85 (d, J = 6.3 Hz, 3H).
1967	490.32	1.50	
1968	548.23	2.62	
1969	538.40	2.66	1H NMR (400 MHz, DMSO-d6) δ 12.43 (s, 1H), 7.78 (d, J = 8.0 Hz, 1H), 7.62 (q, J = 7.4, 6.9 Hz, 3H), 7.38 (t, J = 7.9 Hz, 1H), 7.21 (dd, J = 7.6, 5.4 Hz, 2H), 7.00 (dd, J = 8.2, 2.3 Hz, 1H), 6.69 (d, J = 8.4 Hz, 1H), 6.48 (s, 1H), 3.81 (d, J = 6.6 Hz, 2H), 2.64 - 2.57 (m, 2H), 2.22 (dt, J = 13.4, 7.1 Hz, 1H), 2.04 (dq, J = 13.4, 6.7 Hz, 1H), 1.88 (dd, J = 11.9, 5.6 Hz, 1H), 1.62 (s, 3H), 1.61 (s, 3H), 1.42 (t, J = 12.0 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.88 (d, J = 6.2 Hz, 3H).
1970	528.40	1.62	
1971	528.42	2.68	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1972	511.31	2.45	
1973	583.70	1.68	
1974	472.50	1.58	1H NMR (400 MHz, DMSO) δ 12.35 (s, 1H), 7.58 (d, J = 7.1 Hz, 2H), 7.14 (d, J = 7.1 Hz, 1H), 6.77 - 6.56 (m, 2H), 6.42 (s, 2H), 2.79 (s, 1H), 2.66 (dd, J = 12.3, 10.5 Hz, 1H), 1.69 (d, J = 20.9 Hz, 2H), 1.56 (s, 3H), 1.51 (s, 3H), 1.27 (s, 9H), 1.17 - 1.05 (m, 1H), 0.73 - 0.63 (m, 1H), 0.58 - 0.47 (m, 1H), 0.40 (dd, J = 8.2, 3.9 Hz, 1H), 0.24 (dd, J = 9.0, 4.1 Hz, 1H), 0.07 (dd, J = 9.0, 4.0 Hz, 1H).
1975	542.30	1.65	1H NMR (400 MHz, DMSO-d6) δ 12.45 (s, 1H), 8.24 (dd, J = 4.7, 1.9 Hz, 1H), 8.20 (d, J = 2.8 Hz, 1H), 8.03 (dd, J = 7.8, 1.9 Hz, 1H), 6.95 (d, J = 8.2 Hz, 1H), 6.77 (dd, J = 7.8, 4.8 Hz, 1H), 6.69 (broad s, 2H), 6.14 (d, J = 2.7 Hz, 1H), 3.89 (s, 2H), 2.45 (d, J = 8.9 Hz, 2H), 2.12 (br s, 1H), 1.85 (dd, J = 11.9, 5.4 Hz, 1H), 1.55 (s, 3H), 1.53 (s, 3H), 1.43 - 1.33 (m, 1H), 1.00 (s, 9H), 0.74 (d, J = 6.2 Hz, 3H).
1976	567.00	2.19	1H NMR (400 MHz, DMSO-d6) δ 8.85 (s, 1H), 8.67 (d, J = 4.7 Hz, 1H), 8.43 (d, J = 7.9 Hz, 1H), 8.30 (d, J = 9.0 Hz, 1H), 7.84 (d, J = 8.0 Hz, 1H), 7.53 - 7.42 (m, 1H), 7.28 (d, J = 8.0 Hz, 1H), 6.87 (d, J = 8.7 Hz, 1H), 5.31 (p, J = 6.3 Hz, 1H), 2.41 (s, 2H), 2.25 (s, 3H), 2.18 (dt, J = 16.2, 7.6 Hz, 1H), 1.88 (dd, J = 12.1, 5.8 Hz, 1H), 1.55 (d, J = 4.4 Hz, 6H), 1.42 (t, J = 12.0 Hz, 1H), 1.31 (d, J = 6.1 Hz, 6H), 0.74 (d, J = 6.3 Hz, 3H).
1977	550.33	2.61	
1978	508.31	2.63	
1979	538.30	1.77	
1980	497.20	1.30	
1981	521.61	1.73	
1982	506.25	1.26	
1983	514.31	2.67	
1984	609.60	1.41	
1985	529.32	1.98	
1986	542.23	2.86	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
1987	534.12	2.79	
1988	487.30	2.88	
1989	518.22	2.19	
1990	495.40	1.54	1H NMR (400 MHz, DMSO-d6) δ 12.59 (s, 1H), 12.53 (d, J = 6.3 Hz, 1H), 8.25 (dd, J = 7.3, 2.2 Hz, 1H), 7.82 (dt, J = 6.5, 3.4 Hz, 1H), 7.72 (d, J = 7.8 Hz, 1H), 7.19 - 7.06 (m, 3H), 6.64 (d, J = 7.8 Hz, 1H), 6.45 (t, J = 6.8 Hz, 1H), 2.79 - 2.69 (m, 2H), 2.29 (s, 3H), 2.24 (s, 4H), 1.84 (dd, J = 11.9, 5.7 Hz, 1H), 1.49 (d, J = 6.2 Hz, 6H), 1.41 (t, J = 12.0 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).
1991	508.32	1.12	
1992	487.30	2.39	
1993	556.00	1.85	
1994	526.18	1.38	
1995	570.20	1.86	
1996	500.51	2.65	
1997	497.50	1.07	1H NMR (400 MHz, DMSO-d6) δ 7.94 (s, 1H), 7.87 (d, J = 7.9 Hz, 1H), 7.84 - 7.74 (m, 1H), 7.66 - 7.53 (m, 1H), 7.39 (t, J = 7.7 Hz, 1H), 7.29 (d, J = 7.6 Hz, 1H), 6.97 (d, J = 7.7 Hz, 1H), 6.19 - 5.99 (m, 1H), 5.26 - 5.08 (m, 1H), 4.54 (s, 2H), 3.18 - 3.09 (m, 1H), 3.05 (t, J = 10.8 Hz, 1H), 2.20 - 2.08 (m, 1H), 1.81 (dd, J = 11.2, 5.6 Hz, 1H), 1.60 (d, J = 16.0 Hz, 6H), 1.37 (t, J = 11.9 Hz, 1H), 0.86 (d, J = 6.2 Hz, 3H).
1998	532.16	1.49	1H NMR (400 MHz, DMSO-d6) δ 12.51 (s, 1H), 8.26 (dd, J = 4.8, 1.9 Hz, 1H), 8.07 (dd, J = 7.9, 1.8 Hz, 1H), 7.80 (dd, J = 8.0, 1.7 Hz, 1H), 7.76 - 7.59 (m, 2H), 7.34 (dd, J = 8.0, 1.7 Hz, 1H), 6.90 - 6.65 (m, 3H), 3.96 (d, J = 1.6 Hz, 3H), 2.16 (s, 1H), 1.86 (dd, J = 11.9, 5.6 Hz, 1H), 1.61 (s, 3H), 1.58 (s, 3H), 1.41 (t, J = 12.2 Hz, 1H), 0.76 (d, J = 6.4 Hz, 3H).
1999	496.19	1.21	
2000	499.40	1.55	
2001	499.31	2.89	
2002	552.40	1.87	1H NMR (400 MHz, DMSO-d6) δ 7.58

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(s, 1H), 7.52 - 7.33 (m, 2H), 7.25 (s, 1H), 7.02 (s, 1H), 6.43 (d, J = 22.8 Hz, 2H), 6.11 (s, 2H), 3.55 (q, J = 7.0 Hz, 4H), 3.09 - 2.90 (m, 2H), 2.26 (s, 3H), 2.19 - 2.05 (m, 2H), 1.82 (dd, J = 12.1, 5.7 Hz, 1H), 1.63 (s, 3H), 1.59 (s, 3H), 1.38 (t, J = 12.0 Hz, 1H), 1.15 (t, J = 6.9 Hz, 6H), 0.85 (d, J = 6.4 Hz, 3H).
2003	506.21	2.00	
2004	486.40	2.18	
2005	509.21	1.46	
2006	532.16	1.46	
2007	517.15	1.70	
2008	528.21	1.28	
2009	682.20	2.26	
2010	553.40	2.43	1H NMR (400 MHz, DMSO-d6) δ 12.48 (s, 1H), 7.80 (d, J = 7.9 Hz, 1H), 7.63 (t, J = 7.8 Hz, 1H), 7.41 (s, 1H), 7.27 (d, J = 7.9 Hz, 1H), 7.20 (d, J = 7.2 Hz, 1H), 7.16 (s, 1H), 6.68 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 4.05 (d, J = 6.7 Hz, 2H), 2.60 (q, J = 10.6, 8.9 Hz, 2H), 2.44 (s, 3H), 2.28 - 2.12 (m, 1H), 2.05 (dp, J = 13.3, 6.6 Hz, 1H), 1.88 (dd, J = 11.9, 5.6 Hz, 1H), 1.61 (s, 3H), 1.59 (s, 3H), 1.42 (t, J = 12.1 Hz, 1H), 0.97 (d, J = 6.7 Hz, 6H), 0.88 (d, J = 6.2 Hz, 3H).
2011	512.51	2.69	
2012	524.52	2.10	
2013	511.31	2.08	
2014	560.25	1.88	1H NMR (400 MHz, DMSO-d6) δ 12.47 (s, 1H), 8.42 (d, J = 2.4 Hz, 1H), 7.96 (d, J = 2.4 Hz, 1H), 7.78 (d, J = 8.2 Hz, 2H), 7.70 (d, J = 8.3 Hz, 2H), 7.67 - 7.61 (m, 1H), 7.37 (d, J = 16.5 Hz, 1H), 7.22 (d, J = 7.1 Hz, 1H), 7.19 (d, J = 16.5 Hz, 1H), 6.69 (d, J = 8.4 Hz, 1H), 6.54 (s, 2H), 2.69 - 2.59 (m, 1H), 2.59 - 2.52 (m, 1H), 2.29 - 2.11 (m, 1H), 1.85 (dd, J = 11.9, 5.6 Hz, 1H), 1.53 (s, 6H), 1.38 (d, J = 12.1 Hz, 1H), 0.86 (d, J = 6.3 Hz, 3H).
2015	485.20	2.31	
2016	490.32	1.47	
2017	494.31	1.52	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2018	528.19	1.54	
2019	515.51	3.21	
2020	494.30	1.64	1H NMR (400 MHz, DMSO-d6) δ 12.25 (s, 1H), 7.66 - 7.55 (m, 2H), 7.17 (dd, J = 7.3, 0.7 Hz, 1H), 6.66 (dd, J = 8.4, 0.7 Hz, 1H), 6.44 (s, 2H), 5.02 (p, J = 6.2 Hz, 1H), 2.56 (d, J = 10.6 Hz, 1H), 2.44 (t, J = 8.7 Hz, 1H), 2.25 - 2.15 (m, 1H), 1.88 (ddq, J = 24.1, 11.8, 6.1, 5.5 Hz, 2H), 1.51 (s, 3H), 1.44 (s, 3H), 1.37 (t, J = 12.0 Hz, 1H), 1.24 (d, J = 6.4 Hz, 3H), 0.89 (dd, J = 6.8, 4.5 Hz, 6H), 0.85 (d, J = 6.3 Hz, 3H).
2021	511.18	1.33	
2022	472.50	1.55	1H NMR (400 MHz, CDCl3) δ 15.20 (s, 1H), 8.47 (d, J = 8.2 Hz, 1H), 7.66 - 7.58 (m, 1H), 7.41 (dd, J = 12.8, 7.7 Hz, 2H), 7.12 (d, J = 8.6 Hz, 1H), 4.23 (s, 1H), 3.83 (d, J = 6.6 Hz, 1H), 2.53 - 2.37 (m, 1H), 2.22 - 2.03 (m, 1H), 1.80 (dd, J = 18.4, 9.3 Hz, 1H), 1.56 (s, 3H), 1.27 (d, J = 13.6 Hz, 9H), 1.25 - 1.16 (m, 3H), 0.70 (d, J = 4.1 Hz, 1H), 0.53 (d, J = 7.6 Hz, 2H), 0.15 (s, 2H).
2023	542.23	2.65	
2024	488.20	2.49	
2025	474.99	2.98	
2026	472.29	1.98	
2027	613.50	1.72	1H NMR (400 MHz, DMSO-d6) δ 12.26 (s, 1H), 7.66 - 7.57 (m, 1H), 7.29 (s, 1H), 7.18 (d, J = 7.2 Hz, 1H), 6.67 (d, J = 8.3 Hz, 1H), 6.46 (s, 2H), 5.54 (s, 1H), 4.82 (h, J = 6.1 Hz, 1H), 4.12 - 3.84 (m, 2H), 3.67 - 3.51 (m, 2H), 2.56 (d, J = 10.5 Hz, 1H), 2.41 - 2.24 (m, 2H), 2.24 - 2.11 (m, 1H), 1.83 (dd, J = 11.8, 5.6 Hz, 1H), 1.54 (s, 3H), 1.52 (s, 3H), 1.41 - 1.32 (m, 2H), 1.31 (s, 9H), 1.23 (d, J = 6.3 Hz, 3H), 1.21 (d, J = 6.3 Hz, 3H), 0.85 (d, J = 6.3 Hz, 3H).
2028	501.30	1.62	
2029	542.29	1.62	
2030	472.29	2.57	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2031	496.24	1.57	
2032	489.40	2.07	
2033	514.21	1.40	
2034	496.60	1.57	
2035	520.17	1.49	
2036	512.50	1.56	1H NMR (400 MHz, DMSO-d6) δ 12.48 (broad s, 1H), 8.27 (d, J = 2.5 Hz, 1H), 8.25 (dd, J = 4.8, 1.9 Hz, 1H), 8.03 (dd, J = 7.8, 1.9 Hz, 1H), 7.86 (d, J = 8.2 Hz, 1H), 7.08 (d, J = 8.2 Hz, 1H), 6.77 (dd, J = 7.8, 4.8 Hz, 1H), 6.69 (br s, 2H), 6.41 (d, J = 2.6 Hz, 1H), 2.52 (likely d, 2H, partially overlapped with DMSO), 2.46 (d, J = 8.8 Hz, 2H), 2.13 (br m, 1H), 1.95 (dt, J = 13.5, 6.8 Hz, 1H), 1.85 (dd, J = 12.0, 5.6 Hz, 1H), 1.56 (s, 3H), 1.54 (s, 3H), 1.39 (t, J = 12.1 Hz, 1H), 0.93 (d, J = 2.2 Hz, 3H), 0.92 (d, J = 2.1 Hz, 3H), 0.74 (d, J = 6.3 Hz, 3H).
2037	495.40	1.55	
2038	501.41	2.98	
2039	553.40	2.25	1H NMR (400 MHz, DMSO) δ 12.55 (s, 1H), 7.96 (d, J = 8.2 Hz, 1H), 7.86 (d, J = 8.2 Hz, 1H), 7.68 - 7.60 (m, 1H), 7.56 (s, 1H), 7.49 (d, J = 9.6 Hz, 1H), 7.20 (d, J = 7.3 Hz, 1H), 7.01 - 6.84 (m, 1H), 6.72 (d, J = 8.4 Hz, 1H), 6.55 (s, 2H), 5.62 (d, J = 1.7 Hz, 1H), 3.85 (d, J = 6.6 Hz, 2H), 2.66 (ddd, J = 14.3, 7.4, 1.8 Hz, 1H), 2.03 (dd, J = 13.2, 6.6 Hz, 1H), 1.95 (dd, J = 12.1, 7.5 Hz, 1H), 1.38 - 1.28 (m, 4H), 1.25 (s, 3H), 0.99 (d, J = 6.7 Hz, 6H), 0.93 (d, J = 6.9 Hz, 3H).
2040	534.22	1.01	
2041	526.02	2.29	1H NMR (400 MHz, DMSO) δ 12.56 (s, 1H), 9.42 (s, 1H), 8.15 - 8.13 (dd, 1H), 7.78 (t, J = 7.80, 1H), 7.67 - 7.64 (d, 1H), 7.42 - 7.37 (m, 2H), 7.33 - 7.30 (dd, 1H), 7.0 - 6.97 (d, 1H), 6.83 - 6.80 (d, 1H), 6.53 - 6. 50 (dd, 1H), 3.72 (s, 3H), 3.70 (s, 3H), 1.99 - 1.94 (m, 1H), 1.80 - 1.73 (m, 1H), 1.51 (s, 3H), 1.46 (s, 3H), 1.32 (t, J = 11.91,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 0.77 - 0.74 (d, 3H)
2042	533.50	1.47	
2043	496.30	1.53	1H NMR (400 MHz, DMSO-d6) δ 12.51 (s, 1H), 11.99 (s, 1H), 7.81 (d, J = 8.2 Hz, 1H), 7.75 (d, J = 8.5 Hz, 1H), 7.63 (dd, J = 8.4, 7.3 Hz, 1H), 7.34 (d, J = 8.2 Hz, 1H), 7.21 (d, J = 7.3 Hz, 1H), 6.76 - 6.65 (m, 3H), 2.68 (d, J = 8.7 Hz, 2H), 2.27 (s, 3H), 2.32 - 2.14 (m, 1H), 2.07 (s, 2H), 1.90 (dd, J = 11.9, 5.6 Hz, 1H), 1.52 (d, J = 3.1 Hz, 6H), 1.44 (t, J = 12.1 Hz, 1H), 0.88 (d, J = 6.3 Hz, 3H).
2044	458.18	2.35	
2045	476.25	1.38	
2046	500.18	1.40	
2047	516.40	1.23	1H NMR (400 MHz, Methanol-d4) δ 8.26 - 8.11 (m, 2H), 7.89 (dd, J = 12.7, 7.8 Hz, 1H), 6.90 - 6.73 (m, 2H), 3.28 - 3.13 (m, 1H), 3.07 - 2.93 (m, 1H), 2.92 - 2.75 (m, 1H), 2.57 - 2.40 (m, 1H), 2.09 - 1.96 (m, 1H), 1.94 - 1.81 (m, 1H), 1.77 - 1.35 (m, 12H), 1.17 - 0.89 (m, 9H).
2048	534.22	1.08	
2049	544.33	2.68	
2050	502.33	1.44	1H NMR (400 MHz, DMSO) δ 12.06 (s, 1H), 8.15 (s, 3H), 7.73 (dd, J = 7.7, 1.8 Hz, 1H), 7.67 (d, J = 8.0 Hz, 1H), 7.40 - 7.34 (m, 1H), 7.21 (d, J = 8.0 Hz, 1H), 7.12 (d, J = 7.8 Hz, 1H), 7.09 - 7.03 (m, 1H), 4.11 (q, J = 7.0 Hz, 2H), 3.88 - 3.79 (m, 1H), 3.76 (dd, J = 10.2, 6.6 Hz, 1H), 3.72 - 3.62 (m, 1H), 3.58 (dd, J = 15.0, 8.8 Hz, 1H), 3.41 (dd, J = 10.2, 4.8 Hz, 1H), 3.09 (t, J = 10.1 Hz, 1H), 2.98 - 2.89 (m, 1H), 2.35 (dd, J = 10.9, 5.8 Hz, 1H), 2.28 - 2.16 (m, 1H), 1.96 (ddd, J = 17.2, 12.6, 6.5 Hz, 2H), 1.61 (s, 3H), 1.58 (s, 3H), 1.52 (d, J = 12.1 Hz, 1H), 1.34 (t, J = 6.9 Hz, 3H), 1.04 (d, J = 6.3 Hz, 3H).
2051	529.36	1.13	1H NMR (400 MHz, DMSO) δ 12.59 (s, 1H), 12.53 (s, 1H), 8.25 (dd, J = 7.3, 2.2 Hz, 1H), 7.82 (s, 1H), 7.70 (d, J =

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			8.0 Hz, 1H), 7.49 (dd, J = 9.9, 3.2 Hz, 1H), 7.25 - 7.17 (m, 2H), 7.14 (dd, J = 9.1, 4.7 Hz, 1H), 6.49 - 6.42 (m, 1H), 4.09 (q, J = 7.0 Hz, 2H), 2.81 - 2.68 (m, 2H), 2.23 (s, 1H), 1.87 (dd, J = 11.8, 5.5 Hz, 1H), 1.55 (s, 6H), 1.43 (t, J = 12.1 Hz, 1H), 1.34 (t, J = 6.9 Hz, 3H), 0.91 (d, J = 6.3 Hz, 3H).
2052	498.53	1.08	
2053	458.49	2.14	
2054	505.20	1.37	
2055	512.51	2.62	1H NMR (400 MHz, DMSO) δ 12.69 (s, 1H), 8.16 - 8.14 (dd, 1H), 7.88 (t, J = 8.02, 1H), 7.68 - 7.65 (d, 1H), 7.50 - 7.47 (dd, 1H), 7.06 - 7.03 (dd, 1H), 6.61 - 6.58 (dd, 1H), 4.95 (m, 1H), 3.60 (m, 1H), 3.46 - 3.39 (m, 1H), 2.11 (m, 5H), 1.87 - 1.77 (m, 1H), 1.51 - 1.29 (m, 7H), 0.78 - 0.75 (d, 1.5H), 0.71 - 0.68 (d, 1.5H)
2056	494.30	1.06	
2057	501.11	3.27	
2058	570.20	1.99	
2059	527.22	3.17	
2060	538.33	2.62	
2061	535.02	2.71	
2062	556.20	1.96	1H NMR (400 MHz, DMSO) δ 12.66 (s, 1H), 7.79 (d, J = 7.8 Hz, 1H), 7.63 (t, J = 10.7 Hz, 1H), 7.45 (d, J = 11.6 Hz, 3H), 7.20 (d, J = 6.5 Hz, 1H), 6.90 (d, J = 10.7 Hz, 1H), 6.70 (d, J = 7.1 Hz, 1H), 6.44 (s, 2H), 3.84 (d, J = 6.5 Hz, 2H), 3.50 (s, 1H), 3.38 (s, 1H), 3.12 (d, J = 13.0 Hz, 1H), 2.96 (s, 1H), 2.08 - 1.98 (m, 1H), 1.80 (s, 2H), 1.41 - 1.37 (m, 1H), 1.30 - 1.22 (m, 1H), 1.00 (d, J = 6.7 Hz, 7H), 0.86 (d, J = 6.5 Hz, 4H), 0.76 (d, J = 6.6 Hz, 3H).
2063	498.34	1.82	
2064	514.31	2.39	
2065	526.18	1.43	
2066	500.23	1.42	
2067	499.40	1.50	
2068	467.19	1.43	1H NMR (400 MHz, DMSO) δ 10.68 (s, 1H), 8.32 - 8.30 (dd, 1H), 8.18 - 8.16

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(dd, 1H), 8.11 - 8.08 (d, 1H), 8.01 (t, J = 7.91, 1H), 7.84 - 7.75 (m, 2H), 7.66 - 7.63 (dd, 2H), 7.06 (t, J = 6.26, 1H), 6.62 - 6.59 (dd, 1 H), 2.46 - 2.43(d, 1H), 1.93 - 1.89 (m, 1H), 1.78 - 1.72 (m, 1H), 1.44 (s, 3H), 1.43 (s, 3H), 1.27 (t, J = 11.91, 1H), 0.68 - 0.66 (d, 3H)
2069	574.40	1.51	
2070	542.29	1.53	
2071	528.22	1.53	
2072	524.22	2.30	1H NMR (400 MHz, DMSO) δ 12.49 (s, 1H), 8.15 - 8.13 (dd, 1H), 7.69 - 7.66 (dd, 1H), 7.60 (t, J = 7.80, 1H), 7.25 - 7.15 (m, 4H), 6.85 - 6.82 (d, 2H), 6.74 - 6.71 (d, 1H), 6.53 - 6.50 (dd, 1H), 3.71 (s, 3H), 3.45 - 3.36 (m, 2H), 2.79 (t, J = 7.47, 2H), 2.12 - 2.08 (m, 1H), 1.85 - 1.78 (m, 1H), 1.50 (s, 3H), 1.48 (s, 3H), 1.36 (t, J = 11.65, 1H), 0.82 - 0.80 (d, 3H)
2073	495.30	1.53	
2074	517.32	1.36	
2075	461.19	2.68	
2076	492.40	2.00	
2077	540.40	1.46	1H NMR (400 MHz, Methanol-d4) δ 8.50 (dd, J = 7.7, 1.8 Hz, 1H), 8.27 (d, J = 8.0 Hz, 1H), 8.13 (dd, J = 5.9, 1.8 Hz, 1H), 7.27 (d, J = 8.0 Hz, 1H), 7.00 (dd, J = 7.7, 5.8 Hz, 1H), 3.74 (t, J = 10.0 Hz, 1H), 3.33 - 3.25 (m, 1H, obscured by solvent peak), 2.88 - 2.68 (m, 2H), 2.28 - 2.18 (m, 2H), 2.14 - 2.06 (m, 4H), 1.92 - 1.82 (m, 1H), 1.75 - 1.60 (m, 2H), 1.57 - 1.45 (m, 6H), 1.36 (s, 3H), 1.21 (d, J = 6.6 Hz, 3H).
2078	550.33	2.02	
2079	583.30	1.59	
2080	497.40	1.36	1H NMR (400 MHz, DMSO-d6) δ 12.54 (s, 1H), 8.24 (dd, J = 7.2, 2.1 Hz, 1H), 8.05 - 7.97 (m, 2H), 7.82 (s, 1H), 7.72 (d, J = 8.0 Hz, 1H), 7.12 (d, J = 8.1 Hz, 1H), 7.09 - 7.00 (m, 2H), 6.44 (t, J = 6.8 Hz, 1H), 3.81 (s, 2H), 2.76 - 2.66

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(m, 2H), 2.22 (s, 1H), 1.88 (dd, J = 11.9, 5.6 Hz, 1H), 1.61 (s, 6H), 1.44 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.2 Hz, 3H).
2081	553.20	2.26	
2082	481.30	1.56	1H NMR (400 MHz, DMSO-d6) δ 10.99 (s, 1H), 8.22 - 8.18 (m, 2H), 8.06 (t, J = 8.13, 1H), 7.81 - 7.62 (m, 4H), 7.00 - 6.98 (d, 1H), 6.65 - 6.62 (dd, 1H), 2.30 (s, 3H), 1.87 - 1.72 (m, 2H), 1.45 (s, 3H), 1.42 (s, 3H), 1. 26 (t, J = 11.87, 1H), 0.65 - 0.63 (d, 3H)
2083	523.32	2.14	
2084	499.40	1.59	
2085	538.29	1.23	
2086	511.30	2.13	1H NMR (400 MHz, DMSO) δ 8.65 (d, J = 2.2 Hz, 1H), 8.09 (d, J = 2.3 Hz, 1H), 7.57 (d, J = 7.6 Hz, 1H), 7.49 - 7.36 (m, 1H), 6.98 (dd, J = 7.5, 2.8 Hz, 2H), 6.47 (d, J = 8.2 Hz, 1H), 6.17 (s, 2H), 3.92 (s, 3H), 3.04 - 2.85 (m, 2H), 2.20 (s, 3H), 2.09 (d, J = 17.6 Hz, 1H), 1.81 (dd, J = 11.6, 5.5 Hz, 1H), 1.61 (s, 3H), 1.56 (s, 3H), 1.36 (t, J = 12.0 Hz, 1H), 0.82 (d, J = 6.3 Hz, 3H).
2087	549.43	2.09	
2088	496.24	1.27	
2089	538.34	1.88	
2090	524.52	2.54	
2091	527.16	1.45	
2092	450.50	1.84	1H NMR (400 MHz, DMSO-d6) δ 12.55 (s, 1H), 8.02 (ddt, J = 9.9, 7.2, 1.6 Hz, 4H), 7.77 - 7.70 (m, 2H), 7.70 - 7.62 (m, 2H), 7.53 - 7.46 (m, 2H), 7.46 - 7.40 (m, 1H), 7.21 (d, J = 8.0 Hz, 1H), 2.45 (t, J = 10.3 Hz, 1H), 2.36 (dd, J = 10.2, 7.1 Hz, 1H), 2.19 - 2.08 (m, 1H), 1.83 (dd, J = 12.0, 5.6 Hz, 1H), 1.58 (d, J = 15.4 Hz, 6H), 1.38 (t, J = 12.1 Hz, 1H), 0.67 (d, J = 6.3 Hz, 3H).
2093	494.31	1.69	
2094	520.32	2.62	
2095	538.34	1.84	1H NMR (400 MHz, DMSO-d6) δ 12.46 (s, 1H), 8.46 (s, 1H), 7.84 (d, J = 2.5 Hz, 1H), 7.63 (t, J = 7.9 Hz, 1H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			7.43 (d, J = 12.2 Hz, 2H), 7.22 (d, J = 7.2 Hz, 1H), 6.98 (d, J = 8.5 Hz, 1H), 6.68 (d, J = 8.5 Hz, 1H), 6.48 (s, 2H), 3.97 (t, J = 6.4 Hz, 2H), 3.35 (d, J = 7.2 Hz, 1H), 2.55 (d, J = 8.4 Hz, 2H), 2.23 (s, 3H), 1.84 (dd, J = 11.8, 5.7 Hz, 1H), 1.76 (p, J = 7.0 Hz, 2H), 1.54 (s, 3H), 1.52 (s, 3H), 1.39 (t, J = 12.0 Hz, 1H), 1.02 (t, J = 7.4 Hz, 3H), 0.86 (d, J = 6.3 Hz, 3H).
2096	491.40	1.17	
2097	484.30	1.41	1H NMR (400 MHz, DMSO-d6) δ 12.42 (s, 1H), 8.23 (dd, J = 4.8, 1.8 Hz, 1H), 8.03 (d, J = 7.9 Hz, 1H), 7.60 (dd, J = 7.9, 3.2 Hz, 1H), 6.76 (dd, J = 7.8, 4.8 Hz, 1H), 6.55 (dd, J = 7.9, 5.9 Hz, 1H), 3.24 - 3.14 (m, 1H), 2.46 - 2.39 (m, 2H, obscured by solvent peak), 2.31 - 2.24 (m, 1H), 2.20 - 2.04 (m, 1H), 1.89 - 1.67 (m, 3H), 1.59 - 1.54 (m, 1H), 1.52 (d, J = 5.1 Hz, 6H), 1.49 - 0.95 (m, 7H), 0.74 (d, J = 6.3 Hz, 3H).
2098	506.21	2.66	
2099	517.32	2.55	
2100	492.30	1.57	1H NMR (400 MHz, DMSO) δ 12.28 (s, 1H), 7.66 - 7.50 (m, 2H), 7.17 (d, J = 7.2 Hz, 1H), 6.66 (d, J = 8.3 Hz, 1H), 5.45 - 5.31 (m, 1H), 2.58 - 2.53 (m, 1H), 2.47 - 2.40 (m, 1H), 2.20 (dp, J = 18.4, 6.1 Hz, 1H), 1.99 - 1.79 (m, 4H), 1.78 - 1.65 (m, 4H), 1.65 - 1.55 (m, 2H), 1.53 (s, 3H), 1.46 (s, 3H), 1.37 (t, J = 12.0 Hz, 1H), 0.85 (d, J = 6.2 Hz, 3H).
2101	498.30	1.37	
2102	484.30	2.09	
2103	528.17	1.51	
2104	546.23	1.23	1H NMR (400 MHz, DMSO) δ 12.45 (s, 1H), 7.99 (s, 1H), 7.94 (d, J = 8.5 Hz, 1H), 7.79 (d, J = 8.0 Hz, 1H), 7.63 (t, J = 7.9 Hz, 1H), 7.54 (d, J = 8.5 Hz, 1H), 7.22 (t, J = 8.0 Hz, 2H), 6.69 (d, J = 8.4 Hz, 1H), 6.50 (s, 2H), 2.63 - 2.56 (m, 2H), 2.21 (q, J = 9.7, 7.7 Hz, 2H), 1.88 (dd, J = 11.8, 5.5 Hz, 1H), 1.60 (d, J =

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			5.4 Hz, 6H), 1.42 (t, J = 12.0 Hz, 1H), 0.88 (d, J = 6.1 Hz, 3H).
2105	502.26	1.41	
2106	520.32	2.21	
2107	548.23	2.68	
2108	509.21	2.57	
2109	473.19	2.72	
2110	494.31	1.45	
2111	547.33	2.45	
2112	536.50	1.40	
2113	532.50	2.29	
2114	512.30	1.18	1H NMR (400 MHz, Methanol-d4) δ 7.94 (d, J = 8.0 Hz, 1H), 7.65 - 7.59 (m, 2H), 7.41 (d, J = 8.0 Hz, 1H), 7.35 (d, J = 7.2 Hz, 1H), 6.82 (d, J = 2.5 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 3.99 (d, J = 7.3 Hz, 2H), 3.03 (t, J = 9.3 Hz, 1H), 2.91 (t, J = 10.3 Hz, 1H), 2.41 (s, 1H), 2.29 - 2.16 (m, 1H), 1.97 (dd, J = 13.3, 7.1 Hz, 1H), 1.63 - 1.53 (m, 7H), 1.34 (d, J = 9.5 Hz, 1H), 1.03 (d, J = 6.4 Hz, 3H), 0.93 (d, J = 6.6 Hz, 6H).
2115	539.23	2.51	
2116	458.18	2.27	
2117	514.40	2.02	1H NMR (400 MHz, DMSO) δ 7.77 (d, J = 7.9 Hz, 1H), 7.64 (dd, J = 8.3, 7.4 Hz, 1H), 7.48 - 7.40 (m, 2H), 7.32 (d, J = 7.9 Hz, 1H), 7.23 - 7.15 (m, 1H), 6.90 (d, J = 10.8 Hz, 1H), 6.69 (d, J = 8.4 Hz, 1H), 6.48 (s, 2H), 3.85 (dd, J = 14.4, 7.6 Hz, 4H), 3.30 (dd, J = 8.9, 5.8 Hz, 2H), 2.72 - 2.59 (m, 1H), 2.03 (d, J = 6.7 Hz, 1H), 1.15 (d, J = 6.8 Hz, 3H), 1.00 (d, J = 6.7 Hz, 6H).
2118	547.33	2.32	
2119	522.42	2.59	
2120	516.36	1.53	
2121	557.20	1.71	1H NMR (400 MHz, DMSO) δ 8.53 (s, 1H), 7.73 (s, 3H), 7.70 - 7.63 (m, 1H), 7.59 (d, J = 9.7 Hz, 1H), 7.21 (d, J = 7.2 Hz, 1H), 7.07 (d, J = 10.6 Hz, 1H), 6.73 (d, J = 8.4 Hz, 1H), 3.86 (d, J = 6.6 Hz, 2H), 2.86 - 2.74 (m, 1H), 2.69 (t, J = 10.9 Hz, 1H), 2.34 - 2.18 (m, 1H), 2.13 - 1.99 (m, 1H), 1.95 (dd, J = 11.9, 5.5

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			Hz, 1H), 1.63 (d, J = 23.9 Hz, 6H), 1.50 (t, J = 12.2 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.90 (d, J = 6.3 Hz, 3H).
2122	538.33	2.56	
2123	534.42	2.52	
2124	514.15	1.42	
2125	505.22	1.38	
2126	530.24	1.57	
2127	494.31	1.53	
2128	518.30	1.93	1H NMR (400 MHz, DMSO) δ 12.55 (s, 1H), 7.87 (d, J = 8.0 Hz, 1H), 7.67 (dd, J = 15.7, 7.4 Hz, 1H), 7.57 - 7.33 (m, 3H), 7.21 (d, J = 7.2 Hz, 1H), 6.92 (dt, J = 10.7, 2.2 Hz, 1H), 6.76 - 6.66 (m, 1H), 6.51 (s, 2H), 5.38 (ddd, J = 61.0, 5.7, 2.8 Hz, 1H), 4.12 (ddd, J = 17.3, 11.5, 5.4 Hz, 2H), 3.85 (d, J = 6.5 Hz, 4H), 2.04 (dt, J = 13.3, 6.6 Hz, 1H), 1.00 (d, J = 6.7 Hz, 6H).
2129	571.38	1.58	
2130	475.29	2.83	
2131	474.09	1.63	
2132	480.28	0.99	
2133	480.50	2.09	1H NMR (400 MHz, DMSO-d6) δ 13.25 (s, 1H), 12.66 (s, 1H), 8.04 - 7.99 (m, 2H), 7.82 (d, J = 8.2 Hz, 1H), 7.77 - 7.70 (m, 2H), 7.70 - 7.63 (m, 2H), 7.36 (d, J = 8.3 Hz, 1H), 7.19 (d, J = 7.2 Hz, 1H), 6.84 - 6.78 (m, 1H), 2.62 - 2.51 (m, 2H), 2.19 (s, 3H), 2.14 (q, J = 5.8 Hz, 1H), 1.89 (dd, J = 11.9, 5.5 Hz, 1H), 1.50 (d, J = 3.3 Hz, 6H), 1.41 (t, J = 12.1 Hz, 1H), 0.66 (d, J = 6.3 Hz, 3H).
2134	568.33	1.62	
2135	462.39	1.74	
2136	516.41	2.64	1H NMR (400 MHz, DMSO-d6) δ 12.67 (s, 1H), 9.43 (s, 1H), 8.24 - 8.09 (m, 3H), 7.96 - 7.94 (dd, 1H), 7.86 (t, J = 8.02, 1H), 7.70 - 7.67 (d, 1H), 7.60 - 7.43 (m, 5H), 7.27 - 7.24 (d, 1H), 6.45 - 6.42 (dd, 1H), 2.07 - 1 .91 (m, 1H), 1.77 - 1.73 (m, 1H), 1.45 (s, 3H), 1.43 (s, 3H), 1.33 (t, J = 11.91, 1H), 0.79 - 0.77 (d, 3H)

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2137	512.51	2.56	
2138	529.20	1.64	
2139	516.20	1.31	1H NMR (400 MHz, DMSO) δ 12.48 (s, 1H), 7.70 - 7.52 (m, 2H), 7.17 (dd, J = 7.3, 0.5 Hz, 1H), 6.70 (dd, J = 9.7, 8.5 Hz, 2H), 5.19 (s, 2H), 3.84 (td, J = 11.1, 3.7 Hz, 2H), 3.38 - 3.18 (m, 2H), 2.84 (t, J = 10.0 Hz, 1H), 2.68 - 2.55 (m, 1H), 2.07 (s, 4H), 1.81 (t, J = 6.4 Hz, 1H), 1.65 (dd, J = 30.6, 10.5 Hz, 2H), 1.55 (d, J = 8.6 Hz, 6H), 1.49 (s, 3H), 1.28 (d, J = 6.6 Hz, 10H).
2140	525.42	1.65	
2141	534.60	1.23	
2142	522.42	2.85	1H NMR (400 MHz, DMSO) δ 12.63 (s, 1H), 9.52 (s, 1H), 8.13 - 8.11 (dd, 1H), 8.01 - 7.98 (d, 1H), 7.81 (t, J = 7.80, 1H), 7.71 - 7.68 (dd, 1H), 7.45 - 7.42 (d, 1H), 7.36 (s, 1H), 7.17 (t, J = 7.92, 1H), 7.06 - 6.97 (m, 2H), 6.52 - 6.49 (dd, 1H), 1.86 - 1.70 (m, 2H), 1.45 (s, 3H), 1.41 (s, 3H), 1.28 (m, 10H), 0.74 - 0.72 (d, 3H)
2143	550.33	2.55	
2144	535.50	1.76	1H NMR (400 MHz, DMSO-d6) δ 12.65 (d, J = 4.1 Hz, 1H), 12.53 (s, 1H), 8.40 (s, 1H), 8.34 (d, J = 8.0 Hz, 1H), 8.25 (dd, J = 7.3, 2.2 Hz, 1H), 7.78 (ddt, J = 24.0, 15.6, 7.9 Hz, 4H), 7.35 - 7.30 (m, 1H), 6.54 - 6.35 (m, 1H), 2.82 - 2.63 (m, 2H), 2.24 (s, 1H), 1.90 (dd, J = 12.0, 5.7 Hz, 1H), 1.61 (d, J = 3.3 Hz, 6H), 1.46 (t, J = 12.3 Hz, 1H), 0.91 (t, J = 7.0 Hz, 3H).
2145	501.11	2.34	
2146	500.21	2.30	
2147	471.39	2.56	
2148	497.30	1.49	
2149	528.19	1.49	1H NMR (400 MHz, DMSO-d6) δ 12.58 (s, 1H), 8.26 (dd, J = 4.8, 1.8 Hz, 1H), 8.07 (dd, J = 7.8, 1.9 Hz, 1H), 7.80 (d, J = 7.9 Hz, 1H), 7.44 (dd, J = 6.3, 3.2 Hz, 1H), 7.24 (dd, J = 11.1, 9.0 Hz, 1H), 7.09 (dd, J = 7.9, 2.2 Hz, 1H), 7.01 (dt, J = 9.0, 3.5 Hz, 1H), 6.80 (dd,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			J = 7.9, 4.8 Hz, 3H), 4.04 (q, J = 6.9 Hz, 2H), 2.55 (dd, J = 10.1, 7.3 Hz, 2H), 2.15 (td, J = 11.6, 10.3, 5.8 Hz, 1H), 1.84 (dd, J = 11.8, 5.6 Hz, 1H), 1.57 (s, 3H), 1.54 (s, 3H), 1.40 (t, J = 12.0 Hz, 1H), 1.33 (t, J = 6.9 Hz, 3H), 0.76 (d, J = 6.3 Hz, 3H).
2150	446.40	1.27	1H NMR (400 MHz, Acetonitrile-d3) δ 8.27 (d, J = 8.0 Hz, 1H), 8.24 (dd, J = 7.7, 1.8 Hz, 1H), 8.09 (dd, J = 5.3, 1.8 Hz, 1H), 7.20 (d, J = 8.0 Hz, 1H), 6.81 (dd, J = 7.7, 5.3 Hz, 1H), 3.76 (dd, J = 11.5, 8.8 Hz, 1H), 3.24 (dd, J = 11.5, 8.2 Hz, 1H), 2.81 - 2.71 (m, 1H), 2.69 (d, J = 7.2 Hz, 2H), 2.23 - 2.11 (m, 3H), 1.86 (dd, J = 12.7, 10.2 Hz, 1H), 1.49 (s, 3H), 1.29 (s, 3H), 1.20 (d, J = 6.7 Hz, 3H), 0.93 (d, J = 6.7 Hz, 6H).
2151	531.12	2.71	
2152	488.50	2.10	
2153	539.50	1.87	
2154	495.40	1.59	1H NMR (400 MHz, DMSO-d6) δ 12.55 (s, 1H), 12.52 (d, J = 6.4 Hz, 1H), 8.24 (dd, J = 7.3, 2.1 Hz, 1H), 7.83 (s, 2H), 7.77 (d, J = 8.1 Hz, 1H), 7.72 (d, J = 8.0 Hz, 1H), 7.24 (d, J = 8.0 Hz, 1H), 7.14 (d, J = 8.0 Hz, 1H), 6.44 (t, J = 6.9 Hz, 1H), 2.77 - 2.64 (m, 2H), 2.38 - 2.13 (m, 8H), 1.88 (dd, J = 12.1, 5.7 Hz, 1H), 1.61 (s, 6H), 1.44 (t, J = 12.4 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).
2155	536.30	2.01	1H NMR (400 MHz, DMSO) δ 12.65 (s, 1H), 7.96 (d, J = 8.0 Hz, 1H), 7.67 (dd, J = 8.4, 7.3 Hz, 1H), 7.59 - 7.46 (m, 3H), 7.22 (dd, J = 7.3, 0.6 Hz, 1H), 6.93 (dt, J = 10.8, 2.3 Hz, 1H), 6.76 - 6.68 (m, 1H), 6.54 (s, 2H), 4.15 (t, J = 12.5 Hz, 4H), 3.85 (d, J = 6.5 Hz, 2H), 2.05 (dt, J = 13.3, 4.4 Hz, 1H), 1.00 (d, J = 6.7 Hz, 6H).
2156	516.36	1.55	
2157	506.21	1.34	
2158	512.16	1.32	
2159	503.30	1.51	1H NMR (400 MHz, DMSO-d6) δ 12.64 (s, 1H), 12.54 (d, J = 6.0 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 8.25 (dd, J = 7.2, 2.3 Hz, 1H), 7.91 (td, J = 8.9, 6.7 Hz, 1H), 7.83 (dd, J = 7.1, 4.6 Hz, 1H), 7.77 - 7.71 (m, 1H), 7.37 (ddd, J = 11.9, 9.3, 2.6 Hz, 1H), 7.25 (td, J = 8.5, 2.6 Hz, 1H), 6.96 (dd, J = 7.8, 2.5 Hz, 1H), 6.45 (t, J = 6.8 Hz, 1H), 2.81 - 2.63 (m, 2H), 2.30 - 2.14 (m, 1H), 1.91 - 1.81 (m, 1H), 1.54 (d, J = 1.9 Hz, 6H), 1.43 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.4 Hz, 3H).
2160	495.40	2.16	
2161	501.11	2.53	
2162	503.40	1.52	1H NMR (400 MHz, DMSO-d6) δ 12.67 (s, 1H), 12.55 (s, 1H), 8.25 (dd, J = 7.3, 2.2 Hz, 1H), 7.84 (dt, J = 7.5, 3.7 Hz, 1H), 7.78 (d, J = 7.9 Hz, 1H), 7.65 (ddt, J = 8.2, 6.7, 1.6 Hz, 1H), 7.55 - 7.45 (m, 1H), 7.35 (tdd, J = 8.1, 5.1, 1.5 Hz, 1H), 7.00 (dd, J = 7.9, 2.3 Hz, 1H), 6.45 (t, J = 6.8 Hz, 1H), 2.81 - 2.68 (m, 2H), 2.25 (tt, J = 11.7, 6.4 Hz, 1H), 1.87 (dd, J = 11.9, 5.6 Hz, 1H), 1.54 (s, 6H), 1.44 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).
2163	508.16	1.42	
2164	496.28	0.93	
2165	519.12	2.58	
2166	480.10	2.02	
2167	447.08	2.52	
2168	510.41	2.20	1H NMR (400 MHz, DMSO-d6) δ 12.49 (s, 1H), 8.16 - 8.14 (dd, 1H), 7.68 - 7.60 (m, 3H), 7.23 - 7.20 (d, 1H), 7.16 - 7.13 (d, 1H), 6.97 - 6.93 (m, 2H), 6.79 - 6.73 (m 2H), 6.57 - 6.54 (dd, 1H), 4.47 (s, 2H), 3.66 (s, 3H) , 2.44 - 2.33 (m, 1H), 2.05 - 1.98 (m, 1H), 1.82 - 1.75 (m, 1H), 1.50 (s, 3H), 1.47 (s, 3H), 1.33 (t, J = 12.09, 1H), 0.76 - 0.74 (d, 3H)
2169	525.42	2.44	1H NMR (400 MHz, DMSO-d6) δ 12.82 (s, 1H), 8.19 - 8.17 (dd, 1H), 8.03 (t, J = 7.80, 1H), 7.77 - 7.74 (d, 2H), 7.16 - 7.13 (d, 1H), 7.09 (s, 1H), 7.05 - 7.02 (dd, 1H), 6.83 - 6.80 (d, 1H), 6.64 - 6.61 (dd, 1H), 5.99 (s , 1H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			5.94 (s, 1H), 5.26, (s, 1H), 5.25 (s, 1H), 2.27 (s, 1H), 2.24 (s, 1H), 2.09 - 2.01 (m, 1H), 1.82 - 1.75 (m, 1H), 1.51 (s, 3H), 1.46 (s, 3H), 1.35 (t, J = 11.86, 1H), 0.67 - 0.65 (d, 3H)
2170	539.40	2.55	1H NMR (400 MHz, DMSO-d6) δ 8.63 (d, J = 2.4 Hz, 1H), 8.09 (d, J = 2.6 Hz, 1H), 7.60 (bs, 1H), 7.49 (bs, 1H), 7.01 (bs, 2H), 6.52 (bs, 1H), 6.24 (bs, 2H), 5.32 (p, J = 6.1 Hz, 1H), 3.01 - 2.73 (m, 2H), 2.17 (s, 3H), 2.15 - 2.05 (m, 1H), 1.82 (dd, J = 11.8, 5.6 Hz, 1H), 1.61 (s, 3H), 1.57 (s, 3H), 1.42 - 1.33 (m, 1H), 1.32 (d, J = 6.2 Hz, 6H), 0.82 (d, J = 6.3 Hz, 3H).
2171	570.50	2.19	1H NMR (400 MHz, DMSO) δ 12.51 (s, 1H), 7.77 (d, J = 7.9 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.49 (d, J = 2.0 Hz, 1H), 7.44 (ddd, J = 10.1, 2.2, 1.4 Hz, 1H), 7.30 (d, J = 7.9 Hz, 1H), 7.22 (dd, J = 7.3, 0.6 Hz, 1H), 6.93 - 6.83 (m, 1H), 6.72 (dd, J = 8.4, 0.6 Hz, 1H), 6.48 (s, 2H), 3.84 (d, J = 6.6 Hz, 2H), 3.23 (t, J = 5.4 Hz, 2H), 3.14 - 3.04 (m, 2H), 2.05 (dd, J = 11.8, 5.1 Hz, 1H), 1.72 (s, 2H), 1.57 - 1.49 (m, 2H), 1.30 - 1.20 (m, 2H), 0.99 (d, J = 6.7 Hz, 6H), 0.83 (s, 6H).
2172	508.31	1.55	
2173	515.51	3.19	
2174	497.30	1.48	
2175	550.33	1.25	
2176	481.50	1.52	
2177	526.02	2.42	
2178	556.20	1.96	
2179	526.32	2.25	
2180	522.40	1.86	
2181	487.30	2.37	
2182	533.22	2.72	
2183	500.23	1.49	
2184	480.50	1.57	
2185	492.18	1.47	
2186	495.40	2.60	
2187	522.42	2.86	
2188	459.39	2.57	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2189	514.17	1.42	
2190	524.52	2.32	
2191	528.50	1.50	1H NMR (400 MHz, DMSO-d6) δ 12.46 (broad s, 1H), 8.25 (dd, J = 4.8, 1.8 Hz, 1H), 8.20 (d, J = 2.8 Hz, 1H), 8.03 (dd, J = 7.9, 1.8 Hz, 1H), 7.84 (d, J = 8.2 Hz, 1H), 6.94 (d, J = 8.3 Hz, 1H), 6.77 (dd, J = 7.8, 4.8 Hz, 1H), 6.70 (br s overlapped with previous signal, 2H), 6.12 (d, J = 2.7 Hz, 1H), 3.98 (d, J = 6.6 Hz, 2H), 2.45 (d, J = 8.7 Hz, 2H), 2.20-2.01 (m, 2H), 1.85 (dd, J = 11.9, 5.6 Hz, 1H), 1.55 (s, 3H), 1.53 (s, 3H), 1.38 (t, J = 12.2 Hz, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.74 (d, J = 6.3 Hz, 3H).
2192	544.40	1.15	1H NMR (400 MHz, DMSO-d6) δ 12.45 (s, 1H), 8.23 (dd, J = 4.8, 1.9 Hz, 1H), 8.19 (d, J = 2.8 Hz, 1H), 8.02 (dd, J = 7.8, 1.9 Hz, 1H), 7.83 (d, J = 8.3 Hz, 1H), 6.93 (d, J = 8.2 Hz, 1H), 6.76 (dd, J = 7.8, 4.7 Hz, 1H), 6.67 (broad s, 2H), 6.11 (d, J = 2.8 Hz, 1H), 4.57 (br s, 1H), 4.19 (ddd, J = 9.9, 5.9, 2.4 Hz, 1H), 4.03 (ddd, J = 9.8, 6.5, 3.2 Hz, 1H), 3.41 (br s, 2H), 2.47 (d, J = 8.4 Hz, 2H), 2.19 - 2.06 (m, 1H), 2.06 - 1.92 (m, 1H), 1.85 (dd, J = 11.9, 5.5 Hz, 1H), 1.55 (s, 3H), 1.53 (s, 3H), 1.38 (t, J = 12.1 Hz, 1H), 0.96 (d, J = 6.9 Hz, 3H), 0.74 (d, J = 6.3 Hz, 3H).
2193	519.42	2.24	
2194	540.20	1.67	
2195	526.13	1.39	
2196	470.30	1.25	1H NMR (400 MHz, DMSO-d6) δ 8.27 (s, 1H), 7.69 (dd, J = 8.5, 7.3 Hz, 1H), 7.62 - 7.54 (m, 1H), 7.38 - 7.30 (m, 2H), 7.26 (d, J = 7.3 Hz, 1H), 6.76 (d, J = 8.4 Hz, 1H), 3.84 (s, 3H), 3.09 - 2.98 (m, 1H), 2.77 (t, J = 9.9 Hz, 1H), 2.34 - 2.29 (m, 1H), 1.93 (dd, J = 11.9, 6.0 Hz, 1H), 1.57 (d, J = 13.2 Hz, 6H), 1.47 (t, J = 12.1 Hz, 1H), 0.93 (d, J = 6.3 Hz, 3H).
2197	460.29	2.34	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2198	537.43	1.62	
2199	490.32	1.45	
2200	432.37	2.23	
2201	584.40	2.16	1H NMR (400 MHz, DMSO) δ 7.55 (d, J = 7.7 Hz, 1H), 7.46 (s, 1H), 7.38 (dd, J = 18.8, 10.8 Hz, 2H), 7.06 (d, J = 7.7 Hz, 1H), 7.02 (d, J = 7.3 Hz, 1H), 6.79 (d, J = 10.7 Hz, 1H), 6.42 (d, J = 8.2 Hz, 1H), 6.01 (s, 2H), 3.82 (d, J = 6.6 Hz, 2H), 3.39 (t, J = 10.7 Hz, 1H), 3.23 - 3.14 (m, 1H), 2.05 (dt, J = 13.4, 6.8 Hz, 1H), 1.86 (d, J = 11.5 Hz, 1H), 1.79 (s, 1H), 1.61 (d, J = 22.2 Hz, 6H), 1.45 (t, J = 11.4 Hz, 1H), 1.36 (d, J = 6.8 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.86 (dd, J = 16.6, 6.5 Hz, 6H).
2202	500.21	2.74	
2203	496.60	1.72	
2204	614.20	2.11	1H NMR (400 MHz, DMSO) δ 12.60 (s, 1H), 7.78 (d, J = 7.9 Hz, 1H), 7.65 (t, J = 7.8 Hz, 1H), 7.50 (d, J = 9.2 Hz, 2H), 7.37 (d, J = 7.9 Hz, 1H), 7.22 (d, J = 7.3 Hz, 1H), 6.89 (d, J = 10.7 Hz, 1H), 6.71 (d, J = 8.5 Hz, 1H), 6.48 (s, 2H), 4.51 (dd, J = 7.7, 3.8 Hz, 1H), 3.84 (d, J = 6.4 Hz, 2H), 2.96 - 2.72 (m, 2H), 2.27 - 2.10 (m, 1H), 2.09 - 1.98 (m, 1H), 1.97 - 1.86 (m, 1H), 1.80 (qd, J = 10.5, 5.7 Hz, 2H), 1.29 (s, 9H), 1.00 (d, J = 6.7 Hz, 6H).
2205	498.53	1.05	
2206	510.41	2.30	
2207	511.30	1.48	1H NMR (400 MHz, DMSO-d6) δ 12.52 (d, J = 10.1 Hz, 2H), 8.24 (dd, J = 7.2, 2.1 Hz, 1H), 8.02 - 7.93 (m, 2H), 7.82 (t, J = 6.6 Hz, 1H), 7.71 (d, J = 8.1 Hz, 1H), 7.11 (d, J = 8.1 Hz, 1H), 7.06 - 6.95 (m, 2H), 6.44 (t, J = 7.0 Hz, 1H), 4.09 (q, J = 7.0 Hz, 2H), 2.71 (dd, J = 12.1, 8.5 Hz, 2H), 2.23 (s, 1H), 1.88 (dd, J = 11.8, 5.6 Hz, 1H), 1.61 (s, 6H), 1.44 (t, J = 12.1 Hz, 1H), 1.34 (t, J = 6.9 Hz, 3H), 0.91 (d, J = 6.3 Hz, 3H).
2208	535.32	2.08	
2209	598.30	1.63	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2210	558.25	1.85	
2211	527.22	3.22	
2212	529.20	0.81	1H NMR (400 MHz, DMSO-d6) δ 12.31 (s, 1H), 8.87 - 8.78 (m, 2H), 7.92 (dd, J = 19.3, 6.1 Hz, 2H), 7.69 (dd, J = 10.6, 1.5 Hz, 1H), 7.60 (ddd, J = 8.4, 7.3, 3.4 Hz, 1H), 7.20 - 7.12 (m, 1H), 6.66 (dt, J = 8.4, 0.9 Hz, 1H), 6.44 (s, 1H), 6.14 (q, J = 6.7 Hz, 1H), 2.50 - 2.33 (m, 2H), 2.20 - 2.03 (m, 1H), 1.80 - 1.62 (m, 4H), 1.43 (d, J = 19.0 Hz, 3H), 1.35 - 1.12 (m, 1H), 0.86 - 0.73 (m, 6H).
2213	570.20	1.97	
2214	510.41	2.57	
2215	485.20	1.23	1H NMR (400 MHz, DMSO-d6) δ 8.25 (dd, J = 4.8, 1.8 Hz, 1H), 8.05 (dd, J = 7.8, 1.8 Hz, 1H), 7.77 (d, J = 7.8 Hz, 1H), 7.01 - 6.55 (m, 4H), 2.16 (s, 1H), 1.83 (dd, J = 12.1, 5.7 Hz, 1H), 1.48 (d, J = 4.1 Hz, 6H), 1.36 (t, J = 12.1 Hz, 1H), 0.75 (d, J = 6.3 Hz, 3H).
2216	614.20	2.11	1H NMR (400 MHz, DMSO) δ 12.60 (s, 1H), 7.78 (d, J = 7.9 Hz, 1H), 7.65 (t, J = 7.8 Hz, 1H), 7.50 (d, J = 8.8 Hz, 2H), 7.37 (d, J = 7.9 Hz, 1H), 7.22 (d, J = 7.2 Hz, 1H), 6.89 (d, J = 10.8 Hz, 1H), 6.71 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 4.63 - 4.34 (m, 1H), 3.84 (d, J = 6.5 Hz, 2H), 2.83 (t, J = 5.9 Hz, 2H), 2.27 - 2.11 (m, 1H), 2.10 - 1.98 (m, 1H), 1.93 (tt, J = 10.8, 5.6 Hz, 1H), 1.86 - 1.69 (m, 2H), 1.29 (s, 9H), 1.00 (d, J = 6.7 Hz, 6H).
2217	534.12	2.88	
2218	486.40	2.53	
2219	515.20	1.65	
2220	526.34	1.63	
2221	476.19	2.36	
2222	515.20	1.57	
2223	550.22	1.10	
2224	554.20	2.12	1H NMR (400 MHz, DMSO) δ 12.59 (s, 1H), 7.73 (d, J = 7.9 Hz, 1H), 7.64 (t, J = 6.7 Hz, 1H), 7.46 (s, 2H), 7.33 (d, J = 7.7 Hz, 1H), 7.22 (d, J = 7.1 Hz, 1H), 6.89 (d, J = 10.8 Hz, 1H), 6.70 (d, J =

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			8.3 Hz, 1H), 6.48 (s, 2H), 3.84 (d, J = 6.3 Hz, 2H), 3.52 (d, J = 11.3 Hz, 2H), 3.40 (d, J = 11.3 Hz, 3H), 2.30 (d, J = 4.4 Hz, 2H), 2.24 (dd, J = 12.3, 5.3 Hz, 1H), 1.92 (d, J = 4.5 Hz, 1H), 1.25 (d, J = 9.3 Hz, 1H), 1.04 - 0.94 (m, 6H), 0.74 (d, J = 6.2 Hz, 3H).
2225	486.33	1.45	
2226	529.32	1.56	
2227	538.32	1.22	
2228	551.60	2.40	
2229	494.28	1.50	
2230	466.30	1.38	1H NMR (400 MHz, DMSO-d6) δ 12.26 (s, 1H), 7.66 - 7.55 (m, 2H), 7.17 (dd, J = 7.3, 0.7 Hz, 1H), 6.66 (dd, J = 8.4, 0.8 Hz, 1H), 6.41 (s, 2H), 5.26 (p, J = 6.1 Hz, 1H), 2.60 - 2.51 (m, 1H), 2.49 - 2.40 (m, 1H), 2.33 (p, J = 1.8 Hz, 0H), 2.20 (s, 1H), 1.84 (dd, J = 11.8, 5.6 Hz, 1H), 1.51 (s, 3H), 1.45 (s, 3H), 1.43 - 1.30 (m, 4H), 1.29 (d, J = 6.2 Hz, 3H), 0.85 (d, J = 6.3 Hz, 3H).
2231	512.20	1.35	
2232	496.40	1.20	1H NMR (400 MHz, DMSO-d6) δ 12.40 (s, 1H), 7.76 (d, J = 8.0 Hz, 1H), 7.63 (dd, J = 8.4, 7.3 Hz, 1H), 7.25 (dd, J = 5.1, 2.9 Hz, 2H), 7.20 (d, J = 7.2 Hz, 1H), 7.08 (d, J = 8.1 Hz, 1H), 6.84 - 6.19 (m, 5H), 2.60 (d, J = 8.8 Hz, 2H), 2.29 (s, 3H), 2.26 - 2.14 (m, 1H), 1.87 (dd, J = 11.9, 5.6 Hz, 1H), 1.65 - 1.53 (m, 6H), 1.42 (t, J = 12.0 Hz, 1H), 0.88 (d, J = 6.2 Hz, 3H).
2233	527.22	2.22	
2234	551.50	1.82	1H NMR (400 MHz, DMSO-d6) δ 12.62 (s, 1H), 12.54 (s, 1H), 8.28 - 8.22 (m, 1H), 8.16 (d, J = 8.4 Hz, 2H), 7.84 (d, J = 6.9 Hz, 1H), 7.77 (d, J = 7.9 Hz, 1H), 7.49 (d, J = 8.4 Hz, 2H), 7.21 (d, J = 8.1 Hz, 1H), 6.45 (t, J = 6.7 Hz, 1H), 2.75 (q, J = 10.2 Hz, 2H), 2.24 (d, J = 12.0 Hz, 1H), 1.89 (dd, J = 12.2, 5.5 Hz, 1H), 1.61 (s, 6H), 1.45 (t, J = 11.9 Hz, 1H), 0.91 (d, J = 6.2 Hz, 3H).
2235	511.30	1.52	1H NMR (400 MHz, DMSO-d6) δ

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			12.59 (s, 1H), 12.53 (d, J = 6.2 Hz, 1H), 8.24 (dd, J = 7.2, 2.1 Hz, 1H), 7.83 (td, J = 6.2, 2.1 Hz, 1H), 7.74 (d, J = 8.0 Hz, 1H), 7.60 (dd, J = 7.3, 1.4 Hz, 2H), 7.38 (t, J = 8.0 Hz, 1H), 7.19 (d, J = 8.0 Hz, 1H), 7.01 - 6.96 (m, 1H), 6.45 (t, J = 6.8 Hz, 1H), 4.10 (q, J = 7.0 Hz, 2H), 2.74 (dd, J = 16.1, 9.7 Hz, 2H), 2.24 (s, 1H), 1.93 - 1.85 (m, 1H), 1.62 (d, J = 1.4 Hz, 6H), 1.45 (t, J = 12.1 Hz, 1H), 1.36 (t, J = 7.0 Hz, 3H), 0.91 (d, J = 6.3 Hz, 3H).
2236	502.50	1.66	
2237	574.60	1.59	
2238	542.50	1.44	1H NMR (400 MHz, DMSO-d6) δ 12.52 (s, 1H), 8.24 (dd, J = 4.8, 1.8 Hz, 1H), 8.03 (dd, J = 7.9, 1.8 Hz, 1H), 7.83 (d, J = 8.3 Hz, 1H), 6.95 (d, J = 8.3 Hz, 1H), 6.77 (dd, J = 7.8, 4.8 Hz, 1H), 6.69 (broad s, 2H), 5.83 (s, 1H), 3.90 (d, J = 6.6 Hz, 2H), 2.52 (s, 3H), 2.14 (br s, 1H), 2.02 (hept, J = 6.6 Hz, 1H), 1.85 (dd, J = 12.0, 5.6 Hz, 1H), 1.51 (s, 3H), 1.49 (s, 3H), 1.36 (t, J = 11.9 Hz, 1H), 1.24 (br s, 1H), 0.95 (d, J = 6.7 Hz, 6H), 0.88-0.82 (m, 1H), 0.74 (d, J = 6.3 Hz, 3H).
2239	497.51	2.12	1H NMR (400 MHz, DMSO-d6) δ 12.83 (s, 1H), 8.18 - 8.12 (m, 2H), 7.88 - 7.85 (d, 1H), 7.50 - 7.47 (d, 1H), 7.31 - 7.28 (d, 1H), 7.14 - 6.99 (m, 4H), 6.63 - 6.60 (dd, 1H), 3.78 (s, 3H), 2.30 - 2.21 (m, 1H), 1.89 - 1.82 (m, 1H), 1.51 (s, 3H), 1.47 (s, 3H), 1.46 - 1.35 (m, 1H), 0.89 - 0.87 (d, 3H)
2240	488.33	1.38	
2241	524.24	1.55	
2242	486.20	1.58	
2243	707.50	1.57	
2244	497.20	1.30	
2245	550.22	1.10	
2246	430.40	1.80	
2247	527.30	1.47	
2248	514.40	1.44	1H NMR (400 MHz, DMSO-d6) δ 12.40 (s, 1H), 8.23 (dd, J = 4.8, 1.8 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 8.02 (dd, J = 7.8, 1.9 Hz, 1H), 8.00 (d, J = 1.2 Hz, 1H), 7.80 (d, J = 8.3 Hz, 1H), 6.90 (d, J = 8.3 Hz, 1H), 6.76 (dd, J = 7.8, 4.7 Hz, 1H), 6.68 (broad s, 2H), 4.29 (q, J = 7.0 Hz, 2H), 2.45 (d, J = 8.7 Hz, 2H), 2.12 (br s, 1H), 1.94 (d, J = 0.9 Hz, 3H), 1.84 (dd, J = 12.0, 5.6 Hz, 1H), 1.55 (s, 3H), 1.54 (s, 3H), 1.40-1.33 (m, 4H), 0.74 (d, J = 6.3 Hz, 3H).
2249	525.42	2.42	
2250	472.29	2.32	
2251	518.22	2.48	
2252	487.30	2.84	
2253	545.23	2.34	
2254	537.43	2.28	
2255	492.40	2.73	
2256	525.30	1.57	
2257	482.24	0.90	
2258	570.00	3.40	1H NMR (400 MHz, DMSO-d6) δ 7.67 - 7.54 (m, 2H), 7.49 (t, J = 1.8 Hz, 1H), 7.40 (dt, J = 10.2, 1.8 Hz, 1H), 7.23 (d, J = 8.0 Hz, 1H), 7.06 (d, J = 7.3 Hz, 1H), 6.89 (dt, J = 10.7, 2.3 Hz, 1H), 6.70 (d, J = 8.4 Hz, 1H), 6.64 (s, 2H), 3.84 (d, J = 6.7 Hz, 2H), 3.43 (s, 3H), 2.82 (s, 1H), 2.64 (s, 1H), 2.34 - 2.22 (m, 1H), 2.05 (hept, J = 6.6 Hz, 1H), 1.91 (dd, J = 11.9, 5.7 Hz, 1H), 1.62 (d, J = 4.4 Hz, 6H), 1.46 (t, J = 12.1 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.94 (d, J = 6.3 Hz, 3H).
2259	502.24	1.76	
2260	536.22	2.32	
2261	549.43	2.69	
2262	528.12	2.54	
2263	549.43	2.30	
2264	515.21	2.70	
2265	440.40	1.17	
2266	516.20	1.31	1H NMR (400 MHz, DMSO) δ 12.48 (s, 1H), 7.70 - 7.52 (m, 2H), 7.17 (dd, J = 7.3, 0.5 Hz, 1H), 6.70 (dd, J = 9.7, 8.5 Hz, 2H), 5.19 (s, 2H), 3.84 (td, J = 11.1, 3.7 Hz, 2H), 3.38 - 3.18 (m, 2H), 2.84 (t, J = 10.0 Hz, 1H), 2.68 - 2.55

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			(m, 1H), 2.07 (s, 4H), 1.81 (t, J = 6.4 Hz, 1H), 1.65 (dd, J = 30.6, 10.5 Hz, 2H), 1.55 (d, J = 8.6 Hz, 6H), 1.49 (s, 3H), 1.28 (d, J = 6.6 Hz, 10H).
2267	539.60	1.90	1H NMR (400 MHz, DMSO-d6) δ 12.59 (s, 1H), 12.53 (d, J = 6.4 Hz, 1H), 8.24 (dd, J = 7.3, 2.1 Hz, 1H), 7.82 (dt, J = 6.5, 3.5 Hz, 1H), 7.74 (d, J = 8.0 Hz, 1H), 7.65 - 7.58 (m, 2H), 7.37 (td, J = 8.4, 3.1 Hz, 1H), 7.20 (d, J = 8.0 Hz, 1H), 7.02 - 6.96 (m, 1H), 6.45 (t, J = 6.8 Hz, 1H), 3.81 (d, J = 6.6 Hz, 2H), 2.79 - 2.69 (m, 2H), 2.24 (s, 1H), 2.05 (dt, J = 13.2, 6.6 Hz, 1H), 1.89 (dd, J = 11.8, 5.6 Hz, 1H), 1.62 (d, J = 1.9 Hz, 6H), 1.45 (t, J = 12.0 Hz, 1H), 0.99 (dd, J = 6.8, 2.9 Hz, 6H), 0.91 (d, J = 6.4 Hz, 3H).
2268	497.13	1.40	
2269	568.20	2.00	1H NMR (400 MHz, DMSO) δ 12.50 (s, 1H), 7.81 (d, J = 8.0 Hz, 1H), 7.63 (t, J = 7.8 Hz, 1H), 7.53 (s, 1H), 7.47 (d, J = 9.9 Hz, 1H), 7.33 (d, J = 8.0 Hz, 1H), 7.20 (d, J = 7.3 Hz, 1H), 6.92 (d, J = 10.7 Hz, 1H), 6.69 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 3.84 (d, J = 6.5 Hz, 2H), 3.61 (d, J = 9.7 Hz, 1H), 3.49 (q, J = 9.8 Hz, 1H), 2.71 - 2.51 (m, 2H), 2.38 - 2.19 (m, 1H), 2.03 (dd, J = 13.2, 6.7 Hz, 2H), 1.98 - 1.86 (m, 1H), 1.86 - 1.73 (m, 2H), 1.61 (dt, J = 22.9, 10.5 Hz, 2H), 0.98 (d, J = 6.7 Hz, 6H), 0.91 (d, J = 6.4 Hz, 3H).
2270	531.12	2.72	
2271	610.30	2.23	
2272	502.52	1.05	
2273	515.21	2.62	1H NMR (400 MHz, DMSO) δ 12.81 (s, 1H), 8.19 - 8.17 (dd, 1H), 8.06 (t, J = 7.91, 1H), 7.81 - 7.66 (m, 3H), 7.43 - 7.34 (m, 1H), 7.29 - 7.09 (m, 3H), 6.64 - 6.61 (dd, 1H), 5.44 (s, 2H), 2.37 - 2.14 (m, 2H), 2.00 - 1 .94 (m, 1H), 1.80 - 1.73 (m, 1H), 1.49 (s, 3H), 1.45 (s, 3H), 1.30 (t, J = 11.86, 1H), 0.67 - 0.65 (d, 3H)
2274	517.32	1.37	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2275	466.29	1.87	
2276	527.40	1.62	
2277	444.60	1.40	
2278	529.20	1.71	1H NMR (400 MHz, DMSO) δ 12.62 (s, 1H), 12.53 (d, J = 5.8 Hz, 1H), 8.24 (dd, J = 7.3, 2.2 Hz, 1H), 7.83 (dd, J = 6.4, 4.3 Hz, 1H), 7.75 (d, J = 8.0 Hz, 1H), 7.47 (d, J = 1.9 Hz, 1H), 7.41 (d, J = 10.2 Hz, 1H), 7.24 (d, J = 8.0 Hz, 1H), 6.88 (dt, J = 10.8, 2.3 Hz, 1H), 6.45 (t, J = 6.8 Hz, 1H), 4.12 (q, J = 7.0 Hz, 2H), 2.74 (dd, J = 17.6, 10.2 Hz, 2H), 2.25 (s, 1H), 1.89 (dd, J = 11.9, 5.7 Hz, 1H), 1.60 (d, J = 1.9 Hz, 6H), 1.45 (t, J = 12.1 Hz, 1H), 1.35 (t, J = 7.0 Hz, 3H), 0.91 (d, J = 6.3 Hz, 3H).
2279	566.30	1.82	1H NMR (400 MHz, DMSO) δ 8.70 (s, 1H), 7.99 (s, 1H), 7.56 (d, J = 7.7 Hz, 1H), 7.44 (t, J = 7.8 Hz, 1H), 6.97 (dd, J = 17.9, 7.5 Hz, 2H), 6.45 (d, J = 8.2 Hz, 1H), 6.12 (s, 2H), 3.11 - 2.91 (m, 4H), 2.83 (s, 3H), 2.30 (s, 3H), 2.13 (dq, J = 17.3, 9.3, 8.4 Hz, 1H), 1.90 (t, J = 6.8 Hz, 1H), 1.81 (dd, J = 11.6, 5.5 Hz, 1H), 1.62 (s, 3H), 1.57 (s, 3H), 1.37 (t, J = 11.9 Hz, 1H), 0.88 - 0.81 (m, 10H).
2280	516.23	1.54	
2281	494.30	1.08	1H NMR (400 MHz, DMSO-d6) δ 12.29 (s, 1H), 7.66 - 7.57 (m, 2H), 7.17 (d, J = 7.3 Hz, 1H), 6.66 (d, J = 8.3 Hz, 1H), 6.41 (s, 2H), 5.55 - 5.48 (m, 1H), 3.94 - 3.83 (m, 2H), 3.83 - 3.76 (m, 2H), 2.56 (d, J = 10.6 Hz, 1H), 2.44 (t, J = 8.7 Hz, 1H), 2.32 - 2.16 (m, 1H), 2.10 - 2.02 (m, 1H), 1.84 (dd, J = 11.8, 5.6 Hz, 1H), 1.53 (s, 3H), 1.44 (s, 3H), 1.37 (t, J = 12.0 Hz, 1H), 0.85 (d, J = 6.3 Hz, 3H).
2282	502.31	2.63	1H NMR (400 MHz, DMSO-d6) δ 12.93 (s, 1H), 10.03 (s, 1H), 8.16 - 8.14 (dd, 1H), 7.93 (t, J = 7.80, 1H), 7.74 - 7.71 (dd, 1H), 7.61 - 7.58 (d, 1H), 7.54 - 7.49 (m, 2H), 7.13 - 7.10 (d, 1H), 6.80 - 6.72 (m, 1H), 6.56 -

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			6.53 (dd, 1H), 2.04 - 1.93 (m, 1H), 1.81 - 1.74 (m, 1H), 1.45 (s, 6H), 1.32 (t, J = 11.87, 1H), 0.73 - 0.71 (d, 3H)
2283	524.22	1.65	
2284	503.30	1.56	1H NMR (400 MHz, DMSO-d6) δ 12.62 (s, 1H), 12.53 (d, J = 6.1 Hz, 1H), 8.24 (dd, J = 7.3, 2.2 Hz, 1H), 8.04 (ddd, J = 12.0, 8.3, 2.2 Hz, 1H), 7.92 (dt, J = 8.8, 3.1 Hz, 1H), 7.83 (dt, J = 7.0, 3.5 Hz, 1H), 7.76 (d, J = 8.0 Hz, 1H), 7.57 (dt, J = 10.5, 8.6 Hz, 1H), 7.23 (d, J = 7.9 Hz, 1H), 6.45 (t, J = 6.8 Hz, 1H), 2.73 (dd, J = 9.6, 7.8 Hz, 2H), 2.30 - 2.16 (m, 1H), 1.89 (dd, J = 12.0, 5.5 Hz, 1H), 1.60 (d, J = 2.0 Hz, 6H), 1.45 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).
2285	550.33	1.49	
2286	537.43	3.02	
2287	494.50	1.63	
2288	509.22	1.23	
2289	570.30	2.21	1H NMR (400 MHz, DMSO) δ 12.52 (s, 1H), 7.78 (d, J = 7.9 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.49 (d, J = 1.9 Hz, 1H), 7.43 (dd, J = 9.7, 1.7 Hz, 1H), 7.30 (d, J = 7.9 Hz, 1H), 7.20 (dd, J = 7.3, 0.6 Hz, 1H), 6.88 (dt, J = 10.7, 2.3 Hz, 1H), 6.69 (dd, J = 8.4, 0.6 Hz, 1H), 6.46 (s, 2H), 3.83 (d, J = 6.5 Hz, 2H), 3.47 (d, J = 7.2 Hz, 2H), 2.68 (d, J = 7.2 Hz, 3H), 2.05 (dt, J = 13.3, 4.3 Hz, 1H), 1.76 (s, 1H), 1.61 (d, J = 11.7 Hz, 5H), 1.12 (t, J = 9.5 Hz, 3H), 0.99 (d, J = 6.7 Hz, 6H), 0.95 - 0.80 (m, 2H).
2290	498.23	1.27	
2291	502.26	1.39	
2292	534.12	2.70	
2293	527.22	2.36	
2294	496.30	1.44	
2295	501.30	1.46	1H NMR (400 MHz, DMSO-d6) δ 13.20 (s, 1H), 11.69 (s, 1H), 8.00 - 7.88 (m, 1H), 7.78 - 7.61 (m, 2H), 7.55 - 7.40 (m, 1H), 7.25 (d, J = 8.0 Hz, 1H), 7.09 - 7.00 (m, 1H), 6.85 (dd, J = 8.9, 5.1 Hz, 1H), 6.27 - 6.13 (m, 1H), 3.23 -

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			3.07 (m, 2H), 2.22 - 2.11 (m, 1H), 1.87 (dd, J = 11.8, 5.7 Hz, 1H), 1.50 (d, J = 3.4 Hz, 6H), 1.41 (t, J = 12.1 Hz, 1H), 0.88 (d, J = 6.1 Hz, 3H).
2296	534.20	1.70	
2297	548.23	2.70	
2298	501.11	2.43	
2299	506.21	2.30	
2300	497.30	1.62	
2301	522.30	2.22	
2302	532.22	1.41	
2303	512.23	1.56	
2304	491.23	1.61	
2305	474.39	2.52	
2306	539.40	2.16	1H NMR (400 MHz, DMSO-d6) δ 7.63 (s, 1H), 7.49 (s, 1H), 7.36 (s, 1H), 7.13 (d, J = 6.9 Hz, 1H), 7.06 (s, 2H), 6.52 (s, 1H), 6.23 (s, 2H), 5.26 (p, J = 6.2 Hz, 1H), 2.91 (s, 1H), 2.42 (s, 3H), 2.18 (t, J = 7.4 Hz, 1H), 1.83 (dd, J = 11.8, 5.6 Hz, 1H), 1.61 (s, 3H), 1.56 (s, 3H), 1.47 (q, J = 6.6, 6.1 Hz, 1H), 1.38 (t, J = 11.9 Hz, 1H), 1.30 (d, J = 6.2 Hz, 6H), 0.83 (d, J = 5.9 Hz, 3H).
2307	460.29	2.51	
2308	471.39	2.51	
2309	527.22	2.12	
2310	510.41	2.25	
2311	474.40	1.00	
2312	500.30	1.62	1H NMR (400 MHz, DMSO-d6) δ 12.45 (s, 1H), 8.30 (d, J = 2.2 Hz, 1H), 7.72 (d, J = 2.3 Hz, 1H), 7.68 - 7.53 (m, 2H), 7.51 - 7.43 (m, 1H), 7.46 - 7.34 (m, 2H), 7.21 (d, J = 7.3 Hz, 1H), 6.68 (d, J = 8.4 Hz, 1H), 6.48 (s, 2H), 2.22 (s, 1H), 1.86 (dd, J = 11.9, 5.6 Hz, 1H), 1.55 (d, J = 10.4 Hz, 6H), 1.41 (t, J = 12.0 Hz, 1H), 0.87 (d, J = 6.3 Hz, 3H).
2313	564.42	1.95	
2314	482.20	1.34	
2315	541.30	1.54	1H NMR (400 MHz, DMSO-d6) δ 12.64 - 12.32 (m, 2H), 8.22 (dd, J = 7.2, 2.2 Hz, 1H), 7.85 - 7.77 (m, 1H), 7.60 (d, J = 7.7 Hz, 1H), 6.53 (d, J = 7.8

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			Hz, 1H), 6.43 (t, J = 6.8 Hz, 1H), 2.81 - 2.72 (m, 1H), 2.67 (d, J = 8.7 Hz, 2H), 2.48 - 2.39 (m, obscured by solvent peak),2.21 (s, 1H), 2.07 (s, 1H), 1.84 (dd, J = 11.8, 5.6 Hz, 1H), 1.78 - 1.60 (m, 6H), 1.51 (d, J = 10.8 Hz, 6H), 1.39 (t, J = 12.1 Hz, 1H), 0.89 (d, J = 6.2 Hz, 3H).
2316	509.21	2.17	
2317	457.28	2.40	
2318	460.29	2.31	
2319	510.19	1.02	
2320	481.30	2.30	
2321	550.33	2.40	
2322	536.50	2.01	
2323	544.27	1.17	
2324	474.09	2.01	
2325	610.70	1.93	
2326	588.30	1.32	1H NMR (400 MHz, DMSO-d6) δ 12.50 (s, 1H), 8.29 - 8.18 (m, 2H), 8.07 (dd, J = 7.8, 1.8 Hz, 1H), 7.86 (d, J = 8.3 Hz, 1H), 6.94 (d, J = 8.3 Hz, 1H), 6.90 - 6.63 (m, 0H), 6.79 (dd, J = 7.8, 4.8 Hz, 1H), 6.14 (d, J = 2.8 Hz, 1H), 4.36 - 4.29 (m, 2H), 3.79 - 3.71 (m, 2H), 3.62 - 3.54 (m, 2H), 3.52 - 3.47 (m, 2H), 3.43 (q, J = 7.0 Hz, 2H), 2.46 (d, J = 8.8 Hz, 2H), 2.13 (m, J = 5.8 Hz, 1H), 1.85 (dd, J = 11.9, 5.6 Hz, 1H), 1.55 (s, 3H), 1.53 (s, 3H), 1.44 - 1.33 (m, 1H), 1.09 (t, J = 7.0 Hz, 3H), 0.74 (d, J = 6.3 Hz, 3H).
2327	488.29	1.36	
2328	508.31	1.79	
2329	503.21	2.21	
2330	472.29	2.57	
2331	580.00	3.15	
2332	485.20	2.20	
2333	518.22	1.47	
2334	532.16	1.44	
2335	561.14	2.31	
2336	487.30	2.89	
2337	534.25	1.83	
2338	506.21	2.08	
2339	472.30	1.46	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2340	484.30	2.52	
2341	502.31	1.84	
2342	445.28	2.29	
2343	490.30	2.55	
2344	625.60	1.71	1H NMR (400 MHz, DMSO-d6) δ 12.29 (s, 1H), 7.65 - 7.59 (m, 1H), 7.33 - 7.24 (m, 1H), 7.22 - 7.14 (m, 1H), 6.67 (d, J = 8.4 Hz, 1H), 5.53 (s, 1H), 4.19 - 4.02 (m, 2H), 2.54 (s, 1H), 2.31 (s, 2H), 2.28 (d, J = 5.3 Hz, 1H), 2.19 (d, J = 6.7 Hz, 1H), 1.83 (dd, J = 11.9, 5.7 Hz, 1H), 1.54 (s, 3H), 1.52 (s, 3H), 1.40 - 1.32 (m, 2H), 1.31 (s, 9H), 1.18 (s, 2H), 1.02 (s, 9H), 0.98 (s, 1H), 0.85 (d, J = 6.2 Hz, 3H). (amide rotamers)
2345	498.53	1.09	
2346	474.09	2.70	
2347	561.60	2.32	
2348	499.40	1.59	
2349	500.17	1.29	
2350	453.98	1.90	
2351	474.30	1.15	1H NMR (400 MHz, DMSO-d6) δ 12.45 (s, 1H), 8.24 (dd, J = 5.0, 1.9 Hz, 1H), 8.04 (d, J = 7.7 Hz, 1H), 7.67 (d, J = 7.7 Hz, 1H), 6.77 (dd, J = 7.8, 4.8 Hz, 2H), 6.68 (d, J = 7.8 Hz, 2H), 4.21 (dd, J = 38.7, 11.0 Hz, 1H), 4.02 (d, J = 11.4 Hz, 1H), 3.63 - 3.44 (m, 1H), 2.47 - 2.36 (m, 2H), 2.18 - 2.04 (m, 1H), 2.04 - 1.91 (m, 1H), 1.91 - 1.73 (m, 2H), 1.72 - 1.58 (m, 1H), 1.53 (s, 2H), 1.51 - 1.42 (m, 6H), 1.43 - 1.24 (m, 2H), 0.73 (d, J = 6.1 Hz, 3H).
2352	519.04	1.49	
2353	524.52	2.49	
2354	462.39	1.84	
2355	514.17	1.43	
2356	515.22	1.77	
2357	526.32	2.22	
2358	554.70	1.81	1H NMR (400 MHz, DMSO-d6) δ 12.51 (s, 1H), 8.24 (d, J = 2.8 Hz, 1H), 8.22 (broad s, 1H), 8.00 (d, J = 7.7 Hz, 1H), 7.85 (d, J = 8.2 Hz, 1H), 7.00 (d, J = 8.2 Hz, 1H), 6.78 - 6.70 (m, 1H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			6.65 (br s, 2H), 6.16 (d, J = 2.7 Hz, 1H), 3.90 (s, 2H), 3.50 (br d, J = 9.9 Hz, 1H), 3.39 (br d, J = 8.8 Hz, 1H), 2.48-2.38 (br m, 2H), 2.26 (dd, J = 11.8, 5.2 Hz, 1H), 1.92-1.90 (br m, 2H), 1.80-1.75 (br m, 2H), 1.61 (br s, 1H), 1.51 (t, J = 11.5 Hz, 1H), 1.01 (s, 9H), 0.78 (d, J = 6.4 Hz, 3H).
2359	495.40	1.34	
2360	515.40	1.63	1H NMR (400 MHz, DMSO-d6) δ 12.52 (s, 2H), 8.23 (d, J = 7.1 Hz, 1H), 8.18 (d, J = 2.8 Hz, 1H), 7.87 - 7.71 (m, 2H), 6.86 (d, J = 8.2 Hz, 1H), 6.44 (broad s, 1H), 6.07 (d, J = 2.7 Hz, 1H), 4.85 (sep, J = 6.2 Hz, 1H), 2.71 (br s, 2H), 2.22 (br s, 1H), 1.88 (dd, J = 12.0, 5.6 Hz, 1H), 1.56 (s, 3H), 1.55 (s, 3H), 1.43 (t, J = 12.0 Hz, 1H), 1.33 (d, J = 2.9 Hz, 3H), 1.32 (d, J = 2.9 Hz, 3H), 0.89 (d, J = 6.3 Hz, 3H).
2361	484.30	1.74	1H NMR (400 MHz, DMSO-d6) δ 7.96 (s, 1H), 7.51 (s, 1H), 7.39 (t, J = 7.8 Hz, 1H), 7.00 (d, J = 7.2 Hz, 1H), 6.41 (d, J = 8.2 Hz, 1H), 5.98 (s, 3H), 5.88 (s, 1H), 3.05 - 2.96 (m, 2H), 2.24 (d, J = 19.4 Hz, 2H), 2.11 (s, 3H), 1.64 (s, 2H), 1.48 (d, J = 5.0 Hz, 7H), 0.97 (d, J = 6.4 Hz, 4H), 0.86 (d, J = 6.3 Hz, 4H).
2362	552.60	1.24	
2363	481.23	1.10	
2364	525.30	2.04	1H NMR (400 MHz, DMSO-d6) δ 12.49 (s, 1H), 7.79 (d, J = 8.0 Hz, 1H), 7.62 (t, J = 7.8 Hz, 1H), 7.41 (s, 1H), 7.26 (d, J = 7.9 Hz, 1H), 7.20 (d, J = 7.4 Hz, 1H), 7.14 (s, 1H), 6.67 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 4.32 (q, J = 7.0 Hz, 2H), 2.71 - 2.58 (m, 2H), 2.45 (s, 3H), 2.29 - 2.13 (m, 1H), 1.87 (dd, J = 11.9, 5.7 Hz, 1H), 1.60 (s, 3H), 1.59 (s, 2H), 1.42 (t, J = 12.1 Hz, 1H), 1.33 (t, J = 7.0 Hz, 3H), 0.88 (d, J = 6.3 Hz, 3H).
2365	549.43	2.57	
2366	550.30	1.72	
2367	500.20	1.04	
2368	508.32	1.13	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2369	511.15	1.63	
2370	562.40	1.58	
2371	445.28	2.37	
2372	502.50	1.59	
2373	492.30	1.51	1H NMR (400 MHz, DMSO-d6) δ 12.26 (s, 1H), 7.65 - 7.55 (m, 2H), 7.17 (dd, J = 7.3, 0.7 Hz, 1H), 6.66 (dd, J = 8.5, 0.8 Hz, 1H), 6.45 (s, 2H), 4.88 - 4.57 (m, 1H), 2.63 - 2.51 (m, 1H), 2.43 (t, J = 8.8 Hz, 1H), 2.17 (s, 1H), 1.82 (dd, J = 11.9, 5.6 Hz, 1H), 1.49 - 1.26 (m, 10H), 1.19 (dtd, J = 13.5, 9.0, 8.5, 5.1 Hz, 1H), 0.85 (d, J = 6.3 Hz, 3H), 0.62 - 0.44 (m, 2H), 0.32 - 0.21 (m, 2H).
2374	569.30	2.39	1H NMR (400 MHz, MeOD) δ 7.87 (d, J = 8.1 Hz, 1H), 7.76 (d, J = 8.1 Hz, 1H), 7.70 (t, J = 7.9 Hz, 1H), 7.54 (s, 1H), 7.42 (t, J = 7.8 Hz, 2H), 6.83 (d, J = 8.4 Hz, 1H), 6.75 (d, J = 10.5 Hz, 1H), 3.82 (d, J = 6.5 Hz, 2H), 3.67 (dd, J = 12.3, 6.3 Hz, 1H), 2.66 (t, J = 12.5 Hz, 1H), 2.11 (dt, J = 13.2, 6.6 Hz, 1H), 1.69 (dd, J = 12.5, 6.3 Hz, 1H), 1.57 - 1.40 (m, 2H), 1.18 (s, 3H), 1.05 (d, J = 6.9 Hz, 9H), 0.81 (s, 3H), 0.75 (s, 3H).
2375	556.30	2.43	1H NMR (400 MHz, DMSO) δ 12.84 - 11.49 (m, 1H), 8.30 (d, J = 8.2 Hz, 1H), 8.26 - 8.05 (m, 0H), 7.66 (t, J = 7.9 Hz, 1H), 7.56 (s, 1H), 7.50 (d, J = 9.8 Hz, 1H), 7.27 (d, J = 8.2 Hz, 1H), 7.19 (d, J = 7.2 Hz, 1H), 6.91 (d, J = 10.7 Hz, 1H), 6.71 (d, J = 8.4 Hz, 1H), 6.59 (s, 2H), 3.84 (d, J = 6.5 Hz, 2H), 3.35 (d, J = 6.4 Hz, 2H), 2.10 - 1.97 (m, 1H), 1.64 (d, J = 21.5 Hz, 6H), 1.24 - 1.04 (m, 3H), 1.00 (d, J = 6.7 Hz, 6H), 0.94 (d, J = 10.5 Hz, 2H).
2376	540.20	2.11	1H NMR (400 MHz, DMSO) δ 12.50 (s, 1H), 7.76 (d, J = 7.8 Hz, 1H), 7.63 (dd, J = 8.3, 7.4 Hz, 1H), 7.46 - 7.38 (m, 2H), 7.34 (d, J = 7.9 Hz, 1H), 7.26 - 7.20 (m, 1H), 6.88 (dt, J = 10.8, 2.3 Hz, 1H), 6.71 - 6.67 (m, 1H), 6.45 (s, 2H), 4.21 (d, J = 12.9 Hz, 1H), 3.83 (d,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			J = 6.6 Hz, 2H), 2.60 (t, J = 11.5 Hz, 1H), 2.47 - 2.39 (m, 1H), 2.10 - 1.95 (m, 1H), 1.86 (dd, J = 12.8, 6.3 Hz, 1H), 1.73 (d, J = 12.8 Hz, 1H), 1.58 (d, J = 10.8 Hz, 1H), 1.17 (dd, J = 15.0, 8.4 Hz, 2H), 0.99 (d, J = 6.7 Hz, 6H), 0.69 (dt, J = 9.3, 5.7 Hz, 1H), -0.11 (t, J = 7.8 Hz, 1H).
2377	541.20	1.63	1H NMR (400 MHz, Methanol-d4) δ 9.19 (s, 1H), 8.80 (d, J = 5.0 Hz, 1H), 8.48 (d, J = 8.1 Hz, 1H), 8.00 (d, J = 8.0 Hz, 1H), 7.66 (dd, J = 8.1, 4.9 Hz, 1H), 7.47 (s, 1H), 7.43 (d, J = 8.1 Hz, 1H), 7.35 (d, J = 10.2 Hz, 1H), 6.75 (d, J = 10.9 Hz, 1H), 3.81 (d, J = 6.5 Hz, 2H), 2.98 (t, J = 9.1 Hz, 1H), 2.86 (t, J = 10.0 Hz, 1H), 2.42 (s, 1H), 2.05 (ddd, J = 19.8, 13.3, 7.1 Hz, 3H), 1.58 (d, J = 17.4 Hz, 6H), 1.03 (dd, J = 7.5, 3.4 Hz, 6H), 0.94 (d, J = 6.4 Hz, 3H).
2378	528.12	2.64	
2379	510.17	1.51	
2380	510.15	1.46	
2381	534.12	2.50	
2382	506.21	2.34	
2383	529.32	1.55	1H NMR (400 MHz, DMSO) δ 12.56 (s, 1H), 10.51 (s, 1H), 8.18 - 8.16 (dd, 1H), 7.79 - 7.69 (m, 2H), 7.26 - 7.23 (d, 1H), 7.13 - 7.10 (d, 1H), 6.63 - 6.60 (dd, 1H), 4.44 - 4.35 (m, 1H), 3.05 (m, 2H), 2.83 - 2.80 (m, 2 H), 2.72 (s, 3H), 2.70 (s, 3H), 2.23 - 2.12 (m, 1H), 1.83 - 1.73 (m, 3H), 1.60 (m, 3H), 1.52 (s, 3H), 1.49 (s, 3H), 1.37 (t, J = 11.98, 1H), 1.09 (m, 2H), 0.79 - 0.77 (d, 3H)
2384	511.25	1.47	
2385	526.02	2.69	
2386	500.60	1.53	
2387	523.32	1.90	
2388	532.21	1.49	
2389	506.21	1.37	1H NMR (400 MHz, DMSO) δ 9.03 - 9.00 (d, 1H), 8.68 (s, 1H), 8.40 - 8.30 (m, 2H), 8.18 - 8.16 (dd, 1H), 8.08 - 8.05 (dd, 1H), 8.01 (s, 1H), 7.87 - 7.84 (dd, 1H), 7.63 - 7.60 (dd, 1H), 6.66 -

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			6.63 (dd, 1H), 2.28 - 2. 24 (d, 2H), 1.65 - 1.47 (m, 2H), 1.47 (s, 3H), 1.40 (s, 3H), 1.17 (t, J = 11.09, 1H), 0.15 - 0.13 (d, 3H)
2390	549.13	2.70	
2391	494.20	2.42	
2392	534.30	1.67	
2393	462.39	1.80	
2394	498.50	1.58	
2395	522.30	1.86	
2396	475.29	1.93	
2397	534.12	2.50	
2398	549.50	2.03	
2399	500.19	1.68	
2400	547.33	2.26	
2401	519.27	1.44	
2402	549.13	2.81	
2403	528.19	1.46	
2404	569.50	1.59	
2405	441.40	1.13	1H NMR (400 MHz, DMSO-d6) δ 12.90 (s, 1H), 12.57 (s, 1H), 8.27 (dd, J = 7.3, 2.2 Hz, 1H), 8.17 (s, 1H), 7.85 (s, 1H), 7.73 (d, J = 8.0 Hz, 1H), 7.62 - 7.48 (m, 2H), 7.20 (t, J = 7.4 Hz, 1H), 6.46 (t, J = 6.8 Hz, 1H), 3.01 (t, J = 8.5 Hz, 1H), 2.76 (t, J = 10.1 Hz, 1H), 2.30 - 2.17 (m, 1H), 1.88 (dd, J = 11.8, 5.7 Hz, 1H), 1.63 (s, 6H), 1.44 (t, J = 12.1 Hz, 1H), 0.92 (d, J = 6.3 Hz, 3H).
2406	542.20	1.66	1H NMR (400 MHz, DMSO-d6) δ 8.25 (dd, J = 4.8, 1.8 Hz, 1H), 8.05 (dd, J = 7.8, 1.8 Hz, 1H), 7.92 (d, J = 1.0 Hz, 1H), 7.84 (d, J = 8.2 Hz, 1H), 7.71 (s, 1H), 7.07 (d, J = 8.3 Hz, 1H), 6.78 (dd, J = 7.8, 4.8 Hz, 2H), 2.46 (s, 1H), 2.13 (s, 1H), 1.86 (dd, J = 12.0, 5.7 Hz, 1H), 1.56 (d, J = 4.4 Hz, 6H), 1.39 (t, J = 12.2 Hz, 1H), 0.98 (s, 9H), 0.75 (d, J = 6.2 Hz, 3H).
2407	549.43	2.09	
2408	513.41	2.62	
2409	499.31	2.49	
2410	525.30	1.62	
2411	481.50	1.83	
2412	461.19	2.69	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2413	510.17	1.46	
2414	476.29	1.42	1H NMR (400 MHz, DMSO) δ 12.09 (s, 1H), 8.14 (s, 3H), 7.92 (d, J = 8.0 Hz, 1H), 7.84 (dd, J = 10.4, 2.0 Hz, 1H), 7.75 (d, J = 7.9 Hz, 1H), 7.56 (td, J = 8.0, 6.2 Hz, 1H), 7.33 - 7.25 (m, 2H), 3.78 (ddd, J = 16.8, 10.7, 6.2 Hz, 2H), 3.71 - 3.62 (m, 1H), 3.62 - 3.53 (m, 1H), 3.41 (dd, J = 10.0, 4.7 Hz, 1H), 3.10 (t, J = 10.2 Hz, 1H), 2.99 - 2.91 (m, 1H), 2.36 (dd, J = 13.6, 9.2 Hz, 1H), 2.28 - 2.16 (m, 1H), 1.97 (dd, J = 12.2, 5.8 Hz, 2H), 1.68 (s, 3H), 1.64 (s, 3H), 1.56 (dd, J = 15.4, 8.1 Hz, 1H), 1.05 (d, J = 6.3 Hz, 3H).
2415	523.32	2.92	
2416	538.17	1.49	
2417	645.50	1.63	1H NMR (400 MHz, DMSO-d6) δ 12.28 (s, 1H), 7.65 - 7.57 (m, 1H), 7.25 (ddd, J = 34.6, 18.8, 7.3 Hz, 7H), 6.67 (d, J = 8.3 Hz, 1H), 5.50 (d, J = 19.3 Hz, 1H), 4.26 - 3.97 (m, 2H), 3.81 (s, 1H), 3.77 (d, J = 7.4 Hz, 1H), 3.73 - 3.51 (m, 3H), 2.30 - 2.04 (m, 2H), 1.82 (dd, J = 12.0, 5.7 Hz, 1H), 1.53 (s, 3H), 1.51 (s, 3H), 1.37 (d, J = 10.9 Hz, 2H), 1.27 (s, 9H), 1.17 (s, 1H), 0.84 (d, J = 6.3 Hz, 3H). (rotamers)
2418	517.14	1.14	
2419	471.30	1.19	
2420	497.51	2.15	
2421	516.41	2.36	
2422	515.20	1.60	
2423	543.43	1.87	
2424	481.40	1.50	
2425	516.20	1.31	1H NMR (400 MHz, DMSO) δ 12.47 (s, 1H), 7.70 - 7.52 (m, 2H), 7.17 (dd, J = 7.3, 0.5 Hz, 1H), 6.78 - 6.59 (m, 2H), 5.14 (s, 2H), 3.84 (d, J = 3.5 Hz, 2H), 3.30 (d, J = 10.7 Hz, 2H), 2.84 (t, J = 10.0 Hz, 1H), 2.68 - 2.55 (m, 1H), 2.07 (s, 4H), 1.87 - 1.73 (m, 1H), 1.65 (dd, J = 30.6, 10.5 Hz, 2H), 1.55 (d, J = 8.6 Hz, 6H), 1.49 (s, 3H), 1.28 (d, J = 6.6 Hz, 10H).

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2426	496.60	1.33	
2427	549.60	2.10	
2428	511.31	2.44	1H NMR (400 MHz, DMSO) δ 12.79 (s, 1H), 8.19 - 8.17 (dd, 1H), 8.01 (t, J = 7.80, 1H), 7.80 - 7.74 (m, 2H), 7.50 - 7.47 (d, 2H), 7.14 - 7.11 (d, 1H), 6.82 - 6.79 (d, 2H), 6.64 - 6.61 (dd, 1H), 5.35 - 5.22 (m, 2H), 3 .60 (s, 3H), 2.18 - 1.87 (m, 3H), 1.77 - 1.70 (m, 1H), 1.49 (s, 3H), 1.44 (s, 3H), 1.31- 1.21 (m, 1H), 0.60 - 0.58 (d, 3H)
2429	500.20	1.70	
2430	405.36	1.83	
2431	514.21	1.67	
2432	540.40	1.36	1H NMR (400 MHz, Methanol-d4) δ 8.55 (dd, J = 7.6, 1.7 Hz, 1H), 8.38 (dd, J = 7.2, 2.1 Hz, 1H), 8.11 (dd, J = 6.0, 1.7 Hz, 1H), 7.45 (d, J = 8.1 Hz, 1H), 7.08 - 6.99 (m, 1H), 3.91 (dd, J = 11.5, 8.5 Hz, 1H), 3.47 - 3.34 (m, 1H), 3.14 - 3.05 (m, 1H), 2.91 - 2.75 (m, 1H), 2.48 - 2.32 (m, 0H), 2.33 - 2.17 (m, 3H), 1.98 - 1.76 (m, 7H), 1.55 (s, 3H), 1.33 (s, 3H), 1.24 (d, J = 6.6 Hz, 3H).
2433	524.52	1.75	
2434	459.39	2.51	
2435	484.24	0.94	
2436	526.40	1.22	1H NMR (400 MHz, DMSO-d6) δ 12.53 (s, 1H), 8.25 (dd, J = 4.8, 1.9 Hz, 1H), 8.05 (dd, J = 7.7, 1.9 Hz, 1H), 7.74 (d, J = 7.8 Hz, 1H), 7.24 (dd, J = 7.7, 1.8 Hz, 1H), 7.16 (t, J = 7.9 Hz, 1H), 7.13 - 7.06 (m, 2H), 6.78 (dd, J = 7.8, 4.8 Hz, 3H), 3.85 (s, 3H), 2.53 (s, 2H), 2.15 (s, 1H), 1.82 (dd, J = 11.9, 5.6 Hz, 1H), 1.53 (s, 3H), 1.50 (s, 3H), 1.38 (t, J = 12.1 Hz, 1H), 0.76 (d, J = 6.3 Hz, 3H).
2437	491.40	1.17	1H NMR (400 MHz, DMSO-d6) δ 12.58 (s, 1H), 8.26 (dd, J = 4.8, 1.9 Hz, 1H), 8.24 - 8.15 (m, 2H), 8.05 (dd, J = 7.9, 1.8 Hz, 1H), 8.01 - 7.93 (m, 2H), 7.83 (d, J = 8.0 Hz, 1H), 7.34 (d, J = 8.0 Hz, 1H), 6.90 - 6.52 (m, 3H), 2.60 -

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			2.40 (m, 24H), 2.22 - 2.05 (m, 1H), 1.86 (dd, J = 11.9, 5.6 Hz, 1H), 1.60 (s, 3H), 1.56 (s, 3H), 1.40 (t, J = 12.1 Hz, 1H), 0.76 (d, J = 6.3 Hz, 3H).
2438	481.30	1.50	
2439	587.25	2.39	
2440	511.16	1.69	1H NMR (400 MHz, DMSO-d6) δ 12.73 - 12.37 (m, 1H), 8.25 (dd, J = 7.2, 2.2 Hz, 1H), 7.85 - 7.78 (m, 1H), 7.74 (d, J = 7.9 Hz, 1H), 7.67 (d, J = 1.6 Hz, 1H), 7.57 - 7.51 (m, 1H), 7.24 (d, J = 7.8 Hz, 1H), 7.19 (d, J = 8.0 Hz, 1H), 6.46 (t, J = 6.8 Hz, 1H), 3.85 (s, 3H), 2.79 - 2.66 (m, 2H), 2.29 - 2.16 (m, 4H), 1.89 (dd, J = 11.7, 5.5 Hz, 1H), 1.63 (d, J = 2.7 Hz, 6H), 1.49 - 1.41 (m, 1H), 0.91 (d, J = 6.3 Hz, 3H).
2441	533.22	1.45	
2442	516.41	2.47	
2443	510.41	1.84	
2444	516.19	1.61	
2445	467.40	1.13	1H NMR (400 MHz, DMSO-d6) δ 12.59 (s, 1H), 12.53 (d, J = 6.3 Hz, 1H), 8.24 (dd, J = 7.2, 2.2 Hz, 1H), 8.08 - 8.02 (m, 2H), 7.83 (dt, J = 6.6, 3.3 Hz, 1H), 7.76 (d, J = 8.0 Hz, 1H), 7.54 - 7.46 (m, 2H), 7.46 - 7.40 (m, 1H), 7.19 (d, J = 8.0 Hz, 1H), 6.45 (t, J = 6.8 Hz, 1H), 2.82 - 2.63 (m, 2H), 2.30 - 2.18 (m , 1H), 1.89 (dd, J = 11.8, 5.5 Hz, 1H), 1.62 (s, 6H), 1.45 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).
2446	500.50	1.27	
2447	502.48	1.03	
2448	538.38	1.61	
2449	517.32	1.62	1H NMR (400 MHz, DMSO-d6) δ 12.92 (s, 1H), 10.37 (s, 1H), 9.22 - 9.21 (d, 1H), 9.08 - 9.07 (d, 1H), 8.13 - 8.11 (dd, 1H), 8.02 - 7.93 (m, 2H), 7.87 - 7.84 (d, 1H), 7.81 - 7.78 (dd, 1H), 7.63 - 7.44 (m, 3H), 7.28 - 7.2 5 (d, 1H), 6.49 - 6.46 (dd, 1H), 2.46 - 2.42 (m, 2H), 1.63 - 1.57 (m, 1H), 1.50 - 1.43 (m, 1H), 1.38 (s, 3H), 1.23 -

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1.13 (m, 1H), 1.12 (s, 3H), 0.61 - 0.59 (d, 3H)
2450	524.26	1.52	
2451	556.00	1.85	
2452	502.31	2.82	
2453	476.19	2.13	
2454	498.15	1.43	
2455	532.50	2.52	
2456	499.30	1.56	1H NMR (400 MHz, DMSO-d6) δ 12.58 (s, 1H), 12.53 (d, J = 6.2 Hz, 1H), 8.25 (dd, J = 7.2, 2.2 Hz, 1H), 8.02 - 7.94 (m, 1H), 7.91 (td, J = 5.5, 2.7 Hz, 1H), 7.88 - 7.80 (m, 1H), 7.74 (d, J = 8.0 Hz, 1H), 7.32 - 7.21 (m, 1H), 7.16 (d, J = 8.0 Hz, 1H), 6.45 (t, J = 6.8 Hz, 1H), 2.76 - 2.65 (m, 2H), 2.31 (d, J = 1.8 Hz, 3H), 1.89 (dd, J = 11.9, 5.6 Hz, 1H), 1.61 (s, 6H), 1.45 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).
2457	532.16	1.46	1H NMR (400 MHz, DMSO-d6) δ 12.55 (s, 1H), 8.26 (dd, J = 4.9, 1.8 Hz, 1H), 8.07 (dd, J = 7.9, 1.8 Hz, 1H), 7.90 - 7.69 (m, 3H), 7.30 (d, J = 8.0 Hz, 1H), 6.80 (dd, J = 7.8, 4.9 Hz, 3H), 3.99 (d, J = 1.0 Hz, 3H), 2.17 (dq, J = 15.9, 6.5 Hz, 1H), 1.86 (dd, J = 12.0, 5.6 Hz, 1H), 1.60 (s, 3H), 1.56 (s, 3H), 1.41 (t, J = 12.1 Hz, 1H), 0.76 (d, J = 6.3 Hz, 3H).
2458	542.40	1.77	
2459	529.20	1.62	
2460	481.30	2.30	
2461	501.41	2.35	
2462	517.32	2.56	
2463	522.35	1.85	1H NMR (400 MHz, DMSO-d6) δ 12.48 (s, 1H), 8.50 (d, J = 2.5 Hz, 1H), 7.89 (d, J = 2.4 Hz, 1H), 7.63 (t, J = 7.8 Hz, 1H), 7.58 (d, J = 8.4 Hz, 2H), 7.46 (d, J = 8.4 Hz, 2H), 7.22 (d, J = 7.2 Hz, 1H), 6.68 (d, J = 8.3 Hz, 1H), 6.49 (s, 2H), 2.56 (d, J = 8.8 Hz, 2H), 2.28 - 2.10 (m, 1H), 1.85 (dd, J = 11.9, 5.6 Hz, 1H), 1.55 (s, 3H), 1.53 (s, 3H), 1.40 (t, J = 12.1 Hz, 1H), 1.32 (s, 9H), 0.87 (d, J = 6.1 Hz, 3H).

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2464	460.29	1.76	
2465	524.52	2.33	
2466	535.32	2.20	
2467	447.30	1.39	1H NMR (400 MHz, DMSO-d6) δ 12.50 (s, 1H), 12.44 (s, 1H), 8.22 (dd, J = 7.2, 2.1 Hz, 1H), 7.81 (d, J = 5.8 Hz, 1H), 7.57 (d, J = 7.7 Hz, 1H), 6.47 - 6.37 (m, 2H), 2.66 (d, J = 8.8 Hz, 2H), 2.48 - 2.38 (m, 2H), 2.25 - 2.14 (m, 1H), 2.14 (s, 1H), 2.13 - 2.03 (m, 1H), 1.83 (dd, J = 11.8, 5.6 Hz, 1H), 1.52 (d, J = 11.0 Hz, 6H), 1.39 (t, J = 12.0 Hz, 1H), 0.93 - 0.83 (m, 9H).
2468	532.41	1.35	
2469	511.16	1.64	
2470	509.21	2.73	
2471	455.40	1.22	1H NMR (400 MHz, DMSO-d6) δ 12.89 (s, 1H), 12.57 (s, 1H), 8.27 (dd, J = 7.2, 2.1 Hz, 1H), 8.09 (s, 1H), 7.85 (dd, J = 6.4, 2.2 Hz, 1H), 7.49 (s, 1H), 7.48 - 7.39 (m, 2H), 6.46 (dd, J = 7.2, 6.4 Hz, 1H), 3.04 - 2.95 (m, 1H), 2.76 (t, J = 10.1 Hz, 1H), 2.40 (s, 3H), 2.31 - 2.18 (m, 1H), 1.87 (dd, J = 11.9, 5.7 Hz, 1H), 1.61 (d, J = 3.8 Hz, 6H), 1.44 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).
2472	490.32	1.50	
2473			1H NMR (400 MHz, DMSO-d6) δ 12.62 (s, 1H), 7.73 (d, J = 11.4 Hz, 1H), 7.64 (t, J = 7.9 Hz, 1H), 7.35 (s, 1H), 7.23 (dd, J = 15.3, 8.6 Hz, 2H), 6.96 (dt, J = 10.8, 2.4 Hz, 1H), 6.70 (d, J = 8.5 Hz, 1H), 6.53 (s, 1H), 3.82 (d, J = 6.6 Hz, 2H), 2.69 - 2.52 (m, 2H), 2.22 (s, 1H), 2.05 (hept, J = 6.6 Hz, 1H), 1.87 (dd, J = 11.9, 5.6 Hz, 1H), 1.54 (d, J = 6.7 Hz, 6H), 1.41 (t, J = 12.1 Hz, 1H), 0.98 (d, J = 6.6 Hz, 6H), 0.87 (d, J = 6.3 Hz, 3H).
2474	514.21	1.42	
2475	588.50	1.62	
2476	550.33	2.40	
2477	582.60	1.82	
2478	505.22	1.36	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2479	570.40	2.14	1H NMR (400 MHz, DMSO) δ 12.46 (s, 1H), 7.76 (d, J = 7.9 Hz, 1H), 7.64 (dd, J = 8.4, 7.3 Hz, 1H), 7.51 (d, J = 2.0 Hz, 1H), 7.47 - 7.40 (m, 1H), 7.30 (d, J = 7.9 Hz, 1H), 7.22 (dd, J = 7.3, 0.6 Hz, 1H), 6.89 (dt, J = 10.7, 2.3 Hz, 1H), 6.71 (dd, J = 8.4, 0.6 Hz, 1H), 6.50 (s, 2H), 3.89 - 3.75 (m, 3H), 3.15 (dd, J = 9.8, 4.4 Hz, 1H), 2.75 (ddd, J = 18.0, 13.7, 7.5 Hz, 2H), 2.03 (td, J = 13.3, 6.6 Hz, 2H), 1.68 - 1.47 (m, 3H), 1.34 (d, J = 7.5 Hz, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.88 (dt, J = 23.3, 11.7 Hz, 7H).
2480	483.40	1.38	1H NMR (400 MHz, DMSO-d6) δ 12.67 (s, 1H), 12.53 (s, 1H), 11.94 (s, 1H), 8.25 (dd, J = 7.3, 2.1 Hz, 1H), 7.89 - 7.81 (m, 2H), 7.78 (d, J = 8.2 Hz, 1H), 7.34 (d, J = 8.2 Hz, 1H), 7.26 (ddd, J = 8.5, 7.2, 1.7 Hz, 1H), 6.95 - 6.86 (m, 2H), 6.50 - 6.41 (m, 1H), 2.85 - 2.74 (m, 2H), 2.32 - 2.18 (m, 1H), 1.91 (dd, J = 11.9, 5.6 Hz, 1H), 1.54 (d, J = 5.3 Hz, 6H), 1.46 (t, J = 12.0 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).
2481	540.20	1.67	
2482	514.21	1.43	
2483	509.50	1.84	1H NMR (400 MHz, DMSO-d6) δ 12.56 (s, 1H), 12.55 - 12.49 (m, 1H), 8.24 (dd, J = 7.2, 2.0 Hz, 1H), 7.97 (d, J = 8.0 Hz, 2H), 7.83 (d, J = 6.5 Hz, 1H), 7.74 (d, J = 8.1 Hz, 1H), 7.36 (d, J = 8.0 Hz, 2H), 7.14 (d, J = 8.0 Hz, 1H), 6.44 (t, J = 6.8 Hz, 1H), 2.93 (p, J = 6.9 Hz, 1H), 2.78 - 2.69 (m, 2H), 2.24 (s, 1H), 1.88 (dd, J = 11.9, 5.5 Hz, 1H), 1.61 (s, 6H), 1.44 (t, J = 12.0 Hz, 1H), 1.23 (d, J = 6.9 Hz, 6H), 0.91 (d, J = 6.2 Hz, 3H).
2484	491.20	2.10	
2485	497.30	1.33	1H NMR (400 MHz, DMSO-d6) δ 12.58 (s, 2H), 8.24 (dd, J = 7.2, 2.2 Hz, 1H), 7.83 (s, 1H), 7.72 - 7.64 (m, 2H), 7.38 (td, J = 7.9, 1.8 Hz, 1H), 7.14 (d, J = 8.4 Hz, 1H), 7.05 (dd, J = 8.2, 6.5 Hz, 2H), 6.45 (t, J = 6.7 Hz, 1H), 3.82 (s, 3H), 2.78 - 2.66 (m, 2H), 2.22 (d, J =

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			12.7 Hz, 1H), 1.85 (dd, J = 12.0, 5.7 Hz, 1H), 1.54 (s, 6H), 1.42 (t, J = 12.0 Hz, 1H), 0.91 (d, J = 6.2 Hz, 3H).
2486	509.21	2.53	1H NMR (400 MHz, DMSO-d6) δ 12.85 (s, 1H), 8.21 - 8.15 (m, 2H), 7.91 - 7.88 (d, 1H), 7.44 - 7.26 (m, 5H), 7.03 - 7.01 (dd, 1H), 6.61 - 6.58 (dd, 1H), 3.05 - 2.94 (m, 1H), 2.63 - 2.59 (m, 1H), 2.26 - 2.19 (m, 1H), 1.90 - 1.83 (m, 1H), 1.51 (s, 3H), 1.47 (s, 3H), 1.47 - 1.36 (m, 1H), 1.12 - 1.09 (d, 3H), 1.10 - 1.07 (d, 3H), 0.89 - 0.87 (d, 3H)
2487	499.20	0.64	
2488	530.22	2.14	
2489	494.28	1.47	
2490	490.32	1.45	
2491	499.40	1.51	
2492	569.30	2.41	1H NMR (400 MHz, MeOD) δ 7.87 (d, J = 8.2 Hz, 1H), 7.76 (d, J = 8.2 Hz, 1H), 7.70 (dd, J = 8.3, 7.4 Hz, 1H), 7.54 (s, 1H), 7.46 - 7.38 (m, 2H), 6.83 (d, J = 8.4 Hz, 1H), 6.75 (dt, J = 10.5, 2.3 Hz, 1H), 3.82 (d, J = 6.5 Hz, 2H), 3.67 (dd, J = 12.3, 6.4 Hz, 1H), 2.66 (t, J = 12.5 Hz, 1H), 2.11 (dt, J = 13.3, 6.7 Hz, 1H), 1.69 (dd, J = 12.6, 6.3 Hz, 1H), 1.49 (dd, J = 38.8, 13.1 Hz, 2H), 1.18 (d, J = 5.6 Hz, 3H), 1.10 - 1.00 (m, 9H), 0.81 (s, 3H), 0.75 (s, 3H).
2493	524.52	2.39	
2494	510.16	1.42	
2495	518.22	2.30	
2496	533.31	1.15	1H NMR (400 MHz, DMSO) δ 12.62 (s, 1H), 12.54 (s, 1H), 11.31 (s, 1H), 8.25 (dd, J = 7.3, 2.1 Hz, 1H), 7.83 (s, 1H), 7.79 (d, J = 7.9 Hz, 1H), 7.54 - 7.49 (m, 1H), 7.47 (t, J = 5.8 Hz, 2H), 7.23 - 7.18 (m, 1H), 7.11 (d, J = 7.9 Hz, 1H), 6.70 (s, 1H), 6.45 (t, J = 6.9 Hz, 1H), 2.84 - 2.70 (m, 2H), 2.25 (s, 1H), 1.87 (dd, J = 11.5, 5.3 Hz, 1H), 1.59 (d, J = 3.9 Hz, 6H), 1.45 (t, J = 11.7 Hz, 1H), 0.92 (d, J = 6.3 Hz, 3H)..
2497	569.30	1.39	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2498	509.50	1.84	
2499	510.31	1.62	
2500	552.60	1.33	
2501	467.19	2.63	1H NMR (400 MHz, DMSO) δ 12.23 (s, 1H), 8.23 - 8.21 (dd, 1H), 7.90 - 7.87 (dd, 1H), 7.74 - 7.62 (m, 4H), 7.48 (t, J = 7.47, 1H), 7.25 - 7.22 (d, 1H), 7.06 - 7.03 (dd, 1H), 6.74 - 6.71 (d, 1H), 6.55 - 6.49 (m, 3H), 1.56 - 1.53 (d, 3H)
2502	508.31	2.50	
2503	531.12	2.53	
2504	550.33	1.74	
2505	498.53	1.09	
2506	517.32	1.87	
2507	471.39	2.22	
2508	599.60	2.25	
2509	467.19	2.14	
2510	538.31	1.68	
2511	548.30	2.27	
2512	523.36	1.74	
2513	502.31	2.51	
2514	551.30	1.60	1H NMR (400 MHz, DMSO-d6) δ 12.63 (s, 1H), 12.55 (s, 1H), 8.25 (dd, J = 7.3, 2.1 Hz, 1H), 7.83 (dd, J = 6.4, 2.2 Hz, 1H), 7.75 (d, J = 7.8 Hz, 1H), 7.71 (dd, J = 7.2, 2.3 Hz, 1H), 7.60 - 7.43 (m, 3H), 6.84 (d, J = 7.8 Hz, 1H), 6.45 (t, J = 6.8 Hz, 1H), 2.81 - 2.65 (m, 2H), 2.30 - 2.14 (m, 1H), 1.85 (dd, J = 11.9, 5.6 Hz, 1H), 1.51 (d, J = 8.1 Hz, 6H), 1.43 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).
2515	524.34	1.50	
2516	525.30	1.30	1H NMR (400 MHz, DMSO-d6) δ 12.62 - 12.42 (m, 2H), 8.24 (dd, J = 7.2, 2.1 Hz, 1H), 7.87 - 7.78 (m, 1H), 7.70 (d, J = 8.0 Hz, 1H), 7.61 - 7.51 (m, 2H), 7.10 (d, J = 8.0 Hz, 1H), 6.95 (d, J = 8.4 Hz, 1H), 6.44 (t, J = 6.8 Hz, 1H), 4.28 (s, 4H), 2.79 - 2.68 (m, 2H), 1.88 (dd, J = 11.5, 5.4 Hz, 1H), 1.60 (s, 6H), 1.44 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.2 Hz, 3H).
2517	509.21	1.51	
2518	524.22	2.59	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2519	454.30	1.27	
2520	652.60	1.67	
2521	533.22	1.61	
2522	524.19	1.36	
2523	535.32	2.19	
2524	499.31	2.49	
2525	502.33	1.46	
2526	510.41	2.54	
2527	515.30	1.53	
2528	433.27	2.30	
2529	504.11	2.45	
2530	511.15	1.59	
2531	537.43	1.46	
2532	556.20	2.00	1H NMR (400 MHz, DMSO) δ 12.66 (s, 1H), 7.81 (d, J = 7.9 Hz, 1H), 7.65 (t, J = 7.8 Hz, 1H), 7.50 - 7.41 (m, 3H), 7.22 (d, J = 7.3 Hz, 1H), 6.91 (d, J = 10.9 Hz, 1H), 6.72 (d, J = 8.4 Hz, 1H), 6.47 (s, 1H), 3.85 (d, J = 6.5 Hz, 2H), 3.51 (d, J = 8.5 Hz, 1H), 3.40 (d, J = 12.8 Hz, 1H), 3.13 (d, J = 11.8 Hz, 1H), 2.97 (d, J = 6.0 Hz, 1H), 2.06 (d, J = 9.3 Hz, 1H), 1.89 - 1.72 (m, 2H), 1.42 (s, 2H), 1.00 (d, J = 6.7 Hz, 6H), 0.87 (d, J = 6.6 Hz, 3H), 0.77 (d, J = 6.6 Hz, 3H).
2533	517.32	2.55	
2534	565.50	1.97	
2535	404.46	1.96	
2536	544.33	2.76	
2537	559.04	2.32	
2538	598.50	2.21	1H NMR (400 MHz, DMSO) δ 7.54 (d, J = 7.6 Hz, 1H), 7.46 (s, 1H), 7.38 (dd, J = 17.3, 9.4 Hz, 2H), 7.05 (dd, J = 14.0, 7.6 Hz, 2H), 6.79 (d, J = 10.6 Hz, 1H), 6.42 (d, J = 8.0 Hz, 1H), 5.99 (s, 2H), 3.82 (d, J = 6.7 Hz, 2H), 3.59 (t, J = 10.8 Hz, 1H), 3.08 - 2.98 (m, 1H), 2.05 (dt, J = 13.7, 6.8 Hz, 2H), 1.58 (dd, J = 26.1, 16.0 Hz, 8H), 0.99 (d, J = 6.7 Hz, 6H), 0.87 (d, J = 24.3 Hz, 9H).
2539	550.26	1.03	
2540	624.30	0.89	1H NMR (300 MHz, CDCl3) δ 8.73 (d, J = 8.2 Hz, 1H), 8.28 (d, J = 8.1 Hz, 1H), 8.04 (t, J = 7.9 Hz, 1H), 7.98 - 7.89 (m, 2H), 7.56 (dd, J = 8.2, 1.9 Hz, 1H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			7.51 (d, J = 2.4 Hz, 2H), 4.21 (d, J = 6.1 Hz, 2H), 4.10 (t, J = 6.4 Hz, 2H), 2.73 (t, J = 6.9 Hz, 2H), 2.64 - 2.47 (m, 4H), 2.11 (d, J = 2.8 Hz, 2H), 2.00 - 1.85 (m, 4H), 1.34 (d, J = 6.6 Hz, 6H), 1.25 (s, 6H), 1.11 (t, J = 7.4 Hz, 5H), 0.93 - 0.79 (m, 3H).
2541	525.40	1.55	
2542	485.50	1.52	
2543	546.27	1.16	
2544	488.20	2.47	
2545	530.14	1.78	
2546	533.22	1.98	
2547	524.30	1.28	
2548	454.30	1.27	
2549	525.60	1.72	1H NMR (400 MHz, DMSO-d6) δ 12.59 (s, 1H), 12.53 (s, 1H), 8.24 (d, J = 7.4 Hz, 1H), 7.83 (d, J = 6.0 Hz, 1H), 7.74 (d, J = 7.9 Hz, 1H), 7.62 (s, 1H), 7.58 (d, J = 7.5 Hz, 1H), 7.37 (t, J = 8.0 Hz, 1H), 7.19 (d, J = 8.0 Hz, 1H), 7.00 - 6.93 (m, 1H), 6.45 (t, J = 7.0 Hz, 1H), 4.66 (p, J = 6.1 Hz, 1H), 2.75 (q, J = 9.8, 9.0 Hz, 2H), 2.24 (s, 1H), 1.89 (dd, J = 11.9, 5.5 Hz, 1H), 1.62 (s, 6H), 1.45 (t, J = 12.1 Hz, 1H), 1.31 (d, J = 6.0 Hz, 6H), 0.91 (d, J = 6.3 Hz, 3H).
2550	512.51	1.80	
2551	497.51	2.16	1H NMR (400 MHz, DMSO-d6) δ 12.90 (s, 1H), 8.21 - 8.15 (m, 2H), 7.92 - 7.89 (d, 1H), 7.50 - 7.47 (dd, 1H) 7.39 - 7.33 (m, 2H), 6.87 - 6.84 (dd, 1H), 6.75 (s, 1H), 6.75 - 6.72 (d, 1H), 6.61 - 6.58 (dd, 1H), 3.73 (s, 3H), 2.27 - 2.18 (m, 1H), 1.89 - 1.82 (m, 1H), 1.51 (s, 3H), 1.47 (s, 3H), 1.47 - 1.35 (m, 1H), 0.89 - 0.87 (d, 3H)
2552	568.20	2.00	
2553	458.40	1.26	
2554	490.30	1.86	1H NMR (400 MHz, Methanol-d4) δ 8.27 (d, J = 2.8 Hz, 1H), 7.77 (d, J = 8.3 Hz, 1H), 7.00 (d, J = 8.3 Hz, 1H), 6.00 (d, J = 2.8 Hz, 1H), 3.90 (s, 2H), 3.15 - 3.04 (m, 2H), 2.41 (dd, J = 11.2, 5.6 Hz, 1H), 1.98 (dd, J = 11.9, 5.8 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 1.67 (d, J = 13.8 Hz, 6H), 1.61 (d, J = 12.2 Hz, 1H), 1.36 (ddd, J = 9.5, 7.0, 4.7 Hz, 1H), 1.28 - 1.13 (m, 2H), 1.09 (d, J = 6.4 Hz, 4H), 1.04 (s, 9H).
2555	482.60	1.46	
2556	511.33	0.90	
2557	510.30	1.54	
2558	552.39	1.97	
2559	502.50	1.61	
2560	542.32	3.22	1H NMR (400 MHz, DMSO-d6) δ 12.57 (s, 1H), 8.06 - 7.96 (m, 2H), 7.78 - 7.70 (m, 2H), 7.67 (dd, J = 8.4, 6.8 Hz, 2H), 7.51 - 7.45 (m, 1H), 7.40 (dt, J = 10.1, 1.8 Hz, 1H), 7.27 (d, J = 8.0 Hz, 1H), 6.89 (dt, J = 10.8, 2.2 Hz, 1H), 3.83 (d, J = 6.6 Hz, 2H), 2.07 (ddt, J = 23.2, 13.3, 6.4 Hz, 2H), 1.83 (dd, J = 11.9, 5.7 Hz, 1H), 1.57 (d, J = 18.0 Hz, 6H), 1.38 (t, J = 12.1 Hz, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.66 (d, J = 6.3 Hz, 3H).
2561	444.38	2.06	
2562	514.20	1.38	
2563	532.32	2.58	1H NMR (400 MHz, DMSO-d6) δ 12.86 (s, 1H), 8.23 - 8.13 (m, 2H), 7.92 - 7.89 (d, 1H), 7.69 - 7.65 (d, 2H), 7.49 - 7.38 (m, 4H), 7.28 - 7.24 (d, 2H), 6.57 - 6.54 (dd, 1H), 6.28 (t, J = 2.20, 2H), 2.28 - 2.18 (m, 1H), 1. 89 - 1.82 (m, 1H), 1.51 (s, 3H), 1.47 (s, 3H), 1.46 - 1.34 (m, 1H), 0.90 - 0.87 (d, 3H)
2564	511.09	1.41	
2565	529.32	1.96	
2566	484.50	1.52	
2567	533.30	1.52	
2568	570.40	1.30	1H NMR (400 MHz, DMSO-d6) δ 12.40 (s, 1H), 7.69 (d, J = 8.0 Hz, 1H), 7.62 (dd, J = 8.5, 7.3 Hz, 1H), 7.19 (d, J = 7.0 Hz, 1H), 6.85 (d, J = 8.1 Hz, 1H), 6.80 (dq, J = 4.3, 1.8 Hz, 1H), 6.68 (d, J = 8.4 Hz, 1H), 5.70 (s, 1H), 4.20 (q, J = 2.4 Hz, 2H), 3.39 - 3.32 (m, 2H), 2.55 (d, J = 8.7 Hz, 2H), 2.27 (s, 2H), 2.18 (dd, J = 12.7, 6.4 Hz, 1H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1.84 (dd, J = 11.8, 5.6 Hz, 1H), 1.53 (s, 6H), 1.40 (d, J = 12.2 Hz, 1H), 1.26 (s, 9H), 0.86 (d, J = 6.3 Hz, 3H).
2569	509.21	1.98	
2570	502.29	1.47	
2571	530.22	2.49	
2572	480.10	2.68	
2573	610.20	2.23	
2574	430.27	1.81	
2575	525.37	1.08	
2576	553.40	2.74	1H NMR (400 MHz, DMSO) δ 8.61 (d, J = 2.3 Hz, 1H), 8.08 (d, J = 2.5 Hz, 1H), 7.56 (d, J = 7.7 Hz, 1H), 7.47 (t, J = 7.8 Hz, 1H), 6.97 (dd, J = 7.4, 3.2 Hz, 2H), 6.49 (d, J = 8.3 Hz, 1H), 6.23 (s, 2H), 5.16 (h, J = 6.3 Hz, 1H), 2.97 (t, J = 10.6 Hz, 1H), 2.91 - 2.81 (m, 1H), 2.17 (s, 3H), 2.09 (qt, J = 11.8, 7.9, 6.2 Hz, 1H), 1.81 (dd, J = 11.5, 5.6 Hz, 1H), 1.75 - 1.64 (m, 2H), 1.61 (s, 3H), 1.56 (s, 3H), 1.36 (t, J = 12.0 Hz, 1H), 1.28 (d, J = 6.1 Hz, 3H), 0.92 (t, J = 7.4 Hz, 3H), 0.80 (d, J = 6.2 Hz, 3H).
2577	472.50	1.57	1H NMR (400 MHz, DMSO) δ 12.35 (s, 1H), 7.59 (d, J = 7.7 Hz, 2H), 7.14 (d, J = 6.7 Hz, 1H), 6.73 - 6.58 (m, 2H), 6.43 (s, 2H), 2.79 (s, 1H), 2.62 (s, 1H), 1.80 - 1.59 (m, 2H), 1.56 (s, 3H), 1.51 (s, 3H), 1.27 (s, 9H), 1.13 - 1.03 (m, 1H), 0.71 - 0.58 (m, 1H), 0.56 - 0.46 (m, 1H), 0.46 - 0.33 (m, 1H), 0.24 (td, J = 9.2, 5.0 Hz, 1H), 0.12 - 0.01 (m, 1H).
2578	543.40	1.92	1H NMR (400 MHz, DMSO-d6) δ 12.52 (s, 2H), 8.23 (d, J = 7.4 Hz, 1H), 8.20 (d, J = 2.7 Hz, 1H), 7.87 - 7.74 (m, 2H), 6.88 (d, J = 8.2 Hz, 1H), 6.51 - 6.38 (m, 1H), 6.13 (d, J = 2.7 Hz, 1H), 3.89 (s, 2H), 2.71 (broad d, J = 8.6 Hz, 2H), 2.22 (br s, 1H), 1.88 (dd, J = 12.0, 5.6 Hz, 1H), 1.56 (s, 3H), 1.55 (s, 3H), 1.43 (t, J = 12.1 Hz, 1H), 1.00 (s, 9H), 0.89 (d, J = 6.3 Hz, 3H).
2579	510.41	2.16	
2580	510.11	2.33	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2581	570.20	1.88	1H NMR (400 MHz, DMSO) δ 8.41 (d, J = 8.2 Hz, 1H), 8.24 (d, J = 8.3 Hz, 1H), 7.65 - 7.49 (m, 3H), 7.19 (d, J = 7.2 Hz, 1H), 7.01 (d, J = 10.9 Hz, 1H), 6.65 (d, J = 8.3 Hz, 1H), 6.30 (s, 1H), 3.87 (d, J = 6.5 Hz, 2H), 3.82 - 3.74 (m, 1H), 3.18 (d, J = 10.9 Hz, 1H), 2.06 (dt, J = 13.2, 6.6 Hz, 2H), 2.01 - 1.59 (m, 4H), 1.58 (s, 3H), 1.13 (s, 3H), 0.99 (t, J = 6.9 Hz, 9H).
2582	555.10	1.47	1H NMR (400 MHz, DMSO) δ 12.41 (s, 0H), 3.32 - 3.32 (m, 6H), 8.66 (d, J = 2.4 Hz, 2H), 8.14 (d, J = 2.4 Hz, 2H), 7.68 (s, 2H), 7.55 (s, 1H), 7.12 (s, 4H), 6.60 (s, 1H), 6.33 (s, 0H), 4.52 - 4.39 (m, 5H), 3.74 - 3.65 (m, 5H), 2.86 - 2.61 (m, 1H), 2.21 (s, 7H), 2.19 - 2.10 (m, 5H), 1.85 (dd, J = 11.7, 5.6 Hz, 2H), 1.61 (s, 7H), 1.58 (s, 7H), 1.40 (t, J = 11.9 Hz, 2H), 0.86 (d, J = 6.3 Hz, 8H).
2583	509.20	1.57	
2584	491.30	1.51	
2585	515.21	2.65	
2586	521.30	1.72	1H NMR (400 MHz, DMSO) δ 7.62 (t, J = 7.9 Hz, 1H), 7.51 (d, J = 14.0 Hz, 1H), 7.16 (d, J = 7.3 Hz, 1H), 6.67 (d, J = 8.3 Hz, 1H), 3.73 (t, J = 4.9 Hz, 2H), 3.48 - 3.42 (m, 2H), 3.33 (s, 2H), 2.70 - 2.54 (m, 2H), 2.27 - 2.15 (m, 1H), 1.85 (dd, J = 11.9, 5.9 Hz, 1H), 1.51 - 1.32 (m, 8H), 1.18 (d, J = 10.2 Hz, 7H), 0.88 (d, J = 6.2 Hz, 3H).
2587	490.32	1.52	
2588	586.35	1.88	
2589	582.20	2.05	
2590	496.27	1.11	
2591	519.12	2.49	
2592	496.60	1.45	
2593	534.22	1.07	
2594	554.50	1.78	1H NMR (400 MHz, DMSO-d6) δ 12.51 (broad s, 1H), 8.25-8.23 (m, 2H), 8.02 (dd, J = 7.8, 1.9 Hz, 1H), 7.87 (d, J = 8.2 Hz, 1H), 7.01 (d, J = 8.2 Hz, 1H), 6.76 (dd, J = 7.8, 4.7 Hz, 1H),

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			6.67 (br s, 2H), 6.16 (d, J = 2.7 Hz, 1H), 3.90 (s, 2H), 3.49 (br d, J = 9.9 Hz, 1H), 3.39 (br d, J = 8.9 Hz, 1H), 2.47 - 2.30 (m, 2H), 2.26 (dd, J = 11.8, 5.2 Hz, 1H), 1.92-1.90 (m, 2H), 1.85 - 1.72 (m, 2H), 1.61 (br s, 1H), 1.51 (t, J = 11.5 Hz, 1H), 1.01 (s, 9H), 0.78 (d, J = 6.4 Hz, 3H).
2595	537.43	2.64	
2596	651.50	1.82	
2597	470.49	1.34	
2598	519.12	2.54	1H NMR (400 MHz, DMSO-d6) δ 12.92 (s, 1H), 8.27 - 8.17 (m, 2H), 7.97 - 7.94 (d, 1H), 7.59 - 7.38 (m, 5H), 6.65 - 6.62 (dd, 1H), 2.63 - 2.54 (m, 1H), 2.28 - 2.20 (m, 1H), 1.89 - 1.82 (m, 1H), 1.50 (s, 3H), 1.46 (s, 3H) , 1.40- 1.35 (m, 1H), 0.89 - 0.87 (d, 3H)
2599	458.18	1.80	
2600	546.13	2.70	
2601	541.33	1.65	
2602	476.19	2.38	
2603	542.20	1.85	
2604	540.20	1.67	1H NMR (400 MHz, DMSO) δ 12.50 (s, 1H), 8.25 (dd, J = 4.8, 2.1 Hz, 2H), 8.03 (dd, J = 7.8, 1.8 Hz, 1H), 7.88 (d, J = 8.3 Hz, 1H), 7.01 (d, J = 8.2 Hz, 1H), 6.75 (dt, J = 26.8, 13.4 Hz, 3H), 6.15 (d, J = 2.7 Hz, 1H), 3.99 (d, J = 6.6 Hz, 2H), 3.49 (d, J = 9.5 Hz, 1H), 3.38 (d, J = 8.7 Hz, 1H), 2.40 (dt, J = 27.2, 9.9 Hz, 2H), 2.26 (dd, J = 11.8, 4.8 Hz, 1H), 2.10 - 2.01 (m, 1H), 1.91 (dd, J = 15.4, 8.5 Hz, 2H), 1.78 (dd, J = 12.4, 7.5 Hz, 2H), 1.67 - 1.56 (m, 1H), 1.52 (t, J = 11.5 Hz, 1H), 0.98 (d, J = 6.7 Hz, 6H), 0.78 (d, J = 6.3 Hz, 3H).
2605	486.33	1.53	
2606	515.40	1.46	
2607	524.23	1.48	
2608	560.27	1.30	
2609	458.18	2.31	
2610	505.01	2.51	1H NMR (400 MHz, DMSO) δ 12.82 (s, 1H), 8.19 - 8.17 (dd, 1H), 8.05 (t, J

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			= 7.91, 1H), 7.79 - 7.72 (m, 2H), 7.56 - 7.53 (d, 2H), 7.42 - 7.39 (d, 2H), 7.21 - 7.18 (d, 1H), 6.64 - 6.61 (dd, 1H), 5.38 (s, 2H), 4.18 (s, 1H), 2.26 - 2.10 (m, 2H), 2.00 - 1.89 (m, 1H), 1.79 - 1.72 (m, 1H), 1.50 (s, 3H), 1.45 (s, 3H), 1.28 (t, J = 11.86, 1H), 0.61 - 0.59 (d, 3H)
2611	517.32	2.42	
2612	569.70	1.44	
2613	498.53	1.07	
2614	512.18	1.25	
2615	438.07	1.77	
2616	498.50	1.58	
2617	419.16	2.05	
2618	549.43	2.73	
2619	481.30	2.47	
2620	529.32	1.56	
2621	537.43	1.79	
2622	557.40	1.39	1H NMR (400 MHz, DMSO-d6) δ 12.39 (s, 1H), 7.69 (d, J = 8.1 Hz, 1H), 7.62 (t, J = 7.1 Hz, 1H), 7.19 (d, J = 7.1 Hz, 1H), 6.91 (s, 1H), 6.76 (s, 1H), 6.68 (d, J = 8.5 Hz, 1H), 4.80 (h, J = 6.1 Hz, 1H), 4.35 (d, J = 16.5 Hz, 2H), 3.49 (t, J = 5.9 Hz, 2H), 2.55 (d, J = 8.9 Hz, 1H), 2.35 - 2.27 (m, 2H), 2.26 - 2.08 (m, 1H), 1.85 (dd, J = 11.9, 5.6 Hz, 1H), 1.53 (d, J = 2.6 Hz, 6H), 1.39 (t, J = 12.1 Hz, 1H), 1.21 (d, J = 1.8 Hz, 1H), 1.19 (s, 3H), 1.18 (s, 3H), 0.87 (d, J = 6.3 Hz, 3H).
2623	524.57	1.04	
2624	557.40	1.82	
2625	513.41	2.78	
2626	495.40	1.62	1H NMR (400 MHz, DMSO-d6) δ 12.57 (s, 1H), 12.52 (s, 1H), 8.24 (dd, J = 7.3, 2.2 Hz, 1H), 7.83 (d, J = 7.4 Hz, 1H), 7.73 (d, J = 8.0 Hz, 1H), 7.14 (d, J = 8.0 Hz, 1H), 7.07 (s, 1H), 6.44 (t, J = 6.9 Hz, 1H), 2.77 - 2.63 (m, 2H), 2.33 (s, 6H), 2.29 - 2.15 (m, 1H), 1.88 (dd, J = 11.7, 5.5 Hz, 1H), 1.61 (s, 6H), 1.51 - 1.37 (m, 2H), 0.91 (d, J = 6.3 Hz, 3H).
2627	539.23	2.51	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2628	515.21	1.38	
2629	539.50	1.78	1H NMR (400 MHz, DMSO-d6) δ 12.55 (d, J = 13.7 Hz, 2H), 8.24 (dd, J = 7.3, 2.1 Hz, 1H), 7.85 - 7.78 (m, 1H), 7.73 (dd, J = 7.7, 1.8 Hz, 1H), 7.69 (d, J = 8.0 Hz, 1H), 7.35 (ddd, J = 8.6, 7.3, 1.8 Hz, 1H), 7.18 (d, J = 8.0 Hz, 1H), 7.11 (dd, J = 8.4, 1.1 Hz, 1H), 7.08 - 7.01 (m, 1H), 6.44 (t, J = 6.8 Hz, 1H), 3.84 (dd, J = 6.4, 2.2 Hz, 2H), 2.76 - 2.65 (m, 2H), 2.23 (s, 1H), 2.10 - 1.98 (m, 1H), 1.86 (dd, J = 11.8, 5.6 Hz, 1H), 1.55 (d, J = 4.8 Hz, 6H), 1.48 - 1.38 (m, 1H), 0.99 (d, J = 6.7 Hz, 6H), 0.91 (d, J = 6.3 Hz, 3H).
2630	487.30	2.91	
2631	578.20	2.10	
2632	528.12	2.32	
2633	595.70	1.57	1H NMR (400 MHz, DMSO-d6) δ 12.40 (s, 1H), 7.69 (dd, J = 8.1, 2.7 Hz, 1H), 7.62 (t, 1H), 7.19 (d, J = 7.4 Hz, 1H), 6.95 (d, J = 8.1 Hz, 0.5 H), 6.86 (d, J = 8.1 Hz, 0.5 H), 6.83 - 6.74 (m, 1H), 6.68 (d, J = 8.5 Hz, 1H), 4.43 (d, J = 11.6 Hz, 2H), 2.55 (d, J = 8.9 Hz, 1H), 2.34 (d, J = 6.2 Hz, 1H), 2.25 (d, J = 6.7 Hz, 3H), 2.20 (s, 1H), 1.85 (ddd, J = 11.9, 8.8, 5.6 Hz, 1H), 1.70 - 1.58 (m, 6H), 1.55 (two singlets, 3H), 1.53 ((two singlets, 3H), 1.39 (td, J = 12.2, 3.8 Hz, 2H), 1.30 - 1.05 (m, 4H), 1.05 - 0.88 (m, 3H), 0.87 (d, J = 6.3 Hz, 3H).
2634	573.20	1.73	1H NMR (400 MHz, Methanol-d4) δ 7.69 (s, 1H), 7.32 (t, J = 7.9 Hz, 1H), 7.26 (s, 1H), 7.08 (dd, J = 7.3, 0.8 Hz, 1H), 6.62 - 6.51 (m, 2H), 6.44 (dd, J = 19.3, 9.7 Hz, 2H), 3.58 (s, 1H), 3.49 (d, J = 6.4 Hz, 2H), 2.52 (t, J = 0.8 Hz, 1H), 2.45 (s, 1H), 1.90 (s, 1H), 1.82 - 1.69 (m, 2H), 1.53 (s, 1H), 1.04 (s, 3H), 0.92 (s, 6H), 0.74 (d, J = 6.7 Hz, 6H).
2635	509.30	1.63	
2636	506.19	1.26	
2637	497.51	2.10	
2638	498.50	1.65	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2639	466.29	2.26	1H NMR (400 MHz, DMSO-d6) δ 12.75 (s, 1H), 9.62 (s, 1H), 8.14 - 8.12 (dd, 1H), 7.87 - 7.70 (m, 4H), 7.47 - 7.44 (d, 1H), 7.26 (t, J = 7.91, 2H), 7.09 - 7.06 (d, 1H), 6.94 (t, J = 7.36, 1H), 6.53 - 6.50 (dd, 1H), 2.47 - 2.43 (m, 1H), 1.87 - 1.70 (m, 2H), 1.45 (s, 3H), 1.41 (s, 3H), 1.29 (t, J = 11.86, 1H), 0.73 - 0.71 (d, 3H)
2640	516.41	2.36	
2641	526.32	2.54	
2642	538.33	2.48	
2643	532.32	1.45	
2644	516.30	1.29	
2645	545.23	2.66	
2646	497.22	1.31	1H NMR (400 MHz, DMSO-d6) δ 8.27 (ddd, J = 11.0, 5.1, 2.6 Hz, 2H), 8.06 (dt, J = 7.8, 2.3 Hz, 1H), 7.82 (dd, J = 7.9, 2.9 Hz, 1H), 7.58 (s, 1H), 7.39 (s, 1H), 7.34 (dd, J = 7.8, 2.9 Hz, 1H), 6.97 - 6.56 (m, 4H), 3.97 - 3.85 (m, 3H), 2.59 (s, 2H), 2.16 (s, 1H), 1.85 (s, 1H), 1.58 (dd, J = 15.1, 2.8 Hz, 6H), 1.40 (t, J = 12.3 Hz, 1H), 0.76 (dd, J = 6.3, 3.0 Hz, 3H).
2647	486.33	1.80	1H NMR (400 MHz, DMSO-d6) δ 12.41 (s, 1H), 8.16 (d, J = 2.3 Hz, 1H), 7.68 (d, J = 2.3 Hz, 1H), 7.62 (dd, J = 8.4, 7.3 Hz, 1H), 7.20 (d, J = 7.2 Hz, 1H), 6.68 (d, J = 8.2 Hz, 1H), 6.50 (s, 2H), 6.35 - 6.27 (m, 1H), 6.16 - 6.07 (m, 1H), 2.56 (td, J = 10.3, 7.0 Hz, 2H), 2.17 (q, J = 7.1 Hz, 3H), 1.82 (dd, J = 11.9, 5.7 Hz, 1H), 1.60 (dh, J = 13.4, 6.8, 6.4 Hz, 1H), 1.50 (s, 3H), 1.48 (s, 3H), 1.39 - 1.29 (m, 3H), 0.90 (d, J = 6.6 Hz, 6H), 0.85 (d, J = 6.3 Hz, 3H).
2648	459.09	2.52	
2649	550.24	1.80	
2650	569.50	1.41	
2651	525.20	1.44	
2652	503.40	1.60	1H NMR (400 MHz, DMSO-d6) δ 12.66 (s, 1H), 12.54 (d, J = 6.2 Hz, 1H), 8.25 (dd, J = 7.3, 2.1 Hz, 1H), 7.83 (td, J = 6.4, 2.2 Hz, 1H), 7.79 -

Cmpd No	LC/MS M+1	LC/RT min	NMR
			7.70 (m, 3H), 7.36 - 7.28 (m, 2H), 6.45 (t, J = 6.8 Hz, 1H), 2.81 - 2.68 (m, 2H), 2.24 (dq, J = 11.6, 6.2 Hz, 1H), 1.90 (dd, J = 11.9, 5.6 Hz, 1H), 1.45 (t, J = 12.1 Hz, 1H), 0.91 (d, J = 6.3 Hz, 3H).
2653	500.12	1.43	
2654	521.22	2.44	
2655	547.60	1.65	
2656	501.11	1.72	
2657	605.32	1.71	
2658	513.20	0.64	
2659	491.20	2.48	
2660	498.53	1.05	
2661	574.40	1.55	
2662	540.43	2.40	
2663	535.02	2.76	
2664	526.18	1.32	
2665	528.19	1.48	1H NMR (400 MHz, DMSO-d6) δ 12.53 (s, 1H), 8.25 (dd, J = 4.8, 1.8 Hz, 1H), 8.06 (dd, J = 7.8, 1.8 Hz, 1H), 7.92 - 7.69 (m, 2H), 7.62 (ddd, J = 8.5, 4.3, 2.1 Hz, 1H), 7.37 - 7.17 (m, 2H), 6.78 (dd, J = 7.8, 4.8 Hz, 3H), 4.19 (q, J = 7.0 Hz, 2H), 2.54 (d, J = 10.2 Hz, 2H), 2.15 (s, 1H), 1.85 (dd, J = 11.7, 5.6 Hz, 1H), 1.61 (s, 3H), 1.57 (s, 3H), 1.38 (t, J = 6.9 Hz, 4H), 0.76 (d, J = 6.3 Hz, 3H).
2666	488.33	1.33	
2667	496.27	1.33	1H NMR (400 MHz, DMSO) δ 12.51 (s, 1H), 8.25 (dd, J = 4.8, 1.9 Hz, 1H), 8.06 (d, J = 7.9 Hz, 1H), 8.04 - 7.98 (m, 2H), 7.74 (d, J = 8.0 Hz, 1H), 7.18 (d, J = 8.0 Hz, 1H), 7.05 (dd, J = 9.5, 2.5 Hz, 2H), 6.78 (dd, J = 7.8, 4.8 Hz, 1H), 3.81 (s, 3H), 2.56 - 2.51 (m, 2H), 2.15 (s, 1H), 1.85 (dd, J = 11.9, 5.7 Hz, 1H), 1.60 (s, 3H), 1.58 (s, 3H), 1.40 (t, J = 12.1 Hz, 1H), 0.76 (d, J = 6.3 Hz, 3H).
2668	498.24	1.23	
2669	669.52	0.76	1H NMR (300 MHz, CDCl3) δ 10.43 (s, 1H), 8.57 (d, J = 8.0 Hz, 1H), 7.86 (d, J = 7.2 Hz, 2H), 7.72 (d, J = 4.1 Hz, 1H), 7.66 (s, 1H), 7.62 (s, 1H), 7.58 (d, J =

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			4.4 Hz, 1H), 7.56 - 7.56 (m, 1H), 7.55 (s, 1H), 7.09 (s, 1H), 7.01 (d, J = 9.1 Hz, 1H), 6.77 - 6.70 (m, 1H), 6.66 (d, J = 10.3 Hz, 1H), 4.74 (s, 2H), 3.60 (d, J = 6.4 Hz, 2H), 2.28 (s, 6H), 2.10 - 1.96 (m, 1H), 0.98 (d, J = 6.7 Hz, 6H).
2670	610.20	0.83	1H NMR (400 MHz, Methanol-d4) δ 8.67 (dd, J = 4.8, 1.9 Hz, 1H), 8.48 (dd, J = 8.0, 1.8 Hz, 1H), 8.32 (d, J = 2.8 Hz, 1H), 8.09 (dd, J = 8.5, 1.6 Hz, 1H), 7.66 (dd, J = 8.3, 1.5 Hz, 1H), 7.61 (dd, J = 8.0, 4.7 Hz, 1H), 6.01 (d, J = 2.8 Hz, 1H), 3.92 (d, J = 1.8 Hz, 2H), 3.76 - 3.65 (m, 4H), 3.34 (d, J = 1.9 Hz, 5H), 2.04 (p, J = 5.9 Hz, 4H), 1.85 - 1.70 (m, 2H), 1.03 (d, J = 1.9 Hz, 9H).
2671	691.40	0.83	1H NMR (300 MHz, CDCl3) δ 8.47 (d, J = 8.1 Hz, 1H), 7.76 - 7.68 (m, 1H), 7.64 (d, J = 7.3 Hz, 1H), 7.52 (d, J = 8.1 Hz, 1H), 7.47 (s, 2H), 7.04 - 6.95 (m, 3H), 6.76 (d, J = 8.2 Hz, 1H), 6.63 (dt, J = 10.3, 2.2 Hz, 1H), 3.56 (t, J = 7.0 Hz, 2H), 2.13 - 2.06 (m, 6H), 1.01 (t, J = 6.6 Hz, 6H), 0.85 (t, J = 6.7 Hz, 3H), 0.47 (s, 1H).
2672	474.50	1.18	
2673	557.40	2.15	1H NMR (400 MHz, DMSO-d6) δ 11.58 (s, 1H), 7.89 (s, 1H), 7.62 - 7.53 (m, 1H), 7.11 (d, J = 7.3 Hz, 1H), 6.88 - 6.79 (m, 2H), 6.75 - 6.69 (m, 1H), 6.66 (d, J = 8.3 Hz, 1H), 6.52 (s, 2H), 3.83 (t, J = 9.1 Hz, 1H), 3.74 (d, J = 6.5 Hz, 2H), 3.12 (t, J = 10.4 Hz, 1H), 2.47 - 2.35 (m, 1H), 2.02 (dd, J = 12.8, 6.3 Hz, 2H), 1.61 (d, J = 12.2 Hz, 1H), 1.56 (s, 3H), 1.45 (s, 3H), 1.10 (d, J = 6.4 Hz, 3H), 0.96 (d, J = 6.8 Hz, 6H).
2674	480.20	1.97	1H NMR (400 MHz, Methanol-d4) δ 8.50 (d, J = 2.8 Hz, 1H), 7.95 - 7.87 (m, 2H), 7.84 (dd, J = 8.1, 1.2 Hz, 1H), 7.46 - 7.38 (m, 2H), 7.36 (dd, J = 7.8, 1.8 Hz, 1H), 7.30 (dd, J = 8.2, 1.2 Hz, 1H), 6.94 - 6.88 (m, 1H), 3.18 - 3.01 (m, 2H), 2.51 - 2.35 (m, 1H), 2.01 (dd, J = 11.8, 5.8 Hz, 1H), 1.76 - 1.66 (m,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			6H), 1.63 (d, J = 12.0 Hz, 1H), 1.39 (dt, J = 8.7, 5.3 Hz, 1H), 1.26 - 1.14 (m, 2H), 1.10 (dd, J = 8.7, 5.4 Hz, 4H).
2675	474.50	1.17	
2676	466.20	1.78	1H NMR (400 MHz, Methanol-d4) δ 8.60 (d, J = 2.7 Hz, 1H), 8.22 (d, J = 8.3 Hz, 1H), 7.97 - 7.90 (m, 2H), 7.66 (d, J = 8.3 Hz, 1H), 7.48 - 7.40 (m, 2H), 7.39 - 7.33 (m, 1H), 6.94 (d, J = 2.7 Hz, 1H), 4.20 (q, J = 6.1 Hz, 2H), 3.09 (tt, J = 8.1, 4.7 Hz, 1H), 2.30 - 2.19 (m, 2H), 1.90 - 1.79 (m, 2H), 1.33 - 1.25 (m, 8H), 1.18 - 1.10 (m, 2H).
2677	562.60	1.75	
2678	478.40	1.25	
2679	556.40	1.47	
2680	574.40	1.27	
2681	626.20	1.12	
2682	582.30	1.12	
2683	568.40	1.00	
2684	552.40	1.14	
2685	538.30	0.97	
2686	539.30	1.04	
2687	541.30	1.04	
2688	555.00	2.01	1H NMR (400 MHz, DMSO) δ 12.32 (s, 1H), 8.85 (d, J = 2.4 Hz, 1H), 8.33 - 8.26 (m, 1H), 7.89 (d, J = 8.6 Hz, 1H), 7.72 (d, J = 7.9 Hz, 1H), 7.20 (d, J = 8.0 Hz, 1H), 6.77 (s, 1H), 6.14 (d, J = 8.6 Hz, 1H), 5.39 - 5.25 (m, 1H), 3.82 (s, 3H), 2.15 (s, 1H), 1.85 (dd, J = 11.8, 5.6 Hz, 1H), 1.59 (d, J = 11.4 Hz, 6H), 1.41 (t, J = 12.2 Hz, 1H), 1.32 (d, J = 6.2 Hz, 6H), 0.76 (d, J = 6.3 Hz, 3H).
2689	540.40	1.18	
2690	540.40	1.62	
2691	526.40	1.47	
2692	558.50	1.30	
2693	544.50	1.26	
2694	556.50	1.83	
2695	496.30	0.64	1H NMR (400 MHz, Methanol-d4) δ 8.66 (dd, J = 7.8, 1.8 Hz, 1H), 8.35 (d, J = 2.8 Hz, 1H), 8.24 - 8.13 (m, 2H), 7.49 (d, J = 8.5 Hz, 1H), 7.16 - 7.07 (m, 1H), 6.31 - 6.23 (m, 1H), 3.24 (d, J

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			= 9.6 Hz, 1H), 3.10 (t, J = 10.1 Hz, 1H), 2.56 (d, J = 11.8 Hz, 1H), 2.13 (dd, J = 12.4, 6.6 Hz, 1H), 2.01 (dq, J = 11.6, 3.6 Hz, 1H), 1.72 (t, J = 11.9 Hz, 1H), 1.56 (dd, J = 27.3, 5.2 Hz, 6H), 1.38 - 1.22 (m, 1H), 1.08 (d, J = 6.4 Hz, 3H), 1.00 (ddt, J = 8.4, 4.5, 2.2 Hz, 3H), 0.82 (ddd, J = 7.2, 4.9, 2.8 Hz, 3H).
2696	532.20	0.69	1H NMR (400 MHz, Methanol-d4) δ 8.59 - 8.49 (m, 2H), 8.28 - 8.20 (m, 1H), 8.09 (dd, J = 10.7, 5.2 Hz, 1H), 8.02 - 7.91 (m, 2H), 7.61 - 7.36 (m, 5H), 7.11 - 6.94 (m, 2H), 3.11 - 2.89 (m, 2H), 2.48 (d, J = 15.9 Hz, 1H), 2.06 (dd, J = 11.8, 6.4 Hz, 1H), 1.63 (dd, J = 14.0, 5.6 Hz, 6H), 1.00 (d, J = 6.9 Hz, 3H).
2697	551.50	1.74	1H NMR (400 MHz, DMSO-d6) δ 12.48 (broad s, 1H), 8.24 (dd, J = 4.8, 1.9 Hz, 1H), 8.20 (d, J = 2.8 Hz, 1H), 8.02 (dd, J = 7.8, 1.8 Hz, 1H), 7.85 (d, J = 8.2 Hz, 1H), 6.95 (d, J = 8.3 Hz, 1H), 6.76 (dd, J = 7.8, 4.8 Hz, 1H), 6.70 (br s, 2H), 6.14 (d, J = 2.7 Hz, 1H), 3.88 (s, 2H), 2.43 (d, J = 8.8 Hz, 2H), 2.13 (br s, 1H), 1.84 (dd, J = 11.9, 5.5 Hz, 1H), 1.54 (s, 3H), 1.53 (s, 3H), 1.37 (t, J = 12.2 Hz, 1H), 0.73 (d, J = 6.2 Hz, 3H).
2698	420.30	0.94	1H NMR (400 MHz, DMSO) δ 7.92 (dd, J = 4.8, 1.9 Hz, 1H), 7.80 (dd, J = 7.5, 1.9 Hz, 1H), 7.50 (d, J = 8.0 Hz, 1H), 6.52 (dd, J = 7.5, 4.8 Hz, 1H), 6.27 (s, 2H), 5.80 (d, J = 8.0 Hz, 1H), 3.74 (s, 3H), 2.90 (t, J = 10.8 Hz, 1H), 2.79 (dd, J = 10.7, 7.2 Hz, 1H), 2.14 - 2.00 (m, 1H), 1.76 (dd, J = 11.4, 5.7 Hz, 1H), 1.51 (d, J = 2.7 Hz, 6H), 1.31 (t, J = 12.0 Hz, 1H), 0.76 (d, J = 6.4 Hz, 3H).
2699	478.30	1.06	
2700	522.30	1.07	
2701	565.30	2.34	
2702	563.36	2.73	1H NMR (400 MHz, DMSO-d6) δ 12.46 (s, 1H), 7.78 (d, J = 8.0 Hz, 1H),

Cmpd No.	LC/MS M+1	LC/RT min	NMR
			7.63 (dd, J = 8.4, 7.3 Hz, 1H), 7.53 - 7.45 (m, 1H), 7.40 (dt, J = 10.0, 2.0 Hz, 1H), 7.26 (d, J = 8.0 Hz, 1H), 7.20 (d, J = 7.3 Hz, 1H), 6.89 (dt, J = 10.7, 2.3 Hz, 1H), 6.68 (d, J = 8.4 Hz, 1H), 6.49 (s, 2H), 3.83 (s, 2H), 2.76 - 2.54 (m, 3H), 2.22 (s, 1H), 1.88 (dd, J = 11.9, 5.7 Hz, 1H), 1.60 (d, J = 5.7 Hz, 6H), 1.42 (t, J = 12.1 Hz, 1H), 0.88 (d, J = 6.2 Hz, 3H).
2703	482.30	1.56	
2704	528.50	1.60	
2705	528.50	1.59	
2706	547.00	3.25	1H NMR (400 MHz, DMSO-d6) δ 12.57 (s, 1H), 8.07 - 7.95 (m, 2H), 7.78 - 7.70 (m, 2H), 7.70 - 7.62 (m, 2H), 7.50 - 7.45 (m, 1H), 7.40 (ddd, J = 10.2, 2.4, 1.3 Hz, 1H), 7.27 (d, J = 8.0 Hz, 1H), 6.89 (dt, J = 10.7, 2.3 Hz, 1H), 3.82 (s, 2H), 2.43 (t, J = 10.3 Hz, 1H), 2.36 (dd, J = 10.2, 7.1 Hz, 1H), 2.11 (ddt, J = 17.5, 11.9, 6.2 Hz, 1H), 1.83 (dd, J = 11.8, 5.6 Hz, 1H), 1.59 (s, 3H), 1.55 (s, 3H), 1.38 (t, J = 12.1 Hz, 1H), 0.66 (d, J = 6.3 Hz, 3H).
2707			
2708			1H NMR (300 MHz, DMSO) δ 8.55 (d, J = 2.2 Hz, 1H), 8.17 (d, J = 7.9 Hz, 1H), 8.02 (dd, J = 8.8, 2.5 Hz, 1H), 7.76 (dd, J = 8.4, 1.6 Hz, 3H), 7.72 - 7.65 (m, 2H), 7.64 - 7.57 (m, 2H), 7.56 (s, 2H), 7.24 (d, J = 6.9 Hz, 1H), 6.75 (dd, J = 18.5, 8.4 Hz, 2H), 5.26 (dt, J = 12.3, 6.2 Hz, 2H), 2.14 (s, 6H), 1.27 (d, J = 6.2 Hz, 6H).
2709	520.50	1.59	
2710	534.40	1.28	
2711	480.00	0.53	1H NMR (400 MHz, Methanol-d4) δ 8.84 - 8.78 (m, 1H), 8.74 - 8.67 (m, 2H), 8.62 - 8.57 (m, 1H), 8.04 (ddd, J = 7.4, 5.9, 1.3 Hz, 1H), 7.91 (d, J = 8.1 Hz, 1H), 7.45 - 7.39 (m, 2H), 3.34 (s, 1H), 3.18 - 3.06 (m, 2H), 2.51 - 2.39 (m, 1H), 2.07 - 1.99 (m, 1H), 1.72 (d, J = 14.8 Hz, 6H), 1.65 (d, J = 12.2 Hz,

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
			1H), 1.45 - 1.38 (m, 1H), 1.27 - 1.15 (m, 2H), 1.12 (d, J = 6.4 Hz, 3H).
2712	566.54	2.88	
2713	567.44	3.07	
2714	581.25	2.98	
2715	501.41	3.05	
2716	517.32	3.33	1H NMR (400 MHz, DMSO) δ 12.70 (s, 1H), 8.01 (t, J = 8.02, 1H), 7.74 - 7.67 (m, 2H), 7.13 - 7.10 (d, 1H), 6.67 - 6.64 (d, 1H), 4.34 - 4.20 (m, 2H), 2.43 - 2.40 (m, 2H), 2.14 (m, 1H), 1.85 - 1.69 (m, 2H), 1.62 (q, J = 6.48, 2H), 1.54 (s, 3H), 1.51 (s, 3H), 1.35 (t, J = 11.97, 1H), 1.26 (s, 9H), 0.92 - 0.90 (d, 6H), 0.77 - 0.75 (d, 3H)
2717	533.22	2.62	
2718	535.02	2.64	
2719	519.12	2.66	
2720	535.02	2.62	1H NMR (400 MHz, DMSO) δ 8.03 - 7.95 (m, 2H), 7.77 - 7.74 (d, 1H), 7.59 - 7.56 (d, 1H), 7.42 (t, J = 8.13, 1H), 7.34 - 7.31 (dd, 1H), 7.27 - 7.24 (d, 1H), 7.08 (m, 1H), 6.46 - 6.43 (dd, 1H), 2.89 (m, 2H), 2.21 - 2. 15 (m, 1H), 1.84 - 1.77 (m, 1H), 1.50 (s, 3H), 1.48 (s, 3H), 1.36 (t, J = 12.08, 1H), 0.90 - 0.88 (d, 3H)
2721	495.10	2.62	
2722	540.43	2.26	
2723	486.40	2.71	
2724	446.18	2.32	
2725	492.10	2.18	
2726	446.18	2.32	
2727	509.21	2.29	
2728	525.42	2.34	
2729	538.33	2.22	
2730	509.21	2.01	
2731	482.20	1.92	
2732	488.20	2.10	
2733	521.22	2.00	
2734	476.19	2.10	
2735	495.40	1.35	
2736	502.31	1.84	
2737	523.32	2.55	
2738	550.33	1.73	

Cmpd. No.	LC/MS M+1	LC/RT min	NMR
2739	539.23	2.70	
2740	513.41	2.63	
2741	443.18	2.20	
2742	434.17	2.31	
2743	581.25	2.23	
2744	528.12	2.51	
2745	474.09	2.26	
2746	460.29	2.33	
2747	472.29	2.31	

ASSAYS

PROTOCOL 1

Assays for Detecting and Measuring F508del Potentiation Properties of Compounds

Membrane potential optical methods for assaying F508del modulation properties of compounds

The assay utilizes fluorescent voltage sensing dyes to measure changes in membrane potential using a fluorescent plate reader (e.g., FLIPR III, Molecular Devices, Inc.) as a readout for increase in functional F508del in NIH 3T3 cells. The driving force for the response is the creation of a chloride ion gradient in conjunction with channel activation by a single liquid addition step after the cells have previously been treated with compounds and subsequently loaded with a voltage sensing dye.

Identification of Potentiator Compounds

To identify potentiators of F508del, a double-addition HTS assay format was developed. This HTS assay utilizes fluorescent voltage sensing dyes to measure changes in membrane potential on the FLIPR III as a measurement for increase in gating (conductance) of F508del in temperature-corrected F508del NIH 3T3 cells. The driving force for the response is a Cl^- ion gradient in conjunction with channel activation with forskolin in a single liquid addition step using a fluorescent plate reader such as FLIPR III after the cells have previously been treated with potentiator compounds (or DMSO vehicle control) and subsequently loaded with a redistribution dye.

Solutions

Bath Solution #1: (in mM) NaCl 160, KCl 4.5, $CaCl_2$ 2, $MgCl_2$ 1, HEPES 10, pH 7.4 with NaOH.

Chloride-free bath solution: Chloride salts in Bath Solution #1 (above) are substituted with gluconate salts.

Cell Culture

NIH3T3 mouse fibroblasts stably expressing F508del are used for optical measurements of membrane potential. The cells are maintained at 37 °C in 5% CO_2 and 90 % humidity in Dulbecco's modified Eagle's medium supplemented with 2 mM glutamine, 10 % fetal bovine serum, 1 X NEAA, β-ME, 1 X pen/strep, and 25 mM HEPES in 175 cm^2 culture flasks. For all optical assays, the cells were seeded at ~20,000/well in 384-well matrigel-coated plates and cultured for 2 hrs at 37 °C before culturing at 27 °C for 24 hrs. for the potentiator assay. For the correction assays, the cells are cultured at 27 °C or 37 °C with and without compounds for 16 – 24 hours.

Electrophysiological Assays for assaying F508del modulation properties of compounds.

Ussing Chamber Assay

Ussing chamber experiments were performed on polarized airway epithelial cells expressing F508del to further characterize the F508del augmenters or inducers identified in the optical assays. Non-CF and CF airway epithelia were isolated from bronchial tissue, cultured as previously described (Galietta, L.J.V., Lantero, S., Gazzolo, A., Sacco, O., Romano, L., Rossi, G.A., & Zegarra-Moran, O. (1998) *In Vitro Cell. Dev. Biol.* **34**, 478-481), and plated onto Costar® Snapwell™ filters that were precoated with NIH3T3-conditioned media. After four days the apical media was removed and the cells were grown at an air liquid interface for >14 days prior to use. This resulted in a monolayer of fully differentiated columnar cells that were ciliated, features that are characteristic of airway epithelia. Non-CF HBE were isolated from non-smokers that did not have any known lung disease. CF-HBE were isolated from patients homozygous for F508del.

HBE grown on Costar® Snapwell™ cell culture inserts were mounted in an Using chamber (Physiologic Instruments, Inc., San Diego, CA), and the transepithelial resistance and

short-circuit current in the presence of a basolateral to apical Cl^- gradient (I_{SC}) were measured using a voltage-clamp system (Department of Bioengineering, University of Iowa, IA). Briefly, HBE were examined under voltage-clamp recording conditions (V_{hold} = 0 mV) at 37 °C. The basolateral solution contained (in mM) 145 NaCl, 0.83 K_2HPO_4, 3.3 KH_2PO_4, 1.2 $MgCl_2$, 1.2 $CaCl_2$, 10 Glucose, 10 HEPES (pH adjusted to 7.35 with NaOH) and the apical solution contained (in mM) 145 NaGluconate, 1.2 $MgCl_2$, 1.2 $CaCl_2$, 10 glucose, 10 HEPES (pH adjusted to 7.35 with NaOH).

Identification of Potentiator Compounds

Typical protocol utilized a basolateral to apical membrane Cl^- concentration gradient. To set up this gradient, normal ringers was used on the basolateral membrane, whereas apical NaCl was replaced by equimolar sodium gluconate (titrated to pH 7.4 with NaOH) to give a large Cl^- concentration gradient across the epithelium. Forskolin (10 μM) and all test compounds were added to the apical side of the cell culture inserts. The efficacy of the putative F508del potentiators was compared to that of the known potentiator, genistein.

Patch-clamp Recordings

Total Cl^- current in F508del-NIH3T3 cells was monitored using the perforated-patch recording configuration as previously described (Rae, J., Cooper, K., Gates, P., & Watsky, M. (1991) *J. Neurosci. Methods* **37**, 15-26). Voltage-clamp recordings were performed at 22 °C using an Axopatch 200B patch-clamp amplifier (Axon Instruments Inc., Foster City, CA). The pipette solution contained (in mM) 150 *N*-methyl-D-glucamine (NMDG)-Cl, 2 $MgCl_2$, 2 $CaCl_2$, 10 EGTA, 10 HEPES, and 240 μg/mL amphotericin-B (pH adjusted to 7.35 with HCl). The extracellular medium contained (in mM) 150 NMDG-Cl, 2 $MgCl_2$, 2 $CaCl_2$, 10 HEPES (pH adjusted to 7.35 with HCl). Pulse generation, data acquisition, and analysis were performed using a PC equipped with a Digidata 1320 A/D interface in conjunction with Clampex 8 (Axon Instruments Inc.). To activate F508del, 10 μM forskolin and 20 μM genistein were added to the bath and the current-voltage relation was monitored every 30 sec.

Identification of Potentiator Compounds

The ability of F508del potentiators to increase the macroscopic F508del Cl^- current ($I_{F508del}$) in NIH3T3 cells stably expressing F508del was also investigated using perforated-patch-

recording techniques. The potentiators identified from the optical assays evoked a dose-dependent increase in $I\Delta_{F508}$ with similar potency and efficacy observed in the optical assays. In all cells examined, the reversal potential before and during potentiator application was around -30 mV, which is the calculated E_{Cl} (-28 mV).

Cell Culture

NIH3T3 mouse fibroblasts stably expressing F508del are used for whole-cell recordings. The cells are maintained at 37 °C in 5% CO_2 and 90 % humidity in Dulbecco's modified Eagle's medium supplemented with 2 mM glutamine, 10 % fetal bovine serum, 1 X NEAA, β-ME, 1 X pen/strep, and 25 mM HEPES in 175 cm^2 culture flasks. For whole-cell recordings, 2,500 - 5,000 cells were seeded on poly-L-lysine-coated glass coverslips and cultured for 24 - 48 hrs at 27 °C before use to test the activity of potentiators; and incubated with or without the correction compound at 37 °C for measuring the activity of correctors.

Single-channel recordings

Gating activity of wt-CFTR and temperature-corrected F508del expressed in NIH3T3 cells was observed using excised inside-out membrane patch recordings as previously described (Dalemans, W., Barbry, P., Champigny, G., Jallat, S., Dott, K., Dreyer, D., Crystal, R.G., Pavirani, A., Lecocq, J-P., Lazdunski, M. (1991) *Nature* 354, 526 – 528) using an Axopatch 200B patch-clamp amplifier (Axon Instruments Inc.). The pipette contained (in mM): 150 NMDG, 150 aspartic acid, 5 $CaCl_2$, 2 $MgCl_2$, and 10 HEPES (pH adjusted to 7.35 with Tris base). The bath contained (in mM): 150 NMDG-Cl, 2 $MgCl_2$, 5 EGTA, 10 TES, and 14 Tris base (pH adjusted to 7.35 with HCl). After excision, both wt- and F508del were activated by adding 1 mM Mg-ATP, 75 nM of the catalytic subunit of cAMP-dependent protein kinase (PKA; Promega Corp. Madison, WI), and 10 mM NaF to inhibit protein phosphatases, which prevented current rundown. The pipette potential was maintained at 80 mV. Channel activity was analyzed from membrane patches containing ≤ 2 active channels. The maximum number of simultaneous openings determined the number of active channels during the course of an experiment. To determine the single-channel current amplitude, the data recorded from 120 sec of F508del activity was filtered "off-line" at 100 Hz and then used to construct all-point amplitude histograms that were fitted with multigaussian functions using Bio-Patch Analysis software (Bio-Logic Comp. France). The total microscopic current and open probability (P_o)

were determined from 120 sec of channel activity. The P_o was determined using the Bio-Patch software or from the relationship $P_o = I/i(N)$, where I = mean current, i = single-channel current amplitude, and N = number of active channels in patch.

Cell Culture

NIH3T3 mouse fibroblasts stably expressing F508del are used for excised-membrane patch-clamp recordings. The cells are maintained at 37 °C in 5% CO_2 and 90 % humidity in Dulbecco's modified Eagle's medium supplemented with 2 mM glutamine, 10 % fetal bovine serum, 1 X NEAA, β-ME, 1 X pen/strep, and 25 mM HEPES in 175 cm^2 culture flasks. For single channel recordings, 2,500 - 5,000 cells were seeded on poly-L-lysine-coated glass coverslips and cultured for 24 - 48 hrs at 27 °C before use.

PROTOCOL 2

Assays for Detecting and Measuring F508del Correction Properties of Compounds

Membrane potential optical methods for assaying F508del modulation properties of compounds.

The optical membrane potential assay utilized voltage-sensitive FRET sensors described by Gonzalez and Tsien (*See* Gonzalez, J. E. and R. Y. Tsien (1995) "Voltage sensing by fluorescence resonance energy transfer in single cells" Biophys J 69(4): 1272-80, and Gonzalez, J. E. and R. Y. Tsien (1997) "Improved indicators of cell membrane potential that use fluorescence resonance energy transfer" Chem Biol 4(4): 269-77) in combination with instrumentation for measuring fluorescence changes such as the Voltage/Ion Probe Reader (VIPR) (See, Gonzalez, J. E., K. Oades, et al. (1999) "Cell-based assays and instrumentation for screening ion-channel targets" Drug Discov Today 4(9): 431-439).

These voltage sensitive assays are based on the change in fluorescence resonant energy transfer (FRET) between the membrane-soluble, voltage-sensitive dye, $DiSBAC_2(3)$, and a fluorescent phospholipid, CC2-DMPE, which is attached to the outer leaflet of the plasma membrane and acts as a FRET donor. Changes in membrane potential (V_m) cause the negatively charged $DiSBAC_2(3)$ to redistribute across the plasma membrane and the amount of energy transfer from CC2-DMPE changes accordingly. The changes in fluorescence emission were

monitored using VIPR™ II, which is an integrated liquid handler and fluorescent detector designed to conduct cell-based screens in 96- or 384-well microtiter plates.

Identification of Correction Compounds

To identify small molecules that correct the trafficking defect associated with F508del; a single-addition HTS assay format was developed. The cells were incubated in serum-free medium for 16 hrs at 37 °C in the presence or absence (negative control) of test compound. As a positive control, cells plated in 384-well plates were incubated for 16 hrs at 27 °C to "temperature-correct" F508del. The cells were subsequently rinsed 3X with Krebs Ringers solution and loaded with the voltage-sensitive dyes. To activate F508del, 10 μM forskolin and the CFTR potentiator, genistein (20 μM), were added along with Cl^--free medium to each well. The addition of Cl^--free medium promoted Cl^- efflux in response to F508del activation and the resulting membrane depolarization was optically monitored using the FRET-based voltage-sensor dyes.

Identification of Potentiator Compounds

To identify potentiators of F508del, a double-addition HTS assay format was developed. During the first addition, a Cl^--free medium with or without test compound was added to each well. After 22 sec, a second addition of Cl^--free medium containing 2 - 10 μM forskolin was added to activate F508del. The extracellular Cl^- concentration following both additions was 28 mM, which promoted Cl^- efflux in response to F508del activation and the resulting membrane depolarization was optically monitored using the FRET-based voltage-sensor dyes.

Solutions

Bath Solution #1:	(in mM) NaCl 160, KCl 4.5, $CaCl_2$ 2, $MgCl_2$ 1, HEPES 10, pH 7.4 with NaOH.
Chloride-free bath solution:	Chloride salts in Bath Solution #1 (above) are substituted with gluconate salts.
CC2-DMPE:	Prepared as a 10 mM stock solution in DMSO and stored at -20°C.

$DiSBAC_2(3)$: Prepared as a 10 mM stock in DMSO and stored at -20°C.

Cell Culture

NIH3T3 mouse fibroblasts stably expressing F508del are used for optical measurements of membrane potential. The cells are maintained at 37 °C in 5% CO_2 and 90 % humidity in Dulbecco's modified Eagle's medium supplemented with 2 mM glutamine, 10 % fetal bovine serum, 1 X NEAA, β-ME, 1 X pen/strep, and 25 mM HEPES in 175 cm^2 culture flasks. For all optical assays, the cells were seeded at 30,000/well in 384-well matrigel-coated plates and cultured for 2 hrs at 37 °C before culturing at 27 °C for 24 hrs for the potentiator assay. For the correction assays, the cells are cultured at 27 °C or 37 °C with and without compounds for 16 - 24 hours.

Electrophysiological Assays for assaying F508del modulation properties of compounds

Ussing Chamber Assay

Using chamber experiments were performed on polarized epithelial cells expressing F508del to further characterize the F508del augmenters or inducers identified in the optical assays. FRT ^{F508del} epithelial cells grown on Costar Snapwell cell culture inserts were mounted in an Ussing chamber (Physiologic Instruments, Inc., San Diego, CA), and the monolayers were continuously short-circuited using a Voltage-clamp System (Department of Bioengineering, University of Iowa, IA, and, Physiologic Instruments, Inc., San Diego, CA). Transepithelial resistance was measured by applying a 2-mV pulse. Under these conditions, the FRT epithelia demonstrated resistances of 4 KΩ/ cm^2 or more. The solutions were maintained at 27 °C and bubbled with air. The electrode offset potential and fluid resistance were corrected using a cell-free insert. Under these conditions, the current reflects the flow of Cl^- through F508del expressed in the apical membrane. The I_{SC} was digitally acquired using an MP100A-CE interface and AcqKnowledge software (v3.2.6; BIOPAC Systems, Santa Barbara, CA).

Identification of Correction Compounds

Typical protocol utilized a basolateral to apical membrane Cl^- concentration gradient. To set up this gradient, normal ringer was used on the basolateral membrane, whereas apical

NaCl was replaced by equimolar sodium gluconate (titrated to pH 7.4 with NaOH) to give a large Cl^- concentration gradient across the epithelium. All experiments were performed with intact monolayers. To fully activate F508del, forskolin (10 μM) and the PDE inhibitor, IBMX (100 μM), were applied followed by the addition of the CFTR potentiator, genistein (50 μM).

As observed in other cell types, incubation at low temperatures of FRT cells stably expressing F508del increases the functional density of CFTR in the plasma membrane. To determine the activity of correction compounds, the cells were incubated with 10 μM of the test compound for 24 hours at 37°C and were subsequently washed 3X prior to recording. The cAMP- and genistein-mediated I_{SC} in compound-treated cells was normalized to the 27°C and 37°C controls and expressed as percentage activity. Preincubation of the cells with the correction compound significantly increased the cAMP- and genistein-mediated I_{SC} compared to the 37°C controls.

Identification of Potentiator Compounds

Typical protocol utilized a basolateral to apical membrane Cl^- concentration gradient. To set up this gradient, normal ringers was used on the basolateral membrane and was permeabilized with nystatin (360 μg/ml), whereas apical NaCl was replaced by equimolar sodium gluconate (titrated to pH 7.4 with NaOH) to give a large Cl^- concentration gradient across the epithelium. All experiments were performed 30 min after nystatin permeabilization. Forskolin (10 μM) and all test compounds were added to both sides of the cell culture inserts. The efficacy of the putative F508del potentiators was compared to that of the known potentiator, genistein.

Solutions

Basolateral solution (in mM):	NaCl (135), $CaCl_2$ (1.2), $MgCl_2$ (1.2), K_2HPO_4 (2.4), $KHPO_4$ (0.6), *N*-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES) (10), and dextrose (10). The solution was titrated to pH 7.4 with NaOH.
Apical solution (in mM):	Same as basolateral solution with NaCl replaced with Na Gluconate (135).

Cell Culture

Fisher rat epithelial (FRT) cells expressing F508del (FRT ^{F508del}) were used for Ussing chamber experiments for the putative F508del augmenters or inducers identified from our optical assays. The cells were cultured on Costar Snapwell cell culture inserts and cultured for five days at 37 °C and 5% CO_2 in Coon's modified Ham's F-12 medium supplemented with 5% fetal calf serum, 100 U/ml penicillin, and 100 µg/ml streptomycin. Prior to use for characterizing the potentiator activity of compounds, the cells were incubated at 27 °C for 16 - 48 hrs to correct for the F508del. To determine the activity of corrections compounds, the cells were incubated at 27 °C or 37 °C with and without the compounds for 24 hours.

Whole-cell recordings

The macroscopic F508del current ($I_{F508del}$) in temperature- and test compound-corrected NIH3T3 cells stably expressing F508del were monitored using the perforated-patch, whole-cell recording. Briefly, voltage-clamp recordings of $I_{F508del}$ were performed at room temperature using an Axopatch 200B patch-clamp amplifier (Axon Instruments Inc., Foster City, CA). All recordings were acquired at a sampling frequency of 10 kHz and low-pass filtered at 1 kHz. Pipettes had a resistance of 5 – 6 MΩ when filled with the intracellular solution. Under these recording conditions, the calculated reversal potential for Cl^- (E_{Cl}) at room temperature was -28 mV. All recordings had a seal resistance > 20 GΩ and a series resistance < 15 MΩ. Pulse generation, data acquisition, and analysis were performed using a PC equipped with a Digidata 1320 A/D interface in conjunction with Clampex 8 (Axon Instruments Inc.). The bath contained < 250 µl of saline and was continuously perifused at a rate of 2 ml/min using a gravity-driven perfusion system.

Identification of Correction Compounds

To determine the activity of correction compounds for increasing the density of functional F508del in the plasma membrane, we used the above-described perforated-patch-recording techniques to measure the current density following 24-hr treatment with the correction compounds. To fully activate F508del, 10 µM forskolin and 20 µM genistein were added to the cells. Under our recording conditions, the current density following 24-hr incubation at 27°C was higher than that observed following 24-hr incubation at 37 °C. These

results are consistent with the known effects of low-temperature incubation on the density of F508del in the plasma membrane. To determine the effects of correction compounds on CFTR current density, the cells were incubated with 10 μM of the test compound for 24 hours at 37°C and the current density was compared to the 27°C and 37°C controls (% activity). Prior to recording, the cells were washed 3X with extracellular recording medium to remove any remaining test compound. Preincubation with 10 μM of correction compounds significantly increased the cAMP- and genistein-dependent current compared to the 37°C controls.

Identification of Potentiator Compounds

The ability of F508del potentiators to increase the macroscopic F508del Cl^- current ($I_{F508del}$) in NIH3T3 cells stably expressing F508del was also investigated using perforated-patch-recording techniques. The potentiators identified from the optical assays evoked a dose-dependent increase in $I_{F508del}$ with similar potency and efficacy observed in the optical assays. In all cells examined, the reversal potential before and during potentiator application was around -30 mV, which is the calculated E_{Cl} (-28 mV).

Solutions

Intracellular solution (in mM):	Cs-aspartate (90), CsCl (50), $MgCl_2$ (1), HEPES (10), and 240 μg/ml amphotericin-B (pH adjusted to 7.35 with CsOH).
Extracellular solution (in mM):	*N*-methyl-D-glucamine (NMDG)-Cl (150), $MgCl_2$ (2), $CaCl_2$ (2), HEPES (10) (pH adjusted to 7.35 with HCl).

Cell Culture

NIH3T3 mouse fibroblasts stably expressing F508del are used for whole-cell recordings. The cells are maintained at 37 °C in 5% CO_2 and 90 % humidity in Dulbecco's modified Eagle's medium supplemented with 2 mM glutamine, 10 % fetal bovine serum, 1 X NEAA, β-ME, 1 X pen/strep, and 25 mM HEPES in 175 cm^2 culture flasks. For whole-cell recordings, 2,500 - 5,000 cells were seeded on poly-L-lysine-coated glass coverslips and cultured for 24 - 48 hrs at 27 °C before use to test the activity of potentiators; and incubated with or without the correction compound at 37 °C for measuring the activity of correctors.

Single-channel recordings

The single-channel activities of temperature-corrected F508del stably expressed in NIH3T3 cells and activities of potentiator compounds were observed using excised inside-out membrane patch. Briefly, voltage-clamp recordings of single-channel activity were performed at room temperature with an Axopatch 200B patch-clamp amplifier (Axon Instruments Inc.). All recordings were acquired at a sampling frequency of 10 kHz and low-pass filtered at 400 Hz. Patch pipettes were fabricated from Corning Kovar Sealing #7052 glass (World Precision Instruments, Inc., Sarasota, FL) and had a resistance of 5 - 8 MΩ when filled with the extracellular solution. The F508del was activated after excision, by adding 1 mM Mg-ATP, and 75 nM of the cAMP-dependent protein kinase, catalytic subunit (PKA; Promega Corp. Madison, WI). After channel activity stabilized, the patch was perifused using a gravity-driven microperfusion system. The inflow was placed adjacent to the patch, resulting in complete solution exchange within 1 - 2 sec. To maintain F508del activity during the rapid perifusion, the nonspecific phosphatase inhibitor F^- (10 mM NaF) was added to the bath solution. Under these recording conditions, channel activity remained constant throughout the duration of the patch recording (up to 60 min). Currents produced by positive charge moving from the intra- to extracellular solutions (anions moving in the opposite direction) are shown as positive currents. The pipette potential (V_p) was maintained at 80 mV.

Channel activity was analyzed from membrane patches containing ≤ 2 active channels. The maximum number of simultaneous openings determined the number of active channels during the course of an experiment. To determine the single-channel current amplitude, the data recorded from 120 sec of F508del activity was filtered "off-line" at 100 Hz and then used to construct all-point amplitude histograms that were fitted with multigaussian functions using Bio-Patch Analysis software (Bio-Logic Comp. France). The total microscopic current and open probability (P_o) were determined from 120 sec of channel activity. The P_o was determined using the Bio-Patch software or from the relationship $P_o = I/i(N)$, where I = mean current, i = single-channel current amplitude, and N = number of active channels in patch.

Solutions

Extracellular solution (in mM): NMDG (150), aspartic acid (150), $CaCl_2$ (5), $MgCl_2$ (2), and HEPES (10) (pH adjusted to 7.35 with Tris base).

Intracellular solution (in mM): NMDG-Cl (150), $MgCl_2$ (2), EGTA (5), TES (10), and Tris base (14) (pH adjusted to 7.35 with HCl).

Cell Culture

NIH3T3 mouse fibroblasts stably expressing F508del are used for excised-membrane patch-clamp recordings. The cells are maintained at 37 °C in 5% CO_2 and 90 % humidity in Dulbecco's modified Eagle's medium supplemented with 2 mM glutamine, 10 % fetal bovine serum, 1 X NEAA, β-ME, 1 X pen/strep, and 25 mM HEPES in 175 cm^2 culture flasks. For single channel recordings, 2,500 - 5,000 cells were seeded on poly-L-lysine-coated glass coverslips and cultured for 24 - 48 hrs at 27 °C before use.

The compounds of formula I are useful as modulators of CFTR activity. Table 3 below illustrates the EC30 and relative efficacy of the compounds of Table 1. In Table 3 below, the following meanings apply. EC30: "+++" means < 3 uM; "++" means between 3 uM and 10 uM; "+" means greater than 10 uM.

Table 3.

Cmpd. No.	EC30 (μM)
1	+++
2	+++
3	+++
4	+
5	++
6	+++
7	+++
8	+++
9	+++
10	+++
11	+++
12	+++
13	++

Cmpd. No.	EC30 (μM)
14	+++
15	+++
16	+++
17	+++
18	+++
19	+++
20	+++
21	+++
22	+++
23	++
24	++
25	++
26	++

Cmpd. No.	EC30 (μM)
27	+++
28	+++
29	+++
30	+++
31	++
32	++
33	+++
34	+++
35	+++
36	+++
37	+++
38	+++
39	++
40	+++
41	+++
42	++
43	+++
44	++
45	+++
46	++
47	+++
48	++
49	+++
50	+++
51	+++
52	+++
53	+++
54	++
55	++
56	+++
57	+++
58	++
59	+++
60	+++
61	+++
62	+++
63	+++
64	++
65	++
66	+++
67	++

Cmpd. No.	EC30 (μM)
68	+++
69	+++
70	+++
71	+++
72	++
73	+++
74	+++
75	++
76	+++
77	+++
78	+++
79	++
80	+++
81	+++
82	+++
83	+++
84	+++
85	++
86	++
87	+++
88	+++
89	+++
90	+++
91	++
92	+++
93	+++
94	+++
95	+++
96	+++
97	+++
98	+++
99	+++
100	+++
101	+++
102	+++
103	+++
104	+++
105	+++
106	+++
107	+++
108	+++

Cmpd. No.	EC30 (μM)
109	+++
110	+++
111	+++
112	+++
113	++
114	+++
115	+++
116	++
117	+++
118	++
119	+++
120	+++
121	++
122	++
123	+++
124	++
125	++
126	+++
127	+++
128	+++
129	++
130	+++
131	+++
132	+++
133	+++
134	+++
135	+++
136	+++
137	+++
138	++
139	+++
140	+++
141	+++
142	+++
143	+++
144	+++
145	+++
146	+++
147	+++
148	+++
149	+++

Cmpd. No.	EC30 (μM)
150	+++
151	+++
152	+++
153	+++
154	+++
155	++
156	+++
157	++
158	+++
159	+++
160	+++
161	+++
162	+++
163	+++
164	+++
165	+++
166	+++
167	++
168	+++
169	+++
170	+++
171	+++
172	+++
173	++
174	++
175	++
176	+++
177	++
178	+++
179	+++
180	+++
181	+++
182	+++
183	+++
184	++
185	+++
186	+++
187	++
188	+++
189	+++
190	+++

Cmpd. No.	EC30 (μM)
191	+++
192	+++
193	+++
194	+++
195	+++
196	+++
197	+++
198	+++
199	+++
200	++
201	+++
202	+++
203	+++
204	+++
205	++
206	+++
207	+++
208	+++
209	+++
210	+++
211	++
212	+++
213	+++
214	+++
215	+++
216	+++
217	+++
218	++
219	+++
220	+++
221	+++
222	+++
223	+++
224	+++
225	+++
226	+++
227	++
228	+++
229	+++
230	++
231	+++

Cmpd. No.	EC30 (μM)
232	+++
233	+++
234	+++
235	+++
236	++
237	++
238	+++
239	+++
240	++
241	+++
242	+++
243	+++
244	+++
245	++
246	++
247	+++
248	+++
249	+++
250	+++
251	+++
252	+++
253	+++
254	++
255	+++
256	+++
257	+++
258	++
259	+++
260	++
261	+++
262	+++
263	+++
264	+++
265	+++
266	+++
267	+++
268	+++
269	+++
270	+++
271	+++
272	++

Cmpd No.	EC30 (μM)
273	+++
274	+++
275	+++
276	+++
277	+++
278	++
279	+++
280	+++
281	+++
282	++
283	+++
284	+++
285	+++
286	+++
287	+++
288	+++
289	+++
290	+++
291	++
292	+++
293	+++
294	+++
295	++
296	+++
297	++
298	++
299	+++
300	+++
301	++
302	++
303	+++
304	+++
305	+++
306	+++
307	+++
308	+++
309	+++
310	+++
311	+++
312	+++
313	+++

Cmpd No.	EC30 (μM)
314	+++
315	+++
316	+++
317	+++
318	+++
319	++
320	+++
321	+++
322	+++
323	+++
324	++
325	+++
326	+++
327	+++
328	+++
329	+++
330	+++
331	+++
332	+++
333	+++
334	++
335	+++
336	+++
337	+++
338	+++
339	+++
340	+++
341	+++
342	+++
343	+++
344	+++
345	+++
346	+++
347	+++
348	+++
349	++
350	+++
351	+++
352	+++
353	+++
354	+++

Cmpd. No.	EC30 (μM)
355	+++
356	+++
357	++
358	+++
359	+++
360	+++
361	+++
362	+++
363	+++
364	+++
365	+++
366	+++
367	+++
368	+++
369	+++
370	+++
371	+++
372	++
373	++
374	+++
375	+++
376	+++
377	+++
378	+++
379	+++
380	+++
381	++
382	+++
383	+++
384	+++
385	+++
386	+++
387	+++
388	++
389	+++
390	+++
391	+
392	++
393	++
394	+++
395	+++

Cmpd. No.	EC30 (μM)
396	+++
397	+++
398	+++
399	+++
400	++
401	+++
402	+++
403	++
404	+++
405	+++
406	+++
407	++
408	+++
409	+
410	+++
411	+++
412	+++
413	+++
414	+++
415	+++
416	+++
417	++
418	+++
419	+++
420	+++
421	+++
422	+++
423	+++
424	+++
425	+++
426	+++
427	+++
428	+++
429	+++
430	+++
431	++
432	+++
433	+++
434	+++
435	+++
436	+++

Cmpd. No.	EC30 (μM)
437	+++
438	+++
439	+++
440	+++
441	+++
442	+++
443	+
444	++
445	+
446	+++
447	+++
448	+++
449	+
450	+++
451	++
452	+++
453	+++
454	+++
455	+++
456	+++
457	+++
458	+++
459	+++
460	++
461	++
462	+++
463	++
464	+++
465	++
466	+++
467	++
468	+++
469	++
470	+++
471	+++
472	+
473	+++
474	+++
475	+++
476	++
477	+++

Cmpd. No.	EC30 (μM)
478	++
479	+++
480	+++
481	+++
482	+++
483	+++
484	
485	+++
486	++
487	++
488	+++
489	+++
490	+++
491	+++
492	+++
493	+++
494	+++
495	++
496	+++
497	+++
498	+++
499	++
500	+++
501	+++
502	+++
503	+++
504	+++
505	++
506	+++
507	+++
508	++
509	+++
510	+++
511	+++
512	++
513	++
514	+
515	+++
516	+++
517	++
518	+++

Cmpd No.	EC30 (μM)
519	++
520	+++
521	++
522	+++
523	+++
524	++
525	++
526	+++
527	+++
528	+++
529	++
530	+++
531	+++
532	++
533	+++
534	++
535	+++
536	+++
537	+++
538	+++
539	+
540	+
541	+++
542	+++
543	++
544	+++
545	++
546	++
547	+++
548	+++
549	+++
550	++
551	+++
552	++
553	+++
554	+++
555	+++
556	+
557	+++
558	+++
559	+++

Cmpd No.	EC30 (μM)
560	+++
561	+++
562	+++
563	+++
564	++
565	+
566	+
567	++
568	+++
569	+++
570	+++
571	++
572	+++
573	+++
574	++
575	+++
576	+++
577	++
578	+++
579	+++
580	+++
581	+++
582	++
583	+++
584	++
585	+++
586	+++
587	+++
588	++
589	+++
590	+++
591	+++
592	+++
593	+++
594	++
595	+++
596	++
597	+++
598	++
599	+++
600	+++

Cmpd. No.	EC30 (μM)
601	+++
602	+++
603	+++
604	+++
605	+++
606	++
607	+++
608	++
609	+++
610	+++
611	+++
612	+++
613	+++
614	+++
615	+++
616	+++
617	+++
618	++
619	+++
620	+++
621	+++
622	+++
623	++
624	++
625	+++
626	+++
627	+
628	+++
629	+++
630	+++
631	+++
632	++
633	+++
634	+++
635	+
636	++
637	++
638	++
639	+++
640	++
641	+++

Cmpd. No.	EC30 (μM)
642	+++
643	++
644	+++
645	++
646	++
647	+++
648	+++
649	++
650	++
651	+++
652	++
653	+
654	+++
655	+++
656	+++
657	+++
658	+++
659	++
660	+
661	+++
662	+
663	+++
664	+++
665	+++
666	++
667	+++
668	+++
669	+++
670	++
671	+++
672	+++
673	+++
674	+++
675	+++
676	+++
677	++
678	+++
679	+++
680	+++
681	+++
682	+++

Cmpd. No.	EC30 (μM)
683	+++
684	+++
685	+++
686	+++
687	++
688	+++
689	+++
690	++
691	+++
692	+++
693	+++
694	+++
695	+++
696	+++
697	+++
698	
699	+++
700	+++
701	+++
702	++
703	+++
704	++
705	+++
706	+++
707	+
708	+++
709	+++
710	+++
711	+++
712	+++
713	++
714	+++
715	+++
716	+++
717	+++
718	+++
719	+++
720	+++
721	++
722	+
723	++

Cmpd. No.	EC30 (μM)
724	+++
725	+++
726	++
727	+
728	++
729	++
730	+++
731	+++
732	+++
733	+++
734	+++
735	+++
736	+++
737	+++
738	+++
739	++
740	++
741	+++
742	+++
743	+++
744	+++
745	++
746	+++
747	+++
748	++
749	+++
750	+++
751	+++
752	++
753	++
754	+++
755	++
756	+++
757	+++
758	++
759	+++
760	++
761	+++
762	++
763	+++
764	+++

Cmpd. No.	EC30 (μM)
765	++
766	++
767	+++
768	++
769	+++
770	++
771	+++
772	+++
773	+++
774	+++
775	+++
776	+++
777	+++
778	+++
779	+++
780	+++
781	+
782	+++
783	+++
784	+++
785	+++
786	+
787	+++
788	+++
789	+++
790	+++
791	++
792	++
793	+++
794	+++
795	+++
796	+++
797	+
798	+++
799	+
800	++
801	+
802	+++
803	+++
804	+++
805	+++

Cmpd. No.	EC30 (μM)
806	++
807	+++
808	++
809	++
810	++
811	+++
812	++
813	+++
814	+++
815	++
816	++
817	+++
818	+++
819	+
820	+
821	++
822	+++
823	+
824	++
825	++
826	+++
827	+++
828	+++
829	+++
830	+++
831	+++
832	+++
833	+++
834	+++
835	++
836	++
837	+++
838	++
839	++
840	+++
841	+++
842	+++
843	++
844	++
845	+++
846	+++

Cmpd No.	EC30 (μM)
847	+++
848	+++
849	+
850	+++
851	+++
852	++
853	+++
854	++
855	+
856	++
857	+++
858	++
859	+++
860	++
861	++
862	+++
863	++
864	+++
865	+++
866	+++
867	+++
868	+++
869	+++
870	++
871	+++
872	++
873	+++
874	+++
875	+++
876	+++
877	+++
878	++
879	+
880	++
881	+++
882	+++
883	++
884	++
885	++
886	+++
887	+++

Cmpd No.	EC30 (μM)
888	++
889	++
890	+++
891	+++
892	++
893	+++
894	+++
895	+++
896	+++
897	+++
898	+++
899	+++
900	+++
901	++
902	+++
903	++
904	+++
905	+++
906	+++
907	+++
908	+
909	+++
910	+++
911	+++
912	+++
913	++
914	+++
915	+++
916	+++
917	++
918	+++
919	+++
920	+++
921	++
922	+++
923	+++
924	+++
925	+++
926	+++
927	++
928	+++

Cmpd No.	EC30 (μM)
929	++
930	++
931	+++
932	+
933	+++
934	+++
935	++
936	++
937	+++
938	++
939	+++
940	+++
941	+++
942	+++
943	+
944	+++
945	+++
946	++
947	+++
948	+++
949	+++
950	+++
951	+++
952	+++
953	+++
954	+++
955	++
956	+++
957	+++
958	+++
959	+++
960	++
961	+++
962	++
963	++
964	+++
965	+++
966	+
967	+++
968	+++
969	+++

Cmpd No.	EC30 (μM)
970	
971	+++
972	+++
973	+++
974	+++
975	++
976	++
977	++
978	+++
979	+++
980	++
981	++
982	++
983	++
984	+++
985	+++
986	++
987	+++
988	+++
989	+++
990	+++
991	+++
992	+++
993	+++
994	+++
995	+++
996	+++
997	++
998	++
999	++
1000	+++
1001	++
1002	+++
1003	+
1004	+++
1005	+++
1006	+++
1007	+++
1008	+++
1009	+++
1010	+++

Cmpd. No.	EC30 (μM)
1011	+++
1012	+++
1013	+++
1014	+++
1015	+++
1016	+++
1017	++
1018	++
1019	+++
1020	+++
1021	+++
1022	++
1023	+++
1024	++
1025	+++
1026	++
1027	++
1028	++
1029	++
1030	+++
1031	+++
1032	+++
1033	+++
1034	+++
1035	+++
1036	+
1037	++
1038	++
1039	+++
1040	+++
1041	+++
1042	+++
1043	+++
1044	+++
1045	++
1046	++
1047	+++
1048	+++
1049	+++
1050	++
1051	+++

Cmpd. No.	EC30 (μM)
1052	+++
1053	++
1054	+++
1055	++
1056	+++
1057	+
1058	+++
1059	+++
1060	+++
1061	+++
1062	+++
1063	+++
1064	+++
1065	+++
1066	++
1067	+++
1068	+++
1069	+++
1070	+++
1071	+++
1072	++
1073	+
1074	++
1075	+++
1076	+++
1077	+++
1078	++
1079	+++
1080	+++
1081	++
1082	+
1083	+++
1084	++
1085	+++
1086	++
1087	+++
1088	+++
1089	+++
1090	+++
1091	+++
1092	++

Cmpd. No.	EC30 (μM)
1093	++
1094	+++
1095	++
1096	++
1097	+++
1098	+++
1099	+++
1100	+++
1101	+++
1102	+++
1103	+++
1104	+++
1105	+
1106	+
1107	+++
1108	+++
1109	++
1110	+++
1111	+++
1112	++
1113	+++
1114	
1115	+
1116	++
1117	+++
1118	+++
1119	++
1120	+++
1121	+++
1122	++
1123	+++
1124	+++
1125	++
1126	+
1127	+++
1128	++
1129	++
1130	++
1131	+++
1132	+++
1133	+++

Cmpd. No.	EC30 (μM)
1134	+++
1135	+++
1136	+++
1137	++
1138	+++
1139	+++
1140	+++
1141	++
1142	++
1143	++
1144	+++
1145	+++
1146	++
1147	++
1148	+++
1149	+++
1150	+++
1151	++
1152	+
1153	+++
1154	++
1155	++
1156	+++
1157	+++
1158	+++
1159	+++
1160	+++
1161	+++
1162	+++
1163	+++
1164	++
1165	++
1166	+++
1167	+++
1168	+++
1169	++
1170	+++
1171	+++
1172	++
1173	+++
1174	+++

Cmpd. No.	EC30 (μM)
1175	+++
1176	+++
1177	++
1178	+++
1179	++
1180	+++
1181	+++
1182	++
1183	+
1184	+++
1185	+++
1186	+++
1187	+++
1188	+++
1189	+++
1190	+++
1191	++
1192	+++
1193	+++
1194	+++
1195	+
1196	++
1197	+++
1198	+++
1199	++
1200	++
1201	+++
1202	+++
1203	+
1204	+
1205	+++
1206	++
1207	+++
1208	+++
1209	+++
1210	+++
1211	++
1212	+++
1213	++
1214	++
1215	++

Cmpd. No.	EC30 (μM)
1216	+++
1217	+
1218	+++
1219	++
1220	+++
1221	+++
1222	+++
1223	+++
1224	++
1225	+++
1226	+++
1227	+++
1228	++
1229	+++
1230	++
1231	+++
1232	+++
1233	++
1234	+++
1235	+++
1236	++
1237	++
1238	+++
1239	+++
1240	+++
1241	+++
1242	+++
1243	+++
1244	+++
1245	+
1246	+++
1247	+++
1248	+++
1249	+++
1250	+++
1251	++
1252	+++
1253	++
1254	+++
1255	+++
1256	++

Cmpd. No.	EC30 (μM)
1257	++
1258	+++
1259	+++
1260	+++
1261	+++
1262	++
1263	+++
1264	+++
1265	++
1266	+++
1267	+++
1268	+++
1269	+++
1270	+++
1271	+++
1272	+++
1273	+++
1274	+++
1275	+++
1276	++
1277	+++
1278	+
1279	+++
1280	+++
1281	++
1282	+++
1283	+++
1284	+++
1285	+++
1286	+++
1287	+++
1288	++
1289	+++
1290	+++
1291	++
1292	+
1293	++
1294	+++
1295	+++
1296	+
1297	+++

Cmpd. No.	EC30 (μM)
1298	++
1299	++
1300	+++
1301	++
1302	++
1303	+++
1304	+++
1305	+++
1306	+++
1307	++
1308	+++
1309	+++
1310	+++
1311	+++
1312	++
1313	+++
1314	+++
1315	+++
1316	+++
1317	+++
1318	+++
1319	+++
1320	+++
1321	++
1322	++
1323	+
1324	+++
1325	+++
1326	++
1327	+++
1328	+++
1329	++
1330	+++
1331	+
1332	+++
1333	+++
1334	+++
1335	+++
1336	+++
1337	+++
1338	++

Cmpd. No.	EC30 (μM)
1339	+++
1340	+++
1341	+++
1342	+++
1343	+++
1344	++
1345	+++
1346	+++
1347	+++
1348	+++
1349	+++
1350	+
1351	++
1352	+++
1353	+
1354	+++
1355	++
1356	+++
1357	+++
1358	+++
1359	
1360	+++
1361	+++
1362	+++
1363	+++
1364	+++
1365	+++
1366	+++
1367	+++
1368	+++
1369	+++
1370	+++
1371	+++
1372	+++
1373	+++
1374	+++
1375	++
1376	+++
1377	+++
1378	+++
1379	+++

Cmpd. No.	EC30 (μM)
1380	+++
1381	++
1382	+
1383	+++
1384	+++
1385	+++
1386	++
1387	+++
1388	+++
1389	+++
1390	+++
1391	+++
1392	+++
1393	+++
1394	+++
1395	+++
1396	+++
1397	+++
1398	++
1399	+++
1400	+++
1401	+++
1402	+++
1403	+++
1404	+++
1405	+++
1406	+++
1407	+++
1408	+++
1409	+++
1410	+++
1411	+++
1412	+++
1413	+++
1414	+++
1415	+++
1416	+++
1417	++
1418	+++
1419	+++
1420	++

Cmpd. No.	EC30 (μM)
1421	+++
1422	+++
1423	+++
1424	+++
1425	+++
1426	+++
1427	+++
1428	+++
1429	+++
1430	+++
1431	+++
1432	+++
1433	++
1434	+++
1435	+++
1436	++
1437	++
1438	+++
1439	++
1440	+++
1441	++
1442	+++
1443	+++
1444	+++
1445	+++
1446	+++
1447	+++
1448	
1449	+++
1450	+++
1451	+++
1452	+++
1453	+++
1454	+++
1455	+++
1456	+++
1457	+++
1458	+++
1459	+++
1460	+++
1461	+++

Cmpd. No.	EC30 (μM)
1462	+++
1463	+++
1464	+++
1465	+++
1466	++
1467	+++
1468	+++
1469	++
1470	+++
1471	+++
1472	+++
1473	+++
1474	+++
1475	+++
1476	+++
1477	+++
1478	+++
1479	+++
1480	+++
1481	+++
1482	+++
1483	+++
1484	+++
1485	+++
1486	+++
1487	+++
1488	+++
1489	+++
1490	+++
1491	++
1492	++
1493	+++
1494	+++
1495	+++
1496	+++
1497	+++
1498	+++
1499	++
1500	+++
1501	+++
1502	+++

Cmpd. No.	EC30 (μM)
1503	++
1504	+++
1505	+++
1506	+++
1507	++
1508	+++
1509	+++
1510	+++
1511	+++
1512	+++
1513	+++
1514	+++
1515	+++
1516	+++
1517	+++
1518	+++
1519	+++
1520	+++
1521	+++
1522	+++
1523	+++
1524	+++
1525	+++
1526	+++
1527	+++
1528	++
1529	+++
1530	+++
1531	++
1532	+++
1533	+++
1534	++
1535	+++
1536	++
1537	+++
1538	+
1539	+++
1540	+
1541	+++
1542	+++
1543	

Cmpd. No.	EC30 (μM)
1544	+++
1545	+++
1546	+++
1547	++
1548	+++
1549	+++
1550	++
1551	+++
1552	+++
1553	+++
1554	+++
1555	+++
1556	+++
1557	+++
1558	++
1559	+++
1560	+++
1561	+++
1562	+++
1563	+++
1564	++
1565	+
1566	+++
1567	+++
1568	+++
1569	+++
1570	+++
1571	+++
1572	+++
1573	+++
1574	++
1575	++
1576	+++
1577	+++
1578	+++
1579	+++
1580	+++
1581	+++
1582	+++
1583	++
1584	+++

Cmpd. No.	EC30 (μM)
1585	+++
1586	+++
1587	+++
1588	+++
1589	+++
1590	+++
1591	+++
1592	+++
1593	+++
1594	+++
1595	+++
1596	+++
1597	+++
1598	+++
1599	+++
1600	+++
1601	+++
1602	+++
1603	++
1604	+++
1605	+++
1606	+++
1607	++
1608	+++
1609	+++
1610	+++
1611	+++
1612	+++
1613	+++
1614	+++
1615	+++
1616	+++
1617	+++
1618	+++
1619	+++
1620	+++
1621	+++
1622	+++
1623	+++
1624	+++
1625	++

Cmpd. No.	EC30 (μM)
1626	+++
1627	+++
1628	+++
1629	+++
1630	+++
1631	++
1632	++
1633	+++
1634	++
1635	+++
1636	++
1637	+++
1638	+++
1639	+++
1640	+++
1641	+++
1642	+++
1643	++
1644	+++
1645	+++
1646	+++
1647	+++
1648	+++
1649	++
1650	+++
1651	++
1652	+++
1653	+++
1654	+++
1655	+++
1656	+++
1657	+++
1658	++
1659	+++
1660	+++
1661	+++
1662	++
1663	+++
1664	+++
1665	+++
1666	+++

Cmpd. No.	EC30 (μM)
1667	++
1668	+++
1669	+++
1670	++
1671	+++
1672	+++
1673	+++
1674	+++
1675	++
1676	+++
1677	+++
1678	+
1679	+++
1680	+++
1681	+++
1682	+++
1683	+++
1684	+++
1685	+++
1686	+++
1687	+++
1688	+++
1689	+++
1690	+++
1691	++
1692	++
1693	+++
1694	+++
1695	+++
1696	+++
1697	+++
1698	+++
1699	+++
1700	+++
1701	+++
1702	+++
1703	+++
1704	+++
1705	+++
1706	++
1707	++

Cmpd. No.	EC30 (μM)
1708	+++
1709	+++
1710	+++
1711	+++
1712	+++
1713	+++
1714	++
1715	+++
1716	++
1717	+++
1718	+++
1719	+++
1720	+++
1721	+++
1722	+++
1723	+++
1724	+++
1725	++
1726	+++
1727	+++
1728	+++
1729	+++
1730	+++
1731	+++
1732	
1733	++
1734	+++
1735	+++
1736	++
1737	+++
1738	+++
1739	+++
1740	+++
1741	+++
1742	+++
1743	+++
1744	++
1745	+++
1746	+++
1747	+++
1748	+++

Cmpd. No.	EC30 (μM)
1749	+++
1750	+++
1751	+++
1752	++
1753	++
1754	+++
1755	++
1756	+++
1757	+++
1758	+++
1759	+++
1760	++
1761	++
1762	+++
1763	+++
1764	++
1765	+++
1766	+++
1767	+++
1768	++
1769	++
1770	+++
1771	+++
1772	++
1773	+++
1774	+++
1775	+++
1776	+++
1777	+++
1778	+++
1779	+++
1780	++
1781	+++
1782	+++
1783	+++
1784	+++
1785	+++
1786	++
1787	++
1788	+++
1789	+++

Cmpd. No.	EC30 (μM)
1790	+++
1791	+++
1792	+++
1793	+++
1794	++
1795	+++
1796	+++
1797	+++
1798	+++
1799	+++
1800	+++
1801	+++
1802	+++
1803	+++
1804	+++
1805	++
1806	+++
1807	+++
1808	+++
1809	+++
1810	+++
1811	+++
1812	++
1813	+++
1814	+++
1815	+++
1816	+++
1817	+++
1818	+++
1819	++
1820	++
1821	+++
1822	+++
1823	+++
1824	+++
1825	+++
1826	+++
1827	+++
1828	+++
1829	+++
1830	+++

Cmpd. No.	EC30 (μM)
1831	+++
1832	+++
1833	+++
1834	+++
1835	+++
1836	+++
1837	+++
1838	+++
1839	+++
1840	+++
1841	+++
1842	+++
1843	+++
1844	+++
1845	+++
1846	+++
1847	+++
1848	+++
1849	++
1850	+++
1851	+++
1852	+++
1853	++
1854	+++
1855	+++
1856	+++
1857	+++
1858	++
1859	+++
1860	+++
1861	+++
1862	+++
1863	+++
1864	+++
1865	++
1866	+++
1867	+++
1868	+++
1869	+++
1870	++
1871	+++

Cmpd. No.	EC30 (μM)
1872	+++
1873	+++
1874	+++
1875	++
1876	+++
1877	++
1878	++
1879	+++
1880	++
1881	+++
1882	+
1883	+++
1884	+++
1885	+++
1886	++
1887	+++
1888	++
1889	+++
1890	++
1891	+++
1892	+++
1893	+++
1894	+++
1895	+++
1896	+++
1897	+++
1898	++
1899	+++
1900	++
1901	+++
1902	+++
1903	+++
1904	+++
1905	+++
1906	+++
1907	+++
1908	+++
1909	+++
1910	+++
1911	+++
1912	+++

Cmpd. No.	EC30 (μM)
1913	+++
1914	+++
1915	+++
1916	+++
1917	+++
1918	+++
1919	+++
1920	++
1921	+++
1922	+++
1923	++
1924	+++
1925	+++
1926	+++
1927	+++
1928	++
1929	+++
1930	+++
1931	+++
1932	+++
1933	+++
1934	+++
1935	+++
1936	+++
1937	++
1938	+++
1939	++
1940	+++
1941	+++
1942	++
1943	++
1944	+++
1945	+++
1946	+++
1947	++
1948	+++
1949	+++
1950	+++
1951	+++
1952	+++
1953	+++

Cmpd. No.	EC30 (μM)
1954	+++
1955	+++
1956	+++
1957	++
1958	+++
1959	+++
1960	+++
1961	+++
1962	+++
1963	+++
1964	+++
1965	+++
1966	+++
1967	++
1968	+++
1969	+++
1970	+++
1971	+++
1972	+++
1973	+++
1974	+++
1975	+++
1976	+++
1977	+++
1978	+++
1979	+++
1980	+++
1981	+++
1982	++
1983	+++
1984	++
1985	++
1986	+++
1987	+++
1988	+++
1989	+++
1990	+++
1991	+++
1992	+++
1993	+++
1994	+++

Cmpd. No.	EC30 (μM)
1995	+++
1996	+++
1997	++
1998	+++
1999	++
2000	+++
2001	+++
2002	+++
2003	+++
2004	+++
2005	++
2006	+++
2007	+++
2008	++
2009	+++
2010	+++
2011	+++
2012	+++
2013	+++
2014	+++
2015	+++
2016	++
2017	+++
2018	+++
2019	+++
2020	+++
2021	++
2022	+++
2023	+++
2024	++
2025	+++
2026	+++
2027	+++
2028	+++
2029	+++
2030	+++
2031	+++
2032	++
2033	+++
2034	+++
2035	+++

Cmpd. No.	EC30 (μM)
2036	+++
2037	+++
2038	+++
2039	+++
2040	+++
2041	+++
2042	+++
2043	+++
2044	+++
2045	++
2046	+++
2047	++
2048	+++
2049	+++
2050	+++
2051	+++
2052	+++
2053	+++
2054	+++
2055	+++
2056	++
2057	+++
2058	+++
2059	+++
2060	+++
2061	+++
2062	+++
2063	+++
2064	+++
2065	+++
2066	+++
2067	+++
2068	+++
2069	+++
2070	+++
2071	+++
2072	++
2073	+++
2074	++
2075	+++
2076	+++

Cmpd. No.	EC30 (μM)
2077	+++
2078	++
2079	+++
2080	++
2081	+++
2082	++
2083	+++
2084	+++
2085	+++
2086	+++
2087	+++
2088	++
2089	+++
2090	+++
2091	+++
2092	+++
2093	+++
2094	+++
2095	+++
2096	+++
2097	+++
2098	+++
2099	+++
2100	+++
2101	+++
2102	+++
2103	+++
2104	+++
2105	+++
2106	+++
2107	+++
2108	+++
2109	+++
2110	+
2111	+++
2112	++
2113	+++
2114	+++
2115	+++
2116	+++
2117	+++

Cmpd. No.	EC30 (μM)
2118	+++
2119	+++
2120	+++
2121	+++
2122	+++
2123	+++
2124	+++
2125	+++
2126	+++
2127	+++
2128	++
2129	+++
2130	+++
2131	+
2132	+++
2133	+++
2134	+++
2135	++
2136	+++
2137	+++
2138	+++
2139	+++
2140	++
2141	+++
2142	+++
2143	+++
2144	+++
2145	+++
2146	+++
2147	+++
2148	++
2149	+++
2150	+++
2151	+++
2152	++
2153	+++
2154	+++
2155	++
2156	+++
2157	+
2158	++

Cmpd. No.	EC30 (μM)
2159	+++
2160	++
2161	+++
2162	+++
2163	+++
2164	+++
2165	+++
2166	++
2167	+++
2168	+++
2169	+++
2170	+++
2171	+++
2172	+++
2173	+++
2174	++
2175	++
2176	+++
2177	+++
2178	+++
2179	+++
2180	+++
2181	+++
2182	+++
2183	+++
2184	+++
2185	++
2186	+++
2187	+++
2188	+++
2189	+++
2190	+++
2191	+++
2192	+++
2193	+++
2194	+++
2195	+++
2196	+++
2197	+++
2198	+++
2199	++

Cmpd. No.	EC30 (μM)
2200	++
2201	+++
2202	+++
2203	+++
2204	+++
2205	+++
2206	+++
2207	+++
2208	+++
2209	+++
2210	+++
2211	+++
2212	++
2213	+++
2214	+++
2215	++
2216	+++
2217	+++
2218	+++
2219	++
2220	++
2221	++
2222	++
2223	+++
2224	+++
2225	++
2226	++
2227	+++
2228	+++
2229	++
2230	+++
2231	+++
2232	++
2233	+++
2234	+++
2235	+++
2236	+++
2237	+++
2238	+++
2239	+++
2240	++

Cmpd No.	EC30 (μM)
2241	+++
2242	+++
2243	+++
2244	+++
2245	+++
2246	+++
2247	++
2248	+++
2249	+++
2250	+++
2251	+++
2252	+++
2253	++
2254	++
2255	+++
2256	+++
2257	+++
2258	+++
2259	+++
2260	+++
2261	+++
2262	+++
2263	+++
2264	+++
2265	+++
2266	+++
2267	+++
2268	+++
2269	+++
2270	+++
2271	+++
2272	+++
2273	+++
2274	++
2275	++
2276	+++
2277	+++
2278	+++
2279	+++
2280	+++
2281	++

Cmpd No.	EC30 (μM)
2282	+++
2283	+++
2284	+++
2285	+++
2286	+++
2287	+++
2288	+++
2289	+++
2290	++
2291	+++
2292	+++
2293	+++
2294	++
2295	++
2296	+++
2297	+++
2298	+++
2299	+++
2300	++
2301	+++
2302	+++
2303	++
2304	+++
2305	+++
2306	+++
2307	+++
2308	+++
2309	+++
2310	+++
2311	+++
2312	+++
2313	+++
2314	++
2315	+++
2316	+++
2317	+++
2318	+++
2319	++
2320	+++
2321	++
2322	+++

Cmpd. No.	EC30 (μM)
2323	++
2324	++
2325	+++
2326	+++
2327	++
2328	++
2329	+++
2330	+++
2331	+++
2332	+++
2333	++
2334	+++
2335	++
2336	+++
2337	+++
2338	+++
2339	+++
2340	+++
2341	+++
2342	+++
2343	+++
2344	+++
2345	+++
2346	+++
2347	+++
2348	+++
2349	++
2350	++
2351	+++
2352	+++
2353	+++
2354	++
2355	+++
2356	+++
2357	+++
2358	+++
2359	++
2360	++
2361	+++
2362	+++
2363	++

Cmpd. No.	EC30 (μM)
2364	+++
2365	+++
2366	+++
2367	+++
2368	+++
2369	+++
2370	+++
2371	+++
2372	+++
2373	+++
2374	+++
2375	++
2376	+++
2377	+++
2378	+++
2379	+++
2380	+++
2381	+++
2382	+++
2383	++
2384	+++
2385	+++
2386	+++
2387	++
2388	+++
2389	++
2390	+++
2391	+++
2392	+++
2393	++
2394	+++
2395	+++
2396	++
2397	+++
2398	+++
2399	+++
2400	++
2401	+++
2402	+++
2403	+++
2404	+++

Cmpd. No.	EC30 (μM)
2405	+++
2406	+++
2407	++
2408	+++
2409	+++
2410	+++
2411	+++
2412	+++
2413	+++
2414	++
2415	+++
2416	+++
2417	+++
2418	+++
2419	++
2420	++
2421	+++
2422	++
2423	+++
2424	+++
2425	+++
2426	+++
2427	+++
2428	+++
2429	+++
2430	++
2431	+++
2432	+++
2433	++
2434	+++
2435	+++
2436	+++
2437	+++
2438	+++
2439	+++
2440	+++
2441	++
2442	++
2443	+++
2444	++
2445	+++

Cmpd. No.	EC30 (μM)
2446	+++
2447	+++
2448	+++
2449	+++
2450	+++
2451	+++
2452	+++
2453	+++
2454	++
2455	+++
2456	+++
2457	+++
2458	+++
2459	+++
2460	+++
2461	+++
2462	+++
2463	+++
2464	++
2465	+++
2466	+++
2467	+++
2468	+++
2469	++
2470	+++
2471	++
2472	++
2473	+++
2474	+++
2475	+++
2476	+++
2477	+++
2478	+++
2479	+++
2480	++
2481	+++
2482	+++
2483	+++
2484	+++
2485	+++
2486	+++

Cmpd. No.	EC30 (μM)
2487	+++
2488	+++
2489	++
2490	+++
2491	+++
2492	+++
2493	+++
2494	+++
2495	+++
2496	+++
2497	+++
2498	+++
2499	+++
2500	+++
2501	++
2502	+++
2503	+++
2504	++
2505	+++
2506	++
2507	+++
2508	+++
2509	+++
2510	+++
2511	+++
2512	+++
2513	++
2514	+++
2515	+++
2516	++
2517	+++
2518	+++
2519	+++
2520	++
2521	+++
2522	+++
2523	++
2524	+++
2525	++
2526	+++
2527	+++

Cmpd. No.	EC30 (μM)
2528	+++
2529	+++
2530	+++
2531	+++
2532	+++
2533	+++
2534	+++
2535	++
2536	+++
2537	++
2538	+++
2539	+++
2540	+++
2541	+++
2542	+++
2543	+++
2544	+
2545	+++
2546	+++
2547	+++
2548	+++
2549	+++
2550	++
2551	+++
2552	+++
2553	+++
2554	+++
2555	+++
2556	++
2557	+++
2558	+++
2559	+++
2560	+++
2561	+++
2562	+++
2563	+++
2564	+++
2565	++
2566	+++
2567	+++
2568	++

Cmpd No.	EC30 (μM)
2569	+++
2570	++
2571	+++
2572	+++
2573	+++
2574	++
2575	+++
2576	+++
2577	+++
2578	+++
2579	+++
2580	+++
2581	+++
2582	+++
2583	++
2584	++
2585	+++
2586	++
2587	++
2588	+++
2589	+++
2590	+++
2591	+++
2592	+++
2593	+++
2594	+++
2595	+++
2596	+++
2597	++
2598	+++
2599	++
2600	+++
2601	++
2602	++
2603	+++
2604	+++
2605	++
2606	+++
2607	+++
2608	+++
2609	+++

Cmpd No.	EC30 (μM)
2610	+++
2611	+++
2612	++
2613	+++
2614	++
2615	++
2616	+++
2617	++
2618	+++
2619	+++
2620	++
2621	++
2622	+++
2623	++
2624	++
2625	+++
2626	+++
2627	+++
2628	+++
2629	+++
2630	+++
2631	+++
2632	+++
2633	++
2634	+++
2635	+++
2636	+
2637	+++
2638	+++
2639	+++
2640	+++
2641	+++
2642	+++
2643	+++
2644	+++
2645	+++
2646	+++
2647	+++
2648	+++
2649	+++
2650	++

Cmpd. No.	EC30 (μM)
2651	+++
2652	+++
2653	+++
2654	+++
2655	+++
2656	+++
2657	++
2658	++
2659	++
2660	+++
2661	+++
2662	++
2663	+++
2664	+++
2665	+++
2666	++
2667	+++
2668	++
2669	+++
2670	+++
2671	+++
2672	+++
2673	+++
2674	+++
2675	+++
2676	+++
2677	+++
2678	++
2679	++
2680	++
2681	+++
2682	+++
2683	+++
2684	+++
2685	+++
2686	+++
2687	++
2688	+++
2689	++
2690	+++
2691	+++

Cmpd. No.	EC30 (μM)
2692	++
2693	++
2694	+++
2695	+++
2696	+++
2697	+++
2698	+++
2699	+++
2700	+++
2701	+++
2702	+++
2703	+++
2704	+++
2705	+++
2706	+++
2707	+++
2708	+++
2709	+++
2710	++
2711	++
2712	+++
2713	+++
2714	+++
2715	+++
2716	+++
2717	+++
2718	+++
2719	+++
2720	+++
2721	+++
2722	+++
2723	+++
2724	+++
2725	+++
2726	+++
2727	+++
2728	+++
2729	+++
2730	++
2731	++
2732	++


Cmpd. No.	EC30 (μM)
2733	++
2734	++
2735	++
2736	++
2737	+++
2738	+++
2739	+++
2740	+++
2741	++
2742	++
2743	+++
2744	+++
2745	+++
2746	+++
2747	+++

Other Embodiments

All publications and patents referred to in this disclosure are incorporated herein by reference to the same extent as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Should the meaning of the terms in any of the patents or publications incorporated by reference conflict with the meaning of the terms used in this disclosure, the meaning of the terms in this disclosure are intended to be controlling. Furthermore, the foregoing discussion discloses and describes merely exemplary embodiments of the invention. One skilled in the art will readily recognize from such discussion and from the accompanying drawings and claims, that various changes, modifications and variations can be made therein without departing from the spirit and scope of the invention as defined in the following claims.

CLAIMS

1. A compound of formula I:



I,

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring A is a C6-C10 aryl ring; C3-C10 cycloalkyl ring; or a C3-C14 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring B is a C3-C10 cycloalkyl ring; a C6-C10 aryl ring; or a C4-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C3-C14 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, NR, O, or S; or a C3-C10 cycloalkyl ring;

W is O, NR, or S;

X is O or NR;

Y is independently CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

Z is NR or CHR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-

C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy;C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; OH; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

n is 0, 1, 2 or 3;

o is 0, 1, 2, 3, 4, or 5;



p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5;



---- is a single bond or a double bond;


provided that the moieties containing ring B and ring C are substituted at adjacent positions on ring A.

2. The compound of claim 1, wherein ring A is a pyridyl, indole, indoline, isoindoline, pyrazole, pyrimidine, phenyl, quinoline, 5,6,7,8-tetrahydroquinoline, 1,2,3,4-tetrahydroisoquinoline, pyrrolidine, aza-indole, pyrrole, oxazole, pyrazine, triazole, benzimidazole, indazole, or imidazole ring.

3. The compound of claim 1 or 2, wherein ring A is selected from  ,  , or .

4. The compound of any one of claims 1 to 3, wherein ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, thiophene, dihydrobenzofuran, pyrazine, indazole, thiazole, pyridine-4(1H)-one, pyrrolidinone, or quinoline.


5. The compound of any one of claims 1 to 4, wherein ring B is selected from  ,  .



6. The compound of any one of claims 1 to 5, wherein ring C is cycloalkyl, pyrrolidine, piperidine, or azetidine.

7. The compound of claim 6, wherein ring C is pyrrolidine.

8. The compound of any one of claims 1 to 6, wherein ring C is



9. The compound of any one of claims 1 to 8, wherein Y is O.

10. The compound of any one of claims 1 to 8, wherein Y is CH_2.

11. The compound of any one of claims 1 to 8, wherein Y is CH(C1-C6 aliphatic).

12. The compound of any one of claims 1 to 8, wherein Y is $CH(CH_3)$.


13. The compound of any one of claims 1 to 8, wherein Y is $CH(CH_2CH_3)$.


14. The compound of any one of claims 1 to 8, wherein Y is N(C1-C6 aliphatic).


15. The compound of any one of claims 1 to 8, wherein Y is $N(CH_3)$.


16. The compound of any one of claims 1 to 15, wherein R_1 is halo, CN, C1-C6 aliphatic, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, azetidine, or thiophene ring, wherein all rings may be substituted with halo, C1-C6 aliphatic, C1-C6 alkoxy, C1-C6 fluoroaliphatic, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, C3-C10 heterocycloalkyl, or a (C1-C8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

17. The compound of any one of claims 1 to 16, wherein R_1 is selected from CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$,












18. The compound of any one of claims 1 to 17, wherein R_2 is selected from halo, OH, CN, azide, amino, C1-C6 aliphatic or fluoroaliphatic, C1-C6 alkoxy or fluoroalkoxy, C3-C10 mono- or bicyclic heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or a (C1-C8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.


19. The compound of any one of claims 1 to 18, wherein R_2 is selected from Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, CH_3, CH_2OH, CH_2CH_3, $CH(CH_3)_2$, CHF_2, OCH_3, OCF_3, $OCHF_2$, $OCH(CH_3)_2$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$, 



, or CO_2H.

20. The compound of any one of claims 1 to 19, wherein R_3 is selected from halo, CN, C1-C6 aliphatic or fluoroaliphatic, C1-C6 alkoxy, or C3-C10 mono- or bicyclic heteroaryl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR.

21. The compound of any one of claims 1 to 20, wherein R_3 is selected from Cl, F, CN, CH_3, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CO_2CH_3, tBu, $=CH_2$, =O, 



22. The compound of any one of claims 1 to 21, wherein o is 0.

23. The compound of any one of claims 1 to 21, wherein o is 1.

24. The compound of any one of claims 1 to 21, wherein o is 2.

25. The compound of any one of claims 1 to 24, wherein n is 0.

26. The compound of any one of claims 1 to 24, wherein n is 1.


27. The compound of any one of claims 1 to 24, wherein n is 2.

28. The compound of any one of claims 1 to 27, wherein p is 0.

29. The compound of any one of claims 1 to 27, wherein p is 1.

30. The compound of any one of claims 1 to 27, wherein p is 2.

31. The compound of claim 1, wherein the compound is of formula Ia:



Ia

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C3-C14 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 cycloalkyl ring;

X is O or NR;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;



n is 0, 1, or 2;

o is 0, 1, 2, 3, 4, or 5;



p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.

32. The compound of claim 31, wherein ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, thiophene, dihydrobenzofuran, pyrazine, indazole, thiazole, pyridine-4(1H)-one, pyrrolidinone, or quinoline.

33. The compound of claim 31 or 32, wherein ring B is selected from  ,  , , or .

34. The compound of any one of claims 31 to 33, wherein ring C is cycloalkyl, pyrrolidine, or piperidine.

35. The compound of any one of claims 31 to 34, wherein ring C is  , 



36. The compound of any one of claims 31 to 35, wherein Y is O.

37. The compound of any one of claims 31 to 35, wherein Y is CH_2.


38. The compound of any one of claims 31 to 35, wherein Y is CH(C1-C6 aliphatic).


39. The compound of any one of claims 31 to 35, wherein Y is $CH(CH_3)$.


40. The compound of any one of claims 31 to 35, wherein Y is $CH(CH_2CH_3)$.

41. The compound of any one of claims 31 to 40, wherein R_1 is halo, CN, C1-C6 aliphatic, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, or thiophene ring, wherein all rings may be substituted with halo, C1-C6 aliphatic, C1-C6 alkoxy, C1-C6 fluoroaliphatic, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, C3-C10 heterocycloalkyl, or a (C1-C8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.




42. The compound of any one of claims 31 to 41, wherein R_1 is selected from CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$,






















43. The compound of any one of claims 31 to 42, wherein R_2 is selected from halo, OH, CN, azide, amino, C1-C6 aliphatic or fluoroaliphatic, C1-C6 alkoxy or fluoroalkoxy, C3-C10 mono- or bicyclic heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or a (C1-C8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

44. The compound of any one of claims 31 to 43, wherein R_2 is selected from Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, CH_3, CH_2OH, CH_2CH_3, $CH(CH_3)_2$, CHF_2, OCH_3, OCF_3, $OCHF_2$, $OCH(CH_3)_2$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  ,  ,  , or CO_2H.

45. The compound of any one of claims 31 to 44, wherein R_3 is selected from halo, CN, C1-C6 aliphatic or fluoroaliphatic, C1-C6 alkoxy, or C3-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR.

46. The compound of any one of claims 31 to 45, wherein R_3 is selected from Cl, F, CN, CH_3, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CO_2CH_3, tBu, $=CH_2$, =O,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  ,  , or  .

47. The compound of any one of claims 31 to 46, wherein o is 0.

48. The compound of any one of claims 31 to 46, wherein o is 1.

49. The compound of any one of claims 31 to 48, wherein n is 0.

50. The compound of any one of claims 31 to 48, wherein n is 1.

51. The compound of any one of claims 31 to 48, wherein n is 2.

52. The compound of any one of claims 31 to 51, wherein p is 0.


53. The compound of any one of claims 31 to 51, wherein p is 1.

54. The compound of any one of claims 31 to 51, wherein p is 2.

55. The compound of any one of claims 31 to 54, wherein ring B is phenyl.

56. The compound of any one of claims 31 to 55, wherein ring B is independently phenyl or pyridyl.

57. The compound of claim 1, wherein the compound is of formula Ib:



Ib

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C3-C14 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 mono- or bicyclic cycloalkyl ring;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;


n is 0, 1, 2 or 3;


o is 0, 1, 2, 3, 4, or 5;

p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5.


58. The compound of claim 57, wherein ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, thiophene, dihydrobenzofuran, pyrazine, indazole, thiazole, pyridine-4(1H)-one, pyrrolidinone, or quinoline.

59. The compound of claim 57 or 58, wherein ring B is selected from 



60. The compound of any one of claims 57 to 59, wherein ring C is cycloalkyl or pyrrolidine.

61. The compound of any one of claims 57 to 60, wherein ring C is



62. The compound of any one of claims 57 to 61, wherein Y is O.

63. The compound of any one of claims 57 to 61, wherein Y is CH_2.


64. The compound of any one of claims 57 to 61, wherein Y is CH(C1-C6 aliphatic).


65. The compound of any one of claims 57 to 61, wherein Y is $CH(CH_3)$.


66. The compound of any one of claims 57 to 61, wherein Y is $CH(CH_2CH_3)$.


67. The compound of any one of claims 57 to 66, wherein R_1 is halo, CN, C1-C6 aliphatic, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, or thiophene ring, wherein all rings may be substituted with halo, C1-C6 aliphatic, C1-C6 alkoxy, C1-C6 fluoroaliphatic, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, C3-C10 heterocycloalkyl, or a (C1-C8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

68. The compound of any one of claims 57 to 67, wherein R_1 is selected from CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$,













69. The compound of any one of claims 57 to 68, wherein R_2 is selected from halo, OH, CN, azide, amino, C1-C6 aliphatic or fluoroaliphatic, C1-C6 alkoxy or fluoroalkoxy, C3-C10 mono- or bicyclic heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or a (C1-C8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

70. The compound of any one of claims 57 to 69, wherein R_2 is selected from Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, CH_3, CH_2OH, CH_2CH_3, $CH(CH_3)_2$, CHF_2, OCH_3, OCF_3, $OCHF_2$, $OCH(CH_3)_2$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  ,  , or CO_2H.

71. The compound of any one of claims 57 to 70, wherein R_3 is selected from halo, CN, C1-C6 aliphatic or fluoroaliphatic, C1-C6 alkoxy, or C3-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR.

72. The compound of any one of claims 57 to 71, wherein R_3 is selected from Cl, F, CN, CH_3, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CO_2CH_3, tBu, $=CH_2$, =O, 



73. The compound of any one of claims 57 to 72, wherein o is 0.

74. The compound of any one of claims 57 to 72, wherein o is 1.

75. The compound of any one of claims 57 to 72, wherein o is 2.

76. The compound of any one of claims 57 to 75, wherein n is 0.

77. The compound of any one of claims 57 to 75, wherein n is 1.

78. The compound of any one of claims 57 to 75, wherein n is 2.

79. The compound of any one of claims 57 to 78, wherein p is 0.


80. The compound of any one of claims 57 to 78, wherein p is 1.

81. The compound of any one of claims 57 to 78, wherein p is 2.

82. The compound of any one of claims 57 to 81, wherein ring B is pyridyl.

83. The compound of any one of claims 57 to 82, wherein ring B is phenyl.

84. The compound of claim 1, wherein the compound is of formula Ic:



Ic

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C3-C14 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 cycloalkyl ring;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;


n is 0, 1, 2 or 3;

o is 0, 1, 2, 3, 4, or 5;


p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5.

85. The compound of claim 84, wherein ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, thiophene, dihydrobenzofuran, pyrazine, indazole, thiazole, pyridine-4(1H)-one, pyrrolidinone, or quinoline.

86. The compound of claim 84 or 85, wherein ring B is selected from 

87. The compound of any one of claims 84 to 86, wherein ring C is cycloalkyl, or pyrrolidine,.

88. The compound of any one of claims 84 to 87, wherein ring C is 



89. The compound of any one of claims 84 to 88, wherein Y is O.

90. The compound of any one of claims 84 to 88, wherein Y is CH_2.


91. The compound of any one of claims 84 to 88, wherein Y is CH(C1-C6 aliphatic).


92. The compound of any one of claims 84 to 88, wherein Y is $CH(CH_3)$.


93. The compound of any one of claims 84 to 88, wherein Y is $CH(CH_2CH_3)$.


94. The compound of any one of claims 84 to 93, wherein R_1 is halo, CN, C1-C6 aliphatic, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, or thiophene ring, wherein all rings may be substituted with halo, C1-C6 aliphatic, C1-C6 alkoxy, C1-C6 fluoroaliphatic, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, C3-C10 heterocycloalkyl, or a (C1-C8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

95. The compound of any one of claims 84 to 94, wherein R_1 is selected from CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$,












96. The compound of any one of claims 84 to 95, wherein R_2 is selected from halo, OH, CN, azide, amino, C1-C6 aliphatic or fluoroaliphatic, C1-C6 alkoxy or fluoroalkoxy, C3-C10 mono- or bicyclic heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or a (C1-C8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

97. The compound of any one of claims 84 to 96, wherein R_2 is selected from Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, CH_3, CH_2OH, CH_2CH_3, $CH(CH_3)_2$, CHF_2, OCH_3, OCF_3, $OCHF_2$, $OCH(CH_3)_2$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$, 


, or CO_2H.

98. The compound of any one of claims 84 to 97, wherein R_3 is selected from halo, CN, C1-C6 aliphatic or fluoroaliphatic, C1-C6 alkoxy, or C3-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR.

99. The compound of any one of claims 84 to 98, wherein R_3 is selected from Cl, F, CN, CH_3, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CO_2CH_3, tBu, $=CH_2$, =O,  , or .

100. The compound of any one of claims 84 to 99, wherein o is 0.

101. The compound of any one of claims 84 to 99, wherein o is 1.

102. The compound of any one of claims 84 to 99, wherein o is 2.

103. The compound of any one of claims 84 to 102, wherein n is 0.

104. The compound of any one of claims 84 to 102, wherein n is 1.

105. The compound of any one of claims 84 to 102, wherein n is 2.


106. The compound of any one of claims 84 to 105, wherein p is 0.

107. The compound of any one of claims 84 to 105, wherein p is 1.

108. The compound of any one of claims 84 to 105, wherein p is 2.

109. The compound of any one of claims 84 to 108, wherein ring B is phenyl.

110. The compound of claim 1, wherein the compound is of formula Id:



Id

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring B is a C6-C10 aryl ring or C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR;

Ring C is a C3-C14 heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently N, O, or S, or a C3-C10 cycloalkyl ring;

Y is CRR, CO, O, S, SO, SO_2, S(O)NH or NR;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C1-C6 alkyl or fluoroalkyl; C1-C6 alkoxy or fluoroalkoxy; C1-C6 alkenyl; C1-C6 alkynyl; (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C1-C6 alkyl or fluoroalkyl;C1-C6 alkoxy or fluoroalkoxy;C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C13 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; CO_2R; C1-C6 alkyl or fluoroalkyl; C1-C6 alkenyl; C1-C6 alkynyl; C1-C6 alkoxy or fluoroalkoxy; or C6-C10 aryl; C3-C10 heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; or a (C1-C9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C6-C10 aryl, C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C3-C10 cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C1-C6 alkyl; C1-C6 alkyl; C1-C6 alkenyl; C1-C6 alkynyl; C6-C10 aryl; C3-C10 heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C3-C10 cycloalkyl;


n is 0, 1, 2 or 3;


o is 0, 1, 2, 3, 4, or 5;

p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5.


111. The compound of claim 110, wherein ring B is phenyl, pyridyl, pyridine-2(1*H*)-one, pyrazole, indole, thiophene, dihydrobenzofuran, pyrazine, indazole, thiazole, pyridine-4(1H)-one, pyrrolidinone, or quinoline.

112. The compound of claim 110 or 111, wherein ring B is selected from 



113. The compound of any one of claims 110 to 112, wherein ring C is cycloalkyl, or pyrrolidine.

114. The compound of any one of claims 110 to 113, wherein ring C is



115. The compound of any one of claims 110 to 114, wherein Y is O.


116. The compound of any one of claims 110 to 114, wherein Y is CH_2.


117. The compound of any one of claims 110 to 114, wherein Y is CH(C1-C6 aliphatic).


118. The compound of any one of claims 110 to 114, wherein Y is $CH(CH_3)$.


119. The compound of any one of claims 110 to 114, wherein Y is $CH(CH_2CH_3)$.

120. The compound of any one of claims 110 to 119, wherein R_1 is halo, CN, C1-C6 aliphatic, C1-C6 alkoxy, C3-C8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, or thiophene ring, wherein all rings may be substituted with halo, C1-C6 aliphatic, C1-C6 alkoxy, C1-C6 fluoroaliphatic, C1-C6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C3-C10 heteroaryl, C3-C10 heterocycloalkyl, or a (C1-C8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.



121. The compound of any one of claims 110 to 120, wherein R_1 is selected from CH_3, Cl, F, CN, OCH_3, CF_3, CH_2CH_3, tBu, $CH(CH_3)_2$, 









122. The compound of any one of claims 110 to 121, wherein R_2 is selected from halo, OH, CN, azide, amino, C1-C6 aliphatic or fluoroaliphatic, C1-C6 alkoxy or fluoroalkoxy, C3-C10 mono- or bicyclic heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or a (C1-C8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

123. The compound of any one of claims 110 to 122, wherein R_2 is selected from Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, CH_3, CH_2OH, CH_2CH_3, $CH(CH_3)_2$, CHF_2, OCH_3, OCF_3, $OCHF_2$, $OCH(CH_3)_2$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  ,  , or CO_2H.

124. The compound of any one of claims 110 to 123, wherein R_3 is selected from halo, CN, C1-C6 aliphatic or fluoroaliphatic, C1-C6 alkoxy, or C3-C10 mono- or bicyclic heteroaryl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR.

125. The compound of any one of claims 110 to 124, wherein R_3 is selected from Cl, F, CN, CH_3, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CO_2CH_3, tBu, $=CH_2$, =O, 



126. The compound of any one of claims 110 to 125, wherein o is 0.

127. The compound of any one of claims 110 to 125, wherein o is 1.

128. The compound of any one of claims 110 to 125, wherein o is 2.

129. The compound of any one of claims 110 to 128, wherein n is 0.

130. The compound of any one of claims 110 to 128, wherein n is 1.

131. The compound of any one of claims 110 to 128, wherein n is 2.

132. The compound of any one of claims 110 to 131, wherein p is 0.


133. The compound of any one of claims 110 to 131, wherein p is 1.

134. The compound of any one of claims 110 to 131, wherein p is 2.

135. The compound of any one of claims 110 to 134, wherein ring B is phenyl.

136. The compound of any one of claims 110 to 135, wherein ring B is pyridyl.

137. A compound of formula I:



I,

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring A is pyridine ;

Ring B is phenyl ;

Ring C is pyrrolidine ;

W is O;

X is O;

$(Y)_o$ is a bond ;

Z is NH ;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C_1-C_6 alkyl or fluoroalkyl; C_1-C_6 alkoxy or fluoroalkoxy; C_1-C_6 alkenyl; C_1-C_6 alkynyl; (C_1-C_9 alkylene)- R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C_6-C_{10} aryl; C_3-C_{10} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C_3-C_{10} cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C_1-C_6 alkyl or fluoroalkyl; C_1-C_6 alkoxy or fluoroalkoxy; C_1-C_6 alkenyl; C_1-C_6 alkynyl; C_6-C_{10} aryl; C_3-C_{13} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C_3-C_{10} cycloalkyl; or a (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; OH; CO_2R; C_1-C_6 alkyl or fluoroalkyl; C_1-C_6 alkenyl; C_1-C_6 alkynyl; C_1-C_6 alkoxy or fluoroalkoxy; or C_6-C_{10} aryl; C_3-C_{10} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C_3-C_{10} cycloalkyl; or a (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a = CH_2 or =O group;

R_4 is H; azide; CF_3; CHF_2 ; OR; CCH; CO_2R; OH; C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C_3-C_{10} cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2;C_1-C_6 alkyl; C_1-C_6 alkyl; C_1-C_6 alkenyl; C_1-C_6 alkynyl; C_6-C_{10} aryl; C_3-C_{10} heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C_3-C_{10} cycloalkyl;

n is 0, 1, 2 or 3;

p is 0, 1, 2, or 3; and


q is 0, 1, 2, 3, 4, or 5;


each ---- is independently a single bond or a double bond;


provided that the moieties containing ring B and ring C are substituted at adjacent positions on ring A.


138. The compound of claim 137, wherein R_1 is halo, CN, C_1-C_6 aliphatic, C_1-C_6 alkoxy, C_3-C_8 cycloalkyl, or a phenyl, pyridyl, pyrimidine, indole, aza-indole, azetidine, or thiophene ring, wherein all rings may be substituted with halo, C_1-C_6 aliphatic, C_1-C_6 alkoxy, C_1-C_6 fluoroaliphatic, C_1-C_6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C_3-C_{10} heteroaryl, C_3-C_{10} heterocycloalkyl, or a (C_1-C_8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

139. The compound of claim 137, wherein R_1 is selected from CH_3 , Cl, F, CN, OCH_3 , CF_3 , CH_2, CH_3 , tBu, $CH(CH_3)_2$,











140. The compound of claim 137, wherein R_2 is halo, OH, CN, azide, amino, C_1-C_6 aliphatic or fluoroaliphatic, C_1-C_6 alkoxy or fluoroalkoxy, C_3-C_{10} mono- or bicyclic heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR; or a (C_1-C_8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

141. The compound of claim 137, wherein R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, CH_3, CH_2OH, CH_2CH_3, $CH(CH_3)_2$, CHF_2, OCH_3, OCF_3, $OCHF_2$, $OCH(CH_3)_2$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  or CO_2H.

142. The compound of claim 137, wherein R_3 is halo, CN, C_1-C_6 aliphatic or fluoroaliphatic, C_1-C_6 alkoxy, or C_3-C_{10} mono- or bicyclic heteroaryl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR.

143. The compound of claim 137, wherein R_3 is Cl, F, CN, CH_3, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CO_2CH_3, tBu, $=CH_2$, ,



144. The compound of claim 137, wherein n is 1.


145. The compound of claim 137, wherein n is 2.

146. The compound of claim 137, wherein p is 0.

147. The compound of claim 137, wherein p is 1.

148. The compound of claim 137, wherein p is 2.

149. A pharmaceutical composition comprising a compound of formula I:



I,

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring A is pyridine;

Ring B is phenyl;

Ring C is pyrrolidine;

W is O;

X is O;

$(Y)_o$ is a bond;

Z is NH;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C_1-C_6 alkyl or fluoroalkyl; C_1-C_6 alkoxy or fluoroalkoxy; C_1-C_6 alkenyl; C_1-C_6 alkynyl; (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C_6-C_{10} aryl; C_3-C_{10} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C_3-C_{10} cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C_1-C_6 alkyl or fluoroalkyl;C_1-C_6 alkoxy or fluoroalkoxy;C_1-C_6 alkenyl; C_1-C_6 alkynyl; C_6-C_{10} aryl; C_3-C_{13} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C_3-C_{10} cycloalkyl; or a (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; OH; CO_2R; C_1-C_6 alkyl or fluoroalkyl; C_1-C_6 alkenyl; C_1-C_6 alkynyl; C_1-C_6 alkoxy or fluoroalkoxy; or C_6-C_{10} aryl; C_3-C_{10} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C_3-C_{10} cycloalkyl; or a (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C_3-C_{10} cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2C_1-C_6 alkyl; C_1-C_6 alkyl; C_1-C_6 alkenyl; C_1-C_6 alkynyl; C_6-C_{10} aryl; C_3-C_{10} heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C_3-C_{10} cycloalkyl;

n is 0, 1, 2 or 3;

p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5;

each ---- is independently a single bond or a double bond;

provided that the moieties containing ring B and ring C are substituted at adjacent positions on ring A; and

a pharmaceutically acceptable carrier.


150. The pharmaceutical composition of claim 149, further comprising one or more additional therapeutic agent(s).

151. The pharmaceutical composition of claim 150, wherein the additional therapeutic agent is a mucolytic agent, bronchodilator, an antibiotic, an anti-infective agent, a CFTR modulator, or an anti-inflammatory agent.

152. The pharmaceutical composition of claim 150, wherein the additional therapeutic agent is a CFTR modulator.

153. The pharmaceutical composition of claim 150, wherein the additional therapeutic agent is a CFTR corrector.


154. The pharmaceutical composition of claim 150, wherein the additional therapeutic agent is



,

or pharmaceutically acceptable salt thereof.


155. The pharmaceutical composition of claim 150, wherein the additional therapeutic agent is



,

or pharmaceutically acceptable salt thereof.

156. The pharmaceutical composition of claim 150, wherein the additional therapeutic agent is




,

or pharmaceutically acceptable salt thereof.

157. The pharmaceutical composition of claim 150, wherein the additional therapeutic agent is a CFTR potentiator.


158. The pharmaceutical composition of claim 150, wherein the additional therapeutic agent is




,



or pharmaceutically acceptable salt thereof.



159. The pharmaceutical composition of claim 150, wherein the additional therapeutic agents are a CFTR corrector and a CFTR potentiator.

160. The pharmaceutical composition of claim 150, wherein the additional therapeutic agents are  and




, or pharmaceutically acceptable salts thereof.

161. The pharmaceutical composition of claim 150, wherein the additional therapeutic agents are  and  , or pharmaceutically acceptable salts thereof.

162. The pharmaceutical composition of claim 150, wherein the additional therapeutic agents are  and  , or pharmaceutically acceptable salts thereof.

163. A kit comprising a compound of formula I:



I,

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring A is pyridine;

Ring B is phenyl;

Ring C is pyrrolidine;

W is O;

X is O;

$(Y)_o$ is a bond;

Z is NH;

R_1 is halo; CN; F_5S; SiR_3; OH; NRR; C_1-C_6 alkyl or fluoroalkyl; C_1-C_6 alkoxy or fluoroalkoxy; C_1-C_6 alkenyl; C_1-C_6 alkynyl; (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR; C_6-C_{10} aryl; C_3-C_{10} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C_3-C_{10} cycloalkyl;

R_2 is halo; OH; NRR; azide; CN; CO_2R; C_1-C_6 alkyl or fluoroalkyl; C_1-C_6 alkoxy or fluoroalkoxy; C_1-C_6 alkenyl; C_1-C_6 alkynyl; C_6-C_{10} aryl; C_3-C_{13} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C_3-C_{10} cycloalkyl; or a (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo; CN; OH; CO_2R; C_1-C_6 alkyl or fluoroalkyl; C_1-C_6 alkenyl; C_1-C_6 alkynyl; C_1-C_6 alkoxy or fluoroalkoxy; or C_6-C_{10} aryl; C_3-C_{10} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C_3-C_{10} cycloalkyl; or a (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H; azide; CF_3; CHF_2; OR; CCH; CO_2R; OH; C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; C_3-C_{10} cycloalkyl; NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H; OH; CO_2H; CO_2; C_1-C_6 alkyl; C_1-C_6 alkyl; C_1-C_6 alkenyl; C_1-C_6 alkynyl; C_6-C_{10} aryl; C_3-C_{10} heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR; or C_3-C_{10} cycloalkyl;

n is 0, 1, 2 or 3;

p is 0, 1, 2, or 3; and

q is 0, 1, 2, 3, 4, or 5;

each ---- is independently a single bond or a double bond;

provided that the moieties containing ring B and ring C are substituted at adjacent positions on ring A.


164. The kit of claim 163, further comprising one or more additional therapeutic agent(s).

165. The kit of claim 164, wherein the additional therapeutic agent is a mucolytic agent, bronchodilator, an antibiotic, an anti-infective agent, a CFTR modulator, or an anti-inflammatory agent.

166. The kit of claim 164, wherein the additional therapeutic agent is a CFTR modulator.

167. The kit of claim 164, wherein the additional therapeutic agent is a CFTR corrector.


168. The kit of claim 164, wherein the additional therapeutic agent is



,

or a pharmaceutically acceptable salt thereof.


169. The kit of claim 164, wherein the additional therapeutic agent is



,

or a pharmaceutically acceptable salt thereof.


170. The kit of claim 164, wherein the additional therapeutic agent is



or a pharmaceutically acceptable salt thereof.

171. The kit of claim 164, wherein the additional therapeutic agent is a CFTR potentiator.


172. The kit of claim 164, wherein the additional therapeutic agent is




or a pharmaceutically acceptable salt thereof.

173. The kit of claim 164, wherein the additional therapeutic agents are a CFTR corrector and a CFTR potentiator.

174. The kit of claim 164, wherein the additional therapeutic agents are





and





, or pharmaceutically acceptable salts thereof.

175. The kit of claim 164, wherein the additional therapeutic agents are


and

, or pharmaceutically acceptable salts thereof.

176. The kit of claim 164, wherein the additional therapeutic agents are



and

, or pharmaceutically acceptable salts thereof.

177. The kit of claim 164, wherein the compound of claim 1 and the one or more additional therapeutic agent(s) are in separate containers.


178. The kit of claim 164, wherein the compound of claim 1 and the one or more additional therapeutic agent(s) are in the same container.

179. The kit of claim 164, wherein the container(s) is a bottle, vial, or blister pack, or combination thereof.

180. A compound, wherein the compound has the following formula:



181. A pharmaceutically acceptable salt of a compound having the following formula:



182. A pharmaceutical composition comprising:

(i) a compound according to claim 180; and


(ii) a pharmaceutically acceptable carrier.

183. A pharmaceutical composition comprising:

(i) a pharmaceutically acceptable salt according to claim 181; and


(ii) a pharmaceutically acceptable carrier.

184. The pharmaceutical composition of claim 182, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

185. The pharmaceutical composition of claim 182, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

186. The pharmaceutical composition of claim 182, further comprising additional therapeutic agents having the following structural formulae:




and




, or pharmaceutically acceptable salts thereof.

187. The pharmaceutical composition of claim 183, further comprising an additional therapeutic agent having the following structural formula:



or a pharmaceutically acceptable salt thereof.


188. The pharmaceutical composition of claim 183, further comprising an additional therapeutic agent having the following structural formula:




,

or a pharmaceutically acceptable salt thereof.

189. The pharmaceutical composition of claim 183, further comprising additional therapeutic agents having the following structural formulae:




and




, or pharmaceutically acceptable salts thereof.

190. A compound, wherein the compound has the following formula:



.

191. A pharmaceutically acceptable salt of a compound having the following formula:



192. A pharmaceutical composition comprising:

(i) a compound according to claim 190; and


(ii) a pharmaceutically acceptable carrier.

193. A pharmaceutical composition comprising:

(i) a pharmaceutically acceptable salt according to claim 191; and


(ii) a pharmaceutically acceptable carrier.

194. The pharmaceutical composition of claim 192, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

195. The pharmaceutical composition of claim 192, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

196. The pharmaceutical composition of claim 192, further comprising additional therapeutic agents having the following structural formulae:




and




, or pharmaceutically acceptable salts thereof.

197. The pharmaceutical composition of claim 193, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

198. The pharmaceutical composition of claim 193, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

199. The pharmaceutical composition of claim 193, further comprising additional therapeutic agents having the following structural formulae:




and




, or pharmaceutically acceptable salts thereof.

200. A compound, wherein the compound has the following formula:



201. A pharmaceutically acceptable salt of a compound having the following formula:



202. A pharmaceutical composition comprising:

(i) a compound according to claim 200; and


(ii) a pharmaceutically acceptable carrier.

203. A pharmaceutical composition comprising:

(i) a pharmaceutically acceptable salt according to claim 201; and


(ii) a pharmaceutically acceptable carrier.

204. The pharmaceutical composition of claim 202, further comprising an additional therapeutic agent having the following structural formula:





or a pharmaceutically acceptable salt thereof.

205. The pharmaceutical composition of claim 202, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

206. The pharmaceutical composition of claim 202, further comprising additional therapeutic agents having the following structural formulae:

 and  , or


pharmaceutically acceptable salts thereof.

207. The pharmaceutical composition of claim 203, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

208. The pharmaceutical composition of claim 203, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

209. The pharmaceutical composition of claim 203, further comprising additional therapeutic agents having the following structural formulae:




and




, or pharmaceutically acceptable salts thereof.

210. A compound, wherein the compound has the following formula:



211. A pharmaceutically acceptable salt of a compound having the following formula:



212. A pharmaceutical composition comprising:

(i) a compound according to claim 210; and


(ii) a pharmaceutically acceptable carrier.

213. A pharmaceutical composition comprising:

(i) a pharmaceutically acceptable salt according to claim 211; and


(ii) a pharmaceutically acceptable carrier.

214. The pharmaceutical composition of claim 212, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

215. The pharmaceutical composition of claim 212, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

216. The pharmaceutical composition of claim 212, further comprising additional therapeutic agents having the following structural formulae:




and




, or pharmaceutically acceptable salts thereof.

217. The pharmaceutical composition of claim 213, further comprising an additional therapeutic agent having the following structural formula:



or a pharmaceutically acceptable salt thereof.


218. The pharmaceutical composition of claim 213, further comprising an additional therapeutic agent having the following structural formula:




,

or a pharmaceutically acceptable salt thereof.

219. The pharmaceutical composition of claim 213, further comprising additional therapeutic agents having the following structural formulae:




and



, or pharmaceutically acceptable salts thereof.

220. A compound of formula I:



I,

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring A is pyridine;

Ring B is pyridine;

Ring C is pyrrolidine;

W is O;

X is O;

$(Y)_o$ is a bond;

Z is NH ;

R_1 is halo, CN, F_5S, SiR_3, OH, NRR, C_1-C_6 alkyl or fluoroalkyl, C_1-C_6 alkoxy or fluoroalkoxy, C_1-C_6 alkenyl, C_1-C_6 alkynyl, (C_1-C_9 alkylene)- R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, or C_3-C_{10} cycloalkyl;

R_2 is halo, OH, NRR, azide, CN, CO_2R, C_1-C_6 alkyl or fluoroalkyl, C_1-C_6 alkoxy or fluoroalkoxy, C_1-C_6 alkenyl, C_1-C_6 alkynyl, C_6-C_{10} aryl, C_3-C_{13} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, C_3-C_{10} cycloalkyl, or a (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo, CN, OH, CO_2R, C_1-C_6 alkyl or fluoroalkyl, C_1-C_6 alkenyl, C_1-C_6 alkynyl, C_1-C_6 alkoxy or fluoroalkoxy, C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, C_3-C_{10} cycloalkyl, or a (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a = CH_2 or =O group;

R_4 is H, azide, CF_3, CHF_2, OR, CCH, CO_2R, OH, C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, C_3-C_{10} cycloalkyl, NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H, OH, CO_2H, CO_2, C_1-C_6 alkyl, C_1-C_6 alkyl, C_1-C_6 alkenyl, C_1-C_6 alkynyl, C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, or C_3-C_{10} cycloalkyl;

n is 0, 1, 2, or 3;

p is 0, 1, 2, or 3;


q is 0, 1, 2, 3, 4, or 5; and


---- is independently a single bond or a double bond;


provided that the moieties containing ring B and ring C are substituted at adjacent positions on ring A.


221. The compound of claim 220, wherein R_1 is halo, CN, C_1-C_6 aliphatic, C_1-C_6 alkoxy, C_3-C_8 cycloalkyl, phenyl, pyridyl, pyrimidine, indole, aza-indole, azetidine, or thiophene ring, wherein all rings may be substituted with halo, C_1-C_6 aliphatic, C_1-C_6 alkoxy, C_1-C_6 fluoroaliphatic, C_1-C_6 fluoroalkoxy, OH, CH_2OH, CH_2OCH_3, CN, CO_2H, amino, amido, C_3-C_{10} heteroaryl, C_3-C_{10} heterocycloalkyl, or a (C_1-C_8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2 or NR.

222. The compound of claim 220, wherein R_1 is selected from CH_3, Cl, F, CN, OCH_3, CF_3, CH_2, CH_3, tBu, $CH(CH_3)_2$,











223. The compound of claim 220, wherein R_2 is halo, OH, CN, azide, amino, C_1-C_6 aliphatic or fluoroaliphatic, C_1-C_6 alkoxy or fluoroalkoxy, C_3-C_{10} mono- or bicyclic heterocyclic ring wherein up to 4 carbon atoms may be replaced with O, S, N, or NR, or a (C_1-C_8 aliphatic)-R_4 wherein up to three CH_2 units may be replaced with O, CO, S, SO, SO_2, or NR.

224. The compound of claim 220, wherein R_2 is Cl, F, OH, CN, N_3, NH_2, $NH(CH_3)$, $N(CH_3)_2$, CH_3, CH_2OH, CH_2CH_3, $CH(CH_3)_2$, CHF_2, OCH_3, OCF_3, $OCHF_2$, $OCH(CH_3)_2$, $C(O)CH_3$, CH_2CH_2OH, CH_2NH_2, $NH(CH_2)_2OH$, $NH(CH_2)_2N(CH_3)_2$, $NH(CH_2)_2NH_2$, $NH(CH_2)_3NH_2$, $NH(CH_2)_2OCH_3$, $NHCH(CH_3)_2$,  , or CO_2H.

225. The compound of claim 220, wherein R_3 is halo, CN, C_1-C_6 aliphatic or fluoroaliphatic, C_1-C_6 alkoxy, or C_3-C_{10} mono- or bicyclic heteroaryl or heterocycloalkyl wherein up to 4 carbon atoms may be replaced by O, S, N, or NR.

226. The compound of claim 220, wherein R_3 is Cl, F, CN, CH_3, OCH_3, CF_3, CH_2CH_3, CH_2CF_3, $CH_2CH_2CH_3$, OCH_2CH_3, CH_2OCH_3, $CH(CH_3)_2$, CCH, CO_2CH_3, tBu, $=CH_2$,



227. The compound of claim 220, wherein n is 1.


228. The compound of claim 220, wherein n is 2.

229. The compound of claim 220, wherein p is 0.

230. The compound of claim 220, wherein p is 1.

231. The compound of claim 220, wherein p is 2.

232. A pharmaceutical composition comprising a compound of formula I:



I,

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring A is pyridine;

Ring B is pyridine;

Ring C is pyrrolidine;

W is O;

X is O;

$(Y)_o$ is a bond;

Z is NH;

R_1 is halo, CN, F_5S, SiR_3, OH, NRR, C_1-C_6 alkyl or fluoroalkyl, C_1-C_6 alkoxy or fluoroalkoxy, C_1-C_6 alkenyl, C_1-C_6 alkynyl, (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, or C_3-C_{10} cycloalkyl;

R_2 is halo, OH, NRR, azide, CN, CO_2R, C_1-C_6 alkyl or fluoroalkyl, C_1-C_6 alkoxy or fluoroalkoxy, C_1-C_6 alkenyl, C_1-C_6 alkynyl, C_6-C_{10} aryl, C_3-C_{13} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, C_3-C_{10} cycloalkyl, or a (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo, CN, OH, CO_2R, C_1-C_6 alkyl or fluoroalkyl, C_1-C_6 alkenyl, C_1-C_6 alkynyl, C_1-C_6 alkoxy or fluoroalkoxy, or C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, C_3-C_{10} cycloalkyl, or a (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a $=CH_2$ or =O group;

R_4 is H, azide, CF_3, CHF_2, OR, CCH, CO_2R, OH, C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, C_3-C_{10} cycloalkyl, NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H, OH, CO_2H, CO_2C_1-C_6 alkyl, C_1-C_6 alkyl, C_1-C_6 alkenyl, C_1-C_6 alkynyl, C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, or C_3-C_{10} cycloalkyl;

n is 0, 1, 2, or 3;

p is 0, 1, 2, or 3;

q is 0, 1, 2, 3, 4, or 5; and

---- is independently a single bond or a double bond;

provided that the moieties containing ring B and ring C are substituted at adjacent positions on ring A; and

a pharmaceutically acceptable carrier.


233. The pharmaceutical composition of claim 232, further comprising one or more additional therapeutic agent(s).

234. The pharmaceutical composition of claim 233, wherein the additional therapeutic agent is a mucolytic agent, bronchodilator, an antibiotic, an anti-infective agent, a CFTR modulator, or an anti-inflammatory agent.

235. The pharmaceutical composition of claim 233, wherein the additional therapeutic agent is a CFTR modulator.

236. The pharmaceutical composition of claim 233, wherein the additional therapeutic agent is a CFTR corrector.


237. The pharmaceutical composition of claim 233, wherein the additional therapeutic agent is



,

or a pharmaceutically acceptable salt thereof.


238. The pharmaceutical composition of claim 233, wherein the additional therapeutic agent is



,

or a pharmaceutically acceptable salt thereof.

239. The pharmaceutical composition of claim 233, wherein the additional therapeutic agent is




,

or a pharmaceutically acceptable salt thereof.

240. The pharmaceutical composition of claim 233, wherein the additional therapeutic agent is a CFTR potentiator.


241. The pharmaceutical composition of claim 233, wherein the additional therapeutic agent is




,



or a pharmaceutically acceptable salt thereof.



242. The pharmaceutical composition of claim 233, wherein the additional therapeutic agents are a CFTR corrector and a CFTR potentiator.

243. The pharmaceutical composition of claim 233, wherein the additional therapeutic agents are  and




, or pharmaceutically acceptable salts thereof.

244. The pharmaceutical composition of claim 233, wherein the additional therapeutic agents are  and  , or pharmaceutically acceptable salts thereof.

245. The pharmaceutical composition of claim 233, wherein the additional therapeutic agents are  and  , or pharmaceutically acceptable salts thereof.

246. A kit comprising a compound of formula I:



I,

or a pharmaceutically acceptable salt thereof, wherein, independently for each occurrence:

Ring A is pyridine;

Ring B is pyridine;

Ring C is pyrrolidine;

W is O;

X is O;

$(Y)_o$ is a bond;

Z is NH;

R_1 is halo, CN, F_5S, SiR_3, OH, NRR, C_1-C_6 alkyl or fluoroalkyl, C_1-C_6 alkoxy or fluoroalkoxy, C_1-C_6 alkenyl, C_1-C_6 alkynyl, (C_1-C_9 alkylene)- R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR, C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, or C_3-C_{10} cycloalkyl;

R_2 is halo, OH, NRR, azide, CN, CO_2R, C_1-C_6 alkyl or fluoroalkyl, C_1-C_6 alkoxy or fluoroalkoxy, C_1-C_6 alkenyl, C_1-C_6 alkynyl, C_6-C_{10} aryl, C_3-C_{13} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, C_3-C_{10} cycloalkyl, or a (C_1-C_9 alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_2 groups taken together may form a $=CH_2$ or =O group;

R_3 is halo, CN, OH, CO_2R, C_1-C_6 alkyl or fluoroalkyl, C_1-C_6 alkenyl, C_1-C_6 alkynyl, C_1-C_6 alkoxy or fluoroalkoxy, C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocyclic ring wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, C_3-C_{10} cycloalkyl, or a (C_1-C_9

alkylene)-R_4 wherein up to four CH_2 units are independently replaced with O, CO, S, SO, SO_2 or NR;

or two R_3 groups taken together may form a = CH_2 or =O group;

R_4 is H, azide, CF_3, CHF_2, OR, CCH, CO_2R, OH, C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, C_3-C_{10} cycloalkyl, NRR, NRCOR, CONRR, CN, halo, or SO_2R;

R is independently H, OH, CO_2H, CO_2, C_1-C_6 alkyl, C_1-C_6 alkyl, C_1-C_6 alkenyl, C_1-C_6 alkynyl, C_6-C_{10} aryl, C_3-C_{10} heteroaryl or heterocycloalkyl wherein anywhere from 1 to 4 ring atoms are independently O, S, N, or NR, or C_3-C_{10} cycloalkyl;

n is 0, 1, 2, or 3;

p is 0, 1, 2, or 3;

q is 0, 1, 2, 3, 4, or 5; and

---- is independently a single bond or a double bond;

provided that the moieties containing ring B and ring C are substituted at adjacent positions on ring A;


and one or more additional therapeutic agent(s).

247. The kit of claim 246, wherein the additional therapeutic agent is a mucolytic agent, bronchodilator, an antibiotic, an anti-infective agent, a CFTR modulator, or an anti-inflammatory agent.

248. The kit of claim 246, wherein the additional therapeutic agent is a CFTR modulator.


249. The kit of claim 246, wherein the additional therapeutic agent is a CFTR corrector.

250. The kit of claim 246, wherein the additional therapeutic agent is




or a pharmaceutically acceptable salt thereof.

251. The kit of claim 246, wherein the additional therapeutic agent is



or a pharmaceutically acceptable salt thereof.


252. The kit of claim 246, wherein the additional therapeutic agent is



or a pharmaceutically acceptable salt thereof.

253. The kit of claim 246, wherein the additional therapeutic agent is a CFTR potentiator.



254. The kit of claim 246, wherein the additional therapeutic agent is



or a pharmaceutically acceptable salt thereof.



255. The kit of claim 246, wherein the additional therapeutic agents are a CFTR corrector and a CFTR potentiator.

256. The kit of claim 246, wherein the additional therapeutic agents are

 and  , or



pharmaceutically acceptable salts thereof.

257. The kit of claim 246, wherein the additional therapeutic agents are

 and  , or

pharmaceutically acceptable salts thereof.

258. The kit of claim 246, wherein the additional therapeutic agents are

 and  , or


pharmaceutically acceptable salts thereof.

259. The kit of claim 246, wherein the compound of claim 1 and the one or more additional therapeutic agent(s) are in separate containers.


260. The kit of claim 246, wherein the compound of claim 1 and the one or more additional therapeutic agent(s) are in the same container.

261. The kit of claim 246, wherein the container(s) is a bottle, vial, or blister pack, or combination thereof.

262. A compound, wherein the compound has the following formula:



263. A pharmaceutically acceptable salt of a compound having the following formula:



264. A pharmaceutical composition comprising:

(i) a compound according to claim 262; and


(ii) a pharmaceutically acceptable carrier.

265. A pharmaceutical composition comprising:

(i) a pharmaceutically acceptable salt according to claim 263; and


(ii) a pharmaceutically acceptable carrier.

266. The pharmaceutical composition of claim 264, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

267. The pharmaceutical composition of claim 264, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

268. The pharmaceutical composition of claim 264, further comprising additional therapeutic agents having the following structural formulae:




and




, or pharmaceutically acceptable salts thereof.

269. The pharmaceutical composition of claim 265, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

270. The pharmaceutical composition of claim 265, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

271. The pharmaceutical composition of claim 265, further comprising additional therapeutic agents having the following structural formulae:




and




, or pharmaceutically acceptable salts thereof.

272. A compound, wherein the compound has the following formula:



273. A pharmaceutically acceptable salt of a compound having the following formula:



274. A pharmaceutical composition comprising:

(i) a compound according to claim 272; and


(ii) a pharmaceutically acceptable carrier.

275. A pharmaceutical composition comprising:

(i) a pharmaceutically acceptable salt according to claim 273; and


(ii) a pharmaceutically acceptable carrier.

276. The pharmaceutical composition of claim 274, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

277. The pharmaceutical composition of claim 274, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

278. The pharmaceutical composition of claim 274, further comprising additional therapeutic agents having the following structural formulae:




and



, or


pharmaceutically acceptable salts thereof.

279. The pharmaceutical composition of claim 275, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

280. The pharmaceutical composition of claim 275, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

281. The pharmaceutical composition of claim 275, further comprising additional therapeutic agents having the following structural formulae:




and




, or pharmaceutically acceptable salts thereof.

282. A compound, wherein the compound has the following formula:



283. A pharmaceutically acceptable salt of a compound having the following formula:



284. A pharmaceutical composition comprising:

(i) a compound according to claim 282; and


(ii) a pharmaceutically acceptable carrier.

285. A pharmaceutical composition comprising:

(i) a pharmaceutically acceptable salt according to claim 283; and


(ii) a pharmaceutically acceptable carrier.

286. The pharmaceutical composition of claim 284, further comprising an additional therapeutic agent having the following structural formula:





or a pharmaceutically acceptable salt thereof.

287. The pharmaceutical composition of claim 284, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

288. The pharmaceutical composition of claim 284, further comprising additional therapeutic agents having the following structural formulae:

 and  , or


pharmaceutically acceptable salts thereof.

289. The pharmaceutical composition of claim 285, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

290. The pharmaceutical composition of claim 285, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

291. The pharmaceutical composition of claim 285, further comprising additional therapeutic agents having the following structural formulae:




and




, or pharmaceutically acceptable salts thereof.

292. A compound, wherein the compound has the following formula:



293. A pharmaceutically acceptable salt of a compound having the following formula:



294. A pharmaceutical composition comprising:

(i) a compound according to claim 292; and


(ii) a pharmaceutically acceptable carrier.

295. A pharmaceutical composition comprising:

(i) a pharmaceutically acceptable salt according to claim 293; and


(ii) a pharmaceutically acceptable carrier.

296. The pharmaceutical composition of claim 294, further comprising an additional therapeutic agent having the following structural formula:





or a pharmaceutically acceptable salt thereof.

297. The pharmaceutical composition of claim 294, further comprising an additional therapeutic agent having the following structural formula:




or a pharmaceutically acceptable salt thereof.

298. The pharmaceutical composition of claim 294, further comprising additional therapeutic agents having the following structural formulae:

 and  , or


pharmaceutically acceptable salts thereof.

299. The pharmaceutical composition of claim 295, further comprising an additional therapeutic agent having the following structural formula:



or a pharmaceutically acceptable salt thereof.


300. The pharmaceutical composition of claim 295, further comprising an additional therapeutic agent having the following structural formula:




,

or a pharmaceutically acceptable salt thereof.

301. The pharmaceutical composition of claim 295, further comprising additional therapeutic agents having the following structural formulae:















and



























, or













pharmaceutically acceptable salts thereof.













302. A compound or a pharmaceutically acceptable salt thereof, wherein the compound is selected from:













1	2	3
		
4	5	6
		
7	8	0
		
10	11	12
		













13	14	15
		
16	17	18
		
19	20	21
		
22	23	24
		













25	26	27
		
28	29	30
		
31	32	33
		
34	35	36
		













37	38	39
		
40	41	42
		
43	44	45
		
46	47	48
		













49	50	51
		
52	53	54
		
55	56	57
		
58	59	60
		













61	62	63
		
64	65	66
		
67	68	69
		
70	71	72
		













73	74	75
		
76	77	78
		
79	80	81
		
82	83	84
		













85	86	87
		
88	89	90
		
91	92	93
		
94	95	96
		













97	98	99
		
100	101	102
		
103	104	105
		
106	107	108
		













109	110	111
		
112	113	114
		
115	116	117
		
118	119	120
		













121	122	123
		
124	125	126
		
127	128	129
		
130	131	132
		













133	134	135
		
136	137	138
		
139	140	141
		
142	143	144
		













145	146	147
		
148	149	150
		
151	152	153
		
154	155	156
		













157	158	159
		
160	161	162
		
163	164	165
		
166	167	168
		













169	170	171
		
172	173	174
		
175	176	177
		
178	179	180
		













181	182	183
		
184	185	186
		
187	188	189
		
190	191	192
		













193	194	195
		
196	197	198
		
199	200	201
		
202	203	204
		













205	206	207
		
208	209	210
		
211	212	213
		
214	215	216
		













217	218	219
		
220	221	222
		
223	224	225
		
226	227	228
		













229	230	231
		
232	233	234
		
235	236	237
		
238	239	240
		













241	242	243
		
244	245	246
		
247	248	249
		
250	251	252
		













253	254	255
		
256	257	258
		
259	260	261
		
262	263	264
		













265	266	267
		
268	269	270
		
271	272	273
		
274	275	276
		













277	278	279
		
280	281	282
		
283	284	285
		
286	287	288
		













289	290	291
		
292	293	294
		
295	296	297
		
298	299	300
		













301	302	303
		
304	305	306
		
307	308	309
		
310	311	312
		













313	314	315
		
316	317	318
		
319	320	321
		
322	323	324
		













325	326	327
		
328	329	330
		
331	332	333
		
334	335	336
		













337	338	339
		
340	341	342
		
343	344	345
		
346	347	348
		













349	350	351
		
352	353	354
		
355	356	357
		
358	359	360
		













361	362	363
		
364	366	366
		
367	368	369
		
370	371	372
		













373	374	375
		
376	377	378
		
379	380	381
		
382	383	384
		













385	386	387
		
388	389	390
		
391	392	393
		
394	395	396
		













397	398	399
		
400	401	402
		
403	404	405
		
406	407	408
		













409	410	411
		
412	413	414
		
415	416	417
		
418	419	420
		













421	422	423
		
424	425	426
		
427	428	429
		
430	431	432
		













433	434	435
		
436	437	438
		
439	440	441
		
442	443	444
		













445	446	447
		
448	449	450
		
451	452	453
		
454	455	456
		













457	458	459
		
460	461	462
		
463	464	465
		
466	467	468
		













469	470	471
		
472	473	474
		
475	476	477
		
478	479	480
		













481	482	483
		
484	485	486
		
487	488	489
		
490	491	492
		













493	494	495
		
496	497	498
		
499	500	501
		
502	503	504
		













505	506	507
		
508	509	510
		
511	512	513
		
514	515	516
		













517	518	519
		
520	521	522
		
523	524	525
		
526	527	528
		













529	530	531
		
532	533	534
		
535	536	537
		
538	539	540□
		













541	542	543
		
544	545	546
		
547	548	549
		
550	551	552
		













553	554	555
		
556	557	558
		
559	560	561
		
562	563	564
		













565	566	567
		
568	569	570
		
571	572	573
		
574	575	576
		













577	578	579
		
580	581	582
		
583	584	585
		
586	587	588
		













589	590	591
		
592	593	594
		
595	596	597
		
598	599	600
		













601	602	603
		
604	605	606
		
607	608	609
		
610	611	612
		













613	614	615
		
616	617	618
		
619	620	621
		
622	623	624
		













625	626	627
		
628	629	630
		
631	632	633
		
634	635	636
		













637	636	639
		
640	641	642
		
643	644	645
		
646	647	648
		













649	650	651
		
652	653	654
		
655	656	657
		
658	659	660
		













661	662	663
		
664	665	666
		
667	668	669
		
670	671	672
		













673	674	675
		
676	677	678
		
679	680	681
		
682	683	684
		













685	686	687
		
688	689	690
		
691	692	693
		
694	695	696
		













697	698	699
		
700	701	702
		
703	704	705
		
706	707	708
		













709	710	711
		
712	713	714
		
715	716	717
		
718	719	720
		













721	722	723
		
724	725	726
		
727	728	729
		
730	731	732
		













733	734	735
		
736	737	738
		
739	740	741
		
742	743	744
		













745	746	747
		
748	749	750
		
751	752	753
		
754	755	756
		













757	758	759
		
760	761	762
		
763	764	765
		
766	767	766
		













769	770	771
		
772	773	774
		
775	776	777
		
778	779	780
		













781	782	783
		
784	785	786
		
787	788	789
		
790	791	792
		













793	794	795
		
796	797	798
		
799	800	801
		
802	803	804
		













805	806	807
		
808	809	810
		
811	812	813
		
814	815	816
		













817	818	819
		
820	821	822
		
823	824	825
		
826	827	828
		













829	830	831
		
832	833	834
		
835	836	837
		
838	839	840
		













841	842	843
		
844	845	846
		
847	848	849
		
850	851	852
		













853	854	855
		
856	857	858
		
859	860	861
		
862	863	864
		













865	866	867
		
868	869	870
		
871	872	873
		
874	875	876
		













877	878	879
		
880	881	882
		
883	884	885
		
886	887	888
		













889	890	891
		
892	893	894
		
895	896	897
		
898	899	900
		













901	902	903
		
904	905	906
		
907	908	909
		
910	911	912
		













913	914	915
		
916	917	918
		
919	920	921
		
922	923	924
		













925	926	927
		
928	929	930
		
931	932	933
		
934	935	936
		













937	938	939
		
940	941	942
		
943	944	945
		
940	947	943
		













949	950	951
		
952	953	954
		
955	956	957
		
958	959	960
		













961	962	963
		
964	965	966
		
967	968	969
		
970	971	972
		













973	974	975
		
976	977	978
		
979	980	981
		
982	983	984
		













985	986	987
		
988	989	990
		
991	992	993
		
994	995	990
		













997	998	999
		
1000	1001	1002
		
1003	1004	1005
		
1006	1007	1008
		













1009	1010	1011
		
1012	1013	1014
		
1015	1016	1017
		
1018	1019	1020
		













1021	1022	1023
		
1024	1025	1026
		
1027	1028	1029
		
1030	1031	1032
		













1033	1034	1035
		
1036	1037	1038
		
1039	1040	1041
		
1042	1043	1044
		













1045	1046	1047
		
1048	1049	1050
		
1051	1052	1053
		
1054	1055	1056
		













1057	1058	1059
		
1060	1061	1062
		
1063	1064	1065
		
1066	1067	1063
		













1069	1070	1071
		
1072	1073	1074
		
1075	1076	1077
		
1078	1079	1080
		













1081	1082	1083
		
1084	1085	1086
		
1087	1088	1089
		
1090	1091	1092
		













1093	1094	1095
		
1095	1097	1093
		
1099	1100	1101
		
1102	1103	1104
		













1105	1106	1107
		
1108	1109	1110
		
1111	1112	1113
		
1114	1115	1116
		













1117	1118	1119
		
1120	1121	1122
		
1123	1124	1125
		
1126	1127	1128
		













1129	1130	1131
		
1132	1133	1134
		
1135	1136	1137
		
1138	1139	1140
		













1141	1142	1143
		
1144	1145	1146
		
1147	1148	1149
		
1150	1151	1152
		













1153	1154	1155
		
1156	1157	1158
		
1159	1160	1161
		
1162	1163	1164
		













1165	1166	1167
		
1168	1169	1170
		
1171	1172	1173
		
1174	1175	1176
		













1177	1178	1179
		
1180	1181	1182
		
1183	1184	1185
		
1186	1187	1188
		













1189	1190	1191
		
1192	1193	1194
		
1195	1196	1197
		
1198	1199	1200
		













1201	1202	1203
		
1204	1205	1206
		
1207	1208	1209
		
1210	1211	1212
		













1213	1214	1215
		
1216	1217	1218
		
1219	1220	1221
		
1222	1223	1224
		













1225	1226	1227
		
1228	1229	1230
		
1231	1232	1233
		
1234	1235	1236
		













1237	1238	1239
		
1240	1241	1242
		
1243	1244	1245
		
1246	1247	1248
		













1249	1250	1251
		
1252	1253	1254
		
1255	1256	1257
		
1258	1259	1260
		













1261	1262	1263
		
1264	1265	1266
		
1267	1268	1269
		
1270	1271	1272
		













1273	1274	1275
		
1276	1277	1278
		
1279	1280	1281
		
1282	1283	1284
		













1285	1286	1287
		
1288	1289	1290
		
1291	1292	1293
		
1294	1295	1296
		













1297	1298	1299
		
1300	1301	1302
		
1303	1304	1305
		
1306	1307	1308
		













1309	1310	1311
		
1312	1313	1314
		
1315	1316	1317
		
1318	1319	1320
		













1321	1322	1323
		
1324	1325	1326
		
1327	1328	1329
		
1330	1331	1332
		













1333	1334	1335
		
1336	1337	1338
		
1339	1340	1341
		
1342	1343	1344
		













1345	1346	1347
		
1348	1349	1350
		
1351	1352	1353
		
1354	1355	1356
		













1357	1358	1359
		
1360	1361	1362
		
1363	1364	1365
		
1366	1367	1368
		













1369	1370	1371
		
1372	1373	1374
		
1375	1376	1377
		
1378	1379	1380
		













1381	1382	1383
		
1384	1385	1386
		
1387	1388	1389
		
1390	1391	1392
		













1393	1394	1395
		
1396	1397	1398
		
1399	1400	1401
		
1402	1403	1404
		













1405	1406	1407
		
1408	1409	1410
		
1411	1412	1413
		
1414	1415	1416
		













1417	1418	1419
		
1420	1421	1422
		
1423	1424	1425
		
1426	1427	1428
		













1429	1430	1431
		
1432	1433	1434
		
1435	1436	1437
		
1438	1439	1440
		













1441	1442	1443
		
1444	1445	1446
		
1447	1448	1449
		
1450	1451	1452
		













1453	1454	1455
		
1456	1457	1458
		
1459	1460	1461
		
1462	1463	1464
		













1465	1466	1467
		
1468	1469	1470
		
1471	1472	1473
		
1474	1475	1476
		













1477	1478	1479
		
1480	1481	1482
		
1483	1484	1485
		
1486	1487	1488
		













1489	1490	1491
		
1492	1493	1494
		
1495	1496	1497
		
1498	1499	1500
		













1501	1502	1503
		
1504	1505	1506
		
1507	1508	1509
		
1510	1511	1512
		













1513	1514	1515
		
1516	1517	1518
		
1519	1520	1521
		
1522	1523	1524
		













1525	1526	1527
		
1528	1529	1530
		
1531	1532	1533
		
1534	1535	1536
		













1537	1538	1539
		
1540	1541	1542
		
1543	1544	1545
		
1543	1547	1548
		













1549	1550	1551
		
1552	1553	1554
		
1555	1556	1557
		
1558	1559	1560
		













1561	1562	1563
		
1564	1565	1566
		
1567	1568	1569
		
1570	1571	1572
		










1573	1574	1575
		
1576	1577	1578
		
1579	1580	1581
		
1582	1583	1584
		













1585	1586	1587
		
1588	1589	1590
		
1591	1592	1593
		
1594	1595	1596
		













1597	1598	1599
		
1600	1601	1602
		
1603	1604	1605
		
1606	1607	1608
		













1609	1610	1611
		
1612	1613	1614
		
1615	1616	1617
		
1618	1619	1620
		













1621	1622	1623
		
1624	1625	1626
		
1627	1628	1629
		
1630	1631	1632
		













1633	1634	1635
		
1636	1637	1638
		
1639	1640	1641
		
1642	1643	1644













		
1645	1646	1647
		
1648	1649	1650
		
1651	1652	1653
		
1654	1655	1656













		
1657	1658	1659
		
1660	1661	1662
		
1663	1664	1665
		
1666	1667	1668













		
1669	1670	1671
		
1672	1673	1674
		
1675	1676	1677
		
1678	1679	1680


		
1681	1682	1683
		
1684	1685	1686
		
1687	1688	1689
		
1690	1691	1692


		
1693	1694	1695
		
1696	1697	1698
		
1699	1700	1701
		
1702	1703	1704


		
1705	1706	1707
		
1708	1709	1710
		
1711	1712	1713
		
1714	1715	1716










		
1717	1718	1719
		
1720	1721	1722
		
1723	1724	1725
		
1726	1727	1728













		
1729	1730	1731
		
1732	1733	1734
		
1735	1736	1737
		
1738	1739	1740























































1741	1742	1743
		
1744	1745	1746
		
1747	1748	1749
		




1750	1751	1752
		
1753	1754	1755
		
1756	1757	1758
		
1759	1760	1761
		




1762	1763	1764
		
1765	1766	1767
		
1768	1769	1770
		
1771	1772	1773
		




1774	1775	1776
		
1777	1778	1779
		
1780	1781	1782
		
1783	1784	1785
		




1786	1787	1788
		
1789	1790	1791
		
1792	1793	1794
		
1795	1796	1797
		













1798	1799	1800
		
1801	1802	1803
		
1804	1805	1806
		
1807	1808	1809
		




1810	1811	1812
		










1813	1814	1815
		




1816	1817	1818
		




1819	1820	1821
		




1822	1823	1824
		
1825	1826	1827
		
1828	1829	1830
		
1831	1832	1833
		




1834	1835	1836
		













1837	1838	1839
		
1840	1841	1842
		
1843	1844	1845
		













1846	1847	1848
		




1849	1850	1851
		




1852	1853	1854
		




1855	1856	1857
		




1858	1859	1860
		
1861	1862	1863
		
1864	1865	1866
		
1867	1868	1869
		













1870	1871	1872
		
1873	1874	1875
		
1876	1877	1878
		
1879	1880	1881
		













1882	1883	1884
		













1885	1886	1887
		













1888	1889	1890
		













1891	1892	1893
		













1894	1895	1896
		
1897	1898	1899
		
1900	1901	1902
		
1903	1904	1905
		
1906	1907	1908


		
1909	1910	1911
		
1912	1913	1914
		
1915	1916	1917
		
1918	1919	1920


		
1921	1922	1923
		
1924	1925	1926
		
1927	1928	1929
		
1930	1931	1932


		
1933	1934	1935
		
1936	1937	1938
		
1939	1940	1941
		
1942	1943	1944










		
1945	1946	1947
		
1948	1949	1950
		
1951	1952	1953
		
1954	1955	1956













		
1957	1958	1959
		
1960	1961	1962
		
1963	1964	1965
		
1966	1967	1968





































1969	1970	1971
		
1972	1973	1974
		
1975	1976	1977
		




1978	1979	1980
		
1981	1982	1983
		
1984	1985	1986
		
1987	1988	1989
		




1990	1991	1992
		
1993	1994	1995
		
1996	1997	1998
		
1999	2000	2001
		




2002	2003	2004
		
2005	2006	2007
		
2008	2009	2010
		
2011	2012	2013
		




2014	2015	2016
		




2017	2018	2019
		

2020	2021	2022
		













2023	2024	2025
		







2026	2027	2028
		







2029	2030	2031
		













2032	2033	2034
		







2035	2036	2037







		
2038	2039	2040
		
2041	2042	2043
		
2044	2045	2046
		
2047	2048	2049













		
2050	2051	2052
		













2053	2054	2055
		
2056	2057	2058
		
2059	2060	2061













		
2062	2063	2064
		
2065	2066	2067
		
2068	2069	2070
		
2071	2072	2073













		
2074	2075	2076
		













2077	2078	2079
		
2080	2081	2082
		













2083	2084	2085
		
2086	2087	2088
		
2089	2090	2091
		
2092	2093	2094
		













2095	2096	2097
		
2098	2099	2100
		
2101	2102	2103
		
2104	2105	2106
		













2107	2108	2109
		
2110	2111	2112
		
2113	2114	2115
		
2116	2117	2118
		













2119	2120	2121
		
2122	2123	2124
		
2125	2126	2127
		
2128	2129	2130
		







2131	2132	2133
		
2134	2135	2136
		
2137	2138	2139
		
2140	2141	2142
		







2143	2144	2145
		
2146	2147	2148
		
2149	2150	2151
		
2152	2153	2154
		







2155	2156	2157
		
2158	2159	2160
		
2161	2162	2163
		
2164	2165	2166
		







2167	2168	2169
		
2170	2171	2172
		
2173	2174	2175
		
2176	2177	2178
		













2179	2180	2181
		
2182	2183	2184
		
2185	2186	2187
		
2188	2189	2190
		







2191	2192	2193
		
2194	2195	2196
		







2197	2198	2199
		
2200	2201	2202
		







2203	2204	2205
		
2206	2207	2208
		




2209	2210	2211
		
2212	2213	2214
		













2215	2216	2217
		
2218	2219	2220
		
2221	2222	2223
		
2224	2225	2226
		













2227	2228	2229
		
2230	2231	2232
		










2233	2234	2235
		
2236	2237	2238
		




2239	2240	2241
		
2242	2243	2244
		










2245	2246	2247
		
2248	2249	2250




		
2251	2252	2253
		
2254	2255	2256
		
2257	2258	2259
		
2260	2261	2262













		
2263	2264	2265
		
2266	2267	2268
		
2260	2270	2271
		
2273	2273	2274













		
2275	2276	2277
		
2278	2279	2280
		













2281	2282	2283
		
2284	2285	2286













		
2287	2288	2289
		
2290	2291	2292
		













2293	2294	2295
		
2296	2297	2298













		
2299	2300	2301
		
2302	2303	2304
		
2305	2306	2307
		
2308	2309	2310













		
2311	2312	2313
		
2314	2315	2316
		
2317	2318	2319
		
2320	2321	2322













		
2323	2324	2325
		
2326	2327	2328
		
2329	2330	2331
		
2332	2333	2334













		
2335	2336	2337
		
2338	2339	2340
		
2341	2342	2343
		
2344	2345	2346










		
2347	2348	2349
		
2350	2351	2352
		
2353	2354	2355
		
2356	2357	2358




		
2359	2360	2361
		
2362	2363	2364
		
2365	2366	2367
		
2368	2369	2370













		
2371	2372	2373
		
2374	2375	2376
		
2377	2378	2379
		
2380	2381	2382










		
2383	2384	2385
		
2386	2387	2388
		
2389	2390	2391
		
2392	2393	2394




		
2395	2396	2397
		
2398	2399	2400
		
2401	2402	2403
		
2404	2405	2406













		
2407	2408	2409
		
2410	2411	2412
		













2413	2414	2415
		
2416	2417	2418













		
2419	2420	2421
		
2422	2423	2424
		
2425	2426	2427
		
2428	2429	2430













		
2431	2432	2433
		
2434	2435	2436
		













2437	2438	2439
		
2440	2441	2442













		
2443	2444	2445
		
2446	2447	2448
		
2449	2450	2451
		
2452	2453	2454













		
2455	2456	2457
		
2458	2459	2460
		
2461	2462	2463
		
2464	2465	2466













		
2467	2468	2469
		
2470	2471	2472
		
2473	2474	2475
		
2476	2477	2478













		
2479	2480	2481
		
2482	2483	2484
		
2485	2486	2487
		
2488	2489	2490













		
2491	2492	2493
		
2494	2495	2496
		
2497	2498	2499
		
2500	2501	2502










		
2503	2504	2505
		
2506	2507	2508
		
2509	2510	2511
		
2512	2513	2514




		
2515	2516	2517
		
2518	2519	2520
		
2521	2522	2523
		
2524	2525	2526













		
2527	2528	2529
		
2530	2531	2532
		
2533	2534	2535
		
2536	2537	2538













		
2539	2540	2541
		
2542	2543	2544
		
2545	2546	2547
		
2548	2549	2550













		
2551	2552	2553
		
2554	2555	2556
		
2557	2558	2559
		
2560	2561	2562













		
2563	2564	2565
		
2566	2567	2568
		













2569	2570	2571
		













2572	2573	2574
		
2575	2576	2577
		
2578	2579	2580
		
2581	2582	2583
		










2584	2585	2586
		
2587	2588	2589
		
2590	2591	2592
		
2593	2594	2595
		




2596	2597	2598
		
2599	2600	2601
		
2602	2603	2604
		
2605	2606	2607
		













2608	2609	2610
		
2611	2612	2613
		
2614	2615	2616
		
2617	2618	2619
		










2620	2621	2622
		
2623	2624	2625
		
2626	2627	2628
		
2629	2630	2631
		




2632	2633	2634
		
2635	2636	2637
		
2638	2639	2640
		
2641	2642	2643
		










2644	2645	2646
		
2647	2648	2649
		
2650	2651	2652
		




2653	2654	2655
		










2656	2657	2658
		
2659	2660	2661
		
2662	2663	2664
		
2665	2666	2667
		




2668	2669	2670
		
2671	2672	2673
		
2674	2675	2676
		










2677	2678	2679
		




2680	2681	2682
		
2683	2684	2685
		
2686	2687	2688
		










2689	2690	2691
		




2692	2693	2694
		
2695	2696	2697
		
2698	2699	2700
		













2701	2702	2703
		









2704	2705	2706
		
2707	2708	2709
		
2710	2711	2712
		

2713	2714	2715
		

2716	2717	2718
		
2719	2720	2721
		
2722	2723	2724
		

2725	2726	2727
		

2728	2729	2730
		
2731	2732	2733
		
2734	2735	2736
		
2737	2738	2739
		

2740	2741	2742
		
2743	2744	2745
		
2746	2747	
		

303. A pharmaceutical composition comprising the compound of claim 302 and a pharmaceutically acceptable carrier.


304. The pharmaceutical composition of claim 303, further comprising one or more additional therapeutic agent(s).

305. The pharmaceutical composition of claim 304, wherein the additional therapeutic agent is selected from a mucolytic agent, bronchodilator, an antibiotic, an anti-infective agent, a CFTR modulator, or an anti-inflammatory agent.

306. The pharmaceutical composition of claim 304, wherein the additional therapeutic agent is a CFTR modulator.


307. The pharmaceutical composition of claim 304, wherein the additional therapeutic agent is a CFTR corrector.

308. The pharmaceutical composition of claim 304, wherein the additional therapeutic agent is




or pharmaceutically acceptable salt thereof.

309. The pharmaceutical composition of claim 304, wherein the additional therapeutic agent is



or pharmaceutically acceptable salt thereof.


310. The pharmaceutical composition of claim 304, wherein the additional therapeutic agent is



or pharmaceutically acceptable salt thereof.

311. The pharmaceutical composition of claim 304, wherein the additional therapeutic agent is a CFTR potentiator.



312. The pharmaceutical composition of claim 304, wherein the additional therapeutic agent is







,

or pharmaceutically acceptable salt thereof.

313. The pharmaceutical composition of claim 304, wherein the additional therapeutic agents are a CFTR corrector and a CFTR potentiator.

314. The pharmaceutical composition of claim 304, wherein the additional therapeutic agents are  and  , or pharmaceutically acceptable salts thereof.

315. The pharmaceutical composition of claim 304, wherein the additional therapeutic agents are  and  , or pharmaceutically acceptable salts thereof.

316. The pharmaceutical composition of claim 304, wherein the additional therapeutic agents are  and  , or pharmaceutically acceptable salts thereof.

317. A compound of claim 302 for use in treating cystic fibrosis in a patient.


318. The compound of claim 317, further comprising an additional therapeutic agent(s).

319. The compound of claim 318, wherein the additional therapeutic agent is selected from a mucolytic agent, bronchodilator, an antibiotic, an anti-infective agent, a CFTR modulator, or an anti-inflammatory agent.

320. The compound of claim 318, wherein the additional therapeutic agent is a CFTR modulator.

321. The compound of claim 318, wherein the additional therapeutic agent is a CFTR corrector.


322. The compound of claim 318, wherein the additional therapeutic agent is



,

or a pharmaceutically acceptable salt thereof.


323. The compound of claim 318, wherein the additional therapeutic agent is



,

or a pharmaceutically acceptable salt thereof.

324. The compound of claim 318, wherein the additional therapeutic agent is




,

or a pharmaceutically acceptable salt thereof.

325. The compound of claim 318, wherein the additional therapeutic agent is a CFTR potentiator.

326. The compound of claim 318, wherein the additional therapeutic agent is



or a pharmaceutically acceptable salt thereof.

327. The compound of any one of claims 317 to 326, wherein the patient is homozygous in the F508del mutation.

328. The compound of any one of claims 317 to 326, wherein the patient is heterozygous in the F508del mutation.

329. A kit comprising the compound of claim 303 or the pharmaceutical composition of any one of claims 303 to 316, and instructions for use therof.


330. The kit of claim 329, further comprising one or more additional therapeutic agent(s).

331. The kit of claim 330, wherein the additional therapeutic agent is selected from a mucolytic agent, bronchodilator, an antibiotic, an anti-infective agent, a CFTR modulator, or an anti-inflammatory agent.

332. The kit of claim 330, wherein the additional therapeutic agent is a CFTR modulator.


333. The kit of claim 330, wherein the additional therapeutic agent is a CFTR corrector.

334. The kit of claim 330, wherein the additional therapeutic agent is




or a pharmaceutically acceptable salt thereof.

335. The kit of claim 330, wherein the additional therapeutic agent is



or a pharmaceutically acceptable salt thereof.


336. The kit of claim 330, wherein the additional therapeutic agent is



or a pharmaceutically acceptable salt thereof.

337. The kit of claim 330, wherein the additional therapeutic agent is a CFTR potentiator.



338. The kit of claim 330, wherein the additional therapeutic agent is





or a pharmaceutically acceptable salt thereof.

339. The kit of claim 330, wherein the additional therapeutic agents are a CFTR corrector and a CFTR potentiator.



340. The kit of claim 330, wherein the additional therapeutic agents are


and

, or pharmaceutically acceptable salts thereof.

341. The kit of claim 330, wherein the additional therapeutic agents are


and

, or pharmaceutically acceptable salts thereof.

342. The kit of claim 330, wherein the additional therapeutic agents are


and

, or pharmaceutically acceptable salts thereof.


343. The kit of any one of claims 329 to 342, wherein the compound of claim 302 or the pharmaceutical composition of any one of claims 303 to 317 and the one or more additional therapeutic agent(s) are in separate containers.

344. The kit of any one of claims 329 to 342, wherein the compound of claim 302 or the pharmaceutical composition of any one of claims 303 to 317 and the one or more additional therapeutic agent(s) are in the same container.

345. The kit of claim 343 or 344, wherein the container is a bottle, vial, or blister pack, or combination thereof.

Abstract

The present invention features a compound of formula I:



I,

or a pharmaceutically acceptable salt thereof, where R_1, R_2, R_3, W, X, Y, Z, n, o, p, and q are defined herein, for the treatment of CFTR mediated diseases, such as cystic fibrosis. The present invention also features pharmaceutical compositions, method of treating, and kits thereof.

Figure 1

Mutation cDNA Name	Mutation Protein Name	Mutation Legacy Name
c.1A>G	p.? (unknown)	M1V
c.54-5940_273+10250del21kb	p.Ser18ArgfsX16	CFTRdele2,3
c.91C>T	p.Arg31Cys	R31C
c.115C>T	p.Gln39X	Q39X
c.137C>A	p.Ala46Asp	A46D
c.165-1G>A	No protein name	297-1G->A
c.166G>A	p.Glu56Lys	E56K
c.174_175insA	p.Arg59LysfsX10	306insA
c.178G>T	p.Glu60X	E60X
c.200C>T	p.Pro67Leu	P67L
c.220C>T	p.Arg74Trp	R74W
c.223C>T	p.Arg75X	R75X
c.224G>A	p.Arg75Gln	R75Q
c.254G>A	p.Gly85Glu	G85E
c.262_263delTT	p.Leu88IlefsX22	394delTT
c.273+1G>A	No protein name	405+1G->A
c.274-1G>A	No protein name	406-1G->A
c.274G>A	p.Glu92Lys	E92K
c.274G>T	p.Glu92X	E92X
c.292C>T	p.Gln98X	Q98X
c.313delA	p.Ile105SerfsX2	444delA
c.325_327delTATinsG	p.Tyr109GlyfsX4	457TAT->G
c.328G>C	p.Asp110His	D110H
c.349C>T	p.Arg117Cys	R117C
c.350G>A	p.Arg117His	R117H
c.366T>A	p.Tyr122X	Y122X
c.442delA	p.Ile148LeufsX5	574delA
c.443T>C	p.Ile148Thr	I148T
c.489+1G>T	No protein name	621+1G->T

Figure 1 Continued

Mutation cDNA Name	Mutation Protein Name	Mutation Legacy Name
c.531delT	p.Ile177MetfsX12	663delT
c.532G>A	p.Gly178Glu	G178R
c.543_546delTAGT	p.Leu183PhefsX5	675del4
c.579+1G>T	No protein name	711+1G->T
c.579+3A>G	No protein name	711+3A->G
c.579+5G>A	No protein name	711+5G->A
c.580-1G>T	No protein name	712-1G->T
c.595C>T	p.His199Tyr	H199Y
c.613C>T	p.Pro205Ser	P205S
c.617T>G	p.Leu206Trp	L206W
c.658C>T	p.Gln220X	Q220X
c.680T>G	p.Leu227Arg	L227R
c.720_741delAGGGAGAATGATGATGAAGTAC	p.Gly241GlufsX13	852del22
c.828C>A	p.Cys276X	C276X
c.948delT	p.Phe316LeufsX12	1078delT
c.988G>T	p.Gly330X	G330X
c.1000C>T	p.Arg334Trp	R334W
c.1007T>A	p.Ile336Lys	I336K
c.1013C>T	p.Thr338Ile	T338I
c.1021T>C	p.Ser341Pro	S341P
c.1022_1023insTC	p.Phe342HisfsX28	1154insTC
c.1040G>A	p.Arg347His	R347H
c.1040G>C	p.Arg347Pro	R347P
c.1055G>A	p.Arg352Gln	R352Q
c.[1075C>A;1079C>A]	p.[Gln359Lys;Thr360Lys]	Q359K/T360K

Figure 1 Continued

Mutation cDNA Name	Mutation Protein Name	Mutation Legacy Name
c.1081delT	p.Trp361GlyfsX8	1213delT
c.1116+1G>A	No protein name	1248+1G->A
c.1127_1128insA	p.Gln378AlafsX4	1259insA
c.1153_1154insAT	p.Asn386IlefsX3	1288insTA
c.1202G>A or c.1203G>A	p.Trp401X	W401X
c.1209+1G>A	No protein name	1341+1G->A
c.1210-12[5]	No protein name	5T
c.1210-12[7]	No protein name	7T
c.1240C>T	p.Gln414X	Q414X
c.1329_1330insAGAT	p.Ile444ArgfsX3	1461ins4
c.1340delA	p.Lys447ArgfsX2	1471delA
c.1364C>A	p.Ala455Glu	A455E
c.1393-1G>A	No protein name	1525-1G->A
c.1397C>A or c.1397C>G	p.Ser466X	S466X
c.1400T>C	p.Leu467Pro	L467P
c.1408A>G	p.Met470Val	M470V
c.1418delG	p.Gly473GlufsX54	1548delG
c.1466C>A	p.Ser489X	S489X
c.1475C>T	p.Ser492Phe	S492F
c.1477C>T	p.Gln493X	Q493X
c.1519_1521delATC	p.Ile507del	I507del
c.1521_1523delCTT	p.Phe508del	F508del
c.1545_1546delTA	p.Tyr515X	1677delTA
c.1558G>T	p.Val520Phe	V520F
c.1573C>T	p.Gln525X	Q525X
c.1585-8G>A	No protein name	1717-8G->A
c.1585-1G>A	No protein name	1717-1G->A
c.1624G>T	p.Gly542X	G542X
c.1645A>C or c.1647T>G	p.Ser549Arg	S549R

Figure 1 Continued

Mutation cDNA Name	Mutation Protein Name	Mutation Legacy Name
c.1646G>A	p.Ser549Asn	S549N
c.1650delA	p.Gly551ValfsX8	1782delA
c.1651G>A	p.Gly551Ser	G551S
c.1652G>A	p.Gly551Asp	G551D
c.1654C>T	p.Gln552X	Q552X
c.1657C>T	p.Arg553X	R553X
c.1673T>C	p.Leu558Ser	L558S
c.1675G>A	p.Ala559Thr	A559T
c.1679G>A	p.Arg560Lys	R560K
c.1679G>C	p.Arg560Thr	R560T
c.1679+1G>C	No protein name	1811+1G->C
c.1679+1.6kbA>G	No protein name	1811+1.6kbA->G
c.1680-1G>A	No protein name	1812-1G->A
c.1682C>A	p.Ala561Glu	A561E
c.1692delA	p.Asp565MetfsX7	1824delA
c.1705T>G	p.Tyr569Asp	Y569D
c.1727G>C	p.Gly576Ala	G576A
c.1736A>G	p.Asp579Gly	D579G
c.1753G>T	p.Glu585X	E585X
c.1766+1G>A	No protein name	1898+1G->A
c.1766+1G>C	No protein name	1898+1G->C
c.1766+3A>G	No protein name	1898+3A->G
c.1841A>G	p.Asp614Gly	D614G
c.1923_1931del9insA	p.Ser641ArgfsX5	2055del9->A

Figure 1 Continued

Mutation cDNA Name	Mutation Protein Name	Mutation Legacy Name
c.1973_1985del13insAGAAA	p.Arg658LysfsX4	2105-2117del13insAGAAA
c.1986_1989delAACT	p.Thr663ArgfsX8	2118del4
c.2002C>T	p.Arg668Cys	R668C
c.2012delT	p.Leu671X	2143delT
c.2051_2052delAAinsG	p.Lys684SerfsX38	2183AA->G*
c.2051_2052delAAinsG	p.Lys684SerfsX38	2183delAA->G*
c.2052_2053insA	p.Gln685ThrfsX4	2184insA
c.2052delA	p.Lys684AsnfsX38	2184delA
c.2125C>T	p.Arg709X	R709X
c.2128A>T	p.Lys710X	K710X
c.2175_2176insA	p.Glu726ArgfsX4	2307insA
c.2195T>G	p.Leu732X	L732X
c.2215delG	p.Val739TyrfsX16	2347delG
c.2260G>A	p.Val754Met	V754M
c.2290C>T	p.Arg764X	R764X
c.2353C>T	p.Arg785X	R785X
c.2374C>T	p.Arg792X	R792X
c.2424_2425insAT	p.Ser809IlefsX13	2556insAT
c.2453delT	p.Leu818TrpfsX3	2585delT
c.2462_2463delGT	p.Ser821ArgfsX4	No legacy name
c.2464G>T	p.Glu822X	E822X
c.2490+1G>A	No protein name	2622+1G->A
c.2491G>T	p.Glu831X	E831X
c.2537G>A or c.2538G>A	p.Trp846X	W846X
c.2547C>A	p.Tyr849X	Y849X
c.2551C>T	p.Arg851X	R851X
c.2583delT	p.Phe861LeufsX3	2711delT

Figure 1 Continued

Mutation cDNA Name	Mutation Protein Name	Mutation Legacy Name
c.2657+2_2657+3insA	No protein name	2789+2insA
c.2657+5G>A	No protein name	2789+5G->A
c.2658-1G>C	No protein name	2790-1G->C
c.2668C>T	p.Gln890X	Q890X
c.2735C>A	p.Ser912X	S912X
c.2737_2738insG		2869insG
c.2739T>A	p.Tyr913X	Y913X
c.2764_2765insAG	p.Val922GlufsX2	2896insAG
c.2780T>C	p.Leu927Pro	L927P
c.2834C>T	p.Ser945Leu	S945L
c.2875delG	p.Ala959HisfsX9	3007delG
c.2908G>C	p.Gly970Arg	G970R
c.2930C>T	p.Ser977Phe	S977F
c.2988G>A	No protein name	3120G->A
c.2988+1G>A	No protein name	3120+1G->A
c.2989-977_3367+248del	No protein name	3121-977_3499+248del2515
c.2989-1G>A	No protein name	3121-1G->A
c.2991G>C	p.Leu997Phe	L997F
c.3002_3003delTG	p.Val1001AspfsX45	3132delTG
c.3080T>C	p.Ile1027Thr	I1027T
c.3140-26A>G	No protein name	3272-26A->G
c.3154T>G	p.Phe1052Val	F1052V
c.3160C>G	p.His1054Asp	H1054D
c.3181G>C	p.Gly1061Arg	G1061R
c.3194T>C	p.Leu1065Pro	L1065P
c.3196C>T	p.Arg1066Cys	R1066C
c.3197G>A	p.Arg1066His	R1066H

Figure 1 Continued

Mutation cDNA Name	Mutation Protein Name	Mutation Legacy Name
c.3205G>A	p.Gly1069Arg	G1069R
c.3208C>T	p.Arg1070Trp	R1070W
c.3209G>A	p.Arg1070Gln	R1070Q
c.3222T>A	p.Phe1074Leu	F1074L
c.3230T>C	p.Leu1077Pro	L1077P
c.3266G>A	p.Trp1089X	W1089X
c.3276C>A or c.3276C>G	p.Tyr1092X	Y1092X
c.3302T>A	p.Met1101Lys	M1101K
c.3310G>T	p.Glu1104X	E1104X
c.3454G>C	p.Asp1152His	D1152H
c.3472C>T	p.Arg1158X	R1158X
c.3484C>T	p.Arg1162X	R1162X
c.3485G>T	p.Arg1162Leu	R1162L
c.3528delC	p.Lys1177SerfsX15	3659delC
c.3535_3536insTCAA	p.Thr1179IlefsX17	3667ins4
c.3587C>G	p.Ser1196X	S1196X
c.3605delA	p.Asp1202AlafsX9	3737delA
c.3611G>A or c.3612G>A	p.Trp1204X	W1204X
c.3659delC	p.Thr1220LysfsX8	3791delC
c.3691delT	p.Ser1231ProfsX4	3821delT
c.3700A>G	p.Ile1234Val	I1234V
c.3705T>G	p.Ser1235Arg	S1235R
c.3717+12191C>T	No protein name	3849+10kbC->T

Figure 1 Continued

Mutation cDNA Name	Mutation Protein Name	Mutation Legacy Name
c.3718-1G>A	No protein name	3850-1G->A
c.3731G>A	p.Gly1244Glu	G1244E
c.3744delA	p.Lys1250ArgfsX9	3876delA
c.3752G>A	p.Ser1251Asn	S1251N
c.3763T>C	p.Ser1255Pro	S1255P
c.3764C>A	p.Ser1255X	S1255X
c.3773_3774insT	p.Leu1258PhefsX7	3905insT
c.3808G>A	p.Asp1270Asn	D1270N
c.3846G>A	p.Trp1282X	W1282X
c.3873+1G>A	No protein name	4005+1G->A
c.3883delA	p.Ile1295PhefsX33	4015delA
c.3884_3885insT	p.Ser1297PhefsX5	4016insT
c.3909C>G	p.Asn1303Lys	N1303K
c.3937C>T	p.Gln1313X	Q1313X
c.3964-78_4242+577del	NULL	CFTRdele22,23
c.4046G>A	p.Gly1349Asp	G1349D
c.4077_4080delTGTTinsAA	No protein name	4209TGTT->AA
c.4111G>T	p.Glu1371X	E1371X
c.4196_4197delTC	p.Cys1400X	4326delTC
c.4234C>T	p.Gln1412X	Q1412X
c.4242+1G>T	No protein name	4374+1G->T
c.4251delA	p.Glu1418ArgfsX14	4382delA
c.4296_4297insGA	p.Ser1435GlyfsX14	4428insGA