

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 August 2011 (25.08.2011)

(10) International Publication Number
WO 2011/103524 A2

(51) International Patent Classification:

C07D 487/14 (2006.01) *A61K 31/4025* (2006.01)
C07D 403/12 (2006.01) *A61P 37/00* (2006.01)
C07D 403/06 (2006.01) *A61P 29/00* (2006.01)
C07D 403/14 (2006.01) *A61P 37/08* (2006.01)
A61K 31/407 (2006.01)

[PL/PL]; Kotarbinskiego, 46, PL-91-163 Lodz (PL).
KACZMAREK, Krzysztof [PL/PL]; Michala, 8, 9,
PL-93-324 Lodz (PL).

(21) International Application Number:

PCT/US2011/025571

(22) International Filing Date:

19 February 2011 (19.02.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

P.390493 19 February 2010 (19.02.2010) PL

(74) Agent: FEIGELSON, Daniel; 7 Oppenheimer St., 76701 Rehovot (IL).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: CYCLIC TETRAPEPTIDES AND THERAPEUTIC APPLICATIONS THEREOF



(57) Abstract: There are provided compounds of formula I wherein k, m, n, p, R, R', R'', R'''', R3 and R4 are as defined in the application. Other embodiments are also disclosed.

Cyclic tetrapeptides and therapeutic applications thereof

BACKGROUND

[0001] Immunosuppressive drugs are commonly used in transplantation and in treatment of autoimmune diseases. Production of these drugs is expensive, and the most frequently used of these drugs, namely cyclosporine A, tacrolimus and rapamycin, exhibit undesirable side-effects. The search for new immunosuppressive drugs devoid of side-effects, particularly in the class of natural peptide immunoregulators and their analogues, represents a serious challenge for medicinal chemistry.

[0002] Cyclolinopeptide A (CLA), a very hydrophobic cyclic nonapeptide, was first isolated from linen seeds in 1959. CLA is strongly immunosuppressive, with a potency comparable to that of cyclosporine A (CsA). The mechanism of action of CLA was shown to be similar to that of CsA, i.e. CLA formed a complex with cyclophilin A causing inactivation of calcineurin, albeit at much lower affinity (Gaymes et al., *Febs Lett*, 1997, 418, 224-227). CLA inhibited both humoral and cellular immune response and graft-versus-host reaction; prolonged survival of allogeneic skin grafts; tempered post-adjuvant polyarthritis in rats and hemolytic anemia of New Zealand Black mice; and, similarly to CsA, inhibited IL1 and IL-2 production. Unfortunately, the high hydrophobicity of CLA presents an obstacle for the potential application of the compound in therapy.

[0003] Linear CLA analogues containing alanine residue in successive positions of the peptide chain were found to be immunosuppressive (Wieczorek et al., *Arch Immunol Ther Exp*, **1992**, 40, 213-216). It was also found that the activity of linear CLA analogues gradually decreased with shortening of the peptide chain from the N-terminus, at the same time showing an increase of activity for C-terminal tetra- and tripeptides (Siemion et al., *Arch Immunol Ther Exp*, **1994**, 42, 459-465). The introduction of a single, hydrophilic threonine residue into the CLA molecule did not result in improved solubility in water. However, an improvement in solubility was achieved by the introduction of a sulphonic group in the *para*-position of the phenyl ring of one or two phenylalanine residues, without loss of biological activity (Siemion et al., *Arch Immunol Ther Exp*, **1992**, 40, 257-261; Cebrat et al., *J Peptide Res.*, **1997**, 49, 415-420). In addition, it has been observed that the inclusion of tetrapeptidic (Pro-Pro-Phe-Phe) or tripeptidic (Pro-Phe-Phe) fragments in longer linear peptides chains seem to have significance for immunosuppressive activity (Wieczorek et al., *Arch Immunol Ther Exp*, **1993**, 41, 291-296; Cebrat et al., *Pol. J Chem.*, **1997**, 71, 1401).


[0004] A series of analogues in which the *cis*-peptide bond between proline residues was replaced with 1,5-disubstituted tetrazole ring (a good mimetic of amide bonds in *cis* configuration) showed immunosuppressive activity comparable to CsA. (Karczmarek et al., *Biopolymers*, **2002**, 63, 343-357).

[0005] Synthetic CLA analogues in which leucine residues in position 5 and/or 8 were replaced with their hydroxymethyl analogue displayed a four-fold increase in solubility in water in comparison to CLA, but also showed a 25% diminution in biological activity compared to native CLA (Zubrzak et al., *Biopolymers (Peptide Science)*, **2005**, 80, 347-356).

[0006] A series of nine CLA analogues was obtained by replacement of CLA proline residues with β^2 -isoproline and β^3 -homoproline. In comparison to CsA, these CLA analogues displayed strong inhibitory properties in the cellular immune response. The majority of these analogs were practically devoid of cell toxicity (Katarzyńska et al., *J Pept Sci*, **2009**, 14, 1283-1294).

BRIEF DESCRIPTION OF THE INVENTION

[0007] There are provided in accordance with an embodiment of the present invention compounds having the formula I:



wherein

[0008] k, m, n and p are each independently 0, 1 or 2;

[0009] R and R' are each independently selected from H and C₁₋₃ alkyl, or, when taken together, R and R' are -CR¹R^{1'}-X-CH₂-, wherein CR¹R^{1'} is attached to the backbone nitrogen, R¹ and R^{1'} are each independently selected from H and C₁₋₃ alkyl, and X is selected from -CH₂- , -CH₂CH₂- , -CH(OH)-, -O-, -S- and -NH-;

[0010] R'' and R''' are each independently selected from H and C₁₋₃ alkyl, or, when taken together, R'' and R''' are -CR²R^{2'}-X'-CH₂-, wherein CR²R^{2'} is attached to the backbone nitrogen, R² and R^{2'} are each independently selected from H and C₁₋₃ alkyl, and X' is selected from -CH₂- , -CH₂CH₂- , -CH(OH)-, -O-, -S- and -NH-; and

[0011] R³ and R⁴ are each independently selected from aryl, substituted aryl, heteroaryl and substituted heteroaryl;


[0012] or a pharmaceutically acceptable salt thereof.

[0013] In some embodiments, at least one of R³ and R⁴ is phenyl. In some embodiments, at least one of R³ and R⁴ is 4-hydroxyphenyl. In some embodiments, at least one of R³ and R⁴ is 4-t-butoxyphenyl. In some embodiments, at least one of R³ and R⁴ is 2-indolyl. In some embodiments, R³ and R⁴ are both phenyl. In some embodiments, one of R³ and R⁴ is phenyl and the other of R³ and R⁴ is 4-hydroxyphenyl. In some embodiments, one of R³ and R⁴ is phenyl and the other of R³ and R⁴ is 4-t-butoxyphenyl. In some embodiments, one of R³ and R⁴ is phenyl and the other of R³ and R⁴ is 2-indolyl. In some embodiments, the carbon to which -CH₂-R³ is attached has absolute (R)-stereochemistry. In some embodiments, the carbon to which -CH₂-R³ is attached has absolute (S)-stereochemistry. In some embodiments, the carbon to which -CH₂-R⁴ is attached has absolute (R)-stereochemistry. In some embodiments, the carbon to which -CH₂-R⁴ is attached has absolute (S)-stereochemistry. In some embodiments, one of k, m, n and p is 1, and the remainder of k, m, n and p are 0. In some embodiments, two of k, m, n and p are 1, and the remainder of k, m, n and p are 0. In some embodiments, at least one of k and m is not 0. In some embodiments, at least one of n and p is not 0. In some embodiments, at least one of k and m is not 0 and at least one of n and p is not 0. In some embodiments, both k and n are 0. In some embodiments, both k and n are 0, one of m and p is 0, and the other of m and p is 1. In some embodiments, both k and n are 0 and both m and p are 1. In some embodiments, all four amino acids are L-amino acids. In some embodiments, three of the amino acids are L-amino acids and one of the amino acids is a D-amino acid. In some embodiments, two of the amino acids are L-amino acids and two of the amino acids are D-amino acids. In some embodiments, one of the amino acids is an L-amino acid and three of the amino acids are D-amino acids. In some embodiments, all four amino acids are D-amino acids.


[0014] In some embodiments, R and R' are taken together to form -(CH₂)₃-, i.e. R and R' are taken together to form -CR¹R¹'-X-CH₂- wherein R¹ and R¹' are both H and X is CH₂. In some embodiments, R'' and R''' are taken together to form -(CH₂)₃-, i.e. R'' and R''' are taken together to form -CR²R²'-X'-CH₂- wherein R² and R²' are both H and X' is CH₂. In some embodiments, R and R' are taken together to form -(CH₂)₄-, i.e. R and R' are taken together to form -CR¹R¹'-X-CH₂- wherein R¹ and R¹' are both H and X is (CH₂)₂. In some embodiments, R'' and R''' are taken together to form -(CH₂)₄-, i.e. R'' and R''' are taken together to form -CR²R²'-X-CH₂- wherein R² and R²' are both H and X is (CH₂)₂. In some embodiments, R and

R' are taken together to form -CH₂-CH(OH)-CH₂-, i.e. R and R' are taken together to form -CR¹R¹-X-CH₂- wherein R¹ and R¹' are both H and X is CH(OH). In some embodiments, R'' and R''' are taken together to form -CH₂-CH(OH)-CH₂-, i.e. R'' and R''' are taken together to form -CR²R²-X'-CH₂- wherein R² and R²' are both H and X' is CH(OH). In some embodiments in which R and R' are taken together, the carbon at which R' is attached has absolute (S)-stereochemistry. In some embodiments in which R and R' are taken together, the carbon at which R' is attached has absolute (R)-stereochemistry. In some embodiments in which R'' and R''' are taken together, the carbon at which R''' is attached has absolute (S)-stereochemistry. In some embodiments in which R'' and R''' are taken together, the carbon at which R''' is attached has absolute (R)-stereochemistry.


[0015] In some embodiments, the compound is selected from the group consisting of:




I-1 (cyclic(-Pro-Pro- β^3 hoPhe-Phe-)),






I-2 (cyclic(-Pro-Pro-Phe- β^3 hoPhe-)),



I-3 (cyclic(-Pro-Pro- β^3 hoPhe- β^3 hoPhe-)),







I-4 (cyclic(-Pro-Pro- β^3 hoPhe-Trp-)),

**I-5** (cyclic(-Pro-Pro- β^3 hoPhe-Tyr-)),**I-6** (cyclic(-Pro-Pro- β^3 hoPhe-Tyr(t-Bu-))),**I-7** (cyclic(-Pro-Pip- β^3 hoPhe-Phe-)) and **I-8** (cyclic(Pro-Hyp- β^3 hoPhe-Phe-)).



In some embodiments, the compound is a compound of formula I-1. In some embodiments, the compound is a compound of formula I-2. In some embodiments, the compound is a compound of formula I-3. In some embodiments, the compound is a compound of formula I-4. In some embodiments, the compound is a compound of formula I-5. In some embodiments, the compound is a compound of formula I-6. In some embodiments, the compound is a compound of formula I-7. In some embodiments, the compound is a compound of formula I-8.

[0016] In some embodiments, the compound is selected from the group consisting of:



((cyclic(-(L)-Pro-(L)-Pro-(L)- β^3 hoPhe-(L)-Phe-)),((cyclic(-(L)-Pro-(L)-Pro-(L)-Phe-(L)- β^3 hoPhe-))),



(cyclic(-(L)-Pro-(L)-Pro-(L)- β^3 hoPhe-(L)- β^3 hoPhe-)), (cyclic(-(L)-Pro-(L)-Pro-(L)- β^3 HoPhe-(D)-Phe-)),





(cyclic-(D)-Pro-(D)-Pro-(L)- β^3 HoPhe-(D)-Phe-), (cyclic-(D)-Pro-(L)-Pro-(L)- β^3 HoPhe-(L)-Phe-)),




(cyclic-(L)-Pro-(L)-Pro-(D)- β^3 HoPhe-(L)-Phe-),

(cyclic-(L)-Pro-(D)-Pro-(L)- β^3 HoPhe-(L)-Phe-),




(cyclic-(D)-Pro-(D)-Pro-(D)- β^3 HoPhe-(D)-Phe-),


(cyclic-(D)-Pro-(D)-Pro-(L)- β^3 HoPhe-(L)-Phe-),




(cyclic(-(L)-Pro-(L)-Pro-(L)- β^3 HoPhe-(L)-Trp-)),




(c(-(L)-Pro-(L)-Pro-(L)- β^3 HoPhe-(L)-Tyr-)),



(cyclic[-(L)-Pro-(L)-Pro-(L)- β^3 HoPhe-(L)-Tyr(tBu)-]),



(cyclic(-(L)-Pro-(L)-Pip-(L)- β^3 HoPhe-(L)-Phe-)),



and

(cyclic(-(L)-Pro-(L)-*t*-Hyp-(L)- β^3 HoPhe-(L)-Phe-)). In some

embodiments, the compound is a compound of formula I-A. In some embodiments, the compound is a compound of formula I-B. In some embodiments, the compound is a compound of formula I-C. In some embodiments, the compound is a compound of formula I-D. In some embodiments, the compound is a compound of formula I-E. In some embodiments, the compound is a compound of formula I-F. In some embodiments, the compound is a compound of formula I-G. In some embodiments, the compound is a compound of formula I-H. In some embodiments, the compound is a compound of formula I-I. In some embodiments, the compound is a compound of formula I-J. In some embodiments, the compound is a compound of formula I-K. In some embodiments, the compound is a compound of formula I-L. In some embodiments, the compound is a compound of formula I-M. In some embodiments, the compound is a compound of formula I-N. In some embodiments, the compound is a compound of formula I-O. In some embodiments, the compound is a compound of formula I-P.

[0017] In some embodiments, one or more amino groups in the compound of Formula I are in protected form.

[0018] There is also provided, in accordance with an embodiment of the invention, a pharmaceutical composition comprising a compound of formula I and a pharmaceutically acceptable carrier, excipient or diluent therefor. In some embodiments, the compound is a compound of formula I-1. In some embodiments, the compound is a compound of formula I-2. In some embodiments, the compound is a compound of formula I-3. In some embodiments, the compound is a compound of formula I-4. In some embodiments, the compound is a compound of formula I-5. In some embodiments, the compound is a compound of formula I-6. In some embodiments, the compound is a compound of formula I-7. In some embodiments, the compound is a compound of formula I-8. In some embodiments, the compound is a compound of formula I-A. In some embodiments, the compound is a compound of formula I-B. In some embodiments, the compound is a compound of formula I-C. In some embodiments, the compound is a compound of formula I-D. In some embodiments, the compound is a compound of formula I-E. In some embodiments, the compound is a compound of formula I-F. In some embodiments, the compound is a compound of formula I-G. In some embodiments, the compound is a compound of formula I-H. In some embodiments, the compound is a compound of formula I-I. In some embodiments, the compound is a compound of formula I-J. In some embodiments, the compound is a compound of formula I-K. In some embodiments, the compound is a compound of formula I-L. In some embodiments, the compound is a compound of formula I-M. In some embodiments, the compound is a compound of formula I-N. In some embodiments, the compound is a compound of formula I-O. In some embodiments, the compound is a compound of formula I-P.

[0019] There is also provided, in accordance with an embodiment of the invention, a method of suppressing immune response in a patient, comprising administering to a patient in need thereof an efficacious amount of a compound of formula I. In some embodiments, the immune response which is suppressed is inflammation. In some embodiments the immune response which is suppressed is transplant rejection. In some embodiments, the compound is a compound of formula I-1. In some embodiments, the compound is a compound of formula I-2. In some embodiments, the compound is a compound of formula I-3. In some embodiments, the compound is a compound of formula I-4. In some embodiments, the compound is a compound of formula I-5. In some embodiments, the compound is a compound of formula I-6. In some embodiments, the compound is a compound of formula I-7. In some embodiments, the compound is a compound of formula I-8. In some embodiments, the compound is a compound

of formula I-A. In some embodiments, the compound is a compound of formula I-B. In some embodiments, the compound is a compound of formula I-C. In some embodiments, the compound is a compound of formula I-D. In some embodiments, the compound is a compound of formula I-E. In some embodiments, the compound is a compound of formula I-F. In some embodiments, the compound is a compound of formula I-G. In some embodiments, the compound is a compound of formula I-H. In some embodiments, the compound is a compound of formula I-I. In some embodiments, the compound is a compound of formula I-J. In some embodiments, the compound is a compound of formula I-K. In some embodiments, the compound is a compound of formula I-L. In some embodiments, the compound is a compound of formula I-M. In some embodiments, the compound is a compound of formula I-N. In some embodiments, the compound is a compound of formula I-O. In some embodiments, the compound is a compound of formula I-P.


[0020] There is also provided, in accordance with an embodiment of the invention, a method of treating or preventing an immune-mediated disease or condition in a patient, comprising administering to a patient in need thereof an efficacious amount of a compound of formula I. There is also provided, in accordance with an embodiment of the invention, a method for lowering the toxicity profile of a second drug, comprising administering a compound of formula I in conjunction with the second drug. In some embodiments, the immune-mediated disease or condition is selected from the group consisting of auto-immune diseases, inflammation processes, transplant rejection, and allergic reactions. In some embodiments, the immune-mediated disease or condition is selected from Psoriasis, lichen planus and other papulosquamous disorders. In some embodiments, the immune-mediated disease or condition is selected from eczema and dermatitis. In some embodiments, the eczema or dermatitis is selected from eczema, atopic eczema, seborrheic dermatitis, and pompholyx. In some embodiments, the immune-mediated disease or condition is a skin reaction to sunlight. In some embodiments, the immune-mediated disease or condition selected from non-specific skin irritation and insect bite. In some embodiments, the immune-mediated disease or condition is a urticaria. In some embodiments, the immune-mediated disease or condition is selected from the group consisting of a primary skin tumor (e.g. melanoma); rheumatoid arthritis (both autoimmune and elicited by infection); Crohn's disease; inflammatory bowel disease; irritable bowel syndrome; a neurodegenerative disease (e.g. multiple sclerosis); Parkinson's disease; Graft-versus-Host reaction; severe psoriasis; and atopic dermatitis. In some embodiments, the compound of formula I is administered in conjunction with a chemotherapeutic drug in order to reduce the toxic effects of the chemotherapeutic drug. In some embodiments, the compound

is a compound of formula I-1. In some embodiments, the compound is a compound of formula I-2. In some embodiments, the compound is a compound of formula I-3. In some embodiments, the compound is a compound of formula I-4. In some embodiments, the compound is a compound of formula I-5. In some embodiments, the compound is a compound of formula I-6. In some embodiments, the compound is a compound of formula I-7. In some embodiments, the compound is a compound of formula I-8. In some embodiments, the compound is a compound of formula I-A. In some embodiments, the compound is a compound of formula I-B. In some embodiments, the compound is a compound of formula I-C. In some embodiments, the compound is a compound of formula I-D. In some embodiments, the compound is a compound of formula I-E. In some embodiments, the compound is a compound of formula I-F. In some embodiments, the compound is a compound of formula I-G. In some embodiments, the compound is a compound of formula I-H. In some embodiments, the compound is a compound of formula I-I. In some embodiments, the compound is a compound of formula I-J. In some embodiments, the compound is a compound of formula I-K. In some embodiments, the compound is a compound of formula I-L. In some embodiments, the compound is a compound of formula I-M. In some embodiments, the compound is a compound of formula I-N. In some embodiments, the compound is a compound of formula I-O. In some embodiments, the compound is a compound of formula I-P.

[0021] There is also provided, in accordance with an embodiment of the invention, a kit, comprising a compound of formula I and instructions for using the compound to (a) suppress an immune response in a patient, (b) treat or prevent an immune-mediated disease or condition in a patient, or (c) lower the toxicity profile of a second drug. In some embodiments, the immune response is inflammation. In some embodiments the immune response is transplant rejection. In some embodiments, the immune-mediated disease or condition is selected from the group consisting of auto-immune diseases, inflammation processes, transplant rejection, allergic reactions. In some embodiments, the immune-mediated disease or condition is selected from Psoriasis, lichen planus and other papulosquamous disorders. In some embodiments, the immune-mediated disease or condition is selected from eczema and dermatitis. In some embodiments, the eczema or dermatitis is selected from eczema, atopic eczema, seborrheic dermatitis, and pompholyx. In some embodiments, the immune-mediated disease or condition is a skin reaction to sunlight. In some embodiments, the immune-mediated disease or condition selected from non-specific skin irritation and insect bite. In some embodiments, the immune-mediated disease or condition is a urticaria. In some embodiments, the immune-mediated disease or condition is selected from the group consisting of a primary skin tumor

(e.g. melanoma); rheumatoid arthritis (both autoimmune and elicited by infection); Crohn's disease; inflammatory bowel disease; irritable bowel syndrome; a neurodegenerative disease (e.g. multiple sclerosis); Parkinson's disease; Graft-versus-Host reaction; severe psoriasis; and atopic dermatitis. In some embodiments, the instructions instruct administering the compound of formula I in conjunction with a chemotherapeutic drug in order to reduce the toxic effects of the chemotherapeutic drug. In some embodiments, the compound is a compound of formula I-1. In some embodiments, the compound is a compound of formula I-2. In some embodiments, the compound is a compound of formula I-3. In some embodiments, the compound is a compound of formula I-4. In some embodiments, the compound is a compound of formula I-5. In some embodiments, the compound is a compound of formula I-6. In some embodiments, the compound is a compound of formula I-7. In some embodiments, the compound is a compound of formula I-8. In some embodiments, the compound is a compound of formula I-A. In some embodiments, the compound is a compound of formula I-B. In some embodiments, the compound is a compound of formula I-C. In some embodiments, the compound is a compound of formula I-D. In some embodiments, the compound is a compound of formula I-E. In some embodiments, the compound is a compound of formula I-F. In some embodiments, the compound is a compound of formula I-G. In some embodiments, the compound is a compound of formula I-H. In some embodiments, the compound is a compound of formula I-I. In some embodiments, the compound is a compound of formula I-J. In some embodiments, the compound is a compound of formula I-K. In some embodiments, the compound is a compound of formula I-L. In some embodiments, the compound is a compound of formula I-M. In some embodiments, the compound is a compound of formula I-N. In some embodiments, the compound is a compound of formula I-O. In some embodiments, the compound is a compound of formula I-P.









[0022] There is also provided, in accordance with an embodiment of the invention, a method for making a compound of formula I, comprising cyclizing a compound having the formula II-1, II-2, II-3 or II-4, wherein R, R', R'', R''', R³, R⁴, k, m, n and p are as defined in formula I (collectively referred to hereinafter as compounds of formula II):





















to the corresponding compound of formula I. In some embodiments, the method further comprises synthesizing the compound of formula II-1, II-2, II-3 or II-4. In some embodiments, the compound of formula II-1, II-2, II-3 or II-4 is formed by solid-phase synthesis.







[0023] In accordance with embodiments of the invention, there are also provided compounds of formulae II-1, II-2, II-3 and II-4 *per se*, as well as protected versions of these compounds (e.g. in which one or more amino groups, such as the N-terminal amino group or a side chain amino group, are protected, e.g. by tert-butoxycarbonyl) and these compounds, in protected or unprotected form, when bound to a solid-phase resin. Hereinafter, unless specified otherwise or illogical in the given context, when reference is made to a compound of formula II or a sub-genus or sub-species thereof, such reference is intended to include such compound in a form in which it is (a)(i) at least partly protected or (a)(ii) completely un- or deprotected; (b)(i) bound to a resin (1) directly or (2) through a linker, or (b)(ii) not bound to a resin; or a combination of conditions (a) and (b). Furthermore, as depicted herein, for the sake of convenience, free, non-protected linear peptides are shown as neutral molecules, viz. having H₂N- at the N-terminus and -COOH at the C-terminus; however, it will be appreciated that the actual charge on these moieties, as well as on any ionizable side chain moieties (e.g. carboxylic acid or amine moieties in the side chains) will depend on the pH of surroundings, and will not necessarily be as shown.

[0024] In some embodiments, the compound of formula II is selected from the group consisting of:

II-4-a (H-Pro- β^3 hoPhe-Phe-Pro-OH),II-1-a (H-Pro-Pro- β^3 hoPhe-Phe-OH),II-2-a (H₂N-Phe-Pro-Pro- β^3 hoPhe-OH),II-3-a (H₂N- β^3 hoPhe-Phe-Pro-Pro-OH),II-4-b (H-Pro-Pro-Phe- β^3 hoPhe-OH),II-1-b (H-Pro-Pro-Phe- β^3 hoPhe-OH),

II-2-b ($\text{H}_2\text{N}-\beta^3\text{hoPhe-Pro-Pro-Phe-OH}$),**II-3-b** ($\text{H}_2\text{N-Phe}-\beta^3\text{hoPhe-Pro-Pro-OH}$),**II-4-c** ($\text{H-Pro}-\beta^3\text{hoPhe}-\beta^3\text{hoPhe-Pro-OH}$),**II-1-c** ($\text{H-Pro-Pro}-\beta^3\text{hoPhe}-\beta^3\text{hoPhe-OH}$),**II-2-c** ($\text{H}_2\text{N}-\beta^3\text{hoPhe-Pro-Pro}-\beta^3\text{hoPhe-OH}$), **II-3-c** ($\text{H}_2\text{N}-\beta^3\text{hoPhe}-\beta^3\text{hoPhe-Pro-Pro-OH}$),**II-4-d** ($\text{H-Pro-Pro}-\beta^3\text{hoPhe-Trp-OH}$),**II-1-d** ($\text{H-Pro-Pro}-\beta^3\text{hoPhe-Trp-OH}$),**II-2-d** ($\text{H-Pro-Pro}-\beta^3\text{hoPhe-Trp-OH}$),**II-3-d** ($\text{H}_2\text{N}-\beta^3\text{hoPhe-Trp-Pro-Pro-OH}$),


**II-4-e** (H-Pro- β^3 hoPhe-Tyr-Pro-OH),**II-1-e** (H-Pro-Pro- β^3 hoPhe-Tyr-OH),**II-2-e** (H₂N-Tyr-Pro-Pro- β^3 hoPhe-OH),**II-3-e** (H₂N- β^3 hoPhe-Tyr-Pro-Pro-OH),**II-4-f** (H-Pro- β^3 hoPhe-Tyr(t-Bu)-Pro-OH),**II-1-f** (H-Pro-Pro- β^3 hoPhe-Tyr(t-Bu)-OH),**II-2-f** (H₂N-Tyr(t-Bu)-Pro-Pro- β^3 hoPhe-OH),**II-3-f** (H₂N- β^3 hoPhe-Tyr(t-Bu)-Pro-Pro-OH),




II-4-g (H-Pip- β^3 hoPhe-Phe-Pro-OH),**II-1-g** (H-Pro-Pip- β^3 hoPhe-Phe-OH),**II-2-g** (H₂N-Phe-Pro-Pip- β^3 hoPhe-OH),**II-3-g** (H₂N- β^3 hoPhe-Phe-Pro-Pip-OH),**II-4-h** (H-Hyp- β^3 hoPhe-Phe-Pro-OH) **II-1-h** (H-Pro-Hyp- β^3 hoPhe-Phe-OH)**II-2-h** (H₂N-Phe-Pro-Hyp- β^3 hoPhe-OH), and**II-3-h** (H₂N- β^3 hoPhe-Phe-Pro-Hyp-OH).


In some embodiments the compound is a compound of formula II-4-a. In some embodiments, the compound is a compound of formula II-1-a. In some embodiments, the compound is a compound of formula II-2-a. In some embodiments, the compound is a compound of formula II-3-a. In some embodiments, the compound is a compound of formula II-4-b. In some embodiments, the compound is a compound of formula II-1-b. In some embodiments, the compound is a compound of formula II-2-b. In some embodiments, the compound is a compound of formula II-3-b. In some embodiments, the compound is a compound of formula II-4-c. In some embodiments, the compound is a compound of formula II-1-c. In some embodiments, the compound is a compound of formula II-2-c. In some embodiments, the compound is a compound of formula II-3-c. In some embodiments, the compound is a compound of formula II-4-d. In some embodiments, the compound is a compound of formula

II-1-d. In some embodiments, the compound is a compound of formula II-2-d. In some embodiments, the compound is a compound of formula II-3-d. In some embodiments, the compound is a compound of formula II-4-e. In some embodiments, the compound is a compound of formula II-1-e. In some embodiments, the compound is a compound of formula II-2-e. In some embodiments, the compound is a compound of formula II-3-e. In some embodiments, the compound is a compound of formula II-4-f. In some embodiments, the compound is a compound of formula II-1-f. In some embodiments, the compound is a compound of formula II-2-f. In some embodiments, the compound is a compound of formula II-3-f. In some embodiments, the compound is a compound of formula II-4-g. In some embodiments, the compound is a compound of formula II-1-g. In some embodiments, the compound is a compound of formula II-2-g. In some embodiments, the compound is a compound of formula II-3-g. In some embodiments, the compound is a compound of formula II-4-h. In some embodiments, the compound is a compound of formula II-1-h. In some embodiments, the compound is a compound of formula II-2-h. In some embodiments, the compound is a compound of formula II-3-h.

[0025] In some embodiments the compound of formula II is selected from the group consisting of:









II-C-1

II-C-2




II-C-3

II-C-4




II-D-1

II-D-2




II-D-3

II-D-4




II-E-1

II-E-2




II-E-3

II-E-4




II-F-1

II-F-2




II-F-3

II-F-4




II-G-1

II-G-2




II-G-3

II-G-4




II-H-1


II-H-1



II-H-3


II-H-4






II-L-1

II-L-1'




II-L-3

II-L-4




II-M-1


II-M-2



II-M-3

II-M-4





[0026] In some embodiments, the compound is a compound of formula II-A-1. In some embodiments, the compound is a compound of formula II-A-2. In some embodiments, the compound is a compound of formula II-A-3. In some embodiments, the compound is a compound of formula II-A-4. In some embodiments, the compound is a compound of formula II-B-1. In some embodiments, the compound is a compound of formula II-B-2. In some embodiments, the compound is a compound of formula II-B-3. In some embodiments, the compound is a compound of formula II-B-4. In some embodiments, the compound is a compound of formula II-C-1. In some embodiments, the compound is a compound of formula II-C-2. In some embodiments, the compound is a compound of formula II-C-3. In some embodiments, the compound is a compound of formula II-C-4. In some embodiments, the compound is a compound of formula II-D-1. In some embodiments, the compound is a compound of formula II-D-2. In some embodiments, the compound is a compound of formula II-D-3. In some embodiments, the compound is a compound of formula II-D-4. In some embodiments, the compound is a compound of formula II-E-1. In some embodiments, the compound is a compound of formula II-E-2. In some embodiments, the compound is a compound of formula II-E-3. In some embodiments, the compound is a compound of formula II-E-4. In some

DETAILED DESCRIPTION

[0027] It has been found that compounds of formula I exhibit immunosuppressive and/or anti-inflammatory activity, while at the same time exhibiting less toxicity, than some known compounds. Thus compounds of formula I may be useful as immunosuppressive and/or anti-inflammatory agents. As used herein, the term “immune-mediated” refers to a disease or condition in which the body’s immune system overreacts and/or attacks the body.

[0028] Throughout this specification the terms and substituents retain their definitions.

[0029] “Alkyl” is intended to include linear, branched, or cyclic saturated hydrocarbon structures and combinations thereof. Lower alkyl refers to alkyl groups of from 1 to 6 carbon atoms. Examples of lower alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, s-and t-butyl and the like. Preferred alkyl groups are those of C₂₀ or below. Cycloalkyl is a subset of alkyl and includes cyclic hydrocarbon groups of from 3 to 8 carbon atoms. Examples of cycloalkyl groups include c-propyl, c-butyl, c-pentyl, norbornyl and the like.

[0030] C₁ to C₂₀ hydrocarbon includes alkyl, cycloalkyl, polycycloalkyl, alkenyl, alkynyl, aryl and combinations thereof. Examples include benzyl, phenethyl, cyclohexylmethyl, camphoryl and naphthylethyl. The term “carbocycle” is intended to include ring systems consisting entirely of carbon but of any oxidation state. Thus (C₃-C₁₀) carbocycle refers to such systems as cyclopropane, benzene and cyclohexene; (C₈-C₁₂) carbopolycycle refers to such systems as norbornane, decalin, indane and naphthalene.

[0031] Alkoxy or alkoxy refers to groups of from 1 to 8 carbon atoms of a straight, branched, cyclic configuration and combinations thereof attached to the parent structure through an oxygen atom. Examples include methoxy, ethoxy, propoxy, isopropoxy, cyclopropoxy, cyclohexoxy and the like. Lower-alkoxy refers to groups containing one to four carbons.

[0032] Oxaalkyl refers to alkyl residues in which one or more carbons has been replaced by oxygen. Examples include methoxypropoxy, 3,6,9-trioxadecyl and the like.

[0033] Acyl refers to groups of from 1 to 8 carbon atoms of a straight, branched, cyclic configuration, saturated, unsaturated and aromatic and combinations thereof, attached to the parent structure through a carbonyl functionality. One or more carbons in the acyl residue may be replaced by nitrogen, oxygen or sulfur as long as the point of attachment to the parent remains at the carbonyl. Examples include acetyl, benzoyl, propionyl, isobutyryl, *t*-butoxycarbonyl, benzyloxycarbonyl and the like. Lower-acyl refers to groups containing one to four carbons.

[0034] Aryl means 6-membered aromatic ring; a bicyclic 9- or 10-membered aromatic ring system; or a tricyclic 13- or 14-membered aromatic ring system. The aromatic 6- to 14-membered carbocyclic rings include, *e.g.*, benzene, naphthalene, indane, tetralin, and fluorene.

[0035] Heteroaryl mean a 5- or 6-membered heteroaromatic ring containing 1-3 heteroatoms selected from O, N, or S; a bicyclic 9- or 10-membered heteroaromatic ring system containing 1-3 heteroatoms selected from O, N, or S; or a tricyclic 13- or 14-membered heteroaromatic ring system containing 1-3 heteroatoms selected from O, N, or S. The 5- to 10-membered aromatic heterocyclic rings include, *e.g.*, imidazole, pyridine, indole, thiophene, benzopyranone, thiazole, furan, benzimidazole, quinoline, isoquinoline, quinoxaline, pyrimidine, pyrazine, tetrazole and pyrazole.

[0036] Arylalkyl refers to a substituent in which an aryl residue is attached to the parent structure through alkyl. Examples are benzyl, phenethyl and the like. Heteroarylalkyl refers to a substituent in which a heteroaryl residue is attached to the parent structure through alkyl. Examples include, *e.g.*, pyridinylmethyl, pyrimidinylethyl and the like.

[0037] Heterocycle means a cycloalkyl or aryl residue in which from one to three carbons is replaced by a heteroatom selected from the group consisting of N, O and S. The nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. Examples of heterocycles include pyrrolidine, pyrazole, pyrrole, indole, quinoline, isoquinoline, tetrahydroisoquinoline, benzofuran, benzodioxan, benzodioxole (commonly referred to as methylenedioxyphenyl, when occurring as a substituent), tetrazole, morpholine, thiazole, pyridine, pyridazine, pyrimidine, thiophene, furan, oxazole, oxazoline, isoxazole, dioxane, tetrahydrofuran and the like. It is to be noted that heteroaryl is a subset of heterocycle in which the heterocycle is aromatic. Examples of heterocycl residues additionally include piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxo-pyrrolidinyl, 2-oxoazepinyl, azepinyl, 4-piperidinyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, pyrazinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolyl, quinuclidinyl, isothiazolidinyl, benzimidazolyl, thiadiazolyl, benzopyranyl, benzothiazolyl, tetrahydrofuryl, tetrahydropyranyl, thienyl, benzothienyl, thiamorpholinyl, thiamorpholinylsulfoxide, thiamorpholinylsulfone, oxadiazolyl, triazolyl and tetrahydroquinolinyl.

[0038] Substituted alkyl, aryl, cycloalkyl, heterocycl etc. refer to alkyl, aryl, cycloalkyl, or heterocycl wherein up to three H atoms in each residue are replaced with alkyl, halogen, haloalkyl, hydroxy, loweralkoxy, carboxy, carboalkoxy (also referred to as alkoxy carbonyl), carboxamido (also referred to as alkylaminocarbonyl), cyano, carbonyl, nitro, amino,

alkylamino, dialkylamino, mercapto, alkylthio, sulfoxide, sulfone, acylamino, amidino, phenyl, benzyl, heteroaryl, phenoxy, benzyloxy, or heteroaryloxy.

[0039] The term "halogen" means fluorine, chlorine, bromine or iodine.

[0040] The following abbreviations and terms have the indicated meanings throughout:

Boc	=	t-butyloxy carbonyl
c-	=	cyclo
DCM	=	dichloromethane = methylene chloride = CH ₂ Cl ₂
DIEA	=	N,N-diisopropylethyl amine
DIPEA	=	diisopropylethylamine
DMF	=	N,N-dimethylformamide
Fmoc	=	9-fluorenylmethoxycarbonyl
HATU	=	O-(7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HBTU	=	O-(benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HOAc	=	acetic acid
HOAt	=	1-hydroxy-7-azabenzotriazole
HOBt	=	1-hydroxybenzotriazole
Hyp	=	4-hydroxyproline
Me	=	methyl
Pip	=	Pipecolic acid
Phe	=	phenylalanine
Pro	=	proline
PyBOP	=	O-(benzotriazol-1-yl)-trispyrrolidinephosphonium hexafluorophosphate
rt	=	room temperature
TBTU	=	O-(benzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate
TFA	=	trifluoroacetic acid
<i>t</i> -Hyp	=	<i>trans</i> -4-hydroxyproline
Trp	=	tryptophan
Tyr	=	tyrosine
Tyr(tBu)	=	(O-tert-butyl) tyrosine

[0041] Furthermore, a comprehensive list of abbreviations utilized by organic chemists (i.e. persons of ordinary skill in the art) appears in the first issue of each volume of the Journal of Organic Chemistry. The list, which is typically presented in a table entitled “Standard List of Abbreviations” is incorporated herein by reference. In addition, with respect to non-naturally occurring amino acids in which an additional methylene group is present in the backbone, the notation “ β^3 -Ho-” (or “beta 3-homo-”) is used herein to refer to an amino acid having an extra methylene (-CH₂-) in the backbone between the side-chain bearing carbon atom and the terminal nitrogen atom; the notation “ β^2 -Ho-” (or “beta 2-homo-”) is used herein to refer to an amino acid having an extra methylene in the backbone between the side-chain bearing carbon atom and the terminal carbon atom.

[0042] Embodiments of the present invention include compounds of formula I in the form of salts, in particular acid addition salts. Suitable salts include those formed with both organic and inorganic acids. Such acid addition salts will normally be pharmaceutically acceptable, although salts of non-pharmaceutically acceptable salts may be of utility in the preparation and purification of the compound in question. Thus, preferred salts include those formed from hydrochloric, hydrobromic, sulphuric, citric, tartaric, phosphoric, lactic, pyruvic, acetic, succinic, oxalic, fumaric, maleic, oxaloacetic, methanesulphonic, ethanesulphonic, p-toluenesulphonic, benzenesulphonic and isethionic acids. Salts of the compounds of formula I can be made by reacting the appropriate compound in the form of the free base with the appropriate acid.

[0043] The compounds of formula I in accordance with the embodiments of the invention are cyclic tetrapeptides. The synthesis of these peptides may be accomplished by cyclizing corresponding linear peptides that are themselves synthesized using methodologies known in the art; see, for example, Merrifield, J. Am. Chem. Soc., **85**:2149 (1964); Houghten, Proc. Natl. Acad. Sci. USA, **82**:5132 (1985); Kelley & Winkler in *Genetic Engineering Principles and Methods*, Setlow, J. K., ed., Plenum Press, N.Y., vol. 12, pp 1-19 (1990); Stewart & Young, *Solid Phase Peptide Synthesis*, Pierce Chemical Co., Rockford, Ill. (1984); Mergler et al. (1988) Tetrahedron Letters **29**:4005-4008; Mergler et al. (1988) Tetrahedron Letters **29**:4009-4012; Kamber et al. (eds), *Peptides, Chemistry and Biology*, ESCOM, Leiden (1992) pp. 525-526; Riniker et al. (1993) Tetrahedron Letters **49**:9307-9320; Lloyd-Williams et al. (1993) Tetrahedron Letters **49**:11065-11133; Andersson et al. (2000) Biopolymers **55**:227-250; Bray, Nature Reviews **2**:587-593 (2003), U.S. Pat. Nos. 4,105,603, 3,972,859, 3,842,067, 3,862,925, 6,015,881, 6,197,927, and 7,439,222. Such synthesis may be accomplish via liquid-phase or solid-phase synthesis, or by a combination of both, as is known in the art.

[0044] Liquid phase methods (often referred to as solution phase methods) of synthesis carry out all reactions in a homogeneous phase. Successive amino acids are coupled in solution until the desired peptide material is formed. During synthesis, successive intermediate peptides are purified by precipitation and/or washes.

[0045] In solid phase peptide synthesis (SPPS), a first amino acid or peptide group is bound to an insoluble support, such as a resin. Successive amino acids or peptide groups are added to the first amino acid or peptide group until the peptide material of interest is formed. The product of solid phase synthesis is thus a peptide bound to an insoluble support. Peptides synthesized via SPPS techniques are then cleaved from the resin, and the cleaved peptide is isolated.

[0046] More specifically, solid phase synthesis begins at the carboxy terminus of the putative peptide by coupling a protected amino acid to an inert solid support. The inert solid support can be any macromolecule capable of serving as an anchor for the C-terminus of the initial amino acid. Typically, the macromolecular support is a cross-linked polymeric resin (e.g. a polyamide or polystyrene resin) as shown in FIGS. 1-1 and 1-2, on pages 2 and 4 of Stewart & Young, *supra*. In some cases, the C-terminal amino acid is coupled to a polystyrene resin to form a benzyl ester. A macromolecular support is selected such that the peptide anchor link is stable under the conditions used to deprotect the α -amino group of the blocked amino acids in peptide synthesis. If a base-labile α -protecting group is used, then it is desirable to use an acid-labile link between the peptide and the solid support. For example, an acid-labile ether resin is effective for base-labile Fmoc-amino acid peptide synthesis as described on page 16 of Stewart & Young, *supra*. Alternatively, a peptide anchor link and α -protecting group that are differentially labile to acidolysis can be used. For example, an aminomethyl resin such as the phenylacetamidomethyl (Pam) resin works well in conjunction with Boc-amino acid peptide synthesis as described on pages 11-12 of Stewart & Young, *supra*. Guiller et al., *Chem Rev.* **2000**, *100*, 2091-2157, reviewed linkers and cleavage strategies in solid-phase organic synthesis and combinatorial chemistry, including peptide synthesis.

[0047] After the initial amino acid is coupled to an inert solid support, the α -amino protecting group of the initial amino acid is removed with, for example, trifluoroacetic acid (TFA) in methylene chloride and neutralizing in, for example, triethylamine (TEA). Following deprotection of the initial amino acid's α -amino group, the next α -amino and sidechain protected amino acid in the synthesis is added. The remaining α -amino and, if necessary, side chain protected amino acids are then coupled sequentially in the desired order by condensation to obtain an intermediate compound connected to the solid support. Alternatively, some amino

acids may be coupled to one another to form a fragment of the desired peptide followed by addition of the peptide fragment to the growing solid phase peptide chain

[0048] The condensation reaction between two amino acids, or an amino acid and a peptide, or a peptide and a peptide can be carried out according to the usual condensation methods such as the axide method, mixed acid anhydride method, DCC (N,N'-dicyclohexylcarbodiimide) or DIC (N,N'-diisopropylcarbodiimide) methods, active ester method, p-nitrophenyl ester method, BOP (benzotriazole-1-yl-oxy-tris [dimethylamino] phosphonium hexafluorophosphate) method, N-hydroxysuccinic acid imido ester method, etc, and Woodward reagent K method.

[0049] It is common in the chemical syntheses of peptides to protect any reactive side-chain groups of the amino acids with suitable protecting groups. Ultimately, these protecting groups are removed after the desired polypeptide chain has been sequentially assembled. Also common is the protection of the α -amino group on an amino acid or peptide fragment while the C-terminal carboxy group of the amino acid or peptide fragment reacts with the free N-terminal amino group of the growing solid phase polypeptide chain, followed by the selective removal of the α -amino protecting group to permit the addition of the next amino acid or peptide fragment to the solid phase polypeptide chain. Accordingly, it is common in polypeptide synthesis that an intermediate compound is produced which contains each of the amino acid residues located in the desired sequence in the peptide chain wherein individual residues still carry side-chain protecting groups. These protecting groups can be removed substantially at the same time to produce the desired polypeptide product following removal from the solid phase.

[0050] α - and ω -amino side chains can be protected, for example, with benzyloxycarbonyl (abbreviated Z), isonicotinyloxycarbonyl (iNoc), o-chlorobenzyloxycarbonyl [Z(2Cl) or 2-Cl-Z], p-nitrobenzyloxycarbonyl [Z(NO₂)], p-methoxybenzyloxycarbonyl [Z(OMe)], t-butoxy-carbonyl (Boc), t-amyoxy-carbonyl (Aoc), isobornyloxycarbonyl, adamantlyoxy-carbonyl (Adoc), 2-(4-biphenyl)-2-propyloxycarbonyl (Bpoc), 9-fluorenylmethoxycarbonyl (Fmoc), methylsulfonyethoxycarbonyl (Msc), trifluoroacetyl, phthalyl (Pht), formyl (For), 2-nitro-phenylsulphenyl (Nps), diphenylphosphinothiyl (Ppt), and dimethylphosphinothiyl (Mpt) groups, and the like. Additional examples of side chain protecting groups include acetyl (Ac), benzoyl (Bz), tert butyl (t-Bu), triphenylmethyl (trityl, Trt), tetrahydropyranyl, benzyl (Bzl), 2,6-dichlorobenzyl, nitro, p-toluenesulfonyl (Tos), xanthyl (Xan), benzyl, methyl, ethyl, and t-butyl ester, and aromatic or aliphatic urethan-type protecting groups, photolabile groups such as nitro veratryl oxycarbonyl (Nvoc), and fluoride labile groups such as trimethylsilylethyoxy-carbonyl (TEOC).

[0051] Examples of amino terminal protecting groups (also referred to herein as N-terminal protecting groups) include: (1) acyl-type protecting groups, such as formyl, acrylyl (Acr), benzoyl (Bz) and acetyl (Ac); (2) aromatic urethan-type protecting groups, such as benzyloxy-carbonyl (Z) and substituted Z, such as p-chlorobenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl, p-methoxybenzyloxycarbonyl; (3) aliphatic urethan protecting groups, such as t-butyloxycarbonyl (BOC), diisopropylmethoxycarbonyl, isopropylloxycarbonyl, ethoxycarbonyl, allyloxycarbonyl; (4) cycloalkyl urethan-type protecting groups, such as 9-fluorenyl-methyloxycarbonyl (Fmoc), cyclopentyloxycarbonyl, adamantlyloxycarbonyl, and cyclohexyloxycarbonyl; and (5) thiourethan-type protecting groups, such as phenylthiocarbonyl. Preferred protecting groups include 9-fluorenylmethyloxycarbonyl (Fmoc), 2-(4-biphenylyl)-propyl(2)oxycarbonyl (Bpoc), 2-phenylpropyl(2)-oxycarbonyl (Poc), and t-butyloxycarbonyl (Boc).

[0052] Protective groups for the carboxy functional group are exemplified by benzyl ester (OBzl), cyclohexyl ester (Chx), 4-nitrobenzyl ester (ONb), t-butyl ester (Obut), 4-pyridylmethyl ester (OPic), and the like. It is often desirable that specific amino acids such as arginine, cysteine, and serine possessing a functional group other than amino and carboxyl groups are protected by a suitable protective group. For example, the guanidino group of arginine may be protected with nitro, p-toluenesulfonyl, benzyloxycarbonyl, adamantlyloxycarbonyl, p-methoxybenzesulfonyl, 4-methoxy-2,6-dimethylbenzenesulfonyl (Nds), 1,3,5-trimethylphenylsulfonyl (Mts), and the like. The thiol group of cysteine can be protected with p-methoxybenzyl, trityl, and the like.

[0053] While it may be possible for the compounds of formula I to be administered as the raw chemical, it is preferable to present them as a pharmaceutical composition. According to a further aspect, there are provided in accordance with embodiments of the present invention a pharmaceutical composition comprising a compound of formula I or a pharmaceutically acceptable salt or solvate thereof, together with one or more pharmaceutically carriers thereof and optionally one or more other therapeutic ingredients. The carrier(s) must be “acceptable” in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

[0054] The formulations include those suitable for oral, parenteral (including subcutaneous, intradermal, intramuscular, intravenous and intraarticular), rectal and topical (including dermal, buccal, sublingual and intraocular) administration. The most suitable route may depend upon the condition and disorder of the recipient. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in

the art of pharmacy. Methods for making such formulations include the step of bringing into association a compound of formula I or a pharmaceutically acceptable salt or solvate thereof (“active ingredient”) with the carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

[0055] Formulations in accordance with embodiments of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

[0056] A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, lubricating, surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide sustained, delayed or controlled release of the active ingredient therein.

[0057] Formulations for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient. Formulations for parenteral administration also include aqueous and non-aqueous sterile suspensions, which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of a sterile liquid carrier, for example saline, phosphate-buffered saline (PBS) or the like, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

[0058] Formulations for rectal administration may be presented as a suppository with the usual carriers such as cocoa butter or polyethylene glycol.

[0059] Formulations for topical administration in the mouth, for example buccally or sublingually, include lozenges comprising the active ingredient in a flavoured basis such as

sucrose and acacia or tragacanth, and pastilles comprising the active ingredient in a basis such as gelatin and glycerin or sucrose and acacia.

[0060] Preferred unit dosage formulations are those containing an effective dose, as hereinbelow recited, or an appropriate fraction thereof, of the active ingredient.

[0061] It should be understood that in addition to the ingredients particularly mentioned above, the formulations in accordance with embodiments of this invention may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavoring agents.

[0062] As stated, in accordance with embodiments of the invention, compounds in accordance with embodiments of the invention may be used for the treatment or prevention of certain diseases or conditions. The term “preventing” as used herein refers to administering a medicament beforehand to forestall or obtund an attack. The person of ordinary skill in the medical art (to which the present method of use claims are directed) recognizes that the term “prevent” is not an absolute term. In the medical art it is understood to refer to the prophylactic administration of a drug to substantially diminish the likelihood or seriousness of a condition, and this is the sense intended in applicants’ claims. The reader’s attention is directed to the Physician’s Desk Reference, a standard text in the field, in which the term “prevent” occurs hundreds of times. No person of skill in the medical art construes the term in an absolute sense. Similarly, where it is stated that compounds in accordance with embodiments of the invention may be used to suppress an immune response, “suppress” will be understood to include reducing the degree of the response, and not necessarily absolutely preventing the response.

[0063] It may be found upon examination that compounds that are not presently excluded from the claims are not patentable to the inventors in this application. In that case, the exclusion of species and genera in applicants’ claims are to be considered artifacts of patent prosecution and not reflective of the inventors’ concept or description of their invention. The invention, in a composition aspect, is all compounds of formulae I and II, except those that are in the public’s possession.

[0064] Embodiments of the invention will be better understood with reference to the figures, in which:

[0065] Fig. 1 shows antigen-specific increase of the ear thickness derived from the experiment described in Example 2;

[0066] Fig. 2 shows total numbers of cells in the parotid lymph nodes observed in the experiment described in Example 2;

[0067] Fig. 3 shows the content of viable and dead cells in the parotid lymph nodes described in Example 2;

[0068] Fig. 4 presents number and participation of cell types in the draining lymph nodes registered in the experiment described in Example 2;

[0069] Fig. 5 presents toxicity of the compound of formula IA against mononuclear cells from human blood in as described in Example 3;

[0070] Fig. 6 presents effects of the intraperitoneal administration of the compound of formula IA on the humoral immune response of mice to sheep erythrocytes as described in Example 4;

[0071] Fig. 7 shows effects of the intraperitoneal administration of the compound of formula IA on the cellular immune response to ovalbumin as described in Example 5;

[0072] Figs. 8A and 8B show the effects of peptides in accordance with embodiments of the invention on PMBC survival *in vitro*;

[0073] Figs. 9A and 9B show the effects of peptides in accordance with embodiments of the invention on PHA-induced PBMC proliferation *in vitro*;

[0074] Figs. 10A and 10B show the effects of several compounds on changes in ear thickness in response to antigenic challenge, as described in Example 7;

[0075] Fig. 11 shows the permeability of capillary blood vessels in the Evans blue test, as described in Example 7;

[0076] Fig. 12 shows the total number of cells in the draining lymph nodes, as described in Example 7;

[0077] Fig. 13 shows the effects of the compounds on the numbers of circulating leukocytes, as described in Example 7;

[0078] Fig. 14 which shows the types of leukocytes present in different cases, as described in Example 7; and

[0079] Fig. 15 provides morphometric data on the number and composition of cells in mouse auricles, as described in Example 7.

Example 1. Synthesis of Cyclic Tetrapeptides

[0080] Cyclic tetrapeptides according to embodiments of the invention can be synthesized by the use of known peptide synthesis methods, such as solution-phase or solid-phase methods. In general the synthesis involves two consecutive steps: (1) synthesis of a linear tetrapeptide and (2) cyclization to yield the cyclic tetrapeptide. The linear tetrapeptide may be prepared in protected form and then deprotected prior to cyclization.

[0081] To illustrate, the synthesis of linear tetrapeptides on a solid support (Fmoc-L-phenylalanine attached to a Wang-type resin, or either L-beta-3-homophenylalanine or L-proline attached to a 2-chloro trityl resin) was conducted according to the following protocol:

1. The resin was swelled in dimethylformamide (DMF) (0.25 mmol, 10 ml/g resin) for 15 min.
2. The Fmoc group was removed with a 20% solution of piperidine in DMF (2 x 20 min).
3. The resin was washed with DMF (3 x 2 min)
4. The resin was washed with methanol (MeOH) (3 x 2 min)
5. The resin was washed with dichloromethane (DCM) (3 x 2 min)
6. Amino acid or peptide amino groups on the resin were acylated by shaking with a mixture of Fmoc-protected amino acid (4 eq), HBTU or TBTU (4 eq) and DIPEA (4 eq) dissolved in anhydrous DMF (4 ml/mmol), for 20 hours.
7. The resin was washed with DCM (3 x 2 min)
8. The resin was washed with MeOH (3 x 2 min)
9. The resin was washed with DCM (3 x 2 min)
10. The Kaiser test (for all amino acids except proline) was used to estimate if all amino groups were acylated.

[0082] If the result of the Kaiser test was negative, the resin was washed with DMF (1 x 2 min.) and a new coupling cycle started from point 2 of the protocol. If the result of the Kaiser test was positive, acylation was repeated with half the amount of the reagent used for the first coupling. In the case of acylation of proline, the Kaiser test is not sensitive enough to determine the degree of acylation, and for that reason in the case of proline the coupling procedure was repeated with the half amount of reagents. In the case of repeated acylation, the following washings were carried out:

- 7A. Washing with DCM (3 x 2 min)
- 8A. Washing with MeOH (3 x 2 min)
- 9A. Washing with DCM (3 x 2 min)

[0083] After the last coupling the resin was washed according to points 3-5 of the protocol. The Fmoc group was removed from the peptide as described in point 2, and the resin was washed again with DMF, MeOH and DCM as in points 3-5. Before the cleavage of the peptide from the polymeric support, the resin was dried overnight in a desiccator over KOH pellets under reduced pressure at room temperature.

[0084] Peptides were cleaved from the dried, Wang-type resin with a mixture of trifluoroacetic acid /water /triisopropylsilane 95:2.5:2.5 (v/v/v; 10 ml/1g of peptidyl-resin). The solution obtained was partially evaporated at room temperature under reduced pressure and peptide was

precipitated with 10 volumes of ether. After being filtered off, the crude peptide was dissolved in 0.05 M aqueous HCl and evaporated to dryness. The residue was dissolved in water and lyophilized.

[0085] Peptides were cleaved from 2-Cl-trityl resin by treatment with a 1:3:1 mixture of acetic acid / dichloromethane / trifluoroethanol (v/v/v; 10 ml/1g of peptidyl-resin). The resulting cleavage solution plus collected washings were filtered, evaporated to dryness at ambient temperature and reduced pressure, and the residue dissolved in a minimal volume of DCM, diluted with 20 volumes of hexane and re-evaporated (twice). Crude detached peptide was dissolved in water and lyophilized.

[0086] After lyophilized peptide was dried in a desiccator under vacuum over KOH and P₂O₅, it was ready for cyclisation.

[0087] In a typical synthesis following the above protocol, starting with Fmoc-Phe attached to the Wang resin (278 mg, 0.2 mmol, 0.72 mmol/g) and after sequential coupling of Fmoc-L- β^3 -homoPhe-OH, Fmoc-Pro-OH and again Fmoc-Pro-OH, the yield after lyophilization was 91 mg (82%) of crude linear peptide with a purity of 94% (HPLC).

[0088] Crude peptides were cyclized in DCM solution (peptide concentration 2×10^{-4} millimoles/Liter) with the aid of PyBOP/HOAt/2,4,6-collidine (3:3:5), with disappearance of linear precursors being traced by injecting samples of reaction solutions on an analytical reverse-phase (RP) HPLC column. At the end of the cyclization, the solution was evaporated to dryness under reduced pressure and the residue was partitioned between ethyl acetate (1000 ml of solvent per 1 millimole of peptide and 0.5 N HCl in water (100 ml of solution per 1 millimole of peptide). Organic phase was washed consecutively with 0.5 N HCl in water (2x), water (1x), 1M NaHCO₃ (3x) and water (1x). Organic solvent was removed under reduced pressure and the residual solid was dissolved in dioxane and freeze-dried. The crude product was purified on a preparative Vydac C₁₈ or Kromasil C₈ reversed-phase column (250 mm/22 mm, 100A, 10 μ m) using an elution gradient of solvent B (0.038% TFA in 82% acetonitrile/water) in solvent A (0.05% TFA in water).

[0089] As an example of a cyclization reaction, a solution of H-Pro-Pro- β^3 -hPhe-Phe-OH x HCl (55.7 mg, 0.1 mmol) in 500 ml DCM was treated with PyBOP (157 mg, 0.3 mmol) and HOAt (40 mg, 0.3 mmol) in the presence of 2,4,6-collidine (67.5 μ L, 0.5mmol), yielding after work-up and purification 13.5 mg (27%) of the cyclic tetrapeptide with purity 99% as determined by HPLC.

[0090] In this manner, compounds of formulae I-A through I-P were synthesized. The following Table 1 provides some data for the compounds of formulae I-A through I-P.

TABLE 1

No.	MS (MH ⁺)/MW calculated	HPLC (Vydac C ₁₈ 250 x 4.6 mm, 5μm, 300A, 1mL/min, 45-75% B in 15 min. A:0.05% TFA/water, B:0.038%TFA/82%Acetonitrile/water)	
		t _R (min)	Purity (% , λ 214nm)
I-A	503.45/502.6	4.79	99.35
I-B	503.47/502.6	5.96	99.00
I-C	517.49/516.6	5.26	99.00
I-D	503.25/502.6	5.10	98.12
I-E	503.20/502.6	5.83	97.15
I-F	503.25/502.6	6.70	99.08
I-G	503.25/502.6	5.87	98.71
I-H	503.26/502.6	6.23	98.79
I-J	503.30/502.6	4.80	100.0
I-K	503.27/502.6	4.65	97.44
I-L	542.1/541.6	4.72	99.47
I-M	519.3/518.6	5.81	100.0
I-N	575.3/574.7	7.25	98.43
I-O	517.3/516.6	6.52	97.05
I-P	MS+Na ⁺ = 541.47, MS+K ⁺ = 557.44/ 518.25	9.15 (C-8 Kromasil col., gradient 40- 80 % B in 15 min)	99.00

Example 2. Therapeutic efficacy of the peptide as a 0.1% ointment in the model of contact sensitivity to oxazolone in BALB/c mice

[0091] The aim of this experiment was to verify the therapeutic action of compound I-A and its toxicity in a generally accepted animal model. In the experiment described below, the compound of formula I-A was applied as a therapeutic preparation in the form of 0.1% wt/wt ointment based on a commonly used pharmaceutical vehicle, namely an ointment composed of 50% vaseline and 50% lanoline. The usefulness of the preparation in reduction of the effector phase of the contact sensitivity to oxazolone in mice, in comparison with reference preparations such as tacrolimus (Protopic[®]) and pimecrolimus (Elidel[®]), widely used for treatment of skin diseases, was studied.

Materials and Methods

[0092] Mice: BALB/c female mice, 8-10 week-old, delivered by the Institute of Laboratory Medicine, Łódź, Poland, were used for the study. The mice were fed a commercial, pelleted food and water *ad libitum*. The local ethics committee approved the study.

[0093] Reagents: Water-in-oil cream and ointment were delivered by Nepentes. Cyclic tetrapeptide (compound I-A) was synthesized as described above. Protopic® (tacrolimus) was purchased from Astellas, Ireland; Elidel® (pimecrolimus) was purchased from Novartis; Hydrocortisonum® (hydrocortisone) was purchased from Aflofarm Farmacja Polska, Poland. DMSO was obtained from Fluka; oxazolone, acetone, Evans blue, Giemsa, May-Grünwald, and formalin were from Sigma.

[0094] *Contact sensitivity to oxazolone:* The test was performed according to Noonan et al. (*Int. Arch. Allergy Appl. Immunol.*, 1978, **56**, 523-532), with some modifications. Mice were shaved on the abdomen (2 × 2 cm area) and after 24 h 100 µl of 0.5% oxazolone in acetone was applied to the shaved area. The contact sensitivity reaction was elicited 5 days later by application of 50 µl of 1% oxazolone in acetone on both sides of the ears. Ear edema was measured 48 h later using a spring caliper. The results were presented as antigen-specific increase of ear thickness (i.e. the background (BG) ear thickness of mice was subtracted from the measured thickness).

[0095] *Application of compounds:* In the experiment shown, the compound of formula I-A was applied topically as a 0.1% ointment on both sides of the ears (total volume of 50 µl per ear), at 24 h and 26 h after elicitation of the reaction with the second dose of oxazolone. Reference compounds were used in a similar fashion in the form of commercially available preparations.

[0096] *Determination of lymph node cell numbers:* Superficial parotid, mandibular and accessory mandibular lymph nodes were isolated, homogenized by pressing against a stainless screen into phosphate buffered solution (PBS), washed twice and re-suspended in PBS containing 0.2% Trypan blue. The total and nonviable cell numbers were determined using a light microscope and Bürker's hemocytometer. Mice treated only with the eliciting dose of antigen served as a background control.

[0097] *Determination of circulating leukocyte number and blood picture:* Mice were subjected to halothane anesthesia and bled from the retro-orbital plexus, followed by cervical dislocation. The number of blood leukocytes was determined by dilution of blood in Türk's solution and counting the cells in a hemocytometer. Blood smears were prepared on microscope glass, dried and stained with Giemsa and May-Grünwald reagents. The smears were subsequently reviewed histologically. The circulating leukocyte numbers were presented per 1 mm³ and the

blood cell compositions as a percentage of a given cell type. Mice treated only with the eliciting dose of antigen served as a background control.

[0098] Histological analysis: The auricles were fixed in 4% formalin solution for 48 h, washed for 24 h, dehydrated in an alcohol series and embedded in paraffin. The paraffin blocks were sliced in a Micron HM310 microtome into 6 μm sections. The sections were stained with haematoxylin and eosin and with toluidine blue. The histological analysis was performed in a Nikon Eclipse 801 light microscope. On the histological slides containing cross-sections of auricles, the morphometric estimations of neutrophils, macrophages, lymphocytes and mast cells in the perivascular and subepithelial connective tissue were performed. The cells were counted on the area of 0.07 mm^2 at 400 \times magnification. Morphometric analysis was done with the aid of imagine software NIS-Elements (Nikon). For every examined group, 25 enumerations of neutrophils, macrophages, lymphocytes and mast cells were carried out.

[0099] Statistics: The results are presented as mean values \pm standard error (SE). Brown-Forsyth's test was used to determine the homogeneity of variance between groups. When the variance was homogenous, analysis of variance (one-way ANOVA) was applied, followed by *post hoc* comparisons with the Tukey's test to estimate the significance of the differences between groups. Nonparametric data were evaluated with the Kruskal-Wallis' analysis of variance, as indicated in the text. Significance was determined at $p < 0.05$. Statistical analysis was performed using STATISTICA 7 for Windows. The statistical analysis applies to all data shown in this description.

Results:

[00100] The data included in Fig.1 show the therapeutic efficacy of compound of formula I-A (labeled "4B8M" in Fig. 1 and subsequent figures) and the reference preparations in mice with fully developed contact sensitivity reaction to oxazolone. The preparations were applied topically as described in the Methods. Fig. 1 presents only antigen-specific increases of the ear thickness (as a result of subtracting background values measured in non-sensitized mice which were given only the eliciting dose of antigen). Compound I-A caused about 80% inhibition of the ear edema; Protopic[®] and Elidel[®] respectively caused about 30% and 50% inhibition.

[00101] The intensity of inflammatory processes in the ears should correlate with cell numbers in the draining lymph nodes. Therefore, inhibition of inflammation should be associated with a decrease of cell number in the draining lymph nodes. Fig. 2 shows that both the compound of formula I-A as well as Elidel[®] decreased the numbers of lymphocytes in the

draining lymph nodes to the level registered in non-sensitized mice. However, in mice treated with Protopic® the number of lymph node cells was similar to that in untreated mice.

[00102] Fig. 3 shows the proportions of viable and dead cells in the draining lymph nodes, expressed in percentages. The compound of formula I-A exhibits a negligible toxic effect as compared to the control, non-treated mice. A higher toxic effect is caused by Elidel®, and Protopic® is exceptionally toxic with regard to lymph node cells.

[00103] Complementary information regarding the therapeutic efficacy of the preparations may be derived from histological analysis of the cell number and composition in the inflamed auricles. Fig. 4 depicts the numbers and participation of basic cell types involved in the local inflammatory process. In Fig. 4, Mast = mastocytes, L = lymphocytes, MØ = macrophages, Ne = neutrophils. The auricles from untreated mice (K +) are characterized by a high infiltration of neutrophils. Application of the compound of formula I-A almost entirely reversed the changes observed in control mice (normalization of neutrophil number with some increase in the macrophage content). Protopic®, in turn, caused some changes in the proportion of respective cell types with no reduction of the total cell infiltrate. Elidel® caused a moderate diminution of the total cell number.

Example 3. The toxicity of compound I-A versus that of cyclolinopeptide against human blood mononuclear cells

[00104] For evaluation of toxicity of the compound of formula I-A, human peripheral mononuclear blood cells (PBMC) were chosen. This fraction consists of approximately 80% lymphocytes and 20% monocytes. As a reference compound, cyclolinopeptide (CLA) was selected since the compound of formula I-A shares a part of the sequence of CLA. CLA exhibits immunosuppressive properties comparable to that of cyclosporine A, but is less toxic.

Materials and Methods

[00105] *The cytotoxic test:* Venous blood from a single donor was taken into heparinized syringes, diluted twice with phosphate buffered saline (PBS) and applied onto Lymphoprep® (Polfa, Kutno, Poland) (density of 1.077 g/ml). After centrifugation at 1200 x g for 20 min, the mononuclear cells from the interphase were harvested and washed 3 times with PBS. The cells were re-suspended in a standard culture medium consisting of RPMI-1640 medium, L-glutamine, sodium pyruvate, 2-mercaptoethanol, 100 µg/ml each of streptomycin and penicillin, and 10% fetal calf serum. The cells were distributed to in 96-well flat-bottom culture plates at density of 2 x 10⁵/100 µl. The compounds (formula I-A and CLA) were

initially dissolved in DMSO (5 mg/500 μ l) and subsequently in the culture medium. DMSO, appropriately diluted with the culture medium, was used as a control. After 24 h incubation in a cell culture incubator, cell viability was determined by a colorimetric method (Hansen et al., *J Immunol Methods*, **1989**, 119, 203-210).

[00106] The results are shown in Fig. 5, presented as mean optical density values from quadruplicate wells (cell cultures) \pm SE. As can be seen in Fig. 5, the compound of formula I-A (listed as “4B8M”) did not show appreciable toxicity in the concentration range of 10 to 100 μ g/ml. CLA, on the other hand, showed a statistically significant cytotoxic effect at 40 μ g/ml.

Example 4. Effect of peptide on the humoral immune response to SRBC *in vivo*

[00107] CBA female mice, 8-12 weeks old, were delivered by The Institute of Laboratory Medicine, Łódź, Poland. The mice had free access to water and pelleted food. The local ethics committee approved the study. Sheep erythrocytes (SRBC) were delivered by Wroclaw University of Life and Environmental Sciences, Poland, and maintained on RPMI-1640 medium.

[00108] *The primary humoral immune response to SRBC *in vivo*:* Mice were immunized with 0.2 ml of 5% SRBC suspension (0.5 ml of SRBC pellet re-suspended to a volume of 10 ml of 0.9% NaCl), intraperitoneally. After 4 days the number of antibody-forming cells (AFC) in the spleens was determined using an assay of local hemolysis in agar gel (per Mishell et al., *J Exp Med*, **1967**, 126, 423-442). The results are presented in Fig. 6 as a mean value of 5 mice \pm standard error and expressed as AFC number per 10^6 of viable splenocytes.

[00109] Mice were immunized with SRBC as described above and after 2 h were given 10 or 100 μ g of the compound of formula I-A. Cyclosporin A (CsA) served as a reference drug. The number of antibody-forming cells to SRBC was measured after 4 days. As shown in Fig. 6, the compound of formula IA was more inhibitory at both doses than CsA.

Example 5. Effect of peptide on the cellular immune response *in vivo* to ovalbumin

[00110] Male CBA mice 8-12 weeks old were delivered by The Institute of Laboratory Medicine, Łódź, Poland. The mice had free access to water and pelleted food. Ovoalbumin was from Sigma and the adjuvants from Difco.

[00111] *Delayed type hypersensitivity (DTH) test:* Mice were sensitized subcutaneously with 5 μ g of ovalbumin (OVA) emulsified in Freund’s complete adjuvant in the tail base. After 4 days the mice were challenged with 50 μ g of OVA in Freund’s incomplete adjuvant in the hind footpads. Following the next 24 hours, the footpad thickness was measured using a caliper.

Controls (background response mice) were not sensitized but received the challenging dose of OVA. The compound of formula I-A and the reference compound were administered to mice in two 100 µg intraperitoneal doses, 2 h before and 24 h after the sensitizing dose of antigen. The results, presented in Fig. 7 as a mean value of antigen-specific increase of footpad thickness measured in 5 mice and expressed in DTH units (one DTH unit = 10^{-2} cm) \pm standard error, show that the compound of formula I-A, given in two doses, 2 h and 24 h after immunization, inhibited the delayed-type hypersensitivity reaction to OVA. That suppressive action was stronger than those of CLA and CsA.

Example 6

[00112] Cyclic tetrapeptides were tested *in vitro* for their effects on phytohemagglutinin A (PHA)-induced proliferation of human peripheral blood mononuclear cells (PBMC) and for lipopolysaccharide (LPS)-induced production of tumor necrosis factor alpha (TNF- α) production by whole blood cell cultures at 1-100 µg/ml concentration range. Compounds were also tested for cell toxicity at 1-100 µg/ml concentration range against human PBMC.

Materials and Methods:

[00113] Reagents: RPMI-1640 medium (Cibi/Life Technologies, UK), fetal calf serum (FCS, Gibco), DMSO, phytohemagglutinin A (PHA), lipopolysaccharide (LPS) from *E. coli* strain O111:B4 (Sigma), 93-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT), SDS and DMF (Sigma). The culture medium consisted of RPMI-1640, 10% addition of FCS, L-glutamine, sodium pyruvate, 2-mercaptoethanol and antibiotics (streptomycin and penicillin). Cyclic tetrapeptides were initially dissolved in DMSO (5 mg/ml), the dissolved in the culture medium to the desired concentration.

[00114] Isolation of PBMC: Venous blood was taken from a single donor (a male, 62-years old) into heparinized syringes and diluted twice with phosphate-buffered saline (PBS). PBMC were isolated by centrifugation on Ficoll-uroponine gradient (density 1.077g/ml) (Lymphoprep; PAA Laboratories), at 800×g for 20min at 4°C. The interphase cells, consisting of lymphocytes (20%) and monocytes (80%) were then washed three times with Hanks' medium and re-suspended in the culture medium at density of 2×10^6 cells/ml.

[00115] The proliferative response of PBMC to PHA: The isolated PBMC were distributed into 96-well flat-bottom plates in 100 µl aliquots (2×10^5 cells/well). PHA was used at concentration of 5 µg/ml. The compounds were tested at concentrations of 1, 10 and 100 µg/ml. DMSO at appropriate dilutions served as control. After a four-day incubation in a cell

culture incubator, the proliferative response of cells was determined by the colorimetric MTT method (Hansen et al., J. Immunol. Methods, 1989, pp. 203-210). The data are presented as a mean OD value from quadruplicate wells \pm standard error (SE). The cultures “Control (-)” contained no mitogen (PHA). The cultures “Control (PHA)” contained PHA but not cyclic tetrapeptides.

[00116] Toxicity test: PBMC, at a density of 2×10^5 /100 μ l/well, re-suspended in the culture medium, were cultured for 24 h in a cell culture incubator with the cyclic tetrapeptides at concentrations of 1, 10 and 100 μ g/ml concentrations. Cell survival was determined by the MTT colorimetric method (Hansen et al., J. Immunol. Methods, 1989, pp. 203-210). The data are presented as a mean OD value from quadruplicate wells \pm standard error (SE). The cultures “Control (-)” contained only cells in the culture medium.

[00117] The determination of TNF alpha activity (per Espevik et al., J. Immunol. Methods, 95 (1986):99-103): Human whole blood was diluted 10-fold with RPMI-1640 medium and distributed to 24-well culture plates in 1ml aliquots. LPS was added to the culture at a concentration of 1 μ g/ml. The studied peptides were used at concentrations of 1, 10 and 100 μ g/ml. After overnight incubation, the supernatants were harvested and frozen at -20°C until cytokine determination. TNF- α activity was determined using a bioassay. Briefly, WEHI 164.13 cells (ATCC CRL 1751) were seeded at a concentration of 2×10^4 cells/well in quadruplicate. Increasing dilutions of the assayed supernatant were mixed with the target cells in the presence of actinomycin D (1 μ g/ml). After 20 h of incubation, MTT was added into the wells, and the cultures were incubated for an additional 4 h. Next, a lysing buffer (20% SDS with 50% DMF, pH 4.7) was added and the optical density at 550 nm with the reference wavelength of 630 nm in a Dynatech 5000 spectrophotometer was measured after 24 h. The detection limit of the assay was about 2.5 pg/ml. One unit of TNF- α activity was defined as an inverse of supernatant dilution where 50% cell death took place. The cultures labeled “Control (-)” contained no LPS. The cultures labeled “Control (LPS)” contained LPS and none of the studied compounds. Statistical analysis was not applied since the data derive from single cultures (wells).

[00118] Colorimetric MTT assay for cell growth and kill: The assay was performed per Hansen et al., J. Immunol. Methods, 1989, 119 pp. 203-210. Briefly, 25 μ l of MTT (5 mg/ml) stock solution was added per well at the end of cell incubation and the plates were incubated for 3 h in a cell culture incubator. Then, 100 μ l of the extraction buffer (20% SDS with 50% DMF, pH 4.7) was added. After additional overnight incubation, the optical density was measured at 550 nm (Dynatech 5000).

[00119] Where applicable, results are presented as mean values \pm standard error (SE). Brown-Forsyth's test was used to determine the homogeneity of variance between groups. When the variance was homogenous, analysis of variance (one-way ANOVA) was applied, followed by post hoc comparisons with the Tukey's test to estimate the significance of the differences between groups. Significance was determined at $P < 0.05$. Statistical analysis was performed using STATISTICA 6.1 for Windows.

Results

[00120] Effects of the compounds on survival of PBMC: The effects of the peptides on PMBC survival in 24h culture are presented in Figs 8A and 8B. Peptide 4B8M (compound I-A) was included as a reference compound. Appropriate dilutions of DMSO were added to control cultures. The results showed no signs of toxicity of the compounds in the studied concentration range. In the figures, P01 = compound I-D, P02 = compound I-E, P03 = compound I-F, P04 = compound I-G, P05 = compound I-H, P06 = compound I-J, P07 = compound I-K, P08 = compound I-L, P10a = compound I-M, P10b = compound I-N, and P11 = compound I-O.

[00121] Figs. 8A and 8B show the effects of the tested peptides on the survival of PBMC. Fig 8A: Statistics (all comparisons vs. DMSO at appropriate dilutions): 100 μ g/ml: 4B8M NS ($P = 0.9999$); P01 NS ($P = 1.0000$); P02 NS ($P = 1.0000$); P03 NS ($P = 1.0000$); P04 NS ($P = 0.9047$); P05 NS ($P = 1.0000$); P06 NS ($P = 0.9999$); P07 NS ($P = 1.0000$); 10 μ g/ml: 4B8M NS ($P = 1.0000$); P01 NS ($P = 1.0000$); P02 NS ($P = 1.0000$); P03 NS ($P = 1.0000$); P04 NS ($P = 0.9999$); P05 NS ($P = 1.0000$); P06 NS ($P = 1.0000$); P07 NS ($P = 1.0000$); 1 μ g/ml: 4B8M NS ($P = 1.0000$); P01 NS ($P = 1.0000$); P02 NS ($P = 1.0000$); P03 NS ($P = 1.0000$); P04 NS ($P = 1.0000$); P05 NS ($P = 1.0000$); P06 NS ($P = 0.8253$); P07 NS ($P = 1.0000$) (ANOVA). Fig. 8B: Statistics (all comparisons vs. DMSO at appropriate dilutions): 100 μ g/ml: 4B8M NS ($P = 0.0669$); P08 NS ($P = 0.9957$); P10b NS ($P = 1.0000$); 10 μ g/ml: 4B8M NS ($P = 0.9999$); P08 NS ($P = 1.0000$); P10b NS ($P = 1.0000$); 1 μ g/ml: 4B8M NS ($P = 0.3176$); P08 NS ($P = 0.9999$); P10b NS ($P = 1.0000$) (ANOVA).

[00122] Effects of the peptides on PHA-induced PMBC proliferation: The effects of the peptides on the proliferative response of PMBC are presented in Figs. 9A and 9B. Peptide 4B8M was included as a reference compound. Appropriate dilutions of DMSO were added to control cultures.

[00123] Fig. 9A: Effects of the peptides on PHA-induced PBMC proliferation: Statistics (all comparisons vs. DMSO at appropriate dilutions): 1 μ g/ml: 4B8M NS ($P = 0.9995$); P01 NS

(P=1.0000); P02 NS (P=1.0000); P03 NS (P=1.0000); P04 NS (P=0.9047); P05 NS (P=0.5198); P06 NS (P=1.0000); P07 NS (P=0.1445); 10 μ g/ml: 4B8M NS (P=1.0000); P01 NS (P=0.9999); P02 NS (P=1.0000); P03 NS (P=0.9930); P04 NS (P=0.4297); P05 NS (P=1.0000); P06 NS (P=1.0000); P07 NS (P=0.8647); 100 μ g/ml: 4B8M NS (P=1.0000); P01 NS (P=1.0000); P02 NS (P=0.9982); P03 P=0.0001; P04 NS (P=0.9970); P05 NS (P=0.2037); P06 NS (P=0.1257); P07 NS (P=1.0000) (ANOVA).

[00124] Fig. 9B: Effects of the peptides on PHA-induced PBMC proliferation: Statistics (all comparisons vs. DMSO at appropriate dilutions): 1 μ g/ml: 4B8M NS (P=0.9919); P08 NS (P=0.9999); P10b NS (P=1.0000); 10 μ g/ml: 4B8M NS (P=1.0000); P08 NS (P=1.0000); P10b NS (P=0.2763); 100 μ g/ml: 4B8M NS (P=0.4941); P08 NS (P=1.0000); P10b NS (P=0.9933) (ANOVA).

[00125] Effects of the peptides on LPS-induced TNF- α production in whole blood cell cultures: The effects of the peptides on LPS-induced TNF- α production in whole blood cell cultures are presented in Tables 2A and 2B. Peptide 4B8M was included as a reference compound. Appropriate dilutions of DMSO were added to control cultures.

Table 2A.
Effects of peptides on LPS-induced TNF- α production

Compound	compound concentration (μ g/ml)	TNF- α (pg/ml)	% inhibition vs. DMSO
Control (-)	-	92	-
Control (LPS)	-	6668	-
DMSO Only	1	8167	-
	10	8006	-
	100	196	-
4B8M (compound I-A)	1	7364	9.8
	10	7176	10.4
	100	178	9.2
P01 (compound I-D)	1	7016	14.1
	10	5757	28.1
	100	141	28.1
P02 (compound I-E)	1	6614	19.0
	10	6346	20.7
	100	176	10.2
P03 (compound I-F)	1	6052	25.9
	10	5222	34.8
	100	76	61.2
P04 (compound I-G)	1	5463	33.1
	10	5302	33.8

	100	58	70.4
P05 (compound I-H)	1	7123	12.8
	10	7203	10.0
	100	110	43.9
P06 (compound I-J)	1	8033	1.6
	10	7658	4.4
	100	203	0.0
P07 (compound I-L)	1	6453	21.0
	10	7284	9.0
	100	163	16.8

Table 2B.
Effects of the peptides on LPS-induced TNF alpha production

Compound	compound concentration (μg/ml)	TNF alpha (pg/ml)	% of inhibition vs. DMSO
Control (-)	-	223	-
Control (LPS)	-	4260	-
DMSO Only	1	4550	-
	10	4351	-
	100	854	-
4B8M (compound I-A)	1	4097	10.0
	10	4416	-
	100	979	-
P08 (compound I-M)	1	4039	11.2
	10	3363	22.7
	100	269	68.5
P10b (compound I-O)	1	3792	16.7
	10	3739	14.0
	100	282	67.0

Example 7. Inhibitory effect of Compound I-A on toluene diisocyanate-induced ear inflammation in mice

[00126] The efficacy of the compound I-A in suppressing ear inflammation in BALB/c mice which was induced with toluene diisocyanate (TDI). Commercially available Protopic® (tacrolimus) and Elidel® (pimecrolimus) served as reference drugs.

Materials and Methods

[00127] Mice: BALB/c female mice, 8-10 weeks old, were delivered by the Institute of Laboratory Medicine, Łódź, Poland. The mice were fed a commercial, pelleted food and water *ad libitum*. The local ethics committee approved the study.

[00128] Reagents. Compound I-A was synthesized as described above; Protopic® (tacrolimus) was from Astellas, Ireland; Elidel® (pimecrolimus) from Novartis; DMSO from Fluka; TDI, acetone, Evans blue, Trypan blue, Giemsa, May-Grünwald, haematoxylin, eosin, toluidine blue and formalin were from Sigma.

[00129] Immune response to TDI. The test was performed according to Yamamoto, *Eur. J. Pharmacol.*, 2006, **550**, 166-172, with minor modifications. Mice were shaved on the abdomen (2×2 cm area) and after 24 h 100 µl of 3% TDI in acetone was applied through 3 consecutive days. After 14 days the reaction was elicited by application of 50 µl of 0.3% TDI on both sides of the ears. The procedure was repeated 5 times every 3 days. Ear thickness was measured using a spring caliper (Mitutoyo) 5 h and 24 h after each challenge with TDI.

[00130] Application of compounds. Compound I-A was applied topically in the form of 0.1% ointment on both sides of the ears (total volume of 100 µl – 50 µl per ear), one hour after each challenge with TDI. The reference drugs were applied in a similar way.

[00131] Determination of lymph node cell numbers. Superficial parotid, mandibular and accessory mandibular lymph nodes were isolated, homogenized by pressing against a stainless screen into PBS, washed 2 × with PBS and re-suspended in PBS containing 0.2% Trypan blue. The total and nonviable cell numbers were counted using a light microscope and Bürker's hemocytometer.

[00132] Determination of circulating leukocyte number and blood picture. Mice were subjected to halothane anesthesia and bled from the retro-orbital plexus, followed by the cervical dislocation. The number of blood leukocytes was determined by dilution of blood in Türk's solution and counting the cells in a hemocytometer. Blood smears were prepared on microscope glass, dried and stained with Giemsa and May-Grünwald reagents. The smears were subsequently reviewed histologically. The cell numbers were presented per 1 µl and the blood cell compositions as a percentage of a given cell type.

[00133] Evans blue test. Mice were given 1 mg of Evans blue in 0.2 ml of 0.9% NaCl, intravenously. After 30 min mice were sacrificed, the ears were cut off, weighed and immersed in 50 µl of 1M KOH for 18 h at 37°C. The dye was extracted from the ears using 450 µl of 0.2 M phosphate acid and acetone (5:13 ratio). The samples were centrifuged at 3,000 rpm for 15 min. The optical densities (OD) of the supernatants were measured at 630 nm. The amount of Evans blue (µg/ml) was determined based on a standard curve. The results were presented as the amount of Evans blue per 100 mg of wet tissue. Mice treated only with the eliciting dose of antigen served as a background control.

[00134] Histological analysis. The auricles were fixed in 4% formalin solution for 48 h, washed for 24 h, dehydrated in an alcohol series and embedded in paraffin. The paraffin blocks were sliced in a Micron HM310 microtome into 6 μm sections. The sections were stained with haematoxylin and eosin and with toluidine blue. Histological analysis was performed using a Nikon Eclipse 801 light microscope. Morphometric estimations of neutrophils, macrophages, lymphocytes and mast cells in the perivascular and subepithelial connective tissue were performed on the histological slides containing cross-sections of auricles. Cells were counted on an area of 0.07 mm^2 at 400 \times magnification. Morphometric analysis was performed using an imagine software NIS-Elements (Nikon). For every preparation examined, 25 enumerations of neutrophils, macrophages, lymphocytes and mast cells were carried out.

[00135] Statistics. The results in Figs. 10A and 10B are presented as mean values \pm standard error (SE). Brown-Forsyth's test was used to determine the homogeneity of variance between the groups. When the variance was homogenous, analysis of variance (one-way ANOVA) was applied, followed by *post hoc* comparisons with the Tukey's test to estimate the significance of the differences between groups. Nonparametric data were evaluated with the Kruskal-Wallis' analysis of variance, as indicated in the text. Significance was determined at $P<0.05$. Statistical analysis was performed using STATISTICA 7 for Windows.

Results

[00136] Effects of the compounds on the ear thickness. The humoral immune response to TDI was elicited as described in the Methods. Mice were treated topically with the tetrapeptide (formula I-A, labeled 4B8M in Figs. 10A and 10B) and the reference compounds one hour after each challenge with antigen. The effects of the treatments are presented in Figs. 10A and 10B, which show ear thickness measure 5 h (Fig. 10A) and 24 h (Fig. 10B) after administration on the day of the test indicated in the figure. Control responses to TDI gradually elevated after each antigen challenge (best seen in the 5 h measurement). The results showed differentiated efficacy of the compounds in reducing the ear swelling.

[00137] Fig. 10A: Day 14: Control vs 4B8M $P=0.0005$; Control vs Protopic[®] $P=0.0002$; Control vs Elidel[®] $P=0.0152$; 4B8M vs Protopic[®] NS; 4B8M vs Elidel[®] NS; Day 17: Control vs 4B8M $P=0.0001$; Control vs Protopic[®] $P=0.0001$; Control vs Elidel[®] $P=0.0377$; 4B8M vs Protopic[®] NS; 4B8M vs Elidel[®] $P=0.0002$; Day 20: Control vs 4B8M $P=0.0001$; Control vs Protopic[®] $P=0.0001$; Control vs Elidel[®] NS; 4B8M vs Protopic[®] NS; 4B8M vs Elidel[®] $P=0.0001$; Day 23: Control vs 4B8M $P=0.0001$; Control vs Protopic[®] $P=0.0001$; Control vs

Elidel® NS; 4B8M vs Protopic® NS; 4B8M vs Elidel® P=0.0001; Day 27: Control vs 4B8M P=0.0001; Control vs Protopic® P=0.0001; Control vs Elidel® P=0.0001; 4B8M vs Protopic® NS; 4B8M vs Elidel® P=0.0001 (ANOVA). Fig. 10B: Day 14: Control vs 4B8M P=0.0001; Control vs Protopic® P=0.0001; Control vs Elidel® P=0.0004; 4B8M vs Protopic® NS; 4B8M vs Elidel® NS; Day 17: Control vs 4B8M P=0.0001; Control vs Protopic® P=0.0023; Control vs Elidel® P=0.0004; 4B8M vs Protopic® NS; 4B8M vs Elidel® NS; Day 20: Control vs 4B8M P=0.0006; Control vs Protopic® P=0.0003; Control vs Elidel® P=0.0156; 4B8M vs Protopic® NS; 4B8M vs Elidel® NS; Day 23: Control vs 4B8M P=0.0001; Control vs Protopic® P=0.0001; Control vs Elidel® P=0.0039; 4B8M vs Protopic® NS; 4B8M vs Elidel® P=0.0027; Day 27: Control vs 4B8M P=0.0001; Control vs Protopic® P=0.0001; Control vs Elidel® P=0.0023; 4B8M vs Protopic® NS; 4B8M vs Elidel® P=0.0016 (ANOVA).

[00138] Effects of the compounds on permeability of skin vessels: The permeability of capillary blood vessels is presented in Fig. 11, which shows the permeability of capillary blood vessels in the Evans blue test. The procedure, as described above, was performed 24 h after the fifth challenge with TDI (on day 28). As shown in the Fig. 11, the rates of blood vessel permeability were strictly correlated with the effects of the compounds on ear thickness in the respective mouse groups. Statistics: BG vs Control P = 0.0248; Control vs 4B8M (I-A) P = 0.030; Control vs Protopic® NS; Control vs Elidel® NS; 4B8M (I-A) vs Protopic® NS; 4B8M (I-A) vs Elidel® NS (ANOVA).

[00139] Effects of the compounds on number of cells in draining lymph nodes: Fig. 12 shows the total number of cells in the draining lymph nodes. As shown in Fig. 12, the treatment of mice with compound I-A resulted in a reduction of the lymph node cell numbers almost to the background levels (non-sensitized mice). Statistics: BG vs Control P = 0.0001; Control vs 4B8M (I-A) P = 0.0001; Control vs Protopic® NS; Control vs Elidel® NS; 4B8M (I-A) vs Protopic® P = 0.0183; 4B8M (I-A) vs Elidel® P = 0.0001 (ANOVA of Kruskal-Wallis).

[00140] Effects of the compounds on the numbers of circulating leukocytes. Fig. 13 shows the effects of the compounds on the numbers of circulating leukocytes; the application of the studied preparations lowered the numbers of circulating leukocytes to the levels observed in control, unsensitized mice. Statistics: BG vs Control P = 0.0001; Control vs 4B8M (I-A) P = 0.0001; Control vs Protopic® P = 0.0001; Control vs Elidel® P = 0.0001; 4B8M (I-A) vs Protopic® NS; 4B8M (I-A) vs Elidel® NS (ANOVA).

[00141] Effects of compounds on blood cell composition: The blood composition in control mice with fully developed reaction to TDI was characterized by an increased content of neutrophils and eosinophils compared to control, background mice (Fig. 14, which shows a

breakdown of the types of leukocytes in each case). The blood picture was normalized upon application of 4B8M (I-A) (a reduction of neutrophil and eosinophil contents) but not following administration of Protopic® or Elidel®. Statistics: Bands (B): BG vs Control NS; Control vs 4B8M (I-A) NS; Control vs Protopic® P = 0.0500; Control vs Elidel® P = 0.0500; 4B8M (I-A) vs Protopic® NS; 4B8M vs Elidel® NS (ANOVA of Kruskal-Wallis); Segments (S): BG vs Control P = 0.0131; Control vs 4B8M (I-A) NS; Control vs Protopic® NS; Control vs Elidel® NS; 4B8M (I-A) vs Protopic® P = 0.0163; 4B8M (I-A) vs Elidel® NS (ANOVA of Kruskal-Wallis); Eosinophils (E): BG vs Control P = 0.0001; Control vs 4B8M (I-A) P = 0.0001; Control vs Protopic® NS; Control vs Elidel® NS; 4B8M (I-A) vs Protopic® NS; 4B8M vs Elidel® P = 0.0146 (ANOVA of Kruskal-Wallis); Lymphocytes (L): BG vs Control P = 0.0001; Control vs 4B8M (I-A) P = 0.0043; Control vs Protopic® NS; Control vs Elidel® NS; 4B8M (I-A) vs Protopic® P = 0.0345; 4B8M (I-A) vs Elidel® NS (ANOVA of Kruskal-Wallis).

[00142] The effects of the compounds on cell composition in the auricles: Fig. 15 provides morphometric data on the number and composition of cells in the auricles. The composition of cell types within the auricles differed among the studied mouse groups is presented in Fig. 15. The predominant, residing cell types in control non-sensitized mice are mastocytes and neutrophils (10 and 5 cells per the analyzed area, respectively). In sensitized, control mice, untreated with the therapeutics, the number of mastocytes increased twofold and neutrophils almost 5-fold (20 and 23 cells, respectively). Protopic® and Compound I-A were effective in reducing the cell numbers to 14 and 14.8. Statistics: Ne (neutrophils): BG vs Control P = 0.0001; Control vs 4B8M (I-A) P = 0.0151; Control vs Protopic® P = 0.0001; Control vs Elidel® NS; 4B8M vs Protopic® NS; 4B8M (I-A) vs Elidel® P = 0.0003 (ANOVA of Kruskal-Wallis); MØ (macrophages): BG vs Control P = 0.0144; Control vs 4B8M (I-A) NS; Control vs Protopic® NS; Control vs Elidel® P=0.0255; 4B8M vs Protopic® NS; 4B8M (I-A) vs Elidel® P = 0.0031 (ANOVA of Kruskal-Wallis); L (lymphocytes): BG vs Control NS; Control vs 4B8M (I-A) NS; Control vs Protopic® NS; Control vs Elidel® P = 0.0023; 4B8M vs Protopic® NS; 4B8M (I-A) vs Elidel® P = 0.0001 (Anova of Kruskal-Wallis); Mast (mastocytes): BG vs Control P = 0.0001; Control vs 4B8M (I-A) NS; Control vs Protopic® P = 0.0001; Control vs Elidel® NS; 4B8M (I-A) vs Protopic® NS; 4B8M (I-A) vs Elidel® P = 0.0001 (ANOVA of Kruskal-Wallis).

Example 8 – In vitro tests of Compounds I-B and I-C

Methods

[00143] Proliferative response of splenocytes to concanavalin A (ConA): Spleens were pressed against a plastic screen into 0.83% NH₄Cl solution to lyse erythrocytes (5 min

incubation at room temperature). The cells were then washed twice with Hanks' medium, passed through glass wool column to remove debris, and re-suspended in the culture medium, referred to below as the "culture medium", consisting of RPMI-1640, supplemented with 10% of fetal calf serum, L-glutamine, sodium pyruvate, 2-mercaptoethanol, streptomycin and penicillin (100 µg/ml). The cells were then distributed into 96-well flat-bottom tissue culture plates (Nunc) at a density of 2×10^5 cells/100µl/well. Con A (2.5 µg/ml) was added to induce cell proliferation. The compounds were added to the cultures at doses of 1 – 100 µg/ml. After a three-day incubation, cell proliferation was determined using the colorimetric MTT method (Hansen MB, *J Immunol Methods*, 1989, 119, 203-210). The results were presented as the mean optical density (OD) at 550 nm ± SE from quadriplicate determinations (wells).

[00144] Secondary humoral immune response in vitro to sheep erythrocytes (SRBC): Mice were sensitized intraperitoneally with 0.2 ml of 5% (v/v) SRBC suspension. After four days spleens from these mice were isolated and single cell suspensions prepared by homogenization in PBS solution. After washing the cells in PBS by centrifugation, the cells were re-suspended in the culture medium at a density of 5×10^6 cells/ml. The cells were subsequently distributed to 24-well culture plates in 1 ml aliquots and 0.05 ml of 0.005% SRBC was added as the antigen to each well. The compounds were added to the cultures at the beginning of the four-day incubation period at concentration ranges of 1-100 µg/ml. The number of antibody-forming cells (AFC) in the cultures was determined using a method of local hemolysis in agar gel according to Mishell et al., *J Exp Med*, 1967, 126, 423-442.

[00145] Toxicity test: Splenocytes, at a density of 2×10^5 cells/100µl/well, re-suspended in the culture medium, were cultured for 24 h in a cell culture incubator with the compounds (1-100 µg/ml). Cell survival was determined by MTT colorimetric method. The results were presented as mean optical density (OD) at 550 nm from 4 wells. The viability of cells in respective compound concentrations was compared to appropriate DMSO control groups (100% survival), corresponding to respective compound concentrations.

Results

[00146] At 50-100 µg/ml concentrations, compound I-C showed a strong inhibitory effect on concanavalin A-induced mouse splenocyte proliferation. At 100µg/m, this compound showed 70% toxicity to splenocytes. At 10 µg/m and 100 µg/m, compound I-C showed 33% and 80% suppression, respectively, in the model of *in vitro* humoral immune response to SRBC in mouse splenocyte cultures . In the model of delayed type hypersensitivity to ovalbumin,


compound I-C showed 26.9% inhibition at the dose of 100 μ g, compared to 72.7% inhibition by compound I-A.

[00147] At 100 μ g/ml concentration, compound I-B demonstrated a weak antiproliferative effect on concanavalin A-induced splenocyte proliferation; no such effect was observed at lower concentrations. Compound I-B had 30% toxicity at this concentration.

[00148] Although the foregoing invention has been described in some detail for purposes of illustration, it will be readily apparent to one skilled in the art that changes and modifications may be made without departing from the scope of the invention described herein.

WHAT IS CLAIMED IS:

1. A compound having the formula I:



wherein

k, m, n and p are each independently 0, 1 or 2;

R and R' are each independently selected from H and C₁₋₃ alkyl, or, when taken together, R and R' are -CR¹R^{1'}-X-CH₂-, wherein CR¹R^{1'} is attached to the backbone nitrogen, R¹ and R^{1'} are each independently selected from H and C₁₋₃ alkyl, and X is selected from -CH₂-, -CH₂CH₂-, -CH(OH)-, -O-, -S- and -NH-;

R'' and R''' are each independently selected from H and C₁₋₃ alkyl, or, when taken together, R'' and R''' are -CR²R^{2'}-X'-CH₂-, wherein CR²R^{2'} is attached to the backbone nitrogen, R² and R^{2'} are each independently selected from H and C₁₋₃ alkyl, and X' is selected from -CH₂-, -CH₂CH₂-, -CH(OH)-, -O-, -S- and -NH-; and


R³ and R⁴ are each independently selected from aryl, substituted aryl, heteroaryl and substituted heteroaryl;

or a pharmaceutically acceptable salt thereof.


2. A compound according to claim 1 wherein at least one of R³ and R⁴ is phenyl.
3. A compound according to claim 1 or claim 2 wherein at least one of R³ and R⁴ is 4-hydroxyphenyl.
4. A compound according to claim 1 or claim 2 wherein at least one of R³ and R⁴ is 4-*t*-butoxyphenyl.
5. A compound according to claim 1 or claim 2 wherein at least one of R³ and R⁴ is 2-indolyl.
6. A compound according to claim 1 or claim 2 wherein R³ and R⁴ are both phenyl.
7. A compound according to claim 1 or claim 2 wherein one of R³ and R⁴ is phenyl and the other of R³ and R⁴ is 4-hydroxyphenyl.

8. A compound according to claim 1 or claim 2 wherein one of R³ and R⁴ is phenyl and the other of R³ and R⁴ is 4-*t*-butoxyphenyl.
9. A compound according to claim 1 or claim 2 wherein one of R³ and R⁴ is phenyl and the other of R³ and R⁴ is 2-indolyl.
10. A compound according to any of claims 1 to 9 wherein the carbon to which -CH₂-R³ is attached has absolute (*R*)-stereochemistry.
11. A compound according to any of claims 1 to 9 wherein the carbon to which -CH₂-R³ is attached has absolute (*S*)-stereochemistry.
12. A compound according to any of claims 1 to 11 wherein the carbon to which -CH₂-R⁴ is attached has absolute (*R*)-stereochemistry.
13. A compound according to any of claims 1 to 11 wherein the carbon to which -CH₂-R⁴ is attached has absolute (*S*)-stereochemistry.
14. A compound according to any of claims 1 to 13 wherein one of k, m, n and p is 1, and the remainder of k, m, n and p are 0.
15. A compound according to any of claims 1 to 13 wherein two of k, m, n and p are 1, and the remainder of k, m, n and p are 0.
16. A compound according to any of claims 1 to 15 wherein at least one of k and m is not 0.
17. A compound according to any of claims 1 to 15 wherein at least one of n and p is not 0.
18. A compound according to any of claims 1 to 17 wherein at least one of k and m is not 0 and at least one of n and p is not 0.
19. A compound according to any of claims 1 to 15 wherein both k and n are 0.
20. A compound according to any of claims 1 to 14 wherein both k and n are 0, one of m and p is 0, and the other of m and p is 1.
21. A compound according to claim 15 wherein both k and n are 0 and both m and p are 1.
22. A compound according to any of claims 1 to 21 wherein all four amino acids are L-amino acids.
23. A compound according to any of claims 1 to 21 wherein three of the amino acids are L-amino acids and one of the amino acids is a D-amino acid.
24. A compound according to any of claims 1 to 21 wherein two of the amino acids are L-amino acids and two of the amino acids are D-amino acids.
25. A compound according to any of claims 1 to 21 wherein one of the amino acids is an L-amino acid and three of the amino acids are D-amino acids.
26. A compound according to any of claims 1 to 21 wherein all four amino acids are D-amino acids.


27. A compound according to any of claims 1 to 26 wherein R and R' are taken together to form -(CH₂)₃-.
28. A compound according to any of claims 1 to 27 wherein R'' and R''' are taken together to form -(CH₂)₃-.
29. A compound according to any of claims 1 to 26 wherein R and R' are taken together to form -(CH₂)₄-.
30. A compound according to any of claims 1 to 27 wherein R'' and R''' are taken together to form -(CH₂)₄-.
31. A compound according to any of claims 1 to 27 wherein R and R' are taken together to form -CH₂-CH(OH)-CH₂-.
32. A compound according to claim 31 wherein the carbon to which the OH is attached has absolute (S)- stereochemistry.
33. A compound according to claim 31 wherein the carbon to which the OH is attached has absolute (R)- stereochemistry.
34. A compound according to any of claims 1 to 27 wherein R'' and R''' are taken together to form -CH₂-CH(OH)-CH₂-.
35. A compound according to claim 34 wherein the carbon to which the OH is attached has absolute (S)- stereochemistry.
36. A compound according to claim 34 wherein the carbon to which the OH is attached has absolute (R)- stereochemistry.
37. A compound according to any of claims 1 to 21 and 27 to 36 wherein R and R' are taken together and the carbon at which R' is attached has absolute (S)-stereochemistry.
38. A compound according to any of claims 1 to 21 and 27 to 36 wherein R and R' are taken together and the carbon at which R' is attached has absolute (R)-stereochemistry.
39. A compound according to any of claims 1 to 21 and 27 to 36 wherein R'' and R''' are taken together and the carbon at which R''' is attached has absolute (S)- stereochemistry.
40. A compound according to any of claims 1 to 21 and 27 to 36 wherein R'' and R''' are taken together and the carbon at which R''' is attached has absolute (R)- stereochemistry.
41. A compound according to any claim 1 wherein the compound is selected from the group consisting of:




I-1,




I-2,




I-3,




I-4,




I-5,



I-6,




I-7 and






I-8.


42. A compound of claim 41 which is of formula I-1.

43. A compound of claim 41 which is of formula I-2.
44. A compound of claim 41 which is of formula I-3.
45. A compound of claim 41 which is of formula I-4.
46. A compound of claim 41 which is of formula I-5.
47. A compound of claim 41 which is of formula I-6.
48. A compound of claim 41 which is of formula I-7.
49. A compound of claim 41 which is of formula I-8.
50. A compound according to claim 1 or claim 41 which is selected from the group



consisting of:





51. A compound according to claim 50 which is of formula I-A.
52. A compound according to claim 50 which is of formula I-B.
53. A compound according to claim 50 which is of formula I-C.
54. A compound according to claim 50 which is of formula I-D.
55. A compound according to claim 50 which is of formula I-E.
56. A compound according to claim 50 which is of formula I-F.
57. A compound according to claim 50 which is of formula I-G.
58. A compound according to claim 50 which is of formula I-H.


59. A compound according to claim 50 which is of formula I-I.
60. A compound according to claim 50 which is of formula I-J.
61. A compound according to claim 50 which is of formula I-K.
62. A compound according to claim 50 which is of formula I-L.
63. A compound according to claim 50 which is of formula I-M.
64. A compound according to claim 50 which is of formula I-N.
65. A compound according to claim 50 which is of formula I-O.
66. A compound according to claim 50 which is of formula I-P.
67. A compound according to any of claims 1 to 66 wherein one or more amino groups in the compound of Formula I are in protected form.
68. A pharmaceutical composition comprising a compound of formula I according to any of claims 1 to 66 and a pharmaceutically acceptable carrier, excipient or diluent therefor.
69. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-1.
70. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-2.
71. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-3.
72. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-4.
73. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-5.
74. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-6.
75. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-7.
76. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-8.
77. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-A.
78. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-B.

79. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-C.
80. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-D.
81. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-E.
82. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-F.
83. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-G.
84. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-H.
85. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-I.
86. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-J.
87. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-K.
88. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-L.
89. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-M.
90. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-N.
91. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-O.
92. A pharmaceutical composition according to claim 68 wherein the compound is a compound of formula I-P.
93. A method of suppressing immune response in a patient, comprising administering to a patient in need thereof an efficacious amount of a compound of formula I.
94. A method according to claim 93 wherein the immune response which is suppressed is inflammation.
95. A method according to claim 93 wherein the immune response which is suppressed is transplant rejection.

96. A method of treating or preventing an immune-mediated disease or condition in a patient, comprising administering to a patient in need thereof an efficacious amount of a compound of formula I.
97. A method according to claim 96 wherein the immune-mediated disease or condition is selected from the group consisting of auto-immune diseases, inflammation processes, transplant rejection, and allergic reactions.
98. A method according to claim 96 wherein the immune-mediated disease or condition is selected from Psoriasis, lichen planus and other papulosquamous disorders.
99. A method according to claim 96 wherein the immune-mediated disease or condition is selected from eczema and dermatitis.
100. A method according to claim 99 wherein the eczema or dermatitis is selected from eczema, atopic eczema, seborrheic dermatitis, and pompholyx.
101. A method according to claim 96 wherein the immune-mediated disease or condition is a skin reaction to light.
102. A method according to claim 96 wherein the immune-mediated disease or condition selected from non-specific skin irritation and insect bite.
103. A method according to claim 96 wherein the immune-mediated disease or condition is a urticaria.
104. A method according to claim 96 wherein the immune-mediated disease or condition is selected from the group consisting of a primary skin tumor (e.g. melanoma); rheumatoid arthritis (both autoimmune and elicited by infection); Crohn's disease; inflammatory bowel disease; irritable bowel syndrome; a neurodegenerative disease (e.g. multiple sclerosis); Parkinson's disease; Graft-versus-Host reaction; severe psoriasis; and atopic dermatitis.
105. A method for lowering the toxicity profile of a drug, comprising administering a compound of formula I in conjunction with said drug.
106. A method according to claim 105 wherein the compound of formula I is administered in conjunction with a chemotherapeutic drug.
107. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-1.
108. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-2.
109. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-3.


110. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-4.
111. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-5.
112. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-6.
113. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-7.
114. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-8.
115. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-A.
116. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-B.
117. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-C.
118. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-D.
119. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-E.
120. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-F.
121. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-G.
122. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-H.
123. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-I.
124. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-J.
125. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-K.
126. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-L.


127. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-M.
128. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-N.
129. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-O.
130. A method according to any of claims 93 to 106 wherein the compound is a compound of formula I-P.
131. A kit, comprising a compound of formula I according to any of claims 1 to 58 and instructions for using the compound (a) suppress an immune response in a patient, (b) treat or prevent an immune-mediated disease or condition in a patient, or (c) lower the toxicity profile of a second drug.
132. A kit according to claim 131 wherein the immune response is selected from the group consisting of inflammation, transplant rejection, and infection.
133. A kit according to claim 131 wherein the immune-mediated disease or condition is selected from the group consisting of auto-immune diseases, inflammation processes, transplant rejection, and allergic reactions.
134. A kit according to claim 131 wherein the immune-mediated disease or condition selected from the group consisting of Psoriasis, lichen planus and other papulosquamous disorders; eczema and dermatitis; a skin reaction to light; non-specific skin irritation and insect bite; a urticaria; a primary skin tumor (e.g. melanoma); rheumatoid arthritis (both autoimmune and elicited by infection); Crohn's disease; inflammatory bowel disease; irritable bowel syndrome; a neurodegenerative disease (e.g. multiple sclerosis); Parkinson's disease; Graft-versus-Host reaction; severe psoriasis; and atopic dermatitis.
135. A kit according to claim 134 wherein the eczema or dermatitis is selected from eczema, atopic eczema, seborrheic dermatitis, and pompholyx.
136. A kit according to any of claims 131 to 135 wherein the instructions instruct to administer the compound of formula I in conjunction with a chemotherapeutic drug.
137. A method for making a compound of formula I as defined in claim 1, comprising cyclizing a compound having the formula II-1, II-2, II-3 or II-4 (collectively referred to hereinafter as compounds of formula II):




to the corresponding compound of formula I.

138. A method according to claim 137 which further comprises synthesizing the compound of formula II-1, II-2, II-3 or II-4.
139. A method according to claim 138 wherein the compound of formula II-1, II-2, II-3 or II-4 is synthesized by solid-phase synthesis.
140. A compound selected from the group consisting of formulae II-1, II-2, II-3 and II-4.
141. A compound according to claim 140 which is a compound of formula II-1.
142. A compound according to claim 140 which is a compound of formula II-2.
143. A compound according to claim 140 which is a compound of formula II-3.
144. A compound according to claim 140 which is a compound of formula II-4A.
145. A compound according to claim 140 which is selected from the group consisting of







II-2-a,




II-3-a




II-4-b,




II-1-b,




II-2-b,




II-3-b,




II-4-c,




II-1-c,







II-2-e,




II-3-e,




II-4-f,




II-1-f,




II-2-f,




II-3-f,



II-4-g,



II-1-g,



146. A compound according to claim 145 which is of formula II-1-a.
147. A compound according to claim 145 which is of formula II-2-a.
148. A compound according to claim 145 which is of formula II-3-a.
149. A compound according to claim 145 which is of formula II-4-a.
150. A compound according to claim 145 which is of formula II-1-b.
151. A compound according to claim 145 which is of formula II-2-b.
152. A compound according to claim 145 which is of formula II-3-b.
153. A compound according to claim 145 which is of formula II-4-b.
154. A compound according to claim 145 which is of formula II-1-c.
155. A compound according to claim 145 which is of formula II-2-c.
156. A compound according to claim 145 which is of formula II-3-c.
157. A compound according to claim 145 which is of formula II-4-c.

158. A compound according to claim 145 which is of formula II-1-d.
159. A compound according to claim 145 which is of formula II-2-d.
160. A compound according to claim 145 which is of formula II-3-d.
161. A compound according to claim 145 which is of formula II-4-d.
162. A compound according to claim 145 which is of formula II-1-e.
163. A compound according to claim 145 which is of formula II-2-e.
164. A compound according to claim 145 which is of formula II-3-e.
165. A compound according to claim 145 which is of formula II-4-e.
166. A compound according to claim 145 which is of formula II-1-f.
167. A compound according to claim 145 which is of formula II-2-f.
168. A compound according to claim 145 which is of formula II-3-f.
169. A compound according to claim 145 which is of formula II-4-f.
170. A compound according to claim 145 which is of formula II-1-g.
171. A compound according to claim 145 which is of formula II-2-g.
172. A compound according to claim 145 which is of formula II-3-g.
173. A compound according to claim 145 which is of formula II-4-g.
174. A compound according to claim 145 which is of formula II-1-h.
175. A compound according to claim 145 which is of formula II-2-h.
176. A compound according to claim 145 which is of formula II-3-h.
177. A compound according to claim 145 which is of formula II-4-h.
178. A compound according to claim 140 which is selected from the group consisting of formulae II-A-1, II-A-2, II-A-3, II-A-4, II-B-1, II-B-2, II-B-3, II-B-4, II-C-1, II-C-2, II-C-3, II-C-4, II-D-1, II-D-2, II-D-3, II-D-4, II-E-1, II-E-2, II-E-3, II-E-4, II-F-1, II-F-2, II-F-3, II-F-4, II-G-1, II-G-2, II-G-3, II-G-4, II-H-1, II-H-2, II-H-3, II-H-4, II-J-1, II-J-2, II-J-3, II-J-4, II-K-1, II-K-2, II-K-3, II-K-4, II-L-1, II-L-2, II-L-3, II-L-4, II-M-1, II-M-2, II-M-3, II-M-4, II-N-1, II-N-2, II-N-3, II-N-4, II-O-1, II-O-2, II-O-3, II-O-4, II-P-1, II-P-2, II-P-3 and II-P-4.
179. A compound according to claim 178 which is of formula II-A-1.
180. A compound according to claim 178 which is of formula II-A-2.
181. A compound according to claim 178 which is of formula II-A-3.
182. A compound according to claim 178 which is of formula II-A-4.
183. A compound according to claim 178 which is of formula II-B-1.
184. A compound according to claim 178 which is of formula II-B-2.
185. A compound according to claim 178 which is of formula II-B-3.

186. A compound according to claim 178 which is of formula II-B-4.
187. A compound according to claim 178 which is of formula II-C-1.
188. A compound according to claim 178 which is of formula II-C-2.
189. A compound according to claim 178 which is of formula II-C-3.
190. A compound according to claim 178 which is of formula II-C-4.
191. A compound according to claim 178 which is of formula II-D-1.
192. A compound according to claim 178 which is of formula II-D-2.
193. A compound according to claim 178 which is of formula II-D-3.
194. A compound according to claim 178 which is of formula II-D-4.
195. A compound according to claim 178 which is of formula II-E-1.
196. A compound according to claim 178 which is of formula II-E-2.
197. A compound according to claim 178 which is of formula II-E-3.
198. A compound according to claim 178 which is of formula II-E-4.
199. A compound according to claim 178 which is of formula II-F-1.
200. A compound according to claim 178 which is of formula II-F-2.
201. A compound according to claim 178 which is of formula II-F-3.
202. A compound according to claim 178 which is of formula II-F-4.
203. A compound according to claim 178 which is of formula II-G-1.
204. A compound according to claim 178 which is of formula II-G-2.
205. A compound according to claim 178 which is of formula II-G-3.
206. A compound according to claim 178 which is of formula II-G-4.
207. A compound according to claim 178 which is of formula II-H-1.
208. A compound according to claim 178 which is of formula II-H-2.
209. A compound according to claim 178 which is of formula II-H-3.
210. A compound according to claim 178 which is of formula II-H-4.
211. A compound according to claim 178 which is of formula II-J-1.
212. A compound according to claim 178 which is of formula II-J-2.
213. A compound according to claim 178 which is of formula II-J-3.
214. A compound according to claim 178 which is of formula II-J-4.
215. A compound according to claim 178 which is of formula II-K-1.
216. A compound according to claim 178 which is of formula II-K-2.
217. A compound according to claim 178 which is of formula II-K-3.
218. A compound according to claim 178 which is of formula II-K-4.
219. A compound according to claim 178 which is of formula II-L-1.

220. A compound according to claim 178 which is of formula II-L-2.
221. A compound according to claim 178 which is of formula II-L-3.
222. A compound according to claim 178 which is of formula II-L-4.
223. A compound according to claim 178 which is of formula II-M-1.
224. A compound according to claim 178 which is of formula II-M-2.
225. A compound according to claim 178 which is of formula II-M-3.
226. A compound according to claim 178 which is of formula II-M-4.
227. A compound according to claim 178 which is of formula II-N-1.
228. A compound according to claim 178 which is of formula II-N-2.
229. A compound according to claim 178 which is of formula II-N-3.
230. A compound according to claim 178 which is of formula II-N-4.
231. A compound according to claim 178 which is of formula II-O-1.
232. A compound according to claim 178 which is of formula II-O-2.
233. A compound according to claim 178 which is of formula II-O-3.
234. A compound according to claim 178 which is of formula II-O-4.
235. A compound according to claim 178 which is of formula II-P-1.
236. A compound according to claim 178 which is of formula II-P-2.
237. A compound according to claim 178 which is of formula II-P-3.
238. A compound according to claim 178 which is of formula II-P-4.
239. A compound according to any of claims 140 to 238 in which one or more amino groups is in protected form.
240. A compound according to any of claims 140 to 238 in which no amino groups are in protected form.
241. A compound according to any of claims 140 to 240 which is bound to a solid-phase resin.
242. A compound according to claim 241 which is bound to the solid-phase resin directly.
243. A compound according to claim 241 which is bound to the solid-phase resin through a linker.

1/8




FIG. 1




FIG. 2




Fig. 3

2/8




Fig. 4




Fig 5.

3/8




Fig. 6.




Fig. 7.




Fig. 8A




Fig. 8B

5/8




Fig. 9A




Fig. 9B

6/8




Fig. 10A




Fig. 10B

7/8




Fig. 11




Fig. 12




Fig. 13

8/8




Fig. 14




Fig. 15