



(86) Date de dépôt PCT/PCT Filing Date: 2002/06/05

(87) Date publication PCT/PCT Publication Date: 2002/12/12

(45) Date de délivrance/Issue Date: 2009/08/11

(85) Entrée phase nationale/National Entry: 2003/12/02

(86) N° demande PCT/PCT Application No.: GB 2002/002517

(87) N° publication PCT/PCT Publication No.: 2002/098426

(30) Priorités/Priorities: 2001/06/06 (GB0113708.2);
2001/08/10 (US60/311,502)

(51) Cl.Int./Int.Cl. *C07D 413/06* (2006.01),
A61K 31/472 (2006.01), *A61K 31/4725* (2006.01),
A61P 11/06 (2006.01), *A61P 29/00* (2006.01),
C07D 217/04 (2006.01), *C07D 401/06* (2006.01),
C07D 413/14 (2006.01)


(72) Inventeurs/Inventors:
FENTON, GARRY, GB;
HARRIS, NEIL VICTOR, GB

(73) Propriétaire/Owner:
AVENTIS PHARMA LIMITED, GB

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L., S.R.L.

(54) Titre : TETRAHYDROISOQUINOLINES SUBSTITUEES DESTINEES AU TRAITEMENT DES MALADIES
INFLAMMATOIRES

(54) Title: SUBSTITUTED TETRAHYDROISOQUINOLINES FOR USE IN THE TREATMENT OF INFLAMMATORY
DISEASES



(I)

(57) Abrégé/Abstract:

The invention is directed to physiologically active compounds of formula (I): wherein: R¹ represents optionally substituted aryl, optionally substituted heteroaryl, R³NH-Ar¹-L²- or R³-NH-C(=O)-NH-Ar²-L²-; R³ represents aryl or heteroaryl; Ar¹ represents a saturated, partially saturated or fully unsaturated 8- to 10-membered bicyclic ring system containing at least one heteroatom

(57) Abrégé(suite)/Abstract(continued):

selected from O, S or n; Ar² represents arylidiyl or heteroaryldiyl; L¹ represents a linkage, such as an alkylene linkage; L² represents an alkylene chain linkage; R² represents hydrogen, halogen, C₁₋₄ alkyl or C₁₋₄ alkoxy; and Y is carboxy or an acid bioisostere; but excluding compounds where an oxygen, nitrogen or sulfur atom is attached directly to a carbon carbon multiple bond of an alkenylene or alkynylene residue; and the corresponding N-oxides and ester prodrugs thereof, and the pharmaceutically acceptable salts and solvates of such compounds, and the N-oxides and ester prodrugs thereof. Such compounds have valuable pharmaceutical properties, in particular the ability to regulate the interaction of VCAM-1 and fibronectin with the integrin VLA-4($\alpha 4 \beta 1$).

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 December 2002 (12.12.2002)

PCT

(10) International Publication Number
WO 02/098426 A1

(51) International Patent Classification⁷: **A61K 31/472**,
31/4725, A61P 29/00, C07D 217/16, 413/06, 413/14

(21) International Application Number: PCT/GB02/02517

(22) International Filing Date: 5 June 2002 (05.06.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0113708.2 6 June 2001 (06.06.2001) GB
60/311,502 10 August 2001 (10.08.2001) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **AVENTIS PHARMA LIMTIED** [GB/GB]; Aventis House, 50 Kings Hill Avenue, West Malling, Kent ME19 4AH (GB).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FENTON, Garry** [GB/GB]; Aventis Pharma Limited, Aventis House, 50 Kings Hill Avenue, West Malling, Kent ME19 4AH (GB). **HARRIS, Neil, Victor** [GB/GB]; Aventis Pharma Limited, Aventis House, 50 Kings Hill Avenue, West Malling, Kent ME19 4AH (GB).


Published:

— with international search report
— with amended claims

(74) Agent: **JONES, Stephen, Anthony**; Adamson Jones, Broadway Business Centre, 32a Stoney Street, Nottingham NG1 1LL (GB).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED TETRAHYDROISOQUINOLINES FOR USE IN THE TREATMENT OF INFLAMMATORY DISEASES



(57) Abstract: The invention is directed to physiologically active compounds of formula (I); wherein: R¹ represents optionally substituted aryl, optionally substituted heteroaryl, R³NH-Ar¹-L²- or R³-NH-C(=O)-NH-Ar²-L²-; R³ represents aryl or heteroaryl; Ar¹ represents a saturated, partially saturated or fully unsaturated 8- to 10-membered bicyclic ring system containing at least one heteroatom selected from O, S or n; Ar² represents aryldiyl or heteroaryldiyl; L¹ represents a linkage, such as an alkylene linkage; L² represents an alkylene chain linkage; R² represents hydrogen, halogen, C₁₋₄alkyl or C₁₋₄alkoxy; and Y is carboxy or an acid bioisostere; but excluding compounds where an oxygen, nitrogen or sulfur atom is attached directly to a carbon carbon multiple bond of an alkenylene or alkynylene residue; and the corresponding N-oxides and ester prodrugs thereof, and the pharmaceutically acceptable salts and solvates of such compounds, and the N-oxides and ester prodrugs thereof. Such compounds have valuable pharmaceutical properties, in particular the ability to regulate the interaction of VCAM-1 and fibronectin with the integrin VLA-4(α4β1).

SUBSTITUTED TETRAHYDROISOQUINOLINES FOR USE IN THE TREATMENT OF INFLAMMATORY DISEASES

This invention is directed to tetrahydroisoquinoline derivatives, their preparation, pharmaceutical compositions containing these compounds, and their pharmaceutical use in the treatment of disease states capable of being modulated by the inhibition of cell adhesion.

Cell adhesion is a process by which cells associate with each other, migrate towards a specific target or localise within the extra-cellular matrix. Many of the cell-cell and cell-extracellular matrix interactions are mediated by protein ligands (e.g., fibronectin, VCAM-1 and vitronectin) and their integrin receptors [e.g., $\alpha 5 \beta 1$ (VLA-5), $\alpha 4 \beta 1$ (VLA-4) and $\alpha V \beta 3$]. Recent studies have shown these interactions to play an important part in many physiological (e.g., embryonic development and wound healing) and pathological conditions (e.g., tumour-cell invasion and metastasis, inflammation, atherosclerosis and autoimmune disease).

A wide variety of proteins serve as ligands for integrin receptors. In general, the proteins recognised by integrins fall into one of three classes: extracellular matrix proteins, plasma proteins and cell surface proteins. Extracellular matrix proteins such as collagen fibronectin, fibrinogen, laminin, thrombospondin and vitronectin bind to a number of integrins. Many of the adhesive proteins also circulate in plasma and bind to activated blood cells. Additional components in plasma that are ligands for integrins include fibrinogen and factor X. Cell bound complement C3bi and several transmembrane proteins, such as Ig-like cell adhesion molecule (ICAM-1,2,3) and vascular cell adhesion molecule (VCAM-1), which are members of the Ig superfamily, also serve as cell-surface ligands for some integrins.

Integrins are heterodimeric cell surface receptors consisting of two subunits called α and β . There are at least fifteen different α -subunits ($\alpha 1$ - $\alpha 9$, α -L, α -M, α -X, α -IIb, α -V and α -E) and at least seven different β ($\beta 1$ - $\beta 7$) subunits. The integrin family can be subdivided into classes based on the β subunits, which can be associated with one or more α -subunits. The most widely distributed integrins belong to the $\beta 1$ class, also known as the very late antigens (VLA). The second class of integrins are leukocyte specific receptors and consist of one of three α -subunits (α -L, α -M or α -X) complexed with the $\beta 2$ protein. The cytoadhesins α -IIb $\beta 3$ and α -V $\beta 3$, constitute the third class of integrins.

The present invention principally relates to agents which modulate the interaction of the ligand VCAM-1 with its integrin receptor $\alpha 4 \beta 1$ (VLA-4), which is expressed on numerous hematopoietic


cells and established cell lines, including hematopoietic precursors, peripheral and cytotoxic T lymphocytes, B lymphocytes, monocytes, thymocytes and eosinophils.

The integrin $\alpha 4\beta 1$ mediates both cell-cell and cell-matrix interactions. Cells expressing $\alpha 4\beta 1$ bind to the carboxy-terminal cell binding domain (CS-1) of the extracellular matrix protein fibronectin, to the cytokine-inducible endothelial cell surface protein VCAM-1, and to each other to promote homotypic aggregation. The expression of VCAM-1 by endothelial cells is upregulated by proinflammatory cytokines such as INF- γ , TNF- α , IL-1 β and IL-4.

Regulation of $\alpha 4\beta 1$ mediated cell adhesion is important in numerous physiological processes, including T-cell proliferation, B-cell localisation to germinal centres, and adhesion of activated T-cells and eosinophils to endothelial cells. Evidence for the involvement of VLA-4/VCAM-1 interaction in various disease processes such as melanoma cell division in metastasis, T-cell infiltration of synovial membranes in rheumatoid arthritis, autoimmune diabetes, colitis and leukocyte penetration of the blood-brain barrier in experimental autoimmune encephalomyelitis, atherosclerosis, peripheral vascular disease, cardiovascular disease and multiple sclerosis, has been accumulated by investigating the role of the peptide CS-1 (the variable region of fibronectin to which $\alpha 4\beta 1$ binds via the sequence Leu-Asp-Val) and antibodies specific for VLA-4 or VCAM-1 in various in vitro and in vivo experimental models of inflammation. For example, in a Streptococcal cell wall-induced experimental model of arthritis in rats, intravenous administration of CS-1 at the initiation of arthritis suppresses both acute and chronic inflammation (S.M.Wahl et al., J.Clin.Invest., 1994, 94, pages 655-662). In the oxazalone-sensitised model of inflammation (contact hypersensitivity response) in mice, intravenous administration of anti- $\alpha 4$ specific monoclonal antibodies significantly inhibited (50-60% reduction in the ear swelling response) the efferent response (P.L.Chisholm et al. J.Immunol., 1993, 23, pages 682-688). In a sheep model of allergic bronchoconstriction, HP1/2, an anti- $\alpha 4$ monoclonal antibody given intravenously or by aerosol, blocked the late response and the development of airway hyperresponsiveness (W.M. Abraham et al. J. Clin. Invest., 1994, 93 pages 776-787).

We have now found a novel group of tetrahydroisoquinolines which have valuable pharmaceutical properties, in particular the ability to regulate the interaction of VCAM-1 and fibronectin with the integrin VLA-4 ($\alpha 4\beta 1$).

Thus, in one aspect, the present invention is directed to aza-bicycles of general formula (I):-



(I)

wherein:-

- 5 R^1 represents optionally substituted aryl, optionally substituted heteroaryl, $R^3\text{NH-Ar}^1\text{-L}^2\text{-}$ or $R^3\text{-NH-C(=O)-NH-Ar}^2\text{-L}^2\text{-}$;
 R^2 represents hydrogen, halogen, C_{1-4} alkyl or C_{1-4} alkoxy;
 R^3 represents optionally substituted aryl or optionally substituted heteroaryl;
 R^4 is alkyl, aryl, cycloalkyl, heteroaryl or heterocycloalkyl, or alkyl substituted by aryl, an acidic
 10 functional group, cycloalkyl, heteroaryl, heterocycloalkyl, $-\text{S(O)}_m\text{R}^5$, $-\text{C(=O)-NY}^3\text{Y}^4$ or $-\text{NY}^3\text{Y}^4$;
 R^5 represents alkyl, alkenyl, alkynyl, aryl, arylalkyl, arylalkenyl, arylalkynyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkenyl, cycloalkylalkynyl, cycloalkenyl, cycloalkenylalkyl, heteroaryl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, heterocycloalkyl or heterocycloalkylalkyl;
 R^6 is hydrogen, alkyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl,
 15 heterocycloalkyl or heterocycloalkylalkyl;
 R^7 is hydrogen, R^5 or alkyl substituted with alkoxy, cycloalkyl, hydroxy, mercapto, alkylthio or $-\text{NY}^3\text{Y}^4$;
 R^8 is hydrogen or C_{1-4} alkyl;
 R^9 and R^{11} are each independently selected from hydrogen or a group consisting of amino acid side
 20 chains, an acidic functional group, R^5 , $-\text{C(=O)-R}^5$, or $-\text{C(=O)-NY}^3\text{Y}^4$, or alkyl substituted by an acidic functional group or by R^5 , $-\text{NY}^3\text{Y}^4$, $-\text{NH-C(=O)-R}^5$, $-\text{C(=O)-R}^{12}\text{-NH}_2$, $-\text{C(=O)-Ar}^2\text{-NH}_2$, $-\text{C(=O)-R}^{12}\text{-CO}_2\text{H}$, or $-\text{C(=O)-NY}^3\text{Y}^4$;
 or R^7 and R^9 together with the atoms to which they attached form a 3- to 6-membered heterocycloalkyl ring;
 25 R^{10} represents C_{1-6} alkylene, optionally substituted by R^4 ;
 R^{12} is an alkylene chain, an alkenylene chain, or an alkynylene chain;
 R^{13} is alkyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl or heterocycloalkylalkyl;

Ar¹ represents a saturated, partially saturated or fully unsaturated 8 to 10 membered bicyclic ring system containing at least one heteroatom selected from O, S or N, optionally substituted by one or more aryl group substituents;

Ar² represents arylidiyl or heteroaryldiyl;

5 L¹ represents

- (i) a direct bond;
- (ii) an alkenylene, alkylene or alkynylene linkage each optionally substituted by (a) an acidic functional group, cyano, oxo, -S(O)_mR⁴, R⁵, -C(=O)-R⁵, -C(=O)-OR⁵, -N(R⁶)-C(=O)-R⁴, -N(R⁶)-C(=O)-OR⁴, -N(R⁶)-SO₂-R⁴, -NY³Y⁴ or
 10 -[C(=O)-N(R⁷)-C(R⁸)(R⁹)]_p-C(=O)-NY³Y⁴, or by (b) alkyl substituted by an acidic functional group, or by S(O)_mR⁴, -C(=O)-NY³Y⁴ or -NY³Y⁴;
- (iii) a -[C(=O)-N(R⁷)-C(R⁸)(R⁹)]_p- linkage;
- (iv) a -Z¹-R¹⁰- linkage;
- (v) a -R¹⁰-Z¹-R¹⁰- linkage;
- 15 (vi) a -C(R⁸)(R¹¹)-[C(=O)-N(R⁷)-C(R⁸)(R⁹)]_p- linkage; or
- (vii) a -L³-L⁴-L⁵- linkage;

L² represents an alkylene chain;

L³ and L⁵ each independently represent a direct bond or an alkylene chain;

L⁴ represents a cycloalkylene or heterocycloalkylene linkage;

20 Y¹ and Y² are independently hydrogen, alkenyl, alkyl, aryl, arylalkyl, cycloalkyl, heteroaryl or heteroarylalkyl; or the group -NY¹Y² may form a cyclic amine;

Y³ and Y⁴ are independently hydrogen, alkenyl, alkyl, alkynyl, aryl, cycloalkenyl, cycloalkyl, heteroaryl, heterocycloalkyl, or alkyl substituted by alkoxy, aryl, cyano, cycloalkyl, heteroaryl, heterocycloalkyl, hydroxy, oxo, -NY¹Y², or one or more -CO₂R⁶ or -C(=O)-NY¹Y² groups; or the

25 group -NY³Y⁴ may form a 5- to 7-membered cyclic amine which (i) may be optionally substituted with one or more substituents selected from alkoxy, carboxamido, carboxy, hydroxy, oxo (or a 5-, 6- or 7-membered cyclic acetal derivative thereof), R⁷; (ii) may also contain a further heteroatom selected from O, S, SO₂, or NY⁵; and (iii) may also be fused to additional aryl, heteroaryl, heterocycloalkyl or cycloalkyl rings to form a bicyclic or tricyclic ring system;

30 Y⁵ is hydrogen, alkyl, aryl, arylalkyl, -C(=O)-R¹³, -C(=O)-OR¹³ or -SO₂R¹³;

- 5 -

Z^1 is O, $S(O)_n$, NR^8 , SO_2NR^8 , $C(=O)NR^8$ or $C(=O)$;

Y is carboxy or an acid bioisostere;

m is an integer 1 or 2;

n is zero or an integer 1 or 2; and

5 p is zero or an integer 1 to 4;

but excluding compounds where an oxygen, nitrogen or sulfur atom is attached directly to a carbon carbon multiple bond of an alkenylene or alkynylene residue;

and the corresponding N-oxides, and their prodrugs; and pharmaceutically acceptable salts and solvates (e.g., hydrates) of such compounds and their N-oxides and prodrugs.

10

In the present specification, the term "compounds of the invention", and equivalent expressions, are meant to embrace compounds of general formula (I) as hereinbefore described, which expression includes the prodrugs, protected derivatives of compounds of formula (I) containing one or more acidic functional groups and/or amino-acid side chains, the pharmaceutically acceptable salts, and the solvates, e.g., hydrates, where the context so permits. Similarly, reference to intermediates, whether or not they themselves are claimed, is meant to embrace their salts, and solvates, where the context so permits. For the sake of clarity, particular instances when the context so permits are sometimes indicated in the text, but these instances are purely illustrative and it is not intended to exclude other instances when the context so permits.

20

As used above, and throughout the description of the invention, the following terms, unless otherwise indicated, shall be understood to have the following meanings:-

"Patient" includes both human and other mammals.

25

"Acid bioisostere" means a group which has chemical and physical similarities producing broadly similar biological properties to a carboxy group

30

Examples of suitable acid bioisosteres include: $-C(=O)-NHOH$, $-C(=O)-CH_2OH$, $-C(=O)-CH_2SH$, $-C(=O)-NH-CN$, sulfo, phosphono, alkylsulfonylcarbamoyl, tetrazolyl, arylsulfonylcarbamoyl, heteroarylsulfonylcarbamoyl, N-methoxycarbamoyl, 3-hydroxy-3-cyclobutene-1,2-dione,

35

3,5-dioxo-1,2,4-oxadiazolidinyl or heterocyclic phenols such as 3-hydroxyisoxazolyl and 3-hydroxy-1-methylpyrazolyl.

"Acidic functional group" means a group with an acidic hydrogen within it. The "corresponding protected derivatives" are those where the acidic hydrogen atom has been replaced with a suitable protecting group. For suitable protecting groups see T.W. Greene and P.G.M. Wuts in "Protective Groups in Organic Chemistry" John Wiley and Sons, 1991. Exemplary acidic functional groups include carboxyl (and acid bioisosteres), hydroxy, mercapto and imidazole. Exemplary protected derivatives include esters of carboxy groups (i.e. $-\text{CO}_2\text{R}^{13}$), ethers of hydroxy groups (i.e. $-\text{OR}^{13}$), thioethers of mercapto groups (i.e. $-\text{SR}^{13}$), and N-benzyl derivatives of imidazoles.

"Acyl" means an H-CO- or alkyl-CO- group in which the alkyl group is as described herein.

"Acylamino" is an acyl-NH- group wherein acyl is as defined herein.

"Alkenyl" means an aliphatic hydrocarbon group containing a carbon-carbon double bond and which may be straight or branched having about 2 to about 15 carbon atoms in the chain. Preferred alkenyl groups have 2 to about 12 carbon atoms in the chain; and more preferably about 2 to about 4 carbon atoms in the chain. "Branched", as used herein and throughout the text, means that one or more lower alkyl groups such as methyl, ethyl or propyl are attached to a linear chain; here a linear alkenyl chain. "Lower alkenyl" means about 2 to about 4 carbon atoms in the chain which may be straight or branched. Exemplary alkenyl groups include ethenyl, propenyl, n-butenyl, i-butenyl, 3-methylbut-2-enyl, n-pentenyl, heptenyl, octenyl, cyclohexylbutenyl and decenyl.

"Alkenylene" means an aliphatic bivalent radical derived from a straight or branched alkenyl group, in which the alkenyl group is as described herein. Exemplary alkenylene radicals include vinylene and propylene.

"Alkoxy" means an alkyl-O- group in which the alkyl group is as described herein. Exemplary alkoxy groups include methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy and heptoxy.

"Alkoxycarbonyl" means an alkyl-O-CO- group in which the alkyl group is as described herein. Exemplary alkoxycarbonyl groups include methoxy- and ethoxycarbonyl.

"Alkyl" means, unless otherwise specified, an aliphatic hydrocarbon group which may be straight or branched having about 1 to about 15 carbon atoms in the chain optionally substituted by alkoxy or by one or more halogen atoms. Particular alkyl groups have from 1 to about 6 carbon atoms. "Lower alkyl" as a group or part of a lower alkoxy, lower alkylthio, lower alkylsulfinyl or lower alkylsulfonyl group means unless otherwise specified, an aliphatic hydrocarbon group which may be straight or branched having about 1 to about 4 carbon atoms in the chain. Exemplary alkyl groups include methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, 3-pentyl, heptyl, octyl, nonyl, decyl and dodecyl.

10 "Alkylene" means an aliphatic bivalent radical derived from a straight or branched alkyl group, in which the alkyl group is as described herein. Exemplary alkylene radicals include methylene, ethylene and trimethylene.

"Alkylenedioxy" means an -O-alkyl-O- group in which the alkyl group is as defined above. Exemplary alkylenedioxy groups include methylenedioxy and ethylenedioxy.

"Alkylsulfinyl" means an alkyl-SO- group in which the alkyl group is as previously described. Preferred alkylsulfinyl groups are those in which the alkyl group is C₁₋₄alkyl.

20 "Alkylsulfonyl" means an alkyl-SO₂- group in which the alkyl group is as previously described. Preferred alkylsulfonyl groups are those in which the alkyl group is C₁₋₄alkyl.

"Alkylsulfonylcarbamoyl" means an alkyl-SO₂-NH-C(=O)- group in which the alkyl group is as previously described. Preferred alkylsulfonylcarbamoyl groups are those in which the alkyl group is C₁₋₄alkyl.

"Alkylthio" means an alkyl-S- group in which the alkyl group is as previously described. Exemplary alkylthio groups include methylthio, ethylthio, isopropylthio and heptylthio.

30 "Alkynyl" means an aliphatic hydrocarbon group containing a carbon-carbon triple bond and which may be straight or branched having about 2 to about 15 carbon atoms in the chain. Preferred alkynyl groups have 2 to about 12 carbon atoms in the chain; and more preferably about 2 to about 4 carbon atoms in the chain. Exemplary alkynyl groups include ethynyl, propynyl, n-butynyl, i-butynyl, 3-methylbut-2-ynyl, and n-pentynyl.

"Alkynylene" means an aliphatic bivalent radical derived from a straight or branched alkynyl group, in which the alkynyl group is as described herein. Exemplary alkynylene radicals include ethynylene and propynylene.

5 "Amino acid side chains" means the substituent found on the carbon between the amino and carboxy groups in α -amino acids. For examples of "corresponding protected derivatives" of amino acid side chains, see T.W. Greene and P.G.M. Wuts in "Protective Groups in Organic Chemistry" John Wiley and Sons, 1991.

10 "Aroyl" means an aryl-CO- group in which the aryl group is as described herein. Exemplary aroyl groups include benzoyl and 1- and 2-naphthoyl.

"Aroylamino" is an aroyl-NH- group wherein aroyl is as previously defined.

15 "Aryl" as a group or part of a group denotes: (i) an optionally substituted monocyclic or multicyclic aromatic carbocyclic moiety of about 6 to about 14 carbon atoms, such as phenyl or naphthyl; or (ii) an optionally substituted partially saturated multicyclic aromatic carbocyclic moiety in which an aryl and a cycloalkyl or cycloalkenyl group are fused together to form a cyclic structure, such as a tetrahydronaphthyl, indenyl or indanyl ring. Aryl groups may be substituted with one or more aryl
 20 group substituents which may be the same or different, where "aryl group substituent" includes, for example, acyl, acylamino, alkoxy, alkoxycarbonyl, alkylenedioxy, alkylsulfinyl, alkylsulfonyl, alkylthio, aroyl, aroylamino, aryl, arylalkyloxy, arylalkyloxycarbonyl, arylalkylthio, aryloxy, aryloxycarbonyl, arylsulfinyl, arylsulfonyl, arylthio, carboxy, cyano, halo, heteroaroyl, heteroaryl, heteroarylalkyloxy, heteroaroylamino, heteroaryloxy, hydroxy, nitro, trifluoromethyl, Y^1Y^2N- ,
 25 Y^1Y^2NCO- , $Y^1Y^2NSO_2-$, $Y^1Y^2N-C_{2-6}alkylene-Z^2-$ {where Z^2 is O, NR^8 or $S(O)_n$ }, $alkylC(=O)-Y^1N-$, $alkylSO_2-Y^1N-$ or alkyl optionally substituted with aryl, heteroaryl, hydroxy, or Y^1Y^2N- .

"Arylalkenyl" means an aryl-alkenyl- group in which the aryl and alkenyl are as previously described.

30 Preferred arylalkenyls contain a lower alkenyl moiety. Exemplary arylalkenyl groups include styryl and phenylallyl.

"Arylalkyl" means an aryl-alkyl- group in which the aryl and alkyl moieties are as previously described. Preferred arylalkyl groups contain a C_{1-4} alkyl moiety. Exemplary arylalkyl groups

35 include benzyl, 2-phenethyl and naphthlenemethyl.

"Arylalkyloxy" means an arylalkyl-O- group in which the arylalkyl groups is as previously described. Exemplary arylalkyloxy groups include benzyloxy and 1- or 2-naphthalenemethoxy.

5 "Arylalkyloxycarbonyl" means an arylalkyl-O-CO- group in which the arylalkyl groups is as previously described. An exemplary arylalkyloxycarbonyl group is benzyloxycarbonyl.

"Arylalkylthio" means an arylalkyl-S- group in which the arylalkyl group is as previously described. An exemplary arylalkylthio group is benzylthio.

10

"Arylalkynyl" means an aryl-alkynyl- group in which the aryl and alkynyl are as previously described. Exemplary arylalkynyl groups include phenylethynyl and 3-phenylbut-2-ynyl.

"Aryldiyl" means an optionally substituted bivalent radical derived from an aryl group. Exemplary
15 arylene groups include optionally substituted phenylene, naphthylene and indanylene. When Ar² is arylene this may particularly represent an optionally substituted phenylene. Suitable substituents include one or more "aryl group substituents" as defined above, particularly halogen, methyl or methoxy.

20 "Aryloxy" means an aryl-O- group in which the aryl group is as previously described. Exemplary aryloxy groups include optionally substituted phenoxy and naphthoxy.

"Aryloxycarbonyl" means an aryl-O-C(=O)- group in which the aryl group is as previously described. Exemplary aryloxycarbonyl groups include phenoxycarbonyl and naphthoxycarbonyl.

25

"Arylsulfinyl" means an aryl-SO- group in which the aryl group is as previously described.

"Arylsulfonyl" means an aryl-SO₂- group in which the aryl group is as previously described.

30 "Arylsulfonylcarbamoyl" means an aryl-SO₂-NH-C(=O)- group in which the aryl group is as previously described.

"Arylthio" means an aryl-S- group in which the aryl group is as previously described. Exemplary arylthio groups include phenylthio and naphthylthio.

35

"Azaheteroaryl" means an aromatic carbocyclic moiety of about 5 to about 10 ring members in which one of the ring members is nitrogen and the other ring members are chosen from carbon, oxygen, sulfur, or nitrogen. Examples of azaheteroaryl groups include pyridyl, pyrimidinyl, quinolinyl, isoquinolinyl, quinazolinyl, imidazolyl, and benzimidazolyl.

5

"Azaheteroaryldiyl" means an optionally substituted bivalent radical derived from an azaheteroaryl group.

10

"Cyclic amine" means a 3 to 8 membered monocyclic cycloalkyl ring system where one of the ring carbon atoms is replaced by nitrogen and which (i) may optionally contain an additional heteroatom selected from O, S or NY⁶ (where Y⁶ is hydrogen, alkyl, arylalkyl, and aryl) and (ii) may be fused to additional aryl or heteroaryl ring to form a bicyclic ring system. Exemplary cyclic amines include pyrrolidine, piperidine, morpholine, piperazine, indoline and pyrindoline.

15

"Cycloalkenyl" means a non-aromatic monocyclic or multicyclic ring system containing at least one carbon-carbon double bond and having about 3 to about 10 carbon atoms. Exemplary monocyclic cycloalkenyl rings include cyclopentenyl, cyclohexenyl or cycloheptenyl.

20

"Cycloalkenylalkyl" means a cycloalkenyl-alkyl- group in which the cycloalkenyl and alkyl moieties are as previously described. Exemplary cycloalkenylalkyl groups include cyclopentenylmethyl, cyclohexenylmethyl or cycloheptenylmethyl.

25

"Cycloalkenylene" means a bivalent radical derived from an unsaturated monocyclic hydrocarbon of about 3 to about 10 carbon atoms by removing a hydrogen atom from each of two different carbon atoms of the ring. Exemplary cycloalkenylene radicals include cyclopentenylene and cyclohexenylene.

30

"Cycloalkyl" means a saturated monocyclic or bicyclic ring system of about 3 to about 10 carbon atoms optionally substituted by oxo. Exemplary monocyclic cycloalkyl rings include C₃₋₈cycloalkyl rings such as cyclopropyl, cyclopentyl, cyclohexyl and cycloheptyl.

"Cycloalkylalkenyl" means a cycloalkyl-alkenyl- group in which the cycloalkyl and alkenyl moieties are as previously described. Exemplary monocyclic cycloalkylalkenyl groups include cyclopentylvinylene and cyclohexylvinylene.

"Cycloalkylalkyl" means a cycloalkyl-alkyl- group in which the cycloalkyl and alkyl moieties are as previously described. Exemplary monocyclic cycloalkylalkyl groups include cyclopropylmethyl, cyclopentylmethyl, cyclohexylmethyl and cycloheptylmethyl.

5 "Cycloalkylalkynyl" means a cycloalkyl-alkynyl- group in which the cycloalkyl and alkynyl moieties are as previously described. Exemplary monocyclic cycloalkylalkynyl groups include cyclopropylethynyl, cyclopentylethynyl and cyclohexylethynyl.

10 "Cycloalkylene" means a bivalent radical derived from a saturated monocyclic hydrocarbon of about 3 to about 10 carbon atoms by removing a hydrogen atom from each of two different carbon atoms of the ring. Exemplary cycloalkenylene radicals include cyclopentylene and cyclohexylene.

"Halo" or "halogen" means fluoro, chloro, bromo, or iodo. Preferred are fluoro or chloro.

15 "Heteroaroyl" means a heteroaryl-C(=O)- group in which the heteroaryl group is as described herein. Exemplary groups include pyridylcarbonyl.

"Heteroaroylamino" means a heteroaroyl-NH- group in which the heteroaryl moiety are as previously described.

20

"Heteroaryl" as a group or part of a group denotes: (i) an optionally substituted aromatic monocyclic or multicyclic organic moiety of about 5 to about 10 ring members in which one or more of the ring members is/are element(s) other than carbon, for example nitrogen, oxygen or sulfur (examples of such groups include benzimidazolyl, benzthiazolyl, furyl, imidazolyl, indolyl, indolizinyl, isoxazolyl, 25 isoquinolinyl, isothiazolyl, oxadiazolyl, pyrazinyl, pyridazinyl, pyrazolyl, pyridyl, pyrimidinyl, pyrrolyl, quinazolinyl, quinolinyl, 1,3,4-thiadiazolyl, thiazolyl, thienyl and triazolyl groups, optionally substituted by one or more aryl group substituents as defined above); (ii) an optionally substituted partially saturated multicyclic heterocarbocyclic moiety in which a heteroaryl and a cycloalkyl or cycloalkenyl group are fused together to form a cyclic structure (examples of such groups include 30 pyrindanyl groups). Optional substituents include one or more "aryl group substituents" as defined above.

"Heteroarylalkenyl" means a heteroaryl-alkenyl- group in which the heteroaryl and alkenyl moieties are as previously described. Preferred heteroarylalkenyl groups contain a lower alkenyl moiety.

35 Exemplary heteroarylalkenyl groups include pyridylethenyl and pyridylallyl.

"Heteroarylalkyl" means a heteroaryl-alkyl- group in which the heteroaryl and alkyl moieties are as previously described. Preferred heteroarylalkyl groups contain a C₁₋₄alkyl moiety. Exemplary heteroarylalkyl groups include pyridylmethyl.

5 "Heteroarylalkyloxy" means an heteroarylalkyl-O- group in which the heteroarylalkyl group is as previously described. Exemplary heteroaryloxy groups include optionally substituted pyridylmethoxy.

"Heteroarylalkynyl" means a heteroaryl-alkynyl- group in which the heteroaryl and alkynyl moieties are as previously described. Exemplary heteroarylalkenyl groups include pyridylethynyl and
10 3-pyridylbut-2-ynyl.

"Heteroaryldiyl" means a bivalent radical derived from an aromatic monocyclic or multicyclic organic moiety of about 5 to about 10 ring members in which one or more of the ring members is/are element(s) other than carbon, for example nitrogen, oxygen or sulfur, and optionally substituted by one
15 or more "aryl group substituents" as defined above. When Ar² is an optionally substituted heteroaryldiyl group this may particularly represent an optionally substituted "azaheteroaryldiyl" group.

"Heteroaryloxy" means an heteroaryl-O- group in which the heteroaryl group is as previously
20 described. Exemplary heteroaryloxy groups include optionally substituted pyridyloxy.

"Heteroarylsulfonylcarbamoyl" means a heteroaryl-SO₂-NH-C(=O)- group in which the heteroaryl group is as previously described.

25 "Heterocycloalkyl" means: (i) a cycloalkyl group of about 3 to 7 ring members which contains one or more heteroatoms selected from O, S or NY⁶ and which may optionally be substituted by oxo; (ii) an optionally substituted partially saturated multicyclic heterocarbocyclic moiety in which an aryl (or heteroaryl ring) and a heterocycloalkyl group are fused together to form a cyclic structure (examples of such groups include chromanyl, dihydrobenzofuranyl, indolynyl and pyrindolynyl groups).

30 "Heterocycloalkylalkyl" means a heterocycloalkyl-alkyl- group in which the heterocycloalkyl and alkyl moieties are as previously described.

"Heterocycloalkylene" means a bivalent radical derived from a saturated monocyclic hydrocarbon of
35 about 5 to about 7 atoms, which contains one or more heteroatoms selected from O, S or NY⁶ and is

optionally substituted by oxo, by removing a hydrogen atom from each of two different carbon atoms of the ring, or when NY⁶ is NH by removing a hydrogen atom from one carbon atom of the ring and a hydrogen atom from the NH, or when the ring contains two NY⁶ heteroatoms and NY⁶ is NH by removing a hydrogen atom from both nitrogen atoms. When L¹ is a heterocycloalkylene group this may particularly represent a bivalent radical derived pyrrolidine, especially 3,4-pyrrolidinediyl.

"Prodrug" means a compound which is convertible in vivo by metabolic means (e.g., by hydrolysis) to a compound of formula (I), including N-oxides thereof. For example an ester of a compound of formula (I) containing a hydroxy group may be convertible by hydrolysis in vivo to the parent molecule. Alternatively an ester of a compound of formula (I) containing a carboxy group may be convertible by hydrolysis in vivo to the parent molecule.

Suitable esters of compounds of formula (I) containing a hydroxy group, are for example acetates, citrates, lactates, tartrates, malonates, oxalates, salicylates, propionates, succinates, fumarates, maleates, methylene-bis-β-hydroxynaphthoates, gentisates, isethionates, di-*p*-toluoyltartrates, methanesulfonates, ethanesulfonates, benzenesulfonates, *p*-toluenesulfonates, cyclohexylsulfamates and quinate.

Suitable esters of compounds of formula (I) containing a carboxy group, are for example those described by F.J.Leinweber, Drug Metab. Res., 1987, 18, page 379.

Suitable esters of compounds of formula (I) containing both a carboxy group and a hydroxy group within the moiety -L¹-Y, include lactones, formed by loss of water between said carboxy and hydroxy groups. Examples of lactones include caprolactones and butyrolactones.

An especially useful class of esters of compounds of formula (I) containing a hydroxy group, may be formed from acid moieties selected from those described by Bundgaard et. al., J. Med. Chem., 1989, 32, page 2503-2507, and include substituted (aminomethyl)-benzoates, for example dialkylamino-methylbenzoates in which the two alkyl groups may be joined together and/or interrupted by an oxygen atom or by an optionally substituted nitrogen atom, e.g., an alkylated nitrogen atom, more especially (morpholino-methyl)benzoates, e.g., 3- or 4-(morpholinomethyl)-benzoates, and (4-alkylpiperazin-1-yl)benzoates, e.g., 3- or 4-(4-alkylpiperazin-1-yl)benzoates.

Where the compound of the invention contains a carboxy group, or a sufficiently acidic bioisostere, base addition salts may be formed and are simply a more convenient form for use; and in practice, use

of the salt form inherently amounts to use of the free acid form. The bases which can be used to prepare the base addition salts include preferably those which produce, when combined with the free acid, pharmaceutically acceptable salts, that is, salts whose cations are non-toxic to the patient in pharmaceutical doses of the salts, so that the beneficial inhibitory effects inherent in the free base are not vitiated by side effects ascribable to the cations. Pharmaceutically acceptable salts, including those derived from alkali and alkaline earth metal salts, within the scope of the invention include those derived from the following bases: sodium hydride, sodium hydroxide, potassium hydroxide, calcium hydroxide, aluminium hydroxide, lithium hydroxide, magnesium hydroxide, zinc hydroxide, ammonia, ethylenediamine, N-methyl-glucamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chlorprocaine, diethanolamine, procaine, N-benzylphenethylamine, diethylamine, piperazine, tris(hydroxymethyl)aminomethane, tetramethylammonium hydroxide, and the like.

Some of the compounds of the present invention are basic, and such compounds are useful in the form of the free base or in the form of a pharmaceutically acceptable acid addition salt thereof.

Acid addition salts are a more convenient form for use; and in practice, use of the salt form inherently amounts to use of the free base form. The acids which can be used to prepare the acid addition salts include preferably those which produce, when combined with the free base, pharmaceutically acceptable salts, that is, salts whose anions are non-toxic to the patient in pharmaceutical doses of the salts, so that the beneficial inhibitory effects inherent in the free base are not vitiated by side effects ascribable to the anions. Although pharmaceutically acceptable salts of said basic compounds are preferred, all acid addition salts are useful as sources of the free base form even if the particular salt, per se, is desired only as an intermediate product as, for example, when the salt is formed only for purposes of purification, and identification, or when it is used as intermediate in preparing a pharmaceutically acceptable salt by ion exchange procedures. Pharmaceutically acceptable salts within the scope of the invention include those derived from mineral acids and organic acids, and include hydrohalides, e.g., hydrochlorides and hydrobromides, sulfates, phosphates, nitrates, sulfamates, acetates, citrates, lactates, tartrates, malonates, oxalates, salicylates, propionates, succinates, fumarates, maleates, methylene-bis-b-hydroxynaphthoates, gentisates, isethionates, di-*p*-toluoyltartrates, methane-sulfonates, ethanesulfonates, benzenesulfonates, *p*-toluenesulfonates, cyclohexylsulfamates and quinate.

As well as being useful in themselves as active compounds, salts of compounds of the invention are useful for the purposes of purification of the compounds, for example by exploitation of the solubility


differences between the salts and the parent compounds, side products and/or starting materials by techniques well known to those skilled in the art.

With reference to formula (I) above, the following are particular and preferred groupings:


5

R^1 may particularly represent a group R^3 -NH-Ar¹-L²- in which: L² is a straight or branched C₁₋₆alkylene chain, more particularly a straight C₁₋₄alkylene chain such as methylene or ethylene,

preferably methylene; Ar¹ is an 8 to 10 membered bicyclic system




in which (i) ring



is a 5 or 6 membered optionally substituted heterocycle, preferably a 5 membered

10


heteroaryl ring, (ii) ring



is a 5 or 6 membered optionally substituted heterocycle or an

optionally substituted benzene ring, preferably a benzene ring, (iii) each ring is optionally substituted by one or more "aryl group substituents" as defined above, (iv) the two rings are joined together by a


carbon-carbon linkage or a carbon-nitrogen linkage, and



is preferably optionally

substituted benzoxazolyl or optionally substituted benzimidazolyl, each [more particularly ring

15



] optionally substituted by one or more "aryl group substituents" as defined above [examples

of particular aryl group substituents include C₁₋₄alkyl (e.g., methyl), C₁₋₄alkoxy (e.g., methoxy), amino, halogen, hydroxy, C₁₋₄alkylthio, C₁₋₄alkylsulfinyl, C₁₋₄alkylsulfonyl, nitro or

trifluoromethyl]; and R³ is optionally substituted aryl, such as a 2-substituted phenyl, [examples of particular aryl group substituents include C₁₋₄alkyl (e.g., methyl), C₁₋₄alkoxy (e.g., methoxy), halo

20

(e.g., fluoro or chloro) and Y¹Y²N- (e.g., dimethylamino)], and is preferably 2-methylphenyl.

R^1 may also particularly represent a group R^3 -NH-C(=O)-NH-Ar²-L²- in which L² is a straight or branched C₁₋₆alkylene chain, more particularly a straight C₁₋₄alkylene chain such as methylene or

ethylene, preferably methylene; Ar² is an optionally substituted phenylene, such as optionally

25


substituted *m*- or *p*-phenylene, preferably optionally substituted *p*-phenylene, more preferably a 3-

substituted *p*-phenylene, in which the substituent is ortho to the R³-NH-C(=O)-NH- group, (preferred

optional substituents include halo, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkylthio, C₁₋₄alkylsulfinyl and C₁₋₄alkylsulfonyl, especially chloro, methyl, ethyl, methoxy, methylthio, methylsulfinyl and methylsulfonyl), or Ar² is an optionally substituted heteroaryldiyl, such as optionally substituted azaheteroaryldiyl, (e.g., optionally substituted pyridinediyl, preferably a *p*-pyridinediyl), where the
 5 optional substituents include C₁₋₄alkyl and C₁₋₄alkoxy, especially methyl and methoxy, more preferably a pyridine-2,5-diyl, in which the R³-NH-C(=O)-NH- group is adjacent to the pyridyl nitrogen atom, and which is substituted in the 4- or 6-position with a methyl or methoxy group; and R³ is an optionally substituted aryl group (such as optionally substituted phenyl) or an optionally substituted heteroaryl (such as optionally substituted pyridyl), and is preferably 2- or 3-methyl(or
 10 methoxy)phenyl, more preferably 2-methylphenyl, or 3-methyl-2-pyridyl.


R² may particularly represent hydrogen.

L¹ may particularly represent an optionally substituted alkylene linkage especially optionally
 15 substituted ethylene or propylene, preferably optionally substituted ethylene. Preferred optional substituents include C₁₋₄alkyl, aryl, heteroaryl, -N(R⁶)-C(=O)-R⁴, -N(R⁶)-C(=O)-OR⁴, -N(R⁶)-SO₂-R⁴, -NY³Y⁴ and -[C(=O)-N(R⁷)-C(R⁸)(R⁹)]_p-C(=O)-NY³Y⁴ or alkyl substituted by an acidic functional group, -C(=O)-NY³Y⁴ or -NY³Y⁴. In one preferred embodiment L¹ is a group



20 C₁₋₄alkyl, or where R¹⁴ is hydrogen and R¹⁵ represents aryl, heteroaryl, -N(R⁶)-C(=O)-R⁴, -N(R⁶)-C(=O)-OR⁴, -N(R⁶)-SO₂-R⁴, -NY³Y⁴ or -[C(=O)-N(R⁷)-C(R⁸)(R⁹)]_p-C(=O)-NY³Y⁴, or alkyl substituted by an acidic functional group, -C(=O)-NY³Y⁴ or -NY³Y⁴], and is more preferably a group $\begin{array}{c} \text{---CH---CH}_2\text{---} \\ | \\ \text{R}^{15} \end{array}$, particularly $\begin{array}{c} \text{---CH---CH}_2\text{---} \\ | \\ \text{R}^{15} \end{array}$ [where R¹⁵ represents hydrogen, C₁₋₄alkyl,

aryl, heteroaryl, -N(R⁶)-C(=O)-R⁴, -N(R⁶)-C(=O)-OR⁴, -N(R⁶)-SO₂-R⁴ or -NY³Y⁴ or alkyl
 25 substituted by carboxy, -OH, -OR¹³ or -C(=O)-NY³Y⁴]. In another preferred embodiment L¹ is a




C₁₋₄alkyl, or where R¹⁴ is hydrogen and R¹⁶ represents aryl, heteroaryl, -N(R⁶)-C(=O)-R⁴,
 -N(R⁶)-C(=O)-OR⁴, -N(R⁶)-SO₂-R⁴, -NY³Y⁴ or -[C(=O)-N(R⁷)-C(R⁸)(R⁹)]_p-C(=O)-NY³Y⁴, or
 alkyl substituted by an acidic functional group, -C(=O)-NY³Y⁴ or -NY³Y⁴], and is more preferably a
 group $\text{---CH}_2\text{---CH---}$, particularly $\text{---CH}_2\text{---CH---}$ [where R¹⁶ represents -N(R⁶)-C(=O)-R⁴, or
 R^{16} R^{16}

5 -N(R⁶)-SO₂-R⁴].

Y may particularly represent carboxy.

10 It is to be understood that this invention covers all appropriate combinations of the particular and preferred groupings referred to herein.

A particular group of compounds of the invention are compounds of formula (Ia):-



15

(Ia)

in which R², R³, L¹, L² and Y are as hereinbefore defined, X is O or NR¹⁸ (where R¹⁸ is hydrogen or
 C₁₋₄alkyl), and R¹⁷ is hydrogen or an aryl group substituent, and the corresponding N-oxides, and
 their prodrugs; and pharmaceutically acceptable salts and solvates (e.g., hydrates) of such compounds
 20 and their N-oxides and prodrugs.

Compounds of formula (Ia) in which R³ represents optionally substituted aryl, especially 2-substituted
 phenyl, are preferred. Preferred optional substituents include C₁₋₄alkyl (e.g., methyl), C₁₋₄alkoxy
 (e.g., methoxy), halo (e.g., fluoro or chloro) and Y¹Y²N- (e.g., dimethylamino). R³ especially
 25 represents ortho-tolyl.

Compounds of formula (Ia) in which R¹⁷ represents hydrogen, halo (e.g., chloro), C₁₋₄alkyl (e.g.,
 methyl or ethyl) or C₁₋₄alkoxy (e.g., methoxy) are preferred.

Compounds of formula (Ia) in which L^2 represents a straight or branched C_{1-6} alkylene chain, especially a straight or branched C_{1-4} alkylene chain, more especially methylene, are preferred.

- 5 Compounds of formula (Ia) in which R^2 represents hydrogen are preferred.

Compounds of formula (Ia) in which L^1 represents an optionally substituted alkylene linkage, particularly optionally substituted ethylene or optionally substituted propylene, especially ethylene optionally substituted with C_{1-4} alkyl, aryl, heteroaryl, $-N(R^7)-C(=O)-R^8$, $-N(R^7)-C(=O)-OR^8$,

- 10 $-N(R^7)-SO_2-R^8$, $-NY^1Y^2$ or $-[C(=O)-N(R^9)-C(R^5)(R^{10})]_p-C(=O)-NY^1Y^2$, or alkyl substituted by an acidic functional group, $-C(=O)-NY^1Y^2$ or $-NY^1Y^2$, are preferred. In one preferred embodiment L^1 is

a group $\begin{array}{c} R^{14} \\ | \\ -C-CH_2- \\ | \\ R^{15} \end{array}$ [where R^{14} is hydrogen or C_{1-4} alkyl (e.g., methyl) and R^{15} represents

hydrogen or C_{1-4} alkyl, or where R^{14} is hydrogen and R^{15} represents aryl, heteroaryl,

$-N(R^7)-C(=O)-R^8$, $-N(R^7)-C(=O)-OR^8$, $-N(R^7)-SO_2-R^8$, $-NY^1Y^2$ or

- 15 $-[C(=O)-N(R^9)-C(R^5)(R^{10})]_p-C(=O)-NY^1Y^2$, or alkyl substituted by an acidic functional group, $-C(=O)-NY^1Y^2$ or $-NY^1Y^2$], and is more preferably a group $\begin{array}{c} -CH-CH_2- \\ | \\ R^{15} \end{array}$, particularly

$\begin{array}{c} -CH-CH_2- \\ | \\ R^{15} \end{array}$ [where R^{15} represents hydrogen, C_{1-4} alkyl, aryl, heteroaryl, $-N(R^7)-C(=O)-R^8$,

$-N(R^7)-C(=O)-OR^8$, $-N(R^7)-SO_2-R^8$ or $-NY^1Y^2$ or alkyl substituted by carboxy, $-OH$, $-OR^{13}$ or

$-C(=O)-NY^1Y^2$]. In another preferred embodiment L^1 is a group $\begin{array}{c} R^{14} \\ | \\ -CH_2-C- \\ | \\ R^{16} \end{array}$ [where R^{14} is


- 20 hydrogen or C_{1-4} alkyl (e.g., methyl) and R^{16} represents C_{1-4} alkyl, or where R^{14} is hydrogen and R^{16} represents aryl, heteroaryl, $-N(R^7)-C(=O)-R^8$, $-N(R^7)-C(=O)-OR^8$, $-N(R^7)-SO_2-R^8$, $-NY^1Y^2$ or $-[C(=O)-N(R^9)-C(R^5)(R^{10})]_p-C(=O)-NY^1Y^2$, or alkyl substituted by an acidic functional group,

-C(=O)-NY¹Y² or -NY¹Y²], and is more preferably a group $\text{---CH}_2\text{---CH---}$, particularly

$\text{---CH}_2\text{---CH---}$ [where R¹⁶ represents -N(R⁶)-C(=O)-R⁴, or -N(R⁶)-SO₂-R⁴].

Compounds of formula (Ia) in which Y represents carboxy are preferred.

5



the ring 5 or 6 position when X is NR¹⁸ and R¹⁸ is C₁₋₄alkyl.


The group $\text{---L}^1\text{---Y}$ may preferably be attached at position 6 or 7 of the tetrahydroisoquinoline ring.

10

A preferred group of compounds of the invention are compounds of formula (Ia) in which:- R² is hydrogen; R³ is optionally substituted aryl (especially ortho-tolyl); R¹⁷ is hydrogen, chloro, methyl, ethyl or methoxy; L¹ is a $\text{---CH---CH}_2\text{---}$ group particularly a $\text{---CH---CH}_2\text{---}$ group, where R¹⁵

represents hydrogen, C₁₋₄alkyl, aryl, heteroaryl, -N(R⁷)-C(=O)-R⁸, -N(R⁷)-C(=O)-OR⁸,


15 -N(R⁷)-SO₂-R⁸ or -NY¹Y², or alkyl substituted by carboxy, -OH, -Z¹R¹³, -C(=O)-NY¹Y² or -NY¹Y²; L² is a straight or branched C₁₋₄alkylene chain, especially methylene; X is O; Y is carboxy;



$\text{---L}^1\text{---Y}$ is attached at position 6 or 7 of the tetrahydroisoquinoline ring; and the corresponding N-oxides, and their prodrugs; and pharmaceutically acceptable salts and solvates (e.g., hydrates) of such compounds and their N-oxides and prodrugs.


20

Another preferred group of compounds of the invention are compounds of formula (Ia) in which:- R^2 is hydrogen; R^3 is optionally substituted aryl (especially ortho-tolyl); R^{17} is hydrogen, chloro, methyl, ethyl or methoxy; L^1 is a $\text{---CH}_2\text{---CH---}$ group, particularly $\text{---CH}_2\text{---CH---}$ [where R^{16} represents $\text{---N(R}^7\text{)---C(=O)---R}^8$, or $\text{---N(R}^7\text{)---SO}_2\text{---R}^8$]; L^2 is a straight or branched C_{1-4} alkylene chain, especially



ring 6 position; and the group $\text{---L}^1\text{---Y}$ is attached at position 6 or 7 of the tetrahydroisoquinoline ring; and the corresponding N-oxides, and their prodrugs; and pharmaceutically acceptable salts and solvates (e.g., hydrates) of such compounds and their N-oxides and prodrugs.

- 10 Another preferred group of compounds of the invention are compounds of formula (Ia) in which:- R^2 is hydrogen; R^3 is optionally substituted aryl (especially ortho-tolyl); R^{17} is hydrogen, chloro, methyl, ethyl or methoxy; L^1 is a $\text{---CH---CH}_2\text{---}$ group particularly a $\text{---CH---CH}_2\text{---}$ group, where R^{15}
- 15 $\text{---N(R}^7\text{)---C(=O)---R}^8$, $\text{---N(R}^7\text{)---C(=O)---OR}^8$, $\text{---N(R}^7\text{)---SO}_2\text{---R}^8$ or $\text{---NY}^1\text{Y}^2$, or alkyl substituted by carboxy, ---OH , ---OR^{13} , $\text{---C(=O)---NY}^1\text{Y}^2$ or $\text{---NY}^1\text{Y}^2$; L^4 is a straight or branched C_{1-4} alkylene chain, (especially methylene); X is NR^{18}




5 or 6 position; and the group $\text{---L}^1\text{---Y}$ is attached at position 6 or 7 of the tetrahydroisoquinoline ring; and the corresponding N-oxides, and their prodrugs; and pharmaceutically acceptable salts and solvates (e.g., hydrates) of such compounds and their N-oxides and prodrugs.

- 20 Another preferred group of compounds of the invention are compounds of formula (Ia) in which:- R^2 is hydrogen; R^3 is optionally substituted aryl (especially ortho-tolyl); R^{17} is hydrogen, chloro, methyl,

ethyl or methoxy; L^1 is a $\text{---CH}_2\text{---CH---}$ group, particularly $\text{---CH}_2\text{---CH---}$ [where R^{16} represents


$\text{---N}(R^7)\text{---C(=O)---R}^8$, or $\text{---N}(R^7)\text{---SO}_2\text{---R}^8$]; L^2 is a straight or branched C_{1-4} alkylene chain, (especially methylene); X is NR^{18} (especially NH); Y is carboxy; the group



5 is attached at position 6 or 7 of the tetrahydroisoquinoline ring; and the corresponding N-oxides, and their prodrugs; and pharmaceutically acceptable salts and solvates (e.g., hydrates) of such compounds and their N-oxides and prodrugs.

Another particular group of compounds of the invention are compounds of formula (Ib):-

10



15 in which R^2 , R^3 , Ar^2 , L^1 , L^2 and Y are as hereinbefore defined, and the corresponding N-oxides, and their prodrugs; and pharmaceutically acceptable salts and solvates (e.g., hydrates) of such compounds and their N-oxides and prodrugs.

20 Compounds of formula (Ib) in which R^3 represents an optionally substituted aryl group, particularly an optionally substituted phenyl group, such as a 2-substituted phenyl, especially 2-methylphenyl, are preferred.

Compounds of formula (Ib) in which R^3 represents an optionally substituted heteroaryl group, particularly an optionally substituted pyridyl, such as optionally substituted 2-pyridyl, especially 3-methyl-2-pyridyl, are also preferred.

25

Compounds of formula (Ib) in which Ar^2 represents an optionally substituted phenylene, especially optionally substituted *m*- or *p*-phenylene, more especially optionally substituted *p*-phenylene, are

preferred. Compounds of formula (Ib) in which Ar² represents 3-substituted *p*-phenylene, in which the substituent is ortho to the R³-NH-C(=O)-NH- group, are particularly preferred. Preferred optional substituents include halo, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkylthio, C₁₋₄alkylsulfinyl and C₁₋₄alkylsulfonyl, especially chloro, methyl, ethyl, methoxy, methylthio, methylsulfinyl and methylsulfonyl.

Compounds of formula (Ib) in which Ar² is an optionally substituted heteroaryldiyl, such as optionally substituted pyridinediyl, particularly a *p*-pyridinediyl, more particularly a pyridine-2,5-diyl, in which the R³-NH-C(=O)-NH- group is adjacent to the pyridyl nitrogen atom, and which is substituted in the 4- or 6-position, are also preferred. Preferred optional substituents include C₁₋₄alkyl and C₁₋₄alkoxy, especially methyl and methoxy.

Compounds of formula (Ib) in which L² represents a straight or branched C₁₋₆alkylene chain, more particularly a straight C₁₋₄alkylene chain such as methylene or ethylene, especially methylene, are preferred.

Compounds of formula (Ib) in which R² represents hydrogen are preferred.

Compounds of formula (Ib) in which L¹ represents an optionally substituted alkylene linkage, particularly optionally substituted ethylene or optionally substituted propylene, especially ethylene optionally substituted with C₁₋₄alkyl, aryl, heteroaryl, -N(R⁷)-C(=O)-R⁸, -N(R⁷)-C(=O)-OR⁸, -N(R⁷)-SO₂-R⁸, -NY¹Y² or -[C(=O)-N(R⁹)-C(R⁵)(R¹⁰)]_p-C(=O)-NY¹Y², or alkyl substituted by an acidic functional group, -C(=O)-NY¹Y² or -NY¹Y², are preferred. In one preferred embodiment L¹ is

a group $\begin{array}{c} \text{R}^{14} \\ | \\ \text{---C---CH}_2\text{---} \\ | \\ \text{R}^{15} \end{array}$ [where R¹⁴ is hydrogen or C₁₋₄alkyl (e.g., methyl) and R¹⁵ represents

hydrogen or C₁₋₄alkyl, or where R¹⁴ is hydrogen and R¹⁵ represents aryl, heteroaryl, -N(R⁷)-C(=O)-R⁸, -N(R⁷)-C(=O)-OR⁸, -N(R⁷)-SO₂-R⁸, -NY¹Y² or -[C(=O)-N(R⁹)-C(R⁵)(R¹⁰)]_p-C(=O)-NY¹Y², or alkyl substituted by an acidic functional group, -C(=O)-NY¹Y² or -NY¹Y²], and is more preferably a group $\begin{array}{c} \text{---CH---CH}_2\text{---} \\ | \\ \text{R}^{15} \end{array}$, particularly

— $\underset{\text{R}^{15}}{\text{CH}}\text{—CH}_2\text{—}$ [where R¹⁵ represents hydrogen, C₁₋₄alkyl, aryl, heteroaryl, -N(R⁷)-C(=O)-R⁸,

-N(R⁷)-C(=O)-OR⁸, -N(R⁷)-SO₂-R⁸ or -NY¹Y² or alkyl substituted by carboxy, -OH, -OR¹³ or

-C(=O)-NY¹Y²]. In another preferred embodiment L¹ is a group $\text{—CH}_2\text{—}\underset{\text{R}^{16}}{\overset{\text{R}^{14}}{\text{C}}}\text{—}$ [where R¹⁴ is

hydrogen or C₁₋₄alkyl (e.g., methyl) and R¹⁶ represents C₁₋₄alkyl, or where R¹⁴ is hydrogen and R¹⁶

5 represents aryl, heteroaryl, -N(R⁷)-C(=O)-R⁸, -N(R⁷)-C(=O)-OR⁸, -N(R⁷)-SO₂-R⁸, -NY¹Y² or

-[C(=O)-N(R⁹)-C(R⁵)(R¹⁰)]_p-C(=O)-NY¹Y², or alkyl substituted by an acidic functional group,

-C(=O)-NY¹Y² or -NY¹Y²], and is more preferably a group $\text{—CH}_2\text{—}\underset{\text{R}^{16}}{\text{CH}}\text{—}$, particularly

$\text{—CH}_2\text{—}\underset{\text{R}^{16}}{\text{CH}}\text{—}$ [where R¹⁶ represents -N(R⁷)-C(=O)-R⁸, or -N(R⁷)-SO₂-R⁸].

10 Compounds of formula (Ib) in which Y represents carboxy are preferred.

A preferred group of compounds of the invention are compounds of formula (Ib) in which:- R² is hydrogen; R³ is a 2-substituted phenyl [especially 2-methyl(or methoxy)phenyl]; Ar² is optionally substituted *m*- or *p*-phenylene (especially 3-chloro-*p*-phenylene, 3-methyl-*p*-phenylene, 3-ethyl-*p*-

15 phenylene, 3-methoxy-*p*-phenylene, 3-methylthio-*p*-phenylene, 3-methylsulfinyl-*p*-phenylene and 3-methylsulfonyl-*p*-phenylene) or optionally substituted *p*-pyridinediyl [especially 4(or 6)-methyl(or methoxy)-*p*-pyridine-2,5-diyl]; L¹ is a $\text{—}\underset{\text{R}^{15}}{\text{CH}}\text{—CH}_2\text{—}$ group particularly a $\text{—}\underset{\text{R}^{15}}{\text{CH}}\text{—CH}_2\text{—}$ group,

where R¹⁵ represents hydrogen, C₁₋₄alkyl, aryl, heteroaryl, -N(R⁷)-C(=O)-R⁸, -N(R⁷)-C(=O)-OR⁸, -N(R⁷)-SO₂-R⁸ or -NY¹Y², or alkyl substituted by carboxy, -OH, -OR¹³, -C(=O)-NY¹Y²; L²

20 represents a straight or branched C₁₋₆alkylene chain, especially methylene; and Y represents carboxy; and the corresponding N-oxides, and their prodrugs; and pharmaceutically acceptable salts and solvates (e.g., hydrates) of such compounds and their N-oxides and prodrugs.

Another preferred group of compounds of the invention are compounds of formula (Ib) in which:- R² is hydrogen; R³ is a 2-substituted phenyl [especially 2-methyl(or methoxy)phenyl]; Ar² is optionally substituted *m*- or *p*-phenylene (especially 3-chloro-*p*-phenylene, 3-methyl-*p*-phenylene, 3-methoxy-*p*-phenylene, 3-methylthio-*p*-phenylene, 3-methylsulfinyl-*p*-phenylene and 3-methylsulfonyl-*p*-phenylene) or optionally substituted *p*-pyridinediyl [especially 4(or 6)-methyl(or methoxy)-*p*-pyridine-2,5-diyl]; L¹ is a $\text{---CH}_2\text{---CH---}$ group, particularly $\text{---CH}_2\text{---CH---}$ [where R¹⁶ represents

-N(R⁷)-C(=O)-R⁸, or -N(R⁷)-SO₂-R⁸]; L² represents a straight or branched C₁₋₆alkylene chain, especially methylene; and Y represents carboxy; and the corresponding N-oxides, and their prodrugs; and pharmaceutically acceptable salts and solvates (e.g., hydrates) of such compounds and their

Particular compounds of the invention are selected from the compounds formed by joining the acyl carbon atom (C*) of one of the fragments (A1 to A18) shown in Table 1 to the nitrogen atom (N*) of one of the tetrahydroisoquinoline fragments (B1 to B3) shown in Table 2, and joining the carbon atom (C*) of the phenyl ring in one of the tetrahydroisoquinoline fragments (B1 to B3) shown in Table 2 to the carbon atom (C*) of one of the acidic fragments (C1 to C28) depicted in Table 3.

Table 1

A1		A2	
A3		A4	
A5		A6	
A7		A8	


A9		A10	
A11		A12	
A13		A14	
A15		A16	
A17		A18	
A19		A20	

Table 2

B1		B2	
B3			


Table 3

C1	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ CH_3 \end{array}$	C2	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ CH_2CH_3 \end{array}$
C3	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ CH(CH_3)_2 \end{array}$	C4	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ CH_2CH(CH_3)_2 \end{array}$
C5	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ C(CH_3)_3 \end{array}$	C6	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ \text{C}_6\text{H}_5 \end{array}$
C7	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ \text{C}_6\text{H}_4 \\ \\ F \end{array}$	C8	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ \text{C}_5\text{H}_4\text{N} \end{array}$
C9	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ \text{C}_5\text{H}_4\text{N} \end{array}$	C10	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ \text{C}_5\text{H}_4\text{N} \end{array}$
C11	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ \text{C}_4\text{H}_3\text{O} \end{array}$	C12	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ \text{C}_4\text{H}_3\text{O} \end{array}$
C13	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ \text{C}_4\text{H}_3\text{S} \end{array}$	C14	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ \text{C}_4\text{H}_3\text{S} \end{array}$
C15	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ \text{C}_5\text{H}_9\text{N} \end{array}$	C16	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ \text{C}_5\text{H}_9\text{O} \end{array}$
C17	$\begin{array}{c} *CH-CH-CO_2H \\ \quad \\ C \\ \\ H_2 \end{array}$	C18	$\begin{array}{c} \text{C}_4\text{H}_7 \\ \\ *C-CH_2-CO_2H \end{array}$

C19		C20	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ CH_2CO_2H \end{array}$
C21	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ CH_2 \\ \\ C=O \\ \\ N \\ \\ \text{Cyclopentyl} \end{array}$	C22	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ OMe \end{array}$
C23	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ HN \\ \\ C=O \\ \\ \text{Phenyl} \end{array}$	C24	$\begin{array}{c} *CH_2-CH-CO_2H \\ \\ HN \\ \\ C=O \\ \\ \text{2,6-dichlorophenyl} \end{array}$
C25	$\begin{array}{c} *CH_2-CH-CO_2H \\ \\ HN \\ \\ C=O \\ \\ \text{2-chloro-3-methylphenyl} \end{array}$	C26	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ HN \\ \\ C=O \\ \\ \text{Pyridin-2-yl} \end{array}$
C27	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ HN \\ \\ C=O \\ \\ \text{Thiophen-2-yl} \end{array}$	C28	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ HN \\ \\ C=O \\ \\ \text{4-methylisoxazole-3-yl} \end{array}$
C29	$\begin{array}{c} *CH-CH_2-CO_2H \\ \\ \text{Cyclohexyl} \end{array}$	C30	$\begin{array}{c} *C=CH-CO_2H \\ \\ CH_3 \end{array}$

Compounds in accordance with the present invention may consist of any combination of the above-identified "A", "B", and "C" fragments.

Thus, for example, a compound denoted as A1-B1-C1 is the product of the combination of group A1 in Table 1 and B1 in Table 2 and C1 in Table 3, namely



5

Particular compounds of the invention, as exemplified below, are selected from the following:

3-(((4-methyl-2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-butanoic acid, Example 1(a) [A3-B1-C1];

10 3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-butanoic acid, Example 1(b) [A1-B2-C1];

3-phenyl-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-propanoic acid, Example 1(c) [A1-B1-C6];

15 3-cyclohexyl-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-propanoic acid, Example 1(d) [A1-B1-C29];

3-(pyrid-4-yl)-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid, Example 1(e) [A1-B2-C10];

3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-8-yl)-but-2-enoic acid, Example 1(f) [A1-B3-C30];

20 3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-8-yl)-butanoic acid, Example 1(g) [A1-B3-C1];

3-{2-[(2-o-tolylamino-benzoxazol-6-yl)-carbonyl]-1,2,3,4-tetrahydro-isoquinolin-6-yl}-butanoic acid, Example 1(h) [A19-B1-C1];

25 {5-(3-methoxy-4-[3-(2-methylphenyl)ureido]-phenylacetyl-amino)-1,2,3,4-tetrahydro-isoquinolin-8-yl}-butanoic acid, Example 1(i) [A18-B3-C1];

2-(2,6-dichloro-benzoylamino)-3-[2-(2,6-dichloro-benzoyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl]-propionic acid, Example 1(j) [A20-B2-C24];

3-phenyl-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid, Example 1(k) [A1-B2-C6];

3-{(2-*o*-tolylamino-benzoxazol-6-yl)-acetyl}-1,2,3,4-tetrahydro-isoquinolin-6-yl}-butanoic acid,
Example 1(l) [A1-B1-C1];

3-(pyrid-4-yl)-3-{{{(2-*o*-tolylamino-benzoxazol-6-yl)-acetyl}-1,2,3,4-tetrahydro-isoquinolin-7-yl}-
propanoic acid, enantiomer A, Example 1(m) [A1-B2-C10];

5 3-(pyrid-4-yl)-3-{{{(2-*o*-tolylamino-benzoxazol-6-yl)-acetyl}-1,2,3,4-tetrahydro-isoquinolin-7-yl}-
propanoic acid, enantiomer B, Example 1(n) [A1-B2-C10];

and the corresponding N-oxides, and their prodrugs; and pharmaceutically acceptable salts and solvates
(e.g., hydrates) of such compounds and their N-oxides and prodrugs.

10 Of these, preferred compounds of the invention include:

3-{{{(2-*o*-tolylamino-benzoxazol-6-yl)-acetyl}-1,2,3,4-tetrahydro-isoquinolin-7-yl}-butanoic acid,
Example 1(b) [A1-B2-C1];

3-phenyl-3-{{{(2-*o*-tolylamino-benzoxazol-6-yl)-acetyl}-1,2,3,4-tetrahydro-isoquinolin-6-yl}-propanoic
acid, Example 1(c) [A1-B1-C6];

15 3-(pyrid-4-yl)-3-{{{(2-*o*-tolylamino-benzoxazol-6-yl)-acetyl}-1,2,3,4-tetrahydro-isoquinolin-7-yl}-
propanoic acid, Example 1(e) [A1-B2-C10];

and the corresponding N-oxides, and their prodrugs; and pharmaceutically acceptable salts and solvates
(e.g., hydrates) of such compounds and their N-oxides and prodrugs.

20 The compounds of the invention exhibit useful pharmacological activity and accordingly are
incorporated into pharmaceutical compositions and used in the treatment of patients suffering from
certain medical disorders. The present invention thus provides, according to a further aspect,
compounds of the invention and compositions containing compounds of the invention for use in
therapy.

25 Compounds within the scope of the present invention block the interaction of the ligand VCAM-1 to its
integrin receptor VLA-4 ($\alpha 4\beta 1$) according to tests described in the literature and described in vitro and
in vivo procedures hereinafter, and which tests results are believed to correlate to pharmacological
activity in humans and other mammals. Thus, in a further embodiment, the present invention provides
30 compounds of the invention and compositions containing compounds of the invention for use in the
treatment of a patient suffering from, or subject to, conditions which can be ameliorated by the
administration of an inhibitor of $\alpha 4\beta 1$ mediated cell adhesion. For example, compounds of the
present invention are useful in the treatment of inflammatory diseases, for example joint inflammation,
including arthritis, rheumatoid arthritis and other arthritic conditions such as rheumatoid spondylitis,
35 gouty arthritis, traumatic arthritis, rubella arthritis, psoriatic arthritis and osteoarthritis. Additionally,
the compounds may be useful in the treatment of acute synovitis, autoimmune diabetes, autoimmune

encephalomyelitis, collitis, atherosclerosis, peripheral vascular disease, cardiovascular disease, multiple sclerosis, asthma, psoriasis restenosis, myocarditis, inflammatory bowel disease and melanoma cell division in metastasis.

- 5 A special embodiment of the therapeutic methods of the present invention is the treating of asthma.

Another special embodiment of the therapeutic methods of the present invention is the treating of joint inflammation.

- 10 Another special embodiment of the therapeutic methods of the present invention is the treating of inflammatory bowel disease.

- According to a further feature of the invention there is provided a method for the treatment of a human or animal patient suffering from, or subject to, conditions which can be ameliorated by the
15 administration of an inhibitor of the interaction of the ligand VCAM-1 to its integrin receptor VLA-4 ($\alpha 4\beta 1$), for example conditions as hereinbefore described, which comprises the administration to the patient of an effective amount of compound of the invention or a composition containing a compound of the invention. "Effective amount" is meant to describe an amount of compound of the present invention effective in inhibiting the interaction of the ligand VCAM-1 to its integrin receptor VLA-4
20 ($\alpha 4\beta 1$), and thus producing the desired therapeutic effect.

References herein to treatment should be understood to include prophylactic therapy as well as treatment of established conditions.

- 25 The present invention also includes within its scope pharmaceutical compositions comprising at least one of the compounds of the invention in association with a pharmaceutically acceptable carrier or excipient.

- Compounds of the invention may be administered by any suitable means. In practice compounds of the
30 present invention may generally be administered parenterally, topically, rectally, orally or by inhalation, especially by the oral route.

- Compositions according to the invention may be prepared according to the customary methods, using one or more pharmaceutically acceptable adjuvants or excipients. The adjuvants comprise, inter alia,
35 diluents, sterile aqueous media and the various non-toxic organic solvents. The compositions may be presented in the form of tablets, pills, granules, powders, aqueous solutions or suspensions, injectable

solutions, elixirs or syrups, and can contain one or more agents chosen from the group comprising sweeteners, flavourings, colourings, or stabilisers in order to obtain pharmaceutically acceptable preparations. The choice of vehicle and the content of active substance in the vehicle are generally determined in accordance with the solubility and chemical properties of the active compound, the particular mode of administration and the provisions to be observed in pharmaceutical practice. For example, excipients such as lactose, sodium citrate, calcium carbonate, dicalcium phosphate and disintegrating agents such as starch, alginic acids and certain complex silicates combined with lubricants such as magnesium stearate, sodium lauryl sulfate and talc may be used for preparing tablets. To prepare a capsule, it is advantageous to use lactose and high molecular weight polyethylene glycols. When aqueous suspensions are used they can contain emulsifying agents or agents which facilitate suspension. Diluents such as sucrose, ethanol, polyethylene glycol, propylene glycol, glycerol and chloroform or mixtures thereof may also be used.

For parenteral administration, emulsions, suspensions or solutions of the products according to the invention in vegetable oil, for example sesame oil, groundnut oil or olive oil, or aqueous-organic solutions such as water and propylene glycol, injectable organic esters such as ethyl oleate, as well as sterile aqueous solutions of the pharmaceutically acceptable salts, are used. The solutions of the salts of the products according to the invention are especially useful for administration by intramuscular or subcutaneous injection. The aqueous solutions, also comprising solutions of the salts in pure distilled water, may be used for intravenous administration with the proviso that their pH is suitably adjusted, that they are judiciously buffered and rendered isotonic with a sufficient quantity of glucose or sodium chloride and that they are sterilised by heating, irradiation or microfiltration.

For topical administration, gels (water or alcohol based), creams or ointments containing compounds of the invention may be used. Compounds of the invention may also be incorporated in a gel or matrix base for application in a patch, which would allow a controlled release of compound through the transdermal barrier.

For administration by inhalation compounds of the invention may be dissolved or suspended in a suitable carrier for use in a nebuliser or a suspension or solution aerosol, or may be absorbed or adsorbed onto a suitable solid carrier for use in a dry powder inhaler.

Solid compositions for rectal administration include suppositories formulated in accordance with known methods and containing at least one compound of the invention.

The percentage of active ingredient in the compositions of the invention may be varied, it being necessary that it should constitute a proportion such that a suitable dosage shall be obtained.

Obviously, several unit dosage forms may be administered at about the same time. The dose employed will be determined by the physician, and depends upon the desired therapeutic effect, the route of administration and the duration of the treatment, and the condition of the patient. In the adult, the doses are generally from about 0.001 to about 50, preferably about 0.001 to about 5, mg/kg body weight per day by inhalation, from about 0.01 to about 100, preferably 0.1 to 70, more especially 0.5 to 10, mg/kg body weight per day by oral administration, and from about 0.001 to about 10, preferably 0.01 to 1, mg/kg body weight per day by intravenous administration. In each particular case, the doses will be determined in accordance with the factors distinctive to the subject to be treated, such as age, weight, general state of health and other characteristics which can influence the efficacy of the medicinal product.

The compounds according to the invention may be administered as frequently as necessary in order to obtain the desired therapeutic effect. Some patients may respond rapidly to a higher or lower dose and may find much weaker maintenance doses adequate. For other patients, it may be necessary to have long-term treatments at the rate of 1 to 4 doses per day, in accordance with the physiological requirements of each particular patient. Generally, the active product may be administered orally 1 to 4 times per day. Of course, for some patients, it will be necessary to prescribe not more than one or two doses per day.

Compounds of the invention may be prepared by the application or adaptation of known methods, by which is meant methods used heretofore or described in the literature, for example those described by R.C.Larock in Comprehensive Organic Transformations, VCH publishers, 1989.

In the reactions described hereinafter it may be necessary to protect reactive functional groups, for example hydroxy, amino, imino, thio or carboxy groups, where these are desired in the final product, to avoid their unwanted participation in the reactions. Conventional protecting groups may be used in accordance with standard practice, for examples see T.W. Greene and P.G.M. Wuts in "Protective Groups in Organic Chemistry" John Wiley and Sons, 1991.

Compounds of formula (I), wherein R^1 , R^2 and L^1 are as hereinbefore defined, and where Y is carboxy, may be prepared by hydrolysis of esters of formula (I), wherein R^1 , R^2 and L^1 are as hereinbefore defined and Y is a $-CO_2R^{19}$ group (in which R^{19} is alkyl, alkenyl or arylalkyl). The hydrolysis may conveniently be carried out by alkaline hydrolysis using a base, such as an alkali metal

hydroxide, e.g., lithium hydroxide, or an alkali metal carbonate, e.g., potassium carbonate, in the presence of an aqueous/organic solvent mixture, using organic solvents such as dioxan, tetrahydrofuran or methanol, at a temperature from about ambient to about reflux. The hydrolysis of the esters may also be carried out by acid hydrolysis using an inorganic acid, such as hydrochloric acid, in the presence of an aqueous/inert organic solvent mixture, using organic solvents such as dioxan or tetrahydrofuran, at a temperature from about 50°C to about 80°C.

As another example compounds of formula (I), wherein R^1 , R^2 and L^1 are as hereinbefore defined, and where Y is carboxy, may be prepared by acid catalysed removal of the tert-butyl group of tert-butyl esters of formula (I), wherein R^1 , R^2 and L^1 are as hereinbefore defined and Y is a $-CO_2R^{19}$ group (in which R^{19} is tert-butyl), using standard reaction conditions, for example reaction with trifluoroacetic acid at a temperature at about room temperature.


As another example compounds of formula (I), wherein R^1 , R^2 and L^1 are as hereinbefore defined, and where Y is carboxy, may be prepared by hydrogenation of compounds of formula (I) wherein R^1 , R^2 and L^1 are as hereinbefore defined and Y is a $-CO_2R^{19}$ group (in which R^{19} is arylmethyl, e.g., benzyl). The reaction may be carried out in the presence of ammonium formate and a suitable metal catalyst, e.g., palladium, supported on an inert carrier such as carbon, preferably in a solvent such as methanol or ethanol and at a temperature at about reflux temperature. The reaction may alternatively be carried out in the presence of a suitable metal catalyst, e.g., platinum or palladium optionally supported on an inert carrier such as carbon, preferably in a solvent such as methanol or ethanol. This reaction is most suitable for compounds of formula (I) where L^1 does not contain carbon-carbon multiple bonds.

In a process A compounds of formula (I), containing an amide bond may be prepared by coupling of an acid [or an acid halide (or anhydride)] with an amine to give an amide bond using standard peptide coupling procedures as described hereinafter.

As an example of process A, esters of formula (I), wherein R^1 , R^2 and L^1 are as hereinbefore defined, and Y is a $-CO_2R^{19}$ group (in which R^{19} is as hereinbefore defined), may be prepared by reacting a compound of formula (II):-



wherein R^{20} is optionally substituted aryl, optionally substituted heteroaryl, $R^3NH-Ar^1-L^2$ - or $R^3-NH-C(=O)-NH-Ar^2-L^2$ -, and X^1 is a hydroxy group, a halogen atom, preferably chlorine, or $-O-C(=O)-R^{20}$ with an amine of formula (III):-



(III)


wherein R^2 , R^{19} and L^1 are as hereinbefore defined. When X^1 is a hydroxy group the reaction may be carried out using standard peptide coupling procedures for example coupling in the presence of O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate and triethylamine (or diisopropylethylamine) in tetrahydrofuran (or dimethylformamide), at room temperature. When X^1 is a halogen atom the acylation reaction may be carried out with the aid of a base, such as pyridine, preferably in a solvent such as tetrahydrofuran and at a temperature at about room temperature. When X^1 is $-O-C(=O)-R^{20}$ the reaction may be carried out in an inert solvent, such as dichloromethane, optionally in the presence of a base, such as triethylamine, and at a temperature at about room temperature.

Esters of formula (I), wherein R^1 and R^2 are as hereinbefore defined, Y is a $-CO_2R^{19}$ group (in which R^{19} is as hereinbefore defined) and L^1 contains a $-N(R^6)-C(=O)-R^4$ group (in which R^4 and R^6 are as hereinbefore defined) may be similarly prepared from the corresponding esters of formula (I) where L^1 contains a $-NHR^6$ group (in which R^6 is as hereinbefore defined) by reaction with a compound of formula $R^4-C(=O)-X^1$ wherein R^4 and X^1 are as hereinbefore defined.

Esters of formula (I), wherein R^1 and R^2 are as hereinbefore defined, Y is a $-CO_2R^{19}$ group (in which R^{19} is as hereinbefore defined) and L^1 contains a $-N(R^6)-SO_2-R^4$ group (in which R^4 and R^6 are as hereinbefore defined), may be prepared from the corresponding esters of formula (I) where L^1 contains a $-NHR^6$ group (in which R^6 is as hereinbefore defined) by reaction with sulfonyl chlorides of formula R^4-SO_2Cl wherein R^4 is as hereinbefore defined. The reaction is preferably carried out with the aid of a base, such as a tertiary amine, for example triethylamine, preferably in a solvent such as tetrahydrofuran and at a temperature from about $0^\circ C$ to about room temperature.

Esters of formula (I), wherein R^1 and R^2 are as hereinbefore defined, Y is a $-\text{CO}_2\text{R}^{19}$ group (in which R^{19} is as hereinbefore defined) and L^1 contains a $-\text{N}(\text{R}^6)-\text{C}(=\text{O})-\text{OR}^4$ group (in which R^4 and R^6 are as hereinbefore defined), may be prepared from the corresponding derivatives of formula (I) where L^1 contains a $-\text{NHR}^6$ group (in which R^6 is as hereinbefore defined) by reaction with compounds of formula $\text{R}^4\text{O}-\text{C}(=\text{O})-\text{X}^1$ wherein R^4 and X^1 are as hereinbefore defined, in the presence of a suitable base, such as triethylamine or pyridine, and at a temperature from about 0°C to about room temperature.

10 Esters of formula (I), wherein R^1 and R^2 are as hereinbefore defined, Y is a $-\text{CO}_2\text{R}^{19}$ group (in which R^{19} is alkyl) and L^1 is $\text{---CH(NH}_2\text{)---CH}_2\text{---}$ (or $\text{---CH(NH}_2\text{)---CH}_2\text{---}$) may be prepared by hydrogenation of the corresponding derivatives of formula (I), where L^1 is $\text{---CH(N(CH}_2\text{Ph)}_2\text{)---CH}_2\text{---}$ (or




15 suitable metal catalyst, e.g., palladium, supported on an inert carrier such as carbon, at a temperature at about 60°C . The reaction may conveniently be carried out in the presence of a suitable metal catalyst, e.g., platinum or palladium optionally supported on an inert carrier such as carbon, preferably in a solvent such as methanol or ethanol.

20 Esters of formula (I), wherein R^1 and R^2 are as hereinbefore defined, Y is a $-\text{CO}_2\text{R}^{19}$ group (in which R^{19} is as hereinbefore defined) and L^1 is a $\text{---CH(NH}_2\text{)---CH}_2\text{---}$ (or $\text{---CH(NH}_2\text{)---CH}_2\text{---}$) linkage, may


also be obtained from the racemic mixture following standard recrystallisation of a suitable salt (for example recrystallisation of the tartrate salt), or by the application of standard enzymatic resolution procedures (for example those described by Soloshonok, V. A., et.al., Tetrahedron: Asymmetry 6 (1995) 7, 1601-1610).

Esters of formula (I), wherein R^1 and R^2 are as hereinbefore defined, Y is a $-\text{CO}_2\text{R}^{19}$ group (in which R^{19} is as hereinbefore defined) and L^1 is a $-\text{CH}-\text{CH}_2-$ or



may be prepared by reacting an ester of formula (I), wherein R^1 and R^2 are as hereinbefore defined, Y is a $-\text{CO}_2\text{R}^{19}$ group (in which R^{19} is as hereinbefore defined) and L^1 is a $-\text{CH}=\text{CH}-$ linkage, with an alkali metal hydride, such as sodium hydride, in an inert solvent, e.g., tetrahydrofuran, and at a temperature at about room temperature, and subsequent reaction with the anion derived from treating dibenzylamine, or (*S*)-N-benzyl- α -methylbenzylamine, with butyllithium, at a temperature at about -78°C .

- 10 Esters of formula (I), wherein R^1 and R^2 are as hereinbefore defined, Y is a $-\text{CO}_2\text{R}^{19}$ group (in which R^{19} is as hereinbefore defined) and L^1 is alkenylene, alkynylene or cycloalkenylene in which the aliphatic carbon-carbon multiple bond is attached directly to the phenyl moiety in formula (I), may be prepared by coupling of compounds of formula (IV):-



wherein R^1 and R^2 are as hereinbefore defined and X^2 is a halogen, preferably bromine or iodine, atom with a compound of formula (V):-



wherein R^{19} is as hereinbefore defined and R^{21} is alkenyl, alkynyl or cycloalkenyl. When X^2 is a bromine or iodine atom the reaction may be conveniently carried out in the presence of palladium acetate, a triarylphosphine, such as tri-*o*-tolylphosphine, and a tertiary amine, such as tributylamine, at a temperature up to about 110°C . This reaction is particularly suitable for the preparation of esters of formula (I) in which L^1 is vinylene. When X^2 is a chlorine atom the reaction may be conveniently

carried out in the presence of sodium iodide, nickel bromide, palladium(0) bis(dibenzylideneacetone), a triarylphosphine, such as tri-*o*-tolylphosphine, and a tertiary amine, such as tributylamine, at a temperature up to about 110°C.

- 5 According to a further feature of the present invention, compounds of the invention may be prepared by interconversion of other compounds of the invention.

For example compounds of formula (I), wherein R¹, R² and L¹ are as hereinbefore defined and Y is a group -C(=O)-NHOH, may be prepared by reacting compounds of formula (I), wherein R¹, R² and L¹ are as hereinbefore defined and Y is carboxy, with hydroxylamine using standard peptide coupling
10 procedures such as treatment with a carbodiimide, for example dicyclohexylcarbodiimide, in the presence of triethylamine, in an inert solvent such as dichloromethane or tetrahydrofuran and at a temperature at about room temperature. The coupling may also be carried out using 1-hydroxybenzotriazole and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide in dichloromethane at room
15 temperature. The preparation may also be carried out using an O-protected hydroxylamine such as O-(trimethylsilyl)hydroxylamine, O-(*t*-butyldimethylsilyl)-hydroxylamine, or O-(tetrahydropyranyl)hydroxylamine followed by treatment with acid.

As another example of the interconversion process, compounds of formula (I) containing sulfoxide
20 linkages may be prepared by the oxidation of corresponding compounds containing -S- linkages. For example, the oxidation may conveniently be carried out by means of reaction with a peroxyacid, e.g., 3-chloroperbenzoic acid, preferably in an inert solvent, e.g., dichloromethane, preferably at or near room temperature, or alternatively by means of potassium hydrogen peroxomonosulfate in a medium such as aqueous methanol, buffered to about pH5, at temperatures between about 0°C and room
25 temperature. This latter method is preferred for compounds containing an acid-labile group.

As another example of the interconversion process, compounds of formula (I) containing sulfone linkages may be prepared by the oxidation of corresponding compounds containing -S- or sulfoxide linkages. For example, the oxidation may conveniently be carried out by means of reaction with a
30 peroxyacid, e.g., 3-chloroperbenzoic acid, preferably in an inert solvent, e.g., dichloromethane, preferably at or near room temperature.

As another example of the interconversion process, compounds of formula (I), wherein R¹, R² and Y are as hereinbefore defined, and L¹ is optionally substituted alkylene, may be prepared by
35 hydrogenation of the corresponding compounds of formula (I) in which L¹ is the corresponding

optionally substituted alkenylene. The hydrogenation may be carried out using hydrogen (optionally under pressure) in the presence of a suitable metal catalyst, e.g., platinum or palladium optionally supported on an inert carrier such as carbon, preferably in a solvent such as methanol or ethanol, and at a temperature at about room temperature.

5

As another example of the interconversion process, compounds of formula (I) wherein R^1 and R^2 are as hereinbefore defined, L^1 is a $-\text{CH}-\text{CH}_2-$ linkage and Y is carboxy, may be prepared by (i)



reacting an ester of formula (I) wherein R^1 and R^2 are as hereinbefore defined, L^1 is a $-\text{CH}=\text{CH}-$ linkage and Y is $-\text{CO}_2\text{R}^{19}$ (in which R^{19} is as hereinbefore defined) with dimethyl malonate, in the

10 presence of an alkali metal alkoxide, such as sodium methoxide, in methanol and at a temperature at about reflux temperature and (ii) treatment of the resulting compounds of formula (I) wherein R^1 and R^2 are as hereinbefore defined, L^1 is a $-\text{CH}-\text{CH}_2-$ linkage and Y is $-\text{CO}_2\text{R}^{19}$ with hydrochloric

$$\begin{array}{c} | \\ \text{CH}(\text{CO}_2\text{Me})_2 \end{array}$$

acid at reflux temperature.

15 As another example of the interconversion process, compounds of the invention containing a heterocyclic group wherein the hetero atom is a nitrogen atom may be oxidised to their corresponding N-oxides. The oxidation may conveniently be carried out by means of reaction with a mixture of hydrogen peroxide and an organic acid, e.g., acetic acid, preferably at or above room temperature, for example at a temperature of about 60-90°C. Alternatively, the oxidation may be carried out by

20 reaction with a peracid, for example peracetic acid or *m*-chloroperoxybenzoic acid, in an inert solvent such as chloroform or dichloromethane, at a temperature from about room temperature to reflux, preferably at elevated temperature. The oxidation may alternatively be carried out by reaction with hydrogen peroxide in the presence of sodium tungstate at temperatures between room temperature and about 60°C.

25

It will be appreciated that compounds of the present invention may contain asymmetric centres. These asymmetric centres may independently be in either the R or S configuration. It will be apparent to those skilled in the art that certain compounds of the invention may also exhibit geometrical isomerism. It is to be understood that the present invention includes individual geometrical isomers and

30 stereoisomers and mixtures thereof, including racemic mixtures, of compounds of formula (I) hereinabove. Such isomers can be separated from their mixtures, by the application or adaptation of

known methods, for example chromatographic techniques and recrystallisation techniques, or they are separately prepared from the appropriate isomers of their intermediates.

According to a further feature of the invention, acid addition salts of the compounds of this invention
5 may be prepared by reaction of the free base with the appropriate acid, by the application or adaptation of known methods. For example, the acid addition salts of the compounds of this invention may be prepared either by dissolving the free base in water or aqueous alcohol solution or other suitable solvents containing the appropriate acid and isolating the salt by evaporating the solution, or by reacting the free base and acid in an organic solvent, in which case the salt separates directly or can be
10 obtained by concentration of the solution.

The acid addition salts of the compounds of this invention can be regenerated from the salts by the application or adaptation of known methods. For example, parent compounds of the invention can be regenerated from their acid addition salts by treatment with an alkali, e.g., aqueous sodium bicarbonate
15 solution or aqueous ammonia solution.

Compounds of this invention can be regenerated from their base addition salts by the application or adaptation of known methods. For example, parent compounds of the invention can be regenerated from their base addition salts by treatment with an acid, e.g., hydrochloric acid.
20

Compounds of the present invention may be conveniently prepared, or formed during the process of the invention, as solvates (e.g., hydrates). Hydrates of compounds of the present invention may be conveniently prepared by recrystallisation from an aqueous/organic solvent mixture, using organic solvents such as dioxan, tetrahydrofuran or methanol.
25


According to a further feature of the invention, base addition salts of the compounds of this invention may be prepared by reaction of the free acid with the appropriate base, by the application or adaptation of known methods. For example, the base addition salts of the compounds of this invention may be prepared either by dissolving the free acid in water or aqueous alcohol solution or other suitable
30 solvents containing the appropriate base and isolating the salt by evaporating the solution, or by reacting the free acid and base in an organic solvent, in which case the salt separates directly or can be obtained by concentration of the solution.

The starting materials and intermediates may be prepared by the application or adaptation of known
35 methods, for example methods as described in the Reference Examples or their obvious chemical equivalents.

Compounds of formula (II) wherein R^{20} is R^3 -NH-C(=O)-NH-Ar²-L²- group (in which R^3 , Ar² and L² are as hereinbefore defined) may be prepared by the application or adaptation of methods described in the specification of International Patent Application Publication No. WO 96/22966.


5

Acids of formula (II) wherein R^{20} is R^3 -NH-Ar¹-L²- (in which R^3 and L⁴ are as defined above, Ar¹

is  (in which R^{17} is as hereinbefore defined) and X¹ is a hydroxy group may be

prepared by reaction of compounds of formula (1):-

10




wherein R^{17} and L¹ are as hereinbefore defined, R^{16} is C₁₋₄alkyl and X is O, with isothiocyanates of formula R^3 -N=S=O (in which R^3 is as hereinbefore defined) in ethanol and at room temperature, followed by reaction with a carbodiimide, such as dicyclohexylcarbodiimide or diisopropylcarbodiimide in ethanol and at a temperature from about room temperature to about reflux temperature, and subsequent hydrolysis using standard conditions, for example those described hereinbefore.

15

Acids of formula (II) wherein R^{18} is R^3 -NH-Ar¹-L²- (in which R^3 and L² are as hereinbefore defined,

20

Ar¹ is  (in which R is as hereinbefore defined) and X¹ is hydroxy may be similarly


prepared from compounds of formula (1) wherein R^{17} , L¹ and R^{19} are as hereinbefore defined and X is NH.

Acid chlorides of formula (II) wherein R^{20} is as hereinbefore defined and X¹ is a chlorine atom may be prepared from the corresponding acids of formula (II) wherein R^{20} is as hereinbefore defined and

25

X^1 is hydroxy, by the application of standard procedures for the conversion of acids to acid chlorides for example by reaction with oxalyl chloride.


Compounds of formula (III) wherein R^2 , R^{19} and L^1 are as defined hereinbefore, may be prepared by acid catalysed removal of the protecting group in compounds of formula (2):-



wherein R^2 , R^{19} and L^1 are as defined hereinbefore and R^{22} is a suitable protecting group such as -
 10 COO^tBu or $-C(=O)-CF_3$. When R^{22} is $-COO^tBu$ the reaction may conveniently be carried out using trifluoroacetic acid in an inert solvent, such as dichloromethane, and at a temperature at about $0^\circ C$.
 When R^{22} is $-C(=O)-CF_3$ the reaction may conveniently be carried out using a base such as sodium ethoxide in ethanol and at a temperature at about room temperature.

15 Compounds of formula (III) wherein R^2 is as defined hereinbefore, R^{19} is alkyl and L^1 is alkylene or cycloalkylene may be prepared by hydrogenation of compounds of formula (2),
 wherein R^2 is as defined hereinbefore, R^{19} is alkyl, R^{22} is H [or a protecting group (e.g., benzyl) that is conveniently removed during the hydrogenation] and L^1 is alkenylene, alkynylene or cycloalkenylene. The hydrogenation may be carried out using hydrogen (optionally under pressure) in
 20 the presence of a suitable catalyst, e.g., palladium hydroxide supported on an inert carrier such as carbon, in acetic acid and optionally in the presence of a co-solvent such as methanol or ethanol, and at a temperature at about room temperature.

Compounds of formula (III) wherein R^2 is as defined hereinbefore, R^{19} is alkyl and L^1 is alkylene
 25 may be prepared by hydrogenation of isoquinolines of formula (3):-



wherein R^2 is as defined hereinbefore, R^{19} is alkyl and L^1 is alkenylene. The hydrogenation may be carried out using hydrogen (optionally under pressure) in the presence of a suitable catalyst, e.g., platinum oxide, in ethanol and in the presence of an acid such as hydrochloric acid, and at a temperature at about room temperature.


5

Compounds of formula (2) wherein R^2 is as defined hereinbefore, R^{19} is alkyl, R^{22} is a suitable protecting group, such as tertiary-butyloxycarbonyl, and L^1 is alkylene or cycloalkylene may be prepared by reduction compounds of formula (2) wherein R^2 is as defined hereinbefore, R^{19} is alkyl, R^{22} is a suitable protecting group, such as tertiary-butyloxycarbonyl, and L^1 is alkenylene, alkynylene or cycloalkenylene. The reduction may be carried out in the presence of ammonium formate and a suitable metal catalyst, e.g., palladium, supported on an inert carrier such as carbon, preferably in a solvent such as methanol or ethanol and at a temperature at about reflux temperature.

10

15

Compounds of formula (2) wherein R^2 is as defined hereinbefore, R^{19} is alkyl, R^{22} is a protecting group (e.g., benzyl or tertiary-butyloxycarbonyl) and L^1 is optionally substituted alkenylene, optionally substituted alkynylene or cycloalkenylene in which the carbon-carbon multiple bond is attached directly to the phenyl moiety in formula (2), may be prepared by reaction compounds of formula (4):-



(4)


20

wherein R^2 and R^{22} are as just defined and X^3 is an iodine, or preferably a bromine, atom, with a compound of formula (V) wherein R^{19} and R^{21} are as hereinbefore defined using standard Heck reaction conditions, for example reaction in the presence of palladium acetate, triphenylphosphine and tributylamine in an inert solvent, such as dimethylformamide, and at a temperature up to about 155°C.

25


Compounds of formula (3), wherein R^2 and R^{19} are as defined hereinbefore and L^1 is optionally substituted alkenylene in which the carbon-carbon double bond is attached directly to the phenyl moiety in formula (3), may be similarly prepared from compounds of formula (5):-

- 43 -



wherein R^2 and X^3 are as hereinbefore defined.

- 5 Compounds of formula (4) wherein R^2 is hydrogen, X^3 is as hereinbefore defined and R^{22} is a suitable protecting group {e.g., $CF_3C(=O)-$ } may be prepared by reaction of compounds of formula (6):-




- 10 wherein R^{22} and X^3 are as hereinbefore defined, with paraformaldehyde in the presence of acetic acid and sulphuric acid at a temperature at about room temperature.

Compounds of formula (6) wherein R^{22} is $CF_3C(=O)-$ and X^3 is as hereinbefore defined, may be prepared by reaction of the corresponding compounds of formula (6) in which R^{22} is hydrogen with
 15 trifluoroacetic anhydride in the presence of a base, such as 2,6-lutidine, in an inert solvent, such as dichloromethane, and at room temperature.

Compounds of formula (6) wherein R^{22} is hydrogen and X^3 is as hereinbefore defined, may be prepared by reduction of compounds of formula (7):-

20



- wherein X^3 is as hereinbefore defined. The reduction may conveniently be carried out with sodium borohydride in the presence of cobalt chloride, in an inert solvent, such as methanol, and at a
 25 temperature from about $0^\circ C$ to about $10^\circ C$.

Compounds of formula (III), (2) or (3) wherein L^1 is $\text{---CH}_2\text{---CH---}$ may be prepared by standard




methodology for the preparation of α -amino-acids for example those described in Organic Syntheses Based On Name Reactions and Unnamed Reactions, A.Hassner and C.Stumer, Pergamon, pages 275 and 374.

5

Compounds of formula (2) wherein L^1 is $\text{---CH}_2\text{---CH---}$ may be prepared by reaction of compounds



of formula (8):-



(8)

10

wherein R^2 and R^{22} are as defined hereinbefore and X^4 is a bromine or chlorine atom with the anion derived from reaction of (2R) -(-)-2,5-dihydro-3,6-dimethoxy-2-isopropylpyrazine with butyllithium according to the method described by D. L. Boger and D. Yohannes, J. Org. Chem. [JOCEAH], 1990, 55, for the preparation of compound 31 on page 6010.

15

Compounds of formula (2) wherein L^1 is a $\text{---CH---CH}_2\text{---}$ linkage, may be prepared by reacting



compounds of formula (2) wherein L^1 is a ---CH=CH--- linkage, with dimethyl malonate, in the presence of an alkali metal alkoxide, such as sodium methoxide, in methanol and at a temperature at about reflux temperature.

20

The present invention is further Exemplified but not limited by the following illustrative Examples and Reference Examples.

25

High Pressure Liquid Chromatography/ Mass Spectrometry (LC/MS) conditions for determination of retention times (R_T) were as follows: 3 micron Luna C18 (2) HPLC column (30mm x 4.6mm)

operated under gradient elution conditions with mixtures of (A) water containing 0.1% formic acid and (B) acetonitrile containing 0.1% formic acid as the mobile phase gradient : 0.00 minutes, 95%A:5%B;

0.50 minutes, 95%A:5%B; 4.50 minutes, 5%A:95%B; 5.00 minutes, 5%A:95%B; 5.50 minutes, 95%A:5%B; flow rate 2ml/minute with approximately 200 μ l/minute split to the Mass Spectrometer; injection volume 10-40 μ l; in line Diode Array (220-450nm), in line Evaporative light scattering (ELS) detection ELS - temperature 50°C, Gain 8 - 1.8ml/minute; Source temperature 150°C.

5

EXAMPLE 1

(a) 3-(((4-Methyl-2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-butanoic acid

A stirred solution of 3-(((4-methyl-2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-butanoic acid ethyl ester [0.1g, Reference Example 1(a)] in ethanol (20mL), under argon and at room temperature, was treated with sodium hydroxide solution (1mL, 1M) and then heated at reflux temperature for 2.5 hours. The reaction mixture was evaporated to low volume (approximately 2mL) and then treated with water (12mL). This mixture was acidified to pH 1 by addition of hydrochloric acid (1.75mL, 1M) and cooled in ice. The resulting solid was filtered, then washed three times with water (3mL) and then dried at 60°C under vacuum to give the title compound as a white solid. LC-MS: $R_T = 3.53$ minutes; MS(ES) = 498 (M+H)⁺.

(b) By proceeding in a similar manner to Example 1(a) but using 3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-butanoic acid ethyl ester {Reference Example 1(b)} there was prepared 3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-butanoic acid as a white solid. LC-MS: $R_T = 3.36$ minutes; MS(ES) = 484 (M+H)⁺.

(c) By proceeding in a similar manner to Example 1(a) but using 3-phenyl-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-propanoic acid ethyl ester {Reference Example 1(c)} there was prepared 3-phenyl-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-propanoic acid as a white solid. LC-MS: $R_T = 3.57$ minutes; MS(ES) = 546 (M+H)⁺.

(d) By proceeding in a similar manner to Example 1(a) but using 3-cyclohexyl-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-propanoic acid ethyl ester [Reference Example 1(d)] there was prepared 3-cyclohexyl-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-propanoic acid as a white solid. LC-MS: $R_T = 3.98$ minutes; MS(ES) = 552 (M+H)⁺.

(e) By proceeding in a similar manner to Example 1(a) but using a mixture of 3-(pyrid-4-yl)-3-
{((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl}-propanoic acid ethyl
ester trifluoroacetate and 3-(pyrid-4-yl)-3-{((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-
tetrahydro-isoquinolin-7-yl}-propanoic acid methyl ester trifluoroacetate {Reference Example 1(e)}
5 and subjecting the product to preparative HPLC under gradient elution conditions with mixtures of
acetonitrile and water containing 0.1% trifluoroacetic acid (1:4 initially then a gradient of 1%
acetonitrile /minute) there was prepared 3-(pyrid-4-yl)-3-{((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-
1,2,3,4-tetrahydro-isoquinolin-7-yl}-propanoic acid as a white solid. LC-MS: $R_T = 2.45$ minutes;
MS(ES) = 547 (M+H)⁺.

(f) By proceeding in a similar manner to Example 1(a) but using 3-{((2-*o*-tolylamino-benzoxazol-
6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-8-yl}-but-2-enoic acid ethyl ester {Reference Example
1(f)} there was prepared 3-{((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-
8-yl}-but-2-enoic acid as a white solid. LC-MS: $R_T = 3.38$ minutes; MS(ES) = 482 (M+H)⁺.

(g) By proceeding in a similar manner to Example 1(a) but using 3-{((2-*o*-tolylamino-benzoxazol-
6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-8-yl}-butanoic acid ethyl ester {Reference Example 1(g)}
and subjecting the product to preparative HPLC on a Hypersil Elite column (10cm x 2.1cm) under
gradient elution conditions with mixtures of acetonitrile and water containing 0.1% trifluoroacetic acid
20 (3:10 initially then a gradient of 1% acetonitrile /minute) with a flow rate of 5mL/minute there was
prepared 3-{((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-8-yl}-butanoic
acid as a white solid. LC-MS: $R_T = 3.42$ minutes; MS(ES) = 484(M+H)⁺.

(h) By proceeding in a similar manner to Example 1(a) but using 3-{2-[(2-*o*-tolylamino-
25 benzoxazol-6-yl)-carbonyl]-1,2,3,4-tetrahydro-isoquinolin-6-yl}-butanoic acid ethyl ester {Reference
Example 1(h)} there was prepared 3-{2-[(2-*o*-tolylamino-benzoxazol-6-yl)-carbonyl]-1,2,3,4-
tetrahydro-isoquinolin-6-yl}-butanoic acid as a white solid. LC-MS: $R_T = 3.36$ minutes; MS(ES) =
470(M+H)⁺, 468(M-H)⁻.

(i) By proceeding in a similar manner to Example 1(a) but using {5-(3-methoxy-4-[3-(2-
30 methylphenyl)ureido]phenylacetylamino)-1,2,3,4-tetrahydro-isoquinolin-8-yl}-butanoic acid ethyl ester
{Reference Example 1(i)} there was prepared {5-(3-methoxy-4-[3-(2-methylphenyl)ureido]-
phenylacetylamino)-1,2,3,4-tetrahydro-isoquinolin-8-yl}-butanoic acid as a white solid.

(j) By proceeding in a similar manner to Example 1(a) but using 2-(2,6-dichloro-benzoylamino)-3-[2-(2,6-dichloro-benzoyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl]-propionic acid methyl ester (Reference Example 9) there was prepared 2-(2,6-dichloro-benzoylamino)-3-[2-(2,6-dichloro-benzoyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl]-propionic acid as a pale yellow solid.

5

(k) By proceeding in a similar manner to Example 1(a) but using 3-phenyl-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl}-propanoic acid ethyl ester {Reference Example 1(j)} there was prepared 3-phenyl-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl}-propanoic acid as a white solid. LC-MS: $R_T = 3.62$ minutes, MS(ES) =

10 $546(M+H)^+$, $544(M-H)^-$.

(l) By proceeding in a similar manner to Example 1(a) but using 3-((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl}-butanoic acid ethyl ester {Reference Example 1(k)} there was prepared 3-((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl}-
 15 butanoic acid as a white solid, m.p. softens from 100°C and melts 123-140°C.

(m) By proceeding in a similar manner to Example 1(a) but using 3-(pyrid-4-yl)-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid ethyl ester, enantiomer A {Reference Example 16(a)} there was prepared 3-(pyrid-4-yl)-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid, enantiomer A. LC-
 20 MS: $R_T = 2.45$ minutes, MS(ES) = $547(M+H)^+$, $545(M-H)^-$.

(n) By proceeding in a similar manner to Example 1(a) but using 3-(pyrid-4-yl)-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid ethyl ester, enantiomer B {Reference Example 16(b)} there was prepared 3-(pyrid-4-yl)-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid, enantiomer B. LC-
 25 MS: $R_T = 2.45$ minutes, MS(ES) = $547(M+H)^+$, $545(M-H)^-$.

REFERENCE EXAMPLE 1

30 (a) 3-(((4-Methyl-2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-butanoic acid ethyl ester

A stirred solution of (4-methyl-2-(2-*o*-tolylamino)-benzoxazol-6-yl)-acetic acid (0.3g, prepared according to the procedure described for Reference Example 9(b) in International Patent Application Publication No. WO 00/49005) in dry dimethylformamide (7.5mL), under an argon atmosphere, was
 35 treated with O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (0.7g) then

with di-isopropylethylamine (0.8mL). After stirring at room temperature for 10 minutes the mixture was treated with a solution of ethyl 3-(1,2,3,4-tetrahydroisoquinolin-6-yl)-butanoate hydrochloride {0.38g, Reference Example 2(a)} in dry dimethylformamide (7.5mL) followed by di-isopropylethylamine (0.4mL). The resulting dark solution was stirred at room temperature overnight then evaporated. The residual dark oil was treated with water (20mL) and the mixture was acidified to pH 1 by addition of hydrochloric acid (5mL, 1M) and then extracted with ethyl acetate (25mL). The organic extract was washed with water (20mL) then with brine (15mL) then dried over magnesium sulfate and then evaporated. The residue was subjected to preparative HPLC on a Hypersil Elite column (10cm x 2.1cm) under gradient elution conditions with mixtures of acetonitrile and water containing 0.1% trifluoroacetic acid (45:55 initially, rising at a rate of 1% acetonitrile/minute) with a flow rate of 5mL/minute to give the title compound as an amber coloured glass. LC-MS: $R_T = 4.14$ minutes; $MS(ES) = 526 (M+H)^+$, $MS(ES^-) = 526 (M-H)^-$.

(b) By proceeding in a similar manner to Reference Example 1(a) but using 2-*o*-tolylaminobenzoxazole-6-acetic acid (prepared according to the procedure described for Reference Example 4 in International Patent Application Publication No. WO 00/49005) and ethyl 3-(1,2,3,4-tetrahydroisoquinolin-7-yl)-butanoate {Reference Example 4(a)} there was prepared 3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-butanoic acid ethyl ester as a clear oil. LC-MS: $R_T = 3.93$ minutes; $MS(ES) = 512 (M+H)^+$, $MS(ES^-) = 510 (M-H)^-$.

(c) and (d) By proceeding in a similar manner to Reference Example 1(a) but using 2-*o*-tolylaminobenzoxazole-6-acetic acid and a mixture of 3-phenyl-3-(1,2,3,4-tetrahydroisoquinolin-6-yl)-propanoic acid ethyl ester hydrochloride and 3-cyclohexyl-3-(1,2,3,4-tetrahydroisoquinolin-6-yl)-propanoic acid ethyl ester hydrochloride {Reference Example 2(b)} and subjecting the product to preparative chromatography on a Hypersil Elite column (10cm x 2.1cm) under gradient elution conditions with mixtures of acetonitrile and water containing 0.1% trifluoroacetic acid (50:50 initially, rising at a rate of 1% acetonitrile/minute) with a flow rate of 5mL/minute there was prepared 3-phenyl-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-propanoic acid ethyl ester, Reference Example 1(c), as a pale yellow gum {LC-MS: $R_T = 4.11$ minutes; $MS(ES) = 574 (M+H)^+$ } and 3-cyclohexyl-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-propanoic acid ethyl ester, Reference Example 1(d), as a pale yellow gum {LC-MS: $R_T = 4.61$ minutes; $MS(ES) = 580 (M+H)^+$ }.

(e) By proceeding in a similar manner to Reference Example 1(a) but using 2-*o*-tolylaminobenzoxazole-6-acetic acid and a mixture of 3-(pyrid-4-yl)-3-(1,2,3,4-tetrahydro-isoquinolin-

7-yl)-propanoic acid methyl ester and 3-(pyrid-4-yl)-3-(1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid ethyl ester {Reference Example 4(b)} there was prepared a mixture of 3-(pyrid-4-yl)-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid ethyl ester trifluoroacetate and 3-(pyrid-4-yl)-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid methyl ester trifluoroacetate {Reference Example 1(e)}.

(f) and (g) By proceeding in a similar manner to Reference Example 1(a) but using 2-o-tolylaminobenzoxazole-6-acetic acid and a mixture of 3-(1,2,3,4-tetrahydroisoquinolin-8-yl)-but-2-enoic acid ethyl ester hydrochloride and 3-methyl-3-(1,2,3,4-tetrahydroisoquinolin-8-yl)-propanoic acid ethyl ester hydrochloride {Reference Example 2(c)} and subjecting the product to preparative chromatography on a Hypersil Elite column (10cm x 2.1cm) under gradient elution conditions with mixtures of acetonitrile and water containing 0.1% trifluoroacetic acid (35:65 initially, rising at a rate of 1% acetonitrile/minute) with a flow rate of 5mL/minute there was prepared 3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-8-yl)-but-2-enoic acid ethyl ester, Reference Example 1(f), {LC-MS: $R_T = 3.98$ minutes; MS(ES) = 510 (M+H)⁺} and 3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-8-yl)-butanoic acid ethyl ester, Reference Example 1(g), {LC-MS: $R_T = 3.92$ minutes; MS(ES) = 512 (M+H)⁺}.

(h) By proceeding in a similar manner to Reference Example 1(a) but using 2-o-tolylaminobenzoxazole-6-carboxylic acid there was prepared 3-{2-[(2-o-tolylamino-benzoxazol-6-yl)-carbonyl]-1,2,3,4-tetrahydro-isoquinolin-6-yl}-butanoic acid ethyl ester as a pale amber coloured glass. LC-MS: $R_T = 3.97$ minutes; MS(ES) = 498(M+H)⁺, 496(M-H)⁻.

(i) By proceeding in a similar manner to Reference Example 1(a) but using {5-(3-methoxy-4-[3-(2-methylphenyl)ureido]phenyl)acetic acid (prepared as described in Example 21 of International Patent Application Publication No. WO 96/22966) there was prepared {5-(3-methoxy-4-[3-(2-methylphenyl)ureido]phenyl)acetyl-amino)-1,2,3,4-tetrahydro-isoquinolin-8-yl}-butanoic acid ethyl ester as a fawn coloured glass. TLC: $R_F = 0.08$ (silica plates, cyclohexane:ethylacetate, 1:1).

(j) By proceeding in a similar manner to Reference Example 1(a) but using 2-o-tolylaminobenzoxazole-6-acetic acid and 3-phenyl-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)-propanoic acid ethyl ester {Reference Example 4(c)} there was prepared 3-phenyl-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid ethyl ester as a pale yellow gum. LC-MS: $R_T = 4.17$ minutes; MS(ES) = 574(M+H)⁺, 572(M-H)⁻.

(k) By proceeding in a similar manner to Reference Example 1(a) but using 2-*o*-tolylaminobenzoxazole-6-acetic acid there was prepared 3-((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl}-butanoic acid ethyl ester as a clear gum.

5

REFERENCE EXAMPLE 2

(a) 3-(1,2,3,4-Tetrahydroisoquinolin-6-yl)-butanoic acid ethyl ester hydrochloride

A solution of 3-(isoquinolin-6-yl)-but-2-enoic acid ethyl ester {0.5g, Reference Example 3(a)} in ethanol (30mL) was treated with concentrated hydrochloric acid (1.7mL). This stirred solution was
10 flushed with nitrogen then treated with platinum oxide (0.07g) and then hydrogenated in the presence of hydrogen at 2 bar for 19 hours. The reaction mixture was filtered through Hyflo supercel and the filter pad was washed three times with ethanol (20mL). The combined filtrate plus washings were evaporated to give the title compound as a pale green solid. MS(ES) = 248 (M+H)⁺.

15 (b) By proceeding in a similar manner to Reference Example 2(a) but using 3-(isoquinolin-6-yl)-3-phenyl-prop-2-enoic acid ethyl ester {Reference Example 3(b)} there was prepared a mixture of 3-phenyl-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)-propanoic acid ethyl ester hydrochloride {LC-MS: R_T = 2.36 minutes; MS(ES) = 310 (M+H)⁺} and ethyl 3-cyclohexyl-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)-propanoate hydrochloride {LC-MS: R_T = 2.73 minutes; MS(ES) = 316 (M+H)⁺} as an amber oil which
20 was used without further purification.

(c) By proceeding in a similar manner to Reference Example 2(a) but using 3-(isoquinolin-8-yl)-but-2-enoic acid ethyl ester hydrochloride {Reference Example 3(c)} there was prepared a mixture of 3-(1,2,3,4-tetrahydroisoquinolin-8-yl)-but-2-enoic acid ethyl ester hydrochloride and 3-methyl-3-(1,2,3,4-tetrahydroisoquinolin-8-yl)-propanoic acid ethyl ester hydrochloride as a green glass.
25

REFERENCE EXAMPLE 3

(a) 3-(Isoquinolin-6-yl)-but-2-enoic acid ethyl ester

A stirred solution of 6-bromoisquinoline (1.75g) in dry dimethylformamide (15mL), under argon, was
30 treated with ethyl crotonate(1.7mL) then with palladium (II) acetate (0.14g), then with tri-(*o*-tolyl)phosphine (0.3g) and then with tributylamine (8mL). The suspension was stirred, under argon, at 140°C for 3.5 hours, then stood at room temperature for 3 days and then evaporated. The residual dark oil was treated with ethyl acetate (50mL) and the resulting solution was washed with water (30mL) then twice with brine (25mL), then dried over magnesium sulfate and then evaporated. The residue
35 was subjected to flash chromatography on silica eluting initially with cyclohexane and then with a

mixture of cyclohexane and ethyl acetate (9:1,v/v) to give the title compound as an amber coloured oil which solidified on cooling. MS(ES) = 242(M+H)⁺.

(b) By proceeding in a similar manner to Reference Example 3(a) but using ethyl trans-cinnamate there was prepared 3-(isoquinolin-6-yl)-3-phenyl-prop-2-enoic acid ethyl ester as a viscous amber oil. MS(ES) = 304(M+H)⁺.

(c) By proceeding in a similar manner to Reference Example 3(a) but using 8-bromoisquinoline there was prepared 3-(isoquinolin-8-yl)-but-2-enoic acid ethyl ester hydrochloride. LC-MS: R_T = 2.243 minutes; MS(ES) = 242 (M+H)⁺.

REFERENCE EXAMPLE 4

(a) 3-(1,2,3,4-Tetrahydroisoquinolin-7-yl)-butanoic acid ethyl ester

A stirred solution of 3-(2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-butanoic acid ethyl ester {0.5g, Reference Example 5(a)} in ethanol (30mL), under argon, was treated with sodium ethoxide (0.25g). After stirring at room temperature for 1 hour the reaction mixture was evaporated. The residue was treated with ethyl acetate (30mL) and the resulting solution was washed twice with saturated ammonium chloride solution (15mL), then dried over magnesium sulfate and then evaporated to give the title compound as a yellow gum. MS(ES) = 248(M+H)⁺.

(b) By proceeding in a similar manner to Reference Example 4(a) but using 3-(pyrid-4-yl)-3-(2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-propanoic acid methyl ester {Reference Example 5(b)}, methanol and sodium methoxide there was prepared a mixture of 3-(pyrid-4-yl)-3-(1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid methyl ester and 3-(pyrid-4-yl)-3-(1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid ethyl ester as a yellow oil which was used without further purification.

(c) By proceeding in a similar manner to Reference Example 4(a) but using 3-phenyl-3-(2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-propanoic acid ethyl ester {Reference Example 5(c)} there was prepared 3-phenyl-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)-propanoic acid ethyl ester as a fawn coloured gum. LC-MS: R_T = 2.21 minutes; MS(ES) = 310 (M+H)⁺.

REFERENCE EXAMPLE 5

(a) 3-(2-Trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-butanoic acid ethyl ester

A mixture of 3-(2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-acrylic acid ethyl ester {1.5g, Reference Example 6(a)}, industrial methylated spirits (80mL) and palladium on carbon (0.15g, 10%) was stirred under hydrogen for 18 hours. The reaction mixture was filtered through diatomaceous earth and the filter pad was washed with industrial methylated spirits. The combined filtrate and washings were evaporated to give the title compound as a colourless oil.

(b) By proceeding in a similar manner to Reference Example 5(a) but using 3-(pyrid-4-yl)-3-(2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-prop-2-enoic acid methyl ester {Reference Example 6(b)} there was prepared 3-(pyrid-4-yl)-3-(2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-propanoic acid methyl ester. LC-MS: $R_T = 2.33$ minutes; $MS(ES) = 393 (M+H)^+$.

(c) By proceeding in a similar manner to Reference Example 5(a) but using 3-phenyl-3-(2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-prop-2-enoic acid ethyl ester {Reference Example 6(c)} there was prepared 3-phenyl-3-(2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-propanoic acid ethyl ester as a yellow gum. LC-MS: $R_T = 4.20$ minutes; $MS(ES) = 406 (M+H)^+$, $428(M+Na)^+$.

REFERENCE EXAMPLE 6

(a) 3-(2-Trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-acrylic acid ethyl ester

A solution of 7-bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinoline (2.6g, Reference Example 7), tri-*o*-tolylphosphine (0.37g), palladium (II) acetate, ethyl acrylate (1.6mL) and tributylamine (8.01mL) in dry dimethylformamide (25mL), under argon, was heated at 155°C for 3 hours then stood at room temperature overnight. The reaction mixture was evaporated and the resulting black oil was partitioned between ethyl acetate (200mL) and hydrochloric acid (200mL, 2M). The organic phase was washed with hydrochloric acid (200mL, 2M), then with water (100mL), then with brine (100mL), then dried over magnesium sulfate and then evaporated. The residual yellow oil was subjected to flash chromatography on silica under gradient elution conditions with a mixture of ethyl acetate and pentane (from 3:97 to 10:90, v/v) to give the title compound. LC-MS: $R_T = 4.05$ minutes; no molecular ion observed.

(b) By proceeding in a similar manner to Reference Example 6(a) but using 3-(pyrid-4-yl)-prop-2-enoic acid methyl ester and subjecting the product to reverse phase chromatography on a Hypersil Elite C18 column (10cm x 2.1cm) under gradient elution conditions with mixtures of acetonitrile and water containing 0.1% trifluoroacetic acid (25:75 initially, rising at a rate of 1% acetonitrile/minute) there was prepared 3-(pyrid-4-yl)-3-(2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-prop-2-enoic acid methyl ester. LC-MS: $R_T = 2.63$ minutes; $MS(ES) = 391 (M+H)^+$.

(c) By proceeding in a similar manner to Reference Example 6(a) but using ethyl *trans*-cinnamate and subjecting the product to flash chromatography on silica eluting initially with cyclohexane and then with a mixture of cyclohexane and ethyl acetate (9:1, v/v) there was prepared 3-phenyl-3-(2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-prop-2-enoic acid ethyl ester as an amber oil which solidified on standing.

REFERENCE EXAMPLE 7

7-Bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinoline

Concentrated sulfuric acid (35mL) was added to glacial acetic acid (45mL) with stirring and ice-cooling. After stirring for 30 minutes the mixture was treated with N-2-(4-bromophenyl)ethyl trifluoroacetamide (10.4g, Reference Example 8) and then with paraformaldehyde (1.75g). This mixture was allowed to warm to room temperature and after stirring for a further 24 hours the pale brown solution was treated with a further aliquot of paraformaldehyde (0.8g). After stirring for a further 24 hours the reaction mixture was treated with iced-water (500mL) and then extracted three times with ethyl acetate (200mL). The combined extracts were washed twice with water (200mL), then four times with aqueous sodium bicarbonate solution (200mL), then twice with water (200mL), then dried over magnesium sulfate and then evaporated. The residual yellow oil was subjected to flash chromatography on silica eluting with a mixture of ethyl acetate and pentane (3:97,v/v) to give the title compound as a white crystalline solid. LC-MS: $R_T = 4.76$ minutes; no molecular ion observed.

REFERENCE EXAMPLE 8

N-2-(4-Bromophenyl)ethyl trifluoroacetamide

A stirred solution of 2-(4-bromophenyl)ethylamine (7.9g, prepared according to the procedure described by in J.Org.Chem., 1990, page 4530) and 2,6-lutidine (10.2mL) in dry dichloromethane (150mL), cooled in an ice-bath, was treated dropwise with trifluoroacetic anhydride (6.21mL). After stirring for 2 hours and standing at room temperature for 72 hours the reaction mixture was treated with water (100mL) then stirred for a further 20 minutes. The organic phase was separated and then washed twice with hydrochloric acid (150mL), then with water (100mL), then with saturated aqueous sodium bicarbonate solution, then with water (100mL), then dried over magnesium sulfate and then evaporated to give the title compound as a yellow solid. MS(ES) 294 and 296 (M-H)⁻.

REFERENCE EXAMPLE 9

(a) 2-(2,6-Dichloro-benzoylamino)-3-[2-(2,6-dichloro-benzoyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl]-propionic acid methyl ester

A stirred solution of 2-amino-3-[2-(2,6-dichloro-benzoyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl]-propionic acid methyl ester (1.18g, Reference Example 10) in dry pyridine (50mL), under nitrogen and at room temperature, was treated dropwise with 2,6-dichlorobenzoyl chloride (0.67g). The mixture was stirred at room temperature for 25 minutes and then at 65°C for 40 minutes. The resulting dark red solution was cooled to room temperature and then evaporated. The residual red oil was dissolved in dichloromethane and this solution was washed with aqueous sodium bicarbonate solution (5%), then with water, then dried over magnesium sulfate and then evaporated. The residue was coated onto silica (8.5g) using dichloromethane and then subjected to flash chromatography on silica eluting with a mixture of ethyl acetate and heptane (1:1, v/v) to give the title compound.

(b) By proceeding in a similar manner to Reference Example 9(a) but using 2-*tert*-butyloxycarbonylamino-1,2,3,4-tetrahydro-isoquinolin-7-yl]-propionic acid methyl ester {Reference Example 11(a)} there was prepared 2-*tert*-butyloxycarbonylamino-3-[2-(2,6-dichloro-benzoyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl]-propionic acid methyl ester as a pale yellow solid.

REFERENCE EXAMPLE 10

2-Amino-3-[2-(2,6-dichloro-benzoyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl]-propionic acid methyl ester

A solution of 2-*tert*-butyloxycarbonylamino-3-[2-(2,6-dichloro-benzoyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl]-propionic acid methyl ester {1.7g, Reference Example 9(b)} in a mixture of trifluoroacetic acid (10mL) and dichloromethane (30mL), under nitrogen, was stirred at room temperature for 1.5 hours then evaporated. The residue was dissolved in dichloromethane and this solution was washed with aqueous sodium bicarbonate solution, then with water, then dried over magnesium sulfate and then evaporated to give the title compound (1.18g) which was used without further purification.

REFERENCE EXAMPLE 11

2-*tert*-Butyloxycarbonylamino-1,2,3,4-tetrahydro-isoquinolin-7-yl]-propionic acid methyl ester

A solution of 2-*tert*-butyloxycarbonylamino-3-[2-trifluoroacetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl]-propionic acid methyl ester (286g, Reference Example 12) in methanol (120mL) was treated with potassium carbonate solution (10mL, 1%) at room temperature. After stirring for 1.5 hours the reaction mixture was treated with a further aliquot of potassium carbonate solution (10mL, 1%), then a further aliquot (10mL) after 2.5 hours and then a further aliquot (30mL) after 5.5 hours. The reaction mixture was then stirred at room temperature overnight and then evaporated. The residue was treated with dichloromethane and water and the aqueous phase was extracted three times with dichloromethane. The combined organics were washed with water, then dried over magnesium sulfate and then evaporated to give the title compound (2.14g).

- (b) By proceeding in a similar manner to Reference Example 11(a) but using 3-(pyrid-4-yl)-3-(2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-propanoic acid methyl ester {Reference Example 5(b)} there was prepared 3-(pyrid-4-yl)-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)-propanoic acid methyl ester as a yellow oil. LC-MS: $R_T = 0.64$ minutes; MS(ES) = 297(M+H)⁺.

REFERENCE EXAMPLE 12

2-tert-Butyloxycarbonylamino-3-[2-trifluoroacetyl]-1,2,3,4-tetrahydro-isoquinolin-7-yl]-propionic acid methyl ester

- 10 A solution of 2-*tert*-butyloxycarbonylamino-3-[2-trifluoroacetyl]-1,2,3,4-tetrahydro-isoquinolin-7-yl]-acrylic acid methyl ester (2.8g, Reference Example 13) in industrial methylated spirits (200mL), under nitrogen, was treated with palladium on carbon (0.38g, 10%) and then hydrogen. After 2 hours an additional aliquot of palladium on carbon (0.4g, 10%) was added to the reaction mixture and this mixture was stirred at room temperature overnight and then filtered through Celite®. The filtrate was
15 evaporated to give the title compound (2.86g) as a colourless gum. LC-MS: 430(M)⁺.

REFERENCE EXAMPLE 13

2-tert-Butyloxycarbonylamino-3-[2-trifluoroacetyl]-1,2,3,4-tetrahydro-isoquinolin-7-yl]-acrylic acid methyl ester

- 20 A mixture of 7-bromo-[2-trifluoroacetyl]-1,2,3,4-tetrahydro-isoquinoline (4.29g, Reference Example 7), N-(*tert*-butyloxycarbonyl)-dehydroalanine methyl ester (3.64g, Reference Example 14), bis(*tri-o*-tolylphosphine)palladium (II) chloride 0.109g), *tri-o*-tolylphosphine (0.212g), triethylamine (7.7mL) and dimethylformamide (100mL), under nitrogen, was heated at 97°C for 4 hours. then left at room temperature for 3 days and then evaporated. The residual red-orange syrup was partitioned between
25 ethyl acetate and water. The organic phase was dried over sodium sulfate and evaporated. The residue was subjected to flash chromatography on silica eluting with a mixture of ethyl acetate and heptane (1:2, v/v) to give the title compound (2.9g) as a yellow solid.

REFERENCE EXAMPLE 14

N-(tert-Butyloxycarbonyl)-dehydroalanine methyl ester

A solution of N-(*tert*-butyloxycarbonyl)-DL-serine methyl ester {5.48g, Reference Example 15(a)} in tetrahydrofuran (60mL) was treated with triethylamine (3.48mL) followed by carbonyldiimidazole (4.05g). The mixture was stirred at room temperature overnight and then evaporated. The residue was extracted several times with *tert-butyl* methyl ether. The combined extracts were evaporated to give a

colourless oil which was subjected to flash chromatography on silica eluting with toluene to give the title compound (3.64g) as a colourless oil.

REFERENCE EXAMPLE 15

5 (a) N-(tert-Butyloxycarbonyl)-DL-serine methyl ester

A solution of DL-serine methyl ester hydrochloride (10g) in dry dimethylformamide (100mL) was treated with triethylamine (18mL) followed by a solution of di-tert-butyl dicarbonate (14g) in dry dimethylformamide (50mL). After stirring at room temperature overnight the reaction mixture was evaporated. The residue was dissolved in ethyl acetate and this solution was washed twice with water
10 and then evaporated. The resulting colourless oil (14g) was dissolved in ethyl acetate and this solution was dried over sodium sulfate and then evaporated. The residue was subjected to flash chromatography on silica eluting with a mixture of ethyl acetate and heptane (2:3, v/v) to give the title compound (6.55g) as a colourless oil.

15 (b) By proceeding in a similar manner to Reference Example 15(a) but using 3-(pyrid-4-yl)-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)-propionic acid methyl ester {Reference Example 11(b)} and carrying out the reaction in tetrahydrofuran there was prepared 3-[2-(tert-butyloxycarbonylamino)-1,2,3,4-tetrahydro-isoquinolin-7-yl]-3-(pyrid-4-yl)-propionic acid methyl ester as a red oil. LC-MS: $R_T = 2.61$ minutes, $MS(ES) = 397(M+H)^+$.

20

REFERENCE EXAMPLE 16

(a) 3-(Pyrid-4-yl)-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid ethyl ester, enantiomer A

A solution of 2-o-tolylaminobenzoxazole-6-acetic acid (0.124g) and diisopropylethylamine (0.17g) in
25 dimethylformamide (2mL) was treated with a solution of O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (0.183g) in dimethylformamide. After 10 minutes at room temperature this yellow solution was added dropwise to a rapidly stirred solution of 3-(1,2,3,4-tetrahydro-isoquinolin-7-yl)-3-(pyrid-4-yl)-propionic acid ethyl ester, enantiomer A {0.219mmoles, Reference Example 17(a)} in dimethylformamide. After stirring at room temperature for 2.5 hours the
30 reaction mixture was evaporated. The residual oil was dissolved in a mixture of acetonitrile, water and trifluoroacetic acid (25:75:0.1) and treated with dimethylformamide and trifluoroacetic acid to bring to pH2. This mixture was subjected to preparative HPLC {Hypersil Elite C18 column, 10cm x 2.1cm, using a mixture of acetonitrile, water and trifluoroacetic acid (25:75:0.1) as the mobile phase followed by a 1% acetonitrile/minute gradient) to give the title compound. LC-MS: $MS(ES) = 575(M+H)^+$,
35 $573(M-H)^-$.

(b) By proceeding in a similar manner to Reference Example 16(a) but using 3-(1,2,3,4-tetrahydro-isoquinolin-7-yl)-3-(pyrid-4-yl)-propionic acid ethyl ester, enantiomer B {Reference Example 17(b)} there was prepared 3-(pyrid-4-yl)-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-
5 isoquinolin-7-yl}-propanoic acid ethyl ester, enantiomer B. LC-MS:MS(ES) = 575(M+H)⁺, 573(M-H)⁻.

REFERENCE EXAMPLE 17

(a) 3-(1,2,3,4-Tetrahydro-isoquinolin-7-yl)-3-(pyrid-4-yl)-propionic acid ethyl ester, enantiomer A

10 A solution of 3-(1,2,3,4-tetrahydro-isoquinolin-7-yl)-3-(pyrid-4-yl)-propionic acid, enantiomer A {Reference Example 18(a)} in ethanol (50mL) was treated with concentrated hydrochloric acid (two drops) and then heated at reflux temperature, under nitrogen, for 8 hours. The pH of the reaction mixture was adjusted to ~5 by dropwise addition of sodium bicarbonate solution (5%) and this mixture was then evaporated to give the title compound which was used without further purification.

15 (b) By proceeding in a similar manner to Reference Example 17(a) but using 3-(1,2,3,4-tetrahydro-isoquinolin-7-yl)-3-(pyrid-4-yl)-propionic acid, enantiomer B {Reference Example 18(b)} there was prepared 3-(1,2,3,4-tetrahydro-isoquinolin-7-yl)-3-(pyrid-4-yl)-propionic acid ethyl ester, enantiomer
B which was used without further purification.

REFERENCE EXAMPLE 18

3-(1,2,3,4-Tetrahydro-isoquinolin-7-yl)-3-(pyrid-4-yl)-propionic acid, enantiomer A

25 A solution of 3-(1,2,3,4-tetrahydro-isoquinolin-7-yl)-3-(pyrid-4-yl)-propionic acid (S)-(-)- α -methylbenzyl amide, diastereoisomer A {0.092g, Reference Example 19(a)} in hydrochloric acid (10mL, 6N), under nitrogen, was heated at reflux temperature for 5 hours, then left at room temperature for 3 days, then heated at reflux temperature for 6.5 hours, then left at 100° overnight and then evaporated to give the title compound as a colourless solid. This material was used without further purification.

30 (b) By proceeding in a similar manner to Reference Example 16(a) but using 3-(1,2,3,4-tetrahydro-isoquinolin-7-yl)-3-(pyrid-4-yl)-propionic acid (S)-(-)- α -methylbenzyl amide, diastereoisomer B {Reference Example 19(b)} there was prepared 3-(1,2,3,4-tetrahydro-isoquinolin-7-yl)-3-(pyrid-4-yl)-propionic acid ethyl ester, enantiomer B.

REFERENCE EXAMPLE 19

(a) 3-(1,2,3,4-Tetrahydro-isoquinolin-7-yl)-3-(pyrid-4-yl)-propionic acid (S)-(-)- α -methylbenzyl amide, diastereoisomer A, and

(b) 3-(1,2,3,4-Tetrahydro-isoquinolin-7-yl)-3-(pyrid-4-yl)-propionic acid (S)-(-)- α -methylbenzyl amide, diastereoisomer B

A solution of 3-[2-(*tert*-butoxycarbonylamino)-1,2,3,4-tetrahydro-isoquinolin-7-yl]-3-(pyrid-4-yl)-propionic acid (0.6g, Reference Example 20) in dry dimethylformamide (15mL), under an argon atmosphere, was treated with di-isopropylethylamine (0.61g) then with O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (0.66g). After stirring at room temperature for a few minutes the mixture was treated with (S)-(-)- α -methylbenzylamine (0.19g) and stirring was continued at room temperature overnight. The reaction mixture was evaporated and the residue was partitioned between dichloromethane and aqueous sodium carbonate solution (2%). The organic phase was washed with water and then evaporated. The residue (0.952g) was subjected preparative reverse phase HPLC {Hypersil Elite C18 column, gradient elution using a mixture of 20% acetonitrile and 80% water containing 0.1% trifluoroacetic acid as the mobile phase with a 1% acetonitrile/minute gradient} followed by preparative HPLC {Dynamax silica column (30cm x 1.14cm), 60Å/8µm, using a mixture of methanol, 1,2-dichloroethane, n-heptane and triethylamine (1:650:350:0.25) with a flow rate of 15mL/minute} to give 3-(pyrid-4-yl)-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid ethyl ester, diastereoisomer A {106mg, Reference Example 19(a)} as the first to elute and 3-(pyrid-4-yl)-3-(((2-*o*-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl)-propanoic acid ethyl ester, diastereoisomer B {92mg, Reference Example 19(b)} as the second to elute.

REFERENCE EXAMPLE 20

3-[2-(*tert*-Butyloxycarbonylamino)-1,2,3,4-tetrahydro-isoquinolin-7-yl]-3-(pyrid-4-yl)-propionic acid

A solution of 3-[2-(*tert*-butoxycarbonylamino)-1,2,3,4-tetrahydro-isoquinolin-7-yl]-3-(pyrid-4-yl)-propionic acid methyl ester {1.46g, Reference Example 15(b)} in ethanol (100mL) was treated with sodium hydroxide solution (6mL, 1N). This mixture was heated at 80°C for 9 hours and then evaporated. The residue was partitioned between ethyl acetate and water which had been adjusted to pH 6 by addition of dilute hydrochloric acid. the aqueous phase was extracted twice with ethyl acetate and the combined organics were washed with water and then evaporated to give the title compound (0.6g) which was used without further purification.

IN VITRO AND IN VIVO TEST PROCEDURES

1. Inhibitory effects of compounds on VLA4 dependent cell adhesion to Fibronectin and VCAM.

1.1 Metabolic labelling of RAMOS cells.

RAMOS cells (a pre-B cell line from ECACC, Porton Down, UK) are cultured in RPMI culture
 5 medium (Gibco®, UK) supplemented with 5% foetal calf serum (FCS, Gibco®, UK). Prior to assay the
 cells are suspended at a concentration of 0.5×10^6 cells/ml RPMI and labelled with 400µCi/100mls of
 $[^3\text{H}]$ -methionine (Amersham®, UK) for 18 hours at 37°C.

1.2 96 well plate preparation for adhesion assay.

10 Cytostar plates (Amersham®, UK) were coated with 50µl/well of either 3µg/ml human soluble VCAM-1
 (R&D Systems Ltd., UK) or 28.8µg/ml human tissue Fibronectin (Sigma®, UK). In control non-specific
 binding wells 50µl phosphate buffered saline was added. The plates were then left to dry in an
 incubator at 25°C, overnight. The next day the plates were blocked with 200µl/well of Pucks buffer
 (Gibco®, UK) supplemented with 1% BSA (Sigma®, UK). The plates were left at room temperature in
 15 the dark for 2 hours. The blocking buffer was then disposed of and the plates dried by inverting the
 plate and gently tapping it on a paper tissue. 50µl/well of 3.6% dimethyl sulfoxide in Pucks buffer
 supplemented with 5mM manganese chloride (to activate the integrin receptor Sigma®, UK) and 0.2%
 BSA (Sigma®, UK), was added to the appropriate control test binding and non-specific binding assay
 wells in the plate. 50µl/well of the test compounds at the appropriate concentrations diluted in 3.6%
 20 dimethyl sulfoxide in Pucks buffer supplemented with 5mM manganese chloride and 0.2% BSA, was
 added to the test wells.

Metabolically labelled cells were suspended at 4×10^6 cells/ml in Pucks buffer that was supplemented
 with manganese chloride and BSA as above. 50µl/well of cells in 3.6% dimethyl sulfoxide in Pucks
 buffer and supplements was added to all plate wells.

25 The same procedure exists for plates coated with either VCAM-1 or fibronectin and data is determined
 for compound inhibition of cell binding to both substrates.

1.3 Performance of assay and data analysis.

The plates containing cells in control or compound test wells are incubated in the dark at room
 temperature for 1 hour.

30 The plates are then counted on a Wallac Microbeta scintillation counter (Wallac, UK) and the captured
 data processed in Microsoft® Excel (Microsoft®, US). The data was expressed as an IC_{50} , namely the
 concentration of inhibitor at which 50% of control binding occurs. The percentage binding is
 determined from the equation:

35
$$\{[(C_{\text{TB}} - C_{\text{NS}}) - (C_{\text{I}} - C_{\text{NS}})] / (C_{\text{TB}} - C_{\text{NS}})\} \times 100 = \% \text{ binding}$$

where C_{TB} are the counts bound to fibronectin (or VCAM-1) coated wells without inhibitor present, C_{NS} are the counts present in wells without substrate, and C_I are the counts present in wells containing a cell adhesion inhibitor.

5

Compound data of this invention is expressed for IC_{50} s for inhibition of cell adhesion to both fibronectin and VCAM-1. Particular compounds of the invention inhibit cell adhesion to fibronectin and VCAM-1 with IC_{50} 's in the range 100 micromolar to 77 nanomolar. Preferred compounds of the invention inhibit cell adhesion to fibronectin with IC_{50} 's below 100 nanomolar.

10

2. Inhibition of antigen-induced airway inflammation in the mouse and rat.

2.1 Sensitization of the animals.

Rats (Brown Norway, Harland Olac, UK) are sensitized on days 0, 12 and 21 with ovalbumin (100 μ g, intraperitoneally [i.p], Sigma, UK) administered with aluminium hydroxide adjuvant (100mg, i.p., Sigma, UK) in saline (1ml, i.p.).

In addition mice (C57) are sensitized on days 0 and 12 with ovalbumin (10 μ g, i.p.) administered with aluminium hydroxide adjuvant (20mg, i.p.) in saline (0.2ml, i.p.).

2.2 Antigen challenge.

Rats are challenged on any one day between days 28-38, while mice are challenged on any one day between days 20-30.

The animals are challenged by exposure for 30 minutes (rats) or 1 hour (mice) to an aerosol of ovalbumin (10g / l) generated by an ultrasonic nebulizer (deVilbiss Ultraneb, US) and passed into an exposure chamber.

2.3 Treatment protocols.

Animals are treated as required before or after antigen challenge. The aqueous-soluble compounds of this invention can be prepared in water (for oral, p.o. dosing) or saline (for intratracheal, i.t. dosing).

Non-soluble compounds are prepared as suspensions by grinding and sonicating the solid in 0.5 % methyl cellulose / 0.2 % polysorbate 80 in water (for p.o. dosing, both Merck UK Ltd., UK) or saline (for i.t. dosing). Dose volumes are: for rats 1ml / kg, p.o. or 0.5ml / kg, i.t.; for mice 10ml / kg, p.o. or 1ml / kg, i.t.

2.4 Assessment of airway inflammation.

35

The cell accumulation in the lung is assessed 24 hours after challenge (rats) or 48-72 hours after challenge (mice). The animals are euthanized with sodium pentobarbitone (200mg/kg, i.p., Pasteur Merieux, France) and the trachea is immediately cannulated. Cells are recovered from the airway lumen by bronchoalveolar lavage (BAL) and from the lung tissue by enzymatic (collagenase, Sigma®, UK) disaggregation as follows.

BAL is performed by flushing the airways with 2 aliquots (each 10 ml/kg) RPMI 1640 medium (Gibco®, UK) containing 10 % foetal calf serum (FCS, Serotec Ltd., UK). The recovered BAL aliquots are pooled and cell counts made as described below.

Immediately after BAL, the lung vasculature is flushed with RPMI 1640 / FCS to remove the blood pool of cells. The lung lobes are removed and cut into 0.5 mm pieces. Samples (rats: 400mg; mice: 150mg) of homogenous lung tissue are incubated in RPMI 1640 / FCS with collagenase (20 U/ml for 2 hours, then 60 U/ml for 1 hour, 37°C) to disaggregate cells from the tissue. Recovered cells are washed in RPMI 1640 / FCS.

Counts of total leukocytes recovered from the airway lumen and the lung tissue are made with an automated cell counter (Cobas Argos, US). Differential counts of eosinophils, neutrophils and mononuclear cells are made by light microscopy of cytocentrifuge preparations stained with Wright-Giemsa stain (Sigma®, UK). T cells are counted by flow cytometry (EPICS XL, Coulter® Electronics, US) using fluophore-labelled antibodies against CD2 (a pan-T cell marker used to quantify total T cells), CD4, CD8 and CD25 (a marker of activated T cells). All antibodies were supplied by Serotec Ltd., UK).

2.5 Data analysis.


The cell data was expressed as mean cell numbers in unchallenged, challenged and vehicle treated, and challenged and compound treated groups, including the standard error of the means. Statistical analysis of the difference among treatment groups was evaluated using one-way analysis of variance via the Mann-Whitney test. Where $p < 0.05$, no statistical significance existed.

- 62 -

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A compound of formula (I):

5



(I)

wherein:

10 R^1 represents optionally substituted phenyl, $R^3\text{NH-Ar}^1\text{-L}^2\text{-}$ or $R^3\text{-NH-C(=O)-NH-Ar}^2\text{-L}^2\text{-}$;

R^2 represents hydrogen, halogen, C_{1-4} alkyl or C_{1-4} alkoxy;

R^3 represents optionally substituted aryl or optionally substituted heteroaryl;

15 L^1 represents a C_{2-15} alkenylene, C_{1-15} alkylene or C_{2-15} alkynylene linkage each optionally substituted by (a) carboxy, hydroxy, mercapto, cyano, oxo, $\text{S(O)}_m\text{R}^4$, R^5 , -C(=O)-R^5 , -C(=O)-OR^5 , $\text{-N(R}^6\text{)-C(=O)-R}^4$, $\text{-N(R}^6\text{)-C(=O)-OR}^4$, $\text{N(R}^6\text{)-SO}_2\text{-R}^4$, NY^3Y^4 or $\text{-[C(=O)-N(R}^7\text{)-C(R}^8\text{)(R}^9\text{)]}_p\text{-C(=O)-NY}^3\text{Y}^4$; or by (b) C_{1-15} alkyl substituted by carboxy, hydroxy, mercapto, imidazolyl, $\text{S(O)}_m\text{R}^4$, $\text{-C(=O)-NY}^3\text{Y}^4$ or $\text{-NY}^3\text{Y}^4$;

20 R^4 is C_{1-15} alkyl, aryl, C_{3-10} cycloalkyl, heteroaryl or hetero- C_{3-10} -cycloalkyl, or C_{1-15} alkyl substituted by aryl, an acidic functional group, selected from carboxyl, carboxyl bioisosteres, hydroxy, mercapto and imidazole C_{3-10} cycloalkyl, heteroaryl, hetero- C_{3-10} -cycloalkyl, $\text{-S(O)}_m\text{R}^5$, $\text{-C(=O) NY}^3\text{Y}^4$ or $\text{-NY}^3\text{Y}^4$;

25 R^5 represents C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, aryl, aryl- C_{1-15} -alkyl, aryl- C_{2-15} -alkenyl, aryl- C_{2-15} -alkynyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkyl- C_{1-15} -alkyl, C_{3-10} cycloalkyl- C_{2-15} -alkenyl, C_{3-10} cycloalkyl- C_{2-15} -alkynyl, C_{3-10} cycloalkenyl, C_{3-10} cycloalkenyl- C_{1-15} -alkyl, heteroaryl, heteroaryl- C_{1-15} -alkyl, heteroaryl- C_{2-15} -alkenyl, heteroaryl- C_{2-15} -alkynyl, hetero- C_{3-10} -cycloalkyl or hetero- C_{3-10} -cycloalkyl- C_{1-15} -alkyl;

- 63 -

R^6 is hydrogen, C_{1-15} alkyl, aryl, aryl- C_{1-15} -alkyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkyl- C_{1-15} -alkyl, heteroaryl, heteroaryl- C_{1-15} -alkyl, hetero- C_{3-10} -cycloalkyl or hetero- C_{3-10} -cycloalkyl- C_{1-15} -alkyl;

R^7 is hydrogen, R^5 or C_{1-15} alkyl substituted with C_{1-15} alkoxy, C_{3-10} cycloalkyl, hydroxy, mercapto, C_{1-15} alkylthio, or $-NY^3Y^4$;

R^8 is hydrogen or C_{1-4} alkyl;

R^9 is selected from hydrogen or a group consisting of amino acid side chains, an acidic functional group, R^5 , $-C(=O)-R^5$, or $-C(=O)-NY^3Y^4$, or C_{1-15} alkyl substituted by an acidic functional group or by R^5 , $-NY^3Y^4$, $-NH-C(=O)-R^5$, $-C(=O)-R^{12}-NH_2$, $-C(=O)-Ar^2-NH_2$, $-C(=O)-R^{12}-CO_2H$, or $-C(=O)-NY^3Y^4$;

R^{12} is a C_{1-15} alkylene chain, a C_{2-15} alkenylene chain, or a C_{2-15} alkynylene chain;

R^{13} is C_{1-15} alkyl, aryl, aryl- C_{1-15} -alkyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkyl- C_{1-15} -alkyl, heteroaryl, heteroaryl- C_{1-15} -alkyl, hetero- C_{3-10} -cycloalkyl or hetero- C_{3-10} -cycloalkyl- C_{1-15} -alkyl;

Ar^1 represents a saturated, partially saturated or fully unsaturated 8- to 10-membered bicyclic ring system containing at least one heteroatom selected from O, S or N, optionally substituted by one or more aryl group substituents;

Ar^2 represents optionally substituted arylidiyl or heteroaryldiyl;

L^2 represents a C_{1-15} alkylene chain;

Y^1 and Y^2 are independently hydrogen, C_{2-15} alkenyl, C_{1-15} alkyl, aryl, aryl- C_{1-15} -alkyl, C_{3-10} cycloalkyl, heteroaryl or heteroaryl- C_{1-15} -alkyl;

Y^3 and Y^4 are independently hydrogen, C_{2-15} alkenyl, C_{1-15} alkyl, C_{2-15} alkynyl, aryl, C_{3-10} cycloalkenyl, C_{3-10} cycloalkyl, heteroaryl, hetero- C_{3-10} -cycloalkyl, or C_{1-15} alkyl substituted by C_{1-15} alkoxy, aryl, cyano, C_{3-10} cycloalkyl, heteroaryl, hetero- C_{3-10} -cycloalkyl, hydroxy, oxo, $-NY^1Y^2$, or one or more $-CO_2R^6$ or $-C(=O)-NY^1Y^2$ groups;

Y is carboxy;

m is an integer 1 or 2;

p is zero or an integer 1 to 4;

q is an integer 1 to 4; and

- 64 -

wherein (i) "aryl" as a group or part of a group denotes an optionally substituted monocyclic or multicyclic aromatic carbocyclic moiety of 6 to 14 carbon atoms or an optionally substituted partially saturated multicyclic aromatic carbocyclic moiety in which an aryl and a cycloalkyl or cycloalkenyl group are fused together to form a cyclic structure;

(ii) "heteroaryl" as a group or part of a group denotes an optionally substituted aromatic monocyclic or multicyclic organic moiety of 5 to 10 ring members in which one or more of the ring members is/are element(s) selected from oxygen, sulfur and nitrogen, or an optionally substituted partially saturated multicyclic heterocarbo-cyclic moiety in which a heteroaryl and a cycloalkyl or cycloalkenyl group are fused together to form a cyclic structure; and

(iii) aryl group substituents optionally present on such an aryl or heteroaryl group are selected from the group consisting of H-C(=O)-, C₁₋₁₅alkylC(=O)-, H-C(=O)-NH-, C₁₋₁₅alkylC(=O)-NH-, C₁₋₁₅alkoxy, C₁₋₁₅alkoxycarbonyl, C₁₋₁₅alkylenedioxy, C₁₋₁₅alkylsulfinyl, C₁₋₁₅alkylsulfonyl, C₁₋₁₅alkylthio, aroyl, aroylamino, aryl, aryl-C₁₋₁₅-alkyloxy, aryl-C₁₋₁₅-alkyloxycarbonyl, aryl-C₁₋₁₅-alkylthio, aryloxy, aryloxycarbonyl, arylsulfinyl, arylsulfonyl, arylthio, carboxy, cyano, halo, heteroaroyl, heteroaryl, heteroaryl-C₁₋₁₅-alkyloxy, heteroaroylamino, heteroaryloxy, hydroxy, nitro, trifluoromethyl, Y¹Y²N-, Y¹Y²NCO-, Y¹Y²NSO₂-, Y¹Y²N-C₂₋₆alkylene-Z²-, C₁₋₁₅alkylC(=O)-Y¹N-, C₁₋₁₅alkyl-SO₂-Y¹N- or C₁₋₁₅alkyl optionally substituted with aryl, heteroaryl, hydroxy, or Y¹Y²N-,

where Z² is O, NR⁸ or S(O)_n and n is 0, 1 or 2.

but excluding compounds where an oxygen, nitrogen or sulfur atom is attached directly to a carbon-carbon multiple bond of a C₂₋₁₅alkenylene or C₂₋₁₅alkynylene residue;


or the pharmaceutically acceptable salts or solvates thereof, or the N-oxides and ester prodrugs of the compounds of formula (I) or pharmaceutically acceptable salts or solvates thereof.


30

2. A compound according to claim 1 in which R¹ represents a group R³-NH-Ar¹-L²- in which:

- 65 -

L^2 is a straight or branched C_{1-6} alkylene chain;

Ar^1 is an 8- to 10-membered bicyclic system  in which (i) ring  is


a 5- or 6-membered optionally substituted heterocycle, (ii) ring  is a 5- or

6-membered optionally substituted heterocycle or an optionally substituted


5 benzene ring, and (iii) the two rings are joined together by a carbon-carbon linkage or a carbon-nitrogen linkage; and


R^3 is an optionally substituted aryl.

3. A compound according to claim 2 in which  is a 5-membered

optionally substituted heterocycle, ring  is an optionally substituted

10 benzene ring, and the two rings are joined together by a carbon-carbon linkage.

4. A compound according to claim 2 in which  is an optionally substituted benzoxazolyl or an optionally substituted benzimidazolyl, each in which the benzene ring contains the optional substituents.

15 5. A compound according to claim 2 in which ring  is a benzene ring optionally substituted by one of C_{1-4} alkyl, C_{1-4} alkoxy, amino, halogen, hydroxy, C_{1-4} alkylthio, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, nitro or trifluoromethyl.

20 6. A compound according to any one of claims 2 to 5 in which R^3 represents a 2-substituted phenyl.

7. A compound according to claim 6 in which R^3 represents 2-methylphenyl.

25 8. A compound according to claim 1 in which R^1 represents a group R^3 $NH-C(=O)NH-Ar^2-L^2$ - in which:

L^2 is a straight or branched C_{1-6} alkylene chain;

Ar^2 is an optionally substituted phenylene or an optionally substituted

- 66 -

heteroaryldiyl; and

R^3 is an optionally substituted aryl group or an optionally substituted heteroaryl group.

5 9. A compound according to claim 8 in which Ar^2 is an optionally substituted m- or p-phenylene.

10. A compound according to claim 9 in which Ar^2 is a 3-substituted p-phenylene in which the substituent is ortho to the $R^3-NH-C(=O)-NH-$ group.

10

11. A compound according to claim 8 in which Ar^2 is an optionally substituted azaheteroaryldiyl.


12. A compound according to claim 11 in which Ar^2 is pyridine-2,5-diyl, in
15 which the $R^3-NH-C(=O)-NH-$ group is adjacent to the pyridyl nitrogen atom, and which is substituted in the 4- or 6-position with a methyl or methoxy group.

13. A compound according to any one of claims 8 to 12 in which R^3 is 2- or
20 3-methyl(or methoxy)phenyl.

14. A compound according to any one of claims 8 to 12 in which R^3 is an optionally substituted pyridyl.

25 15. A compound according to claim 14 in which R^3 is 3-methyl-2-pyridyl.

16. A compound according to claim 1 of formula (Ia):



- 67 -

in which R^2 , R^3 , L^1 , L^2 and Y are as defined in any one of claims 1 to 15, X is
 O or NR^{18} , where R^{18} is hydrogen or C_{1-4} alkyl, and R^{17} is hydrogen or an aryl
 group substituent, or the pharmaceutically acceptable salts and solvates
 5 thereof or the N-oxides and ester prodrugs of the compounds of formula (Ia)
 or pharmaceutically acceptable salts and solvates thereof.


17. A compound according to claim 16 in which R^3 represents an optionally
 substituted aryl.

10

18. A compound according to claim 16 or 17 in which R^{17} represents
 hydrogen, halo, C_{1-4} alkyl, or C_{1-4} alkoxy.

19. A compound according to any one of claims 16 to 18 in which L^2
 15 represents a straight or branched C_{1-6} alkylene chain.

20. A compound according to any one of claims 16 to 19 in which the


group  is attached at the ring 6 position or at the ring 5
 or 6 position when X is NR^{18} and R^{18} is C_{1-4} alkyl.

20

21. A compound according to any one of claims 16 to 20 in which the
 group $-L^1-Y$ is attached at position 6 or 7 of the tetrahydroisoquinoline ring.

22. A compound according to claim 1 of formula (Ib):

25



- 68 -

in which R^2 , R^3 , Ar^2 , L^1 , L^2 and Y are as defined in any one of claims 1 to 21, and the corresponding N-oxides and ester prodrugs thereof, and the pharmaceutically acceptable salts and solvates of such compounds, and the N-oxides and ester prodrugs thereof.

5

23. A compound according to claim 22 in which R^3 represents 2-methylphenyl or 3-methyl-2pyridyl.

24. A compound according to any one of claims 22 to 23 in which Ar^2 represents an optionally substituted phenylene or optionally substituted heteroaryldiyl.

25. A compound according to claim 24 in which Ar^2 is p-phenylene or phenylene substituted in the 3-position by halo, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} alkylthio, C_{1-4} alkylsulfinyl or C_{1-4} alkylsulfonyl.

26. A compound according to claim 24 in which Ar^2 is pyridine-2,5-diyl, in which the R^3 -NH-C(=O)-NH- group is adjacent to the pyridyl nitrogen atom, and which is substituted in the 4- or 6-position with a methyl or methoxy group.

27. A compound according to any one of claims 22 to 26 in which L^2 represents a straight or branched C_{1-6} alkylene chain.


28. A compound according to any one of claims 1 to 27 in which L^1 represents a C_{1-4} alkylene linkage optionally substituted by C_{1-4} alkyl, aryl, heteroaryl, $-N(R^6)-C(=O)-R^4$, $-N(R^6)C(=O)-OR^4$, $-N(R^6)-SO_2-R^4$, $-NY^3Y^4$ or $-[C(=O)-N(R^7)-C(R^8)(R^9)]_p-C(=O)-NY^3Y^4$, or C_{1-15} alkyl substituted by carboxy, hydroxy, mercapto, imidazolyl, $-C(=O)-NY^3Y^4$ or $-NY^3Y^4$.

30

29. A compound according to claim 28 in which the C_{1-4} alkylene linkage is an ethylene linkage.


- 69 -

30. A compound according to claim 29 in which L^1 represents a group




, where R^{14} is hydrogen or C_{1-4} alkyl and R^{15} represents hydrogen or C_{1-4} alkyl; or where R^{14} is hydrogen and R^{15} represents aryl, heteroaryl,
 5 $-N(R^6)-C(=O)-R^4$, $-N(R^6)-C(=O)-OR^4$, $-N(R^6)SO_2-R^4$, $-NY^3Y^4$ or $-[C(=O)-N(R^7)-C(R^8)(R^9)]_p-C(=O)-NY^3Y^4$, or C_{1-15} alkyl substituted by carboxy, hydroxy, mercapto, imidazolyl, $-C(=O)-NY^3Y^4$ or $-NY^3Y^4$.

31. A compound according to claim 29 in which L^1 represents a group



10 , where R^{14} is hydrogen or C_{1-4} alkyl and R^{16} represents C_{1-4} alkyl; or where R^{14} is hydrogen and R^{16} represents aryl, heteroaryl, $-N(R^6)-C(=O)-R^4$, $-N(R^6)-C(=O)-OR^4$, $-N(R^6)-SO_2-R^4$, $-NY^3Y^4$ or $-[C(=O)-N(R^7)-C(R^8)(R^9)]_p-C(=O)-NY^3Y^4$, or C_{1-15} alkyl substituted by carboxy, hydroxy, mercapto, imidazolyl, $-C(=O)-NY^3Y^4$ or $-NY^3Y^4$.

15 32. A compound according to claim 31 in which L^1 is a group



33. A compound according to claim 32 where R^{16} represents $N(R^6)-C(=O)-R^4$, or $-N(R^6)-SO_2-R^4$.

20 34. A compound according to claim 1 selected from:

3-(((4-methyl-2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-butanoic acid;

3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7yl)-butanoic acid;

25 3-phenyl-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl)-propanoic acid;

- 70 -

3-cyclohexyl-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl}-propanoic acid;

3-(pyrid-4-yl)-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl}-propanoic acid;

5 3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-8yl}-but-2-enoic acid;

3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-8yl)-butanoic acid;

10 3-{2-[(2-o-tolylamino-benzoxazol-6-yl)-carbonyl]-1,2,3,4-tetrahydro-isoquinolin-6-yl}-butanoic acid;

{5-(3-methoxy-4-[3-(2-methylphenyl)ureido]-phenylacetyl-amino)-1,2,3,4-tetrahydro-isoquinolin-8-yl}-butanoic acid;

3-phenyl-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydroisoquinolin-7-yl}-propanoic acid;

15 3-{(2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-6yl}-butanoic acid;

3-(pyrid-4-yl)-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl}-propanoic acid, enantiomer R;

20 3-(pyrid-4-yl)-3-(((2-o-tolylamino-benzoxazol-6-yl)-acetyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl}-propanoic acid, enantiomer S;

or the pharmaceutically acceptable salts or solvates thereof or the N-oxides and ester prodrugs of these compounds or the pharmaceutically acceptable salts thereof.

25 35. A pharmaceutical composition comprising a compound according to any one of claims 1-34 or a corresponding N-oxide or ester prodrug thereof, or a pharmaceutically acceptable salt or solvate of such a compound, or an N-oxide or ester prodrug thereof, in association with a pharmaceutically acceptable carrier or excipient.

30

36. A compound or composition according to claim 1 or 35, respectively, for use in the treatment of inflammatory diseases.

- 71 -

37. A compound or composition according to claim 1 or 35, respectively, for use in the treatment of asthma.

