



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 7/00, A61K 39/12	A1	(11) International Publication Number: WO 94/18311 (43) International Publication Date: 18 August 1994 (18.08.94)
(21) International Application Number: PCT/US94/00951 (22) International Filing Date: 26 January 1994 (26.01.94) (30) Priority Data: 08/014,915 8 February 1993 (08.02.93) US (71) Applicants: MILES INC. [US/US]; One Mellon Center, 500 Grant Street, Pittsburgh, PA 15219-2502 (US). IOWA STATE UNIVERSITY RESEARCH FOUNDATION INC. [US/US]; 214 O & L, Ames, IA 50011 (US). (72) Inventors: SANDERSON, Thomas; 4901 Searle Parkway, Skokie, IL 60077 (US). MCGINLEY, Michael, J.; 7703 Monrovia Street, Lenexa, KS 66216 (US). ZIMMERMAN, Jeffrey, J.; R.R. #3, Ames, IA 50010 (US). HILL, Howard, T.; Route 1, Box 6, Cambridge, IA 50046 (US). MEETZ, Michael, C.; R.R. #2, Box 24, Nevada, IA 50501 (US). PIRTLE, Eugene, C.; 810 Brookridge Avenue, Ames, IA 50010 (US). SWENSON, Sabrina, L.; 11000 Northwest 166th Avenue, Madrid, IA 50158 (US). SHIBLEY, George, P.; 4149 W. 128 Terrace, Leawood, KS 66209 (US). (74) Agent: MUELLER, Gerhard; Bayer Aktiengesellschaft, Konzernverwaltung RP, Patentabteilung, D-51368 Leverkusen (DE).		(81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FI, HU, JP, KR, KZ, LK, NO, NZ, PL, RO, RU, SK, UA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PROCESS FOR GROWING PORCINE REPRODUCTIVE AND RESPIRATORY SYNDROME VIRUS AND ITS USE IN VACCINES		
(57) Abstract A process for growing a PRRSV in a tissue culture which is susceptible to infection to replicate the virus to an amount sufficient to protect animals against PRRS or be used in diagnosing PRRS or identifying the molecular structure of PRRSV for development of recombinant products, comprising inoculating the virus onto the tissue culture and harvesting the replicated virus.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

-1-

PROCESS FOR GROWING PORCINE REPRODUCTIVE AND RESPIRATORY SYNDROME VIRUS AND ITS USE IN VACCINES.

BACKGROUND OF THE INVENTION

Field of the Invention: The present invention relates to a porcine reproductive and respiratory syndrome virus (PRRSV) isolate. More specifically, the present invention relates to diagnostic and protective antigens and vaccine for porcine reproductive and respiratory disease of pigs, and the methods of making and using the same.

Brief Description of the Prior Art: Porcine reproductive and respiratory syndrome is rapidly emerging as an economically devastating disease problem for U.S. and European swine producers. The syndrome was first described in the U.S. in 1987, with similar descriptions surfacing in Europe and Canada, two to three years later. The disease syndrome has been referred to by many different names including mystery swine disease, abortus blau, blue eared pig disease, porcine epidemic abortion and respiratory syndrome (PEARS), swine infertility and respiratory syndrome (SIRS). The name porcine reproductive and respiratory syndrome (PRRS) is employed hereinafter.

The etiological agent capable of reproducing the disease syndrome has been identified as a small enveloped spherical RNA virus, with an average virion diameter of 62 nm and a 25-30 nm core surrounded by an envelope. This virus has been tentatively grouped as a member of the genus Arterivirus within the Togaviridae family. The porcine reproductive and respiratory syndrome virus (PRRSV) isolate, described herein, fits this tentative classification and has been shown to reproduce the disease syndrome.

-2-

The disease syndrome associated with PRRSV is characterized by acute and chronic reproductive failure in adult female swine and severe to mild respiratory disease in young pigs. Reproductive failure is characterized by late term abortions
5 resulting in increased incidence of mummified, stillborn and weak born pigs with markedly reduced chances for survival. Chronic problems with delayed return to estrus in infected sows has also been described. Respiratory disease sequela range from marked fever and interstitial pneumonitis to mild upper
10 respiratory signs (i.e. sneezing, coughing, and nasal or ocular discharge) in young pigs. Recent (1990) serological and herd history studies indicate that at least 50% of U.S. swine herds have been exposed to PRRSV or have experienced reproductive failure and respiratory disease indicative of PRRSV infection.

15 Due to the recent emergence of this disease syndrome, studies on the pathogenesis, epidemiology and control of disease have been limited. As would be realized, an efficient propagation and processing of the antigens comprising PRRSV, will facilitate the development of a PRRSV vaccine as an aid in
20 the prevention and control of porcine reproductive and respiratory syndrome.

SUMMARY OF THE INVENTION

In accordance with the foregoing, the present invention encompasses a virus isolate designated as ISU-P which is useful
25 in the preparation of antigens for the diagnosis of porcine reproductive and respiratory syndrome and the induction of a protective immune response to porcine reproductive and respiratory syndrome virus (PRRSV). Also encompassed by the invention are the processes for making and using the antigens.

30 In a present embodiment of the invention, PRRSV isolate can be obtained by a process comprising isolating the PRRSV by co-cultivation of 10% weight/volume (w/v) lung homogenates from

-3-

virus infected pigs with primary porcine alveolar macrophage or continuous cell line cultures at 35°C.

5 The present invention further encompasses propagation of the virus to high titer in certain cell lines such as an African Green Monkey Kidney continuous cell line (MA 104) and a unique cloned derivative of the same (9009B).

Also encompassed by the invention are processes for preparing and using live virus or killed virus antigens (conventional or recombinant) and the vaccines resulting therefrom by combining an immunologically effective amount of
10 the virus with diluent and/or adjuvant, respectively.

It has been found that direct experimental immunization of sows with live virus or contact with live virus immunized sows prevented reproductive failure following an intranasal chal-
15 lenge with virulent PRRSV.

It has also been found that inactivated, adjuvanted PRRSV vaccines protected pigs from PRRS infection post challenge as measured by lack of virus replication in vaccinates as compared with non-vaccinated control pigs.

20 DETAILED DESCRIPTION OF THE INVENTION

As set forth above, the present invention is directed to the isolation of the PRRSV virus isolate ISU-P for the purpose of obtaining viral antigens for use in vaccines and diagnostic assays. The ISU-P isolate was obtained from the lung tissue of
25 a weak born pig. Illustratively, a 10% w/v tissue homogenate from lung, spleen, and lymph node was prepared in Minimal Essential Media supplemented with antibiotics (MEM+A) to minimize the potential for contamination. The crude homogenate was clarified by centrifugation at 4°C for 15 minutes at
30 1500 x g. Clarified supernatant was diluted 1:5 in MEM+A and 1.0mL was adsorbed onto a 48 hour old confluent monolayer of MA 104 cells or primary alveolar macrophages in 25 cm² flasks. Following a 2 hour adsorption period at 35°C monolayers were washed twice

with MEM+A, and 5.0 mL of MEM+A supplemented with 5.0% gamma-irradiated fetal calf serum was added. Virus isolation flasks were incubated at 35°C in a 5% CO₂ atmosphere and observed daily for evidence of cytopathic effect (CPE). The
5 PRRSV ISU-P cytopathic effect began with the rounding of small foci of 5 to 10 cells and progressed as many cells became pyknotic and detached from the substrate over the course of 4 to 7 days post infection.

The ISU-P isolate was confirmed as a PRRSV isolate by
10 fluorescent antibody staining with PRRSV group specific monoclonal antibody, designated in the public domain as SDOW17. In addition, negative stained virus infected cells were examined by transmission electron microscopy for the presence of virus particles. The virus particles observed were
15 spherical and enveloped with an average virion diameter of 65 nm and core diameter of 27 nm. Tissue culture virus characterized in this manner was used to experimentally reproduce late term abortion and reproductive failure in pregnant sows (Table 1). Table 1 shows that pregnant gilts
20 exposed to either ISU-P-infected lung homogenates (PRRS-hm) or ISU-P grown in a cell culture (PRRS-tc) demonstrate typical disease signs of porcine reproductive and respiratory syndrome (PRRS). In the PRRS-hm groups there were no normal pigs farrowed. In the PRRS-tc groups only 4 of 33 pigs were
25 farrowed normal. In comparison, 29 of 30 (96.7%) of the non-exposed pigs were normal at farrowing. The fact that ISU-P, even after purification in tissue culture, can reproduce the disease of PRRS confirms that this isolate is correctly classified as a member of the genus Arterivirus within the
30 Togaviridae family. This isolate has also been found to produce a mild interstitial pneumonia in experimentally infected pigs.

It is a feature of the invention that the PRRSV in accordance with the invention can be grown in certain tissue culture cells which effectively replicate the virus to a level

sufficient enough to provide the antigenic mass necessary to be incorporated in vaccines which can protect pigs and/or pregnant gilts or sows from PRRS challenge. Illustratively, the PRRSV can be grown in a tissue culture cell designated MA 104 by Whittaker BioProducts. Virus propagation experiments, employing identical culture conditions, with MA 104 cells obtained from three different sources indicate that a high degree of variability exists among MA 104 cells in terms of the ability to support PRRSV replication to high titer. One MA 104 cell culture designated MA 104(M) by Miles Inc., Agriculture Division, Animal Health Products Group, and used in the studies described herein, consistently supported PRRSV replication to high titer. Furthermore, a cell line clone, generated at Miles Inc., Agriculture Division, Animal Health Products Group, which derived from MA 104(M), designated 9009B, supports replication of PRRSV to titers higher than the parental cell line. Therefore, the preferred tissue culture cell for growth of ISU-P is 9009B. The following more fully illustrates cell culture propagation of the PRRSV ISU-P.

Tissue culture cells [MA 104 (M) and 9009B] were grown and maintained in Dulbecco's modified minimum essential media, high glucose (DMEM/HG), supplemented with 5% fetal calf serum and buffered with 1.3g/L sodium bicarbonate. Cell passages were made with sufficient cell counts to attain a 80-100% confluent monolayer in 48 hours. PRRSV ISU-P virus stocks were diluted in DMEM/HG + 5% fetal calf serum buffered to pH 6.8 with PIPES. Virus was inoculated onto the cell sheet at a low multiplicity of infection (MOI) and incubated at 36°C. Infected cultures were observed daily for CPE for 4-7 days. Infected cultures were harvested when 90-100% CPE was evident. Maximum virus antigenic mass was obtained after one -70°C freeze/thaw cycle. Mean virus titers resulting from propagation in MA 104(M) cells ranged from $10^{3.0}$ to $10^{4.5}$ fluorescent antibody infectious dose 50 /mL (FAID 50 /mL). Virus titers achieved by propagation in the 9009B cell clone ranges from $10^{5.5}$ to $10^{6.5}$ FAID 50 /mL.

-6-

Titers of this magnitude make it possible to incorporate sufficient antigenic mass into commercial scale PRRSV vaccine formulations of greater than 50L in volume.

As would be realized by those skilled in the art, once the PRRSV can be propagated to high levels of antigenic mass, derivatives of this virus including subunits thereof which are effective in accordance with the invention, can be obtained by means known to the art such as extraction from the virus. Additionally, the protective antigens can be identified at the molecular level and reproduced and expressed using recombinant technology.

Viral fluids for vaccine formulation were inactivated by two methods. The inactivants chosen were formalin and binary ethylenimine (BEI). Inactivation was performed by standard methods described more fully herein. Formalin inactivation was accomplished by mixing viral fluids with stock 37% formaldehyde to a final formalin concentration of 0.05%. The formalin-virus mixture was held at room temperature (approximately 25°C) with constant mixing for 24 hours. Samples were taken at times 0, 8, 12 and 24 hours and assayed for live virus in 9009B cells. Cytopathic effect and fluorescent staining with group specific monoclonal antibody detected live PRRSV only at time = 0 hours.

Inactivation with BEI was accomplished by combining stock 0.1 M BEI (20.5 g/L 2-bromo-ethylamine HBR in 0.175 N NaOH) with viral fluids to a final concentration of 1.0 mM BEI. Inactivation was performed by holding the BEI-virus mixture at room temperature with constant mixing for 48 hours. Virus inactivation was halted by the addition of 1.0 M sodium thiosulfate to a final concentration of 0.1 mM with mixing for 2 hours. Samples were taken at times 0, 2, 4, 8, 12, 24 and 48 hours and assayed for live virus as described above. Live PRRSV was detected only at times = 0, 2 and 4 hours.

In the preparation of a modified live or attenuated vaccine of the invention, the PRRSV is altered so that it can still infect the host in a limited manner but cannot cause

-7-

signs of disease in the host. Normally, such alteration of the virus takes place by passaging it through a non-host cell such as African Green Monkey Kidney cells. Initially, the virus may not grow well or even not at all. In the latter case, it may be necessary to force the virus to adapt to growth in cell culture. This can be done by adding the virus to the cell culture at a high or low multiplicity of infection (MOI). Alternately, the virus can be force-passaged or blind passaged along with the cells until enough cytopathic effect is observed, thus indicating that virus is being adapted. As adaptation occurs, the virus titer will increase as is observed when the PRRSV isolate ISU-P is passaged from the MA 104(M) cell line to the 9009B cell line. Table 1 shows that the tissue culture passaged PRRSV ISU-P (PRRS-tc) may be less virulent than that isolated from lung homogenates, a typical result of adaptation. Data disclosed in Table 2 indicate that the method of making vaccines in accordance with this invention are especially effective in conferring protection against PRRS.

The inactivated PRRSV ISU-P was adjuvanted as follows. Inactivated fluids were adjuvanted with adjuvants selected from the group consisting of Freund's complete adjuvant (FCA), Freund's incomplete adjuvant (FIA), a carbopol based adjuvant, and Diamond Scientific Adjuvant B (Adj B). These adjuvants represent the major types used in vaccine manufacture. Oil-based adjuvants are represented by FCA, FIA and ADJ B. Aqueous-based adjuvants are represented by carbopol. The inactivated PRRSV ISU-P may optionally contain a suitable preservative.

Vaccine formulations with FCA and FIA were prepared by emulsifying the inactivated virus, typically in equal volumes of inactivated virus fluid and adjuvant using an 18 guage double luer lock apparatus and two syringes. The fluids were expressed repeatedly through the assembly between syringes until a thickened semi-solid emulsion was formed. The

emulsions were loaded directly into syringes for administration.

Vaccine formulations with Adj B were prepared by mixing stock Adj B with an equal volume of inactivated PRRSV ISU-P viral fluid. Adjuvant and virus was mixed by stirring at room temperature for 1 hour. Adjuvanted vaccine was aseptically transferred to sterile multidose vials and stored at 4°C prior to administration.

Carbopol-based vaccine formulations were prepared by mixing stock carbopol adjuvant with inactivated viral fluids to a final adjuvant concentration of 10% v/v. Adjuvant and virus was mixed by stirring at room temperature for 4 hours. Adjuvanted vaccine was aseptically transferred to sterile multidose vials and stored at 4°C prior to administration.

PRRSV ISU-P virus at a preinactivation titer as low as $10^{5.0}$ FAID₅₀/mL was used in the various vaccine formulations described.

In order to show that PRRSV isolate ISU-P could protect against porcine reproduction and respiratory syndrome, two vaccination/challenge studies were conducted. In the first study, 4 gilts were immunized with live PRRSV ISU-P by oronasal exposure to virus via either a direct route or contact with diseased pigs. Two gilts of the same age were left unimmunized (unexposed) and were housed separately to serve as controls. The direct-exposure vaccinates (#57 and #58) received 3.0 mL of virus inoculum intranasally. They seroconverted by 14 days post exposure according to an Indirect Fluorescent Antibody Test. The contact-exposed gilts (#476 and #480) were immunized by the oronasal route after being placed in close contact with diseased pigs. These pigs seroconverted by 90 days post exposure. Seroconversion is indicated by an IFA titer greater than 1:20.

After the immunized gilts seroconverted, their estrus was synchronized and they were bred. Pregnancy was confirmed by ultrasound. On gestation day 90, vaccinate and control

pregnant gilts received 3.0 mL of PRRS challenge virus at a titer of 1×10^4 FAID₅₀, intranasally.

Control gilts #52 and #477 demonstrated early signs of disease in that they had reduced appetites and elevated body temperatures (103-104°F) for two to three days post challenge. No clinical signs of disease were noted in the ISU-P exposed (vaccinated) gilts. These results demonstrate protection from the clinical respiratory disease syndrome.

All gilts were held until pigs were weaned (4 weeks post farrowing). Table 2 shows the farrowing results from this experiment.

Farrowing results were recorded as to gestation length (gestation is normally shortened by PRRS infection), stillbirths or dead fetuses, weak pigs and mummies. These farrowing difficulties are all indicative of PRRS infection in pregnant gilts or sows. Additionally, the number of normal pigs were noted.

Neither of the control gilts farrowed normal pigs (0 of 22 pigs were normal). Seven (7) of 22 pigs were born dead and 15 of 22 were weak born pigs. The length of gestation of the control gilts was shortened by 2.5 days which is typical of PRRS in pregnant swine. In the immunized group (n = 50) 40 pigs or 80% were born normal. Six (6) of 50 pigs were born dead and 4 of 50 fetuses appeared mummified at farrowing. There were no weak pigs in the vaccinate group. It is clear that gilts previously exposed to live PRRSV ISU-P which are bred and then challenged are protected from challenge as demonstrated by lack of direct clinical signs of disease in the gilts and by protection of the fetuses which they carry.

As would be realized, since the IFA test reagents which were used to determine the serostatus of the gilts in this study were derived from PRRSV ISU-P and since there is a direct correlation between the serostatus of the gilts and protection from PRRS, PRRSV ISU-P isolate can be useful in diagnostic tests such as Indirect Fluorescent Antibody tests as

-10-

demonstrated herein, ELISAs, serum neutralization assays, Direct Fluorescent Antibody assays and other art-known diagnostic assays. Hence, a diagnostic assay derived from antigen obtained from PRRSV ISU-P can be prepared and used in accordance with the invention.

In the second vaccination/challenge study, young pigs (4 to 6 weeks of age) were vaccinated with several inactivated, adjuvanted PRRSV ISU-P vaccines. A vaccine antigen pool was prepared by inactivation of the PRRSV ISU-P with BEI as described earlier and adjuvanted with either Carbopol, FCA/FIA or Adj B. A second vaccine antigen pool was prepared by inactivation of the PRRSV ISU-P with formalin as described earlier and adjuvanted with the same 3 adjuvants. Three mock vaccines were produced in order to make sure that the adjuvants and cells could not stimulate a false immunological response. Mock vaccines were prepared by adjuvanting uninfected tissue culture cells. Two pigs were vaccinated with each vaccine. Vaccines were administered subcutaneously on days 0, 21, 42 and 64. Seven days after the last vaccination the pigs were challenged intranasally with $10^{3.5}$ FAID₅₀ of virulent live PRRSV. All pigs were observed daily post challenge for clinical signs of disease. Additionally, blood samples were drawn and processed for recovery of virus in tissue culture. Live virus found in the blood stream post challenge signifies that a viremia has occurred.

Serum neutralization titers were evaluated at the beginning of the study, on the day of challenge and at 19 days post challenge. It should be noted that the serum neutralization test is less sensitive than the IFA test used in the previous study. Therefore, one cannot compare these titer results with the previous titer results.

Table 3 lists the results of the challenge. No clinical signs of disease were observed in vaccinated or control pigs. Additionally, neither vaccinates nor controls developed serum neutralization titers during the vaccination period. This

-11-

indicates that there was no exposure to live virus during this time period (controls remained negative) and also indicates that the vaccines did not stimulate a humoral response high enough to be measured by a serum neutralization test. The fact
5 that serum neutralization titer response post challenge was significant indicates that all pigs were effectively challenged with the live PRRS virus. A verification of this challenge exposure is the percentage of control pigs demonstrating viremia. All but one of the control pigs (87.5%)
10 became infected. The immune response in vaccinated pigs was so strong that they were able to block viremia. Only 1 of 12 of the vaccinated pigs (8.3%) demonstrated viremia. Therefore, the vaccines protected 91.7% of the pigs.

Without being bound to any particular theory of the
15 invention, it is believed that force-passaging of the original PRRSV isolate through a specially-adapted clone of MA104(M) African Green Monkey Cells (9009B) may have successfully altered one or more epitopes of the virus resulting in enhanced immunogenicity of this virus so as to allow production of
20 vaccine in a field where other scientists have failed.

25

30

TABLE 1: Reproduction of late term abortion and reproduction failure in sows exposed to PRRSV isolate ISU-P as compared to non-exposed control sows.

I. PRRS Exposed Group						
Sow #	Gestation Length(a)	Virus Preparation	Stillbirths	Weak	Mummies	Normal
57	112 days	PRRS-hm ^b	6	8	0	0
54	108 days	PRRS-hm	6	1	0	0
649	112 days	PRRS-hm	3	7	0	0
58	112 days	PRRS-hm ^c	4	4	3	0
166	112 days	PRRS-tc ^d	0	2	6	1
164	113 days	PRRS-tc ^e	2	2	5	1
80	110 days	PRRS-tc	2	2	2	2
Totals			23	26	16	4

a = Normal swine gestation period is 114

b = PRRSV positive gnotobiotic pig lung homogenate inoculum

c = Tissue culture derived PRRSV, $10^{3.0}$ Fluorescent antibody infectious dose 50 per ml (FAID 50/ml)

d = tissue culture derived PRRSV, $10^{4.0}$ FAID50/ml

e = tissue culture derived PRRSV, $10^{5.0}$ FAID50/ml

II. Non-Exposed Control Group

Sow #	Gestation Length(a)	Virus Preparation	Stillbirths	Weak	Mummies	Normal
52	114 days	na	0	0	0	11
56	115 days	na	0	0	0	12
163	116 days	na	1	0	0	7
Totals			1	0	0	30

TABLE 2: Farrowing Results of ISU-P Immunized and Non-immunized Gilts

GILT NO.	IFA TITER	GESTATION LENGTH	STILLBIRTHS OR DEAD	WEAK PIGS	MUMMIES	NORMAL
57v	+	113 days	2*	0	0	7
58v	+	114 days	4	0	3	11
476v	+	113 days	0	0	1	10
480v	+	114 days	0	0	0	12
52 C	-	111 days	5	7	0	0
477	-	112 days	2	8	0	0
TOTALS			6 vacc. 7 cont.	0 vacc. 15 cont.	4 vacc. 0 cont.	40 vacc. 0 cont.

+ indicates an IFA titer of > 20
 - indicates an IFA titer of < 20

V indicates vaccinates
 C indicates controls

TABLE 3: PRRSV inactivated vaccine trial summary.

<u>Vaccine Group</u>	<u>n</u>	<u>Mean SN Each Challenge</u>	<u>Mean SN Post-Challenge^a</u>	<u>Virus Isolation</u>
BEI, PRRS-Carbopol	n = 2	<1:2	1:82	neg/neg
BEI, PRRS-FCA/FIA	n = 2	<1:2	1:4	neg/neg
BEI, PRRS- AdjB	n = 2	<1:2	1:5	neg/neg
Form., PRRS-Carbopol	n = 2	<1:2	1:23	neg/pos
Form., PRRS-FCA/FIA	n = 2	<1:2	1:2	neg/neg
Form., PRRS-AdjB	n = 2	1:3	1:8	neg/neg
<u>Mock Vaccine Group</u>				
Mock AG - carbopol	n = 2	<1:2	<1:2	pos/pos
Mock AG - FCA/FIA	n = 2	<1:2	<1:2	pos/pos
Mock AG - AdjB	n = 2	<1:2	<1:2	pos/pos

a = SN response on day 19 post-challenge, termination of the experiment.

b = Virus isolation from plasma at anytime during the 19 day post-challenge test period.
 Virus isolation detected by CPE in 9009B cells and confirmed by FA staining of the same with PRRSV group specific monoclonal antibody SDOW17.

Although the invention has been described in detail in the foregoing for the purpose of illustration, it is to be understood that such detail is solely for that purpose and that variations can be made therein by those skilled in the art without departing from the spirit and scope of the invention except as it may be limited by the claims.

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page _____, line _____	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet ☐	
Name of depositary institution AMERICAN TYPE CULTURE COLLECTION	
Address of depositary institution (including postal code and country) 12301 Parklawn Drive Rockville, MD 20652 USA	
Date of deposit March 16, 1993	Accession Number ATCC VR 2402 and ATCC CRL 11302
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet ☐	
None	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
All	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only ☒ This sheet was received with the international application Authorized officer	For International Bureau use only ☐ This sheet was received by the International Bureau on: Authorized officer

WHAT IS CLAIMED:

1. A process for growing a PRRSV in a tissue culture which is susceptible to infection to replicate the virus to an amount sufficient to protect animals against PRRS or be used in diagnosing PRRS or identifying the molecular structure of PRRSV for development of recombinant products, comprising inoculating the virus onto the tissue culture and harvesting the replicated virus.

2. The process of Claim 1 wherein the tissue culture comprises a cell line.

3. The process of Claim 2 wherein the tissue culture is an African Green Monkey Kidney cell line.

4. The process of Claim 3 wherein the African Green Monkey Kidney cell line is of the type designated as MA 104(M).

5. The process of Claim 4 wherein the cell line is a clone of the African Green Monkey cell line which clone is designated as 9009B.

6. A tissue culture containing the PRRSV as recited in Claim 1.

7. A tissue culture containing the PRRSV as recited in Claim 2.

8. A tissue culture containing the PRRSV as recited in Claim 3.

9. A tissue culture containing PRRSV produced according to Claim 1 on a commercial scale of >25L at a FAID₅₀ titer of $>10^{5.0}$ /mL.

10. A process for preparing a vaccine effective in protecting pigs against PRRS comprising providing PRRSV as recited in Claim 1, releasing the virus from the tissue culture cells and adjusting antigenic mass by dilution, concentration or extraction to produce an immunologically effective amount of the antigenic mass for a modified live (attenuated), inactivated, subunit or recombinant formulation.

11. A method of adapting the PRRSV to produce a modified live or attenuated vaccine by force-passaging the PRRSV through

non-host cells and cell lines until the virus no longer produces signs of PRRS.

12. A vaccine which is prepared by the process of Claim 10, wherein the PRRSV is modified live (attenuated) and is in a liquid, frozen or dessicated form.

13. The vaccine prepared using the process of Claim 9, wherein the PRRSV is inactivated and adjuvanted and optionally contains a suitable preservative.

14. The vaccine of Claim 13, wherein the adjuvant is either oil-based or aqueous-based.

15. The vaccine of Claim 13 wherein the adjuvant is selected from the group consisting of Freund's Complete Adjuvant, Freund's Incomplete Adjuvant, a carbopol based adjuvant and Diamond Scientific Adjuvant B.

16. The vaccine of Claim 13 wherein the inactivating agent is Binary ethyleneimine, Beta propiolactone or formalin.

17. The vaccine prepared according to the process of Claim 10, wherein the tissue culture is a cell line of the type classified as African Green Monkey cells.

18. The vaccine prepared according to the process of Claim 10, wherein the African Green Monkey cell is a cell line clone designated 9009B.

19. A vaccine prepared according to the process of Claim 10, wherein the antigenic mass is inactivated, modified live (attenuated), extracted and used as a subunit or obtained via recombinant technology.

20. A vaccine prepared by the process of Claim 10, wherein the PRRSV comprises a plurality of PRRSV isolates.

21. A vaccine prepared by the process of Claim 10, wherein the PRRSV is an isolate designated as ISU-P.

22. A method of altering the epitopes of the PRRSV to enhance immunogenicity by force-passaging the virus through an African Green Monkey Kidney cell line designated 9009B.

23. A vaccine prepared from the PRRSV of Claim 22.

INTERNATIONAL SEARCH REPORT

Internal Application No
PCT/US 94/00951

A. CLASSIFICATION OF SUBJECT MATTER
IPC 5 C12N7/00 A61K39/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,92 21375 (STICHTING CENTRAAL DIERGENEESKUNDIG INSTITUT) 10 December 1992 see page 2, line 14 - line 19 see page 3, line 13 - line 23 see page 8, line 3 - line 9 see page 9, line 35 - page 11, line 12 see page 13, line 22 - page 14, line 30 see claims 1-20 see figure 1 --- -/--	1,2,6,7, 9-19,23

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

24 May 1994

Date of mailing of the international search report

14.06.94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Fernandez y Branas, F

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 94/00951

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO,A,93 03760 (COLLINS, J.) 4 March 1993 see page 2, line 18 - page 3, line 17 see page 5, line 5 - page 8, line 7 see page 9, line 17 - page 10, line 24 see page 13, line 24 - page 16, line 7 see page 17, line 10 - line 33 see page 19, line 31 - page 20, line 11 see page 21, line 1 - line 8 see page 24, line 9 - page 25, line 3 see claims 36,37 ---	1-4, 6-19,23
E	EP,A,0 595 436 (SOLVAY ANIMAL HEALTH INC.) 4 May 1994 see page 7, line 6 - page 10, line 24 see page 11, line 45 - page 14, line 4 see page 19 - page 22; example II see page 41; example VIII ---	1-4, 6-19,23
X,P	EP,A,0 529 584 (BOEHRINGER INGELHEIM ANIMAL HEALTH INC.) 3 March 1993 see the whole document ---	1-4, 6-12,17, 19,21,22
X,P	EP,A,0 541 418 (RHONE MERIEUX S.A.) 12 May 1993 see the whole document ---	1,2,6,7, 9-14, 17-19,23
X,P	WO,A,93 07898 (AKZO N.V.) 29 April 1993 see the whole document ---	1,2,6,7, 9-19,23
X,P	ARCHIVES OF VIROLOGY vol. 133, no. 3-4 , 1993 , NEW YORK pages 477 - 483 KIM, H.S. ET AL 'Enhanced replication of porcine reproductive and respiratory syndrome (PRRS) virus in a homogeneous subpopulation of ma-104 cell line' see the whole document ---	1-4,6-9
X	AMERICAN JOURNAL OF VETERINARY RESEARCH vol. 53, no. 4 , 1992 , CHICAGO pages 485 - 488 CHRISTIANSON W.T. 'Experimental reproduction of swine infertility and respiratory syndrome in pregnant cows' see the whole document -----	1,2,6,7, 9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 94/00951

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9221375	10-12-92	AU-A- 2196592 EP-A- 0587780	08-01-93 23-03-94
WO-A-9303760	04-03-93	AU-A- 2508492 CA-A- 2076744 EP-A- 0529584 AU-A- 2696792 WO-A- 9306211	16-03-93 27-02-93 03-03-93 27-04-93 01-04-93
EP-A-0595436	04-05-94	AU-B- 5037593	12-05-94
EP-A-0529584	03-03-93	AU-A- 2508492 CA-A- 2076744 WO-A- 9303760	16-03-93 27-02-93 04-03-93
EP-A-0541418	12-05-93	FR-A- 2682966 CA-A- 2081638	30-04-93 30-04-93
WO-A-9307898	29-04-93	AU-A- 2684792	21-05-93