



- (51) **International Patent Classification:**
C12N 1/20 (2006.01) *C07K 14/32* (2006.01)
C07K 14/195 (2006.01)
- (21) **International Application Number:**
PCT/EP2012/060624
- (22) **International Filing Date:**
5 June 2012 (05.06.2012)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
11352006.8 6 June 2011 (06.06.2011) EP
- (71) **Applicants** (*for all designated States except US*): **COM-MISSARIAT À L'ÉNERGIE ATOMIQUE ET AUX ÉNERGIES ALTERNATIVES** [FR/FR]; 25, rue Leblanc, Bâtiment "Le Ponant D", F-75015 Paris (FR). **UNIVERSITE D'AVIGNON ET DES PAYS DE VAUCLUSE** [FR/FR]; 74, rue Louis Pasteur, F-84029 Avignon cedex 1 (FR).
- (72) **Inventors; and**
- (75) **Inventors/Applicants** (*for US only*): **ARMENGAUD, Jean** [FR/FR]; 26, rue Camp de Bataille, F-30400 Villeneuve-lez-Avignon (FR). **CLAIR, Gérémy** [FR/FR]; 11, rue du Chêne Vert, F-30400 Villeneuve-lez-Avignon (FR). **DUPORT, Catherine** [FR/FR]; 989, Rte de Cavaillon, F-84250 Le Thor (FR).
- (74) **Agents:** **AUGARDE, Eric** et al.; Brevalex, 56, boulevard de l'Embouchure, B.P. 27519, F-31075 Toulouse Cedex 2 (FR).
- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- *with international search report (Art. 21(3))*
 - *with sequence listing part of description (Rule 5.2(a))*

(54) **Title:** A METHOD FOR ENHANCING THE FERMENTATIVE POTENTIAL AND GROWTH RATE OF MICROORGANISMS UNDER ANAEROBIOSIS

(57) **Abstract:** The present invention concerns a method for enhancing the growth rate, the biomass production and/or the fermentative potential of an anaerobic microorganism, comprising the steps consisting in:(a) modifying said anaerobic microorganism by inactivating the *ohrR* gene, the *ohrR* gene product or homolog thereof;(b) culturing the modified anaerobic microorganism obtained in step (a) under low-ORP anaerobic conditions.



**A METHOD FOR ENHANCING THE FERMENTATIVE POTENTIAL
AND GROWTH RATE OF MICROORGANISMS UNDER ANAEROBIOSIS**

DESCRIPTION

5 TECHNICAL FIELD

The present invention belongs to the technical field of methods using microorganisms and, in particular, of methods using genetically modified microorganisms that have enhanced fermentation activity and/or enhanced growth rate in selected particular culture conditions.

The present invention proposes a method that results in a genetically modified strain exhibiting higher anaerobic growth rate using basically the same culture condition as the wild-type strain. More particularly, the present invention method is based on the abolishment of the synthesis of a protein that is substantially similar to *Bacillus cereus* OhrR (for "Organic hydroperoxide resistance Regulator").

The present invention relates to a method for increasing the fermentative potential and/or the growth rate of a microorganism under low Oxido-Reduction Potential (low-ORP) anaerobic culture conditions.

25 STATE OF THE PRIOR ART

Nowadays many microorganisms are implemented for different industrial applications. Fields of application are covering industrial processes related to human or animal health (red biotechnology) ; agro-industries, agronomy and farming (green

biotechnology) ; marine and aquatic processes (blue technology) ; chemical industry (white biotechnology) ; energy industry and, in particular, biomass conversion field ; environmental processes such as bioremediation,
5 water and industrial fluid purification, CO₂ captation, etc... Most of these processes use the ability of microorganisms to produce cell biomass, proteins and/or primary/secondary metabolites.

As for examples, microorganisms can be used in
10 the production of food products such as wine or beer ; in the production of biofuels, notably by the conversion of sugar into ethanol ; in the pharmaceutical field to produce active ingredients such as antibiotics or insulin; in the production of
15 valuable chemicals or in the destruction of hazardous and/or polluting chemicals.

As a consequence, there is a real need to propose methods that make it possible to obtain an enhanced production and/or an enhanced production yield
20 in the processes using microorganisms.

This technical problem can be solved by selecting appropriate culture conditions. Thus, international application WO 2009/101328 [1] proposes a method for culturing microorganisms in order to
25 increase their biomass and their metabolic activity. The method disclosed consists in a combined adjustment of the Oxidation-Reduction Potential (ORP) and the dissolved oxygen. Indeed, the inventors of this application have noted that using a strain of
30 *Lactococcus lactis* as microorganisms and adjusting simultaneously the redox potential within the range

extending from -400 to 0 mV and the dissolved oxygen value within the range extending from 1 to 30%, large gains are obtained in biomass and acidifying activity. Nevertheless, the main drawbacks of the process disclosed in [1] concern the fact that it can only be implemented with facultative anaerobe or aerobe strains because of the presence of dissolved oxygen and that it requires the continuous adjustment of parameters which influence each other.

Alternatively, the above-identified technical problem can be solved by identifying and using microorganisms which present enhanced production, enhanced production yield and/or enhanced growth rate, when compared to the microorganisms implemented until now in the above-listed processes. Most of the time, these microorganisms are genetically modified.

Among the different studies already disclosed on such microorganisms, one can cite the patent applications FR 2 635 788 [2] and FR 2 778 922 [3]. The former concerns the use of a nucleic acid comprising a functional nucleotide sequence contained in a 14 kb fragment obtained from the chromosome of a *Bacillus subtilis* strain having a *sacU^h* mutation. Once introduced into *B. subtilis* or into any other microorganism, said sequence is capable of exploiting to its advantage the information carried by this fragment, of inducing therein an overproduction of proteins or of conferring on it a *Deg^h* phenotype [2]. The patent application FR 2 778 922 discloses the use of a polynucleotide comprising all or part of at least one regulatory sequence of a *Clostridium* toxin

synthesis operon to modify *Clostridium* or *Escherichia coli* and to promote the synthesis of a homologous or heterologous protein therein [3].

International application WO 2011/018116 [4] proposes a metabolically enhanced photoautotrophic, ethanol producing host cell comprising : (1) at least two first metabolic enhancements reducing the enzymatic activity or affinity of at least two endogenous host cell enzymes involved in acetate and lactate fermentation, the first metabolic enhancements resulting in an enhanced level of biosynthesis of acetaldehyde, pyruvate, acetyl-CoA or precursors thereof compared to the respective wild type host cell, (2) at least one second metabolic enhancement different from the first metabolic enhancement comprising an overexpressed enzyme for the formation of ethanol. The host cell implemented in this method is a prokaryotic cell and advantageously a cyanobacterial cell.

Moreover, international application WO 2011/006136 [5] proposes a method for producing ethanol, ethanol derivatives, ethanol conjugates and/or other organic chemical intermediates using biological system. This latter consists in a microorganism strain and, in particular, a *Sacharomyces cerevisiae* strain or a bacterium that includes an alteration that adds or increases a phosphogluconate dehydratase activity, a 2-keto-3-deoxygluconate-6-phosphate aldolase activity and a xylose isomerase activity.

It must be pointed out that some of the genetically-modified microorganisms used in the above-cited prior art only make it possible to enhance the

production of particular proteins and/or of particular primary/secondary metabolites. Other genetically-modified microorganisms require several genetic modification steps to be prepared.

5

DISCUSSION OF THE INVENTION

The present invention based on unexpected data recorded by the inventors makes it possible to solve the above-identified technical problem and some drawbacks of the prior art methods.

Indeed, the present inventors have worked on *Bacillus cereus* that is a motile facultative anaerobe responsible of diarrhoeal disease in humans. To survive in the gastrointestinal tract and initiate infection, this microorganism is able to adapt to variable oxygen and oxidoreduction potential (ORP) conditions by modulating its central metabolic pathways [6].

Consequently, the present inventors have been aimed at elucidating the regulatory mechanisms tuning these pathways. They have shown that a dominant component of these regulatory mechanisms involves whole proteome remodeling that restricts fermentative potential while boosting motility, respiratory potential and oxidative stress resistance. Specifically, gene inactivation studies and high-throughput proteome analyses have identified organic hydroxyperoxide resistance (Ohr)-homologous proteins, OhrA (for "Organic hydroxyperoxide resistance A") and OhrR, as key players of the proteome remodeling occurring in response to oxygen availability and extracellular ORP changes.

In addition and unexpectedly, the present inventors have noted that disruption of *ohrR* gene leads to a significant increase ($56\% \pm 10\%$) of the growth rate of fermentative cells under low-ORP anaerobic conditions compared with wild-type cells and no significant difference can be observed under high-ORP anaerobiosis and aerobiosis. On the contrary, the growth rate of the $\Delta ohrA$ mutant is significantly lower than that of the wild-type under aerobiosis ($-29\% \pm 3\%$) and in a lesser extent under anaerobiosis ($-22\% \pm 2\%$).

Thanks to these results, the inventors have identified that the combination of a particular microorganism i.e. a $\Delta ohrR$ mutant and of particular culture conditions i.e. low-ORP anaerobic conditions makes it possible to obtain a strain with enhanced growth rate and enhanced fermentative potential, when compared with the growth rate and the fermentative potential of the wild-type strain, i.e. the non mutated strain, grown in the same culture conditions.

A key advantage of this modified strain compared with the wild-type strain is that the anaerobic growth rate in low-ORP condition can be significantly increased without modifying substantially the growth conditions, resulting in the production of higher levels of biomass in less time. Another aspect of the invention is to achieve higher growth rate of the modified strain in low-ORP anaerobic conditions compared with high-ORP anaerobic conditions.

In addition, since *B. cereus* OhrR homologues are present in many microorganisms used or susceptible to be used in industrial applications, the process

described here can be applied to many microorganisms in order to improve the production of cell biomass, proteins, or primary/secondary metabolites; or increase the yield of any process into which these micro-organisms are used.

The inventors have also noted that *ohrR* disruption strongly decreases the abundance level of twelve flagellar and chemotaxis-related proteins besides *OhrA*. Thus the *ohrR* mutant presents a decreased motility which can be an additional advantage when using the strain in decontamination process. Indeed, in such processes, strains which stay where they have been inoculated are preferred.

Thus, the present invention concerns a method for enhancing the growth rate, the biomass production and/or the fermentative potential of an anaerobic microorganism, comprising the steps consisting in :

a) modifying said anaerobic microorganism by inactivating the *ohrR* gene, the *ohrR* gene product or homolog thereof ;

b) culturing the modified anaerobic microorganism obtained in step (a) under low-ORP anaerobic conditions ;

in consideration of which (or in view of which or thereby) the growth rate, the biomass production and/or the fermentative potential of said modified microorganism is enhanced when compared with the growth rate, the biomass production and/or the fermentative potential of the corresponding non modified microorganism cultured under the same conditions.

For microorganisms, growth results in an increase in the number of individuals. This increase comes from one cell division into two new cells. In the
5 present invention, the growth rate can be determined by different experimental methods.

Indeed, the growth rate can be determined by counting the number of microorganisms. More particularly, a volume v of the microorganism culture
10 is taken after a time t of culture under low-ORP anaerobic conditions in a medium m , the number of microorganisms is manually counted in this volume v thanks to a microscope (Petroff-Hausser counting chamber) or thanks to electronic counters such as
15 Coulter counter.

The growth rate can also be determined by measuring the turbidity of the microorganism culture. More particularly, a volume v of the microorganism culture is taken after a time t of culture under low-
20 ORP anaerobic conditions in a medium m . The turbidity of this volume v is measured thanks to a turbidimeter or a colorimeter typically at a wavelength of 560 nm.

Alternatively, the growth rate can be determined by plate counting of CFU (for "Colony
25 Forming Unit"). In practice, a volume v of the microorganism culture optionally diluted is spread on the surface of solidified nutrient medium for example agar with 4% glucose, which when incubated, leads to each cell developing into a visible colony through
30 repeated fission. After incubation during a time t , all

colonies formed are counted and the number of CFU in the volume v is calculated.

When comparing the growth rate of the modified microorganism implemented in the present invention with the non modified corresponding microorganism, it is clear that both growth rates must be measured thanks to the same method, in an identical volume v of microorganism culture after an identical time t of culture under identical low-ORP anaerobic conditions in an identical medium m , modified and non modified microorganisms having been seeded in the medium m at the same concentration.

The biomass production can be determined by measuring the dry weight of the microorganism culture. More particularly, a volume v of the microorganism culture is taken after a time t of culture under low-ORP anaerobic conditions in a medium m . This volume v is dried for example at 110°C during 12 h, then the dried sample thus obtained is weighed.

When comparing the biomass production of the modified microorganism implemented in the present invention with the non modified corresponding microorganism, it is clear that both biomass productions must be measured in an identical volume v of microorganism culture after an identical time t of culture under identical low-ORP anaerobic conditions in an identical medium m , modified and non modified microorganisms having been seeded in the medium m at the same concentration.

In addition, once the biomass production has been obtained, the specific growth rate dy/dt can be calculated from the growth curve which was modeled using the modified Gompertz equation :

5
$$y = A \exp\{-\exp[(\mu_m \cdot e/A) (\lambda - t) + 1]\},$$

wherein y is the biomass production (in $g.l^{-1}$), μ_m the maximum specific growth rate (in h^{-1}), λ the lag time (in h), A and e , the mathematical parameters describing the sigmoidal growth curve and t the time
10 (in h).

The fermentative potential is representative of the catabolic activity of the anaerobic microorganism. The fermentative potential can be determined by
15 measuring the quantity of a particular element in the microorganism culture. Said particular element which can be a metabolite must be representative of the fermentative activity of the microorganism. In other words, the particular element is a molecule produced by
20 the microorganism when submitted to the low-ORP anaerobic conditions. As for examples, one can cite ethanol, acetate, formate, lactate, acetoin, butanediol and CO_2 . Clearly, this particular element will depend on the microorganism implemented in the present
25 invention.

More particularly, a volume v of the microorganism culture is taken after a time t of culture under low-ORP anaerobic conditions in a medium m and the quantity of the particular element is
30 quantified in the volume v . This quantification is

carried out by using commercial kits or methods well-known by the one skilled in the art.

When comparing the fermentative potential of the modified microorganism implemented in the present invention with the non modified corresponding microorganism, it is clear that both fermentative potentials must be measured in an identical volume v of microorganism culture after an identical time t of culture under identical low-ORP anaerobic conditions in an identical medium m , modified and non modified microorganisms having been seeded in the medium m at the same concentration.

The one skilled in the art will choose the appropriate time t and the appropriate volume v to conduct any comparison between the modified microorganism and the non modified microorganism, according to these microorganisms and the medium implemented.

In the context of the present invention, only one of the parameters chosen from the growth rate, the biomass production and the fermentative potential can be enhanced in the modified microorganism when compared to the non modified microorganism.

Alternatively, two of these parameters, i.e. growth rate and biomass production; growth rate and fermentative potential; biomass production and fermentative potential, can be enhanced in the modified microorganism when compared to the non modified microorganism.

Alternatively again, the three parameters can be enhanced in the modified microorganism when compared to the non modified microorganism.

The term "enhanced" means that the growth rate, the biomass production or the fermentative potential is increased and, in particular, significantly increased when the value (V_1) of one of these parameters determined for the modified microorganism is compared to the value (V_2) of the corresponding parameter determined for the non modified corresponding microorganism. This increase (I) of formula $(I) = [(V_1 - V_2) / V_2 \times 100]$ can be superior or equal to 10%, notably superior or equal to 20%, in particular, superior or equal to 40% and, more particularly, superior or equal to 50%.

An anaerobic microorganism is a microorganism that thrives in oxygen-deprived areas and its growth in oxygen-proficient areas is severely limited.

The anaerobic microorganism which can be used in the method according to the present invention can be a facultative anaerobic microorganism or an obligate anaerobic microorganism. A facultative anaerobic microorganism makes ATP by aerobic respiration if oxygen is present but is also capable of switching to fermentation in the absence of oxygen or to anaerobic respiration in the absence of oxygen and in the presence of at least one external electron acceptor such as nitrates. In contrast, an obligate anaerobic microorganism is incapable of aerobic respiration.

The anaerobic microorganism which can be used in the method according to the present invention can be any natural, genetically-modified or synthetic, eukaryotic or prokaryotic, anaerobic microorganism. Advantageously, the anaerobic microorganism which can be used in the method according to the present invention can be any natural, genetically-modified or synthetic, prokaryotic, anaerobic microorganism. For example, a suitable anaerobic microorganism can be an anaerobic bacteria and notably a facultative anaerobic bacteria or an obligate anaerobic bacteria.

By "genetically-modified microorganism", it must be understood that the microorganism can have already been genetically modified before the inactivation of the *ohrR* gene, of the *ohrR* gene product or of the homolog thereof. As used herein, the expressions "genetically-modified" and "genetically-engineered" are equivalent and can be used interchangeably.

This genetic modification can consist in inserting a gene or a particular polynucleotide sequence in the microorganism to confer a particular property to the microorganism such as the capacity of utilizing carbohydrates or to activate the activity for which this gene or this particular polynucleotide sequence encodes, such as the expression of an enzyme. For example, this gene or this particular polynucleotide sequence encodes enzymes in a particular metabolic pathway such as in the metabolic pathway of ethanol. In this example, this gene or this particular polynucleotide sequence encodes an enzyme that

metabolizes pentose and/or hexose sugars. Alternatively, this genetic modification can consist in the inactivation of a competing pathway such as a competing pathway for carbon flow by inactivating various genes. Additionally, this genetic modification can consist in conferring or improving the tolerance to a particular element, such as, for example, butanol tolerance, through chemical mutagenesis and mutant screening.

By "synthetic microorganism", it must be understood a genome-reduced host constructed and engineered by synthetic biology. This synthetic microorganism is obtained by introducing selected genetic information from source organism(s) into a genetically friendlier host, also designed as "chassis" or "recipient". Particular examples of processes suitable to prepare such synthetic microorganism can be found in Gao et al., 2010 [7] and in Gibson et al., 2010 [8].

A natural, genetically-modified or a synthetic anaerobic microorganism suitable for the method according to the present invention presents an *ohrR* gene or a homolog thereof. This gene is designed as "native gene" or "endogenous gene". Alternatively, the selected genetic information from source organism(s) introduced into the recipient host to prepare a synthetic anaerobic microorganism can lack an *ohrR* gene or a homolog thereof. In this alternative, the step (a) of the method according to the present invention is carried out while preparing the synthetic anaerobic microorganism.

Suitable natural or genetically-modified microorganisms for the method of the present invention include, but are not limited to, members of the genera : Abiotrophia, Acetivibrio, Acetobacter, 5 Acetohalobium, Achromobacter, Acidaminococcus, Acidiphilium, Acidithiobacillus, Acidobacterium, Acidothermus, Acidovorax, Acinetobacter, Actinomyces, Actinosynnema, Aerococcus, Aeromicrobium, Aeromonas, Afipia, Agrobacterium, Alcanivorax, Algoriphagus, 10 Alicyclophilus, Alicyclobacillus, Alkaliphilus, alpha, Alteromonadales, Alteromonas, Aminobacterium, Aminomonas, Amycolatopsis, Amycolicicoccus, Anabaena, Anaerococcus, Anaerofustis, Anaerolinea, Anaeromyxobacter, Anaerostipes, Anaerotruncus, 15 Anoxybacillus, Archaeoglobus, Arcobacter, Aromatoleum, Arthrobacter, Asticcacaulis, Atopobium, Aurantimonas, Azoarcus, Azorhizobium, Azospirillum, Azotobacter, Bacillus, Bacteriovorax, Bacterium, Bacteroides, Bdellovibrio, Beijerinckia, Beta, Beutenbergia, 20 Bilophila, Blastopirellula, Blautia, Bordetella, Brachyspira, Bradyrhizobium, Brevibacillus, Brevibacterium, Brevundimonas, Brucella, Bryantella, Bulleidia, Burkholderia, Burkholderiales, Butyrivibrio, Caldicellulosiruptor, Candidatus, Capnocytophaga, 25 Carboxydotherrmus, Cardiobacterium, Carnobacterium, Catenibacterium, Catenulispora, Caulobacter, Cellulomonas, Cellulophaga, Cellvibrio, Chitinophaga, Chlorobaculum, Chlorobium, Chloroflexus, Chloroherpeton, Chromobacterium, Chromohalobacter, 30 Chryseobacterium, Chthoniobacter, Citreicella, Citrobacter, Citromicrobium, Clavibacter, Clostridia,

Clostridiales, Clostridium, Collinsella, Colwellia,
Comamonas, Conexibacter, Congregibacter, Coprobacillus,
Coproccoccus, Coprothermobacter, Coraliomargarita,
Corynebacterium, Coxiella, Croceibacter, Cronobacter,
5 Cryptobacterium, Cupriavidus, Curvibacter, Cyanothece,
Cytophaga, Deinococcus, Delftia, delta, Denitrovibrio,
Desmospora, Desulfarculus, Desulfatibacillum,
Desulfitobacterium, Desulfobacca, Desulfobacterium,
Desulfococcus, Desulfohalobium, Desulfomicrobium,
10 Desulfotalea, Desulfotomaculum, Desulfovibrio,
Desulfurispirillum, Desulfuromonas, Dethiobacter,
Dethiosulfovibrio, Dichelobacter, Dickeya,
Dictyoglomus, Dietzia, Dinoroseobacter, Dokdonia,
Dorea, Dyadobacter, Dysgonomonas, Edwardsiella,
15 Eggerthella, Eikenella, Elusimicrobium, Enhydrobacter,
Enterobacter, Enterococcus, Eremococcus, Erwinia,
Erysipelotrichaceae, Erythrobacter, Escherichia,
Ethanoligenens, Eubacterium, Exiguobacterium,
Faecalibacterium, Ferrimonas, Ferroglobus,
20 Fervidobacterium, Fibrobacter, Finegoldia,
Flavobacteria, Flavobacteriaceae, Flavobacteriales,
Flavobacterium, Fluviicola, Frankia, Fulvimarina,
Fusobacterium, gamma, Gemella, Gemmata, Gemmatimonas,
Geobacillus, Geobacter, Geodermatophilus, Glaciecola,
25 Gloeobacter, Gluconacetobacter, Gordonina,
Gordonibacter, Gramella, Granulicatella, Grimontia,
Hahella, Halalkalicoccus, Halanaerobium, Haliangium,
Halomonas, Heliobacterium, Herbaspirillum,
Herminiimonas, Herpetosiphon, Hirschia, Hoeflea,
30 Holdemania, Hylemonella, Hyphomicrobium, Hyphomonas,
Idiomarina, Ilyobacter, Intraspangium, Isosphaera,

Janibacter, Jannaschia, Janthinobacterium, Jonesia,
 Jonquetella, Kineococcus, Kitasatospora, Klebsiella,
 Kluyvera, Kocuria, Kordia, Kosmotoga, Kribbella,
 Krokinobacter, Ktedonobacter, Labrenzia,
 5 Lachnospiraceae, Lacinutrix, Lactobacillus,
 Lactococcus, Lautropia, Leadbetterella,
 Leeuwenhoekella, Legionella, Leptospira,
 Leptospirillum, Leptothrix, Leptotrichia, Leuconostoc,
 Limnobacter, Listeria, Lutiella, Lysinibacillus,
 10 Macrococcus, Magnetococcus, Magnetospirillum, Mahella,
 Maribacter, Maricaulis, Marinithermus, Marinobacter,
 Marinomonas, Mariprofundus, Maritimibacter, Marivirga,
 Megamonas, Megasphaera, Meiothermus, Mesorhizobium,
 Metallosphaera, Methanobacterium, Methanobrevibacter,
 15 Methanocella, Methanococcus, Methanohalophilus,
 Methanoplanus, Methanosarcina, Methanosphaera,
 Methanosphaerula, Methanospirillum, Methylibium,
 Methylobacter, Methylobacterium, Methylocella,
 Methylocystis, Methyломicrobium, Methylophaga,
 20 Methylophilales, Methylosinus, Methylothermus,
 Methyloversatilis, Methylovorus, Microbacterium,
 Micrococcus, Micromonospora, Microscilla, Mitsuokella,
 Moorella, Moritella, Mucilaginibacter, Mycobacterium,
 Mycoplasma, Myxococcus, Nakamurella, Natranaerobius,
 25 Neisseria, Nitrobacter, Nitrosococcus, Nocardia,
 Nocardioideae, Nocardioideae, Nocardiosis, Nostoc,
 Novosphingobium, Oceanibulbus, Oceanicaulis,
 Oceanicola, Oceanobacillus, Oceanobacter,
 Oceanospirillum, Ochrobactrum, Octadecabacter,
 30 Odoribacter, Oenococcus, Oligotropha, Olsenella,
 Oribacterium, Oscillatoria, Oscillochloris,

Oxalobacter, Oxalobacteraceae, Paenibacillus,
 Paludibacter, Pantoea, Parabacteroides, Paracoccus,
 Paraprevotella, Parasutterella, Parvibaculum,
 Parvularcula, Pectobacterium, Pediococcus, Pedobacter,
 5 Pelagibaca, Pelobacter, Pelotomaculum,
 Peptostreptococcus, Petrotoga, Phaeobacter,
 Phenylobacterium, Photobacterium, Photorhabdus,
 Physcomitrella, Picrophilus, Pirellula, Planctomyces,
 Planococcus, Plautia, Polaribacter, Polaromonas,
 10 polymorphum, Polynucleobacter, Populus, Prevotella,
 Propionibacterium, Prosthecochloris, Providencia,
 Pseudoalteromonas, Pseudomonas, Pseudonocardia,
 Pseudoramibacter, Pseudovibrio, Pseudoxanthomonas,
 Psychroflexus, Psychromonas, Pusillimonas,
 15 Pyramidobacter, Rahnella, Ralstonia, Reinekea,
 Renibacterium, Rhizobium, Rhodobacter, Rhodobacterales,
 Rhodococcus, Rhodoferax, Rhodopirellula,
 Rhodopseudomonas, Rhodospirillum, Rhodothermus,
 Ricinus, Robiginitalea, Roseburia, Roseibium,
 20 Roseiflexus, Roseobacter, Roseomonas, Roseovarius,
 Rothia, Rubrivivax, Rubrobacter, Ruegeria,
 Ruminococcus, Saccharomonospora, Saccharopolyspora,
 Sagittula, Salinispora, Salmonella, Sanguibacter,
 Sebalbella, Segniliparus, Selenomonas, Serratia,
 25 Shewanella, Shigella, Shuttleworthia, Sideroxydans,
 Silicibacter, Simonsiella, Sinorhizobium, Slackia,
 Sodalis, Solibacillus, Solobacterium, Sorangium,
 Sordaria, Sphaerobacter, Sphingobacterium, Sphingobium,
 Sphingomonas, Sphingopyxis, Spirochaeta, Spirosoma,
 30 Stackebrandtia, Staphylococcus, Stappia, Starkeya,
 Stenotrophomonas, Stigmatella, Streptococcus,

Streptomyces, Streptosporangium, Subdoligranulum,
 Sulfitobacter, Sulfolobus, Sulfuricurvum, Sutterella,
 Symbiobacterium, Synechococcus, Synechocystis,
 Synergistetes, synthetic, Syntrophobacter,
 5 Syntrophobotulus, Syntrophomonas, Syntrophothermus,
 Syntrophus, Tepidanaerobacter, Teredinibacter,
 Terriglobus, Thalassiobium, Thauera, Thermaerobacter,
 Thermincola, Thermoanaerobacter, Thermoanaerobacterium,
 Thermobaculum, Thermobifida, Thermobispora,
 10 Thermomicrobium, Thermomonospora, Thermoplasma,
 Thermosediminibacter, Thermosipho, Thermotoga,
 Thermotogales, Thioalkalivibrio, Thiobacillus,
 Thiomonas, Tolumonas, Treponema, Tsukamurella,
 Turicibacter, Variovorax, Veillonella,
 15 Verminephrobacter, Verrucomicrobiae, Verrucomicrobium,
 Verrucosipora, Vibrio, Vibrionales, Victivallis,
 Weissella, Wolinella, Xanthobacter, Xanthomonas,
 Xenopus, Xenorhabdus, Xylella, Yersinia and
 Zunongwangia. Indeed, in each above-listed genus, at
 20 least one representative comprises a close OhrR
 homologue identified using the OhrR amino acid sequence
 from *Bacillus cereus* by two iterations of Position
 Specific Iterated-BLAST (PSI-BLAST) (p value < 1e-10).

25 Based on published results, the *ohrA* gene is
 known in bacteria as a gene coding a thiol peroxidase
 that catalyses the reduction of an organic
 hydroxyperoxide to its corresponding less harmful
 alcohol. The expression of this gene is strongly and
 30 selectively induced by organic peroxides. This
 regulation requires OhrR, an organic peroxide-inducible

negative regulator belonging to the MarR family of repressor proteins [9].

In *Bacillus cereus* ATCC14579, the gene *ohrR* codes for the protein OhrR, the amino acid sequence of which can be obtained in the NCBI entrez database under the accession number NP_834186.1 (http://www.ncbi.nlm.nih.gov/protein/NP_834186.1) and is the following:

```
MTEDSLHLDNQLCFSIYACSREVTRFYRPYLEEMGITYPQYITLLVL
10 WEQDGLTVKEIGERLFLDSGTLTPMLKRMESLNLVKRVRSKEDERKVCIELTEQG
KDLQDKACSLPTTMATNLGITEQEYRSLLIQLNKLIETMKTINDRKGE (SEQ
ID NO: 1 in the appended sequence listing).
```

The *ohrR* gene product refers not only to the polypeptide or protein OhrR but also to the corresponding mRNA.

By « homolog thereof » it must be understood in the present invention not only a homolog of the gene *ohrR* of *Bacillus cereus* and, in particular, of *Bacillus cereus* ATCC14579 but also a homolog of the gene product of *Bacillus cereus* and, in particular, of *Bacillus cereus* ATCC14579.

As used in the present invention, the terms "homolog" and "homologous" refer to the relationship between genes, nucleic acids, ribonucleic acids or proteins that possess a common evolutionary origin and/or a common function, including homologous genes, nucleic acids, ribonucleic acids or proteins from different species. Such proteins and their encoding nucleic acid sequences have sequence homology, as reflected by their sequence identity, whether in terms

of percent identity or similarity, or the presence of specific residues or motifs at conserved positions.

In the present application, a homolog to *ohrR* gene is a gene coding for an amino acid sequence identified or detected using the *OhrR* amino acid sequence from *Bacillus cereus* by two iterations of Position Specific Iterated-BLAST (PSI-BLAST) (p value < 1e-10). Similarly, a homolog to the *ohrR* gene product is an amino acid sequence or a RNA coding for an amino acid sequence, wherein these amino acid sequences are identified or detected using the *OhrR* amino acid sequence from *Bacillus cereus* by two iterations of Position Specific Iterated-BLAST (PSI-BLAST) (p value < 1e-10). In other words, when two iterations of Position Specific Iterated-BLAST (PSI-BLAST) (p value < 1e-10) using the *OhrR* amino acid sequence from *Bacillus cereus* are implemented, the amino acid sequence of the *ohrR* gene product or coded by the *ohrR* gene product (RNA) or by the *ohrR* gene already present in the database used or just added therein is identified. Any database known by the one skilled in the art can be used for these iterations of PSI-BLAST (p value < 1e-10). For example, one can cite the NCBI non-redundant protein sequences database (NCBI nr), the reference proteins database (refseq_protein), the UniProtKB/Swiss-Prot database (swissprot), the metagenomic proteins database (NCBI env_nr).

In a specific embodiment, two genes or (ribo)nucleic acid sequences are homologous or substantially identical when at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least

70%, at least 80%, at least 90% or at least 95%, of the (ribo)nucleotides match over the defined length of the (ribo)nucleic acid sequence, as determined by sequence comparison algorithms, such as BLAST, FASTA or DNA Strider. Sequences that are homologous can be identified by comparing the sequences using standard software available in sequence databases, or in a Southern hybridization experiment under, for example, stringent conditions. The appropriate stringency conditions for hybridizing nucleic acids in terms of temperature and of ionic strength depends on the length of the (ribo)nucleic acids and the degree of complementation. Such conditions can be easily determined by the one skilled in the art.

Similarly, in another specific embodiment, two amino acid sequences are homologous or substantially identical when at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90% or at least 95% of the amino acids are identical. Advantageously, the similar or homologous amino acid sequences are identified by alignment using, for example, sequence comparison algorithms, such as BLAST, FASTA or DNA Strider.

Thus, a homolog of the *ohrR* gene is a gene or a nucleic acid presenting an identity in the nucleotide sequence of at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90% or at least 95% with the *ohrR* gene nucleotide sequence of *Bacillus cereus* ATCC14579.

In addition, a homolog of the *ohrR* gene product is a ribonucleic acid or a polypeptide presenting an

identity in the ribonucleotide or in the amino acid sequence of at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90% or at least 95% with the *ohrR* gene product
5 ribonucleotide or amino acid sequence of *Bacillus cereus* ATCC14579.

It must be reminded that percent identity between two (ribo)nucleotide or amino acid sequences is generally determined by aligning the residues of the
10 two sequences to optimize the number of identical (ribo)nucleotides or amino acids along the lengths of their sequences; gaps in either or both sequences are permitted in making the alignment in order to optimize the number of identical (ribo)nucleotides or amino
15 acids, although the (ribo)nucleotides or amino acids in each sequence must nonetheless remain in their proper order.

Advantageously, identical sequences are or encode for proteins having identical or similar
20 function. In the present case, identical sequences should be or encode a repressor of *OhrA* or of a gene homolog to *OhrA*.

In addition, when percentage of sequence identity is used in reference to proteins, it is
25 recognized that residue positions which are not identical often differ by conservative amino acid substitutions, where amino acid residues are substituted for other amino acid residues with similar chemical properties, such as charge, hydrophilicity or
30 hydrophobicity, and therefore do not change the functional properties of the molecule. When sequences

differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the substitution. Sequences that differ by such conservative substitutions are said to have "sequence similarity" or "similarity". Means for making this adjustment are well known to the one skilled in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity.

Conservative substitution tables providing functionally similar amino acids are well known in the art. Indeed, the following eight groups each contain amino acids that are conservative substitutions for one another: (1) Alanine (A), Glycine (G); (2) Aspartic acid (D), Glutamic acid (E); (3) Asparagine (N), Glutamine (Q); (4) Arginine (R), Lysine (K); (5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); (6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W); (7) Serine (S), Threonine (T); and (8) Cysteine (C), Methionine (M).

When comparing (ribo)nucleic acids, similarity can also be determined. In this case, the similarity is appreciated at the codon level. Two codons are similar, if they are different codons specifying the same amino acid or if they are different codons specifying two similar amino acids.

In the present invention, a homolog of the *ohrR* gene is a gene or a nucleic acid presenting a similarity in the nucleotide sequence of at least 50%, at least 60%, at least 70%, at least 80%, at least 90%

or at least 95% with the *ohrR* gene nucleotide sequence of *Bacillus cereus* ATCC14579.

In addition, a homolog of the *ohrR* gene product is a ribonucleic acid or a polypeptide presenting a
5 similarity in the ribonucleotide or in the amino acid sequence of at least 50%, at least 60%, at least 70%, at least 80%, at least 90% or at least 95% with the *ohrR* gene product ribonucleotide or amino acid sequence of *Bacillus cereus* ATCC14579.

10 Advantageously, in the present invention, a homolog of the *ohrR* gene or of the *ohrR* gene product is a negative regulator belonging to the MarR family and is related to organic peroxide sensing, i.e. which is organic peroxide-inducible. This latter property can be
15 checked by a disk diffusion assay well-known by the one skilled in the art and, particularly, by the disk diffusion assay using H₂O₂ as indicated chemical disclosed by Fuangthong et al., 2001 [10].

As for non exhaustive examples of OhrR homologs
20 identified as previously explained i.e. by two iterations of PSI-BLAST (p value < 1e-10), one can cite the protein of *Corynebacterium pseudogenitalium* ATCC 33035, the accession number of which is ZP_07715254.1 ; the protein of *Pseudomonas syringae* pv. *mori* str.
25 301020, the accession number of which is EGH21983.1 ; the protein of *Streptomyces sviveus* ATCC 29083, the accession number of which is ZP_06917482.1 ; the protein of *Arthrobacter aurescens* TC1, the accession number of which is YP 947034.1 ; the protein of
30 *Myxococcus xanthus* DK1622, the accession number of which is YP_628922.1 ; and the protein of *Clostridium*

botulinum B str. Eklund 17B, the accession number of which is YP_001887666.1.

The step (a) of the method according to the present invention consists in inactivating the *ohrR* gene or a homolog thereof or in inactivating the *ohrR* gene product or a homolog thereof.

Any process known by the one skilled in the art to inactivate a gene or a gene product can be utilized in the present invention. This latter employs routine techniques in the field of molecular biology. Basic texts disclosing the general methods useful for practicing the present invention include Sambrook and Russell, 2001 [11].

As used herein, the expression "inactivation of a gene" covers not only the inactivation *stricto sensu* but also refers to the alteration; partial, substantial, or complete deletion; silence or down-regulation of this gene. The terms and expressions "deletion", "partial deletion", "substantial deletion", "complete deletion", "eliminate", "elimination" and "knockout" can be utilized interchangeably.

In the context of the present invention, the inactivation of *ohrR* gene includes a structural alteration in the *ohrR* gene or locus that reduces or eliminates the production or activity of the *ohrR* gene product, i.e. the RNA or the polypeptide. In another embodiment, the genetically modified microorganism prepared at step (a) includes a structural alteration in a part of the microorganism genome different from the *ohrR* gene or locus that directly or indirectly

reduces or eliminates the production or activity of the *ohrR* gene product, such as a structural alteration in a regulatory region that is not positioned within the *ohrR* gene or locus. By "*ohrR* locus", it must be understood a chromosomal locus comprising the *ohrR* gene.

The structural alteration can include, for example, a deletion, insertion or substitution in the *ohrR* gene, or other region that affects the production of the *ohrR* gene product. In a particularly preferred embodiment, the structural alteration includes a deletion, insertion or substitution in the coding region of the *ohrR* gene or locus. Deletion refers to the elimination of at least one nucleotide in a sequence. Insertion refers to the addition of at least one nucleotide in a sequence. Substitution refers to the change of at least one nucleotide in a sequence. A substitution can create a point mutation. In addition, a deletion, insertion or substitution can create a codon stop in the *ohrR* gene.

For example, the step (a) of the present invention method can provide a genetically modified microorganism that has an *ohrR* gene or locus which has been modified by deletion of a segment of the DNA of the *ohrR* gene or locus or insertion of a foreign DNA in the *ohrR* gene or locus or both.

Concerning the inactivation of the *ohrR* gene, a deletion of all or part of the *ohrR* gene or in the different part of the modified microorganism provides maximum stability because there is no opportunity for a reverse mutation to restore function.

In the present invention, the inactivation of the *ohrR* gene product can be obtained by introducing an exogenous agent into the microorganism to yield a modified microorganism, wherein the exogenous agent reduces or eliminates the production or activity of the *ohrR* gene product. The present invention is not limited by the nature of the exogenous agent supplied. This latter can be an interfering RNA (iRNA), an antisense DNA (asDNA), an antisense RNA (asRNA), double-stranded RNA (dsRNA), a ribozyme, a riboswitch, a DNA aptamer, a RNA aptamer, a peptide aptamer, a molecular inhibitor, a nucleic acid coding thereof or a vector or a plasmid comprising or operably encoding thereof.

A "molecular inhibitor" refers to a natural or synthetic molecule which inhibits the production or activity of the *ohrR* gene product. This molecular inhibitor can be a small organic molecule; a metabolite; a protein or peptide inhibitor such as a linear or cyclic peptide or a peptidomimetic; an antibody or a fragment thereof directed against the polypeptide OhrR or against particular epitope present therein. A fragment of an antibody is advantageously chosen from Fab fragment, F(ab')₂ fragment, Fv fragment, scFv fragment and diabody.

Indeed, a way the activity of the OhrR polypeptide can be reduced is by interfering with the mRNA transcription product of the *ohrR* gene. For example, an exogenous agent such as a ribozyme or an expression vector operably encoding a ribozyme can be delivered to the microorganism to cleave the target mRNA, thereby reducing or eliminating production of the

polypeptide. AsDNA, asRNA and dsRNA may also be used to interfere with translation.

In addition, one can inactivate the mRNA transcription product of the *ohrR* gene by adding a
5 riboswitch before the translation start. Indeed, riboswitch RNA are able to sense specific metabolites and accordingly regulate production of proteins (see [12] and [13]).

10 Alternatively, the OhrR polypeptide or the OhrR protein can also be the target of the inactivation. Antibodies or antibody-like molecules such as peptide aptamers can be introduced into the microorganism to abolish the activity of the translated polypeptide, as
15 can protein or peptide inhibitors such as linear or cyclic peptides, peptidomimetics, or small organic molecules.

The exogenous agent can be delivered or introduced into the microorganism by any routine
20 methods known by the one skilled in the art such as an electroporation, a heat shock or/and a chemical process.

All the above explanations and details given for the inactivation of the *ohrR* gene or of the *ohrR*
25 gene product apply to a gene homologous to the *ohrR* gene or to the gene product of a gene homologous to the *ohrR* gene *mutatis mutandis*. As a consequence, the expressions used hereinafter "*ohrR* gene (homolog)" means either "*ohrR* gene" or "*ohrR* gene homolog" and
30 "*ohrR* gene (homolog) product" means either "*ohrR* gene product" or "*ohrR* gene homolog product".

The step (b) of the method according to the present invention consists in culturing the modified microorganism in particular culture conditions for the phenotype due to said modification (growth rate, biomass production and/or fermentative potential enhancement) to be expressed. These particular culture conditions are low-ORP anaerobic conditions.

The expression "growth conditions" and "culture conditions" are equivalent and can be used interchangeably. Similarly, the expression "growth medium" and "culture medium" are equivalent and can be used interchangeably.

The expression "anaerobic conditions" as used herein, means growth conditions in the absence of oxygen and notably refers to the absence of oxygen in the culture medium. In other words, the set-point dissolved oxygen tension value (pO_2) is below 0.2% in growth medium and, in particular, of 0% in growth medium. A pO_2 below 0.2% takes into account the traces of oxygen that can always be found in growth medium even under anaerobic conditions. Such pO_2 values can be achieved by continuously flushing the medium with either pure N_2 gas or with pure H_2 gas, optionally previously filtered.

The expressions "low-ORP", "low oxidation-reduction potential", "low oxidoreduction potential" and "low redox potential" are equivalent and can be used interchangeably. As used herein, the ORP and notably the low-ORP means the initial ORP and notably the initial low-ORP i.e. the ORP measured in the

culture medium just before seeding or inoculating the modified microorganism in said medium. The ORP of a culture medium can be measured by any process known by the one skilled in the art. This measurement can
5 utilize a sensor immersed in the culture medium or a redox-combined electrode such as the redox-combined electrode AgCl of Mettler Toledo and the measured values have to be corrected according to the reference electrode value, both electrodes being immersed in the
10 culture medium.

In addition, as used herein, a low-ORP means a ORP inferior or equal to -100 mV, notably inferior or equal to -200 mV and, in particular, inferior or equal to -300 mV. A low-ORP can be obtain in the growth
15 medium by introducing in said medium any known reducing agent in a appropriate quantity the one skilled in the art can calculate or determine by iteration. Various compounds may be used as a reducing agent. Among these, mention may be made of sulfites, sodium thioglycolate,
20 dithiothreitol, ascorbic acid (vitamin C) and ammonia but also of reducing gases such as hydrogen or gaseous mixtures containing hydrogen.

In step (b) of the method according to the
25 present invention, the modified microorganism can be cultured in a medium according to the usual method used for culturing microorganisms. This medium can be a commercial medium or an extemporaneously prepared medium.

As a medium used for culturing, the general nutritional medium containing a carbon source, a nitrogen source and inorganic salts can be used.

Any carbon source which can be assimilated by the modified microorganism is used. Examples include carbohydrates such as glucose, fructose, sucrose, molasses containing them; starch or starch hydrolysate; cellulose, lignocellulose or hemicellulose; one-carbon substrates; organic acids such as acetic acid or propionic acid; and alcohols such as ethanol and propanol.

Examples of suitable nitrogen source include ammonia; ammonium salts of inorganic acids or organic acids such as ammonium chloride, ammonium sulfate, ammonium acetate and ammonium phosphate; other nitrogen-containing compounds; peptone; meat extract; yeast extract; corn steep liquor; casein hydrolysate; soybean meal and soybean meal hydrolysate; and various cells obtained by fermentation and their digested products.

Examples of suitable inorganic salts include potassium dihydrogen phosphate, dipotassium hydrogen phosphate, magnesium phosphate, magnesium sulfate, sodium chloride, ferrous sulfate, manganese sulfate, copper sulfate and calcium carbonate.

The culturing temperature is typically from 15 to 40°C, and the pH during the culturing is typically maintained at 3.0 to 9.0. The pH can be adjusted using, for example, an inorganic or organic acid, an alkali solution, urea, calcium carbonate or ammonia. It must be pointed out that these general culture conditions

can be slightly modified by the one skilled in the art relative to the microorganism cultured.

Advantageously, the cultured modified microorganisms are preferably in a suspension state. A
5 "suspension" of cells is to be broadly understood as including all types of suspended or dispersed cell cultures; the term "suspension state" is thus used to distinguish cells that are not cultured in a liquid medium, such as cells cultured by way of adhering on a
10 Petri dish. Thus, the term "suspension" includes both freely dispersed cells and agglomerated cells, regardless of whether agglomeration occurs spontaneously.

In addition, culture methods that can be
15 utilized in the step (b) of the method according to the present invention include, but are not limited to, a batch culture, a continuous culture, or a "fed-batch" culture.

A "continuous (cell) culture" is a cell culture
20 characterized by both a continuous inflow of a liquid nutrient feed and a continuous liquid outflow. Alternatively, a continuous culture may constitute a "perfusion culture" in which case the liquid outflow contains culture medium that is substantially free of
25 cells, or substantially lower cell concentration than that in the bioreactor. In a perfusion culture, cells can be retained by, for example, filtration, centrifugation, or sedimentation.

A "fed-batch" culture is a batch cell culture
30 to which substrate, in either solid or concentrated liquid form, is added either periodically or

continuously during the run. Just as in a batch culture, a fed-batch culture is initiated by inoculating cells to the medium, but, in contrast to a batch culture, there is a subsequent inflow of nutrients, such as by way of a concentrated nutrient feed. In contrast to a continuous culture there is no systematic removal of culture fluid or cells from a fed-batch culture.

Thus, the present invention concerns a method for enhancing the growth rate, the biomass production and/or the fermentative potential of an anaerobic microorganism, comprising the steps consisting in :

a) modifying said anaerobic microorganism by inactivating the *ohrR* gene, the *ohrR* gene product or homolog thereof,

a homolog to *ohrR* gene or to the *ohrR* gene product being a gene coding for an amino acid sequence, an amino acid sequence or a RNA coding for an amino acid sequence, wherein said amino acid sequence is detected using the OhrR amino acid sequence from *Bacillus cereus* by two iterations of Position Specific Iterated-BLAST (PSI-BLAST) (p value < 1e-10) ;

b) culturing the modified anaerobic microorganism obtained in step (a) under low-ORP anaerobic conditions with a ORP inferior or equal to -100 mV ;

thereby the growth rate, the biomass production and/or the fermentative potential of said modified microorganism is enhanced when compared with the growth rate, the biomass production and/or the fermentative

potential of the corresponding non modified microorganism cultured under the same conditions.

The present invention provides a novel method
5 for enhancing the anaerobic growth rate, the biomass production and/or the fermentative potential of microorganisms. These results with this increase can make it possible to obtain better yields for various and numerous industrial processes based on the use of a
10 modified microorganism as previously defined in low-ORP anaerobic culture conditions.

Fields of application of the present invention are covering industrial processes related to human and animal health; agronomy and farming, agro-industries;
15 marine and aquatic processes; chemical industry; energy industry and environmental processes.

As a consequence, the present invention concerns the use of a modified microorganism as previously defined, in low-ORP anaerobic culture
20 conditions, in an industrial process related to human and animal health; agronomy and farming, agro-industries; marine and aquatic processes; chemical industry; energy industry and environmental processes.

More particularly, the present invention
25 concerns the use of a modified microorganism as previously defined, in low-ORP anaerobic culture conditions, in the production of food products ; in the production of biofuels ; in the pharmaceutical field to produce active ingredients; in the production of
30 valuable chemicals or in the destruction of hazardous and/or polluting chemicals.

Other characteristics and advantages of the present invention will additionally be apparent to the one skilled in the art on reading the examples below, which are given as an illustration and not a limitation, with reference to the attached figures.

SHORT DESCRIPTION OF THE DRAWINGS

Figure 1 presents the physiological parameters for controlled batch cultures of the $\Delta ohrR$ mutant and its parent strain *B. cereus* ATCC14579 (wt).

Figure 2 presents the specific uptake of glucose of the $\Delta ohrR$ mutant and its parent strain *B. cereus* ATCC14579 (wt).

DETAILED DISCUSSION OF PARTICULAR EMBODIMENTS

I. Construction of $\Delta ohrR$ mutant.

B. cereus strains implemented in the experimental part are ATCC 14579 strains.

The $\Delta ohrR$ cell mutant was constructed as disclosed hereinafter.

A *Bgl*II-SalI DNA fragment of 1,489 bp encompassing the *ohrA* and *ohrR* ORFs was amplified by PCR using chromosomal DNA as template and the primer pairs *ohrmutF* (5'-AGATCTTCTTATGAATCTGACAATCGGG-3', SEQ ID NO: 2 in the appended sequence listing) plus *ohrmutR* (5'-GTCGACCCGATAAAATTCCGAAAGGG-3', SEQ ID NO: 3 in the appended sequence listing).

The amplified DNA fragment was cloned into pCRXL-TOPO (Invitrogen). The resulting pCRXLohr plasmid was then digested with *HpaI* and *BsgI*. A 1.5 kb *SmaI* fragment containing the entire spectinomycin resistance gene *spc* was purified from the pDIA plasmid. This purified DNA fragment was ligated into *BsgI*-digested pCRXLohr. The resulting plasmid was digested with *EcoRI* and *BglIII*. The resulting *ohrR-spc* fragment was subsequently inserted between the corresponding pMAD sites.

The resulting plasmid was introduced into *B. cereus* strain by electroporation. The *ohrR* gene was deleted by a double crossover event.

Chromosomal allele exchange was confirmed by PCR with oligonucleotide primers located upstream and downstream of the DNA regions used for allelic exchange. To confirm the nonpolar effect of the spectinomycin resistance cassette insertion in *ohrR* (which precedes *ohrA* in the *ohrR-ohrA* operon), fragments corresponding partially to *ohrA* and *ohrA-ohrR* intergenic region mRNA were amplified from total RNA isolated from Δ *ohrR* cells as described above, thus confirming correct expression of the gene.

25 II. Analytical procedures and physiological parameters.

B. cereus growth was monitored spectrophotometrically at 560 nm and calibrated against cell dry weight measurements, as previously described. Specific growth rate (μ) was determined using the modified Gompertz equation. Glucose, lactate, ethanol,

formate, acetate and succinate concentrations were determined using commercial enzymatic kits (Diffchamb, Lyon, France; R-Biopharm, Saint- Didier au Mont-d'Or, France; Roche, Meylan, France).

5 Specific glucose consumption rate, defined as the differential change in glucose concentration with time, was calculated from the equation $q_{\text{glucose}} = \mu/Y_x$, where μ is the specific growth rate [h^{-1}] and Y_x , the biomass yield [$\text{g. mol carbon substrate}^{-1}$].

10

III. Growth conditions.

Wild-type and ΔohrR *B. cereus* ATCC 14579 strains were grown at 37°C in synthetic MOD medium supplemented with 30 mM glucose as carbon source as
15 previously described [14].

The pH was maintained at 7.2 by automatic addition of 2N KOH. The temperature and pH regulated batch was equipped with a Mettler Toledo polarographic oxygen electrode coupled with feedback regulation to
20 maintain the set-point dissolved oxygen tension value ($p\text{O}_2$) via air sparging and agitation speed. The bioreactor was spurge with air alone to set a $P\text{O}_2$ value of 100%. A $p\text{O}_2$ of 0% was obtained by continuously flushing the medium at 20 ml/h with either pure N_2 gas
25 or with pure H_2 gas previously filtered through a Hungate column.

Oxidoreduction potential (ORP) was measured using a redox-combined electrode (AgCl, Mettler Toledo) and the values were corrected according to the
30 reference electrode value ($E_{\text{ref}} = + 200 \text{ mV}$ at 37°C).

Each bioreactor was inoculated with a subculture grown overnight under aerobiosis or anaerobiosis in MOD medium supplemented with carbon source and the required electron acceptors. A sample of
5 the overnight culture was diluted in fresh medium to achieve an initial optical density of the culture at 560 nm of 0.02. Cells were harvested by centrifugation at the early-exponential growth phase, i.e. 1.5 h after reaching the maximal growth rate (when $\mu = 80 (\pm 10) \%$
10 of $\mu_{\max}$,) and immediately frozen at -80°C until analysis. Supernatants were kept to perform metabolite and glucose assays.

IV. Results.

15 The disruption of *ohrR* led to a significant increase (~1.6-fold) of the growth rate of fermentative cells under low-ORP conditions compared with wild-type cells (Table 1 hereinafter ; Figure 1).

No significant differences between the two
20 strains could be observed under high-ORP anaerobiosis and aerobiosis. The higher growth rate under low-ORP conditions results from a higher carbon flow through glycolysis, as reflected by the higher glucose uptake rate (Figure 2) and subsequently from a higher ATP
25 production (Table 1).

Table 1: Physiological parameters for controlled batch cultures of the $\Delta ohrR$ mutant and its parent strain *B. cereus* ATCC14579 (wt).

Growth	Anaerobic fermentative				Aerobic	
iORP	- 410 mV		- 10 mV		+ 140 mV	
Strain	WT	$\Delta ohrR$	WT	$\Delta ohrR$	WT	$\Delta ohrR$
$\mu_{\max} (h^{-1})$	0.75 ± 0.05	1.17 ± 0.01	0.80 ± 0.07	0.86 ± 0.02	1.26 ± 0.12	1.22 ± 0.18
Y_x	21 ± 1	21 ± 1	23 ± 1	23 ± 1	8 ± 1	8 ± 1
q_{glucose}	35 ± 2	55 ± 5	35 ± 2	38 ± 2	16 ± 2	15 ± 2
Y_{Lactate}	0.68 ± 0.05	0.66 ± 0.15	0.48 ± 0.12	0.63 ± 0.15	nz	nz
Y_{Acetate}	0.12 ± 0.02	0.20 ± 0.04	0.14 ± 0.04	0.15 ± 0.04	0.14 ± 0.01	0.20 ± 0.04
Y_{Formate}	0.05 ± 0.01	0.13 ± 0.07	0.04 ± 0.02	0.05 ± 0.03	nz	nz
Y_{Ethanol}	0.070 ± 0.005	0.10 ± 0.02	0.054 ± 0.006	0.07 ± 0.01	nz	nz
Y_{ATP}	1.98	1.62	2.00	nd	nd	nd
NADH recovery	1.2	1.2	1.5	1.5	nd	nd

Cells were grown under H_2 and N_2 -generated anaerobiosis (iORP = -410 mV and -10 mV, respectively; pO_2 = 0%) and full aerobiosis (iORP = +140 mV; pO_2 = 100%;). Data are the means of triplicate measures obtained from three independent cultures.

Abbreviations: $\mu_{\max}$, specific growth rate determined by measurement of the dry weight ; Y_x , biomass yield on glucose (gram of biomass per gram of glucose consumed); q_{glucose} , millimoles of glucose consumed per milligram of biomass per hour; Y_{lactate} , Y_{acetate} , Y_{ethanol} and Y_{formate} , moles of metabolite produced per moles of glucose consumed measured at $\mu_{\max}$; **nz**, near zero; **nd**, not determined. NADH recovery was calculated at the ratio of pathways producing NADH versus those consuming NADH (producing NAD^+).

The data obtained by the inventors also indicate that disruption of *ohrR* allows i) lactate production to reach its apogee in terms of quantity under high-ORP fermentative conditions and ii) formate, ethanol and acetate to reach their apogee in terms of quantity under low-ORP conditions.

The high-throughput proteomic data obtained by the inventors (not described here) point at i) the role of OhrR in proteome remodeling when oxygen is lacking, and more specifically when anoxic-ORP is decreasing and, ii) its regulatory impact on activity of main central metabolism pathways. Through the control of proteome change, OhrR strangles the fermentative potential of *B. cereus* and restricts cell growth under low-ORP anaerobiosis while ensuring a high synthesis of chemotaxis and motility proteins. Inactivation of the *ohrR* gene or its product results in higher fermentative potential under these conditions.

REFERENCES

1. International application WO 2009/101328 in the name of L'AIR LIQUIDE SOCIETE ANONYME POUR L'ETUDE
5 ET L'EXPLOITATION DES PROCEDES GEORGES CLAUDE and published on August 20, 2009.
2. Patent application FR 2 635 788 in the names of Institut Pasteur and Centre National de la Recherche
10 Scientifique and published on March 2, 1990.
3. Patent application FR 2 778 922 in the name of Institut Pasteur and published on November 26, 1999.
- 15 4. International application WO 2011/018116 in the name of ALGENOL BIOFUELS Inc. and published on February 17, 2011.
- 20 5. International application WO 2011/006136 in the name of VERDEZYNE Inc. and published on January 13, 2011.
- 25 6. Article by Duport *et al.*, 2006, J. Bacteriol., vol. 188, pages 6640-6651.
7. Article by Gao *et al.*, 2010, Protein Cell, vol. 1, pages 621-626.
- 30 8. Article by Gibson *et al.*, 2010, Science, vol. 329, pages 52-56.

9. Article by Duarte and Latour, 2010, Molecular BioSystems, vol. 6, pages 316-323.

10. Article by Fuangthong *et al.*, 2001, J. Bacteriol., vol. 183, pages 4134-4141.

11. Sambrook and Russell, 2001, Molecular Cloning, A Laboratory Manual (3rd edition).

10. 12. Article by Dambach and Winkler, 2009, Curr. Opin. Microbiol., vol. 12, pages 161-169.

13. Article by Henkin, 2008, Genes Dev., vol. 22, pages 3383-3390.

15

14. Article by Clair *et al.*, 2010, Mol. Cell. Proteomics, vol. 9, pages 1486-1498.

CLAIMS

1) Method for enhancing the growth rate, the biomass production and/or the fermentative potential of an anaerobic microorganism, comprising the steps consisting in :

a) modifying said anaerobic microorganism by inactivating the *ohrR* gene, the *ohrR* gene product or homolog thereof,

10 a homolog to *ohrR* gene or to the *ohrR* gene product being a gene or a RNA coding for an amino acid sequence or an amino acid sequence, wherein said amino acid sequence is detected using the OhrR amino acid sequence from *Bacillus cereus* by two iterations of Position Specific Iterated-BLAST (PSI-BLAST) (p value < 1e-10) ;

b) culturing the modified anaerobic microorganism obtained in step (a) under low-ORP anaerobic conditions with a ORP inferior or equal to -100 mV ;

20 thereby the growth rate, the biomass production and/or the fermentative potential of said modified microorganism is enhanced when compared with the growth rate, the biomass production and/or the fermentative potential of the corresponding non modified microorganism cultured under the same conditions.

2) Method according to claim 1, wherein a homolog of the *ohrR* gene or of the *ohrR* gene product is a negative regulator belonging to the MarR family and is related to organic peroxide sensing.

30

3) Method according to claim 1 or 2, wherein said anaerobic microorganism is a facultative anaerobic microorganism or an obligate anaerobic microorganism.

5

4) Method according to any one of claims 1 to 3, wherein said anaerobic microorganism is a natural, genetically-modified or synthetic, eukaryotic or prokaryotic, anaerobic microorganism.

10

5) Method according to any one of claims 1 to 4, wherein a homolog of the *ohrR* gene is a gene or a nucleic acid presenting an identity in the nucleotide sequence of at least 20%, at least 30%, at least 40%,
15 at least 50%, at least 60%, at least 70%, at least 80%, at least 90% or at least 95% with the *ohrR* gene nucleotide sequence of *Bacillus cereus* ATCC14579.

6) Method according to any one of claims 1 to
20 4, wherein a homolog of the *ohrR* gene product is a ribonucleic acid or a polypeptide presenting an identity in the ribonucleotide or in the amino acid sequence of at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%,
25 at least 90% or at least 95% with the *ohrR* gene product ribonucleotide or amino acid sequence of *Bacillus cereus* ATCC14579.

7) Method according to any one of claims 1 to
30 6, wherein the inactivation of said *ohrR* gene (homolog) includes a structural alteration in the *ohrR* gene

(homolog) or locus that reduces or eliminates the production or activity of the *ohrR* gene (homolog) product.

5 8) Method according to any one of claims 1 to 6, wherein the inactivation of said *ohrR* gene (homolog) includes a structural alteration in a part of the microorganism genome different from the *ohrR* gene (homolog) or locus that directly or indirectly reduces
10 or eliminates the production or activity of the *ohrR* gene (homolog) product.

 9) Method according to any one of claims 1 to 6, wherein the inactivation of said *ohrR* gene (homolog)
15 product is obtained by introducing an exogenous agent into the microorganism to yield a modified microorganism, wherein the exogenous agent reduces or eliminates the production or activity of the *ohrR* gene (homolog) product.

20

 10) Method according to claim 9, wherein said exogenous agent is chosen from an interfering RNA (iRNA), an antisense DNA (asDNA), an antisense RNA (asRNA), double-stranded RNA (dsRNA), a ribozyme, a
25 riboswitch, a DNA aptamer, a RNA aptamer, a peptide aptamer, a molecular inhibitor, a nucleic acid coding thereof or a vector or a plasmid comprising or operably encoding thereof.

30 11) Method according to any one of the preceding claims, wherein said ORP is inferior or equal

to -200 mV and, in particular, inferior or equal to -300 mV.

12) Use of a modified microorganism as defined
5 in any one of claims 1, 3 or 4, in low-ORP anaerobic culture conditions with a ORP inferior or equal to -100 mV, in an industrial process related to human and animal health; agronomy and farming, agro-industries; marine and aquatic processes; chemical industry; energy
10 industry and environmental processes.

13) Use of a modified microorganism as defined
in any one of claims 1, 3 or 4, in low-ORP anaerobic culture conditions with a ORP inferior or equal to
15 -100 mV, in the production of food products ; in the production of biofuels ; in the pharmaceutical field to produce active ingredients; in the production of valuable chemicals or in the destruction of hazardous and/or polluting chemicals.




FIG.1




FIG.2

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/060624

A. CLASSIFICATION OF SUBJECT MATTER
INV. C12N1/20 C07K14/195 C07K14/32
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	CLAIR GÉRÉMY ET AL: "Restricting Fermentative Potential by Proteome Remodeling: AN ADAPTIVE STRATEGY EVIDENCED IN BACILLUS CEREUS.", MOLECULAR & CELLULAR PROTEOMICS : MCP JUN 2012 LNKD- PUBMED:22232490, vol. 11, no. 6, 9 January 2012 (2012-01-09), XP009161847, ISSN: 1535-9484 the whole document ----- -/--	1-13



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

10 August 2012

Date of mailing of the international search report

21/08/2012

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Giebeler, Katharina

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/060624

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ARNTJEN B ET AL: "Impact of MarA Point Mutations on Growth Rate and Susceptibilities to Different Antibiotics of in vitro Generated Mutants of E. coli with Mar Phenotype", INTERNATIONAL JOURNAL OF MEDICAL MICROBIOLOGY, URBAN UND FISCHER, DE, vol. 299, no. Suppl. 1, 1 September 2009 (2009-09-01), page 82, XP009153673, ISSN: 1438-4221 the whole document	1-13
A	D. A. CANO ET AL: "Salmonella enterica Serovar Typhimurium Response Involved in Attenuation of Pathogen Intracellular Proliferation", INFECTION AND IMMUNITY, vol. 69, no. 10, 1 October 2001 (2001-10-01), pages 6463-6474, XP055011125, ISSN: 0019-9567, DOI: 10.1128/IAI.69.10.6463-6474.2001 the whole document abstract	1-13
A	WO 2007/090123 A2 (UNIV CHICAGO [US]; CHEN PENG; HE CHUAN) 9 August 2007 (2007-08-09) the whole document page 6, line 5 - line 7	1-13
A	WO 2007/011845 A2 (BASF AG [DE]; ZELDER OSKAR [DE]; HEROLD ANDREA [DE]; KLOPPROGGE CORINN) 25 January 2007 (2007-01-25) the whole document page 32, line 1 - line 2	1-13
A	ELLISON D W ET AL: "Regulation of virulence by members of the MarR/SlyA family", CURRENT OPINION IN MICROBIOLOGY, CURRENT BIOLOGY LTD, GB, vol. 9, no. 2, 1 April 2006 (2006-04-01), pages 153-159, XP025174542, ISSN: 1369-5274, DOI: 10.1016/J.MIB.2006.02.003 [retrieved on 2006-04-01] the whole document	1-13
	----- -/--	

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2012/060624

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	M. FUANGTHONG ET AL: "OhrR Is a Repressor of ohrA, a Key Organic Hydroperoxide Resistance Determinant in Bacillus subtilis", JOURNAL OF BACTERIOLOGY, vol. 183, no. 14, 15 July 2001 (2001-07-15), pages 4134-4141, XP055011126, ISSN: 0021-9193, DOI: 10.1128/JB.183.14.4134-4141.2001 the whole document	1-13
A	----- MINSUN HONG ET AL: "Structure of an OhrR-ohrA Operator Complex Reveals the DNA Binding Mechanism of the MarR Family", MOLECULAR CELL, vol. 20, no. 1, 1 October 2005 (2005-10-01), pages 131-141, XP055011130, ISSN: 1097-2765, DOI: 10.1016/j.molcel.2005.09.013 the whole document	1-13
A	----- S. SOONSANGA ET AL: "Mutational Analysis of Active Site Residues Essential for Sensing of Organic Hydroperoxides by Bacillus subtilis OhrR", JOURNAL OF BACTERIOLOGY, vol. 189, no. 19, 1 October 2007 (2007-10-01), pages 7069-7076, XP055011132, ISSN: 0021-9193, DOI: 10.1128/JB.00879-07 the whole document	1-13
A	----- DUARTE VICTOR ET AL: "PerR vs OhrR: selective peroxide sensing in Bacillus subtilis", MOLECULAR BIOSYSTEMS, ROYAL SOCIETY OF CHEMISTRY, GB, vol. 6, no. 2, 1 January 2010 (2010-01-01), pages 316-323, XP009153352, ISSN: 1742-206X cited in the application the whole document	1-13
A	----- ROBERT G. MARTIN ET AL: "Transcriptional and translational regulation of the marRAB multiple antibiotic resistance operon in Escherichia coli", MOLECULAR MICROBIOLOGY, vol. 53, no. 1, 1 July 2004 (2004-07-01), pages 183-191, XP055011134, ISSN: 0950-382X, DOI: 10.1111/j.1365-2958.2004.04080.x the whole document	1-13
	----- -/--	

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2012/060624

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	G. CLAIR ET AL: "Expanding the Known Repertoire of Virulence Factors Produced by <i>Bacillus cereus</i> through Early Secretome Profiling in Three Redox Conditions", MOLECULAR & CELLULAR PROTEOMICS, vol. 9, no. 7, 1 July 2010 (2010-07-01), pages 1486-1498, XP055011137, ISSN: 1535-9476, DOI: 10.1074/mcp.M000027-MCP201 cited in the application the whole document -----	1-13

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP2012/060624

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:
 - a. (means)
☐ on paper
☒ in electronic form
 - b. (time)
☒ in the international application as filed
☐ together with the international application in electronic form
☐ subsequently to this Authority for the purpose of search
2. ☐ In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2012/060624

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2007090123 A2	09-08-2007	US 2007248581 A1	25-10-2007
		WO 2007090123 A2	09-08-2007

WO 2007011845 A2	25-01-2007	CA 2615419 A1	25-01-2007
		EP 1907557 A2	09-04-2008
		JP 2009501547 A	22-01-2009
		US 2009221027 A1	03-09-2009
		WO 2007011845 A2	25-01-2007
