



(86) Date de dépôt PCT/PCT Filing Date: 2004/10/15
(87) Date publication PCT/PCT Publication Date: 2005/05/06
(45) Date de délivrance/Issue Date: 2013/06/04
(85) Entrée phase nationale/National Entry: 2006/04/11
(86) N° demande PCT/PCT Application No.: EP 2004/011630
(87) N° publication PCT/PCT Publication No.: 2005/039548
(30) Priorité/Priority: 2003/10/15 (US60/512,038)


(51) Cl.Int./Int.Cl. *A61K 31/00* (2006.01),
A61K 31/13 (2006.01), *A61P 25/28* (2006.01),
A61P 35/00 (2006.01)

(72) Inventeurs/Inventors:
SCHILLING, STEPHAN, DE;
HOFFMANN, TORSTEN, DE;
NIESTROJ, ANDRE JOHANNES, DE;
DEMUTH, HANS-ULRICH, DE;
HEISER, ULRICH, DE

(73) Propriétaire/Owner:
PROBIODRUG AG, DE

(74) Agent: NORTON ROSE CANADA
S.E.N.C.R.L.,S.R.L./LLP

(54) Titre : UTILISATION D'INHIBITEURS DE LA GLUTAMYL CYCLASE DANS LE TRAITEMENT DE LA DEMENCE
BRITANNIQUE FAMILIALE ET DE LA DEMENCE DANOISE FAMILIALE
(54) Title: USE OF GLUTAMYL CYCLASE INHIBITORS IN THE TREATMENT OF FAMILIAL BRITISH DEMENTIA AND
FAMILIAL DANISH DEMENTIA



formula 1

(57) Abrégé/Abstract:

The present invention provides novel physiological substrates of mammalian glutaminyl cyclase (QC, EC 2.3.2.5), new effectors of QC and the use of such effectors and pharmaceutical compositions comprising such effectors for the treatment of diseases that

(57) **Abrégé(suite)/Abstract(continued):**


can be treated by modulation of QC-activity, e.g. diseases selected from the group consisting of duodenal cancer with or w/o *Helicobacter pylori* infections, colorectal cancer, Zollinger-Ellison syndrome, Familial British Dementia and Familial Danish Dementia. (see formula 1)

ABSTRACT

5

The present invention provides novel physiological substrates of mammalian glutaminy cyclase (QC, EC 2.3.2.5), new effectors of QC and the use of such effectors and pharmaceutical compositions comprising such effectors for the treatment of diseases that can be treated by modulation of QC-activity, e.g. diseases selected from the group consisting of duodenal cancer with or w/o *Helicobacter pylori* infections, colorectal cancer, Zollinger-Ellison syndrome, Familial British Dementia and Familial Danish Dementia.

15



USE OF GLUTAMYL CYCLASE INHIBITORS IN THE TREATMENT OF FAMILIAL BRITISH DEMENTIA AND FAMILIAL DANISH DEMENTIA

Field of the invention

5 The invention relates to glutamyl cyclase (QC, EC 2.3.2.5) that catalyzes the intramolecular cyclization of N-terminal glutamine residues into pyroglutamic acid (5-oxo-proline, pGlu*) under liberation of ammonia and the intramolecular cyclization of N-terminal glutamate residues into pyroglutamic acid under liberation of water.

10 The present invention identifies mammalian QCs as metalloenzymes, provides novel physiological substrates of QC in mammals, novel effectors of QC and the use of effectors of QC and pharmaceutical compositions comprising effectors of QC for the treatment of conditions that can be treated by modulation of QC-activity. Additionally, it is shown that metal interaction is a useful approach for development of QC inhibitors.

15 In a preferred embodiment, the present invention provides the use of effectors of QC activity in combination with inhibitors of DP IV or DP IV-like enzymes for the treatment or alleviation of conditions that can be treated by modulation of QC- and DP IV-activity.

20 A screening method is also provided for the identification and selection of effectors of QC activity.

Background

25 Glutamyl cyclase (QC, EC 2.3.2.5) catalyzes the intramolecular cyclization of N-terminal glutamine residues into pyroglutamic acid (pGlu*) liberating ammonia. A QC was first isolated by Messer from the latex of the tropical plant *Carica papaya* in 1963 (Messer, M. 1963 *Nature* 4874, 1299). 24 years later, a corresponding enzymatic activity was discovered in animal pituitary (Busby, W. H. J. et al. 1987 *J Biol Chem* 30 262, 8532-8536; Fischer, W. H. and Spiess, J. 1987 *Proc Natl Acad Sci U S A* 84, 3628-3632). For the mammalian QC, the conversion of Gln into pGlu by QC could be shown for the precursors of TRH and GnRH (Busby, W. H. J. et al. 1987 *J Biol Chem*

262, 8532-8536; Fischer, W. H. and Spiess, J. 1987 *Proc Natl Acad Sci U S A* 84, 3628-3632). In addition, initial localization experiments of QC revealed a co-localization with its putative products of catalysis in bovine pituitary, further improving the suggested function in peptide hormone synthesis (Bockers, T. M. et al. 1995 *J Neuroendocrinol* 7, 445-453). In contrast, the physiological function of the plant QC is less clear. In case of the enzyme from *C. papaya*, a role in the plant defense against pathogenic microorganisms was suggested (El Moussaoui, A. et al. 2001 *Cell Mol Life Sci* 58, 556-570). Putative QCs from other plants were identified by sequence comparisons recently (Dahl, S. W. et al. 2000 *Protein Expr Purif* 20, 27-36). The physiological function of these enzymes, however, is still ambiguous.

The QCs known from plants and animals show a strict specificity for L-Glutamine in the N-terminal position of the substrates and their kinetic behavior was found to obey the Michaelis-Menten equation (Pohl, T. et al. 1991 *Proc Natl Acad Sci U S A* 88, 10059-10063; Consalvo, A. P. et al. 1988 *Anal Biochem* 175, 131-138; Gololobov, M. Y. et al. 1996 *Biol Chem Hoppe Seyler* 377, 395-398). A comparison of the primary structures of the QCs from *C. papaya* and that of the highly conserved QC from mammals, however, did not reveal any sequence homology (Dahl, S. W. et al. 2000 *Protein Expr Purif* 20, 27-36). Whereas the plant QCs appear to belong to a new enzyme family (Dahl, S. W. et al. 2000 *Protein Expr Purif* 20, 27-36), the mammalian QCs were found to have a pronounced sequence homology to bacterial aminopeptidases (Bateman, R. C. et al. 2001 *Biochemistry* 40, 11246-11250), leading to the conclusion that the QCs from plants and animals have different evolutionary origins.

EP 02 011 349.4 discloses polynucleotides encoding insect glutaminyl cyclase, as well as polypeptides encoded thereby. This application further provides host cells comprising expression vectors comprising polynucleotides of the invention. Isolated polypeptides and host cells comprising insect QC are useful in methods of screening for agents that reduce glutaminyl cyclase activity. Such agents are described as useful as pesticides.

Alzheimer's disease (AD) is characterized by abnormal accumulation of extracellular amyloidotic plaques closely associated with dystrophic neurones, reactive astrocytes

and microglia (Terry, R. D. and Katzman, R. 1983 *Ann Neurol* 14, 497-506; Glenner, G. G. and Wong, C. W. 1984 *Biochem Biophys Res Comm* 120, 885-890; Intagaki, S. et al. 1989 *J Neuroimmunol* 24, 173-182; Funato, H. et al. 1998 *Am J Pathol* 152, 983-992; Selkoe, D. J. 2001 *Physiol Rev* 81, 741-766). Amyloid- β ($A\beta$) peptides are
5 the primary components of senile plaques and are considered to be directly involved in the pathogenesis and progression of AD, a hypothesis supported by genetic studies (Glenner, G. G. and Wong, C. W. 1984 *Biochem Biophys Res Comm* 120, 885-890; Borchelt, D. R. et al. 1996 *Neuron* 17, 1005-1013; Lemere, C. A. et al. 1996 *Nat Med* 2, 1146-1150; Mann, D. M. and Iwatsubo, T. 1996 *Neurodegeneration* 5, 115-120; Citron, M. et al. 1997 *Nat Med* 3, 67-72; Selkoe, D. J. 2001 *Physiol Rev* 81, 741-766). $A\beta$ is generated by proteolytic processing of the β -amyloid precursor protein (APP) (Kang, J. et al. 1987 *Nature* 325, 733-736; Selkoe, D. J. 1998 *Trends Cell Biol* 8, 447-453), which is sequentially cleaved by β -secretase at the N-terminus and by γ -secretase at the C-terminus of $A\beta$ (Haass, C. and Selkoe, D. J. 1993 *Cell* 15 75, 1039-1042; Simons, M. et al. 1996 *J Neurosci* 16 899-908). In addition to the dominant $A\beta$ peptides starting with L-Asp at the N-terminus ($A\beta$ 1-42/40), a great heterogeneity of N-terminally truncated forms occurs in senile plaques. Such shortened peptides are reported to be more neurotoxic *in vitro* and to aggregate more rapidly than the full-length isoforms (Pike, C. J. et al. 1995 *J Biol Chem* 270 20 23895-23898). N-truncated peptides are known to be overproduced in early onset familial AD (FAD) subjects (Saido, T. C. et al. 1995 *Neuron* 14, 457-466; Russo, C. et al. 2000 *Nature* 405, 531-532), to appear early and to increase with age in Down's syndrome (DS) brains (Russo, C. et al. 1997 *FEBS Lett* 409, 411-416, Russo, C. et al. 2001 *Neurobiol Dis* 8, 173-180; Tekirian, T. L. et al. 1998 *J Neuropathol Exp Neurol* 25 57, 76-94). Finally, their amount reflects the progressive severity of the disease (Russo, C. et al. 1997 *FEBS Lett* 409, 411-416). Additional post-translational processes may further modify the N-terminus by isomerization or racemization of the aspartate at position 1 and 7 and by cyclization of glutamate at residues 3 and 11. Pyroglutamate-containing isoforms at position 3 [pGlu³] $A\beta$ (3-40/42) represent the
30 prominent forms – approximately 50 % of the total $A\beta$ amount – of the N-truncated species in senile plaques (Mori, H. et al. 1992 *J Biol Chem* 267, 17082-17086, Saido, T. C. et al. 1995 *Neuron* 14, 457-466; Russo, C. et al. 1997 *FEBS Lett* 409, 411-416; Tekirian, T. L. et al. 1998 *J Neuropathol Exp Neurol* 57, 76-94; Geddes, J. W. et al.

1999 *Neurobiol Aging* 20, 75-79; Harigaya, Y. et al. 2000 *Biochem Biophys Res Commun* 276, 422-427) and they are also present in pre-amyloid lesions (Lalowski, M. et al. 1996 *J Biol Chem* 271, 33623-33631). The accumulation of [pGlu³]A β (3-40/42) peptides is likely due to the structural modification that enhances aggregation and confers resistance to most aminopeptidases (Saïdo, T. C. et al. 1995 *Neuron* 14, 457-466 ; Tekirian, T. L. et al. 1999 *J Neurochem* 73, 1584-1589). This evidence provides clues for a pivotal role of [pGlu³]A β (3-40/42) peptides in AD pathogenesis. However, relatively little is known about their neurotoxicity and aggregation properties (He, W. and Barrow, C. J. 1999 *Biochemistry* 38, 10871-10877; Tekirian, T. L. et al. 1999 *J Neurochem* 73, 1584-1589). Moreover, the action of these isoforms on glial cells and the glial response to these peptides are completely unknown, although activated glia is strictly associated to senile plaques and might actively contribute to the accumulation of amyloid deposits. In recent studies, the toxicity, aggregation properties and catabolism of A β (1-42), A β (1-40), [pGlu³]A β (3-42) and [pGlu³]A β (3-40) peptides were investigated in neuronal and glial cell cultures, and it was shown that pyroglutamate modification exacerbates the toxic properties of A β -peptides and also inhibits their degradation by cultured astrocytes. Shirotani et al. (2002) investigated the generation of [pGlu³]A β peptides in primary cortical neurons infected by Sindbis virus *in vitro*. They constructed amyloid precursor protein complementary DNAs, which encoded a potential precursor for [pGlu³]A β by amino acid substitution and deletion. For one artificial precursor starting with a N-terminal glutamine residue instead of glutamate in the natural precursor, a spontaneous conversion or an enzymatic conversion by glutaminyl cyclase to pyroglutamate was suggested. The cyclization mechanism of N-terminal glutamate at position 3 in the natural precursor of [pGlu³]A β was not determined *in vivo* (Shirotani, K. et al. 2002 *Neurosci Lett* 327, 25-28)

Familial British Dementia (FBD) and Familial Danish Dementia (FDD) are early-onset autosomal dominant disorders characterized by progressive cognitive impairment, spasticity and cerebellar ataxia (Ghiso, J. et al. 2000, *Ann N Y Acad Sci* 903, 129-137; Vidal, R. et al. 1999, *Nature* 399, 776-781; Vidal, R. et al. 2004, *J Neuropathol Exp Neurol* 63, 787-800). Similar to Alzheimers disease, widespread parenchymal and vascular amyloid deposits are formed in patients accompanied by Hippocampal

neurodegeneration, complement and glial activation (Rostagno, A. et al. 2002, *J Biol Chem* 277, 49782-49790). The diseases are caused by different mutations in the BRI gene (SwissProt Q9Y287) leading to an open reading frame that is 11 amino acids longer compared to wild type BRI. In case of FBD, the change in the ORF is caused
5 by a mutation in the stop codon of BRI (BRI-L), whereas in FDD a ten-nucleotide duplication-insertion leads to a larger BRI (BRI-D) (Ghisso J. et al. 2001 *Amyloid* 8, 277-284; Rostagno, A. et al. 2002 *J Biol Chem* 277, 49782-49790). BRI, a class 2 transmembran protein encoded on chromosome 13, has shown to be processed by furin and other prohormone convertases in the C-terminal region, releasing a 23
10 amino acids long peptide (Kim, S. H. et al. 2000 *Ann N Y Acad Sci* 920, 93-99; Kim, S. H. et al. 2002 *J Biol Chem* 277, 1872-1877). Cleavage of the mutant BRI proteins BRI-D and BRI-L leads to generation of peptides (ABri and ADan, both 34 amino acids) that are prone to aggregation causing non-fibrillar deposits as well as amyloid fibrils (El Agnaf, O. M. et al. 2004 *Protein Pept Lett* 11, 207-212; El Agnaf, O. M. et al. 2001 *Biochemistry* 40, 3449-3457; El Agnaf, O. M. et al. 2001 *J Mol Biol* 310, 157-168; Srinivasan et al. 2003 *J Mol Biol* 333, 1003-1023). The ADan and ABri peptides are identical in their N-terminal 22 amino acids, but contain distinct C-terminal regions. The C-terminal parts have shown to be required for fibril formation and neurotoxicity (El Agnaf, O. M. et al. 2004 *Protein Pept Lett* 11, 207-212).

20 It has been shown that the N-Terminus of the ABri and ADan peptides is blocked by pyroglutamyl formation. According to pyroglutamyl formation at the N-terminus of A β in Alzheimers disease, pGlu is formed from glutamic acid (Ghisso J. et al. 2001 *Amyloid* 8; Saido et al. 1995 *Neuron* 14, 457-466). Pyroglutamyl formation, in turn, stabilizes the peptides towards degradation by most aminopeptidases thus provoking
25 the progression of the diseases. Aggregate formation has been shown to proceed extracellularly but also in the secretory pathway of the cells (Kim et al. 2002 *J Biol Chem* 277, 1872-1877). Therefore, suppression of pGlu formation at the N-terminus of neurotoxic ABri and ADan peptides by inhibition of glutaminy and glutamate cyclases represents a new approach to treat FBD and FDD.

30 Dipeptidyl peptidase IV (DP IV) is a post-proline (to a lesser extent post-alanine, post-serine or post-glycine) cleaving serine protease found in various tissues of the body including kidney, liver, and intestine and cleaves N-terminal dipeptides from a peptide chain. Recently it was shown that DP IV plays an important role in

neuropeptide metabolism, T-cell activation, attachment of cancer cells to the endothelium and the entry of HIV into lymphoid cells. See therefore WO 02/34242, WO 02/34243, WO 03/002595 and WO 03/002596.

- 5 The DP IV inhibitors disclosed in WO 99/61431 comprise an amino acid residue and a thiazolidine or pyrrolidine group, and salts thereof, especially L-*threo*-isoleucyl thiazolidine, L-*allo*-isoleucyl thiazolidine, L-*threo*-isoleucyl pyrrolidine, L-*allo*-isoleucyl thiazolidine, L-*allo*-isoleucyl pyrrolidine.
- 10 Further examples of low molecular weight dipeptidyl peptidase IV inhibitors are agents such as tetrahydroisoquinolin-3-carboxamide derivatives, N-substituted 2-cyanopyrroles and -pyrrolidines, N-(N'-substituted glycyl)-2-cyanopyrrolidines, N-(substituted glycyl)-thiazolidines, N-(substituted glycyl)-4-cyanothiazolidines, amino-acyl-borono-prolyl-inhibitors, cyclopropyl-fused pyrrolidines and heterocyclic
15 compounds. Inhibitors of dipeptidyl peptidase IV are described in US 6,380,398, US 6,011,155; US 6,107,317; US 6,110,949; US 6,124,305; US 6,172,081; WO 95/15309, WO 99/61431, WO 99/67278, WO 99/67279, DE 198 34 591, WO 97/40832, DE 196 16 486 C 2, WO 98/19998, WO 00/07617, WO 99/38501, WO 99/46272, WO 99/38501, WO 01/68603, WO 01/40180, WO 01/81337, WO
20 01/81304, WO 01/55105, WO 02/02560 and WO 02/14271, WO 02/04610, WO 02/051836, WO 02/068420, WO 02/076450; WO 02/083128, WO 02/38541, WO 03/000180, WO 03/000181, WO 03/000250, WO 03/002530, WO 03/002531, WO 03/002553, WO 03/002593, WO 03/004496, WO 03/024942 and WO 03/024965.

Summary of the invention

The present invention provides novel physiological substrates of QC in mammals,
30 selected from the group consisting of Glu¹-ABri, Glu¹-ADan, Gln³-Aβ(3-40/42), and Gln¹-Gastrins (17 and 34) and the use of effectors of QC and pharmaceutical compositions comprising effectors of QC for the treatment of conditions that can be treated by modulation of QC activity, preferably selected from the group consisting of

duodenal cancer with or w/o *Helicobacter pylori* infections, colorectal cancer, Zollinger-Ellison syndrome, Familial British Dementia and Familial Danish Dementia.


It was shown by inhibition studies that human QC is a metal-dependent transferase.

- 5 QC apoenzyme could be reactivated most efficiently by zinc ions, and the metal-binding motif of zinc-dependent aminopeptidases is also present in human QC. Compounds interacting with the active-site bound metal are potent inhibitors of QC.

- 10 Unexpectedly, it was shown that recombinant human QC as well as QC-activity from brain extracts catalyze both, the *N*-terminal glutaminyl as well as glutamate cyclization. Most striking is the finding, that cyclase-catalyzed Glu¹-conversion is favored around pH 6.0 while Gln¹-conversion to pGlu-derivatives occurs with a pH-optimum of around 8.0. Since the formation of pGlu-A β -related peptides can therefore be suppressed by inhibition of recombinant human QC and QC-activity
15 from pig pituitary extracts, the enzyme QC is, according to the present invention, a target in drug development for treatment of Alzheimer's disease.

- The present invention provides pharmaceutical compositions for parenteral, enteral or oral administration, comprising at least one effector of QC optionally in
20 combination with customary carriers and/or excipients; or comprising at least one effector of QC in combination with at least one DP IV-inhibitor, optionally in combination with customary carriers and/or excipients.

- The present invention provides QC-inhibitors which can be described generally by the formula 1 or the pharmaceutically acceptable salts thereof, including all
25 stereoisomers:



formula 1

wherein $R^1 - R^6$ are independently H or a branched or unbranched alkyl chain, a branched or unbranched alkenyl chain, a branched or unbranched alkynyl chain, carbocyclic, aryl, heteroaryl, heterocyclic, aza-amino acid, amino acid or a mimetic thereof, peptide or a mimetic thereof; all of the above residues optionally being substituted, and
n can be 0-2.

Brief description of the drawings

Further understanding of these and other aspects of the present invention will be had by reference to the figures wherein:

Figure 1 shows progress curves of the cyclization of H-Gln-Ala-OH, catalyzed by human QC, monitoring the decrease in absorbance at 340 nm. The samples contained 0.3 mM NADH/ H^+ , 14 mM α -Ketoglutaric acid, 30 U/ml glutamic dehydrogenase and 1 mM H-Gln-Ala-OH. From curve A-D, varying concentrations of QC were applied: A, 10 mU/ml, B, 5 mU/ml, C, 2.5 mU/ml. In case of curve D, QC was omitted. A linear relationship was obtained between the QC concentration and the observed activity (inset).

Figure 2 shows the pH dependence of human and papaya (inset) QC, determined under first-order rate conditions using Gln- β NA as substrate. In case of human QC, a buffer system providing a constant ionic strength according to Ellis and Morrison was used, consisting of 25 mM MES, 25 mM acetic acid and 50 mM Tris (Ellis, K. J. and Morrison, J. F. 1982 *Methods Enzymol.* 87, 405-426). Due to a slightly inhibiting effect of Tris, papaya QC was investigated using a 50 mM Mops buffer. The ionic strength was adjusted to 0.05 M by addition of NaCl. The rate profiles were evaluated by fitting to a model that is based on dissociating groups. In case of papaya QC, a pK_a of 7.13 ± 0.03 was obtained by fitting of the data to a single dissociation model.

Figure 3 shows the effect of the pH on the stability of the QC from Papaya latex and human QC. An enzyme stock solution was diluted 20-fold in 0.1 M buffer of various

pH values (pH 4-7 sodium citrate, pH 7-10 sodium phosphate). Enzyme solutions were incubated at 30 °C for 30 min and subsequently enzymatic activity was analyzed according to the standard protocol.

5 **Figure 4** shows the comparison of the specificity constant k_{cat}/K_M for a set of substrates containing glutamate in the second amino acid position. Whereas an increase in specificity of human QC was detected from the di- to the tetrapeptides, no change was observed in case of papaya QC. The data presented here is a replot of the parameters given in Table 3.

10

Figure 5 shows the formation of pGlu-Lys(pGlu)-Arg-Leu-Ala-NH₂ from H-Gln-Lys(Gln)-Arg-Leu-Ala-NH₂, catalyzed by human QC. Substrate conversion is monitored by a time-dependent change in the m/z ratio due to the expulsion of ammonia. The sample composition was 0.5 mM substrate, 38 nM QC in 40 mM
15 Tris/HCl, pH 7.7. At the times indicated, samples were removed from the assay tube, mixed with matrix solution (1:1 v/v) and subsequently the mass spectra recorded. A very similar dependence was observed in case of papaya QC.

Figure 6 shows the formation of pGlu-Phe-Lys-Ala-Glu-NH₂ from H-Gln(NMe)-Phe-Lys-Ala-Glu-NH₂ catalyzed by papaya QC. Substrate conversion is monitored by a
20 time-dependent change in the m/z ratio due to the expulsion of methylamine. The sample composition was 0.5 mM substrate, 0.65 µM papaya QC in 40 mM Tris/HCl, pH 7.7. At the times indicated, samples were removed from the assay tube, mixed with matrix solution (1:1 v/v) and subsequently the mass spectra recorded. No
25 substrate conversion was observed in samples without papaya QC or by applying up to 1.5 µM human QC to the substrate (not shown).

Figure 7 shows the formation of Gln³-Aβ(3-11)a from Gln³-Aβ(1-11)a catalysed by DPIV. At the times indicated, samples were removed from the assay tube, mixed with
30 matrix solution (1:1 v/v) and subsequently the mass spectra recorded.

Figure 8 shows the prevention of the cleavage of Gln³-Aβ(1-11)a by the DP IV-inhibitor Val-Pyrrolidide (Val-Pyrr). At the times indicated, samples were removed

from the assay tube, mixed with matrix solution (1:1 v/v) and subsequently the mass spectra recorded.

Figure 9 shows the formation of [pGlu³]Aβ(3-11)a from Gln³-Aβ(3-11)a catalyzed by QC. At the times indicated, samples were removed from the assay tube, mixed with matrix solution (1:1 v/v) and subsequently the mass spectra recorded.

Figure 10 shows the inhibition of the formation of [pGlu³]Aβ(3-11)a from Gln³-Aβ(3-11)a by the QC-inhibitor 1,10-phenanthroline. At the times indicated, samples were removed from the assay tube, mixed with matrix solution (1:1 v/v) and subsequently the mass spectra recorded.

Figure 11 shows the formation of [pGlu³]Aβ(3-11)a from Gln³-Aβ(1-11)a after consecutive catalysis by DP IV and QC. At the times indicated, samples were removed from the assay tube, mixed with matrix solution (1:1 v/v) and subsequently the mass spectra recorded.

Figure 12 shows the inhibition of [pGlu³]Aβ(3-11)a formation from Gln³-Aβ(1-11)a by the QC-inhibitor 1,10-phenanthroline in the presence of catalytically active DP IV and QC. At the times indicated, samples were removed from the assay tube, mixed with matrix solution (1:1 v/v) and subsequently the mass spectra recorded.

Figure 13 shows the reduction of [pGlu³]Aβ(3-11)a formation from Gln³-Aβ(1-11)a by the DP IV-inhibitor Val-Pyrr in the presence of catalytically active DP IV and QC. At the times indicated, samples were removed from the assay mixture, mixed with matrix solution (1:1 v/v) and subsequently the mass spectra recorded.

Figure 14 shows the formation of [pGlu³]Aβ(3-11)a from Gln³-Aβ(1-11)a after consecutive catalysis by aminopeptidase(s) and QC that are present in porcine pituitary homogenate. At the times indicated, samples were removed from the assay tube, mixed with matrix solution (1:1 v/v) and subsequently the mass spectra recorded.

Figure 15 A and B show Mass spectra of A β (3-11)a and A β (3-21)a incubated with recombinant human QC, that was boiled for 10 min before use. **C and D** show Mass spectra of A β (3-11)a and A β (3-21)a in presence of active human QC resulting in the formation of [pGlu³]A β (3-11)a and [pGlu³]A β (3-21)a, respectively. **E and F** show

5 Mass spectra of A β (3-11)a and A β (3-21)a in presence of active QC and 5 mM Benzimidazole suppressing the formation of [pGlu³]formation.

Figure 16 shows reaction rates of papaya QC- catalyzed Glu- β NA-conversion plotted against the substrate concentration. The initial rates were measured in 0.1 M pyrophosphate buffer, pH 6.1 (squares), 0.1 M phosphate buffer, pH 7.5 (circles) and

10 0.1 M borate buffer, pH 8.5 (triangles). The kinetic parameters were as follows: $K_M = 1.13 \pm 0.07$ mM, $k_{cat} = 1.13 \pm 0.04$ min⁻¹ (pH 6.1); $K_M = 1.45 \pm 0.03$ mM, $k_{cat} = 0.92 \pm 0.01$ min⁻¹ (pH 7.5); $K_M = 1.76 \pm 0.06$ mM, $k_{cat} = 0.56 \pm 0.01$ min⁻¹ (pH 8.5).

15 **Figure 17** shows the pH-dependence of the conversion of Gln- β NA (circles) and Glu- β NA (squares), determined under first-order rate-law conditions ($S \ll K_M$). Substrate concentration was 0.01 mM and 0.25 mM, respectively. For both determinations, a three-component buffer system was applied consisting of 0.05 M acetic acid, 0.05 M pyrophosphoric acid and 0.05 M Tricine. All buffers were adjusted to equal

20 conductivity by addition of NaCl, in order to avoid differences in ionic strength. The data were fitted to equations that account for two dissociating groups revealing pK_a -values of 6.91 ± 0.02 and 9.5 ± 0.1 for Gln- β NA and 4.6 ± 0.1 and 7.55 ± 0.02 for Glu- β NA. The pK_a -values of the respective substrate amino groups, determined by titration, were 6.97 ± 0.01 (Gln- β NA) and 7.57 ± 0.05 (Glu- β NA). All determinations

25 were carried out at 30 °C.

Figure 18 shows progress curves of human QC-catalyzed cyclization of H-Gln-AMC in presence of imidazole, dipicolinic acid and in absence of an inhibitory compound. The hyperbolic shape of the curve in presence of dipicolinic acid indicates metal ion

30 removal from the active site of QC.

Figure 19 shows the time-dependent inactivation of QC by the heterocyclic chelator 1,10-phenanthroline. After incubation of the QC-enzyme with the inhibitor in absence

of substrate (continuous line), a reduced enzymatic activity was observed compared to samples that were not preincubated with inhibitor (dotted trace), indicating metal ion removal from the active site of QC.

5 **Figure 20** shows the reactivation of human QC with monovalent- and divalent metal ions. QC was inactivated by addition of 2 mM dipicolinic acid in 50 mM Bis-Tris, pH 6.8. Subsequently, the enzyme was subjected to dialysis against 50 mM Bis-Tris, pH 6.8, containing 1.0 mM EDTA. Reactivation of the enzymes was achieved by incubation of the inactivated enzyme sample with metal ions at a concentration of
10 0.5 mM, in presence of 0.5 mM EDTA in order to avoid an unspecific reactivation by traces of metal ions present in buffer solutions. Controls are given by enzyme samples that were not inactivated, but also dialyzed against EDTA solution as the inactivated enzyme (+EDTA) and enzyme samples that were dialyzed against buffer solutions without added EDTA (-EDTA).

15 **Figure 21** Sequence alignment of human QC (hQC) and other M28 family members of the metallopeptidase Clan MH. Multiple sequence alignment was performed using ClustalW at ch.EMBnet.org with default settings. The conservation of the zinc-ion ligating residues is shown for human QC (hQC; GenBank X71125), the Zn-
20 dependent aminopeptidase from *Streptomyces griseus* (SGAP; Swiss-Prot P80561), and within the N-acetylated-alpha-linked acidic dipeptidase (NAALADase I) domain (residues 274 – 587) of the human Glutamate carboxypeptidase II (hGCP II; Swiss-Prot Q04609). The amino acids involved in metal binding are typed in bold and underlined. In case of human QC, these residues are the putative counterparts to the
25 peptidases.

Figure 22 shows the pH-dependence of inhibition of murine QC by cysteamine (squares), dimethyl-cysteamine (circles) and mercaptoethanol (triangles). Points were fitted to equations that account for one dissociating group. The curves reveal
30 different shapes, indicating that the dependence is due to alterations in the protonation state of the inhibitor. The kinetically determined pK_a -values for cysteamine (8.71 ± 0.07) and dimethyl-cysteamine (8.07 ± 0.03) match well with those obtained from literature data for the amino group (8.6 and 7.95, respectively) (Buist, G. J. and Lucas, H. J. 1957 *J Am Chem Soc* 79, 6157; Edsall, J. T. *Biophysical*

Chemistry, Academic Press, Inc., New York, 1958). Accordingly, the pH-dependence of mercaptoethanol possesses a slope of unity, since it does not carry a dissociative group in the pH-range investigated.

5 **Figure 23** shows a Lineweaver-Burk plot of the kinetic data obtained for conversion of Gln-AMC (0.25, 0.125, 0.063, 0.031 mM), catalyzed by human QC in presence of various concentrations of cysteamine (0, 0.25, 0.5, 1 mM). The data were fitted according to competitive inhibition. The determined K_i -value was 0.037 ± 0.001 mM.

10 **Peptide Sequences**

The peptides mentioned and used herein have the following sequences:

A β (1-42):

15 Asp-Ala-Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val-Ile-Ala

A β (1-40):

20 Asp-Ala-Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val

A β (3-42):

25 Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val-Ile-Ala

A β (3-40):

Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val

A β (1-11)a:

30 Asp-Ala-Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-NH₂

A β (3-11)a:

35 Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-NH₂

A β (1-21)a:

40 Asp-Ala-Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-NH₂

A β (3-21)a:

Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-NH₂

Gln³-A β (3-40):

Gln-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-
Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val

Gln³-Aβ(3-21)a:

5 Gln-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-
NH₂

Gln³-Aβ(1-11)a:

10 Asp-Ala-Gln-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-NH₂

Gln³-Aβ(3-11)a:

Gln-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-NH₂

ABri

15 pGlu-Ala-Ser-Asn-Cys-Phe-Ala-Ile-Arg-His-Phe-Glu-Asn-Lys-Phe-Ala-Val-Glu-Thr-
Leu-Ile-Cys-Ser-Arg-Thr-Val-Lys-Lys-Asn-Ile-Ile-Glu-Glu-Asn

Glu¹-ABri

20 Glu-Ala-Ser-Asn-Cys-Phe-Ala-Ile-Arg-His-Phe-Glu-Asn-Lys-Phe-Ala-Val-Glu-Thr-
Leu-Ile-Cys-Ser-Arg-Thr-Val-Lys-Lys-Asn-Ile-Ile-Glu-Glu-Asn

ADan

pGlu-Ala-Ser-Asn-Cys-Phe-Ala-Ile-Arg-His-Phe-Glu-Asn-Lys-Phe-Ala-Val-Glu-Thr-
Leu-Ile-Cys-Phe-Asn-Leu-Phe-Leu-Asn-Ser-Gln-Glu-Lys-His-Tyr

25

Glu¹-ADan

Glu-Ala-Ser-Asn-Cys-Phe-Ala-Ile-Arg-His-Phe-Glu-Asn-Lys-Phe-Ala-Val-Glu-Thr-
Leu-Ile-Cys-Phe-Asn-Leu-Phe-Leu-Asn-Ser-Gln-Glu-Lys-His-Tyr

30

Detailed description of the invention

The present invention provides effectors of glutaminyl cyclase (QC) for

- a) the treatment of diseases in mammals that can be treated by modulation of
QC activity *in vivo* and/or
- 35 b) the modulation of physiological processes based on the action of pGlu-
containing peptides caused by modulation of QC activity.

Furthermore, the present invention provides compounds for the inhibition of glutaminy cyclase (QC, EC 2.3.2.5) and/or QC-like enzymes in a mammal and the use of inhibitors of QC activity for the treatment of pathological conditions related to QC activity.

5

The present invention also provides a new method for the treatment of Alzheimer's disease and Down Syndrome. The N-termini of amyloid β -peptides deposited in Alzheimer's disease and Down syndrome brain bear pyroglutamic acid. The pGlu formation is an important event in the development and progression in the disease, since the modified amyloid β -peptides show an enhanced tendency to β -amyloid aggregation and toxicity, likely worsening the onset and progression of the disease (Russo, C. et al. 2002 *J Neurochem* 82,1480-1489).

In contrast, in the natural A β -peptides (3-40/42), glutamic acid is present as an N-terminal amino acid. An enzymatic conversion of Glu to pGlu was not known to date. Moreover, spontaneous cyclization of Glu-peptides to pGlu-peptides has not been observed as yet. Therefore, one aspect of the present invention was to determine the role of QC in Alzheimer's disease and Down Syndrome. This aspect was addressed by the synthesis of A β (3-11)a and A β (1-11)a, containing the amino acid glutamine instead of glutamic acid at position three, the determination of the substrate characteristics of these modified amyloid β -peptides against QC, DP IV and DP IV-like enzymes and aminopeptidases and the use of inhibitors of QC to prevent the formation of pGlu from a N-terminal glutaminy residue of the amyloid β -derived peptides (1-11) and (3-11). The results are shown in example 8. The applied method is described in example 3.

To date, there are no hints indicating an involvement of QC in the progression of the diseases, because glutamic acid is the N-terminal amino acid in A β (3-40/42, or 11-40/42). But, QC is the only known enzyme capable of forming pGlu at the N-terminus of peptides. Other aspects of the present invention concern the following findings and discoveries:

30

- a) In addition to glutamine, QC catalyzes the cyclization of glutamic acid into pyroglutamic acid at very low rates,
- b) Glutamic acid of APP or its subsequently formed amyloid β -peptides is converted into glutamine post-translationally by an unknown enzymatic activity and in a second step, QC catalyzes the cyclization of glutamine into pyroglutamic acid after processing of the amyloid β -peptide N-terminus,
- c) Glutamic acid is converted into glutamine post-translationally by a chemical catalysis or autocatalysis and subsequently, QC catalyzes the cyclization of glutamine to pyroglutamic acid after processing of the amyloid β -peptide N-terminus,
- d) There are mutations in the APP gene, which encode the amyloid β -protein, leading to Gln instead of Glu in position 3. After translation and processing of the N-terminus, QC catalyzes the cyclization of glutamine to pyroglutamic acid,
- e) Glutamine is incorporated into the nascent peptide chain of APP, due to a malfunction of an unknown enzymatic activity and subsequently, QC catalyzes the cyclization of N-terminal glutamine to pyroglutamic acid after processing of the amyloid β -peptide N-terminus.

QC is involved in the critical step in all five cases listed above, namely the formation of pyroglutamic acid that favors the aggregation of amyloid β -peptides. Thus, an inhibition of QC leads to a prevention of the precipitation of the plaque-forming A β (3-40/42) or A β (11-40/42), causing the onset and progression of Alzheimer's disease and Down Syndrome, independently of the mechanism by which cyclization occurs.

Glutamate is found in positions 3, 11 and 22 of the amyloid β -peptide. Among them the mutation from glutamic acid (E) to glutamine (Q) in position 22 (corresponding to amyloid precursor protein APP 693, Swissprot P05067) has been described as the so called Dutch type cerebroarterial amyloidosis mutation.

The β -amyloid peptides with a pyroglutamic acid residue in position 3, 11 and/or 22 have been described to be more cytotoxic and hydrophobic than $A\beta(1-40/42/43)$ (Saïdo, T.C. 2000 *Medical Hypotheses* 54: 427-429).

5 The multiple N-terminal variations can be generated by the β -secretase enzyme β -site amyloid precursor protein-cleaving enzyme (BACE) at different sites (Huse, J. T. et al. 2002 *J Biol Chem* 277, 16278-16284), and/or by aminopeptidase processing. In all cases, cyclization can take place according to a)-e) as described above.

10 So far, there was no evidence supporting the enzymatic conversion of Glu^1 -peptides into pGlu-peptides by an unknown glutamyl cyclase (EC) corresponding to pathway a) (Garden, R. W. et al. 1999 *J Neurochem* 72, 676-681; Hosoda, R. et al. 1998 *J Neuropathol Exp Neurol* 57, 1089-1095). To date, no such enzyme activity has been identified, capable to cyclize Glu^1 -peptides which are protonated N-terminally and possess a negatively charged Glu^1 γ -carboxylate moiety under mildly alkaline pH-
15 conditions.

QC-activity against Gln^1 -substrates is dramatically reduced below pH 7.0. In contrast, it appears that Glu^1 -conversion can occur at acidic reaction conditions (Iwatsubo, T. et al. 1996 *Am J Pathol* 149, 1823-1830; Russo, C. et al. 1997 *FEBS Lett* 409, 411-
20 416; Russo, C. et al. 2001 *Neurobiol Dis* 8, 173-180; Tekirian, T. L. et al. 1998 *J Neuropathol Exp Neurol* 57, 76-94; Russo, C. et al. 2002 *J Neurochem* 82, 1480-1489; Hosoda, R. et al. 1998 *J Neuropathol Exp Neurol* 57, 1089-1095; Garden, R. W. et al. 1999 *J Neurochem* 72, 676-681).

25 According to the present invention it was investigated whether QC is able to recognize and to turnover amyloid- β derived peptides under mild acidic conditions. Therefore, the peptides Gln^3 - $A\beta(1-11)a$, $A\beta(3-11)a$, Gln^3 - $A\beta(3-11)a$, $A\beta(3-21)a$, Gln^3 - $A\beta(3-21)a$ and Gln^3 - $A\beta(3-40)$ as potential substrates of the enzyme were synthesized and investigated. These sequences were chosen for mimicking natural N-terminally
30 and C-terminally truncated Glu^3 - $A\beta$ peptides and Gln^3 - $A\beta$ peptides which could occur due to posttranslational Glu-amidation.

In the present invention it was shown that papaya and human QC catalyze both glutamyl and glutamyl cyclization. Apparently, the primary physiological function of QC is to finish hormone maturation in endocrine cells by glutamine cyclization prior or during the hormone secretion process. Such secretory vesicles are known to be acidic in pH. Thus, a side activity of the enzyme in the narrow pH-range from 5.0 to 7.0 could be its newly discovered glutamyl cyclase activity transforming also Glu-A β peptides. However, due to the much slower occurring Glu-cyclization compared to Gln-conversion, it is questionable whether the glutamyl cyclization plays a significant physiological role. In the pathology of neurodegenerative disorders, however, the glutamyl cyclization is of relevance.

Investigating the pH-dependency of this enzymatic reaction, we found that the unprotonated *N*-terminus was essential for the cyclization of Gln¹-peptides and accordingly that the pK_a-value of the substrate was identical to the pK_a-value for QC-catalysis (see Figure 17). Thus, QC stabilizes the intramolecular nucleophilic attack of the unprotonated α -amino moiety on the γ -carbonyl carbon electrophilically activated by amidation (Scheme 1).

In contrast to the monovalent charge present on *N*-terminal glutamine containing peptides, the *N*-terminal Glu-residue in Glu-containing peptides is predominantly bivalently charged around neutral pH. Glutamate exhibits pK_a-values of about 4.2 and 7.5 for the γ -carboxylic and for the α -amino moiety, respectively. I.e. at neutral pH and above, although the α -amino nitrogen is in part or fully unprotonated and nucleophilic, the γ -carboxylic group is unprotonated, and so exercising no electrophilic carbonyl activity. Hence, intramolecular cyclization is impossible.

However, in the pH-range of about 5.2-6.5, between their respective pK_a-values, the two functional groups are present both in non-ionized forms, in concentrations of about 1-10% (-NH₂) or 10-1% (-COOH) of total *N*-terminal Glu-containing peptide. As a result, over a mildly acidic pH-range species of *N*-terminal Glu-peptides are present which carry both groups uncharged, and, therefore, it is possible that QC could stabilize the intermediate of intramolecular cyclization to pGlu-peptide. I.e. if the γ -carboxylic group is protonated, the carbonyl carbon is electrophilic enough to allow nucleophilic attack by the unprotonated α -amino group. At this pH the hydroxyl ion functions as a leaving group (Scheme 3). These assumptions are corroborated by the pH-dependence data obtained for the QC catalyzed conversion of Glu- β NA (see

example 11). In contrast to glutamine conversion of Gln- β NA by QC, the pH-optimum of catalysis shifts to the acidic range around pH 6.0, i.e. the pH-range, in which substrate molecule species are simultaneously abundant carrying a protonated γ -carboxyl and unprotonated α -amino group. Furthermore, the kinetically determined
5 pK_a-value of 7.55 ± 0.02 is in excellent agreement with that of the α -amino group of Glu- β NA, determined by titration (7.57 ± 0.05).

Physiologically, at pH 6.0 the second-order rate constant (or specificity constant, k_{cat}/K_M) of the QC-catalyzed glutamate cyclization might be in the range of 8,000fold slower than the one for glutamine cyclization (Figure 17). However, the nonenzymatic
10 turnover of both model substrates Glu- β NA and Gln- β NA is negligible, which conforms to the observed negligible pGlu-peptide formation in the present invention. Hence, for the pGlu-formation by QC an acceleration of at least 10^8 can be estimated from the ratio of the enzymatic versus non-enzymatic rate constants (comparing the second-order rate constants for the enzyme catalysis with the respective
15 nonenzymatic cyclization first-order rate constants, the catalytic proficiency factor is 10^9 - 10^{10} M^{-1} for the Gln- and the Glu-conversion, respectively). The conclusion from these data is, that *in vivo* only an enzymatic path resulting in pGlu-formations seems conceivable.

Since QC is highly abundant in the brain and taking into account the high turnover
20 rate of 0.9 min^{-1} recently found for the maturation of $30 \text{ }\mu\text{M}$ of (Gln-)TRH-like peptide (Prokai, L. et al. 1999 *J Med Chem* 42, 4563-4571), one can predict a cyclization half-life of about 100 hours for an appropriate glutamate-substrate, similar reaction conditions provided. Moreover, given compartmentalization and localization of brain QC/EC in the secretory pathway, the actual *in vivo* enzyme and substrate
25 concentrations and reaction conditions might be even more favorable for the enzymatic cyclization in the intact cell. And, if *N*-terminal Glu is transformed to Gln a much more rapid pGlu-formation mediated by QC could be expected. *In vitro*, both reactions were suppressed by applying inhibitors of QC/EC-activity (Figures 9, 10 and 15).

30 In summary, the present invention shows that human QC, which is highly abundant in the brain, is a catalyst to the formation of the amyloidogenic pGlu-A β peptides from Glu-A β and Gln-A β precursors which make up more than 50% of the plaque deposits found in Alzheimer's Disease. These findings identify QC/EC as a player in senile

plaque formation and thus as a novel drug target in the treatment of Alzheimer's Disease.

In a second embodiment of the present invention, it was found that amyloid β -
5 derived peptides are a substrate of dipeptidyl peptidase IV (DP IV) or DP IV-like
enzymes, preferably dipeptidyl peptidase II (DPII). DP IV, DP II or other DP IV-like
enzymes release a dipeptide from the N-terminus of the modified amyloid β -peptide
(1-11) generating amyloid β -peptide (3-11) with glutamine as the N-terminal amino
acid residue. The results are shown in example 8.

10 Prior to cleavage by DP II, DPIV or other DP IV-like enzymes, the peptide bond
between aspartic acid (residue 1 of amyloid β -peptide) and alanine (residue 2 of
amyloid β -peptide) may be isomerised yielding an isoaspartyl residue as described in
the literature (Kuo, Y.-M., Emmerling, M. R., Woods, A. S., Cotter, R. J., Roher, A. E.
(1997) *BBRC* 237, 188-191; Shimizu, T., Watanabe, A., Ogawara, M., Mori, H. and
15 Shirasawa, T. (2000) *Arch. Biochem. Biophys.* 381, 225-234).

These isoaspartyl residues render the amyloid β -peptide resistant against
aminopeptidase degradation and consequently the core plaques contain high
amounts of isoAsp¹-amyloid β -peptides, which suggests a reduced turnover at the N-
terminus.

20 However, in the present invention it is demonstrated for the first time, that the N-
terminal dipeptide H-isoAsp¹-Ala²-OH can be released by dipeptidyl peptidases
especially under acidic conditions. Furthermore, it was shown that isomerization can
precede cleavage by β -secretase too, and that isomerization may accelerate
proteolytic processing, thus leading to liberation of an N-terminal isoaspartyl bond of
25 isoAsp¹-amyloid β -peptides which subsequently is subject to turnover by DP II, DPIV
or DP IV-like enzymes (Momand, J. and Clarke, S. 1987 *Biochemistry* 26, 7798-
7805; Kuo, Y.-M. et al. 1997 *BBRC* 237, 188-191). Accordingly, inhibition of
isoaspartyl formation may lead to the reduction of cleavage by β -secretase and, in
turn, to a reduced formation of amyloid β -peptides. In addition, blockage of the
30 isoAsp¹-amyloid β -peptide turnover by inhibition of DP II, DPIV or DP IV-like enzymes
would prevent the exposure of Glu³-A β to QC/EC-catalyzed formation of [pGlu³]A β .

In a third embodiment of the present invention, a combination of inhibitors of DP IV-activity and of inhibitors of QC can be used for the treatment of Alzheimer's disease and Down Syndrome.

- 5 The combined effect of DP IV and/or DP IV-like enzymes and of QC is illustrated as follows:
- a) DP IV and/or DP IV-like enzymes cleave $A\beta(1-40/42)$, a dipeptide comprising H-Asp-Ala-OH and $A\beta(3-40/42)$ are released,
 - b) In a side reaction, QC catalyzes the cyclization of glutamic acid to
10 pyroglutamic acid at very low rates,
 - c) Glutamic acid is converted into glutamine at the N-terminus post-translationally by an unknown enzymatic activity and subsequently, QC catalyzes the cyclization of glutamine into pyroglutamic acid after processing of the amyloid β -peptide N-terminus,
 - d) Glutamic acid is converted into glutamine post-translationally by a
15 chemical catalysis or autocatalysis and in a second step, QC catalyzes the cyclization of glutamine into pyroglutamic acid after processing of the amyloid β -peptide N-terminus,
 - e) There are mutations in the APP gene, which encode the amyloid β -
20 protein, leading to Gln instead of Glu in position 3 of $A\beta$. After translation and processing of the N-terminus, QC catalyzes the cyclization of glutamine to pyroglutamic acid,
 - f) Glutamine is incorporated into the nascent peptide chain of APP,
25 due to a malfunction of an unknown enzymatic activity and subsequently, QC catalyzes the cyclization of N-terminally glutamine to pyroglutamic acid after processing of the amyloid β -peptide N-terminus,

30 The N-terminal Gln-exposure to QC-activity can be also triggered by different peptidase activities. Aminopeptidases can remove sequentially Asp and Ala from the N-terminus of $A\beta(1-40/42)$, thus unmasking amino acid three that is prone to cyclization. Dipeptidyl peptidases, such as DP I, DP II, DP IV, DP 8, DP 9 and DP 10,

remove the dipeptide Asp-Ala in one step. Hence, inhibition of aminopeptidase- or dipeptidylpeptidase-activity is useful to prevent the formation of A β (3-40/42).

5 The combined effect of inhibitors of DP IV and/or DP IV-like enzymes and of activity lowering effectors of QC is illustrated in the following way:

- a) The inhibitors of DP IV and/or DP IV-like enzymes inhibit the conversion of A β (1-40/42) to A β (3-40/42).
- 10 b) An N-terminal exposure of glutamic acid is thereby prevented and no conversion to glutamine, either by enzymatic or by chemical catalysis, subsequently leading to pyroglutamic acid formation, is possible.
- 15 c) Inhibitors of QC prevent in addition the formation pyroglutamic acid from any residual modified A β (3-40/42) molecules and those modified A β (3-40/42) molecules, which are generated by mutations of the APP gene.

20 Within the present invention, a similar combined action of DP IV or DP IV-like enzymes and QC was demonstrated for further peptide hormones, such as glucagon, CC chemokines and substance P.

25 Glucagon is a 29-amino acid polypeptide released from pancreatic islet alpha-cells that acts to maintain euglycemia by stimulating hepatic glycogenolysis and gluconeogenesis. Despite its importance, there remains controversy about the mechanisms responsible for glucagon clearance in the body. Pospisilik et al. assessed the enzymatic metabolism of glucagon using sensitive mass spectrometric techniques to identify the molecular products. Incubation of glucagon with purified porcine dipeptidyl peptidase IV (DP IV) yielded sequential production of glucagon3-29 and glucagons(5-29). In human serum, degradation to glucagons(3-29) was 30 rapidly followed by N-terminal cyclization of glucagon, preventing further DP IV-mediated hydrolysis. Bioassay of glucagon, following incubation with purified DP IV or normal rat serum demonstrated a significant loss of hyperglycemic activity, while a similar incubation in DP IV-deficient rat serum did not show any loss of glucagon

bioactivity. Degradation, monitored by mass spectrometry and bioassay, was blocked by the specific DP IV inhibitor, isoleucyl thiazolidine. These results identify DP IV as a primary enzyme involved in the degradation and inactivation of glucagon. These findings have important implications for the determination of glucagon levels in human plasma (Pospisilik A. et al. 2001 *Regul Pept* 96, 133-41).

Human Monocyte Chemotactic Protein 2 (MCP-2) has originally been isolated from stimulated osteosarcoma cells as a chemokine coproduced with MCP-1 and MCP-3. Von Coillie et al. (Van Coillie, E. et al. 1998 *Biochemistry* 37, 12672-12680) cloned a 5'-end extended MCP-2 cDNA from a human testis cDNA library. It encoded a 76 residue MCP-2 protein, but differed from the reported bone marrow-derived MCP-2 cDNA sequence in codon 46, which coded for a Lys instead of a Gln. This MCP-2Lys46 variant, caused by a single nucleotide polymorphism (SNP), was biologically compared with MCP-2Gln46. The coding regions were sub-cloned into the bacterial expression vector pHEN1, and after transformation of *Escherichia coli*, the two MCP-2 protein variants were recovered from the periplasm. Edman degradation revealed a Gln residue at the NH₂ terminus instead of a pGlu. rMCP-2Gln46 and rMCP-2Lys46 and the NH₂-terminal cyclic counterparts were tested on monocytic cells in calcium mobilization and chemotaxis assays. No significant difference in biological activity was observed between the rMCP-2Gln46 and rMCP-2Lys46 isoforms. However, for both MCP-2 variants the NH₂-terminal pyroglutamate was shown to be essential for chemotaxis, but not for calcium mobilization. NH₂-terminal truncation of rMCP-2Lys46 by the serine protease CD26/dipeptidyl peptidase IV (CD26/DPP IV) resulted in the release of the NH₂-terminal Gln-Pro dipeptide, whereas synthetic MCP-2 with an amino-terminal pGlu remained unaffected. CD26/DPP IV-clipped rMCP-2Lys46(3-76) was almost completely inactive in both chemotaxis and signaling assays. These observations indicated that the NH₂-terminal pGlu in MCP-2 is necessary for chemotactic activity but also that it protects the protein against degradation by CD26/DPP IV (van Coillie, E. et al. 1998, *Biochemistry* 37, 12672-80).

Within the present invention, it was determined by LC/MS-analysis that the formation of the N-terminal pyroglutamate residue determined in glucagon(3-29) (Pospisilik et al., 2001), and in MCP-2 isoforms (van Coillie et al., 1998), is catalyzed by QC.

In addition, it was proven by LC/MS-investigation that after N-terminal DP IV-catalyzed removal of the two dipeptides Lys-Pro and Arg-Pro from substance P the remaining [Gln⁵]substanceP5-11 is transformed by QC to [pGlu⁵]substanceP5-11.

5

DP IV inhibitors are disclosed in WO 99/61431. In particular, DP IV inhibitors are disclosed comprising an amino acid residue and a thiazolidine or pyrrolidine group, and salts thereof, especially L-threo-isoleucyl thiazolidine, L-allo-isoleucyl thiazolidine, L-threo-isoleucyl pyrrolidine, L-allo-isoleucyl thiazolidine, L-allo-isoleucyl pyrrolidine, and salts thereof.

Further examples of low molecular weight dipeptidyl peptidase IV inhibitors are agents such as tetrahydroisoquinolin-3-carboxamide derivatives, N-substituted 2-cyanopyrroles and -pyrrolidines, N-(N'-substituted glycy)-2-cyanopyrrolidines, N-(substituted glycy)-thiazolidines, N-(substituted glycy)-4-cyanothiazolidines, aminoacyl-borono-prolyl-inhibitors, cyclopropyl-fused pyrrolidines and heterocyclic compounds. Inhibitors of dipeptidyl peptidase IV are described in US 6,380,398, US 6,011,155; US 6,107,317; US 6,110,949; US 6,124,305; US 6,172,081; WO 95/15309, WO 99/61431, WO 99/67278, WO 99/67279, DE 198 34 591, WO 97/40832, DE 196 16 486 C 2, WO 98/19998, WO 00/07617, WO 99/38501, WO 99/46272, WO 99/38501, WO 01/68603, WO 01/40180, WO 01/81337, WO 01/81304, WO 01/55105, WO 02/02560 and WO 02/14271, WO 02/04610, WO 02/051836, WO 02/068420, WO 02/076450; WO 02/083128, WO 02/38541, WO 03/000180, WO 03/000181, WO 03/000250, WO 03/002530, WO 03/002531, WO 03/002553, WO 03/002593, WO 03/004496, WO 03/024942 and WO 03/024965.

Preferred for the use in combination with effectors of QC are DPIV inhibitors such as NVP-DPP728A (1-[[[2-{ {5-cyanopyridin-2-yl}amino}ethyl]amino]acetyl]-2-cyano-(S)-pyrrolidine) (Novartis) as disclosed by Hughes et al. 1999 *Biochemistry* 38 11597-11603, LAF-237 (1-[(3-hydroxy-adamant-1-ylamino)-acetyl]-pyrrolidine-2(S)-carbonitrile); disclosed by Hughes et al., Meeting of the American Diabetes Association 2002, Abstract no. 272 (Novartis), TSL-225 (tryptophyl-1,2,3,4-


tetrahydroisoquinoline-3-carboxylic acid), disclosed by Yamada et al. 1998 *Bioorg Med Chem Lett* 8 , 1537-1540, 2-cyanopyrrolidides and 4-cyanopyrrolidides as disclosed by Asworth et al. 1996 *Bioorg Med Chem Lett* 6, , 1163-1166 and 2745-2748 , FE-999011, disclosed by Sudre et al. 2002 *Diabetes* 51, 1461-1469 (Ferring)
5 and the compounds disclosed in WO 01/34594 (Guilford), employing dosages as set out in the above references.

More preferred DP IV inhibitors for the use in combination with effectors of QC are dipeptide compounds in which the amino acid is preferably selected from a natural
10 amino acid, such as, for example, leucine, valine, glutamine, glutamic acid, proline, isoleucine, asparagines and aspartic acid. The dipeptide-like compounds used according to the invention exhibit at a concentration (of dipeptide compounds) of 10 μ M, a reduction in the activity of plasma dipeptidyl peptidase IV or DP-IV-analogous enzyme activities of at least 10 %, especially of at least 40 %. Frequently a reduction
15 in activity of at least 60 % or at least 70 % is also desired *in vivo*. Preferred compounds may also exhibit a reduction in activity of a maximum of 20 % or 30 %.

Preferred dipeptide compounds are N-valyl prolyl, O-benzoyl hydroxylamine, alanyl pyrrolidine, isoleucyl thiazolidine like L-allo-isoleucyl thiazolidine, L-threo-isoleucyl
20 pyrrolidine and salts thereof, especially the fumaric salts, and L-allo-isoleucyl pyrrolidine and salts thereof. Especially preferred compounds are glutaminy pyrrolidine and glutaminy thiazolidine, H-Asn-pyrrolidine, H-Asn-thiazolidine, H-Asp-pyrrolidine, H-Asp-thiazolidine, H-Asp(NHOH)-pyrrolidine, H-Asp(NHOH)-thiazolidine, H-Glu-pyrrolidine, H-Glu-thiazolidine, H-Glu(NHOH)-pyrrolidine, H-Glu(NHOH)-
25 thiazolidine, H-His-pyrrolidine, H-His-thiazolidine, H-Pro-pyrrolidine, H-Pro-thiazolidine, H-Ile-azididine, H-Ile-pyrrolidine, H-L-*allo*-Ile-thiazolidine, H-Val-pyrrolidine and H-Val-thiazolidine and pharmaceutically acceptable salts thereof. These compounds are described in WO 99/61431 and EP 1 304 327.

30 Furthermore, the present invention provides for the use of effectors of QC in combination with substrate-like peptide compounds useful for competitive modulation of dipeptidyl peptidase IV catalysis. Preferred peptide compounds are 2-Amino octanoic acid-Pro-Ile, Abu-Pro-Ile, Aib-Pro-Ile, Aze-Pro-Ile, Cha-Pro-Ile, Ile-Hyp-Ile, Ile-Pro-*allo*-Ile, Ile-Pro-*t*-butyl-Gly, Ile-Pro-Val, Nle-Pro-Ile, Nva-Pro-Ile, Orn-Pro-Ile,

Phe-Pro-Ile, Phg-Pro-Ile, Pip-Pro-Ile, Ser(Bzl)-Pro-Ile, Ser(P)-Pro-Ile, Ser-Pro-Ile, *t*-butyl-Gly-Pro-D-Val, *t*-butyl-Gly-Pro-Gly, *t*-butyl-Gly-Pro-Ile, *t*-butyl-Gly-Pro-Ile-amide, *t*-butyl-Gly-Pro-*t*-butyl-Gly, *t*-butyl-Gly-Pro-Val, Thr-Pro-Ile, Tic-Pro-Ile, Trp-Pro-Ile, Tyr(P)-Pro-Ile, Tyr-Pro-*allo*-Ile, Val-Pro-*allo*-Ile, Val-Pro-*t*-butyl-Gly, Val-Pro-Val and
 5 pharmaceutically acceptable salts thereof, wherein *t*-butyl-Gly is defined as



10 and Ser(Bzl) and Ser(P) are defined as benzyl-serine and phosphoryl-serine, respectively. Tyr(P) is defined as phosphoryl-tyrosine. These compounds are disclosed in WO 03/002593.

Further preferred DP IV-inhibitors, which can be used according to the present
 15 invention in combination with effectors of QC, are peptidylketones, e.g.

- 2-Methylcarbonyl-1-N-[(L)-Alanyl-(L)-Valinyl]-(2S)-pyrrolidine hydrobromide, 2-Methylcarbonyl-1-N-[(L)-Valinyl-(L)-Prolyl-(L)-Valinyl]-(2S)-pyrrolidine hydrobromide,
- 20 • 2-[(Acetyl-oxy-methyl)carbonyl]-1-N-[(L)-Alanyl-(L)-Valinyl]-(2S)-pyrrolidine hydrobromide,
- 2-[Benzoyl-oxy-methyl)carbonyl]-1-N-[(L)-Alanyl-(L)-Valinyl]-(2S)-pyrrolidine hydrobromide,
- 25 • 2-[(2,6-Dichlorobenzyl)thiomethyl)carbonyl]-1-N-[(L)-Alanyl-(L)-Valinyl]-(2S)-pyrrolidine,
- 2-[Benzoyl-oxy-methyl)carbonyl]-1-N-[Glycyl-(L)-Valinyl]-(2S)-pyrrolidine hydrobromide,
- 2-[(1,3-Thiazolethiazol-2-yl)carbonyl]-1-N-[(L)-Alanyl-(L)-Valinyl]-(2S)-pyrrolidine trifluoroacetate,

- 2-[(Benzothiazolethiazol-2-yl)carbonyl]-1-N-[N-{(L)-Alanyl}-(L)-Valinyl]-(2S)-pyrrolidin trifluoroacetate,
- 2-[-Benzothiazolethiazol-2-yl)carbonyl]-1-N-[{(L)-Alanyl}-Glycyl]-(2S)-pyrrolidine trifluoroacetate,
- 5 • 2-[(Pyridin-2-yl)carbonyl]-1-N-[N-{(L)-Alanyl}-(L)-Valinyl]-(2S)-pyrrolidine trifluoroacetate

and other pharmaceutically acceptable salts thereof. These compounds are disclosed in WO 03/033524.

10 Further, according to the present invention substituted aminoketones can be used in combination with effectors of QC. Preferred substituted aminoketones are

- 1-cyclopentyl-3-methyl-1-oxo-2-pentanaminium chloride,
- 1-cyclopentyl-3-methyl-1-oxo-2-butanaminium chloride,
- 1-cyclopentyl-3,3-dimethyl-1-oxo-2-butanaminium chloride,
- 15 • 1-cyclohexyl-3,3-dimethyl-1-oxo-2-butanaminium chloride,
- 3-(cyclopentylcarbonyl)-1,2,3,4-tetrahydroisoquinolinium chloride,
- N-(2-cyclopentyl-2-oxoethyl)cyclohexanaminium chloride

and other pharmaceutically acceptable salts thereof.

20 Among the rare group of proline-specific proteases, DP IV was originally believed to be the only membrane-bound enzyme specific for proline as the penultimate residue at the amino-terminus of the polypeptide chain. However, other molecules, even those structurally non-homologous with the DP IV but bearing corresponding enzyme activity, have been identified. DP IV-like enzymes, which have been identified so far,

25 include e.g. fibroblast activation protein α , dipeptidyl peptidase IV β , dipeptidyl aminopeptidase-like protein, N-acetylated α -linked acidic dipeptidase, quiescent cell proline dipeptidase, dipeptidyl peptidase II, attractin and dipeptidyl peptidase IV related protein (DPP 8), DPL1 (DPX, DP6), DPL2 and DPP 9 described in review articles by Sedo & Malik (Sedo and Malik 2001, *Biochim Biophys Acta*, 36506, 1-10)

30 and Abbott and Gorrell (Abbott, C.A. and Gorrell, M.D. 2002 In: Langner & Ansorge (ed.), *Ecto-peptidases*. Kluwer Academic/Plenum Publishers, New York, 171-195). Recently, the cloning and characterization of dipeptidyl peptidase 10 (DPP 10) was

reported (Qi, S.Y. et al. Biochemical Journal Immediate Publication. Published on 28 Mar 2003 as manuscript BJ20021914).

Effectors, as that term is used herein, are defined as molecules that bind to enzymes and increase or decrease their activity *in vitro* and/or *in vivo*. Some enzymes have binding sites for small molecules that affect their catalytic activity; a stimulator molecule is called an activator. Enzymes may even have multiple sites for recognizing more than one activator or inhibitor. Enzymes can detect concentrations of a variety of molecules and use that information to vary their own activities.

Effectors can modulate enzymatic activity because enzymes can assume both active and inactive conformations: activators are positive effectors, inhibitors are negative effectors. Effectors act not only at the active sites of enzymes, but also at regulatory sites, or allosteric sites, terms used to emphasize that the regulatory site is an element of the enzyme distinct from the catalytic site and to differentiate this form of regulation from competition between substrates and inhibitors at the catalytic site (Darnell, J., Lodish, H. and Baltimore, D. 1990, Molecular Cell Biology 2nd Edition, Scientific American Books, New York, page 63).

Preferred effectors according to the present invention are inhibitors of QC-and EC-activity. Most preferred are competitive inhibitors of QC-and EC-activity.

Where appropriate, activators of QC- and EC-activity are preferred.

In the peptides of the present invention, each amino acid residue is represented by a one-letter or a three-letter designation, corresponding to the trivial name of the amino acid, in accordance with the following conventional list:


<u>Amino Acid</u>	<u>One-Letter Symbol</u>	<u>Three-Letter Symbol</u>
Alanine	A	Ala
Arginine	R	Arg
Asparagine	N	Asn
Aspartic acid	D	Asp
Cysteine	C	Cys

	Glutamine	Q	Gln
	Glutamic acid	E	Glu
	Glycine	G	Gly
	Histidine	H	His
5	Isoleucine	I	Ile
	Leucine	L	Leu
	Lysine	K	Lys
	Methionine	M	Met
	Phenylalanine	F	Phe
10	Proline	P	Pro
	Serine	S	Ser
	Threonine	T	Thr
	Tryptophan	W	Trp
	Tyrosine	Y	Tyr
15	Valine	V	Val

The term "QC" as used herein comprises glutaminy cyclase (QC) and QC-like enzymes. QC and QC-like enzymes have identical or similar enzymatic activity, further defined as QC activity. In this regard, QC-like enzymes can fundamentally differ in their molecular structure from QC.


The term "QC activity" as used herein is defined as intramolecular cyclization of N-terminal glutamine residues into pyroglutamic acid (pGlu*) or of N-terminal L-homoglutamine or L- β -homoglutamine to a cyclic pyro-homoglutamine derivative under liberation of ammonia. See schemes 1 and 2.

Scheme 1: Cyclization of glutamine by QC



Scheme 2: Cyclization of L-homoglutamine by QC

5



The term "EC" as used herein comprises the side activity of QC and QC-like enzymes as glutamate cyclase (EC), further defined as EC activity.


10

The term "EC activity" as used herein is defined as intramolecular cyclization of N-terminal glutamate residues into pyroglutamic acid (pGlu*) by QC. See scheme 3.

The term "metal-dependent enzyme" as used herein is defined as enzyme(s) that require a bound metal ion in order to fulfill their catalytic function and/or require a bound metal ion in order to form the catalytically active structure.

15

Scheme 3: *N*-terminal cyclization of uncharged glutamyl peptides by QC (EC)



Another aspect of the present invention is the identification of new physiological substrates of QC. These were identified by performing cyclization experiments with mammalian peptides as described in example 5. Human QC and papaya QC were isolated as described in example 1. The applied methods are described in example 2, and the peptide synthesis employed is outlined in example 6. The results of the study are shown in Table 1.

Table 1: New physiological substrates of glutaminy cyclase

Substrate	Human QC			Papaya QC		
	K_M (μM)	k_{cat} (s^{-1})	k_{cat}/K_M ($\text{mM}^{-1} \text{s}^{-1}$)	K_M (μM)	k_{cat} (s^{-1})	k_{cat}/K_M ($\text{mM}^{-1} \text{s}^{-1}$)
[Gln ¹]-Gastrin	31 ± 1	54.1 ± 0.6	1745.2 ± 36.9	34 ± 2	25.8 ± 0.5	759 ± 30
[Gln ¹]-Neurotensin	37 ± 1	48.8 ± 0.4	1318.9 ± 24.8	40 ± 3	35.7 ± 0.9	893 ± 44
[Gln ¹]-FPP	87 ± 2	69.6 ± 0.3	800.0 ± 14.9	232 ± 9	32.5 ± 0.4	140 ± 4
[Gln ¹]-TRH	90 ± 4	82.8 ± 1.2	920.0 ± 27.6	n.d.	n.d.	n.d.
[Gln ¹]-GnRH	53 ± 3	69.2 ± 1.1	1305.7 ± 53.2	169 ± 9	82.5 ± 1.9	488.2 ± 14.8
[Gln ³]-glucagon(3-29)			*			*
[Gln ⁵]-substance P(5-11)			*			*

* determined qualitatively by MALDI-TOF experiments

All analyses were performed in the optimal range of activity and stability of either human or plant QC, as demonstrated in example 4.

The amino acid sequences of physiological active peptides having a glutamine residue at the N-terminus and being therefore substrates for the QC enzyme are listed in Table 2.

Table 2: Amino acid sequences of physiological active peptides with a glutamine residue at the N-terminus, that is converted posttranslationally into pyroglutamic acid (pGlu)

Peptide	Amino acid sequence	Function
Gastrin 17 Swiss-Prot: P01350	QGPWL EEEEEAYGWM DF (amide)	Gastrin stimulates the stomach mucosa to produce and secrete hydrochloric acid and the pancreas to secrete its digestive enzymes. It also stimulates smooth muscle contraction and increases blood circulation and water secretion in the stomach and intestine.
Neurotensin Swiss-Prot: P30990	QLYENKPRRP YIL	Neurotensin plays an endocrine or paracrine role in the regulation of fat metabolism. It causes contraction of smooth muscle.
FPP	QEP amide	A tripeptide related to thyrotrophin releasing hormone (TRH), is found in seminal plasma. Recent evidence obtained <i>in vitro</i> and <i>in vivo</i> showed that FPP plays an important role in regulating sperm fertility.
TRH Swiss-Prot: P20396	QHP amide	TRH functions as a regulator of the biosynthesis of TSH in the anterior pituitary gland and as a neurotransmitter/neuromodulator in the central and peripheral nervous systems.
GnRH Swiss-Prot: P01148	QHWSYGL RP(G) amide	Stimulates the secretion of gonadotropins; it stimulates the secretion of both luteinizing and follicle-stimulating hormones.
CCL16 (small	QPKVPEW VNTPSTCCLK YYEKVLPRL VVGYSKALNC	Shows chemotactic activity for lymphocytes and

inducible cytokine A16) Swiss-Prot: O15467	HLPALFVTK RNREVCTNPN DDWVQEYIKD PNLPLLPTRN LSTVKIITAK NGQPQLLSQ	monocytes but not neutrophils. Also shows potent myelosuppressive activity, suppresses proliferation of myeloid progenitor cells. Recombinant SCYA16 shows chemotactic activity for monocytes and THP-1 monocytes, but not for resting lymphocytes and neutrophils. Induces a calcium flux in THP-1 cells that were desensitized by prior expression to RANTES.
CCL8 (small inducible cytokine A8) Swiss-Prot: P80075	QPDSVSI PITCCFNVIN RKIPQRLS YTRITNIQCP KEAVIFKTKR GKEVCADPKE RWVRDSMKHL DQIFQNLKP	Chemotactic factor that attracts monocytes, lymphocytes, basophils and eosinophils. May play a role in neoplasia and inflammatory host responses. This protein can bind heparin.
CCL2 (small inducible cytokine A2) Swiss-Prot: P13500	QPDAINA PVTCCYNFTN RKISVQRLAS YRRITSSKCP KEAVIFKTIV AKEICADPKQ KWVQDSMDHL DKQTQTPKT	Chemotactic factor that attracts monocytes and basophils but not neutrophils or eosinophils. Augments monocyte anti-tumor activity. Has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis or atherosclerosis. May be involved in the recruitment of monocytes into the arterial wall during the disease process of atherosclerosis. Binds to CCR2 and CCR4.
CCL18 (small inducible cytokine A18) Swiss-Prot: P55774	QVGTKNKELC CLVYTSWQIP QKFVDYSET SPQCPKPGVI LLTKRGRQIC ADPNKKWVQK YISDLKLNA	Chemotactic factor that attracts lymphocytes but not monocytes or granulocytes. May be involved in B cell migration into B cell follicles in lymph nodes. Attracts naive T lymphocytes toward dendritic cells and activated

		macrophages in lymph nodes, has chemotactic activity for naive T cells, CD4+ and CD8+ T cells and thus may play a role in both humoral and cell-mediated immunity responses.
Fractalkine (neurotactin) Swiss-Prot: P78423	QHHGVT KCNITCSKMT SKIPVALLIH YQQNQASCGK RAIILETRQH RLFCADPKEQ WVKDAMQHLD RQAAALTRNG GTFEKQIGEV KPRTPPAAGG MDESVVLEPE ATGESSSLEP TPSSQEAQRA LGTSPELPTG VTGSSGTRL PTPKAQDGGP VGTELFRVPP VSTAATWQSS APHQPGPSLW AEAKTSEAPS TQDPSTQAST ASSPAPEENA PSEGQRVWGQ GQSPRPENSL EREEMGPVPA HTDAFQDWGP GSMAGVSVVP VSSEGTPSRE PVASGSWTPK AEEPIHATMD PQRLGVLITP VPDAQAATRR QAVGLLAFLG LLFCLGVAMF TYQSLQGCPR KMAGEMAEGL RYIPRSCGSN SYVLVPV	The soluble form is chemotactic for T cells and monocytes, but not for neutrophils. The membrane-bound form promotes adhesion of those leukocytes to endothelial cells. May play a role in regulating leukocyte adhesion and migration processes at the endothelium. Binds to cx3cr1.
CCL7 (small inducible cytokine A7) Swiss-Prot: P80098	QPVGIN TSTCCYRFIN KKIPKQRLES YRRTTSSHCP REAVIFKTKL DKEICADPTQ KWVQDFMKHL DKKTQTPKL	Chemotactic factor that attracts monocytes and eosinophils, but not neutrophils. Augments monocyte anti-tumor activity. Also induces the release of gelatinase B. This protein can bind heparin. Binds to CCR1, CCR2 and CCR3.
Orexin^A (Hypocretin-1) Swiss-Prot O43612	QPLPDCCRQK TCSCRLYELL HGAGNHAAGI LTL	Neuropeptide that plays a significant role in the regulation of food intake and sleep-wakefulness, possibly by coordinating the complex behavioral and physiologic responses of these complementary homeostatic functions. It plays also a broader role in the homeostatic regulation of energy metabolism,

		autonomic function, hormonal balance and the regulation of body fluids. Orexin-A binds to both OX1R and OX2R with a high affinity.
Substance P	RPK PQQFFGLM (cyclization of Gln ⁵ after cleavage of residues 1-4)	Belongs to the tachykinins. Tachykinins are active peptides which excite neurons, evoke behavioral responses, are potent vasodilators and secretagogues, and contract (directly or indirectly) many smooth muscles.

In a fourth embodiment, the peptides Gln¹-Gastrin (17 and 34 amino acids in length), Gln¹-Neurotensin and Gln¹-FPP were identified as new physiological substrates of QC. Gastrin, Neurotensin and FPP comprise a pGlu residue in their N-terminal position. This N-terminal pGlu residue was shown to be formed from N-terminal glutamine by QC catalysis for all peptides. As a result, these peptides are activated in terms of their biological function upon conversion of the glutamine residue at the N-terminus to pGlu.

Transepithelial transducing cells, particularly the gastrin (G) cell, co-ordinate gastric acid secretion with the arrival of food in the stomach. Recent work showed that multiple active products are generated from the gastrin precursor, and that there are multiple control points in gastrin biosynthesis. Biosynthetic precursors and intermediates (progastrin and Gly-gastrins) are putative growth factors; their products, the amidated gastrins, regulate epithelial cell proliferation, the differentiation of acid-producing parietal cells and histamine-secreting enterochromaffin-like (ECL) cells, and the expression of genes associated with histamine synthesis and storage in ECL cells, as well as acutely stimulating acid secretion. Gastrin also stimulates the production of members of the epidermal growth factor (EGF) family, which in turn inhibit parietal cell function but stimulate the growth of surface epithelial cells. Plasma gastrin concentrations are elevated in subjects with

Helicobacter pylori, who are known to have increased risk of duodenal ulcer disease and gastric cancer (Dockray, G.J. 1999 *J Physiol* 15, 315-324).

The peptide hormone gastrin, released from antral G cells, is known to stimulate the synthesis and release of histamine from ECL cells in the oxyntic mucosa via CCK-2 receptors. The mobilized histamine induces acid secretion by binding to the H(2) receptors located on parietal cells. Recent studies suggest that gastrin, in both its fully amidated and less processed forms (progastrin and glycine-extended gastrin), is also a growth factor for the gastrointestinal tract. It has been established that the major trophic effect of amidated gastrin is for the oxyntic mucosa of the stomach, where it causes increased proliferation of gastric stem cells and ECL cells, resulting in increased parietal and ECL cell mass. On the other hand, the major trophic target of the less processed gastrin (e.g. glycine-extended gastrin) appears to be the colonic mucosa (Koh, T.J. and Chen, D. 2000 *Regul Pept* 9, 337-44).

In a fifth embodiment, the present invention provides the use of activity increasing effectors of QC for the stimulation of gastrointestinal tract cell proliferation, especially gastric mucosal cell proliferation, epithelial cell proliferation, the differentiation of acid-producing parietal cells and histamine-secreting enterochromaffin-like (ECL) cells, and the expression of genes associated with histamine synthesis and storage in ECL cells, as well as for the stimulation of acute acid secretion in mammals by maintaining or increasing the concentration of active [pGlu¹] Gastrin.

In a sixth embodiment, the present invention provides the use of activity decreasing effectors of QC for the treatment of duodenal ulcer disease, gastric cancer with or w/o *Helicobacter pylori*, colorectal cancer, and Zollinger-Ellison Syndrome in mammals by decreasing the conversion rate of inactive Gln¹-Gastrin to active [pGlu¹]Gastrin.

Neurotensin (NT) is a neuropeptide implicated in the pathophysiology of schizophrenia that specifically modulates neurotransmitter systems previously demonstrated to be misregulated in this disorder. Clinical studies in which cerebrospinal fluid (CSF) NT concentrations have been measured revealed a subset

of schizophrenic patients with decreased CSF NT concentrations that are restored by effective antipsychotic drug treatment. The involvement of NT systems in the mechanism of action of antipsychotic drugs is known. The behavioral and biochemical effects of centrally administered NT resemble those of systemically administered antipsychotic drugs, and antipsychotic drugs increase NT neurotransmission. Consequently, NT functions as an endogenous antipsychotic. Moreover, typical and atypical antipsychotic drugs differentially alter NT neurotransmission in nigrostriatal and mesolimbic dopamine terminal regions, and these effects are predictive of side effect liability and efficacy, respectively (Binder, E. B. et al. 2001 *Biol Psychiatry* 50, 856-872).

In a seventh embodiment, the present invention provides the use of activity increasing effectors of QC for the preparation of antipsychotic drugs and/or for the treatment of schizophrenia in mammals. The effectors of QC either maintain or increase the concentration of active [pGlu¹]neurotensin.

Fertilization promoting peptide (FPP), a tripeptide related to thyrotrophin releasing hormone (TRH), is found in seminal plasma. Recent evidence obtained *in vitro* and *in vivo* showed that FPP plays an important role in regulating sperm fertility. Specifically, FPP initially stimulates nonfertilizing (uncapacitated) spermatozoa to "switch on" and become fertile more quickly, but then arrests capacitation so that spermatozoa do not undergo spontaneous acrosome loss and therefore do not lose fertilizing potential. These responses are mimicked, and indeed augmented, by adenosine, known to regulate the adenylyl cyclase (AC)/cAMP signal transduction pathway. Both FPP and adenosine have been shown to stimulate cAMP production in uncapacitated cells but inhibit it in capacitated cells, with FPP receptors somehow interacting with adenosine receptors and G proteins to achieve regulation of AC. These events affect the tyrosine phosphorylation state of various proteins, some being important in the initial "switching on," others possibly being involved in the acrosome reaction itself. Calcitonin and angiotensin II, also found in seminal plasma, have similar effects *in vitro* on uncapacitated spermatozoa and can augment responses to FPP. These molecules have similar effects *in vivo*, affecting fertility by stimulating and then maintaining fertilizing potential. Either reductions in the

availability of FPP, adenosine, calcitonin, and angiotensin II or defects in their receptors contribute to male infertility (Fraser, L. R. and Adeoya-Osiguwa, S. A. 2001 *Vitam Horm* 63, 1-28).

- 5 In an eighth embodiment, the present invention provides the use of activity lowering effectors of QC for the preparation of fertilization prohibitive drugs and/or for the preparation of drugs, which reduce the fertility in mammals. The activity lowering effectors of QC decrease the concentration of active [pGlu¹]FPP, leading to a prevention of sperm capacitation and deactivation of sperm cells. In contrast it could
10 be shown that activity increasing effectors of QC are able to stimulate fertility in males and to treat infertility.

In a ninth embodiment, further physiological substrates of QC were identified within the present invention. These are Gln¹-CCL2, Gln¹-CCL7, Gln¹-CCL8, Gln¹-CCL16,
15 Gln¹-CCL18 and Gln¹-fractalkine. For details see Table 2. These polypeptides play an important role in pathophysiological conditions, such as suppression of proliferation of myeloid progenitor cells, neoplasia, inflammatory host responses, cancer, psoriasis, rheumatoid arthritis, atherosclerosis, humoral and cell-mediated immunity responses, leukocyte adhesion and migration processes at the endothelium
20 and inflammatory processes related to Alzheimers disease, FBD and FDD.

Several cytotoxic T lymphocyte peptide-based vaccines against hepatitis B, human immunodeficiency virus and melanoma were recently studied in clinical trials. One interesting melanoma vaccine candidate alone or in combination with other tumor
25 antigens, is the decapeptide ELA. This peptide is a Melan-A/MART-1 antigen immunodominant peptide analog, with an N-terminal glutamic acid. It has been reported that the amino group and gamma-carboxylic group of glutamic acids, as well as the amino group and gamma-carboxamide group of glutamines, condense easily to form pyroglutamic derivatives. To overcome this stability problem, several peptides
30 of pharmaceutical interest have been developed with a pyroglutamic acid instead of N-terminal glutamine or glutamic acid, without loss of pharmacological properties. Unfortunately compared with ELA, the pyroglutamic acid derivative (PyrELA) and also the N-terminal acetyl-capped derivative (AcELA) failed to elicit cytotoxic T

lymphocyte (CTL) activity. Despite the apparent minor modifications introduced in PyrELA and AcELA, these two derivatives probably have lower affinity than ELA for the specific class I major histocompatibility complex. Consequently, in order to conserve full activity of ELA, the formation of PyrELA must be avoided (Beck, A. et al. 2001, *J Pept Res* 57, 528-38.). Recently, it was found that also the enzyme glutaminy cyclase (QC) is overexpressed in melanomas (Ross, D. T. et al. 2000, *Nat Genet* 24, 227-35.).

In a tenth embodiment, the present invention provides the use of effectors of QC for the preparation of a medicament for the treatment of pathophysiological conditions, such as suppression of proliferation of myeloid progenitor cells, neoplasia, inflammatory host responses, cancer, malign metastasis, melanoma, psoriasis, rheumatoid arthritis, atherosclerosis, impaired humoral and cell-mediated immunity responses, leukocyte adhesion and migration processes at the endothelium and inflammatory processes related to Alzheimers disease, FBD and FDD.

In an eleventh embodiment, Gln¹-orexin A was identified as a physiological substrate of QC within the present invention. Orexin A is a neuropeptide that plays a significant role in the regulation of food intake and sleep-wakefulness, possibly by coordinating the complex behavioral and physiologic responses of these complementary homeostatic functions. It plays also a role in the homeostatic regulation of energy metabolism, autonomic function, hormonal balance and the regulation of body fluids.

In a twelfth embodiment, the present invention provides the use of effectors of QC for the preparation of a medicament for the treatment of impaired food intake and sleep-wakefulness, impaired homeostatic regulation of energy metabolism, impaired autonomic function, impaired hormonal balance and impaired regulation of body fluids.

Polyglutamine expansions in several proteins lead to neurodegenerative disorders, such as Parkinson disease and Kennedy's disease. The mechanism therefore remains largely unknown. The biochemical properties of polyglutamine repeats suggest one possible explanation: endolytic cleavage at a glutaminy-glutaminy bond followed by pyroglutamate formation may contribute to the pathogenesis

through augmenting the catabolic stability, hydrophobicity, amyloidogenicity, and neurotoxicity of the polyglutaminy proteins (Saido, T. 2000 *Med Hypotheses* 54, 427-9).

- 5 In a thirteenth embodiment, the present invention provides therefore the use of effectors of QC for the preparation of a medicament for the treatment of Parkinson disease and Huntington's disease.

10 In a fourteenth embodiment, the present invention provides a general way to reduce or inhibit the enzymatic activity of QC. Examples of inhibitory compounds are also provided.

Inhibition of a mammalian QC was only detected initially for 1,10-phenanthroline and reduced 6-methylpterin (Busby, W. H. J. et al. 1987 *J Biol Chem* 262, 8532-8536).
15 EDTA did not inhibit QC, thus it was concluded that QC is not a metal-dependent enzyme (Busby, W. H. J. et al. 1987 *J Biol Chem* 262, 8532-8536, Bateman, R. C. J. et al. 2001 *Biochemistry* 40, 11246-11250, Booth, R. E. et al. 2004 *BMC Biology* 2). In the present invention, however, it is shown that human QC and other animal QCs are metal-dependent enzymes, as revealed by the inhibition characteristics of QC by
20 1,10-phenanthroline, dipicolinic acid, 8-hydroxy-quinoline and other chelators (Figures 18, 19) and by the reactivation of QC by transition metal ions (Figure 20). Finally, the metal dependence is outlined by a sequence comparison to other metal-dependent enzymes, showing a conservation of the chelating amino acid residues also in human QC (Figure 21). The interaction of compounds with the active-site
25 bound metal ion represents a general way to reduce or inhibit QC activity.

In the present invention it is shown that imidazole derivatives are potent inhibitors of QC. Using the continuous assay (for details see example 2), many imidazole derivatives were analyzed concerning their ability to inhibit the human QC as a member of the highly conserved mammalian QCs.

30

Thus, the present invention provides imidazole derivatives and histidine and its derivatives as activity reducing effectors of QC and their characteristics in terms of inhibition type and potency. Structures and K_i -values are shown in tables 3 and 4. The results are described in detail in example 7.

Table 3: Inhibitory constants of imidazole derivatives in the human QC catalyzed reaction. Determinations were performed at 30 °C in 0.05 M Tris-HCl pH 8.0, containing 5 mM EDTA.

5


Compound	K _i -value (mM)	Structure
core structures		
imidazole	0.103 ±0.004	
benzimidazole	0.138 ±0.005	
N-1 derivatives		
1-benzylimidazole	0.0071 ±0.0003	
1-methylimidazole	0.030 ±0.001	
1-vinylimidazole	0.049 ±0.002	
oxalic acid diimidazolidide	0.078 ±0.002	
N-acetylimidazole	0.107 ±0.003	
N-(trimethylsilyl)-imidazole	0.167 ±0.007	
N-benzoylimidazole	0.174 ±0.007	
1-(2-oxo-2-phenyl-ethyl)-imidazole	0.184 ±0.005	
1-(3-aminopropyl)-imidazole	0.41 ±0.01	
1-phenylimidazole	no inhibition	
1,1'-sulfonyldiimidazole	no inhibition	
C-4(5) derivatives		
N-omega-acetylhistamine	0.017 ±0.001	
L-histidinamide	0.56 ±0.04	
H-His-Trp-OH	0.60 ±0.03	
L-histidinol	1.53 ±0.12	
L-histidine	4.4 ±0.2	
4-imidazole-carboxaldehyde	7.6 ±0.7	
imidazole-4-carbonic acid methylester	14.5 ±0.6	
L-histamine	0.85 ±0.04	
C-4,5 derivatives		
5-hydroxymethyl-4-methyl-imidazole	0.129 ±0.005	
4-amino-imidazole-5-carbonic acid amide	15.5 ±0.5	
4,5-diphenyl-imidazole	no inhibition	
4,5-dicyanoimidazole	no inhibition	
C-2 derivatives		
2-methyl-benzylimidazole	0.165 ±0.004	
2-ethyl-4-methyl-imidazole	0.58 ±0.04	
2-aminobenzimidazole	1.8 ±0.1	

2-chloro-1H-benzimidazole


no inhibition

Others


3-(1H-imidazol-1-yl)-1-(3-methylbenzo[b]thiophene-2-yl)propan-1-one

0.0025 $\pm$ 0.0001


4-[(1-methyl-1H-imidazol-5-yl)methyl]-3-propyldihydrofuran-2-(3H)-one

0.0067 $\pm$ 0.0003


4-[2-(1H-imidazol-1-yl)ethoxy]benzoic acid

0.0034 $\pm$ 0.0001




3-[3-(1H-imidazol-1-yl)propyl]-2-thioxoimidazolidin-4-one

0.00081 $\pm$ 0.00001

5-nitro-2-[2-([3-(1H-imidazol-1-yl)propyl]amino)carbonyl]phenyl]furanamide

0.0066 $\pm$ 0.0004

N-(4-chlorophenyl)-N'-[2-(1H-imidazol-1-yl)ethyl]thiourea

0.00165 $\pm$ 0.000070,0322 $\pm$ 0,0007

n.d.

Imidazo<1.5a>pyridine



0.0356 $\pm$ 0.0005Methyl (2*S*)-2-[[[(2*S*)-2-amino-5-(1*H*-imidazol-1-ylamino)-5-oxopentanoyl]amino]-3-methylbutanoate0.164 $\pm$ 0.004

Table 4: QC inhibition by L-histamine and its two biological metabolites (also known as *te/e*-methylhistamine).


Compound	K _i value (mM)	Structure
L-histamine	0.85 $\pm$ 0.04	
3-methyl-4-(β -aminoethyl)-imidazole	0.120 $\pm$ 0.004	
1-methyl-4-(β -aminoethyl)-imidazole	n.i.	

In a fifteenth embodiment, new inhibitors of QC and QC-like enzymes, based on cysteamine, are provided.

Besides imidazole derivatives and hydroxamates, thiol reagents are frequently described as inhibitors of metal-dependent enzymes (Lowther, W. T. and Matthews,

B. W. 2002 *Chem Rev* 102, 4581-4607; Lipscomb, W. N. and Sträter, N. 1996 *Chem Rev* 96, 2375-2433). Additionally, thiol peptides were described as inhibitors of a QC-related aminopeptidase of the Clan MH (Huntington, K. M. et al. 1999 *Biochemistry* 38, 15587-15596). Although these inhibitors are inactive with regard to mammalian QCs, it was possible to isolate cysteamine derivatives as potent competitive QC-inhibitors (Figure 23).

The present invention provides QC-inhibitors which can be described generally by the formula 1 or the pharmaceutically acceptable salts thereof, including all stereoisomers:



formula 1

wherein R¹ – R⁶ are independently H or a branched or unbranched alkyl chain, a branched or unbranched alkenyl chain, a branched or unbranched alkynyl chain, carbocyclic, aryl, heteroaryl, heterocyclic, aza-amino acid, amino acid or a mimetic thereof, peptide or a mimetic thereof; all of the above residues optionally being substituted, and

n is 0, 1 or 2, preferably 1, most preferably 0.

Throughout the description and the claims the expression "alkyl" can denote a C₁₋₅₀ alkyl group, preferably a C₁₋₃₀ alkyl group, especially a C₁₋₁₂ or C₁₋₈ alkyl group; for example, an alkyl group may be a methyl, ethyl, propyl, isopropyl or butyl group. The expression "alk", for example in the expression "alkoxy", and the expression "alkan", for example in the expression "alkanoyl", are defined as for "alkyl"; aromatic ("aryl") compounds are preferably substituted or optionally unsubstituted phenyl, benzyl, naphthyl, biphenyl or anthracene groups, which preferably have at least 8 C atoms; the expression "alkenyl" can denote a C₂₋₁₀ alkenyl group, preferably a C₂₋₆ alkenyl group, which has the double bond(s) at any desired location and may be substituted or unsubstituted; the expression "alkynyl" can denote a C₂₋₁₀ alkynyl group, preferably

a C₂₋₆ alkynyl group, which has the triple bond(s) at any desired location and may be substituted or unsubstituted; the expression "substituted" or substituent can denote any desired substitution by one or more, preferably one or two, alkyl, alkenyl, alkynyl, mono- or multi-valent acyl, alkanoyl, alkoxyalkanoyl or alkoxyalkyl groups; the afore-
5 mentioned substituents may in turn have one or more (but preferably zero) alkyl, alkenyl, alkynyl, mono- or multi-valent acyl, alkanoyl, alkoxyalkanoyl or alkoxyalkyl groups as side groups.

Throughout the description and the claims the expression "acyl" can denote a C₁₋₂₀
10 acyl residue, preferably a C₁₋₈ acyl residue and especially preferred a C₁₋₄ acyl residue; and "carbocyclic" can denote a C₃₋₁₂ carbocyclic residue, preferably a C₄, C₅ or C₆ carbocyclic residue. "Heteroaryl" is defined as an aryl residue, wherein 1 to 4, and more preferably 1, 2 or 3 ring atoms are replaced by heteroatoms like N, S or O. "Heterocyclic" is defined as a cycloalkyl residue, wherein 1, 2 or 3 ring atoms are
15 replaced by heteroatoms like N, S or O.



"Peptide mimetics" per se are known to a person skilled in the art. They are preferably defined as compounds which have a secondary structure like a peptide and optionally further structural characteristics; their mode of action is largely similar or identical to the mode of action of the native peptide; however, their activity (e.g. as
20 an antagonist or inhibitor) can be modified as compared with the native peptide, especially vis à vis receptors or enzymes. Moreover, they can imitate the effect of the native peptide (agonist). Examples of peptide mimetics are scaffold mimetics, non-peptidic mimetics, peptoides, peptide nucleic acids, oligopyrrolinones, vinylogpeptides and oligocarbamates. For the definitions of these peptide mimetics
25 see Lexikon der Chemie, Spektrum Akademischer Verlag Heidelberg, Berlin, 1999.

An "aza-amino acid" is defined as an amino acid where the chiral α -CH group is replaced by a nitrogen atom, whereas an "aza-peptide" is defined as a peptide, in which the chiral α -CH group of one or more amino acid residues in the peptide chain
30 is replaced by a nitrogen atom.

The role of the thiol and amino group is outlined by a comparison of the inhibitory potency of dimethylcysteamine, cysteamine, mercaptoethanol, ethylenediamine,

ethanolamine as shown in table 5. Only compounds bearing amino and thiol group were potent, loss or modification of either group led to a decrease in inhibitory power. Moreover, the pH-dependence of inhibition of murine QC by cysteamine, dimethyl-cysteamine and mercaptoethanol revealed differences, indicating that the protonation state of the inhibitor influences inhibitor binding to the active site (Figure 22).

Table 5 Comparison of the potency of cysteamine-derived compounds to inhibit QC (n.i., no inhibition detected at a concentration of 5 mM, pH=8.0, 30°C and a substrate concentration of 1 K_M in the sample, n.d., not determined)

Compound	K_i -value (mM)	Structure
cysteamine	0.043 ± 0.002	
dimethyl-cysteamine	0.0190 ± 0.0002	
diethyl-cysteamine	0.0109 ± 0.0004	
mercaptoethanol	3.91 ± 0.08	
ethylmercaptane	n.d.	
ethylenediamine	n.i.	
ethanolamine	n.i.	

Surprisingly, during characterization of the enzymatic activity it was discovered that besides an N-terminal glutamyl residue, β -homo-glutamyl residues at the N-terminus also fulfill properties as substrates of QCs from plants and mammals. The N-terminal β -homo-glutamyl residue was converted into a five-membered lactam ring by catalysis of human and papaya QC, respectively. The results are described in

example 5. The applied method is illustrated in example 2 and the peptide synthesis was performed as described in example 6.

Another preferred embodiment of the present invention comprises screening
5 methods for effectors of QC.

A preferred screening method for identifying activity modifying effectors of QC from a group of compounds comprises the steps of:

- 10 a) Contacting said compounds with QC under conditions which would permit binding therebetween;
- b) Adding a substrate of QC;
- c) Monitoring the conversion of the substrate or optionally measuring the residual QC activity; and
- 15 d) Calculating changes in the substrate conversion and/or enzyme activity of QC to identify an activity modifying effector.

Another preferred screening method relates to a method for the identification and selection of effectors which interact directly or indirectly with the active-site bound
20 metal ion of QC and comprises the following steps:

- a) Contacting said compounds with QC under conditions which would permit binding therebetween;
- b) Adding a substrate of QC which is subject to conversion by QC;
- c) Monitoring the conversion of the substrate or optionally measuring the
25 residual QC activity; and
- d) Calculating changes in the substrate conversion and/or enzyme activity of QC wherein changes may be used to identify an activity modifying effector of QC.

30 Preferred for the use in the above described screening methods are mammalian QC or *Papaya* QC. Especially preferred is mammalian QC, since the effectors identified by these screening methods shall be used for the treatment of diseases in mammals, especially in humans.

The agents selected by the above described screening methods can work by decreasing the conversion of at least one substrate of QC (negative effectors, inhibitors), or by increasing the conversion of at least one substrate of QC (positive effectors, activators).

5

The compounds of the present invention can be converted into acid addition salts, especially pharmaceutically acceptable acid addition salts.

10

The salts of the compounds of the invention may be in the form of inorganic or organic salts.

15

20

The compounds of the present invention can be converted into and used as acid addition salts, especially pharmaceutically acceptable acid addition salts. The pharmaceutically acceptable salt generally takes a form in which a basic side chain is protonated with an inorganic or organic acid. Representative organic or inorganic acids include hydrochloric, hydrobromic, perchloric, sulfuric, nitric, phosphoric, acetic, propionic, glycolic, lactic, succinic, maleic, fumaric, malic, tartaric, citric, benzoic, mandelic, methanesulfonic, hydroxyethanesulfonic, benzenesulfonic, oxalic, pamoic, 2-naphthalenesulfonic, p-toulenesulfonic, cyclohexanesulfamic, salicylic, saccharinic or trifluoroacetic acid. All pharmaceutically acceptable acid addition salt forms of the compounds of the present invention are intended to be embraced by the scope of this invention.

25

In view of the close relationship between the free compounds and the compounds in the form of their salts, whenever a compound is referred to in this context, a corresponding salt is also intended, provided such is possible or appropriate under the circumstances.

30

Where the compounds according to this invention have at least one chiral center, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as diastereomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention. Furthermore, some of the crystalline forms of the compounds may exist as polymorphs and as such are intended to be included in the

present invention. In addition, some of the compounds may form solvates with water (i.e. hydrates) or common organic solvents, and such solvates are also intended to be encompassed within the scope of this invention.

- 5 The compounds, including their salts, can also be obtained in the form of their hydrates, or include other solvents used for their crystallization.

10 In a further embodiment, the present invention provides a method of preventing or treating a condition mediated by modulation of the QC enzyme activity in a subject in need thereof which comprises administering any of the compounds of the present invention or pharmaceutical compositions thereof in a quantity and dosing regimen therapeutically effective to treat the condition. Additionally, the present invention includes the use of the compounds of this invention, and their corresponding pharmaceutically acceptable acid addition salt forms, for the
15 preparation of a medicament for the prevention or treatment of a condition mediated by modulation of the QC activity in a subject. The compound may be administered to a patient by any conventional route of administration, including, but not limited to, intravenous, oral, subcutaneous, intramuscular, intradermal, parenteral and combinations thereof.

20 In a further preferred form of implementation, the invention relates to pharmaceutical compositions, that is to say, medicaments, that contain at least one compound of the invention or salts thereof, optionally in combination with one or more pharmaceutically acceptable carriers and/or solvents.

25 The pharmaceutical compositions may, for example, be in the form of parenteral or enteral formulations and contain appropriate carriers, or they may be in the form of oral formulations that may contain appropriate carriers suitable for oral administration. Preferably, they are in the form of oral formulations.

30 The effectors of QC activity administered according to the invention may be employed in pharmaceutically administrable formulations or formulation complexes as inhibitors of QC- and EC-activity, preferably competitive inhibitors, or in combination with enzyme inhibitors, competitive enzyme inhibitors, substrates,

pseudosubstrates, inhibitors of QC expression, binding proteins or antibodies of those enzyme proteins that reduce the QC protein concentration in mammals. The compounds of the invention make it possible to adjust treatment individually to patients and diseases, it being possible, in particular, to avoid individual intolerances, allergies and side-effects.

The compounds also exhibit differing degrees of activity as a function of time. The doctor providing treatment is thereby given the opportunity to respond differently to the individual situation of patients: he is able to adjust precisely, on the one hand, the speed of the onset of action and, on the other hand, the duration of action and especially the intensity of action.

A preferred treatment method according to the invention represents a new approach for the prevention or treatment of a condition mediated by modulation of the QC enzyme activity in mammals. It is advantageously simple, susceptible of commercial application and suitable for use, especially in the treatment of diseases that are based on unbalanced concentration of physiological active QC substrates, e.g. listed in Tables 1 and 2, in mammals and especially in human medicine.

The compounds may be advantageously administered, for example, in the form of pharmaceutical preparations that contain the active ingredient in combination with customary additives like diluents, excipients and/or carriers known from the prior art. For example, they can be administered parenterally (for example *i.v.* in physiological saline solution) or enterally (for example orally, formulated with customary carriers).

Depending upon their endogenous stability and their bioavailability, one or more doses of the compounds can be given per day in order to achieve the desired normalisation of the blood glucose values. For example, such a dosage range in humans may be in the range of from about 0.01 mg to 250.0 mg per day, preferably in the range of from about 0.01 to 100 mg of compound per kilogram of body weight.

By administering effectors of QC activity to a mammal it could be possible to prevent or alleviate or treat conditions selected from Alzheimer's disease, Down Syndrome, Familial British Dementia (FBD), Familial Danish Dementia (FDD), ulcer disease and

gastric cancer with or w/o *Helicobacter pylori* infections, colorectal cancer, Zollinger-Ellison Syndrome, pathogenic psychotic conditions, schizophrenia, infertility, neoplasia, inflammatory host responses, cancer, psoriasis, rheumatoid arthritis, atherosclerosis, impaired humoral and cell-mediated immune responses, leukocyte
5 adhesion and migration processes in the endothelium, impaired food intake, sleep-wakefulness, impaired homeostatic regulation of energy metabolism, impaired autonomic function, impaired hormonal balance and impaired regulation of body fluids.

10 Further, by administering effectors of QC activity to a mammal it could be possible to stimulate gastrointestinal tract cell proliferation, preferably proliferation of gastric mucosal cells, epithelial cells, acute acid secretion and the differentiation of acid producing parietal cells and histamine-secreting enterochromaffin-like cells.

15 In addition, administration of QC inhibitors to mammals may lead to a loss of sperm cell function thus suppressing male fertility. Thus, the present invention provides a method for the regulation and control of male fertility and the use of activity lowering effectors of QC for the preparation of contraceptive medicaments for males.

20 Furthermore, by administering effectors of QC activity to a mammal it may be possible to suppress the proliferation of myeloid progenitor cells.

The compounds used according to the invention can accordingly be converted in a manner known *per se* into conventional formulations, such as, for example, tablets,
25 capsules, dragées, pills, suppositories, granules, aerosols, syrups, liquid, solid and cream-like emulsions and suspensions and solutions, using inert, non-toxic, pharmaceutically suitable carriers and additives or solvents. In each of those formulations, the therapeutically effective compounds are preferably present in a concentration of approximately from 0.1 to 80 % by weight, more preferably from 1 to
30 50 % by weight, of the total mixture, that is to say, in amounts sufficient for the mentioned dosage latitude to be obtained.

The substances can be used as medicaments in the form of dragées, capsules, bitable capsules, tablets, drops, syrups or also as suppositories or as nasal sprays.

The formulations may be advantageously prepared, for example, by extending the active ingredient with solvents and/or carriers, optionally with the use of emulsifiers and/or dispersants, it being possible, for example, in the case where water is used as diluent, for organic solvents to be optionally used as auxiliary solvents.

Examples of excipients useful in connection with the present invention include: water, non-toxic organic solvents, such as paraffins (for example natural oil fractions), vegetable oils (for example rapeseed oil, groundnut oil, sesame oil), alcohols (for example ethyl alcohol, glycerol), glycols (for example propylene glycol, polyethylene glycol); solid carriers, such as, for example, natural powdered minerals (for example highly disperse silica, silicates), sugars (for example raw sugar, lactose and dextrose); emulsifiers, such as non-ionic and anionic emulsifiers (for example polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, alkylsulphonates and arylsulphonates), dispersants (for example lignin, sulphite liquors, methylcellulose, starch and polyvinylpyrrolidone) and lubricants (for example magnesium stearate, talcum, stearic acid and sodium lauryl sulphate) and optionally flavourings.

Administration may be carried out in the usual manner, preferably enterally or parenterally, especially orally. In the case of enteral administration, tablets may contain in addition to the mentioned carriers further additives such as sodium citrate, calcium carbonate and calcium phosphate, together with various additives, such as starch, preferably potato starch, gelatin and the like. Furthermore, lubricants, such as magnesium stearate, sodium lauryl sulphate and talcum, can be used concomitantly for tableting. In the case of aqueous suspensions and/or elixirs intended for oral administration, various taste correctives or colourings can be added to the active ingredients in addition to the above-mentioned excipients.

In the case of parenteral administration, solutions of the active ingredients using suitable liquid carriers can be employed. In general, it has been found advantageous to administer, in the case of intravenous administration, amounts of approximately from 0.01 to 2.0 mg/kg, preferably approximately from 0.01 to 1.0 mg/kg, of body weight per day to obtain effective results and, in the case of enteral administration,

the dosage is approximately from 0.01 to 2 mg/kg, preferably approximately from 0.01 to 1 mg/kg, of body weight per day.

It may nevertheless be necessary in some cases to deviate from the stated amounts, depending upon the body weight of the experimental animal or the patient or upon the type of administration route, but also on the basis of the species of animal and its individual response to the medicament or the interval at which administration is carried out. Accordingly, it may be sufficient in some cases to use less than the above-mentioned minimum amount, while, in other cases, the mentioned upper limit will have to be exceeded. In cases where relatively large amounts are being administered, it may be advisable to divide those amounts into several single doses over the day. For administration in human medicine, the same dosage latitude is provided. The above remarks apply analogously in that case.

15 Examples of pharmaceutical formulations

1. Capsules containing 100 mg of a compound of the invention per capsule:

For approximately 10,000 capsules a solution of the following composition is prepared:

compound of the invention	1.0 kg
glycerol	0.5 kg
polyethylene glycol	3.0 kg
25 water	<u>0.5 kg</u>
	5.0 kg

The solution is introduced into soft gelatin capsules in a manner known *per se*. The capsules are suitable for chewing or swallowing.

30

2. Tablets or coated tables or dragées containing 100 mg of a compound of the invention:

The following amounts refer to the preparation of 100,000 tablets:

compound of the invention,

finely ground	10.0 kg
glucose	4.35 kg
5 lactose	4.35 kg
starch	4.50 kg
cellulose, finely ground	4.50 kg

The above constituents are mixed and then provided with a solution prepared from

10

polyvinylpyrrolidone	2.0 kg
polysorbate	0.1 kg
and water	approx. 5.0 kg

15 and granulated in a manner known *per se* by grating the moist mass and, after the addition of 0.2 kg of magnesium stearate, drying it. The finished tablet mixture of 30.0 kg is processed to form convex tablets weighing 300 mg. Ideally, the tablets can be coated or sugar-coated in a manner known *per se*.

20 The pharmaceutical compositions defined throughout the specification advantageously contain a combination of at least one effector of QC activity and at least one DP IV inhibitor. Such pharmaceutical compositions are especially useful for the treatment of Alzheimer's Disease and Down Syndrome.

25

Example 1: Preparation of Human and Papaya QC

Host strains and media

30 *Pichia pastoris* strain X33 (AOX1, AOX2), used for the expression of human QC was grown, transformed and analyzed according to the manufacturer's instructions (Invitrogen). The media required for *P. pastoris*, i.e. buffered glycerol (BMGY)

complex or methanol (BMMY) complex medium, and the fermentation basal salts medium were prepared according to the manufacturer's recommendations.

Molecular cloning of plasmid vectors encoding the human QC

5 All cloning procedures were done applying standard molecular biology techniques. For expression in yeast, the vector pPICZ α B (Invitrogen) was used. The pQE-31 vector (Qiagen) was used to express the human QC in *E. coli*. The cDNA of the mature QC starting with codon 38 was fused in frame with the plasmid encoded 6 $\times$ histidine tag. After amplification utilizing the primers pQCyc-1 and pQCyc-2 and
10 subcloning, the fragment was inserted into the expression vector employing the restriction sites of *Sph*I and *Hind*III.

Transformation of *P. pastoris* and mini-scale expression

Plasmid DNA was amplified in *E. coli* JM109 and purified according to the
15 recommendations of the manufacturer (Qiagen). In the expression plasmid used, pPICZ α B, three restriction sites are provided for linearization. Since *Sac*I and *Bst*XI cut within the QC cDNA, *Pme*I was chosen for linearization. 20-30 μ g plasmid DNA was linearized with *Pme*I, precipitated by ethanol, and dissolved in sterile, deionized water. 10 μ g of the DNA was then applied for transformation of competent *P. pastoris*
20 cells by electroporation according to the manufacturer's instructions (BioRad). Selection was done on plates containing 150 μ g/ml Zeocin. One transformation using the linearized plasmid yielded several hundred transformants.

In order to test the recombinant yeast clones for QC expression, recombinants were
25 grown for 24 h in 10 ml conical tubes containing 2 ml BMGY. Afterwards, the yeast was centrifuged and resuspended in 2 ml BMMY containing 0.5 % methanol. This concentration was maintained by addition of methanol every 24 h up to 72 h. Subsequently, QC activity in the supernatant was determined. The presence of the fusion protein was confirmed by western blot analysis using an antibody directed
30 against the 6 $\times$ histidine tag (Qiagen). Clones that displayed the highest QC activity were chosen for further experiments and fermentation.

Large-scale expression in a fermenter

Expression of the QC was performed in a 5 l reactor (Biostat B, B. Braun biotech), essentially as described in the "*Pichia* fermentation process guidelines" (Invitrogen).

5 Briefly, the cells were grown in the fermentation basal salts medium supplemented with trace salts, and with glycerol as the sole carbon source (pH 5.5). During an initial batch phase for about 24 h and a subsequent fed-batch phase for about 5 h, cell mass was accumulated. Once a cell wet weight of 200 g/l was achieved, induction of QC expression was performed using methanol applying a three-step feeding profile
10 for an entire fermentation time of approximately 60 h. Subsequently, cells were removed from the QC-containing supernatant by centrifugation at 6000×g, 4 °C for 15 min. The pH was adjusted to 6.8 by addition of NaOH, and the resultant turbid solution was centrifuged again at 37000×g, 4 °C for 40 min. In cases of continued turbidity, an additional filtration step was applied using a cellulose membrane (pore
15 width 0.45 µm).

Purification of 6×histidine tagged QC expressed in *P. pastoris*

The His-tagged QC was first purified by immobilized metal affinity chromatography (IMAC). In a typical purification, 1000 ml of culture supernatant were applied to a
20 Ni²⁺-loaded Chelating Sepharose FF column (1.6 × 20 cm, Pharmacia), that was equilibrated with 50 mM phosphate buffer, pH 6.8, containing 750 mM NaCl, at a flow rate of 5 ml/min. After washing with 10 column volumes of equilibration buffer and 5 column volumes of equilibration buffer containing 5 mM histidine, the bound protein was eluted by a shift to 50 mM phosphate buffer, pH 6.8, containing 150 mM NaCl
25 and 100 mM histidine. The resulting eluate was dialyzed against 20 mM Bis-Tris/HCl, pH 6.8, at 4 °C overnight. Subsequently, the QC was further purified by anion exchange chromatography on a Mono Q6 column (BioRad), equilibrated with dialysis buffer. The QC-containing fraction was loaded onto the column using a flow rate of 4 ml/min. The column was then washed with equilibration buffer containing 100 mM
30 NaCl. The elution was performed by two gradients resulting in equilibration buffer containing 240 mM and 360 mM NaCl in 30 or 5 column volumes, respectively. Fractions of 6 ml were collected and the purity was analyzed by SDS-PAGE.

Fractions containing homogenous QC were pooled and concentrated by ultrafiltration. For long-term storage (-20°C), glycerol was added to a final concentration of 50 %. Protein was quantified according to the methods of Bradford or Gill and von Hippel (Bradford, M. M. 1976 *Anal Biochem* 72, 248-254; Gill, S. C. and von Hippel, P. H. 1989 *Anal Biochem* 182, 319-326.).

Expression and purification of QC in *E. coli*

The construct encoding the QC was transformed into M15 cells (Qiagen) and grown on selective LB agar plates at 37°C . Protein expression was carried out in LB medium containing 1 % glucose and 1 % ethanol at room temperature. When the culture reached an OD_{600} of approximately 0.8, expression was induced with 0.1 mM IPTG overnight. After one cycle of freezing and thawing, cells were lysed at 4°C by addition of 2.5 mg/ml lysozyme in 50 mM phosphate buffer, pH 8.0, containing 300 mM NaCl and 2 mM histidine for approximately 30 min. The solution was clarified by centrifugation at $37000\times g$, 4°C for 30 min, followed by a filtration applying a glass frit (DNA separation) and two additional filtration steps applying cellulose filters for crude and fine precipitates. The supernatant (approx. 500 ml) was applied onto a Ni^{2+} -affinity column (1.6 x 20 cm) at a flow rate of 1 ml/min. Elution of QC was carried out with 50 mM phosphate buffer containing 150 mM NaCl and 100 mM histidine. The QC-containing fraction was concentrated by ultrafiltration.

Purification of QC from papaya latex

QC from papaya latex was prepared using the BioCAD 700E (Perseptive Biosystems, Wiesbaden, Germany) with a modified version of a previously reported method (Zerhouni, S. et al. 1989 *Biochim Biophys Acta* 138, 275-290). 50 g latex was dissolved in water and centrifugated as described therein. Inactivation of proteases was performed with S-methyl methanethiosulfonate, and the resultant crude extract was dialyzed. After dialysis, the entire supernatant was loaded onto a (2152.5 cm i.d.) SP Sepharose Fast Flow column, equilibrated with 100 mM sodium acetate buffer, pH 5.0 (flow rate 3 ml/min). Elution was performed in three steps by increasing sodium acetate buffer concentration at a flow rate of 2 ml/min. The first step was a linear gradient from 0.1 to 0.5 M acetate buffer in 0.5 column volumes.

The second step was a linear increase in buffer concentration from 0.5 to 0.68 M in four column volumes. During the last elution step, one column volume of 0.85 M buffer was applied. Fractions (6 ml) containing the highest enzymatic activity were pooled. Concentration and buffer changes to 0.02 M Tris/HCl, pH 8.0 were performed
5 via ultrafiltration (Amicon; molecular mass cut-off of the membrane 10 kDa).

Ammonium sulfate was added to the concentrated papaya enzyme, obtained from the ion exchange chromatography step to a final concentration of 2 M. This solution was applied onto a (2152.5 cm i.d.) Butyl Sepharose*4 Fast Flow column (flow rate
10 1.3 ml/min), equilibrated with 2 M ammonium sulfate, 0.02 M Tris/HCl, pH 8.0. Elution was performed in three steps with decreasing concentrations of ammonium sulfate. During the first step a linear gradient from 2 to 0.6 M ammonium sulfate, 0.02 M Tris/HCl, pH 8.0 was applied for 0.5 column volumes at a flow rate of 1.3 ml/min. The second step was a linear gradient from 0.6 to 0 M ammonium sulfate, 0.02 M
15 Tris/HCl, pH 8.0, in 5 column volumes at a flow rate of 1.5 ml/min. The last elution step was carried out by applying 0.02 M Tris/HCl at pH 8.0 for 2 column volumes at a flow rate of 1.5 ml/min. All fractions containing QC activity were pooled and concentrated by ultrafiltration. The resultant homogenous QC was stored at -70°C. Final protein concentrations were determined using the method of Bradford,
20 compared to a standard curve obtained with bovine serum albumin.

Example 2: Assays for glutaminy cyclase activity

25 Fluorometric assays

All measurements were performed with a BioAssay Reader HTS-7000Plus for microplates (Perkin Elmer) at 30 °C. QC activity was evaluated fluorometrically using H-Gln- β NA. The samples consisted of 0.2 mM fluorogenic substrate, 0.25 U pyroglutamyl aminopeptidase (Unizyme, Hørsholm, Denmark) in 0.2 M Tris/HCl, pH
30 8.0 containing 20 mM EDTA and an appropriately diluted aliquot of QC in a final volume of 250 μ l. Excitation/emission wavelengths were 320/410 nm. The assay reactions were initiated by addition of glutaminy cyclase. QC activity was determined from a standard curve of β -naphthylamine under assay conditions. One unit is

***Trade-mark**

defined as the amount of QC catalyzing the formation of 1 μ mol pGlu- β NA from H-Gln- β NA per minute under the described conditions.

In a second fluorometric assay, QC activity was determined using H-Gln-AMC as substrate. Reactions were carried out at 30°C utilizing the NOVOStar reader for microplates (BMG labtechnologies). The samples consisted of varying concentrations of the fluorogenic substrate, 0.1 U pyroglutamyl aminopeptidase (Qiagen) in 0.05 M Tris/HCl, pH 8.0 containing 5 mM EDTA and an appropriately diluted aliquot of QC in a final volume of 250 μ l. Excitation/emission wavelengths were 380/460 nm. The assay reactions were initiated by addition of glutaminy cyclase. QC activity was determined from a standard curve of 7-amino-4-methylcoumarin under assay conditions. The kinetic data were evaluated using GraFit software.

Spectrophotometric assay of QC

This novel assay was used to determine the kinetic parameters for most of the QC substrates. QC activity was analyzed spectrophotometrically using a continuous method, that was derived by adapting a previous discontinuous assay (Bateman, R. C. J. 1989 *J Neurosci Methods* 30, 23-28) utilizing glutamate dehydrogenase as auxiliary enzyme. Samples consisted of the respective QC substrate, 0.3 mM NADH, 14 mM α -Ketoglutaric acid and 30 U/ml glutamate dehydrogenase in a final volume of 250 μ l. Reactions were started by addition of QC and pursued by monitoring of the decrease in absorbance at 340 nm for 8-15 min. Typical time courses of product formation are presented in Figure 1.

The initial velocities were evaluated and the enzymatic activity was determined from a standard curve of ammonia under assay conditions. All samples were measured at 30°C, using either the SPECTRAFluor Plus* or the Sunrise* (both from TECAN) reader for microplates. Kinetic data was evaluated using GraFit* software.

*Trade-mark

Inhibitor assay

For inhibitor testing, the sample composition was the same as described above, except of the putative inhibitory compound added. For a rapid test of QC-inhibition, samples contained 4 mM of the respective inhibitor and a substrate concentration at 1 K_M . For detailed investigations of the inhibition and determination of K_i -values, influence of the inhibitor on the auxiliary enzymes was investigated first. In every case, there was no influence on either enzyme detected, thus enabling the reliable determination of the QC inhibition. The inhibitory constant was evaluated by fitting the set of progress curves to the general equation for competitive inhibition using GraFit software.

Example 3: MALDI-TOF mass spectrometry

Matrix-assisted laser desorption/ionization mass spectrometry was carried out using the Hewlett-Packard G2025 LD-TOF System with a linear time of flight analyzer. The instrument was equipped with a 337 nm nitrogen laser, a potential acceleration source (5 kV) and a 1.0 m flight tube. Detector operation was in the positive-ion mode and signals were recorded and filtered using LeCroy 9350M digital storage oscilloscope linked to a personal computer. Samples (5 μ l) were mixed with equal volumes of the matrix solution. For matrix solution we used DHAP/DAHC, prepared by solving 30 mg 2',6'-dihydroxyacetophenone (Aldrich) and 44 mg diammonium hydrogen citrate (Fluka) in 1 ml acetonitrile/0.1% TFA in water (1/1, v/v). A small volume ($\approx$ 1 μ l) of the matrix-analyte-mixture was transferred to a probe tip and immediately evaporated in a vacuum chamber (Hewlett-Packard G2024A sample prep accessory) to ensure rapid and homogeneous sample crystallization.

For long-term testing of Glu¹-cyclization, A β -derived peptides were incubated in 100 μ l 0.1 M sodium acetate buffer, pH 5.2 or 0.1 M Bis-Tris buffer, pH 6.5 at 30°C. Peptides were applied in 0.5 mM [A β 3-11a] or 0.15 mM [A β 3-21a] concentrations, and 0.2 U QC was added all 24 hours. In case of A β (3-21)a, the assays contained 1 % DMSO. At different times, samples were removed from the assay tube, peptides extracted using ZipTips (Millipore) according to the manufacturer's recommendations, mixed with matrix solution (1:1 v/v) and subsequently the mass spectra recorded. Negative controls did either contain no QC or heat deactivated

enzyme. For the inhibitor studies the sample composition was the same as described above, with exception of the inhibitory compound added (5 mM benzimidazole or 2 mM 1,10-phenanthroline).

5

Example 4: pH dependence

The pH-dependence of catalysis of human and papaya QC was investigated under first-order rate conditions, thus reflecting the impact of the proton concentration on the specificity constant k_{cat}/K_M . For this purpose, the coupled enzymatic assay using pyroglutamyl aminopeptidase as auxiliary enzyme and Gln- β NA as substrate was used. Pyroglutamyl aminopeptidase was shown to be active and stable between pH 5.5-8.5 (Tsuru, D. et al. 1978 *J Biochem (Tokyo)* 84, 467-476). Hence, the assay enabled the study of QC catalysis in this pH-region. The rate profiles obtained were fit to classical bell shaped curves, as shown in Figure 2. The human QC bears a very narrow pH-dependence with an optimum at about pH 7.8-8.0. The rate tended to decrease at more basic pH. This is in contrast to the rate profile observed with papaya QC, which showed no drop in activity up to pH 8.5 (Figure 2, inset). However, both enzymes had their optimum of specificity at pH 8. Surprisingly, evaluation of the curves revealed identical pK_a -values in the acidic range of 7.17 ± 0.02 and 7.15 ± 0.02 for human and papaya QC, respectively.

The reduction of the activity of human QC at basic pH-values was obviously due to dissociation of a group with a pK_a of approximately 8.5. In case of papaya QC, there was no excessive data point collection in the basic pH-range possible to enable a reliable determination of the second pK_a value. This is supported by fitting of the data to a single dissociation model, resulting in an almost identical pK_a -value (pK_a 7.13 ± 0.03) compared to fitting the data to a double dissociation model. This indicates that both pK_a -values are fairly separated..

30

pH stability

The stability of the glutamyl cyclases was investigated by incubating the plant and animal enzymes at 30 °C for 30 min at different pH values between pH 4-10. Afterwards, QC activity was determined under standard conditions. The results are shown in Figure 3.

The QC from papaya latex was stable in the pH-range studied, without an obvious trend for instability in the acidic or basic range. In contrast, human QC only showed a comparable stability in the pH-range between 7 and 8.5, exhibiting a remarkable instability at pH values above pH 8.5 and below pH 6. Thus, the region around pH 8 seems to be optimal for activity and stability of plant and human QC and a suitable pH-value for performing a substrate specificity comparison of the QCs.

Example 5: Determination of substrate specificity of QC

Spectrophotometric assay

The continuous spectrophotometric assay was performed as described in example 2. Accordingly, QC activity is reflected by a decrease in absorbance at 340 nm caused by the ammonia release and subsequent consumption of NADH/H⁺ due to formation of glutamate from α -ketoglutaric acid. As shown in Figure 1, linear progress curves were monitored and there was a linear relationship between the measured activity and the concentration of QC. Furthermore, the kinetic parameters obtained for H-Gln-Gln-OH using the continuous assay presented here (Table 6) were in good agreement with those obtained using the discontinuous method ($K_M = 175 \pm 18 \mu M$, $k_{cat} = 21.3 \pm 0.6 s^{-1}$). In addition, the kinetic parameters for conversion of the substrates H-Gln-Ala-OH, H-Gln-Glu-OH, H-Gln-Gln-OH, H-Gln-OtBu and H-Gln-NH₂ by papaya QC shown in Table 1 correspond well to those determined using a direct method at pH 8.8 and 37 °C (Gololobov, M. Y. et al. 1996 *Biol Chem Hoppe Seyler* 377, 395-398). Hence, it is quite obvious that the novel continuous assay provides reliable results.

Di-, Tri- and Dipeptide-surrogates

Utilizing the novel continuous assay described above, about 30 compounds were tested as potential substrates of QC from *C. papaya* and human. The results are displayed in Table 6. By comparison of the specificities it was shown, that nearly all of the short peptide substrates are more efficiently converted by papaya QC compared to the human enzyme. Interestingly, for both enzymes substrates with large hydrophobic residues in the second position are the most potent ones, as shown by the specificities of H-Gln-Tyr-Ala-OH, H-Gln-Phe-Ala-NH₂ and H-Gln-Trp-Ala-NH₂ compared to those of other tripeptides or by the reactivities of the chromophoric substrates H-Gln-AMC, H-Gln- β NA and H-Gln-Tyr-OH in comparison to dipeptide substrates. For papaya QC, this finding is in agreement with earlier results showing that the specificity is in correlation with the size of the second amino acid residue (Gololobov, M. Y. et al. 1996 *Biol Chem Hoppe Seyler* 377, 395-398). The only striking difference in specificity of the plant and animal QC was observed in case of H-Gln-OtBu. Whereas the ester was converted by papaya QC with similar specificity compared to dipeptide substrates, it was converted about one order of magnitude slower by human QC.

Oligopeptides

Besides several dipeptides and tripeptides, a number of oligopeptides was tested upon conversion by papaya and human QC (Table 6). Interestingly, the overall difference in the specificities between human and plant QC for a set of tetrapeptides was not that large as it was observed for dipeptide and tripeptide substrates. This indicates that the amino acids in the 3rd and 4th position still affect the kinetic behavior especially of human QC. An exception, however, comprise the peptides with a proline residue in the second amino acid position which show noticeably reduced k_{cat}/K_M values in a set of tetrapeptides of the structure H-Gln-X_{aa}-Tyr-Phe-NH₂ (Table 6). The reduction in specificity was more pronounced for human QC, leading to an approximately 8-fold difference in the k_{cat}/K_M -value as compared to papaya QC.

Slightly reduced specificities of human QC were also observed for conversion of substrates with a positively charged amino acid C-terminal of glutamine, as indicated by the specificities for H-Gln-Arg-Tyr-Phe-NH₂, H-Gln-Arg-Tyr-Phe-NH₂ and H-Gln-

Lys-Arg-Leu-NH₂ as compared to other tetrapeptides. Apparently, the reduced specificity was mainly due to a smaller turnover number. This effect was not the case for the plant enzyme.

5

Table 6: Kinetic evaluation of peptide substrates of human and Papaya QC

(n.r., not reactive; n.i., no inhibition; n.d., not determined; *, for substrate inhibition)

Substrate	Human QC				Papaya QC			
	K _M (μM)	k _{cat} (s ⁻¹)	K _i * (mM)	k _{cat} /K _M (mM ⁻¹ s ⁻¹)	K _M (μM)	k _{cat} (s ⁻¹)	K _i * (mM)	k _{cat} /K _M (mM ⁻¹ s ⁻¹)
H-Gln-OH	n.r.	n.r.	n.d.	n.r.	n.d.	n.d.	n.d.	0.23 ±0.1
H-Gln-AMC	54 ±2	5.3 ±0.1	n.d.	98 ±2	42 ±1	39.4 ±0.4	n.d.	938 ±13
H-Gln-βNA	70 ±3	20.6 ±0.5	1.21 ±0.07	294 ±6	38 ±3	51.4 ±1.4	1.20 ±0.08	1353 ±70
H-Gln-OtBu	1235 ±74	6.7 ±0.2	n.i.	5.4 ±0.2	223 ±9	49.4 ±0.6	n.i.	222 ±6
H-Gln-NH ₂	409 ±40	12.8 ±0.5	n.i.	31 ±2	433 ±13	44.8 ±0.4	n.i.	103 ±2
H-Gln-Gly-OH	247 ±10	13.2 ±0.2	n.i.	53 ±1	641 ±20	45.8 ±0.4	n.i.	71 ±2
H-Gln-Ala-OH	232 ±5	57.2 ±0.4	n.i.	247 ±4	158 ±8	69.8 ±1.0	n.i.	442 ±16
H-Gln-Gln-OH	148 ±5	20.7 ±0.2	n.i.	140 ±2	44 ±3	43.2 ±0.7	n.i.	982 ±51
H-Gln-Glu-OH	359 ±10	24.7 ±0.2	n.i.	58 ±1	106 ±5	50.3 ±0.6	n.i.	475 ±17
H-Gln-Val-OH	196 ±5	17.2 ±0.1	n.i.	88 ±2	n.d.	n.d.	n.i.	n.d.
H-Gln-Tyr-OH	211 ±5	94 ±1	n.i.	446 ±6	n.d.	n.d.	n.i.	n.d.
H-Gln-Glu-Tyr-NH ₂	79 ±2	45.1 ±0.4	n.i.	524 ±8	103 ±4	53.6 ±0.7	n.i.	520 ±13
H-Gln-Gly-Pro-OH	130 ±5	25.3 ±0.2	n.i.	195 ±7	333 ±15	41.7 ±0.5	n.i.	125 ±4
H-Gln-Tyr-Ala-OH	101 ±4	125 ±1	n.i.	930 ±27	63 ±3	104.0 ±1.0	n.i.	1650 ±63
H-Gln-Phe-Ala-NH ₂	69 ±3	109 ±1	n.i.	1811 ±64	111 ±5	132.1 ±0.6	n.i.	1190 ±48
H-Gln-Trp-Ala-NH ₂	50 ±2	47.0 ±0.7	n.i.	940 ±24	78 ±5	151.8 ±2.6	n.i.	1946 ±91
H-Gln-Arg-Gly-Ile-NH ₂	143 ±4	33.5 ±0.4	n.i.	234 ±4	123 ±10	49.2 ±1.7	n.i.	400 ±19
H-Gln-Asn-Gly-Ile-NH ₂	172 ±5	56.6 ±0.5	n.i.	329 ±7	153 ±9	51.4 ±0.9	n.i.	336 ±14
H-Gln-Ser-Tyr-Phe-NH ₂	55 ±3	52.8 ±0.8	n.i.	960 ±38	135 ±6	64.9 ±1.0	n.i.	481 ±14
H-Gln-Arg-Tyr-Phe-NH ₂	55 ±2	29.6 ±0.3	n.i.	538 ±14	124 ±6	48.9 ±0.7	n.i.	394 ±13
H-Gln-Pro-Tyr-Phe-NH ₂	1889 ±152	31.7 ±1.2	n.i.	17 ±1	149 ±14	18.8 ±0.6	n.i.	126 ±8
H-Gln-His-Tyr-Phe-NH ₂	68 ±3	55.4 ±0.7	n.i.	815 ±26	92 ±7	75.9 ±1.4	n.i.	825 ±48
H-Gln-Gln-Tyr-Phe-NH ₂	41 ±2	41.4 ±0.4	n.i.	1010 ±40	45 ±2	52.9 ±0.7	n.i.	1176 ±37
H-Gln-Glu-Tyr-Phe-NH ₂	47 ±4	46 ±1	n.i.	979 ±62	100 ±4	54.6 ±0.6	n.i.	546 ±16
H-Gln-Glu-Ala-Ala-NH ₂	77 ±4	46 ±1	n.i.	597 ±18	102 ±4	53.7 ±0.6	n.i.	526 ±15
H-Gln-Glu-Tyr-Ala-NH ₂	69 ±2	42.1 ±0.4	n.i.	610 ±12	113 ±5	44.7 ±0.5	n.i.	396 ±13
H-Gln-Glu-Ala-Phe-NH ₂	39 ±3	39 ±1	n.i.	1000 ±51	81 ±3	48.5 ±0.45	n.i.	599 ±17
H-Gln-Glu-Asp-Leu-NH ₂	55 ±2	45.8 ±0.5	n.i.	833 ±21	107 ±6	58.5 ±0.4	n.i.	547 ±27
H-Gln-Lys-Arg-Leu-NH ₂	54 ±3	33.4 ±0.5	n.i.	619 ±25	118 ±6	48.2 ±0.8	n.i.	408 ±14

10 The results obtained with the tetrapeptides give also rise to another conclusion. As already pointed out, papaya QC showed a higher selectivity for dipeptides. For some of the tetrapeptides, however, higher specificity-constants were observed with human

QC, as shown in Figure 4 providing a plot of the data given in Table 6, for a set of peptides containing glutamate in the second amino acid position. Furthermore, as the chain length increases from di- to tetrapeptides, the selectivity of human QC increases, in contrast to the results obtained with papaya QC. Additionally, the highest selectivity of human QC was recorded for the peptides containing bulky hydrophobic residues in the 3rd and 4th amino acid position, which indicate hydrophobic interactions with the enzyme. By comparison of the kinetic parameters for the respective peptides, the changes seem to be mainly due to lower K_M -values, the turnover numbers for conversion of the peptides were found to be similar. Thus, the higher selectivity of human QC for longer peptides is considered to be the result of tighter binding of the more hydrophobic substrates to the enzyme.

The differences between human and plant QC observed with peptides containing hydrophobic amino acids in the 3rd and 4th position becomes also evident by a comparison of the specificity constants of the enzymes towards H-Gln-Arg-Gly-Ile-NH₂ and H-Gln-Arg-Tyr-Phe-NH₂ or H-Gln-Gln-OH and H-Gln-Gln-Tyr-Phe-OH.

Human QC was also found to be more selective for homologous substrates containing N-terminal Gln and an increasing number of C-terminal Ala residues (Table 7). While the selectivity of human QC increased with substrate length, there was no such a trend with the papaya QC. Since human QC was less specific for a peptide containing a Ser residue in the sequence, also the nature of the side chain seems to be of importance (table 6).

Table 7: Influence of substrate length on the activity of human and Papaya QC

Substrate	Human QC			Papaya QC		
	K_M (μ M)	k_{cat} (s^{-1})	k_{cat}/K_M ($mM^{-1} s^{-1}$)	K_M (μ M)	k_{cat} (s^{-1})	k_{cat}/K_M ($mM^{-1} s^{-1}$)
H-Gln-Ala -NH ₂	155 $\pm$ 9	40.1 $\pm$ 0.9	259 $\pm$ 9	212 $\pm$ 21	62.8 $\pm$ 3.0	296 $\pm$ 15
H-Gln-Ala-Ala-NH ₂	87 $\pm$ 3	76.3 $\pm$ 0.7	877 $\pm$ 22	164 $\pm$ 6	83.2 $\pm$ 1.0	507 $\pm$ 12
H-Gln-Ala-Ala-Ala-Ala-NH ₂	65 $\pm$ 3	60.5 $\pm$ 0.7	1174 $\pm$ 43	197 $\pm$ 8	74.6 $\pm$ 1.0	379 $\pm$ 10
H-Gln-Ala-Ala-Ser-Ala-Ala-NH ₂	79 $\pm$ 6	55.3 $\pm$ 1.6	700 $\pm$ 33	216 $\pm$ 6	78.5 $\pm$ 1.0	363 $\pm$ 5

Influence of ionic strength on catalysis

Another parameter that was investigated concerning its influence on the substrate specificity was ionic strength. For that purpose, the kinetic parameters for cyclization of several substrates were determined in presence and absence of 0.5 M KCl (Table 8). Surprisingly, the selectivity for substrates with uncharged backbone did not change significantly by addition of the salt in case of QC from papaya latex and human QC. The specificity constants of the human QC for H-Gln-Ala-OH and H-Gln-Glu-OH, however, decreased by addition of KCl. As indicated by the individual kinetic parameters, this effect was due to an increasing K_M and an only slightly decreasing k_{cat} -value. In case of papaya QC, there was no effect on either parameter detected. The effect seemed not to be due to the negatively charged substrate as such, since unchanged parameters were found for the negatively charged peptide H-Gln-Glu-Asp-Leu-NH₂. An interesting effect of the salt addition was found for the positively charged substrates H-Gln-Arg-Gly-Ile-NH₂ and H-Gln-Lys-Arg-Leu-NH₂. In case of plant and human QC, a positive effect on catalysis was determined mainly due to a smaller K_M value and a slightly higher turnover number.

Table 8: Influence of ionic strength on catalysis of human and Papaya QC

	Substrate	0.05 M Tricine-NaOH, pH 8.0				0.05 M Tricine-NaOH, pH 8.0, 0.5 M KCl			
		K_M (mM)	k_{cat} (s ⁻¹)	k_{cat}/K_M (mM ⁻¹ s ⁻¹)	K_i (mM)	K_M (mM)	k_{cat} (s ⁻¹)	k_{cat}/K_M (mM ⁻¹ s ⁻¹)	K_i (mM)
Papaya QC	H-Gln-NH ₂	0.434 ±0.015	43.4 ±0.4	100 ±3	n.i.	0.446 ±0.010	45.2 ±0.3	101 ±2	n.i.
	H-Gln-βNA	0.036 ±0.002	48.8 ±1.0	1356 ±50	1.14 ±0.05	0.032 ±0.002	47.2 ±0.8	1475 ±70	1.33 ±0.07
	H-Gln-Ala-OH	0.137 ±0.007	69.7 ±.9	509 ±19	n.i.	0.143 ±0.005	68.1 ±0.6	480 ±12	n.i.
	H-Gln-Glu-OH	0.098 ±0.005	45.0 ±0.5	459 ±18	n.i.	0.094 ±0.003	44.4 ±0.3	472 ±12	n.i.
	H-Gln-Trp-Ala-NH ₂	0.079 ±0.005	138 ±3	1747 ±73	n.i.	0.072 ±0.004	133 ±3	1847 ±61	n.i.
	H-Gln-Arg-Gly-Ile-NH ₂	0.106 ±0.008	52.9 ±1.2	499 ±26	n.i.	0.065 ±0.005	48.4 ±1.0	745 ±42	n.i.
	H-Gln-Lys-Arg-Leu-NH ₂	0.102 ±0.007	50 ±1	493 ±22	n.i.	0.053 ±0.002	58.1 ±0.7	1096 ±28	n.i.
	H-Gln-Glu-Asp-Leu-NH ₂	0.109 ±0.005	52.4 ±0.7	481 ±16	n.i.	0.094 ±0.003	53.6 ±0.5	570 ±13	n.i.
Human QC		0.05 M Tris-HCl, pH 8.0				0.05 M Tris-HCl, pH 8.0, 0.5 M KCl			
	H-Gln-NH ₂	0.442 ±0.030	12.8 ±0.3	29±1	n.i.	0.401 ±0.014	12.2 ±0.1	30±1	n.i.
	H-Gln-βNA	0.076 ±0.004	21.7 ±0.5	285±8	1.39 ±0.08	0.063 ±0.003	20.0 ±0.4	318 ±9	0.97 ±0.04
	H-Gln-Ala-OH	0.269 ±0.007	54.4 ±0.5	202±3	n.i.	0.357 ±0.012	47.6 ±0.6	133 ±3	n.i.
	H-Gln-Glu-OH	0.373 ±0.015	21.4 ±0.3	57±2	n.i.	0.607 ±0.036	18.9 ±0.5	31 ±1	n.i.

H-Gln-Trp-Ala-NH ₂	0.054 ±0.003	50.8 ±0.6	941 ±41	n.i.	0.056 ±0.002	50.0 ±0.4	893±25	n.i.
H-Gln-Arg-Gly-Ile-NH ₂	0.166 ±0.013	31 ±1	187 ±9	n.i.	0.091 ±0.005	29.8 ±0.5	327 ±12	n.i.
H-Gln-Lys-Arg-Leu-NH ₂	0.051 ±0.003	29.4 ±0.5	577 ±24	n.i.	0.034 ±0.001	31.6 ±0.3	929 ±19	n.i.
H-Gln-Glu-Asp-Leu-NH ₂	0.060 ±0.002	46.6 ±0.5	777 ±18	n.i.	0.061 ±0.002	45.6 ±0.5	748 ±16	n.i.

Physiological substrates

- 5 In earlier studies, conversion of [Gln¹]-TRH and [Gln¹]-GnRH by QC was already shown for the QC from bovine and porcine pituitary (Busby, W. H. J. et al. 1987 *J Biol Chem* 262, 8532-8536; Fischer, W. H. and Spiess, J. 1987 *Proc Natl Acad Sci U S A* 84, 3628-3632). In addition to these already investigated pituitary hormones, three potential physiological substrates of human QC were synthesized and tested
- 10 upon conversion, namely Gln¹-Gastrin, Gln¹-Neurotensin, and Gln¹-FPP. The kinetic parameters for their conversion are listed in Table 1. Interestingly, the glutaminyl peptides are converted to the respective pyroglutamyl peptides with increasing specificity constants depending on their size, i.e., the first largest peptide pro-gastrin with 17 amino acids followed by pro-neurotensin, pro-GnRH, pro-TRH and pro-FPP.
- 15 These findings correspond to the data obtained with the synthetic peptides.

Surprisingly, the longer substrates are also converted with higher selectivity by the plant enzyme, a result that contrasts in part with the findings for the shorter oligopeptides. Possibly, there are secondary binding interactions between substrate

20 and enzyme far distant from the active site.

Peptides comprising modified amino acids

In order to further investigate the specificity and selectivity of the QCs, peptides were synthesized comprising either a modified N-terminal glutaminyl residue or a modified

25 amino acid in the second position. The conversion of these peptides was investigated qualitatively utilizing MALDI-TOF mass spectrometry (see also example 3). Due to the cyclization of the glutaminyl residue or its analog, respectively, a mass difference of the substrate and the product of catalysis is detected. In cases of ammonia

liberation of one mole per mole of substrate, the conversion was also analyzed quantitatively using the spectrophotometric assay.

H-Gln-Lys(Gln)-Arg-Leu-Ala-NH₂. This N-terminally branched peptide, comprising two glutamyl residues at the N-terminus that are bound to a lysyl residue via a peptide- and partial isopeptide bond, was converted by human (Figure 5) and papaya QC (not shown) in an apparently identical manner. Both glutamyl residues were converted into pyroglutamic acid, without any detectable preference for a distinct residue, as indicated by the consistent substrate conversion (Figure 5). Thus, the selectivity of the QCs for the differently bound glutamyl residues differs not fundamentally.

H-Gln(NMe)-Phe-Lys-Ala-Glu-NH₂. The methylated glutamyl residue was only converted into a pyroglutamyl residue by papaya QC (Figure 6). Additionally, an inhibition of the human QC by the peptide was not detected, indicating that the methylated residue is not recognized by human QC.

H-Glu(OMe)-βNA and H-Glu-βNA. Neither of these compounds were converted by papaya or human QC. These fluorogenic substrates were analyzed fluorometrically, utilizing pyroglutamyl aminopeptidase as auxiliary enzyme. The O-methylated glutamate residue, however, showed a remarkable instability in both, Tris and Tricine buffers tending to a non-enzymatically catalyzed cyclization. Furthermore, activity of both QCs against H-Gln-AMC as substrate was not inhibited by the longer peptides H-Glu(OMe)-Phe-Lys-Arg-Leu-Ala-NH₂ or H-Glu-Phe-Lys-Arg-Leu-Ala-NH₂, indicating that glutamic acid or derivatives are not recognized by both QC forms. Furthermore, the result implies that not only the negative charge of the glutamic acid residue is the reason for the repulsion of the peptide from the active site.

H-Gln-cyclo(Nε-Lys-Arg-Pro-Ala-Gly-Phe). The conversion of *H-Gln-cyclo(Nε-Lys-Arg-Pro-Ala-Gly-Phe)*, which contains an intramolecular partial isopeptide bond was analyzed quantitatively, revealing K_M -values of $240 \pm 14 \mu\text{M}$ and $133 \pm 5 \mu\text{M}$ for human and papaya QC, respectively. Due to the higher turnover number of

conversion by papaya QC ($49.4 \pm 0.6 \text{ s}^{-1}$) compared to human QC ($22.8 \pm 0.6 \text{ s}^{-1}$), the plant enzyme exhibits with $372 \pm 9 \text{ mM}^{-1} \text{ min}^{-1}$ an approximately 4-fold higher $k_{\text{cat}}/K_{\text{M}}$ -value than the human QC. Thus, the specificity constant is in case of the papaya QC only slightly smaller compared to substrates having a similar size, such as H-Gln-Ala-
5 Ala-Ser-Ala-Ala-NH₂. The $k_{\text{cat}}/K_{\text{M}}$ -value for human QC, however, was found with $95 \pm 3 \text{ mM}^{-1} \text{ s}^{-1}$ to be approximately one order of magnitude smaller in comparison with substrates of similar size (Table 5).

H-βhomoGln-Phe-Lys-Arg-Leu-Ala-NH₂. The N-terminal β-homoglutaminyl
10 residue was converted into a five-membered lactam ring by catalysis of human and papaya QC, respectively. The concomitant liberation of ammonia was analyzed spectrophotometrically and by MALDI-tof analysis as described before. There was no liberation of ammonia detected when QC was omitted or boiled, indicating a specific catalysis of the cyclization. Interestingly, the QC from *C. papaya* ($K_{\text{M}} = 3.1 \pm 0.3 \text{ mM}$,
15 $k_{\text{cat}} = 4.0 \pm 0.4 \text{ s}^{-1}$) and human ($K_{\text{M}} = 2.5 \pm 0.2 \text{ mM}$, $k_{\text{cat}} = 3.5 \pm 0.1 \text{ s}^{-1}$) catalyze the conversion of this peptide with almost identical $k_{\text{cat}}/K_{\text{M}}$ values of 1.4 ± 0.1 and $1.3 \pm 0.1 \text{ mM}^{-1} \text{ s}^{-1}$, respectively. Thus, the cyclization of the β-homoglutamine residue is catalyzed with an approximately 1000-fold reduced efficiency compared to peptides of similar size containing a glutaminyl residue at their N-terminus. This shows that the
20 constitution of the α-carbon of the substrate is important for substrate recognition by the QC forms, but not essential. The essential requirement for being a substrate is a γ-amide group and an unprotonated N-terminal amino group in distance and angle prone for cyclization, a requirement that is fulfilled by N-terminal glutaminyl and β-homo-glutaminyl residues.

Example 6: Synthesis of the QC substrates

Oligopeptides. Peptides were synthesized semiautomatically in 0.5 mmol scale using a peptide synthesizer (Labortec SP650, Bachem, Switzerland) as previously described (Schilling, S. et al. 2002 *Biochemistry* 41, 10849-10857). Longer peptides were synthesized in 25 μ mol scale using the automated Symphony peptide synthesizer (Rainin Instrument Co.) as described (Manhart, S. et al. 2003 *Biochemistry* 42, 3081-3088). For all peptide couplings modified Fmoc-protocols of solid-phase peptide synthesis were employed using 2-(1H-Benzotriazole-1-yl)-1,1,3,3,-tetramethyluronium tetrafluoroborate (TBTU; Novabiochem)/base (diisopropyl ethylamine or N-methyl- morpholine; Merck) or in case of difficult couplings *N*-[(Dimethylamino)-1*H*-1,2,3,-triazolo[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethanaminium hexafluorophosphate *N*-oxide (4,5) (HATU; Applied Biosystems)/ diisopropyl ethylamine as activating reagents were used. After cleavage from the resin by trifluoroacetic acid (TFA; Merck) containing cocktail, the crude peptides were purified by preparative HPLC with acid free solvents in order to avoid further cyclization of the N-terminal glutamine. Preparative HPLC was performed with a linear gradient of acetonitrile (Merck) in water (5-40 % or 65 % acetonitrile over 40 min) on a 250-21 Luna RP18 column (Phenomenex). To confirm peptide purity and identity analytical HPLC and ESI-MS was employed.

Glu(NH-NH₂)-Ser-Pro-Thr-Ala-NH₂. The linear precursor peptide (Fmoc-Glu-Ser-Pro-Thr-Ala-NH₂) was synthesized according to standard Fmoc-procedures (Schilling, S. et al. 2002 *Biochemistry* 41, 10849-10857) on Rink amide MBHA resin (Novabiochem). After cleavage of the Fmoc-protected peptide from the resin, the peptide was precipitated with diethyl ether (Merck), filtered and dried. HMBA-AM resin (1.16 mmol/g, Novabiochem) was used for coupling of the γ -carboxylic acid group of glutamic acid of the precursor peptide (3 eq.) in dichloromethane (DCM, Merck). Dicyclohexylcarbodiimide (DCC, Serva) (4 eq.) and dimethylaminopyridine (DMAP, Aldrich) (0.1 eq) were used as coupling reagents. After 12 hours the resin was filtered, washed with DCM and the reaction was repeated. After deprotection of the N-terminal Fmoc-group by employing 20% piperidine in DMF (3 $\times$ 5 min) the peptide resin was treated with a 5% hydrazine solution (20 ml/g) for 1.5 hours. The

resin was filtered and washed with dimethylformamide (DMF, Roth, Germany) and TFA. Following evaporation, the crude peptide was precipitated with ether giving 76 % yield.

5 *H-Gln-Lys(Gln)-Arg-Leu-Ala-NH₂*. The linear peptide was synthesized according to standard Fmoc/^tBu-procedure on Rink amide MBHA (Schilling, S. et al. 2002 *Biochemistry* 41, 10849-10857) using Fmoc-Lys(Fmoc)-OH as penultimate amino acid coupling. After deprotection of the two amino protecting groups of lysine with 20 % piperidine (Merck) in DMF, 4 eq. Fmoc-Gln(Trt)-OH were coupled.
10 Standard cleavage procedure resulted in 95% yield.

H-Gln(NMe)-Phe-Lys-Ala-Glu-NH₂. Fmoc-Gln(NMe)-OH was synthesized starting from Fmoc-Glu-OtBu loaded on Fmoc-MI-AM (Novabiochem) resin. After swelling with DCM, the resin (0.5 g) was washed with DMF and deprotected with
15 20% piperidine solution in DMF. The resin was given into 5 ml DMF and 5 eq. Fmoc-Glu-OtBu, 5 eq. HATU and 10 eq. DIPEA were added subsequently and shaken for 6 hours. After filtration and washing, the product was cleaved according to standard TFA cleavage conditions. The peptide *H-Gln(NMe)-Phe-Lys-Ala-Glu-NH₂* was synthesized as described (Schilling, S. et al. 2002 *Biochemistry* 41, 10849-10857).
20 Fmoc-Gln(NMe)-OH was coupled with HATU/DIPEA overnight. Standard cleavage procedure resulted in 78% of the crude peptide.

H-Glu(OMe)-β-naphthylamide, *H-Gln-Val-OH*, *H-Gln-Tyr-OH*. Boc-protected dipeptides were synthesized applying standard mixed anhydride procedure by using
25 isobutyl chlorocarbonate (Merck). The C-terminal methylesters Boc-Gln-Tyr-OMe and Boc-Gln-Val-OMe were saponified by 1N NaOH in dioxane. The Boc-protected peptides were deprotected by HCl/dioxane solution for 10 min. After evaporation the residue was crystallized with several solvents giving 60-70% of a solid compound.

30 *H-Gln-cyclo(Nε-Lys-Arg-Pro-Ala-Gly-Phe)*. The linear precursor Boc-Gln(Trt)-Lys-Arg(Pmc)-Ala-Gly-Phe-OH was synthesized on acid sensitive 2-chlorotrityl resin. Coupling was carried out using a standard protocol of Fmoc/^tBu-strategy using Fmoc-Lys(Mtt)-OH. After cleavage with 3% TFA solution in DCM (10 times 5 min), the solution was neutralized with 10% pyridine (Merck) in methanol (MeOH; Merck),

washed 3 times with DCM and MeOH, evaporated to 5 % of the volume and the crude peptide was precipitated with icecold water. Following, the crude peptide was cyclized using DCC/*N*-hydroxybenzotriazole (HOBt; Aldrich) activation. The crude peptide was dissolved in dry dichloromethane (0.2 mmol/50 ml), 0.2 mmol *N*-methylmorpholine and 0.4 mmol 1-hydroxybenzotriazole were added. This solution was added dropwise to a solution of 0.4 mmol dicyclohexylcarbodiimide in 250 ml dichloromethane at 0°C. The reaction was completed by stirring overnight at room temperature. After filtration of *N,N'*-dicyclohexylurea, the solvent was removed by evaporation. The residue was dissolved in ethyl acetate and washed several times with 1N HCl, saturated solution of NaHCO₃ and water. The solution was dried over anhydrous Na₂SO₄, filtered and evaporated to dryness in vacuo.

Example 7: Characterization of effectors of QC

Imidazole derivatives

Imidazole and benzimidazole derivatives carrying substituents in different positions of the 5-membered ring were tested as inhibitors of QC (Table 3). The constitution of the numbers refers to the imidazole ring. The applied methods are described in example 2.

C-4(5) and C-4,5 derivatives. The compounds carrying substitutions in either in the constitutionally equivalent 4- or 5- position of the imidazole ring or in both positions showed a diminished potency for inhibition of human QC. The only exception, however, comprised *N*- ω -acetylated histamine that proved to be one of the most potent inhibitory compounds. Small substituents in these positions had only little effect on binding as indicated by the similar inhibition constant of 5-hydroxymethyl-4-methyl-imidazole compared to imidazole. Larger and more bulky groups attached to these sites diminished or abolished binding of the compound by the enzyme. Some of the other substituents tested are known to exert negative inductive or mesomeric effects that are capable to reduce the electron density in the imidazole ring, which also contributes to poorer binding constants. The difference in the *K_i*-values of L-histidine and histidinamide also indicate some influence of the charge on binding.

Evidence for electrostatic repulsion of charged substrates were already shown in the substrate specificity studies, i.e. glutaminamide was readily converted to products by human QC, but no reactivity was observed for free glutamine as substrate.

5 *C-2 derivatives.* All derivatives tested inhibited QC more weakly as imidazole. Any substitution bigger than a proton hinders proper QC-binding. Only due to the methyl group in 2-methyl-benzimidazole, the inhibition constant drops about one order of magnitude. A very similar relation was shown by comparison of the K_i -values for benzimidazole and 2-amino-benzimidazole. Additionally, the results indicate that
10 the influence is not related to electronic alterations.

N-1 derivatives. Among the imidazole derivatives tested on inhibition of human QC, most compounds that had improved K_i -values compared to imidazole showed alterations at one nitrogen atom. These compounds also contained one of the most
15 effective QC inhibitors, 1-benzylimidazole. Interestingly, only little alterations of this structure led to a loss of inhibitory quality, as can be seen for 1-benzoylimidazole and phenylimidazole, which was inactive under the experimental conditions. Also in this case, the observed changes seemed not to be only caused by a reduced electron density of the imidazole ring due to the negative mesomeric effect of the Phenyl
20 group, because also the bulky trimethyl-silyl group, exhibiting a positive inductive effect showed reduced binding compared to other residues. Interestingly, one of the less effective compounds of this group was 1-aminopropyl-imidazole. The small efficacy of this compound is caused by the basic amino group, since the sterically similar compounds 1-methylimidazole and 1-vinylimidazole showed improved binding
25 to the active site. Thus, the positively charged amino group accounts for the smaller K_i -value, a result that is corroborated by a comparison of the K_i -values of N- ω -acetylated histamine (Table 3) and histamine (Table 4).

Effect of 3,4 and 3,5 derivatization. The imidazole derivatives that contained
30 substituents in positions 4(5) or both were shown to have a restricted efficiency for binding to the enzyme. The effect of the specific substitutions were specified by comparison of the inhibitory constants of L-histamine and the two intermediates in the biological degradation of histamine, 3-methyl-4-histamine and 3-methyl-5-histamine (Table 4). L-Histamine revealed a K_i value that was about one order of magnitude

smaller compared to its acetylated counterpart. Methylation of one nitrogen resulted in a considerable improvement of efficacy in case of 3-methyl-4-histamine. Methylation leading to 3-methyl-5-histamine, however, resulted in a complete loss of inhibitory activity. Thus, the observed effect seems to be mainly caused by a sterical hindrance of binding due to the derivatisation of the carbon adjacent to the basic nitrogen. Presumably, the basic nitrogen plays a key role for binding to the enzyme.

Example 8: Formation of A β (3-40/42) derivatives

The measurements were carried out with two short N-terminal peptide sequences of A β (3-40/42), Gln³-A β (1-11)a (sequence: DAQFRHDSGYE) and Gln³-A β (3-11)a, which contain a glutamine instead of an glutamic acid residue in the third position. Cleavage by DP IV and cyclization of the N-terminal glutamine residue by QC of the two peptides was tested using MALDI-TOF mass spectrometry. Measurements were carried out using purified DP IV (porcine kidney) or crude porcine pituitary homogenate as sources of QC as well as for both enzymes for measurements of consecutive catalysis.

Results

1. *Formation of Gln³-A β (3-11)a from Gln³-A β 1-11a catalysed by DP IV and its prevention by the DP IV-inhibitor Val-Pyrrolidide (Val-Pyrr)*

DP IV or DP IV-like activity is cleaving Gln³-A β (1-11)a under formation of Gln³-A β (3-11)a (Figure 7). The residue in the third position is uncovered by this cleavage and becomes therefore accessible for modification by other enzymes, i.e. QC. As expected, catalysis can be completely prevented by Val-Pyrr (Figure 8).

2. *Formation of pGlu³-A β (3-11)a from Gln³-A β (3-11)a by catalysis of QC in pituitary homogenate and prevention by 1,10-phenanthroline*

Glutamyl cyclase present in the homogenate of porcine pituitary catalyzes conversion of Gln³-A β (3-11)a to [pGlu³]A β (3-11)a (Figure 9). Formation of [pGlu³]A β (3-11)a was inhibited by addition of 1,10-phenanthroline (Figure 10).

3. *Consecutive catalysis of DPIV and QC resulting in formation of [pGlu³]A β (3-11)a and prevention by Val-Pyrr and 1,10-phenanthroline*

Formation of [pGlu³]A β (3-11)a from Gln³-A β (1-11)a takes place after consecutive
5 catalysis by DP IV and QC, measured in crude homogenate of porcine pituitary with
added DPIV from porcine kidney (Figure 11). [pGlu³]A β (3-11)a was not formed when
the QC-inhibitor 1,10-phenanthroline (Figure 12) or the DP IV-inhibitor Val-Pyrr was
added (Figure 13). The slight appearance of [pGlu³]A β (3-11)a is due to
10 aminopeptidase cleavage and following cyclization of the glutamine residue, also
indicated by formation of Gln³-A β (2-11)a.

4. *Formation of [pGlu³]A β (3-11)a in crude pituitary homogenate by catalysis of aminopeptidase(s)*

15 Due to the formation of [pGlu³]A β (3-11)a that was not dependent on DPIV catalysis,
degradation of Gln³-A β (1-11)a was investigated in crude pituitary homogenate
without added DPIV (Figure 14). As expected from the data in section 4, formation of
[pGlu³]A β (3-11)a was observed. The data show that the degradation of Gln³-A β (1-
11)a may also be catalyzed by aminopeptidase(s), resulting in [pGlu³]A β (3-11)a.
20 Hence, the results show that pyroglutamyl formation is an endpoint of N-terminal
peptide degradation in this tissue, further supporting the role of QC in plaque
formation.

25 **Example 9: Turnover of Gln³-A β (3-11)a; (3-21)a and (3-40) by recombinant human QC**

All Gln³-A β derived peptides tested were efficiently converted by human QC into the
corresponding pyroglutamyl forms (Table 9). Due to the poor solubility of Gln³-A β (3-
21)a and Gln³-A β (3-40) in aqueous solution, the determinations were carried out in
30 presence of 1% DMSO. The better solubility of Gln³-A β (3-11)a, however, allowed the
kinetic analysis of the QC-catalyzed turnover in presence and absence of DMSO
(Table 8). Taken together, the investigation of the A β peptides as QC-substrates with
chain-length of 8, 18 and 37 amino acids (see Table 9) confirmed the observation

that human QC-activity increases with the length of its substrates. Accordingly, Gln¹-gastrin, Gln¹-neurotensin, Gln¹-GnRH are among the best QC-substrates taking the specificity constants into account. Similarly, Gln³-A β (3-40) and glucagon, the largest QC-substrates investigated thus far, exhibited high second order rate constants (449 mM⁻¹s⁻¹ and 526 mM⁻¹s⁻¹ respectively) even in presence of 1% DMSO (Table 9).

Interestingly, the kinetic parameters for the conversion of the investigated amyloid peptides did not change dramatically with increasing size, suggesting only moderate effects of the C-terminal part of A β on QC catalysis. Therefore, due to better solubility and experimental handling, the further investigations concerning N-terminal aminopeptidase processing of these peptides were performed using the smaller fragments of A β , Gln³-A β (1-11)a, Gln³-A β (3-11)a and A β (3-11)a.

Table 9: Kinetic parameters for conversion of N-terminally Gln-containing peptides by recombinant human QC in buffer solution containing 1% DMSO

Peptide	K _M (μ M)	k _{cat} (s ⁻¹)	k _{cat} /K _M (mM ⁻¹ s ⁻¹)
[Gln ³]A β 3-11a	87 $\pm$ 3 [#]	55 $\pm$ 1 [#]	632 $\pm$ 10 [#]
[Gln ³]A β 3-11a	155 $\pm$ 4	41.4 $\pm$ 0.4	267 $\pm$ 4
[Gln ³]A β 3-21a	162 $\pm$ 12	62 $\pm$ 3	383 $\pm$ 10
[Gln ³]A β 3-40	89 $\pm$ 10	40 $\pm$ 2	449 $\pm$ 28
Glucagon(3-29)	19 $\pm$ 1	10.0 $\pm$ 0.2	526 $\pm$ 17

[#] Determined in absence of DMSO

Example 10: Turnover of A β (3-11)a and A β (3-21)a by recombinant human QC

The incubation of A β (3-11)a and A β (3-21)a in presence of QC revealed that in contrast to previous work, glutamate-containing peptides can also serve as QC-substrates (Figures 15 C and D). The QC-catalyzed formation of [pGlu³]A β (3-11)a and [pGlu³]A β (3-21)a was investigated at pH 5.2 and 6.5, respectively. If the QC-inhibitor benzimidazole was added to the solution before starting the assay by addition of QC, substrate conversion resulting in [pGlu³]A β (3-11)a or [pGlu³]A β (3-21)a was suppressed (Figures 15 E and F). If QC was boiled before addition, formation of the pGlu-peptides was negligible (Figures 15 A and B).

Example 11: pH-dependency of the papaya QC-catalyzed cyclization of Gln- β NA and Glu- β NA

5 Papaya QC converted Glu- β NA in a concentration range up to 2 mM (which was limited by substrate solubility) in accordance with Michaelis-Menten kinetics (Figure 16). Inspection of turnover versus substrate concentration diagrams for the QC-catalyzed conversion of Glu- β NA, studied between pH 6.1 and 8.5, revealed that for
10 this Glu-substrate both parameters, K_M and k_{cat} , changed in a pH-dependent manner (Figure 16). This is in contrast to the previously described QC-catalyzed glutamine cyclization, for which only changes in K_M were observed over the given pH range (Gololobov, M. Y. et al. 1994 *Arch Biochem Biophys* 309, 300-307).

Subsequently, to study the impact of the proton concentration during Glu- and Gln-cyclization, the pH-dependence of cyclization of Glu- β NA and Gln- β NA under first-
15 order rate-law conditions (i.e. substrate concentrations far below K_M -values) was investigated (Figure 17). The cyclization of glutamine has a pH-optimum at pH 8.0, in contrast to the cyclization of glutamic acid which showed a pH-optimum of pH 6.0. While the specificity constants at the respective pH-optima differ approximately
20 80,000-fold, the ratio of QC versus EC activity around pH 6.0, is only about 8,000.

The nonenzymatic pGlu-formation from Gln- β NA investigated at pH 6.0, was followed for 4 weeks and revealed a first-order rate constant of $1.2 \cdot 10^{-7} \text{ s}^{-1}$. However, during the same time period, no pGlu- β NA was formed from Glu- β NA, allowing to estimate a limiting rate constant for turnover of $1.0 \cdot 10^{-9} \text{ s}^{-1}$.

Example 12: Enzyme Inactivation/Reactivation Procedures

25 An aliquot of human QC (0.1-0.5 mg, 1 mg/ml) was inactivated overnight by dialysis against a 3000-fold excess of 5 mM 1,10-phenanthroline or 5 mM dipicolinic acid in 0.05 M Bis-Tris/HCl, pH 6.8. Subsequently, the inactivating agent was carefully
30 removed by dialysis (3 cycles, 2000-fold excess) of the samples against 0.05 M Bis-Tris/HCl, pH 6.8, containing 1 mM EDTA. Reactivation experiments were performed at room temperature for 15 minutes using Zn^{++} , Mn^{++} , Ni^{++} , Ca^{++} , K^+ and Co^{++} ions at concentrations of 1.0, 0.5, 0.25 mM in 0.025 M Bis-Tris, pH 6.8 containing 0.5 mM

EDTA. QC activity assays were performed in 0.05 M Tris/HCl, pH 8.0, containing 2 mM EDTA, in order to avoid a rapid reactivation by traces of metal ions present in buffer solutions.

5 The inhibition of porcine QC by 1,10-phenanthroline has already been described (Busby, W. H. J. et al. 1987 *J Biol Chem* 262, 8532-8536, Bateman, R. C. J. et al. 2001 *Biochemistry* 40, 11246-11250). However, the fact that EDTA has been shown to have an activating effect on QC catalysis suggested that inhibition by phenanthroline is not due to metal chelation (Busby, W. H. J. et al. 1987 *J Biol Chem* 262, 8532-8536, Bateman, R. C. J. et al. 2001 *Biochemistry* 40, 11246-11250). Also, 10 in addition to being inhibited by 1,10-phenanthroline, human QC catalyzed substrate cyclization was abolished in presence of dipicolinic acid and 8-hydroxyquinoline, other inhibitors of metalloenzymes. These chelators inhibited QC in a competitive and time-dependent manner, i.e., already competitively inhibited initial activity was found to be further reduced after prolonged incubation with the compounds (Figure 15 18, 19). Interestingly, EDTA did not show remarkable inhibition regardless of incubation time or under any conditions.

Human QC was almost completely inactivated after extensive dialysis against 5 mM 1,10-phenanthroline or 5 mM dipicolinic acid. After repeated dialysis overnight against chelator-free buffer solutions, QC activity was partially reactivated up to 50- 20 60%. However, when dialyzed against buffers containing 1 mM EDTA, no reactivation was observed.

Near-total restoration of QC activity after inactivation by either dipicolinic acid or 1,10-phenanthroline was achieved by incubating the protein for 10 minutes with 0.5 mM ZnSO₄ in presence of 0.5 mM EDTA (Figure 20). Partial restoration of QC activity 25 was similarly obtained using Co⁺⁺ and Mn⁺⁺ ions for reactivation. Even in the presence of 0.25 mM Zn⁺⁺ a reactivation up to 25% of the original activity was possible. No reactivation was observed applying Ni⁺⁺, Ca⁺⁺ or K⁺ ions. Similarly, incubation of fully active QC with these ions had no effect on the enzyme activity.

78a

SEQUENCE LISTING

<110> Probiodrug AG

<120> Use of effectors of Glutaminy1 and Glutamate Cyclases

<130> 17771-2CA

<140> Corresponding to PCT/EP2004/011630

<141> 2004-10-15

<150> US 60/512,038

<151> 2003-10-15

<160> 27

<170> PatentIn version 3.1

<210> 1

<211> 42

<212> PRT

<213> Homo sapiens

<400> 1

Asp	Ala	Glu	Phe	Arg	His	Asp	Ser	Gly	Tyr	Glu	Val	His	His	Gln	Lys
1				5				10						15	

Leu	Val	Phe	Phe	Ala	Glu	Asp	Val	Gly	Ser	Asn	Lys	Gly	Ala	Ile	Ile
			20					25					30		

Gly	Leu	Met	Val	Gly	Gly	Val	Val	Ile	Ala
		35				40			

<210> 2

<211> 40

<212> PRT

<213> Homo sapiens

<400> 2

Asp	Ala	Glu	Phe	Arg	His	Asp	Ser	Gly	Tyr	Glu	Val	His	His	Gln	Lys
1				5				10						15	

Leu	Val	Phe	Phe	Ala	Glu	Asp	Val	Gly	Ser	Asn	Lys	Gly	Ala	Ile	Ile
			20					25					30		

Gly	Leu	Met	Val	Gly	Gly	Val	Val
		35				40	

78b

<210> 3
 <211> 40
 <212> PRT
 <213> Homo sapiens

<400> 3
 Glu Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys Leu Val
 1 5 10 15
 Phe Phe Ala Glu Asp Val Gly Ser Asn Lys Gly Ala Ile Ile Gly Leu
 20 25 30
 Met Val Gly Gly Val Val Ile Ala
 35 40

<210> 4
 <211> 38
 <212> PRT
 <213> Homo sapiens

<400> 4
 Glu Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys Leu Val
 1 5 10 15
 Phe Phe Ala Glu Asp Val Gly Ser Asn Lys Gly Ala Ile Ile Gly Leu
 20 25 30
 Met Val Gly Gly Val Val
 35

<210> 5
 <211> 11
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic peptide

<220>
 <221> MOD_RES
 <222> (11)..(11)
 <223> AMIDATION

<400> 5
 Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu
 1 5 10

78c

<210> 6
<211> 9
<212> PRT
<213> Artificial sequence

<220>
<223> synthetic peptide

<220>
<221> MOD_RES
<222> (9)..(9)
<223> AMIDATION

<400> 6
Glu Phe Arg His Asp Ser Gly Tyr Glu
1 5

<210> 7
<211> 21
<212> PRT
<213> Artificial sequence

<220>
<223> synthetic peptide

<220>
<221> MOD_RES
<222> (21)..(21)
<223> AMIDATION

<400> 7
Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys
1 5 10 15

Leu Val Phe Phe Ala
20

<210> 8
<211> 19
<212> PRT
<213> Artificial sequence

<220>
<223> synthetic peptide

78d

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 8
Glu Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys Leu Val
1 5 10 15

Phe Phe Ala

<210> 9
<211> 38
<212> PRT
<213> Artificial sequence

<220>
<223> Synthetic peptide

<400> 9
Gln Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys Leu Val
1 5 10 15

Phe Phe Ala Glu Asp Val Gly Ser Asn Lys Gly Ala Ile Ile Gly Leu
20 25 30

Met Val Gly Gly Val Val
35

<210> 10
<211> 19
<212> PRT
<213> Artificial sequence

<220>
<223> Synthetic peptide

<220>
<221> MOD_RES
<222> (19)..(19)
<223> AMIDATION

<400> 10
Gln Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys Leu Val
1 5 10 15

Phe Phe Ala

78e

<210> 11
 <211> 11
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic peptide

<220>
 <221> MOD_RES
 <222> (11)..(11)
 <223> AMIDATION

<400> 11
 Asp Ala Gln Phe Arg His Asp Ser Gly Tyr Glu
 1 5 10

<210> 12
 <211> 9
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic peptide

<220>
 <221> MOD_RES
 <222> (9)..(9)
 <223> AMIDATION

<400> 12
 Gln Phe Arg His Asp Ser Gly Tyr Glu
 1 5

<210> 13
 <211> 17
 <212> PRT
 <213> Homo sapiens

<220>
 <221> MOD_RES
 <222> (17)..(17)
 <223> AMIDATION

<400> 13
 Gln Gly Pro Trp Leu Glu Glu Glu Glu Ala Tyr Gly Trp Met Asp
 1 5 10 15

78f

Phe

<210> 14
 <211> 13
 <212> PRT
 <213> Homo sapiens

<400> 14
 Gln Leu Tyr Glu Asn Lys Pro Arg Arg Pro Tyr Ile Leu
 1 5 10

<210> 15
 <211> 10
 <212> PRT
 <213> Homo sapiens

<220>
 <221> MOD_RES
 <222> (10)..(10)
 <223> AMIDATION

<400> 15
 Gln His Trp Ser Tyr Gly Leu Arg Pro Gly
 1 5 10

<210> 16
 <211> 97
 <212> PRT
 <213> Homo sapiens

<400> 16
 Gln Pro Lys Val Pro Glu Trp Val Asn Thr Pro Ser Thr Cys Cys Leu
 1 5 10 15
 Lys Tyr Tyr Glu Lys Val Leu Pro Arg Arg Leu Val Val Gly Tyr Arg
 20 25 30
 Lys Ala Leu Asn Cys His Leu Pro Ala Ile Ile Phe Val Thr Lys Arg
 35 40 45
 Asn Arg Glu Val Cys Thr Asn Pro Asn Asp Asp Trp Val Gln Glu Tyr
 50 55 60
 Ile Lys Asp Pro Asn Leu Pro Leu Leu Pro Thr Arg Asn Leu Ser Thr
 65 70 75 80
 Val Lys Ile Ile Thr Ala Lys Asn Gly Gln Pro Gln Leu Leu Asn Ser
 85 90 95

78g

Gln

<210> 17
 <211> 76
 <212> PRT
 <213> Homo sapiens

<400> 17
 Gln Pro Asp Ser Val Ser Ile Pro Ile Thr Cys Cys Phe Asn Val Ile
 1 5 10 15
 Asn Arg Lys Ile Pro Ile Gln Arg Leu Glu Ser Tyr Thr Arg Ile Thr
 20 25 30
 Asn Ile Gln Cys Pro Lys Glu Ala Val Ile Phe Lys Thr Lys Arg Gly
 35 40 45
 Lys Glu Val Cys Ala Asp Pro Lys Glu Arg Trp Val Arg Asp Ser Met
 50 55 60
 Lys His Leu Asp Gln Ile Phe Gln Asn Leu Lys Pro
 65 70 75

<210> 18
 <211> 76
 <212> PRT
 <213> Homo sapiens

<400> 18
 Gln Pro Asp Ala Ile Asn Ala Pro Val Thr Cys Cys Tyr Asn Phe Thr
 1 5 10 15
 Asn Arg Lys Ile Ser Val Gln Arg Leu Ala Ser Tyr Arg Arg Ile Thr
 20 25 30
 Ser Ser Lys Cys Pro Lys Glu Ala Val Ile Phe Lys Thr Ile Val Ala
 35 40 45
 Lys Glu Ile Cys Ala Asp Pro Lys Gln Lys Trp Val Gln Asp Ser Met
 50 55 60
 Asp His Leu Asp Lys Gln Thr Gln Thr Pro Lys Thr
 65 70 75

<210> 19
 <211> 68
 <212> PRT
 <213> Homo sapiens

78h

<400> 19

Gln Val Gly Thr Asn Lys Glu Leu Cys Cys Leu Val Tyr Thr Ser Trp
 1 5 10 15

Gln Ile Pro Gln Lys Phe Ile Val Asp Tyr Ser Glu Thr Ser Pro Gln
 20 25 30

Cys Pro Lys Pro Gly Val Ile Leu Leu Thr Lys Arg Gly Arg Gln Ile
 35 40 45

Cys Ala Asp Pro Asn Lys Lys Trp Val Gln Lys Tyr Ile Ser Asp Leu
 50 55 60

Lys Leu Asn Ala
 65

<210> 20

<211> 373

<212> PRT

<213> Homo sapiens

<400> 20

Gln His His Gly Val Thr Lys Cys Asn Ile Thr Cys Ser Lys Met Thr
 1 5 10 15

Ser Lys Ile Pro Val Ala Leu Leu Ile His Tyr Gln Gln Asn Gln Ala
 20 25 30

Ser Cys Gly Lys Arg Ala Ile Ile Leu Glu Thr Arg Gln His Arg Leu
 35 40 45

Phe Cys Ala Asp Pro Lys Glu Gln Trp Val Lys Asp Ala Met Gln His
 50 55 60

Leu Asp Arg Gln Ala Ala Ala Leu Thr Arg Asn Gly Gly Thr Phe Glu
 65 70 75 80

Lys Gln Ile Gly Glu Val Lys Pro Arg Thr Thr Pro Ala Ala Gly Gly
 85 90 95

Met Asp Glu Ser Val Val Leu Glu Pro Glu Ala Thr Gly Glu Ser Ser
 100 105 110

Ser Leu Glu Pro Thr Pro Ser Ser Gln Glu Ala Gln Arg Ala Leu Gly
 115 120 125

Thr Ser Pro Glu Leu Pro Thr Gly Val Thr Gly Ser Ser Gly Thr Arg
 130 135 140

Leu Pro Pro Thr Pro Lys Ala Gln Asp Gly Gly Pro Val Gly Thr Glu
 145 150 155 160

Leu Phe Arg Val Pro Pro Val Ser Thr Ala Ala Thr Trp Gln Ser Ser
 165 170 175

78i

Ala Pro His Gln Pro Gly Pro Ser Leu Trp Ala Glu Ala Lys Thr Ser
 180 185 190
 Glu Ala Pro Ser Thr Gln Asp Pro Ser Thr Gln Ala Ser Thr Ala Ser
 195 200 205
 Ser Pro Ala Pro Glu Glu Asn Ala Pro Ser Glu Gly Gln Arg Val Trp
 210 215 220
 Gly Gln Gly Gln Ser Pro Arg Pro Glu Asn Ser Leu Glu Arg Glu Glu
 225 230 235 240
 Met Gly Pro Val Pro Ala His Thr Asp Ala Phe Gln Asp Trp Gly Pro
 245 250 255
 Gly Ser Met Ala His Val Ser Val Val Pro Val Ser Ser Glu Gly Thr
 260 265 270
 Pro Ser Arg Glu Pro Val Ala Ser Gly Ser Trp Thr Pro Lys Ala Glu
 275 280 285
 Glu Pro Ile His Ala Thr Met Asp Pro Gln Arg Leu Gly Val Leu Ile
 290 295 300
 Thr Pro Val Pro Asp Ala Gln Ala Ala Thr Arg Arg Gln Ala Val Gly
 305 310 315 320
 Leu Leu Ala Phe Leu Gly Leu Leu Phe Cys Leu Gly Val Ala Met Phe
 325 330 335
 Thr Tyr Gln Ser Leu Gln Gly Cys Pro Arg Lys Met Ala Gly Glu Met
 340 345 350
 Ala Glu Gly Leu Arg Tyr Ile Pro Arg Ser Cys Gly Ser Asn Ser Tyr
 355 360 365
 Val Leu Val Pro Val
 370

<210> 21
 <211> 76
 <212> PRT
 <213> Homo sapiens

<400> 21
 Gln Pro Val Gly Ile Asn Thr Ser Thr Thr Cys Cys Tyr Arg Phe Ile
 1 5 10 15
 Asn Lys Lys Ile Pro Lys Gln Arg Leu Glu Ser Tyr Arg Arg Thr Thr
 20 25 30
 Ser Ser His Cys Pro Arg Glu Ala Val Ile Phe Lys Thr Lys Leu Asp
 35 40 45

78j

Lys Glu Ile Cys Ala Asp Pro Thr Gln Lys Trp Val Gln Asp Phe Met
 50 . 55 60

Lys His Leu Asp Lys Lys Thr Gln Thr Pro Lys Leu
 65 70 75

<210> 22
 <211> 33
 <212> PRT
 <213> Homo sapiens

<400> 22
 Gln Pro Leu Pro Asp Cys Cys Arg Gln Lys Thr Cys Ser Cys Arg Leu
 1 5 10 15

Tyr Glu Leu Leu His Gly Ala Gly Asn His Ala Ala Gly Ile Leu Thr
 20 25 30

Leu

<210> 23
 <211> 11
 <212> PRT
 <213> Homo sapiens

<400> 23
 Arg Pro Lys Pro Gln Gln Phe Phe Gly Leu Met
 1 5 10

<210> 24
 <211> 34
 <212> PRT
 <213> Homo sapiens

<220>
 <221> MOD_RES
 <222> (1)..(1)
 <223> PYRROLIDONE CARBOXYLIC ACID

<400> 24
 Glu Ala Ser Asn Cys Phe Ala Ile Arg His Phe Glu Asn Lys Phe Ala
 1 5 10 15

Val Glu Thr Leu Ile Cys Ser Arg Thr Val Lys Lys Asn Ile Ile Glu
 20 25 30

Glu Arg

78k

<210> 25
<211> 34
<212> PRT
<213> Homo sapiens

<400> 25
Glu Ala Ser Asn Cys Phe Ala Ile Arg His Phe Glu Asn Lys Phe Ala
1 5 10 15
Val Glu Thr Leu Ile Cys Ser Arg Thr Val Lys Lys Asn Ile Ile Glu
20 25 30
Glu Arg

<210> 26
<211> 34
<212> PRT
<213> Homo sapiens

<220>
<221> MOD_RES
<222> (1)..(1)
<223> PYRROLIDONE CARBOXYLIC ACID


<400> 26
Glu Ala Ser Asn Cys Phe Ala Ile Arg His Phe Glu Asn Lys Phe Ala
1 5 10 15
Val Glu Thr Leu Ile Cys Ser Arg Thr Val Lys Lys Asn Ile Ile Glu
20 25 30
Glu Arg

<210> 27
<211> 34
<212> PRT
<213> Homo sapiens

<400> 27
Glu Ala Ser Asn Cys Phe Ala Ile Arg His Phe Glu Asn Lys Phe Ala
1 5 10 15
Val Glu Thr Leu Ile Cys Ser Arg Thr Val Lys Lys Asn Ile Ile Glu
20 25 30
Glu Arg

CLAIMS

1. Use of an inhibitor of glutamyl cyclase (QC) for the preparation of a medicament for the treatment of a disease selected from the group consisting of Familial British Dementia and Familial Danish Dementia, characterized in that the conversion of the glutamic acid to a pyroglutamic acid residue at the N-terminus of at least one QC-substrate selected from Glu¹-ADan and Glu¹-ABri is inhibited.
2. The use according to claim 1, wherein the inhibitor of QC has the general formula 1:




formula 1


including all pharmaceutically acceptable salts or stereoisomers thereof,

wherein

$R^1 - R^6$ are independently H or a branched or unbranched alkyl chain, a branched or unbranched alkenyl chain, a branched or unbranched alkynyl chain, carbocyclic, aryl, heteroaryl, heterocyclic, aza-amino acid, amino acid or a mimetic thereof, peptide or a mimetic thereof; all of the above residues optionally being substituted, and n is 0, or 2.

3. The use according to any one of claims 1 and 2, wherein the inhibitor of QC is a compound selected from





4. The use according to any one of claims 1 to 3, wherein the inhibitor of QC is used in combination with at least one customary carrier and/or excipient.
5. The use according to any one of claims 1 to 4, wherein the inhibitor of QC is used in combination with a DP IV inhibitor.
6. The use according to claim 5, wherein the DP IV-inhibitor is selected from the group consisting of L-threo-isoleucyl thiazolidine, L-allo-isoleucyl thiazolidine, L-threo-isoleucyl pyrrolidine, L-allo-isoleucyl pyrrolidine, NVP-DPP728A (1-[[[2-[{5-cyanopyridin-2-yl}amino]ethyl]amino]acetyl]-2-cyano-(S)-pyrrolidine), LAF-237 (1-[(3-hydroxy-adamant-1-ylamino)-acetyl]-pyrrolidine-2(S)-carbonitrile); TSL-225 (tryptophyl-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid), FE-999011, N-valyl prolyl, O-benzoyl hydroxylamine, alanyl pyrrolidine, H-Asn-pyrrolidine, H-Asn-thiazolidine, H-Asp-pyrrolidine, H-Asp-thiazolidine, H-Asp(NHOH)-pyrrolidine, H-Asp(NHOH)-thiazolidine, H-Glu-pyrrolidine, H-Glu-thiazolidine, H-Glu(NHOH)-pyrrolidine, H-Glu(NHOH)-thiazolidine, H-His-pyrrolidine, H-His-thiazolidine, H-Pro-pyrrolidine, H-Pro-thiazolidine, H-Ile-azididine, H-Ile-pyrrolidine, H-L-allo-Ile-thiazolidine, H-Val-pyrrolidine and H-Val-thiazolidine, 2-Amino octanoic acid-Pro-Ile, Abu-Pro-Ile, Aib-Pro-Ile, Aze-Pro-Ile, Cha-Pro-Ile, Ile-Hyp-Ile, Ile-Pro-allo-Ile, Ile-Pro-t-butyl-Gly, Ile-Pro-Val, Nle-Pro-Ile, Nva-Pro-Ile, Orn-Pro-Ile, Phe-Pro-Ile, Phg-Pro-Ile, Pip-Pro-Ile, Ser(Bzl)-Pro-Ile, Ser(P)-Pro-Ile, Ser-Pro-Ile, t-butyl-Gly-Pro-D-Val, t-butyl-Gly-Pro-Gly, t-butyl-Gly-Pro-Ile, t-butyl-Gly-Pro-Ile-amide, t-butyl-Gly-Pro-t-butyl-Gly, t-butyl-Gly-Pro-Val, Thr-Pro-Ile, Tic-Pro-Ile, Trp-Pro-Ile, Tyr(P)-Pro-Ile, Tyr-Pro-allo-Ile, Val-Pro-allo-Ile, Val-Pro-t-butyl-Gly, Val-Pro-Val, and a pharmaceutical acceptable salt thereof.
7. The use according to claim 5 , wherein the DP IV-inhibitor is selected from the group consisting of

2-Methylcarbonyl-1-N-[(L)-Alanyl-(L)-Valinyl]-(2S)-pyrrolidine hydrobromide,
 2-Methylcarbonyl-1-N-[(L)-Valinyl-(L)-Prolyl-(L)-Valinyl]-(2S)-pyrrolidine
 hydrobromide,
 2-[(Acetyl-oxy-methyl)carbonyl]-1-N-[(L)-Alanyl-(L)-Valinyl]-(2S)-pyrrolidine
 hydrobromide,
 2-[Benzoyl-oxy-methyl]carbonyl]-1-N-[(L)-Alanyl]-(L)-Valinyl]-(2S)-pyrrolidine
 hydrobromide,
 2-[(2,6-Dichlorobenzyl)thiomethyl]carbonyl]-1-N-[(L)-Alanyl]-(L)-Valinyl]-(2S)-
 pyrrolidine,
 2-[Benzoyl-oxy-methyl]carbonyl]-1-N-[Glycyl-(L)-Valinyl]-(2S)-pyrrolidine
 hydrobromide,
 2-[(1,3-Thiazolethiazol-2-yl)carbonyl]-1-N-[(L)-Alanyl]-(L)-Valinyl]-(2S)-pyrrolidine
 trifluoroacetate,
 2-[(Benzothiazolethiazol-2-yl)carbonyl]-1-N-[N-[(L)-Alanyl]-(L)-Valinyl]-(2S)-
 pyrrolidine trifluoroacetate,
 2-[(Benzothiazolethiazol-2-yl)carbonyl]-1-N-[(L)-Alanyl]-Glycyl]-(2S)-pyrrolidine
 trifluoroacetate,
 2-[(Pyridin-2-yl)carbonyl]-1-N-[N-[(L)-Alanyl]-(L)-Valinyl]-(2S)-pyrrolidine
 trifluoroacetate,
 1-cyclopentyl-3-methyl-1-oxo-2-pentanaminium chloride,
 1-cyclopentyl-3-methyl-1-oxo-2-butanaminium chloride,
 1-cyclopentyl-3,3-dimethyl-1-oxo-2-butanaminium chloride,
 1-cyclohexyl-3,3-dimethyl-1-oxo-2-butanaminium chloride,
 3-(cyclopentylcarbonyl)-1,2,3,4-tetrahydroisoquinolinium chloride, and
 N-(2-cyclopentyl-2-oxoethyl)cyclohexanaminium chloride,
 and other pharmaceutical acceptable salts thereof.

8. The use according to any one of claims 1 to 4, wherein the inhibitor of QC is
 used in combination with an inhibitor of a DP IV-like enzyme selected from the group
 consisting of fibroblast activation protein α , dipeptidyl peptidase IV β , dipeptidyl
 aminopeptidase-like protein, N-acetylated α -linked acidic dipeptidase, quiescent cell
 proline dipeptidase, dipeptidyl peptidase II, attractin, dipeptidyl peptidase IV related
 protein (DPP 8), DPL1 (DPX, DP6), DPL2, DPP 9 and dipeptidyl peptidase 10.

9. The use according to any one of claims 1 to 8, wherein the inhibitor of QC is a competitive inhibitor.
10. The use according to any one of claims 1 to 9, wherein the glutaminyI cyclase inhibitor is adapted for parenteral, enteral or oral administration.
11. The use according to any one of claims 1 to 9, wherein the glutaminyI cyclase inhibitor is adapted for oral administration.

Figure 1




Figure 2




Figure 3






Figure 4




5/23

Figure 5




6/23

Figure 6




7/23

Figure 7




8/23

Figure 8




9/23

Figure 9




10/23

Figure 10



11/23

Figure 11



12/23

Figure 12





Figure 13



14/23

Figure 14




Figure 15




Figure 16




Figure 17




Figure 18





Figure 19



20/23

Figure 20




Figure 21

hQC	MAGGRHRRRVGTLHLLLLVAALPWASRGVSPSASAWPEEKNYHQPAILNSSALRQIAEGT
SGAP	-----APDIPLANVKAHLTQLS
hGCP II	-----ANEYAYRRGIAEAVGLPSIPVHPIGYYDAQ-K
	* * : : .
hQC	SISEMWQNDLQPLLIERYPGSPGSYAARQHIMQRIQRLQADWVLEIDTFLSQTTPYGYRSF
SGAP	TIAAN--NGGN-----RAHGRPGYKASVDYVKAKLD--AAGYTTTLQQFTSGGATGYNLI
hGCP II	LLEKM--GGSAP---PDSSWRGSLKVPYNVGPFTGNFSTQKVKMHIHSTNEVTRIYNVI
	: . . . : : . . . * . :
hQC	SNIISTLNPTAKRHLVLACHYDSKYFSHWNNRVFVGATDSAVPCAMMLELAR--ALDKK
SGAP	ANWPG-GDP--NKVLMAGAHLD--VSSG-----AGINDNGSGSAAVLETAL--AVSRA
hGCP II	GTLRGAVEP--DRYVILGGHRDS-----W----VFGGIDPQSGAAVVHEIVRSFGTLKKE
	. . . : * : : . * * * * . * : * . : : :
hQC	LLSLKTVSDSKPDLQLIFFDGEAFLHWSPQDSLYGSRHLAAKMASTPHPPGARGTSQ
SGAP	GY--Q-----PDKHLRFAWWGAEEELGLIGS---KFY-----VNNLPSADR--SK
hGCP II	GW--R-----PRRTILFASWDAEEFGLLGS---TEW-----AEENSR-LLQ
	: * : : : . * * * * . : * :
hQC	LHGMDLLVLLDLIGAPNPTFPNFF--PNSARWFERLQAIEHELH--ELGLLKDHSLEGR
SGAP	LAG--YLNFDMIGSPNPGYFVYDDDPVIEKTFKNYFAGLNVPT--EIETEGDGRSDHA
hGCP II	ERG-VAYINADSSIEGNYTLRVDCT-PLMYSLVHNLTKELKSPDEGFEGKSLYESWTKKS
	* : * * * . . . : * : .
hQC	Y---FQNYSY---G-G---VIQDD-HIPFLRRGVP-VLHLIPSPFPEVWHTMDDNEE
SGAP	P---FKNVGVP--VG-G---LFTGAGYTKSAAQAQK-WGGTAGQAFDRCYHSSCDSLS
hGCP II	PSPEFSGMPRISKLGSGNDFEVFFQRLGIASGRARYTKNWETNKFSGYP-LYHSVYETYE
	* . . * * . : : : * : : .
hQC	NLDESTID-N-LNKILQVFVLEYLHL----
SGAP	NINDTALDRNSDAAAHAIWTLSSGTGEPPT
hGCP II	-LVEKFYD--PMFKYHLTVAQVRGGMVFEL
	: : . *

Figure 22






Figure 23





formula 1