



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07K 14/805, A61K 38/42, C07K 14/76, 16/00, A61K 9/127, A61J 1/05</p>	A1	<p>(11) International Publication Number: WO 95/05397</p> <p>(43) International Publication Date: 23 February 1995 (23.02.95)</p>					
<p>(21) International Application Number: PCT/US94/09246</p> <p>(22) International Filing Date: 16 August 1994 (16.08.94)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>08/107,543</td> <td>16 August 1993 (16.08.93)</td> <td>US</td> </tr> <tr> <td>08/291,590</td> <td>15 August 1994 (15.08.94)</td> <td>US</td> </tr> </table> <p>(71)(72) Applicant and Inventor: HSIA, Jen-Chang [CA/US]; 135 Starcrest, Irvine, CA 92715 (US).</p> <p>(74) Agents: MULVILLE, Kurt, T. et al.; Lyon & Lyon, First Interstate World Center, Suite 4700, 633 West Fifth Street, Los Angeles, CA 90071-2066 (US).</p>	08/107,543	16 August 1993 (16.08.93)	US	08/291,590	15 August 1994 (15.08.94)	US	<p>(81) Designated States: CA, CN, DE, GB, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
08/107,543	16 August 1993 (16.08.93)	US					
08/291,590	15 August 1994 (15.08.94)	US					
<p>(54) Title: COMPOSITION AND METHODS USING NITROXIDES TO AVOID OXYGEN TOXICITY</p> <p>(57) Abstract</p> <p>This invention relates to the technical field of compositions and processes to alleviate oxygen toxicity based on the addition of nitroxides to physiologically compatible macromolecules. In particular, hemoglobin-based red cell substitutes are described featuring stable nitroxide free radicals for use in cell-free hemoglobin solutions, encapsulated hemoglobin solutions, stabilized hemoglobin solutions, polymerized hemoglobin solutions, conjugated hemoglobin solutions, nitroxide-labelled albumin, and nitroxide-labelled immunoglobulin. The formulations described herein interact with free radicals, act as antioxidant enzyme-mimics, and alleviate oxidative stress and oxygen-related toxicity.</p>							

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

COMPOSITION AND METHODS USING NITROXIDES TO AVOID OXYGEN TOXICITY.

TECHNICAL FIELD

This invention relates to the use of nitroxide compounds that are used with macromolecules, including hemoglobin, albumin, immunoglobulins and liposomes to alleviate the toxic effects of oxygen-related species in a living organism. In particular, this invention discloses compounds and methods featuring nitroxides associated with physiologically compatible cell-free and encapsulated hemoglobin solutions for use as a red cell substitute and nitroxides associated with other physiologically compatible macromolecules for alleviation and prevention of damage and oxidative stress caused by free radicals.

BACKGROUND ART

Although the physiological mechanisms of oxygen metabolism have been known for many years, an understanding of the role played by oxidative stress in physiology and medicine is not well understood. The mechanism by which free radicals contribute to a variety of types of physiological damage has also been studied in connection with oxidative stress and its toxic effects. However, the development of methods and compounds to combat oxidative stress or toxicity associated with oxygen-related species has enjoyed limited success. The difficulties encountered in creating a blood substitute and are an acute example of the difficulty in preventing or alleviating oxygen toxicity.

Scientists and physicians have struggled for decades to produce a blood substitute that could be safely transfused into humans. Persistent blood shortages and the problems of

incompatible blood types, cross-matching, and the communication of disease have led to a broad-based effort by private industry, universities, and governments to discover a formulation that would allow a large volume of a blood substitute to be safely transfused without significant physiological side effects. At present, several companies are conducting clinical trials on experimental blood substitutes. However, unexpected adverse physiological reactions and the inherent complexity of the research and development process have impeded progress through the regulatory approval stage and have prevented the introduction of a clinically useful blood substitute.

A Research Advisory Committee of the United States Navy issued a report in August 1992 outlining the efforts by several groups to produce a blood substitute, assessing the status of those efforts, and generally describing the toxicity problems encountered. The Naval Research Advisory Committee Report reflects the current consensus in the scientific community that even though the existing blood substitute products, often termed "hemoglobin-based oxygen carriers" (HBOC), have demonstrated efficacy in oxygen transport, certain toxicity issues are unresolved. The adverse transfusion reactions that have been observed in clinical studies of existing hemoglobin-based oxygen carriers (HBOC) include systemic hypertension and vasoconstriction. These adverse reactions have forced a number of pharmaceutical companies to abandon their clinical trials or to proceed at low dosage levels.

The toxicity problem in the existing hemoglobin-based blood substitutes has been given a high priority by the United States Government. The Naval Research Committee recommendation has been implemented by the National Institute of Health in the form of a Request For Proposal (PA-93-23) on the subject of "Hemoglobin-Based Oxygen Carriers: Mechanism

of Toxicity." Therefore, the medical and scientific community suffers from an acute and pressing need for a blood substitute that may be infused without the side effects observed with the existing hemoglobin-based oxygen carriers.

The red blood cells are the major component of blood and contain the body's oxygen transport system. It has long been recognized that the most important characteristic of a blood substitute is the ability to carry oxygen. The red blood cells are able to carry oxygen because the primary component of the red cells is hemoglobin, which functions as the oxygen carrier. Most of the products undergoing clinical testing as blood substitutes contain hemoglobin that has been separated from the red blood cell membranes and the remaining constituents of the red blood cells and has been purified to remove essentially all contaminants. However, when hemoglobin is removed from the red cells and placed in solution in its native form, it is unstable and rapidly dissociates into its constituent subunits. For this reason, the hemoglobin used in a hemoglobin-based oxygen carrier (HBOC) must be stabilized to prevent dissociation in solution. Substantial expenditures in scientific labor and capital were necessary to develop hemoglobin-based products that are stable in solution, and which are stabilized in such a way that the oxygen transport function is not impaired. The ability of the existing hemoglobin-based oxygen carriers to transport oxygen has been well established (See United States Patent Nos. 3,925,344; 4,001,200; 4,001,401; 4,053,590; 4,061,736; 4,136,093; 4,301,144; 4,336,248; 4,376,095; 4,377,512; 4,401,652; 4,473,494; 4,473,496; 4,600,531; 4,584,130; 4,857,636; 4,826,811; 4,911,929 and 5,061,688).


In the body, hemoglobin in the red cells binds oxygen molecules as the blood passes through the lungs and delivers the oxygen molecules throughout the body to meet the demands of the body's normal metabolic function. However, the

atmospheric oxygen that most living beings must breathe to survive is a scientific and medical paradox. On the one hand, almost all living organisms require oxygen for life. On the other hand, a variety of toxic oxygen-related chemical species are produced during normal oxygen metabolism.

With respect to oxidative stress resulting from the transportation of oxygen by hemoglobin, it is known that in the process of transporting oxygen, the hemoglobin (Hb) molecule can itself be oxidized by the oxygen (O_2) molecule it is carrying. This auto-oxidation reaction produces two undesirable products: met-hemoglobin (met-Hb) and the superoxide anion ($\cdot O_2^-$). The chemical reaction may be written as follows:



The superoxide anion ($\cdot O_2^-$) is an oxygen molecule that carries an additional electron and a negative charge. The superoxide anion is highly reactive and toxic. In the case of oxygen transport by hemoglobin, potentially damaging oxidative stress originates with the superoxide anion being generated by the auto-oxidation of hemoglobin and results from the subsequent conversion of the superoxide anion to toxic hydrogen peroxide in the presence of the enzyme superoxide dismutase (SOD) by the following reaction:




The presence of the superoxide anion and hydrogen peroxide in

the red blood cells is believed to be the major source of oxidative stress to the red cells.

Apart from oxygen transport by the hemoglobin contained therein, a less recognized characteristic of the red cells is that they contain a specific set of enzymes which are capable of detoxifying oxygen-related chemical species produced as by-products of oxygen metabolism. Without the protection of these specific enzyme systems, autoxidation of hemoglobin would lead to deterioration and destruction of the red cells. In the body, however, the reserve capacity of the enzyme systems in the red cells protects the body from oxygen toxicity by converting the superoxide anion generated during normal metabolism to non-toxic species and thereby controls the level of oxidative stress. However, if this enzyme system breaks down, the integrity of the red cells will be damaged. A lesion of the gene that produces one of the enzymes in the protective system in the red blood cells will cause an observable pathological condition. For example, glucose-6-phosphate dehydrogenase deficiency, a genetic disorder of red cells, is responsible for hydrogen peroxide induced hemolytic anemia. This disorder is due to the inability of the affected cells to maintain NAD(P)H levels sufficient for the reduction of oxidized glutathione resulting in inadequate detoxification of hydrogen peroxide through glutathione peroxidase (P. Hochstein, Free Radical Biology & Medicine, 5:387 (1988)).

The protective enzyme system of the red blood cells converts the toxic superoxide anion molecule to a non-toxic form in a two-step chemical pathway. The first step of the pathway is the conversion of the superoxide anion to hydrogen peroxide by the enzyme superoxide dismutase (SOD) (See Equation [2]). Because hydrogen peroxide is also toxic to cells, the red cells contain another enzyme, catalase, which converts hydrogen peroxide to water as the second step of the pathway (See Equation [3]).



Red cells are also capable of detoxifying hydrogen peroxide and other toxic organoperoxides using the enzyme glutathione peroxidase which reacts with glutathione to convert hydrogen peroxide and organoperoxides to water. Red cells also contain an enzyme to prevent the build up of the met-hemoglobin produced by the auto-oxidation of hemoglobin. The enzyme met-hemoglobin reductase converts met-hemoglobin back to the native form of hemoglobin. Therefore, in the body, the toxic effects of the auto-oxidation of hemoglobin are prevented by specific enzyme-based reaction pathways that eliminate the unwanted by-products of oxygen metabolism.

The enzymatic oxygen detoxification functions of superoxide dismutase, catalase, and glutathione peroxidase that protect red blood cells from oxygen toxicity during normal

oxygen transport do not exist in the hemoglobin-based oxygen carriers (HBOC) developed to date. Without the oxygen detoxification function, the safety of the existing HBOC solutions will suffer due to the presence of toxic oxygen-related species.

The principle method by which the existing HBOC solutions are manufactured is through the removal of hemoglobin from the red cells and subsequent purification to remove all non-hemoglobin proteins and other impurities that may cause an adverse reaction during transfusion (See United States Patent Nos. 4,780,210; 4,831,012; and 4,925,574). The substantial destruction or removal of the oxygen detoxification enzyme systems is an unavoidable result of the existing isolation and purification processes that yield the purified hemoglobin used in most HBOCs. Alternatively, instead of isolating and purifying hemoglobin from red cells, pure hemoglobin has been produced using recombinant techniques. However, recombinant human hemoglobin is also highly purified and does not contain the oxygen detoxification systems found in the red cells. Thus, the development of sophisticated techniques to create a highly purified hemoglobin solution is a mixed blessing because the purification processes remove the detrimental impurities and the beneficial oxygen detoxification enzymes normally present in the red cells and ultimately contributes to oxygen-related toxicity.

One of the observed toxic side effects of the existing

HBOCs is vasoconstriction or hypertension. It is well known that the enzyme superoxide dismutase (SOD) in vitro will rapidly scavenge the superoxide anion and prolong the vasorelaxant effect of nitric oxide (NO). Nitric oxide is a molecule that has recently been discovered to be the substance previously known only as the "endothelium-derived relaxing factor" (EDRF). The prolongation of the vasorelaxant effect of nitric oxide by SOD has been ascribed to the ability of SOD to prevent the reaction between the superoxide anion and nitric oxide. (M.E. Murphy et. al., Proc. Natl. Acad. Sci. USA 88:10860 (1991); Ignarro et.al. J. Pharmacol. Exp. Ther. 244: 81 (1988); Rubanyi Am. J. Physiol. 250: H822 (1986); Gryglewski et.al. Nature 320: 454 (1986)).

However, in vivo, the inactivation of EDRF by the superoxide anion has not been observed and is generally not thought to be likely. Nevertheless, certain pathophysiological conditions that impair SOD activity could result in toxic effects caused by the superoxide anion (Ignarro L.J. Annu. Rev. Pharmacol. Toxicol. 30:535 (1990)). The hypertensive effect observed in preclinical animal studies of the existing HBOC solutions suggests that the concentration of superoxide anion in large volume transfusions of the existing HBOCs is the cause for the destruction of EDRF and the observed vasoconstriction and systemic hypertension.

It is, therefore, important to delineate the hypertensive effect resulting from the reaction of the superoxide anion

with nitric oxide (NO) from that resulting from extravasation and the binding of NO by hemoglobin. Upon transfusion of an HBOC, the hemoglobin can also depress the vasorelaxant action of nitric oxide by reacting with nitric oxide to yield the corresponding nitrosyl-heme (NO-heme) adduct. In particular, deoxy-hemoglobin is known to bind nitric oxide with an affinity which is several orders of magnitude higher than that of carbon monoxide. These hemoglobin-NO interactions have been used to assay for nitric oxide and to study the biological activity of nitric oxide. For example, the antagonism of the vasorelaxant effect of nitric oxide by hemoglobin appears to be dependent on the cell membrane permeability of hemoglobin. In intact platelets, hemoglobin did not reverse the effect of L-arginine which is the precursor of nitric oxide. In contrast, in the cytosol of lysed platelets, hemoglobin is the most effective inhibitor of L-arginine induced cyclic-GMP formation mediated by nitric oxide. These experiments demonstrated that the hemoglobin did not penetrate the platelet membrane effectively. (Radomski et.al. Br. J. Pharmacol. 101:325 (1990)). Therefore, one of the desired characteristics of the HBOCs is to eliminate the interaction of nitric oxide with hemoglobin. Hemoglobin is also known to antagonize both endothelium-dependent vascular relaxation (Martin W. et. al. J. Pharmacol. Exp. Ther. 232: 708 (1985)) as well as NO-elicited vascular smooth muscle relaxation (Grueter C.A. et. al. J.

Cyclic. Nucleotide Res. 5:211 (1979)). Attempts have been made to limit the extravasation and hypertensive effect of hemoglobin by chemically stabilizing, polymerizing, encapsulating, or conjugating the hemoglobin in the HBOCs to prolong the circulation time. Therefore, although the current HBOCs are relatively membrane impermeable and able to transport oxygen, the HBOC solutions do not have the capability of preventing the reaction between superoxide anion and nitric oxide when transfused.

An ideal solution to the toxicity problems of the existing blood substitutes would be a hemoglobin-based formulation that combines the oxygen-transport function of the existing HBOCs with the oxygen detoxification function of the red cells. However, a simple addition of the enzyme superoxide dismutase (SOD) into an existing HBOC solution would not be desirable because, by reducing the concentration of superoxide anion, the reaction whereby hemoglobin is oxidized to met-hemoglobin would be encouraged, leading to an undesirable build-up of met-hemoglobin (See Equation [1]). Also, it is not desirable to encourage the conversion of the superoxide anion to hydrogen peroxide in a hemoglobin solution because the hydrogen peroxide is toxic and reactive and will decompose to toxic hydroxyl radicals or form other toxic organoperoxides during storage.

This invention contemplates the use of stable nitroxide free radicals, hereafter referred to as "nitroxide(s)", to

provide the oxygen detoxification function of the red cells to hemoglobin-based blood substitutes and to alleviate oxidative stress to avoid biological damage associated with free radical toxicity, including inflammation, post-ischemic reperfusion injury, ionizing radiation, and the aging process. Nitroxides are stable free radicals that are shown to have antioxidant catalytic activities which mimic those of superoxide dismutase (SOD), and which when existing *in vivo*, can interact with other substances to perform catalase-mimic activity. In the past, nitroxides have been used in electron spin resonance spectroscopy as "spin labels" for studying conformational and motional characteristics of biomacromolecules. Nitroxides have also been used to detect reactive free radical intermediates because their chemical structure provides a stable unpaired electron with well defined hyperfine interactions. In addition, nitroxides have been observed to act as enzyme mimics; certain low molecular weight nitroxides have been identified to mimic the activity of superoxide dismutase (SOD). (A. Samuni et. al. J. Biol. Chem. 263:17921 (1988)) and catalase (R.J. Mehlhorn et. al., Free Rad. Res. Comm., 17:157 (1992)). Numerous studies also show that nitroxides that are permeable to cell membranes are capable of short-term protection of mammalian cells against cytotoxicity from superoxide anion generated by hypoxanthine/xanthine oxidase and from hydrogen peroxide exposure.

With regard to safety *in vivo*, relatively high levels of

nitroxide are expected to be well tolerated as nitroxides are known to be relatively safe: for example, the maximum tolerated intraperitoneal dose of TEMPO in mice is 275 mg/kg and the LD₅₀ is 341 mg/kg. Further, a macromolecule-bound nitroxide will be safer than a free nitroxide. The utility of nitroxide-labelled macromolecule as an antioxidant enzyme mimic, therefore, lies in the possibility of achieving high nitroxide levels (and hence activity) with acceptable safety.

Most of the nitroxides studied to date in living organisms have been relatively low molecular weight compounds which can easily permeate across cell membranes into body tissues. The nitroxides used as enzyme mimics, pursuant to this invention, are associated with biological and synthetic macromolecules which may be infused and may remain confined to the vascular compartment. In the preferred embodiments of this invention, nitroxides are covalently attached to macromolecules to alleviate free radical toxicity while confining the nitroxide to the location, i.e., the vascular compartment, where their utility in reacting with free radicals is optimized.

A variety of techniques have been described to covalently attach a nitroxide to biomacromolecules, including hemoglobin, albumin, immunoglobulins, and liposomes. See e.g., McConnell et. al., Quart. Rev. Biophys. 3:p.91 (1970); Hamilton et. al., "Structural Chemistry and Molecular Biology" A. Rich et. al., eds. W.H. Freeman, San Francisco,

p.115 (1968); Griffith et. al., Acc. Chem. Res. 2:p.17 (1969); Smith I.C.P. "Biological Applications of Electron Spin Resonance Spectroscopy" Swartz, H.M. et. al., eds., Wiley/Interscience, New York p.483 (1972). Although selected nitroxides have been covalently bound to hemoglobin molecules for the purpose of studying cooperative oxygen binding mechanisms of hemoglobin, nitroxides have not been used in connection with hemoglobin that is specially formulated for use with blood substitutes. Experimental results are presented below to demonstrate that nitroxides may be attached to stabilized, polymerized, conjugated and encapsulated hemoglobin for use as a blood substitute because the nitroxide reacts with free radicals. The interaction of nitroxide-labelled hemoglobin with free radicals also suggests that other biologically compatible macromolecules with a substantial plasma half-life may be labelled with nitroxides to advantageously provide resistance to or protection from oxidative stress or toxicity caused by free radical chemical species.

As noted above, it is known that nitroxides can be chemically bound to biological macromolecules, including hemoglobin, serum albumin, immunoglobulins, and liposomes. However, this work has generally used nitroxides simply as molecular probes for biophysical research; nitroxide-labeled macromolecules have not been specially formulated for use as therapeutic substances.

With respect to the macromolecules described here, two techniques for binding the nitroxides to a macromolecule, often known as "labelling strategies" are possible. The significance of specific labelling lies in the micro-environment in which the nitroxide is bound to the macromolecule and the nitroxide's resulting catalytic activity. Specific labelling at a particular ligand binding site or sites will yield a homogeneous product with a more consistent binding site micro-environment and thus a more reliable compound in terms of the catalytic specificity and activity of the nitroxide.

Based on the experimental results presented here involving the infusion of nitroxide-labelled HBOC, the reaction of small and large molecular weight nitroxides with free radicals has been observed *in vitro* and *in vivo* in the vascular compartment. Based on these studies, the reaction mechanism whereby nitroxide-labelled HBOC participates in the oxidation/reduction reaction of free radicals demonstrates the capability to formulate novel HBOC compounds and other nitroxide-labelled macromolecules to detoxify free radicals.

DISCLOSURE OF INVENTION

This invention discloses compositions and methods using stable nitroxides in connection with biological macromolecules and, in particular, with hemoglobin-containing solutions. In particular embodiments, stable nitroxides are used

with hemoglobin-containing solutions to create several formulations for a blood substitute that will possess the oxygen detoxification function of the red cells. These formulations may be described herein as hemoglobin-based red cell substitutes (HRCs) because the oxygen transport capability of the hemoglobin-based oxygen carriers (HBOC) is enhanced by providing the oxygen detoxification function of the body's red cells. The nitroxide-labelled albumin and nitroxide-labelled immunoglobulins also provide oxygen detoxification by providing a nitroxide covalently attached to a macromolecule possessing a variety of characterized ligand binding sites that is both stable and non-toxic *in vivo*, and which has a substantial plasma half-life.

In addition, this invention describes nitroxides that are covalently attached to the inner surface of a container or are attached to an insoluble matrix housed in a filter to be used with an existing HBOC to scavenge toxic oxygen-related compounds before infusion into a patient. The HRCs formulations described herein will alleviate the oxidative stress originating from the generation of the superoxide anion in the existing HBOC solutions, and upon transfusion, will diminish the destruction of nitric oxide, the endothelium-derived relaxing factor (EDRF). If the destruction of EDRF is prevented, the problem of vasoconstriction and systemic hypertension that are observed when the existing HBOC solutions are infused into a patient will be substantially

alleviated.

The HRCS formulations and nitroxide-labelled macromolecules described below retard the formation of toxic oxygen-related species by causing a nitroxide to function as a "superoxide oxidase," an enzyme-like reaction not known to occur in the red cells. In these HRCS formulations, the nitroxide prevents the accumulation of the undesirable superoxide anion generated from the auto-oxidation of hemoglobin (See Equation [1]). The nitroxide-labelled albumin and immunoglobulins similarly function as antioxidant enzyme mimics whose function remains localized in the vascular and interstitial compartments.

In the "superoxide oxidase" reaction, the superoxide anion is oxidized back into molecular oxygen without proceeding to the formation of hydrogen peroxide. This is accomplished in part by creating a storage condition wherein the concentration of nitroxide greatly exceeds that of oxygen. Used in the manner disclosed herein, the nitroxide prevents the cascade of undesirable oxidative reactions that begin with the formation of the superoxide anion. Furthermore, the physiologically compatible HRCS solutions described here will offer advantages over the existing HBOC solutions because the nitroxide will mimic the enzymatic activity of superoxide dismutase and catalase after the formulations described herein are infused into a patient.

Preferred compositions using nitroxide in connection with

a physiologically compatible hemoglobin solutions include: 1) nitroxide attached to a storage container or covalently attached to an insoluble matrix used as a filter, 2) nitroxide covalently linked to hemoglobin that is stabilized by chemical or recombinant cross-linking, 3) nitroxide covalently linked to polymerized hemoglobin, in particular, in 2, 4, and 8 molar equivalents of nitroxide, 4) nitroxide co-encapsulated with hemoglobin inside a liposome or intercalated into a liposome membrane, (5) nitroxide covalently bound to conjugated hemoglobin, (6) nitroxide covalently bound to albumin, and (7) nitroxide covalently bound to immunoglobulins.

Also, HRCS based on the co-encapsulation of a membrane-impermeable nitroxide with hemoglobin is disclosed. The methods of preparing the above formulations are described as are methods for producing nitroxide-containing filters and containers for the storage and detoxification of an existing HBOC.

Preferred compositions using nitroxide in connection with human serum albumin or recombinant albumin include:

1) non-specific labelling of albumin with nitroxide (e.g., 4-(2-bromoacetamido)-TEMPO at high nitroxide to albumin ratios; and

2) specific labelling of albumin at specific ligand binding sites.

Preferred compositions using nitroxide in connection with

immunoglobulins include a nitroxide labelled hapten or antigen bound to an immunoglobulin specific for the hapten or antigen.

BRIEF DESCRIPTION OF DRAWINGS

Figures 1A and 1B show the electron spin resonance spectra of 4-amino-TEMPO labelled o-raffinose polymerized hemoglobin recorded on (A) day 1 and (B) day 30 (TEMPO: 2,2,6,6 tetramethylpiperidine-1-oxyl). Figure 1C is the spectra of the sample in Figure 1A diluted with equal volume of unlabelled hemoglobin recorded on day 1. Figure 1D is the sample in Figure 1C recorded on day 30.

Figures 2A and 2B are, respectively, the electron spin resonance spectra demonstrating covalent attachment of 4-(2-bromoacetamido)-TEMPO to ω -aminoethyl-agarose and 4-amino-TEMPO to 1,4-bis(2:3-Epoxypropoxy) butane-activated agarose.

Figures 3A and 3B, respectively, are electron spin resonance spectra demonstrating successful covalent attachment of 4-(2-Bromoacetamido)-TEMPO and 3-maleimido-PROXYL to 3,5-bis-bromosilicyl-bisfumarate (DBBF) cross-linked or diaspirin cross-linked human hemoglobin.

Figure 4A is an ESR spectra of 4-(2-bromoacetamido)-TEMPO. Figure 4B is an ESR spectra of 4-(2-bromoacetamido)-TEMPO-labelled HBOC. Figure 4C is an ESR spectra of $^{15}\text{ND}_{17}$ TEMPOL in Lactated Ringer's solution recorded at room temperature.

Figure 5 is an ESR spectra of 4-(2-bromoacetamido)-TEMPO-labelled HBOC with different molar ratios of nitroxides to Hb; Fig. 5A 2:1, Fig. 5B 4:1 and Fig. 5C 8:1. The instrument sensitivity were decreased proportionately from Fig. 5A to Fig. 5B to Fig. 5C to record the spectra so that the center peak (Mo) would be shown to have similar peak height.

Figure 6 is an ESR spectrum of a mixture of 4-(2-bromoacetamido)-TEMPO labelled HBOC and $^{15}\text{ND}_{17}$ -TEMPOL wherein the center peak (see down arrow) of the former and the high field peak (see up-arrow) of the latter were adjusted to similar intensity. This is a superimposition of ESR spectrum from Fig. 4B and Fig. 4C.

Figure 7 is a plasma half-life of 4-(2-bromoacetamido)-TEMPO-labelled HBOC in mouse. Fig. 7A is the ESR spectrum of the nitroxide signal recorded from the mouse tail approximately 10 minutes after i.v. infusion of 0.5 ml of the sample shown in Fig. 6. Fig. 7B is the time dependent (scan time 30 minutes) decrease in the center peak (Mo) signal intensity of Fig. 7A recorded at 10 times of the instrument sensitivity. Fig. 7C is a continuation of Fig. 7B at the end of its scan.

Figure 8 is a plasma half-life of a mixture of 4-(2-bromoacetamido)-TEMPO-labelled HBOC (8g/dl of Hb and 8:1 TEMPO to Hb) and $^{15}\text{ND}_{17}$ TEMPOL (0.5 ml in a 32 g. mouse) recorded from the mouse tail with a cannula for immediate recording of the infused nitroxides. The ESR spectrum of the sample prior to injection is shown in Fig. 6. Fig. 8A is a

series of 5 ESR spectrum recorded at 0.5 minute intervals, the magnetic field strength was increased by 2 Gauss in between each scan to display the decrease in signal intensity as a function of time. Fig. 8B is the continuation from Fig. 8A of repeated recording of a series of 6 ESR spectrum at the same time intervals except that the magnetic field strength was decreased by 2 Gauss in between each scan.

Figures 9A and 9B, respectively, are electron spin resonance spectra demonstrating 4-amino-TEMPO labelled and o-raffinose cross-linked and polymerized human hemoglobin and 3-maleimido-PROXYL labelled DBBF-hemoglobin polymerized with glutaldehyde.

Figures 10A, 10B, and 10C respectively, are electron spin resonance spectra of liposome encapsulated human hemoglobin containing (A) 3-DOXYL-cholestane (B) 16-DOXYL-stearic acid and (C) both 3-DOXYL-cholestane and 16-DOXYL-stearate.

Figure 11 is the electron spin resonance spectrum of nitroxide-labelled hemoglobin labelled with 4-amino-TEMPO and conjugated with dextran.

Figure 12 is an embodiment of a filter cartridge that contains a solid matrix to which a nitroxide is bound and through which a hemoglobin-containing solution may be passed.

INDUSTRIAL APPLICABILITY

The term "hemoglobin" is used generally herein to describe oxy-, carboxy-, carbonmonoxy-, and deoxy-hemoglobin


except as otherwise noted by the context of the description. The hemoglobin used with this invention may be human, recombinant or animal in origin and is obtained and purified by known techniques. The hemoglobin may be covalently bound to pyridoxal groups of pyridoxal-5'-phosphate or ring opened adenosine triphosphate (o-ATP) by reaction with the aldehyde groups and cross-linked derivatives of hemoglobin. The cross-linked derivatives may include polyfunctional, heterobifunctional and homobifunctional cross-linking reagents such as dialdehyde, polyaldehyde, diepoxide, polyepoxide, activated polycarboxyl and dicarboxyl groups, for example, 3,5-bis-bromosilicyl-bisfumarate, and TEMPO succinate or TOPS See (U.S. Patent No. 4,240,797) cyclodextrans and their anionic (e.g., sulfate) cross-linked hemoglobin as well as polymerized hemoglobin. All hemoglobin solutions described herein for use with this invention are physiologically compatible. The hemoglobin solutions are cell-free to remove pyrogens, endotoxins, and other contaminants.

The term "nitroxide" is used herein to describe stable nitroxide free radicals, their precursors, and their derivatives thereof including the corresponding hydroxylamine derivative where the oxygen atoms are replaced with a hydroxyl group and exist in hydrogen halide form. For the purposes of this invention, the chloride salt form of the hydroxylamine derivatives are preferred.

In the nitroxides described here, the unpaired electron


of a nitroxide is stable in part because the nitrogen nucleus is attached to two carbon atoms which are substituted with strong electron donors. With the partial negative charge on the oxygen of the N-O bond, the two adjacent carbon atoms together localize the unpaired electron on the nitrogen nucleus.

Nitroxides may have either a heterocyclic or linear structure. The fundamental criterion is a stable free radical. Structurally, nitroxides of the following formula are preferred where $R_1 - R_4$ are electron donors and A is the remaining members of a heterocyclic ring.




In these heterocyclic structures, "A" represents the remaining carbon atoms of a 5-membered (pyrrolidinyl or PROXYL with one double bond, i.e., pyrroline) or a 6-membered (piperidinyl or TEMPO) heterocyclic structure and in which one carbon atoms may be substituted with an oxygen atom (oxazolinyl or DOXYL) and certain hydrogen atoms may be substituted with up to two bromine atoms. In such heterocyclic structures, stable isotopes may be utilized (e.g., N_{15} ,


deuterium). Substitution at the α carbons should be such that the unpaired electron is maintained substantially in a πp orbital configuration. R_1 through R_4 are alkyls (straight and branched chain) or aryl groups but are preferred to be methyl or ethyl groups. The substituent groups on the alpha carbons in any nitroxide should be strong electron donors to enhance stability, thus methyl (CH_3) groups or ethyl (C_2H_5) groups are preferred although other longer carbon chain species could be used. In practice, stearic considerations may limit the scope of the nitroxide compounds that are practical and economical. The preferred nitroxides used with this invention include nitroxides having the following structure:



DOXYL
(4,4-Dimethyl-3-oxazolinyl-oxy)
(4,4-Dimethyloxazolidine-N-oxy)




PROXYL
(2,2,5,5-Tetramethyl-1-pyrrolidinyl-oxy)
(2,2,5,5-Tetramethylpyrrolidine-N-oxy)



TEMPO
(2,2,6,6-Tetramethyl-1-piperidinyl-oxy)
(2,2,6,6-Tetramethylpiperidine-N-oxy)

As is apparent from the above, most suitable nitroxide compounds may be represented basically by the structural formula



assuming that the R group is selected from among the configurations which preserve the stability of the free radical.

The nitroxides which can be employed in this invention are structurally diverse because the requisite property of the nitroxides is their ability to influence the course of the superoxide anion cascade in HRCS by mimicking the superoxide oxidase, superoxide dismutase, and catalase activities without substantially being consumed in the process. Although a wide variety of nitroxides may be used with this invention, the nitroxide should be physiologically acceptable at a minimum concentration required to alleviate oxygen toxicity in the HRCS. In assessing an operative species, it is noteworthy that the relatively low toxicity of nitroxides has encouraged their development as contrasting agent in NMR imaging (See U.S. Patent Nos. 4,834,964; 4,863,717; 5,104,641).

A number of methods for isolating and purifying hemoglobin solutions such that they are physiologically compatible are known to those skilled in the art. Typically, purified hemoglobin compositions contain at least 99% hemoglobin by weight of total protein, a total phospholipid content of less than about 3ug/ml, less than 1ug/ml of either phosphatidylserine or phosphatidylethanolamine and an inactive heme pigment of less than 6%. The purified hemoglobin solutions which are useful in this invention can be prepared using a variety of conventional techniques, including but are not limited to, those disclosed in Cheung et. al., Anal Biochem 137:481-484 (1984), De Venuto et. al., J. Lab. Clin. Med.

89:509-516 (1977), and Lee, et. al., Vith International Symposium on Blood Substitutes, San Diego, CA Mar. 17-20 Abstract H51 (1993).

The purified hemoglobin solutions used in this invention should be essentially free of oxygen. Hemoglobin in solution may be deoxygenated by admixture with a chemical reducing agent which causes the hemoglobin to release oxygen and to be maintained in a substantially deoxygenated state. A preferred method for deoxygenating a hemoglobin solution is performed by exposing a hemoglobin solution to an inert, essentially oxygen-free gas, such as nitrogen or carbon monoxide to cause removal of bound oxygen from the hemoglobin and conversion of the hemoglobin in solution to a form such as deoxy-hemoglobin or carbonmonoxy-hemoglobin that lacks oxygen. Alternatively, hemoglobin may be exposed to a vacuum or gas through a membrane that is permeable to oxygen yet impermeable to hemoglobin. For example, a hemoglobin solution may be passed through a diffusion cell having a membrane wall along which hemoglobin flows and through which oxygen is capable of passing, but hemoglobin is not. Inert gas is circulated along the side of the membrane wall opposite the hemoglobin solution causing the removal of oxygen and the conversion of the hemoglobin in solution to the deoxygenated state. Preferably, the deoxy-hemoglobin is maintained in an essentially oxygen-free environment during nitroxide labelling, cross-linking, polymerization, or

conjugation.


After removal of any bound oxygen, a nitroxide is covalently attached to the hemoglobin. Normally at least one, and preferably more than one, nitroxide will be covalently attached to a single hemoglobin molecule. The nitroxide may be covalently attached to the hemoglobin at any of several sites on the hemoglobin molecule including:

- (a) at one or more of the free sulfhydro (-SH) groups, for example, at the β - 93 site;
- (b) at any reactive amino (-NH₂) groups, for example, in the DPG site at Val-1 of the β -chain and/or lysine-82 of the β -chain and/or lysine-99 of the α -chain;
- (c) at any non-specific surface amino (-NH₂) or carboxyl (-COOH) group;

A nitroxide may also be bound to any residual aldehyde, epoxy, or activated carboxyl groups of a divalent- or a multivalent-cross-linker involved in the cross-linking and polymerization of hemoglobin or at any residual reactive groups on an agent such as dextran (Dx) or hydroxyethylstarch (HES) or polyoxyethylene (POE) used to conjugate hemoglobin.

As described in Equation [1], above, during the storage period, the hemoglobin in an HBOC solution is slowly auto-oxidized by oxygen to form met-hemoglobin and the superoxide anion. However, during the storage of the HRCS that are the subject of this invention, the superoxide anion thus formed

will reduce the nitroxide to a hydroxylamine derivative, and the superoxide anion will be oxidized to form molecular oxygen by the following reaction.




The conversion of superoxide anion to molecular oxygen described in Equation [4] prevents the accumulation of superoxide anion and the subsequent formation of hydrogen peroxide. This activity, described herein as a "superoxide oxidase" activity, will be most effective when the initial oxygen content in the composition is kept to a minimum, the composition is stored in an essentially oxygen free environment and the nitroxide concentration is sufficient to prevent the formation of superoxide anion and hydrogen peroxide. Therefore, storage of the HRCS in an essentially oxygen-free container is preferred.

Container systems that permit a solution to be stored in an oxygen free environment are well known because many non-hemoglobin based intravenous solutions are sensitive to oxygen. For example, a glass container that is purged of oxygen during the filling and sealing process may be used. Also, flexible plastic containers are available that may be enclosed in an overwrap to seal against oxygen. Basically, any container that prevents oxygen from interacting with hemoglobin in solution may be used.


To demonstrate the "superoxide oxidase" activity of a nitroxide, samples of nitroxide-labelled hemoglobin in solution are kept in an accelerated oxidative storage condition and the redox state of the nitroxide is studied over time by electron spin resonance spectroscopy. For example, an o-raffinose polymerized hemoglobin solution that has been labelled with 4-amino-TEMPO is stored in its oxygenated state in a sealed glass container (Figure 1A). In such a state, the rate of superoxide anion and met-hemoglobin formation in solution is sufficiently rapid that the conversion of the nitroxide to its hydroxylamine derivatives may be conveniently monitored (See Equation [4] and compare Figures 1A and 1B). Equation 4 represents that the conversion of nitroxide to its diamagnetic hydroxyl derivative is coupled to the conversion of the superoxide anion back to molecular oxygen. The experimental evidence in support of such a conversion is shown in Figures 1A and 1B. The electron spin resonance spectrum of TEMPO covalently attached to the hemoglobin (Figure 1A) was converted to its diamagnetic derivatives which result in the complete disappearance of the resonance peaks after storage of the sample for 30 days at 4°C (Figure 1B). The nitroxide is considered to have performed a "superoxide oxidase"-like activity when it is converted to its hydroxylamine derivative in the presence of hemoglobin.

The "superoxide dismutase" activity of a nitroxide in an

HBOC solution is demonstrated by showing the reconversion of the hydroxylamine derivative back to a nitroxide (See Equation [5] together with Equation [4]). Knowing that under the experimental conditions described in Figures 1A and 1B the nitroxide is fully converted to hydroxylamine (See Equation [4]), the nitroxide may be regenerated by simply providing more superoxide anion as shown in Equation 5. To demonstrate this reaction mechanism, the relative concentration of hemoglobin (and thus superoxide anion) to the nitroxide is increased by diluting the sample in Figure 1A with an equal volume of unlabelled hemoglobin. A comparison of Figures 1A and 1C shows an approximate 50% reduction of the signal intensity of the nitroxide due to the dilution effect. On the other hand, after 30 days of cold storage at 4°C, the nitroxide was partially regenerated (See Figure 1D) as predicted by Equation [5]. This observation is consistent with the reconversion of the hydroxylamine derivative to nitroxide coupled with the formation of hydrogen peroxide from superoxide anion.



Summing equations [4] and [5] results in:



which demonstrates that the nitroxide acts as a low molecular weight, metal-free, SOD mimic in "HBOC" solutions. The detection of electron spin resonance spectrum of the nitroxide (in Figure 1D) is consistent with the reaction of superoxide anion with the hydroxylamine ($\text{R}-\text{N}(\text{OH})_2$) resulting in the formation of nitroxide ($\text{R}-\text{N}(\text{O})$) and hydrogen peroxide (H_2O_2). Recently, oxoammonium cation has been proposed to be involved as one intermediate in the nitroxide catalyzed dismutation of superoxide. (Krishna et al., Proc. Nat. Acad. Sci. USA 89 5537-5541 (1992)).

The number of nitroxide molecules per hemoglobin molecule may be in the range of approximately 1-40 and for specific labelling is most preferably about 2. However, the nitroxide-hemoglobin ratio should be kept to a minimum due to pharmacokinetic, toxicological and immunological considerations. For example, a nitroxide such as 3-maleimido-PROXYL is covalently bound to hemoglobin in solution by first preparing a 100mM solution of the nitroxide in ethanol as the carrier solvent. Two(2) molar equivalents of the nitroxide to hemoglobin was added directly with mixing to a DCL-Hb (8g/dl) in Lactated Ringers. The reaction mixture was allowed to react at 22°C with agitation until greater than 90% of the nitroxide was covalently linked to the DCL-Hb, usually within one hour. The unreacted nitroxide was then removed with a cross-flow membrane filtration system having a molecular-weight cut-off of 30,000 daltons by washing three(3) times

with 10 volumes of Lactated Ringers. The retantate hemoglobin concentration is adjusted to between 7-14 g/dl, sterile filtered, and stored at 4°C. After transfusion, when the HRCS is fully oxygenated, the nitroxide is expected to function as a SOD-mimic and secondly as a catalase-mimic. As an SOD-mimic it dismutates the superoxide anion to hydrogen peroxide (See Equation [2]) and consequently protect against the destruction of nitric oxide in the endothelium to prevent vasoconstriction. As a catalase-mimic it prevents hydrogen peroxide toxicity by converting the latter to harmless water (See Equation [3]).

As noted above, nitroxides have been covalently bound to hemoglobin to study the cooperative oxygen binding properties of the hemoglobin molecule itself. However, nitroxides have not been used with stabilized, i.e., cross-linked, or polymerized, encapsulated, or conjugated hemoglobin solution that are physiologically compatible. The known chemistry of hemoglobin and nitroxides suggests that it is possible to perform similar nitroxide-labelling of hemoglobin that has been chemically cross-linked or cross-linked through recombinant techniques by selecting an available site on the hemoglobin molecule that is not blocked by the particular compound used to stabilize, polymerize, or conjugate the hemoglobin molecule(s). Because certain of the stabilized and polymerized forms of hemoglobin described below are currently involved in clinical trials, the attachment of

nitroxides to these stabilized and polymerized hemoglobin-based oxygen carriers is described below in the context of the second, third, and fifth preferred embodiments to demonstrate that the oxygen detoxification function of this invention is applicable to the existing hemoglobin solutions.

The nitroxide-labeling technology demonstrated here in the example of nitroxide-HBOC is readily applied to the production of other nitroxide-labelled macromolecules with useful antioxidant enzyme-mimetic activities, for example nitroxide-labelled serum albumin and nitroxide-labelled immunoglobulin. Forms of serum albumin which can readily be labelled by nitroxide by this technology are monomeric (normal) albumin, and albumin homodimers, oligomers, and aggregates (microspheres).

The antioxidant enzyme mimic effect of a nitroxide-labelled macromolecule, hemoglobin, or the other appropriate macromolecules described here, has utility in other applications, medical or otherwise, where antioxidant catalysis is useful.

MODES FOR CARRYING OUT THE INVENTION

I. First Preferred Embodiment -- Nitroxide Labelled Containers and Filters for HBOC

It is possible to provide the oxygen-detoxification function of this invention to existing HBOC solutions without chemically modifying the hemoglobin in these formulations. By covalently attaching nitroxides to a surface inside the

vessel in which the HBOC is stored, this invention may alleviate the adverse physiological effects caused by oxygen toxicity that are observed with the existing formulations.

The container used with the hemoglobin-containing solutions that are the subject of this invention should be physiologically compatible having similar characteristics as any container to be used with intravenous fluids. Typically, glass or a biocompatible plastic is suitable. For the embodiments of the invention where a solution containing hemoglobin is placed in a container for any length of time, the container should be oxygen free and capable of being sealed to exclude oxygen. With a glass container, a traditional stopper and closure means is sufficient. However, some of the flexible plastic containers currently available are oxygen permeable. In this case, a foil overwrap or other sealing mechanism may be used to prevent oxygen from contacting the hemoglobin in solution.

To apply a nitroxide to an inner surface of a container, a non-leaching layer of a nitroxide polymer or a nitroxide-doped copolymer is coated directly on the inner surface. Nitroxide-containing polymers can be created by a number of techniques based on generally known principles of polymerization as long as the stability of the free radical is maintained in the polymerization process.

Also, the interior surface of an HBOC container may be modified to contain a coating layer of a substance that can

bind a nitroxide, such as hydrophilic hydrazide groups which will react with the ketone or the aldehyde group of a nitroxide to form stable hydrazone derivatives. The coating reaction is straight forward. For example, the nitroxide (100mM) in acetate buffer at pH 5.0 is added to a hydrazide activated plastic container to facilitate the formation of a hydrazone linkage.

Once the container is prepared, a physiologically compatible solution is added. This solution may be a stabilized and purified HBOC or the HRCS disclosed herein, but could also include any intravenous colloid or crystalloid solution that is desirable to co-infuse with hemoglobin. The solution is then maintained in an essentially oxygen-free environment.

In addition to treating a surface inside a container, a filter-type cartridge, with a luer lock inlet and outlet, containing a gel or solid matrix upon which a nitroxide is immobilized may be used to remove reactive oxygen-derived reactive species while the hemoglobin solution passes through the cartridge. In these applications, the nitroxide is bound to a soft- or hard-gel matrix through which the HBOC passes, functioning essentially as a sterile in-line filter, prior to infusion. A variety of methods to attach small ligands, such as nitroxide, to a solid matrix are well known in the art of affinity chromatography, as are the techniques to chemically modify glass and plastic surfaces. Several types of matrices that are compatible with sterile solutions are known includ

ing agarose gel, polysulfone-based material, latex, and others.

In the filter cartridge approach, the solid matrix is covalently linked with a nitroxide and contained in a filter housing or other such apparatus such that a hemoglobin solution can flow through the apparatus and be brought into contact with a nitroxide while being infused into a patient. A practical approach is to use a commonly available activated agarose gel as the matrix and contain the gel in a sterile luer lock cartridge. The cartridge is then simply inserted in the fluid administration line during the transfusion of a solution containing hemoglobin. In practice, the structure that comprises the filter housing in which the nitroxide and through which hemoglobin is passed can be provided by a variety of known structures. Referring now to Figure 12, housing 1 contains a nitroxide-labelled agarose gel. For example, a 4-bromoacetamido-TEMPO labelled ω -aminohexyl-agarose (See Figure 2A) a 1,4-bis(2,3-epoxypropoxy)butane agarose coupled with 4-amino-TEMPO (See Figure 2B).

During the transfusion, the intravenous transfusion line containing a hemoglobin solution would be connected to the luer inlet 2 allowed to enter the housing 1 wherein the hemoglobin solution would encounter the nitroxide bound to the matrix 4 to remove the toxic oxygen-related species. The hemoglobin solution would then pass out of the cartridge through the luer outlet 3 and would be directly transfused

into a patient. The electron resonance spectrum of 4-amino-TEMPO labelled epoxy-agarose is shown in Figure 2A. Alternatively, an ω -aminohexyl-agarose may be reacted with 4-(2-bromoacetamido)-TEMPO to form TEMPO labelled agarose. The electron spin resonance spectrum is shown in Figure 2B. An alternative would be to couple the 4-carboxyl-TEMPO to the amino-agarose with carbodiimide via a carboamide linkage. Conversely, the 4-amino-TEMPO is readily coupled to the carboxyl group on an agarose gel using carbodiimide, for example, 1-ethyl-3-(3-dimethylamino-propyl)carbodiimide.

The cartridge labelled with 4-amino-TEMPO prepared by circulating a 100mM 4-amino-TEMPO (Sigma Chem. Co.) in a Lactated Ringers solution through an aldehyde AvidChrom Cartridge (Unisyn Tech. Inc.) at room temperature for one hour followed by the reduction with sodium cyanoborohydride for six (6) hours. The interior of the cartridge housing is thoroughly washed with Lactated Ringers.

The cartridge labelled with 3-amino-PROXYL may be similarly prepared by substituting 4-amino-TEMPO with 3-amino-PROXYL according to the procedure described above.

II. Second Preferred Embodiment -- Nitroxide-Labelled

Stabilized Hemoglobin

To prevent dissociation of hemoglobin into its constituent subunits, hemoglobin is intramolecularly stabilized by chemical or recombinant cross-linking of its subunits. "Stabilized" hemoglobin is referred herein to describe

hemoglobin monomers that are stabilized by chemical or recombinant cross-linking and also to describe dimers, trimers, and larger oligomers whose constituent hemoglobin molecules are stabilized by cross-linking with cyclodextrans and their sulfated derivatives.


A preferred technique and best mode for attaching nitroxide to stabilized hemoglobin is by the covalent attachment of the nitroxide to the β -93 sulfhydryl groups of the two β -chains of stabilized hemoglobin. Although specific labelling at the β -93 site has been demonstrated on native human hemoglobin for conformational studies (See review by McConnell et. al., *Quart. Rev. Biophys.* 3:91 (1970)), such a specific labelling of cross-linked hemoglobin has not been reported. As noted above, several types of hemoglobin-based oxygen carriers have been developed that are stabilized through chemical cross-linking with DBBF, diaspirin cross-linked hemoglobin and hemoglobin that is stabilized and oligomerized with o-raffinose.

The ring opened sugars described in my United States Patent No. 4,857,636 yield polyvalent aldehydes derived from disaccharides, oligosaccharides, or, preferably, trisaccharides such as o-raffinose. These compounds function both to provide intramolecular stabilization (cross-linking) and intermolecular polymerization. By controlling the reaction disclosed in my earlier patent, the polyvalent aldehydes may be used to produce "stabilized" hemoglobin as

defined above without polymerization. In another case, a nitroxide may be covalently bound to the stabilized hemoglobin or the polymerized hemoglobin. Therefore, the hemoglobin-based solutions that are stabilized using the polyvalent aldehydes are considered in the present embodiment as a "stabilized" hemoglobin and in the subsequent embodiment as a polymerized hemoglobin.


To demonstrate, that the β -93 site of the chemically modified hemoglobin has not been rendered sterically inaccessible for nitroxide attachment, results are presented to confirm that a nitroxide may be covalently bound to the β -93 site of DBBF-Hb.

In this embodiment, DBBF-Hb is reacted with two types of nitroxides (TEMPO and PROXYL) which contain two types of sulfhydryl group specific functional groups and have the following structural formula:



(I)

4-(2-Bromoacetamido)-TEMPO



(II)

3-Maleimido-PROXYL

DBBF-Hb is prepared by cross-linking purified deoxygenated hemoglobin in solution with bis(3,5 dibromosalicyl)fumarate by known techniques, and the resulting product is purified by column chromatography. The covalent attachment of 3-maleimido (2,2,5,5 -tetramethyl pyrrolidine-N-Oxyl) [3-maleimido-PROXYL] is accomplished by adding 2 molar equivalents of this nitroxide using methanol as the carrier solvent at a concentration of approximately 100mM of 3-maleimido-PROXYL to 1ml of DBBF-Hb at a concentration of approximately 8 g/dl in Lactate Ringers. The DBBF-Hb is allowed to react at 22-23°C for approximately 30 minutes with mixing. The extent of cross-linking is estimated from the percent disappearance of the electron spin resonance signal intensity of the unreacted nitroxide. To remove the unreacted nitroxide, the reaction mixture was washed three (3) times with a 10 volume excess of Lactated Ringers using a Filtron stire cell with a 30 kilodalton cut-off nominal molecular weight limits (NMWL) polyethylene sulfone (PES) membrane (Filtron Technology Co.). The electron spin resonance measurements of the nitroxide-labelled hemoglobin was recorded with a Bruker ESR spectrometer. Figure 3A shows the electron spin resonance spectra of 4-(2-bromoacetamido)-TEMPO labelled DBBF-Hb. The electron spin resonance spectrum of DBBF-Hb that is similarly labelled with 3-maleimido-PROXYL is shown in Figure 3B.

In this embodiment, the nitroxide is covalently linked to

the lone sulfhydro group on the two β -globin chains of hemoglobin. Thus, the nitroxide to hemoglobin-bound oxygen ratio is approximately 200 to 1 at 99.00% deoxy-hemoglobin because there are two nitroxides attached to the two β -globin chains of the hemoglobin. After transfusion, however, the deoxygenated HRCS picks up oxygen in the lung and the nitroxide to hemoglobin-bound oxygen ratio becomes approximately 1 to 2 at 100% oxygenation because there are four oxygen molecules bound to the four globin chains of the hemoglobin with the two nitroxides remaining on the β -globin chains.

Using a hemoglobin-to-nitroxide ratio of 1:2, greater than 90% of the nitroxide is covalently attached to the DBBF-Hb. DBBF-Hb may also be covalently labelled with a spacer group (e.g., an extra methyl group) between the maleimido and PROXYL moieties (i.e., 3-maleimidomethyl-PROXYL) which would exhibit a resonance spectrum similar to that of Figure 3B. It is noteworthy that other nitroxides may be covalently attached to specific amino-groups in the DPG binding site (e.g., β -Val-1 β -Lys-82 and α -Lys-99) or may be attached to the remaining 40-plus surface lysine ϵ -amino groups on hemoglobin. Isothiocyanate derivatives of the TEMPO and PROXYL nitroxides are also reactive with the amino group. For example, 4-isothiocyanate-TEMPO may be added to hemoglobin at a molar ratio of approximately 10:1. Resonance spectrum (not shown) of hemoglobin labelled with this

nitroxide at other sites is similar to that shown in Figure 3A.

The ability to attach nitroxides at several sites of DBBF-Hb suggests that recombinant hemoglobin that is stabilized with alpha-globin dimers (D. Looker et.al. NATURE 356:258 (1992)) may be similarly labelled with a nitroxide. It is also possible to prepare a DBBF analogue of a nitroxide labelled cross-linking agent such as a TEMPO labelled succinate (See U.S. Patent No. 4,240,797).

Figure 4 is ESR spectra of (A) 2-(bromoacetamido)-TEMPO, (B) 2-(bromoacetamido)-TEMPO-labelled HBOC and (C) $^{15}\text{ND}_{17}$ TEMPOL (TEMPOL: 4-hydroxy-2,2,6,6-tetramethylpiperidine-1-oxyl) in Lactated Ringer's solution recorded at room temperature. The difference in Figure 4A and 4B represents the difference in the mobility of a small molecular weight nitroxide to that of a nitroxide covalently attached to a macromolecule such as hemoglobin. Figure 4C shows that a stable isotope nitrogen ^{15}N with a nuclear spin of 1/2 yields two resonance peaks and that natural-isotopic ^{14}N with a nuclear spin of 1 yields three resonance peaks (compare 4A to 4C). In the set of experiments described here the separation of these resonance peaks is used to demonstrate the enzyme-mimic and *in vivo* and *in vitro* oxidation/reduction reactions of small and macromolecular weight nitroxides.

Nitroxide labelled HBOC with different molar ratios of nitroxide to hemoglobin are prepared as follows. 2, 4, and

8 molar equivalents of 4-(2-bromoacetamido)-TEMPO, were added as solid powder directly into three separate 15 ml Vacutainers in a clean hood. After replacing the rubber septum, 4-(2-bromoacetamido)-TEMPO was subsequently dissolved in 200 ul chloroform. The Vacutainers were then connected to high vacuum (5 mm Hg) via a 27 gauge needle through the rubber septum and the chloroform was removed leaving a thin film of 4-(2-bromoacetamido)-TEMPO coating the lower half of the Vacutainer. After introducing the appropriate amount of HBOC via sterile transfer through the rubber septum, the solutions were allowed to react at room temperature with intermittent vortex mixing at approximately 5 minute intervals for 1/2 hour (not all solids were dissolved in the 4 and 8 molar ratios of nitroxide to hemoglobin), the Vacutainers were then left at 4 degrees C in a refrigerator over night. Vortex mixing at room temperature was resumed the next morning for another 1/2 hour until all solids of 4-(2-bromoacetamido)-TEMPO had visually disappeared from the surface of the Vacutainer.

The reaction mixtures and the control, were then transferred to a sterile dialyzing tube and dialyzed against Lactated Ringers until no unlabelled free 4-(2-bromoacetamido)-TEMPO electron spin resonance (ESR) signals could be detected. The ESR spectra of 4-(2-bromoacetamido)-TEMPO-labelled HBOC at 2, 4, and 8 molar equivalents 4-(2-bromoacetamido)-TEMPO to Hb are shown in Figures 5A-5C respective-

ly. At 2 molar equivalents of 4-(2-bromoacetamido)-TEMPO to hemoglobin, the ESR spectra are essentially the same with or without dialysis indicating the covalent labeling is quantitative. The two SH- groups on the beta globulin chains appear to be the site of covalent attachment in the case of HBOC (this can be confirmed by selective blocking of 4-(2-bromoacetamido)-TEMPO labeling with N-ethyl-maleimide or globulin chain analysis by reverse phase HPLC). It is noteworthy that the ESR signal intensity (peak Mo) ratios for 2, 4, and 8 are in approximately the same ratio as the spectra were recorded at proportionately decreasing instrument sensitivity.

Furthermore, it is expected that more 4-(2-bromoacetamido)-TEMPO could be attached to Hb at even higher molar ratios, for example as radiation-protective agents *in vivo*.

The preferred molar of nitroxide to hemoglobin in the blood substitute formulations is 8:1 as described below.

Referring to Figure 6, an ESR spectrum of a mixture of 4-(2-bromoacetamide)-TEMPO-labelled HBOC and $^{15}\text{ND}_{17}$ -TEMPOL wherein the center peak of the 4-(2-bromoacetamido)-TEMPO (indicated by down arrow) and the high field peak of $^{15}\text{ND}_{17}$ -TEMPOL (indicated by the up-arrow) were adjusted to similar intensity.

The separation of the resonance peaks permits the simultaneous monitoring of free radical or enzyme mimic activities involving the small molecular weight nitroxide (TEMPOL) and

its macromolecular conjugate in both *in vitro* and *in vivo* (murine) reactions. For example, the *in vivo* plasma half-life of the two nitroxides was compared by referring to the unique spectral characteristics of the different nitroxides. Specifically, the *in vivo* ESR studies of hemoglobin-based solutions, on the mouse were performed using a nitroxide to hemoglobin ratio of 8:1 (see Fig. 5C) to take advantage of its high ESR signal intensity. First, the approximate plasma half-life of a small molecular weight nitroxide ($^{15}\text{ND}_{17}$ -TEMPOL see Fig. 4C) and a large molecular weight 4-(2-bromoacetamido)-TEMPOL-labelled HBOC (see Fig. 4B) are determined by preparing a mixture of the two and adjusting the ESR signal intensity to be approximately the same (see Fig. 6). 0.5 ml of the mixture was injected under anesthesia into a distended mouse tail vein under a heat lamp. The mouse tail was inserted into an ESR cavity and the spectrum was recorded within 10 min. after injection (see Fig. 7A).

Referring to Figure 7A, 7B, and 7C the $^{15}\text{ND}_{17}$ -TEMPOL signal could not be detected, however, the 4-(2-bromoacetamido)-TEMPO-labelled HBOC was clearly resolved (see Fig. 7B and 7C for plasma half-life studies where 7C is a continuation of 7B). Since the vasoconstrictive effect of HBOC is reported to be fully developed during the first 5-15 min. of bolus injection of an HBOC in rats, the participation of the nitroxide-labelled HBOC in free radical redox-reactions immediately after transfusion in a mouse was measured. The

tail vein of female CH3 mouse was cannulated under anesthesia with 80% nitrous oxide, 20% oxygen, and 3% isofluorane. Under a heat lamp the mouse tail vein became visibly distended, a cannula consisting of a 30 gauge hypodermic needle attached to a one foot length of polyethylene tubing was inserted into the tail vein and held in place by cyanoacrylate glue. For *in vivo* ESR measurements, the cannulated mouse was transferred under anesthesia to a 50 ml conical centrifuge tube modified to allow the tail to protrude from the conical end and to allow a continuous flow of anesthetic gas from the opening end of the tube. The tail was inserted into a plastic tube which was then fitted into a TE 102 cavity. The cannula was flushed periodically with heparin (100 unit/ml) to ensure patency. The cannula was near the root of the tail and was kept outside of the ESR cavity so that a pure signal from the tail could be measured immediately after bolus injection. 0.5 ml of samples (see Fig. 8) were injected via the cannula and the spectrometer was set for a repeat scanning mode at 1/2 min. intervals (see Figs. 8A and 8B). In Fig. 8A the magnetic field was increased by two Gauss, and in Fig. 8B the magnetic field was decreased by two Gauss, to superimpose the resonance spectra. The $^{15}\text{ND}_{17}$ -TEMPOL signal disappeared within 2.5 minutes after injection. During the same time period the 4-(2-bromoacetamido)-TEMPOL-labelled HBOC also decreased at a similar rate.

However, the nitroxide-HBOC signal were shown to be

stable in plasma (Fig. 8B). Therefore, Fig. 8B together with results from Fig. 7 show that the nitroxide-labelled to macromolecules such as HBOC has considerably longer plasma half-life as compared to small molecular weight nitroxide (e.g., $^{15}\text{ND}_{17}$ -TEMPOL).

The observed nature of the free radical reaction involves two pathways:

1. the rapid phase appears to involve the free radical (e.g. superoxide) oxidation of the nitroxide to its oxoammonium cation intermediate followed by the reduction of the oxoammonium cation to its stable hydroxylamine derivative of the nitroxide. Such reduction involves the participation of either one or two reducing equivalents (e.g. NADH) present in the vascular compartment. The reduction of nitroxide to its hydroxylamine would lead to a rapid reduction in ESR signal intensity, in the case of 8:1 molar ratio of 4-(2-bromoacetamido)-TEMPO-labelled HBOC represents approximately 25% of the 4-(2-bromoacetamido)-TEMPO on the HBOC. This phase involves both small molecule and macromolecular nitroxide.
2. the slow phase appears (see Figure 8B) to represent the antioxidant enzyme-mimic activities of the remaining 75% of 4-(2-bromoacetamido)-TEMPO on the HBOC in accordance with the reaction mechanism

wherein the nitroxide is involved in the cyclic-free radical reactions for example the SOD-mimic reaction. Where the nitroxide free radical is essentially unconsumed as a SOD-mimic, the slow rate of decrease of the ESR signal intensity can be attributable primarily to the reaction mechanism described above and secondarily to the decrease in HBOC concentration as it is slowly eliminated from the vascular compartment as a function of its plasma half-life.

This result demonstrates the utility of nitroxide labelled HBOC in detoxifying free radicals *in vivo*. This utility is defined in terms of providing short term (in minutes) scavenging of free radicals and persistent (in hours) protection against oxidant reactions by nitroxides acting as enzyme mimics *in vivo*.

Based on the analysis of the spectra in Figure 8, the oxidation/reduction (redox) cycling reactions involve approximately 73% of 4-(2-bromoacetamido)-TEMPO-labelled HBOC remaining in its free radical state. This indicates that TEMPOL participates in *in vivo* redox-reactions within the confines of the vascular space.

III. Third Preferred Embodiment -- Nitroxide-Labelled Polymers of Stabilized Hemoglobin

While it is possible to produce dimers of stabilized hemoglobin from cross-linked monomers, it is also possible to

produce hemoglobin polymers from stabilized or native hemoglobin. Solutions of hemoglobin polymers contain a mixture of monomers, dimers, trimers, tetramers, and other oligomers. Solutions containing polymerized hemoglobins used as an HBOC generally have longer plasma circulation times and higher oxygen carrying capacities as compared to stabilized monomeric hemoglobin. Such polymerized hemoglobin may be prepared by a number of pathways using several different polymerizing agents. (See, U.S. Pat. Nos. 4,001,200, 4,857,636, and 4,826,811). The preferred method of introducing a nitroxide to a solution of polymerized hemoglobin is again by covalently attaching a nitroxide to the β -93 sulfhydryl groups of the two β -globin chains of hemoglobin. These sulfhydryl groups are not known to be involved in the stabilization or polymerization processes. Consequently, the nitroxide is preferably covalently attached to hemoglobin before the stabilization and polymerization of the hemoglobin monomers.

For example, nitroxide is covalently attached to DBBF-Hb according to the procedure described in the second embodiment above, followed by polymerization with glutaldehyde according to the procedure described in Sehgal et. al. U. S. Patent No. 4,826,811. Figure 4B is an electron spin resonance spectra of the DBBF-Hb labelled with 3-maleimido-PROXYL and polymerized with glutaldehyde. Similarly, DBBF-Hb that is polymerized with glutaldehyde may be labelled with 4-(2-bromo-

acetamido)-TEMPO by the same method.

Using a similar approach, a polymerized hemoglobin intermediate, such as a glutaldehyde-polymerized, an o-raffinose-polymerized, or an o-cyclodextran-polymerized hemoglobin intermediate that contains unreacted aldehyde groups, may be used for covalent attachment of either 4-amino-TEMPO or 3-amino-PROXYL via reductive amination to yield a nitroxide-labelled hemoglobin polymer. With reductive amination, the sequence and timing of the reaction are important. The 4-amino-TEMPO is added to glutaldehyde-polymerized hemoglobin after completion of polymerization, but prior to the reduction reaction that results in covalent attachment of the nitroxide to the polymerized hemoglobin. Likewise, the nitroxide labelling of a o-raffinose polymerized hemoglobin may be accomplished by the addition of either 4-amino-TEMPO or 3-amino-PROXYL prior to reductive amination. For example, 4-amino-TEMPO labelled o-raffinose polymerized hemoglobin is prepared according to the procedure described in my United States Patent No. 4,857,636 except that 6 molar equivalents of 4-amino-TEMPO is added after the completion of the polymerization and prior to the reduction with 20 molar excess of borane dimethylamine complex. As described therein, hemoglobin may be cross-linked and polymerized using polyvalent aldehydes derived from disaccharides or ring-opened sugars including, oligosaccharides, or preferably, trisaccharides such as o-raffinose. Likewise,

monosaccharides may be used to stabilize and polymerize hemoglobin although the higher molecular weight sugars are preferred. The resonance spectrum of a dialyzed and washed o-raffinose polymerized hemoglobin labelled with 4-amino-TEMPO was shown in Figure 9A.

To increase the yield of hemoglobin oligomers (Hb_n where $n=2-4$) of the polymerized hemoglobin, it is desirable to increase the valance of the polyaldehyde of the cross-linker, with the use of α -cyclodextran, β -cyclodextran, and γ -cyclodextran, as well as their sulfate derivatives which represents 6-, 7-, and 8-cyclized glucose molecules, the ring opened α -cyclodextran, β -cyclodextran, and γ -cyclodextran have 12, 14, and 16 reactive aldehyde groups respectively. These ring-opened cross-linkers can be used to cross-link and polymerize hemoglobin to produce polymerized hemoglobin which is rich in oligomers. The unreacted aldehyde, as described above, may be utilized to covalently attached to an amino-nitroxide, for example, 4-amino-TEMPO or 3-amino-PROXYL.

Furthermore, the ring-opened sulfate derivatives, for example, the sulfated α -cyclodextran will be an effective cross-linker for two additional reasons: (1) the sulfate groups will mimic the activity of DPG in lowering the oxygen affinity of the cross-linked hemoglobin, thus improving oxygen transport properties, and (2) the sulfate groups will serve as affinity labels which will complex multiple (e.g., $n=2-4$) hemoglobins to initially form a "cluster." Once the

"cluster" complex is formed, the aldehyde groups on the cyclodextran will be brought to close proximity with the NH₂ groups within the DPG binding sites, thus promoting the covalent intra-subunit and inter-molecular cross-linking of hemoglobin resulting in an increased yield of hemoglobin oligomers. In addition to antioxidant enzyme-mimic activities, the ring-opened cyclodextran polymerized and nitroxide-labelled hemoglobin will also have improved yield and composition as compared to o-raffinose and glutaldehyde polymerized hemoglobin.

IV. Fourth Preferred Embodiment -- Nitroxide-Labelled Liposome-Encapsulated Hemoglobin

Liposomes are particles which are formed from the aggregation of amphophilic molecules to form a bilayer structure in a hollow spherical shape with the polar sides facing an internal water compartment and external bulk water. Several acceptable methods for forming liposomes are known in the art. Typically, molecules form liposomes in aqueous solution like dipalmitoyl phosphatidylcholine. Liposomes may be formulated with cholesterol for added stability and may include other materials such as neutral lipids, and surface modifiers such as positively or negatively charged compounds. The preferred liposomes are small unilamellar-bilayered spherical shells.

A method for encapsulating hemoglobin in a liposome is also known (See Farmer et. al., United States Patent No.

4,911,921). For the purpose of this invention, a number of approaches may be used to introduce the nitroxide-based oxygen detoxification function to a solution of liposome-encapsulated hemoglobin. For example, it is possible to use nitroxide-labelled native hemoglobin, or a nitroxide-labelled stabilized hemoglobin as disclosed above, as the starting material and then performing the process of liposome encapsulation of the nitroxide-labelled hemoglobin by known techniques. In the present embodiment, purified hemoglobin may also be coencapsulated with a membrane impermeable nitroxide such as TEMPO-choline chloride disclosed for a spin membrane immunoassay in Hsia et. al. United States Patent No. 4,235,792.

Also, any purified hemoglobin may be encapsulated with a liposome comprised of nitroxide-labelled fatty acids (e.g., 7-DOXYL-stearate, 12-DOXYL-stearic acid, and 16-DOXYL-stearate), cholestane, an analogue of cholesterol (e.g., 3-DOXYL-cholestane), or phospholipid (e.g., 12-DOXYL-stearate-labelled phosphatidylcholine). The preparation of hemoglobin encapsulated in a liposome comprised of 3-DOXYL-cholestane labelled may be prepared by a method analogous to that described in Tabushi et. al., (J. Am. Chem. Soc. 106: 219 (1984)). A 5ml chloroform solution containing lipid compositions, including DOXYL labelled stearic acid and/or cholestane, as specified below were first dried in a stream of nitrogen to remove the solvent. Next, the residues were

dried in vacuo and the resulting film was suspended in 2ml of hemoglobin (24g/dl) in a Lactated Ringers solution. The lipid concentration in the dispersion is 100 mM. The liposome encapsulated hemoglobin is then rotated and incubated preferably at 37°C until all lipids are dispersed to form multilamellar vesicles. The resulting solution containing multilamellar liposome encapsulated hemoglobin and free unencapsulated hemoglobin is then forced through a microfluidizer to form 0.2 micron liposomes according to the procedure of Cook et.al. (See U.S. Pat.4,533,254). The molar ratio of dipalmitoyl phosphatidylcholine: cholesterol: dipalmitidyl phosphatidic acid: 3-DOXYL-cholestane in the liposome is 0.5:0.4:0.02:0.07. The resonance spectrum of the resulting 3-DOXYL-cholestane labelled liposome-encapsulated hemoglobin is shown in Figure 10A. In this configuration, the nitroxide is intercalated into the liposome membrane and can be found at both the inner and outer surface of the lipid bilayer water interface. Substituting the 3-DOXYL-cholestane with 16-DOXYL-stearic acid in the lipid composition shown in Figure 10A results in an electron resonance spectrum shown in Figure 10B. The mobility of the nitroxide as reflected from the resonance spectrum is consistent with the interpretation that the DOXYL-moiety of the stearic acid is located predominately in the hydrophobic interior of the lipid bilayer. With the addition of both the 3-DOXYL-cholestane and 16-DOXYL-stearate to the lipid composition at the same molar

ratio, the resonance spectrum of the double nitroxide labelled liposome encapsulated hemoglobin is shown in Figure 10C. The resonance spectrum of Figure 10C is a composite of Figures 10A and 10B because the nitroxides in this embodiment are located at both the membrane-water interface and its hydrophobic lipid bilayer interior. By placing the nitroxide in both locations, this embodiment provides the oxygen detoxification function at both the lipid bilayer hydrophobic interior and the membrane-water interface thus providing the added benefit of an additional reserve of oxygen-detoxification capacity for the encapsulated hemoglobin.

V. Fifth Preferred Embodiment -- Nitroxide-Labelled Conjugated Hemoglobin

A physiologically compatible solution of conjugated hemoglobin is produced by forming a conjugate of hemoglobin and a biocompatible macromolecule used as a plasma expander. Plasma expanders, such as dextran (Dx), polyoxyethylene (POE), hydroxyethyl starch (HES), are used to increase the circulation half life of hemoglobin in the body. In this state, the hemoglobin molecules together with the biocompatible macromolecule are collectively referred to as a hemoglobin conjugate. There are a number of convenient methods to incorporate a nitroxide into a hemoglobin conjugate. For example, one may simply substitute the hemoglobin to be conjugated with a nitroxide-labelled hemoglobin such as TEMPO labelled DBBF-Hb. This can be accomplished by substi-

tuting hemoglobin or pyridoxylated hemoglobin with 3-maleimido-PROXYL-DBBF-Hb or 4-(2-bromoacetamido)-TEMPO-DBBF-Hb in the preparation of conjugated hemoglobin.

4-Amino-TEMPO labelled dextran conjugated hemoglobin is prepared in accord with the procedure described by Tam et al. (Proc. Natl. Acad. Sci., 73:2128 (1976)). Initially, an 8% hemoglobin solution in 0.15 M NaCl and 5 mM phosphate buffer, pH 7.4 is conjugated to periodate-oxidized dextran to form a Schiff-base intermediate. Twenty molar equivalents of 4-amino-TEMPO is added to hemoglobin to form the Schiff-base between the nitroxide and the remaining reactive aldehyde groups on the dextran. After a 30 minute of incubation at 4°C, a 50 molar equivalent of dimethylamine borane in water is added. The solution is incubated for a further 2 hours at 4°C. Afterwards, the solution is dialyzed, reconstituted with Lactate Ringers buffer and sterile filtered with Filtron membrane filtration units (Filtron Technology Co.). The electron spin resonance spectrum of the 4-amino-TEMPO labelled dextran-conjugated hemoglobin is a sharp asymmetric triplet reflecting a high degree of motional freedom (See Figure 11). The increased mobility of the TEMPO covalently attached to the Dextran is consistent with the nitroxide linked to a flexible polysaccharide dextran chain as compared to that of a tightly folded hemoglobin molecule (See Figures 3A and 3B). Thus, resonance spectrum in Figure 11 demonstrates that a novel nitroxide labelled dextran conjugated

hemoglobin has been prepared.

VI. Sixth Preferred Embodiment -- Enzyme-Mimetic Activity of Nitroxide-Labelled Albumin

As noted above, nitroxides (e.g., TEMPOL) have been shown to have catalytic activity which mimics that of SOD, the metalloenzyme which dismutates superoxide to hydrogen peroxide. Furthermore, in biological systems nitroxides can interact with peroxidases and pseudoperoxidases to achieve an activity mimicking that of catalase, the enzyme which converts hydrogen peroxide to oxygen. The biological effects of nitroxide include contributing to protection against cytotoxicity of reactive oxygen species, apparently by reducing oxidative stress, for example, nitroxides protect against damage by ionizing radiation *in vivo* and *in vitro*. Thus, nitroxides when administered *in vivo*, display complex antioxidant enzyme-mimetic activities.

When injected intravenously, TEMPOL has been shown to have very short plasma half-life. Due to its molecular size and charge characteristics, it readily leaves the vascular space. In certain medical applications, other than those involving HBOC or HRCS, however, it may be desirable to have an antioxidant enzyme mimic which persists in the vascular space. This may be achieved by attaching a nitroxide compound to a macromolecule other than hemoglobin and which is biologically safe and has a desirable plasma half-life.

An example of such a desirable macromolecules is human serum albumin (HSA).

Serum albumin is a plasma protein with multiple ligand-binding sites and is the transport protein for many ligands in the blood. Nitroxides can bind specifically to human serum albumin at a number of specific ligand binding sites, or non-specifically.

Nitroxide-labelled albumin may be used *in vivo* to provide protection against cellular damage by reactive oxygen species. Nitroxide-albumin may be used either alone or in combination with a low molecular weight nitroxide compound, e.g., TEMPOL. Nitroxide-labelled albumin is also available as an "improved" version of albumin (i.e., improved by having antioxidant activity) with utility in any application where albumin is now conventionally used, including as a parenteral colloid solution, in biomaterials, in biocompatible surface coatings, etc.

The albumin may be obtained from plasma or may be produced by recombinant genetic means. HSA may be used in a variety of forms, including monomers (normal plasma form), homodimers, oligomers, and aggregates (microspheres). Specific labelling of the albumin with a nitroxide may be achieved at several binding sites, including bilirubin, FFA, indole, or Cu⁺⁺ binding site by using nitroxide compounds which have been activated in order to confer upon them binding specificity of the relevant site on the protein. A

preferred example is 2, 2, 6, 6-tetramethyl-1-oxyl-4-piperidylidene succinate (TOPS) nitroxide covalently bound to the primary bilirubin-binding site of HSA. Non-specific labeling of albumin may be achieved at approximately 50 accessible amino groups.

VII. Seventh Preferred Embodiment -- Nitroxide-Labelled Immunoglobulin

As in the above embodiments, certain nitroxides have been shown to have very short plasma half-life when injected intravenously. Due to the desire to have an antioxidant enzyme mimic with a long plasma half-life, a nitroxide compound may be attached to an immunoglobulin to provide long-lasting antioxidant enzyme mimic activity.

Immunoglobulins are a class of plasma proteins produced in the B-cells of the immune system and which are characterized by two specific ligand binding sites (the antigen-binding sites). Nitroxides have been used in the past as probes in research on hapten-binding specificity and affinity of immunoglobulins during the primary and secondary immune response.

As with the above-embodiment describing nitroxide-labelled albumin, the nitroxide-labelling technology demonstrated above in the example of nitroxide-HBOC is readily applied to the production of nitroxide-labelled immunoglobulins.

Nitroxide-labelled immunoglobulins may be used *in vivo* to

provide protection against cellular damage by reactive oxygen species. Nitroxide-immunoglobulin may be used either alone or in combination with a low molecular weight nitroxide compound to provide extended antioxidant activity with an extended plasma half-life.

Nitroxide-labelled immunoglobulin may be prepared by specific labelling of the immunoglobulin itself or by covalently labelling at a hapten-binding site. To avoid clearance of the nitroxide-labelled immunoglobulin as part of the body's natural immune response, one may use immunoglobulin fragments, for example, (Fab)₂, produced by cleaving the immunoglobulin according to known techniques with non-specific-labelling, a preferred molar ratio of nitroxide:-immunoglobulin is up to 60:1.

Although the invention has been illustrated by the specific embodiments described above, due to variety of forms of physiologically compatible hemoglobin and the structural diversity of the stable nitroxide-free radicals, a number of variations on the above embodiments are possible without departing from the basic spirit of the invention as described by the following claims.

I Claim as follows:

1. A physiologically compatible hemoglobin-based solution characterized by
stabilized hemoglobin, and
5 a nitroxide covalently bound to the stabilized hemoglobin.


2. The hemoglobin-based solution of Claim 1 wherein the hemoglobin is stabilized by cross-linking.

3. The hemoglobin-based solution of Claim 1 wherein the
10 hemoglobin is recombinant and stabilized with alpha-globin dimers.

4. The hemoglobin-based solution of Claim 2 wherein the hemoglobin is cross-linked using a polyvalent aldehyde derived from a ring-opened linear sugar.

- 15 5. The hemoglobin-based solution of Claim 2 wherein the hemoglobin is cross-linked using a polyvalent aldehyde derived from a ring-opened cyclic sugar.


6. The hemoglobin-based solution of Claims 1, 2, 3, 4, or 5, wherein the nitroxide has the structure



where $R_1 - R_4$ are alkyl groups of 1-4 carbon atoms and A is the remaining members of a 5-membered ring.

- 5 7. The hemoglobin-based solution of Claim 6 wherein the nitroxide is 2,2,5,5-tetramethylpyrrolidine-N-oxyl.

8. The hemoglobin-based solution of Claim 1, 2, 3, 4, or 5 wherein the nitroxide has the structure



where $R_1 - R_4$ are alkyl groups of 1-4 carbon atoms and
10 A is the remaining members of a 6-membered ring.

9. The hemoglobin-based solution of Claim 8 wherein the nitroxide is 2,2,6,6-tetramethylpiperdine-N-oxyl.


10. A physiologically compatible hemoglobin-based solution characterized by

5 a polymer hemoglobin, and
a nitroxide covalently bound to the polymerized hemoglobin.

11. The hemoglobin-based solution of Claim 10 wherein the hemoglobin polymer comprises a polyvalent aldehyde derived
10 from a ring-opened linear sugar.

12. The hemoglobin-based solution of Claim 11 wherein the hemoglobin polymer comprises a polyvalent aldehyde derived from a ring-opened cyclic sugar.


13. The hemoglobin-based solution of Claims 10, 11, or 12
15 wherein the nitroxide has the structure



where $R_1 - R_4$ are alkyl groups of 1-4 carbon atoms and A is the remaining members of a 5-membered ring.

14. The hemoglobin-based colution of Claim 13 wherein the nitroxide is 2,2,5,5-tetramethylpyrrolidine-N-oxyl.

5 15. The hemoglobin-based solution of Claims 10, 11, or 12 wherein the nitroxide has the structure



where $R_1 - R_4$ are alkyl groups of 1-4 carbon atoms and A is the remaining members of a 6-membered ring.


16. The hemoglobin-based solution of Claim 15 wherein the
10 nitroxide is 2,2,6,6-tetramethylpiperdine-N-oxyl.

17. A physiologically compatible hemoglobin-based solution characterized by

a physiologically compatible liposome,

hemoglobin encapsulated in the liposome, and
a nitroxide encapsulated in the liposome.


18. The hemoglobin-based solution of Claim 17 wherein the nitroxide is intercalated in the liposome membrane.
- 5 19. The hemoglobin-based solution of Claim 18 wherein the hemoglobin is stabilized by cross-linking or polymerization.
20. The hemoglobin-based solution of Claim 17 wherein the liposome is comprised of nitroxide-labelled fatty acids,
10 cholestone, or phospholipids.
21. The hemoglobin-based solution of Claims 17, 18, 19, or 20 wherein the nitroxide has the structure



where $R_1 - R_4$ are alkyl groups of 1-4 carbon atoms and
A is the remaining members of a 5-membered ring.

22. The hemoglobin-based solution of Claim 21 wherein the nitroxide is 2,2,5,5-tetramethylpyrrolidine-N-oxyl.

23. The hemoglobin-based solution of Claims 17, 18, 19, or 20 wherein the nitroxide has the structure




5 where $R_1 - R_4$ are alkyl groups of 1-4 carbon atoms and
A is the remaining members of a 6-membered ring.

24. The hemoglobin-based solution of Claim 23 wherein the nitroxide is 2,2,6,6-tetramethylpiperdine-N-oxyl.

25. A physiologically compatible hemoglobin-based solution
10 comprising

stabilized hemoglobin covalently bound to a nitroxide and forming a conjugate with albumin, dextran, polyoxyethylene, or hydroxyethyl starch.


26. The conjugated hemoglobin-based solution of Claim 25 wherein the nitroxide has the structure



where $R_1 - R_4$ are alkyl groups of 1-4 carbon atoms and A is the remaining members of a 5-membered ring.

5 27. The conjugated hemoglobin-based solution of Claim 26 wherein the nitroxide is 2,2,5,5,-tetramethylpyrrolidine-N-oxyl.

28. The conjugated hemoglobin-based solution of Claim 25 wherein the nitroxide has the structure



10 where $R_1 - R_4$ are alkyl groups of 1-4 carbon atoms and A is the remaining members of a 6-membered ring.

29. The hemoglobin conjugate of Claim 28 wherein the nitroxide is 2,2,6,6-tetramethylpiperdine-N-oxyl.

30. A method to produce a physiologically compatible solution containing stabilized hemoglobin characterized by the steps of:

- 5 covalently binding a nitroxide to stabilized hemoglobin, and
containing the stabilized hemoglobin solution in an environment that is substantially free of oxygen.

31. The method of Claim 30 wherein the nitroxide is covalently bound to a site on the stabilized hemoglobin
10 comprising a sulfhydro group, a β 93 site, or a reactive amino group.

32. A method to produce a physiologically compatible solution containing a biologically compatible macromolecule characterized by:

- 15 providing a physiologically compatible solution containing a biologically compatible macromolecule covalently bound to a nitroxide where the nitroxide is reactive with free radical species in the vascular compartment of an organism.

20 33. The method of Claim 32 wherein the macromolecule is selected from among the group consisting of hemoglobin, albumin, or immunoglobulin.

34. The method of Claim 33 wherein the macromolecule is stabilized hemoglobin.

35. The method of Claim 33 wherein the macromolecule is polymerized hemoglobin.

5 36. A method to produce a liposome-encapsulated hemoglobin solution characterized by

encapsulating a nitroxide and hemoglobin inside a liposome.


37. The method of Claim 36 wherein the nitroxide is
10 covalently bound to the hemoglobin.

38. A container for physiologically compatible hemoglobin solutions characterized by

a nitroxide coated on a surface inside the container.

39. The container of Claim 38 wherein the nitroxide is
15 coated on the surface of a matrix located inside the container.


40. The container of Claims 38 or 39 wherein the nitroxide has the structure



where $R_1 - R_4$ are alkyl groups of 1-4 carbon atoms and A is the remaining members of a 5-membered ring.

5 41. The container of Claim 40 wherein the nitroxide is 2,2,5,5-tetramethylpyrrolidine-N-oxyl.



42. The container of Claim 38 or 39 wherein the nitroxide has the structure






where $R_1 - R_4$ are alkyl groups of 1-4 carbon atoms and
10 A is the remaining members of a 6-membered ring.




43. The container of Claim 42 wherein the nitroxide is 2,2,6,6-tetramethylpiperdine-N-oxyl.



44. A filter having a housing for physiologically compatible hemoglobin-based solutions characterized by
5 a nitroxide bound to a matrix contained within the housing.















FIG. 6






FIG. 8A




FIG. 8B




FIG. 9A







FIG. 9B






Figure 11




FIG. 12

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 94/09246

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C07K14/805 A61K38/42 C07K14/76 C07K16/00 A61K9/127 A61J1/05		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07K A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	FREE RADICAL RESEARCH COMMUNICATIONS, vol.17, no.3, 1992, LONDON, GB pages 157 - 175 R.J. MEHLHORN ET AL. 'NITROXIDE-STIMULATED H2O2 DECOMPOSITION BY PEROXIDASES AND PSEUDOPEROXIDASES.' cited in the application see the whole document ---	1-44
A	NATURE., vol.220, 23 November 1968, LONDON GB pages 787 - 788 H.M. MCCONNELL ET AL. 'SPIN-LABELLED HAEMOGLOBIN AND THE HAEM-HAEM INTERACTION.' see the whole document ---	1,6-10, 13-16, 21-44
	-/--	
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents :		
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed		"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family
Date of the actual completion of the international search	Date of mailing of the international search report	
16 December 1994	28.12.94	
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax (+31-70) 340-3016	Authorized officer Ryckebosch, A	

INTERNATIONAL SEARCH REPORT

Inter. Application No
PCT/US 94/09246

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 290 252 (J.-C. HSIA) 9 November 1988 cited in the application see claims -----	1-5, 11, 12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 94/09246

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0290252	09-11-88	AU-B- 610883	30-05-91
		AU-A- 1563588	10-11-88
		CA-A- 1306583	18-08-92
		DE-A- 3877811	11-03-93
		ES-T- 2053729	01-08-94
		JP-B- 5032372	14-05-93
		JP-A- 63297330	05-12-88
		US-A- 4857636	15-08-89
		US-A- 5189146	23-02-93
		US-A- 5364932	15-11-94
