

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Oversættelse af europæisk patentskrift

(10) **DK/EP 3626725 T3**

-
- (51) Int.Cl.: *C 07 J 9/00 (2006.01)* *A 61 K 31/575 (2006.01)* *A 61 K 35/00 (2006.01)*
A 61 P 1/00 (2006.01) *A 61 P 1/04 (2006.01)* *A 61 P 1/16 (2006.01)*
A 61 P 3/00 (2006.01) *A 61 P 3/04 (2006.01)* *A 61 P 3/06 (2006.01)*
A 61 P 3/10 (2006.01) *A 61 P 5/38 (2006.01)* *A 61 P 5/50 (2006.01)*
A 61 P 9/00 (2006.01) *A 61 P 9/10 (2006.01)* *A 61 P 13/12 (2006.01)*
A 61 P 17/02 (2006.01) *A 61 P 19/02 (2006.01)* *A 61 P 35/00 (2006.01)*
A 61 P 37/02 (2006.01) *A 61 P 43/00 (2006.01)* *A 61 P 21/00 (2006.01)*
A 61 P 25/00 (2006.01) *A 61 P 29/00 (2006.01)* *A 61 P 31/00 (2006.01)*
A 61 P 31/14 (2006.01) *A 61 P 31/18 (2006.01)* *A 61 P 31/20 (2006.01)*
C 07 J 31/00 (2006.01) *C 07 J 41/00 (2006.01)*
- (45) Oversættelsen bekendtgjort den: **2023-02-27**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2022-11-30**
- (86) Europæisk ansøgning nr.: **19204247.1**
- (86) Europæisk indleveringsdag: **2015-05-28**
- (87) Den europæiske ansøgnings publiceringsdag: **2020-03-25**
- (30) Prioritet: **2014-05-29 IT FI20140130**
- (62) Stamansøgningsnr: **15731260.4**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Bar Pharmaceuticals S.r.l., Via Gramsci, 88/A, 42124 Reggio Nell'Emilia (RE), Italien**
- (72) Opfinder: **Angela, ZAMPELLA, Via Luigia Sanfelice, 71, 80127 NAPOLI, Italien**
Stefano, FIORUCCI, Via Assisi, 70, 06100 PERUGIA, Italien
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**
- (54) Benævnelse: **CHOLAN-DERIVATER TIL ANVENDELSE VED BEHANDLINGEN OG/ELLER FOREBYGGELSEN AF FXR- OG TGR5/GPBAR1-MEDIEREDE SYGDOMME**
- (56) Fremdragne publikationer:
CLAUDIO D'AMORE ET AL: "Design, Synthesis, and Biological Evaluation of Potent Dual Agonists of Nuclear and Membrane Bile Acid Receptors", JOURNAL OF MEDICINAL CHEMISTRY, vol. 57, no. 3, 13 February 2014 (2014-02-13), pages 937-954, XP055165457, ISSN: 0022-2623, DOI: 10.1021/jm401873d
SATO HIROYUKI ET AL: "Novel potent and selective bile acid derivatives as TGR5 agonists: biological screening, structure-activity relationships, and molecular modeling studies", JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, US, vol. 51, no. 6, 27 March 2008 (2008-03-27), pages 1831-1841, XP002520339, ISSN: 0022-2623, DOI: 10.1021/JM7015864 [retrieved on 2008-02-29]
YUSUKE IGUCHI ET AL: "Effects of Chemical Modification of Ursodeoxycholic Acid on TGR5 Activation", BIOLOGICAL & PHARMACEUTICAL BULLETIN (OF JAPAN), vol. 34, no. 1, 1 January 2011 (2011-01-01), pages 1-7, XP55594216, JP ISSN: 0918-6158, DOI: 10.1248/bpb.34.1

Fortsættes ...

CARMEN FESTA ET AL: "Exploitation of Cholane Scaffold for the Discovery of Potent and Selective Farnesoid X Receptor (FXR) and G-Protein Coupled Bile Acid Receptor 1 (GP-BAR1) Ligands", JOURNAL OF MEDICINAL CHEMISTRY, vol. 57, no. 20, 23 October 2014 (2014-10-23), pages 8477-8495, XP055165458, ISSN: 0022-2623, DOI: 10.1021/jm501273r

VALENTINA SEPE ET AL: "Modification on Ursodeoxycholic Acid (UDCA) Scaffold. Discovery of Bile Acid Derivatives As Selective Agonists of Cell-Surface G-Protein Coupled Bile Acid Receptor 1 (GP-BAR1)", JOURNAL OF MEDICINAL CHEMISTRY, vol. 57, no. 18, 7 September 2014 (2014-09-07), pages 7687-7701, XP55549277, US ISSN: 0022-2623, DOI: 10.1021/jm500889f

DESCRIPTION

FIELD OF THE INVENTION

[0001] The present invention relates to compounds having cholane scaffolds, said compounds for use in the treatment and/or prevention of FXR and TGR5/GPBAR1 mediated diseases.


STATE OF THE ART

[0002] Bile acids (BAs) are signaling molecules interacting with two type of dedicated cellular receptors, intracellular nuclear receptors and cell-surface receptors. Nuclear receptors include farnesoid X receptor (FXR), identified as the endogenous bile acid sensor (Makishima et al. Science 1999, 284, 1362; Parks et al. Science 1999, 284, 1365).

[0003] Highly expressed in entero-hepatic tissues (liver and intestine), FXR regulates bile acid homeostasis, metabolic pathways also including lipid and glucose homeostasis (Zhang et al. Proc. Natl. Acad. Sci. USA 2006, 103, 1006). Additionally FXR agonists provides anti-inflammatory and anti-fibrotic and anticancer effects (Renga et al. FASEB J. 2012, 26, 3021-3031).

[0004] Bile acid cell-surface receptor (GPBAR1, M-BAR1, GP-BAR1, TGR5) belongs to the rhodopsin-like superfamily of G protein coupled receptors (Takeda et al. FEBS Lett. 2002, 520, 97; Kawamata et al. J. Biol. Chem. 2003, 278, 9435).

[0005] Ligand binding to TGR5/GPBAR1 results in elevation of intracellular cAMP levels with consequently activation of a signaling cascade. GPBAR1 is highly expressed in the liver and in the intestine but also muscles, brain, adipose tissue, macrophages and endothelial cells. In muscle and brown adipose tissue, TGR5/GPBAR1 increases energy expenditure and oxygen consumption (Watanabe et al. Nature 2006, 439, 484) in entero-endocrine L cells, TGR5/GPBAR1 activation stimulates the secretion of glucagon-like peptide (GLP)-1, an incretin that improves pancreas insulin release, thus regulating glucose blood levels, gastrointestinal motility and appetite (Thomas, et al. Cell. Metab. 2009, 10, 167).



[0006] Chemically BAs are truncated cholesterol side chain derivatives. Their molecular repertoire is generated firstly in the liver with the production of primary bile acids, cholic acid (CA) and chenodeoxycholic acid (CDCA). Microbio-transformation in the intestine generates secondary bile acids, deoxycholic acid (DCA) and lithocholic acid (LCA). In human body bile acids are conjugated to glycine and taurine. The activity towards the two BA receptors is structure dependent with CDCA the most potent endogenous FXR activator, and LCA and TLCA the strongest natural agonists of TGR5/GPBAR1.

[0007] Cholestatic pruritus has been noted as a severe side-effect associated with the use of FXR agonists in PBC and a recent study indicated TGR5/GPBAR1 as the molecular target involved in the development of this side effect (Alemi et al. J. Clin. Invest. 2013, 123, 1513-1530).

[0008] WO2013192097 describes 6- α -ethyl-chenodeoxycholic acid (6-ECDCA), a potent and selective FXR agonist endowed with anticholestatic effect. WO2008002573 describes bile acid derivatives as FXR ligands for the prevention or treatment of FXR-mediated diseases or conditions.

[0009] WO2010014836 and Sato H. (J Med Chem. 2008, 51, 4849) describes TGR5 modulators.

[0010] D'Amore C. et al. (J. Med. Chem. 2014, 57, 937) describes Design, synthesis, and biological evaluation of GP-BAR1/FXR dual agonists. D'Amore et al. describes compounds BAR502 and BAR504 as synthesis intermediates.

[0011] Iguchi Y. et al. (J Lipid Res. 2010, 51, 1432) describes bile alcohols function as the ligands of TGR5.

[0012] Compounds BAR107 is disclosed as synthesis intermediate by Kihira K. et al. (Steroids 1992, 57(4), 193-198).

[0013] Swaan P. W. et al. (J. Comp.-Aid. Mol. Des. 1997, 11, 581-588) in a molecular modeling of the intestinal bile acid carrier tested ursocolate (therein compound 15, herein BARn406) among a set of bile acid-conjugates. BARn406 resulted to have an undetectable ability to inhibit taurocholic acid transport in CaCo-2 cells.


[0014] Burns et al. (Steroids 2011, 76(3), 291-300) describes synthesis and olfactory activity of unnatural, sulfated 5-bile acid derivatives in the sea lamprey (*Petromyzon marinus*). Therein disclosed compound 9e (herein compound BAR407) did not to elicit an olfactory response.

[0015] Aim of the present invention is the identification of novel compounds containing the cholane chemical scaffold and that modulate FXR and/or TGR5/GPBAR1.

SUMMARY OF THE INVENTION

[0016] Subject-matter of the present invention is a compound chosen from the following: BAR704, BARn704, BAR710, BARn710 and BAR711, as defined herein.

[0017] These compounds have the following formulae, including inorganic and organic pharmaceutically acceptable salts and solvates thereof:




[0018] Compounds as above described have been found to be FXR or/and TGR5/GPBAR1 modulators and are therefore useful for the treatment of FXR and TGR5/GPBAR1 mediated diseases.

[0019] Therefore for an aspect the present invention relates to the above compounds for use as a medicament.


[0020] For a further aspect the present invention relates to a compound of the invention as above described for use in the prevention and/or treatment of gastrointestinal disorders, liver diseases, cardiovascular diseases, atherosclerosis, metabolic diseases, infectious diseases, cancer, renal disorders, inflammatory disorders, and neurological disorders (such as stroke), said compound of the invention as above described.

DETAILED DESCRIPTION OF THE INVENTION

[0021] According to the invention are provided the compounds BAR704, BARn704, BAR710, BARn710 and BAR711 as defined above. Also provided are the compounds illustrated below, wherein of these, compound BAR712 is not a compound according to the invention.




[0022] Provided herein is also the following compound, which does not form part of the present invention:




BAR802

[0023] Provided herein are also the following compounds, which do not form part of the present invention:



[0024] Provided herein are also the following compounds. Compounds BAR704, BARn704, BAR710, BARn710 and BAR711 are part of the present invention, whereas compounds BAR712 and BAR802 are not part of the present invention:



[0025] The compounds according to the invention, have been found to be highly selective FXR or TGR5/GPBAR1 modulators or dual FXR and TGR5/GPBAR1 modulators and are therefore useful as medicaments in particular for use in the prevention and/or treatment of gastrointestinal disorders, liver diseases, cardiovascular diseases, atherosclerosis, metabolic diseases, metabolic disorders, infectious diseases, cancer, renal disorders, inflammatory disorders, and neurological disorders such as stroke.

[0026] In certain embodiments the liver disease is selected in the group consisting of chronic liver diseases including primary biliary cirrhosis (PBC), cerebrotendinous xanthomatosis (CTX), primary sclerosing cholangitis (PSC), drug induced cholestasis, intrahepatic cholestasis of pregnancy, parenteral nutrition associated cholestasis, bacterial overgrowth and sepsis associated cholestasis, autoimmune hepatitis, chronic viral hepatitis, alcoholic liver disease, nonalcoholic fatty liver disease (NAFLD), nonalcoholic steatohepatitis (NASH), liver transplant associated graft versus host disease, living donor transplant, liver regeneration, congenital hepatic fibrosis, granulomatous liver disease, intra- or extrahepatic malignancy, Wilson's

disease, hemochromatosis, and alpha 1-antitrypsin deficiency.

[0027] In certain embodiments the gastrointestinal disease is selected in the group consisting of inflammatory bowel disease (IBD) (including Crohn's disease, ulcerative colitis and undetermined colitis), irritable bowel syndrome (IBS), bacterial overgrowth, acute and chronic pancreatitis, malabsorption, post-radiation colitis, and microscopic colitis.

[0028] In certain embodiments the renal disease is selected in the group consisting of diabetic nephropathy, hypertensive nephropathy, chronic glomerular disease, including chronic glomerulonephritis and chronic transplant glomerulopathy, chronic tubulointerstitial diseases and vascular disorders of the kidney.

[0029] In certain embodiments the cardiovascular disease is selected in the group consisting of atherosclerosis, dyslipidemia, hypercholesterolemia, hypertriglyceridemia, hypertension also known as arterial hypertension, inflammatory heart disease including myocarditis and endocarditis, ischemic heart disease stable angina, unstable angina, myocardial infarction, cerebrovascular disease including ischemic stroke, pulmonary heart disease including pulmonary hypertension, peripheral artery disease (PAD), also known as peripheral vascular disease (PVD) peripheral artery occlusive disease, and peripheral obliterative arteriopathy.

[0030] In certain embodiments the metabolic disease is selected in the group consisting of insulin resistance, metabolic syndrome, Type I and Type II diabetes, hypoglycemia, disorders of adrenal cortex including adrenal cortex insufficiency.

[0031] In certain embodiments metabolic disorder is selected in the group consisting of obesity and conditions associated to bariatric surgery.

[0032] In certain embodiments cancer is selected in the group of liver cancer, bile duct cancers, pancreatic cancer, gastric cancer, colon-rectal cancer, breast cancer, ovary cancer and condition associated with chemotherapy resistance.

[0033] In certain embodiments infectious disorder is selected in the group of human immunodeficiency associated disease (AIDS) and related disorders, virus B and Virus C infection.

[0034] In certain embodiments inflammatory disorder is selected in the group of rheumatoid arthritis, fibromyalgia, Sjogren's syndrome, scleroderma, Behcet's syndrome, vasculitis and systemic lupus erythematosus.

[0035] The data on the activity of certain compounds of the invention and other compounds disclosed herein on FXR and TGR5/GPBAR1 are described in the following table. In this table, activities for compounds of the invention and other compounds disclosed herein on FXR and GPBAR1 was compared to those of reference compounds: i.e. CDCA for FXR and TLCA for TGR5/GPBAR1. Each compound was tested at the concentration of 10 microM and

transactivation activity of CDCA on FXR and TLCA on CRE (i.e. TGR5/GPBAR1) was considered equal to 100%.

Table 1

Compounds of formula (I)	FXR	GPBAR1
	(% of activity in comparison to 10 μ M CDCA)	(% of activity in comparison to 10 μ M TLCA)
BAR106	0	19.0 $\pm$ 1.3
BAR107	1.8 $\pm$ 0.3	9.9 $\pm$ 1.7
BAR305	0	23.9 $\pm$ 4.0
BAR304	0	55.1 $\pm$ 12.5
BAR402	336.4 $\pm$ 20.8	28.0 $\pm$ 1.8
BARn406	27.0 $\pm$ 1.5	4.8 $\pm$ 1.8
BAR407	177.1 $\pm$ 3.5	37.3 $\pm$ 2.5
BAR501	9.9 $\pm$ 0.1	64.5 $\pm$ 0.5
BAR501-6a	15.4 $\pm$ 1.2	46.6 $\pm$ 6.7
BARn501	8.5 $\pm$ 1.4	83.1 $\pm$ 7.4
BAR502	263.0 $\pm$ 32.0	74.5 $\pm$ 6.4
BAR503	68.8 $\pm$ 26.6	59.8 $\pm$ 0.1
BAR504	488.5 $\pm$ 17.5	103.0 $\pm$ 12.1
BAR504-6b	32.4 $\pm$ 14.1	75.3 $\pm$ 3.4
BAR506	411.5 $\pm$ 36.5	80.9 $\pm$ 9.5
BAR701	101.3 $\pm$ 10.1	50.2 $\pm$ 2.3
BARn701	90 $\pm$ 22	55 $\pm$ 1.0
BAR701 solf	229 $\pm$ 2.0	63.5 $\pm$ 0.5
BAR702	11.7 $\pm$ 0.8	35.8 $\pm$ 1.2
BAR703	8.6 $\pm$ 0.8	47.7 $\pm$ 1.0
BAR704	220.5 $\pm$ 37.5	27.9 $\pm$ 6.3
BARn704	202.5 $\pm$ 1.5	64 $\pm$ 2.0
BAR705	43.7 $\pm$ 3.9	67.0 $\pm$ 9.9
BAR706	153 $\pm$ 9.0	66.05 $\pm$ 7.6
BARn706	197.5 $\pm$ 4.5	69.5 $\pm$ 6.5
BARn706solf	120 $\pm$ 9.5	71.5 $\pm$ 0.5
BAR707	92.5 $\pm$ 7.5	65 $\pm$ 6.0
BAR708	6.4 $\pm$ 0.8	57.6 $\pm$ 4.6
BAR709	33.7 $\pm$ 2.0	54.2 $\pm$ 0.3
BAR710	179 $\pm$ 43	49.5 $\pm$ 0.5

Compounds of formula (I)	FXR	GPBAR1
	(% of activity in comparison to 10 μ M CDCA)	(% of activity in comparison to 10 μ M TLCA)
BARn710	142 $\pm$ 1.0	75 $\pm$ 8.0
BAR712	33.0 $\pm$ 0.15	49.7 $\pm$ 1.2
BAR802	11.5 $\pm$ 8.5	56 $\pm$ 5.0
BAR803	122.5 $\pm$ 5.5	80.5 $\pm$ 5.0
BAR804	196.5 $\pm$ 6.5	52.5 $\pm$ 3.5

[0036] For one aspect, the present invention relates to compounds of the invention wherein the compounds are FXR and TGR5/GPBAR1 dual agonists. A compound, not according to the present invention, but which has such dual agonist activity is BAR502. Surprisingly, BAR502 does not induce itching when administered to animals rendered cholestatic by administration of ANIT or Estrogen. In cholestatic syndromes, body accumulation of bile acids is thought to cause itching. Recently, TGR5/GPBAR1 shown to mediate itching caused by intradermal administration of DCA and LCA (Alemi et al. J. Clin. Invest. 2013, 123, 1513-1530). In clinical trials, administration of patients suffering from primary biliary cirrhosis (PBC) with obeticholic acid has resulted in severe itching in approximately 80% of patients. One specific and surprising advantage of BAR502 is that this agent do not induce itching when administered to animals rendered cholestatic by administration of α -naphthyl-isothiocyanate (ANIT) or 17 α -ethynylestradiol (two validated model of cholestasis). In these experimental setting BAR502 administration increases survival, attenuates serum alkaline phosphatase levels and robustly modulates the liver expression of canonical FXR target genes including OSTa, BSEP, SHP and MDR1, without inducing pruritus. In the 17 α -ethynylestradiol model, BAR502 attenuates cholestasis and reshapes bile acid pool without inducing itching, demonstrating that in models of non-obstructive cholestasis, BAR502 attenuates liver injury without causing itching.

[0037] In one aspect, the present invention relates to compounds of the invention wherein the compounds are high selective FXR agonists without effects on GPBAR1 when administered alone but effective in inhibiting GPBAR1 activation caused by TLCA (10 μ M)), thus behaving as GPBAR1 antagonists. In one aspect, the present invention relates to compounds of the invention wherein the compounds are high selective GPBAR1 agonists without effects on FXR. In one aspect, the present invention relates to compounds of the invention wherein the compounds are high selective GPBAR1 antagonists without effects on FXR.


[0038] The present invention could be better understood in light of the examples and experimental section below. The following examples are examples of the present invention only in as far as they relate to the synthesis and testing of the compounds BAR704, BARn704, BAR710, BARn710 and BAR711. The synthesis and testing of other compounds in the following examples is only for the purposes of comparison.

EXPERIMENTAL SECTION

Chemistry

EXAMPLE 1. Preparation of compounds of formula (I) wherein $R_2=H$ EXAMPLE 1A. Synthesis of bis-*homoursodeoxycholate* derivatives

[0039] A four-steps reaction sequence on 1, including protection of alcoholic functions at C3 and C7, reduction of the side chain methyl ester, and subsequent one pot Swern oxidation/Wittig C2 homologation gave the protected methyl ester of $\Delta^{24,25}$ -bis-*homo*UDCA. Side chain double bond hydrogenation and alcoholic function deprotection gave bis-*homo*UDCA methyl ester 4, that was used as starting material in the preparation of BAR305 and its corresponding alcohol, BAR304, through treatment with LiOH and LiBH₄, respectively.



a) 2,6-lutidine, *t*-butyldimethylsilyl trifluoromethanesulfonate, CH₂Cl₂, 0 °C; b) LiBH₄, MeOH dry, THF, 0 °C, quantitative yield over two steps; c) DMSO, oxalyl chloride, TEA dry, CH₂Cl₂, -78 °C then methyl(triphenylphosphoranylidene)acetate, 76%; d) H₂, Pd(OH)₂/C Degussa type, THF/MeOH 1:1, quantitative yield; e) HCl 37%, MeOH, quantitative yield; f) NaOH 5% in MeOH/H₂O 1:1 v/v, 60%; g) LiBH₄, MeOH dry, THF, 0 °C, 77%.

Step a,b) Preparation of 3 α , 7 β -di(*tert*-butyldimethylsilyloxy)-5 β -cholan-24-ol (2)

[0040] Compound 1 (1.2 g, 3 mmol) was protected at the two alcoholic function following the same synthetic procedure described in J. Med. Chem. 2014, 57, 937 to obtain 1.9 g of methyl 3 α , 7 β -di(*tert*-butyldimethylsilyloxy)-5 β -cholan-24-oate (quantitative yield) in the form of colorless needles, that was subjected to next step without any purification.

[0041] Methanol (850 μ L, 21 mmol) and LiBH₄ (10.5 mL, 2M in THF, 21 mmol) were added to a solution of methyl ester (1.9 g, 3 mmol) in dry THF (30 mL) at 0 °C following the same synthetic procedure described in J. Med. Chem. 2014, 57, 937. Purification by silica gel (hexane/ethyl acetate 99:1 and 0.5% TEA) gave 2 as a white solid (1.8 g, quantitative yield).

Step c) One pot preparation of methyl 3 α , 7 β -di(*tert*-butyldimethylsilyloxy)-25, 26-

bishomo-5 β -chol-24-en-26-oate (3).

[0042] DMSO (2.1 mL, 30 mmol) was added dropwise for 15 min to a solution of oxalyl chloride (7.5 mL, 15 mmol) in dry dichloromethane (30 mL) at -78 °C under argon atmosphere. After 30 min a solution of 2 (1.8 g, 3 mmol) in dry CH₂Cl₂ was added via cannula and the mixture was stirred at -78 °C for 30 min. Et₃N (2.5 mL, 18 mmol) was added dropwise. After 1 h methyl(triphenylphosphoranylidene)acetate (2.0 g, 6 mmol) was added and the mixture was allowed to warm to room temperature. NaCl saturated solution was added and the aqueous phase was extracted with diethyl ether (3×100 mL). The combined organic phases were washed with water, dried (Na₂SO₄) and concentrated. Purification by silica gel (hexane-ethyl acetate 95:5 and 0.5% TEA) gave compound 3 as a colorless oil (1.5 g, 76%).

[0043] Step d) Preparation of methyl 3 α , 7 β -di(tert-butyldimethylsilyloxy)-25, 26-bishomo-5 β -cholan-26-oate. A solution of compound 3 (1.5 g, 2.3 mmol) in THF dry/MeOH dry (25 mL/25 mL, v/v) was hydrogenated in presence of Pd(OH)₂ 5% wt on activated carbon Degussa type (20 mg) following the same synthetic procedure described in J. Med. Chem. 2014, 57, 937 affording methyl 3 α , 7 β -di(tert-butyldimethylsilyloxy)-25, 26-bishomo-5 β -cholan-26-oate (1.5 g, quantitative yield) that was subjected to step e) without purification.

[0044] Step e) Preparation of methyl 3 α , 7 β -dihydroxy-25, 26-bishomo-5 β -cholan-26-oate (4)

[0045] Methyl 3 α , 7 β -di(tert-butyldimethylsilyloxy)-25, 26-bishomo-5 β -cholan-26-oate (1.5 g) was dissolved in methanol (70 mL). At the solution HCl (2 mL, 37% v/v) was added following the same synthetic procedure described in J. Med. Chem. 2014, 57, 937 affording 4 as colorless amorphous solid (1.0 g, quantitative yield).

[0046] Step f) Preparation of 3 α , 7 β -dihydroxy-25, 26-bishomo-5 β -cholan-26-oic acid (BAR305). A portion of compound 4 (430 mg, 1 mmol) was hydrolyzed with NaOH (400 mg, 10 mmol) in a solution of MeOH: H₂O 1:1 v/v (20 mL) for 4 h at reflux. An analytic sample was purified by HPLC on a Nucleodur 100-5 C18 (5 μ m; 4.6 mm i.d. × 250 mm) with MeOH/H₂O (95:5) as eluent (flow rate 1 mL/min) (t_R=5 min). BAR305: C₂₆H₄₄O₄

[0047] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 3.47 (2H, m, H-3 and H-7), 2.27 (2H, t, *J* = 7.2 Hz, H₂-25), 0.96 (3H, s, H₃-19), 0.94 (3H, d, *J* = 6.5 Hz, H₃-21), 0.70 (3H, s, H₃-18).

[0048] The ¹³C NMR was recorded on Varian Inova 100 MHz, using CD₃OD as solvent: δ 178.2, 72.1, 71.9, 57.6, 56.7, 44.8, 44.5, 44.0, 41.6, 40.7, 38.6, 38.0, 36.9, 36.8, 36.1, 35.3, 35.2, 30.9, 29.8, 27.9, 26.7, 26.6, 23.9, 22.4, 19.3, 12.7.

[0049] Step g) 25, 26-bishomo-5 β -cholan-3 α , 7 β , 26-triol (BAR304). Compound 4 (500 mg, 1.2 mmol) was reduced in the same operative condition described in step b). Purification by silica gel (CH₂Cl₂/methanol 9:1) gave BAR304 as a colorless oil (375 mg, 77%). An analytic sample was purified by HPLC on a Nucleodur 100-5 C18 (5 μ m; 4.6 mm i.d. $\times$ 250 mm) with MeOH/H₂O (85:15) as eluent (flow rate 1 mL/min) (t_R=9 min).


[0050] BAR 304: C₂₆H₄₆O₃

[0051] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 3.53 (2H, t, J = 6.5 Hz, H₂-26), 3.48 (2H, m, H-3 and H-7), 0.95 (3H, s, H₃-19), 0.93 (3H, d, J = 6.5 Hz, H₃-21), 0.70 (3H, s, H₃-18).

[0052] The ¹³C NMR was recorded on Varian Inova 100 MHz, using CD₃OD as solvent: δ 72.1, 71.9, 63.0, 57.5, 56.7, 44.7, 44.4, 44.0, 41.6, 40.7, 38.5, 37.9, 37.2, 37.0, 36.1, 35.2, 33.7, 30.9, 29.8, 27.9, 27.4, 27.1, 23.9, 22.4, 19.4, 12.7.

EXAMPLE 1B. Synthesis of 3 α ,7 β -dihydroxy-24-nor-5 β -cholan-23-yl-23-sodium sulfate (BAR106) and 3 α ,7 β -dihydroxy-24-nor-5 β -cholan-23-ol (BAR107)

[0053] BAR106 was prepared starting from UDCA by a reaction sequence comprising permylation at the hydroxyl groups, Beckmann one carbon degradation at C24 and transformation of the C23 carboxyl group into the corresponding methyl ester intermediate. Protection at the hydroxyl groups at C-3 and C-7 as silyl ethers, reduction at C23 methyl ester, sulfation at C23 primary alcoholic function and finally deprotection furnished crude BAR106 as ammonium salt. Purification on Amberlite and then by HPLC gave title BAR106 as sodium salt.



a) HCOOH, HClO₄, 96%; b) TFA, trifluoroacetic anhydride, NaNO₂, 96%; c) KOH 30% in MeOH/H₂O 1:1 v/v, 97%; d) p-TsOH, MeOH dry, 98%; e) 2,6-lutidine, t-butyldimethylsilyl trifluoromethanesulfonate, CH₂Cl₂, 0 °C, 88%; f) LiBH₄, MeOH dry, THF, 0 °C, quantitative yield; g) Et₃N.SO₃, DMF, 95 °C; h) HCl 37%, MeOH, then Amberlite CG-120, MeOH, 86% over two steps; i) LiBH₄, MeOH dry, THF, 0 °C, quantitative yield.

[0054] Steps a,d) Preparation of methyl 3 α ,7 β -dihydroxy-24-nor-5 β -cholan-23-oate (6).

Ursodeoxycholic acid (2.0 g, 5.1 mmol) was transformed in methyl 3 α ,7 β -dihydroxy-24-*nor*-5 β -cholan-23-oate (6, 1.6 g, 87%) following the same synthetic procedure described in J. Med. Chem. 2014, 57, 937.

Step e) Preparation of methyl 3 α , 7 β -di(*tert*-butyldimethylsilyloxy)-5 β -cholan-24-oate

[0055] Compound 6 (1.2 g, 3.0 mmol) was protected at the hydroxyl groups in the same operative condition described in example 1A step a). Purification by flash chromatography on silica gel using hexane/ethyl acetate 9:1 and 0.5% of triethylamine as eluent, gave protected methyl ester (1.6 g, 88%).

Step f) Preparation of 3 α , 7 β -di(*tert*-butyldimethylsilyloxy)-5 β -cholan-24-ol (7)

[0056] Side chain methyl ester (818 mg, 1.3 mmol) was reduced in the same operative condition described in example 1A step b). Purification by flash chromatography on silica gel using hexane/ethyl acetate 98:2 and 0.5% of triethylamine as eluent, gave 7 (770 mg, quantitative yield).

Steps g, h) Preparation of 3 α ,7 β -dihydroxy-24-*nor*-5 β -cholan-23-yl-23-sodium sulfate (BAR106)

[0057] The triethylamine-sulfur trioxide complex (2.0 g, 11 mmol) was added to a solution of 7 (660 mg, 1.1 mmol) in DMF dry (25 mL) following the same synthetic procedure described in J. Med. Chem. 2014, 57, 937. HPLC on a Nucleodur 100-5 C18 (5 μ m; 10 mm i.d. $\times$ 250 mm) with MeOH/H₂O (65:35) as eluent (flow rate 3 mL/min), gave 442 mg (86% over two steps) of BAR106 (t_R =8.4 min).

[0058] BAR 106: C₂₃H₃₉NaO₆S

[0059] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 4.04 (2H, m, H₂-23), 3.48 (2H, m, H-3 and H-7), 1.00 (3H, d, J = 6.5 Hz, H₃-21), 0.97 (3H, s, H₃-19), 0.72 (3H, s, H₃-18).

Step i) Preparation of 3 α , 7 β -dihydroxy-24-*nor*-5 β -cholan-23-ol (BAR107) Compound 6 was transformed in BAR107 in the same operative condition described in step f.


[0060] BAR107: C₂₃H₄₀O₃

[0061] The ^1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 3.60 (1H, m, H-7), 3.51 (1H, m, H-3), 3.50 (2H, m, H_2 -23), 0.97 (3H, d, ovl, H_3 -21), 0.96 (3H, s, H_3 -19), 0.72 (3H, s, H_3 -18).

[0062] The ^{13}C NMR was recorded on Varian Inova 100 MHz, using CD_3OD as solvent: δ 72.1, 71.9, 60.8, 57.5, 57.1, 44.8, 44.5, 44.0, 41.6, 40.7, 39.9, 38.6, 38.0, 36.1, 35.2, 34.1, 31.0, 29.8, 27.9, 23.9, 22.4, 19.5, 12.6;

Example 1C. Synthesis of 7 β -hydroxy-5 β -cholan-24-yl-24-sodium sulfate (BAR402)

[0063] Tosylation and elimination at C-3 hydroxyl group on methyl ester 8 followed by double bond reduction, subsequent LiBH_4 treatment and regioselective sulfation at C-24 primary hydroxyl group gave BAR402.



a) p-TsCl, pyridine, quantitative yield; b) LiBr, Li_2CO_3 , DMF, reflux, c) H_2 , $\text{Pd}(\text{OH})_2$, THF/MeOH 1:1, room temperature, quantitative yield; d) LiBH_4 , MeOH dry, THF, 0 °C, 79%; e) $\text{Et}_3\text{N} \cdot \text{SO}_3$, DMF, 95 °C.

Steps a-c) Preparation of methyl 7-keto-5 β -cholan-24-oate (9)

[0064] To a solution of 8 (965 mg, 2.5 mmol) in dry pyridine (100 mL), tosyl chloride (4.7 g, 25.0 mmol) was added, and the mixture was stirred at room temperature for 4 h. It was poured into cold water (150 mL) and extracted with CH_2Cl_2 (3 $\times$ 150 mL). The combined organic layer was washed with saturated NaHCO_3 solution (150 mL), and water (150 mL), and then dried over anhydrous MgSO_4 and evaporated in vacuo to give 1.4 g of methyl 3 α -tosyloxy-7-keto-5 β -cholan-24-oate (quantitative yield). Lithium bromide (434 mg, 5.0 mmol) and lithium carbonate (370 mg, 5.0 mmol) were added to a solution of 3 α -tosyloxy-7-keto-5 β -cholan-24-oate (1.4 g, 2.5 mmol) in dry DMF (30 mL), and the mixture was refluxed for 2 h. After cooling to room temperature, the mixture was slowly poured into 10% HCl solution (20 mL) and extracted with CH_2Cl_2 (3 $\times$ 50 mL). The combined organic layer was washed successively with water, saturated NaHCO_3 solution and water, and then dried over anhydrous MgSO_4 and evaporated to dryness to give 965 mg of oleos residue (quantitative yield), that was subjected to next step without any purification. Hydrogenation on $\text{Pd}(\text{OH})_2$ in the same operative condition described in example 1A, step d furnished 975 mg of 9 (quantitative yield), that was subjected to next step without any purification.

Step d) Preparation of 5 β -cholan-7 α ,24-diol

[0065] LiBH₄ treatment on compound 9 in the same operative condition described in example 1A step b and purification by silica gel (ethyl acetate-hexane, 85:15) gave 5 β -cholan-7 α ,24-diol as a white solid (714 mg, 79%).

Step e) Preparation of 7 α -hydroxy-5 β -cholan-24-yl-24-sodium sulfate (BAR402).

[0066] Sulfation on C24 was performed in the same operative conditions described in example 1B step g) to give crude BAR402 as ammonium salt. RP18/HPLC on a Nucleodur 100-5 C18 (5 μ m; 10 mm i.d. $\times$ 250 mm) with MeOH/H₂O (90:10) as eluent (flow rate 3 mL/min) afforded BAR402 (t_R = 6.6 min) as sodium salt.


[0067] BAR402: C₂₄H₄₁NaO₅S

[0068] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 3.96 (2H, t, J = 6.6 Hz, H₂-24), 3.78 (1H, br s, H-7), 0.96 (3H, d, J = 6.5 Hz, H₃-21), 0.92 (3H, s, H₃-19), 0.69 (3H, s, H₃-18);

[0069] The ¹³C NMR was recorded on Varian Inova 100 MHz, using CD₃OD as solvent: δ 69.7, 69.4, 57.7, 51.6, 45.0, 43.8, 41.2, 41.0, 39.0, 37.1, 37.0, 36.3, 34.2, 33.3, 31.7, 29.5, 29.0, 27.3, 24.8, 24.3, 22.7, 21.9, 19.2, 12.3.

EXAMPLE 2. Preparation of compounds of formula (I) wherein R₂=Et or =CH-CH₃**EXAMPLE 2A. Synthesis of 6 β -ethyl-3 α ,7 β -dihydroxy-5 β -cholan-24-ol (BAR501)**

[0070] Methyl ester formation and acetylation at C-3 hydroxyl group on 7-KLCA furnished intermediate **10** in 84% yield over two steps. Aldolic addition to a silyl enol ether intermediate generated **11** that was hydrogenated at the exocyclic double bond (H₂ on Pd(OH)₂) affording **12** in 80% yield over three steps. NaBH₄ treatment in methanol followed by LiBH₄ reduction on the crude reaction product afforded a mixture whose HPLC purification (88% MeOH:H₂O) gave pure BAR501 in a 79% yield respect to its C7 epimer, BAR504-6b.



a) *p*-TsOH, MeOH dry; b) acetic anhydride, pyridine, 84% yield over two steps; c) DIPA, *n*-BuLi, TMSCl, TEA dry, THF dry -78 °C; d) acetaldehyde, BF₃(OEt)₂, CH₂Cl₂, -60 °C, 80% over two steps; e) H₂, Pd(OH)₂, THF/MeOH 1:1, quantitative yield; f) NaBH₄, MeOH; g) LiBH₄, MeOH dry, THF, 0 °C, 79% over two steps.

Steps a-d). Preparation of methyl 3 α -acetoxy-6-ethylidene-7-keto-5 β -cholan-24-oate (11)

[0071] To a solution of 7-ketolithocholic acid (5 g, 12.8 mmol), dissolved in 100 mL of dry methanol was added *p*-toluenesulfonic acid (11 g, 64.1 mmol). The solution was left to stand at room temperature for 2 h. The mixture was quenched by addition of NaHCO₃ saturated solution. After the evaporation of the methanol, the residue was extracted with EtOAc (3×150 mL). The combined extract was washed with brine, dried with Na₂SO₄, and evaporated to give the methyl ester as amorphous solid (5.13 g, quantitative yield).

[0072] At the solution of the methyl ester (5.13 g, 12.7 mmol) in dry pyridine (100 mL), an excess of acetic anhydride (8.4 mL, 89 mmol) was added. When the reaction was complete, the pyridine was concentrated under vacuum. The residue was poured into cold water (100 mL) and extracted with AcOEt (3×150 mL). The combined organic phases were dried (Na₂SO₄) and concentrated to give a residue that was further purified by flash chromatography on silica gel using hexane/ethyl acetate 8:2 and 0.5% of triethylamine as eluent (4.8 g of **10** as a white solid, 84% yield over two steps).

[0073] To a solution of diisopropylamine (23 mL, 0.16 mol) in dry THF (50 mL) was added dropwise a solution of *n*-butyllithium (60 mL, 2.5 M in hexane, 0.15 mol) at -78 °C. After 30 min, trimethylchlorosilane (27.1 mL, 0.21 mol) was added. After additional 30 min, a solution of compound **10** (4.8 g, 10.7 mmol) in dry THF (70 mL) was added. The reaction was stirred at -78 °C for an additional 45 min and then triethylamine (54 mL, 0.38 mol) was added. After 1 h, the reaction mixture was allowed to warm to -20 °C, treated with aqueous saturated solution of NaHCO₃ (100 mL) and brought up to room temperature in 2 h. The aqueous phase was extracted with ethyl acetate (3×50 mL). The combined organic phases were washed then with saturated solution of NaHCO₃, water and brine. After drying over anhydrous Na₂SO₄, the residue was evaporated under vacuum to give 6 g of yellow residue, that was diluted in dry CH₂Cl₂ (50 mL) and cooled at -78 °C. At this stirred solution acetaldehyde (3 mL, 53 mmol) and BF₃·OEt₂ (13.5 mL, 0.107 mol) were added dropwise. The reaction mixture was stirred for 2 h at -60 °C and allowed to warm to room temperature. The mixture was quenched with saturated aqueous solution of NaHCO₃ and extracted with CH₂Cl₂. The combined organic phases were washed with brine, dried over anhydrous Na₂SO₄ and concentrated under vacuum.

[0074] Purification by silica gel (hexane-ethyl acetate 9:1 and 0.5% TEA) gave compound **11**

(4.1 g, 80%). NMR analysis demonstrated a diastomeric ratio E/Z >95%. The *E* configuration at the exocyclic double bond was established by dipolar coupling H₃-26 (δ 1.67)/H-5 (δ 2.62) in Noesy spectrum (400 MHz, mixing time 400 ms).

(*E*)-3 α -acetoxy-6-ethylidene-7-keto-5 β -cholan-24-oate (11): C₂₉H₄₄O₅

[0075] The ¹H NMR was recorded on Varian Inova 400 MHz, using CDCl₃ as solvent: δ 6.16 (1H, q, *J* = 7.0 Hz, H-25), 4.74 (1H, m, H-3), 3.64 (3H, s, COOCH₃), 2.62 (1H, dd, *J* = 13.0, 3.6 Hz, H-5), 1.98 (3H, s, COCH₃), 1.67 (3H, d, *J* = 7.0 Hz, H₃-26), 1.00 (3H, s, H₃-19), 0.92 (3H, d, *J* = 6.0 Hz, H₃-21), 0.67 (3H, s, H₃-18).

[0076] The ¹³C NMR was recorded on Varian Inova 100 MHz, using CDCl₃ as solvent: δ 204.5, 174.6, 170.7, 143.1, 130.2, 72.5, 54.5, 51.4, 50.7, 48.6, 45.2, 43.5, 39.1, 38.9, 35.1, 34.9, 34.1, 33.4, 31.0, 30.9, 28.4, 25.9 (2C), 22.8, 21.4, 21.2, 18.4, 12.7, 12.2.

Steps e) Preparation of methyl 3 α -acetoxy-6 β -ethyl-7-keto-5 β -cholan-24-oate (12).

[0077] A solution of **11** (4.0 g, 8.5 mmol) in THF dry/MeOH dry (100 mL, 1:1 v/v) was hydrogenated in presence of Pd(OH)₂ 20% wt on activated carbon (100 mg) degussa type. The mixture was transferred to a standard PARR apparatus and flushed with nitrogen and then with hydrogen several times. The apparatus was shacked under 344,738 Pa (50 psi) of H₂. The reaction was stirred at room temperature for 8 h.

[0078] The catalyst was filtered through Celite, and the recovered filtrate was concentrated under vacuum to give **12** (4.0 g, quantitative yield).

Methyl 3 α -acetoxy-6 β -ethyl-7-keto-5 β -cholan-24-oate (12): C₂₉H₄₆O₅

[0079] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 4.65 (1H, m, H-3), 3.66 (3H, s, COOCH₃), 2.56 (1H, t, *J* = 11.5 Hz, H-8), 2.35 (1H, m, H-23a), 2.22 (1H, m, H-23b), 1.99 (3H, s, COCH₃), 1.22 (3H, s, H₃-19), 0.92 (3H, d, *J* = 6.3 Hz, H₃-21), 0.83 (3H, t, *J* = 7.2 Hz, H₃-26), 0.67 (3H, s, H₃-18).

[0080] The ¹³C NMR was recorded on Varian Inova 100 MHz, using CD₃OD as solvent: δ 214.7, 174.3, 170.2, 72.6, 61.7, 54.8, 51.3, 49.0, 48.5, 45.3, 42.7, 42.3 (2C), 38.6, 35.4, 35.1, 35.0, 31.0, 30.8, 28.0 (2C), 26.4, 25.7, 24.7, 21.3, 21.1, 18.2, 12.9, 11.9. The β configuration of ethyl group at C-6 was determined by dipolar couplings H₃-26 (δ 0.83)/ H₃-19 (δ 1.22) and

H-8 (δ 2.56)/H-25 (δ 1.83) in Noesy spectrum (400 MHz, mixing time 400 ms).

Steps f,g) Preparation of 6 β -ethyl-3 α ,7 β -dihydroxy-5 β -cholan-24-ol (BAR501).

[0081] To a methanol solution of compound **12** (1.18 g, 2.5 mmol), a large excess of NaBH₄ was added at 0 °C. The mixture was left at room temperature for 2 h and then water and MeOH were added dropwise during a period of 15 min at 0 °C with effervescence being observed. After evaporation of the solvents, the residue was diluted with water and extracted with AcOEt (3×50 mL). The combined extract was washed with brine, dried with Na₂SO₄, and evaporated to give 1.3 g of a crude residue that was subjected to the next step without further purification. The crude residue was treated with LiBH₄ (2 M in THF) in the same operative condition described in example 1A step b). HPLC purification on a Nucleodur 100-5 C18 (5 μ m; 10 mm i.d. × 250 mm) with MeOH/H₂O (88:12) as eluent (flow rate 3 mL/min), gave 802 mg of BAR501 (79%, t_R = 11 min).

[0082] Alternatively step f was performed with Ca(BH₄)₂, produced in situ.

[0083] To a solution of compound **12** (500 mg, 1.05 mmol) and absolute ethanol (4 mL), at 0 °C, CaCl₂ (466 mg, 4.2 mmol) was added. At the same solution was added a solution of NaBH₄ (159 mg, 4.2 mmol) in absolute ethanol (4 mL). After 4 h at -5 °C, MeOH was added dropwise. Then after evaporation of the solvents, the residue was diluted with water and extracted with AcOEt (3×50 mL). The combined extract was washed with brine, dried with Na₂SO₄, and evaporated to give 500 mg of a crude residue that was subjected to the step g without further purification.

[0084] BAR501: C₂₆H₄₆O₃

[0085] The ¹H NMR was recorded on Varian Inova 700 MHz, using CD₃OD as solvent: δ 3.74 (1H, dd, J = 10.3, 6.0 Hz, H-7), 3.51 (1H, ovl, H-3), 3.49 (2H, ovl, H₂-24), 1.00 (3H, s, H₃-19), 0.97 (3H, d, J = 6.5 Hz, H₃-21), 0.96 (3H, t, J = 7.6 Hz, H₃-26), 0.72 (3H, s, H₃-18).

[0086] The ¹³C NMR was recorded on Varian Inova 175 MHz, using CD₃OD as solvent: δ 75.3, 71.9, 63.6, 57.5, 56.5, 51.6, 45.7, 44.9, 42.1, 41.5, 40.4, 40.3, 37.1, 35.8, 32.4, 30.7, 30.3, 29.7, 29.6, 28.3, 26.2, 23.4, 22.1, 19.4, 14.8, 12.7.

Example 2B. Preparation of 6 β -ethyl-3 α ,7 α -dihydroxy-5 β -cholan-24-ol (BAR504-6b)


[0087] BAR504-6b was prepared as described in the Example 2A (t_R = 20.4 min). **BAR504-6b:** C₂₆H₄₆O₃

[0088] The ^1H NMR was recorded on Varian Inova 700 MHz, using CD_3OD as solvent: δ 3.60 (1H, s, H-7), 3.51 (2H, m, H₂-24), 3.35 (1H, ovl, H-3), 2.30 (1H, q, J = 13.5 Hz, H-4a), 0.97 (3H, d, J = 6.8 Hz, H₃-21), 0.95 (3H, t, J = 7.3 Hz, H₃-26), 0.94 (3H, s, H₃-19), 0.70 (3H, s, H₃-18).

[0089] The ^{13}C NMR was recorded on Varian Inova 175 MHz, using CD_3OD as solvent: δ 71.9, 71.8, 62.7, 56.8, 51.7, 50.5, 46.7, 42.5, 41.4, 40.1, 36.6, 36.4, 36.2, 36.0, 33.2, 32.4, 30.1, 29.5, 28.8, 28.5, 25.3, 23.9, 20.7, 18.4, 13.7, 11.4.

EXAMPLE 2C. Synthesis of 6 α -ethyl-3 α , 7 α -dihydroxy-24-nor-5 β -cholan-23-ol (BAR502), 6 β -ethyl-3 α , 7 β -dihydroxy-24-nor-5 β -cholan-23-ol (BARn501) and 6 β -ethyl-3 α , 7 α -dihydroxy-24-nor-5 β -cholan-23-ol (BARn504-6b)

[0090] 7-KLCA (1g, 2.56 mmol) was subjected to Beckmann degradation at C24 and methylation at C-23 furnishing **13** in 66% yield. Acetylation at C-3 and alkylation furnished **14** that was hydrogenated affording **15**. MeONa/MeOH treatment gave concomitant hydrolysis at C-3 and epimerization at C-6. Simultaneous reduction at C-23 methyl ester function and at C-7 carbonyl group furnished BAR502 in 89% yield. Intermediate **15** (250 mg, 0.54 mmol) was also used as starting material in the preparation of BARn501 and BARn504-6b.



a) HCOOH , HClO_4 ; b) TFA, trifluoroacetic anhydride, NaNO_2 ; c) KOH 30% in $\text{MeOH}/\text{H}_2\text{O}$ 1:1 v/v, 88% over three steps; d) p-TsOH, MeOH dry; e) acetic anhydride, pyridine; f) DIPA, *n*-BuLi, TMSCl, TEA dry, THF dry -78°C ; g) acetaldehyde, $\text{BF}_3(\text{OEt})_2$, CH_2Cl_2 , -60°C , 60% over four steps; h) H_2 Pd(OH)₂, THF/MeOH 1:1, quantitative yield; i) MeONa, MeOH; j) LiBH_4 , MeOH, THF dry, 0°C , 70% over two steps; k) NaBH_4 , MeOH dry, 0°C ; l) LiBH_4 , MeOH, THF dry, 0°C , 77% over two steps.

Steps a-d) Preparation of methyl 7-keto-24-nor-LCA (13)

[0091] Compound **13** (660 mg, 1.69 mmol, 66% over four steps) was prepared from 7-KLCA in the same operative condition described in example 1B, steps a-d).

[0092] **Steps e-h) Preparation of methyl 3 α -acetoxy-6 β -ethyl-7-keto-24-nor-5 β -cholan-23-oate (15).** Compound **13** (660 mg, 1.69 mmol) was subjected to the same operative condition described in example 2A, steps b-d to obtain 603 mg of **14** (78% over three steps). NMR analysis demonstrated a diastomeric ratio E/Z >95%. The *E* configuration at the exocyclic double bond was established by dipolar coupling H₃-25 (δ 1.67)/H-5 (δ 2.61) in Noesy spectrum (400 MHz, mixing time 400 ms).

(*E*-3 α -acetoxy-6-ethylidene-7-keto-24-nor-5 β -cholan-23-oate (14): C₂₈H₄₂O₅)

[0093] The ¹H NMR was recorded on Varian Inova 400 MHz, using CDCl₃ as solvent: δ 6.17 (1H, q, *J* = 7.2 Hz, H-24), 4.75 (1H, m, H-3), 3.64 (3H, s, COOCH₃), 2.61 (1H, dd, *J* = 13.1, 4.0 Hz, H-5), 1.98 (3H, s, COCH₃), 1.67 (3H, d, *J* = 7.2 Hz, H₃-25), 1.00 (3H, s, H₃-19), 0.97 (3H, d, *J* = 6.8 Hz, H₃-21), 0.67 (3H, s, H₃-18).

[0094] The ¹³C NMR was recorded on Varian Inova 100 MHz, using CDCl₃ as solvent: δ 204.5, 174.2, 170.5, 143.0, 130.6, 72.5, 54.7, 51.4, 50.7, 48.6, 45.3, 43.7, 41.5, 39.1, 38.8, 34.6, 34.2, 33.6, 33.4, 28.5, 25.9 (2C), 22.8, 21.3 (2C), 19.7, 12.7, 12.1. Hydrogenation on Pd(OH)₂ in the same operative condition described in example 2A, step e, furnished 600 mg of **15** (quantitative yield).

[0095] The β configuration of ethyl group at C-6 was determined by dipolar couplings H₃-25 (δ 0.83)/ H₃-19 (δ 1.22) in Noesy spectrum (400 MHz, mixing time 400 ms). **3 α -acetoxy-6 β -ethyl-7-keto-24-nor-5 β -cholan-23-oate (15): C₂₈H₄₄O₅** The ¹H NMR was recorded on Varian Inova 400 MHz, using CDCl₃ as solvent: δ 4.65 (1H, m, H-3), 3.67 (3H, s, COOCH₃), 2.60 (1H, t, *J* = 11.2 Hz, H-8), 2.43 (1H, dd, *J* = 14.2, 2.6 Hz, H-22a), 1.98 (3H, s, COCH₃), 1.88 (1H, m ovl, H-6), 1.22 (3H, s, H₃-19), 0.98 (3H, d, *J* = 6.4 Hz, H₃-21), 0.83 (3H, t, *J* = 7.0 Hz, H₃-25), 0.70 (3H, s, H₃-18).

[0096] The ¹³C NMR was recorded on Varian Inova 100 MHz, using CDCl₃ as solvent: δ 215.3, 174.0, 170.5, 72.8, 61.9, 55.0, 51.4, 49.2, 48.7, 45.5, 42.9, 42.6, 41.4, 38.7 (2C), 35.6, 35.3, 34.9, 28.3 (2C), 26.5, 25.9, 24.8, 21.4, 21.3, 19.6, 13.0, 12.1.

Steps i,j) Preparation of 6 α -ethyl-3 α , 7 α -dihydroxy-24-nor-5 β -cholan-23-ol (BAR502)

[0097] To a solution of compound **15** (450 mg, 1.0 mmol) and dry methanol (4 mL), MeONa (20 mL, 0.5 M in MeOH, 10 mmol) was added. After 24 h, H₂O was added dropwise. Then after evaporation of the solvents, the residue was diluted with water and extracted with AcOEt (3×50 mL). The combined extract was washed with water, dried with Na₂SO₄, and evaporated to give **16** that was subjected to the step g without further purification.

Methyl 6 α -ethyl-3 α -hydroxy-7-keto-24-nor-5 β -cholan-23-oate (16): C₂₆H₄₂O₄

[0098] The ¹H NMR was recorded on Varian Inova 400 MHz, using CDCl₃ as solvent: δ 3.64 (3H, s, COOCH₃), 3.45 (1H, m, H-3), 2.83 (1H, q, *J* = 7.3 Hz, H-6), 2.51 (1H, t, *J* = 11.2 Hz, H-8), 2.45 (1H, dd, *J* = 14.5, 3.2 Hz, H-22a), 1.26 (3H, s, H₃-19), 0.98 (3H, d, *J* = 6.6 Hz, H₃-21), 0.81 (3H, t, *J* = 7.0 Hz, H₃-25), 0.73 (3H, s, H₃-18).

[0099] The ¹³C NMR was recorded on Varian Inova 100 MHz, using CDCl₃ as solvent: δ 214.9, 175.4, 71.6, 56.2, 53.2, 52.0, 51.9, 51.0, 50.5, 45.2, 43.8, 42.2, 40.2, 36.7, 35.3, 34.8, 32.5, 30.5, 29.4, 25.6, 24.0, 22.9, 20.1, 20.0, 12.6, 12.4.

[0100] Compound **16** was subjected to LiBH₄ reduction in the same operative condition described in example 1A, step g. Silica gel chromatography eluting with hexane/EtOAc 6:4 afforded BAR502 (274 mg, 70% over two steps). An analytic sample was obtained by HPLC on a Nucleodur 100-5 C18 (5 μ m; 4.6 mm i.d. × 250 mm) with MeOH/H₂O (88:12) as eluent (flow rate 1 mL/min, *t_R*=10.8 min).

[0101] BAR502: C₂₅H₄₄O₃

[0102] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 3.65 (1H, s, H-7), 3.61 (1H, m, H-23a), 3.53 (1H, m, H-23b) 3.31 (1H, m, H-3), 0.97 (3H, d, *J* = 6.6 Hz, H₃-21), 0.92 (3H, s, H₃-19), 0.91 (3H, t, *J* = 7.0 Hz, H₃-25), 0.71 (3H, s, H₃-18).

[0103] The ¹³C NMR was recorded on Varian Inova 100 MHz, using CD₃OD as solvent: δ 73.2, 71.1, 60.7, 57.7, 51.4, 46.9, 43.8, 42.9, 41.3, 40.9, 39.8, 36.7, 36.5, 34.6, 34.5, 34.2, 31.2, 29.4, 24.5, 23.7, 23.4, 21.8, 19.3, 12.1, 11.9.

[0104] Steps k,l). Preparation of 6 β -ethyl-3 α , 7 β -dihydroxy-24-nor-5 β -cholan-23-ol (BARn501) and 6 β -ethyl-3 α , 7 α -dihydroxy-24-nor-5 β -cholan-23-ol (BARn504-6b). Compound **15** (100 mg, 0.22 mmol) was subjected to the same operative condition described in example 2A, steps f-g. HPLC purification on a Nucleodur 100-5 C18 (5 μ m; 10 mm i.d. × 250 mm) with MeOH/H₂O (86:14) as eluent (flow rate 3 mL/min), gave 47 mg of BARn501 (54%,

$t_R = 11$ min) and 20 mg of BARn504-6b (23%, $t_R = 15$ min).

[0105] BARn501: $C_{25}H_{44}O_3$

[0106] The 1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 3.73 (1H, dd, $J = 10.5, 5.5$ Hz, H-7), 3.61 (1H, m, H-23a), 3.51 (1H, m, ovl, H-23b), 3.51 (1H, m, ovl, H-3), 0.98 (3H, d, ovl, H_3 -21), 0.97 (3H, s, H_3 -19), 0.96 (3H, t, ovl, H_3 -25), 0.70 (3H, s, H_3 -18).

[0107] The ^{13}C NMR was recorded on Varian Inova 100 MHz, using CD_3OD as solvent: δ 75.2, 71.8, 60.8, 57.5, 56.6, 51.5, 45.5, 44.8, 42.0, 41.4, 40.7, 40.3, 39.9, 36.9, 36.0, 34.2, 30.5, 29.6, 28.3, 26.2, 23.4, 22.0, 19.4, 14.7, 12.9.


[0108] BARn504-6b: $C_{25}H_{44}O_3$

[0109] The 1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 3.63 (1H, m, H-23a), 3.60 (1H, m, H-7), 3.55 (1H, m, H-23b), 3.37 (1H, m, H-3), 2.30 (1H, q, $J = 12.5$ Hz, H-4a), 0.97 (3H, d, $J = 6.6$ Hz, H_3 -21), 0.95 (3H, s, H_3 -19), 0.95 (3H, t, $J = 7.0$ Hz, H_3 -25), 0.72 (3H, s, H_3 -18).

[0110] The ^{13}C NMR was recorded on Varian Inova 100 MHz, using CD_3OD as solvent: δ 72.8, 72.7, 60.8, 57.9, 52.7, 51.4, 47.5, 43.7, 42.3, 41.0, 39.9, 37.5, 37.3, 36.7, 34.2, 33.3, 31.0, 29.6, 29.4, 26.2, 24.8, 21.6, 19.3, 14.5, 12.1.

EXAMPLE 2D. Synthesis of 6-ethylidene-3 α ,7 β -dihydroxy-5 β -cholan-24-ol (BAR503), 6 α -ethyl-3 α ,7 β -dihydroxy-5 β -cholan-24-ol (BAR501-6a), 6-ethylidene-3 α ,7 β -dihydroxy-24-nor-5 β -cholan-23-ol (BARn503) and 6 α -ethyl-3 α ,7 β -dihydroxy-24-nor-5 β -cholan-23-ol (BARn501-6a)

[0111] Intermediate **11** was subjected to $NaBH_4$ reduction followed by treatment with $LiBH_4$. Alternatively $LiAlH_4$ treatment proceeded in a straightforward manner affording the concomitant reduction at C-24 and C-7. BAR503 was also used as starting material for BAR501-6a by hydrogenation on $Pd(OH)_2$ catalyst. The same synthetic protocol was performed on intermediate **14** producing the corresponding 23-derivatives, BARn503 and BARn501-6a.



a) $NaBH_4$, MeOH; b) $LiBH_4$, MeOH dry, THF, 0 °C, 85% over two steps; c) H_2 , $Pd(OH)_2$,

THF:MeOH 1:1 v/v.

Steps a,b). Preparation of 6-ethylidene-3 α , 7 β -dihydroxy-5 β -cholan-24-ol (BAR503) and 6-ethylidene-3 α ,7 β -dihydroxy-24-*nor*-5 β -cholan-23-ol (BARn503).

[0112] Compound **11** (1 g, 2.11 mmol) was subjected to the same operative condition described in example 2A, steps f, g. HPLC purification on a Nucleodur 100-5 C18 (5 μ m; 10 mm i.d. $\times$ 250 mm) with MeOH/H₂O (88:12) as eluent (flow rate 3 mL/min), gave 727 mg of BAR503 (85% over two steps, t_R = 9.2 min). Alternatively LiAlH₄ treatment on **11** furnished BAR503.

[0113] **BAR503:** C₂₆H₄₄O₃

[0114] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 5.66 (1H, q, J = 6.9 Hz, H-25), 3.90 (1H, d, J = 9.8 Hz, H-7), 3.55 (1H, m, H-3), 3.50 (2H, m, H₂-24), 2.50 (1H, dd, J = 4.0, 13.1 Hz, H-5), 1.62 (3H, d, J = 6.9 Hz, H₃-26), 0.97 (3H, d, J = 6.8 Hz, H₃-21), 0.81 (3H, s, H₃-19), 0.70 (3H, s, H₃-18).

[0115] The ¹³C NMR was recorded on Varian Inova 100 MHz, using CD₃OD as solvent: δ 142.7, 114.5, 73.4, 71.1, 63.6, 57.1, 56.1, 45.2, 44.9, 44.2, 40.7, 40.2, 36.3, 36.2, 35.9, 34.7, 32.4, 30.2, 29.5, 28.8, 27.4, 22.6, 21.5, 18.5, 11.8, 11.7.

[0116] The same synthetic protocol was performed on intermediate **14**. HPLC purification on a Nucleodur 100-5 C18 (5 μ m; 10 mm i.d. $\times$ 250 mm) with MeOH/H₂O (86:14) as eluent (flow rate 3 mL/min), gave BARn503 (t_R = 8 min).

[0117] **BARn503:** C₂₅H₄₂O₃

[0118] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 5.66 (1H, q, J = 6.8 Hz, H-24), 3.92 (1H, d, J = 9.9 Hz, H-7), 3.60 (1H, m, H-23a), 3.56 (1H, m, H-3), 3.55 (1H, m, H-23b), 2.52 (1H, dd, J = 3.7, 13.2 Hz, H-5), 1.63 (3H, d, J = 6.8 Hz, H₃-25), 0.98 (3H, d, J = 6.5 Hz, H₃-21), 0.95 (3H, s, H₃-19), 0.71 (3H, s, H₃-18).

[0119] The ¹³C NMR was recorded on Varian Inova 100 MHz, using CD₃OD as solvent: δ 143.7, 115.4, 74.1, 71.8, 60.8, 58.0, 57.1, 46.1, 45.9, 45.1, 41.6, 41.1, 39.9, 37.0, 36.4, 35.8, 34.1, 30.9, 29.8, 28.1, 23.5, 22.5, 19.5, 12.7, 12.6.

Step c). Preparation of 6 α -ethyl-3 α ,7 β -dihydroxy-5 β -cholan-24-ol (BAR501-6a) and 6 α -ethyl-3 α ,7 β -dihydroxy-24-*nor*-5 β -cholan-23-ol (BARn501-6a)

[0120] **BAR503** (350 mg, 0.86 mmol) was subjected to the same operative condition described in example 2A step e, obtaining BAR501-6a in quantitative yield.

[0121] **BAR501-6a**: $C_{26}H_{46}O_3$

[0122] The 1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 3.50 (2H, t, $J = 6.8$ Hz, H_3-24), 3.44 (1H, m, $H-3$), 3.07 (1H, t, $J = 9.8$ Hz, $H-7$), 0.96 (3H, d, $J = 6.8$ Hz, H_3-21), 0.95 (3H, s, H_3-19), 0.86 (3H, t, $J = 7.4$ Hz, H_3-26), 0.71 (3H, s, H_3-18).

[0123] The ^{13}C NMR was recorded on Varian Inova 100 MHz, using CD_3OD as solvent: δ 76.5, 72.3, 63.6, 57.9, 57.3, 46.3, 45.0, 44.8, 41.8, 41.0, 39.9, 37.0, 36.4, 35.5, 33.3, 31.3, 31.0, 30.3, 29.8, 27.8, 24.3, 22.5, 22.0, 19.3, 12.8, 11.8.

[0124] **BARn503** was subjected to the same operative condition described in example 2A step e, obtaining **BARn501-6a** in quantitative yield.


[0125] **BARn501-6a**: $C_{25}H_{44}O_3$

[0126] The 1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 3.62 (1H, m, $H-23a$), 3.54 (1H, m, $H-23b$), 3.45 (1H, m, $H-3$), 3.08 (1H, t, $J = 9.8$ Hz, $H-7$), 0.97 (3H, d, $J = 6.5$ Hz, H_3-21), 0.95 (3H, s, H_3-19), 0.86 (3H, t, $J = 7.4$ Hz, H_3-25), 0.73 (3H, s, H_3-18).

[0127] The ^{13}C NMR was recorded on Varian Inova 100 MHz, using CD_3OD as solvent: δ 76.4, 72.5, 60.8, 57.9, 57.2, 46.2, 45.1, 44.7, 41.8, 41.2, 40.0, 39.8, 36.5, 35.6, 34.2, 31.2, 30.9, 29.9, 27.9, 24.1, 22.7, 22.0, 19.5, 12.7, 11.7.

EXAMPLE 2E Synthesis of 6 α -ethyl-3 α ,7 α -dihydroxy-24-nor-5 β -cholan-23-nitrile (**BAR506**)

[0128] 7-KLCA was transformed in nitrile **17** following the same synthetic procedure described in Example 1B steps a-b. Alkylation followed by double bond reduction and epimerization at C-6 in the same operative condition described in example 2A steps c-d and example 2C step i, respectively furnished **18**. $LiBH_4$ treatment as in example 2C step j afforded the desired 7 α hydroxyl group in **BAR506**.



a) HCOOH, HClO₄; b) TFA, trifluoroacetic anhydride, NaNO₂; c) DIPA, *n*-BuLi, TMSCl, TEA dry, THF dry -78 °C; d) acetaldehyde, BF₃(OEt)₂, CH₂Cl₂, -60 °C; e) H₂, Pd(OH)₂, THF/MeOH 1:1; f) MeONa, MeOH; g) LiBH₄, MeOH, THF dry, 0 °C


[0129] **BAR506:** C₂₅H₄₁NO₂

[0130] The ¹H NMR was recorded on Varian Inova 700 MHz, using CD₃OD as solvent: δ 3.66 (1H, br s, H-7), 3.31 (1H, ovl, H-3), 2.46 (1H, dd, *J* = 3.8, 16.9 Hz, H-22a), 2.34 (1H, dd, *J* = 7.4, 16.9 Hz, H-22b), 1.16 (3H, d, *J* = 6.5 Hz, H₃-21), 0.91 (3H, t, *J* = 7.5 Hz, H₃-25), 0.92 (3H, s, H₃-19), 0.73 (3H, s, H₃-18).

[0131] The ¹³C NMR was recorded on Varian Inova 175 MHz, using CD₃OD as solvent: δ 120.3, 72.9, 70.9, 56.1, 51.5, 46.7, 43.4, 42.9, 41.4, 40.2, 36.5, 36.2, 34.3 (2C), 34.2, 30.7, 29.2, 24.9, 24.4, 23.4, 23.3, 21.9, 18.5, 12.1, 11.6.

EXAMPLE 2F. Synthesis of 6α-ethyl-7α-hydroxy-5β-cholan-24-ol (BAR701), 6α-ethyl-7α-hydroxy-5β-cholan-24-yl 24-sodium sulfate (BAR701solf), 6β-ethyl-7β-hydroxy-5β-cholan-24-ol (BAR702), 6α-ethyl-7β-hydroxy-5β-cholan-24-ol (BAR703), 6α-ethyl-7α-hydroxy-5β-cholan-24-oic acid (BAR704), 6α-ethyl-7α-hydroxy-5β-cholan-24-oyl taurine sodium sulfate (BART704), 6β-ethyl-7α-hydroxy-5β-cholan-24-ol (BAR705) and 6α-ethyl-7β-hydroxy-5β-cholan-24-oic acid (BAR711)

[0132] Compound **12** was treated with MeONa in methanol to obtain deacetylation at C-3 and inversion at C-6. Tosylation, elimination and hydrogenation of the double bond on ring A gave **20**. Hydrolysis at methyl ester function followed by LiBH₄ treatment furnished BAR704 in high chemical yield. Intermediate **20** was also used as starting material for BAR701. Sulfation on C-24 on a small aliquot of BAR701 furnished BAR701solf.



a) MeONa, MeOH; b) *p*-TsCl, pyridine, quantitative yield over two steps; c) LiBr, O₂CO₃, DMF, reflux, d) H₂, Pd(OH)₂, THF/MeOH 1:1, room temperature, 88% over two steps; e) NaOH, MeOH:H₂O 1:1 v/v, 82%; f) LiBH₄, MeOH dry, THF, 0 °C, 83%; g) DMT-MM, Et₃N, taurine, DMF dry; h) LiBH₄, MeOH dry, THF, 0 °C, 77 %; i) Et₃N·SO₃, DMF, 95 °C.

Steps a-f) Preparation of 6α-ethyl-7α-hydroxy-5β-cholan-24-oic acid (BAR704).

[0133] Compound **12** (500 mg, 1.05 mmol) was treated with MeONa (2.1 mL, 0.5 M in MeOH,

1.05 mmol) in MeOH (5 mL) overnight in the same operative condition of Example 2C step i. Tosylation on the crude reaction product in the same operative condition of Example 1C, step a, furnished **19** (620 mg, quantitative yield over two steps). Intermediate **19** (500 mg, 0.85 mmol) was subjected to the same operative condition of Example 1C, steps b,c, to obtain 312 mg of **20** (88% over two steps). Compound **20** (200 mg, 0.48 mmol) was hydrolyzed with NaOH (96 mg, 2.4 mmol) in a solution of MeOH:H₂O 1:1 v/v (10 mL) in the same operative condition of Example 1A step f. Crude carboxylic acid intermediate (190 mg, 0.47 mmol) was treated with LiBH₄ (1.65 mL, 2 M in THF, 3.3 mmol) and MeOH (133 μ L, 3.3 mmol) in THF dry (5 mL). Purification by silica gel (CH₂Cl₂-MeOH 99:1) furnished 157 mg of BAR704 (83%). In same embodiments LiBH₄ treatment after alkaline hydrolysis produced small amounts (about 10%) of 6 α -ethyl-7 β -hydroxy-5 β -cholan-24-oic acid (BAR711) that was isolated by HPLC purification on a Nucleodur 100-5 C18 (5 μ m; 10 mm i.d. $\times$ 250 mm) with MeOH/H₂O (88:12) as eluent (flow rate 3 mL/min, t_R = 16 min).

[0134] BAR704: C₂₆H₄₄O₃

[0135] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 3.65 (1H, br s, H-7), 2.34 (1H, m, H-23a), 2.20 (1H, m, H-23b), 0.96 (3H, d, J = 6.3 Hz, H₃-21), 0.92 (3H, s, H₃-19), 0.89 (3H, t, J = 7.4 Hz, H₃-26), 0.70 (3H, s, H₃-18). The ¹³C NMR was recorded on Varian Inova 100 MHz, using CD₃OD as solvent: δ 178.0, 71.6, 57.4, 51.7, 48.7, 43.8, 43.3, 41.5, 41.1, 39.3, 37.4, 36.8, 34.6, 32.5 (2C), 29.3, 28.8, 25.1, 24.6 (2C), 23.5, 22.5, 22.0, 18.8, 12.2, 12.1.

[0136] BAR711: C₂₆H₄₄O₃

[0137] The ¹H NMR was recorded on Varian Inova 500 MHz, using CD₃OD as solvent: δ 3.08 (1H, t, J = 9.6 Hz, H-7), 2.32 (1H, m, H-23a), 2.20 (1H, m, H-23b), 0.96 (3H, d, J = 6.2 Hz, H₃-21), 0.95 (3H, s, H₃-19), 0.85 (3H, t, J = 7.0 Hz, H₃-26), 0.70 (3H, s, H₃-18).

Step g) Preparation of 6 α -ethyl-7 α -hydroxy-5 β -cholan-24-oyl taurine sodium sulfate (BART704)

[0138] An aliquot of BAR704 (10 mg, 0.024 mmol) in DMF dry (5 mL) was treated with DMT-MM (20.5 mg, 0.07 mmol) and triethylamine (83 μ L, 0.6 mmol) and the mixture was stirred at room temperature for 10 min. Then to the mixture was added taurine (18 mg, 0.14 mmol). After 3 h, the reaction mixture was concentrated under vacuo and dissolved in water (5 mL). Purification on C18 silica gel column and then HPLC on a Nucleodur 100-5 C18 (5 μ m; 10 mm i.d. $\times$ 250 mm) with MeOH/H₂O (83:17) as eluent (flow rate 3 mL/min), gave 4.5 mg BART704 (t_R = 10 min).

[0139] BART704: $C_{28}H_{48}NNaO_5S$

[0140] The 1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 3.65 (1H, br s, H-7), 3.58 (2H, t, $J = 7.0$ Hz, CH_2-N), 2.96 (2H, t, $J = 9.6$ Hz, CH_2-S), 2.25 (1H, m, H-23a), 2.10 (1H, m, H-23b), 0.97 (3H, d, $J = 6.4$ Hz, H_3-21), 0.92 (3H, s, H_3-19), 0.89 (3H, t, $J = 7.1$ Hz, H_3-26), 0.70 (3H, s, H_3-18).

Step h) Preparation of 6 α -ethyl-7 α -hydroxy-5 β -cholan-24-ol (BAR701)

[0141] Compound **20** (100 mg, 0.24 mmol) was treated in the same operative condition of Example 2C step j. HPLC purification on a Nucleodur 100-5 C18 (5 μ m; 10 mm i.d. $\times$ 250 mm) with MeOH/ H_2O (92:8) as eluent (flow rate 3 mL/min), gave 64 mg of BAR701 ($t_R = 31$ min) and a small amount of 6 α -ethyl-7 β -hydroxy-5 β -cholan-24-ol (BAR703) (8 mg, $t_R = 24.8$ min).

[0142] BAR701: $C_{26}H_{46}O_2$

[0143] The 1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 3.65 (1H, br s, H-7), 3.51 (2H, m, H_2-24), 0.97 (3H, d, $J = 6.3$ Hz, H_3-21), 0.92 (3H, s, H_3-19), 0.89 (3H, t, $J = 7.3$ Hz, H_3-26), 0.71 (3H, s, H_3-18).

[0144] The ^{13}C NMR was recorded on Varian Inova 100 MHz, using CD_3OD as solvent: δ 71.6, 63.6, 57.6, 51.8, 48.7, 43.7, 43.3, 41.5, 41.1, 39.3, 37.5, 37.0, 34.6, 33.2, 30.3, 29.4, 28.8, 25.1, 24.6 (2C), 23.5, 22.5, 22.0, 19.2, 12.3, 12.1.

[0145] BAR703 $C_{26}H_{46}O_2$

[0146] The 1H NMR was recorded on Varian Inova 500 MHz, using CD_3OD as solvent: δ 3.51 (2H, m, H_2-24), 3.07 (1H, t, $J = 10.0$ Hz, H-7), 0.96 (3H, d, $J = 6.6$ Hz, H_3-21), 0.84 (3H, t, $J = 7.0$ Hz, H_3-26), 0.95 (3H, s, H_3-19), 0.71 (3H, s, H_3-18).

[0147] The ^{13}C NMR was recorded on Varian Inova 100 MHz, using CD_3OD as solvent: δ 76.3, 63.6, 57.9, 56.8, 46.3, 45.3, 45.0, 44.7, 41.7, 41.1, 38.8, 37.0, 36.3, 33.3, 30.3, 28.1, 27.9 (2C), 25.0 (2C), 22.0, 21.9 (2C), 19.4, 12.7, 11.6.

Step i) Preparation of 6 α -ethyl-7 α -hydroxy-5 β -cholan-24-yl 24-sodium sulfate (BAR701solf)

[0148] Sulfation on C-24 on a small aliquot of BAR701 was performed in the same operative


conditions described in example 1B step g) to give crude BAR701solf as ammonium salt. RP18/HPLC on a Nucleodur 100-5 C18 (5 μ m; 10 mm i.d. $\times$ 250 mm) with MeOH/H₂O (82:18) as eluent (flow rate 3 mL/min) afforded BAR701solf (t_R = 14.2 min) as sodium salt.

[0149] BAR701solf: C₂₆H₄₅ NaO₅S

[0150] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 3.96 (2H, t, J = 6.3 Hz, H₂-24), 3.64 (1H, br s, H-7), 0.96 (3H, d, J = 6.6 Hz, H₃-21), 0.91 (3H, s, H₃-19), 0.88 (3H, t, J = 7.4 Hz, H₃-26), 0.69 (3H, s, H₃-18).

Preparation of 6 β -ethyl-7 β -hydroxy-5 β -cholan-24-ol (BAR702) and 6 β -ethyl-7 α -hydroxy-5 β -cholan-24-ol (BAR705)

[0151] Compound **12** (500 mg, 1.05 mmol) was treated with MeONa (2.1 mL, 0.5 M in MeOH, 1.05 mmol) in MeOH (10 mL) in the same operative condition of Example 2C, step i, except for reaction time (2 h). Tosylation on the crude reaction product in the same operative condition of Example 1C, step a, furnished **21** (620 mg, quantitative yield over two steps). Intermediate **21** (600 mg, 1.02 mmol) was subjected to the same operative condition of Example 1C, steps b,c, to obtain 400 mg of **22** (94%). Compound **22** (350 mg, 0.84 mmol) was reduced with NaBH₄/LiBH₄ in the same operative condition of Example 2A steps f,g. HPLC purification on a Nucleodur 100-5 C18 (5 μ m; 10 mm i.d. $\times$ 250 mm) with MeOH/H₂O (92:8) as eluent (flow rate 3 mL/min), furnished 180 mg of BAR702 (t_R = 25 min) and 75.4 mg of BAR705 (t_R = 13 min). Alternatively step e was performed with Ca(BH₄)₂, produced in situ.



a) MeONa, MeOH; b) p-TsCl, pyridine, quantitative yield over two steps; c) LiBr, O₂CO₃, DMF, reflux; d) H₂, Pd(OH)₂, THF/MeOH 1:1, room temperature, 94% over two steps; e) NaBH₄, MeOH; f) LiBH₄, MeOH dry, THF, 0 °C, 78% over two steps.

[0152] BAR702: C₂₆H₄₆O₂

[0153] The ¹H NMR was recorded on Varian Inova 700 MHz, using CD₃OD as solvent: δ 3.67 (1H, dd, J = 8.7, 4.7 Hz, H-7), 3.51 (2H, m, H₂-24), 0.98 (3H, s, H₃-19), 0.97 (3H, d, J = 6.6 Hz, H₃-21), 0.96 (3H, t, J = 7.4 Hz, H₃-26), 0.71 (3H, s, H₃-18).

[0154] The ¹³C NMR was recorded on Varian Inova 175 MHz, using CD₃OD as solvent: δ 75.5, 63.8, 57.6, 56.5, 44.2, 43.7, 42.8, 41.0, 40.9, 40.8, 38.2 (2C), 36.9, 34.4, 32.8, 29.7, 28.9,

27.0, 26.1, 24.7, 22.2, 22.0 (2C), 19.2, 13.9, 12.3.


[0155] BAR705: C₂₆H₄₆O₂

[0156] The ¹H NMR was recorded on Varian Inova 700 MHz, using CD₃OD as solvent: □ 3.59 (1H, br s, H-7), 3.51 (2H, m, H₂-24), 2.23 (1H, dq, *J* = 13.9, 4.0 Hz, H-4a), 0.97 (3H, d, *J* = 6.6, H₃-21), 0.95 (3H, t, *J* = 7.1 Hz, H₃-26), 0.94 (3H, s, H₃-19), 0.70 (3H, s, H₃-18).

[0157] The ¹³C NMR was recorded on Varian Inova 175 MHz, using CD₃OD as solvent: □ 73.1, 63.2, 57.3, 52.8, 51.4, 49.4, 43.8, 41.3, 39.7, 37.4 (2C), 37.2, 34.2, 32.6 (2C), 29.9, 29.5, 28.9, 28.3, 27.3, 24.5, 21.7, 21.2, 18.8, 14.3, 12.3.

EXAMPLE 2G. Synthesis of 6α-ethyl-3β,7α-dihydroxy-5β-cholan-24-oic acid (BAR710), 6α-ethyl-3β,7α-dihydroxy-5β-cholan-24-oyl taurine sodium sulfate (BART710), 6α-ethyl-3β,7α-dihydroxy-5β-cholan-24-ol (BAR706), 6α-ethyl-3β,7α-dihydroxy-5β-cholan-24-yl 24-sodium sulfate (BAR706solf), 6α-ethyl-3β,7β-dihydroxy-5β-cholan-24-ol (BAR707), 6β-ethyl-3β,7β-dihydroxy-5β-cholan-24-ol (BAR708) and 6β-ethyl-3β,7α-dihydroxy-5β-cholan-24-ol (BAR709) and 6α-ethyl-3β,7β-dihydroxy-5β-cholan-24-oic acid (BAR712).

[0158] In a convergent protocol inversion at C-3 on derivative **19** followed by treatment with MeONa/MeOH gave **23** that was used as starting material in the synthesis of BAR706, BAR706solf, BAR710 and BART710. Inversion at C-3 followed by reduction at C-7 and C-24, produced BAR708 and BAR709.



a) CH₃COOK, DMF:H₂O 5:1 v/v; b) NaOMe, MeOH, 74 % over two steps; c) NaOH, MeOH:H₂O 1:1 v/v; d) LiBH₄, MeOH dry, THF, 0 °C, 65% over two steps; e) DMT-MM, Et₃N, taurine, DMF dry; f) LiBH₄, MeOH dry, THF, 0 °C, 58%; g) Et₃N.SO₃, DMF, 95 °C; h) CH₃COOK, DMF:H₂O 5:1; i) NaBH₄, MeOH; j) LiBH₄, MeOH dry, THF, 0 °C, 74% over three steps.

Steps a-d) Preparation of 6α-ethyl-3β,7α-dihydroxy-5β-cholan-24-oic acid (BAR710)

[0159] A solution of **21** (600 mg, 1.0 mmol) and CH₃COOK (98 mg, 1.0 mmol) dissolved in water (2 mL) and N,N'-dimethylformamide (DMF, 10 mL) was refluxed for 2 h. The solution was cooled at room temperature and then ethyl acetate and water were added. The separated aqueous phase was extracted with ethyl acetate (3 × 30 mL). The combined organic phases

were washed with water, dried (Na_2SO_4) and evaporated to dryness to give 600 mg of mixture. Purification by silica gel (hexane/ethyl acetate 8:2 and 0.5% TEA) gave 350 mg of oleos oil. C-6 inversion in the same operative condition as described in Example 2C step i, furnished **23** (320 mg, 74% over two steps) that was subjected to hydrolysis followed to LiBH_4 treatment as described in Example 2F, steps e,f. HPLC purification on a Nucleodur 100-5 C18 (5 μm ; 10 mm i.d. $\times$ 250 mm) with $\text{MeOH}/\text{H}_2\text{O}$ (88:12) as eluent (flow rate 3 mL/min), gave 208 mg of BAR710 (65%, t_R = 11 min). Alternatively inversion at C-3 on **21** followed by alkaline hydrolysis and then LiBH_4 treatment afforded BAR710 in a straightforward manner.

[0160] In same embodiments LiBH_4 treatment after alkaline hydrolysis produced small amounts (about 10%) of 6 α -ethyl-3 β ,7 β -dihydroxy-5 β -cholan-24-oic acid (BAR712) that was isolated by HPLC purification on a Nucleodur 100-5 C18 (5 μm ; 10 mm i.d. $\times$ 250 mm) with $\text{MeOH}/\text{H}_2\text{O}$ (88:12) as eluent (flow rate 3 mL/min, t_R = 8 min)

[0161] BAR710: $\text{C}_{26}\text{H}_{44}\text{O}_4$

[0162] The ^1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 3.97 (1H, br s, H-3), 3.67 (1H, br s, H-7), 2.33 (1H, m, H-23a), 2.21 (1H, m, H-23b), 0.96 (3H, d, J = 6.5 Hz, H_3 -21), 0.94 (3H, s, H_3 -19), 0.91 (3H, t, J = 7.6 Hz, H_3 -26), 0.70 (3H, s, H_3 -18).

[0163] The ^{13}C NMR was recorded on Varian Inova 100 MHz, using CD_3OD as solvent: δ 178.3, 71.3, 67.5, 57.4, 51.7, 43.8, 42.8, 41.5, 41.2, 41.0, 37.0, 36.7, 33.8, 32.4, 32.0, 31.1 (2C), 29.3, 28.3, 24.6, 24.2, 23.3, 22.2, 18.8, 12.3, 12.2.

[0164] BAR712: $\text{C}_{26}\text{H}_{44}\text{O}_4$

[0165] The ^1H NMR was recorded on Varian Inova 500 MHz, using CD_3OD as solvent: δ 4.01 (1H, br s, H-3), 3.06 (1H, t, J = 9.7 Hz, H-7), 2.32 (1H, m, H-23a), 2.19 (1H, m, H-23b), 0.97 (3H, s, H_3 -19), 0.96 (3H, d, ovl, H_3 -21), 0.87 (3H, t, J = 7.7 Hz, H_3 -26), 0.71 (3H, s, H_3 -18).

Step e) Preparation of 6 α -ethyl-3 β ,7 α -dihydroxy-5 β -cholan-24-oyl taurine sodium sulfate (BART710)

[0166] An aliquot of BAR710 (10 mg) was treated in the same operative condition of Example 2F step g.

[0167] BART710: $\text{C}_{28}\text{H}_{48}\text{NNaO}_6\text{S}$

[0168] The ^1H NMR was recorded on Varian Inova 500 MHz, using CD_3OD as solvent: δ 3.97

(1H, br s, H-3), 3.67 (1H, br s, H-7), 3.59 (2H, t, $J = 6.8$ Hz, CH₂-N), 2.96 (2H, t, $J = 6.8$ Hz, CH₂-S), 0.97 (3H, d, $J = 6.4$ Hz, H₃-21), 0.95 (3H, s, H₃-19), 0.91 (3H, t, $J = 7.1$ Hz, H₃-26), 0.70 (3H, s, H₃-18).

Steps f) Preparation of 6 α -ethyl-3 β ,7 α -dihydroxy-5 β -cholan-24-ol (BAR706) and 6 α -ethyl-3 β ,7 β -dihydroxy-5 β -cholan-24-ol (BAR707)

[0169] Intermediate 23 (500 mg, 1.16 mmol) was treated with LiBH₄ (4 mL, 8.1 mmol) and MeOH (327 μ L, 8.1 mmol) in THF dry (10 mL) as described in Example 2C, step j. HPLC purification on a Nucleodur 100-5 C18 (5 μ m; 10 mm i.d. $\times$ 250 mm) with MeOH/H₂O (88:12) as eluent (flow rate 3 mL/min), gave BAR706 (250 mg, $t_R = 12.6$ min) and a small amount of 6 α -ethyl-3 β ,7 β -dihydroxy-5 β -cholan-24-ol (BAR707) (23 mg, $t_R = 8.2$ min).

[0170] BAR706: C₂₆H₄₆O₃

[0171] The ¹H NMR was recorded on Varian Inova 500 MHz, using CD₃OD as solvent: δ 3.97 (1H, br s, H-3), 3.66 (1H, br s, H-7), 3.51 (2H, m, H₂-24), 0.96 (3H, d, $J = 6.6$ Hz, H₃-21), 0.94 (3H, s, H₃-19), 0.91 (3H, t, $J = 7.5$ Hz, H₃-26), 0.70 (3H, s, H₃-18). The ¹³C NMR was recorded on Varian Inova 125 MHz, using CD₃OD as solvent: δ 71.4, 67.4, 63.6, 57.6, 51.7, 43.7, 42.8, 41.5, 41.2, 41.1, 37.1 (2C), 33.8, 33.2, 31.3 (2C), 30.3, 29.4, 28.3, 24.6, 24.2, 23.3, 22.3, 19.2, 12.7, 12.1.

[0172] BAR707: C₂₆H₄₆O₃

[0173] The ¹H NMR was recorded on Varian Inova 500 MHz, using CD₃OD as solvent: δ 4.01 (1H, br s, H-3), 3.51 (2H, m, H₂-24), 3.05 (1H, t, $J = 9.7$ Hz, H-7), 0.97 (3H, s, H₃-19), 0.96 (3H, d, $J = 6.4$ Hz, H₃-21), 0.88 (3H, t, $J = 7.6$ Hz, H₃-26), 0.72 (3H, s, H₃-18).

[0174] The ¹³C NMR was recorded on Varian Inova 100 MHz, using CD₃OD as solvent: δ 76.3, 67.1, 63.6, 57.8, 56.8, 45.0, 44.8, 44.5, 41.7, 40.3, 39.1, 37.0, 35.9, 33.0, 31.2, 30.2, 29.8, 28.4, 28.0, 27.9, 24.8, 22.9, 21.8, 19.3, 12.8, 11.6.

Step g) Preparation of 6 α -ethyl-3 β ,7 α -dihydroxy-5 β -cholan-24-yl 24-sodium sulfate (BAR706solf)

[0175] Sulfation on C-24 on a small aliquot of BAR706 was performed in the same operative conditions described in example 1B step g.

[0176] BAR706solf: $C_{26}H_{46}NaO_6S$

[0177] The 1H NMR was recorded on Varian Inova 500 MHz, using CD_3OD as solvent: δ 3.97 (1H, br s ovl, H-3), 3.96 (2H, t ovl, H_2-24), 3.65 (1H, br s, H-7), 0.96 (3H, d, $J = 6.6$ Hz, H_3-21), 0.94 (3H, s, H_3-19), 0.90 (3H, t, $J = 7.5$ Hz, H_3-26), 0.70 (3H, s, H_3-18).

Steps h,i,j) Preparation of 6 β -ethyl-3 β ,7 β -dihydroxy-5 β -cholan-24-ol (BAR708) and 6 β -ethyl-3 β ,7 α -dihydroxy-5 β -cholan-24-ol (BAR709).

[0178] Compound 19 was treated in the same operative condition of step a. $NaBH_4/LiBH_4$ reduction of 100 mg (0.23 mmol) in the same operative conditions of Example 2A, steps f,g, afforded a mixture whose HPLC purification (88% MeOH:H₂O) gave pure 6 β -ethyl-3 β ,7 β -dihydroxy-5 β -cholan-24-ol (BAR708) (48.3 mg, $t_R = 11$ min) and 6 β -ethyl-3 β ,7 α -dihydroxy-5 β -cholan-24-ol (BAR709) (20.7 mg, $t_R = 13$ min). Alternatively step i was performed with $Ca(BH_4)_2$, produced in situ.

[0179] BAR708: $C_{26}H_{46}O_3$

[0180] The 1H NMR was recorded on Varian Inova 700 MHz, using CD_3OD as solvent: δ 3.59 (1H, br s, H-3), 3.57 (1H, dd, $J = 12.6, 2.3$ Hz, H-7), 3.51 (2H, m, H_2-24), 0.98 (3H, s, H_3-19), 0.96 (3H, ovl, H_3-21), 0.96 (3H, t, ovl, H_3-26), 0.70 (3H, s, H_3-18).

[0181] The ^{13}C NMR was recorded on Varian Inova 175 MHz, using CD_3OD as solvent: δ 75.2, 68.3, 63.6, 58.3, 57.1, 45.7 (2C), 44.2, 41.8 (2C), 41.2, 40.0, 37.0, 35.9, 33.3, 31.1, 30.3, 29.4 (2C), 26.6 (2C), 23.2 (2C), 19.3, 13.0, 12.3.

[0182] BAR709: $C_{26}H_{46}O_3$


[0183] The 1H NMR was recorded on Varian Inova 700 MHz, using CD_3OD as solvent: δ 3.91 (1H, br s, H-3), 3.60 (1H, br s, H-7), 3.51 (2H, m, H_2-24), 2.45 (1H, t, $J = 13.3$ Hz, H-4a), 0.97 (3H, s, H_3-19), 0.97 (3H, ovl, H_3-21), 0.95 (3H, t, $J = 7.4$ Hz, H_3-26), 0.71 (3H, s, H_3-18).

[0184] The ^{13}C NMR was recorded on Varian Inova 175 MHz, using CD_3OD as solvent: δ 72.8, 67.4, 63.4, 57.2, 51.3, 51.2, 43.2, 41.6, 40.5, 37.3, 37.1 (2C), 36.9, 34.0, 33.3, 32.1, 30.3, 29.3, 28.9, 28.6, 26.3, 24.9, 22.0, 19.3, 13.8, 12.1.

EXAMPLE 2H. Synthesis of 6 α -ethyl-7 α -hydroxy-24-nor-5 β -cholan-23-oic acid (BARn704), 6 α -ethyl-7 α -hydroxy-24-nor-5 β -cholan-23-oyl taurine sodium sulfate (BARTn704), 6 α -ethyl-7 α -hydroxy-24-nor-5 β -cholan-23-ol (BARn701) and 6 α -ethyl-7 α -

hydroxy-24-nor-5 β -cholan-23-yl 23-sodium sulfate (BARn701solf)

[0185] BARn704, BARTn704, BARn701 and BARn701solf were prepared starting from **15** and following the same synthetic protocol described for their C24 homologues (Example 2F, steps a-i).



a) MeONa, MeOH; b) p-TsCl, pyridine, 73% over two steps; c) LiBr, Li₂CO₃, DMF, reflux, d) H₂, Pd(OH)₂, THF/MeOH 1:1, room temperature, quantitative yield over two steps; e) NaOH, MeOH:H₂O 1:1 v/v, 80%; f) LiBH₄, MeOH dry, THF, 0 °C, 92%; g) DMT-MM, Et₃N, taurine, DMF dry; h) LiBH₄, MeOH dry, THF, 0 °C, 70%; i) Et₃N.SO₃, DMF, 95 °C.

[0186] BARn704: C₂₅H₄₂O₃

[0187] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 3.64 (1H, br s, H-7), 2.41 (1H, dd, *J* = 11.0, 2.6 Hz, H-22a), 1.00 (3H, d, *J* = 6.0 Hz, H₃-21), 0.90 (3H, s, H₃-19), 0.87 (3H, t, *J* = 7.4 Hz, H₃-25), 0.71 (3H, s, H₃-18). The ¹³C NMR was recorded on Varian Inova 100 MHz, using CD₃OD as solvent: δ 178.9, 71.6, 57.5, 51.7, 48.6, 43.8, 43.4, 43.3, 41.5, 40.9, 39.1, 37.4, 35.2, 34.6, 29.4, 28.8, 25.0, 24.6 (2C), 23.5, 22.4, 22.0, 20.1, 12.2, 12.1.

[0188] BARTn704: C₂₇H₄₆NNaO₅S

[0189] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 3.64 (1H, br s, H-7), 3.59 (2H, t, *J* = 6.8 Hz, CH₂-N), 2.96 (2H, t, *J* = 6.8 Hz, CH₂-S), 2.40 (1H, dd, *J* = 11.0, 2.8 Hz, H-22a), 1.00 (3H, d, *J* = 6.0 Hz, H₃-21), 0.89 (3H, s, H₃-19), 0.86 (3H, t, *J* = 7.4 Hz, H₃-25), 0.70 (3H, s, H₃-18).

[0190] BARn701: C₂₅H₄₄O₂


[0191] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 3.66 (1H, br s, H-7), 3.61 (1H, m, H-23a), 3.54 (1H, m, H-23b), 0.96 (3H, d, *J* = 6.7 Hz, H₃-21), 0.91 (3H, s, H₃-19), 0.89 (3H, t, *J* = 7.3 Hz, H₃-25), 0.70 (3H, s, H₃-18). The ¹³C NMR was recorded on Varian Inova 100 MHz, using CD₃OD as solvent: δ 71.5, 60.8, 57.8, 51.7, 48.6, 43.8, 43.3, 41.5, 41.1, 39.9, 39.2, 37.4, 34.5, 34.2, 29.4, 28.8, 25.0, 24.6 (2C), 23.5, 22.5, 22.0, 19.4, 12.2, 12.1.

[0192] **BARn701solf**: $C_{25}H_{43}NaO_5S$

[0193] The 1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 4.02 (2H, m, H_{2-23}), 3.65 (1H, br s, H-7), 0.97 (3H, d, $J = 6.7$ Hz, H_{3-21}), 0.91 (3H, s, H_{3-19}), 0.88 (3H, t, $J = 7.5$ Hz, H_{3-25}), 0.70 (3H, s, H_{3-18}).

EXAMPLE 2I. Synthesis of 6 α -ethyl-3 β ,7 α -dihydroxy-24-nor-5 β -cholan-23-oic acid (BARn710), 6 α -ethyl-3 β ,7 α -dihydroxy-24-nor-5 β -cholan-23-oyl taurine sodium sulfate (BARTn710), 6 α -ethyl-3 β ,7 α -dihydroxy-24-nor-5 β -cholan-23-ol (BARn706), and 6 α -ethyl-3 β ,7 α -dihydroxy-24-nor-5 β -cholan-23-yl 23-sodium sulfate (BARn706solf)

[0194]



a) CH_3COOK , $DMF:H_2O$ 5:1 v/v; b) $NaOH$, $MeOH:H_2O$ 1:1 v/v; c) $LiBH_4$, $MeOH$ dry, THF, 0 °C, 58% over two steps; d) DMT-MM, Et_3N , taurine, DMF dry; e) $LiBH_4$, $MeOH$ dry, THF, 0 °C, 57%; f) $Et_3N \cdot SO_3$, DMF , 95 °C.

[0195] BARn710, BARTn710, BARn706 and BARn706solf were prepared starting from **24** and following the same synthetic protocol described for their C24 homologues (Example 2G, steps c-g).

[0196] **BARn710**: $C_{25}H_{42}O_4$

[0197] The 1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 3.97 (1H, br s, H-3), 3.66 (1H, br s, H-7), 2.42 (1H, dd, $J = 11.3, 3.3$ Hz, H-22a), 1.02 (3H, d, $J = 6.0$ Hz, H_{3-21}), 0.94 (3H, s, H_{3-19}), 0.91 (3H, t, $J = 7.3$ Hz, H_{3-25}), 0.73 (3H, s, H_{3-18}).

[0198] The ^{13}C NMR was recorded on Varian Inova 100 MHz, using CD_3OD as solvent: δ 177.7, 71.3, 67.4, 57.4, 51.8, 43.8 (2C), 42.8, 41.5, 41.2, 41.0, 37.0, 35.1, 33.8, 31.3, 31.2, 29.4, 28.3, 24.6, 24.2, 23.3, 22.2, 20.0, 12.2, 12.1.

[0199] **BARTn710**: $C_{27}H_{46}NNaO_6S$

[0200] The 1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 3.97 (1H, br s, H-3), 3.66 (1H, br s, H-7), 3.59 (2H, t, $J = 6.8$ Hz, CH_2-N), 2.96 (2H, t, $J = 6.8$ Hz, CH_2-S), 2.42 (1H, dd, $J = 11.3, 3.3$ Hz, H-22a), 1.00 (3H, d, $J = 6.3$ Hz, H_{3-21}), 0.94 (3H, s, H_{3-19}), 0.91 (3H, t, $J = 7.3$ Hz, H_{3-25}), 0.73 (3H, s, H_{3-18}).

19), 0.90 (3H, t, $J = 7.0$ Hz, H₃-25), 0.70 (3H, s, H₃-18).

[0201] BARn706: C₂₅H₄₄O₃

[0202] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 3.97 (1H, br s, H-3), 3.67 (1H, br s, H-7), 3.61 (1H, m, H-23a), 3.55 (1H, m, H-23b), 0.97 (3H, d, $J = 6.6$ Hz, H₃-21), 0.95 (3H, s, H₃-19), 0.91 (3H, t, $J = 7.4$ Hz, H₃-25), 0.71 (3H, s, H₃-18).


[0203] The ¹³C NMR was recorded on Varian Inova 100 MHz, using CD₃OD as solvent: δ 71.4, 67.4, 60.8, 57.9, 51.8, 43.8, 42.8, 41.5, 41.2, 41.1, 39.9, 37.0, 34.2, 33.8, 31.3, 31.2, 29.4, 28.3, 24.6, 24.2, 23.3, 22.3, 19.4, 12.2, 12.1.

[0204] BARn706solf: C₂₅H₄₃NaO₆S

[0205] The ¹H NMR was recorded on Varian Inova 400 MHz, using CD₃OD as solvent: δ 4.05 (2H, m, H₂-23), 3.97 (1H, br s, H-3), 3.66 (1H, br s, H-7), 1.00 (3H, d, $J = 6.0$ Hz, H₃-21), 0.94 (3H, s, H₃-19), 0.91 (3H, t, $J = 6.9$ Hz, H₃-25), 0.71 (3H, s, H₃-18).

EXAMPLE 2J. Preparation of 6 α -ethyl-3 α , 7 α -dihydroxy-25, 26-bishomo-5 β -cholan-26-oic acid (BAR802), 6 α -ethyl-3 α , 7 α -dihydroxy-25, 26-bishom0-5p-cholan-26-ol (BAR803), and 6 α -ethyl-3 α , 7 α -dihydroxy-25, 26-bishomo-5 β -cholan-26-yl-26-sodium sulfate (BAR804)

[0206]



a) 2,6-lutidine, t-butyldimethylsilyl trifluoromethanesulfonate, CH₂Cl₂, 0 °C; b) LiBH₄, MeOH dry, THF, 0 °C, 68% over two steps; c) DMSO, oxalyl chloride, TEA dry, CH₂Cl₂, -78 °C then methyl(triphenylphosphoranylidene)acetate, 79%; d) H₂, Pd(OH)₂/C Degussa type, THF/MeOH 1:1, quantitative yield; e) HCl 37%, MeOH, 87%; f) NaOH 5% in MeOH/H₂O 1:1 v/v, 89%; g) LiBH₄, MeOH dry, THF, 0 °C, 78%; h) Et₃N.SO₃, DMF, 95 °C, 25 %.

[0207] BAR802-804 were prepared following the same synthetic protocols as in Example 1A, steps a-g. Sulfation on a small aliquot of BAR803 in the same operative conditions of Example 1B step g afforded BAR804.

[0208] BAR802: C₂₈H₄₈O₄

[0209] The ^1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 3.66 (1H, br s, H-7), 3.31 (1H, m ovl, H-3), 2.24 (2H, t, $J = 7.3$ Hz, H_2 -25), 0.95 (3H, d, $J = 6.4$ Hz, H_3 -21), 0.92 (3H, s, H_3 -19), 0.91 (3H, t, $J = 6.9$ Hz, H_3 -28), 0.70 (3H, s, H_3 -18).

[0210] The ^{13}C NMR was recorded on Varian Inova 175 MHz, using CD_3OD as solvent: δ 186.7, 73.3, 71.3, 57.7, 51.7, 47.0, 43.7, 43.1, 41.6, 41.1, 37.1, 36.9, 36.8, 36.6, 34.5, 34.4 (2C), 31.3, 29.4, 27.0 (2C), 24.6, 23.8, 23.5, 22.0, 19.2, 12.2, 12.0.

[0211] BAR803: $\text{C}_{28}\text{H}_{50}\text{O}_3$

[0212] The ^1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 3.64 (1H, br s, H-7), 3.53 (2H, t, $J = 6.6$ Hz, H_2 -26), 3.30 (1H, m ovl, H-3), 0.94 (3H, d, $J = 6.7$ Hz, H_3 -21), 0.91 (3H, s, H_3 -19), 0.90 (3H, t, $J = 7.0$ Hz, H_3 -28), 0.68 (3H, s, H_3 -18).

[0213] The ^{13}C NMR was recorded on Varian Inova 175 MHz, using CD_3OD as solvent: δ 73.3, 71.3, 63.1, 57.7, 51.7, 47.0, 43.7, 43.1, 41.6, 41.1, 37.1, 36.8 (2C), 36.6, 34.5, 34.4, 33.7, 31.3, 29.5, 27.4, 27.0, 24.6, 23.8, 23.5, 22.0, 19.3, 12.3, 12.0. **BAR804:** $\text{C}_{28}\text{H}_{49}\text{NaO}_6\text{S}$

[0214] The ^1H NMR was recorded on Varian Inova 400 MHz, using CD_3OD as solvent: δ 3.99 (2H, t, $J = 6.6$ Hz, H_2 -26), 3.65 (1H, br s, H-7), 3.31 (1H, m ovl, H-3), 0.94 (3H, d, $J = 6.2$ Hz, H_3 -21), 0.91 (3H, s, H_3 -19), 0.90 (3H, t, $J = 7.0$ Hz, H_3 -28), 0.69 (3H, s, H_3 -18).

[0215] Biological Activities. Activity of selected compounds was tested in vitro using a whole cell model transfected with a reporter genes to establish selectivity of compounds shown in table 1 toward FXR and TGR5/GPBAR1 in comparison with chenodeoxycholic acid (CDCA) and TLCA. CDCA is a primary bile acid that functions as an endogenous ligand for FXR, while TLCA is a physiological ligand for TGR5/GPBAR1. In this assay, HepG2 cells (a liver-derived cell line) were cultured at 37 °C in minimum essential medium with Earl's salts containing 10% fetal bovine serum (FBS), 1% L-glutamine, and 1% penicillin/streptomycin. HEK-293T cells were cultured at 37 °C in D-MEM containing 10% fetal bovine serum (FBS), 1% L-glutamine, and 1% penicillin/streptomycin. The transfection experiments were performed using Fugene HD according to manufactured specifications. Cells were plated in a 24-well plate at 5×10^4 cells/well. For FXR mediated transactivation, HepG2 cells were transfected with 100 ng of pSG5-FXR, 100 ng of pSG5-RXR, 100 ng of pGL4.70 a vector encoding the human Renilla gene and 250 ng of the reporter vector p(hsp27)-TK-LUC containing the FXR response element IR1 cloned from the promoter of heat shock protein 27 (hsp27).

[0216] For GPBAR1 mediated transactivation, HEK-293T cells were transfected with 200 ng of pGL4.29, a reporter vector containing a cAMP response element (CRE) that drives the transcription of the luciferase reporter gene luc2P, with 100 ng of pCMVSPORT6-human

GPBAR1, and with 100 ng of pGL4.70 a vector encoding the human Renilla gene. In control experiments HEK-293T cells were transfected only with vectors pGL4.29 and pGL4.70 to exclude any possibility that compounds could activate the CRE in a GPBAR1 independent manner. At 24 h post-transfection, cells were stimulated for 18 h with 10 μ M TLCA as a control agent or putative GPBAR1 agonists as the same concentration. After treatments, cells were lysed in 100 μ L of lysis buffer (25 mM Tris-phosphate, pH 7.8; 2 mM DTT; 10% glycerol; 1% Triton X-100), and 20 μ L of cellular lysate was assayed for luciferase activity using the luciferase assay system. Luminescence was measured using Glomax 20/20 luminometer. Luciferase activities were normalized against Renilla activities. Antagonism against FXR of GPBAR1/TGR5 was measured as percent of activity in transactivation assay using activity of TLCA as example of agonism.

[0217] Animals and protocols. GPBAR1 null mice (GPBAR1-B6=GPBAR12/2 mice, generated directly into C57BL/6NCrl background), and congenic littermates on C57BL/6NCrl were housed under controlled temperatures (22 °C) and photoperiods (12:12-hour light/dark cycle), allowed unrestricted access to standard mouse chow and tap water and allowed to acclimate to these conditions for at least 5 days before inclusion in an experiment.

[0218] Scratching test. Male GPBAR1^{-/-} mice and their congenic littermates (8-12 weeks of age) were used for this studies. The fur at the base of the neck was shaved, and mice were placed in individual cylinders on a glass shelf. A circumference of approx. 0.5 cm of diameter was drawn in the neck and test agents injected in this area. Mice were acclimatized to the experimental room, restraint apparatus and investigators for 2 h periods on 2 successive days before experiments. Scratching behavior was quantified by 2 observers unaware of tested agents or genotypes. A scratch was defined as lifting the hind limb to the injection site and then a placing of the paw on the floor, regardless of the number of strokes. If counts differed by greater than 5 scratches over a 30-minute period, both observers reevaluated the records. Results were expressed as the number of scratching events during 30 or 60 min of observation. Tested agents were: DCA (25 μ g), TLCA (25 μ g), UDCA (25 μ g), and BAR502 (25 μ g), or with betulinic acid (50 μ g), oleanolic acid (50 μ g). LCA and DCA were dissolved in DMSO and the other agents in 0.9% NaCl (10 μ L). In another experimental setting GPBAR1^{-/-} mice and their congenic littermates were administered alpha-naphthylisothiocyanate (ANIT) (25 mg/kg, *per os*) dissolved in olive oil or olive oil alone (control mice) or with the combination of ANIT plus BAR502 (15 mg/Kg once a day, *per os*) for 10 days. At day 5 spontaneous scratching was evaluated for 60 min and after subcutaneous injection of 25 μ g DCA. Serum levels of total bilirubin, aspartate aminotransferase (AST) and alkaline phosphatase were measured by routine clinical chemistry testing performed on a Hitachi 717 automatic analyzer. For the estrogen model, wild type C57BL6 mice were administered 10 mg/Kg i.p. with 17 α -Ethinylestradiol (17 α E₂) dissolved in PEG or PEG alone (control mice) or the combination of 17 α E₂ and BAR502 (15 mg/Kg daily, *per os*) for 8 days. At the end of the study the spontaneous scratching and scratching induced by s.c. injection of 25 μ g DCA was recorded. Gallbladder weight and serum levels of bilirubin and alkaline phosphatase were also measured. Throughout the studies animals were visually assessed at least twice a day from

Monday to Friday and once a day over the week end by investigators and by highly trained animal facility personnel's including animal facility's veterinarian. Animals were weighted daily and sacrificed at indicated time points or when their clinical conditions become critical as assessed by a reduction of body weight higher than 25% of basal body weight in 7 days. In addition, animals were sacrificed when at the daily evaluation they demonstrate inability to rise or ambulate. Mice were euthanized by an overdose of sodium pentobarbital (>100 mg/kg i.p.).

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.


Patent documents cited in the description

- [WO2013192097A](#) [\[0008\]](#)
- [WO2008002573A](#) [\[0008\]](#)
- [WO2010014836A](#) [\[0009\]](#)

Non-patent literature cited in the description

- **MAKISHIMA et al.** Science, 1999, vol. 284, 1362- [\[0002\]](#)
- **PARKS et al.** Science, 1999, vol. 284, 1365- [\[0002\]](#)
- **ZHANG et al.** Proc. Natl. Acad. Sci. USA, 2006, vol. 103, 1006- [\[0003\]](#)
- **RENGA et al.** FASEB J., 2012, vol. 26, 3021-3031 [\[0003\]](#)
- **TAKEDA et al.** FEBS Lett., 2002, vol. 520, 97- [\[0004\]](#)
- **KAWAMATA et al.** J. Biol. Chem., 2003, vol. 278, 9435- [\[0004\]](#)
- **WATANABE et al.** Nature, 2006, vol. 439, 484- [\[0005\]](#)
- **THOMAS et al.** Cell. Metab., 2009, vol. 10, 167- [\[0005\]](#)
- **ALEMI et al.** J. Clin. Invest., 2013, vol. 123, 1513-1530 [\[0007\]](#) [\[0036\]](#)
- **SATO H.** J Med Chem., 2008, vol. 51, 4849- [\[0009\]](#)
- **D'AMORE C. et al.** J. Med. Chem., 2014, vol. 57, 937- [\[0010\]](#)


- IGUCHI Y. et al. J Lipid Res., 2010, vol. 51, 1432- [0011]
- KIHARA K. et al. Steroids, 1992, vol. 57, 4193-198 [0012]
- SWAAN P. W. et al. J. Comp.-Aid. Mol. Des., 1997, vol. 11, 581-588 [0013]
- BURNS et al. Steroids, 2011, vol. 76, 3291-300 [0014]
- J. Med. Chem., 2014, vol. 57, 937- [0040] [0041] [0043] [0045] [0057]

Patentkrav**1.** Forbindelse valgt fra gruppen bestående af

5

2. Forbindelsen ifølge krav 1, hvor forbindelsen er 6 α -ethyl-7 α -hydroxy-5 β -cholan-24-syre.**3.** Forbindelse til anvendelse som et medikament, idet nævnte forbindelse er valgt

10 fra gruppen bestående af:

**4.** Forbindelse til anvendelse ifølge krav 3, hvor nævnte forbindelse er 6 α -ethyl-7 α -hydroxy-5 β -cholan-24-syre.**5.** Forbindelse til anvendelse ifølge krav 3, som FXR- og/eller TGR5/GPBAR1-modulator, ved forebyggelsen og/eller behandlingen af gastrointestinale lidelser, leversygdomme, kardiovaskulære sygdomme, aterosklerose, metaboliske sygdomme, infektionssygdomme, kræft, nyrelidelser, inflammatoriske lidelser og neurologiske lidelser.**6.** Farmaceutisk sammensætning omfattende en forbindelse ifølge krav 1 og mindst en anden farmaceutisk bestanddel.

25