

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 200191431 B2**
(10) Patent No. **785384**

(54) Title
Environmental stress responsive promoter

(51)⁶ International Patent Classification(s)
C12N 15/11 (2006.01)⁸BMEP **C12N**
C12N 15/29 (2006.01) 15/29
C12N 15/11 20060101ALI2006040
20060101AFI2006040 ⁸BMEP

(21) Application No: 200191431 (22) Application Date: 2001 .11.21

(30) Priority Data

(31) Number	(32) Date	(33) Country
356652/2000	2000 .11.22	JP
309984/2001	2001 .10.05	JP

(43) Publication Date : 2002 .05.23
(43) Publication Journal Date : 2002 .05.23
(44) Accepted Journal Date : 2007 .03.29

(71) Applicant(s)
Toyota Jidosha Kabushiki Kaisha, Riken

(72) Inventor(s)
Kazuo Shinozaki, Tokihiko Nanjo, Motoaki Seki

(74) Agent/Attorney
F B Rice and Co,Level 23,44 Market Street,SYDNEY NSW 2000

(56) Related Art
WO 2002/016655 A2
EP 1033405 A2

ABSTRACT

The present invention provides a stress responsive promoter.

The environmental stress responsive promoter of the present invention comprises DNA of the following (a), (b) or (c):

- (a) DNA consisting of any nucleotide sequence selected from SEQ ID NOS: 1 to 18;
- (b) DNA consisting of a nucleotide sequence comprising a deletion, substitution or addition of one or more nucleotides relative to any nucleotide sequence selected from SEQ ID NOS: 1 to 18, and functioning as an environmental stress responsive promoter; and
- (c) DNA hybridizing under stringent conditions to DNA consisting of any nucleotide sequence selected from SEQ ID NOS: 1 to 18, and functioning as an environmental stress responsive promoter.



AUSTRALIA
Patents Act 1990

**Toyota Jidosha Kabushiki Kaisha;
Riken**

**COMPLETE SPECIFICATION
STANDARD PATENT**

Invention Title:

Environmental stress responsive promoter

The following statement is a full description of this invention
including the best method of performing it known to us:-

ENVIRONMENTAL STRESS RESPONSIVE PROMOTER

FIELD OF THE INVENTION

The present invention relates to an environmental stress responsive promoter.

BACKGROUND OF THE INVENTION

5 By means of gene sequencing projects, large quantities of genomic and cDNA sequences have been determined for a number of organisms, and in a plant model, *Arabidopsis thaliana*, the complete genomic sequences of two chromosomes have been determined (Lin, X. et al., (1999) Nature 402, 761-768.; Mayer, K. et al., (1999) Nature 402, 769-777.)

The EST (expressed sequence tag) project has also contributed greatly to the discovery of expression genes (Hofte, H. et al., (1993) Plant J. 4, 1051-1061.; Newman, T. et al., (1994) Plant Physiol. 106, 1241-1255.; Cooke, R. et al., (1996) Plant J. 9, 101-124. Asamizu, E. et al., (2000) DNA Res. 7, 175-180.) For example, dbEST (the EST database of the National Center for Biotechnology Information (NCBI)) comprises partial cDNA sequences, in which more than a half (about 28,000 genes) of the total gene complement is represented (as estimated from the gene content of the completely sequenced *Arabidopsis thaliana* chromosome 2 (Lin, X. et al., (1999) Nature 402, 761-768.))

20 In recent years, microarray (DNA chip) technology has become a useful tool for analysis of genome-scale gene expression (Schena, M. et al., (1995) Science 270, 467-470.; Eisen, M. B. and Brown, P. O. (1999) Methods Enzymol. 303, 179-205.) In

this DNA chip-based technology, a cDNA sequence is arrayed on a glass slide at a density of more than 1,000 genes/cm². The thus arrayed cDNA sequence is hybridized simultaneously to a two-color fluorescently labeled cDNA probe pair of different cell or tissue type RNA samples, so as to allow direct and large-scale comparative analysis of gene expression. This technology was first demonstrated by analyzing 48 Arabidopsis genes in respect of differential expression in roots and shoots (Schena, M. et al., (1995) Science 270, 467-470.) Furthermore, microarrays were used to study 1,000 clones randomly selected from a human cDNA library for identification of novel genes responding to heat shock and protein kinase C activation (Schena M. et al., (1996) Proc. Natl. Acad. Sci. USA, 93, 10614-10619.)

In another study, expression profiles of inflammatory disease-related genes were analyzed under various induction conditions by this DNA chip-based method (Heller, R. A. et al., (1997) Proc. Natl. Acad. Sci. USA, 94, 2150-2155.) Moreover, the yeast genome of more than 6,000 coding sequences has also been analyzed in respect of dynamic expression by the use of microarrays (DeRisi, J. L. et al., (1997) Science 278, 680-686.; Wodicka, L. et al., (1997) Nature Biotechnol. 15, 1359-1367.)

In plant science, however, only a few reports regarding microarray analysis have been published (Schena, M. et al., (1995) Science 270, 467-470.; Ruan, Y. et al., (1998) Plant J. 15, 821-833.; Aharoni. A. et al., (2000) Plant Cell 12, 647-661.; Reymond, P. et al., (2000) Plant Cell 12, 707-719.)

Plant growth is greatly affected by environmental stresses such as drought, high salinity and low temperature. Among these stresses, drought or water deficiency is the most severe limiting factor for plant growth and crop production. Drought stress induces various biochemical and physiological responses in plants.

Plants acquire responsivity and adaptability to these stresses to survive under

stress conditions. Recently, a number of genes responding to drought at a transcriptional level have been described (Bohnert, H.J. et al., (1995) *Plant Cell* 7, 1099-1111.; Ingram, J., and Bartels, D. (1996) *Plant Mol. Biol.* 47, 377-403.; Bray, E. A. (1997) *Trends Plant Sci.* 2, 48-54.; Shinozaki, K., and Yamaguchi-Shinozaki, K. (1997) *Plant Physiol.* 115, 327-334.; Shinozaki, K., and Yamaguchi-Shinozaki, K. (1999). Molecular responses to drought stress. Molecular responses to cold, drought, heat and salt stress in higher plants. Edited by Shinozaki, K. and Yamaguchi-Shinozaki, K. R. G. Landes Company.; Shinozaki, K., and Yamaguchi-Shinozaki, K. (2000) *Curr. Opin. Plant Biol.* 3, 217-223.)

Stress-inducible genes have been used to improve stress tolerance of plants by gene transfer (Holmberg, N., and Bulow, L. (1998) *Trends Plant Sci.* 3, 61-66.; Bajaj, S. et al., (1999) *Mol. Breed.* 5, 493-503.) It is important to analyze the functions of stress-inducible genes not only to understand the molecular mechanisms of stress tolerance and responses of higher plants, but also to improve the stress tolerance of crops by gene manipulation.

DRE/CRT (dehydration-responsive element/C-repeat sequence) has been identified as an important *cis*-acting element in drought-, high salt-, and cold stress-responsive gene expression in an ABA-independent manner (ABA refers to abscisic acid which is a kind of plant hormone and which acts as a signal transmission factor of seed dormancy and environmental stress) (Yamaguchi-Shinozaki, K., and Shinozaki, K. (1994) *Plant Cell* 6, 251-264.; Thomashow, M.F. et al., (1999) *Plant Mol. Biol.* 50, 571-599.; Shinozaki, K., and Yamaguchi-Shinozaki, K. (2000) *Curr. Opin. Plant Biol.* 3, 217-223.) Transcription factors (DREB/CBF) involved in DRE/CRT-responsive gene expression have been cloned (Stockinger, E.J. et al., (1997) *Proc. Natl. Acad. Sci. USA* 94, 1035-1040.; Liu, Q. et al., (1998) *Plant Cell* 10, 1391-1406.; Shinwari, Z.K. et al., (1998) *Biochem. Biophys. Res. Commun.* 250, 161-170.; Gilmour, S.J. et al., (1998) *Plant J.* 16, 433-443.) DREB1/CBFs are

considered to function in cold-responsive gene expression, whereas DREB2s are involved in drought-responsive gene expression. Strong tolerance to freezing stress was observed in transgenic Arabidopsis plants that overexpress CBF1 (*DREB1B*) cDNA under the control of a cauliflower mosaic virus (CaMV) 35S promoter (Jaglo-
 5 Ottosen, K.R. et al, (1998) Science 280, 104-106).

The present inventors have reported that overexpression of the DREB1A (*CBF3*) cDNA molecules in transgenic plants under the control of a CaMV 35S promoter or a stress-inducible rd29A promoter gave rise to strong constitutive
 10 expression of the stress-inducible DREB1A target genes and increased tolerance to freezing, drought, and salt stresses (Liu, Q. et al, (1998) Plant Cell 10, 1391-1406; Kasuga, M. et al, (1999) Nature Biotechnol. 17, 287-291). Furthermore, the present inventors have already identified six DREB1A target genes such as *rd9A/lti78/cor78*, *kin1*, *kin2/cor6.6*, *cor15a*, *rd17/cor47* and *erd10* (Kasuga, M. et al, (1999) Nature
 15 Biotechnol. 17, 287-291). However, it is not well clarified how overexpression of the *DREB1A* cDNA molecules in transgenic plants increases stress tolerance to freezing, drought and salt. To study the molecular mechanisms of drought and freezing tolerance, it is important to identify and analyse as many genes controlled by DREB1A as possible.

20 Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a context for the present invention. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed before the priority date of each claim
 25 of this application.

SUMMARY OF THE INVENTION

The present invention is directed to providing an environmental stress responsive promoter.

30 Through intensive studies directed toward the above, the present inventors have succeeded in identifying a novel DREB1A target gene and isolating a promoter region thereof by applying cDNA microarray analysis, thereby completing the present invention.

That is to say, the present invention is an environmental stress responsive promoter comprising DNA of the following (a), (b) or (c):

- (a) DNA consisting of any nucleotide sequence selected from SEQ ID NOS: 1 to 18;
- 5 (b) DNA consisting of a nucleotide sequence comprising a deletion, substitution or addition of one or more nucleotides relative to any nucleotide sequence selected from SEQ ID NOS: 1 to 18, and functioning as an environmental stress responsive promoter; and
- (c) DNA hybridizing under stringent conditions to DNA consisting of any
10 nucleotide sequence selected from SEQ ID NOS: 1 to 18, and functioning as an environmental stress responsive promoter.

The environmental stress is at least one selected from the group consisting of cold stress, drought stress, salt stress and high photo stress.

- Moreover, the present invention is an expression vector comprising the above
15 promoter, or the expression vector further comprising a desired gene.

Furthermore, the present invention is a transformant comprising the above expression vector.

Still further, the present invention is a transgenic plant (e.g. a plant body, plant organ, plant tissue or plant culture cell) comprising the above expression vector.

- Moreover, the present invention is a method for producing a stress-resistant
20 plant, which comprises culturing or cultivating the above transgenic plant.

According to the invention there is also provided an environmental stress responsive promoter comprising DNA of the following (a), (b) or (c):

- (a) DNA consisting of a nucleotide sequence SEQ ID NO: 6,
- 25 (b) DNA consisting of a nucleotide sequence comprising a deletion, substitution or addition of one or more nucleotides relative to nucleotide sequence SEQ ID NO: 6, and functioning as an environmental stress responsive promoter; and
- (c) DNA hybridizing under stringent conditions to DNA consisting of
30 nucleotide sequence SEQ ID NO: 6, and functioning as an environmental stress responsive promoter.

According to the invention there is also provided an expression vector comprising the promoter according to the invention.

- According to the invention there is also provided a transformant comprising the
35 expression vector according to the invention.

According to the invention there is also provided a transgenic plant comprising the expression vector according to the invention.

According to the invention there is also provided a method for producing a stress-resistant plant, which comprises culturing or cultivating the transgenic plant
5 according to the invention.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

10

BRIEF DESCRIPTION OF THE DRAWINGS



5A

Figure 1 is a photograph showing results of cDNA microarray analysis of gene expression under cold stress.

Figure 2 is a figure showing strategy for the identification of drought- or cold-inducible genes and DREB1A target genes.

Figure 3 is a photograph showing a comparison of cDNA microarray and Northern Blot analysis for new DREB1A target genes and a DREB1A gene.

Figure 4 is a figure showing results of classification of the identified drought- or cold-inducible genes into four groups on the basis of RNA gel blot and microarray analyses.

Figure 5 shows the relation between cold treatment period and expression rate regarding FL3-5A3.

Figure 6 shows the relation between dehydration treatment period and expression rate regarding FL3-5A3.

Figure 7 shows the relation between high salt treatment period and expression rate regarding FL3-5A3.

Figure 8 shows the relation between cold treatment period and expression rate regarding FL5-2H15.

Figure 9 shows the relation between dehydration treatment period and expression rate regarding FL5-2H15.

Figure 10 shows the relation between high salt treatment period and

expression rate regarding FL5-2H15.

Figure 11 shows the relation between dehydration treatment period and expression rate regarding FL5-3M24.

5

Figure 12 shows the relation between high salt treatment period and expression rate regarding FL5-3M24.

Figure 13 shows the relation between cold treatment period and expression rate regarding FL5-90.

10

Figure 14 shows the relation between cold treatment period and expression rate regarding FL5-2I22.




Figure 15 shows the relation between dehydration treatment period and expression rate regarding FL5-2I22.



Figure 16 shows the relation between high salt treatment period and expression rate regarding FL5-2I22.



Figure 17 shows the relation between dehydration treatment period and expression rate regarding FL6-55.



Figure 18 shows the relation between high salt treatment period and expression rate regarding FL6-55.

25

Figure 19 shows the relation between dehydration treatment period and expression rate regarding FL1-159.

Figure 20 shows the relation between dehydration treatment period and expression rate regarding FL5-2D23.

Figure 21 shows the relation between high salt treatment period and expression rate regarding FL5-2D23.

Figure 22 shows the relation between dehydration treatment period and expression rate regarding FL05-08-P24.

Figure 23 shows the relation between dehydration treatment period and expression rate regarding FL05-09-G08.

Figure 24 shows the relation between dehydration treatment period and expression rate regarding FL05-09-P10.

Figure 25 shows the relation between ABA treatment period and expression rate regarding FL05-09-P10.

Figure 26 shows the relation between high salt treatment period and expression rate regarding FL05-10-N02.

Figure 27 shows the relation between dehydration treatment period and expression rate regarding FL05-18-I12.

Figure 28 shows the relation between high salt treatment period and expression rate regarding FL05-18-I12.

Figure 29 shows the relation between ABA treatment period and expression rate regarding FL05-18-I12.

Figure 30 shows the relation between dehydration treatment period and expression rate regarding FL05-21-F13.

5 Figure 31 shows the relation between cold treatment period and expression rate regarding FL05-21-F13.

Figure 32 shows the relation between dehydration treatment period and expression rate regarding FL06-10-C16.

10 Figure 33 shows the relation between high salt treatment period and expression rate regarding FL06-10-C16.

Figure 34 shows the relation between ABA treatment period and expression rate regarding FL06-10-C16.

15 Figure 35 shows the relation between dehydration treatment period and expression rate regarding FL06-15-P15.

Figure 36 shows the relation between high salt treatment period and expression rate regarding FL06-15-P15.

Figure 37 shows the relation between ABA treatment period and expression rate regarding FL06-15-P15.

25 Figure 38 shows the relation between dehydration treatment period and expression rate regarding FL08-10-E21.

Figure 39 shows the relation between high salt treatment period and

expression rate regarding FL09-11-P10.

DETAILED DESCRIPTION OF THE INVENTION

5

Hereinafter, the present invention is described in detail.

10

15

20

25

30

35

Applying the biotinylated CAP trapper method (Carninci. P. et al., (1996) Genomics, 37, 327-336.), the present inventors constructed full-length cDNA libraries from Arabidopsis plants under different conditions, such as drought-treated and cold-treated plants, etc. (Seki. M. et al., (1998) Plant J. 15, 707-720.) Using about 1,300 full-length cDNA molecules and about 7,000 full-length cDNA molecules which both contained stress-inducible genes, the present inventors prepared an Arabidopsis full-length cDNA microarray for each case. In addition to these drought- and cold-inducible full-length cDNA molecules, the present inventors also prepared another cDNA microarray, using a DREB1A target gene, a transcriptional regulator for controlling expression of a stress-responsive gene. Thereafter, expression patterns of genes under drought- and cold-stresses were monitored to exhaustively analyze stress-responsive genes. As a result, novel environmental stress responsive genes, that is, 44 drought-inducible genes and 19 cold-inducible genes were isolated from a full-length cDNA microarray containing about 1,300 full-length cDNA molecules. 30 out of the 44 drought-inducible genes and 10 out of the 19 cold-inducible genes, were novel stress-inducible genes. Moreover, it was found that 12 stress-inducible genes were DREB1A target genes, and 6 of these 12 were novel genes. Furthermore, as a result of this analysis, 301 drought-inducible genes, 54 cold-inducible genes and 211 high salt-inducible genes were isolated from a cDNA microarray containing about 7,000 full-length cDNA molecules.

Thereafter, promoter regions were successfully isolated from these environmental stress responsive genes.

As stated above, a full-length cDNA microarray is a useful tool for analysis of the expression manner of *Arabidopsis thaliana* drought- and cold-stress inducible genes, and analysis of the target gene of a stress-related transcriptional regulator.

1. Isolation of promoter

The promoter of the present invention is a *cis*-element existing upstream of a gene encoding a stress-responsive protein expressed by environmental stresses such as cold-, drought- and high salt-stresses, and the *cis*-element has a function of binding to a transcriptional factor to activate transcription of a gene existing downstream. Examples of such *cis*-elements include a drought stress responsive element (DRE; dehydration-responsive element), an abscisic acid responsive element (ABRE), and a cold stress responsive element, etc. Examples of genes encoding proteins binding to these elements include a DRE binding protein 1A gene (referred to also as a DREB1A gene), a DRE binding protein 1C gene (referred to also as a DREB1C gene), a DRE binding protein 2A gene (referred to also as a DREB2A gene) and a DRE binding protein 2B gene (referred to also as a DREB2B gene), etc.

For isolation of the promoter of the present invention, first, stress responsive genes are isolated using a microarray. For preparation of a microarray, there can be used about 1,300 cDNA molecules in total, being genes isolated from *Arabidopsis* full-length cDNA libraries, RD (responsive to dehydration) genes, ERD (early responsive to dehydration) genes, *kin1* genes, *kin2* genes, *cor15a* genes, α -tubulin genes as an internal standard, and as negative controls, epsilon subunit (nAChRE) genes of a mouse acetylcholine nicotinate receptor and homologous genes of a mouse glucocorticoid receptor.

As a microarray used to isolate the promoter of the present invention, there can be used about 7,000 cDNA molecules in total, being genes isolated from

Arabidopsis full-length cDNA libraries, RD (responsive to dehydration) genes, ERD (early responsive to dehydration) genes, and PCR amplification fragments as an internal standard obtained from λ control template DNA fragments (TX803, Takara Shuzo), and as negative controls, epsilon subunit (nAChRE) genes of a mouse acetylcholine nicotinate receptor and homologous genes of a mouse glucocorticoid receptor.

A plasmid DNA extracted with a plasmid preparation device (Kurabo) is sequenced by sequence analysis, using a DNA sequencer (ABI PRISM 3700, PE Applied Biosystems, CA, USA). Based on the GenBank/EMBL database, homology detection of the obtained sequence is carried out with the BLAST program.

After poly A selection, reverse transcription is carried out to synthesize double-stranded DNA molecules, and a cDNA molecule is inserted into a vector.

The cDNA molecule inserted into a vector for preparation of cDNA libraries is amplified by PCR, using primers complementary to sequences of vectors on both sides of the cDNA molecule. Examples of such vectors include λ ZAPII and λ PS, etc.

A microarray can be prepared according to ordinary methods and so the method is not particularly limited. For example, using a gene tip microarray stamp machine, GTMASS SYSTEM (Nippon Laser & Electronics Lab.), the above obtained PCR product is loaded from a microtiter plate and spotted on a micro slide glass at regular intervals. Then, to prevent expression of non-specific signals, the slide is immersed into a blocking solution.

Examples of plant materials include plant strains obtained by destroying specific genes as well as wild type plants, and there can be used a transgenic plant, into which cDNA of DREB1A is introduced. Examples of plant varieties include

Arabidopsis thaliana, tobacco and rice, etc., and *Arabidopsis thaliana* is preferable.


Drought- and cold-stress treatments can be carried out according to a known method (Yamaguchi-Shinozaki, K., and Shinozaki, K. (1994) Plant Cell 6, 251-264.)

5

After performing the stress treatments, plant bodies (wild type plants and DREB1A overexpression transformants) are sampled, and are subjected to cryopreservation with liquid nitrogen. The wild type plants and the DREB1A overexpression transformants are used for an experiment to identify DREB1A target genes. According to a known method or using a kit, mRNA is isolated from plant bodies and purified.

10

In the presence of Cy3 dUTP or Cy5 dUTP for labeling (Amersham Pharmacia), each of the mRNA samples is subjected to reverse transcription and then used for hybridization.



After hybridization, the microarray is scanned with a scanning laser microscope. As a program for analyzing data of a microarray, Imagen Ver 2.0 (BioDiscovery) and QuantArray (GSI Lumonics), etc. can be used.



After scanning, genes of interest are isolated by preparation of a plasmid comprising the gene.



Determination of a promoter region is carried out by analysis of the nucleotide sequences of the above isolated genes, followed by the use of a gene analysis program based on the genomic information of database (GenBank/EMBL, ABRC). The isolated genes can be classified into ones having both drought- and cold-stress inductivity, ones specific for drought stress inductivity, and ones specific for cold stress inductivity (Figure 4). According to a gene analysis program, 18 types of genes

25

(FL3-5A3, FL5-2H15, FL5-3M24, FL5-90, FL5-2I22, FL6-55, FL1-159, FL5-2D23, FL05-08P-24, FL05-09-G08, FL05-09-P10, FL05-10-N02, FL05-18-I12, FL05-21-F13, FL06-10-C16, FL06-15-P15, FL08-10-E21 and FL09-11-P10) are identified from the above genes. The promoter regions of these genes are shown in SEQ ID NOS: 1 to 18, respectively.

As long as the promoter of the present invention acts as an environmental stress responsive promoter, it may be a promoter having a nucleotide sequence comprising a deletion, substitution or addition of one or more nucleotides, preferably one or several nucleotides (e.g. 1 to 10 nucleotides, preferably 1 to 5 nucleotides) relative to any nucleotide sequence selected from SEQ ID NOS: 1 to 18. Furthermore, the promoter of the present invention also includes DNA hybridizing under stringent conditions to the DNA comprising any nucleotide sequence selected from SEQ ID NOS: 1 to 18 and further acting as an environmental stress responsive promoter.

Once the nucleotide sequence of the promoter of the present invention is determined, then the promoter itself can be obtained by chemical synthesis, PCR using a cloned probe as a template, or hybridization, using as a probe, DNA fragments having the nucleotide sequence. Furthermore, a mutant of the present promoter, which has functions equivalent to those of a non-mutated promoter, can also be synthesized by a site-directed mutagenesis, etc.

To introduce a mutation into a promoter sequence, the known methods such as the Kunkel method and Gapped duplex method, or an equivalent method, can be applied. Introduction of a mutation can be carried out, for example, using a kit for introducing mutant (e.g. Mutant-K (Takara) and Mutant-G (Takara)) by a site-directed mutagenesis or using the LA PCR in vitro Mutagenesis series kit (Takara).

The term "function as an environmental stress responsive promoter" is used

herein to mean a function of binding RNA polymerase to a promoter to allow initiation of transcription, when the promoter is exposed to a specific environmental stress condition.

5 The term "environmental stress" is used generally to mean an abiotic stress such as drought stress, cold stress, high salt stress, high photo stress, etc. The term "drought" is used herein to mean a water deficient state, while the term "cold" is used herein to mean a state of being exposed to a lower temperature than the optimum living temperature of each organism variety (e.g., in the case of *Arabidopsis thaliana*, it is continuously exposed at -20 to +21°C for 1 hour to several weeks). The term "high salt" is used herein to mean a state after continuous treatment with NaCl having a concentration of 50mM to 600mM for 0.5 hours to several weeks. The term "high photo stress" is used herein to mean a state wherein a strong light greater than its photosynthetic ability is applied to a plant, and an example is application of a strong light of more than 5,000 to 10,000 lx. With regard to these environmental stresses, one kind of stress may be loaded, or several kinds of stresses may be loaded.

10 The plant promoter of the present invention includes a promoter comprising an addition of a nucleotide sequence which increases translation efficiency at the 3'-terminus of any nucleotide sequence of SEQ ID NOS: 1 to 18, or a promoter retaining a promoter activity thereof while deleting a 5'-terminus thereof.

15 Furthermore, the promoter of the present invention includes DNA which hybridizes under stringent conditions to DNA consisting of any nucleotide sequence selected from SEQ ID NOS: 1 to 18, and functions as an environmental stress responsive promoter. The term "stringent conditions" used herein means sodium concentration of 25 to 500mM, preferably 25 to 300mM, and temperature of 42°C to 68°C, preferably 42°C to 65°C. More specifically, such conditions are 5×SSC (83mM NaCl, 83mM sodium citrate) and temperature of 42°C.

2. Construction of expression vector

The expression vector of the present invention can be obtained by ligation (insertion) of the promoter of the present invention to an appropriate vector. As long as a vector into which the promoter of the present invention is inserted is capable of replicating in a host, it is not particularly limited, and examples of vectors include a plasmid, a shuttle vector and a helper plasmid, etc.

Examples of plasmid DNA include a plasmid derived from *Escherichia coli* (e.g. pBR322, pBR325, pUC118, pUC119, pUC18, pUC19 and pBluescript, etc.), a plasmid derived from *Bacillus subtilis* (e.g. pUB110 and pTP5, etc.), and a plasmid derived from yeast (e.g. YEp13 and YCp50, etc.), and examples of phage DNA include λ phage (e.g. Charon4A, Charon21A, EMBL3, EMBL4, λ gt10, λ gt11 and λ ZAP, etc.) Further, animal virus vectors such as a retrovirus and a vaccinia virus, and insect virus vectors such as a baculovirus can also be used.

To insert the promoter of the present invention into a vector, there is applied a method in which, first, the purified DNA is cleaved with suitable restriction enzymes, and, next, the obtained DNA fragment is inserted into the restriction enzyme site of a suitable vector DNA or a multi-cloning site so as to ligate to the vector.

In the present invention, in order to express a desired gene, the desired gene can be further inserted into the above expression vector. The technique involving insertion of a desired gene is the same as the method involving insertion of a promoter into a vector. A desired gene is not particularly limited, and examples of the gene include genes shown in Table 2 and the known genes other than those, etc.

In a case where the promoter of the present invention is used with a reporter gene, e.g. a GUS gene widely used in plant science, ligated to a 3'-terminus thereof, the

strength of the promoter can easily be determined by examining a GUS activity. As a reporter gene, not only a GUS gene but also luciferase and a green fluorescent protein can be used.

5

Thus, various types of vectors can be used in the present invention. Further, there can be prepared a product by connecting a desired gene of interest to the promoter of the present invention in a sense or antisense direction, and thereafter such product can be inserted into a vector called a binary vector, such as pBI101 (Clontech).

10

3. Preparation of transformant

The transformant of the present invention can be obtained by introduction of the expression vector of the present invention into a host. A host herein is not particularly limited, as long as it can express a promoter or gene of interest, a plant being preferable. Where a host is a plant, a transformant plant (a transgenic plant) can be obtained as follows.

15

A plant to be transformed in the present invention means any of an entire plant, a plant organ (e.g. a leaf, a petal, a stem, a root, a seed, etc.), a plant tissue (e.g., an epidermis, a phloem, a parenchyma, a xylem, a vascular bundle, etc.) and a plant culture cell. Examples of plants used for transformation include plants belonging to Brassicaceae, Gramineae, Solanaceae and Leguminosae, etc. (see below), but are not limited thereto.

20

25

Brassicaceae: *Arabidopsis thaliana*

Gramineae: *Nicotiana tabacum*

Solanaceae: *Zea mays*, *Oryza sativa*

Leguminosae: *Glycine max*

The above recombinant vector can be introduced into a plant by ordinary

transformation methods such as electroporation, Agrobacterium method, particle gun method, PEG, etc.

For example, where electroporation is applied, using an electroporation device equipped with a pulse controller, a vector is processed under conditions of a voltage of 500 to 1,600V, 25 to 1,000 μ F and 20 to 30msec, and a gene is introduced into a host.

Where a particle gun method is applied, a plant body, plant organ or plant tissue may be used as is, after preparation of a section, or a protoplast may be prepared. The thus prepared sample can be processed with a gene-introduction device (e.g. PDS-1000/He, Bio-Rad, etc.) Conditions for processing depend on a plant or sample, but generally, a pressure of about 1,000 to 1,800psi and a distance of 5 to 6cm are applied as processing conditions.

A gene of interest can be introduced into a plant by using a plant virus as a vector. Examples of available plant viruses include a cauliflower mosaic virus. That is, first, a virus genome is inserted into a vector derived from *Escherichia coli* to prepare a recombinant, and then the gene of interest is inserted into the virus genome. The thus modified virus genome is cut from the recombinant with restriction enzymes, and inoculated into a plant host, so that a gene of interest can be introduced therein.

When bacteria belonging to *Agrobacterium* are transfected to a plant, the bacteria introduce a portion of plasmid DNA thereof into a plant genome. In a method involving the use of Ti plasmid of *Agrobacterium*, using such a character, a gene of interest is introduced into a plant host. Among bacteria belonging to *Agrobacterium*, *Agrobacterium tumefaciens* transfects to a plant and forms therein a tumor called a crown gall, whereas *Agrobacterium rhizogenes* transfects to a plant to generate hairy roots. These phenomena originate from a cause whereby a region called a T-DNA region (a transferred DNA region) located on a plasmid called a Ti plasmid or Ri

plasmid existing in each bacterium is transferred into a plant and incorporated into a plant genome at a time of transfection.

5 By inserting DNA to be incorporated into a plant genome, into the T-DNA region of a Ti or Ri plasmid, DNA of interest can be incorporated into a plant genome, when *Agrobacterium* bacteria are transfected to a plant host.

10 Tumoral tissues, shoots and hairy roots obtained as a result of transformation can directly be used for cell culture, tissue culture or organ culture, and according to the previously known plant tissue culture method, a plant body can be regenerated by administration of a plant hormone (e.g. auxin, cytokinin, gibberellin, abscisic acid, ethylene, brassinoride, etc.) in a suitable concentration.

15 The vector of the present invention cannot only be introduced into the above-stated plant hosts, but can also be introduced into bacteria belonging to *Escherichia* such as *Escherichia coli*, *Bacillus* such as *Bacillus subtilis* and *Pseudomonas* such as *Pseudomonas putida*; yeast such as *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe*; animal cells such as COS cell and CHO cell; and insect cells such as Sf9 cell, so that a transformant can be obtained. Where a bacterium such as *Escherichia coli* or yeast is used as a host, it is preferable that the recombinant vector of the present invention is capable of self-replicating in the bacterium and, at the same time, is also comprised of the promoter of the present invention, a ribosome binding sequence, a gene of interest and a transcription termination sequence. Furthermore, it may also comprise a gene for controlling a promoter.

20

A method for introduction of a recombinant vector into bacteria is not particularly limited, as long as it is a method for introduction of DNA into bacteria. For example, a method involving the use of calcium ions and an electroporation method

can be applied.

S

Where yeast is used as a host, *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe*, etc. can be used. A method for introduction of a recombinant vector into yeast is not particularly limited, as long as it is a method for introduction of DNA into yeast, and examples of such methods include electroporation, spheroplast method, lithium phosphate method, etc.

to

Where an animal cell is used as a host, a monkey COS-7 cell, Vero, a Chinese hamster ovary cell (a CHO cell), a mouse L cell, etc. are used. Examples of methods for introduction of a recombinant vector into an animal cell include electroporation, calcium phosphate method, lipofection method, etc.

15
16
17
18
19

Where an insect cell is a host, a Sf9 cell and the like can be used. Examples of methods for introduction of a recombinant vector into an insect cell include calcium phosphate method, lipofection method, electroporation, etc.

20
21
22
23

24
25

Confirmation regarding whether a gene is incorporated into a host or not can be carried out by methods such as PCR, Southern hybridization, Northern hybridization, etc. For example, DNA is prepared from a transformant, and DNA specific primers are designed for use with PCR. PCR is carried out under the same conditions as used for preparation of the above plasmid. Thereafter, the obtained amplified product is subjected to agarose gel electrophoresis, polyacrylamide gel electrophoresis or capillary electrophoresis, so that the product is stained with ethidium bromide or a SYBR Green solution, etc. Then, it is confirmed that the amplified product was transformed, by detecting the product as a single band. Or, the amplified product can also be detected by PCR, using primers stained with fluorescent dye or the like beforehand. Furthermore, there may also be adopted a method in which the amplified product is bound to a solid phase such as a microplate and confirmed by fluorescent or enzymic

reaction, etc.

4. Production of plant

In the present invention, a transformed plant body can be regenerated from the above transformed plant cell and the like. An adaptable regeneration method is one in which transformed cells are transferred to and cultured in media with different types of hormones and concentrations to promote nucellar embryony, thereby obtaining an entire plant body. Examples of applicable media include an LS medium and an MS medium, etc.

The "method for producing a plant body" of the present invention comprises processes of, introducing a plant expression vector, into which the above plant promoter is inserted, into a host cell to obtain a transformed plant cell; regenerating a transformed plant body from the transformed plant cell; obtaining plant seeds from the resulting transformed plant body; and producing a plant body from the plant seed.

To obtain a plant seed from a transformed plant body, for example, a transformed plant body is collected from a rooting medium and transferred to a pot with water-containing soil. Then, the transformed plant body is grown at constant temperature to form flowers, thereby finally obtaining seeds. To produce a plant body from a seed, for example, after a seed formed in a transformed plant body has matured, the seed is isolated and implanted in water-containing soil, followed by growing at constant temperature under illumination. The thus bred plant becomes an environmental stress-resistant plant corresponding to the stress responsivity of a promoter introduced.

EXAMPLES

The present invention is further described in the following examples. The


examples are not intended to limit the scope of the invention.

Example 1. Isolation of promoter

1. Materials and methods


S (1) Arabidopsis cDNA clone

For preparation of a microarray, there were used about 1,300 cDNA molecules in total, which are genes isolated from Arabidopsis full-length cDNA libraries, RD (responsive to dehydration) genes, ERD (early responsive to dehydration) genes, *kin1* genes, *kin2* genes, *cor15a* genes, and α -tubulin genes as an internal standard, and as negative controls, epsilon subunit (nAChRE) genes of a mouse acetylcholine nicotinate receptor and homologous genes of a mouse glucocorticoid receptor.



Positive control: drought-inducible genes (responsive-to-dehydration genes: rd, and early responsive-to-dehydration genes: erd)

Internal standard: α -tubulin genes



Negative control: for analysis of non-specific hybridization, acetylcholine nicotinate receptor ϵ subunit (nAChRE) genes and mouse glucocorticoid receptor homolog genes, which do not substantially have homology with any given sequence in Arabidopsis database.



Furthermore, for preparation of a microarray, there were used about 7,000 cDNA molecules in total, which are genes isolated from Arabidopsis full-length cDNA libraries, RD (responsive to dehydration) genes, ERD (early responsive to dehydration) genes, and PCR amplification fragments (hereinafter referred to as PCR fragments) as an internal standard obtained from λ control template DNA fragments (TX803, Takara Shuzo), and as negative controls, epsilon subunit (nAChRE) genes of a mouse

acetylcholine nicotinate receptor and homologous genes of a mouse glucocorticoid receptor.

Positive control: drought-inducible genes (responsive-to-dehydration genes: rd, and early responsive-to-dehydration genes: erd)

Internal standard: PCR fragments

Negative control: for analysis of non-specific hybridization, acetylcholine nicotinate receptor ϵ subunit (nAChRE) genes and mouse glucocorticoid receptor homolog genes, which do not substantially have homology with any given sequence in the Arabidopsis database.

(2) Arabidopsis full-length cDNA microarray

According to the biotinylated CAP trapper method, the present inventor has constructed a full-length cDNA library from Arabidopsis plant bodies under different conditions (e.g. dehydration treatment, cold treatment and untreatment in various growth stages from budding to mature seeds). The present inventor has independently isolated each of about 1,300 and about 7,000 Arabidopsis full-length cDNA molecules from a full-length cDNA library. According to the known method (Eisen and Brown, 1999), cDNA fragments amplified by PCR were arranged on a slide glass. The present inventor has prepared both a full-length cDNA microarray containing about 1,300 Arabidopsis full-length cDNA molecules and another full-length cDNA microarray containing about 7,000 Arabidopsis full-length cDNA molecules, which comprise the genes stated below.

(3) Isolation of drought-, cold- and high salt-inducible genes using cDNA microarray


In this example, using a full-length cDNA microarray containing about 1,300 Arabidopsis full-length cDNA molecules, drought- and cold-inducible genes were

isolated. Further, using a full-length cDNA microarray containing about 7,000 Arabidopsis full-length cDNA molecules, drought-, cold- and high salt-inducible genes were isolated.

S

Both Cy3 and Cy5 fluorescent labeled probes of drought treated, cold treated and untreated plants were mixed, and the obtained mixture was hybridized to a full-length cDNA microarray containing about 1,300 Arabidopsis full-length cDNA molecules. Figure 1 shows an image of the cDNA microarray. By the double labeling of a pair of cDNA probes, wherein one mRNA sample is labeled with Cy3-dUTP and the other mRNA sample is labeled with Cy5-dUTP, simultaneous hybridization to DNA elements on a microarray becomes possible and direct assay of gene expression level under two different conditions (that is, stressed and unstressed) is facilitated. The Cy3 and Cy5 emissions of each DNA element on the hybridized microarray was scanned using two different laser channels. Thereafter, the intensity rate of the two fluorescent signals of each DNA element was determined as a relative value, and then the change of differential expression of genes was determined, which was shown as a cDNA spot on a microarray. In this example, there was used, as an internal control gene, an α -tubulin gene, the expression level of which remains almost constant under two different experimental conditions.

10



In the case of a full-length cDNA microarray containing about 7,000 Arabidopsis full-length cDNA molecules, Cy3 and Cy5 fluorescent labeled probes of each of a dehydration treated plant, a cold treated plant, a high salt -inducible gene and an unstressed plant were mixed and the obtained mixture hybridized. A PCR fragment was used as an internal control gene in this cDNA microarray.

Figure 2 shows the identification process of drought- or cold-inducible genes in a full-length cDNA microarray containing about 1,300 Arabidopsis full-length cDNA molecules. Furthermore, in the case of a full-length cDNA microarray containing

about 7,000 Arabidopsis full-length cDNA molecules also, the identification of drought-, cold- or high salt-inducible genes was performed according to the same process as shown in Figure 2.

5 1) Both mRNA molecules derived from a dehydration- or cold-treated plant and mRNA molecules derived from an unstressed wild type plant were used to prepare both Cy3- and Cy5-labeled cDNA probes. These cDNA probes were mixed, and then hybridized to a cDNA microarray. In this example, there was used, as an internal control gene, an α -tubulin gene, the expression level of which remains almost constant under two
10 different experimental conditions. A gene having more than double the expression level (drought/unstressed or cold/unstressed) of an α -tubulin gene was defined as a drought- or cold-inducible gene (Figure 2).

15 2) Both mRNA molecules derived from a 35S:DREB1A transgenic plant and mRNA molecules derived from an unstressed wild type plant were used to prepare both Cy3- and Cy5-labeled cDNA probes. These cDNA probes were mixed, and then hybridized to a cDNA microarray. In this example, there was used, as an internal control gene, an α -tubulin gene, the expression level of which remains almost constant under two
20 different experimental conditions. A gene having an expression level in a 35S:DREB1A transgenic plant of more than double its expression level in an unstressed wild type plant was defined as a DREB1A target gene (Figure 2).

25 Both mRNA molecules derived from a dehydration- or cold-treated plant and mRNA molecules derived from an unstressed wild type plant were used to prepare both Cy3- and Cy5-labeled cDNA probes. These cDNA probes were mixed, and then hybridized to a cDNA microarray. To evaluate reproducibility of microarray analysis, the same experiment was repeated five times. When the same mRNA sample was hybridized to various types of microarrays, a good correlation was observed. A gene having more than double the expression level (drought/unstressed or cold/unstressed) of

an α -tubulin gene was defined as a drought- or cold-inducible gene (Figure 2).

(4) Analysis of sequence

To perform homology detection of gene sequences, a plasmid DNA extracted with a plasmid preparation device (NA 100, Kurabo) was used for sequence analysis. A DNA sequence was determined by dye terminator cycle sequencing method, using a DNA sequencer (ABI PRISM 3700, PE Applied Biosystems, CA, USA). Based on the GenBank/EMBL database, homology detection of sequences was carried out using the BLAST program.

(5) Amplification of cDNA

As a vector for preparation of cDNA libraries, λ ZAPII (Carninci et al., 1996) was used. The cDNA inserted into a vector for libraries was amplified by PCR, using primers complementary to vector sequences located on both sides of the cDNA.

The sequences of primers are as follows:

FL forward 1224: 5'-CGCCAGGGTTTCCCAGTCACGA (SEQ ID NO: 19)

FL reverse 1233: 5'-AGCGGATAACAATTCACACAGGA (SEQ ID NO: 20)

As a template, a plasmid (1 to 2ng) was added to 100 μ l of PCR mixture (0.25mM dNTP, 0.2 μ M PCR primers, 1 X Ex Taq buffer, 1.25 U Ex Taq polymerase (Takara Shuzo)). PCR was performed under the following conditions: initial reaction at 94°C for 3 minutes, 35 cycles of 95°C for 1 minute, 60°C for 30 seconds and 72°C for 3 minutes, and the final reaction at 72°C for 3 minutes. After precipitation of a PCR product with ethanol, the precipitate was dissolved into 25 μ l of 3 X SSC. 0.7% agarose gel electrophoresis was performed to confirm the quality of the obtained DNA and amplification efficiency of PCR.

(6) Production of cDNA microarray

Using a gene tip microarray stamp machine, GTMASS SYSTEM (Nippon Laser & Electronics Lab.), 0.5 μ l of PCR product (100 to 500ng/ml) was loaded from a 384-well microtiter plate, and 5nl each of the product was spotted on 6 micro slide glasses (S7444, Matsunami) coated with poly-L-lysine at a space of 280 μ m. To spot an equivalent amount of DNA, after printing, slides were wetted in a beaker containing hot distilled water and then dried at 100°C for 3 seconds. Thereafter, the slides were placed on a slide rack, and the rack was placed into a glass chamber. Then, a blocking solution (containing 15ml of 1M sodium borate salt (pH8.0), 5.5g of succinic anhydrous compound (Wako), and 335ml of 1-methyl-2-pyrrolidone (Wako)) was poured into the glass chamber. After shaking the glass chamber containing the slides rack up and down 5 times, it was further shaken gently for 15 minutes. Thereafter, the slide rack was transferred to a glass chamber containing boiling water and shaken 5 times, followed by being left at rest for 2 minutes. Then, the slide rack was transferred to a glass chamber containing 95% ethanol and shaken 5 times, followed by centrifugation (800rpm) for 30 minutes.

(7) Plant materials and isolation of RNA

As plant materials, there were used both a wild type *Arabidopsis thaliana* plant body which was seeded on an agar medium and grown for 3 weeks (Yamaguchi-Shinozaki and Shinozaki, 1994), and an *Arabidopsis thaliana* (Colombian species) plant body, into which DREB1A cDNA (Kasuga et al., 1999) connected to 35S promoter of a cauliflower mosaic virus was introduced. Drought- and cold-stress treatments were performed by the method of Yamaguchi-Shinozaki and Shinozaki (1994). That is to say, a plant body pulled out of an agar medium was placed on a filter and then dehydration treatment was carried out under conditions of 22°C and 60% relative humidity. Cold treatment was carried out by transferring the plant grown at 22°C into conditions of 4°C. High salt-stress treatment was carried out by water-culturing in a solution containing 250mM NaCl.

After wild type plant bodies were exposed to stress-treatment for 2 or 10 hours, they were subjected to sampling, and then subjected to cryopreservation with liquid nitrogen. Both wild type and DREB1A overexpression type transformants, which were cultured in an agar medium without kanamycin, were used for an experiment for identification of a DREB1A target gene. Stress treatment was not performed for DREB1A overexpression type transformants. The total RNA was isolated from plant bodies using ISOGEN (Nippon gene, Tokyo, Japan), and then mRNA was isolated and purified using an Oligotex-dT30 mRNA purification kit (Takara, Tokyo, Japan).

(8) Fluorescent labeling of probe

In the presence of Cy3 dUTP or Cy5 dUTP (Amersham Pharmacia), each of the mRNA samples was reverse transcribed. The composition of the reverse transcription buffer (30 μ l) is as follows.

poly (A) ⁺ RNA with 6 μ g oligo (dT) 18-mer	1 μ g
10mM DTT	
500 μ M dATP, dCTP and dGTP	
200 μ M dTTP	
100 μ M Cy3 dUTP or Cy5 dUTP	
400 units of SuperScript II reverse transcription enzymes (Life Technologies)	
<u>1 X Superscript first strand buffer (Life Technologies)</u>	
	30 μ l in total

After reaction at 42°C for 1 hour, two samples (one sample labeled with Cy3 and the other sample labeled with Cy5) were mixed, and 15 μ l of 0.1M NaOH and 1.5 μ l of 20mM EDTA were added thereto and the mixture was treated at 70°C for 10 minutes. Then, 15 μ l of 0.1M HCl was further added thereto, and the sample then

transferred to a Micro con 30 micro concentrator (Amicon). 400 μ l of TE buffer was added, followed by centrifugation so that the amount of the buffer became 10 to 20 μ l, and then an effluent was thrown away. After 400 μ l of TE buffer and 20 μ l of 1mg/ml human Cot-1 DNA (Gibco BRL) were added thereto, the mixture was subjected to centrifugation again. The labeled samples were collected by centrifugation, and several μ l of distilled water were added thereto. To the obtained probes, 2 μ l of 10 μ g/ μ l yeast tRNA, 2 μ l of 1 μ g/ μ l pd(A)₁₂₋₁₈ (Amersham Pharmacia), 3.4ml of 20 X SSC and 0.6 μ l of 10% SDS were added. Furthermore, the samples were denatured at 100°C for 1 minute and placed at room temperature for 30 minutes, and then used for hybridization.

(9) Microarray hybridization and scanning

Using benchtop micro centrifuge, a probe was subjected to high-speed centrifugation for 1 minute. To avoid generation of bubbles, the probe was placed in the center of an array, and a cover slip was placed thereon. Four drops of 5 μ l of 3 X SSC were dropped on a slide glass and a chamber was kept at suitable humidity to prevent dehydration of the probe during hybridization. The slide glass was placed into a cassette for hybridization (THC-1, BM machine) and sealed, followed by treatment at 65°C for 12 to 16 hours. The slide glass was taken out of the cassette and placed on a cassette rack, and a cover slip was carefully removed therefrom in solution 1 (2 X SSC, 0.1% SDS). Thereafter, the rack was shaken to wash, and transferred into solution 2 (1 X SSC) to wash for 2 minutes. Then, the rack was further transferred into solution 3 (0.2 X SSC) and left for 2 minutes, and then subjected to centrifugation (800rpm, 1 min) for drying.

Using a scanning laser microscope (ScanArray4000; GSI Lumonics, Watertown, MA), a microarray was scanned in a resolution of 10 μ m per pixel. As a program for analyzing the microarray data, Imogene Ver 2.0 (BioDiscovery) and Quant Array (GSI Lumonics) were used.

(10) Northern analysis

Using total RNA, Northern analysis was carried out (Yamaguchi-Shinozaki and Shinozaki, 1994). DNA fragments isolated from *Arabidopsis thaliana* full-length cDNA libraries by PCR were used as probes for Northern hybridization.

(11) Determination of promoter region

Based on the genomic information of *Arabidopsis thaliana* in database (GenBank/EMBL, ABRC), a promoter region was analyzed using the BLAST program for gene analysis.

2. Results

(1) Stress-inducible gene

Fluorescent-labeled cDNA was prepared from mRNA, which were isolated from unstressed *Arabidopsis thaliana* plants, by reverse transcription in the presence of Cy5-dUTP. From cold-treated plants (2 hours), the second probes labeled with Cy3-dUTP were prepared. Both types of probes were simultaneously hybridized to a cDNA microarray comprising about 1,300 *Arabidopsis thaliana* cDNA clones, and then a pseudo color image was created (Figure 1).

Genes induced and inhibited by cold stress are represented by a red signal and a green signal, respectively. Genes which expressed at almost an equivalent level in both treatments are represented by a yellow signal. The strength of each spot corresponds to the absolute value of the expression level of each gene. It is shown that a cold-inducible gene (rd29A) is represented by a red signal whereas an α -tubulin gene (an internal control) is represented by a yellow signal.

By means of cDNA microarray analysis, the total 44 drought-inducible genes

5

Figure 1. The four types of stimuli used in the experiment. The stimuli were presented in a 2 × 2 grid. The top row shows the two types of stimuli used in the first experiment (Experiment 1) and the bottom row shows the two types of stimuli used in the second experiment (Experiment 2). The stimuli were presented in a 2 × 2 grid.

	Number of genes
Drought-inducible gene	44
New drought-inducible gene	30
Cold-inducible gene	19
New cold-inducible gene	10
DREB1A target gene	12
New DREB1A target gene	6

Table 2. Drought- and Cold-inducible Genes, and DREB1A Target Genes Identified by cDNA Microarray Analysis

Gene	Accession	Coded Protein/Other Features	Drought (2hr)		Cold (2hr)		35S:DREB1A	
			Ratio	New or Reported	Ratio	New or Reported	Ratio	New or Reported
rd29A	D13044	Hydrophilic protein	6.4	Reported	5.1	Reported	7.9	Reported
cor15a	U01377	-	5.0	Reported	3.4	Reported	8.1	Reported
kin2	X55053	-	5.8	Reported	2.9	Reported	4.9	Reported
erd10	D17714	Group II LEA protein	8.0	Reported	4.6	Reported	3.5	Reported
kin1	X51474	-	2.9	Reported	2.0	Reported	3.4	Reported
rd17	AB004872	Group II LEA protein	6.4	Reported	4.6	Reported	4.6	Reported
rd20	-	Ca-binding EF hand protein	5.1	Reported	n.d.	-	n.d.	-
erd7	-	-	3.8	Reported	2.3	Reported	n.d.	-
erd4	-	Membrane protein	2.6	Reported	2.2	Reported	2.5	New
erd3	-	-	2.6	Reported	n.d.	-	n.d.	-
FL3-519	D17715	Group II LEA protein	3.5	Reported	1.5	-	1.2	-
FL3-3A1	D13042	Thiol protease	2.8	Reported	1.9	-	1.5	-
FL5-1F23	D32138	Δ^1 -pyrroline-5-carboxylate synthetase (AtP5CS)	2.8	Reported	1.5	-	n.d.	-
FL2-1F6	D01113	Unidentified seed protein	2.2	Reported	1.1	-	0.5	-
FL3-5A3	AC006438	Putative cold acclimation protein	6.2	New	2.3	New	3.4	New
FL6-55	X91919	LEA 76 type 1 protein	2.9	New	n.d.	-	n.d.	-
FL5-77	AF121355	Peroxisomal protein TRX1	2.2	New	1.9	-	3.0	New
FL3-5J1	AF057137	Gamma tonoplast intrinsic protein 2 (TIP2)	2.0	New	1.2	-	1.3	-
FL5-1N11	M80567	Non-specific lipid transfer protein (LTP1)	2.7	New	1.3	-	1.2	-
FL5-95	-	Rice glyceraldehyde 3-phosphate dehydrogenase 1 homolog	2.3	New	2.8	New	n.d.	-
FL5-2H15	T45998(EST)	-	2.1	New	n.d.	-	1.4	-
FL5-2C24	AC005770	Putative water channel protein	2.4	New	n.d.	-	1.4	-
FL5-2G21	AF034255	Reversibly glycosylated polypeptide-3 (RGP)	2.1	New	n.d.	-	1.3	-
FL5-1A9	-	Nodulin-like protein homolog	2.9	New	2.1	New	0.8	-
FL5-94	X58107	Enolase	2.0	New	1.8	-	2.3	New
FL5-3J4	-	Heat shock protein dnaJ homolog	2.8	New	1.4	-	n.d.	-
FL5-3M24	-	LEA protein SAG21 homolog	2.3	New	2.2	New	1.0	-
FL5-1C3	H37392(EST)	-	3.0	New	n.d.	-	n.d.	-
FL5-2I23	D14442	Ascorbate peroxidase	2.1	New	1.1	-	1.0	-
FL1-159	AB015098	HVA22 homolog	3.7	New	3.8	New	1.9	-
FL3-27	-	Cysteine proteinase inhibitor homolog	2.2	New	n.d.	-	2.2	New
FL5-2I22	X80342	DC 1.2 homolog	2.6	New	2.9	New	2.1	New
FL5-1C20	-	Major latex protein type I homolog	2.0	New	1.4	-	1.8	-
FL2-1H6	-	<i>Brassica napus</i> jasmonate-inducible protein homolog	2.4	New	1.3	-	0.9	-
FL5-2E17	-	<i>Brassica napus</i> beta-glucosidase homolog	2.3	New	1.1	-	1.0	-
FL3-3B1	AC006403	Hypothetical protein	2.7	New	1.4	-	1.1	-
FL5-3E18	M80567	Aquaporin homolog	2.0	New	1.1	-	0.9	-
FL5-3A15	X94248	Ferritin	2.8	New	2.1	New	0.9	-
FL2-56	AF104330	Glycine-rich protein 3 short isoform (GRP35)	2.6	New	1.4	-	1.6	-
FL5-2D23	-	T20517 (EST) homolog	2.6	New	1.3	-	n.d.	-
FL3-2C6	Z35474	Thioredoxin	2.3	New	1.0	-	1.9	-
FL5-1P10	AC004044	Putative photosystem I reaction center subunit II precursor	2.1	New	1.4	-	1.2	-
FL2-5G7	U43147	Catalase 3 (CAT3)	2.4	New	1.2	-	1.6	-
FL2-1C1	Z9734	Cysteine proteinase homolog	3.0	New	1.0	-	1.7	-
DREB1A	AB007787	EREBP/AP2 protein	n.d.	-	6.3	Reported	5.8	-
FL2-5A4	AB010259	DEAD box ATPase/RNA helicase protein (DRH1)	n.d.	-	2.1	New	n.d.	-
FL5-90	AJ250341	β -amylase	n.d.	-	4.4	New	1.2	-
FL5-3P12	D63510	EXGT-A2	1.2	-	3.2	New	0.8	-

In Table 2, genes which are drought- and cold-inducible, and DREB1A target genes (35S:DREB1A) are rd29, cor15A, kin2, erd10, kin1, rd17, erd4, FL3-5A3, FL5-77, FL5-94, FL3-27 and FL5-2I22. Genes which are drought- and cold inducible but are not DREB1A target genes, are FL5-2024, FL5-1A9, FL5-3M24 and FL5-3A15.

Specifically drought-inducible genes are rd20, FL6-55, FL5-3J4, FL2-56 and FL5-2D23. Specifically cold-inducible genes are DREB1A and FL5-90. The results of classification of these genes are shown in Figure 4.

Moreover, in Table 2, the “Coded Protein/Other Features” column shows putative functions of a gene product, which are predicted from sequence homology.

The “Ratio” in the “Drought” column is obtained by the following equation (FI: fluorescence intensity):

$$\text{Ratio} = \left[\frac{\text{(the FI of each cDNA under drought condition)}}{\text{(the FI of each cDNA under unstressed condition)}} \right] \div \left[\frac{\text{(the FI of } \alpha\text{-tubulin under drought condition)}}{\text{(the FI of } \alpha\text{-tubulin under unstressed condition)}} \right]$$

The “Ratio” in the “Cold” column is obtained by the following equation (FI: fluorescence intensity):

$$\text{Ratio} = \left[\frac{\text{(the FI of each cDNA under cold condition)}}{\text{(the FI of each cDNA under unstressed condition)}} \right] \div \left[\frac{\text{(the FI of } \alpha\text{-tubulin under cold condition)}}{\text{(the FI of } \alpha\text{-tubulin under unstressed condition)}} \right]$$

The “Ratio” in the “35S:DREB1A” column is obtained by the following equation (FI: fluorescence intensity):

$$\text{Ratio} = \left[\frac{\text{(the FI of each cDNA of 35S:DREB1A plants)}}{\text{(the FI of each cDNA of wild type plants)}} \right] \div \left[\frac{\text{(the FI of } \alpha\text{-tubulin of 35S:DREB1A plants)}}{\text{(the FI of } \alpha\text{-tubulin of wild type plants)}} \right]$$

With regard to the term “New or Reported” in the “Drought” column, where a

gene has not been reported as a drought-inducible gene, it is shown as "New", and where it has been reported as a drought-inducible gene, it is shown as "Reported". The same applies for a cold-inducible gene in the "Cold" column and a DREB1A target gene in the "35S:DREB1A" column.

5

Among the genes described in Table 2, 14 genes (cor15A, kin1, kin2, rd17, rd19A, rd20, rd22, rd29A, erd3, erd4, erd7, erd10, erd14, AtP5CS) have previously been reported as drought-inducible genes (Bohnert, H.J. et al., (1995) Plant Cell 7, 1099-1111.; Ingram, J., and Bartels, D. (1996) Plant Mol. Biol. 47, 377-403.; Bray, E. A. (1997) Trends Plant Sci. 2, 48-54.; Shinozaki, K., and Yamaguchi-Shinozaki, K. (1997) Plant Physiol. 115, 327-334.; Shinozaki, K., and Yamaguchi-Shinozaki, K. (1999). Molecular responses to drought stress. Molecular responses to cold, drought, heat and salt stress in higher plants. Edited by Shinozaki, K. and Yamaguchi-Shinozaki, K., R. G. Landes Company.; Shinozaki, K., and Yamaguchi-Shinozaki, K. (2000) Curr. Opin. Plant Biol. 3, 217-223.; Tajiri, T. et al., (1999) Plant Cell Physiol. 40, 119-123.; Takahashi, S. et al., (2000) Plant Cell Physiol. 41, 898-903.)

to

75

80

85

90

From these results, it is shown that the cDNA microarray system of the present inventors has functioned appropriately to find stress-inducible genes. In the remaining 30 new drought-inducible genes, there were found cDNA molecules (FL3-5A3, FL6-55, FL5-1N11, FL5-2024, FL5-2H15 and FL1-159) which show sequence identity with putative cold acclimation protein (Accession No. AC006438), LEA 76 type 1 protein (Accession No. X91919), non-specific lipid transfer protein (LTP1; Accession No. M80567) putative water channel protein (Accession No. AC005770), T45998 EST, and HVA22 homolog (Accession No. AB015098).

Moreover, the total 19 cold-inducible genes have been identified by cDNA microarray analysis (Tables 1 and 2). Nine of these genes have been reported as cold-inducible genes, rd29A, cor15a, kin1, kin2, rd17, erd10, erd7 and erd4 (Kiyosue et

al., 1994; Shinozaki and Yamaguchi-Shinozaki, 1997, 1999, 2000; Taji et al., 1999; Thomashow, 1999) and a DREB1A gene (Lui et al., 1998).

Likewise, in the remaining new cold-inducible genes, there were found cDNA molecules (FL3-5A3, FL5-3A15, FL5-3P12, FL5-90, FL5-2I22 and FL1-159) which show sequence identity with putative cold acclimation protein (Accession No. AC006438), ferritin (Accession No. X94248), EXGT-A2 (Accession No. D63510), β -amylase (Accession No. AJ250341), DC 1.2 homolog (Accession No. X80342), and HVA22 homolog (Accession No. AB015098), and also found cDNA molecules (FL5-1A9, FL5-95 and FL5-3M24) which show sequence similarity with Nodulin-like protein (Accession No. CAA22576), rice glyoxalase I (Accession No. AB017042) and LEA protein homolog (SAG21; Accession No. AF053065).

Furthermore, the present inventors have identified stress-inducible genes controlled by a DREB1A transcription factor, using a full-length cDNA microarray. Figure 2 shows a procedure for identification of DREB1A target genes. The mRNA prepared from a transgenic *Arabidopsis thaliana* plant (a 35S:DREB1A transgenic plant), which overexpresses DREB1A cDNA under the control of a CaMV 35S promoter, and the mRNA prepared from a wild type control plant were used to prepare Cy3-labeled and Cy5-labeled cDNA probes, respectively. These cDNA probes were mixed and then hybridized to a cDNA microarray. A gene having more than a two-fold greater expression level in a 35S:DREB1A transgenic plant than in a wild type control plant was defined as a DREB1A target gene.

The total 12 DREB1A target genes have been identified by cDNA microarray analysis (Tables 1 and 2). Six of these genes have been reported as DREB1A target genes, rd29A/cor78, cor15a, kin1, kin2, rd17/cor47 and erd10 (Kasuga et al., 1999). Likewise, in the remaining 6 new DREB1A target genes, there were found cDNA molecules (FL3-5A3, FL5-2I22, FL5-94 and FL5-77) which show sequence identity

with putative cold acclimation protein (Accession No. AC006438), DC 1.2 homolog (Accession No. X80342), enolase (Accession No. X58107) and peroxiredoxin TPX1 (Accession No. AF121355), and also found a cDNA molecule (FL3-27) showing sequence similarity with a cowpea cysteine proteinase inhibitor (Accession No. Z21954) and erd4 cDNA (Kiyosue et al., 1994; Taji et al., 1999).

The identified drought- or cold-inducible genes were classified into the following 3 groups (Figure 4):

- 1) Drought- and cold-inducible genes
- 2) Specifically drought-inducible genes
- 3) Specifically cold-inducible genes

With regard to the following 21 genes, it was difficult to classify them into the above 3 groups and so they did not undergo such a classification:

erd3, FL3-519, FL3-3A1, FL5-1F23, FL2-1F6, FL3-5J1, FL5-1N11, FL5-2H15, FL5-2G21, FL5-2I23, FL5-1C20, FL2-1H6, FL5-2E17, FL3-3B1, FL5-3E18, FL3-2C6, FL5-1P10, FL2-5G7, FL2-1C1, FL2-5A4, and FL5-3P12

As a result, the identified genes were classified into 20 drought- and cold-inducible genes, 5 specifically drought-inducible genes and 2 specifically cold-inducible genes. Thereafter, the drought- and cold-inducible genes were classified into two groups:

- 1) DREB1A target genes
- 2) Genes other than DREB1A target genes

Thus, 16 drought- and cold-inducible genes were classified into 12 DREB1A target genes and 4 genes other than DREB1A target genes.

(2) RNA gel blot analysis

In cDNA gel blot analysis, it is important to extract appropriate data with minimum effort. The present inventors evaluated the effectiveness of cDNA microarray analysis by the following method.

First, 80 genes having more than double the expression ratio (drought 2 hours/unstressed) of α -tubulin were identified. The 80 putative drought-inducible genes were subjected to Northern blot analysis, 44 of which were identified as actual genes. The disparity between the results from microarray analysis and those from Northern blot analysis was caused by (1) low expression of genes, (2) high background, (3) dusts or scratches on a cDNA spot, and (4) a bad cDNA probe with a low specific activity. Accordingly, the above experimental data were marked and a half thereof was excluded from the following analysis. After the data processing, 44 drought-inducible genes, 19 cold-inducible genes and 12 DREB1A target genes were finally identified based on cDNA microarray analysis. Thereafter, RNA gel blot analysis was carried out to confirm the obtained results using a cDNA microarray. The result of expression data obtained by microarray analysis that 44 drought-inducible genes, 19 cold-inducible genes and 12 DREB1A target genes were identified, was consistent with the result obtained by Northern blot analysis.

Figure 3 shows a comparison between the result of microarray analysis and that of Northern blot analysis in respect of 6 new DREB1A target genes (FL3-5A3, FL3-27, FL5-2I22, FL5-94, FL5-77 and erd4). All of the 6 genes were induced by drought- and cold-treatments and overexpressed in 35S:DREB1A plants under unstressed conditions.

Samples derived from drought-treated wild type plants (dehydration for 2 hours or 10 hours (left on a filter paper)), cold-treated wild type plants (cooling at 4°C for 2 hours or 10 hours), or untreated 35S:DREB1A transgenic plants (35S:DREB1A control) were subjected to fluorescent labeling with Cy3-dUTP, whereas samples

derived from untreated wild type plants (control) were subjected to fluorescent labeling with Cy5-dUTP. These samples were hybridized to a cDNA microarray followed by scanning to calculate a relative expression ratio and shown in a figure (Figure 3). The figure shows Northern blot analysis images of drought- and cold-treated wild type plants and 35S:DREB1A transgenic plants. The full-length cDNA sequences of two DREB1A target genes (FL3-5A3 and FL3-27) are independently registered with GenBank, EMBL and DDBJ database under Accession Nos. AB044404 and AB044405, respectively.

Likewise, using a full-length cDNA microarray containing about 7,000 Arabidopsis full-length cDNA molecules, 301 drought-inducible genes, 54 cold-inducible genes and 211 high salt stress-inducible genes were isolated.

(3) Identification of promoter region

As a result of identification of promoter regions, there were obtained the promoter regions of 8 types of genes (FL3-5A3, FL5-2H15, FL5-3M24, FL5-90, FL5-2I22, FL6-55, FL1-159 and FL5-2D23) obtained in a full-length cDNA microarray containing about 1,300 Arabidopsis full-length cDNA molecules. The sequences of these promoters are shown in SEQ ID NOS: 1 to 8.

Gene Name	Promoter Region Sequence
FL3-5A3	SEQ ID NO: 1
FL5-2H15	SEQ ID NO: 2
FL5-3M24	SEQ ID NO: 3
FL5-90	SEQ ID NO: 4
FL5-2I22	SEQ ID NO: 5
FL6-55	SEQ ID NO: 6
FL1-159	SEQ ID NO: 7
FL5-2D23	SEQ ID NO: 8

As a result of the identification of promoter regions, there were obtained the promoter regions of 10 types of genes (FL05-08-P24, FL05-09-G08, FL05-09-P10, FL05-10-N02, FL05-18-I12, FL05-21-F13, FL06-10-C16, FL06-15-P15, FL08-10-E21 and FL09-11-P10) obtained in a full-length cDNA microarray containing about 7,000 Arabidopsis full-length cDNA molecules. The sequences of these promoters are shown in SEQ ID NOS: 9 to 18.

Gene Name	Promoter Region Sequence
FL05-08-P24	SEQ ID NO: 9
FL05-09-G08	SEQ ID NO: 10
FL05-09-P10	SEQ ID NO: 11
FL05-10-N02	SEQ ID NO: 12
FL05-18-I12	SEQ ID NO: 13
FL05-21-F13	SEQ ID NO: 14
FL06-10-C16	SEQ ID NO: 15
FL06-15-P15	SEQ ID NO: 16
FL08-10-E21	SEQ ID NO: 17
FL09-11-P10	SEQ ID NO: 18

It is known that a conserved sequence "PyACGTG (G or T)C" (wherein Py represents a pyrimidine base, that is, C or T) acts as an ABA-responsive element (ABRE) in many ABA-responsive genes (Ingram, J., and Bartels, D. (1996) *Plant Mol. Biol.* 47, 377-403.; Bray, E. A. (1997) *Trends Plant Sci.* 2, 48-54.; Shinozaki, K., and Yamaguchi-Shinozaki, K. (1999). *Molecular responses to drought stress. Molecular responses to cold, drought, heat and salt stress in higher plants.* Edited by Shinozaki, K. and Yamaguchi-Shinozaki, K., R. G. Landes Company.) Furthermore, it is also known that a conserved sequence consisting of 9 nucleotides "TACCGACAT" (a dehydration-responsive element: DRE) is essential to the induction control of rd29A

expression under drought-, cold- and high salt-stress conditions, but does not function as an ABA-responsive element (ABRE) (Yamaguchi-Shinozaki, K., and Shinozaki, K. (1994) Plant Cell 6, 251-264). It is also known that CRT or LTRE, which is DRE or DRE-related core motifs (CCCAG), exists in the regions of drought- and cold-inducible genes (e.g. kin1, kin2, rd17/cor47 and cor15a) (see Table 3, Baker, S. S. et al., (1994) Plant Mol. Biol. 24, 701-713.; Wang, H. et al., (1995) Plant Mol. Biol. 28, 605-617.; Iwasaki, T. et al., (1997) Plant Physiol. 115, 1287.)

Table 3. ABRE, DRE, and CCGAC Core Sequences Observed in the Promoter Regions of the DREB1A Target Genes Identified by cDNA Microarray Analysis a)

Gene	ABRE (PyACGTG(T/G)C)	DRE (TACCGACAT)	CCGAC Core Motif (CCGAC)
rd29A	TACGTGTC(-45 to -38) ^{b)}	TACCGACAT(-148 to -140) TACCGACAT(-206 to -198)	AGCCGACAC(-111 to -103) GACCGACTA(-256 to -248)
cor15A	CACGTGGC(-132 to -125)	-	GGCCGACCT(-184 to -176) GGCCGACAT(-361 to -353) AACCGACAA(-416 to -424)
kin1	-	TACCGACAT(-120 to -112)	ATCCGACAT(-720 to -712)
kin2	CACGTGGC(-68 to -61)	TACCGACAT(-127 to -119)	CCCGGACGC(-403 to -395)
rd17	TACGTGTC(-920 to -913)	-	TACCGACTT(-162 to -154) AGCCGACCA(-967 to -959) GACCGACAT(-997 to -989)
erd10	CACGTGGC(-838 to -831)	-	GACCGACGT(-198 to -190) ^{c)} GACCGACCG(-202 to -194) ^{c)} CACCGACCG(-206 to -198) ^{c)} GACCGACAT(-999 to -991)
FL3-5A3	CACGTGGC(-74 to -67)	TACCGACAT(-415 to -407)	TGCCGACAT(-806 to -798)
FL3-27	-	TACCGACAT(-89 to -81)	-
FL5-2122	-	-	TACCGACTC(-191 to -183) TACCGACTA(-266 to -258) TGCCGACCT(-418 to -410) ACCCGACTA(-695 to -703) GACCGACGT(-786 to -778)
FL5-77	-	-	CCCGGACTA(-315 to -307)
FL5-94	-	-	TACCGACTA(-190 to -198) TTCCGACTA(-260 to -268) ATCCGACGC(-630 to -622)

In Table 3, a), b) and c) are defined as follows:

- a): ABRE, DRE and CCGAC core sequences refer to sequences observed at 1,000bp upstream of the 5'-terminus of the longest cDNA isolated.
- b): Figures in parentheses represent nucleotides initiating at the 5'-termini of the

isolated cDNA. A minus sign means that a nucleotide exists upstream of the 5'-terminus of a putative transcription initiation site.

c): Each of these sequences consisting of 9 nucleotides comprises a CCGAC core motif, and the sequences overlap with one another.

5

The present inventors identified 12 DREB1A target genes by cDNA microarray analysis. A DRE sequence of 9bp is observed in the promoter regions of genes corresponding to the cDNA molecules of FL3-5A3 and FL3-27 (Table 3). A core, sequence "CCGAC" is observed in the promoter regions of genes corresponding to the cDNA molecules of FL3-5A3, FL5-2I22, FL5-77 and FL5-94 (Table 3). Almost all of the drought- and cold-inducible genes are DREB1A target genes, each of which contains a DRE/CRT *cis*-acting element in a promoter thereof (Table 3 and Figure 4). An ABRE sequence ("PyACGTG(G or T)C") was observed in 6 promoter regions among the identified 12 DREB1A target genes (Table 3). This shows that many drought- and cold-inducible genes are controlled by both ABA-dependent and ABA-independent routes. However, some drought- and cold-inducible genes (FL5-3M24, FL5-3A15, FL5-1A9 and FL5-2024) did not increase in 35S:DREB1A transgenic plants (Figure 4). This shows that these genes are not DREB1A target genes. A core sequence "CCGAC" was not observed in a region 2,000 bp upstream of the 5'-terminus of cDNA of FL5-3M24. This result shows a possibility that a new *cis*-acting element associated with expression of drought- and cold-inducible genes exists in the promoter region of a FL5-3M24 gene.

(4) Relation between various stress treatment periods of time and expression ratio

With regard to 18 types of stress-inducible genes isolated as above, the results of analysis of the relation between various stress treatment periods of time and expression ratio are shown in the following Figures 5 to 39.

Gene Name	Stress Result (Figures showing the results)

5	FL3-5A3	Cold (Fig. 5), Drought (Fig. 6) and High Salt (Fig. 7)
	FL5-2H15	Cold (Fig. 8), Drought (Fig. 9) and High Salt (Fig. 10)
	FL5-3M24	Drought (Fig. 11) and High Salt (Fig. 12)
	FL5-90	Cold (Fig. 13)
	FL5-2I22	Cold (Fig. 14), Drought (Fig. 15) and High Salt (Fig. 16)
	FL6-55	Drought (Fig. 17) and High Salt (Fig. 18)
	FL1-159	Drought (Fig. 19)
	FL5-2D23	Drought (Fig. 20) and High Salt (Fig. 21)
	FL05-08-P24	Drought (Fig. 22)
	FL05-09-G08	Drought (Fig. 23)
	FL05-09-P10	Drought (Fig. 24) and ABA (Fig. 25)
	FL05-10-N02	High Salt (Fig. 26)
	FL05-18-I12	Drought (Fig. 27), High Salt (Fig. 28) and ABA (Fig. 29)
	FL05-21-F13	Drought (Fig. 30) and Cold (Fig. 31)
	FL06-10-C16	Drought (Fig. 32), High Salt (Fig. 33) and ABA (Fig. 34)
10	FL06-15-P15	Drought (Fig. 35), High Salt (Fig. 36) and ABA (Fig. 37)
	FL08-10-E21	Drought (Fig. 38)
	FL-9-11-P10	High Salt (Fig. 39)

As shown in Figures 5 to 39, the stress-inducible genes isolated by the method of the present invention are different profiles, but are expressions induced by the addition of various types of stresses.

Effect of the Invention

According to the present invention, a stress responsive promoter is provided. The promoter of the present invention is useful in that it can be used for molecular breeding of environmental stress-resistant plants.

EDITORIAL NOTE - No. 91431/01

Pages 43 and 44 are claims pages, they follow the sequence listing, pages
45 to 72

SEQUENCE LISTING

<110> RIKEN

TOYOTA MOTOR CORPORATION

<120> A STRESS-RESPONSIVE PROMOTER

<130> xxxxx

<150> JP 2001-309984

<151> 2001-10-05

<150> JP 2000-356652

<151> 2000-11-22

PatentIn Ver. 2.0

<160> 20

<170> PatentIn Ver. 2.0

1

<210> 1

<211> 2072

DNA

<212> DNA

<213> Arabidopsis thaliana

1

<400> 1

tgggcctttg atccacatga cgaatcatata gtcttttct tcaacacaaa acacttgaag 60
tgtaatttat caactaaact ctattccagc aaaattatit caaaattaga tataatgggag 120
aatittttaca ggccattcta gattttcatt aacgctgtga ttatccglt aataagtacg 180
attaagagia aaacgacgtc gtttgcaatt gaaacaaaaa tccccaattc agaaataaga 240

aaccctagat cctcaaatct ctctctcttt tctctctcga attcactcct gtattttgaa 300
atcaaagtga atlgcaaggc acaaagttca tgcittaaag cactgcaaat ggcaagtc 360
tgalittat ggcatttttc ctgtttttgt ttcaattgca aacgacgtcg ttttgccttt 420
aaacggcttc attaacggat aaatcacatc ttctaattgc tctgtttatt cagttgatgt 480
tgtactaat ttggctctgt caacgaaagt tagtgatttt atctgaagtc aacacgtaat 540
tgaigatttt aatcacaaag tacaagttt atgtttaaag caccacaaat tgcgaagtcg 600
tgalttaaat gacatttttc cgttctcgtt taaactaagt ttctattatt taatalggat 660
attgiggagc gtagtgcag cactcagcag ctiaattaac gtgatactct ttaaccgagt 720
aacaactagt gaggtcatat tcaggaccga tccaacaata ttgagggttt tactccaagt 780
aaaaatttag ttttattttt aattatcata aacgacataa atataatatg gaaagatcac 840
aaalacigat taaaaactaa aatcalcaaa acgaaaaaga aaaaagaaaa aattgggttc 900
aacctctcag agttattaaa catittaggt tttaggctta aatctttaaa aaaaatcaga 960
actgaaaaac gaaaaatctt aattttattt tggactctga ttcatagctt atgtcgctta 1020
tgtagttagt ctagggaaga atctgtattt cgttaccgta atgagagttc gatactctct 1080
tacttgtagc gattctggag catgttacct tttttctttt cgttcaacaa caactttaat 1140
atggtaaaac aaaaatttatt ttattttggc tggctctact caagacaaat cttctgcga 1200
catcacataa tcatattaaa aaccataact tctgccactc tgtttttttt tttttttgta 1260
accattaaat gattggattt tgaatccatc catctgattt tttagctcaa caatttactt 1320
gcacattttc tatttgggtt tattttatct tagttacata taigattatc gaactagtat 1380
ctctttataa ttaagtattt ttctattttt tttaatttta gatttttgtg aattcattta 1440
cagtagaaaa ctglaaaacc atatggtcta attatagaat gaaaacttca acgaatccat 1500
acaacttatt ggctaaaat atataaatctg ctgaagcat attgttatta tttagttgga 1560
tttgacgac tctgacttta atgtataccg acatacccta tgatttagat gttgattttt 1620
cccattctta atatatccat gtaagagat tccaccaata catactaat tatttgcatt 1680
giaataaata ttatcatiaa aaaaaaat acactggacag ctggctcgtc ccattgtttc 1740
ttacgtccac caattacatt tgltaaagca aacttattag aacgttcatt tgtgagaagt 1800
tgggtctgac atgtgtctaa ggctatgtc agaaatcgga ttagcttatt aagtaacta 1860
tactataatc ttgttaatat agataaata tctagttcgt ccaaattaaa ctattttcat 1920
aacgtccacg tggcglaaac gtaatccatc agtcacttgt aatacttta taaccaaagt 1980

cttccaacac attcatcacc atctatctac tctttactct ctctctctct cacatcaatt 2040
attcatagtt ctctctcttc cggcaagaaa ag 2072

<210> 2

<211> 2000

<212> DNA

<213> Arabidopsis thaliana

<400> 2

ctctgttcc atgatcatat ataaaaggaa agagattgat tcagctaacc aaattcataa 60
acaaggataa gatatacaca actctaaacc tagctcaaac taigaacaat agaggataac 120
cacttgaac ctaattttaa tcagaagatt ttagtacttg tgctgcttat tcaaaattcc 180
tttcccaaa aaaaaaagt ttgacttacc tttigaaaga cgaacgtatc cactatggaa 240
caatctttcg taatctcact ttctagtttg gtatttataa aaccacaaat ttgattgtca 300
aattaaatac aataagagat tcctaaaccc agaattcaat tgcgaaatca aacagagacg 360
atacagagag agcgagagag cgagagagag agagtgaatt acgtgagaac gaatcggata 420
aacggaaatt tccagaagcc ggaatgagtt cccgctattg tgatgagtaa caaaacaaat 480
tcgggagttg tctatttaat acttatcaat agcccatgaa cgcagagacg tatggacgcg 540
agaatcgtaa gtacgtgaca cgtgtaaacc agtgacgtca ccgctctcgt cacaaccccg 600
ctagtattgt cctcttctat ttatctcac acaatcatgt taggttatca tagtaccact 660
aaacatttat gtgagaatgg atggatatta gacaaccgac gacaaataaa acaaacaaaa 720
gtagatagac gaccataaat taaactataa acgaagattg agggaaaaaa tcattggttt 780
agattttcga acaaatatcg tgaagagttt ttcattctgt agtttttgta attaatggig 840
tatgaaaaat cattggittg ttiagtigaa ttaaatttca glaaacggit tagagtatga 900
accactcaa taaagatttg actgccaagc gttactggig tciggtggat tgaatagtta 960
agagataaag gtttgggcca ttatgtggtt ttatttaaag ttgaaggcti catittattaa 1020
ttccttagag ctgatggcta atgtacaacg aatttgattc aggagtaaga taaacggtac 1080
acccaaacac gcgacgcgag cgaagaagac gatgacgaac aaggagaaaag ctcgagagag 1140
aaggagaaaa agaatgcagg agatctctct ccttcgaact attccttact ctgaccacca 1200

taggatgaa gagacctaat ttgtccaatt gtctcggtt ctttgtatcc taagatctat 1260
 catcatttac tctcgagat tctcaattc acagcattcg atgtgttaat ctctaatttc 1320
 acttgcttaa atccatttct tagtgctgt tacagtgtt gatgagttc tattttctc 1380
 tatttgggag ttggagacig aattgglat gtgtttgta cttacagatt tgtgtttcac 1440
 taaacaaaa gaatttagag aaattgaaaa tgatgaagta tgtgtctctc aagccctctt 1500
 ttgaatttg actgttcatt aggcgtcaaa tgattttta ggtgggtgtc ttgtgaaaa 1560
 gtacagtag tgactgttc aaaccgctgg attgattcg agattgcaag acagcttgcg 1620
 gtacacgat tgacgtgtt tcttacagc agaaacgtg atgtgtgtc tgaagcagt 1680
 aaatcttga ggcaccaaga agaaggtctc aaggtttatt tcatcaact tgatgtcaca 1740
 gactcttct cgattagaga gtgtgtgtc tggcttaagc aaacatttg aggttagat 1800
 attctctaa gatccttct gattctaat ctggaggat agttctaag tttctttaga 1860
 ttgaattac atgcatttgt gtgtctagt ttttcaggtg aataatgcag gtgttaacta 1920
 caatctcgcc tcagataata cgttgaatt tctgaaaca gtataatca ctaactacca 1980
 aggaacaaa aacatgacaa 2000

<210> 3

<211> 2000

<212> DNA

<213> Arabidopsis thaliana

<400> 3

catatacaca ccgcctaaca ttacagcctg tccccatccc ctatctttat tttcttttt 60
 ggctcattta gcttcaactt tttttaatag gtgtgattag aacatactaa aaaaaacata 120
 tgttttaact tatattatat gaataaagtg ttgtttttaa tagtttatat ttctattaca 180
 agtacaatat ataggatatt ttgtttaaaca attttaaat taaaagatat tgtgagtaca 240
 aatcattaaag cttaacacaa atgttcattg ctctttaatt cttacaatcg gaaaatttgg 300
 acaaaaaatc tagactaatc tccaaaacat tgtgaaaaat tgggggtttt ttttatcatt 360
 taaatgtct atagtgttca atcattatta aaaaaaatg tttttgttg aaattttaaa 420
 ttataaaaaa aaaacacgtt tatacataaa atgcaaaaac tttctttaa ccaaattaca 480

gagaaaggct atcaccatcc ttgtcttga tccaaccaca ctgtaatcc caaaatata 540
acttaaaacta taclatacta acatgaattg gttatagat tatctatcta gttagctaca 600
taattcataf aattagtttt ttaaaatata ctatgtaat gatagtaact aaaaaatata 660
ctatctcgag aaatccaacc aaaatgctat gatagtatat aaaataaaat gaaagaaaga 720
ggaaccaagt tgtgaaacca ttaccaaga agaagaagaa aaaaacaaga tggctcgagtc 780
tcgtgatagt atgaagcigg ggccgaaacc acgaatcaaa gaaccatttg acgatgcaag 840
tggcatcgat taaccaatca atagaaactc caccgctca aaatatcata gtgctattta 900
atttttcata ctatcatlla ttgtaggcta taacataitt ttgtatigta gatggacggc 960
taglacacgc aagtagtca ataaactttt aacaaatggc atcttgacat gcttaaacac 1020
attttataat tcgaacctca ctttgcttac ttacacacg tctacttata gtatgagc 1080
atcaatcatt gaaccgttcg tataatataa ttttttttt gtaatacgii ataattagaa 1140
agtaatggaa aataagttga aacacaaatt ttcaacgttt ttactaagat ttaattttac 1200
aattttcaaa atataaccga attattttaa tegtatttaa ttataatta attttctaaa 1260
aaaaaacaca gattaaaaaa atattttaa actaatcata aatatattaa tttttgtaat 1320
ttatattttt ttataacttt aaactaataa tattcaatta ttatctttaa agtttaca 1380
ttatcaataa taactaataa ataatgcata aaaacttttt aaaaattaac tttttgaaac 1440
aaattttttt ctaaaaaatt aaataataaa gaacagaaaa aatattattc aactaaaaat 1500
attggataat aactctgaaa gtttgcaatt ttcaataaaa tcatgtattt tttttgtttg 1560
acataaaaca ttaatttccc ctgtaaaac acacatttaa aatctaacta tatttaaaca 1620
aaaaaaaaaa gtccaatttt gactttaatt atataaaaaa agcaaagcta attatgggtc 1680
aaaagactca aaggccagaa ttgacgcagc cgtttatgag aagtgagaag acaatttcgg 1740
agtcacttcc ttattttcct ctcttcttc actctctcct atataaacct tctctctc 1800
ttctctctct ctatctctt caaaccattt tcgaaagcct tgagagagag aacacagacg 1860
ataccaactt tcttcaactt cgttgccgca gtataattat ctatctctc ggatatatct 1920
ctcttctcgc ggccgcgaca agaagctaca agaataaaaa gtctgttttc tctctttca 1980
agaaaccact tacttcgaaa 2000

<210> 4

<211> 2000

<212> DNA

<213> *Arabidopsis thaliana*

<400> 4

agcaagtagg ttacgtggaa ggaagatc aacgcacaaa aagagaaaca ctcaacgtta 60
agggaccaac cacaatctt ggattttgc tgaatgcagtg tgtgaaaaca ggiccacacg 120
ctttgtactt ttttcatcat cggctcacia taaagctcaa gcttttccat ggacagagat 180
aagcaagaaa aagatattaac attgttcacc tcaaaataat ttaggaaaat gacaatgaga 240
tttttacctc aaaagccaaa atcaaatct acccattctt tcactcgtca caatcatgag 300
tttcttacia tgccttacct agagttaga gacatatct cacaacaaaa gcttgtgtct 360
atttcaaatt cttagccgta gatgtcaca catgcataa tcattgaaa cagagcaaca 420
caggaaacca agcatatgta tctagatata cttagcaaga cataactata gcttttgaat 480
caacataggg attaatgata gagaatgagg aagctcaaga ttttatact agtttcttac 540
aaaacaaatt tctctctaac tgcacaaa ccaattagga ttgaagagc gtaccgttt 600
gagtcattgt ccaatgtct ccccccct tctacattt ttagcctgt gaataaaagc 660
acaagccaaa atgagaaggt gccaaaggcg ataaggatca atttctacca ttcaaaaaac 720
taatggtagg aattagaac gagagaaaac tacttgttga ggaaatagcc aaaagcgcaa 780
tctctgtcac ctgaataaag accaaaccgt cactttcaat gattcagcaa gaaaagaga 840
gagagagaga gagagattct ctataacatt tgagtcgaca tggattctaa tgcatacaaa 900
gtcatctcca ataaacaaac acttgaaact cacatggcta atagaacaag atcaaacct 960
taagtattaa gcattacaga cactactggc taacttttga cacatgttct taagtaacat 1020
agtatcaata tccgtgaac acatgaaca cacacaaca gggttaatg catcaaagtc 1080
ctgttatttc catatacaa catatttcat ttacaacag aatgcagcat tcaggcagtc 1140
caaatggaaa ggttgacaaa aaaaataat ctgttaact tacatatag gcagaatgia 1200
ataaccagcg aagaaaaaa cagaataaac agatcaatga gtaatgata aaaaaagtc 1260
acaagaatg tgccacagtg aacaagagg ccatgagaag aaattttcaa agaaaatatt 1320
agcatgttta gaatttttg ggtcaatgga tctgtcagct gcttagtttg aaaacacaaa 1380
tcttacagga aggaaagtc aagaaaaa aaataagcaa agttaatagc caccacaaga 1440
aatttcatac agaataatt aaatcgttg acttatctt ttattcaac taaaatcaag 1500

agaacttaat aattttcagc cacacgaacc atgtgttcaa agccaaaggt gagaagccaa 1560
aattatcagc ttaictccat taacaaggga aaagcaagac tagatttaag agttctctgt 1620
aactaaaaac tgcaggagtg agtaagtaaa taattcacca acaggaaaac aaaactcaat 1680
tatctatagc tgaatacaca tglaaalgag aatttattaa ctaaaacatc ttcctttgta 1740
actgaigtga catttacaat ttticatitt gaggtgtaag aaccgtgtga caagtgaaa 1800
ggttaaaata agcaaccttt gtgatatttt ctctccactt ttgaaattg ggtctccaaa 1860
ccacagccaa tcaatatctt ttataaatac aaacacacaa acagcatctt tctctcaaac 1920
acaacatat ctctatcaa acaccaacag ctctatctc tacttcattt ctcatcataa 1980
caaagagaga gaaaaaaact 2000

<210> 5

<211> 2000

<212> DNA

<213> Arabidopsis thaliana



<400> 5

tttctggaac acgttcacat gccgataatt attgttaagt tgtttgcata ttttattatg 60
tgcccttggtt gtgggatttt cgctaatttc tttctctggt ttttcgttct cctcttccct 120
caattatttt ctcaacttcc tatteggacca aagcattaat tctacacacg acagtaaagg 180
accactcaa acaataattt gttttcaact ctctcagaat atattttagt gacactaaaa 240
tagcataatc taaacacaac agaggccaac gactctgaat gatccgtacc gcagtggaaa 300
taacagtccg gttatgtcgg taggcatalat aagccactac ttcatattca atttctcaac 360
tttgtgttac ttagtctgta tctatcagt ttatcatcac ttttaattgt taccatatag 420
tatctgtaaa aactaattaa tatitatgaa tcatgctcat attttcaaat cttagagcta 480
ctcaacagaa cacacacaca tacagtcaaa attttatact acataatttt tgttttaagt 540
ttacagitat aaaccaaaca aatcacctca caaaagtata tagttatacc aaaccaagca 600
ccicataaat gaaaacaaaa ttaaaaataa taatatatat aataagaaat attcatcaat 660
actaacgttc aatttttagta aattggcaaa aataaaaat tagtatcact tccagagatt 720
acattattat ttgggtgaatt tggcaaaaat aaaatatctt gttgggtgat gccttgctgg 780

gacgacittg ttattaccg tcataitgia gtaacctaatt ttattacct aacattcatt 840
tgaacattaa aaacattaat gtcatitgat tggagagti tattaatgac cattatttcc 900
ttatttccaa acctaataaa cccattatit ccttatatta aattatggct tgattgggtc 960
aatitctttc tagctctttc tgttccgic ttgatcgaa taaaaacgac aaatgttatt 1020
tctaccggia tccgattatg tttaatatat tatgcattga tatttcacgc tacgatctgg 1080
accgacgitt agtltggacc ctgtcacgag ttttatctat gacggaccaa aagaaaaact 1140
aattaaagca gaalatcaaa gttagtcggg taagtccaa ctaataaccg agccctgttt 1200
ggctttgagc attagtagta atcaaagtaa acataatgaa gaaaagtatt tataatctat 1260
tataacaga ttaatatgta cgattccaga ttctatitai gtttctcgta tglaaaacat 1320
ataalgigaa cctacttaaa ctatactaag actattagti tggtttcaaa ttagattaat 1380
tagtttaacc aaactgaaca tatccctaga ttgtaaaagi ctaatatatc cactaattat 1440
ttttgcttgc cgacctctaa tccacaacti ttgttttct catttccact gattttcaac 1500
attccaccgg gtgttaccaa cagtaacata catacagta attttttcaa cactccaccg 1560
catgttacca atagtagcaa aacctcgcac aaaaatttgt accgactata aatattcaac 1620
tccaactctt tctcttccac aacttcaaat atctctccaa cactaaaata attgtaccga 1680
ctctctctct ctcctctctc gtaacaaaaa aatcaccaat ggcaaaaaca tatctctttg 1740
tactctctc aatctctat ctcttaccac tggagctcac ggcgccacc gcagcttcc 1800
agaccggagc ttcaaaaaa gccataaact tcatccaatc ttcttgcaaa accaccacat 1860
acctgtcctt atgtgtccac tcactctcgc tctaagcaaa cgacatccaa acaagcccta 1920
aacgttttagc tgagaccgct atagccgtga cactaagccg agcccaatcc acgaagctct 1980
tcgtctcgcg tctaacacgt 2000

<210> 6

<211> 2000

<212> DNA

<213> Arabidopsis thaliana

<400> 6

attatattta taataacatg attctatttc aatcgagtat gtaaaaaaag taaaaatttg 60

atcgattaat catgtagtaa tctcgtgttt tttccaaca aataattgat taaaagttaa 120
 aaacataatca taaatcaaaa ccgaatgagt aatccggatt gagctagaa ttacaagtta 180
 agttatcgt catacgggtg taatcatala tttttgttag aatccittaag agttgagaac 240
 tticaagaac agttaagttt aataatatt taatctttgt ataagagatt tcttcattct 300
 tctaaaacca caacaccttg tgatcgacaa ttcatacgt catatttgag atttaatttg 360
 ttttcttat ataagaaatc aataatctta tggatgatga ctaatttatt atataaaaac 420
 ccttactagg ggtatataa gacacgtaga aatgtctcca acatacagta ctataccaca 480
 aaatcagaa attgaacaa ccaaagcaat cagaaaacaa taitcatttt tgtttatttg 540
 gaaaagattg agccaatatt tcttaataa ctccttttta gataatcttg gaaccgggct 600
 tgttcgagcg ggtcgtgcca gccgggctgc tagggctgtt ggtagccgga ctcatccga 660
 gagtgttctt cacagcatcg gcgtctctt gtgcatgtt ctacttctt ccaccgtct 720
 acactccaaa atcataatta atgalataa tigtctataa ttaataataa tgaacaaaat 780
 tttgagaag aatcgtaac ctgttgatg acactagagg cttagaataa tagggaagga 840
 tigtctgat catgctcga gtgtaactgt gactgtgagt ggggtgtatg atcagcattc 900
 tggttcattg cgtgtgatac attgttcaag tactcttctt tcttcagcta taaataaaca 960
 tatatttgat ggttcattca tcacaaaat cagatttga aaattgcaa atttcagatc 1020
 atataagcaa aacaaatgtc gctattttgt ggtgtgtatg tacttgaact tgaccgtga 1080
 ctctccgcgc gcgtggctc aactgcggc tcgacgacat ctcttgggt gttctgtgaa 1140
 atgtcttaat taattaagat tgaagcttg tagaatactt gaagggattt gatatgcatt 1200
 ataattglat ttatataatt agaaagtga tatttttagta aaatcciaaa tctaagcatt 1260
 acactaacac gtggaaaata acataaccatt gacgatgac atggctaatt tttgtggag 1320
 gtgaatagtt tgaggattta ttaccctaac gtgtcttggc caagaagta agtaggatga 1380
 caggcaatag gaagatctta aactttttt tccggtgaca attatttatg actttttatt 1440
 gttgtcaaaa aataatatt caglaalata tcaataacga atacaataa aactcatccg 1500
 atcgatttct aagaatttat agctatatta aaattactt gaatccatgt aagaattgtg 1560
 tattggctt ttttagaaaa aagtaataat ctatgcagta atggcgggtg cataatata 1620
 gccitgagta gatgaatac caalatcaag ataacgtgag tcaccacgtg tctaacatct 1680
 tccgtagctc cgtttttacc atgacgtgc acatagatat aggtcatcat gaaaacgaga 1740
 aaactaactt taacactcgc acataactcc aagtttcgaa acttcttcac atcaacctaa 1800

tcggggcacg tacctacaca ccctgcgcga aacigcaaca cctatcttgi tctctcgccg 1860
 accaagacit gctataaata acctigacta acgagtcgga gacaaticac agttccaaac 1920
 acacaaaaaa cacaagaict aaaaaaaaaa gctttatca tttagaaaaa ttgggttctg 1980
 aatttcttgc aagagtgaag 2000

<210> 7

<211> 2000


<212> DNA

<213> Arabidopsis thaliana

<400> 7

tgaacaaaat taatgcagaa tttagtggag gatcagttgt gattttcctt cggtaaaatc 60
 attttagatt attgacattt gaattgttca atttcaaatt tcaaataatta taattcaatt 120
 caaagactag ctaagttttt aaaagtaatt agtgagagat ctacacagtga tataaaccac 180
 attcatttta attcatagca aagtaaatgg taagaaatga aaatggaaga aagagaggag 240
 agtagttcac acgaacatct gaaggagaag ttgagagagc ttgaggagga atggacagcc 300
 atgaaaacag gtaaaaactc atctgcggtt tcatggatta ccgtggaaga tgctcttgag 360
 tatctcgaaa acctctccag aaacctaatg ctctctctcc aacacaaacc aaaggcagag 420
 atgatacaag aaalatcgcc attaaggaga aaactcttcc acgactctga tgatgatgat 480
 caaactaaga agacgacatt gccttctcat tcatcttgtt ggctctcgaa tctgacgagt 540
 tgcagtgata ccacgaaagc aaagaagaag acaacgatta gaagatttgt tctgttaca 600
 atgggtctgc ttgtgtcatg gggttttagt gtgttgatga accactttga tcatctctcc 660
 atgaatacac aaatcattac ttgtgttctt acttgatatt ttattttatt ttgggttgag 720
 aatgttcatc aatctcaata aataaaaaaa gtttagtttt gtattattac ctctcgccct 780
 ctcttttaat tcatlgaacc tcttcaactt tggctgtaac tatgttggtt aaaacagiat 840
 aagtaatgia tgtatttatg gttttaggaa aatcttaca aataatactc tatcattgga 900
 tgaagaagca aaaactacac cctctacat cttttctatt cataaacttt ttgggggaaa 960
 gaactacatt ggtgggtgatt aggtgcatcg gtgacactaa tgttgaaaat atgttttgt 1020
 taaaattagt atgggttaaag ctaatcattt agttgagcaa aaaaagatgc taatcattta 1080

aafagaaaat ggactcaaac aaaagaagc cacacatca atgttaccat ctttatiggg 1140
ttttcatitt tggaagctat ttttatgaat ttgcagtggt ttgttgccca aaagaaaaaa 1200
agcataatc caaglatatc tttcatittg aatccatcag tcataatggt tacttgaatc 1260
aactatittgt tgggttttac gaagttacti gttagaaga ttcaaaatat atataattag 1320
actatcaaaag gaacaaaaga atgtaattgg gtcataactt ttccgaagc ttacgtgact 1380
tgggagaaaa tcaaacccat tagtttcacc cttaatcaaa ctaatcaca attttgaact 1440
ttgcaagaa ggtaaaataa taatcagcaa aaagattaat aaaatgaca aagtaatctt 1500
ttaaaagatt aataaagaaa taaagaaatg acataagctc tcaccgaccg acgtcttgct 1560
tttttattc ttaaggtcgg tgccatgctg acgcttcaa gcttagaatc tcgccagctg 1620
tgggataaac taaccgaacc caatcgaacc agacagacca aagacagacc agaccggact 1680
agtgtatgt cggcatcagc gtcttagtac gacgttaatt aaaaaataa aaataaata 1740
ttattatga taitccacca ataagagcca atgaaaagtg tglgaacaa ccaacttctg 1800
tagcaaaccg taccatttct ctcttagata cctttactt ccttctataa attaggttct 1860
tcctttcaca ttcttttgc ttcttcttct acataaaatt gatcgcctc gaaatttaca 1920
cagacttgic gctctctctt tctctcttat ttcacagagc ccctaacaat ttccaaaaat 1980
cttgcaagg aagaaaaaca 2000



<210> 8

<211> 2000



<212> DNA

<213> Arabidopsis thaliana




<400> 8



aaatatcatc atttataca aatttatgat atataaata tcattatcat tttaaaataa 60
cactaaaaa atcagtgcaa ttccgtataa tcttcataig aaactgcaag aatccttggt 120
gttccaaaaga ttatgatgat tailgtgagc ttgatattat ggtttccatc ttgtcaaaaa 180
atggctttta taaatggtag agctttctta gtlttctta agatatacac aacaggttac 240
aaatataatt tatttttgc taaaagaaata gttatagtaa acttgtaaaa actactacaa 300
ctattttatg gaaaalgtct taaagctgtt tcaactttta tacacattac tgcatttact 360

attctgaaat caatttttaa ttaaggctct tglactataa ggaatccctt aaataaatat 420
 aaatattaac aataaaaaaa tccaaaaatt cagaatgcaa tggctttaca attacaataa 480
 taatgtatit gattgtacia aaaaatatac ttaaatactc accaatatc ttttataata 540
 taatccacaa attctggaa gaggaaaaaac aagtaaatgc aagtcacatt tcacatttca 600
 aattaccaac taccaactac aaactacttc cgcgtctcaa attgaaccac taattaatca 660
 ctactttata tgcctatcat tcttttctt tcattacaat tctagaataa tgactlaacc 720
 aaaattcata caaataaaaac aatattttgg ctcccttcac caggctaaga attttggat 780
 tgaaaaatca agtacttcat aagaaactgt tggaatcatt taaaatttgt gatcatatat 840
 aaaaaaattt attcatcttt atatttaaga gtttaaaaac tgcaactttt gtttttcttt 900
 cactaagctt tatggccaca gtaattaaa agcagaatga aggtggctca atggaaaagg 960
 agaaltgat tgggctagt gggagagttc tgatgtctag tgttgggtac acgtgtccgt 1020
 cagttacaca tagcaataaa tcagacggca tgcattatt caaatctagt tcacatagta 1080
 cgactaatag ctgataaaat aatgattata cagcatatga attatgaatt caaaaaaaa 1140
 aaaaaattga aaatgttaag gagatgctat attttacaaa attcatcgca atgctttcta 1200
 ctaatttgcg aagtggtctt ctccagttag tctgtctgat tccaagcgat attattaaat 1260
 ctftgaagcat cgtctaaaagc attatagctt aagataacca aatgtttatt aaaaacacct 1320
 agtgaattt ttaaatlaaa acaattttga tatctttgta atatctaata ctactcttc 1380
 tgtgtctaaa aggatttaatt ttcaaaaatt tcacacatat taaaaaaaa aaaaaattac 1440
 tagctaaaca attttcaata atcataaaac aatagtaact taataatttt tttttatttt 1500
 caaaatagtc ctccaagttt acaattcatt ttagtattat aatcaacaaa atttgtatta 1560
 aaaagtltga aaattaatct ttgttgaaca aaaaaatcta gaaatcattt tttagaatta 1620
 gagagaggtt tgataaaaaa aaataaaaaa aatagagag aggtagtlaca tactaaacga 1680
 tgtgatacia ctattgacaa aatcttaatt ctccagtttag tagaataaac tagaaggaat 1740
 gaatgaagla aatgcgaatc caactactaa caaaccttac ttagtcatca tattttccca 1800
 tatgaaatcc ctatataaac ccatcalcat ctcccacttt ttcatatcc aacaacaaaa 1860
 acataagcia agaaaacgaa actcaactaa ttttgttata acccaaaaag aagttcaaac 1920
 acaatggctt tgcctttgag gtcttcaca tgccttggtt taacgggtgt catagtgtga 1980
 tcagtcgatg ctgcaatctc 2000



<210> 9

<211> 2000

<212> DNA

<213> *Arabidopsis thaliana*

<400> 9

tcaatttgag acgcggaagt agttttagt tggtagttgg taatttgaaa tgtgaaatgt 60
gacttgcaat tacttgtttt tccctctcca gaaatttgtg gatttatat taaaagataa 120
ttggtagia ttttaaggat atttttagta caatcaaata cattattatt gtaattgtaa 180
gaccattgca ttctgaaatt ttggattttt ttattgttaa tatttatatt tatttaaagg 240
attccttata gtacaaggac cttaattaaa aattgatttc agaatagtaa atgcagtaat 300
gtgtataaaa gtgaaacag cttaagaca ttttcataa aatagtgtga gtagtttta 360
caagtttact ataactatit cttttagcaa aaataaaita tatttgtaac ctgttgigta 420
tatcttaagg aaaactaaga aagctctacc atttataaaa gccatttttt gacaagatgg 480
aaaccataaa tcaaagctca caataalcat cataatcttt ggaacaccaa ggattcttgc 540
agtttcatat gaagattata cggaattgca ctgattatit tagtgttatt ttaaaatgat 600
aatgatattt tatatatcat aaatttgata taaatgatga tttttacag aatgaaatgt 660
tgttatatcat actataccit aagagcatat aacaigatat aattgctaca ttagtgtcgg 720
tagaggtttg tgtagicaaa ttgigactat taagtactac acatgtgaga catttatcaa 780
tcttgtaaga tgttgcttga cggtttttgg ctgatagtat ccaacaccaa acatcaacct 840
ttcacgtccc aagctctcta agacatcgga gaagcatcat cacatgcgta cgtagaaaagg 900
aaacacaaaa gttaaagaca ttltgtaata tgtgtaacta tgagacatat atgagactgc 960
atgtgactat gagttatata tggcctatag agcgatagtg ttatggagat tatattgcat 1020
tcatgctatg tactgcataa gacgaattcg gtttcgtgta tttattcccg cttagtcta 1080
gaattggitt caaglatagt tagaaaaaga ctgttttttt cgtcctatta tggttcgtat 1140
acaaaattat gcaatgaatt gatgiggcat tcacgtaaaa atcttcttgc agggctataa 1200
aacgtatttt tggcgtacga ttataataaa ttttgttatt attcagatta tttattaaat 1260
cttttttgtt gttagacaaac gcatttatgt taatgattat acgaaatac gagaaaaact 1320
ccatatagca cctctatcat tcaattatgt agtaagtcac aaaatgtgaa ctttgcta 1380

tataatggcc ataattagta aacgigaagg gagaattgtg atgtttagtg ttggatacac 1440
 gigtctatca gcgcacatc gcatttaatt ggacggtaatt acatatatag tacacataat 1500
 agctgacatt tcalctacaa cgaagataca aattttttaa cgattttctc ttgictacta 1560
 cagtataaca tcgctctaaa catgctaatt aaaaaaaca gtttaattaa ctgcagataa 1620
 atgaatatat attttttaac agtatagctg tatatatatt gacgtattgg tctttgtaca 1680
 ttctgaaaac tgalactaatt gtttacataa ttacatatga ctggagaaac taattatttt 1740
 ggtttagtaga agaaacataa ctacatatga ctggagaaac taattatttt cgtttagtaga 1800
 agaaactaga aagaatgcat gaagtaaatg caaaaccaac tactaacaaa ccctacttag 1860
 tcatcatitt tccccatc gaaatcccta tataaaccca tcatcatctc ccacttttct 1920
 cacatccaac aacaaaacac aagcgtagaa aacaaaactc aactaattgt gttatcacc 1980
 aaaagagaag agcaaacaca 2000

<210> 10


<211> 2000

<212> DNA

<213> Arabidopsis thaliana

<400> 10

ttttcattga ttacatttac aaatttttgg ttcatgttat tagtaaaata gatgtttatg 60
 aacaaaattg agctaacata tgacctctca taatatattg ttgatatcag tttaaatatc 120
 aatacattaa taataccggg aaaaaaaatc attcccaagt ttgttaaaga agatgttcat 180
 ataatttaag tttaaaaggg tataaccaac aaatgatta caaatgcgtc gatgggataa 240
 ataatagata tgagagaacc atacaaatat gatagtgaac aaaataaaat ctagtataat 300
 tatgggtcaa aaaaacataa aaagagggga aaagaaaaga taaagggaaa atcaaaaggg 360
 atgaggcaaa atagagaglt tcaataaaga taaatttggg tgatggtagac ggaataaattc 420
 ttcatgtttt tegttttgac catcttttcc ttlatittctc actttaattt tcttttattt 480
 ctctcgtgtt ttattcccat aaaattttta aaatttaacta gaacaagca caaaaatggg 540
 caatcttcca aatatactaa aatttttctat tataattaac aaaataaaaa agagagggaa 600



acigctaata tcggagtc attccttgg agaagagtaa agtgggcaaa caaaaaaaaa 660
ggaactgagc tgattcgatt ctctttggga cccccatcca tgcataccct ctctcatagt 720
ttcatitttac ttttatgttt attatitttac aagattaagi aattagcttt tttaacigca 780
agcataacaa ataacaaaac ctctattacg gtctcaaagc ctcaaaaatg tttttaaaca 840
tagtaaaaagg tttcgttcaa tacaatttga aaaaggaaaa aacaaaaaca atttcttac 900
aaatagatgg ccaccaacga ttaattaita cgttcaaact tataattggg atgactcggg 960
ggttttcgca tggcgtttgi tccaaaagag tgccataatt ttcattttgtt ctatataatg 1020
ttttaatcca taacacgttt tggatcaatg ttttcaagta atcaaagtga aagtagagaa 1080
tataatcaatt ggtgaaaatt ttgaaatgc aatgtaattc tttttaagtg gatccattga 1140
tcaaggaggc cactaaaatt tgcctgaggc ttggtagtgg gticatttgi gtttgataaa 1200
tacacttttt agtactcacc cccctttttt gtctgacacc gtatattgat attggttaca 1260
tctaaatttg tcaacttgti gtagcagta tagttgtctc agtgtgtcta tagactatag 1320
ttaaatgtia attagtagta ttatataaat actaattaca tcaatlaaat atttgcata 1380
gattcacaca aggcaaatit ccttggctaa ttgtcacact atttatgtga aggtgagtag 1440
ttaagctaa taacctcgaa gtgtcttggg ttatatgaga tcaagtttgg ctcttctaag 1500
tgaagataat aatttaaaga gtattttaat cgtgataatc taaatgttag ttgatagtt 1560
atacgtcagc gaacatgtac cgaccgacgg ccgatticca tgatacaata aaaaaatgaa 1620
aattagcaat ctataacat ggcacgacga tacttatggi gttgccacgi agcaagcacc 1680
ttccctttaa ccatgaggig tgcacacctc agaggacaac acggacaaga ccgagaaacc 1740
gcataactaac gcttgcaacg taaagcaca caccctaaaa ctctattttt tttaattttc 1800
ttttaacctc aacacaccac gcagctatac acgtttcaat gtgctaatac gtgtcttctt 1860
ctccgccgac caagacacac tacaatgtc ccgatggttt ggggagacaa atcacagttt 1920
ctactacaac aacaaatact ttacgaaaa aagcttttaa aacttatcta gtttcgtttg 1980
atacacattt gaagaaagaa 2000

<210> 11

<211> 2000

<212> DNA

<213> Arabidopsis thaliana

<400> 11

attaccataa tcataatc atgggtccga ctatatggct cgagtcctta tegtltatit 60
ttatttctta aatcatlgac ttataaacg cccatgagca ttaaataatt tgtctcaaaa 120
actcgaaaga aaatcgttat tgaatatggt ttaataacca aatctaatcc ttccgactc 180
atattggitt cattataaca cttaacaagt aaaagaattt tactcgactg cticagccac 240
aatgcggtag aagtcgcaa aggaccattc ttgatcgica ttttttctt ctaagcgitt 300
gcgtagcaa tticagcca tagcttccaa gtccttcgcg tctacctctc ctitaccctt 360
tgcctcgctc cataatlaag talaccaaac aaaaaaiga taattttatt aatgaagtga 420
acatcaatga aagacgtgat ttgggggag aaagagaaag gacgaaccgg latlgagagc 480
ttgaccata ttgtgataac ttctctctc ttgtttttt gcataatgta acaaatitga 540
aggataatat tgagaaataa attttgaga cgttgacttg tgacttgtga ggtaggatca 600
gtatcttgcc atttctgat cgataagcca cgtaccgaa gcgtgacagt ttaacttaag 660
gatctagaga aaaagtgtga gctatctcat ttcatattt acatatagca tataaatact 720
attttattt cctttgttta cagtttaata gttttagact cacaatatga cttttgaaag 780
ccaaaatgia gcagaacaat taagtitttt taatatctct ttgaaatgc aaatatgttt 840
ttaacctata taactggcaa catiaaaagg aagagattct catgatttgc aaaatatcaa 900
ggtagaaaag gcatgtaat ctttatitaa tatatatctt tgtttaaaac cacattggag 960
ataattgacc aaaaaaata caattitgtc aaaaagctta aattattgga ttgttaatga 1020
gcataiggaa gttlgacact tggagaaaat tcacatgtca ccacatagag ataagtatt 1080
tctagtcca ctaacacgat tacatttaca cccattccc tgcctgacct tgagtcactc 1140
ccatctctt ctttgactc ttgcgagaca cttagattta gaacaacaaa tgagttaag 1200
taagtaclaa gcctatgaaa aaaaaaaaag gtagaataag cttaagaatc cagcaacatt 1260
catggtatca ttgccattt atatcttaaa cgtgaagaa ataaagagag aaaggttatg 1320
aatctcttt gicaatagat atcatiggaa ctaagcgta tgaattctt ttgtcaatag 1380
gctaaaaacc gttttataa ttittacaaga aacatgatt ttccataata ttgaaacttt 1440
atatattct taatactta ggcaattata aattattcta tgtttgaat atttatitaa 1500
aatttttga gaataclaga acatataact ttccaagtc tagtgatccc ggctacggca 1560

taaattcata aattgatatt ctctccctcg cctccaaga aggaaatgac ccataaatgt 840
 ttgttccctt aatctcaaag caaaacaaca cagaaaatct tccatacctt caaggttgag 900
 actagcatct gcttcttctt ttctagaaaa cacctgctcg atactcactt gcttcgtcac 960
 aaccttcgtt tcggaattct tcctagtcag taactccaca gattctgtct caagaaccac 1020
 aatgtcgtca ccgtcatcat caggcaaagg aagagacgaa atgacttcag attccctata 1080
 aggactatca tcacttgaag atgatagcga caacaatgaa glagtcattg gtccttgcgt 1140
 aaaaaacca tccactttca atgttlaatt tctatataat tcaaacaaat atataaaaata 1200
 gaaacagaaa tcagattgaa tctggaacct cgttaagagcg aagccaatct ggagatccct 1260
 ctctlagagct gctgtcctata atttcacttc ttctagagctc aaaaacaaca tctaagcttc 1320
 ttcccgactt caatatccac tcaattacaa caaaatcagg tcgtacattc aggatttact 1380
 tcgcaatata aaaaccctaa aaagagatct taaaagctat aaattcgatt gaaaatcaca 1440
 tgaaatcgaa gctcacatga agaagaagaa gacaaaattg agtgattgac gaatctaatg 1500
 aggttaaaaa ccggcgagga gacgtggaaa aggcaatcga atgactgag gcctgaggga 1560
 agaagcaatg atttcgccgg cggagagatt gaagtgtgtg aaagacgcgg gaagaagatt 1620
 ggggcataaa ctgttcttgc gattttccaa gtttttggtc acaagagcaa tactatcccc 1680
 aacgtcgacg taaaatttac taaaagccca atatgtttta atagaaaaat taacataaag 1740
 cccaaattcg atgtaatggg ctctcatctt tcaggaaaaa ttaacattag cccagatatg 1800
 ccaatagagt agagataatg acgagtgacg acataaaaaa aggcacgcaa agaataataat 1860
 attaatitct tgggtggacaa gcaigtgtga ccattctcat ttcttctgc ccaaacitai 1920
 atttggatg ctcttcctac atatagacaa gacaatggta attcgacaat tcatcaagtc 1980
 tcaaatctaa ataagtaaac 2000

.
 .
 .
 .
 .

.
 .
 .

.
 .
 .

.
 .
 .

<210> 13
 <211> 2000
 <212> DNA
 <213> Arabidopsis thaliana

<400> 13

agaaacggca gicccacgag tatctgtata accattcatt gaccagctga atggtacttc 60
 ttgcataatc aaaactataa ttgcctggat aatgcaactc acctatttac atgaagaaa 120
 ctacagttaa caagtticca aattgtcttg gagaacaat gctacgcata caaatgggtg 180
 aataagaagg ctataagag atatatgaa acctgaacaa gagctgcacc agccaaagca 240
 gacaccactc caatgatatt catggccgtg gaatagtaaa acttcacata tgaagaaaa 300
 caatgctaga agaataataa agcaaatcca taaactaatt ctctccacg aacatatata 360
 ccttaagtag cagaagtaac aaagtggag catacaaaag tacattcatt ttacagaaa 420
 cagctccact gtataaaaag aagggttaagc agactagtca atctaagaa tggagaagca 480
 aaagcagcaa tctaacctga aaacaagcat ccctagatgc catllacgat agagaaagag 540
 agccattgat gcatgtagga gagtcatggc aaaacaatca itaaagagac ggagaacaaa 600
 gattgaatgt atctcttgg ataaacatag tagacttaat gcccaccaig gcaccigaaa 660
 tacgtaagga tacaagaaga gatacatca ctgaattcac taaaalcact cccatcataa 720
 acagacaaat caaaacaaa atattgttca ttaacatcgt aaatggacaa cgcttgacaa 780
 cgctacaaa tacttgatga aaagagataa gaaataagtc ttcatttata ccacatcagt 840
 cttcacgtag atgatcaaga cgatccccag atgacaalg tagagaacac caaaaagaat 900
 ctgcaaatg taagagagta gaattgagac aaatgaagcg atggatggat caaatccgc 960
 agctaacca aagataggag aticaaagag agggaaatgc aagcacctga gcagggtaga 1020
 cttcccccc agttaagtc tgcactgcag agtaaacata gaggaaacct gctggataaa 1080
 ccaatgccc tgtatgcct ticaagttcc catagctct ctctccccg agaaccac 1140
 taacctgccc caattagata aatttcactt aaagactact agtttacttt gaagaaacca 1200
 cagttgacca ttgacacta ccgtgacat glaagcatcc caatcgatct tegtgtctgt 1260
 taaaggcaaa tgacgaaatt tggtttaaaa cctatgcaaa tgcacatttc gagggatacg 1320
 caagcaaaaa aaaaaaggag gtaaatagct ctacatgga acataggcga taataagtc 1380
 gactaggatt gcatcagcaa gaattagcgc gaatgcgaat ggaactgcag gtttttgaa 1440
 tagatcggtat cgattcgtct cctccccag ccgaaggcta cgagaagctc tcaaactcgc 1500
 cggtagtag gcgcccgcga tgaacacaga gcaaatcgca tcagcgtcta gccaacgccg 1560
 cgtaacagac aactacttc atattactac tcttctaatt agcccaaatt aaatgagcct 1620
 attgggcttc ttgcttagt cgggttagag cccaatgtt gtatttttt ttaataatgc 1680
 aaaagtatta agcgataaat aaataagcat cgcaatcgtc ccaaactgt gtgtatgcat 1740

cagacatgag catatagagi aagcacgigi ccacacitit tcacaaagit alctaaaaac 1800
 aaaaaacaat taattagcat tcgacgtgia catatcactc gccacgtgia caagagccit 1860
 ggccititig cttctcttc ttgtctatta atatcatctc ctgattatac tctctitiga 1920
 ccaagctgct tcttctccat ctatccaaca tctcaaatct cagtaatctt gttctctcga 1980
 ttctgtitig gacgtitigia 2000

<210> 14

<211> 2000

<212> DNA

<213> Arabidopsis thaliana

<400> 14

ttcggagaaa atgcagctaa cgatcagg aaaagtgatt ggtattgtgg gagtgacagt 60
 agaagatgaa gaggaagcaa tctggggaa gccaaagatcc catcaccgtt gagctctctc 120
 tctctctctc tctctctcgt cgtgtggatc tctctctcga tctctctttg agccgcaaca 180
 actgatgagt tctctcagtc tctctctctt tctctctctt ctctcttttc ctctctctct 240
 tctctctctt aaaagggcta agttttttta aaaaatgtct accaaactga tacaagagg 300
 aagaagaaga aacactctgt actttttttt ttaacaagag aagaaaatgaa aacaagatg 360
 tgaatttaag aaaaaaata atggtctgac attacaaaac tgaatttaat catcatitga 420
 aaacatttaa tcatitgata aaatctgtaa cattaccccc ttaactttct tccacctaa 480
 cctttgtatc attgtttgtt ctctctatit tataacattc cttaaattat tgaattaaaa 540
 ctatctaaat tctgttaatt actcattgtt aatccatttt gtccactgga aatglaaatt 600
 ttactgtctt attttgcaa acactctiga attttttttt attttttttc tggtaataa 660
 gacacaaaca tgttggatta taaactattg tgaatcatac tctgttcggt aatgttttca 720
 atatcaaat tgaatgttgg tgaatggatt tgtcaacatc tctttttttt ttgttttaat 780
 cctttgctaa gttzagtcgg ccacatata attggttacg atttcatag gtctttctctc 840
 acgccagaag tgtgttttta ttgtttgat tgagttatta attattggaa gcttttcttt 900
 caagcaaagt aaaatgcgta ataattgatta gtcacatcca atggtagtc agtctattac 960

accgttaac aagctctggt catataattt ttttattttt ggaactaaca cttattagtt 1020
 taggtttcca tcacctattt aaticgiaat tcittatacat gcatataata gagatacata 1080
 tatacaaat ttatgatcatt ttigcacaac atgtgaicgc attcattagt atgcattatg 1140
 cgaaaacctc gacgcgcaaa agacacgtaa tagctaataa ttttactcat ttataatgat 1200
 tgaagcaaga cgaaaacaac aacatatata tcaaattgta aactagatat ttcttaaaag 1260
 tgaaaaaaaaa caaagaataa taaaggacaa ttttgagtca gtcctttaat attaaaacat 1320
 atatacataa ataagcacia acgtgggtac ctgtcttcac gcaatgtgga ctttagttta 1380
 tctaatcaaa atcaaaaata aaggtglaal agttctcgtc atttttcaaa ttttaaaaat 1440
 cagaaccaag tgatttttgt ttgagtattg atccattgtt taaacaattt aacacagtat 1500
 atacgtctct tgagatgttg acatgatgat aaaatcacgag atcgtctctt ggttttcgaa 1560
 ttttgaactt taatagtttt tttttttagg gaaactttta tagttgttta tcataagatt 1620
 agtcacctaa tggttacgtt gcagtaccga accaattttt tacccttttt tctaaatgtg 1680
 gtctgtggcat aatttccaaa agagatccaa aacccggttt gctcaactga taagccggtc 1740
 ggttctggtt tgaaaaacaa gaaataatct gaaagtgtga aacagcaacg tgtctcggtg 1800
 ttatcatgac caccitgccac ctcatcacg tcggtcattt tgtcgtttca cggttcacgc 1860
 tctagacacg tgcctcgtcc ccacatgac ttctcgtgcc gactcgttc gctttgcaaa 1920
 ctcaaacatg tgtgtatgtg taagtttcat cctaataagc atctcttacc acattaattt 1980
 aaaacaaaga aaacatcaaa 2000

.
 .
 .
 .
 .

.
 .
 .

<210> 15

.
 .
 .

<211> 2000

<212> DNA

.
 .
 .

<213> Arabidopsis thaliana

<400> 15

atgcccglia cttgatccca caggggtata attcactgaa ggaatttttc ccaacctga 60
 aaatgagacc aaatccacaa tgcctgaatg tagcttgtct agagcggcag glaacctgat 120
 taaatgcaac cacaaaacta tctcagcttg tctttgtttt gcgccttttg ttccatcca 180

tggtcgggtg aaaaagtga aatcccaaaa tgcagaaaga atatatgctt gcaaaaccag 240
 aaagggatgc tgcgtctaaa gctaagatgg aagctgatgc atccacaact atcgaagaag 300
 gtccactcca tgaigataat gagtggaca taaggtaata actcagtgaa tiagttaaaa 360
 atcaaaatit gtatititgt tatcatatit tctctitcc agccatitit gcaacaagaa 420
 atcgtatgtg tatctcatgt ctitttatca gigtgtcga tgaatgaaaat gagaaggata 480
 ctacaaaagc tgcctcaagt tcaggttact taaacttggg actctatgtg ctatccatgt 540
 gaaaatgtat caaalataag tccgtgtctt ctgactgtta tctaaattga tccgtgttta 600
 tgtgtagata cattaccga agggcttact cgggagcttc cagttgcaga tgagtatgag 660
 aaagcaatag ccatagcctc agggtcagga gaaacagagg aggaagatga ccttgaagat 720
 ctgaagaaac agctcgaagc tctcaacgct gcttgagtct ttatgagga gaaaaaatag 780
 ttgttttagc atatttatac atgaaagagg ctacaagtc aatttgtat gcaaaaagag 840
 tgtgttagga ttggtaaatt ggaiccatg gacatataaa cgtttttgat gatgactct 900
 cccgtttttt tagttgttac caagacaaa aagatacaat ttgaagatg ttcatggat 960
 ttattgtcat aatcatccga actagaactt agaaattcaa attagaaaaa gattttgtcc 1020
 aatgatgatg atccaagttt gglttgaat glgtaacca gcttttttg aggtaagata 1080
 ttaattgcgc ggggaaatca actagaagc gltgaatcg cgaattacaa ttacgtttt 1140
 ggaaggagaa ctgtctttt ttccctttt alatccctaa aatactttt tgtacattca 1200
 ttatatacag gaaaaacaga gtgttaacca aattattaca gagatattca aagatattc 1260
 aacctalcaa ctaaagatga cgtgtatatt gagaatggca attgcgttta tgagtgaggaa 1320
 accaaattat ttcatataca tgatctttag gaataccgct aataagagat tactaatcac 1380
 tatttgaggt ttaattttaa atctttgtg tccgtattt gaatatctg ttttagctaa 1440
 atcagctcca attttccca gttatcggtt gtcgaattca tattactgtt aatttagtta 1500
 acttccgaag actattagta catgattctt caacaagca aataaagaaa agaactttga 1560
 tgtcgttctc tcaacaattc aacaaaactt ggltcgcagc actcgaccgt gtgagatgtt 1620
 tgagaagica gaacaccaca cgtgtcgaca taaagccgc accacgcacg ctittcgtct 1680
 ggaatatacc aaattacttt acagaaatca ttgtattgt ttgttaataa aaactttgaa 1740
 ccactttgat tactatatta ttttttga atttattctg gtgacattaa tgttctatcc 1800
 tccacgcgtt tctcataaac cgacttggct tctttatit tctttatct ttctctatat 1860
 atttacacac acacgcattt gtgcagagat attcaagat tagaaacatt ctgatatag 1920

acaaaaaaca ttttcagac acaaaaalcat aaaatcttgg cctttagtag atcaaagttc 1980
 ttacattaa tcgttagaag 2000

<210> 16

<211> 2000

<212> DNA

<213> Arabidopsis thaliana

<400> 16

.....

.....

.....

ttagctgggg caggtcggct tgatttcatt gaggatctgc ttgagcatca gaagacactt 60
 ccacaaggta ggctgaggg ttcatgtgtt aggattataa tgttgtatgg caaggcigga 120
 atgacgaagc aagcacttga tactttcttc aatatggatt tatacggttg caaaagatct 180
 gtttaagtcct tcaatgccgc tcttcaagtc ttatctttca atcctgatct ccacaccatc 240
 tgggagtcc ttacgacgc cccatcgaag talggcattg atattgatgc ggtttctttt 300
 aacatigcca tlaaatcttt ctgtgagctg ggtattcttg acggggccta tatggcaatg 360
 agggagatgg agaaatcagg gttaacacca galgtagtta cataactac acttatatcg 420
 gcgttataca agcacgagag gtgtgtgatt ggaaatggat tgggaaacct tatggctc 480
 aagggtgtga agccgaacct caccactttt aatgttagga ttcagttttt agtgaatagg 540
 agacgagctt gggatgcaaa tgalctgttg ctgttgatgc ctaaactcca ggtggaaccg 600
 gacagcataa catataacat ggttatcaaa gggtttttcc ttgcgagggt cctgacatg 660
 gcggaaagag ttatatacgc gatgcatggt aagggttaca aaccaattt gaagatatac 720
 cagacgatga ttactatctt gtgcaaggca ggaaattttg atttggctta tacaatgtgc 780
 aaggatttga tgagaaagaa gtggtaccca aacttggaca cagttagat gtgtctcaa 840
 gggcttgtga aaaaggggca gcttgalcag gctaagtcga tcatggagtt agttcacaga 900
 agagtlacc ctttcaggtc aaaacagttg ctctccctta agtccatttt gtaatgttta 960
 catgaaatgg aacaaatttg ctacatgtac acgacaatgc ggccaaaaga attctttctt 1020
 tctttctcct gagtttattt tgtatagctg tgggtggaact ggaaatggga ttagcctca 1080
 tattaggagg cacacttate ttccaatcag attcaactat ttgttgttca atgtgaataa 1140

agcttaggat gatttggaaa ctccaatgga gggatacca gtaagtcag gaigcctcta 1200
 caatcttaag ttgtgatagc acagaactag ttacccttt gtaatgtac ttacaaaaca 1260
 tgttgttgat ttacgtgga gagcaaatg gcatacaca agcagggacc atgtgtgaa 1320
 gaaacaaaa ggaacttcgt ctgacgatgc aaaagctctt gagcacctgg aaaaatgagt 1380
 tatgttgtaa gcttcacta gccaaatfaa aagcactaaa catatacaa aggatatcaa 1440
 cgttgatagc tttttctga tatagcacga atgtaaaaa cagagaatag aacagagaca 1500
 acattaccag caatgaatat ttttattatt atgtcgaata aatacatttt cccacaagt 1560
 acagagcaaa caaaattcac atatttttgt ctattacaci ttgtgtcag actgacttgt 1620
 taactagatt tatggcggg gattataaca ttgtaacttt aaatatcttt taacgtaaga 1680
 aaaaaaac caaaattggt gtttattttt aaaaataaaa atttgagggg gcatgtatgt 1740
 ttgtttagt gattcacgtg ttcttttgt ctgaaaattc tgaattgat gattttgaaa 1800
 aaggaaaaaa agatcagagg tatctcaat tccaaaagga tgaatgatac aactcatata 1860
 tacaagatta tagtaaaaag aaggatgca tataaagaaa catagacatc atcgccatcc 1920
 atctcagat cagttccgta gtttactgga tcacgacaca cacacatata cactgtaca 1980
 tataagccat tcttatcaaa 2000

<210> 17
 <211> 2000
 <212> DNA
 <213> Arabidopsis thaliana

<400> 17

cactttgac gggaatcaa gcatcttag aagcgacgga gaagtcgaac aagattcaca 60
 tcgtcgatt ttgaatcgtt caaggatatac aatggcctgc tcttcttcaa gctctagcta 120
 ctgtacttc ttgtaaacc acataaatcc gggctcggg tatacccgct ccatctctcg 180
 gtgaatctcc ggaaccgtcg ttaatcgcca ccggaaccg cctccgtgat ttgccaagg 240
 ttctggatct gaatttcgat ttatcccaa ttctactcc catacattta ctaacgggt 300
 caagttccg ggtcgaccgg gatgaagtac tggccgtgaa ttcatgtct cagctctaca 360

aattactcga cgagacgccg acgatagttg acaccgcact acggctcgcc aaatcgttga 420
accggagggt cgtacacttc ggagaatcag aagtagcctt aaaccgggtc ggttcgccta 480
accgggtaaa gaacgcgctt caattctatt ccgcggtttt cgaatccctt gaaccgaact 540
tggggcgiga ttccgaggag agagttagag ttgagcgaga gttgttcggc cggagaatct 600
cgggtttgat tggaccggag aaaaccggaa ttcatagaga aagaatggaa gagaaagagc 660
aatggcgggt attaatggag aatgccggtt ttgaalcggt taagctgagt aattacgcag 720
tgagccaagc gaagattcta ttgtggaatt acaattacag caatttgtat tcaattgttg 780
aatctaaagc tggcttcac tccttggcct ggaacgattt acctctcttc actctttctt 840
ccctggcgata accaaaccaa accgatccgg tattcttagt ttgtttttgt tttcaatgtt 900
atttttgggt agacaaatat tcaattgtta atatactccg tggtcagagt gttttgtttt 960
tccttttagt cgaacgttga attaatcag gggtagggtt tgaattctct gaaccttaag 1020
tgttttttgg taacatcatt tggatttgag aactagggtt aaaaactggg cttagctctg 1080
ttgtttcttc attagataat ttaactgggt ttgcttcttt atttttgggt ggataaagtg 1140
accggttctg gtcactgttt tgagatgtaa ttactatttc ataaaattag gaagttgaaa 1200
gccaaatata ttgttaacta ctcttttatt tgaattttg ctcaaaaaag tgatgaaaag 1260
tagttttgat atatgaatat ctaccattat acataagtat atctgaacat ggtacaactt 1320
atgaaagcta aatgtcaata ctgcaaaga tataacaaat acaagttaca tgaataagag 1380
atgtgtgttg aatttataag tgcattttc ttttcacttt aaaacaaact tcatcttctt 1440
ttgtttctta tgtgtcaaag ttgccacagt tgcctctatt gagtctttca glgtcagctt 1500
cagtcactgt actgatttta cttttttttg ttgagtggtc caatgatgac atcactccca 1560
cgctctccat ccgtcttctt ttaacggta cgtggctccc accctctttt ctgatgtctt 1620
ttaccgactt gttctagccc aacttacttg ggccatttag attttttggg ggcccaagtt 1680
gctaaaagag gatttatcat agaaatctga acccgltgca gcgctcaaca catgtcacag 1740
tcctgacaaa cacttatca aatccttgtt aagtcgcc accgtgcacc agagcaccac 1800
gaggcaaaat ctgacagga caccgtcgta ctattatgtc ggaagacagg aaagcttaat 1860
taagcttaaa cctgacgtat ttaacttcgt taactctacc ttactaaagg gttttaattt 1920
aaaacttacc atctctctgt aagaataaaa actacttact ctataaattt aagcttcaag 1980
aaacctccaa aagcagagaa 2000

<210> 18
<211> 2000
<212> DNA
<213> *Arabidopsis thaliana*


<400> 18

gggggttcca caatgcattg attgagttg tgcigatgca gttacaactt gcttccgtct 60
ttttcacgtt tcagctcggg acaaaaacac attactacgg aaggacattg tttcatggag 120
ggcgagagta tagaggaact ggtcgagggt ttgtggctt ccacgcgaaa ttgctgaga 180
actataggtt ctatcccggt agtcactttg tgaaggcat tgaacttatg attcttctac 240
tggatataca gatctttggt caatcalata ggggtgtggt tacttatatt ctgatcactg 300
tttcgataat gttcatgggt gtgacatggc tcttgcctc ttctctttc aatccgtctg 360
gttttgagtg gcaaaagatt gtgatgact ggacagactg gaacaaatgg atttataacc 420
gaggaggaat tgggtccctt cctgagaaga gtgggaatc ttgtgggag aaagagttag 480
aacatcttag acatcagggt gttcgttgta tcaccttga gatctttttg gccttgcgt 540
tcttcattct tcaaatgga ctgtgtatc atctcagtac ctccaagga aaaaatcaga 600
gcttttgggt aagctaacct taaactact tccctgtgaa tggatgaaact ctgtattatt 660
gaacgataat gactctgttt tcaggtttat ggagcgtcat ggtttgtgat ctatttcatt 720
ctactgatgt tgaaggatg aatctttaag cacattcaca ttcaatttc ttgccaatc 780
ttaacattgt gttgccttct tctatgtaac gatcattgca gggtttgggc gttggaaggc 840
gaagattcag cacgaatttc cagcttctat tcagaatcat aaagggtctt gtgttctca 900
catttgtggc aattctcata acttctctcg cacttctctt gataacaatc aaagacttat 960
tcacttgcgt gcttgccttc atgccaactg gatggggaat gcttctcgta agtcttctc 1020
ctccatcaaa ctctctatatt tgtatctaag ataccagttt tgtctgggtt ttgtgacagt 1080
tgctttgttt tgcctccac gtcagattgc acaagcgtgt aagcctctga tccaacagct 1140
agggatatgg tcatcagtta ggacgctagc tcgaggctac gagatcgtga tgggcttgc 1200
tctgttcaca ccagttgcgt ttttggcctg gtttctttt gtgtctgagt tccagacaag 1260
aatgtctttt aaccaagcct ttagtagagg tctacaaatc tctgtattc ttgttggtca 1320

aagaaaagac cggcttcaa aaaacaagga gtgacatgta cacatgtaaa tggcgtatgt 1380
gtagatttaa tggtaacig ttattgttc ctccaagt aatctatagt tagaaagttc 1440
cigtaacitt catagatata tgttaattat ttccatttc cataagttcg cgtccattat 1500
aatagaaagt gcatgagacg tatcttttc atatggaaga cgaaaacatt ggttciccat 1560
tctaaaagtc aaaaagaaaa tgtatcac ttgaactttt ccacataggi catgataaag 1620
acaggactaa tcaatcattc attgttgata atggctcaa ttacctgttg ctagtattt 1680
ctacatttgt atagtgaat actgttttt ttccctctc caaataaaaa ctacagtact 1740
aaaagaaagt tggtaatcat aaaagtata aaactcaat aaaagcaatt actctgttc 1800
ttttcactc ttattttata tgactgaaat gaaattgtct atatgtgtt ttgtttgt 1860
tgttcattga tatgtaaca tcaaatctgc aaaaaacaac cgcaagtttg gattatacg 1920
atgaagcgtt tatctagtga gcgtcacgc agacgtcaca agttatgttg tacaatttg 1980
cgtaigtgtg atgaatcgc 2000



<210> 19
<211> 23
<212> DNA
<213> Artificial Sequence



<220>
<223> Description of Artificial Sequence: synthetic DNA



<400> 19
cgccagggtt ttccagta cga 23

<210> 20
<211> 24
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: synthetic DNA

<400> 20

agcggataac aatttcacac agga

24

5
3
5
3
5
3

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. An environmental stress responsive promoter comprising DNA of the following (a), (b) or (c):
 - 5 (a) DNA consisting of nucleotide sequence SEQ ID NO: 6,
 - (b) DNA consisting of a nucleotide sequence comprising a deletion, substitution or addition of one or more nucleotides relative to nucleotide sequence SEQ ID NO: 6, and functioning as an environmental stress responsive promoter; and
 - 10 (c) DNA hybridizing under stringent conditions to DNA consisting of nucleotide sequence SEQ ID NO: 6, and functioning as an environmental stress responsive promoter.
- 15 2. The promoter according to claim 1, wherein the environmental stress is at least one selected from the group consisting of cold stress, drought stress, salt stress and high photo stress.
3. An expression vector comprising the promoter according to either claim 1 or claim 2.
- 20 4. The expression vector according to claim 3, into which a desired gene is further incorporated.
- 25 5. A transformant comprising the expression vector according to either claim 3 or claim 4.
6. A transgenic plant comprising the expression vector according to claim 3 or claim 4.
- 30 7. The transgenic plant according to claim 6, wherein the plant is a plant body, plant organ, plant tissue or plant culture cell.
8. A method for producing a stress-resistant plant, which comprises culturing or cultivating the transgenic plant according to either claim 6 or claim 7.
- 35

9. An environmental stress responsive promoter according to claim 1, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.
- 5 10. An expression vector according to claim 3, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.
11. A transformant according to claim 5, substantially as herein described with
10 reference to any one or more of the Examples and/or accompanying Figures.
12. A transgenic plant according to claim 6, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.
- 15 13. A method according to claim 8, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.

Dated this twenty-third day of May 2006

Toyota Jidosha Kabushiki Kaisha and
Riken
Patent Attorneys for the Applicant:

F B RICE & CO

21 11 01 91431

FIG.1







FIG.2



1. $\frac{\text{WT (Dehydration- or cold-treated)}}{\text{WT (unstressed)}} > 2$
Drought- or cold-inducible genes
- $\frac{\text{WT (Dehydration- or cold-treated)}}{\text{WT (unstressed)}} = 1$
(Control; α -tubulin)
2. $\frac{\text{35S:DREB1A transgenic plants (unstressed)}}{\text{WT (unstressed)}} > 2$
DREB1A target genes
- $\frac{\text{35S:DREB1A transgenic plants (unstressed)}}{\text{WT (unstressed)}} = 1$
(Control; α -tubulin)

FIG.3



3/39

21 11 01 91431

FIG.4




FIG.5




FIG.6




FIG.7




FIG.8




FIG.9





FIG.10



10/39

FIG.11




FIG.12





FIG.13



13/39

FIG.14




FIG.15




FIG.16




FIG.17





FIG.18



18/39

FIG.19




FIG.20




FIG.21




FIG.22




FIG.23




FIG.24




FIG.25




FIG.26




FIG.27




FIG.28




FIG.29




FIG.30




FIG.31




FIG.32





FIG.33



33/39

FIG.34




FIG.35





FIG.36



36/39

FIG.37




FIG.38




FIG.39

