

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2007253803 B2

(54) Title **RNAi modulation of Aha and therapeutic uses thereof**

(51) International Patent Classification(s)
A61K 31/70 (2006.01) **C12N 15/113** (2010.01)
A61K 48/00 (2006.01)

(21) Application No: **2007253803** (22) Date of Filing: **2007.05.18**

(87) WIPO No: **WO07/137156**

(30) Priority Data

(31) Number **60/801,840** (32) Date **2006.05.19** (33) Country **US**

(43) Publication Date: **2007.11.29**
(44) Accepted Journal Date: **2012.05.24**

(71) Applicant(s)
The Scripps Research Institute;Alnylam Pharmaceuticals, Inc.

(72) Inventor(s)
Bramlage, Birgit;Balch, William;Vornlocher, Hans-Peter;Tan, Pamela;Constien, Rainer

(74) Agent / Attorney
FB Rice, Level 23 44 Market Street, Sydney, NSW, 2000

(56) Related Art
WO 2003/070918 A2
WO 2005/044981 A2
HARBORTH J et al. Antisense & Nucleic Acid Drug Development. 2003. 13(2): 83-105
HARST ANJA et al. The Biochemical Journal. 2005. 387(Pt 3): 789-796

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



PCT

(43) International Publication Date
29 November 2007 (29.11.2007)

(10) International Publication Number
WO 2007/137156 A3

(51) International Patent Classification:
A61K 31/70 (2006.01)

(74) Agents: MYERS, Louis et al.; Lowrie, Lando & Anastasi, LLP, One Main Street, Eleventh Floor, Cambridge, Massachusetts 02142 (US).

(21) International Application Number:
PCT/US2007/069229

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date: 18 May 2007 (18.05.2007)

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(25) Filing Language: English

Published:

(26) Publication Language: English

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(30) Priority Data:
60/801,840 19 May 2006 (19.05.2006) US

(88) Date of publication of the international search report:
23 October 2008

(71) Applicants (for all designated States except US): ALNYLAM PHARMACEUTICALS, INC. [US/US]; 300 Third Street, Cambridge, Massachusetts 02142 (US). THE SCRIPPS RESEARCH INSTITUTE [US/US]; 10550 North Torrey Pines Road, La Jolla, CA 92037 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): CONSTIEN, Rainer [DE/DE]; Kalte Marter 8, 95326 Kulmbach (DE). BRAMLAGE, Birgit [DE/DE]; Johann-Eck-Str. 11, 95326 Kulmbach (DE). TAN, Pamela [DE/DE]; Kalte Marter 8, 95326 Kulmbach (DE). VORNLOCHER, Hans-Peter [DE/DE]; Albert-Einstein-Ring 43a, 95448 Bayreuth (DE). BALCH, William [US/US]; 4037 Grayson Drive, San Diego, CA 92130 (US).

(54) Title: RNAI MODULATION OF AHA AND THERAPEUTIC USES THEREOF

(57) Abstract: The invention relates to a double-stranded ribonucleic acid (dsRNA) for inhibiting the expression of an Aha gene (Ahal gene), comprising an antisense strand having a nucleotide sequence which is less than 30 nucleotides in length, generally 19-25 nucleotides in length, and which is substantially complementary to at least a part of an Aha gene. The invention also relates to a pharmaceutical composition comprising the dsRNA together with a pharmaceutically acceptable carrier; methods for treating diseases caused by Ahal expression and the expression of an Aha gene using the pharmaceutical composition; and methods for inhibiting the expression of an Aha gene in a cell.

WO 2007/137156 A3

RNAi MODULATION OF AHA AND THERAPEUTIC USES THEREOF

Cross-Reference to Related Applications

This application is the Australian national phase filing of International application No. PCT/US20067/069229 filed May 18 2007 which claims priority to United States Serial No. 60/801,840 filed May 19 2006, the contents of which are incorporated herein in their entirety.

Field of the Invention

The present invention concerns methods of treatment using modulators of the gene Activator of Heat Shock Protein 90 ATPase (Aha). More specifically, the invention concerns methods of treating disorders associated with undesired Aha activity, by administering short interfering RNA which down-regulate the expression of Aha, and agents useful therein.

Background of the Invention

Activator of Heat Shock Protein 90 ATPase 1 (herein: Aha1) is an activator of the ATPase-activity of Hsp90 and is able to stimulate the inherent activity of yeast Hsp90 by 12-fold and human Hsp90 by 50-fold (Panaretou, B., et al., Mol. Cell 2002, 10:1307–1318). Biochemical studies have shown that Aha1 binds to the middle region of Hsp90 (Panaretou et al., 2002, *supra*, Lotz, G. P., et al., J. Biol. Chem. 2003, 278:17228–17235), and recent structural studies of the Aha1-Hsp90 core complex suggest that the co-chaperone promotes a conformational switch in the middle segment catalytic loop (370–390) of Hsp90 that releases the catalytic Arg380 and facilitates its interaction with ATP in the N-terminal nucleotide-binding domain (Meyer, P., et al., EMBO J. 2004, 23:511–519).

The molecular chaperone Heat shock protein 90 (Hsp90) is responsible for the *in vivo* activation or maturation of specific client proteins (Picard, D., Cell Mol. Life Sci. 2002, 59:1640–1648; Pearl, L. H., and Prodromou, C., Adv. Protein Chem. 2002, 59:157–185; Pratt, W. B., and Toft, D. O., Exp. Biol. Med. 2003, 228:111–133; Prodromou, C., and Pearl, L. H., Curr. Cancer Drug Targets 2003, 3:301–323). Crucial to such activation is the essential ATPase

activity of Hsp90 (Panaretou, B., et al., EMBO J. 1998, 17:4829–4836), which drives a conformational cycle involving transient association of the N-terminal nucleotide-binding domains within the Hsp90 dimer (Prodromou, C., et al., EMBO J. 2000, 19:4383–4392).

As a molecular chaperone, HSP90 promotes the maturation and maintains the stability of a large number of conformationally labile client proteins, most of which are involved in biologic processes that are often deranged within tumor cells, such as signal transduction, cell-cycle progression and apoptosis. As a result, and in contrast to other molecular targeted therapeutics, inhibitors of HSP90 achieve promising anticancer activity through simultaneous disruption of many oncogenic substrates within cancer cells (Whitesell L, and Dai C., Future Oncol. 2005; 1:529-540; WO 03/067262). Furthermore, HSP90 has been implicated in the degradation of Cystic Fibrosis Transmembrane Conductance Regulator (CFTR). Mutations in the CFTR gene lead to defective folding and ubiquitination of the protein as a consequence of HSP90 ATPase activity. Following ubiquitination, CFTR is degraded before it can reach its site of activity. Lack of active CFTR then leads to the development of cystic fibrosis in human subjects having such mutation. Therefore, the inhibition of HSP90 activity may be beneficial for subjects suffering from cancer or Cystic Fibrosis.

Hsp90 constitutes about 1-2% of total cellular protein (Pratt, W. B., Annu. Rev. Pharmacol. Toxicol. 1997, 37:297-326), and the inhibition of such large amounts of protein by means of an antagonist or inhibitor would potentially require the introduction of excessive amounts of the inhibitor or antagonist into a cell. An alternative approach is the inhibition of activators of HSP90's ATPase activity, such as Aha1, which are present in smaller amounts. By downregulating the amount of Aha1 present in the cell, the activity of HSP90 may be lowered substantially.

Significant sequence homology exists between *Homo sapiens* (NM_012111.1), *Mus musculus* (NM_146036.1) and *Pan troglodytes* (XM_510094.1) Aha 1. A clear *Rattus norvegicus* homologue of Aha 1 has not been identified; however, there is a *Rattus norvegicus* (XM_223680.3) gene which has been termed activator of heat shock protein ATPase homolog 2 (Ahsa 2) on the basis of its sequence homology to yeast Ahsa 2. Its sequence is homologous to *Mus musculus* RIKEN cDNA 1110064P04 gene (NM_172391.3), which is in turn similar in

sequence to *Mus musculus* Aha 1 except for N-terminal truncation. A *Homo sapiens* Ahsa 2 (NM_152392.1) has also been predicted, but sequence homology is limited. The functions of these latter three genes have not been sufficiently elucidated. However, there exists one region in which all of the above sequences are identical, and which may be used as the target for RNAi agents. It may be advantageous to inhibit the activity of more than one Aha gene.

Recently, double-stranded RNA molecules (dsRNA) have been shown to block gene expression in a highly conserved regulatory mechanism known as RNA interference (RNAi). WO 99/32619 (Fire et al.) discloses the use of a dsRNA of at least 25 nucleotides in length to inhibit the expression of genes in *C. elegans*. dsRNA has also been shown to degrade target

RNA in other organisms, including plants (see, e.g., WO 99/53050, Waterhouse et al.; and WO 99/61631, Heifetz et al.), *Drosophila* (see, e.g., Yang, D., et al., *Curr. Biol.* (2000) 10:1191-1200), and mammals (see WO 00/44895, Limmer; and DE 101 00 586.5, Kreutzer et al.). This natural mechanism has now become the focus for the development of a new class of pharmaceutical agents for treating disorders that are caused by the aberrant or unwanted regulation of a gene.

Despite significant advances in the field of RNAi and advances in the treatment of pathological processes mediated by HSP90, there remains a need for an agent that can selectively and efficiently attenuate HSP90 ATPase activity using the cell's own RNAi machinery. Such agent shall possess both high biological activity and in vivo stability, and shall effectively inhibit expression of a target Aha gene, such as Ahal, for use in treating pathological processes mediated directly or indirectly by Aha expression, e.g. Ahal expression.

Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a context for the present invention. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed before the priority date of each claim of this application.

Summary of the Invention

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

The invention provides double-stranded ribonucleic acid (dsRNA), as well as compositions and methods for inhibiting the expression of an Aha gene in a cell or mammal using such dsRNA. The invention also provides compositions and methods for treating pathological conditions and diseases mediated by the expression of an Aha gene, such as in cancer or cystic fibrosis. The dsRNA of the invention comprises an RNA strand (the antisense strand) having a region which is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and is substantially complementary to at least part of an mRNA transcript of an Aha gene.

In one example, the present invention provides a double-stranded ribonucleic acid (dsRNA) for inhibiting the expression of a human Aha gene in a cell e.g., an Aha1 gene such as from humans, wherein said dsRNA comprises at least two sequences that are complementary to each other and wherein a sense strand comprises a first sequence and an antisense strand comprises a second sequence comprising a region of complementarity which is substantially complementary to at least a part of a mRNA encoding an Aha gene, and wherein said region of complementarity is less than 30 nucleotides in length and wherein the dsRNA effects cleavage of an mRNA encoding an Aha gene within the target sequence of a second dsRNA having a sense strand set forth in SEQ ID NO: 51 and an antisense strand complementary to the latter sense strand and having a sequence set forth in SEQ ID NO: 52. For example, the ds RNA is AL-DP-7324 consisting of a sense strand having the sequence set forth in SEQ ID NO: 51 and an antisense strand having the sequence set forth in SEQ ID NO: 51. The dsRNA may, upon contact with a cell expressing the Aha gene e.g., HeLa cells or MLE12 cells, said dsRNA inhibits expression of said Aha gene in said cell by at least 20% compared to the inhibition of said Aha gene produced by a control siRNA duplex having a sequence that is unrelated to said Aha gene. The dsRNA may comprise at least one modified nucleotide.

In another example, the present invention also provides a vector that encodes at least one strand of the dsRNA according to any example hereof, and preferably encodes both strands of the dsRNA.

In another example, the present invention also provides a cell comprising the dsRNA or the vector according to any example hereof.

In another example, the present invention also provides a pharmaceutical composition comprising the dsRNA or the vector according to any example hereof and a pharmaceutically acceptable carrier or excipient. The pharmaceutical composition may comprise more than one dsRNA described herein.

In another example, the present invention also provides a method performed *in vitro* or *in vivo*, the method comprising introducing into the cell the double-stranded ribonucleic acid (dsRNA) or vector or pharmaceutical composition according to any example hereof and maintaining the cell for a time sufficient to obtain degradation of a mRNA transcript of an Aha gene, thereby inhibiting expression of an Aha gene in the cell. The method may be performed *in vitro* or *in vivo*.

In another example, the present invention also provides a method of treating, preventing or managing a pathological process mediated by Aha expression comprising administering to a patient in need of such treatment, prevention or management a therapeutically or prophylactically effective amount of the dsRNA or the vector or the pharmaceutical composition according to any example hereof. In a related example, the present invention also provides a use of the ds RNA or the vector or the pharmaceutical composition according to any example hereof in medicine *e.g.*, in the preparation of a medicament for treatment of Cystic Fibrosis or cancer or a tumor metastasis or a complication of Cystic Fibrosis mediated by expression of an Ahal gene of humans or a complication of a cancer mediated by expression of an Ahal gene of humans.

This invention also provides additional double-stranded ribonucleic acid (dsRNA) molecules for inhibiting the expression of an Aha gene. The dsRNA comprises at least two sequences that are complementary to each other. The dsRNA comprises a sense strand

comprising a first sequence and an antisense strand comprising a second sequence. The antisense strand comprises a nucleotide sequence which is substantially complementary to at least part of an mRNA encoding an Aha gene, and the region of complementarity is less than 30 nucleotides in length, generally 19-24 nucleotides in length. The dsRNA effects cleavage of an mRNA

5 encoding an Aha gene within the target sequence of a second dsRNA having a sense strand chosen from the group of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 27, SEQ ID NO: 29, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 35, SEQ ID NO: 37, SEQ ID NO: 39, SEQ ID NO: 43, SEQ ID NO: 45, SEQ ID NO: 47, SEQ ID NO: 49, SEQ ID NO: 51, SEQ ID NO: 53, SEQ ID NO: 55, SEQ ID NO: 57, SEQ ID NO: 59, SEQ ID NO: 61, SEQ ID NO: 63, SEQ ID NO: 65, SEQ ID NO: 67, SEQ ID NO: 69, SEQ ID NO: 71, SEQ ID NO: 73, SEQ ID NO: 75, SEQ ID NO: 77, SEQ ID NO: 79, SEQ ID NO: 81, SEQ ID NO: 83, SEQ ID NO: 85, SEQ ID NO: 87, SEQ ID NO: 89, SEQ ID NO: 91, SEQ ID NO: 93, SEQ ID NO: 95, SEQ ID NO: 97, SEQ ID NO: 99, SEQ ID NO: 101, SEQ ID NO: 103, SEQ ID NO: 105, SEQ ID NO: 107, SEQ ID NO: 109, SEQ ID NO: 111, SEQ ID NO: 113, SEQ ID NO: 115, SEQ ID NO: 117, SEQ ID NO: 119, SEQ ID NO: 121, SEQ ID NO: 123, SEQ ID NO: 125, SEQ ID NO: 127, SEQ ID NO: 129, SEQ ID NO: 131, SEQ ID NO: 133, SEQ ID NO: 135, SEQ ID NO: 137, SEQ ID NO: 139, SEQ ID NO: 141, SEQ ID NO: 143, SEQ ID NO: 145, SEQ ID NO: 147, SEQ ID NO: 149, SEQ ID NO: 151, SEQ ID NO: 153, SEQ ID NO: 155, SEQ ID NO: 157, SEQ ID NO: 159, SEQ ID NO: 163, SEQ ID NO: 165, SEQ ID NO: 167, SEQ ID NO: 169, SEQ ID NO: 171, SEQ ID NO: 173, SEQ ID NO: 175, SEQ ID NO: 177, SEQ ID NO: 179, SEQ ID NO: 181, and SEQ ID NO: 183, and an antisense strand complementary to the latter sense strand and chosen from the group of SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34, SEQ ID NO: 36, SEQ ID NO: 38, SEQ ID NO: 40, SEQ ID NO: 44, SEQ ID NO: 46, SEQ ID NO: 48, SEQ ID NO: 50, SEQ ID NO: 52, SEQ ID NO: 54, SEQ ID NO: 56, SEQ ID NO: 58, SEQ ID NO: 60, SEQ ID NO: 62, SEQ ID NO: 64, SEQ ID NO: 66, SEQ ID NO: 68, SEQ ID NO: 70, SEQ ID NO: 72, SEQ ID NO: 74, SEQ ID NO: 76, SEQ ID NO: 78, SEQ ID NO: 80, SEQ ID NO: 82, SEQ ID NO: 84, SEQ ID NO: 86, SEQ ID NO: 88, SEQ ID NO: 90, SEQ ID NO: 92, SEQ ID NO: 94, SEQ ID NO: 96, SEQ ID NO: 98, SEQ ID NO: 100, SEQ ID NO: 102, SEQ

104, SEQ ID NO: 106, SEQ ID NO: 108, SEQ ID NO: 110, SEQ ID NO: 112, SEQ ID
NO: 114, SEQ ID NO: 116, SEQ ID NO: 118, SEQ ID NO: 120, SEQ ID NO: 122, SEQ ID NO:
124, SEQ ID NO: 126, SEQ ID NO: 128, SEQ ID NO: 130, SEQ ID NO: 132, SEQ ID NO: 134,
SEQ ID NO: 136, SEQ ID NO: 138, SEQ ID NO: 140, SEQ ID NO: 142, SEQ ID NO: 144, SEQ
5 ID NO: 146, SEQ ID NO: 148, SEQ ID NO: 150, SEQ ID NO: 152, SEQ ID NO: 154, SEQ ID
NO: 156, SEQ ID NO: 158, SEQ ID NO: 160, SEQ ID NO: 164, SEQ ID NO: 166, SEQ ID NO:
168, SEQ ID NO: 170, SEQ ID NO: 172, SEQ ID NO: 174, SEQ ID NO: 176, SEQ ID NO: 178,
SEQ ID NO: 180, SEQ ID NO: 182, and SEQ ID NO: 184 (see Table 1 and Table 2). The Aha
gene is preferably an Aha1 gene, and more preferably a Homo sapiens Aha1 gene. The dsRNA,
10 upon contacting with a cell expressing the Aha gene, may inhibit the expression of the Aha gene
in said cell by at least 20%, or at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%,
85%, 90% or 95%, e.g. in HeLa and/or MLE 12 cells. The dsRNA may be different from said
second dsRNA, but may have at least 5, at least 10, at least 15, at least 18, or at least 20
contiguous nucleotides per strand in common with one of the above named nucleotide
15 sequences.

Preferably, the second dsRNA is chosen from the group of AL-DP-7301, AL-DP-7308,
AL-DP-7318, AL-DP-7320, AL-DP-7322, AL-DP-7324, AL-DP-7325, AL-DP-7326, AL-DP-
7327, AL-DP-7329, AL-DP-7331, AL-DP-7333, AL-DP-7340, AL-DP-7342, AL-DP-7303, AL-
DP-7305, AL-DP-7307, AL-DP-7309, AL-DP-7316, and AL-DP-7337, AL-DP-7304, AL-DP-
20 7312, AL-DP-7339, AL-DP-7344, AL-DP-7306, AL-DP-7317, AL-DP-7346, AL-DP-7310, AL-
DP-7323, AL-DP-7335, AL-DP-7338, AL-DP-7341, AL-DP-7302, AL-DP-7315, AL-DP-7328,
AL-DP-7330, AL-DP-7336, AL-DP-7345, AL-DP-9250, AL-DP-9251, AL-DP-9252, AL-DP-
9253, AL-DP-9254, AL-DP-9255, AL-DP-9256, AL-DP-9257, AL-DP-9258, AL-DP-9259, AL-
DP-9260, AL-DP-9261, AL-DP-9262, AL-DP-9263, AL-DP-9264, AL-DP-9265, AL-DP-9266,
25 AL-DP-9267, AL-DP-9268, AL-DP-9269, AL-DP-9270, AL-DP-9271, AL-DP-9272, AL-DP-
9273, AL-DP-9274, AL-DP-9275, AL-DP-9276, AL-DP-9277, AL-DP-9279, AL-DP-9280, AL-
DP-9281, AL-DP-9282, AL-DP-9283, AL-DP-9284, AL-DP-9285, AL-DP-9286, AL-DP-9287,
AL-DP-9288, and AL-DP-9289 (see Table 1 and Table 2).

Alternatively, the dsRNA itself may be chosen from the group of AL-DP-7301, AL-DP-
30 7308, AL-DP-7318, AL-DP-7320, AL-DP-7322, AL-DP-7324, AL-DP-7325, AL-DP-7326, AL-

DP-7327, AL-DP-7329, AL-DP-7331, AL-DP-7333, AL-DP-7340, AL-DP-7342, AL-DP-7303, AL-DP-7305, AL-DP-7307, AL-DP-7309, AL-DP-7316, and AL-DP-7337, AL-DP-7304, AL-DP-7312, AL-DP-7339, AL-DP-7344, AL-DP-7306, AL-DP-7317, AL-DP-7346, AL-DP-7310, AL-DP-7323, AL-DP-7335, AL-DP-7338, AL-DP-7341, AL-DP-7302, AL-DP-7315, AL-DP-5 7328, AL-DP-7330, AL-DP-7336, AL-DP-7345, AL-DP-9250, AL-DP-9251, AL-DP-9252, AL-DP-9253, AL-DP-9254, AL-DP-9255, AL-DP-9256, AL-DP-9257, AL-DP-9258, AL-DP-9259, AL-DP-9260, AL-DP-9261, AL-DP-9262, AL-DP-9263, AL-DP-9264, AL-DP-9265, AL-DP-10 9266, AL-DP-9267, AL-DP-9268, AL-DP-9269, AL-DP-9270, AL-DP-9271, AL-DP-9272, AL-DP-9273, AL-DP-9274, AL-DP-9275, AL-DP-9276, AL-DP-9277, AL-DP-9279, AL-DP-9280, AL-DP-9281, AL-DP-9282, AL-DP-9283, AL-DP-9284, AL-DP-9285, AL-DP-9286, AL-DP-9287, AL-DP-9288, and AL-DP-9289 (see Table 1 and Table 2).

The dsRNA may comprise at least one modified nucleotide. Preferably, the modified nucleotide is chosen from the group of: a 2'-O-methyl modified nucleotide, a nucleotide comprising a 5'-phosphorothioate group, and a terminal nucleotide linked to a cholesteryl derivative or dodecanoic acid bisdecylamide group. Alternatively, the modified nucleotide is chosen from the group of: a 2'-deoxy-2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a locked nucleotide, an abasic nucleotide, 2'-amino-modified nucleotide, 2'-alkyl-modified nucleotide, morpholino nucleotide, a phosphoramidate, and a non-natural base comprising nucleotide.

20 In another aspect, the invention provides an isolated cell comprising one of the dsRNAs of the invention. The cell is generally a mammalian cell, such as a human cell. Other embodiments of the cell comprising a dsRNA of the invention are as provided for other aspects of the invention above.

25 In yet another aspect, a pharmaceutical composition for inhibiting the expression of an Aha gene in an organism is provided, comprising a dsRNA and a pharmaceutically acceptable carrier, wherein the dsRNA comprises at least two sequences that are complementary to each other and wherein a sense strand comprises a first sequence and an antisense strand comprises a second sequence comprising a region of complementarity which is substantially complementary to at least a part of a mRNA encoding an Aha gene, and wherein said region of complementarity

is less than 30 nucleotides in length, and wherein the dsRNA effects cleavage of an mRNA encoding an Aha gene within the target sequence of a second dsRNA having a sense strand chosen from the group of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 27, SEQ ID NO: 29, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 35, SEQ ID NO: 37, SEQ ID NO: 39, SEQ ID NO: 43, SEQ ID NO: 45, SEQ ID NO: 47, SEQ ID NO: 49, SEQ ID NO: 51, SEQ ID NO: 53, SEQ ID NO: 55, SEQ ID NO: 57, SEQ ID NO: 59, SEQ ID NO: 61, SEQ ID NO: 63, SEQ ID NO: 65, SEQ ID NO: 67, SEQ ID NO: 69, SEQ ID NO: 71, SEQ ID NO: 73, SEQ ID NO: 75, SEQ ID NO: 77, SEQ ID NO: 79, SEQ ID NO: 81, SEQ ID NO: 83, SEQ ID NO: 85, SEQ ID NO: 87, SEQ ID NO: 89, SEQ ID NO: 91, SEQ ID NO: 93, SEQ ID NO: 95, SEQ ID NO: 97, SEQ ID NO: 99, SEQ ID NO: 101, SEQ ID NO: 103, SEQ ID NO: 105, SEQ ID NO: 107, SEQ ID NO: 109, SEQ ID NO: 111, SEQ ID NO: 113, SEQ ID NO: 115, SEQ ID NO: 117, SEQ ID NO: 119, SEQ ID NO: 121, SEQ ID NO: 123, SEQ ID NO: 125, SEQ ID NO: 127, SEQ ID NO: 129, SEQ ID NO: 131, SEQ ID NO: 133, SEQ ID NO: 135, SEQ ID NO: 137, SEQ ID NO: 139, SEQ ID NO: 141, SEQ ID NO: 143, SEQ ID NO: 145, SEQ ID NO: 147, SEQ ID NO: 149, SEQ ID NO: 151, SEQ ID NO: 153, SEQ ID NO: 155, SEQ ID NO: 157, SEQ ID NO: 159, SEQ ID NO: 163, SEQ ID NO: 165, SEQ ID NO: 167, SEQ ID NO: 169, SEQ ID NO: 171, SEQ ID NO: 173, SEQ ID NO: 175, SEQ ID NO: 177, SEQ ID NO: 179, SEQ ID NO: 181, and SEQ ID NO: 183, and an antisense strand complementary to the latter sense strand and chosen from the group of SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34, SEQ ID NO: 36, SEQ ID NO: 38, SEQ ID NO: 40, SEQ ID NO: 44, SEQ ID NO: 46, SEQ ID NO: 48, SEQ ID NO: 50, SEQ ID NO: 52, SEQ ID NO: 54, SEQ ID NO: 56, SEQ ID NO: 58, SEQ ID NO: 60, SEQ ID NO: 62, SEQ ID NO: 64, SEQ ID NO: 66, SEQ ID NO: 68, SEQ ID NO: 70, SEQ ID NO: 72, SEQ ID NO: 74, SEQ ID NO: 76, SEQ ID NO: 78, SEQ ID NO: 80, SEQ ID NO: 82, SEQ ID NO: 84, SEQ ID NO: 86, SEQ ID NO: 88, SEQ ID NO: 90, SEQ ID NO: 92, SEQ ID NO: 94, SEQ ID NO: 96, SEQ ID NO: 98, SEQ ID NO: 100, SEQ ID NO: 102, SEQ ID NO: 104, SEQ ID NO: 106, SEQ ID NO: 108, SEQ ID NO: 110, SEQ ID NO: 112, SEQ ID NO: 114, SEQ ID NO: 116, SEQ ID NO: 118, SEQ ID NO: 120, SEQ ID NO: 122, SEQ ID NO: 124, SEQ ID NO: 126, SEQ ID NO: 128, SEQ ID NO: 130, SEQ ID NO: 132, SEQ ID NO: 134,

SEQ ID NO: 136, SEQ ID NO: 138, SEQ ID NO: 140, SEQ ID NO: 142, SEQ ID NO: 144, SEQ ID NO: 146, SEQ ID NO: 148, SEQ ID NO: 150, SEQ ID NO: 152, SEQ ID NO: 154, SEQ ID NO: 156, SEQ ID NO: 158, SEQ ID NO: 160, SEQ ID NO: 164, SEQ ID NO: 166, SEQ ID NO: 168, SEQ ID NO: 170, SEQ ID NO: 172, SEQ ID NO: 174, SEQ ID NO: 176, SEQ ID NO: 178, 5 SEQ ID NO: 180, SEQ ID NO: 182, and SEQ ID NO: 184 (see Table 1 and Table 2). Therein, the Aha gene may be an Aha1 gene, and preferably a *Homo sapiens* Aha1 gene. The dsRNA comprised in the pharmaceutical composition may, upon contact with a cell expressing said Aha gene, inhibit the expression of said Aha gene in said cell by at least 20%, or at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 85%, 90% or 95%, e.g. in HeLa and/or MLE 12 10 cells. The dsRNA may be different from said second dsRNA, but may have at least 5, at least 10, at least 15, at least 18, or at least 20 contiguous nucleotides per strand in common with one of the above named nucleotide sequences.

Preferably, the second dsRNA is chosen from the group of AL-DP-7301, AL-DP-7308, AL-DP-7318, AL-DP-7320, AL-DP-7322, AL-DP-7324, AL-DP-7325, AL-DP-7326, AL-DP- 15 7327, AL-DP-7329, AL-DP-7331, AL-DP-7333, AL-DP-7340, AL-DP-7342, AL-DP-7303, AL-DP-7305, AL-DP-7307, AL-DP-7309, AL-DP-7316, and AL-DP-7337, AL-DP-7304, AL-DP-7312, AL-DP-7339, AL-DP-7344, AL-DP-7306, AL-DP-7317, AL-DP-7346, AL-DP-7310, AL-DP-7323, AL-DP-7335, AL-DP-7338, AL-DP-7341, AL-DP-7302, AL-DP-7315, AL-DP-7328, 20 AL-DP-7330, AL-DP-7336, AL-DP-7345, AL-DP-9250, AL-DP-9251, AL-DP-9252, AL-DP-9253, AL-DP-9254, AL-DP-9255, AL-DP-9256, AL-DP-9257, AL-DP-9258, AL-DP-9259, AL-DP-9260, AL-DP-9261, AL-DP-9262, AL-DP-9263, AL-DP-9264, AL-DP-9265, AL-DP-9266, 25 AL-DP-9267, AL-DP-9268, AL-DP-9269, AL-DP-9270, AL-DP-9271, AL-DP-9272, AL-DP-9273, AL-DP-9274, AL-DP-9275, AL-DP-9276, AL-DP-9277, AL-DP-9279, AL-DP-9280, AL-DP-9281, AL-DP-9282, AL-DP-9283, AL-DP-9284, AL-DP-9285, AL-DP-9286, AL-DP-9287, AL-DP-9288, and AL-DP-9289 (see Table 1 and Table 2).

Alternatively, the dsRNA comprised in the pharmaceutical composition itself may be chosen from the group of AL-DP-7301, AL-DP-7308, AL-DP-7318, AL-DP-7320, AL-DP-7322, AL-DP-7324, AL-DP-7325, AL-DP-7326, AL-DP-7327, AL-DP-7329, AL-DP-7331, AL-DP-7333, AL-DP-7340, AL-DP-7342, AL-DP-7303, AL-DP-7305, AL-DP-7307, AL-DP-7309, AL-DP-7316, and AL-DP-7337, AL-DP-7304, AL-DP-7312, AL-DP-7339, AL-DP-7344, AL-DP- 30

7306, AL-DP-7317, AL-DP-7346, AL-DP-7310, AL-DP-7323, AL-DP-7335, AL-DP-7338, AL-DP-7341, AL-DP-7302, AL-DP-7315, AL-DP-7328, AL-DP-7330, AL-DP-7336, AL-DP-7345, AL-DP-9250, AL-DP-9251, AL-DP-9252, AL-DP-9253, AL-DP-9254, AL-DP-9255, AL-DP-9256, AL-DP-9257, AL-DP-9258, AL-DP-9259, AL-DP-9260, AL-DP-9261, AL-DP-9262, AL-5 DP-9263, AL-DP-9264, AL-DP-9265, AL-DP-9266, AL-DP-9267, AL-DP-9268, AL-DP-9269, AL-DP-9270, AL-DP-9271, AL-DP-9272, AL-DP-9273, AL-DP-9274, AL-DP-9275, AL-DP-9276, AL-DP-9277, AL-DP-9279, AL-DP-9280, AL-DP-9281, AL-DP-9282, AL-DP-9283, AL-DP-9284, AL-DP-9285, AL-DP-9286, AL-DP-9287, AL-DP-9288, and AL-DP-9289 (see Table 1 and Table 2).

10 The dsRNA comprised in the pharmaceutical composition may comprise at least one modified nucleotide. Preferably, said modified nucleotide is chosen from the group of: a 2'-O-methyl modified nucleotide, a nucleotide comprising a 5'-phosphorothioate group, and a terminal nucleotide linked to a cholesteryl derivative or dodecanoic acid bisdecylamide group. Alternatively, said modified nucleotide is chosen from the group of: a 2'-deoxy-2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a locked nucleotide, an abasic nucleotide, 15 2'-amino-modified nucleotide, 2'-alkyl-modified nucleotide, morpholino nucleotide, a phosphoramidate, and a non-natural base comprising nucleotide.

In yet another aspect, a method for inhibiting the expression of an Aha gene in a cell is provided, the method comprising:

20 (a) introducing into the cell a double-stranded ribonucleic acid (dsRNA), wherein the dsRNA comprises at least two sequences that are complementary to each other and wherein a sense strand comprises a first sequence and an antisense strand comprises a second sequence comprising a region of complementarity which is substantially complementary to at least a part of a mRNA encoding Aha1, and wherein said region of complementarity is less than 30 nucleotides in length and wherein the dsRNA effects cleavage of an mRNA encoding an Aha gene within the target sequence of a second dsRNA having a sense strand chosen from the group of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 27, SEQ ID NO: 29, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 35, SEQ ID NO: 37, SEQ ID NO: 39, SEQ ID NO: 43, SEQ ID NO: 45, SEQ ID NO: 47, SEQ ID NO: 49, SEQ ID NO: 51,

SEQ ID NO: 53, SEQ ID NO: 55, SEQ ID NO: 57, SEQ ID NO: 59, SEQ ID NO: 61, SEQ ID NO: 63, SEQ ID NO: 65, SEQ ID NO: 67, SEQ ID NO: 69, SEQ ID NO: 71, SEQ ID NO: 73, SEQ ID NO: 75, SEQ ID NO: 77, SEQ ID NO: 79, SEQ ID NO: 81, SEQ ID NO: 83, SEQ ID NO: 85, SEQ ID NO: 87, SEQ ID NO: 89, SEQ ID NO: 91, SEQ ID NO: 93, SEQ ID NO: 95, 5 SEQ ID NO: 97, SEQ ID NO: 99, SEQ ID NO: 101, SEQ ID NO: 103, SEQ ID NO: 105, SEQ ID NO: 107, SEQ ID NO: 109, SEQ ID NO: 111, SEQ ID NO: 113, SEQ ID NO: 115, SEQ ID NO: 117, SEQ ID NO: 119, SEQ ID NO: 121, SEQ ID NO: 123, SEQ ID NO: 125, SEQ ID NO: 127, SEQ ID NO: 129, SEQ ID NO: 131, SEQ ID NO: 133, SEQ ID NO: 135, SEQ ID NO: 137, SEQ ID NO: 139, SEQ ID NO: 141, SEQ ID NO: 143, SEQ ID NO: 145, SEQ ID NO: 147, SEQ 10 ID NO: 149, SEQ ID NO: 151, SEQ ID NO: 153, SEQ ID NO: 155, SEQ ID NO: 157, SEQ ID NO: 159, SEQ ID NO: 163, SEQ ID NO: 165, SEQ ID NO: 167, SEQ ID NO: 169, SEQ ID NO: 171, SEQ ID NO: 173, SEQ ID NO: 175, SEQ ID NO: 177, SEQ ID NO: 179, SEQ ID NO: 181, and SEQ ID NO: 183, and an antisense strand complementary to the latter sense strand and chosen from the group of SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ 15 ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34, SEQ ID NO: 36, SEQ ID NO: 38, SEQ ID NO: 40, SEQ ID NO: 44, SEQ ID NO: 46, SEQ ID NO: 48, SEQ ID NO: 50, SEQ ID NO: 52, SEQ ID NO: 54, SEQ ID NO: 56, SEQ ID NO: 58, SEQ ID NO: 60, SEQ ID NO: 62, SEQ ID NO: 64, SEQ ID NO: 66, SEQ ID NO: 68, SEQ ID NO: 70, SEQ ID NO: 72, SEQ ID NO: 74, SEQ ID NO: 76, SEQ ID NO: 78, SEQ ID NO: 80, SEQ ID NO: 82, SEQ 20 ID NO: 84, SEQ ID NO: 86, SEQ ID NO: 88, SEQ ID NO: 90, SEQ ID NO: 92, SEQ ID NO: 94, SEQ ID NO: 96, SEQ ID NO: 98, SEQ ID NO: 100, SEQ ID NO: 102, SEQ ID NO: 104, SEQ ID NO: 106, SEQ ID NO: 108, SEQ ID NO: 110, SEQ ID NO: 112, SEQ ID NO: 114, SEQ ID NO: 116, SEQ ID NO: 118, SEQ ID NO: 120, SEQ ID NO: 122, SEQ ID NO: 124, SEQ ID 25 NO: 126, SEQ ID NO: 128, SEQ ID NO: 130, SEQ ID NO: 132, SEQ ID NO: 134, SEQ ID NO: 136, SEQ ID NO: 138, SEQ ID NO: 140, SEQ ID NO: 142, SEQ ID NO: 144, SEQ ID NO: 146, SEQ ID NO: 148, SEQ ID NO: 150, SEQ ID NO: 152, SEQ ID NO: 154, SEQ ID NO: 156, SEQ ID NO: 158, SEQ ID NO: 160, SEQ ID NO: 164, SEQ ID NO: 166, SEQ ID NO: 168, SEQ ID NO: 170, SEQ ID NO: 172, SEQ ID NO: 174, SEQ ID NO: 176, SEQ ID NO: 178, SEQ ID NO: 180, SEQ ID NO: 182, and SEQ ID NO: 184; and 30 (b) maintaining the cell produced in step (a) for a time sufficient to obtain degradation of the

mRNA transcript of an Aha gene, thereby inhibiting expression of an Aha gene in the cell. The Aha gene is preferably an Aha1 gene, and more preferably a homo sapiens Aha1 gene. The dsRNA may be different from said second dsRNA, but may have at least 5, at least 10, at least 15, at least 18, or at least 20 contiguous nucleotides per strand in common with one of the above 5 named nucleotide sequences.

Preferably, the second dsRNA is chosen from the group of AL-DP-7301, AL-DP-7308, AL-DP-7318, AL-DP-7320, AL-DP-7322, AL-DP-7324, AL-DP-7325, AL-DP-7326, AL-DP-7327, AL-DP-7329, AL-DP-7331, AL-DP-7333, AL-DP-7340, AL-DP-7342, AL-DP-7303, AL-DP-7305, AL-DP-7307, AL-DP-7309, AL-DP-7316, and AL-DP-7337, AL-DP-7304, AL-DP-10 7312, AL-DP-7339, AL-DP-7344, AL-DP-7306, AL-DP-7317, AL-DP-7346, AL-DP-7310, AL-DP-7323, AL-DP-7335, AL-DP-7338, AL-DP-7341, AL-DP-7302, AL-DP-7315, AL-DP-7328, AL-DP-7330, AL-DP-7336, AL-DP-7345, AL-DP-9250, AL-DP-9251, AL-DP-9252, AL-DP-9253, AL-DP-9254, AL-DP-9255, AL-DP-9256, AL-DP-9257, AL-DP-9258, AL-DP-9259, AL-DP-9260, AL-DP-9261, AL-DP-9262, AL-DP-9263, AL-DP-9264, AL-DP-9265, AL-DP-9266, 15 AL-DP-9267, AL-DP-9268, AL-DP-9269, AL-DP-9270, AL-DP-9271, AL-DP-9272, AL-DP-9273, AL-DP-9274, AL-DP-9275, AL-DP-9276, AL-DP-9277, AL-DP-9279, AL-DP-9280, AL-DP-9281, AL-DP-9282, AL-DP-9283, AL-DP-9284, AL-DP-9285, AL-DP-9286, AL-DP-9287, AL-DP-9288, and AL-DP-9289 (see Table 1 and Table 2).

Alternatively, the dsRNA itself is chosen from the group of AL-DP-7301, AL-DP-7308, 20 AL-DP-7318, AL-DP-7320, AL-DP-7322, AL-DP-7324, AL-DP-7325, AL-DP-7326, AL-DP-7327, AL-DP-7329, AL-DP-7331, AL-DP-7333, AL-DP-7340, AL-DP-7342, AL-DP-7303, AL-DP-7305, AL-DP-7307, AL-DP-7309, AL-DP-7316, and AL-DP-7337, AL-DP-7304, AL-DP-7312, AL-DP-7339, AL-DP-7344, AL-DP-7306, AL-DP-7317, AL-DP-7346, AL-DP-7310, AL-DP-7323, AL-DP-7335, AL-DP-7338, AL-DP-7341, AL-DP-7302, AL-DP-7315, AL-DP-7328, 25 AL-DP-7330, AL-DP-7336, AL-DP-7345, AL-DP-9250, AL-DP-9251, AL-DP-9252, AL-DP-9253, AL-DP-9254, AL-DP-9255, AL-DP-9256, AL-DP-9257, AL-DP-9258, AL-DP-9259, AL-DP-9260, AL-DP-9261, AL-DP-9262, AL-DP-9263, AL-DP-9264, AL-DP-9265, AL-DP-9266, AL-DP-9267, AL-DP-9268, AL-DP-9269, AL-DP-9270, AL-DP-9271, AL-DP-9272, AL-DP-9273, AL-DP-9274, AL-DP-9275, AL-DP-9276, AL-DP-9277, AL-DP-9279, AL-DP-9280, AL- 30 DP-9281, AL-DP-9282, AL-DP-9283, AL-DP-9284, AL-DP-9285, AL-DP-9286, AL-DP-9287,

AL-DP-9288, and AL-DP-9289. Preferably, the method is performed in vitro. Other embodiments of the method for inhibiting the expression of an Aha gene in a cell are as provided for other aspects of the invention above.

In yet another aspect, a method of treating, preventing or managing pathological

5 processes mediated by Aha expression is provided, comprising administering to a patient in need of such treatment, prevention or management a therapeutically or prophylactically effective amount of a dsRNA, wherein the dsRNA comprises at least two sequences that are complementary to each other and wherein a sense strand comprises a first sequence and an antisense strand comprises a second sequence comprising a region of complementarity which is
10 substantially complementary to at least a part of a mRNA encoding Aha1, and wherein said region of complementarity is less than 30 nucleotides in length and wherein the dsRNA effects cleavage of an mRNA encoding an Aha gene within the target sequence of a second dsRNA having a sense strand chosen from the group of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID
15 NO: 21, SEQ ID NO: 23, SEQ ID NO: 27, SEQ ID NO: 29, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 35, SEQ ID NO: 37, SEQ ID NO: 39, SEQ ID NO: 43, SEQ ID NO: 45, SEQ ID NO: 47, SEQ ID NO: 49, SEQ ID NO: 51, SEQ ID NO: 53, SEQ ID NO: 55, SEQ ID NO: 57, SEQ ID NO: 59, SEQ ID NO: 61, SEQ ID NO: 63, SEQ ID NO: 65, SEQ ID NO: 67, SEQ ID NO: 69, SEQ ID NO: 71, SEQ ID NO: 73, SEQ ID NO: 75, SEQ ID NO: 77, SEQ ID NO: 79,
20 SEQ ID NO: 81, SEQ ID NO: 83, SEQ ID NO: 85, SEQ ID NO: 87, SEQ ID NO: 89, SEQ ID NO: 91, SEQ ID NO: 93, SEQ ID NO: 95, SEQ ID NO: 97, SEQ ID NO: 99, SEQ ID NO: 101, SEQ ID NO: 103, SEQ ID NO: 105, SEQ ID NO: 107, SEQ ID NO: 109, SEQ ID NO: 111, SEQ ID NO: 113, SEQ ID NO: 115, SEQ ID NO: 117, SEQ ID NO: 119, SEQ ID NO: 121, SEQ ID NO: 123, SEQ ID NO: 125, SEQ ID NO: 127, SEQ ID NO: 129, SEQ ID NO: 131, SEQ ID NO:
25 133, SEQ ID NO: 135, SEQ ID NO: 137, SEQ ID NO: 139, SEQ ID NO: 141, SEQ ID NO: 143, SEQ ID NO: 145, SEQ ID NO: 147, SEQ ID NO: 149, SEQ ID NO: 151, SEQ ID NO: 153, SEQ ID NO: 155, SEQ ID NO: 157, SEQ ID NO: 159, SEQ ID NO: 163, SEQ ID NO: 165, SEQ ID NO: 167, SEQ ID NO: 169, SEQ ID NO: 171, SEQ ID NO: 173, SEQ ID NO: 175, SEQ ID NO: 177, SEQ ID NO: 179, SEQ ID NO: 181, and SEQ ID NO: 183, and an antisense strand
30 complementary to the latter sense strand and chosen from the group of SEQ ID NO: 6, SEQ ID

NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34, SEQ ID NO: 36, SEQ ID NO: 38, SEQ ID NO: 40, SEQ ID NO: 44, SEQ ID NO: 46, SEQ ID NO: 48, SEQ ID NO: 50, SEQ ID NO: 52, SEQ ID NO: 54, SEQ ID NO: 56, SEQ ID NO: 58, SEQ ID NO: 60, SEQ ID NO: 62, SEQ ID NO: 64, SEQ ID NO: 66, SEQ ID NO: 68, SEQ ID NO: 70, SEQ ID NO: 72, SEQ ID NO: 74, SEQ ID NO: 76, SEQ ID NO: 78, SEQ ID NO: 80, SEQ ID NO: 82, SEQ ID NO: 84, SEQ ID NO: 86, SEQ ID NO: 88, SEQ ID NO: 90, SEQ ID NO: 92, SEQ ID NO: 94, SEQ ID NO: 96, SEQ ID NO: 98, SEQ ID NO: 100, SEQ ID NO: 102, SEQ ID NO: 104, SEQ ID NO: 106, SEQ ID NO: 108, SEQ ID NO: 110, SEQ ID NO: 112, SEQ ID NO: 114, SEQ ID NO: 116, SEQ ID NO: 118, SEQ ID NO: 120, SEQ ID NO: 122, SEQ ID NO: 124, SEQ ID NO: 126, SEQ ID NO: 128, SEQ ID NO: 130, SEQ ID NO: 132, SEQ ID NO: 134, SEQ ID NO: 136, SEQ ID NO: 138, SEQ ID NO: 140, SEQ ID NO: 142, SEQ ID NO: 144, SEQ ID NO: 146, SEQ ID NO: 148, SEQ ID NO: 150, SEQ ID NO: 152, SEQ ID NO: 154, SEQ ID NO: 156, SEQ ID NO: 158, SEQ ID NO: 160, SEQ ID NO: 164, SEQ ID NO: 166, SEQ ID NO: 168, SEQ ID NO: 170, SEQ ID NO: 172, SEQ ID NO: 174, SEQ ID NO: 176, SEQ ID NO: 178, SEQ ID NO: 180, SEQ ID NO: 182, and SEQ ID NO: 184. The dsRNA may be different from said second dsRNA, but may have at least 5, at least 10, at least 15, at least 18, or at least 20 contiguous nucleotides per strand in common with one of the above named nucleotide sequences.

20 Preferably, the second dsRNA is chosen from the group of AL-DP-7301, AL-DP-7308, AL-DP-7318, AL-DP-7320, AL-DP-7322, AL-DP-7324, AL-DP-7325, AL-DP-7326, AL-DP-7327, AL-DP-7329, AL-DP-7331, AL-DP-7333, AL-DP-7340, AL-DP-7342, AL-DP-7303, AL-DP-7305, AL-DP-7307, AL-DP-7309, AL-DP-7316, and AL-DP-7337, AL-DP-7304, AL-DP-7312, AL-DP-7339, AL-DP-7344, AL-DP-7306, AL-DP-7317, AL-DP-7346, AL-DP-7310, AL-25 DP-7323, AL-DP-7335, AL-DP-7338, AL-DP-7341, AL-DP-7302, AL-DP-7315, AL-DP-7328, AL-DP-7330, AL-DP-7336, AL-DP-7345, AL-DP-9250, AL-DP-9251, AL-DP-9252, AL-DP-9253, AL-DP-9254, AL-DP-9255, AL-DP-9256, AL-DP-9257, AL-DP-9258, AL-DP-9259, AL-DP-9260, AL-DP-9261, AL-DP-9262, AL-DP-9263, AL-DP-9264, AL-DP-9265, AL-DP-9266, AL-DP-9267, AL-DP-9268, AL-DP-9269, AL-DP-9270, AL-DP-9271, AL-DP-9272, AL-DP-30 9273, AL-DP-9274, AL-DP-9275, AL-DP-9276, AL-DP-9277, AL-DP-9279, AL-DP-9280, AL-

DP-9281, AL-DP-9282, AL-DP-9283, AL-DP-9284, AL-DP-9285, AL-DP-9286, AL-DP-9287, AL-DP-9288, and AL-DP-9289 (see Table 1 and Table 2).

Alternatively, the dsRNA itself is chosen from the group of AL-DP-7301, AL-DP-7308, AL-DP-7318, AL-DP-7320, AL-DP-7322, AL-DP-7324, AL-DP-7325, AL-DP-7326, AL-DP-
5 7327, AL-DP-7329, AL-DP-7331, AL-DP-7333, AL-DP-7340, AL-DP-7342, AL-DP-7303, AL-DP-7305, AL-DP-7307, AL-DP-7309, AL-DP-7316, and AL-DP-7337, AL-DP-7304, AL-DP-7312, AL-DP-7339, AL-DP-7344, AL-DP-7306, AL-DP-7317, AL-DP-7346, AL-DP-7310, AL-DP-7323, AL-DP-7335, AL-DP-7338, AL-DP-7341, AL-DP-7302, AL-DP-7315, AL-DP-7328, AL-DP-7330, AL-DP-7336, AL-DP-7345, AL-DP-9250, AL-DP-9251, AL-DP-9252, AL-DP-
10 9253, AL-DP-9254, AL-DP-9255, AL-DP-9256, AL-DP-9257, AL-DP-9258, AL-DP-9259, AL-DP-9260, AL-DP-9261, AL-DP-9262, AL-DP-9263, AL-DP-9264, AL-DP-9265, AL-DP-9266, AL-DP-9267, AL-DP-9268, AL-DP-9269, AL-DP-9270, AL-DP-9271, AL-DP-9272, AL-DP-9273, AL-DP-9274, AL-DP-9275, AL-DP-9276, AL-DP-9277, AL-DP-9279, AL-DP-9280, AL-DP-9281, AL-DP-9282, AL-DP-9283, AL-DP-9284, AL-DP-9285, AL-DP-9286, AL-DP-9287, AL-DP-9288, and AL-DP-9289. Other embodiments of the method comprising administering a
15 dsRNA of the invention are as provided for other aspects of the invention above.

In yet another aspect, a vector for inhibiting the expression of an Aha gene in a cell is provided, said vector comprising a regulatory sequence operably linked to a nucleotide sequence that encodes at least one strand of a dsRNA, wherein one of the strands of said dsRNA is substantially complementary to at least a part of a mRNA encoding Aha1 and wherein said dsRNA is less than 30 base pairs in length and wherein the dsRNA effects cleavage of an mRNA encoding an Aha gene within the target sequence of a second dsRNA having a sense strand chosen from the group of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 27, SEQ ID NO: 29, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 35, SEQ ID NO: 37, SEQ ID NO: 39, SEQ ID NO: 43, SEQ ID NO: 45, SEQ ID NO: 47, SEQ ID NO: 49, SEQ ID NO: 51, SEQ ID NO: 53, SEQ ID NO: 55, SEQ ID NO: 57, SEQ ID NO: 59, SEQ ID NO: 61, SEQ ID NO: 63, SEQ ID NO: 65, SEQ ID NO: 67, SEQ ID NO: 69, SEQ ID NO: 71, SEQ ID NO: 73, SEQ ID NO: 75, SEQ ID NO: 77, SEQ ID NO: 79, SEQ ID NO: 81, SEQ ID NO: 83, SEQ ID NO: 85, SEQ ID NO: 87, SEQ ID NO: 89, SEQ ID NO: 91, SEQ ID NO:
20
25
30

93, SEQ ID NO: 95, SEQ ID NO: 97, SEQ ID NO: 99, SEQ ID NO: 101, SEQ ID NO: 103, SEQ ID NO: 105, SEQ ID NO: 107, SEQ ID NO: 109, SEQ ID NO: 111, SEQ ID NO: 113, SEQ ID NO: 115, SEQ ID NO: 117, SEQ ID NO: 119, SEQ ID NO: 121, SEQ ID NO: 123, SEQ ID NO: 125, SEQ ID NO: 127, SEQ ID NO: 129, SEQ ID NO: 131, SEQ ID NO: 133, SEQ ID NO: 135, 5 SEQ ID NO: 137, SEQ ID NO: 139, SEQ ID NO: 141, SEQ ID NO: 143, SEQ ID NO: 145, SEQ ID NO: 147, SEQ ID NO: 149, SEQ ID NO: 151, SEQ ID NO: 153, SEQ ID NO: 155, SEQ ID NO: 157, SEQ ID NO: 159, SEQ ID NO: 163, SEQ ID NO: 165, SEQ ID NO: 167, SEQ ID NO: 169, SEQ ID NO: 171, SEQ ID NO: 173, SEQ ID NO: 175, SEQ ID NO: 177, SEQ ID NO: 179, SEQ ID NO: 181, and SEQ ID NO: 183, and an antisense strand complementary to the latter 10 sense strand and chosen from the group of SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34, SEQ ID NO: 36, SEQ ID NO: 38, SEQ ID NO: 40, SEQ ID NO: 44, SEQ ID NO: 46, SEQ ID NO: 48, SEQ ID NO: 50, SEQ ID NO: 52, SEQ ID NO: 54, SEQ ID NO: 56, SEQ ID NO: 58, SEQ 15 ID NO: 60, SEQ ID NO: 62, SEQ ID NO: 64, SEQ ID NO: 66, SEQ ID NO: 68, SEQ ID NO: 70, SEQ ID NO: 72, SEQ ID NO: 74, SEQ ID NO: 76, SEQ ID NO: 78, SEQ ID NO: 80, SEQ ID NO: 82, SEQ ID NO: 84, SEQ ID NO: 86, SEQ ID NO: 88, SEQ ID NO: 90, SEQ ID NO: 92, SEQ ID NO: 94, SEQ ID NO: 96, SEQ ID NO: 98, SEQ ID NO: 100, SEQ ID NO: 102, SEQ ID NO: 104, SEQ ID NO: 106, SEQ ID NO: 108, SEQ ID NO: 110, SEQ ID NO: 112, SEQ ID 20 NO: 114, SEQ ID NO: 116, SEQ ID NO: 118, SEQ ID NO: 120, SEQ ID NO: 122, SEQ ID NO: 124, SEQ ID NO: 126, SEQ ID NO: 128, SEQ ID NO: 130, SEQ ID NO: 132, SEQ ID NO: 134, SEQ ID NO: 136, SEQ ID NO: 138, SEQ ID NO: 140, SEQ ID NO: 142, SEQ ID NO: 144, SEQ ID NO: 146, SEQ ID NO: 148, SEQ ID NO: 150, SEQ ID NO: 152, SEQ ID NO: 154, SEQ ID NO: 156, SEQ ID NO: 158, SEQ ID NO: 160, SEQ ID NO: 164, SEQ ID NO: 166, SEQ ID NO: 168, SEQ ID NO: 170, SEQ ID NO: 172, SEQ ID NO: 174, SEQ ID NO: 176, SEQ ID NO: 178, 25 SEQ ID NO: 180, SEQ ID NO: 182, and SEQ ID NO: 184. The dsRNA may be different from said second dsRNA, but may have at least 5, at least 10, at least 15, at least 18, or at least 20 contiguous nucleotides per strand in common with one of the above named nucleotide sequences.

Preferably, the second dsRNA is chosen from the group of AL-DP-7301, AL-DP-7308, AL-DP-7318, AL-DP-7320, AL-DP-7322, AL-DP-7324, AL-DP-7325, AL-DP-7326, AL-DP-7327, AL-DP-7329, AL-DP-7331, AL-DP-7333, AL-DP-7340, AL-DP-7342, AL-DP-7303, AL-DP-7305, AL-DP-7307, AL-DP-7309, AL-DP-7316, and AL-DP-7337, AL-DP-7304, AL-DP-5 7312, AL-DP-7339, AL-DP-7344, AL-DP-7306, AL-DP-7317, AL-DP-7346, AL-DP-7310, AL-DP-7323, AL-DP-7335, AL-DP-7338, AL-DP-7341, AL-DP-7302, AL-DP-7315, AL-DP-7328, AL-DP-7330, AL-DP-7336, AL-DP-7345, AL-DP-9250, AL-DP-9251, AL-DP-9252, AL-DP-9253, AL-DP-9254, AL-DP-9255, AL-DP-9256, AL-DP-9257, AL-DP-9258, AL-DP-9259, AL-DP-9260, AL-DP-9261, AL-DP-9262, AL-DP-9263, AL-DP-9264, AL-DP-9265, AL-DP-9266, 10 AL-DP-9267, AL-DP-9268, AL-DP-9269, AL-DP-9270, AL-DP-9271, AL-DP-9272, AL-DP-9273, AL-DP-9274, AL-DP-9275, AL-DP-9276, AL-DP-9277, AL-DP-9279, AL-DP-9280, AL-DP-9281, AL-DP-9282, AL-DP-9283, AL-DP-9284, AL-DP-9285, AL-DP-9286, AL-DP-9287, AL-DP-9288, and AL-DP-9289 (see Table 1 and Table 2). Other embodiments of the vector of the invention are as provided for other aspects of the invention above.

15 In yet another aspect, an isolated cell comprising the above vector is provided. Other embodiments of the cell comprising a vector of the invention are as provided for other aspects of the invention above.

20 **Table 1: RNAi agents for the down-regulation of homo sapiens (NM_012111.1), mus musculus (NM_146036.1) and pan troglodytes (XM_510094.1) Aha 1, and minimal off-target interactions in rat cells; AL-DP-7561, AL-DP-7562, AL-DP-7563 and AL-DP-7564 are additionally cross-reactive to mus musculus (NM_172391.3) and rattus norvegicus (XM_223680.3) Aha 2**

Duplex identifier	Sense strand sequence ¹	SEQ ID NO:	Antisense strand sequence ¹	SEQ ID NO:
AL-DP-7299	auugguccacggauaagcuTT	1	agcuuauccguggaccaaauTT	2
AL-DP-7300	gugaguaagcuugauggagTT	3	cuccaucaagcuacucacTT	4
AL-DP-7301	aguaaaauccccacuuguTT	5	acaagugggauuuugacuTT	6
AL-DP-7302	aaaucucguggccuuuaugTT	7	cauuuaggccacgagauuuTT	8
AL-DP-7303	gagauuagugugagccuugTT	9	caaggcucacacuauucuTT	10
AL-DP-7304	aaucucguggccuuuaugaTT	11	ucauuaaggccacgagauuTT	12
AL-DP-7305	agauuagugugagccuugCTT	13	gcaaggcucacacuauucuTT	14
AL-DP-7306	cggcggacgcccaccaacgTT	15	cguugguggcguccgcccTT	16
AL-DP-7307	ggcggacgcccaccaacguTT	17	gacguugguggcguccgcccTT	18
AL-DP-7308	ggcggacgcccaccaacguTT	19	acguugguggcguccgcccTT	20

Duplex identifier	Sense strand sequence ¹	SEQ ID NO:	Antisense strand sequence ¹	SEQ ID NO:
AL-DP-7309	caacguacaacaacuggcacTT	21	gugccaguuguugacguugTT	22
AL-DP-7310	gcgggcggacggcaccaacTT	23	guugguggcguccgccccgTT	24
AL-DP-7311	aucucguggccuuuaugaaTT	25	uucauuaaggccacgagauTT	26
AL-DP-7312	acguacaacaacuggcacugTT	27	cagugccaguuguugacguuTT	28
AL-DP-7313	accaacguacaacaacuggcTT	29	gccaguuguugacguugguTT	30
AL-DP-7314	acgcugggaucguggaggagTT	31	cuccuccacgauccagcguTT	32
AL-DP-7315	agacccacgcuggaucgugTT	33	cacgauccagcugggguuTT	34
AL-DP-7316	gaccacgcuggaucguggTT	35	ccacgauccagcugggguuTT	36
AL-DP-7317	gaauuuuacauacagcaccuTT	37	aggugcugauguaauuuuucTT	38
AL-DP-7318	gggaaauuuuacauacagcaccTT	39	ggugcugauguaauuuuuccTT	40
AL-DP-7319	uggaaauuuuacauacagcacTT	41	gugcugauguaauuuuuccaTT	42
AL-DP-7320	ccaacguacaacaacuggcaTT	43	ugccaguuguugacguuggTT	44
AL-DP-7321	aaguggggugagggagaccTT	45	ggucuccucaccccacuuTT	46
AL-DP-7322	acacaaaucucguggccuuTT	47	aaggccacgagauuuuguguTT	48
AL-DP-7323	accacgcuggaucguggaTT	49	uccacgauccagcugggguTT	50
AL-DP-7324	gagucaaaaaucccacuugTT	51	caaguggggauuuuugacuTT	52
AL-DP-7325	gagcucuauagaguguuaTT	53	uaaacacucuauagagcucTT	54
AL-DP-7326	ggcagcggguacuacuuugaTT	55	ucaaaguaguaccgcugccTT	56
AL-DP-7327	gacacaaaucucguggccuTT	57	aggccacgagauuuugugucTT	58
AL-DP-7328	agcgggcggacgcaccaaaTT	59	uugguggcguccgccccguTT	60
AL-DP-7329	caaaaaucccccacuuguaagTT	61	cuuacaaguggggauuuuugTT	62
AL-DP-7330	gagaccccacgcuggaucguTT	63	acgauccagcugggguuucTT	64
AL-DP-7331	gagccuugccaaaagaugagTT	65	cucaucuuuggcaaggcucTT	66
AL-DP-7332	ugacacaaaucucguggccTT	67	ggccacgagauuuugugucATT	68
AL-DP-7333	ggagcucuauagaguguuTT	69	aaacacucuauagagcuccTT	70
AL-DP-7334	cccacgcuggaucguggagTT	71	cuccacgauccagcugggguTT	72
AL-DP-7335	gauccccacuuuugucugauTT	73	aucagacaaaauuuuggggauTT	74
AL-DP-7336	gagaucccaacuuuugucugTT	75	cagacaaaauuuuggggauuucTT	76
AL-DP-7337	agccugacacaaaucucguTT	77	acgagauuuugugucaggcucTT	78
AL-DP-7338	agaucccaacuuuugucugaTT	79	ucagacacaaaauuuuggggauuTT	80
AL-DP-7339	aggagacccacgcuggauTT	81	auccagcugggguucccuTT	82
AL-DP-7340	gagggagacccacgcuggaTT	83	uccagcugggguucccuuTT	84
AL-DP-7341	gccaagugggugagggagTT	85	cuccucacccacuuggcTT	86
AL-DP-7342	uggcagcggguacuacuuugTT	87	caaaguaguaccgcugccTT	88
AL-DP-7343	ugagggagacccacgcuggTT	89	ccagcugggguucccuuTT	90
AL-DP-7344	aguggagauuaugugugacTT	91	gcucacacuauacuccacuTT	92
AL-DP-7345	aggagcucuauagaguguTT	93	aacacucuauagagcuccuTT	94
AL-DP-7346	agcggguacuacuuuugagggTT	95	cccucaaaguaguaccgcuTT	96
AL-DP-7561	cgcugggaucguggaggagcTT	97	gcuccuccacgauccagcTT	98
AL-DP-7562	gcuggauctguggaggagcgTT	99	cgccuccuccacgauccagcTT	100
AL-DP-7563	cuggauctguggaggagcgTT	101	ccgcuccuccacgauccagcTT	102
AL-DP-7564	uggaucguggaggagcgTT	103	cccgcuccuccacgauccaTT	104

¹ Capital letters = desoxyribonucleotides; small letters = ribonucleotides

Table 2: RNAi agents for the down-regulation of homo sapiens (NM_012111.1), mus musculus (NM_146036.1) and pan troglodytes (XM_510094.1) Aha 1, and minimal off-target interactions in human cells

Duplex identifier	Sense strand sequence ¹	SEQ ID NO:	Antisense strand sequence ¹	SEQ ID NO:
AL-DP-9250	gccugacacaaaucucguugTT	105	cacgagauuugugugucaggcTT	106
AL-DP-9251	ccugacacaaaucucguuggTT	107	ccacgagauuugugugucaggTT	108
AL-DP-9252	acgcccaccaacguacaacaTT	109	uuguugacguugguggcguTT	110
AL-DP-9253	agcucuauagaguguuuacTT	111	guaaacacucuauagagcuTT	112
AL-DP-9254	ggcuggcagcgguacuacTT	113	guaguaccgcugccagcccTT	114
AL-DP-9255	cuggcagcgguacuacuuuTT	115	aaaguaguaccgcugccaggTT	116
AL-DP-9256	ggaugaaguggagauuaguTT	117	acuaauccacuacuucauccTT	118
AL-DP-9257	accagaggagcucuauagaTT	119	ucuauagagcuccucugguTT	120
AL-DP-9258	aaggaggagauuagugugagTT	121	cucacacuaauccacuuuTT	122
AL-DP-9259	gaggagcucuauagaguguTT	123	acacucuauagagcuccuTT	124
AL-DP-9260	gggagacccacgcuggaucTT	125	gauccagcgugggucucccTT	126
AL-DP-9261	ugagccugacacaaaucucTT	127	gagauuugugugucaggcuTT	128
AL-DP-9262	gcggacgccaccaacgucaTT	129	ugacguugguggcguccgcTT	130
AL-DP-9263	cggacgccaccaacgucaaTT	131	uugacguugguggcguccgTT	132
AL-DP-9264	gaaguggagauuagugugaTT	133	ucacacuaauccacuuuTT	134
AL-DP-9265	cucguggccuuuaaugaaggTT	135	ccuucauuaggccacgagTT	136
AL-DP-9266	ucguggccuuuaaugaaggTT	137	uccuucauuaggccacgatTT	138
AL-DP-9267	aauggggaaauuuuacaucagcTT	139	gcugauguaaaauucccauuTT	140
AL-DP-9268	ggaauuuacaucagcaccTT	141	gggugcugauguaauuuccTT	142
AL-DP-9269	ggagauuagugugagccuuTT	143	aaggcucacacuaauccuccTT	144
AL-DP-9270	cacaaaucucguggccuuTT	145	uaaggccacgagauuugugTT	146
AL-DP-9271	acaaaucucguggccuuuaTT	147	uuuaggccacgagauuuguTT	148
AL-DP-9272	ggagacccacgcuggaucgTT	149	cgauccagcgugggucuccTT	150
AL-DP-9273	ggacgccaccaacgucaacTT	151	guugacguugguggcguccTT	152
AL-DP-9274	gaugaaguggagauuagugTT	153	cacuaauccacuuucauTT	154
AL-DP-9275	gugagccuugccaaagaaugTT	155	caucuuuggcaaggcucactTT	156
AL-DP-9276	caaugaauggagagucaguTT	157	acugacucuccaaaucauugTT	158
AL-DP-9277	auuagugugagccuugccaaTT	159	uggcaaggcucacacuaauTT	160
AL-DP-9278	agaugagccugacacaaaTT	161	auuugugucaggcucaucuTT	162
AL-DP-9279	uagugugagccuugccaaTT	163	uuuggcaaggcucacacuaTT	164
AL-DP-9280	uuugccaccaucaccuugaTT	165	ucaagguguguggggcaaaTT	166
AL-DP-9281	acggagagagaugcuucaaTT	167	uugaagcaucucucuccgTT	168
AL-DP-9282	cggagagagagaugcuucaaTT	169	uuuugaagcaucucucuccgTT	170
AL-DP-9283	aaaauccccacuuguaagaTT	171	ucuuacaaguggggauuuuTT	172
AL-DP-9284	aucccccaauuuugucugaugTT	173	caucagacaaaauuggggauTT	174
AL-DP-9285	ucaaaaauccccacuuguaaaTT	175	uuacaaguggggauuuuugaTT	176
AL-DP-9286	aaauccccacuuguaagauTT	177	aucuuacaaguggggauuuuTT	178
AL-DP-9287	uccccaauuuugucugaugaTT	179	ucaucagacaaaauuggggaaTT	180
AL-DP-9288	augccaaguggggugaggTT	181	ccucaccccacuuggccauTT	182

Duplex identifier	Sense strand sequence ¹	SEQ ID NO:	Antisense strand sequence ¹	SEQ ID NO:
AL-DP-9289	ggagucaaaauccccacuuTT	183	aaguggggauuuugacuccTT	184

¹ Capital letters = desoxyribonucleotides; small letters = ribonucleotides

Brief Description of the Figures

No Figures are presented.

Detailed Description of the Invention

5 The invention provides double-stranded ribonucleic acid (dsRNA), as well as compositions and methods for inhibiting the expression of an Aha gene in a cell or mammal using the dsRNA. The invention also provides compositions and methods for treating pathological conditions and diseases in a mammal caused by the expression of an Aha gene using dsRNA. dsRNA directs the sequence-specific degradation of mRNA through a process
10 known as RNA interference (RNAi).

The dsRNA of the invention comprises an RNA strand (the antisense strand) having a region which is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and is substantially complementary to at least part of an mRNA transcript of an Aha gene. The use of these dsRNAs enables the targeted degradation of mRNAs of genes that are implicated in
15 replication and or maintenance of cancer cells in mammals, and/or in the degradation of misfolded Cystic Fibrosis Transmembrane Conductance Regulator (CFTR). Using cell-based and animal assays, the present inventors have demonstrated that very low dosages of these dsRNA can specifically and efficiently mediate RNAi, resulting in significant inhibition of expression of an Aha gene. Thus, the methods and compositions of the invention comprising
20 these dsRNAs are useful for treating pathological processes mediated by Aha expression, e.g. cancer and/or cystic fibrosis, by targeting a gene involved in protein degradation.

The following detailed description discloses how to make and use the dsRNA and compositions containing dsRNA to inhibit the expression of an Aha gene, as well as compositions and methods for treating diseases and disorders caused by the expression of an Aha
25 gene, such as cancer and/or cystic fibrosis. The pharmaceutical compositions of the invention

comprise a dsRNA having an antisense strand comprising a region of complementarity which is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and is substantially complementary to at least part of an RNA transcript of an Aha gene, together with a pharmaceutically acceptable carrier.

5 Accordingly, certain aspects of the invention provide pharmaceutical compositions comprising the dsRNA of the invention together with a pharmaceutically acceptable carrier, methods of using the compositions to inhibit expression of an Aha gene, and methods of using the pharmaceutical compositions to treat diseases caused by expression of an Aha gene.

Definitions

10 For convenience, the meaning of certain terms and phrases used in the specification, examples, and appended claims, are provided below. If there is an apparent discrepancy between the usage of a term in other parts of this specification and its definition provided in this section, the definition in this section shall prevail.

"G," "C," "A", "T" and "U" (irrespective of whether written in capital or small letters) 15 each generally stand for a nucleotide that contains guanine, cytosine, adenine, thymine, and uracil as a base, respectively. However, it will be understood that the term "ribonucleotide" or "nucleotide" can also refer to a modified nucleotide, as further detailed below, or a surrogate replacement moiety. The skilled person is well aware that guanine, cytosine, adenine, thymine, and uracil may be replaced by other moieties without substantially altering the base pairing 20 properties of an oligonucleotide comprising a nucleotide bearing such replacement moiety. For example, without limitation, a nucleotide comprising inosine as its base may base pair with nucleotides containing adenine, cytosine, or uracil. Hence, nucleotides containing uracil, guanine, or adenine may be replaced in the nucleotide sequences of the invention by a nucleotide containing, for example, inosine. Sequences comprising such replacement moieties are 25 embodiments of the invention.

As used herein, "Aha gene" refers to Activator of Heat Shock Protein 90 ATPase genes. "Ahal" refers to Activator of Heat Shock Protein 90 ATPase 1 genes, non-exhaustive examples of which are found under Genbank accession numbers NM_012111.1 (*Homo sapiens*),

NM_146036.1 (*Mus musculus*), and XM_510094.1 (*Pan troglodytes*). "Aha2" refers to putative Activator of Heat Shock Protein 90 ATPase 2 genes, also known Ahsa2, non-exhaustive examples of which may be found under Genbank accession numbers NM_172391.3 (*Mus musculus*) and XM_223680.3 (*Rattus norvegicus*).

5 As used herein, "target sequence" refers to a contiguous portion of the nucleotide sequence of an mRNA molecule formed during the transcription of an Aha gene, including mRNA that is a product of RNA processing of a primary transcription product. The target sequence of any given RNAi agent of the invention means an mRNA-sequence of X nucleotides that is targeted by the RNAi agent by virtue of the complementarity of the antisense strand of the
10 RNAi agent to such sequence and to which the antisense strand may hybridize when brought into contact with the mRNA, wherein X is the number of nucleotides in the antisense strand plus the number of nucleotides in a single-stranded overhang of the sense strand, if any.

15 As used herein, the term "strand comprising a sequence" refers to an oligonucleotide comprising a chain of nucleotides that is described by the sequence referred to using the standard nucleotide nomenclature.

20 As used herein, and unless otherwise indicated, the term "complementary," when used to describe a first nucleotide sequence in relation to a second nucleotide sequence, refers to the ability of an oligonucleotide or polynucleotide comprising the first nucleotide sequence to hybridize and form a duplex structure under certain conditions with an oligonucleotide or polynucleotide comprising the second nucleotide sequence, as will be understood by the skilled person. Such conditions can, for example, be stringent conditions, where stringent conditions may include: 400 mM NaCl, 40 mM PIPES pH 6.4, 1 mM EDTA, 50°C or 70°C for 12-16 hours followed by washing. Other conditions, such as physiologically relevant conditions as may be encountered inside an organism, can apply. The skilled person will be able to determine the set
25 of conditions most appropriate for a test of complementarity of two sequences in accordance with the ultimate application of the hybridized nucleotides.

This includes base-pairing of the oligonucleotide or polynucleotide comprising the first nucleotide sequence to the oligonucleotide or polynucleotide comprising the second nucleotide

sequence over the entire length of the first and second nucleotide sequence. Such sequences can be referred to as "fully complementary" with respect to each other herein. However, where a first sequence is referred to as "substantially complementary" with respect to a second sequence herein, the two sequences can be fully complementary, or they may form one or more, but 5 generally not more than 4, 3 or 2 mismatched base pairs upon hybridization, while retaining the ability to hybridize under the conditions most relevant to their ultimate application. However, where two oligonucleotides are designed to form, upon hybridization, one or more single stranded overhangs, such overhangs shall not be regarded as mismatches with regard to the determination of complementarity. For example, a dsRNA comprising one oligonucleotide 21 10 nucleotides in length and another oligonucleotide 23 nucleotides in length, wherein the longer oligonucleotide comprises a sequence of 21 nucleotides that is fully complementary to the shorter oligonucleotide, may yet be referred to as "fully complementary" for the purposes of the invention.

"Complementary" sequences, as used herein, may also include, or be formed entirely 15 from, non-Watson-Crick base pairs and/or base pairs formed from non-natural and modified nucleotides, in as far as the above requirements with respect to their ability to hybridize are fulfilled.

The terms "complementary", "fully complementary" and "substantially complementary" herein may be used with respect to the base matching between the sense strand and the antisense 20 strand of a dsRNA, or between the antisense strand of a dsRNA and a target sequence, as will be understood from the context of their use.

As used herein, a polynucleotide which is "substantially complementary to at least part of" a messenger RNA (mRNA) refers to a polynucleotide which is substantially complementary to a contiguous portion of the mRNA of interest (e.g., encoding Aha1). For example, a 25 polynucleotide is complementary to at least a part of an Aha1 mRNA if the sequence is substantially complementary to a non-interrupted portion of an mRNA encoding Aha1.

The term "double-stranded RNA" or "dsRNA", as used herein, refers to a complex of ribonucleic acid molecules, having a duplex structure comprising two anti-parallel and

substantially complementary, as defined above, nucleic acid strands. The two strands forming the duplex structure may be different portions of one larger RNA molecule, or they may be separate RNA molecules. Where the two strands are part of one larger molecule, and therefore are connected by an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5 5'end of the respective other strand forming the duplex structure, the connecting RNA chain is referred to as a “hairpin loop”. Where the two strands are connected covalently by means other than an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'end of the respective other strand forming the duplex structure, the connecting structure is referred to as a “linker”. The RNA strands may have the same or a different number of nucleotides. The 10 maximum number of base pairs is the number of nucleotides in the shortest strand of the dsRNA minus any overhangs that are present in the duplex. In addition to the duplex structure, a dsRNA may comprise one or more nucleotide overhangs.

As used herein, a “nucleotide overhang” refers to the unpaired nucleotide or nucleotides that protrude from the duplex structure of a dsRNA when a 3'-end of one strand of the dsRNA 15 extends beyond the 5'-end of the other strand, or vice versa. “Blunt” or “blunt end” means that there are no unpaired nucleotides at that end of the dsRNA, i.e., no nucleotide overhang. A “blunt ended” dsRNA is a dsRNA that has no nucleotide overhang at either end of the molecule.

The term “antisense strand” refers to the strand of a dsRNA which includes a region that is substantially complementary to a target sequence. As used herein, the term “region of 20 complementarity” refers to the region on the antisense strand that is substantially complementary to a sequence, for example a target sequence, as defined herein. Where the region of complementarity is not fully complementary to the target sequence, the mismatches are most tolerated in the terminal regions and, if present, are generally in a terminal region or regions, *e.g.*, within 6, 5, 4, 3, or 2 nucleotides of the 5' and/or 3' terminus. Most preferably, the 25 mismatches are located within 6, 5, 4, 3, or 2 nucleotides of the 5' terminus of the antisense strand and/or the 3' terminus of the sense strand.

The term “sense strand,” as used herein, refers to the strand of a dsRNA that includes a region that is substantially complementary to a region of the antisense strand.

“Introducing into a cell”, when referring to a dsRNA, means facilitating uptake or absorption into the cell, as is understood by those skilled in the art. Absorption or uptake of dsRNA can occur through unaided diffusive or active cellular processes, or by auxiliary agents or devices. The meaning of this term is not limited to cells in vitro; a dsRNA may also be 5 "introduced into a cell", wherein the cell is part of a living organism. In such instance, introduction into the cell will include the delivery to the organism. For example, for in vivo delivery, dsRNA can be injected into a tissue site or administered systemically. In vitro introduction into a cell includes methods known in the art such as electroporation and lipofection.

10 The terms “silence” and “inhibit the expression of”, in as far as they refer to an Aha gene, e.g. an Aha1 gene, herein refer to the at least partial suppression of the expression of an Aha gene, e.g. an Aha1 gene, as manifested by a reduction of the amount of mRNA transcribed from an Aha gene which may be isolated from a first cell or group of cells in which an Aha gene is transcribed and which has or have been treated such that the expression of an Aha gene is 15 inhibited, as compared to a second cell or group of cells substantially identical to the first cell or group of cells but which has or have not been so treated (control cells). Preferably, the cells are HeLa or MLE 12 cells. The degree of inhibition is usually expressed in terms of

$$\frac{(\text{mRNA in control cells}) - (\text{mRNA in treated cells})}{(\text{mRNA in control cells})} \cdot 100\%$$

Alternatively, the degree of inhibition may be given in terms of a reduction of a 20 parameter that is functionally linked to Aha gene transcription, e.g. the amount of protein encoded by an Aha gene which is secreted by a cell, or found in solution after lysis of such cells, or the number of cells displaying a certain phenotype, e.g. apoptosis or cell surface CFTR. In principle, Aha gene silencing may be determined in any cell expressing the target, either constitutively or by genomic engineering, and by any appropriate assay. However, when a 25 reference is needed in order to determine whether a given dsRNA inhibits the expression of an Aha gene by a certain degree and therefore is encompassed by the instant invention, the assays provided in the Examples below shall serve as such reference.

For example, in certain instances, expression of an Aha gene, e.g. an Aha1 gene, is suppressed by at least about 20%, 25%, 35%, or 50% by administration of the double-stranded oligonucleotide of the invention. In some embodiment, an Aha gene, e.g. an Aha1 gene, is suppressed by at least about 60%, 70%, or 80% by administration of the double-stranded oligonucleotide of the invention. In some embodiments, an Aha gene, e.g. an Aha1 gene, is suppressed by at least about 85%, 90%, or 95% by administration of the double-stranded oligonucleotide of the invention. Table 6 provides values for inhibition of Aha1 expression using various dsRNA molecules of the invention.

As used herein in the context of Aha expression, e.g. Aha1 expression, the terms "treat", "treatment", and the like, refer to relief from or alleviation of pathological processes mediated by Aha expression. In the context of the present invention insofar as it relates to any of the other conditions recited herein below (other than pathological processes mediated by Aha expression), the terms "treat", "treatment", and the like mean to relieve or alleviate at least one symptom associated with such condition, or to slow or reverse the progression of such condition.

As used herein, the phrases "therapeutically effective amount" and "prophylactically effective amount" refer to an amount that provides a therapeutic benefit in the treatment, prevention, or management of pathological processes mediated by Aha expression or an overt symptom of pathological processes mediated by Aha expression. The specific amount that is therapeutically effective can be readily determined by ordinary medical practitioner, and may vary depending on factors known in the art, such as, e.g. the type of pathological processes mediated by Aha expression, the patient's history and age, the stage of pathological processes mediated by Aha expression, and the administration of other anti-pathological processes mediated by Aha expression agents.

As used herein, a "pharmaceutical composition" comprises a pharmacologically effective amount of a dsRNA and a pharmaceutically acceptable carrier. As used herein, "pharmacologically effective amount," "therapeutically effective amount" or simply "effective amount" refers to that amount of an RNA effective to produce the intended pharmacological, therapeutic or preventive result. For example, if a given clinical treatment is considered effective when there is at least a 25% reduction in a measurable parameter associated with a

disease or disorder, a therapeutically effective amount of a drug for the treatment of that disease or disorder is the amount necessary to effect at least a 25% reduction in that parameter.

The term “pharmaceutically acceptable carrier” refers to a carrier for administration of a therapeutic agent. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The term specifically excludes cell culture medium. For drugs administered orally, pharmaceutically acceptable carriers include, but are not limited to pharmaceutically acceptable excipients such as inert diluents, disintegrating agents, binding agents, lubricating agents, sweetening agents, flavoring agents, coloring agents and preservatives. Suitable inert diluents include sodium and calcium carbonate, sodium and calcium phosphate, and lactose, while corn starch and alginic acid are suitable disintegrating agents. Binding agents may include starch and gelatin, while the lubricating agent, if present, will generally be magnesium stearate, stearic acid or talc. If desired, the tablets may be coated with a material such as glyceryl monostearate or glyceryl distearate, to delay absorption in the gastrointestinal tract.

As used herein, a “transformed cell” is a cell into which a vector has been introduced from which a dsRNA molecule may be expressed.

Double-stranded ribonucleic acid (dsRNA)

In one embodiment, the invention provides double-stranded ribonucleic acid (dsRNA) molecules for inhibiting the expression of an Aha gene, e.g. an Aha1 gene, in a cell or mammal, wherein the dsRNA comprises an antisense strand comprising a region of complementarity which is complementary to at least a part of an mRNA formed in the expression of an Aha gene, e.g. an Aha1 gene, and wherein the region of complementarity is less than 30 nucleotides in length, generally 19-24 nucleotides in length. The dsRNA may be identical to one of the dsRNAs shown in Table 1 and Table 2, or it may effect cleavage of an mRNA encoding an Aha gene within the target sequence of one of the dsRNAs shown in Table 1 and Table 2. Preferably, the dsRNA has at least 5, at least 10, at least 15, at least 18, or at least 20 contiguous nucleotides per strand in common with at least one strand, but preferably both strands, of one of the dsRNAs shown in Table 1 and Table 2. Alternative dsRNAs that target elsewhere in the target sequence

of one of the dsRNAs provided in Table 1 and Table 2 can readily be determined using the target sequence and the flanking Aha1 sequence.

The dsRNA comprises two RNA strands that are sufficiently complementary to hybridize to form a duplex structure. One strand of the dsRNA (the antisense strand) comprises a region of complementarity that is substantially complementary, and generally fully complementary, to a target sequence, derived from the sequence of an mRNA formed during the expression of an Aha gene, the other strand (the sense strand) comprises a region which is complementary to the antisense strand, such that the two strands hybridize and form a duplex structure when combined under suitable conditions. Generally, the duplex structure is between 15 and 30, more generally between 18 and 25, yet more generally between 19 and 24, and most generally between 19 and 21 base pairs in length. Similarly, the region of complementarity to the target sequence is between 15 and 30, more generally between 18 and 25, yet more generally between 19 and 24, and most generally between 19 and 21 nucleotides in length. The dsRNA of the invention may further comprise one or more single-stranded nucleotide overhang(s). The dsRNA can be synthesized by standard methods known in the art as further discussed below, e.g., by use of an automated DNA synthesizer, such as are commercially available from, for example, Biosearch, Applied Biosystems, Inc. In a preferred embodiment, an Aha gene is the human Aha1 gene. In specific embodiments, the first strand of the dsRNA comprises the sense sequences of the RNAi agents AL-DP-7301 - AL-DP-7346 and AL-DP-7561 - AL-DP-7564 (Table 1), and AL-DP-9250 - AL-DP-9289 (Table 2), and the second sequence is selected from the group consisting of the antisense sequences of AL-DP-7301 - AL-DP-7346 and AL-DP-7561 - AL-DP-7564 (Table 1) and AL-DP-9250 - AL-DP-9289 (Table 2).

In further embodiments, the dsRNA comprises at least one nucleotide sequence selected from the groups of sequences provided above for the RNAi agents AL-DP-7301 - AL-DP-7346 and AL-DP-7561 - AL-DP-7564 (Table 1) and AL-DP-9250 - AL-DP-9289 (Table 2). In other embodiments, the dsRNA comprises at least two sequences selected from this group, wherein one of the at least two sequences is complementary to another of the at least two sequences, and one of the at least two sequences is substantially complementary to a sequence of an mRNA generated in the expression of an Aha gene, e.g. an Aha1 gene. Generally, the dsRNA comprises two oligonucleotides, wherein one oligonucleotide may be described as the sense strand in one

of the RNAi agents AL-DP-7301 - AL-DP-7346 and AL-DP-7561 - AL-DP-7564 (Table 1), and AL-DP-9250 - AL-DP-9289 (Table 2), and the second oligonucleotide may be described as the antisense strand in one of the RNAi agents AL-DP-7301 - AL-DP-7346 and AL-DP-7561 - AL-DP-7564 (Table 1), and AL-DP-9250 - AL-DP-9289 (Table 2).

5 The skilled person is well aware that dsRNAs comprising a duplex structure of between 20 and 23, but specifically 21, base pairs have been hailed as particularly effective in inducing RNA interference (Elbashir et al., EMBO 2001, 20:6877-6888). However, others have found that shorter or longer dsRNAs can be effective as well. In the embodiments described above, by virtue of the nature of the oligonucleotide sequences provided for the RNAi agents AL-DP-7301
10 - AL-DP-7346 and AL-DP-7561 - AL-DP-7564 (Table 1), and AL-DP-9250 - AL-DP-9289 (Table 2), the dsRNAs of the invention can comprise at least one strand of a length of minimally 21 nt. It can be reasonably expected that shorter dsRNAs comprising one of the sequences provided herein for the RNAi agents AL-DP-7301 - AL-DP-7346 and AL-DP-7561 - AL-DP-7564 (Table 1), and AL-DP-9250 - AL-DP-9289 (Table 2), minus only a few nucleotides on one
15 or both ends may be similarly effective as compared to the dsRNAs described above. Hence, dsRNAs comprising a partial sequence of at least 15, 16, 17, 18, 19, 20, or more contiguous nucleotides from one of the sequences of the RNAi agents AL-DP-7301 - AL-DP-7346 and AL-DP-7561 - AL-DP-7564 (Table 1), and AL-DP-9250 - AL-DP-9289 (Table 2), and differing in their ability to inhibit the expression of an Aha gene, e.g. an Aha1 gene, in a FACS assay as
20 described herein below by not more than 5, 10, 15, 20, 25, or 30 % inhibition from a dsRNA comprising the full sequence, are contemplated by the invention.

Further dsRNAs that cleave within the target sequence of the RNAi agents AL-DP-7301 - AL-DP-7346 and AL-DP-7561 - AL-DP-7564 (Table 1), and AL-DP-9250 - AL-DP-9289 (Table 2), can readily be made using the Aha1 gene sequence and the respective target sequence. The
25 RNAi agents provided in Table 1 and Table 2 identify a site in the Aha1 mRNA that is susceptible to RNAi based cleavage. As such the present invention includes RNAi agents that target within the sequence targeted by one of the agents of the present invention. As used herein a dsRNA is said to target within the sequence of a second dsRNA if the dsRNA cleaves the message anywhere within the mRNA that is complementary to the antisense strand of the second
30 dsRNA. Such a dsRNA will generally have least 5, at least 10, at least 15, at least 18, or at least

20 contiguous nucleotides from one of the sequences provided in Table 1 and Table 2 coupled to additional nucleotide sequences taken from the region contiguous to the selected sequence in an mRNA encoding an Aha gene. For example, the 3'-most 15 nucleotides of the target sequence of AL-DP-7301 combined with the next 6 nucleotides from the target Aha1 gene produces a single 5 strand agent of 21 nucleotides that is based on one of the sequences provided in Table 1 and Table 2.

Preferably, the second dsRNA is chosen from the group of dsRNAs having a certain activity in inhibiting the expression of an Aha gene in a suitable assay, such as the assays described herein. Consequently, in certain preferred embodiments, the second dsRNA is chosen 10 from the group of AL-DP-7301, AL-DP-7308, AL-DP-7318, AL-DP-7320, AL-DP-7322, AL-DP-7324, AL-DP-7325, AL-DP-7326, AL-DP-7327, AL-DP-7329, AL-DP-7331, AL-DP-7333, AL-DP-7340, AL-DP-7342, AL-DP-7303, AL-DP-7305, AL-DP-7307, AL-DP-7309, AL-DP-7316, and AL-DP-7337, AL-DP-7304, AL-DP-7312, AL-DP-7339, AL-DP-7344, AL-DP-7306, AL-DP-7317, AL-DP-7346, AL-DP-7310, AL-DP-7323, AL-DP-7335, AL-DP-7338, AL-DP-15 7341, AL-DP-7302, AL-DP-7315, AL-DP-7328, AL-DP-7330, AL-DP-7336, AL-DP-7345, AL-DP-9250, AL-DP-9251, AL-DP-9252, AL-DP-9253, AL-DP-9254, AL-DP-9255, AL-DP-9256, AL-DP-9257, AL-DP-9258, AL-DP-9259, AL-DP-9260, AL-DP-9261, AL-DP-9262, AL-DP-9263, AL-DP-9264, AL-DP-9265, AL-DP-9266, AL-DP-9267, AL-DP-9268, AL-DP-9269, AL-DP-9270, AL-DP-9271, AL-DP-9272, AL-DP-9273, AL-DP-9274, AL-DP-9275, AL-DP-9276, 20 AL-DP-9277, AL-DP-9279, AL-DP-9280, AL-DP-9281, AL-DP-9282, AL-DP-9283, AL-DP-9284, AL-DP-9285, AL-DP-9286, AL-DP-9287, AL-DP-9288, and AL-DP-9289.

The dsRNA of the invention can contain one or more mismatches to the target sequence. In a preferred embodiment, the dsRNA of the invention contains no more than 3 mismatches. If the antisense strand of the dsRNA contains mismatches to a target sequence, it is preferable that 25 the area of mismatch not be located in the center of the region of complementarity. If the antisense strand of the dsRNA contains mismatches to the target sequence, it is preferable that the mismatch be restricted to 5 nucleotides from either end, for example 5, 4, 3, 2, or 1 nucleotide from either the 5' or 3' end of the region of complementarity, and preferably from the 5'-end. For example, for a 23 nucleotide dsRNA strand which is complementary to a region of an 30 Aha gene, the dsRNA generally does not contain any mismatch within the central 13 nucleotides.

In another embodiment, the antisense strand of the dsRNA does not contain any mismatch in the region from positions 1, or 2, to positions 9, or 10, of the antisense strand (counting 5'-3'). The methods described within the invention can be used to determine whether a dsRNA containing a mismatch to a target sequence is effective in inhibiting the expression of an Aha gene.

5 Consideration of the efficacy of dsRNAs with mismatches in inhibiting expression of an Aha gene is important, especially if the particular region of complementarity in an Aha gene is known to have polymorphic sequence variation within the population.

In one embodiment, at least one end of the dsRNA has a single-stranded nucleotide overhang of 1 to 4, generally 1 or 2 nucleotides. dsRNAs having at least one nucleotide 10 overhang have unexpectedly superior inhibitory properties than their blunt-ended counterparts. Moreover, the present inventors have discovered that the presence of only one nucleotide 15 overhang strengthens the interference activity of the dsRNA, without affecting its overall stability. dsRNA having only one overhang has proven particularly stable and effective *in vivo*, as well as in a variety of cells, cell culture mediums, blood, and serum. Generally, the single- 20 stranded overhang is located at the 3'-terminal end of the antisense strand or, alternatively, at the 3'-terminal end of the sense strand. The dsRNA may also have a blunt end, generally located at the 5'-end of the antisense strand. Such dsRNAs have improved stability and inhibitory activity, thus allowing administration at low dosages, i.e., less than 5 mg/kg body weight of the recipient per day. Generally, the antisense strand of the dsRNA has a nucleotide overhang at the 3'-end, 25 and the 5'-end is blunt. In another embodiment, one or more of the nucleotides in the overhang is replaced with a nucleoside thiophosphate.

In yet another embodiment, the dsRNA is chemically modified to enhance stability. The nucleic acids of the invention may be synthesized and/or modified by methods well established in the art, such as those described in "Current protocols in nucleic acid chemistry", Beaucage, 25 S.L. et al. (Edrs.), John Wiley & Sons, Inc., New York, NY, USA, which is hereby incorporated herein by reference. Specific examples of preferred dsRNA compounds useful in this invention include dsRNAs containing modified backbones or no natural internucleoside linkages. As defined in this specification, dsRNAs having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. 30 For the purposes of this specification, and as sometimes referenced in the art, modified dsRNAs

that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

Preferred modified dsRNA backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

Representative U.S. patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S. Pat. Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,195; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,316; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050, each of which is herein incorporated by reference

Preferred modified dsRNA backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatoms and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts.

Representative U.S. patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S. Pat. Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134;

5,216,141; 5,235,033; 5,64,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and, 5,677,439, each of which is herein incorporated by reference.

5 In other preferred dsRNA mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an dsRNA mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar
10 backbone of an dsRNA is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative U.S. patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further
15 teaching of PNA compounds can be found in Nielsen et al., *Science*, 1991, 254, 1497-1500.

Most preferred embodiments of the invention are dsRNAs with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular CH_2NHCH_2 , $\text{CH}_2\text{N}(\text{CH}_3)\text{OCH}_2$ [known as a methylene (methylimino) or MMI backbone], $\text{CH}_2\text{O.N}(\text{CH}_3)\text{CH}_2$, $\text{CH}_2\text{N}(\text{CH}_3)\text{N}(\text{CH}_3)\text{CH}_2$ and $\text{N}(\text{CH}_3)\text{CH}_2\text{CH}_2$ [wherein the native
20 phosphodiester backbone is represented as O.P.O.CH_2] of the above-referenced U.S. Pat. No. 5,489,677, and the amide backbones of the above-referenced U.S. Pat. No. 5,602,240. Also preferred are dsRNAs having morpholino backbone structures of the above-referenced U.S. Pat. No. 5,034,506.

Modified dsRNAs may also contain one or more substituted sugar moieties. Preferred
25 dsRNAs comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Particularly preferred are O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. Other preferred dsRNAs

comprise one of the following at the 2' position: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an dsRNA, or a group for improving the pharmacodynamic properties of an dsRNA, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., Helv. Chim. Acta, 1995, 78, 486-504) i.e., an alkoxy-alkoxy group. A further preferred modification includes 2'-dimethylaminoxyethoxy, i.e., a 5 O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE, as described in examples hereinbelow, and 10 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'- DMAEOE), i.e., 2'-O-CH₂-O-CH₂-N(CH₃)₂, also described in examples hereinbelow.

Other preferred modifications include 2'-methoxy (2'-OCH₃), 2'-aminopropoxy (2'-OCH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications may also be made at other 15 positions on the dsRNA, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked dsRNAs and the 5' position of 5' terminal nucleotide. DsRNAs may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative U.S. patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S. Pat. Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 20 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

DsRNAs may also include nucleobase (often referred to in the art simply as "base") 25 modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives 30 of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and

cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl anal other 8-substituted adenines and guanines, 5-halo, particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-daazaadenine and 3-deazaguanine and 3-deazaadenine.

Further nucleobases include those disclosed in U.S. Pat. No. 3,687,808, those disclosed in The Concise Encyclopedia Of Polymer Science And Engineering, pages 858-859, Kroschwitz, J. L, ed. John Wiley & Sons, 1990, these disclosed by Englisch et al., Angewandte Chemie, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y S., Chapter 15, DsRNA

10 Research and Applications, pages 289-302, Crooke, S. T. and Lebleu, B., Ed., CRC Press, 1993.

Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and 0-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2.degree. C. (Sanghvi, Y. S., Crooke, S. T. and Lebleu, B., Eds., DsRNA Research and Applications, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative U.S. patents that teach the preparation of certain of the above noted

20 modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. Pat. No. 3,687,808, as well as U.S. Pat. Nos. 4,845,205; 5,130,30; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; and 5,681,941, each of which is herein incorporated by reference, and U.S. Pat. No. 5,750,692, also herein incorporated by reference.

25 Another modification of the dsRNAs of the invention involves chemically linking to the dsRNA one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the dsRNA. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., Proc. Natl. Acid. Sci. USA, 199, 86, 6553-6556), cholic acid (Manoharan et al., Biorg. Med. Chem. Let., 1994 4 1053-1060), a thioether, e.g., beryl-S-tritylthiol (Manoharan et al., Ann. N.Y. Acad. Sci., 1992, 660, 306-309; Manoharan et al., Biorg.

Med. Chem. Let., 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., Nucl. Acids Res., 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., EMBO J, 1991, 10, 1111-1118; Kabanov et al., FEBS Lett., 1990, 259, 327-330; Svinarchuk et al., Biochimie, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or 5 triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654; Shea et al., Nucl. Acids Res., 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., Nucleosides & Nucleotides, 1995, 14, 969-973), or adamantan acetic acid (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., Biochim. Biophys. Acta, 1995, 1264, 229-237), or an 10 octadecylamine or hexylamino-carbonyloxycholesterol moiety (Crooke et al., J. Pharmacol. Exp. Ther., 1996, 277, 923-937).

Representative U.S. patents that teach the preparation of such dsRNA conjugates include, but are not limited to, U.S. Pat. Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 15 5,545,730; 5,552,538; 5,578,717, 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 20 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241, 5,391,723; 5,416,203, 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, each of which is herein incorporated by reference.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an dsRNA. The present invention also includes 25 dsRNA compounds which are chimeric compounds. "Chimeric" dsRNA compounds or "chimeras," in the context of this invention, are dsRNA compounds, particularly dsRNAs, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an dsRNA compound. These dsRNAs typically contain at least one region wherein the dsRNA is modified so as to confer upon the dsRNA increased resistance to 30 nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target

nucleic acid. An additional region of the dsRNA may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of 5 dsRNA inhibition of gene expression. Consequently, comparable results can often be obtained with shorter dsRNAs when chimeric dsRNAs are used, compared to phosphorothioate deoxydsRNAs hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

10 In certain instances, the dsRNA may be modified by a non-ligand group. A number of non-ligand molecules have been conjugated to dsRNAs in order to enhance the activity, cellular distribution or cellular uptake of the dsRNA, and procedures for performing such conjugations are available in the scientific literature. Such non-ligand moieties have included lipid moieties, such as cholesterol (Letsinger et al., Proc. Natl. Acad. Sci. USA, 1989, 86:6553), cholic acid 15 (Manoharan et al., Bioorg. Med. Chem. Lett., 1994, 4:1053), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., Ann. N.Y. Acad. Sci., 1992, 660:306; Manoharan et al., Bioorg. Med. Chem. Lett., 1993, 3:2765), a thiocholesterol (Oberhauser et al., Nucl. Acids Res., 1992, 20:533), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., EMBO J., 1991, 10:111; Kabanov et al., FEBS Lett., 1990, 259:327; Svinarchuk et al., Biochimie, 1993, 75:49), a 20 phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., Tetrahedron Lett., 1995, 36:3651; Shea et al., Nucl. Acids Res., 1990, 18:3777), a polyamine or a polyethylene glycol chain (Manoharan et al., Nucleosides & Nucleotides, 1995, 14:969), or adamantane acetic acid (Manoharan et al., Tetrahedron Lett., 1995, 36:3651), a palmityl moiety (Mishra et al., Biochim. Biophys. Acta, 25 1995, 1264:229), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., J. Pharmacol. Exp. Ther., 1996, 277:923). Representative United States patents that teach the preparation of such dsRNA conjugates have been listed above. Typical conjugation protocols involve the synthesis of dsRNAs bearing an aminolinker at one or more positions of the sequence. The amino group is then reacted with the molecule being conjugated using appropriate 30 coupling or activating reagents. The conjugation reaction may be performed either with the

dsRNA still bound to the solid support or following cleavage of the dsRNA in solution phase. Purification of the dsRNA conjugate by HPLC typically affords the pure conjugate.

Vector encoded RNAi agents

The dsRNA of the invention can also be expressed from recombinant viral vectors 5 intracellularly in vivo. The recombinant viral vectors of the invention comprise sequences encoding the dsRNA of the invention and any suitable promoter for expressing the dsRNA sequences. Suitable promoters include, for example, the U6 or H1 RNA pol III promoter sequences and the cytomegalovirus promoter. Selection of other suitable promoters is within the skill in the art. The recombinant viral vectors of the invention can also comprise inducible or 10 regulatable promoters for expression of the dsRNA in a particular tissue or in a particular intracellular environment. The use of recombinant viral vectors to deliver dsRNA of the invention to cells in vivo is discussed in more detail below.

dsRNA of the invention can be expressed from a recombinant viral vector either as two 15 separate, complementary RNA molecules, or as a single RNA molecule with two complementary regions.

Any viral vector capable of accepting the coding sequences for the dsRNA molecule(s) to be expressed can be used, for example vectors derived from adenovirus (AV); adeno-associated virus (AAV); retroviruses (e.g., lentiviruses (LV), Rhabdoviruses, murine leukemia virus); herpes virus, and the like. The tropism of viral vectors can be modified by pseudotyping the vectors 20 with envelope proteins or other surface antigens from other viruses, or by substituting different viral capsid proteins, as appropriate.

For example, lentiviral vectors of the invention can be pseudotyped with surface proteins from vesicular stomatitis virus (VSV), rabies, Ebola, Mokola, and the like. AAV vectors of the invention can be made to target different cells by engineering the vectors to express different 25 capsid protein serotypes. For example, an AAV vector expressing a serotype 2 capsid on a serotype 2 genome is called AAV 2/2. This serotype 2 capsid gene in the AAV 2/2 vector can be replaced by a serotype 5 capsid gene to produce an AAV 2/5 vector. Techniques for constructing AAV vectors which express different capsid protein serotypes are within the skill in the art; see,

e.g., Rabinowitz J E et al. (2002), J Virol 76:791-801, the entire disclosure of which is herein incorporated by reference.

Selection of recombinant viral vectors suitable for use in the invention, methods for inserting nucleic acid sequences for expressing the dsRNA into the vector, and methods of delivering the viral vector to the cells of interest are within the skill in the art. See, for example, Dornburg R (1995), Gene Therap. 2: 301-310; Eglitis M A (1988), Biotechniques 6: 608-614; Miller A D (1990), Hum Gene Therap. 1: 5-14; Anderson W F (1998), Nature 392: 25-30; and Rubinson D A et al., Nat. Genet. 33: 401-406, the entire disclosures of which are herein incorporated by reference.

Preferred viral vectors are those derived from AV and AAV. In a particularly preferred embodiment, the dsRNA of the invention is expressed as two separate, complementary single-stranded RNA molecules from a recombinant AAV vector comprising, for example, either the U6 or H1 RNA promoters, or the cytomegalovirus (CMV) promoter.

A suitable AV vector for expressing the dsRNA of the invention, a method for constructing the recombinant AV vector, and a method for delivering the vector into target cells, are described in Xia H et al. (2002), Nat. Biotech. 20: 1006-1010.

Suitable AAV vectors for expressing the dsRNA of the invention, methods for constructing the recombinant AV vector, and methods for delivering the vectors into target cells are described in Samulski R et al. (1987), J. Virol. 61: 3096-3101; Fisher K J et al. (1996), J. Virol. 70: 520-532; Samulski R et al. (1989), J. Virol. 63: 3822-3826; U.S. Pat. No. 5,252,479; U.S. Pat. No. 5,139,941; International Patent Application No. WO 94/13788; and International Patent Application No. WO 93/24641, the entire disclosures of which are herein incorporated by reference.

Pharmaceutical compositions comprising dsRNA

In one embodiment, the invention provides pharmaceutical compositions comprising a dsRNA, as described herein, and a pharmaceutically acceptable carrier. The pharmaceutical composition comprising the dsRNA is useful for treating a disease or disorder associated with

the expression or activity of an Aha gene, such as pathological processes mediated by Aha1 expression. Such pharmaceutical compositions are formulated based on the mode of delivery. One example is compositions that are formulated for systemic administration via parenteral delivery.

5 The pharmaceutical compositions of the invention are administered in dosages sufficient to inhibit expression of an Aha gene. The present inventors have found that, because of their improved efficiency, compositions comprising the dsRNA of the invention can be administered at surprisingly low dosages. A maximum dosage of 5 mg dsRNA per kilogram body weight of recipient per day is sufficient to inhibit or completely suppress expression of an Aha gene.

10 In general, a suitable dose of dsRNA will be in the range of 0.01 microgram to 5.0 milligrams per kilogram body weight of the recipient per day, generally in the range of 1 microgram to 1 mg per kilogram body weight per day. The pharmaceutical composition may be administered once daily, or the dsRNA may be administered as two, three, or more sub-doses at appropriate intervals throughout the day or even using continuous infusion or delivery through a 15 controlled release formulation. In that case, the dsRNA contained in each sub-dose must be correspondingly smaller in order to achieve the total daily dosage. The dosage unit can also be compounded for delivery over several days, e.g., using a conventional sustained release formulation which provides sustained release of the dsRNA over a several day period. Sustained release formulations are well known in the art and are particularly useful for vaginal delivery of 20 agents, such as could be used with the agents of the present invention. In this embodiment, the dosage unit contains a corresponding multiple of the daily dose.

25 The skilled artisan will appreciate that certain factors may influence the dosage and timing required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of a composition can include a single treatment or a series of treatments. Estimates of effective dosages and in vivo half-lives for the individual dsRNAs encompassed by the invention can be made using conventional methodologies or on the basis of in vivo testing using an appropriate animal model, as described elsewhere herein.

Advances in mouse genetics have generated a number of mouse models for the study of various human diseases, such as pathological processes mediated by Aha expression. Such models are used for in vivo testing of dsRNA, as well as for determining a therapeutically effective dose.

5 The present invention also includes pharmaceutical compositions and formulations which include the dsRNA compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical, pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, 10 intranasal, epidermal and transdermal, oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and 15 powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful. Preferred topical formulations include those in which the dsRNAs of the invention are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Preferred lipids and liposomes include neutral (e.g. 20 dioleoylphosphatidyl ethanolamine = DOPE, dimyristoylphosphatidyl choline = DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol = DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl = DOTAP and dioleoylphosphatidyl ethanolamine = DOTMA). DsRNAs of the invention may be encapsulated within liposomes or may form 25 complexes thereto, in particular to cationic liposomes. Alternatively, dsRNAs may be complexed to lipids, in particular to cationic lipids. Preferred fatty acids and esters include but are not limited arachidonic acid, oleic acid, eicosanoic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a C₁₋₁₀ alkyl ester (e.g. isopropylmyristate IPM), monoglyceride, diglyceride 30 or pharmaceutically acceptable salt thereof. Topical formulations are described in detail in U.S.

patent application Ser. No. 09/315,298 filed on May 20, 1999 which is incorporated herein by reference in its entirety.

Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, 5 capsules, gel capsules, sachets, tablets or minitablets. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable. Preferred oral formulations are those in which dsRNAs of the invention are administered in conjunction with one or more penetration enhancers, surfactants, and chelators. Preferred surfactants include fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Preferred bile acids/salts include chenodeoxycholic acid (CDCA) and ursodeoxychenodeoxycholic acid (UDCA), cholic acid, dehydrocholic acid, deoxycholic acid, glucholic acid, glycholic acid, glycocodeoxycholic acid, taurocholic acid, taurodeoxycholic acid, sodium tauro-24,25-dihydro-fusidate and sodium glycodihydrofusidate. Preferred fatty acids include arachidonic acid, undecanoic acid, oleic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, 10 tricaprate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a monoglyceride, a diglyceride or a pharmaceutically acceptable salt thereof (e.g. sodium). Also preferred are combinations of penetration enhancers, for 15 example, fatty acids/salts in combination with bile acids/salts. A particularly preferred combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether. DsRNAs of the invention may be delivered orally, in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. DsRNA complexing agents include poly-amino acids; polyimines; polyacrylates; polyalkylacrylates, polyoxethanes, polyalkylcyanoacrylates; cationized gelatins, albumins, starches, acrylates, polyethyleneglycols (PEG) and starches; 20 polyalkylcyanoacrylates; DEAE-derivatized polyimines, pollulans, celluloses and starches. Particularly preferred complexing agents include chitosan, N-trimethylchitosan, poly-L-lysine, polyhistidine, polyornithine, polyspermines, protamine, polyvinylpyridine, polythiodiethylaminomethylethylene P(TDAE), polyaminostyrene (e.g. p-amino), poly(methylcyanoacrylate), poly(ethylcyanoacrylate), poly(butylcyanoacrylate), 25 poly(isobutylcyanoacrylate), poly(isohexylcynaoacrylate), DEAE-methacrylate, DEAE- 30 poly(isobutylcyanoacrylate), poly(isohexylcynaoacrylate), DEAE-methacrylate, DEAE-

hexylacrylate, DEAE-acrylamide, DEAE-albumin and DEAE-dextran, polymethylacrylate, polyhexylacrylate, poly(D,L-lactic acid), poly(DL-lactic-co-glycolic acid (PLGA), alginate, and polyethyleneglycol (PEG). Oral formulations for dsRNAs and their preparation are described in detail in U.S. application. Ser. No. 08/886,829 (filed Jul. 1, 1997), Ser. No. 09/108,673 (filed Jul. 5, 1998), Ser. No. 09/256,515 (filed Feb. 23, 1999), Ser. No. 09/082,624 (filed May 21, 1998) and Ser. No. 09/315,298 (filed May 20, 1999), each of which is incorporated herein by reference in their entirety.

Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents 10 and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

Pharmaceutical compositions of the present invention include, but are not limited to, 15 solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into 20 association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

The compositions of the present invention may be formulated into any of many possible 25 dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The 5 preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

Emulsions

The compositions of the present invention may be prepared and formulated as emulsions. 10 Emulsions are typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1 μm in diameter (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage* 15 *Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pa., 1985, p. 301). Emulsions are often biphasic systems comprising two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be of either the water-in-oil (w/o) or the oil-in-water (o/w) variety. When an aqueous phase is finely 20 divided into and dispersed as minute droplets into a bulk oily phase, the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase, the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases, and the active drug which may be present as a solution in either the aqueous 25 phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary 30 emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion

enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous phase provides an o/w/o emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase

5 and maintained in this form through the means of emulsifiers or the viscosity of the formulation.

Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers,

10 absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*,

Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger, in

15 *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199).

Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the

20 hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting

surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

25 Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include

polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

5 A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in Pharmaceutical Dosage Forms, Lieberman, 10 Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose ethers, 15 and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external phase.

Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these formulations 20 often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated 25 hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in Pharmaceutical

Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of ease of formulation, as well as efficacy from an absorption and bioavailability standpoint (Rosoff, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 5 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

10 In one embodiment of the present invention, the compositions of dsRNAs and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems 15 that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: Controlled Release of Drugs: Polymers and Aggregate Systems, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). 20 Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon 25 tails of the surfactant molecules (Schott, in Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pa., 1985, p. 271).

The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker 30 (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, in Pharmaceutical

Dosage Forms, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

5 Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750),
10 decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules.
15 Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C₈-C₁₂) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters,
20 fatty alcohols, polyglycolized glycerides, saturated polyglycolized C₈-C₁₀ glycerides, vegetable oils and silicone oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al.,
25 Pharmaceutical Research, 1994, 11, 1385-1390; Ritschel, Meth. Find. Exp. Clin. Pharmacol., 1993, 13, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity
30 (Constantinides et al., Pharmaceutical Research, 1994, 11, 1385; Ho et al., J. Pharm. Sci., 1996,

85, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or dsRNAs. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications.

5 It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of dsRNAs and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of dsRNAs and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

Microemulsions of the present invention may also contain additional components and 10 additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the dsRNAs and nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories.surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., Critical Reviews in 15 Therapeutic Drug Carrier Systems, 1991, p. 92). Each of these classes has been discussed above.

Liposomes

There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity 20 and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

Liposomes are unilamellar or multilamellar vesicles which have a membrane formed 25 from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages *in vivo*.

In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

5 Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important
10 considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

15 Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes and as the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

20 Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

25 Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis

Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex.

The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., Biochem. Biophys. Res. Commun., 1987, 147, 980-985).

5 Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the
10 target cells (Zhou et al., Journal of Controlled Release, 1992, 19, 269-274).

One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while
15 anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

Several studies have assessed the topical delivery of liposomal drug formulations to the
20 skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner et al., Journal of Drug Targeting, 1992, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the
25 liposomal formulation was superior to aqueous administration (du Plessis et al., Antiviral Research, 1992, 18, 259-265).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and

cholesterol. Non-ionic liposomal formulations comprising Novasome.TM. I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome.TM. II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective 5 in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu et al. S.T.P. Pharma. Sci., 1994, 4, 6, 466).

Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized 10 lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G_m1, or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, 15 sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., FEBS Letters, 1987, 223, 42; Wu et al., Cancer Research, 1993, 53, 3765).

Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (Ann. N.Y. Acad. Sci., 1987, 507, 64) reported the ability of 20 monosialoganglioside G_m1, galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (Proc. Natl. Acad. Sci. U.S.A., 1988, 85, 6949). U.S. Pat. No. 4,837,028 and WO 88/04924, both to Allen et al., disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside G_m1 or a galactocerebroside sulfate ester. U.S. Pat. No. 5,543,152 (Webb et al.) discloses liposomes 25 comprising sphingomyelin. Liposomes comprising 1,2-sn-dimyristoylphosphat-idylcholine are disclosed in WO 97/13499 (Lim et al.).

Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (Bull. Chem. Soc. Jpn., 1980, 53, 2778) described liposomes comprising a nonionic detergent, 2C₁₂I₅G, that contains a

PEG moiety. Illum et al. (FEBS Lett., 1984, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Pat. Nos. 4,426,330 and 4,534,899). Klibanov et al. (FEBS Lett., 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (Biochimica et Biophysica Acta, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle et al. (U.S. Pat. Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Pat. No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Pat. No. 5,225,212 (both to Martin et al.) and in WO 94/20073 (Zalipsky et al.) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al). U.S. Pat. No. 5,540,935 (Miyazaki et al.) and U.S. Pat. No. 5,556,948 (Tagawa et al.) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

20 A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry et al. discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Pat. No. 5,264,221 to Tagawa et al. discloses protein-bonded liposomes and asserts that the contents of such liposomes may include dsRNA. U.S. Pat. No. 5,665,710 to Rahman et al. describes certain methods of encapsulating oligodeoxynucleotides in liposomes. 25 WO 97/04787 to Love et al. discloses liposomes comprising dsRNAs targeted to the raf gene.

Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the 30 environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores

in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as 5 effective as subcutaneous injection of a solution containing serum albumin.

Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the 10 "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, N.Y., 1988, p. 285).

If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a 15 wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this 20 class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates 25 and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

5 If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, N.Y., 1988, p. 285).

10 Penetration Enhancers

In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly dsRNAs, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-15 lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., 20 *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, 25 with the result that absorption of dsRNAs through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee et al., *Critical Reviews*

in Therapeutic Drug Carrier Systems, 1991, p.92); and perfluorochemical emulsions, such as FC-43 (Takahashi et al., J. Pharm. Pharmacol., 1988, 40, 252).

Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C₁ - C₁₀ alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and di-glycerides thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., Critical Reviews in Therapeutic Drug 5 Carryier Systems, 1991, p.92; Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1-33; El Hariri et al., J. Pharm. Pharmacol., 1992, 44, 651-654).

Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's The Pharmacological Basis of Therapeutics, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 15 1996, pp. 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), gluchoic acid (sodium gluchoate), glycholic acid (sodium glycocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodieshydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., Critical Reviews in Therapeutic Drug Carrier 20 Systems, 1991, page 92; Swinyard, Chapter 39 In: Remington's Pharmaceutical Sciences, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, Pa., 1990, pages 782-783; Muranishi, Critical 25 Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1-33; Yamamoto et al., J. Pharm. Exp. Ther., 1992, 263, 25; Yamashita et al., J. Pharm. Sci., 1990, 79, 579-583).

Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of dsRNAs through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the 5 added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, J. Chromatogr., 1993, 618, 315-339). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), N-acyl derivatives of collagen, lauroth-9 and N-amino 10 acyl derivatives of beta-diketones (enamines)(Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, page 92; Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1-33; Buur et al., J. Control Rel., 1990, 14, 43-51).

Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as 15 chelating agents or as surfactants but that nonetheless enhance absorption of dsRNAs through the alimentary mucosa (Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1-33). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, page 92); and non-steroidal anti-inflammatory agents such as 20 diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., J. Pharm. Pharmacol., 1987, 39, 621-626).

Agents that enhance uptake of dsRNAs at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Pat. No. 5,705,188), cationic glycerol derivatives, and 25 polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of dsRNAs.

Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

Carriers

Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (i.e., does not possess biological activity per se) but is recognized 5 as a nucleic acid by in vivo processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, 10 presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate dsRNA in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'isothiocyanostilbene-2,2'-disulfonic acid (Miyao et al., Antisense Res. Dev., 1995, 5, 115-121; Takakura et al., Antisense & Nucl. Acid Drug Dev., 1996, 6, 177-183).

15 Excipients

In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the 20 desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); 25 lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (e.g., starch, sodium starch glycolate, etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, 5 amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or 10 solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with 15 nucleic acids can be used.

Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt 15 solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Pharmaceutical compositions for the delivery to the respiratory tract

Another aspect of the invention provides for the delivery of iRNA agents to the respiratory tract, particularly for the treatment of cystic fibrosis. The respiratory tract includes the upper airways, including the oropharynx and larynx, followed by the lower airways, which 20 include the trachea followed by bifurcations into the bronchi and bronchioli. The upper and lower airways are called the conductive airways. The terminal bronchioli then divide into respiratory bronchioli which then lead to the ultimate respiratory zone, the alveoli, or deep lung. The deep lung, or alveoli, are the primary target of inhaled therapeutic aerosols for systemic delivery of iRNA agents.

25 Pulmonary delivery compositions can be delivered by inhalation by the patient of a dispersion so that the composition, preferably the iRNA agent, within the dispersion can reach the lung where it can, for example, be readily absorbed through the alveolar region directly into

blood circulation. Pulmonary delivery can be effective both for systemic delivery and for localized delivery to treat diseases of the lungs.

Pulmonary delivery can be achieved by different approaches, including the use of nebulized, aerosolized, micellular and dry powder-based formulations; administration by inhalation may be oral and/or nasal. Delivery can be achieved with liquid nebulizers, aerosol-based inhalers, and dry powder dispersion devices. Metered-dose devices are preferred. One of the benefits of using an atomizer or inhaler is that the potential for contamination is minimized because the devices are self contained. Dry powder dispersion devices, for example, deliver drugs that may be readily formulated as dry powders. An iRNA composition may be stably stored as lyophilized or spray-dried powders by itself or in combination with suitable powder carriers. The delivery of a composition for inhalation can be mediated by a dosing timing element which can include a timer, a dose counter, time measuring device, or a time indicator which when incorporated into the device enables dose tracking, compliance monitoring, and/or dose triggering to a patient during administration of the aerosol medicament.

Examples of pharmaceutical devices for aerosol delivery include metered dose inhalers (MDIs), dry powder inhalers (DPIs), and air-jet nebulizers. Exemplary delivery systems by inhalation which can be readily adapted for delivery of the subject iRNA agents are described in, for example, U.S. Pat. Nos. 5,756,353; 5,858,784; and PCT applications WO98/31346; WO98/10796; WO00/27359; WO01/54664; WO02/060412. Other aerosol formulations that may be used for delivering the iRNA agents are described in U.S. Pat. Nos. 6,294,153; 6,344,194; 6,071,497, and PCT applications WO02/066078; WO02/053190; WO01/60420; WO00/66206. Further, methods for delivering iRNA agents can be adapted from those used in delivering other oligonucleotides (e.g., an antisense oligonucleotide) by inhalation, such as described in Templin et al., *Antisense Nucleic Acid Drug Dev*, 2000, 10:359-68; Sandrasagra et al., *Expert Opin Biol Ther*, 2001, 1:979-83; Sandrasagra et al., *Antisense Nucleic Acid Drug Dev*, 2002, 12:177-81.

The delivery of the inventive agents may also involve the administration of so called "pro-drugs", i.e. formulations or chemical modifications of a therapeutic substance that require some form of processing or transport by systems innate to the subject organism to release the therapeutic substance, preferably at the site where its action is desired; this latter embodiment

may be used in conjunction with delivery of the respiratory tract, but also together with other embodiments of the present invention. For example, the human lungs can remove or rapidly degrade hydrolytically cleavable deposited aerosols over periods ranging from minutes to hours. In the upper airways, ciliated epithelia contribute to the "mucociliary excalator" by which 5 particles are swept from the airways toward the mouth. Pavia, D., "Lung Mucociliary Clearance," in *Aerosols and the Lung: Clinical and Experimental Aspects*, Clarke, S. W. and Pavia, D., Eds., Butterworths, London, 1984. In the deep lungs, alveolar macrophages are capable of phagocytosing particles soon after their deposition. Warheit et al. *Microscopy Res. Tech.*, 26: 412-422 (1993); and Brain, J. D., "Physiology and Pathophysiology of Pulmonary 10 Macrophages," in *The Reticuloendothelial System*, S. M. Reichard and J. Filkins, Eds., Plenum, New. York., pp. 315-327, 1985.

In preferred embodiments, particularly where systemic dosing with the iRNA agent is desired, the aerosoled iRNA agents are formulated as microparticles. Microparticles having a diameter of between 0.5 and ten microns can penetrate the lungs, passing through most of the 15 natural barriers. A diameter of less than ten microns is required to bypass the throat; a diameter of 0.5 microns or greater is required to avoid being exhaled.

Other Components

The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage 20 levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. 25 However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings

and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The 5 suspension may also contain stabilizers.

Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more dsRNA agents and (b) one or more other chemotherapeutic agents which function by a non-RNA interference mechanism. Examples of such chemotherapeutic agents include but are not limited to daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, 10 idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bis-chloroethylnitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone, tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylcyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 15 5-azacytidine, hydroxyurea, deoxycoformycin, 4-hydroxyperoxycyclophosphoramide, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUDR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, The Merck Manual of Diagnosis and Therapy, 15th Ed. 1987, pp. 1206-1228, Berkow et al., eds., Rahway, N.J. When 20 used with the compounds of the invention, such chemotherapeutic agents may be used individually (e.g., 5-FU and oligonucleotide), sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide). Anti-inflammatory drugs, including but not limited to nonsteroidal anti- 25 inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, The Merck Manual of Diagnosis and Therapy, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-dsRNA 30 chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the 5 therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds which exhibit high therapeutic indices are preferred.

The data obtained from cell culture assays and animal studies can be used in formulation a range of dosage for use in humans. The dosage of compositions of the invention lies generally within a range of circulating concentrations that include the ED50 with little or no toxicity. The 10 dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range of the compound or, when appropriate, of the polypeptide product of a target sequence (e.g., achieving 15 a decreased concentration of the polypeptide) that includes the IC50 (i.e., the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

In addition to their administration individually or as a plurality, as discussed above, the 20 dsRNAs of the invention can be administered in combination with other known agents effective in treatment of pathological processes mediated by Aha expression. In any event, the administering physician can adjust the amount and timing of dsRNA administration on the basis of results observed using standard measures of efficacy known in the art or described herein.

Methods for treating diseases caused by expression of an Aha gene

25 The invention relates in particular to the use of a dsRNA or a pharmaceutical composition prepared therefrom for the treatment of Cystic Fibrosis. Owing to the inhibitory effect on Aha1 expression, an dsRNA according to the invention or a pharmaceutical composition prepared therefrom can enhance the quality of life of Cystic Fibrosis patients.

Furthermore, the invention relates to the use of a dsRNA or a pharmaceutical composition of the invention aimed at the treatment of cancer, e.g., for inhibiting tumor growth and tumor metastasis. For example, the dsRNA or a pharmaceutical composition prepared therefrom may be used for the treatment of solid tumors, like breast cancer, lung cancer, head and neck cancer, brain cancer, abdominal cancer, colon cancer, colorectal cancer, esophagus cancer, gastrointestinal cancer, glioma, liver cancer, tongue cancer, neuroblastoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, retinoblastoma, Wilm's tumor, multiple myeloma and for the treatment of skin cancer, like melanoma, for the treatment of lymphomas and blood cancer. The invention further relates to the use of an dsRNA according to 5 the invention or a pharmaceutical composition prepared therefrom for inhibiting Aha1 expression and/or for inhibiting accumulation of ascites fluid and pleural effusion in different types of cancer, e.g., breast cancer, lung cancer, head cancer, neck cancer, brain cancer, abdominal cancer, colon cancer, colorectal cancer, esophagus cancer, gastrointestinal cancer, glioma, liver cancer, tongue cancer, neuroblastoma, osteosarcoma, ovarian cancer, pancreatic 10 cancer, prostate cancer, retinoblastoma, Wilm's tumor, multiple myeloma, skin cancer, melanoma, lymphomas and blood cancer. Owing to the inhibitory effect on Aha1 expression, an dsRNA according to the invention or a pharmaceutical composition prepared therefrom can 15 enhance the quality of life of cancer patients.

The invention furthermore relates to the use of an dsRNA or a pharmaceutical composition thereof, e.g., for treating Cystic Fibrosis or cancer or for preventing tumor metastasis, in combination with other pharmaceuticals and/or other therapeutic methods, e.g., with known pharmaceuticals and/or known therapeutic methods, such as, for example, those which are currently employed for treating Cystic Fibrosis or cancer and/or for preventing tumor metastasis. Where the pharmaceutical composition aims for the treatment of Cystic fibrosis, 20 preference is given to a combination with daily chest physiotherapy, orally applied pancreatic enzymes, daily oral or inhaled antibiotics to counter lung infection, inhaled anti-asthma therapy, corticosteroid tablets, dietary vitamin supplements, especially A and D, inhalation of PulmozymeTM, medicines to relieve constipation or to improve the activity of the enzyme 25 supplements, insulin for CF-related diabetes, medication for CF-associated liver disease, and 30 oxygen to help with breathing.

Where the pharmaceutical composition aims for the treatment of cancer and/or for preventing tumor metastasis, preference is given to a combination with radiation therapy and chemotherapeutic agents, such as cisplatin, cyclophosphamide, 5-fluorouracil, adriamycin, daunorubicin or tamoxifen.

5 The invention can also be practiced by including with a specific RNAi agent another anti-cancer chemotherapeutic agent, such as any conventional chemotherapeutic agent. The combination of a specific binding agent with such other agents can potentiate the chemotherapeutic protocol. Numerous chemotherapeutic protocols will present themselves in the mind of the skilled practitioner as being capable of incorporation into the method of the

10 invention. Any chemotherapeutic agent can be used, including alkylating agents, antimetabolites, hormones and antagonists, radioisotopes, as well as natural products. For example, the compound of the invention can be administered with antibiotics such as doxorubicin and other anthracycline analogs, nitrogen mustards such as cyclophosphamide, pyrimidine analogs such as 5-fluorouracil, cisplatin, hydroxyurea, taxol and its natural and synthetic derivatives, and the like. As another example, in the case of mixed tumors, such as adenocarcinoma of the breast, where the tumors include gonadotropin-dependent and gonadotropin-independent cells, the compound can be administered in conjunction with leuprolide or goserelin (synthetic peptide analogs of LH-RH). Other antineoplastic protocols include the use of a tetracycline compound with another treatment modality, e.g., surgery, radiation, etc., also referred to herein as "adjunct 15 antineoplastic modalities." Thus, the method of the invention can be employed with such conventional regimens with the benefit of reducing side effects and enhancing efficacy.

20

Methods for inhibiting expression of an Aha gene

In yet another aspect, the invention provides a method for inhibiting the expression of an Aha gene in a mammal. The method comprises administering a composition of the invention to 25 the mammal such that expression of the target Aha gene, e.g. Aha1, is silenced. Because of their high specificity, the dsRNAs of the invention specifically target RNAs (primary or processed) of the target Aha gene. Compositions and methods for inhibiting the expression of these Aha genes using dsRNAs can be performed as described elsewhere herein.

In one embodiment, the method comprises administering a composition comprising a dsRNA, wherein the dsRNA comprises a nucleotide sequence which is complementary to at least a part of an RNA transcript of an Aha gene, e.g. Aha1, of the mammal to be treated. When the organism to be treated is a mammal such as a human, the composition may be administered by 5 any means known in the art including, but not limited to oral or parenteral routes, including intravenous, intramuscular, subcutaneous, transdermal, airway (aerosol), nasal, rectal, vaginal and topical (including buccal and sublingual) administration. In preferred embodiments, the compositions are administered by intravenous infusion or injection.

Unless otherwise defined, all technical and scientific terms used herein have the same 10 meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, 15 including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

EXAMPLES

Gene Walking of an Aha gene

siRNA design was carried out to identify siRNAs targeting Aha1. The mRNA sequences 20 of *Homo sapiens* (NM_012111.1), *mus musculus* (NM_146036.1) and *pan troglodytes* (XM_510094.1) Aha 1 were examined by computer analysis to identify homologous sequences of 19 or 21 nucleotides that yield RNAi agents cross-reactive between these three species. Of those identified, 48 such sequences were selected for minimal off-target interactions in rats (at 25 least 3 mismatches to any other gene in the rat genome, or at least two mismatches to any other gene in the rat genome, wherein one of said at least two mismatches is located in a position complementary to position 9 or 10 of the antisense strand of the corresponding RNAi agent, counting 5' to 3') and the corresponding dsRNAs synthesized for screening (AL-DP-7301 - AL-DP-7346, see Table 1). AL-DP-7561, AL-DP-7562, AL-DP-7563 and AL-DP-7564 which are

additionally cross-reactive to *mus musculus* (NM_172391.3) and *rattus norvegicus* (XM_223680.3) Aha 2, were also synthesized and screened. In addition, a further 40 sequences were selected for minimal predicted off-target interactions in humans (at least 3 mismatches to any other gene in the human genome, or at least two mismatches to any other gene in the human

5 genome, wherein one of said at least two mismatches is located in a position complementary to position 9 or 10 of the antisense strand of the corresponding RNAi agent, counting 5' to 3') and the corresponding dsRNAs synthesized for screening (AL-DP-9250 - AL-DP-9289, see Table 2). 17 sequences were identified as belonging to both sets (AL-DP-7301, AL-DP-7304, AL-DP-7305, AL-DP-7307, AL-DP-7310, AL-DP-7312, AL-DP-7315, AL-DP-7316, AL-DP-7317, AL-10 DP-7323, AL-DP-7324, AL-DP-7332, AL-DP-7336, AL-DP-7337, AL-DP-7338, AL-DP-7342, and AL-DP-7344.

dsRNA synthesis

Source of reagents

Where the source of a reagent is not specifically given herein, such reagent may be 15 obtained from any supplier of reagents for molecular biology at a quality/purity standard for application in molecular biology.

siRNA synthesis


Single-stranded RNAs were produced by solid phase synthesis on a scale of 1 μ mole using an Expedite 8909 synthesizer (Applied Biosystems, Applera Deutschland GmbH, 20 Darmstadt, Germany) and controlled pore glass (CPG, 500 \AA , Proligo Biochemie GmbH, Hamburg, Germany) as solid support. RNA and RNA containing 2'-*O*-methyl nucleotides were generated by solid phase synthesis employing the corresponding phosphoramidites and 2'-*O*-methyl phosphoramidites, respectively (Proligo Biochemie GmbH, Hamburg, Germany). These building blocks were incorporated at selected sites within the sequence of the 25 oligoribonucleotide chain using standard nucleoside phosphoramidite chemistry such as described in Current protocols in nucleic acid chemistry, Beaucage, S.L. et al. (Edrs.), John Wiley & Sons, Inc., New York, NY, USA. Phosphorothioate linkages were introduced by replacement of the iodine oxidizer solution with a solution of the Beaucage reagent (Chruachem

Ltd, Glasgow, UK) in acetonitrile (1%). Further ancillary reagents were obtained from Mallinckrodt Baker (Griesheim, Germany).

Deprotection and purification of the crude oligoribonucleotides by anion exchange HPLC were carried out according to established procedures. Yields and concentrations were determined
5 by UV absorption of a solution of the respective RNA at a wavelength of 260 nm using a spectral photometer (DU 640B, Beckman Coulter GmbH, Unterschleißheim, Germany). Double stranded RNA was generated by mixing an equimolar solution of complementary strands in annealing buffer (20 mM sodium phosphate, pH 6.8; 100 mM sodium chloride), heated in a water bath at 85 - 90°C for 3 minutes and cooled to room temperature over a period of 3 - 4
10 hours. The annealed RNA solution was stored at -20 °C until use.

For the synthesis of 3'-cholesterol-conjugated siRNAs (herein referred to as -Chol-3'), an appropriately modified solid support was used for RNA synthesis. The modified solid support was prepared as follows:


Diethyl-2-azabutane-1,4-dicarboxylate AA



15

A 4.7 M aqueous solution of sodium hydroxide (50 mL) was added into a stirred, ice-cooled solution of ethyl glycinate hydrochloride (32.19 g, 0.23 mole) in water (50 mL). Then, ethyl acrylate (23.1 g, 0.23 mole) was added and the mixture was stirred at room temperature until completion of the reaction was ascertained by TLC. After 19 h the solution was partitioned
20 with dichloromethane (3 x 100 mL). The organic layer was dried with anhydrous sodium sulfate, filtered and evaporated. The residue was distilled to afford AA (28.8 g, 61%).

3-{Ethoxycarbonylmethyl-[6-(9H-fluoren-9-ylmethoxycarbonyl-amino)-hexanoyl]-amino}-propionic acid ethyl ester **AB**




AB

5 Fmoc-6-amino-hexanoic acid (9.12 g, 25.83 mmol) was dissolved in dichloromethane (50 mL) and cooled with ice. Diisopropylcarbodiimide (3.25 g, 3.99 mL, 25.83 mmol) was added to the solution at 0°C. It was then followed by the addition of Diethyl-azabutane-1,4-dicarboxylate (5 g, 24.6 mmol) and dimethylamino pyridine (0.305 g, 2.5 mmol). The solution was brought to room temperature and stirred further for 6 h. Completion of the reaction was ascertained by TLC.

10 The reaction mixture was concentrated under vacuum and ethyl acetate was added to precipitate diisopropyl urea. The suspension was filtered. The filtrate was washed with 5% aqueous hydrochloric acid, 5% sodium bicarbonate and water. The combined organic layer was dried over sodium sulfate and concentrated to give the crude product which was purified by column chromatography (50 % EtOAc/Hexanes) to yield 11.87 g (88%) of AB.


15 3-[(6-Amino-hexanoyl)-ethoxycarbonylmethyl-amino]-propionic acid ethyl ester **AC**



AC

3-{Ethoxycarbonylmethyl-[6-(9H-fluoren-9-ylmethoxycarbonylamino)-hexanoyl]-amino}-propionic acid ethyl ester AB (11.5 g, 21.3 mmol) was dissolved in 20% piperidine in dimethylformamide at 0°C. The solution was continued stirring for 1 h. The reaction mixture was concentrated under vacuum, water was added to the residue, and the product was extracted with ethyl acetate. The crude product was purified by conversion into its hydrochloride salt.

3-({6-[17-(1,5-Dimethyl-hexyl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-3-yloxycarbonylamino]-hexanoyl}ethoxycarbonylmethyl-amino)-propionic acid ethyl ester **AD**




5

AD


The hydrochloride salt of 3-[(6-Amino-hexanoyl)-ethoxycarbonylmethyl-amino]-propionic acid ethyl ester AC (4.7 g, 14.8 mmol) was taken up in dichloromethane. The suspension was cooled to 0°C on ice. To the suspension diisopropylethylamine (3.87 g, 5.2 mL, 30 mmol) was added. To the resulting solution cholesteryl chloroformate (6.675 g, 14.8 mmol) was added. The reaction mixture was stirred overnight. The reaction mixture was diluted with dichloromethane and washed with 10% hydrochloric acid. The product was purified by flash chromatography (10.3 g, 92%).

1-{6-[17-(1,5-Dimethyl-hexyl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a] phenanthren-3-yloxycarbonylamino]-hexanoyl}-4-oxo-15 pyrrolidine-3-carboxylic acid ethyl ester **AE**

**AE**

Potassium t-butoxide (1.1 g, 9.8 mmol) was slurried in 30 mL of dry toluene. The mixture was cooled to 0°C on ice and 5 g (6.6 mmol) of diester AD was added slowly with stirring within 20 mins. The temperature was kept below 5°C during the addition. The stirring was continued for 30 mins at 0°C and 1 mL of glacial acetic acid was added, immediately followed by 4 g of 5 NaH₂PO₄·H₂O in 40 mL of water. The resultant mixture was extracted twice with 100 mL of dichloromethane each and the combined organic extracts were washed twice with 10 mL of phosphate buffer each, dried, and evaporated to dryness. The residue was dissolved in 60 mL of toluene, cooled to 0°C and extracted with three 50 mL portions of cold pH 9.5 carbonate buffer. The aqueous extracts were adjusted to pH 3 with phosphoric acid, and extracted with five 40 mL 10 portions of chloroform which were combined, dried and evaporated to dryness. The residue was purified by column chromatography using 25% ethylacetate/hexane to afford 1.9 g of b-ketoester (39%).

[6-(3-Hydroxy-4-hydroxymethyl-pyrrolidin-1-yl)-6-oxo-hexyl]-carbamic acid 17-(1,5-dimethyl-hexyl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-15 cyclopenta[a]phenanthren-3-yl ester **AF**




AF

Methanol (2 mL) was added dropwise over a period of 1 h to a refluxing mixture of b-ketoester AE (1.5 g, 2.2 mmol) and sodium borohydride (0.226 g, 6 mmol) in tetrahydrofuran (10 20 mL). Stirring was continued at reflux temperature for 1 h. After cooling to room temperature, 1 N HCl (12.5 mL) was added, the mixture was extracted with ethylacetate (3 x 40 mL). The combined ethylacetate layer was dried over anhydrous sodium sulfate and concentrated under

vacuum to yield the product which was purified by column chromatography (10% MeOH/CHCl₃) (89%).


5 (6-{3-[Bis-(4-methoxy-phenyl)-phenyl-methoxymethyl]-4-hydroxy-pyrrolidin-1-yl}-6-oxo-hexyl)-carbamic acid 17-(1,5-dimethyl-hexyl)-10,13-dimethyl-
2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-3-yl ester **AG**



AG

10 Diol AF (1.25 gm 1.994 mmol) was dried by evaporating with pyridine (2 x 5 mL) *in vacuo*. Anhydrous pyridine (10 mL) and 4,4'-dimethoxytritylchloride (0.724 g, 2.13 mmol) were added with stirring. The reaction was carried out at room temperature overnight. The reaction was quenched by the addition of methanol. The reaction mixture was concentrated under vacuum and to the residue dichloromethane (50 mL) was added. The organic layer was washed with 1M aqueous sodium bicarbonate. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated. The residual pyridine was removed by evaporating with toluene. The crude product was purified by column chromatography (2% MeOH/Chloroform, R_f = 0.5 in 5% MeOH/CHCl₃) (1.75 g, 95%).


Succinic acid mono-(4-[bis-(4-methoxy-phenyl)-phenyl-methoxymethyl]-1-{6-[17-(1,5-dimethyl-hexyl)-10,13-dimethyl 2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H cyclopenta[a]phenanthren-3-yloxycarbonylamino]-hexanoyl}-pyrrolidin-3-yl) ester **AH**



5

AH

Compound AG (1.0 g, 1.05 mmol) was mixed with succinic anhydride (0.150 g, 1.5 mmol) and DMAP (0.073 g, 0.6 mmol) and dried in a vacuum at 40°C overnight. The mixture was dissolved in anhydrous dichloroethane (3 mL), triethylamine (0.318 g, 0.440 mL, 3.15 mmol) was added and the solution was stirred at room temperature under argon atmosphere for 10 16 h. It was then diluted with dichloromethane (40 mL) and washed with ice cold aqueous citric acid (5 wt%, 30 mL) and water (2 X 20 mL). The organic phase was dried over anhydrous sodium sulfate and concentrated to dryness. The residue was used as such for the next step.

Cholesterol derivatised CPG **AI**

15

AI

Succinate AH (0.254 g, 0.242 mmol) was dissolved in a mixture of dichloromethane/acetonitrile (3:2, 3 mL). To that solution DMAP (0.0296 g, 0.242 mmol) in acetonitrile (1.25 mL), 2,2'-Dithio-bis(5-nitropyridine) (0.075 g, 0.242 mmol) in acetonitrile/dichloroethane (3:1, 1.25 mL) were added successively. To the resulting solution 5 triphenylphosphine (0.064 g, 0.242 mmol) in acetonitrile (0.6 ml) was added. The reaction mixture turned bright orange in color. The solution was agitated briefly using a wrist-action shaker (5 mins). Long chain alkyl amine-CPG (LCAA-CPG) (1.5 g, 61 mM) was added. The suspension was agitated for 2 h. The CPG was filtered through a sintered funnel and washed with acetonitrile, dichloromethane and ether successively. Unreacted amino groups were masked 10 using acetic anhydride/pyridine. The achieved loading of the CPG was measured by taking UV measurement (37 mM/g).

The synthesis of siRNAs bearing a 5'-12-dodecanoic acid bisdecylamide group (herein referred to as "5'-C32-") or a 5'-cholesteryl derivative group (herein referred to as "5'-Chol-") was performed as described in WO 2004/065601, except that, for the cholesteryl derivative, the 15 oxidation step was performed using the Beaucage reagent in order to introduce a phosphorothioate linkage at the 5'-end of the nucleic acid oligomer.

Nucleic acid sequences are represented below using standard nomenclature, and specifically the abbreviations of Table 2.

20 **Table 3: Abbreviations of nucleotide monomers used in nucleic acid sequence representation. It will be understood that these monomers, when present in an oligonucleotide, are mutually linked by 5'-3'-phosphodiester bonds.**

Abbreviation ^a	Nucleotide(s)
A, a	2'-deoxy-adenosine-5'-phosphate, adenosine-5'-phosphate
C, c	2'-deoxy-cytidine-5'-phosphate, cytidine-5'-phosphate
G, g	2'-deoxy-guanosine-5'-phosphate, guanosine-5'-phosphate
T, t	2'-deoxy-thymidine-5'-phosphate, thymidine-5'-phosphate
U, u	2'-deoxy-uridine-5'-phosphate, uridine-5'-phosphate
N, n	any 2'-deoxy-nucleotide/nucleotide (G, A, C, or T, g, a, c or u)
Am	2'-O-methyladenosine-5'-phosphate
Cm	2'-O-methylcytidine-5'-phosphate
Gm	2'-O-methylguanosine-5'-phosphate

Abbreviation ^a	Nucleotide(s)
Tm	2'- <i>O</i> -methyl-thymidine-5'-phosphate
Um	2'- <i>O</i> -methyluridine-5'-phosphate
Af	2'-fluoro-2'-deoxy-adenosine-5'-phosphate
Cf	2'-fluoro-2'-deoxy-cytidine-5'-phosphate
Gf	2'-fluoro-2'-deoxy-guanosine-5'-phosphate
Tf	2'-fluoro-2'-deoxy-thymidine-5'-phosphate
Uf	2'-fluoro-2'-deoxy-uridine-5'-phosphate
<u>A</u> , <u>C</u> , <u>G</u> , <u>T</u> , <u>U</u> , <u>a</u> , <u>c</u> , <u>g</u> , <u>t</u> , <u>u</u>	underlined: nucleoside-5'-phosphorothioate
<u>am</u> , <u>cm</u> , <u>gm</u> , <u>tm</u> , <u>um</u>	underlined: 2- <i>O</i> -methyl-nucleoside-5'-phosphorothioate

^acapital letters represent 2'-deoxyribonucleotides (DNA), lower case letters represent ribonucleotides (RNA)

dsRNA expression vectors

In another aspect of the invention, Aha1 specific dsRNA molecules that modulate Aha1 gene expression activity are expressed from transcription units inserted into DNA or RNA vectors (see, e.g., Couture, A, et al., *TIG*. (1996), **12**:5-10; Skillem, A., et al., International PCT Publication No. WO 00/22113, Conrad, International PCT Publication No. WO 00/22114, and Conrad, US Pat. No. 6,054,299). These transgenes can be introduced as a linear construct, a circular plasmid, or a viral vector, which can be incorporated and inherited as a transgene integrated into the host genome. The transgene can also be constructed to permit it to be inherited as an extrachromosomal plasmid (Gassmann, et al., *Proc. Natl. Acad. Sci. USA* (1995) **92**:1292).

The individual strands of a dsRNA can be transcribed by promoters on two separate expression vectors and co-transfected into a target cell. Alternatively each individual strand of the dsRNA can be transcribed by promoters both of which are located on the same expression plasmid. In a preferred embodiment, a dsRNA is expressed as an inverted repeat joined by a linker polynucleotide sequence such that the dsRNA has a stem and loop structure.

The recombinant dsRNA expression vectors are generally DNA plasmids or viral vectors. dsRNA expressing viral vectors can be constructed based on, but not limited to, adeno-

associated virus (for a review, see Muzyczka, et al., *Curr. Topics Micro. Immunol.* (1992) 158:97-129)); adenovirus (see, for example, Berkner, et al., *BioTechniques* (1998) 6:616), Rosenfeld et al. (1991, *Science* 252:431-434), and Rosenfeld et al. (1992), *Cell* 68:143-155)); or alphavirus as well as others known in the art. Retroviruses have been used to introduce a variety 5 of genes into many different cell types, including epithelial cells, in vitro and/or in vivo (see, e.g., Eglitis, et al., *Science* (1985) 230:1395-1398; Danos and Mulligan, *Proc. Natl. Acad. Sci. USA* (1998) 85:6460-6464; Wilson et al., 1988, *Proc. Natl. Acad. Sci. USA* 85:3014-3018; Armentano et al., 1990, *Proc. Natl. Acad. Sci. USA* 87:61416145; Huber et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:8039-8043; Ferry et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:8377-8381; 10 Chowdhury et al., 1991, *Science* 254:1802-1805; van Beusechem. et al., 1992, *Proc. Nad. Acad. Sci. USA* 89:7640-19 ; Kay et al., 1992, *Human Gene Therapy* 3:641-647; Dai et al., 1992, *Proc. Natl. Acad. Sci. USA* 89:10892-10895; Hwu et al., 1993, *J. Immunol.* 150:4104-4115; U.S. Patent No. 4,868,116; U.S. Patent No. 4,980,286; PCT Application WO 89/07136; PCT Application WO 89/02468; PCT Application WO 89/05345; and PCT Application WO 15 92/07573). Recombinant retroviral vectors capable of transducing and expressing genes inserted into the genome of a cell can be produced by transfecting the recombinant retroviral genome into suitable packaging cell lines such as PA317 and Psi-CRIP (Comette et al., 1991, *Human Gene Therapy* 2:5-10; Cone et al., 1984, *Proc. Natl. Acad. Sci. USA* 81:6349). Recombinant adenoviral vectors can be used to infect a wide variety of cells and tissues in susceptible hosts 20 (e.g., rat, hamster, dog, and chimpanzee) (Hsu et al., 1992, *J. Infectious Disease*, 166:769), and also have the advantage of not requiring mitotically active cells for infection.

The promoter driving dsRNA expression in either a DNA plasmid or viral vector of the invention may be a eukaryotic RNA polymerase I (e.g. ribosomal RNA promoter), RNA polymerase II (e.g. CMV early promoter or actin promoter or U1 snRNA promoter) or generally 25 RNA polymerase III promoter (e.g. U6 snRNA or 7SK RNA promoter) or a prokaryotic promoter, for example the T7 promoter, provided the expression plasmid also encodes T7 RNA polymerase required for transcription from a T7 promoter. The promoter can also direct transgene expression to the pancreas (see, e.g. the insulin regulatory sequence for pancreas (Bucchini et al., 1986, *Proc. Natl. Acad. Sci. USA* 83:2511-2515)).

In addition, expression of the transgene can be precisely regulated, for example, by using an inducible regulatory sequence and expression systems such as a regulatory sequence that is sensitive to certain physiological regulators, e.g., circulating glucose levels, or hormones (Docherty et al., 1994, FASEB J. 8:20-24). Such inducible expression systems, suitable for the 5 control of transgene expression in cells or in mammals include regulation by ecdysone, by estrogen, progesterone, tetracycline, chemical inducers of dimerization, and isopropyl-beta-D 1 - thiogalactopyranoside (EPTG). A person skilled in the art would be able to choose the appropriate regulatory/promoter sequence based on the intended use of the dsRNA transgene.

Generally, recombinant vectors capable of expressing dsRNA molecules are delivered as 10 described below, and persist in target cells. Alternatively, viral vectors can be used that provide for transient expression of dsRNA molecules. Such vectors can be repeatedly administered as necessary. Once expressed, the dsRNAs bind to target RNA and modulate its function or expression. Delivery of dsRNA expressing vectors can be systemic, such as by intravenous or intramuscular administration, by administration to target cells ex-planted from the patient 15 followed by reintroduction into the patient, or by any other means that allows for introduction into a desired target cell.

dsRNA expression DNA plasmids are typically transfected into target cells as a complex with cationic lipid carriers (e.g. Oligofectamine) or non-cationic lipid-based carriers (e.g. Transit-TKOTM). Multiple lipid transfections for dsRNA-mediated knockdowns targeting 20 different regions of a single Aha1 gene or multiple Aha1 genes over a period of a week or more are also contemplated by the invention. Successful introduction of the vectors of the invention into host cells can be monitored using various known methods. For example, transient transfection can be signaled with a reporter, such as a fluorescent marker, such as Green Fluorescent Protein (GFP). Stable transfection of ex vivo cells can be ensured using markers 25 that provide the transfected cell with resistance to specific environmental factors (e.g., antibiotics and drugs), such as hygromycin B resistance.

The Aha1 specific dsRNA molecules can also be inserted into vectors and used as gene therapy vectors for human patients. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (see U.S. Patent 5,328,470) or by

stereotactic injection (see e.g., Chen et al. (1994) Proc. Natl. Acad. Sci. USA 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from 5 recombinant cells, e.g., retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

Aha1 siRNA in vitro screening

Single Dose Screen in HeLa and MLE 12 cells

HeLa cells were obtained from American Type Culture Collection (Rockville, MD, cat.

10 No. HB-8065) and cultured in Ham's F12 (Biochrom AG, Berlin, Germany, cat. No. FG0815) supplemented to contain 10% fetal calf serum (FCS) (Biochrom AG, Berlin, Germany, cat. No. S0115), Penicillin 100 U/ml, Streptomycin 100 µg/ml (Biochrom AG, Berlin, Germany, cat. No. A2213) at 37°C in an atmosphere with 5% CO₂ in a humidified incubator (Heraeus HERAcell, Kendro Laboratory Products, Langenselbold, Germany).

15 MLE 12 cells were obtained from American Type Culture Collection (Rockville, MD, cat. No. CRL-2110) and cultured in HITES Medium (1:1 mix Dulbecco's MEM (Biochrom AG, Berlin, Germany, cat. No: F0435) + Ham's F12 (Biochrom AG, Berlin, Germany, cat. No: FG0815)) supplemented to contain 2% fetal calf serum (FCS) (Biochrom AG, Berlin, Germany, cat. No. S0115), Penicillin 100 U/ml, Streptomycin 100 µg/ml (Biochrom AG, Berlin, Germany, 20 cat. No. A2213), 4 mM L-Glutamin (Biochrom AG, Berlin, Germany, cat. No: K0282), 1 x Insulin/Transferrin/Na-Selenit (Gibco: 51500-056), 10 nM Hydrocortisone (Sigma Munich, Germany, cat.No: H6909), 10 nM β-Estradiol (Sigma Munich, Germany, cat.No: E2257), and 10 mM HEPES (USB Europe GmbH, Staufen, Germany cat. No.: 16926) at 37°C in an atmosphere with 5% CO₂ in a humidified incubator (Heraeus HERAcell, Kendro Laboratory 25 Products, Langenselbold, Germany).

Transfection and mRNA quantification

For transfection with siRNA, HeLa and MLE12 cells were seeded at a density of 2.0×10^4 cells/well in 96-well plates and transfected directly. Transfection of siRNA (30nM) was carried out with lipofectamine 2000 (Invitrogen GmbH, Karlsruhe, Germany, cat. No. 11668-019) as described by the manufacturer. 24 hours after transfection cells were lysed and Aha1 mRNA levels were quantified with the Quantigene Explore Kit (Genospectra, Dumbarton Circle Fremont, USA, cat. No. QG-000-02) according to the manufacturer's protocol. Aha1 mRNA levels were normalized to GAPDH mRNA. Readings were obtained in quadruplicates for each siRNA. siRNA duplexes unrelated to Aha1 gene were used as control. The activity of a given Aha1 specific siRNA duplex was expressed as percent Aha1 mRNA concentration in treated cells relative to Aha1 mRNA concentration in cells treated with the control siRNA duplex.

Table 4: Probe sequences used with Quantigene Explore Kit (Genospectra) in quantification of Homo sapiens (hs) Aha1

FPL Name	Function	Sequence
hsAha1 001	CE	GATGTAAATTCCCATTGCTTCTCTTTCTCTTGAAAGAAAGT
hsAha1 002	CE	TGAACTCTGTTGAGGGTGCCTTCTCTTGAAAGAAAGT
hsAha1 003	CE	GGGTCTACTGACTCTCATTGTTCTCTTGAAAGAAAGT
hsAha1 004	CE	CCTTGCCTCAGTTCTCTTGAAAGAAAGT
hsAha1 005	CE	GGTTTTGAAGGAGCAGGCTTAGTTCTCTTGAAAGAAAGT
hsAha1 006	LE	ACGCTTTTCATCAGACAAATTTTTAGGCATAGGACCGTGTCT
hsAha1 007	LE	GCTCACACTAATCTCCACTTCATCCTTTAGGCATAGGACCGTGTCT
hsAha1 008	LE	TCATTAAGGCCACGAGATTGTTTTAGGCATAGGACCGTGTCT
hsAha1 009	LE	TAGGTAAGATCATGCCCTGGGTTTTAGGCATAGGACCGTGTCT
hsAha1 010	LE	ACTCCAACAGGTCTGCCCTGTTTAGGCATAGGACCGTGTCT
hsAha1 011	BL	GTCAGGCTCATTTGGCAAG
hsAha1 012	BL	TAGAAGTTCACCCCTCTCCT
hsAha1 013	BL	AGTGCTGGCTGCCCACT

Table 5: Probe sequences used with Quantigene Explore Kit (Genospectra) in quantification of Mus musculus (mm) Aha1

FPL Name	Function	Sequence
mmAhsa 1001	CE	CTCGAACGGCCAGGAACATTTCTCTTGAAAGAAAGT
mmAhsa 1002	CE	GCACTTGCCCTCTCATTCTATTTCTCTTGAAAGAAAGT
mmAhsa 1003	CE	TTGATGGATGCCTCCCATTTCTCTTGAAAGAAAGT
mmAhsa 1004	CE	AACTCTGTCTTGAGGGTGCCTGATTTCTCTTGAAAGAAAGT
	CE	TTTGGCCTGGCTTTGAATTCTCTTGAAAGAAAGT

FPL Name	Function	Sequence
	LE	CAAGCTTGTTCACCTCGGTACACCTCTTTAGGCATAGGACCCGTGTCT
	LE	CCTGACTTAGAGGTACCTGTCCAGTTTTAGGCATAGGACCCGTGTCT
	LE	GATTCCACATGTCCTTGTACTGCACTTTTTAGGCATAGGACCCGTGTCT
	LE	ATTTCATCAGACAAATTGGGTTTTAGGCATAGGACCCGTGTCT
	LE	TAATCTCCACTTCATCCACGCTTTTTAGGCATAGGACCCGTGTCT
	LE	TTTCACCCCGTCTCCTCATTTTAGGCATAGGACCCGTGTCT
	LE	ACTGTGGGCAAGATCATGCCCTGAGTATTTAGGCATAGGACCCGTGTCT
	BL	AAGATAAGTTGCCTTCCTGTTG
	BL	CAGTTGATGGTCCACTCATAGAAG
	BL	CATCTTGGCAAGGCTCACAC
	BL	TTAAGGCCACGAGATTGTGTCAGGCT
	BL	GTAAATTCCCCTGCTTCTCTCAGAAG
	BL	CACTGGATCTACTGACTCTCCATTG
	BL	CTCAGTCTTAGTGCTGGCTGGCC
	BL	GGAGCAGACTTAGCCTTGCAAGT

Dose-Response Curves in HeLa cells

Transfection and mRNA quantification: For transfection with siRNA, HeLa cells were seeded at a density of 2.0×10^4 cells/well in 96-well plates and transfected directly. Transfection 5 of siRNA was carried out with lipofectamine 2000 (Invitrogen GmbH, Karlsruhe, Germany, cat. No. 11668-019) as described by the manufacturer. siRNAs were concentrated from 30 nM in 3 fold dilutions to 14 pM. 24 hours after transfection Hela cells were lysed and Aha1 mRNA levels were quantified with the Quantigene Explore Kit (Genospectra, Dumbarton Circle Fremont, USA, cat. No. QG-000-02) according to the protocol. Aha1 mRNA levels were normalized to 10 GAP-DH mRNA. For each siRNA four individual datapoints were collected. siRNA duplexes unrelated to Aha1 gene were used as control. The activity of a given Aha1 specific siRNA duplex was expressed as percent Aha1 mRNA concentration in treated cells relative to Aha1 mRNA concentration in cells treated with the control siRNA duplex. XL-fit was used to calculate IC₅₀ values.

15 Table 6 provides values for inhibition of Aha1 expression using various dsRNA molecules of the invention.

Table 6: Residual Aha1 mRNA in % of control in HeLa and MLE12 cells treated with 30 nM solutions of various RNAi agents specific for Aha1, and IC₅₀ for selected RNAi agents determined in HeLa cells

Duplex identifier	HeLa cells, residual mRNA [%]	IC ₅₀ in HeLa cells [nM]	MLE12 cells, residual mRNA [%]
AL-DP-7299	19 ± 3		105 ± 17
AL-DP-7300	7 ± 2		94 ± 13
AL-DP-7301	3 ± 1	0,035	14 ± 3
AL-DP-7302	17 ± 5		61 ± 16
AL-DP-7303	5 ± 2		23 ± 5
AL-DP-7304	7 ± 3		30 ± 7
AL-DP-7305	6 ± 2		26 ± 6
AL-DP-7306	27 ± 8		45 ± 11
AL-DP-7307	16 ± 6	1,1	27 ± 8
AL-DP-7308	13 ± 5	0,21	20 ± 7
AL-DP-7309	10 ± 3	0,36	22 ± 8
AL-DP-7310	51 ± 13		59 ± 13
AL-DP-7311	4 ± 3	0,07	16 ± 4
AL-DP-7312	14 ± 3		40 ± 8
AL-DP-7313	63 ± 14		80 ± 10
AL-DP-7314	97 ± 21		88 ± 10
AL-DP-7315	76 ± 24		77 ± 10
AL-DP-7316	7 ± 2	0,34	23 ± 6
AL-DP-7317	11 ± 3		44 ± 12
AL-DP-7318	4 ± 2	0,29	16 ± 3
AL-DP-7319	38 ± 7		73 ± 21
AL-DP-7320	16 ± 4	0,07	15 ± 5
AL-DP-7321	130 ± 36		81 ± 16
AL-DP-7322	4 ± 2	0,045	10 ± 3
AL-DP-7323	24 ± 6		55 ± 11
AL-DP-7324	3 ± 2	0,089	12 ± 4
AL-DP-7325	5 ± 2	0,3	12 ± 4
AL-DP-7326	3 ± 1	0,27	19 ± 7
AL-DP-7327	3 ± 1	0,08	13 ± 7
AL-DP-7328	49 ± 14		67 ± 10
AL-DP-7329	6 ± 2	0,2	18 ± 5
AL-DP-7330	64 ± 19		78 ± 10

Duplex identifier	HeLa cells, residual mRNA [%]	IC ₅₀ in HeLa cells [nM]	MLE12 cells, residual mRNA [%]
AL-DP-7331	5 ± 2	0,55	13 ± 5
AL-DP-7332	95 ± 20		82 ± 15
AL-DP-7333	2 ± 1	0,27	9 ± 3
AL-DP-7334	94 ± 17		83 ± 19
AL-DP-7335	11 ± 5		57 ± 11
AL-DP-7336	22 ± 4		63 ± 12
AL-DP-7337	6 ± 2	0,29	20 ± 5
AL-DP-7338	39 ± 6		56 ± 10
AL-DP-7339	10 ± 1		35 ± 6
AL-DP-7340	8 ± 2	0,61	19 ± 6
AL-DP-7341	17 ± 4		55 ± 16
AL-DP-7342	6 ± 4	0,5	15 ± 3
AL-DP-7343	26 ± 4		103 ± 19
AL-DP-7344	5 ± 2		38 ± 11
AL-DP-7345	53 ± 22		63 ± 15
AL-DP-7346	22 ± 4		44 ± 11
AL-DP-9250	4 ± 1		
AL-DP-9251	51 ± 9		
AL-DP-9252	19 ± 2		
AL-DP-9253	11 ± 1		
AL-DP-9254	7 ± 1		
AL-DP-9255	5 ± 0		
AL-DP-9256	5 ± 0		
AL-DP-9257	7 ± 0		
AL-DP-9258	9 ± 1		
AL-DP-9259	7 ± 1		
AL-DP-9260	15 ± 3		
AL-DP-9261	21 ± 2		
AL-DP-9262	24 ± 4		
AL-DP-9263	25 ± 6		
AL-DP-9264	7 ± 2		
AL-DP-9265	8 ± 1		
AL-DP-9266	11 ± 2		
AL-DP-9267	45 ± 4		
AL-DP-9268	9 ± 1		

Duplex identifier	HeLa cells, residual mRNA [%]	IC ₅₀ in HeLa cells [nM]	MLE12 cells, residual mRNA [%]
AL-DP-9269	5 ± 1		
AL-DP-9270	6 ± 1		
AL-DP-9271	6 ± 2		
AL-DP-9272	26 ± 8		
AL-DP-9273	11 ± 1		
AL-DP-9274	7 ± 1		
AL-DP-9275	8 ± 1		
AL-DP-9276	4 ± 1		
AL-DP-9277	10 ± 1		
AL-DP-9278	2 ± 0		
AL-DP-9279	3 ± 0		
AL-DP-9280	12 ± 1		
AL-DP-9281	8 ± 2		
AL-DP-9282	3 ± 0		
AL-DP-9283	6 ± 1		
AL-DP-9284	39 ± 2		
AL-DP-9285	4 ± 1		
AL-DP-9286	61 ± 11		
AL-DP-9287	3 ± 1		
AL-DP-9288	27 ± 5		
AL-DP-9289	6 ± 1		

In summary, AL-DP-7301, AL-DP-7308, AL-DP-7318, AL-DP-7320, AL-DP-7322, AL-DP-7324, AL-DP-7325, AL-DP-7326, AL-DP-7327, AL-DP-7329, AL-DP-7331, AL-DP-7333, AL-DP-7340, and AL-DP-7342 inhibited Aha1 expression by at least 80% in both HeLa and

5 MLE12 cells, AL-DP-7303, AL-DP-7305, AL-DP-7307, AL-DP-7309, AL-DP-7316, and AL-DP-7337 inhibited Aha1 expression by at least 80% in HeLa cells and by at least 70% in MLE12 cells, AL-DP-7304, AL-DP-7312, AL-DP-7339, and AL-DP-7344 inhibited Aha1 expression by at least 80% in HeLa cells and by at least 60% in MLE12 cells, AL-DP-7306, AL-DP-7317, and AL-DP-7346 inhibited Aha1 expression by at least 70% in HeLa cells and by at least 50% in

10 MLE12 cells, AL-DP-7310, AL-DP-7323, AL-DP-7335, AL-DP-7338, and AL-DP-7341 inhibited Aha1 expression by at least 40% in both HeLa and MLE12 cells, and AL-DP-7302,

AL-DP-7315, AL-DP-7328, AL-DP-7330, AL-DP-7336, and AL-DP-7345, inhibited Aha1 expression by at least 20% in both HeLa and MLE12 cells.

In addition, AL-DP-9250, AL-DP-9252, AL-DP-9253, AL-DP-9254, AL-DP-9255, AL-DP-9256, AL-DP-9257, AL-DP-9258, AL-DP-9259, AL-DP-9260, AL-DP-9264, AL-DP-9265, 5 AL-DP-9266, AL-DP-9268, AL-DP-9269, AL-DP-9270, AL-DP-9271, AL-DP-9273, AL-DP-9274, AL-DP-9275, AL-DP-9276, AL-DP-9277, AL-DP-9279, AL-DP-9280, AL-DP-9281, AL-DP-9282, AL-DP-9283, AL-DP-9285, AL-DP-9287, and AL-DP-9289 inhibited Aha1 expression by at least 80% in HeLa cells, AL-DP-9261, AL-DP-9262, AL-DP-9263, AL-DP-9272, and AL-DP-9288 inhibited Aha1 expression by at least 70% in HeLa cells, AL-DP-9263 10 inhibited Aha1 expression by at least 60% in HeLa cells, AL-DP-9267 inhibited Aha1 expression by at least 50% in HeLa cells, AL-DP-9251 inhibited Aha1 expression by at least 40% in HeLa cells, and AL-DP-9286 inhibited Aha1 expression by at least 30% in HeLa cells.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A double-stranded ribonucleic acid (dsRNA) for inhibiting the expression of a human Aha gene in a cell, wherein said dsRNA comprises at least two sequences that are complementary to each other and wherein a sense strand comprises a first sequence and an antisense strand comprises a second sequence comprising a region of complementarity which is substantially complementary to at least a part of a mRNA encoding an Aha gene, and wherein said region of complementarity is less than 30 nucleotides in length and wherein the dsRNA effects cleavage of an mRNA encoding an Aha gene within a target sequence of a second dsRNA having a sense strand set forth in SEQ ID NO: 51 and an antisense strand complementary to the latter sense strand and having a sequence set forth in SEQ ID NO: 52.
2. The dsRNA of claim 1, wherein said dsRNA is AL-DP-7324 consisting of a sense strand having the sequence set forth in SEQ ID NO: 51 and an antisense strand having the sequence set forth in SEQ ID NO: 52.
3. The dsRNA of claim 1 or 2, wherein said Aha gene is an Ahal gene.
4. The dsRNA according to any one of claims 1 to 3, wherein the Aha gene is an Ahal gene of humans.
5. The dsRNA according to any one of claims 1 to 4, wherein, upon contact with a cell expressing the Aha gene, said dsRNA inhibits expression of said Aha gene in said cell by at least 20% compared to the inhibition of said Aha gene produced by a control siRNA duplex having a sequence that is unrelated to said Aha gene.
6. The dsRNA of claim 5, wherein the cells expressing the Aha gene are HeLa cells or MLE12 cells.
7. The dsRNA according to any one of claims 1 to 6, wherein said dsRNA comprises at least one modified nucleotide.
8. The dsRNA of claim 7, wherein the modified nucleotide is a 2'-O-methyl modified nucleotide, a nucleotide comprising a 5'-phosphorothioate group, a terminal nucleotide linked to a cholestryl derivative or dodecanoic acid bisdecylamide group, a 2'-deoxy-2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a locked nucleotide, an abasic

nucleotide, a 2'-amino-modified nucleotide, a 2'-alkyl-modified nucleotide, a morpholino nucleotide, a phosphoramidate, or a non-natural base comprising nucleotide.

9. A vector for inhibiting the expression of an Aha gene in a cell, said vector comprising a regulatory sequence operably linked to a nucleotide sequence that encodes at least one strand of the dsRNA according to any one of claims 1 to 8.
10. The vector according to claim 9, wherein the nucleotide sequence encodes both strands of the dsRNA.
11. A cell comprising the dsRNA according to any one of claims 1 to 8 or the vector according to claim 9 or 10.
12. A pharmaceutical composition for inhibiting the expression of an Aha gene in an organism, wherein said pharmaceutical composition comprises the dsRNA according to any one of claims 1 to 8 or the vector according to claim 9 or 10 and a pharmaceutically acceptable carrier or excipient.
13. The pharmaceutical composition according to claim 12, wherein said composition further comprises a dsRNA comprising at least two sequences that are complementary to each other and wherein a sense strand comprises a first sequence and an antisense strand comprises a second sequence comprising a region of complementarity which is substantially complementary to at least a part of a mRNA encoding an Aha gene, and wherein said region of complementarity is less than 30 nucleotides in length, and wherein the dsRNA effects cleavage of an mRNA encoding an Aha gene within the target sequence of a second dsRNA having a sense strand sequence selected from the group of SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 27, SEQ ID NO: 29, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 35, SEQ ID NO: 37, SEQ ID NO: 39, SEQ ID NO: 43, SEQ ID NO: 45, SEQ ID NO: 47, SEQ ID NO: 49, SEQ ID NO: 53, SEQ ID NO: 55, SEQ ID NO: 57, SEQ ID NO: 59, SEQ ID NO: 61, SEQ ID NO: 63, SEQ ID NO: 65, SEQ ID NO: 67, SEQ ID NO: 69, SEQ ID NO: 71, SEQ ID NO: 73, SEQ ID NO: 75, SEQ ID NO: 77, SEQ ID NO: 79, SEQ ID NO: 81, SEQ ID NO: 83, SEQ ID NO: 85, SEQ ID NO: 87, SEQ ID NO: 89, SEQ ID NO: 91, SEQ ID NO: 93, SEQ ID NO: 95, SEQ ID NO: 97, SEQ ID NO: 99, SEQ ID NO: 101, SEQ ID NO: 103, SEQ ID NO:

105, SEQ ID NO: 107, SEQ ID NO: 109, SEQ ID NO: 111, SEQ ID NO: 113, SEQ ID NO: 115, SEQ ID NO: 117, SEQ ID NO: 119, SEQ ID NO: 121, SEQ ID NO: 123, SEQ ID NO: 125, SEQ ID NO: 127, SEQ ID NO: 129, SEQ ID NO: 131, SEQ ID NO: 133, SEQ ID NO: 135, SEQ ID NO: 137, SEQ ID NO: 139, SEQ ID NO: 141, SEQ ID NO: 143, SEQ ID NO: 145, SEQ ID NO: 147, SEQ ID NO: 149, SEQ ID NO: 151, SEQ ID NO: 153, SEQ ID NO: 155, SEQ ID NO: 157, SEQ ID NO: 159, SEQ ID NO: 163, SEQ ID NO: 165, SEQ ID NO: 167, SEQ ID NO: 169, SEQ ID NO: 171, SEQ ID NO: 173, SEQ ID NO: 175, SEQ ID NO: 177, SEQ ID NO: 179, SEQ ID NO: 181, and SEQ ID NO: 183, and an antisense strand sequence complementary to the latter sense strand and selected from the group of SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34, SEQ ID NO: 36, SEQ ID NO: 38, SEQ ID NO: 40, SEQ ID NO: 44, SEQ ID NO: 46, SEQ ID NO: 48, SEQ ID NO: 50, SEQ ID NO: 54, SEQ ID NO: 56, SEQ ID NO: 58, SEQ ID NO: 60, SEQ ID NO: 62, SEQ ID NO: 64, SEQ ID NO: 66, SEQ ID NO: 68, SEQ ID NO: 70, SEQ ID NO: 72, SEQ ID NO: 74, SEQ ID NO: 76, SEQ ID NO: 78, SEQ ID NO: 80, SEQ ID NO: 82, SEQ ID NO: 84, SEQ ID NO: 86, SEQ ID NO: 88, SEQ ID NO: 90, SEQ ID NO: 92, SEQ ID NO: 94, SEQ ID NO: 96, SEQ ID NO: 98, SEQ ID NO: 100, SEQ ID NO: 102, SEQ ID NO: 104, SEQ ID NO: 106, SEQ ID NO: 108, SEQ ID NO: 110, SEQ ID NO: 112, SEQ ID NO: 114, SEQ ID NO: 116, SEQ ID NO: 118, SEQ ID NO: 120, SEQ ID NO: 122, SEQ ID NO: 124, SEQ ID NO: 126, SEQ ID NO: 128, SEQ ID NO: 130, SEQ ID NO: 132, SEQ ID NO: 134, SEQ ID NO: 136, SEQ ID NO: 138, SEQ ID NO: 140, SEQ ID NO: 142, SEQ ID NO: 144, SEQ ID NO: 146, SEQ ID NO: 148, SEQ ID NO: 150, SEQ ID NO: 152, SEQ ID NO: 154, SEQ ID NO: 156, SEQ ID NO: 158, SEQ ID NO: 160, SEQ ID NO: 164, SEQ ID NO: 166, SEQ ID NO: 168, SEQ ID NO: 170, SEQ ID NO: 172, SEQ ID NO: 174, SEQ ID NO: 176, SEQ ID NO: 178, SEQ ID NO: 180, SEQ ID NO: 182, and SEQ ID NO: 184.

14. A method for inhibiting the expression of an Aha gene in a cell, the method comprising:
 - (a) introducing into the cell the double-stranded ribonucleic acid (dsRNA) according to any one of claims 1 to 8 or the vector according to claim 9 or 10 or the pharmaceutical composition according to claim 12 or 13; and
 - (b) maintaining the cell produced in step (a) for a time sufficient to obtain degradation of

the mRNA transcript of an Aha gene, thereby inhibiting expression of an Aha gene in the cell.

15. The method according to claim 14 when performed *in vitro*.
16. The method according to claim 14 when performed *in vivo*.
17. A method of treating, preventing or managing a pathological process mediated by Aha expression comprising administering to a patient in need of such treatment, prevention or management a therapeutically or prophylactically effective amount of the dsRNA according to any one of claims 1 to 8 or the vector according to claim 9 or 10 or the pharmaceutical composition according to claim 12 or 13.
18. The method according to claim 17 wherein the pathological process mediated by Aha expression is Cystic Fibrosis or cancer or a tumor metastasis or a complication of Cystic Fibrosis mediated by expression of an Ahal gene of humans or a complication of a cancer mediated by expression of an Ahal gene of humans.
19. Use of the dsRNA according to any one of claims 1 to 8 or the vector according to claim 9 or 10 or the pharmaceutical composition according to claim 12 or 13 in medicine.
20. Use of the dsRNA according to any one of claims 1 to 8 or the vector according to claim 9 or 10 in the preparation of a medicament for treatment of Cystic Fibrosis or cancer or a tumor metastasis or a complication of Cystic Fibrosis mediated by expression of an Ahal gene of humans or a complication of a cancer mediated by expression of an Ahal gene of humans.

21. The dsRNA according to any one of claims 1 to 8 or the vector according to claim 9 or 10 or the cell according to claim 11 or the pharmaceutical composition according to claim 12 or 13 or the method according to any one of claims 14 to 16 or the method according to claim 17 or 18 or the use according to claim 19 or the use according to claim 20 substantially as hereinbefore described with reference to the accompanying Examples.

DATED this TWENTIETH day of APRIL, 2012

Alnylam Pharmaceuticals, Inc

-and-

The Scripps Research Institute

By the Patent Attorneys for the Applicants:

FB Rice