




(86) Date de dépôt PCT/PCT Filing Date: 2014/03/12
(87) Date publication PCT/PCT Publication Date: 2014/09/25
(85) Entrée phase nationale/National Entry: 2015/08/13
(86) N° demande PCT/PCT Application No.: US 2014/024524
(87) N° publication PCT/PCT Publication No.: 2014/150914
(30) Priorités/Priorities: 2013/03/15 (US61/793,439);
2013/07/30 (US61/859,883)

(51) Cl.Int./Int.Cl. *A01H 5/00* (2006.01),
C12N 15/82 (2006.01)
(71) Demandeur/Applicant:
PIONEER HI-BRED INTERNATIONAL, INC., US
(72) Inventeurs/Inventors:
CONG, RUTH, US;
HOU, JINGTONG, US;
HOU, ZHENGLIN, US;
PATTEN, PHILLIP A., US;
YAMAMOTO, TAKASHI, US
(74) Agent: TORYS LLP

(54) Titre : POLYPEPTIDES PHI-4 ET LEURS PROCEDES D'UTILISATION
(54) Title: PHI-4 POLYPEPTIDES AND METHODS FOR THEIR USE

Fig. 4



(57) **Abrégé/Abstract:**

Compositions and methods for controlling pests are provided. The methods involve transforming organisms with a nucleic acid sequence encoding an insecticidal protein. In particular, the nucleic acid sequences are useful for preparing plants and microorganisms that possess insecticidal activity. Thus, transformed bacteria, plants, plant cells, plant tissues and seeds are provided. Compositions are insecticidal nucleic acids and proteins of bacterial species. The sequences find use in the construction of expression vectors for subsequent transformation into organisms of interest, as probes for the isolation of other homologous (or partially homologous) genes. The insecticidal proteins find use in controlling, inhibiting growth or killing lepidopteran, coleopteran, dipteran, fungal, hemipteran, and nematode pest populations and for producing compositions with insecticidal activity.

(51) International Patent Classification:
C12N 15/82 (2006.01)

(21) International Application Number:
PCT/US2014/024524

(22) International Filing Date:
12 March 2014 (12.03.2014)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/793,439 15 March 2013 (15.03.2013) US
61/859,883 30 July 2013 (30.07.2013) US

(71) Applicant: **PIONEER HI-BRED INTERNATIONAL, INC.** [US/US]; 7100 N.W. 62nd Avenue, Johnston, Iowa 50131-1014 (US).

(72) Inventors; and

(71) Applicants : **CONG, Ruth** [US/US]; Pioneer Hi-Bred International, Inc., 7250 N.W. 62nd Avenue, Johnston, Iowa 50131-0552 (US). **HOU, Jingtong** [CN/US]; Pioneer Hi-Bred International, Inc., 7250 N.W. 62nd Avenue, Johnston, Iowa 50131-0552 (US). **HOU, Zhenglin** [US/US]; Pioneer Hi-Bred International, Inc., 7250 N.W. 62nd Avenue, Johnston, Iowa 50131-0552 (US). **PATTEN, Phillip A.** [US/US]; Pioneer Hi-Bred International, Inc., 7250

N.W. 62nd Avenue, Johnston, Iowa 50131-0552 (US). **YAMAMOTO, Takashi** [US/US]; Pioneer Hi-Bred International, Inc., 7250 N.W. 62nd Avenue, Johnston, Iowa 50131-0552 (US).


(74) Agent: **BAUER, S. Christoper**; Pioneer Hi-Bred International, Inc., 7250 N.W. 62nd Avenue, Johnston, Iowa 50131-0552 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

[Continued on next page]

(54) Title: PHI-4 POLYPEPTIDES AND METHODS FOR THEIR USE



(57) Abstract: Compositions and methods for controlling pests are provided. The methods involve transforming organisms with a nucleic acid sequence encoding an insecticidal protein. In particular, the nucleic acid sequences are useful for preparing plants and microorganisms that possess insecticidal activity. Thus, transformed bacteria, plants, plant cells, plant tissues and seeds are provided. Compositions are insecticidal nucleic acids and proteins of bacterial species. The sequences find use in the construction of expression vectors for subsequent transformation into organisms of interest, as probes for the isolation of other homologous (or partially homologous) genes. The insecticidal proteins find use in controlling, inhibiting growth or killing lepidopteran, coleopteran, dipteran, fungal, hemipteran, and nematode pest populations and for producing compositions with insecticidal activity.

WO 2014/150914 A3



TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, —
KM, ML, MR, NE, SN, TD, TG).

*before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))*

Declarations under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii))*

(88) Date of publication of the international search report:

13 November 2014

Published:

— *with international search report (Art. 21(3))*

PHI-4 POLYPEPTIDES AND METHODS FOR THEIR USE

REFERENCE TO SEQUENCE LISTING SUBMITTED ELECTRONICALLY

The official copy of the sequence listing is submitted electronically via EFS-Web
5 as an ASCII formatted sequence listing with a file named
“5414WOPCT_Sequence_Listing” created on March 04, 2014, and having a size of 5,935
kilobytes and is filed concurrently with the specification. The sequence listing contained
in this ASCII formatted document is part of the specification and is herein incorporated by
reference in its entirety.

10

FIELD

This disclosure relates to the field of molecular biology. Provided are novel genes
that encode pesticidal proteins. These pesticidal proteins and the nucleic acid sequences
encoding them are useful in preparing pesticidal formulations and in the production of
15 transgenic pest-resistant plants.

BACKGROUND

Biological control of insect pests of agricultural significance using a microbial
agent, such as fungi, bacteria or another species of insect affords an environmentally
20 friendly and commercially attractive alternative to synthetic chemical pesticides.
Generally speaking, the use of biopesticides presents a lower risk of pollution and
environmental hazards, and biopesticides provide greater target specificity than is
characteristic of traditional broad-spectrum chemical insecticides. In addition,
biopesticides often cost less to produce and thus improve economic yield for a wide
25 variety of crops.

Certain species of microorganisms of the genus *Bacillus* are known to possess
pesticidal activity against a range of insect pests including Lepidoptera, Diptera,
Coleoptera, Hemiptera and others. *Bacillus thuringiensis* (*Bt*) and *Bacillus popilliae* are
among the most successful biocontrol agents discovered to date. Insect pathogenicity
30 has also been attributed to strains of *B. larvae*, *B. lentimorbus*, *B. sphaericus* and *B.*
cereus. Microbial insecticides, particularly those obtained from *Bacillus* strains, have
played an important role in agriculture as alternatives to chemical pest control.

Crop plants have been developed with enhanced insect resistance by genetically
engineering crop plants to produce pesticidal proteins from *Bacillus*. For example, corn
35 and cotton plants have been genetically engineered to produce pesticidal proteins
isolated from strains of *Bt*. These genetically engineered crops are now widely used in

agriculture and have provided the farmer with an environmentally friendly alternative to traditional insect-control methods. While they have proven to be very successful commercially, these genetically engineered, insect-resistant crop plants provide resistance to only a narrow range of the economically important insect pests. In some cases, insects can develop resistance to different insecticidal compounds, which raises the need to identify alternative biological control agents for pest control.

Accordingly, there remains a need for new pesticidal proteins with different ranges of insecticidal activity against insect pests, e.g., insecticidal proteins which are active against a variety of insects in the order Coleoptera. In addition, there remains a need for biopesticides having activity against a variety of insect pests that have developed resistance to existing pesticides.

SUMMARY

In one aspect compositions and methods for conferring pesticidal activity to bacteria, plants, plant cells, tissues and seeds are provided. Compositions include nucleic acid molecules encoding sequences for pesticidal and insecticidal polypeptides, vectors comprising those nucleic acid molecules, and host cells comprising the vectors. Compositions also include the pesticidal polypeptide sequences and antibodies to those polypeptides. The nucleic acid sequences can be used in DNA constructs or expression cassettes for transformation and expression in organisms, including microorganisms and plants. The nucleotide or amino acid sequences may be synthetic sequences that have been designed for expression in an organism including, but not limited to, a microorganism or a plant. Compositions also comprise transformed bacteria, plants, plant cells, tissues and seeds.

In particular, isolated or recombinant nucleic acid molecules are provided encoding PHI-4 polypeptides including amino acid substitutions, amino acid deletions, amino acid insertions, and fragments thereof, and combinations thereof. Additionally, amino acid sequences corresponding to the PHI-4 polypeptides are encompassed. Nucleic acid sequences that are complementary to a nucleic acid sequence of the embodiments or that hybridize to a sequence of the embodiments are also encompassed.

In another aspect methods are provided for producing the polypeptides and for using those polypeptides for controlling, inhibiting growth or killing a Lepidopteran, Coleopteran, nematode, fungi, Hemipteran and/or Dipteran pests. The transgenic plants of the embodiments express one or more of the pesticidal sequences disclosed herein. In various embodiments, the transgenic plant further comprises one or more additional genes for insect resistance, for example, one or more additional genes for controlling

coleopteran, lepidopteran, hemipteran or nematode pests. It will be understood by one of skill in the art that the transgenic plant may comprise any gene imparting an agronomic trait of interest.

5 In another aspect methods for detecting the nucleic acids and polypeptides of the embodiments in a sample are also included. A kit for detecting the presence of a PHI-4 polypeptide or detecting the presence of a nucleotide sequence encoding a PHI-4 polypeptide in a sample is provided. The kit is provided along with all reagents and control samples necessary for carrying out a method for detecting the intended agent, as well as instructions for use.

10 In another aspect the compositions and methods of the embodiments are useful for the production of organisms with enhanced pest resistance or tolerance. These organisms and compositions comprising the organisms are desirable for agricultural purposes. The compositions of the embodiments are also useful for generating altered or improved proteins that have pesticidal activity or for detecting the presence of PHI-4
15 polypeptides or nucleic acids in products or organisms.

The following embodiments are encompassed by the present disclosure.

Embodiment 1 is a PHI-4 polypeptide having improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35).

20 Embodiment 2 is the PHI-4 polypeptide of embodiment 1, wherein the insecticidal activity is increased about 1.5 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 3 is the PHI-4 polypeptide of embodiment 1, wherein the insecticidal activity is increased about 2 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

25 Embodiment 4 is the PHI-4 polypeptide of embodiment 1, wherein the insecticidal activity is increased about 2.5 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 5 is the PHI-4 polypeptide of embodiment 1, wherein the insecticidal activity is increased about 3 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 6 is the PHI-4 polypeptide of embodiment 1, wherein the insecticidal activity is increased about 5 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

30 Embodiment 7 is the PHI-4 polypeptide of any one of embodiments 1-6, wherein the improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35) is against Western Corn Root Worm (WCRW) larvae.

Embodiment 8 is the PHI-4 polypeptide of any one of embodiments 1-7, wherein the improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35) is quantified as
35 a Mean FAE Index.

Embodiment 9 is the PHI-4 polypeptide of any one of embodiments 1-7, wherein the improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35) is quantified as an EC50 value.

Embodiment 10 is the PHI-4 polypeptide of any one of embodiments 1-7, wherein
5 the improved activity compared to AXMI-205 (SEQ ID NO: 35) is quantified as a Mean Deviation Score.

Embodiment 11 is the PHI-4 polypeptide of any one of embodiments 1-10, wherein the PHI-4 polypeptide comprises one or more amino acid substitutions compared to the native amino acid at position 40, 42, 43, 46, 52, 97, 98, 99, 145, 150, 151, 153, 163, 171,
10 172, 182, 196, 206, 210, 216, 220, 278, 283, 289, 293, 328, 333, 334, 336, 338, 339, 342, 346, 354, 355, 370, 389, 393, 396, 401, 402, 403, 410, 412, 416, 417, 426, 442, 447, 452, 454, 455, 457, 461, 462, 500, 509, 520 or 527 of SEQ ID NO: 35.

Embodiment 12 is the PHI-4 polypeptide of embodiment 11, wherein the amino acid at position 40 is Leu or Ile; the amino acid at position 42 is Asp or Asn; the amino
15 acid at position 43 is Phe or Glu; the amino acid at position 46 is Glu or Asn; the amino acid at position 52 is Ile or Val; the amino acid at position 97 is Arg, Asp, Glu or Asn; the amino acid at position 98 is Tyr or Phe; the amino acid at position 99 is Lys or Leu; the amino acid at position 145 is Leu or Val; the amino acid at position 150 is Arg or Gln; the amino acid at position 151 is Asp or Ser; the amino acid at position 153 is Leu or Ile; the
20 amino acid at position 163 is Leu or Val; the amino acid at position 171 is Tyr or Phe; the amino acid at position 172 is Ile or Leu; the amino acid at position 182 is Asp or Gln; the amino acid at position 196 is Gln or Asn; the amino acid at position 206 is Tyr or Phe; the amino acid at position 210 is Val or Ile; the amino acid at position 216 is Glu or Gln; the amino acid at position 220 is Glu, Gln, His or Asp; the amino acid at position 278 is Glu or
25 Asn; the amino acid at position 283 is Ile or Val; the amino acid at position 289 is Lys, Gln or Leu; the amino acid at position 293 is Arg, Gln or Glu; the amino acid at position 328 is Lys or Glu; the amino acid at position 333 is Ser, Lys or Val; the amino acid at position 334 is Gly, Lys or Arg; the amino acid at position 336 is Gly or Ala; the amino acid at position 338 is Ser or Val; the amino acid at position 339 is Glu, Asn or Gln; the amino
30 acid at position 342 is Ala or Ser; the amino acid at position 346 is Pro or Ala; the amino acid at position 354 is Met or Leu; the amino acid at position 355 is Val or Ile; the amino acid at position 370 is His or Arg; the amino acid at position 389 is Trp or Leu; the amino acid at position 393 is Trp or Leu; the amino acid at position 396 is Ala, Leu, Lys, Thr or Gly; the amino acid at position 401 is Ser, His, Gly, Lys or Pro; the amino acid at position
35 402 is Lys, His, Gly or Trp; the amino acid at position 403 is Asp or Tyr; the amino acid at position 410 is Ile or Val; the amino acid at position 412 is Pro or Ala; the amino acid at

position 416 is Arg or Glu; the amino acid at position 417 is Ala or Ser; the amino acid at position 426 is Thr or Ser; the amino acid at position 442 is Gln or Glu; the amino acid at position 447 is Asp or Lys; the amino acid at position 452 is Gln or Lys; the amino acid at position 454 is Arg or Gln; the amino acid at position 455 is Val or Ile; the amino acid at position 457 is Trp or Asn; the amino acid at position 461 is Thr or Ser; the amino acid at position 462 is Gly or Ala; the amino acid at position 500 is Arg or Gln; the amino acid at position 509 is Lys or Gln; the amino acid at position 520 is Lys, Glu or Gln; and the amino acid at position 527 is Gln or Lys.

Embodiment 13 is the PHI-4 polypeptide of embodiment 11 or 12, further comprising one or more amino acid substitutions at position 86, 359, 464, 465, 466, 467, 468, 499 or 517.

Embodiment 14 is the PHI-4 polypeptide of embodiment 13, wherein the amino acid at position 86 is Glu or Thr; the amino acid at position 359 is Gly or Ala; the amino acid at position 464 is Arg, Ala, Lys, Asp or Asn; the amino acid at position 465 is Lys or Met, the amino acid at position 467 is Val, Ala, Leu or Thr; the amino acid at position 468 is Ser or Leu; the amino acid at position 499 is Glu or Ala or the amino acid at position 517 is Glu or Arg.

Embodiment 15 is the PHI-4 polypeptide of any one of embodiments 1-12, having 1 to 54 amino acid substitutions compared to SEQ ID NO: 35.

Embodiment 16 is the PHI-4 polypeptide of any one of embodiments 1-12, having 1 to 27 amino acid substitutions compared to SEQ ID NO: 35.

Embodiment 17 is the PHI-4 polypeptide of any one of embodiments 1-12, having 1 to 20 amino acid substitutions compared to SEQ ID NO: 35.

Embodiment 18 is the PHI-4 polypeptide of any one of embodiments 1-12, having 1 to 15 amino acid substitutions compared to SEQ ID NO: 35.

Embodiment 19 is the PHI-4 polypeptide of any one of embodiments 13 or 14, comprising 2 to 54 amino acid substitutions compared to SEQ ID NO: 35.

Embodiment 20 is the PHI-4 polypeptide of any one of embodiments 13 or 14, comprising 2 to 27 amino acid substitutions compared to SEQ ID NO: 35.

Embodiment 21 is the PHI-4 polypeptide of any one of embodiments 13 or 14, comprising 2 to 20 amino acid substitutions compared to SEQ ID NO: 35.

Embodiment 22 is the PHI-4 polypeptide of any one of embodiments 13 or 14, comprising 2 to 15 amino acid substitutions compared to SEQ ID NO: 35.

Embodiment 23 is the PHI-4 polypeptide of any one of embodiments 1-22, wherein the PHI-4 polypeptide has at least 80% identity to SEQ ID NO: 35.

Embodiment 24 is the PHI-4 polypeptide of any one of embodiments 1-22, wherein the PHI-4 polypeptide has at least 90% identity to SEQ ID NO: 35.

Embodiment 25 is the PHI-4 polypeptide of any one of embodiments 1-22, wherein the PHI-4 polypeptide has at least 95% identity to SEQ ID NO: 35.

5 Embodiment 26 is the PHI-4 polypeptide of any one of embodiments 1-22, wherein the PHI-4 polypeptide has at least 97% identity to SEQ ID NO: 35.

Embodiment 27 is a PHI-4 polypeptide, comprising at least one amino acid substitution at a residue relative to SEQ ID NO: 35 in a structural domain selected from:

- 10 a hydrophilic residue;
- a residue in a membrane insertion initiation loop;
- a residue in a receptor binding loop; and
- a residue in a protease sensitive region,

wherein the PHI-4 polypeptide has increased insecticidal activity compared to the polypeptide of SEQ ID NO: 35.

15 Embodiment 28 is the PHI-4 polypeptide of embodiment 27, wherein the hydrophilic residues are Asp, Glu, Lys, Arg, His, Ser, Thr, Tyr, Trp, Asn, Gln, and Cys.

Embodiment 29 is the PHI-4 polypeptide of embodiment 27 or 28, wherein the membrane insertion loops are between about the amino acid at position 92 (Val) and about the amino acid at position 101 (Ala) and between about the amino acid at position 20 211 (Gly) and about the amino acid at position 220 (Glu), relative to SEQ ID NO: 35.

Embodiment 30 is the PHI-4 polypeptide of embodiment 29, wherein the membrane insertion initiation loop residue is selected from position 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220 of SEQ ID NO: 35.

25 Embodiment 31 is the PHI-4 polypeptide of any one of embodiments 27-30, wherein the receptor binding loops are between about the amino acid at position 332 (Asp) and about the amino acid at position 340 (Asp), between about the amino acid at position 395 (Asp) and about the amino acid at position 403 (Asp), between amino acid at position 458 (Asp) and about the amino acid at position 466 (Asp) relative to SEQ ID NO: 30 35.

Embodiment 32 is the PHI-4 polypeptide of embodiment 31, wherein the receptor binding loop residue is selected from positions 332, 333, 334, 335, 336, 337, 338, 339, 340, 395, 396, 397, 398, 399, 400, 401, 402, 403, 458, 459, 460, 461, 462, 463, 464, 465, 466 of SEQ ID NO: 35.

35 Embodiment 33 is the PHI-4 polypeptide of any one of embodiments 27-32, wherein the protease sensitive region residue is selected from between about the amino

acid at position 305 (Lys) and about the amino acid at position 316 (Lys) and from about the amino acid at position 500 (Arg) and about the amino acid at position 535 (Lys) relative to SEQ ID NO: 35.

Embodiment 34 is the PHI-4 polypeptide of embodiment 27, wherein the protease
5 is trypsin.

Embodiment 35 is the PHI-4 polypeptide of any one of embodiments 27-34, wherein the PHI-4 polypeptide has at least 80% sequence identity to SEQ ID NO: 35.

Embodiment 36 is the PHI-4 polypeptide of any one of embodiments 27-34, wherein the PHI-4 polypeptide has at least 90% sequence identity to SEQ ID NO: 35.

10 Embodiment 37 is the PHI-4 polypeptide of any one of embodiments 27-34, wherein the PHI-4 polypeptide has at least 95% sequence identity to SEQ ID NO: 35.

Embodiment 38 is the PHI-4 polypeptide of any one of embodiments 27-34, wherein the PHI-4 polypeptide has at least 97% sequence identity to SEQ ID NO: 35.

15 Embodiment 39 is a PHI-4 polypeptide, comprising an amino acid sequence of the formula,

	5	10	15
	Met Xaa Ser Ala Ala Asn Ala Gly Xaa	Leu Gly Asn Leu Xaa Gly	
	20	25	30
	Xaa Thr Ser Xaa Gly Met Xaa Tyr Xaa	Val Asn Gly Leu Tyr Ala	
20	35	40	45
	Ser Pro Glu Ser Leu Xaa Gly Gln Pro	Leu Phe Xaa Xaa Gly Gly	
	50	55	60
	Xaa Leu Asp Ser Xaa Xaa Ile Glu Gly	Xaa Xaa Xaa Xaa Phe Pro	
	65	70	75
25	Xaa Ser Met His Val His Thr Tyr Phe	His Ser Asp Xaa Xaa Gln	
	80	85	90
	Xaa Val Ser Xaa Xaa Ile Xaa Xaa Xaa	Arg Xaa Xaa Xaa Ser Xaa	
	95	100	105
	His Val Gly Xaa Ser Gly Xaa Xaa Xaa	Leu Phe Ser Xaa Ser Xaa	
30	110	115	120
	Ser Val Asp Xaa Thr Thr Xaa Xaa Gln	Gln Leu Xaa Glu Ile Thr	
	125	130	135
	Xaa Ser Ser Thr Arg Glu Xaa His Val	Leu Trp Tyr Ile Ser Leu	
	140	145	150
35	Pro Gly Ala Ala Thr Leu Xaa Ser Met	Leu Xaa Xaa Xaa Xaa Xaa	
	155	160	165
	Xaa Asp Xaa Xaa Xaa Pro Asn Met Xaa	Ala Met Xaa Leu Phe Xaa	

WO 2014/150914

PCT/US2014/024524

					170					175					180
	Xaa	Xaa	Gly	Pro	Xaa	Xaa	Xaa	Xaa	Xaa	Ala	Ala	Val	Gly	Gly	Arg
					185					190					195
	Leu	Xaa	Xaa	Xaa	Xaa	Ala	Ser	Lys	Xaa	Leu	Xaa	Met	Xaa	Ser	Ser
5					200					205					210
	Xaa	Ser	Leu	Ser	Thr	Thr	Xaa	Xaa	Xaa	Ser	Xaa	Xaa	Ala	Xaa	Xaa
					215					220					225
	Gly	Glu	Ile	Xaa	Ile	Xaa	His	Gly	Ser	Xaa	Met	Glu	Lys	Gln	Val
					230					235					240
10	Asn	Ser	Phe	Xaa	Xaa	Xaa	Ser	Thr	Ile	Arg	Xaa	Thr	Ala	Thr	Gly
					245					250					255
	Gly	Lys	Pro	Gly	Xaa	Thr	Xaa	Arg	Ile	Leu	His	Gly	Pro	Asp	Ser
					260					265					270
	Xaa	Xaa	Ala	Phe	Ser	Xaa	Trp	Ala	Xaa	Ser	Leu	Leu	Xaa	Tyr	Ala
15					275					280					285
	Thr	Leu	Met	Asp	Phe	Xaa	Thr	Xaa	Ser	Leu	Xaa	Xaa	Ile	Xaa	Ala
					290					295					300
	Leu	Xaa	Asp	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Glu	Xaa	Xaa	Xaa	Ala	Xaa
					305					310					315
20	Pro	Xaa	Xaa	Met	Xaa	Xaa	Ser	Gln	Xaa	Ser	Ile	Pro	Xaa	Val	Asp
					320					325					330
	Xaa	Val	Leu	Leu	Met	Asp	Ala	Arg	Pro	Pro	Met	Val	Xaa	Ala	Gly
					335					340					345
	Xaa	Asp	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Asp	Xaa	Xaa	Xaa	Xaa	Xaa
25					350					355					360
	Xaa	Ser	Thr	Ser	Xaa	Xaa	Tyr	Lys	Xaa	Xaa	Gly	Gln	Phe	Xaa	Gln
					365					370					375
	Arg	Xaa	His	Xaa	Ser	Val	Ala	Asp	Gly	His	Xaa	Pro	Ile	Xaa	Xaa
					380					385					390
30	Asp	Leu	Phe	Asp	Xaa	Gly	Xaa	Xaa	Xaa	Xaa	Pro	Val	Gly	Xaa	Gln
					395					400					405
	Xaa	Val	Trp	Asp	Xaa	Xaa	Xaa	Xaa	Gly	Lys	Xaa	Xaa	Xaa	Tyr	Xaa
					410					415					420
	Cys	Trp	Arg	Xaa	Xaa	Xaa	Xaa	Gln	Gly	Tyr	Xaa	Xaa	Xaa	Gly	Asp
35					425					430					435
	Val	Xaa	Met	Leu	Ala	Xaa	Ser	Gly	Tyr	Asn	Pro	Pro	Asn	Leu	Pro
					440					445					450

Xaa Xaa Xaa Cys Xaa His Xaa Ser Leu Xaa Ala Xaa Xaa Xaa Thr
 455 460 465
 Leu Xaa Xaa Xaa Xaa Trp Xaa Asp Xaa Xaa Xaa Xaa Xaa Xaa
 470 475 480
 5 Xaa Val Ser Leu Trp Xaa Xaa Gly Ala Ala Gly Ala Val Ala Ser
 485 490 495
 Ser Cys Phe Ala Gly Val Pro Asn Tyr Asn Asn Pro Pro Asn Ser
 500 505 510
 Gly Xaa Ile Xaa Xaa Leu Xaa Gly Ser Ile Ala Cys Val Xaa Thr
 10 515 520 525
 Ser Ala Ile Ala Ser Met Xaa Xaa Met Xaa Ser Met Leu Ser Xaa
 530 535
 His Xaa Gly Met Glu Ala Met Met Ser Lys Leu (SEQ ID NO: 3),
 wherein

- 15 Xaa at position 2 is Ala or Arg; Xaa at position 9 is Gln, Lys or Glu; Xaa at position 14 is Pro or Ala; Xaa at position 16 is Val or Asp; Xaa at position 19 is Met or Leu; Xaa at position 22 is Gly or Ser; Xaa at position 24 is Asp, Asn or Gln; Xaa at position 36 is Leu or Met; Xaa at position 42 is Asp, Asn or Gln; Xaa at position 43 is Phe or Glu; Xaa at position 46 is Glu, Asp, Asn or Gly; Xaa at position 50 is Ile or Val; Xaa at position 51 is Glu or Gln; Xaa at position 55 is Arg or Lys; Xaa at position 56 is Ser or Thr; Xaa at position 57 is Tyr or Phe; Xaa at position 58 is Thr or Ser; Xaa at position 61 is Arg, Lys or Glu; Xaa at position 73 is Phe or Tyr; Xaa at position 74 is Lys, Glu, Gly, Arg, Met, Leu, His or Asp; Xaa at position 76 is Asp or Gln; Xaa at position 79 is Lys or Glu; Xaa at position 80 is Glu or Ser; Xaa at position 82 is Glu, Ile, Leu, Tyr or Gln; Xaa at position 83 is Glu or Gln; Xaa at position 84 is Tyr or Phe; Xaa at position 86 is Glu or Gln; Xaa at position 87 is Lys or Gln; Xaa at position 88 is Met, Ile or Leu; Xaa at position 90 is Gln or Glu; Xaa at position 94 is Val or Ile; Xaa at position 97 is Arg, Asn, Asp, Glu, Gln, Gly, Ser, Ile, Phe, His, Lys, Thr, Asn, Tyr, Trp, Pro, Cys, Ala, Met, Val or Leu; Xaa at position 98 is Tyr or Phe; Xaa at position 99 is Lys, Leu, Tyr, Ile, Met, Phe, Cys, Val or Asn; Xaa at position 103 is Ala or Gly; Xaa at position 105 is Leu or Ile; Xaa at position 109 is Phe, Lys, Gly, Met, Ser, Asp, Asn, Glu, Cys, Ala or Arg; Xaa at position 112 is Thr or Ser; Xaa at position 113 is Asp, Glu or Met; Xaa at position 117 is Thr or Ser; Xaa at position 121 is Tyr or Phe; Xaa at position 127 is Ala or Thr; Xaa at position 142 is Arg or Glu; Xaa at position 146 is Arg or Gln; Xaa at position 147 is Arg, Glu or Gln; Xaa at position 148 is Asp, Phe, Pro, Val, Glu, His, Trp, Ala, Arg, Leu, Ser, Gln or Gly; Xaa at position 149 is Phe or Val; Xaa at position 150 is Arg, Gln, Glu or Asn; Xaa at position 151 is Asp, Ser, Ala, Asn, Trp, Val, Gln, Cys, Met, Leu, Arg or Glu; Xaa at position 153 is Leu or Ile; Xaa at

position 154 is Asn or Asp; Xaa at position 155 is Asn or Lys; Xaa at position 159 is Pro or Asp; Xaa at position 162 is Glu, Asp, Gln, Asn or Leu; Xaa at position 165 is Lys, Glu, Gln, Pro, Thr, Ala, Leu, Gly, Asp, Val, His, Ile, Met, Trp, Phe, Tyr or Arg; Xaa at position 166 is Arg or Gln; Xaa at position 167 is Tyr, Trp or Cys; Xaa at position 170 is Tyr or His;

5 Xaa at position 171 is Tyr or Phe; Xaa at position 172 is Ile, Leu or Val; Xaa at position 173 is Ser or Ala; Xaa at position 174 is Glu, Gln, Asn, Lys, Val or Ser; Xaa at position 182 is Asp or Gln; Xaa at position 183 is Tyr or Val; Xaa at position 184 is Ser or Thr; Xaa at position 185 is Ala or Ser; Xaa at position 189 is Thr, Lys or Ile; Xaa at position 191 is Lys or Gln; Xaa at position 193 is Asp or Asn; Xaa at position 196 is Gln, Lys, Asn, Asp,

10 Glu, Ala, Ile or Arg; Xaa at position 202 is Ala or Val; Xaa at position 203 is Glu, Thr or His; Xaa at position 204 is Met or Ala; Xaa at position 206 is Tyr or Phe; Xaa at position 207 is Lys or Gln; Xaa at position 209 is Leu or Pro; Xaa at position 210 is Val or Ile; Xaa at position 214 is Lys, Ser or Gln; Xaa at position 216 is Glu, Gln, Phe, Val, Tyr or Arg; Xaa at position 220 is Glu, His, Asp, Thr, Tyr, Val, Ser, Gln, Arg, Trp, Met, Ala, Phe, Ile,

15 Leu, Cys or Asn; Xaa at position 229 is Arg or Glu; Xaa at position 230 is Ser or Glu; Xaa at position 231 is Asn or Ser; Xaa at position 236 is Leu or Pro; Xaa at position 245 is Met or Leu; Xaa at position 247 is Asp or Tyr; Xaa at position 256 is Gln, Lys or Glu; Xaa at position 257 is Gln, Ile, Glu, Cys, Ser, His, Trp or Met; Xaa at position 261 is Gln, Glu, Lys or Ala; Xaa at position 264 is Glu or Gln; Xaa at position 268 is Asp or Asn; Xaa at

20 position 276 is Ser or Ala; Xaa at position 278 is Glu, Asn or Gln; Xaa at position 281 is Gln, Lys or Glu; Xaa at position 282 is Pro or Gly; Xaa at position 284 is Trp or Arg; Xaa at position 287 is Ala or Cys; Xaa at position 289 is Lys, Leu, Val, Pro, Glu, Gln, Tyr, Thr, Asp, Phe, Ser, Met, Arg, Trp, Ile, His, Asn, Cys, Gly or Ala; Xaa at position 291 is Glu or Gln; Xaa at position 292 is Arg or Gln; Xaa at position 293 is Arg, Glu or Gln; Xaa at

25 position 294 is Val or Ala; Xaa at position 296 is Leu or Ile; Xaa at position 297 is Glu or Gln; Xaa at position 298 is Asp or Gln; Xaa at position 300 is Phe or Tyr; Xaa at position 302 is Glu or Gln; Xaa at position 303 is Phe or Tyr; Xaa at position 305 is Lys, Gln, Ala, Ile, Met, Asn, Thr or Val; Xaa at position 306 is Gln or Lys; Xaa at position 309 is Gln, Lys or Glu; Xaa at position 313 is Lys, Gln or Arg; Xaa at position 316 is Lys or Gln; Xaa at

30 position 328 is Lys, Glu or Gln; Xaa at position 331 is Glu, Asn or Gln; Xaa at position 333 is Ser, Arg, Gly, Lys, Val, Asn, Ala, His, Gln, Thr, Asp, Ile, Leu, Cys or Glu; Xaa at position 334 is Gly, Arg, Lys, Ile or Trp; Xaa at position 335 is Ser or Ala; Xaa at position 336 is Gly or Ala; Xaa at position 337 is Ala, Val or Gly; Xaa at position 338 is Ser, His, Val, Lys, Ala, Gly, Thr, Ile, Glu, Met, Arg, Pro, Asp, Asn or Leu; Xaa at position 339 is Glu, Asn,

35 Gln, Ile, Pro, Met, Ser, Ala, Cys, Phe, Val, Leu, Asp, Trp, His or Arg; Xaa at position 341 is Leu or Val; Xaa at position 342 is Ala, Ser or Val; Xaa at position 343 is Val or Ile; Xaa

at position 344 is Phe or Trp; Xaa at position 345 is Asn or His; Xaa at position 346 is Pro
 or Ala; Xaa at position 350 is Asn or Ser; Xaa at position 351 is Gly or Val; Xaa at position
 354 is Met or Leu; Xaa at position 355 is Val, Ile or Leu; Xaa at position 359 is Gly or Ala;
 Xaa at position 362 is Asn or Ser; Xaa at position 364 is Ala or Ser; Xaa at position 371 is
 5 Ala, Gly or Thr; Xaa at position 374 is Phe or Ile; Xaa at position 375 is Lys or Arg; Xaa at
 position 380 is Leu or Gly; Xaa at position 382 is Val, Asp or Leu; Xaa at position 383 is
 Leu, Ile or Val; Xaa at position 384 is Lys, Ala or Gly; Xaa at position 385 is Ala or Gly;
 Xaa at position 389 is Trp or Tyr; Xaa at position 391 is Arg, Leu, Glu, Gln, Asp or His;
 Xaa at position 395 is Asp or Cys; Xaa at position 396 is Ala, Leu, Lys, Asn, Gly, Ile, Met,
 10 Arg, Tyr, Gln, His or Thr; Xaa at position 397 is Gly, Arg or Ala; Xaa at position 398 is Ser,
 Gln or Cys; Xaa at position 401 is Ser, His, Pro, Gly, Lys, Val, Arg, Ile, Asn, Phe, Thr, Ala,
 Asp, Met, Gln or Glu; Xaa at position 402 is Lys, Phe, His, Arg, Trp, Gly, Asn, Leu, Tyr,
 Thr, Val, Met, Pro or Ala; Xaa at position 403 is Asp, Tyr, Trp, Phe or Glu; Xaa at position
 405 is Ala or Ser; Xaa at position 409 is Ala or Pro; Xaa at position 410 is Ile or Val; Xaa
 15 at position 411 is Pro or Ala; Xaa at position 412 is Pro or Ala; Xaa at position 416 is Arg,
 Glu or Gln; Xaa at position 417 is Ala, Ser or Cys; Xaa at position 418 is Leu or Met; Xaa
 at position 422 is Met or Val; Xaa at position 426 is Thr or Ser; Xaa at position 436 is Asp
 or Lys; Xaa at position 437 is Tyr or Val; Xaa at position 438 is Val or Arg; Xaa at position
 440 is Val or Leu; Xaa at position 442 is Gln, Lys or Glu; Xaa at position 445 is Cys, Leu
 20 or Thr; Xaa at position 447 is Asp, Lys, Tyr, Ser, Glu, Ile, Gly, Pro, Leu, Phe, Trp or Thr;
 Xaa at position 448 is Val or Ala; Xaa at position 449 is Gln or Glu; Xaa at position 452 is
 Gln, Lys or Glu, Ala; Xaa at position 453 is Asn or Asp; Xaa at position 454 is Arg, Tyr,
 Met, Ser, Val, Ile, Lys, Phe, Trp, Gln, Gly, His, Asp, Leu, Thr, Pro or Asn; Xaa at position
 455 is Val or Ile; Xaa at position 457 is Trp or Asn; Xaa at position 459 is Lys, Met, Val,
 25 Trp, Gln, Ile, Thr, Ser, His, Cys, Tyr, Pro, Asn, Ala, Arg or Glu; Xaa at position 460 is Gly
 or Ala; Xaa at position 461 is Thr or Ser; Xaa at position 462 is Gly or Ala; Xaa at position
 463 is Ala, Ser or Gly; Xaa at position 464 is Arg, Gly, His, Gln, Thr, Phe, Ala, Asp, Ser or
 Lys; Xaa at position 465 is Lys, Asn, Val, Met, Pro, Gly, Arg, Thr, His, Cys, Trp, Phe or
 Leu; Xaa at position 466 is Asp or Arg; Xaa at position 471 is Gln, Lys, Glu or Met; Xaa at
 30 position 472 is Pro or Ser; Xaa at position 497 is Asp or Gln; Xaa at position 499 is Glu or
 Gln; Xaa at position 500 is Arg, Gln or Lys; Xaa at position 502 is Arg, Glu or Gln; Xaa at
 position 509 is Lys, Gln, Glu or Ala; Xaa at position 517 is Gln, Cys, Asn, Val or Pro; Xaa
 at position 518 is Glu or Gln; Xaa at position 520 is Lys, Gln, Glu, His or Ala; Xaa at
 position 525 is Gln or Lys; and Xaa at position 527 is Gln, Lys, Pro, Cys, Glu, Ser, His,
 35 Phe or Trp; wherein one or more amino acid(s) designated by Xaa in SEQ ID NO: 3 is an
 amino acid different from the corresponding amino acid of SEQ ID NO: 35 and wherein the

PHI-4 polypeptide has increased insecticidal activity compared to the polypeptide of SEQ ID NO: 35.

Embodiment 40 is a PHI-4 polypeptide, comprising an amino acid sequence of the formula

5					5					10					15
	Met	Xaa	Ser	Ala	Ala	Asn	Ala	Gly	Gln	Leu	Gly	Asn	Leu	Pro	Gly
					20					25					30
	Val	Thr	Ser	Met	Gly	Met	Gly	Tyr	Xaa	Val	Asn	Gly	Leu	Tyr	Ala
					35					40					45
	10	Ser	Pro	Glu	Ser	Leu	Leu	Gly	Gln	Pro	Leu	Phe	Xaa	Xaa	Gly
				50					55					60	
	Xaa	Leu	Asp	Ser	Ile	Glu	Ile	Glu	Gly	Arg	Ser	Tyr	Thr	Phe	Pro
					65					70					75
	Arg	Ser	Met	His	Val	His	Thr	Tyr	Phe	His	Ser	Asp	Phe	Xaa	Gln
	15					80					85				
Asp		Val	Ser	Xaa	Glu	Ile	Xaa	Glu	Tyr	Arg	Glu	Lys	Met	Ser	Gln
					95					100					105
	His	Val	Gly	Val	Ser	Gly	Xaa	Xaa	Xaa	Leu	Phe	Ser	Ala	Ser	Leu
					110					115					120
	20	Ser	Val	Asp	Xaa	Thr	Thr	Thr	Asp	Gln	Gln	Leu	Thr	Glu	Ile
				125					130					135	
	Tyr	Ser	Ser	Thr	Arg	Glu	Ala	His	Val	Leu	Trp	Tyr	Ile	Ser	Leu
					140					145					150
	Pro	Gly	Ala	Ala	Thr	Leu	Arg	Ser	Met	Leu	Arg	Xaa	Xaa	Phe	Xaa
	25					155					160				
Xaa		Asp	Xaa	Asn	Asn	Pro	Asn	Met	Pro	Ala	Met	Xaa	Leu	Phe	Xaa
					170					175					180
	Xaa	Tyr	Gly	Pro	Tyr	Xaa	Ile	Ser	Xaa	Ala	Ala	Val	Gly	Gly	Arg
					185					190					195
	30	Leu	Xaa	Tyr	Ser	Ala	Ala	Ser	Lys	Thr	Leu	Lys	Met	Asp	Ser
				200					205					210	
	Xaa	Ser	Leu	Ser	Thr	Thr	Ala	Xaa	Met	Ser	Xaa	Lys	Ala	Leu	Val
					215					220					225
	Gly	Glu	Ile	Lys	Ile	Xaa	His	Gly	Ser	Xaa	Met	Glu	Lys	Gln	Val
	35					230					235				
Asn		Ser	Phe	Arg	Ser	Asn	Ser	Thr	Ile	Arg	Leu	Thr	Ala	Thr	Gly
					245					250					255

	Gly	Lys	Pro	Gly	Met	Thr	Xaa	Arg	Ile	Leu	His	Gly	Pro	Asp	Ser	
					260					265					270	
	Xaa	Xaa	Ala	Phe	Ser	Xaa	Trp	Ala	Glu	Ser	Leu	Leu	Asp	Tyr	Ala	
					275					280					285	
5	Thr	Leu	Met	Asp	Phe	Ser	Thr	Xaa	Ser	Leu	Xaa	Pro	Ile	Trp	Ala	
					290					295					300	
	Leu	Ala	Asp	Xaa	Pro	Glu	Arg	Xaa	Val	Glu	Leu	Glu	Asp	Ala	Phe	
					305					310					315	
	Pro	Glu	Phe	Met	Lys	Gln	Ser	Gln	Gln	Ser	Ile	Pro	Xaa	Val	Asp	
10					320					325					330	
	Lys	Val	Leu	Leu	Met	Asp	Ala	Arg	Pro	Pro	Met	Val	Xaa	Ala	Gly	
					335					340					345	
	Glu	Asp	Xaa	Xaa	Ser	Xaa	Ala	Xaa	Xaa	Asp	Leu	Ala	Xaa	Phe	Asn	
					350					355					360	
15	Xaa	Ser	Thr	Ser	Asn	Gly	Tyr	Lys	Met	Xaa	Gly	Gln	Phe	Xaa	Gln	
					365					370					375	
	Arg	Asn	His	Ala	Ser	Val	Ala	Asp	Gly	His	Ala	Pro	Ile	Phe	Lys	
					380					385					390	
	Asp	Leu	Phe	Asp	Leu	Gly	Val	Leu	Lys	Ala	Pro	Val	Gly	Trp	Gln	
20					395					400					405	
	Xaa	Val	Trp	Asp	Asp	Xaa	Gly	Ser	Gly	Lys	Xaa	Xaa	Xaa	Tyr	Ala	
					410					415					420	
	Cys	Trp	Arg	Ala	Ile	Xaa	Xaa	Gln	Gly	Tyr	Xaa	Xaa	Xaa	Gly	Asp	
					425					430					435	
25	Val	Met	Met	Leu	Ala	Xaa	Ser	Gly	Tyr	Asn	Pro	Pro	Asn	Leu	Pro	
					440					445					450	
	Asp	Tyr	Val	Cys	Xaa	His	Gln	Ser	Leu	Cys	Ala	Xaa	Val	Gln	Thr	
					455					460					465	
	Leu	Xaa	Asn	Xaa	Xaa	Trp	Trp	Asp	Xaa	Gly	Xaa	Xaa	Xaa	Xaa	Xaa	
30					470					475					480	
	Asp	Val	Ser	Leu	Trp	Xaa	Xaa	Gly	Ala	Ala	Gly	Ala	Val	Ala	Ser	
					485					490					495	
	Ser	Cys	Phe	Ala	Gly	Val	Pro	Asn	Tyr	Asn	Asn	Pro	Pro	Asn	Ser	
					500					505					510	
35	Gly	Asp	Ile	Glu	Xaa	Leu	Arg	Gly	Ser	Ile	Ala	Cys	Val	Xaa	Thr	
					515					520					525	
	Ser	Ala	Ile	Ala	Ser	Met	Gln	Glu	Met	Xaa	Ser	Met	Leu	Ser	Gln	

530

535

His Xaa Gly Met Glu Ala Met Met Ser Lys Leu (SEQ ID NO: 4),

wherein

Xaa at position 2 is Ala or Arg; Xaa at position 24 is Asp or Asn; Xaa at position 42 is Asp
 5 or Asn; Xaa at position 43 is Phe or Glu; Xaa at position 46 is Glu or Asn; Xaa at position
 74 is Lys, Glu or Gly; Xaa at position 79 is Lys or Glu; Xaa at position 82 is Glu, Ile, Leu or
 Tyr; Xaa at position 97 is Arg, Asn, Asp, Glu, Gln, Gly, Ser, Ile, Phe, His, Lys, Thr, Asn,
 Tyr, Trp, Pro, Cys, Ala, Met, Val or Leu; Xaa at position 98 is Tyr or Phe; Xaa at position
 99 is Lys, Leu, Tyr, Ile or Met; Xaa at position 109 is Phe, Lys, Gly, Met, Ser, Asp or Asn;
 10 Xaa at position 147 is Arg or Glu; Xaa at position 148 is Asp, Phe or Pro; Xaa at position
 150 is Arg, Gln, Glu or Asn; Xaa at position 151 is Asp, Ser, Ala or Asn; Xaa at position
 153 is Leu or Ile; Xaa at position 162 is Glu, Asp, Gln, Asn or Leu; Xaa at position 165 is
 Lys, Glu or Gln; Xaa at position 166 is Arg or Gln; Xaa at position 171 is Tyr or Phe; Xaa
 at position 174 is Glu, Gln, Asn, Lys, Val or Ser; Xaa at position 182 is Asp or Gln; Xaa at
 15 position 196 is Gln, Lys, Asn or Asp; Xaa at position 203 is Glu, Thr or His; Xaa at position
 206 is Tyr or Phe; Xaa at position 216 is Glu or Gln; Xaa at position 220 is Glu, His, Asp,
 Thr, Tyr, Val, Ser or Gln; Xaa at position 247 is Asp or Tyr; Xaa at position 256 is Gln or
 Lys; Xaa at position 257 is Gln or Ile; Xaa at position 261 is Gln, Glu, Lys or Ala; Xaa at
 position 278 is Glu or Asn; Xaa at position 281 is Gln, Lys or Glu; Xaa at position 289 is
 20 Lys, Leu, Val, Pro, Glu, Gln, Tyr, Thr or Asp; Xaa at position 293 is Arg, Glu or Gln; Xaa at
 position 313 is Lys or Gln; Xaa at position 328 is Lys, Glu or Gln; Xaa at position 333 is
 Ser, Gly, Lys, Val or Asn; Xaa at position 334 is Gly, Arg, Lys or Ile; Xaa at position 336 is
 Gly or Ala; Xaa at position 338 is Ser, His, Val, Lys or Ala; Xaa at position 339 is Glu, Asn,
 Ile or Pro; Xaa at position 343 is Val or Ile; Xaa at position 346 is Pro or Ala; Xaa at
 25 position 355 is Val or Ile; Xaa at position 359 is Gly or Ala; Xaa at position 391 is Arg,
 Leu, Glu, Gln, Asp or His; Xaa at position 396 is Ala, Leu, Lys, Asn, Gly or Thr; Xaa at
 position 401 is Ser, His, Pro, Gly, Lys, Val or Arg; Xaa at position 402 is Lys, Phe, His,
 Arg, Gly, Trp, Thr, Asn, Tyr or Met; Xaa at position 403 is Asp or Tyr; Xaa at position 411
 is Pro or Ala; Xaa at position 412 is Pro or Ala; Xaa at position 416 is Arg or Glu; Xaa at
 30 position 417 is Ala or Ser; Xaa at position 418 is Leu or Met; Xaa at position 426 is Thr or
 Ser; Xaa at position 440 is Val or Leu; Xaa at position 447 is Asp, Lys, Tyr, Ser, Glu or Ile;
 Xaa at position 452 is Gln, Lys or Glu; Xaa at position 454 is Arg, Tyr, Met, Ser, Val, Ile,
 Lys, Phe, Trp or Gln; Xaa at position 455 is Val or Ile; Xaa at position 459 is Lys, Met, Val,
 Trp, Gln, Ile or Tyr; Xaa at position 461 is Thr or Ser; Xaa at position 462 is Gly or Ala;
 35 Xaa at position 463 is Ala or Ser; Xaa at position 464 is Arg, Gly, His, Ala, Asp, Ser or
 Lys; Xaa at position 465 is Lys, Asn, Val, Met, Pro, Gly or Arg; Xaa at position 471 is Gln,

Lys, Glu or Met; Xaa at position 472 is Pro or Ser; Xaa at position 500 is Arg or Gln; Xaa at position 509 is Lys, Gln or Ala; Xaa at position 520 is Lys, Gln, Glu, His or Ala; and Xaa at position 527 is Gln, Lys, Pro, Cys or Glu; wherein one or more amino acid(s) designated by Xaa in SEQ ID NO: 4 is an amino acid different from the corresponding amino acid of SEQ ID NO: 35 and wherein the PHI-4 polypeptide has increased insecticidal activity compared to the polypeptide of SEQ ID NO: 35.

Embodiment 41 is the PHI-4 polypeptide of embodiment 39 or 40, further comprising one or more amino acid substitutions at position 86, 359, 464, 465, 466, 467, 468, 499 or 517 of SEQ ID NO: 3 or SEQ ID NO: 4.

Embodiment 42 is the PHI-4 polypeptide of embodiment 41, wherein the amino acid at position 86 is Glu or Thr; the amino acid at position 359 is Gly or Ala; the amino acid at position 464 is Arg, Ala, Lys, Asp or Asn; the amino acid at position 465 is Lys or Met, the amino acid at position 467 is Val, Ala, Leu or Thr; the amino acid at position 468 is Ser or Leu; the amino acid at position 499 is Glu or Ala or the amino acid at position 517 is Glu or Arg.

Embodiment 43 is the PHI-4 polypeptide of embodiment 39-42, further comprising one or more conservative amino acid substitution, insertion of one or more amino acids, deletion of one or more amino acids, and combinations thereof.

Embodiment 44 is the PHI-4 polypeptide of any one of embodiments 39 - 43, wherein the insecticidal activity is increased about 1.5 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 45 is the PHI-4 polypeptide of any one of embodiments 39 - 43, wherein the insecticidal activity is increased about 2 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 46 is the PHI-4 polypeptide of any one of embodiments 39 - 43, wherein the insecticidal activity is increased about 2.5 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 47 is the PHI-4 polypeptide of any one of embodiments 39 - 43, wherein the insecticidal activity is increased about 3 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 48 is the PHI-4 polypeptide of any one of embodiments 39 - 43, wherein the insecticidal activity is increased about 5 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 49 is the PHI-4 polypeptide of any one of embodiments 39-48, wherein the improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35) is against Western Corn Root Worm (WCRW) larvae.

Embodiment 50 is the PHI-4 polypeptide of any one of embodiments 39-49, wherein the improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35) is quantified as a Mean FAE Index.

Embodiment 51 is the PHI-4 polypeptide of any one of embodiments 39-49, wherein the improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35) is quantified as an EC50 value.

Embodiment 52 is the PHI-4 polypeptide of any one of embodiments 39-49, wherein the improved activity compared to AXMI-205 (SEQ ID NO: 35) is quantified as a Mean Deviation Score.

Embodiment 53 is the PHI-4 polypeptide of any one of embodiments 39-49, having 1 to 54 amino acid substitutions at a position(s) designated as Xaa in SEQ ID NO: 3 or 4.

Embodiment 54 is the PHI-4 polypeptide of any one of embodiments 39-49, having 1 to 27 amino acid substitutions at a position(s) designated as Xaa in SEQ ID NO: 3 or 4.

Embodiment 55 is the PHI-4 polypeptide of any one of embodiments 39-49, having 1 to 20 amino acid substitutions at a position(s) designated as Xaa in SEQ ID NO: 3 or 4.

Embodiment 56 is the PHI-4 polypeptide of any one of embodiments 39-49, having 1 to 15 amino acid substitutions at a position(s) designated as Xaa in SEQ ID NO: 3 or 4.

Embodiment 57 is the PHI-4 polypeptide of any one of embodiments 1-56, wherein 1-25 amino acids are deleted from the N-terminus of the PHI-4 polypeptide and/or C-terminus of the PHI-4 polypeptide.

Embodiment 58 is the PHI-4 polypeptide of any one of embodiments 1-53, wherein 1-20 amino acids are deleted from the C-terminus of the PHI-4 polypeptide.

Embodiment 59 is a polynucleotide encoding a PHI-4 polypeptide, wherein the PHI-4 polypeptide has improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 60 is the polynucleotide of embodiment 59, wherein the insecticidal activity of the PHI-4 polypeptide is increased about 1.5 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 61 is the polynucleotide of embodiment 59, wherein the insecticidal activity of the PHI-4 polypeptide is increased about 2 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 62 is the polynucleotide of embodiment 59, wherein the insecticidal activity of the PHI-4 polypeptide is increased about 2.5 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 63 is the polynucleotide of embodiment 59, wherein the insecticidal activity of the PHI-4 polypeptide is increased about 3 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

5 Embodiment 64 is the polynucleotide of embodiment 59, wherein the insecticidal activity of the PHI-4 polypeptide is increased about 5 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 65 is the polynucleotide of any one of embodiments 59-64, wherein the improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35) is against Western Corn Root Worm (WCRW) larvae.

10 Embodiment 66 is the polynucleotide of any one of embodiments 59-64, wherein the improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35) is quantified as a Mean FAE Index.

Embodiment 67 is the polynucleotide of any one of embodiments 59-64, wherein the improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35) is quantified as
15 an EC50 value.

Embodiment 68 is the polynucleotide of any one of embodiments 59-64, wherein the improved activity compared to AXMI-205 (SEQ ID NO: 35) is quantified as a Mean Deviation Score.

Embodiment 69 is the polynucleotide of any one of embodiments 59-68, wherein
20 the PHI-4 polypeptide comprises one or more amino acid substitutions compared to the native amino acid at position 40, 42, 43, 46, 52, 97, 98, 99, 145, 150, 151, 153, 163, 171, 172, 182, 196, 206, 210, 216, 220, 278, 283, 289, 293, 328, 333, 334, 336, 338, 339, 342, 346, 354, 355, 370, 389, 393, 396, 401, 402, 403, 410, 412, 416, 417, 426, 442, 447, 452, 454, 455, 457, 461, 462, 500, 509, 520 or 527 of SEQ ID NO: 35.

25 Embodiment 70 is the polynucleotide of any one of embodiments 69, wherein the amino acid at position 40 is Leu or Ile; the amino acid at position 42 is Asp or Asn; the amino acid at position 43 is Phe or Glu; the amino acid at position 46 is Glu or Asn; the amino acid at position 52 is Ile or Val; the amino acid at position 97 is Arg, Asp, Glu or Asn; the amino acid at position 98 is Tyr or Phe; the amino acid at position 99 is Lys or
30 Leu; the amino acid at position 145 is Leu or Val; the amino acid at position 150 is Arg or Gln; the amino acid at position 151 is Asp or Ser; the amino acid at position 153 is Leu or Ile; the amino acid at position 163 is Leu or Val; the amino acid at position 171 is Tyr or Phe; the amino acid at position 172 is Ile or Leu; the amino acid at position 182 is Asp or Gln; the amino acid at position 196 is Gln or Asn; the amino acid at position 206 is Tyr or
35 Phe; the amino acid at position 210 is Val or Ile; the amino acid at position 216 is Glu or Gln; the amino acid at position 220 is Glu, Gln, His or Asp; the amino acid at position 278

is Glu or Asn; the amino acid at position 283 is Ile or Val; the amino acid at position 289 is Lys, Gln or Leu; the amino acid at position 293 is Arg, Gln or Glu; the amino acid at position 328 is Lys or Glu; the amino acid at position 333 is Ser, Lys or Val; the amino acid at position 334 is Gly, Lys or Arg; the amino acid at position 336 is Gly or Ala; the amino acid at position 338 is Ser or Val; the amino acid at position 339 is Glu, Asn or Gln; the amino acid at position 342 is Ala or Ser; the amino acid at position 346 is Pro or Ala; the amino acid at position 354 is Met or Leu; the amino acid at position 355 is Val or Ile; the amino acid at position 370 is His or Arg; the amino acid at position 389 is Trp or Leu; the amino acid at position 393 is Trp or Leu; the amino acid at position 396 is Ala, Leu, Lys, Thr or Gly; the amino acid at position 401 is Ser, His, Gly, Lys or Pro; the amino acid at position 402 is Lys, His, Gly or Trp; the amino acid at position 403 is Asp or Tyr; the amino acid at position 410 is Ile or Val; the amino acid at position 412 is Pro or Ala; the amino acid at position 416 is Arg or Glu; the amino acid at position 417 is Ala or Ser; the amino acid at position 426 is Thr or Ser; the amino acid at position 442 is Gln or Glu; the amino acid at position 447 is Asp or Lys; the amino acid at position 452 is Gln or Lys; the amino acid at position 454 is Arg or Gln; the amino acid at position 455 is Val or Ile; the amino acid at position 457 is Trp or Asn; the amino acid at position 461 is Thr or Ser; the amino acid at position 462 is Gly or Ala; the amino acid at position 500 is Arg or Gln; the amino acid at position 509 is Lys or Gln; the amino acid at position 520 is Lys, Glu or Gln; and the amino acid at position 527 is Gln or Lys.

Embodiment 71 is the polynucleotide of embodiment 69 or 70, wherein the PHI-4 polypeptide further comprises one or more amino acid substitutions compared to the native amino acid at position 86, 359, 464, 465, 466, 467, 468, 499 or 517 of SEQ ID NO: 35.

Embodiment 72 is the polynucleotide of embodiment 71, wherein the amino acid at position 86 is Glu or Thr; the amino acid at position 359 is Gly or Ala; the amino acid at position 464 is Arg, Ala, Lys, Asp or Asn; the amino acid at position 465 is Lys or Met, the amino acid at position 467 is Val, Ala, Leu or Thr; the amino acid at position 468 is Ser or Leu; the amino acid at position 499 is Glu or Ala or the amino acid at position 517 is Glu or Arg.

Embodiment 73 is the polynucleotide of any one of embodiments 59-69 and 72, wherein the PHI-4 polypeptide has 1 to 54 amino acid substitutions compared to SEQ ID NO: 35.

Embodiment 74 is the polynucleotide of any one of embodiments 59-70, wherein the PHI-4 polypeptide has 1 to 27 amino acid substitutions compared to SEQ ID NO: 2.

Embodiment 75 is the polynucleotide of any one of embodiments 59-70, wherein the PHI-4 polypeptide has 1 to 20 amino acid substitutions compared to SEQ ID NO: 35.

Embodiment 76 is the polynucleotide of any one of embodiments 59-70, wherein the PHI-4 polypeptide has 1 to 15 amino acid substitutions compared to SEQ ID NO: 35.

5 Embodiment 77 is the polynucleotide of embodiments 71 or 72, wherein the PHI-4 polypeptide has 2 to 54 amino acid substitutions compared to SEQ ID NO: 35.

Embodiment 78 is the polynucleotide of embodiments 71 or 72, wherein the PHI-4 polypeptide has 2 to 27 amino acid substitutions compared to SEQ ID NO: 35.

10 Embodiment 79 is the polynucleotide of embodiments 71 or 72, wherein the PHI-4 polypeptide has 2 to 20 amino acid substitutions compared to SEQ ID NO: 35.

Embodiment 80 is the polynucleotide of embodiments 71 or 72, wherein the PHI-4 polypeptide has 2 to 15 amino acid substitutions compared to SEQ ID NO: 35.

Embodiment 81 is the polynucleotide of any one of embodiments 59-80, wherein the PHI-4 polypeptide has at least 80% identity to SEQ ID NO: 35.

15 Embodiment 82 is the polynucleotide of any one of embodiments 59-80, wherein the PHI-4 polypeptide has at least 90% identity to SEQ ID NO: 35.

Embodiment 83 is the polynucleotide of any one of embodiments 59-80, wherein the PHI-4 polypeptide has at least 95% identity to SEQ ID NO: 35.

20 Embodiment 84 is the polynucleotide of any one of embodiments 59-80, wherein the PHI-4 polypeptide has at least 97% identity to SEQ ID NO: 35.

Embodiment 85 is a polynucleotide encoding a PHI-4 polypeptide, wherein the PHI-4 polypeptide has at least one amino acid substitution at a residue relative to SEQ ID NO: 35 in a structural domain selected from:

- 25
- a hydrophilic residue;
 - a residue in a membrane insertion initiation loop;
 - a residue in a receptor binding loop; and
 - a residue in a protease sensitive region,

wherein the PHI-4 polypeptide has increased insecticidal activity compared to the polypeptide of SEQ ID NO: 35.

30 Embodiment 86 is the polynucleotide of embodiment 85, wherein the hydrophilic residues are Asp, Glu, Lys, Arg, His, Ser, Thr, Tyr, Trp, Asn, Gln, and Cys.

Embodiment 87 is the polynucleotide of embodiment 85 or 86, wherein the membrane insertion loops are between about the amino acid at position 92 (Val) and about the amino acid at position 101 (Ala) and between about the amino acid 211 (Gly) and about the amino acid 220 (Glu), relative to SEQ ID NO: 35.

35

Embodiment 88 is the polynucleotide of embodiment 87, wherein the PHI-4 polypeptide has one or more amino substitution compared to the native amino acid at position 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 211, 212, 213, 214, 215, 216, 217, 218, 219, and 220 of SEQ ID NO: 35.

5 Embodiment 89 is the polynucleotide of any one of embodiments 85, 86, 87 or 88, wherein the receptor binding loops are between about the amino acid at position 332 (Asp) and about the amino acid at position 340 (Asp), between about the amino acid at position 395 (Asp) and about the amino acid at position 403 (Asp), and between about the amino acid at position 458 (Asp) and about the amino acid 466 (Asp), relative to SEQ ID
10 NO: 35.

Embodiment 90 is the polynucleotide of embodiment 89, wherein the PHI-4 polypeptide has one or more amino substitution compared to the native amino acid at position 332, 333, 334, 335, 336, 337, 338, 339, 340, 395, 396, 397, 398, 399, 400, 401, 402, 403, 458, 459, 460, 461, 462, 463, 464, 465, 466 of SEQ ID NO: 35.

15 Embodiment 91 is the polynucleotide of any one of embodiments 85, 86, 87, 88, 89 or 90, wherein the protease sensitive region residue is selected from between about the amino acid at position 305 (Lys) and about the amino acid at position 316 (Lys) and/or between about the amino acid at position 500 (Arg) and about the amino acid at position 535 (Lys) relative to SEQ ID NO: 35.

20 Embodiment 92 is the polynucleotide of any one of embodiments 85, 86, 87, 88, 89, 90 or 91, wherein the protease is trypsin.

Embodiment 93 is the polynucleotide of any one of embodiments 85-92, wherein the PHI-4 polypeptide has at least 80% sequence identity to SEQ ID NO: 35.

25 Embodiment 94 is the polynucleotide of any one of embodiments 85-92, wherein the PHI-4 polypeptide has at least 90% sequence identity to SEQ ID NO: 35.

Embodiment 95 is the polynucleotide of any one of embodiments 85-92, wherein the PHI-4 polypeptide has at least 95% sequence identity to SEQ ID NO: 35.

Embodiment 96 is the polynucleotide of any one of embodiments 85-92, wherein the PHI-4 polypeptide has at least 97% sequence identity to SEQ ID NO: 35.

30 Embodiment 97 is a polynucleotide encoding a PHI-4 polypeptide, wherein the PHI-4 polypeptide comprises an amino acid sequence of the formula,

	5	10	15
	Met Xaa Ser Ala Ala Asn Ala Gly Xaa	Leu Gly Asn Leu Xaa Gly	
	20	25	30
35	Xaa Thr Ser Xaa Gly Met Xaa Tyr Xaa	Val Asn Gly Leu Tyr Ala	
	35	40	45

WO 2014/150914

PCT/US2014/024524

	Ser	Pro	Glu	Ser	Leu	Xaa	Gly	Gln	Pro	Leu	Phe	Xaa	Xaa	Gly	Gly
					50					55					60
	Xaa	Leu	Asp	Ser	Xaa	Xaa	Ile	Glu	Gly	Xaa	Xaa	Xaa	Xaa	Phe	Pro
					65					70					75
5	Xaa	Ser	Met	His	Val	His	Thr	Tyr	Phe	His	Ser	Asp	Xaa	Xaa	Gln
					80					85					90
	Xaa	Val	Ser	Xaa	Xaa	Ile	Xaa	Xaa	Xaa	Arg	Xaa	Xaa	Xaa	Ser	Xaa
					95					100					105
	His	Val	Gly	Xaa	Ser	Gly	Xaa	Xaa	Xaa	Leu	Phe	Ser	Xaa	Ser	Xaa
10					110					115					120
	Ser	Val	Asp	Xaa	Thr	Thr	Xaa	Xaa	Gln	Gln	Leu	Xaa	Glu	Ile	Thr
					125					130					135
	Xaa	Ser	Ser	Thr	Arg	Glu	Xaa	His	Val	Leu	Trp	Tyr	Ile	Ser	Leu
					140					145					150
15	Pro	Gly	Ala	Ala	Thr	Leu	Xaa	Ser	Met	Leu	Xaa	Xaa	Xaa	Xaa	Xaa
					155					160					165
	Xaa	Asp	Xaa	Xaa	Xaa	Pro	Asn	Met	Xaa	Ala	Met	Xaa	Leu	Phe	Xaa
					170					175					180
	Xaa	Xaa	Gly	Pro	Xaa	Xaa	Xaa	Xaa	Xaa	Ala	Ala	Val	Gly	Gly	Arg
20					185					190					195
	Leu	Xaa	Xaa	Xaa	Xaa	Ala	Ser	Lys	Xaa	Leu	Xaa	Met	Xaa	Ser	Ser
					200					205					210
	Xaa	Ser	Leu	Ser	Thr	Thr	Xaa	Xaa	Xaa	Ser	Xaa	Xaa	Ala	Xaa	Xaa
					215					220					225
25	Gly	Glu	Ile	Xaa	Ile	Xaa	His	Gly	Ser	Xaa	Met	Glu	Lys	Gln	Val
					230					235					240
	Asn	Ser	Phe	Xaa	Xaa	Xaa	Ser	Thr	Ile	Arg	Xaa	Thr	Ala	Thr	Gly
					245					250					255
	Gly	Lys	Pro	Gly	Xaa	Thr	Xaa	Arg	Ile	Leu	His	Gly	Pro	Asp	Ser
30					260					265					270
	Xaa	Xaa	Ala	Phe	Ser	Xaa	Trp	Ala	Xaa	Ser	Leu	Leu	Xaa	Tyr	Ala
					275					280					285
	Thr	Leu	Met	Asp	Phe	Xaa	Thr	Xaa	Ser	Leu	Xaa	Xaa	Ile	Xaa	Ala
					290					295					300
35	Leu	Xaa	Asp	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Glu	Xaa	Xaa	Xaa	Ala	Xaa
					305					310					315
	Pro	Xaa	Xaa	Met	Xaa	Xaa	Ser	Gln	Xaa	Ser	Ile	Pro	Xaa	Val	Asp

		320		325		330									
	Xaa	Val	Leu	Leu	Met	Asp	Ala	Arg	Pro	Pro	Met	Val	Xaa	Ala	Gly
		335		340		345									
	Xaa	Asp	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Asp	Xaa	Xaa	Xaa	Xaa	Xaa
5		350		355		360									
	Xaa	Ser	Thr	Ser	Xaa	Xaa	Tyr	Lys	Xaa	Xaa	Gly	Gln	Phe	Xaa	Gln
		365		370		375									
	Arg	Xaa	His	Xaa	Ser	Val	Ala	Asp	Gly	His	Xaa	Pro	Ile	Xaa	Xaa
		380		385		390									
10	Asp	Leu	Phe	Asp	Xaa	Gly	Xaa	Xaa	Xaa	Xaa	Pro	Val	Gly	Xaa	Gln
		395		400		405									
	Xaa	Val	Trp	Asp	Xaa	Xaa	Xaa	Xaa	Gly	Lys	Xaa	Xaa	Xaa	Tyr	Xaa
		410		415		420									
	Cys	Trp	Arg	Xaa	Xaa	Xaa	Xaa	Gln	Gly	Tyr	Xaa	Xaa	Xaa	Gly	Asp
15		425		430		435									
	Val	Xaa	Met	Leu	Ala	Xaa	Ser	Gly	Tyr	Asn	Pro	Pro	Asn	Leu	Pro
		440		445		450									
	Xaa	Xaa	Xaa	Cys	Xaa	His	Xaa	Ser	Leu	Xaa	Ala	Xaa	Xaa	Xaa	Thr
		455		460		465									
20	Leu	Xaa	Xaa	Xaa	Xaa	Trp	Xaa	Asp	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
		470		475		480									
	Xaa	Val	Ser	Leu	Trp	Xaa	Xaa	Gly	Ala	Ala	Gly	Ala	Val	Ala	Ser
		485		490		495									
	Ser	Cys	Phe	Ala	Gly	Val	Pro	Asn	Tyr	Asn	Asn	Pro	Pro	Asn	Ser
25		500		505		510									
	Gly	Xaa	Ile	Xaa	Xaa	Leu	Xaa	Gly	Ser	Ile	Ala	Cys	Val	Xaa	Thr
		515		520		525									
	Ser	Ala	Ile	Ala	Ser	Met	Xaa	Xaa	Met	Xaa	Ser	Met	Leu	Ser	Xaa
		530		535											
30	His	Xaa	Gly	Met	Glu	Ala	Met	Met	Ser	Lys	Leu	(SEQ ID NO: 3),			

wherein

Xaa at position 2 is Ala or Arg; Xaa at position 9 is Gln, Lys or Glu; Xaa at position 14 is Pro or Ala; Xaa at position 16 is Val or Asp; Xaa at position 19 is Met or Leu; Xaa at position 22 is Gly or Ser; Xaa at position 24 is Asp, Asn or Gln; Xaa at position 36 is Leu or Met; Xaa at position 42 is Asp, Asn or Gln; Xaa at position 43 is Phe or Glu; Xaa at position 46 is Glu, Asp, Asn or Gly; Xaa at position 50 is Ile or Val; Xaa at position 51 is Glu or Gln; Xaa at position 55 is Arg or Lys; Xaa at position 56 is Ser or Thr; Xaa at

position 57 is Tyr or Phe; Xaa at position 58 is Thr or Ser; Xaa at position 61 is Arg, Lys or Glu; Xaa at position 73 is Phe or Tyr; Xaa at position 74 is Lys, Glu, Gly, Arg, Met, Leu, His or Asp; Xaa at position 76 is Asp or Gln; Xaa at position 79 is Lys or Glu; Xaa at position 80 is Glu or Ser; Xaa at position 82 is Glu, Ile, Leu, Tyr or Gln; Xaa at position 83
 5 is Glu or Gln; Xaa at position 84 is Tyr or Phe; Xaa at position 86 is Glu or Gln; Xaa at position 87 is Lys or Gln; Xaa at position 88 is Met, Ile or Leu; Xaa at position 90 is Gln or Glu; Xaa at position 94 is Val or Ile; Xaa at position 97 is Arg, Asn, Asp, Glu, Gln, Gly, Ser, Ile, Phe, His, Lys, Thr, Asn, Tyr, Trp, Pro, Cys, Ala, Met, Val or Leu; Xaa at position 98 is Tyr or Phe; Xaa at position 99 is Lys, Leu, Tyr, Ile, Met, Phe, Cys, Val or Asn; Xaa at
 10 position 103 is Ala or Gly; Xaa at position 105 is Leu or Ile; Xaa at position 109 is Phe, Lys, Gly, Met, Ser, Asp, Asn, Glu, Cys, Ala or Arg; Xaa at position 112 is Thr or Ser; Xaa at position 113 is Asp, Glu or Met; Xaa at position 117 is Thr or Ser; Xaa at position 121 is Tyr or Phe; Xaa at position 127 is Ala or Thr; Xaa at position 142 is Arg or Glu; Xaa at position 146 is Arg or Gln; Xaa at position 147 is Arg, Glu or Gln; Xaa at position 148 is
 15 Asp, Phe, Pro, Val, Glu, His, Trp, Ala, Arg, Leu, Ser, Gln or Gly; Xaa at position 149 is Phe or Val; Xaa at position 150 is Arg, Gln, Glu or Asn; Xaa at position 151 is Asp, Ser, Ala, Asn, Trp, Val, Gln, Cys, Met, Leu, Arg or Glu; Xaa at position 153 is Leu or Ile; Xaa at position 154 is Asn or Asp; Xaa at position 155 is Asn or Lys; Xaa at position 159 is Pro or Asp; Xaa at position 162 is Glu, Asp, Gln, Asn or Leu; Xaa at position 165 is Lys, Glu,
 20 Gln, Pro, Thr, Ala, Leu, Gly, Asp, Val, His, Ile, Met, Trp, Phe, Tyr or Arg; Xaa at position 166 is Arg or Gln; Xaa at position 167 is Tyr, Trp or Cys; Xaa at position 170 is Tyr or His; Xaa at position 171 is Tyr or Phe; Xaa at position 172 is Ile, Leu or Val; Xaa at position 173 is Ser or Ala; Xaa at position 174 is Glu, Gln, Asn, Lys, Val or Ser; Xaa at position 182 is Asp or Gln; Xaa at position 183 is Tyr or Val; Xaa at position 184 is Ser or Thr; Xaa
 25 at position 185 is Ala or Ser; Xaa at position 189 is Thr, Lys or Ile; Xaa at position 191 is Lys or Gln; Xaa at position 193 is Asp or Asn; Xaa at position 196 is Gln, Lys, Asn, Asp, Glu, Ala, Ile or Arg; Xaa at position 202 is Ala or Val; Xaa at position 203 is Glu, Thr or His; Xaa at position 204 is Met or Ala; Xaa at position 206 is Tyr or Phe; Xaa at position 207 is Lys or Gln; Xaa at position 209 is Leu or Pro; Xaa at position 210 is Val or Ile; Xaa
 30 at position 214 is Lys, Ser or Gln; Xaa at position 216 is Glu, Gln, Phe, Val, Tyr or Arg; Xaa at position 220 is Glu, His, Asp, Thr, Tyr, Val, Ser, Gln, Arg, Trp, Met, Ala, Phe, Ile, Leu, Cys or Asn; Xaa at position 229 is Arg or Glu; Xaa at position 230 is Ser or Glu; Xaa at position 231 is Asn or Ser; Xaa at position 236 is Leu or Pro; Xaa at position 245 is Met or Leu; Xaa at position 247 is Asp or Tyr; Xaa at position 256 is Gln, Lys or Glu; Xaa at
 35 position 257 is Gln, Ile, Glu, Cys, Ser, His, Trp or Met; Xaa at position 261 is Gln, Glu, Lys or Ala; Xaa at position 264 is Glu or Gln; Xaa at position 268 is Asp or Asn; Xaa at

position 276 is Ser or Ala; Xaa at position 278 is Glu, Asn or Gln; Xaa at position 281 is
 Gln, Lys or Glu; Xaa at position 282 is Pro or Gly; Xaa at position 284 is Trp or Arg; Xaa
 at position 287 is Ala or Cys; Xaa at position 289 is Lys, Leu, Val, Pro, Glu, Gln, Tyr, Thr,
 Asp, Phe, Ser, Met, Arg, Trp, Ile, His, Asn, Cys, Gly or Ala; Xaa at position 291 is Glu or
 5 Gln; Xaa at position 292 is Arg or Gln; Xaa at position 293 is Arg, Glu or Gln; Xaa at
 position 294 is Val or Ala; Xaa at position 296 is Leu or Ile; Xaa at position 297 is Glu or
 Gln; Xaa at position 298 is Asp or Gln; Xaa at position 300 is Phe or Tyr; Xaa at position
 302 is Glu or Gln; Xaa at position 303 is Phe or Tyr; Xaa at position 305 is Lys, Gln, Ala,
 Ile, Met, Asn, Thr or Val; Xaa at position 306 is Gln or Lys; Xaa at position 309 is Gln, Lys
 10 or Glu; Xaa at position 313 is Lys, Gln or Arg; Xaa at position 316 is Lys or Gln; Xaa at
 position 328 is Lys, Glu or Gln; Xaa at position 331 is Glu, Asn or Gln; Xaa at position 333
 is Ser, Arg, Gly, Lys, Val, Asn, Ala, His, Gln, Thr, Asp, Ile, Leu, Cys or Glu; Xaa at position
 334 is Gly, Arg, Lys, Ile or Trp; Xaa at position 335 is Ser or Ala; Xaa at position 336 is
 Gly or Ala; Xaa at position 337 is Ala, Val or Gly; Xaa at position 338 is Ser, His, Val, Lys,
 15 Ala, Gly, Thr, Ile, Glu, Met, Arg, Pro, Asp, Asn or Leu; Xaa at position 339 is Glu, Asn,
 Gln, Ile, Pro, Met, Ser, Ala, Cys, Phe, Val, Leu, Asp, Trp, His or Arg; Xaa at position 341
 is Leu or Val; Xaa at position 342 is Ala, Ser or Val; Xaa at position 343 is Val or Ile; Xaa
 at position 344 is Phe or Trp; Xaa at position 345 is Asn or His; Xaa at position 346 is Pro
 or Ala; Xaa at position 350 is Asn or Ser; Xaa at position 351 is Gly or Val; Xaa at position
 20 354 is Met or Leu; Xaa at position 355 is Val, Ile or Leu; Xaa at position 359 is Gly or Ala;
 Xaa at position 362 is Asn or Ser; Xaa at position 364 is Ala or Ser; Xaa at position 371 is
 Ala, Gly or Thr; Xaa at position 374 is Phe or Ile; Xaa at position 375 is Lys or Arg; Xaa at
 position 380 is Leu or Gly; Xaa at position 382 is Val, Asp or Leu; Xaa at position 383 is
 Leu, Ile or Val; Xaa at position 384 is Lys, Ala or Gly; Xaa at position 385 is Ala or Gly;
 25 Xaa at position 389 is Trp or Tyr; Xaa at position 391 is Arg, Leu, Glu, Gln, Asp or His;
 Xaa at position 395 is Asp or Cys; Xaa at position 396 is Ala, Leu, Lys, Asn, Gly, Ile, Met,
 Arg, Tyr, Gln, His or Thr; Xaa at position 397 is Gly, Arg or Ala; Xaa at position 398 is Ser,
 Gln or Cys; Xaa at position 401 is Ser, His, Pro, Gly, Lys, Val, Arg, Ile, Asn, Phe, Thr, Ala,
 Asp, Met, Gln or Glu; Xaa at position 402 is Lys, Phe, His, Arg, Trp, Gly, Asn, Leu, Tyr,
 30 Thr, Val, Met, Pro or Ala; Xaa at position 403 is Asp, Tyr, Trp, Phe or Glu; Xaa at position
 405 is Ala or Ser; Xaa at position 409 is Ala or Pro; Xaa at position 410 is Ile or Val; Xaa
 at position 411 is Pro or Ala; Xaa at position 412 is Pro or Ala; Xaa at position 416 is Arg,
 Glu or Gln; Xaa at position 417 is Ala, Ser or Cys; Xaa at position 418 is Leu or Met; Xaa
 at position 422 is Met or Val; Xaa at position 426 is Thr or Ser; Xaa at position 436 is Asp
 35 or Lys; Xaa at position 437 is Tyr or Val; Xaa at position 438 is Val or Arg; Xaa at position
 440 is Val or Leu; Xaa at position 442 is Gln, Lys or Glu; Xaa at position 445 is Cys, Leu

or Thr; Xaa at position 447 is Asp, Lys, Tyr, Ser, Glu, Ile, Gly, Pro, Leu, Phe, Trp or Thr; Xaa at position 448 is Val or Ala; Xaa at position 449 is Gln or Glu; Xaa at position 452 is Gln, Lys or Glu, Ala; Xaa at position 453 is Asn or Asp; Xaa at position 454 is Arg, Tyr, Met, Ser, Val, Ile, Lys, Phe, Trp, Gln, Gly, His, Asp, Leu, Thr, Pro or Asn; Xaa at position 5 455 is Val or Ile; Xaa at position 457 is Trp or Asn; Xaa at position 459 is Lys, Met, Val, Trp, Gln, Ile, Thr, Ser, His, Cys, Tyr, Pro, Asn, Ala, Arg or Glu; Xaa at position 460 is Gly or Ala; Xaa at position 461 is Thr or Ser; Xaa at position 462 is Gly or Ala; Xaa at position 463 is Ala, Ser or Gly; Xaa at position 464 is Arg, Gly, His, Gln, Thr, Phe, Ala, Asp, Ser or Lys; Xaa at position 465 is Lys, Asn, Val, Met, Pro, Gly, Arg, Thr, His, Cys, Trp, Phe or 10 Leu; Xaa at position 466 is Asp or Arg; Xaa at position 471 is Gln, Lys, Glu or Met; Xaa at position 472 is Pro or Ser; Xaa at position 497 is Asp or Gln; Xaa at position 499 is Glu or Gln; Xaa at position 500 is Arg, Gln or Lys; Xaa at position 502 is Arg, Glu or Gln; Xaa at position 509 is Lys, Gln, Glu or Ala; Xaa at position 517 is Gln, Cys, Asn, Val or Pro; Xaa at position 518 is Glu or Gln; Xaa at position 520 is Lys, Gln, Glu, His or Ala; Xaa at 15 position 525 is Gln or Lys; and Xaa at position 527 is Gln, Lys, Pro, Cys, Glu, Ser, His, Phe or Trp; wherein one or more amino acid(s) designated by Xaa in SEQ ID NO: 3 is an amino acid different from the corresponding amino acid of SEQ ID NO: 35 and wherein the PHI-4 polypeptide has increased insecticidal activity compared to the polypeptide of SEQ ID NO: 35.

Embodiment 98 is a polynucleotide encoding a PHI-4 polypeptide, wherein the PHI-4 polypeptide comprises an amino acid sequence of the formula,

	5	10	15
	Met Xaa Ser Ala Ala Asn Ala Gly Gln	Leu Gly Asn Leu Pro Gly	
	20	25	30
25	Val Thr Ser Met Gly Met Gly Tyr Xaa	Val Asn Gly Leu Tyr Ala	
	35	40	45
	Ser Pro Glu Ser Leu Leu Gly Gln Pro	Leu Phe Xaa Xaa Gly Gly	
	50	55	60
	Xaa Leu Asp Ser Ile Glu Ile Glu Gly	Arg Ser Tyr Thr Phe Pro	
30	65	70	75
	Arg Ser Met His Val His Thr Tyr Phe	His Ser Asp Phe Xaa Gln	
	80	85	90
	Asp Val Ser Xaa Glu Ile Xaa Glu Tyr	Arg Glu Lys Met Ser Gln	
	95	100	105
35	His Val Gly Val Ser Gly Xaa Xaa Xaa	Leu Phe Ser Ala Ser Leu	
	110	115	120
	Ser Val Asp Xaa Thr Thr Thr Asp Gln	Gln Leu Thr Glu Ile Thr	

WO 2014/150914

PCT/US2014/024524

					125					130					135
	Tyr	Ser	Ser	Thr	Arg	Glu	Ala	His	Val	Leu	Trp	Tyr	Ile	Ser	Leu
					140					145					150
	Pro	Gly	Ala	Ala	Thr	Leu	Arg	Ser	Met	Leu	Arg	Xaa	Xaa	Phe	Xaa
5					155					160					165
	Xaa	Asp	Xaa	Asn	Asn	Pro	Asn	Met	Pro	Ala	Met	Xaa	Leu	Phe	Xaa
					170					175					180
	Xaa	Tyr	Gly	Pro	Tyr	Xaa	Ile	Ser	Xaa	Ala	Ala	Val	Gly	Gly	Arg
					185					190					195
10	Leu	Xaa	Tyr	Ser	Ala	Ala	Ser	Lys	Thr	Leu	Lys	Met	Asp	Ser	Ser
					200					205					210
	Xaa	Ser	Leu	Ser	Thr	Thr	Ala	Xaa	Met	Ser	Xaa	Lys	Ala	Leu	Val
					215					220					225
	Gly	Glu	Ile	Lys	Ile	Xaa	His	Gly	Ser	Xaa	Met	Glu	Lys	Gln	Val
15					230					235					240
	Asn	Ser	Phe	Arg	Ser	Asn	Ser	Thr	Ile	Arg	Leu	Thr	Ala	Thr	Gly
					245					250					255
	Gly	Lys	Pro	Gly	Met	Thr	Xaa	Arg	Ile	Leu	His	Gly	Pro	Asp	Ser
					260					265					270
20	Xaa	Xaa	Ala	Phe	Ser	Xaa	Trp	Ala	Glu	Ser	Leu	Leu	Asp	Tyr	Ala
					275					280					285
	Thr	Leu	Met	Asp	Phe	Ser	Thr	Xaa	Ser	Leu	Xaa	Pro	Ile	Trp	Ala
					290					295					300
	Leu	Ala	Asp	Xaa	Pro	Glu	Arg	Xaa	Val	Glu	Leu	Glu	Asp	Ala	Phe
25					305					310					315
	Pro	Glu	Phe	Met	Lys	Gln	Ser	Gln	Gln	Ser	Ile	Pro	Xaa	Val	Asp
					320					325					330
	Lys	Val	Leu	Leu	Met	Asp	Ala	Arg	Pro	Pro	Met	Val	Xaa	Ala	Gly
					335					340					345
30	Glu	Asp	Xaa	Xaa	Ser	Xaa	Ala	Xaa	Xaa	Asp	Leu	Ala	Xaa	Phe	Asn
					350					355					360
	Xaa	Ser	Thr	Ser	Asn	Gly	Tyr	Lys	Met	Xaa	Gly	Gln	Phe	Xaa	Gln
					365					370					375
	Arg	Asn	His	Ala	Ser	Val	Ala	Asp	Gly	His	Ala	Pro	Ile	Phe	Lys
35					380					385					390
	Asp	Leu	Phe	Asp	Leu	Gly	Val	Leu	Lys	Ala	Pro	Val	Gly	Trp	Gln
					395					400					405

Xaa Val Trp Asp Asp Xaa Gly Ser Gly Lys Xaa Xaa Xaa Tyr Ala
 410 415 420
 Cys Trp Arg Ala Ile Xaa Xaa Gln Gly Tyr Xaa Xaa Xaa Gly Asp
 425 430 435
 5 Val Met Met Leu Ala Xaa Ser Gly Tyr Asn Pro Pro Asn Leu Pro
 440 445 450
 Asp Tyr Val Cys Xaa His Gln Ser Leu Cys Ala Xaa Val Gln Thr
 455 460 465
 Leu Xaa Asn Xaa Xaa Trp Trp Asp Xaa Gly Xaa Xaa Xaa Xaa Xaa
 10 470 475 480
 Asp Val Ser Leu Trp Xaa Xaa Gly Ala Ala Gly Ala Val Ala Ser
 485 490 495
 Ser Cys Phe Ala Gly Val Pro Asn Tyr Asn Asn Pro Pro Asn Ser
 500 505 510
 15 Gly Asp Ile Glu Xaa Leu Arg Gly Ser Ile Ala Cys Val Xaa Thr
 515 520 525
 Ser Ala Ile Ala Ser Met Gln Glu Met Xaa Ser Met Leu Ser Gln
 530 535
 His Xaa Gly Met Glu Ala Met Met Ser Lys Leu (SEQ ID NO: 4),

20 wherein

Xaa at position 2 is Ala or Arg; Xaa at position 24 is Asp or Asn; Xaa at position 42 is Asp
 or Asn; Xaa at position 43 is Phe or Glu; Xaa at position 46 is Glu or Asn; Xaa at position
 74 is Lys, Glu or Gly; Xaa at position 79 is Lys or Glu; Xaa at position 82 is Glu, Ile, Leu or
 Tyr; Xaa at position 97 is Arg, Asn, Asp, Glu, Gln, Gly, Ser, Ile, Phe, His, Lys, Thr, Asn,
 25 Tyr, Trp, Pro, Cys, Ala, Met, Val or Leu; Xaa at position 98 is Tyr or Phe; Xaa at position
 99 is Lys, Leu, Tyr, Ile or Met; Xaa at position 109 is Phe, Lys, Gly, Met, Ser, Asp or Asn;
 Xaa at position 147 is Arg or Glu; Xaa at position 148 is Asp, Phe or Pro; Xaa at position
 150 is Arg, Gln, Glu or Asn; Xaa at position 151 is Asp, Ser, Ala or Asn; Xaa at position
 153 is Leu or Ile; Xaa at position 162 is Glu, Asp, Gln, Asn or Leu; Xaa at position 165 is
 30 Lys, Glu or Gln; Xaa at position 166 is Arg or Gln; Xaa at position 171 is Tyr or Phe; Xaa
 at position 174 is Glu, Gln, Asn, Lys, Val or Ser; Xaa at position 182 is Asp or Gln; Xaa at
 position 196 is Gln, Lys, Asn or Asp; Xaa at position 203 is Glu, Thr or His; Xaa at position
 206 is Tyr or Phe; Xaa at position 216 is Glu or Gln; Xaa at position 220 is Glu, His, Asp,
 Thr, Tyr, Val, Ser or Gln; Xaa at position 247 is Asp or Tyr; Xaa at position 256 is Gln or
 35 Lys; Xaa at position 257 is Gln or Ile; Xaa at position 261 is Gln, Glu, Lys or Ala; Xaa at
 position 278 is Glu or Asn; Xaa at position 281 is Gln, Lys or Glu; Xaa at position 289 is
 Lys, Leu, Val, Pro, Glu, Gln, Tyr, Thr or Asp; Xaa at position 293 is Arg, Glu or Gln; Xaa at

position 313 is Lys or Gln; Xaa at position 328 is Lys, Glu or Gln; Xaa at position 333 is Ser, Gly, Lys, Val or Asn; Xaa at position 334 is Gly, Arg, Lys or Ile; Xaa at position 336 is Gly or Ala; Xaa at position 338 is Ser, His, Val, Lys or Ala; Xaa at position 339 is Glu, Asn, Ile or Pro; Xaa at position 343 is Val or Ile; Xaa at position 346 is Pro or Ala; Xaa at position 355 is Val or Ile; Xaa at position 359 is Gly or Ala; Xaa at position 391 is Arg, Leu, Glu, Gln, Asp or His; Xaa at position 396 is Ala, Leu, Lys, Asn, Gly or Thr; Xaa at position 401 is Ser, His, Pro, Gly, Lys, Val or Arg; Xaa at position 402 is Lys, Phe, His, Arg, Gly, Trp, Thr, Asn, Tyr or Met; Xaa at position 403 is Asp or Tyr; Xaa at position 411 is Pro or Ala; Xaa at position 412 is Pro or Ala; Xaa at position 416 is Arg or Glu; Xaa at position 417 is Ala or Ser; Xaa at position 418 is Leu or Met; Xaa at position 426 is Thr or Ser; Xaa at position 440 is Val or Leu; Xaa at position 447 is Asp, Lys, Tyr, Ser, Glu or Ile; Xaa at position 452 is Gln, Lys or Glu; Xaa at position 454 is Arg, Tyr, Met, Ser, Val, Ile, Lys, Phe, Trp or Gln; Xaa at position 455 is Val or Ile; Xaa at position 459 is Lys, Met, Val, Trp, Gln, Ile or Tyr; Xaa at position 461 is Thr or Ser; Xaa at position 462 is Gly or Ala; Xaa at position 463 is Ala or Ser; Xaa at position 464 is Arg, Gly, His, Ala, Asp, Ser or Lys; Xaa at position 465 is Lys, Asn, Val, Met, Pro, Gly or Arg; Xaa at position 471 is Gln, Lys, Glu or Met; Xaa at position 472 is Pro or Ser; Xaa at position 500 is Arg or Gln; Xaa at position 509 is Lys, Gln or Ala; Xaa at position 520 is Lys, Gln, Glu, His or Ala; and Xaa at position 527 is Gln, Lys, Pro, Cys or Glu; wherein one or more amino acid(s) designated by Xaa in SEQ ID NO: 4 is an amino acid different from the corresponding amino acid of SEQ ID NO: 35 and wherein the PHI-4 polypeptide has increased insecticidal activity compared to the polypeptide of SEQ ID NO: 35.

Embodiment 99 is the polynucleotide of embodiments 97 or 98, wherein the PHI-4 polypeptide further comprises one or more amino acid substitutions at position 86, 359, 464, 465, 466, 467, 468, 499 or 517 of SEQ ID NO: 3 or SEQ ID NO: 4.

Embodiment 100 is the polynucleotide of embodiment 99, wherein the amino acid at position 86 is Glu or Thr; the amino acid at position 359 is Gly or Ala; the amino acid at position 464 is Arg, Ala, Lys, Asp or Asn; the amino acid at position 465 is Lys or Met, the amino acid at position 467 is Val, Ala, Leu or Thr; the amino acid at position 468 is Ser or Leu; the amino acid at position 499 is Glu or Ala or the amino acid at position 517 is Glu or Arg.

Embodiment 101 is the polynucleotide of any one of embodiments 97-100, wherein the PHI-4 polypeptide further comprises one or more conservative amino acid substitution, insertion of one or more amino acids, deletion of one or more amino acids, and combinations thereof.

Embodiment 102 is the polynucleotide of any one of embodiments 97-101, wherein the insecticidal activity of the PHI-4 polypeptide is increased about 1.5 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

5 Embodiment 103 is the polynucleotide of any one of embodiments 97-101, wherein the insecticidal activity of the PHI-4 polypeptide is increased about 2 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 104 is the polynucleotide of any one of embodiments 97-101, wherein the insecticidal activity of the PHI-4 polypeptide is increased about 2.5 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

10 Embodiment 105 is the polynucleotide of any one of embodiments 97-101, wherein the insecticidal activity of the PHI-4 polypeptide is increased about 3 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 106 is the polynucleotide of any one of embodiments 97-101, wherein the insecticidal activity of the PHI-4 polypeptide is increased about 5 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

Embodiment 107 is the polynucleotide of any one of embodiments 97-106, wherein the improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35) is against Western Corn Root Worm (WCRW) larvae.

20 Embodiment 108 is the polynucleotide of any one of embodiments 97-107, wherein the improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35) is quantified as a Mean FAE Index.

Embodiment 109 is the polynucleotide of any one of embodiments 97-107, wherein the improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35) is quantified as an EC50 value.

25 Embodiment 110 is the polynucleotide of any one of embodiments 97-107, wherein the improved activity compared to AXMI-205 (SEQ ID NO: 35) is quantified as a Mean Deviation Score.

Embodiment 111 is the polynucleotide of any one of embodiments 97-110, wherein the PHI-4 polypeptide has 1 to 54 amino acid substitutions at a position(s) designated as Xaa in SEQ ID NO: 3 or SEQ ID NO: 4.

Embodiment 112 is the polynucleotide of any one of embodiments 97-110, wherein the PHI-4 polypeptide has 1 to 27 amino acid substitutions at a position(s) designated as Xaa in SEQ ID NO: 3 or SEQ ID NO: 4.

35 Embodiment 113 is the polynucleotide of any one of embodiments 97-110, wherein the PHI-4 polypeptide has 1 to 20 amino acid substitutions at a position(s) designated as Xaa in SEQ ID NO: 3 or SEQ ID NO: 4.

Embodiment 114 is the polynucleotide of any one of embodiments 97-110, wherein the PHI-4 polypeptide has 1 to 15 amino acid substitutions at a position(s) designated as Xaa in SEQ ID NO: 3 or SEQ ID NO: 4.

Embodiment 115 is the polynucleotide of any one of embodiments 97-114, wherein 1-25 amino acids are deleted from the N-terminus of the PHI-4 polypeptide and/or C-terminus of the PHI-4 polypeptide.

Embodiment 116 is the polynucleotide of any one of embodiments 97-114, wherein 1-20 amino acids are deleted from the C-terminus of the PHI-4 polypeptide.

Embodiment 117 is a composition, comprising an insecticidally-effective amount of the PHI-4 polypeptide of any one of embodiments 1-58.

Embodiment 118 is a method of inhibiting growth or killing an insect pest, comprising contacting the insect pest with the composition of embodiment 117.

Embodiment 119 is a method for controlling an insect pest population resistant to a pesticidal protein, comprising contacting the resistant insect pest population with the composition of embodiment 117.

Embodiment 120 is the method of controlling an insect pest population resistant to an pesticidal protein of embodiment 119, wherein the pesticidal protein is selected from Cry1Ac, Cry1Ab, Cry1A.105, Cry1Ac, Cry1F, Cry1Fa2, Cry1F, Cry2Ab, Cry3A, mCry3A, Cry3Bb1, Cry34Ab1, Cry35Ab1, Vip3A, Cry9c, eCry3.1Ab and CBI-Bt.

Embodiment 121 is a transgenic plant or progeny thereof, comprising the polynucleotide of any one of embodiments 59-116. Embodiment 122 is the transgenic plant or progeny thereof of embodiment 121, wherein the transgenic plant is a monocotyledon.

Embodiment 123 is the transgenic plant or progeny thereof of embodiment 122, wherein the plant is selected from barley, corn, oat, rice, rye, sorghum, turf grass, sugarcane, wheat, alfalfa, banana, broccoli, bean, cabbage, canola, carrot, cassava, cauliflower, celery, citrus, cotton, a cucurbit, eucalyptus, flax, garlic, grape, onion, lettuce, pea, peanut, pepper, potato, poplar, pine, sunflower, safflower, soybean, strawberry, sugar beet, sweet potato, tobacco, tomato ornamental, shrub, nut, chickpea, pigeon pea, millets, hops and pasture grasses.

Embodiment 124 is the transgenic plant or progeny thereof of embodiment 123, further comprising one or more additional transgenic traits.

Embodiment 125 is the transgenic plant of embodiment 124, wherein the one or more additional transgenic trait is selected from insect resistance, herbicide resistance, fungal resistance, viral resistance, stress tolerance, disease resistance, male sterility, stalk strength, increased yield, modified starches, improved oil profile, balanced amino

acids, high lysine or methionine, increased digestibility, improved fiber quality, flowering, ear and seed development, enhancement of nitrogen utilization efficiency, altered nitrogen responsiveness, drought resistance or tolerance, cold resistance or tolerance, salt resistance or tolerance and increased yield under stress.

5 Embodiment 126 is seed, grain or processed product thereof of the transgenic plant of any one of embodiments 121-126, wherein the seed, grain or processed product thereof comprises the polynucleotide of any one of embodiments 121-125.

 Embodiment 127 is an expression cassette, comprising the polynucleotide of any one of embodiments 59-116 operably linked to one or more regulatory sequences
10 directing expression of the PHI-4 polypeptide.

 Embodiment 128 is a transgenic plant or plant cell, comprising the expression cassette of embodiment 127.

 Embodiment 129 is seed, grain or processed product thereof of the transgenic plant of embodiment 128, wherein the seed, grain or processed product thereof
15 comprises the recombinant nucleic acid molecule of embodiment of 1 and the recombinant nucleic acid molecule of 24.

 Embodiment 130 is the seed of embodiment 129, wherein one or more seed treatment has been applied to the seed.

 Embodiment 131 is a method for expressing in a plant a polynucleotide encoding
20 an insecticidal protein, comprising

 (a) inserting into a plant cell the polynucleotide of any one of embodiment 59-116 encoding the PHI-4 polypeptide;

 (b) obtaining a transformed plant cell comprising the nucleic acid sequence of step (a); and

25 (c) generating from the transformed plant cell a plant capable of expressing the polynucleotide encoding the PHI-4 polypeptide.

 Embodiment 132 is a method for protecting a plant from an insect pest, comprising expressing in the plant or cell thereof, an insecticidally-effective amount of the PHI-4 polypeptide of any one of embodiments 1-58.

30 Embodiment 133 is a method for controlling an insect pest population, comprising contacting the insect pest population with an insecticidally-effective amount of the PHI-4 polypeptide of any one of embodiments 1-58.

 Embodiment 134 is a method of inhibiting growth or killing an insect pest, comprising contacting the insect pest with a composition comprising an insecticidally-
35 effective amount of the the PHI-4 polypeptide of any one of embodiments 1-58.

Embodiment 135 is a method for controlling an insect pest population resistant to a pesticidal protein, comprising contacting the insect pest population with an insecticidally-effective amount of the PHI-4 polypeptide of any one of embodiments 1-58.

Embodiment 136 is a fusion protein comprising the PHI-4 polypeptide of any one of embodiments 1-58.

Embodiment 137 is a fusion protein comprising the PHI-4 polypeptide of any one of embodiments 1-58 and a maltose binding protein.

Embodiment 138 is the fusion protein of embodiment 137, wherein the maltose binding protein has an amino acid sequence of SEQ ID NO: 1516 or SEQ ID NO: 1517.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows FAE analysis of MPB::PHI-4-SFR12-004 (SEQ ID NO: 31). The reference protein is MBP::PHI-4 polypeptide of (SEQ ID NO: 6). The % Response is given on the y axis. The dose of the toxin fragment is given on the x axis in parts per million (ppm). Half dose (+) and full dose (*) indicate a mean response from six replicate wells at the indicated concentration. All data are from a single experiment.

Figure 2 shows EC50 analysis of MBP::PHI-4 polypeptide of SEQ ID NO: 6 and MPB::PHI-4-SFR12-004 (SEQ ID NO: 35). The % Response is given on the y axis. The dose of the toxin fragment is given on the x axis in parts per million (ppm). The % Response is given on the y axis. The dose of the toxin fragment is given on the x axis in parts per million (ppm). The protein concentration (toxin portion of the protein only) is given on the x axis. Each symbol indicates a mean response from twenty-four replicate wells at the indicated concentration. All data are from a single experiment. triangles: MBP::PHI-4 (SEQ ID NO: 6); circles: MBP::PHI-4-SFR12-004 (SEQ ID NO: 31).

Figure 3 shows the amino acids sequence of the C-terminal portion of the PHI-4 polypeptide (SEQ ID NO: 2) is given. Three stretches of sequence corresponding to nine amino acid motifs that align to the putative sugar binding loop motif D-X-G-(S/T)-G-X₃-D (SEQ ID NO: 40) are highlighted in grey.

Figure 4 shows EC50 data for SEQ ID NO: 610, SEQ ID NO: 595, SEQ ID NO: 584, SEQ ID NO: 591, SEQ ID NO: 576, SEQ ID NO: 73, SEQ ID NO: 74, SEQ ID NO: 75, SEQ ID NO: 79, SEQ ID NO: 81, SEQ ID NO: 150, SEQ ID NO: 150, SEQ ID NO: 149, SEQ ID NO: 167, SEQ ID NO: 167, SEQ ID NO: 164, SEQ ID NO: 164, SEQ ID NO: 170, SEQ ID

NO: 170, SEQ ID NO: 795, SEQ ID NO: 794, SEQ ID NO: 784, SEQ ID NO: 799, SEQ ID NO: 785, SEQ ID NO: 788, SEQ ID NO: 786, SEQ ID NO: 796, SEQ ID NO: 787. The x-axis corresponds to the mean fold improvement in EC50, relative to MBP::PHI-4 fusion (SEQ ID NO: 6) from at least three independent EC50 measurements. The y-axis corresponds to the mean fold improvement of the mean FAE Index from at least three independent measurements.

Figure 5. T0 seedlings in the V3 – V4 growth stage were challenged as described (Oleson J. et al *J. Economic Entomology* **98**:1-8; 2005) and root nodal injury scores were recorded. The groups indicated along the x axis were either controls or were transformed with a plant vector of Example 18: 1 - untransformed control; 2 - positive control transgenic expressing positive control protein; 3 - AXMI-205 polypeptide (SEQ ID NO: 22); 4 - PHI-4-B09 polypeptide (SEQ ID NO: 23); 5 - PHI-4-D09 polypeptide (SEQ ID NO: 24); 6 - PHI-4-H08 polypeptide (SEQ ID NO: 25).

DETAILED DESCRIPTION

It is to be understood that this disclosure is not limited to the particular methodology, protocols, cell lines, genera, and reagents described, as such may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present disclosure.

As used herein the singular forms "a", "and", and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a cell" includes a plurality of such cells and reference to "the protein" includes reference to one or more proteins and equivalents thereof known to those skilled in the art, and so forth. All technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this disclosure belongs unless clearly indicated otherwise.

The present disclosure is drawn to compositions and methods for controlling pests. The methods involve transforming organisms with a nucleic acid sequence encoding a PHI-4 polypeptide. In particular, the nucleic acid sequences of the embodiments are useful for preparing plants and microorganisms possessing pesticidal activity. Thus, transformed bacteria, plants, plant cells, plant tissues and seeds are provided. Compositions are pesticidal nucleic acids and proteins of bacterial species. The nucleic acid sequences find use in the construction of expression vectors for subsequent transformation into organisms of interest, as probes for the isolation of other homologous (or partially homologous) genes, and for the generation of altered PHI-4

polypeptides by methods known in the art, such as site directed mutagenesis, domain swapping or DNA shuffling. The PHI-4 polypeptides find use in controlling, inhibiting growth or killing Lepidopteran, Coleopteran, Dipteran, fungal, Hemipteran, and nematode pest populations and for producing compositions with pesticidal activity. Insect pests of interest include, but are not limited to, the superfamily of stink bugs and other related insects including, but not limited to, species belonging to the family *Pentatomidae* (*Nezara viridula*, *Halyomorpha halys*, *Piezodorus guildini*, *Euschistus servus*, *Acrosternum hilare*, *Euschistus heros*, *Euschistus tristigma*, *Acrosternum hilare*, *Dichelops furcatus*, *Dichelops melacanthus*, and *Bagrada hilaris* (Bagrada Bug)), the family *Plataspidae* (*Megacopta cribraria* - Bean plataspid), and the family *Cydnidae* (*Scaptocoris castanea* - Root stink bug) and Lepidoptera species including but not limited to: diamond-back moth, e.g., *Helicoverpa zea* Boddie; soybean looper, e.g., *Pseudoplusia includens* Walker and velvet bean caterpillar e.g., *Anticarsia gemmatilis* Hübner.

By “pesticidal toxin” or “pesticidal protein” is intended a toxin that has toxic activity against one or more pests, including, but not limited to, members of the Lepidoptera, Diptera, Hemiptera and Coleoptera orders or the Nematoda phylum or a protein that has homology to such a protein. Pesticidal proteins have been isolated from organisms including, for example, *Bacillus* sp., *Pseudomonas* sp., *Photorhabdus* sp., *Xenorhabdus* sp., *Clostridium bifermentans* and *Paenibacillus popilliae*. Pesticidal proteins include but are not limited to: insecticidal proteins from *Pseudomonas* sp. such as PSEEN3174 (Monalysin, (2011) *PLoS Pathogens*, 7:1-13) from *Pseudomonas protegens* strain CHA0 and Pf-5 (previously *fluorescens*) (Pechy-Tarr, (2008) *Environmental Microbiology* 10:2368-2386; GenBank Accession No. EU400157); from *Pseudomonas Taiwanensis* (Liu, et al., (2010) *J. Agric. Food Chem.* 58:12343-12349) and from *Pseudomonas pseudoalcaligenes* (Zhang, et al., (2009) *Annals of Microbiology* 59:45-50 and Li, et al., (2007) *Plant Cell Tiss. Organ Cult.* 89:159-168); insecticidal proteins from *Photorhabdus* sp. and *Xenorhabdus* sp. (Hinchliffe, et al., (2010) *The Open Toxinology Journal* 3:101-118 and Morgan, et al., (2001) *Applied and Envir. Micro.* 67:2062-2069), US Patent Number 6,048,838, and US Patent Number 6,379,946; and δ -endotoxins including, but not limited to, the cry1, cry2, cry3, cry4, cry5, cry6, cry7, cry8, cry9, cry10, cry11, cry12, cry13, cry14, cry15, cry16, cry17, cry18, cry19, cry20, cry21, cry22, cry23, cry24, cry25, cry26, cry27, cry 28, cry 29, cry 30, cry31, cry32, cry33, cry34, cry35, cry36, cry37, cry38, cry39, cry40, cry41, cry42, cry43, cry44, cry45, cry 46, cry47, cry49, cry 51, cry 52, cry 53, cry54, cry55, cry56, cry57, cry58, cry59, cry60, cry61, cry62, cry63, cry64, cry65, cry66, cry67, cry68, cry69, cry70, cry71, cry72, and cry73 classes of δ -endotoxin genes and the *B. thuringiensis* cytolytic cyt1 and cyt2 genes. Members of these classes of *B.*

thuringiensis insecticidal proteins include, but are not limited to Cry1Aa1 (Accession # AAA22353); Cry1Aa2 (Accession # Accession # AAA22552); Cry1Aa3 (Accession # BAA00257); Cry1Aa4 (Accession # CAA31886); Cry1Aa5 (Accession # BAA04468); Cry1Aa6 (Accession # AAA86265); Cry1Aa7 (Accession # AAD46139); Cry1Aa8
5 (Accession # I26149); Cry1Aa9 (Accession # BAA77213); Cry1Aa10 (Accession # AAD55382); Cry1Aa11 (Accession # CAA70856); Cry1Aa12 (Accession # AAP80146); Cry1Aa13 (Accession # AAM44305); Cry1Aa14 (Accession # AAP40639); Cry1Aa15 (Accession # AAY66993); Cry1Aa16 (Accession # HQ439776); Cry1Aa17 (Accession # HQ439788); Cry1Aa18 (Accession # HQ439790); Cry1Aa19 (Accession # HQ685121);
10 Cry1Aa20 (Accession # JF340156); Cry1Aa21 (Accession # JN651496); Cry1Aa22 (Accession # KC158223); Cry1Ab1 (Accession # AAA22330); Cry1Ab2 (Accession # AAA22613); Cry1Ab3 (Accession # AAA22561); Cry1Ab4 (Accession # BAA00071); Cry1Ab5 (Accession # CAA28405); Cry1Ab6 (Accession # AAA22420); Cry1Ab7 (Accession # CAA31620); Cry1Ab8 (Accession # AAA22551); Cry1Ab9 (Accession #
15 CAA38701); Cry1Ab10 (Accession # A29125); Cry1Ab11 (Accession # I12419); Cry1Ab12 (Accession # AAC64003); Cry1Ab13 (Accession # AAN76494); Cry1Ab14 (Accession # AAG16877); Cry1Ab15 (Accession # AAO13302); Cry1Ab16 (Accession # AAK55546); Cry1Ab17 (Accession # AAT46415); Cry1Ab18 (Accession # AAQ88259); Cry1Ab19 (Accession # AAW31761); Cry1Ab20 (Accession # ABB72460); Cry1Ab21
20 (Accession # ABS18384); Cry1Ab22 (Accession # ABW87320); Cry1Ab23 (Accession # HQ439777); Cry1Ab24 (Accession # HQ439778); Cry1Ab25 (Accession # HQ685122); Cry1Ab26 (Accession # HQ847729); Cry1Ab27 (Accession # JN135249); Cry1Ab28 (Accession # JN135250); Cry1Ab29 (Accession # JN135251); Cry1Ab30 (Accession # JN135252); Cry1Ab31 (Accession # JN135253); Cry1Ab32 (Accession # JN135254);
25 Cry1Ab33 (Accession # AAS93798); Cry1Ab34 (Accession # KC156668); Cry1Ab-like (Accession # AAK14336); Cry1Ab-like (Accession # AAK14337); Cry1Ab-like (Accession # AAK14338); Cry1Ab-like (Accession # ABG88858); Cry1Ac1 (Accession # AAA22331); Cry1Ac2 (Accession # AAA22338); Cry1Ac3 (Accession # CAA38098); Cry1Ac4 (Accession # AAA73077); Cry1Ac5 (Accession # AAA22339); Cry1Ac6 (Accession #
30 AAA86266); Cry1Ac7 (Accession # AAB46989); Cry1Ac8 (Accession # AAC44841); Cry1Ac9 (Accession # AAB49768); Cry1Ac10 (Accession # CAA05505); Cry1Ac11 (Accession # CAA10270); Cry1Ac12 (Accession # I12418); Cry1Ac13 (Accession # AAD38701); Cry1Ac14 (Accession # AAQ06607); Cry1Ac15 (Accession # AAN07788); Cry1Ac16 (Accession # AAU87037); Cry1Ac17 (Accession # AAX18704); Cry1Ac18
35 (Accession # AAY88347); Cry1Ac19 (Accession # ABD37053); Cry1Ac20 (Accession # ABB89046); Cry1Ac21 (Accession # AAY66992); Cry1Ac22 (Accession # ABZ01836);

Cry1Ac23 (Accession # CAQ30431); Cry1Ac24 (Accession # ABL01535); Cry1Ac25
 (Accession # FJ513324); Cry1Ac26 (Accession # FJ617446); Cry1Ac27 (Accession #
 FJ617447); Cry1Ac28 (Accession # ACM90319); Cry1Ac29 (Accession # DQ438941);
 Cry1Ac30 (Accession # GQ227507); Cry1Ac31 (Accession # GU446674); Cry1Ac32
 5 (Accession # HM061081); Cry1Ac33 (Accession # GQ866913); Cry1Ac34 (Accession #
 HQ230364); Cry1Ac35 (Accession # JF340157); Cry1Ac36 (Accession # JN387137);
 Cry1Ac37 (Accession # JQ317685); Cry1Ad1 (Accession # AAA22340); Cry1Ad2
 (Accession # CAA01880); Cry1Ae1 (Accession # AAA22410); Cry1Af1 (Accession #
 AAB82749); Cry1Ag1 (Accession # AAD46137); Cry1Ah1 (Accession # AAQ14326);
 10 Cry1Ah2 (Accession # ABB76664); Cry1Ah3 (Accession # HQ439779); Cry1Ai1
 (Accession # AAO39719); Cry1Ai2 (Accession # HQ439780); Cry1A-like (Accession #
 AAK14339); Cry1Ba1 (Accession # CAA29898); Cry1Ba2 (Accession # CAA65003);
 Cry1Ba3 (Accession # AAK63251); Cry1Ba4 (Accession # AAK51084); Cry1Ba5
 (Accession # ABO20894); Cry1Ba6 (Accession # ABL60921); Cry1Ba7 (Accession #
 15 HQ439781); Cry1Bb1 (Accession # AAA22344); Cry1Bb2 (Accession # HQ439782);
 Cry1Bc1 (Accession # CAA86568); Cry1Bd1 (Accession # AAD10292); Cry1Bd2
 (Accession # AAM93496); Cry1Be1 (Accession # AAC32850); Cry1Be2 (Accession #
 AAQ52387); Cry1Be3 (Accession # ACV96720); Cry1Be4 (Accession # HM070026);
 Cry1Bf1 (Accession # CAC50778); Cry1Bf2 (Accession # AAQ52380); Cry1Bg1
 20 (Accession # AAO39720); Cry1Bh1 (Accession # HQ589331); Cry1Bi1 (Accession #
 KC156700); Cry1Ca1 (Accession # CAA30396); Cry1Ca2 (Accession # CAA31951);
 Cry1Ca3 (Accession # AAA22343); Cry1Ca4 (Accession # CAA01886); Cry1Ca5
 (Accession # CAA65457); Cry1Ca6 [1] (Accession # AAF37224); Cry1Ca7 (Accession #
 AAG50438); Cry1Ca8 (Accession # AAM00264); Cry1Ca9 (Accession # AAL79362);
 25 Cry1Ca10 (Accession # AAN16462); Cry1Ca11 (Accession # AAX53094); Cry1Ca12
 (Accession # HM070027); Cry1Ca13 (Accession # HQ412621); Cry1Ca14 (Accession #
 JN651493); Cry1Cb1 (Accession # M97880); Cry1Cb2 (Accession # AAG35409);
 Cry1Cb3 (Accession # ACD50894); Cry1Cb-like (Accession # AAX63901); Cry1Da1
 (Accession # CAA38099); Cry1Da2 (Accession # I76415); Cry1Da3 (Accession #
 30 HQ439784); Cry1Db1 (Accession # CAA80234); Cry1Db2 (Accession # AAK48937);
 Cry1Dc1 (Accession # ABK35074); Cry1Ea1 (Accession # CAA37933); Cry1Ea2
 (Accession # CAA39609); Cry1Ea3 (Accession # AAA22345); Cry1Ea4 (Accession #
 AAD04732); Cry1Ea5 (Accession # A15535); Cry1Ea6 (Accession # AAL50330); Cry1Ea7
 (Accession # AAW72936); Cry1Ea8 (Accession # ABX11258); Cry1Ea9 (Accession #
 35 HQ439785); Cry1Ea10 (Accession # ADR00398); Cry1Ea11 (Accession # JQ652456);
 Cry1Eb1 (Accession # AAA22346); Cry1Fa1 (Accession # AAA22348); Cry1Fa2

(Accession # AAA22347); Cry1Fa3 (Accession # HM070028); Cry1Fa4 (Accession # HM439638); Cry1Fb1 (Accession # CAA80235); Cry1Fb2 (Accession # BAA25298); Cry1Fb3 (Accession # AAF21767); Cry1Fb4 (Accession # AAC10641); Cry1Fb5 (Accession # AAO13295); Cry1Fb6 (Accession # ACD50892); Cry1Fb7 (Accession # ACD50893); Cry1Ga1 (Accession # CAA80233); Cry1Ga2 (Accession # CAA70506); Cry1Gb1 (Accession # AAD10291); Cry1Gb2 (Accession # AAO13756); Cry1Gc1 (Accession # AAQ52381); Cry1Ha1 (Accession # CAA80236); Cry1Hb1 (Accession # AAA79694); Cry1Hb2 (Accession # HQ439786); Cry1H-like (Accession # AAF01213); Cry1Ia1 (Accession # CAA44633); Cry1Ia2 (Accession # AAA22354); Cry1Ia3 (Accession # AAC36999); Cry1Ia4 (Accession # AAB00958); Cry1Ia5 (Accession # CAA70124); Cry1Ia6 (Accession # AAC26910); Cry1Ia7 (Accession # AAM73516); Cry1Ia8 (Accession # AAK66742); Cry1Ia9 (Accession # AAQ08616); Cry1Ia10 (Accession # AAP86782); Cry1Ia11 (Accession # CAC85964); Cry1Ia12 (Accession # AAV53390); Cry1Ia13 (Accession # ABF83202); Cry1Ia14 (Accession # ACG63871); Cry1Ia15 (Accession # FJ617445); Cry1Ia16 (Accession # FJ617448); Cry1Ia17 (Accession # GU989199); Cry1Ia18 (Accession # ADK23801); Cry1Ia19 (Accession # HQ439787); Cry1Ia20 (Accession # JQ228426); Cry1Ia21 (Accession # JQ228424); Cry1Ia22 (Accession # JQ228427); Cry1Ia23 (Accession # JQ228428); Cry1Ia24 (Accession # JQ228429); Cry1Ia25 (Accession # JQ228430); Cry1Ia26 (Accession # JQ228431); Cry1Ia27 (Accession # JQ228432); Cry1Ia28 (Accession # JQ228433); Cry1Ia29 (Accession # JQ228434); Cry1Ia30 (Accession # JQ317686); Cry1Ia31 (Accession # JX944038); Cry1Ia32 (Accession # JX944039); Cry1Ia33 (Accession # JX944040); Cry1Ib1 (Accession # AAA82114); Cry1Ib2 (Accession # ABW88019); Cry1Ib3 (Accession # ACD75515); Cry1Ib4 (Accession # HM051227); Cry1Ib5 (Accession # HM070028); Cry1Ib6 (Accession # ADK38579); Cry1Ib7 (Accession # JN571740); Cry1Ib8 (Accession # JN675714); Cry1Ib9 (Accession # JN675715); Cry1Ib10 (Accession # JN675716); Cry1Ib11 (Accession # JQ228423); Cry1Ic1 (Accession # AAC62933); Cry1Ic2 (Accession # AAE71691); Cry1Id1 (Accession # AAD44366); Cry1Id2 (Accession # JQ228422); Cry1Ie1 (Accession # AAG43526); Cry1Ie2 (Accession # HM439636); Cry1Ie3 (Accession # KC156647); Cry1Ie4 (Accession # KC156681); Cry1If1 (Accession # AAQ52382); Cry1Ig1 (Accession # KC156701); Cry1I-like (Accession # AAC31094); Cry1I-like (Accession # ABG88859); Cry1Ja1 (Accession # AAA22341); Cry1Ja2 (Accession # HM070030); Cry1Ja3 (Accession # JQ228425); Cry1Jb1 (Accession # AAA98959); Cry1Jc1 (Accession # AAC31092); Cry1Jc2 (Accession # AAQ52372); Cry1Jd1 (Accession # CAC50779); Cry1Ka1 (Accession # AAB00376); Cry1Ka2 (Accession # HQ439783); Cry1La1 (Accession # AAS60191); Cry1La2 (Accession #

HM070031); Cry1Ma1 (Accession # FJ884067); Cry1Ma2 (Accession # KC156659);
 Cry1Na1 (Accession # KC156648); Cry1Nb1 (Accession # KC156678); Cry1-
 like (Accession # AAC31091); Cry2Aa1 (Accession # AAA22335); Cry2Aa2 (Accession #
 AAA83516); Cry2Aa3 (Accession # D86064); Cry2Aa4 (Accession # AAC04867);
 5 Cry2Aa5 (Accession # CAA10671); Cry2Aa6 (Accession # CAA10672); Cry2Aa7
 (Accession # CAA10670); Cry2Aa8 (Accession # AAO13734); Cry2Aa9 (Accession #
 AAO13750); Cry2Aa10 (Accession # AAQ04263); Cry2Aa11 (Accession # AAQ52384);
 Cry2Aa12 (Accession # ABI83671); Cry2Aa13 (Accession # ABL01536); Cry2Aa14
 (Accession # ACF04939); Cry2Aa15 (Accession # JN426947); Cry2Ab1 (Accession #
 10 AAA22342); Cry2Ab2 (Accession # CAA39075); Cry2Ab3 (Accession # AAG36762);
 Cry2Ab4 (Accession # AAO13296); Cry2Ab5 (Accession # AAQ04609); Cry2Ab6
 (Accession # AAP59457); Cry2Ab7 (Accession # AAZ66347); Cry2Ab8 (Accession #
 ABC95996); Cry2Ab9 (Accession # ABC74968); Cry2Ab10 (Accession # EF157306);
 Cry2Ab11 (Accession # CAM84575); Cry2Ab12 (Accession # ABM21764); Cry2Ab13
 15 (Accession # ACG76120); Cry2Ab14 (Accession # ACG76121); Cry2Ab15 (Accession #
 HM037126); Cry2Ab16 (Accession # GQ866914); Cry2Ab17 (Accession # HQ439789);
 Cry2Ab18 (Accession # JN135255); Cry2Ab19 (Accession # JN135256); Cry2Ab20
 (Accession # JN135257); Cry2Ab21 (Accession # JN135258); Cry2Ab22 (Accession #
 JN135259); Cry2Ab23 (Accession # JN135260); Cry2Ab24 (Accession # JN135261);
 20 Cry2Ab25 (Accession # JN415485); Cry2Ab26 (Accession # JN426946); Cry2Ab27
 (Accession # JN415764); Cry2Ab28 (Accession # JN651494); Cry2Ac1 (Accession #
 CAA40536); Cry2Ac2 (Accession # AAG35410); Cry2Ac3 (Accession # AAQ52385);
 Cry2Ac4 (Accession # ABC95997); Cry2Ac5 (Accession # ABC74969); Cry2Ac6
 (Accession # ABC74793); Cry2Ac7 (Accession # CAL18690); Cry2Ac8 (Accession #
 25 CAM09325); Cry2Ac9 (Accession # CAM09326); Cry2Ac10 (Accession # ABN15104);
 Cry2Ac11 (Accession # CAM83895); Cry2Ac12 (Accession # CAM83896); Cry2Ad1
 (Accession # AAF09583); Cry2Ad2 (Accession # ABC86927); Cry2Ad3 (Accession #
 CAK29504); Cry2Ad4 (Accession # CAM32331); Cry2Ad5 (Accession # CAO78739);
 Cry2Ae1 (Accession # AAQ52362); Cry2Af1 (Accession # ABO30519); Cry2Af2
 30 (Accession # GQ866915); Cry2Ag1 (Accession # ACH91610); Cry2Ah1 (Accession #
 EU939453); Cry2Ah2 (Accession # ACL80665); Cry2Ah3 (Accession # GU073380);
 Cry2Ah4 (Accession # KC156702); Cry2Ai1 (Accession # FJ788388); Cry2Aj (Accession
 #); Cry2Ak1 (Accession # KC156660); Cry2Ba1 (Accession # KC156658); Cry3Aa1
 (Accession # AAA22336); Cry3Aa2 (Accession # AAA22541); Cry3Aa3 (Accession #
 35 CAA68482); Cry3Aa4 (Accession # AAA22542); Cry3Aa5 (Accession # AAA50255);
 Cry3Aa6 (Accession # AAC43266); Cry3Aa7 (Accession # CAB41411); Cry3Aa8

(Accession # AAS79487); Cry3Aa9 (Accession # AAW05659); Cry3Aa10 (Accession # AAU29411); Cry3Aa11 (Accession # AAW82872); Cry3Aa12 (Accession # ABY49136); Cry3Ba1 (Accession # CAA34983); Cry3Ba2 (Accession # CAA00645); Cry3Ba3 (Accession # JQ397327); Cry3Bb1 (Accession # AAA22334); Cry3Bb2 (Accession # AAA74198); Cry3Bb3 (Accession # I15475); Cry3Ca1 (Accession # CAA42469); Cry4Aa1 (Accession # CAA68485); Cry4Aa2 (Accession # BAA00179); Cry4Aa3 (Accession # CAD30148); Cry4Aa4 (Accession # AFB18317); Cry4A-like (Accession # AAY96321); Cry4Ba1 (Accession # CAA30312); Cry4Ba2 (Accession # CAA30114); Cry4Ba3 (Accession # AAA22337); Cry4Ba4 (Accession # BAA00178); Cry4Ba5 (Accession # CAD30095); Cry4Ba-like (Accession # ABC47686); Cry4Ca1 (Accession # EU646202); Cry4Cb1 (Accession # FJ403208); Cry4Cb2 (Accession # FJ597622); Cry4Cc1 (Accession # FJ403207); Cry5Aa1 (Accession # AAA67694); Cry5Ab1 (Accession # AAA67693); Cry5Ac1 (Accession # I34543); Cry5Ad1 (Accession # ABQ82087); Cry5Ba1 (Accession # AAA68598); Cry5Ba2 (Accession # ABW88931); Cry5Ba3 (Accession # AFJ04417); Cry5Ca1 (Accession # HM461869); Cry5Ca2 (Accession # ZP_04123426); Cry5Da1 (Accession # HM461870); Cry5Da2 (Accession # ZP_04123980); Cry5Ea1 (Accession # HM485580); Cry5Ea2 (Accession # ZP_04124038); Cry6Aa1 (Accession # AAA22357); Cry6Aa2 (Accession # AAM46849); Cry6Aa3 (Accession # ABH03377); Cry6Ba1 (Accession # AAA22358); Cry7Aa1 (Accession # AAA22351); Cry7Ab1 (Accession # AAA21120); Cry7Ab2 (Accession # AAA21121); Cry7Ab3 (Accession # ABX24522); Cry7Ab4 (Accession # EU380678); Cry7Ab5 (Accession # ABX79555); Cry7Ab6 (Accession # ACI44005); Cry7Ab7 (Accession # ADB89216); Cry7Ab8 (Accession # GU145299); Cry7Ab9 (Accession # ADD92572); Cry7Ba1 (Accession # ABB70817); Cry7Bb1 (Accession # KC156653); Cry7Ca1 (Accession # ABR67863); Cry7Cb1 (Accession # KC156698); Cry7Da1 (Accession # ACQ99547); Cry7Da2 (Accession # HM572236); Cry7Da3 (Accession # KC156679); Cry7Ea1 (Accession # HM035086); Cry7Ea2 (Accession # HM132124); Cry7Ea3 (Accession # EEM19403); Cry7Fa1 (Accession # HM035088); Cry7Fa2 (Accession # EEM19090); Cry7Fb1 (Accession # HM572235); Cry7Fb2 (Accession # KC156682); Cry7Ga1 (Accession # HM572237); Cry7Ga2 (Accession # KC156669); Cry7Gb1 (Accession # KC156650); Cry7Gc1 (Accession # KC156654); Cry7Gd1 (Accession # KC156697); Cry7Ha1 (Accession # KC156651); Cry7Ia1 (Accession # KC156665); Cry7Ja1 (Accession # KC156671); Cry7Ka1 (Accession # KC156680); Cry7Kb1 (Accession # BAM99306); Cry7La1 (Accession # BAM99307); Cry8Aa1 (Accession # AAA21117); Cry8Ab1 (Accession # EU044830); Cry8Ac1 (Accession # KC156662); Cry8Ad1 (Accession # KC156684); Cry8Ba1 (Accession # AAA21118); Cry8Bb1 (Accession # CAD57542);

Cry8Bc1 (Accession # CAD57543); Cry8Ca1 (Accession # AAA21119); Cry8Ca2
 (Accession # AAR98783); Cry8Ca3 (Accession # EU625349); Cry8Ca4 (Accession #
 ADB54826); Cry8Da1 (Accession # BAC07226); Cry8Da2 (Accession # BD133574);
 Cry8Da3 (Accession # BD133575); Cry8Db1 (Accession # BAF93483); Cry8Ea1
 5 (Accession # AAQ73470); Cry8Ea2 (Accession # EU047597); Cry8Ea3 (Accession #
 KC855216); Cry8Fa1 (Accession # AAT48690); Cry8Fa2 (Accession # HQ174208);
 Cry8Fa3 (Accession # AFH78109); Cry8Ga1 (Accession # AAT46073); Cry8Ga2
 (Accession # ABC42043); Cry8Ga3 (Accession # FJ198072); Cry8Ha1 (Accession #
 AAW81032); Cry8Ia1 (Accession # EU381044); Cry8Ia2 (Accession # GU073381);
 10 Cry8Ia3 (Accession # HM044664); Cry8Ia4 (Accession # KC156674); Cry8Ib1 (Accession
 # GU325772); Cry8Ib2 (Accession # KC156677); Cry8Ja1 (Accession # EU625348);
 Cry8Ka1 (Accession # FJ422558); Cry8Ka2 (Accession # ACN87262); Cry8Kb1
 (Accession # HM123758); Cry8Kb2 (Accession # KC156675); Cry8La1 (Accession #
 GU325771); Cry8Ma1 (Accession # HM044665); Cry8Ma2 (Accession # EEM86551);
 15 Cry8Ma3 (Accession # HM210574); Cry8Na1 (Accession # HM640939); Cry8Pa1
 (Accession # HQ388415); Cry8Qa1 (Accession # HQ441166); Cry8Qa2 (Accession #
 KC152468); Cry8Ra1 (Accession # AFP87548); Cry8Sa1 (Accession # JQ740599);
 Cry8Ta1 (Accession # KC156673); Cry8-like (Accession # FJ770571); Cry8-like
 (Accession # ABS53003); Cry9Aa1 (Accession # CAA41122); Cry9Aa2 (Accession #
 20 CAA41425); Cry9Aa3 (Accession # GQ249293); Cry9Aa4 (Accession # GQ249294);
 Cry9Aa5 (Accession # JX174110); Cry9Aa like (Accession # AAQ52376); Cry9Ba1
 (Accession # CAA52927); Cry9Ba2 (Accession # GU299522); Cry9Bb1 (Accession #
 AAV28716); Cry9Ca1 (Accession # CAA85764); Cry9Ca2 (Accession # AAQ52375);
 Cry9Da1 (Accession # BAA19948); Cry9Da2 (Accession # AAB97923); Cry9Da3
 25 (Accession # GQ249293); Cry9Da4 (Accession # GQ249297); Cry9Db1 (Accession #
 AAX78439); Cry9Dc1 (Accession # KC156683); Cry9Ea1 (Accession # BAA34908);
 Cry9Ea2 (Accession # AAO12908); Cry9Ea3 (Accession # ABM21765); Cry9Ea4
 (Accession # ACE88267); Cry9Ea5 (Accession # ACF04743); Cry9Ea6 (Accession #
 ACG63872); Cry9Ea7 (Accession # FJ380927); Cry9Ea8 (Accession # GQ249292);
 30 Cry9Ea9 (Accession # JN651495); Cry9Eb1 (Accession # CAC50780); Cry9Eb2
 (Accession # GQ249298); Cry9Eb3 (Accession # KC156646); Cry9Ec1 (Accession #
 AAC63366); Cry9Ed1 (Accession # AAX78440); Cry9Ee1 (Accession # GQ249296);
 Cry9Ee2 (Accession # KC156664); Cry9Fa1 (Accession # KC156692); Cry9Ga1
 (Accession # KC156699); Cry9-like (Accession # AAC63366); Cry10Aa1 (Accession #
 35 AAA22614); Cry10Aa2 (Accession # E00614); Cry10Aa3 (Accession # CAD30098);
 Cry10Aa4 (Accession # AFB18318); Cry10A-like (Accession # DQ167578); Cry11Aa1

(Accession # AAA22352); Cry11Aa2 (Accession # AAA22611); Cry11Aa3 (Accession # CAD30081); Cry11Aa4 (Accession # AFB18319); Cry11Aa-like (Accession # DQ166531); Cry11Ba1 (Accession # CAA60504); Cry11Bb1 (Accession # AAC97162); Cry11Bb2 (Accession # HM068615); Cry12Aa1 (Accession # AAA22355); Cry13Aa1 (Accession # AAA22356); Cry14Aa1 (Accession # AAA21516); Cry14Ab1 (Accession # KC156652); Cry15Aa1 (Accession # AAA22333); Cry16Aa1 (Accession # CAA63860); Cry17Aa1 (Accession # CAA67841); Cry18Aa1 (Accession # CAA67506); Cry18Ba1 (Accession # AAF89667); Cry18Ca1 (Accession # AAF89668); Cry19Aa1 (Accession # CAA68875); Cry19Ba1 (Accession # BAA32397); Cry19Ca1 (Accession # AFM37572); Cry20Aa1 (Accession # AAB93476); Cry20Ba1 (Accession # ACS93601); Cry20Ba2 (Accession # KC156694); Cry20-like (Accession # GQ144333); Cry21Aa1 (Accession # I32932); Cry21Aa2 (Accession # I66477); Cry21Ba1 (Accession # BAC06484); Cry21Ca1 (Accession # JF521577); Cry21Ca2 (Accession # KC156687); Cry21Da1 (Accession # JF521578); Cry22Aa1 (Accession # I34547); Cry22Aa2 (Accession # CAD43579); Cry22Aa3 (Accession # ACD93211); Cry22Ab1 (Accession # AAK50456); Cry22Ab2 (Accession # CAD43577); Cry22Ba1 (Accession # CAD43578); Cry22Bb1 (Accession # KC156672); Cry23Aa1 (Accession # AAF76375); Cry24Aa1 (Accession # AAC61891); Cry24Ba1 (Accession # BAD32657); Cry24Ca1 (Accession # CAJ43600); Cry25Aa1 (Accession # AAC61892); Cry26Aa1 (Accession # AAD25075); Cry27Aa1 (Accession # BAA82796); Cry28Aa1 (Accession # AAD24189); Cry28Aa2 (Accession # AAG00235); Cry29Aa1 (Accession # CAC80985); Cry30Aa1 (Accession # CAC80986); Cry30Ba1 (Accession # BAD00052); Cry30Ca1 (Accession # BAD67157); Cry30Ca2 (Accession # ACU24781); Cry30Da1 (Accession # EF095955); Cry30Db1 (Accession # BAE80088); Cry30Ea1 (Accession # ACC95445); Cry30Ea2 (Accession # FJ499389); Cry30Fa1 (Accession # ACI22625); Cry30Ga1 (Accession # ACG60020); Cry30Ga2 (Accession # HQ638217); Cry31Aa1 (Accession # BAB11757); Cry31Aa2 (Accession # AAL87458); Cry31Aa3 (Accession # BAE79808); Cry31Aa4 (Accession # BAF32571); Cry31Aa5 (Accession # BAF32572); Cry31Aa6 (Accession # BAI44026); Cry31Ab1 (Accession # BAE79809); Cry31Ab2 (Accession # BAF32570); Cry31Ac1 (Accession # BAF34368); Cry31Ac2 (Accession # AB731600); Cry31Ad1 (Accession # BAI44022); Cry32Aa1 (Accession # AAG36711); Cry32Aa2 (Accession # GU063849); Cry32Ab1 (Accession # GU063850); Cry32Ba1 (Accession # BAB78601); Cry32Ca1 (Accession # BAB78602); Cry32Cb1 (Accession # KC156708); Cry32Da1 (Accession # BAB78603); Cry32Ea1 (Accession # GU324274); Cry32Ea2 (Accession # KC156686); Cry32Eb1 (Accession # KC156663); Cry32Fa1 (Accession # KC156656); Cry32Ga1 (Accession # KC156657); Cry32Ha1 (Accession # KC156661); Cry32Hb1 (Accession # KC156666); Cry32Ia1

(Accession # KC156667); Cry32Ja1 (Accession # KC156685); Cry32Ka1 (Accession # KC156688); Cry32La1 (Accession # KC156689); Cry32Ma1 (Accession # KC156690); Cry32Mb1 (Accession # KC156704); Cry32Na1 (Accession # KC156691); Cry32Oa1 (Accession # KC156703); Cry32Pa1 (Accession # KC156705); Cry32Qa1 (Accession # KC156706); Cry32Ra1 (Accession # KC156707); Cry32Sa1 (Accession # KC156709); Cry32Ta1 (Accession # KC156710); Cry32Ua1 (Accession # KC156655); Cry33Aa1 (Accession # AAL26871); Cry34Aa1 (Accession # AAG50341); Cry34Aa2 (Accession # AAK64560); Cry34Aa3 (Accession # AAT29032); Cry34Aa4 (Accession # AAT29030); Cry34Ab1 (Accession # AAG41671); Cry34Ac1 (Accession # AAG50118); Cry34Ac2 (Accession # AAK64562); Cry34Ac3 (Accession # AAT29029); Cry34Ba1 (Accession # AAK64565); Cry34Ba2 (Accession # AAT29033); Cry34Ba3 (Accession # AAT29031); Cry35Aa1 (Accession # AAG50342); Cry35Aa2 (Accession # AAK64561); Cry35Aa3 (Accession # AAT29028); Cry35Aa4 (Accession # AAT29025); Cry35Ab1 (Accession # AAG41672); Cry35Ab2 (Accession # AAK64563); Cry35Ab3 (Accession # AY536891); Cry35Ac1 (Accession # AAG50117); Cry35Ba1 (Accession # AAK64566); Cry35Ba2 (Accession # AAT29027); Cry35Ba3 (Accession # AAT29026); Cry36Aa1 (Accession # AAK64558); Cry37Aa1 (Accession # AAF76376); Cry38Aa1 (Accession # AAK64559); Cry39Aa1 (Accession # BAB72016); Cry40Aa1 (Accession # BAB72018); Cry40Ba1 (Accession # BAC77648); Cry40Ca1 (Accession # EU381045); Cry40Da1 (Accession # ACF15199); Cry41Aa1 (Accession # BAD35157); Cry41Ab1 (Accession # BAD35163); Cry41Ba1 (Accession # HM461871); Cry41Ba2 (Accession # ZP_04099652); Cry42Aa1 (Accession # BAD35166); Cry43Aa1 (Accession # BAD15301); Cry43Aa2 (Accession # BAD95474); Cry43Ba1 (Accession # BAD15303); Cry43Ca1 (Accession # KC156676); Cry43Cb1 (Accession # KC156695); Cry43Cc1 (Accession # KC156696); Cry43-like (Accession # BAD15305); Cry44Aa (Accession # BAD08532); Cry45Aa (Accession # BAD22577); Cry46Aa (Accession # BAC79010); Cry46Aa2 (Accession # BAG68906); Cry46Ab (Accession # BAD35170); Cry47Aa (Accession # AAY24695); Cry48Aa (Accession # CAJ18351); Cry48Aa2 (Accession # CAJ86545); Cry48Aa3 (Accession # CAJ86546); Cry48Ab (Accession # CAJ86548); Cry48Ab2 (Accession # CAJ86549); Cry49Aa (Accession # CAH56541); Cry49Aa2 (Accession # CAJ86541); Cry49Aa3 (Accession # CAJ86543); Cry49Aa4 (Accession # CAJ86544); Cry49Ab1 (Accession # CAJ86542); Cry50Aa1 (Accession # BAE86999); Cry50Ba1 (Accession # GU446675); Cry50Ba2 (Accession # GU446676); Cry51Aa1 (Accession # ABI14444); Cry51Aa2 (Accession # GU570697); Cry52Aa1 (Accession # EF613489); Cry52Ba1 (Accession # FJ361760); Cry53Aa1 (Accession # EF633476); Cry53Ab1 (Accession # FJ361759); Cry54Aa1 (Accession # ACA52194); Cry54Aa2 (Accession # GQ140349); Cry54Ba1

(Accession # GU446677); Cry55Aa1 (Accession # ABW88932); Cry54Ab1 (Accession # JQ916908); Cry55Aa2 (Accession # AAE33526); Cry56Aa1 (Accession # ACU57499); Cry56Aa2 (Accession # GQ483512); Cry56Aa3 (Accession # JX025567); Cry57Aa1 (Accession # ANC87261); Cry58Aa1 (Accession # ANC87260); Cry59Ba1 (Accession # JN790647); Cry59Aa1 (Accession # ACR43758); Cry60Aa1 (Accession # ACU24782); Cry60Aa2 (Accession # EAO57254); Cry60Aa3 (Accession # EEM99278); Cry60Ba1 (Accession # GU810818); Cry60Ba2 (Accession # EAO57253); Cry60Ba3 (Accession # EEM99279); Cry61Aa1 (Accession # HM035087); Cry61Aa2 (Accession # HM132125); Cry61Aa3 (Accession # EEM19308); Cry62Aa1 (Accession # HM054509); Cry63Aa1 (Accession # BAI44028); Cry64Aa1 (Accession # BAJ05397); Cry65Aa1 (Accession # HM461868); Cry65Aa2 (Accession # ZP_04123838); Cry66Aa1 (Accession # HM485581); Cry66Aa2 (Accession # ZP_04099945); Cry67Aa1 (Accession # HM485582); Cry67Aa2 (Accession # ZP_04148882); Cry68Aa1 (Accession # HQ113114); Cry69Aa1 (Accession # HQ401006); Cry69Aa2 (Accession # JQ821388); Cry69Ab1 (Accession # JN209957); Cry70Aa1 (Accession # JN646781); Cry70Ba1 (Accession # ADO51070); Cry70Bb1 (Accession # EEL67276); Cry71Aa1 (Accession # JX025568); Cry72Aa1 (Accession # JX025569); and Cry73Aa (Accession # AEH76822); Cyt1Aa (GenBank Accession # X03182), Cyt1Ab (GenBank Accession # X98793), Cyt1B (GenBank Accession # U37196), Cyt2A (GenBank Accession # Z14147), Cyt2B (GenBank Accession # U52043).

Examples of δ -endotoxins also include but are not limited to Cry1A proteins of US Patent Numbers 5,880,275 and 7,858,849; a DIG-3 or DIG-11 toxin (N-terminal deletion of α -helix 1 and/or α -helix 2 variants of cry proteins such as Cry1A, Cry3A) of US Patent Numbers 8,304,604 and 8,304,605, Cry1B of US Patent Application Serial Number 10/525,318; Cry1C of US Patent Number 6,033,874; Cry1F of US Patent Numbers 5,188,960, 6,218,188; Cry1A/F chimeras of US Patent Numbers 7,070,982; 6,962,705 and 6,713,063; a Cry2 protein such as Cry2Ab protein of US Patent Number 7,064,249; a Cry3A protein including but not limited to an engineered hybrid insecticidal protein (eHIP) created by fusing unique combinations of variable regions and conserved blocks of at least two different Cry proteins (US Patent Application Publication Number 2010/0017914); a Cry4 protein; a Cry5 protein; a Cry6 protein; Cry8 proteins of US Patent Numbers 7,329,736, 7,449,552, 7,803,943, 7,476,781, 7,105,332, 7,378,499 and 7,462,760; a Cry9 protein such as members of the Cry9A, Cry9B, Cry9C, Cry9D, Cry9E, and Cry9F families; a Cry15 protein of Naimov, *et al.*, (2008) *Applied and Environmental Microbiology* 74:7145–7151; a Cry22, a Cry34Ab1 protein of US Patent Numbers 6,127,180, 6,624,145 and 6,340,593; a CryET33 and cryET34 protein of US

Patent Numbers 6,248,535, 6,326,351, 6,399,330, 6,949,626, 7,385,107 and 7,504,229; a CryET33 and CryET34 homologs of US Patent Publication Number 2006/0191034, 2012/0278954, and PCT Publication Number WO 2012/139004; a Cry35Ab1 protein of US Patent Numbers 6,083,499, 6,548,291 and 6,340,593; a Cry46 protein, a Cry 51 protein, a Cry binary toxin; a TIC901 or related toxin; TIC807 of US 2008/0295207; ET29, ET37, TIC809, TIC810, TIC812, TIC127, TIC128 of PCT US 2006/033867; AXMI-027, AXMI-036, and AXMI-038 of US Patent Number 8,236,757; AXMI-031, AXMI-039, AXMI-040, AXMI-049 of US7,923,602; AXMI-018, AXMI-020, and AXMI-021 of WO 2006/083891; AXMI-010 of WO 2005/038032; AXMI-003 of WO 2005/021585; AXMI-008 of US 2004/0250311; AXMI-006 of US 2004/0216186; AXMI-007 of US 2004/0210965; AXMI-009 of US 2004/0210964; AXMI-014 of US 2004/0197917; AXMI-004 of US 2004/0197916; AXMI-028 and AXMI-029 of WO 2006/119457; AXMI-007, AXMI-008, AXMI-0080rf2, AXMI-009, AXMI-014 and AXMI-004 of WO 2004/074462; AXMI-150 of US Patent Number 8,084,416; AXMI-205 of US20110023184; AXMI-011, AXMI-012, AXMI-013, AXMI-015, AXMI-019, AXMI-044, AXMI-037, AXMI-043, AXMI-033, AXMI-034, AXMI-022, AXMI-023, AXMI-041, AXMI-063, and AXMI-064 of US 2011/0263488; AXMI-R1 and related proteins of US 2010/0197592; AXMI221Z, AXMI222z, AXMI223z, AXMI224z and AXMI225z of WO 2011/103248; AXMI218, AXMI219, AXMI220, AXMI226, AXMI227, AXMI228, AXMI229, AXMI230, and AXMI231 of WO11/103247; AXMI-115, AXMI-113, AXMI-005, AXMI-163 and AXMI-184 of US Patent Number 8,334,431; AXMI-001, AXMI-002, AXMI-030, AXMI-035, and AXMI-045 of US 2010/0298211; AXMI-066 and AXMI-076 of US20090144852; AXMI128, AXMI130, AXMI131, AXMI133, AXMI140, AXMI141, AXMI142, AXMI143, AXMI144, AXMI146, AXMI148, AXMI149, AXMI152, AXMI153, AXMI154, AXMI155, AXMI156, AXMI157, AXMI158, AXMI162, AXMI165, AXMI166, AXMI167, AXMI168, AXMI169, AXMI170, AXMI171, AXMI172, AXMI173, AXMI174, AXMI175, AXMI176, AXMI177, AXMI178, AXMI179, AXMI180, AXMI181, AXMI182, AXMI185, AXMI186, AXMI187, AXMI188, AXMI189 of US Patent Number 8,318,900; AXMI079, AXMI080, AXMI081, AXMI082, AXMI091, AXMI092, AXMI096, AXMI097, AXMI098, AXMI099, AXMI100, AXMI101, AXMI102, AXMI103, AXMI104, AXMI107, AXMI108, AXMI109, AXMI110, AXMI111, AXMI112, AXMI114, AXMI116, AXMI117, AXMI118, AXMI119, AXMI120, AXMI121, AXMI122, AXMI123, AXMI124, AXMI1257, AXMI1268, AXMI127, AXMI129, AXMI164, AXMI151, AXMI161, AXMI183, AXMI132, AXMI138, AXMI137 of US 2010/0005543; cry proteins such as Cry1A and Cry3A having modified proteolytic sites of US Patent Number 8,319,019; a Cry1Ac, Cry2Aa and Cry1Ca toxin protein from *Bacillus thuringiensis* strain VBTS 2528 of US Patent Application Publication Number 2011/0064710. Other Cry proteins are well known

to one skilled in the art (see, Crickmore, *et al.*, "Bacillus thuringiensis toxin nomenclature" (2011), at lifesci.sussex.ac.uk/home/Neil_Crickmore/Bt/ which can be accessed on the world-wide web using the "www" prefix). The insecticidal activity of Cry proteins is well known to one skilled in the art (for review, see, van Frannkenhuyzen, (2009) *J. Invert. Path.* 101:1-16). The use of Cry proteins as transgenic plant traits is well known to one skilled in the art and Cry-transgenic plants including but not limited to Cry1Ac, Cry1Ac+Cry2Ab, Cry1Ab, Cry1A.105, Cry1F, Cry1Fa2, Cry1F+Cry1Ac, Cry2Ab, Cry3A, mCry3A, Cry3Bb1, Cry34Ab1, Cry35Ab1, Vip3A, mCry3A, Cry9c and CBI-Bt have received regulatory approval (see, Sanahuja, (2011) *Plant Biotech Journal* 9:283-300 and the CERA (2010) GM Crop Database Center for Environmental Risk Assessment (CERA), ILSI Research Foundation, Washington D.C. at cera-gmc.org/index.php?action=gm_crop_database which can be accessed on the world-wide web using the "www" prefix). More than one pesticidal proteins well known to one skilled in the art can also be expressed in plants such as Vip3Ab & Cry1Fa (US2012/0317682), Cry1BE & Cry1F (US2012/0311746), Cry1CA & Cry1AB (US2012/0311745), Cry1F & CryCa (US2012/0317681), Cry1DA & Cry1BE (US2012/0331590), Cry1DA & Cry1Fa (US2012/0331589), Cry1AB & Cry1BE (US2012/0324606), and Cry1Fa & Cry2Aa, Cry1I or Cry1E (US2012/0324605). Pesticidal proteins also include insecticidal lipases including lipid acyl hydrolases of US Patent Number 7,491,869, and cholesterol oxidases such as from *Streptomyces* (Purcell et al. (1993) *Biochem Biophys Res Commun* 15:1406-1413). Pesticidal proteins also include VIP (vegetative insecticidal proteins) toxins of US Patent Numbers 5,877,012, 6,107,279 6,137,033, 7,244,820, 7,615,686, and 8,237,020 and the like. Other VIP proteins are well known to one skilled in the art (see, lifesci.sussex.ac.uk/home/Neil_Crickmore/Bt/vip.html which can be accessed on the world-wide web using the "www" prefix). Pesticidal proteins also include toxin complex (TC) proteins, obtainable from organisms such as *Xenorhabdus*, *Photorhabdus* and *Paenibacillus* (see, US Patent Numbers 7,491,698 and 8,084,418). Some TC proteins have "stand alone" insecticidal activity and other TC proteins enhance the activity of the stand-alone toxins produced by the same given organism. The toxicity of a "stand-alone" TC protein (from *Photorhabdus*, *Xenorhabdus* or *Paenibacillus*, for example) can be enhanced by one or more TC protein "potentiators" derived from a source organism of a different genus. There are three main types of TC proteins. As referred to herein, Class A proteins ("Protein A") are stand-alone toxins. Class B proteins ("Protein B") and Class C proteins ("Protein C") enhance the toxicity of Class A proteins. Examples of Class A proteins are TcbA, TcdA, XptA1 and XptA2. Examples of Class B proteins are TcaC, TcdB, XptB1Xb and XptC1Wi. Examples of Class C proteins are TccC, XptC1Xb and

XptB1Wi. Pesticidal proteins also include spider, snake and scorpion venom proteins. Examples of spider venom peptides include but not limited to lycotoxin-1 peptides and mutants thereof (US Patent Number 8,334,366).

In some embodiments the PHI-4 polypeptides include amino acid sequences deduced from the full-length nucleic acid sequences disclosed herein, and amino acid sequences that are shorter than the full-length sequences, either due to the use of an alternate downstream start site or due to processing that produces a shorter protein having pesticidal activity. Processing may occur in the organism the protein is expressed in or in the pest after ingestion of the protein.

Thus, provided herein are novel isolated or recombinant nucleic acid sequences that confer pesticidal activity. Also provided are the amino acid sequences of PHI-4 polypeptides. The protein resulting from translation of these PHI-4 polypeptide genes allows cells to control or kill pests that ingest it.

Nucleic Acid Molecules, and Variants and Fragments Thereof

One aspect pertains to isolated or recombinant nucleic acid molecules comprising nucleic acid sequences encoding PHI-4 polypeptides and polypeptides or biologically active portions thereof, as well as nucleic acid molecules sufficient for use as hybridization probes to identify nucleic acid molecules encoding proteins with regions of sequence homology. As used herein, the term “nucleic acid molecule” is intended to include DNA molecules (e.g., recombinant DNA, cDNA, genomic DNA, plastid DNA, mitochondrial DNA) and RNA molecules (e.g., mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

An “isolated” nucleic acid molecule (or DNA) is used herein to refer to a nucleic acid sequence (or DNA) that is no longer in its natural environment, for example in vitro. A “recombinant” nucleic acid molecule (or DNA) is used herein to refer to a nucleic acid sequence (or DNA) that is in a recombinant bacterial or plant host cell. In some embodiments, an “isolated” or “recombinant” nucleic acid is free of sequences (preferably protein encoding sequences) that naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For purposes of the disclosure, “isolated” or “recombinant” when used to refer to nucleic acid molecules excludes isolated chromosomes. For example, in various embodiments, the recombinant nucleic acid molecule encoding a PHI-4 polypeptide can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of

nucleic acid sequences that naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived.

In some embodiments a nucleic acid molecule encoding the PHI-4 polypeptide is a non-genomic nucleic acid sequence. As used herein a “non-genomic nucleic acid sequence “or “non-genomic nucleic acid molecule” refers to a nucleic acid molecule that has one or more change in the nucleic acid sequence compared to a native or genomic nucleic acid sequence. In some embodiments the change to a native or genomic nucleic acid molecule includes but is not limited to: changes in the nucleic acid sequence due to the degeneracy of the genetic code; codon optimization of the nucleic acid sequence for expression in plants; changes in the nucleic acid sequence to introduce at least one amino acid substitution, insertion, deletion and/or addition compared to the native or genomic sequence; removal of one or more intron associated with the genomic nucleic acid sequence; insertion of one or more heterologous introns; deletion of one or more upstream or downstream regulatory regions associated with the genomic nucleic acid sequence; insertion of one or more heterologous upstream or downstream regulatory regions; deletion of the 5' and/or 3' untranslated region associated with the genomic nucleic acid sequence; insertion of a heterologous 5' and/or 3' untranslated region; and modification of a polyadenylation site. In some embodiments the non-genomic nucleic acid molecule is a cDNA. In some embodiments the non-genomic nucleic acid molecule is a synthetic nucleic acid sequence.

A variety of polynucleotides encoding a PHI-4 polypeptide(s) or related proteins are contemplated. Such polynucleotides are useful for production of PHI-4 polypeptides in host cells when operably linked to suitable promoter, transcription termination and/or polyadenylation sequences. Such polynucleotides are also useful as probes for isolating homologous or substantially homologous polynucleotides encoding PHI-4 polypeptides or related proteins.

The present disclosure provides isolated or recombinant polynucleotides that encode any of the PHI-4 polypeptides disclosed herein. Those having ordinary skill in the art will readily appreciate that due to the degeneracy of the genetic code, a multitude of nucleotide sequences encoding β -glucosidase polypeptides of the present disclosure exist. Table 1 is a Codon Table that provides the synonymous codons for each amino acid. For example, the codons AGA, AGG, CGA, CGC, CGG, and CGU all encode the amino acid arginine. Thus, at every position in the nucleic acids of the disclosure where an arginine is specified by a codon, the codon can be altered to any of the corresponding codons described above without altering the encoded polypeptide. It is understood that U in an RNA sequence corresponds to T in a DNA sequence.

Table 1

Alanine	Ala	A	GCA GCC GCG GCU
Cysteine	Cys	C	UGC UGU
Aspartic acid	Asp	D	GAC GAU
Glutamic acid	Glu	E	GAA GAG
Phenylalanine	Phe	F	UUC UUU
Glycine	Gly	G	GGA GGC GGG GGU
Histidine	His	H	CAC CAU
Isoleucine	Ile	I	AUA AUC AUU
Lysine	Lys	K	AAA AAG
Leucine	Leu	L	UUA UUG CUA CUC CUG CUU
Methionine	Met	M	AUG
Asparagine	Asn	N	AAC AAU
Proline	Pro	P	CCA CCC CCG CCU
Glutamine	Gln	Q	CAA CAG
Arginine	Arg	R	AGA AGG CGA CGC CGG CGU
Serine	Ser	S	AGC AGU UCA UCC UCG UCU
Threonine	Thr	T	ACA ACC ACG ACU
Valine	Val	V	GUA GUC GUG GUU
Tryptophan	Trp	W	UGG
Tyrosine	Tyr	Y	UAC UAU

Such “silent variations” are one species of “conservative” variation. One of ordinary skill in the art will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine) can be modified by standard techniques to encode a functionally identical polypeptide. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide is implicit in any described sequence. The disclosure contemplates and provides each and every possible variation of nucleic acid sequence encoding a polypeptide of the disclosure that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code (set forth in Table 1), as applied to the polynucleotide sequences of the present disclosure.

A group of two or more different codons that, when translated in the same context, all encode the same amino acid, are referred to herein as “synonymous codons.” Polynucleotides encoding PHI-4 polypeptides of the present disclosure may be codon optimized for expression in a particular host organism by modifying the polynucleotides to

conform with the optimum codon usage of the desired host organism. Those having ordinary skill in the art will recognize that tables and other references providing preference information for a wide range of organisms are readily available.

Polynucleotides encoding a PHI-4 polypeptide can also be synthesized de novo
5 from a PHI-4 polypeptide sequence. The sequence of the polynucleotide gene can be deduced from a PHI-4 polypeptide sequence through use of the genetic code. Computer programs such as "BackTranslate" (GCG™ Package, Acclerys, Inc. San Diego, Calif.) can be used to convert a peptide sequence to the corresponding nucleotide sequence encoding the peptide. Examples of PHI-4 polypeptide sequences that can be used to
10 obtain corresponding nucleotide encoding sequences include, but are not limited to, the PHI-4 polypeptide sequence of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526. Furthermore, synthetic PHI-4 polynucleotide sequences of the disclosure can be designed so that they will be expressed in plants. US Patent Number 5,500,365 describes a method for synthesizing
15 plant genes to improve the expression level of the protein encoded by the synthesized gene. This method relates to the modification of the structural gene sequences of the exogenous transgene, to cause them to be more efficiently transcribed, processed, translated and expressed by the plant. Features of genes that are expressed well in plants include elimination of sequences that can cause undesired intron splicing or
20 polyadenylation in the coding region of a gene transcript while retaining substantially the amino acid sequence of the toxic portion of the insecticidal protein. A similar method for obtaining enhanced expression of transgenes in monocotyledonous plants is disclosed in US Patent Number 5,689,052.

In some embodiments the nucleic acid molecule encoding a PHI-4 polypeptide is a
25 polynucleotide having the sequence set forth in SEQ ID NO: 1, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NOs: 24-30, SEQ ID NOs: 1163-1505 and variants, fragments and complements thereof. By "complement" is intended a nucleic acid sequence that is sufficiently complementary to a given nucleic acid sequence such that it can hybridize to the given nucleic acid sequence to thereby form a stable duplex. In some embodiments
30 the nucleic acid molecule encoding a PHI-4 polypeptide is a nucleic acid molecule having the sequence set forth in SEQ ID NO: 1, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NOs: 24-30 and SEQ ID NOs: 1163-1505. The corresponding amino acid sequences for the insecticidal protein encoded by these nucleic acid sequences are set forth in SEQ ID NO: 1, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NOs: 24-30 and SEQ ID NOs: 1163-1505.

35 In some embodiments the nucleic acid molecule encoding a PHI-4 polypeptide is a polynucleotide having a nucleotide sequence encoding a polypeptide comprising an

amino acid sequence having at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or greater sequence identity to the amino acid sequence of SEQ ID NO: 35, wherein the polypeptide has pesticidal activity. In some embodiments the nucleic acid molecule encoding a PHI-4 polypeptide is a polynucleotide having a nucleotide sequence encoding a polypeptide comprising an amino acid sequence having 80% and 95% identity, to the amino acid sequence of SEQ ID NO: 35, wherein the polypeptide has pesticidal activity.

In some embodiments the nucleic acid molecule encoding a PHI-4 polypeptide is a polynucleotide having a nucleotide sequence encoding a polypeptide comprising an amino acid sequence having at least 80% identity, to any one of the amino acid sequence of SEQ ID NOs: 51-1162 SEQ ID NOs: 1518-1526 wherein the polypeptide has pesticidal activity.

In some embodiments the nucleic acid molecule encoding a PHI-4 polypeptide is a polynucleotide having a nucleotide sequence encoding a polypeptide comprising an amino acid sequence of SEQ ID NO: 3, wherein Xaa at position 2 is Ala or Arg; Xaa at position 9 is Gln, Lys or Glu; Xaa at position 14 is Pro or Ala; Xaa at position 16 is Val or Asp; Xaa at position 19 is Met or Leu; Xaa at position 22 is Gly or Ser; Xaa at position 24 is Asp, Asn or Gln; Xaa at position 36 is Leu or Met; Xaa at position 42 is Asp, Asn or Gln; Xaa at position 43 is Phe or Glu; Xaa at position 46 is Glu, Asp, Asn or Gly; Xaa at position 50 is Ile or Val; Xaa at position 51 is Glu or Gln; Xaa at position 55 is Arg or Lys; Xaa at position 56 is Ser or Thr; Xaa at position 57 is Tyr or Phe; Xaa at position 58 is Thr or Ser; Xaa at position 61 is Arg, Lys or Glu; Xaa at position 73 is Phe or Tyr; Xaa at position 74 is Lys, Glu, Gly, Arg, Met, Leu, His or Asp; Xaa at position 76 is Asp or Gln; Xaa at position 79 is Lys or Glu; Xaa at position 80 is Glu or Ser; Xaa at position 82 is Glu, Ile, Leu, Tyr or Gln; Xaa at position 83 is Glu or Gln; Xaa at position 84 is Tyr or Phe; Xaa at position 86 is Glu or Gln; Xaa at position 87 is Lys or Gln; Xaa at position 88 is Met, Ile or Leu; Xaa at position 90 is Gln or Glu; Xaa at position 94 is Val or Ile; Xaa at position 97 is Arg, Asn, Asp, Glu, Gln, Gly, Ser, Ile, Phe, His, Lys, Thr, Asn, Tyr, Trp, Pro, Cys, Ala, Met, Val or Leu; Xaa at position 98 is Tyr or Phe; Xaa at position 99 is Lys, Leu, Tyr, Ile, Met, Phe, Cys, Val or Asn; Xaa at position 103 is Ala or Gly; Xaa at position 105 is Leu or Ile; Xaa at position 109 is Phe, Lys, Gly, Met, Ser, Asp, Asn, Glu, Cys, Ala or Arg; Xaa at position 112 is Thr or Ser; Xaa at position 113 is Asp, Glu or Met; Xaa at position 117 is Thr or Ser; Xaa at position 121 is Tyr or Phe; Xaa at position 127 is Ala or Thr; Xaa at position 142 is Arg or Glu; Xaa at position 146 is Arg or Gln; Xaa at position 147 is Arg, Glu or Gln; Xaa at position 148 is Asp, Phe, Pro, Val, Glu, His, Trp, Ala, Arg, Leu, Ser, Gln or Gly; Xaa at position 149 is Phe or Val; Xaa at position 150 is Arg, Gln,

Glu or Asn; Xaa at position 151 is Asp, Ser, Ala, Asn, Trp, Val, Gln, Cys, Met, Leu, Arg or
 Glu; Xaa at position 153 is Leu or Ile; Xaa at position 154 is Asn or Asp; Xaa at position
 155 is Asn or Lys; Xaa at position 159 is Pro or Asp; Xaa at position 162 is Glu, Asp, Gln,
 Asn or Leu; Xaa at position 165 is Lys, Glu, Gln, Pro, Thr, Ala, Leu, Gly, Asp, Val, His, Ile,
 5 Met, Trp, Phe, Tyr or Arg; Xaa at position 166 is Arg or Gln; Xaa at position 167 is Tyr,
 Trp or Cys; Xaa at position 170 is Tyr or His; Xaa at position 171 is Tyr or Phe; Xaa at
 position 172 is Ile, Leu or Val; Xaa at position 173 is Ser or Ala; Xaa at position 174 is
 Glu, Gln, Asn, Lys, Val or Ser; Xaa at position 182 is Asp or Gln; Xaa at position 183 is
 Tyr or Val; Xaa at position 184 is Ser or Thr; Xaa at position 185 is Ala or Ser; Xaa at
 10 position 189 is Thr, Lys or Ile; Xaa at position 191 is Lys or Gln; Xaa at position 193 is
 Asp or Asn; Xaa at position 196 is Gln, Lys, Asn, Asp, Glu, Ala, Ile or Arg; Xaa at position
 202 is Ala or Val; Xaa at position 203 is Glu, Thr or His; Xaa at position 204 is Met or Ala;
 Xaa at position 206 is Tyr or Phe; Xaa at position 207 is Lys or Gln; Xaa at position 209 is
 Leu or Pro; Xaa at position 210 is Val or Ile; Xaa at position 214 is Lys, Ser or Gln; Xaa at
 15 position 216 is Glu, Gln, Phe, Val, Tyr or Arg; Xaa at position 220 is Glu, His, Asp, Thr,
 Tyr, Val, Ser, Gln, Arg, Trp, Met, Ala, Phe, Ile, Leu, Cys or Asn; Xaa at position 229 is Arg
 or Glu; Xaa at position 230 is Ser or Glu; Xaa at position 231 is Asn or Ser; Xaa at
 position 236 is Leu or Pro; Xaa at position 245 is Met or Leu; Xaa at position 247 is Asp or
 Tyr; Xaa at position 256 is Gln, Lys or Glu; Xaa at position 257 is Gln, Ile, Glu, Cys, Ser,
 20 His, Trp or Met; Xaa at position 261 is Gln, Glu, Lys or Ala; Xaa at position 264 is Glu or
 Gln; Xaa at position 268 is Asp or Asn; Xaa at position 276 is Ser or Ala; Xaa at position
 278 is Glu, Asn or Gln; Xaa at position 281 is Gln, Lys or Glu; Xaa at position 282 is Pro
 or Gly; Xaa at position 284 is Trp or Arg; Xaa at position 287 is Ala or Cys; Xaa at position
 289 is Lys, Leu, Val, Pro, Glu, Gln, Tyr, Thr, Asp, Phe, Ser, Met, Arg, Trp, Ile, His, Asn,
 25 Cys, Gly or Ala; Xaa at position 291 is Glu or Gln; Xaa at position 292 is Arg or Gln; Xaa
 at position 293 is Arg, Glu or Gln; Xaa at position 294 is Val or Ala; Xaa at position 296 is
 Leu or Ile; Xaa at position 297 is Glu or Gln; Xaa at position 298 is Asp or Gln; Xaa at
 position 300 is Phe or Tyr; Xaa at position 302 is Glu or Gln; Xaa at position 303 is Phe or
 Tyr; Xaa at position 305 is Lys, Gln, Ala, Ile, Met, Asn, Thr or Val; Xaa at position 306 is
 30 Gln or Lys; Xaa at position 309 is Gln, Lys or Glu; Xaa at position 313 is Lys, Gln or Arg;
 Xaa at position 316 is Lys or Gln; Xaa at position 328 is Lys, Glu or Gln; Xaa at position
 331 is Glu, Asn or Gln; Xaa at position 333 is Ser, Arg, Gly, Lys, Val, Asn, Ala, His, Gln,
 Thr, Asp, Ile, Leu, Cys or Glu; Xaa at position 334 is Gly, Arg, Lys, Ile or Trp; Xaa at
 position 335 is Ser or Ala; Xaa at position 336 is Gly or Ala; Xaa at position 337 is Ala, Val
 35 or Gly; Xaa at position 338 is Ser, His, Val, Lys, Ala, Gly, Thr, Ile, Glu, Met, Arg, Pro, Asp,
 Asn or Leu; Xaa at position 339 is Glu, Asn, Gln, Ile, Pro, Met, Ser, Ala, Cys, Phe, Val,

Leu, Asp, Trp, His or Arg; Xaa at position 341 is Leu or Val; Xaa at position 342 is Ala,
 Ser or Val; Xaa at position 343 is Val or Ile; Xaa at position 344 is Phe or Trp; Xaa at
 position 345 is Asn or His; Xaa at position 346 is Pro or Ala; Xaa at position 350 is Asn or
 Ser; Xaa at position 351 is Gly or Val; Xaa at position 354 is Met or Leu; Xaa at position
 5 355 is Val, Ile or Leu; Xaa at position 359 is Gly or Ala; Xaa at position 362 is Asn or Ser;
 Xaa at position 364 is Ala or Ser; Xaa at position 371 is Ala, Gly or Thr; Xaa at position
 374 is Phe or Ile; Xaa at position 375 is Lys or Arg; Xaa at position 380 is Leu or Gly; Xaa
 at position 382 is Val, Asp or Leu; Xaa at position 383 is Leu, Ile or Val; Xaa at position
 384 is Lys, Ala or Gly; Xaa at position 385 is Ala or Gly; Xaa at position 389 is Trp or Tyr;
 10 Xaa at position 391 is Arg, Leu, Glu, Gln, Asp or His; Xaa at position 395 is Asp or Cys;
 Xaa at position 396 is Ala, Leu, Lys, Asn, Gly, Ile, Met, Arg, Tyr, Gln, His or Thr; Xaa at
 position 397 is Gly, Arg or Ala; Xaa at position 398 is Ser, Gln or Cys; Xaa at position 401
 is Ser, His, Pro, Gly, Lys, Val, Arg, Ile, Asn, Phe, Thr, Ala, Asp, Met, Gln or Glu; Xaa at
 position 402 is Lys, Phe, His, Arg, Trp, Gly, Asn, Leu, Tyr, Thr, Val, Met, Pro or Ala; Xaa
 15 at position 403 is Asp, Tyr, Trp, Phe or Glu; Xaa at position 405 is Ala or Ser; Xaa at
 position 409 is Ala or Pro; Xaa at position 410 is Ile or Val; Xaa at position 411 is Pro or
 Ala; Xaa at position 412 is Pro or Ala; Xaa at position 416 is Arg, Glu or Gln; Xaa at
 position 417 is Ala, Ser or Cys; Xaa at position 418 is Leu or Met; Xaa at position 422 is
 Met or Val; Xaa at position 426 is Thr or Ser; Xaa at position 436 is Asp or Lys; Xaa at
 20 position 437 is Tyr or Val; Xaa at position 438 is Val or Arg; Xaa at position 440 is Val or
 Leu; Xaa at position 442 is Gln, Lys or Glu; Xaa at position 445 is Cys, Leu or Thr; Xaa at
 position 447 is Asp, Lys, Tyr, Ser, Glu, Ile, Gly, Pro, Leu, Phe, Trp or Thr; Xaa at position
 448 is Val or Ala; Xaa at position 449 is Gln or Glu; Xaa at position 452 is Gln, Lys or Glu,
 Ala; Xaa at position 453 is Asn or Asp; Xaa at position 454 is Arg, Tyr, Met, Ser, Val, Ile,
 25 Lys, Phe, Trp, Gln, Gly, His, Asp, Leu, Thr, Pro or Asn; Xaa at position 455 is Val or Ile;
 Xaa at position 457 is Trp or Asn; Xaa at position 459 is Lys, Met, Val, Trp, Gln, Ile, Thr,
 Ser, His, Cys, Tyr, Pro, Asn, Ala, Arg or Glu; Xaa at position 460 is Gly or Ala; Xaa at
 position 461 is Thr or Ser; Xaa at position 462 is Gly or Ala; Xaa at position 463 is Ala,
 Ser or Gly; Xaa at position 464 is Arg, Gly, His, Gln, Thr, Phe, Ala, Asp, Ser or Lys; Xaa at
 30 position 465 is Lys, Asn, Val, Met, Pro, Gly, Arg, Thr, His, Cys, Trp, Phe or Leu; Xaa at
 position 466 is Asp or Arg; Xaa at position 471 is Gln, Lys, Glu or Met; Xaa at position 472
 is Pro or Ser; Xaa at position 497 is Asp or Gln; Xaa at position 499 is Glu or Gln; Xaa at
 position 500 is Arg, Gln or Lys; Xaa at position 502 is Arg, Glu or Gln; Xaa at position 509
 is Lys, Gln, Glu or Ala; Xaa at position 517 is Gln, Cys, Asn, Val or Pro; Xaa at position
 35 518 is Glu or Gln; Xaa at position 520 is Lys, Gln, Glu, His or Ala; Xaa at position 525 is
 Gln or Lys; and Xaa at position 527 is Gln, Lys, Pro, Cys, Glu, Ser, His, Phe or Trp; and

having one or more amino acid substitutions at positions designated as Xaa in SEQ ID NO: 3 and wherein the PHI-4 polypeptide has increased insecticidal activity compared to the polypeptide of SEQ ID NO: 35.

In some embodiments the nucleic acid molecule encoding a PHI-4 polypeptide is a polynucleotide having a nucleotide sequence encoding a polypeptide comprising an amino acid sequence of SEQ ID NO: 4, wherein Xaa at position 2 is Ala or Arg; Xaa at position 24 is Asp or Asn; Xaa at position 42 is Asp or Asn; Xaa at position 43 is Phe or Glu; Xaa at position 46 is Glu or Asn; Xaa at position 74 is Lys, Glu or Gly; Xaa at position 79 is Lys or Glu; Xaa at position 82 is Glu, Ile, Leu or Tyr; Xaa at position 97 is Arg, Asn, Asp, Glu, Gln, Gly, Ser, Ile, Phe, His, Lys, Thr, Asn, Tyr, Trp, Pro, Cys, Ala, Met, Val or Leu; Xaa at position 98 is Tyr or Phe; Xaa at position 99 is Lys, Leu, Tyr, Ile or Met; Xaa at position 109 is Phe, Lys, Gly, Met, Ser, Asp or Asn; Xaa at position 147 is Arg or Glu; Xaa at position 148 is Asp, Phe or Pro; Xaa at position 150 is Arg, Gln, Glu or Asn; Xaa at position 151 is Asp, Ser, Ala or Asn; Xaa at position 153 is Leu or Ile; Xaa at position 162 is Glu, Asp, Gln, Asn or Leu; Xaa at position 165 is Lys, Glu or Gln; Xaa at position 166 is Arg or Gln; Xaa at position 171 is Tyr or Phe; Xaa at position 174 is Glu, Gln, Asn, Lys, Val or Ser; Xaa at position 182 is Asp or Gln; Xaa at position 196 is Gln, Lys, Asn or Asp; Xaa at position 203 is Glu, Thr or His; Xaa at position 206 is Tyr or Phe; Xaa at position 216 is Glu or Gln; Xaa at position 220 is Glu, His, Asp, Thr, Tyr, Val, Ser or Gln; Xaa at position 247 is Asp or Tyr; Xaa at position 256 is Gln or Lys; Xaa at position 257 is Gln or Ile; Xaa at position 261 is Gln, Glu, Lys or Ala; Xaa at position 278 is Glu or Asn; Xaa at position 281 is Gln, Lys or Glu; Xaa at position 289 is Lys, Leu, Val, Pro, Glu, Gln, Tyr, Thr or Asp; Xaa at position 293 is Arg, Glu or Gln; Xaa at position 313 is Lys or Gln; Xaa at position 328 is Lys, Glu or Gln; Xaa at position 333 is Ser, Gly, Lys, Val or Asn; Xaa at position 334 is Gly, Arg, Lys or Ile; Xaa at position 336 is Gly or Ala; Xaa at position 338 is Ser, His, Val, Lys or Ala; Xaa at position 339 is Glu, Asn, Ile or Pro; Xaa at position 343 is Val or Ile; Xaa at position 346 is Pro or Ala; Xaa at position 355 is Val or Ile; Xaa at position 359 is Gly or Ala; Xaa at position 391 is Arg, Leu, Glu, Gln, Asp or His; Xaa at position 396 is Ala, Leu, Lys, Asn, Gly or Thr; Xaa at position 401 is Ser, His, Pro, Gly, Lys, Val or Arg; Xaa at position 402 is Lys, Phe, His, Arg, Gly, Trp, Thr, Asn, Tyr or Met; Xaa at position 403 is Asp or Tyr; Xaa at position 411 is Pro or Ala; Xaa at position 412 is Pro or Ala; Xaa at position 416 is Arg or Glu; Xaa at position 417 is Ala or Ser; Xaa at position 418 is Leu or Met; Xaa at position 426 is Thr or Ser; Xaa at position 440 is Val or Leu; Xaa at position 447 is Asp, Lys, Tyr, Ser, Glu or Ile; Xaa at position 452 is Gln, Lys or Glu; Xaa at position 454 is Arg, Tyr, Met, Ser, Val, Ile, Lys, Phe, Trp or Gln; Xaa at position 455 is Val or Ile; Xaa at position 459 is Lys, Met, Val, Trp, Gln, Ile or Tyr; Xaa at

position 461 is Thr or Ser; Xaa at position 462 is Gly or Ala; Xaa at position 463 is Ala or Ser; Xaa at position 464 is Arg, Gly, His, Ala, Asp, Ser or Lys; Xaa at position 465 is Lys, Asn, Val, Met, Pro, Gly or Arg; Xaa at position 471 is Gln, Lys, Glu or Met; Xaa at position 472 is Pro or Ser; Xaa at position 500 is Arg or Gln; Xaa at position 509 is Lys, Gln or Ala; Xaa at position 520 is Lys, Gln, Glu, His or Ala; and Xaa at position 527 is Gln, Lys, Pro, Cys or Glu; and having one or more amino acid substitutions at positions designated as Xaa in SEQ ID NO: 4 and wherein the PHI-4 polypeptide has increased insecticidal activity compared to the polypeptide of SEQ ID NO: 35.

In some embodiments exemplary nucleic acid molecules encode a PHI-4 of any one of SEQ ID NOs: 51-1162 and SEQ ID NOs: 1518-1526 as well as amino acid substitutions, amino acid deletions, insertions and fragments thereof and combinations thereof.

In some embodiments the nucleic acid molecules encode a PHI-4 polypeptide of Table 3, Table 4, Table 5, Table 6, Table 7, Table 8, and Table 9 and combinations of the amino acid substitutions thereof and amino acid deletions and/or insertions thereof.

Also provided are nucleic acid molecules that encode transcription and/or translation products that are subsequently spliced to ultimately produce functional PHI-4. Splicing can be accomplished in vitro or in vivo, and can involve cis- or trans-splicing. The substrate for splicing can be polynucleotides (e.g., RNA transcripts) or polypeptides. An example of cis-splicing of a polynucleotide is where an intron inserted into a coding sequence is removed and the two flanking exon regions are spliced to generate a PHI-4 encoding sequence. An example of trans splicing would be where a polynucleotide is encrypted by separating the coding sequence into two or more fragments that can be separately transcribed and then spliced to form the full-length pesticidal encoding sequence. The use of a splicing enhancer sequence, which can be introduced into a construct, can facilitate splicing either in cis or trans-splicing of polypeptides (US Patent Numbers 6,365,377 and 6,531,316). Thus, in some embodiments the polynucleotides do not directly encode a full-length PHI-4, but rather encode a fragment or fragments of a PHI-4. These polynucleotides can be used to express a functional PHI-4 through a mechanism involving splicing, where splicing can occur at the level of polynucleotide (e.g., intron/exon) and/or polypeptide (e.g., intein/extein). This can be useful, for example, in controlling expression of pesticidal activity, since functional pesticidal polypeptide will only be expressed if all required fragments are expressed in an environment that permits splicing processes to generate functional product. In another example, introduction of one or more insertion sequences into a polynucleotide can facilitate recombination with a low homology polynucleotide; use of an intron or intein for the insertion sequence

facilitates the removal of the intervening sequence, thereby restoring function of the encoded variant.

Nucleic acid molecules that are fragments of these nucleic acid sequences encoding PHI-4 are also encompassed by the embodiments. By “fragment” is intended a portion of the nucleic acid sequence encoding a PHI-4. A fragment of a nucleic acid sequence may encode a biologically active portion of a PHI-4 polypeptide or it may be a fragment that can be used as a hybridization probe or PCR primer using methods disclosed below. Nucleic acid molecules that are fragments of a nucleic acid sequence encoding a PHI-4 comprise at least about 50, 100, 200, 300, 400, 500, 600 or 700, contiguous nucleotides or up to the number of nucleotides present in a full-length nucleic acid sequence encoding a PHI-4 disclosed herein, depending upon the intended use. By “contiguous” nucleotides is intended nucleotide residues that are immediately adjacent to one another. Fragments of the nucleic acid sequences of the embodiments will encode protein fragments that retain the biological activity of the PHI-4 and, hence, retain insecticidal activity. By “retains activity” is intended that the fragment will encode a polypeptide having at least about 30%, at least about 50%, at least about 70%, 80%, 90%, 95% or higher of the insecticidal activity of the full-length PHI-4. In one embodiment, the insecticidal activity is Lepodoptera activity. In another embodiment, the insecticidal activity is Hemiptera activity.

In some embodiments a fragment of a nucleic acid sequence encoding a PHI-4 encoding a biologically active portion of a protein will encode at least about 15, 25, 30, 50, 75, 100, 125, 150, 175, 200 or 250, contiguous amino acids or up to the total number of amino acids present in a full-length PHI-4 of the embodiments. In some embodiments, the fragment is an N-terminal or a C-terminal truncation of at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25 or more amino acids relative to SEQ ID NO: 35, SEQ ID NOs: 51-1162, SEQ ID NOs: 1518-1526 or variants thereof, e.g., by proteolysis, insertion of a start codon, deletion of the codons encoding the deleted amino acids with the concomitant insertion of a stop codon or by insertion of a stop codon in the coding sequence. In some embodiments, the fragments encompassed herein result from the removal of the N-terminal 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34 or more amino acids relative to SEQ ID NO: 35, SEQ ID NOs: 51-1162, SEQ ID NOs: 1518-1526 or variants thereof, e.g., by proteolysis or by insertion of a start codon in the coding sequence.

In some embodiments the PHI-4 are encoded by a nucleic acid sequence sufficiently identical to the nucleic acid sequence of SEQ ID NO: 1, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NOS: 24-30. By “sufficiently identical” is intended an amino acid or

nucleic acid sequence that has at least about 60% or 65% sequence identity, about 70% or 75% sequence identity, about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or greater sequence identity compared to a reference sequence using one of the alignment programs described herein using standard parameters. In some embodiments the sequence homology identity is against the full length sequence of the polynucleotide encoding a PHI-4 or against the full length sequence of a PHI-4 polypeptide. In some embodiments the polynucleotide encoding the PHI-4 has at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or greater sequence identity compared to SEQ ID NO: 1, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NOS: 24-30. One of skill in the art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleic acid sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning, and the like.

To determine the percent identity of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., $\text{percent identity} = \frac{\text{number of identical positions}}{\text{total number of positions (e.g., overlapping positions)}} \times 100$). In one embodiment, the two sequences are the same length. In another embodiment, the comparison is across the entirety of the reference sequence (e.g., across the entirety of SEQ ID NO: 35 or across the entirety of one of SEQ ID NO: 1, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NOS: 24-30). The percent identity between two sequences can be determined using techniques similar to those described below, with or without allowing gaps. In calculating percent identity, typically exact matches are counted.

The determination of percent identity between two sequences can be accomplished using a mathematical algorithm. A non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, (1990) *Proc. Natl. Acad. Sci. USA* 87:2264, modified as in Karlin and Altschul, (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877. Such an algorithm is incorporated into the BLASTN and BLASTX programs of Altschul, *et al.*, (1990) *J. Mol. Biol.* 215:403. BLAST nucleotide searches can be performed with the BLASTN program, score=100, wordlength=12, to obtain nucleic acid sequences homologous to pesticidal-like nucleic acid molecules. BLAST protein searches can be performed with the BLASTX program, score=50, wordlength=3, to obtain amino acid sequences homologous to pesticidal protein molecules. To obtain gapped alignments for comparison purposes, Gapped

BLAST (in BLAST 2.0) can be utilized as described in Altschul, *et al.*, (1997) *Nucleic Acids Res.* 25:3389. Alternatively, PSI-Blast can be used to perform an iterated search that detects distant relationships between molecules. See, Altschul, *et al.*, (1997) *supra*. When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters
5 of the respective programs (e.g., BLASTX and BLASTN) can be used. Alignment may also be performed manually by inspection.

Another non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the ClustalW algorithm (Higgins, *et al.*, (1994) *Nucleic Acids Res.* 22:4673-4680). ClustalW compares sequences and aligns the entirety of the amino
10 acid or DNA sequence and thus can provide data about the sequence conservation of the entire amino acid sequence. The ClustalW algorithm is used in several commercially available DNA/amino acid analysis software packages, such as the ALIGNX® module of the Vector NTI® Program Suite (Invitrogen Corporation, Carlsbad, Calif.). After alignment of amino acid sequences with ClustalW, the percent amino acid identity can be assessed.
15 A non-limiting example of a software program useful for analysis of ClustalW alignments is GENEDOC™. GENEDOC™ (Karl Nicholas) allows assessment of amino acid (or DNA) similarity and identity between multiple proteins. Another non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, (1988) *CABIOS* 4:11-17. Such an algorithm is incorporated into the ALIGN
20 program (version 2.0), which is part of the GCG Wisconsin Genetics Software Package, Version 10 (available from Accelrys, Inc., 9685 Scranton Rd., San Diego, Calif., USA). When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4 can be used.

Another non-limiting example of a mathematical algorithm utilized for the
25 comparison of sequences is the algorithm of Needleman and Wunsch, (1970) *J. Mol. Biol.* 48(3):443-453, used GAP Version 10 software to determine sequence identity or similarity using the following default parameters: % identity and % similarity for a nucleic acid sequence using GAP Weight of 50 and Length Weight of 3 and the nwsgapdna.cmpii scoring matrix; % identity or % similarity for an amino acid sequence using GAP weight of
30 8 and length weight of 2, and the BLOSUM62 scoring program. Equivalent programs may also be used. By “equivalent program” is intended any sequence comparison program that, for any two sequences in question, generates an alignment having identical nucleotide residue matches and an identical percent sequence identity when compared to the corresponding alignment generated by GAP Version 10.

35 The embodiments also encompass nucleic acid molecules encoding variants of PHI-4 polypeptide. “Variants” of the PHI-4 polypeptide encoding nucleic acid sequences

include those sequences that encode the PHI-4 polypeptides disclosed herein but that differ conservatively because of the degeneracy of the genetic code as well as those that are sufficiently identical as discussed above. Naturally occurring allelic variants can be identified with the use of well-known molecular biology techniques, such as polymerase chain reaction (PCR) and hybridization techniques as outlined below. Variant nucleic acid sequences also include synthetically derived nucleic acid sequences that have been generated, for example, by using site-directed mutagenesis but which still encode the PHI-4 polypeptides disclosed as discussed below.

The skilled artisan will further appreciate that changes can be introduced by mutation of the nucleic acid sequences thereby leading to changes in the amino acid sequence of the encoded PHI-4 polypeptides, without altering the biological activity of the proteins. Thus, variant nucleic acid molecules can be created by introducing one or more nucleotide substitutions, nucleotide additions and/or nucleotide deletions into the corresponding nucleic acid sequence disclosed herein, such that one or more amino acid substitutions, amino acid additions or amino acid deletions are introduced into the encoded protein. Mutations can be introduced by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Such variant nucleic acid sequences are also encompassed by the present disclosure.

Alternatively, variant nucleic acid sequences can be made by introducing mutations randomly along all or part of the coding sequence, such as by saturation mutagenesis and the resultant mutants can be screened for ability to confer pesticidal activity to identify mutants that retain activity. Following mutagenesis, the encoded protein can be expressed recombinantly, and the activity of the protein can be determined using standard assay techniques.

The polynucleotides of the disclosure and fragments thereof are optionally used as substrates for a variety of recombination and recursive recombination reactions, in addition to standard cloning methods as set forth in, e.g., Ausubel, Berger and Sambrook, i.e., to produce additional pesticidal polypeptide homologues and fragments thereof with desired properties. A variety of such reactions are known, including those developed by the inventors and their co-workers. Methods for producing a variant of any nucleic acid listed herein comprising recursively recombining such polynucleotide with a second (or more) polynucleotide, thus forming a library of variant polynucleotides are also embodiments of the disclosure, as are the libraries produced, the cells comprising the libraries, and any recombinant polynucleotide produced by such methods. Additionally, such methods optionally comprise selecting a variant polynucleotide from such libraries

based on pesticidal activity, as is wherein such recursive recombination is done in vitro or in vivo.

A variety of diversity generating protocols, including nucleic acid recursive recombination protocols are available and fully described in the art. The procedures can be used separately, and/or in combination to produce one or more variants of a nucleic acid or set of nucleic acids, as well as variants of encoded proteins. Individually and collectively, these procedures provide robust, widely applicable ways of generating diversified nucleic acids and sets of nucleic acids (including, e.g., nucleic acid libraries) useful, e.g., for the engineering or rapid evolution of nucleic acids, proteins, pathways, cells and/or organisms with new and/or improved characteristics.

While distinctions and classifications are made in the course of the ensuing discussion for clarity, it will be appreciated that the techniques are often not mutually exclusive. Indeed, the various methods can be used singly or in combination, in parallel or in series, to access diverse sequence variants.

The result of any of the diversity generating procedures described herein can be the generation of one or more nucleic acids, which can be selected or screened for nucleic acids with or which confer desirable properties or that encode proteins with or which confer desirable properties. Following diversification by one or more of the methods herein or otherwise available to one of skill, any nucleic acids that are produced can be selected for a desired activity or property, e.g. pesticidal activity or such activity at a desired pH, etc. This can include identifying any activity that can be detected, for example, in an automated or automatable format, by any of the assays in the art, see, e.g., discussion of screening of insecticidal activity, *infra*. A variety of related (or even unrelated) properties can be evaluated, in serial or in parallel, at the discretion of the practitioner.

Descriptions of a variety of diversity generating procedures for generating modified nucleic acid sequences, e.g., those coding for polypeptides having pesticidal activity or fragments thereof, are found in the following publications and the references cited therein: Soong, *et al.*, (2000) *Nat Genet* 25(4):436-439; Stemmer, *et al.*, (1999) *Tumor Targeting* 4:1-4; Ness *et al.* (1999) *Nat Biotechnol* 17:893-896; Chang *et al.* (1999) *Nat Biotechnol* 17:793-797; Minshull and Stemmer, (1999) *Curr Opin Chem Biol* 3:284-290; Christians, *et al.*, (1999) *Nat Biotechnol* 17:259-264; Crameri, *et al.*, (1998) *Nature* 391:288-291; Crameri, *et al.*, (1997) *Nat Biotechnol* 15:436-438; Zhang, *et al.*, (1997) *PNAS USA* 94:4504-4509; Patten, *et al.*, (1997) *Curr Opin Biotechnol* 8:724-733; Crameri, *et al.*, (1996) *Nat Med* 2:100-103; Crameri, *et al.*, (1996) *Nat Biotechnol* 14:315-319; Gates, *et al.*, (1996) *J Mol Biol* 255:373-386; Stemmer, (1996) "Sexual PCR and Assembly PCR" In:

The Encyclopedia of Molecular Biology. VCH Publishers, New York. pp. 447-457; Crameri and Stemmer, (1995) *BioTechniques* 18:194-195; Stemmer, *et al.*, (1995) *Gene*, 164:49-53; Stemmer, (1995) *Science* 270:1510; Stemmer, (1995) *Bio/Technology* 13:549-553; Stemmer, (1994) *Nature* 370:389-391 and Stemmer, (1994) *PNAS USA* 91:10747-10751.

5 Mutational methods of generating diversity include, for example, site-directed mutagenesis (Ling, *et al.*, (1997) *Anal Biochem* 254(2):157-178; Dale, *et al.*, (1996) *Methods Mol Biol* 57:369-374; Smith, (1985) *Ann Rev Genet* 19:423-462; Botstein and Shortle, (1985) *Science* 229:1193-1201; Carter, (1986) *Biochem J* 237:1-7 and Kunkel, (1987) "The efficiency of oligonucleotide directed mutagenesis" in *Nucleic Acids &*
 10 *Molecular Biology* (Eckstein and Lilley, eds., Springer Verlag, Berlin)); mutagenesis using uracil containing templates (Kunkel, (1985) *PNAS USA* 82:488-492; Kunkel, *et al.*, (1987) *Methods Enzymol* 154:367-382 and Bass, *et al.*, (1988) *Science* 242:240-245); oligonucleotide-directed mutagenesis (Zoller and Smith, (1983) *Methods Enzymol* 100:468-500; Zoller and Smith, (1987) *Methods Enzymol* 154:329-350; Zoller and Smith, (1982) *Nucleic Acids Res* 10:6487-6500), phosphorothioate-modified DNA mutagenesis (Taylor, *et al.*, (1985) *Nucl Acids Res* 13:8749-8764; Taylor, *et al.*, (1985) *Nucl Acids Res* 13:8765-8787 (1985); Nakamaye and Eckstein (1986) *Nucl Acids Res* 14:9679-9698; Sayers, *et al.*, (1988) *Nucl Acids Res* 16:791-802 and Sayers, *et al.*, (1988) *Nucl Acids Res* 16: 803-814); mutagenesis using gapped duplex DNA (Kramer, *et al.*, (1984) *Nucl*
 15 *Acids Res* 12:9441-9456; Kramer and Fritz, (1987) *Methods Enzymol* 154:350-367; Kramer, *et al.*, (1988) *Nucl Acids Res* 16:7207 and Fritz, *et al.*, (1988) *Nucl Acids Res* 16:6987-6999).

Additional suitable methods include point mismatch repair (Kramer, *et al.*, (1984) *Cell* 38:879-887), mutagenesis using repair-deficient host strains (Carter, *et al.*, (1985) *Nucl Acids Res* 13:4431-4443 and Carter, (1987) *Methods in Enzymol* 154:382-403), deletion mutagenesis (Eghtedarzadeh and Henikoff, (1986) *Nucl Acids Res* 14:5115), restriction-selection and restriction-purification (Wells, *et al.*, (1986) *Phil Trans R Soc Lond A* 317:415-423), mutagenesis by total gene synthesis (Nambiar, *et al.*, (1984) *Science* 223:1299-1301; Sakamar and Khorana, (1988) *Nucl Acids Res* 14:6361-6372; Wells, *et al.*, (1985) *Gene* 34:315-323 and Grundström, *et al.*, (1985) *Nucl Acids Res* 13:3305-3316), double-strand break repair (Mandecki, (1986) *PNAS USA*, 83:7177-7181 and Arnold, (1993) *Curr Opin Biotech* 4:450-455). Additional details on many of the above methods can be found in *Methods Enzymol* Volume 154, which also describes useful controls for trouble-shooting problems with various mutagenesis methods.

35 Additional details regarding various diversity generating methods can be found in the following US Patents, PCT Publications and Applications and EPO Publications: US

Patent Number 5,723,323, US Patent Number 5,763,192, US Patent Number 5,814,476, US Patent Number 5,817,483, US Patent Number 5,824,514, US Patent Number 5,976,862, US Patent Number 5,605,793, US Patent Number 5,811,238, US Patent Number 5,830,721, US Patent Number 5,834,252, US Patent Number 5,837,458, WO
 5 1995/22625, WO 1996/33207, WO 1997/20078, WO 1997/35966, WO 1999/41402, WO 1999/41383, WO 1999/41369, WO 1999/41368, EP 752008, EP 0932670, WO 1999/23107, WO 1999/21979, WO 1998/31837, WO 1998/27230, WO 1998/27230, WO 2000/00632, WO 2000/09679, WO 1998/42832, WO 1999/29902, WO 1998/41653, WO 1998/41622, WO 1998/42727, WO 2000/18906, WO 2000/04190, WO 2000/42561, WO
 10 2000/42559, WO 2000/42560, WO 2001/23401, and PCT/US01/06775.

The nucleotide sequences of the embodiments can also be used to isolate corresponding sequences from other organisms, particularly other bacteria. In this manner, methods such as PCR, hybridization and the like can be used to identify such sequences based on their sequence homology to the sequences set forth herein.
 15 Sequences that are selected based on their sequence identity to the entire sequences set forth herein or to fragments thereof are encompassed by the embodiments. Such sequences include sequences that are orthologs of the disclosed sequences. The term "orthologs" refers to genes derived from a common ancestral gene and which are found in different species as a result of speciation. Genes found in different species are
 20 considered orthologs when their nucleotide sequences and/or their encoded protein sequences share substantial identity as defined elsewhere herein. Functions of orthologs are often highly conserved among species.

In a PCR approach, oligonucleotide primers can be designed for use in PCR reactions to amplify corresponding DNA sequences from cDNA or genomic DNA
 25 extracted from any organism of interest. Methods for designing PCR primers and PCR cloning are generally known in the art and are disclosed in Sambrook, *et al.*, (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York), hereinafter "Sambrook". See also, Innis, *et al.*, eds. (1990) *PCR Protocols: A Guide to Methods and Applications* (Academic Press, New York); Innis and
 30 Gelfand, eds. (1995) *PCR Strategies* (Academic Press, New York); and Innis and Gelfand, eds. (1999) *PCR Methods Manual* (Academic Press, New York). Known methods of PCR include, but are not limited to, methods using paired primers, nested primers, single specific primers, degenerate primers, gene-specific primers, vector-specific primers, partially-mismatched primers, and the like.

35 To identify potential PHI-4 polypeptides from bacterial collections, the bacterial cell lysates can be screened with antibodies generated against a PHI-4 polypeptide using

Western blotting and/or ELISA methods. This type of assays can be performed in a high throughput fashion. Positive samples can be further analyzed by various techniques such as antibody based protein purification and identification. Methods of generating antibodies are well known in the art as discussed infra.

5 Alternatively, mass spectrometry based protein identification method can be used to identify homologs of a PHI-4 polypeptide using protocols in the literatures (Patterson, (1998), 10(22):1-24, Current Protocol in Molecular Biology published by John Wiley & Son Inc). Specifically, LC-MS/MS based protein identification method is used to associate the MS data of given cell lysate or desired molecular weight enriched samples (excised from
10 SDS-PAGE gel of relevant molecular weight bands to a PHI-4 polypeptide) with sequence information of a PHI-4 polypeptide and its homologs. Any match in peptide sequences indicates the potential of having the homologous proteins in the samples. Additional techniques (protein purification and molecular biology) can be used to isolate the protein and identify the sequences of the homologs.

15 In hybridization methods, all or part of the pesticidal nucleic acid sequence can be used to screen cDNA or genomic libraries. Methods for construction of such cDNA and genomic libraries are generally known in the art and are disclosed in Sambrook and Russell, (2001), *supra*. The so-called hybridization probes may be genomic DNA fragments, cDNA fragments, RNA fragments or other oligonucleotides, and may be
20 labeled with a detectable group such as ³²P or any other detectable marker, such as other radioisotopes, a fluorescent compound, an enzyme or an enzyme co-factor. Probes for hybridization can be made by labeling synthetic oligonucleotides based on the known PHI-4 polypeptide-encoding nucleic acid sequence disclosed herein. Degenerate primers designed on the basis of conserved nucleotides or amino acid residues in the nucleic acid
25 sequence or encoded amino acid sequence can additionally be used. The probe typically comprises a region of nucleic acid sequence that hybridizes under stringent conditions to at least about 12, at least about 25, at least about 50, 75, 100, 125, 150, 175 or 200 consecutive nucleotides of nucleic acid sequence encoding a PHI-4 polypeptide of the disclosure or a fragment or variant thereof. Methods for the preparation of probes for
30 hybridization are generally known in the art and are disclosed in Sambrook and Russell, (2001), *supra*, herein incorporated by reference.

For example, an entire nucleic acid sequence, encoding a PHI-4 polypeptide, disclosed herein or one or more portions thereof, may be used as a probe capable of specifically hybridizing to corresponding nucleic acid sequences encoding PHI-4
35 polypeptide-like sequences and messenger RNAs. To achieve specific hybridization under a variety of conditions, such probes include sequences that are unique and are

preferably at least about 10 nucleotides in length or at least about 20 nucleotides in length. Such probes may be used to amplify corresponding pesticidal sequences from a chosen organism by PCR. This technique may be used to isolate additional coding sequences from a desired organism or as a diagnostic assay to determine the presence of coding sequences in an organism. Hybridization techniques include hybridization screening of plated DNA libraries (either plaques or colonies; see, for example, Sambrook, *et al.*, (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.).

Hybridization of such sequences may be carried out under stringent conditions. By “stringent conditions” or “stringent hybridization conditions” is intended conditions under which a probe will hybridize to its target sequence to a detectably greater degree than to other sequences (e.g., at least 2-fold over background). Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences that are 100% complementary to the probe can be identified (homologous probing). Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing). Generally, a probe is less than about 1000 nucleotides in length, preferably less than 500 nucleotides in length.

Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37°C., and a wash in 1× to 2×SSC (20×SSC=3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55°C. Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1.0 M NaCl, 1% SDS at 37°C., and a wash in 0.5× to 1×SSC at 55 to 60°C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37°C., and a wash in 0.1×SSC at 60 to 65°C. Optionally, wash buffers may comprise about 0.1% to about 1% SDS. Duration of hybridization is generally less than about 24 hours, usually about 4 to about 12 hours.

Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA-DNA hybrids, the T_m can be approximated from the equation of Meinkoth and Wahl, (1984) *Anal. Biochem.* 138:267-284: $T_m = 81.5^\circ\text{C} + 16.6 (\log M) + 0.41 (\% \text{ GC}) - 0.61 (\% \text{ form}) -$

500/L; where M is the molarity of monovalent cations, % GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. The T_m is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe. T_m is reduced by about 1°C for each 1% of mismatching; thus, T_m, hybridization, and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with ≥90% identity are sought, the T_m can be decreased 10°C. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3 or 4°C lower than the thermal melting point (T_m); moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9 or 10°C lower than the thermal melting point (T_m); low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15 or 20°C lower than the thermal melting point (T_m). Using the equation, hybridization and wash compositions, and desired T_m, those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45°C (aqueous solution) or 32°C (formamide solution), it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen, (1993) *Laboratory Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 (Elsevier, N.Y.); and Ausubel, *et al.*, eds. (1995) *Current Protocols in Molecular Biology*, Chapter 2 (Greene Publishing and Wiley-Interscience, New York). See, Sambrook, *et al.*, (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.).

Proteins and Variants and Fragments Thereof

PHI-4 polypeptides are encompassed by the disclosure. By “PHI-4 polypeptide” or “PHI-4 protein” as used herein interchangeably is intended a polypeptide that has increased insecticidal activity against one or more insect pests of the Lepidoptera and/or Coleoptera orders compared to the protein of SEQ ID NO: 35, and is sufficiently identical to the protein of SEQ ID NO: 35. A variety of PHI-4 polypeptides are contemplated.

The Western Corn Rootworm active protein AXMI-205 (SEQ ID NO: 35) encoded by the polynucleotide of SEQ ID NO: 34 was identified from the *Chromobacterium* Strain ATX 2024 (US20110023184). Synthetic genes encoding AXMI-205; AXMI-205 variants having a truncation of the last 10 and 20 amino acids from the C-terminus (SEQ ID NO:

36 and SEQ ID NO: 37); and alanine scanning at every other residue from residue 307-536 of AXMI-205 (SEQ ID NO: 35), with S307A, D315A, V317A, S349A, G351A, K353A, V355A, D395A, G399A, W407A, G419A, P435A, S443A, K465A, V467A, F483A, P487A, S495A, D497A, E499A, K509A, and 1513A identified as having WCRW activity; are disclosed in US20110023184. AXMI-205 variants evo 24 (E499A); evo25 (V467A - SEQ ID NO: 41); evo30 (V467L - SEQ ID NO: 42); PMlibl PoolIG2_p2al 1 (S468L - SEQ ID NO: 45); PMlibl PoolIG2_plcl (V467T - SEQ ID NO: 46), PMlibl PoolIG2_pla4 (R464N - SEQ ID NO: 47), evo34 (E86T - SEQ ID NO: 43) and evo35 (Q517R - SEQ ID NO: 44) having increased insecticidal activity are disclosed in WO2013/016617.

As used herein, the terms "protein," "peptide molecule" or "polypeptide" includes any molecule that comprises five or more amino acids. It is well known in the art that protein, peptide or polypeptide molecules may undergo modification, including post-translational modifications, such as, but not limited to, disulfide bond formation, glycosylation, phosphorylation or oligomerization. Thus, as used herein, the terms "protein," "peptide molecule" or "polypeptide" includes any protein that is modified by any biological or non-biological process. The terms "amino acid" and "amino acids" refer to all naturally occurring L-amino acids.

A "recombinant protein" is used to refer to a protein that is no longer in its natural environment, for example in vitro or in a recombinant bacterial or plant host cell. A PHI-4 polypeptide that is substantially free of cellular material includes preparations of protein having less than about 30%, 20%, 10% or 5% (by dry weight) of non-pesticidal protein (also referred to herein as a "contaminating protein").

By "improved activity" or "increased activity" is intended an increase of at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 100%, at least about 110%, at least about 120%, at least about 130%, at least about 140%, at least about 150%, at least about 160%, at least about 170%, at least about 180%, at least about 190%, at least about 200%, at least about 210% at least about 220%, at least about 230%, at least about 240%, at least about 250%, at least about 260%, at least about 270%, at least about 280%, at least about 290%, at least about 300%, at least about 310%, at least about 320%, at least about 330%, at least about 340%, at least about 350%, at least about 360%, at least about 370%, at least about 380%, at least about 390%, at least about 400%, at least about 410%, at least about 420%, at least about 430%, at least about 440%, at least about 450%, at least about 460%, at least about 470%, at least about 480%, at least about 490%, at least about 500%, at least

about 510%, at least about 520%, at least about 530%, at least about 540%, at least about 550%, at least about 560%, at least about 570%, at least about 580%, at least about 590%, at least about 600%, at least about 650%, at least about 700%, at least about 750%, at least about 800%, at least about 850%, at least about 900%, at least about 950%, at least about 1000% or higher or at least about 1.1-fold, at least about 1.2-fold, at least about 1.3-fold, at least about 1.4-fold or at least about 1.5-fold, at least about 1.6-fold, at least about 1.7-fold, at least about 1.8-fold, at least about 1.9-fold, at least about 2-fold, at least about 2.1-fold, at least about 2.2-fold, at least about 2.3-fold, at least about 2.4-fold, at least about 2.5-fold, at least about 2.6-fold, at least about 2.7-fold, at least about 2.8-fold, at least about 2.9-fold, at least about 3-fold, at least about 3.1-fold, at least about 3.2-fold, at least about 3.3-fold, at least about 3.4-fold, at least about 3.5-fold, at least about 3.6-fold, at least about 3.7-fold, at least about 3.8-fold, at least about 3.9-fold, at least about 4-fold, at least about 4.1-fold, at least about 4.2-fold, at least about 4.3-fold, at least about 4.4-fold, at least about 4.5-fold, at least about 4.6-fold, at least about 4.7-fold, at least about 4.8-fold, at least about 4.9-fold, at least about 5-fold, at least about 5.1-fold, at least about 5.2-fold, at least about 5.3-fold, at least about 5.4-fold, at least about 5.5-fold, at least about 5.6-fold, at least about 5.7-fold, at least about 5.8-fold, at least about 5.9-fold, at least about 6-fold, at least about 6.1-fold, at least about 6.2-fold, at least about 6.3-fold, at least about 6.4-fold, at least about 6.5-fold, at least about 6.6-fold, at least about 6.7-fold, at least about 6.8-fold, at least about 6.9-fold, at least about 7-fold, at least about 7.1-fold, at least about 7.2-fold, at least about 7.3-fold, at least about 7.4-fold, at least about 7.5-fold, at least about 7.6-fold, at least about 7.7-fold, at least about 7.8-fold, at least about 7.9-fold, at least about 8-fold, at least about 8.1-fold, at least about 8.2-fold, at least about 8.3-fold, at least about 8.4-fold, at least about 8.5-fold, at least about 8.6-fold, at least about 8.7-fold, at least about 8.8-fold, at least about 8.9-fold, at least about 9-fold, at least about 9.1-fold, at least about 9.2-fold, at least about 9.3-fold, at least about 9.4-fold, at least about 9.5-fold, at least about 9.6-fold, at least about 9.7-fold, at least about 9.8-fold, at least about 9.9-fold, at least about 10-fold or higher increase in the pesticidal activity of the variant protein relative to the pesticidal activity of AXMI-205 (SEQ ID NO: 35).

In some embodiments, the improvement consists of a decrease in the EC₅₀ of at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 100%, at least about 110%, at least about 120%, at least about 130%, at least about 140%, at least about 150%, at least about 160%, at least about 170%, at least about 180%, at least

about 190%, at least about 200%, at least about 210% at least about 220%, at least
about 230%, at least about 240%, at least about 250%, at least about 260%, at least
about 270%, at least about 280%, at least about 290%, at least about 300%, at least
about 310%, at least about 320%, at least about 330%, at least about 340%, at least
5 about 350%, at least about 360%, at least about 370%, at least about 380%, at least
about 390%, at least about 400%, at least about 410%, at least about 420%, at least
about 430%, at least about 440%, at least about 450%, at least about 460%, at least
about 470%, at least about 480%, at least about 490%, at least about 500%, at least
about 510%, at least about 520%, at least about 530%, at least about 540%, at least
10 about 550%, at least about 560%, at least about 570%, at least about 580%, at least
about 590%, at least about 600%, at least about 650%, at least about 700%, at least
about 750%, at least about 800%, at least about 850%, at least about 900%, at least
about 950%, at least about 1000% or higher or at least about 1.1-fold, at least about 1.2-
fold, at least about 1.3-fold, at least about 1.4-fold or at least about 1.5-fold, at least about
15 1.6-fold, at least about 1.7-fold, at least about 1.8-fold, at least about 1.9-fold, at least
about 2-fold, at least about 2.1-fold, at least about 2.2-fold, at least about 2.3-fold, at least
about 2.4-fold, at least about 2.5-fold, at least about 2.6-fold, at least about 2.7-fold, at
least about 2.8-fold, at least about 2.9-fold, at least about 3 –fold, at least about 3.1-fold,
at least about 3.2-fold, at least about 3.3-fold, at least about 3.4-fold, at least about 3.5-
20 fold, at least about 3.6-fold, at least about 3.7-fold, at least about 3.8-fold, at least about
3.9-fold, at least about 4-fold, at least about 4.1-fold, at least about 4.2-fold, at least about
4.3-fold, at least about 4.4-fold, at least about 4.5-fold, at least about 4.6-fold, at least
about 4.7-fold, at least about 4.8-fold, at least about 4.9-fold, at least about 5-fold, at least
about 5.1-fold, at least about 5.2-fold, at least about 5.3-fold, at least about 5.4-fold, at
25 least about 5.5-fold, at least about 5.6-fold, at least about 5.7-fold, at least about 5.8-fold,
at least about 5.9-fold, at least about 6-fold, at least about 6.1-fold, at least about 6.2-fold,
at least about 6.3-fold, at least about 6.4-fold, at least about 6.5-fold, at least about 6.6-
fold, at least about 6.7-fold, at least about 6.8-fold, at least about 6.9-fold, at least about
7-fold, at least about 7.1-fold, at least about 7.2-fold, at least about 7.3-fold, at least about
30 7.4-fold, at least about 7.5-fold, at least about 7.6-fold, at least about 7.7-fold, at least
about 7.8-fold, at least about 7.9-fold, at least about 8-fold, at least about 8.1-fold, at least
about 8.2-fold, at least about 8.3-fold, at least about 8.4-fold, at least about 8.5-fold, at
least about 8.6-fold, at least about 8.7-fold, at least about 8.8-fold, at least about 8.9-fold,
at least about 9-fold, at least about 9.1-fold, at least about 9.2-fold, at least about 9.3-fold,
35 at least about 9.4-fold, at least about 9.5-fold, at least about 9.6-fold, at least about 9.7-
fold, at least about 9.8-fold, at least about 9.9-fold, at least about 10-fold or greater

reduction in the EC50 of the PHI-4 polypeptide relative to the pesticidal activity of AXMI-205 (SEQ ID NO: 35).

In some embodiments the EC50 of the PHI-4 polypeptide is <100 ppm, <90 ppm, <80 ppm, <70 ppm, <60 ppm, <50 ppm, <45 ppm, <40 ppm, <35 ppm, <30 ppm, <25 ppm, <20 ppm, <19 ppm, <18 ppm, <17 ppm, <16 ppm, <15 ppm, <14 ppm, <13 ppm, <12 ppm, <11 ppm, <10 ppm, <9 ppm, <8 ppm, <7 ppm, <6 ppm, <5 ppm, <4 ppm, <3 ppm, <2 ppm, <1 ppm, <0.9 ppm, <0.8 ppm, <0.7 ppm, <0.6 ppm, <0.5 ppm, <0.4 ppm, <0.3 ppm, <0.2 ppm or <0.1 ppm.

In some embodiments, the improvement consists of an increase in the Mean FAE Index of at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 100%, at least about 110%, at least about 120%, at least about 130%, at least about 140%, at least about 150%, at least about 160%, at least about 170%, at least about 180%, at least about 190%, at least about 200%, at least about 210% at least about 220%, at least about 230%, at least about 240%, at least about 250%, at least about 260%, at least about 270%, at least about 280%, at least about 290%, at least about 300%, at least about 310%, at least about 320%, at least about 330%, at least about 340%, at least about 350%, at least about 360%, at least about 370%, at least about 380%, at least about 390%, at least about 400%, at least about 410%, at least about 420%, at least about 430%, at least about 440%, at least about 450%, at least about 460%, at least about 470%, at least about 480%, at least about 490%, at least about 500%, at least about 510%, at least about 520%, at least about 530%, at least about 540%, at least about 550%, at least about 560%, at least about 570%, at least about 580%, at least about 590%, at least about 600%, at least about 650%, at least about 700%, at least about 750%, at least about 800%, at least about 850%, at least about 900%, at least about 950%, at least about 1000% or higher or at least about 1.1-fold, at least about 1.2-fold, at least about 1.3-fold, at least about 1.4-fold or at least about 1.5-fold, at least about 1.6-fold, at least about 1.7-fold, at least about 1.8-fold, at least about 1.9-fold, at least about 2-fold, at least about 2.1-fold, at least about 2.2-fold, at least about 2.3-fold, at least about 2.4-fold, at least about 2.5-fold, at least about 2.6-fold, at least about 2.7-fold, at least about 2.8-fold, at least about 2.9-fold, at least about 3 –fold, at least about 3.1-fold, at least about 3.2-fold, at least about 3.3-fold, at least about 3.4-fold, at least about 3.5-fold, at least about 3.6-fold, at least about 3.7-fold, at least about 3.8-fold, at least about 3.9-fold, at least about 4-fold, at least about 4.1-fold, at least about 4.2-fold, at least about 4.3-fold, at least about 4.4-fold, at least about 4.5-fold, at least

about 4.6-fold, at least about 4.7-fold, at least about 4.8-fold, at least about 4.9-fold, at least about 5-fold, at least about 5.1-fold, at least about 5.2-fold, at least about 5.3-fold, at least about 5.4-fold, at least about 5.5-fold, at least about 5.6-fold, at least about 5.7-fold, at least about 5.8-fold, at least about 5.9-fold, at least about 6-fold, at least about 6.1-fold, at least about 6.2-fold, at least about 6.3-fold, at least about 6.4-fold, at least about 6.5-fold, at least about 6.6-fold, at least about 6.7-fold, at least about 6.8-fold, at least about 6.9-fold, at least about 7-fold, at least about 7.1-fold, at least about 7.2-fold, at least about 7.3-fold, at least about 7.4-fold, at least about 7.5-fold, at least about 7.6-fold, at least about 7.7-fold, at least about 7.8-fold, at least about 7.9-fold, at least about 8-fold, at least about 8.1-fold, at least about 8.2-fold, at least about 8.3-fold, at least about 8.4-fold, at least about 8.5-fold, at least about 8.6-fold, at least about 8.7-fold, at least about 8.8-fold, at least about 8.9-fold, at least about 9-fold, at least about 9.1-fold, at least about 9.2-fold, at least about 9.3-fold, at least about 9.4-fold, at least about 9.5-fold, at least about 9.6-fold, at least about 9.7-fold, at least about 9.8-fold, at least about 9.9-fold, at least about 10-fold or higher increase in the Mean FAE Index of the PHI-4 polypeptide relative to the pesticidal activity of AXMI-205 (SEQ ID NO: 35).

In some embodiments, the improvement consists of an increase in the Mean Deviation Score of at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 100%, at least about 110%, at least about 120%, at least about 130%, at least about 140%, at least about 150%, at least about 160%, at least about 170%, at least about 180%, at least about 190%, at least about 200%, at least about 210%, at least about 220%, at least about 230%, at least about 240%, at least about 250%, at least about 260%, at least about 270%, at least about 280%, at least about 290%, at least about 300%, at least about 310%, at least about 320%, at least about 330%, at least about 340%, at least about 350%, at least about 360%, at least about 370%, at least about 380%, at least about 390%, at least about 400%, at least about 410%, at least about 420%, at least about 430%, at least about 440%, at least about 450%, at least about 460%, at least about 470%, at least about 480%, at least about 490%, at least about 500%, at least about 510%, at least about 520%, at least about 530%, at least about 540%, at least about 550%, at least about 560%, at least about 570%, at least about 580%, at least about 590%, at least about 600%, at least about 650%, at least about 700%, at least about 750%, at least about 800%, at least about 850%, at least about 900%, at least about 950%, at least about 1000% or higher or at least about 1.1-fold, at least about 1.2-fold, at least about 1.3-fold, at least about 1.4-fold or at least about

1.5-fold, at least about 1.6-fold, at least about 1.7-fold, at least about 1.8-fold, at least about 1.9-fold, at least about 2-fold, at least about 2.1-fold, at least about 2.2-fold, at least about 2.3-fold, at least about 2.4-fold, at least about 2.5-fold, at least about 2.6-fold, at least about 2.7-fold, at least about 2.8-fold, at least about 2.9-fold, at least about 3 –fold, at least about 3.1-fold, at least about 3.2-fold, at least about 3.3-fold, at least about 3.4-fold, at least about 3.5-fold, at least about 3.6-fold, at least about 3.7-fold, at least about 3.8-fold, at least about 3.9-fold, at least about 4-fold, at least about 4.1-fold, at least about 4.2-fold, at least about 4.3-fold, at least about 4.4-fold, at least about 4.5-fold, at least about 4.6-fold, at least about 4.7-fold, at least about 4.8-fold, at least about 4.9-fold, at least about 5-fold, at least about 5.1-fold, at least about 5.2-fold, at least about 5.3-fold, at least about 5.4-fold, at least about 5.5-fold, at least about 5.6-fold, at least about 5.7-fold, at least about 5.8-fold, at least about 5.9-fold, at least about 6-fold, at least about 6.1-fold, at least about 6.2-fold, at least about 6.3-fold, at least about 6.4-fold, at least about 6.5-fold, at least about 6.6-fold, at least about 6.7-fold, at least about 6.8-fold, at least about 6.9-fold, at least about 7-fold, at least about 7.1-fold, at least about 7.2-fold, at least about 7.3-fold, at least about 7.4-fold, at least about 7.5-fold, at least about 7.6-fold, at least about 7.7-fold, at least about 7.8-fold, at least about 7.9-fold, at least about 8-fold, at least about 8.1-fold, at least about 8.2-fold, at least about 8.3-fold, at least about 8.4-fold, at least about 8.5-fold, at least about 8.6-fold, at least about 8.7-fold, at least about 8.8-fold, at least about 8.9-fold, at least about 9-fold, at least about 9.1-fold, at least about 9.2-fold, at least about 9.3-fold, at least about 9.4-fold, at least about 9.5-fold, at least about 9.6-fold, at least about 9.7-fold, at least about 9.8-fold, at least about 9.9-fold, at least about 10-fold or higher increase in the Mean Deviation Score of the PHI-4 polypeptide relative to the pesticidal activity of AXMI-205 (SEQ ID NO: 35).

In some embodiments the improved activity of the PHI-4 polypeptide is relative to the pesticidal activity of AXMI-205(evo25) (SEQ ID NO: 41), AXMI-205(evo30) (SEQ ID NO: 42), Axmi205 PMlibl PoolIG2_p2a1 1 (mutation S468L; SEQ ID NO: 45), Axmi205 PMlibl PoolIG2_plcl (mutation V467T; SEQ ID NO: 46), Axmi205 PMlibl PoolIG2_pla4 (mutation R464N; SEQ ID NO: 47), AXMI-205(evo34) (SEQ ID NO: 43) or AXMI-205(evo35) (SEQ ID NO: 44).

“Mean FAE Index” (MFI) refers to the mean of multiple FAEGN an arithmetic mean of FAEGN. As used herein, the “Mean Deviation Score” refers to the arithmetic mean of multiple Deviation Scores.

In some embodiments the PHI-4 polypeptides have increased insecticidal activity against one or more insect pests of the order Coleoptera.

In some embodiments the PHI-4 polypeptides have increased insecticidal activity against one or more insect pests of the corn rootworm complex (western corn rootworm, *Diabrotica virgifera*; northern corn rootworm, *D. barberi*; and the Mexican corn rootworm, *D. virgifera zeae*).

5 In some embodiments the PHI-4 polypeptides have increased insecticidal activity against *Diabrotica virgifera* larvae - Western Corn Root Worm (WCRW).

“Fragments” or “biologically active portions” include polypeptide fragments comprising amino acid sequences sufficiently identical to a PHI-4 polypeptide and that exhibit insecticidal activity. “Fragments” or “biologically active portions” include
 10 polypeptide fragments comprising amino acid sequences sufficiently identical to the amino acid sequence set forth in SEQ ID NO: 35, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526 and that exhibit insecticidal activity. A biologically active portion of a PHI-4 polypeptide can be a polypeptide that is, for example, 10, 25, 50, 100, 150, 200, 250, 300, 350, 400, 450, 500, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527,
 15 528, 529, 530, 531, 532, 533, 534 or 535 amino acids in length. Such biologically active portions can be prepared by recombinant techniques and evaluated for insecticidal activity. As used here, a fragment comprises at least 8 contiguous amino acids of a PHI-4 polypeptide. In some embodiments a fragment comprises at least 8 contiguous amino acids of SEQ ID NO: 2, SEQ ID NOs: 51-1162, or SEQ ID NOs: 1518-1526. In some
 20 embodiments a fragment comprises at least 8 contiguous amino acids of SEQ ID NO: 2 or SEQ ID NOs: 51-1162, or SEQ ID NOs: 1518-1526. The embodiments encompass other fragments, however, such as any fragment in the protein greater than about 10, 20, 30, 50, 100, 150, 200, 250 or more amino acids.

In some embodiments, the fragment is an N-terminal and/or a C-terminal
 25 truncation of at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25 or more amino acids relative to SEQ ID NO: 2, SEQ ID NO: 35, SEQ ID NOs: 51-1162, SEQ ID NOs: 1518-1526 or variants thereof e.g., by proteolysis, by insertion of a start codon, by deletion of the codons encoding the deleted amino acids and concomitant insertion of a start codon and/or insertion of a stop codon. In some embodiments, the
 30 fragments encompassed herein result from the removal of the C-terminal 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28 or 29 amino acids relative to SEQ ID NO: 35, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, SEQ ID NOs: 1518-1526, and variants thereof by proteolysis or by insertion of a start codon, by deletion of the codons encoding the deleted amino acids and
 35 concomitant insertion of a start codon. In particular embodiments the proteolytic cleavage site is between Lys at 520 and Ser at 521 Ser or Lys at 313 and Val at 314 of SEQ ID NO:

35 or variants thereof. It is well known in the art that polynucleotide encoding the truncated PHI-4 polypeptide can be engineered to add a start codon at the N-terminus such as ATG encoding methionine or methionine followed by an alanine. It is also well known in the art that depending on what host the PHI-4 polypeptide is expressed in the methionine may be partially of completed processed off.

In some embodiments fragments, biologically active portions, of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, SEQ ID NOs: 1518-1526 as well as amino acid substitutions, amino acid deletions and/or insertions thereof are also provided, and may be used to practice the methods of the disclosure.

In some embodiments PHI-4 polypeptides are provided having one or more amino acid substitution compared to AXMI-205 (SEQ ID NO: 35). In some embodiments PHI-4 polypeptides are provided having amino acid substitutions at solvent exposed surface residues to modify the protein characteristics of AXMI-205 (SEQ ID NO: 35), including but not limited to the ionic polarity of the protein surface. In some embodiments PHI-4 polypeptides are provided having amino acid substitutions at hydrophilic residues such as Asp, Glu, Lys, Arg, His, Ser, Thr, Tyr, Trp, Asn, Gln, and Cys. In some embodiments the PHI-4 polypeptides are provided having amino acid substitutions changing a Lysine or Arginine to a Glutamine, Glutamic Acid, Asparagine or Glutamic Acid; changing a Glutamic Acid or Aspartic Acid to a Lysine, Asparagine or Glutamine; and changing a Glutamine to a Asparagine or Lysine.

In some embodiments PHI-4 polypeptides are provided having amino acid substitutions at residues in a membrane insertion loop. In some embodiments PHI-4 polypeptides are provided having amino acid substitutions in a membrane insertion loop between about amino acid at position 92 (Val) and 101 (Ala) and/or at position 211 (Gly) and 220 (Glu) relative to SEQ ID NO: 35.

In some embodiments PHI-4 polypeptides are provided having amino acid substitutions at residues and receptor binding loops. In some embodiments PHI-4 polypeptides are provided having amino acid substitutions at residues and receptor binding loops between about amino acid 332 (Asp) and 340 (Asp), 395 (Asp) and 403 (Asp) , 458 (Asp) and 466 (Asp) relative to SEQ ID NO: 35.

In some embodiments PHI-4 polypeptides are provided having amino acid substitutions at residues in a protease sensitive region. In some embodiments PHI-4 polypeptides are provided having amino acid substitutions at residues in a protease sensitive region from about amino acid residues between 305 (Lys) and 316 (Lys) and 500 (Arg) and 535 (Lys) relative to SEQ ID NO: 35.

By variants is intended proteins or polypeptides having an amino acid sequence that is at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the parental amino acid sequence. In some embodiments a PHI-4 polypeptide
 5 has at least about 60%, 65%, about 70%, 75%, at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater identity across the entire length of the amino acid sequence of SEQ ID NO: 2, SEQ ID NOs: 51-1162, SEQ ID NOs: 1518-1526. In some embodiments a PHI-4 polypeptide has at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%,
 10 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater identity across the entire length of the amino acid sequence of any one of SEQ ID NO: 2, SEQ ID NOs: 51-1162, or SEQ ID NOs: 1518-1526.

In some embodiments a PHI-4 polypeptide comprises an amino acid sequence having at least 80% identity, to the amino acid sequence of any one of SEQ ID NO: 2,
 15 SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162 or SEQ ID NO: 1518-1526, wherein the polypeptide has insecticidal activity.

In some embodiments a PHI-4 polypeptide comprises an amino acid sequence having at least 90% identity to the amino acid sequence of any one of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162 or SEQ ID NO: 1518-1526, wherein
 20 the polypeptide has insecticidal activity.

In some embodiments a PHI-4 polypeptide comprises an amino acid sequence having at least 95% identity to the amino acid sequence of any one of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162 or SEQ ID NO: 1518-1526 wherein
 the polypeptide has insecticidal activity.

25 In some embodiments a PHI-4 polypeptide comprises an amino acid sequence having at least 97% identity to the amino acid sequence of any one of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162 or SEQ ID NO: 1518-1526, wherein the polypeptide has insecticidal activity.

In some embodiments a PHI-4 polypeptide comprises an amino acid sequence of
 30 SEQ ID NO: 3, wherein Xaa at position 2 is Ala or Arg; Xaa at position 9 is Gln, Lys or Glu; Xaa at position 14 is Pro or Ala; Xaa at position 16 is Val or Asp; Xaa at position 19 is Met or Leu; Xaa at position 22 is Gly or Ser; Xaa at position 24 is Asp, Asn or Gln; Xaa at position 36 is Leu or Met; Xaa at position 42 is Asp, Asn or Gln; Xaa at position 43 is Phe or Glu; Xaa at position 46 is Glu, Asp, Asn or Gly; Xaa at position 50 is Ile or Val; Xaa
 35 at position 51 is Glu or Gln; Xaa at position 55 is Arg or Lys; Xaa at position 56 is Ser or Thr; Xaa at position 57 is Tyr or Phe; Xaa at position 58 is Thr or Ser; Xaa at position 61

is Arg, Lys or Glu; Xaa at position 73 is Phe or Tyr; Xaa at position 74 is Lys, Glu, Gly, Arg, Met, Leu, His or Asp; Xaa at position 76 is Asp or Gln; Xaa at position 79 is Lys or Glu; Xaa at position 80 is Glu or Ser; Xaa at position 82 is Glu, Ile, Leu, Tyr or Gln; Xaa at position 83 is Glu or Gln; Xaa at position 84 is Tyr or Phe; Xaa at position 86 is Glu or Gln; Xaa at position 87 is Lys or Gln; Xaa at position 88 is Met, Ile or Leu; Xaa at position 90 is Gln or Glu; Xaa at position 94 is Val or Ile; Xaa at position 97 is Arg, Asn, Asp, Glu, Gln, Gly, Ser, Ile, Phe, His, Lys, Thr, Asn, Tyr, Trp, Pro, Cys, Ala, Met, Val or Leu; Xaa at position 98 is Tyr or Phe; Xaa at position 99 is Lys, Leu, Tyr, Ile, Met, Phe, Cys, Val or Asn; Xaa at position 103 is Ala or Gly; Xaa at position 105 is Leu or Ile; Xaa at position 109 is Phe, Lys, Gly, Met, Ser, Asp, Asn, Glu, Cys, Ala or Arg; Xaa at position 112 is Thr or Ser; Xaa at position 113 is Asp, Glu or Met; Xaa at position 117 is Thr or Ser; Xaa at position 121 is Tyr or Phe; Xaa at position 127 is Ala or Thr; Xaa at position 142 is Arg or Glu; Xaa at position 146 is Arg or Gln; Xaa at position 147 is Arg, Glu or Gln; Xaa at position 148 is Asp, Phe, Pro, Val, Glu, His, Trp, Ala, Arg, Leu, Ser, Gln or Gly; Xaa at position 149 is Phe or Val; Xaa at position 150 is Arg, Gln, Glu or Asn; Xaa at position 151 is Asp, Ser, Ala, Asn, Trp, Val, Gln, Cys, Met, Leu, Arg or Glu; Xaa at position 153 is Leu or Ile; Xaa at position 154 is Asn or Asp; Xaa at position 155 is Asn or Lys; Xaa at position 159 is Pro or Asp; Xaa at position 162 is Glu, Asp, Gln, Asn or Leu; Xaa at position 165 is Lys, Glu, Gln, Pro, Thr, Ala, Leu, Gly, Asp, Val, His, Ile, Met, Trp, Phe, Tyr or Arg; Xaa at position 166 is Arg or Gln; Xaa at position 167 is Tyr, Trp or Cys; Xaa at position 170 is Tyr or His; Xaa at position 171 is Tyr or Phe; Xaa at position 172 is Ile, Leu or Val; Xaa at position 173 is Ser or Ala; Xaa at position 174 is Glu, Gln, Asn, Lys, Val or Ser; Xaa at position 182 is Asp or Gln; Xaa at position 183 is Tyr or Val; Xaa at position 184 is Ser or Thr; Xaa at position 185 is Ala or Ser; Xaa at position 189 is Thr, Lys or Ile; Xaa at position 191 is Lys or Gln; Xaa at position 193 is Asp or Asn; Xaa at position 196 is Gln, Lys, Asn, Asp, Glu, Ala, Ile or Arg; Xaa at position 202 is Ala or Val; Xaa at position 203 is Glu, Thr or His; Xaa at position 204 is Met or Ala; Xaa at position 206 is Tyr or Phe; Xaa at position 207 is Lys or Gln; Xaa at position 209 is Leu or Pro; Xaa at position 210 is Val or Ile; Xaa at position 214 is Lys, Ser or Gln; Xaa at position 216 is Glu, Gln, Phe, Val, Tyr or Arg; Xaa at position 220 is Glu, His, Asp, Thr, Tyr, Val, Ser, Gln, Arg, Trp, Met, Ala, Phe, Ile, Leu, Cys or Asn; Xaa at position 229 is Arg or Glu; Xaa at position 230 is Ser or Glu; Xaa at position 231 is Asn or Ser; Xaa at position 236 is Leu or Pro; Xaa at position 245 is Met or Leu; Xaa at position 247 is Asp or Tyr; Xaa at position 256 is Gln, Lys or Glu; Xaa at position 257 is Gln, Ile, Glu, Cys, Ser, His, Trp or Met; Xaa at position 261 is Gln, Glu, Lys or Ala; Xaa at position 264 is Glu or Gln; Xaa at position 268 is Asp or Asn; Xaa at position 276 is Ser or Ala; Xaa at position 278 is Glu, Asn or Gln; Xaa at position

281 is Gln, Lys or Glu; Xaa at position 282 is Pro or Gly; Xaa at position 284 is Trp or Arg;
 Xaa at position 287 is Ala or Cys; Xaa at position 289 is Lys, Leu, Val, Pro, Glu, Gln, Tyr,
 Thr, Asp, Phe, Ser, Met, Arg, Trp, Ile, His, Asn, Cys, Gly or Ala; Xaa at position 291 is Glu
 or Gln; Xaa at position 292 is Arg or Gln; Xaa at position 293 is Arg, Glu or Gln; Xaa at
 5 position 294 is Val or Ala; Xaa at position 296 is Leu or Ile; Xaa at position 297 is Glu or
 Gln; Xaa at position 298 is Asp or Gln; Xaa at position 300 is Phe or Tyr; Xaa at position
 302 is Glu or Gln; Xaa at position 303 is Phe or Tyr; Xaa at position 305 is Lys, Gln, Ala,
 Ile, Met, Asn, Thr or Val; Xaa at position 306 is Gln or Lys; Xaa at position 309 is Gln, Lys
 or Glu; Xaa at position 313 is Lys, Gln or Arg; Xaa at position 316 is Lys or Gln; Xaa at
 10 position 328 is Lys, Glu or Gln; Xaa at position 331 is Glu, Asn or Gln; Xaa at position 333
 is Ser, Arg, Gly, Lys, Val, Asn, Ala, His, Gln, Thr, Asp, Ile, Leu, Cys or Glu; Xaa at position
 334 is Gly, Arg, Lys, Ile or Trp; Xaa at position 335 is Ser or Ala; Xaa at position 336 is
 Gly or Ala; Xaa at position 337 is Ala, Val or Gly; Xaa at position 338 is Ser, His, Val, Lys,
 Ala, Gly, Thr, Ile, Glu, Met, Arg, Pro, Asp, Asn or Leu; Xaa at position 339 is Glu, Asn,
 15 Gln, Ile, Pro, Met, Ser, Ala, Cys, Phe, Val, Leu, Asp, Trp, His or Arg; Xaa at position 341
 is Leu or Val; Xaa at position 342 is Ala, Ser or Val; Xaa at position 343 is Val or Ile; Xaa
 at position 344 is Phe or Trp; Xaa at position 345 is Asn or His; Xaa at position 346 is Pro
 or Ala; Xaa at position 350 is Asn or Ser; Xaa at position 351 is Gly or Val; Xaa at position
 354 is Met or Leu; Xaa at position 355 is Val, Ile or Leu; Xaa at position 359 is Gly or Ala;
 20 Xaa at position 362 is Asn or Ser; Xaa at position 364 is Ala or Ser; Xaa at position 371 is
 Ala, Gly or Thr; Xaa at position 374 is Phe or Ile; Xaa at position 375 is Lys or Arg; Xaa at
 position 380 is Leu or Gly; Xaa at position 382 is Val, Asp or Leu; Xaa at position 383 is
 Leu, Ile or Val; Xaa at position 384 is Lys, Ala or Gly; Xaa at position 385 is Ala or Gly;
 Xaa at position 389 is Trp or Tyr; Xaa at position 391 is Arg, Leu, Glu, Gln, Asp or His;
 25 Xaa at position 395 is Asp or Cys; Xaa at position 396 is Ala, Leu, Lys, Asn, Gly, Ile, Met,
 Arg, Tyr, Gln, His or Thr; Xaa at position 397 is Gly, Arg or Ala; Xaa at position 398 is Ser,
 Gln or Cys; Xaa at position 401 is Ser, His, Pro, Gly, Lys, Val, Arg, Ile, Asn, Phe, Thr, Ala,
 Asp, Met, Gln or Glu; Xaa at position 402 is Lys, Phe, His, Arg, Trp, Gly, Asn, Leu, Tyr,
 Thr, Val, Met, Pro or Ala; Xaa at position 403 is Asp, Tyr, Trp, Phe or Glu; Xaa at position
 30 405 is Ala or Ser; Xaa at position 409 is Ala or Pro; Xaa at position 410 is Ile or Val; Xaa
 at position 411 is Pro or Ala; Xaa at position 412 is Pro or Ala; Xaa at position 416 is Arg,
 Glu or Gln; Xaa at position 417 is Ala, Ser or Cys; Xaa at position 418 is Leu or Met; Xaa
 at position 422 is Met or Val; Xaa at position 426 is Thr or Ser; Xaa at position 436 is Asp
 or Lys; Xaa at position 437 is Tyr or Val; Xaa at position 438 is Val or Arg; Xaa at position
 35 440 is Val or Leu; Xaa at position 442 is Gln, Lys or Glu; Xaa at position 445 is Cys, Leu
 or Thr; Xaa at position 447 is Asp, Lys, Tyr, Ser, Glu, Ile, Gly, Pro, Leu, Phe, Trp or Thr;

Xaa at position 448 is Val or Ala; Xaa at position 449 is Gln or Glu; Xaa at position 452 is Gln, Lys or Glu, Ala; Xaa at position 453 is Asn or Asp; Xaa at position 454 is Arg, Tyr, Met, Ser, Val, Ile, Lys, Phe, Trp, Gln, Gly, His, Asp, Leu, Thr, Pro or Asn; Xaa at position 455 is Val or Ile; Xaa at position 457 is Trp or Asn; Xaa at position 459 is Lys, Met, Val, Trp, Gln, Ile, Thr, Ser, His, Cys, Tyr, Pro, Asn, Ala, Arg or Glu; Xaa at position 460 is Gly or Ala; Xaa at position 461 is Thr or Ser; Xaa at position 462 is Gly or Ala; Xaa at position 463 is Ala, Ser or Gly; Xaa at position 464 is Arg, Gly, His, Gln, Thr, Phe, Ala, Asp, Ser or Lys; Xaa at position 465 is Lys, Asn, Val, Met, Pro, Gly, Arg, Thr, His, Cys, Trp, Phe or Leu; Xaa at position 466 is Asp or Arg; Xaa at position 471 is Gln, Lys, Glu or Met; Xaa at position 472 is Pro or Ser; Xaa at position 497 is Asp or Gln; Xaa at position 499 is Glu or Gln; Xaa at position 500 is Arg, Gln or Lys; Xaa at position 502 is Arg, Glu or Gln; Xaa at position 509 is Lys, Gln, Glu or Ala; Xaa at position 517 is Gln, Cys, Asn, Val or Pro; Xaa at position 518 is Glu or Gln; Xaa at position 520 is Lys, Gln, Glu, His or Ala; Xaa at position 525 is Gln or Lys; and Xaa at position 527 is Gln, Lys, Pro, Cys, Glu, Ser, His, Phe or Trp; and having one or more amino acid substitutions at positions designated as Xaa in SEQ ID NO: 3; and amino acid substitutions, deletions, insertions, and combinations thereof and fragments thereof; and wherein the PHI-4 polypeptide has increased insecticidal activity compared to SEQ ID NO: 2.

In some embodiments a PHI-4 polypeptide comprises an amino acid sequence of SEQ ID NO: 4 having 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60 or 61 amino acid substitutions, in any combination, at residues designated by Xaa in SEQ ID NO: 4 compared to the native amino acid at the corresponding position of SEQ ID NO: 2.

In some embodiments a PHI-4 polypeptide comprises an amino acid sequence of SEQ ID NO: 4 having 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 amino acid substitutions, in any combination, at residues designated by Xaa in SEQ ID NO: 4 compared to the native amino acid at the corresponding position of SEQ ID NO: 2.

In some embodiments a PHI-4 polypeptide comprises an amino acid sequence having at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or greater sequence identity to the amino acid sequence of SEQ ID NO: 4.

In some embodiments a PHI-4 polypeptide comprises an amino acid sequence of SEQ ID NO: 4, wherein Xaa at position 2 is Ala or Arg; Xaa at position 24 is Asp or Asn; Xaa at position 42 is Asp or Asn; Xaa at position 43 is Phe or Glu; Xaa at position 46 is Glu or Asn; Xaa at position 74 is Lys, Glu or Gly; Xaa at position 79 is Lys or Glu; Xaa at

position 82 is Glu, Ile, Leu or Tyr; Xaa at position 97 is Arg, Asn, Asp, Glu, Gln, Gly, Ser, Ile, Phe, His, Lys, Thr, Asn, Tyr, Trp, Pro, Cys, Ala, Met, Val or Leu; Xaa at position 98 is Tyr or Phe; Xaa at position 99 is Lys, Leu, Tyr, Ile or Met; Xaa at position 109 is Phe, Lys, Gly, Met, Ser, Asp or Asn; Xaa at position 147 is Arg or Glu; Xaa at position 148 is Asp, Phe or Pro; Xaa at position 150 is Arg, Gln, Glu or Asn; Xaa at position 151 is Asp, Ser, Ala or Asn; Xaa at position 153 is Leu or Ile; Xaa at position 162 is Glu, Asp, Gln, Asn or Leu; Xaa at position 165 is Lys, Glu or Gln; Xaa at position 166 is Arg or Gln; Xaa at position 171 is Tyr or Phe; Xaa at position 174 is Glu, Gln, Asn, Lys, Val or Ser; Xaa at position 182 is Asp or Gln; Xaa at position 196 is Gln, Lys, Asn or Asp; Xaa at position 203 is Glu, Thr or His; Xaa at position 206 is Tyr or Phe; Xaa at position 216 is Glu or Gln; Xaa at position 220 is Glu, His, Asp, Thr, Tyr, Val, Ser or Gln; Xaa at position 247 is Asp or Tyr; Xaa at position 256 is Gln or Lys; Xaa at position 257 is Gln or Ile; Xaa at position 261 is Gln, Glu, Lys or Ala; Xaa at position 278 is Glu or Asn; Xaa at position 281 is Gln, Lys or Glu; Xaa at position 289 is Lys, Leu, Val, Pro, Glu, Gln, Tyr, Thr or Asp; Xaa at position 293 is Arg, Glu or Gln; Xaa at position 313 is Lys or Gln; Xaa at position 328 is Lys, Glu or Gln; Xaa at position 333 is Ser, Gly, Lys, Val or Asn; Xaa at position 334 is Gly, Arg, Lys or Ile; Xaa at position 336 is Gly or Ala; Xaa at position 338 is Ser, His, Val, Lys or Ala; Xaa at position 339 is Glu, Asn, Ile or Pro; Xaa at position 343 is Val or Ile; Xaa at position 346 is Pro or Ala; Xaa at position 355 is Val or Ile; Xaa at position 359 is Gly or Ala; Xaa at position 391 is Arg, Leu, Glu, Gln, Asp or His; Xaa at position 396 is Ala, Leu, Lys, Asn, Gly or Thr; Xaa at position 401 is Ser, His, Pro, Gly, Lys, Val or Arg; Xaa at position 402 is Lys, Phe, His, Arg, Gly, Trp, Thr, Asn, Tyr or Met; Xaa at position 403 is Asp or Tyr; Xaa at position 411 is Pro or Ala; Xaa at position 412 is Pro or Ala; Xaa at position 416 is Arg or Glu; Xaa at position 417 is Ala or Ser; Xaa at position 418 is Leu or Met; Xaa at position 426 is Thr or Ser; Xaa at position 440 is Val or Leu; Xaa at position 447 is Asp, Lys, Tyr, Ser, Glu or Ile; Xaa at position 452 is Gln, Lys or Glu; Xaa at position 454 is Arg, Tyr, Met, Ser, Val, Ile, Lys, Phe, Trp or Gln; Xaa at position 455 is Val or Ile; Xaa at position 459 is Lys, Met, Val, Trp, Gln, Ile or Tyr; Xaa at position 461 is Thr or Ser; Xaa at position 462 is Gly or Ala; Xaa at position 463 is Ala or Ser; Xaa at position 464 is Arg, Gly, His, Ala, Asp, Ser or Lys; Xaa at position 465 is Lys, Asn, Val, Met, Pro, Gly or Arg; Xaa at position 471 is Gln, Lys, Glu or Met; Xaa at position 472 is Pro or Ser; Xaa at position 500 is Arg or Gln; Xaa at position 509 is Lys, Gln or Ala; Xaa at position 520 is Lys, Gln, Glu, His or Ala; and Xaa at position 527 is Gln, Lys, Pro, Cys or Glu; and having one or more amino acid substitutions at positions designated as Xaa in SEQ ID NO: 4; and amino acid substitutions, deletions, insertions, and combinations thereof and

fragments thereof; and wherein the PHI-4 polypeptide has increased insecticidal activity compared to SEQ ID NO: 2.

In some embodiments a PHI-4 polypeptide comprises an amino acid sequence of SEQ ID NO: 3 having 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75 or 76 amino acid substitutions, in any combination, at residues designated by Xaa in SEQ ID NO: 3 compared to the native amino acid at the corresponding position of SEQ ID NO: 2.

In some embodiments PHI-4 polypeptide comprises an amino acid sequence of SEQ ID NO: 3 having 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53 or 54 amino acid substitutions, in any combination, at residues designated by Xaa in SEQ ID NO: 3 compared to the native amino acid at the corresponding position of SEQ ID NO: 2.

In some embodiments PHI-4 polypeptide comprises an amino acid sequence of SEQ ID NO: 3 having 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 11, 12, 13, 14 or 15 amino acid substitutions, in any combination, at residues designated by Xaa in SEQ ID NO: 3 compared to the native amino acid at the corresponding position of SEQ ID NO: 2.

In some embodiments a PHI-4 polypeptide comprises an amino acid sequence having at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or greater sequence identity to the amino acid sequence of SEQ ID NO: 3.

In some embodiments a PHI-4 polypeptide comprises one or more amino acid substitutions compared to the native amino acid at position 40, 42, 43, 46, 52, 97, 98, 99, 145, 150, 151, 153, 163, 171, 172, 182, 196, 206, 210, 216, 220, 278, 283, 289, 293, 328, 333, 334, 336, 338, 339, 342, 346, 354, 355, 370, 389, 393, 396, 401, 402, 403, 410, 412, 416, 417, 426, 442, 447, 452, 454, 455, 457, 461, 462, 500, 509, 520 or 527 of SEQ ID NO: 35.

In some embodiments a PHI-4 polypeptide comprises one or more amino acid substitutions compared to the native amino acid at position 40, 42, 43, 46, 52, 97, 98, 99, 145, 150, 151, 153, 163, 171, 172, 182, 196, 206, 210, 216, 220, 278, 283, 289, 293, 328, 333, 334, 336, 338, 339, 342, 346, 354, 355, 370, 389, 393, 396, 401, 402, 403, 410, 412, 416, 417, 426, 442, 447, 452, 454, 455, 457, 461, 462, 500, 509, 520 or 527 of SEQ ID NO: 35 wherein the amino acid at position 40 is Leu or Ile; the amino acid at position 42 is Asp or Asn; the amino acid at position 43 is Phe or Glu; the amino acid at position 46 is

Glu or Asn; the amino acid at position 52 is Ile or Val; the amino acid at position 97 is Arg, Asp, Glu or Asn; the amino acid at position 98 is Tyr or Phe; the amino acid at position 99 is Lys or Leu; the amino acid at position 145 is Leu or Val; the amino acid at position 150 is Arg or Gln; the amino acid at position 151 is Asp or Ser; the amino acid at position 153 is Leu or Ile; the amino acid at position 163 is Leu or Val; the amino acid at position 171 is Tyr or Phe; the amino acid at position 172 is Ile or Leu; the amino acid at position 182 is Asp or Gln; the amino acid at position 196 is Gln or Asn; the amino acid at position 206 is Tyr or Phe; the amino acid at position 210 is Val or Ile; the amino acid at position 216 is Glu or Gln; the amino acid at position 220 is Glu, Gln, His or Asp; the amino acid at position 278 is Glu or Asn; the amino acid at position 283 is Ile or Val; the amino acid at position 289 is Lys, Gln or Leu; the amino acid at position 293 is Arg, Gln or Glu; the amino acid at position 328 is Lys or Glu; the amino acid at position 333 is Ser, Lys or Val; the amino acid at position 334 is Gly, Lys or Arg; the amino acid at position 336 is Gly or Ala; the amino acid at position 338 is Ser or Val; the amino acid at position 339 is Glu, Asn or Gln; the amino acid at position 342 is Ala or Ser; the amino acid at position 346 is Pro or Ala; the amino acid at position 354 is Met or Leu; the amino acid at position 355 is Val or Ile; the amino acid at position 370 is His or Arg; the amino acid at position 389 is Trp or Leu; the amino acid at position 393 is Trp or Leu; the amino acid at position 396 is Ala, Leu, Lys, Thr or Gly; the amino acid at position 401 is Ser, His, Gly, Lys or Pro; the amino acid at position 402 is Lys, His, Gly or Trp; the amino acid at position 403 is Asp or Tyr; the amino acid at position 410 is Ile or Val; the amino acid at position 412 is Pro or Ala; the amino acid at position 416 is Arg or Glu; the amino acid at position 417 is Ala or Ser; the amino acid at position 426 is Thr or Ser; the amino acid at position 442 is Gln or Glu; the amino acid at position 447 is Asp or Lys; the amino acid at position 452 is Gln or Lys; the amino acid at position 454 is Arg or Gln; the amino acid at position 455 is Val or Ile; the amino acid at position 457 is Trp or Asn; the amino acid at position 461 is Thr or Ser; the amino acid at position 462 is Gly or Ala; the amino acid at position 500 is Arg or Gln; the amino acid at position 509 is Lys or Gln; the amino acid at position 520 is Lys, Glu or Gln; and the amino acid at position 527 is Gln or Lys., and amino acid deletions, insertions, and combinations thereof and fragments thereof.

In some embodiments the PHI-4 polypeptide comprising one or more amino acid substitutions at position 86, 359, 399, 464, 465, 466, 467, 468, 499 or 517.

In some embodiments the PHI-4 polypeptide comprising one or more amino acid substitutions at position 86, 359, 399, 464, 465, 466, 467, 468, 499 or 517, wherein the amino acid at position 86 is Glu or Thr; the amino acid at position 359 is Gly or Ala; the amino acid at position 399 is Gly or Ala; the amino acid at position 464 is Arg, Ala, Lys,

Asp or Asn; the amino acid at position 465 is Lys or Met, the amino acid at position 467 is Val, Ala, Leu or Thr; the amino acid at position 468 is Ser or Leu; the amino acid at position 499 is Glu or Ala or the amino acid at position 517 is Glu or Arg.

5 In some embodiments exemplary PHI-4 polypeptides are encoded by the polynucleotide sequence set forth in SEQ ID NO: 1, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NOs: 24-30, SEQ ID NOs. 1163-1505, and SEQ ID NOs. 1527-1535.

10 In some embodiments a PHI-4 polypeptide includes variants where an amino acid that is part of a proteolytic cleavage site is changed to another amino acid to eliminate or alter the proteolytic cleavage at that site. In some embodiments the proteolytic cleavage is by a protease in the insect gut. In other embodiments the proteolytic cleavage is by a plant protease in the transgenic plant.

15 In some embodiments exemplary PHI-4 polypeptides are the polypeptides shown in Table 3, Table 4, Table 5, Table 6, Table 7, Table 8, and Table 9 and combinations of the amino substitutions thereof as well as amino acid deletions, and or insertions and fragments thereof.

20 In some embodiments the PHI-4 polypeptide is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the nucleic acid molecule of SEQ ID NO: 1, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NOs: 24-30, SEQ ID NOs. 1163-1505, and SEQ ID NOs. 1527-1535. Variants include polypeptides that differ in amino acid sequence due to mutagenesis. Variant proteins encompassed by the disclosure are biologically active, that is they continue to possess the desired biological activity (i.e. pesticidal activity) of the native protein. By "retains activity" is intended that the variant will have at least about 30%, at least about 50%, at least about 70% or at least about 80% of the insecticidal activity of the native protein. In some embodiments, the variants may
25 have improved activity over the native protein.

Bacterial genes quite often possess multiple methionine initiation codons in proximity to the start of the open reading frame. Often, translation initiation at one or more of these start codons will lead to generation of a functional protein. These start codons can include ATG codons. However, bacteria such as *Bacillus* sp. also recognize
30 the codon GTG as a start codon, and proteins that initiate translation at GTG codons contain a methionine at the first amino acid. On rare occasions, translation in bacterial systems can initiate at a TTG codon, though in this event the TTG encodes a methionine. Furthermore, it is not often determined a priori which of these codons are used naturally in the bacterium. Thus, it is understood that use of one of the alternate methionine codons
35 may also lead to generation of pesticidal proteins. These pesticidal proteins are encompassed in the present disclosure and may be used in the methods of the present

disclosure. It will be understood that, when expressed in plants, it will be necessary to alter the alternate start codon to ATG for proper translation.

In another aspect the PHI-4 polypeptide may be expressed as a precursor protein with an intervening sequence that catalyzes multi-step, post translational protein splicing.

5 Protein splicing involves the excision of an intervening sequence from a polypeptide with the concomitant joining of the flanking sequences to yield a new polypeptide (Chong, *et al.*, (1996) *J. Biol. Chem.* 271:22159-22168). This intervening sequence or protein splicing element, referred to as inteins, which catalyze their own excision through three coordinated reactions at the N-terminal and C-terminal splice junctions: an acyl

10 rearrangement of the N-terminal cysteine or Serine; a transesterification reaction between the two termini to form a branched ester or thioester intermediate and peptide bond cleavage coupled to cyclization of the intein C-terminal asparagine to free the intein (Evans, *et al.*, (2000) *J. Biol. Chem.* 275:9091-9094. The elucidation of the mechanism of protein splicing has led to a number of intein-based applications (Comb, *et al.*, US Patent

15 Number 5,496,714; Comb, *et al.*, US Patent Number 5,834,247; Camarero and Muir, (1999) *J. Amer. Chem. Soc.* 121:5597-5598; Chong, *et al.*, (1997) *Gene* 192:271-281, Chong, *et al.*, (1998) *Nucleic Acids Res.* 26:5109-5115; Chong, *et al.*, (1998) *J. Biol. Chem.* 273:10567-10577; Cotton, *et al.*, (1999) *J. Am. Chem. Soc.* 121:1100-1101; Evans, *et al.*, (1999) *J. Biol. Chem.* 274:18359-18363; Evans, *et al.*, (1999) *J. Biol. Chem.*

20 274:3923-3926; Evans, *et al.*, (1998) *Protein Sci.* 7:2256-2264; Evans, *et al.*, (2000) *J. Biol. Chem.* 275:9091-9094; Iwai and Pluckthun, (1999) *FEBS Lett.* 459:166-172; Mathys, *et al.*, (1999) *Gene* 231:1-13; Mills, *et al.*, (1998) *Proc. Natl. Acad. Sci. USA* 95:3543-3548; Muir, *et al.*, (1998) *Proc. Natl. Acad. Sci. USA* 95:6705-6710; Otomo, *et al.*, (1999) *Biochemistry* 38:16040-16044; Otomo, *et al.*, (1999) *J. Biomol. NMR* 14:105-114; Scott, *et al.*, (1999) *Proc. Natl. Acad. Sci. USA* 96:13638-13643; Severinov and Muir, (1998) *J. Biol. Chem.* 273:16205-16209; Shingledecker, *et al.*, (1998) *Gene* 207:187-195; Southworth, *et al.*, (1998) *EMBO J.* 17:918-926; Southworth, *et al.*, (1999) *Biotechniques* 27:110-120; Wood, *et al.*, (1999) *Nat. Biotechnol.* 17:889-892; Wu, *et al.*, (1998a) *Proc. Natl. Acad. Sci. USA* 95:9226-9231; Wu, *et al.*, (1998b) *Biochim Biophys Acta* 1387:422-

30 432; Xu, *et al.*, (1999) *Proc. Natl. Acad. Sci. USA* 96:388-393; Yamazaki, *et al.*, (1998) *J. Am. Chem. Soc.* 120:5591-5592). For the application of inteins in plant transgenes see Yang, J, *et al.*, (*Transgene Res* 15:583-593 (2006)) and Evans, *et al.*, (*Annu. Rev. Plant Biol.* 56:375-392, (2005)).

In another aspect the PHI-4 polypeptide may be encoded by two separate genes

35 where the intein of the precursor protein comes from the two genes, referred to as a split-intein and the two portions of the precursor are joined by a peptide bond formation. This

peptide bond formation is accomplished by intein-mediated trans-splicing. For this purpose, a first and a second expression cassette comprising the two separate genes further code for inteins capable of mediating protein trans-splicing. By trans-splicing, the proteins and polypeptides encoded by the first and second fragments may be linked by peptide bond formation. Trans-splicing inteins may be selected from the nucleolar and organellar genomes of different organisms including eukaryotes, archaeobacteria and eubacteria. Inteins that may be used for are listed at neb.com/neb/inteins.html, which can be accessed on the world-wide web using the "www" prefix). The nucleotide sequence coding for an intein may be split into a 5' and a 3' part that code for the 5' and the 3' part of the intein, respectively. Sequence portions not necessary for intein splicing (e.g., homing endonuclease domain) may be deleted. The intein coding sequence is split such that the 5' and the 3' parts are capable of trans-splicing. For selecting a suitable splitting site of the intein coding sequence, the considerations published by Southworth, *et al.*, (1998) *EMBO J.* 17:918-926 may be followed. In constructing the first and the second expression cassette, the 5' intein coding sequence is linked to the 3' end of the first fragment coding for the N-terminal part of the PHI-4 polypeptide and the 3' intein coding sequence is linked to the 5' end of the second fragment coding for the C-terminal part of the PHI-4 polypeptide.

In general, the trans-splicing partners can be designed using any split intein, including any naturally-occurring or artificially-split split intein. Several naturally-occurring split inteins are known, for example: the split intein of the DnaE gene of *Synechocystis sp.* PCC6803 (see, Wu, *et al.*, (1998) *Proc Natl Acad Sci USA* 95(16):9226-31 and Evans, *et al.*, (2000) *J Biol Chem* 275(13):9091-4 and of the DnaE gene from *Nostoc punctiforme* (see, Iwai, *et al.*, (2006) *FEBS Lett* 580(7):1853-8). Non-split inteins have been artificially split in the laboratory to create new split inteins, for example: the artificially split Ssp DnaB intein (see, Wu, *et al.*, (1998) *Biochim Biophys Acta* 1387:422-32) and split Sce VMA intein (see, Brenzel, *et al.*, (2006) *Biochemistry* 45(6):1571-8) and an artificially split fungal mini-intein (see, Elleuche, *et al.*, (2007) *Biochem Biophys Res Commun* 355(3):830-4). There are also intein databases available that catalogue known inteins (see, for example the online-database available at: bioinformatics.weizmann.ac.il/~pietro/inteins/Inteinstable.html, which can be accessed on the world-wide web using the "www" prefix).

Naturally-occurring non-split inteins may have endonuclease or other enzymatic activities that can typically be removed when designing an artificially-split split intein. Such mini-inteins or minimized split inteins are well known in the art and are typically less than 200 amino acid residues long (see, Wu, *et al.*, (1998) *Biochim Biophys Acta*

1387:422-32). Suitable split inteins may have other purification enabling polypeptide elements added to their structure, provided that such elements do not inhibit the splicing of the split intein or are added in a manner that allows them to be removed prior to splicing. Protein splicing has been reported using proteins that comprise bacterial intein-like (BIL) domains (see, Amitai, *et al.*, (2003) *Mol Microbiol* 47:61-73) and hedgehog (Hog) auto-processing domains (the latter is combined with inteins when referred to as the Hog/intein superfamily or HINT family (see, Dassa, *et al.*, (2004) *J Biol Chem.* 279 32001-7) and domains such as these may also be used to prepare artificially-split inteins. In particular, non-splicing members of such families may be modified by molecular biology methodologies to introduce or restore splicing activity in such related species. Recent studies demonstrate that splicing can be observed when a N-terminal split intein component is allowed to react with a C-terminal split intein component not found in nature to be its "partner"; for example, splicing has been observed utilizing partners that have as little as 30 to 50% homology with the "natural" splicing partner (see, Dassa, *et al.*, (2007) *Biochemistry* 46(1):322-30). Other such mixtures of disparate split intein partners have been shown to be unreactive one with another (see, Brenzel, *et al.*, 2006 *Biochemistry* 45(6):1571-8). However, it is within the ability of a person skilled in the relevant art to determine whether a particular pair of polypeptides is able to associate with each other to provide a functional intein, using routine methods and without the exercise of inventive skill.

In another aspect the PHI-4 polypeptide is a circular permuted variant. In certain embodiments the PHI-4 polypeptide is a circular permuted variant of the polypeptide of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526. The development of recombinant DNA methods has made it possible to study the effects of sequence transposition on protein folding, structure and function. The approach used in creating new sequences resembles that of naturally occurring pairs of proteins that are related by linear reorganization of their amino acid sequences (Cunningham, *et al.*, (1979) *Proc. Natl. Acad. Sci. U.S.A.* 76:3218-3222; Teather and Erfle, (1990) *J. Bacteriol.* 172:3837-3841; Schimming, *et al.*, (1992) *Eur. J. Biochem.* 204:13-19; Yamiuchi and Minamikawa, (1991) *FEBS Lett.* 260:127-130; MacGregor, *et al.*, (1996) *FEBS Lett.* 378:263-266). The first in vitro application of this type of rearrangement to proteins was described by Goldenberg and Creighton (*J. Mol. Biol.* 165:407-413, 1983). In creating a circular permuted variant a new N-terminus is selected at an internal site (breakpoint) of the original sequence, the new sequence having the same order of amino acids as the original from the breakpoint until it reaches an amino acid that is at or near the original C-terminus. At this point the new sequence is joined,

either directly or through an additional portion of sequence (linker), to an amino acid that is at or near the original N-terminus and the new sequence continues with the same sequence as the original until it reaches a point that is at or near the amino acid that was N-terminal to the breakpoint site of the original sequence, this residue forming the new C-terminus of the chain. The length of the amino acid sequence of the linker can be selected empirically or with guidance from structural information or by using a combination of the two approaches. When no structural information is available, a small series of linkers can be prepared for testing using a design whose length is varied in order to span a range from 0 to 50 Å and whose sequence is chosen in order to be consistent with surface exposure (hydrophilicity, Hopp and Woods, (1983) *Mol. Immunol.* 20:483-489; Kyte and Doolittle, (1982) *J. Mol. Biol.* 157:105-132; solvent exposed surface area, Lee and Richards, (1971) *J. Mol. Biol.* 55:379-400) and the ability to adopt the necessary conformation without deranging the configuration of the pesticidal polypeptide (conformationally flexible; Karplus and Schulz, (1985) *Naturwissenschaften* 72:212-213. Assuming an average of translation of 2.0 to 3.8 Å per residue, this would mean the length to test would be between 0 to 30 residues, with 0 to 15 residues being the preferred range. Exemplary of such an empirical series would be to construct linkers using a cassette sequence such as Gly-Gly-Gly-Ser repeated n times, where n is 1, 2, 3 or 4. Those skilled in the art will recognize that there are many such sequences that vary in length or composition that can serve as linkers with the primary consideration being that they be neither excessively long nor short (cf., Sandhu, (1992) *Critical Rev. Biotech.* 12:437-462); if they are too long, entropy effects will likely destabilize the three-dimensional fold, and may also make folding kinetically impractical, and if they are too short, they will likely destabilize the molecule because of torsional or steric strain. Those skilled in the analysis of protein structural information will recognize that using the distance between the chain ends, defined as the distance between the c-alpha carbons, can be used to define the length of the sequence to be used or at least to limit the number of possibilities that must be tested in an empirical selection of linkers. They will also recognize that it is sometimes the case that the positions of the ends of the polypeptide chain are ill-defined in structural models derived from x-ray diffraction or nuclear magnetic resonance spectroscopy data, and that when true, this situation will therefore need to be taken into account in order to properly estimate the length of the linker required. From those residues whose positions are well defined are selected two residues that are close in sequence to the chain ends, and the distance between their c-alpha carbons is used to calculate an approximate length for a linker between them. Using the calculated length as a guide, linkers with a range of number of residues (calculated using 2 to 3.8 Å per

residue) are then selected. These linkers may be composed of the original sequence, shortened or lengthened as necessary, and when lengthened the additional residues may be chosen to be flexible and hydrophilic as described above; or optionally the original sequence may be substituted for using a series of linkers, one example being the Gly-Gly-Gly-Ser cassette approach mentioned above; or optionally a combination of the original sequence and new sequence having the appropriate total length may be used. Sequences of pesticidal polypeptides capable of folding to biologically active states can be prepared by appropriate selection of the beginning (amino terminus) and ending (carboxyl terminus) positions from within the original polypeptide chain while using the linker sequence as described above. Amino and carboxyl termini are selected from within a common stretch of sequence, referred to as a breakpoint region, using the guidelines described below. A novel amino acid sequence is thus generated by selecting amino and carboxyl termini from within the same breakpoint region. In many cases the selection of the new termini will be such that the original position of the carboxyl terminus immediately preceded that of the amino terminus. However, those skilled in the art will recognize that selections of termini anywhere within the region may function, and that these will effectively lead to either amino acid deletions or additions to the amino or carboxyl portions of the new sequence. It is a central tenet of molecular biology that the primary amino acid sequence of a protein dictates folding to the three-dimensional structure necessary for expression of its biological function. Methods are known to those skilled in the art to obtain and interpret three-dimensional structural information using x-ray diffraction of single protein crystals or nuclear magnetic resonance spectroscopy of protein solutions. Examples of structural information that are relevant to the identification of breakpoint regions include the location and type of protein secondary structure (alpha and 3-10 helices, parallel and anti-parallel beta sheets, chain reversals and turns, and loops; Kabsch and Sander, (1983) *Biopolymers* 22:2577-2637; the degree of solvent exposure of amino acid residues, the extent and type of interactions of residues with one another (Chothia, (1984) *Ann. Rev. Biochem.* 53:537-572) and the static and dynamic distribution of conformations along the polypeptide chain (Alber and Mathews, (1987) *Methods Enzymol.* 154:511-533). In some cases additional information is known about solvent exposure of residues; one example is a site of post-translational attachment of carbohydrate which is necessarily on the surface of the protein. When experimental structural information is not available or is not feasible to obtain, methods are also available to analyze the primary amino acid sequence in order to make predictions of protein tertiary and secondary structure, solvent accessibility and the occurrence of turns and loops. Biochemical methods are also sometimes applicable for empirically

determining surface exposure when direct structural methods are not feasible; for example, using the identification of sites of chain scission following limited proteolysis in order to infer surface exposure (Gentile and Salvatore, (1993) *Eur. J. Biochem.* 218:603-621). Thus using either the experimentally derived structural information or predictive methods (e.g., Srinivisan and Rose, (1995) *Proteins: Struct., Funct. & Genetics* 22:81-99) the parental amino acid sequence is inspected to classify regions according to whether or not they are integral to the maintenance of secondary and tertiary structure. The occurrence of sequences within regions that are known to be involved in periodic secondary structure (alpha and 3-10 helices, parallel and anti-parallel beta sheets) are regions that should be avoided. Similarly, regions of amino acid sequence that are observed or predicted to have a low degree of solvent exposure are more likely to be part of the so-called hydrophobic core of the protein and should also be avoided for selection of amino and carboxyl termini. In contrast, those regions that are known or predicted to be in surface turns or loops, and especially those regions that are known not to be required for biological activity, are the preferred sites for location of the extremes of the polypeptide chain. Continuous stretches of amino acid sequence that are preferred based on the above criteria are referred to as a breakpoint region. Polynucleotides encoding circular permuted PHI-4 polypeptides with new N-terminus/C-terminus which contain a linker region separating the original C-terminus and N-terminus can be made essentially following the method described in Mullins, *et al.*, (1994) *J. Am. Chem. Soc.* 116:5529-5533. Multiple steps of polymerase chain reaction (PCR) amplifications are used to rearrange the DNA sequence encoding the primary amino acid sequence of the protein. Polynucleotides encoding circular permuted PHI-4 polypeptides with new N-terminus/C-terminus which contain a linker region separating the original C-terminus and N-terminus can be made based on the tandem-duplication method described in Horlick, *et al.*, (1992) *Protein Eng.* 5:427-431. Polymerase chain reaction (PCR) amplification of the new N-terminus/C-terminus genes is performed using a tandemly duplicated template DNA.

In another aspect fusion proteins are provided that include within its amino acid sequence an amino acid sequence comprising a PHI-4 polypeptide including but not limited to the polypeptide of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526 and active fragments thereof.

In another aspect fusion proteins are provided comprising a PHI-4 polypeptide and a second pesticidal polypeptide such a Cry protein. Methods for design and construction of fusion proteins (and polynucleotides encoding same) are known to those of skill in the art. Polynucleotides encoding a PHI-4 polypeptide may be fused to signal sequences

which will direct the localization of the PHI-4 polypeptide to particular compartments of a prokaryotic or eukaryotic cell and/or direct the secretion of the PHI-4 polypeptide of the embodiments from a prokaryotic or eukaryotic cell. For example, in *E. coli*, one may wish to direct the expression of the protein to the periplasmic space. Examples of signal sequences or proteins (or fragments thereof) to which the PHI-4 polypeptide may be fused in order to direct the expression of the polypeptide to the periplasmic space of bacteria include, but are not limited to, the *pelB* signal sequence, the maltose binding protein (MBP) signal sequence, MBP, the *ompA* signal sequence, the signal sequence of the periplasmic *E. coli* heat-labile enterotoxin B-subunit, and the signal sequence of alkaline phosphatase. Several vectors are commercially available for the construction of fusion proteins which will direct the localization of a protein, such as the pMAL series of vectors (particularly the pMAL-p series) available from New England Biolabs. In a specific embodiment, the PHI-4 polypeptide may be fused to the *pelB* pectate lyase signal sequence to increase the efficiency of expression and purification of such polypeptides in Gram-negative bacteria (see, US Patent Numbers 5,576,195 and 5,846,818). Plant plastid transit peptide/polypeptide fusions are well known in the art (see, US Patent Number 7,193,133). Apoplast transit peptides such as rice or barley alpha-amylase secretion signal are also well known in the art. The plastid transit peptide is generally fused N-terminal to the polypeptide to be targeted (e.g., the fusion partner). In one embodiment, the fusion protein consists essentially of the peptide transit plastid and the PHI-4 polypeptide to be targeted. In another embodiment, the fusion protein comprises the peptide transit plastid and the polypeptide to be targeted. In such embodiments, the plastid transit peptide is preferably at the N-terminus of the fusion protein. However, additional amino acid residues may be N-terminal to the plastid transit peptide providing that the fusion protein is at least partially targeted to a plastid. In a specific embodiment, the plastid transit peptide is in the N-terminal half, N-terminal third or N-terminal quarter of the fusion protein. Most or all of the plastid transit peptide is generally cleaved from the fusion protein upon insertion into the plastid. The position of cleavage may vary slightly between plant species, at different plant developmental stages, as a result of specific intercellular conditions or the particular combination of transit peptide/fusion partner used. In one embodiment, the plastid transit peptide cleavage is homogenous such that the cleavage site is identical in a population of fusion proteins. In another embodiment, the plastid transit peptide is not homogenous, such that the cleavage site varies by 1-10 amino acids in a population of fusion proteins. The plastid transit peptide can be recombinantly fused to a second protein in one of several ways. For example, a restriction endonuclease recognition site can be introduced into the nucleotide sequence

of the transit peptide at a position corresponding to its C-terminal end and the same or a compatible site can be engineered into the nucleotide sequence of the protein to be targeted at its N-terminal end. Care must be taken in designing these sites to ensure that the coding sequences of the transit peptide and the second protein are kept "in frame" to allow the synthesis of the desired fusion protein. In some cases, it may be preferable to remove the initiator methionine codon of the second protein when the new restriction site is introduced. The introduction of restriction endonuclease recognition sites on both parent molecules and their subsequent joining through recombinant DNA techniques may result in the addition of one or more extra amino acids between the transit peptide and the second protein. This generally does not affect targeting activity as long as the transit peptide cleavage site remains accessible and the function of the second protein is not altered by the addition of these extra amino acids at its N-terminus. Alternatively, one skilled in the art can create a precise cleavage site between the transit peptide and the second protein (with or without its initiator methionine) using gene synthesis (Stemmer, *et al.*, (1995) *Gene* 164:49-53) or similar methods. In addition, the transit peptide fusion can intentionally include amino acids downstream of the cleavage site. The amino acids at the N-terminus of the mature protein can affect the ability of the transit peptide to target proteins to plastids and/or the efficiency of cleavage following protein import. This may be dependent on the protein to be targeted. See, e.g., Comai, *et al.*, (1988) *J. Biol. Chem.* 263(29):15104-9.

In some embodiments fusion proteins are provided comprising a PHI-4 polypeptide, a pesticidal protein such as a cry protein, and an amino acid linker.

In some embodiments fusion proteins are provided represented by a formula selected from the group consisting of

$R^1\text{-L-R}^2$, $R^2\text{-L-R}^1$, $R^1\text{-R}^2$ or $R^2\text{-R}^1$

where R^1 is a PHI-4 polypeptide, R^2 is a pesticidal protein with a different but complementary activity to the PHI-4 polypeptide, including but not limited to cry proteins; a polypeptide that increases the solubility and/or stability of the PHI-4 polypeptide; or a transit peptide or leader sequence. The R^1 polypeptide is fused either directly or through a linker segment to the R^2 polypeptide. The term "directly" defines fusions in which the polypeptides are joined without a peptide linker. Thus L represents a chemical bond or polypeptide segment to which both R^1 and R^2 are fused in frame, most commonly L is a linear peptide to which R^1 and R^2 are bound by amide bonds linking the carboxy terminus of R^1 to the amino terminus of L and carboxy terminus of L to the amino terminus of R^2 . By "fused in frame" is meant that there is no translation termination or disruption between the reading frames of R^1 and R^2 . The linking group (L) is generally a polypeptide of

between 1 and 500 amino acids in length. The linkers joining the two molecules are preferably designed to (1) allow the two molecules to fold and act independently of each other, (2) not have a propensity for developing an ordered secondary structure which could interfere with the functional domains of the two proteins, (3) have minimal hydrophobic or charged characteristic which could interact with the functional protein domains and (4) provide steric separation of R^1 and R^2 such that R^1 and R^2 could interact simultaneously with their corresponding receptors on a single cell. Typically surface amino acids in flexible protein regions include Gly, Asn and Ser. Virtually any permutation of amino acid sequences containing Gly, Asn and Ser would be expected to satisfy the above criteria for a linker sequence. Other neutral amino acids, such as Thr and Ala, may also be used in the linker sequence. Additional amino acids may also be included in the linkers due to the addition of unique restriction sites in the linker sequence to facilitate construction of the fusions.

In some embodiments the linkers comprise sequences selected from the group of formulas: $(Gly_3Ser)_n$, $(Gly_4Ser)_n$, $(Gly_5Ser)_n$, $(Gly_nSer)_n$ or $(AlaGlySer)_n$ where n is an integer. One example of a highly-flexible linker is the (GlySer)-rich spacer region present within the pIII protein of the filamentous bacteriophages, e.g., bacteriophages M13 or fd (Schaller, *et al.*, 1975). This region provides a long, flexible spacer region between two domains of the pIII surface protein. Also included are linkers in which an endopeptidase recognition sequence is included. Such a cleavage site may be valuable to separate the individual components of the fusion to determine if they are properly folded and active in vitro. Examples of various endopeptidases include, but are not limited to, Plasmin, Enterokinase, Kallikrein, Urokinase, Tissue Plasminogen activator, clostripain, Chymosin, Collagenase, Russell's Viper Venom Protease, Postproline cleavage enzyme, V8 protease, Thrombin and factor Xa. In some embodiments the linker comprises the amino acids EEKKN from the multi-gene expression vehicle (MGEV), which is cleaved by vacuolar proteases as disclosed in US 2007/0277263. In other embodiments, peptide linker segments from the hinge region of heavy chain immunoglobulins IgG, IgA, IgM, IgD or IgE provide an angular relationship between the attached polypeptides. Especially useful are those hinge regions where the cysteines are replaced with serines. Preferred linkers of the present disclosure include sequences derived from murine IgG gamma 2b hinge region in which the cysteines have been changed to serines. The fusion proteins are not limited by the form, size or number of linker sequences employed and the only requirement of the linker is that functionally it does not interfere adversely with the folding and function of the individual molecules of the fusion.

In another embodiment chimeric PHI-4 polypeptide are provided that are created through joining two or more portions of genes, which originally encoded separate insecticidal proteins from different species, to create a chimeric gene. The translation of the chimeric gene results in a single chimeric pesticidal polypeptide with regions, motifs or domains derived from each of the original polypeptides. In certain embodiments the chimeric protein comprises portions, motifs or domains of PHI-4 polypeptides in any combination. In certain embodiments the chimeric insecticidal polypeptide includes but not limited to the polypeptides of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526.

It is recognized that DNA sequences may be altered by various methods, and that these alterations may result in DNA sequences encoding proteins with amino acid sequences different than that encoded by the wild-type (or native) pesticidal protein. These proteins may be altered in various ways including amino acid substitutions, amino acid deletions, amino acid truncations, and insertions of one or more amino acids, including up to 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155 or more amino acid substitutions, amino acid deletions and/or insertions or combinations thereof compared to any one of SEQ ID NO: 35, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526. In some embodiments a PHI-4 polypeptide comprises the deletion of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28 or more amino acids from the C-terminus of the PHI-4 polypeptide relative to the amino acid position of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526. Methods for such manipulations are generally known in the art. For example, amino acid sequence variants of a PHI-4 polypeptide can be prepared by mutations in the DNA. This may also be accomplished by one of several forms of mutagenesis and/or in directed evolution. In some aspects, the changes encoded in the amino acid sequence will not substantially affect the function of the protein. Such variants will possess the desired pesticidal activity. However, it is understood that the ability of a PHI-4 polypeptide to confer pesticidal activity may be improved by the use of such techniques upon the compositions of this disclosure.

For example, conservative amino acid substitutions may be made at one or more, predicted, nonessential amino acid residues. A “nonessential” amino acid residue is a residue that can be altered from the wild-type sequence of a PHI-4 polypeptide without altering the biological activity. A “conservative amino acid substitution” is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art.

These families include: amino acids with basic side chains (e.g., lysine, arginine, histidine); acidic side chains (e.g., aspartic acid, glutamic acid); polar, negatively charged residues and their amides (e.g., aspartic acid, asparagine, glutamic, acid, glutamine; uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, 5 tyrosine, cysteine); small aliphatic, nonpolar or slightly polar residues (e.g., Alanine, serine, threonine, proline, glycine); nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan); large aliphatic, nonpolar residues (e.g., methionine, leucine, isoleucine, valine, cystine); beta-branched side chains (e.g., threonine, valine, isoleucine); aromatic side chains (e.g., tyrosine, phenylalanine, 10 tryptophan, histidine); large aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan).

Amino acid substitutions may be made in nonconserved regions that retain function. In general, such substitutions would not be made for conserved amino acid residues or for amino acid residues residing within a conserved motif, where such 15 residues are essential for protein activity. Examples of residues that are conserved and that may be essential for protein activity include, for example, residues that are identical between all proteins contained in an alignment of similar or related toxins to the sequences of the embodiments (e.g., residues that are identical in an alignment of homologous proteins). Examples of residues that are conserved but that may allow 20 conservative amino acid substitutions and still retain activity include, for example, residues that have only conservative substitutions between all proteins contained in an alignment of similar or related toxins to the sequences of the embodiments (e.g., residues that have only conservative substitutions between all proteins contained in the alignment homologous proteins). However, one of skill in the art would understand that functional 25 variants may have minor conserved or nonconserved alterations in the conserved residues. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the protein of interest may be found in the model of Dayhoff, *et al.*, (1978) *Atlas of Protein Sequence and Structure* (Natl. Biomed. Res. Found., Washington, D.C.), herein incorporated by reference.

30 In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a protein is generally understood in the art (Kyte and Doolittle, (1982) *J Mol Biol.* 157(1):105-32). It is accepted that the relative hydropathic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines 35 the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens and the like.

It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydropathic index or score and still result in a protein with similar biological activity, i.e., still obtain a biological functionally equivalent protein. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics (Kyte and Doolittle, *ibid*). These are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9) and arginine (-4.5). In making such changes, the substitution of amino acids whose hydropathic indices are within +2 is preferred, those which are within +1 are particularly preferred and those within +0.5 are even more particularly preferred.

It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. US Patent Number 4,554,101, states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein.

As detailed in US Patent Number 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0.+0.1); glutamate (+3.0.+0.1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5.+0.1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4).

Alternatively, alterations may be made to the protein sequence of many proteins at the amino or carboxy terminus without substantially affecting activity. This can include amino acid insertions, amino acid deletions or amino acid alterations introduced by modern molecular methods, such as PCR, including PCR amplifications that alter or extend the protein coding sequence by virtue of inclusion of amino acid encoding sequences in the oligonucleotides utilized in the PCR amplification. Alternatively, the protein sequences added can include entire protein-coding sequences, such as those used commonly in the art to generate protein fusions. Such fusion proteins are often used to (1) increase expression of a protein of interest (2) introduce a binding domain, enzymatic activity or epitope to facilitate either protein purification, protein detection or other experimental uses known in the art (3) target secretion or translation of a protein to a subcellular organelle, such as the periplasmic space of Gram-negative bacteria, mitochondria or chloroplasts of plants or the endoplasmic reticulum of eukaryotic cells, the latter of which often results in glycosylation of the protein.

In specific embodiments, the substitution is an alanine for the native amino acid at the recited position(s). Also encompassed are the nucleic acid sequence(s) encoding the variant protein or polypeptide.

Variant nucleotide and amino acid sequences of the disclosure also encompass
5 sequences derived from mutagenic and recombinogenic procedures such as DNA shuffling. With such a procedure, one or more different PHI-4 polypeptide coding regions can be used to create a new PHI-4 polypeptide possessing the desired properties. In this manner, libraries of recombinant polynucleotides are generated from a population of related sequence polynucleotides comprising sequence regions that have substantial
10 sequence identity and can be homologously recombined in vitro or in vivo. For example, using this approach, sequence motifs encoding a domain of interest may be shuffled between a pesticidal gene and other known pesticidal genes to obtain a new gene coding for a protein with an improved property of interest, such as an increased insecticidal activity. Strategies for such DNA shuffling are known in the art. See, for example,
15 Stemmer, (1994) *Proc. Natl. Acad. Sci. USA* 91:10747-10751; Stemmer, (1994) *Nature* 370:389-391; Crameri, *et al.*, (1997) *Nature Biotech.* 15:436-438; Moore, *et al.*, (1997) *J. Mol. Biol.* 272:336-347; Zhang, *et al.*, (1997) *Proc. Natl. Acad. Sci. USA* 94:4504-4509; Crameri, *et al.*, (1998) *Nature* 391:288-291 and US Patent Numbers 5,605,793 and 5,837,458.

20 Domain swapping or shuffling is another mechanism for generating altered PHI-4 polypeptides. Domains may be swapped between PHI-4 polypeptides, resulting in hybrid or chimeric toxins with improved pesticidal activity or target spectrum. Methods for generating recombinant proteins and testing them for pesticidal activity are well known in the art (see, for example, Naimov, *et al.*, (2001) *Appl. Environ. Microbiol.* 67:5328-5330; de Maagd, *et al.*, (1996) *Appl. Environ. Microbiol.* 62:1537-1543; Ge, *et al.*, (1991) *J. Biol. Chem.* 266:17954-17958; Schnepf, *et al.*, (1990) *J. Biol. Chem.* 265:20923-20930; Rang, *et al.*, 91999) *Appl. Environ. Microbiol.* 65:2918-2925).

Antibodies

30 Antibodies to a PHI-4 polypeptide of the embodiments or to variants or fragments thereof, are also encompassed. Methods for producing antibodies are well known in the art (see, for example, Harlow and Lane, (1988) *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.; US Patent Number 4,196,265).

A kit for detecting the presence of a PHI-4 polypeptide or detecting the presence
35 of a nucleotide sequence encoding a PHI-4 polypeptide, in a sample is provided. In one embodiment, the kit provides antibody-based reagents for detecting the presence of a

PHI-4 polypeptide in a tissue sample. In another embodiment, the kit provides labeled nucleic acid probes useful for detecting the presence of one or more polynucleotides encoding PHI-4 polypeptide(s). The kit is provided along with appropriate reagents and controls for carrying out a detection method, as well as instructions for use of the kit.”

5

Receptor identification and isolation

Receptors to the PHI-4 polypeptide of the embodiments or to variants or fragments thereof, are also encompassed. Methods for identifying receptors are well known in the art (see, Hofmann, *et al.*, (1988) *Eur. J. Biochem.* 173:85-91; Gill, *et al.*, (1995) *J. Biol. Chem.* 272:277-2782) can be employed to identify and isolate the receptor that recognizes the PHI-4 polypeptides using the brush-border membrane vesicles from susceptible insects. In addition to the radioactive labeling method listed in the cited literatures, PHI-4 polypeptide can be labeled with fluorescent dye and other common labels such as streptavidin. Brush-border membrane vesicles (BBMV) of susceptible insects such as soybean looper and stink bugs can be prepared according to the protocols listed in the references and separated on SDS-PAGE gel and blotted on suitable membrane. Labeled PHI-4 polypeptides can be incubated with blotted membrane of BBMV and labeled the PHI-4 polypeptides can be identified with the labeled reporters. Identification of protein band(s) that interact with the PHI-4 polypeptides can be detected by N-terminal amino acid gas phase sequencing or mass spectrometry based protein identification method (Patterson, (1998) 10(22):1-24, *Current Protocol in Molecular Biology* published by John Wiley & Son Inc). Once the protein is identified, the corresponding gene can be cloned from genomic DNA or cDNA library of the susceptible insects and binding affinity can be measured directly with the PHI-4 polypeptides. Receptor function for insecticidal activity by the PHI-4 polypeptides can be verified by accomplished by RNAi type of gene knock out method (Rajagopal, *et al.*, (2002) *J. Biol. Chem.* 277:46849-46851).

Nucleotide Constructs, Expression Cassettes and Vectors

The use of the term "nucleotide constructs" herein is not intended to limit the embodiments to nucleotide constructs comprising DNA. Those of ordinary skill in the art will recognize that nucleotide constructs particularly polynucleotides and oligonucleotides composed of ribonucleotides and combinations of ribonucleotides and deoxyribonucleotides may also be employed in the methods disclosed herein. The nucleotide constructs, nucleic acids, and nucleotide sequences of the embodiments additionally encompass all complementary forms of such constructs, molecules and sequences. Further, the nucleotide constructs, nucleotide molecules and nucleotide

sequences of the embodiments encompass all nucleotide constructs, molecules and sequences which can be employed in the methods of the embodiments for transforming plants including, but not limited to, those comprised of deoxyribonucleotides, ribonucleotides and combinations thereof. Such deoxyribonucleotides and
5 ribonucleotides include both naturally occurring molecules and synthetic analogues. The nucleotide constructs, nucleic acids, and nucleotide sequences of the embodiments also encompass all forms of nucleotide constructs including, but not limited to, single-stranded forms, double-stranded forms, hairpins, stem-and-loop structures and the like.

A further embodiment relates to a transformed organism such as an organism
10 selected from plant and insect cells, bacteria, yeast, baculovirus, protozoa, nematodes and algae. The transformed organism comprises a DNA molecule of the embodiments, an expression cassette comprising the DNA molecule or a vector comprising the expression cassette, which may be stably incorporated into the genome of the transformed organism.

15 The sequences of the embodiments are provided in DNA constructs for expression in the organism of interest. The construct will include 5' and 3' regulatory sequences operably linked to a sequence of the embodiments. The term "operably linked" as used herein refers to a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence
20 corresponding to the second sequence. Generally, operably linked means that the nucleic acid sequences being linked are contiguous and where necessary to join two protein coding regions in the same reading frame. The construct may additionally contain at least one additional gene to be cotransformed into the organism. Alternatively, the additional gene(s) can be provided on multiple DNA constructs.

25 Such a DNA construct is provided with a plurality of restriction sites for insertion of the PHI-4 polypeptide gene sequence to be under the transcriptional regulation of the regulatory regions. The DNA construct may additionally contain selectable marker genes.

The DNA construct will generally include in the 5' to 3' direction of transcription: a transcriptional and translational initiation region (i.e., a promoter), a DNA sequence of the
30 embodiments and a transcriptional and translational termination region (i.e., termination region) functional in the organism serving as a host. The transcriptional initiation region (i.e., the promoter) may be native, analogous, foreign or heterologous to the host organism and/or to the sequence of the embodiments. Additionally, the promoter may be the natural sequence or alternatively a synthetic sequence. The term "foreign" as used
35 herein indicates that the promoter is not found in the native organism into which the promoter is introduced. Where the promoter is "foreign" or "heterologous" to the

sequence of the embodiments, it is intended that the promoter is not the native or naturally occurring promoter for the operably linked sequence of the embodiments. As used herein, a chimeric gene comprises a coding sequence operably linked to a transcription initiation region that is heterologous to the coding sequence. Where the promoter is a native or natural sequence, the expression of the operably linked sequence is altered from the wild-type expression, which results in an alteration in phenotype.

In some embodiments the DNA construct may also include a transcriptional enhancer sequence. As used herein, the term an "enhancer" refers to a DNA sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Various enhancers are known in the art including for example, introns with gene expression enhancing properties in plants (US Patent Application Publication Number 2009/0144863, the ubiquitin intron (i.e., the maize ubiquitin intron 1 (see, for example, NCBI sequence S94464)), the omega enhancer or the omega prime enhancer (Gallie, *et al.*, (1989) *Molecular Biology of RNA* ed. Cech (Liss, New York) 237-256 and Gallie, *et al.*, (1987) *Gene* 60:217-25), the CaMV 35S enhancer (see, e.g., Benfey, *et al.*, (1990) *EMBO J.* 9:1685-96) and the enhancers of US Patent Number 7,803,992 may also be used, each of which is incorporated by reference. The above list of transcriptional enhancers is not meant to be limiting. Any appropriate transcriptional enhancer can be used in the embodiments.

The termination region may be native with the transcriptional initiation region, may be native with the operably linked DNA sequence of interest, may be native with the plant host or may be derived from another source (i.e., foreign or heterologous to the promoter, the sequence of interest, the plant host or any combination thereof).

Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. See also, Guerineau, *et al.*, (1991) *Mol. Gen. Genet.* 262:141-144; Proudfoot, (1991) *Cell* 64:671-674; Sanfacon, *et al.*, (1991) *Genes Dev.* 5:141-149; Mogen, *et al.*, (1990) *Plant Cell* 2:1261-1272; Munroe, *et al.*, (1990) *Gene* 91:151-158; Ballas, *et al.*, (1989) *Nucleic Acids Res.* 17:7891-7903 and Joshi, *et al.*, (1987) *Nucleic Acid Res.* 15:9627-9639.

Where appropriate, a nucleic acid may be optimized for increased expression in the host organism. Thus, where the host organism is a plant, the synthetic nucleic acids can be synthesized using plant-preferred codons for improved expression. See, for example, Campbell and Gowri, (1990) *Plant Physiol.* 92:1-11 for a discussion of host-preferred codon usage. For example, although nucleic acid sequences of the embodiments may be expressed in both monocotyledonous and dicotyledonous plant

species, sequences can be modified to account for the specific codon preferences and GC content preferences of monocotyledons or dicotyledons as these preferences have been shown to differ (Murray *et al.* (1989) *Nucleic Acids Res.* 17:477-498). Thus, the maize-preferred codon for a particular amino acid may be derived from known gene sequences from maize. Maize codon usage for 28 genes from maize plants is listed in Table 4 of Murray, *et al.*, *supra*. Methods are available in the art for synthesizing plant-preferred genes. See, for example, US Patent Numbers 5,380,831, and 5,436,391 and Murray, *et al.*, (1989) *Nucleic Acids Res.* 17:477-498, herein incorporated by reference.

Where appropriate, a nucleic acid may be optimized for increased expression in the host organism. Thus, where the host organism is a plant, the synthetic nucleic acids can be synthesized using plant-preferred codons for improved expression. See, for example, Campbell and Gowri, (1990) *Plant Physiol.* 92:1-11 for a discussion of host-preferred codon usage. For example, although nucleic acid sequences of the embodiments may be expressed in both monocotyledonous and dicotyledonous plant species, sequences can be modified to account for the specific codon preferences and GC content preferences of monocotyledons or dicotyledons as these preferences have been shown to differ (Murray *et al.* (1989) *Nucleic Acids Res.* 17:477-498). Thus, the maize-preferred codon for a particular amino acid may be derived from known gene sequences from maize. Maize codon usage for 28 genes from maize plants is listed in Table 4 of Murray, *et al.*, *supra*. Methods are available in the art for synthesizing plant-preferred genes. See, for example, US Patent Numbers 5,380,831, and 5,436,391 and Murray, *et al.*, (1989) *Nucleic Acids Res.* 17:477-498, and Liu H et al. *Mol Bio Rep* 37:677-684, 2010, herein incorporated by reference. A *Zea mays* codon usage table can be also found at kazusa.or.jp/codon/cgi-bin/showcodon.cgi?species=4577, which can be accessed using the www prefix. Table 2 shows a maize optimal codon analysis (adapted from Liu H et al. *Mol Bio Rep* 37:677-684, 2010).

Table 2

Amino Acid	Codon	High Count	RSCU	Low Count	RSCU	Amino Acid	Codon	High Count	RSCU	Low Count	RSCU
Phe	UUU	115	0.04	2,301	1.22	Ala	GCU	629	0.17	3,063	1.59
	UUC*	5,269	1.96	1,485	0.78		GCC*	8,057	2.16	1,136	0.59
Ser	UCU	176	0.13	2,498	1.48		GCA	369	0.1	2,872	1.49
	UCC*	3,489	2.48	1,074	0.63		GCG*	5,835	1.57	630	0.33
	UCA	104	0.07	2,610	1.54	Tyr	UAU	71	0.04	1,632	1.22
	UCG*	1,975	1.4	670	0.4		UAC*	3,841	1.96	1,041	0.78
	AGU	77	0.05	1,788	1.06	His	CAU	131	0.09	1,902	1.36
	AGC*	2,617	1.86	1,514	0.89		CAC*	2,800	1.91	897	0.64
Leu	UUA	10	0.01	1,326	0.79	Cys	UGU	52	0.04	1,233	1.12
	UUG	174	0.09	2,306	1.37		UGC*	2,291	1.96	963	0.88

	CUU	223	0.11	2,396	1.43	Gln	CAA	99	0.05	2,312	1.04
	CUC*	5,979	3.08	1,109	0.66		CAG*	3,557	1.95	2,130	0.96
	CUA	106	0.05	1,280	0.76	Arg	CGU	153	0.12	751	0.74
	CUG*	5,161	2.66	1,646	0.98		CGC*	4,278	3.25	466	0.46
Pro	CCU	427	0.22	1,900	1.47		CGA	92	0.07	659	0.65
	CCC*	3,035	1.59	601	0.47		CGG*	1,793	1.36	631	0.62
	CCA	311	0.16	2,140	1.66		AGA	83	0.06	1,948	1.91
	CCG*	3,846	2.02	513	0.4		AGG*	1,493	1.14	1,652	1.62
Ile	AUU	138	0.09	2,388	1.3	Asn	AAU	131	0.07	3,074	1.26
	AUC*	4,380	2.85	1,353	0.74		AAC*	3,814	1.93	1,807	0.74
	AUA	88	0.06	1,756	0.96	Lys	AAA	130	0.05	3,215	0.98
Thr	ACU	136	0.09	1,990	1.43		AAG*	5,047	1.95	3,340	1.02
	ACC*	3,398	2.25	991	0.71	Asp	GAU	312	0.09	4,217	1.38
	ACA	133	0.09	2,075	1.5		GAC*	6,729	1.91	1,891	0.62
	ACG*	2,378	1.57	495	0.36	Gly	GGU	363	0.13	2,301	1.35
Val	GUU	182	0.07	2,595	1.51		GGC*	7,842	2.91	1,282	0.75
	GUC*	4,584	1.82	1,096	0.64		GGA	397	0.15	2,044	1.19
	GUA	74	0.03	1,325	0.77		GGG*	2,186	0.81	1,215	0.71
	GUG*	5,257	2.08	1,842	1.07	Glu	GAA	193	0.06	4,080	1.1
							GAG*	6,010	1.94	3,307	0.9

Codon usage was compared using Chi squared contingency test to identify optimal codons.

Codons that occur significantly more often ($P \leq 0.01$) are indicated with an asterisk.

- 5 A *Glycine max* codon usage table is shown in Table 3 and can also be found at kazusa.or.jp/codon/cgi-bin/showcodon.cgi?species=3847&aa=1&style=N, which can be accessed using the www prefix.

Table 3

TTT	F	21.2	(10493)	TCT	S	18.4	(9107)
TTC	F	21.2	(10487)	TCC	S	12.9	(6409)
TTA	L	9.2	(4545)	TCA	S	15.6	(7712)
TTG	L	22.9	(11340)	TCG	S	4.8	(2397)
CTT	L	23.9	(11829)	CCT	P	18.9	(9358)
CTC	L	17.1	(8479)	CCC	P	10.1	(5010)
CTA	L	8.5	(4216)	CCA	P	19.1	(9461)
CTG	L	12.7	(6304)	CCG	P	4.7	(2312)
ATT	I	25.1	(12411)	ACT	T	17.1	(8490)
ATC	I	16.3	(8071)	ACC	T	14.3	(7100)
ATA	I	12.9	(6386)	ACA	T	14.9	(7391)
ATG	M	22.7	(11218)	ACG	T	4.3	(2147)
GTT	V	26.1	(12911)	GCT	A	26.7	(13201)
GTC	V	11.9	(5894)	GCC	A	16.2	(8026)
GTA	V	7.7	(3803)	GCA	A	21.4	(10577)
GTG	V	21.4	(10610)	GCG	A	6.3	(3123)
TAT	Y	15.7	(7779)	TGT	C	8.1	(3995)
TAC	Y	14.9	(7367)	TGC	C	8.0	(3980)
TAA	*	0.9	(463)	TGA	*	1.0	(480)

TAG	*	0.5	(263)	TGG	W	13.0	(6412)
CAT	H	14.0	(6930)	CGT	R	6.6	(3291)
CAC	H	11.6	(5759)	CGC	R	6.2	(3093)
CAA	Q	20.5	(10162)	CGA	R	4.1	(2018)
CAG	Q	16.2	(8038)	CGG	R	3.1	(1510)
AAT	N	22.4	(11088)	AGT	S	12.6	(6237)
AAC	N	22.8	(11284)	AGC	S	11.3	(5594)
AAA	K	26.9	(13334)	AGA	R	14.8	(7337)
AAG	K	35.9	(17797)	AGG	R	13.3	(6574)
GAT	D	32.4	(16040)	GGT	G	20.9	(10353)
GAC	D	20.4	(10097)	GGC	G	13.4	(6650)
GAA	E	33.2	(16438)	GGA	G	22.3	(11022)
GAG	E	33.2	(16426)	GGG	G	13.0	(6431)

In some embodiments the recombinant nucleic acid molecule encoding a PHI-4 polypeptide has maize optimized codons.

Additional sequence modifications are known to enhance gene expression in a cellular host. These include elimination of sequences encoding spurious polyadenylation signals, exon-intron splice site signals, transposon-like repeats, and other well-characterized sequences that may be deleterious to gene expression. The GC content of the sequence may be adjusted to levels average for a given cellular host, as calculated by reference to known genes expressed in the host cell. The term "host cell" as used herein refers to a cell which contains a vector and supports the replication and/or expression of the expression vector is intended. Host cells may be prokaryotic cells such as *E. coli* or eukaryotic cells such as yeast, insect, amphibian or mammalian cells or monocotyledonous or dicotyledonous plant cells. An example of a monocotyledonous host cell is a maize host cell. When possible, the sequence is modified to avoid predicted hairpin secondary mRNA structures.

The expression cassettes may additionally contain 5' leader sequences. Such leader sequences can act to enhance translation. Translation leaders are known in the art and include: picornavirus leaders, for example, EMCV leader (Encephalomyocarditis 5' noncoding region) (Elroy-Stein, *et al.*, (1989) *Proc. Natl. Acad. Sci. USA* 86:6126-6130); potyvirus leaders, for example, TEV leader (Tobacco Etch Virus) (Gallie, *et al.*, (1995) *Gene* 165(2):233-238), MDMV leader (Maize Dwarf Mosaic Virus), human immunoglobulin heavy-chain binding protein (BiP) (Macejak, *et al.*, (1991) *Nature* 353:90-94); untranslated leader from the coat protein mRNA of alfalfa mosaic virus (AMV RNA 4) (Jobling, *et al.*, (1987) *Nature* 325:622-625); tobacco mosaic virus leader (TMV) (Gallie, *et al.*, (1989) in *Molecular Biology of RNA*, ed. Cech (Liss, New York), pp. 237-256) and maize chlorotic mottle virus leader (MCMV) (Lommel, *et al.*, (1991) *Virology* 81:382-385).

See also, Della-Cioppa, *et al.*, (1987) *Plant Physiol.* 84:965-968. Such constructs may also contain a "signal sequence" or "leader sequence" to facilitate co-translational or post-translational transport of the peptide to certain intracellular structures such as the chloroplast (or other plastid), endoplasmic reticulum or Golgi apparatus.

5 By "signal sequence" is intended a sequence that is known or suspected to result in cotranslational or post-translational peptide transport across the cell membrane. In eukaryotes, this typically involves secretion into the Golgi apparatus, with some resulting glycosylation. Insecticidal toxins of bacteria are often synthesized as protoxins, which are protolytically activated in the gut of the target pest (Chang, (1987) *Methods Enzymol.*
10 153:507-516). In some embodiments, the signal sequence is located in the native sequence or may be derived from a sequence of the embodiments. By "leader sequence" is intended any sequence that when translated, results in an amino acid sequence sufficient to trigger co-translational transport of the peptide chain to a subcellular organelle. Thus, this includes leader sequences targeting transport and/or glycosylation
15 by passage into the endoplasmic reticulum, passage to vacuoles, plastids including chloroplasts, mitochondria and the like. Nuclear-encoded proteins targeted to the chloroplast thylakoid lumen compartment have a characteristic bipartite transit peptide, composed of a stromal targeting signal peptide and a lumen targeting signal peptide. The stromal targeting information is in the amino-proximal portion of the transit peptide. The
20 lumen targeting signal peptide is in the carboxyl-proximal portion of the transit peptide, and contains all the information for targeting to the lumen. Recent research in proteomics of the higher plant chloroplast has achieved in the identification of numerous nuclear-encoded lumen proteins (Kieselbach et al. *FEBS LETT* 480:271-276, 2000; Peltier et al. *Plant Cell* 12:319-341, 2000; Bricker et al. *Biochim. Biophys Acta* 1503:350-356, 2001),
25 the lumen targeting signal peptide of which can potentially be used in accordance with the present disclosure. About 80 proteins from *Arabidopsis*, as well as homologous proteins from spinach and garden pea, are reported by Kieselbach et al., *Photosynthesis Research*, 78:249-264, 2003. In particular, table 2 of this publication, which is incorporated into the description herewith by reference, discloses 85 proteins from the
30 chloroplast lumen, identified by their accession number (see also US Patent Application Publication 2009/09044298). In addition, the recently published draft version of the rice genome (Goff et al, *Science* 296:92-100, 2002) is a suitable source for lumen targeting signal peptide which may be used in accordance with the present disclosure.

Suitable chloroplast transit peptides (CTP) are well known to one skilled in the art
35 also include chimeric CTPs comprising but not limited to, an N-terminal domain, a central domain or a C-terminal domain from a CTP from *Oryza sativa* 1-deoxy-D xyulose-5-

Phosphate Synthase, *Oryza sativa*-Superoxide dismutase, *Oryza sativa*-soluble starch synthase, *Oryza sativa*-NADP-dependent Malic acid enzyme, *Oryza sativa*-Phospho-2-dehydro-3-deoxyheptonate Aldolase 2, *Oryza sativa*-L-Ascorbate peroxidase 5, *Oryza sativa*-Phosphoglucan water dikinase, *Zea Mays* ssRUBISCO, *Zea Mays*-beta-glucosidase, *Zea Mays*-Malate dehydrogenase, *Zea Mays* Thioredoxin M-type US Patent Application Publication 2012/0304336).

The PHI-4 polypeptide gene to be targeted to the chloroplast may be optimized for expression in the chloroplast to account for differences in codon usage between the plant nucleus and this organelle. In this manner, the nucleic acids of interest may be synthesized using chloroplast-preferred codons. See, for example, US Patent Number 5,380,831, herein incorporated by reference.

In preparing the expression cassette, the various DNA fragments may be manipulated so as to provide for the DNA sequences in the proper orientation and, as appropriate, in the proper reading frame. Toward this end, adapters or linkers may be employed to join the DNA fragments or other manipulations may be involved to provide for convenient restriction sites, removal of superfluous DNA, removal of restriction sites or the like. For this purpose, *in vitro* mutagenesis, primer repair, restriction, annealing, resubstitutions, e.g., transitions and transversions, may be involved.

A number of promoters can be used in the practice of the embodiments. The promoters can be selected based on the desired outcome. The nucleic acids can be combined with constitutive, tissue-preferred, inducible or other promoters for expression in the host organism. Suitable constitutive promoters for use in a plant host cell include, for example, the core promoter of the Rsyn7 promoter and other constitutive promoters disclosed in WO 1999/43838 and US Patent Number 6,072,050; the core CaMV 35S promoter (Odell, *et al.*, (1985) *Nature* 313:810-812); rice actin (McElroy, *et al.*, (1990) *Plant Cell* 2:163-171); ubiquitin (Christensen, *et al.*, (1989) *Plant Mol. Biol.* 12:619-632 and Christensen, *et al.*, (1992) *Plant Mol. Biol.* 18:675-689); pEMU (Last, *et al.*, (1991) *Theor. Appl. Genet.* 81:581-588); MAS (Velten, *et al.*, (1984) *EMBO J.* 3:2723-2730); ALS promoter (US Patent Number 5,659,026) and the like. Other constitutive promoters include, for example, those discussed in US Patent Numbers 5,608,149; 5,608,144; 5,604,121; 5,569,597; 5,466,785; 5,399,680; 5,268,463; 5,608,142 and 6,177,611.

Depending on the desired outcome, it may be beneficial to express the gene from an inducible promoter. Of particular interest for regulating the expression of the nucleotide sequences of the embodiments in plants are wound-inducible promoters. Such wound-inducible promoters, may respond to damage caused by insect feeding, and include potato proteinase inhibitor (pin II) gene (Ryan, (1990) *Ann. Rev. Phytopath.*

28:425-449; Duan, *et al.*, (1996) *Nature Biotechnology* 14:494-498); *wun1* and *wun2*, US Patent Number 5,428,148; *win1* and *win2* (Stanford, *et al.*, (1989) *Mol. Gen. Genet.* 215:200-208); *systemin* (McGurl, *et al.*, (1992) *Science* 225:1570-1573); *WIP1* (Rohmeier, *et al.*, (1993) *Plant Mol. Biol.* 22:783-792; Eckelkamp, *et al.*, (1993) *FEBS Letters* 323:73-76); *MPI* gene (Corderok, *et al.*, (1994) *Plant J.* 6(2):141-150) and the like, herein incorporated by reference.

Additionally, pathogen-inducible promoters may be employed in the methods and nucleotide constructs of the embodiments. Such pathogen-inducible promoters include those from pathogenesis-related proteins (PR proteins), which are induced following infection by a pathogen; e.g., PR proteins, SAR proteins, beta-1,3-glucanase, chitinase, etc. See, for example, Redolfi, *et al.*, (1983) *Neth. J. Plant Pathol.* 89:245-254; Uknes, *et al.*, (1992) *Plant Cell* 4:645-656 and Van Loon, (1985) *Plant Mol. Virol.* 4:111-116. See also, WO 1999/43819, herein incorporated by reference.

Of interest are promoters that are expressed locally at or near the site of pathogen infection. See, for example, Marineau, *et al.*, (1987) *Plant Mol. Biol.* 9:335-342; Matton, *et al.*, (1989) *Molecular Plant-Microbe Interactions* 2:325-331; Somsisch, *et al.*, (1986) *Proc. Natl. Acad. Sci. USA* 83:2427-2430; Somsisch, *et al.*, (1988) *Mol. Gen. Genet.* 2:93-98 and Yang, (1996) *Proc. Natl. Acad. Sci. USA* 93:14972-14977. See also, Chen, *et al.*, (1996) *Plant J.* 10:955-966; Zhang, *et al.*, (1994) *Proc. Natl. Acad. Sci. USA* 91:2507-2511; Warner, *et al.*, (1993) *Plant J.* 3:191-201; Siebertz, *et al.*, (1989) *Plant Cell* 1:961-968; US Patent Number 5,750,386 (nematode-inducible) and the references cited therein. Of particular interest is the inducible promoter for the maize PRms gene, whose expression is induced by the pathogen *Fusarium moniliforme* (see, for example, Cordero, *et al.*, (1992) *Physiol. Mol. Plant Path.* 41:189-200).

Chemical-regulated promoters can be used to modulate the expression of a gene in a plant through the application of an exogenous chemical regulator. Depending upon the objective, the promoter may be a chemical-inducible promoter, where application of the chemical induces gene expression or a chemical-repressible promoter, where application of the chemical represses gene expression. Chemical-inducible promoters are known in the art and include, but are not limited to, the maize *ln2-2* promoter, which is activated by benzenesulfonamide herbicide safeners, the maize *GST* promoter, which is activated by hydrophobic electrophilic compounds that are used as pre-emergent herbicides, and the tobacco *PR-1a* promoter, which is activated by salicylic acid. Other chemical-regulated promoters of interest include steroid-responsive promoters (see, for example, the glucocorticoid-inducible promoter in Schena, *et al.*, (1991) *Proc. Natl. Acad. Sci. USA* 88:10421-10425 and McNellis, *et al.*, (1998) *Plant J.* 14(2):247-257) and

tetracycline-inducible and tetracycline-repressible promoters (see, for example, Gatz, *et al.*, (1991) *Mol. Gen. Genet.* 227:229-237 and US Patent Numbers 5,814,618 and 5,789,156), herein incorporated by reference.

Tissue-preferred promoters can be utilized to target enhanced PHI-4 polypeptide expression within a particular plant tissue. Tissue-preferred promoters include those discussed in Yamamoto, *et al.*, (1997) *Plant J.* 12(2):255-265; Kawamata, *et al.*, (1997) *Plant Cell Physiol.* 38(7):792-803; Hansen, *et al.*, (1997) *Mol. Gen Genet.* 254(3):337-343; Russell, *et al.*, (1997) *Transgenic Res.* 6(2):157-168; Rinehart, *et al.*, (1996) *Plant Physiol.* 112(3):1331-1341; Van Camp, *et al.*, (1996) *Plant Physiol.* 112(2):525-535; Canevascini, *et al.*, (1996) *Plant Physiol.* 112(2):513-524; Yamamoto, *et al.*, (1994) *Plant Cell Physiol.* 35(5):773-778; Lam, (1994) *Results Probl. Cell Differ.* 20:181-196; Orozco, *et al.*, (1993) *Plant Mol Biol.* 23(6):1129-1138; Matsuoka, *et al.*, (1993) *Proc Natl. Acad. Sci. USA* 90(20):9586-9590 and Guevara-Garcia, *et al.*, (1993) *Plant J.* 4(3):495-505. Such promoters can be modified, if necessary, for weak expression.

Leaf-preferred promoters are known in the art. See, for example, Yamamoto, *et al.*, (1997) *Plant J.* 12(2):255-265; Kwon, *et al.*, (1994) *Plant Physiol.* 105:357-67; Yamamoto, *et al.*, (1994) *Plant Cell Physiol.* 35(5):773-778; Gotor, *et al.*, (1993) *Plant J.* 3:509-18; Orozco, *et al.*, (1993) *Plant Mol. Biol.* 23(6):1129-1138 and Matsuoka, *et al.*, (1993) *Proc. Natl. Acad. Sci. USA* 90(20):9586-9590.

Root-preferred or root-specific promoters are known and can be selected from the many available from the literature or isolated *de novo* from various compatible species. See, for example, Hire, *et al.*, (1992) *Plant Mol. Biol.* 20(2):207-218 (soybean root-specific glutamine synthetase gene); Keller and Baumgartner, (1991) *Plant Cell* 3(10):1051-1061 (root-specific control element in the GRP 1.8 gene of French bean); Sanger, *et al.*, (1990) *Plant Mol. Biol.* 14(3):433-443 (root-specific promoter of the mannopine synthase (MAS) gene of *Agrobacterium tumefaciens*) and Miao, *et al.*, (1991) *Plant Cell* 3(1):11-22 (full-length cDNA clone encoding cytosolic glutamine synthetase (GS), which is expressed in roots and root nodules of soybean). See also, Bogusz, *et al.*, (1990) *Plant Cell* 2(7):633-641, where two root-specific promoters isolated from hemoglobin genes from the nitrogen-fixing nonlegume *Parasponia andersonii* and the related non-nitrogen-fixing nonlegume *Trema tomentosa* are described. The promoters of these genes were linked to a β -glucuronidase reporter gene and introduced into both the nonlegume *Nicotiana tabacum* and the legume *Lotus corniculatus*, and in both instances root-specific promoter activity was preserved. Leach and Aoyagi, (1991) describe their analysis of the promoters of the highly expressed rolC and rolD root-inducing genes of *Agrobacterium rhizogenes* (see, *Plant Science* (Limerick) 79(1):69-76). They concluded that enhancer and tissue-

preferred DNA determinants are dissociated in those promoters. Teeri, *et al.*, (1989) used gene fusion to lacZ to show that the *Agrobacterium* T-DNA gene encoding octopine synthase is especially active in the epidermis of the root tip and that the TR2' gene is root specific in the intact plant and stimulated by wounding in leaf tissue, an especially desirable combination of characteristics for use with an insecticidal or larvicidal gene (see, *EMBO J.* 8(2):343-350). The TR1' gene fused to *nptII* (neomycin phosphotransferase II) showed similar characteristics. Additional root-preferred promoters include the VfENOD-GRP3 gene promoter (Kuster, *et al.*, (1995) *Plant Mol. Biol.* 29(4):759-772) and rolB promoter (Capana, *et al.*, (1994) *Plant Mol. Biol.* 25(4):681-691. See also, US Patent Numbers 5,837,876; 5,750,386; 5,633,363; 5,459,252; 5,401,836; 5,110,732 and 5,023,179.

"Seed-preferred" promoters include both "seed-specific" promoters (those promoters active during seed development such as promoters of seed storage proteins) as well as "seed-germinating" promoters (those promoters active during seed germination). See, Thompson, *et al.*, (1989) *BioEssays* 10:108, herein incorporated by reference. Such seed-preferred promoters include, but are not limited to, Cim1 (cytokinin-induced message); cZ19B1 (maize 19 kDa zein); and milps (myo-inositol-1-phosphate synthase) (see, US Patent Number 6,225,529, herein incorporated by reference). Gamma-zein and Glb-1 are endosperm-specific promoters. For dicots, seed-specific promoters include, but are not limited to, Kunitz trypsin inhibitor 3 (KTI3) (Jofuku, K.D. and Goldberg, R.B. *Plant Cell* 1:1079-1093, 1989), bean β -phaseolin, napin, β -conglycinin, glycinin 1, soybean lectin, cruciferin, and the like. For monocots, seed-specific promoters include, but are not limited to, maize 15 kDa zein, 22 kDa zein, 27 kDa zein, g-zein, waxy, shrunken 1, shrunken 2, globulin 1, etc. See also, WO 2000/12733, where seed-preferred promoters from *end1* and *end2* genes are disclosed; herein incorporated by reference. In dicots, seed specific promoters include but are not limited to seed coat promoter from *Arabidopsis*, pBAN; and the early seed promoters from *Arabidopsis*, p26, p63, and p63tr (US Patent Numbers 7,294,760 and 7,847,153). A promoter that has "preferred" expression in a particular tissue is expressed in that tissue to a greater degree than in at least one other plant tissue. Some tissue-preferred promoters show expression almost exclusively in the particular tissue.

Where low level expression is desired, weak promoters will be used. Generally, the term "weak promoter" as used herein refers to a promoter that drives expression of a coding sequence at a low level. By low level expression at levels of about 1/1000 transcripts to about 1/100,000 transcripts to about 1/500,000 transcripts is intended. Alternatively, it is recognized that the term "weak promoters" also encompasses

promoters that drive expression in only a few cells and not in others to give a total low level of expression. Where a promoter drives expression at unacceptably high levels, portions of the promoter sequence can be deleted or modified to decrease expression levels.

5 Such weak constitutive promoters include, for example the core promoter of the Rsyn7 promoter (WO 1999/43838 and US Patent Number 6,072,050), the core 35S CaMV promoter, and the like. Other constitutive promoters include, for example, those disclosed in US Patent Numbers 5,608,149; 5,608,144; 5,604,121; 5,569,597; 5,466,785; 5,399,680; 5,268,463; 5,608,142 and 6,177,611, herein incorporated by reference.

10 The above list of promoters is not meant to be limiting. Any appropriate promoter can be used in the embodiments.

 Generally, the expression cassette will comprise a selectable marker gene for the selection of transformed cells. Selectable marker genes are utilized for the selection of transformed cells or tissues. Marker genes include genes encoding antibiotic resistance, such as those encoding neomycin phosphotransferase II (NEO) and hygromycin phosphotransferase (HPT), as well as genes conferring resistance to herbicidal compounds, such as glufosinate ammonium, bromoxynil, imidazolinones and 2,4-dichlorophenoxyacetate (2,4-D). Additional examples of suitable selectable marker genes include, but are not limited to, genes encoding resistance to chloramphenicol (Herrera Estrella, *et al.*, (1983) *EMBO J.* 2:987-992); methotrexate (Herrera Estrella, *et al.*, (1983) *Nature* 303:209-213 and Meijer, *et al.*, (1991) *Plant Mol. Biol.* 16:807-820); streptomycin (Jones, *et al.*, (1987) *Mol. Gen. Genet.* 210:86-91); spectinomycin (Bretagne-Sagnard, *et al.*, (1996) *Transgenic Res.* 5:131-137); bleomycin (Hille, *et al.*, (1990) *Plant Mol. Biol.* 7:171-176); sulfonamide (Guerineau, *et al.*, (1990) *Plant Mol. Biol.* 15:127-136); bromoxynil (Stalker, *et al.*, (1988) *Science* 242:419-423); glyphosate (Shaw, *et al.*, (1986) *Science* 233:478-481 and US Patent Application Serial Numbers 10/004,357 and 10/427,692); phosphinothricin (DeBlock, *et al.*, (1987) *EMBO J.* 6:2513-2518). See generally, Yarranton, (1992) *Curr. Opin. Biotech.* 3:506-511; Christopherson, *et al.*, (1992) *Proc. Natl. Acad. Sci. USA* 89:6314-6318; Yao, *et al.*, (1992) *Cell* 71:63-72; Reznikoff, (1992) *Mol. Microbiol.* 6:2419-2422; Barkley, *et al.*, (1980) in *The Operon*, pp. 177-220; Hu, *et al.*, (1987) *Cell* 48:555-566; Brown, *et al.*, (1987) *Cell* 49:603-612; Figge, *et al.*, (1988) *Cell* 52:713-722; Deuschle, *et al.*, (1989) *Proc. Natl. Acad. Sci. USA* 86:5400-5404; Fuerst, *et al.*, (1989) *Proc. Natl. Acad. Sci. USA* 86:2549-2553; Deuschle, *et al.*, (1990) *Science* 248:480-483; Gossen, (1993) Ph.D. Thesis, University of Heidelberg; Reines, *et al.*, (1993) *Proc. Natl. Acad. Sci. USA* 90:1917-1921; Labow, *et al.*, (1990) *Mol. Cell. Biol.* 10:3343-3356; Zambretti, *et al.*, (1992) *Proc. Natl. Acad. Sci. USA* 89:3952-3956; Baim, *et al.*, (1991) *Proc. Natl. Acad. Sci. USA* 88:5072-5076;

Wyborski, *et al.*, (1991) *Nucleic Acids Res.* 19:4647-4653; Hillenand-Wissman, (1989) *Topics Mol. Struc. Biol.* 10:143-162; Degenkolb, *et al.*, (1991) *Antimicrob. Agents Chemother.* 35:1591-1595; Kleinschmidt, *et al.*, (1988) *Biochemistry* 27:1094-1104; Bonin, (1993) Ph.D. Thesis, University of Heidelberg; Gossen, *et al.*, (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Oliva, *et al.*, (1992) *Antimicrob. Agents Chemother.* 36:913-919; Hlavka, *et al.*, (1985) *Handbook of Experimental Pharmacology*, Vol. 78 (Springer-Verlag, Berlin) and Gill, *et al.*, (1988) *Nature* 334:721-724. Such disclosures are herein incorporated by reference.

The above list of selectable marker genes is not meant to be limiting. Any selectable marker gene can be used in the embodiments.

Plant Transformation

The methods of the embodiments involve introducing a polypeptide or polynucleotide into a plant. "Introducing" is intended to mean presenting to the plant the polynucleotide or polypeptide in such a manner that the sequence gains access to the interior of a cell of the plant. The methods of the embodiments do not depend on a particular method for introducing a polynucleotide or polypeptide into a plant, only that the polynucleotide or polypeptides gains access to the interior of at least one cell of the plant. Methods for introducing polynucleotide or polypeptides into plants are known in the art including, but not limited to, stable transformation methods, transient transformation methods and virus-mediated methods.

"Stable transformation" is intended to mean that the nucleotide construct introduced into a plant integrates into the genome of the plant and is capable of being inherited by the progeny thereof. "Transient transformation" is intended to mean that a polynucleotide is introduced into the plant and does not integrate into the genome of the plant or a polypeptide is introduced into a plant. By "plant" is intended whole plants, plant organs (e.g., leaves, stems, roots, etc.), seeds, plant cells, propagules, embryos and progeny of the same. Plant cells can be differentiated or undifferentiated (e.g. callus, suspension culture cells, protoplasts, leaf cells, root cells, phloem cells, and pollen).

Transformation protocols as well as protocols for introducing nucleotide sequences into plants may vary depending on the type of plant or plant cell, i.e., monocot or dicot, targeted for transformation. Suitable methods of introducing nucleotide sequences into plant cells and subsequent insertion into the plant genome include microinjection (Crossway, *et al.*, (1986) *Biotechniques* 4:320-334), electroporation (Riggs, *et al.*, (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-5606), *Agrobacterium*-mediated transformation (US Patent Numbers 5,563,055 and 5,981,840), direct gene transfer

(Paszkowski, *et al.*, (1984) *EMBO J.* 3:2717-2722) and ballistic particle acceleration (see, for example, US Patent Numbers 4,945,050; 5,879,918; 5,886,244 and 5,932,782; Tomes, *et al.*, (1995) in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg and Phillips, (Springer-Verlag, Berlin) and McCabe, *et al.*, (1988) *Biotechnology* 6:923-926) and Lecl transformation (WO 2000/28058). For potato transformation see, Tu, *et al.*, (1998) *Plant Molecular Biology* 37:829-838 and Chong, *et al.*, (2000) *Transgenic Research* 9:71-78. Additional transformation procedures can be found in Weissinger, *et al.*, (1988) *Ann. Rev. Genet.* 22:421-477; Sanford, *et al.*, (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou, *et al.*, (1988) *Plant Physiol.* 87:671-674 (soybean); McCabe, *et al.*, (1988) *Bio/Technology* 6:923-926 (soybean); Finer and McMullen, (1991) *In Vitro Cell Dev. Biol.* 27P:175-182 (soybean); Singh, *et al.*, (1998) *Theor. Appl. Genet.* 96:319-324 (soybean); Datta, *et al.*, (1990) *Biotechnology* 8:736-740 (rice); Klein, *et al.*, (1988) *Proc. Natl. Acad. Sci. USA* 85:4305-4309 (maize); Klein, *et al.*, (1988) *Biotechnology* 6:559-563 (maize); US Patent Numbers 5,240,855; 5,322,783 and 5,324,646; Klein, *et al.*, (1988) *Plant Physiol.* 91:440-444 (maize); Fromm, *et al.*, (1990) *Biotechnology* 8:833-839 (maize); Hooykaas-Van Slogteren, *et al.*, (1984) *Nature (London)* 311:763-764; US Patent Number 5,736,369 (cereals); Bytebier, *et al.*, (1987) *Proc. Natl. Acad. Sci. USA* 84:5345-5349 (Liliaceae); De Wet, *et al.*, (1985) in *The Experimental Manipulation of Ovule Tissues*, ed. Chapman, *et al.*, (Longman, New York), pp. 197-209 (pollen); Kaeppler, *et al.*, (1990) *Plant Cell Reports* 9:415-418 and Kaeppler, *et al.*, (1992) *Theor. Appl. Genet.* 84:560-566 (whisker-mediated transformation); D'Halluin, *et al.*, (1992) *Plant Cell* 4:1495-1505 (electroporation); Li, *et al.*, (1993) *Plant Cell Reports* 12:250-255 and Christou and Ford, (1995) *Annals of Botany* 75:407-413 (rice); Osjoda, *et al.*, (1996) *Nature Biotechnology* 14:745-750 (maize via *Agrobacterium tumefaciens*); all of which are herein incorporated by reference.

In specific embodiments, the sequences of the embodiments can be provided to a plant using a variety of transient transformation methods. Such transient transformation methods include, but are not limited to, the introduction of the PHI-4 polypeptide or variants and fragments thereof directly into the plant or the introduction of the PHI-4 polypeptide transcript into the plant. Such methods include, for example, microinjection or particle bombardment. See, for example, Crossway, *et al.*, (1986) *Mol Gen. Genet.* 202:179-185; Nomura, *et al.*, (1986) *Plant Sci.* 44:53-58; Hepler, *et al.*, (1994) *Proc. Natl. Acad. Sci.* 91:2176-2180 and Hush, *et al.*, (1994) *The Journal of Cell Science* 107:775-784, all of which are herein incorporated by reference. Alternatively, the PHI-4 polypeptide polynucleotide can be transiently transformed into the plant using techniques known in the art. Such techniques include viral vector system and the precipitation of the

polynucleotide in a manner that precludes subsequent release of the DNA. Thus, transcription from the particle-bound DNA can occur, but the frequency with which it is released to become integrated into the genome is greatly reduced. Such methods include the use of particles coated with polyethylimine (PEI; Sigma #P3143).

5 Methods are known in the art for the targeted insertion of a polynucleotide at a specific location in the plant genome. In one embodiment, the insertion of the polynucleotide at a desired genomic location is achieved using a site-specific recombination system. See, for example, WO 1999/25821, WO 1999/25854, WO 1999/25840, WO 1999/25855 and WO 1999/25853, all of which are herein incorporated
10 by reference. Briefly, the polynucleotide of the embodiments can be contained in transfer cassette flanked by two non-identical recombination sites. The transfer cassette is introduced into a plant have stably incorporated into its genome a target site which is flanked by two non-identical recombination sites that correspond to the sites of the transfer cassette. An appropriate recombinase is provided and the transfer cassette is
15 integrated at the target site. The polynucleotide of interest is thereby integrated at a specific chromosomal position in the plant genome.

Plant transformation vectors may be comprised of one or more DNA vectors needed for achieving plant transformation. For example, it is a common practice in the art to utilize plant transformation vectors that are comprised of more than one contiguous
20 DNA segment. These vectors are often referred to in the art as “binary vectors”. Binary vectors as well as vectors with helper plasmids are most often used for *Agrobacterium*-mediated transformation, where the size and complexity of DNA segments needed to achieve efficient transformation is quite large, and it is advantageous to separate functions onto separate DNA molecules. Binary vectors typically contain a plasmid vector
25 that contains the cis-acting sequences required for T-DNA transfer (such as left border and right border), a selectable marker that is engineered to be capable of expression in a plant cell, and a “gene of interest” (a gene engineered to be capable of expression in a plant cell for which generation of transgenic plants is desired). Also present on this plasmid vector are sequences required for bacterial replication. The cis-acting sequences
30 are arranged in a fashion to allow efficient transfer into plant cells and expression therein. For example, the selectable marker gene and the pesticidal gene are located between the left and right borders. Often a second plasmid vector contains the trans-acting factors that mediate T-DNA transfer from *Agrobacterium* to plant cells. This plasmid often contains the virulence functions (Vir genes) that allow infection of plant cells by
35 *Agrobacterium*, and transfer of DNA by cleavage at border sequences and vir-mediated DNA transfer, as is understood in the art (Hellens and Mullineaux, (2000) *Trends in Plant*

Science 5:446-451). Several types of *Agrobacterium* strains (e.g. LBA4404, GV3101, EHA101, EHA105, etc.) can be used for plant transformation. The second plasmid vector is not necessary for transforming the plants by other methods such as microprojection, microinjection, electroporation, polyethylene glycol, etc.

5 In general, plant transformation methods involve transferring heterologous DNA into target plant cells (e.g., immature or mature embryos, suspension cultures, undifferentiated callus, protoplasts, etc.), followed by applying a maximum threshold level of appropriate selection (depending on the selectable marker gene) to recover the transformed plant cells from a group of untransformed cell mass. Following integration of
10 heterologous foreign DNA into plant cells, one then applies a maximum threshold level of appropriate selection in the medium to kill the untransformed cells and separate and proliferate the putatively transformed cells that survive from this selection treatment by transferring regularly to a fresh medium. By continuous passage and challenge with appropriate selection, one identifies and proliferates the cells that are transformed with
15 the plasmid vector. Molecular and biochemical methods can then be used to confirm the presence of the integrated heterologous gene of interest into the genome of the transgenic plant.

Explants are typically transferred to a fresh supply of the same medium and cultured routinely. Subsequently, the transformed cells are differentiated into shoots after
20 placing on regeneration medium supplemented with a maximum threshold level of selecting agent. The shoots are then transferred to a selective rooting medium for recovering rooted shoot or plantlet. The transgenic plantlet then grows into a mature plant and produces fertile seeds (e.g., Hiei, *et al.*, (1994) *The Plant Journal* 6:271-282; Ishida, *et al.*, (1996) *Nature Biotechnology* 14:745-750). Explants are typically transferred
25 to a fresh supply of the same medium and cultured routinely. A general description of the techniques and methods for generating transgenic plants are found in Ayres and Park, (1994) *Critical Reviews in Plant Science* 13:219-239 and Bommineni and Jauhar, (1997) *Maydica* 42:107-120. Since the transformed material contains many cells; both transformed and non-transformed cells are present in any piece of subjected target callus
30 or tissue or group of cells. The ability to kill non-transformed cells and allow transformed cells to proliferate results in transformed plant cultures. Often, the ability to remove non-transformed cells is a limitation to rapid recovery of transformed plant cells and successful generation of transgenic plants.

The cells that have been transformed may be grown into plants in accordance with
35 conventional ways. See, for example, McCormick, *et al.*, (1986) *Plant Cell Reports* 5:81-84. These plants may then be grown, and either pollinated with the same transformed

strain or different strains and the resulting hybrid having constitutive or inducible expression of the desired phenotypic characteristic identified. Two or more generations may be grown to ensure that expression of the desired phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure that expression of the
5 desired phenotypic characteristic has been achieved.

The nucleotide sequences of the embodiments may be provided to the plant by contacting the plant with a virus or viral nucleic acids. Generally, such methods involve incorporating the nucleotide construct of interest within a viral DNA or RNA molecule. It is recognized that the recombinant proteins of the embodiments may be initially synthesized
10 as part of a viral polyprotein, which later may be processed by proteolysis *in vivo* or *in vitro* to produce the desired PHI-4 polypeptide. It is also recognized that such a viral polyprotein, comprising at least a portion of the amino acid sequence of a PHI-4 polypeptide of the embodiments, may have the desired pesticidal activity. Such viral polyproteins and the nucleotide sequences that encode for them are encompassed by the
15 embodiments. Methods for providing plants with nucleotide constructs and producing the encoded proteins in the plants, which involve viral DNA or RNA molecules are known in the art. See, for example, US Patent Numbers 5,889,191; 5,889,190; 5,866,785; 5,589,367 and 5,316,931, herein incorporated by reference.

Methods for transformation of chloroplasts are known in the art. See, for example,
20 Svab, *et al.*, (1990) *Proc. Natl. Acad. Sci. USA* 87:8526-8530; Svab and Maliga, (1993) *Proc. Natl. Acad. Sci. USA* 90:913-917; Svab and Maliga, (1993) *EMBO J.* 12:601-606. The method relies on particle gun delivery of DNA containing a selectable marker and targeting of the DNA to the plastid genome through homologous recombination. Additionally, plastid transformation can be accomplished by transactivation of a silent
25 plastid-borne transgene by tissue-preferred expression of a nuclear-encoded and plastid-directed RNA polymerase. Such a system has been reported in McBride, *et al.*, (1994) *Proc. Natl. Acad. Sci. USA* 91:7301-7305.

The embodiments further relate to plant-propagating material of a transformed plant of the embodiments including, but not limited to, seeds, tubers, corms, bulbs, leaves,
30 and cuttings of roots and shoots.

The embodiments may be used for transformation of any plant species, including, but not limited to, monocots and dicots. Examples of plants of interest include, but are not limited to, corn (*Zea mays*), *Brassica* sp. (e.g., *B. napus*, *B. rapa*, *B. juncea*), particularly those *Brassica* species useful as sources of seed oil, alfalfa (*Medicago sativa*), rice (*Oryza sativa*), rye (*Secale cereale*), sorghum (*Sorghum bicolor*, *Sorghum vulgare*), millet (e.g.,
35 pearl millet (*Pennisetum glaucum*), proso millet (*Panicum miliaceum*), foxtail millet (*Setaria*

italica), finger millet (*Eleusine coracana*), sunflower (*Helianthus annuus*), safflower (*Carthamus tinctorius*), wheat (*Triticum aestivum*), soybean (*Glycine max*), tobacco (*Nicotiana tabacum*), potato (*Solanum tuberosum*), peanuts (*Arachis hypogaea*), cotton (*Gossypium barbadense*, *Gossypium hirsutum*), sweet potato (*Ipomoea batatas*), cassava (*Manihot esculenta*), coffee (*Coffea* spp.), coconut (*Cocos nucifera*), pineapple (*Ananas comosus*), citrus trees (*Citrus* spp.), cocoa (*Theobroma cacao*), tea (*Camellia sinensis*), banana (*Musa* spp.), avocado (*Persea americana*), fig (*Ficus casica*), guava (*Psidium guajava*), mango (*Mangifera indica*), olive (*Olea europaea*), papaya (*Carica papaya*), cashew (*Anacardium occidentale*), macadamia (*Macadamia integrifolia*), almond (*Prunus amygdalus*), sugar beets (*Beta vulgaris*), sugarcane (*Saccharum* spp.), oats, barley, vegetables, ornamentals and conifers.

Vegetables include tomatoes (*Lycopersicon esculentum*), lettuce (e.g., *Lactuca sativa*), green beans (*Phaseolus vulgaris*), lima beans (*Phaseolus limensis*), peas (*Lathyrus* spp.), and members of the genus *Cucumis* such as cucumber (*C. sativus*), cantaloupe (*C. cantalupensis*), and musk melon (*C. melo*). Ornamentals include azalea (*Rhododendron* spp.), hydrangea (*Macrophylla hydrangea*), hibiscus (*Hibiscus rosasanensis*), roses (*Rosa* spp.), tulips (*Tulipa* spp.), daffodils (*Narcissus* spp.), petunias (*Petunia hybrida*), carnation (*Dianthus caryophyllus*), poinsettia (*Euphorbia pulcherrima*), and chrysanthemum. Conifers that may be employed in practicing the embodiments include, for example, pines such as loblolly pine (*Pinus taeda*), slash pine (*Pinus elliotii*), ponderosa pine (*Pinus ponderosa*), lodgepole pine (*Pinus contorta*) and Monterey pine (*Pinus radiata*); Douglas-fir (*Pseudotsuga menziesii*); Western hemlock (*Tsuga canadensis*); Sitka spruce (*Picea glauca*); redwood (*Sequoia sempervirens*); true firs such as silver fir (*Abies amabilis*) and balsam fir (*Abies balsamea*); and cedars such as Western red cedar (*Thuja plicata*) and Alaska yellow-cedar (*Chamaecyparis nootkatensis*). Plants of the embodiments include crop plants (for example, corn, alfalfa, sunflower, *Brassica*, soybean, cotton, safflower, peanut, sorghum, wheat, millet, tobacco, etc.), such as corn and soybean plants.

Turf grasses include, but are not limited to: annual bluegrass (*Poa annua*); annual ryegrass (*Lolium multiflorum*); Canada bluegrass (*Poa compressa*); Chewing's fescue (*Festuca rubra*); colonial bentgrass (*Agrostis tenuis*); creeping bentgrass (*Agrostis palustris*); crested wheatgrass (*Agropyron desertorum*); fairway wheatgrass (*Agropyron cristatum*); hard fescue (*Festuca longifolia*); Kentucky bluegrass (*Poa pratensis*); orchardgrass (*Dactylis glomerata*); perennial ryegrass (*Lolium perenne*); red fescue (*Festuca rubra*); redtop (*Agrostis alba*); rough bluegrass (*Poa trivialis*); sheep fescue (*Festuca ovina*); smooth brome grass (*Bromus inermis*); tall fescue (*Festuca arundinacea*); timothy (*Phleum pratense*); velvet bentgrass (*Agrostis canina*); weeping alkaligrass (*Puccinellia distans*); western

wheatgrass (*Agropyron smithii*); Bermuda grass (*Cynodon* spp.); St. Augustine grass (*Stenotaphrum secundatum*); zoysia grass (*Zoysia* spp.); Bahia grass (*Paspalum notatum*); carpet grass (*Axonopus affinis*); centipede grass (*Eremochloa ophiuroides*); kikuyu grass (*Pennisetum clandestinum*); seashore paspalum (*Paspalum vaginatum*); blue gramma
 5 (*Bouteloua gracilis*); buffalo grass (*Buchloe dactyloids*); sideoats gramma (*Bouteloua curtipendula*).

Plants of interest include grain plants that provide seeds of interest, oil-seed plants, and leguminous plants. Seeds of interest include grain seeds, such as corn, wheat, barley, rice, sorghum, rye, millet, *etc.* Oil-seed plants include cotton, soybean,
 10 safflower, sunflower, *Brassica*, maize, alfalfa, palm, coconut, flax, castor, olive *etc.* Leguminous plants include beans and peas. Beans include guar, locust bean, fenugreek, soybean, garden beans, cowpea, mungbean, lima bean, fava bean, lentils, chickpea, *etc.*

Evaluation of Plant Transformation

15 Following introduction of heterologous foreign DNA into plant cells, the transformation or integration of heterologous gene in the plant genome is confirmed by various methods such as analysis of nucleic acids, proteins and metabolites associated with the integrated gene.

PCR analysis is a rapid method to screen transformed cells, tissue or shoots for
 20 the presence of incorporated gene at the earlier stage before transplanting into the soil (Sambrook and Russell, (2001) *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY). PCR is carried out using oligonucleotide primers specific to the gene of interest or *Agrobacterium* vector background, *etc.*

25 Plant transformation may be confirmed by Southern blot analysis of genomic DNA (Sambrook and Russell, (2001) *supra*). In general, total DNA is extracted from the transformant, digested with appropriate restriction enzymes, fractionated in an agarose gel and transferred to a nitrocellulose or nylon membrane. The membrane or “blot” is then probed with, for example, radiolabeled ³²P target DNA fragment to confirm the
 30 integration of introduced gene into the plant genome according to standard techniques (Sambrook and Russell, (2001) *supra*).

In Northern blot analysis, RNA is isolated from specific tissues of transformant, fractionated in a formaldehyde agarose gel, and blotted onto a nylon filter according to standard procedures that are routinely used in the art (Sambrook and Russell, (2001)
 35 *supra*). Expression of RNA encoded by the pesticidal gene is then tested by hybridizing

the filter to a radioactive probe derived from a pesticidal gene, by methods known in the art (Sambrook and Russell, (2001) *supra*).

Western blot, biochemical assays and the like may be carried out on the transgenic plants to confirm the presence of protein encoded by the pesticidal gene by
5 standard procedures (Sambrook and Russell, 2001, *supra*) using antibodies that bind to one or more epitopes present on the PHI-4 polypeptide.

Stacking of traits in transgenic plant

Transgenic plants may comprise a stack of one or more insecticidal
10 polynucleotides disclosed herein with one or more additional polynucleotides resulting in the production or suppression of multiple polypeptide sequences. Transgenic plants comprising stacks of polynucleotide sequences can be obtained by either or both of traditional breeding methods or through genetic engineering methods. These methods include, but are not limited to, breeding individual lines each comprising a polynucleotide
15 of interest, transforming a transgenic plant comprising a gene disclosed herein with a subsequent gene, and co- transformation of genes into a single plant cell. As used herein, the term “stacked” includes having the multiple traits present in the same plant (i.e., both traits are incorporated into the nuclear genome, one trait is incorporated into the nuclear genome and one trait is incorporated into the genome of a plastid or both traits
20 are incorporated into the genome of a plastid). In one non-limiting example, “stacked traits” comprise a molecular stack where the sequences are physically adjacent to each other. A trait, as used herein, refers to the phenotype derived from a particular sequence or groups of sequences. Co-transformation of genes can be carried out using single transformation vectors comprising multiple genes or genes carried separately on multiple
25 vectors. If the sequences are stacked by genetically transforming the plants, the polynucleotide sequences of interest can be combined at any time and in any order. The traits can be introduced simultaneously in a co-transformation protocol with the polynucleotides of interest provided by any combination of transformation cassettes. For example, if two sequences will be introduced, the two sequences can be contained in
30 separate transformation cassettes (trans) or contained on the same transformation cassette (cis). Expression of the sequences can be driven by the same promoter or by different promoters. In certain cases, it may be desirable to introduce a transformation cassette that will suppress the expression of the polynucleotide of interest. This may be combined with any combination of other suppression cassettes or overexpression
35 cassettes to generate the desired combination of traits in the plant. It is further recognized that polynucleotide sequences can be stacked at a desired genomic location

using a site-specific recombination system. See, for example, WO 1999/25821, WO 1999/25854, WO 1999/25840, WO 1999/25855 and WO 1999/25853, all of which are herein incorporated by reference.

In some embodiments the polynucleotides encoding the PHI-4 polypeptides disclosed herein, alone or stacked with one or more additional insect resistance traits can be stacked with one or more additional input traits (e.g., herbicide resistance, fungal resistance, virus resistance or stress tolerance, disease resistance, male sterility, stalk strength, and the like) or output traits (e.g., increased yield, modified starches, improved oil profile, balanced amino acids, high lysine or methionine, increased digestibility, improved fiber quality, drought resistance, and the like). Thus, the polynucleotide embodiments can be used to provide a complete agronomic package of improved crop quality with the ability to flexibly and cost effectively control any number of agronomic pests.

Transgenes useful for stacking include but are not limited to:

1. Transgenes that Confer Resistance to Insects or Disease and that Encode:

(A) Plant disease resistance genes. Plant defenses are often activated by specific interaction between the product of a disease resistance gene (R) in the plant and the product of a corresponding avirulence (Avr) gene in the pathogen. A plant variety can be transformed with cloned resistance gene to engineer plants that are resistant to specific pathogen strains. See, for example, Jones, *et al.*, (1994) *Science* 266:789 (cloning of the tomato Cf-9 gene for resistance to *Cladosporium fulvum*); Martin, *et al.*, (1993) *Science* 262:1432 (tomato Pto gene for resistance to *Pseudomonas syringae* pv. tomato encodes a protein kinase); Mindrinos, *et al.*, (1994) *Cell* 78:1089 (*Arabidopsis* RSP2 gene for resistance to *Pseudomonas syringae*), McDowell and Woffenden, (2003) *Trends Biotechnol.* 21(4):178-83 and Toyoda, *et al.*, (2002) *Transgenic Res.* 11(6):567-82. A plant resistant to a disease is one that is more resistant to a pathogen as compared to the wild type plant.

(B) Genes encoding a *Bacillus thuringiensis* protein, a derivative thereof or a synthetic polypeptide modeled thereon. See, for example, Geiser, *et al.*, (1986) *Gene* 48:109, who disclose the cloning and nucleotide sequence of a Bt delta-endotoxin gene. Moreover, DNA molecules encoding delta-endotoxin genes can be purchased from American Type Culture Collection (Rockville, Md.), for example, under ATCC Accession Numbers 40098, 67136, 31995 and 31998. Other non-limiting examples of *Bacillus thuringiensis* transgenes being genetically engineered are given in the following patents and patent applications and hereby are incorporated by reference for this purpose: US

Patent Numbers 5,188,960; 5,689,052; 5,880,275; 5,986,177; 6,023,013, 6,060,594, 6,063,597, 6,077,824, 6,620,988, 6,642,030, 6,713,259, 6,893,826, 7,105,332; 7,179,965, 7,208,474; 7,227,056, 7,288,643, 7,323,556, 7,329,736, 7,449,552, 7,468,278, 7,510,878, 7,521,235, 7,544,862, 7,605,304, 7,696,412, 7,629,504, 7,705,216, 7,772,465, 7,790,846, 5 7,858,849 and WO 1991/14778; WO 1999/31248; WO 2001/12731; WO 1999/24581 and WO 1997/40162.

Genes encoding pesticidal proteins may also be stacked including but are not limited to: insecticidal proteins from *Pseudomonas* sp. such as PSEEN3174 (Monalysin, (2011) *PLoS Pathogens*, 7:1-13), from *Pseudomonas protegens* strain CHA0 and Pf-5 10 (previously *fluorescens*) (Pechy-Tarr, (2008) *Environmental Microbiology* 10:2368-2386: GenBank Accession No. EU400157); from *Pseudomonas Taiwanensis* (Liu, *et al.*, (2010) *J. Agric. Food Chem.* 58:12343-12349) and from *Pseudomonas pseudoalcaligenes* (Zhang, *et al.*, (2009) *Annals of Microbiology* 59:45-50 and Li, *et al.*, (2007) *Plant Cell Tiss. Organ Cult.* 89:159-168); insecticidal proteins from *Photorhabdus* sp. and 15 *Xenorhabdus* sp. (Hinchliffe, *et al.*, (2010) *The Open Toxinology Journal* 3:101-118 and Morgan, *et al.*, (2001) *Applied and Envir. Micro.* 67:2062-2069), US Patent Number 6,048,838, and US Patent Number 6,379,946; a PIP-1 polypeptide of US Serial Number 13792861; an AfIP-1A and/or AfIP-1B polypeptide of US Serial Number 13/800233; a PHI-4 polypeptide of US Serial Number 13/839702; and δ -endotoxins including, but not 20 limited to, the Cry1, Cry2, Cry3, Cry4, Cry5, Cry6, Cry7, Cry8, Cry9, Cry10, Cry11, Cry12, Cry13, Cry14, Cry15, Cry16, Cry17, Cry18, Cry19, Cry20, Cry21, Cry22, Cry23, Cry24, Cry25, Cry26, Cry27, Cry 28, Cry 29, Cry 30, Cry31, Cry32, Cry33, Cry34, Cry35, Cry36, Cry37, Cry38, Cry39, Cry40, Cry41, Cry42, Cry43, Cry44, Cry45, Cry 46, Cry47, Cry49, Cry 51, Cry 52, Cry 53, Cry54, Cry55, Cry56, Cry57, Cry58, Cry59, Cry60, Cry61, Cry62, 25 Cry63, Cry64, Cry65, Cry66, Cry67, Cry68, Cry69, Cry70, Cry71, Cry72, And Cry73 classes of δ -endotoxin genes and the *B. thuringiensis* cytolytic Cyt1 and Cyt2 genes. Members of these classes of *B. thuringiensis* insecticidal proteins include, but are not limited to Cry1Aa1 (Accession # AAA22353); Cry1Aa2 (Accession # Accession # AAA22552); Cry1Aa3 (Accession # BAA00257); Cry1Aa4 (Accession # CAA31886); 30 Cry1Aa5 (Accession # BAA04468); Cry1Aa6 (Accession # AAA86265); Cry1Aa7 (Accession # AAD46139); Cry1Aa8 (Accession # I26149); Cry1Aa9 (Accession # BAA77213); Cry1Aa10 (Accession # AAD55382); Cry1Aa11 (Accession # CAA70856); Cry1Aa12 (Accession # AAP80146); Cry1Aa13 (Accession # AAM44305); Cry1Aa14 (Accession # AAP40639); Cry1Aa15 (Accession # AAY66993); Cry1Aa16 (Accession # HQ439776); Cry1Aa17 (Accession # HQ439788); Cry1Aa18 (Accession # HQ439790); 35 Cry1Aa19 (Accession # HQ685121); Cry1Aa20 (Accession # JF340156); Cry1Aa21

(Accession # JN651496); Cry1Aa22 (Accession # KC158223); Cry1Ab1 (Accession # AAA22330); Cry1Ab2 (Accession # AAA22613); Cry1Ab3 (Accession # AAA22561); Cry1Ab4 (Accession # BAA00071); Cry1Ab5 (Accession # CAA28405); Cry1Ab6 (Accession # AAA22420); Cry1Ab7 (Accession # CAA31620); Cry1Ab8 (Accession # AAA22551); Cry1Ab9 (Accession # CAA38701); Cry1Ab10 (Accession # A29125); Cry1Ab11 (Accession # I12419); Cry1Ab12 (Accession # AAC64003); Cry1Ab13 (Accession # AAN76494); Cry1Ab14 (Accession # AAG16877); Cry1Ab15 (Accession # AAO13302); Cry1Ab16 (Accession # AAK55546); Cry1Ab17 (Accession # AAT46415); Cry1Ab18 (Accession # AAQ88259); Cry1Ab19 (Accession # AAW31761); Cry1Ab20 (Accession # ABB72460); Cry1Ab21 (Accession # ABS18384); Cry1Ab22 (Accession # ABW87320); Cry1Ab23 (Accession # HQ439777); Cry1Ab24 (Accession # HQ439778); Cry1Ab25 (Accession # HQ685122); Cry1Ab26 (Accession # HQ847729); Cry1Ab27 (Accession # JN135249); Cry1Ab28 (Accession # JN135250); Cry1Ab29 (Accession # JN135251); Cry1Ab30 (Accession # JN135252); Cry1Ab31 (Accession # JN135253); Cry1Ab32 (Accession # JN135254); Cry1Ab33 (Accession # AAS93798); Cry1Ab34 (Accession # KC156668); Cry1Ab-like (Accession # AAK14336); Cry1Ab-like (Accession # AAK14337); Cry1Ab-like (Accession # AAK14338); Cry1Ab-like (Accession # ABG88858); Cry1Ac1 (Accession # AAA22331); Cry1Ac2 (Accession # AAA22338); Cry1Ac3 (Accession # CAA38098); Cry1Ac4 (Accession # AAA73077); Cry1Ac5 (Accession # AAA22339); Cry1Ac6 (Accession # AAA86266); Cry1Ac7 (Accession # AAB46989); Cry1Ac8 (Accession # AAC44841); Cry1Ac9 (Accession # AAB49768); Cry1Ac10 (Accession # CAA05505); Cry1Ac11 (Accession # CAA10270); Cry1Ac12 (Accession # I12418); Cry1Ac13 (Accession # AAD38701); Cry1Ac14 (Accession # AAQ06607); Cry1Ac15 (Accession # AAN07788); Cry1Ac16 (Accession # AAU87037); Cry1Ac17 (Accession # AAX18704); Cry1Ac18 (Accession # AAY88347); Cry1Ac19 (Accession # ABD37053); Cry1Ac20 (Accession # ABB89046); Cry1Ac21 (Accession # AAY66992); Cry1Ac22 (Accession # ABZ01836); Cry1Ac23 (Accession # CAQ30431); Cry1Ac24 (Accession # ABL01535); Cry1Ac25 (Accession # FJ513324); Cry1Ac26 (Accession # FJ617446); Cry1Ac27 (Accession # FJ617447); Cry1Ac28 (Accession # ACM90319); Cry1Ac29 (Accession # DQ438941); Cry1Ac30 (Accession # GQ227507); Cry1Ac31 (Accession # GU446674); Cry1Ac32 (Accession # HM061081); Cry1Ac33 (Accession # GQ866913); Cry1Ac34 (Accession # HQ230364); Cry1Ac35 (Accession # JF340157); Cry1Ac36 (Accession # JN387137); Cry1Ac37 (Accession # JQ317685); Cry1Ad1 (Accession # AAA22340); Cry1Ad2 (Accession # CAA01880); Cry1Ae1 (Accession # AAA22410); Cry1Af1 (Accession # AAB82749); Cry1Ag1 (Accession # AAD46137); Cry1Ah1 (Accession # AAQ14326); Cry1Ah2 (Accession # ABB76664);

Cry1Ah3 (Accession # HQ439779); Cry1Ai1 (Accession # AAO39719); Cry1Ai2
 (Accession # HQ439780); Cry1A-like (Accession # AAK14339); Cry1Ba1 (Accession #
 CAA29898); Cry1Ba2 (Accession # CAA65003); Cry1Ba3 (Accession # AAK63251);
 Cry1Ba4 (Accession # AAK51084); Cry1Ba5 (Accession # ABO20894); Cry1Ba6
 5 (Accession # ABL60921); Cry1Ba7 (Accession # HQ439781); Cry1Bb1 (Accession #
 AAA22344); Cry1Bb2 (Accession # HQ439782); Cry1Bc1 (Accession # CAA86568);
 Cry1Bd1 (Accession # AAD10292); Cry1Bd2 (Accession # AAM93496); Cry1Be1
 (Accession # AAC32850); Cry1Be2 (Accession # AAQ52387); Cry1Be3 (Accession #
 ACV96720); Cry1Be4 (Accession # HM070026); Cry1Bf1 (Accession # CAC50778);
 10 Cry1Bf2 (Accession # AAQ52380); Cry1Bg1 (Accession # AAO39720); Cry1Bh1
 (Accession # HQ589331); Cry1Bi1 (Accession # KC156700); Cry1Ca1 (Accession #
 CAA30396); Cry1Ca2 (Accession # CAA31951); Cry1Ca3 (Accession # AAA22343);
 Cry1Ca4 (Accession # CAA01886); Cry1Ca5 (Accession # CAA65457); Cry1Ca6 [1]
 (Accession # AAF37224); Cry1Ca7 (Accession # AAG50438); Cry1Ca8 (Accession #
 15 AAM00264); Cry1Ca9 (Accession # AAL79362); Cry1Ca10 (Accession # AAN16462);
 Cry1Ca11 (Accession # AAX53094); Cry1Ca12 (Accession # HM070027); Cry1Ca13
 (Accession # HQ412621); Cry1Ca14 (Accession # JN651493); Cry1Cb1 (Accession #
 M97880); Cry1Cb2 (Accession # AAG35409); Cry1Cb3 (Accession # ACD50894);
 Cry1Cb-like (Accession # AAX63901); Cry1Da1 (Accession # CAA38099); Cry1Da2
 20 (Accession # I76415); Cry1Da3 (Accession # HQ439784); Cry1Db1 (Accession #
 CAA80234); Cry1Db2 (Accession # AAK48937); Cry1Dc1 (Accession # ABK35074);
 Cry1Ea1 (Accession # CAA37933); Cry1Ea2 (Accession # CAA39609); Cry1Ea3
 (Accession # AAA22345); Cry1Ea4 (Accession # AAD04732); Cry1Ea5 (Accession #
 A15535); Cry1Ea6 (Accession # AAL50330); Cry1Ea7 (Accession # AAW72936);
 25 Cry1Ea8 (Accession # ABX11258); Cry1Ea9 (Accession # HQ439785); Cry1Ea10
 (Accession # ADR00398); Cry1Ea11 (Accession # JQ652456); Cry1Eb1 (Accession #
 AAA22346); Cry1Fa1 (Accession # AAA22348); Cry1Fa2 (Accession # AAA22347);
 Cry1Fa3 (Accession # HM070028); Cry1Fa4 (Accession # HM439638); Cry1Fb1
 (Accession # CAA80235); Cry1Fb2 (Accession # BAA25298); Cry1Fb3 (Accession #
 30 AAF21767); Cry1Fb4 (Accession # AAC10641); Cry1Fb5 (Accession # AAO13295);
 Cry1Fb6 (Accession # ACD50892); Cry1Fb7 (Accession # ACD50893); Cry1Ga1
 (Accession # CAA80233); Cry1Ga2 (Accession # CAA70506); Cry1Gb1 (Accession #
 AAD10291); Cry1Gb2 (Accession # AAO13756); Cry1Gc1 (Accession # AAQ52381);
 Cry1Ha1 (Accession # CAA80236); Cry1Hb1 (Accession # AAA79694); Cry1Hb2
 35 (Accession # HQ439786); Cry1H-like (Accession # AAF01213); Cry1Ia1 (Accession #
 CAA44633); Cry1Ia2 (Accession # AAA22354); Cry1Ia3 (Accession # AAC36999);

Cry1Ia4 (Accession # AAB00958); Cry1Ia5 (Accession # CAA70124); Cry1Ia6 (Accession # AAC26910); Cry1Ia7 (Accession # AAM73516); Cry1Ia8 (Accession # AAK66742); Cry1Ia9 (Accession # AAQ08616); Cry1Ia10 (Accession # AAP86782); Cry1Ia11 (Accession # CAC85964); Cry1Ia12 (Accession # AAV53390); Cry1Ia13 (Accession # ABF83202); Cry1Ia14 (Accession # ACG63871); Cry1Ia15 (Accession # FJ617445); Cry1Ia16 (Accession # FJ617448); Cry1Ia17 (Accession # GU989199); Cry1Ia18 (Accession # ADK23801); Cry1Ia19 (Accession # HQ439787); Cry1Ia20 (Accession # JQ228426); Cry1Ia21 (Accession # JQ228424); Cry1Ia22 (Accession # JQ228427); Cry1Ia23 (Accession # JQ228428); Cry1Ia24 (Accession # JQ228429); Cry1Ia25 (Accession # JQ228430); Cry1Ia26 (Accession # JQ228431); Cry1Ia27 (Accession # JQ228432); Cry1Ia28 (Accession # JQ228433); Cry1Ia29 (Accession # JQ228434); Cry1Ia30 (Accession # JQ317686); Cry1Ia31 (Accession # JX944038); Cry1Ia32 (Accession # JX944039); Cry1Ia33 (Accession # JX944040); Cry1Ib1 (Accession # AAA82114); Cry1Ib2 (Accession # ABW88019); Cry1Ib3 (Accession # ACD75515); Cry1Ib4 (Accession # HM051227); Cry1Ib5 (Accession # HM070028); Cry1Ib6 (Accession # ADK38579); Cry1Ib7 (Accession # JN571740); Cry1Ib8 (Accession # JN675714); Cry1Ib9 (Accession # JN675715); Cry1Ib10 (Accession # JN675716); Cry1Ib11 (Accession # JQ228423); Cry1Ic1 (Accession # AAC62933); Cry1Ic2 (Accession # AAE71691); Cry1Id1 (Accession # AAD44366); Cry1Id2 (Accession # JQ228422); Cry1Ie1 (Accession # AAG43526); Cry1Ie2 (Accession # HM439636); Cry1Ie3 (Accession # KC156647); Cry1Ie4 (Accession # KC156681); Cry1If1 (Accession # AAQ52382); Cry1Ig1 (Accession # KC156701); Cry1I-like (Accession # AAC31094); Cry1I-like (Accession # ABG88859); Cry1Ja1 (Accession # AAA22341); Cry1Ja2 (Accession # HM070030); Cry1Ja3 (Accession # JQ228425); Cry1Jb1 (Accession # AAA98959); Cry1Jc1 (Accession # AAC31092); Cry1Jc2 (Accession # AAQ52372); Cry1Jd1 (Accession # CAC50779); Cry1Ka1 (Accession # AAB00376); Cry1Ka2 (Accession # HQ439783); Cry1La1 (Accession # AAS60191); Cry1La2 (Accession # HM070031); Cry1Ma1 (Accession # FJ884067); Cry1Ma2 (Accession # KC156659); Cry1Na1 (Accession # KC156648); Cry1Nb1 (Accession # KC156678); Cry1-like (Accession # AAC31091); Cry2Aa1 (Accession # AAA22335); Cry2Aa2 (Accession # AAA83516); Cry2Aa3 (Accession # D86064); Cry2Aa4 (Accession # AAC04867); Cry2Aa5 (Accession # CAA10671); Cry2Aa6 (Accession # CAA10672); Cry2Aa7 (Accession # CAA10670); Cry2Aa8 (Accession # AAO13734); Cry2Aa9 (Accession # AAO13750); Cry2Aa10 (Accession # AAQ04263); Cry2Aa11 (Accession # AAQ52384); Cry2Aa12 (Accession # ABI83671); Cry2Aa13 (Accession # ABL01536); Cry2Aa14 (Accession # ACF04939); Cry2Aa15 (Accession # JN426947); Cry2Ab1 (Accession # AAA22342); Cry2Ab2

(Accession # CAA39075); Cry2Ab3 (Accession # AAG36762); Cry2Ab4 (Accession # AAO13296); Cry2Ab5 (Accession # AAQ04609); Cry2Ab6 (Accession # AAP59457); Cry2Ab7 (Accession # AAZ66347); Cry2Ab8 (Accession # ABC95996); Cry2Ab9 (Accession # ABC74968); Cry2Ab10 (Accession # EF157306); Cry2Ab11 (Accession # CAM84575); Cry2Ab12 (Accession # ABM21764); Cry2Ab13 (Accession # ACG76120); Cry2Ab14 (Accession # ACG76121); Cry2Ab15 (Accession # HM037126); Cry2Ab16 (Accession # GQ866914); Cry2Ab17 (Accession # HQ439789); Cry2Ab18 (Accession # JN135255); Cry2Ab19 (Accession # JN135256); Cry2Ab20 (Accession # JN135257); Cry2Ab21 (Accession # JN135258); Cry2Ab22 (Accession # JN135259); Cry2Ab23 (Accession # JN135260); Cry2Ab24 (Accession # JN135261); Cry2Ab25 (Accession # JN415485); Cry2Ab26 (Accession # JN426946); Cry2Ab27 (Accession # JN415764); Cry2Ab28 (Accession # JN651494); Cry2Ac1 (Accession # CAA40536); Cry2Ac2 (Accession # AAG35410); Cry2Ac3 (Accession # AAQ52385); Cry2Ac4 (Accession # ABC95997); Cry2Ac5 (Accession # ABC74969); Cry2Ac6 (Accession # ABC74793); Cry2Ac7 (Accession # CAL18690); Cry2Ac8 (Accession # CAM09325); Cry2Ac9 (Accession # CAM09326); Cry2Ac10 (Accession # ABN15104); Cry2Ac11 (Accession # CAM83895); Cry2Ac12 (Accession # CAM83896); Cry2Ad1 (Accession # AAF09583); Cry2Ad2 (Accession # ABC86927); Cry2Ad3 (Accession # CAK29504); Cry2Ad4 (Accession # CAM32331); Cry2Ad5 (Accession # CAO78739); Cry2Ae1 (Accession # AAQ52362); Cry2Af1 (Accession # ABO30519); Cry2Af2 (Accession # GQ866915); Cry2Ag1 (Accession # ACH91610); Cry2Ah1 (Accession # EU939453); Cry2Ah2 (Accession # ACL80665); Cry2Ah3 (Accession # GU073380); Cry2Ah4 (Accession # KC156702); Cry2Ai1 (Accession # FJ788388); Cry2Aj (Accession #); Cry2Ak1 (Accession # KC156660); Cry2Ba1 (Accession # KC156658); Cry3Aa1 (Accession # AAA22336); Cry3Aa2 (Accession # AAA22541); Cry3Aa3 (Accession # CAA68482); Cry3Aa4 (Accession # AAA22542); Cry3Aa5 (Accession # AAA50255); Cry3Aa6 (Accession # AAC43266); Cry3Aa7 (Accession # CAB41411); Cry3Aa8 (Accession # AAS79487); Cry3Aa9 (Accession # AAW05659); Cry3Aa10 (Accession # AAU29411); Cry3Aa11 (Accession # AAW82872); Cry3Aa12 (Accession # ABY49136); Cry3Ba1 (Accession # CAA34983); Cry3Ba2 (Accession # CAA00645); Cry3Ba3 (Accession # JQ397327); Cry3Bb1 (Accession # AAA22334); Cry3Bb2 (Accession # AAA74198); Cry3Bb3 (Accession # I15475); Cry3Ca1 (Accession # CAA42469); Cry4Aa1 (Accession # CAA68485); Cry4Aa2 (Accession # BAA00179); Cry4Aa3 (Accession # CAD30148); Cry4Aa4 (Accession # AFB18317); Cry4A-like (Accession # AAY96321); Cry4Ba1 (Accession # CAA30312); Cry4Ba2 (Accession # CAA30114); Cry4Ba3 (Accession # AAA22337); Cry4Ba4 (Accession # BAA00178); Cry4Ba5 (Accession # CAD30095);

Cry4Ba-like (Accession # ABC47686); Cry4Ca1 (Accession # EU646202); Cry4Cb1
 (Accession # FJ403208); Cry4Cb2 (Accession # FJ597622); Cry4Cc1 (Accession #
 FJ403207); Cry5Aa1 (Accession # AAA67694); Cry5Ab1 (Accession # AAA67693);
 Cry5Ac1 (Accession # I34543); Cry5Ad1 (Accession # ABQ82087); Cry5Ba1 (Accession
 5 # AAA68598); Cry5Ba2 (Accession # ABW88931); Cry5Ba3 (Accession # AFJ04417);
 Cry5Ca1 (Accession # HM461869); Cry5Ca2 (Accession # ZP_04123426); Cry5Da1
 (Accession # HM461870); Cry5Da2 (Accession # ZP_04123980); Cry5Ea1 (Accession #
 HM485580); Cry5Ea2 (Accession # ZP_04124038); Cry6Aa1 (Accession # AAA22357);
 Cry6Aa2 (Accession # AAM46849); Cry6Aa3 (Accession # ABH03377); Cry6Ba1
 10 (Accession # AAA22358); Cry7Aa1 (Accession # AAA22351); Cry7Ab1 (Accession #
 AAA21120); Cry7Ab2 (Accession # AAA21121); Cry7Ab3 (Accession # ABX24522);
 Cry7Ab4 (Accession # EU380678); Cry7Ab5 (Accession # ABX79555); Cry7Ab6
 (Accession # ACI44005); Cry7Ab7 (Accession # ADB89216); Cry7Ab8 (Accession #
 GU145299); Cry7Ab9 (Accession # ADD92572); Cry7Ba1 (Accession # ABB70817);
 15 Cry7Bb1 (Accession # KC156653); Cry7Ca1 (Accession # ABR67863); Cry7Cb1
 (Accession # KC156698); Cry7Da1 (Accession # ACQ99547); Cry7Da2 (Accession #
 HM572236); Cry7Da3 (Accession # KC156679); Cry7Ea1 (Accession # HM035086);
 Cry7Ea2 (Accession # HM132124); Cry7Ea3 (Accession # EEM19403); Cry7Fa1
 (Accession # HM035088); Cry7Fa2 (Accession # EEM19090); Cry7Fb1 (Accession #
 20 HM572235); Cry7Fb2 (Accession # KC156682); Cry7Ga1 (Accession # HM572237);
 Cry7Ga2 (Accession # KC156669); Cry7Gb1 (Accession # KC156650); Cry7Gc1
 (Accession # KC156654); Cry7Gd1 (Accession # KC156697); Cry7Ha1 (Accession #
 KC156651); Cry7Ia1 (Accession # KC156665); Cry7Ja1 (Accession # KC156671);
 Cry7Ka1 (Accession # KC156680); Cry7Kb1 (Accession # BAM99306); Cry7La1
 25 (Accession # BAM99307); Cry8Aa1 (Accession # AAA21117); Cry8Ab1 (Accession #
 EU044830); Cry8Ac1 (Accession # KC156662); Cry8Ad1 (Accession # KC156684);
 Cry8Ba1 (Accession # AAA21118); Cry8Bb1 (Accession # CAD57542); Cry8Bc1
 (Accession # CAD57543); Cry8Ca1 (Accession # AAA21119); Cry8Ca2 (Accession #
 AAR98783); Cry8Ca3 (Accession # EU625349); Cry8Ca4 (Accession # ADB54826);
 30 Cry8Da1 (Accession # BAC07226); Cry8Da2 (Accession # BD133574); Cry8Da3
 (Accession # BD133575); Cry8Db1 (Accession # BAF93483); Cry8Ea1 (Accession #
 AAQ73470); Cry8Ea2 (Accession # EU047597); Cry8Ea3 (Accession # KC855216);
 Cry8Fa1 (Accession # AAT48690); Cry8Fa2 (Accession # HQ174208); Cry8Fa3
 (Accession # AFH78109); Cry8Ga1 (Accession # AAT46073); Cry8Ga2 (Accession #
 35 ABC42043); Cry8Ga3 (Accession # FJ198072); Cry8Ha1 (Accession # AAW81032);
 Cry8Ia1 (Accession # EU381044); Cry8Ia2 (Accession # GU073381); Cry8Ia3 (Accession

HM044664); Cry8Ia4 (Accession # KC156674); Cry8Ib1 (Accession # GU325772); Cry8Ib2 (Accession # KC156677); Cry8Ja1 (Accession # EU625348); Cry8Ka1 (Accession # FJ422558); Cry8Ka2 (Accession # ACN87262); Cry8Kb1 (Accession # HM123758); Cry8Kb2 (Accession # KC156675); Cry8La1 (Accession # GU325771);

5 Cry8Ma1 (Accession # HM044665); Cry8Ma2 (Accession # EEM86551); Cry8Ma3 (Accession # HM210574); Cry8Na1 (Accession # HM640939); Cry8Pa1 (Accession # HQ388415); Cry8Qa1 (Accession # HQ441166); Cry8Qa2 (Accession # KC152468); Cry8Ra1 (Accession # AFP87548); Cry8Sa1 (Accession # JQ740599); Cry8Ta1 (Accession # KC156673); Cry8-like (Accession # FJ770571); Cry8-like (Accession #

10 ABS53003); Cry9Aa1 (Accession # CAA41122); Cry9Aa2 (Accession # CAA41425); Cry9Aa3 (Accession # GQ249293); Cry9Aa4 (Accession # GQ249294); Cry9Aa5 (Accession # JX174110); Cry9Aa like (Accession # AAQ52376); Cry9Ba1 (Accession # CAA52927); Cry9Ba2 (Accession # GU299522); Cry9Bb1 (Accession # AAV28716); Cry9Ca1 (Accession # CAA85764); Cry9Ca2 (Accession # AAQ52375); Cry9Da1

15 (Accession # BAA19948); Cry9Da2 (Accession # AAB97923); Cry9Da3 (Accession # GQ249293); Cry9Da4 (Accession # GQ249297); Cry9Db1 (Accession # AAX78439); Cry9Dc1 (Accession # KC156683); Cry9Ea1 (Accession # BAA34908); Cry9Ea2 (Accession # AAO12908); Cry9Ea3 (Accession # ABM21765); Cry9Ea4 (Accession # ACE88267); Cry9Ea5 (Accession # ACF04743); Cry9Ea6 (Accession # ACG63872);

20 Cry9Ea7 (Accession # FJ380927); Cry9Ea8 (Accession # GQ249292); Cry9Ea9 (Accession # JN651495); Cry9Eb1 (Accession # CAC50780); Cry9Eb2 (Accession # GQ249298); Cry9Eb3 (Accession # KC156646); Cry9Ec1 (Accession # AAC63366); Cry9Ed1 (Accession # AAX78440); Cry9Ee1 (Accession # GQ249296); Cry9Ee2 (Accession # KC156664); Cry9Fa1 (Accession # KC156692); Cry9Ga1 (Accession #

25 KC156699); Cry9-like (Accession # AAC63366); Cry10Aa1 (Accession # AAA22614); Cry10Aa2 (Accession # E00614); Cry10Aa3 (Accession # CAD30098); Cry10Aa4 (Accession # AFB18318); Cry10A-like (Accession # DQ167578); Cry11Aa1 (Accession # AAA22352); Cry11Aa2 (Accession # AAA22611); Cry11Aa3 (Accession # CAD30081); Cry11Aa4 (Accession # AFB18319); Cry11Aa-like (Accession # DQ166531); Cry11Ba1

30 (Accession # CAA60504); Cry11Bb1 (Accession # AAC97162); Cry11Bb2 (Accession # HM068615); Cry12Aa1 (Accession # AAA22355); Cry13Aa1 (Accession # AAA22356); Cry14Aa1 (Accession # AAA21516); Cry14Ab1 (Accession # KC156652); Cry15Aa1 (Accession # AAA22333); Cry16Aa1 (Accession # CAA63860); Cry17Aa1 (Accession # CAA67841); Cry18Aa1 (Accession # CAA67506); Cry18Ba1 (Accession # AAF89667);

35 Cry18Ca1 (Accession # AAF89668); Cry19Aa1 (Accession # CAA68875); Cry19Ba1 (Accession # BAA32397); Cry19Ca1 (Accession # AFM37572); Cry20Aa1 (Accession #

AAB93476); Cry20Ba1 (Accession # ACS93601); Cry20Ba2 (Accession # KC156694);
 Cry20-like (Accession # GQ144333); Cry21Aa1 (Accession # I32932); Cry21Aa2
 (Accession # I66477); Cry21Ba1 (Accession # BAC06484); Cry21Ca1 (Accession #
 JF521577); Cry21Ca2 (Accession # KC156687); Cry21Da1 (Accession # JF521578);
 5 Cry22Aa1 (Accession # I34547); Cry22Aa2 (Accession # CAD43579); Cry22Aa3
 (Accession # ACD93211); Cry22Ab1 (Accession # AAK50456); Cry22Ab2 (Accession #
 CAD43577); Cry22Ba1 (Accession # CAD43578); Cry22Bb1 (Accession # KC156672);
 Cry23Aa1 (Accession # AAF76375); Cry24Aa1 (Accession # AAC61891); Cry24Ba1
 (Accession # BAD32657); Cry24Ca1 (Accession # CAJ43600); Cry25Aa1 (Accession #
 10 AAC61892); Cry26Aa1 (Accession # AAD25075); Cry27Aa1 (Accession # BAA82796);
 Cry28Aa1 (Accession # AAD24189); Cry28Aa2 (Accession # AAG00235); Cry29Aa1
 (Accession # CAC80985); Cry30Aa1 (Accession # CAC80986); Cry30Ba1 (Accession #
 BAD00052); Cry30Ca1 (Accession # BAD67157); Cry30Ca2 (Accession # ACU24781);
 Cry30Da1 (Accession # EF095955); Cry30Db1 (Accession # BAE80088); Cry30Ea1
 15 (Accession # ACC95445); Cry30Ea2 (Accession # FJ499389); Cry30Fa1 (Accession #
 ACI22625); Cry30Ga1 (Accession # ACG60020); Cry30Ga2 (Accession # HQ638217);
 Cry31Aa1 (Accession # BAB11757); Cry31Aa2 (Accession # AAL87458); Cry31Aa3
 (Accession # BAE79808); Cry31Aa4 (Accession # BAF32571); Cry31Aa5 (Accession #
 BAF32572); Cry31Aa6 (Accession # BAI44026); Cry31Ab1 (Accession # BAE79809);
 20 Cry31Ab2 (Accession # BAF32570); Cry31Ac1 (Accession # BAF34368); Cry31Ac2
 (Accession # AB731600); Cry31Ad1 (Accession # BAI44022); Cry32Aa1 (Accession #
 AAG36711); Cry32Aa2 (Accession # GU063849); Cry32Ab1 (Accession # GU063850);
 Cry32Ba1 (Accession # BAB78601); Cry32Ca1 (Accession # BAB78602); Cry32Cb1
 (Accession # KC156708); Cry32Da1 (Accession # BAB78603); Cry32Ea1 (Accession #
 25 GU324274); Cry32Ea2 (Accession # KC156686); Cry32Eb1 (Accession # KC156663);
 Cry32Fa1 (Accession # KC156656); Cry32Ga1 (Accession # KC156657); Cry32Ha1
 (Accession # KC156661); Cry32Hb1 (Accession # KC156666); Cry32Ia1 (Accession #
 KC156667); Cry32Ja1 (Accession # KC156685); Cry32Ka1 (Accession # KC156688);
 Cry32La1 (Accession # KC156689); Cry32Ma1 (Accession # KC156690); Cry32Mb1
 30 (Accession # KC156704); Cry32Na1 (Accession # KC156691); Cry32Oa1 (Accession #
 KC156703); Cry32Pa1 (Accession # KC156705); Cry32Qa1 (Accession # KC156706);
 Cry32Ra1 (Accession # KC156707); Cry32Sa1 (Accession # KC156709); Cry32Ta1
 (Accession # KC156710); Cry32Ua1 (Accession # KC156655); Cry33Aa1 (Accession #
 AAL26871); Cry34Aa1 (Accession # AAG50341); Cry34Aa2 (Accession # AAK64560);
 35 Cry34Aa3 (Accession # AAT29032); Cry34Aa4 (Accession # AAT29030); Cry34Ab1
 (Accession # AAG41671); Cry34Ac1 (Accession # AAG50118); Cry34Ac2 (Accession #

AAK64562); Cry34Ac3 (Accession # AAT29029); Cry34Ba1 (Accession # AAK64565); Cry34Ba2 (Accession # AAT29033); Cry34Ba3 (Accession # AAT29031); Cry35Aa1 (Accession # AAG50342); Cry35Aa2 (Accession # AAK64561); Cry35Aa3 (Accession # AAT29028); Cry35Aa4 (Accession # AAT29025); Cry35Ab1 (Accession # AAG41672);

5 Cry35Ab2 (Accession # AAK64563); Cry35Ab3 (Accession # AY536891); Cry35Ac1 (Accession # AAG50117); Cry35Ba1 (Accession # AAK64566); Cry35Ba2 (Accession # AAT29027); Cry35Ba3 (Accession # AAT29026); Cry36Aa1 (Accession # AAK64558); Cry37Aa1 (Accession # AAF76376); Cry38Aa1 (Accession # AAK64559); Cry39Aa1 (Accession # BAB72016); Cry40Aa1 (Accession # BAB72018); Cry40Ba1 (Accession #

10 BAC77648); Cry40Ca1 (Accession # EU381045); Cry40Da1 (Accession # ACF15199); Cry41Aa1 (Accession # BAD35157); Cry41Ab1 (Accession # BAD35163); Cry41Ba1 (Accession # HM461871); Cry41Ba2 (Accession # ZP_04099652); Cry42Aa1 (Accession # BAD35166); Cry43Aa1 (Accession # BAD15301); Cry43Aa2 (Accession # BAD95474); Cry43Ba1 (Accession # BAD15303); Cry43Ca1 (Accession # KC156676); Cry43Cb1

15 (Accession # KC156695); Cry43Cc1 (Accession # KC156696); Cry43-like (Accession # BAD15305); Cry44Aa (Accession # BAD08532); Cry45Aa (Accession # BAD22577); Cry46Aa (Accession # BAC79010); Cry46Aa2 (Accession # BAG68906); Cry46Ab (Accession # BAD35170); Cry47Aa (Accession # AAY24695); Cry48Aa (Accession # CAJ18351); Cry48Aa2 (Accession # CAJ86545); Cry48Aa3 (Accession # CAJ86546);

20 Cry48Ab (Accession # CAJ86548); Cry48Ab2 (Accession # CAJ86549); Cry49Aa (Accession # CAH56541); Cry49Aa2 (Accession # CAJ86541); Cry49Aa3 (Accession # CAJ86543); Cry49Aa4 (Accession # CAJ86544); Cry49Ab1 (Accession # CAJ86542); Cry50Aa1 (Accession # BAE86999); Cry50Ba1 (Accession # GU446675); Cry50Ba2 (Accession # GU446676); Cry51Aa1 (Accession # ABI14444); Cry51Aa2 (Accession #

25 GU570697); Cry52Aa1 (Accession # EF613489); Cry52Ba1 (Accession # FJ361760); Cry53Aa1 (Accession # EF633476); Cry53Ab1 (Accession # FJ361759); Cry54Aa1 (Accession # ACA52194); Cry54Aa2 (Accession # GQ140349); Cry54Ba1 (Accession # GU446677); Cry55Aa1 (Accession # ABW88932); Cry54Ab1 (Accession # JQ916908); Cry55Aa2 (Accession # AAE33526); Cry56Aa1 (Accession # ACU57499); Cry56Aa2

30 (Accession # GQ483512); Cry56Aa3 (Accession # JX025567); Cry57Aa1 (Accession # ANC87261); Cry58Aa1 (Accession # ANC87260); Cry59Ba1 (Accession # JN790647); Cry59Aa1 (Accession # ACR43758); Cry60Aa1 (Accession # ACU24782); Cry60Aa2 (Accession # EAO57254); Cry60Aa3 (Accession # EEM99278); Cry60Ba1 (Accession # GU810818); Cry60Ba2 (Accession # EAO57253); Cry60Ba3 (Accession # EEM99279);

35 Cry61Aa1 (Accession # HM035087); Cry61Aa2 (Accession # HM132125); Cry61Aa3 (Accession # EEM19308); Cry62Aa1 (Accession # HM054509); Cry63Aa1 (Accession #

BAI44028); Cry64Aa1 (Accession # BAJ05397); Cry65Aa1 (Accession # HM461868); Cry65Aa2 (Accession # ZP_04123838); Cry66Aa1 (Accession # HM485581); Cry66Aa2 (Accession # ZP_04099945); Cry67Aa1 (Accession # HM485582); Cry67Aa2 (Accession # ZP_04148882); Cry68Aa1 (Accession # HQ113114); Cry69Aa1 (Accession # HQ401006); Cry69Aa2 (Accession # JQ821388); Cry69Ab1 (Accession # JN209957); Cry70Aa1 (Accession # JN646781); Cry70Ba1 (Accession # ADO51070); Cry70Bb1 (Accession # EEL67276); Cry71Aa1 (Accession # JX025568); Cry72Aa1 (Accession # JX025569); and Cry73Aa (Accession # AEH76822).

Examples of δ -endotoxins also include but are not limited to Cry1A proteins of US Patent Numbers 5,880,275 and 7,858,849; a DIG-3 or DIG-11 toxin (N-terminal deletion of α -helix 1 and/or α -helix 2 variants of Cry proteins such as Cry1A) of US Patent Numbers 8,304,604 and 8,304,605, Cry1B of US Patent Application Serial Number 10/525,318; Cry1C of US Patent Number 6,033,874; Cry1F of US Patent Numbers 5,188,960, 6,218,188; Cry1A/F chimeras of US Patent Numbers 7,070,982; 6,962,705 and 6,713,063); a Cry2 protein such as Cry2Ab protein of US Patent Number 7,064,249); a Cry3A protein including but not limited to an engineered hybrid insecticidal protein (eHIP) created by fusing unique combinations of variable regions and conserved blocks of at least two different Cry proteins (US Patent Application Publication Number 2010/0017914); a Cry4 protein; a Cry5 protein; a Cry6 protein; Cry8 proteins of US Patent Numbers 7,329,736, 7,449,552, 7,803,943, 7,476,781, 7,105,332, 7,378,499 and 7,462,760; a Cry9 protein such as such as members of the Cry9A, Cry9B, Cry9C, Cry9D, Cry9E, and Cry9F families; a Cry15 protein of Naimov, *et al.*, (2008) *Applied and Environmental Microbiology* 74:7145–7151; a Cry22, a Cry34Ab1 protein of US Patent Numbers 6,127,180, 6,624,145 and 6,340,593; a CryET33 and CryET34 protein of US Patent Numbers 6,248,535, 6,326,351, 6,399,330, 6,949,626, 7,385,107 and 7,504,229; a CryET33 and CryET34 homologs of US Patent Publication Number 2006/0191034, 2012/0278954, and PCT Publication Number WO 2012/139004; a Cry35Ab1 protein of US Patent Numbers 6,083,499, 6,548,291 and 6,340,593; a Cry46 protein, a Cry 51 protein, a Cry binary toxin; a TIC901 or related toxin; TIC807 of US 2008/0295207; ET29, ET37, TIC809, TIC810, TIC812, TIC127, TIC128 of PCT US 2006/033867; AXMI-027, AXMI-036, and AXMI-038 of US Patent Number 8,236,757; AXMI-031, AXMI-039, AXMI-040, AXMI-049 of US 7,923,602; AXMI-018, AXMI-020, and AXMI-021 of WO 2006/083891; AXMI-010 of WO 2005/038032; AXMI-003 of WO 2005/021585; AXMI-008 of US 2004/0250311; AXMI-006 of US 2004/0216186; AXMI-007 of US 2004/0210965; AXMI-009 of US 2004/0210964; AXMI-014 of US 2004/0197917; AXMI-004 of US 2004/0197916; AXMI-028 and AXMI-029 of WO 2006/119457; AXMI-007, AXMI-008,

AXMI-0080rf2, AXMI-009, AXMI-014 and AXMI-004 of WO 2004/074462; AXMI-150 of US Patent Number 8,084,416; AXMI-205 of US20110023184; AXMI-011, AXMI-012, AXMI-013, AXMI-015, AXMI-019, AXMI-044, AXMI-037, AXMI-043, AXMI-033, AXMI-034, AXMI-022, AXMI-023, AXMI-041, AXMI-063, and AXMI-064 of US 2011/0263488; AXMI-5 R1 and related proteins of US 2010/0197592; AXMI221Z, AXMI222z, AXMI223z, AXMI224z and AXMI225z of WO 2011/103248; AXMI218, AXMI219, AXMI220, AXMI226, AXMI227, AXMI228, AXMI229, AXMI230, and AXMI231 of WO11/103247; AXMI-115, AXMI-113, AXMI-005, AXMI-163 and AXMI-184 of US Patent Number 8,334,431; AXMI-001, AXMI-002, AXMI-030, AXMI-035, and AXMI-045 of US 2010/0298211; AXMI-066 10 and AXMI-076 of US20090144852; AXMI128, AXMI130, AXMI131, AXMI133, AXMI140, AXMI141, AXMI142, AXMI143, AXMI144, AXMI146, AXMI148, AXMI149, AXMI152, AXMI153, AXMI154, AXMI155, AXMI156, AXMI157, AXMI158, AXMI162, AXMI165, AXMI166, AXMI167, AXMI168, AXMI169, AXMI170, AXMI171, AXMI172, AXMI173, AXMI174, AXMI175, AXMI176, AXMI177, AXMI178, AXMI179, AXMI180, AXMI181, 15 AXMI182, AXMI185, AXMI186, AXMI187, AXMI188, AXMI189 of US Patent Number 8,318,900; AXMI079, AXMI080, AXMI081, AXMI082, AXMI091, AXMI092, AXMI096, AXMI097, AXMI098, AXMI099, AXMI100, AXMI101, AXMI102, AXMI103, AXMI104, AXMI107, AXMI108, AXMI109, AXMI110, AXMI111, AXMI112, AXMI114, AXMI116, AXMI117, AXMI118, AXMI119, AXMI120, AXMI121, AXMI122, AXMI123, AXMI124, 20 AXMI1257, AXMI1268, AXMI127, AXMI129, AXMI164, AXMI151, AXMI161, AXMI183, AXMI132, AXMI138, AXMI137 of US 2010/0005543; Axmi115 variants of US2013097728Cry proteins such as Cry1A and Cry3A having modified proteolytic sites of US Patent Number 8,319,019; and a Cry1Ac, Cry2Aa and Cry1Ca toxin protein from *Bacillus thuringiensis* strain VBTS 2528 of US Patent Application Publication Number 25 2011/0064710. Other Cry proteins are well known to one skilled in the art (see, Crickmore, *et al.*, "Bacillus thuringiensis toxin nomenclature" (2011), at lifesci.sussex.ac.uk/home/Neil_Crickmore/Bt/ which can be accessed on the world-wide web using the "www" prefix). The insecticidal activity of Cry proteins is well known to one skilled in the art (for review, see, van Frannkenhuyzen, (2009) *J. Invert. Path.* 101:1-16).

30 The use of Cry proteins as transgenic plant traits is well known to one skilled in the art and Cry-transgenic plants including but not limited to Cry1Ac, Cry1Ac+Cry2Ab, Cry1Ab, Cry1A.105, Cry1F, Cry1Fa2, Cry1F+Cry1Ac, Cry2Ab, Cry3A, mCry3A, Cry3Bb1, Cry34Ab1, Cry35Ab1, Vip3A, mCry3A, Cry9c and CBI-Bt have received regulatory approval (see, Sanahuja, (2011) *Plant Biotech Journal* 9:283-300 and the CERA (2010) 35 GM Crop Database Center for Environmental Risk Assessment (CERA), ILSI Research Foundation, Washington D.C. at cera-gmc.org/index.php?action=gm_crop_database

which can be accessed on the world-wide web using the "www" prefix). More than one pesticidal proteins well known to one skilled in the art can also be expressed in plants such as Vip3Ab & Cry1Fa (US2012/0317682), Cry1BE & Cry1F (US2012/0311746), Cry1CA & Cry1AB (US2012/0311745), Cry1F & CryCa (US2012/0317681), Cry1DA & Cry1BE (US2012/0331590), Cry1DA & Cry1Fa (US2012/0331589), Cry1AB & Cry1BE (US2012/0324606), and Cry1Fa & Cry2Aa, Cry1I or Cry1E (US2012/0324605). Pesticidal proteins also include insecticidal lipases including lipid acyl hydrolases of US Patent Number 7,491,869, and cholesterol oxidases such as from *Streptomyces* (Purcell et al. (1993) *Biochem Biophys Res Commun* 15:1406-1413). Pesticidal proteins also include VIP (vegetative insecticidal proteins) toxins of US Patent Numbers 5,877,012, 6,107,279, 6,137,033, 7,244,820, 7,615,686, and 8,237,020, and the like. Other VIP proteins are well known to one skilled in the art (see, lifesci.sussex.ac.uk/home/Neil_Crickmore/Bt/vip.html which can be accessed on the world-wide web using the "www" prefix). Pesticidal proteins also include toxin complex (TC) proteins, obtainable from organisms such as *Xenorhabdus*, *Photorhabdus* and *Paenibacillus* (see, US Patent Numbers 7,491,698 and 8,084,418). Some TC proteins have "stand alone" insecticidal activity and other TC proteins enhance the activity of the stand-alone toxins produced by the same given organism. The toxicity of a "stand-alone" TC protein (from *Photorhabdus*, *Xenorhabdus* or *Paenibacillus*, for example) can be enhanced by one or more TC protein "potentiators" derived from a source organism of a different genus. There are three main types of TC proteins. As referred to herein, Class A proteins ("Protein A") are stand-alone toxins. Class B proteins ("Protein B") and Class C proteins ("Protein C") enhance the toxicity of Class A proteins. Examples of Class A proteins are TcbA, TcdA, XptA1 and XptA2. Examples of Class B proteins are TcaC, TcdB, XptB1Xb and XptC1Wi. Examples of Class C proteins are TccC, XptC1Xb and XptB1Wi. Pesticidal proteins also include spider, snake and scorpion venom proteins. Examples of spider venom peptides include but are not limited to lycotoxin-1 peptides and mutants thereof (US Patent Number 8,334,366).

(C) A polynucleotide encoding an insect-specific hormone or pheromone such as an ecdysteroid and juvenile hormone, a variant thereof, a mimetic based thereon or an antagonist or agonist thereof. See, for example, the disclosure by Hammock, *et al.*, (1990) *Nature* 344:458, of baculovirus expression of cloned juvenile hormone esterase, an inactivator of juvenile hormone.

(D) A polynucleotide encoding an insect-specific peptide which, upon expression, disrupts the physiology of the affected pest. For example, see the disclosures of, Regan, (1994) *J. Biol. Chem.* 269:9 (expression cloning yields DNA coding for insect diuretic

hormone receptor); Pratt, *et al.*, (1989) *Biochem. Biophys. Res. Comm.* 163:1243 (an allostatin is identified in *Diploptera puntata*); Chattopadhyay, *et al.*, (2004) *Critical Reviews in Microbiology* 30(1):33-54; Zjawiony, (2004) *J Nat Prod* 67(2):300-310; Carlini and Grossi-de-Sa, (2002) *Toxicon* 40(11):1515-1539; Ussuf, *et al.*, (2001) *Curr Sci.* 80(7):847-853 and Vasconcelos and Oliveira, (2004) *Toxicon* 44(4):385-403. See also, US Patent Number 5,266,317 to Tomalski, *et al.*, who disclose genes encoding insect-specific toxins.

(E) A polynucleotide encoding an enzyme responsible for a hyperaccumulation of a monoterpene, a sesquiterpene, a steroid, hydroxamic acid, a phenylpropanoid derivative or another non-protein molecule with insecticidal activity.

(F) A polynucleotide encoding an enzyme involved in the modification, including the post-translational modification, of a biologically active molecule; for example, a glycolytic enzyme, a proteolytic enzyme, a lipolytic enzyme, a nuclease, a cyclase, a transaminase, an esterase, a hydrolase, a phosphatase, a kinase, a phosphorylase, a polymerase, an elastase, a chitinase and a glucanase, whether natural or synthetic. See, PCT Application WO 1993/02197 in the name of Scott, *et al.*, which discloses the nucleotide sequence of a callase gene. DNA molecules which contain chitinase-encoding sequences can be obtained, for example, from the ATCC under Accession Numbers 39637 and 67152. See also, Kramer, *et al.*, (1993) *Insect Biochem. Molec. Biol.* 23:691, who teach the nucleotide sequence of a cDNA encoding tobacco hookworm chitinase and Kawalleck, *et al.*, (1993) *Plant Molec. Biol.* 21:673, who provide the nucleotide sequence of the parsley ubi4-2 polyubiquitin gene, and US Patent Numbers 6,563,020; 7,145,060 and 7,087,810.

(G) A polynucleotide encoding a molecule that stimulates signal transduction. For example, see the disclosure by Botella, *et al.*, (1994) *Plant Molec. Biol.* 24:757, of nucleotide sequences for mung bean calmodulin cDNA clones, and Griess, *et al.*, (1994) *Plant Physiol.* 104:1467, who provide the nucleotide sequence of a maize calmodulin cDNA clone.

(H) A polynucleotide encoding a hydrophobic moment peptide. See, PCT Application WO 1995/16776 and US Patent Number 5,580,852 disclosure of peptide derivatives of Tachyplesin which inhibit fungal plant pathogens) and PCT Application WO 1995/18855 and US Patent Number 5,607,914 (teaches synthetic antimicrobial peptides that confer disease resistance).

(I) A polynucleotide encoding a membrane permease, a channel former or a channel blocker. For example, see the disclosure by Jaynes, *et al.*, (1993) *Plant Sci.*

89:43, of heterologous expression of a cecropin-beta lytic peptide analog to render transgenic tobacco plants resistant to *Pseudomonas solanacearum*.

(J) A gene encoding a viral-invasive protein or a complex toxin derived therefrom. For example, the accumulation of viral coat proteins in transformed plant cells imparts
 5 resistance to viral infection and/or disease development effected by the virus from which the coat protein gene is derived, as well as by related viruses. See, Beachy, *et al.*, (1990) *Ann. Rev. Phytopathol.* 28:451. Coat protein-mediated resistance has been conferred upon transformed plants against alfalfa mosaic virus, cucumber mosaic virus, tobacco streak virus, potato virus X, potato virus Y, tobacco etch virus, tobacco rattle virus and
 10 tobacco mosaic virus. *Id.*

(K) A gene encoding an insect-specific antibody or an immunotoxin derived therefrom. Thus, an antibody targeted to a critical metabolic function in the insect gut would inactivate an affected enzyme, killing the insect. Cf. Taylor, *et al.*, Abstract #497, SEVENTH INT'L SYMPOSIUM ON MOLECULAR PLANT-MICROBE INTERACTIONS
 15 (Edinburgh, Scotland, 1994) (enzymatic inactivation in transgenic tobacco via production of single-chain antibody fragments).

(L) A gene encoding a virus-specific antibody. See, for example, Tavladoraki, *et al.*, (1993) *Nature* 366:469, who show that transgenic plants expressing recombinant antibody genes are protected from virus attack.

(M) A polynucleotide encoding a developmental-arrestive protein produced in
 20 nature by a pathogen or a parasite. Thus, fungal endo alpha-1,4-D-polygalacturonases facilitate fungal colonization and plant nutrient release by solubilizing plant cell wall homo-alpha-1,4-D-galacturonase. See, Lamb, *et al.*, (1992) *Bio/Technology* 10:1436. The cloning and characterization of a gene which encodes a bean endopolygalacturonase-
 25 inhibiting protein is described by Toubart, *et al.*, (1992) *Plant J.* 2:367.

(N) A polynucleotide encoding a developmental-arrestive protein produced in nature by a plant. For example, Logemann, *et al.*, (1992) *Bio/Technology* 10:305, have shown that transgenic plants expressing the barley ribosome-inactivating gene have an increased resistance to fungal disease.

(O) Genes involved in the Systemic Acquired Resistance (SAR) Response and/or
 30 the pathogenesis related genes. Briggs, (1995) *Current Biology* 5(2), Pieterse and Van Loon, (2004) *Curr. Opin. Plant Bio.* 7(4):456-64 and Somssich, (2003) *Cell* 113(7):815-6.

(P) Antifungal genes (Cornelissen and Melchers, (1993) *Pl. Physiol.* 101:709-712 and Parijs, *et al.*, (1991) *Planta* 183:258-264 and Bushnell, *et al.*, (1998) *Can. J. of Plant Path.* 20(2):137-149. Also see, US Patent Application Serial Numbers 09/950,933;
 35 11/619,645; 11/657,710; 11/748,994; 11/774,121 and US Patent Numbers 6,891,085 and

7,306,946. LysM Receptor-like kinases for the perception of chitin fragments as a first step in plant defense response against fungal pathogens (US 2012/0110696).

(Q) Detoxification genes, such as for fumonisin, beauvericin, moniliformin and zearalenone and their structurally related derivatives. For example, see, US Patent
5 Numbers 5,716,820; 5,792,931; 5,798,255; 5,846,812; 6,083,736; 6,538,177; 6,388,171 and 6,812,380.

(R) A polynucleotide encoding a Cystatin and cysteine proteinase inhibitors. See, US Patent Number 7,205,453.

(S) Defensin genes. See, WO 2003/000863 and US Patent Numbers 6,911,577;
10 6,855,865; 6,777,592 and 7,238,781.

(T) Genes conferring resistance to nematodes. See, e.g., PCT Application WO 1996/30517; PCT Application WO 1993/19181, WO 2003/033651 and Urwin, *et al.*, (1998) *Planta* 204:472-479, Williamson, (1999) *Curr Opin Plant Bio.* 2(4):327-31; US Patent Numbers 6,284,948 and 7,301,069 and miR164 genes (WO 2012/058266).

(U) Genes that confer resistance to Phytophthora Root Rot, such as the Rps 1, Rps 1-a, Rps 1-b, Rps 1-c, Rps 1-d, Rps 1-e, Rps 1-k, Rps 2, Rps 3-a, Rps 3-b, Rps 3-c, Rps 4, Rps 5, Rps 6, Rps 7 and other Rps genes. See, for example, Shoemaker, *et al.*, Phytophthora Root Rot Resistance Gene Mapping in Soybean, Plant Genome IV Conference, San Diego, Calif. (1995).
15

(V) Genes that confer resistance to Brown Stem Rot, such as described in US Patent Number 5,689,035 and incorporated by reference for this purpose.
20

(W) Genes that confer resistance to Colletotrichum, such as described in US Patent Application Publication US 2009/0035765 and incorporated by reference for this purpose. This includes the Rcg locus that may be utilized as a single locus conversion.
25

2. Transgenes that Confer Resistance to a Herbicide, for Example:

(A) A polynucleotide encoding resistance to a herbicide that inhibits the growing point or meristem, such as an imidazolinone or a sulfonylurea. Exemplary genes in this category code for mutant ALS and AHAS enzyme as described, for example, by Lee, *et al.*, (1988) *EMBO J.* 7:1241 and Miki, *et al.*, (1990) *Theor. Appl. Genet.* 80:449, respectively. See also, US Patent Numbers 5,605,011; 5,013,659; 5,141,870; 5,767,361; 5,731,180; 5,304,732; 4,761,373; 5,331,107; 5,928,937 and 5,378,824; US Patent Application Serial Number 11/683,737 and International Publication WO 1996/33270.
30

(B) A polynucleotide encoding a protein for resistance to Glyphosate (resistance imparted by mutant 5-enolpyruvyl-3-phosphikimate synthase (EPSP) and aroA genes, respectively) and other phosphono compounds such as glufosinate (phosphinothricin
35

acetyl transferase (PAT) and *Streptomyces hygroscopicus* phosphinothricin acetyl transferase (bar) genes), and pyridinoxy or phenoxy proprionic acids and cyclohexones (ACCase inhibitor-encoding genes). See, for example, US Patent Number 4,940,835 to Shah, *et al.*, which discloses the nucleotide sequence of a form of EPSPS which can confer glyphosate resistance. US Patent Number 5,627,061 to Barry, *et al.*, also describes genes encoding EPSPS enzymes. See also, US Patent Numbers 6,566,587; 6,338,961; 6,248,876 B1; 6,040,497; 5,804,425; 5,633,435; 5,145,783; 4,971,908; 5,312,910; 5,188,642; 5,094,945, 4,940,835; 5,866,775; 6,225,114 B1; 6,130,366; 5,310,667; 4,535,060; 4,769,061; 5,633,448; 5,510,471; Re. 36,449; RE 37,287 E and 5,491,288 and International Publications EP 1173580; WO 2001/66704; EP 1173581 and EP 1173582, which are incorporated herein by reference for this purpose. Glyphosate resistance is also imparted to plants that express a gene encoding a glyphosate oxidoreductase enzyme as described more fully in US Patent Numbers 5,776,760 and 5,463,175, which are incorporated herein by reference for this purpose. In addition glyphosate resistance can be imparted to plants by the over expression of genes encoding glyphosate N-acetyltransferase. See, for example, US Patent Numbers 7,462,481; 7,405,074 and US Patent Application Publication Number US 2008/0234130. A DNA molecule encoding a mutant *aroA* gene can be obtained under ATCC Accession Number 39256, and the nucleotide sequence of the mutant gene is disclosed in US Patent Number 4,769,061 to Comai. EP Application Number 0 333 033 to Kumada, *et al.*, and US Patent Number 4,975,374 to Goodman, *et al.*, disclose nucleotide sequences of glutamine synthetase genes which confer resistance to herbicides such as L-phosphinothricin. The nucleotide sequence of a phosphinothricin-acetyl-transferase gene is provided in EP Application Numbers 0 242 246 and 0 242 236 to Leemans, *et al.*; De Greef, *et al.*, (1989) *Bio/Technology* 7:61, describe the production of transgenic plants that express chimeric bar genes coding for phosphinothricin acetyl transferase activity. See also, US Patent Numbers 5,969,213; 5,489,520; 5,550,318; 5,874,265; 5,919,675; 5,561,236; 5,648,477; 5,646,024; 6,177,616 B1 and 5,879,903, which are incorporated herein by reference for this purpose. Exemplary genes conferring resistance to phenoxy proprionic acids and cyclohexones, such as sethoxydim and haloxyfop, are the Acc1-S1, Acc1-S2 and Acc1-S3 genes described by Marshall, *et al.*, (1992) *Theor. Appl. Genet.* 83:435.

(C) A polynucleotide encoding a protein for resistance to herbicide that inhibits photosynthesis, such as a triazine (*psbA* and *gs+* genes) and a benzonitrile (nitrilase gene). Przibilla, *et al.*, (1991) *Plant Cell* 3:169, describe the transformation of *Chlamydomonas* with plasmids encoding mutant *psbA* genes. Nucleotide sequences for

nitrilase genes are disclosed in US Patent Number 4,810,648 to Stalker and DNA molecules containing these genes are available under ATCC Accession Numbers 53435, 67441 and 67442. Cloning and expression of DNA coding for a glutathione S-transferase is described by Hayes, *et al.*, (1992) *Biochem. J.* 285:173.

5 (D) A polynucleotide encoding a protein for resistance to Acetohydroxy acid synthase, which has been found to make plants that express this enzyme resistant to multiple types of herbicides, has been introduced into a variety of plants (see, e.g., Hattori, *et al.*, (1995) *Mol Gen Genet.* 246:419). Other genes that confer resistance to herbicides include: a gene encoding a chimeric protein of rat cytochrome P4507A1 and
10 yeast NADPH-cytochrome P450 oxidoreductase (Shiota, *et al.*, (1994) *Plant Physiol* 106:17), genes for glutathione reductase and superoxide dismutase (Aono, *et al.*, (1995) *Plant Cell Physiol* 36:1687) and genes for various phosphotransferases (Datta, *et al.*, (1992) *Plant Mol Biol* 20:619).

(E) A polynucleotide encoding resistance to a herbicide targeting
15 Protoporphyrinogen oxidase (protox) which is necessary for the production of chlorophyll. The protox enzyme serves as the target for a variety of herbicidal compounds. These herbicides also inhibit growth of all the different species of plants present, causing their total destruction. The development of plants containing altered protox activity which are resistant to these herbicides are described in US Patent Numbers 6,288,306 B1;
20 6,282,837 B1 and 5,767,373 and International Publication WO 2001/12825.

(F) The aad-1 gene (originally from *Sphingobium herbicidovorans*) encodes the aryloxyalkanoate dioxygenase (AAD-1) protein. The trait confers tolerance to 2,4-dichlorophenoxyacetic acid and aryloxyphenoxypropionate (commonly referred to as "fop" herbicides such as quizalofop) herbicides. The aad-1 gene, itself, for herbicide tolerance
25 in plants was first disclosed in WO 2005/107437 (see also, US 2009/0093366). The aad-12 gene, derived from *Delftia acidovorans*, which encodes the aryloxyalkanoate dioxygenase (AAD-12) protein that confers tolerance to 2,4-dichlorophenoxyacetic acid and pyridyloxyacetate herbicides by deactivating several herbicides with an aryloxyalkanoate moiety, including phenoxy auxin (e.g., 2,4-D, MCPA), as well as
30 pyridyloxy auxins (e.g., fluroxypyr, triclopyr).

(G) A polynucleotide encoding a herbicide resistant dicamba monooxygenase disclosed in US Patent Application Publication 2003/0135879 for imparting dicamba tolerance;

(H) A polynucleotide molecule encoding bromoxynil nitrilase (Bxn) disclosed in US
35 Patent Number 4,810,648 for imparting bromoxynil tolerance;

(l) A polynucleotide molecule encoding phytoene (crtl) described in Misawa, *et al.*, (1993) *Plant J.* 4:833-840 and in Misawa, *et al.*, (1994) *Plant J.* 6:481-489 for norflurazon tolerance.

5 3. Transgenes that Confer or Contribute to an Altered Grain Characteristic

Such as:

(A) Altered fatty acids, for example, by

(1) Down-regulation of stearyl-ACP to increase stearic acid content of the plant. See, Knultzon, *et al.*, (1992) *Proc. Natl. Acad. Sci. USA* 89:2624 and WO 1999/64579
10 (Genes to Alter Lipid Profiles in Corn).

(2) Elevating oleic acid via FAD-2 gene modification and/or decreasing linolenic acid via FAD-3 gene modification (see, US Patent Numbers 6,063,947; 6,323,392; 6,372,965 and WO 1993/11245).

(3) Altering conjugated linolenic or linoleic acid content, such as in WO
15 2001/12800.

(4) Altering LEC1, AGP, Dek1, Superal1, mi1 ps, various lpa genes such as lpa1, lpa3, hpt or hgg. For example, see, WO 2002/42424, WO 1998/22604, WO 2003/011015, WO 2002/057439, WO 2003/011015, US Patent Numbers 6,423,886, 6,197,561, 6,825,397 and US Patent Application Publication Numbers US 2003/0079247,
20 US 2003/0204870 and Rivera-Madrid, *et al.*, (1995) *Proc. Natl. Acad. Sci.* 92:5620-5624.

(5) Genes encoding delta-8 desaturase for making long-chain polyunsaturated fatty acids (US Patent Numbers 8,058,571 and 8,338,152), delta-9 desaturase for lowering saturated fats (US Patent Number 8,063,269), Primula Δ 6-desaturase for improving omega-3 fatty acid profiles.

25 (6) Isolated nucleic acids and proteins associated with lipid and sugar metabolism regulation, in particular, lipid metabolism protein (LMP) used in methods of producing transgenic plants and modulating levels of seed storage compounds including lipids, fatty acids, starches or seed storage proteins and use in methods of modulating the seed size, seed number, seed weights, root length and leaf size of plants (EP 2404499).

30 (7) Altering expression of a High-Level Expression of Sugar-Inducible 2 (HSI2) protein in the plant to increase or decrease expression of HSI2 in the plant. Increasing expression of HSI2 increases oil content while decreasing expression of HSI2 decreases abscisic acid sensitivity and/or increases drought resistance (US Patent Application Publication Number 2012/0066794).

35 (8) Expression of cytochrome b5 (Cb5) alone or with FAD2 to modulate oil content in plant seed, particular to increase the levels of omega-3 fatty acids and improve the ratio

of omega-6 to omega-3 fatty acids (US Patent Application Publication Number 2011/0191904).

(9) Nucleic acid molecules encoding wrinkled1-like polypeptides for modulating sugar metabolism (US Patent Number 8,217,223).

5 (B) Altered phosphorus content, for example, by the

(1) Introduction of a phytase-encoding gene would enhance breakdown of phytate, adding more free phosphate to the transformed plant. For example, see, Van Hartingsveldt, *et al.*, (1993) *Gene* 127:87, for a disclosure of the nucleotide sequence of an *Aspergillus niger* phytase gene.

10 (2) Modulating a gene that reduces phytate content. In maize, this, for example, could be accomplished, by cloning and then re-introducing DNA associated with one or more of the alleles, such as the LPA alleles, identified in maize mutants characterized by low levels of phytic acid, such as in WO 2005/113778 and/or by altering inositol kinase activity as in WO 2002/059324, US Patent Application Publication Number 2003/0009011,
15 WO 2003/027243, US Patent Application Publication Number 2003/0079247, WO 1999/05298, US Patent Number 6,197,561, US Patent Number 6,291,224, US Patent Number 6,391,348, WO 2002/059324, US Patent Application Publication Number 2003/0079247, WO 1998/45448, WO 1999/55882, WO 2001/04147.

(C) Altered carbohydrates affected, for example, by altering a gene for an enzyme
20 that affects the branching pattern of starch or, a gene altering thioredoxin such as NTR and/or TRX (see, US Patent Number 6,531,648, which is incorporated by reference for this purpose) and/or a gamma zein knock out or mutant such as cs27 or TUSC27 or en27 (see, US Patent Number 6,858,778 and US Patent Application Publication Number 2005/0160488, US Patent Application Publication Number 2005/0204418, which are
25 incorporated by reference for this purpose). See, Shiroza, *et al.*, (1988) *J. Bacteriol.* 170:810 (nucleotide sequence of *Streptococcus* mutant fructosyltransferase gene), Steinmetz, *et al.*, (1985) *Mol. Gen. Genet.* 200:220 (nucleotide sequence of *Bacillus subtilis* levansucrase gene), Pen, *et al.*, (1992) *Bio/Technology* 10:292 (production of transgenic plants that express *Bacillus licheniformis* alpha-amylase), Elliot, *et al.*, (1993)
30 *Plant Molec. Biol.* 21:515 (nucleotide sequences of tomato invertase genes), Sogaard, *et al.*, (1993) *J. Biol. Chem.* 268:22480 (site-directed mutagenesis of barley alpha-amylase gene) and Fisher, *et al.*, (1993) *Plant Physiol.* 102:1045 (maize endosperm starch branching enzyme II), WO 1999/10498 (improved digestibility and/or starch extraction through modification of UDP-D-xylose 4-epimerase, Fragile 1 and 2, Ref1, HCHL, C4H),
35 US Patent Number 6,232,529 (method of producing high oil seed by modification of starch levels (AGP)). The fatty acid modification genes mentioned herein may also be used to

affect starch content and/or composition through the interrelationship of the starch and oil pathways.

(D) Altered antioxidant content or composition, such as alteration of tocopherol or tocotrienols. For example, see, US Patent Number 6,787,683, US Patent Application Publication Number 2004/0034886 and WO 2000/68393 involving the manipulation of antioxidant levels and WO 2003/082899 through alteration of a homogentisate geranyl geranyl transferase (hggt).

(E) Altered essential seed amino acids. For example, see, US Patent Number 6,127,600 (method of increasing accumulation of essential amino acids in seeds), US Patent Number 6,080,913 (binary methods of increasing accumulation of essential amino acids in seeds), US Patent Number 5,990,389 (high lysine), WO 1999/40209 (alteration of amino acid compositions in seeds), WO 1999/29882 (methods for altering amino acid content of proteins), US Patent Number 5,850,016 (alteration of amino acid compositions in seeds), WO 1998/20133 (proteins with enhanced levels of essential amino acids), US Patent Number 5,885,802 (high methionine), US Patent Number 5,885,801 (high threonine), US Patent Number 6,664,445 (plant amino acid biosynthetic enzymes), US Patent Number 6,459,019 (increased lysine and threonine), US Patent Number 6,441,274 (plant tryptophan synthase beta subunit), US Patent Number 6,346,403 (methionine metabolic enzymes), US Patent Number 5,939,599 (high sulfur), US Patent Number 5,912,414 (increased methionine), WO 1998/56935 (plant amino acid biosynthetic enzymes), WO 1998/45458 (engineered seed protein having higher percentage of essential amino acids), WO 1998/42831 (increased lysine), US Patent Number 5,633,436 (increasing sulfur amino acid content), US Patent Number 5,559,223 (synthetic storage proteins with defined structure containing programmable levels of essential amino acids for improvement of the nutritional value of plants), WO 1996/01905 (increased threonine), WO 1995/15392 (increased lysine), US Patent Application Publication Number 2003/0163838, US Patent Application Publication Number 2003/0150014, US Patent Application Publication Number 2004/0068767, US Patent Number 6,803,498, WO 2001/79516.

4. Genes that Control Male-Sterility:

There are several methods of conferring genetic male sterility available, such as multiple mutant genes at separate locations within the genome that confer male sterility, as disclosed in US Patent Numbers 4,654,465 and 4,727,219 to Brar, *et al.*, and chromosomal translocations as described by Patterson in US Patent Numbers 3,861,709 and 3,710,511. In addition to these methods, Albertsen, *et al.*, US Patent Number

5,432,068, describe a system of nuclear male sterility which includes: identifying a gene which is critical to male fertility; silencing this native gene which is critical to male fertility; removing the native promoter from the essential male fertility gene and replacing it with an inducible promoter; inserting this genetically engineered gene back into the plant; and
5 thus creating a plant that is male sterile because the inducible promoter is not “on” resulting in the male fertility gene not being transcribed. Fertility is restored by inducing or turning “on”, the promoter, which in turn allows the gene that confers male fertility to be transcribed.

(A) Introduction of a deacetylase gene under the control of a tapetum-specific
10 promoter and with the application of the chemical N-Ac-PPT (WO 2001/29237).

(B) Introduction of various stamen-specific promoters (WO 1992/13956, WO 1992/13957).

(C) Introduction of the barnase and the barstar gene (Paul, *et al.*, (1992) *Plant Mol. Biol.* 19:611-622).

15 For additional examples of nuclear male and female sterility systems and genes, see also, US Patent Numbers 5,859,341; 6,297,426; 5,478,369; 5,824,524; 5,850,014 and 6,265,640, all of which are hereby incorporated by reference.

5. Genes that create a site for site specific DNA integration.

20 This includes the introduction of FRT sites that may be used in the FLP/FRT system and/or Lox sites that may be used in the Cre/Loxp system. For example, see, Lyznik, *et al.*, (2003) *Plant Cell Rep* 21:925-932 and WO 1999/25821, which are hereby incorporated by reference. Other systems that may be used include the Gin recombinase of phage Mu (Maeser, *et al.*, (1991) Vicki Chandler, The Maize Handbook ch. 118
25 (Springer-Verlag 1994), the Pin recombinase of E. coli (Enomoto, *et al.*, 1983) and the R/RS system of the pSRi plasmid (Araki, *et al.*, 1992).

6. Genes that affect abiotic stress resistance

30 Including but not limited to flowering, ear and seed development, enhancement of nitrogen utilization efficiency, altered nitrogen responsiveness, drought resistance or tolerance, cold resistance or tolerance and salt resistance or tolerance and increased yield under stress.

(A) For example, see: WO 2000/73475 where water use efficiency is altered through alteration of malate; US Patent Numbers 5,892,009, 5,965,705, 5,929,305,
35 5,891,859, 6,417,428, 6,664,446, 6,706,866, 6,717,034, 6,801,104, WO 2000/060089, WO 2001/026459, WO 2001/035725, WO 2001/034726, WO 2001/035727, WO

2001/036444, WO 2001/036597, WO 2001/036598, WO 2002/015675, WO 2002/017430, WO 2002/077185, WO 2002/079403, WO 2003/013227, WO 2003/013228, WO 2003/014327, WO 2004/031349, WO 2004/076638, WO 199809521.

5 (B) WO 199938977 describing genes, including CBF genes and transcription factors effective in mitigating the negative effects of freezing, high salinity and drought on plants, as well as conferring other positive effects on plant phenotype.

(C) US Patent Application Publication Number 2004/0148654 and WO 2001/36596 where abscisic acid is altered in plants resulting in improved plant phenotype such as increased yield and/or increased tolerance to abiotic stress.

10 (D) WO 2000/006341, WO 2004/090143, US Patent Numbers 7,531,723 and 6,992,237 where cytokinin expression is modified resulting in plants with increased stress tolerance, such as drought tolerance, and/or increased yield. Also see, WO 2002/02776, WO 2003/052063, JP 2002/281975, US Patent Number 6,084,153, WO 2001/64898, US Patent Number 6,177,275 and US Patent Number 6,107,547 (enhancement of nitrogen
15 utilization and altered nitrogen responsiveness).

(E) For ethylene alteration, see, US Patent Application Publication Number 2004/0128719, US Patent Application Publication Number 2003/0166197 and WO 2000/32761.

20 (F) For plant transcription factors or transcriptional regulators of abiotic stress, see, e.g., US Patent Application Publication Number 2004/0098764 or US Patent Application Publication Number 2004/0078852.

(G) Genes that increase expression of vacuolar pyrophosphatase such as AVP1 (US Patent Number 8,058,515) for increased yield; nucleic acid encoding a HSFA4 or a HSFA5 (Heat Shock Factor of the class A4 or A5) polypeptides, an oligopeptide
25 transporter protein (OPT4-like) polypeptide; a plastochron2-like (PLA2-like) polypeptide or a Wuschel related homeobox 1-like (WOX1-like) polypeptide (U. Patent Application Publication Number US 2011/0283420).

(H) Down regulation of polynucleotides encoding poly (ADP-ribose) polymerase (PARP) proteins to modulate programmed cell death (US Patent Number 8,058,510) for
30 increased vigor.

(I) Polynucleotide encoding DTP21 polypeptides for conferring drought resistance (US Patent Application Publication Number US 2011/0277181).

(J) Nucleotide sequences encoding ACC Synthase 3 (ACS3) proteins for modulating development, modulating response to stress, and modulating stress tolerance
35 (US Patent Application Publication Number US 2010/0287669).

(K) Polynucleotides that encode proteins that confer a drought tolerance phenotype (DTP) for conferring drought resistance (WO 2012/058528).

(L) Tocopherol cyclase (TC) genes for conferring drought and salt tolerance (US Patent Application Publication Number 2012/0272352).

5 (M) CAAX amino terminal family proteins for stress tolerance (US Patent Number 8,338,661).

(N) Mutations in the SAL1 encoding gene have increased stress tolerance, including increased drought resistant (US Patent Application Publication Number 2010/0257633).

10 (O) Expression of a nucleic acid sequence encoding a polypeptide selected from the group consisting of: GRF polypeptide, RAA1-like polypeptide, SYR polypeptide, ARKL polypeptide, and YTP polypeptide increasing yield-related traits (US Patent Application Publication Number 2011/0061133).

(P) Modulating expression in a plant of a nucleic acid encoding a Class III
15 Trehalose Phosphate Phosphatase (TPP) polypeptide for enhancing yield-related traits in plants, particularly increasing seed yield (US Patent Application Publication Number 2010/0024067).

Other genes and transcription factors that affect plant growth and agronomic traits such as yield, flowering, plant growth and/or plant structure, can be introduced or
20 introgressed into plants, see e.g., WO 1997/49811 (LHY), WO 1998/56918 (ESD4), WO 1997/10339 and US Patent Number 6,573,430 (TFL), US Patent Number 6,713,663 (FT), WO 1996/14414 (CON), WO 1996/38560, WO 2001/21822 (VRN1), WO 2000/44918 (VRN2), WO 1999/49064 (GI), WO 2000/46358 (FR1), WO 1997/29123, US Patent Number 6,794,560, US Patent Number 6,307,126 (GAI), WO 1999/09174 (D8 and Rht)
25 and WO 2004/076638 and WO 2004/031349 (transcription factors).

7. Genes that confer increased yield

(A) A transgenic crop plant transformed by a 1-AminoCyclopropane-1-Carboxylate Deaminase-like Polypeptide (ACCDP) coding nucleic acid, wherein
30 expression of the nucleic acid sequence in the crop plant results in the plant's increased root growth, and/or increased yield, and/or increased tolerance to environmental stress as compared to a wild type variety of the plant (US Patent Number 8,097,769).

(B) Over-expression of maize zinc finger protein gene (Zm-ZFP1) using a seed preferred promoter has been shown to enhance plant growth, increase kernel number and total kernel weight per plant (US Patent Application Publication Number 2012/0079623).

(C) Constitutive over-expression of maize lateral organ boundaries (LOB) domain protein (Zm-LOBDP1) has been shown to increase kernel number and total kernel weight per plant (US Patent Application Publication Number 2012/0079622).

(D) Enhancing yield-related traits in plants by modulating expression in a plant of a nucleic acid encoding a VIM1 (Variant in Methylation 1)-like polypeptide or a VTC2-like (GDP-L-galactose phosphorylase) polypeptide or a DUF1685 polypeptide or an ARF6-like (Auxin Responsive Factor) polypeptide (WO 2012/038893).

(E) Modulating expression in a plant of a nucleic acid encoding a Ste20-like polypeptide or a homologue thereof gives plants having increased yield relative to control plants (EP 2431472).

(F) Genes encoding nucleoside diphosphatase kinase (NDK) polypeptides and homologs thereof for modifying the plant's root architecture (US Patent Application Publication Number 2009/0064373).

8. Genes that confer plant digestibility.

(A) Altering the level of xylan present in the cell wall of a plant by modulating expression of xylan synthase (US 8,173,866).

In some embodiment the stacked trait may be a trait or event that has received regulatory approval including but not limited to the events in Table 2A -1F.

Table 2A *Helianthus annuus* Sunflower

Event	Company	Description
X81359	BASF Inc.	Tolerance to imidazolinone herbicides by selection of a naturally occurring mutant.

Table 2B *Medicago sativa* Alfalfa

Event	Company	Description
J101, J163	Monsanto Company and Forage Genetics International	Glyphosate herbicide tolerant alfalfa (lucerne) produced by inserting a gene encoding the enzyme 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) from the CP4 strain of <i>Agrobacterium tumefaciens</i> .

Table 2C *Oryza sativa* Rice

Event	Company	Description
CL121, CL141, CFX51	BASF Inc.	Tolerance to the imidazolinone herbicide, imazethapyr, induced by chemical mutagenesis of the acetolactate synthase (ALS) enzyme using ethyl methanesulfonate (EMS).
IMINTA-1, IMINTA-4	BASF Inc.	Tolerance to imidazolinone herbicides induced by chemical mutagenesis of the acetolactate synthase (ALS) enzyme using sodium azide.

Event	Company	Description
LLRICE06, LLRICE62	Aventis CropScience	Glufosinate ammonium herbicide tolerant rice produced by inserting a modified phosphinothricin acetyltransferase (PAT) encoding gene from the soil bacterium <i>Streptomyces hygroscopicus</i> .
LLRICE601	Bayer CropScience (Aventis CropScience(AgrEvo))	Glufosinate ammonium herbicide tolerant rice produced by inserting a modified phosphinothricin acetyltransferase (PAT) encoding gene from the soil bacterium <i>Streptomyces hygroscopicus</i> .
PWC16	BASF Inc.	Tolerance to the imidazolinone herbicide, imazethapyr, induced by chemical mutagenesis of the acetolactate synthase (ALS) enzyme using ethyl methanesulfonate (EMS).

Table 2D *Glycine max* L. Soybean

Event	Company	Description
A5547-127	Bayer CropScience (Aventis CropScience (AgrEvo))	Glufosinate ammonium herbicide tolerant soybean produced by inserting a modified phosphinothricin acetyltransferase (PAT) encoding gene from the soil bacterium <i>Streptomyces viridochromogenes</i> .
BPS-CV127-9	BASF Inc.	The introduced <i>csr1-2</i> gene from <i>Arabidopsis thaliana</i> encodes an acetohydroxyacid synthase protein that confers tolerance to imidazolinone herbicides due to a point mutation that results in a single amino acid substitution in which the serine residue at position 653 is replaced by asparagine (S653N).
DP-305423	Pioneer Hi-Bred International Inc.	High oleic acid soybean produced by inserting additional copies of a portion of the omega-6 desaturase encoding gene, <i>gm-fad2-1</i> resulting in silencing of the endogenous omega-6 desaturase gene (<i>FAD2-1</i>).
DP356043	Pioneer Hi-Bred International Inc.	Soybean event with two herbicide tolerance genes: glyphosate N-acetyltransferase, which detoxifies glyphosate, and a modified acetolactate synthase (ALS) gene which is tolerant to ALS-inhibiting herbicides.
G94-1, G94-19, G168	DuPont Canada Agricultural Products	High oleic acid soybean produced by inserting a second copy of the fatty acid desaturase (<i>GmFad2-1</i>) encoding gene from soybean, which resulted in "silencing" of the endogenous host gene.
GTS 40-3-2	Monsanto Company	Glyphosate tolerant soybean variety produced by inserting a modified 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) encoding gene from the soil bacterium <i>Agrobacterium tumefaciens</i> .
GU262	Bayer CropScience (Aventis CropScience(AgrEvo))	Glufosinate ammonium herbicide tolerant soybean produced by inserting a modified phosphinothricin acetyltransferase (PAT) encoding gene from the soil bacterium <i>Streptomyces viridochromogenes</i> .
MON87701	Monsanto Company	Resistance to lepidopteran pests of soybean including velvetbean caterpillar (<i>Anticarsia gemmatilis</i>) and soybean looper (<i>Pseudoplusia includens</i>).

Event	Company	Description
MON87701 x MON89788	Monsanto Company	Glyphosate herbicide tolerance through expression of the EPSPS encoding gene from <i>A. tumefaciens</i> strain CP4, and resistance to lepidopteran pests of soybean including velvetbean caterpillar (<i>Anticarsia gemmatilis</i>) and soybean looper (<i>Pseudoplusia includens</i>) via expression of the Cry1Ac encoding gene from <i>B. thuringiensis</i> .
MON89788	Monsanto Company	Glyphosate-tolerant soybean produced by inserting a modified 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) encoding <i>aroA</i> (<i>epsps</i>) gene from <i>Agrobacterium tumefaciens</i> CP4.
OT96-15	Agriculture & Agri-Food Canada	Low linolenic acid soybean produced through traditional cross-breeding to incorporate the novel trait from a naturally occurring <i>fan1</i> gene mutant that was selected for low linolenic acid.
W62, W98	Bayer CropScience (Aventis CropScience(AgrEvo))	Glufosinate ammonium herbicide tolerant soybean produced by inserting a modified phosphinothricin acetyltransferase (PAT) encoding gene from the soil bacterium <i>Streptomyces hygroscopicus</i> .

Table 2E *Zea mays* L. Maize

Event	Company	Description
176	Syngenta Seeds, Inc.	Insect-resistant maize produced by inserting the <i>cry1Ab</i> gene from <i>Bacillus thuringiensis</i> subsp. <i>kurstaki</i> . The genetic modification affords resistance to attack by the European corn borer (ECB).
3751IR	Pioneer Hi-Bred International Inc.	Selection of somaclonal variants by culture of embryos on imidazolinone containing media.
676, 678, 680	Pioneer Hi-Bred International Inc.	Male-sterile and glufosinate ammonium herbicide tolerant maize produced by inserting genes encoding DNA adenine methylase and phosphinothricin acetyltransferase (PAT) from <i>Escherichia coli</i> and <i>Streptomyces viridochromogenes</i> , respectively.
B16 (DLL25)	Dekalb Genetics Corporation	Glufosinate ammonium herbicide tolerant maize produced by inserting the gene encoding phosphinothricin acetyltransferase (PAT) from <i>Streptomyces hygroscopicus</i> .
BT11 (X4334CBR, X4734CBR)	Syngenta Seeds, Inc.	Insect-resistant and herbicide tolerant maize produced by inserting the <i>cry1Ab</i> gene from <i>Bacillus thuringiensis</i> subsp. <i>kurstaki</i> , and the phosphinothricin N-acetyltransferase (PAT) encoding gene from <i>S. viridochromogenes</i> .
BT11 x GA21	Syngenta Seeds, Inc.	Stacked insect resistant and herbicide tolerant maize produced by conventional cross breeding of parental lines BT11 (OECD unique identifier: SYN-BTO11-1) and GA21 (OECD unique identifier: MON-00021-9).
BT11 x MIR162	Syngenta Seeds, Inc.	Stacked insect resistant and herbicide tolerant maize produced by conventional cross breeding of parental lines BT11 (OECD unique identifier: SYN-BTO11-1) and MIR162 (OECD unique identifier: SYN-IR162-4). Resistance to the European Corn Borer and tolerance to the herbicide glufosinate ammonium (Liberty) is derived from BT11, which contains the <i>cry1Ab</i> gene from <i>Bacillus thuringiensis</i> subsp. <i>kurstaki</i> , and the phosphinothricin N-acetyltransferase (PAT) encoding gene from <i>S. viridochromogenes</i> . Resistance to other lepidopteran pests, including <i>H. zea</i> , <i>S. frugiperda</i> , <i>A. ipsilon</i> , and <i>S. albicosta</i> , is derived from MIR162, which contains the <i>vip3Aa</i> gene from <i>Bacillus thuringiensis</i> strain AB88.
BT11 x MIR162 x MIR604	Syngenta Seeds, Inc.	<i>Bacillus thuringiensis</i> Cry1Ab delta-endotoxin protein and the genetic material necessary for its production (via elements of vector pZO1502) in Event Bt11 corn (OECD Unique Identifier: SYN-BTO11-1) x <i>Bacillus thuringiensis</i> Vip3Aa20 insecticidal protein and the genetic material necessary for its production (via elements of vector pNOV1300) in Event MIR162 maize (OECD Unique Identifier: SYN-IR162-4) x modified Cry3A protein and the genetic material necessary for its production (via elements of vector pZM26) in Event MIR604 corn (OECD Unique Identifier: SYN-IR604-5).

Event	Company	Description
BT11 x MIR162 x MIR604 x GA21	Syngenta Seeds, Inc.	Resistance to coleopteran pests, particularly corn rootworm pests (<i>Diabrotica</i> spp.) and several lepidopteran pests of corn, including European corn borer (ECB, <i>Ostrinia nubilalis</i>), corn earworm (CEW, <i>Helicoverpa zea</i>), fall army worm (FAW, <i>Spodoptera frugiperda</i>), and black cutworm (BCW, <i>Agrotis ipsilon</i>); tolerance to glyphosate and glufosinate-ammonium containing herbicides.
BT11 x MIR604	Syngenta Seeds, Inc.	Stacked insect resistant and herbicide tolerant maize produced by conventional cross breeding of parental lines BT11 (OECD unique identifier: SYN-BTO11-1) and MIR604 (OECD unique identifier: SYN-IR6O5-5). Resistance to the European Corn Borer and tolerance to the herbicide glufosinate ammonium (Liberty) is derived from BT11, which contains the <i>cry1Ab</i> gene from <i>Bacillus thuringiensis</i> subsp. kurstaki, and the phosphinothricin N-acetyltransferase (PAT) encoding gene from <i>S. viridochromogenes</i> . Corn rootworm-resistance is derived from MIR604 which contains the <i>mcry3A</i> gene from <i>Bacillus thuringiensis</i> .
BT11 x MIR604 x GA21	Syngenta Seeds, Inc.	Stacked insect resistant and herbicide tolerant maize produced by conventional cross breeding of parental lines BT11 (OECD unique identifier: SYN-BTO11-1), MIR604 (OECD unique identifier: SYN-IR6O5-5) and GA21 (OECD unique identifier: MON-OOO21-9). Resistance to the European Corn Borer and tolerance to the herbicide glufosinate ammonium (Liberty) is derived from BT11, which contains the <i>cry1Ab</i> gene from <i>Bacillus thuringiensis</i> subsp. kurstaki, and the phosphinothricin N-acetyltransferase (PAT) encoding gene from <i>S. viridochromogenes</i> . Corn rootworm-resistance is derived from MIR604 which contains the <i>mcry3A</i> gene from <i>Bacillus thuringiensis</i> . Tolerance to glyphosate herbicide is derived from GA21 which contains a modified EPSPS gene from maize.
CBH-351	Aventis CropScience	Insect-resistant and glufosinate ammonium herbicide tolerant maize developed by inserting genes encoding Cry9C protein from <i>Bacillus thuringiensis</i> subsp. tolworthi and phosphinothricin acetyltransferase (PAT) from <i>Streptomyces hygroscopicus</i> .
DAS-06275-8	DOW AgroSciences LLC	Lepidopteran insect resistant and glufosinate ammonium herbicide-tolerant maize variety produced by inserting the <i>cry1F</i> gene from <i>Bacillus thuringiensis</i> var. <i>aizawai</i> and the phosphinothricin acetyltransferase (PAT) from <i>Streptomyces hygroscopicus</i> .
DAS-59122-7	DOW AgroSciences LLC and Pioneer Hi-Bred International Inc.	Corn rootworm-resistant maize produced by inserting the <i>cry34Ab1</i> and <i>cry35Ab1</i> genes from <i>Bacillus thuringiensis</i> strain PS149B1. The PAT encoding gene from <i>Streptomyces viridochromogenes</i> was introduced as a selectable marker.

Event	Company	Description
DAS-59122-7 x NK603	DOW AgroSciences LLC and Pioneer Hi-Bred International Inc.	Stacked insect resistant and herbicide tolerant maize produced by conventional cross breeding of parental lines DAS-59122-7 (OECD unique identifier: DAS-59122-7) with NK603 (OECD unique identifier: MON-OO6O3-6). Corn rootworm-resistance is derived from DAS-59122-7 which contains the <i>cry34Ab1</i> and <i>cry35Ab1</i> genes from <i>Bacillus thuringiensis</i> strain PS149B1. Tolerance to glyphosate herbicide is derived from NK603.
DAS-59122-7 x TC1507 x NK603	DOW AgroSciences LLC and Pioneer Hi-Bred International Inc.	Stacked insect resistant and herbicide tolerant maize produced by conventional cross breeding of parental lines DAS-59122-7 (OECD unique identifier: DAS-59122-7) and TC1507 (OECD unique identifier: DAS-O15O7-1) with NK603 (OECD unique identifier: MON-OO6O3-6). Corn rootworm-resistance is derived from DAS-59122-7 which contains the <i>cry34Ab1</i> and <i>cry35Ab1</i> genes from <i>Bacillus thuringiensis</i> strain PS149B1. Lepidopteran resistance and tolerance to glufosinate ammonium herbicide is derived from TC1507. Tolerance to glyphosate herbicide is derived from NK603.
DBT418	Dekalb Genetics Corporation	Insect-resistant and glufosinate ammonium herbicide tolerant maize developed by inserting genes encoding Cry1AC protein from <i>Bacillus thuringiensis</i> subsp kurstaki and phosphinothricin acetyltransferase (PAT) from <i>Streptomyces hygroscopicus</i>
DK404SR	BASF Inc.	Somaclonal variants with a modified acetyl-CoA-carboxylase (ACCase) were selected by culture of embryos on sethoxydim enriched medium.
Event 3272	Syngenta Seeds, Inc.	Maize line expressing a heat stable alpha-amylase gene <i>amy797E</i> for use in the dry-grind ethanol process. The phosphomannose isomerase gene from <i>E.coli</i> was used as a selectable marker.
Event 98140	Pioneer Hi-Bred International Inc.	Maize event expressing tolerance to glyphosate herbicide, via expression of a modified bacterial glyphosate N-acetyltransferase, and ALS-inhibiting herbicides, vial expression of a modified form of the maize acetolactate synthase enzyme.
EXP1910IT	Syngenta Seeds, Inc. (formerly Zeneca Seeds)	Tolerance to the imidazolinone herbicide, imazethapyr, induced by chemical mutagenesis of the acetolactate synthase (ALS) enzyme using ethyl methanesulfonate (EMS).
GA21	Syngenta Seeds, Inc. (formerly Zeneca Seeds)	Introduction, by particle bombardment, of a modified 5-enolpyruvyl shikimate-3-phosphate synthase (EPSPS), an enzyme involved in the shikimate biochemical pathway for the production of the aromatic amino acids.
GA21 x MON810	Monsanto Company	Stacked insect resistant and herbicide tolerant corn hybrid derived from conventional cross-breeding of the parental lines GA21 (OECD identifier: MON-OOO21-9) and MON810 (OECD identifier: MON-OO81O-6).

IT	Pioneer Hi-Bred International Inc.	Tolerance to the imidazolinone herbicide, imazethapyr, was obtained by in vitro selection of somaclonal variants.
LY038	Monsanto Company	Altered amino acid composition, specifically elevated levels of lysine, through the introduction of the <i>cordapA</i> gene, derived from <i>Corynebacterium glutamicum</i> , encoding the enzyme dihydrodipicolinate synthase (cDHDPS).
MIR162	Syngenta Seeds, Inc.	Insect-resistant maize event expressing a Vip3A protein from <i>Bacillus thuringiensis</i> and the <i>Escherichia coli</i> PMI selectable marker
MIR604	Syngenta Seeds, Inc.	Corn rootworm resistant maize produced by transformation with a modified <i>cry3A</i> gene. The phosphomannose isomerase gene from <i>E.coli</i> was used as a selectable marker.
MIR604 x GA21	Syngenta Seeds, Inc.	Stacked insect resistant and herbicide tolerant maize produced by conventional cross breeding of parental lines MIR604 (OECD unique identifier: SYN-IR605-5) and GA21 (OECD unique identifier: MON-00021-9). Corn rootworm-resistance is derived from MIR604 which contains the <i>mcry3A</i> gene from <i>Bacillus thuringiensis</i> . Tolerance to glyphosate herbicide is derived from GA21.
MON80100	Monsanto Company	Insect-resistant maize produced by inserting the <i>cry1Ab</i> gene from <i>Bacillus thuringiensis</i> subsp. <i>kurstaki</i> . The genetic modification affords resistance to attack by the European corn borer (ECB).
MON802	Monsanto Company	Insect-resistant and glyphosate herbicide tolerant maize produced by inserting the genes encoding the Cry1Ab protein from <i>Bacillus thuringiensis</i> and the 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) from <i>A. tumefaciens</i> strain CP4.
MON809	Pioneer Hi-Bred International Inc.	Resistance to European corn borer (<i>Ostrinia nubilalis</i>) by introduction of a synthetic <i>cry1Ab</i> gene. Glyphosate resistance via introduction of the bacterial version of a plant enzyme, 5-enolpyruvyl shikimate-3-phosphate synthase (EPSPS).
MON810	Monsanto Company	Insect-resistant maize produced by inserting a truncated form of the <i>cry1Ab</i> gene from <i>Bacillus thuringiensis</i> subsp. <i>kurstaki</i> HD-1. The genetic modification affords resistance to attack by the European corn borer (ECB).
MON810 x LY038	Monsanto Company	Stacked insect resistant and enhanced lysine content maize derived from conventional cross-breeding of the parental lines MON810 (OECD identifier: MON-00810-6) and LY038 (OECD identifier: REN-00038-3).
MON810 x MON88017	Monsanto Company	Stacked insect resistant and glyphosate tolerant maize derived from conventional cross-breeding of the parental lines MON810 (OECD identifier: MON-00810-6) and MON88017 (OECD identifier: MON-88017-3). European corn borer (ECB) resistance is derived from a truncated form of the <i>cry1Ab</i> gene from <i>Bacillus thuringiensis</i> subsp. <i>kurstaki</i> HD-1 present in MON810. Corn rootworm resistance is derived from the <i>cry3Bb1</i> gene from <i>Bacillus thuringiensis</i> subspecies <i>kumamotoensis</i> strain EG4691 present in MON88017. Glyphosate tolerance is derived from a 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) encoding gene from <i>Agrobacterium tumefaciens</i> strain CP4 present in MON88017.
MON832	Monsanto Company	Introduction, by particle bombardment, of glyphosate oxidase (GOX) and a modified 5-enolpyruvyl shikimate-3-phosphate synthase (EPSPS), an enzyme involved in the shikimate biochemical pathway for the production of the aromatic amino acids.
MON863	Monsanto Company	Corn root worm resistant maize produced by inserting the <i>cry3Bb1</i> gene from <i>Bacillus thuringiensis</i> subsp. <i>kumamotoensis</i> .

MON863 x MON810	Monsanto Company	Stacked insect resistant corn hybrid derived from conventional cross-breeding of the parental lines MON863 (OECD identifier: MON-OO863-5) and MON810 (OECD identifier: MON-OO810-6)
MON863 x MON810 x NK603	Monsanto Company	Stacked insect resistant and herbicide tolerant corn hybrid derived from conventional cross-breeding of the stacked hybrid MON-OO863-5 x MON-OO810-6 and NK603 (OECD identifier: MON-OO603-6).
MON863 x NK603	Monsanto Company	Stacked insect resistant and herbicide tolerant corn hybrid derived from conventional cross-breeding of the parental lines MON863 (OECD identifier: MON-OO863-5) and NK603 (OECD identifier: MON-OO603-6).
MON87460	Monsanto Company	MON 87460 was developed to provide reduced yield loss under water-limited conditions compared to conventional maize. Efficacy in MON 87460 is derived by expression of the inserted <i>Bacillus subtilis</i> cold shock protein B (CspB).
MON88017	Monsanto Company	Corn rootworm-resistant maize produced by inserting the <i>cry3Bb1</i> gene from <i>Bacillus thuringiensis</i> subspecies <i>kumamotoensis</i> strain EG4691. Glyphosate tolerance derived by inserting a 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) encoding gene from <i>Agrobacterium tumefaciens</i> strain CP4.
MON89034	Monsanto Company	Maize event expressing two different insecticidal proteins from <i>Bacillus thuringiensis</i> providing resistance to number of lepidopteran pests.
MON89034 x MON88017	Monsanto Company	Stacked insect resistant and glyphosate tolerant maize derived from conventional cross-breeding of the parental lines MON89034 (OECD identifier: MON-89034-3) and MON88017 (OECD identifier: MON-88017-3). Resistance to Lepidopteran insects is derived from two <i>cry</i> genes present in MON89043. Corn rootworm resistance is derived from a single <i>cry</i> genes and glyphosate tolerance is derived from the 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) encoding gene from <i>Agrobacterium tumefaciens</i> present in MON88017.
MON89034 x NK603	Monsanto Company	Stacked insect resistant and herbicide tolerant maize produced by conventional cross breeding of parental lines MON89034 (OECD identifier: MON-89034-3) with NK603 (OECD unique identifier: MON-OO603-6). Resistance to Lepidopteran insects is derived from two <i>cry</i> genes present in MON89043. Tolerance to glyphosate herbicide is derived from NK603.
MON89034 x TC1507 x MON88017 x DAS-59122-7	Monsanto Company and Mycogen Seeds c/o Dow AgroSciences LLC	Stacked insect resistant and herbicide tolerant maize produced by conventional cross breeding of parental lines: MON89034, TC1507, MON88017, and DAS-59122. Resistance to the above-ground and below-ground insect pests and tolerance to glyphosate and glufosinate-ammonium containing herbicides.
MS3	Bayer CropScience (Aventis CropScience(AgrEvo))	Male sterility caused by expression of the barnase ribonuclease gene from <i>Bacillus amyloliquefaciens</i> ; PPT resistance was via PPT-acetyltransferase (PAT).
MS6	Bayer CropScience (Aventis CropScience(AgrEvo))	Male sterility caused by expression of the barnase ribonuclease gene from <i>Bacillus amyloliquefaciens</i> ; PPT resistance was via PPT-acetyltransferase (PAT).

NK603	Monsanto Company	Introduction, by particle bombardment, of a modified 5-enolpyruvyl shikimate-3-phosphate synthase (EPSPS), an enzyme involved in the shikimate biochemical pathway for the production of the aromatic amino acids.
NK603 x MON810	Monsanto Company	Stacked insect resistant and herbicide tolerant corn hybrid derived from conventional cross-breeding of the parental lines NK603 (OECD identifier: MON-OO6O3-6) and MON810 (OECD identifier: MON-OO81O-6).
NK603 x T25	Monsanto Company	Stacked glufosinate ammonium and glyphosate herbicide tolerant maize hybrid derived from conventional cross-breeding of the parental lines NK603 (OECD identifier: MON-OO6O3-6) and T25 (OECD identifier: ACS-ZM003-2).
T14, T25	Bayer CropScience (Aventis CropScience(AgrEvo))	Glufosinate herbicide tolerant maize produced by inserting the phosphinothricin N-acetyltransferase (PAT) encoding gene from the aerobic actinomycete <i>Streptomyces viridochromogenes</i> .
T25 x MON810	Bayer CropScience (Aventis CropScience(AgrEvo))	Stacked insect resistant and herbicide tolerant corn hybrid derived from conventional cross-breeding of the parental lines T25 (OECD identifier: ACS-ZMOO3-2) and MON810 (OECD identifier: MON-OO81O-6).
TC1507	Mycogen (c/o Dow AgroSciences); Pioneer (c/o DuPont)	Insect-resistant and glufosinate ammonium herbicide tolerant maize produced by inserting the <i>cry1F</i> gene from <i>Bacillus thuringiensis</i> var. <i>aizawai</i> and the phosphinothricin N-acetyltransferase encoding gene from <i>Streptomyces viridochromogenes</i> .
TC1507 x DAS-59122-7	DOW AgroSciences LLC and Pioneer Hi-Bred International Inc.	Stacked insect resistant and herbicide tolerant maize produced by conventional cross breeding of parental lines TC1507 (OECD unique identifier: DAS-O15O7-1) with DAS-59122-7 (OECD unique identifier: DAS-59122-7). Resistance to lepidopteran insects is derived from TC1507 due the presence of the <i>cry1F</i> gene from <i>Bacillus thuringiensis</i> var. <i>aizawai</i> . Corn rootworm-resistance is derived from DAS-59122-7 which contains the <i>cry34Ab1</i> and <i>cry35Ab1</i> genes from <i>Bacillus thuringiensis</i> strain PS149B1. Tolerance to glufosinate ammonium herbicide is derived from TC1507 from the phosphinothricin N-acetyltransferase encoding gene from <i>Streptomyces viridochromogenes</i> .
TC1507 x NK603	DOW AgroSciences LLC	Stacked insect resistant and herbicide tolerant corn hybrid derived from conventional cross-breeding of the parental lines 1507 (OECD identifier: DAS-O15O7-1) and NK603 (OECD identifier: MON-OO6O3-6).

Table 2F *Triticum aestivum* Wheat

Event	Company	Description
AP205CL	BASF Inc.	Selection for a mutagenized version of the enzyme acetohydroxyacid synthase (AHAS), also known as acetolactate synthase (ALS) or acetolactate pyruvate- lyase.
AP602CL	BASF Inc.	Selection for a mutagenized version of the enzyme acetohydroxyacid synthase (AHAS), also known as acetolactate synthase (ALS) or acetolactate pyruvate- lyase.
BW255-2, BW238-3	BASF Inc.	Selection for a mutagenized version of the enzyme acetohydroxyacid synthase (AHAS), also known as acetolactate synthase (ALS) or acetolactate pyruvate- lyase.
BW7	BASF Inc.	Tolerance to imidazolinone herbicides induced by chemical mutagenesis of the acetohydroxyacid synthase (AHAS) gene using sodium azide.
MON71800	Monsanto Company	Glyphosate tolerant wheat variety produced by inserting a modified 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) encoding gene from the soil bacterium <i>Agrobacterium tumefaciens</i> , strain CP4.
SWP965001	Cyanamid Crop Protection	Selection for a mutagenized version of the enzyme acetohydroxyacid synthase (AHAS), also known as acetolactate synthase (ALS) or acetolactate pyruvate- lyase.
Teal 11A	BASF Inc.	Selection for a mutagenized version of the enzyme acetohydroxyacid synthase (AHAS), also known as acetolactate synthase (ALS) or acetolactate pyruvate- lyase.
A2704-12, A2704-21, A5547-35	Bayer CropScience (Aventis CropScience (AgrEvo))	Glufosinate ammonium herbicide tolerant soybean produced by inserting a modified phosphinothricin acetyltransferase (PAT) encoding gene from the soil bacterium <i>Streptomyces viridochromogenes</i> .

5

Other events with regulatory approval are well known to one skilled in the art and can be found at the Center for Environmental Risk Assessment (ceramc.org/?action=gm_crop_database, which can be accessed using the www prefix) and at the International Service for the Acquisition of Agri-Biotech Applications (isaaa.org/gmapprovaldatabase/default.asp, which can be accessed using the www prefix).

10

Gene silencing

In some embodiments the stacked trait may be in the form of silencing of one or more polynucleotides of interest resulting in suppression of one or more target pest

15

polypeptides. In some embodiments the silencing is achieved through the use of a suppression DNA construct.

In some embodiments one or more of the PHI-4 polypeptides or fragments or variants thereof may be stacked with one or more polynucleotides encoding one or more polypeptides having insecticidal activity or agronomic traits as set forth supra and optionally may further include one or more polynucleotides providing for gene silencing of one or more target polynucleotides as discussed infra.

“Suppression DNA construct” is a recombinant DNA construct which when transformed or stably integrated into the genome of the plant, results in “silencing” of a target gene in the plant. The target gene may be endogenous or transgenic to the plant. “Silencing,” as used herein with respect to the target gene, refers generally to the suppression of levels of mRNA or protein/enzyme expressed by the target gene, and/or the level of the enzyme activity or protein functionality. The term “suppression” includes lower, reduce, decline, decrease, inhibit, eliminate and prevent. “Silencing” or “gene silencing” does not specify mechanism and is inclusive, and not limited to, anti-sense, cosuppression, viral-suppression, hairpin suppression, stem-loop suppression, RNAi-based approaches and small RNA-based approaches.

A suppression DNA construct may comprise a region derived from a target gene of interest and may comprise all or part of the nucleic acid sequence of the sense strand (or antisense strand) of the target gene of interest. Depending upon the approach to be utilized, the region may be 100% identical or less than 100% identical (e.g., at least 50% or any integer between 51% and 100% identical) to all or part of the sense strand (or antisense strand) of the gene of interest.

Suppression DNA constructs are well-known in the art, are readily constructed once the target gene of interest is selected, and include, without limitation, cosuppression constructs, antisense constructs, viral-suppression constructs, hairpin suppression constructs, stem-loop suppression constructs, double-stranded RNA-producing constructs, and more generally, RNAi (RNA interference) constructs and small RNA constructs such as siRNA (short interfering RNA) constructs and miRNA (microRNA) constructs.

“Antisense inhibition” refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein.

“Antisense RNA” refers to an RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target isolated nucleic acid fragment (US Patent Number 5,107,065). The complementarity of an

antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns or the coding sequence.

“Cosuppression” refers to the production of sense RNA transcripts capable of suppressing the expression of the target protein. “Sense” RNA refers to RNA transcript
5 that includes the mRNA and can be translated into protein within a cell or in vitro. Cosuppression constructs in plants have been previously designed by focusing on overexpression of a nucleic acid sequence having homology to a native mRNA, in the sense orientation, which results in the reduction of all RNA having homology to the overexpressed sequence (see, Vaucheret, et al. (1998) *Plant J.* 16:651-659 and Gura,
10 (2000) *Nature* 404:804-808).

Another variation describes the use of plant viral sequences to direct the suppression of proximal mRNA encoding sequences (PCT Publication WO 1998/36083).

Recent work has described the use of “hairpin” structures that incorporate all or part, of an mRNA encoding sequence in a complementary orientation that results in a
15 potential “stem-loop” structure for the expressed RNA (PCT Publication Number WO 1999/53050). In this case the stem is formed by polynucleotides corresponding to the gene of interest inserted in either sense or anti-sense orientation with respect to the promoter and the loop is formed by some polynucleotides of the gene of interest, which do not have a complement in the construct. This increases the frequency of
20 cosuppression or silencing in the recovered transgenic plants. For review of hairpin suppression see, Wesley, et al., (2003) *Methods in Molecular Biology, Plant Functional Genomics: Methods and Protocols* 236:273-286.

A construct where the stem is formed by at least 30 nucleotides from a gene to be suppressed and the loop is formed by a random nucleotide sequence has also effectively
25 been used for suppression (WO 1999/61632).

The use of poly-T and poly-A sequences to generate the stem in the stem-loop structure has also been described (WO 2002/00894).

Yet another variation includes using synthetic repeats to promote formation of a stem in the stem-loop structure. Transgenic organisms prepared with such recombinant
30 DNA fragments have been shown to have reduced levels of the protein encoded by the nucleotide fragment forming the loop as described in PCT Publication Number WO 2002/00904.

RNA interference refers to the process of sequence-specific post-transcriptional gene silencing in animals mediated by short interfering RNAs (siRNAs) (Fire, et al., (1998)
35 *Nature* 391:806). The corresponding process in plants is commonly referred to as post-transcriptional gene silencing (PTGS) or RNA silencing and is also referred to as quelling

in fungi. The process of post-transcriptional gene silencing is thought to be an evolutionarily-conserved cellular defense mechanism used to prevent the expression of foreign genes and is commonly shared by diverse flora and phyla (Fire, *et al.*, (1999) *Trends Genet.* 15:358). Such protection from foreign gene expression may have evolved
5 in response to the production of double-stranded RNAs (dsRNAs) derived from viral infection or from the random integration of transposon elements into a host genome via a cellular response that specifically destroys homologous single-stranded RNA of viral genomic RNA. The presence of dsRNA in cells triggers the RNAi response through a mechanism that has yet to be fully characterized.

10 The presence of long dsRNAs in cells stimulates the activity of a ribonuclease III enzyme referred to as dicer. Dicer is involved in the processing of the dsRNA into short pieces of dsRNA known as short interfering RNAs (siRNAs) (Berstein, *et al.*, (2001) *Nature* 409:363). Short interfering RNAs derived from dicer activity are typically about 21 to about 23 nucleotides in length and comprise about 19 base pair duplexes (Elbashir, *et al.*, (2001) *Genes Dev.* 15:188). Dicer has also been implicated in the excision of 21- and
15 22-nucleotide small temporal RNAs (stRNAs) from precursor RNA of conserved structure that are implicated in translational control (Hutvagner, *et al.*, (2001) *Science* 293:834). The RNAi response also features an endonuclease complex, commonly referred to as an RNA-induced silencing complex (RISC), which mediates cleavage of single-stranded RNA
20 having sequence complementarity to the antisense strand of the siRNA duplex. Cleavage of the target RNA takes place in the middle of the region complementary to the antisense strand of the siRNA duplex (Elbashir, *et al.*, (2001) *Genes Dev.* 15:188). In addition, RNA interference can also involve small RNA (e.g., miRNA) mediated gene silencing, presumably through cellular mechanisms that regulate chromatin structure and thereby
25 prevent transcription of target gene sequences (see, e.g., Allshire, (2002) *Science* 297:1818-1819; Volpe, *et al.*, (2002) *Science* 297:1833-1837; Jenuwein, (2002) *Science* 297:2215-2218; and Hall, *et al.*, (2002) *Science* 297:2232-2237). As such, miRNA molecules of the disclosure can be used to mediate gene silencing via interaction with RNA transcripts or alternately by interaction with particular gene sequences, wherein such
30 interaction results in gene silencing either at the transcriptional or post-transcriptional level.

Methods and compositions are further provided which allow for an increase in RNAi produced from the silencing element. In such embodiments, the methods and compositions employ a first polynucleotide comprising a silencing element for a target
35 pest sequence operably linked to a promoter active in the plant cell; and, a second polynucleotide comprising a suppressor enhancer element comprising the target pest

sequence or an active variant or fragment thereof operably linked to a promoter active in the plant cell. The combined expression of the silencing element with suppressor enhancer element leads to an increased amplification of the inhibitory RNA produced from the silencing element over that achievable with only the expression of the silencing element alone. In addition to the increased amplification of the specific RNAi species itself, the methods and compositions further allow for the production of a diverse population of RNAi species that can enhance the effectiveness of disrupting target gene expression. As such, when the suppressor enhancer element is expressed in a plant cell in combination with the silencing element, the methods and composition can allow for the systemic production of RNAi throughout the plant; the production of greater amounts of RNAi than would be observed with just the silencing element construct alone; and, the improved loading of RNAi into the phloem of the plant, thus providing better control of phloem feeding insects by an RNAi approach. Thus, the various methods and compositions provide improved methods for the delivery of inhibitory RNA to the target organism. See, for example, US 2009/0188008.

As used herein, a “suppressor enhancer element” comprises a polynucleotide comprising the target sequence to be suppressed or an active fragment or variant thereof. It is recognized that the suppressor enhancer element need not be identical to the target sequence, but rather, the suppressor enhancer element can comprise a variant of the target sequence, so long as the suppressor enhancer element has sufficient sequence identity to the target sequence to allow for an increased level of the RNAi produced by the silencing element over that achievable with only the expression of the silencing element. Similarly, the suppressor enhancer element can comprise a fragment of the target sequence, wherein the fragment is of sufficient length to allow for an increased level of the RNAi produced by the silencing element over that achievable with only the expression of the silencing element.

It is recognized that multiple suppressor enhancer elements from the same target sequence or from different target sequences or from different regions of the same target sequence can be employed. For example, the suppressor enhancer elements employed can comprise fragments of the target sequence derived from different region of the target sequence (i.e., from the 3'UTR, coding sequence, intron, and/or 5'UTR). Further, the suppressor enhancer element can be contained in an expression cassette, as described elsewhere herein, and in specific embodiments, the suppressor enhancer element is on the same or on a different DNA vector or construct as the silencing element. The suppressor enhancer element can be operably linked to a promoter as disclosed herein. It is recognized that the suppressor enhancer element can be expressed constitutively or

alternatively, it may be produced in a stage-specific manner employing the various inducible or tissue-preferred or developmentally regulated promoters that are discussed elsewhere herein.

In specific embodiments, employing both a silencing element and the suppressor enhancer element the systemic production of RNAi occurs throughout the entire plant. In further embodiments, the plant or plant parts of the disclosure have an improved loading of RNAi into the phloem of the plant than would be observed with the expression of the silencing element construct alone and, thus provide better control of phloem feeding insects by an RNAi approach. In specific embodiments, the plants, plant parts, and plant cells of the disclosure can further be characterized as allowing for the production of a diversity of RNAi species that can enhance the effectiveness of disrupting target gene expression.

In specific embodiments, the combined expression of the silencing element and the suppressor enhancer element increases the concentration of the inhibitory RNA in the plant cell, plant, plant part, plant tissue or phloem over the level that is achieved when the silencing element is expressed alone.

As used herein, an "increased level of inhibitory RNA" comprises any statistically significant increase in the level of RNAi produced in a plant having the combined expression when compared to an appropriate control plant. For example, an increase in the level of RNAi in the plant, plant part or the plant cell can comprise at least about a 1%, about a 1%-5%, about a 5%-10%, about a 10%-20%, about a 20%-30%, about a 30%-40%, about a 40%-50%, about a 50%-60%, about 60-70%, about 70%-80%, about a 80%-90%, about a 90%-100% or greater increase in the level of RNAi in the plant, plant part, plant cell or phloem when compared to an appropriate control. In other embodiments, the increase in the level of RNAi in the plant, plant part, plant cell or phloem can comprise at least about a 1 fold, about a 1 fold-5 fold, about a 5 fold-10 fold, about a 10 fold-20 fold, about a 20 fold-30 fold, about a 30 fold-40 fold, about a 40 fold-50 fold, about a 50 fold-60 fold, about 60 fold-70 fold, about 70 fold-80 fold, about a 80 fold-90 fold, about a 90 fold-100 fold or greater increase in the level of RNAi in the plant, plant part, plant cell or phloem when compared to an appropriate control. Examples of combined expression of the silencing element with suppressor enhancer element for the control of Stinkbugs and Lygus can be found in US 2011/0301223 and US 2009/0192117.

Some embodiments relate to down-regulation of expression of target genes in insect pest species by interfering ribonucleic acid (RNA) molecules. WO 2007/074405 describes methods of inhibiting expression of target genes in invertebrate pests including Colorado potato beetle. WO 2005/110068 describes methods of inhibiting expression of

target genes in invertebrate pests including in particular Western corn rootworm as a means to control insect infestation. Furthermore, WO 2009/091864 describes compositions and methods for the suppression of target genes from insect pest species including pests from the *Lygus* genus. Nucleic acid molecules including RNAi for

5 targeting the vacuolar ATPase H subunit, useful for controlling a coleopteran pest population and infestation as described in US Patent Application Publication 2012/0198586. WO 2012/055982 describes ribonucleic acid (RNA or double stranded RNA) that inhibits or down regulates the expression of a target gene that encodes: an

10 insect ribosomal protein such as the ribosomal protein L19, the ribosomal protein L40 or the ribosomal protein S27A; an insect proteasome subunit such as the Rpn6 protein, the Pros 25, the Rpn2 protein, the proteasome beta 1 subunit protein or the Pros beta 2 protein; an insect β -coatamer of the COPI vesicle, the γ -coatamer of the COPI vesicle, the β' - coatamer protein or the ζ -coatamer of the COPI vesicle; an insect Tetraspanine 2 A protein which is a putative transmembrane domain protein; an insect protein belonging to

15 the actin family such as Actin 5C; an insect ubiquitin-5E protein; an insect Sec23 protein which is a GTPase activator involved in intracellular protein transport; an insect crinkled protein which is an unconventional myosin which is involved in motor activity; an insect crooked neck protein which is involved in the regulation of nuclear alternative mRNA splicing; an insect vacuolar H⁺-ATPase G-subunit protein; and an insect Tbp-1 such as

20 Tat-binding protein. US Patent Application Publications 2012/029750, US 20120297501, and 2012/0322660 describe interfering ribonucleic acids (RNA or double stranded RNA) that functions upon uptake by an insect pest species to down-regulate expression of a target gene in said insect pest, wherein the RNA comprises at least one silencing element wherein the silencing element is a region of double-stranded RNA comprising annealed

25 complementary strands, one strand of which comprises or consists of a sequence of nucleotides which is at least partially complementary to a target nucleotide sequence within the target gene. US Patent Application Publication 2012/0164205 describe potential targets for interfering double stranded ribonucleic acids for inhibiting invertebrate pests including: a Chd3 Homologous Sequence, a Beta-Tubulin Homologous Sequence,

30 a 40 kDa V-ATPase Homologous Sequence, a EF1 α Homologous Sequence, a 26S Proteasome Subunit p28 Homologous Sequence, a Juvenile Hormone Epoxide Hydrolase Homologous Sequence, a Swelling Dependent Chloride Channel Protein Homologous Sequence, a Glucose-6-Phosphate 1-Dehydrogenase Protein Homologous Sequence, an Act42A Protein Homologous Sequence, a ADP-Ribosylation Factor 1 Homologous

35 Sequence, a Transcription Factor IIB Protein Homologous Sequence, a Chitinase Homologous Sequences, a Ubiquitin Conjugating Enzyme Homologous Sequence, a

Glyceraldehyde-3-Phosphate Dehydrogenase Homologous Sequence, an Ubiquitin B Homologous Sequence, a Juvenile Hormone Esterase Homolog, and an Alpha Tubulin Homologous Sequence.

5 Use in Pesticidal Control

General methods for employing strains comprising a nucleic acid sequence of the embodiments or a variant thereof, in pesticide control or in engineering other organisms as pesticidal agents are known in the art. See, for example US Patent Number 5,039,523 and EP 0480762A2.

10 Microorganism hosts that are known to occupy the "phytosphere" (phylloplane, phyllosphere, rhizosphere, and/or rhizoplane) of one or more crops of interest may be selected. These microorganisms are selected so as to be capable of successfully competing in the particular environment with the wild-type microorganisms, provide for stable maintenance and expression of the gene expressing the PHI-4 polypeptide, and
15 desirably, provide for improved protection of the pesticide from environmental degradation and inactivation.

Such microorganisms include bacteria, algae, and fungi. Of particular interest are microorganisms such as bacteria, e.g., *Pseudomonas*, *Erwinia*, *Serratia*, *Klebsiella*, *Xanthomonas*, *Streptomyces*, *Rhizobium*, *Rhodopseudomonas*, *Methylius*,
20 *Agrobacterium*, *Acetobacter*, *Lactobacillus*, *Arthrobacter*, *Azotobacter*, *Leuconostoc*, and *Alcaligenes*, fungi, particularly yeast, e.g., *Saccharomyces*, *Cryptococcus*, *Kluyveromyces*, *Sporobolomyces*, *Rhodotorula*, and *Aureobasidium*. Of particular interest are such phytosphere bacterial species as *Pseudomonas syringae*, *Pseudomonas fluorescens*, *Pseudomonas chlororaphis*, *Serratia marcescens*, *Acetobacter xylinum*,
25 *Agrobacteria*, *Rhodopseudomonas spheroides*, *Xanthomonas campestris*, *Rhizobium melioli*, *Alcaligenes entrophus*, *Clavibacter xyli* and *Azotobacter vinelandii* and phytosphere yeast species such as *Rhodotorula rubra*, *R. glutinis*, *R. marina*, *R. aurantiaca*, *Cryptococcus albidus*, *C. diffluens*, *C. laurentii*, *Saccharomyces rosei*, *S. pretoriensis*, *S. cerevisiae*, *Sporobolomyces roseus*, *S. odoratus*, *Kluyveromyces veronae*,
30 and *Aureobasidium pollulans*. Of particular interest are the pigmented microorganisms. Host organisms of particular interest include yeast, such as *Rhodotorula* spp., *Aureobasidium* spp., *Saccharomyces* spp. (such as *S. cerevisiae*), *Sporobolomyces* spp., phylloplane organisms such as *Pseudomonas* spp. (such as *P. aeruginosa*, *P. fluorescens*, *P. chlororaphis*), *Erwinia* spp., and *Flavobacterium* spp., and other such
35 organisms, including *Agrobacterium tumefaciens*, *E. coli*, *Bacillus subtilis*, *Bacillus cereus* and the like.

Genes encoding the PHI-4 polypeptides of the embodiments can be introduced into microorganisms that multiply on plants (epiphytes) to deliver PHI-4 polypeptides to potential target pests. Epiphytes, for example, can be gram-positive or gram-negative bacteria.

5 Root-colonizing bacteria, for example, can be isolated from the plant of interest by methods known in the art. Specifically, a *Bacillus cereus* strain that colonizes roots can be isolated from roots of a plant (see, for example, Handelsman, *et al.*, (1991) *Appl. Environ. Microbiol.* 56:713-718). Genes encoding the PHI-4 polypeptides of the embodiments can be introduced into a root-colonizing *Bacillus cereus* by standard
10 methods known in the art.

Genes encoding PHI-4 polypeptides can be introduced, for example, into the root-colonizing *Bacillus* by means of electro transformation. Specifically, genes encoding the PHI-4 polypeptides can be cloned into a shuttle vector, for example, pHT3101 (Lerecius, *et al.*, (1989) *FEMS Microbiol. Letts.* 60:211-218). The shuttle vector pHT3101 containing
15 the coding sequence for the particular PHI-4 polypeptide gene can, for example, be transformed into the root-colonizing *Bacillus* by means of electroporation (Lerecius, *et al.*, (1989) *FEMS Microbiol. Letts.* 60:211-218).

Expression systems can be designed so that PHI-4 polypeptides are secreted outside the cytoplasm of gram-negative bacteria, such as *E. coli*, for example.
20 Advantages of having PHI-4 polypeptides secreted are: (1) avoidance of potential cytotoxic effects of the PHI-4 polypeptide expressed; and (2) improvement in the efficiency of purification of the PHI-4 polypeptide, including, but not limited to, increased efficiency in the recovery and purification of the protein per volume cell broth and decreased time and/or costs of recovery and purification per unit protein.

25 PHI-4 polypeptides can be made to be secreted in *E. coli*, for example, by fusing an appropriate *E. coli* signal peptide to the amino-terminal end of the PHI-4 polypeptide. Signal peptides recognized by *E. coli* can be found in proteins already known to be secreted in *E. coli*, for example the OmpA protein (Ghrayeb, *et al.*, (1984) *EMBO J.* 3:2437-2442). OmpA is a major protein of the *E. coli* outer membrane, and thus its signal
30 peptide is thought to be efficient in the translocation process. Also, the OmpA signal peptide does not need to be modified before processing as may be the case for other signal peptides, for example lipoprotein signal peptide (Duffaud, *et al.*, (1987) *Meth. Enzymol.* 153:492).

PHI-4 polypeptides of the embodiments can be fermented in a bacterial host and
35 the resulting bacteria processed and used as a microbial spray in the same manner that *Bt* strains have been used as insecticidal sprays. In the case of a PHI-4 polypeptide that

is secreted from *Bacillus*, the secretion signal is removed or mutated using procedures known in the art. Such mutations and/or deletions prevent secretion of the PHI-4 polypeptide into the growth medium during the fermentation process. The PHI-4 polypeptides are retained within the cell, and the cells are then processed to yield the encapsulated PHI-4 polypeptides. Any suitable microorganism can be used for this purpose. *Pseudomonas* has been used to express *Bt* toxins as encapsulated proteins and the resulting cells processed and sprayed as an insecticide (Gaertner, *et al.*, (1993), in: *Advanced Engineered Pesticides*, ed. Kim).

Alternatively, the PHI-4 polypeptides are produced by introducing a heterologous gene into a cellular host. Expression of the heterologous gene results, directly or indirectly, in the intracellular production and maintenance of the pesticide. These cells are then treated under conditions that prolong the activity of the toxin produced in the cell when the cell is applied to the environment of target pest(s). The resulting product retains the toxicity of the toxin. These naturally encapsulated PHI-4 polypeptides may then be formulated in accordance with conventional techniques for application to the environment hosting a target pest, e.g., soil, water, and foliage of plants. See, for example EPA 0192319, and the references cited therein.

Pesticidal Compositions

In some embodiments the active ingredients can be applied in the form of compositions and can be applied to the crop area or plant to be treated, simultaneously or in succession, with other compounds. These compounds can be fertilizers, weed killers, cryoprotectants, surfactants, detergents, pesticidal soaps, dormant oils, polymers, and/or time-release or biodegradable carrier formulations that permit long-term dosing of a target area following a single application of the formulation. They can also be selective herbicides, chemical insecticides, virucides, microbicides, amoebicides, pesticides, fungicides, bacteriocides, nematocides, molluscicides or mixtures of several of these preparations, if desired, together with further agriculturally acceptable carriers, surfactants or application-promoting adjuvants customarily employed in the art of formulation. Suitable carriers and adjuvants can be solid or liquid and correspond to the substances ordinarily employed in formulation technology, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, binders or fertilizers. Likewise the formulations may be prepared into edible "baits" or fashioned into pest "traps" to permit feeding or ingestion by a target pest of the pesticidal formulation.

Methods of applying an active ingredient or an agrochemical composition that contains at least one of the PHI-4 polypeptides produced by the bacterial strains include

leaf application, seed coating and soil application. The number of applications and the rate of application depend on the intensity of infestation by the corresponding pest.

The composition may be formulated as a powder, dust, pellet, granule, spray, emulsion, colloid, solution or such like, and may be prepared by such conventional means
5 as desiccation, lyophilization, homogenation, extraction, filtration, centrifugation, sedimentation or concentration of a culture of cells comprising the polypeptide. In all such compositions that contain at least one such pesticidal polypeptide, the polypeptide may be present in a concentration of from about 1% to about 99% by weight.

Lepidopteran, dipteran, heteropteran, nematode, hemiptera or coleopteran pests
10 may be killed or reduced in numbers in a given area by the methods of the disclosure or may be prophylactically applied to an environmental area to prevent infestation by a susceptible pest. Preferably the pest ingests or is contacted with, a pesticidally-effective amount of the polypeptide. By "pesticidally-effective amount" is intended an amount of the pesticide that is able to bring about death to at least one pest or to noticeably reduce
15 pest growth, feeding or normal physiological development. This amount will vary depending on such factors as, for example, the specific target pests to be controlled, the specific environment, location, plant, crop or agricultural site to be treated, the environmental conditions, and the method, rate, concentration, stability, and quantity of application of the pesticidally-effective polypeptide composition. The formulations may
20 also vary with respect to climatic conditions, environmental considerations, and/or frequency of application and/or severity of pest infestation.

The pesticide compositions described may be made by formulating either the bacterial cell, crystal and/or spore suspension or isolated protein component with the desired agriculturally-acceptable carrier. The compositions may be formulated prior to
25 administration in an appropriate means such as lyophilized, freeze-dried, desiccated or in an aqueous carrier, medium or suitable diluent, such as saline or other buffer. The formulated compositions may be in the form of a dust or granular material or a suspension in oil (vegetable or mineral) or water or oil/water emulsions or as a wettable powder or in combination with any other carrier material suitable for agricultural application. Suitable
30 agricultural carriers can be solid or liquid and are well known in the art. The term "agriculturally-acceptable carrier" covers all adjuvants, inert components, dispersants, surfactants, tackifiers, binders, etc. that are ordinarily used in pesticide formulation technology; these are well known to those skilled in pesticide formulation. The formulations may be mixed with one or more solid or liquid adjuvants and prepared by
35 various means, e.g., by homogeneously mixing, blending and/or grinding the pesticidal composition with suitable adjuvants using conventional formulation techniques. Suitable

formulations and application methods are described in US Patent Number 6,468,523, herein incorporated by reference. The plants can also be treated with one or more chemical compositions, including one or more herbicide, insecticides or fungicides. Exemplary chemical compositions include: Fruits/Vegetables Herbicides: Atrazine, Bromacil, Diuron, Glyphosate, Linuron, Metribuzin, Simazine, Trifluralin, Fluazifop, Glufosinate, Halo sulfuron Gowan, Paraquat, Propyzamide, Sethoxydim, Butafenacil, Halosulfuron, Indaziflam; Fruits/Vegetables Insecticides: Aldicarb, *Bacillus thuriengensis*, Carbaryl, Carbofuran, Chlorpyrifos, Cypermethrin, Deltamethrin, Diazinon, Malathion, Abamectin, Cyfluthrin/beta-cyfluthrin, Esfenvalerate, Lambda-cyhalothrin, Acequinocyl, Bifenazate, Methoxyfenozide, Novaluron, Chromafenozide, Thiacloprid, Dinotefuran, Fluacrypyrim, Tolfeppyrad, Clothianidin, Spirodiclofen, Gamma-cyhalothrin, Spiromesifen, Spinosad, Rynaxypyr, Cyazypyr, Spinoteram, Triflumuron, Spirotetramat, Imidacloprid, Flubendiamide, Thiodicarb, Metaflumizone, Sulfoxaflor, Cyflumetofen, Cyanopyrafen, Imidacloprid, Clothianidin, Thiamethoxam, Spinotoram, Thiodicarb, Flonicamid, Methiocarb, Emamectin-benzoate, Indoxacarb, Forthiazate, Fenamiphos, Cadusaphos, Pyriproxifen, Fenbutatin-oxid, Hexthiazox, Methomyl, 4-[[[(6-Chlorpyridin-3-yl)methyl](2,2-difluorethyl)amino]furan-2(5H)-on; Fruits/Vegetables Fungicides: Carbendazim, Chlorothalonil, EBDCs, Sulphur, Thiophanate-methyl, Azoxystrobin, Cymoxanil, Fluazinam, Fosetyl, Iprodione, Kresoxim-methyl, Metalaxyl/mefenoxam, Trifloxystrobin, Ethaboxam, Iprovalicarb, Trifloxystrobin, Fenhexamid, Oxpoconazole fumarate, Cyazofamid, Fenamidone, Zoxamide, Picoxystrobin, Pyraclostrobin, Cyflufenamid, Boscalid; Cereals Herbicides: Isoproturon, Bromoxynil, Ioxynil, Phenoxies, Chlorsulfuron, Clodinafop, Diclofop, Diflufenican, Fenoxaprop, Florasulam, Fluoroxypyr, Metsulfuron, Triasulfuron, Flucarbazone, Iodosulfuron, Propoxycarbazone, Picolinafen, Mesosulfuron, Beflubutamid, Pinoxaden, Amidosulfuron, Thifensulfuron Methyl, Tribenuron, Flupyrsulfuron, Sulfosulfuron, Pyrasulfotole, Pyroxsulam, Flufenacet, Tralkoxydim, Pyroxasulfon; Cereals Fungicides: Carbendazim, Chlorothalonil, Azoxystrobin, Cyproconazole, Cyprodinil, Fenpropimorph, Epoxiconazole, Kresoxim-methyl, Quinoxifen, Tebuconazole, Trifloxystrobin, Simeconazole, Picoxystrobin, Pyraclostrobin, Dimoxystrobin, Prothioconazole, Fluoxastrobin; Cereals Insecticides: Dimethoate, Lambda-cyhalothrin, Deltamethrin, alpha-Cypermethrin, beta-cyfluthrin, Bifenthrin, Imidacloprid, Clothianidin, Thiamethoxam, Thiacloprid, Acetamiprid, Dinotefuran, Clorphyriphos, Metamidophos, Oxidemethon-methyl, Pirimicarb, Methiocarb; Maize Herbicides: Atrazine, Alachlor, Bromoxynil, Acetochlor, Dicamba, Clopyralid, (S-) Dimethenamid, Glufosinate, Glyphosate, Isoxaflutole, (S-)Metolachlor, Mesotrione, Nicosulfuron, Primisulfuron, Rimsulfuron, Sulcotrione, Foramsulfuron, Topramezone,

Tembotrione, Saflufenacil, Thien carbazon, Flufenacet, Pyroxasulfon; Maize Insecticides:
 Carbofuran, Chlorpyrifos, Bifenthrin, Fipronil, Imidacloprid, Lambda-Cyhalothrin,
 Tefluthrin, Terbufos, Thiamethoxam, Clothianidin, Spiromesifen, Flubendiamide,
 Triflumuron, Rynaxypyr, Deltamethrin, Thiodicarb, β -Cyfluthrin, Cypermethrin, Bifenthrin,
 5 Lufenuron, Triflumoron, Tefluthrin, Tebupirimphos, Ethiprole, Cyazypyr, Thiacloprid,
 Acetamiprid, Dinotofuran, Avermectin, Methiocarb, Spirodiclofen, Spirotetramat; Maize
Fungicides: Fenitropan, Thiram, Prothioconazole, Tebuconazole, Trifloxystrobin; Rice
Herbicides: Butachlor, Propanil, Azimsulfuron, Bensulfuron, Cyhalofop, Daimuron,
 Fentrazamide, Imazosulfuron, Mefenacet, Oxaziclomefone, Pyrazosulfuron, Pyributicarb,
 10 Quinclorac, Thiobencarb, Indanofan, Flufenacet, Fentrazamide, Halosulfuron,
 Oxaziclomefone, Benzobicyclon, Pyriftalid, Penoxsulam, Bispyribac, Oxadiargyl,
 Ethoxysulfuron, Pretilachlor, Mesotrione, Tefuryltrione, Oxadiazon, Fenoxaprop,
 Pyrimisulfan; Rice Insecticides: Diazinon, Fenitrothion, Fenobucarb, Monocrotophos,
 Benfuracarb, Buprofezin, Dinotefuran, Fipronil, Imidacloprid, Isoprocarb, Thiacloprid,
 15 Chromafenozide, Thiacloprid, Dinotefuran, Clothianidin, Ethiprole, Flubendiamide,
 Rynaxypyr, Deltamethrin, Acetamiprid, Thiamethoxam, Cyazypyr, Spinosad, Spinotoram,
 Emamectin-Benzoate, Cypermethrin, Chlorpyrifos, Cartap, Methamidophos, Etofenprox,
 Triazophos, 4-[[[(6-Chlorpyridin-3-yl)methyl](2,2-difluorethyl)amino]furan-2(5H)-on,
 Carbofuran, Benfuracarb; Rice Fungicides: Thiophanate-methyl, Azoxystrobin,
 20 Carpropamid, Edifenphos, Ferimzone, Iprobenfos, Isoprothiolane, Pencycuron,
 Probenazole, Pyroquilon, Tricyclazole, Trifloxystrobin, Diclocymet, Fenoxanil,
 Simeconazole, Tiadinil; Cotton Herbicides: Diuron, Fluometuron, MSMA, Oxyfluorfen,
 Prometryn, Trifluralin, Carfentrazone, Clethodim, Fluazifop-butyl, Glyphosate,
 Norflurazon, Pendimethalin, Pyriothobac-sodium, Trifloxysulfuron, Tepraloxymid,
 25 Glufosinate, Flumioxazin, Thidiazuron; Cotton Insecticides: Acephate, Aldicarb,
 Chlorpyrifos, Cypermethrin, Deltamethrin, Malathion, Monocrotophos, Abamectin,
 Acetamiprid, Emamectin Benzoate, Imidacloprid, Indoxacarb, Lambda-Cyhalothrin,
 Spinosad, Thiodicarb, Gamma-Cyhalothrin, Spiromesifen, Pyridalyl, Flonicamid,
 Flubendiamide, Triflumuron, Rynaxypyr, Beta-Cyfluthrin, Spirotetramat, Clothianidin,
 30 Thiamethoxam, Thiacloprid, Dinotofuran, Flubendiamide, Cyazypyr, Spinosad,
 Spinotoram, gamma Cyhalothrin, 4-[[[(6-Chlorpyridin-3-yl)methyl](2,2-
 difluorethyl)amino]furan-2(5H)-on, Thiodicarb, Avermectin, Flonicamid, Pyridalyl,
 Spiromesifen, Sulfoxaflor, Profenophos, Thiazophos, Endosulfan; Cotton Fungicides:
 Etridiazole, Metalaxyl, Quintozene; Soybean Herbicides: Alachlor, Bentazone, Trifluralin,
 35 Chlorimuron-Ethyl, Cloransulam-Methyl, Fenoxaprop, Fomesafen, Fluazifop, Glyphosate,
 Imazamox, Imazaquin, Imazethapyr, (S-)Metolachlor, Metribuzin, Pendimethalin,

Tepraloxydim, Glufosinate; Soybean Insecticides: Lambda-cyhalothrin, Methomyl, Parathion, Thiocarb, Imidacloprid, Clothianidin, Thiamethoxam, Thiacloprid, Acetamiprid, Dinetofuran, Flubendiamide, Rynaxypyr, Cyazypyr, Spinosad, Spinotoram, Emamectin-Benzoate, Fipronil, Ethiprole, Deltamethrin, β -Cyfluthrin, gamma and lambda Cyhalothrin,
 5 4-[[[(6-Chlorpyridin-3-yl)methyl](2,2-difluorethyl)amino]furan-2(5H)-on, Spirotetramat, Spinodiclofen, Triflumuron, Flonicamid, Thiodicarb, beta-Cyfluthrin; Soybean Fungicides: Azoxystrobin, Cyproconazole, Epoxiconazole, Flutriafol, Pyraclostrobin, Tebuconazole, Trifloxystrobin, Prothioconazole, Tetraconazole; Sugarbeet Herbicides: Chloridazon, Desmedipham, Ethofumesate, Phenmedipham, Triallate, Clopyralid, Fluazifop, Lenacil,
 10 Metamitron, Quinmerac, Cycloxydim, Triflurosulfuron, Tepraloxydim, Quizalofop; Sugarbeet Insecticides: Imidacloprid, Clothianidin, Thiamethoxam, Thiacloprid, Acetamiprid, Dinetofuran, Deltamethrin, β -Cyfluthrin, gamma/lambda Cyhalothrin, 4-[[[(6-Chlorpyridin-3-yl)methyl](2,2-difluorethyl)amino]furan-2(5H)-on, Tefluthrin, Rynaxypyr, Cyaxypyr, Fipronil, Carbofuran; Canola Herbicides: Clopyralid, Diclofop, Fluazifop, Glufosinate,
 15 Glyphosate, Metazachlor, Trifluralin Ethametsulfuron, Quinmerac, Quizalofop, Clethodim, Tepraloxydim; Canola Fungicides: Azoxystrobin, Carbendazim, Fludioxonil, Iprodione, Prochloraz, Vinclozolin; Canola Insecticides: Carbofuran, Organophosphates, Pyrethroids, Thiacloprid, Deltamethrin, Imidacloprid, Clothianidin, Thiamethoxam, Acetamiprid, Dinetofuran, β -Cyfluthrin, gamma and lambda Cyhalothrin, tau-Fluvalerate, Ethiprole,
 20 Spinosad, Spinotoram, Flubendiamide, Rynaxypyr, Cyazypyr, 4-[[[(6-Chlorpyridin-3-yl)methyl](2,2-difluorethyl)amino]furan-2(5H)-on.

In some embodiments the herbicide is Atrazine, Bromacil, Diuron, Chlorsulfuron, Metsulfuron, Thifensulfuron Methyl, Tribenuron, Acetochlor, Dicamba, Isoxaflutole, Nicosulfuron, Rimsulfuron, Pyriithiobac-sodium, Flumioxazin, Chlorimuron-Ethyl,
 25 Metribuzin, Quizalofop, S-metolachlor, Hexazinne or combinations thereof.

In some embodiments the insecticide is Esfenvalerate, Chlorantraniliprole, Methomyl, Indoxacarb, Oxamyl or combinations thereof.

Pesticidal and insecticidal activity

30 "Pest" includes but is not limited to, insects, fungi, bacteria, nematodes, mites, ticks, and the like. Insect pests include insects selected from the orders Coleoptera, Diptera, Hymenoptera, Lepidoptera, Mallophaga, Homoptera, Hemiptera, Orthoptera, Thysanoptera, Dermaptera, Isoptera, Anoplura, Siphonaptera, Trichoptera, etc., particularly Lepidoptera, and Hemiptera.

35 Those skilled in the art will recognize that not all compounds are equally effective against all pests. Compounds of the embodiments display activity against insect pests,

which may include economically important agronomic, forest, greenhouse, nursery, ornamentals, food and fiber, public and animal health, domestic and commercial structure, household and stored product pests.

Larvae of the order Lepidoptera include, but are not limited to, armyworms, cutworms, loopers, and heliothines in the family Noctuidae *Spodoptera frugiperda* JE Smith (fall armyworm); *S. exigua* Hübner (beet armyworm); *S. litura* Fabricius (tobacco cutworm, cluster caterpillar); *Mamestra configurata* Walker (bertha armyworm); *M. brassicae* Linnaeus (cabbage moth); *Agrotis ipsilon* Hufnagel (black cutworm); *A. orthogonia* Morrison (western cutworm); *A. subterranea* Fabricius (granulate cutworm); *Alabama argillacea* Hübner (cotton leaf worm); *Trichoplusia ni* Hübner (cabbage looper); *Pseudoplusia includens* Walker (soybean looper); *Anticarsia gemmatilis* Hübner (velvetbean caterpillar); *Hypena scabra* Fabricius (green cloverworm); *Heliothis virescens* Fabricius (tobacco budworm); *Pseudaletia unipuncta* Haworth (armyworm); *Athetis mindara* Barnes and McDunnough (rough skinned cutworm); *Euxoa messoria* Harris (darksided cutworm); *Earias insulana* Boisduval (spiny bollworm); *E. vittella* Fabricius (spotted bollworm); *Helicoverpa armigera* Hübner (American bollworm); *H. zea* Boddie (corn earworm or cotton bollworm); *Melanchra picta* Harris (zebra caterpillar); *Egira (Xylomyges) curialis* Grote (citrus cutworm); borers, casebearers, webworms, coneworms, and skeletonizers from the family Pyralidae *Ostrinia nubilalis* Hübner (European corn borer); *Amyelois transitella* Walker (naval orangeworm); *Anagasta kuehniella* Zeller (Mediterranean flour moth); *Cadra cautella* Walker (almond moth); *Chilo suppressalis* Walker (rice stem borer); *C. partellus*, (sorghum borer); *Corcyra cephalonica* Stainton (rice moth); *Crambus caliginosellus* Clemens (corn root webworm); *C. teterrellus* Zincken (bluegrass webworm); *Cnaphalocrocis medinalis* Guenée (rice leaf roller); *Desmia funeralis* Hübner (grape leaffolder); *Diaphania hyalinata* Linnaeus (melon worm); *D. nitidalis* Stoll (pickleworm); *Diatraea grandiosella* Dyar (southwestern corn borer), *D. saccharalis* Fabricius (surgarcane borer); *Eoreuma loftini* Dyar (Mexican rice borer); *Ephestia elutella* Hübner (tobacco (cacao) moth); *Galleria mellonella* Linnaeus (greater wax moth); *Herpetogramma licarsisalis* Walker (sod webworm); *Homoeosoma electellum* Hulst (sunflower moth); *Elasmopalpus lignosellus* Zeller (lesser cornstalk borer); *Achroia grisella* Fabricius (lesser wax moth); *Loxostege sticticalis* Linnaeus (beet webworm); *Orthaga thyrisalis* Walker (tea tree web moth); *Maruca testulalis* Geyer (bean pod borer); *Plodia interpunctella* Hübner (Indian meal moth); *Scirpophaga incertulas* Walker (yellow stem borer); *Udea rubigalis* Guenée (celery leaftier); and leafrollers, budworms, seed worms, and fruit worms in the family Tortricidae *Acleris gloverana* Walsingham (Western blackheaded budworm); *A. variana* Fernald (Eastern blackheaded budworm); *Archips*

argyrospila Walker (fruit tree leaf roller); *A. rosana* Linnaeus (European leaf roller); and other *Archips* species, *Adoxophyes orana* Fischer von Rösslerstamm (summer fruit tortrix moth); *Cochylis hospes* Walsingham (banded sunflower moth); *Cydia latiferreana* Walsingham (filbertworm); *C. pomonella* Linnaeus (coding moth); *Platynota flavedana* Clemens (variegated leafroller); *P. stultana* Walsingham (omnivorous leafroller); *Lobesia botrana* Denis & Schiffermüller (European grape vine moth); *Spilonota ocellana* Denis & Schiffermüller (eyespotted bud moth); *Endopiza viteana* Clemens (grape berry moth); *Eupoecilia ambiguella* Hübner (vine moth); *Bonagota salubricola* Meyrick (Brazilian apple leafroller); *Grapholita molesta* Busck (oriental fruit moth); *Suleima helianthana* Riley (sunflower bud moth); *Argyrotaenia* spp.; *Choristoneura* spp..

Selected other agronomic pests in the order Lepidoptera include, but are not limited to, *Alsophila pometaria* Harris (fall cankerworm); *Anarsia lineatella* Zeller (peach twig borer); *Anisota senatoria* J.E. Smith (orange striped oakworm); *Antheraea pernyi* Guérin-Méneville (Chinese Oak Tussock Moth); *Bombyx mori* Linnaeus (Silkworm); *Bucculatrix thurberiella* Busck (cotton leaf perforator); *Colias eurytheme* Boisduval (alfalfa caterpillar); *Datana integerrima* Grote & Robinson (walnut caterpillar); *Dendrolimus sibiricus* Tschetwerikov (Siberian silk moth), *Ennomos subsignaria* Hübner (elm spanworm); *Erannis tiliaria* Harris (linden looper); *Euproctis chrysorrhoea* Linnaeus (browntail moth); *Harrisina americana* Guérin-Méneville (grapeleaf skeletonizer); *Hemileuca oliviae* Cockrell (range caterpillar); *Hyphantria cunea* Drury (fall webworm); *Keiferia lycopersicella* Walsingham (tomato pinworm); *Lambdina fiscellaria fiscellaria* Hulst (Eastern hemlock looper); *L. fiscellaria lugubrosa* Hulst (Western hemlock looper); *Leucoma salicis* Linnaeus (satin moth); *Lymantria dispar* Linnaeus (gypsy moth); *Manduca quinquemaculata* Haworth (five spotted hawk moth, tomato hornworm); *M. sexta* Haworth (tomato hornworm, tobacco hornworm); *Operophtera brumata* Linnaeus (winter moth); *Paleacrita vernata* Peck (spring cankerworm); *Papilio cresphontes* Cramer (giant swallowtail, orange dog); *Phryganidia californica* Packard (California oakworm); *Phyllocnistis citrella* Stainton (citrus leafminer); *Phyllonorycter blancardella* Fabricius (spotted tentiform leafminer); *Pieris brassicae* Linnaeus (large white butterfly); *P. rapae* Linnaeus (small white butterfly); *P. napi* Linnaeus (green veined white butterfly); *Platyptilia carduidactyla* Riley (artichoke plume moth); *Plutella xylostella* Linnaeus (diamondback moth); *Pectinophora gossypiella* Saunders (pink bollworm); *Pontia protodice* Boisduval & Leconte (Southern cabbageworm); *Sabulodes aegrotata* Guenée (omnivorous looper); *Schizura concinna* J.E. Smith (red humped caterpillar); *Sitotroga cerealella* Olivier (Angoumois grain moth); *Thaumetopoea pityocampa* Schiffermüller (pine processionary caterpillar); *Tineola bisselliella* Hummel (webbing clothesmoth); *Tuta*

absoluta Meyrick (tomato leafminer); *Yponomeuta padella* Linnaeus (ermine moth); *Heliothis subflexa* Guenée; *Malacosoma* spp. and *Orgyia* spp.

Of interest are larvae and adults of the order Coleoptera including weevils from the families Anthribidae, Bruchidae, and Curculionidae (including, but not limited to:

5 *Anthonomus grandis* Boheman (boll weevil); *Lissorhoptrus oryzophilus* Kuschel (rice water weevil); *Sitophilus granarius* Linnaeus (granary weevil); *S. oryzae* Linnaeus (rice weevil); *Hypera punctata* Fabricius (clover leaf weevil); *Cylindrocopturus adspersus* LeConte (sunflower stem weevil); *Smicronyx fulvus* LeConte (red sunflower seed weevil); *S. sordidus* LeConte (gray sunflower seed weevil); *Sphenophorus maidis* Chittenden

10 (maize billbug)); flea beetles, cucumber beetles, rootworms, leaf beetles, potato beetles, and leafminers in the family Chrysomelidae (including, but not limited to: *Leptinotarsa decemlineata* Say (Colorado potato beetle); *Diabrotica virgifera virgifera* LeConte (western corn rootworm); *D. barberi* Smith & Lawrence (northern corn rootworm); *D. undecimpunctata howardi* Barber (southern corn rootworm); *Chaetocnema pulicaria*

15 Melsheimer (corn flea beetle); *Phyllotreta cruciferae* Goeze (Crucifer flea beetle); *Phyllotreta striolata* (stripped flea beetle); *Colaspis brunnea* Fabricius (grape colaspis); *Oulema melanopus* Linnaeus (cereal leaf beetle); *Zygogramma exclamationis* Fabricius (sunflower beetle)); beetles from the family Coccinellidae (including, but not limited to: *Epilachna varivestis* Mulsant (Mexican bean beetle)); chafers and other beetles from the

20 family Scarabaeidae (including, but not limited to: *Popillia japonica* Newman (Japanese beetle); *Cyclocephala borealis* Arrow (northern masked chafer, white grub); *C. immaculata* Olivier (southern masked chafer, white grub); *Rhizotrogus majalis* Razoumowsky (European chafer); *Phyllophaga crinita* Burmeister (white grub); *Ligyrus gibbosus* De Geer (carrot beetle)); carpet beetles from the family Dermestidae; wireworms

25 from the family Elateridae, *Eleodes* spp., *Melanotus* spp.; *Conoderus* spp.; *Limonius* spp.; *Agriotes* spp.; *Ctenicera* spp.; *Aeolus* spp.; bark beetles from the family Scolytidae and beetles from the family Tenebrionidae.

Adults and immatures of the order Diptera are of interest, including leafminers *Agromyza parvicornis* Loew (corn blotch leafminer); midges (including, but not limited to:

30 *Contarinia sorghicola* Coquillett (sorghum midge); *Mayetiola destructor* Say (Hessian fly); *Sitodiplosis mosellana* Géhin (wheat midge); *Neolasioptera murtfeldtiana* Felt, (sunflower seed midge)); fruit flies (Tephritidae), *Oscinella frit* Linnaeus (fruit flies); maggots (including, but not limited to: *Delia platura* Meigen (seedcorn maggot); *D. coarctata* Fallen (wheat bulb fly); and other *Delia* spp., *Meromyza americana* Fitch (wheat stem maggot);

35 *Musca domestica* Linnaeus (house flies); *Fannia canicularis* Linnaeus, *F. femoralis* Stein (lesser house flies); *Stomoxys calcitrans* Linnaeus (stable flies)); face flies, horn flies,

blow flies, *Chrysomya* spp.; *Phormia* spp.; and other muscoid fly pests, horse flies *Tabanus* spp.; bot flies *Gastrophilus* spp.; *Oestrus* spp.; cattle grubs *Hypoderma* spp.; deer flies *Chrysops* spp.; *Melophagus ovinus* Linnaeus (keds); and other *Brachycera*, mosquitoes *Aedes* spp.; *Anopheles* spp.; *Culex* spp.; black flies *Prosimulium* spp.;
 5 *Simulium* spp.; biting midges, sand flies, sciarids, and other *Nematocera*.

Included as insects of interest are adults and nymphs of the orders Hemiptera and Homoptera such as, but not limited to, adelgids from the family Adelgidae, plant bugs from the family Miridae, cicadas from the family Cicadidae, leafhoppers, *Empoasca* spp.; from the family Cicadellidae, planthoppers from the families Cixiidae, Flatidae, Fulgoroidea,
 10 Issidae and Delphacidae, treehoppers from the family Membracidae, psyllids from the family Psyllidae, whiteflies from the family Aleyrodidae, aphids from the family Aphididae, phylloxera from the family Phylloxeridae, mealybugs from the family Pseudococcidae, scales from the families Asterolecanidae, Coccidae, Dactylopiidae, Diaspididae, Eriococcidae, Ortheziidae, Phoenicococcidae and Margarodidae, lace bugs from the
 15 family Tingidae, stink bugs from the family Pentatomidae, cinch bugs, *Blissus* spp.; and other seed bugs from the family Lygaeidae, spittlebugs from the family Cercopidae squash bugs from the family Coreidae, and red bugs and cotton stainers from the family Pyrrhocoridae.

Agronomically important members from the order Homoptera further include, but
 20 are not limited to: *Acyrtosiphon pisum* Harris (pea aphid); *Aphis craccivora* Koch (cowpea aphid); *A. fabae* Scopoli (black bean aphid); *A. gossypii* Glover (cotton aphid, melon aphid); *A. maidiradicis* Forbes (corn root aphid); *A. pomi* De Geer (apple aphid); *A. spiraecola* Patch (spirea aphid); *Aulacorthum solani* Kaltenbach (foxglove aphid); *Chaetosiphon fragaefolii* Cockerell (strawberry aphid); *Diuraphis noxia*
 25 Kurdjumov/Mordvilko (Russian wheat aphid); *Dysaphis plantaginea* Paaserini (rosy apple aphid); *Eriosoma lanigerum* Hausmann (woolly apple aphid); *Brevicoryne brassicae* Linnaeus (cabbage aphid); *Hyalopterus pruni* Geoffroy (mealy plum aphid); *Lipaphis erysimi* Kaltenbach (turnip aphid); *Metopolophium dirrhodum* Walker (cereal aphid); *Macrosiphum euphorbiae* Thomas (potato aphid); *Myzus persicae* Sulzer (peach-potato
 30 aphid, green peach aphid); *Nasonovia ribisnigri* Mosley (lettuce aphid); *Pemphigus* spp. (root aphids and gall aphids); *Rhopalosiphum maidis* Fitch (corn leaf aphid); *R. padi* Linnaeus (bird cherry-oat aphid); *Schizaphis graminum* Rondani (greenbug); *Sipha flava* Forbes (yellow sugarcane aphid); *Sitobion avenae* Fabricius (English grain aphid); *Therioaphis maculata* Buckton (spotted alfalfa aphid); *Toxoptera aurantii* Boyer de
 35 Fonscolombe (black citrus aphid); and *T. citricida* Kirkaldy (brown citrus aphid); *Adelges* spp. (adelgids); *Phylloxera devastatrix* Pergande (pecan phylloxera); *Bemisia tabaci*

Gennadius (tobacco whitefly, sweetpotato whitefly); *B. argentifolii* Bellows & Perring (silverleaf whitefly); *Dialeurodes citri* Ashmead (citrus whitefly); *Trialeurodes abutiloneus* (bandedwinged whitefly) and *T. vaporariorum* Westwood (greenhouse whitefly); *Empoasca fabae* Harris (potato leafhopper); *Laodelphax striatellus* Fallen (smaller brown planthopper); *Macrolestes quadrilineatus* Forbes (aster leafhopper); *Nephotettix cincticeps* Uhler (green leafhopper); *N. nigropictus* Stål (rice leafhopper); *Nilaparvata lugens* Stål (brown planthopper); *Peregrinus maidis* Ashmead (corn planthopper); *Sogatella furcifera* Horvath (white-backed planthopper); *Sogatodes orizicola* Muir (rice delphacid); *Typhlocyba pomaria* McAtee (white apple leafhopper); *Erythroneoura* spp. (grape leafhoppers); *Magicicada septendecim* Linnaeus (periodical cicada); *Icerya purchasi* Maskell (cottony cushion scale); *Quadraspidiotus perniciosus* Comstock (San Jose scale); *Planococcus citri* Risso (citrus mealybug); *Pseudococcus* spp. (other mealybug complex); *Cacopsylla pyricola* Foerster (pear psylla); *Trioza diospyri* Ashmead (persimmon psylla).

Agronomically important species of interest from the order Hemiptera include, but are not limited to: *Acrosternum hilare* Say (green stink bug); *Anasa tristis* De Geer (squash bug); *Blissus leucopterus leucopterus* Say (chinch bug); *Corythuca gossypii* Fabricius (cotton lace bug); *Cyrtopeltis modesta* Distant (tomato bug); *Dysdercus suturellus* Herrich-Schäffer (cotton stainer); *Euschistus servus* Say (brown stink bug); *E. variolarius* Palisot de Beauvois (one-spotted stink bug); *Graptostethus* spp. (complex of seed bugs); *Leptoglossus corculus* Say (leaf-footed pine seed bug); *Lygus lineolaris* Palisot de Beauvois (tarnished plant bug); *L. hesperus* Knight (Western tarnished plant bug); *L. pratensis* Linnaeus (common meadow bug); *L. rugulipennis* Poppius (European tarnished plant bug); *Lygocoris pabulinus* Linnaeus (common green capsid); *Nezara viridula* Linnaeus (southern green stink bug); *Oebalus pugnax* Fabricius (rice stink bug); *Oncopeltus fasciatus* Dallas (large milkweed bug); *Pseudatomoscelis seriatus* Reuter (cotton fleahopper).

Furthermore, embodiments may be effective against Hemiptera such, *Calocoris norvegicus* Gmelin (strawberry bug); *Orthops campestris* Linnaeus; *Plesiocoris rugicollis* Fallen (apple capsid); *Cyrtopeltis modestus* Distant (tomato bug); *Cyrtopeltis notatus* Distant (suckfly); *Spanagonicus albofasciatus* Reuter (whitemarked fleahopper); *Diaphnocoris chlorionis* Say (honeylocust plant bug); *Labopidicola allii* Knight (onion plant bug); *Pseudatomoscelis seriatus* Reuter (cotton fleahopper); *Adelphocoris rapidus* Say (rapid plant bug); *Poecilocapsus lineatus* Fabricius (four-lined plant bug); *Nysius ericae* Schilling (false chinch bug); *Nysius raphanus* Howard (false chinch bug); *Nezara viridula* Linnaeus (Southern green stink bug); *Eurygaster* spp.; *Coreidae* spp.; *Pyrrhocoridae* spp.; *Tinidae* spp.; *Blotomatidae* spp.; *Reduviidae* spp.; and *Cimicidae* spp.

Also included are adults and larvae of the order Acari (mites) such as *Aceria tosichella* Keifer (wheat curl mite); *Petrobia latens* Müller (brown wheat mite); spider mites and red mites in the family Tetranychidae, *Panonychus ulmi* Koch (European red mite); *Tetranychus urticae* Koch (two spotted spider mite); (*T. mcdanieli* McGregor (McDaniel mite); *T. cinnabarinus* Boisduval (carmine spider mite); *T. turkestan* Ugarov & Nikolski (strawberry spider mite); flat mites in the family Tenuipalpidae, *Brevipalpus lewisi* McGregor (citrus flat mite); rust and bud mites in the family Eriophyidae and other foliar feeding mites and mites important in human and animal health, i.e. dust mites in the family Epidermoptidae, follicle mites in the family Demodicidae, grain mites in the family Glycyphagidae, ticks in the order Ixodidae. *Ixodes scapularis* Say (deer tick); *I. holocyclus* Neumann (Australian paralysis tick); *Dermacentor variabilis* Say (American dog tick); *Amblyomma americanum* Linnaeus (lone star tick); and scab and itch mites in the families Psoroptidae, Pyemotidae, and Sarcoptidae.

Insect pests of the order Thysanura are of interest, such as *Lepisma saccharina* Linnaeus (silverfish); *Thermobia domestica* Packard (firebrat).

Additional arthropod pests covered include: spiders in the order Araneae such as *Loxosceles reclusa* Gertsch & Mulaik (brown recluse spider); and the *Latrodectus mactans* Fabricius (black widow spider); and centipedes in the order Scutigleromorpha such as *Scutigera coleoptrata* Linnaeus (house centipede).

Insect pest of interest include the superfamily of stink bugs and other related insects including but not limited to species belonging to the family Pentatomidae (*Nezara viridula*, *Halyomorpha halys*, *Piezodorus guildini*, *Euschistus servus*, *Acrosternum hilare*, *Euschistus heros*, *Euschistus tristigmus*, *Acrosternum hilare*, *Dichelops furcatus*, *Dichelops melacanthus*, and *Bagrada hilaris* (Bagrada Bug)), the family Plataspididae (*Megacopta cribraria* - Bean plataspid), and the family Cydnidae (*Scaptocoris castanea* - Root stink bug); and Lepidoptera species including but not limited to: diamond-back moth, e.g., *Helicoverpa zea* Boddie; soybean looper, e.g., *Pseudoplusia includens* Walker; and velvet bean caterpillar e.g., *Anticarsia gemmatilis* Hübner.

Methods for measuring pesticidal activity are well known in the art. See, for example, Czapla and Lang, (1990) *J. Econ. Entomol.* 83:2480-2485; Andrews, et al., (1988) *Biochem. J.* 252:199-206; Marrone, et al., (1985) *J. of Economic Entomology* 78:290-293 and US Patent Number 5,743,477, all of which are herein incorporated by reference in their entirety. Generally, the protein is mixed and used in feeding assays. See, for example Marrone, et al., (1985) *J. of Economic Entomology* 78:290-293. Such assays can include contacting plants with one or more pests and determining the plant's ability to survive and/or cause the death of the pests.

Nematodes include parasitic nematodes such as root-knot, cyst, and lesion nematodes, including *Heterodera* spp., *Meloidogyne* spp., and *Globodera* spp.; particularly members of the cyst nematodes, including, but not limited to, *Heterodera glycines* (soybean cyst nematode); *Heterodera schachtii* (beet cyst nematode);
 5 *Heterodera avenae* (cereal cyst nematode); and *Globodera rostochiensis* and *Globodera pailida* (potato cyst nematodes). Lesion nematodes include *Pratylenchus* spp.

Seed Treatment

To protect and to enhance yield production and trait technologies, seed treatment
 10 options can provide additional crop plan flexibility and cost effective control against insects, weeds and diseases. Seed material can be treated, typically surface treated, with a composition comprising combinations of chemical or biological herbicides, herbicide safeners, insecticides, fungicides, germination inhibitors and enhancers, nutrients, plant growth regulators and activators, bactericides, nematocides, avicides and/or
 15 molluscicides. These compounds are typically formulated together with further carriers, surfactants or application-promoting adjuvants customarily employed in the art of formulation. The coatings may be applied by impregnating propagation material with a liquid formulation or by coating with a combined wet or dry formulation. Examples of the various types of compounds that may be used as seed treatments are provided in The
 20 Pesticide Manual: A World Compendium, C.D.S. Tomlin Ed., Published by the British Crop Production Council, which is hereby incorporated by reference.

Some seed treatments that may be used on crop seed include, but are not limited to, one or more of abscisic acid, acibenzolar-S-methyl, avermectin, amitrol, azaconazole, azospirillum, azadirachtin, azoxystrobin, *Bacillus* spp. (including one or more of *cereus*,
 25 *firmus*, *megaterium*, *pumilis*, *sphaericus*, *subtilis* and/or *thuringiensis* species), *bradyrhizobium* spp. (including one or more of *betae*, *canariense*, *elkanii*, *iriomotense*, *japonicum*, *liaonigense*, *pachyrhizi* and/or *yuanmingense*), captan, carboxin, chitosan, clothianidin, copper, cyazypyr, difenoconazole, etidiazole, fipronil, fludioxonil, fluoxastrobin, fluquinconazole, flurazole, fluxofenim, harpin protein, imazalil, imidacloprid,
 30 ipconazole, isoflavenoids, lipo-chitooligosaccharide, mancozeb, manganese, maneb, mefenoxam, metalaxyl, metconazole, myclobutanil, PCNB, penflufen, penicillium, penthiopyrad, permethrine, picoxystrobin, prothioconazole, pyraclostrobin, rynaxypyr, S-metolachlor, saponin, sedaxane, TCMTB, tebuconazole, thiabendazole, thiamethoxam, thiocarb, thiram, tolclofos-methyl, triadimenol, trichoderma, trifloxystrobin, triticonazole
 35 and/or zinc. PCNB seed coat refers to EPA registration number 00293500419, containing quintozen and terrazole. TCMTB refers to 2-(thiocyanomethylthio) benzothiazole.

Seed varieties and seeds with specific transgenic traits may be tested to determine which seed treatment options and application rates may complement such varieties and transgenic traits in order to enhance yield. For example, a variety with good yield potential but head smut susceptibility may benefit from the use of a seed treatment
5 that provides protection against head smut, a variety with good yield potential but cyst nematode susceptibility may benefit from the use of a seed treatment that provides protection against cyst nematode, and so on. Likewise, a variety encompassing a transgenic trait conferring insect resistance may benefit from the second mode of action conferred by the seed treatment, a variety encompassing a transgenic trait conferring
10 herbicide resistance may benefit from a seed treatment with a safener that enhances the plants resistance to that herbicide, etc. Further, the good root establishment and early emergence that results from the proper use of a seed treatment may result in more efficient nitrogen use, a better ability to withstand drought and an overall increase in yield potential of a variety or varieties containing a certain trait when combined with a seed
15 treatment.

Methods for inhibiting growth or killing an insect pest and controlling an insect population

In some embodiments methods are provided for inhibiting growth or killing an
20 insect pest, comprising contacting the insect pest with an insecticidally-effective amount of a recombinant PHI-4 polypeptide. In some embodiments methods are provided for inhibiting growth or killing an insect pest, comprising contacting the insect pest with an insecticidally-effective amount of a recombinant pesticidal protein of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526 or a
25 variant thereof.

In some embodiments methods are provided for controlling an insect pest population, comprising contacting the insect pest population with an insecticidally-effective amount of a recombinant PHI-4 polypeptide. In some embodiments methods are provided for controlling an insect pest population, comprising contacting the insect pest
30 population with an insecticidally-effective amount of a recombinant pesticidal protein of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526 or a variant thereof. As used herein, by “controlling a pest population” or “controls a pest” is intended any effect on a pest that results in limiting the damage that the pest causes. Controlling a pest includes, but is not limited to, killing the pest,
35 inhibiting development of the pest, altering fertility or growth of the pest in such a manner that the pest provides less damage to the plant, decreasing the number of offspring

produced, producing less fit pests, producing pests more susceptible to predator attack or deterring the pests from eating the plant.

In some embodiments methods are provided for controlling an insect pest population resistant to a pesticidal protein, comprising contacting the insect pest population with an insecticidally-effective amount of a recombinant PHI-4 polypeptide. In some embodiments methods are provided for controlling an insect pest population resistant to a pesticidal protein, comprising contacting the insect pest population with an insecticidally-effective amount of a recombinant pesticidal protein of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526 or a variant thereof.

In some embodiments methods are provided for protecting a plant from an insect pest, comprising expressing in the plant or cell thereof a recombinant PHI-4 polypeptide. In some embodiments methods are provided for protecting a plant from an insect pest, comprising expressing in the plant or cell thereof a recombinant pesticidal protein of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526 or variants thereof.

Insect Resistance Management (IRM) Strategies

Expression of *B. thuringiensis* δ -endotoxins in transgenic corn plants has proven to be an effective means of controlling agriculturally important insect pests (Perlak, *et al.*, 1990; 1993). However, insects have evolved that are resistant to *B. thuringiensis* δ -endotoxins expressed in transgenic plants. Such resistance, should it become widespread, would clearly limit the commercial value of germplasm containing genes encoding such *B. thuringiensis* δ -endotoxins.

One way to increasing the effectiveness of the transgenic insecticides against target pests and contemporaneously reducing the development of insecticide-resistant pests is to use provide non-transgenic (i.e., non-insecticidal protein) refuges (a section of non-insecticidal crops/ corn) for use with transgenic crops producing a single insecticidal protein active against target pests. The United States Environmental Protection Agency (epa.gov/oppbppdl/biopesticides/pips/bt_corn_refuge_2006.htm, which can be accessed using the www prefix) publishes the requirements for use with transgenic crops producing a single Bt protein active against target pests. In addition, the National Corn Growers Association, on their website: (ncga.com/insect-resistance-management-fact-sheet-bt-corn, which can be accessed using the www prefix) also provides similar guidance regarding refuge requirements. Due to losses to insects within the refuge area, larger refuges may reduce overall yield.

Another way of increasing the effectiveness of the transgenic insecticides against target pests and contemporaneously reducing the development of insecticide-resistant pests would be to have a repository of insecticidal genes that are effective against groups of insect pests and which manifest their effects through different modes of action.

5 Expression in a plant of two or more insecticidal compositions toxic to the same insect species, each insecticide being expressed at efficacious levels would be another way to achieve control of the development of resistance. This is based on the principle that evolution of resistance against two separate modes of action is far more unlikely than only one. Roush for example, outlines two-toxin strategies, also called "pyramiding" or
10 "stacking," for management of insecticidal transgenic crops. (The Royal Society. Phil. Trans. R. Soc. Lond. B. (1998) 353:777-1786). Stacking or pyramiding of two different proteins each effective against the target pests and with little or no cross-resistance can allow for use of a smaller refuge. The U.S. Environmental Protection Agency requires significantly less (generally 5%) structured refuge of non-Bt corn be planted than for
15 single trait products (generally 20%). There are various ways of providing the IRM effects of a refuge, including various geometric planting patterns in the fields and in-bag seed mixtures, as discussed further by Roush.

In some embodiments the PHI-4 polypeptides of the disclosure are useful as an insect resistance management strategy in combination (i.e., pyramided) with other
20 pesticidal proteins include but are not limited to Bt toxins, *Xenorhabdus* sp. or *Photorhabdus* sp. insecticidal proteins, and the like.

Provided are methods of controlling Coleoptera insect infestation(s) in a transgenic plant that promote insect resistance management, comprising expressing in the plant at least two different insecticidal proteins having different modes of action.

25 In some embodiments the methods of controlling Coleoptera insect infestation in a transgenic plant and promoting insect resistance management the at least one of the insecticidal proteins comprise a PHI-4 polypeptide insecticidal to insects in the order Coleoptera.

In some embodiments the methods of controlling Coleoptera insect infestation in a
30 transgenic plant and promoting insect resistance management the at least one of the insecticidal proteins comprises a protein of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526 or variants thereof, insecticidal to insects in the order Coleoptera.

In some embodiments the methods of controlling Coleoptera insect infestation in a
35 transgenic plant and promoting insect resistance management comprise expressing in the

transgenic plant a PHI-4 polypeptide and a cry protein insecticidal to insects in the order Coleoptera having different modes of action.

In some embodiments the methods of controlling Coleoptera insect infestation in a transgenic plant and promoting insect resistance management comprise in the transgenic
5 plant a protein of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526 or variants thereof and a cry protein insecticidal to insects in the order Coleoptera having different modes of action.

Also provided are methods of reducing likelihood of emergence of Coleoptera insect resistance to transgenic plants expressing in the plants insecticidal proteins to
10 control the insect species, comprising expression of a PHI-4 polypeptide insecticidal to the insect species in combination with a second insecticidal protein to the insect species having different modes of action.

Also provided are methods of reducing likelihood of emergence of Coleoptera insect resistance to transgenic plants expressing in the plants insecticidal proteins to
15 control the insect species, comprising expression of a protein of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526 or variants thereof, insecticidal to the insect species in combination with a second insecticidal protein to the insect species having different modes of action.

Also provided are means for effective Coleoptera insect resistance management
20 of transgenic plants, comprising co-expressing at high levels in the plants two or more insecticidal proteins toxic to Coleoptera insects but each exhibiting a different mode of effectuating its inhibiting growth or killing activity, wherein the two or more insecticidal proteins comprise a PHI-4 polypeptide and a cry protein. Also provided are means for effective Coleoptera insect resistance management of transgenic plants, comprising co-
25 expressing at high levels in the plants two or more insecticidal proteins toxic to Coleoptera insects but each exhibiting a different mode of effectuating its inhibiting growth or activity, wherein the two or more insecticidal proteins comprise a protein of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526 or variants thereof and a cry protein.

30 In addition, methods are provided for obtaining regulatory approval for planting or commercialization of plants expressing proteins insecticidal to insects in the order Coleoptera, comprising the step of referring to, submitting or relying on insect assay binding data showing that the PHI-4 polypeptide does not compete with binding sites for cry proteins in such insects. In addition, methods are provided for obtaining regulatory
35 approval for planting or commercialization of plants expressing proteins insecticidal to insects in the order Coleoptera, comprising the step of referring to, submitting or relying

on insect assay binding data showing that the protein of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NOs: 51-1162, and SEQ ID NOs: 1518-1526 or variant thereof does not compete with binding sites for cry proteins in such insects.

5 **Methods for Increasing Plant Yield**

Methods for increasing plant yield are provided. The methods comprise providing a plant or plant cell expressing a polynucleotide encoding the pesticidal polypeptide sequence disclosed herein and growing the plant or a seed thereof in a field infested with a pest against which the polypeptide has pesticidal activity. In some embodiments, the polypeptide has pesticidal activity against a lepidopteran, coleopteran, dipteran, hemipteran or nematode pest, and the field is infested with a lepidopteran, hemipteran, coleopteran, dipteran or nematode pest.

As defined herein, the “yield” of the plant refers to the quality and/or quantity of biomass produced by the plant. By “biomass” is intended any measured plant product. An increase in biomass production is any improvement in the yield of the measured plant product. Increasing plant yield has several commercial applications. For example, increasing plant leaf biomass may increase the yield of leafy vegetables for human or animal consumption. Additionally, increasing leaf biomass can be used to increase production of plant-derived pharmaceutical or industrial products. An increase in yield can comprise any statistically significant increase including, but not limited to, at least a 1% increase, at least a 3% increase, at least a 5% increase, at least a 10% increase, at least a 20% increase, at least a 30%, at least a 50%, at least a 70%, at least a 100% or a greater increase in yield compared to a plant not expressing the pesticidal sequence.

In specific methods, plant yield is increased as a result of improved pest resistance of a plant expressing a PHI-4 polypeptide disclosed herein. Expression of the PHI-4 polypeptide results in a reduced ability of a pest to infest or feed on the plant, thus improving plant yield.

These and other changes may be made to the disclosure in light of the above detailed description. In general, in the following claims, the terms used should not be construed to limit the disclosure to the specific embodiments disclosed in the specification and the claims.

The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, manuals, books or other disclosures) in the Background, Detailed Description, and Examples is herein incorporated by reference in their entireties.

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the subject disclosure, and are not intended to limit the scope of what is regarded as the invention. Efforts have been made to ensure accuracy with respect to the numbers used (e.g. amounts, temperature, concentrations, etc.) but some experimental errors and deviations should be allowed for. Unless otherwise indicated, parts are parts by weight, molecular weight is average molecular weight; temperature is in degrees centigrade; and pressure is at or near atmospheric.

10 EXPERIMENTALS

Example 1: Generating PHI-4 genes

Polynucleotides having single codon substitutions compared to the PHI-4 polypeptide of SEQ ID NO: 1 were generated. As described in the examples below, the corresponding PHI-4 polypeptides were expressed, purified and assayed for WCRW insecticidal activity in order to assess the corresponding activity diversity. A reverse mutagenesis primer and a complementary forward mutagenesis primer were designed to create the desired amino acid substitution(s) at the site(s) of interest. Typically the mutagenesis primer was between 30 to 45 bases in length with two or more bases, usually 10 to 15, on both sides of the site of interest. In the case of saturation mutagenesis, degenerated primers that cover all possible amino acid residues were used. Unless otherwise noted, the mutagenic reactions were carried out using Agilent's QuikChange™ Lightning Site-Directed Mutagenesis kit. Materials provided in the kit are QuikChange™ Lightning Enzyme, 10X QuikChange™ Lightning Buffer, dNTP mix, QuikSolution™ reagent and Dpn1 restriction enzyme according to the manufactures directions.

PCR amplifications were typically carried out with Expand™ High Fidelity PCR system (Roche, Switzerland) in 50 ul containing 50–100 ng templates, 0.4–2 µM primer pair, 200 µM dNTPs and 2 Units of DNA polymerase. The mutagenesis reaction was initiated by pre-heating the reaction mixture to 94°C for 3 min, followed by 16 cycles of the following cycling program: 94°C for 1 min, 52°C for 1 min and 68°C for 8, 12, 16 or 24 min according to the length of template. The mutagenesis reaction was completed by incubation at 68°C for 1 h. The PCR-amplification products were evaluated by agarose gel electrophoresis. The PCR products were purified by QIAquick™ PCR purification kit (Qiagen, Germany) and further treated with the restriction enzyme Dpn1. An aliquot of 1 µl of this PCR product was typically transformed into BL21(DE3) cells and inoculated on

Luria–Bertani (LB) plate containing 100 µg/ml ampicillin. About 10 colonies in the case of single amino acid mutation or 48 or more colonies for saturation mutagenesis were selected and plasmid DNA was isolated for sequencing. Two step sequencing was used, first for specific mutation site(s) with one sequencing primer followed by full length sequence confirmation with multiple sequencing primers.

Example 2: Purification of MBP::PHI-4 fusion polypeptides

Polynucleotides encoding PHI-4 polypeptides were expressed in a modified pMAL vector (New England Bio Lab) as a fusion (i.e. MBP::PHI-4; SEQ ID NO: 6) with MBP (maltose binding protein). The pMAL vector was modified to attach a 6X His tag to the N-terminus of MBP. In order to clone the polynucleotide encoding the MBP::PHI-4 fusion protein (SEQ ID NO: 6), Sph1 and BamH1 sites were engineered in the vector at the cloning site. The polynucleotide (SEQ ID NO: 1) encoding the PHI-4 polypeptide (SEQ ID NO: 2) was amplified with a forward primer (SEQ ID NO: 32) overlapping the Sph1 site and a reverse primer (SEQ ID NO: 33) overlapping the BamH1 site. This PCR product was digested with Sph1 and BamH1 and cloned into pMAL that was pre-cut with the same enzymes. The forward primer was designed such that polynucleotides encoding both MBP and PHI-4 polypeptide (SEQ ID NO: 2) within the MBP::PHI-4 gene (SEQ ID NO: 5) were ligated in frame. The plasmid containing the polynucleotide (SEQ ID NO: 5) encoding the MBP::PHI-4 polypeptide (SEQ ID NO: 6) was transformed into *E. coli* BL21(DE3) cells. The BL21(DE3) cells were grown in MagicMedia™ (Life Technologies) in either 96 deep well plates or flasks in a shaker running at 250 rpm at 37°C for 8 hrs. followed by 16°C for 48-60 hrs. During the 16°C incubation, the MBP::PHI-4 polypeptide fusion protein accumulated in the BL21(DE3) cells as a soluble protein.

In order to purify the MBP::PHI-4 fusion protein (SEQ ID NO: 6), the *E. coli* cells were harvested by centrifugation and treated in a lysozyme solution consisting of 2 mg/ml lysozyme in 50 ml sodium phosphate buffer at pH 8.0 containing 300 mM NaCl, 2 U/ml endonuclease (Epicentre) and 5 mM MgCl₂ for 3 hrs. at 37°C with gentle shaking. The lysozyme treated *E. coli* cells were then disrupted with 1% Triton X100 and clear lysate containing the MBP::PHI-4 proteins were prepared by centrifugation at 4000 rpm, 30 min (96 well plates) or 9000 rpm (flask produced samples). His tagged MBP-PHI-4 polypeptide fusion proteins were purified from the clear lysates by affinity chromatography using NiNTA agarose (catalog #30450; Qiagen) following the manufacturer's standard procedure. For high throughput protein purification, Pall 96 deep well filter plates (Pall Corporation; Catalogue #5051) were used for the affinity chromatography. The purified MBP::PHI-4 polypeptide fusion protein was eluted from NiNTA agarose and passed

through Sephadex G25 to change the phosphate buffer to 25 mM HEPES-NaOH, pH 8 and used in insect bioassays for determining the insecticidal activity against Western Corn Rootworm (WCRW). Calipar GXII capillary electrophoresis with a protein chip (Agilent; catalogue #P/N760499) was used to determine the MBP::PHI-4 polypeptide concentrations. The protein analysis was repeated at least 3 times until the final concentration was within the predetermined deviation (less than 10%). Unless otherwise noted, the PHI-4 polypeptides disclosed herein were expressed, purified and assayed for WCRW insecticidal activity as maltose binding protein fusions (i.e. MBP::PHI-4; SEQ ID NO: 6) as described above.

Example 3: Determination of WCRW insecticidal activity of MBP::PHI-4 (SEQ ID NO: 6) and MBP::PHI-4-SFR12-004 (SEQ ID NO: 31) polypeptides.

The activity of MBP::PHI-4 (SEQ ID NO: 6) and MBP::PHI-4-SFR12-004 (SEQ ID NO: 31; Example 8) polypeptides against WCRW (western corn rootworm, *Diabrotica virgifera virgifera*) was determined in an artificial diet feeding assay essentially as described by Cong, R. et al. (Proceedings of the 4th Pacific Rim Conferences on Biotechnology of *Bacillus thuringiensis* and its environmental impact, pp.118-123, ed. by R. J. Akhurst, C. E. Beard and P. Hughes, published in 2002, Canberra, Australia). The assays were conducted on an artificial diet containing dilutions of these polypeptides. The MBP::PHI-4-SFR12-004 polypeptide fusion (SEQ ID NO: 31) and MBP::PHI-4 (SEQ ID NO: 6) fusions were prepared as above, and 10 μ L of protein samples were mixed with 50 μ L of molten (40-50°C) artificial insect diet especially prepared for *Diabrotica* sp. with low temperature melting agarose, whey protein and wheat germ. The diet-PHI-4 polypeptide mixture was placed in each well of a 96 well micro-titer plate. Four or more neonate WCRW larvae were placed in each well to feed for 4 days at 25°C. The response of insects towards the proteins was scored using a 0-3 numerical scoring system based on the size and mortality of the larvae in each well. If no response (or normal growth) was seen, a score of 0 was given. When the growth was slightly retarded, a score of 1 was given. A score of 2 meant that the larvae were severely retarded in growth (close to neonate size). A score of 3 meant death to all the larvae in the well. The percent response (% Response) for each treatment was calculated by dividing the total score, a sum of scores from replicated wells for each treatment, by the total highest possible scores and multiplying by 100 to yield "% Response". For example, if one treatment (one sample, one dose) had 6 replicated wells, the total highest possible score would be 3 X 6 = 18. An observed set of scores of 3, 2, 2, 3, 2, 2 for six wells at a given dose for a given variant would result in (14/18)X100 = 78% Response.

Fast Activity Evaluation (FAE) analysis: The PHI-4 polypeptides at two concentrations were assayed along with 4 doses (two-fold dilutions bracketing the EC50) of the reference MBP::PHI-4 fusion protein (SEQ ID NO: 6) within one 96-well assay plate. The concentrations of the PHI-4 polypeptides were within the 4 doses of the reference protein concentrations, preferably around the middle point of the 4 dose concentrations. Each sample plate contained the reference MBP::PHI-4 protein (SEQ ID NO: 6) in a significant number of wells such as 16 wells in 4 separate doses. In each plate, up to 80 MBP::PHI-4 polypeptide variants were included and assayed for activity comparison with the reference PHI-4 polypeptide protein. From a sample plate, 10 ul of samples from each well were picked by multi-channel pipette and dispensed in one assay plate containing 50 ul molten diet in each well and mixed on a shaker. This process of producing the assay plate was repeated as many as 6 times or more to produce a desired number of replicate assay plates. After the diet was solidified and cooled to 4°C, neonate WCRW larvae were placed in each well, the plate was sealed with perforated Mylar film and incubated in a constant temperature incubator at 25°C. After 4 days, the insect responses were scored under a magnifying glass. The sigmoid dose-response values (Responses) were converted to linear probit dose-response values using SAS-JMP®, Generalized Linear Model, Binomial Response, Probit). The response for each protein in replicates was summed, this sum was compared with the probit dose-response line of the activity reference protein and the nominal fold improvement in potency was calculated. This nominal fold improvement estimated for a given dose in a given experiment is defined as the Fast Activity Evaluation Guide Number (FAEGN). For example, if a PHI-4 polypeptide showed a certain % response value at 40 ppm and comparison to the Probit curve indicated that the same response is predicted for the reference protein at 100 ppm, then the FAE Guide Number is 2.5 (100/40). According to this analysis, the PHI-4 polypeptide is nominally 2.5 times more potent than the reference PHI-4 polypeptide protein. The FAE assay was typically done with 2 different doses of PHI-4 polypeptides at a time and performed in three independent experiments, generating 6 FAEGN for each mutant in a typical FAE evaluation. The mean FAEGN is calculated to yield the Mean FAE Index (MFI). As used herein “Mean FAE Index” (MFI) refers to the mean of multiple FAEGN (typically 6 or more); unless otherwise indicated MFI is understood to be an arithmetic mean of FAEGN. For each protein, a two sided t-test was done comparing the multiple FAEGN from the clone with FAEGN values from the reference protein (typically 48-96 FAEGN of SEQ ID NO: 6). The two-sided t-test was done between these 48-96 FAEGN associated with the reference protein and the 6 or more FAEGN associated with the variant of interest. The Bonferroni correction was used to evaluate p-values (number

of novel proteins/alpha) for the hypothesis that a given variant is significantly different in insecticidal activity compared to the reference protein, MBP::PHI-4 fusion protein (SEQ ID NO: 6) unless otherwise noted. The Bonferroni correction was used to evaluate *p*-values (number of novel proteins/alpha) for the hypothesis that a given variant is significantly different in insecticidal activity compared to the reference protein, MBP::PHI-4 fusion protein (SEQ ID NO: 6) unless otherwise noted.

Effective Concentration₅₀ (EC50) evaluation: Variants of particular interest were assayed at higher power in more extensive dose response curves to more accurately estimate the EC50s. The preparation of dose response, infestation, incubation and scoring were as described for the FAE assay format. EC50 determinations were typically carried out with a no insecticidal protein control plus seven two-fold dilutions bracketing the expected EC50 (typically 100 ppm for MBP::PHI-4; SEQ ID NO: 6) and 24 or more replicate wells within a given experiment. As used herein, the EC50 is defined as the predicted point with 50% response in the scoring scheme. It is a combination of growth or feeding inhibition and lethal responses. In order to determine EC50 values, each treatment (one dose) was repeated 6 or more, usually 24, times.

Data from exemplary FAE and EC50 determinations are given in Figures 1 and 2 respectively. MBP::PHI-4-SFR12-004 (SEQ ID NO: 31; Example 8) is a variant with improved insecticidal activity. Figure 1 shows the primary data for a typical FAE assay. Proteins were purified and quantified as described above. The % response in a typical FAE assay is given on the y axis. The protein concentration (toxin portion of the protein only) is given on the x axis. The geometric mean FAE Index in this instance is 4.0. Figure 2 shows the data for a typical EC50 measurement. Proteins were purified and quantified as described above. The fractional response (multiply by 100 to get "% Response") is given on the y axis. The inferred EC50s are 245 ppm (MBP::PHI-1 of SEQ ID NO: 6) and 48 ppm (SFR12-004; SEQ ID NO: 31). The data indicate that the insecticidal activity of MBP::PHI-4-SFR12-004 (SEQ ID NO: 31) is increased relative to that of MBP::PHI-4 (SEQ ID NO: 6).

The MBP::PHI-4 fusion is rapidly cleaved in the presence of insect gut fluid to yield MBP and mature PHI-4 protein and it is believed that the insecticidal activity is due to the cleaved toxin molecules. The MBP::PHI-4 fusion protein (SEQ ID NO: 6) was digested with 1/100 (w/w) Factor Xa (New England Biolabs) at 25° C overnight and the PHI-4 polypeptide was purified by Superdex 200 column chromatography utilizing the size difference and a weak affinity of MBP to Superdex. The MPB::PHI-4 fusion was also cleaved with trypsin and the mature PHI-4 polypeptide (derived from SEQ ID NO: 6) was purified. The mature, purified PHI-4 polypeptide derived from Factor Xa or trypsin

cleavage has a MW of ~60 kDa as measured on SDS-PAGE gels and is fully reactive to polyclonal antisera that also react with the MBP::PHI-4 fusion protein. The EC50 of this "mature" PHI-4 polypeptide fragment is within experimental error of the EC50 of the MBP::PHI-4 parental protein (SEQ ID NO: 6, as calculated on the basis of ppm associated with the toxin fragment and excluding ppm associated with the MBP domain of SEQ ID NO: 6.

Example 4: Single Amino Acid PHI-4 Polypeptide Variants (S1-S4)

A set of 209 PHI-4 polypeptide single amino acid substitution variants spread across both the N and C terminal portions of the protein were made and characterized (Table 8; MUT ID: 1-209). The mutagenesis template was the polynucleotide of SEQ ID NO: 5 encoding PHI-4 (SEQ ID NO: 6) as an MBP fusion. The mutations were made using the QuickChange kit (Agilent; Catalogue #200524) essentially as described in Example 1. The particular amino acid substitutions relative to PHI-4 (SEQ ID NO: 2) are as indicated in Table 8. For example in Table 8, the PHI-4 polypeptide identified as MUT ID: 1 has a valine substituted for the native amino acid alanine at position 202 of PHI-4 (SEQ ID NO: 2) and referred to "A202V". The polypeptide variant for which activity is reported was prepared as an MBP fusion that is identical to SEQ ID NO: 6 except for this single amino acid substitution. In a similar manner, MUT ID: 1-872 (Table 8) are all made in the context of SEQ ID NO: 6; MUT ID: 873-910 (Table 8) are made in the context of SEQ ID NO: 8; MUT ID: 911-1135 (Table 8) are made in the context of SEQ ID NO: 10. All polypeptides of Table 8 were expressed and purified as MBP fusions as described in Example 2. The PHI-4 polypeptides were expressed as MBP fusions and purified as described in Example 2. The assay protocol for WCRW insecticidal activity of the PHI-4 polypeptides was essentially as described for FAE assays in Example 3 using SEQ ID NO: 6 as the reference protein. For the analysis of the data a "Mean Deviation Score" was calculated rather than a mean FAE Index. This is a related metric that is calculated as follows. Data from four two-fold dilutions of MBP::PHI-4 (typically about 25, 50, 100 and 200 ppm final concentration of PHI-4 fragment in artificial insect diet) is plotted on a Probit plot. The difference between response expected for MBP::PHI-4 (SEQ ID NO: 6) at a given dose, based on the Probit curve, and that observed for the mean score for a PHI-4 polypeptide variant at that given dose is calculated and this difference is defined as the Deviation Score. A negative Deviation Score indicates that the response is lower than that which is expected for the relevant parental backbone (typically MBP::PHI-4; SEQ ID NO: 6) at the same concentration and indicates that the variant is nominally less active than the parental backbone. A positive Deviation Score indicates that the variant is

5 nominally more potent than the parental backbone. As used herein, the Mean Deviation Score refers to the arithmetic mean of multiple Deviation Scores which are typically derived from independent experiments. The Mean Deviation Score is used to estimate rank order of activities associated with a set of variants within a given experiment. The Mean Deviation Scores for the 209 variants of this example are given in Table 8 and is typically an average of at least three independent Deviation Score measurements.

Example 5: Single amino acid substitution mutants #2 (SFR)

A BLAST search revealed that AXMI-205 is a bacterial perforin-like
 10 protein. Perforin proteins from *Clavibacter michiganensis* (GenBank Accession number: YP_001223127; SEQ ID NO: 49), *Laccaria bicolor* (GenBank Accession number: XP_001885969; SEQ ID NO: 48), *Marinomonas sp.* (GenBank Accession number: ZP_01077945; SEQ ID NO: 38), *Nematostella vectensis* (GenBank Accession number: XP_001617993; SEQ ID NO: 50), *Photorhabdus luminescens* (GenBank
 15 Accession number: NP_928713; SEQ ID NO: 1507), and *Serratia proteamaculans* (GenBank Accession number: YP_00147861; SEQ ID NO: 1506) were found to be homologous to AXMI-205. The N-terminal region of up to 311 amino acid of AXMI-205 (SEQ ID NO: 35) is highly homologous to those perforin proteins. Among those perforin proteins, the 3D X-ray structure of *Photorhabdus* perforin-like protein has been published
 20 (Science 317, 1548-1551, 2007; PDB ID: 2QP2; SEQ ID NO: 1508). According to the leading theory of perforin mode of action, the protein inserts 5 alpha helices of its N-terminal region into the target host membrane to form a large size pore (*Proc. Natl. Acad. Sci. USA* 102, 600-605, 2005; *Immunology Today*, 16, 194-201, 1995). In the *Photorhabdus* 2QP2 structure, there are two loops between Alpha C and D and Alpha I
 25 and J are considered to be the site for initiating the membrane insertion of those 5 alpha helices. These loops are called membrane insertion initiation loops. The primary sequences of those perforin proteins listed below were aligned using Vector NTI Align X function in order to identify the membrane insertion loop sequences of AXMI-205 (SEQ ID NO: 35). Amino acid sequences of V92-A103 and G211-E220 of SEQ ID NO: 6 were
 30 aligned with the membrane insertion initiation loops identified in the 2QP2 structure. A number of acidic, basic and other hydrophilic amino acids were found in the AXMI-205 (SEQ ID NO: 35) membrane insertion loops indicating these loops are exposed to the solvent.

Based on this homology model and prior mutation-activity relationship data,
 35 another 664 single amino acid substitution PHI-4 polypeptide variants were made in order to assess the sequence-insecticidal activity-relationships at the selected positions. This

set of 664 point mutations (Table 8; MUT ID: 210-872) were made with MBP::PHI-1 polypeptide of SEQ ID NO: 5 as the DNA template. Mutations were made according to the method of Example 1. The mutagenesis oligonucleotides used to create an exemplary mutant (PHI-4-R97D; SEQ ID NO: 7) are SEQ ID NO: 13 and SEQ ID NO: 14.

5 The other PHI-4 polypeptides (Table 8; MUT ID: 210-872) were made in the same manner using mutagenesis oligonucleotides designed to create the selected substitutions at the desired residues of the protein. The resulting PHI-4 polypeptides were purified as MBP fusions and the activity measured in the FAE assay format or EC50 assay format as described in Examples 2 & 3. The Mean FAE Indices associated with these 664 PHI-4
10 polypeptides are given in lines 210-872 of Table 8.

Example 6: SFR16 point mutants

Another set of 38 single amino acid substitution PHI-4 polypeptide variants (Table 8; MUT ID: 873-910) were made with MBP::PHI-4-R97D (SEQ ID NO: 7; Example 5) as
15 the backbone. The substitutions were selected based on the sequence-activity relationships inferred from the PHI-4 polypeptide variants of the preceding Examples. Mutations were made according to the method of Example 1. Variants were purified as MBP fusions and activity measured in FAE assay format as described in Examples 2 & 3. The Mean FAE Indices associated with these mutants are given in lines 873-910 of Table
20 8. The reference for the Mean FAE Index in this example is SEQ ID NO: 8.

Example 7: PSR3 mutants

The C-terminal domain of AXMI-205 shares significant homology with proteins that have β -prism 3D structural folding patterns. In the typical β -prism, the oligosaccharide
25 binding site is almost always in a cavity formed between the apex and belt-loops from the same Greek-key motif. The primary amino-acid sequence motif has been identified as G-X₃-D (SEQ ID NO: 39) in banana lectin, which has two binding sites (Meagher JL et al *Glycobiology* 15:1033-42; 2005). The primary sequence of the PHI-4 polypeptide of SEQ ID NO: 2 contains 3 such motifs. The canonical structure of these motifs is indicated in
30 Figure 3. In addition, the G-X₃-D motif (SEQ ID NO: 39) can be extended toward the N-terminal direction into the D-X-G-[S/T]-G-X₃-D motif (SEQ ID NO: 40), which are present 1, 2 or 3 times in 24 proteins that are orthologous to the C-terminal portion of AXMI-205 (GenBank accession numbers: gi|136474758; gi|136444345; gi|136141087;
gi|143658948; gi|142085802; gi|135275135; gi|138446054; gi|294814724; gi|170109524;
35 gi|156316804; gi|156377786; gi|170109526; gi|77456557; gi|1209377; gi|302823768; gi|302532087; gi|256764986; gi|302787479; gi|302823738; gi|169762636; gi|302766657;

gi|270056485; gi|302792467; gi|238488445;). Each of the 3 loops in the PHI-4 polypeptide of SEQ ID NO: 2 has potential to bind an oligosaccharide, a putative binding receptor present in WCRW mid-gut cell membrane surface.

As indicated in Table 8, 225 PHI-4 polypeptide variants (MUT IDs: 911-1135) were made to introduce an additional amino acid substitution into the PHI-4 polypeptide of PHI-4-D09 (SEQ ID NO: 10) using MBP::PHI-4-D09 (SEQ ID NO: 9) as the DNA template. The PHI-4-D09 backbone contains the following substitutions relative to SEQ ID NO: 2: L40I, Y98F, L145V, L163V, I172L, V355I, and P412A (numbers are relative to the PHI-4 polypeptide backbone of SEQ ID NO: 2. Mutagenesis was done by a modification of the method of Dominy et al (Methods in Molecular Biology, Vol. 235:209-223; 2003). Briefly, "NNK" mutagenesis at position 396 was done as follows. A pMAL vector encoding SEQ ID NO: 9 was amplified by inverse PCR for 20 cycles using SEQ ID NO: 15 & SEQ ID NO: 16. The PCR product was diluted 10-fold, subjected to one additional round of amplification using SEQ ID NO: 16 & SEQ ID NO: 17. The PCR product was purified on QuiaQuick™ column, phosphorylated with T4 polynucleotide kinase, circularized with T4 DNA ligase and transformed into *E. coli* BL21(DE3) cells. Candidate colonies were amplified by colony PCR and the PCR product was sequenced first with a single primer to confirm the presence of the desired mutation and subsequently sequenced fully with multiple primers to identify clones with no additional mutations. All other PHI-4 polypeptides of this example (Table 8, lines 911-1135) were made by a similar manner using mutagenesis oligonucleotides designed to create the selected substitutions at the desired residues of the protein. Positions with multiple desired mutations were made with degenerate forward primers whereas positions with only one desired mutation were made with non-degenerate primers. Clones with the desired sequences were used to express protein essentially as described in Example 2. Protein purification, activity measurements and statistical analysis was done essentially as described in example 3. The Mean FAE Index reflects the fold difference relative to PHI-4 polypeptide of SEQ ID NO: 6. The Mean EC50 of PHI-4-D09 (SEQ ID NO: 10) was measured at high statistical power and is 1.3-fold improved relative to MBP::PHI-4 (SEQ ID NO: 6). PHI-4 polypeptides of this example with Mean FAE Index > 1.3x are deemed nominally improved relative to the parental backbone (MBP::PHI-4-D09; SEQ ID NO: 10) and diversity meeting this criterion was used for production of subsequent combinatorial mutants.

Example 8: Identification of combinatorial mutants of PHI-4 polypeptides with improved insecticidal activity as measured in an artificial insect diet feeding assay

Activity diversity identified in Examples 4-7 was used to create 192 combinatorial PHI-4 polypeptide variants (SEQ ID NO: 51-242; Table 9). The PHI-4 polypeptide variants were made by sequential point mutagenesis by the method of Example 1. In all cases, the indicated PHI-4 polypeptide variants were made as MBP fusions with the same linker as is indicated in SEQ ID NO: 6. The MBP::PHI-4 polypeptides were expressed and purified as indicated in Example 2 & 3. Purified MBP::PHI-4 polypeptides were assayed in FAE assays to derive a Mean FAE Index or in EC50 assays as indicated in Example 3. Thirty-seven exemplary active PHI-4 polypeptide variants with increased Mean FAE Indices are given in Table 3, along with the sequence variation relative to SEQ ID NO: 6. The substitutions relative to PHI-4 polypeptide of SEQ ID NO: 2 are given in the right-most column. All proteins were expressed and purified as MBP fusion proteins. The reference protein for the Mean FAE Index is MBP::PHI-4 (SEQ ID NO: 6). The functional data on all of the PHI-4 polypeptides of this example is given in lines 51-242 of Table 9.

Table 3

Ex. #	SEQ ID NO:	Alias	FAE	p value	Mutation List (vs. SEQ ID NO: 2)
8	148	SFR11-001	27.4	1.36E-13	R097D, K099L, E220D, K289L, R293Q
8	225	SFR17-013	20.9	0.001577	D042N, E046N, R097D, K099L, E220D, K289L, R293Q, S333K, G336A, S401H, K402H
8	226	SFR17-019	20.1	1.25E-07	D042N, E046N, R097D, K099L, E220D, K289L, R293Q, S333K, G336A, V355I, S401H, K402H, P412A
8	227	SFR17-014	19.5	1.48E-09	D042N, E046N, R097D, K099L, E220D, K289L, S333K, G336A, S401H, K402H, P412A
8	228	SFR17-011	19.0	5.42E-09	D042N, E046N, R097D, K099L, E220D, K289L, S333K, G336A, S401H, K402H
8	229	SFR17-005	17.1	6.82E-10	R097D, K099L, E220D, K289L, R293Q, S401H, K402H, P412A
8	73	SFR10-032	17.1	1.48E-19	R097D, S333K, G336A
8	230	SFR17-018	16.5	7.08E-09	R097D, K099L, E220D, K289L, R293Q, S333K, G336A, V355I, S401H, K402H, P412A
8	149	SFR11-012	14.8	8.29E-15	R097D, K099L
8	231	SFR17-009	13.5	7.19E-16	D042N, E046N, R097D, K099L, E220D, K289L, R293Q, V355I, S401H, K402H, P412A
8	232	SFR17-006	13.1	1.63E-10	D042N, E046N, R097D, K099L, E220D, K289L, R293Q, S401H, K402H, P412A

8	233	SFR17-016	12.9	5.18E-12	D042N, E046N, R097D, K099L, E220D, K289L, R293Q, S333K, G336A, S401H, K402H, P412A
8	150	SFR11-005	12.8	2.12E-10	R097D, K099L, E220D, K289L
8	74	SFR10-042	12.0	7.73E-11	R097D, S401H
8	234	SFR17-012	11.7	1.82E-07	R097D, K099L, E220D, K289L, R293Q, S333K, G336A, S401H, K402H
8	235	SFR17-004	11.3	7.41E-06	D042N, E046N, R097D, K099L, E220D, K289L, S401H, K402H, P412A
8	236	SFR17-017	10.5	2.1E-15	D042N, E046N, R097D, K099L, E220D, K289L, S333K, G336A, V355I, S401H, K402H, P412A
8	151	SFR11-014	9.9	2.08E-09	R097D, K289L
8	237	SFR17-003	9.7	1E-04	D042N, E046N, R097D, K099L, E220D, K289L, R293Q, S401H, K402H
8	183	SFR13-035	9.3	1.5E-11	R097D, K099L, E220D, K289L, V355I, A396T, P412A, K520E
8	75	SFR10-72	8.9	0.003487	R097D, S333V, K520E, Q527K
8	238	SFR17-001	8.7	0.010038	D042N, E046N, R097D, K099L, E220D, K289L, S401H, K402H
8	76	SFR10-056	8.4	2.98E-05	R097D, G462A, R464A, K465M
8	77	SFR10-036	8.2	1.37E-05	R097D, S333K, G336A, E339N
8	239	SFR17-015	8.2	4.75E-21	R097D, K099L, E220D, K289L, R293Q, S333K, G336A, S401H, K402H, P412A
8	152	SFR11-015	8.2	1.76E-07	R097D, R293Q
8	153	SFR11-010	8.1	1.7E-06	R097D, E220D, K289L
8	240	SFR17-002	7.6	5.14E-09	R097D, K099L, E220D, K289L, R293Q, S401H, K402H
8	78	SFR10-039	7.2	9.2E-12	R097D, S333V, G336A, S338V
8	79	SFR10-82	6.7	8.17E-06	R097D, S401H, K402H, K520E, Q527K
8	80	SFR10-045	6.6	2.09E-14	R097D, S401H, K402H
8	81	SFR10-87	6.4	2.15E-06	R097D, S401G, K402H, K520E, Q527K
8	82	SFR10-060	6.3	2.89E-08	R097D, G462A, R464K, K465M
8	51	SFR5-014	6.2	1.65E-05	R097D, R293Q, R416E, K520E
8	184	SFR13-018	6.1	1.84E-13	R097D, K099L, E220D, K289L, V355I, S401G, P412A, K520E
8	154	SFR11-013	5.5	1.54E-09	R097D, E220D
8	241	SFR17-007	5.5	5.11E-22	D042N, E046N, R097D, K099L, E220D, K289L, V355I, S401H, K402H, P412A

Example 9: Identification of combinatorial mutants of PHI-4 polypeptides with improved insecticidal activity as measured in an artificial WCRW insect diet feeding assay.

Activity data from 315 PHI-4 polypeptide combinatorial variants (SEQ ID NOs: 243-558) is provided in Table 9. Libraries were initially prepared by incorporation of diversity into SEQ ID NO: 5. The diversity was largely derived from that described in Example 4. Oligonucleotides encoding diversity at forty positions were incorporated into SEQ ID NO: 5 in a DNA shuffling reaction essentially as described (Ness *et al.*, *Nature Biotechnol.* **20**, 1251; 2002). Briefly, oligonucleotides typically of 30-45 bases in length encoding the diversity elements of interest were mixed at 0.02 - 2 micromolar each in the presence of an appropriate concentration of fragments of SEQ ID NO: 5. This reaction was assembled, rescued and cloned essentially as described for synthetic genes in Example 10 and as described (Ness *et al.*, *Nature Biotechnol.* **20**, 1251; 2002). Improved variants from initial libraries were subjected to family DNA shuffling essentially as described (A. Crameri, *et al Nature* **391**, 288; 1998). This family shuffled library was screened by methods similar to those described in Example 3 (FAE) and to those described in Example 4 (Mean Deviation Score). Selected, improved PHI-4 polypeptide variants from the second round of DNA shuffling were further diversified by recombining N terminal and C terminal domains of elite clones using the method of splicing by overlap extension (R. Horton, *et al.*, *Gene* **77**:61-68; 1989) to yield novel variants. All variants were purified by the method of Example 2 and assayed by the method of Mean Deviations of Example 4. PHI-4 polypeptides variants identified are given in Table 9 (SEQ ID NO: 243-558).

Example 10: Identification of combinatorial mutants of PHI-4 polypeptides with improved insecticidal activity as measured in an artificial WCRW insect diet feeding assay

A set of 158 PHI-4 polypeptide combinatorial variants (SEQ ID NO: 559-716) were prepared by total gene synthesis, essentially as described by Stemmer *et al* (*Gene* **164**:49-53; 1995). An additional treatment was implemented as described (Saaem, I. *et al, Nucleic Acids Research*, Published Nov. 29, 2011, **1-8**). Briefly, in a typical gene synthesis reaction a set of oligonucleotides of 120 bases each encoding both top and bottom strands of the target gene were designed. Complementary oligos typically overlap by 54-65 nucleotides. Oligos to make synthetic genes are combined such that a final concentration of each oligo is approximately 0.05 - 1 uM. Gene assembly is typically done with Herculase II (Agilent) using the following cycling program: 98°C 3 min followed

by (96°C x 30 sec, 40°C x 30 sec, 72°C x 30 sec) x 24 cycles. The initial PCR is then used as template for a second PCR in which a second pair of primers is used to amplify the fully synthetic gene product. Typical PCR conditions for the second synthetic gene reaction were 98°C 3 min followed by (96°C 30 sec, 50°C 30 sec, 72°C 35 sec), x 24 cycles. Reactions were analyzed by 1% E-gels (Invitrogen). A subsequent treatment consisting of a re-annealing step, treatment with Cel1 (Transgenomics; Catalogue #706020) and subsequent amplification (25 cycles) was done essentially as described (Saaem, I. *et al*, *Nucleic Acids Research*, Published Nov. 29, 2011, **1-8**). A third and final amplification of the synthetic gene was done with similar PCR conditions in a single cycle.

The product of this reaction was purified and ligated by Gibson ligation method (New England Biolabs; Catalogue ## E2611L) to an appropriate vector transformed to chemical competent BL21(DE3) cells. Sequence verified clones (Lines 559-716 of Table 9) comprising the PHI-4 polypeptide were expressed as MBP fusions, purified and assayed essentially as described in Examples 2 and 3. Table 4 shows the SEQ ID NOs: and substitutions relative to the PHI-4 polypeptide of SEQ ID NO: 2 for twenty active variants. The mean FAE Index is calculated relative to the MBP::PHI-4 backbone (SEQ ID NO: 6).

Table 4

Ex. #	SEQ ID NO:	Alias	FAE	p value	Mutation List(vs. SEQ ID NO: 2)
10	559	PSR1-1-076	25.6	3.26E-05	D042N, Y098F, L145V, L153I, I172L, Y206F, I283V, V355I, G359A, W389L, I410V, A417S
10	560	PSR1-1-074	22.7	4.19E-05	D042N, Y098F, I283V, V355I
10	561	PSR1-2-145	15.9	7.67E-06	E046N, Y098F, L145V, Y171F, I172L, D182Q, E278N, V355I, I410V, A417S, Q442E, V455I
10	562	PSR1-2-082	11.6	1.46E-18	F043E, Y098F, L145V, Y171F, I172L, Y206F, E278N, E339Q, V355I, V455I, W457N
10	563	PSR1-2-088	11.0	3.91E-05	F043E, Y098F, L145V, Y171F, I172L, Y206F, E278N, V355I, Q442E, V455I, W457N
10	564	PSR1-2-094	10.1	1.07E-10	D042N, F043E, Y098F, L145V, Y171F, I172L, E278N, M354L, V355I, V455I, W457N
10	565	PSR1-2-110	9.7	2.8E-08	D042N, F043E, Y098F, L145V, Y171F, I172L, E278N, V355I, I410V, Q442E, V455I
10	566	PSR1-1-073	8.5	0.003166	D042N, Y098F, I283V, V355I, A417S
10	567	PSR1-2-091	7.2	1.03E-10	D042N, F043E, Y098F, L145V, Y171F, I172L, V355I, Q442E, V455I, W457N

10	568	PSR1- 2-149	7.0	8.39E- 08	F043E, Y098F, L145V, Y171F, I172L, Y206F, E278N, V355I, A417S, V455I, W457N
10	569	PSR1- 2-087	6.9	3.93E- 10	D042N, F043E, Y098F, L145V, Y171F, I172L, E278N, V355I, Q442E, V455I, W457N
10	570	PSR1- 2-158	6.9	1.01E- 07	D042N, R097N, Y098F, L145V, Y171F, I172L, V355I, I410V, V455I
10	571	PSR1- 2-086	6.7	2.02E- 05	E046N, Y098F, L145V, Y171F, I172L, D182Q, E278N, V355I, Q442E, V455I, W457N
10	572	PSR1- 1-053	6.5	0.01072 6	D042N, I052V, Y098F, L145V, I172L, Y206F, I283V, V355I, H370R, I410V, P412A, A417S, T426S, T461S
10	573	PSR1- 2-096	6.2	5.28E- 07	D042N, F043E, Y098F, L145V, Y171F, I172L, V210I, I283V, M354L, V355I, V455I, W457N
10	574	PSR1- 2-135	6.1	2.92E- 11	D042N, R097N, Y098F, L145V, Y171F, I172L, V355I, H370R, Q442E, V455I
10	575	PSR1- 1-014	5.9	0.01827 6	D042N, R097N, Y098F, L145V, I172L, I283V, V355I, I410V, Q442E, V455I
10	576	PSR1- 2-141	5.8	5.02E- 09	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, Q442E, V455I
10	577	PSR1- 1-006	5.6	0.01725 2	E046N, Y098F, L145V, L163V, I172L, Y206F, V210I, E339Q, V355I, A417S
10	578	PSR1- 2-095	5.3	7.35E- 05	F043E, Y098F, L145V, Y171F, I172L, Y206F, E278N, M354L, V355I, V455I, W457N

Example 11: Identification of combinatorial mutants of PHI-4 polypeptides with improved insecticidal activity as measured in an artificial insect diet feeding assay

5 Sixty-six PHI-4 polypeptide variants (SEQ ID NO: 717-783), containing permutations of a number of substitutions, were made by total gene synthesis, essentially as described in Example 10. The substitutions were done in the context of a backbone (PSR1-2-105; SEQ ID NO: 584 from Table 9) containing the following substitutions relative to PHI-4 polypeptide of SEQ ID NO: 2: E46N, R97N, Y98F, L145V, Y171F, I172L, 10 V355I, I410V, V455I, and W457N. The resulting MBP::PHI-4 polypeptide fusion proteins were expressed, purified, assayed for insecticidal activity on WCRW larvae and analyzed for insecticidal relative to MBP::PHI-4 (SEQ ID NO: 6) using the Mean FAE Index metric as described in Examples 2 & 3. Table 5 shows the Mean FAE Indices, SEQ ID NOs: and substitutions relative to PHI-4 polypeptide of SEQ ID NO: 2 for twenty active PHI-4 15 polypeptide variants of this example. The Mean FAE Index was calculated relative to the MBP::PHI-4 backbone (SEQ ID NO: 6). The insecticidal activities of the PHI-4

polypeptides in Table 5 (Mean FAE Index) reflect the arithmetic means of three independent experiments and are expressed as fold WCRW insecticidal activity improvement of the PHI-4 polypeptide variants relative to MBP::PHI-4 (SEQ ID NO: 6). As indicated, the mean FAE Indices range from 0.26x to >8x (fold improvement relative to MBP::PHI-4). The majority of the PHI-4 polypeptides have increased insecticidal activity relative to MBP::PHI-4 (FAE>1). The p values indicate that the measured differences relative to MBP::PHI-4 (SEQ ID NO: 6) are highly significant.

10

Table 5

Ex. #	SEQ ID NO:	Alias	FAE	Mutation List(vs. SEQ ID NO: 2)
11	717	PSR7-141	10.0	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396K, S401K, D403Y, I410V, V455I, W457N
11	718	PSR7-63	8.4	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396L, S401H, K402G, I410V, V455I, W457N
11	719	PSR7-89	8.1	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396L, S401H, D403Y, I410V, V455I, W457N
11	720	PSR7-94	6.6	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396G, S401K, I410V, V455I, W457N
11	721	PSR7-106	6.1	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396K, D403Y, I410V, V455I, W457N
11	722	PSR7-96	6.1	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396L, S401H, K402W, D403Y, I410V, V455I, W457N
11	723	PSR7-100	5.0	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396K, K402H, I410V, V455I, W457N
11	724	PSR7-148	4.7	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396G, S401K, K402G, D403Y, I410V, V455I, W457N
11	725	PSR7-98	4.7	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396K, S401H, K402W, D403Y, I410V, V455I, W457N
11	726	PSR7-113	4.5	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396K, S401K, I410V, V455I, W457N
11	727	PSR7-121	4.5	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, S401H, I410V, V455I, W457N
11	728	PSR7-7	4.5	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396K, S401H, K402H, I410V, V455I, W457N
11	729	PSR7-86	4.2	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396K, S401G, K402H, I410V, V455I, W457N

11	730	PSR7-155	3.7	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, S401H, K402G, D403Y, I410V, V455I, W457N
11	731	PSR7-116	3.6	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, S401K, K402W, D403Y, I410V, V455I, W457N
11	732	PSR7-95	3.4	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, S401G, K402H, D403Y, I410V, V455I, W457N
11	733	PSR7-90	3.3	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396L, K402G, I410V, V455I, W457N
11	734	PSR7-97	3.3	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396G, K402H, I410V, V455I, W457N
11	735	PSR7-64	2.8	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, A396G, S401H, K402W, D403Y, I410V, V455I, W457N
11	736	PSR7-93	2.8	E046N, R097N, Y098F, L145V, Y171F, I172L, V355I, S401H, K402W, D403Y, I410V, V455I, W457N

Example 12: Identification of combinatorial mutants of PHI-4 polypeptides with improved insecticidal activity as measured in an artificial WCRW insect diet feeding assay

Thirty six PHI-4 polypeptide variants (Table 9; SEQ ID NO: 784-819), containing permutations of eleven substitutions (E220D, G336A, K099L, K289L, K402H, K520E, P412A, R097D, S333K, S401H, V355I; numbering scheme as per SEQ ID NO: 2; all variants made as MBP fusions as indicated in Table 9), were made using site directed mutagenesis as described in Example 1 and PHI-4 polypeptide variants were expressed, purified, assayed for insecticidal activity on WCRW larvae and analyzed for insecticidal activity relative to PHI-4 polypeptide of SEQ ID NO: 2 as described in Examples 2 & 3. Table 6 shows the Mean FAE Indices, SEQ ID NOs: and substitutions relative to the PHI-4 polypeptide of SEQ ID NO: 2 for twenty active PHI-4 polypeptide variants of this example. The mean FAE Index is calculated relative to the MBP::PHI-4 backbone (SEQ ID NO: 6). The mean FAE indices reflect the arithmetic means of three independent experiments. As indicated, the mean FAE indices range from 4.5 to >8. The p values indicate that the measured differences relative to MBP::PHI-4 (SEQ ID NO: 6) are highly significant.

Table 6

Ex. #	SEQ ID NO:	Alias	FAE	Mutation List (vs. SEQ ID NO: 2)
12	784	SFR15-009	18.8	R097D,K099L,E220D,K289L,S333K,G336A,S401H,K402H,K520E
12	785	SFR15-019	8.0	R097D,K099L,E220D,K289L,S401H,K402H,P412A
12	786	SFR15-021	7.9	R097D,K099L,E220D,K289L,S333K,G336A,S401H,K402H
12	787	SFR15-033	7.9	R097D,K099L,E220D,K289L,S333K,V355I,S401H,K402H
12	788	SFR15-020	7.3	R097D,K099L,E220D,K289L,V355I,S401H,K402H,P412A
12	789	SFR15-027	7.0	R097D,K099L,E220D,K289L,S333K,V355I,S401H
12	790	SFR15-036	6.6	R097D,K099L,E220D,K289L,V355I,K520E
12	791	SFR15-007	6.6	R097D,K099L,E220D,K289L,S401H,K402H,P412A,K520E
12	792	SFR15-017	5.5	R097D,K099L,E220D,K289L,S401H,K402H
12	793	SFR15-015	5.5	R097D,K099L,E220D,K289L,S333K,G336A,P412A
12	794	SFR15-005	5.4	R097D,K099L,E220D,K289L,S401H,K402H,K520E
12	795	SFR15-001	5.3	R097D,K099L,E220D,K289L,S333K,G336A,K520E
12	796	SFR15-030	5.2	R097D,K099L,E220D,K289L,V355I
12	797	SFR15-025	5.1	R097D,K099L,S333K,G336A,S401H,K402H,K520E
12	798	SFR15-011	5.0	R097D,K099L,E220D,K289L,S333K,G336A,S401H,K402H,P412A,K520E
12	799	SFR15-012	4.9	R097D,K099L,E220D,K289L,S333K,G336A,V355I,S401H,K402H,P412A,K520E
12	800	SFR15-029	4.7	R097D,K099L,E220D,K289L,S333K,V355I,S401H,P412A
12	801	SFR15-016	4.7	R097D,K099L,E220D,K289L,S333K,G336A,V355I,P412A
12	802	SFR15-010	4.5	R097D,K099L,E220D,K289L,S333K,G336A,V355I,S401H,K402H,K520E
12	803	SFR15-003	4.5	R097D,K099L,E220D,K289L,S333K,G336A,P412A,K520E

Example 13: Accordance between FAE and EC50 assays.

5 Experimental data on Mean FAE Index and mean EC50 for twenty-five PHI-4 polypeptide variants is given in Figure 4. PHI-4 polypeptide variants were first tested in the FAE assay and then selected ones were retested in a multiple EC50 assays. In general, the fold improvement in mean FAE Index is modestly larger than the fold improvement that was subsequently measured in Mean EC50 measurements. This

10 overall trend is as expected from the phenomenon of regression toward the mean (International Journal of Epidemiology 2005; 34:215–220). All 25 PHI-4 polypeptides

elected for retesting in EC50 assays repeated as being significantly improved. Figure 4 shows the EC50 measurements for representative variants from Table 9 (SEQ ID NO: 610, SEQ ID NO: 595, SEQ ID NO: 584, SEQ ID NO: 591, SEQ ID NO: 576, SEQ ID NO: 73, SEQ ID NO: 74, SEQ ID NO: 75, SEQ ID NO: 79, SEQ ID NO: 81, SEQ ID NO: 150, SEQ ID NO: 150, SEQ ID NO: 149, SEQ ID NO: 167, SEQ ID NO: 167, SEQ ID NO: 164, SEQ ID NO: 164, SEQ ID NO: 170, SEQ ID NO: 170, SEQ ID NO: 795, SEQ ID NO: 794, SEQ ID NO: 784, SEQ ID NO: 799, SEQ ID NO: 785, SEQ ID NO: 788, SEQ ID NO: 786, SEQ ID NO: 796, SEQ ID NO: 787).

10 **Example 14: Combinatorial substitutions**

Example 11 yielded numerous PHI-4 polypeptide variants with improved insecticidal activity based on combinatorial substitutions of the three lectin-like motifs described in Example 7. A combinatorial library was prepared of 120 genes based on this diversity as follows. SEQ ID NOs: 760 & 761 each contains unique substitutions in loop 1. SEQ ID NOs: 717-726; 728-732; 734-737 & 760 contain unique substitutions in loop 2. SEQ ID NOs: 761 and 758 contain unique substitutions in loop 3. Gene synthesis was used to create a combinatorial library of these loop sequences essentially as described in Example 10. These genes can be expressed and assayed for activity on WCRW larvae by the methods described above.

20

Example 15: Mutagenesis of putative protease sensitive sites of PHI-4 polypeptides

Trypsin was used to identify the site(s) where proteases possibly attack (Protease Accessible Sites) the PHI-4 polypeptide of SEQ ID NO: 2. The PHI-4 polypeptide of SEQ ID NO: 2 in 50mM Tris-HCl, pH8 was mixed with 1/50 (weight/weight) trypsin and incubated for 1 hr. at 37°C. It was found that protein was relatively resistant to trypsin, with no immediate digestion down to the small fragments, but produced a 55 kDa major band and 24 kDa minor band by SDS-PAGE analysis after the incubation. These two bands were excised from the gel and analyzed by mass spectrometry and N-terminal sequencing. The N-terminal sequencing revealed SAANAGQLGN (amino acids 3 - 12 of SEQ ID NO: 2) for the 55 kDa protein indicating that only two amino acid residues, Methionine and Alanine were missing from the N-terminal sequence. The mass-spectrometry, however, showed a loss of C-terminal sequence from Ser at 521 to Leu at 536 of SEQ ID NO: 2. This indicates that trypsin digested the PHI-4 polypeptide protein at the C-terminal side of Lys at 520 of SEQ ID NO: 2. The N-terminal sequence of the 24 kDa band was VDKVLLMD (amino acids 314 to 321 of SEQ ID NO: 2). The mass-spectrometry analysis on the 24 kDa fragment confirmed the C-terminal region of PHI-4

35

polypeptide of SEQ ID NO: 2 starting with Val at 314 relative to SEQ ID NO: 2 as shown by N-terminal sequencing and ending at Lys at 520 relative to SEQ ID NO: 2. This indicated that trypsin digested the PHI-4 polypeptide of SEQ ID NO: 2 at the C-terminal side of Lys at 313 of SEQ ID NO: 2.

5 From this experiment, it was found that there are at least two protease accessible sites, Lys at 313 and Lys at 520 SEQ ID NO: 6. These two sites were mutated to other amino acid residues by saturation mutagenesis and it was found that mutations at Lys at 313 and Lys at 520 of SEQ ID NO: 2 increase the insecticidal activity significantly. For example, the activity of the PHI-4 polypeptide variant, K313Q (MUT ID: 889), was
10 enhanced 2.3 fold over the activity of PHI-4 polypeptide of SEQ ID NO: 2 as measured in an FAE assay (Table 8). The activity of the PHI-4 polypeptide K520Q (MUT ID: 881) was increased 3.1 fold. Activity increases were also found in combinations with other mutations. For example, the activity of the PHI-4 polypeptide having the R097D and K520E substitutions (SEQ ID NO: 52) is 3.5 fold higher than that of PHI-4-R097D (MUT
15 ID: 8) alone by FAE assay.

Example 16: Saturation mutagenesis of amino acid residues selected by site directed single amino acid mutagenesis

Certain amino acid residues showed activity changes when mutated by site
20 directed single amino acid mutagenesis. Those residues were “Selected” for saturation mutagenesis. For the purposes of this example, “Selected” can refer to single amino acid mutations that affect the activity, positively or negatively, relative to the parental backbone in which they were made. More specifically substitutions of Table 8 with Mean FAE Indices of <0.7 or of >1.3 are deemed “Selected”. For example, Selected amino acid
25 residues were found by performing site directed mutagenesis at certain residues such as Arg and Lys. These basic amino acid residues were mutated to either acidic (Asp, Glu) or neutral, polar (e.g.: Asn and Gln) residues, and the activity of those mutants was determined by the FAE insect assay individually. Acidic amino acid residues such as Asp and Glu were changed to basic (e.g.: Arg, Lys) or neutral, polar (e.g.: Asn and Gln)
30 residues and the mutant activity was determined. Neutral, polar amino acid residues such as Gln and Asn were mutated to either acidic (Asp, Glu) or basic (e.g.: Arg, Lys) amino acid residues to see if the activity of those mutants were changed positively (for example mean FAE Index >1.3 relative to the reference protein) or negatively (for example mean FAE Index <0.7 relative to the reference sequence). Another example of finding Selected
35 amino acid residues is based on the sequence-function relationship. Since AXMI-205 is a member of the perforin family it is possible to identify amino acid residues of PHI-4

polypeptide of SEQ ID NO: 2 which are involved in the mode of action elements such as membrane insertion initiation and receptor binding loops. Amino acid residues found in those regions were considered Selected for saturation mutagenesis in this Example. One can use the alanine scanning to empirically define Selected residues. This technique was
5 used to find additional Selected amino acid residues in the putative receptor binding loops.

After any amino acid residues were determined Selected, those residues were subjected to saturation or near saturation mutagenesis to produce a set of up to 19 mutants for each site (20 all possible amino acids minus the amino acid found in the wild
10 type). The insecticidal activity of all these mutants was determined by the FAE insect assay. Saturation mutagenesis of the Selected amino acid residues, was useful for identifying substitutions with Mean FAE Indices of >1 , in many cases >1.3 . When the activity of one single amino acid mutation was found to be positive by showing increased activity over the PHI-4 polypeptide of SEQ ID NO: 2, the saturation mutagenesis enabled
15 us to find other mutation(s) that showed further increased activity. For example, while the FAE Index of E082Q (MUT ID: 370) was positive (1.37), the saturation mutagenesis at this site revealed other mutations showing much higher FAE Indices. For example, the index of E082I (MUT ID: 219) was 7.80 and that of E082L (MUT ID: 259) was 2.71 indicating that PHI-4 polypeptide of SEQ ID NO: 2 hydrophobic residues are beneficial at
20 this site as far as its insecticidal activity is concerned.

Other Selected amino acid substitutions resulted in decreased activity. When these sites were examined further by saturation mutagenesis, substitutions with Mean FAE Indices of >1 were observed. For example, the FAE Indices of K099Q (MUT ID: 677), K099E (MUT ID: 715) were 0.34 and 0.26, respectively. This shows that Lysine at
25 this site is functionally involved in activity and that alternative substitutions may result in improved activity. In this example, the saturation mutagenesis revealed substitutions with Mean FAE Index > 1 . For example, the substitution K099L (MUT ID: 299) has a Mean FAE Index of 5.72 (Table 8). Similar instances were found across the entire PHI-4 polypeptide of SEQ ID NO: 2, for example those indicated in Table 7. Table 7 shows the
30 Mean FAE Indices for nine pairs of substitutions. All data is from Table 8.

Table 7

Substitution A			Substitution B		
MUT ID	Substitution	mean FAE Index	MUT ID	Substitution	mean FAE Index
570	K074Q	0.7	215	K074E	12.40
596	E203Q	0.59	288	E203T	2.18
800	R235Q	0.06	497	R235K	1.34
906	K313E	0.14	889	K313Q	2.29
838	D395Q	0.03	832	D395R	1.60
784	S398A	0.11	342	S398Q	1.51
629	K402Q	0.47	216	K402F	10.20
842	D403Q	0.03	251	D403Y	3.03
611	D447Q	0.54	211	D447K	31.70

5 The serine at position 98 of SEQ ID NO: 2 was selected by alanine scanning amino acid residues found in a region of the protein that is suspected overlap with the receptor binding loops. Point mutants with improved potency may then be used to prepare and screen combinatorial libraries based on that diversity.

Taking mean FAE Index <0.7 as a definition of Selected, the following positions are deemed Selected: P14, D24, Q38, E53, R55, R61, Q75, D76, E83, E118, E126, 10 D152, R166, K188, K191, D193, K242, P243, R248, D254, L266, D268, A270, D274, D298, K313, D315, K316, D321, V343, S349, Q360, R361, D368, I373, D376, F378, D379, D394, Y404, Q413, N430, Q449, D497, R500, S504. Saturation or near saturation mutagenesis at these positions can be performed by the method of Example 7 or equivalent methods and purified and screened the variant proteins for activity by the 15 methods of Example 2 & 3 or equivalent methods.

Example 17: Transgenic expression and activity evaluation

The polynucleotides encoding PHI-4 polypeptides of SEQ ID NOs: 22-25 were cloned under control of the maize ubiquitin promoter (Christensen and Quail, (1996) 20 *Transgenic Research* 5:213-218) into a standard vector suitable for transformation of maize by *Agrobacterium*. Transgenic maize plants were produced by the method of Example 20. Selected T0 plants were tested for susceptibility to WCRW feeding by challenging T0 plants with WCRW larvae. After 19-21 days of challenge, the roots were visually examined and root nodal injury scores were recorded as described (Oleson J. et al 25 *J. Economic Entomology* **98**:1-8; 2005). Root nodal injury scores are indicated in Figure 5. The data support the conclusion that the three PHI-4 polypeptide variants

provide measurable in planta efficacy for protection of maize transgenic plants against WCRW. Figure 5 shows the T0 seedlings in the V3 – V4 growth stage were challenged as described (Oleson J. et al *J. Economic Entomology* **98**:1-8; 2005) and root nodal injury scores were recorded.

5

Example 18: In planta expression of fusion proteins

Localization of the protein can also play an important role in *in planta* accumulation. One can direct proteins such as PHI-4 polypeptides to the chloroplast using a chloroplast targeting peptide (CTP). Additionally, one can direct expression to the apoplastic space using fusions to peptides such as the barley alpha amylase-derived peptide (BAA; SEQ ID NO: 1513). One may also direct transgenically expressed proteins for retention in the endoplasmic reticulum by fusing to both BAA and the sequence "KDEL" (SEQ ID NO: 1514). Proteins can also be directed to the vacuolar space by fusion with the C terminal peptide from plant defensins such as the maize defensin 20 C-terminal propeptide (SEQ ID NO: 511). Other functionally equivalent gene elements may be combined in a similar manner. One may also direct expression specifically to the roots with root-specific promoters. Each of these modifications may be made separately or in combination and any given combination of elements to improve accumulation of protein in plant tissue or in functionally improved efficacy of the expressed protein.

20

Example 19: Transformation of Maize by Particle Bombardment and Regeneration of Transgenic Plants

Immature maize embryos from greenhouse donor plants are bombarded with a DNA molecule containing the PHI-4 polypeptide of nucleotide sequence (e.g., SEQ ID NO: 1) operably linked to an ubiquitin promoter and the selectable marker gene PAT (Wohlleben, *et al.*, (1988) *Gene* 70: 25-37), which confers resistance to the herbicide Bialaphos. Alternatively, the selectable marker gene is provided on a separate DNA molecule. Transformation is performed as follows. Media recipes follow below.

30

Preparation of Target Tissue

The ears are husked and surface sterilized in 30% CLOROX™ bleach plus 0.5% Micro detergent for 20 minutes, and rinsed two times with sterile water. The immature embryos are excised and placed embryo axis side down (scutellum side up), 25 embryos per plate, on 560Y medium for 4 hours and then aligned within the 2.5 cm target zone in preparation for bombardment.

35

Preparation of DNA

A plasmid vector comprising a nucleotide sequence (e.g., SEQ ID NO: 1) operably linked to an ubiquitin promoter is made. For example, a suitable transformation vector comprises a UBI1 promoter from *Zea mays*, a 5' UTR from UBI1 and a UBI1 intron, in combination with a PinII terminator. The vector additionally contains a PAT selectable marker gene driven by a CAMV35S promoter and includes a CAMV35S terminator. Optionally, the selectable marker can reside on a separate plasmid. A DNA molecule comprising a toxin nucleotide sequence as well as a PAT selectable marker is precipitated onto 1.1 μm (average diameter) tungsten pellets using a CaCl_2 precipitation procedure as follows:

- 100 μL prepared tungsten particles in water
- 10 μL (1 μg) DNA in Tris EDTA buffer (1 μg total DNA)
- 100 μL 2.5 M CaCl_2
- 10 μL 0.1 M spermidine

Each reagent is added sequentially to a tungsten particle suspension, while maintained on the multitube vortexer. The final mixture is sonicated briefly and allowed to incubate under constant vortexing for 10 minutes. After the precipitation period, the tubes are centrifuged briefly, liquid removed, washed with 500 mL 100% ethanol, and centrifuged for 30 seconds. Again the liquid is removed, and 105 μL 100% ethanol is added to the final tungsten particle pellet. For particle gun bombardment, the tungsten/DNA particles are briefly sonicated and 10 μL spotted onto the center of each macrocarrier and allowed to dry about 2 minutes before bombardment.

Particle Gun Treatment

The sample plates are bombarded at level #4 in particle gun #HE34-1 or #HE34-2. All samples receive a single shot at 650 PSI, with a total of ten aliquots taken from each tube of prepared particles/DNA.

Subsequent Treatment

Following bombardment, the embryos are kept on 560Y medium for 2 days, then transferred to 560R selection medium containing 3 mg/liter Bialaphos, and subcultured every 2 weeks. After approximately 10 weeks of selection, selection-resistant callus clones are transferred to 288J medium to initiate plant regeneration. Following somatic embryo maturation (2-4 weeks), well-developed somatic embryos are transferred to medium for germination and transferred to the lighted culture room. Approximately 7-10

days later, developing plantlets are transferred to 272V hormone-free medium in tubes for 7-10 days until plantlets are well established. Plants are then transferred to inserts in flats (equivalent to 2.5" pot) containing potting soil and grown for 1 week in a growth chamber, subsequently grown an additional 1-2 weeks in the greenhouse, then transferred to
 5 classic 600 pots (1.6 gallon) and grown to maturity. Plants are monitored and scored for expression of the toxin by assays known in the art or as described above.

Bombardment and Culture Media

Bombardment medium (560Y) comprises 4.0 g/L N6 basal salts (SIGMA C-1416),
 10 1.0 mL/L Eriksson's Vitamin Mix (1000x SIGMA-1511), 0.5 mg/L thiamine HCl, 120.0 g/L sucrose, 1.0 mg/L 2,4-D and 2.88 g/L L-proline (brought to volume with deionized H₂O following adjustment to pH 5.8 with KOH); 2.0 g/L Gelrite™ (added after bringing to volume with dl H₂O); and 8.5 mg/L silver nitrate (added after sterilizing the medium and cooling to room temperature). Selection medium (560R) comprises 4.0 g/L N6 basal salts
 15 (SIGMA C-1416), 1.0 mL/L Eriksson's Vitamin Mix (1000x SIGMA-1511), 0.5 mg/L thiamine HCl, 30.0 g/L sucrose, and 2.0 mg/L 2,4-D (brought to volume with dl H₂O following adjustment to pH 5.8 with KOH); 3.0 g/L Gelrite™ (added after bringing to volume with dl H₂O); and 0.85 mg/L silver nitrate and 3.0 mg/L Bialaphos (both added after sterilizing the medium and cooling to room temperature).

20 Plant regeneration medium (288J) comprises 4.3 g/L MS salts (GIBCO 11117-074), 5.0 mL/L MS vitamins stock solution (0.100 g nicotinic acid, 0.02 g/L thiamine HCl, 0.10 g/L pyridoxine HCl, and 0.40 g/L Glycine brought to volume with polished D-I H₂O) (Murashige and Skoog, (1962) *Physiol. Plant.* 15:473), 100 mg/L myo-inositol, 0.5 mg/L zeatin, 60 g/L sucrose, and 1.0 mL/L of 0.1 mM abscisic acid (brought to volume with
 25 polished dl H₂O after adjusting to pH 5.6); 3.0 g/L Gelrite™ (added after bringing to volume with dl H₂O); and 1.0 mg/L indoleacetic acid and 3.0 mg/L Bialaphos (added after sterilizing the medium and cooling to 60 °C).

Hormone-free medium (272V) comprises 4.3 g/L MS salts (GIBCO 11117-074), 5.0 mL/L MS vitamins stock solution (0.100 g/L nicotinic acid, 0.02 g/L thiamine HCl, 0.10
 30 g/L pyridoxine HCl, and 0.40 g/L Glycine brought to volume with polished dl H₂O), 0.1 g/L myo-inositol, and 40.0 g/L sucrose (brought to volume with polished dl H₂O after adjusting pH to 5.6); and 6 g/L Bacto-agar (added after bringing to volume with polished dl H₂O), sterilized and cooled to 60°C.

35 **Example 20: *Agrobacterium*-Mediated Transformation of Maize and Regeneration of Transgenic Plants**

For *Agrobacterium*-mediated transformation of maize with a toxin nucleotide sequence (e.g., SEQ ID NO: 1), the method of Zhao can be used (US Patent Number 5,981,840 and PCT Patent Publication Number WO 1998/32326; the contents of which are hereby incorporated by reference). Briefly, immature embryos are isolated from

5 maize and the embryos contacted with a suspension of *Agrobacterium* under conditions whereby the bacteria are capable of transferring the nucleotide sequence (e.g. SEQ ID NO: 1) to at least one cell of at least one of the immature embryos (step 1: the infection step). In this step the immature embryos can be immersed in an *Agrobacterium* suspension for the initiation of inoculation. The embryos are co-cultured for a time with

10 the *Agrobacterium* (step 2: the co-cultivation step). The immature embryos can be cultured on solid medium following the infection step. Following this co-cultivation period an optional "resting" step is contemplated. In this resting step, the embryos are incubated in the presence of at least one antibiotic known to inhibit the growth of *Agrobacterium* without the addition of a selective agent for plant transformants (step 3: resting step). The

15 immature embryos can be cultured on solid medium with antibiotic, but without a selecting agent, for elimination of *Agrobacterium* and for a resting phase for the infected cells. Next, inoculated embryos are cultured on medium containing a selective agent and growing transformed callus is recovered (step 4: the selection step). The immature embryos are cultured on solid medium with a selective agent resulting in the selective

20 growth of transformed cells. The callus is then regenerated into plants (step 5: the regeneration step), and calli grown on selective medium can be cultured on solid medium to regenerate the plants.

Table Legends

25 Table 8. The definitions of the column headings are as follows: "MUT ID", a unique identifier for each substitutions; "Backbone", the SEQ ID corresponding to the polypeptide backbone in which the substitution was made; "Position", amino acid position according to the numbering convention of SEQ ID NO: 2, "Ref. A.A.", the standard single letter code for the amino acid present in the backbone sequence at the indicated position; "Substitution",

30 the standard single letter code for the amino acid present in the mutant sequence at the indicated position; "FAE", the arithmetic Mean FAE Index as further defined in Example 3; "p-value" the calculated p value associated with the hypothesis that the variant polypeptide is significantly different than the reference protein used in that particular FAE assay, as defined further in Example 3; "EC50 (ppm)", EC50 as defined in example 3 with

35 the EC50 dose given in ppm for the toxin portion of the sample; "Deviation", Mean Deviation Score as defined in Example 4; "Example #", the example number

corresponding to the creation of the variant. The reference protein against which the variant protein is compared is: (MUT IDs: 1-872 and 911-1135) used SEQ ID NO: 6 as the reference protein; (MUT IDs: 873-910) used SEQ ID NO: 8 as the reference protein.

Table 8

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
1	PHI-4	202	A	V			154	0.45	4
2	PHI-4	342	A	S			129	0.32	4
3	PHI-4	417	A	S			96	-0.13	4
4	PHI-4	24	D	N			96	0.23	4
5	PHI-4	42	D	N			100	0.62	4
6	PHI-4	331	E	D			122	-0.02	4
7	PHI-4	109	F	I			202		4
8	PHI-4	300	F	Y			132	0.26	4
9	PHI-4	359	G	A			98	-0.06	4
10	PHI-4	52	I	V			202		4
11	PHI-4	133	I	L			124	-0.21	4
12	PHI-4	172	I	L			123	0.26	4
13	PHI-4	283	I	V			187		4
14	PHI-4	410	I	V			100	0.05	4
15	PHI-4	40	L	I			113		4
16	PHI-4	145	L	V			102		4
17	PHI-4	153	L	I			79	0.22	4
18	PHI-4	163	L	V			104		4
19	PHI-4	296	L	I			116	0.36	4
20	PHI-4	418	L	M			84	0.1	4
21	PHI-4	154	N	D			108	0.01	4
22	PHI-4	346	P	A			85		4
23	PHI-4	411	P	A			91	0.06	4
24	PHI-4	412	P	A			79	0.2	4
25	PHI-4	34	S	A			176		4
26	PHI-4	78	S	G			144	-0.23	4
27	PHI-4	335	S	T			160	-0.26	4
28	PHI-4	426	T	S			93	-0.13	4
29	PHI-4	461	T	S			54		4
30	PHI-4	343	V	I			89	0.39	4
31	PHI-4	355	V	I			78	0.39	4
32	PHI-4	392	V	I			121		4
33	PHI-4	421	V	L			106		4
34	PHI-4	440	V	L			75	0.26	4
35	PHI-4	456	W	Y			199		4
36	PHI-4	98	Y	F			71		4
37	PHI-4	121	Y	F			117	0.11	4
38	PHI-4	206	Y	F			82	0.51	4
39	PHI-4	337	A	G				-0.1	4
40	PHI-4	364	A	S				0.09	4

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
41	PHI-4	371	A	G				0.02	4
42	PHI-4	371	A	T				0.24	4
43	PHI-4	385	A	G				0.08	4
44	PHI-4	385	A	P				-0.23	4
45	PHI-4	396	A	E				-0.56	4
46	PHI-4	405	A	S				0.09	4
47	PHI-4	405	A	W				-0.38	4
48	PHI-4	409	A	P				0.25	4
49	PHI-4	417	A	C				0.11	4
50	PHI-4	445	C	L				0.16	4
51	PHI-4	445	C	T				0.06	4
52	PHI-4	331	E	N				0.24	4
53	PHI-4	339	E	N				0.04	4
54	PHI-4	339	E	Q				0.34	4
55	PHI-4	344	F	W				0.16	4
56	PHI-4	374	F	I				0.1	4
57	PHI-4	378	F	I				-0.78	4
58	PHI-4	351	G	V				0.1	4
59	PHI-4	397	G	R				0.29	4
60	PHI-4	428	G	S				-0.17	4
61	PHI-4	373	I	V				-0.52	4
62	PHI-4	375	K	R				0.19	4
63	PHI-4	384	K	A				0.05	4
64	PHI-4	341	L	V				0.19	4
65	PHI-4	380	L	G				0.17	4
66	PHI-4	383	L	I				0.07	4
67	PHI-4	383	L	V				0.17	4
68	PHI-4	354	M	L				0.29	4
69	PHI-4	422	M	V				0.05	4
70	PHI-4	345	N	H				0.08	4
71	PHI-4	362	N	S				0.2	4
72	PHI-4	430	N	D				-0.7	4
73	PHI-4	453	N	D				0.02	4
74	PHI-4	372	P	L				-0.13	4
75	PHI-4	390	Q	D				-0.36	4
76	PHI-4	452	Q	G				-0.1	4
77	PHI-4	391	R	L				0.16	4
78	PHI-4	333	S	E				-0.34	4
79	PHI-4	333	S	R				0.14	4
80	PHI-4	349	S	F				-0.53	4
81	PHI-4	349	S	P				-0.9	4
82	PHI-4	398	S	G				-0.71	4
83	PHI-4	427	S	N				-0.08	4
84	PHI-4	427	S	T				-0.22	4
85	PHI-4	343	V	F				-0.81	4
86	PHI-4	355	V	L				0.09	4
87	PHI-4	382	V	D				0.23	4

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
88	PHI-4	382	V	L				0.16	4
89	PHI-4	387	V	T				-0.33	4
90	PHI-4	392	V	L				-0.26	4
91	PHI-4	438	V	R				0.28	4
92	PHI-4	455	V	I				0.81	4
93	PHI-4	389	W	F				-0.06	4
94	PHI-4	389	W	Y				0.19	4
95	PHI-4	457	W	N				0.34	4
96	PHI-4	352	Y	C				-0.34	4
97	PHI-4	352	Y	F				-0.2	4
98	PHI-4	404	Y	F				-0.04	4
99	PHI-4	404	Y	G				-0.7	4
100	PHI-4	429	Y	E				-0.36	4
101	PHI-4	437	Y	I				-0.01	4
102	PHI-4	437	Y	V				0.05	4
103	PHI-4	30	A	C				-0.08	4
104	PHI-4	103	A	G				0.03	4
105	PHI-4	127	A	T				0.1	4
106	PHI-4	185	A	S				0.28	4
107	PHI-4	238	A	T				-0.24	4
108	PHI-4	263	A	S				-0.01	4
109	PHI-4	270	A	P				-0.6	4
110	PHI-4	287	A	C				0.37	4
111	PHI-4	182	D	Q				0.5	4
112	PHI-4	193	D	N				0.32	4
113	PHI-4	268	D	N				0.08	4
114	PHI-4	46	E	D				0.47	4
115	PHI-4	46	E	N				0.67	4
116	PHI-4	80	E	S				0.27	4
117	PHI-4	83	E	S				-0.62	4
118	PHI-4	162	E	D				0.28	4
119	PHI-4	278	E	N				0.61	4
120	PHI-4	43	F	E				0.53	4
121	PHI-4	73	F	Y				0.01	4
122	PHI-4	149	F	A				-0.05	4
123	PHI-4	149	F	V				0.25	4
124	PHI-4	303	F	Y				0.08	4
125	PHI-4	22	G	S				0.07	4
126	PHI-4	50	I	V				0.31	4
127	PHI-4	119	I	N				-0.22	4
128	PHI-4	213	I	L				-0.05	4
129	PHI-4	207	K	A				-0.4	4
130	PHI-4	214	K	S				0.11	4
131	PHI-4	36	L	M				0.12	4
132	PHI-4	100	L	F				-0.31	4
133	PHI-4	105	L	I				0.22	4
134	PHI-4	141	L	H				-0.34	4

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
135	PHI-4	181	L	A				-0.33	4
136	PHI-4	266	L	I				-0.73	4
137	PHI-4	266	L	V				-0.47	4
138	PHI-4	19	M	I				-0.04	4
139	PHI-4	19	M	L				0.31	4
140	PHI-4	88	M	I				0.2	4
141	PHI-4	88	M	L				0	4
142	PHI-4	204	M	A				0.1	4
143	PHI-4	245	M	L				0.34	4
144	PHI-4	155	N	K				0.11	4
145	PHI-4	231	N	S				0.32	4
146	PHI-4	14	P	A				0.13	4
147	PHI-4	159	P	D				0.2	4
148	PHI-4	243	P	L				-0.54	4
149	PHI-4	282	P	G				0.26	4
150	PHI-4	55	R	K				0.3	4
151	PHI-4	61	R	K				0.24	4
152	PHI-4	97	R	N				0.59	4
153	PHI-4	292	R	Q				0.27	4
154	PHI-4	56	S	T				0.36	4
155	PHI-4	173	S	A				0.29	4
156	PHI-4	184	S	T				0.27	4
157	PHI-4	219	S	N				-0.4	4
158	PHI-4	230	S	E				0.11	4
159	PHI-4	276	S	A				0.36	4
160	PHI-4	279	S	P				-0.35	4
161	PHI-4	58	T	S				0.09	4
162	PHI-4	112	T	S				0.04	4
163	PHI-4	117	T	S				0.18	4
164	PHI-4	189	T	K				0.08	4
165	PHI-4	94	V	I				0.18	4
166	PHI-4	210	V	I				0.41	4
167	PHI-4	57	Y	F				0.26	4
168	PHI-4	167	Y	W				0.08	4
169	PHI-4	170	Y	H				0.23	4
170	PHI-4	171	Y	F				1.29	4
171	PHI-4	183	Y	V				0.05	4
172	PHI-4	186	A	V				-0.34	4
173	PHI-4	342	A	V				0.24	4
174	PHI-4	445	C	S				-0.24	4
175	PHI-4	321	D	E				-0.27	4
176	PHI-4	46	E	G				0.02	4
177	PHI-4	222	E	G				-0.34	4
178	PHI-4	297	E	G				-0.42	4
179	PHI-4	344	F	Y				-0.07	4
180	PHI-4	483	F	S				-0.19	4
181	PHI-4	66	H	R				0	4

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
182	PHI-4	441	H	R				-0.13	4
183	PHI-4	172	I	V				0.07	4
184	PHI-4	384	K	G				0.19	4
185	PHI-4	465	K	E				-0.19	4
186	PHI-4	209	L	P				0.4	4
187	PHI-4	236	L	P				0.44	4
188	PHI-4	144	M	I				-0.03	4
189	PHI-4	158	M	R				-0.4	4
190	PHI-4	12	N	D				-0.28	4
191	PHI-4	155	N	D				-0.02	4
192	PHI-4	350	N	S				0.08	4
193	PHI-4	14	P	L				-1.16	4
194	PHI-4	115	Q	L				-0.14	4
195	PHI-4	306	Q	L				-0.39	4
196	PHI-4	309	Q	R				-0.3	4
197	PHI-4	134	S	G				-0.21	4
198	PHI-4	195	S	N				-0.02	4
199	PHI-4	504	S	C				-0.59	4
200	PHI-4	189	T	I				0.26	4
201	PHI-4	233	T	A				-0.33	4
202	PHI-4	16	V	D				0.03	4
203	PHI-4	294	V	A				0.03	4
204	PHI-4	355	V	G				-0.31	4
205	PHI-4	438	V	A				-0.15	4
206	PHI-4	448	V	A				0.05	4
207	PHI-4	284	W	R				0.11	4
208	PHI-4	84	Y	F				0.35	4
209	PHI-4	167	Y	C				0.1	4
210	PHI-4	97	R	D	125.8	0.01			5
211	PHI-4	447	D	K	31.7	0.598			5
212	PHI-4	334	G	R	25.2	0			5
213	PHI-4	527	Q	K	25	0			5
214	PHI-4	109	F	K	14.3	0			5
215	PHI-4	74	K	E	12.4	0.152			5
216	PHI-4	402	K	F	10.2	0			5
217	PHI-4	336	G	A	8	0			5
218	PHI-4	527	Q	P	7.8	0			5
219	PHI-4	82	E	I	7.8	0.086			5
220	PHI-4	109	F	G	6.8	0.151			5
221	PHI-4	97	R	E	6.7	0			5
222	PHI-4	220	E	H	6.6	0			5
223	PHI-4	165	K	E	6.6	0			5
224	PHI-4	289	K	L	6.6	0			5
225	PHI-4	454	R	Y	6.4	0			5
226	PHI-4	109	F	M	6.4	0			5
227	PHI-4	247	D	Y	6.1	0			5
228	PHI-4	454	R	M	5.8	0			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
229	PHI-4	99	K	L	5.7	0			5
230	PHI-4	289	K	V	5.1	0			5
231	PHI-4	109	F	S	5	0.374			5
232	PHI-4	289	K	P	5	0			5
233	PHI-4	454	R	S	4.6	0			5
234	PHI-4	220	E	D	4.4	0			5
235	PHI-4	334	G	K	4.3	0.395			5
236	PHI-4	459	K	M	4.2	0			5
237	PHI-4	97	R	Q	4	0			5
238	PHI-4	454	R	V	4	0.001			5
239	PHI-4	517	Q	I	4	0.006			5
240	PHI-4	99	K	Y	4	0			5
241	PHI-4	256	Q	K	4	0.192			5
242	PHI-4	109	F	D	3.9	0.596			5
243	PHI-4	220	E	T	3.8	0			5
244	PHI-4	196	Q	K	3.7	0.08			5
245	PHI-4	517	Q	F	3.7	0.001			5
246	PHI-4	79	K	E	3.6	0.006			5
247	PHI-4	454	R	I	3.5	0.002			5
248	PHI-4	454	R	K	3.5	0.001			5
249	PHI-4	289	K	E	3.3	0.994			5
250	PHI-4	74	K	G	3.3	0			5
251	PHI-4	403	D	Y	3.3	0			5
252	PHI-4	166	R	Q	3.3	0.992			5
253	PHI-4	517	Q	K	3.3	0.162			5
254	PHI-4	447	D	Y	3.2	0			5
255	PHI-4	289	K	Q	2.8	0.028			5
256	PHI-4	454	R	F	2.8	0.011			5
257	PHI-4	220	E	Y	2.7	0.003			5
258	PHI-4	447	D	S	2.7	0			5
259	PHI-4	82	E	L	2.7	0.881			5
260	PHI-4	196	Q	N	2.7	0			5
261	PHI-4	216	E	Q	2.7	0.017			5
262	PHI-4	334	G	I	2.6	0.893			5
263	PHI-4	151	D	S	2.6	0.001			5
264	PHI-4	454	R	W	2.6	0.005			5
265	PHI-4	165	K	Q	2.6	0.878			5
266	PHI-4	459	K	V	2.6	0.024			5
267	PHI-4	148	D	F	2.6	0.001			5
268	PHI-4	220	E	V	2.5	0.016			5
269	PHI-4	454	R	Q	2.5	0.068		0.26	5
270	PHI-4	447	D	E	2.5	0			5
271	PHI-4	527	Q	C	2.5	0.001			5
272	PHI-4	196	Q	D	2.5	0.004			5
273	PHI-4	82	E	Y	2.5	0.938			5
274	PHI-4	527	Q	E	2.4	0.423			5
275	PHI-4	402	K	H	2.4	0.016			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
276	PHI-4	459	K	W	2.4	0.043			5
277	PHI-4	459	K	Q	2.3	0.116			5
278	PHI-4	289	K	Y	2.3	0.027			5
279	PHI-4	99	K	I	2.3	0.024			5
280	PHI-4	289	K	T	2.3	0.031			5
281	PHI-4	220	E	S	2.3	0.037			5
282	PHI-4	459	K	I	2.3	0.049			5
283	PHI-4	462	G	A	2.3	0.017			5
284	PHI-4	99	K	M	2.2	0.025			5
285	PHI-4	289	K	D	2.2	0.032			5
286	PHI-4	109	F	N	2.2	0.061			5
287	PHI-4	220	E	Q	2.2	0.166			5
288	PHI-4	203	E	T	2.2	0.973			5
289	PHI-4	257	Q	I	2.1	0.013			5
290	PHI-4	203	E	H	2.1	0.948			5
291	PHI-4	151	D	A	2.1	0.017			5
292	PHI-4	447	D	I	2.1	0.017			5
293	PHI-4	97	R	G	2.1	0.068			5
294	PHI-4	151	D	N	2	0.026			5
295	PHI-4	148	D	P	2	0.001			5
296	PHI-4	97	R	S	2	0.106		0.45	5
297	PHI-4	151	D	W	2	0.042			5
298	PHI-4	257	Q	E	2	0.543			5
299	PHI-4	109	F	E	2	0.926			5
300	PHI-4	527	Q	S	2	0.282			5
301	PHI-4	403	D	W	2	0.001			5
302	PHI-4	518	E	Q	1.9	0.377			5
303	PHI-4	460	G	A	1.9	0.124			5
304	PHI-4	499	E	Q	1.9	0.395			5
305	PHI-4	148	D	V	1.9	0.003			5
306	PHI-4	148	D	E	1.9	0.004			5
307	PHI-4	459	K	T	1.9	0.232			5
308	PHI-4	289	K	F	1.9	0.158			5
309	PHI-4	289	K	S	1.9	0.149			5
310	PHI-4	151	D	V	1.9	0.079			5
311	PHI-4	402	K	R	1.9	0.247		0.08	5
312	PHI-4	196	Q	E	1.8	0.482			5
313	PHI-4	525	Q	K	1.8	0.647			5
314	PHI-4	289	K	M	1.8	0.189			5
315	PHI-4	302	E	Q	1.8	0.505			5
316	PHI-4	403	D	F	1.8	0.009			5
317	PHI-4	148	D	H	1.8	0.011			5
318	PHI-4	165	K	P	1.8	0.231			5
319	PHI-4	459	K	S	1.8	0.323			5
320	PHI-4	24	D	Q	1.8	0.766			5
321	PHI-4	151	D	Q	1.7	0.766			5
322	PHI-4	289	K	R	1.7	0.256			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
323	PHI-4	196	Q	A	1.7	0.142			5
324	PHI-4	459	K	H	1.7	0.408			5
325	PHI-4	454	R	G	1.7	0.407			5
326	PHI-4	395	D	C	1.7	0.807			5
327	PHI-4	220	E	R	1.7	0.414			5
328	PHI-4	99	K	F	1.6	0.363			5
329	PHI-4	289	K	W	1.6	0.457			5
330	PHI-4	527	Q	H	1.6	0.251			5
331	PHI-4	220	E	W	1.6	0.508			5
332	PHI-4	9	Q	K	1.6	0.756			5
333	PHI-4	99	K	C	1.6	0.454			5
334	PHI-4	309	Q	K	1.6	0.737			5
335	PHI-4	148	D	W	1.6	0.054			5
336	PHI-4	216	E	F	1.6	0.451			5
337	PHI-4	99	K	V	1.6	0.487			5
338	PHI-4	454	R	H	1.6	0.577			5
339	PHI-4	309	Q	E	1.5	0.704			5
340	PHI-4	165	K	T	1.5	0.519			5
341	PHI-4	165	K	A	1.5	0.529			5
342	PHI-4	398	S	Q	1.5	0.523			5
343	PHI-4	454	R	D	1.5	0.628			5
344	PHI-4	142	R	E	1.5	0.934			5
345	PHI-4	449	Q	E	1.5	0.713			5
346	PHI-4	196	Q	I	1.5	0.425			5
347	PHI-4	278	E	Q	1.5	0.065			5
348	PHI-4	502	R	E	1.5	0.962			5
349	PHI-4	165	K	L	1.5	0.602			5
350	PHI-4	90	Q	E	1.5	0.721			5
351	PHI-4	399	G	A	1.5	0.587			5
352	PHI-4	447	D	G	1.5	0.157		0.12	5
353	PHI-4	151	D	C	1.5	0.485			5
354	PHI-4	289	K	I	1.5	0.636			5
355	PHI-4	459	K	C	1.5	0.73			5
356	PHI-4	220	E	M	1.4	0.611			5
357	PHI-4	454	R	L	1.4	0.721			5
358	PHI-4	459	K	Y	1.4	0.736			5
359	PHI-4	442	Q	K	1.4	0.751			5
360	PHI-4	99	K	N	1.4	0.687			5
361	PHI-4	402	K	W	1.4	0.758			5
362	PHI-4	216	E	V	1.4	0.658			5
363	PHI-4	165	K	G	1.4	0.731			5
364	PHI-4	214	K	Q	1.4	0.701			5
365	PHI-4	165	K	D	1.4	0.741			5
366	PHI-4	165	K	V	1.4	0.744			5
367	PHI-4	220	E	A	1.4	0.694			5
368	PHI-4	289	K	H	1.4	0.774			5
369	PHI-4	165	K	H	1.4	0.785			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
370	PHI-4	82	E	Q	1.4	0.875		-0.77	5
371	PHI-4	109	F	C	1.4	0.764			5
372	PHI-4	220	E	F	1.4	0.726			5
373	PHI-4	51	E	Q	1.4	0.853			5
374	PHI-4	459	K	P	1.3	0.888			5
375	PHI-4	447	D	P	1.3	0.328			5
376	PHI-4	165	K	I	1.3	0.831			5
377	PHI-4	220	E	I	1.3	0.764			5
378	PHI-4	459	K	N	1.3	0.905			5
379	PHI-4	454	R	T	1.3	0.926			5
380	PHI-4	87	K	Q	1.3	0.688			5
381	PHI-4	402	K	G	1.3	0.96			5
382	PHI-4	289	K	N	1.3	0.91			5
383	PHI-4	148	D	A	1.3	0.428			5
384	PHI-4	216	E	Y	1.3	0.857			5
385	PHI-4	306	Q	K	1.3	0.669			5
386	PHI-4	148	D	R	1.3	0.572			5
387	PHI-4	151	D	M	1.3	0.852			5
388	PHI-4	257	Q	C	1.3	0.848			5
389	PHI-4	9	Q	E	1.3	0.687			5
390	PHI-4	402	K	N	1.2	0.927			5
391	PHI-4	148	D	L	1.2	0.614			5
392	PHI-4	148	D	S	1.2	0.628			5
393	PHI-4	436	D	K	1.2	0.755			5
394	PHI-4	459	K	A	1.2	0.886			5
395	PHI-4	447	D	L	1.2	0.673			5
396	PHI-4	454	R	P	1.2	0.981			5
397	PHI-4	398	S	C	1.2	0.984			5
398	PHI-4	76	D	Q	1.2	0.755			5
399	PHI-4	220	E	L	1.2	0.994			5
400	PHI-4	165	K	M	1.2	0.903			5
401	PHI-4	196	Q	R	1.2	0.819			5
402	PHI-4	402	K	L	1.2	0.848			5
403	PHI-4	220	E	C	1.2	0.994			5
404	PHI-4	403	D	E	1.2	0.986			5
405	PHI-4	220	E	N	1.2	0.836			5
406	PHI-4	42	D	Q	1.2	0.754			5
407	PHI-4	165	K	W	1.2	0.873			5
408	PHI-4	466	D	R	1.2	0.795			5
409	PHI-4	517	Q	C	1.2	0.789			5
410	PHI-4	256	Q	E	1.2	0.618			5
411	PHI-4	517	Q	N	1.2	0.785			5
412	PHI-4	148	D	Q	1.2	0.754			5
413	PHI-4	517	Q	V	1.2	0.777			5
414	PHI-4	83	E	Q	1.2	0.668			5
415	PHI-4	165	K	F	1.2	0.836			5
416	PHI-4	447	D	F	1.2	0.912			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
417	PHI-4	109	F	A	1.2	0.71			5
418	PHI-4	86	E	Q	1.2	0.565			5
419	PHI-4	61	R	E	1.2	0.567			5
420	PHI-4	151	D	L	1.2	0.891			5
421	PHI-4	305	K	Q	1.1	0.553			5
422	PHI-4	257	Q	S	1.1	0.729			5
423	PHI-4	74	K	R	1.1	0.782			5
424	PHI-4	257	Q	H	1.1	0.731			5
425	PHI-4	447	D	W	1.1	0.933			5
426	PHI-4	402	K	Y	1.1	0.74			5
427	PHI-4	257	Q	W	1.1	0.72			5
428	PHI-4	517	Q	P	1.1	0.831			5
429	PHI-4	331	E	Q	1.1	0.661			5
430	PHI-4	334	G	W	1.1	0.635			5
431	PHI-4	335	S	A	1.1	0.754			5
432	PHI-4	297	E	Q	1.1	0.497			5
433	PHI-4	264	E	Q	1.1	0.658			5
434	PHI-4	447	D	T	1.1	0.767			5
435	PHI-4	229	R	E	1.1	0.494			5
436	PHI-4	298	D	Q	1.1	0.752			5
437	PHI-4	527	Q	F	1.1	0.763			5
438	PHI-4	74	K	M	1.1	0.695			5
439	PHI-4	289	K	C	1.1	0.677			5
440	PHI-4	113	D	E	1.1	0.877			5
441	PHI-4	257	Q	M	1.1	0.651			5
442	PHI-4	207	K	Q	1.1	0.463			5
443	PHI-4	454	R	N	1.1	0.624			5
444	PHI-4	502	R	Q	1.1	0.454			5
445	PHI-4	402	K	T	1.1	0.62			5
446	PHI-4	151	D	R	1.1	0.692			5
447	PHI-4	527	Q	W	1.1	0.616			5
448	PHI-4	109	F	R	1	0.682			5
449	PHI-4	151	D	E	1	0.655			5
450	PHI-4	289	K	G	1	0.617			5
451	PHI-4	74	K	L	1	0.611			5
452	PHI-4	146	R	Q	1	0.424			5
453	PHI-4	113	D	M	1	0.736			5
454	PHI-4	74	K	H	1	0.636			5
455	PHI-4	191	K	Q	1	0.649			5
456	PHI-4	291	E	Q	1	0.424			5
457	PHI-4	148	D	G	1	0.724			5
458	PHI-4	165	K	Y	1	0.595			5
459	PHI-4	74	K	D	1	0.582			5
460	PHI-4	402	K	V	1	0.556			5
461	PHI-4	165	K	R	1	0.575			5
462	PHI-4	497	D	Q	1	0.75			5
463	PHI-4	397	G	A	1	0.568			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
464	PHI-4	289	K	A	1	0.558			5
465	PHI-4	517	Q	E	1	0.509			5
466	PHI-4	360	Q	E	1	0.523			5
467	PHI-4	254	D	K	1	0.75			5
468	PHI-4	402	K	S	1	0.511			5
469	PHI-4	288	D	Q	1	0.75			5
470	PHI-4	454	R	C	1	0.51			5
471	PHI-4	306	Q	E	1	0.547			5
472	PHI-4	82	E	V	1	0.626			5
473	PHI-4	113	D	N	1	0.538			5
474	PHI-4	447	D	M	1	0.498			5
475	PHI-4	447	D	H	1	0.497			5
476	PHI-4	151	D	G	1	0.494			5
477	PHI-4	214	K	E	1	0.339			5
478	PHI-4	148	D	C	1	0.506			5
479	PHI-4	216	E	T	1	0.601			5
480	PHI-4	87	K	E	0.9	0.329			5
481	PHI-4	165	K	N	0.9	0.467			5
482	PHI-4	151	D	K	0.9	0.749			5
483	PHI-4	113	D	S	0.9	0.441			5
484	PHI-4	402	K	D	0.9	0.415			5
485	PHI-4	459	K	R	0.9	0.42		0.24	5
486	PHI-4	74	K	F	0.9	0.427			5
487	PHI-4	361	R	E	0.9	0.304			5
488	PHI-4	165	K	C	0.9	0.421			5
489	PHI-4	72	D	Q	0.9	0.748			5
490	PHI-4	257	Q	V	0.9	0.441			5
491	PHI-4	220	E	K	0.9	0.535			5
492	PHI-4	334	G	H	0.9	0.575			5
493	PHI-4	113	D	H	0.9	0.394			5
494	PHI-4	517	Q	R	0.9	0.421			5
495	PHI-4	436	D	Q	0.9	0.748			5
496	PHI-4	235	R	P	0.9	0.396			5
497	PHI-4	235	R	K	0.9	0.748		-0.01	5
498	PHI-4	79	K	Q	0.9	0.277			5
499	PHI-4	527	Q	Y	0.9	0.414			5
500	PHI-4	113	D	A	0.9	0.326			5
501	PHI-4	216	E	S	0.9	0.501			5
502	PHI-4	257	Q	L	0.9	0.409			5
503	PHI-4	46	E	Q	0.9	0.627			5
504	PHI-4	447	D	N	0.9	0.369			5
505	PHI-4	247	D	E	0.9	0.321			5
506	PHI-4	72	D	K	0.9	0.748			5
507	PHI-4	517	Q	H	0.9	0.404			5
508	PHI-4	74	K	W	0.9	0.361			5
509	PHI-4	442	Q	E	0.9	0.48		0.3	5
510	PHI-4	229	R	Q	0.9	0.258			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
511	PHI-4	527	Q	R	0.9	0.388			5
512	PHI-4	74	K	V	0.9	0.367			5
513	PHI-4	288	D	K	0.9	0.747			5
514	PHI-4	48	D	Q	0.9	0.747			5
515	PHI-4	517	Q	S	0.9	0.378			5
516	PHI-4	402	K	A	0.9	0.333			5
517	PHI-4	216	E	A	0.8	0.452			5
518	PHI-4	216	E	G	0.8	0.45			5
519	PHI-4	113	D	Y	0.8	0.302			5
520	PHI-4	402	K	M	0.8	0.313			5
521	PHI-4	82	E	P	0.8	0.63			5
522	PHI-4	525	Q	E	0.8	0.469			5
523	PHI-4	99	K	R	0.8	0.312			5
524	PHI-4	257	Q	G	0.8	0.35			5
525	PHI-4	48	D	K	0.8	0.746			5
526	PHI-4	305	K	E	0.8	0.226			5
527	PHI-4	216	E	W	0.8	0.417			5
528	PHI-4	207	K	E	0.8	0.22			5
529	PHI-4	447	D	V	0.8	0.266			5
530	PHI-4	196	Q	F	0.8	0.337			5
531	PHI-4	216	E	K	0.8	0.406			5
532	PHI-4	151	D	H	0.8	0.263			5
533	PHI-4	74	K	S	0.8	0.285			5
534	PHI-4	90	Q	K	0.8	0.466			5
535	PHI-4	459	K	G	0.8	0.282			5
536	PHI-4	454	R	E	0.8	0.215			5
537	PHI-4	216	E	R	0.8	0.391		0.04	5
538	PHI-4	74	K	Y	0.8	0.276			5
539	PHI-4	99	K	S	0.8	0.277			5
540	PHI-4	248	R	Q	0.8	0.201			5
541	PHI-4	142	R	Q	0.8	0.193			5
542	PHI-4	82	E	F	0.8	0.615			5
543	PHI-4	461	T	A	0.8	0.245			5
544	PHI-4	33	E	Q	0.8	0.185			5
545	PHI-4	193	D	Q	0.8	0.745			5
546	PHI-4	332	D	N	0.8	0.226			5
547	PHI-4	42	D	K	0.8	0.745			5
548	PHI-4	74	K	A	0.8	0.238			5
549	PHI-4	257	Q	K	0.7	0.461			5
550	PHI-4	148	D	M	0.7	0.117			5
551	PHI-4	527	Q	G	0.7	0.272			5
552	PHI-4	527	Q	V	0.7	0.276			5
553	PHI-4	379	D	Q	0.7	0.745			5
554	PHI-4	146	R	E	0.7	0.607			5
555	PHI-4	527	Q	T	0.7	0.266			5
556	PHI-4	454	R	A	0.7	0.213			5
557	PHI-4	216	E	D	0.7	0.306			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
558	PHI-4	402	K	C	0.7	0.203			5
559	PHI-4	517	Q	T	0.7	0.256			5
560	PHI-4	257	Q	P	0.7	0.256			5
561	PHI-4	220	E	G	0.7	0.3			5
562	PHI-4	99	K	H	0.7	0.2			5
563	PHI-4	55	R	E	0.7	0.153			5
564	PHI-4	340	D	N	0.7	0.179			5
565	PHI-4	466	D	N	0.7	0.185			5
566	PHI-4	257	Q	A	0.7	0.238			5
567	PHI-4	99	K	T	0.7	0.191			5
568	PHI-4	74	K	T	0.7	0.189			5
569	PHI-4	113	D	L	0.7	0.081			5
570	PHI-4	74	K	Q	0.7	0.602			5
571	PHI-4	527	Q	I	0.7	0.238			5
572	PHI-4	99	K	A	0.7	0.186			5
573	PHI-4	517	Q	W	0.7	0.235			5
574	PHI-4	126	E	Q	0.7	0.459		-0.15	5
575	PHI-4	196	Q	G	0.7	0.231			5
576	PHI-4	274	D	K	0.7	0.744			5
577	PHI-4	517	Q	G	0.7	0.229			5
578	PHI-4	99	K	W	0.7	0.175			5
579	PHI-4	394	D	Q	0.7	0.743			5
580	PHI-4	216	E	I	0.7	0.257			5
581	PHI-4	517	Q	M	0.7	0.214			5
582	PHI-4	334	G	S	0.7	0.507			5
583	PHI-4	517	Q	D	0.7	0.207			5
584	PHI-4	196	Q	L	0.7	0.21			5
585	PHI-4	298	D	K	0.6	0.742			5
586	PHI-4	74	K	C	0.6	0.141			5
587	PHI-4	395	D	S	0.6	0.538			5
588	PHI-4	74	K	I	0.6	0.139			5
589	PHI-4	527	Q	M	0.6	0.189			5
590	PHI-4	113	D	Q	0.6	0.742			5
591	PHI-4	340	D	A	0.6	0.142			5
592	PHI-4	216	E	H	0.6	0.216			5
593	PHI-4	402	K	I	0.6	0.13			5
594	PHI-4	235	R	S	0.6	0.123			5
595	PHI-4	75	Q	K	0.6	0.422			5
596	PHI-4	203	E	Q	0.6	0.588		-0.37	5
597	PHI-4	527	Q	A	0.6	0.167			5
598	PHI-4	517	Q	A	0.6	0.163			5
599	PHI-4	361	R	Q	0.6	0.094			5
600	PHI-4	466	D	P	0.6	0.075			5
601	PHI-4	193	D	K	0.6	0.741			5
602	PHI-4	458	D	V	0.6	0.075			5
603	PHI-4	268	D	Q	0.6	0.741			5
604	PHI-4	61	R	Q	0.6	0.086			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
605	PHI-4	151	D	Y	0.6	0.068			5
606	PHI-4	196	Q	Y	0.6	0.146			5
607	PHI-4	196	Q	V	0.5	0.145			5
608	PHI-4	527	Q	D	0.5	0.144			5
609	PHI-4	166	R	E	0.5	0.084			5
610	PHI-4	334	G	A	0.5	0.477			5
611	PHI-4	447	D	Q	0.5	0.74			5
612	PHI-4	196	Q	C	0.5	0.137			5
613	PHI-4	466	D	K	0.5	0.74			5
614	PHI-4	74	K	P	0.5	0.083			5
615	PHI-4	196	Q	M	0.5	0.127			5
616	PHI-4	99	K	D	0.5	0.083			5
617	PHI-4	82	E	D	0.5	0.141			5
618	PHI-4	216	E	M	0.5	0.136			5
619	PHI-4	235	R	H	0.5	0.074			5
620	PHI-4	340	D	P	0.5	0.078			5
621	PHI-4	216	E	L	0.5	0.134			5
622	PHI-4	458	D	G	0.5	0.119			5
623	PHI-4	403	D	S	0.5	0.074			5
624	PHI-4	360	Q	K	0.5	0.335			5
625	PHI-4	216	E	P	0.5	0.127			5
626	PHI-4	257	Q	Y	0.5	0.11			5
627	PHI-4	118	E	Q	0.5	0.061			5
628	PHI-4	379	D	K	0.5	0.739			5
629	PHI-4	402	K	Q	0.5	0.061			5
630	PHI-4	254	D	Q	0.5	0.739			5
631	PHI-4	257	Q	T	0.5	0.107			5
632	PHI-4	458	D	P	0.5	0.04			5
633	PHI-4	466	D	W	0.5	0.036			5
634	PHI-4	403	D	L	0.5	0.037			5
635	PHI-4	196	Q	H	0.5	0.101			5
636	PHI-4	517	Q	L	0.4	0.099			5
637	PHI-4	196	Q	S	0.4	0.099			5
638	PHI-4	257	Q	F	0.4	0.096			5
639	PHI-4	196	Q	T	0.4	0.096			5
640	PHI-4	235	R	G	0.4	0.055			5
641	PHI-4	332	D	S	0.4	0.005			5
642	PHI-4	152	D	Q	0.4	0.738			5
643	PHI-4	368	D	Q	0.4	0.738			5
644	PHI-4	257	Q	R	0.4	0.092			5
645	PHI-4	315	D	K	0.4	0.738			5
646	PHI-4	99	K	P	0.4	0.052			5
647	PHI-4	109	F	P	0.4	0.103			5
648	PHI-4	113	D	T	0.4	0.004			5
649	PHI-4	334	G	M	0.4	0.447			5
650	PHI-4	75	Q	E	0.4	0.315			5
651	PHI-4	274	D	Q	0.4	0.738			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
652	PHI-4	402	K	E	0.4	0.048			5
653	PHI-4	191	K	E	0.4	0.048			5
654	PHI-4	332	D	E	0.4	0.004			5
655	PHI-4	517	Q	Y	0.4	0.085			5
656	PHI-4	340	D	I	0.4	0.025			5
657	PHI-4	459	K	D	0.4	0.05			5
658	PHI-4	76	D	K	0.4	0.738			5
659	PHI-4	216	E	C	0.4	0.093			5
660	PHI-4	340	D	R	0.4	0.048			5
661	PHI-4	466	D	Y	0.4	0.025			5
662	PHI-4	340	D	G	0.4	0.046			5
663	PHI-4	376	D	Q	0.4	0.737			5
664	PHI-4	203	E	C	0.4	0.088			5
665	PHI-4	235	R	N	0.4	0.042			5
666	PHI-4	247	D	S	0.4	0.076			5
667	PHI-4	332	D	F	0.4	0.021			5
668	PHI-4	216	E	N	0.4	0.083			5
669	PHI-4	527	Q	N	0.4	0.076			5
670	PHI-4	340	D	C	0.4	0.043			5
671	PHI-4	38	Q	E	0.4	0.297			5
672	PHI-4	152	D	K	0.4	0.737			5
673	PHI-4	413	Q	E	0.4	0.316			5
674	PHI-4	247	D	T	0.4	0.002			5
675	PHI-4	235	R	V	0.3	0.034			5
676	PHI-4	268	D	K	0.3	0.736			5
677	PHI-4	99	K	Q	0.3	0.034			5
678	PHI-4	235	R	D	0.3	0.032			5
679	PHI-4	394	D	K	0.3	0.736			5
680	PHI-4	458	D	C	0.3	0.001			5
681	PHI-4	340	D	Y	0.3	0.015			5
682	PHI-4	24	D	K	0.3	0.736			5
683	PHI-4	466	D	G	0.3	0.001			5
684	PHI-4	466	D	C	0.3	0.014			5
685	PHI-4	402	K	P	0.3	0.013			5
686	PHI-4	340	D	L	0.3	0.031			5
687	PHI-4	334	G	Y	0.3	0.421			5
688	PHI-4	334	G	Q	0.3	0.421			5
689	PHI-4	334	G	L	0.3	0.42			5
690	PHI-4	398	S	V	0.3	0.013			5
691	PHI-4	334	G	N	0.3	0.42			5
692	PHI-4	398	S	H	0.3	0.012			5
693	PHI-4	113	D	R	0.3	0.001			5
694	PHI-4	235	R	T	0.3	0.026			5
695	PHI-4	334	G	C	0.3	0.417			5
696	PHI-4	248	R	E	0.3	0.028			5
697	PHI-4	235	R	I	0.3	0.025			5
698	PHI-4	398	S	R	0.3	0.011			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
699	PHI-4	334	G	F	0.3	0.455			5
700	PHI-4	113	D	P	0.3	0.001			5
701	PHI-4	109	F	L	0.3	0.052			5
702	PHI-4	203	E	S	0.3	0.412			5
703	PHI-4	109	F	Y	0.3	0.412			5
704	PHI-4	466	D	E	0.3	0.001			5
705	PHI-4	55	R	Q	0.3	0.025			5
706	PHI-4	203	E	V	0.3	0.411			5
707	PHI-4	398	S	L	0.3	0.01			5
708	PHI-4	332	D	G	0.3	0.001			5
709	PHI-4	203	E	G	0.3	0.41			5
710	PHI-4	466	D	I	0.3	0.009			5
711	PHI-4	466	D	A	0.3	0			5
712	PHI-4	235	R	A	0.3	0.021			5
713	PHI-4	340	D	V	0.3	0.022			5
714	PHI-4	466	D	F	0.3	0.009			5
715	PHI-4	99	K	E	0.3	0.023			5
716	PHI-4	188	K	E	0.3	0.022			5
717	PHI-4	235	R	C	0.2	0.019			5
718	PHI-4	235	R	F	0.2	0.018			5
719	PHI-4	340	D	W	0.2	0.02			5
720	PHI-4	235	R	L	0.2	0.018			5
721	PHI-4	113	D	W	0.2	0			5
722	PHI-4	459	K	E	0.2	0.541			5
723	PHI-4	334	G	V	0.2	0.4			5
724	PHI-4	113	D	V	0.2	0			5
725	PHI-4	113	D	G	0.2	0			5
726	PHI-4	235	R	M	0.2	0.015			5
727	PHI-4	403	D	N	0.2	0.012			5
728	PHI-4	203	E	N	0.2	0.396			5
729	PHI-4	340	D	F	0.2	0.006			5
730	PHI-4	340	D	S	0.2	0.005			5
731	PHI-4	247	D	P	0.2	0			5
732	PHI-4	203	E	I	0.2	0.035			5
733	PHI-4	148	D	K	0.2	0.733			5
734	PHI-4	466	D	T	0.2	0			5
735	PHI-4	332	D	H	0.2	0.005			5
736	PHI-4	458	D	A	0.2	0			5
737	PHI-4	235	R	E	0.2	0.016			5
738	PHI-4	340	D	K	0.2	0.733			5
739	PHI-4	466	D	V	0.2	0			5
740	PHI-4	109	F	V	0.2	0.032			5
741	PHI-4	398	S	T	0.2	0.389			5
742	PHI-4	99	K	G	0.2	0.012			5
743	PHI-4	466	D	L	0.2	0			5
744	PHI-4	340	D	T	0.2	0.013			5
745	PHI-4	332	D	A	0.2	0			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
746	PHI-4	403	D	R	0.2	0.013			5
747	PHI-4	113	D	K	0.2	0.733			5
748	PHI-4	403	D	A	0.2	0.013			5
749	PHI-4	109	F	Q	0.2	0.03			5
750	PHI-4	332	D	C	0.2	0			5
751	PHI-4	340	D	M	0.2	0.004			5
752	PHI-4	203	E	R	0.2	0.029			5
753	PHI-4	235	R	Y	0.2	0.011			5
754	PHI-4	196	Q	P	0.2	0.029			5
755	PHI-4	247	D	W	0.2	0			5
756	PHI-4	203	E	A	0.2	0.384			5
757	PHI-4	466	D	H	0.2	0			5
758	PHI-4	203	E	F	0.1	0.027			5
759	PHI-4	449	Q	K	0.1	0.206			5
760	PHI-4	334	G	T	0.1	0.382			5
761	PHI-4	413	Q	K	0.1	0.223			5
762	PHI-4	38	Q	K	0.1	0.242			5
763	PHI-4	82	E	R	0.1	0.381			5
764	PHI-4	340	D	H	0.1	0.003			5
765	PHI-4	235	R	W	0.1	0.009			5
766	PHI-4	398	S	P	0.1	0.026			5
767	PHI-4	458	D	S	0.1	0		-0.49	5
768	PHI-4	247	D	N	0.1	0			5
769	PHI-4	466	D	S	0.1	0			5
770	PHI-4	395	D	I	0.1	0.379			5
771	PHI-4	466	D	Q	0.1	0.732			5
772	PHI-4	220	E	P	0.1	0.01			5
773	PHI-4	340	D	Q	0.1	0.732			5
774	PHI-4	247	D	G	0.1	0			5
775	PHI-4	203	E	Y	0.1	0.025			5
776	PHI-4	334	G	E	0.1	0.378			5
777	PHI-4	247	D	A	0.1	0.025			5
778	PHI-4	458	D	F	0.1	0.003			5
779	PHI-4	403	D	C	0.1	0			5
780	PHI-4	403	D	I	0.1	0.003			5
781	PHI-4	403	D	V	0.1	0.009			5
782	PHI-4	403	D	G	0.1	0.009			5
783	PHI-4	203	E	K	0.1	0.373			5
784	PHI-4	398	S	A	0.1	0.372			5
785	PHI-4	340	D	E	0.1	0.008			5
786	PHI-4	82	E	H	0.1	0.371			5
787	PHI-4	203	E	W	0.1	0.37			5
788	PHI-4	447	D	R	0.1	0			5
789	PHI-4	109	F	W	0.1	0.37			5
790	PHI-4	247	D	V	0.1	0			5
791	PHI-4	203	E	L	0.1	0.369			5
792	PHI-4	332	D	T	0.1	0			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
793	PHI-4	332	D	I	0.1	0.002			5
794	PHI-4	82	E	G	0.1	0.019			5
795	PHI-4	247	D	L	0.1	0			5
796	PHI-4	368	D	K	0.1	0.731			5
797	PHI-4	82	E	M	0.1	0.018			5
798	PHI-4	247	D	H	0.1	0			5
799	PHI-4	458	D	N	0.1	0.019			5
800	PHI-4	235	R	Q	0.1	0.008			5
801	PHI-4	398	S	D	0.1	0.017			5
802	PHI-4	332	D	Y	0.1	0			5
803	PHI-4	247	D	F	0.1	0.018			5
804	PHI-4	332	D	V	0.1	0			5
805	PHI-4	395	D	A	0.1	0.017			5
806	PHI-4	334	G	P	0.1	0.362			5
807	PHI-4	53	E	Q	0.1	0.008			5
808	PHI-4	97	R	T	0.1	0.016			5
809	PHI-4	458	D	I	0.1	0			5
810	PHI-4	398	S	N	0.1	0.016			5
811	PHI-4	109	F	H	0.1	0.361			5
812	PHI-4	403	D	P	0	0			5
813	PHI-4	497	D	K	0	0.73			5
814	PHI-4	458	D	M	0	0.017			5
815	PHI-4	458	D	T	0	0.017			5
816	PHI-4	447	D	A	0	0			5
817	PHI-4	395	D	N	0	0.404	180	-0.91	5
818	PHI-4	247	D	R	0	0			5
819	PHI-4	82	E	T	0	0.36			5
820	PHI-4	395	D	Y	0	0.36			5
821	PHI-4	398	S	E	0	0.36			5
822	PHI-4	332	D	Q	0	0.73			5
823	PHI-4	403	D	M	0	0			5
824	PHI-4	247	D	Q	0	0.73			5
825	PHI-4	398	S	I	0	0.015			5
826	PHI-4	458	D	Y	0	0.016			5
827	PHI-4	398	S	M	0	0.015			5
828	PHI-4	403	D	H	0	0			5
829	PHI-4	82	E	N	0	0.358			5
830	PHI-4	403	D	T	0	0			5
831	PHI-4	247	D	M	0	0.016			5
832	PHI-4	395	D	R	0	0.015			5
833	PHI-4	398	S	Y	0	0.401			5
834	PHI-4	395	D	F	0	0.357			5
835	PHI-4	395	D	M	0	0.357			5
836	PHI-4	395	D	P	0	0.357			5
837	PHI-4	97	R	A	0	0.014			5
838	PHI-4	395	D	Q	0	0.73		-0.82	5
839	PHI-4	332	D	L	0	0			5

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
840	PHI-4	395	D	W	0	0.357			5
841	PHI-4	332	D	P	0	0			5
842	PHI-4	403	D	Q	0	0.73			5
843	PHI-4	458	D	L	0	0			5
844	PHI-4	398	S	K	0	0.356			5
845	PHI-4	458	D	E	0	0.015			5
846	PHI-4	203	E	P	0	0.014			5
847	PHI-4	395	D	T	0	0.356			5
848	PHI-4	82	E	C	0	0.356			5
849	PHI-4	395	D	G	0	0.355			5
850	PHI-4	97	R	V	0	0.014			5
851	PHI-4	332	D	W	0	0			5
852	PHI-4	395	D	L	0	0.355			5
853	PHI-4	332	D	M	0	0			5
854	PHI-4	398	S	W	0	0.014			5
855	PHI-4	395	D	E	0	0.355			5
856	PHI-4	458	D	R	0	0.015			5
857	PHI-4	321	D	K	0	0.729			5
858	PHI-4	82	E	K	0	0.013			5
859	PHI-4	82	E	A	0	0.353			5
860	PHI-4	332	D	R	0	0			5
861	PHI-4	458	D	H	0	0			5
862	PHI-4	398	S	F	0	0.353			5
863	PHI-4	395	D	K	0	0.729			5
864	PHI-4	395	D	H	0	0.353			5
865	PHI-4	203	E	M	0	0.353			5
866	PHI-4	458	D	W	0	0.014			5
867	PHI-4	403	D	K	0	0.729			5
868	PHI-4	247	D	K	0	0.729			5
869	PHI-4	376	D	K	0	0.729			5
870	PHI-4	321	D	Q	0	0.729			5
871	PHI-4	332	D	K	0	0.729			5
872	PHI-4	315	D	Q					5
873	SFR16	293	R	E	19.5	0.045			6
874	SFR16	416	R	E	18	0.039			6
875	SFR16	328	K	E	17.8	0.06			6
876	SFR16	500	R	Q	17.1	0.061			6
877	SFR16	452	Q	K	8.8	0.014			6
878	SFR16	293	R	Q	6.9	0			6
879	SFR16	150	R	Q	4.2	0			6
880	SFR16	471	Q	K	4	0.14			6
881	SFR16	261	Q	E	3.1	0.011			6
882	SFR16	520	K	Q	3.1	0.006			6
883	SFR16	471	Q	E	3.1	0.01			6
884	SFR16	147	R	E	2.7	0.038			6
885	SFR16	520	K	E	2.6	0.888			6
886	SFR16	509	K	Q	2.5	0.053			6

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
887	SFR16	162	E	Q	2.5	0.058			6
888	SFR16	281	Q	K	2.4	0.415			6
889	SFR16	452	Q	E	2.4	0.13			6
890	SFR16	313	K	Q	2.3	0.204			6
891	SFR16	328	K	Q	2.2	0.194			6
892	SFR16	391	R	E	2.1	0.812			6
893	SFR16	281	Q	E	2.1	0.535			6
894	SFR16	391	R	Q	2.1	0.253			6
895	SFR16	174	E	Q	2	0.138			6
896	SFR16	391	R	D	1.8	0.768			6
897	SFR16	147	R	Q	1.5	0.712			6
898	SFR16	261	Q	K	1.4	0.739			6
899	SFR16	313	K	R	1.4	0.758			6
900	SFR16	316	K	Q	1.3	0.685			6
901	SFR16	150	R	E	1.2	0.677			6
902	SFR16	500	R	K	1.2	0.754			6
903	SFR16	509	K	E	1.1	0.537			6
904	SFR16	416	R	Q	1	0.416			6
905	SFR16	316	K	E	0.5	0.061			6
906	SFR16	242	K	R	0.4	0.738			6
907	SFR16	313	K	E	0.1	0.013			6
908	SFR16	242	K	E	0.1	0.011			6
909	SFR16	242	K	Q	0.1	0.011			6
910	SFR16	500	R	E	0.1	0.01			6
911	PSR3	339	E	N	343.6	0			7
912	PSR3	401	S	H	32.8	0			7
913	PSR3	401	S	P	21.4	0			7
914	PSR3	339	E	I	14.8	0			7
915	PSR3	465	K	N	14.7	0			7
916	PSR3	402	K	R	11.7	0			7
917	PSR3	402	K	G	8.3	0			7
918	PSR3	333	S	G	7.4	0			7
919	PSR3	464	R	D	7.1	0			7
920	PSR3	402	K	W	6.8	0			7
921	PSR3	401	S	G	6.2	0			7
922	PSR3	465	K	V	5	0.002			7
923	PSR3	402	K	H	4.7	0			7
924	PSR3	396	A	L	4.6	0			7
925	PSR3	338	S	H	4.6	0.001			7
926	PSR3	464	R	K	4.3	0			7
927	PSR3	465	K	M	4.3	0			7
928	PSR3	333	S	K	4.3	0			7
929	PSR3	464	R	A	4	0			7
930	PSR3	396	A	K	3.9	0			7
931	PSR3	338	S	V	3.2	0			7
932	PSR3	333	S	V	3.1	0.005			7
933	PSR3	401	S	K	3.1	0			7

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
934	PSR3	339	E	P	3	0.026			7
935	PSR3	465	K	P	2.9	0.071			7
936	PSR3	396	A	N	2.8	0.042			7
937	PSR3	402	K	T	2.8	0.06			7
938	PSR3	464	R	S	2.7	0.031			7
939	PSR3	463	A	S	2.7	0.083			7
940	PSR3	459	K	Y	2.6	0.051			7
941	PSR3	333	S	N	2.6	0.051			7
942	PSR3	396	A	G	2.5	0.057			7
943	PSR3	464	R	G	2.5	0.079			7
944	PSR3	338	S	K	2.5	0.008			7
945	PSR3	402	K	N	2.3	0.127			7
946	PSR3	465	K	G	2.3	0.268			7
947	PSR3	402	K	Y	2.2	0.158			7
948	PSR3	401	S	V	2.2	0.024			7
949	PSR3	464	R	H	2.2	0.175			7
950	PSR3	465	K	R	2.1	0.246			7
951	PSR3	401	S	R	2.1	0.206			7
952	PSR3	402	K	M	2.1	0.242			7
953	PSR3	338	S	A	2	0.376			7
954	PSR3	464	R	N	2	0.293			7
955	PSR3	459	K	E	2	0.298			7
956	PSR3	465	K	T	2	0.401			7
957	PSR3	333	S	A	2	0.082			7
958	PSR3	338	S	G	1.9	0.456			7
959	PSR3	339	E	M	1.9	0.368			7
960	PSR3	396	A	I	1.9	0.479			7
961	PSR3	338	S	T	1.8	0.182			7
962	PSR3	402	K	P	1.8	0.587			7
963	PSR3	459	K	P	1.8	0.466			7
964	PSR3	333	S	H	1.8	0.561			7
965	PSR3	396	A	M	1.8	0.213			7
966	PSR3	333	S	Q	1.7	0.254			7
967	PSR3	339	E	S	1.6	0.359			7
968	PSR3	401	S	I	1.6	0.753			7
969	PSR3	338	S	I	1.6	0.756			7
970	PSR3	401	S	N	1.6	0.38			7
971	PSR3	333	S	T	1.5	0.833			7
972	PSR3	339	E	A	1.5	0.837			7
973	PSR3	339	E	C	1.5	0.485			7
974	PSR3	401	S	F	1.5	0.846			7
975	PSR3	463	A	G	1.5	0.761			7
976	PSR3	464	R	Q	1.5	0.931			7
977	PSR3	396	A	R	1.5	0.538			7
978	PSR3	339	E	F	1.5	0.805			7
979	PSR3	338	S	E	1.5	0.907			7
980	PSR3	339	E	V	1.4	0.674			7

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
981	PSR3	465	K	H	1.4	0.933			7
982	PSR3	465	K	C	1.4	0.935			7
983	PSR3	339	E	L	1.4	0.741			7
984	PSR3	401	S	T	1.4	0.753			7
985	PSR3	401	S	A	1.4	0.962			7
986	PSR3	459	K	R	1.4	0.962			7
987	PSR3	333	S	D	1.4	0.972			7
988	PSR3	339	E	Q	1.3	0.842			7
989	PSR3	464	R	T	1.3	0.975			7
990	PSR3	338	S	M	1.3	0.857			7
991	PSR3	338	S	R	1.3	0.89			7
992	PSR3	333	S	I	1.3	0.904			7
993	PSR3	465	K	W	1.3	0.864			7
994	PSR3	338	S	P	1.3	0.921			7
995	PSR3	333	S	L	1.3	0.847			7
996	PSR3	459	K	H	1.3	0.824			7
997	PSR3	338	S	D	1.3	0.979			7
998	PSR3	465	K	F	1.2	0.881			7
999	PSR3	402	K	F	1.2	0.793			7
1000	PSR3	339	E	D	1.2	0.795			7
1001	PSR3	337	A	V	1.2	0.899			7
1002	PSR3	338	S	N	1.2	0.883			7
1003	PSR3	465	K	A	1.2	0.796			7
1004	PSR3	396	A	Y	1.1	0.747			7
1005	PSR3	401	S	D	1.1	0.701			7
1006	PSR3	333	S	C	1.1	0.741			7
1007	PSR3	339	E	W	1.1	0.681			7
1008	PSR3	333	S	E	1.1	0.658			7
1009	PSR3	337	A	G	1.1	0.656			7
1010	PSR3	459	K	W	1.1	0.7			7
1011	PSR3	401	S	M	1.1	0.694			7
1012	PSR3	401	S	Q	1.1	0.635			7
1013	PSR3	465	K	L	1.1	0.634			7
1014	PSR3	396	A	Q	1.1	0.697			7
1015	PSR3	402	K	A	1.1	0.629			7
1016	PSR3	401	S	E	1.1	0.674			7
1017	PSR3	333	S	R	1.1	0.649			7
1018	PSR3	339	E	H	1.1	0.642			7
1019	PSR3	338	S	L	1.1	0.636			7
1020	PSR3	396	A	H	1	0.562			7
1021	PSR3	464	R	F	1	0.587			7
1022	PSR3	402	K	L	1	0.549			7
1023	PSR3	339	E	R	1	0.549			7
1024	PSR3	465	K	I	1	0.575			7
1025	PSR3	464	R	I	1	0.553			7
1026	PSR3	402	K	D	1	0.54			7
1027	PSR3	333	S	F	0.9	0.509			7

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
1028	PSR3	465	K	Q	0.9	0.634			7
1029	PSR3	465	K	D	0.9	0.488			7
1030	PSR3	396	A	T	0.9	0.409			7
1031	PSR3	465	K	Y	0.9	0.479			7
1032	PSR3	464	R	V	0.9	0.467			7
1033	PSR3	338	S	Q	0.9	0.382			7
1034	PSR3	401	S	L	0.9	0.434			7
1035	PSR3	459	K	V	0.9	0.457			7
1036	PSR3	464	R	L	0.9	0.447			7
1037	PSR3	465	K	S	0.9	0.409			7
1038	PSR3	339	E	Y	0.9	0.407			7
1039	PSR3	459	K	Q	0.9	0.422			7
1040	PSR3	333	S	M	0.8	0.395			7
1041	PSR3	463	A	I	0.8	0.411			7
1042	PSR3	401	S	Y	0.8	0.318			7
1043	PSR3	396	A	V	0.8	0.287			7
1044	PSR3	463	A	C	0.8	0.382			7
1045	PSR3	339	E	G	0.8	0.27			7
1046	PSR3	400	K	D	0.8	0.27			7
1047	PSR3	400	K	Y	0.8	0.267			7
1048	PSR3	338	S	F	0.8	0.259			7
1049	PSR3	459	K	C	0.8	0.361			7
1050	PSR3	396	A	S	0.8	0.249			7
1051	PSR3	400	K	H	0.8	0.23			7
1052	PSR3	400	K	R	0.8	0.228			7
1053	PSR3	459	K	T	0.7	0.328			7
1054	PSR3	402	K	I	0.7	0.308			7
1055	PSR3	464	R	Y	0.7	0.314			7
1056	PSR3	464	R	P	0.7	0.298			7
1057	PSR3	339	E	K	0.7	0.188			7
1058	PSR3	402	K	Q	0.7	0.273			7
1059	PSR3	333	S	P	0.7	0.169			7
1060	PSR3	400	K	N	0.7	0.167			7
1061	PSR3	333	S	Y	0.7	0.256			7
1062	PSR3	400	K	L	0.6	0.242			7
1063	PSR3	459	K	S	0.6	0.24			7
1064	PSR3	333	S	W	0.6	0.246			7
1065	PSR3	339	E	T	0.6	0.237			7
1066	PSR3	400	K	M	0.6	0.13			7
1067	PSR3	396	A	F	0.6	0.235			7
1068	PSR3	401	S	W	0.6	0.233			7
1069	PSR3	459	K	G	0.6	0.233			7
1070	PSR3	402	K	C	0.6	0.226			7
1071	PSR3	465	K	E	0.6	0.099			7
1072	PSR3	400	K	T	0.6	0.108			7
1073	PSR3	396	A	P	0.6	0.208			7
1074	PSR3	338	S	Y	0.6	0.206			7

MUT ID NO:	Back-bone	Position	Ref A.A.	Substitution	FAE	P-value	EC50 (ppm)	Deviation	Example #
1075	PSR3	459	K	I	0.6	0.203			7
1076	PSR3	459	K	A	0.6	0.201			7
1077	PSR3	396	A	E	0.6	0.198			7
1078	PSR3	459	K	F	0.6	0.198			7
1079	PSR3	400	K	S	0.5	0.195			7
1080	PSR3	338	S	W	0.5	0.188			7
1081	PSR3	463	A	V	0.5	0.189			7
1082	PSR3	459	K	L	0.5	0.175			7
1083	PSR3	400	K	A	0.5	0.064			7
1084	PSR3	400	K	Q	0.5	0.056			7
1085	PSR3	402	K	V	0.4	0.146			7
1086	PSR3	459	K	N	0.4	0.14			7
1087	PSR3	400	K	I	0.4	0.047			7
1088	PSR3	400	K	C	0.4	0.13			7
1089	PSR3	464	R	W	0.4	0.129			7
1090	PSR3	402	K	S	0.4	0.124			7
1091	PSR3	459	K	M	0.4	0.117			7
1092	PSR3	396	A	C	0.4	0.112			7
1093	PSR3	396	A	D	0.4	0.032			7
1094	PSR3	402	K	E	0.4	0.114			7
1095	PSR3	400	K	V	0.3	0.085			7
1096	PSR3	459	K	D	0.3	0.079			7
1097	PSR3	400	K	E	0.2	0.02			7
1098	PSR3	463	A	F	0.2	0.066			7
1099	PSR3	463	A	L	0.2	0.065			7
1100	PSR3	337	A	Q	0.2	0.01			7
1101	PSR3	463	A	M	0.2	0.059			7
1102	PSR3	337	A	F	0.2	0.009			7
1103	PSR3	464	R	E	0.2	0.014			7
1104	PSR3	400	K	G	0.2	0.059			7
1105	PSR3	338	S	C	0.2	0.054			7
1106	PSR3	337	A	S	0.2	0.008			7
1107	PSR3	337	A	T	0.2	0.008			7
1108	PSR3	463	A	T	0.2	0.057			7
1109	PSR3	463	A	N	0.1	0.052			7
1110	PSR3	401	S	C	0.1	0.008			7
1111	PSR3	400	K	F	0.1	0.049			7
1112	PSR3	400	K	W	0.1	0.049			7
1113	PSR3	337	A	C	0.1	0.052			7
1114	PSR3	337	A	W	0.1	0.047			7
1115	PSR3	396	A	W	0.1	0.051			7
1116	PSR3	463	A	D	0.1	0.046			7
1117	PSR3	337	A	Y	0.1	0.006			7
1118	PSR3	463	A	Q	0.1	0.043			7
1119	PSR3	463	A	W	0.1	0.042			7
1120	PSR3	337	A	P	0.1	0.005			7
1121	PSR3	337	A	H	0.1	0.004			7

MUT ID NO:	Back-bone	Posit-ion	Ref A.A.	Substi-tution	FAE	P-value	EC50 (ppm)	Devia-tion	Example #
1122	PSR3	337	A	K	0.1	0.043			7
1123	PSR3	337	A	L	0.1	0.004			7
1124	PSR3	463	A	H	0.1	0.038			7
1125	PSR3	337	A	I	0.1	0.004			7
1126	PSR3	337	A	D	0.1	0.004			7
1127	PSR3	337	A	N	0.1	0.004			7
1128	PSR3	463	A	K	0.1	0.037			7
1129	PSR3	337	A	M	0	0.003			7
1130	PSR3	400	K	P	0	0.037			7
1131	PSR3	463	A	R	0	0.032			7
1132	PSR3	463	A	Y	0	0.031			7
1133	PSR3	337	A	R	0	0.035			7
1134	PSR3	463	A	P	0	0.031			7
1135	PSR3	337	A	E	0	0.035			7

Table 9. The definitions of the column headings are as follows: “SEQ ID NO:”, a unique identifier for each DNA or amino acid sequence; “Trivial Name”, a trivial but unique name for each DNA or protein sequence; “Reference Protein”, the SEQ ID NO corresponding to the reference protein used in the FAE assays for each variant; “FAE”, the arithmetic Mean FAE Index as further defined in Example 3; “p-value” the calculated p value associated with the hypothesis that the variant polypeptide is significantly different than the reference protein used in that particular FAE assay, as defined further in Example 3; “EC50 Fold”, the fold increase in potency as defined by ratio of reference EC50 to variant EC50 is calculated in each of multiple experiments and then the mean of these independently measured numbers is reported as “EC50 Fold”; “Deviation”, Mean Deviation Score as defined in Example 4; “Example #”, the example number corresponding to the creation of the variant. The reference protein against which the variant protein is compared is: (MUT IDs: 1-872 and 911-1135) used SEQ ID NO: 6 as the reference protein; (MUT IDs: 873-910) used SEQ ID NO: 8 as the reference protein.

Table 9

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
51	SFR5-014	6.2	0			8
52	SFR5-001	3.8	0			8
53	SFR5-007	3.6	0			8
54	SFR5-010	3.3	0			8
55	SFR5-009	3.3	0			8
56	SFR5-003	3.2	0			8
57	SFR5-017	3.1	0			8

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
58	SFR5-006	2.4	0.015			8
59	SFR5-013	2.2	0.046			8
60	SFR5-016	2.1	0.057			8
61	SFR5-004	1.9	0.139			8
62	SFR5-005	1.9	0.146			8
63	SFR5-020	1.7	0.328			8
64	SFR5-002	1.5	0.487			8
65	SFR5-008	1.4	0.927			8
66	SFR5-015	1.3	0.753			8
67	SFR09-007			3.3		8
68	SFR09-005			2.4		8
69	SFR09-002			2.2		8
70	SFR09-004			2.1		8
71	SFR09-006			2		8
72	SFR09-003			2		8
73	SFR10-032	17.1	0			8
74	SFR10-042	12	0			8
75	SFR10-72	8.9	0.003			8
76	SFR10-056	8.4	0			8
77	SFR10-036	8.2	0			8
78	SFR10-039	7.2	0			8
79	SFR10-82	6.7	0			8
80	SFR10-045	6.6	0			8
81	SFR10-87	6.4	0			8
82	SFR10-060	6.3	0			8
83	SFR10-052	5.3	0			8
84	SFR10-059	4.9	0			8
85	SFR10-84	4.6	0.025			8
86	SFR10-031	4.4	0			8
87	SFR10-054	4.2	0			8
88	SFR10-064	3.9	0			8
89	SFR10-76	3.8	0			8
90	SFR10-008	3.7	0			8
91	SFR10-035	3.7	0			8
92	SFR10-015	3.5	0			8
93	SFR10-71	3.5	0			8
94	SFR10-74	3.4	0			8
95	SFR10-047	3.4	0			8
96	SFR10-043	3.4	0.006			8
97	SFR10-055	2.9	0			8
98	SFR10-065	2.8	0			8
99	SFR10-041	2.8	0			8
100	SFR10-83	2.8	0.006			8
101	SFR10-002	2.7	0.001			8
102	SFR10-046	2.5	0.002			8
103	SFR10-037	2.5	0.002			8
104	SFR10-048	2.5	0.002			8

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
105	SFR10-78	2.4	0			8
106	SFR10-050	2.4	0			8
107	SFR10-058	2.4	0.017			8
108	SFR10-020	2.4	0.009			8
109	SFR10-057	2.3	0.018			8
110	SFR10-040	2.3	0.008			8
111	SFR10-75	2.3	0			8
112	SFR10-73	2.3	0			8
113	SFR10-049	2.2	0.044			8
114	SFR10-003	2.2	0.022			8
115	SFR10-038	2.2	0.015			8
116	SFR10-024	2.2	0.022			8
117	SFR10-79	2.1	0.006			8
118	SFR10-81	2	0.001			8
119	SFR10-77	2	0.006			8
120	SFR10-88	1.9	0.205			8
121	SFR10-051	1.9	0.109			8
122	SFR10-016	1.9	0.12			8
123	SFR10-025	1.8	0.152			8
124	SFR10-017	1.6	0.364			8
125	SFR10-018	1.6	0.393			8
126	SFR10-004	1.6	0.39			8
127	SFR10-029	1.6	0.395			8
128	SFR10-053	1.6	0.434			8
129	SFR10-006	1.5	0.637			8
130	SFR10-80	1.4	0.113			8
131	SFR10-021	1.4	0.697			8
132	SFR10-009	1.4	0.726			8
133	SFR10-007	1.4	0.728			8
134	SFR10-030	1.3	0.965			8
135	SFR10-014	1.2	0.79			8
136	SFR10-044	1.2	0.8			8
137	SFR10-89	1.1	0.673			8
138	SFR10-013	1.1	0.596			8
139	SFR10-011	1.1	0.588			8
140	SFR10-010	1	0.457			8
141	SFR10-023	0.9	0.356			8
142	SFR10-022	0.9	0.288			8
143	SFR10-027	0.9	0.264			8
144	SFR10-005	0.8	0.233			8
145	SFR10-019	0.7	0.148			8
146	SFR10-028	0.7	0.093			8
147	SFR10-026	0.6	0.079			8
148	SFR11-001	27.4	0			8
149	SFR11-012	14.8	0			8
150	SFR11-005	12.8	0			8
151	SFR11-014	9.9	0			8

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
152	SFR11-015	8.2	0			8
153	SFR11-010	8.1	0			8
154	SFR11-013	5.5	0			8
155	SFR11-004	4.3	0			8
156	SFR11-009	4	0			8
157	SFR11-002	4	0			8
158	SFR11-011	3.9	0			8
159	SFR11-007	3.8	0.007			8
160	SFR11-008	3.1	0			8
161	SFR11-006	2.3	0.012			8
162	SFR12-028			7.6		8
163	SFR12-022			5.7		8
164	SFR12-004			5		8
165	SFR12-006			4.9		8
166	SFR12-015			4.6		8
167	SFR12-001			4.6		8
168	SFR12-014			4.6		8
169	SFR12-002			4.5		8
170	SFR12-005			4.4		8
171	SFR12-017			4.2		8
172	SFR12-018			3.9		8
173	SFR12-003			3.9		8
174	SFR12-032			3.5		8
175	SFR12-016			3.2		8
176	SFR12-029			3.2		8
177	SFR12-011			3.2		8
178	SFR12-007			3		8
179	SFR12-010			3		8
180	SFR12-012			2.8		8
181	SFR12-009			2.8		8
182	SFR12-031			2.3		8
183	SFR13-035	9.3	0			8
184	SFR13-018	6.1	0			8
185	SFR13-039	5.4	0			8
186	SFR13-008	5.3	0			8
187	SFR13-012	5.3	0			8
188	SFR13-036	5.2	0			8
189	SFR13-009	4.8	0			8
190	SFR13-025	4.8	0			8
191	SFR13-033	4.7	0			8
192	SFR13-038	4.5	0			8
193	SFR13-003	4.5	0			8
194	SFR13-021	4.3	0			8
195	SFR13-030	4.3	0			8
196	SFR13-017	4.2	0			8
197	SFR13-004	4.2	0			8
198	SFR13-006	4.1	0			8

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
199	SFR13-026	4.1	0			8
200	SFR13-016	4.1	0			8
201	SFR13-031	4.1	0			8
202	SFR13-034	3.7	0			8
203	SFR13-007	3.7	0			8
204	SFR13-029	3.7	0			8
205	SFR13-024	3.6	0			8
206	SFR13-020	3.4	0			8
207	SFR13-001	3	0			8
208	SFR13-005	3	0			8
209	SFR13-037	2.8	0			8
210	SFR13-032	2.8	0			8
211	SFR13-027	2.5	0			8
212	SFR13-028	2.4	0			8
213	SFR13-010	2.4	0			8
214	SFR13-011	2.2	0			8
215	SFR13-019	2.2	0			8
216	SFR13-023	2	0			8
217	SFR13-022	1.3	0.322			8
218	SFR14-004			3.3		8
219	SFR14-007			3.2		8
220	SFR14-008			3.2		8
221	SFR14-005			2.8		8
222	SFR14-001			2.8		8
223	SFR14-002			2		8
224	SFR14-003			1.3		8
225	SFR17-013	20.9	0.002			8
226	SFR17-019	20.1	0			8
227	SFR17-014	19.5	0			8
228	SFR17-011	19	0			8
229	SFR17-005	17.1	0			8
230	SFR17-018	16.5	0			8
231	SFR17-009	13.5	0			8
232	SFR17-006	13.1	0			8
233	SFR17-016	12.9	0			8
234	SFR17-012	11.7	0			8
235	SFR17-004	11.3	0			8
236	SFR17-017	10.5	0			8
237	SFR17-003	9.7	0			8
238	SFR17-001	8.7	0.01			8
239	SFR17-015	8.2	0			8
240	SFR17-002	7.6	0			8
241	SFR17-007	5.5	0			8
242	SFR17-008	4.7	0			8
243	P053168-D-01_S03721995				0.39	9
244	P053168-A-10_S03723031				0.34	9
245	P053168-D-06_S03722432				0.26	9

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
246	P053168-F-07_S03722451				0.25	9
247	P053168-F-10_S03723024				0.24	9
248	P053169-E-10_S03723571				0.23	9
249	P053168-B-03_S03722281				0.19	9
250	P053168-E-10_S03722994				0.19	9
251	P053168-G-09_S03723000				0.17	9
252	P053168-H-05_S03722286				0.16	9
253	P053168-F-04_S03722292				0.15	9
254	P053168-A-03_S03722326				0.14	9
255	P053168-C-04_S03722283				0.14	9
256	P053168-D-07_S03722516				0.14	9
257	P053168-A-12_S03722983				0.14	9
258	P053169-H-01_S03723182				0.13	9
259	P053168-H-11_S03723019				0.13	9
260	P053168-A-02_S03723761				0.12	9
261	P053168-C-10_S03722993				0.12	9
262	P053168-A-07_S03722449				0.12	9
263	P053168-B-08_S03722529				0.1	9
264	P053168-B-12_S03722998				0.09	9
265	P053169-E-12_S03723656				0.08	9
266	P053168-A-08_S03722523				0.07	9
267	P053168-B-04_S03722320				0.07	9
268	P053168-B-10_S03722972				0.07	9
269	P053169-G-11_S03723594				0.07	9
270	P053168-D-11_S03722996				0.06	9
271	P053169-G-01_S03723175				0.06	9
272	P053168-E-05_S03722315				0.06	9
273	P053168-F-08_S03722955				0.05	9
274	P053168-D-05_S03722300				0.04	9
275	P053169-H-02_S03723149				0.03	9
276	P053168-C-06_S03722354				0.02	9
277	P053169-D-06_S03723260				0.02	9
278	P053169-B-08_S03723469				0.02	9
279	P053169-G-08_S03723462				0.01	9
280	P053168-G-04_S03722314				0.01	9
281	P053168-H-08_S03722875				0.01	9
282	P053169-B-12_S03723595				0.01	9
283	P053169-D-01_S03723219				-0.01	9
284	P053168-G-05_S03722346				-0.01	9
285	P053168-A-09_S03722919				-0.01	9
286	P053168-A-04_S03722290				-0.01	9
287	P053169-F-11_S03723652				-0.02	9
288	P053169-H-12_S03723599				-0.03	9
289	P053168-F-01_S03723758				-0.03	9
290	P053169-H-04_S03723234				-0.04	9
291	P053168-A-06_S03722294				-0.05	9
292	P053169-G-04_S03723174				-0.05	9

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
293	P053169-A-11_S03723493				-0.06	9
294	P053169-B-09_S03723434				-0.06	9
295	P053169-G-12_S03723613				-0.06	9
296	P053168-B-02_S03723770				-0.06	9
297	P053168-G-02_S03722325				-0.06	9
298	P053168-H-06_S03722508				-0.07	9
299	P053168-C-09_S03723029				-0.08	9
300	P053169-E-11_S03723504				-0.08	9
301	P053168-F-06_S03722352				-0.08	9
302	P053168-D-09_S03722970				-0.08	9
303	P053169-D-08_S03723410				-0.08	9
304	P053169-F-03_S03723179				-0.08	9
305	P053169-G-10_S03723505				-0.09	9
306	P053168-D-12_S03723072				-0.09	9
307	P053169-H-09_S03723489				-0.09	9
308	P053169-A-02_S03723190				-0.09	9
309	P053169-B-02_S03723176				-0.1	9
310	P053169-B-05_S03723255				-0.1	9
311	P053169-H-06_S03723276				-0.11	9
312	P053169-A-03_S03723185				-0.11	9
313	P053168-D-10_S03723046				-0.11	9
314	P053168-E-11_S03722989				-0.12	9
315	P053169-F-10_S03723498				-0.12	9
316	P053169-G-05_S03723273				-0.12	9
317	P053168-H-09_S03722964				-0.12	9
318	P053168-E-07_S03722450				-0.12	9
319	P053168-G-01_S03723771				-0.12	9
320	P053169-F-09_S03723457				-0.13	9
321	P053169-B-01_S03723205				-0.13	9
322	P053168-F-09_S03723030				-0.14	9
323	P053169-C-10_S03723416				-0.14	9
324	P053169-A-09_S03723486				-0.14	9
325	P053168-C-11_S03723040				-0.15	9
326	P053169-D-09_S03723487				-0.15	9
327	P053169-A-06_S03723289				-0.16	9
328	P053169-C-04_S03723203				-0.17	9
329	P053169-C-12_S03723581				-0.17	9
330	P053169-C-03_S03723201				-0.17	9
331	P053169-C-01_S03723212				-0.17	9
332	P053169-F-05_S03723296				-0.18	9
333	P053169-B-04_S03723195				-0.18	9
334	P053169-C-09_S03723456				-0.18	9
335	P053169-E-05_S03723236				-0.18	9
336	P053169-D-02_S03723169				-0.19	9
337	P053169-D-11_S03723568				-0.19	9
338	P053169-E-04_S03723145				-0.19	9
339	P053169-F-01_S03723167				-0.19	9

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
340	P053169-B-07_S03723314				-0.19	9
341	P053168-G-12_S03723112				-0.21	9
342	P053168-H-03_S03722342				-0.21	9
343	P053169-E-01_S03723146				-0.22	9
344	P053169-B-03_S03723193				-0.22	9
345	P053168-C-02_S03723834				-0.22	9
346	P053169-G-06_S03723253				-0.22	9
347	P053169-C-02_S03723148				-0.23	9
348	P053169-B-06_S03723231				-0.23	9
349	P053169-G-09_S03723466				-0.24	9
350	P053169-H-11_S03723608				-0.24	9
351	P053168-H-02_S03722266				-0.24	9
352	P053169-H-05_S03723280				-0.25	9
353	P053169-E-02_S03723214				-0.27	9
354	P053168-H-12_S03723098				-0.27	9
355	P053169-B-10_S03723467				-0.27	9
356	P053168-E-12_S03723094				-0.28	9
357	P053168-F-11_S03722997				-0.29	9
358	P053169-H-07_S03723341				-0.29	9
359	P053169-F-06_S03723313				-0.3	9
360	P053168-E-09_S03722977				-0.3	9
361	P053168-G-11_S03723004				-0.31	9
362	P053169-E-09_S03723450				-0.31	9
363	P053168-A-11_S03722967				-0.31	9
364	P053169-D-05_S03723295				-0.33	9
365	P053169-F-04_S03723166				-0.34	9
366	P053169-F-08_S03723446				-0.4	9
367	P053169-A-04_S03723180				-0.4	9
368	P053169-E-08_S03723424				-0.43	9
369	P053168-G-08_S03722933				-0.45	9
370	P053169-C-05_S03723285				-0.46	9
371	P053168-B-01_S03722061				-0.47	9
372	P053168-E-02_S03722184				-0.54	9
373	P053168-G-10_S03722966				-0.55	9
374	P053168-D-08_S03722699				-0.55	9
375	P053168-G-06_S03722434				-0.55	9
376	P053169-D-07_S03723336				-0.55	9
377	P053168-C-01_S03722075				-0.55	9
378	P053168-C-07_S03722502				-0.57	9
379	P053169-C-07_S03723247				-0.63	9
380	P053168-E-08_S03722830				-0.75	9
381	P053168-B-09_S03722914				-0.79	9
382	P053168-F-12_S03723066				-0.96	9
383	P053168-D-02_S03722190				-0.97	9
384	P053169-G-07_S03723338				-1.19	9
385	P053169-F-12_S03723612				-1.33	9
386	P053169-H-08_S03723485				-1.33	9

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
387	P053168-G-07_S03722459				-1.36	9
388	P053169-E-03_S03723164				-1.41	9
389	P053364-A-09_P053168-B-07 P053169-H-01				0.55	9
390	P053364-G-01_P053168-D-03 P053169-D-12				0.51	9
391	P053364-H-01_P053168-B-07 P053169-E-12				0.5	9
392	P053364-A-01_P053168-B-07 P053169-D-12				0.46	9
393	P053364-H-08_P053168-B-06 P053169-H-03				0.42	9
394	P053364-G-05_P053168-D-03 P053169-H-03				0.41	9
395	P053364-A-08_P053168-B-07 P053169-A-05				0.41	9
396	P053364-C-01_P053168-H-01 P053169-D-12				0.39	9
397	P053364-F-01_P053168-E-03 P053169-D-12				0.37	9
398	P053364-C-05_P053168-H-01 P053169-H-03				0.36	9
399	P053364-B-11_P053168-B-06 Axmi205				0.36	9
400	P053364-A-05_P053168-B-07 P053169-H-03				0.34	9
401	P053364-F-10_P053168-B-05 Axmi205				0.33	9
402	P053364-C-09_P053168-H-01 P053169-H-01				0.31	9
403	P053364-B-01_P053168-B-05 P053169-D-12				0.3	9
404	P053364-B-09_P053168-B-05 P053169-H-01				0.29	9
405	P053364-A-03_P053168-B-07 P053169-E-10				0.29	9
406	P053364-A-02_P053168-B-07 P053169-A-10				0.28	9
407	P053364-B-04_P053168-B-05 P053169-A-12				0.28	9
408	P053364-G-10_P053168-H-01 Axmi205				0.27	9
409	P053364-D-01_P053168-H-04 P053169-D-12				0.27	9
410	P053364-G-03_P053168-D-03 P053169-E-10				0.25	9
411	P053364-B-10_P053168-B-05 P053169-A-07				0.24	9
412	P053364-A-10_P053168-B-07 P053169-A-07				0.24	9
413	P053364-E-01_P053168-F-05 P053169-D-12				0.23	9
414	P053364-H-03_P053168-H-01 P053169-E-12				0.23	9
415	P053364-H-10_P053168-H-04 Axmi205				0.23	9
416	P053364-B-05_P053168-B-05 P053169-H-03				0.21	9
417	P053364-F-02_P053168-E-03 P053169-A-10				0.2	9
418	P053364-E-11_P053168-D-03 Axmi205				0.19	9
419	P053364-C-03_P053168-H-01 P053169-E-10				0.19	9
420	P053364-C-02_P053168-H-01 P053169-A-10				0.18	9
421	P053364-B-03_P053168-B-05 P053169-E-10				0.18	9
422	P053364-D-09_P053168-H-04 P053169-H-01				0.16	9
423	P053364-H-02_P053168-B-05 P053169-E-12				0.15	9
424	P053364-G-08_P053168-B-06 P053169-A-10				0.15	9
425	P053364-F-04_P053168-E-03 P053169-A-12				0.15	9
426	P053364-C-08_P053168-H-01 P053169-A-05				0.15	9
427	P053364-F-11_P053168-E-01 Axmi205				0.14	9
428	P053364-E-10_P053168-B-07 Axmi205				0.14	9
429	P053364-H-09_P053168-A-10 P053169-H-03				0.13	9
430	P053364-G-02_P053168-D-03 P053169-A-10				0.13	9
431	P053364-E-12_Axmi205 P053169-A-05				0.13	9
432	P053364-D-03_P053168-H-04 P053169-E-10				0.12	9
433	P053364-F-07_P053168-B-06 P053169-D-12				0.12	9

SEQ ID NO:	Trivial Name	F AE	P-value	EC50 Fold	Deviation	Example #
434	P053364-B-02_P053168-B-05 P053169-A-10				0.1	9
435	P053364-F-08_P053168-B-06 P053169-E-10				0.1	9
436	P053364-A-04_P053168-B-07 P053169-A-12				0.1	9
437	P053364-E-02_P053168-F-05 P053169-A-10				0.1	9
438	P053364-D-11_P053168-F-05 Axmi205				0.1	9
439	P053364-G-04_P053168-D-03 P053169-A-12				0.09	9
440	P053364-C-10_P053168-H-01 P053169-A-07				0.09	9
441	P053364-C-11_P053168-E-03 Axmi205				0.09	9
442	P053364-G-12_Axmi205 P053169-A-07				0.07	9
443	P053364-G-07_P053168-A-10 P053169-D-12				0.06	9
444	P053364-A-07_P053168-B-07 P053169-G-02				0.05	9
445	P053364-C-12_Axmi205 P053169-H-03				0.03	9
446	P053364-H-11_Axmi205 P053169-A-10				0.02	9
447	P053364-D-02_P053168-H-04 P053169-A-10				0.01	9
448	P053364-B-07_P053168-B-05 P053169-G-02				0.01	9
449	P053364-E-05_P053168-F-05 P053169-H-03				0	9
450	P053364-F-09_P053168-B-06 P053169-A-12				-0.01	9
451	P053364-H-07_P053168-E-01 P053169-H-03				-0.01	9
452	P053364-B-08_P053168-B-05 P053169-A-05				-0.02	9
453	P053364-G-09_P053168-A-10 P053169-E-10				-0.02	9
454	P053364-C-04_P053168-H-01 P053169-A-12				-0.03	9
455	P053364-G-11_Axmi205 P053169-D-12				-0.04	9
456	P053364-F-05_P053168-E-03 P053169-H-03				-0.05	9
457	P053364-D-12_Axmi205 P053169-G-02				-0.05	9
458	P053364-H-05_P053168-A-10 P053169-A-12				-0.08	9
459	P053364-E-03_P053168-F-05 P053169-E-10				-0.09	9
460	P053364-C-07_P053168-H-01 P053169-G-02				-0.09	9
461	P053364-D-08_P053168-H-04 P053169-A-05				-0.1	9
462	P053364-H-04_P053168-H-04 P053169-E-12				-0.11	9
463	P053364-D-05_P053168-H-04 P053169-H-03				-0.12	9
464	P053364-D-10_P053168-H-04 P053169-A-07				-0.2	9
465	P053364-E-07_P053168-E-01 P053169-D-12				-0.22	9
466	P053364-D-07_P053168-H-04 P053169-G-02				-0.28	9
467	P053364-B-12_Axmi205 P053169-A-12				-0.3	9
468	P053364-F-03_P053168-E-03 P053169-E-10				-0.32	9
469	P053364-D-04_P053168-H-04 P053169-A-12				-0.45	9
470	P053364-E-08_P053168-E-01 P053169-A-10				-0.47	9
471	P053364-E-09_P053168-E-01 P053169-A-12				-0.49	9
472	P052569-H-01_S03656474				0.8	9
473	P052569-H-12_S03657014				0.79	9
474	P052569-A-08_S03656726				0.76	9
475	P052569-G-12_S03657008				0.76	9
476	P052569-G-11_S03656954				0.59	9
477	P052569-F-06_S03656660				0.46	9
478	P052569-H-04_S03656592				0.46	9
479	P052569-E-12_S03656987				0.43	9
480	P052569-A-07_S03656670				0.42	9

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
481	P052569-H-11_S03656955				0.4	9
482	P052569-F-12_S03657007				0.38	9
483	P052569-C-09_S03656798				0.37	9
484	P052570-E-01_S03657050				0.36	9
485	P052569-E-09_S03656809				0.35	9
486	P052569-D-08_S03656755				0.34	9
487	P052570-H-01_S03657072				0.32	9
488	P052570-D-06_S03657721				0.32	9
489	P052569-D-10_S03656861				0.3	9
490	P052569-F-10_S03656870				0.29	9
491	P052569-E-11_S03656903				0.28	9
492	P052569-B-06_S03656630				0.24	9
493	P052570-C-09_S03657822				0.2	9
494	P052569-C-11_S03656884				0.2	9
495	P052569-C-12_S03656973				0.2	9
496	P052570-H-04_S03657568				0.19	9
497	P052569-A-11_S03656874				0.16	9
498	P052569-A-09_S03656786				0.12	9
499	P052569-A-12_S03656968				0.08	9
500	P052570-B-03_S03657111				0.08	9
501	P052570-A-03_S03657109				0.08	9
502	P052570-E-09_S03657840				0.05	9
503	P052569-C-03_S03656567				0	9
504	P052569-A-10_S03656822				-0.02	9
505	P052569-B-05_S03656597				-0.03	9
506	P052569-G-07_S03656711				-0.04	9
507	P052569-D-11_S03656887				-0.05	9
508	P052569-C-01_S03656390				-0.08	9
509	P052569-F-08_S03656761				-0.1	9
510	P052569-E-03_S03656569				-0.1	9
511	P052569-E-01_S03656453				-0.11	9
512	P052569-D-06_S03656641				-0.12	9
513	P052569-E-04_S03656588				-0.13	9
514	P052570-H-09_S03657859				-0.13	9
515	P052569-G-09_S03656819				-0.14	9
516	P052570-B-01_S03657015				-0.15	9
517	P052569-C-02_S03656518				-0.15	9
518	P052569-G-05_S03656616				-0.18	9
519	P052569-B-02_S03656510				-0.21	9
520	P052569-G-02_S03656544				-0.25	9
521	P052569-F-05_S03656612				-0.25	9
522	P052569-B-03_S03656560				-0.26	9
523	P052570-E-04_S03657538				-0.27	9
524	P052570-G-03_S03657389				-0.27	9
525	P052569-A-06_S03656628				-0.27	9
526	P052570-E-05_S03657649				-0.27	9
527	P052569-F-07_S03656686				-0.28	9

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
528	P052569-B-10_S03656825				-0.29	9
529	P052569-C-05_S03656602				-0.32	9
530	P052569-E-05_S03656609				-0.36	9
531	P052570-G-01_S03657062				-0.39	9
532	P052570-F-03_S03657348				-0.41	9
533	P052570-F-09_S03657844				-0.41	9
534	P052569-C-04_S03656585				-0.44	9
535	P052569-F-01_S03656460				-0.44	9
536	P052570-F-04_S03657542				-0.45	9
537	P052569-A-04_S03656581				-0.47	9
538	P052569-H-06_S03656667				-0.48	9
539	P052570-H-08_S03657813				-0.49	9
540	P052569-F-04_S03656589				-0.49	9
541	P052569-D-01_S03656438				-0.5	9
542	P052569-H-02_S03656556				-0.56	9
543	P052569-C-10_S03656837				-0.57	9
544	P052570-A-09_S03657814				-0.62	9
545	P052570-A-07_S03657735				-0.64	9
546	P052569-H-08_S03656779				-0.66	9
547	P052569-D-02_S03656527				-0.69	9
548	P052570-C-07_S03657741				-0.8	9
549	P052569-D-03_S03656568				-0.86	9
550	P052569-B-04_S03656583				-0.91	9
551	P052569-G-08_S03656766				-0.92	9
552	P052570-E-10_S03657894				-1.04	9
553	P052569-G-01_S03656472				-1.11	9
554	P052569-D-07_S03656678				-1.18	9
555	P052570-A-10_S03657861				-1.23	9
556	P052569-B-07_S03656671				-1.38	9
557	P052570-B-06_S03657713				-1.47	9
558	P052570-A-06_S03657712				-1.83	9
559	PSR1-1-076	25.6	0			10
560	PSR1-1-074	22.7	0			10
561	PSR1-2-145	15.9	0			10
562	PSR1-2-082	11.6	0			10
563	PSR1-2-088	11	0			10
564	PSR1-2-094	10.1	0			10
565	PSR1-2-110	9.7	0			10
566	PSR1-1-073	8.5	0.003			10
567	PSR1-2-091	7.2	0			10
568	PSR1-2-149	7	0			10
569	PSR1-2-087	6.9	0			10
570	PSR1-2-158	6.9	0			10
571	PSR1-2-086	6.7	0			10
572	PSR1-1-053	6.5	0.011			10
573	PSR1-2-096	6.2	0			10
574	PSR1-2-135	6.1	0			10

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
575	PSR1-1-014	5.9	0.018			10
576	PSR1-2-141	5.8	0	2.5		10
577	PSR1-1-006	5.6	0.017			10
578	PSR1-2-095	5.3	0			10
579	PSR1-2-097	5.2	0			10
580	PSR1-1-039	5	0.046			10
581	PSR1-1-049	4.8	0.07			10
582	PSR1-2-127	4.6	0			10
583	PSR1-2-109	4.5	0			10
584	PSR1-2-105	4.5	0	1.9		10
585	PSR1-2-150	4.4	0			10
586	PSR1-1-061	4.3	0.118			10
587	PSR1-1-028	4.1	0.164			10
588	PSR1-2-128	4	0			10
589	PSR1-2-151	4	0			10
590	PSR1-1-045	3.9	0.185			10
591	PSR1-2-113	3.9	0	1.8		10
592	PSR1-1-005	3.9	0.001			10
593	PSR1-1-036	3.8	0.229			10
594	PSR1-1-003	3.7	0.241			10
595	PSR1-2-098	3.7	0	2.1		10
596	PSR1-2-138	3.7	0			10
597	PSR1-2-107	3.7	0			10
598	PSR1-2-143	3.6	0			10
599	PSR1-1-001	3.5	0.282			10
600	PSR1-2-102	3.5	0			10
601	PSR1-2-093	3.5	0			10
602	PSR1-1-051	3.5	0.302			10
603	PSR1-2-101	3.5	0			10
604	PSR1-1-052	3.4	0.317			10
605	PSR1-1-033	3.4	0.324			10
606	PSR1-1-072	3.3	0.341			10
607	PSR1-2-081	3.3	0.001			10
608	PSR1-1-065	3.3	0.383			10
609	PSR1-1-012	3.2	0.371			10
610	PSR1-2-089	3.2	0	2.7		10
611	PSR1-1-063	3.2	0.399			10
612	PSR1-1-004	3.2	0.388			10
613	PSR1-1-048	3.1	0.415			10
614	PSR1-1-058	3.1	0.425			10
615	PSR1-1-057	3.1	0.446			10
616	PSR1-1-060	3.1	0.442			10
617	PSR1-2-142	3	0.001			10
618	PSR1-2-136	3	0			10
619	PSR1-2-083	2.8	0.001			10
620	PSR1-1-066	2.7	0.575			10
621	PSR1-2-111	2.7	0.002			10

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
622	PSR1-2-130	2.7	0.002			10
623	PSR1-2-112	2.7	0.022			10
624	PSR1-1-059	2.6	0.634			10
625	PSR1-1-041	2.6	0.655			10
626	PSR1-1-030	2.6	0.656			10
627	PSR1-1-043	2.5	0.665			10
628	PSR1-1-011	2.5	0.677			10
629	PSR1-1-021	2.5	0.684			10
630	PSR1-1-077	2.5	0.699			10
631	PSR1-2-121	2.5	0.006			10
632	PSR1-2-114	2.4	0.01			10
633	PSR1-2-154	2.3	0.032			10
634	PSR1-1-068	2.3	0.785			10
635	PSR1-2-134	2.3	0.017			10
636	PSR1-1-019	2.3	0.787			10
637	PSR1-2-084	2.3	0.019			10
638	PSR1-1-018	2.2	0.833			10
639	PSR1-1-013	2.2	0.837			10
640	PSR1-1-015	2.2	0.837			10
641	PSR1-1-042	2.2	0.838			10
642	PSR1-2-090	2.2	0.035			10
643	PSR1-1-020	2.1	0.861			10
644	PSR1-1-056	2.1	0.865			10
645	PSR1-2-157	2.1	0.055			10
646	PSR1-1-055	2.1	0.902			10
647	PSR1-1-008	2.1	0.906			10
648	PSR1-2-147	2	0.075			10
649	PSR1-2-117	2	0.097			10
650	PSR1-1-034	2	0.911			10
651	PSR1-2-137	2	0.092			10
652	PSR1-1-009	2	0.922			10
653	PSR1-1-046	2	0.956			10
654	PSR1-2-099	1.9	0.153			10
655	PSR1-1-031	1.9	0.979			10
656	PSR1-1-067	1.9	0.996			10
657	PSR1-2-106	1.9	0.17			10
658	PSR1-1-075	1.9	0.99			10
659	PSR1-2-085	1.8	0.216			10
660	PSR1-1-029	1.7	0.933			10
661	PSR1-2-120	1.7	0.334			10
662	PSR1-1-002	1.7	0.899			10
663	PSR1-1-024	1.6	0.879			10
664	PSR1-1-079	1.6	0.878			10
665	PSR1-2-144	1.6	0.4			10
666	PSR1-2-122	1.6	0.454			10
667	PSR1-2-156	1.6	0.46			10
668	PSR1-2-104	1.6	0.447			10

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
669	PSR1-1-026	1.6	0.854			10
670	PSR1-1-071	1.5	0.832			10
671	PSR1-1-054	1.5	0.833			10
672	PSR1-1-064	1.5	0.83			10
673	PSR1-2-153	1.5	0.536			10
674	PSR1-2-146	1.5	0.59			10
675	PSR1-1-032	1.5	0.814			10
676	PSR1-1-050	1.5	0.81			10
677	PSR1-2-140	1.5	0.597			10
678	PSR1-2-125	1.5	0.59			10
679	PSR1-2-092	1.5	0.607			10
680	PSR1-1-044	1.5	0.798			10
681	PSR1-1-010	1.5	0.791			10
682	PSR1-1-070	1.4	0.766			10
683	PSR1-2-116	1.4	0.773			10
684	PSR1-1-035	1.4	0.745			10
685	PSR1-1-016	1.4	0.74			10
686	PSR1-1-022	1.3	0.729			10
687	PSR1-2-152	1.3	0.881			10
688	PSR1-2-129	1.2	0.962			10
689	PSR1-2-115	1.2	0.91			10
690	PSR1-2-100	1.2	0.845			10
691	PSR1-1-037	1.1	0.606			10
692	PSR1-2-123	1.1	0.649			10
693	PSR1-1-038	1	0.57			10
694	PSR1-2-118	1	0.544			10
695	PSR1-1-027	1	0.567			10
696	PSR1-2-124	1	0.541			10
697	PSR1-1-069	1	0.564			10
698	PSR1-2-148	1	0.523			10
699	PSR1-2-103	0.9	0.35			10
700	PSR1-2-126	0.9	0.336			10
701	PSR1-1-017	0.8	0.501			10
702	PSR1-1-023	0.8	0.499			10
703	PSR1-2-131	0.8	0.294			10
704	PSR1-2-155	0.8	0.256			10
705	PSR1-2-132	0.7	0.17			10
706	PSR1-2-139	0.6	0.144			10
707	PSR1-2-108	0.6	0.111			10
708	PSR1-2-133	0.6	0.11			10
709	PSR1-2-080	0.4	0.051			10
710	PSR1-1-062	0.3	0.315			10
711	PSR1-1-078	0.3	0.303			10
712	PSR1-2-119	0.3	0.017			10
713	PSR1-1-025	0.2	0.277			10
714	PSR1-1-040	0.2	0.263			10
715	PSR1-1-047	0.1	0.251			10

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
716	PSR1-1-007		0.204			10
717	PSR7-141	10				11
718	PSR7-63	8.4				11
719	PSR7-89	8.1				11
720	PSR7-94	6.6				11
721	PSR7-106	6.1				11
722	PSR7-96	6.1				11
723	PSR7-100	5				11
724	PSR7-148	4.7				11
725	PSR7-98	4.7				11
726	PSR7-113	4.5				11
727	PSR7-121	4.5				11
728	PSR7-7	4.5				11
729	PSR7-86	4.2				11
730	PSR7-155	3.7				11
731	PSR7-116	3.6				11
732	PSR7-95	3.4				11
733	PSR7-90	3.3				11
734	PSR7-97	3.3				11
735	PSR7-64	2.8				11
736	PSR7-93	2.8				11
737	PSR7-101	2.7				11
738	PSR7-112	2.5				11
739	PSR7-133	2.4				11
740	PSR7-110	2.3				11
741	PSR7-111	2.1				11
742	PSR7-103	2.1				11
743	PSR7-32	2.1				11
744	PSR7-91	1.9				11
745	PSR7-124	1.9				11
746	PSR7-146	1.7				11
747	PSR7-136	1.6				11
748	PSR7-41	1.5				11
749	PSR7-159	1.5				11
750	PSR7-138	1.3				11
751	PSR7-156	1.1				11
752	PSR7-135	1				11
753	PSR7-122	1				11
754	PSR7-134	0.8				11
755	PSR7-92	0.8				11
756	PSR7-154	0.6				11
757	PSR7-115	0.5				11
758	PSR8-64	15				11
759	PSR8-11	7.9				11
760	PSR8-40	7.5				11
761	PSR8-58	6.6				11
762	PSR8-70	3.3				11

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
763	PSR8-69	3.1				11
764	PSR8-49	2.9				11
765	PSR8-55	2.4				11
766	PSR8-43	2.1				11
767	PSR8-28	2				11
768	PSR8-47	2				11
769	PSR8-31	1.6				11
770	PSR8-33	1.3				11
771	PSR8-37	1.2				11
772	PSR8-23	1.1				11
773	PSR8-4	1.1				11
774	PSR8-50	1				11
775	PSR8-39	0.9				11
776	PSR8-42	0.9				11
777	PSR8-9	0.9				11
778	PSR8-63	0.7				11
779	PSR8-8	0.6				11
780	PSR8-17	0.6				11
781	PSR8-34	0.5				11
782	PSR8-53	0.3				11
783	PSR8-44	0.3				11
784	SFR15-009	18.8	0	6.3		12
785	SFR15-019	8	0	6.8		12
786	SFR15-021	7.9	0	6.9		12
787	SFR15-033	7.9	0	4.8		12
788	SFR15-020	7.3	0	8.5		12
789	SFR15-027	7	0			12
790	SFR15-036	6.6	0			12
791	SFR15-007	6.6	0			12
792	SFR15-017	5.5	0			12
793	SFR15-015	5.5	0			12
794	SFR15-005	5.4	0	5.4		12
795	SFR15-001	5.3	0	4.7		12
796	SFR15-030	5.2	0	3.2		12
797	SFR15-025	5.1	0			12
798	SFR15-011	5	0			12
799	SFR15-012	4.9	0	6.6		12
800	SFR15-029	4.7	0			12
801	SFR15-016	4.7	0			12
802	SFR15-010	4.5	0			12
803	SFR15-003	4.5	0			12
804	SFR15-023	4.4	0			12
805	SFR15-006	4.3	0			12
806	SFR15-028	4	0			12
807	SFR15-026	4	0			12
808	SFR15-032	3.9	0			12
809	SFR15-013	3.7	0			12

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
810	SFR15-004	3.7	0			12
811	SFR15-018	3.5	0			12
812	SFR15-034	3.5	0			12
813	SFR15-008	3.5	0			12
814	SFR15-022	3.5	0			12
815	SFR15-035	3.4	0			12
816	SFR15-031	3.3	0			12
817	SFR15-002	3.1	0			12
818	SFR15-014	2.4	0			12
819	SFR15-024	2.1	0			12
820	SFR12-035	12.37				
821	SFR12-037	9.61				
822	SFR12-038	12.63				
823	SFR12-040	9.05				
824	SFR17-020	27.97				
825	SFR17-021	5.86				
826	SFR17-022	7.98				
827	SFR18-001	7.54				
828	SFR18-002	7.25				
829	SFR18-003	5.94				
830	SFR18-004	7.2				
831	SFR18-005	8.63				
832	SFR18-006	6.94				
833	SFR18-007	11.53				
834	SFR18-009	7.31				
835	SFR19-001	21.43				
836	SFR19-002	11.63				
837	SFR19-003	10				
838	SFR19-004	18.62				
839	SFR19-005	9.79				
840	SFR19-007	15.94				
841	SFR19-008	11.06				
842	SFR19-009	27.29				
843	SFR19-010	12.85				
844	SFR19-011	18.24				
845	SFR19-012	10.92				
846	SFR19-013	9.13				
847	SFR19-014	14.13				
848	SFR19-015	11.32				
849	SFR19-016	20.96				
850	SFR19-017	12.41				
851	SFR19-018	6.18				
852	SFR19-019	8.49				
853	SFR19-020	11.78				
854	SFR19-021	23.45				
855	SFR19-022	13.51				
856	SFR19-023	9.31				

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
857	SFR19-024	12.15				
858	SFR19-025	15.25				
859	SFR19-026	10.59				
860	SFR19-028	6.97				
861	SFR19-031	6.39				
862	SFR19-032	6.54				
863	SFR19-034	6.71				
864	SFR19-035	20.38				
865	SFR19-037	7.3				
866	SFR19-038	9.24				
867	SFR19-039	6.83				
868	SFR19-041	14.94				
869	SFR19-044	13.24				
870	SFR19-045	9.01				
871	SFR19-046	7.13				
872	SFR19-047	16.25				
873	SFR19-048	19.11				
874	SFR19-049	16.73				
875	SFR19-051	8.65				
876	SFR19-052	9.84				
877	SFR19-053	9.75				
878	SFR19-054	10.39				
879	SFR19-056	10.21				
880	SFR19-057	15.88				
881	SFR19-063	10.27				
882	SFR19-065	8.49				
883	SFR19-071	9.78				
884	SFR19-072	9.81				
885	SFR19-075	17.4				
886	SFR19-077	37.92				
887	SFR19-079	37.38				
888	SFR19-080	18.2				
889	SFR20-001	7.45				
890	SFR20-007	6.03				
891	SFR20-010	8.12				
892	SFR20-011	7.15				
893	SFR20-015	6.65				
894	SFR20-016	6.07				
895	SFR20-018	7.13				
896	SFR20-019	9.3				
897	SFR20-020	16.62				
898	SFR20-021	7.98				
899	SFR20-022	6.66				
900	SFR20-024	7.44				
901	SFR20-026	7.96				
902	SFR20-028	11.67				
903	SFR20-029	7.72				

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
904	SFR20-030	6.57				
905	SFR20-031	5.97				
906	SFR20-033	6.64				
907	SFR20-036	7.39				
908	SFR20-044	8.65				
909	SFR20-045	14.21				
910	SFR20-046	15.77				
911	SFR20-047	17.88				
912	SFR20-048	6.82				
913	SFR20-049	13.4				
914	SFR20-051	6.06				
915	SFR20-053	11.18				
916	SFR20-054	12.84				
917	SFR20-056	10.69				
918	SFR20-057	7.2				
919	SFR20-064	9.57				
920	SFR20-066	5.93				
921	SFR20-074	27.75				
922	SFR20-075	14.4				
923	SFR20-076	12.11				
924	SFR21-001	18.77				
925	SFR21-004	9.04				
926	SFR21-007	10.77				
927	SFR21-008	16.87				
928	SFR21-010	11.23				
929	SFR21-011	12.73				
930	SFR21-012	13.63				
931	SFR21-013	8.61				
932	SFR21-014	9.57				
933	SFR21-015	9.74				
934	SFR21-018	9.47				
935	SFR21-024	13.82				
936	SFR22-031	21.7		21.71		
937	SFR22-032	17.97		17.98		
938	SFR23-001	31.56		31.58		
939	SFR23-002	20.38		20.39		
940	SFR23-003	14.13		14.14		
941	SFR23-004	20.27		20.28		
942	SFR23-005	12.83		12.83		
943	SFR23-006	13.49		13.49		
944	SFR23-007	24.97		24.99		
945	SFR23-008	12.02		12.03		
946	SFR23-009	11.6		11.61		
947	SFR23-010	22.49		22.5		
948	SFR23-012	27.88		27.89		
949	SFR23-013	8.05		8.06		
950	SFR24-006	22.58		22.59		

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
951	SFR24-007	9.44		9.44		
952	SFR24-008	20.56		20.57		
953	SFR24-015	5.87		5.87		
954	SFR24-016	18.63		18.64		
955	SFR24-017	6.73		6.74		
956	PSR10_SB-003	34.27		6.12		
957	PSR10_SA-009	31.62		6.17		
958	PSR10_SB-041	29.57		7.42		
959	PSR10_SA-077	27.77		6.79		
960	PSR10_SA-080	26.44		12.65		
961	PSR10_SA-021	25.6		10.8		
962	PSR10_SA-061	24.99		11.01		
963	PSR10_SB-001	23.63		9.38		
964	PSR10_SA-002	23.63				
965	PSR10_SB-011	22.63				
966	PSR10_SA-067	22.46				
967	PSR10_SA-041	22.01				
968	PSR10_SA-065	22		8.01		
969	PSR10_SB-044	21.57				
970	PSR10_SA-089	21.47				
971	PSR10_SB-033	21.1				
972	PSR10_SA-085	20.89				
973	PSR10_SB-015	20.68				
974	PSR10_SB-121	19.58				
975	PSR10_SA-001	19.11				
976	PSR10_SA-040	19.1				
977	PSR10_SB-028	19.09				
978	PSR10_SB-045	18.89		6.99		
979	PSR10_SB-037	18.68				
980	PSR10_SB-013	18.68				
981	PSR10_SA-070	18.58				
982	PSR10_SB-022	18.32				
983	PSR10_SA-020	18.1				
984	PSR10_SA-100	18.07				
985	PSR10_SB-052	17.8				
986	PSR10_SB-118	17.75				
987	PSR10_SA-019	17.41				
988	PSR10_SB-089	17.36				
989	PSR10_SB-043	17.29				
990	PSR10_SB-110	17.07				
991	PSR10_SB-039	16.79				
992	PSR10_SA-101	16.66				
993	PSR10_SA-062	16.53				
994	PSR10_SA-010	16.52				
995	PSR10_SA-022	16.46				
996	PSR10_SA-047	16.28				
997	PSR10_SB-070	16				

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
998	PSR10_SA-017	15.89				
999	PSR10_SB-020	15.72				
1000	PSR10_SA-003	15.25				
1001	PSR10_SB-064	15.25				
1002	PSR10_SB-069	15.12				
1003	PSR10_SA-081	15.02				
1004	PSR10_SB-106	14.91				
1005	PSR10_SB-012	14.88				
1006	PSR10_SB-073	14.87				
1007	PSR10_SA-006	14.86				
1008	PSR10_SB-068	14.82				
1009	PSR10_SB-063	14.65				
1010	PSR10_SB-055	14.62				
1011	PSR10_SA-050	14.62				
1012	PSR10_SB-024	14.59				
1013	PSR10_SB-014	14.58				
1014	PSR10_SB-081	14.57				
1015	PSR10_SA-069	14.49				
1016	PSR10_SA-084	14.44				
1017	PSR10_SB-108	14.42				
1018	PSR10_SB-062	14.31				
1019	PSR10_SB-077	14.31				
1020	PSR10_SB-102	14.31				
1021	PSR10_SB-046	14.28				
1022	PSR10_SB-059	14.28				
1023	PSR10_SB-016	14.21				
1024	PSR10_SB-101	14.16				
1025	PSR10_SA-005	14.07		7.58		
1026	PSR10_SB-067	14.06				
1027	PSR10_SB-002	14.06				
1028	PSR10_SA-024	14.02				
1029	PSR10_SB-082	13.96				
1030	PSR10_SB-095	13.93				
1031	PSR10_SB-065	13.82				
1032	PSR10_SB-104	13.63				
1033	PSR10_SB-117	13.57				
1034	PSR10_SB-057	13.47				
1035	PSR10_SA-090	13.3				
1036	PSR10_SB-019	13.23				
1037	PSR10_SB-038	13.21				
1038	PSR10_SA-042	13.09				
1039	PSR10_SB-010	12.93				
1040	PSR10_SB-103	12.85				
1041	PSR10_SA-096	12.77				
1042	PSR10_SA-068	12.74				
1043	PSR10_SB-026	12.68				
1044	PSR10_SB-097	12.68				

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
1045	PSR10_SB-009	12.59				
1046	PSR10_SA-004	12.43				
1047	PSR10_SA-064	12.33				
1048	PSR10_SB-025	12.3				
1049	PSR10_SB-088	12.19				
1050	PSR10_SB-090	12.18				
1051	PSR10_SB-036	12.15				
1052	PSR10_SA-045	12.14				
1053	PSR10_SB-049	12.14				
1054	PSR10_SB-021	12.13				
1055	PSR10_SB-030	12.07				
1056	PSR10_SA-029	12.04				
1057	PSR10_SB-066	11.97				
1058	PSR10_SA-025	11.84				
1059	PSR10_SB-119	11.81				
1060	PSR10_SB-076	11.79				
1061	PSR10_SA-058	11.68				
1062	PSR10_SB-120	11.63				
1063	PSR10_SB-004	11.61				
1064	PSR10_SB-114	11.57				
1065	PSR10_SA-082	11.57				
1066	PSR10_SB-053	11.55				
1067	PSR10_SB-051	11.46				
1068	PSR10_SB-098	11.35				
1069	PSR10_SA-118	11.3				
1070	PSR10_SA-097	11.26				
1071	PSR10_SB-099	11.21				
1072	PSR10_SA-099	11.17				
1073	PSR10_SB-083	11.16				
1074	PSR10_SA-060	11.16				
1075	PSR10_SA-095	11.13				
1076	PSR10_SB-112	11.09				
1077	PSR10_SA-102	11.08				
1078	PSR10_SB-007	11.02				
1079	PSR10_SB-113	10.99				
1080	PSR10_SB-084	10.98				
1081	PSR10_SB-115	10.98				
1082	PSR10_SA-007	10.95				
1083	PSR10_SA-087	10.94				
1084	PSR10_SB-111	10.93				
1085	PSR10_SA-030	10.84				
1086	PSR10_SB-006	10.82				
1087	PSR10_SB-105	10.8				
1088	PSR10_SB-075	10.74				
1089	PSR10_SA-086	10.65				
1090	PSR10_SB-040	10.59				
1091	PSR10_SB-060	10.5				

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
1092	PSR10_SA-098	10.45				
1093	PSR10_SA-011	10.42				
1094	PSR10_SA-023	10.41				
1095	PSR10_SB-017	10.41		6.94		
1096	PSR10_SB-116	10.36				
1097	PSR10_SB-029	10.35				
1098	PSR10_SA-038	10.3				
1099	PSR10_SA-033	10.25				
1100	PSR10_SA-093	10.18				
1101	PSR10_SA-057	10.16				
1102	PSR10_SB-092	10.09				
1103	PSR10_SB-078	10.05				
1104	PSR10_SB-023	10.02				
1105	PSR10_SB-027	9.95				
1106	PSR10_SA-037	9.93				
1107	PSR10_SB-042	9.92				
1108	PSR10_SB-080	9.9				
1109	PSR10_SB-031	9.88				
1110	PSR10_SB-107	9.87				
1111	PSR10_SB-050	9.84				
1112	PSR10_SB-005	9.81				
1113	PSR10_SA-107	9.79				
1114	PSR10_SB-074	9.74				
1115	PSR10_SA-018	9.58				
1116	PSR10_SB-054	9.53				
1117	PSR10_SB-035	9.48				
1118	PSR10_SA-106	9.21				
1119	PSR10_SA-120	9.13				
1120	PSR10_SB-091	8.84				
1121	PSR10_SB-058	8.76				
1122	PSR10_SA-026	8.7				
1123	PSR10_SA-052	8.69				
1124	PSR10_SA-078	8.66				
1125	PSR10_SB-061	8.63				
1126	PSR10_SA-110	8.59				
1127	PSR10_SB-048	8.54				
1128	PSR10_SB-032	8.49				
1129	PSR10_SB-008	8.45				
1130	PSR11-2-016	32.82				
1131	PSR11-2-077	30.06				
1132	PSR11-2-061	29.88				
1133	PSR11-2-072	29.67				
1134	PSR11-2-038	29.12				
1135	PSR11-2-008	27.23				
1136	PSR11-2-059	27.22				
1137	PSR11-2-063	25.89				
1138	PSR11-2-009	25				

SEQ ID NO:	Trivial Name	FAE	P-value	EC50 Fold	Deviation	Example #
1139	PSR11-2-065	21.97				
1140	PSR11-2-022	20.39				
1141	PSR11_188	18.95				
1142	PSR11-2-070	18.79				
1143	PSR11-2-024	18.6				
1144	PSR11-2-051	17.84				
1145	PSR11_004	16.81				
1146	PSR11_065	16.55				
1147	PSR11_071	15.61				
1148	PSR11-2-068	15.48				
1149	PSR11_120	14.6				
1150	PSR11_002	14.22				
1151	PSR11-2-020	14.15				
1152	PSR11-2-053	13.8				
1153	PSR11-2-067	13.16				
1154	PSR11-2-021	13.1				
1155	PSR11-2-047	11.41				
1156	PSR11_069	11.3				
1157	PSR11-2-033	10.94				
1158	PSR11_010	10.22				
1159	PSR11-2-044	10.2				
1160	PSR11-2-004	9.91				
1161	PSR11-2-025	8.62				
1162	PSR11-2-054	8.45				
1518	SFR12-020			5.66		
1519	SFR12-033			9.74		
1520	SFR12-034			8.87		
1521	SFR12-036			9.11		
1522	SFR18-008			5.36		
1523	SFR18-010			5.67		
1524	SFR18-011			5.81		
1525	SFR21-016			6.61		
1526	SFR21-017			7.89		

THAT WHICH IS CLAIMED:

1. A PHI-4 polypeptide having improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35).
- 5 2. The PHI-4 polypeptide of claim 1, wherein the PHI-4 polypeptide comprises one or more amino acid substitutions compared to the native amino acid at position 40, 42, 43, 46, 52, 97, 98, 99, 145, 150, 151, 153, 163, 171, 172, 182, 196, 206, 210, 216, 220, 278, 283, 289, 293, 328, 333, 334, 336, 338, 339, 342, 346, 354, 355, 370, 389, 393, 396, 401, 402, 403, 410, 412, 416, 417, 426, 442, 447, 452, 454, 455, 457, 461, 462, 500, 509, 520
10 or 527 of SEQ ID NO: 35.
3. The PHI-4 polypeptide of claim 2, wherein the amino acid at position 40 is Leu or Ile; the amino acid at position 42 is Asp or Asn; the amino acid at position 43 is Phe or Glu; the amino acid at position 46 is Glu or Asn; the amino acid at position 52 is Ile or Val; the amino acid at position 97 is Arg, Asp, Glu or Asn; the amino acid at position 98 is Tyr
15 or Phe; the amino acid at position 99 is Lys or Leu; the amino acid at position 145 is Leu or Val; the amino acid at position 150 is Arg or Gln; the amino acid at position 151 is Asp or Ser; the amino acid at position 153 is Leu or Ile; the amino acid at position 163 is Leu or Val; the amino acid at position 171 is Tyr or Phe; the amino acid at position 172 is Ile or Leu; the amino acid at position 182 is Asp or Gln; the amino acid at position 196 is Gln or
20 Asn; the amino acid at position 206 is Tyr or Phe; the amino acid at position 210 is Val or Ile; the amino acid at position 216 is Glu or Gln; the amino acid at position 220 is Glu, Gln, His or Asp; the amino acid at position 278 is Glu or Asn; the amino acid at position 283 is Ile or Val; the amino acid at position 289 is Lys, Gln or Leu; the amino acid at position 293 is Arg, Gln or Glu; the amino acid at position 328 is Lys or Glu; the amino acid at position
25 333 is Ser, Lys or Val; the amino acid at position 334 is Gly, Lys or Arg; the amino acid at position 336 is Gly or Ala; the amino acid at position 338 is Ser or Val; the amino acid at position 339 is Glu, Asn or Gln; the amino acid at position 342 is Ala or Ser; the amino acid at position 346 is Pro or Ala; the amino acid at position 354 is Met or Leu; the amino acid at position 355 is Val or Ile; the amino acid at position 370 is His or Arg; the amino
30 acid at position 389 is Trp or Leu; the amino acid at position 393 is Trp or Leu; the amino acid at position 396 is Ala, Leu, Lys, Thr or Gly; the amino acid at position 401 is Ser, His, Gly, Lys or Pro; the amino acid at position 402 is Lys, His, Gly or Trp; the amino acid at position 403 is Asp or Tyr; the amino acid at position 410 is Ile or Val; the amino acid at position 412 is Pro or Ala; the amino acid at position 416 is Arg or Glu; the amino acid at
35 position 417 is Ala or Ser; the amino acid at position 426 is Thr or Ser; the amino acid at position 442 is Gln or Glu; the amino acid at position 447 is Asp or Lys; the amino acid at

position 452 is Gln or Lys; the amino acid at position 454 is Arg or Gln; the amino acid at position 455 is Val or Ile; the amino acid at position 457 is Trp or Asn; the amino acid at position 461 is Thr or Ser; the amino acid at position 462 is Gly or Ala; the amino acid at position 500 is Arg or Gln; the amino acid at position 509 is Lys or Gln; the amino acid at position 520 is Lys, Glu or Gln; and the amino acid at position 527 is Gln or Lys.

4. The PHI-4 polypeptide of claim 1, wherein the PHI-4 polypeptide comprises an amino acid sequence of the formula,

				5					10					15	
	Met	Xaa	Ser	Ala	Ala	Asn	Ala	Gly	Xaa	Leu	Gly	Asn	Leu	Xaa	Gly
10				20					25					30	
	Xaa	Thr	Ser	Xaa	Gly	Met	Xaa	Tyr	Xaa	Val	Asn	Gly	Leu	Tyr	Ala
				35					40					45	
	Ser	Pro	Glu	Ser	Leu	Xaa	Gly	Gln	Pro	Leu	Phe	Xaa	Xaa	Gly	Gly
				50					55					60	
15	Xaa	Leu	Asp	Ser	Xaa	Xaa	Ile	Glu	Gly	Xaa	Xaa	Xaa	Xaa	Phe	Pro
				65					70					75	
	Xaa	Ser	Met	His	Val	His	Thr	Tyr	Phe	His	Ser	Asp	Xaa	Xaa	Gln
				80					85					90	
	Xaa	Val	Ser	Xaa	Xaa	Ile	Xaa	Xaa	Xaa	Arg	Xaa	Xaa	Xaa	Ser	Xaa
20				95					100					105	
	His	Val	Gly	Xaa	Ser	Gly	Xaa	Xaa	Xaa	Leu	Phe	Ser	Xaa	Ser	Xaa
				110					115					120	
	Ser	Val	Asp	Xaa	Thr	Thr	Xaa	Xaa	Gln	Gln	Leu	Xaa	Glu	Ile	Thr
				125					130					135	
25	Xaa	Ser	Ser	Thr	Arg	Glu	Xaa	His	Val	Leu	Trp	Tyr	Ile	Ser	Leu
				140					145					150	
	Pro	Gly	Ala	Ala	Thr	Leu	Xaa	Ser	Met	Leu	Xaa	Xaa	Xaa	Xaa	Xaa
				155					160					165	
	Xaa	Asp	Xaa	Xaa	Xaa	Pro	Asn	Met	Xaa	Ala	Met	Xaa	Leu	Phe	Xaa
30				170					175					180	
	Xaa	Xaa	Gly	Pro	Xaa	Xaa	Xaa	Xaa	Xaa	Ala	Ala	Val	Gly	Gly	Arg
				185					190					195	
	Leu	Xaa	Xaa	Xaa	Xaa	Ala	Ser	Lys	Xaa	Leu	Xaa	Met	Xaa	Ser	Ser
				200					205					210	

WO 2014/150914

PCT/US2014/024524

	Xaa	Ser	Leu	Ser	Thr	Thr	Xaa	Xaa	Xaa	Ser	Xaa	Xaa	Ala	Xaa	Xaa
					215					220					225
	Gly	Glu	Ile	Xaa	Ile	Xaa	His	Gly	Ser	Xaa	Met	Glu	Lys	Gln	Val
					230					235					240
5	Asn	Ser	Phe	Xaa	Xaa	Xaa	Ser	Thr	Ile	Arg	Xaa	Thr	Ala	Thr	Gly
					245					250					255
	Gly	Lys	Pro	Gly	Xaa	Thr	Xaa	Arg	Ile	Leu	His	Gly	Pro	Asp	Ser
					260					265					270
	Xaa	Xaa	Ala	Phe	Ser	Xaa	Trp	Ala	Xaa	Ser	Leu	Leu	Xaa	Tyr	Ala
10					275					280					285
	Thr	Leu	Met	Asp	Phe	Xaa	Thr	Xaa	Ser	Leu	Xaa	Xaa	Ile	Xaa	Ala
					290					295					300
	Leu	Xaa	Asp	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Glu	Xaa	Xaa	Xaa	Ala	Xaa
					305					310					315
15	Pro	Xaa	Xaa	Met	Xaa	Xaa	Ser	Gln	Xaa	Ser	Ile	Pro	Xaa	Val	Asp
					320					325					330
	Xaa	Val	Leu	Leu	Met	Asp	Ala	Arg	Pro	Pro	Met	Val	Xaa	Ala	Gly
					335					340					345
	Xaa	Asp	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Asp	Xaa	Xaa	Xaa	Xaa	Xaa
20					350					355					360
	Xaa	Ser	Thr	Ser	Xaa	Xaa	Tyr	Lys	Xaa	Xaa	Gly	Gln	Phe	Xaa	Gln
					365					370					375
	Arg	Xaa	His	Xaa	Ser	Val	Ala	Asp	Gly	His	Xaa	Pro	Ile	Xaa	Xaa
					380					385					390
25	Asp	Leu	Phe	Asp	Xaa	Gly	Xaa	Xaa	Xaa	Xaa	Pro	Val	Gly	Xaa	Gln
					395					400					405
	Xaa	Val	Trp	Asp	Xaa	Xaa	Xaa	Xaa	Gly	Lys	Xaa	Xaa	Xaa	Tyr	Xaa
					410					415					420
	Cys	Trp	Arg	Xaa	Xaa	Xaa	Xaa	Gln	Gly	Tyr	Xaa	Xaa	Xaa	Gly	Asp
30					425					430					435
	Val	Xaa	Met	Leu	Ala	Xaa	Ser	Gly	Tyr	Asn	Pro	Pro	Asn	Leu	Pro
					440					445					450
	Xaa	Xaa	Xaa	Cys	Xaa	His	Xaa	Ser	Leu	Xaa	Ala	Xaa	Xaa	Xaa	Thr
					455					460					465

Leu Xaa Xaa Xaa Xaa Trp Xaa Asp Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 470 475 480

Xaa Val Ser Leu Trp Xaa Xaa Gly Ala Ala Gly Ala Val Ala Ser
 485 490 495

5 Ser Cys Phe Ala Gly Val Pro Asn Tyr Asn Asn Pro Pro Asn Ser
 500 505 510

Gly Xaa Ile Xaa Xaa Leu Xaa Gly Ser Ile Ala Cys Val Xaa Thr
 515 520 525

Ser Ala Ile Ala Ser Met Xaa Xaa Met Xaa Ser Met Leu Ser Xaa
 10 530 535

His Xaa Gly Met Glu Ala Met Met Ser Lys Leu (SEQ ID NO: 3),
 wherein

Xaa at position 2 is Ala or Arg;

Xaa at position 9 is Gln, Lys or Glu;

15 Xaa at position 14 is Pro or Ala;

Xaa at position 16 is Val or Asp;

Xaa at position 19 is Met or Leu;

Xaa at position 22 is Gly or Ser;

Xaa at position 24 is Asp, Asn or Gln;

20 Xaa at position 36 is Leu or Met;

Xaa at position 42 is Asp, Asn or Gln;

Xaa at position 43 is Phe or Glu;

Xaa at position 46 is Glu, Asp, Asn or Gly;

Xaa at position 50 is Ile or Val;

25 Xaa at position 51 is Glu or Gln;

Xaa at position 55 is Arg or Lys;

Xaa at position 56 is Ser or Thr;

Xaa at position 57 is Tyr or Phe;

Xaa at position 58 is Thr or Ser;

30 Xaa at position 61 is Arg, Lys or Glu;

Xaa at position 73 is Phe or Tyr;

Xaa at position 74 is Lys, Glu, Gly, Arg, Met, Leu, His or
 Asp;

Xaa at position 76 is Asp or Gln;

Xaa at position 79 is Lys or Glu;
Xaa at position 80 is Glu or Ser;
Xaa at position 82 is Glu, Ile, Leu, Tyr or Gln;
Xaa at position 83 is Glu or Gln;
5 Xaa at position 84 is Tyr or Phe;
Xaa at position 86 is Glu or Gln;
Xaa at position 87 is Lys or Gln;
Xaa at position 88 is Met, Ile or Leu;
Xaa at position 90 is Gln or Glu;
10 Xaa at position 94 is Val or Ile;
Xaa at position 97 is Arg, Asn, Asp, Glu, Gln, Gly, Ser,
Ile, Phe, His, Lys, Thr, Asn, Tyr, Trp, Pro, Cys, Ala, Met,
Val or Leu;
Xaa at position 98 is Tyr or Phe;
15 Xaa at position 99 is Lys, Leu, Tyr, Ile, Met, Phe, Cys, Val
or Asn;
Xaa at position 103 is Ala or Gly;
Xaa at position 105 is Leu or Ile;
Xaa at position 109 is Phe, Lys, Gly, Met, Ser, Asp, Asn,
20 Glu, Cys, Ala or Arg;
Xaa at position 112 is Thr or Ser;
Xaa at position 113 is Asp, Glu or Met;
Xaa at position 117 is Thr or Ser;
Xaa at position 121 is Tyr or Phe;
25 Xaa at position 127 is Ala or Thr;
Xaa at position 142 is Arg or Glu;
Xaa at position 146 is Arg or Gln;
Xaa at position 147 is Arg, Glu or Gln;
Xaa at position 148 is Asp, Phe, Pro, Val, Glu, His, Trp,
30 Ala, Arg, Leu, Ser, Gln or Gly;
Xaa at position 149 is Phe or Val;
Xaa at position 150 is Arg, Gln, Glu or Asn;
Xaa at position 151 is Asp, Ser, Ala, Asn, Trp, Val, Gln,
Cys, Met, Leu, Arg or Glu;

- Xaa at position 153 is Leu or Ile;
Xaa at position 154 is Asn or Asp;
Xaa at position 155 is Asn or Lys;
Xaa at position 159 is Pro or Asp;
- 5 Xaa at position 162 is Glu, Asp, Gln, Asn or Leu;
Xaa at position 165 is Lys, Glu, Gln, Pro, Thr, Ala, Leu, Gly, Asp, Val, His, Ile, Met, Trp, Phe, Tyr or Arg;
Xaa at position 166 is Arg or Gln;
Xaa at position 167 is Tyr, Trp or Cys;
- 10 Xaa at position 170 is Tyr or His;
Xaa at position 171 is Tyr or Phe;
Xaa at position 172 is Ile, Leu or Val;
Xaa at position 173 is Ser or Ala;
Xaa at position 174 is Glu, Gln, Asn, Lys, Val or Ser;
- 15 Xaa at position 182 is Asp or Gln;
Xaa at position 183 is Tyr or Val;
Xaa at position 184 is Ser or Thr;
Xaa at position 185 is Ala or Ser;
Xaa at position 189 is Thr, Lys or Ile;
- 20 Xaa at position 191 is Lys or Gln;
Xaa at position 193 is Asp or Asn;
Xaa at position 196 is Gln, Lys, Asn, Asp, Glu, Ala, Ile or Arg;
Xaa at position 202 is Ala or Val;
- 25 Xaa at position 203 is Glu, Thr or His;
Xaa at position 204 is Met or Ala;
Xaa at position 206 is Tyr or Phe;
Xaa at position 207 is Lys or Gln;
Xaa at position 209 is Leu or Pro;
- 30 Xaa at position 210 is Val or Ile;
Xaa at position 214 is Lys, Ser or Gln;
Xaa at position 216 is Glu, Gln, Phe, Val, Tyr or Arg;
Xaa at position 220 is Glu, His, Asp, Thr, Tyr, Val, Ser, Gln, Arg, Trp, Met, Ala, Phe, Ile, Leu, Cys or Asn;

- Xaa at position 229 is Arg or Glu;
Xaa at position 230 is Ser or Glu;
Xaa at position 231 is Asn or Ser;
Xaa at position 236 is Leu or Pro;
5 Xaa at position 245 is Met or Leu;
Xaa at position 247 is Asp or Tyr;
Xaa at position 256 is Gln, Lys or Glu;
Xaa at position 257 is Gln, Ile, Glu, Cys, Ser, His, Trp or Met;
10 Xaa at position 261 is Gln, Glu, Lys or Ala;
Xaa at position 264 is Glu or Gln;
Xaa at position 268 is Asp or Asn;
Xaa at position 276 is Ser or Ala;
Xaa at position 278 is Glu, Asn or Gln;
15 Xaa at position 281 is Gln, Lys or Glu;
Xaa at position 282 is Pro or Gly;
Xaa at position 284 is Trp or Arg;
Xaa at position 287 is Ala or Cys;
Xaa at position 289 is Lys, Leu, Val, Pro, Glu, Gln, Tyr,
20 Thr, Asp, Phe, Ser, Met, Arg, Trp, Ile, His, Asn, Cys, Gly or Ala;
Xaa at position 291 is Glu or Gln;
Xaa at position 292 is Arg or Gln;
Xaa at position 293 is Arg, Glu or Gln;
25 Xaa at position 294 is Val or Ala;
Xaa at position 296 is Leu or Ile;
Xaa at position 297 is Glu or Gln;
Xaa at position 298 is Asp or Gln;
Xaa at position 300 is Phe or Tyr;
30 Xaa at position 302 is Glu or Gln;
Xaa at position 303 is Phe or Tyr;
Xaa at position 305 is Lys, Gln, Ala, Ile, Met, Asn, Thr or Val;
Xaa at position 306 is Gln or Lys;

Xaa at position 309 is Gln, Lys or Glu;
Xaa at position 313 is Lys, Gln or Arg;
Xaa at position 316 is Lys or Gln;
Xaa at position 328 is Lys, Glu or Gln;
5 Xaa at position 331 is Glu, Asn or Gln;
Xaa at position 333 is Ser, Arg, Gly, Lys, Val, Asn, Ala,
His, Gln, Thr, Asp, Ile, Leu, Cys or Glu;
Xaa at position 334 is Gly, Arg, Lys, Ile or Trp;
Xaa at position 335 is Ser or Ala;
10 Xaa at position 336 is Gly or Ala;
Xaa at position 337 is Ala, Val or Gly;
Xaa at position 338 is Ser, His, Val, Lys, Ala, Gly, Thr,
Ile, Glu, Met, Arg, Pro, Asp, Asn or Leu;
Xaa at position 339 is Glu, Asn, Gln, Ile, Pro, Met, Ser,
15 Ala, Cys, Phe, Val, Leu, Asp, Trp, His or Arg;
Xaa at position 341 is Leu or Val;
Xaa at position 342 is Ala, Ser or Val;
Xaa at position 343 is Val or Ile;
Xaa at position 344 is Phe or Trp;
20 Xaa at position 345 is Asn or His;
Xaa at position 346 is Pro or Ala;
Xaa at position 350 is Asn or Ser;
Xaa at position 351 is Gly or Val;
Xaa at position 354 is Met or Leu;
25 Xaa at position 355 is Val, Ile or Leu;
Xaa at position 359 is Gly or Ala;
Xaa at position 362 is Asn or Ser;
Xaa at position 364 is Ala or Ser;
Xaa at position 371 is Ala, Gly or Thr;
30 Xaa at position 374 is Phe or Ile;
Xaa at position 375 is Lys or Arg;
Xaa at position 380 is Leu or Gly;
Xaa at position 382 is Val, Asp or Leu;
Xaa at position 383 is Leu, Ile or Val;

- Xaa at position 384 is Lys, Ala or Gly;
Xaa at position 385 is Ala or Gly;
Xaa at position 389 is Trp or Tyr;
Xaa at position 391 is Arg, Leu, Glu, Gln, Asp or His;
5 Xaa at position 395 is Asp or Cys;
Xaa at position 396 is Ala, Leu, Lys, Asn, Gly, Ile, Met, Arg, Tyr, Gln, His or Thr;
Xaa at position 397 is Gly, Arg or Ala;
Xaa at position 398 is Ser, Gln or Cys;
10 Xaa at position 401 is Ser, His, Pro, Gly, Lys, Val, Arg, Ile, Asn, Phe, Thr, Ala, Asp, Met, Gln or Glu;
Xaa at position 402 is Lys, Phe, His, Arg, Trp, Gly, Asn, Leu, Tyr, Thr, Val, Met, Pro or Ala;
Xaa at position 403 is Asp, Tyr, Trp, Phe or Glu;
15 Xaa at position 405 is Ala or Ser;
Xaa at position 409 is Ala or Pro;
Xaa at position 410 is Ile or Val;
Xaa at position 411 is Pro or Ala;
Xaa at position 412 is Pro or Ala;
20 Xaa at position 416 is Arg, Glu or Gln;
Xaa at position 417 is Ala, Ser or Cys;
Xaa at position 418 is Leu or Met;
Xaa at position 422 is Met or Val;
Xaa at position 426 is Thr or Ser;
25 Xaa at position 436 is Asp or Lys;
Xaa at position 437 is Tyr or Val;
Xaa at position 438 is Val or Arg;
Xaa at position 440 is Val or Leu;
Xaa at position 442 is Gln, Lys or Glu;
30 Xaa at position 445 is Cys, Leu or Thr;
Xaa at position 447 is Asp, Lys, Tyr, Ser, Glu, Ile, Gly, Pro, Leu, Phe, Trp or Thr;
Xaa at position 448 is Val or Ala;
Xaa at position 449 is Gln or Glu;

Xaa at position 452 is Gln, Lys or Glu, Ala;
Xaa at position 453 is Asn or Asp;
Xaa at position 454 is Arg, Tyr, Met, Ser, Val, Ile, Lys,
Phe, Trp, Gln, Gly, His, Asp, Leu, Thr, Pro or Asn;
5 Xaa at position 455 is Val or Ile;
Xaa at position 457 is Trp or Asn;
Xaa at position 459 is Lys, Met, Val, Trp, Gln, Ile, Thr,
Ser, His, Cys, Tyr, Pro, Asn, Ala, Arg or Glu;
Xaa at position 460 is Gly or Ala;
10 Xaa at position 461 is Thr or Ser;
Xaa at position 462 is Gly or Ala;
Xaa at position 463 is Ala, Ser or Gly;
Xaa at position 464 is Arg, Gly, His, Gln, Thr, Phe, Ala,
Asp, Ser or Lys;
15 Xaa at position 465 is Lys, Asn, Val, Met, Pro, Gly, Arg,
Thr, His, Cys, Trp, Phe or Leu;
Xaa at position 466 is Asp or Arg;
Xaa at position 471 is Gln, Lys, Glu or Met;
Xaa at position 472 is Pro or Ser;
20 Xaa at position 497 is Asp or Gln;
Xaa at position 499 is Glu or Gln;
Xaa at position 500 is Arg, Gln or Lys;
Xaa at position 502 is Arg, Glu or Gln;
Xaa at position 509 is Lys, Gln, Glu or Ala;
25 Xaa at position 517 is Gln, Cys, Asn, Val or Pro;
Xaa at position 518 is Glu or Gln;
Xaa at position 520 is Lys, Gln, Glu, His or Ala;
Xaa at position 525 is Gln or Lys; and
Xaa at position 527 is Gln, Lys, Pro, Cys, Glu, Ser, His,
30 Phe or Trp; wherein one or more amino acid(s) designated by
Xaa in SEQ ID NO: 3 is an amino acid different from the
corresponding amino acid of SEQ ID NO: 35 and wherein the
PHI-4 polypeptide has increased insecticidal activity
compared to the polypeptide of SEQ ID NO: 35.

5. A PHI-4 polypeptide, comprising an amino acid sequence of the formula

					5					10					15
	Met	Xaa	Ser	Ala	Ala	Asn	Ala	Gly	Gln	Leu	Gly	Asn	Leu	Pro	Gly
					20					25					30
5	Val	Thr	Ser	Met	Gly	Met	Gly	Tyr	Xaa	Val	Asn	Gly	Leu	Tyr	Ala
					35					40					45
	Ser	Pro	Glu	Ser	Leu	Leu	Gly	Gln	Pro	Leu	Phe	Xaa	Xaa	Gly	Gly
					50					55					60
	Xaa	Leu	Asp	Ser	Ile	Glu	Ile	Glu	Gly	Arg	Ser	Tyr	Thr	Phe	Pro
10					65					70					75
	Arg	Ser	Met	His	Val	His	Thr	Tyr	Phe	His	Ser	Asp	Phe	Xaa	Gln
					80					85					90
	Asp	Val	Ser	Xaa	Glu	Ile	Xaa	Glu	Tyr	Arg	Glu	Lys	Met	Ser	Gln
					95					100					105
15	His	Val	Gly	Val	Ser	Gly	Xaa	Xaa	Xaa	Leu	Phe	Ser	Ala	Ser	Leu
					110					115					120
	Ser	Val	Asp	Xaa	Thr	Thr	Thr	Asp	Gln	Gln	Leu	Thr	Glu	Ile	Thr
					125					130					135
	Tyr	Ser	Ser	Thr	Arg	Glu	Ala	His	Val	Leu	Trp	Tyr	Ile	Ser	Leu
20					140					145					150
	Pro	Gly	Ala	Ala	Thr	Leu	Arg	Ser	Met	Leu	Arg	Xaa	Xaa	Phe	Xaa
					155					160					165
	Xaa	Asp	Xaa	Asn	Asn	Pro	Asn	Met	Pro	Ala	Met	Xaa	Leu	Phe	Xaa
					170					175					180
25	Xaa	Tyr	Gly	Pro	Tyr	Xaa	Ile	Ser	Xaa	Ala	Ala	Val	Gly	Gly	Arg
					185					190					195
	Leu	Xaa	Tyr	Ser	Ala	Ala	Ser	Lys	Thr	Leu	Lys	Met	Asp	Ser	Ser
					200					205					210
	Xaa	Ser	Leu	Ser	Thr	Thr	Ala	Xaa	Met	Ser	Xaa	Lys	Ala	Leu	Val
30					215					220					225
	Gly	Glu	Ile	Lys	Ile	Xaa	His	Gly	Ser	Xaa	Met	Glu	Lys	Gln	Val
					230					235					240
	Asn	Ser	Phe	Arg	Ser	Asn	Ser	Thr	Ile	Arg	Leu	Thr	Ala	Thr	Gly
					245					250					255

	Gly	Lys	Pro	Gly	Met	Thr	Xaa	Arg	Ile	Leu	His	Gly	Pro	Asp	Ser	
					260					265					270	
	Xaa	Xaa	Ala	Phe	Ser	Xaa	Trp	Ala	Glu	Ser	Leu	Leu	Asp	Tyr	Ala	
					275					280					285	
5	Thr	Leu	Met	Asp	Phe	Ser	Thr	Xaa	Ser	Leu	Xaa	Pro	Ile	Trp	Ala	
					290					295					300	
	Leu	Ala	Asp	Xaa	Pro	Glu	Arg	Xaa	Val	Glu	Leu	Glu	Asp	Ala	Phe	
					305					310					315	
10	Pro	Glu	Phe	Met	Lys	Gln	Ser	Gln	Gln	Ser	Ile	Pro	Xaa	Val	Asp	
					320					325					330	
	Lys	Val	Leu	Leu	Met	Asp	Ala	Arg	Pro	Pro	Met	Val	Xaa	Ala	Gly	
					335					340					345	
	Glu	Asp	Xaa	Xaa	Ser	Xaa	Ala	Xaa	Xaa	Asp	Leu	Ala	Xaa	Phe	Asn	
					350					355					360	
15	Xaa	Ser	Thr	Ser	Asn	Gly	Tyr	Lys	Met	Xaa	Gly	Gln	Phe	Xaa	Gln	
					365					370					375	
	Arg	Asn	His	Ala	Ser	Val	Ala	Asp	Gly	His	Ala	Pro	Ile	Phe	Lys	
					380					385					390	
20	Asp	Leu	Phe	Asp	Leu	Gly	Val	Leu	Lys	Ala	Pro	Val	Gly	Trp	Gln	
					395					400					405	
	Xaa	Val	Trp	Asp	Asp	Xaa	Gly	Ser	Gly	Lys	Xaa	Xaa	Xaa	Tyr	Ala	
					410					415					420	
	Cys	Trp	Arg	Ala	Ile	Xaa	Xaa	Gln	Gly	Tyr	Xaa	Xaa	Xaa	Gly	Asp	
					425					430					435	
25	Val	Met	Met	Leu	Ala	Xaa	Ser	Gly	Tyr	Asn	Pro	Pro	Asn	Leu	Pro	
					440					445					450	
	Asp	Tyr	Val	Cys	Xaa	His	Gln	Ser	Leu	Cys	Ala	Xaa	Val	Gln	Thr	
					455					460					465	
30	Leu	Xaa	Asn	Xaa	Xaa	Trp	Trp	Asp	Xaa	Gly	Xaa	Xaa	Xaa	Xaa	Xaa	
					470					475					480	
	Asp	Val	Ser	Leu	Trp	Xaa	Xaa	Gly	Ala	Ala	Gly	Ala	Val	Ala	Ser	
					485					490					495	
	Ser	Cys	Phe	Ala	Gly	Val	Pro	Asn	Tyr	Asn	Asn	Pro	Pro	Asn	Ser	
					500					505					510	

Xaa at position 206 is Tyr or Phe;

- Xaa at position 216 is Glu or Gln;
Xaa at position 220 is Glu, His, Asp, Thr, Tyr, Val, Ser or Gln;
Xaa at position 247 is Asp or Tyr;
5 Xaa at position 256 is Gln or Lys;
Xaa at position 257 is Gln or Ile;
Xaa at position 261 is Gln, Glu, Lys or Ala;
Xaa at position 278 is Glu or Asn;
Xaa at position 281 is Gln, Lys or Glu;
10 Xaa at position 289 is Lys, Leu, Val, Pro, Glu, Gln, Tyr, Thr or Asp;
Xaa at position 293 is Arg, Glu or Gln;
Xaa at position 313 is Lys or Gln;
Xaa at position 328 is Lys, Glu or Gln;
15 Xaa at position 333 is Ser, Gly, Lys, Val or Asn;
Xaa at position 334 is Gly, Arg, Lys or Ile;
Xaa at position 336 is Gly or Ala;
Xaa at position 338 is Ser, His, Val, Lys or Ala;
Xaa at position 339 is Glu, Asn, Ile or Pro;
20 Xaa at position 343 is Val or Ile;
Xaa at position 346 is Pro or Ala;
Xaa at position 355 is Val or Ile;
Xaa at position 359 is Gly or Ala;
Xaa at position 391 is Arg, Leu, Glu, Gln, Asp or His;
25 Xaa at position 396 is Ala, Leu, Lys, Asn, Gly or Thr;
Xaa at position 401 is Ser, His, Pro, Gly, Lys, Val or Arg;
Xaa at position 402 is Lys, Phe, His, Arg, Gly, Trp, Thr, Asn, Tyr or Met;
Xaa at position 403 is Asp or Tyr;
30 Xaa at position 411 is Pro or Ala;
Xaa at position 412 is Pro or Ala;
Xaa at position 416 is Arg or Glu;
Xaa at position 417 is Ala or Ser;
Xaa at position 418 is Leu or Met;

Xaa at position 426 is Thr or Ser;

Xaa at position 440 is Val or Leu;

Xaa at position 447 is Asp, Lys, Tyr, Ser, Glu or Ile;

Xaa at position 452 is Gln, Lys or Glu;

5 Xaa at position 454 is Arg, Tyr, Met, Ser, Val, Ile, Lys, Phe, Trp or Gln;

Xaa at position 455 is Val or Ile;

Xaa at position 459 is Lys, Met, Val, Trp, Gln, Ile or Tyr;

Xaa at position 461 is Thr or Ser;

10 Xaa at position 462 is Gly or Ala;

Xaa at position 463 is Ala or Ser;

Xaa at position 464 is Arg, Gly, His, Ala, Asp, Ser or Lys;

Xaa at position 465 is Lys, Asn, Val, Met, Pro, Gly or Arg;

Xaa at position 471 is Gln, Lys, Glu or Met;

15 Xaa at position 472 is Pro or Ser;

Xaa at position 500 is Arg or Gln;

Xaa at position 509 is Lys, Gln or Ala;

Xaa at position 520 is Lys, Gln, Glu, His or Ala; and

20 Xaa at position 527 is Gln, Lys, Pro, Cys or Glu; wherein one or more amino acid(s) designated by Xaa in SEQ ID NO: 4 is an amino acid different from the corresponding amino acid of SEQ ID NO: 35 and wherein the PHI-4 polypeptide has increased insecticidal activity compared to the polypeptide of SEQ ID NO: 35.

25 6. The PHI-4 polypeptide of any one of claims 2 to 5, further comprising one or more amino acid substitutions at position 86, 359, 399, 464, 465, 466, 467, 468, 499 or 517 of SEQ ID NO: 35.

30 7. The PHI-4 polypeptide of claim 6, wherein the amino acid at position 86 is Glu or Thr; the amino acid at position 359 is Gly or Ala; the amino acid at position 399 is Gly or Ala; the amino acid at position 464 is Arg, Ala, Lys, Asp or Asn; the amino acid at position 465 is Lys or Met, the amino acid at position 467 is Val, Ala, Leu or Thr; the amino acid at position 468 is Ser or Leu; the amino acid at position 499 is Glu or Ala or the amino acid at position 517 is Glu or Arg.

8. The PHI-4 polypeptide of claim 1, wherein the PHI-polypeptide comprises at least one amino acid substitution at a residue relative to SEQ ID NO: 35 in a structural domain selected from:

a hydrophilic residue;

5 a residue in a membrane insertion initiation loop;

a residue in a receptor binding loop; and

a residue in a protease sensitive region,

wherein the PHI-4 polypeptide has increased insecticidal activity compared to the polypeptide of SEQ ID NO: 35.

10 9. The PHI-4 polypeptide of claim 8, wherein the hydrophilic residues are Asp, Glu, Lys, Arg, His, Ser, Thr, Tyr, Trp, Asn, Gln, and Cys.

10. The PHI-4 polypeptide of claim 8 or 9, wherein the membrane insertion loops are between about amino acid at position 92 (Val) and 101 (Ala) and at position 211 (Gly) and 220 (Glu) relative to SEQ ID NO: 35.

15 11. The PHI-4 polypeptide of any one of claims 8 to 10, wherein the receptor binding loops are between about amino acid 332 (Asp) and 340 (Asp), 395 (Asp) and 403 (Asp) , 458 (Asp) and 466 (Asp) relative to SEQ ID NO: 35.

12. The PHI-4 polypeptide of claim 11, wherein the receptor binding loop residue is selected from positions 332, 333, 334, 335, 336, 337, 338, 339, 340, 395, 396, 397, 398, 20 399, 400, 401, 402, 403, 458, 459, 460, 461, 462, 463, 464, 465, 466 of SEQ ID NO: 35.

13. The PHI-4 polypeptide of any one of claims 8 to 12, wherein the protease sensitive region residue is selected from about amino acid residues between 305 (Lys) and 316 (Lys) and 500 (Arg) and 535 (Lys) relative to SEQ ID NO: 35.

14. The PHI-4 polypeptide of any one of claims 1 to 13, wherein the PHI-4 polypeptide 25 comprises 1 to 54 amino acid substitutions compared to SEQ ID NO: 35.

15. The PHI-4 polypeptide of any one of claims 1 to 13, wherein the PHI-4 polypeptide comprises 1 to 15 amino acid substitutions compared to SEQ ID NO: 35.

16. The PHI-4 polypeptide of any one of claims 1 to 13, wherein the PHI-4 polypeptide has at least 80% identity to SEQ ID NO: 35.

30 17. The PHI-4 polypeptide of any one of claims 1 to 16, wherein the insecticidal activity is increased about 1.5 fold or greater compared to AXMI-205 (SEQ ID NO: 35).

18. The PHI-4 polypeptide of any one of claims 1 to 17, wherein the improved insecticidal activity compared to AXMI-205 (SEQ ID NO: 35) is against Western Corn Root Worm (WCRW) larvae.

19. The PHI-4 polypeptide of any one of claims 1-18, wherein 1-25 amino acids are deleted from the N-terminus of the PHI-4 polypeptide and/or C-terminus of the PHI-4 polypeptide.
20. A polynucleotide encoding a PHI-4 polypeptide of any one of claims 1 to 19.
- 5 21. A composition, comprising an insecticidally-effective amount of the PHI-4 polypeptide of any one of claims 1-19.
22. A method of inhibiting growth or killing an insect pest, comprising contacting the insect pest with the composition of claim 21.
23. A method for controlling an insect pest population resistant to a pesticidal protein,
10 comprising contacting the resistant insect pest population with the composition of claim 21.
24. The method of controlling an insect pest population resistant to an pesticidal protein of claim 23, wherein the pesticidal protein is selected from Cry1Ac, Cry1Ab, Cry1A.105, Cry1Ac, Cry1F, Cry1Fa2, Cry1F, Cry2Ab, Cry3A, mCry3A, Cry3Bb1, Cry34Ab1, Cry35Ab1, Vip3A, Cry9c, eCry3.1Ab and CBI-Bt.
- 15 25. A transgenic plant or progeny thereof, comprising the polynucleotide of claim 20.
26. The transgenic plant or progeny thereof of claim 25, wherein the transgenic plant is a monocotyledon.
27. The transgenic plant or progeny thereof of claim 25, wherein the plant is selected
20 from barley, corn, oat, rice, rye, sorghum, turf grass, sugarcane, wheat, alfalfa, banana, broccoli, bean, cabbage, canola, carrot, cassava, cauliflower, celery, citrus, cotton, a cucurbit, eucalyptus, flax, garlic, grape, onion, lettuce, pea, peanut, pepper, potato, poplar, pine, sunflower, safflower, soybean, strawberry, sugar beet, sweet potato, tobacco, tomato ornamental, shrub, nut, chickpea, pigeon pea, millets, hops and pasture
25 grasses.
28. The transgenic plant or progeny thereof of claim 25, further comprising one or more additional transgenic traits.
29. The transgenic plant of claim 28, wherein the one or more additional transgenic trait is selected from insect resistance, herbicide resistance, fungal resistance, viral
30 resistance, stress tolerance, disease resistance, male sterility, stalk strength, increased yield, modified starches, improved oil profile, balanced amino acids, high lysine or methionine, increased digestibility, improved fiber quality, flowering, ear and seed development, enhancement of nitrogen utilization efficiency, altered nitrogen responsiveness, drought resistance or tolerance, cold resistance or tolerance, salt
35 resistance or tolerance and increased yield under stress.

30. Seed, grain or processed product thereof of the transgenic plant of any one of claims 25 to 29, wherein the seed, grain or processed product thereof comprises the polynucleotide.

31. An expression cassette, comprising the polynucleotide of claim 20 operably linked to one or more regulatory sequences directing expression of the PHI-4 polypeptide.

32. The seed of claim 30, wherein one or more seed treatment has been applied to the seed.

33. A method for expressing in a plant a polynucleotide encoding an insecticidal protein, comprising

(a) inserting into a plant cell the polynucleotide of claim 20 encoding the PHI-4 polypeptide;

(b) obtaining a transformed plant cell comprising the nucleic acid sequence of step (a); and

(c) generating from the transformed plant cell a plant capable of expressing the polynucleotide encoding the PHI-4 polypeptide.

34. A method for protecting a plant from an insect pest, comprising expressing in the plant or cell thereof, an insecticidally-effective amount of the PHI-4 polypeptide of any one of claims 1 to 19.

35. A method for controlling an insect pest population, comprising contacting the insect pest population with an insecticidally-effective amount of the PHI-4 polypeptide of any one of claims 1 to 19.

36. A method of inhibiting growth or killing an insect pest, comprising contacting the insect pest with a composition comprising an insecticidally-effective amount of the PHI-4 polypeptide of any one of claims 1 to 19.

37. A method for controlling an insect pest population resistant to a pesticidal protein, comprising contacting the insect pest population with an insecticidally-effective amount of the PHI-4 polypeptide of any one of claims 1 to 19.

38. A fusion protein comprising the PHI-4 polypeptide of any one of claims 1 to 19.

Fig. 1




Fig. 2




Fig. 3

1 50

PHI-4c-ter (1) MVKAGED**DSGSGASEDLA**VFNPTSTNGYKMGVGFQQRNHASVADGHAPIFK

51 100

PHI-4c-ter (51) DLFDLGVLKAPVGWQRVWDD**DAGSGKSKDYA**CWRAIPPQGYRALGDVMMLA

101 150

PHI-4c-ter (101) TSGYNPPNLPDYVCVHQSLCADVQTLQNRVWW**DKGTGARKDVS**LLWQPGAA

151 175

PHI-4c-ter (151) GAVASSCFAGVPNYPNPPNSGDIER

Fig. 4




Fig. 5




Fig. 4

