



US011168311B2

(12) **United States Patent**
Seehra et al.(10) **Patent No.:** **US 11,168,311 B2**(45) **Date of Patent:** ***Nov. 9, 2021**(54) **METHODS FOR TREATING ANEMIA IN A SUBJECT IN NEED THEREOF**(71) Applicant: **Acceleron Pharma Inc.**, Cambridge, MA (US)(72) Inventors: **Jasbir Seehra**, Lexington, MA (US); **Robert Scott Pearsall**, Woburn, MA (US); **Ravindra Kumar**, Acton, MA (US)(73) Assignee: **ACCELERON PHARMA INC.**, Cambridge, MA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **16/795,083**(22) Filed: **Feb. 19, 2020**(65) **Prior Publication Data**

US 2020/0208124 A1 Jul. 2, 2020

Related U.S. Application Data

(60) Continuation of application No. 16/455,301, filed on Jun. 27, 2019, which is a division of application No. 15/336,363, filed on Oct. 27, 2016, now Pat. No. 10,377,996, which is a continuation of application No. 13/750,249, filed on Jan. 25, 2013, now Pat. No. 9,505,813, which is a continuation of application No. 13/247,748, filed on Sep. 28, 2011, now Pat. No. 8,361,957, which is a division of application No. 12/583,177, filed on Aug. 13, 2009, now Pat. No. 8,058,229.

(60) Provisional application No. 61/189,094, filed on Aug. 14, 2008.

(51) **Int. Cl.****C07K 14/475** (2006.01)**C12N 9/12** (2006.01)**A61K 38/17** (2006.01)**C07K 14/47** (2006.01)**C12Q 1/6897** (2018.01)(52) **U.S. Cl.**CPC **C12N 9/12** (2013.01); **A61K 38/179** (2013.01); **A61K 38/1796** (2013.01); **C07K 14/47** (2013.01); **C07K 14/475** (2013.01); **C12Q 1/6897** (2013.01); **C12Y 207/1103** (2013.01); **C07K 2319/30** (2013.01)(58) **Field of Classification Search**CPC **C07K 14/495**
See application file for complete search history.(56) **References Cited****U.S. PATENT DOCUMENTS**4,973,577 A 11/1990 Vale, Jr. et al.
5,118,667 A 6/1992 Adams et al.5,658,876 A 8/1997 Crowley et al.
5,703,043 A 12/1997 Celeste et al.
5,760,010 A 6/1998 Klein
5,808,007 A 9/1998 Lee et al.
5,824,637 A 10/1998 Crowley et al.
5,847,078 A 12/1998 Eto et al.
5,885,794 A 3/1999 Mathews et al.
6,004,780 A 12/1999 Soppet et al.
6,034,062 A 3/2000 Thies et al.
6,093,547 A 7/2000 Jin et al.
6,132,988 A 10/2000 Sugino et al.
6,162,896 A 12/2000 Mathews et al.
6,287,816 B1 9/2001 Rosen et al.
6,440,930 B1 8/2002 Rinella, Jr.
6,451,334 B2 9/2002 Perrine
6,537,966 B1 3/2003 Duan et al.
6,548,634 B1 4/2003 Ballinger et al.
6,599,876 B2 7/2003 Kojima
6,605,699 B1 8/2003 Ni et al.
6,632,180 B1 10/2003 Laragh
6,656,475 B1 12/2003 Lee et al.
6,656,708 B1 12/2003 Yu et al.
6,692,925 B1 2/2004 Miyazono et al.
6,696,260 B1 2/2004 Lee et al.
6,777,205 B1 8/2004 Carcagno et al.
6,835,544 B2 12/2004 Mathews et al.
6,891,082 B2 5/2005 Lee et al.
6,914,128 B1 7/2005 Salfeld et al.
7,052,873 B2 5/2006 Tsuchiya
7,192,717 B2 3/2007 Hill et al.
7,202,210 B2 4/2007 Wolfman et al.
7,261,893 B2 8/2007 Veldman et al.
7,320,789 B2 1/2008 Dunham et al.
7,560,441 B2 7/2009 Wolfman et al.
7,612,041 B2 11/2009 Knopf et al.

(Continued)

FOREIGN PATENT DOCUMENTSEP 1 174 149 A1 1/2002
EP 1 362 062 A1 11/2003

(Continued)

OTHER PUBLICATIONS

Attie et al., "A phase 1 study of ACE-536, a regulator of erythroid differentiation, in healthy volunteers," American Journal of Hematology, vol. 89(7): 766-770 (2014).

(Continued)

Primary Examiner — Gyan Chandra

(74) Attorney, Agent, or Firm — White & Case LLP

(57) **ABSTRACT**

In certain aspects, the present invention provides compositions and methods for increasing red blood cell and/or hemoglobin levels in vertebrates, including rodents and primates, and particularly in humans.

25 Claims, 21 Drawing Sheets**(13 of 21 Drawing Sheet(s) Filed in Color)****Specification includes a Sequence Listing.**

(56)

References Cited

U.S. PATENT DOCUMENTS

7,709,605 B2	5/2010	Knopf et al.	2007/0172956 A1	7/2007	Magari et al.
7,842,663 B2	11/2010	Knopf et al.	2007/0184052 A1	8/2007	Lin et al.
7,893,213 B2	2/2011	Mathews et al.	2007/0249022 A1	10/2007	Knopf et al.
7,919,296 B2	4/2011	Wang	2007/0275895 A1	11/2007	Duan et al.
7,947,646 B2	5/2011	Sun et al.	2007/0292885 A1	12/2007	Bejanin et al.
7,951,771 B2	5/2011	Knopf et al.	2008/0021104 A1	1/2008	Tarallo
7,960,343 B2	6/2011	Knopf et al.	2008/0075692 A1	3/2008	Perrine
7,968,091 B2	6/2011	Woolf et al.	2008/0089897 A1	4/2008	Wolfman
7,988,973 B2	8/2011	Sherman	2008/0102065 A1	5/2008	Borges et al.
8,007,809 B2	8/2011	Sherman	2008/0139590 A1	6/2008	Qian et al.
8,058,229 B2	11/2011	Seehra et al.	2008/0261879 A1	10/2008	Melton et al.
8,067,360 B2	11/2011	Knopf et al.	2009/0005308 A1	1/2009	Knopf et al.
8,110,355 B2	2/2012	Atwood et al.	2009/0017019 A1	1/2009	Shields et al.
8,124,830 B2	2/2012	Lee et al.	2009/0047281 A1	2/2009	Sherman
8,128,933 B2	3/2012	Knopf et al.	2009/0074768 A1	3/2009	Knopf et al.
8,138,142 B2	3/2012	Seehra et al.	2009/0087433 A1	4/2009	Wolfman et al.
8,173,601 B2	5/2012	Knopf et al.	2009/0098113 A1	4/2009	Knopf et al.
8,178,488 B2	5/2012	Knopf et al.	2009/0099086 A1	4/2009	Knopf et al.
8,216,997 B2	7/2012	Seehra et al.	2009/0118188 A1	5/2009	Knopf et al.
8,252,900 B2	8/2012	Knopf et al.	2009/0142333 A1	6/2009	Knopf et al.
8,293,236 B2	10/2012	Lin et al.	2009/0148436 A1	6/2009	LaVallie et al.
8,293,881 B2	10/2012	Seehra et al.	2009/0163417 A1	6/2009	Sherman
8,309,082 B2	11/2012	Han et al.	2009/0202471 A1	8/2009	Khetani et al.
8,343,933 B2	1/2013	Knopf et al.	2009/0226460 A1	9/2009	Phillips et al.
8,367,611 B2	2/2013	Knopf et al.	2010/0008918 A1	1/2010	Sherman et al.
8,388,968 B2	3/2013	Berger et al.	2010/0015144 A1	1/2010	Sherman et al.
8,410,043 B2	4/2013	Sun et al.	2010/0028331 A1	2/2010	Sherman et al.
8,435,948 B2	5/2013	Zaidi et al.	2010/0028332 A1	2/2010	Sherman et al.
8,501,678 B2	8/2013	Sun et al.	2010/0068215 A1	3/2010	Seehra et al.
8,629,109 B2	1/2014	Knopf et al.	2010/0113327 A1	5/2010	Van Leeuwen et al.
8,637,023 B2	1/2014	Lin et al.	2010/0125099 A1	5/2010	't Hoen et al.
8,703,694 B2	4/2014	Knopf et al.	2010/0183624 A1	7/2010	Seehra et al.
8,703,927 B2	4/2014	Seehra et al.	2010/0272734 A1	10/2010	Berger et al.
8,716,459 B2	5/2014	Sun et al.	2010/0279409 A1	11/2010	Robson et al.
8,753,627 B2	6/2014	Han et al.	2010/0316644 A1	12/2010	Seehra et al.
8,765,663 B2	7/2014	Seehra et al.	2011/0038831 A1	2/2011	Seehra et al.
8,822,411 B2	9/2014	Lee et al.	2011/0070233 A1	3/2011	Seehra et al.
8,865,168 B2	10/2014	Lin et al.	2011/0092670 A1	4/2011	Knopf et al.
8,895,016 B2	11/2014	Sherman et al.	2011/0129469 A1	6/2011	Koncarevic et al.
8,987,203 B2	3/2015	Van Leeuwen et al.	2011/0135638 A1	6/2011	Seehra et al.
8,999,917 B2	4/2015	Sun et al.	2011/0218147 A1	9/2011	Knopf et al.
9,353,356 B2	5/2016	Knopf et al.	2011/0286998 A1	11/2011	Gregory et al.
9,526,759 B2	12/2016	Knopf et al.	2011/0293526 A1	12/2011	Plikus et al.
9,595,813 B2	3/2017	Raring et al.	2012/0003218 A1	1/2012	Sherman et al.
9,850,298 B2	12/2017	Attie	2012/0015877 A1	1/2012	Seehra et al.
10,377,996 B2	8/2019	Seehra et al.	2012/0052067 A1	3/2012	Sherman
2001/0039036 A1	11/2001	Mathews et al.	2012/0148588 A1	6/2012	Knopf et al.
2003/0082233 A1	5/2003	Lyons et al.	2012/0156204 A1	6/2012	Seehra et al.
2003/0083251 A1	5/2003	Westenfelder	2012/0232021 A1	9/2012	Martini et al.
2003/0106093 A1	6/2003	Narva et al.	2012/0237521 A1	9/2012	Berger et al.
2003/0118556 A1	6/2003	Kaspar et al.	2013/0004489 A1	1/2013	Knopf et al.
2003/0119158 A1	6/2003	Malvar et al.	2013/0065299 A1	3/2013	Knopf et al.
2003/0144203 A1	7/2003	Bowen	2013/0071393 A1	3/2013	Seehra et al.
2003/0215913 A1	11/2003	Alvarez et al.	2013/0108650 A1	5/2013	Kumar et al.
2003/0224397 A1	12/2003	Lowman et al.	2013/0177559 A1	7/2013	Seehra et al.
2003/0224501 A1	12/2003	Young et al.	2013/0184210 A1	7/2013	Knopf et al.
2004/0033511 A1	2/2004	Pfizenmaier et al.	2013/0195862 A1	8/2013	Knopf et al.
2004/0132675 A1	7/2004	Kuo et al.	2013/0225484 A1	8/2013	Sun et al.
2004/0138129 A1	7/2004	MacLeod	2013/0243743 A1	9/2013	Seehra et al.
2004/0197828 A1	10/2004	Gaddy	2013/0244324 A1	9/2013	Seehra et al.
2004/0209805 A1	10/2004	Phillips et al.	2013/0287765 A1	10/2013	Zaidi et al.
2004/0223966 A1	11/2004	Wolfman et al.	2013/0344091 A1	12/2013	Berger et al.
2005/0014733 A1	1/2005	Whittemore et al.	2014/0056902 A1	2/2014	Shimizu et al.
2005/0106148 A1	5/2005	Kay et al.	2014/0079700 A1	3/2014	Knopf et al.
2005/0197292 A1	9/2005	Smithson et al.	2014/0194355 A1	7/2014	Sun et al.
2005/0239070 A1	10/2005	Von Knebel-Doeberitz et al.	2014/0220033 A1	8/2014	Han et al.
2005/0244867 A1	11/2005	Soppet et al.	2014/0303068 A1	10/2014	O'Hehir et al.
2005/0257278 A1	11/2005	Lee et al.	2014/0348827 A1	11/2014	Sun et al.
2006/0068468 A1	3/2006	Knopf et al.	2015/0072927 A1	3/2015	Lin et al.
2006/0172347 A1	8/2006	Mellor et al.	2015/0086556 A1	3/2015	Han et al.
2006/0178316 A1	8/2006	Klaus et al.	2015/0111248 A1	4/2015	Bancel et al.
2006/0210657 A1	9/2006	Chou	2015/0139983 A1	5/2015	Karni et al.
2007/0048830 A1	3/2007	Gilbert et al.	2015/0231206 A1	8/2015	Sun et al.
2007/0117130 A1	5/2007	Han et al.	2015/0258120 A1	9/2015	Zarnitsyn et al.
2007/0149455 A1	6/2007	Wolfman et al.	2015/0266950 A1	9/2015	Sung et al.
			2015/0276766 A1	10/2015	Sung et al.

(56)

References Cited**U.S. PATENT DOCUMENTS**

2015/0283209 A1 10/2015 Sung et al.
 2015/0328249 A1 11/2015 Gonzalez-Cadavid et al.

FOREIGN PATENT DOCUMENTS

EP 1 416 273 A1 5/2004
 EP 2468290 A1 6/2012
 JP 2007-099764 A 4/2007
 WO WO-92/04913 A1 4/1992
 WO WO-92/20793 A1 11/1992
 WO WO-93/00432 A1 1/1993
 WO WO-94/15965 A1 7/1994
 WO WO-94/17784 A1 8/1994
 WO WO-94/26893 A1 11/1994
 WO WO-95/10611 A1 4/1995
 WO WO-95/29685 A1 11/1995
 WO WO-97/023613 A2 7/1997
 WO WO-98/18926 A1 5/1998
 WO WO-99/06559 A1 2/1999
 WO WO-00/18932 A2 4/2000
 WO WO-00/25807 A1 5/2000
 WO WO-0025808 A1 5/2000
 WO WO-00/43781 A2 7/2000
 WO WO-00/62809 A1 10/2000
 WO WO-01/36001 A2 5/2001
 WO WO-01/43763 A1 6/2001
 WO WO-01/87329 A1 11/2001
 WO WO-02/10214 A2 2/2002
 WO WO-02/22680 A2 3/2002
 WO WO-02/40501 A2 5/2002
 WO WO-2002/036152 A1 5/2002
 WO WO-02/043759 A2 6/2002
 WO WO-02/074340 A1 9/2002
 WO WO-02/085306 A2 10/2002
 WO WO-02/094852 A2 11/2002
 WO WO-03/006057 A1 1/2003
 WO WO-03/053219 A2 7/2003
 WO WO-03/072808 A1 9/2003
 WO WO-2003/087162 A2 10/2003
 WO WO-2004/012759 A2 2/2004
 WO WO-2004/016639 A1 2/2004
 WO WO-2004/034962 A2 4/2004
 WO WO-2004/039948 A2 5/2004
 WO WO-2004/069237 A1 8/2004
 WO WO-2004/073633 A2 9/2004
 WO WO-2004/076481 A2 9/2004
 WO WO-2004/086953 A2 10/2004
 WO WO-2004/092199 A2 10/2004
 WO WO-04/108157 A2 12/2004
 WO WO-2005/003158 A2 1/2005
 WO WO-2005/009460 A2 2/2005
 WO WO-2005/014650 A2 2/2005
 WO WO-2005/028517 A2 3/2005
 WO WO-2005/053795 A2 6/2005
 WO WO-2005/070967 A2 8/2005
 WO WO-2005/094871 A2 10/2005
 WO WO-2005/097825 A2 10/2005
 WO WO-2005/113590 A2 12/2005
 WO WO-2006/002387 A2 1/2006
 WO WO-2006/012527 A1 2/2006
 WO WO-2006/012627 A2 2/2006
 WO WO-2006/020884 A2 2/2006
 WO WO-2006/039400 A2 4/2006
 WO WO-2006/063101 A2 6/2006
 WO WO-2006/083183 A1 8/2006
 WO WO-2006/088972 A2 8/2006
 WO WO-2007/038703 A2 4/2007
 WO WO-2007/042169 A2 4/2007
 WO WO-2007/053775 A1 5/2007
 WO WO-2007/062188 A2 5/2007
 WO WO-2007/067616 A2 6/2007
 WO WO-2007/071023 A1 6/2007
 WO WO-2007/075702 A2 7/2007
 WO WO-2007/076127 A2 7/2007
 WO WO-2007/087505 A2 8/2007

WO WO-2007/101060 A2 9/2007
 WO WO-2008/015383 A2 2/2008
 WO WO-2008/031061 A2 3/2008
 WO WO-2008/060139 A1 5/2008
 WO WO-2008/072723 A1 6/2008
 WO WO-2008/073292 A2 6/2008
 WO WO-2008/076437 A2 6/2008
 WO WO-2008/094708 A2 8/2008
 WO WO-2008/097541 A2 8/2008
 WO WO-2008/100384 A2 8/2008
 WO WO-2008/109167 A2 9/2008
 WO WO-2006/115274 12/2008
 WO WO-2008/151078 A1 12/2008
 WO WO-2009/009059 A1 1/2009
 WO WO-2009/019504 A1 2/2009
 WO WO-2009/019505 A2 2/2009
 WO WO-2009/021747 A2 2/2009
 WO WO-2009/025651 A1 2/2009
 WO WO-2009/058346 A1 5/2009
 WO WO-2009/070243 A2 6/2009
 WO WO-2009/114180 A1 9/2009
 WO WO-2009/137075 A1 11/2009
 WO WO-2009/137613 A2 11/2009
 WO WO-2009/158015 A2 12/2009
 WO WO-2009/158025 A2 12/2009
 WO WO-2009/158033 A2 12/2009
 WO WO-2010/019261 A1 2/2010
 WO WO-2010/083034 A1 7/2010
 WO WO-2010/125003 A1 11/2010
 WO WO-2010/151426 A1 12/2010
 WO WO-2010/144452 A1 12/2010
 WO WO-2011/020045 A1 2/2011
 WO WO-2011/031901 A1 3/2011
 WO WO-2012/027065 A2 3/2012
 WO WO-2013/006437 A1 1/2013
 WO WO-2013/059347 A1 4/2013
 WO WO-2013/063536 A1 5/2013
 WO WO-2014/066487 A2 5/2014
 WO WO-2014/064292 A1 5/2014
 WO WO-2014/152940 A1 9/2014
 WO WO-2015/017576 A1 2/2015
 WO WO-2015/022658 A2 2/2015
 WO WO-2015/089575 A1 6/2015
 WO WO-2015/108972 A1 7/2015
 WO WO-2015/111008 A2 7/2015
 WO WO-2015/161220 A1 10/2015
 WO WO-2015/152183 A1 10/2015
 WO WO-2015/162590 A1 10/2015
 WO WO-2015/192127 A2 12/2015
 WO WO-2016/183280 A1 11/2016

OTHER PUBLICATIONS

Beijer et al., "A role of active brown adipose tissue in cancer cachexia?", *Oncology Reviews*, vol. 6(1): 88-94; E11 (2012).
 Bradley et al., "Visions & Reflections (Minireview) Myostatin as a therapeutic target for musculoskeletal disease", *Cellular and Molecular Life Sciences*, vol. 65: 2119-2124 (2008).
 Camaschella, Clara, "Hereditary Sideroblastic Anemias: Pathophysiology, Diagnosis, and Treatment," *Seminars in Hematology*, vol. 46(4): 371-377 (2009).
 Cesari et al., "Bone density and hemoglobin levels in older persons: results from the InCHIANTI study," *Osteoporos Int.*, vol. 16: 691-699 (2005).
 Chitturi, Shivakumar, "Treatment for nonalcoholic fatty liver disease," *Therapeutic Advances in Gastroenterology*, vol. 1(3): 173-189 (2008).
 Cypess and Kahn, "Brown fat as a therapy for obesity and diabetes," *Current Opinion in Endocrinology Diabetes, and Obesity*, vol. 17(2): 143-149 (2010).
 Cypess et al., "Identification and Importance of Brown Adipose Tissue in Adult Humans," *the New England Journal of Medicine*, vol. 260(15): 1509-1517 (2009).
 Fabbrini et al., "Obesity and Nonalcoholic Fatty Liver Disease: Biochemical, Metabolic, and Clinical Implications," *Hepatology Reviews*, vol. 51(2): 679-689 (2010).

(56)

References Cited

OTHER PUBLICATIONS

- Halpern et al., "Anemia, costs and mortality in Chronic Obstructive Pulmonary Disease," Cost Effectiveness and Resource Allocation, vol. 4: 17-24 (2006).
- Heuser and Ganser, "Recombinant human erythropoietin in the treatment of nonrenal anemia," *Ann. Hematology*, vol. 85:69-78 (2006).
- Marchesini et al., "Nonalcoholic Fatty Liver Disease: A Feature of the Metabolic Syndrome," *Diabetes*, vol. 50: 1844-1850 (2001).
- Mathews, et al., "Characterization of Type II Activin Receptors," *the Journal of Biological Chemistry*, vol. 268(25): 19013-19018 (1993).
- Matsuzaki et al., "Regulation of Endocytosis of Activin Type II Receptors by a Novel PDZ Protein through RAL/RAL-binding Protein 1-dependent Pathway," *the Journal of Biological Chemistry*, vol. 277(21): 19008-19018 (2002).
- Okusuri 110, Nov. 2, 2001 (in Japanese); Partial Translation: Ingredient (common name): Sulpiride; Outline: This is a drug strengthening the stomach mucosa.
- Shankar, Vijay, "Soft Tissue Adipose Tissue Normal Brown Fat," *Hematopathology, PathologyOutlines.com*, 3 pages (2018).
- Sheftel et al., "Mitochondrial Iron Metabolism and Sideroblastic Anemia," *Acta Haematol*, vol. 122: 120-133 (2009).
- Tsuchida, et al., "Cloning and characterization of a transmembrane serine kinase that acts as an activin type I receptor," *Proceedings of the National Academy of Science*, vol. 90: 11242-11246 (1993).
- Vogiatzi et al., "Bone Disease in Thalassemia: A Frequent and Still Unresolved Problem," *Journal of Bone and Mineral Research*, vol. 24: 543-557 (2008).
- Xu et al., "Treatment with Panax Ginseng Antagonizes the Estrogen Decline in Ovariectomized Mice," *Int. Journal of Molecular Sciences*, vol. 15: 7827-7840 (2014).
- Yip, Ray, "Significance of an abnormally low or high hemoglobin concentration during pregnancy: special consideration of iron nutrition," *Am. J. Clin. Nutr.* vol. 72 A7(suppl): 272S-279S (2000).
- Yoshinaga et al., "N-Cadherin is Regulated by Activin A and Associated with Tumor Aggressiveness in Esophageal Carcinoma," *Clinical Cancer Research*, vol. 10: 5702-5707 (2004).
- Yujian et al., "Advances in the Research of Inhibin, Activin and FS in Femal Reproductive Physiology," *Foreign Medical Science Section Family Planning*, vol. 23(3): 134-136 (2004).
- Walker, et al., "Structural basis for potency differences between GDF8 and GDF11," *BMC Biology*, vol. 15(1): 19 (2017).
- "Anti-human Activin RIIA Antibody," R&D Systems, Catalog No. AF340 (Feb. 14, 2006).
- Abaza, M.S.I., et al., "Effects of Amino acid Substitutions Outside an Antigenic Site," *J. Protein Chem.*, 11(5):433-444 (1992).
- Abbotec: ACTR-IIA Antibody: Catalog No. 251303 (<http://www.abbotec.com>) Jun. 3, 2010.
- Abrahams, B. and Ertel, S., 'Acceleron Pharma at Wells Fargo Healthcare; Conference—Final', published on Jun 17, 2014, Fair Disclosure Wire (Quarterly; Earnings Reports), Accession No. 32U3101469591FDW.
- Acceleron Pharma Presents Positive Phase 1 Results Demonstrating ACE-011 Increases Markers of Bone Formation, Acceleron Pharma, pp. 1-2, <www.acceleronpharma.com/contents/news/press-releases/detail.jsp/q/news-id/47> Downloaded from the Internet on Feb. 17, 2009.
- Acceleron, 'Corporate Overview', considered published in Jul. 31, 2014; Retrieved on Aug. 20, 2015 from the Internet.
- Acceleron, 'Review of the Data Presented at the European Hematology Association; 19th Annual Meeting', considered published on Jun. 16, 2014; Retrieved on Aug. 20, 2015 from the Internet.
- Acta Cryst., "The CCP4 suite: programs for protein crystallography: Collaborative Computational Project, No. 4," D50: 760-763 (1994).
- ACTR-II (149/1): sc-57022, Santa Cruz Biotechnology, Inc.: <http://datasheets.scbt.com/sc-57022.pdf>, Dated Jun. 3, 2010.
- ACTR-II (D-15): sc-5669, Santa Cruz Biotechnology, Inc.: <http://datasheets.scbt.com/sc-5669.pdf>, dated Jun. 3, 2010.
- ACTR-II (H-65): sc-25451, Santa Cruz Biotechnology, Inc.: <http://datasheets.scbt.com/sc-25451.pdf>, dated Jun. 3, 2010.
- ACTR-IIA (A-24): sc-130679, Santa Cruz Biotechnology, Inc.: <http://datasheets.scbt.com/sc-130679.pdf>, dated Jun. 3, 2010.
- ACTR-IIA (N-17): sc-5667, Santa Cruz Biotechnology, Inc.: <http://datasheets.scbt.com/sc-5667.pdf>, dated Jun. 3, 2010.
- Akel et al, Neutralization of Autocrine Transforming Growth Factor- β in Human Cord Blood CD34+CD38-Lin-Cells Promotes Stem-Cell-Factor-Mediated Erythropoietin-Independent Early Erythroid Progenitor Development and Reduces Terminal Differentiation. *Stem Cells*, 21:557-567 (2003).
- Akpan, I., et al., "The effects of a soluble activin type IIB receptor on obesity and insulin sensitivity," *International Journal of Obesity*, 33(11):1265-1273 (2009).
- Allendorph, G.P., et al., "Structure of the ternary signaling complex of a TGF- β superfamily member," *PNAS*, 103(20):7643-7648 (2006).
- Anonymous "Ferritin" www.webmd.com/a-to-z-guides/ferritin? page=2 originally published 2008.
- Anonymous "Iron and Thalassemia," Accessed on the Internet Apr. 3, 2014 at <sickle.bwh.harvard.edu/thaliron.html>. Published Aug. 25, 1997.
- Anonymous "Learning about Thalassemia" <<http://www.genome.gov/10001221>> Accessed on Internet Jul. 9, 2013.
- Anti-ActRIIA Antibodies: Commercial Monoclonal Antibodies Against Human ActRIIA (2010).
- Antibodies for ACVR2A: <<http://www.genecards.org/cgi-bin/carddisp.pl?gene=Acvr2a>> (Jun. 8, 2010).
- Attie et al., "A Single Ascending-Dose Study of Muscle Regulator Ace-031 in Healthy volunteers," *Muscle & Nerve*, pp. 1-8 (2012).
- Banks, G.B., et al., "The Value of Mammalian Models for Duchenne Muscular Dystrophy in Developing Therapeutic Strategies," *Current Topics in Developmental Biology*, 84:431-453 (2008).
- Benny Klimek, Margaret E., et al., "Acute inhibition of myostatin-family proteins preserves skeletal muscle in mouse models of cancer cachexia," *Biochemical and Biophysical Research Communications*, 391:1548-1554 (2010).
- Berenson, J.R., "Multiple Myeloma," *Multiple Myeloma: Plasma Cell Disorders*: Merck Manual Professional, pp. 1-5, Jul. 2008.
- Beutler et al., *Williams Hematology*, 6th Edition. McGraw Hill, p. 561, published 2001.
- Bhatia et al., Protein Glycosylation: Implications for in Vivo Functions and Therapeutic Applications. *Advances in Biochemical Engineering/Biotechnology*, vol. 64: 155-201 (1998).
- Binkert, et al., "Cloning, sequence analysis and expression of a cDNA encoding a novel insulin-like growth factor binding protein (IGFBP-2)," *the EMBO Journal*, 8(9):2497-2502 (1989).
- Biosis Accession No. 2015:276893 & Piga, A. et al., 'ACE-536 Increases Hemoglobin and Decreases Transfusion Burden and Serum Ferritin in Adults with Beta-Thalassemia: Preliminary Results from a Phase 2 Study', *Blood*, vol. 124(21): p. 53 (2014).
- Bodey, B., et al., "Failure of Cancer Vaccines: The Significant Limitations of this Approach to Immunotherapy," *Anticancer Research*, 20:2665-2676 (2000).
- Bogdanovich, S., et al., "Functional improvement of dystrophic muscle by myostatin blockade," *Nature*, 420:418-421 (2002).
- Bork, "Powers and Pitfalls in Sequence Analysis: The 70% Hurdle," *Genome Research*, 10:398-400 (2000).
- Bowie, J.U., et al., "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science*, 257:1306-1310 (1990).
- Broxmeyer, H.E., et al, "Selective and indirect modulation of human multipotential and erythroid hematopoietic progenitor cell proliferation by recombinant human activin and inhibin," *Proc. Natl. Acad. Sci. USA*, 85:9052-9056 (1988).
- Burdette et al., Activin A mediates growth inhibition and cell cycle arrest through Smads in human breast cancer cells. *Cancer Research*, 65(17):7968-7975; Abstract (2005).
- Burgess, W.H., et al., "Possible Dissociation of the Heparin-binding and Mitogenic Activities of Heparin-binding (Acidic Fibroblast) Growth Factor-1 from Its Receptor-binding Activities by Site-directed Mutagenesis of a Single Lysine Residue," *J. Cell Biol.*, 111:2129-2138 (1990).

(56)

References Cited

OTHER PUBLICATIONS

- Cadena, S.M., et al., "Administration of a Soluble Activin Type IIB Receptor Promotes Skeletal Muscle Growth Independent of Fiber Type," *Journal of Applied Physiology*, 109:635-642 (2010).
- Cannon and Nedergaard, "Neither fat nor flesh," *Nature*, vol. 454(7207):947-948 (2008).
- Caricasole, A. A. D., et al., "Human Growth-Differentiation Factor 3 (HGDF3): Developmental Regulation in Human Teratocarcinoma Cell Lines and Expression in Primary Testicular Germ Cell Tumours," *Oncogene*, 16:95-103 (1998).
- Carrancio, S. et al. "An activin receptor IIA ligand trap promotes erythropoiesis resulting in a rapid induction of red blood cells and haemoglobin," *British Journal of Haematology*, vol. 165: 870-882 (2014).
- Casset et al., "A Peptide mimetic of an anti-CD4 monoclonal antibody by rational design," *Biochemical and Biophysical Research Communications*, vol. 307:198-205 (2003).
- CDR Definitions from Handbook of Therapeutic Antibodies, (2007).
- Centrella et al., "Activin-A Binding and Biochemical Effects in Osteoblast-Enriched Cultures from Fetal-Rat Parietal Bone," *Molecular and Cellular Biology*, 11(1):250-58 (1991).
- Chamberlain, R.S., et al., "Innovations and strategies for the development of anticancer vaccines," *Expert Opinion on Pharmacotherapy*, 1(4):603-614 (2000).
- Chamow, S.M., and Ashkenazi, A., "Immunoadhesins: Principles and Applications," *TibTech*, 14: 52-60 (1996).
- Chang, Sam S., "Exploring the Effects of Luteinizing Hormone-Releasing Hormone Agonist Therapy on Bone Health: Implications in the Management of Prostate Cancer," *Urology*, vol. 52: 29-35 (2003).
- Chantry et al., "Inhibiting Activin-A Signaling Stimulates Bone Formation and Prevents Cancer-Induced Bone Destruction in Vivo," *Journal of Bone and Mineral Research*, vol. 25(12): 2357-2370 (2010).
- Chapman, B., et al., "Effect of intron A from human cytomegalovirus (Towne) immediate-early gene on heterologous expression in mammalian cells," *Nucleic Acids Research*, 19(14):3979-3986 (1991).
- Chardès et al., "Efficient amplification and direct sequencing of mouse variable regions from any immunoglobulin gene family," *FEBS Lett.* vol. 452(3): 386-394 (1999).
- Chavez-Tapia, Norberto-C et al., "Insulin sensitizers in treatment of nonalcoholic fatty liver disease: Systematic review," *World Journal of Gastroenterology*, vol. 12(48): 7826-7831 (2006).
- Chen et al., "Selection and Analysis of an Optimized Anti-VEGF Antibody: Crystal Structure of an Affinity-matured Fab in Complex with Antigen," *J. Mol. Biol.*, vol. 293: 865-881 (1999).
- Chen, Y.G., et al. "Regulation of Cell Proliferation, Apoptosis, and Carcinogenesis by Activin," *Exp. Biol. Med.*, 227(2):75-87 (2002).
- Cirillo et al., "Hematocrit, blood pressure, and hypertension. The Gubbio Population Study," *Hypertension*, 20(3):319-326 (1992).
- Coerver, et al., "Activin Signaling through Activin Receptor Type II Causes the Cachexia-Like Symptoms in Inhibin-Deficient Mice," *10(5):534-543* (1996).
- Collins, C.D., "Multidisciplinary Symposium: Haematological Malignancies," *Cancer Imaging* 5:S119-S126 (2005).
- Colman, P.M., et al., "Effects of amino acid sequence changes on antibody-antigen interactions," *Research of Immunology*, 145(1):33-36 (1994).
- Crisan et al., "A Reservoir of Brown Adipocyte Progenitors in Human Skeletal Muscle," *Stem Cells*, vol. 26(9):2425-2433 (2008).
- Daluisi et al., "Bone Morphogenetic Protein-3 is a Negative Regulator of Bone Density," *Nature Genetics*, 27:84-88 (2001).
- Database Geneseq [Online], "Variable heavy chain of anti-human Fas ligand antibody NOK-4," retrieved from EBI accession No. GSP:AAW00829; Database accession No. AAW00829; abstract, sequence (1997).
- Database Geneseq [Online], "Monoclonal antibody 10D4 HMGB1 V κ domain," retrieved from EBI accession No. GSP:ADY85028, Database accession No. GSP:ADY85028; abstract, sequence (2005).
- Datta-Mannan et al, Addendum to "An Engineered Human Follistatin Variant: Insights into the Pharmacokinetic and Pharmacodynamic Relationships of a Novel Molecule with Broad Therapeutic Potential," the *Journal of Pharmacology and Experimental Therapeutics*, 1 page (2013).
- Deal, C., "Potential New Drug Targets for Osteoporosis," *Nature Clinical Practice*, 5(1):20-27 (2009).
- Deconinck, N., et al., "Pathophysiology of Duchenne Muscular Dystrophy: Current Hypotheses," *Pediatr. Neurol.*, 36:1-7 (2007).
- del Re et al., "Reconstitution and Analysis of Soluble Inhibin and Activin Receptor Complexes in a Cell-free System," the *Journal of Biological Chemistry*, 279(51):53126-53135 (2004).
- Delogu, G., et al., "DNA vaccine combinations expressing either tissue plasminogen activator signal sequence fusion proteins or ubiquitin-conjugated antigens induce sustained protective immunity in a mouse model of pulmonary tuberculosis," *Infection and Immunity*, 70(1):292-302 (2002).
- DePaolo, L.V., et al., "Passive Immunoneutralization with a Monoclonal Antibody Reveals a Role for Endogenous Activin-B in Mediating FSH Hypersecretion during Estrus and Following Ovariectomy of Hypophysectomized, Pituitary-Grafted Rats," *Endocrinology*, 130(3):1741-1743 (1992).
- Donald et al., "SDR: a database of predicted specificity-determining residues in proteins," *Nucleic Acids Research*, vol. 37: D191-D194 (2009).
- Donaldson et al., GenBank: BAA06548.1: activin typell A receptor precursor [*Homo sapiens*] (1992).
- Donaldson, et al., "Activin and inhibin binding to the soluble extracellular domain of activin receptor II", *Endocrinology* 140(4):1760-1766 (1999).
- Donaldson, et al., "Molecular Cloning and Binding Properties of the Human Type II Activin Receptor", *Biochemical and Biophysical Research Communications*, 184(1):310-316 (1992).
- Dussiot et al., "An activin receptor IIA ligand trap corrects ineffective erythropoiesis in B-thalassemia," *Nature Medicine*, vol. 20: 398-407 (2014).
- Ear et al., "RAP-011 Efficiently Rescues Erythropoiesis in Zebrafish Models of Diamond Blackfan Anemia," 55 ASH Annual Meeting and Exposition. Abstract #3702 (2013).
- Eijken, M., "The Activin A-Follistatin System: Potent Regulator of Human Extracellular Matrix Mineralization," the *FASEB Journal*, 21:2949-2960 (2007).
- Elliot et al., "Enhancement of therapeutic protein in vivo activities through glycoengineering," *Nature Biotechnology*, vol. 21: 414-421 (2003).
- Fafioffe, et al., "Activin and inhibin receptor gene expression in the ewe pituitary throughout the oestrous cycle," *Journal of Endocrinology*, 182:55-68 (2004).
- Fajardo, R. J., et al., "Treatment with a Soluble Receptor for Activin Improves Bone Mass and Structure in the Axial and Appendicular Skeleton of Female Cynomolgus Macaques (*Macaca fascicularis*)," *Bone*, 46:64-71 (2010).
- Fan, et al., "Preclinical evaluation of Hematide, a novel erythropoiesis stimulating agent, for the treatment of anemia," *Experimental Hematology* 34, pp. 1303-1311 (2006).
- Farmer, Stephen R., "Brown Fat and Skeletal Muscle: Unlikely Cousins?," *Cell*, vol. 134(5):726-727 (2008).
- Ferguson et al., "The role of effectors of the activin signalling pathway, activin receptors IIA and IIB, and Smad2, in patterning of tooth development," *Development*, vol. 128: 4605-4613 (2001).
- Foucar, K., Myelodysplastic/Myeloproliferative Neoplasms, *Am J Clin Pathol*, vol. 132: 281-289 (2009).
- Fournier et al., "Blockade of the activin receptor IIB activates functional brown adipogenesis and thermogenesis by inducing mitochondrial oxidative metabolism," *Mol. Cell. Biol.* vol. 32(14): 2871-2879 (2012).
- Frankel et al., "Characterization of diphtheria fusion proteins targeted to the human interleukin-3 receptor," *Protein Engineering*, vol. 13(8): 575-581 (2000).
- Frigon, N.L., et al, "Regulation of Globin Gene Expression in Human K562 Cells by Recombinant Activin A," *Blood*, 79(3):765-772 (1992).

(56)

References Cited

OTHER PUBLICATIONS

- Fuller et al., "Activin A is an Essential Cofactor for Osteoclast Induction," *Biochemical and Biophysical Research Communications*, 268:2-7 (2000).
- Funaba et al., "Expression and Localization of Activin Receptors During Endochondral Bone Development," *European Journal of Endocrinology*, 144:63-71 (2001).
- Gaddy-Kurten et al., "Inhibin Suppresses and Activin Stimulates Osteoblastogenesis and Osteoclastogenesis in Murine Bone Marrow Cultures," *Endocrinology*, 143(1):74-83 (2002).
- Gamer et al., "BMP-3 is a Novel Inhibitor of Both Activin and BMP-4 Signaling in *Xenopus* Embryos," *Developmental Biology*, 285:156-168 (2005).
- Ge, G., et al., "GDF11 Forms a Bone Morphogenetic Protein 1-Activated Latent Complex That Can Modulate Nerve Growth Factor-Induced Differentiation of PC12 Cells," *Molecular and Cellular Biology*, 25(14):5846-5858 (2005).
- GenBank NM_001106, *Homo sapiens* activin a receptor, type IIB (ACVR2B), mRNA, <http://www.ncbi.nlm.nih.gov/entrez/viewer.fcgi?db=nucleotide&val=116734707> (Downloaded on Jan. 29, 2007).
- GenBank NP_001607.1, Activin A Type II receptor precursor [*Homo sapiens*], <http://www.ncbi.nlm.nih.gov/protein/4501897?sat=34&satkey=10571517> (Apr. 22, 2005); downloaded Nov. 24, 2015).
- Gilbert, R., et al., "Prolonged dystrophin expression and functional correction of mdx mouse muscle following gene transfer with a helper-dependent (guttated) adenovirencoding murine dystrophin," *Human Molecular Genetics*, 12(11):1287-1299 (2003).
- Gilchrist, A., et al., "Antagonists of the Receptor-G Protein Interface Block Gi-coupled Signal Transduction," *Journal of Biological Chemistry*, the American Society of Biological Chemists, Inc., 273(24):14912-14919 (1998).
- Gonzalez-Cadavid, N.F., et al., "Organization of the human myostatin gene and expression in healthy men and HIV-infected men with muscle wasting," *PNAS*, 95:14938-14943 (1998).
- Gray, et al., "Identification of a binding site on the type II activin receptor for activin and inhibin," *Journal of Biological Chemistry*, 275(5):3206-3212(2000).
- Greenspan, N.S., et al., "Defining epitopes: It's not as easy as it seems," *Nature Biotechnology*, 17:936-937 (1999).
- Greenwald, et al., "Characterization of the Extracellular Ligand-Binding Domain of the Type II Activin Receptor," *Biochemistry*, 37(47):16711-16718 (1998).
- Greenwald, et al., "The BMP7/ActRII Extracellular Domain Complex Provides New Insights into the Cooperative Nature of Receptor Assembly," *Molecular Cell*, 11:605-617 (2003).
- Greenwald, J., et al., "Three-finger toxin fold for the extracellular ligand-binding domain of the type II activin receptor serine kinase," *Nature Structural Biology*, 6(1):18-22 (1999).
- Gregoriadis, G., et al., "Polysialic acids: potential in drug delivery," *FEBS*, 315(3):271-276 (1993).
- Guerra, Amaliris et al., "Lack of GDF11 Does not Ameliorate Erythropoiesis in β -Thalassemia and Does not Prevent the Activity of the Trap-Ligand RAP-536," *Blood*, 2018, 132:165 (6 pages).
- Guo, et al., Protein Tolerance to Random Amino Acid Change. *Proc. Natl. Acad. Sci. USA*, 101(25):9205-9210 (Jun. 22, 2004). Epub Jun. 14, 2004.
- Gupta, V. et al., "Transforming Growth Factor- β Superfamily: Evaluation as Breast Cancer Biomarkers and Preventive Agents," *Current Cancer Drug Targets*, 4:165-182 (2004).
- Gura, T., "Cancer Models: Systems for Identifying New Drugs are Often Faulty," *Science*, 278(5340):1041-1042 (1997).
- Haidar et al., "Paraspinal extramedullary hematopoiesis in patients with thalassemia intermedia," *Eur Spine J.*, vol. 19: 871-878 (2010).
- Hamrick et al., "Bone Mineral Content and Density in the Humerus of Adult Myostatin-Deficient Mice," *Calcified Tissue International*, 71:63-68 (2002).
- Hamrick, "Increased Bone Mineral Density in the Femora of GDF8 Knockout Mice," *the Anatomical Record, Part A* 272A:388-391 (2003).
- Hamrick, M.W., et al., "Femoral Morphology and Cross-sectional Geometry of Adult Myostatin-deficient Mice," *Bone*, 27(3):343-349 (2000).
- Harousseau et al., "Multiple Myeloma," *American Society of Hematology*, pp. 237-256 (2004).
- Harrison, C.A., et al., "An Activin Mutant with Disrupted ALK4 Binding Blocks Signaling via Type II Receptors," *the Journal of Biological Chemistry*, 279(27):28036-28044 (2004).
- Harrison, et al., "Antagonists of activin signaling: mechanisms and potential biological applications," *Trends in Endocrinology and Metabolism*, 16(2):73-78 (2005).
- Hashimoto et al., "Functional Regulation of Osteoblastic Cells by the Interaction of Activin-A with Follistatin," *the Journal of Biological Chemistry*, 267(7):4999-5004 (1992).
- Hemmatti-Brivanlou, A., et al., "A truncated activin receptor inhibits mesoderm induction and formation of axial structures in *Xenopus* embryos," *Nature*, 359:609-614 (1992).
- Herbert, W.J., et al., *The Dictionary of Immunology*, Academic Press, 3rd Edition, London, pp. 58-59 (1985).
- Hilden, K., et al., "Expression of Type II Activin Receptor Genes During Differentiation of Human K562 Cells and cDNA Cloning of the Human Type IIB Activin Receptor," *Blood*, 83(8):2163-2170 (1994).
- Hill, J.J., et al., "Regulation of Myostatin in Vivo by Growth and Differentiation Factor-Associated Serum Protein-1: A Novel Protein with Protease Inhibitor and Follistatin Domains," *Molecular Endocrinology*, 17(6):1144-1154 (2003).
- Holm et al., "Functional mapping and single chain construction of the anti-cytokeratin 8 monoclonal antibody TS1," *ScienceDirect; Molecular Immunology*, vol. 44(6): 1075-1084 (2007).
- Hsieh, Matthew M, et al., "HIF-prolyl hydroxylase inhibition results in endogenous erythropoietin induction, erythrocytosis, and modest fetal hemoglobin expression in rhesus macaques," *Blood*, 110(6):2140-2147 (2007).
- Human Activin RIIA Antibody, R&D Systems, Tools for Cell Biology Research, Catalog No. MAB340 (Rev Mar. 22, 2011).
- Ikenoue et al., "Inhibitory Effects of Activin-A on Osteoblast Differentiation During Cultures of Fetal Rat Calvarial Cells," *Journal of Cellular Biochemistry*, 75:206-214 (1999).
- International Search Report, PCT/US2010/045509, dated Nov. 23, 2010.
- Ito et al., "Presence of activin signal transduction in normal ovarian cells and epithelial ovarian carcinoma," *British Journal of Cancer*, vol. 82(8): 1415-1420 (2000).
- Kaiser, J., "First Pass at Cancer Genome Reveals Complex Landscape," *Science*, 313:1370 (2006).
- Kanemitsu, Fusae, "Clinical application of subforms of creatine kinase MM and macro creatine kinases," *Journal of Chromatography*, vol. 526: 423-438 (1990).
- Kaspar, B.K., et al., "Retrograde Viral Delivery of IGF-1 Prolongs Survival in a Mouse ALS Model," *Science*, 301:839-842 (2003).
- Keutmann et al., "The Role of Follistatin Domains in Follistatin Biological Action," *Molecular Endocrinology*, Jan. 18(1) pp. 228-240 (2003).
- Kim, et al., "Type IIa IgG-Fc Fusion Protein, Increases Hemoglobin and Hematocrit Levels in Postmenopausal Healthy Women," *Blood*, 112(11):1316 (2008).
- Knight, "Roles of Inhibins, Activins, and Follistatin in the Female Reproductive System," *Frontiers in Neuroendocrinology*, 17:476-509 (1996).
- Koncarevic et al., "A Soluble Activin Receptor Type IIB Prevents the Effects of Androgen Deprivation on Body Composition and Bone Health," *Endocrinology*, vol. 151(9): 4289-4300 (2010).
- Kos et al., "Activin type II receptors in embryonic dorsal root ganglion neurons of the chicken," *J. Neurobiol.*, vol. 47(2): 93-108 (2001).
- Kosaki, R., et al., "Left-Right Axis Malformations Associated With Mutations in ACVR2B, the Gene for Human Activin Receptor Type IIB," *American Journal of Medical Genetics*, 82:70-76 (1999).
- Koseki, et al., "Role of TCF- β Family in Osteoclastogenesis Induced by RANKL," *Cellular Signaling*, 14:31-36 (2002).
- Krag, T.O.B., et al., "Heregulin ameliorates the dystrophic phenotype in mdx mice," *PNAS*, 101(38):13856-13860 (2004).

(56)

References Cited

OTHER PUBLICATIONS

- Krnet, J., et al., "Dissociation of Angiogenesis and Tumorigenesis in Follistatin- and Activin-Expressing Tumors," *Cancer Research*, 66(11):5686-5695 (2006).
- Krystal et al., Transforming Growth Factor β 1 is an Inducer of Erythroid Differentiation. *J. Exp. Med.* vol. 180 pp. 851-860 (1994).
- Kubane, B., "Introduction: The Role of the Microenvironment and Cytokines on the Modulation of Erythropoiesis," *Annals New York Academy of Sciences*, pp. 257-258 (1994).
- Kumar and Tiwari, "Iron Overload in Beta Thalassemia Major and Intermedia Patients," *Maedica—Journal of Clinical Medicine*, vol. 8(4): 328-332 (2013).
- Kumar, G. Ed., "The Illustrated Guide to Bone Marrow Diagnosis Second Edition," originally published 2003.
- Kumar, T.R., et al., "Regulation of FSH β and GnRH Receptor Gene Expression in Activin Receptor II Knockout Male Mice," *Mol. Cell. Endocrinol.*, 212(1-2):19-27 (2003).
- Kunihro, T., et al., "Regulation of Muscle Mass and Hepatic Steatosis by Follistatin-derived Myostatin Inhibitors," *Making Muscle in the Embryo and Adult: a joint meeting of Frontiers in Myogenesis and Skeletal Muscle Stem and Satellite Cells*, New York, NY, p. 45 (Abstract) (1990).
- Kuntz, "Structure-based strategies for drug design and discovery," *Science*, 257(5073):1078-1082 (1992).
- Kwiatkowski, J.L. et al., "Iron chelation therapy in sickle-cell disease and other transfusion-dependent anemias," *Hematol Oncol Clin N Am.*, vol. 18: 1355-1377 (2004).
- Lazar, E., et al., "Transforming Growth Factor α : Mutation of Aspartic Acid 47 and Leucine 48 Results in Different Biological Activities," *Molecular and Cellular Biology*, 8(3):1247-1252 (1988).
- Lazar, Mitchell A., "How Now, Brown Fat?" *Science*, vol. 321(5892): 1048-1049 (2008).
- Lebrun, J.J., et al., "Activin and Inhibin Have Antagonistic Effects on Ligand-Dependent Heteromerization of the Type I and Type II Activin Receptors and Human Erythroid Differentiation," *Molecular and Cellular Biology*, 17(3):1682-1691 (1997).
- Lee et al., "Regulation of Muscle Growth by Multiple Ligands Signaling Through Activin Type II Receptors," *PNAS* 102(50):18117-18122 (2005).
- Lee, et al., "Regulation of Myostatin Activity and Muscle Growth," *PNAS*, 98(16):9306-9311 (2001).
- Leto et al., "Activin A Circulating Levels in Patients with Bone Metastasis from Breast or Prostate Cancer," *Clin Exp Metastasis*, 23(2):117-122 (2006).
- Li, Q., et al., "Prevention of cachexia-like syndrome development and reduction of tumor progression in inhibin-deficient mice following administration of a chimeric activin receptor type II-murine Fc protein," *Molecular Human Reproduction*, 13(9):675-683 (2007).
- Lifespan Biosciences, Activin Receptor Type 2A (ACVR2A) Mouse anti-Human Monoclonal Antibody—LS-C33835—LifeSpan Biosciences, (2010).
- Liu et al., "Characterization of isoforms of activin receptor-interacting protein 2 that augment activin signaling," *Journal of Endocrinology*, vol. 189: 409-421 (2006).
- Lotinun, S., et al., "A Soluble Activin Receptor Type IIA Fusion Protein (ACE-011) Increases Bone Mass Via a Dual Anabolic-Antiresorptive Effect in Cynomolgus Monkeys," *Bone*, 46:1082-1088 (2010).
- Lu, S., et al., "Simian Immunodeficiency Virus DNA Vaccine Trial in Macaques," *Journal of Virology*, 70(6):3978-3991 (1996).
- Ludlow, H., et al., "Development of a new antibody to the human inhibin/activin β B subunit and its application to improved inhibin B ELISAs," *J. Immunol. Methods*, 329:102-111 (2008).
- Ma, "Animal Models of Disease," *Modern Drug Discovery*, 30-36 (2004).
- MacCallum et al., "Antibody-antigen Interactions: Contact Analysis and Binding site Topography," *J. Mol. Biol.*, vol. 262: 732-745 (1996).
- MacLennan et al., "Multiple Myeloma," *BMJ*, vol. 308:1033-1036 (1994).
- Maguer-Satta, V., et al., "A Novel Role for Fibronectin Type 1 Domain in the Regulation of Human Hematopoietic Cell Adhesiveness Through Binding to Follistatin Domains of FLRG and Follistatin," *Experimental Cell Research*, Academic Press, 312(4):434-442 (2006).
- Maguer-Satta, V., et al., "Regulation of human erythropoiesis by activin A, BMP2, and BMP4, members of the TGF β family," *Experimental Cell Research*, 282:110-120 (2003).
- Maguer-Satta, V., et al., "FLRG, Member of the Follistatin Family, a New Player in Hematopoiesis," *Molecular and Cellular Endocrinology*, Elsevier Ireland Ltd., 225(1-2):109-118 (2004).
- Mallat et al., "Potential mechanisms for renal damage in beta-thalassemia," *J. Nephrol.*, vol. 26(5): 821-828 (2013).
- Marri, R. et al., "Human biochemistry, Moscow, Mir", vol. 1: 34 (1993).
- Martens et al., "Inhibin Interferes with Activin Signaling at the Level of the Activin Receptor Complex in Chinese Hamster Ovary Cells," *Endocrinology*, vol. 138(7): 2928-2936 (1997).
- Mathews, L.S., et al., "Expression Cloning of an Activin Receptor, a Predicted Transmembrane Serine Kinase," *Cell*, 65(6):973-982 (1991).
- Matzuk et al., "Cloning of the human activin receptor cDNA reveals high evolutionary conservation," *Biochim Biophys Acta*, 1130(1):105-108 (1992).
- Matzuk et al., "Different phenotypes for mice deficient in either activins or activin receptor type II," *Nature*, 374:356-360 (1995).
- McCarthy et al., Monoclonal antibodies that recognize the type-2 activin receptor, ACTR2, Hybridoma, vol. 13(3):199-203 (1994) (abstract).
- McNally, E.M., "Powerful Genes—Myostatin Regulation of Human Muscle Mass," *N. Engl. J. Med.*, 350(26):2642-2644 (2004).
- McPherron and Lee, "Suppression of body fat accumulation in myostatin-deficient mice," *the Journal of Clinical Investigation*, vol. 109(5):595-601 (2002).
- McPherron, A.C., et al., "GDF-3 and GDF-9: Two New Members of the Transforming Growth Factor- β Superfamily Containing a Novel Pattern of Cysteines," *Journal of Endocrinology*, 268(5):3444-3449 (1993).
- McPherron, A.C., et al., "Regulation of Skeletal Muscle Mass in Mice by a New TGF- β Superfamily Member," *Nature*, 387:83-90 (1997).
- McPherson, S.J., et al., "Growth inhibitory response to activin A and B by human prostate tumour cell lines LNCaP and DU146," *Journal of Endocrinology*, 154:535-545 (1997).
- Menstruation: Absent Periods (Amenorrhea), Website downloaded on Jun. 14, 2010, <http://adam.about.com/reports/000101_2.htm? p=1> (11 pages).
- Merck Manual. Iron-Utilization Anemias (Sideroblastic Anemias), pp. 1150-1151 (1992).
- Merck Manuals Online Medical Library (online). Anemia of Chronic Disease, Jun. 10, 2008. Downloaded from the internet on Jan. 5, 2010. <<http://web.archive.org/web/20080610070226/http://www.merck.com/mmpe/sec11/ch130/ch130d.html>> pp. 1-2.
- Merck Manuals Online Medical Library (online). Iron Deficiency Anemia, Jun. 10, 2008. Downloaded from the internet on Jan. 5, 2010. <<http://web.archive.org/web/20080610070221/http://www.merck.com/mmpe/sec11/ch130/ch130d.html>> pp. 1-4.
- Meriggiola et al., "Follistatin Decreases Activin-Stimulated FSH Secretion with no Effect on GnRH-Stimulated FSH Secretion in Prepubertal Male Monkeys," *Endocrinology*, 134(4):1967-1970 (1994).
- Mickle, et al., "Genotype-Phenotype Relationships in Cystic Fibrosis," *Med. Clin. North Am.*, 84(3):597-607 (2000).
- Miller et al., Ligand binding to proteins: the binding landscape model. *Protein Sci.*, 6(10):2166-79 (1997).
- Miura, P., et al., "Utrophin upregulation for treating Duchenne or Becker muscular dystrophy: how close are we?," *Trends in Molecular Medicine*, 12(3):122-129 (2006).
- Monoclonal Anti-human Activin RII Antibody, R&D Systems, Catalog No. MAB3391 (Feb. 18, 2009).
- Moore et al., "Molecular Basis of Bone Morphogenetic Protein-15 Signaling in Granulosa Cells," *the Journal of Biological Chemistry*, vol. 278(1): 304-310 (2003).

(56)

References Cited

OTHER PUBLICATIONS

- Morrison et al., "A soluble activin type IIB receptor improves function in a mouse model of amyotrophic lateral sclerosis," *Experimental Neurology*, vol. 217:258-268 (2009).
- Mosekilde, L., et al., "Emerging Anabolic Treatments in Osteoporosis," *Current Drug Safety*, 6:62-74 (2011).
- Murase et al., "Possible Involvement of Protein Kinases and Smad2 Signaling Pathways on Osteoclast Differentiation Enhanced by Activin A," *Journal of Cellular Physiology*, 188:236-242 (2001).
- Murata, T., et al., "Anti-activin A Antibody (IgY) Specifically Neutralizes Various Activin A Activities," *Proceedings of the Society for Experimental Biology & Medicine*, 211(1):100-107 (1996).
- Nagamine et al., "Immunohistochemical Detection of Activin A, Follistatin, and Activin Receptors during Fracture Healing in the Rat," *Journal of Orthopaedic Research*, 16:314-321 (1998).
- Nakamura, K., et al., "Effect of Erythroid Differentiation Factor on Maintenance of Human Hematopoietic Cells in Co-cultures with Allogenic Stromal Cells," *Biochemical and Biophysical Research Communications*, 194(3):1103-1110 (1993).
- Nemeth, E., "Hepcidin in β -thalassemia," *Annals of the New York Academy of Sciences*, vol. 1202: 31-35. Published Aug. 2, 2010.
- Ngo, J.T., et al., "Computational Complexity, Protein Structure Prediction, and the Levinthal Paradox," the Protein Folding Problem and Tertiary Structure Prediction, Merz et al., eds., Birkhauser, Boston, 433 and 492-495 (1994).
- NIH website downloaded May 28, 2014 from: web.archive.org/web/20030409091558/http://www.cc.nih.gov/ccc/patient_education/pepubs/subq.pdf; Patient Information Publications: Giving a Subcutaneous Injection (6 pages total).
- Nolan, V.G., et al, 'Sickle Cell Leg Ulcers: Associations with Haemolysis and SNPs in Klotho, TEK and Genes of the TGF- β /BMP Pathway:—Sickle Cell Leg Ulcers, Genetics and Haemolysis', *British Journal of Haematology*, 133(5), pp. 570-578 (2006).
- Ogawa et al., "Bovine Bone Activin Enhances Bone Morphogenetic Protein-Induced Ectopic Bone Formation," *the Journal of Biological Chemistry*, 267(20):14233-14237 (1992).
- Oh, S.P., et al., "Activin type IIA and IIB receptors mediate Gdf11 signaling in axial vertebral patterning," *Genes & Development*, 16:2749-2754 (2002).
- Oue et al., "Effect of Local Injection of Activin A on Bone Formation in Newborn Rats," *Bone*, 15(3):361-366 (1994).
- Padlan et al., "Identification of specificity-determining residues in antibodies," *the FASEB Journal*, vol. 9:133-139 (1995).
- Pak et al., "Suppression of hepcidin during anemia requires erythropoietic activity," *Blood*, vol. 108(12): 3730-3735 (2006).
- Pakula and Sauer, "Genetic analysis of protein stability and function," *Annu. Rev. Genet.*, vol. 23: 289-310 (1989).
- Patel, K., et al., "The function of Myostatin and strategies of Myostatin blockade—new hope for therapies aimed at promoting growth of skeletal muscle," *Neuromuscular Disorders*, 15:117-126 (2005).
- Paul, William E., *Fundamental Immunology*, 3rd edition, Raven Press, New York, 1003: 292-295 (1999).
- Paulson, Robert F., "Targeting a new regulator of erythropoiesis to alleviate anemia," *Nature Medicine, News and Views*, vol. 20(4) (2 pages) (2014).
- Pearsall et al., An investigative pharmacology study of a GDF-8 (myostatin) inhibitor, ACE-031, in the common Marmoset (*Calithrix jacchus*), Database Biosis, Biosciences Information Service, Accession No. PREV201200750016; *Faseb Journal*, vol. 22, Experimental Biology Annual Meeting, San Diego, CA (Apr. 5-9, 2008) (Abstract).
- Pearsall, et al., "A Soluble Activin Receptor Type IIA (ACTRIIA) Acts as a Novel Bone Anabolic Agent," the Official Journal of the European Calcified Tissue Society, 34th European Symposium on Calcified Tissues, (May 2007).
- Pearsall, et al., "Treatment with a Soluble Activin Type II Receptor Reverses Bone Loss in Ovariectomized Mice," *Journal of Bone and Mineral Research* 2006 Abstracts, 21(1):s1-s530 (2006).
- Pearsall, R.S., et al., "A soluble activin Type IIA receptor induces bone formation and improves skeletal integrity," *PNAS*, 105(9):7082-7087 (2008).
- Pennucci et al., Multiplexed evaluation of a cell-based assay for the detection of antidrug neutralizing antibodies to Panitumumab in human serum using automated fluorescent microscopy, *J. Biomol. Screen.* vol. 15: 644-652 (2010).
- Perrien, D. S., et al., "Inhibin A is an Endocrine Stimulator of Bone Mass and Strength," *Endocrinology*, 148(4):1654-1665 (2007).
- Phillips, A.J., "The challenge of gene therapy and DNA delivery," *J. Pharm. Pharmacology*, 53:1169-1174 (2001).
- Pirollo, K.F., et al., "Targeted Delivery of Small Interfering RNA: Approaching Effective Cancer Therapies," *Cancer Res.*, 68(5):1247-1250 (2008).
- Qi, et al., "Blockade of type β transforming growth factor signaling prevents liver fibrosis and dysfunction in the rat," *PNAS*, 96:2345-2349 (1999).
- R&D Systems 339-RB/CF, Recombinant Human Activin RIIB/Fc Chimera (Aug. 27, 2003).
- R&D Systems 340-R2, Recombinant Human Activin RIIA/Fc Chimera (Aug. 27, 2003).
- R&D Systems Catalogue No. AF339 Datasheet: Human Activin RIIB Antibody [retrieved on Feb. 13, 2013] Retrieved from the Internet: <http://www.rndsystems.com/pdf/af339.pdf>.
- R&D Systems, "Antibody Reference Guide and Catalog Instructions," [retrieved on Feb. 13, 2013]; http://web.archive.org/web/20090220022132/http://rndsystems.com/DAM_public/5658.pdf; published Mar. 14, 2009 as per the Wayback Engine. See, in particular: p. 3.
- Raju, T.S., "Glycosylation in the Fc domain of IgG increases resistance to proteolytic cleavage by papain," *Biochemical and Biophysical Research Communications*, 341:797-803 (2006).
- Rebbapragada, et al., "Myostatin Signals Through a Transforming Growth Factor β -Like Signaling Pathway to Block Adipogenesis," *Molecular and Cellular Biology*, 23(20):7230-7242 (2003).
- Reis, F.M., et al., "Activin, Inhibin and the Human Breast," *Molecular and Cellular Endocrinology*, 225:77-82 (2004).
- Risbridger, G.P., et al., "Activins and Inhibins in Endocrine and Other Tumors," *Endocrine Reviews*, 22(6):836-858 (2001).
- Robinson, G.W., et al., "Inhibins and Activins Regulate Mammary Epithelial Cell Differentiation Through Mesenchymal-epithelial Interactions," *Development*, 124:2701-2708 (1997).
- Rodriguez, J.E.S., et al., "Enhanced Osteoclastogenesis Causes Osteopenia in Twisted Gastrulation-Deficient Mice Through Increased BMP Signaling," *J. Bone Miner. Res.*, 24:1917-1926 (2009).
- Rosenzweig et al., "Cloning and characterization of a human type II receptor for bone morphogenetic proteins," *PNAS*, 92:7632-7636 (1995).
- Ruckle et al., "Single-Dose, Randomized, Double-Blind, Placebo-Controlled Study of ACE-011 (ACTRIIA-IgG1) in Postmenopausal Women," *Journal of Bone and Mineral Research*, vol. 24(4), pp. 744-752 (2009).
- Rund and Rachmilewitz, "Medical Progress Beta-Thalassemia," *N. England J. Medicine*, vol. 35: 1135-1146 (2005).
- Ruzek et al. Minimal Effects on Immune Parameters Following Chronic Anti-TGF- β Monoclonal Antibody Administration to Normal Mice. *Immunopharmacology and Immunotoxicology*, 25(2):235-257 (2003).
- Sakai et al., "Activin Enhances Osteoclast-Like Cell Formation in Vitro," *Biochemical and Biophysical Research Communications*, 195(1):39-46 (1993).
- Sakai et al., "Activin Increases Bone Mass and Mechanical Strength of Lumbar Vertebrae in Aged Ovariectomized Rats," *Bone*, 27(1):91-96 (2000).
- Sakai et al., "Activin release from bone coupled to bone resorption in organ culture of neonatal mouse calvaria," *Bone*, 26(3):235-240 (2000).
- Sakai et al., "Involvement of Activin in the Regulation of Bone Metabolism," *Molecular and Cellular Endocrinology*, 180:183-188 (2001).
- Sakai et al., "Local Administration of Activin Promotes Fracture Healing in the Rat Fibula Fracture Model," *Bone*, 25(2):191-196 (1999).

(56)

References Cited

OTHER PUBLICATIONS

- Sakai et al., the Measurement of Activin/EDF in Mouse Serum: Evidence for Extragonadal Production. *Biochemical and Biophysical Research Communications*, 188(2):921-926 (1992).
- Sakai, et al., "Osteogenic Activity of Activin in Young Normal Rats and Young Adult and Aged Rats after Ovarectomy," *Bone* 23:(Suppl.) 467 (1998).
- Sako, D., et al., "Characterization of the Ligand Binding Functionality of the Extracellular Domain of Activin Receptor Type IIB," *The Journal of Biological Chemistry*, 285(27):21037-21048 (2010).
- Satoh, et al., "Hemodynamic changes by recombinant erythropoietin therapy in hemodialyzed patients," *Hypertension*, 15(3):262-266 (1990).
- Schmelzer, C.H., et al., "Purification and Characterization of Recombinant Human Activin B," *Biochimica et Biophysica Acta*, 1039(2):135-141 (1990).
- Schuelke, M., et al., "Myostatin Mutation Associated with Gross Muscle Hypertrophy in a Child," *New England Journal of Medicine*, 350(26):2682-2688 (2004).
- Seale et al., "PRDM16 controls a brown fat/skeletal muscle switch," *Nature*, vol. 454(7207):961-967 (2008).
- Shao, L., et al., "Effect of Activin A on Globin Gene Expression in Purified Human Erythroid Progenitors," *Blood*, 79(3):773-781 (1992).
- Shao, L., et al., "Efficient synthesis of globoside and isogloboside tetrasaccharides by using beta(1->3)N-acetylgalactosaminyltransferase/UDP-N-acetylglucosamine C4 epimerase fusion protein," *Chem Commun.*: 1422-1423 (2003).
- Shapiro et al., "Side Effects of Adjuvant Treatment of Breast Cancer," *New England Journal of Medicine*, vol. 344: 1997-2008 (2001).
- Shav-Tal, Y., et al., "The Role of Activin A in Regulation of Hemopoiesis," *Stem Cells*, 20:493-500 (2002).
- Shi et al., "Energy Balance, Myostatin, and GILZ: Factors Regulating Adipocyte Differentiation in Belly and Bone," *PPAR Research*, pp. 1-12 (2007).
- Shiozaki, M., et al., "Activin A: A Commitment Factor in Erythroid Differentiation," *Biochemical and Biophysical Research Communications*, 242:631-635 (1998).
- Shiozaki, M., et al., "Evidence for the participation of endogenous activin A/erythroid differentiation factor in the regulation of erythropoiesis," *Proc. Natl. Acad. Sci. USA*, 89:1553-1556 (1992).
- Shiozaki, M., et al., "In Vivo Treatment With Erythroid Differentiation Factor (EDF / Activin A) Increases Erythroid Precursors (CFU-E and BFU-E) in Mice," *Biochemical and Biophysical Research Communications*, 165(3):1155-1161 (1989).
- Shoji et al., "Identification and Characterization of a PDZ Protein That Interacts with Activin Type II Receptors," *The Journal of Biological Chemistry*, vol. 275(8): 5485-5492 (2000).
- Shuto et al., "Osteoblasts Express Types I and II Activin Receptors During Early Intramembranous and Endochondral Bone Formation," *Journal of Bone Mineral Research*, 12(3):403-411 (1997).
- Sirskiy et al., "Detection of influenza A and B neutralizing antibodies in vaccinated ferrets and macaques using specific biotin-streptavidin conjugated antibodies," *J. Virol. Methods*, vol. 163: 459-464 (2010).
- Smith, L. et al., "The analysis of doxorubicin resistance in human breast cancer cells using antibody microarrays," *Mol. Cancer Therapy*, vol. 5: 2115-2120 (2006).
- Smith, L. et al., "The Status, Quality, and Expansion of the NIH Full-Length cDNA Project: The Mammalian Gene Collection (MGC)," *Genome Res.*, vol. 14(10b): 2127-2127 (2004).
- Song, J., et al., "The Type II Activin Receptors are Essential for Egg Cylinder Growth, Gastrulation, and Rostral Head Development in Mice," *Development Biology*, 213:157-169 (1999).
- Springer, et al., "Seventh European Congress on Clinical and Economic Aspects of Osteoporosis and Osteoarthritis," *Osteoporosis International*, 18(1):S29-S75 (2007).
- Sun Shuhan et al., "Chromosome, Gene, and Disease," *Science Press* (2009).
- Sun, et al., "FSH Directly Regulates Bone Mass," *Cell*, 125:247-260 (2006).
- Suragani et al., "4236 ACE-536, a Modified Type II Activin Receptor Increases Red Blood Cells in Vivo by Promoting Maturation of Late Stage Erythroblasts," 52nd ASH Annual Meeting and Expositions, Orange County Convention Center, Orlando, FL Dec. 4-7, 2010.
- Suragani et al., "Transforming growth factor- β superfamily ligand trap ACE-536 corrects anemia by promoting late-state erythropoiesis," *Letters, Nature Medicine*, Advance Online Publication (44 pages) (2014).
- Swanson et al., "Use of Biosensors to Monitor the Immune Response," *Biologics*, vol. 109: 71-78 (2000).
- Swanson, S. J., "New Technologies for the Detection of Antibodies to Therapeutic Proteins," *Immunogenicity of Therapeutics Biological Products*, vol. 112: 127-133 (2003).
- Tanno, T. and Miller, J.L., "Iron Loading and Overloading due to Ineffective Erythropoiesis," *Advances in Hematology*, Article ID 358283, Chapter 2 (Abstract) (2010).
- The Merck Manual of Diagnosis and Therapy, 17th Edition. *Nyelodysplastic Syndrome*, pp. 865 and 963-955 (1999).
- The website downloaded Oct. 28, 2014 from the Multiple Myeloma Research Foundation, themmr.org/multiple-myeloma/symptoms/bone-lesions/, 2 pages total.
- Thompson, et al., "Structures of an ActRIIB: activin A complex reveal a novel binding mode for TGF-beta ligand: receptor interactions," *EMBO* 22(7):1555-1566 (2003).
- Thompson, T.B., et al., "Beta A versus beta B: is it merely a matter of express?," *Molecular and Cellular Endocrinology*, 225:9-17 (2004).
- Thorpe and Swanson, "Current methods for Detecting Antibodies against Erythropoietin and Other Recombinant Proteins," *Clinical and Diagnostic Laboratory Immunology*, vol. 12(1): 28-39 (2005).
- Tinsley, J., et al., "Expression of full-length utrophin prevents muscular dystrophy in mdx mice," *Nature Medicine*, 4(12):1441-1444 (1998).
- Tisdale, M.J., "Cachexia in Cancer Patients," *Nat. Rev. Cancer*, 2:862-871 (2002).
- Tokuriki, N., et al., "Stability effects of mutations and protein evolvability," *Current Opinion in Structural Biology*, 19:596-604 (2009).
- Trivedi, R., et al., "Investigational Anabolic Therapies for Osteoporosis," *Expert Opin. Investig. Drugs*, 19(8):995-1005 (2010).
- Truska et al., "Bone morphogenetic proteins 2, 4, and 9 stimulate murine hepcidin 1 expression independently of Hfe, transferrin receptor 2 (Tfr2), and IL-6," *PNAS*, vol. 103(27):10289-10293 (2006).
- Tseng, Yu-Hua et al., "New role of bone morphogenetic protein 7 in brown adipogenesis and energy expenditure," *Nature: International Weekly Journal of Science (and Supplementary Information)*, vol. 454(7207): 1000-1004 (2008).
- Tsuchida, et al., "Activin isoforms signal through type I receptor serine/threonine kinase ALK7," *Molecular and Cellular Endocrinology*, 220:59-65 (2004).
- Tu, P., et al., "Genetic Disruption of Myostatin Reduces the Development of Proatherogenic Dyslipidemia and Atherogenic Lesions in Ldlr Null Mice," *Diabetes*, 58:1739-1748 (2009).
- Type 2 Diabetes, *PubMed Health, Diseases and Conditions*, U.S. National Library of Medicine, Bethesda, MD (online), Jun. 28, 2011 [retrieved on Jun. 6, 2012]. Retrieved from the Internet<URL:<http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0001356/>>.
- Ukkola, et al., "Adiponectin: A Link Between Excess Adiposity and Associated Comorbidities?," *Journal of Molecular Medicine*, 80(11):696-702 (2002).
- US Biological Technical Data Sheet for A0856-10A, accessed on Feb. 20, 2013.
- US Biological, Activin Receptor Type IIA (RIIA) A0856-05E www.usbio.net/technical-sheet.php?item=A0856-05E dated Jun. 8, 2010.
- Utzschneider, et al., "The Role of Insulin Resistance in Nonalcoholic Fatty Liver Disease," *J. Clin. Endocrinol. Metab.*, 91(12):4753-4761 (Dec. 2006). Epub Sep. 12, 2006.
- Vajdos et al., "Comprehensive Functional Maps of the Antigen-binding Site of an Anti-ErbB2 Antibody Obtained with Shotgun Scanning mutagenesis," *J. Mol. Biol.*, vol. 320(2): 415-428 (2002).

(56)

References Cited**OTHER PUBLICATIONS**


- Vallet, S., et al., "Activin A promotes multiple myeloma-induced osteolysis and is a promising target for myeloma bone disease," PNAS, 107(11):5124-5129 (2010).
- Vidal, L., et al., "Making sense of antisense," European Journal of Cancer, 41:2812-2818 (2005).
- Wagner, K.R., et al., "A Phase I/II trial of MYO-029 in Adult Subjects with Muscular Dystrophy," Ann. Neurol., 63:561-571 (2008).
- Wagner, K.R., et al., "Loss of Myostatin Attenuates Severity of Muscular Dystrophy in mdx Mice," Ann. Neurol., 52:832-836 (2002).
- Wagner, K.R., et al., "Muscle regeneration in the prolonged absence of myostatin," PNAS, 102(7):2519-2524 (2005).
- Walsh, F. S., et al., "Myostatin: a modulator of skeletal-muscle stem cells," Biochemical Society Transactions, 33(Pt.6):1513-1517 (2005).
- Wang, et al., A single amino acid determines lysophospholipid specificity of the S1P1 (EDG1) and LPA1 (EDG2) phospholipid growth factor receptors. JBC 276:49213-49220 (2001).
- Wang, W., et al., "GDF-3 is an adipogenic cytokine under high fat dietary condition," Biochemical and Biophysical Research Comm., 321(4):1024-1031 (2004).
- Ward, R., "An update on disordered iron metabolism and iron overload," Hematology, vol. 15(5):311-317 (2010).
- Ware, Russell E., "How I use hydroxyurea to treat young patients with sickle cell anemia," Blood, vol. 115(26): 5300-5311 (2010).
- Weber, et al., A silent H-bond can be mutationally activated for high-affinity interaction of BMP-2 and activin type IIB receptor, BMC Structural Biology.
- Wells, J.A., "Additivity of Mutational Effects in Proteins," Biochemistry, 29(37):8509-8517 (1990).
- Welt, et al., "Activin: an endocrine or paracrine agent?," European Journal of Endocrinology 139:469-471 (1998).
- Wiater, et al., "Inhibin is an Antagonist of Bone Morphogenetic Protein Signaling," The Journal of Biological Chemistry, 278(10):7934-7941 (2003).
- Wolfman, N.M., et al., "Activation of latent myostatin by the BMP-1/tolloid family of metalloproteinases," PNAS, 100(26):15842-15846 (2003).
- Wong et al., "Validation parameters for a novel biosensor assay which simultaneously measures serum concentrations of a humanized monoclonal antibody and detects induced antibodies," Journal of Immunological Methods, vol. 209: 1-15 (1997).
- Yamato et al., "Induction of apoptosis in Myeloma Cells with Activin A," Japanese Journal of Clinical Hematology; 37th Annual Meeting, Symposium 3, Apoptosis in Blood Disorders, 37:7, pp. 564-567 (2012). (translated).
- Yokota, T., et al., "Isolation and characterization of a mouse cDNA clone that expresses mast-cell growth-factor activity in monkey cells," Proc. Natl. Acad. Sci. USA, 81:1070-1074 (1984).
- Yu et al., "Specific roles of activin/inhibin in human erythropoiesis in vitro," Annals New York Academy of Sciences, 20(10):1243-1246 (1991).
- Yu, J., et al., "Importance of FSH-releasing protein and inhibin in erythrodifferentiation," Nature, 330:765-767 (1987).
- Zatz et al., "Serum creatine-kinase (CK) and pyruvate-kinase (PK) activities in Duchenne (DMD) as compared with Becker (BMD) muscular dystrophy," Journal of the Neurological Sciences, vol. 102: 190-196 (1991).
- Zhang et al., Effects of Activin A on the Activities of the Mouse Peritoneal Macrophages, Cellular & Molecular Immunology, vol. 2(1): 63-67 (2005).
- Zhao, B., et al., "Transgenic expression of myostatin propeptide prevents diet-induced obesity and insulin resistance," Biochemical and Biophysical Research Communications, 337:248-255 (2005).
- Bando, et al., The Journal of Therapy, vol. 89(8): 2499-2504 (2007).
- "Method of examination of anemia of patients with endocrine disease" (translated).
- Kawai et al, Primary care for anemia and polycythemia, the Journal of Therapy, vol. 84(2): 223-229 (2002).
- Levine et al., "Treatment of Acute Postoperative Anemia with Recombinant Human Erythropoietin," The Journal of Trauma, vol. 29(8): 1134-1139 (1989).

ActRIIa ILGRSETQEC IFPMANWEED RTNQTGVFPC YGDKDEPRHC ERTWKNISSGS
ActRIIb GRGEAETREC IITYNANWELE RTNQSGLERC EGEQDKRLHC YASWRNSSGT

IEIVNQGQWL DDINCYDPTD CVERKDSPEV YFCCCEGNNC NEKFSYFFEM
IELVKKGCWL DDFNCDYRQE CVATEENPQV YFCCCEGNFC NERFTHLPEA

EVTQPTSNPV TPKPPT
GGPEVTYEPP PTAPT

FIGURE 1



1 MDAMKRGLCC VLLLCGAVFV SPGASRGEA ETRECIYYNA NWELERTNQS
51 GLERCEGEQD KRLHCYASWR NSSGTIELVK KGCWDDFNC YDRQECVATE
101 ENPQVYFCCC EGNFCNERFT HLPEAGGPEV TYEPPPTAPT GGGTHTCPFC
151 PAPELLGGPS VFLEPPKPKD TLMISRTPEV TCVVVDVSHE DPEVKFNWYV
201 DGVEVHNAKT KPREEQYNST YRVSVLTVL HQDWLNGKEY KCKVSNKALP
251 APIEKTISKA KGQPREPQVY TLPPSREEMT KNQVSLTCLV KGFYPSDIAV
301 EWESNGQPEN NYKTTTPVLD SDGSFFLYSK LTVDKSRWQQ GNVFSCSVMH
351 EALHNHYTQK SLSLSPGK

FIGURE 3

1 ATGGATGCAA TGAAGAGAGG GCTCTGCTGT GTGCTGCTGC TGTGTGGAGC
 TACCTACGTT ACTTCTCTCC CGAGACGACA CACGACGACG ACACACCTCG

51 AGTCTTCGTT TCGCCCGGCG CCTCTGGGCG TGGGGAGGCT GAGACACGGG
 TCAGAAGCAA AGCGGGCCGC GGAGACCCGC ACCCTCCGA CTCTGTGCCC

101 AGTGCATCTA CTACAACGCC AACTGGGAGC TGGAGCGCAC CAACCAGAGC
 TCACGTAGAT GATGTTGCGG TTGACCCTCG ACCTCGCGTG GTTGGTCTCG

151 GGCCTGGAGC GCTGCGAAGG CGAGCAGGAC AAGCGGCTGC ACTGCTACGC
 CCGGACCTCG CGACGCTTCC GCTCGTCCTG TTCGCCGACG TGACGATGCG

201 CTCCTGGGCG AACAGCTCTG GCACCATCGA GCTCGTGAAG AAGGGCTGCT
 GAGGACCGCG TTGTCGAGAC CGTGGTAGCT CGAGCACTTC TTCCCGACGA

251 GGGATGATGA CTTCAACTGC TACGATAGGC AGGAGTGTGT GGCCACTGAG
 CCCTACTACT GAAGTTGACG ATGCTATCCG TCCTCACACA CCGGTGACTC

301 GAGAACCCCC AGGTGTACTT CTGCTGCTGT GAAGGCAACT TCTGCAACGA
 CTCTTGGGGG TCCACATGAA GACGACGACA CTTCCGTGA AGACGTTGCT

351 GCGCTTCACT CATTTGCCAG AGGCTGGGGG CCCGGAAGTC ACGTACGASC
 CGCGAAGTGA GTAAACGGTC TCCGACCCCC GGGCCTTCAG TGCATGCTCG

401 CACCCCGGAC AGCCCCCACC GGTGGTGGAA CTCACACATG CCCACCGTGC
 GTGGGGGCTG TCGGGGGTGG CCACCACCTT GAGTGTGTAC GGGTGGCAGC

451 CCAGCACCTG AACTCCTGGG GGGACCGTCA GTCTTCCTCT TCCCCCAAA
 GGTCGTGGAC TTGAGGACCC CCCTGGCAGT CAGAAGGAGA AGGGGGSTTT

501 ACCCAAGGAC ACCCTCATGA TCTCCCGGAC CCCTGAGGTC ACATGCGTGG
 TGGGTTCCTG TGGGAGTACT AGAGGGCCTG GGGACTCCAG TGTACGCACC

551 TGGTGGACGT GAGCCACGAA GACCCTGAGG TCAAGTTCAA CTGGTACGTG
 ACCACCTGCA CTCGGTGCTT CTGGGACTCC AGTTCAAGTT GACCATGCAC

601 GACGGCGTGG AGGTGCATAA TGCCAAGACA AAGCCGCGGG AGGAGCAGTA
 CTGCCGCACC TCCACGATT ACGGTTCTGT TTCGGCGCCC TCCTCGTCAT

651 CAACAGCACG TACCGTGTGG TCAGCGTCCT CACCGTCCTG CACCAGGACT
 GTGTCGTGC ATGGCACACC AGTCGCAGGA GTGGCAGGAC GTGGTCTTGA

701 GGCTGAATGG CAAGGAGTAC AAGTGCAAGG TCTCCAACAA AGCCCTCCCA
 CCGACTTACC GTTCCTCATG TTCACGTTCC AGAGGTTGTT TCGGGAGGGT

751 GCCCCCATCG AGAAAACCAT CTCCAAAGCC AAAGGGCAGC CCCGAGAACC
 CGGGGGTAGC TCTTTTGGTA GAGGTTTCGG TTTCCCGTCG GGGCTCTTGG

FIGURE 4A

801 ACAGGTGTAC ACCCTGCCCC CATCCCGGGA GGAGATGACC AAGAACCAGG
TGTCCACATG TGGGACGGGG GTAGGGCCCT CCTCTACTGG TTCTTGGTCC

851 TCAGCCTGAC CTGCCTGGTC AAAGGCTTCT ATCCCAGCGA CATCGCCGTG
AGTCGGACTG GACGGACCAG TTTCCGAAGA TAGGGTCGCT GTAGCGGCAC

901 GAGTGGGAGA GCAATGGGCA GCCGGAGAAC AACTACAAGA CCACGCCTCC
CTCACCTCTT CGTTACCCGT CGGCCTCTTG TTGATGTTCT GGTGCGGAGG

951 CGTGCTGGAC TCCGACGGCT CCTTCTTCCT CTATAGCAAG CTCACCGTGG
GCACGACCTG AGGCTGCCGA GGAAGAAGGA GATATCGTTC GAGTGGCACC

1001 ACAAGAGCAG GTGGCAGCAG GGAACGTCT TCTCATGCTC CGTGATGCAT
TGTTCTCGTC CACCGTCGTC CCCTTGCAGA AGAGTACGAG GCACCTACGTA

1051 GAGGCTCTGC ACAACCACTA CACGCAGAAG AGCCTCTCCC TGTCCCCGGG
CTCCGAGACG TGTTGGTGAT GTGCGTCTTC TCGGAGAGGG ACAGGGGCCC

1101 TAAATGA (SEQ ID NO:25)
ATTTACT (SEQ ID NO:33)

FIGURE 4B

1 MDAMKRGGLCC VLLLCGAVFV SPGAAITREC IYYNANWELE RTNQSGLERC
51 EGEQDKRLHC YASWRNSSGT IELVKKGCWDDFNCYDRQE CVATEENPQV
101 YFCCCEGNFC NERFTHLPEA GGPEVTYEPP PTGGGTHTCP PCPAPELLGG
151 PSVFLFPPKP KDTLMIS RTP EVTCVVVDVS HEDPEVKFNW YVDGVEVHNA
201 KTKPREEQYN STYRVVSVLT VLNQDNLNGK EYKCKVSNKA LPAPIEKTIS
251 KAKGQPREPQ VYTLPPSREE MTKNQVSLTLC LVKGFYPSDI AVEWESNGQP
301 ENNYKTTTPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT
351 QKSLSLSPGK (SEQ ID NO: 26)

FIGURE 5

1 ATGGATGCAA TGAAGAGAGG GCTCTGCTGT GTGCTGCTGC TGTGTGGAGC
TACCTACGTT ACTTCTCTCC CGAGACGACA CACGACGACG ACACACCTCG

51 AGTCTTCGTT TCGCCCGGCG E T R E C I Y Y
TCAGAAGCAA AGCGGGCCGC GCGCACTCTG TGCCCTCAGC TAGATGATGT

101 N A N W E L E R T N Q S G L E R C
ACGCCAACTG GGAGCTGGAG CGCACCAACC AGAGCGGCCT GGAGCGCTGC
TGCGGTTGAC CCTCGACCTC GCGTGGTTGG TCTCGCCGGA CCTCGCGACG

151 E G E Q D K R L H C Y A S W R N S
GAAGGCGAGC AGGACAAGCG GCTGCACTGC TACGCCTCCT GGCGCAACAG
CTTCGCTCG TCCTGTTGCG CGACGTGACG ATGCGGAGGA CCGCGTTGTC

201 S G T I E L V K K G C W D D D F
CTCTGGCACC ATCGAGCTCG TGAAGAAGGG CTGCTGGGAC GATGACTTCA
GAGACCGTGG TAGCTCGAGC ACTTCTTCCC GACGACCCTG CTACTGAAGT

251 N C Y D R Q E C V A T E E N P Q V
ACTGCTACGA TAGGCAGGAG TGTGTGGCCA CTGAGGAGAA CCCCCAGGTG
TGACGATGCT ATCCGTCTCT ACACACCGGT GACTCCTCTT GGGGGTCCAC

301 Y F C C C E G N F C W E R F T H L
TACTTCTGCT GCTGTGAAGG CAACTTCTGC AACGAGCGCT TCACTCATT
ATGAAGACGA CGACACTTCC GTTGAAGACG TTGCTCGCGA AGTGAGTAAA

351 P E A G G P E V T Y E F P F T
GCCAGAGGCT GGGGGCCCCG AAGTCACGTA CGAGCCACCC CCGACAGGTG
CGGTCTCCGA CCCCCGGGCC TTCAGTGCAT GCTCGGTGGG GGCTGTCCAC

401 GTGGAAGTCA CACATGCCCA CCGTGCCAG CACCTGAAGT CCTGGGGGGA
CACCTTGAGT GTGTACGGGT GGCACGGGTC GTGGACTTGA GGACCCCCCT

451 CCGTCAGTCT TCCTCTTCCC CCCAAAACCC AAGGACACCC TCATGATCTC
GGCAGTCAGA AGGAGAAGGG GGGTTTTGGG TTCCTGTGGG AGTACTAGAG

501 CCGGACCCCT GAGGTCACAT GCGTGGTGGT GGACGTGAGC CACGAAGACC
GGCCTGGGGA CTCCAGTGTA CGCACACCA CCTGCACTCG GTGCTTCTGG

551 CTGAGGTCAA GTTCAACTGG TACGTGGACG GCGTGGAGGT GCATAATGCC
GACTCCAGTT CAAGTTGACC ATGCACCTGC CGCACCTCCA CGTATTACGG

FIGURE 6A

601 AAGACAAAGC CGCGGGAGGA GCAGTACAAC AGCACGTACC GTGTGGTCAG
TTCTGTTTCG GCGCCCTCCT CGTCATGTTG TCGTGCA TGG CACACCAGTC

651 CGTCCTCACC GTCCTGCACC AGGACTGGCT GAATGGCAAG GAGTACAAGT
GCAGGAGTGG CAGGACGTGG TCCTGACCGA CTTACCGTTC CTCATGTTCA

701 GCAAGGTCTC CAACAAAGCC CTCCCAGCCC CCATCGAGAA AACCATCTCC
CGTTCCAGAG GTTGTTTCGG GAGGGTCGGG GGTAGCTCTT TTGGTAGAGG

751 AAAGCCAAAG GGCAGCCCCG AGAACCACAG GTGTACACCC TGCCCCATC
TTTCGGTTTC CCGTCGGGGC TCTTGGTGTC CACATGTGGG ACGGGGGTAG

801 CCGGGAGGAG ATGACCAAGA ACCAGGTCAG CCTGACCTGC CTGGTCAAAG
GGCCCTCCTC TACTGTTTCT TGGTCCAGTC GGACTGGACG GACCAGTTTC

851 GCTTCTATCC CAGCGACATC GCGTGGAGT GGGAGAGCAA TGGGSCAGCG
CGAAGATAGG GTCGCTGTAG CCGCACCTCA CCCTCTCGTT ACCCGTCGGC

901 GAGAACAAC TACAAGACCAC GCGTCCCGTG CTGGACTCCG ACGGCTCCTT
CTCTTGTTGA TGTTCTGGTG CCGAGGGCAC GACCTGAGGC TGCCGAGSAA

951 CTTCCTCTAT AGCAAGCTCA CCGTGGACAA GAGCAGGTGG CAGCAGGGGA
GAAGGAGATA TCGTTCGAGT GGCACCTGTT CTCGTCCACC GTCGTCCCT

1001 ACGTCTTCTC ATGCTCCGTG ATGCATGAGG CTCTGCACAA CCACTACACG
TGCAGAAGAG TACGAGGCAC TACGTACTCC GAGACGTGTT GGTGATGTGC

1051 CAGAAGAGCC TCTCCCTGTC CCCGGGTAAA TGA (SEQ ID NO: 27)
GTCTTCTCGG AGAGGGACAG GGGCCCATTT ACT (SEQ ID NO: 34)

FIGURE 6B

1 ETRECIYYNA NWELERTNQS GLERCEGEQD KRLHCYASWR NSSGTIELVK
51 KGCWDDDFNC YDRQECVATE ENPQVYFCCC EGNFCNERFT HLPEAGGPEV
101 TYEPPPTGGG THTCPPCPAP ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV
151 VVDVSHEDPE VKFNWYVDGV EVHNAKTKPR EEQYNSTYRV VSVLTVLHQD
201 WLNGKEYKCK VSNKALPAPI EKTISKAKGQ PREPQVYTLF PSREEMTKNQ
251 VSLTCLVKGF YPSDIAVEWE SNGQPENNYK TTPFVLDSDG SFFLYSKLTV
301 DKSRWQQGNV FSCSVMHEAL HNHYTQKSLS LSPGK (SEQ ID NO: 28)

FIGURE 7

1 ETRECIYYNA NWELERTNQS GLERCEGEQD KRLHCYASWR NSSGTIELVK
51 KGCWDDDFNC YDRQECVATE ENPQVYFCCC EGNFCNERFT HLPEAGGPEV
101 TYEPPPT (SEQ ID NO: 29)

FIGURE 8

1 ATGGATGCAA TGAAGAGAGG GCTCTGCTGT GTGCTGCTGC TGTGTGGAGC
TACCTACGTT ACTTCTCTCC CGAGACGACA CACGACGACG ACACACCTCG

51 AGTCTTCGTT TCGCCCGGCG CCGCCGAAAC CCGGATGT ATTTATTACA
TCAGAAGCAA AGCGGGCCGC GSCGGCTTTG GCGCCTTACA TAAATAATGT

101 N A N W E L E R T N Q S G L E R C
ATGCTAATTG GGAATCTGA CCGACGAACC AATCGGCT GAAAGCTG
TACGATTAAC CTTGAGCTT GCCTGCTTGG TTAGGCCCGA GCTTGCCACA

151 E G E Q D K R L H C Y A S W R N S
GAAGGGAAC AGGAATAACG CTTCAATGC TATGCTCTT GGAGAACG
CTCCCCCTTG TCCTATTTGC GSAGGTAACG ATACGCAGCA CCTCCTTGAG

201 S G T I E L V K K G C W D D D F
CTCAGGAC ATGAATCTG TAAAGAAAGG TGTGTTGGAC GAAGATTCA
GAGGCCCTGC TAACTTGACC AGTTCTTTCC CACGACCCTG CTGCTAAAGT

251 N C Y D R Q E C V A T E E N P Q V
ATGTATTGA GGCAGGA TGTGTGCA CAGAGAGAA CCGAGGTG
TAACAATACT GCGGTCCTT ACACAGCGCT GGCTTCTCTT AGGCGTCCAG

301 Y F C C C E G N F C N E R F T H L
TATTTCTGTT GTGCGAGGG GAATTTCTGT AATGAAGGT TTACACCT
ATAAAGACAA CAACGCTCCC CTTAAAGACA TTACTTGCCA AATGGGTGGA

351 P E A G G P E V T Y E P P P T
CCCGAAGCG GCGGGCCCG AGGTACCTA GAGCCCG CCGACGGTG
GGGGCTTCGG CCGCCCGGGC TCACTGGAT ACTTGGGGC GGGTGGCCAC

401 GTGGAAGTCA CACATGCCCA CCGTGCCAG CACCTGAAGT CCTGGGGGGA
CACCTTGAGT GTGTACGGGT GGCACGGGTC GTGGAAGTGA GGACCCCCCT

451 CCGTCAGTCT TCCTCTTCCC CCCAAAACCC AAGGACACCC TCATGATCTC
GGCAGTCAGA AGGAGAAGGG GGGTTTGGG TTCCTGTGGG AGTACTAGAG

501 CCGGACCCCT GAGGTACAT GCGTGGTGGT GGACGTGAGC CACGAAGACC
GGCCTGGGGA CTCCAGTGTA CACACCACCA CCTGCACTCG GTGCTTCTGG

551 CTGAGGTCAA GTTCAACTGG TACGTGGACG GCGTGGAGGT GCATAATGCC
GACTCCAGTT CAAGTTGACC ATGCACCTGC CGCACCTCCA CGTATTACGG

601 AAGACAAAGC CCGGGGAGGA GCAGTACAAC AGCAGTACC GTGTGGTACG
TTCTGTTTCG GCGCCCTCCT CGTGATGTTG TCGTGCATGG CACACCAAGT

651 CGTCCTCACC GTCTGCACC AGGACTGGCT GAATGGCAAG GAGTACAAGT
GCAGGAGTGG CAGGACGTGG TCCTGACCGA CTACCGTTC CTGATGTTCA

FIGURE 9A

701 GCAAGGTCTC CAACAAAGCC CTCCCAGCCC CCATCGAGAA AACCATCTCC
CGTTCCAGAG GTTGTTTCGG GAGGGTCGGG GGTAGCTCTT TTGGTAGAGG

751 AAAGCCAAAG GGCAGCCCCG AGAACCACAG GTGTACACCC TGCCCCCATC
TTTCGGTTTC CCGTCGGGGC TCTTGGTGTC CACATGTGGG ACGGGGGTAG

801 CCGGGAGGAG ATGACCAAGA ACCAGGTCAG CCTGACCTGC CTGGTCAAAG
GGCCCTCCTC TACTGGTTCT TGGTCCAGTC GGACTGGACG GACCAGTTTC

851 GCTTCTATCC CAGCGACATC GCCGTGGAGT GGGAGAGCAA TGGGCAGCCG
CGAAGATAGG GTCGCTGTAG CGGCACCTCA CCCTCTCGTT ACCCGTCGGC

901 GAGAACAAC TACAAGACCAC GCCTCCCGTG CTGGACTCCG ACGGCTCCTT
CTCTTGTTGA TGTTCTGGTG CGGAGGGCAC GACCTGAGGC TGCCGAGGAA

951 CTTCCTCTAT AGCAAGCTCA CCGTGGACAA GAGCAGGTGG CAGCAGGGGA
GAAGGAGATA TCGTTCGAGT GGCACCTGTT CTCGTCCACC GTCGTCCCTT

1001 ACGTCTTCTC ATGCTCCGTG ATGCATGAGG CTCTGCACAA CCACTACACG
TGCAGAAGAG TACGAGGCAC TACGTACTCC GAGACGTGTT GGTGATGTGC

1051 CAGAAGAGCC TCTCCCTGTC CCCGGGTAAA TGA (SEQ ID NO: 30)
GTCTTCTCGG AGAGGGACAG GGGCCCATTT ACT (SEQ ID NO: 35)

FIGURE 9B

GAAAC CCGGAACTGT ATTAATACA ATGCTAAATG GGAAGTGA CAGACGAACC
AATCCGGCT CAGACGCTGT GAGGGGAAC AGGAATAACG CCTCCATTGC TATGCGTCGT
GGAGGAACTC CTCGGGACG ATGAAGTGG TCAAGAAAGG GTGCTGGGAC GAGGATTTCA
ATTGTTATGA CCGGCAGGAA TGTGTGCGA CAGAGAGAA TCCGAGGT TATTTCTGTT
GTGCGAGGG AATTTCTGT AATGAACGCT TTACCACTT CCGGAAGCT GGCGGCCCG
ACGTACCTA TGAACCCG CCGAC (SEQ ID NO: 31)

FIGURE 10




FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14




FIGURE 15




FIGURE 16




FIGURE 17




FIGURE 18

1

METHODS FOR TREATING ANEMIA IN A SUBJECT IN NEED THEREOF

RELATED APPLICATIONS

This application is a continuation of U.S. application Ser. No. 16/455,301, filed Jun. 27, 2019, which is a divisional of U.S. application Ser. No. 15/336,363, filed Oct. 27, 2016 (now U.S. Pat. No. 10,377,996), which is a continuation of U.S. application Ser. No. 13/750,249, filed Jan. 25, 2013 (now U.S. Pat. No. 9,505,813), which is a continuation of U.S. application Ser. No. 13/247,748, filed Sep. 28, 2011 (now U.S. Pat. No. 8,361,957), which is a divisional of U.S. application Ser. No. 12/583,177, filed Aug. 13, 2009 (now U.S. Pat. No. 8,058,229), which claims the benefit of U.S. Provisional Application Ser. No. 61/189,094, filed Aug. 14, 2008. The specifications of each of the foregoing applications are incorporated herein by reference in their entirety.

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Feb. 19, 2020, is named 1848179-0002-040-110_Seq.txt, and is 71,080 bytes in size.

BACKGROUND OF THE INVENTION

The mature red blood cell, or erythrocyte, is responsible for oxygen transport in the circulatory systems of vertebrates. Red blood cells carry high concentrations of hemoglobin, a protein that binds oxygen in the lungs at relatively high partial pressure of oxygen (pO_2) and delivers oxygen to areas of the body with a relatively low pO_2 .

Mature red blood cells are produced from pluripotent hematopoietic stem cells in a process termed erythropoiesis. Postnatal erythropoiesis occurs primarily in the bone marrow and in the red pulp of the spleen. The coordinated action of various signaling pathways control the balance of cell proliferation, differentiation, survival and death. Under normal conditions, red blood cells are produced at a rate that maintains a constant red cell mass in the body, and production may increase or decrease in response to various stimuli, including increased or decreased oxygen tension or tissue demand. The process of erythropoiesis begins with the formation of lineage committed precursor cells and proceeds through a series of distinct precursor cell types. The final stages of erythropoiesis occur as reticulocytes are released into the bloodstream and lose their mitochondria and ribosomes while assuming the morphology of mature red blood cell. An elevated level of reticulocytes, or an elevated reticulocyte:erythrocyte ratio, in the blood is indicative of increased red blood cell production rates.

Erythropoietin (Epo) is widely recognized as the most significant positive regulator of erythropoiesis in post-natal vertebrates. Epo regulates the compensatory erythropoietic response to reduced tissue oxygen tension (hypoxia) and low red blood cell levels or low hemoglobin levels. In humans, elevated Epo levels promote red blood cell formation by stimulating the generation of erythroid progenitors in the bone marrow and spleen. In the mouse, Epo enhances erythropoiesis primarily in the spleen.

Various forms of recombinant Epo are used by physicians to increase red blood cell levels in a variety of clinical settings, and particularly for the treatment of anemia. Anemia is a broadly-defined condition characterized by lower

2

than normal levels of hemoglobin or red blood cells in the blood. In some instances, anemia is caused by a primary disorder in the production or survival of red blood cells. More commonly, anemia is secondary to diseases of other systems (Weatherall & Provan (2000) *Lancet* 355, 1169-1175). Anemia may result from a reduced rate of production or increased rate of destruction of red blood cells or by loss of red blood cells due to bleeding. Anemia may result from a variety of disorders that include, for example, chronic renal failure, chemotherapy treatment, myelodysplastic syndrome, rheumatoid arthritis, and bone marrow transplantation.

Treatment with Epo typically causes a rise in hemoglobins by about 1-3 g/dL in healthy humans over a period of weeks. When administered to anemic individuals, this treatment regimen often provides substantial increases in hemoglobin and red blood cell levels and leads to improvements in quality of life and prolonged survival. Epo is not uniformly effective, and many individuals are refractory to even high doses (Horl et al. (2000) *Nephrol Dial Transplant* 15, 43-50). Over 50% of patients with cancer have an inadequate response to Epo, approximately 10% with end-stage renal disease are hyporesponsive (Glaspy et al. (1997) *J Clin Oncol* 15, 1218-1234; Demetri et al. (1998) *J Clin Oncol* 16, 3412-3425), and less than 10% with myelodysplastic syndrome respond favorably (Estey (2003) *Curr Opin Hematol* 10, 60-67). Several factors, including inflammation, iron and vitamin deficiency, inadequate dialysis, aluminum toxicity, and hyperparathyroidism may predict a poor therapeutic response, the molecular mechanisms of resistance to Epo are as yet unclear.

Thus, it is an object of the present disclosure to provide alternative compositions and methods for increasing red blood cell levels in patients.

SUMMARY OF THE INVENTION

In part, the disclosure demonstrates that GDF Traps may be used to increase red blood cell and hemoglobin levels. Variant ActRIIB polypeptides having a significantly decreased affinity for activin (e.g., activin A and/or activin B) relative to other ActRIIB ligands, such as GDF11 and/or myostatin, are referred to as GDF Traps. ActRIIB variants described herein are GDF Traps unless otherwise stated. In particular, the disclosure demonstrates that a GDF Trap which is a soluble form of ActRIIB polypeptide having an acidic residue at position 79 of SEQ ID NO: 1, when administered in vivo, increases red blood cell levels in the blood. Therefore, in certain embodiments, the disclosure provides methods for using GDF Traps to increase red blood cell and hemoglobin levels in patients and to treat disorders associated with low red blood cell or hemoglobin levels in patients in need thereof. As described in U.S. patent application Ser. No. 12/012,652, incorporated by reference herein, GDF Traps can be used to increase muscle mass and decrease fat mass.

In certain aspects, the present disclosure provides GDF Traps that are variant ActRIIB polypeptides, including ActRIIB polypeptides having amino- and carboxy-terminal truncations and sequence alterations. Optionally, GDF Traps of the invention may be designed to preferentially antagonize one or more ligands of ActRIIB receptors, such as GDF8 (also called myostatin), GDF11, Nodal, and BMP7 (also called OP-1). Examples of GDF Traps include a set of variants derived from ActRIIB that have greatly diminished affinity for activin. These variants exhibit desirable effects on red blood cells while reducing effects on other tissues.

Examples of such variants include those having an acidic amino acid (e.g., aspartic acid, D, or glutamic acid, E) at the position corresponding to position 79 of SEQ ID NO. 1. In certain embodiments, the GDF Trap polypeptide comprises an amino acid sequence that comprises, consists of, or consists essentially of, the amino acid sequence of SEQ ID NO: 7, 26, 28, 29, 32, 37 or 38, and polypeptides that are at least 80%, 85%, 90%, 95%, 97%, 98%, or 99% identical to any of the foregoing.

In certain aspects, the disclosure provides pharmaceutical preparations comprising a GDF Trap that binds to an ActRIIB ligand such as GDF8, GDF11, activin (e.g., activin B), BMP7 or nodal, and a pharmaceutically acceptable carrier. Optionally, the GDF Trap binds to an ActRIIB ligand with a K_d less than 10 micromolar, less than 1 micromolar, less than 100 nanomolar, less than 10 nanomolar, or less than 1 nanomolar. Optionally, the GDF Trap inhibits ActRIIB signaling, such as intracellular signal transduction events triggered by an ActRIIB ligand. A GDF Trap for use in such a preparation may be any of those disclosed herein, including, for example, GDF Traps having an amino acid sequence selected from SEQ ID NOs: 2, 3, 7, 11, 26, 28, 29, 32, 37, 38 or 40, or GDF Traps having an amino acid sequence that is at least 80%, 85%, 90%, 95%, 97% or 99% identical to an amino acid sequence selected from SEQ ID NOs: 2, 3, 7, 11, 26, 28, 29, 32, 37, 38 or 40, or GDF Traps having an amino acid sequence that is at least 80%, 85%, 90%, 95%, 97% or 99% identical to an amino acid sequence selected from SEQ ID NOs: 2, 3, 7, 11, 26, 28, 29, 32, 37, 38 or 40 wherein the position corresponding to L79 in SEQ ID NO: 1 is an acidic amino acid. A preferred GDF Trap for use in such a preparation consists of, or consists essentially of, the amino acid sequence of SEQ ID NO: 26. A GDF Trap may include a functional fragment of a natural ActRIIB polypeptide, such as one comprising at least 10, 20 or 30 amino acids of a sequence selected from SEQ ID NOs: 2, 3, 7, 11, 26, 28, 29, 32, 37, 38 or 40 or a sequence of SEQ ID NO: 2, lacking the C-terminal 1, 2, 3, 4, 5 or 10 to 15 amino acids and lacking 1, 2, 3, 4 or 5 amino acids at the N-terminus. A preferred polypeptide will comprise a truncation relative to SEQ ID NO: 2 or 40 of between 2 and 5 amino acids at the N-terminus and no more than 3 amino acids at the C-terminus. A GDF Trap may include one or more alterations in the amino acid sequence of an ActRIIB polypeptide (e.g., in the ligand-binding domain) relative to a naturally occurring ActRIIB polypeptide. The alteration in the amino acid sequence may, for example, alter glycosylation of the polypeptide when produced in a mammalian, insect or other eukaryotic cell or alter proteolytic cleavage of the polypeptide relative to the naturally occurring ActRIIB polypeptide.

A GDF Trap may be a fusion protein that has, as one domain, an ActRIIB polypeptide (e.g., a ligand-binding domain of an ActRIIB with one or more sequence variations) and one or more additional domains that provide a desirable property, such as improved pharmacokinetics, easier purification, targeting to particular tissues, etc. For example, a domain of a fusion protein may enhance one or more of in vivo stability, in vivo half life, uptake/administration, tissue localization or distribution, formation of protein complexes, multimerization of the fusion protein, and/or purification. GDF Trap fusion proteins may include an immunoglobulin Fc domain (wild-type or mutant) or a serum albumin. In certain embodiments, a GDF Trap fusion comprises a relatively unstructured linker positioned between the Fc domain and the extracellular ActRIIB domain. This unstructured linker may correspond to the roughly 15 amino acid unstructured region at the C-terminal

end of the extracellular domain of ActRIIB (the "tail"), or it may be an artificial sequence of between 3 and 5, 15, 20, 30, 50 or more amino acids that are relatively free of secondary structure. A linker may be rich in glycine and proline residues and may, for example, contain repeating sequences of threonine/serine and glycines (e.g., TG4 (SEQ ID NO: 13) or SG₄ (SEQ ID NO: 14) singlets or repeats) or a series of three glycines. A fusion protein may include a purification subsequence, such as an epitope tag, a FLAG tag, a polyhistidine sequence, and a GST fusion. In certain embodiments, a GDF Trap fusion comprises a leader sequence. The leader sequence may be a native ActRIIB leader sequence or a heterologous leader sequence. In certain embodiments, the leader sequence is a Tissue Plasminogen Activator (TPA) leader sequence. In an embodiment, a GDF Trap fusion protein comprises an amino acid sequence as set forth in the formula A-B-C. The B portion is an N- and C-terminally truncated ActRIIB polypeptide consisting of the amino acid sequence corresponding to amino acids 25-131 of SEQ ID NO: 2 or 40. The A and C portions may be independently zero, one or more than one amino acids, and both A and C portions are heterologous to B. The A and/or C portions may be attached to the B portion via a linker sequence.

Optionally, a GDF Trap includes a variant ActRIIB polypeptide having one or more modified amino acid residues selected from: a glycosylated amino acid, a PEGylated amino acid, a farnesylated amino acid, an acetylated amino acid, a biotinylated amino acid, an amino acid conjugated to a lipid moiety, and an amino acid conjugated to an organic derivatizing agent. A pharmaceutical preparation may also include one or more additional compounds such as a compound that is used to treat an ActRIIB-associated disorder. Preferably, a pharmaceutical preparation is substantially pyrogen free. In general, it is preferable that a GDF Trap be expressed in a mammalian cell line that mediates suitably natural glycosylation of the GDF Trap so as to diminish the likelihood of an unfavorable immune response in a patient. Human and CHO cell lines have been used successfully, and it is expected that other common mammalian expression vectors will be useful.

In certain aspects, the disclosure provides packaged pharmaceuticals comprising a pharmaceutical preparation described herein and labeled for use in increasing red blood cell levels in a human.

In certain aspects, the disclosure provides GDF Traps which are soluble ActRIIB polypeptides comprising an altered ligand-binding (e.g., GDF8-binding) domain. GDF Traps with altered ligand-binding domains may comprise, for example, one or more mutations at amino acid residues such as E37, E39, R40, K55, R56, Y60, A64, K74, W78, L79, D80, F82 and F101 of human ActRIIB (numbering is relative to SEQ ID NO: 1). Optionally, the altered ligand-binding domain can have increased selectivity for a ligand such as GDF8/GDF11 relative to a wild-type ligand-binding domain of an ActRIIB receptor. To illustrate, these mutations are demonstrated herein to increase the selectivity of the altered ligand-binding domain for GDF11 (and therefore, presumably, GDF8) over activin: K74Y, K74F, K74I, L79D, L79E, and D80I. The following mutations have the reverse effect, increasing the ratio of activin binding over GDF11: D54A, K55A, L79A and F82A. The overall (GDF11 and activin) binding activity can be increased by inclusion of the "tail" region or, presumably, an unstructured linker region, and also by use of a K74A mutation. Other mutations that caused an overall decrease in ligand binding affinity, include: R40A, E37A, R56A, W78A, D80K, D80R, D80A, D80G, D80F, D80M and D80N. Mutations may be com-

bined to achieve desired effects. For example, many of the mutations that affect the ratio of GDF11:Activin binding have an overall negative effect on ligand binding, and therefore, these may be combined with mutations that generally increase ligand binding to produce an improved binding protein with ligand selectivity. In an exemplary embodiment, a GDF Trap is an ActRIIB polypeptide comprising an L79D or L79E mutation, optionally in combination with additional amino acid substitutions, additions or deletions.

Optionally, a GDF Trap comprising an altered ligand-binding domain has a ratio of K_d for activin binding to K_d for GDF8 binding that is at least 2, 5, 10, or even 100 fold greater relative to the ratio for the wild-type ligand-binding domain. Optionally, the GDF Trap comprising an altered ligand-binding domain has a ratio of IC_{50} for inhibiting activin to IC_{50} for inhibiting GDF8/GDF11 that is at least 2, 5, 10, or even 100 fold greater relative to the wild-type ActRIIB ligand-binding domain. Optionally, the GDF Trap comprising an altered ligand-binding domain inhibits GDF8/GDF11 with an IC_{50} at least 2, 5, 10, or even 100 times less than the IC_{50} for inhibiting activin. These GDF Traps can be fusion proteins that include an immunoglobulin Fc domain (either wild-type or mutant). In certain cases, the subject soluble GDF Traps are antagonists (inhibitors) of GDF8 and/or GDF11.

Other GDF Traps are contemplated, such as the following. A GDF Trap fusion protein comprising a portion derived from the ActRIIB sequence of SEQ ID NO: 1 or 39 and a second polypeptide portion, wherein the portion derived from ActRIIB corresponds to a sequence beginning at any of amino acids 21-29 of SEQ ID NO: 1 or 39 (optionally beginning at 22-25 of SEQ ID NO: 1 or 39) and ending at any of amino acids 109-134 of SEQ ID NO: 1 or 39, and wherein the GDF Trap fusion protein inhibits signaling by activin, myostatin and/or GDF11 in a cell-based assay. The GDF Trap fusion protein above, wherein the portion derived from ActRIIB corresponds to a sequence beginning at any of amino acids 20-29 of SEQ ID NO: 1 or 39 (optionally beginning at 22-25 of SEQ ID NO: 1 or 39) and ending at any of amino acids 109-133 of SEQ ID NO: 1 or 39. The GDF Trap fusion protein above, wherein the portion derived from ActRIIB corresponds to a sequence beginning at any of amino acids 20-24 of SEQ ID NO: 1 or 39 (optionally beginning at 22-25 of SEQ ID NO: 1 or 39) and ending at any of amino acids 109-133 of SEQ ID NO: 1 or 39. The GDF Trap fusion protein above, wherein the portion derived from ActRIIB corresponds to a sequence beginning at any of amino acids 21-24 of SEQ ID NO: 1 or 39 and ending at any of amino acids 109-134 of SEQ ID NO: 1 or 39. The GDF Trap fusion protein above, wherein the portion derived from ActRIIB corresponds to a sequence beginning at any of amino acids 20-24 of SEQ ID NO: 1 or 39 and ending at any of amino acids 118-133 of SEQ ID NO: 1 or 39. The GDF Trap fusion protein above, wherein the portion derived from ActRIIB corresponds to a sequence beginning at any of amino acids 21-24 of SEQ ID NO: 1 or 39 and ending at any of amino acids 118-134 of SEQ ID NO: 1 or 39. The GDF Trap fusion protein above, wherein the portion derived from ActRIIB corresponds to a sequence beginning at any of amino acids 20-24 of SEQ ID NO: 1 or 39 and ending at any of amino acids 128-133 of SEQ ID NO: 1 or 39. The GDF Trap fusion protein above, wherein the portion derived from ActRIIB corresponds to a sequence beginning at any of amino acids 20-24 of SEQ ID NO: 1 or 39 and ending at any of amino acids 128-133 of SEQ ID NO: 1 or 39. The GDF Trap fusion protein above, wherein the portion derived from

ActRIIB corresponds to a sequence beginning at any of amino acids 21-29 of SEQ ID NO: 1 or 39 and ending at any of amino acids 118-134 of SEQ ID NO: 1 or 39. The GDF Trap fusion protein above, wherein the portion derived from ActRIIB corresponds to a sequence beginning at any of amino acids 20-29 of SEQ ID NO: 1 or 39 and ending at any of amino acids 118-133 of SEQ ID NO: 1 or 39. The GDF Trap fusion protein above, wherein the portion derived from ActRIIB corresponds to a sequence beginning at any of amino acids 21-29 of SEQ ID NO: 1 or 39 and ending at any of amino acids 128-134 of SEQ ID NO: 1 or 39. The GDF Trap fusion protein above, wherein the portion derived from ActRIIB corresponds to a sequence beginning at any of amino acids 20-29 of SEQ ID NO: 1 and ending at any of amino acids 128-133 of SEQ ID NO: 1 or 39. Surprisingly, constructs beginning at 22-25 of SEQ ID NO: 1 or 39 have activity levels greater than proteins having the full extracellular domain of human ActRIIB. In a preferred embodiment, the GDF Trap fusion protein comprises, consists essentially of, or consists of, an amino acid sequence beginning at amino acid position 25 of SEQ ID NO: 1 or 39 and ending at amino acid position 131 of SEQ ID NO: 1 or 39. In another preferred embodiment, the GDF Trap polypeptide consists of, or consists essentially of, the amino acid sequence of SEQ ID NO: 7, 26, 28, 29, 32, 37 or 38. Any of the above GDF Trap fusion proteins may be produced as a homodimer. Any of the above GDF Trap fusion proteins may have a heterologous portion that comprises a constant region from an IgG heavy chain, such as an Fc domain. Any of the above GDF Trap fusion proteins may comprise an acidic amino acid at the position corresponding to position 79 of SEQ ID NO: 1, optionally in combination with one or more additional amino acid substitutions, deletions or insertions relative to SEQ ID NO: 1.

Other GDF Trap proteins are contemplated, such as the following. A GDF Trap protein comprising an amino acid sequence that is at least 80% identical to the sequence of amino acids 29-109 of SEQ ID NO: 1 or 39, wherein the position corresponding to 64 of SEQ ID NO: 1 is an R or K, and wherein the GDF Trap protein inhibits signaling by activin, myostatin and/or GDF11 in a cell-based assay. The GDF Trap protein above, wherein at least one alteration with respect to the sequence of SEQ ID NO: 1 or 39 is positioned outside of the ligand binding pocket. The GDF Trap protein above, wherein at least one alteration with respect to the sequence of SEQ ID NO: 1 or 39 is a conservative alteration positioned within the ligand binding pocket. The GDF Trap protein above, wherein at least one alteration with respect to the sequence of SEQ ID NO: 1 or 39 is an alteration at one or more positions selected from the group consisting of K74, R40, Q53, K55, F82 and L79. The GDF Trap protein above, wherein the protein comprises at least one N-X-S/T sequence at a position other than an endogenous N-X-S/T sequence of ActRIIB, and at a position outside of the ligand binding pocket.

Other GDF Traps are contemplated, such as the following. A GDF Trap protein comprising an amino acid sequence that is at least 80% identical to the sequence of amino acids 29-109 of SEQ ID NO: 1 or 39, and wherein the protein comprises at least one N-X-S/T sequence at a position other than an endogenous N-X-S/T sequence of ActRIIB, and at a position outside of the ligand binding pocket. The GDF Trap protein above, wherein the GDF Trap protein comprises an N at the position corresponding to position 24 of SEQ ID NO: 1 or 39 and an S or T at the position corresponding to position 26 of SEQ ID NO: 1 or 39, and wherein the GDF Trap inhibits signaling by activin, myostatin and/or GDF11 in a cell-based

assay. The GDF Trap above, wherein the GDF Trap protein comprises an R or K at the position corresponding to position 64 of SEQ ID NO: 1 or 39. The GDF Trap above, wherein the ActRIIB protein comprises a D or E at the position corresponding to position 79 of SEQ ID NO: 1 or 39 and wherein the GDF Trap inhibits signaling by activin, myostatin and/or GDF11 in a cell-based assay. The GDF Trap above, wherein at least one alteration with respect to the sequence of SEQ ID NO: 1 or 39 is a conservative alteration positioned within the ligand binding pocket. The GDF Trap above, wherein at least one alteration with respect to the sequence of SEQ ID NO: 1 or 39 is an alteration at one or more positions selected from the group consisting of K74, R40, Q53, K55, F82 and L79. The GDF Trap above, wherein the protein is a fusion protein further comprising a heterologous portion. Any of the above GDF Trap fusion proteins may be produced as a homodimer. Any of the above GDF Trap fusion proteins may have a heterologous portion that comprises a constant region from an IgG heavy chain, such as an Fc domain.

In certain aspects, the disclosure provides nucleic acids encoding a GDF Trap polypeptide. An isolated polynucleotide may comprise a coding sequence for a soluble GDF Trap polypeptide, such as described above. For example, an isolated nucleic acid may include a sequence coding for a GDF Trap comprising an extracellular domain (e.g., ligand-binding domain) of an ActRIIB polypeptide having one or more sequence variations and a sequence that would code for part or all of the transmembrane domain and/or the cytoplasmic domain of an ActRIIB polypeptide, but for a stop codon positioned within the transmembrane domain or the cytoplasmic domain, or positioned between the extracellular domain and the transmembrane domain or cytoplasmic domain. For example, an isolated polynucleotide coding for a GDF Trap may comprise a full-length ActRIIB polynucleotide sequence such as SEQ ID NO: 4 having one or more variations, or a partially truncated version, said isolated polynucleotide further comprising a transcription termination codon at least six hundred nucleotides before the 3'-terminus or otherwise positioned such that translation of the polynucleotide gives rise to an extracellular domain optionally fused to a truncated portion of a full-length ActRIIB Nucleic acids disclosed herein may be operably linked to a promoter for expression, and the disclosure provides cells transformed with such recombinant polynucleotides. Preferably the cell is a mammalian cell such as a CHO cell.

In certain aspects, the disclosure provides methods for making a GDF Trap polypeptide. Such a method may include expressing any of the nucleic acids (e.g., SEQ ID NO: 5, 25, 27, 30 or 31) disclosed herein in a suitable cell, such as a Chinese hamster ovary (CHO) cell. Such a method may comprise: a) culturing a cell under conditions suitable for expression of the GDF Trap polypeptide, wherein said cell is transformed with a GDF Trap expression construct; and b) recovering the GDF Trap polypeptide so expressed. GDF Trap polypeptides may be recovered as crude, partially purified or highly purified fractions using any of the well known techniques for obtaining protein from cell cultures.

In certain aspects, a GDF Trap polypeptide disclosed herein may be used in a method for promoting red blood cell production or increasing red blood cell levels in a subject. In certain embodiments, the disclosure provides methods for treating a disorder associated with low red blood cell counts or low hemoglobin levels (e.g., an anemia), or to promote red blood cell production, in patients in need thereof. A method may comprise administering to a subject in need

thereof an effective amount of a GDF Trap polypeptide. In certain aspects, the disclosure provides uses of GDF Trap polypeptides for making a medicament for the treatment of a disorder or condition as described herein.

In certain aspects, the disclosure provides methods for administering a GDF Trap polypeptide to a patient. In part, the disclosure demonstrates that GDF Trap polypeptides can be used to increase red blood cell and hemoglobin levels. GDF Trap polypeptides may also be used for treating or preventing other therapeutic uses such as promoting muscle growth. In certain instances, when administering a GDF Trap polypeptide for promoting muscle growth, it may be desirable to monitor the effects on red blood cells during administration of the GDF Trap polypeptide, or to determine or adjust the dosing of the GDF Trap polypeptide, in order to reduce undesired effects on red blood cells. For example, increases in red blood cell levels, hemoglobin levels, or hematocrit levels may cause increases in blood pressure.

BRIEF DESCRIPTION OF THE DRAWINGS

The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

FIG. 1 shows an alignment of the extracellular domains of human ActRIIA (SEQ ID NO: 15) and human ActRIIB (SEQ ID NO: 2) with the residues that are deduced herein, based on composite analysis of multiple ActRIIB and ActRIIA crystal structures to directly contact ligand (the ligand binding pocket) indicated with boxes.

FIG. 2 shows a multiple sequence alignment of various vertebrate ActRIIB proteins and human ActRIIA (SEQ ID NOs: 16-23).

FIG. 3 shows the full amino acid sequence for the GDF Trap ActRIIB(L79D 20-134)-hFc (SEQ ID NO: 43), including the TPA leader sequence (double underlined), ActRIIB extracellular domain (residues 20-134 in SEQ ID NO: 1; underlined), and hFc domain. The aspartate substituted at position 79 in the native sequence is double underlined and highlighted, as is the glycine revealed by sequencing to be the N-terminal residue in the mature fusion protein.

FIGS. 4A and 4B show a nucleotide sequence encoding ActRIIB(L79D 20-134)-hFc. SEQ ID NO: 25 corresponds to the sense strand, and SEQ ID NO: 33 corresponds to the antisense strand. The TPA leader (nucleotides 1-66) is double underlined, and the ActRIIB extracellular domain (nucleotides 76-420) is underlined.

FIG. 5 shows the full amino acid sequence for the truncated GDF Trap ActRIIB(L79D 25-131)-hFc (SEQ ID NO: 26), including the TPA leader (double underlined), truncated ActRIIB extracellular domain (residues 25-131 in SEQ ID NO: 1; underlined), and hFc domain. The aspartate substituted at position 79 in the native sequence is double underlined and highlighted, as is the glutamate revealed by sequencing to be the N-terminal residue in the mature fusion protein.

FIGS. 6A and 6B show a nucleotide sequence encoding ActRIIB(L79D 25-131)-hFc. SEQ ID NO: 27 corresponds to the sense strand, and SEQ ID NO: 34 corresponds to the antisense strand. The TPA leader (nucleotides 1-66) is double underlined, and the truncated ActRIIB extracellular domain (nucleotides 76-396) is underlined. The amino acid sequence for the ActRIIB extracellular domain (residues 25-131 in SEQ ID NO: 1) is also shown (SEQ ID NO: 44).

FIG. 7 shows the amino acid sequence for the truncated GDF Trap ActRIIB(L79D 25-131)-hFc without a leader

(SEQ ID NO: 28). The truncated ActRIIB extracellular domain (residues 25-131 in SEQ ID NO: 1) is underlined. The aspartate substituted at position 79 in the native sequence is double underlined and highlighted, as is the glutamate revealed by sequencing to be the N-terminal residue in the mature fusion protein.

FIG. 8 shows the amino acid sequence for the truncated GDF Trap ActRIIB(L79D 25-131) without the leader, hFc domain, and linker (SEQ ID NO: 29). The aspartate substituted at position 79 in the native sequence is underlined and highlighted, as is the glutamate revealed by sequencing to be the N-terminal residue in the mature fusion protein.

FIGS. 9A and 9B show an alternative nucleotide sequence encoding ActRIIB(L79D 25-131)-hFc. SEQ ID NO: 30 corresponds to the sense strand, and SEQ ID NO: 35 corresponds to the antisense strand. The TPA leader (nucleotides 1-66) is double underlined, the truncated ActRIIB extracellular domain (nucleotides 76-396) is underlined, and substitutions in the wildtype nucleotide sequence of the extracellular domain are double underlined and highlighted (compare with SEQ ID NO: 27, FIGS. 6A and 6B). The amino acid sequence for the ActRIIB extracellular domain (residues 25-131 in SEQ ID NO: 1) is also shown (SEQ ID NO: 44).

FIG. 10 shows nucleotides 76-396 (SEQ ID NO: 31) of the alternative nucleotide sequence shown in FIGS. 9A and 9B (SEQ ID NO: 30). The same nucleotide substitutions indicated in FIGS. 9A and 9B are also underlined and highlighted here. SEQ ID NO: 31 encodes only the truncated ActRIIB extracellular domain (corresponding to residues 25-131 in SEQ ID NO: 1) with a L79D substitution, e.g., ActRIIB(L79D 25-131).

FIG. 11 shows the effect of ActRIIB(L79D 25-131)-hFc on hemoglobin concentration in a mouse model of chemotherapy-induced anemia. Data are means $\pm$ SEM. **, P<0.01 vs. paclitaxel at the same time point. This GDF Trap offset the anemia induced by paclitaxel treatment.

FIG. 12 shows the effect of ActRIIB(L79D 25-131)-hFc on red blood cell (RBC) levels in a unilaterally nephrectomized (NEPHX) mouse model of chronic kidney disease. Data are means $\pm$ SEM. ***, P<0.001 vs. baseline. This GDF Trap reversed the nephrectomy-induced anemia observed in control mice.

FIG. 13 shows the effect of ActRIIB(L79D 25-131)-hFc on red blood cell (RBC), hemoglobin (HGB), and hematocrit (HCT) levels in a unilaterally nephrectomized (NEPHX) mouse model of chronic kidney disease. Data are mean changes from baseline over 4 weeks ($\pm$ SEM). *, P<0.05; **, P<0.01; ***, P<0.001 vs. NEPHX controls. This GDF Trap prevented the nephrectomy-associated decline in these erythrocytic parameters, increasing each by a magnitude similar to that in kidney-intact (sham) mice.

FIG. 14 shows the effect of ActRIIB(L79D 25-131)-hFc on red blood cell (RBC) levels in a rat model of anemia induced by acute blood loss. Blood removal occurred on Day -1, with dosing on Days 0 and 3. Data are means $\pm$ SEM. **, P<0.01; ***, P<0.001 vs. vehicle at same time point. This GDF Trap improved the rate and extent of recovery from blood-loss-induced anemia.

FIG. 15 shows the effect of treatment with ActRIIB(L79D 20-134)-hFc (gray) or ActRIIB(L79D 25-131)-hFc (black) on the absolute change in red blood cell concentration from baseline in cynomolgus monkey. VEH=vehicle. Data are means $\pm$ SEM. n=4-8 per group.

FIG. 16 shows the effect of treatment with ActRIIB(L79D 20-134)-hFc (gray) or ActRIIB(L79D 25-131)-hFc (black)

on the absolute change in hematocrit from baseline in cynomolgus monkey. VEH=vehicle. Data are means $\pm$ SEM. n=4-8 per group.

FIG. 17 shows the effect of treatment with ActRIIB(L79D 20-134)-hFc (gray) or ActRIIB(L79D 25-131)-hFc (black) on the absolute change in hemoglobin concentration from baseline in cynomolgus monkey. VEH=vehicle. Data are means $\pm$ SEM. n=4-8 per group.

FIG. 18 shows the effect of treatment with ActRIIB(L79D 20-134)-hFc (gray) or ActRIIB(L79D 25-131)-hFc (black) on the absolute change in circulating reticulocyte concentration from baseline in cynomolgus monkey. VEH=vehicle. Data are means $\pm$ SEM. n=4-8 per group.

DETAILED DESCRIPTION OF THE INVENTION

1. Overview

The transforming growth factor-beta (TGF-beta) superfamily contains a variety of growth factors that share common sequence elements and structural motifs. These proteins are known to exert biological effects on a large variety of cell types in both vertebrates and invertebrates. Members of the superfamily perform important functions during embryonic development in pattern formation and tissue specification and can influence a variety of differentiation processes, including adipogenesis, myogenesis, chondrogenesis, cardiogenesis, hematopoiesis, neurogenesis, and epithelial cell differentiation. The family is divided into two general branches: the BMP/GDF and the TGF-beta/Activin/BMP10 branches, whose members have diverse, often complementary effects. By manipulating the activity of a member of the TGF-beta family, it is often possible to cause significant physiological changes in an organism. For example, the Piedmontese and Belgian Blue cattle breeds carry a loss-of-function mutation in the GDF8 (also called myostatin) gene that causes a marked increase in muscle mass. Grobet et al., *Nat Genet.* 1997, 17(1):71-4. Furthermore, in humans, inactive alleles of GDF8 are associated with increased muscle mass and, reportedly, exceptional strength. Schuelke et al., *N Engl J Med* 2004, 350:2682-8.

TGF- β signals are mediated by heteromeric complexes of type I and type II serine/threonine kinase receptors, which phosphorylate and activate downstream Smad proteins upon ligand stimulation (Massague, 2000, *Nat. Rev. Mol. Cell Biol.* 1:169-178). These type I and type II receptors are transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling. Type II receptors are required for binding ligands and for expression of Type I receptors. Type I and II activin receptors form a stable complex after ligand binding, resulting in phosphorylation of Type I receptors by Type II receptors.

Two related Type II receptors (ActRII), ActRIIA and ActRIIB, have been identified as the Type II receptors for activins (Mathews and Vale, 1991, *Cell* 65:973-982; Attisano et al., 1992, *Cell* 68: 97-108). Besides activins, ActRIIA and ActRIIB can biochemically interact with several other TGF- β family proteins, including BMP7, Nodal, GDF8, and GDF11 (Yamashita et al., 1995, *J. Cell Biol.* 130:217-226; Lee and McPherron, 2001, *Proc. Natl. Acad. Sci.* 98:9306-9311; Yeo and Whitman, 2001, *Mol. Cell* 7: 949-957; Oh et al., 2002, *Genes Dev.* 16:2749-54). ALK4 is the primary type I receptor for activins, particularly for activin A, and ALK-7 may serve as a receptor for activins as

well, particularly for activin B. In certain embodiments, the present invention relates to antagonizing a ligand of ActRIIB receptors (also referred to as an ActRIIB ligand) with a subject GDF Trap polypeptide. Exemplary ligands of ActRIIB receptors include some TGF- β family members, such as activin, Nodal, GDF8, GDF11, and BMP7.

Activins are dimeric polypeptide growth factors that belong to the TGF-beta superfamily. There are three principal activin forms (A, B, and AB) that are homo/heterodimers of two closely related β subunits ($\beta_A\beta_A$, $\beta_B\beta_B$, and $\beta_A\beta_B$, respectively). The human genome also encodes an activin C and an activin E, which are primarily expressed in the liver, and heterodimeric forms containing β_C or β_E are also known. In the TGF-beta superfamily, activins are unique and multifunctional factors that can stimulate hormone production in ovarian and placental cells, support neuronal cell survival, influence cell-cycle progress positively or negatively depending on cell type, and induce mesodermal differentiation at least in amphibian embryos (DePaolo et al., 1991, *Proc Soc Exp Biol Med.* 198:500-512; Dyson et al., 1997, *Curr Biol.* 7:81-84; Woodruff, 1998, *Biochem Pharmacol.* 55:953-963). Moreover, erythroid differentiation factor (EDF) isolated from the stimulated human monocytic leukemic cells was found to be identical to activin A (Murata et al., 1988, *PNAS*, 85:2434). It has been suggested that activin A promotes erythropoiesis in the bone marrow. In several tissues, activin signaling is antagonized by its related heterodimer, inhibin. For example, during the release of follicle-stimulating hormone (FSH) from the pituitary, activin promotes FSH secretion and synthesis, while inhibin prevents FSH secretion and synthesis. Other proteins that may regulate activin bioactivity and/or bind to activin include follistatin (FS), follistatin-related protein (FSRP) and α_2 -macroglobulin.

Nodal proteins have functions in mesoderm and endoderm induction and formation, as well as subsequent organization of axial structures such as heart and stomach in early embryogenesis. It has been demonstrated that dorsal tissue in a developing vertebrate embryo contributes predominantly to the axial structures of the notochord and pre-chordal plate while it recruits surrounding cells to form non-axial embryonic structures. Nodal appears to signal through both type I and type II receptors and intracellular effectors known as Smad proteins. Recent studies support the idea that ActRIIA and ActRIIB serve as type II receptors for Nodal (Sakuma et al., *Genes Cells.* 2002, 7:401-12). It is suggested that Nodal ligands interact with their co-factors (e.g., *cripto*) to activate activin type I and type II receptors, which phosphorylate Smad2. Nodal proteins are implicated in many events critical to the early vertebrate embryo, including mesoderm formation, anterior patterning, and left-right axis specification. Experimental evidence has demonstrated that Nodal signaling activates pAR3-Lux, a luciferase reporter previously shown to respond specifically to activin and TGF-beta. However, Nodal is unable to induce pTlx2-Lux, a reporter specifically responsive to bone morphogenetic proteins. Recent results provide direct biochemical evidence that Nodal signaling is mediated by both activin-TGF-beta pathway Smads, Smad2 and Smad3. Further evidence has shown that the extracellular *cripto* protein is required for Nodal signaling, making it distinct from activin or TGF-beta signaling.

Growth and Differentiation Factor-8 (GDF8) is also known as myostatin. GDF8 is a negative regulator of skeletal muscle mass. GDF8 is highly expressed in the developing and adult skeletal muscle. The GDF8 null mutation in transgenic mice is characterized by a marked hypertrophy

and hyperplasia of the skeletal muscle (McPherron et al., *Nature*, 1997, 387:83-90). Similar increases in skeletal muscle mass are evident in naturally occurring mutations of GDF8 in cattle (Ashmore et al., 1974, *Growth*, 38:501-507; Swatland and Kieffer, *J. Anim. Sci.*, 1994, 38:752-757; McPherron and Lee, *Proc. Natl. Acad. Sci. USA*, 1997, 94:12457-12461; and Kambadur et al., *Genome Res.*, 1997, 7:910-915) and, strikingly, in humans (Schuelke et al., *N Engl J Med* 2004; 350:2682-8). Studies have also shown that muscle wasting associated with HIV-infection in humans is accompanied by increases in GDF8 protein expression (Gonzalez-Cadavid et al., *PNAS*, 1998, 95:14938-43). In addition, GDF8 can modulate the production of muscle-specific enzymes (e.g., creatine kinase) and modulate myoblast cell proliferation (WO 00/43781). The GDF8 propeptide can noncovalently bind to the mature GDF8 domain dimer, inactivating its biological activity (Miyazono et al. (1988) *J. Biol. Chem.*, 263: 6407-6415; Wakefield et al. (1988) *J. Biol. Chem.*, 263: 7646-7654; and Brown et al. (1990) *Growth Factors*, 3: 35-43). Other proteins which bind to GDF8 or structurally related proteins and inhibit their biological activity include follistatin, and potentially, follistatin-related proteins (Gamer et al. (1999) *Dev. Biol.*, 208: 222-232).

Growth and Differentiation Factor-11 (GDF11), also known as BMP11, is a secreted protein (McPherron et al., 1999, *Nat. Genet.* 22: 260-264). GDF11 is expressed in the tail bud, limb bud, maxillary and mandibular arches, and dorsal root ganglia during mouse development (Nakashima et al., 1999, *Mech. Dev.* 80: 185-189). GDF11 plays a unique role in patterning both mesodermal and neural tissues (Gamer et al., 1999, *Dev Biol.*, 208:222-32). GDF11 was shown to be a negative regulator of chondrogenesis and myogenesis in developing chick limb (Gamer et al., 2001, *Dev Biol.* 229:407-20). The expression of GDF11 in muscle also suggests its role in regulating muscle growth in a similar way to GDF8. In addition, the expression of GDF11 in brain suggests that GDF11 may also possess activities that relate to the function of the nervous system. Interestingly, GDF11 was found to inhibit neurogenesis in the olfactory epithelium (Wu et al., 2003, *Neuron.* 37:197-207). Hence, GDF11 may have in vitro and in vivo applications in the treatment of diseases such as muscle diseases and neurodegenerative diseases (e.g., amyotrophic lateral sclerosis).

Bone morphogenetic protein (BMP7), also called osteogenic protein-1 (OP-1), is well known to induce cartilage and bone formation. In addition, BMP7 regulates a wide array of physiological processes. For example, BMP7 may be the osteoinductive factor responsible for the phenomenon of epithelial osteogenesis. It is also found that BMP7 plays a role in calcium regulation and bone homeostasis. Like activin, BMP7 binds to Type II receptors, ActRIIA and ActRIIB. However, BMP7 and activin recruit distinct Type I receptors into heteromeric receptor complexes. The major BMP7 Type I receptor observed was ALK2, while activin bound exclusively to ALK4 (ActRIIB). BMP7 and activin elicited distinct biological responses and activated different Smad pathways (Macias-Silva et al., 1998, *J Biol Chem.* 273:25628-36).

As demonstrated herein, a GDF Trap polypeptide, which is a variant ActRIIB polypeptide (ActRIIB), is more effective at increasing red blood cell levels in vivo as compared to a wild-type soluble ActRIIB polypeptide and has beneficial effects in a variety of models for anemias. It should be noted that hematopoiesis is a complex process, regulated by a variety of factors, including erythropoietin, G-CSF and iron homeostasis. The terms "increase red blood cell levels"

and “promote red blood cell formation” refer to clinically observable metrics, such as hematocrit, red blood cell counts and hemoglobin measurements, and are intended to be neutral as to the mechanism by which such changes occur.

In addition to stimulating red blood cell levels, GDF Trap polypeptides are useful for a variety of therapeutic applications, including, for example, promoting muscle growth (see PCT Publication Nos. WO 2006/012627 and WO 2008/097541, which are hereby incorporated by reference in their entirety). In certain instances, when administering a GDF Trap polypeptide for the purpose of increasing muscle, it may be desirable to reduce or minimize effects on red blood cells. By monitoring various hematologic parameters in patients being treated with, or who are candidates for treatment with, a GDF Trap polypeptide, appropriate dosing (including amounts and frequency of administration) may be determined based on an individual patient’s needs, baseline hematologic parameters, and purpose for treatment. Furthermore, therapeutic progress and effects on one or more hematologic parameters over time may be useful in managing patients being dosed with a GDF Trap polypeptide by facilitating patient care, determining appropriate maintenance dosing (both amounts and frequency), etc.

The terms used in this specification generally have their ordinary meanings in the art, within the context of this invention and in the specific context where each term is used. Certain terms are discussed below or elsewhere in the specification, to provide additional guidance to the practitioner in describing the compositions and methods of the invention and how to make and use them. The scope or meaning of any use of a term will be apparent from the specific context in which the term is used.

“About” and “approximately” shall generally mean an acceptable degree of error for the quantity measured given the nature or precision of the measurements. Typically, exemplary degrees of error are within 20 percent (%), preferably within 10%, and more preferably within 5% of a given value or range of values.

Alternatively, and particularly in biological systems, the terms “about” and “approximately” may mean values that are within an order of magnitude, preferably within 5-fold and more preferably within 2-fold of a given value. Numerical quantities given herein are approximate unless stated otherwise, meaning that the term “about” or “approximately” can be inferred when not expressly stated.

The methods of the invention may include steps of comparing sequences to each other, including wild-type sequence to one or more mutants (sequence variants). Such comparisons typically comprise alignments of polymer sequences, e.g., using sequence alignment programs and/or algorithms that are well known in the art (for example, BLAST, FASTA and MEGALIGN, to name a few). The skilled artisan can readily appreciate that, in such alignments, where a mutation contains a residue insertion or deletion, the sequence alignment will introduce a “gap” (typically represented by a dash, or “A”) in the polymer sequence not containing the inserted or deleted residue.

“Homologous,” in all its grammatical forms and spelling variations, refers to the relationship between two proteins that possess a “common evolutionary origin,” including proteins from superfamilies in the same species of organism, as well as homologous proteins from different species of organism. Such proteins (and their encoding nucleic acids) have sequence homology, as reflected by their sequence similarity, whether in terms of percent identity or by the presence of specific residues or motifs and conserved positions.

The term “sequence similarity,” in all its grammatical forms, refers to the degree of identity or correspondence between nucleic acid or amino acid sequences that may or may not share a common evolutionary origin.

However, in common usage and in the instant application, the term “homologous,” when modified with an adverb such as “highly,” may refer to sequence similarity and may or may not relate to a common evolutionary origin.

2. GDF Trap Polypeptides

In certain aspects, the invention relates to GDF Trap polypeptides, e.g., soluble variant ActRIIB polypeptides, including, for example, fragments, functional variants, and modified forms of ActRIIB polypeptides. In certain embodiments, the GDF Trap polypeptides have at least one similar or same biological activity as a corresponding wild-type ActRIIB polypeptide. For example, a GDF Trap polypeptide of the invention may bind to and inhibit the function of an ActRIIB ligand (e.g., activin A, activin AB, activin B, Nodal, GDF8, GDF11 or BMP7). Optionally, a GDF Trap polypeptide increases red blood cell levels. Examples of GDF Trap polypeptides include human ActRIIB precursor polypeptides (SEQ ID NO: 1 or 39) having one or more sequence variations, and soluble human ActRIIB polypeptides (e.g., SEQ ID NOs: 2, 3, 7, 11, 26, 28, 29, 32, 37, 38, 40 and 41) having one or more sequence variations. A GDF Trap refers to an ActRIIB polypeptide having a decreased affinity for activin relative to other ActRIIB ligands, including for example GDF11 and/or myostatin.

As used herein, the term “ActRIIB” refers to a family of activin receptor type IIb (ActRIIB) proteins from any species and variants derived from such ActRIIB proteins by mutagenesis or other modification. Reference to ActRIIB herein is understood to be a reference to any one of the currently identified forms. Members of the ActRIIB family are generally transmembrane proteins, composed of a ligand-binding extracellular domain with a cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine kinase activity. Amino acid sequences of human ActRIIA soluble extracellular domain (provided for comparison) and ActRIIB soluble extracellular domain are illustrated in FIG. 1.

The term “ActRIIB polypeptide” includes polypeptides comprising any naturally occurring polypeptide of an ActRIIB family member as well as any variants thereof (including mutants, fragments, fusions, and peptidomimetic forms) that retain a useful activity. See, for example, WO 2006/012627. For example, ActRIIB polypeptides include polypeptides derived from the sequence of any known ActRIIB having a sequence at least about 80% identical to the sequence of an ActRIIB polypeptide, and optionally at least 85%, 90%, 95%, 97%, 99% or greater identity. For example, an ActRIIB polypeptide may bind to and inhibit the function of an ActRIIB protein and/or activin. An ActRIIB polypeptide which is a GDF Trap may be selected for activity in promoting red blood cell formation in vivo. Examples of ActRIIB polypeptides include human ActRIIB precursor polypeptide (SEQ ID NO: 1 and 39) and soluble human ActRIIB polypeptides (e.g., SEQ ID NO: 2, 3, 7, 11, 26, 28, 29, 32, 37, 38, 40 and 41). Numbering of amino acids for all ActRIIB-related polypeptides described herein is based on the numbering for SEQ ID NO:1, unless specifically designated otherwise.

15

The human ActRIIB precursor protein sequence is as follows:

(SEQ ID NO: 1)

MTAPWVALALLWGSLWPGSGRGEAETRECIYYNANWELERTNSGLER

CEGEQDKRLHCYASWRNS**SGTIELVKKGCWLDDFNCYDRQECVATEEN**

PQVYFCCCEGNFCNERFTHLPEAGGPEVTYEPPTAPTLLTVLAYSLL

PIGGLSLIVLLAFWMYRHRKPPYGHVDIHEDPGPPPPSPVLGLKPLQL

LEIKARGRFGCVWKAQLMNDFAVKIFPLQDKQSWQSEREIFSTPGMK

HENLLQFIAAEDRGSNLEVELWLITAFHDKGSLTDYLGKNIITWNELC

HVAETMSRGLSYLHEDVPWCRGEGHKPSIAHRDFKSKNVLLKSDLTAV

LADFLAVRFEPGKPPGDTHGQVGTTRYMAPEVLEGAINFQRDAFLRI

DMYAMGLVLWELVSRCKAADGPVDEYMLPFEEIQQHPSLEELQEVVV

HKKMRPTIKDHWLKHPLGLAQLCVTIEECWDHDAEARLSAGCVEERVSL

IRRSVNGTSDCLVSLVTSVTNVDLPPKESSI

The signal peptide is single underlined; the extracellular domain is in bold and the potential N-linked glycosylation sites are in boxes.

A form with an alanine at position 64 is also reported in the literature, as follows:

(SEQ ID NO: 39)

MTAPWVALALLWGSLWPGSGRGEAETRECIYYNANWELERTNSGLER

CEGEQDKRLHCYASWANS**SGTIELVKKGCWLDDFNCYDRQECVATEEN**

PQVYFCCCEGNFCNERFTHLPEAGGPEVTYEPPTAPTLLTVLAYSLLPI

GGLSLIVLLAFWMYRHRKPPYGHVDIHEDPGPPPPSPVLGLKPLQLLEIK

ARGRFGCVWKAQLMNDFAVKIFPLQDKQSWQSEREIFSTPGMKHENLLQ

FIAAEKRGSNLEVELWLITAFHDKGSLTDYLGKNIITWNELCVHAETMSR

GLSYLHEDVPWCRGEGHKPSIAHRDFKSKNVLLKSDLTAVLADFLAVRF

EPGKPPGDTHGQVGTTRYMAPEVLEGAINFQRDAFLRIDMYAMGLVLWEL

VSRCKAADGPVDEYMLPEEIIQQHPSLEELQEVVVHKKMRPTIKDHWLKH

PGLAQLCVTIEECWDHDAEARLSAGCVEERVSLIRRSVNGTSDCLVSLV

TSVTNVDLPPKESSI

The human ActRIIB soluble (extracellular), processed polypeptide sequence is as follows:

(SEQ ID NO: 2)

GRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWRNSSG

TIELVKKGCWLDDFNCYDRQECVATEENPQVYFCCCEGNFCNERFTHLP

EAGGPEVTYEPPTAPT

The alternative form with an A64 is as follows:

(SEQ ID NO: 40)

GRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWRNSSG

TIELVKKGCWLDDFNCYDRQECVATEENPQVYFCCCEGNFCNERFTHLP

EAGGPEVTYEPPTAPT

16

In some conditions, the protein may be produced with an "SGR . . ." sequence at the N-terminus. The C-terminal "tail" of the extracellular domain is underlined. The sequence with the "tail" deleted (a Δ15 sequence) is as follows:

(SEQ ID NO: 3)

GRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWRNSSG

TIELVKKGCWLDDFNCYDRQECVATEENPQVYFCCCEGNFCNERFTHLP

EA

The alternative form with an A64 is as follows:

(SEQ ID NO: 41)

GRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWRNSSG

TIELVKKGCWLDDFNCYDRQECVATEENPQVYFCCCEGNFCNERFTHLP

EA

In some conditions, the protein may be produced with an "SGR . . ." sequence at the N-terminus. The nucleic acid sequence encoding a human ActRIIB precursor protein is as follows: (nucleotides 5-1543 of Genbank entry NM_001106)(the sequence as shown provides an alanine at position 64, and may be modified to provide an arginine instead)

(SEQ ID NO: 4)

ATGACGGCGCCTGGGTGGCCCTCGCCCTCCTCTGGGGATCGCTGTGGC

CCGGCTCTGGGCGTGGGAGGCTGAGACACGGGAGTGCATCTACTACAA

CGCCAACTGGGAGCTGGAGCGCACCAACAGAGCGCCTGGAGCGCTGC

GAAGCGAGCAGGACAAGCGCTGCACTGCTACGCCTCTGGGCCAACA

GCTCTGGCACCATCGAGCTCGTGAAGAAGGCTGCTGGCTAGATGACTT

CAACTGCTACGATAGGCAGGAGTGTGTGGCCACTGAGGAGAACCCCCAG

GTGTACTTCTGCTGTGTGAAGGCAACTTCTGCAACGAGCGCTTCACTC

ATTTGCCAGAGGCTGGGGGCCCGAAGTACGTACGAGCCACCCCCGAC

AGCCCCCACCCTGCTCAGGTGCTGGCCTACTACTGCTGCCCATCGGG

GGCCTTTCCCTCATCGTCTGCTGGCCTTTTGATGTACCGGCATCGCA

AGCCCCCTACGGTCATGTGGACATCCATGAGGACCTTGGGCCCTCCACC

ACCATCCCCCTCTGGTGGGCTGAAGCCACTGCAGCTGCTGGAGATCAAG

GCTCGGGGCGCTTTGGCTGTGTCTGGAAGGCCAGCTCATGAATGACT

TTGTAGCTGTCAAGATCTTCCCACTCCAGGACAAGCAGTCGTGGCAGAG

TGAACGGGAGATCTTCAACACCTGGCATGAAGCACGAGAACCTGCTA

CAGTTCATTGCTGCCGAGAAGCGAGGCTCCAACCTCGAAGTAGAGCTGT

GGCTCATCACGGCCTTCCATGACAAGGGCTCCCTCACGGATTACCTCAA

GGGGAACATCATCATGGAACGAACTGTGTCTAGTAGAGAGACGATG

TCACAGAGGCTCTCATACCTGCATGAGGATGTGCCTGGTCCCTGGCG

AGGGCCACAAGCCGTCTATTGCCACAGGGACTTTAAAGTAAGAATGT

ATTGCTGAAGAGCGACCTCACAGCCGTGCTGGCTGACTTTGGCTTGGCT

GTTTCGATTGAGCCAGGGAACCTCCAGGGGACACCCACGGACAGGTAG

17

-continued

GCACGAGACGGTACATGGCTCTGAGGTGCTCGAGGGAGCCATCAACTT
 CCAGAGAGATGCCCTTCCTGCGCATTGACATGTATGCCATGGGGTTGGTG
 CTGTGGGAGCTTGTGTCTCGCTGCAAGGCTGCAGACGGACCCGTGGATG
 AGTACATGCTGCCCTTTGAGGAAGAGATTGGCCAGCACCCCTTCGTGGGA
 GGAGCTGCAGGAGGTGGTGGTGACACAAGAAGATGAGGCCACCATTAA
 GATCACTGGTTGAAACACCCGGGCTGCGCCAGCTTTGTGTGACCATCG
 AGGAGTGCTGGGACCATGATGCAGAGGCTCGCTTGTCCGCGGGCTGTGT
 GGAGGAGCGGGTGTCCCTGATTGCGAGGTGCGTCAACGGCACTACCTCG
 GACTGTCTCGTTTCCCTGGTGACCTCTGTCAACCAATGTGGACCTGCCCC
 CTAAGAGTCAAGCATCTAA

The nucleic acid sequence encoding a human ActRIIA soluble (extracellular) polypeptide is as follows (the sequence as shown provides an alanine at position 64, and may be modified to provide an arginine instead):

(SEQ ID NO: 5)
 GGGCGTGGGGAGGCTGAGACACGGAGTGCATCTACTACAACGCCAACT
 GGGAGCTGGAGCGCACCAACAGAGCGGCTGGAGCGCTGCGAAGGCGA
 GCAGGACAAGCGGCTGCACTGCTACGCCTCTGGGCCAACAGCTCTGGC
 ACCATCGAGCTCGTGAAGAAGGGCTGCTGGCTAGATGACTTCAACTGCT
 ACGATAGGCAGGAGTGTGTGGCCACTGAGGAGAACCCCCAGGTGTACTT
 CTGCTGCTGTGAAGCAACTTCTGCAACGAGCGCTTCACTCATTTGCCA
 GAGGCTGGGGGCCCGGAAGTCACGTACGAGCCACCCCGACAGCCCCCA
 CC

In a specific embodiment, the invention relates to GDF Trap polypeptides which are variant forms of soluble ActRIIB polypeptides. As described herein, the term “soluble ActRIIB polypeptide” generally refers to polypeptides comprising an extracellular domain of an ActRIIB protein. The term “soluble ActRIIB polypeptide,” as used herein, includes any naturally occurring extracellular domain of an ActRIIB protein as well as any variants thereof (including mutants, fragments and peptidomimetic forms) that retain a useful activity. For example, the extracellular domain of an ActRIIB protein binds to a ligand and is generally soluble. Examples of soluble ActRIIB polypeptides include ActRIIB soluble polypeptides (e.g., SEQ ID NOs: 22, 3, 7, 11, 26, 28, 29, 32, 37, 38, 40 and 41). Other examples of soluble ActRIIB polypeptides comprise a signal sequence in addition to the extracellular domain of an ActRIIB protein, see Example 1. The signal sequence can be a native signal sequence of an ActRIIB, or a signal sequence from another protein, such as a tissue plasminogen activator (TPA) signal sequence or a honey bee melittin (HBM) signal sequence.

The disclosure identifies functionally active portions and variants of ActRIIB. Applicants have ascertained that an Fc fusion protein having the sequence disclosed by Hilden et al. (Blood. 1994 Apr. 15; 83(8):2163-70), which has an Alanine at the position corresponding to amino acid 64 of SEQ ID NO: 1 (A64), has a relatively low affinity for activin and GDF-11. By contrast, the same Fc fusion protein with an Arginine at position 64 (R64) has an affinity for activin and GDF-11 in the low nanomolar to high picomolar range.

18

Therefore, a sequence with an R64 is used as the wild-type reference sequence for human ActRIIB in this disclosure.

Attisano et al. (Cell. 1992 Jan. 10; 68(1):97-108) showed that a deletion of the proline knot at the C-terminus of the extracellular domain of ActRIIB reduced the affinity of the receptor for activin. An ActRIIB-Fc fusion protein containing amino acids 20-119 of SEQ ID NO: 1, “ActRIIB(20-119)-Fc”, has reduced binding to GDF-11 and activin relative to an ActRIIB(20-134)-Fc, which includes the proline knot region and the complete juxtamembrane domain. However, an ActRIIB(20-129)-Fc protein retains similar but somewhat reduced activity relative to the wild type, even though the proline knot region is disrupted. Thus, ActRIIB extracellular domains that stop at amino acid 134, 133, 132, 131, 130 and 129 are all expected to be active, but constructs stopping at 134 or 133 may be most active. Similarly, mutations at any of residues 129-134 are not expected to alter ligand binding affinity by large margins. In support of this, mutations of P129 and P130 do not substantially decrease ligand binding. Therefore, a GDF Trap polypeptide which is an ActRIIB-Fc fusion protein may end as early as amino acid 109 (the final cysteine), however, forms ending at or between 109 and 119 are expected to have reduced ligand binding. Amino acid 119 is poorly conserved and so is readily altered or truncated. Forms ending at 128 or later retain ligand binding activity. Forms ending at or between 119 and 127 will have an intermediate binding ability. Any of these forms may be desirable to use, depending on the clinical or experimental setting.

At the N-terminus of ActRIIB, it is expected that a protein beginning at amino acid 29 or before will retain ligand binding activity. Amino acid 29 represents the initial cysteine. An alanine to asparagine mutation at position 24 introduces an N-linked glycosylation sequence without substantially affecting ligand binding. This confirms that mutations in the region between the signal cleavage peptide and the cysteine cross-linked region, corresponding to amino acids 20-29 are well tolerated. In particular, constructs beginning at position 20, 21, 22, 23 and 24 will retain activity, and constructs beginning at positions 25, 26, 27, 28 and 29 are also expected to retain activity. Data shown in the Examples demonstrates that, surprisingly, a construct beginning at 22, 23, 24 or 25 will have the most activity.

Taken together, an active portion of ActRIIB comprises amino acids 29-109 of SEQ ID NO: 1, and GDF Trap constructs may, for example, comprise a portion of ActRIIB beginning at a residue corresponding to amino acids 20-29 of SEQ ID NO: 1 or 39 and ending at a position corresponding to amino acids 109-134 of SEQ ID NO: 1 or 39. Other examples include constructs that begin at a position from 20-29 or 21-29 and end at a position from 119-134, 119-133, 129-134, or 129-133 of SEQ ID NO: 1 or 39. Other examples include constructs that begin at a position from 20-24 (or 21-24, or 22-25) and end at a position from 109-134 (or 109-133), 119-134 (or 119-133) or 129-134 (or 129-133) of SEQ ID NO: 1 or 39. Variants within these ranges are also contemplated, particularly those having at least 80%, 85%, 90%, 95% or 99% identity to the corresponding portion of SEQ ID NO: 1 or 39. In certain embodiments, the GDF Trap polypeptide comprises, consists essentially of, or consists of, a polypeptide having an amino acid sequence that is at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identical to amino acid residues 25-131 of SEQ ID NO: 1 or 39. In certain embodiments, the GDF Trap polypeptide comprises, consists essentially of, or consists of, a polypeptide having an amino acid sequence that is at least 80%, 85%, 90%, 95%, 96%, 97%,

98%, 99% or 100% identical to SEQ ID NOs: 7, 26, 28, 29, 32, 37 or 38. In preferred embodiments, the GDF Trap polypeptide consists of, or consists essentially of, the amino acid sequence of SEQ ID NO: 7, 26, 28, 29, 32, 37 or 38.

The disclosure includes the results of an analysis of composite ActRIIB structures, shown in FIG. 1, demonstrating that the ligand binding pocket is defined by residues Y31, N33, N35, L38 through T41, E47, E50, Q53 through K55, L57, H58, Y60, S62, K74, W78 through N83, Y85, R87, A92, and E94 through F101. At these positions, it is expected that conservative mutations will be tolerated, although a K74A mutation is well-tolerated, as are R40A, K55A, F82A and mutations at position L79. R40 is a K in *Xenopus*, indicating that basic amino acids at this position will be tolerated. Q53 is R in bovine ActRIIB and K in *Xenopus* ActRIIB, and therefore amino acids including R, K, Q, N and H will be tolerated at this position. Thus, a general formula for a GDF Trap protein is one that comprises amino acids 29-109 of SEQ ID NO: 1 or 39, but optionally beginning at a position ranging from 20-24 or 22-25 and ending at a position ranging from 129-134, and comprising no more than 1, 2, 5, 10 or 15 conservative amino acid changes in the ligand binding pocket, and zero, one or more non-conservative alterations at positions 40, 53, 55, 74, 79 and/or 82 in the ligand binding pocket. Such a protein may retain greater than 80%, 90%, 95% or 99% sequence identity to the sequence of amino acids 29-109 of SEQ ID NO: 1 or 39. Sites outside the binding pocket, at which variability may be particularly well tolerated, include the amino and carboxy termini of the extracellular domain (as noted above), and positions 42-46 and 65-73. An asparagine to alanine alteration at position 65 (N65A) actually improves ligand binding in the A64 background, and is thus expected to have no detrimental effect on ligand binding in the R64 background. This change probably eliminates glycosylation at N65 in the A64 background, thus demonstrating that a significant change in this region is likely to be tolerated. While an R64A change is poorly tolerated, R64K is well-tolerated, and thus another basic residue, such as H may be tolerated at position 64.

ActRIIB is well-conserved across nearly all vertebrates, with large stretches of the extracellular domain conserved completely. Many of the ligands that bind to ActRIIB are also highly conserved. Accordingly, comparisons of ActRIIB sequences from various vertebrate organisms provide insights into residues that may be altered. Therefore, an active, human ActRIIB variant polypeptide useful as a GDF Trap may include one or more amino acids at corresponding positions from the sequence of another vertebrate ActRIIB, or may include a residue that is similar to that in the human or other vertebrate sequence. The following examples illustrate this approach to defining an active ActRIIB variant. L46 is a valine in *Xenopus* ActRIIB, and so this position may be altered, and optionally may be altered to another hydrophobic residue, such as V, I or F, or a non-polar residue such as A. E52 is a K in *Xenopus*, indicating that this site may be tolerant of a wide variety of changes, including polar residues, such as E, D, K, R, H, S, T, P, G, Y and probably A. T93 is a K in *Xenopus*, indicating that a wide structural variation is tolerated at this position, with polar residues favored, such as S, K, R, E, D, H, G, P, G and Y. F108 is a Y in *Xenopus*, and therefore Y or other hydrophobic group, such as I, V or L should be tolerated. E111 is K in *Xenopus*, indicating that charged residues will be tolerated at this position, including D, R, K and H, as well as Q and N. R112 is K in *Xenopus*, indicating that basic residues are tolerated at this position, including R and H. A at position 119 is

relatively poorly conserved, and appears as P in rodents and V in *Xenopus*, thus essentially any amino acid should be tolerated at this position.

The disclosure demonstrates that the addition of a further N-linked glycosylation site (N-X-S/T) increases the serum half-life of an ActRIIB-Fc fusion protein, relative to the ActRIIB(R64)-Fc form. By introducing an asparagine at position 24 (A24N construct), an NXT sequence is created that confers a longer half-life. Other NX(T/S) sequences are found at 42-44 (NQS) and 65-67 (NSS), although the latter may not be efficiently glycosylated with the R at position 64. N-X-S/T sequences may be generally introduced at positions outside the ligand binding pocket defined in FIG. 1. Particularly suitable sites for the introduction of non-endogenous N-X-S/T sequences include amino acids 20-29, 20-24, 22-25, 109-134, 120-134 or 129-134. N-X-S/T sequences may also be introduced into the linker between the ActRIIB sequence and the Fc or other fusion component. Such a site may be introduced with minimal effort by introducing an N in the correct position with respect to a pre-existing S or T, or by introducing an S or T at a position corresponding to a pre-existing N. Thus, desirable alterations that would create an N-linked glycosylation site are: A24N, R64N, S67N (possibly combined with an N65A alteration), E106N, R112N, G120N, E123N, P129N, A132N, R112S and R112T. Any S that is predicted to be glycosylated may be altered to a T without creating an immunogenic site, because of the protection afforded by the glycosylation. Likewise, any T that is predicted to be glycosylated may be altered to an S. Thus the alterations S67T and S44T are contemplated. Likewise, in an A24N variant, an S26T alteration may be used. Accordingly, a GDF Trap may be an ActRIIB variant having one or more additional, non-endogenous N-linked glycosylation consensus sequences.

Position L79 of ActRIIB may be altered to confer altered activin—myostatin (GDF-11) binding properties. L79A or L79P reduces GDF-11 binding to a greater extent than activin binding. L79E or L79D retains GDF-11 binding. Remarkably, the L79E and L79D variants have greatly reduced activin binding. In vivo experiments indicate that these non-activin receptors retain significant ability to increase red blood cells but show decreased effects on other tissues. These data demonstrate the desirability and feasibility for obtaining polypeptides with reduced effects on activin. In exemplary embodiments, the methods described herein utilize a GDF Trap polypeptide which is a variant ActRIIB polypeptide comprising an acidic amino acid (e.g., D or E) at the position corresponding to position 79 of SEQ ID NO: 1 or 39, optionally in combination with one or more additional amino acid substitutions, additions, or deletions.

The variations described may be combined in various ways. Additionally, the results of the mutagenesis program described herein indicate that there are amino acid positions in ActRIIB that are often beneficial to conserve. These include position 64 (basic amino acid), position 80 (acidic or hydrophobic amino acid), position 78 (hydrophobic, and particularly tryptophan), position 37 (acidic, and particularly aspartic or glutamic acid), position 56 (basic amino acid), position 60 (hydrophobic amino acid, particularly phenyl-alanine or tyrosine). Thus, in each of the variants disclosed herein, the disclosure provides a framework of amino acids that may be conserved. Other positions that may be desirable to conserve are as follows: position 52 (acidic amino acid), position 55 (basic amino acid), position 81 (acidic), 98 (polar or charged, particularly E, D, R or K).

In certain embodiments, isolated fragments of ActRIIB polypeptides can be obtained by screening polypeptides

recombinantly produced from the corresponding fragment of the nucleic acid encoding an ActRIIB polypeptide (e.g., SEQ ID NOs: 4 and 5). In addition, fragments can be chemically synthesized using techniques known in the art such as conventional Merrifield solid phase f-Moc or t-Boc chemistry. The fragments can be produced (recombinantly or by chemical synthesis) and tested to identify those peptidyl fragments that can function, for example, as antagonists (inhibitors) or agonists (activators) of an ActRIIB protein or an ActRIIB ligand.

In certain embodiments, GDF Trap polypeptide is a variant ActRIIB polypeptide having an amino acid sequence that is at least 75% identical to an amino acid sequence selected from SEQ ID NOs: 2, 3, 7, 11, 26, 28, 29, 32, 37, 38, 40 or 41. In certain cases, the GDF Trap has an amino acid sequence at least 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100% identical to an amino acid sequence selected from SEQ ID NOs: 2, 3, 7, 11, 26, 28, 29, 32, 37, 38, 40 or 41. In certain embodiments, the GDF Trap comprises, consists essentially of, or consists of, an amino acid sequence at least 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100% identical to an amino acid sequence selected from SEQ ID NOs: 2, 3, 7, 11, 26, 28, 29, 32, 37, 38, 40 or 41, wherein the position corresponding to L79 of SEQ ID NO: 1 is an acidic amino acid (e.g., a D or E amino acid residue).

In certain embodiments, the present invention contemplates making functional variants by modifying the structure of a GDF Trap polypeptide for such purposes as enhancing therapeutic efficacy, or stability (e.g., ex vivo shelf life and resistance to proteolytic degradation in vivo). GDF Trap polypeptides can also be produced by amino acid substitution, deletion, or addition. For instance, it is reasonable to expect that an isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid (e.g., conservative mutations) will not have a major effect on the biological activity of the resulting molecule. Conservative replacements are those that take place within a family of amino acids that are related in their side chains. Whether a change in the amino acid sequence of a GDF Trap polypeptide results in a functional variant can be readily determined by assessing the ability of the GDF Trap polypeptide to produce a response in cells relative to the unmodified GDF Trap polypeptide or a wild-type ActRIIB polypeptide, or to bind to one or more ligands, such as activin, GDF-11 or myostatin as compared to the unmodified GDF Trap polypeptide or a wild-type ActRIIB polypeptide.

In certain specific embodiments, the present invention contemplates making mutations in the extracellular domain (also referred to as ligand-binding domain) of an ActRIIB polypeptide such that the ActRIIB polypeptide has altered ligand-binding activities (e.g., binding affinity or binding specificity). In certain cases, such GDF Trap polypeptides have altered (elevated or reduced) binding affinity for a specific ligand. In other cases, the GDF Trap polypeptides have altered binding specificity for ActRIIB ligands.

For example, the disclosure provides GDF Trap polypeptides that preferentially bind to GDF8/GDF11 relative to activins. The disclosure further establishes the desirability of such polypeptides for reducing off-target effects, although such selective variants may be less desirable for the treatment of severe diseases where very large gains in red blood cell levels may be needed for therapeutic effect and where some level of off-target effect is acceptable. For example, amino acid residues of the ActRIIB protein, such as E39, K55, Y60, K74, W78, D80, and F101, are in the ligand-

binding pocket and mediate binding to its ligands such as activin and GDF8. Thus, the present invention provides a GDF Trap comprising an altered ligand-binding domain (e.g., GDF8-binding domain) of an ActRIIB receptor, which comprises one or more mutations at those amino acid residues. Optionally, the altered ligand-binding domain can have increased selectivity for a ligand such as GDF8 relative to a wild-type ligand-binding domain of an ActRIIB receptor. To illustrate, these mutations increase the selectivity of the altered ligand-binding domain for GDF8 over activin. Optionally, the altered ligand-binding domain has a ratio of K_d for activin binding to K_d for GDF8 binding that is at least 2, 5, 10, or even 100 fold greater relative to the ratio for the wild-type ligand-binding domain. Optionally, the altered ligand-binding domain has a ratio of IC_{50} for inhibiting activin to IC_{50} for inhibiting GDF8 that is at least 2, 5, 10, or even 100 fold greater relative to the wild-type ligand-binding domain. Optionally, the altered ligand-binding domain inhibits GDF8 with an IC_{50} at least 2, 5, 10, or even 100 times less than the IC_{50} for inhibiting activin.

As a specific example, the positively-charged amino acid residue Asp (D80) of the ligand-binding domain of ActRIIB can be mutated to a different amino acid residue to produce a GDF Trap polypeptide that preferentially binds to GDF8, but not activin. Preferably, the D80 residue is changed to an amino acid residue selected from the group consisting of: an uncharged amino acid residue, a negative amino acid residue, and a hydrophobic amino acid residue. As a further specific example, the hydrophobic residue, L79, can be altered to the acidic amino acids aspartic acid or glutamic acid to greatly reduce activin binding while retaining GDF11 binding. As will be recognized by one of skill in the art, most of the described mutations, variants or modifications may be made at the nucleic acid level or, in some cases, by post translational modification or chemical synthesis. Such techniques are well known in the art.

In certain embodiments, the present invention contemplates GDF Trap polypeptides having specific mutations in ActRIIB so as to alter the glycosylation of the ActRIIB polypeptide. Exemplary glycosylation sites in GDF Trap polypeptides are illustrated in FIG. 1 (e.g., the underlined NX(S/T) sites). Such mutations may be selected so as to introduce or eliminate one or more glycosylation sites, such as O-linked or N-linked glycosylation sites. Asparagine-linked glycosylation recognition sites generally comprise a tripeptide sequence, asparagine-X-threonine (where "X" is any amino acid) which is specifically recognized by appropriate cellular glycosylation enzymes. The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the wild-type ActRIIB polypeptide (for O-linked glycosylation sites). A variety of amino acid substitutions or deletions at one or both of the first or third amino acid positions of a glycosylation recognition site (and/or amino acid deletion at the second position) results in non-glycosylation at the modified tripeptide sequence. Another means of increasing the number of carbohydrate moieties on a GDF Trap polypeptide is by chemical or enzymatic coupling of glycosides to the GDF Trap polypeptide. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine; (b) free carboxyl groups; (c) free sulfhydryl groups such as those of cysteine; (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline; (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan; or (f) the amide group of glutamine. These methods are described in WO 87/05330 and in Aplin and Wriston (1981) CRC Crit. Rev. Biochem., pp. 259-306,

incorporated by reference herein. Removal of one or more carbohydrate moieties present on a GDF Trap polypeptide may be accomplished chemically and/or enzymatically. Chemical deglycosylation may involve, for example, exposure of the GDF Trap polypeptide to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the amino acid sequence intact. Chemical deglycosylation is further described by Hakimuddin et al. (1987) Arch. Biochem. Biophys. 259:52 and by Edge et al. (1981) Anal. Biochem. 118:131. Enzymatic cleavage of carbohydrate moieties on GDF Trap polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al. (1987) Meth. Enzymol. 138:350. The sequence of a GDF Trap polypeptide may be adjusted, as appropriate, depending on the type of expression system used, as mammalian, yeast, insect and plant cells may all introduce differing glycosylation patterns that can be affected by the amino acid sequence of the peptide. In general, GDF Trap polypeptides for use in humans will be expressed in a mammalian cell line that provides proper glycosylation, such as HEK293 or CHO cell lines, although other mammalian expression cell lines are expected to be useful as well.

This disclosure further contemplates a method of generating variants, particularly sets of combinatorial variants of a GDF Trap polypeptide, including, optionally, truncation variants; pools of combinatorial mutants are especially useful for identifying GDF Trap sequences. The purpose of screening such combinatorial libraries may be to generate, for example, GDF Trap polypeptide variants which have altered properties, such as altered pharmacokinetics, or altered ligand binding. A variety of screening assays are provided below, and such assays may be used to evaluate variants. For example, a GDF Trap polypeptide variant may be screened for the ability to bind to an ActRIIB polypeptide, to prevent binding of an ActRIIB ligand to an ActRIIB polypeptide or to interfere with signaling caused by an ActRIIB ligand.

The activity of a GDF Trap polypeptide or its variants may also be tested in a cell-based or in vivo assay. For example, the effect of a GDF Trap polypeptide variant on the expression of genes involved in hematopoiesis may be assessed. This may, as needed, be performed in the presence of one or more recombinant ActRIIB ligand proteins (e.g., activin), and cells may be transfected so as to produce a GDF Trap polypeptide and/or variants thereof, and optionally, an ActRIIB ligand. Likewise, a GDF Trap polypeptide may be administered to a mouse or other animal, and one or more blood measurements, such as an RBC count, hemoglobin levels, hematocrit levels, iron stores, or reticulocyte count may be assessed using art recognized methods.

Combinatorially-derived variants can be generated which have a selective potency relative to a reference GDF Trap polypeptide. Such variant proteins, when expressed from recombinant DNA constructs, can be used in gene therapy protocols. Likewise, mutagenesis can give rise to variants which have intracellular half-lives dramatically different than the corresponding unmodified GDF Trap polypeptide. For example, the altered protein can be rendered either more stable or less stable to proteolytic degradation or other processes which result in destruction of, or otherwise inactivation of an unmodified GDF Trap polypeptide. Such variants, and the genes which encode them, can be utilized to alter GDF Trap polypeptide levels by modulating the half-life of the GDF Trap polypeptides. For instance, a short

half-life can give rise to more transient biological effects and, when part of an inducible expression system, can allow tighter control of recombinant GDF Trap polypeptide levels within the cell. In an Fc fusion protein, mutations may be made in the linker (if any) and/or the Fc portion to alter the half-life of the protein.

In certain embodiments, the GDF Trap polypeptides of the invention may further comprise post-translational modifications in addition to any that are naturally present in the ActRIIB polypeptides. Such modifications include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. As a result, GDF Trap polypeptides may contain non-amino acid elements, such as polyethylene glycols, lipids, poly- or mono-saccharide, and phosphates. Effects of such non-amino acid elements on the functionality of a GDF Trap polypeptide may be tested as described herein for other GDF Trap polypeptide variants. When a GDF Trap polypeptide is produced in cells by cleaving a nascent form of the GDF Trap polypeptide, post-translational processing may also be important for correct folding and/or function of the protein. Different cells (such as CHO, HeLa, MDCK, 293, WI38, NIH-3T3 or HEK293) have specific cellular machinery and characteristic mechanisms for such post-translational activities and may be chosen to ensure the correct modification and processing of the GDF Trap polypeptides.

In certain aspects, GDF Trap polypeptides include fusion proteins having at least a portion of an ActRIIB polypeptide and one or more fusion domains. Well known examples of such fusion domains include, but are not limited to, polyhistidine, Glu-Glu, glutathione S transferase (GST), thioredoxin, protein A, protein G, an immunoglobulin heavy chain constant region (e.g., an Fc), maltose binding protein (MBP), or human serum albumin. A fusion domain may be selected so as to confer a desired property. For example, some fusion domains are particularly useful for isolation of the fusion proteins by affinity chromatography. For the purpose of affinity purification, relevant matrices for affinity chromatography, such as glutathione-, amylase-, and nickel- or cobalt-conjugated resins are used. Many of such matrices are available in "kit" form, such as the Pharmacia GST purification system and the QIAexpress™ system (Qiagen) useful with (HIS₆) (SEQ ID NO: 24) fusion partners. As another example, a fusion domain may be selected so as to facilitate detection of the GDF Trap polypeptides. Examples of such detection domains include the various fluorescent proteins (e.g., GFP) as well as "epitope tags," which are usually short peptide sequences for which a specific antibody is available. Well known epitope tags for which specific monoclonal antibodies are readily available include FLAG, influenza virus haemagglutinin (HA), and c-myc tags. In some cases, the fusion domains have a protease cleavage site, such as for Factor Xa or Thrombin, which allows the relevant protease to partially digest the fusion proteins and thereby liberate the recombinant proteins therefrom. The liberated proteins can then be isolated from the fusion domain by subsequent chromatographic separation. In certain preferred embodiments, a GDF Trap polypeptide is fused with a domain that stabilizes the GDF Trap polypeptide in vivo (a "stabilizer" domain). By "stabilizing" is meant anything that increases serum half life, regardless of whether this is because of decreased destruction, decreased clearance by the kidney, or other pharmacokinetic effect. Fusions with the Fc portion of an immunoglobulin are known to confer desirable pharmacokinetic properties on a wide range of proteins. Likewise, fusions to human serum albumin can confer desirable properties. Other types of

25

fusion domains that may be selected include multimerizing (e.g., dimerizing, tetramerizing) domains and functional domains (that confer an additional biological function, such as further increasing red blood cell levels).

As a specific example, the present invention provides a GDF Trap that is an ActRIIB-Fc fusion protein which comprises an extracellular (e.g., ligand-binding) domain of ActRIIB polypeptide fused to an Fc domain. The sequence of an exemplary Fc domain is shown below (SEQ ID NO: 6).

```

THTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVD (A) VSH
EDPEVKFNWYDGVGVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGK
EYCKK (A) VSNKALPVPIEKTISKAKGQPREPQVYTLPPSREEMTKNQV
SLTCLVKGFGFYPSDIAVEWESNGQPENNYKTPPVLDSDGPFPLYSKLTV
DKSRWQQGNVSCVMHEALHN (A) HYTKSLSLSPGK*

```

Optionally, the Fc domain has one or more mutations at residues such as Asp-265, lysine 322, and Asn-434. In certain cases, the mutant Fc domain having one or more of these mutations (e.g., Asp-265 mutation) has reduced ability of binding to the Fcγ receptor relative to a wildtype Fc domain. In other cases, the mutant Fc domain having one or more of these mutations (e.g., Asn-434 mutation) has increased ability of binding to the WIC class I-related Fc-receptor (FcRN) relative to a wildtype Fc domain.

It is understood that different elements of the fusion proteins may be arranged in any manner that is consistent with the desired functionality. For example, a GDF Trap polypeptide may be placed C-terminal to a heterologous domain, or, alternatively, a heterologous domain may be placed C-terminal to a GDF Trap polypeptide. The GDF Trap polypeptide domain and the heterologous domain need not be adjacent in a fusion protein, and additional domains or amino acid sequences may be included C- or N-terminal to either domain or between the domains.

In certain embodiments, a GDF Trap fusion protein comprises an amino acid sequence as set forth in the formula A-B-C. The B portion is an N- and C-terminally truncated ActRIIB polypeptide consisting of the amino acid sequence corresponding to amino acids 26-132 of SEQ ID NO: 26. The A and C portions may be independently zero, one or more than one amino acids, and both the A and C portions when present are heterologous to B. The A and/or C portions may be attached to the B portion via a linker sequence. Exemplary linkers include short polypeptide linkers such as 2-10, 2-5, 2-4, 2-3 Glycine residues, such as, for example, a Gly-Gly-Gly linker. Other suitable linkers are described herein above. In certain embodiments, a GDF Trap fusion protein comprises an amino acid sequence as set forth in the formula A-B-C, wherein A is a leader sequence, B consists of amino acids 26-132 of SEQ ID NO: 26, and C is a polypeptide portion that enhances one or more of in vivo stability, in vivo half life, uptake/administration, tissue localization or distribution, formation of protein complexes, and/or purification. In certain embodiments, a GDF Trap fusion protein comprises an amino acid sequence as set forth in the formula A-B-C, wherein A is a TPA leader sequence, B consists of amino acids 26-132 of SEQ ID NO: 26, and C is an immunoglobulin Fc domain. A preferred GDF Trap fusion protein comprises the amino acid sequence set forth in SEQ ID NO: 26.

In certain embodiments, the GDF Trap polypeptides of the present invention contain one or more modifications that

26

are capable of stabilizing the GDF Trap polypeptides. For example, such modifications enhance the in vitro half life of the GDF Trap polypeptides, enhance circulatory half life of the GDF Trap polypeptides or reducing proteolytic degradation of the GDF Trap polypeptides. Such stabilizing modifications include, but are not limited to, fusion proteins (including, for example, fusion proteins comprising an GDF Trap polypeptide and a stabilizer domain), modifications of a glycosylation site (including, for example, addition of a glycosylation site to a GDF Trap polypeptide), and modifications of carbohydrate moiety (including, for example, removal of carbohydrate moieties from a GDF Trap polypeptide). In the case of fusion proteins, a GDF Trap polypeptide is fused to a stabilizer domain such as an IgG molecule (e.g., an Fc domain). As used herein, the term "stabilizer domain" not only refers to a fusion domain (e.g., Fc) as in the case of fusion proteins, but also includes nonproteinaceous modifications such as a carbohydrate moiety, or nonproteinaceous polymer, such as polyethylene glycol.

In certain embodiments, the present invention makes available isolated and/or purified forms of the GDF Trap polypeptides, which are isolated from, or otherwise substantially free of, other proteins.

In certain embodiments, GDF Trap polypeptides (unmodified or modified) of the invention can be produced by a variety of art-known techniques. For example, such GDF Trap polypeptides can be synthesized using standard protein chemistry techniques such as those described in Bodansky, M. Principles of Peptide Synthesis, Springer Verlag, Berlin (1993) and Grant G. A. (ed.), Synthetic Peptides: A User's Guide, W. H. Freeman and Company, New York (1992). In addition, automated peptide synthesizers are commercially available (e.g., Advanced ChemTech Model 396; Milligen/Biosearch 9600). Alternatively, the GDF Trap polypeptides, fragments or variants thereof may be recombinantly produced using various expression systems (e.g., *E. coli*, Chinese Hamster Ovary (CHO) cells, COS cells, baculovirus) as is well known in the art. In a further embodiment, the modified or unmodified GDF Trap polypeptides may be produced by digestion of recombinantly produced full-length GDF Trap polypeptides by using, for example, a protease, e.g., trypsin, thermolysin, chymotrypsin, pepsin, or paired basic amino acid converting enzyme (PACE). Computer analysis (using a commercially available software, e.g., MacVector, Omega, PCGene, Molecular Simulation, Inc.) can be used to identify proteolytic cleavage sites. Alternatively, such GDF Trap polypeptides may be produced from recombinantly produced full-length GDF Trap polypeptides such as standard techniques known in the art, such as by chemical cleavage (e.g., cyanogen bromide, hydroxylamine).

3. Nucleic Acids Encoding GDF Trap Polypeptides

In certain aspects, the invention provides isolated and/or recombinant nucleic acids encoding any of the GDF Trap polypeptides disclosed herein. SEQ ID NO: 4 encodes a naturally occurring ActRIIB precursor polypeptide, while SEQ ID NO: 5 encodes a soluble ActRIIB polypeptide, and SEQ ID NOS: 25, 27, 30 and 31 encode soluble GDF Traps. The subject nucleic acids may be single-stranded or double stranded. Such nucleic acids may be DNA or RNA molecules. These nucleic acids may be used, for example, in methods for making GDF Trap polypeptides or as direct therapeutic agents (e.g., in a gene therapy approach).

In certain aspects, the subject nucleic acids encoding GDF Trap polypeptides are further understood to include nucleic acids that are variants of SEQ ID NOS: 5, 25, 27, 30 and 31.

Variant nucleotide sequences include sequences that differ by one or more nucleotide substitutions, additions or deletions, such as allelic variants; and will, therefore, include coding sequences that differ from the nucleotide sequence of the coding sequence designated in SEQ ID NOs: 5, 25, 27, 30 and 31.

In certain embodiments, the invention provides isolated or recombinant nucleic acid sequences that are at least 80%, 85%, 90%, 95%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 5, 25, 27, 30 or 31. One of ordinary skill in the art will appreciate that nucleic acid sequences complementary to SEQ ID NO: 5, 25, 27, 30 or 31, and variants of SEQ ID NO: 5, 25, 27, 30 or 31, are also within the scope of this invention. In further embodiments, the nucleic acid sequences of the invention can be isolated, recombinant, and/or fused with a heterologous nucleotide sequence, or in a DNA library.

In other embodiments, nucleic acids of the invention also include nucleotide sequences that hybridize under highly stringent conditions to the nucleotide sequence designated in SEQ ID NO: 5, 25, 27, 30 or 31, complement sequence of SEQ ID NO: 5, 25, 27, 30 or 31, or fragments thereof. As discussed above, one of ordinary skill in the art will understand readily that appropriate stringency conditions which promote DNA hybridization can be varied. For example, one could perform the hybridization at 6.0x sodium chloride/sodium citrate (SSC) at about 45° C., followed by a wash of 2.0xSSC at 50° C. For example, the salt concentration in the wash step can be selected from a low stringency of about 2.0xSSC at 50° C. to a high stringency of about 0.2xSSC at 50° C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22° C., to high stringency conditions at about 65° C. Both temperature and salt may be varied, or temperature or salt concentration may be held constant while the other variable is changed. In one embodiment, the invention provides nucleic acids which hybridize under low stringency conditions of 6xSSC at room temperature followed by a wash at 2xSSC at room temperature.

Isolated nucleic acids which differ from the nucleic acids as set forth in SEQ ID NO: 5, 25, 27, 30 or 31 due to degeneracy in the genetic code are also within the scope of the invention. For example, a number of amino acids are designated by more than one triplet. Codons that specify the same amino acid, or synonyms (for example, CAU and CAC are synonyms for histidine) may result in "silent" mutations which do not affect the amino acid sequence of the protein. In certain embodiments, the GDF Trap polypeptide will be encoded by an alternative nucleotide sequence. Alternative nucleotide sequences are degenerate with respect to the native GDF Trap nucleic acid sequence but still encode for the same fusion protein. In certain embodiments, the GDF Trap having SEQ ID NO: 26 is encoded by an alternative nucleic acid sequence comprising SEQ ID NO: 30. However, it is expected that DNA sequence polymorphisms that do lead to changes in the amino acid sequences of the subject proteins will exist among mammalian cells. One skilled in the art will appreciate that these variations in one or more nucleotides (up to about 3-5% of the nucleotides) of the nucleic acids encoding a particular protein may exist among individuals of a given species due to natural allelic variation. Any and all such nucleotide variations and resulting amino acid polymorphisms are within the scope of this invention.

In certain embodiments, the recombinant nucleic acids of the invention may be operably linked to one or more regulatory nucleotide sequences in an expression construct. Regulatory nucleotide sequences will generally be appro-

priate to the host cell used for expression. Numerous types of appropriate expression vectors and suitable regulatory sequences are known in the art for a variety of host cells. Typically, said one or more regulatory nucleotide sequences may include, but are not limited to, promoter sequences, leader or signal sequences, ribosomal binding sites, transcriptional start and termination sequences, translational start and termination sequences, and enhancer or activator sequences. Constitutive or inducible promoters as known in the art are contemplated by the invention. The promoters may be either naturally occurring promoters, or hybrid promoters that combine elements of more than one promoter. An expression construct may be present in a cell on an episome, such as a plasmid, or the expression construct may be inserted in a chromosome. In a preferred embodiment, the expression vector contains a selectable marker gene to allow the selection of transformed host cells. Selectable marker genes are well known in the art and will vary with the host cell used.

In certain aspects of the invention, the subject nucleic acid is provided in an expression vector comprising a nucleotide sequence encoding a GDF Trap polypeptide and operably linked to at least one regulatory sequence. Regulatory sequences are art-recognized and are selected to direct expression of the GDF Trap polypeptide. Accordingly, the term regulatory sequence includes promoters, enhancers, and other expression control elements. Exemplary regulatory sequences are described in Goeddel; *Gene Expression Technology: Methods in Enzymology*, Academic Press, San Diego, Calif. (1990). For instance, any of a wide variety of expression control sequences that control the expression of a DNA sequence when operatively linked to it may be used in these vectors to express DNA sequences encoding a GDF Trap polypeptide. Such useful expression control sequences, include, for example, the early and late promoters of SV40, tet promoter, adenovirus or cytomegalovirus immediate early promoter, RSV promoters, the lac system, the trp system, the TAC or TRC system, T7 promoter whose expression is directed by T7 RNA polymerase, the major operator and promoter regions of phage lambda, the control regions for fd coat protein, the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the promoters of acid phosphatase, e.g., Pho5, the promoters of the yeast α -mating factors, the polyhedron promoter of the baculovirus system and other sequences known to control the expression of genes of prokaryotic or eukaryotic cells or their viruses, and various combinations thereof. It should be understood that the design of the expression vector may depend on such factors as the choice of the host cell to be transformed and/or the type of protein desired to be expressed. Moreover, the vector's copy number, the ability to control that copy number and the expression of any other protein encoded by the vector, such as antibiotic markers, should also be considered.

A recombinant nucleic acid of the invention can be produced by ligating the cloned gene, or a portion thereof, into a vector suitable for expression in either prokaryotic cells, eukaryotic cells (yeast, avian, insect or mammalian), or both. Expression vehicles for production of a recombinant GDF Trap polypeptide include plasmids and other vectors. For instance, suitable vectors include plasmids of the types: pBR322-derived plasmids, pEMBL-derived plasmids, pEX-derived plasmids, pBTac-derived plasmids and pUC-derived plasmids for expression in prokaryotic cells, such as *E. coli*.

Some mammalian expression vectors contain both prokaryotic sequences to facilitate the propagation of the vector in bacteria, and one or more eukaryotic transcription units

that are expressed in eukaryotic cells. The pcDNA1/amp, pcDNA1/neo, pRc/CMV, pSV2gpt, pSV2neo, pSV2-dhfr, pTk2, pRSVneo, pMSG, pSVT7, pko-neo and pHyg derived vectors are examples of mammalian expression vectors suitable for transfection of eukaryotic cells. Some of these vectors are modified with sequences from bacterial plasmids, such as pBR322, to facilitate replication and drug resistance selection in both prokaryotic and eukaryotic cells. Alternatively, derivatives of viruses such as the bovine papilloma virus (BPV-1), or Epstein-Barr virus (pHEBo, pREP-derived and p205) can be used for transient expression of proteins in eukaryotic cells. Examples of other viral (including retroviral) expression systems can be found below in the description of gene therapy delivery systems. The various methods employed in the preparation of the plasmids and in transformation of host organisms are well known in the art. For other suitable expression systems for both prokaryotic and eukaryotic cells, as well as general recombinant procedures, see *Molecular Cloning A Laboratory Manual*, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press, 1989) Chapters 16 and 17. In some instances, it may be desirable to express the recombinant polypeptides by the use of a baculovirus expression system. Examples of such baculovirus expression systems include pVL-derived vectors (such as pVL1392, pVL1393 and pVL941), pAcUW-derived vectors (such as pAcUW1), and pBlueBac-derived vectors (such as the β -gal containing pBlueBac III).

In a preferred embodiment, a vector will be designed for production of the subject GDF Trap polypeptides in CHO cells, such as a Pcmv-Script vector (Stratagene, La Jolla, Calif.), pcDNA4 vectors (Invitrogen, Carlsbad, Calif.) and pCI-neo vectors (Promega, Madison, Wisc.). As will be apparent, the subject gene constructs can be used to cause expression of the subject GDF Trap polypeptides in cells propagated in culture, e.g., to produce proteins, including fusion proteins or variant proteins, for purification.

This invention also pertains to a host cell transfected with a recombinant gene including a coding sequence (e.g., SEQ ID NO: 4, 5, 25, 27, 30 or 31) for one or more of the subject GDF Trap polypeptides. The host cell may be any prokaryotic or eukaryotic cell. For example, a GDF Trap polypeptide of the invention may be expressed in bacterial cells such as *E. coli*, insect cells (e.g., using a baculovirus expression system), yeast, or mammalian cells. Other suitable host cells are known to those skilled in the art.

Accordingly, the present invention further pertains to methods of producing the subject GDF Trap polypeptides. For example, a host cell transfected with an expression vector encoding a GDF Trap polypeptide can be cultured under appropriate conditions to allow expression of the GDF Trap polypeptide to occur. The GDF Trap polypeptide may be secreted and isolated from a mixture of cells and medium containing the GDF Trap polypeptide. Alternatively, the GDF Trap polypeptide may be retained cytoplasmically or in a membrane fraction and the cells harvested, lysed and the protein isolated. A cell culture includes host cells, media and other byproducts. Suitable media for cell culture are well known in the art. The subject GDF Trap polypeptides can be isolated from cell culture medium, host cells, or both, using techniques known in the art for purifying proteins, including ion-exchange chromatography, gel filtration chromatography, ultrafiltration, electrophoresis, and immunoaffinity purification with antibodies specific for particular epitopes of the GDF Trap polypeptides. In a preferred embodiment, the GDF Trap polypeptide is a fusion protein containing a domain which facilitates its purification.

In another embodiment, a fusion gene coding for a purification leader sequence, such as a poly-(His)/enterokinase cleavage site sequence at the N-terminus of the desired portion of the recombinant GDF Trap polypeptide, can allow purification of the expressed fusion protein by affinity chromatography using a Ni^{2+} metal resin. The purification leader sequence can then be subsequently removed by treatment with enterokinase to provide the purified GDF Trap polypeptide (e.g., see Hochuli et al., (1987) *J. Chromatography* 411:177; and Janknecht et al., *PNAS USA* 88:8972).

Techniques for making fusion genes are well known. Essentially, the joining of various DNA fragments coding for different polypeptide sequences is performed in accordance with conventional techniques, employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel et al., John Wiley & Sons: 1992).

4. Screening Assays

In certain aspects, the present invention relates to the use of the subject GDF Trap polypeptides (e.g., soluble variant ActRIIB polypeptides) to identify compounds (agents) which are agonist or antagonists of ActRIIB polypeptides. Compounds identified through this screening can be tested to assess their ability to modulate red blood cell, hemoglobin and/or reticulocyte levels in vivo or in vitro. These compounds can be tested, for example, in animal models.

There are numerous approaches to screening for therapeutic agents for increasing red blood cell or hemoglobin levels by targeting ActRIIB signaling. In certain embodiments, high-throughput screening of compounds can be carried out to identify agents that perturb ActRIIB-mediated effects on a selected cell line. In certain embodiments, the assay is carried out to screen and identify compounds that specifically inhibit or reduce binding of an ActRIIB polypeptide to its binding partner, such as an ActRIIB ligand (e.g., activin, Nodal, GDF8, GDF11 or BMP7). Alternatively, the assay can be used to identify compounds that enhance binding of an ActRIIB polypeptide to its binding partner such as an ActRIIB ligand. In a further embodiment, the compounds can be identified by their ability to interact with an ActRIIB polypeptide.

A variety of assay formats will suffice and, in light of the present disclosure, those not expressly described herein will nevertheless be comprehended by one of ordinary skill in the art. As described herein, the test compounds (agents) of the invention may be created by any combinatorial chemical method. Alternatively, the subject compounds may be naturally occurring biomolecules synthesized in vivo or in vitro. Compounds (agents) to be tested for their ability to act as modulators of tissue growth can be produced, for example, by bacteria, yeast, plants or other organisms (e.g., natural products), produced chemically (e.g., small molecules, including peptidomimetics), or produced recombinantly. Test compounds contemplated by the present invention include non-peptidyl organic molecules, peptides, polypeptides, peptidomimetics, sugars, hormones, and nucleic acid

molecules. In a specific embodiment, the test agent is a small organic molecule having a molecular weight of less than about 2,000 Daltons.

The test compounds of the invention can be provided as single, discrete entities, or provided in libraries of greater complexity, such as made by combinatorial chemistry. These libraries can comprise, for example, alcohols, alkyl halides, amines, amides, esters, aldehydes, ethers and other classes of organic compounds. Presentation of test compounds to the test system can be in either an isolated form or as mixtures of compounds, especially in initial screening steps. Optionally, the compounds may be optionally derivatized with other compounds and have derivatizing groups that facilitate isolation of the compounds. Non-limiting examples of derivatizing groups include biotin, fluorescein, digoxigenin, green fluorescent protein, isotopes, polyhistidine, magnetic beads, glutathione S transferase (GST), photoactivatable crosslinkers or any combinations thereof.

In many drug screening programs which test libraries of compounds and natural extracts, high throughput assays are desirable in order to maximize the number of compounds surveyed in a given period of time. Assays which are performed in cell-free systems, such as may be derived with purified or semi-purified proteins, are often preferred as "primary" screens in that they can be generated to permit rapid development and relatively easy detection of an alteration in a molecular target which is mediated by a test compound. Moreover, the effects of cellular toxicity or bioavailability of the test compound can be generally ignored in the *in vitro* system, the assay instead being focused primarily on the effect of the drug on the molecular target as may be manifest in an alteration of binding affinity between an ActRIIB polypeptide and its binding partner (e.g., an ActRIIB ligand).

Merely to illustrate, in an exemplary screening assay of the present invention, the compound of interest is contacted with an isolated and purified ActRIIB polypeptide which is ordinarily capable of binding to an ActRIIB ligand, as appropriate for the intention of the assay. To the mixture of the compound and ActRIIB polypeptide is then added to a composition containing an ActRIIB ligand. Detection and quantification of ActRIIB/ActRIIB ligand complexes provides a means for determining the compound's efficacy at inhibiting (or potentiating) complex formation between the ActRIIB polypeptide and its binding protein. The efficacy of the compound can be assessed by generating dose response curves from data obtained using various concentrations of the test compound. Moreover, a control assay can also be performed to provide a baseline for comparison. For example, in a control assay, isolated and purified ActRIIB ligand is added to a composition containing the ActRIIB polypeptide, and the formation of ActRIIB/ActRIIB ligand complex is quantitated in the absence of the test compound. It will be understood that, in general, the order in which the reactants may be admixed can be varied, and can be admixed simultaneously. Moreover, in place of purified proteins, cellular extracts and lysates may be used to render a suitable cell-free assay system.

Complex formation between the ActRIIB polypeptide and its binding protein may be detected by a variety of techniques. For instance, modulation of the formation of complexes can be quantitated using, for example, detectably labeled proteins such as radiolabeled (e.g., ^{32}P , ^{35}S , ^{14}C or ^3H), fluorescently labeled (e.g., FITC), or enzymatically labeled ActRIIB polypeptide or its binding protein, by immunoassay, or by chromatographic detection.

In certain embodiments, the present invention contemplates the use of fluorescence polarization assays and fluorescence resonance energy transfer (FRET) assays in measuring, either directly or indirectly, the degree of interaction between an ActRIIB polypeptide and its binding protein. Further, other modes of detection, such as those based on optical waveguides (PCT Publication WO 96/26432 and U.S. Pat. No. 5,677,196), surface plasmon resonance (SPR), surface charge sensors, and surface force sensors, are compatible with many embodiments of the invention.

Moreover, the present invention contemplates the use of an interaction trap assay, also known as the "two hybrid assay," for identifying agents that disrupt or potentiate interaction between an ActRIIB polypeptide and its binding partner. See for example, U.S. Pat. No. 5,283,317; Zervos et al. (1993) *Cell* 72:223-232; Madura et al. (1993) *J Biol Chem* 268:12046-12054; Bartel et al. (1993) *Biotechniques* 14:920-924; and Iwabuchi et al. (1993) *Oncogene* 8:1693-1696. In a specific embodiment, the present invention contemplates the use of reverse two hybrid systems to identify compounds (e.g., small molecules or peptides) that dissociate interactions between an ActRIIB polypeptide and its binding protein. See for example, Vidal and Legrain, (1999) *Nucleic Acids Res* 27:919-29; Vidal and Legrain, (1999) *Trends Biotechnol* 17:374-81; and U.S. Pat. Nos. 5,525,490; 5,955,280; and 5,965,368.

In certain embodiments, the subject compounds are identified by their ability to interact with an ActRIIB polypeptide. The interaction between the compound and the ActRIIB polypeptide may be covalent or non-covalent. For example, such interaction can be identified at the protein level using *in vitro* biochemical methods, including photo-crosslinking, radiolabeled ligand binding, and affinity chromatography (Jakoby W B et al., 1974, *Methods in Enzymology* 46: 1). In certain cases, the compounds may be screened in a mechanism based assay, such as an assay to detect compounds which bind to an ActRIIB polypeptide. This may include a solid phase or fluid phase binding event. Alternatively, the gene encoding an ActRIIB polypeptide can be transfected with a reporter system (e.g., β -galactosidase, luciferase, or green fluorescent protein) into a cell and screened against the library preferably by a high throughput screening or with individual members of the library. Other mechanism based binding assays may be used, for example, binding assays which detect changes in free energy. Binding assays can be performed with the target fixed to a well, bead or chip or captured by an immobilized antibody or resolved by capillary electrophoresis. The bound compounds may be detected usually using colorimetric or fluorescence or surface plasmon resonance.

5. Exemplary Therapeutic Uses

In certain embodiments, the GDF Trap polypeptides of the present invention can be used to increase red blood cell levels in mammals such as rodents and primates, and particularly human patients. In certain embodiments, the present invention provides methods of treating or preventing anemia in an individual in need thereof by administering to the individual a therapeutically effective amount of a GDF Trap polypeptide. These methods may be used for therapeutic and prophylactic treatments of mammals, and particularly humans.

As used herein, a therapeutic that "prevents" a disorder or condition refers to a compound that, in a statistical sample, reduces the occurrence of the disorder or condition in the treated sample relative to an untreated control sample, or delays the onset or reduces the severity of one or more symptoms of the disorder or condition relative to the

untreated control sample. The term "treating" as used herein includes prophylaxis of the named condition or amelioration or elimination of the condition once it has been established. In either case, prevention or treatment may be discerned in the diagnosis provided by a physician or other health care provider and the intended result of administration of the therapeutic agent.

As shown herein, GDF Trap polypeptides may be used to increase red blood cell, hemoglobin or reticulocyte levels in healthy individuals, and such GDF Trap polypeptides may be used in selected patient populations. Examples of appropriate patient populations include those with undesirably low red blood cell or hemoglobin levels, such as patients having an anemia, and those that are at risk for developing undesirably low red blood cell or hemoglobin levels, such as those patients that are about to undergo major surgery or other procedures that may result in substantial blood loss. In one embodiment, a patient with adequate red blood cell levels is treated with a GDF Trap polypeptide to increase red blood cell levels, and then blood is drawn and stored for later use in transfusions.

GDF Trap polypeptides disclosed herein may be used to increase red blood cell levels in patients having an anemia. When observing hemoglobin levels in humans, a level of less than normal for the appropriate age and gender category may be indicative of anemia, although individual variations are taken into account. For example, a hemoglobin level of 12 g/dl is generally considered the lower limit of normal in the general adult population. Potential causes include blood-loss, nutritional deficits, medication reaction, various problems with the bone marrow and many diseases. More particularly, anemia has been associated with a variety of disorders that include, for example, chronic renal failure, myelodysplastic syndrome, rheumatoid arthritis, bone marrow transplantation. Anemia may also be associated with the following conditions: solid tumors (e.g. breast cancer, lung cancer, colon cancer); tumors of the lymphatic system (e.g. chronic lymphocyte leukemia, non-Hodgkins and Hodgkins lymphomas); tumors of the hematopoietic system (e.g. leukemia, myelodysplastic syndrome, multiple myeloma); radiation therapy; chemotherapy (e.g. platinum containing regimens); inflammatory and autoimmune diseases, including, but not limited to, rheumatoid arthritis, other inflammatory arthritides, systemic lupus erythematosus (SLE), acute or chronic skin diseases (e.g. psoriasis), inflammatory bowel disease (e.g. Crohn's disease and ulcerative colitis); acute or chronic renal disease or failure including idiopathic or congenital conditions; acute or chronic liver disease; acute or chronic bleeding; situations where transfusion of red blood cells is not possible due to patient allo- or auto-antibodies and/or for religious reasons (e.g. some Jehovah's Witnesses); infections (e.g. malaria, osteomyelitis); hemoglobinopathies, including, for example, sickle cell disease, thalassemias; drug use or abuse, e.g. alcohol misuse; pediatric patients with anemia from any cause to avoid transfusion; and elderly patients or patients with underlying cardiopulmonary disease with anemia who cannot receive transfusions due to concerns about circulatory overload.

GDF Trap polypeptides would be appropriate for treating anemias of hypoproliferative bone marrow, which are typically associated with little change in red blood cell (RBC) morphology. Hypoproliferative anemias include: 1) anemia of chronic disease, 2) anemia of kidney disease, and 3) anemia associated with hypometabolic states. In each of these types, endogenous erythropoietin levels are inappropriately low for the degree of anemia observed. Other hypoproliferative anemias include: 4) early-stage iron-defi-

cient anemia, and 5) anemia caused by damage to the bone marrow. In these types, endogenous erythropoietin levels are appropriately elevated for the degree of anemia observed.

The most common type is anemia of chronic disease, which encompasses inflammation, infection, tissue injury, and conditions such as cancer, and is distinguished by both low erythropoietin levels and an inadequate response to erythropoietin in the bone marrow (Adamson, 2008, Harrison's Principles of Internal Medicine, 17th ed.; McGraw Hill, New York, pp 628-634). Many factors can contribute to cancer-related anemia. Some are associated with the disease process itself and the generation of inflammatory cytokines such as interleukin-1, interferon-gamma, and tumor necrosis factor (Bron et al., 2001, Semin Oncol 28(Suppl 8):1-6). Among its effects, inflammation induces the key iron-regulatory peptide hepcidin, thereby inhibiting iron export from macrophages and generally limiting iron availability for erythropoiesis (Ganz, 2007, J Am Soc Nephrol 18:394-400). Blood loss through various routes can also contribute to cancer-related anemia. The prevalence of anemia due to cancer progression varies with cancer type, ranging from 5% in prostate cancer up to 90% in multiple myeloma. Cancer-related anemia has profound consequences for patients, including fatigue and reduced quality of life, reduced treatment efficacy, and increased mortality.

Chronic kidney disease is associated with hypoproliferative anemia that varies in severity with the degree of renal impairment. Such anemia is primarily due to inadequate production of erythropoietin and reduced survival of red blood cells. Chronic kidney disease usually proceeds gradually over a period of years or decades to end-stage (Stage-5) disease, at which point dialysis or kidney transplantation is required for patient survival. Anemia often develops early in this process and worsens as disease progresses. The clinical consequences of anemia of kidney disease are well-documented and include development of left ventricular hypertrophy, impaired cognitive function, reduced quality of life, and altered immune function (Levin et al., 1999, Am J Kidney Dis 27:347-354; Nissenson, 1992, Am J Kidney Dis 20(Suppl 1):21-24; Revicki et al., 1995, Am J Kidney Dis 25:548-554; Gaftor et al., 1994, Kidney Int 45:224-231). As demonstrated by the Applicants in a mouse model of chronic kidney disease (see Example below), a GDF Trap polypeptide can be used to treat anemia of kidney disease.

Many conditions resulting in a hypometabolic rate can produce a mild-to-moderate hypoproliferative anemia. Among such conditions are endocrine deficiency states. For example, anemia can occur in Addison's disease, hypothyroidism, hyperparathyroidism, or males who are castrated or treated with estrogen. Mild-to-moderate anemia can also occur with reduced dietary intake of protein, a condition particularly prevalent in the elderly. Finally, anemia can develop in patients with chronic liver disease arising from nearly any cause (Adamson, 2008, Harrison's Principles of Internal Medicine, 17th ed.; McGraw Hill, New York, pp 628-634).

Anemia resulting from acute blood loss of sufficient volume, such as from trauma or postpartum hemorrhage, is known as acute post-hemorrhagic anemia. Acute blood loss initially causes hypovolemia without anemia since there is proportional depletion of RBCs along with other blood constituents. However, hypovolemia will rapidly trigger physiologic mechanisms that shift fluid from the extravascular to the vascular compartment, which results in hemodilution and anemia. If chronic, blood loss gradually depletes body iron stores and eventually leads to iron deficiency. As demonstrated by the Applicants in a mouse

model (see Example below), a GDF Trap polypeptide can be used to speed recovery from anemia of acute blood loss.

Iron-deficiency anemia is the final stage in a graded progression of increasing iron deficiency which includes negative iron balance and iron-deficient erythropoiesis as intermediate stages. Iron deficiency can result from increased iron demand, decreased iron intake, or increased iron loss, as exemplified in conditions such as pregnancy, inadequate diet, intestinal malabsorption, acute or chronic inflammation, and acute or chronic blood loss. With mild-to-moderate anemia of this type, the bone marrow remains hypoproliferative, and RBC morphology is largely normal; however, even mild anemia can result in some microcytic hypochromic RBCs, and the transition to severe iron-deficient anemia is accompanied by hyperproliferation of the bone marrow and increasingly prevalent microcytic and hypochromic RBCs (Adamson, 2008, Harrison's Principles of Internal Medicine, 17th ed.; McGraw Hill, New York, pp 628-634). Appropriate therapy for iron-deficiency anemia depends on its cause and severity, with oral iron preparations, parenteral iron formulations, and RBC transfusion as major conventional options. An GDF Trap polypeptide could be used to treat chronic iron-deficiency anemias alone or in combination with conventional therapeutic approaches, particularly to treat anemias of multifactorial origin.

Hypoproliferative anemias can result from primary dysfunction or failure of the bone marrow, instead of dysfunction secondary to inflammation, infection, or cancer progression. Prominent examples would be myelosuppression caused by cancer chemotherapeutic drugs or cancer radiation therapy. A broad review of clinical trials found that mild anemia can occur in 100% of patients after chemotherapy, while more severe anemia can occur in up to 80% of such patients (Groopman et al., 1999, J Natl Cancer Inst 91:1616-1634). Myelosuppressive drugs include: 1) alkylating agents such as nitrogen mustards (e.g., melphalan) and nitrosoureas (e.g., streptozocin); 2) antimetabolites such as folic acid antagonists (e.g., methotrexate), purine analogs (e.g., thioguanine), and pyrimidine analogs (e.g., gemcitabine); 3) cytotoxic antibiotics such as anthracyclines (e.g., doxorubicin); 4) kinase inhibitors (e.g., gefitinib); 5) mitotic inhibitors such as taxanes (e.g., paclitaxel) and *vinca* alkaloids (e.g., vinorelbine); 6) monoclonal antibodies (e.g., rituximab); and 7) topoisomerase inhibitors (e.g., topotecan and etoposide). As demonstrated in a mouse model of chemotherapy-induced anemia (see Example below), a GDF Trap polypeptide can be used to treat anemia caused by chemotherapeutic agents and/or radiation therapy.

GDF Trap polypeptides would also be appropriate for treating anemias of disordered RBC maturation, which are characterized in part by undersized (microcytic), oversized (macrocytic), misshapen, or abnormally colored (hypochromic) RBCs.

Patients may be treated with a dosing regimen intended to restore the patient to a target hemoglobin level, usually between about 10 g/dl and about 12.5 g/dl, and typically about 11.0 g/dl (see also Jacobs et al. (2000) Nephrol Dial Transplant 15, 15-19), although lower target levels may cause fewer cardiovascular side effects. Alternatively, hematocrit levels (percentage of the volume of a blood sample occupied by the cells) can be used as a measure for the condition of red blood cells. Hematocrit levels for healthy individuals range from 41 to 51% for adult males and from 35 to 45% for adult females. Target hematocrit levels are usually around 30-33%. Moreover, hemoglobin/hematocrit

levels vary from person to person. Thus, optimally, the target hemoglobin/hematocrit level can be individualized for each patient.

The rapid effect on red blood cell levels of the GDF Trap polypeptides disclosed herein indicate that these agents act by a different mechanism than Epo. Accordingly, these antagonists may be useful for increasing red blood cell and hemoglobin levels in patients that do not respond well to Epo. For example, a GDF Trap polypeptide may be beneficial for a patient in which administration of a normal to increased (>300 IU/kg/week) dose of Epo does not result in the increase of hemoglobin level up to the target level. Patients with an inadequate Epo response are found for all types of anemia, but higher numbers of non-responders have been observed particularly frequently in patients with cancers and patients with end-stage renal disease. An inadequate response to Epo can be either constitutive (i.e. observed upon the first treatment with Epo) or acquired (e.g. observed upon repeated treatment with Epo).

The GDF Trap polypeptides may also be used to treat patients that are susceptible to adverse effects of Epo. The primary adverse effects of Epo are an excessive increase in the hematocrit or hemoglobin levels and polycythemia. Elevated hematocrit levels can lead to hypertension (more particularly aggravation of hypertension) and vascular thrombosis. Other adverse effects of Epo which have been reported, some of which related to hypertension, are headaches, influenza-like syndrome, obstruction of shunts, myocardial infarctions and cerebral convulsions due to thrombosis, hypertensive encephalopathy, and red cell blood cell aplasia (Singibarti, (1994) J. Clin Investig 72(suppl 6), S36-S43; Horl et al. (2000) Nephrol Dial Transplant 15(suppl 4), 51-56; Delanty et al. (1997) Neurology 49, 686-689; Bunn (2002) N Engl J Med 346(7), 522-523).

GDF traps may also be used in combination with Epo and other agents that activate the erythropoietin pathway. In some instances, this may permit lower dosing of each drug in the combination.

In certain embodiments, the present invention provides methods for managing a patient that has been treated with, or is a candidate to be treated with, a GDF Trap polypeptide by measuring one or more hematologic parameters in the patient. The hematologic parameters may be used to evaluate appropriate dosing for a patient who is a candidate to be treated with a GDF Trap polypeptide, to monitor the hematologic parameters during treatment with a GDF Trap polypeptide, to evaluate whether to adjust the dosage during treatment with a GDF Trap polypeptide, and/or to evaluate an appropriate maintenance dose of a GDF Trap polypeptide. If one or more of the hematologic parameters are outside the normal level, dosing with a GDF Trap polypeptide may be reduced, delayed or terminated.

Hematologic parameters that may be measured in accordance with the methods provided herein include, for example, red blood cell levels, blood pressure, iron stores, and other agents found in bodily fluids that correlate with increased red blood cell levels, using art recognized methods. Such parameters may be determined using a blood sample from a patient. Increases in red blood cell levels, hemoglobin levels, and/or hematocrit levels may cause increases in blood pressure.

In one embodiment, if one or more hematologic parameters are outside the normal range, or on the high side of normal, in a patient who is a candidate to be treated with a GDF Trap polypeptide then onset of administration of the GDF Trap polypeptide may be delayed until the hematologic parameters have returned to a normal or acceptable level

either naturally or via therapeutic intervention. For example, if a candidate patient is hypertensive or prehypertensive, then the patient may be treated with a blood pressure lowering agent in order to reduce the patient's blood pressure. Any blood pressure lowering agent appropriate for the individual patient's condition may be used including, for example, diuretics, adrenergic inhibitors (including alpha blockers and beta blockers), vasodilators, calcium channel blockers, angiotensin-converting enzyme (ACE) inhibitors, or angiotensin II receptor blockers. Blood pressure may alternatively be treated using a diet and exercise regimen. Similarly, if a candidate patient has iron stores that are lower than normal, or on the low side of normal, then the patient may be treated with an appropriate regimen of diet and/or iron supplements until the patient's iron stores have returned to a normal or acceptable level. For patients having higher than normal red blood cell levels and/or hemoglobin levels, then administration of the GDF Trap polypeptide may be delayed until the levels have returned to a normal or acceptable level.

In certain embodiments, if one or more hematologic parameters are outside the normal range, or on the high side of normal, in a patient who is a candidate to be treated with a GDF Trap polypeptide then the onset of administration may be not be delayed. However, the dosage amount or frequency of dosing of the GDF Trap polypeptide may be set at an amount that would reduce the risk of an unacceptable increase in the hematologic parameters arising upon administration of the GDF Trap polypeptide. Alternatively, a therapeutic regimen may be developed for the patient that combines a GDF Trap polypeptide with a therapeutic agent that addresses the undesirable level of the hematologic parameter. For example, if the patient has elevated blood pressure, then a therapeutic regimen involving administration of a GDF Trap polypeptide and a blood pressure lowering agent may be designed. For a patient having lower than desired iron stores, a therapeutic regimen of a GDF Trap polypeptide and iron supplementation may be developed.

In one embodiment, baseline parameter(s) for one or more hematologic parameters may be established for a patient who is a candidate to be treated with a GDF Trap polypeptide and an appropriate dosing regimen establish for that patient based on the baseline value(s). Alternatively, established baseline parameters based on a patient's medical history could be used to inform an appropriate GDF Trap polypeptide dosing regimen for a patient. For example, if a healthy patient has an established baseline blood pressure reading that is above the defined normal range it may not be necessary to bring the patient's blood pressure into the range that is considered normal for the general population prior to treatment with the GDF Trap polypeptide. A patient's baseline values for one or more hematologic parameters prior to treatment with a GDF Trap polypeptide may also be used as the relevant comparative values for monitoring any changes to the hematologic parameters during treatment with the GDF Trap polypeptide.

In certain embodiments, one or more hematologic parameters are measured in patients who are being treated with a GDF Trap polypeptide. The hematologic parameters may be used to monitor the patient during treatment and permit adjustment or termination of the dosing with the GDF Trap polypeptide or additional dosing with another therapeutic agent. For example, if administration of a GDF Trap polypeptide results in an increase in blood pressure, red blood cell level, or hemoglobin level, or a reduction in iron stores, then the dose of the GDF Trap polypeptide may be reduced

in amount or frequency in order to decrease the effects of the GDF Trap polypeptide on the one or more hematologic parameters. If administration of a GDF Trap polypeptide results in a change in one or more hematologic parameters that is adverse to the patient, then the dosing of the GDF Trap polypeptide may be terminated either temporarily, until the hematologic parameter(s) return to an acceptable level, or permanently. Similarly, if one or more hematologic parameters are not brought within an acceptable range after reducing the dose or frequency of administration of the GDF Trap polypeptide then the dosing may be terminated. As an alternative, or in addition to, reducing or terminating the dosing with the GDF Trap polypeptide, the patient may be dosed with an additional therapeutic agent that addresses the undesirable level in the hematologic parameter(s), such as, for example, a blood pressure lowering agent or an iron supplement. For example, if a patient being treated with a GDF Trap polypeptide has elevated blood pressure, then dosing with the GDF Trap polypeptide may continue at the same level and a blood pressure lowering agent is added to the treatment regimen, dosing with the GDF Trap polypeptide may be reduced (e.g., in amount and/or frequency) and a blood pressure lowering agent is added to the treatment regimen, or dosing with the GDF Trap polypeptide may be terminated and the patient may be treated with a blood pressure lowering agent.

In certain embodiments, patients being treated with a GDF Trap polypeptide, or candidate patients to be treated with a GDF Trap polypeptide, are patients in need of muscle growth, such as patients suffering from, or at risk of developing, a neuromuscular disorder or musculogenerative disorder. For example, patients or candidate patients may be suffering from, or at risk for developing, Lou Gehrig's disease (ALS), cancer anorexia-cachexia syndrome, muscular dystrophy, muscle atrophy, congestive obstructive pulmonary disease (and muscle wasting associated with COPD), muscle wasting syndrome, sarcopenia, or cachexia. Muscular dystrophy refers to a group of degenerative muscle diseases characterized by gradual weakening and deterioration of skeletal muscles and sometimes the heart and respiratory muscles. Exemplary muscular dystrophies that can be treated with a regimen including the subject GDF Trap polypeptides include: Duchenne Muscular Dystrophy (DMD), Becker Muscular Dystrophy (BMD), Emery-Dreifuss Muscular Dystrophy (EDMD), Limb-Girdle Muscular Dystrophy (LGMD), Facioscapulohumeral Muscular Dystrophy (FSH or FSHD) (also known as Landouzy-Dejerine), Myotonic Dystrophy (MMD) (also known as Steinert's Disease), Oculopharyngeal Muscular Dystrophy (OPMD), Distal Muscular Dystrophy (DD), Congenital Muscular Dystrophy (CMD).

6. Pharmaceutical Compositions

In certain embodiments, compounds (e.g., GDF Trap polypeptides) of the present invention are formulated with a pharmaceutically acceptable carrier. For example, a GDF Trap polypeptide can be administered alone or as a component of a pharmaceutical formulation (therapeutic composition). The subject compounds may be formulated for administration in any convenient way for use in human or veterinary medicine.

In certain embodiments, the therapeutic method of the invention includes administering the composition systemically, or locally as an implant or device. When administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Therapeutically useful agents other than the GDF Trap polypeptides which may also optionally be included in the

composition as described above, may be administered simultaneously or sequentially with the subject compounds (e.g., GDF Trap polypeptides) in the methods of the invention.

Typically, compounds will be administered parenterally. Pharmaceutical compositions suitable for parenteral administration may comprise one or more GDF Trap polypeptides in combination with one or more pharmaceutically acceptable sterile isotonic aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents. Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

Further, the composition may be encapsulated or injected in a form for delivery to a target tissue site (e.g., bone marrow). In certain embodiments, compositions of the present invention may include a matrix capable of delivering one or more therapeutic compounds (e.g., GDF Trap polypeptides) to a target tissue site (e.g., bone marrow), providing a structure for the developing tissue and optimally capable of being resorbed into the body. For example, the matrix may provide slow release of the GDF Trap polypeptides. Such matrices may be formed of materials presently in use for other implanted medical applications.

The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. The particular application of the subject compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium sulfate, tricalciumphosphate, hydroxyapatite, polylactic acid and polyanhydrides. Other potential materials are biodegradable and biologically well defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are non-biodegradable and chemically defined, such as sintered hydroxyapatite, bioglass, aluminates, or other ceramics. Matrices may be comprised of combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalciumphosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability.

In certain embodiments, methods of the invention can be administered for orally, e.g., in the form of capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia) and/or as mouth washes and the like, each containing a predetermined amount of an agent as an active ingredient. An agent may also be administered as a bolus, electuary or paste.

In solid dosage forms for oral administration (capsules, tablets, pills, dragees, powders, granules, and the like), one

or more therapeutic compounds of the present invention may be mixed with one or more pharmaceutically acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose, and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds; (7) wetting agents, such as, for example, cetyl alcohol and glycerol monostearate; (8) absorbents, such as kaolin and bentonite clay; (9) lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and (10) coloring agents. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups, and elixirs. In addition to the active ingredient, the liquid dosage forms may contain inert diluents commonly used in the art, such as water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming, and preservative agents.

Suspensions, in addition to the active compounds, may contain suspending agents such as ethoxylated isostearyl alcohols, polyoxyethylene sorbitol, and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

The compositions of the invention may also contain adjuvants, such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption, such as aluminum monostearate and gelatin.

It is understood that the dosage regimen will be determined by the attending physician considering various factors which modify the action of the subject compounds of the invention (e.g., GDF Trap polypeptides). The various factors include, but are not limited to, the patient's red blood cell count, hemoglobin level or other diagnostic assessments, the desired target red blood cell count, the patient's age, sex, and diet, the severity of any disease that may be contributing to a depressed red blood cell level, time of administration, and other clinical factors. The addition of other known growth factors to the final composition may also affect the dosage.

Progress can be monitored by periodic assessment of red blood cell and hemoglobin levels, as well as assessments of reticulocyte levels and other indicators of the hematopoietic process.

In certain embodiments, the present invention also provides gene therapy for the in vivo production of GDF Trap polypeptides. Such therapy would achieve its therapeutic effect by introduction of the GDF Trap polynucleotide sequences into cells or tissues having the disorders as listed above. Delivery of GDF Trap polynucleotide sequences can be achieved using a recombinant expression vector such as a chimeric virus or a colloidal dispersion system. Preferred for therapeutic delivery of GDF Trap polynucleotide sequences is the use of targeted liposomes.

Various viral vectors which can be utilized for gene therapy as taught herein include adenovirus, herpes virus, vaccinia, or an RNA virus such as a retrovirus. The retroviral vector may be a derivative of a murine or avian retrovirus. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), and Rous Sarcoma Virus (RSV). A number of additional retroviral vectors can incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. Retroviral vectors can be made target-specific by attaching, for example, a sugar, a glycolipid, or a protein. Preferred targeting is accomplished by using an antibody. Those of skill in the art will recognize that specific polynucleotide sequences can be inserted into the retroviral genome or attached to a viral envelope to allow target specific delivery of the retroviral vector containing the GDF Trap polynucleotide.

Alternatively, tissue culture cells can be directly transfected with plasmids encoding the retroviral structural genes gag, pol and env, by conventional calcium phosphate transfection. These cells are then transfected with the vector plasmid containing the genes of interest. The resulting cells release the retroviral vector into the culture medium.

Another targeted delivery system for GDF Trap polynucleotides is a colloidal dispersion system. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. The preferred colloidal system of this invention is a liposome. Liposomes are artificial membrane vesicles which are useful as delivery vehicles in vitro and in vivo. RNA, DNA and intact virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (see e.g., Fraley, et al., Trends Biochem. Sci., 6:77, 1981). Methods for efficient gene transfer using a liposome vehicle, are known in the art, see e.g., Mannino, et al., Biotechniques, 6:682, 1988. The composition of the liposome is usually a combination of phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations.

Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebroside, and gangliosides. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine, and distearoylphosphatidylcholine. The targeting of liposomes is also possible based

on, for example, organ-specificity, cell-specificity, and organelle-specificity and is known in the art.

EXEMPLIFICATION

The invention now being generally described, it will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain embodiments and embodiments of the present invention, and are not intended to limit the invention.

Example 1. Generation of a GDF Trap

Applicants constructed a GDF Trap as follows. A polypeptide having a modified extracellular domain of ActRIIB with greatly reduced activin A binding relative to GDF11 and/or myostatin (as a consequence of a leucine-to-aspartate substitution at position 79 in SEQ ID NO: 1) was fused to a human or mouse Fc domain with a minimal linker (three glycine amino acids) in between. The constructs are referred to as ActRIIB(L79D 20-134)-hFc and ActRIIB(L79D 20-134)-mFc, respectively. Alternative forms with a glutamate rather than an aspartate at position 79 performed similarly (L79E). Alternative forms with an alanine rather than a valine at position 226 with respect to SEQ ID NO: 7, below were also generated and performed equivalently in all respects tested. The aspartate at position 79 (relative to SEQ ID NO: 1, or position 60 relative to SEQ ID NO: 7) is highlighted in gray below. The valine at position 226 relative to SEQ ID NO: 7 is also highlighted in gray below.

The GDF Trap ActRIIB(L79D 20-134)-hFc is shown below as purified from CHO cell lines (SEQ ID NO: 7).

```
GRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWRNSSG
TIELVKKGCDWDDFNCYDRQECVATEENPQVYFCCCEGNFCNERFTHLP
EAGGPEVTYEPPTAPTGGGTHTCPPCPAPPELLGGPSVFLFPPKPKDTL
MISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY
RVVSVLTVLHQDWLNGKEYKCKVSNKALPWPPIEKTISKAKGQPREPQVY
TLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVL
DSGSGFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNNHYTQKSLSLSPGK
```

The ActRIIB-derived portion of the GDF Trap has an amino acid sequence set forth below (SEQ ID NO: 32), and that portion could be used as a monomer or as a non-Fc fusion protein as a monomer, dimer or greater order complex.

```
(SEQ ID NO: 32)
GRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWRNSS
GTIELVKKGCDWDDFNCYDRQECVATEENPQVYFCCCEGNFCNERFTH
LPEAGGPEVTYEPPTAPT
```

43

The GDF Trap protein was expressed in CHO cell lines. Three different leader sequences were considered:

(i) Honey bee melittin (HBML): (SEQ ID NO: 8) 5
MKFLVNVAVFMVVIISYIYA

(ii) Tissue Plasminogen Activator (TPA): (SEQ ID NO: 9)
MDAMKRGLCCVLLLCGAVFVSP 10

(iii) Native: (SEQ ID NO: 10)
MTAPWVALALLWGSLCAGS.

The selected form employs the TPA leader and has the following unprocessed amino acid sequence:

(SEQ ID NO: 11)
MDAMKRGLCCVLLLCGAVFVSPGASGRGEAETRECIYYNANWELERTNQ
SGLERCEGEQDKRLHCYASWRNSSGTIELVKKGCWDDDFNCYDRQECVA
TEENPQVYFCCCEGNFCNERFTHLPEAGGPEVTYEPPTAPTGGGTHTC
PCPAPPELLGGPSVFLFPKPKDTLMISRTPEVTCVVVDVSHEDPEVKF
NWYVDGVEVHNAAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVLS
NKAALPVPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFY
PSDIAVEWESNGQPENNYKTTPPVLDSDGSFPLYSKLTVDKSRWQQGNV
FSCSVMHLEALHNHYTQKSLSLSPGK

This polypeptide is encoded by the following nucleic acid sequence (SEQ ID NO:12):

A TGGATGCAAT GAAGAGAGGG CTCTGCTGTG TGCTGCTGCT
GTGTGGAGCA GTCTTCGTTT CGCCCGGCGC CTCTGGGCGT
GGGGAGGCTG AGACACGGGA GTGCATCTAC TACAACGCCA
ACTGGGAGCT GGAGCGCACC AACAGAGCG GCCTGGAGCG
CTGCGAAGGC GAGCAGGACA AGCGGCTGCA CTGCTACGCC
TCCTGGCGCA ACAGCTCTGG CACCATCGAG CTCGTGAAGA
AGGGCTGCTG GGACGATGAC TTCAACTGCT ACGATAGGCA
GGAGTGTGTG GCCACTGAGG AGAACCCCA GGTGTACTTC
TGCTGTGTG AAGGCAACTT CTGCAACGAG CGCTTCACTC
ATTTGCCAGA GGCTGGGGGC CCGGAAGTCA CGTACGAGCC
ACCCCGACA GCCCCACCG GTGGTGAAC TCACACATGC
CCACCGTGCC CAGCACCTGA ACTCCTGGG GGACCGTCAG
TCTTCCTCTT CCCCCAAAA CCAAGGACA CCCTCATGAT
CTCCCGGACC CCTGAGGTCA CATGCGTGGT GGTGGACGTG
AGCCACGAAG ACCCTGAGGT CAAGTTCAAC TGTACGTGG
ACGGCGTGGA GGTGCATAAT GCCAAGACAA AGCCGCGGGA
GGAGCAGTAC AACAGCACGT ACCGTGTGGT CAGCGTCTTC
ACCGTCTGC ACCAGGACTG GCTGAATGGC AAGGAGTACA
AGTGCAAGGT CTCCAACAAA GCCCTCCAG TCCCCATCGA
GAAAACCATC TCCAAGCCA AAGGCGAGCC CCGAGAACCA

44

-continued

CAGGTGTACA CCCTGCCCC ATCCCGGGAG GAGATGACCA
AGAACCAGGT CAGCCTGACC TGCCTGGTCA AAGGCTTCTA
TCCCAGCGAC ATCGCCGTGG AGTGGGAGAG CAATGGGCAG
CCGGAGAACA ACTACAAGAC CACGCCTCCC GTGCTGGACT
CCGACGGCTC CTTCTTCCTC TATAGCAAGC TCACCGTGA
CAAGAGCAGG TGGCAGCAGG GGAACGTCTT CTCATGCTCC
GTGATGCATG AGGCTCTGCA CAACCACTAC ACGCAGAAGA
GCCTCTCCCT GTCTCCGGGT AAATGA

Purification could be achieved by a series of column chromatography steps, including, for example, three or more of the following, in any order: protein A chromatography, Q sepharose chromatography, phenylsepharose chromatography, size exclusion chromatography, and cation exchange chromatography. The purification could be completed with viral filtration and buffer exchange. In an example of a purification scheme, the cell culture medium is passed over a protein A column, washed in 150 mM Tris/NaCl (pH 8.0), then washed in 50 mM Tris/NaCl (pH 8.0) and eluted with 0.1 M glycine, pH 3.0. The low pH eluate is kept at room temperature for 30 minutes as a viral clearance step. The eluate is then neutralized and passed over a Q sepharose ion exchange column and washed in 50 mM Tris pH 8.0, 50 mM NaCl, and eluted in 50 mM Tris pH 8.0, with an NaCl concentration between 150 mM and 300 mM. The eluate is then changed into 50 mM Tris pH 8.0, 1.1 M ammonium sulfate and passed over a phenyl sepharose column, washed, and eluted in 50 mM Tris pH 8.0 with ammonium sulfate between 150 and 300 mM. The eluate is dialyzed and filtered for use.

Additional GDF Traps (ActRIIB-Fc fusion proteins modified so as to reduce the ratio of activin A binding relative to myostatin or GDF11) are described in PCT/US2008/001506 and WO 2006/012627, incorporated by reference herein.

Example 2. Bioassay for GDF-11 and Activin-Mediated Signaling

An A-204 Reporter Gene Assay was used to evaluate the effects of ActRIIB-Fc proteins and GDF Traps on signaling by GDF-11 and Activin A. Cell line: Human Rhabdomyosarcoma (derived from muscle). Reporter vector: pGL3 (CAGA)12 (Described in Dennler et al, 1998, EMBO 17: 3091-3100). The CAGA12 motif is present in TGF-Beta responsive genes (PAI-1 gene), so this vector is of general use for factors signaling through Smad2 and 3.

Day 1: Split A-204 cells into 48-well plate.

Day 2: A-204 cells transfected with 10 ug pGL3(CAGA)12 or pGL3(CAGA)12(10 ug)+pRLCMV (1 ug) and Eugene.

Day 3: Add factors (diluted into medium+0.1% BSA). Inhibitors need to be preincubated with Factors for 1 hr before adding to cells. 6 hrs later, cells rinsed with PBS, and lyse cells.

This is followed by a Luciferase assay. In the absence of any inhibitors, Activin A showed 10 fold stimulation of reporter gene expression and an ED50~2 ng/ml. GDF-11: 16 fold stimulation, ED50: ~1.5 ng/ml.

ActRIIB(20-134) is a potent inhibitor of activin, GDF-8 and GDF-11 activity in this assay. Variants were tested in this assay as well.

45

Example 3. GDF-11 Inhibition by N-Terminal and C-Terminal Truncations

Variants of ActRIIB(20-134)-hFc with truncations at the N-terminus and/or C-terminus were generated and tested for activity as inhibitors of GDF-11 and activin. The activities are shown below (as measured in conditioned media): C-Terminal ActRIIB-hFc Truncations:

	IC50 (ng/mL)	
	GDF-11	Activin
ActRIIB(20-134)-hFc	45	22
ActRIIB(20-132)-hFc	87	32
ActRIIB(20-131)-hFc	120	44
ActRIIB(20-128)-hFc	130	158

As can be seen, truncations of three (ending with . . . PPT), six (ending with . . . YEP) or more amino acids at the C-terminus causes a threefold or greater decrease in the activity of the molecule. The truncation of the final 15 amino acids of the ActRIIB portion causes a greater loss of activity (see WO2006/012627).

Amino terminal truncations were made in the background of an ActRIIB(20-131)-hFc protein. The activities are shown below (as measured in conditioned media): N-Terminal ActRIIB-hFc Truncations:

	IC50 (ng/mL)	
	GDF-11	Activin
ActRIIB(20-131)-hFc (GRG . . .)	183	201
ActRIIB(21-131)-hFc (RGE . . .)	121	325
ActRIIB(22-131)-hFc (GEA . . .)	71	100
ActRIIB (23-131)-hFc (EAE . . .)	60	43
ActRIIB(24-131)-hFc (AET . . .)	69	105

Accordingly, truncations of two, three or four amino acids from the N-terminus lead to the production of a more active protein than the versions with a full-length extracellular domain. Additional experiments show that a truncation of five amino acids, ActRIIB(25-131)-hFc has activity equivalent to the untruncated form, and additional deletions at the N-terminus continue to degrade the activity of the protein. Therefore, optimal constructs will have a C-terminus ending between amino acid 133-134 of SEQ ID NO: 1 and an N-terminus beginning at amino acids 22-24 of SEQ ID NO: 1. An N-terminus corresponding to amino acids 21 or 25 will give activity that is similar to the ActRIIB(20-134)-hFc construct. These truncations may also be used in the context of GDF Traps, such as an L79D or L79E variant.

Example 4. ActRIIB-Fc Variants, Cell-Based Activity

Activity of ActRIIB-Fc proteins and GDF Traps was tested in a cell based assay, as described above. Results are summarized in the table below. Some variants were tested in different C-terminal truncation constructs. As discussed above, truncations of five or fifteen amino acids caused reduction in activity. The GDF Traps (L79D and L79E

46

variants) showed substantial loss of activin binding while retaining almost wild-type inhibition of GDF-11. Soluble ActRIIB-Fc Binding to GDF11 and Activin A:

ActRIIB-Fc Variations	Portion of ActRIIB (corresponds to amino acids of SEQ ID NO: 1)	GDF11 Inhibition Activity	Activin Inhibition Activity
R64	20-134	+++ (approx. 10^{-8} M K_D)	+++ (approx. 10^{-8} M K_D)
A64	20-134	+ (approx. 10^{-6} M K_D)	+ (approx. 10^{-6} M K_D)
R64	20-129	+++	+++
R64 K74A	20-134	++++	++++
R64 A24N	20-134	+++	+++
R64 A24N	20-119	++	++
R64 A24N K74A	20-119	+	+
R64 L79P	20-134	+	+
R64 L79P K74A	20-134	+	+
R64 L79D	20-134	+++	+
R64 L79E	20-134	+++	+
R64K	20-134	+++	+++
R64K	20-129	+++	+++
R64 P129S P130A	20-134	+++	+++
R64N	20-134	+	+

+ Poor activity (roughly 1×10^{-6} K_D)
++ Moderate activity (roughly 1×10^{-7} K_D)
+++ Good (wild-type) activity (roughly 1×10^{-8} K_D)
++++ Greater than wild-type activity

Several variants have been assessed for serum half-life in rats. ActRIIB(20-134)-Fc has a serum half-life of approximately 70 hours. ActRIIB(A24N 20-134)-Fc has a serum half-life of approximately 100-150 hours. The A24N variant has activity in the cell-based assay (above) and in vivo assays (below) that are equivalent to the wild-type molecule. Coupled with the longer half-life, this means that over time an A24N variant will give greater effect per unit of protein than the wild-type molecule. The A24N variant, and any of the other variants tested above, may be combined with the GDF Trap molecules, such as the L79D or L79E variants.

Example 5. GDF-11 and Activin A Binding

Binding of certain ActRIIB-Fc proteins and GDF Traps to ligands was tested in a BiaCore™ assay.

The ActRIIB-Fc variants or wild-type protein were captured onto the system using an anti-hFc antibody. Ligands were injected and flowed over the captured receptor proteins. Results are summarized in the tables, below. Ligand Binding Specificity IIB Variants.

Protein	Kon (1/Ms)	Koff (1/s)	KD (M)
GDF11			
ActRIIB(20-134)-hFc	1.34e-6	1.13e-4	8.42e-11
ActRIIB(A24N 20-134)-hFc	1.21e-6	6.35e-5	5.19e-11
ActRIIB(L79D 20-134)-hFc	6.7e-5	4.39e-4	6.55e-10
ActRIIB(L79E 20-134)-hFc	3.8e-5	2.74e-4	7.16e-10
ActRIIB(R64K 20-134)-hFc	6.77e-5	2.41e-5	3.56e-11
GDF8			
ActRIIB(20-134)-hFc	3.69e-5	3.45e-5	9.35e-11
ActRIIB(A24N 20-134)-hFc			
ActRIIB(L79D 20-134)-hFc	3.85e-5	8.3e-4	2.15e-9
ActRIIB(L79E 20-134)-hFc	3.74e-5	9e-4	2.41e-9
ActRIIB(R64K 20-134)-hFc	2.25e-5	4.71e-5	2.1e-10
ActRIIB(R64K 20-129)-hFc	9.74e-4	2.09e-4	2.15e-9

47

-continued

Protein	Kon (1/Ms)	Koff (1/s)	KD (M)
ActRIIB(P129S, P130R 20-134)-hFc	1.08e-5	1.8e-4	1.67e-9
ActRIIB(K74A 20-134)-hFc	2.8e-5	2.03e-5	7.18e-11
		ActivinA	
ActRIIB(20-134)-hFc	5.94e6	1.59e-4	2.68e-11
ActRIIB(A24N 20-134)-hFc	3.34e6	3.46e-4	1.04e-10
ActRIIB(L79D 20-134)-hFc			Low binding
ActRIIB(L79E 20-134)-hFc			Low binding
ActRIIB(R64K 20-134)-hFc	6.82e6	3.25e-4	4.76e-11
ActRIIB(R64K 20-129)-hFc	7.46e6	6.28e-4	8.41e-11
ActRIIB(P129S, P130R 20-134)-hFc	5.02e6	4.17e-4	8.31e-11

These data confirm the cell based assay data, demonstrating that the A24N variant retains ligand-binding activity that is similar to that of the ActRIIB(20-134)-hFc molecule, and that the L79D or L79E molecule retains myostatin and GDF11 binding but shows markedly decreased (non-quantifiable) binding to Activin A.

Other variants have been generated and tested, as reported in WO2006/012627 (incorporated herein by reference in its entirety) see e.g., pp. 59-60, using ligands coupled to the device and flowing receptor over the coupled ligands. Notably, K74Y, K74F, K74I (and presumably other hydrophobic substitutions at K74, such as K74L), and D80I, cause a decrease in the ratio of Activin A binding to GDF11 binding, relative to the wild-type K74 molecule. A table of data with respect to these variants is reproduced below:

Soluble ActRIIB-Fc Variants Binding to GDF11 and Activin A (BiaCore Assay)

ActRIIB	ActA	GDF11
WT (64A)	KD = 1.8e-7M	KD = 2.6e-7M
	(+)	(+)
WT (64R)	na	KD = 8.6e-8M
		(+++)
+15tail	KD ~2.6 e-8M	KD = 1.9e-8M
	(+++)	(++++)
E37A	*	*
R40A	-	-
D54A	-	*
K55A	++	*
R56A	*	*
K74A	KD = 4.35e-9 M	KD = 5.3e-9M
	+++++	+++++
K74Y	*	--
K74F	*	--
K74I	*	--
W78A	*	*
L79A	+	*
D80K	*	*
D80R	*	*
D80A	*	*
D80F	*	*
D80G	*	*
D80M	*	*
D80N	*	*
D80I	*	--
F82A	++	-

* No observed binding

-- < 1/5 WT binding

- ~1/2 WT binding

+ WT

++ <2x increased binding

+++ ~5x increased binding

++++ ~10x increased binding

+++++ ~40x increased binding

48

Example 6. ActRIIB-hFc Stimulates Erythropoiesis in Non-Human Primates

ActRIIB(20-134)-hFc (IgG1) was administered once a week for 1 month to male and female cynomolgus monkeys by subcutaneous injection. Forty-eight cynomolgus monkeys (24/sex) were assigned to one of four treatment groups (6 animals/sex/group) and were administered subcutaneous injections of either vehicle or ActRIIB-hFc at 3, 10, or 30 mg/kg once weekly for 4 weeks (total of 5 doses). Parameters evaluated included general clinical pathology (hematology, clinical chemistry, coagulation, and urinalysis). ActRIIB-hFc caused statistically significant elevated mean absolute reticulocyte values by day 15 in treated animals. By day 36, ActRIIB-hFc caused several hematological changes, including elevated mean absolute reticulocyte and red blood cell distribution width values and lower mean corpuscular hemoglobin concentration. All treated groups and both sexes were affected. These effects are consistent with a positive effect of ActRIIB-hFc on the release of immature reticulocytes from the bone marrow. This effect was reversed after drug was washed out of the treated animals (by study day 56). Accordingly, we conclude that ActRIIB-hFc stimulates erythropoiesis.

Example 7. ActRIIB-mFc Promotes Aspects of Erythropoiesis in Mice by Stimulation of Splenic Erythropoietic Activities

In this study the effects of the in vivo administration of ActRIIB(20-134)-mFc on the frequency of hematopoietic progenitors in bone marrow and spleen was analyzed. One group of C57BL/6 mice was injected with PBS as a control and a second group of mice administered two doses of ActRIIB-mFc at 10 mg/kg and both groups sacrificed after 8 days. Peripheral blood was used to perform complete blood counts and femurs and spleens were used to perform in vitro clonogenic assays to assess the lymphoid, erythroid and myeloid progenitor cell content in each organ. In the brief time frame of this study, no significant changes were seen in red blood cell, hemoglobin or white blood cell levels in treated mice. In the femurs there was no difference in the nucleated cell numbers or progenitor content between the control and treated groups. In the spleens, the compound treated group experienced a statistically significant increase in the mature erythroid progenitor (CFU-E) colony number per dish, frequency and total progenitor number per spleen. In addition, and increase was seen in the number of myeloid (CFU-GM), immature erythroid (BFU-E) and total progenitor number per spleen.

Animals:

Sixteen C57BL/6 female mice 6-8 weeks of age were used in the study. Eight mice were injected subcutaneously with test compound ActRIIB-mFc at days 1 and 3 at a dose of 10 mg/kg and eight mice were injected subcutaneously with vehicle control, phosphate buffered saline (PBS), at a volume of 100 μ L per mouse. All mice were sacrificed 8 days after first injection in accordance with the relevant Animal Care Guidelines. Peripheral blood (PB) samples from individual animals were collected by cardiac puncture and used for complete blood counts and differential (CBC/Diff). Femurs and spleens were harvested from each mouse.

Tests Performed:

CBC/Diff Counts

PB from each mouse was collected via cardiac puncture and placed into the appropriate microtainer tubes. Samples were sent to CLV for analysis on a CellDyn 3500 counter.

Clonogenic Assays

Clonogenic progenitors of the myeloid, erythroid and lymphoid lineages were assessed using the in vitro methylcellulose-based media systems described below.

Mature Erythroid Progenitors:

Clonogenic progenitors of the mature erythroid (CFU-E) lineages were cultured in MethoCult™ 3334, a methylcellulose-based medium containing recombinant human (rh) Erythropoietin (3 U/mL).

Lymphoid Progenitors:

Clonogenic progenitors of the lymphoid (CFU-pre-B) lineage were cultured in MethoCult® 3630, a methylcellulose-based medium containing rh Interleukin 7 (10 ng/mL).

Myeloid and Immature Erythroid Progenitors:

Clonogenic progenitors of the granulocyte-monocyte (CFU-GM), erythroid (BFU-E) and multipotential (CFU-GEMM) lineages were cultured in MethoCult™ 3434, a methylcellulose-based medium containing recombinant murine (rm) Stem Cell Factor (50 ng/mL), rh Interleukin 6 (10 ng/mL), rm Interleukin 3 (10 ng/mL) and rh Erythropoietin (3 U/mL).

Methods:

Mouse femurs and spleens were processed by standard protocols. Briefly, bone marrow was obtained by flushing the femoral cavity with Iscove's Modified Dulbecco's Media containing 2% fetal bovine serum (IMDM 2% FBS) using a 21 gauge needle and 1 cc syringe. Spleen cells were obtained by crushing spleens through a 70 µm filter and rinsing the filter with IMDM 2% FBS. Nucleated cell counts in 3% glacial acetic acid were then performed on the single cells suspensions using a Neubauer counting chamber so that the total cells per organ could be calculated. To remove contaminating red blood cells, total spleen cells were then diluted with 3 times the volume of ammonium chloride lysis buffer and incubated on ice 10 minutes. The cells were then washed and resuspended in IMDM 2% FBS and a second cell count were performed to determine the cell concentration of cells after lysis.

Cell stocks were made and added to each methylcellulose-based media formulation to obtain the optimal plating concentrations for each tissue in each media formulation. Bone marrow cells were plated at 1×10^5 cells per dish in MethoCult™ 3334 to assess mature erythroid progenitors, 2×10^5 cells per dish in MethoCult™ 3630 to assess lymphoid progenitors and 3×10^4 cells per dish in MethoCult™ 3434 to assess immature erythroid and myeloid progenitors. Spleen cells were plated at 4×10^5 cells per dish in MethoCult™ 3334 to assess mature erythroid progenitors, 4×10^5 cells per dish in MethoCult™ 3630 to assess lymphoid progenitors and 2×10^5 cells per dish in MethoCult™ 3434 to assess immature erythroid and myeloid progenitors. Cultures plated in triplicate dishes were incubated at 37° C., 5% CO₂ until colony enumeration and evaluation was performed by trained personnel. Mature erythroid progenitors were cul-

tured for 2 days, lymphoid progenitors were cultured for 7 days and mature erythroid and myeloid progenitors were cultured for 12 days.

Analysis:

The mean $\pm$ 1 standard deviation was calculated for the triplicate cultures of the clonogenic assays and for the control and treatment groups for all data sets.

Frequency of colony forming cells (CFC) in each tissue was calculated as follows:

Cells plated per dish

Mean CFC scored per dish

Total CFC per femur or spleen was calculated as follows:

Total CFC scored $\times$ nucleated cell count per femur or spleen (following RBC lysis)

Number of nucleated cells cultured

Standard t-tests were performed to assess if there was a differences in the mean number of cells or hematopoietic progenitors between the PBS control mice and compound treated mice. Due to the potential subjectivity of colony enumeration, a p value of less than 0.01 is deemed significant. Mean values ($\pm$ SD) for each group are shown in the tables below.

TABLE

Treatment Group	Hematologic Parameters			
	White Blood Cells (x10 ⁹ /L)	Red Blood Cells (x10 ⁹ /L)	Hemoglobin (g/L)	Hematocrit (L/L)
PBS (n = 8)	9.53 $\pm$ 1.44	10.5 $\pm$ 1.1	160.9 $\pm$ 13.3	0.552 $\pm$ 0.057
ActRIIB-mFc (n = 8)	9.77 $\pm$ 1.19	10.8 $\pm$ 0.3	162.1 $\pm$ 4.1	0.567 $\pm$ 0.019

TABLE

Treatment Group	CFC From Femur and Spleen			
	Total CFC per Femur	Total CFC per Spleen	Total CFU-E per Femur	Total CFU-E per Spleen
PBS (n = 8)	88 $\pm$ 10	54 $\pm$ 14	156 $\pm$ 27	131 $\pm$ 71
ActRIIB-mFc (n = 8)	85 $\pm$ 9	79 $\pm$ 6*	164 $\pm$ 23	436 $\pm$ 86*

* preliminary analysis indicates p < 0.05

Treatment of mice with ActRIIB(20-134)-mFc, in the brief time frame of this study, did not result in significant increases in red blood cell or hemoglobin content. However, the effect on progenitor cell content was notable. In the femurs there was no difference in the nucleated cell numbers or progenitor content between the control and treated groups. In the spleens, the compound treated group experienced a statistically significant increase in the nucleated cell number before red blood cell lysis and in the mature erythroid progenitor (CFU-E) colony number per dish, frequency and total progenitor number per spleen. In addition, an increase was seen in the number of myeloid (CFU-GM), immature erythroid (BFU-E) and total progenitor number per spleen. Accordingly, it is expected that over a longer time course, ActRIIB(20-134)-mFc treatment may result in elevated red blood cell and hemoglobin content.

Example 8: A GDF Trap Increases Red Blood Cell Levels in Vivo

Twelve-week-old male C57BL/6NTac mice were assigned to one of two treatment groups (N=10). Mice were

51

dosed with either vehicle or with a variant ActRIIB polypeptide ("GDF Trap") [ActRIIB(L79D 20-134)-hFc] by subcutaneous injection (SC) at 10 mg/kg twice per week for 4 weeks. At study termination, whole blood was collected by cardiac puncture into EDTA containing tubes and analyzed for cell distribution using an HM2 hematology analyzer (Abaxis, Inc).

Group Designation

Group	N	Mice	Injection	Dose (mg/kg)	Route	Frequency
1	10	C57BL/6	PBS	0	SC	Twice/week
2	10	C57BL/6	GDF Trap [ActRIIB(L79D 20-134)-hFc]	10	SC	Twice/week

Treatment with the GDF Trap did not have a statistically significant effect on the number of white blood cells (WBC) compared to the vehicle controls. Red blood cell (RBC) numbers were increased in the treated group relative to the controls (see table below). Both the hemoglobin content (HGB) and hematocrit (HCT) were also increased due to the additional red blood cells. The average width of the red blood cells (RDWc) was higher in the treated animals, indicating an increase in the pool of immature red blood cells. Therefore, treatment with the GDF Trap leads to increases in red blood cells, with no distinguishable effects on white blood cell populations.

Hematology Results

	RBC 10 ¹² /L	HGB (g/dL)	HCT (%)	RDWc (%)
PBS	10.7 ± 0.1	14.8 ± 0.6	44.8 ± 0.4	17.0 ± 0.1
GDF Trap	12.4 ± 0.4**	17.0 ± 0.7*	48.8 ± 1.8*	18.4 ± 0.2**

* = p < 0.05,
** = p < 0.01

Example 9: A GDF Trap is Superior to ActRIIB-Fc for Increasing Red Blood Cell Levels In Vivo

Nineteen-week-old male C57BL/6NTac mice were randomly assigned to one of three groups. Mice were dosed with vehicle (10 mM Tris Buffered Saline, TBS), wild-type ActRIIB(20-134)-mFc, or GDF trap ActRIIB(L79D 20-134)-hFc by subcutaneous injection twice per week for three weeks. Blood was collected cheek bleed at baseline and after three weeks of dosing and analyzed for cell distribution using a hematology analyzer (HM2, Abaxis, Inc.)

Treatment with ActRIIB-Fc or the GDF trap did not have a significant effect on white blood cell (WBC) numbers compared to vehicle controls. The red blood cell count (RBC), hematocrit (HCT), and hemoglobin levels were all elevated in GDF Trap treated mice compared to either the controls or the wild-type construct (see table below). Therefore, in a direct comparison, the GDF trap promotes increases in red blood cells to a significantly greater extent than a wild-type ActRIIB-Fc protein. In fact, in this experiment, the wild-type ActRIIB-Fc protein did not cause a statistically significant increase in red blood cells, suggesting that longer or higher dosing would be needed to reveal this effect.

52

Hematology Results after Three Weeks of Dosing

	RBC (10 ¹² /ml)	HCT %	HGB g/dL
TBS	11.06 ± 0.46	46.78 ± 1.9	15.7 ± 0.7
ActRIIB-mFc	11.64 ± 0.09	49.03 ± 0.3	16.5 ± 1.5
GDF Trap	13.19 ± 0.2**	53.04 ± 0.8**	18.4 ± 0.3**

** = p < 0.01

Example 10. Generation of a GDF Trap with Truncated ActRIIB Extracellular Domain

As described in Example 1, a GDF Trap referred to as ActRIIB(L79D 20-134)-hFc was generated by N-terminal fusion of TPA leader to the ActRIIB extracellular domain (residues 20-134 in SEQ ID NO: 1) containing a leucine-to-aspartate substitution (at residue 79 in SEQ ID NO: 1) and C-terminal fusion of human Fc domain with minimal linker (three glycine residues) (FIG. 3). A nucleotide sequence corresponding to this fusion protein is shown in FIGS. 4A and 4B.

A GDF Trap with truncated ActRIIB extracellular domain, referred to as ActRIIB(L79D 25-131)-hFc, was generated by N-terminal fusion of TPA leader to truncated extracellular domain (residues 25-131 in SEQ ID NO: 1) containing a leucine-to-aspartate substitution (at residue 79 in SEQ ID NO: 1) and C-terminal fusion of human Fc domain with minimal linker (three glycine residues) (FIG. 5). A nucleotide sequence corresponding to this fusion protein is shown in FIGS. 6A and 6B.

Example 11. Selective Ligand Binding by GDF Trap with Double-Truncated ActRIIB Extracellular Domain

The affinity of GDF Traps and other ActRIIB-hFc proteins for several ligands was evaluated in vitro with a Biacore™ instrument. Results are summarized in the table below. Kd values were obtained by steady-state affinity fit due to the very rapid association and dissociation of the complex, which prevented accurate determination of k_{on} and k_{off}.
Ligand Selectivity of ActRIIB-hFc Variants:

Fusion Construct	Activin A (Kd e-11)	Activin B (Kd e-11)	GDF11 (Kd e-11)
ActRIIB(L79 20-134)-hFc	1.6	1.2	3.6
ActRIIB(L79D 20-134)-hFc	1350.0	78.8	12.3
ActRIIB(L79 25-131)-hFc	1.8	1.2	3.1
ActRIIB(L79D 25-131)-hFc	2290.0	62.1	7.4

The GDF Trap with a truncated extracellular domain, ActRIIB(L79D 25-131)-hFc, equaled or surpassed the ligand selectivity displayed by the longer variant, ActRIIB (L79D 20-134)-hFc, with pronounced loss of activin A and activin B binding and nearly full retention of GDF11 binding compared to ActRIIB-hFc counterparts lacking the L79D substitution. Note that truncation alone (without L79D substitution) did not alter selectivity among the ligands displayed here [compare ActRIIB(L79 25-131)-hFc with ActRIIB(L79 20-134)-hFc].

Example 12. Generation of ActRIIB(L79D 25-131)-hFc with Alternative Nucleotide Sequences

To generate ActRIIB(L79D 25-131)-hFc, the human ActRIIB extracellular domain with an aspartate substitution

at native position 79 (SEQ ID NO: 1) and with N-terminal and C-terminal truncations (residues 25-131 in SEQ ID NO: 1) was fused N-terminally with a TPA leader sequence instead of the native ActRIIB leader and C-terminally with a human Fc domain via a minimal linker (three glycine residues) (FIG. 5). One nucleotide sequence encoding this fusion protein is shown in FIGS. 6A and 6B (SEQ ID NO: 27), and an alternative nucleotide sequence encoding exactly the same fusion protein is shown in FIGS. 9A and 9B (SEQ ID NO: 30). This protein was expressed and purified using the methodology described in Example 1.

Example 13. GDF Trap with a Truncated ActRIIB Extracellular Domain Increases Proliferation of Erythroid Progenitors in Mice

ActRIIB(L79D 25-131)-hFc was evaluated to determine its effect on proliferation of erythroid progenitors. Male C57BL/6 mice (8 weeks old) were treated with ActRIIB (L79D 25-131)-hFc (10 mg/kg, s.c.; n=6) or vehicle (TBS; n=6) on Days 1 and 4, then euthanized on Day 8 for collection of spleens, tibias, femurs, and blood. Cells of the spleen and bone marrow were isolated, diluted in Iscove's modified Dulbecco's medium containing 5% fetal bovine serum, suspended in specialized methylcellulose-based medium, and cultured for either 2 or 12 days to assess levels of clonogenic progenitors at the colony-forming unit-erythroid (CFU-E) and burst forming unit-erythroid (BFU-E) stages, respectively. Methylcellulose-based medium for BFU-E determination (MethoCult M3434, Stem Cell Technologies) included recombinant murine stem cell factor, interleukin-3, and interleukin-6, which were not present in methylcellulose medium for CFU-E determination (MethoCult M3334, Stem Cell Technologies), while both media contained erythropoietin, among other constituents. For both BFU-E and CFU-E, the number of colonies were determined in duplicate culture plates derived from each tissue sample, and statistical analysis of the results was based on the number of mice per treatment group.

Spleen-derived cultures from mice treated with ActRIIB (L79D 25-131)-hFc had twice the number of CFU-E colonies as did corresponding cultures from control mice ($P<0.05$), whereas the number of BFU-E colonies did not differ significantly with treatment in vivo. The number of CFU-E or BFU-E colonies from bone marrow cultures also did not differ significantly with treatment. As expected, increased numbers of CFU-E colonies in spleen-derived cultures were accompanied by highly significant ($P<0.001$) changes in red blood cell level (11.6% increase), hemoglobin concentration (12% increase), and hematocrit level (11.6% increase) at euthanasia in mice treated with ActRIIB (L79D 25-131)-hFc compared to controls. These results indicate that in vivo administration of a GDF Trap with truncated ActRIIB extracellular domain can stimulate proliferation of erythroid progenitors as part of its overall effect to increase red blood cell levels.

Example 14. GDF Trap with a Truncated ActRIIB Extracellular Domain Offsets Chemotherapy-Induced Anemia in Mice

Applicants investigated the effect of ActRIIB(L79D 25-131)-hFc on erythropoietic parameters in a mouse model of chemotherapy-induced anemia based on paclitaxel, which inhibits cell division by blocking microtubule polymeriza-

- 1) paclitaxel (25 mg/kg, i.p.)
- 2) ActRIIB(L79D 25-131)-hFc (10 mg/kg, i.p.)
- 3) paclitaxel+ActRIIB(L79D 25-131)-hFc
- 4) vehicle (TBS).

Paclitaxel was administered on Day 0, while ActRIIB(L79D 25-131)-hFc or vehicle were administered on Days 0 and 3. Blood samples were collected for CBC analysis from separate cohorts on Days 1, 3, and 5, and results for treatment groups 1-3 (above) were expressed as percent difference from vehicle at a given time point. Attrition due to paclitaxel toxicity was an issue in the paclitaxel-only cohort on Day 3 (where n=1); otherwise, n=3-5 per treatment per time point. Compared to vehicle, paclitaxel alone decreased hemoglobin concentration by nearly 13% at Day 5, whereas addition of ActRIIB(L79D 25-131)-hFc prevented this paclitaxel-induced decline (FIG. 11). Similar effects were observed for hematocrit and RBC levels. In the absence of paclitaxel, ActRIIB(L79D 25-131)-hFc increased hemoglobin concentration by 10% compared to vehicle on Days 3 and 5 (FIG. 11). Thus, a GDF Trap with truncated ActRIIB extracellular domain can increase levels of red blood cells sufficiently to offset chemotherapy-induced anemia.

Example 15. GDF Trap with a Truncated ActRIIB Extracellular Domain Reverses Nephrectomy-Induced Anemia in Mice

Applicants investigated the effect of ActRIIB(L79D 25-131)-hFc on anemia in a nephrectomized mouse model of chronic kidney disease. Male C57BL/6 mice (11 weeks old) underwent either a sham operation or a unilateral nephrectomy to reduce the capacity for erythropoietin production. Mice were allowed a week for postsurgical recovery and then treated twice-weekly with ActRIIB(L79D 25-131)-hFc (10 mg/kg, i.p.; n=15 per condition) or vehicle (TBS; n=15 per condition) for a total of 4 weeks. Blood samples were collected before the onset of dosing and after 4 weeks of treatment. Whereas vehicle-treated nephrectomized mice displayed a significant decline in red blood cell number over the 4-week treatment period, treatment with ActRIIB(L79D 25-131)-hFc not only prevented the decline but increased red blood cell levels 17% ($P<0.001$) above baseline (FIG. 12), despite reduced renal capacity for erythropoietin production. In nephrectomized mice, ActRIIB (L79D 25-131)-hFc also generated significant increases from baseline in hemoglobin concentration and hematocrit level and, notably, stimulated each of these erythropoietic parameters to approximately the same extent under nephrectomized conditions as under sham-operated conditions (FIG. 13). Thus, a GDF Trap with truncated ActRIIB extracellular domain can increase red blood cell levels sufficiently to reverse anemia in a model of chronic kidney disease.

Example 16. GDF Trap with a Truncated ActRIIB Extracellular Domain Improves Recovery from Anemia Induced by Blood Loss in Rats

Applicants investigated the effect of ActRIIB(L79D 25-131)-hFc on erythropoietic parameters in a rat model of anemia induced by acute blood loss (acute post-hemorrhagic anemia). Male Sprague-Dawley rats (approximately 300 g) received a chronic jugular catheter at the vendor (Harlan). On Day -1, 20% of total blood volume was withdrawn from each rat over a 5-minute period via the catheter under isoflurane anesthesia. The volume of blood removed was based on a value for total blood volume calculated according

55

to the following relationship derived by Lee and co-workers (J Nucl Med 25:72-76, 1985) for rats with body weight greater than 120 g:

$$\text{Total blood volume (ml)} = 0.062 \times \text{body weight (g)} + 0.0012$$

An equal volume of phosphate-buffered saline was replaced via the catheter at the time of blood removal. Rats were treated with ActRIIB(L79D 25-131)-hFc (10 mg/kg, s.c.; n=5) or vehicle (TBS; n=5) on Days 0 and 3. Blood samples for CBC analysis were removed via the catheter on Days -1 (baseline), 0, 2, 4, and 6.

Control rats responded to 20% blood loss with a drop of nearly 15% in red-blood-cell levels by Day 0. These levels remained significantly lower than baseline on Days 2 and 4, and had not recovered fully by Day 6 (FIG. 14). Although rats treated with ActRIIB(L79D 25-131)-hFc showed a nearly identical drop in red-blood-cell levels after 20% blood loss, these rats then displayed a complete recovery in such levels by Day 2, followed by further elevation on Days 4 and 6, which represents a highly significant improvement over control levels at the corresponding time points (FIG. 14). Similar results were obtained for hemoglobin concentration. These findings demonstrate that a GDF Trap with truncated ActRIIB extracellular domain can produce a faster recovery of red blood cell levels from anemia caused by acute hemorrhage.

Example 17. GDF Trap with Truncated ActRIIB Extracellular Domain Increases Levels of Red Blood Cells in Non-Human Primates

Two GDF Traps, ActRIIB(L79D 20-134)-hFc and ActRIIB(L79D 25-131)-hFc, were evaluated for their ability to stimulate red blood cell production in cynomolgus monkey. Monkeys were treated subcutaneously with GDF Trap (10 mg/kg; n=4 males/4 females), or vehicle (n=2 males/2 females) on Days 1 and 8. Blood samples were collected on Days 1 (pretreatment baseline), 3, 8, 15, 29, and 44, and were analyzed for red blood cell levels (FIG. 15), hematocrit (FIG. 16), hemoglobin levels (FIG. 17), and reticulocyte levels (FIG. 18). Vehicle-treated monkeys exhibited decreased levels of red blood cells, hematocrit, and hemoglobin at all post-treatment time points, an expected effect of repeated blood sampling. In contrast, treatment with ActRIIB(L79D 20-134)-hFc or ActRIIB(L79D 25-131)-hFc increased these parameters by the first post-treatment time point (Day 3) and maintained them at substantially elevated levels for the duration of the study (FIGS. 15-17). Importantly, reticulocyte levels in monkeys treated with ActRIIB (L79D 20-134)-hFc or ActRIIB(L79D 25-131)-hFc were substantially increased at Days 8, 15, and 29 compared to vehicle (FIG. 18). This result demonstrates that GDF Trap treatment increased production of red blood cell precursors, resulting in elevated red blood cell levels.

Taken together, these data demonstrate that truncated GDF Traps, as well as a full-length variants, can be used as selective antagonists of GDF11 and potentially related ligands to increase red blood cell formation in vivo.

Example 18. GDF Trap Derived from ActRIIB5

Others have reported an alternate, soluble form of ActRIIB (designated ActRIIB5), in which exon 4, including

56

the ActRIIB transmembrane domain, has been replaced by a different C-terminal sequence (WO2007/053775).

The sequence of native human ActRIIB5 without its leader is as follows:

(SEQ ID NO: 36)
GRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWRNSSG
TIELVKKGCVLDDDFNCYDRQECVATEENPQVYFCCCEGNFCNERFTHLP
EAGGPEGPWASTTIPSGGPEATAAAGDQGSGALWLCLEGPAAHE

An leucine-to-aspartate substitution, or other acidic substitutions, may be performed at native position 79 (underlined and highlighted) as described to construct the variant ActRIIB5(L79D), which has the following sequence:

(SEQ ID NO: 37)
GRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWRNSSG
TIELVKKGCVLDDDFNCYDRQECVATEENPQVYFCCCEGNFCNERFTHLP
EAGGPEGPWASTTIPSGGPEATAAAGDQGSGALWLCLEGPAAHE

This variant may be connected to human Fc with a TGGG (SEQ ID NO: 42) linker to generate a human ActRIIB5 (L79D)-hFc fusion protein with the following sequence:

(SEQ ID NO: 38)
GRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWRNSSG
TIELVKKGCVLDDDFNCYDRQECVATEENPQVYFCCCEGNFCNERFTHLP
EAGGPEGPWASTTIPSGGPEATAAAGDQGSGALWLCLEGPAAHETGGGTH
TCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEV
KFNWYVDGVEVHNAKTKPREEQYNSTYRVSVLTVHLQDNLNGKEYKCK
VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKG
FYPYSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQG
NVFSCSVMEALHNHYTQKSLSLSPGK

This construct may be expressed in CHO cells.

INCORPORATION BY REFERENCE

All publications and patents mentioned herein are hereby incorporated by reference in their entirety as if each individual publication or patent was specifically and individually indicated to be incorporated by reference.

While specific embodiments of the subject matter have been discussed, the above specification is illustrative and not restrictive. Many variations will become apparent to those skilled in the art upon review of this specification and the claims below. The full scope of the invention should be determined by reference to the claims, along with their full scope of equivalents, and the specification, along with such variations.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 44

<210> SEQ ID NO 1
<211> LENGTH: 512
<212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

Met Thr Ala Pro Trp Val Ala Leu Ala Leu Leu Trp Gly Ser Leu Trp
 1           5           10           15

Pro Gly Ser Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr
 20           25           30

Asn Ala Asn Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg
 35           40           45

Cys Glu Gly Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Arg
 50           55           60

Asn Ser Ser Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp
 65           70           75           80

Asp Phe Asn Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn
 85           90           95

Pro Gln Val Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg
100           105           110

Phe Thr His Leu Pro Glu Ala Gly Gly Pro Glu Val Thr Tyr Glu Pro
115           120           125

Pro Pro Thr Ala Pro Thr Leu Leu Thr Val Leu Ala Tyr Ser Leu Leu
130           135           140

Pro Ile Gly Gly Leu Ser Leu Ile Val Leu Leu Ala Phe Trp Met Tyr
145           150           155           160

Arg His Arg Lys Pro Pro Tyr Gly His Val Asp Ile His Glu Asp Pro
165           170           175

Gly Pro Pro Pro Pro Ser Pro Leu Val Gly Leu Lys Pro Leu Gln Leu
180           185           190

Leu Glu Ile Lys Ala Arg Gly Arg Phe Gly Cys Val Trp Lys Ala Gln
195           200           205

Leu Met Asn Asp Phe Val Ala Val Lys Ile Phe Pro Leu Gln Asp Lys
210           215           220

Gln Ser Trp Gln Ser Glu Arg Glu Ile Phe Ser Thr Pro Gly Met Lys
225           230           235           240

His Glu Asn Leu Leu Gln Phe Ile Ala Ala Glu Lys Arg Gly Ser Asn
245           250           255

Leu Glu Val Glu Leu Trp Leu Ile Thr Ala Phe His Asp Lys Gly Ser
260           265           270

Leu Thr Asp Tyr Leu Lys Gly Asn Ile Ile Thr Trp Asn Glu Leu Cys
275           280           285

His Val Ala Glu Thr Met Ser Arg Gly Leu Ser Tyr Leu His Glu Asp
290           295           300

Val Pro Trp Cys Arg Gly Glu Gly His Lys Pro Ser Ile Ala His Arg
305           310           315           320

Asp Phe Lys Ser Lys Asn Val Leu Leu Lys Ser Asp Leu Thr Ala Val
325           330           335

Leu Ala Asp Phe Gly Leu Ala Val Arg Phe Glu Pro Gly Lys Pro Pro
340           345           350

Gly Asp Thr His Gly Gln Val Gly Thr Arg Arg Tyr Met Ala Pro Glu
355           360           365

Val Leu Glu Gly Ala Ile Asn Phe Gln Arg Asp Ala Phe Leu Arg Ile
370           375           380

Asp Met Tyr Ala Met Gly Leu Val Leu Trp Glu Leu Val Ser Arg Cys
385           390           395           400

```

-continued

Lys Ala Ala Asp Gly Pro Val Asp Glu Tyr Met Leu Pro Phe Glu Glu
 405 410 415
 Glu Ile Gly Gln His Pro Ser Leu Glu Glu Leu Gln Glu Val Val Val
 420 425 430
 His Lys Lys Met Arg Pro Thr Ile Lys Asp His Trp Leu Lys His Pro
 435 440 445
 Gly Leu Ala Gln Leu Cys Val Thr Ile Glu Glu Cys Trp Asp His Asp
 450 455 460
 Ala Glu Ala Arg Leu Ser Ala Gly Cys Val Glu Glu Arg Val Ser Leu
 465 470 475 480
 Ile Arg Arg Ser Val Asn Gly Thr Thr Ser Asp Cys Leu Val Ser Leu
 485 490 495
 Val Thr Ser Val Thr Asn Val Asp Leu Pro Pro Lys Glu Ser Ser Ile
 500 505 510

<210> SEQ ID NO 2
 <211> LENGTH: 115
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala Asn
 1 5 10 15
 Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly
 20 25 30
 Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Arg Asn Ser Ser
 35 40 45
 Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp Asp Phe Asn
 50 55 60
 Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val
 65 70 75 80
 Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His
 85 90 95
 Leu Pro Glu Ala Gly Gly Pro Glu Val Thr Tyr Glu Pro Pro Pro Thr
 100 105 110
 Ala Pro Thr
 115

<210> SEQ ID NO 3
 <211> LENGTH: 100
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala Asn
 1 5 10 15
 Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly
 20 25 30
 Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Arg Asn Ser Ser
 35 40 45
 Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp Asp Phe Asn
 50 55 60
 Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val
 65 70 75 80
 Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His
 85 90 95

-continued

Leu Pro Glu Ala
100

<210> SEQ ID NO 4
<211> LENGTH: 1539
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

```

atgacggcgc cctgggtggc cctcgccctc ctctggggat cgctgtggcc cggctctggg    60
cgtggggagg ctgagacacg ggagtgcac tactacaacg ccaactggga gctggagcgc    120
accaaccaga gcggcctgga gcgctgcgaa ggcgagcagg acaagcggct gcactgctac    180
gcctcctggg ccaacagctc tggcaccatc gagctcgtga agaagggctg ctggctagat    240
gacttcaact gctacgatag gcaggagtgt gtggccactg aggagaacct ccagggttac    300
ttctgctgct gtgaaggcaa cttctgcaac gagcgttca ctcatTTGCC agaggctggg    360
ggcccggaaG tcacttacga gccacccccg acagccccca ccctgctcac ggtgctggcc    420
tactcactgc tgcccatcgg gggcctttcc ctcatcgtcc tgetggcctt ttggatgtac    480
cggcatcgca agcccccta cggtcatgtg gacatccatg aggaccctgg gcctccacca    540
ccatccccctc tggTgggcct gaagccactg cagctgctgg agatcaaggc tcgggggcgc    600
tttgctgtgT tctggaaggc ccagctcatg aatgactttg tagctgtcaa gatcttccca    660
ctccaggaca agcagtcgtg gcagagtgaA cgggagatct tcagcacacc tggcatgaag    720
cacgagaacc tgctacagtt cattgtctgc gagaagcgag gctccaacct cgaagtagag    780
ctgtggctca tcacggcctt ccatgacaag ggctccctca cggattacct caaggggaac    840
atcatcacat ggaacgaact gtgtcatgta gcagagacga tgtcacgagg cctctcatac    900
ctgcatgagg atgtgccctg gtgccgtggc gagggccaca agccgtctat tgccacagg    960
gactttaaaa gtaagaatgt attgctgaag agcgacctca cagccgtgct ggctgacttt 1020
ggcttggtgT ttcgatttga gccagggaAa cctccagggg acaccacagg acaggtaggc 1080
acgagacggT acatggctcc tgaggTgctc gagggagcca tcaacttcca gagagatgcc 1140
ttcctgcgca ttgacatgta tgccatgggg ttggtgctgt gggagcttgt gtctcgtgc 1200
aaggctgcag acggaccctg ggatgagtac atgtgccct ttgaggaaga gattggccag 1260
cacccttctg tggaggagct gcaggaggTg gtggtgcaca agaagatgag gccaccatt 1320
aaagatcact ggttgaaaca cccgggcctg gccagcttt gtgtgaccat cgaggagtgc 1380
tgggaccatg atgcagaggc tcgcttgtcc gcgggctgtg tggaggagcg ggtgtccctg 1440
attcggaggt cggTcaacgg cactacctcg gactgtctcg tttccctggT gacctctgtc 1500
accaatgtgg acctgcccc taaagagtca agcatctaa    1539

```

<210> SEQ ID NO 5
<211> LENGTH: 345
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

```

gggcgtgggg aggctgagac acgggagtgc atctactaca acgccaactg ggagctggag    60
cgcaccaacc agagcggcct ggagcgtgc gaaggcgagc aggacaagcg gctgcactgc    120
tacgcctcct gggccaacag ctctggcacc atcgagctcg tgaagaaggg ctgctggcta    180
gatgacttca actgctacga taggcaggag tgtgtggcca ctgaggagaa cccccaggTg    240

```

-continued

```
tactttctgct gctgtgaagg caacttctgc aacgagcgct tcaactcattt gccagaggct 300
ggggggcccg aagtcacgta cgagccaccc ccgacagccc ccacc 345
```

```
<210> SEQ ID NO 6
<211> LENGTH: 225
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polypeptide
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (43)..(43)
<223> OTHER INFORMATION: Asp or Ala
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (100)..(100)
<223> OTHER INFORMATION: Lys or Ala
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (212)..(212)
<223> OTHER INFORMATION: Asn or Ala
```

```
<400> SEQUENCE: 6
```

```
Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro
1          5          10          15
Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
20        25        30
Arg Thr Pro Glu Val Thr Cys Val Val Val Xaa Val Ser His Glu Asp
35        40        45
Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn
50        55        60
Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val
65        70        75        80
Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu
85        90        95
Tyr Lys Cys Xaa Val Ser Asn Lys Ala Leu Pro Val Pro Ile Glu Lys
100       105       110
Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr
115       120       125
Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr
130       135       140
Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu
145       150       155       160
Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu
165       170       175
Asp Ser Asp Gly Pro Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys
180       185       190
Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
195       200       205
Ala Leu His Xaa His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
210       215       220
Lys
225
```

```
<210> SEQ ID NO 7
<211> LENGTH: 343
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
```

-continued

polypeptide

<400> SEQUENCE: 7

Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala Asn
 1 5 10 15
 Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly
 20 25 30
 Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Arg Asn Ser Ser
 35 40 45
 Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Asp Asp Asp Phe Asn
 50 55 60
 Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val
 65 70 75 80
 Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His
 85 90 95
 Leu Pro Glu Ala Gly Gly Pro Glu Val Thr Tyr Glu Pro Pro Pro Thr
 100 105 110
 Ala Pro Thr Gly Gly Gly Thr His Thr Cys Pro Pro Cys Pro Ala Pro
 115 120 125
 Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
 130 135 140
 Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
 145 150 155 160
 Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp
 165 170 175
 Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr
 180 185 190
 Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
 195 200 205
 Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu
 210 215 220
 Pro Val Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
 225 230 235 240
 Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys
 245 250 255
 Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
 260 265 270
 Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
 275 280 285
 Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
 290 295 300
 Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser
 305 310 315 320
 Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
 325 330 335
 Leu Ser Leu Ser Pro Gly Lys
 340

<210> SEQ ID NO 8

<211> LENGTH: 21

<212> TYPE: PRT

<213> ORGANISM: *Apis mellifera*

<400> SEQUENCE: 8

Met Lys Phe Leu Val Asn Val Ala Leu Val Phe Met Val Val Tyr Ile

-continued

1 5 10 15

Ser Tyr Ile Tyr Ala
20

<210> SEQ ID NO 9
 <211> LENGTH: 22
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown: Tissue Plasminogen
 Activator

<400> SEQUENCE: 9

Met Asp Ala Met Lys Arg Gly Leu Cys Cys Val Leu Leu Leu Cys Gly
 1 5 10 15

Ala Val Phe Val Ser Pro
20

<210> SEQ ID NO 10
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown: Native peptide

<400> SEQUENCE: 10

Met Thr Ala Pro Trp Val Ala Leu Ala Leu Leu Trp Gly Ser Leu Cys
 1 5 10 15

Ala Gly Ser

<210> SEQ ID NO 11
 <211> LENGTH: 368
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polypeptide

<400> SEQUENCE: 11

Met Asp Ala Met Lys Arg Gly Leu Cys Cys Val Leu Leu Leu Cys Gly
 1 5 10 15

Ala Val Phe Val Ser Pro Gly Ala Ser Gly Arg Gly Glu Ala Glu Thr
 20 25 30

Arg Glu Cys Ile Tyr Tyr Asn Ala Asn Trp Glu Leu Glu Arg Thr Asn
 35 40 45

Gln Ser Gly Leu Glu Arg Cys Glu Gly Glu Gln Asp Lys Arg Leu His
 50 55 60

Cys Tyr Ala Ser Trp Arg Asn Ser Ser Gly Thr Ile Glu Leu Val Lys
 65 70 75 80

Lys Gly Cys Trp Asp Asp Asp Phe Asn Cys Tyr Asp Arg Gln Glu Cys
 85 90 95

Val Ala Thr Glu Glu Asn Pro Gln Val Tyr Phe Cys Cys Cys Glu Gly
 100 105 110

Asn Phe Cys Asn Glu Arg Phe Thr His Leu Pro Glu Ala Gly Gly Pro
 115 120 125

Glu Val Thr Tyr Glu Pro Pro Pro Thr Ala Pro Thr Gly Gly Gly Thr
 130 135 140

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser
 145 150 155 160

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg

-continued

165								170					175				
Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro		
			180				185						190				
Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala		
			195				200						205				
Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val		
			210				215						220				
Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr		
			225				230						240				
Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Val	Pro	Ile	Glu	Lys	Thr		
			245				250						255				
Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu		
			260				265						270				
Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys		
			275				280						285				
Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser		
			290				295						300				
Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp		
			305				310						315				
Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser		
			325				330						335				
Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala		
			340				345						350				
Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys		
			355				360						365				

<210> SEQ ID NO 12

<211> LENGTH: 1107

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 12

```

atggatgcaa tgaagagagg gctctgctgt gtgctgctgc tgtgtggagc agtcttcggt      60
tcgcccggcg cctctggggc tggggaggct gagacacggg agtgcattca ctacaacgcc      120
aactgggagc tggagcgcac caaccagagc ggcctggagc gctgcgaagg cgagcaggac      180
aagcggctgc actgctacgc ctctctggcg aacagctctg gcaccatcga gctcgtgaag      240
aagggtctgt gggacgatga cttcaactgc tacgataggc aggagtgtgt ggccactgag      300
gagaaccccc aggtgtactt ctgctgctgt gaaggcaact tctgcaacga gcgcttcact      360
catttgccag aggtgtgggg ccgggaagtc acgtacgagc ccccccgac agccccacc      420
ggtggtggaa ctcacacatg cccaccgtgc ccagcacctg aactcctggg gggaccgtca      480
gtcttcctct tcccccaaaa acccaaggac accctcatga tctcccgga cctgaggtc      540
acatgcgtgg tgggtgaagt gagccacgaa gacctgagg tcaagttcaa ctggtacgtg      600
gacggcgtgg aggtgcataa tgccaagaca aagccgcggg aggagcagta caacagcacg      660
taccgtgtgg tcagcgtctc caccgtctg caccaggact ggctgaatgg caaggagtac      720
aagtgcgaag tctccaacaa agccctccca gtccccatcg agaaaacat ctccaaagcc      780
aaagggcagc ccgagaacc acaggtgtac accctgcccc catcccgga ggagatgacc      840
aagaaccagg tcagcctgac ctgcctggtc aaagcttct atcccagca catcgccgtg      900

```

-continued

```

gagtgggaga gcaatgggca gccggagAAC aactacaaga ccacgcctcc cgtgctggac    960
tccgacggct ccttcttctt ctatagcaag ctacacgtgg acaagagcag gtggcagcag    1020
gggaacgtct tctcatgtct cgtgatgcat gaggtctctgc acaaccacta cacgcagaag    1080
agcctctccc tgtctccggg taaatga                                     1107

```

```

<210> SEQ ID NO 13
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        peptide

```

```

<400> SEQUENCE: 13

```

```

Thr Gly Gly Gly Gly
1           5

```

```

<210> SEQ ID NO 14
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        peptide

```

```

<400> SEQUENCE: 14

```

```

Ser Gly Gly Gly Gly
1           5

```

```

<210> SEQ ID NO 15
<211> LENGTH: 116
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 15

```

```

Ile Leu Gly Arg Ser Glu Thr Gln Glu Cys Leu Phe Phe Asn Ala Asn
1           5           10           15
Trp Glu Lys Asp Arg Thr Asn Gln Thr Gly Val Glu Pro Cys Tyr Gly
20          25          30
Asp Lys Asp Lys Arg Arg His Cys Phe Ala Thr Trp Lys Asn Ile Ser
35          40          45
Gly Ser Ile Glu Ile Val Lys Gln Gly Cys Trp Leu Asp Asp Ile Asn
50          55          60
Cys Tyr Asp Arg Thr Asp Cys Val Glu Lys Lys Asp Ser Pro Glu Val
65          70          75          80
Tyr Phe Cys Cys Cys Glu Gly Asn Met Cys Asn Glu Lys Phe Ser Tyr
85          90          95
Phe Pro Glu Met Glu Val Thr Gln Pro Thr Ser Asn Pro Val Thr Pro
100         105         110
Lys Pro Pro Thr
115

```

```

<210> SEQ ID NO 16
<211> LENGTH: 150
<212> TYPE: PRT
<213> ORGANISM: Rattus sp.

```

```

<400> SEQUENCE: 16

```

```

Met Thr Ala Pro Trp Ala Ala Leu Ala Leu Leu Trp Gly Ser Leu Cys
1           5           10           15

```

-continued

Ala	Gly	Ser	Gly	Arg	Gly	Glu	Ala	Glu	Thr	Arg	Glu	Cys	Ile	Tyr	Tyr
			20					25					30		
Asn	Ala	Asn	Trp	Glu	Leu	Glu	Arg	Thr	Asn	Gln	Ser	Gly	Leu	Glu	Arg
		35					40					45			
Cys	Glu	Gly	Glu	Gln	Asp	Lys	Arg	Leu	His	Cys	Tyr	Ala	Ser	Trp	Pro
	50					55					60				
Asn	Ser	Ser	Gly	Thr	Ile	Glu	Leu	Val	Lys	Lys	Gly	Cys	Trp	Leu	Asp
65					70					75					80
Asp	Phe	Asn	Cys	Tyr	Asp	Arg	Gln	Glu	Cys	Val	Ala	Thr	Glu	Glu	Asn
			85					90						95	
Pro	Gln	Val	Tyr	Phe	Cys	Cys	Cys	Glu	Gly	Asn	Phe	Cys	Asn	Glu	Arg
			100					105					110		
Phe	Thr	His	Leu	Pro	Glu	Pro	Gly	Gly	Pro	Glu	Val	Thr	Tyr	Glu	Pro
		115					120					125			
Pro	Pro	Thr	Ala	Pro	Thr	Leu	Leu	Thr	Val	Leu	Ala	Tyr	Ser	Leu	Leu
	130					135					140				
Pro	Ile	Gly	Gly	Leu	Ser										
145					150										

<210> SEQ ID NO 17
 <211> LENGTH: 150
 <212> TYPE: PRT
 <213> ORGANISM: Sus sp.

<400> SEQUENCE: 17

Met	Thr	Ala	Pro	Trp	Ala	Ala	Leu	Ala	Leu	Leu	Trp	Gly	Ser	Leu	Cys
1				5					10					15	
Val	Gly	Ser	Gly	Arg	Gly	Glu	Ala	Glu	Thr	Arg	Glu	Cys	Ile	Tyr	Tyr
			20					25					30		
Asn	Ala	Asn	Trp	Glu	Leu	Glu	Arg	Thr	Asn	Gln	Ser	Gly	Leu	Glu	Arg
		35					40					45			
Cys	Glu	Gly	Glu	Gln	Asp	Lys	Arg	Leu	His	Cys	Tyr	Ala	Ser	Trp	Arg
	50					55					60				
Asn	Ser	Ser	Gly	Thr	Ile	Glu	Leu	Val	Lys	Lys	Gly	Cys	Trp	Leu	Asp
65					70					75					80
Asp	Phe	Asn	Cys	Tyr	Asp	Arg	Gln	Glu	Cys	Val	Ala	Thr	Glu	Glu	Asn
			85					90						95	
Pro	Gln	Val	Tyr	Phe	Cys	Cys	Cys	Glu	Gly	Asn	Phe	Cys	Asn	Glu	Arg
			100					105					110		
Phe	Thr	His	Leu	Pro	Glu	Ala	Gly	Gly	Pro	Glu	Val	Thr	Tyr	Glu	Pro
		115					120					125			
Pro	Pro	Thr	Ala	Pro	Thr	Leu	Leu	Thr	Val	Leu	Ala	Tyr	Ser	Leu	Leu
	130					135					140				
Pro	Ile	Gly	Gly	Leu	Ser										
145					150										

<210> SEQ ID NO 18
 <211> LENGTH: 150
 <212> TYPE: PRT
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 18

Met	Thr	Ala	Pro	Trp	Ala	Ala	Leu	Ala	Leu	Leu	Trp	Gly	Ser	Leu	Cys
1				5					10					15	
Ala	Gly	Ser	Gly	Arg	Gly	Glu	Ala	Glu	Thr	Arg	Glu	Cys	Ile	Tyr	Tyr
			20					25					30		

-continued

```

Asn Ala Asn Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg
   35                               40               45

Cys Glu Gly Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Arg
   50                               55               60

Asn Ser Ser Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp
   65                               70               75               80

Asp Phe Asn Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn
   85                               90               95

Pro Gln Val Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg
  100                               105              110

Phe Thr His Leu Pro Glu Pro Gly Gly Pro Glu Val Thr Tyr Glu Pro
  115                               120              125

Pro Pro Thr Ala Pro Thr Leu Leu Thr Val Leu Ala Tyr Ser Leu Leu
  130                               135              140

Pro Ile Gly Gly Leu Ser
  145                               150

```

```

<210> SEQ ID NO 19
<211> LENGTH: 150
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 19

```

```

Met Thr Ala Pro Trp Val Ala Leu Ala Leu Leu Trp Gly Ser Leu Cys
  1      5              10              15

Ala Gly Ser Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr
  20      25              30

Asn Ala Asn Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg
  35      40              45

Cys Glu Gly Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Arg
  50      55              60

Asn Ser Ser Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp
  65      70              75               80

Asp Phe Asn Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn
  85      90              95

Pro Gln Val Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg
 100      105             110

Phe Thr His Leu Pro Glu Ala Gly Gly Pro Glu Val Thr Tyr Glu Pro
 115      120             125

Pro Pro Thr Ala Pro Thr Leu Leu Thr Val Leu Ala Tyr Ser Leu Leu
 130      135             140

Pro Ile Gly Gly Leu Ser
 145      150

```

```

<210> SEQ ID NO 20
<211> LENGTH: 150
<212> TYPE: PRT
<213> ORGANISM: Bos sp.

```

```

<400> SEQUENCE: 20

```

```

Met Thr Ala Pro Trp Ala Ala Leu Ala Leu Leu Trp Gly Ser Leu Cys
  1      5              10              15

Ala Gly Ser Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr
  20      25              30

Asn Ala Asn Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg
  35      40              45

```

-continued

Cys Glu Gly Glu Arg Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Arg
 50 55 60
 Asn Ser Ser Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp
 65 70 75 80
 Asp Phe Asn Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn
 85 90 95
 Pro Gln Val Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg
 100 105 110
 Phe Thr His Leu Pro Glu Ala Gly Gly Pro Glu Val Thr Tyr Glu Pro
 115 120 125
 Pro Pro Thr Ala Pro Thr Leu Leu Thr Val Leu Ala Tyr Ser Leu Leu
 130 135 140
 Pro Val Gly Gly Leu Ser
 145 150

<210> SEQ ID NO 21
 <211> LENGTH: 150
 <212> TYPE: PRT
 <213> ORGANISM: *Xenopus* sp.

<400> SEQUENCE: 21

Met Gly Ala Ser Val Ala Leu Thr Phe Leu Leu Leu Ala Thr Phe
 1 5 10 15
 Arg Ala Gly Ser Gly His Asp Glu Val Glu Thr Arg Glu Cys Ile Tyr
 20 25 30
 Tyr Asn Ala Asn Trp Glu Leu Glu Lys Thr Asn Gln Ser Gly Val Glu
 35 40 45
 Arg Leu Val Glu Gly Lys Lys Asp Lys Arg Leu His Cys Tyr Ala Ser
 50 55 60
 Trp Arg Asn Asn Ser Gly Phe Ile Glu Leu Val Lys Lys Gly Cys Trp
 65 70 75 80
 Leu Asp Asp Phe Asn Cys Tyr Asp Arg Gln Glu Cys Ile Ala Lys Glu
 85 90 95
 Glu Asn Pro Gln Val Phe Phe Cys Cys Cys Glu Gly Asn Tyr Cys Asn
 100 105 110
 Lys Lys Phe Thr His Leu Pro Glu Val Glu Thr Phe Asp Pro Lys Pro
 115 120 125
 Gln Pro Ser Ala Ser Val Leu Asn Ile Leu Ile Tyr Ser Leu Leu Pro
 130 135 140
 Ile Val Gly Leu Ser Met
 145 150

<210> SEQ ID NO 22
 <211> LENGTH: 150
 <212> TYPE: PRT
 <213> ORGANISM: *Homo sapiens*

<400> SEQUENCE: 22

Met Gly Ala Ala Ala Lys Leu Ala Phe Ala Val Phe Leu Ile Ser Cys
 1 5 10 15
 Ser Ser Gly Ala Ile Leu Gly Arg Ser Glu Thr Gln Glu Cys Leu Phe
 20 25 30
 Phe Asn Ala Asn Trp Glu Lys Asp Arg Thr Asn Gln Thr Gly Val Glu
 35 40 45
 Pro Cys Tyr Gly Asp Lys Asp Lys Arg Arg His Cys Phe Ala Thr Trp
 50 55 60

-continued

Lys Asn Ile Ser Gly Ser Ile Glu Ile Val Lys Gln Gly Cys Trp Leu
65 70 75 80

Asp Asp Ile Asn Cys Tyr Asp Arg Thr Asp Cys Val Glu Lys Lys Asp
85 90 95

Ser Pro Glu Val Tyr Phe Cys Cys Cys Glu Gly Asn Met Cys Asn Glu
100 105 110

Lys Phe Ser Tyr Phe Pro Glu Met Glu Val Thr Gln Pro Thr Ser Asn
115 120 125

Pro Val Thr Pro Lys Pro Pro Tyr Tyr Asn Ile Leu Leu Tyr Ser Leu
130 135 140

Val Pro Leu Met Leu Ile
145 150

<210> SEQ ID NO 23
 <211> LENGTH: 154
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 consensus polypeptide
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (8)..(8)
 <223> OTHER INFORMATION: Thr or Ala
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (121)..(121)
 <223> OTHER INFORMATION: Pro, Ala, Val or Met

<400> SEQUENCE: 23

Met Thr Ala Pro Trp Ala Ala Xaa Leu Ala Leu Leu Trp Gly Ser Leu
1 5 10 15

Cys Ala Gly Ser Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr
20 25 30

Tyr Asn Ala Asn Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu
35 40 45

Arg Leu Cys Glu Gly Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser
50 55 60

Trp Arg Asn Ser Ser Gly Thr Leu Glu Leu Val Lys Lys Gly Cys Trp
65 70 75 80

Leu Asp Asp Phe Asn Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu
85 90 95

Glu Asn Pro Gln Val Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn
100 105 110

Glu Arg Phe Thr His Leu Pro Glu Xaa Gly Gly Pro Glu Val Thr Tyr
115 120 125

Glu Pro Lys Pro Pro Thr Ala Pro Thr Leu Leu Thr Val Leu Ala Tyr
130 135 140

Ser Leu Leu Pro Ile Gly Gly Leu Ser Met
145 150

<210> SEQ ID NO 24
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 6xHis tag

<400> SEQUENCE: 24

His His His His His His

-continued

```

1           5

<210> SEQ ID NO 25
<211> LENGTH: 1107
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide

<400> SEQUENCE: 25

atggatgcaa tgaagagagg gctctgctgt gtgctgctgc tgtgtggagc agtcttcgtt    60
tcgcccggcg cctctggggc tggggaggct gagacacggg agtgcattcta ctacaacgcc    120
aactggggagc tggagcgcac caaccagagc ggcctgggagc gctgcgaagg cgagcaggac    180
aagcggctgc actgctacgc ctctctggcg aacagctctg gcaccatcga gctcgtgaag    240
aagggctgct gggatgatga cttcaactgc tacgataggc aggagtgtgt ggccactgag    300
gagaaccccc aggtgtactt ctgctgctgt gaaggcaact tctgcaacga gcgcttcact    360
catttgccag aggttggggg ccgggaagtc acgtacgagc ccccccgac agccccacc    420
ggtggtggaa ctcacacatg cccaccgtgc ccagcacctg aactcctggg gggaccgtca    480
gtcttcctct tcccccaaaa acccaaggac accctcatga tctcccgga cctgaggtc    540
acatgcgtgg tgggtgagct gagccacgaa gacctgagg tcaagttcaa ctggtacgtg    600
gacggcgtgg aggtgcataa tgccaagaca aagccgcggg aggagcagta caacagcacg    660
taccgtgtgg tcagcgtcct caccgtcctg caccaggact ggctgaatgg caaggagtac    720
aagtgcgaagg tctccaacaa agccctccca gcccccatcg agaaaaccat ctccaaagcc    780
aaagggcagc cccgagaacc acaggtgtac accctgcccc catcccggga ggagatgacc    840
aagaaccagg tcagcctgac ctgcctggtc aaaggcttct atcccagcga catcgccgtg    900
gagtgggaga gcaatgggca gccggagaac aactacaaga ccacgcctcc cgtgctggac    960
tccgacggct ccttcttct ctatagcaag ctcaccgtgg acaagagcag gtggcagcag    1020
gggaacgtct tctcatgctc cgtgatgcat gaggtcttgc acaaccacta cagcagaag    1080
agcctctccc tgtccccggg taaatga                                     1107

```

```

<210> SEQ ID NO 26
<211> LENGTH: 360
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polypeptide

```

```

<400> SEQUENCE: 26

Met Asp Ala Met Lys Arg Gly Leu Cys Cys Val Leu Leu Leu Cys Gly
1           5           10           15

Ala Val Phe Val Ser Pro Gly Ala Ala Glu Thr Arg Glu Cys Ile Tyr
          20           25           30

Tyr Asn Ala Asn Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu
          35           40           45

Arg Cys Glu Gly Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp
          50           55           60

Arg Asn Ser Ser Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Asp
65           70           75           80

Asp Asp Phe Asn Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu
          85           90           95

```

-continued

Asn Pro Gln Val Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu
 100 105 110
 Arg Phe Thr His Leu Pro Glu Ala Gly Gly Pro Glu Val Thr Tyr Glu
 115 120 125
 Pro Pro Pro Thr Gly Gly Gly Thr His Thr Cys Pro Pro Cys Pro Ala
 130 135 140
 Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
 145 150 155 160
 Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
 165 170 175
 Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
 180 185 190
 Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
 195 200 205
 Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
 210 215 220
 Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala
 225 230 235 240
 Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
 245 250 255
 Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr
 260 265 270
 Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
 275 280 285
 Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
 290 295 300
 Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
 305 310 315 320
 Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
 325 330 335
 Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
 340 345 350
 Ser Leu Ser Leu Ser Pro Gly Lys
 355 360

<210> SEQ ID NO 27
 <211> LENGTH: 1083
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polynucleotide
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (76)..(396)

<400> SEQUENCE: 27

atggatgcaa tgaagagagg gctctgctgt gtgctgctgc tgtgtggagc agtcttcgtt 60
 tcgcccggcg ccgct gag aca cgg gag tgc atc tac tac aac gcc aac tgg 111
 Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala Asn Trp
 1 5 10
 gag ctg gag cgc acc aac cag agc ggc ctg gag cgc tgc gaa ggc gag 159
 Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly Glu
 15 20 25
 cag gac aag cgg ctg cac tgc tac gcc tcc tgg cgc aac agc tct ggc 207
 Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Arg Asn Ser Ser Gly
 30 35 40

-continued

```

acc atc gag ctc gtg aag aag ggc tgc tgg gac gat gac ttc aac tgc      255
Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Asp Asp Asp Phe Asn Cys
45              50              55              60

tac gat agg cag gag tgt gtg gcc act gag gag aac ccc cag gtg tac      303
Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val Tyr
              65              70              75

ttc tgc tgc tgt gaa ggc aac ttc tgc aac gag cgc ttc act cat ttg      351
Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His Leu
              80              85              90

cca gag gct ggg ggc ccg gaa gtc acg tac gag cca ccc ccg aca      396
Pro Glu Ala Gly Gly Pro Glu Val Thr Tyr Glu Pro Pro Pro Thr
              95              100              105

ggtggtggaa ctcacacatg cccaccgtgc ccagcacctg aactcctggg gggaccgtca      456

gtcttctctt tcccccaaaa acccaaggac accctcatga tctcccgga cctgaggtc      516

acatgcgtgg tgggtggacgt gagccacgaa gaccctgagg tcaagttcaa ctggtacgtg      576

gacggcgtgg aggtgcataa tgccaagaca aagccgcggg aggagcagta caacagcacg      636

taccgtgtgg tcagcgtcct caccgtcctg caccaggact ggctgaatgg caaggagtac      696

aagtgcgaagg tctccaacaa agccctccca gcccccatcg agaaaacat ctccaaagcc      756

aaagggcagc cccgagaacc acaggtgtac accctgcccc catcccgga ggagatgacc      816

aagaaccagg tcagcctgac ctgcctggtc aaaggcttct atcccagcga catcgccgtg      876

gagtgggaga gcaatgggca gccggagaa aactacaaga ccacgcctcc cgtgctggac      936

tccgacggct ccttcttct ctatagcaag ctcaccgtgg acaagagcag gtggcagcag      996

gggaacgtct tctcatgtc cgtgatgcat gaggtctctg acaaccacta cacgcagaag    1056

agcctctccc tgtccccggg taaatga                                          1083

```

<210> SEQ ID NO 28

<211> LENGTH: 335

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 28

```

Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala Asn Trp Glu Leu Glu Arg
1              5              10              15

Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly Glu Gln Asp Lys Arg
              20              25              30

Leu His Cys Tyr Ala Ser Trp Arg Asn Ser Ser Gly Thr Ile Glu Leu
              35              40              45

Val Lys Lys Gly Cys Trp Asp Asp Asp Phe Asn Cys Tyr Asp Arg Gln
              50              55              60

Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val Tyr Phe Cys Cys Cys
65              70              75              80

Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His Leu Pro Glu Ala Gly
              85              90              95

Gly Pro Glu Val Thr Tyr Glu Pro Pro Pro Thr Gly Gly Gly Thr His
              100              105              110

Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val
              115              120              125

Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
              130              135              140

```

-continued

```

Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
145                      150                      155                      160

Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
                      165                      170                      175

Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
                      180                      185                      190

Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
                      195                      200                      205

Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
210                      215                      220

Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
225                      230                      235                      240

Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
                      245                      250                      255

Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
                      260                      265                      270

Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
                      275                      280                      285

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
290                      295                      300

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
305                      310                      315                      320

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
                      325                      330                      335

```

```

<210> SEQ ID NO 29
<211> LENGTH: 107
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polypeptide

```

```

<400> SEQUENCE: 29

```

```

Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala Asn Trp Glu Leu Glu Arg
1          5          10          15

Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly Glu Gln Asp Lys Arg
20        25        30

Leu His Cys Tyr Ala Ser Trp Arg Asn Ser Ser Gly Thr Ile Glu Leu
35        40        45

Val Lys Lys Gly Cys Trp Asp Asp Asp Phe Asn Cys Tyr Asp Arg Gln
50        55        60

Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val Tyr Phe Cys Cys Cys
65        70        75        80

Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His Leu Pro Glu Ala Gly
85        90        95

Gly Pro Glu Val Thr Tyr Glu Pro Pro Pro Thr
100      105

```

```

<210> SEQ ID NO 30
<211> LENGTH: 1083
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (76)..(396)

```

-continued

<400> SEQUENCE: 30

```

atggatgcaa tgaagagagg gctctgctgt gtgctgctgc tgtgtggagc agtcttcgtt      60
tcgcccggcg cgcgc gaa acc cgc gaa tgt att tat tac aat gct aat tgg      111
          1      5      10
gaa ctc gaa cgg acg aac caa tcc ggg ctc gaa cgg tgt gag ggg gaa      159
Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly Glu
          15      20      25
cag gat aaa cgc ctc cat tgc tat gcg tgg tgg agg aac tcc tcc ggg      207
Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Arg Asn Ser Ser Gly
          30      35      40
acg att gaa ctg gtc aag aaa ggg tgc tgg gac gac gat ttc aat tgt      255
Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Asp Asp Asp Phe Asn Cys
          45      50      55      60
tat gac cgc cag gaa tgt gtc gcg acc gaa gag aat ccg cag gtc tat      303
Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val Tyr
          65      70      75
ttc tgt tgt tgc gag ggg aat ttc tgt aat gaa cgg ttt acc cac ctc      351
Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His Leu
          80      85      90
ccc gaa gcc ggc ggg ccc gag gtg acc tat gaa ccc ccg ccc acc      396
Pro Glu Ala Gly Gly Pro Glu Val Thr Tyr Glu Pro Pro Pro Thr
          95      100      105
ggtggtggaa ctcacacatg cccaccgtgc ccagcacctg aactcctggg gggaccgtca      456
gtcttcctct tcccccaaaa acccaaggac accctcatga tctcccgga cctgaggtc      516
acatgcgtgg tgggtgaagt gagccacgaa gacctgagg tcaagttcaa ctggtacgtg      576
gacggcgtgg aggtgcataa tgccaagaca aagccgcggg aggagcagta caacagcacg      636
taccgtgtgg tcagcgtcct caccgtcctg caccaggact ggctgaatgg caaggagtac      696
aagtgcgaagg tctccaacaa agccctccca gcccctatcg agaaaaccat ctccaaagcc      756
aaagggcagc cccgagaacc acaggtgtac accctgcccc catcccgga ggagatgacc      816
aagaaccagg tcagcctgac ctgcctggtc aaaggcttct atcccagcga catcgccgtg      876
gagtgaggaga gcaatgggca gccgggagaac aactacaaga ccacgcctcc cgtgctggac      936
tccgacggct cttcttctct ctatagcaag ctcaccgtgg acaagagcag gtggcagcag      996
gggaacgtct tctcatgtc cgtgatgcat gaggtctcgc acaaccacta cagcagaag      1056
agcctctccc tgtccccggg taaatga      1083

```

<210> SEQ ID NO 31

<211> LENGTH: 321

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 31

```

gaaacccgcg aatgtattta ttacaatgct aattgggaac tcgaacggac gaaccaatcc      60
gggctcgaac ggtgtgaggg ggaacaggat aaacgcctcc attgctatgc gtcgtggagg      120
aactcctccg ggacgattga actggtcaag aaaggggtgct gggacgacga tttcaattgt      180
tatgaccgcc aggaatgtgt cgcgaccgaa gagaatccgc aggtctatct ctgttgttgc      240
gaggggaatt tctgtaatga acggtttacc cacctccccg aagccggcgg gcccgaggtg      300
acctatgaac cccgcccac c      321

```

-continued

<210> SEQ ID NO 32
 <211> LENGTH: 115
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 32

Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala Asn
 1 5 10 15
 Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly
 20 25 30
 Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Arg Asn Ser Ser
 35 40 45
 Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Asp Asp Asp Phe Asn
 50 55 60
 Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val
 65 70 75 80
 Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His
 85 90 95
 Leu Pro Glu Ala Gly Gly Pro Glu Val Thr Tyr Glu Pro Pro Pro Thr
 100 105 110
 Ala Pro Thr
 115

<210> SEQ ID NO 33
 <211> LENGTH: 1107
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polynucleotide

<400> SEQUENCE: 33

tacctaagtt acttctctcc cgagacgaca cagcagcagc acacacctcg tcagaagcaa 60
 agcggggcgc ggagaccgc acccctccga ctctgtgccc tcacgtagat gatgttgcg 120
 ttgacctcg acctcgctg gttggtctcg ccggacctcg cgacgcttcc gctcgctctg 180
 ttgcgccagc tgacgatgcg gaggaccgc ttgtcgagac cgtggtagct cgagcacttc 240
 ttcccgacga cctactact gaagttgacg atgctatccg tctcacaca ccggtgactc 300
 ctcttggggg tccacatgaa gacgacgaca cttccgttga agacgttgct cgcgaagtga 360
 gtaaacggtc tccgaccccc gggccttcag tgcattgctg gtgggggctg tggggggtgg 420
 ccaccacctt gagtgtgtac ggttggcacg ggtcgtggac ttgaggaccc cctgggcagt 480
 cagaaggaga aggggggttt tgggttctcg tgggagtact agagggcctg gggactccag 540
 tgtacgcacc accacctgca ctcggtgctt ctgggactcc agttcaagtt gacctgcac 600
 ctgccgcacc tccacgtatt acggttctgt ttcggcgccc tctcgtcat gttgtcgtgc 660
 atggcacacc agtcgcagga gtggcaggac gtggtcctga ccgacttacc gttcctcatg 720
 ttcacgttcc agaggttgtt tcgggagggt cgggggtagc tcttttgta gaggtttcgg 780
 tttcccgctg gggctcttgg tgtccacatg tgggacgggg gtagggccct cctctactgg 840
 ttcttgggtc agtcggactg gacggaccag ttccgaaga tagggctcgt gtagcggcac 900
 ctaccctct cgttaccgt cggcctcttg ttgatgttct ggtcggagg gcacgacctg 960
 aggtgcca ggaagaagga gatatcgtt gagtggcacc tgttctcgtc caccgtcgtc 1020
 cccttgca gaagtagcag gcactacgta ctccgagacg tgttggtgat gtgcgtcttc 1080

-continued

tcggagaggg acagggggccc atttact

1107

<210> SEQ ID NO 34

<211> LENGTH: 1083

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 34

tacctacgtt acttctctcc cgagacgaca cacgacgacg acacacctcg tcagaagcaa	60
agcggggccgc ggcgactctg tgccttcacg tagatgatgt tgcgggtgac cctcgacctc	120
gcgtgggttg tctcgccgga cctcgcgacg cttccgctcg tctgttctgc cgacgtgacg	180
atgcggagga ccgcgttgtc gagaccgtgg tagctcgagc acttcttccc gacgacctg	240
ctactgaagt tgacgatgct atccgtcttc acacaccggt gactcctctt ggggggtccac	300
atgaagacga cgacacttcc gttgaagacg ttgctcgca agtgagtaaa cggctctccga	360
cccccgggcc ttcagtgcac gctcgggtgg ggctgtccac caccttgagt gtgtacgggt	420
ggcacgggtc gtggacttga ggacccccct ggcagtcaga aggagaaggg gggttttggg	480
ttctgtggg agtactagag ggctgggga ctccagtgtc cgcaccacca cctgcactcg	540
gtgcttctgg gactccagtt caagttgacc atgcacctgc cgcacctcca cgtattacgg	600
ttctgtttcg gcgcctctct cgtcatgttg tcgtgcatgg cacaccagtc gcaggagtgg	660
caggacgtgg tcttgaccga cttaccgttc ctcatgttca cgttcagag gttgtttcgg	720
gagggtcggg ggtagctctt ttggtagagg ttctgggttc ccgtcggggc tcttggtgtc	780
cacatgtggg acgggggtag ggccctcttc tactggttct tgggtccagtc ggactggacg	840
gaccagtttc cgaagatagg gtcgctgtag cggcacctca cctctcgtt acccgtcggc	900
ctcttgttga tgttctgggt cggagggcac gacctgaggc tgccgaggaa gaaggagata	960
tcgttcgagt ggcacctgtt ctgcgtccacc gtcgtccctc tgcagaagag tacgaggcac	1020
tacgtactcc gagacgtgtt ggtgatgtgc gtcttctcgg agagggacag gggcccattt	1080
act	1083

<210> SEQ ID NO 35

<211> LENGTH: 1083

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 35

tacctacgtt acttctctcc cgagacgaca cacgacgacg acacacctcg tcagaagcaa	60
agcggggccgc ggcgggtttg ggcgcttaca taaataatgt tacgattaac ccttgagctt	120
gcctgcttgg ttaggccga gcttgccaca ctcccccttg tctattttgc ggaggtaacg	180
atagcagca cctccttgag gaggcctgc taacttgacc agttctttcc cagcacctg	240
ctgctaaagt taacaatact ggcggtcttc acacagcgtt ggcttctctt aggcgtccag	300
ataaagacaa caacgctccc cttaaagaca ttacttgcca aatgggtgga ggggcttcgg	360
ccgccggggc tccactggat acttgggggc ggggtggccac caccttgagt gtgtacgggt	420
ggcacgggtc gtggacttga ggacccccct ggcagtcaga aggagaaggg gggttttggg	480

-continued

```

ttctgtggg agtactagag ggcctgggga ctccagtgtg cgcaccacca cctgcactcg 540
gtgcttcttg gactccagtt caagttgacc atgcacctgc cgcacctcca cgtattacgg 600
ttctgtttcg gcgccctoct cgtcatgttg tcgtgcatgg cacaccagtc gcaggagtgg 660
caggacgtgg tctgaccga cttaccgttc ctcatgttca cgttcagag gttgtttcgg 720
gagggtcggg ggtagctctt ttggtagagg ttccgggttc ccgtcggggc tcttggtgtc 780
cacatgtggg acgggggtag ggcctctctc tactggttct tggtcagtc ggactggacg 840
gaccagtttc cgaagatagg gtcgctgtag cggcacctca ccctctcgtt acccgtcggc 900
ctcttggtga tgttctggtg cggagggcac gacctgaggg tgccgaggaa gaaggagata 960
tcgttcgagt ggcacctgtt ctgcctcacc gtcgtccct tgcagaagag tacgaggcac 1020
tacgtactcc gagacgtgtt ggtgatgtgc gtcttctcgg agagggacag gggcccattt 1080
act 1083

```

<210> SEQ ID NO 36

<211> LENGTH: 141

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 36

```

Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala Asn
1          5          10          15
Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly
20          25          30
Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Arg Asn Ser Ser
35          40          45
Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp Asp Phe Asn
50          55          60
Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val
65          70          75          80
Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His
85          90          95
Leu Pro Glu Ala Gly Gly Pro Glu Gly Pro Trp Ala Ser Thr Thr Ile
100          105          110
Pro Ser Gly Gly Pro Glu Ala Thr Ala Ala Ala Gly Asp Gln Gly Ser
115          120          125
Gly Ala Leu Trp Leu Cys Leu Glu Gly Pro Ala His Glu
130          135          140

```

<210> SEQ ID NO 37

<211> LENGTH: 141

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 37

```

Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala Asn
1          5          10          15
Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly
20          25          30
Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Arg Asn Ser Ser
35          40          45
Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Asp Asp Asp Phe Asn
50          55          60

```

-continued

Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val
 65 70 75 80
 Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His
 85 90 95
 Leu Pro Glu Ala Gly Gly Pro Glu Gly Pro Trp Ala Ser Thr Thr Ile
 100 105 110
 Pro Ser Gly Gly Pro Glu Ala Thr Ala Ala Ala Gly Asp Gln Gly Ser
 115 120 125
 Gly Ala Leu Trp Leu Cys Leu Glu Gly Pro Ala His Glu
 130 135 140

<210> SEQ ID NO 38
 <211> LENGTH: 370
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polypeptide

<400> SEQUENCE: 38

Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala Asn
 1 5 10 15
 Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly
 20 25 30
 Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Arg Asn Ser Ser
 35 40 45
 Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Asp Asp Asp Phe Asn
 50 55 60
 Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val
 65 70 75 80
 Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His
 85 90 95
 Leu Pro Glu Ala Gly Gly Pro Glu Gly Pro Trp Ala Ser Thr Thr Ile
 100 105 110
 Pro Ser Gly Gly Pro Glu Ala Thr Ala Ala Ala Gly Asp Gln Gly Ser
 115 120 125
 Gly Ala Leu Trp Leu Cys Leu Glu Gly Pro Ala His Glu Thr Gly Gly
 130 135 140
 Gly Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
 145 150 155 160
 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
 165 170 175
 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 180 185 190
 Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 195 200 205
 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
 210 215 220
 Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
 225 230 235 240
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
 245 250 255
 Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
 260 265 270
 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu

-continued

275	280	285
Thr Cys Leu Val Lys Gly	Phe Tyr Pro Ser Asp	Ile Ala Val Glu Trp
290	295	300
Glu Ser Asn Gly Gln Pro	Glu Asn Asn Tyr Lys	Thr Thr Pro Pro Val
305	310	315
Leu Asp Ser Asp Gly Ser	Phe Phe Leu Tyr Ser	Lys Leu Thr Val Asp
	325	330
Lys Ser Arg Trp Gln Gln	Gly Asn Val Phe Ser	Cys Ser Val Met His
	340	345
Glu Ala Leu His Asn His	Tyr Thr Gln Lys Ser	Leu Ser Leu Ser Pro
	355	360
Gly Lys		
370		

<210> SEQ ID NO 39
 <211> LENGTH: 512
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 39

Met Thr Ala Pro Trp Val	Ala Leu Ala Leu Leu Trp	Gly Ser Leu Trp
1	5	10
Pro Gly Ser Gly Arg Gly	Glu Ala Glu Thr Arg	Glu Cys Ile Tyr Tyr
	20	25
Asn Ala Asn Trp Glu Leu	Glu Arg Thr Asn Gln Ser	Gly Leu Glu Arg
	35	40
Cys Glu Gly Glu Gln Asp	Lys Arg Leu His Cys	Tyr Ala Ser Trp Ala
	50	55
Asn Ser Ser Gly Thr Ile	Glu Leu Val Lys Lys	Gly Cys Trp Leu Asp
	65	70
Asp Phe Asn Cys Tyr Asp	Arg Gln Glu Cys Val	Ala Thr Glu Glu Asn
	85	90
Pro Gln Val Tyr Phe Cys	Cys Cys Glu Gly Asn	Phe Cys Asn Glu Arg
	100	105
Phe Thr His Leu Pro Glu	Ala Gly Gly Pro Glu	Val Thr Tyr Glu Pro
	115	120
Pro Pro Thr Ala Pro Thr	Leu Leu Thr Val Leu	Ala Tyr Ser Leu Leu
	130	135
Pro Ile Gly Gly Leu Ser	Leu Ile Val Leu Leu	Ala Phe Trp Met Tyr
	145	150
Arg His Arg Lys Pro Pro	Tyr Gly His Val Asp	Ile His Glu Asp Pro
	165	170
Gly Pro Pro Pro Pro Ser	Pro Leu Val Gly Leu	Lys Pro Leu Gln Leu
	180	185
Leu Glu Ile Lys Ala Arg	Gly Arg Phe Gly Cys	Val Trp Lys Ala Gln
	195	200
Leu Met Asn Asp Phe Val	Ala Val Lys Ile Phe	Pro Leu Gln Asp Lys
	210	215
Gln Ser Trp Gln Ser Glu	Arg Glu Ile Phe Ser	Thr Pro Gly Met Lys
	225	230
His Glu Asn Leu Leu Gln	Phe Ile Ala Ala Glu	Lys Arg Gly Ser Asn
	245	250
Leu Glu Val Glu Leu Trp	Leu Ile Thr Ala Phe	His Asp Lys Gly Ser
	260	265
		270

-continued

Leu Thr Asp Tyr Leu Lys Gly Asn Ile Ile Thr Trp Asn Glu Leu Cys
 275 280 285
 His Val Ala Glu Thr Met Ser Arg Gly Leu Ser Tyr Leu His Glu Asp
 290 295 300
 Val Pro Trp Cys Arg Gly Glu Gly His Lys Pro Ser Ile Ala His Arg
 305 310 315 320
 Asp Phe Lys Ser Lys Asn Val Leu Leu Lys Ser Asp Leu Thr Ala Val
 325 330 335
 Leu Ala Asp Phe Gly Leu Ala Val Arg Phe Glu Pro Gly Lys Pro Pro
 340 345 350
 Gly Asp Thr His Gly Gln Val Gly Thr Arg Arg Tyr Met Ala Pro Glu
 355 360 365
 Val Leu Glu Gly Ala Ile Asn Phe Gln Arg Asp Ala Phe Leu Arg Ile
 370 375 380
 Asp Met Tyr Ala Met Gly Leu Val Leu Trp Glu Leu Val Ser Arg Cys
 385 390 395 400
 Lys Ala Ala Asp Gly Pro Val Asp Glu Tyr Met Leu Pro Phe Glu Glu
 405 410 415
 Glu Ile Gly Gln His Pro Ser Leu Glu Glu Leu Gln Glu Val Val Val
 420 425 430
 His Lys Lys Met Arg Pro Thr Ile Lys Asp His Trp Leu Lys His Pro
 435 440 445
 Gly Leu Ala Gln Leu Cys Val Thr Ile Glu Glu Cys Trp Asp His Asp
 450 455 460
 Ala Glu Ala Arg Leu Ser Ala Gly Cys Val Glu Glu Arg Val Ser Leu
 465 470 475 480
 Ile Arg Arg Ser Val Asn Gly Thr Thr Ser Asp Cys Leu Val Ser Leu
 485 490 495
 Val Thr Ser Val Thr Asn Val Asp Leu Pro Pro Lys Glu Ser Ser Ile
 500 505 510

<210> SEQ ID NO 40
 <211> LENGTH: 115
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 40

Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala Asn
 1 5 10 15
 Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly
 20 25 30
 Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Ala Asn Ser Ser
 35 40 45
 Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp Asp Phe Asn
 50 55 60
 Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val
 65 70 75 80
 Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His
 85 90 95
 Leu Pro Glu Ala Gly Gly Pro Glu Val Thr Tyr Glu Pro Pro Pro Thr
 100 105 110
 Ala Pro Thr
 115

<210> SEQ ID NO 41

-continued

<211> LENGTH: 100
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 41

Gly Arg Gly Glu Ala Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala Asn
 1 5 10 15
 Trp Glu Leu Glu Arg Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly
 20 25 30
 Glu Gln Asp Lys Arg Leu His Cys Tyr Ala Ser Trp Ala Asn Ser Ser
 35 40 45
 Gly Thr Ile Glu Leu Val Lys Lys Gly Cys Trp Leu Asp Asp Phe Asn
 50 55 60
 Cys Tyr Asp Arg Gln Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val
 65 70 75 80
 Tyr Phe Cys Cys Cys Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His
 85 90 95
 Leu Pro Glu Ala
 100

<210> SEQ ID NO 42
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 peptide

<400> SEQUENCE: 42

Thr Gly Gly Gly
 1

<210> SEQ ID NO 43
 <211> LENGTH: 368
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polypeptide

<400> SEQUENCE: 43

Met Asp Ala Met Lys Arg Gly Leu Cys Cys Val Leu Leu Leu Cys Gly
 1 5 10 15
 Ala Val Phe Val Ser Pro Gly Ala Ser Gly Arg Gly Glu Ala Glu Thr
 20 25 30
 Arg Glu Cys Ile Tyr Tyr Asn Ala Asn Trp Glu Leu Glu Arg Thr Asn
 35 40 45
 Gln Ser Gly Leu Glu Arg Cys Glu Gly Glu Gln Asp Lys Arg Leu His
 50 55 60
 Cys Tyr Ala Ser Trp Arg Asn Ser Ser Gly Thr Ile Glu Leu Val Lys
 65 70 75 80
 Lys Gly Cys Trp Asp Asp Asp Phe Asn Cys Tyr Asp Arg Gln Glu Cys
 85 90 95
 Val Ala Thr Glu Glu Asn Pro Gln Val Tyr Phe Cys Cys Cys Glu Gly
 100 105 110
 Asn Phe Cys Asn Glu Arg Phe Thr His Leu Pro Glu Ala Gly Gly Pro
 115 120 125
 Glu Val Thr Tyr Glu Pro Pro Pro Thr Ala Pro Thr Gly Gly Gly Thr
 130 135 140

-continued

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser															
145				150				155						160	
Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg															
			165				170						175		
Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro															
			180				185						190		
Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala															
			195				200						205		
Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val															
			210				215						220		
Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr															
			225				230						235		240
Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr															
							245						250		255
Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu															
			260										265		270
Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys															
			275										280		285
Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser															
			290				295						300		
Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp															
			305				310						315		320
Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser															
							325						330		335
Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala															
							340						345		350
Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys															
			355										360		365

<210> SEQ ID NO 44

<211> LENGTH: 107

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 44

Glu Thr Arg Glu Cys Ile Tyr Tyr Asn Ala Asn Trp Glu Leu Glu Arg															
1			5				10						15		
Thr Asn Gln Ser Gly Leu Glu Arg Cys Glu Gly Glu Gln Asp Lys Arg															
			20				25						30		
Leu His Cys Tyr Ala Ser Trp Arg Asn Ser Ser Gly Thr Ile Glu Leu															
			35				40						45		
Val Lys Lys Gly Cys Trp Asp Asp Asp Phe Asn Cys Tyr Asp Arg Gln															
			50				55						60		
Glu Cys Val Ala Thr Glu Glu Asn Pro Gln Val Tyr Phe Cys Cys Cys															
			65				70						75		80
Glu Gly Asn Phe Cys Asn Glu Arg Phe Thr His Leu Pro Glu Ala Gly															
							85						90		95
Gly Pro Glu Val Thr Tyr Glu Pro Pro Pro Thr															
			100										105		

107

We claim:

1. A method of treating anemia in a subject in need thereof, comprising subcutaneously administering to the subject a polypeptide comprising an amino acid sequence that is at least 90% identical to the sequence of amino acids 29-109 of SEQ ID NO: 1, wherein the polypeptide comprises an acidic amino acid at the position corresponding to position 79 of SEQ ID NO: 1, wherein the polypeptide binds to GDF11 and/or myostatin.

2. The method of claim 1, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to the sequence of amino acids 29-109 of SEQ ID NO: 1.

3. The method of claim 2, wherein the acidic amino acid is glutamic acid.

4. The method of claim 2, wherein the acidic amino acid is aspartic acid.

5. The method of claim 4, wherein the polypeptide further comprises a constant region of an immunoglobulin.

6. The method of claim 5, wherein the constant region is derived from an IgG heavy chain.

7. The method of claim 6, wherein the constant region of an immunoglobulin is an Fc domain.

8. The method of claim 7, wherein the polypeptide forms a homodimer.

9. The method of claim 2, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to SEQ ID NO: 28.

10. The method of claim 2, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to SEQ ID NO: 28.

11. The method of claim 2, wherein the polypeptide comprises an amino acid sequence that is at least 99% identical to SEQ ID NO: 28.

12. The method of claim 11, wherein the subject has myelodysplastic syndrome.

13. The method of claim 11, wherein the subject has a thalassemia.

14. The method of claim 2, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 28.

108

15. The method of claim 1, wherein the polypeptide comprises one or more modified amino acid residues selected from: a glycosylated amino acid, a PEGylated amino acid, a farnesylated amino acid, an acetylated amino acid, a biotinylated amino acid, an amino acid conjugated to a lipid moiety, and an amino acid conjugated to an organic derivatizing agent.

16. The method of claim 1, wherein the polypeptide binds to GDF11.

17. The method of claim 1, wherein the polypeptide binds to myostatin.

18. The method of claim 1, wherein the polypeptide binds to myostatin and GDF11.

19. The method of claim 1, wherein the polypeptide inhibits signaling by GDF11 and myostatin in a cell-based assay.

20. The method of claim 1, wherein the polypeptide inhibits signaling by GDF11 in a cell-based assay.

21. The method of claim 1, wherein the polypeptide inhibits signaling by myostatin in a cell-based assay.

22. The method of claim 1, wherein the polypeptide comprises the sequence of amino acids 29-109 of SEQ ID NO: 1, but wherein the polypeptide comprises an acidic amino acid at the position corresponding to position 79 of SEQ ID NO: 1.

23. The method of claim 1, wherein the polypeptide comprises an amino acid sequence that is at least 90% identical to the sequence of amino acids 25-131 of SEQ ID NO: 1.

24. The method of claim 1, wherein the polypeptide comprises an amino acid sequence that is at least 95% identical to the sequence of amino acids 25-131 of SEQ ID NO: 1.

25. The method of claim 1, wherein the polypeptide comprises the sequence of amino acids 25-131 of SEQ ID NO: 1, but wherein the polypeptide comprises an acidic amino acid at the position corresponding to position 79 of SEQ ID NO: 1.

* * * * *