

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 February 2009 (05.02.2009)

PCT

(10) International Publication Number
WO 2009/017831 A1

(51) International Patent Classification:
C07D 213/04 (2006.01) C07D 213/22 (2006.01)

(21) International Application Number:
PCT/US2008/009346

(22) International Filing Date: 31 July 2008 (31.07.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/962,823 1 August 2007 (01.08.2007) US

(71) Applicant (for all designated States except US): SYNTA
PHARMACEUTICALS CORP. [US/US]; 45 Hartwell
Ave., Lexington, MA 02421 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): CHEN, Shoujun
[US/US]; 19 Dunster Road, Bedford, MA 01730 (US).

(74) Agents: HSI, Jeffrey D. et al.; Edwards Angell Palmer &
Dodge LLP, P.O. Box 55874, Boston, MA 02205 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE,
EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID,
IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK,
LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW,
MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL,
NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG,
CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— with international search report



WO 2009/017831 A1

(54) Title: VINYL-ARYL DERIVATIVES FOR INFLAMMATION AND IMMUNE-RELATED USES

(57) Abstract: The invention relates to compounds that are useful as immunosuppressive agents and for treating and preventing inflammatory conditions, allergic disorders, and immune disorders.

**VINYL-ARYL DERIVATIVES FOR INFLAMMATION AND IMMUNE-RELATED
USES**

RELATED APPLICATIONS

5

This application claims the benefit of U.S. Provisional Application No. 60/962,823, filed August 1, 2007, the entire teachings of which are incorporated herein by reference.

FIELD OF THE INVENTION

10

This invention relates to biologically active chemical compounds, namely vinyl-phenyl derivatives that may be used for immunosuppression or to treat or prevent inflammatory conditions, allergic disorders and immune disorders.

BACKGROUND OF THE INVENTION

15

Inflammation is a mechanism that protects mammals from invading pathogens. However, while transient inflammation is necessary to protect a mammal from infection, uncontrolled inflammation causes tissue damage and is the underlying cause of many illnesses. Inflammation is typically initiated by binding of an antigen to T-cell antigen receptor. Antigen binding by a T-cell initiates calcium influx into the cell via calcium ion channels, such as Ca²⁺-release-activated Ca²⁺ channels (CRAC). Calcium ion influx in turn initiates a signaling cascade that leads to activation of these cells and an inflammatory response characterized by cytokine production.

20

25

Interleukin 2 (IL-2) is a cytokine that is secreted by T cells in response to calcium ion influx into the cell. IL-2 modulates immunological effects on many cells of the immune system. For example, it is a potent T cell mitogen that is required for T cell proliferation, promoting their progression from G1 to S phase of the cell cycle; it stimulates the growth of NK cells; and it acts as a growth factor to B cells and stimulates antibody synthesis.

30

IL-2, although useful in the immune response, can cause a variety of problems. IL-2 damages the blood-brain barrier and the endothelium of brain vessels. These effects may be the underlying causes of neuropsychiatric side effects observed under IL-2 therapy, e.g. fatigue, disorientation and depression. It also alters the electrophysiological behavior of neurons.

35

Due to its effects on both T and B cells, IL-2 is a major central regulator of immune responses. It plays a role in inflammatory reactions, tumor surveillance, and hematopoiesis.

It also affects the production of other cytokines, inducing IL-1, TNF- α and TNF- β secretion, as well as stimulating the synthesis of IFN- γ in peripheral leukocytes.

5 T cells that are unable to produce IL-2 become inactive (anergic). This renders them potentially inert to any antigenic stimulation they might receive in the future. As a result, agents which inhibit IL-2 production can be used for immunosuppression or to treat or prevent inflammation and immune disorders. This approach has been clinically validated with immunosuppressive drugs such as cyclosporin, FK506, and RS61443. Despite this proof of concept, agents that inhibit IL-2 production remain far from ideal. Among other problems, 10 efficacy limitations and unwanted side effects (including dose-dependant nephrotoxicity and hypertension) hinder their use.

Over production of proinflammatory cytokines other than IL-2 has also been implicated in many autoimmune diseases. For example, Interleukin 5 (IL-5), a cytokine that increases the 15 production of eosinophils, is increased in asthma. Overproduction of IL-5 is associated with accumulation of eosinophils in the asthmatic bronchial mucosa, a hall mark of allergic inflammation. Thus, patients with asthma and other inflammatory disorders involving the accumulation of eosinophils would benefit from the development of new drugs that inhibit the production of IL-5.

20 Interleukin 4 (IL-4) and interleukin 13 (IL-13) have been identified as mediators of the hypercontractility of smooth muscle found in inflammatory bowel disease and asthma. Thus, patients with asthma and inflammatory bowel disease would benefit from the development of new drugs that inhibit IL-4 and IL-13 production.


25 Granulocyte macrophage-colony stimulating factor (GM-CSF) is a regulator of maturation of granulocyte and macrophage lineage population and has been implicated as a key factor in inflammatory and autoimmune diseases. Anti-GM-CSF antibody blockade has been shown to ameliorate autoimmune disease. Thus, development of new drugs that inhibit the production 30 of GM-CSF would be beneficial to patients with an inflammatory or autoimmune disease.

There is a continuing need for new drugs which overcome one or more of the shortcomings of drugs currently used for immunosuppression or in the treatment or prevention of inflammatory disorders, allergic disorders and autoimmune disorders. Desirable properties of 35 new drugs include efficacy against diseases or disorders that are currently untreatable or poorly treatable, new mechanism of action, oral bioavailability and/or reduced side effects.

SUMMARY OF THE INVENTION

This invention meets the above-mentioned needs by providing certain vinyl-phenyl derivatives that inhibit the activity of CRAC ion channels and inhibit the production of IL-2, IL-4, IL-5, IL-13, GM-CSF, TNF- α , and IFN γ . These compounds are particularly useful for immunosuppression and/or to treat or prevent inflammatory conditions, allergic disorders and immune disorders.

In one embodiment, the invention relates to compounds of structural formula (I):



(I)

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

X_1 and X_2 are, independently, CH or N;

one of X_3 or X_6 is NR_{29} , and the other is CH_2 , CHR_2 , or $C(R_2)_2$;

L is a linker selected from the group consisting of $-NHCH_2-$, $-CH_2NH-$, $-NR_5-C(O)-$, $-C(O)-NR_5-$, $-NR_5-C(S)-$, $-C(S)-NR_5-$, $-NR_5-C(NR_8)-$, $-C(NR_8)-NR_5-$, $-NR_5C(S)NR_5-$, $-NR_5C(NR_8)NR_5-$, or $-NR_5CR^aR^bNR_5-$;

Y is a substituted phenyl, provided that Y is not substituted with $-NHC(O)R_{30}$;

R_1 is H or a substituent;

R_2 , for each occurrence is, independently, a substituent, provided that R_2 is not $=O$ or $=C(R_{28})_2$;

R_4 is a substituent;

R^a and R^b , for each occurrence, are independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, cyano, nitro, halo, $-OR_5$, $-SR_5$, $-NR_6R_7$, $-C(O)NR_6R_7$, $-NR_5C(O)R_5$, $-C(O)R_5$, $-C(O)OR_5$, $-OC(O)R_5$, $-C(O)SR_5$, $-SC(O)R_5$, $-C(S)NR_6R_7$, $-NR_5C(S)R_5$, $-C(S)R_5$, $-C(S)OR_5$, $-OC(S)R_5$, $-C(S)SR_5$, $-SC(S)R_5$, $-C(NR_8)NR_6R_7$, $-NR_5C(NR_8)R_5$, $-C(NR_8)R_5$, $-C(NR_8)OR_5$, $-OC(NR_8)R_5$, $-C(NR_8)SR_5$, $-SC(NR_8)R_5$, $-OC(O)OR_5$, $-OC(O)NR_6R_7$, $-NR_5C(O)OR_5$, $-NR_5C(O)NR_6R_7$, $-SC(O)OR_5$,

SC(O)NR₆R₇, -SC(O)SR₅, -NR₅C(O)SR₅, -OC(O)SR₅, -OC(S)OR₅, -OC(S)NR₆R₇, -NR₅C(S)OR₅, -NR₅C(S)NR₆R₇, -SC(S)OR₅, -SC(S)NR₆R₇, -SC(S)SR₅, -NR₅C(S)SR₅, -OC(S)SR₅, -OC(NR₈)OR₅, -OC(NR₈)NR₆R₇, -NR₅C(NR₈)OR₅, -NR₅C(NR₈)NR₆R₇, -SC(NR₈)OR₅, -SC(NR₈)NR₆R₇, -SC(NR₈)SR₅, -NR₅C(NR₈)SR₅, or -OC(NR₈)SR₅;

5 R₅, for each occurrence, is independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

10 R₆ and R₇, for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R₆ and R₇ taken together with the
15 nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

R₈, for each occurrence, is independently -H, a halo, an alkyl, -OR₅, -NR₆R₇, -C(O)R₅, -C(O)OR₅, or -C(O)NR₆R₇;

R₂₈, for each occurrence is, independently, H or a substituent;

20 R₂₉ is -C(O)R₃₀ or a N-containing 6-membered aromatic ring;

R₃₀ is -H, -OR₅, -SR₅, -NR₆R₇, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an
25 optionally substituted heteraralkyl;

m is 0, 1, or 2; and

n is 0, 1 or 2.

In one aspect, the compound of formula (I) is not:

30 5-[4-(2,6-difluoro-benzoylamino)-phenyl]-4-methyl-2,4-dihydro-2H-pyridine-1-carboxylic acid ethyl ester;

Ethyl 3-(4-(2,6-difluorobenzamido)phenyl)-4-methyl-5,6-dihydropyridine-1(2H)-carboxylate;

Ethyl 4-(4-(2,6-difluorobenzamido)phenyl)-3-methyl-5,6-dihydropyridine-1(2H)-carboxylate;

35 2,6-Difluoro-N-[4-(1-methoxycarbonyl-4-thiazol-2-yl)-1,2,5,6-tetrahydro-pyridin-3-yl]-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-methoxycarbonyl-4-isopropyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-methoxycarbonyl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide;

5 2,6-Difluoro-N-[4-(1-allyloxycarbonyl-3-methyl-1,2,5,6-tetrahydro-pyridin-4-yl)-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-phenyloxycarbonyl-3-methyl-1,2,5,6-tetrahydro-pyridin-4-yl)-phenyl]-benzamide;

10 2,6-Difluoro-N-[4-(1-acetyl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-cyclopropylcarbonyl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide;

2,6-Difluoro-N-{4-[1-(pyridine-3-carbonyl)-cyclopropanecarbonyl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl]-phenyl}-benzamide; or

15 2,6-Difluoro-N-[4-(1-pyrimidin-2-yl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide.

A compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof is particularly useful inhibiting immune cell (e.g., mast cells, T-cells and/or
20 B-cells) activation (e.g., cytokine production and/or proliferation in response to an antigen; and or mast cell degranulation). In particular, a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof can inhibit the production of certain cytokines that regulate immune cell activation. For example, a
25 compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof can inhibit the production of IL-2, IL-4, IL-5, IL-13, GM-CSF, TNF- α , INF- γ or combinations thereof. Moreover, a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof can modulate the activity of one or more ion channel involved in activation of immune cells, such as CRAC ion channels.

30 In one embodiment, compounds of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof are particularly useful for inhibiting mast cell degranulation. Mast cell degranulation has been implicated in allergic reactions.

35 A compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof is particularly useful for immunosuppression or for treating or preventing inflammatory conditions, allergic disorders, and immune disorders.

The invention also encompasses pharmaceutical compositions comprising a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof; and a pharmaceutically acceptable carrier or vehicle. These compositions may further comprise additional agents. These compositions are useful for immunosuppression and treating or preventing inflammatory conditions, allergic disorders and immune disorders.

The invention further encompasses methods for treating or preventing inflammatory conditions, allergic disorders, and immune disorders, comprising administering to a subject in need thereof an effective amount of a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, or a pharmaceutical composition comprising a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof. These methods may also comprise administering to the subject an additional agent separately or in a combination composition with the compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

The invention further encompasses methods for suppressing the immune system of a subject, comprising administering to a subject in need thereof an effective amount of a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, or a pharmaceutical composition comprising a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof. These methods may also comprise administering to the subject an additional agent separately or in a combination composition with the compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

The invention further encompasses methods for inhibiting immune cell activation, including inhibiting proliferation of T cells and/or B cells, *in vivo* or *in vitro* comprising administering to the cell an effective amount of a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof or a pharmaceutical composition comprising a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

The invention further encompasses methods for inhibiting cytokine production in a cell (e.g., IL-2, IL-4, IL-5, IL-13, GM-CSF, TNF- α , and/or INF- γ production) *in vivo* or *in vitro* comprising administering to a cell an effective amount of a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof or a pharmaceutical composition comprising a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

The invention further encompasses methods for modulating ion channel activity (e.g., CRAC channel activity) *in vivo* or *in vitro* comprising administering an effective amount of a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof or a pharmaceutical composition comprising a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

All of the methods of this invention may be practice with a compound of the invention alone, or in combination with other agents, such as other immunosuppressive agents, anti-inflammatory agents, agents for the treatment of allergic disorders or agents for the treatment of immune disorders.

DETAILED DESCRIPTION OF THE INVENTION

15 DEFINITIONS

Unless otherwise specified, the below terms used herein are defined as follows:

As used herein, the term an "aromatic ring" or "aryl" means a monocyclic or polycyclic-aromatic ring or ring radical comprising carbon and hydrogen atoms. Examples of suitable aryl groups include, but are not limited to, phenyl, tolyl, anthaceny, fluorenyl, indenyl, azuleny, and naphthyl, as well as benzo-fused carbocyclic moieties such as 5,6,7,8-tetrahydronaphthyl. An aryl group can be unsubstituted or substituted with one or more substituents (including without limitation alkyl (preferably, lower alkyl or alkyl substituted with one or more halo), hydroxy, alkoxy (preferably, lower alkoxy), alkylsulfanyl, cyano, halo, amino, and nitro. In certain embodiments, the aryl group is a monocyclic ring, wherein the ring comprises 6 carbon atoms.

As used herein, the term "alkyl" means a saturated straight chain or branched non-cyclic hydrocarbon typically having from 1 to 10 carbon atoms. Representative saturated straight chain alkyls include methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl and n-decyl; while saturated branched alkyls include isopropyl, *sec*-butyl, isobutyl, *tert*-butyl, isopentyl, 2-methylbutyl, 3-methylbutyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 2-methylhexyl, 3-methylhexyl, 4-methylhexyl, 5-methylhexyl, 2,3-dimethylbutyl, 2,3-dimethylpentyl, 2,4-dimethylpentyl, 2,3-dimethylhexyl, 2,4-dimethylhexyl, 2,5-dimethylhexyl, 2,2-dimethylpentyl, 2,2-dimethylhexyl, 3,3-dimethylpentyl, 3,3-dimethylhexyl, 4,4-dimethylhexyl, 2-ethylpentyl, 3-ethylpentyl, 2-ethylhexyl, 3-ethylhexyl, 4-ethylhexyl, 2-methyl-2-ethylpentyl, 2-methyl-3-ethylpentyl, 2-

methyl-4-ethylpentyl, 2-methyl-2-ethylhexyl, 2-methyl-3-ethylhexyl, 2-methyl-4-ethylhexyl, 2,2-diethylpentyl, 3,3-diethylhexyl, 2,2-diethylhexyl, 3,3-diethylhexyl and the like. Alkyl groups included in compounds of this invention may be optionally substituted with one or more substituents. Examples of substituents include, but are not limited to, amino, alkylamino, alkoxy, alkylsulfanyl, oxo, halo, acyl, nitro, hydroxyl, cyano, aryl, alkylaryl, aryloxy, arylsulfanyl, arylamino, carbocyclyl, carbocyclyloxy, carbocyclylthio, carbocyclylamino, heterocyclyl, heterocyclyloxy, heterocyclylamino, heterocyclylthio, and the like. In addition, any carbon in the alkyl segment may be substituted with oxygen (=O), sulfur (=S), or nitrogen (=NR²², wherein R²² is -H, an alkyl, acetyl, or aralkyl). Lower alkyls are typically preferred for the compounds of this invention.

The term alkylene refers to an alkyl group or a cycloalkyl group that has two points of attachment to two moieties (e.g., {-CH₂-}, {-CH₂CH₂-},



, etc., wherein the brackets indicate the points of attachment). Alkylene groups may be substituted or unsubstituted with one or more substituents.

An aralkyl group refers to an aryl group that is attached to another moiety via an alkylene linker. Aralkyl groups can be substituted or unsubstituted with one or more substituents.

The term "alkoxy," as used herein, refers to an alkyl group which is linked to another moiety through an oxygen atom. Alkoxy groups can be substituted or unsubstituted with one or more substituents.

The term "alkylsulfanyl," as used herein, refers to an alkyl group which is linked to another moiety through a divalent sulfur atom. Alkylsulfanyl groups can be substituted or unsubstituted with one or more substituents.

The term "arylsulfanyl," as used herein, refers to an aryl group which is linked to another moiety through a divalent sulfur atom. Arylsulfanyl groups can be substituted or unsubstituted with one or more substituents.

The term "alkyl ester" as used herein, refers to a group represented by the formula –
C(O)OR₃₂, wherein R₃₂ is an alkyl group. A lower alkyl ester is a group represented by the
formula –C(O)OR₃₂, wherein R₃₂ is a lower alkyl group.

- 5 The term "heteroalkyl," as used herein, refers to an alkyl group which has one or more
carbons in the alkyl chain replaced with an –O-, -S- or –NR₂₇-, wherein R₂₇ is H or a lower
alkyl. Heteroalkyl groups can be substituted or unsubstituted with one or more substituents.

The term "alkylamino," as used herein, refers to an amino group in which one hydrogen atom
10 attached to the nitrogen has been replaced by an alkyl group. The term "dialkylamino," as
used herein, refers to an amino group in which two hydrogen atoms attached to the nitrogen
have been replaced by alkyl groups, in which the alkyl groups can be the same or different.
Alkylamino groups and dialkylamino groups can be substituted or unsubstituted with one or
more substituents.

15

As used herein, the term "alkenyl" means a straight chain or branched, hydrocarbon radical
typically having from 2 to 10 carbon atoms and having at least one carbon-carbon double
bond. Representative straight chain and branched alkenyls include vinyl, allyl, 1-butenyl,
2-butenyl, isobutylenyl, 1-pentenyl, 2-pentenyl, 3-methyl-1-butenyl, 1-methyl-2-butenyl,
20 2,3-dimethyl-2-butenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 1-heptenyl, 2-heptenyl, 3-heptenyl,
1-octenyl, 2-octenyl, 3-octenyl, 1-nonenyl, 2-nonenyl, 3-nonenyl, 1-decenyl, 2-decenyl, 3-
decenyl and the like. Alkenyl groups can be substituted or unsubstituted with one or more
substituents.

25

As used herein, the term "alkynyl" means a straight chain or branched, hydrocarbon radical
typically having from 2 to 10 carbon atoms and having at least one carbon-carbon triple
bond. Representative straight chain and branched alkynyls include acetylenyl, propynyl,
1-butyne, 2-butyne, 1-pentyne, 2-pentyne, 3-methyl-1-butyne, 4-pentyne, 1-hexyne, 2-
hexyne, 5-hexyne, 1-heptyne, 2-heptyne, 6-heptyne, 1-octyne, 2-octyne, 7-octyne, 1-
30 nonyne, 2-nonyne, 8-nonyne, 1-decyne, 2-decyne, 9-decyne and the like. Alkynyl
groups can be substituted or unsubstituted with one or more substituents.

35

As used herein, the term "cycloalkyl" means a saturated, mono- or polycyclic alkyl radical
typically having from 3 to 14 carbon atoms. Representative cycloalkyls include cyclopropyl,
cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl,
adamantly, decahydronaphthyl, octahydropentalene, bicycle[1.1.1]pentanyl, and the like.
Cycloalkyl groups can be substituted or unsubstituted with one or more substituents.

As used herein, the term "cycloalkenyl" means a cyclic non-aromatic alkenyl radical having at least one carbon-carbon double bond in the cyclic system and typically having from 5 to 14 carbon atoms. Representative cycloalkenyls include cyclopentenyl, cyclopentadienyl, cyclohexenyl, cyclohexadienyl, cycloheptenyl, cycloheptadienyl, cycloheptatrienyl, cyclooctenyl, cyclooctadienyl, cyclooctatrienyl, cyclooctatetraenyl, cyclononenyl, cyclononadienyl, cyclodecenyl, cyclodecadienyl and the like. Cycloalkenyl groups can be substituted or unsubstituted with one or more substituents.

As used herein, the term "heterocyclyl" means a monocyclic or polycyclic heterocyclic ring (typically having 3- to 14-members) which is either a saturated ring or an unsaturated non-aromatic ring. A 3-membered heterocyclyl can contain up to 3 heteroatoms, and a 4- to 14-membered heterocyclyl can contain from 1 to about 8 heteroatoms. Each heteroatom is independently selected from nitrogen, which can be quaternized; oxygen; and sulfur, including sulfoxide and sulfone. The heterocyclyl may be attached via any heteroatom or carbon atom. Representative heterocyclyls include morpholinyl, thiomorpholinyl, pyrrolidinonyl, pyrrolidinyl, piperidinyl, piperazinyl, hydantoinyl, valerolactamyl, oxiranyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, 4H-pyranyl, tetrahydropyridinyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, and the like. A heteroatom may be substituted with a protecting group known to those of ordinary skill in the art, for example, the hydrogen on a nitrogen may be substituted with a tert-butoxycarbonyl group. Furthermore, the heterocyclyl may be optionally substituted with one or more substituents (including without limitation a halo, an alkyl, a haloalkyl, or aryl). Only stable isomers of such substituted heterocyclic groups are contemplated in this definition.

As used herein, the term "heteroaromatic" or "heteroaryl" means a monocyclic or polycyclic heteroaromatic ring (or radical thereof) comprising carbon atom ring members and one or more heteroatom ring members (such as, for example, oxygen, sulfur or nitrogen). Typically, the heteroaromatic ring has from 5 to about 14 ring members in which at least 1 ring member is a heteroatom selected from oxygen, sulfur and nitrogen. In another embodiment, the heteroaromatic ring is a 5 or 6 membered ring and may contain from 1 to about 4 heteroatoms. In another embodiment, the heteroaromatic ring system has a 7 to 14 ring members and may contain from 1 to about 7 heteroatoms. Representative heteroaryls include pyridyl, furyl, thienyl, pyrrolyl, oxazolyl, imidazolyl, indoliziny, thiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, triazolyl, pyridinyl, thiadiazolyl, pyrazinyl, quinolyl, isoquinolyl, indazolyl, benzoxazolyl, benzofuryl, benzothiazolyl, indoliziny, imidazopyridinyl, isothiazolyl, tetrazolyl, benzimidazolyl,

benzoxazolyl, benzothiazolyl, benzothiadiazolyl, benzoxadiazolyl, indolyl, tetrahydroindolyl, azaindolyl, imidazopyridyl, quizaolinyl, purinyl, pyrrolo[2,3]pyrimidyl, pyrazolo[3,4]pyrimidyl or benzo(b)thienyl and the like. Heteroaryl groups may be optionally substituted with one or more substituents

5

A heteroaralkyl group refers to a heteroaryl group that is attached to another moiety via an alkylene linker. Heteroaralkyl groups can be substituted or unsubstituted with one or more substituents.

10 As used herein, the term "N-containing 6-membered aromatic ring" means an optionally substituted 6-membered aromatic ring that contains one or more N atoms. In one embodiment, the ring contains one N atom. In another embodiment, the ring contains two N atoms.

15 As used herein, the term "halogen" or "halo" means -F, -Cl, -Br or -I.

As used herein, the term "haloalkyl" means an alkyl group in which one or more -H is replaced with a halo group. Examples of haloalkyl groups include -CF₃, -CHF₂, -CCl₃, -CH₂CH₂Br, -CH₂CH(CH₂CH₂Br)CH₃, -CHICH₃, and the
20 like.

As used herein, the term "haloalkoxy" means an alkoxy group in which one or more -H is replaced with a halo group. Examples of haloalkoxy groups include -OCF₃ and -OCHF₂.

25 The terms "bioisostere" and "bioisosteric replacement" have the same meanings as those generally recognized in the art. Bioisosteres are atoms, ions, or molecules in which the peripheral layers of electrons can be considered substantially identical. The term bioisostere is usually used to mean a portion of an overall molecule, as opposed to the entire molecule itself. Bioisosteric replacement involves using one bioisostere to replace another with the
30 expectation of maintaining or slightly modifying the biological activity of the first bioisostere. The bioisosteres in this case are thus atoms or groups of atoms having similar size, shape and electron density. Preferred bioisosteres of esters, amides or carboxylic acids are compounds containing two sites for hydrogen bond acceptance. In one embodiment, the ester, amide or carboxylic acid bioisostere is a 5-membered monocyclic heteroaryl ring, such as an optionally
35 substituted 1H-imidazolyl, an optionally substituted oxazolyl, 1H-tetrazolyl, [1,2,4]triazolyl, or an optionally substituted [1,2,4]oxadiazolyl.

As used herein, the terms “subject”, “patient” and “animal”, are used interchangeably and include, but are not limited to, a cow, monkey, horse, sheep, pig, mini pig, chicken, turkey, quail, cat, dog, mouse, rat, rabbit, guinea pig and human. The preferred subject, patient or animal is a human.

5

As used herein, the term “lower” refers to a group having up to four carbon atoms. For example, a “lower alkyl” refers to an alkyl radical having from 1 to 4 carbon atoms, and a “lower alkenyl” or “lower alkynyl” refers to an alkenyl or alkynyl radical having from 2 to 4 carbon atoms, respectively. A lower alkoxy or a lower alkylsulfanyl refers to an alkoxy or an alkylsulfanyl having from 1 to 4 carbon atoms. Lower substituents are typically preferred.

10

Where a particular substituent, such as an alkyl substituent, occurs multiple times in a given structure or moiety, the identity of the substituent is independent in each case and may be the same as or different from other occurrences of that substituent in the structure or moiety.

15

Furthermore, individual substituents in the specific embodiments and exemplary compounds of this invention are preferred in combination with other such substituents in the compounds of this invention, even if such individual substituents are not expressly noted as being preferred or not expressly shown in combination with other substituents.

20

The compounds of the invention are defined herein by their chemical structures and/or chemical names. Where a compound is referred to by both a chemical structure and a chemical name, and the chemical structure and chemical name conflict, the chemical structure is determinative of the compound's identity.

25

Suitable substituents for an alkyl, alkoxy, alkylsulfanyl, alkylamino, dialkylamino, alkylene, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, aralkyl, heteroaryl, and heteroaralkyl groups include any substituent which will form a stable compound of the invention. Examples of substituents for an alkyl, alkoxy, alkylsulfanyl, alkylamino, dialkylamino, alkylene, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, aralkyl, heteroaryl, and heteroaralkyl include an alkyl, an alkoxy, an alkylsulfanyl, an alkylamino, a dialkylamino, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, a heterocyclyl, an aryl, a heteroaryl, an aralkyl, a heteraralkyl, a haloalkyl, $-C(O)NR_{23}R_{24}$, $-NR_{25}C(O)R_{26}$, halo, $-OR_{25}$, cyano, nitro,

30

haloalkoxy, $-C(O)R_{25}$, $-NR_{23}R_{24}$, $-SR_{25}$, $-C(O)OR_{25}$, $-OC(O)R_{25}$, $-NR_{25}C(O)NR_{23}R_{24}$, $-OC(O)NR_{23}R_{24}$, $-NR_{25}C(O)OR_{26}$, $-S(O)_pR_{25}$, or $-S(O)_pNR_{23}R_{24}$, wherein R_{23} and R_{24} , for each occurrence are, independently, H, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, a heterocyclyl, an aryl, a heteroaryl, an aralkyl, or a heteraralkyl; or R_{23} and R_{24}

35

taken together with the nitrogen to which they are attached is a heterocyclyl or a heteroaryl; and R₂₅ and R₂₆ for each occurrence are, independently, H, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, a heterocyclyl, an aryl, a heteroaryl, an aralkyl, or a heteraralkyl;

- 5 In addition, alkyl, cycloalkyl, alkylene, a heterocyclyl, and any saturated portion of a alkenyl, cycloalkenyl, alkynyl, aralkyl, and heteroaralkyl groups, may also be substituted with =O, =S, =N-R₂₂.

10 When a heterocyclyl, heteroaryl, or heteroaralkyl group contains a nitrogen atom, it may be substituted or unsubstituted. When a nitrogen atom in the aromatic ring of a heteroaryl group has a substituent the nitrogen may be a quaternary nitrogen.

15 Choices and combinations of substituents and variables envisioned by this invention are only those that result in the formation of stable compounds. The term "stable", as used herein, refers to compounds which possess stability sufficient to allow manufacture and which maintains the integrity of the compound for a sufficient period of time to be useful for the purposes detailed herein (e.g., therapeutic or prophylactic administration to a subject). Typically, such compounds are stable at a temperature of 40°C or less, in the absence of excessive moisture, for at least one week. Such choices and combinations will be apparent to
20 those of ordinary skill in the art and may be determined without undue experimentation.

Unless indicated otherwise, the compounds of the invention containing reactive functional groups (such as, without limitation, carboxy, hydroxy, and amino moieties) also include protected derivatives thereof. "Protected derivatives" are those compounds in which a
25 reactive site or sites are blocked with one or more protecting groups. Suitable protecting groups for carboxy moieties include benzyl, tert-butyl, and the like. Suitable protecting groups for amino and amido groups include acetyl, tert-butoxycarbonyl, benzyloxycarbonyl, and the like. Suitable protecting groups for hydroxy include benzyl, trimethyl silyl (TMS) and the like. Other suitable protecting groups are well known to those of ordinary skill in the
30 art and include those found in T. W. Greene, *Protecting Groups in Organic Synthesis*, John Wiley & Sons, Inc. 1981, the entire teachings of which are incorporated herein by reference.

As used herein, the term "compound(s) of this invention" and similar terms refers to a
35 compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof and also include protected derivatives thereof.

As used herein and unless otherwise indicated, the term "prodrug" means a derivative of a compound that can hydrolyze, oxidize, or otherwise react under biological conditions (*in vitro* or *in vivo*) to provide a compound of this invention. Prodrugs may only become active upon such reaction under biological conditions, but they may have activity in their unreacted forms.

5 Examples of prodrugs contemplated in this invention include, but are not limited to, analogs or derivatives of compounds of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB),, or Table 1 that comprise biohydrolyzable moieties such as biohydrolyzable amides, biohydrolyzable esters, biohydrolyzable carbamates, biohydrolyzable carbonates, biohydrolyzable ureides, and biohydrolyzable phosphate
10 analogues. Other examples of prodrugs include derivatives of compounds of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB),, or of Table 1 that comprise -NO, -NO₂, -ONO, or -ONO₂ moieties. Prodrugs can typically be prepared using well-known methods, such as those described by 1 BURGER'S MEDICINAL CHEMISTRY AND DRUG DISCOVERY (1995) 172-178, 949-982 (Manfred E. Wolff ed., 5th ed), the entire
15 teachings of which are incorporated herein by reference.

As used herein and unless otherwise indicated, the terms "biohydrolyzable amide", "biohydrolyzable ester", "biohydrolyzable carbamate", "biohydrolyzable carbonate", "biohydrolyzable ureide" and "biohydrolyzable phosphate analogue" mean an amide, ester,
20 carbamate, carbonate, ureide, or phosphate analogue, respectively, that either: 1) does not destroy the biological activity of the compound and confers upon that compound advantageous properties *in vivo*, such as uptake, duration of action, or onset of action; or 2) is itself biologically inactive but is converted *in vivo* to a biologically active compound. Examples of biohydrolyzable amides include, but are not limited to, lower alkyl amides,
25 α -amino acid amides, alkoxyacyl amides, and alkylaminoalkylcarbonyl amides. Examples of biohydrolyzable esters include, but are not limited to, lower alkyl esters, alkoxyacyloxy esters, alkyl acylamino alkyl esters, and choline esters. Examples of biohydrolyzable carbamates include, but are not limited to, lower alkylamines, substituted ethylenediamines, aminoacids, hydroxyalkylamines, heterocyclic and heteroaromatic amines, and polyether
30 amines.

As used herein, the term "pharmaceutically acceptable salt," is a salt formed from an acid and a basic group of one of the compounds of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or of Table 1. Illustrative salts include, but are not limited, to
35 sulfate, citrate, acetate, oxalate, chloride, bromide, iodide, nitrate, bisulfate, phosphate, acid phosphate, isonicotinate, lactate, salicylate, acid citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucaronate,

saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate, and pamoate (*i.e.*,

1,1'-methylene-bis-(2-hydroxy-3-naphthoate)) salts. The term "pharmaceutically acceptable salt" also refers to a salt prepared from a compound of any one of formulas (I), (IIA), (IIB),

5 (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1 having an acidic functional group, such as a carboxylic acid functional group, and a pharmaceutically acceptable inorganic or organic base. Suitable bases include, but are not limited to, hydroxides of alkali metals such as sodium, potassium, and lithium; hydroxides of alkaline earth metal such as calcium and magnesium; hydroxides of other metals, such as aluminum and zinc; ammonia, and organic

10 amines, such as unsubstituted or hydroxy-substituted mono-, di-, or trialkylamines; dicyclohexylamine; tributyl amine; pyridine; N-methyl,N-ethylamine; diethylamine; triethylamine; mono-, bis-, or tris-(2-hydroxy-lower alkyl amines), such as mono-, bis-, or tris-(2-hydroxyethyl)- amine, 2-hydroxy-tert-butylamine, or

tris-(hydroxymethyl)methylamine, N, N,-di-lower alkyl-N-(hydroxy lower alkyl)-amines, such as N,N-dimethyl-N-(2-hydroxyethyl)- amine, or tri-(2-hydroxyethyl)amine;

15 N-methyl-D-glucamine; and amino acids such as arginine, lysine, and the like. The term "pharmaceutically acceptable salt" also refers to a salt prepared from a compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1 having a basic functional group, such as an amino functional group, and a pharmaceutically acceptable

20 inorganic or organic acid. Suitable acids include, but are not limited to, hydrogen sulfate, citric acid, acetic acid, oxalic acid, hydrochloric acid, hydrogen bromide, hydrogen iodide, nitric acid, phosphoric acid, isonicotinic acid, lactic acid, salicylic acid, tartaric acid, ascorbic acid, succinic acid, maleic acid, besylic acid, fumaric acid, gluconic acid, glucaronic acid, saccharic acid, formic acid, benzoic acid, glutamic acid, methanesulfonic acid, ethanesulfonic

25 acid, benzenesulfonic acid, and *p*-toluenesulfonic acid.

When a disclosed compound is named or depicted by structure, it is to be understood that solvates (e.g., hydrates) of the compound or its pharmaceutically acceptable salts are also included. "Solvates" refer to crystalline forms wherein solvent molecules are incorporated

30 into the crystal lattice during crystallization. Solvate may include water or nonaqueous solvents such as ethanol, isopropanol, DMSO, acetic acid, ethanalamine, and EtOAc. Solvates, wherein water is the solvent molecule incorporated into the crystal lattice, are typically referred to as "hydrates". Hydrates include a stoichiometric or non-stoichiometric amount of water bound by non-covalent intermolecular forces.

35 When a disclosed compound is named or depicted by structure, it is to be understood that the compound, including solvates thereof, may exist in crystalline forms, non-crystalline forms or

a mixture thereof. The compounds or solvates may also exhibit polymorphism (i.e. the capacity to occur in different crystalline forms). These different crystalline forms are typically known as “polymorphs.” It is to be understood that when named or depicted by structure, the disclosed compounds and solvates (e.g., hydrates) also include all polymorphs thereof. As
5 used herein, the term “polymorph” means solid crystalline forms of a compound of the present invention or complex thereof. Different polymorphs of the same compound can exhibit different physical, chemical and/or spectroscopic properties. Different physical properties include, but are not limited to stability (e.g., to heat or light), compressibility and density (important in formulation and product manufacturing), and dissolution rates (which
10 can affect bioavailability). Differences in stability can result from changes in chemical reactivity (e.g., differential oxidation, such that a dosage form discolors more rapidly when comprised of one polymorph than when comprised of another polymorph) or mechanical characteristics (e.g., tablets crumble on storage as a kinetically favored polymorph converts to thermodynamically more stable polymorph) or both (e.g., tablets of one polymorph are more
15 susceptible to breakdown at high humidity). Different physical properties of polymorphs can affect their processing. For example, one polymorph might be more likely to form solvates or might be more difficult to filter or wash free of impurities than another due to, for example, the shape or size distribution of particles of it. In addition, one polymorph may spontaneously convert to another polymorph under certain conditions.

20

When a disclosed compound is named or depicted by structure, it is to be understood that clathrates (“inclusion compounds”) of the compound or its pharmaceutically acceptable salts, solvates or polymorphs are also included. As used herein, the term “clathrate” means a
25 compound of the present invention or a salt thereof in the form of a crystal lattice that contains spaces (e.g., channels) that have a guest molecule (e.g., a solvent or water) trapped within.

25

As used herein, the term “asthma” means a pulmonary disease, disorder or condition characterized by reversible airway obstruction, airway inflammation, and increased airway
30 responsiveness to a variety of stimuli.

30

“Immunosuppression” refers to impairment of any component of the immune system resulting in decreased immune function. This impairment may be measured by any conventional means including whole blood assays of lymphocyte function, detection of lymphocyte
35 proliferation and assessment of the expression of T cell surface antigens. The antisheep red blood cell (SRBC) primary (IgM) antibody response assay (usually referred to as the plaque assay) is one specific method. This and other methods are described in Luster, M.I., Portier,

35

C., Pait, D.G., White, K.L., Jr., Gennings, C., Munson, A.E., and Rosenthal, G.J. (1992). "Risk Assessment in Immunotoxicology I: Sensitivity and Predictability of Immune Tests." *Fundam. Appl. Toxicol.*, 18, 200-210. Measuring the immune response to a T-cell dependent immunogen is another particularly useful assay (Dean, J.H., House, R.V., and Luster, M.I. (2001). "Immunotoxicology: Effects of, and Responses to, Drugs and Chemicals." In *Principles and Methods of Toxicology: Fourth Edition* (A.W. Hayes, Ed.), pp. 1415-1450, Taylor & Francis, Philadelphia, Pennsylvania).

The compounds of this invention can be used to treat subjects with immune disorders. As used herein, the term "immune disorder" and like terms means a disease, disorder or condition caused by the immune system of an animal, including autoimmune disorders. Immune disorders include those diseases, disorders or conditions that have an immune component and those that are substantially or entirely immune system-mediated. Autoimmune disorders are those wherein the animal's own immune system mistakenly attacks itself, thereby targeting the cells, tissues, and/or organs of the animal's own body. For example, the autoimmune reaction is directed against the nervous system in multiple sclerosis and the gut in Crohn's disease. In other autoimmune disorders such as systemic lupus erythematosus (lupus), affected tissues and organs may vary among individuals with the same disease. One person with lupus may have affected skin and joints whereas another may have affected skin, kidney, and lungs. Ultimately, damage to certain tissues by the immune system may be permanent, as with destruction of insulin-producing cells of the pancreas in Type 1 diabetes mellitus. Specific autoimmune disorders that may be ameliorated using the compounds and methods of this invention include without limitation, autoimmune disorders of the nervous system (e.g., multiple sclerosis, myasthenia gravis, autoimmune neuropathies such as Guillain-Barré, and autoimmune uveitis), autoimmune disorders of the blood (e.g., autoimmune hemolytic anemia, pernicious anemia, and autoimmune thrombocytopenia), autoimmune disorders of the blood vessels (e.g., temporal arteritis, anti-phospholipid syndrome, vasculitides such as Wegener's granulomatosis, and Behcet's disease), autoimmune disorders of the skin (e.g., psoriasis, dermatitis herpetiformis, pemphigus vulgaris, and vitiligo), autoimmune disorders of the gastrointestinal system (e.g., Crohn's disease, ulcerative colitis, primary biliary cirrhosis, and autoimmune hepatitis), autoimmune disorders of the endocrine glands (e.g., Type 1 or immune-mediated diabetes mellitus, Grave's disease. Hashimoto's thyroiditis, autoimmune oophoritis and orchitis, and autoimmune disorder of the adrenal gland); and autoimmune disorders of multiple organs (including connective tissue and musculoskeletal system diseases) (e.g., rheumatoid arthritis, systemic lupus erythematosus, scleroderma, polymyositis, dermatomyositis, spondyloarthropathies such as ankylosing spondylitis, and Sjogren's syndrome). In addition, other immune system mediated diseases,

such as graft-versus-host disease and allergic disorders, are also included in the definition of immune disorders herein. Because a number of immune disorders are caused by inflammation, there is some overlap between disorders that are considered immune disorders and inflammatory disorders. For the purpose of this invention, in the case of such an overlapping disorder, it may be considered either an immune disorder or an inflammatory disorder. "Treatment of an immune disorder" herein refers to administering a compound or a composition of the invention to a subject, who has an immune disorder, a symptom of such a disease or a predisposition towards such a disease, with the purpose to cure, relieve, alter, affect, or prevent the autoimmune disorder, the symptom of it, or the predisposition towards it.

As used herein, the term "allergic disorder" means a disease, condition or disorder associated with an allergic response against normally innocuous substances. These substances may be found in the environment (such as indoor air pollutants and aeroallergens) or they may be non-environmental (such as those causing dermatological or food allergies). Allergens can enter the body through a number of routes, including by inhalation, ingestion, contact with the skin or injection (including by insect sting). Many allergic disorders are linked to atopy, a predisposition to generate the allergic antibody IgE. Because IgE is able to sensitize mast cells anywhere in the body, atopic individuals often express disease in more than one organ. For the purpose of this invention, allergic disorders include any hypersensitivity that occurs upon re-exposure to the sensitizing allergen, which in turn causes the release of inflammatory mediators. Allergic disorders include without limitation, allergic rhinitis (e.g., hay fever), sinusitis, rhinosinusitis, chronic or recurrent otitis media, drug reactions, insect sting reactions, latex reactions, conjunctivitis, urticaria, anaphylaxis and anaphylactoid reactions, atopic dermatitis, asthma and food allergies.

The compounds of this invention can be used to prevent or to treat subjects with inflammatory disorders. As used herein, an "inflammatory disorder" means a disease, disorder or condition characterized by inflammation of body tissue or having an inflammatory component. These include local inflammatory responses and systemic inflammation. Examples of such inflammatory disorders include: transplant rejection, including skin graft rejection; chronic inflammatory disorders of the joints, including arthritis, rheumatoid arthritis, osteoarthritis and bone diseases associated with increased bone resorption; inflammatory bowel diseases such as ileitis, ulcerative colitis, Barrett's syndrome, and Crohn's disease; inflammatory lung disorders such as asthma, adult respiratory distress syndrome, and chronic obstructive airway disease; inflammatory disorders of the eye including corneal dystrophy, trachoma, onchocerciasis, uveitis, sympathetic ophthalmitis and endophthalmitis; chronic inflammatory

disorders of the gums, including gingivitis and periodontitis; tuberculosis; leprosy; inflammatory diseases of the kidney including uremic complications, glomerulonephritis and nephrosis; inflammatory disorders of the skin including sclerodermatitis, psoriasis and eczema; inflammatory diseases of the central nervous system, including chronic
5 demyelinating diseases of the nervous system, multiple sclerosis, AIDS-related neurodegeneration and Alzheimer's disease, infectious meningitis, encephalomyelitis, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis and viral or autoimmune encephalitis; autoimmune disorders, immune-complex vasculitis, systemic lupus and erythematoses; systemic lupus erythematosus (SLE); and inflammatory diseases of the
10 heart such as cardiomyopathy, ischemic heart disease hypercholesterolemia, atherosclerosis; as well as various other diseases with significant inflammatory components, including preeclampsia; chronic liver failure, brain and spinal cord trauma, and cancer. There may also be a systemic inflammation of the body, exemplified by gram-positive or gram negative shock, hemorrhagic or anaphylactic shock, or shock induced by cancer chemotherapy in
15 response to pro-inflammatory cytokines, e.g., shock associated with pro-inflammatory cytokines. Such shock can be induced, e.g., by a chemotherapeutic agent used in cancer chemotherapy. "Treatment of an inflammatory disorder" herein refers to administering a compound or a composition of the invention to a subject, who has an inflammatory disorder, a symptom of such a disorder or a predisposition towards such a disorder, with the purpose to
20 cure, relieve, alter, affect, or prevent the inflammatory disorder, the symptom of it, or the predisposition towards it.

An "effective amount" is the quantity of compound in which a beneficial outcome is achieved when the compound is administered to a subject or alternatively, the quantity of compound
25 that possess a desired activity in-vivo or in-vitro. In the case of inflammatory disorders and autoimmune disorders, a beneficial clinical outcome includes reduction in the extent or severity of the symptoms associated with the disease or disorder and/or an increase in the longevity and/or quality of life of the subject compared with the absence of the treatment. The precise amount of compound administered to a subject will depend on the type and severity of
30 the disease or condition and on the characteristics of the subject, such as general health, age, sex, body weight and tolerance to drugs. It will also depend on the degree, severity and type of inflammatory disorder, autoimmune disorder, allergic disorder, or the degree of immunosuppression sought. The skilled artisan will be able to determine appropriate dosages depending on these and other factors. Effective amounts of the disclosed compounds
35 typically range between about 1 mg/m² per day and about 10 grams/m² per day, and preferably between 10 mg/m² per day and about 1 gram/m².

The compounds of the invention may contain one or more chiral centers and/or double bonds and, therefore, may exist as stereoisomers, such as double-bond isomers (*i.e.*, geometric isomers), enantiomers, or diastereomers. According to this invention, the chemical structures depicted herein, including the compounds of this invention, encompass all of the

5 corresponding compounds' enantiomers and stereoisomers, that is, both the stereomerically pure form (*e.g.*, geometrically pure, enantiomerically pure, or diastereomerically pure) and enantiomeric, diastereomeric, and geometric isomeric mixtures. In some cases, one enantiomer, diastereomer, or geometric isomer will possess superior activity or an improved toxicity or kinetic profile compared to others. In those cases, such enantiomers,
10 diastereomers, and geometric isomers of a compound of this invention are preferred.

The term "inhibit production of IL-2" and like terms means inhibiting IL-2 synthesis (*e.g.* by inhibiting transcription (mRNA expression), or translation (protein expression)) and/or inhibiting IL-2 secretion in a cell that has the ability to produce and/or secrete IL-2 (*e.g.*, T
15 lymphocyte). Likewise, the term "inhibiting production of IL-4, IL-5, IL-13, GM-CSF, TNF- α or INF- γ means inhibiting the synthesis (*e.g.* by inhibiting transcription, or translation) and/or inhibiting the secretion in a cell that has the ability to produce and/or secrete these cytokines.

20 As used herein, a composition that "substantially" comprises a compound means that the composition contains more than about 80% by weight, more preferably more than about 90% by weight, even more preferably more than about 95% by weight, and most preferably more than about 97% by weight of the compound.

25 As used herein, a composition that is "substantially free" of a compound means that the composition contains less than about 20% by weight, more preferably less than about 10% by weight, even more preferably less than about 5% by weight, and most preferably less than about 3% by weight of the compound.

30 As used herein, a reaction that is "substantially complete" means that the reaction contains more than about 80% by weight of the desired product, more preferably more than about 90% by weight of the desired product, even more preferably more than about 95% by weight of the desired product, and most preferably more than about 97% by weight of the desired product.

35 As used herein, a racemic mixture means about 50% of one enantiomer and about 50% of is corresponding enantiomer relative to all chiral centers in the molecule. The invention encompasses all enantiomerically-pure, enantiomerically-enriched, diastereomerically pure,

diastereomerically enriched, and racemic mixtures of the compounds of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1.

5 Enantiomeric and diastereomeric mixtures can be resolved into their component enantiomers or stereoisomers by well known methods, such as chiral-phase gas chromatography, chiral-phase high performance liquid chromatography, crystallizing the compound as a chiral salt complex, or crystallizing the compound in a chiral solvent. Enantiomers and diastereomers can also be obtained from diastereomerically- or enantiomerically-pure intermediates, reagents, and catalysts by well known asymmetric synthetic methods.

10 When administered to a patient, *e.g.*, to a non-human animal for veterinary use or for improvement of livestock, or to a human for clinical use, the compounds of the invention are typically administered in isolated form or as the isolated form in a pharmaceutical composition. As used herein, "isolated" means that the compounds of the invention are
15 separated from other components of either (a) a natural source, such as a plant or cell, preferably bacterial culture, or (b) a synthetic organic chemical reaction mixture. Preferably, via conventional techniques, the compounds of the invention are purified. As used herein, "purified" means that when isolated, the isolate contains at least 95%, preferably at least 98%, of a single compound of the invention by weight of the isolate.


20 Only those choices and combinations of substituents that result in a stable structure are contemplated. Such choices and combinations will be apparent to those of ordinary skill in the art and may be determined without undue experimentation.

25 The invention can be understood more fully by reference to the following detailed description and illustrative examples, which are intended to exemplify non-limiting embodiments of the invention.

30 **SPECIFIC EMBODIMENTS**

The invention relates to compounds and pharmaceutical compositions that are particularly useful for immunosuppression or to treat or prevent inflammatory conditions, immune disorders, and allergic disorders.

35 In one embodiment, the invention relates to compounds of structural formula (I):



(I)

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

X_1 and X_2 are, independently, CH or N;

5 one of X_3 or X_6 is NR_{29} , and the other is CH_2 , CHR_2 , or $C(R_2)_2$;

L is a linker selected from the group consisting of $-NHCH_2-$, $-CH_2NH-$, $-NR_5-C(O)-$, $-C(O)-NR_5-$, $-NR_5-C(S)-$, $-C(S)-NR_5-$, $-NR_5-C(NR_8)-$, $-C(NR_8)-NR_5-$, $-NR_5C(S)NR_5-$, $-NR_5C(NR_8)NR_5-$, or $-NR_5CR^aR^bNR_5-$;

Y is a substituted phenyl, provided that Y is not substituted with $-NHC(O)R_{30}$;

10 R_1 is H or a substituent;

R_2 , for each occurrence is, independently, a substituent, provided that R_2 is not $=O$ or $=C(R_{28})_2$;

R_4 is a substituent;

15 R^a and R^b , for each occurrence, are independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, cyano, nitro, halo, $-OR_5$, $-SR_5$, $-NR_6R_7$, $-C(O)NR_6R_7$, $-NR_5C(O)R_5$, $-C(O)R_5$, $-C(O)OR_5$, $-OC(O)R_5$, $-C(O)SR_5$, $-SC(O)R_5$, $-C(S)NR_6R_7$, $-NR_5C(S)R_5$, $-C(S)R_5$, $-C(S)OR_5$, $-OC(S)R_5$, $-C(S)SR_5$, $-SC(S)R_5$, $-C(NR_8)NR_6R_7$, $-NR_5C(NR_8)R_5$, $-C(NR_8)R_5$, $-C(NR_8)OR_5$, $-OC(NR_8)R_5$, $-C(NR_8)SR_5$, $-SC(NR_8)R_5$, $-OC(O)OR_5$, $-OC(O)NR_6R_7$, $-NR_5C(O)OR_5$, $-NR_5C(O)NR_6R_7$, $-SC(O)OR_5$, $-SC(O)NR_6R_7$, $-SC(O)SR_5$, $-NR_5C(O)SR_5$, $-OC(O)SR_5$, $-OC(S)OR_5$, $-OC(S)NR_6R_7$, $-NR_5C(S)OR_5$, $-NR_5C(S)NR_6R_7$, $-SC(S)OR_5$, $-SC(S)NR_6R_7$, $-SC(S)SR_5$, $-NR_5C(S)SR_5$, $-OC(S)SR_5$, $-OC(NR_8)OR_5$, $-OC(NR_8)NR_6R_7$, $-NR_5C(NR_8)OR_5$, $-NR_5C(NR_8)NR_6R_7$, $-SC(NR_8)OR_5$, $-SC(NR_8)NR_6R_7$, $-SC(NR_8)SR_5$, $-NR_5C(NR_8)SR_5$, or $-OC(NR_8)SR_5$;

20 R_5 , for each occurrence, is independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

R₆ and R₇, for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R₆ and R₇ taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

R₈, for each occurrence, is independently -H, a halo, an alkyl, -OR₅, -NR₆R₇, -C(O)R₅, -C(O)OR₅, or -C(O)NR₆R₇;

R₂₈, for each occurrence is, independently, H or a substituent;

R₂₉ is -C(O)R₃₀ or a N-containing 6-membered aromatic ring;

R₃₀ is -H, -OR₅, -SR₅, -NR₆R₇, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

m is 0, 1, or 2; and

n is 0, 1 or 2.

In one aspect, the compound of formula (I) is not:

5-[4-(2,6-difluoro-benzoylamino)-phenyl]-4-methyl-2,4-dihydro-2H-pyridine-1-carboxylic acid ethyl ester;

Ethyl 3-(4-(2,6-difluorobenzamido)phenyl)-4-methyl-5,6-dihydropyridine-1(2H)-carboxylate;

Ethyl 4-(4-(2,6-difluorobenzamido)phenyl)-3-methyl-5,6-dihydropyridine-1(2H)-carboxylate;

2,6-Difluoro-N-[4-(1-methoxycarbonyl-4-thiazol-2-yl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-methoxycarbonyl-4-isopropyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-methoxycarbonyl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-allyloxycarbonyl-3-methyl-1,2,5,6-tetrahydro-pyridin-4-yl)-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-phenyloxycarbonyl-3-methyl-1,2,5,6-tetrahydro-pyridin-4-yl)-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-acetyl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-

benzamide;


2,6-Difluoro-N-[4-(1-cyclopropylcarbonyl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide;

2,6-Difluoro-N-{4-[1-(pyridine-3-carbonyl)-cyclopropanecarbonyl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl]-phenyl}-benzamide; or


2,6-Difluoro-N-[4-(1-pyrimidin-2-yl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide.

In another embodiment, the invention relates to compounds represented by formula (IIA) or

(IIB):



or



or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein


R₃₁ is halo, an optionally substituted alkyl, an optionally substituted alkenyl, an

optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

p is 1, 2, 3, 4 or 5; and

R₁, R₃₀, L, X₁, and X₂ are defined as for formula (I).


In another embodiment, the invention relates to compounds represented by formula (IIIA) or (IIB):



or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein R₁, R₃₀, L, X₁, and X₂ are defined as for formula (I) and R₃₁ is defined as for formulas (IIA) and (IIB).

5

In another embodiment, the invention relates to compounds represented by formula (IVA) or (IVB):




or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein each of X₇, X₈, X₉, X₁₀, and X₁₁ are independently CH, CR₂, or N, provided that at least one of X₇, X₈, X₉, X₁₀, or X₁₁ is N;

5 R₃₁ is halo, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

10 p is 1, 2, 3, 4 or 5; and

R₁, L, X₁, and X₂ are defined as for formula (I)

In another embodiment, the invention relates to compounds represented by formula (VA) or (VB):



15

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein R₁, L, X₁, and X₂ are defined as for formula (I) and X₇, X₈, X₉, X₁₀, and X₁₁ are defined as for formulas (IVA) and (IVB).

20

In some embodiments of the compounds represented by formula (I), (IIA), (IIB), (IIIA),

(IIIB), (IVA), (IVB), (VA), or (VB), R_1 is selected from the group consisting of a halo, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl. In one aspect, R_1 is selected from the group consisting of halo or an optionally substituted lower alkyl. In one aspect, R_1 is selected from the group consisting of $-\text{CH}_2\text{OAc}$, CH_3 , or Cl .

In some embodiments of the compounds represented by formula (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), L is $-\text{NHCH}_2-$, $-\text{CH}_2\text{NH}-$, $-\text{NH-C(O)-}$, or $-\text{C(O)-NH-}$. In one aspect, L is $-\text{NHC(O)-}$.

In some embodiments of the compounds represented by formula (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), X_1 and X_2 are both CH .

In some embodiments of the compounds represented by formula (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), X_1 and X_2 are both N .

In some embodiments of the compounds represented by formula (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), X_1 is N and X_2 is CH .

In some embodiments of the compounds represented by formula (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), X_2 is N and X_1 is CH .

In some embodiments of the compounds represented by formula (I), Y is substituted with one to two substituents. In one aspect, the one to two substituents are each independently a lower alkyl or a halo. In another aspect, Y is a difluorophenyl. In a further aspect, Y is 2,6-difluorophenyl.

In some embodiments of the compounds represented by formula (I), X_3 is CH_2 and X_6 is NR_{29} .

In some embodiments of the compounds represented by formula (I), X_6 is CH_2 and X_3 is NR_{29} .

In some embodiments of the compounds represented by formula (I), m is 0.

In some embodiments of the compounds represented by formula (I), Y is a difluorophenyl;

L is -NHC(O)-; R₁ is selected from the group consisting of halo or an optionally substituted lower alkyl; and m is 0.

5 In some embodiments of the compounds represented by formula (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), each R₃₁ is independently a lower alkyl or a halo. In one aspect, R₃₁ is halo. In one aspect, R₃₁ is F.

10 In some embodiments of the compounds represented by formula (IIA), (IIB), (IVA), or (IVB), p is 1 or 2. In one aspect, each R₃₁ is independently a lower alkyl or a halo. In one aspect, R₃₁ is halo. In one aspect, R₃₁ is F.

15 In some embodiments of the compounds represented by formula (IIIA), (IIIB), (VA), or (VB), each R₃₁ is independently a lower alkyl or a halo. In one aspect, each R₃₁ is halo. In another aspect, each R₃₁ is F.

20 In some embodiments of the compounds represented by formula (IIA), (IIB), (IIIA), or (IIIB), R₃₀ is selected from the group consisting of -OR₅, optionally substituted alkyl, optionally substituted aryl, and optionally substituted heteroaryl. In one aspect, R₃₀ is selected from the group consisting of -OR₅ and optionally substituted heteroaryl. In another aspect, R₃₀ is -OR₅ and R₅ is a lower alkyl. In a further aspect, R₃₀ is a pyridine ring.

In some embodiments of the compounds represented by formula (IVA), (IVB), (VA), or (VB), two of X₇, X₈, X₉, X₁₀, and X₁₁ are N. In one aspect, X₇ and X₁₁ are both N.

25 In some embodiments of the compounds represented by formula (IVA), (IVB), (VA), or (VB), one of X₇, X₈, X₉, X₁₀, and X₁₁ is N.

In some embodiments of the compounds represented by formula (IIIA) or (IIIB), each R₃₁ is halo;

30 L is -NHC(O)-; R₁ is selected from the group consisting of halo or an optionally substituted lower alkyl; and R₃₀ is selected from the group consisting of -OR₅ and optionally substituted heteroaryl. In one aspect, each R₃₁ is F. In one aspect, X₁ and X₂ are both CH. In another aspect, X₁ and X₂ are both N. In a further aspect, X₁ is N and X₂ is CH.

35 In some embodiments of the compounds represented by formula (VA), or (VB), each R₃₁ is halo;
L is -NHC(O)-; R₁ is selected from the group consisting of halo or an optionally substituted

lower alkyl; and two of X₇, X₈, X₉, X₁₀, and X₁₁ are N. In one aspect, each R₃₁ is F. In one aspect, X₁ and X₂ are both CH. In another aspect, X₁ and X₂ are both N. In a further aspect, X₁ is N and X₂ is CH.

- 5 In one embodiment, the compounds of the invention are not compounds disclosed in U.S. Patent Application No.11/699,002.

In another embodiment, the invention relates to compounds selected from the group consisting of:

ethyl 3-(6-(2,6-difluorobenzamido)pyridin-3-yl)-4-methyl-5,6-dihydropyridine-1(2H)-carboxylate;
allyl 3-(acetoxymethyl)-4-(4-(2,6-difluorobenzamido)phenyl)-5,6-dihydropyridine-1(2H)-carboxylate;
2,6-difluoro-N-(5-(4-methyl-1-picolinoyl-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)benzamide hydrochloride;

ethyl 3-(acetoxymethyl)-4-(4-(2,6-difluorobenzamido)phenyl)-5,6-dihydropyridine-1(2H)-carboxylate
or

ethyl 4-chloro-3-(4-(2,6-difluorobenzamido)phenyl)-5,6-dihydropyridine-1(2H)-carboxylate;

- 10 or pharmaceutically acceptable salts, solvates, clathrates, or prodrugs thereof.

In another embodiment, the invention relates to 2,6-difluoro-N-(5-(4-methyl-1-(pyrimidin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide hydrochloride or pharmaceutically

- 15 acceptable salts, solvates, clathrates, or prodrugs thereof.

In another embodiment, the invention relates to

2,6-difluoro-N-(5-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)benzamide;
2,6-difluoro-N-(5-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)benzamide hydrochloride;

2,6-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide;

2,4-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide;

2,6-difluoro-N-(5-(4-methyl-1-(pyrazin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide;

N-(5-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide;

2,6-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide hydrochloride;

2-chloro-6-fluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide hydrochloride;

N-(5-(1-(3-chloropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide hydrochloride;

N-(5-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,4-difluorobenzamide;

2,6-difluoro-N-(5-(1-(3-fluoropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide hydrochloride;

2,4-difluoro-N-(5-(1-(3-fluoropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide hydrochloride;

N-(5-(1-(3-cyanopyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide;

N-(5-(1-(3,5-difluoropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide;

N-(5-(1-(3,5-dichloropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide;

N-(5-(1-(3-bromopyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide;

N-(4-(1-(3-aminopyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)-2,6-difluorobenzamide;

4-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)-N-(2,6-difluorobenzyl)aniline;

N-(4-(1-(3-cyanopyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)-2,6-difluorobenzamide;

2,6-difluoro-N-(4-(1-(3-fluoropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide;

N-(4-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)-2,6-difluorobenzamide; or

In another embodiment, the invention relates to

2,6-difluoro-N-(5-(4-methyl-1-(pyrimidin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide hydrochloride

2,6-difluoro-N-(5-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)benzamide;

2,6-difluoro-N-(5-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)benzamide hydrochloride;

2,6-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide;

2,4-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide;

2,6-difluoro-N-(5-(4-methyl-1-(pyrazin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide;

N-(5-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide;

2,6-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide hydrochloride;

2-chloro-6-fluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide hydrochloride;

N-(5-(1-(3-chloropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide hydrochloride;

N-(5-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,4-difluorobenzamide;

2,6-difluoro-N-(5-(1-(3-fluoropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide hydrochloride;

2,4-difluoro-N-(5-(1-(3-fluoropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide hydrochloride;

N-(5-(1-(3-cyanopyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide;

N-(5-(1-(3,5-difluoropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide;

N-(5-(1-(3,5-dichloropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide;

N-(5-(1-(3-bromopyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide;

N-(4-(1-(3-aminopyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)-2,6-difluorobenzamide;

4-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)-N-(2,6-difluorobenzyl)aniline;

N-(4-(1-(3-cyanopyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)-2,6-difluorobenzamide;

2,6-difluoro-N-(4-(1-(3-fluoropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide;

N-(4-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)-2,6-difluorobenzamide; or

2,6-difluoro-N-(4-(4-methyl-1-(pyrazin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide; or pharmaceutically acceptable salts, solvates, clathrates, or prodrugs thereof.

All of the features, specific embodiments and particular substituents disclosed herein may be combined in any combination. Each feature, embodiment or substituent disclosed in this specification may be replaced by an alternative feature, embodiment or substituent serving the same, equivalent, or similar purpose. In the case of chemical compounds, specific values for

variables (e.g., values shown in the exemplary compounds disclosed herein) in any chemical formula disclosed herein can be combined in any combination resulting in a stable structure. Furthermore, specific values (whether preferred or not) for substituents in one type of chemical structure may be combined with values for other substituents (whether preferred or not) in the same or different type of chemical structure. Thus, unless expressly stated otherwise, each feature, embodiment or substituent disclosed is only an example of a generic series of equivalent or similar features, embodiments or substituents.

In another embodiment, the invention relates to pharmaceutical compositions that comprise a compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, as an active ingredient, and a pharmaceutically acceptable carrier or vehicle. The compositions are useful for immunosuppression or to treat or prevent inflammatory conditions, allergic conditions or immune disorders.

In another embodiment, the invention relates to methods for immunosuppression or for treating or preventing inflammatory conditions, immune disorders, or allergic disorders in a patient in need thereof comprising administering an effective amount of a compound represented by any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

In another embodiment, the invention relates to methods for immunosuppression or for treating or preventing inflammatory conditions, immune disorders, or allergic disorders in a patient in need thereof comprising administering an effective amount of a pharmaceutical composition that comprises a compound represented by any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

In another embodiment, compounds of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, are particularly useful inhibiting immune cell (e.g., T-cells and/or B-cells) activation (e.g., activation in response to an antigen) and/or T cell and/or B cell proliferation. Indicators of immune cell activation include secretion of IL-2 by T cells, proliferation of T cells and/or B cells, and the like. In one embodiment, immune cell activation and/or T cell and/or B cell proliferation is inhibited in a mammal (e.g., a human), by administering to the mammal (e.g., human) a compound of any one of formulas (I), (IIA),

(IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

5 In another embodiment, compounds of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, can inhibit the production of certain cytokines that regulate immune cell activation. For example, compounds of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, can inhibit the production of IL-2, IL-4, IL-5, IL-13,
10 GM-CSF, IFN- γ , TNF- α and combinations thereof. In one embodiment, cytokine production is inhibited in a mammal (e.g., a human), by administering to the mammal (e.g., human) a compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

15 In another embodiment, compounds of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, can modulate the activity of one or more ion channel, such as CRAC ion channels, involved in activation of immune cells. In one embodiment, a compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1 can inhibit the influx of calcium ions into an immune cell (e.g., T cells, B
20 cells, and/or mast cells) by inhibiting the action of CRAC ion channels either directly or indirectly. In general, a decrease in I_{CRAC} current upon contacting a cell with a compound is one indicator that the compound inhibitions CRAC ion channels. I_{CRAC} current can be measured, for example, using a patch clamp technique, which is described in more detail in the examples below. In one embodiment, a compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1 modulates an ion channel in a mammal (e.g., a human). In one embodiment, the activity of one or more ion channels is inhibited in a mammal (e.g., a human), by administering to the mammal (e.g., human) a
25 compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.
30

In another embodiment, compounds of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, can inhibit degranulation of mast cell. Inhibition of mast cell
35 degranulation can determined as described in the experimental section herein or by any method known to those skilled in the art. In one embodiment, mast cell degranulation is inhibited in a mammal (e.g., a human), by administering to the mammal (e.g., human) a

compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.







EXEMPLARY COMPOUNDS OF THE INVENTION







5







Exemplary compounds of the invention are depicted in Table 1 below.







Table 1

Compound No.	Structure	Chemical Name
1		ethyl 3-(6-(2,6-difluorobenzamido)pyridin-3-yl)-4-methyl-5,6-dihydropyridine-1(2H)-carboxylate
2		2,6-difluoro-N-(5-(4-methyl-1-(pyrimidin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide hydrochloride
3		allyl 3-(acetoxymethyl)-4-(4-(2,6-difluorobenzamido)phenyl)-5,6-dihydropyridine-1(2H)-carboxylate
4		2,6-difluoro-N-(5-(4-methyl-1-(picolinoyl)-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)benzamide hydrochloride

5		ethyl 3-(acetoxymethyl)-4-(4-(2,6-difluorobenzamido)phenyl)-5,6-dihydropyridine-1(2H)-carboxylate
6		ethyl 4-chloro-3-(4-(2,6-difluorobenzamido)phenyl)-5,6-dihydropyridine-1(2H)-carboxylate
7		2,6-difluoro-N-(5-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)benzamide
8		2,6-difluoro-N-(5-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)benzamide hydrochloride
9		2,6-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide
10		2,4-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide

11		2,6-difluoro-N-(5-(4-methyl-1-(pyrazin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide
12		N-(5-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide
13		2,6-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide hydrochloride
14		2-chloro-6-fluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide hydrochloride
15		N-(5-(1-(3-chloropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide hydrochloride
16		N-(5-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,4-difluorobenzamide

17		2,6-difluoro-N-(5-(1-(3-fluoropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide hydrochloride
18		2,4-difluoro-N-(5-(1-(3-fluoropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide hydrochloride
19		N-(5-(1-(3-cyanopyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide
20		N-(5-(1-(3,5-difluoropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide
21		N-(5-(1-(3,5-dichloropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide
22		N-(5-(1-(3-bromopyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide

23		N-(4-(1-(3-aminopyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)-2,6-difluorobenzamide
24		4-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)-N-(2,6-difluorobenzyl)aniline
25		N-(4-(1-(3-cyanopyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)-2,6-difluorobenzamide
26		2,6-difluoro-N-(4-(1-(3-fluoropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide
27		N-(4-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)-2,6-difluorobenzamide
28		2,6-difluoro-N-(4-(4-methyl-1-(pyrazin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide

MECHANISM OF ACTION

Activation of T-lymphocytes in response to an antigen is dependent on calcium ion oscillations. Calcium ion oscillations in T-lymphocytes are triggered through stimulation of the T-cell antigen receptor, and involve calcium ion influx through the stored-operated Ca²⁺-release-activated Ca²⁺ (CRAC) channel. In addition, antigen induced degranulation of mast cells has also been shown to be initiated by calcium ion in flux. Although the molecular structure of the CRAC ion channel has not been identified, a detailed electrophysiological profile of the channel exist. Thus, modulation (e.g., inhibition) of CRAC ion channels can be measured by measuring modulation (e.g., inhibition) of the I_{CRAC} current. Calcium ion oscillations in T-cells have been implicated in the activation of several transcription factors (e.g., NFAT, Oct/Oap and NFκB) which are critical for T-cell activation (Lewis, *Biochemical Society Transactions* (2003), 31:925-929, the entire teachings of which are incorporated herein by reference). Without wishing to be bound by any theory, it is believed that because the compounds of the invention inhibit the activity of CRAC ion channels, they inhibit immune cell activation.

METHODS OF TREATMENT AND PREVENTION


In accordance with the invention, an effective amount of a compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, and prodrug thereof, or a pharmaceutical composition comprising a compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, and prodrug thereof, is administered to a patient in need of immunosuppression or in need of treatment or prevention of an inflammatory condition, an immune disorder, or an allergic disorder. Such patients may be treatment naïve or may experience partial or no response to conventional therapies.

Responsiveness to immunosuppression or of a particular inflammatory condition, immune disorder, or allergic disorder in a subject can be measured directly (e.g., measuring blood levels of inflammatory cytokines (such as IL-2, IL-4, IL-5, IL-13, GM-CSF, TNF-α, IFN-γ and the like) after administration of a compound of this invention), or can be inferred based on an understanding of disease etiology and progression. The compounds of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof can be assayed *in vitro* or *in vivo*, for the desired therapeutic or prophylactic activity, prior to use in humans. For example, known animal models of inflammatory conditions, immune disorders, or

allergic disorders can be used to demonstrate the safety and efficacy of compounds of this invention.

PREPARATION OF COMPOUNDS OF THE INVENTION

5 General Coupling Procedure for Boronic Esters




(X = halo or OTf)

Scheme I

- 10 To a solution of $Pd(PhCN)_2Cl_2$ (767mg, 2mmol) and 1,4-bis(diphenylphosphino)-butane (dppb) (1.28g, 3mmol) in 100 mL of toluene was added a boronic ester (XXXVII) (22.6 mmol) followed by an alkenyl halide or alkenyl triflate (XXXVI) (22.6 mmol), and sodium carbonate (3.4g, 22.6mmol), followed by 23 mL of water and 2 mL of methanol. This mixture was heated to 100°C and refluxed for 18 hours. After allowing cooling to room temperature,
- 15 the reaction was quenched with water and worked-up with ethyl acetate. The organic solution was then filtered through silica gel, evaporated in vacuo to give an amine derivative (XXXVIII) (see Scheme I) that was used without further purification in the next step.

20 General Procedure for amide coupling reaction



Scheme II

- 25 Amine derivative (XXXVIII) (19.2 mmol) was dissolved in methylene chloride and cooled to 0°C. Pyridine (3.87ml, 48 mmol) was added to the reaction mixture followed by an acyl chloride (XXXIX) (19.2 mmol). This mixture was allowed to warm to room temperature for 35 minutes at which time water was added and the reaction mixture was partitioned between the methylene chloride and water layers. The organic layer was collected and dried over a


drying agent such as NaSO₄, then concentrated. The product was purified using flash chromatography to give a compound of the invention (XL) (see Scheme II).

5 Compounds of the invention in which L is -NHC(S)- or -C(S)NH- can be prepared by treating compounds having an amide linker with Lawesson's reagent.

Compounds of the invention having -CH₂-NH- or -NH-CH₂- linkers can be prepared by contacting compounds having -NHC(S)- or -C(S)NH- linkers with Raney Ni. Alternatively, compounds of the invention having a -CH₂-NH- or -NH-CH₂- linker can be prepared by
 10 reducing a compound having a -C(O)-NH- or -NH-C(O)- linker, respectively, with, for example, sodium borohydride (see U.S. Patent Application No. 10/897,681, filed on July 22, 2004, the entire teachings of which are incorporated herein by reference).

Compounds of the invention having a thiourea linker (-NHC(S)NH-) can be prepared by
 15 treating compounds having a urea linker with Lawesson's reagent.

Compounds of the invention having an amide linker in which the amine group is attached to Y (-C(O)NH-) can be prepared by combining a halo-benzyl chloride or halo-pyridine
 20 carbonyl chloride (LII) with an amine derivative (XXVII) in pyridine to form a halo-amide intermediate (XXVIII). The reaction conditions are similar to those described for the amide coupling reaction above. A solution of Pd(PhCN)₂Cl₂ (767mg, 2mmol) and 1,4-bis(diphenylphosphino)-butane (dppb) (1.28g, 3mmol) in 100 mL of toluene is prepared. A
 cycloalkylene boronic acid ester (XXIX) is added to the solution, followed by the halo-amide intermediate (XXVIII) (22.6 mmol), and sodium carbonate (22.6mmol), followed by 23 mL
 25 of water and 2 mL of methanol. This mixture was heated to 100°C and refluxed for about 18 hours. After allowing cooling to room temperature, the reaction was quenched with water and worked-up with ethyl acetate to form a compound of the invention having a -C(O)NH- linker (XXX) (see Scheme VII).



Scheme VII

PHARMACEUTICAL COMPOSITIONS AND DOSAGE FORMS

Pharmaceutical compositions and dosage forms of the invention comprise one or more active ingredients in relative amounts and formulated in such a way that a given pharmaceutical composition or dosage form can be used for immunosuppression or to treat or prevent
5 inflammatory conditions, immune disorders, and allergic disorders. Preferred pharmaceutical compositions and dosage forms comprise a compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable prodrug, salt, solvate, or clathrate thereof, optionally in combination with one or more additional active agents.

10

Single unit dosage forms of the invention are suitable for oral, mucosal (*e.g.*, nasal, sublingual, vaginal, buccal, or rectal), parenteral (*e.g.*, subcutaneous, intravenous, bolus injection, intramuscular, or intraarterial), or transdermal administration to a patient. Examples of dosage forms include, but are not limited to: tablets; caplets; capsules, such as
15 soft elastic gelatin capsules; cachets; troches; lozenges; dispersions; suppositories; ointments; cataplasms (poultices); pastes; powders; dressings; creams; plasters; solutions; patches; aerosols (*e.g.*, nasal sprays or inhalers); gels; liquid dosage forms suitable for oral or mucosal administration to a patient, including suspensions (*e.g.*, aqueous or non-aqueous liquid suspensions, oil-in-water emulsions, or a water-in-oil liquid emulsions), solutions, and elixirs;
20 liquid dosage forms suitable for parenteral administration to a patient; and sterile solids (*e.g.*, crystalline or amorphous solids) that can be reconstituted to provide liquid dosage forms suitable for parenteral administration to a patient.

20

The composition, shape, and type of dosage forms of the invention will typically vary
25 depending on their use. For example, a dosage form suitable for mucosal administration may contain a smaller amount of active ingredient(s) than an oral dosage form used to treat the same indication. This aspect of the invention will be readily apparent to those skilled in the art. *See, e.g.*, Remington's Pharmaceutical Sciences (1990) 18th ed., Mack Publishing, Easton PA.

30

Typical pharmaceutical compositions and dosage forms comprise one or more excipients. Suitable excipients are well known to those skilled in the art of pharmacy, and non-limiting examples of suitable excipients are provided herein. Whether a particular excipient is suitable
35 for incorporation into a pharmaceutical composition or dosage form depends on a variety of factors well known in the art including, but not limited to, the way in which the dosage form will be administered to a patient. For example, oral dosage forms such as tablets may contain excipients not suited for use in parenteral dosage forms.

35

The suitability of a particular excipient may also depend on the specific active ingredients in the dosage form. For example, the decomposition of some active ingredients can be accelerated by some excipients such as lactose, or when exposed to water. Active ingredients
5 that comprise primary or secondary amines (*e.g.*, N-desmethylvenlafaxine and N,N-didesmethylvenlafaxine) are particularly susceptible to such accelerated decomposition. Consequently, this invention encompasses pharmaceutical compositions and dosage forms that contain little, if any, lactose. As used herein, the term "lactose-free" means that the amount of lactose present, if any, is insufficient to substantially increase the degradation rate
10 of an active ingredient. Lactose-free compositions of the invention can comprise excipients that are well known in the art and are listed, for example, in the U.S. Pharmacopia (USP) SP (XXI)/NF (XVI). In general, lactose-free compositions comprise active ingredients, a binder/filler, and a lubricant in pharmaceutically compatible and pharmaceutically acceptable amounts. Preferred lactose-free dosage forms comprise active ingredients, microcrystalline
15 cellulose, pre-gelatinized starch, and magnesium stearate.

This invention further encompasses anhydrous pharmaceutical compositions and dosage forms comprising active ingredients, since water can facilitate the degradation of some compounds. For example, the addition of water (*e.g.*, 5%) is widely accepted in the
20 pharmaceutical arts as a means of simulating long-term storage in order to determine characteristics such as shelf-life or the stability of formulations over time. *See, e.g.*, Jens T. Carstensen (1995) *Drug Stability: Principles & Practice*, 2d. Ed., Marcel Dekker, NY, NY, 379-80. In effect, water and heat accelerate the decomposition of some compounds. Thus, the effect of water on a formulation can be of great significance since moisture and/or
25 humidity are commonly encountered during manufacture, handling, packaging, storage, shipment, and use of formulations.

Anhydrous pharmaceutical compositions and dosage forms of the invention can be prepared using anhydrous or low moisture containing ingredients and low moisture or low humidity
30 conditions. Pharmaceutical compositions and dosage forms that comprise lactose and at least one active ingredient that comprises a primary or secondary amine are preferably anhydrous if substantial contact with moisture and/or humidity during manufacturing, packaging, and/or storage is expected.

35 An anhydrous pharmaceutical composition should be prepared and stored such that its anhydrous nature is maintained. Accordingly, anhydrous compositions are preferably packaged using materials known to prevent exposure to water such that they can be included

in suitable formulary kits. Examples of suitable packaging include, but are not limited to, hermetically sealed foils, plastics, unit dose containers (*e.g.*, vials), blister packs, and strip packs.

- 5 The invention further encompasses pharmaceutical compositions and dosage forms that comprise one or more compounds that reduce the rate by which an active ingredient will decompose. Such compounds, which are referred to herein as “stabilizer” include, but are not limited to, antioxidants such as ascorbic acid, pH buffers, or salt buffers.
- 10 Like the amounts and types of excipients, the amounts and specific types of active ingredients in a dosage form may differ depending on factors such as, but not limited to, the route by which it is to be administered to patients. However, typical dosage forms of the invention comprise a compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug
- 15 thereof in an amount of from about 1 mg to about 1000 mg, preferably in an amount of from about 50 mg to about 500 mg, and most preferably in an amount of from about 75 mg to about 350 mg. The typical total daily dosage of a compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof can range from about 1 mg to about
- 20 5000 mg per day, preferably in an amount from about 50 mg to about 1500 mg per day, more preferably from about 75 mg to about 1000 mg per day. It is within the skill of the art to determine the appropriate dose and dosage form for a given patient.

ORAL DOSAGE FORMS

- 25 Pharmaceutical compositions of the invention that are suitable for oral administration can be presented as discrete dosage forms, such as, but are not limited to, tablets (*e.g.*, chewable tablets), caplets, capsules, and liquids (*e.g.*, flavored syrups). Such dosage forms contain predetermined amounts of active ingredients, and may be prepared by methods of pharmacy well known to those skilled in the art. *See generally*, Remington's Pharmaceutical Sciences
- 30 (1990) 18th ed., Mack Publishing, Easton PA.

- Typical oral dosage forms of the invention are prepared by combining the active ingredient(s) in an admixture with at least one excipient according to conventional pharmaceutical compounding techniques. Excipients can take a wide variety of forms depending on the form
- 35 of preparation desired for administration. For example, excipients suitable for use in oral liquid or aerosol dosage forms include, but are not limited to, water, glycols, oils, alcohols, flavoring agents, preservatives, and coloring agents. Examples of excipients suitable for use

in solid oral dosage forms (*e.g.*, powders, tablets, capsules, and caplets) include, but are not limited to, starches, sugars, micro-crystalline cellulose, diluents, granulating agents, lubricants, binders, and disintegrating agents.

5 Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid excipients are employed. If desired, tablets can be coated by standard aqueous or nonaqueous techniques. Such dosage forms can be prepared by any of the methods of pharmacy. In general, pharmaceutical compositions and dosage forms are prepared by uniformly and intimately admixing the active ingredients with liquid carriers,
10 finely divided solid carriers, or both, and then shaping the product into the desired presentation if necessary.

For example, a tablet can be prepared by compression or molding. Compressed tablets can be prepared by compressing in a suitable machine the active ingredients in a free-flowing form
15 such as powder or granules, optionally mixed with an excipient. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

Examples of excipients that can be used in oral dosage forms of the invention include, but are
20 not limited to, binders, fillers, disintegrants, and lubricants. Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (*e.g.*, ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium
25 carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, (*e.g.*, Nos. 2208, 2906, 2910), microcrystalline cellulose, and mixtures thereof.

Suitable forms of microcrystalline cellulose include, but are not limited to, the materials sold
30 as AVICEL-PH-101, AVICEL-PH-103 AVICEL RC-581, AVICEL-PH-105 (available from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, PA), and mixtures thereof. One specific binder is a mixture of microcrystalline cellulose and sodium carboxymethyl cellulose sold as AVICEL RC-581. Suitable anhydrous or low moisture excipients or additives include AVICEL-PH-103J and Starch 1500 LM.

35 Examples of fillers suitable for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (*e.g.*, granules or

powder), microcrystalline cellulose, powdered cellulose, dextrans, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The binder or filler in pharmaceutical compositions of the invention is typically present in from about 50 to about 99 weight percent of the pharmaceutical composition or dosage form.

5

Disintegrants are used in the compositions of the invention to provide tablets that disintegrate when exposed to an aqueous environment. Tablets that contain too much disintegrant may disintegrate in storage, while those that contain too little may not disintegrate at a desired rate or under the desired conditions. Thus, a sufficient amount of disintegrant that is neither too
10 much nor too little to detrimentally alter the release of the active ingredients should be used to form solid oral dosage forms of the invention. The amount of disintegrant used varies based upon the type of formulation, and is readily discernible to those of ordinary skill in the art. Typical pharmaceutical compositions comprise from about 0.5 to about 15 weight percent of disintegrant, preferably from about 1 to about 5 weight percent of disintegrant.

15

Disintegrants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other
20 starches, clays, other algins, other celluloses, gums, and mixtures thereof.

25

Lubricants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other glycols, stearic acid,
25 sodium lauryl sulfate, talc, hydrogenated vegetable oil (e.g., peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laureate, agar, and mixtures thereof. Additional lubricants include, for example, a syloid silica gel (AEROSIL 200, manufactured by W.R. Grace Co. of Baltimore, MD), a coagulated aerosol of synthetic silica (marketed by Degussa Co. of Plano, TX), CAB-O-SIL (a pyrogenic
30 silicon dioxide product sold by Cabot Co. of Boston, MA), and mixtures thereof. If used at all, lubricants are typically used in an amount of less than about 1 weight percent of the pharmaceutical compositions or dosage forms into which they are incorporated.

30

CONTROLLED RELEASE DOSAGE FORMS

35

Active ingredients of the invention can be administered by controlled release means or by delivery devices that are well known to those of ordinary skill in the art. Examples include, but are not limited to, those described in U.S. Patent Nos.: 3,845,770; 3,916,899; 3,536,809;

3,598,123; and 4,008,719, 5,674,533, 5,059,595, 5,591,767, 5,120,548, 5,073,543, 5,639,476, 5,354,556, and 5,733,566, each of which is incorporated herein by reference. Such dosage forms can be used to provide slow or controlled-release of one or more active ingredients using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable
5 membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the active ingredients of the invention. The invention thus encompasses single unit dosage forms suitable for oral
10 administration such as, but not limited to, tablets, capsules, gelcaps, and caplets that are adapted for controlled-release.

All controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an
15 optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include extended activity of the drug, reduced dosage frequency, and increased patient compliance. In addition, controlled-release formulations can be used to affect the time of onset of action or other
20 characteristics, such as blood levels of the drug, and can thus affect the occurrence of side (*e.g.*, adverse) effects.

Most controlled-release formulations are designed to initially release an amount of drug (active ingredient) that promptly produces the desired therapeutic effect, and gradually and
25 continually release of other amounts of drug to maintain this level of therapeutic or prophylactic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled-release of an active ingredient can be stimulated by various conditions including, but not limited to, pH,
30 temperature, enzymes, water, or other physiological conditions or compounds.

A particular extended release formulation of this invention comprises a therapeutically or prophylactically effective amount of a compound of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1, or a pharmaceutically acceptable salt, solvate,
35 hydrate, clathrate, or prodrug thereof, in spheroids which further comprise microcrystalline cellulose and, optionally, hydroxypropylmethyl-cellulose coated with a mixture of ethyl cellulose and hydroxypropylmethylcellulose. Such extended release formulations can be

prepared according to U.S. Patent No. 6,274,171, the entire teachings of which are incorporated herein by reference.

5 A specific controlled-release formulation of this invention comprises from about 6% to about 40% a compound of any one of formulas (I), (IIA), (IIB), (IIIA), (IIIB), (IVA), (IVB), (VA), or (VB), or Table 1 by weight, about 50% to about 94% microcrystalline cellulose, NF, by weight, and optionally from about 0.25% to about 1% by weight of hydroxypropyl-methylcellulose, USP, wherein the spheroids are coated with a film coating composition comprised of ethyl cellulose and hydroxypropylmethylcellulose.

10

PARENTERAL DOSAGE FORMS

Parenteral dosage forms can be administered to patients by various routes including, but not limited to, subcutaneous, intravenous (including bolus injection), intramuscular, and intraarterial. Because their administration typically bypasses patients' natural defenses
15 against contaminants, parenteral dosage forms are preferably sterile or capable of being sterilized prior to administration to a patient. Examples of parenteral dosage forms include, but are not limited to, solutions ready for injection, dry products ready to be dissolved or suspended in a pharmaceutically acceptable vehicle for injection, suspensions ready for injection, and emulsions.

20

Suitable vehicles that can be used to provide parenteral dosage forms of the invention are well known to those skilled in the art. Examples include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated
25 Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

30

Compounds that increase the solubility of one or more of the active ingredients disclosed herein can also be incorporated into the parenteral dosage forms of the invention.

TRANSDERMAL, TOPICAL, AND MUCOSAL DOSAGE FORMS

Transdermal, topical, and mucosal dosage forms of the invention include, but are not limited
35 to, ophthalmic solutions, sprays, aerosols, creams, lotions, ointments, gels, solutions, emulsions, suspensions, or other forms known to one of skill in the art. *See, e.g.*, Remington's Pharmaceutical Sciences (1980 & 1990) 16th and 18th eds., Mack Publishing, Easton PA and

Introduction to Pharmaceutical Dosage Forms (1985) 4th ed., Lea & Febiger, Philadelphia. Dosage forms suitable for treating mucosal tissues within the oral cavity can be formulated as mouthwashes or as oral gels. Further, transdermal dosage forms include “reservoir type” or “matrix type” patches, which can be applied to the skin and worn for a specific period of time
5 to permit the penetration of a desired amount of active ingredients.

Suitable excipients (*e.g.*, carriers and diluents) and other materials that can be used to provide transdermal, topical, and mucosal dosage forms encompassed by this invention are well known to those skilled in the pharmaceutical arts, and depend on the particular tissue to which
10 a given pharmaceutical composition or dosage form will be applied. With that fact in mind, typical excipients include, but are not limited to, water, acetone, ethanol, ethylene glycol, propylene glycol, butane-1,3-diol, isopropyl myristate, isopropyl palmitate, mineral oil, and mixtures thereof to form lotions, tinctures, creams, emulsions, gels or ointments, which are non-toxic and pharmaceutically acceptable. Moisturizers or humectants can also be added to
15 pharmaceutical compositions and dosage forms if desired. Examples of such additional ingredients are well known in the art. *See, e.g.*, Remington's Pharmaceutical Sciences (1980 & 1990) 16th and 18th eds., Mack Publishing, Easton PA.

Depending on the specific tissue to be treated, additional components may be used prior to, in
20 conjunction with, or subsequent to treatment with active ingredients of the invention. For example, penetration enhancers can be used to assist in delivering the active ingredients to the tissue. Suitable penetration enhancers include, but are not limited to: acetone; various alcohols such as ethanol, oleyl, and tetrahydrofuryl; alkyl sulfoxides such as dimethyl sulfoxide; dimethyl acetamide; dimethyl formamide; polyethylene glycol; pyrrolidones such
25 as polyvinylpyrrolidone; Kollidon grades (Povidone, Polyvidone); urea; and various water-soluble or insoluble sugar esters such as Tween 80 (polysorbate 80) and Span 60 (sorbitan monostearate).

The pH of a pharmaceutical composition or dosage form, or of the tissue to which the
30 pharmaceutical composition or dosage form is applied, may also be adjusted to improve delivery of one or more active ingredients. Similarly, the polarity of a solvent carrier, its ionic strength, or tonicity can be adjusted to improve delivery. Compounds such as stearates can also be added to pharmaceutical compositions or dosage forms to advantageously alter the hydrophilicity or lipophilicity of one or more active ingredients so as to improve delivery. In
35 this regard, stearates can serve as a lipid vehicle for the formulation, as an emulsifying agent or surfactant, and as a delivery-enhancing or penetration-enhancing agent. Different salts,

hydrates or solvates of the active ingredients can be used to further adjust the properties of the resulting composition.

COMBINATION THERAPY

5 The methods for immunosuppression or for treating or preventing inflammatory conditions, allergic disorders, and immune disorders in a patient in need thereof can further comprise administering to the patient being administered a compound of this invention, an effective amount of one or more other active agents. Such active agents may include those used conventionally for immunosuppression or for inflammatory conditions, allergic disorders, or
10 immune disorders. These other active agents may also be those that provide other benefits when administered in combination with the compounds of this invention. For example, other therapeutic agents may include, without limitation, steroids, non-steroidal anti-inflammatory agents, antihistamines, analgesics, immunosuppressive agents and suitable mixtures thereof. In such combination therapy treatment, both the compounds of this invention and the other
15 drug agent(s) are administered to a subject (e.g., humans, male or female) by conventional methods. The agents may be administered in a single dosage form or in separate dosage forms. Effective amounts of the other therapeutic agents and dosage forms are well known to those skilled in the art. It is well within the skilled artisan's purview to determine the other therapeutic agent's optimal effective-amount range.

20 In one embodiment of the invention where another therapeutic agent is administered to a subject, the effective amount of the compound of this invention is less than its effective amount when the other therapeutic agent is not administered. In another embodiment, the effective amount of the conventional agent is less than its effective amount when the
25 compound of this invention is not administered. In this way, undesired side effects associated with high doses of either agent may be minimized. Other potential advantages (including without limitation improved dosing regimens and/or reduced drug cost) will be apparent to those of skill in the art.

30 In one embodiment relating to autoimmune, allergic and inflammatory conditions, the other therapeutic agent may be a steroid or a non-steroidal anti-inflammatory agent. Particularly useful non-steroidal anti-inflammatory agents, include, but are not limited to, aspirin, ibuprofen, diclofenac, naproxen, benoxaprofen, flurbiprofen, fenoprofen, flubufen, ketoprofen, indoprofen, piroprofen, carprofen, oxaprozin, pramoprofen, muroprofen,
35 trioxaprofen, suprofen, aminoprofen, tiaprofenic acid, fluprofen, bucloxic acid, indomethacin, sulindac, tolmetin, zomepirac, tiopinac, zidometacin, acemetacin, fentiazac, clidanac, oxpinac, mefenamic acid, meclofenamic acid, flufenamic acid, niflumic acid, tolfenamic acid,

diflurisal, flufenisal, piroxicam, sudoxicam, isoxicam; salicylic acid derivatives, including aspirin, sodium salicylate, choline magnesium trisalicylate, salsalate, diflunisal, salicylsalicylic acid, sulfasalazine, and olsalazin; para-aminophenol derivatives including acetaminophen and phenacetin; indole and indene acetic acids, including indomethacin, 5 sulindac, and etodolac; heteroaryl acetic acids, including tolmetin, diclofenac, and ketorolac; anthranilic acids (fenamates), including mefenamic acid, and meclofenamic acid; enolic acids, including oxicams (piroxicam, tenoxicam), and pyrazolidinediones (phenylbutazone, oxyphenthartazone); and alkanones, including nabumetone and pharmaceutically acceptable salts thereof and mixtures thereof. For a more detailed description of the NSAIDs, see Paul 10 A. Insel, *Analgesic-Antipyretic and Antiinflammatory Agents and Drugs Employed in the Treatment of Gout*, in Goodman & Gilman's *The Pharmacological Basis of Therapeutics* 617-57 (Perry B. Molinoff and Raymond W. Ruddon eds., 9th ed 1996) and Glen R. Hanson, *Analgesic, Antipyretic and Anti-Inflammatory Drugs in Remington: The Science and Practice of Pharmacy Vol II* 1196-1221 (A.R. Gennaro ed. 19th ed. 1995) which are hereby 15 incorporated by reference in their entireties.

Of particular relevance to allergic disorders, the other therapeutic agent may be an antihistamine. Useful antihistamines include, but are not limited to, loratadine, cetirizine, 20 fexofenadine, desloratadine, diphenhydramine, chlorpheniramine, chlorcyclizine, pyrilamine, promethazine, terfenadine, doxepin, carbinoxamine, clemastine, tripeleminamine, brompheniramine, hydroxyzine, cyclizine, meclizine, cyproheptadine, phenindamine, acrivastine, azelastine, levocabastine, and mixtures thereof. For a more detailed description of antihistamines, see Goodman & Gilman's *The Pharmacological Basis of Therapeutics* (2001) 651-57, 10th ed).

25 Immunosuppressive agents include glucocorticoids, corticosteroids (such as Prednisone or Solumedrol), T cell blockers (such as cyclosporin A and FK506), purine analogs (such as azathioprine (Imuran)), pyrimidine analogs (such as cytosine arabinoside), alkylating agents (such as nitrogen mustard, phenylalanine mustard, buslfan, and cyclophosphamide), folic acid 30 antagonists (such as aminopterin and methotrexate), antibiotics (such as rapamycin, actinomycin D, mitomycin C, puramycin, and chloramphenicol), human IgG, antilymphocyte globulin (ALG), and antibodies (such as anti-CD3 (OKT3), anti-CD4 (OKT4), anti-CD5, anti-CD7, anti-IL-2 receptor, anti-alpha/beta TCR, anti-ICAM-1, anti-CD20 (Rituxan), anti-IL-12 and antibodies to immunotoxins).

35 The foregoing and other useful combination therapies will be understood and appreciated by those of skill in the art. Potential advantages of such combination therapies include a

different efficacy profile, the ability to use less of each of the individual active ingredients to minimize toxic side effects, synergistic improvements in efficacy, improved ease of administration or use and/or reduced overall expense of compound preparation or formulation.

5

OTHER EMBODIMENTS

The compounds of this invention may be used as research tools (for example, as a positive control for evaluating other potential CRAC inhibitors, or IL-2, IL-4, IL-5, IL-13, GM-CSF, TNF- α , and/or INF- γ inhibitors). These and other uses and embodiments of the compounds and compositions of this invention will be apparent to those of ordinary skill in the art.

10
15

The invention is further defined by reference to the following examples describing in detail the preparation of compounds of the invention. It will be apparent to those skilled in the art that many modifications, both to materials and methods, may be practiced without departing from the purpose and interest of this invention. The following examples are set forth to assist in understanding the invention and should not be construed as specifically limiting the invention described and claimed herein. Such variations of the invention, including the substitution of all equivalents now known or later developed, which would be within the purview of those skilled in the art, and changes in formulation or minor changes in experimental design, are to be considered to fall within the scope of the invention incorporated herein.

20

EXAMPLES

25

EXPERIMENTAL RATIONALE

Without wishing to be bound by theory, it is believed that the compounds of this invention inhibit CRAC ion channels, thereby inhibiting production of IL-2 and other key cytokines involved with inflammatory, allergic and immune responses. The examples that follow demonstrate these properties.

30

MATERIALS AND GENERAL METHODS

Reagents and solvents used below can be obtained from commercial sources such as Aldrich Chemical Co. (Milwaukee, Wisconsin, USA). $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra were recorded on a Varian 300MHz NMR spectrometer. Significant peaks are tabulated in the order: δ (ppm): chemical shift, multiplicity (s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br s, broad singlet), coupling constant(s) in Hertz (Hz) and number of protons.

35


Patch clamp experiments were performed in the tight-seal whole-cell configuration at 21-25°C. High resolution current recordings were acquired by a computer-based patch clamp amplifier system (EPC-9, HEKA, Lambrecht, Germany). Patch pipettes had resistances
5 between 2-4 M Ω after filling with the standard intracellular solution. Immediately following establishment of the whole-cell configuration, voltage ramps of 50-200 ms duration spanning the voltage range of -100 to +100 mV were delivered at a rate of 0.5 Hz over a period of 300-400 seconds. All voltages were corrected for a liquid junction potential of 10 mV between
10 external and internal solutions when using glutamate as the intracellular anion. Currents were filtered at 2.9 kHz and digitized at 10 μ s intervals. Capacitive currents and series resistance were determined and corrected before each voltage ramp using the automatic capacitance compensation of the EPC-9. The low resolution temporal development of membrane currents was assessed by extracting the current amplitude at -80 mV or +80 mV from individual ramp current records.

15

SYNTHESIS OF REPRESENTATIVE EXEMPLARY COMPOUNDS OF THIS INVENTION

In general, the compounds of the invention can be synthesized using methods analogous to
20 those described in U.S. Patent Application Serial No. 10/897,681 and U.S. Patent Application Serial No. 11/233,224, the entire teachings of these patent applications are incorporated herein by reference.

General Suzuki Coupling Procedure for Boronic Esters:




(X = halo or OTf)

5

To a solution of Pd(PhCN)₂Cl₂ (767mg, 2mmol) and 1,4-bis(diphenylphosphino)-butane (dppb) (1.28g, 3mmol) in 100 mL of toluene was added a boronic ester (XXXVII) (22.6 mmol) followed by an alkenyl halide or alkenyl triflate (XXXVI) (22.6 mmol), and sodium carbonate(3.4g, 22.6mmol), followed by 23 mL of water and 2 mL of methanol. This mixture was heated to 100°C and refluxed for 18 hours. After allowing cooling to room temperature, the reaction was quenched with water and worked-up with ethyl acetate. The organic solution was then filtered through silica gel, evaporated in vacuo to give an amine derivative (XXXVIII) that was used without further purification in the next step.

15


General Procedure for amide formation using acyl chloride:



An amine derivative (XXXVIII) (19.2 mmol) was dissolved in methylene chloride and cooled to 0°C. Pyridine (3.87ml, 48 mmol) was added to the reaction mixture followed by an acyl chloride (XXXIX) (19.2 mmol). This mixture was allowed to warm to room temperature for 35 minutes at which time water was added and the reaction mixture was partitioned between the methylene chloride and water layers. The organic layer was collected and dried over a drying agent such as NaSO₃, then concentrated. The product was purified using flash chromatography to give a compound of the invention (XL).

25


General Procedure for Pyridinium Salt formation



A solution of pyridine (200mg, 0.62mmol) was dissolved in dichloromethane(1.5ml). Benzyl bromide (3eq) was added and the room temperature solution was stirred for 24 hours. The solid was filtered and washed with cold dichloromethane to give pure pyridinium salt.

5


General Procedure for Reduction of Pyridinium Salt



The pyridinium salt (crude) was dissolved in methanol and sodium borohydride (113mg, 3eq) was added. This solution was warmed to 35 degrees for 5 hours then cooled to room


10 temperature. The methanol was removed, water and methylene chloride added, and extracted. The 150mg (70%yield) of tetrahydropyridine was produced after column chromatography.

General procedure for Carbamate.




The benzyl amine (31mg, 0.074mmol) was dissolved in methylene chloride(2ml) at room temperature. Ethyl chloroformate (14ul, 2eq) was added to the solution and it was stirred for 3 hours The solvent was removed and the crude mixture was flashed with ethyl acetate/hexane to produce 28.4mg (97%yield) of carbamate.

5 General Procedure for Tetrahydropyridine deprotection



10% Palladium on carbon (10mg, 10%w/w) was added to a methanolic solution (1.5ml) of benzylcarbamate (109mg, 0.247mmol). A balloon of hydrogen was added to the top of the flask and thoroughly stirred for 1 hour. After filtration completely pure free amine was recovered (73mg, 96%yield). No further purification required.

General Procedure for thiazole amine formation.



The amine(725mg, 1.73mmol) was dissolved in methylene chloride(5ml) and cyanogen bromide(549mg,5.19mmol) was added. This reaction was stirred at room temperature for 15 hour then purified by flash chromatography. Pure cyanamide (415mg, 68% yield) was isolated as a white solid.


The cyanamide(55mg, .01556mmol) was dissolved in ethanol/ammonia(2ml, 2M). hydrogen sulfide gas was then bubbled through the reaction solution for 20minutes at which time the bubbling was siesed and the reaction was capped and stirred for 5hours at room temperature.

20 After completion was solvent was removed and the thiourea was recrystallized from ethanol to remove yellow color.

The thiourea was then dissolved in anhydrous THF (5ml) and the bromide (140mg, 0.709mmol) was added and refluxed for 5 hours. The rxn mixture was then quenched with sodium bicarbonate solution and diluted with ethylacetate. After drying over magnesium sulfate, the compound was purified by flash chromatography yielding 39mg of pure thiazole product. The compound was then dissolved in ethanol and excess HCl in ether was added to form the salt. Solid precipitated out of solution at -40°C and was filtered.

Example 1: Ethyl 3-(4-(2,6-difluorobenzamido)phenyl)-4-methyl-5,6-dihydropyridine-

1(2H)-carboxylate



5 Compound A (31mg, 0.074mmol) was dissolved in methylene chloride (2mL) at room temperature. Ethyl chloroformate (14 μ l, 2 eq) was added to the solution and it was heated to 110°C in the microwave for 1 hour. The solvent was removed and the crude mixture was flashed with ethyl acetate/hexane to produce 28.4 mg (97% yield) of Compound 12.


¹H NMR (300 MHz, CDCl₃) δ 7.65 (d, $J=8.1$ Hz, 2H), 7.36-7.48 (m, 1H), 7.19 (d, $J=8.1$ Hz, 2H), 6.997 (t, $J=8.1$ Hz, 1H), 4.00-4.20(m, 3H), 3.59-3.63 (m, 1H), 2.17-2.20 (m, 3H), 1.63 (s, 3H), 1.26 (t, $J=6.3$ Hz, 3H) ppm.

10

ESMS calcd. (C₂₂H₂₂F₂N₂O₃): 400.2; found: 401.2. (M + H).

Example 2: methyl 3-(4-(2,6-difluorobenzamido)phenyl)-4-(thiazol-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (B) and methyl 3-(4-(2,6-difluorobenzamido)phenyl)-4-isopropyl-5,6-dihydropyridine-1(2H)-carboxylate (C)

15




- N*-[4-(4-bromopyridin-3-yl)phenyl]-2,6-difluorobenzamide (e). To a solution of 4-Bromopyridine hydrochloride (500 mg, 2.57 mmol) in THF (15 mL) was added LDA (2.0 N, 2.6 mL, 5.2 mmol) at $-78\text{ }^\circ\text{C}$. After 30 min, the solution of $ZnCl_2$ (1 M in ether, 3.1 mL, 3.1 mmol) in THF (5 mL) was added slowly. The cold bath was removed and the solution was allowed to warm up room temperature. To this solution was added $Pd(PPh_3)_4$ (50 mg) and compound **d** (370 mg, 1.03 mmol). The solution was heated at $65\text{ }^\circ\text{C}$ for 2 hrs before it was diluted with EtOAc (20 mL) and H_2O (10 mL). The organic phase was collected, dried and concentrated. Column chromatography afforded Compound **e** (210 mg, 21 %). 1H NMR (300 MHz, $(CD_3)_2SO$) δ 10.99 (s, 1 H), 8.52 (s, 1 H), 8.40 (d, $J = 5.4$ Hz, 1 H), 7.84-7.78 (m, 3 H), 7.63-7.57 (m, 1 H), 7.47 (d, $J = 8.7$ Hz, 2 H), 7.27 (t, $J = 8.1$ Hz, 2 H); ESMS calcd ($C_{18}H_{11}BrF_2N_2O$): 388.0; found: 389.1 (M+H).
- 15 Example 3: 2,6-Difluoro-*N*-[4-(1-methoxycarbonyl-4-thiazol-2-yl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide

The solution of Compound e (15 mg, 0.04 mmol), 2-thiazolylzinc bromide solution (0.5 M, 0.2 mL, 0.1 mmol), and Pd(PPh₃)₂Cl₂ (5 mg) in THF (4 mL) was heated in microwave synthesizer at 150 °C for 30 min. The solution was concentrated and chromatographed to give Compound f (13 mg, 83%). To the solution of Compound f (13 mg, 0.033 mmol) in MeOH (2 mL) was added benzyl bromide (BnBr) (0.05 mL, 0.4 mmol). After stirring 60 min, the solvent was removed and the residue was washed with EtOAc (5 mL) to remove BnBr. The resulting residue was dissolved in MeOH (4 mL) and added NaBH₄ (10 mg, 0.26 mmol). The solution was passed through a short column of silica gel. The filtrate was concentrated and dissolved in DCM (4 mL). After adding methyl chloroformate (0.1 mL), the solution was stirred overnight before it was concentrated and chromatographed to give Compound 41 (10 mg, 67%). ¹H NMR (300 MHz, CDCl₃) δ 7.79-7.62 (m, 4 H), 7.52-7.38 (m, 1 H), 7.28-7.18 (m, 2 H), 7.09-6.98 (m, 3 H), 4.26-4.18 (m, 2 H), 3.74 (s, 3 H), 2.95-2.88 (m, 2 H), 1.14-1.09 (m, 2 H); ESMS caclcd (C₂₃H₁₉F₂N₃O₃S): 455.1; found: 456.2 (M+H).

Example 4: 2,6-Difluoro-N-[4-(1-methoxycarbonyl-4-isopropyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide


To a solution of isopropylmagnesium chloride (2 M, 0.65 mL, 1.3 mmol) in Et₂O (5 mL) was added ZnCl₂ (1 M, 1.5 mL, 1.5 mmol) at 0 °C. After 30 min, Compound e (50 mg, 0.13 mmol) was added. The solution was stirred at room temperature for 2 hrs before it was quenched with NH₄Cl (5 mL). The mixture was diluted with EtOAc and the organic phase was collected, dried, and concentrated. Column chromatography afforded Compound g (25 mg, 56%). From Compound g, Compound 42 was prepared following the same procedure as described for Compound 41 to give a 35 % yield. ¹H NMR (300 MHz, CDCl₃) δ 7.64-7.59 (m, 3 H), 7.45-7.38 (m, 1 H), 7.19-7.16 (m, 2 H), 7.14-6.97 (m, 2 H), 4.06-3.99 (m, 2 H), 3.73 (s, 3 H), 3.62-3.58 (m, 2 H), 2.22-2.18 (m, 2 H), 1.94-1.89 (m, 1 H), 1.57 (s, 6 H); ESMS caclcd (C₂₃H₂₄F₂N₂O₃): 414.2; found: 415.3 (M+H).

General Procedure for Pyridinium Salt Formation:



A solution of pyridine (200mg, 0.62mmol) was dissolved in dichloromethane (1.5ml). Benzyl bromide (3eq) was added and the room temperature solution was stirred for 24 hours. The solid was filtered and washed with cold dichloromethane to give pure pyridinium salt.


General Procedure for Reduction of Pyridinium Salt:



The pyridinium salt (crude) was dissolved in methanol and sodium borohydride (113 mg, 3 eq) was added. This solution was warmed to 35 °C for 5 hours then cooled to room temperature. The methanol was removed, water and methylene chloride added, and extracted.


5 The 150 mg (70% yield) of tetrahydropyridine was produced after column chromatography.

General procedure for Carbamate Formation:




The Compound D (31mg, 0.074mmol) was dissolved in methylene chloride (2 ml) at room temperature. Ethyl chloroformate (14 μ l, 2eq) was added to the solution and the reaction was stirred for 3 hours. The solvent was removed and the crude mixture was purified by flash chromatograph with ethyl acetate/hexane to produce 28.4mg (97% yield) of Compound E.

5 General Procedure for Tetrahydropyridine Deprotection:



10% Palladium on carbon (10mg, 10%w/w) was added to a methanolic solution (1.5ml) of Compound F (109 mg, 0.247 mmol). A balloon of hydrogen was added to the top of the flask and thoroughly stirred for 1 hour. After filtration pure Compound G was recovered (73 mg, 96% yield). No further purification was required.

General Procedure for Thiazole Amine Formation:




Compound D (725mg, 1.73 mmol) was dissolved in methylene chloride (5 ml) and cyanogen bromide (549 mg, 5.19 mmol) was added. This reaction was stirred at room temperature for 1 hour then purified by flash chromatography. Pure cyanamide (415mg, 68% yield) was isolated as a white solid.

The cyanamide (55 mg, 0.01556 mmol) was dissolved in ethanol/ammonia (2 ml, 2M). Hydrogen sulfide gas was then bubbled through the reaction solution for 20 minutes at which time the bubbling was stopped, and the reaction was capped and allowed to stirred for 5 hours at room temperature. After completion, the solvent was removed and the thiourea was recrystallized from ethanol to remove yellow color.

The thiourea was then dissolved in anhydrous THF (5 ml) and the bromide (140 mg, 0.709 mmol) was added and refluxed for 5 hours. The reaction mixture was then quenched with sodium bicarbonate solution and diluted with ethyl acetate. After drying over magnesium sulfate, the compound was purified by flash chromatography to yield 39 mg of pure Compound H. The hydrochloride salt of Compound H was prepared by dissolving

Compound H in ethanol and excess HCl in ether was added. Solid precipitated out of solution at -40°C and was collected by filtration to yield the hydrochloride salt.

General Procedure for Palladium amine coupling:



- 5 Compound J (200mg, 0.642mmol) was dissolved in toluene/MeOH/H₂O (6ml/0.5ml/1ml) followed by 2-bromo-pyridine (101 mg 0.77mmol), Pd (20 mg, 10% w/w), heated to 170°C for 1 hour. The solvent was removed and it was dissolved in methylene chloride and purified by flash column to give Compound K (20mg, 8% yield).

Example 5: N-[4-(1-pyridin-2-yl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-3-fluoro-isonicotinamide

10

Procedure: General Procedure for palladium amine coupling.


NMR (300 MHz, CDCl₃) 8.67 (d, $J=2.7\text{Hz}$, 1H), 8.65 (dd, $J=1.2, 4.8\text{Hz}$, 1H), 8.40 (d, $J=13.8\text{Hz}$, 1H), 8.21 (ddd, $J=0.9, 1.8, 4.8\text{Hz}$, 1H), 8.04 (dd, $J=4.8, 6.6\text{Hz}$, 1H), 7.68-7.62 (m, 2H), 7.51-7.45 (m, 1H), 7.32-7.26 (m, 1H), 6.66-6.58 (m, 2H), 4.09-4.06 (m, 2H), 3.83 (t, $J=5.7\text{Hz}$, 2H), 2.33 (s, 2H), 1.67 (s, 3H)ppm

15

ESMS calcd. (C₂₃H₂₁FN₄O): 388.2; found: 389.2(M + H).

Example 6: Ethyl 3-(6-(2,6-difluorobenzamido)pyridin-3-yl)-4-methyl-5,6-dihydropyridine-1(2H)-carboxylate

20




25

ESMS calcd. (C₂₁H₂₁F₂N₃O₃): 401.2; found: 402.3(M + H).

Procedure: See general procedure for Carbamate Formation, beginning with N-(5-(1-benzyl-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)-2,6-difluorobenzamide.

Example 7: 2,6-difluoro-N-(5-(4-methyl-1-(pyrimidin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide hydrochloride


Pyrimidyl amine synthesis:



Procedure: The tetrahydropyridine(25mg, 0.076mmol), 2-chloropyrimidine(17.3mg, 0.152mmol), and diisopropylamine(53μl, 0.303mmol) was dissolved in 5ml dichloromethane in a microwave tube. It was heated to 135°C for 1hour. The methylene chloride solution was then loaded on a column. Although the compound was not completely pure, when HCl in ether was added the pure salt fell out of the ethyl acetate solution. 4mg was isolated in a 13% yield.

ESMS calcd. (C₂₁H₁₉ClF₂N₆O): 444.1; found: 408.2(M - Cl).


Example 8: Allyl 4-(4-(2,6-difluorobenzamido)phenyl)-3-methyl-5,6-dihydropyridine-1(2H)-carboxylate



ESMS calcd. (C₂₃H₂₂F₂N₂O₃): 412.2; found: 413.2(M + H).

Procedure: Synthesized using the general thiazole procedure and using allyl chloroformate.

Example 9: 2,6-difluoro-N-(5-(4-methyl-1-picolinoyl-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)benzamide hydrochloride




5

ESMS calcd. ($C_{24}H_{21}ClF_2N_4O_2$): 470.1; found:435.2(M -Cl).

Procedure: See Example of Substitution of amine with Acid chloride/sulfonyl chloride beginning with 2,6-difluoro-N-(5-(4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)benzamide and picolinoyl chloride followed by acid formation using procedure (free base dissolved in ethyl acetate and 2 eq. of HCl in diethyl ether added, the ppt was filtered and dried).

Example 10: Ethyl 3-(acetoxymethyl)-4-(4-(2,6-difluorobenzamido)phenyl)-5,6-dihydropyridine-1(2H)-carboxylate

Procedure




5

Ethyl 1-benzyl-4-(4-(2,6-difluorobenzamido)phenyl)-1,2,5,6-tetrahydropyridine-3-carboxylate was made in the same fashion as the first step of thiazole amin formation. (200mg, 0.42mmol) was dissolved in 5ml of THF. Lithium aluminum hydride (48mg, 1.26mmol) was added as a solid at room temperature and the reaction was stirred for 20 minutes. It was then quenched with water and stirred until the solid was white. After drying with magnesium sulfate the allylic alcohol was dissolved in methylene chloride(5mls). Pyridine(100mg, 1.26mmol) and acetic anhydride(128.5mg, 1.26mmol) were added to the reaction and it was stirred overnight. Quenched with sodium bicarbonate(sat) solution and purified by column chromatography. 50mg of pure acetate was isolated.

15 (1-benzyl-4-(4-(2,6-difluorobenzamido)phenyl)-1,2,5,6-tetrahydropyridin-3-yl)methyl acetate(15mg,0.031mmol) was then dissolved in methylene chloride(5ml) and ethyl chloroformate was added. This reaction was warmed to 40°C for 30 minutes. Quenched with sodium bicarbonate solution and purified by flash chromatography. 5mg(36% yield) was isolated

20 ESMS calcd. (C₂₄H₂₄F₂N₂O₅): 458.2; found:459.2(M + H).

Example 11: ethyl 4-chloro-3-(4-(2,6-difluorobenzamido)phenyl)-5,6-dihydropyridine-1(2H)-carboxylate




ESMS calcd. (C₂₁H₁₉ClF₂N₂O₃): 420.1; found: 421.1(M + H).

Procedure: See general procedure for carbamate formation.

Example 12: 2,6-difluoro-N-(5-(4-methyl-1-(pyrazin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide (Compound 11)

Procedure:




10

The amine(2g, 6.2mmol) was dissolved in dichloromethane(0.1M) and cooled to 0 °C. The allyl chloroformate(1.32ml, 12.4mmol) was added and stirred for 45min. Quenched with sat. sodium bicarbonated solution and the dichloromethane layer was rotovaped onto silica gel. A quick column afforded clean carbamate and removed benzyl chloride. This solid was

15

dissolved in THF and 5ml of pyrrolidine was added along with 10% w/w palladium tetrakis and THF. Heated to reflux for 3hrs. Purified by column on alumina with dichloromethane /methanol. Isolated 1.09g (76%yield) of pure product.



20 General Aryl amine synthesis using KF:

The amine, KF and pyrizine were added to DMSO solution and heated in microwave for 15min at 100 °C. After cooling, water was added to the DMSO solution followed by dichloromethane. The organic layer as separated and evaporated. Isolated pure product was

25

obtained via column chromatography.

General Procedure for Pyrazine amide exchange:


The acetamide was dissolved in dichloromethane and diisopropyl ethylamine was added at
 5 room temperature. The acid chloride was then added and stirred for 2 hours . Quenched with
 saturated solution of sodium bicarbonate. The organic layer was separated and the
 dichloromethane was removed. The residue was dissolved in Methanol and potassium
 carbonate was then added followed by heating to reflux for 1 min, then evaporated onto silica
 gel and purified by flash chromatography.

10 ¹H NMR (300 MHz, CDCl₃) 9.70 (s, 1H), 8.46 (s, 1H), 8.27 (d, J=1.2Hz, 1H), 8.19 (d,
 J=1.2Hz, 1H), 8.08-8.09 (m, 1H), 7.84 (d, J=1.8Hz, 1H), 7.45-7.53 (m, 1H), 7.05 (t, J=6Hz,
 2H), 4.37-4.39 (m, 2H), 3.88 (t, J=4.5Hz, 2H), 2.39-2.41 (m, 2H), 1.86 (s, 3H) ppm ESMS
 calcd. (C₂₁H₁₈F₂N₆O) 408.2; found: 409.2 (M+H).

15 Example 13: 2,6-difluoro-N-(5-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-
 yl)pyridin-2-yl)benzamide (Compound 7)

20 ¹H NMR (300 MHz, CDCl₃) δ 9.01 (s, 1H), 8.37 (d, J=6.3Hz 1H), 8.22 (d, J=2.7Hz, 1H),
 8.01 (s, 1H), 7.65(dd, J=1.5, 6.3Hz, 1H), 7.40-7.52 (m, 2H), 7.00 (t, J=6.0Hz, 2H), 6.60-6.66
 (m, 2H), 4.06 (s, 2H), 3.81 (t, J=4.3Hz, 2H) 2.34 (s, 2H), 1.66 (s, 3H) ppm. ESMS calcd.
 (C₂₃H₂₀F₂N₄O) 406.2; found: 407.2 (M+H).

Procedure.



30 The amine(179mg, 0.544mmol) was dissolved in a solution of acetonitrile(5ml) and
 triethylamine (1ml) was added. At room temperature, the benzyl pyridinium salt(154mg,
 0.544mmol) was added and stirred for 1.5hr. The solvent was evaporated and methanol was
 added along with 5% Pd/C (15mg) and a hydrogen balloon was added to the top. After
 stirring overnight the solvent was removed and the product purified by column
 chromatography using ethyl acetate hexane. The desired product was btained as white solid
 (18mg).

Example 14: 2,6-difluoro-N-(5-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)benzamide hydrochloride (Compound 8)

A solution of the pyridine amine (8mg) was made in methylene chloride and methanol.

- 5 Excess HCl in ether was added and the solvent was then evaporated. The HCl salt was obtained as yellow solid (9mg).

ESMS calcd. (C₂₃H₂₁ClF₂N₄O) 442.1; found: 407.2 (M-Cl).

- 10 **Example 15:** N-(5-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide (Compound 12)

ESMS calcd. (C₂₁H₁₇ClF₂N₆O) 442.1; found: 443.2 (M+H).


See procedure for 2,6-difluoro-N-(5-(4-methyl-1-(pyrazin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide.

15

Example 16: 2,6-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide (Compound 9)

- 20 ESMS calcd. (C₂₄H₂₁F₂N₃O) 405.2; found: 406.2 (M+H).

Procedure:



- 25 The amine(200mg, 0.916mmol) was dissolved in a solution of acetonitrile(5ml) and triethylamine (1ml) was added. At room temperature, the benzyl pyridinium salt(188mg, 0.916mmol) was added and stirred for 1hr. The solvent was evaporated and methanol was added along with 10% Pd/C (20mg) and a hydrogen balloon was added to the top. After stirring 4 hour the solvent was removed and the product purified by column chromatography
- 30 using a mixture of ethyl acetate and hexane. The aryl amine intermediate was obtained as a syruo (187mg). The amine was then coupled with a carboxylic acid using general EDC coupling procedure to form amide.

Example 17: 2,4-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide (Compound 10)

ESMS calcd. (C₂₄H₂₁F₂N₃O) 405.2; found: 406.2 (M+1).

- 5 See synthesis of 2,6-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide

Example 18: 2,6-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide hydrochloride (Compound 13)

10

ESMS calcd. (C₂₃H₂₁Cl₃N₄O) 474.08; found: 439.1 (M-Cl).

2,6-difluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide hydrochloride

- 15 Example 19: 2-chloro-6-fluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide hydrochloride (Compound 14)

ESMS calcd. (C₂₄H₂₂Cl₂FN₃O) 457.1; found: 422.1 (M-Cl)

2-chloro-6-fluoro-N-(4-(4-methyl-1-(pyridin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide hydrochloride

20

Example 20: N-(5-(1-(3-chloropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide hydrochloride (Compound 15)

- 25 ¹H NMR (300 MHz, CDCl₃) 9.53(s, 1H), 8.45 (s, 1H), 8.35 (dd, J=0.9, 5.7Hz, 1H), 8.13 (dd, J=0.9, 4.5, 6.9, Hz, 1Hz), 7.53-7.61(m, 1H), 7.26 (dd, J=4.5, 5.7Hz, 1H), 7.13 (t, J=6Hz, 2H), 4.52 (s, 2H), 3.98 (t, J=4.2Hz, 2H), 2.63-2.64 (m, 2H), 1.95 (s, 3H) ppm ESMS calcd. (C₂₂H₁₉Cl₂F₂N₅O) 477.1; found: 442.1 (M-Cl).

- 30 Example 21: 2N-(5-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,4-difluorobenzamide (Compound 16)

¹H NMR (300 MHz, CDCl₃) 9.67 (d, J=0.9Hz, 1H), 8.95 (d, J=10.8Hz, 1H), 8.31 (d, J=0.9Hz, 1H), 8.25 (dd, J=4.8, 6.9, 13.5Hz, 1Hz), 8.09(d, J=1.8Hz, 1H), 7.85 (d, J=1.8Hz, 1H), 7.07-7.12 (m, 1H), 6.96-7.01 (m, 1H), 4.32-4.34 (m, 2H), 3.74 (t, J=4.2Hz, 2H), 2.48-2.49 (m, 2H),

- 35 1.86 (s, 3H)ppm ESMS calcd. (C₂₁H₁₇ClF₂N₆O) 442.1; found: 443.2 (M+H).

Example 22: 2,6-difluoro-N-(5-(1-(3-fluoropyridin-2-yl)-4-methyl-1,2,5,6-

tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide hydrochloride (Compound 17)

¹H NMR (300 MHz, DMSO d₆) 11.70 (s, 1H), 9.42 (s, 1H), 8.47 (s, 1H), 8.02 (d, *J*=3.6Hz, 1H), 7.54-7.63 (m 2H), 7.26 (t, *J*=6Hz, 2H), 6.80-6.90 (m, 1H), 4.31 (s, 2H), 3.60-3.70 (m, 2H), 2.38-2.39 (m, 2H), 1.83 (s, 3H) ppm . ESMS calcd. (C₂₂H₁₉ClF₃N₅O) 461.1; found: 425.2 (M-Cl)

Example 23: 2,4-difluoro-N-(5-(1-(3-fluoropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide hydrochloride(Compound 18)

10

ESMS calcd. (C₂₂H₁₉ClF₃N₅O) 461.1; found: 425.2 (M-Cl)

See procedure for 2,6-difluoro-N-(5-(4-methyl-1-(pyrazin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide.

15 Example 24: N-(5-(1-(3-cyanopyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide (Compound 19)

¹H NMR (300 MHz, CDCl₃) 9.68(s, 1H), 8.27 (s, 1H), 7.34 (dd, *J*=1.5, 3.6Hz, 1H), 8.16-8.17 (m, 1H), 7.78 (dd, *J*=1.5, 5.7Hz, 1H), 7.43-7.50 (m, 1H), 7.03 (t, *J*=6.0Hz, 2H), 6.73 (dd, *J*=3.6, 5.7Hz, 1H), 4.49-4.50 (m, 2H), 3.97 (t, *J*=4.2Hz, 2H), 2.49-2.50 (m, 2H), 1.82 (s, 3H), ppm ESMS calcd. (C₂₃H₁₈F₂N₆O) 432.2; found: 433.2 (M+H)

20

See procedure for 2,6-difluoro-N-(5-(4-methyl-1-(pyrazin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide

25 Example 25: N-(5-(1-(3,5-difluoropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide (Compound 20)

¹H NMR (300 MHz, CDCl₃) 9.67(s, 1H), 8.45 (s, 1H), 8.23 (d, *J*=1.2Hz, 1H), 7.94 (d, *J*=1.8Hz, 1H), 7.45-7.52 (m, 1H), 7.13 (ddd, *J*=1.8, 5.7, 9.0Hz, 1H), 7.04 (t, *J*=6.0Hz, 2H), 4.25-4.26 (m, 2H), 3.35 (t, *J*=4.2Hz, 2H), 2.41-2.50 (m, 2H), 1.82 (s, 3H), ppm ESMS calcd. (C₂₂H₁₇F₄N₅O) 443.1; found: 474.2 (M+H)

30

Example 26: N-(5-(1-(3,5-dichloropyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide (Compound 21)

35

¹H NMR (300 MHz, CDCl₃) 9.66(s, 1H), 8.48 (s, 1H), 8.22 (d, *J*=1.2Hz, 1H), 8.11 (d,

$J=1.8\text{Hz}$, 1H), 7.60 (d, $J=1.8\text{Hz}$, 1H), 7.44-7.51 (m, 1H), 7.04 (t, $J=6.3\text{Hz}$, 2H), 4.20-4.21 (m, 2H), 3.59 (t, $J=4.2\text{Hz}$, 2H), 2.45-2.46 (m, 2H), 1.83 (s, 3H), ppm ESMS calcd. ($\text{C}_{22}\text{H}_{17}\text{Cl}_2\text{F}_2\text{N}_5\text{O}$) 475.1; found: 476.2 (M+H)

5 Example 27: N-(5-(1-(3-bromopyridin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)-2,6-difluorobenzamide (Compound 22)

^1H NMR (300 MHz, $\text{CDCl}_3\text{-MeOD}$) 9.68 (s, 1H), 8.41-8.44(m, 2H), 8.36 (d, $J=0.9\text{Hz}$, 1H), 8.22(dd, $J=0.9$, 5.7Hz, 1H), 7.46-7.53 (m, 1H), 7.06 (t, $J=6.3\text{Hz}$, 2H), 6.99 (dd, $J=4.5$, 5.7Hz, 1H), 4.78 (s, 2H), 4.07 (t, $J=4.5\text{Hz}$, 2H), 2.70-2.75 (m, 2H), 1.91 (s, 3H) ppm ESMS calcd. ($\text{C}_{22}\text{H}_{18}\text{BrF}_2\text{N}_5\text{O}$) 485.1; found: 486.2 (M+H) See procedure for 2,6-difluoro-N-(5-(4-methyl-1-(pyrazin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide

15 Example 28: N-(4-(1-(3-aminopyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)-2,6-difluorobenzamide (Compound 23)

^1H NMR (400MHz, CDCl_3) δ (ppm), 8.32-6.80(m, 9H), 4.63(s, 2H), 3.39(m, 2H), 2.35(m, 2H), 1.46(s, 3H); ESMS clcd for $\text{C}_{23}\text{H}_{21}\text{F}_2\text{N}_5\text{O}$: 421.2; Found: 422.2 (M+H)⁺.

20 Example 29: 4-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)-N-(2,6-difluorobenzyl)aniline (Compound 24)

^1H NMR (400MHz, CDCl_3) δ (ppm), 8.03(s, 1H), 7.78(s, 1H), 7.32-6.60(m,7H), 4.33(s, 2H), 4.05(s,2H), 3.69(m, 2H), 2.36(m, 2H), 1.55(s, 3H); ESMS clcd for $\text{C}_{23}\text{H}_{21}\text{ClF}_2\text{N}_4\text{O}$: 426.1; Found: 427.1 (M+H)⁺.

Example 30: N-(4-(1-(3-cyanopyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)-2,6-difluorobenzamide (Compound 25)

30 ^1H NMR (400MHz, CDCl_3) δ (ppm), 8.36-6.71(m, 9H), 4.36(s, 2H), 3.04(m, 2H), 2.41(s, 2H), 1.63(s, 3H); ESMS clcd for $\text{C}_{24}\text{H}_{19}\text{F}_2\text{N}_5\text{O}$: 431.2; Found: 432.2(M+H)⁺.

Example 31: 2,6-difluoro-N-(4-(1-(3-fluoropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide (Compound 26)

35

^1H NMR (400MHz, CDCl_3) δ (ppm), 7.91-6.82 (m, 9H), 4.23(s, 2H), 3.70(m, 2H), 2.34(s,

2H), 1.65(s, 3H); ESMS clcd for C₂₃H₁₉F₃N₄O: 424.2; Found: 425.2(M+H)⁺.

Example 32: N-(4-(1-(3-chloropyrazin-2-yl)-4-methyl-1,2,5,6-tetrahydropyridin-3-yl)phenyl)-2,6-difluorobenzamide (Compound 27)

5

¹H NMR (400MHz,CD₃OD) δ (ppm), 8.24-7.02 (m, 9H), 4.14(s, 2H), 3.63(m, 2H), 2.41(s, 2H), 1.63(s, 3H); ESMS clcd for C₂₃H₁₉ClF₂N₄O: 440.1; Found: 441.1(M+H)⁺.

Example 33: 2,6-difluoro-N-(4-(4-methyl-1-(pyrazin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)phenyl)benzamide (Compound 28)

10

¹H NMR (400MHz,CD₃OD) δ (ppm), 8.34-6.97 (m, 9H), 4.22(s, 2H), 3.81(m, 2H), 2.41(s, 2H), 1.65(s, 3H); ESMS clcd for C₂₃H₂₀F₂N₄O: 406.2; Found: 407.2(M+H)⁺.

15

EXAMPLE A: INHIBITION OF IL-2 PRODUCTION

Jurkat cells were placed in a 96 well plate (0.5 million cells per well in 1% FBS medium) then a test compound of this invention was added at different concentrations. After 10 minutes, the cells were activated with PHA (final concentration 2.5 μg/mL) and incubated for 20 hours at 37°C under CO₂. The final volume was 200 μL. Following incubation, the cells were centrifuged and the supernatants collected and stored at -70°C prior to assaying for IL-2 production. A commercial ELISA kit (IL-2 Eli-pair, Diaclone Research, Besancon, France) was used to detect production of IL-2, from which dose response curves were obtained. The IC₅₀ value was calculated as the concentration at which 50% of maximum IL-2 production after stimulation was inhibited versus a non-stimulation control.

20

25

Compound #	IC ₅₀ (nM)
1	30
2	38
3	>1000
4	934
5	>1000
6	66

Inhibition of other cytokines, such as IL-4, IL-5, IL-13, GM-CSF, TNF-α, and INF-γ, can be tested in a similar manner using a commercially available ELISA kit for each cytokine.

EXAMPLE B: PATCH CLAMP STUDIES OF INHIBITION OF I_{CRAC} CURRENT IN RBL CELLS, JURKAT CELLS, AND PRIMARY T CELLS

In general, a whole cell patch clamp method was used to examine the effects of a compound of the invention on a channel that mediates I_{crac} . In such experiments, a baseline measurement was established for a patched cell. Then a compound to be tested was perfused (or puffed) to cells in the external solution and the effect of the compound on I_{crac} was measured. A compound that modulates I_{crac} (e.g., inhibits) is a compound that is useful in the invention for modulating CRAC ion channel activity.

1) RBL cells**Cells**

Rat basophilic leukemia cells (RBL-2H3) are grown in DMEM media supplemented with 10% fetal bovine serum in an atmosphere of 95% air/5% CO_2 . Cells are seeded on glass coverslips 1-3 days before use.

Recording Conditions

Membrane currents of individual cells are recorded using the whole-cell configuration of the patch clamp technique with an EPC10 (HEKA Elektronik, Lambrecht, Germany). Electrodes (2-5 $M\Omega$ in resistance) are fashioned from borosilicate glass capillary tubes (Sutter Instruments, Novato, Ca). The recordings are done at room temperature.

Intracellular pipette solution

The intracellular pipette solution contains Cs-Glutamate 120mM; CsCl 20mM; CsBAPTA 10mM; CsHEPES 10mM; NaCl 8mM; $MgCl_2$ 1mM; IP3 0.02mM; pH=7.4 adjusted with CsOH. The solution is kept on ice and shielded from light before the experiment is preformed.

Extracellular solution

The extracellular solution contains NaCl 138mM; NaHEPES, 10mM; CsCl 10mM; $CaCl_2$ 10mM; Glucose 5.5mM; KCl 5.4mM; KH_2PO_4 0.4mM; $Na_2HPO_4 \cdot H_2O$ 0.3mM at pH=7.4 adjusted with NaOH.

Compound treatment

Each test compound is diluted from a 10 mM stock in series using DMSO. The final DMSO

concentration is kept at 0.1 %.

Experimental procedure

5 I_{CRAC} currents are monitored every 2 seconds using a 50 msec protocol, where the voltage was ramped from -100 mV to +100 mV. The membrane potential is held at 0 mV between the test ramps. In a typical experiment, the peak inward currents are expected to develop within 50-100 seconds. Once the I_{CRAC} currents are stabilized, the cells are perfused with a test compound in the extracellular solution. At the end of an experiment, the remaining I_{CRAC} currents are then challenged with a control compound (SKF96365, 10 μ M) to ensure that the current could still be inhibited.

10 Data analysis

The I_{CRAC} current level is determined by measuring the inward current amplitude at -80 mV of the voltage ramp in an off-line analysis using MATLAB. The I_{CRAC} current inhibition for each concentration is calculated using peak amplitude in the beginning of the experiment from the same cell. The IC_{50} value and Hill coefficient for each compound is estimated by
15 fitting all the individual data points to a single Hill equation.

Results

Compounds of the invention are expected to inhibit I_{CRAC} current in RBL cells.

20 2) Jurkat cells

Cells

Jurkat T cells are grown on glass coverslips, transferred to the recording chamber and kept in a standard modified Ringer's solution of the following composition: NaCl 145mM, KCl 2.8mM, CsCl 10mM, CaCl₂ 10mM, MgCl₂ 2mM, glucose 10mM, HEPES·NaOH 10mM, pH
25 7.2.

Extracellular Solution

The external solution contains 10 mM CaNaR, 11.5 mM glucose and a test compound at various concentrations.

Intracellular Pipette Solution

30 The standard intracellular pipette solution contains: Cs-glutamate 145 mM, NaCl 8 mM, MgCl₂ 1 mM, ATP 0.5 mM, GTP 0.3 mM, pH 7.2 adjusted with CsOH. The solution is supplemented with a mixture of 10 mM Cs-BAPTA and 4.3-5.3 mM CaCl₂ to buffer $[Ca^{2+}]_i$ to resting levels of 100-150 nM.

Patch-clamp recordings

Patch-clamp experiments are performed in the tight-seal whole-cell configuration at 21-25°C. High-resolution current recordings are acquired by a computer-based patch-clamp amplifier system (EPC-9, HEKA, Lambrecht, Germany). Sylgard®- coated patch pipettes typically

- 5 have resistances between 2-4 MΩ after filling with the standard intracellular solution. Immediately following establishment of the whole-cell configuration, voltage ramps of 50 ms duration spanning the voltage range of -100 to +100 mV are delivered from a holding potential of 0 mV at a rate of 0.5 Hz over a period of 300 to 400 seconds. All voltages are corrected for a liquid junction potential of 10 mV between external and internal solutions.
- 10 Currents are filtered at 2.3 kHz and digitized at 100 μs intervals. Capacitive currents and series resistance are determined and corrected before each voltage ramp using the automatic capacitance compensation of the EPC-9.

Data analysis

- The very first ramps before activation of I_{CRAC} (usually 1 to 3) are digitally filtered at 2 kHz, pooled and used for leak-subtraction of all subsequent current records. The low-resolution
- 15 temporal development of inward currents is extracted from the leak-corrected individual ramp current records by measuring the current amplitude at -80 mV or a voltage of choice.

Compounds of the invention are expected to inhibit I_{CRAC} current in Jurkat cells.

20

3) Primary T Cells

Preparation of Primary T Cells

- Primary T cells are obtained from human whole blood samples by adding 100 μL of
- 25 RosetteSep® human T cell enrichment cocktail to 2 mL of whole blood. The mixture is incubated for 20 minutes at room temperature, then diluted with an equal volume of PBS containing 2% FBS. The mixture is layered on top of RosetteSep® DM-L density medium and then centrifuged for 20 minutes at 1200 g at room temperature. The enriched T cells are recovered from the plasma/density medium interface, then washed with PBS containing 2%
- 30 FBS twice, and used in patch clamp experiments following the procedure described for RBL cells.

Compounds of the invention are expected to inhibit I_{CRAC} current in human primary T cells.

- 35 **EXAMPLE C: INHIBITION OF MULTIPLE CYTOKINES IN PRIMARY HUMAN PBMCs**

Peripheral blood mononuclear cells (PBMCs) are stimulated with phytohemagglutinin (PHA) in the presence of varying concentrations of compounds of the invention or cyclosporine A (CsA), a known inhibitor of cytokine production. Cytokine production is measured using commercially available human ELISA assay kits (from Cell Science, Inc.) following the
5 manufacturers instructions.

The compounds of the invention are potent inhibitors of IL-2, and are expected to be potent inhibitors of IL-4, IL-5, IL-13, GM-CSF, INF- α and TNF- γ in primary human PBM cells. In addition, compounds of the invention are not expected to inhibit the anti-inflammatory
10 cytokine, IL-10.

EXAMPLE D: COMPOUNDS OF THE INVENTION ARE POTENT INHIBITORS OF DEGRANULATION IN RBL CELLS

Procedure:

The day before the assay is performed, RBL cells, that had been grown to confluence in a 96 well plate, are incubated at 37°C for at least 2 hours. The medium is replaced in each well with 100 μ L of fresh medium containing 2 μ Lg/mL of anti-DNP IgE.

20 On the following day, the cells are washed once with PRS (2.6 mM glucose and 0.1% BSA) and 160 μ L of PRS was added to each well. A test compound is added to a well in a 20 μ L solution at 10X of the desired concentration and incubated for 20 to 40 minutes at 37°C. 20 μ L of 10X mouse anti-IgE (10 μ L/mL) is added. SKF96365 is used as a positive control. Maximum degranulation typically occurs between 15 to 40 minutes after addition of anti-IgE.

Results:

Compounds of the invention are expected to inhibit degranulation of RBL cells.

EXAMPLE E: COMPOUNDS OF THE INVENTION ARE POTENT INHIBITORS OF CHEMOTAXIS IN T CELLS

T-cell isolation:

25 Twenty ml aliquots of heparinized whole blood (2 pig, 1 human) are subjected to density gradient centrifugation on Ficoll Hypaque. The buffy coat layers representing peripheral blood mononuclear cells (PBMCs) containing lymphocytes and monocytes are washed once,
35 resuspended in 12 ml of incomplete RPMI 1640 and then placed in gelatin-coated T75 culture flasks for 1 hr at 37°C. The non-adherent cells, representing peripheral blood lymphocytes

(PBLs) depleted of monocytes, are resuspended in complete RPMI media and placed in loosely packed activated nylon wool columns that had been equilibrated with warm media. After 1 hr at 37°C, the non-adherent T cell populations are eluted by washing of the columns with additional media. The T cell preparations are centrifuged, resuspended in 5 ml of
5 incomplete RPMI, and counted using a hemocytometer.

Cell migration assay:

Aliquots of each T cell preparation are labeled with Calciem AM (TefLabs) and suspended at a concentration of 2.4×10^6 /ml in HEPES-buffered Hank's Balanced Salt Solution containing
10 1.83 mM CaCl_2 and 0.8 mM MgCl_2 , pH 7.4 (HHBSS). An equal volume of HHBSS containing 0, 20 nM, 200 nM or 2000 nM of compound 1 or 20 nM EDTA is then added and the cells are incubated for 30 min at 37°C. Fifty μl aliquots of the cell suspensions (60,000 cells) are placed on the membrane (pore size 5 μm) of a Neuroprobe ChemoTx 96 well chemotaxis unit that had been affixed over wells containing 10 ng/ml MIP-1 α in HHBSS.
15 The T cells are allowed to migrate for 2 hr at 37°C, after which the apical surface of the membrane is wiped clean of cells. The chemotaxis units are then placed in a CytoFlour 4000 (PerSeptive BioSystems) and the fluorescence of each well is measured (excitation and emission wavelengths of 450 and 530 nm, respectively). The number of migrating cells in each well is determined from a standard curve generated from measuring the fluorescence of
20 serial two-fold dilutions of the labeled cells placed in the lower wells of the chemotaxis unit prior to affixing the membrane.

Results: Compounds of the invention are expected to be inhibitory to the chemotactic response of porcine T cells and in human T cells.

25 All publications, patent applications, patents, and other documents cited herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting in any way.


30 From the foregoing description, it will be apparent that variations and modifications may be made to the invention described herein. Such embodiments are also within the scope of the following claims.

CLAIMS

WE CLAIM:

1. A compound represented by the following structural formula:

5



(I)

or a pharmaceutically acceptable salt thereof, wherein:

X_1 and X_2 are, independently, CH or N;

10

one of X_3 or X_6 is NR_{29} , and the other is CH_2 , CHR_2 , or $\text{C}(\text{R}_2)_2$;

L is a linker selected from the group consisting of $-\text{NHCH}_2-$, $-\text{CH}_2\text{NH}-$, $-\text{NR}_5-\text{C}(\text{O})-$, $-\text{C}(\text{O})-\text{NR}_5-$, $-\text{NR}_5-\text{C}(\text{S})-$, $-\text{C}(\text{S})-\text{NR}_5-$, $-\text{NR}_5-\text{C}(\text{NR}_8)-$, $-\text{C}(\text{NR}_8)-\text{NR}_5-$, $-\text{NR}_5\text{C}(\text{S})\text{NR}_5-$, $-\text{NR}_5\text{C}(\text{NR}_8)\text{NR}_5-$, or $-\text{NR}_5\text{CR}^a\text{R}^b\text{NR}_5-$;

Y is a substituted phenyl, provided that Y is not substituted with $-\text{NHC}(\text{O})\text{R}_{30}$;

15

R_1 is H or a substituent;

R_2 , for each occurrence is, independently, a substituent, provided that R_2 is not $=\text{O}$ or $=\text{C}(\text{R}_{28})_2$;

R_4 is a substituent;

20

R^a and R^b , for each occurrence, are independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, cyano, nitro, halo, $-\text{OR}_5$, $-\text{SR}_5$, $-\text{NR}_6\text{R}_7$, $-\text{C}(\text{O})\text{NR}_6\text{R}_7$, $-\text{NR}_5\text{C}(\text{O})\text{R}_5$, $-\text{C}(\text{O})\text{R}_5$, $-\text{C}(\text{O})\text{OR}_5$, $-\text{OC}(\text{O})\text{R}_5$, $-\text{C}(\text{O})\text{SR}_5$, $-\text{SC}(\text{O})\text{R}_5$, $-\text{C}(\text{S})\text{NR}_6\text{R}_7$, $-\text{NR}_5\text{C}(\text{S})\text{R}_5$, $-\text{C}(\text{S})\text{R}_5$, $-\text{C}(\text{S})\text{OR}_5$, $-\text{OC}(\text{S})\text{R}_5$, $-\text{C}(\text{S})\text{SR}_5$, $-\text{SC}(\text{S})\text{R}_5$, $-\text{C}(\text{NR}_8)\text{NR}_6\text{R}_7$, $-\text{NR}_5\text{C}(\text{NR}_8)\text{R}_5$, $-\text{C}(\text{NR}_8)\text{R}_5$, $-\text{C}(\text{NR}_8)\text{OR}_5$, $-\text{OC}(\text{NR}_8)\text{R}_5$, $-\text{C}(\text{NR}_8)\text{SR}_5$, $-\text{SC}(\text{NR}_8)\text{R}_5$, $-\text{OC}(\text{O})\text{OR}_5$, $-\text{OC}(\text{O})\text{NR}_6\text{R}_7$, $-\text{NR}_5\text{C}(\text{O})\text{OR}_5$, $-\text{NR}_5\text{C}(\text{O})\text{NR}_6\text{R}_7$, $-\text{SC}(\text{O})\text{OR}_5$, $-\text{SC}(\text{O})\text{NR}_6\text{R}_7$, $-\text{SC}(\text{O})\text{SR}_5$, $-\text{NR}_5\text{C}(\text{O})\text{SR}_5$, $-\text{OC}(\text{O})\text{SR}_5$, $-\text{OC}(\text{S})\text{OR}_5$, $-\text{OC}(\text{S})\text{NR}_6\text{R}_7$, $-\text{NR}_5\text{C}(\text{S})\text{OR}_5$, $-\text{NR}_5\text{C}(\text{S})\text{NR}_6\text{R}_7$, $-\text{SC}(\text{S})\text{OR}_5$, $-\text{SC}(\text{S})\text{NR}_6\text{R}_7$, $-\text{SC}(\text{S})\text{SR}_5$, $-\text{NR}_5\text{C}(\text{S})\text{SR}_5$, $-\text{OC}(\text{S})\text{SR}_5$,

30

OC(NR₈)OR₅, -OC(NR₈)NR₆R₇, -NR₅C(NR₈)OR₅, -NR₅C(NR₈)NR₆R₇, -
 SC(NR₈)OR₅, -SC(NR₈)NR₆R₇, -SC(NR₈)SR₅, -NR₅C(NR₈)SR₅, or -OC(NR₈)SR₅;

R₅, for each occurrence, is independently, H, an optionally substituted alkyl,
 an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally
 5 substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally
 substituted heterocyclyl, an optionally substituted aryl, an optionally substituted
 heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

R₆ and R₇, for each occurrence are, independently, H, an optionally
 substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl,
 10 an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an
 optionally substituted heterocyclyl, an optionally substituted aryl, an optionally
 substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted
 heteraralkyl; or R₆ and R₇ taken together with the nitrogen to which they are attached
 are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

R₈, for each occurrence, is independently -H, a halo, an
 15 alkyl, -OR₅, -NR₆R₇, -C(O)R₅, -C(O)OR₅, or -C(O)NR₆R₇;

R₂₈, for each occurrence is, independently, H or a substituent;

R₂₉ is -C(O)R₃₀ or an optionally substituted N-containing 6-membered
 aromatic ring;

R₃₀ is -H, -OR₅, -SR₅, -NR₆R₇, an optionally substituted alkyl, an optionally
 20 substituted alkenyl, an optionally substituted alkynyl, an optionally substituted
 cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted
 heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an
 optionally substituted aralkyl, or an optionally substituted heteraralkyl;

m is 0, 1, or 2;

n is 0, 1 or 2; and

with the proviso that the compound is not:

5-[4-(2,6-difluoro-benzoylamino)-phenyl]-4-methyl-2,4-dihydro-2H-
 pyridine-1-carboxylic acid ethyl ester;

30 Ethyl 3-(4-(2,6-difluorobenzamido)phenyl)-4-methyl-5,6-dihydropyridine-
 1(2H)-carboxylate;

Ethyl 4-(4-(2,6-difluorobenzamido)phenyl)-3-methyl-5,6-dihydropyridine-
 1(2H)-carboxylate;

2,6-Difluoro-N-[4-(1-methoxycarbonyl-4-thiazol-2-yl)-1,2,5,6-tetrahydro-
 35 pyridin-3-yl]-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-methoxycarbonyl-4-isopropyl-1,2,5,6-tetrahydro-

pyridin-3-yl)-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-methoxycarbonyl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-allyloxycarbonyl-3-methyl-1,2,5,6-tetrahydro-pyridin-4-yl)-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-phenyloxycarbonyl-3-methyl-1,2,5,6-tetrahydro-pyridin-4-yl)-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-acetyl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide;

2,6-Difluoro-N-[4-(1-cyclopropylcarbonyl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide;

2,6-Difluoro-N-{4-[1-(pyridine-3-carbonyl)-cyclopropanecarbonyl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl]-phenyl}-benzamide; or

2,6-Difluoro-N-[4-(1-pyrimidin-2-yl-4-methyl-1,2,5,6-tetrahydro-pyridin-3-yl)-phenyl]-benzamide.

2. The compound of Claim 1, wherein Y is substituted with one to two substituents.

3. The compound of Claim 2, wherein the one to two substituents are each independently a lower alkyl or a halo.

4. The compound of Claim 3, wherein Y is a difluorophenyl.



5. The compound of Claim 4, wherein Y is 2,6-difluorophenyl.

6. The compound of Claim 1, wherein L is -NHCH₂-, -CH₂NH-, -NH-C(O)-, or -C(O)-NH-.

7. The compound of Claim 6, wherein L is -NHC(O)-.

8. The compound of Claim 1, wherein R₁ is selected from the group consisting of a halo, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl.

9. The compound of Claim 8, wherein R_1 is selected from the group consisting of halo or an optionally substituted lower alkyl.
10. The compound of Claim 1, wherein X_3 is CH_2 and X_6 is NR_{29} .
- 5 11. The compound of Claim 1, wherein X_6 is CH_2 and X_3 is NR_{29} .
12. The compound of Claim 1, wherein X_1 and X_2 are both CH.
- 10 13. The compound of Claim 1, wherein X_1 and X_2 are both N.
14. The compound of Claim 1, wherein X_1 is N and X_2 is CH.
15. The compound of Claim 1, wherein m is 0.
- 15 16. The compound of Claim 1, 10, 11, 12, 13, or 14 wherein
Y is a difluorophenyl;
L is $-NHC(O)-$;
 R_1 is selected from the group consisting of halo or an optionally substituted
20 lower alkyl; and
m is 0.
17. The compound of Claim 1, wherein the compound is represented by one of the following structures:



wherein

R_{31} is halo, an optionally substituted alkyl, an optionally substituted alkenyl,
 5 an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally
 substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally
 substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl,
 or an optionally substituted heteraralkyl; and

p is 1, 2, 3, 4 or 5.

18. The compound of Claim 17, wherein p is 1 or 2.

19. The compound of Claim 18, wherein each R_{31} is independently a lower alkyl or a
 halo.


20. The compound of Claim 17, wherein L is $-NHCH_2-$, $-CH_2NH-$,
 $-NH-C(O)-$, or $-C(O)-NH-$.

21. The compound of Claim 20, wherein L is $-NHC(O)-$.

22. The compound of Claim 17, wherein R_1 is selected from the group consisting of a
 halo, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally
 substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted
 cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an


optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl.

- 5 23. The compound of Claim 22, wherein R₁ is selected from the group consisting of halo or an optionally substituted lower alkyl.
24. The compound of Claim 17, wherein X₁ and X₂ are both CH.
- 10 25. The compound of Claim 17, wherein X₁ and X₂ are both N.
26. The compound of Claim 17, wherein X₁ is N and X₂ is CH.
- 15 27. The compound of Claim 17, wherein R₃₀ is selected from the group consisting of -OR₅, optionally substituted alkyl, optionally substituted aryl, and optionally substituted heteroaryl.
28. The compound of Claim 27, wherein R₃₀ is selected from the group consisting of -OR₅ and optionally substituted heteroaryl.
- 20 29. The compound of Claim 28, wherein R₃₀ is -OR₅ and R₅ is a lower alkyl.
30. The compound of Claim 28, wherein R₃₀ is a pyridine ring.
- 25 31. The compound of Claim 17, wherein the compound is represented by one of the following structures:




32. The compound of Claim 31, wherein each R_{31} is independently a lower alkyl or a halo.
- 5
33. The compound of Claim 32, wherein each R_{31} is halo.
34. The compound of Claim 33, wherein each R_{31} is F.
- 10
35. The compound of Claim 31, wherein L is $-NHCH_2-$, $-CH_2NH-$, $-NH-C(O)-$, or $-C(O)-NH-$.
36. The compound of Claim 35, wherein L is $-NHC(O)-$.
- 15
37. The compound of Claim 31, wherein R_1 is selected from the group consisting of a halo, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl.
- 20

38. The compound of Claim 37, wherein R_1 is selected from the group consisting of halo or an optionally substituted lower alkyl.
39. The compound of Claim 31, wherein X_1 and X_2 are both CH.
- 5 40. The compound of Claim 31, wherein X_1 and X_2 are both N.
41. The compound of Claim 31, wherein X_1 is N and X_2 is CH.
- 10 42. The compound of Claim 31, wherein R_{30} is selected from the group consisting of - OR_5 , optionally substituted alkyl, optionally substituted aryl, and optionally substituted heteroaryl.
- 15 43. The compound of Claim 42, wherein R_{30} is selected from the group consisting of - OR_5 and optionally substituted heteroaryl.
44. The compound of Claim 43, wherein R_{30} is - OR_5 and R_5 is a lower alkyl.
45. The compound of Claim 43, wherein R_{30} is a pyridine ring.
- 20 46. The compound of Claim 31, 39, 40, or 41, wherein:
each R_{31} is halo;
L is -NHC(O)-;
 R_1 is selected from the group consisting of halo or an optionally substituted
25 lower alkyl; and
 R_{30} is selected from the group consisting of - OR_5 and optionally substituted heteroaryl.
47. The compound of Claim 46, wherein each R_{31} is F.
- 30 48. The compound of Claim 1, wherein the compound is represented by one of the following structures:



or



wherein

each of X₇, X₈, X₉, X₁₀, and X₁₁ are independently CH, CR₂, or N, provided
 5 that at least one of X₇, X₈, X₉, X₁₀, or X₁₁ is N;

R₃₁ is halo, an optionally substituted alkyl, an optionally substituted alkenyl,
 an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally
 substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally
 substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl,
 10 or an optionally substituted heteraralkyl; and

p is 1, 2, 3, 4 or 5.

49. The compound of Claim 48, wherein p is 1 or 2.


15 50. The compound of Claim 49, wherein each R₃₁ is independently a lower alkyl or a halo.

51. The compound of Claim 48, wherein L is -NHCH₂-, -CH₂NH-,
 -NH-C(O)-, or -C(O)-NH-.


20

52. The compound of Claim 51, wherein L is -NHC(O)-.

53. The compound of Claim 48, wherein R_1 is selected from the group consisting of a halo, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl.
54. The compound of Claim 53, wherein R_1 is selected from the group consisting of halo or an optionally substituted lower alkyl.
55. The compound of Claim 48, wherein X_1 and X_2 are both CH.
56. The compound of Claim 48, wherein X_1 and X_2 are both N.
57. The compound of Claim 48, wherein X_1 is N and X_2 is CH.
58. The compound of Claim 48, wherein two of X_7 , X_8 , X_9 , X_{10} , and X_{11} are N.
59. The compound of Claim 48, wherein one of X_7 , X_8 , X_9 , X_{10} , and X_{11} is N.
60. The compound of Claim 48, wherein the compound is represented by one of the following structures:



or



61. The compound of Claim 60, wherein each R_{31} is independently a lower alkyl or a halo.

5

62. The compound of Claim 61, wherein each R_{31} is halo.

63. The compound of Claim 62, wherein each R_{31} is F.

10 64. The compound of Claim 60, wherein L is $-NHCH_2-$, $-CH_2NH-$, $-NH-C(O)-$, or $-C(O)-NH-$.

65. The compound of Claim 64, wherein L is $-NHC(O)-$.

15 66. The compound of Claim 60, wherein R_1 is selected from the group consisting of a halo, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl.

20

67. The compound of Claim 66, wherein R₁ is selected from the group consisting of halo or an optionally substituted lower alkyl.
68. The compound of Claim 60, wherein X₁ and X₂ are both CH.
- 5 69. The compound of Claim 60, wherein X₁ and X₂ are both N.
70. The compound of Claim 60, wherein X₁ is N and X₂ is CH.
- 10 71. The compound of Claim 60, wherein two of X₇, X₈, X₉, X₁₀, and X₁₁ are N.
72. The compound of Claim 60, wherein one of X₇, X₈, X₉, X₁₀, and X₁₁ is N.
73. The compound of Claim 60, 68, 69, or 70, wherein:
- 15 each R₃₁ is halo;
 L is -NHC(O)-;
 R₁ is selected from the group consisting of halo or an optionally substituted lower alkyl; and
 two of X₇, X₈, X₉, X₁₀, and X₁₁ are N.
- 20 74. The compound of Claim 73, wherein each R₃₁ is F.
75. The compound of Claim 1, wherein the compound is selected from the group consisting of:
- ethyl 3-(6-(2,6-difluorobenzamido)pyridin-3-yl)-4-methyl-5,6-dihydropyridine-1(2H)-carboxylate;
- allyl 3-(acetoxymethyl)-4-(4-(2,6-difluorobenzamido)phenyl)-5,6-dihydropyridine-1(2H)-carboxylate;
- 2,6-difluoro-N-(5-(4-methyl-1-picolinoyl-1,2,5,6-tetrahydropyridin-3-yl)pyridin-2-yl)benzamide hydrochloride;
- ethyl 3-(acetoxymethyl)-4-(4-(2,6-difluorobenzamido)phenyl)-5,6-dihydropyridine-1(2H)-carboxylate or
- ethyl 4-chloro-3-(4-(2,6-difluorobenzamido)phenyl)-5,6-dihydropyridine-1(2H)-carboxylate.
- 25

76. The compound of Claim 1, wherein the compound is 2,6-difluoro-N-(5-(4-methyl-1-(pyrimidin-2-yl)-1,2,5,6-tetrahydropyridin-3-yl)pyrazin-2-yl)benzamide hydrochloride.
- 5 77. A pharmaceutical composition, comprising a pharmaceutically acceptable carrier and a compound of any one of Claims 1 through 76.
78. The pharmaceutical composition of Claim 77, further comprising one or more additional therapeutic agents.
- 10 79. The pharmaceutical composition according to Claim 78, wherein the additional therapeutic agent is selected from the group consisting of immunosuppressive agents, anti-inflammatory agents and suitable mixtures thereof.
- 15 80. The pharmaceutical composition of Claim 79, wherein the additional therapeutic agent is selected from the group consisting of steroids, non-steroidal anti-inflammatory agents, antihistamines, analgesics, and suitable mixtures thereof.
81. A method of inhibiting immune cell activation comprising administering to the cell a
20 compound of any one of Claims 1 through 76.
82. The method of Claim 81, wherein immune cell activation is inhibited in a subject by administering the compound to the subject.
- 25 83. The method of Claim 82, wherein the subject is human.
84. A method of inhibiting cytokine production in a cell, comprising administering to the cell a compound of any one of Claims 1 through 76.
- 30 85. The method of Claim 84, wherein cytokine production is inhibited in a subject by administering the compound to the subject.
86. The method of Claim 85, wherein the subject is human.
- 35 87. The method of Claim 85, wherein the cytokine is selected from the group consisting of IL-2, IL-4, IL-5, IL-13, GM-CSF, IFN- γ , TNF- α , and combinations thereof.

88. The method of Claim 87, wherein the cytokine is IL-2.
89. A method of modulating an ion channel in a cell, wherein the ion channel is involved in immune cell activation, comprising administering to the cell a compound of any one of Claims 1 through 76.
90. The method of Claim 89, wherein the ion channel is in a subject and it is modulated by administering the compound to the subject.
91. The method of Claim 90, wherein the subject is human.
92. The method of Claim 90, wherein the ion channel is a Ca²⁺-release-activated Ca²⁺ channel (CRAC).
93. A method of inhibiting T-cell and/or B-cell proliferation in response to an antigen, comprising administering to the cell a compound of any one of Claims 1 through 76.
94. The method of Claim 93, wherein T-cell and/or B-cell proliferation is inhibited in a subject by administering the compound to the subject.
95. The method of Claim 94, wherein the subject is human.
96. A method for treating or preventing an immune disorder in a subject in need thereof, comprising administering to the subject an effective amount of a compound of any one of Claims 1 through 76.
97. The method of Claim 96, wherein the subject is human.
98. The method of Claim 96, wherein the disorder is selected from the group consisting of multiple sclerosis, myasthenia gravis, Guillain-Barré, autoimmune uveitis, autoimmune hemolytic anemia, pernicious anemia, autoimmune thrombocytopenia, temporal arteritis, anti-phospholipid syndrome, vasculitides such as Wegener's granulomatosis, Behcet's disease, psoriasis, dermatitis herpetiformis, pemphigus vulgaris, vitiligo, Crohn's disease, ulcerative colitis, primary biliary cirrhosis, autoimmune hepatitis, Type 1 or immune-mediated diabetes mellitus, Grave's disease, Hashimoto's thyroiditis, autoimmune oophoritis and orchitis, autoimmune disorder of

the adrenal gland, rheumatoid arthritis, systemic lupus erythematosus, scleroderma, polymyositis, dermatomyositis, ankylosing spondylitis, and Sjogren's syndrome.

- 5 99. A method for treating or preventing an inflammatory condition in a subject in need thereof, comprising administering to the subject an effective amount of a compound of any one of Claims 1 through 76.
100. The method of Claim 99, wherein the subject is human.
- 10 101. The method according to Claim 99, wherein the disorder is selected from transplant rejection, skin graft rejection, arthritis, rheumatoid arthritis, osteoarthritis and bone diseases associated with increased bone resorption; inflammatory bowel disease, ileitis, ulcerative colitis, Barrett's syndrome, Crohn's disease; asthma, adult respiratory distress syndrome, chronic obstructive airway disease; corneal dystrophy, 15 trachoma, onchocerciasis, uveitis, sympathetic ophthalmitis, endophthalmitis; gingivitis, periodontitis; tuberculosis; leprosy; uremic complications, glomerulonephritis, nephrosis; sclerodermatitis, psoriasis, eczema; chronic demyelinating diseases of the nervous system, multiple sclerosis, AIDS-related neurodegeneration, Alzheimer's disease, infectious meningitis, encephalomyelitis, 20 Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis viral or autoimmune encephalitis; autoimmune disorders, immune-complex vasculitis, systemic lupus and erythematoses; systemic lupus erythematosus (SLE); cardiomyopathy, ischemic heart disease hypercholesterolemia, atherosclerosis, preeclampsia; chronic liver failure, brain and spinal cord trauma, and cancer.
- 25 102. A method for suppressing the immune system of a subject in need thereof, comprising administering to the subject an effective amount of a compound of any one of Claims 1 through 76.
- 30 103. The method of Claim 102, wherein the subject is human.
104. A method for treating or preventing an allergic disorder in a subject in need thereof, comprising administering to the subject an effective amount of a compound of any one of Claims 1 through 76.
- 35 105. The method of Claim 104, wherein the subject is human.

106. The method of Claim 104, wherein the disorder is allergic rhinitis, sinusitis, rhinosinusitis, chronic otitis media, recurrent otitis media, drug reactions, insect sting reactions, latex reactions, conjunctivitis, urticaria, anaphylaxis reactions, anaphylactoid reactions, atopic dermatitis, asthma, or food allergies.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 08/09346

A. CLASSIFICATION OF SUBJECT MATTER
 IPC(8) - C07D 213/04, 213/22 (2008.04)
 USPC - 546/255, 257
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 USPC- 546/255, 257

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
 USPC- 514/264.1, 293, 557; 546/193 (text search-see search terms below)


Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 PubWEST (USPT, PGPB, EPAB, JPAB) and Google Patent/Scholar
 Search terms: benzamide, pyrimidine, pyridine, pyrazine, inflammation, immune disorder

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2006/0173006 A1 (Sun et al.) 03 August 2006 (03.08.2006) para [0013], [0015]-[0016], [0017], [0019]-[0022], [0024], [0026]-[0027], [0034]	1-76
Y	US 2006/0173021 A1 (Sun et al.) 03 August 2006 (03.08.2006) para [0012]-[0013], [0018], [0074], [0084]	1-76

Further documents are listed in the continuation of Box C.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier application or patent but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 14 October 2008 (14.10.2008)	Date of mailing of the international search report 05 NOV 2008
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US, Commissioner for Patents P.O. Box 1450, Alexandria, Virginia 22313-1450 Facsimile No. 571-273-3201	Authorized officer:  Lee W. Young PCT Helpdesk: 571-272-4300 PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 08/09346

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.: 77-105
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.