



(86) Date de dépôt PCT/PCT Filing Date: 2002/04/05
(87) Date publication PCT/PCT Publication Date: 2002/10/17
(45) Date de délivrance/Issue Date: 2012/12/04
(85) Entrée phase nationale/National Entry: 2003/10/03
(86) N° demande PCT/PCT Application No.: EP 2002/003810
(87) N° publication PCT/PCT Publication No.: 2002/081513
(30) Priorité/Priority: 2001/04/06 (FR01/04712)

(51) Cl.Int./Int.Cl. *C12N 15/12* (2006.01),
C07K 14/47 (2006.01), *C07K 16/18* (2006.01),
C12N 15/63 (2006.01), *C12N 5/10* (2006.01),
C12Q 1/68 (2006.01), *G01N 33/53* (2006.01),
G01N 33/68 (2006.01)

(72) Inventeurs/Inventors:
LEVEILLARD, THIERRY, FR;
SAHEL, JOSE ALAIN, FR;
MOHAND-SAID, SADDEK, FR;
HICKS, DAVID, FR

(73) Propriétaires/Owners:
NOVARTIS AG, CH;
UNIVERSITE DE STRASBOURG, FR

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : PROTEINE ASSOCIEE A UNE MALADIE
(54) Title: DISEASE-ASSOCIATED PROTEIN

(57) **Abrégé/Abstract:**

Disclosed are methods and compositions for early diagnosis, monitoring and treatment of retinal dystrophy, age -related macular degeneration, Bardet-Biedel syndrome, Bassen-kornzweig syndrome, best disease, chroidema, gyrate atrophy, congenital amourosis, refsun syndrome, stargardt disease and Usher syndrome. In particular, the invention relates to a protein, termed 2Rdcvf1," that is differentially transcribed and expressed in subjects suffering from retinal dystrophies and the like, such as retinal dystrophy and age -related macular degeneration compared with non- sufferers, antibodies which recongnize this protein, and methods for diagnosing such conditions.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 October 2002 (17.10.2002)

PCT

(10) International Publication Number
WO 02/081513 A3

(51) International Patent Classification⁷: **C12N 15/12**,
C07K 14/47, C12N 15/63, 5/10, C07K 16/18, C12Q 1/68,
G01N 33/53, 33/68

Bernegger, F-67000 Strasbourg (FR). **MOHAND-SAID, Saddek** [FR/FR]; 14, rue des Horticulteurs, F-67000 Strasbourg (FR). **HICKS, David** [GB/FR]; 4, rue du Maréchal Joffre, F-67000 Strasbourg (FR).

(21) International Application Number: PCT/EP02/03810

(74) Agent: **GROS, Florent**; Novartis AG, Corporate Intellectual Property, Patent & Trademark Department, CH-4002 Basel (CH).

(22) International Filing Date: 5 April 2002 (05.04.2002)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LT, LU, LV, MA, MD, MK, MN, MX, NO, NZ, OM, PH, PL, PT, RO, RU, SE, SG, SI, SK, TJ, TM, TN, TR, TT, UA, US, UZ, VN, YU, ZA, ZW.

(26) Publication Language: English

(30) Priority Data:
01/04712 6 April 2001 (06.04.2001) FR

(71) Applicant (*for all designated States except AT, US*): **NOVARTIS AG** [CH/CH]; Lichtstrasse 35, 4065 Basel (CH).

(84) Designated States (*regional*): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).

(71) Applicant (*for AT only*): **NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT M.B.H.** [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).

Published:

— with international search report

(71) Applicant (*for all designated States except US*): **UNIVERSITE LOUIS PASTEUR** [FR/FR]; 4, rue Blaise-Pascal, F-67000 Strasbourg (FR).

(88) Date of publication of the international search report:
1 May 2003

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **LEVEILLARD, Thierry** [FR/FR]; 17, rue des Champs, F-67201 Eckolsheim (FR). **SAHEL, José, Alain** [FR/FR]; 2, rue

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DISEASE-ASSOCIATED PROTEIN

(57) Abstract: Disclosed are methods and compositions for early diagnosis, monitoring and treatment of retinal dystrophy, age-related macular degeneration, Bardet-Biedel syndrome, Bassen-kornzweig syndrome, best disease, chroidema, gyrate atrophy, congenital amourosis, refsun syndrome, stargardt disease and Usher syndrome. In particular, the invention relates to a protein, termed 2Rdcvf1," that is differentially transcribed and expressed in subjects suffering from retinal dystrophies and the like, such as retinal dystrophy and age-related macular degeneration compared with non-sufferers, antibodies which recognize this protein, and methods for diagnosing such conditions.



WO 02/081513 A3

DISEASE-ASSOCIATED PROTEIN

FIELD OF THE INVENTION

This invention relates to methods and compositions for detection and treatment of retinal degenerative diseases. In particular, the invention relates to a protein that protects against cone degeneration, nucleic acid molecules that encode such a protein, antibodies that recognize the protein, and methods for diagnosing retinal degenerative diseases.

BACKGROUND OF THE INVENTION

Photoreceptors are a specialized subset of retinal neurons, that are responsible for vision. Photoreceptors consist of rods and cones which are the photosensitive cells of the retina. Each rod and cone elaborates a specialized cilium, referred to as an outer segment, that houses the phototransduction machinery. The rods contain a specific light-absorbing visual pigment, rhodopsin. There are three classes of cones in humans, characterized by the expression of distinct visual pigments: the blue cone, green cone and red cone pigments. Each type of visual pigment protein is tuned to absorb light maximally at different wavelengths. The rod rhodopsin mediates scotopic vision (in dim light), whereas the cone pigments are responsible for photopic vision (in bright light). The red, blue and green pigments also form the basis of color vision in humans. The visual pigments in rods and cones respond to light and generate an action potential in the output cells, the rod bipolar neurons, which is then relayed by the retinal ganglion neurons to produce a visual stimulus in the visual cortex.

In humans, a number of diseases of the retina involve the progressive degeneration and eventual death of photoreceptors, leading inexorably to blindness. Degeneration of photoreceptors, such as by inherited retinal dystrophies (e.g., retinitis pigmentosa), age-related macular degeneration and other maculopathies, or retinal detachment, are all characterized by the progressive atrophy and loss of function of photoreceptor outer segments. In addition, death of photoreceptors or loss of photoreceptor function results in partial deafferentation of second order retinal neurons (rod bipolar cells and horizontal cells) in patients with retinal dystrophies, thereby decreasing the overall efficiency of the

propagation of the electrical signal generated by photoreceptors. Secondary glial and pigment epithelium changes secondary to photoreceptors degeneration result in vascular changes leading to ischemia and gliosis. Trophic factors that are capable of rescuing photoreceptors from cell death and/or restoring the function of dysfunctional (atrophic or dystrophic) photoreceptors may represent useful therapies for the treatment of such conditions.

The progression of these conditions points to a sequential loss of the two classes of photoreceptors: initially rods are lost as a direct result of a genetic or environmental or unknown lesion, resulting in night blindness and a reduction in visual field followed inevitably by loss of cones leading to total blindness. Thus, cones die indirectly since they do not express the primary lesion.

Not all of the genes associated with retinal dystrophy have yet been identified. Identification of such genes would make possible to both diagnose the disease and identify effective therapies.

SUMMARY OF THE INVENTION

The invention relates generally to a novel gene family, Rod-derived Cone Viability Factor (Rdcvf). In a first aspect, the invention provides an isolated polypeptide with an amino acid sequence as set forth in SEQ ID NO:2 or SEQ ID NO:4. Such a polypeptide, or fragments thereof, is found in the eye of sufferers of retinal dystrophies to a much lesser extent than in the eye of individuals without retinal dystrophy. Fragments of the isolated polypeptide with an amino acid sequence as set forth in SEQ ID NO:2 or SEQ ID NO:4 will comprise polypeptides comprising from about 5 to 10 amino acids, preferably from about 10 to about 20 amino acids, more preferably from about 20 to about 100 amino acids, and most preferably from about 20 to about 50 amino acids. In accordance with this aspect of the invention there are provided a novel polypeptide of mammalian origin, and in particular of mouse or human origin as well as biologically, diagnostically or therapeutically useful fragments, variants and derivatives thereof, variants and derivatives of the fragments, and analogs of the foregoing. Also within the scope of the present invention are polypeptides that are substantially similar to the

polypeptide with the amino acid sequence as set forth in SEQ ID NO:2 or SEQ ID NO:4 e.g. an amino acid sequence as set forth in SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12 and SEQ ID NO: 14.

In a second aspect, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence as set forth in SEQ ID NO:1 or SEQ ID NO:3. Also within the scope of the present invention are nucleic acids that are substantially similar to the nucleic acid with the nucleotide sequence as set forth in SEQ ID NO:1 or SEQ ID NO:3 e.g. nucleotide sequences as set forth in SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11 or SEQ ID NO: 13. In a preferred embodiment, the invention provides an isolated nucleic acid molecule that encodes for a polypeptide selected from the group consisting of the polypeptides set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 and SEQ ID NO:14, e.g. nucleotides 45-374 of SEQ ID NO:1, nucleotides 26-676 of SEQ ID NO:3, nucleotides 24-353 of SEQ ID NO:5, nucleotides 48-686 of SEQ ID NO:7, nucleotides 265-570 of SEQ ID NO:9, nucleotides 300-770 of SEQ ID NO: 11 or nucleotides 331-738 of SEQ ID NO:13. In a preferred embodiment, the isolated DNA takes the form of a vector molecule comprising the DNA as set forth in SEQ ID NO:1 or SEQ ID NO:3.

A third aspect of the present invention encompasses a method for the diagnosis of retinal dystrophy in a human which includes detecting the decreased transcription of messenger RNA transcribed from Rdcvf1 or Rdcvf2-encoding DNA in the eye from a mammalian organism, preferably a human, where such decreased transcription is diagnostic of the organisms' affliction with retinal dystrophy or pathological aging (ARMD). Another embodiment of the assay aspect of the invention provides a method for the diagnosis of retinal dystrophy in a mammalian organism, preferably a human, which requires measuring the amount of a Rdcvf1 or Rdcvf2 polypeptide or fragments thereof in the eye of a human suspected of suffering from a retinal dystrophy, where the presence of a decreased amount of the polypeptide or fragments thereof, relative to the amount of the polypeptide or fragments thereof in the eye of an individual not suffering from a retinal dystrophy, is diagnostic of the human's suffering from retinal dystrophy.

In accordance with another aspect of the invention there are provided anti-sense polynucleotides that regulate transcription of the Rdcvf1 or Rdcvf2 gene; in another embodiment, double stranded RNA is provided that can regulate the transcription of the Rdcvf1 or Rdcvf2 gene.

Another aspect of the invention provides a process for producing the aforementioned polypeptides, polypeptide fragments, variants and derivatives, fragments of the variants and derivatives, and analogs of the foregoing. In a preferred embodiment of this aspect of the invention there are provided methods for producing the aforementioned Rdcvf1 polypeptides comprising culturing host cells having incorporated therein an expression vector containing an exogenously-derived Rdcvf1 or Rdcvf2 - encoding polynucleotide under conditions sufficient for expression of Rdcvf1 or Rdcvf2 polypeptides in the host and then recovering the expressed polypeptide.

In accordance with another aspect of the invention there are provided products, compositions, processes and methods that utilize the aforementioned polypeptides and polynucleotides for, *inter alia*, research, biological, clinical and therapeutic purposes.

In certain additional preferred aspects of the invention there are provided an antibody or a fragment thereof which specifically binds to a polypeptide that comprises the amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8, i.e., Rdcvf1, or SEQ ID NO: 10, SEQ ID NO: 12 or SEQ ID NO: 14, i.e. Rdcvf2. In certain particularly preferred aspects in this regard, the antibodies are highly selective for mammalian, preferably mouse and in particular human Rdcvf1 or Rdcvf2 polypeptides or portions of such Rdcvf1 or Rdcvf2 polypeptides. In a related aspect, an antibody or fragment thereof is provided that binds to a fragment or portion of the amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 or SEQ ID NO:14.

In another aspect, methods of treating a disease in a subject, where the disease is mediated by or associated with a change in Rdcvf1 or Rdcvf2 gene expression e.g. a decrease in the presence of RDCVF1 or RDCVF2 polypeptide in the eye, by the administration of a therapeutically effective amount of a RDCVF1 or RDCVF2 protein as set out in SEQ ID NO:2 SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10,

SEQ ID NO:12 or SEQ ID NO:14 or a related protein or a fragment or portion thereof to the subject is provided. Also provided are methods for the diagnosis of a disease or condition associated with an decrease in Rdcvf1 or Rdcvf2 gene expression or decrease in the presence of RDCVF1 or RDCVF2 polypeptide in a subject, which comprises utilizing an antibody that binds to a polypeptide with the amino acid sequence set out in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 or SEQ ID NO:14., or a fragment or portion thereof in an immunoassay.

In yet another aspect, the invention provides cells which can be propagated in vitro, preferably vertebrate cells, which are capable upon growth in culture of producing a polypeptide that comprises the amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 or SEQ ID NO:14 or fragments thereof, where the cells contain transcriptional control DNA sequences, other than mouse or human Rdcvf1 or Rdcvf2 transcriptional control sequences, where the transcriptional control sequences control transcription of DNA encoding a polypeptide with the amino acid sequence according to SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 or SEQ ID NO:14 or fragments thereof.

In a related aspect, the present invention provides a method for producing Rdcvf1 or Rdcvf2 polypeptides which comprises culturing a host cell having incorporated therein an expression vector containing an exogenously-derived Rdcvf1 or Rdcvf2 -encoding polynucleotide under conditions sufficient for expression of Rdcvf1 or Rdcvf2 polypeptides in the host cell, thereby causing the production of an expressed polypeptide, and recovering the expressed polypeptide.

In yet another aspect of the present invention there are provided assay methods and kits comprising the components necessary to detect abnormal, e.g. below-normal expression of Rdcvf1 or Rdcvf2 polynucleotides or polypeptides or fragments thereof in body tissue samples derived from a patient, such kits comprising e.g., antibodies that bind to Rdcvf1 or Rdcvf2 or oligonucleotide probes that hybridize with polynucleotides of the invention. In a preferred embodiment, such kits also comprise instructions detailing the procedures by which the kit components are to be used.

31494-12

In another aspect, the invention is directed to an Rdcvf1 or Rdcvf2 polypeptide for use in the treatment of a human or animal body. A related aspect is directed to the use of an Rdcvf1 or Rdcvf2 polypeptide or fragment thereof, nucleotide encoding Rdcvf1 or Rdcvf2 or a fragment thereof, or antibody that binds to Rdcvf1 or Rdcvf2 or a fragment thereof in the manufacture of a medicament to treat a retinal dystrophy.

In another aspect, the invention provides a retinoprotective agent comprising a polypeptide selected from the group of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 or SEQ ID NO:14, and, optionally, a pharmaceutically acceptable carrier. In a related aspect the invention provides a pharmaceutical compositions comprising a Rdcvf1 or Rdcvf2 polypeptide or fragment thereof, nucleotide encoding Rdcvf1 or Rdcvf2 or a fragment thereof, for the treatment of a retinal dystrophy. In another related aspect, the invention provides a pharmaceutical composition comprising a therapeutically effective amount of a polypeptide selected from the group of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 or SEQ ID NO:14 and a pharmaceutically acceptable carrier.

In a related aspect, the invention provides a method for the treatment of retinal dystrophy comprising administering a therapeutically effective amount of a polypeptide selected from the group of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 or SEQ ID NO:14, and pharmaceutically acceptable carrier, to a subject in need.

In another aspect, the invention is directed to methods for the identification of molecules that can bind to Rdcvf1 or Rdcvf2 and/or modulate the activity of Rdcvf1 or Rdcvf2 or molecules that can bind to nucleic acid sequences that modulate the transcription or translation of Rdcvf1 or Rdcvf2. Such methods are disclosed in, e.g., U.S. Patent Nos. 5,541,070; 5,567,317; 5,593,853; 5,670,326; 5,679,582; 5,856,083; 5,858,657; 5,866,341; 5,876,946; 5,989,814; 6,010,861; 6,020,141; 6,030,779; and 6,043,024. Molecules

identified by such methods also fall within the scope of the present invention.

In yet another aspect, the invention is directed to methods for the introduction of nucleic acids of the invention into one or more tissues of a subject in need of treatment

31494-12

with the result that one or more proteins encoded by the nucleic acids are expressed and or secreted by cells within the tissue.

In one aspect, the invention relates to an isolated retinoprotective polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2 or
5 SEQ ID NO:4, or an amino acid sequence at least 99% identical thereto.

In another aspect, the invention relates to an isolated retinoprotective polypeptide consisting of an amino acid sequence as set forth in SEQ ID NO:2 or SEQ ID NO:4, or an amino acid sequence at least 99% identical thereto.

In another aspect, the invention relates to a vertebrate host cell which
10 can be propagated in vitro and which is capable upon growth in culture of producing the polypeptide as described herein, wherein said cell is transformed with at least one transcriptional control sequence, wherein said one or more transcriptional control sequences control transcription of DNA encoding a polypeptide as described herein.

In another aspect, the invention relates to a method for the diagnosis of
15 retinal dystrophy in a mammal which comprises: detecting the decreased transcription of messenger RNA transcribed from a retinoprotective-polypeptide-encoding-polynucleotide comprising a nucleic acid sequence as set forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, a coding region thereof, or a fragment thereof which
20 encodes a polypeptide having retinoprotective activity, in the eye of a mammal, wherein said decreased transcription is diagnostic of said mammal's suffering from retinal dystrophy.

In another aspect, the invention relates to a method for the diagnosis of retinal dystrophy in a mammal which comprises: detecting the amount of a
25 retinoprotective polypeptide comprising an amino acid sequence as set forth in

31494-12

SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, or SEQ ID NO:14, or a fragment thereof having retinoprotective activity, in rod cells of a mammal, wherein the presence of an decreased amount of said polypeptide or fragments thereof, relative to the amount of said polypeptide or fragments thereof in normal eye tissue, is diagnostic of said mammal's suffering from retinal dystrophy.

In another aspect, the invention relates to a purified antibody or a fragment thereof which specifically binds to a retinoprotective polypeptide that consists of an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 or SEQ ID NO:14, or a fragment thereof having retinoprotective activity.

In another aspect, the invention relates to a kit for the diagnosis of retinal dystrophy in a human comprising a means for collecting a sample and an antibody as described herein.

In another aspect, the invention relates to a method for producing a retinoprotective polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, or SEQ ID NO:14, or a fragment thereof having retinoprotective activity, comprising: culturing a host cell having incorporated therein an expression vector containing an exogenously-derived polynucleotide encoding said polypeptide under conditions sufficient for expression of said polypeptide in the host cell, thereby causing the production of an expressed polypeptide; and recovering the polypeptide produced by said cell.

In another aspect, the invention relates to use, for preventing or protecting photoreceptors in retinal cells from degeneration, of a vector comprising a nucleic acid sequence encoding a retinoprotective polypeptide comprising an amino

31494-12

acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, or SEQ ID NO:14, or a fragment thereof having retinoprotective activity, operably linked to a transcription control sequence.

5 In another aspect, the invention relates to a pharmaceutical composition comprising a therapeutically effective amount of a retinoprotective polypeptide comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, or a fragment thereof having retinoprotective activity, and a pharmaceutically acceptable carrier.

10 In another aspect, the invention relates to a pharmaceutical composition comprising a therapeutically effective amount of a retinoprotective nucleic acid selected from the group of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, the coding region thereof, and a fragment thereof which encodes a polypeptide having retinoprotective activity, and a
15 pharmaceutically acceptable carrier.

 In another aspect, the invention relates to use, for the treatment of retinal dystrophy in a subject in need thereof, of a polypeptide comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, or a fragment thereof having retinoprotective activity,
20 and a pharmaceutically acceptable carrier.

 In another aspect, the invention relates to a method of enhancing cone cell survival in vitro comprising contacting the cell with a polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, or a fragment thereof
25 having retinoprotective activity.

31494-12

In another aspect, the invention relates to use, in the manufacture of a medicament for preventing or protecting photoreceptors in retinal cells from degeneration, of a vector comprising a nucleic acid sequence encoding a retinoprotective polypeptide comprising an amino acid sequence as set forth in
5 SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, or SEQ ID NO:14, or a fragment thereof having retinoprotective activity, operably linked to a transcription control sequence.

In another aspect, the invention relates to a vector comprising a nucleic acid sequence encoding a retinoprotective polypeptide comprising an amino acid
10 sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, or SEQ ID NO:14, or a fragment thereof having retinoprotective activity, operably linked to a transcription control sequence, for use in preventing or protecting photoreceptors in retinal cells from degeneration.

In another aspect, the invention relates to use, in the manufacture of a
15 medicament for the treatment of retinal dystrophy in a subject in need thereof, of a polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, or a fragment thereof having retinoprotective activity, and a pharmaceutically acceptable carrier.

20 In another aspect, the invention relates to a polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, or a fragment thereof having retinoprotective activity, for use in the treatment of retinal dystrophy in a subject in need thereof.

25 In another aspect, the invention relates to a nucleic acid comprising a nucleic acid sequence as set forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, the coding region

31494-12

thereof, or a fragment thereof which encodes a polypeptide having retinoprotective activity, for use in the treatment of retinal dystrophy in a subject in need thereof.

In another aspect, the invention relates to a nucleic acid molecule comprising a nucleotide sequence that encodes a polypeptide as set forth in
5 SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, for use in the treatment of retinal dystrophy in a subject in need thereof.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill from the following description. It
10 should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present
15 disclosure.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1: mouse Rdcvf1 nucleotide sequence from the expression cloning and mouse RdCVF1 amino acid sequence.

Figure 2: mouse Rdcvf1L nucleotide sequence and amino acid
20 sequence.

Figure 3: human Rdcvf1 and human Rdcvf1 amino acids sequence.

Figure 4: human Rdcvf1L nucleotide sequence, human Rdcvf1L amino acids sequence.

Figure 5: mouse Rdcvf2 nucleotide sequence and: mouse
25 Rdcvf2 amino acids.

31494-12

Figure 6: mouse Rdcvf2L nucleotide sequence and mouse Rdcvf2L amino acid.

Figure 7: human Rdcvf2 nucleotide sequence, and human Rdcvf2 amino acids sequence.

5 **Figure 8:** depicts an amino acid alignments of the short forms of Rdcvf: (SEQ ID NO:2, 6, 10 and 14) and of the long forms of Rdcvf: SEQ ID NO:4, 8, 12 and 14).

Figure 9: depicts the primers for GST-Rdcvf1.

Figure 10: Multiple Alignment of RDCVF1/RDCVF2.

10 **Figure 11:** Comparison of mouse and human RDCVF2.

Figure 12: Multiple Alignment of mouse Rdcvf2 with EST clones be552141, bi517442, bg707818 and bi603812.

31494-12

Figure 13: Multiple Alignment of Rdcvf1 with EST clones bg299078, ai716631, bg294111, be108041 and bg395178.

Figure 14: EST sequence bg299078 corrected to match Rdcvf1.

Figure 15: EST sequence bg294111 corrected to match Rdcvf1L.

Figure 16: Real-time RT-PCR analysis of the expression of rod arrestin (A) and RdCSF1 (B) in 5 weeks retina C57BL/6@N 5 weeks (green) and C3H/HE@N (red).

Figure 17: RT-PCR analysis showing that Rdcvf2 is expressed in a rod-dependent manner and is expressed in another part of the CNS.

Figure 18: PCR analysis showing that RdCVF1 is expressed in a rod-dependent manner.

DETAILED DESCRIPTION OF THE INVENTION

In practicing the present invention, many conventional techniques in molecular biology, microbiology, and recombinant DNA are used. These techniques are well known and are explained in, for example, Current Protocols in Molecular Biology, Volumes I, II, and III, 1997 (F. M. Ausubel ed.); Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.; DNA Cloning: A Practical Approach, Volumes I and II, 1985 (D. N. Glover ed.); Oligonucleotide Synthesis, 1984 (M. L. Gait ed.); Nucleic Acid Hybridization, 1985, (Hames and Higgins); Transcription and Translation, 1984 (Hames and Higgins eds.); Animal Cell Culture, 1986 (R. I. Freshney ed.); Immobilized Cells and Enzymes, 1986 (IRL Press); Perbal, 1984, A Practical Guide to Molecular Cloning; the series, Methods in Enzymology (Academic Press, Inc.); Gene Transfer Vectors for Mammalian Cells, 1987 (J. H. Miller and M. P. Calos eds., Cold Spring Harbor Laboratory); and Methods in Enzymology Vol. 154 and Vol. 155 (Wu and Grossman, and Wu, eds., respectively).

As used herein, "differentially expressed gene" refers to (a) a gene containing at least one of the DNA sequences disclosed herein (e.g., as shown in Figure 1 and SEQ ID NO:1 or as shown in Figure 2 and SEQ ID NO:3); (b) any DNA sequence that encodes

31494-12

the amino acid sequence encoded by the DNA sequences disclosed herein (e.g., as shown in Figure 1 and SEQ ID NO:2 or as shown in Figure 2 and SEQ ID NO:4); or (c) any DNA sequence that is substantially similar to the coding sequences disclosed herein.

In its broadest sense, the term "substantially similar", when used herein with respect to a nucleotide sequence, means a nucleotide sequence corresponding to a reference nucleotide sequence, wherein the corresponding sequence encodes a polypeptide having substantially the same structure and function as the polypeptide encoded by the reference nucleotide sequence, e.g. where only changes in amino acids not affecting the polypeptide function occur. Desirably the substantially similar nucleotide sequence encodes the polypeptide encoded by the reference nucleotide sequence. The percentage of identity between the substantially similar nucleotide sequence and the reference nucleotide sequence desirably is at least 90%, more preferably at least 95%, still more preferably at least 99%. Sequence comparisons are carried out using a Smith-Waterman sequence alignment algorithm (see e.g. Waterman, M.S. Introduction to Computational Biology: Maps, sequences and genomes. Chapman & Hall. London: 1995. ISBN 0-412-99391-0). The

localS program, version 1.16, is used with following parameters: match: 1, mismatch penalty: 0.33, open-gap penalty: 2, extended-gap penalty: 2. A nucleotide sequence "substantially similar" to reference nucleotide sequence hybridizes to the reference nucleotide sequence in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 2X SSC, 0.1% SDS at 50°C, more desirably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 1X SSC, 0.1% SDS at 50°C, more desirably still in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 0.5X SSC, 0.1% SDS at 50°C, preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 0.1X SSC, 0.1% SDS at 50°C, more preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 0.1X SSC, 0.1% SDS at 65°C, yet still encodes a functionally equivalent gene product.

The differentially expressed genes disclosed herein are expressed in eye tissue and in particular is produced in rod cells however, in a human afflicted with a retinal

dystrophy such as retinitis pigmentosa, age-related macular degeneration, Bardet-Biedel syndrome, Bassen-kornzweig syndrome, best disease, choroidema, gyrate atrophy, congenital amaurosis, refsum syndrome, stargardt disease and Usher syndrome it is produced in decreases amounts relative to, i.e., to a lesser extent than in the corresponding tissues of humans who do not suffer from retinal dystrophy. Messenger RNA transcribed from the differentially expressed genes, and protein translated from such mRNA, is present in rod tissues and/or associated with such tissues in an amount at least about half, preferably at least about five times, more preferably at least amount ten times, most preferably at least about 100 times less than the levels of mRNA and protein found in corresponding tissues found in humans who do not suffer from a retinal dystrophy. Such decreases transcription of Rdcvf1 or Rdcvf2 mRNA is referred to herein as "decreased transcription."

A "host cell," as used herein, refers to a prokaryotic or eukaryotic cell that contains heterologous DNA that has been introduced into the cell by any means, e.g., electroporation, calcium phosphate precipitation, microinjection, transformation, viral infection, and the like.

"Heterologous" as used herein means "of different natural origin" or represent a non-natural state. For example, if a host cell is transformed with a DNA or gene derived from another organism, particularly from another species, that gene is heterologous with respect to that host cell and also with respect to descendants of the host cell which carry that gene. Similarly, heterologous refers to a nucleotide sequence derived from and inserted into the same natural, original cell type, but which is present in a non-natural state, e.g. a different copy number, or under the control of different regulatory elements.

A vector molecule is a nucleic acid molecule into which heterologous nucleic acid may be inserted which can then be introduced into an appropriate host cell. Vectors preferably have one or more origin of replication, and one or more site into which the recombinant DNA can be inserted. Vectors often have convenient means by which cells with vectors can be selected from those without, e.g., they encode drug resistance genes. Common vectors include plasmids, viral genomes, and (primarily in yeast and bacteria) "artificial chromosomes."

“Plasmids” generally are designated herein by a lower case p preceded and/or followed by capital letters and/or numbers, in accordance with standard naming conventions that are familiar to those of skill in the art. Starting plasmids disclosed herein are either commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids by routine application of well known, published procedures. Many plasmids and other cloning and expression vectors that can be used in accordance with the present invention are well known and readily available to those of skill in the art. Moreover, those of skill readily may construct any number of other plasmids suitable for use in the invention. The properties, construction and use of such plasmids, as well as other vectors, in the present invention will be readily apparent to those of skill from the present disclosure.

The term “isolated” means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated, even if subsequently reintroduced into the natural system. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment.

As used herein, the term “transcriptional control sequence” refers to DNA sequences, such as initiator sequences, enhancer sequences, and promoter sequences, which induce, repress, or otherwise control the transcription of protein encoding nucleic acid sequences to which they are operably linked.

As used herein, “Rdcvf1 transcriptional control sequences” or “Rdcvf2 transcriptional control sequences” are any of those transcriptional control sequences normally found associated with a mammalian Rdcvf1 or Rdcvf2 gene, preferably with the Rdcvf2 gene as found in the mouse or human genome.

As used herein, “non-human transcriptional control sequence” is any transcriptional control sequence not found in the human genome.

The term "polypeptide" is used interchangeably herein with the terms "polypeptides" and "protein(s)".

As used herein, a "chemical derivative" of a polypeptide of the invention is a polypeptide of the invention that contains additional chemical moieties not normally a part of the molecule. Such moieties may improve the molecule's solubility, absorption, biological half life, etc. The moieties may alternatively decrease the toxicity of the molecule, eliminate or attenuate any undesirable side effect of the molecule, etc. Moieties capable of mediating such effects are disclosed, for example, in Remington's Pharmaceutical Sciences, 16th ed., Mack Publishing Co., Easton, Pa. (1980).

As used herein, a "neuroprotective agent" is a compound that prevents or protects neuronal cells from degeneration. A "retinoprotective agent" is a compound that prevents or protects retinal cells from degeneration.

The invention includes nucleic acid molecules, preferably DNA molecules, such as (1) an isolated comprising a nucleotide sequence as set forth in SEQ ID NO:1 or SEQ ID NO:3, (2) isolated nucleic acid molecules that comprise nucleic acid sequences that hybridize under high stringency conditions to the isolated DNA as set forth in SEQ ID NO:1 or SEQ ID NO:3, and (3) nucleic acid sequences that hybridize to (1) or (2), above. Such hybridization conditions may be highly stringent or less highly stringent, as described above. In instances wherein the nucleic acid molecules are deoxyoligonucleotides ("oligos"), highly stringent conditions may refer, e.g., to washing in 6X SSC/0.05% sodium pyrophosphate at 37 °C. (for 14-base oligos), 48 °C. (for 17-base oligos), 55 °C. (for 20-base oligos), and 60 °C. (for 23-base oligos). Suitable ranges of such stringency conditions for nucleic acids of varying compositions are described in Krause and Aaronson (1991) Methods in Enzymology, 200:546-556 in addition to Maniatis et al., cited above.

These nucleic acid molecules may act as target gene antisense molecules, useful, for example, in target gene regulation and/or as antisense primers in amplification reactions of target gene nucleic acid sequences. Further, such sequences may be used as part of ribozyme and/or triple helix sequences, also useful for target gene regulation. Still

further, such molecules may be used as components of diagnostic methods whereby the presence of a RdCVF1 or RdCVF2 disease-causing allele, may be detected.

The invention also encompasses (a) vectors that contain any of the foregoing coding sequences (i.e., sense) and/or their complements (i.e., antisense); (b) expression vectors that contain any of the foregoing coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences; and (c) genetically engineered host cells that contain any of the foregoing coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell. As used herein, regulatory elements include but are not limited to inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression.

The invention includes fragments of any of the nucleic acid sequences disclosed herein. Fragments of the full length Rdcvf1 or Rdcvf2 gene may be used as a hybridization probe for a cDNA library to isolate the full length gene and to isolate other genes which have a high sequence similarity to the Rdcvf1 or Rdcvf2 gene and similar biological activity. Probes of this type preferably have at least about 30 bases and may contain, for example, from about 30 to about 50 bases, about 50 to about 100 bases, about 100 to about 200 bases, or more than 200 bases (e.g. 300). The probe may also be used to identify a cDNA clone corresponding to a full length transcript and a genomic clone or clones that contain the complete Rdcvf1 or Rdcvf2 gene including regulatory and promoter regions, exons, and introns. An example of a screen comprises isolating the coding region of the Rdcvf1 or Rdcvf2 gene by using the known DNA sequence to synthesize an oligonucleotide probe or random priming of the isolated sequence disclosed in Figure 1 to 8. Labeled oligonucleotides having a sequence complementary to that of the gene of the present invention are used to screen a library of human cDNA, genomic DNA to determine which individual clones of the library the probe hybridizes to.

In addition to the gene sequences described above, orthologs of such sequences, as may, for example be present in other species, may be identified and may be readily isolated, without undue experimentation, by molecular biological techniques well known in the art. Further, there may exist genes at other genetic loci within the genome that

encode proteins which have extensive homology (homologs) to one or more domains of such gene products. These genes may also be identified via similar techniques. Examples of orthologs or homologues are provided in Figures 8, 10, 11, 12 or 13.

For example, the isolated expressed gene sequence may be labeled and used to screen a cDNA library constructed from mRNA obtained from the organism of interest. Hybridization conditions will be of a lower stringency when the cDNA library was derived from an organism different from the type of organism from which the labeled sequence was derived. Alternatively, the labeled fragment may be used to screen a genomic library derived from the organism of interest, again, using appropriately low stringent conditions. Such low stringency conditions will be well known to those of skill in the art, and will vary predictably depending on the phylogeny of specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook et al. cited above.

Further, a previously unknown expressed gene-type sequence may be isolated by performing PCR using two degenerated oligonucleotide primer pools designed on the basis of amino acid sequences within the gene of interest. The template for the reaction may be cDNA obtained by reverse transcription of mRNA prepared from human or non-human cell lines or tissue known or suspected to express a homologue or splicing variants.

The PCR product may be subcloned and sequenced to ensure that the amplified sequences represent the sequences of a expressed gene-like nucleic acid sequence. The PCR fragment may then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment may be labeled and used to screen a bacteriophage cDNA library. Alternatively, the labeled fragment may be used to screen a genomic library.

PCR technology may also be utilized to isolate full length cDNA sequences. For example, RNA may be isolated, following standard procedures, from an appropriate cellular or tissue source. A reverse transcription reaction may be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed"

with guanines using a standard terminal transferase reaction, the hybrid may be digested with RNAase H, and second strand synthesis may then be primed with a poly-C primer. Thus, cDNA sequences upstream of the amplified fragment may easily be isolated. For a review of cloning strategies which may be used, see e.g., Sambrook et al., 1989, supra.

In cases where the differentially expressed gene identified is the normal, or wild type, gene, this gene may be used to isolate mutant alleles of the gene. Such an isolation is preferable in processes and disorders which are known or suspected to have a genetic basis. Mutant alleles may be isolated from individuals either known or suspected to have a genotype which contributes to disease symptoms. Mutant alleles and mutant allele products may then be utilized in the diagnostic assay systems described below.

A cDNA of the mutant gene may be isolated, for example, by using RT-PCR, a technique which is well known to those of skill in the art. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide (or random hexamers) to mRNA isolated from tissue known or suspected to be expressed in an individual putatively carrying the mutant allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal gene (or by any other means). Using these two primers, the product is then amplified via PCR, cloned into a suitable vector, and subjected to DNA sequence analysis through methods well known to those of skill in the art. By comparing the DNA sequence of the mutant gene to that of the normal gene, the mutation(s) responsible for the loss or alteration of function of the mutant gene product can be ascertained.

Alternatively, a genomic or cDNA library can be constructed and screened using DNA or RNA, from a tissue known to or suspected of expressing the gene of interest in an individual suspected of to carry the mutant allele. The normal gene or any suitable fragment thereof may then be labeled and used as a probe to identify the corresponding mutant allele in the library. The clone containing this gene may then be purified through methods routinely practiced in the art, and subjected to sequence analysis as described above.

Additionally, an expression library can be constructed utilizing DNA isolated from or cDNA synthesized from a tissue known to or suspected of expressing the gene of interest in an individual suspected of to carry the mutant allele. In this manner, gene products made by the putatively mutant tissue may be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against the normal gene product, as described, below. (For screening techniques, see, for example, Harlow, E. and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor.) In cases where the mutation results in an expressed gene product with altered function (e.g., as a result of a missense mutation), a polyclonal set of antibodies are likely to cross-react with the mutant gene product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis as described above.

Differentially expressed gene products include those proteins encoded by nucleotide sequence set forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11 or SEQ ID NO:13, in particular, a polypeptide that is or includes the amino acid sequence set out in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 or SEQ ID NO:14, or fragments thereof.

In addition, expressed gene products may include proteins that represent functionally equivalent gene products. Such an equivalent gene product may contain deletions, additions or substitutions of amino acid residues within the amino acid sequence encoded by the differentially expressed gene sequences described, above, but which result in a silent change, thus producing a functionally equivalent differentially expressed gene product (polymorphisms). Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved and on comparison with amino-acids sequence from other species.

For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and

glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid. "Functionally equivalent," as utilized herein, may refer to a protein or polypeptide capable of exhibiting a substantially similar *in vivo* or *in vitro* activity as the endogenous differentially expressed gene products encoded by the differentially expressed gene sequences described above. "Functionally equivalent" may also refer to proteins or polypeptides capable of interacting with other cellular or extracellular molecules in a manner similar to the way in which the corresponding portion of the endogenous differentially expressed gene product would. For example, a "functionally equivalent" peptide would be able, in an immunoassay, to diminish the binding of an antibody to the corresponding peptide (i.e., the peptidic amino acid sequence of which was modified to achieve the "functionally equivalent" peptide) of the endogenous protein, or to the endogenous protein itself, where the antibody was raised against the corresponding peptide of the endogenous protein. An equimolar concentration of the functionally equivalent peptide will diminish the aforesaid binding of the corresponding peptide by at least about 5%, preferably between about 5% and 10%, more preferably between about 10% and 25%, even more preferably between about 25% and 50%, and most preferably between about 40% and 50%.

The differentially expressed gene products may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing the differentially expressed gene polypeptides and peptides of the invention by expressing nucleic acid encoding differentially expressed gene sequences are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing expressed gene protein coding sequences and appropriate transcriptional/translational control signals. These methods include, for example, *in vitro* recombinant DNA techniques, synthetic techniques and *in vivo* recombination/genetic recombination. See, for example, the techniques described in Sambrook et al., 1989, *supra*, and Ausubel et al., 1989, *supra*. Alternatively, RNA or cDNA capable of encoding expressed gene protein sequences may be chemically synthesized using, for example, synthesizers. See, for example, the techniques described in "Oligonucleotide Synthesis",

31494-12

1984, Gait, M. J. ed., IRL Press, Oxford .

A variety of host-expression vector systems may be utilized to express the differentially expressed gene coding sequences of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express the differentially expressed gene protein of the invention *in situ*. These include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing differentially expressed gene protein coding sequences; yeast (e.g. *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing the differentially expressed gene protein coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing the differentially expressed gene protein coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid transformation vectors (e.g., Ti plasmid) containing differentially expressed gene protein coding sequences; or mammalian cell systems (e.g. COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothioneine promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter; Cytomegalovirus Early gene promoter).

Expression of RDCVF1 or RDCVF2 by a cell from an *Rdcvf1* or *Rdcvf2* gene that is native to a the cell can also be performed. Methods for such expression are detailed in, e.g., U.S. Patents 5,641,670; 5,733,761; 5,968,502; and 5,994,127.

Cells that have

been induced to express RDCVF1 or RDCVF2 by the methods of any of U.S. Patents 5,641,670; 5,733,761; 5,968,502; and 5,994,127 can be implanted into a desired tissue in a living animal in order to increase the local concentration of RDCVF1 or RDCVF2 in the tissue.

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the differentially expressed gene protein being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of antibodies or to screen peptide libraries, for example, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which the differentially expressed gene protein coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke & Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. PGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-asephagarose beads followed by elution in the presence of free glutathione. The PGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene protein can be released from the GST moiety by using these endopeptidases.

Promoter regions can be selected from any desired gene using vectors that contain a reporter transcription unit lacking a promoter region, such as a chloramphenicol acetyl transferase ("cat") or luciferase transcription unit, downstream of restriction site or sites for introducing a candidate promoter fragment; i.e., a fragment that may contain a promoter. As is well known, introduction into the vector of a promoter-containing fragment at the restriction site upstream of the cat or luciferase gene engenders production of CAT or luciferase activity, which can be detected by standard CAT assays or luminometry. Vectors suitable to this end are well known and readily available. Three such vectors are pKK232-8, -pCM7 and pGL3 (Promega, E1751, Genebank Ass n° u47295). Thus, promoters for expression of polynucleotides of the present invention include not only well known and readily available promoters, but also promoters that readily may be obtained by the foregoing technique, using a reporter gene assay.

Among known bacterial promoters suitable for expression of polynucleotides and polypeptides in accordance with the present invention are the *E. coli* lacI and lacZ promoters, the T3 and T7 promoters, the T5 tac promoter, the lambda PR, PL promoters and the trp promoter. Among known eukaryotic promoters suitable in this regard are the CMV immediate early promoter, the HSV thymidine kinase promoter, the early and late SV40 promoters, the promoters of retroviral LTRs, such as those of the Rous sarcoma virus ("RSV"), and metallothionein promoters, such as the mouse metallothionein-I promoter.

In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is one of several insect systems that can be used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The differentially expressed gene coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of differentially expressed gene coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted gene is expressed. (E.g., see Smith et al., 1983, J. Virol. 46: 584; Smith, U.S. Pat. No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the differentially expressed gene coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing differentially expressed gene protein in infected hosts. (e.g., See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals may also be required for efficient translation of inserted differentially expressed gene coding sequences. These signals include the ATG initiation codon and adjacent sequences. In

cases where an entire differentially expressed gene, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of the differentially expressed gene coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals (Kozack sequence) and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner et al., 1987, *Methods in Enzymol.* 153:516-544).

Selection of appropriate vectors and promoters for expression in a host cell is a well known procedure and the requisite techniques for expression vector construction, introduction of the vector into the host and expression in the host per se are routine skills in the art.

Generally, recombinant expression vectors will include origins of replication, a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence, and a selectable marker to permit isolation of vector containing cells after exposure to the vector.

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, etc.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the differentially expressed protein may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the differentially expressed protein. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the differentially expressed protein. These stable cell lines might be used as a way of cellular therapy directly or after encapsulation.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, *Cell* 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, *Proc. Natl. Acad. Sci. USA* 48:2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, *Cell* 22:817) genes can be employed in *tk⁻*, *hgprt⁻* or *aprt⁻* cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for *dhfr*, which confers resistance to methotrexate (Wigler, et al., 1980, *Natl. Acad. Sci. USA* 77:3567; O'Hare, et al., 1981, *Proc. Natl. Acad. Sci. USA* 78:1527); *gpt*, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, *Proc. Natl. Acad. Sci. USA* 78:2072); *neo*, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, *J. Mol. Biol.* 150:1); and *hygro*, which confers resistance to hygromycin (Santerre, et al., 1984, *Gene* 30:147) genes.

An alternative fusion protein system allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, *Proc. Natl. Acad. Sci. USA* 88: 8972-8976). In this system, the gene of interest is subcloned

into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni^{2+} nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

When used as a component in assay systems such as those described below, the differentially expressed protein may be labeled, either directly or indirectly, to facilitate detection of a complex formed between the differentially expressed protein and a test substance. Any of a variety of suitable labeling systems may be used including but not limited to radioisotopes such as ^{125}I ; enzyme labeling systems that generate a detectable calorimetric signal or light when exposed to substrate; and fluorescent labels.

Where recombinant DNA technology is used to produce the differentially expressed protein for such assay systems, it may be advantageous to engineer fusion proteins that can facilitate labeling, immobilization and/or detection.

Indirect labeling involves the use of a protein, such as a labeled antibody, which specifically binds to either a differentially expressed gene product. Such antibodies include but are not limited to polyclonal, monoclonal, chimeric, single chain, Fab fragments and fragments produced by an Fab expression library.

In another embodiment, nucleic acids comprising a sequence encoding RDCVF1 or RDCVF2 protein or functional derivative thereof, are administered to promote cone function, by way of gene therapy. Gene therapy refers to therapy performed by the administration of a nucleic acid to a subject. In this embodiment of the invention, the nucleic acid produces its encoded protein that mediates a therapeutic effect by promoting cone function.

Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

In a preferred aspect, the therapeutic comprises a Rdcvf1 or Rdcvf2 nucleic acid that is part of an expression vector that expresses a RDCVF1 or RDCVF2 protein or fragment or chimeric protein thereof in a suitable host. In particular, such a nucleic acid has a promoter operably linked to the Rdcvf1 or Rdcvf2 coding region, said promoter

31494-12

being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, a nucleic acid molecule is used in which the Rdcvf1 or Rdcvf2 coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the Rdcvf1 or Rdcvf2 nucleic acid (Koller and Smithies, 1989, Proc. Natl. Acad. Sci. USA 86:8932-8935; Zijlstra et al., 1989, Nature 342:435-438).

Delivery of the nucleic acid into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vector, or indirect, in which case, cells are first transformed with the nucleic acid *in vitro*, then transplanted into the patient. These two approaches are known, respectively, as *in vivo* or *ex vivo* gene therapy.

In a specific embodiment, the nucleic acid is directly administered *in vivo*, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, e.g., by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by infection using a defective or attenuated retroviral or other viral vector (adenovirus, adeno-associated virus and lentivirus) (see, e.g., U.S. Pat. No. 4,980,286 and others mentioned *infra*), or by direct injection of naked DNA, or by use of microparticle bombardment (e.g., a gene gun; Biolistic^{*}, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering it in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see e.g., U.S. Patents 5,166,320; 5,728,399; 5,874,297; and 6,030,954)

(which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, a nucleic acid-ligand complex can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted *in vivo* for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Publications WO

*Trade-mark

31494-12

92/06180; WO 92/22635; WO92/20316; WO93/14188; and WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (see, e.g., U.S. Patents 5,413,923; 5,416,260; and 5,574,205; and Zijlstra et al., 1989, Nature 342:435-438).

In a specific embodiment, a viral vector that contains the Rdcvf1 or Rdcvf2 nucleic acid is used. For example, a retroviral vector can be used (see, e.g., U.S. Patents 5,219,740; 5,604,090; and 5,834,182). These retroviral vectors have been modified to delete retroviral sequences that are not necessary for packaging of the viral genome and integration into host cell DNA. The Rdcvf1 or Rdcvf2 nucleic acid to be used in gene therapy is cloned into the vector, which facilitates delivery of the gene into a patient.

Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Methods for conducting adenovirus-based gene therapy are described in, e.g., U.S. Patents 5,824,544; 5,868,040; 5,871,722; 5,880,102; 5,882,877; 5,885,808; 5,932,210; 5,981,225; 5,994,106; 5,994,132; 5,994,134; 6,001,557; and 6,033,8843.

Adeno-associated virus (AAV) has also been proposed for use in gene therapy. Adeno-associated viruses are especially attractive vehicles for delivering genes to the retina. Methods for producing and utilizing AAV are described, e.g., in U.S. Patents 5,173,414; 5,252,479; 5,552,311; 5,658,785; 5,763,416; 5,773,289; 5,843,742; 5,869,040; 5,942,496; and 5,948,675.

Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those

cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient directly or after encapsulation.

In this embodiment, the nucleic acid is introduced into a cell prior to administration in vivo of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a patient by various methods known in the art. In a preferred embodiment, epithelial cells are injected, e.g., subcutaneously. In another embodiment, recombinant skin cells may be applied as a skin graft onto the patient. Recombinant blood cells (e.g., hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood; fetal liver, etc.

In a preferred embodiment, the cell used for gene therapy is autologous to the patient.

In an embodiment in which recombinant cells are used in gene therapy, a Rdcvf1 or Rdcvf2 nucleic acid is introduced into the cells such that it is expressible by the cells or their progeny, and the recombinant cells are then administered in vivo for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem-and/or progenitor cells which can be isolated and maintained in vitro can potentially be used in accordance with this embodiment of the present invention. Such stem cells include but are not limited to hematopoietic stem cells (HSC), stem cells of epithelial tissues such as the skin and the lining of the gut, embryonic heart muscle cells, liver stem cells (see, e.g., WO 94/08598), and neural stem cells (Stemple and Anderson, 1992, *Cell* 71:973-985).

Epithelial stem cells (ESCs) or keratinocytes can be obtained from tissues such as the skin and the lining of the gut by known procedures (Rheinwald, 1980, *Meth. Cell Bio.* 21A:229). In stratified epithelial tissue such as the skin, renewal occurs by mitosis of stem cells within the germinal layer, the layer closest to the basal lamina. Stem cells within the lining of the gut provide for a rapid renewal rate of this tissue. ESCs or keratinocytes obtained from the skin or lining of the gut of a patient or donor can be grown in tissue culture (Pittelkow and Scott, 1986, *Mayo Clinic Proc.* 61:771). If the ESCs are provided by a donor, a method for suppression of host versus graft reactivity (e.g., irradiation, drug or antibody administration to promote moderate immunosuppression) can also be used. Retinal stem cells (Tropepe et al., 2000, *Science*, 287: 2032).

With respect to hematopoietic stem cells (HSC), any technique which provides for the isolation, propagation, and maintenance in vitro of HSC can be used in this embodiment of the invention. Techniques by which this may be accomplished include (a) the isolation and establishment of HSC cultures from bone marrow cells isolated from the future host, or a donor, or (b) the use of previously established long-term HSC cultures, which may be allogeneic or xenogeneic. Non-autologous HSC are used preferably in conjunction with a method of suppressing transplantation immune reactions of the future host/patient. In a particular embodiment of the present invention, human bone marrow cells can be obtained from the posterior iliac crest by needle aspiration (see, e.g., Kodo et al., 1984, *J. Clin. Invest.* 73:1377-1384). In a preferred embodiment of the present

invention, the HSCs can be made highly enriched or in substantially pure form. This enrichment can be accomplished before, during, or after long-term culturing, and can be done by any techniques known in the art. Long-term cultures of bone marrow cells can be established and maintained by using, for example, modified Dexter cell culture techniques (Dexter et al., 1977, J. Cell Physiol. 91:335) or Witlock-Witte culture techniques (Witlock and Witte, 1982, Proc. Natl. Acad. Sci. USA 79:3608-3612).

In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

Described herein are methods for the production of antibodies capable of specifically recognizing one or more differentially expressed gene epitopes. Such antibodies may include, but are not limited to polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. Such antibodies may be used, for example, in the detection of a fingerprint, target, gene in a biological sample, or, alternatively, as a method for the inhibition of abnormal target gene activity. Thus, such antibodies may be utilized as part of disease treatment methods, and/or may be used as part of diagnostic techniques whereby patients may be tested for abnormal levels of Rdcvf1 or Rdcvf2, or for the presence of abnormal forms of Rdcvf1 or Rdcvf2 by sampling the aqueous humor and/or vitreous by methods familiar to those of skill in the art (e.g. Forster, RK, Abbott, RL, Gelender, H. (1980) Management of infectious endophthalmitis Ophthalmology 87,313-319)].

For the production of antibodies to a differentially expressed gene, various host animals may be immunized by injection with a differentially expressed protein, or a portion thereof or by DNA immunization. Such host animals may include but are not limited to rabbits, mice, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide,

surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*.

Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of animals immunized with an antigen, such as target gene product, or an antigenic functional derivative thereof. For the production of polyclonal antibodies, host animals such as those described above, may be immunized by injection with differentially expressed gene product supplemented with adjuvants as also described above.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, may be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to the hybridoma technique of Kohler and Milstein, (1975, *Nature* 256:495-497; and U.S. Pat. No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, *Immunology Today* 4:72; Cole et al., 1983, *Proc. Natl. Acad. Sci. USA* 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, *Monoclonal Antibodies And Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated in vitro or in vivo. Production of high titers of mAbs in vivo makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, *Proc. Natl. Acad. Sci.*, 81:6851-6855; Neuberger et al., 1984, *Nature*, 312:604-608; Takeda et al., 1985, *Nature*, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable or hypervariable region derived from a murine mAb and a human immunoglobulin constant region.

31494-12

Alternatively, techniques described for the production of single chain antibodies (U.S. Pat. No. 4,946,778; Bird, 1988, Science 242:423-426; Huston et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward et al., 1989, Nature 334:544-546) can be adapted to produce differentially expressed gene-single chain antibodies. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

Most preferably, techniques useful for the production of "humanized antibodies" can be adapted to produce antibodies to the polypeptides, fragments, derivatives, and functional equivalents disclosed herein. Such techniques are disclosed in U.S. Patent Nos. 5,932, 448; 5,693,762; 5,693,761; 5,585,089; 5,530,101; 5,910,771; 5,569,825; 5,625,126; 5,633,425; 5,789,650; 5,545,580; 5,661,016; and 5,770,429.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed (Huse et al., 1989, Science, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Particularly preferred, for ease of detection, is the sandwich assay, of which a number of variations exist, all of which are intended to be encompassed by the present invention.

For example, in a typical forward assay, unlabeled antibody is immobilized on a solid substrate and the sample to be tested brought into contact with the bound molecule. After a suitable period of incubation, for a period of time sufficient to allow formation of an antibody-antigen binary complex. At this point, a second antibody, labeled with a reporter molecule capable of inducing a detectable signal, is then added and incubated, allowing time sufficient for the formation of a ternary complex of antibody-antigen-labeled antibody. Any unreacted material is washed away, and the presence of the antigen is determined by observation of a signal, or may be quantitated by comparing with a

control sample containing known amounts of antigen. Variations on the forward assay include the simultaneous assay, in which both sample and antibody are added simultaneously to the bound antibody, or a reverse assay in which the labeled antibody and sample to be tested are first combined, incubated and added to the unlabeled surface bound antibody. These techniques are well known to those skilled in the art, and the possibility of minor variations will be readily apparent. As used herein, "sandwich assay" is intended to encompass all variations on the basic two-site technique. For the immunoassays of the present invention, the only limiting factor is that the unlabeled and the labeled antibodies be an RdCVF1- or RdCVF2-specific antibody.

The most commonly used reporter molecules in this type of assay are either enzymes, fluorophore- or radionuclide-containing molecules. In the case of an enzyme immunoassay an enzyme is conjugated to the second antibody, usually by means of glutaraldehyde or periodate. As will be readily recognized, however, a wide variety of different ligation techniques exist, which are well-known to the skilled artisan. Commonly used enzymes include horseradish peroxidase, glucose oxidase, beta-galactosidase and alkaline phosphatase, among others. The substrates to be used with the specific enzymes are generally chosen for the production, upon hydrolysis by the corresponding enzyme, of a detectable color change. For example, p-nitrophenyl phosphate is suitable for use with alkaline phosphatase conjugates; for peroxidase conjugates, 1,2-phenylenediamine or toluidine are commonly used. It is also possible to employ fluorogenic substrates, which yield a fluorescent product rather than the chromogenic substrates noted above. A solution containing the appropriate substrate is then added to the tertiary complex of antibody-RdCVF1- or RdCVF2-labeled antibody. The substrate reacts with the enzyme linked to the second antibody, giving a qualitative visual signal, which may be further quantitated, usually spectrophotometrically, to give an evaluation of the amount of Rdcvf1 or Rdcvf2 which is present in the serum sample.

Alternately, fluorescent compounds, such as fluorescein and rhodamine, may be chemically coupled to antibodies without altering their binding capacity. When activated by illumination with light of a particular wavelength, the fluorochrome-labeled antibody absorbs the light energy, inducing a state of excitability in the molecule, followed by

emission of the light at a characteristic longer wavelength. The emission appears as a characteristic color visually detectable with a light microscope. Immunofluorescence and EIA techniques are both very well established in the art and are particularly preferred for the present method. However, other reporter molecules, such as radioisotopes, chemiluminescent or bioluminescent molecules may also be employed. It will be readily apparent to the skilled artisan how to vary the procedure to suit the required use.

This invention also relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of the gene encoding a polypeptide selected from the group consisting of the polypeptides set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 and SEQ ID NO:14 which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-expression or altered spatial or temporal expression of the gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents (SSCP), or by direct DNA sequencing (e.g., Myers et al., Science (1985) 230:1242). Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (see Cotton et al., Proc Natl Acad Sci USA (1985) 85: 4397-4401). In another embodiment, an array of oligonucleotides probes comprising the Rdcvf1 or Rdcvf2 nucleotide sequence or

fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see for example: M. Chee et al., Science, Vol 274, pp 610-613 (1996)).

The diagnostic assays offer a process for diagnosing or determining a susceptibility to disease through detection of mutation in the Rdcvf1 or Rdcvf2 gene by the methods described. In addition, such diseases may be diagnosed by methods comprising determining from a sample derived from a subject an abnormal Rdcvf1 or Rdcvf2 expression: Expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, nucleic acid amplification, for instance PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit which comprises:

(a) a polynucleotide of the present invention, preferably the nucleotide sequence encoding a polypeptide selected from the group consisting of the polypeptides set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 and SEQ ID NO:14, or a fragment thereof;

(b) a nucleotide sequence complementary to that of (a);

(c) a polypeptide of the present invention, preferably the polypeptide of or a fragment thereof; or

(d) an antibody to a polypeptide of the present invention, preferably to a polypeptide selected from the group consisting of the polypeptides set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 and SEQ ID NO:14.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, particularly retinitis pigmentosa, age-related macular degeneration, Bardet-Biedel syndrome, Bassen-kornzweig syndrome, best disease, choroidema, gyrate atrophy, congenital amourosis, refsun syndrome, stargardt disease and Usher syndrome. The nucleotide sequences of the present invention are also valuable for chromosome localization. The chromosomal location can be obtained by PCR on DNA prepared from a panel of hybrid cell lines (mouse-hamster). The chromosomal location of the human gene can be predicted from the location of the mouse gene by syntheny (McCarthy et al.(1997), Genome research, 7, 1153). The sequence is specifically targeted to, and can hybridize with, a particular location on an individual chromosome, including a mouse or a human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian Inheritance in Man (available on-line through Johns Hopkins University Welch Medical Library, RetNet). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

An additional embodiment of the invention relates to the administration of a pharmaceutical composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of Rdcvf1 or Rdcvf2, mimetics or agonists. The compositions may be administered alone or in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical

carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

The pharmaceutical compositions encompassed by the invention may be administered by any number of routes: e.g. Transscleral delivery of bioreactive protein to the choroid and retina Ambati et al. (2000) *Investigative Ophthalmology and Visual Science*, 41, 1186. The formulation of topical ophthalmic preparations, including ophthalmic solutions, suspensions and ointments is well known to those skilled in the art (see Remington's *Pharmaceutical Sciences*, 18th Edition, Chapter 86, pages 1581-1592 Mack Publishing Company, 1990). Other modes of administration are available, including intracameral injections (which may be made directly into the anterior chamber or directly into the vitreous chamber), subconjunctival injections and retrobulbar injections, and methods and means for producing ophthalmic preparations suitable for such modes of administration are also well known.

As used in this application, "extraocular" refers to the ocular surface and the (external) space between the eyeball and the eyelid. Examples of extraocular regions include the eyelid fornix or cul-de-sac, the conjunctival surface and the corneal surface. This location is external to all ocular tissue and an invasive procedure is not required to access this region. Examples of extraocular systems include inserts and "topically" applied drops, gels or ointments which may be used to deliver therapeutic material to these regions. Extraocular devices are generally easily removable, even by the patient.

The following patents disclose extraocular systems which are used to administer drugs to the extraocular regions. Higuchi et al. discloses in U.S. Pat. Nos. 3,981,303, 3,986,510 and 3,995,635, a biodegradable ocular insert which contains a drug. The insert can be made in different shapes for retention in the cul-de-sac of the eyeball, the extraocular space between the eyeball and the eyelid. Several common biocompatible polymers are disclosed as suitable for use in fabricating this device. These polymers include zinc alginate, poly (lactic acid), poly (vinyl alcohol), poly (anhydrides) and poly (glycolic acid). The patents also describe membrane coated devices with reduced permeation to the drug and hollow chambers holding the drug formulation.

Theeuwes, U.S. Pat. No. 4,217,898, discloses microporous reservoirs which are used for controlled drug delivery. These devices are placed extraocularly in the ocular cul-de-sac. Among the polymer systems of interest include poly (vinylchloride)-co-poly (vinyl acetate) copolymers. Kaufman discloses in U.S. Pat. Nos. 4,865,846 and 4,882,150 an ophthalmic drug delivery system which contains at least one bio-erodible material or ointment carrier for the conjunctival sac. The patent discloses polymer systems, such as, poly (lactide), poly (glycolide), poly (vinyl alcohol) and cross linked collagen, as suitable delivery systems.

In the presently described use of RDCVF1 or RDCVF2 protein product for the treatment of retinal disease or injury it is also advantageous that a topically applied ophthalmic formulation include an agent to promote the penetration or transport of the therapeutic agent into the eye. Such agents are known in the art. For example, Ke et al., U.S. Pat. No. 5,221,696 disclose the use of materials to enhance the penetration of ophthalmic preparations through the cornea.

Intraocular systems are those systems which are suitable for use in any tissue compartment within, between or around the tissue layers of the eye itself. These locations include subconjunctival (under the ocular mucous membrane adjacent to the eyeball), orbital (behind the eyeball), and intracameral (within the chambers of the eyeball itself). In contrast to extraocular systems, an invasive procedure consisting of injection or implantation is required to access these regions.

The following patents disclose intraocular devices. Wong, U.S. Pat. No. 4,853,224, discloses microencapsulated drugs for introduction into the chamber of the eye. Polymers which are used in this system include polyesters and polyethers. Lee, U.S. Pat. No. 4,863,457, discloses a biodegradable device which is surgically implanted intraocularly for the sustained release of therapeutic agents. The device is designed for surgical implantation under the conjunctiva (mucous membrane of the eyeball). Krezancaki, U.S. Pat. No. 4,188,373, discloses a pharmaceutical vehicle which gels at human body temperature. This vehicle is an aqueous suspension of the drug and gums or cellulose derived synthetic derivatives. Haslam et al. discloses in U.S. Pat. Nos. 4,474,751 and 4,474,752 a polymer-drug system which is

liquid at room temperature and gels at body temperature. Suitable polymers used in this system include polyoxyethylene and polyoxy propylene. Davis et al. disclose in U.S. Pat. No. 5,384,333 a biodegradable injectable drug delivery polymer which provides long term drug release. The drug composition is made up of a pharmaceutically active agent in a biodegradable polymer matrix, where the polymer matrix is a solid at temperatures in the range 20.degree. to 37.degree. C. and is flowable at temperatures in the range 38.degree. to 52.degree. C. The drug delivery polymer is not limited to the delivery of soluble or liquid drug formulations. For example, the polymer can be used as a matrix for stabilizing and retaining at the site of injection drug-containing microspheres, liposomes or other particulate-bound drugs.

A particularly suitable vehicle for intraocular injection is sterile distilled water in which the RDCVF1 or RDCVF2 protein product is formulated as a sterile, isotonic solution, properly preserved. Yet another ophthalmic preparation may involve the formulation of the RDCVF1 or RDCVF2 protein product with an agent, such as injectable microspheres or liposomes, that provides for the slow or sustained release of the protein which may then be delivered as a depot injection. Other suitable means for the intraocular introduction of RDCVF1 or RDCVF2 protein product includes, implantable drug delivery devices or which contain the RDCVF1 or RDCVF2 protein product.

The ophthalmic preparations of the present invention, particularly topical preparations, may include other components, for example ophthalmically acceptable preservatives, tonicity agents, cosolvents, wetting agents, complexing agents, buffering agents, antimicrobials, antioxidants and surfactants, as are well known in the art. For example, suitable tonicity enhancing agents include alkali metal halides (preferably sodium or potassium chloride), mannitol, sorbitol and the like. Sufficient tonicity enhancing agent is advantageously added so that the formulation to be instilled into the eye is hypotonic or substantially isotonic. Suitable preservatives include, but are not limited to, benzalkonium chloride, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid and the like. Hydrogen peroxide may also be used as preservative. Suitable cosolvents include, but are not limited to, glycerin, propylene glycol and polyethylene glycol. Suitable complexing agents include caffeine,

polyvinylpyrrolidone, beta -cyclodextrin or hydroxypropyl- beta -cyclodextrin. Suitable surfactants or wetting agents include, but are not limited to, sorbitan esters, polysorbates such as polysorbate 80, tromethamine, lecithin, cholesterol, tyloxapol and the like. The buffers can be conventional buffers such as borate, citrate, phosphate, bicarbonate, or Tris-HCl.

The formulation components are present in concentrations that are acceptable to the extraocular or intraocular site of administration. For example, buffers are used to maintain the composition at physiological pH or at slightly lower pH, typically within a pH range of from about 5 to about 8. Additional formulation components may include materials which provide for the prolonged ocular residence of the extraocularly administered therapeutic agent so as to maximize the topical contact and promote absorption. Suitable materials include polymers or gel forming materials which provide for increased viscosity of the ophthalmic preparation. Chitosan is a particularly suitable material as an ocular release-rate controlling agent in sustained release liquid ophthalmic drug formulations (see U.S. Pat. No. 5,422,116, Yen, et. al.) The suitability of the formulations of the instant invention for controlled release (e.g., sustained and prolonged delivery) of an ophthalmic treating agent in the eye can be determined by various procedures known in the art, e.g., as described in *Journal of Controlled Release*, 6:367-373, 1987, as well as variations thereof.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of *Remington's Pharmaceutical Sciences* (Maack Publishing Co., Easton, Pa.).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethylcellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid

esters, such as ethyl oleate or triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of Rdcvf1 or Rdcvf2, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models, usually mice, rabbits, dogs, or pigs. The animal model may also be used to determine the appropriate

concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example Rdcvf1 or Rdcvf2 or fragments thereof, antibodies to Rdcvf1, agonists, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from 0.1 to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc. Pharmaceutical formulations suitable for oral administration of proteins are described, e.g., in U.S.

Patents 5,008,114; 5,505,962; 5,641,515; 5,681,811; 5,700,486; 5,766,633; 5,792,451; 5,853,748; 5,972,387; 5,976,569; and 6,051,561.

The following Examples illustrate the present invention, without in any way limiting the scope thereof.

EXAMPLES

1. Expression cloning :

1) Total ARN purification from five weeks normal mouse retinas :

The cDNA library was constructed from retinas of Five weeks old C57BL/6@N mice generally according to the method of Glissin et al (1974), Biochemistry, 13, 2633-2637. Briefly, after killing, animals were enucleated and the eyes were first placed in Phosphate Buffer Saline (PBS) supplemented with 0,1% DiEthyl PyroCarbonate (DEPC). Neural retina was quickly dissected (the retinal pigmented epithelium was omitted in this tissue preparation). Quickly after each dissection, tissues were homogenised in fresh 6 M Guanidinium Chloride. Ten retinas were pooled in 2.4 ml GC in 4 ml sterile tubes, and the tissue was disrupted completely by strong homogenisation for 1 minute at room temperature.

Messenger RNA (mRNA) purification from five weeks normal mouse retina :

mRNA was isolated on oligo-dT coated porous beads (Oligotex, Qiagen) under stringent conditions according to the method of Kuribayashi et al, (1988). Nucleic Acids Res. Symposium series, 19, 61-64. Briefly, 100-150 µg total mouse retina RNA were mixed with 15 µl of oligo-dT beads in Binding Buffer [10 mM Tris pH 7.5 ; 0.3 M NaCl ; 0.1 M EDTA ; 0.5 % w/v Sodium Duodecyl Sulfate (SDS)] and incubated 6 minutes at 65°C in 0.5 l becher of water, then progressively cooled to room temperature for about 3-4 hours, then centrifuged at room temperature to recover the agarose beads. They were then washed twice by incubating 10 minutes in 0.4 ml of (0.1 M Tris pH 7.5 ; 0.1 M NaCl ; 1 mM EDTA ; 0.5 % w/v SDS). Bounded RNA (mRNA) were eluted in two steps with 50

31494-12

μl of 70°C-warmed RNase-free water, precipitated with 10 μl Sodium Acetate pH 5.2 and 0.25 ml ethanol and incubated 12 hours at -70°C. mRNA were collected by centrifugation (1 hour at 15.000 rpm, followed by two washes with 70% ethanol) and resuspended in 20 μl of RNase-free water. mRNA concentration was measured at 260 nm and the absence of rRNA contaminations was checked by gel electrophoresis under denaturing condition as above.

cDNA synthesis :

was carried out according to the method of Okayama and Berg,(1982),Mol Cell Biol., 2, 161-170. First strand synthesis was primed with 2.5 μg of NotI adaptor oligonucleotide (5'TGTTACCAATCTGAAGTGGGAGCGGCCGACAA(T)₁₈ 3') and incubated for 2 hours with 50 units of modified Moloney Murine Leukaemia virus (M-MLV) reverse transcriptase (Superscript II,* Life Technology) under the conditions recommended by the supplier. For second strand synthesis, reaction was incubated 4 hours at 14°C with 4 units of RNaseH and 100 units of DNA polymerase I in SS Buffer [40 mM Tris pH 7.2 ; 85 mM Potassium Chloride ; 4.4 mM Magnesium chloride ; 3 mM DTT ; 5 μg/ml Bovin Serum Albumin (BSA)] in a final volume of 0.25 ml. EcoRI adapters (5'-OH AATTCGGCACGAGG 3'-OH/3'-OH GCCGTGCTCC5'-PO₄) were ligated at both ends of double strand cDNA in 14 hours at 16°C using 40 units of T4 DNA ligase (Promega, Madison, USA) in a total volume of 20 μl under the conditions recommended by the supplier. The products of this reaction are dscDNA that have an EcoRI half site in 5' and NotI half site in 3' that can be oriented in the cloning vector.

Ligation of dscDNA in pcDNA3 :

was performed according to Maniatis T.(1992), Molecular cloning: a laboratory manual, 2nd ed., using 10 μg of pcDNA3 plasmid (Invitrogen) prepared by cutting with EcoRI and NotI (Promega, Madison,USA) under the conditions specified by the supplier.

Propagation of recombinant clones :

*Trade-mark

31494-12

was performed generally according to the method of Birnboim et al (1979), Nucleic Acids Res., 7, 1513-1523. Briefly, In order to make pools of 100 primary clones, we slightly modified the XL1 Gold^{*} (Stratagene) transformation protocol (provided by Stratagene) as followed. After incubation in growth medium, the transformation reaction was brought to 20 % (vol./vol.) glycerol and 8 % (vol./vol.) horse serum albumin (HAS, Life Technologies). HAS and glycerol prevent death followed freezing thawing. A titration was done by plating on agar plates (100 µg/ml ampicillin) increasing volume of each transformation reaction to calculate the volume giving 100 colonies, while the bulk transformation reaction was stored at -80°C. Recombinant plasmids from the library were purified by 96 at once. In order to prepare 96 pools of 100 clones, the calculated volume corresponding to 100 clones was plated on agar and grown 20 hours at 37°C. The DNA was purified directly from colonies striped from agar plates. A stock of each culture at 23% glycerol was stored at -80°C. The DNA was purified using Qiawell^{*} ultra (Qiagen,) using a protocol recommended by the supplier. Typically, 10 µg of purified plasmid were obtained, the concentration of each preparation was measured using optical density at 260 nm. To sub-divide selected pools of 100 into pools of 10, 50 µl of a 1/250.000 dilution of the glycerol stock from the original pool was plate on an agar plate at 100 µg/ml ampicillin. After growth 16 hours at 37°C, individual 160 colonies were replicated on 16 agar plates (10 per plate) and grown 16 hours at 37°C. The 10 colonies from each plate were harvested and grown in liquid medium [Luria Broth (LB), 100 µg/ml ampicillin] for 3 hours at 37°C. A stock of these cultures at 30% glycerol was stored at -80°C. Plasmid DNA was prepared as before. To divide sub-pools of 10 into individual clones, 50 µl of a 1/250.000 dilution of the glycerol stock of the sub-pools of 10 was plated onto agar plate with 100 µg/ml ampicillin. After growth 16 hours at 37°C, 16 individual colonies were picked and grown 16 hours in 2 ml LB 100 µg/ml ampicillin. A stock of these cultures at 30% glycerol was stored at -80°C. Plasmid DNA was prepared as before.

Transient transfection in Cos-1 cells :

was performed using the method of Chen and Okayama, (1987). High-efficiency transformation of mammalian cells by plasmid DNA (Mol Cell Biol., 7, 2745-2752).

*Trade-mark

31494-12

Chick embryo retinal cultures :

The protocol was adapted from Adler and Hatlee [(1989) Science, 243, 391]. Chick embryonic retina (6 days *in ovo*) is dissociated and plated in monolayer culture. Under these culture conditions with the absence of differentiation signals, cones represent 60-80% of cells. We have produced polyclonal antibodies into rabbit against visinin (a chicken cone marker, Genbank accession number M84729) and verified that the proportion of cones in our culture is of 60-80%. The simple environment of our model (chemically defined medium, absence of cell to cell contacts) in addition to the ease and speed of the method make it a very appropriate system to study trophic factors involved in cone survival. Briefly, Retinas from embryos issued from a control isolate of progenitors, are dissected after six days of development *in ovo*, cells dissociated and plated at low density (10^5 cells/cm²). During ten days cell viability (60-80% cones) was followed using LIVE/DEAD assay (Molecular probes, Eugene, USA) an assay that quantifies live and dead cells. The number of cells alive decreases down to 8% of initial cells number after seven days in culture in chemically defined medium. When performed in the presence of conditioned media from COS1 cells transfected with pools of clones from the library, live cells are counted after seven days *in vitro*.

Chicken progenitors (strain 657 red label) were maintained in a separated compartment for the purpose of this experiment in a hatching facility 25 km from the laboratory. Fertilized eggs obtained naturally were collected weekly, and maintained at 17°C (their biological zero) in the laboratory after hatching. Daily, 5 eggs are incubated for 24 hours at 20°C then 136 hours at 37°C with intermittent reversion of the inclination of the eggs in a humidified chamber. The day of the culture, eggs surface are washed with Mucocit-A* then broken, the chicken embryos are transferred in PBS. The stage of development of the each embryo is verified be the 29th by visual comparison to Hamburger and Hamilton (1951), in (Essential Development Biology, Stern and Holland Ed.). Two of the embryos were chosen and enucleated, the eyes transferred in CO₂-independent medium (Life technologies). Retinas were dissected and transferred in Ringer buffer and washed twice. Retinas are cut in small pieces and treated 20 minutes at 37°C with a solution of trypsin

*Trade-mark

31494-12

(0.25 % w/v). The reaction is stopped by addition of culture media (M199, Life Technologies) supplemented with 10 % inactivated FCS. The cell suspension is treated for few minutes in 25 μ l of DNase I (1mg/ml, Sigma). The cell suspension is then washed twice in Chemical Defined Culture Media [CDCM, equal volumes of DMEM and M199 media (Life Technologies) and AB with Supplements (5 μ g/ml Insulin ; 5 μ g/ml Transferring ; 64 nM Progesterone ; 0.1mM Putrescin ; 5ng/ml Selenium ; 3mM Taurine ; 2.7 μ M Cytidine 5'-diphosphoethanolamine ; 5.2 μ M Cytidine 5'-diphosphocholine, 0.2 μ g/ml Hydrocortisone ; 30 nM 3,3'-5-triiodo-L-thyronine ; 1 mM Sodium Pyruvate), 0.3 μ M Prostaglandin D₂ ; 0.1 mg/ml linoleic acid] in order to remove the FCS. The concentration of cells stained with trypan blue is measured with Mallassez'cell and brought to two concentrations (5.6 and 1.12 10⁵ cells/ml) corresponding to the two plating densities (2 and 4 10⁵ cells/cm²).

Conditioned media from Cos-1 transfected cells are thaw on ice and 50 μ l transferred to two 96 wells tissue culture treated black plate (Corning Costar*) that have been coated a solution of 100 μ g/ml Poly-L-lysine (Sigma) according the plan :

First round of screening :

1	1	1	1	2	C	2	2	2	3	3	P
C	3	3	4	4	4	C	4	5	5	5	5
6	C	6	6	6	7	7	C	7	7	8	8
8	8	C	9	9	9	9	10	C	10	10	10
11	11	11	C	11	12	12	12	12	C	13	13
13	13	14	14	C	14	14	15	15	15	C	15
16	16	16	16	17	C	17	17	17	18	18	C
18	18	19	19	19	19	P	20	20	20	C	20

Where numbers refer to n° of pools of 100 clones, C to conditioned media from Cos-1 cells transfected with the empty vector (pcDNA3) and P a positive control (conditioned media transfected with pcDNA-mouseGDNF).

*Trade-mark

Second and third round of screening :

x.(y).01	x.(y).01	x.(y).01	x.(y).01	x.(y).02	C	x.(y).02	x.(y).02	x.(y).02	x.(y).03	x.(y).03	P
C	x.(y).03	x.(y).03	x.(y).04	x.(y).04	x.(y).04	C	x.(y).04	x.(y).05	x.(y).05	x.(y).05	x.(y).05
x.(y).06	C	x.(y).06	x.(y).06	x.(y).06	x.(y).07	x.(y).07	C	x.(y).07	x.(y).07	x.(y).08	x.(y).08
x.(y).08	x.(y).08	C	x.(y).09	x.(y).09	x.(y).09	x.(y).09	x.(y).10	C	x.(y).10	x.(y).10	x.(y).10
x.(y).11	x.(y).11	x.(y).11	C	x.(y).11	x.(y).12	x.(y).12	x.(y).12	x.(y).12	C	x.(y).13	x.(y).13
x.(y).13	x.(y).13	x.(y).14	x.(y).14	C	x.(y).14	x.(y).14	x.(y).15	x.(y).15	x.(y).15	C	x.(y).15
x.(y).16	x.(y).16	x.(y).16	x.(y).16	x.(y)	C	x.(y)	x.(y)	x.(y)	C57	C57	C
C57	C57	C3H	C3H	C3H	C3H	P	0	0	0	C	0

Where x (second round) and y (third round) represent the n° of the pools selected in the first and second round respectively. 01 to 16 represent the sub-pools used. x(y) represents the parental pool from which the 16 pools derived. C is as is first round. P was modified as pCMVScript-CNTF in the second round and progressively to pcDNA-939.09.08 in the third round. 0 represents chick cone cells in CDCM media only. C57 and C3H, conditioned media from explants of retinas prepared as described in (preparing conditioned media from explants of mouse retinas) from C57BL/6@N and C3H/He@N mouse retinas aged of 5 weeks respectively.

50 µl of cell suspensions corresponding to two densities (2 and 4 10⁵ cells/cm²) were added to the well of the two 96 wells plates filled conditioned media with a 8 channel motorized pipette (Biohit) in order to minimized experimental errors. Cells were incubated for 7 days at 37°C in 5% CO₂.

Preparing conditioned media from explants of mouse retinas :

The second and third round screening included positive controls adapted from Mohand-Saïd et al. (1998). Five weeks aged mouse 5C57BL/6@N (wild type) and C3H/He@N (*rd1*) were sacrificed and enucleated. Two retinas were dissected and incubated in 24 hours at 37°C in 5% CO₂ in 1.5 ml of CDCM in 12 wells plates. Conditioned media were

31494-12

recovered and concentrated by a factor 40 by ultra-filtration on Vivaspin* (Sartorius, cutting point 10 kDa). Conditioned media were frizzed in liquid nitrogen and stored in aliquots at -20°C before used. The day of use, conditioned media were thawed on ice, diluted 10 times in CDCM and sterilized by filtration on 0.22 μm filter (Acrodisk 13, Gelman Sciences).

Functional assay, Live/Dead assay :

The functional assay is based on the number of chicken retinal cells alive after 7 days incubation *in vitro*. We used Live/Dead assay kit (Molecular probes, Eugene, USA) that is based on the use of two fluorogenic dyes (Calcein AM* and Ethidium dimer) that stain live and dead cells respectively. A cell that is alive processes a metabolic activity (here an esterase activity) that converts the substrate (Calcein AM) in its fluorescent product emitting at 520 nm. The membrane permeability of a dead cell is altered and permits the DNA staining of the nucleus by Ethidium dimmer emitting at 635 nm. A cell is alive : emitting at 520 nm after excitation at 485 nm, or dead : emitting at 635 nm after excitation at 520 nm. Using epifluorescence microscopy, the two types of fluorescent cells can be visualized separately. After 7 days *in vitro*, cells were incubated for 30 minutes at room temperature in the dark with 2.7 μM Calcein-AM and 0,3 mM Ethidium dimer.

Image acquisition :

Briefly, image acquisition consisted of autofocussing each well, automatic cell counting in two fluorescence's followed by processing of the raw data using specialist software e.g. Metamorph* (Universal Imaging Corporation, West Chester, USA) to obtain digitalised pictures of each well of the plate. We used a inverted microscope (Nikon TE 200) equipped with a mercury epifluorescent lamp with two excitation filters 485 and 520 nm, two emission filters 520 and 635 nm, an object (x10), a computer driven motorized platine (Multicontrol 2000*, Martzauzer* and a CCD camera (Cohu).

To record the plate, it is positioned on the motorised platine, and the focus done manually on the first well and this plane is recorded (z origin). The threshold of the image dead and

*Trade-mark

live is set from the first well. The centre of the first well is adjusted using white light by aligning manually the bottom of the first well to the bottom of the image on the computer monitor, then by aligning the extreme right of the first well to the right of the image on the computer screen and to record the two positions. The centre of the first well is calculated and gives the position of the centre of each well of the plate. We have noticed in the development process that there is a slight higher density of cells at the edge of the well and excluded the edge from the acquisitions. It is important that the image from each well be centred perfectly in order to avoid any misleading results. When set-up, the first scan of the plate executes a recording of dead cells. The dead cells density is the less variable under these conditions. The application executes an auto focusing by taking images at different focal plans and choosing the brightest one, the right focus. This z position is stored and the platine executes programmed movements in the x and y axes taking a total of 4 images that when reconstituted in one image represents 2/3 of the surface of the well. A pile of images from the focus plans is stored for control. The platine executes an auto focusing and four acquisitions for each well of the plate starting by wells A1 to A12, then B12 to B1, C1 to C12 etc... At the end, the platine moves outside the plate in order to overexpose the last well (H1). The scanning of dead cells takes 30 minutes. The second scan (live cells) is executed after switching the filter. This second scan is using the recorded z positions of each well from the dead scan. Four images are taken from each well as for dead cells. At the end of the second scan (22 minutes) dead and live reconstituted images are stored in a file that is named automatically with the date of the day. Cells numbers (dead and live) are counted automatically with pre-set morphometric parameters (average) and displayed on the computer monitor in order to check if the experiment is correct. It is important to check on a daily basis that the number of cells alive is not too high. We have observed that if plated at too important density, the chicken retinal cells survived longer most likely by producing their own survival factor. We screened cells in the absence of this effect. Before scanning the second plate (same experiment with twice density of cells plated), we add an α at the end of the name of the log file from the first plate. Images of each experiment were stored on CD-rom. We have generated a library of more than 250 CD-rom.

Cells counting and selection of the pools :

The cells numbers (live and dead) were counted using images of each experiment stored on CD-rom. The log file from an experiment was first loaded on a computer (counting off-line) and opened using Metamorph software. In a first step, the images corresponding to the 14 wells C (conditioned media from Cos-1 cells transfected with the empty vector) are opened. After adjusting the threshold of the image, the command Integrated Morphometry Analysis is used to measure the distribution of the areas between 10 and 250 of the object (number of live cells for each total area) for these control wells. The distribution follows a Gaussian curve with the maximal number of objects corresponding to an isolated cell. This standard value (SV) is then used to calculate the value of the area above which an object will be counted as two cells (standard object cut, SOC) through our empirical function: $SOC = 29/20.74 \text{ SV}$. The SOC value of each individual plate is used to count the number of live cells of the plate. These numbers are then transferred into a excel table.

For the first round of screening, the value were plotted for each pool as fold difference (increase or decrease live cells numbers) versus the average of the 14 control wells + the standard deviation. In order to redress the variations coming from the position difference within the plate, we calculated the average of the fold difference individually between the 80 wells corresponding to positions where the pools are tested and the 14 wells control for 200 independent plates. In average, the differences observed are due only to position difference and cells numbers were corrected with this coefficient. In order to discriminate in more stringent way the pools to be selected, the live cells counts were plotted as fold difference versus as control all values not exceeding 1.3 but exceeding 0.4. In this manner, all pools were fold difference versus the 14 wells corresponding to empty vector are in the interval of 0.4 to 1.3 are considered as having no effect and used as control. After correction, the fold difference versus control of the two plates was multiplied and the result sorted by decreasing fold difference. The pools on the top of that list were further checked by visual inspection of the graph corresponding to the 20 pools of the experiment (both plates) and of the images of live and dead cells to avoid to screen misleading pools.

31494-12

For the second and third round, the plates screening for sub-pools include additional control. We prepared conditioned media from retina explants of 5 weeks of age. We selected experiments where positive effects were recorded for retina explants from C57BL/6@N and dismissed the others. The results were plotted as fold difference versus the 14 control wells, no recalculation of the control was made. The fold difference versus control of the two plates was multiplied and the results were sorted by decreasing fold difference for the 16 sub-pools.

Isolated cDNA were sequenced using T7 primer (5' GTAATACGACTCACTATAGGGC 3') on a capillary sequencer (CEQ2000^{*}, Beckman Coulter). DNA sequence was compared to databases using Basic Local Alignment Search Tool (BLAST).

Identification of Rdcvf2 and human homologues:

Using the Rdcvf1 sequence (the nucleotide sequence encoding the polypeptides set forth in SEQ ID NO:2 or SEQ ID NO:4) and BLAST, homologous murine and human polypeptides were identified (Figure 8). EST clones with homology to mouse RdCVF2 (GenBank Accession No: bc016199) were identified: GenBank Accession Nos: be552141, bi517442, bg707818, bi603812, ai433287, be088414, bg297383, bg297304 (see also Figure 12). EST clones with homology to mouse Rdcvf1 (SEQ ID NO:1) were identified: GenBank Accession Nos: bg299078, ai716631, bg294111, be108041, bg395178 (see also Figure 13).

Real-time RT-PCR analysis of Rdcvf1 expression :

The retinal expression of Rdcvf1 by mouse C57BL/6@N and C3H/He@N aged as well as congenic C3H (+/+ and rd/rd) of 5 weeks is studied using real-time RT-PCR on a lightcycler (Roche) with sybergreen^{*} PCR kit (Roche). cDNAs are produced by priming with a random hexamer oligonucleotide (pdN6, Amersham), M-MLV reverse transcriptase (superscript II, Life Technologies) and total RNA from mouse retina prepared as in 1). cDNAs are normalized using a ubiquitous messenger Glucose-6-Phosphate DeHydrogenase (G6PDH). 0.2 µl of first strand cDNAs synthesis (an equivalent of 10 ng of total RNA) is amplified with 2 µM of the oligonucleotides of SEQ ID NO:24 and SEQ ID NO:25 in triple in a total volume of 25 µl using the following

*Trade-mark

program : 30 seconds at 95°C, and 35 cycles of a sequence (1 second at 95°C, 18 seconds at 55°C, 10 seconds at 72°C), The analysis (Figure 16) shows that *Rdcvf1* expression decreases after rod degeneration in the *rd1* mouse (C3H/He@N). *Rdcvf1* was also shown to be directly expressed by photoreceptors by real-time RT-PCR using RNA prepared from outer layer of the retina by vibratome sectionning. Products were checked by agarose gel electrophoresis. Similar results are obtained with another pair of *Rdcvf1* specific primers. As a positive control, the expression of rod arrestin (Ass n° M24086) is monitored in the same conditions with primers (5' CTATTACGTCAAGCCTGTAGCC 3' and 5' CATCCTCATCTTTCTTCCCTTC 3'). Confirmation that *Rdcvf1* is the cone protective factor can be obtained by adding a suitable amount of *Rdcvf1* to a retinal explant of 5 week old *rd1* mouse (C3H/He@N). What is a suitable amount can be arrived at by some initial titration experiments. Compare with appropriate controls after 7 days cone survival will be increased.

RT-PCR Analysis of *Rdcvf2*

RT-PCR for *Rdcvf2* expression was carried out using primers 5' GCCAGCGTTTTCTGCCTTTTAC 3' and 5' AAGCCCTGCCTGCTCTAACATC 3'. The analysis shows that *RdCVF2* is expressed in a rod-dependent manner and that *Rdcvf2* expression is not restricted to the retina, but that also other neuronal cells express *Rdcvf2*. (Figure 17), whereas the expression of *Rdcvf1* seems to be restricted to retinal cells.

Live/Death Assays of *Rdcvf1* or *Rdcvf2*

COS-1 cells are transfected with a suitable expression vector carrying the *Rdcvf1* or *Rdcvf2* under control of a inducible promoter. Control cells are transfected with the empty vector. The cells are incubated for a suitable period of time upon induction of *Rdcvf1* or *Rdcvf2* expression. Subsequently, the number of surviving cone cells incubated with conditioned media from COS-1 cells transfected with *Rdcvf1* or *Rdcvf2* and the number of surviving control cells are counted according to the above described method. The cells expressing *Rdcvf1* or *Rdcvf2* show a significantly higher amount of surviving cells.

31494-12

Rod specific factor

Real-time RT-PCR analysis, carried out under standard conditions, of the expression of rod arrestin (control) and Rdcvf1 in 5 weeks retinal explants from C57BL/6@N 5 weeks and C3H/HE@N, using the primers:

SEQ ID NO: 24 : 5' TCTATGTGTCCCAGGACCCTACAG 3'

SEQ ID NO: 25 : 5' TTTATGCACAAGTAGTACCAGGACAG 3'

demonstrates that RdCVF1 is expressed only in the presence of rods (Rod-derivedCVF1).

Production of polyclonal antibodies :

Polyclonal antibodies are prepared by injecting a Glutathion-S-Transferase (GST) purified fusion protein (GST-Rdcvf1) as well as from mouse RdCVF1 peptide sequence amino acids 11 to 32 from SEQ ID N02 (Ab n°2) and peptide sequence amino acids 79 to 96 from SEQ ID N02 (Ab n°3) into rabbit. The fusion construct pGST-Rdcvf1 is prepared by amplification with oligonucleotides of SEQ ID NO:26 and SEQ ID NO:27 using pcDNA-Rdcvf1 as template under standard conditions. The Open Reading Frame (ORF) of Rdcvf1 is cloned in frame into pGex2TK (Pharmacia) between the BamHI and EcoRI restriction sites, and transformed into E. coli [BL21 (DE3) pLysS, Promega] by standard procedure. A single colony is grown in 3 litres of LB liquid media with 100 µg/ml ampicillin at 30°C and protein production is induced by addition of 1 µg/ml isopropylthio-β-D-galactoside (IPTG) and continued for 5 hours at 30°C. Cells are harvested, lysed by sonication and purified onto glutathion sepharose* under standard protocol. The fusion protein is eluted with 10 mM reduced glutathion at room temperature. The Eluted protein is dialyzed into PBS before injection to rabbits. Protein purity is monitored by polyacrylamide gel electrophoresis. Two rabbits are immunized by intradermic injection at 80 sites of 100 µg of purified GST-Rdcvf1. Serum is harvested after 8 weeks.

*Trade-mark

New Seq RdCVF.ST25.txt
SEQUENCE LISTING

<110> UNIVERSITE LOUIS PASTEUR/ NOVARTIS AG

<120> Disease associated protein

<130> 4-31883

<160> 35

<170> PatentIn version 3.1

<210> 1

<211> 468

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (45)..(374)

<223>

<400> 1

atcggatccc tctctggggtc cccagctcct tgcatactgc tacc atg gca tct ctc	56
Met Ala Ser Leu	
1	
ttc tct gga cgc atc ttg atc agg aac aac agc gac cag gat gaa gtg	104
Phe Ser Gly Arg Ile Leu Ile Arg Asn Asn Ser Asp Gln Asp Glu Val	
5 10 15 20	
gag aca gag gca gag ctg agc cgt agg tta gag aat cgt ctg gtg ttg	152
Glu Thr Glu Ala Glu Leu Ser Arg Arg Leu Glu Asn Arg Leu Val Leu	
25 30 35	
ctg ttc ttc ggc gcc ggc gcc tgt ccc cag tgc cag gcc ttt gcc cca	200
Leu Phe Phe Gly Ala Gly Ala Cys Pro Gln Cys Gln Ala Phe Ala Pro	
40 45 50	
gtc ctc aaa gac ttc ttc gtg cgg ctc act gac gag ttc tac gtg ctg	248
Val Leu Lys Asp Phe Phe Val Arg Leu Thr Asp Glu Phe Tyr Val Leu	
55 60 65	
cgg gca gca cag ctg gcc ctg gtc tat gtg tcc cag gac cct aca gag	296
Arg Ala Ala Gln Leu Ala Leu Val Tyr Val Ser Gln Asp Pro Thr Glu	

New Seq RdCVF.ST25.txt

70 75 80

gag caa cag gac ctc ttc ctc agg gac atg cct gaa aaa tgg ctc ttc 344
 Glu Gln Gln Asp Leu Phe Leu Arg Asp Met Pro Glu Lys Trp Leu Phe
 85 90 95 100

ctg ccg ttc cat gat gaa ctg agg agg tga ggccccaggg aagaccaggg 394
 Leu Pro Phe His Asp Glu Leu Arg Arg
 105

agggcttcct ggagaaggca tttccctgga ggtttactgt cctggtacta cttgtgcata 454
 aagaggtatt cctc 468

<210> 2

<211> 109

<212> PRT

<213> Mus musculus

<400> 2

Met Ala Ser Leu Phe Ser Gly Arg Ile Leu Ile Arg Asn Asn Ser Asp
 1 5 10 15

Gln Asp Glu Val Glu Thr Glu Ala Glu Leu Ser Arg Arg Leu Glu Asn
 20 25 30

Arg Leu Val Leu Leu Phe Phe Gly Ala Gly Ala Cys Pro Gln Cys Gln
 35 40 45

Ala Phe Ala Pro Val Leu Lys Asp Phe Phe Val Arg Leu Thr Asp Glu
 50 55 60

Phe Tyr Val Leu Arg Ala Ala Gln Leu Ala Leu Val Tyr Val Ser Gln
 65 70 75 80

Asp Pro Thr Glu Glu Gln Gln Asp Leu Phe Leu Arg Asp Met Pro Glu
 85 90 95

Lys Trp Leu Phe Leu Pro Phe His Asp Glu Leu Arg Arg
 100 105

<210> 3

<211> 764

<212> DNA

<213> Mus musculus

<220>

New Seq RdCVF.ST25.txt

<221> CDS

<222> (26)..(676)

<223>

<400> 3

```

ccccagctcc ttgcatactg ctacc atg gca tct ctc ttc tct gga cgc atc      52
                        Met Ala Ser Leu Phe Ser Gly Arg Ile
                        1      5

ttg atc agg aac aac agc gac cag gat gaa gtg gag aca gag gca gag      100
Leu Ile Arg Asn Asn Ser Asp Gln Asp Glu Val Glu Thr Glu Ala Glu
10      15      20      25

ctg agc cgt agg tta gag aat cgt ctg gtg ttg ctg ttc ttc ggc gcc      148
Leu Ser Arg Arg Leu Glu Asn Arg Leu Val Leu Leu Phe Phe Gly Ala
30      35      40

ggc gcc tgt ccc cag tgc cag gcc ttt gcc cca gtc ctc aaa gac ttc      196
Gly Ala Cys Pro Gln Cys Gln Ala Phe Ala Pro Val Leu Lys Asp Phe
45      50      55

ttc gtg cgg ctc act gac gag ttc tac gtg ctg cgg gca gca cag ctg      244
Phe Val Arg Leu Thr Asp Glu Phe Tyr Val Leu Arg Ala Ala Gln Leu
60      65      70

gcc ctg gtc tat gtg tcc cag gac cct aca gag gag caa cag gac ctc      292
Ala Leu Val Tyr Val Ser Gln Asp Pro Thr Glu Glu Gln Gln Asp Leu
75      80      85

ttc ctc agg gac atg cct gaa aaa tgg ctc ttc ctg ccg ttc cat gat      340
Phe Leu Arg Asp Met Pro Glu Lys Trp Leu Phe Leu Pro Phe His Asp
90      95      100      105

gaa ctg agg agg gac ctc ggg cgc cag ttc tct gtc cgt caa ctg cca      388
Glu Leu Arg Arg Asp Leu Gly Arg Gln Phe Ser Val Arg Gln Leu Pro
110      115      120

gcg gtt gtg gta ctt aag cct ggt ggg gac gtg ctg aca agc gac gcc      436
Ala Val Val Val Leu Lys Pro Gly Gly Asp Val Leu Thr Ser Asp Ala
125      130      135

acg gag gag atc cag cgt ctg gga ccc gcc tgc ttt gcc aac tgg cag      484
Thr Glu Glu Ile Gln Arg Leu Gly Pro Ala Cys Phe Ala Asn Trp Gln
140      145      150

gag gcc gca gag ctc ctg gac cgc agc ttc ctg caa ccg gag gat ttg      532
Glu Ala Ala Glu Leu Leu Asp Arg Ser Phe Leu Gln Pro Glu Asp Leu
155      160      165

gat gag cct gcg cgg cgc agc atc acc gag cct ctg cgc cgt cgc aag      580
Asp Glu Pro Ala Arg Arg Ser Ile Thr Glu Pro Leu Arg Arg Arg Lys
170      175      180      185

tac cga gta gac cgg gat gtc ggc ggg agc ggg gcg aaa cgg cgc gac      628
Tyr Arg Val Asp Arg Asp Val Gly Gly Ser Gly Ala Lys Arg Arg Asp
190      195      200

tct ggt gaa ccc cag ggg gac gcg ggt aca agg gcg gag ctc tgg tga      676
Ser Gly Glu Pro Gln Gly Asp Ala Gly Thr Arg Ala Glu Leu Trp
205      210      215

ctcccagggt aggagtgggg accggagctc tggtgacacc aaagtaccgg tgcacgaccg      736

```

New Seq RdCVF.ST25.txt

aggttgatga ccctcccgaa ggaaccgg

764

<210> 4

<211> 216

<212> PRT

<213> Mus musculus

<400> 4

Met Ala Ser Leu Phe Ser Gly Arg Ile Leu Ile Arg Asn Asn Ser Asp
 1 5 10 15

Gln Asp Glu Val Glu Thr Glu Ala Glu Leu Ser Arg Arg Leu Glu Asn
 20 25 30

Arg Leu Val Leu Leu Phe Phe Gly Ala Gly Ala Cys Pro Gln Cys Gln
 35 40 45

Ala Phe Ala Pro Val Leu Lys Asp Phe Phe Val Arg Leu Thr Asp Glu
 50 55 60

Phe Tyr Val Leu Arg Ala Ala Gln Leu Ala Leu Val Tyr Val Ser Gln
 65 70 75 80

Asp Pro Thr Glu Glu Gln Gln Asp Leu Phe Leu Arg Asp Met Pro Glu
 85 90 95

Lys Trp Leu Phe Leu Pro Phe His Asp Glu Leu Arg Arg Asp Leu Gly
 100 105 110

Arg Gln Phe Ser Val Arg Gln Leu Pro Ala Val Val Val Leu Lys Pro
 115 120 125

Gly Gly Asp Val Leu Thr Ser Asp Ala Thr Glu Glu Ile Gln Arg Leu
 130 135 140

Gly Pro Ala Cys Phe Ala Asn Trp Gln Glu Ala Ala Glu Leu Leu Asp
 145 150 155 160

Arg Ser Phe Leu Gln Pro Glu Asp Leu Asp Glu Pro Ala Arg Arg Ser
 165 170 175

Ile Thr Glu Pro Leu Arg Arg Arg Lys Tyr Arg Val Asp Arg Asp Val
 180 185 190

Gly Gly Ser Gly Ala Lys Arg Arg Asp Ser Gly Glu Pro Gln Gly Asp
 195 200 205

Ala Gly Thr Arg Ala Glu Leu Trp
210 215

<210>	5
<211>	353
<212>	DNA
<213>	Homo sapiens

<220>
<221> CDS
<222> (24) .. (353)
<223>

[illegible]

<210>	6
<211>	109
<212>	PRT
<213>	Homo sapiens

New Seq RdCVF.ST25.txt

<400> 6

Met Ala Ser Leu Phe Ser Gly Arg Ile Leu Ile Arg Asn Asn Ser Asp
 1 5 10 15

Gln Asp Glu Leu Asp Thr Glu Ala Glu Val Ser Arg Arg Leu Glu Asn
 20 25 30

Arg Leu Val Leu Leu Phe Phe Gly Ala Gly Ala Cys Pro Gln Cys Gln
 35 40 45

Ala Phe Val Pro Ile Leu Lys Asp Phe Phe Val Arg Leu Thr Asp Glu
 50 55 60

Phe Tyr Val Leu Arg Ala Ala Gln Leu Ala Leu Val Tyr Val Ser Gln
 65 70 75 80

Asp Ser Thr Glu Glu Gln Gln Asp Leu Phe Leu Lys Asp Met Pro Lys
 85 90 95

Lys Trp Leu Phe Leu Pro Phe Glu Asp Asp Leu Arg Arg
 100 105

<210> 7

<211> 686

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (48)..(686)

<223>

<400> 7

ccggggacca cacgccgcgc tgtccccagc acccaaccca gggtacc atg gcc tcc 56
 Met Ala Ser
 1

ctg ttc tct ggc cgc atc ctg atc cgc aac aat agc gac cag gac gag 104
 Leu Phe Ser Gly Arg Ile Leu Ile Arg Asn Asn Ser Asp Gln Asp Glu
 5 10 15

ctg gat acg gag gct gag gtc agt cgc agg ctg gag aac cgg ctg gtg 152
 Leu Asp Thr Glu Ala Glu Val Ser Arg Arg Leu Glu Asn Arg Leu Val
 20 25 30 35

ctg ctg ttc ttt ggt gct ggg gct tgt cca cag tgc cag gcc ttc gtg 200
 Leu Leu Phe Phe Gly Ala Gly Ala Cys Pro Gln Cys Gln Ala Phe Val
 40 45 50

New Seq RdCVF.ST25.txt

ccc atc ctc aag gac ttc ttc gtg cgg ctc aca gat gag ttc tat gta	248
Pro Ile Leu Lys Asp Phe Phe Val Arg Leu Thr Asp Glu Phe Tyr Val	
55 60 65	
ctg cgg gcg gct cag ctg gcc ctg gtg tac gtg tcc cag gac tcc acg	296
Leu Arg Ala Ala Gln Leu Ala Leu Val Tyr Val Ser Gln Asp Ser Thr	
70 75 80	
gag gag cag cag gac ctg ttc ctc aag gac atg cca aag aaa tgg ctt	344
Glu Glu Gln Gln Asp Leu Phe Leu Lys Asp Met Pro Lys Lys Trp Leu	
85 90 95	
ttc ctg ccc ttt gag gat gat ctg agg agg gac ctc ggg cgc cag ttc	392
Phe Leu Pro Phe Glu Asp Asp Leu Arg Arg Asp Leu Gly Arg Gln Phe	
100 105 110 115	
tca gtg gag cgc ctg ccg gcg gtc gtg gtg ctc aag ccg gac ggg gac	440
Ser Val Glu Arg Leu Pro Ala Val Val Val Leu Lys Pro Asp Gly Asp	
120 125 130	
gtg ctc act cgc gac ggc gcc gac gag atc cag cgc ctg ggc acc gcc	488
Val Leu Thr Arg Asp Gly Ala Asp Glu Ile Gln Arg Leu Gly Thr Ala	
135 140 145	
tgc ttc gcc aac tgg cag gag gcg gcc gag gtg ctg gac cgc aac ttc	536
Cys Phe Ala Asn Trp Gln Glu Ala Ala Glu Val Leu Asp Arg Asn Phe	
150 155 160	
cag ctg cca gag gac ctg gag gac cag gag cca cgg agc ctc acc gag	584
Gln Leu Pro Glu Asp Leu Glu Asp Gln Glu Pro Arg Ser Leu Thr Glu	
165 170 175	
tgc ctg cgc cgc cac aag tac cgc gtg gaa aag gcg gcg cga ggc ggg	632
Cys Leu Arg Arg His Lys Tyr Arg Val Glu Lys Ala Ala Arg Gly Gly	
180 185 190 195	
cgc gac ccc ggg gga ggg ggt ggg gag gag ggc ggg gcc ggg ggg ctg	680
Arg Asp Pro Gly Gly Gly Gly Glu Glu Gly Gly Ala Gly Gly Leu	
200 205 210	
ttc tga	686
Phe	

<210> 8

<211> 212

<212> PRT

<213> Homo sapiens

<400> 8

Met Ala Ser Leu Phe Ser Gly Arg Ile Leu Ile Arg Asn Asn Ser Asp
1 5 10 15

Gln Asp Glu Leu Asp Thr Glu Ala Glu Val Ser Arg Arg Leu Glu Asn
20 25 30

Arg Leu Val Leu Leu Phe Phe Gly Ala Gly Ala Cys Pro Gln Cys Gln

New Seq RdCVF.ST25.txt
40 45

35

Ala Phe Val Pro Ile Leu Lys Asp Phe Phe Val Arg Leu Thr Asp Glu
50 55 60

Phe Tyr Val Leu Arg Ala Ala Gln Leu Ala Leu Val Tyr Val Ser Gln
65 70 75 80

Asp Ser Thr Glu Glu Gln Gln Asp Leu Phe Leu Lys Asp Met Pro Lys
85 90 95

Lys Trp Leu Phe Leu Pro Phe Glu Asp Asp Leu Arg Arg Asp Leu Gly
100 105 110

Arg Gln Phe Ser Val Glu Arg Leu Pro Ala Val Val Val Leu Lys Pro
115 120 125

Asp Gly Asp Val Leu Thr Arg Asp Gly Ala Asp Glu Ile Gln Arg Leu
130 135 140

Gly Thr Ala Cys Phe Ala Asn Trp Gln Glu Ala Ala Glu Val Leu Asp
145 150 155 160

Arg Asn Phe Gln Leu Pro Glu Asp Leu Glu Asp Gln Glu Pro Arg Ser
165 170 175

Leu Thr Glu Cys Leu Arg Arg His Lys Tyr Arg Val Glu Lys Ala Ala
180 185 190

Arg Gly Gly Arg Asp Pro Gly Gly Gly Gly Gly Glu Glu Gly Gly Ala
195 200 205

Gly Gly Leu Phe
210

<210> 9

<211> 600

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (265)..(570)

<223>

New Seq RdCVF.ST25.txt

Phe Glu Val Val Phe Val Ser Ala Asp Gly Ser Ala Glu Glu Met Leu
65 70 75 80

Asp Phe Met Arg Glu Leu His Gly Ser Trp Leu Ala Leu Pro Phe His
85 90 95

Asp Pro Tyr Arg Gln
100

<210> 11

<211> 1164

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (300)..(770)

<223>

<400> 11

ttgactctgg	tgggtagaga	gggtttgcaa	ggcaggataa	aatagagggg	gggagagggt	60
gatggcgtgg	ctctgctttt	tgggtgcggg	caccagctgt	catcgctgct	gtcgcagctt	120
ctggagtggc	cactgtgctc	tctcctccct	tcggctcaag	gtgagctggt	ccagcagaag	180
gcggggctga	gaggcgccta	gtgctgcggg	aggctcagtg	tcattctcca	gctaacaggt	240
ggccgtgcag	cccagggctc	gtctctccac	tgtgtcctct	tcacgccgag	ctcgtggcg	299
atg gtg gac gtg ctg ggc ggg cgg cgc ctg gtg acc cgg gag ggc acg	347					
Met Val Asp Val Leu Gly Gly Arg Arg Leu Val Thr Arg Glu Gly Thr						
1 5 10 15						
gtg gtg gag gcc gag gtg gcg ctg cag aac aag gtg gta gct ttg tac	395					
Val Val Glu Ala Glu Val Ala Leu Gln Asn Lys Val Val Ala Leu Tyr						
20 25 30						
ttt gcg gcg ggc cgg tgc tcg ccc agc cgc gac ttc acg ccg ctg ctc	443					
Phe Ala Ala Gly Arg Cys Ser Pro Ser Arg Asp Phe Thr Pro Leu Leu						
35 40 45						
tgc gac ttc tac acg gag ctg gtg agc gag gcg cgg cgg ccc gct ccc	491					
Cys Asp Phe Tyr Thr Glu Leu Val Ser Glu Ala Arg Arg Pro Ala Pro						
50 55 60						
ttc gag gtg gtt ttc gtg tcg gca gac ggc agt gcg gag gag atg ttg	539					
Phe Glu Val Val Phe Val Ser Ala Asp Gly Ser Ala Glu Glu Met Leu						
65 70 75 80						
gac ttc atg cgc gag ctg cac ggc tcc tgg ctg gca ttg ccc ttc cac	587					
Asp Phe Met Arg Glu Leu His Gly Ser Trp Leu Ala Leu Pro Phe His						
85 90 95						

New Seq RdCVF.ST25.txt

```

gac ccc tac cgg cat gaa ctg aag aag agg tac gaa atc acc gcc atc      635
Asp Pro Tyr Arg His Glu Leu Lys Lys Arg Tyr Glu Ile Thr Ala Ile
100                               105                               110

ccc aag ctg gtg gtc atc aag cag aac gga gct gtc atc acc aac aaa      683
Pro Lys Leu Val Val Ile Lys Gln Asn Gly Ala Val Ile Thr Asn Lys
115                               120                               125

ggg cgg aag cag atc cga gag cgc ggg cta gct tgc ttt cag aac tgg      731
Gly Arg Lys Gln Ile Arg Glu Arg Gly Leu Ala Cys Phe Gln Asn Trp
130                               135                               140

gtg gaa gca gcc gat gtt ttc caa aac ttc tcg ggg tga ccagggcagt      780
Val Glu Ala Ala Asp Val Phe Gln Asn Phe Ser Gly
145                               150                               155

tgctggaagt tcagggcaac tatcttcaaa aagggcttag ctggttccct tctctgctga      840

ggaatgtcat tntagagtca ccatgctgtg acagagagca taaactgctc aggaaagaac      900

tacgtctgcc ccctgtgggt cctagagctc cgttgaatgt ttatttctta cacctttctc      960

caccggtgcc taggatccag gacacatcag ccacgagtta acagaactct atgcaagatg    1020

ctctttccta caggaaattt ctttgataaa ttgacctatg gaggtgatac attttctgat    1080

gacatttttg tgatgctttg gtaaacgtat ttattactcg ggtttgtaga ctgtgtaatt    1140

taataaacca acactcacac tttg                                           1164

```

<210> 12

<211> 156

<212> PRT

<213> Mus musculus

<400> 12

```

Met Val Asp Val Leu Gly Gly Arg Arg Leu Val Thr Arg Glu Gly Thr
1                               5                               10                               15

Val Val Glu Ala Glu Val Ala Leu Gln Asn Lys Val Val Ala Leu Tyr
20                               25                               30

Phe Ala Ala Gly Arg Cys Ser Pro Ser Arg Asp Phe Thr Pro Leu Leu
35                               40                               45

Cys Asp Phe Tyr Thr Glu Leu Val Ser Glu Ala Arg Arg Pro Ala Pro
50                               55                               60

Phe Glu Val Val Phe Val Ser Ala Asp Gly Ser Ala Glu Glu Met Leu
65                               70                               75                               80

Asp Phe Met Arg Glu Leu His Gly Ser Trp Leu Ala Leu Pro Phe His
85                               90                               95

```

New Seq RdCVF.ST25.txt

Asp Pro Tyr Arg His Glu Leu Lys Lys Arg Tyr Glu Ile Thr Ala Ile
 100 105 110

Pro Lys Leu Val Val Ile Lys Gln Asn Gly Ala Val Ile Thr Asn Lys
 115 120 125

Gly Arg Lys Gln Ile Arg Glu Arg Gly Leu Ala Cys Phe Gln Asn Trp
 130 135 140

Val Glu Ala Ala Asp Val Phe Gln Asn Phe Ser Gly
 145 150 155

<210> 13

<211> 1472

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (331)..(738)

<223>

<400> 13

gtgtgggagg ggcgcagttg ggggaggggtg cagagacctg agggcttgag gttgcctggc 60
 tggccccgct cccagaggcg ggtgccgcgc tgtcgcccag gtatctgggg tctctggtgt 120
 ctgagtgtct cattgtcggc gcgaacacaa ttgctccagc cacaggcgag gcctggccaa 180
 ggtgtgggag catctagggc aggtcttgag aggtccagcg cccggtgggtg cggacagagg 240
 cggggcaccg cggcgctcgc cgccgcctcc ccgcaggtga tcacctcct gcaggtgtcc 300
 tcgggtctca ggtggctgcg tgtctgcgcc atg gtt gac att ctg ggc gag cgg 354
 Met Val Asp Ile Leu Gly Glu Arg
 1 5

cac ctg gtg acc tgt aag ggc gcg acg gtg gag gcc gag gcg gcg ctg 402
 His Leu Val Thr Cys Lys Gly Ala Thr Val Glu Ala Glu Ala Ala Leu
 10 15 20

cag aac aag gtg gtg gca ctg tac ttc gcg gcg gcc cgg tgc gcg ccg 450
 Gln Asn Lys Val Val Ala Leu Tyr Phe Ala Ala Ala Arg Cys Ala Pro
 25 30 35 40

agc cgc gac ttc acg ccg ctg ctc tgc gac ttc tat acg gcg ctg gtg 498
 Ser Arg Asp Phe Thr Pro Leu Leu Cys Asp Phe Tyr Thr Ala Leu Val
 45 50 55

gcc gag gcg cgg cgg ccc gcg ccc ttc gaa gtg gtc ttc gtg tca gcc 546
 Ala Glu Ala Arg Arg Pro Ala Pro Phe Glu Val Val Phe Val Ser Ala
 60 65 70

New Seq RdCVF.ST25.txt

gac ggc agc tgc cag gag atg ctg gac ttc atg cgc gag ctg cat ggc 594
 Asp Gly Ser Cys Gln Glu Met Leu Asp Phe Met Arg Glu Leu His Gly
 75 80 85

gcc tgg ctg gcg ctg ccc ttc cac gac ccc tac cgg caa cgg agt ctc 642
 Ala Trp Leu Ala Leu Pro Phe His Asp Pro Tyr Arg Gln Arg Ser Leu
 90 95 100

gct ctg ttg ccc agg ctg gag tgc agt ggc gtg atc tta gct cac tgc 690
 Ala Leu Leu Pro Arg Leu Glu Cys Ser Gly Val Ile Leu Ala His Cys
 105 110 115 120

aac ctt tgc ctc ctg ggt tca agt gat tct cta gcc tta gcc tcc tga 738
 Asn Leu Cys Leu Leu Gly Ser Ser Asp Ser Leu Ala Leu Ala Ser
 125 130 135

gcatctggga ctacagccat tgctgtgaat tacgtgaggg aaagatattg aagaggagtt 798
 ggacactccg agagtgcagc tggttctcccc ccgcaccatc cgtgtcctgc attctgagag 858
 tctgtgctca ttaacaatgt gctgtgacca tgtgactcag caatcctgct gctgggtata 918
 taccgaaag aaaggaaaag gaagccagta tattgaagag gtatctgcac ccccatgttt 978
 attgcagcac tgttcacaac agccaagatt tggaagcaac ctaagtgtcc atcaacagat 1038
 gaatggataa agaaaacgtg gtacatatata acaatggagt actcttcagc cattaataaaa 1098
 atgagattct gtcatttgca ataatataga tggaaaagga ggcccttatg tgaagtgaag 1158
 taagccaggc acagaaagac aaacatcaca tggttctcact tatttgtggg atctaattgat 1218
 caaaacaatt gaactcttgg acatagagag tagaagggtg gttaccagaa gctggaaagg 1278
 aaagtggggg tgggaggaag gtgggaatgg ttaataggta caaaaaata caaagaataa 1338
 ataagaccta atatttgata gcacaacagt gtgactactg tcaataatca tttaattgta 1398
 catttaaaaa taactataat tgcattgttt gtaacacaaa agataaatgc ttgaggagaa 1458
 aaaaaaaaaa aaaa 1472

<210> 14

<211> 135

<212> PRT

<213> Homo sapiens

<400> 14

Met Val Asp Ile Leu Gly Glu Arg His Leu Val Thr Cys Lys Gly Ala
 1 5 10 15

Thr Val Glu Ala Glu Ala Ala Leu Gln Asn Lys Val Val Ala Leu Tyr
 20 25 30

Phe Ala Ala Ala Arg Cys Ala Pro Ser Arg Asp Phe Thr Pro Leu Leu
 35 40 45

New Seq RdCVF.ST25.txt

Cys Asp Phe Tyr Thr Ala Leu Val Ala Glu Ala Arg Arg Pro Ala Pro
50 55 60

Phe Glu Val Val Phe Val Ser Ala Asp Gly Ser Cys Gln Glu Met Leu
65 70 75 80

Asp Phe Met Arg Glu Leu His Gly Ala Trp Leu Ala Leu Pro Phe His
85 90 95

Asp Pro Tyr Arg Gln Arg Ser Leu Ala Leu Leu Pro Arg Leu Glu Cys
100 105 110

Ser Gly Val Ile Leu Ala His Cys Asn Leu Cys Leu Leu Gly Ser Ser
115 120 125

Asp Ser Leu Ala Leu Ala Ser
130 135

<210> 15

<211> 702

<212> DNA

<213> Mus musculus

<400> 15

atcggatcct ctctgggtcc ccagctcctt gcatactgct accatggcat ctctcttctc 60
tggacgcatac ttgatcagga acaacagcga ccaggatgaa gtggagacag aggcagagct 120
gagccgtagg ttagagaatc gtctgggtgtt gctgttcttc ggcgccggcg cctgtcccca 180
gtgccaggcc ttgccccagt cctcaaagac ttcttcgtgc ggctcactga cgagttctac 240
gtgctgcggg cagcacagct ggccctggtc tatgtgtccc aggaccctac agaggagcaa 300
caggacctct tcctcaggga catgcctgaa aaatggctct tcctgccgtc ccactgatga 360
actgaggagg tgaggcccca gggaagacca gggagggtt cctggagaag gcatttccct 420
ggaggtttac tgtcctggta ctacttgtgc actaaagagg tattcctcca caccaaccac 480
aggcgacaac aacacacaag aggtgtccca tccgctcttc catcacagcc cactgacgcc 540
agacagcatc gcgacgctca cggctcagaa aaacacaggt agtctcacag gcctgccatc 600
ctaatactgg ccaccctgag cacaagagcg atggctacaa gcctcaaggc tagaatctaa 660
aaccacgagg tggggaccgt aggccccact ccccgaggagc gc 702

<210> 16

<211> 387

<212> DNA

New Seq RdCVF.ST25.txt

<213> Rattus norvegicus

<400> 16
cggccgctta attaagacgg atccccgact acgtagtcgg gaattcggca cgaggggccc 60
catcttgatc aggaacaaca gcgaccagga tgaagtggag acagaggcag agctgagccg 120
ccggttagag aatcgtcttg tgctactgtt cttcgggtgct ggggcctgtc cccagtgcc 180
ggccttcgcc ccagtcctca aagacttctt cgtgcggctc actgatgagt tctacgtgct 240
acgggcagca cagctggccc tggcttatgt gtcccaggac cctacagagg agcaacagga 300
cctgttcctc cgggacatgc ctgaaaagtg gctcttcctg ccgttccatg atgacctgag 360
gagagacctc gggcgccagt tctccgt 387

<210> 17

<211> 759

<212> DNA

<213> Mus musculus

<400> 17
cagctccttg catactgcta ccatggcatc tctcttctct ggacgcatct tgatcaggaa 60
caacagcgac caggatgaag tggagacaga ggcagagctg agccgtaggt tagagaatcg 120
tctggtgtgc tgttcttcgg cgccggcgcc tgtccccagt gccaggcctt gccccagtcc 180
tcaaagactt cttcgtgcgg ctactgacg agttctacgt gctgcgggca gcacagctgg 240
ccctggtcta tgtgtcccag gaccctacag aggagcaaca ggacctcttc ctcagggaca 300
tgcctgaaaa atggctcttc ctgccgttcc atgatgaact gaggagggac ctcgggcgcc 360
agttctctgt ccgtcaactg ccagcggttg tggacttaa gcctggtggg gacgtgctga 420
caagcgacgc cacggaggag atccagcgtc tgggacccgc ctgctttgcc aactggcagg 480
aggccgcaga gctcctggac cgcagcttcc tgcaaccgga ggatttggat gagcctgcgc 540
ggcgagcat caccgagcct ctgcgccgtc gcaagtaccg agtagaccgg gatgtcggcg 600
ggagcggggc gaaacggcgc gactctggtg aaccccaggg ggacgcgggt acaagggcgg 660
agctctggtg actcccaggg taggagtggg gaccggagct ctggtgacac caaagtaccg 720
gtgcacgacc gaggttgatg accctcccga aggaaccgg 759

<210> 18

<211> 443

<212> DNA

<213> Rattus norvegicus

New Seq RdCVF.ST25.txt

<220>

<221> misc_feature

<222> (46)..(46)

<223> any nucleotide

<400> 18

acgaggtcaa ccttggctac acagggagtc tgaggacagc atgggntaca agaaaccctc	60
tctcaaaacc aaacaaggcc tggcagtact agtgcacttg ggaggcagag gaacaacagc	120
gaccaggatg aagtggagac agaggcagag ctgagccgcc ggtagagaa tcgtcttgtg	180
ctactgttct tcggtgctgg ggcctgtccc cagtgccagg ccttcgcccc agtcctcaaa	240
gacttcttcg tgcggctcac tgatgagttc tacgtgctac gggcagcaca gctggccctg	300
gtctatgtgt cccaggaccc tacagaggag caacaggacc tgttcctccg ggacatgcct	360
gaaaagtggc tcttcctgcc gttccatgat gacctgagga gtaataaaaa ttagaggttg	420
tggctcaaaa aaaaaaaaaa aaa	443

<210> 19

<211> 889

<212> DNA

<213> Homo sapiens

<400> 19

acgccgcgct gtccccagca cccaacccag gttaccatgg cctccctggt ctctggccgc	60
atcctgatcc gcaacaatag cgaccaggac gagctggata cggaggctga ggtcagtcgc	120
aggctggaga accggctggt gctgctgttc tttggtgctg gggcttgtcc acagtgccag	180
gccttcgtgc ccatcctcaa ggacttcttc gtgcggctca cagatgagtt ctatgtactg	240
cgggcggctc agctggccct ggtgtacgtg tcccaggact ccacggagga gcagcaggac	300
ctgttcctca aggacatgcc aaagaaatgg cttttcctgc cctttgagga tgatctgagg	360
agggacctcg ggcgccagtt ctcagtggag cgcctgccgg cggtcgtggt gctcaagccg	420
gacggggacg tgctcactcg cgacggcgcc gacgagatcc agcgcctggg caccgcctgc	480
ttcgccaact ggcaggaggc ggccgaggtg ctggaccgca acttccagct gccagaggac	540
ctggaggacc aggagccacg gagcctcacc gagtgcctgc gccgccacaa gtaccgcgtg	600
gaaaaggcgg cgcgaggcgg cgcgaccggg gggaggggct ggggacggag gccggggccc	660
ggggggctgt actgaccgct ggggtggagca gagggagggg gattggtgga agaacaacaa	720
ccacacgcag ccagcaccag gtatccccgac taggggagac agggcgaaga cctgacccaa	780
agcacaacca ccggggacac taaacgactc aactcaatcc tgtgggcacc aggacaccgc	840

New Seq RdCVF.ST25.txt

aaaaaaaaac aaaaaaagca aaatgcaaaa aaagacagga catacgacg 889

<210> 20

<211> 348

<212> DNA

<213> Homo sapiens

<400> 20

tgtcctcggg tctcaggtgg ctgctgtct ggcctatggt tgacattctg ggcgagcggc 60
acctggtgac ctgtaagggc ggcacggtgg aggccgaggc ggcgctgcag aacaaggtgg 120
tggcactgta cttcgcggcg gcccggtgcg gccgagccg cgacttcacg ccgctgctct 180
gcgacttcta tacggcgctg gtggccgagg cgcggcggcc cgcgcccttc gaagtggctct 240
tcgtgtcagc cgacggcagc tcccaggaga tgctggactt catgcgcgag ctgcatggcg 300
cctggctggc gctgcccttc cacgaccct accggcacca ttgctgtg 348

<210> 21

<211> 340

<212> DNA

<213> Homo sapiens

<400> 21

tgtcctcggg tctcaggtgg ctgctgtct ggcctatggt tgacattctg ggcgagcggc 60
acctggtgac ctgtaagggc ggcacggtgg aggccgaggc ggcgctgcag aacaaggtgg 120
tggcactgta cttcgcggcg gcccggtgcg gccgagccg gattcacgcc gctgctctgc 180
gacttctata cggcgctggt ggccgagcgc ggggccgcgc cttcgaagtg gtcttcgtgt 240
cagccgacgg cagctcccag gagatgctgg acttcatgcg cgagctgatg gcgcctggct 300
ggcgctgcct tccacgaccc ctaccggcac agccggagcc 340

<210> 22

<211> 348

<212> DNA

<213> Homo sapiens

<400> 22

tgtcctcggg tctcaggtgg ctgctgtct ggcctatggt tgacattctg ggcgagcggc 60
acctggtgac ctgtaagggc ggcacggtgg aggccgaggc ggcgctgcag aacaaggtgg 120

New Seq RdCVF.ST25.txt

tggcactgta cttcgcggcg gcccgggtgcg cgccgagccg cgacttcacg ccgctgctct 180
 gcgacttcta tacggcgctg gtggccgagg cgcggcggcc cgcgcccttc gaagtgggtct 240
 tcgtgtcagc cgacggcagc tgccaggaga tgctggactt catgcgcgag ctgcatggcg 300
 cctggctggc gctgcccttc cacgaaccct accggcaacg gagtctcg 348

<210> 23
 <211> 350
 <212> DNA
 <213> Homo sapiens

<400> 23
 tgtcctcggg tctcaggtgg ctgcgtgtct gcgccatggt tgacattctg ggcgagcggc 60
 acctggtgac ctgtaagggc gcgacggtgg aggccgaggc ggcgctgcag aacaaggtgg 120
 tggcactgta cttcgcggcg gcccgggtgcg cgccgagccg cgacttcacg ccgctgctct 180
 gcgacttcta tacggcgctg gtggccgagg cgcggcggcc cgcgcccttc gaagtgggtct 240
 tcgtgtcagc cgacggcagc tgccaggaga tgcttggact tcatgcgcga gctgcattgc 300
 gcctggcttg gcgctgccct tccacgaccc ctaccggcaa cggagtctcg 350

<210> 24
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer
 <400> 24
 tctatgtgtc ccaggaccct acag 24

<210> 25
 <211> 26
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer
 <400> 25
 tttatgcaca agtagtacca ggacag 26

New Seq RdCVF.ST25.txt

<210> 26

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 26

cgggatccat ggcattcttc ttctctggac gc

32

<210> 27

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 27

ggaattctca cctcctcagt tcatcatgga a

31

<210> 28

<211> 50

<212> DNA

<213> Artificial Sequence

<220>

<223> Adaptor

<400> 28

tgttaccaat ctgaagtggg agcggccgac aatttttttt tttttttttt

50

<210> 29

<211> 14

<212> DNA

<213> Artificial Sequence

<220>

New Seq RdCVF.ST25.txt

<223> Adaptor

<400> 29
aattcggcac gagg

14

<210> 30

<211> 10

<212> DNA

<213> Artificial Sequence

<220>

<223> Adaptor

<400> 30
cctcgtgccg

10

<210> 31

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 31
gtaatacgac tcactatagg gc

22

<210> 32

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 32
ctattacgtc aagcctgtag cc

22

<210> 33

<211> 22

<212> DNA

New Seq RdCVF.ST25.txt

<213> Artificial Sequence

<220>

<223> Primer

<400> 33
catcctcatc tttcttccct tc

22

<210> 34

<211> 22

<212> DNA

<213> Artificial

<400> 34
gccagcgttt tctgcctttt ac

22

<210> 35

<211> 22

<212> DNA

<213> Artificial

<400> 35
aagccctgcc tgctctaaca tc

22

31494-12

CLAIMS:

1. An isolated retinoprotective polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2 or SEQ ID NO:4, or an amino acid sequence at least 99% identical thereto.
- 5 2. An isolated retinoprotective polypeptide consisting of an amino acid sequence as set forth in SEQ ID NO:2 or SEQ ID NO:4, or an amino acid sequence at least 99% identical thereto.
3. A vertebrate host cell which can be propagated in vitro and which is capable upon growth in culture of producing the polypeptide according to claim 1,
10 wherein said cell is transformed with at least one transcriptional control sequence, wherein said one or more transcriptional control sequences control transcription of DNA encoding a polypeptide according to claim 1.
4. The vertebrate host cell according to claim 3 wherein said one or more transcriptional control DNA sequences are non-human transcriptional control
15 sequences.
5. A method for the diagnosis of retinal dystrophy in a mammal which comprises: detecting the decreased transcription of messenger RNA transcribed from a retinoprotective-polypeptide-encoding-polynucleotide comprising a nucleic acid sequence as set forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7,
20 SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, a coding region thereof, or a fragment thereof which encodes a polypeptide having retinoprotective activity, in the eye of a mammal, wherein said decreased transcription is diagnostic of said mammal's suffering from retinal dystrophy.
6. A method for the diagnosis of retinal dystrophy in a mammal which
25 comprises: detecting the amount of a retinoprotective polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6,

31494-12

SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, or SEQ ID NO:14, or a fragment thereof having retinoprotective activity, in rod cells of a mammal, wherein the presence of an decreased amount of said polypeptide or fragments thereof, relative to the amount of said polypeptide or fragments thereof in normal eye tissue, is
 5 diagnostic of said mammal's suffering from retinal dystrophy.

7. The method of claim 6, wherein said detecting step comprises contacting said tissue with an antibody which specifically binds to a retinoprotective polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, or
 10 SEQ ID NO:14, or a fragment thereof having retinoprotective activity, and detecting specific binding of said antibody with a polypeptide in said tissue, wherein detection of specific binding to a polypeptide indicates the presence of said polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, or SEQ ID NO:14, or a
 15 fragment thereof having the same activity thereof.

8. A purified antibody or a fragment thereof which specifically binds to a retinoprotective polypeptide that consists of an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 or SEQ ID NO:14, or a fragment thereof having retinoprotective
 20 activity.

9. The antibody fragment according to claim 8 which is an Fab or F(ab')₂ fragment.

10. The antibody according to claim 8 which is a polyclonal antibody.

11. The antibody according to claim 8, which is a monoclonal antibody.

31494-12

12. A kit for the diagnosis of retinal dystrophy in a human comprising a means for collecting a sample and an antibody according to any one of claims 8-11.

13. A method for producing a retinoprotective polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6,
 5 SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, or SEQ ID NO:14, or a fragment thereof having retinoprotective activity, comprising: culturing a host cell having incorporated therein an expression vector containing an exogenously-derived polynucleotide encoding said polypeptide under conditions sufficient for expression of said polypeptide in the host cell, thereby causing the production of an expressed
 10 polypeptide; and recovering the polypeptide produced by said cell.

14. Use, for preventing or protecting photoreceptors in retinal cells from degeneration, of a vector comprising a nucleic acid sequence encoding a retinoprotective polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10,
 15 SEQ ID NO:12, or SEQ ID NO:14, or a fragment thereof having retinoprotective activity, operably linked to a transcription control sequence.

15. The use of claim 14 wherein the vector is derived from an adenovirus or an adeno-associated virus.

16. A pharmaceutical composition comprising a therapeutically effective
 20 amount of a retinoprotective polypeptide comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, or a fragment thereof having retinoprotective activity, and a pharmaceutically acceptable carrier.

17. A pharmaceutical composition comprising a therapeutically effective
 25 amount of a retinoprotective nucleic acid selected from the group of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11,

31494-12

SEQ ID NO:13 the coding region thereof, and a fragment thereof which encodes a polypeptide having retinoprotective activity, and a pharmaceutically acceptable carrier.

18. Use, for the treatment of retinal dystrophy in a subject in need thereof,
5 of a polypeptide comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6,
SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, or a fragment thereof
having retinoprotective activity, and a pharmaceutically acceptable carrier.

19. The use according to claim 18, for the treatment of retinitis pigmentosa,
age-related macular degeneration, Bardet-Biedel syndrome, Bassen-Kornzweig
10 syndrome, best disease, choroidema, gyrate atrophy, congenital amourosis, refsun
syndrome, stargardt disease or Usher syndrome.

20. The use according to claim 18, which decreases the degeneration of
photoreceptors in the retinal cells of the subject.

21. The use according to claim 20, wherein the retinal dystrophy is retinitus
15 pigmentosa, age-related macular degeneration, Bardet-Biedel syndrome,
Bassen-Kornzweig syndrome, best disease, choroidema, gyrate atrophy, congenital
amourosis, refsun syndrome, stargardt disease or Usher syndrome.

22. The use of claim 18, wherein the subject is a human.

23. A method of enhancing cone cell survival in vitro comprising contacting
20 the cell with a polypeptide comprising an amino acid sequence as set forth in
SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10,
SEQ ID NO:12, SEQ ID NO:14 or a fragment thereof having retinoprotective activity.

24. The method of claim 23, wherein the polypeptide is produced
synthetically or by recombinant DNA technology.

31494-12

25. Use, in the manufacture of a medicament for preventing or protecting photoreceptors in retinal cells from degeneration, of a vector comprising a nucleic acid sequence encoding a retinoprotective polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, or SEQ ID NO:14, or a fragment thereof having retinoprotective activity, operably linked to a transcription control sequence.
26. The use of claim 25 wherein the vector is derived from an adenovirus or an adeno-associated virus.
27. A vector comprising a nucleic acid sequence encoding a retinoprotective polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, or SEQ ID NO:14, or a fragment thereof having retinoprotective activity, operably linked to a transcription control sequence, for use in preventing or protecting photoreceptors in retinal cells from degeneration.
28. The vector according to claim 27, wherein the vector is derived from an adenovirus or an adeno-associated virus.
29. Use, in the manufacture of a medicament for the treatment of retinal dystrophy in a subject in need thereof, of a polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, or a fragment thereof having retinoprotective activity, and a pharmaceutically acceptable carrier.
30. The use according to claim 29, wherein the retinal dystrophy is retinitis pigmentosa, age-related macular degeneration, Bardet-Biedel syndrome, Bassen-Kornzweig syndrome, best disease, choroidema, gyrate atrophy, congenital amaurosis, refsum syndrome, stargardt disease or Usher syndrome.

31494-12

31. The use according to claim 29, wherein the medicament decreases the degeneration of photoreceptors in the retinal cells of the subject.

32. The use according to claim 31, wherein the retinal dystrophy is retinitis pigmentosa, age-related macular degeneration, Bardet-Biedel syndrome,

5 Bassen-Kornzweig syndrome, best disease, choroidema, gyrate atrophy, congenital amourosis, refsun syndrome, stargardt disease or Usher syndrome.

33. The use of claim 29, wherein the subject is a human.

34. A polypeptide comprising an amino acid sequence as set forth in

SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10,

10 SEQ ID NO:12, SEQ ID NO:14, or a fragment thereof having retinoprotective activity, for use in the treatment of retinal dystrophy in a subject in need thereof.

35. The polypeptide according to claim 34, for the treatment of retinitis

pigmentosa, age-related macular degeneration, Bardet-Biedel syndrome,

Bassen-Kornzweig syndrome, best disease, choroidema, gyrate atrophy, congenital

15 amourosis, refsun syndrome, stargardt disease or Usher syndrome.

36. The polypeptide according to claim 34, which decreases the

degeneration of photoreceptors in the retinal cells of the subject.

37. The polypeptide according to claim 36, wherein the retinal dystrophy is

retinitis pigmentosa, age-related macular degeneration, Bardet-Biedel syndrome,

20 Bassen-Kornzweig syndrome, best disease, choroidema, gyrate atrophy, congenital amourosis, refsun syndrome, stargardt disease or Usher syndrome.

38. The polypeptide according to claim 34, wherein the subject is a human.

39. A nucleic acid comprising a nucleic acid sequence as set forth in

SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9,

25 SEQ ID NO:11, SEQ ID NO:13, the coding region thereof, or a fragment thereof

31494-12

which encodes a polypeptide having retinoprotective activity, for use in the treatment of retinal dystrophy in a subject in need thereof.

40. A nucleic acid molecule comprising a nucleotide sequence that encodes a polypeptide as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, for use in the treatment of retinal dystrophy in a subject in need thereof.
- 5

Figure 1:**Mouse RdCVF1 cDNA: SEQ ID NO:1**

```

1 ATCGGATCCCTCTCTGGGTCCCCAGCTCCTTGCATACTGCTACCATGGCA 50
51 TCTCTCTTCTCTGGACGCATCTTGATCAGGAACAACAGCGACCAGGATGA 100
101 AGTGGAGACAGAGGCAGAGCTGAGCCGTAGGTTAGAGAATCGTCTGGTGT 150
151 TGCTGTTCTTCGGCGCCGGCGCCTGTCCCCAGTGCCAGGCCTTTGCCCCA 200
201 GTCCTCAAAGACTTCTTCGTGCGGCTCACTGACGAGTTCTACGTGCTGCG 250
251 GGCAGCACAGCTGGCCCTGGTCTATGTGTCCCAGGACCCTACAGAGGAGC 300
301 AACAGGACCTCTTCCTCAGGGACATGCCTGAAAAATGGCTCTTCCTGCCG 350
351 TTCCATGATGAACTGAGGAGGTGAGGCCCCAGGGAAGACCAGGGAGGGCT 400
401 TCCTGGAGAAGGCATTTCCCTGGAGGTTTACTGTCCTGGTACTACTTGTG 450
451 CATAAAGAGGTATTCCTC 468

```

Polypeptide : ATG to TGA: SEQ ID NO:2

```

1 MASLFSGRIL IRNNSDQDEV ETEAELSRRL ENRLVLLFFG AGACPQCQAF
51 APVLKDFEVR LTDEFYVLRA AQLALVYVSQ DPTEEQQDLF LRDMPEKWLF
101 LPFHDELRR

```

Figure 2:**Mouse RdCVF1L cDNA: SEQ ID NO:3**

```

1 CCCCAGCTCC TTGCATACTG CTACCATGGC ATCTCTCTTC TCTGGACGCA
51 TCTTGATCAG GAACAACAGC GACCAGGATG AAGTGGAGAC AGAGGCAGAG
101 CTGAGCCGTA GGTTAGAGAA TCGTCTGGTG TTGCTGTTCT TCGGCGCCGG
151 CGCCTGTCCC CAGTGCCAGG CCTTTGCCCC AGTCCTCAA GACTTCTTCG
201 TGCGGCTCAC TGACGAGTTC TACGTGCTGC GGCAGCACA GCTGGCCCTG
251 GTCTATGTGT CCCAGGACCC TACAGAGGAG CAACAGGACC TCTTCCTCAG
301 GGACATGCCT GAAAAATGGC TCTTCCTGCC GTTCCATGAT GAACTGAGGA
351 GGGACCTCGG GCGCCAGTTC TCTGTCCGTC AACTGCCAGC GGTGTGTGTA
401 CTTAAGCCTG GTGGGGACGT GCTGACAAGC GACGCCACGG AGGAGATCCA
451 GCGTCTGGGA CCCGCCTGCT TTGCCAACTG GCAGGAGGCC GCAGAGCTCC
501 TGGACCGCAG CTTCCTGCAA CCGGAGGATT TGGATGAGCC TGC GCGGCGC
551 AGCATCACCG AGCCTCTGCG CCGTCGCAAG TACCGAGTAG ACCGGGATGT
601 CGGCGGGAGC GGGGCGAAAC GGCGCGACTC TGGTGAACCC CAGGGGGACG
651 CGGGTACAAG GGCGGAGCTC TGGTGACTCC CAGGGTAGGA GTGGGGACCG
701 GAGCTCTGGT GACACCAAAG TACCGGTGCA CGACCGAGGT TGATGACCCT
751 CCCGAAGGAA CCGG

```

Polypeptide : ATG to TGA: SEQ ID NO:4

```

1 MASLFSGRIL IRNNSDQDEV ETEAELSRRL ENRLVLLFFG AGACPQCQAF
51 APVLKDFEVR LTDEFYVLRA AQLALVYVSQ DPTEEQQDLF LRDMPEKWLF
101 LPFHDELRRD LGRQFSVRQL PAVVVLKPGG DVLTSDATEE IQLGPACFA
151 NWQEAAELLD RSFLQPEDLD EPARRSITEP LRRRKYRVDR DVGGSGAKRR
201 DSGEPQGDAG TRAEWL

```

Figure 3:**Human RdCVF1 cDNA: SEQ ID NO:5**

```

1   CCCAGCACCC AACCCAGGTT ACCATGGCCT CCCTGTTCTC TGGCCGCATC
51  CTGATCCGCA ACAATAGCGA CCAGGACGAG CTGGATACGG AGGCTGAGGT
101 CAGTCGCAGG CTGGAGAACC GGCTGGTGCT GCTGTTCTTT GGTGCTGGGG
151 CTTGTCCACA GTGCCAGGCC TTCGTGCCCC TCCTCAAGGA CTTCTTCGTG
201 CGGCTCACAG ATGAGTTCTA TGTACTGCGG GCGGCTCAGC TGGCCCTGGT
251 GTACGTGTCC CAGGACTCCA CGGAGGAGCA GCAGGACCTG TTCCTCAAGG
301 ACATGCCAAA GAAATGGCTT TTCCTGCCCT TTGAGGATGA TCTGAGGAGG
351 TGA

```

Polypeptide : ATG to TGA: SEQ ID NO:6

```

1   MASLFSGRIL IRNNSDQDEL DTEAEVSRRL ENRLVLLFFG AGACPQCQAF
51  VPILKDFEVR LTDEFYVLRA AQLALVYVSQ DSTEEQQDLF LKDMPKKWLF
101 LPFEDDLRR

```

Figure 4:**Human RdCVF1L cDNA: SEQ ID NO:7**

```

1   CCGGGGACCA CACGCCGCGC TGTCCCCAGC ACCCAACCCA GGTTACCATG
51  GCCTCCCTGT TCTCTGGCCG CATCCTGATC CGCAACAATA GCGACCAGGA
101 CGAGCTGGAT ACGGAGGCTG AGGTCAGTCG CAGGCTGGAG AACCGGCTGG
151 TGCTGCTGTT CTTTGGTGCT GGGGCTTGTC CACAGTGCCA GGCCTTCGTG
201 CCCATCCTCA AGGACTTCTT CGTGCGGCTC ACAGATGAGT TCTATGTACT
251 GCGGGCGGCT CAGCTGGCCC TGGTGTACGT GTCCCAGGAC TCCACGGAGG
301 AGCAGCAGGA CCTGTTCTCT AAGGACATGC CAAAGAAATG GCTTTTCCTG
351 CCCTTTGAGG ATGATCTGAG GAGGGACCTC GGGCGCCAGT TCTCAGTGGA
401 GCGCCTGCCG GCGGTCGTGG TGCTCAAGCC GGACGGGGAC GTGCTCACTC
451 GCGACGGCGC CGACGAGATC CAGCGCCTGG GCACCGCCTG CTTGCGCAAC
501 TGGCAGGAGG CGGCCGAGGT GCTGGACCGC AACTTCCAGC TGCCAGAGGA
551 CCTGGAGGAC CAGGAGCCAC GGAGCCTCAC CGAGTGCCTG CGCCGCCACA
601 AGTACCGCGT GGAAAAGGCG GCGCGAGGCG GGCGCGACCC CGGGGGAGGG
651 GGTGGGGAGG AGGGCGGGGC CGGGGGGCTG TTCTGA

```

Polypeptide : ATG to TGA: SEQ ID NO:8

```

1   MASLFSGRIL IRNNSDQDEL DTEAEVSRRL ENRLVLLFFG AGACPQCQAF
51  VPILKDFEVR LTDEFYVLRA AQLALVYVSQ DSTEEQQDLF LKDMPKKWLF
101 LPFEDDLRRD LGRQFSVERL PAVVVLKPDG DVLTRDGADE IQLRGTAFA
151 NWQEAAEVLD RNFQLPEDLE DQEPRSLTEC LRRHKYRVEK AARGGRDPGG
201 GGEEGGAGG LF

```

Figure 5:**Mouse RdCVF2 cDNA: SEQ ID NO:9**

```

1   ATAAAATAGA GGGTGGGAGA GGTGATGGC GTGGCTCTGC TTTTGGTGC
51  GGGGCACCCA GCTGTCATCG CTGCTGTCGC AGCTTCTGGA GTGGCCACTG

101 TGCTCTCTCC TCCCTTCGGC TCAAGGTGAG CTGTTCCAGC AGAAGGCGGG
151 GCTGAGAGGC GCCTAGTGCT GCGGGAGGCT CAGTGTTCATC TTCCAGCTAA
201 CAGGTGGCCG TGCAGCCAG GGCTCGTCTC TCCACTGTGT CCTCTTCACG
251 CCGAGCTCGT GGCGATGGTG GACGTGCTGG GCGGGCGGCG CCTGGTGACC
301 CGGGAGGGCA CCGTGGTGGA GGCCGAGGTG GCGCTGCAGA ACAAGGTGGT
351 AGCTTTGTAC TTTGCGGCGG GCCGGTGCTC GCCCAGCCGC GACTTCACGC
401 CGCTGCTCTG CGACTTCTAC ACGGAGCTGG TGAGCGAGGC GCGGCGGCCC
451 GCTCCCTTCG AGGTGGTTTT CGTGTGCGCA GACGGCAGTG CGGAGGAGAT
501 GTTGGACTTC ATGCGCGAGC TGCACGGCTC CTGGCTGGCA TTGCCCTTCC
551 ACGACCCCTA CCGGCAGTGA GTGGGGACCC AGGGGTCATG GGGCTGGCGC

```

Polypeptide : ATG to TGA: SEQ ID NO:10

```

1   MVDVLGGRRL VTREGTVVEA EVALQNKVVA LYFAAGRCSP SRDFTPLLCD
51  FYTELVSEAR RPAPFEVVFV SADGSAEEML DFMRELHGSW LALPFHDPYR
101 Q

```

Figure 6:**Mouse RdCVF2L cDNA: SEQ ID NO:11**

```

1   TTGACTCTGG TGGGTAGAGA GGGTTTGCAA GGCAGGATAA AATAGAGGGT
51  GGGAGAGGTT GATGGCGTGG CTCTGCTTTT TGGTGCGGGG CACCAGCTGT
101 CATCGCTGCT GTCGCAGCTT CTGGAGTGGC CACTGTGCTC TCTCCTCCCT
151 TCGGCTCAAG GTGAGCTGTT CCAGCAGAAG GCGGGGCTGA GAGGCGCCTA
201 GTGCTGCGGG AGGCTCAGTG TCATCTTCCA GCTAACAGGT GGCCGTGCAG
251 CCCAGGGCTC GTCTCTCCAC TGTGTCCTCT TCACGCCGAG CTCGTGGCGA
301 TGGTGGACGT GCTGGGCGGG CGGCGCCTGG TGACCCGGGA GGGCACGGTG
351 GTGGAGGCCG AGGTGGCGCT GCAGAACAAG GTGGTAGCTT TGTACTTTGC
401 GGCGGGCCGG TGCTCGCCCA GCCGCGACTT CACGCCGCTG CTCTGCGACT
451 TCTACACGGA GCTGGTGAGC GAGGCGCGGC GGCCCGCTCC CTTCGAGGTG
501 GTTTTCGTGT CGGCAGACGG CAGTGCGGAG GAGATGTTGG ACTTCATGCG
551 CGAGCTGCAC GGCTCCTGGC TGGCATTGCC CTTCCACGAC CCCTACCGGC
601 ATGAACTGAA GAAGAGGTAC GAAATCACCG CCATCCCCAA GCTGGTGGTC
651 ATCAAGCAGA ACGGAGCTGT CATCACCAAC AAAGGGCGGA AGCAGATCCG
701 AGAGCGCGGG CTAGCTTGCT TTCAGAACTG GGTGGAAGCA GCCGATGTTT
751 TCCAAAACCTT CTCGGGGTGA CCAGGGCAGT TGCTGGAAGT TCAGGGCAAC
801 TATCTTCAAA AAGGGCTTAG CTGGTTCCCT TCTCTGCTGA GGAATGTCAT
851 TGTAGAGTCA CCATGCTGTG ACAGAGAGCA TAAACTGCTC AGGAAAGAAC
901 TACGTCTGCC CCCTGTGGGT CCTAGAGCTC CGTTGAATGT TTATTTCTTA
951 CACCTTTCTC CACCGGTGCC TAGGATCCAG GACACATCAG CCACGAGTTA
1001 ACAGAACTCT ATGCAAGATG CTCTTTCCTA CAGGAAATTT CTTTGATAAA
1051 TTGACCTATG GAGGTGATAC ATTTTCTGAT GACATTTTGT TGATGCTTTG
1101 GTAAACGTAT TTATTACTCG GGTTTGTAGA CTGTGTAATT TAATAAACCA
1151 AACTCACAC TTTG

```

Polypeptide : ATG to TGA: SEQ ID NO:12

```

1   MVDVLGGRRL VTREGTVVEA EVALQNKVVA LYFAAGRCSP SRDFTPLLCD

```

```

51  FYTELVSEAR RPAPFEVVFV SADGSAEEML DFMRELHGSW LALPFHDPYR
101 HELKKRYEIT AIPKLVVIKQ NGAVITNKGR KQIRERGLAC FQNWVEAADV
151 FQNFSG

```

Figure 7:**Human RdCVF2 cDNA: SEQ ID NO:13**

```

1  GTGTGGGCGG GCGCAGTTG GGGGAGGGTG CAGAGACCTG AGGGCTTGAG
51  GTTGCCCTGGC TGGCCCCGCT CCCAGAGGCG GGTGCCGCGC TGTCGCCCAG
101 GTATCTGGGG TCTCTGGTGT CTGAGTGTCT CATTGTCTGGC GCGAACACAA
151 TTGCTCCAGC CACAGGCGAG GCCTGGCCAA GGTGTGGGCG CATCTAGGGC
201 AGGTCTTGAG AGGTCCAGCG CCCGGTGGTG CGGACAGAGG CGGGGCACCG
251 CGGCGCTCGC CGCCGCCTCC CCGCAGGTGA TCATCCTCCT GCAGGTGTCC
301 TCGGGTCTCA GGTGGCTGCG TGTCTGCGCC ATGGTTGACA TTCTGGGCGA
351 GCGGCACCTG GTGACCTGTA AGGGCGCGAC GGTGGAGGCC GAGGCGGCGC
401 TGCAGAACAA GGTGGTGGCA CTGTACTTCG CGGCGGCCCG GTGCGCGCCG
451 AGCCGCGACT TCACGCCGCT GCTCTGCGAC TTCTATACGG CGCTGGTGGC
501 CGAGGCGCGG CGGCCCGCGC CCTTCGAAGT GGTCTTCGTG TCAGCCGACG
551 GCAGCTGCCA GGAGATGCTG GACTTCATGC GCGAGCTGCA TGGCGCCTGG
601 CTGGCGCTGC CCTTCCACGA CCCCTACCGG CAACGGAGTC TCGCTCTGTT
651 GCCCAGGCTG GAGTGCAGTG GCGTGATCTT AGCTCACTGC AACCTTTGCC
701 TCCTGGGTTC AAGTGATTCT CTAGCCTTAG CCTCCTGAGC ATCTGGGACT
751 ACAGCCATTG CTGTGAATTA CGTGAGGGAA AGATATTGAA GAGGAGTTGG
801 AACTCCGAG AGTGCAGCTG TTCTCCCCC GCACCATCCG TGTCTGTCAT
851 TCTGCGAGTC TGTGCTCATT AACAATGTGC TGTGACCATG TGA CTCAGCA
901 ATCCTGCTGC TGGGTATATA CCCGAAAGAA AGGAAAAGGA AGCCAGTATA
951 TTGAAGAGGT ATCTGCACCC CCATGTTTAT TGCAGCACTG TTCACAACAG
1001 CCAAGATTTG GAAGCAACCT AAGTGTCAT CAACAGATGA ATGGATAAAG
1051 AAAACGTGGT ACATATACAC AATGGAGTAC TCTTCAGCCA TTAAAAAAT
1101 GAGATTCTGT CATTTGCAAT AATATAGATG GAAAAGGAGG CCCTTATGTG
1151 AAGTGAAATA AGCCAGGCAC AGAAAGACAA ACATCACATG TTCTCACTTA
1201 TTTGTGGGAT CTAATGATCA AAACAATTGA ACTCTTGAC ATAGAGAGTA
1251 GAAGGTTGGT TACCAGAAGC TGGAAAGGAA AGTGGGGTTG GGAGGAAGGT
1301 GGGAATGGTT AATAGGTACA AAAAAATACA AAGAATAAAT AAGACCTAAT
1351 ATTTGATAGC ACAACAGTGT GACTACTGTC AATAATCATT TAATTGTACA
1401 TTTAAAAATA ACTATAATTG CATTGTTTGT AACACAAAAG ATAAATGCTT
1451 GAGGAGAAAA AAAAAAAAAA AA

```

Polypeptide : ATG to TGA : SEQ ID NO:14

```

1  MVDILGERHL VTCKGATVEA EAALQNKVVA LYFAAARCAP SRDFTPLLCD
51  FYTALVAEAR RPAPFEVVFV SADGSCQEML DFMRELHGAW LALPFHDPYR
101 QRSLLALLPRL ECSGVILAHC NLCLLGSSDS LALAS

```

Figure 8:**Pileup short sequences :**

```

1 50
mRdCVF1 MASLFSGRIL IRNNSDQDEV ETEAELSRRL ENRLVLLFFG AGACPQCQAF
hRdCVF1 MASLFSGRIL IRNNSDQDEL DTEAEVSRRL ENRLVLLFFG AGACPQCQAF
mRdCVF2 MVDVLGGRRL VTREG....T VVEAEVA..L QNKVVALYFA AGRCSPSRDF
hRdCVF2 MVDILGERHL VTCKG....A TVEAEAA..L QNKVVALYFA AARCAPSRDF

```

	51		100
mRdCVF1	APVLKDFEVR LTDEFYVLRA AQLALVYVSQ DPTEEQQDLF LRDMPEKWLF		
hRdCVF1	VPILKDFEVR LTDEFYVLRA AQLALVYVSQ DSTEEQQDLF LKDMPPKKWLF		
mRdCVF2	TPLLCDFYTE LVSE..ARRP APFEVVFVSA DGSAEEMLDF MRELHGSWLA		
hRdCVF2	TPLLCDFYTA LVAE..ARRP APFEVVFVSA DGSCQEMLDF MRELHGAWLA		
	101		143
mRdCVF1	LPFHDELRR		
hRdCVF1	LPFEDDLRR		
mRdCVF2	LPFHDPYRQ		
hRdCVF2	LPFHDPYRQR SLALLPRLEC SGVILAHCNL CLLGSSDSL LAS		

Pileup long sequences :

	1		50
mRdCVF1L	MASLFSGRIL IRNNSDQDEV ETEAELSRRL ENRLVLLFFG AGACPQCQAF		
hRdCVF1L	MASLFSGRIL IRNNSDQDEL DTEAEVSRRL ENRLVLLFFG AGACPQCQAF		
mRdCVF2L	MVDVLGGRRL VTREG....T VVEAEVA..L QNKVVALYFA AGRCSPSRDF		
hRdCVF2	MVDILGERHL VTCKG....A TVEAEAA..L QNKVVALYFA AARCAPSRDF		
	51		100
mRdCVF1L	APVLKDFEVR LTDEFYVLRA AQLALVYVSQ DPTEEQQDLF LRDMPEKWLF		
hRdCVF1L	VPILKDFEVR LTDEFYVLRA AQLALVYVSQ DSTEEQQDLF LKDMPPKKWLF		
mRdCVF2L	TPLLCDFYTE LVSE..ARRP APFEVVFVSA DGSAEEMLDF MRELHGSWLA		
hRdCVF2	TPLLCDFYTA LVAE..ARRP APFEVVFVSA DGSCQEMLDF MRELHGAWLA		
	101		150
mRdCVF1L	LPFHDELRRD ...LGRQFSV RQLPAVVVLK PGGDVLTSDA TEEIQRLGPA		
hRdCVF1L	LPFEDDLRRD ...LGRQFSV ERLPAVVVLK PDGDVLTRDG ADEIQRLGTA		
mRdCVF2L	LPFHDPYRHE ...LKKRYEI TAIPKLVIK QNGAVITNKG RKQIRERGLA		
hRdCVF2	LPFHDPYRQR SLALLPRLEC SGV...ILAH CNLCLLGSSD SLALAS		
	151		200
mRdCVF1L	CFANWQEAAE LLDRSFLQPE DLDEPARRSI TEPLRRRKYR VDRDVGGSGA		
hRdCVF1L	CFANWQEAAE VLDRNFQLPE DLEDQEPRSL TECLRRHKYR VEKAARGG..		
mRdCVF2L	CFQNWVEAAD VFQNFSG		
hRdCVF2			
	201	219	
mRdCVF1L	KRRDSGEPQG DAGTRAELW		
hRdCVF1L	..RDPGGGGG EEGGAGGLF		
mRdCVF2L			
hRdCVF2			

Figure 9:

SEQ ID NO: 26: 5' cgggatccATGGCATCTCTCTTCTCTGGACGC 3'

SEQ ID NO: 27: 5' ggaattcTCACCTCCTCAGTTCATCATGGAA 3'

Figure 10:

	1					50
RdCVF1	MASLFSGRIL	IRNNSDQDEV	ETEAELSRRL	ENRLVLLFFG	AGACPQCQAF	
RdCVF2	MVDVLGGRRL	V....TREGT	VVEAEVA..L	QNKVVALYFA	AGRCSPSRDF	
	51					100
RdCVF1	APVLKDFVVR	LTDEFYVLRA	AQLALVYVSQ	DPTEEQQDLF	LRDMPEKWLF	
RdCVF2	TPLLCDFYTE	LVSE..ARRP	APFEVVFVSA	DGSAEEMLDF	MRELHGSWLA	
	101					
RdCVF1	LPFHDELRR					
RdCVF2	LPFHDPYRQ					

Figure 11:

bc016199		mouse RdCVF2	
bg707818		human RdCVF2	
	1		50
bc016199	MVDVLGGRRL	VTREGTVVEA	EVALQNKVVA LYFAAGRCSP SRDFTPLLCD
bg707818	MVDILGERHL	VTCKGATVEA	EAALQNKVVA LYFAAARCAP SRDFTPLLCD
	51		100
bc016199	FYTELVSEAR	RPAPFEVVFV	SADGSAEEML DFMRELHGSW LALPFHDPYR
bg707818	FYTALVAEAR	RPAPFEVVFV	SADGSCQEML DFMRELHGAW LALPFHEPYR
	101	120	
bc016199	Q*		
bg707818	Q*SLALLPRL	ECSGVILAH*	

Figure 12:

Name: mRdCVF2	mouse sequence bc016199
Name: be552141	homo sapiens
Name: bi517442	homo sapiens
Name: bg707818	homo sapiens
Name: bi603812	homo sapiens
mRdCVF2	ATGGT GGACGTGCTG
be552141	TGTCCTCGGG TCTCAGGTGG CTGCGTGTCT GCGCCATGGT TGACATTCTG
bi517442	TGTCCTCGGG TCTCAGGTGG CTGCGTGTCT GCGCCATGGT TGACATTCTG
bg707818	TGTCCTCGGG TCTCAGGTGG CTGCGTGTCT GCGCCATGGT TGACATTCTG
bi603812	TGTCCTCGGG TCTCAGGTGG CTGCGTGTCT GCGCCATGGT TGACATTCTG
mRdCVF2	GGCGGGCGGC GCCTGGTGAC CCGGGAGGGC ACGGTGGTGG AGGCCGAGGT
be552141	GGCGAGCGGC ACCTGGTGAC CTGTAAGGGC GCGACGGTGG AGGCCGAGGC
bi517442	GGCGAGCGGC ACCTGGTGAC CTGTAAGGGC GCGACGGTGG AGGCCGAGGC
bg707818	GGCGAGCGGC ACCTGGTGAC CTGTAAGGGC GCGACGGTGG AGGCCGAGGC
bi603812	GGCGAGCGGC ACCTGGTGAC CTGTAAGGGC GCGACGGTGG AGGCCGAGGC
mRdCVF2	GGCGCTGCAG AACAAAGGTGG TAGCTTTGTA CTTTGCGGCG GGCCGGTGCT
be552141	GGCGCTGCAG AACAAAGGTGG TGGCACTGTA CTTTCGCGGCG GCCCGGTGCG
bi517442	GGCGCTGCAG AACAAAGGTGG TGGCACTGTA CTTTCGCGGCG G.CCGGTGCG
bg707818	GGCGCTGCAG AACAAAGGTGG TGGCACTGTA CTTTCGCGGCG GCCCGGTGCG
bi603812	GGCGCTGCAG AACAAAGGTGG TGGCACTGTA CTTTCGCGGCG GCCCGGTGCG

```

mRdCVF2    CGCCCAGCCG CGACTTCACG CCGCTGCTCT GCGACTTCTA CACGGAGCTG
be552141   CGCCGAGCCG CGACTTCACG CCGCTGCTCT GCGACTTCTA TACGGCGCTG
bi517442   CGCCGAGCCG CGA.TTCACG CCGCTGCTCT GCGACTTCTA TACGGCGCTG
bg707818   CGCCGAGCCG CGACTTCACG CCGCTGCTCT GCGACTTCTA TACGGCGCTG
bi603812   CGCCGAGCCG CGACTTCACG CCGCTGCTCT GCGACTTCTA TACGGCGCTG

mRdCVF2    GTGAGCGAGG CGCGGCGGCC CGCTCCCTTC GAGGTGGTTT TCGTGTCGGC
be552141   GTGGCCGAGG CGCGGCGGCC CGCGCCCTTC GAAGTGGTCT TCGTGTCAGC
bi517442   GTGGCCGAG. ...CGCGGGG CCGCGCCTTC GAAGTGGTCT TCGTGTCAGC
bg707818   GTGGCCGAGG CGCGGCGGCC CGCGCCCTTC GAAGTGGTCT TCGTGTCAGC
bi603812   GTGGCCGAGG CGCGGCGGCC CGCGCCCTTC GAAGTGGTCT TCGTGTCAGC

mRdCVF2    AGACGGCAGT GCGGAGGAGA TGT.TGGACT TCATGCGCGA GCTGCACGGC
be552141   CGACGGCAGC TCCCAGGAGA TGC.TGGACT TCATGCGCGA GCTGCATGGC
bi517442   CGACGGCAGC TCCCAGGAGA TGC.TGGACT TCATGCGCGA GCTG.ATGGC
bg707818   CGACGGCAGC TGCCAGGAGA TGC.TGGACT TCATGCGCGA GCTGCATGGC
bi603812   CGACGGCAGC TGCCAGGAGA TGCTTGGACT TCATGCGCGA GCTGCATTGC

mRdCVF2    TCCTGGC.TG GCATTGCCCT TCCACGACCC CTACCGGCAG TGA
be552141   GCCTGGC.TG GCGCTGCCCT TCCACGACCC CTACCGGCAC CATTGCTGTG
bi517442   GCCTGGC.TG GCGCTG.CCT TCCACGACCC CTACCGGCAC AGCCGGAGCC
bg707818   GCCTGGC.TG GCGCTGCCCT TCCACGAACC CTACCGGCAA CGGAGTCTCG
bi603812   GCCTGGCTTG GCGCTGCCCT TCCACGACCC CTACCGGCAA CGGAGTCTCG

```

Figure 13:

pileup

GapWeight: 5
GapLengthWeight: 1

Name: RdCVF1
Name: bg299078
Name: ai716631
Name: bg294111
Name: be108041
Name: bg395178

Rev

```

      1                                     50
RdCVF1  ~~~~~~
bg299078 ~~~~~~
ai716631 ~~~~~~
bg294111 ~~~~~~
be108041 ACGAGGTCAA CCTTGGCTAC ACAGGGAGTC TGAGGACAGC ATGGGNTACA
bg395178 ~~~~~~

      51                                     100
RdCVF1  ~~~~~~
bg299078 CCCCAGCTCC TTGCATACTG CTACCATGGC ATCTCTCTTC TCTGGACGCA
ai716631 TAAGACGGAT CCCCGACTAC GTAGTCGGGA ATTCGGCACG AGGGGCCGCA
bg294111 ~~~~CAGCTCC TTGCATACTG CTACCATGGC ATCTCTCTTC TCTGGACGCA
be108041 AGAAACCCTC TCTCAAAACC AAACAAGGCC TGGCAGTACT AGTGCACTTG
bg395178 TCCCCAGCAC CCAACCCAGG TTACCATGGC CTCCCTGTTC TCTGGCCGCA

      101                                     150
RdCVF1  TCTTGATCAG GAACAACAGC GACCAGGATG AAGTGGAGAC AGAGGCAGAG
bg299078 TCTTGATCAG GAACAACAGC GACCAGGATG AAGTGGAGAC AGAGGCAGAG

```

ai716631	TCTTGATCAG	GAACAACAGC	GACCAGGATG	AAGTGGAGAC	AGAGGCAGAG
bg294111	TCTTGATCAG	GAACAACAGC	GACCAGGATG	AAGTGGAGAC	AGAGGCAGAG
be108041	GGAGGCAGAG	GAACAACAGC	GACCAGGATG	AAGTGGAGAC	AGAGGCAGAG
bg395178	TCCTGATCCG	CAACAATAGC	GACCAGGACG	AGCTGGATAC	GGAGGCTGAG

	151				200
RdCVF1	CTGAGCCGTA	GGTTAGAGAA	TCGTCTGGTG	TTGCTGTTCT	TCGGCGCCGG
bg299078	CTGAGCCGTA	GGTTAGAGAA	TCGTCTGGTG	TTGCTGTTCT	TCGGCGCCGG
ai716631	CTGAGCCGCC	GGTTAGAGAA	TCGTCTTGTG	CTACTGTTCT	TCGGTGCTGG
bg294111	CTGAGCCGTA	GGTTAGAGAA	TCGTCTGGTG	<u>.TGCTGTTCT</u>	TCGGCGCCGG
be108041	CTGAGCCGCC	GGTTAGAGAA	TCGTCTTGTG	<u>CTACTGTTCT</u>	TCGGTGCTGG
bg395178	GTCAGTCGCA	GGCTGGAGAA	CCGGCTGGTG	CTGCTGTTCT	TTGGTGCTGG

	201				250
RdCVF1	CGCCTGTCCC	CAGTGCCAGG	CCTTTGCCCC	AGTCCTCAAA	GACTTCTTCG
bg299078	CGCCTGTCCC	CAGTGCCAGG	<u>CC</u> .TTGCCCC	AGTCCTCAAA	GACTTCTTCG
ai716631	GGCCTGTCCC	CAGTGCCAGG	CCTTCGCCCC	AGTCCTCAAA	GACTTCTTCG
bg294111	CGCCTGTCCC	CAGTGCCAGG	<u>CC</u> .TTGCCCC	AGTCCTCAAA	GACTTCTTCG
be108041	GGCCTGTCCC	CAGTGCCAGG	CCTTCGCCCC	AGTCCTCAAA	GACTTCTTCG
bg395178	GGCTTGTCCA	CAGTGCCAGG	CCTTCGTGCC	CATCCTCAAG	GACTTCTTCG

	251				300
RdCVF1	TGCGGCTCAC	TGACGAGTTC	TACGTGCTGC	GGGCAGCACA	GCTGGCCCTG
bg299078	TGCGGCTCAC	TGACGAGTTC	TACGTGCTGC	GGGCAGCACA	GCTGGCCCTG
ai716631	TGCGGCTCAC	TGATGAGTTC	TACGTGCTAC	GGGCAGCACA	GCTGGCCCTG
bg294111	TGCGGCTCAC	TGACGAGTTC	TACGTGCTGC	GGGCAGCACA	GCTGGCCCTG
be108041	TGCGGCTCAC	TGATGAGTTC	TACGTGCTAC	GGGCAGCACA	GCTGGCCCTG
bg395178	TGCGGCTCAC	AGATGAGTTC	TATGTACTGC	GGGCGGCTCA	GCTGGCCCTG

	301				350
RdCVF1	GTCTATGTGT	CCCAGGACCC	TACAGAGGAG	CAACAGGACC	TCTTCCTCAG
bg299078	GTCTATGTGT	CCCAGGACCC	TACAGAGGAG	CAACAGGACC	TCTTCCTCAG
ai716631	GTCTATGTGT	CCCAGGACCC	TACAGAGGAG	CAACAGGACC	TGTTCCCTCCG
bg294111	GTCTATGTGT	CCCAGGACCC	TACAGAGGAG	CAACAGGACC	TCTTCCTCAG
be108041	GTCTATGTGT	CCCAGGACCC	TACAGAGGAG	CAACAGGACC	TGTTCCCTCCG
bg395178	GTGTACGTGT	CCCAGGACTC	CACGGAGGAG	CAGCAGGACC	TGTTCCCTCAA

	351				400
RdCVF1	GGACATGCCT	GAAAAATGGC	TCTTCCTGCC	GTTCCA.TGA	TGAACTGAGG
bg299078	GGACATGCCT	GAAAAATGGC	TCTTCCTGCC	GTCCCACTGA	TGAACTGAGG
ai716631	GGACATGCCT	GAAAAGTGGC	TCTTCCTGCC	GTTCCA.TGA	TGACCTGAGG
bg294111	GGACATGCCT	GAAAAATGGC	TCTTCCTGCC	GTTCCA.TGA	TGAACTGAGG
be108041	GGACATGCCT	GAAAAGTGGC	TCTTCCTGCC	GTTCCA.TGA	TGACCTGAGG
bg395178	GGACATGCCA	AAGAAATGGC	TTTTCCTGCC	CTTTGA.GGA	TGATCTGAGG

	401				450
RdCVF1	AGGTGA~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg299078	AGGTGAGGCC	CCAGGGAAGA	CCAGGGAGGG	CTTCCTGGAG	AAGGCATTTC
ai716631	AGAGACCTCG	GGCGCCAGTT	CTCCGT~~~~	~~~~~	~~~~~
bg294111	AGGGACCTCG	GGCGCCAGTT	CTCTGTCCGT	CAACTGCCAG	CGGTTGTGGT
be108041	AGTAATAAAA	ATTAGAGGTT	GTGGCTCAAA	AAAAAAAAAA	AAAA~~~~~
bg395178	AGGGACCTCG	GGCGCCAGTT	CTCAGTGGAG	CGCCTGCCCG	CGGTCGTGGT

	451				500
RdCVF1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg299078	CCTGGAGGTT	TACTGTCCTG	GTACTACTTG	TGCACTAAAG	AGGTATTTCCT
ai716631	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg294111	ACTTAAGCCT	GGTGGGGACG	TGCTGACAAG	CGACGCCACG	GAGGAGATCC
be108041	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg395178	GCTCAAGCCG	GACGGGGACG	TGCTCACTCG	CGACGGCGCC	GACGAGATCC

	501				550
RdCVF1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg299078	CCACACCAAC	CACAGGCGAC	AACAACACAC	AAGAGGTGTC	CCATCCGCTC
ai716631	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg294111	AGCGTCTGGG	ACCCGCCTGC	TTTGCCAAC	GGCAGGAGGC	CGCAGAGCTC
be108041	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg395178	AGCGCCTGGG	CACCGCCTGC	TTCGCCAAC	GGCAGGAGGC	GGCCGAGGTG
	551				600
RdCVF1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg299078	TTCCATCACA	GCCCACTGAC	GCCAGACAGC	ATCGCGACGC	TCACGGCTCA
ai716631	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg294111	CTGGACCGCA	GCTTCCTGCA	ACCGGAGGAT	TTGGATGAGC	CTGCGCGGCG
be108041	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg395178	CTGGACCGCA	ACTTCCAGCT	GCCAGAGGAC	CTGGAGGACC	AGGAGCCACG
	601				650
RdCVF1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg299078	GAAAAACACA	GGTAGTCTCA	CAGGCCTGCC	ATCCTAATAC	TGGCCACCCT
ai716631	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg294111	CAGCATCACC	GAGCCTCTGC	GCCGTCGCAA	GTACCGAGTA	GACCGGGATG
be108041	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg395178	GAGCCTCACC	GAGTGCCTGC	GCCGCCACAA	GTACCGCGTG	GAAAAGGCGG
	651				700
RdCVF1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg299078	GAGCACAAGA	GCGATGGCTA	CAAGCCTCAA	GGCTAGAATC	TAAAACCACG
ai716631	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg294111	TCGGCGGGAG	CGGGGCGAAA	CGGCGCGACT	CTGGTGAACC	CCAGGGGGAC
be108041	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg395178	CGCGAGGCGG	CGCGACCCGG	GGGAGGGGCT	GGGGACGGAG	GCCGGGGCCC
	701				750
RdCVF1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg299078	AGGTGGGGAC	CGTAGGCCCC	ACTCCCCGGG	AGCGC~~~~~	~~~~~
ai716631	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg294111	GCGGGTACAA	GGGCGGAGCT	CTGGTGACTC	CCAGGGTAGG	AGTGGGGACC
be108041	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg395178	GGGGGGCTGT	ACTGACCGCT	GGGTGGAGCA	GAGGGAGGGG	GATTGGTGGA
	751				800
RdCVF1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg299078	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
ai716631	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg294111	GGAGCTCTGG	TGACACCAAA	GTACCGGTGC	ACGACCGAGG	TTGATGACCC
be108041	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg395178	AGAACAACAA	CCACACGCAG	CCAGCACCAG	GTATCCCGAC	TAGGGGAGAC
	801				850
RdCVF1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg299078	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
ai716631	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg294111	TCCCGAAGGA	ACCGG~~~~~	~~~~~	~~~~~	~~~~~
be108041	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg395178	AGGGCGAAGA	CCTGACCCAA	AGCACAACCA	CCGGGGACAC	TAAACGACTC
	851				900
RdCVF1	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg299078	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
ai716631	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
bg294111	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~

```

be108041 ~~~~~~ ~~~~~~ ~~~~~~ ~~~~~~ ~~~~~~
bg395178  AACTCAATCC TGTGGGCACC AGGACACCGC AAAAAAAAC AAAAAAGCA

          901                               929
RdCVF1    ~~~~~~ ~~~~~~ ~~~~~~
bg299078  ~~~~~~ ~~~~~~ ~~~~~~
ai716631  ~~~~~~ ~~~~~~ ~~~~~~
bg294111  ~~~~~~ ~~~~~~ ~~~~~~
be108041  ~~~~~~ ~~~~~~ ~~~~~~
bg395178  AAATGCAAAA AAAGACAGGA CATACGACG

```

Figure 14:**BG299078**

```

ATCGGATCCTCTCTGGGTCCCCAGCTCCTTGCATACTGCTACCATGGCATCTCTCTTCTC
1  -----+-----+-----+-----+-----+-----+ 60
TAGCCTAGGAGAGACCCAGGGGTCGAGGAACGTATGACGATGGTACCGTAGAGAGAAGAG

TGGACGCATCTTGATCAGGAACAACAGCGACCAGGATGAAGTGGAGACAGAGGCAGAGCT
61 -----+-----+-----+-----+-----+-----+ 120
ACCTGCGTAGAACTAGTCCTTGTTGTCGCTGGTCCCTACTTCACCTCTGTCTCCGTCTCGA

GAGCCGTAGGTTAGAGAATCGTCTGGTGTGCTGTTCTTCGGCGCCGGCGCCTGTCCCCA
121 -----+-----+-----+-----+-----+-----+ 180
CTCGGCATCCAATCTCTTAGCAGACCACAACGACAAGAAGCCGCGGCCGCGACAGGGGT
      +T
GTGCCAGGCCTTGCCCCAGTCCTCAAAGACTTCTTCGTGCGGCTCACTGACGAGTTCTAC
181 -----+-----+-----+-----+-----+-----+ 240
CACGGTCCGGAACGGGGTCAGGAGTTTCTGAAGAAGCACGCCGAGTGACTGCTCAAGATG

GTGCTGCGGGCAGCACAGCTGGCCCTGGTCTATGTGTCCCAGGACCCTACAGAGGAGCAA
241 -----+-----+-----+-----+-----+-----+ 300
CACGACGCCCCTCGTGTGACCGGGACCAGATACACAGGGTCCTGGGATGTCTCCTCGTT
                                removed
CAGGACCTCTTCCTCAGGGACATGCCTGAAAAATGGCTCTTCCTGCCGTCCCAGTATGA
301 -----+-----+-----+-----+-----+-----+ 360
GTCCTGGAGAAGGAGTCCCTGTACGGACTTTTACCAGAGAAGGACGGCAGGGTGACTACT

ACTGAGGAGGTGAGGCCCCAGGGAAGACCAGGGAGGGCTTCCTGGAGAAGGCATTTCCCT
361 -----+-----+-----+-----+-----+-----+ 420
TGACTCCTCCACTCCGGGGTCCCTTCTGGTCCCTCCCGAAGGACCTCTTCGGTAAAGGGA

GGAGGTTTACTGTCCTGGTACTACTTGTGCACTAAAGAGGTATTCCTCCACACCAACCAC
421 -----+-----+-----+-----+-----+-----+ 480
CCTCCAAATGACAGGACCATGATGAACACGTGATTTCTCCATAAGGAGGTGTGGTTGGTG

AGGCGACAACAACACACAAGAGGTGTCCCATCCGCTCTTCATCACAGCCCACTGACGCC
481 -----+-----+-----+-----+-----+-----+ 540
TCCGCTGTTGTTGTGTGTTCTCCACAGGGTAGGCGAGAAGGTAGTGTCGGGTGACTGCGG

AGACAGCATCGCGACGCTCACGGCTCAGAAAAACACAGGTAGTCTCACAGGCCTGCCATC
541 -----+-----+-----+-----+-----+-----+ 600
TCTGTCTGTAGCGCTGCGAGTGCCGAGTCTTTTTGTGTCCATCAGAGTGTCGGGACGGTAG

CTAATACTGGCCACCCTGAGCACAAGAGCGATGGCTACAAGCCTCAAGGCTAGAATCTAA
601 -----+-----+-----+-----+-----+-----+ 660
GATTATGACCGGTGGGACTCGTGTCTCGCTACCGATGTTTCGGAGTTCCGATCTTAGATT

```

AACCACGAGGTGGGGACCGTAGGCCCCACTCCCCGGGAGCGC
 661 -----+-----+-----+-----+--- 702
 TTGGTGCTCCACCCCTGGCATCCGGGGTGAGGGGCCCTCGCG

Figure 15:**BG294111**

CAGCTCCTTGCATACTGCTACCATGGCATCTCTTCTCTGGACGCATCTTGATCAGGAA
 1 -----+-----+-----+-----+-----+-----+ 60
 GTCGAGGAACGTATGACGATGGTACCGTAGAGAGAAGAGACCTGCGTAGAACTAGTCCTT

CAACAGCGACCAGGATGAAGTGGAGACAGAGGCAGAGCTGAGCCGTAGGTTAGAGAATCG
 61 -----+-----+-----+-----+-----+-----+ 120
 GTTGTGCTGGTCCCTACTTCACCTCTGTCTCCGTCTCGACTCGGCATCCAATCTCTTAGC
 +T +T
 TCTGGTGTGCTGTTCTTCGGCGCCGGCGCCTGTCCCCAGTGCCAGGCCTTGCCCCAGTCC
 121 -----+-----+-----+-----+-----+-----+ 180
 AGACCACACGACAAGAAGCCGCGGCCGCGGACAGGGGTCACGGTCCGGAACGGGGTCAGG

TCAAAGACTTCTTCGTGCGGCTCACTGACGAGTTCTACGTGCTGCGGGCAGCACAGCTGG
 181 -----+-----+-----+-----+-----+-----+ 240
 AGTTTCTGAAGAAGCACGCCGAGTGACTGCTCAAGATGCACGACGCCCGTCGTGTCGACC

CCCTGGTCTATGTGTCCCAGGACCTTACAGAGGAGCAACAGGACCTCTTCCTCAGGGACA
 241 -----+-----+-----+-----+-----+-----+ 300
 GGGACCAGATACACAGGGTCTTGGGATGTCTCCTCGTTGTCTGGAGAAGGAGTCCCTGT

TGCCTGAAAAATGGCTCTTCCTGCCGTTCCATGATGAACTGAGGAGGGACCTCGGGCGCC
 301 -----+-----+-----+-----+-----+-----+ 360
 ACGGACTTTTACCGAGAAGGACGGCAAGGTACTACTTGACTCCTCCCTGGAGCCCGCGG

AGTTCTCTGTCCGTCAACTGCCAGCGGTTGTGGTACTTAAGCCTGGTGGGGACGTGCTGA
 361 -----+-----+-----+-----+-----+-----+ 420
 TCAAGAGACAGGCAGTTGACGGTCGCCAACACCATGAATTGGGACCACCCCTGCACGACT

CAAGCGACGCCACGGAGGAGATCCAGCGTCTGGGACCCGCTTGCTTTGCCAACTGGCAGG
 421 -----+-----+-----+-----+-----+-----+ 480
 GTTCGCTGCGGTGCCTCCTCTAGGTCGCAGACCCTGGGCGGACGAAACGGTTGACCGTCC

AGGCCGCAGAGCTCCTGGACCGCAGCTTCCTGCAACCGGAGGATTTGGATGAGCCTGCGC
 481 -----+-----+-----+-----+-----+-----+ 540
 TCCGGCGTCTCGAGGACCTGGCGTCGAAGGACGTTGGCCTCCTAAACCTACTCGGACGCG

GGCGCAGCATCACCGAGCCTCTGCGCCGTCGCAAGTACCGAGTAGACCGGGATGTGCGCG
 541 -----+-----+-----+-----+-----+-----+ 600
 CCGCGTCGTAGTGGCTCGGAGACGCGGCAGCGTTCATGGCTCATCTGGCCCTACAGCCGC

GGAGCGGGGCGAAACGGCGCGACTCTGGTGAACCCAGGGGGACGCGGGTACAAGGGCGG
 601 -----+-----+-----+-----+-----+-----+ 660
 CCTCGCCCCGCTTTGCCGCGCTGAGACCACTTGGGGTCCCCCTGCGCCCATGTTCCCGCC

AGCTCTGGTGACTCCCAGGGTAGGAGTGGGGACCGGAGCTCTGGTGACACCAAAGTACCG
 661 -----+-----+-----+-----+-----+-----+ 720
 TCGAGACCACTGAGGGTCCCATCCTCACCCCTGGCCTCGAGACCACTGTGGTTTCATGGC

GTGCACGACCGAGGTTGATGACCCTCCCGAAGGAACCGG
 721 -----+-----+-----+-----+-----+ 759
 CACGTGCTGGCTCCAACCTACTGGGAGGGCTTCCTTGGCC

Figure 16:





Figure 17:

Figure 18:

