



- (51) **International Patent Classification:**
C12N 7/01 (2006.01) *A61K 31/7088* (2006.01)
A61K 48/00 (2006.01)
- (21) **International Application Number:**
PCT/US2012/034408
- (22) **International Filing Date:**
20 April 2012 (20.04.2012)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/478,012 21 April 2011 (21.04.2011) US
- (71) **Applicant (for all designated States except US):** **NATIONWIDE CHILDREN'S HOSPITAL, INC.** [US/US]; 700 Children's Drive, Room W172, Columbus, OH 43205 (US).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** **HARPER, Scott** **Quenton** [US/US]; 384 Briarbend Boulevard, Powell, OH 43065 (US). **LIU, Jian** [CN/US]; 2612 Lorain Court, Columbus, OH 43210 (US).
- (74) **Agent:** **NOLAND, Greta E.**; Marshall, Gerstein & Borun LLP, 233 S. Wacker Drive, 6300 Willis Tower, Chicago, IL 60606-6357 (US).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))
- with sequence listing part of description (Rule 5.2(a))

(54) **Title:** RECOMBINANT VIRUS PRODUCTS AND METHODS FOR INHIBITION OF EXPRESSION OF MYOTILIN

(57) **Abstract:** The present invention relates to RNA interference-based methods for inhibiting the expression of the myotilin gene. Recombinant adeno-associated viruses of the invention deliver DNAs encoding microRNAs that knock down the expression of myotilin. The methods have application in the treatment of muscular dystrophies such as Limb Girdle Muscular Dystrophy Type 1A.



RECOMBINANT VIRUS PRODUCTS AND
METHODS FOR INHIBITION OF EXPRESSION OF MYOTILIN

[0001] This application is a continuation-in-part of U.S. Provisional Patent Application No. 61/478,012 filed April 21, 2011, which is incorporated by reference herein in its entirety.

Field of the Invention

[0002] The present invention relates to RNA interference-based methods for inhibiting the expression of the myotilin gene. Recombinant adeno-associated viruses of the invention deliver DNAs encoding microRNAs that knock down the expression of myotilin. The methods have application in the treatment of muscular dystrophies such as Limb Girdle Muscular Dystrophy Type 1A.

Incorporation by Reference of the Sequence Listing

[0003] This application contains, as a separate part of disclosure, a Sequence Listing in computer-readable form (filename: 46208PCT_SeqListing.txt; 1,721,237 bytes – ASCII text file) which is incorporated by reference herein in its entirety.

Background

[0004] Muscular dystrophies (MDs) are a group of genetic diseases. The group is characterized by progressive weakness and degeneration of the skeletal muscles that control movement. Some forms of MD develop in infancy or childhood, while others may not appear until middle age or later. The disorders differ in terms of the distribution and extent of muscle weakness (some forms of MD also affect cardiac muscle), the age of onset, the rate of progression, and the pattern of inheritance.

[0005] One group of MDs is the limb girdle group (LGMD) of MDs. LGMDs are rare conditions and they present differently in different people with respect to age of onset, areas of muscle weakness, heart and respiratory involvement, rate of progression and severity. LGMDs can begin in childhood, adolescence, young adulthood or even later. Both genders are affected equally. LGMDs cause weakness in the shoulder and pelvic girdle, with nearby muscles in the upper legs and arms sometimes also weakening with time. Weakness of the

legs often appears before that of the arms. Facial muscles are usually unaffected. As the condition progresses, people can have problems with walking and may need to use a wheelchair over time. The involvement of shoulder and arm muscles can lead to difficulty in raising arms over head and in lifting objects. In some types of LGMD, the heart and breathing muscles may be involved.

[0006] There are at least nineteen forms of LGMD, and the forms are classified by their associated genetic defects.

Type	Pattern of Inheritance	Gene or Chromosome
LGMD1A	Autosomal dominant	Myotilin gene
LGMD1B	Autosomal dominant	Lamin A/C gene
LGMD1C	Autosomal dominant	Caveolin gene
LGMD1D	Autosomal dominant	Chromosome 7
LGMD1E	Autosomal dominant	Desmin gene
LGMD1F	Autosomal dominant	Chromosome 7
LGMD1G	Autosomal dominant	Chromosome 4
LGMD2A	Autosomal recessive	Calpain-3 gene
LGMD2B	Autosomal recessive	Dysferlin gene
LGMD2C	Autosomal recessive	Gamma-sarcoglycan gene
LGMD2D	Autosomal recessive	Alpha-sarcoglycan gene
LGMD2E	Autosomal recessive	Beta-sarcoglycan gene
LGMD2F	Autosomal recessive	Delta-sarcoglycan gene
LGMD2G	Autosomal recessive	Telethonin gene
LGMD2H	Autosomal recessive	TRIM32
LGMD2I	Autosomal recessive	FKRP gene
LGMD2J	Autosomal recessive	Titin gene
LGMD2K	Autosomal recessive	POMT1 gene
LGMD2L	Autosomal recessive	Fukutin gene

[0007] Specialized tests for LGMD are now available through a national scheme for diagnosis, the National Commissioning Group (NCG).

[0008] LGMD1A is caused by gain-of-function missense mutations in the myotilin (MYOT) gene [Hauser et al., *Am. J. Hum. Genet.*, 71: 1428-1432 (2002); Hauser et al., *Hum. Mol. Genet.*, 9: 2141-2147 (2000); Shalaby et al., *J. Neuropathol. Exp. Neurol.*, 68: 701-707 (2009)]. LGMD1A patients develop proximal leg and arm weakness in early adulthood (25 years is mean onset age), which progresses to the distal limb musculature. At the histological level, patients show myofiber degeneration and size variability, fiber splitting, centrally located myonuclei, autophagic vesicles, and replacement of myofibers with fat and fibrotic tissue, which are all common features of muscular dystrophy. Patients with LGMD1A also develop intramuscular myofibrillar protein aggregates, rimmed vacuoles, and severe Z-disc disorganization (called Z-disc streaming), which completely disrupt the sarcomeric structure. A transgenic mouse model, the T57I mouse model, using a mutant human MYOT allele has been developed [Garvey et al., *Hum. Mol. Genet.* 15: 2348-2362 (2006)]. Importantly, T57I mice recapitulate the progressive histological and functional abnormalities associated with LGMD1A, including reduced muscle size, muscle weakness, intramuscular myofibrillar aggregates, Z-disc streaming, and centrally located myonuclei.

[0009] The myotilin gene encodes a 57 kDa protein expressed primarily in skeletal and cardiac muscle. Myotilin appears to function as a structural component of the Z-disc, and may therefore contribute to sarcomere assembly, actin filament stabilization, and force transmission in striated muscle. Nevertheless, myotilin is not required for normal muscle development or function, since myotilin null mice are overtly and histologically normal. Specifically, mouse muscles lacking myotilin are indistinguishable from wild type in muscle mass, myofiber size, contractile strength (specific force), and sarcolemmal integrity. Moreover, MYOT null mice develop normally, live a normal life span, and show no histological evidence of muscular dystrophy or Z-disc malformations. Mouse and human myotilin transcripts are expressed in the same tissues, have the same genomic structures, and protein sequences are highly conserved (90% identity; 94% similarity), which indicates a conserved functional.

[0010] RNA interference (RNAi) is a mechanism of gene regulation in eukaryotic cells that has been considered for the treatment of various diseases. RNAi refers to post-

transcriptional control of gene expression mediated by microRNAs (miRNAs). The miRNAs are small (21-25 nucleotides), noncoding RNAs that share sequence homology and base-pair with cognate messenger RNAs (mRNAs). The interaction between the miRNAs and mRNAs directs cellular gene silencing machinery to prevent the translation of the mRNAs. The RNAi pathway is summarized in Duan (Ed.), Section 7.3 of Chapter 7 in *Muscle Gene Therapy*, Springer Science+Business Media, LLC (2010). Section 7.4 mentions MYOT RNAi therapy of LGMD1A in mice to demonstrate proof-of-principle for RNAi therapy of dominant muscle disorders.

[0011] As an understanding of natural RNAi pathways has developed, researchers have designed artificial miRNAs for use in regulating expression of target genes for treating disease. As described in Section 7.4 of Duan, *supra*, artificial miRNAs can be transcribed from DNA expression cassettes. The miRNA sequence specific for a target gene is transcribed along with sequences required to direct processing of the miRNA in a cell. Viral vectors such as adeno-associated virus have been used to deliver miRNAs to muscle [Fechner et al., *J. Mol. Med.*, 86: 987-997 (2008)].

[0012] Adeno-associated virus (AAV) is a replication-deficient parvovirus, the single-stranded DNA genome of which is about 4.7 kb in length including 145 nucleotide inverted terminal repeat (ITRs). There are multiple serotypes of AAV. The nucleotide sequences of the genomes of the AAV serotypes are known. For example, the complete genome of AAV-1 is provided in GenBank Accession No. NC_002077; the complete genome of AAV-2 is provided in GenBank Accession No. NC_001401 and Srivastava et al., *J. Virol.*, 45: 555-564 (1983); the complete genome of AAV-3 is provided in GenBank Accession No. NC_1829; the complete genome of AAV-4 is provided in GenBank Accession No. NC_001829; the AAV-5 genome is provided in GenBank Accession No. AF085716; the complete genome of AAV-6 is provided in GenBank Accession No. NC_001862; at least portions of AAV-7 and AAV-8 genomes are provided in GenBank Accession Nos. AX753246 and AX753249, respectively; the AAV-9 genome is provided in Gao et al., *J. Virol.*, 78: 6381-6388 (2004); the AAV-10 genome is provided in *Mol. Ther.*, 13(1): 67-76 (2006); and the AAV-11 genome is provided in *Virology*, 330(2): 375-383 (2004). Cis-acting sequences directing viral DNA replication (rep), encapsidation/packaging and host cell chromosome integration are contained within the AAV ITRs. Three AAV promoters (named p5, p19, and p40 for their relative map locations) drive the expression of the two AAV internal open reading frames encoding rep and cap genes. The two rep promoters (p5 and p19), coupled with the

differential splicing of the single AAV intron (at nucleotides 2107 and 2227), result in the production of four rep proteins (rep 78, rep 68, rep 52, and rep 40) from the rep gene. Rep proteins possess multiple enzymatic properties that are ultimately responsible for replicating the viral genome. The cap gene is expressed from the p40 promoter and it encodes the three capsid proteins VP1, VP2, and VP3. Alternative splicing and non-consensus translational start sites are responsible for the production of the three related capsid proteins. A single consensus polyadenylation site is located at map position 95 of the AAV genome. The life cycle and genetics of AAV are reviewed in Muzyczka, Current Topics in Microbiology and Immunology, 158: 97-129 (1992).

[0013] AAV possesses unique features that make it attractive as a vector for delivering foreign DNA to cells, for example, in gene therapy. AAV infection of cells in culture is noncytopathic, and natural infection of humans and other animals is silent and asymptomatic. Moreover, AAV infects many mammalian cells allowing the possibility of targeting many different tissues *in vivo*. Moreover, AAV transduces slowly dividing and non-dividing cells, and can persist essentially for the lifetime of those cells as a transcriptionally active nuclear episome (extrachromosomal element). The AAV proviral genome is infectious as cloned DNA in plasmids which makes construction of recombinant genomes feasible. Furthermore, because the signals directing AAV replication, genome encapsidation and integration are contained within the ITRs of the AAV genome, some or all of the internal approximately 4.3 kb of the genome (encoding replication and structural capsid proteins, rep-cap) may be replaced with foreign DNA. The rep and cap proteins may be provided in trans. Another significant feature of AAV is that it is an extremely stable and hearty virus. It easily withstands the conditions used to inactivate adenovirus (56o to 65oC for several hours), making cold preservation of AAV less critical. AAV may even be lyophilized. Finally, AAV-infected cells are not resistant to superinfection.

[0014] There remains a need in the art for a treatment for LGMD1A.

Summary

[0015] The present invention provides methods and products for preventing or inhibiting the expression of the MYOT gene. The methods of the invention utilize RNAi to prevent or inhibit the expression of the MYOT gene. The methods involve delivering inhibitory RNAs specific for the MYOT gene to muscle cells. The MYOT inhibitory RNAs contemplated include, but are not limited to, antisense RNAs, small inhibitory RNAs (siRNAs), short

hairpin RNAs (shRNAs) or artificial microRNAs (MYOT miRNAs) that inhibit expression of MYOT. Use of the methods and products is indicated, for example, in preventing or treating LGMD1A. Some embodiments of the invention exploit the unique properties of AAV to deliver DNA encoding MYOT inhibitory RNAs to muscle cells. Other embodiments of the invention utilize other vectors (for example, other viral vectors such as adenovirus, retrovirus, lentivirus, equine-associated virus, alphavirus, pox viruses, herpes virus, polio virus, sindbis virus and vaccinia viruses) to deliver polynucleotides encoding MYOT inhibitory RNAs.

[0016] In one aspect, the invention provides MYOT miRNAs. In another aspect, the invention provides rAAV encoding the MYOT miRNAs wherein the rAAV lack rep and cap genes. In some embodiments, the MYOT miRNA comprises an miRNA antisense guide strand selected from those set out in SEQ ID NO: 7 through SEQ ID NO: 11266. These sequences comprise antisense “guide” strand sequences of the invention of varying sizes. The antisense guide strand is the strand of the mature miRNA duplex that becomes the RNA component of the RNA induced silencing complex ultimately responsible for sequence-specific gene silencing. See Section 7.3 of Duan, *supra*. For example, the first antisense guide strand in SEQ ID NO: 7 corresponds to (is the reverse complement of) the 3' end of the myotilin sequence set out in Figure 1. The second antisense guide strand (SEQ ID NO: 8) is offset one nucleotide from the first and so on. In some embodiments, the GC content of the antisense guide strand is 60% or less, and/or the 5' end of the antisense guide strand is more AU rich while the 3' end is more GC rich. Exemplified MYOT miRNA are encoded by the DNAs set out in SEQ ID NOs: 1, 2, 3 and 4.

[0017] In another aspect, the invention provides a composition comprising the rAAV encoding the MYOT miRNA.

[0018] In yet another aspect, the invention provides a method of preventing or inhibiting expression of the MYOT gene in a cell comprising contacting the cell with a rAAV encoding an MYOT miRNA, wherein the miRNA is encoded by the DNA set out in SEQ ID NO: 1, 2, 3 or 4 and wherein the rAAV lacks rep and cap genes. Expression of MYOT is inhibited by at least 10, 20, 30, 40, 50, 60, 70, 80, 90, 95 or 99 percent.

[0019] In still another aspect, the invention provides a method of delivering DNA encoding the MYOT miRNA set out in SEQ ID NO: 1, 2, 3 or 4 to an animal in need thereof, comprising respectively administering to the animal a rAAV encoding the MYOTmi RNA, wherein the rAAV lacks rep and cap genes.

[0020] In yet another aspect, the invention provides a method of preventing or treating a muscular dystrophy (including, but not limited to, LGMD1A) comprising administering a rAAV encoding an MYOT miRNA, wherein the miRNA is encoded by the DNA set out in SEQ ID NO: 1, 2, 3 or 4 and wherein the rAAV lacks rep and cap genes. “Treating” may include ameliorating one or more symptoms of the muscular dystrophy (such as LGMD1A). Molecular, biochemical, histological, and functional endpoints demonstrate the therapeutic efficacy of MYOT miRNAs. Endpoints contemplated by the invention include one or more of: the reduction or elimination of mutant MYOT protein in affected muscles, MYOT gene knockdown, reduction or elimination of (for example, LGMD1A-associated) pathogenic protein aggregates in muscle, increase in myofiber diameters, and improvement in muscle strength.

Detailed Description

[0021] Recombinant AAV genomes of the invention comprise one or more AAV ITRs flanking a polynucleotide encoding, for example, one or more MYOT miRNAs. The polynucleotide is operatively linked to transcriptional control DNA, specifically promoter DNA that is functional in target. Commercial providers such as Ambion Inc. (Austin, TX), Dharmacon Inc. (Lafayette, CO), InvivoGen (San Diego, CA), and Molecular Research Laboratories, LLC (Herndon, VA) generate custom inhibitory RNA molecules. In addition, commercially kits are available to produce custom siRNA molecules, such as SILENCER™ siRNA Construction Kit (Ambion Inc., Austin, TX) or psiRNA System (InvivoGen, San Diego, CA). Embodiments include a rAAV genome comprising: the DNA set out in SEQ ID NO: 1 encoding the MYOT miRNA (named “miMyoT-1291”), the DNA set out in SEQ ID NO: 2 encoding the MYOT miRNA (named “miMyoT-1321”), the DNA set out in SEQ ID NO: 3 encoding the MYOT miRNA (named “miMyoT-1366”) or the DNA set out in SEQ ID NO: 4 encoding the MYOT miRNA (named “miMyoT-1490”).

[0022] The rAAV genomes of the invention lack AAV rep and cap DNA. AAV DNA in the rAAV genomes may be from any AAV serotype for which a recombinant virus can be derived including, but not limited to, AAV serotypes AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-6, AAV-7, AAV-8, AAV-9, AAV-10 and AAV-11. As noted in the Background section above, the nucleotide sequences of the genomes of various AAV serotypes are known in the art.

[0023] DNA plasmids of the invention comprise rAAV genomes of the invention. The DNA plasmids are transferred to cells permissible for infection with a helper virus of AAV (e.g., adenovirus, E1-deleted adenovirus or herpesvirus) for assembly of the rAAV genome into infectious viral particles. Techniques to produce rAAV particles, in which an AAV genome to be packaged, rep and cap genes, and helper virus functions are provided to a cell are standard in the art. Production of rAAV requires that the following components are present within a single cell (denoted herein as a packaging cell): a rAAV genome, AAV rep and cap genes separate from (i.e., not in) the rAAV genome, and helper virus functions. The AAV rep and cap genes may be from any AAV serotype for which recombinant virus can be derived and may be from a different AAV serotype than the rAAV genome ITRs, including, but not limited to, AAV serotypes AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-6, AAV-7, AAV-8, AAV-9, AAV-10 and AAV-11. Production of pseudotyped rAAV is disclosed in, for example, WO 01/83692 which is incorporated by reference herein in its entirety.

[0024] A method of generating a packaging cell is to create a cell line that stably expresses all the necessary components for AAV particle production. For example, a plasmid (or multiple plasmids) comprising a rAAV genome lacking AAV rep and cap genes, AAV rep and cap genes separate from the rAAV genome, and a selectable marker, such as a neomycin resistance gene, are integrated into the genome of a cell. AAV genomes have been introduced into bacterial plasmids by procedures such as GC tailing (Samulski et al., 1982, Proc. Natl. Acad. Sci. USA, 79:2077-2081), addition of synthetic linkers containing restriction endonuclease cleavage sites (Laughlin et al., 1983, Gene, 23:65-73) or by direct, blunt-end ligation (Senapathy & Carter, 1984, J. Biol. Chem., 259:4661-4666). The packaging cell line is then infected with a helper virus such as adenovirus. The advantages of this method are that the cells are selectable and are suitable for large-scale production of rAAV. Other examples of suitable methods employ adenovirus or baculovirus rather than plasmids to introduce rAAV genomes and/or rep and cap genes into packaging cells.

[0025] General principles of rAAV production are reviewed in, for example, Carter, 1992, Current Opinions in Biotechnology, 1533-539; and Muzyczka, 1992, Curr. Topics in Microbial. and Immunol., 158:97-129). Various approaches are described in Ratschin et al., Mol. Cell. Biol. 4:2072 (1984); Hermonat et al., Proc. Natl. Acad. Sci. USA, 81:6466 (1984); Tratschin et al., Mol. Cell. Biol. 5:3251 (1985); McLaughlin et al., J. Virol., 62:1963 (1988); and Lebkowski et al., 1988 Mol. Cell. Biol., 7:349 (1988). Samulski et al. (1989, J. Virol., 63:3822-3828); U.S. Patent No. 5,173,414; WO 95/13365 and corresponding U.S. Patent No.

5,658,776 ; WO 95/13392; WO 96/17947; PCT/US98/18600; WO 97/09441 (PCT/US96/14423); WO 97/08298 (PCT/US96/13872); WO 97/21825 (PCT/US96/20777); WO 97/06243 (PCT/FR96/01064); WO 99/11764; Perrin et al. (1995) Vaccine 13:1244-1250; Paul et al. (1993) Human Gene Therapy 4:609-615; Clark et al. (1996) Gene Therapy 3:1124-1132; U.S. Patent. No. 5,786,211; U.S. Patent No. 5,871,982; and U.S. Patent. No. 6,258,595. The foregoing documents are hereby incorporated by reference in their entirety herein, with particular emphasis on those sections of the documents relating to rAAV production.

[0026] The invention thus provides packaging cells that produce infectious rAAV. In one embodiment packaging cells may be stably transformed cancer cells such as HeLa cells, 293 cells and PerC.6 cells (a cognate 293 line). In another embodiment, packaging cells are cells that are not transformed cancer cells, such as low passage 293 cells (human fetal kidney cells transformed with E1 of adenovirus), MRC-5 cells (human fetal fibroblasts), WI-38 cells (human fetal fibroblasts), Vero cells (monkey kidney cells) and FRhL-2 cells (rhesus fetal lung cells).

[0027] Recombinant AAV (i.e., infectious encapsidated rAAV particles) of the invention comprise a rAAV genome. Embodiments include, but are not limited to, the rAAV including a genome encoding the MYOT miRNA set out in SEQ ID NO: 1 (named "AAV-U6-miMyoT-1291"), the rAAV including a genome encoding the MYOT miRNA set out in SEQ ID NO: 2 (named "AAV-U6-miMyoT-1321"), the rAAV including a genome encoding the MYOT miRNA set out in SEQ ID NO: 3 (named "AAV-U6-miMyoT-1366") and the rAAV including a genome encoding the MYOT miRNA set out in SEQ ID NO: 4 (named "AAV-U6-miMyoT-1490"). The genomes of the rAAV lack AAV rep and cap DNA, that is, there is no AAV rep or cap DNA between the ITRs of the genomes.

[0028] The rAAV may be purified by methods standard in the art such as by column chromatography or cesium chloride gradients. Methods for purifying rAAV vectors from helper virus are known in the art and include methods disclosed in, for example, Clark et al., Hum. Gene Ther., 10(6): 1031-1039 (1999); Schenpp and Clark, Methods Mol. Med., 69 427-443 (2002); U.S. Patent No. 6,566,118 and WO 98/09657.

[0029] In another embodiment, the invention contemplates compositions comprising rAAV of the present invention. Compositions of the invention comprise rAAV in a pharmaceutically acceptable carrier. The compositions may also comprise other ingredients

such as diluents and adjuvants. Acceptable carriers, diluents and adjuvants are nontoxic to recipients and are preferably inert at the dosages and concentrations employed, and include buffers such as phosphate, citrate, or other organic acids; antioxidants such as ascorbic acid; low molecular weight polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween, pluronics or polyethylene glycol (PEG).

[0030] Titers of rAAV to be administered in methods of the invention will vary depending, for example, on the particular rAAV, the mode of administration, the treatment goal, the individual, and the cell type(s) being targeted, and may be determined by methods standard in the art. Titers of rAAV may range from about 1×10^6 , about 1×10^7 , about 1×10^8 , about 1×10^9 , about 1×10^{10} , about 1×10^{11} , about 1×10^{12} , about 1×10^{13} to about 1×10^{14} or more DNase resistant particles (DRP) per ml. Dosages may also be expressed in units of viral genomes (vg).

[0031] Methods of transducing a target cell with rAAV, *in vivo* or *in vitro*, are contemplated by the invention. The *in vivo* methods comprise the step of administering an effective dose, or effective multiple doses, of a composition comprising a rAAV of the invention to an animal (including a human being) in need thereof. If the dose is administered prior to development of a disorder/disease, the administration is prophylactic. If the dose is administered after the development of a disorder/disease, the administration is therapeutic. In embodiments of the invention, an effective dose is a dose that alleviates (eliminates or reduces) at least one symptom associated with the disorder/disease state being treated, that slows or prevents progression to a disorder/disease state, that slows or prevents progression of a disorder/disease state, that diminishes the extent of disease, that results in remission (partial or total) of disease, and/or that prolongs survival. An example of a disease contemplated for prevention or treatment with methods of the invention is LGMD1A.

[0032] Combination therapies are also contemplated by the invention. Combination as used herein includes both simultaneous treatment or sequential treatments. Combinations of methods of the invention with standard medical treatments (e.g., corticosteroids) are specifically contemplated, as are combinations with novel therapies.

[0033] Administration of an effective dose of the compositions may be by routes standard in the art including, but not limited to, intramuscular, parenteral, intravenous, oral, buccal, nasal, pulmonary, intracranial, intraosseous, intraocular, rectal, or vaginal. Route(s) of administration and serotype(s) of AAV components of the rAAV (in particular, the AAV ITRs and capsid protein) of the invention may be chosen and/or matched by those skilled in the art taking into account the infection and/or disease state being treated and the target cells/tissue(s) that are to express the MYOT miRNAs.

[0034] In particular, actual administration of rAAV of the present invention may be accomplished by using any physical method that will transport the rAAV recombinant vector into the target tissue of an animal. Administration according to the invention includes, but is not limited to, injection into muscle, the bloodstream and/or directly into the liver. Simply resuspending a rAAV in phosphate buffered saline has been demonstrated to be sufficient to provide a vehicle useful for muscle tissue expression, and there are no known restrictions on the carriers or other components that can be co-administered with the rAAV (although compositions that degrade DNA should be avoided in the normal manner with rAAV). Capsid proteins of a rAAV may be modified so that the rAAV is targeted to a particular target tissue of interest such as muscle. See, for example, WO 02/053703, the disclosure of which is incorporated by reference herein. Pharmaceutical compositions can be prepared as injectable formulations or as topical formulations to be delivered to the muscles by transdermal transport. Numerous formulations for both intramuscular injection and transdermal transport have been previously developed and can be used in the practice of the invention. The rAAV can be used with any pharmaceutically acceptable carrier for ease of administration and handling.

[0035] For purposes of intramuscular injection, solutions in an adjuvant such as sesame or peanut oil or in aqueous propylene glycol can be employed, as well as sterile aqueous solutions. Such aqueous solutions can be buffered, if desired, and the liquid diluent first rendered isotonic with saline or glucose. Solutions of rAAV as a free acid (DNA contains acidic phosphate groups) or a pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. A dispersion of rAAV can also be prepared in glycerol, liquid polyethylene glycols and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms. In this connection, the sterile aqueous media

employed are all readily obtainable by standard techniques well-known to those skilled in the art.

[0036] The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating actions of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of a dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal and the like. In many cases it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by use of agents delaying absorption, for example, aluminum monostearate and gelatin.

[0037] Sterile injectable solutions are prepared by incorporating rAAV in the required amount in the appropriate solvent with various other ingredients enumerated above, as required, followed by filter sterilization. Generally, dispersions are prepared by incorporating the sterilized active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying technique that yield a powder of the active ingredient plus any additional desired ingredient from the previously sterile-filtered solution thereof.

[0038] Transduction with rAAV may also be carried out *in vitro*. In one embodiment, desired target muscle cells are removed from the subject, transduced with rAAV and reintroduced into the subject. Alternatively, syngeneic or xenogeneic muscle cells can be used where those cells will not generate an inappropriate immune response in the subject.

[0039] Suitable methods for the transduction and reintroduction of transduced cells into a subject are known in the art. In one embodiment, cells can be transduced *in vitro* by

combining rAAV with muscle cells, *e.g.*, in appropriate media, and screening for those cells harboring the DNA of interest using conventional techniques such as Southern blots and/or PCR, or by using selectable markers. Transduced cells can then be formulated into pharmaceutical compositions, and the composition introduced into the subject by various techniques, such as by intramuscular, intravenous, subcutaneous and intraperitoneal injection, or by injection into smooth and cardiac muscle, using *e.g.*, a catheter.

[0040] Transduction of cells with rAAV of the invention results in sustained expression of MYOT miRNAs. The present invention thus provides methods of administering/delivering rAAV which express MYOT miRNAs to an animal, preferably a human being. These methods include transducing tissues (including, but not limited to, tissues such as muscle, organs such as liver and brain, and glands such as salivary glands) with one or more rAAV of the present invention. Transduction may be carried out with gene cassettes comprising tissue specific control elements. For example, one embodiment of the invention provides methods of transducing muscle cells and muscle tissues directed by muscle specific control elements, including, but not limited to, those derived from the actin and myosin gene families, such as from the myoD gene family [See Weintraub et al., *Science*, 251: 761-766 (1991)], the myocyte-specific enhancer binding factor MEF-2 [Cserjesi and Olson, *Mol Cell Biol* 11: 4854-4862 (1991)], control elements derived from the human skeletal actin gene [Muscat et al., *Mol Cell Biol*, 7: 4089-4099 (1987)], the cardiac actin gene, muscle creatine kinase sequence elements [See Johnson et al., *Mol Cell Biol*, 9:3393-3399 (1989)] and the murine creatine kinase enhancer (mCK) element, control elements derived from the skeletal fast-twitch troponin C gene, the slow-twitch cardiac troponin C gene and the slow-twitch troponin I gene: hypoxia-inducible nuclear factors [Semenza et al., *Proc Natl Acad Sci USA*, 88: 5680-5684 (1991)], steroid-inducible elements and promoters including the glucocorticoid response element (GRE) [See Mader and White, *Proc. Natl. Acad. Sci. USA*, 90: 5603-5607 (1993)], and other control elements.

[0041] Muscle tissue is an attractive target for *in vivo* DNA delivery, because it is not a vital organ and is easy to access. The invention contemplates sustained expression of miRNAs from transduced myofibers.

[0042] By “muscle cell” or “muscle tissue” is meant a cell or group of cells derived from muscle of any kind (for example, skeletal muscle and smooth muscle, *e.g.* from the digestive tract, urinary bladder, blood vessels or cardiac tissue). Such muscle cells may be

differentiated or undifferentiated, such as myoblasts, myocytes, myotubes, cardiomyocytes and cardiomyoblasts.

[0043] The term “transduction” is used to refer to the administration/delivery of MYOT miRNAs to a recipient cell either *in vivo* or *in vitro*, via a replication-deficient rAAV of the invention resulting in expression of a MYOT miRNA by the recipient cell.

[0044] Thus, the invention provides methods of administering an effective dose (or doses, administered essentially simultaneously or doses given at intervals) of rAAV that encode MYOT miRNAs to a patient in need thereof.

Brief Description of the Drawing

[0045] Figure 1 shows target sites in the myotilin sequence (SEQ ID NO: 11266) for exemplified miRNAs.

[0046] Figure 2 sets out sequences of two MYOT-targeted miRNAs. In each panel, the top sequences indicate the DNA templates from which each respective miRNA is transcribed. In the top panel, the DNA template miMYOT.1321 is SEQ ID NO: 2. In the bottom panel, the DNA template miMYOT.1366 is SEQ ID NO: 3. The folded miRNA transcripts are shown as hairpin structures. The miMYOT.1321 folded miRNA is SEQ ID NO: 11268. The miMYOT.1366 folded miRNA is SEQ ID NO: 11271. The mature miMYO.1321 (SEQ ID NO: 11270 which pairs with SEQ ID NO: 11269 in the figure) and miDUX4.1366 (SEQ ID NO: 11273 which pairs with SEQ ID NO: 11272 in the figure) sequences arise following processing in target cells by host miRNA processing machinery (including Drosha, DGCR8, Dicer, and Exportin-5). Sequences shaded in gray indicate sites used for cloning each miRNA into the U6T6 vector. The nucleotides corresponding to the mature miRNA antisense guide strand that ultimately helps catalyze cleavage of the MYOT target mRNA are underlined in the miRNA hairpin portions of this diagram. The gray and black arrowheads indicate Drosha- and Dicer- catalyzed cleavage sites, respectively. The numbers 13, 35, 53, and 75 are provided for orientation. The sequences between (and including) positions 35-53 are derived from the natural human mir-30a sequence, except the A at position 39, which is a G in the normal mir-30a sequence. This was changed to an A to facilitate folding of the miRNA loop, based on *in silico* RNA folding models. The base of the stem (5' of position 13 and 3' of position 75) is also derived from mir-30a structure and sequence with some modifications depending on the primary sequence of the guide strand. Specifically, the

nucleotide at position 13 can vary to help facilitate a required mismatched between the position 13 and 75 nucleotides. This bulged structure is hypothesized to facilitate proper Drosha cleavage.

[0047] Figure 3 shows the effect of MYOT-targeted miRNAs in LGMD1A mice expressing mutant myotilin (MYOT). Figure 3A is a Western blot showing knockdown of mutant myotilin expression in muscle extracts from three-month old LGMD1A mice, where Left (L) = miMYOT treatment side and Right (R) = miGFP control treated side. Figure 3B shows real-time PCR results confirming the Western data.

Examples

[0048] Aspects and embodiments of the invention are illustrated by the following examples. Example 1 describes miRNAs specific for the MYOT gene. Example 2 describes the effect of the miRNAs on the expression of MYOT as measured by real-time PCR. Example 3 describes rAAV encoding the miRNAs. Example 4 describes the effect of the U6T6 expressing the miRNAs on the expression of MYOT as measured by Western blot. Example 5 describes delivery of MYOT miRNA to newborn mice. Example 6 describes delivery of MYOT miRNA to adult mice.

Example 1

MicroRNAs specific for the MYOT gene

[0049] Six DNAs encoding miRNAs specific for the MYOT gene were generated by PCR. The PCR primers used had the following sequences.

Primer 775 (miMyoT-592-Forward) (SEQ ID NO: 11274):

AAAACTCGAGTGAGCGACCTGATTACAATAGCAGTAACTGTAAAGCCACAGATGGG

Primer 776 (miMyoT-592-Reverse) (SEQ ID NO: 11275):

TTTTACTAGTAGGCAGCCTGATTACAATAGCAGTAAACCCATCTGTGGCTTTACAG

Primer 777 (miMyoT-1291-Forward) (SEQ ID NO: 11276):

AAAACTCGAGTGAGCGACTGGATGTCCTTGCAAAGAAGTAAAGCCACAGATGGG

Primer 778 (miMyoT-1291-Reverse) (SEQ ID NO: 11277):

TTTTACTAGTAGGCAGCTGGATGTCCTTGCAAAGAAGTAAAGCCACAGATGGG

Primer 779 (miMyoT-1321-Forward) (SEQ ID NO: 11278):

AAAACTCGAGTGAGCGCGCACCAATGTTTATCTACAACTGTAAAGCCACAGATGGG

Primer 780 (miMyoT-1321-Reverse) (SEQ ID NO: 11279):

TTTTACTAGTAGGCAAGCACCAATGTTTATCTACAAACCCATCTGTGGCTTTACAG

Primer 781 (miMyoT-1366-Forward) (SEQ ID NO: 11280) :

AAAACTCGAGTGAGCGAGGAGATTTCAGTGAACTAGAACTGTAAAGCCACAGATGGG

Primer 782 (miMyoT-1366-Reverse) (SEQ ID NO: 11281):

TTTTACTAGTAGGCAGGGAGATTTCAGTGAACTAGAACCCATCTGTGGCTTTACAG

Primer 783 (miMyoT-1490-Forward) (SEQ ID NO: 11282):

AAAACTCGAGTGAGCGCGAAGAGTTACTTTACTGATAACTGTAAAGCCACAGATGGG

Primer 784 (miMyoT-1490-Reverse) (SEQ ID NO: 11283):

TTTTACTAGTAGGCAGGAAGAGTTACTTTACTGATAACCCATCTGTGGCTTTACAG

Primer 785 (miMyoT-1603-Forward) (SEQ ID NO: 11284):

AAAACTCGAGTGAGCGAGCACGTCCAAACCAAACCTCTTCTGTAAAGCCACAGATGGG

Primer 786 (miMyoT-1603-Reverse) (SEQ ID NO: 11285):

TTTTACTAGTAGGCAGGCACGTCCAAACCAAACCTCTTCCCATCTGTGGCTTTACAG

[0050] DNA encoding a miRNA designated miMyoT-592 was generated using primers 775 and 776. DNA encoding miRNA designated miMyoT-1291 was generated using primers 777 and 778. DNA encoding miRNA designated miMyoT-1321 was generated using primers 779 and 780. DNA encoding miRNA designated miMyoT-1366 was generated using primers 781 and 782. DNA encoding miRNA designated miMyoT-1490 was generated using primers 783 and 784. DNA encoding miRNA designated miMyoT-1603 was generated using primers 785 and 786. The DNAs are set out below, wherein the number in the names indicates the 5' target nucleotide in the myotylin sequence (SEQ ID NO: 11267). See Figure 1 where the target sequences for the miRNAs in the myotilin sequence are underlined.

miMyoT-592

CTCGAGTGAGCGACCTGATTACAATAGCAGTAACTGTAAAGCCACAGATGGGTTTACTGCT
ATTGTAATCAGGCTGCCTACTAGA (SEQ ID NO: 5)

miMyoT-1291

CTCGAGTGAGCGACTGGATGTCCTTGCAAAAGAACTGTAAAGCCACAGATGGGTTATTTTGC
AAGGACATCCAGCTGCCTACTAGA (SEQ ID NO: 1)

miMyoT-1321

CTCGAGTGAGCGCGCACCAATGTTTATCTACAACTGTAAAGCCACAGATGGGTTTGTAGAT
AAACATTGGTGCTTGCCTACTAGA (SEQ ID NO: 2)

miMyoT-1366

CTCGAGTGAGCGAGGAGATTCAGTGAACTAGAACTGTAAAGCCACAGATGGGTTCTAGTTT
CACTGAATCTCCCTGCCTACTAGA (SEQ ID NO: 3)

miMyoT-1490

CTCGAGTGAGCGCGAAGAGTTACTTTACTGATAACTGTAAAGCCACAGATGGGTTATCAGTA
AAGTAACTCTTCCTGCCTACTAGA (SEQ ID NO: 4)

miMyoT-1603

CTCGAGTGAGCGAGCACGTCCAAACCAAACCTCTTCTGTAAAGCCACAGATGGGAAGAGTTTG
GTTTACGTGCCTGCCTACTAGA (SEQ ID NO: 6)

[0051] Figure 2 shows the DNA templates miMyoT.1321 and miMyoT.1366 and their corresponding folded and mature miRNAs.

[0052] One µg of each primer was added to a 1 cycle primer extension reaction: 95°C for 5 min.; 94° C for 2min.; 52° C for 1min.; 72° C for 15min.; and then holding at 4°C. The PCR products were cleaned up with the Qiagen QIAquick PCR Purification kit before being digested overnight with XhoI and SpeI restriction enzymes. The digestion product was then run on a 1.5% TBE gel and the band excised and purified using the Qiagen QIAquick Gel Extraction Kit.

[0053] The PCR products were ligated to a U6T6 vector (via XhoI and XbaI) overnight. This vector contains a mouse U6 promoter and an RNA polymerase III termination signal (6 thymidine nucleotides). miRNAs are cloned into XhoI + XbaI restriction sites located between the 3' end of the U6 promoter and the termination signal (SpeI on the 3' end of the DNA template for each miRNA has complementary cohesive ends with the XbaI site). The ligation product was transformed into chemically competent E-coli cells with a 42°C heat shock and incubated at 37°C shaking for 1 hour before being plated on kanamycin selection

plates. The colonies were allowed to grow overnight at 37°. The following day they were mini-prepped and sequenced for accuracy.

Example 2

Real-time PCR Reaction for Effect of Expression of MYOT miRNAs

[0054] Expression of the MYOT target sequence in the presence of the MYOT miRNAs was assayed. A lipofectamine 2000 transfection was done in C2C12 cells in a 12-well, white-walled assay plate. 52,000 cells were transfected with 100 ng of AAV-CMV-mutMyoT and 1500 ng of one of the U6T6 vectors described in Example 1 containing miRNA-encoding DNA. The assay was performed 48 hours later.

[0055] The media was removed from the cells and 1 µl of Trizol was added per well. Then the cells were resuspended and the lysates were transferred to 1.5 ml EP tubes. Samples were incubated at room temperature for 5 min and 200 µl chloroform was added. The tubes were shaken vigorously for 15 sec, incubated at room temperature for 3 min and centrifuged at 12,000 g for 15 min at 4 °C. Then the aqueous phase was transferred to a fresh tube and 0.5 ml isopropyl alcohol was added. The samples were incubated at room temperature for 10 min and centrifuged at 12,000 g for 10 min at 4°C. The RNA pellet was washed once with 1 ml 75% ethanol and aired dry. 20 µl of RNase-Free water was added to dissolve the pellet and the concentration/purification were measured by Nano-drop. 1.5 µg total RNA was added to cDNA generation reaction: 5°C for 10 min.; 37° C for 120 min.; 85° C for 5 sec and then holding at 4°C. The cDNA products were diluted at 1:10 and 4.5 µl was added to real-time PCR reaction. Human Myotilin was used to check the expression of the MYOT and the relative expression was normalized to mouse GAPDH expression.

[0056] U6T6-miMyoT-592 (SEQ ID NO: 5) showed higher expression of MYOT than U6T6-miGFP control. U6T6-miMyoT-1291 (SEQ ID NO: 1) reduced the expression of MYOT to 60%, U6T6-miMyoT-1321 (SEQ ID NO: 2) reduced the expression of MYOT to 19%, U6T6-miMyoT-1366 (SEQ ID NO: 3) reduced the expression of MYOT to 41.7%, U6T6-miMyoT-1490 (SEQ ID NO: 4) reduced the expression of MYOT to 55.3%, U6T6-miMyoT-1603 (SEQ ID NO: 6) reduced the expression of MYOT to 34.9%, when compared to U6T6-miGFP control.

Example 3

Production of rAAV Encoding MYOT MicroRNAs

[0057] The U6-miMYOT DNAs were cut from U6T6-miMYOT constructs at EcoRI sites and then respectively cloned into AAV6-hrGFPs to generate rAAV-U6-miMyoT vectors. These rAAV vectors express miRNA and hrGFP

Example 4

Western Blot Assay for Effect of

Expression of MYOT miRNAs from U6T6 vectors and rAAV

[0058] The effect of expression of MYOT miRNAs from the U6T6 vectors described in Example 1 and the rAAV described in Example 4 was assayed by Western blot.

[0059] One day before transfection, 293 cells were plated in a 24-well plate at 1.5×10^5 cells/well. The cells were then transfected with U6T6-miMyoT (592, 1291, 1321, 1366, 1490 or 1603) using Lipofectamine 2000 (Invitrogen, Cat. No. 11668-019).

[0060] Forty-eight hours after transfection, cells were collected and washed with cold PBS once. Seventy μ l lysis buffer (137 mM NaCl, 10mM Tris pH=7.4, 1% NP40) were then added. The cells were resuspended completely and incubated on ice for 30 min. The samples were centrifuged for 20 min at 13,000 rpm at 4°C and the supernatant was collected. The cell lysate was diluted 5-fold for the Lowry protein concentration assay (Bio-Rad Dc Protein Assay Reagent A, B, S; Cat. No. 500-0113, 500-0114, 500-115). Twenty μ g of each sample was taken and 2x sample buffer (100 mM Tris pH=6.8, 100 mM DTT, 10% glycerol, 2% SDS, 0.006% bromophenol blue) was added. The samples were boiled for 10 min and then put on ice.

[0061] The samples were loaded onto a 10% polyacrylamide gel (based on 37.5:1 acrylamide:bis acrylamide ratio, Bio-Rad, Cat. No. 161-0158), 15 μ g on a gel for each sample. Proteins were transferred to PVDF membranes at 15 V for 1 h using semi-dry transfer (Trans-Blot SD Semi-Dry Transfer Cell, Bio-Rad, Cat. No. 170-3940). The blots were placed into blocking buffer (5% non-fat dry milk, 30mM Tris pH=7.5, 150mM NaCl, 0.05% Tween-20) and agitated for 1 h at room temperature. The blocking buffer was decanted and anti-myotilin primary antibody solution (rabbit polyclonal generated by Bethyl Laboratories using a peptide corresponding to myotilin residues 473-488) was added and incubated with agitation overnight at 4°C. The membranes were then washed for 30 min, changing the wash

buffer (150 mM NaCl, 30mM Tris pH=7.5, 0.05% Tween-20) every 10 min. Peroxidase-conjugated Goat Anti-Mouse Antibody (Jackson ImmunoReserch, Cat. No. 115-035-146, 1:100,000) was added and incubated at room temperature for 2 h. The membranes were then washed for 30 min, changing the wash buffer every 10 min. The blots were placed in chemiluminescent working solution (Immobilon Weatern Chemiluminescent HRP Substrate, Millipore, Cat. No. WBKLS0500), incubated with agitation for 5 min at room temperature, and then exposed to X-ray film.

[0062] The membranes were washed for 20 min, changing the wash buffer every 10 min. Next, stripping buffer (2% SDS, 62.5 mM Tris pH=6.7, 100mM b-ME) was added to the blots and incubated at 50°C for 30 min. The membranes were washed again for 30 min, changing the wash buffer every 10 min. Then, the membranes were blocked again and re-probed with Anti-GAPDH primary antibody solution (Chemicon, Cat. No. MAB374, 1:200) and peroxidase-conjugated Goat Anti-Mouse Antibody (Jackson ImmunoReserch, Cat. No. 115-035-146, 1:100,000) was used as secondary antibody.

[0063] The film was scanned and the density ratio of MYOT to GAPDH was caculated. Compared to U6T6-miGFP control, the expression of MYOT was higher (1.08) in samples of U6T6-miMyoT-592 (SEQ ID NO: 5) and the the expression of MYOT was reduced to 78.9% by U6T6-miMyoT-1291 (SEQ ID NO: 1), 50.2% by U6T6-miMyoT-1321 (SEQ ID NO: 2), 60.2% by U6T6-miMyoT-1366 (SEQ ID NO: 3), 76.2% by U6T6-miMyoT-1490 (SEQ ID NO: 4), 87% by U6T6-miMyoT-1603 (SEQ ID NO: 6).

[0064] U6T6-miMYOT-1321 most effectively knocked down myotilin expression both in the real-time PCR and western-blot experiments. The knockdown effect by AAV-miMyoT-1321 was also confirmed by western-blot experiment.

Example 5

Delivery to Newborn Mice

[0065] The PCR genotype of newborn pups was determined to identify female WT or T57I MYOT mice (using human MYOT primers and Y chromosome primers). Bilateral intramuscular injections of 5×10^{10} AAV6.miMYOT-1321 or control AAV6.miGFP particles per leg in 1-2 day old mice were sufficient to saturate the lower limb musculature.

[0066] Phenotypic correction was then determined initially by histological analyses. Specifically, 3 months after viral delivery, muscles were harvested and cryopreserved. Ten micron serial cryosections were cut and stained with antibodies to detect myotilin-positive protein aggregates in T57I myofibers. AAV6.miMYOT-1321 muscles had significantly reduced numbers of aggregates per section compared to AAV6.miGFP or untreated controls. In addition, when AAV6.miMYOT-132-treated muscles did show occasional aggregates, they were significantly smaller than those seen in control-treated or untreated T57I animals. AAV6.miMYOT-132 treatment also improved muscle size deficits relative to the control treatment.

[0067] MYOT knockdown was confirmed by Western blot and real-time PCR as shown in Figure 3. The AAV delivered miMYOT-1321 significantly reduced mutant MYOT protein (Figure 3A) and mRNA (Figure 3B) in the muscles.

[0068] These results support therapeutic efficacy. Continuing experiments include determining the functional effects of MYOT knockdown in whole muscles by measuring EDL specific force.

Example 6

Delivery to Adult Mice

[0069] The PCR genotype of weanlings is determined and 12-week old mice which have significant pre-existing LGMD1A-associated pathology are chosen for treatment. 5×10^{11} AAV6 vectors are delivered to lower limb musculature by isolated leg perfusion. Phenotypic correction (including hindlimb grip strength, gross muscle parameters and EDL specific force are then measured using various methods over the following months.

[0070] While the present invention has been described in terms of specific embodiments, it is understood that variations and modifications will occur to those skilled in the art. Accordingly, only such limitations as appear in the claims should be placed on the invention.

[0071] All documents referred to in this application are hereby incorporated by reference in their entirety.

Claims

We claim:

1. A recombinant adeno-associated virus comprising the MYOT miRNA-encoding DNA set out in SEQ ID NO: 1, 2, 3 or 4, wherein the recombinant adeno-associated virus lacks rep and cap genes.
2. A composition comprising the recombinant adeno-associated virus of claim 1.
3. A method of inhibiting expression of the MYOT gene in a cell comprising contacting the cell with a recombinant adeno-associated virus of claim 1 or the composition of claim 2.
4. A method of delivering the MYOT miRNA-encoding DNA set out in SEQ ID NO: 1 to an animal in need thereof, comprising administering to the animal a recombinant adeno-associated virus comprising the MYOT miRNA-encoding DNA set out in SEQ ID NO: 1, wherein the recombinant adeno-associated virus lacks rep and cap genes.
5. A method of delivering the MYOT miRNA-encoding DNA set out in SEQ ID NO: 2 to an animal in need thereof, comprising administering to the animal a recombinant adeno-associated virus comprising the MYOT miRNA-encoding DNA set out in SEQ ID NO: 2, wherein the recombinant adeno-associated virus lacks rep and cap genes.
6. A method of delivering the MYOT miRNA-encoding DNA set out in SEQ ID NO: 3 to an animal in need thereof, comprising administering to the animal a recombinant adeno-associated virus comprising the MYOT miRNA-encoding DNA set out in SEQ ID NO: 3, wherein the recombinant adeno-associated virus lacks rep and cap genes.
7. A method of delivering the MYOT miRNA-encoding DNA set out in SEQ ID NO: 4 to an animal in need thereof, comprising administering to the animal a recombinant adeno-

associated virus comprising the MYOT miRNA-encoding DNA set out in SEQ ID NO: 4, wherein the recombinant adeno-associated virus lacks rep and cap genes.

8. A method of treating limb girdle muscular dystrophy type 1A comprising administering a recombinant adeno-associated virus comprising the MYOT miRNA-encoding DNA set out in SEQ ID NO: 1, wherein the recombinant adeno-associated virus lacks rep and cap genes.

9. A method of treating limb girdle muscular dystrophy type 1A comprising administering a recombinant adeno-associated virus comprising the MYOT miRNA-encoding DNA set out in SEQ ID NO: 2, wherein the recombinant adeno-associated virus lacks rep and cap genes.

10. A method of treating limb girdle muscular dystrophy type 1A comprising administering a recombinant adeno-associated virus comprising the MYOT miRNA-encoding DNA set out in SEQ ID NO: 3, wherein the recombinant adeno-associated virus lacks rep and cap genes.

11. A method of treating limb girdle muscular dystrophy type 1A comprising administering a recombinant adeno-associated virus comprising the MYOT miRNA-encoding DNA set out in SEQ ID NO: 4, wherein the recombinant adeno-associated virus lacks rep and cap genes.

12. The recombinant adeno-associated virus of claim 1 wherein expression of the miRNA-encoding DNA is under the control of a CMV promoter, a muscle creatine kinase (MCK) promoter, an alpha-myosin heavy chain enhancer-/MCK enhancer-promoter (MHCK7) or a desmin promoter.

13. The composition of claim 2 wherein expression of the miRNA-encoding DNA is under the control of a CMV promoter, a muscle creatine kinase (MCK) promoter, an alpha-myosin heavy chain enhancer-/MCK enhancer-promoter (MHCK7) or a desmin promoter.

14. The method of any one of claims 3-11 wherein expression of the miRNA-encoding DNA is under the control of a CMV promoter, a muscle creatine kinase (MCK) promoter, an alpha-myosin heavy chain enhancer-/MCK enhancer-promoter (MHCK7) or a desmin promoter.


Figure 1

Human C450T myotilin sequence (target sites are underlined and highlighted)


GGGAAGGAGATGCCTCTTCCTTCCCTTCAATAGTGGGTAAACCCAGCTGGCA
CCCTCTGGAACCTACGGGAACAATATTCTTCAAGAGAAGGTCACCTCTACCAAAG
CCAGGAGCACAGTATTCTCAGGATCTCAACAAGGAAGAGCAGACCAAGGTTGC
TTCTGATTCTTACAACCTTCCGTAATTCCAGGCTTGTGGCCCCAAATTCAGG
GCCCCACCCTTCCAGGAACAAATCATTATAGTAATAATTTGCCTTCATCTTCC
ATATACCAACTAAGCATGTTTAACTACGAACGTCCAAAACACTTCATCCAGTC
CCAAAACCCATGTGGCTCCAGATTGCAGCCTCCTGGACCAGAAACCTCCAGCT
TCTCTAGCCAGACCAACAGTCTTCCATTATCATCCAGCCCCGCCAGTGTACA
GAGCAAAGATTTTCTGCCTCCTCAATACTGAGCTCTCACATCACCATGTCCTC
CTCTGCTTTCCCTGCTTCTCCCAAGCAGCATGCTGGCTCCAACCCAGGCCAAA
GGGTTACAACCACCTATAACCAGTCCCCAGCCAGCTTCCTCAGCTCCATATTA
CCATCACA592GCCTGATTACAATAGCAGTAAAATCCCTTCCGCTATGGATTCT
CAACTATCAACAGTCCTCAGCTGGCCAACCTATAAATGCAAAGCCATCCCCAAA
CTGCAAATGCTAAGCCCATACCAAGAACTCCTGATCATGAAATACAAGGATCA
AAAGAAGCTTTGATTCAAGATTTGGAAAGAAAGCTGAAATGCAAGGACACCTT
TCTTCATAATGGAAATCAACGTCTAACATATGAAGAGAAGATGGCTCGCAGAT
TGCTAGGACCACAGAATGCAGCTGCTGTGTTTCAAGCTCAGGATGACAGTGGT
GCACAAGACTCGCAGCAACACAACCTCAGAACATGCGCGACTGCAAGTTCCTAC
ATCACAAGTAAGAAGTAGATCAACCTCAAGGGGAGATGTGAATGATCAGGATG
CAATCCAGGAGAAATTTTACCCACCACGTTTCATTCAAGTGCCAGAGAACATG
TCGATTGATGAAGGAAGATTCTGCAGAATGGACTTCAAAGTGAGTGGACTGCC
AGCTCCTGATGTGTCATGGTATCTAAATGGAAGAACAGTTCAATCAGATGATT
TGCACAAAATGATAGTGTCTGAGAAGGGTCTTCATTCATCATCTTTGAAGTA
GTCAGAGCTTCAGATGCAGGGGCTTATGCATGTGTTGCCAAGAATAGAGCAGG
AGAAGCCACCTTCACTGTGCA1291GCTGGATGTCCTTGCAAAAGAACATAAA
AG1321AGCACCAATGTTTATCTACAAACCACAGAGCAAAAAAGTTTTAGA13
66GGGAGATTCAGTGAACTAGAATGCCAGATCTCGGCTATACCTCCACCAAA
GCTTTTCTGGAAAAGAAATAATGAAATGGTACAATTCAACACTGACCGAATAA
GCTTATATCAAGATAACACT1490GGAAGAGTTACTTTACTGATAAAAGATGT
AAACAAGAAAGATGCTGGGTGGTATACTGTGTCAGCAGTTAATGAAGCTGGAG
TGACTACATGTAACACAAGATTAGACGTTAC1603GGCACGTCCAAACCAAAC
TCTTCCAGCTCCTAAGCAGTTACGGGTTCGACCAACATTCAGCAAAATTTTAG
CACTTAATGGGAAAGGTTTGAATGTAAACAAGCTTTTAAACCCAGAAGGAGAA
TTTCAGCGTTTGGCAGCTCAATCTGGACTCTATGAAAGTGAAGAACTTTAATA
ACTTTACCAACATTGGAAAACAGCCAACTACACCATTAGTAATATATTTGATT
ACATTTTTTTTGAATTAATCCATAGCTGTATTAACAGATTATGGTTTTAATTA
GGTAATATAGTTAATATATATTTATAATATTATTTATCCTTTGACTCTTGAC
ATTCTATGTACCCCTCCGATTTGTGAAGCCTACAGGAAATCTGGGTATATGGA
TTTGTAAGTGCAGAAGACTATCTTAAAATACAGGATTTTAAACATTTAAGTCAT
GCACATTTAACAATTACAGGTTATAAATTAGTATCAACTTTTTTAAACACATCT
AATGCTTGTAATAACGTTTACTGGTACTGCTTTCTAAATACTGTTTTTACCCGT
TTTCTCTTGTAGGAATACTAACATGGTATAGATTATCTGAGTGTTCCACAGTT
GTATGTCAAAGAAAATAAAATTCAAATATTTAAAACGGA

2/3

Figure 2




Binds MYOT sequences 1321-1342 from Genbank file: AF144477.1



Binds MYOT sequences 1366-1387 from Genbank file: AF144477.1

3/3

Figure 3**A****B**