

12

**EUROPEAN PATENT APPLICATION**

21 Application number: 85306215.6

51 Int. Cl.4: **C 11 D 3/386**

22 Date of filing: 02.09.85

30 Priority: 12.09.84 US 650617

43 Date of publication of application:  
09.04.86 Bulletin 86/15

84 Designated Contracting States:  
AT BE CH DE FR GB IT LI LU NL SE

71 Applicant: **The Clorox Company**  
**1221 Broadway**  
**Oakland California 94612(US)**

72 Inventor: **Munk, Barbara H.**  
**809 Curtis**  
**Albany California 94706(US)**

74 Representative: **Ford, Michael Frederick et al,**  
**MEWBURN ELLIS & CO. 2/3 Cursitor Street**  
**London EC4A 1BQ(GB)**

54 **Pre-wash compositions containing enzymes.**

57 Compositions effective in removing protein, fat and/or starch based stains as well as oil and grease based stains from fabrics comprise organic solvents admixed with surfactants and an enzyme component wherein the enzymes are dissolved in aqueous solution and the aqueous solution is, in turn, dispersed as reversed micelles within the solvent/surfactant mixture. Encapsulating the enzymes in reversed micelles protects them from degradation by the solvents and thereby prolongs the effectiveness of the enzymes for extended periods of time. The total water content of the compositions is kept below 5 weight% so that the compositions' oil and grease stain removal ability remains unimpaired.

EP 0 177 183 A1

Pre-wash Compositions Containing Enzymes

Modern home laundry cleaning compositions are very reliable and efficient in removing soil from household clothing articles, linens, and the like. Household laundry detergent compositions are formulated to remove soil and stains from various kinds of dirt depositions in the fabric. Generally such formulations effectively remove dirt induced soil and are even effective in removing low level stains or soil from oil or grease sources. However, common laundry detergents are deficient in handling soil resulting from protein sources, e.g., blood, grass; or soil where the proteins are combined with oils or greases from animal or vegetable origin; or soils of heavy grease, fats, or starch origin.

In order to effectively remove these difficult soils, various compositions are being marketed as "pre-washes" or "pre-spotters." Such compositions are applied directly to the difficult stains before the washing process. The "pre-wash" compositions are formulated primarily to remove oily or greasy stains. For this purpose, such compositions rely principally upon organic based solvents such as hydrocarbons, halogenated hydrocarbons etc. in combination with

hydrocarbon compatible surfactants. Such compositions effectively dissolve and/or emulsify oily or greasy stains. These compositions, however, are not nearly as effective in removing stains caused by protein sources  
5 such as blood, and grass; or from combined protein and fat sources such as sebum; or from fats and starches. These sources produce stains which are difficult to remove with solvent/surfactant combinations alone.

In an effort to treat protein, fat and starch  
10 based stains, some producers have resorted to the use of enzyme compositions that are available from various sources. These compositions employ protease, amylase, or lipase enzymes to attack protein, fat and starch based stains and chemically degrade these compounds so  
15 that they can more readily be removed by subsequent or concurrent treatment with conventional surfactants. While enzyme containing compositions are quite proficient in treating such stains, the enzymes themselves are quite susceptible to deactivation if  
20 mixed with other components, especially hydrocarbon solvents.

Thus the "pre-wash" manufacturer faces a dilemma if it is desired to effectively treat both oil and grease based stains as well as protein, fat and/or  
25 starch based stains on the same garments. Since enzymes are incompatible with hydrocarbon solvents, it

has heretofore been impractical to devise a single enzyme-hydrocarbon solvent product that will effectively treat both grease stains and protein, fat and/or starch stains. It is not practical to provide two separate formulations, one enzyme and one solvent, since separate compositions would be more costly; and cause twice the work for the user. Therefore such separate treatment procedures are unacceptable to the consumer.

10           The ideal solution to the problem would be a "pre-wash" composition which would simply and easily treat both grease and protein, fat, and/or starch stains in a single treatment. Such a composition should have a reasonable storage life during which the enzymes would not excessively degrade or deteriorate; it should be quite liquid so that it could be easily applied to the stains, as by spraying; it should be a single phase composition, so that one component would not settle out thereby requiring agitation or shaking to mix the components before use; and it should be dispersible in, or miscible with laundry wash water so that it is easily removed when the fabrics are subsequently washed in a conventional washing machine.

25           The present invention solves the problems enumerated above.

The present invention relates to laundry "pre-wash" compositions; and more particularly to pre-wash compositions that are single phase liquids effective against both oil and grease stains as well as protein, fat and/or starch stains. The invention compositions effectively combine the advantages of both hydrocarbon oil and grease solvents and enzymes in a single liquid composition. The normally incompatible hydrocarbon solvents and enzymes are combined without expressively limiting the products' "shelf-life", i.e., the enzyme component retains an effective activity against stains for extended periods; and the hydrocarbon solvents retain their effectiveness against oil and grease stains.

The enzymes are water soluble and they must remain in aqueous solution to retain their activity against proteins, fats, and/or starches. The hydrocarbon solvents, on the other hand are quite hydrophobic and immiscible in aqueous solutions. The hydrocarbon solvents are effective against oil and grease in the absence of water. Any water present in the "pre-wash" composition tends to interfere with the solvents' ability to remove oil and grease.

The present compositions reconcile these adverse properties by dispensing aqueous solutions of

the enzymes as "reversed micelles" within the hydrocarbon solvent/surfactant medium. The "reversed micelles" are formed by providing suitable surfactants which facilitate encapsulation of an aqueous solution  
5 of the enzymes and subsequently dispersing the enzyme/surfactant mixture in the hydrocarbon solvent medium. The enzymes are dissolved in the aqueous solution; surfactants are added. The surfactants encapsulate the enzymes, and then permit the dispersion  
10 of the enzyme-aqueous portion in the hydrocarbon medium.

In other words, the surfactants effectively facilitate the formation of reversed micelles. In such reversed micelles the hydrophobic tail of the  
15 surfactant dissolves in the hydrocarbon medium, while the hydrophilic head of the surfactant dissolves in the aqueous enzyme solution. Under appropriate conditions minute micelles are formed within the hydrocarbon medium. The enzymes are dissolved within the aqueous  
20 micelle interior where they are effectively protected from attack and degradation by the hydrocarbon solvent.

When the "pre-wash" composition is applied to a fabric, the hydrocarbon component attacks and removes the oil and grease stains without interference from the  
25 aqueous portion of the composition since the aqueous portion is but a minor amount, 5% or less, of the

composition. At the same time, the enzymes are released upon the fabric surface where they can effectively attack the protein, fat, and/or starch stains.

The aqueous enzyme-surfactant "reversed-  
5 micelles" are extremely small and are dissolved within the hydrocarbon medium to form a suspension or dispersion. Thus there is no problem with phase separation. Further, due to the unique fine structure, i.e., reversed micelles, the enzymes in the aqueous  
10 portion are protected from premature degradation by the surrounding hydrocarbon medium.

From a component standpoint the invention compositions comprise a small percentage (by weight) of the enzymes; somewhat larger percentages of enzyme  
15 stabilizers, e.g., sodium chloride, calcium chloride, or triethanolamine, all of which are dissolved in aqueous solution with the total water content being less than 5% of the composition; substantial percentages of one or more surfactants, preferably  
20 nonionic; and a large percentage of hydrocarbon solvent. In addition, small percentages of hydrotropes, such as glycols may also be present. Small amounts of perfumes or dyes may be included for aesthetic purposes.

25 The pre-wash compositions are prepared in a prescribed procedure so as to produce a solution of

reversed micelles having the aqueous enzyme solution held within the interior thereof; and with the micelles dispersed throughout the hydrocarbon solvent-surfactant medium.

5           In general, the compositions may comprise several tenths of a percent of solution (as a glycol/water solution) enzyme; about a tenth of a percent of NaCl (as enzyme stabilizer); less than five percent water; about twenty-five to thirty-five percent  
10 surfactant; about sixty to seventy percent hydrocarbon solvent; about two to three percent of a glycol as a hydrotrope; and if desired, several tenths of percent perfume and coloring material (dye).

          In a preferred composition the water  
15 dissolved enzyme may comprise about 0.2 weight%; about 1.5 weight% of 1M NaCl aqueous solution; about 2.5 weight% of ethylene glycol; about 14.5 weight% of an ethoxy-propyloxy linear alcohol nonionic surfactant; about 14.5 weight% of an ethoxylated linear alcohol  
20 nonionic surfactant; with the remainder being hydrocarbon solvent, e.g., a nominal C-12 to C-16 mixture of isoparaffins. Very small percentages of perfumes or coloring agents may be added.

          It will be understood that the enzyme compo-  
25 nent is present to effectively remove protein based stains, fat based stains, and/or starch based stains.

Of these types of stains the most important are the protein based stains resulting from blood, grass, body fluids, and the like. Therefore, proteases are most desirable in the invention compositions. It should also  
5 be apparent that related enzymes having particular effectiveness against other organic molecules, e.g., starches, fats, etc., may also be included along with the proteases. If desired, amylases, lipases etc. may be combined with the protease enzymes to produce a  
10 pre-wash composition that is effective against a full spectrum of fabric stains. The only precaution to be taken if additional enzymes are introduced, is that such added enzymes be compatible with the protease.

It is most important that the total amount of  
15 water in the compositions be maintained at levels no greater than 5 weight%. Quantities of water greater than 5 weight% may interfere with the ability of the hydrocarbon solvent to effectively remove oil and grease stains.

20 It is an object of the invention to provide improved pre-wash compositions.

It is another object of the invention to provide compositions that are useful for removing both oil and/or grease based stains and protein, fat, and/or  
25 starch based stains from fabrics.

It is another object of the invention to provide compositions wherein enzymes in aqueous solution are encapsulated in surfactants dispersed within a surfactant-hydrocarbon solvent medium to form  
5 reversed micelles therein.

It is still another object of the invention to provide compositions including hydrocarbon solvents, surfactants, and enzymes as active ingredients and wherein the enzymes are protected from the hydrocarbon  
10 solvent by encapsulating the enzymes within surfactants and then dispersed in the solvent to form reversed micelles.

It is yet another object of the invention to provide a method for encapsulating enzymes in aqueous  
15 solution within reversed micelles dispersed in a surfactant-hydrocarbon solvent medium.

Other objects and advantages of the invention will become apparent from the following specification and claims.

Pre-wash compositions that exhibit the ability to simultaneously remove oil/grease stains and protein, fat, and/or starch stains from fabrics are formulated from surfactants, enzymes, and hydrocarbon solvents. While enzymes and hydrocarbon solvents are normally incompatible, the invention compositions overcome this difficulty by enclosing the enzymes in nonionic surfactants, and then dispersing the mixture in hydrocarbon solvents. Reversed micelles are thus formed in the surfactant-hydrocarbon medium. The water content of the compositions is kept very low, i.e., no more than 5 weight%, so that the oil and grease stain removing ability of the hydrocarbon solvent substrate remains unimpaired. At the same time the encapsulated enzymes are released to attack protein stains when the pre-wash composition is applied to soiled fabrics.

More specifically, the pre-wash compositions comprise a hydrocarbon solvent medium in which surfactants are dispersed, as well as an aqueous enzyme solution in the form of reversed micelles. The dispersed surfactants form the "micelle" interface between the aqueous enzyme solution and the surrounding hydrocarbon solvent medium. The hydrocarbon solvent comprises the major component, ideally being present in amounts greater than 60% by weight. The surfactants

comprise the next most abundant component, ideally being present in an aggregate amount in the neighborhood of 30%. It is desirable that the solvent-surfactant ratio be maintained at about 2:1, e.g., if the solvent is about 60%, then the surfactants should be present at about 30%. The total amount of solvent and surfactant is not critical; however, together they should comprise well over 90% of the composition.

10           The component that is critical to the effectiveness of the pre-wash compositions is water. Water comprises the solvent for the enzymes and its presence is necessary for that purpose. However, it has been determined that if water is present in quantities  
15 greater than about 5% by weight, the oil and grease removal efficiency of the hydrocarbon solvent is adversely affected. When the water content rises appreciably above 5%, there is a noticeable decrease in the solvents' ability to thoroughly remove oil and  
20 grease stains. Therefore, the water content is always kept at 5% or less, preferably less.

          The water acts as a solvent for the enzyme component, which is necessary to remove protein, fat and/or starch based stains. The enzymes are dissolved  
25 in the water. The enzymes are stabilized in the aqueous solution by the addition of salt (NaCl). Only

a small percentage, e.g., several (i.e. 1, 2, 3, 4, 5 or 6) tenths of a percent of the enzyme solution is needed to be effective. Higher amounts of enzymes may also be utilized, but will add to the cost of materials.

5 In commercially available enzyme solutions having an enzyme activity equivalent to 8 K.N.P.U. (K.N.P.U. means Kilo Novo Protease Units. There is no industry-wide standard for measuring activity.), about 0.2% by weight is an effective amount. Such enzyme solutions may be

10 further stabilized by the addition of 1-2% of a 1M NaCl solution.

A hydrotrope, such as ethylene glycol can be added in low percentages, e.g., 2-3%, to aid the solubility of the surfactants in the solvent.

15 It will be understood that, except for the limit on water in the compositions, the percentages of all components may be varied over relatively wide ranges, i.e., there is nothing critical about the hydrocarbon solvent, surfactants, enzyme, salt or

20 glycol percentages. On the other hand, experimentation has developed a preferred composition having the following percentages of components:

<u>Component</u>	<u>Wt%</u>
Enzyme (Savinase 8.0L) (Enzyme dissolved in a solution of 80% propylene glycol/20% H <sub>2</sub> O)	0.2
5 NaCl (1M in water)	1.5
Ethylene glycol	2.5
Nonionic surfactants (50-50 mixture of ethoxylated linear alcohols and ethoxylated- propoxylated linear alcohols)	29.5
10 Hydrocarbon solvent (isoparaffins) having nominal 12 to 16 carbons)	66.0
Perfume and dye	remainder.

15                   When produced in accordance with the  
procedure set forth below, the compositions are a clear  
solution with the enzymes encapsulated in reversed  
micelles within the surfactant-hydrocarbon solvent  
medium. The compositions exhibit excellent activity  
20 against both oil and grease stains as well as protein,  
and/or starch and/or fat-based stains. The  
compositions also retain an appreciable portion of the  
enzyme activity when stored for extended periods.

25                   The selection of the enzymes for use in the  
compositions may be made from any number of commercial-  
ly available liquid enzyme solutions that are useful  
against proteins, lipids, and starchy substances. Such  
enzyme solutions are available from several commercial  
sources.

For instance, the commercially available alkaline proteases (preferred in the invention) are derived from various strains of the bacterial *Bacillus subtilis*. These proteases are also known as  
5 subtilisins, and are available under the trademarks Esperase ®, Savinase ®, and Alcalase ®, from Novo Industri A/S, of Bagsvaerd, Denmark; and also under the trademarks Maxatase ® and Maxacal ® from Gist-Brocades N.V of Delft, Netherlands.

10           These commercially available proteases are supplied as aqueous stabilized solutions of the enzyme. The enzymes are generally stabilized by the addition of glycols such as propylene glycol. These solutions are also supplied in various strengths wherein the  
15 strengths are defined by the activity exhibited by the enzyme. Thus as noted above, in the preferred composition the enzyme solution has an activity of 8 K.N.P.U. Of course, enzyme solutions of lesser or greater activity can be utilized in the invention compositions.

20           It should be noted that other enzymes may be used in the compositions in addition to, or in place of, the proteases. Thus lipases effective against fats; or amylases, effective against starches, can also be used in the formulations. Both types of enzymes are  
25 commercially available, e.g. lipases - (see U.S. Patent 3,950,277 column 3, lines 15-55 for a description of

lipase enzymes, their origins, and sources -  
incorporated herein by reference) amylases; - Rapidase  
• from Societa Rapidase, France; and Milezyme • from  
Miles Labs., Elkhurst, Ind.

5                   It has been determined that the addition of  
salt, NaCl, to the enzyme solutions will further  
stabilize the enzymes against degradation in the  
presence of the hydrocarbon solvents of the pre-wash  
compositions. To this end, the compositions also  
10 include aqueous salt solution as stabilizing agent.  
While the exact amounts of the stabilizer is not  
critical, the inclusion of roughly 1.5 weight% of 1M  
NaCl to the aqueous enzyme solution has been found to  
aid in maintaining enzyme activity when the product is  
15 stored.

Surfactants comprise a major component of the  
pre-wash compositions. Surfactants that are compatible  
with the hydrocarbon solvent medium are most necessary;  
and in addition such surfactants must not degrade, or  
20 interfere with the enzymes in the reversed micelles.  
The surfactants are also responsible for the micelle  
formation within the hydrocarbon medium. Nonionic  
surfactants are ideal for the above stated purposes.  
The long chain alcohols such as linear ethoxylated and  
25 linear propoxylated alcohols and mixtures thereof are  
particularly useful in the invention compositions.

Such surfactants are completely compatible with the hydrocarbon solvents; they efficiently form reversed micelles to encapsulate the aqueous enzyme solutions; they do not degrade the enzymes; and they contribute  
5 significantly to the removal of soil from fabrics to which the compositions are applied.

Alkyl ethoxylated and propoxylated alcohols in the nominal C-12 to C-16 range are most preferred. Such surfactants are available as standard articles of  
10 commerce under the name "Biosoft" from the Stepan Chemical Co.; or under the name "Neodol" from the Shell Chemical Co. The "Biosoft" series of nonionic surfactants are ethoxylated and propoxylated fatty alcohols sold in liquid form. The "Neodol" series of  
15 surfactants comprise a large group of nonionic surfactants including ethoxylated long chain alcohols with ethoxy groups ranging from 3 to 12 and the carbon chains from 12 to 15.

All such nonionic surfactants are useful in  
20 the invention compositions, although Neodol 25-3 is preferred. Neodol 25-3 is composed of carbon chains nominally in the C-12 to C-15 range with an average of three ethoxy moieties per mole of alcohol. Biosoft EA-10 is a mixture of ethoxylated and propoxylated long  
25 chain fatty alcohols wherein the carbon chains are

nominally in the C-10 to C-12 range with an average of 7.1 ethoxy and 2 propoxy moieties per mole of alcohol.

Other similar nonionic surfactants can be substituted for the aforementioned surfactants in the compositions so long as they met the criteria set forth above.

The surfactants are included in the compositions in substantial quantities making up somewhat less than one-third of the weight. While the total surfactant percentage is not critical, they should be present in roughly one-half the amount of hydrocarbon solvent. It has been found that this ratio of surfactant to solvent is necessary to achieve good removal of oil and grease based soil wherein both the hydrocarbon solvent and surfactants play key roles.

The hydrocarbon solvent is the major component in the compositions and, in conjunction with the surfactants, is the primary agent for treating oil and grease based stains. The hydrocarbon solvent comprises well over half the compositions by weight; generally the percentages are in the 60-70% range.

The solvent component can be selected from any number of hydrocarbon based oil and grease solvents. Such materials are staple industrial products and may be procured from a number of oil industry sources.

It is necessary however that the solvent component be compatible with the formation of micelles, in this instance reversed micelles, i.e., micelles having an encapsulated aqueous component dispersed within an organic solvent medium. Because of this requirement it is desirable to employ solvents that are low in sulfur, acids, and oxygenated compounds. Pure hydrocarbon solvents, especially paraffinic hydrocarbons having fairly long carbon chains, e.g., C-10 to C-14 are highly preferred for use in the compositions. Such solvents are available from the Exxon Corporation of Houston, Texas under the trademark "Isopar" and "Norpar". Both of these groups of hydrocarbon solvents have a very high ( 98%) normal or isoparaffin content and very low concentrations of sulfur, acids, carbonyls, chlorides, etc. These solvents, or their equivalents from other manufacturers, are the preferred organic solvents for use in the compositions.

In order to produce the compositions wherein the aqueous enzyme component is protected from degradation by the organic solvents, i.e., encapsulation within reversed micelles, it is necessary to proceed according to the following method:

The desired amount of aqueous salt solution is prepared. The aqueous/glycol enzyme solution as

procured from the manufacturer is then mixed into the salt solution. (Gentle agitation is used throughout the production process to thoroughly mix all components. Violent agitation is to be avoided as it may actually degrade the enzymes by a physical shearing of the enzyme molecules.)

A portion, perhaps one half, of the surfactant is then gently mixed into the aqueous enzyme/salt solution. After thorough mixing is obtained, the remainder of the surfactant is then added and thoroughly mixed in. At this stage, the aqueous enzyme/salt/surfactant mixture is a rather viscous clear liquid. The glycol component is then added with continuous mixing; after which, the hydrocarbon solvent is gradually added to form the finished product.

The final product is a clear transparent liquid, principally solvent and surfactant with small amounts ( 5%) of water, and smaller amounts of enzyme and enzyme stabilizer.

Although the preparation procedure seems very straightforward and uncomplicated, it is emphasized that the sequence of addition of components should be closely adhered to. It is particularly important to first mix the components that dissolve in the aqueous phase; secondly mix in the surfactants; and finally add the hydrophobic organic solvent component. Any

variation in the procedure, may fail to produce the desired reversed micelles and the enzymes will be unprotected from the solvent components. Failure to produce the protective micelles will be apparent if a  
5 cloudy solution is produced; or if a precipitate forms upon standing.

The clear liquid product may be packaged in any suitable container and stored for periods of several months and yet retain a good percentage of the  
10 enzyme activity. When used it is preferred to spray the product on the soiled fabrics a few minutes before placing them into a washing machine. Normal laundering procedures may be utilized. Of course, an effort should be made to spray the product directly on the  
15 visibly soiled portions of the fabric as direct contact will ensure full opportunity for attacking the oil/grease, and/or protein, fat or starch stains prior to the laundering process.

When prepared according to the above  
20 directions, the enzyme stability of the compositions as noted above is quite good. In one test over 50% of the enzymes' initial activity remained after six month's storage at 70°F.

In another group of tests loss in enzyme  
25 activity with time was studied for the following

composition made in accordance with the preferred method of the invention:

	<u>Component</u>	<u>Wt. %</u>
	Savinase 8.0L	0.2
5	1M NaCl (aq.)	1.5
	Ethylene glycol	2.5
	Biosoft EA-10	14.65
	Neodol 25-3	14.65
	Isopar L	66.3
10	Perfume	0.2

Samples of the above composition were then stored at various temperature for a period of six months. Table I below sets/forth the results of tests on enzyme activity at the end of the six month period.

15

TABLE I

Enzyme Activity After Six Months Storage

	Storage temperature	Wt. % active enzyme	% activity* remaining
	35°F	0.17	85
20	70°F	0.11	55
	100°F	0.02	10

\*assumes 0.2 wt. % active enzyme in original sample

A series of studies were carried out to evaluate the efficacy of the enzyme containing pre-wash compositions of the invention.

In a first series, a number of compositions were prepared wherein the enzyme content was varied from 0 up to 1.0%. These compositions were then tested for effectiveness against various types of stains, i.e., grass, ballpoint pen ink, dirty motor oil, and azocasein. Laundry detergent and a commercial pre-wash formulation, i.e., Clorox pre-wash, were, in each instance, used for comparison purposes. Table II below sets forth the compositions. Tables III, IV and VII set forth the results of % stain removal as measured by instrument. Tables V, VI, and VIII set forth the results measured by visual grading (on a scale of 1-5, with 5 highest, 1 lowest).

TABLE II  
Formula Composition

<u>Composition</u> <u>(Wt%)</u>	<u>Formula Composition</u>		
	<u>Formula</u> <u>A</u>	<u>Formula</u> <u>B</u>	<u>Formula</u> <u>C</u>
Savinase 8.0L	0-1	0-1	0-1
Salt Water (1M NaCl)	5	5	5
Biosoft EA-4	14-15	13.3-15	13-15
Biosoft EA-10	14-15	13.3-15	13-15
Petro 22*	0	0	1
Ethylene Glycol	0	2.5	2.5
Isopar L	65.3-66.3	65-66.3	64.3-66.3
Perfume	0.2	0.2	0.2

\* 1 alkyl naphthalene sulfonate, Na salt (wetting agent).  
The control prewash is a commercial product with no enzymes and utilized in these tests for comparison purposes.

TABLE III  
PERFORMANCE TESTING  
% STAIN REMOVAL  
AZOCASEIN<sup>1</sup>

FABRIC Basic Formula	COTTON			POLYESTER		
	A	B	C	A	B	C
Wt. % Enzyme						
0	13.6	10.8	6.6	15.0	11.7	12.3
0.1	41.5	36.2	37.9	43.9	45.2	48.2
0.2	39.5	32.4	49.8	47.8	41.2	55.2
0.4	53.8	45.2	46.3	51.5	55.4	57.2
0.8	56.2	59.8	54.5	57.6	59.6	54.7
1.0	55.5	43.0	55.4	58.7	40.7	62.4
Clorox Pre-wash Detergent		17.2			15.7	
		10.3			6.7	
LSD*		17.2			15.8	

\* LEAST SIGNIFICANT DIFFERENCE  
 1 "azocasein" is a model stain designed to measure enzyme activity.

TABLE IV  
PERFORMANCE TESTING  
& STAIN REMOVAL  
GRASS

FABRIC BASIC FORMULA Wt. % Enzyme	COTTON			POLYESTER		
	A	B	C	A	B	C
0	56.8	64.8	58.4	64.2	64.2	59.6
0.1	90.9	92.3	90.0	89.9	93.6	92.6
0.2	91.7	92.6	93.5	92.9	94.5	96.2
0.4	93.5	94.5	92.2	96.0	95.6	95.5
0.8	92.2	94.4	93.8	95.3	97.6	96.1
1.0	92.6	93.3	94.0	96.1	97.1	97.0
Clorox Pre-wash Detergent		56.6 42.2			60.3 62.7	
LSD		4.5			5.8	

TABLE V  
PERFORMANCE TESTING  
VISUAL GRADING  
GRASS

FABRIC BASIC FORMULA Wt. % Enzyme	COTTON			POLYESTER		
	A	B	C	A	B	C
0	1.5	1.5	1.6	1.0	1.1	1.0
0.1	3.4	3.8	3.2	1.8	2.1	2.0
0.2	3.9	3.6	4.1	2.2	2.3	2.3
0.4	4.1	4.2	4.1	2.7	2.4	2.4
0.8	4.0	4.1	4.3	2.7	2.7	2.8
1.0	4.0	3.6	4.1	2.9	2.9	3.1
Clorox Pre-wash Detergent		1.4 1.3			1.0 1.0	
LSD		0.2			0.2	

TABLE VI

PERFORMANCE TESTING  
VISUAL GRADING  
BALLPOINT INK

FABRIC BASIC FORMULA Wt. % Enzyme	COTTON			POLYESTER		
	A	B	C	A	B	C
0	3.1	2.4	2.6	1.1	1.4	1.1
0.1	2.0	2.7	2.5	1.3	1.1	1.2
0.2	2.8	2.3	3.4	1.2	1.2	1.1
0.4	2.9	3.1	3.0	1.1	1.0	1.4
0.8	1.7	3.3	2.3	1.1	1.1	1.2
1.0	3.2	2.5	2.6	1.5	1.0	1.5
Clorox Pre-wash Detergent	3.3	1.0		2.0	1.1	
LSD		0.4			0.1	

TABLE VII  
PERFORMANCE TESTING  
& STAIN REMOVAL  
DIRTY MOTOR OIL

FABRIC BASIC FORMULA Wt. % Enzyme	COTTON			POLYESTER		
	A	B	C	A	B	C
	0	92.8	94.2	92.3	78.7	82.1
0.1	93.6	92.1	92.3	84.2	71.0	75.3
0.2	93.6	93.4	92.2	84.7	79.3	80.5
0.4	94.9	94.7	92.3	82.0	78.3	83.1
0.8	93.4	93.6	92.5	84.1	74.8	70.8
1.0	93.7	85.4	93.0	84.0	75.5	81.6
Clorox Pre-wash Detergent		94.5 47.1			85.4 6.3	
LSD		3.8			9.2	

TABLE VIII

PERFORMANCE TESTING  
VISUAL GRADING  
DIRTY MOTOR OIL

FABRIC BASIC FORMULA Wt. & Enzyme	COTTON			POLYESTER		
	A	B	C	A	B	C
0	3.9	3.7	3.6	2.7	1.8	3.0
0.1	3.6	3.7	3.9	2.7	2.1	2.2
0.2	3.5	3.5	3.8	2.3	2.0	2.2
0.4	3.8	3.8	3.6	2.8	2.5	2.5
0.8	3.4	3.5	3.2	2.7	2.4	1.6
1.0	3.8	3.9	3.3	2.5	2.2	2.3
Clorox Pre-wash Detergent		4.0 1.4			2.3 1.0	
LSD		0.3			0.4	

A review of the data in Tables III, IV, V, VI, VII, and VIII above will make it apparent that as little as 0.1% added enzyme produces a significant increase in the ability of the compositions to remove  
5 azocasein and grass stains.

In a second series of tests, another group of invention compositions were prepared and tested along with Clorox pre-wash and laundry detergent against a broad range of stains. In this second series, the  
10 enzyme content was kept constant at 0.2%. Table IX sets forth the various compositions. Tables X, XI, and XII set forth the results when the formulations were tested against the indicated stains on 100% cotton fabric, 65/35 polyester/cotton fabric, and 100%  
15 polyester fabric.

TABLE IX

Formula Composition

	Formula 1	Formula 2	Formula 3	Formula 4
	wt%	wt%	wt%	wt%
Savinase 8.0L	0.2	0.2	0.2	0.2
Salt Water (1MNaCl)	1.5	3.0	1.5	3.0
Water	0	0	0	0
Ethylene Glycol	2.5	2.5	0	0
Petro 22	0	0	0	0
Biosoft EA-4	14.65	13.9	15.0	15.0
Biosoft EA-10	14.65	13.9	15.0	15.0
Isopar L	66.3	66.3	68.1	66.6
Perfume	0.2	0.2	0.2	0.2

TABLE X  
PERFORMANCE EVALUATION  
100% COTTON

INSTRUMENTAL GRADING	FORMULA 1		FORMULA 2		FORMULA 3		FORMULA 4		CLOROX PRE-WASH		DETERGENT		LSD
	HOT	WARM	HOT	WARM	HOT	WARM	HOT	WARM	HOT	WARM	HOT	WARM	
Chocolate Ice Cream	91.2	91.3	91.2	92.0	92.1	92.6	91.2	91.9	90.9	92.0	90.6	90.5	1.1
Axocasein	36.1	33.8	42.0	31.9	41.1	37.8	39.9	34.1	-0.5	-1.0	4.9	-3.0	4.8
Egg	99.2	99.2	98.9	98.7	99.1	98.8	99.1	98.8	98.8	98.6	98.3	98.7	0.4
Grass	95.2	93.2	95.7	93.5	95.4	95.6	95.0	93.1	67.9	65.4	56.9	59.0	2.3
Spaghetti Sauce	94.2	91.9	92.7	91.8	92.8	93.8	92.2	91.9	89.4	89.3	86.1	86.3	1.3
Dirty Motor Oil	94.8	94.6	94.6	92.5	93.9	94.4	93.4	93.9	93.4	93.2	42.7	41.6	2.3
Blood	98.2	98.5	98.2	98.4	98.0	98.0	97.8	97.8	94.6	97.0	97.0	96.8	0.6
VISUAL GRADING													
STAINS:													
Grass	4.5	4.1	4.4	4.3	4.4	4.4	4.2	4.3	2.4	2.4	1.8	2.1	0.2
Spaghetti Sauce	3.5	3.3	3.6	3.6	3.5	3.8	3.5	3.7	2.9	3.2	2.6	2.6	0.2
Dirty Motor Oil	4.2	4.4	4.4	4.2	4.3	4.4	4.3	4.2	4.2	3.9	1.1	1.2	0.2
Blood	4.1	4.3	4.3	4.2	4.3	4.1	4.0	4.1	2.8	3.7	3.8	3.8	0.2
Ballpoint Ink	3.0	2.4	3.2	2.9	2.6	2.2	2.2	2.5	2.9	3.7	1.5	1.4	0.2

TABLE XI  
PERFORMANCE EVALUATION  
65/35 POLYESTER/COTTON

INSTRUMENTAL GRADING	FORMULA 1		FORMULA 2		FORMULA 3		FORMULA 4		CLOROX PRE-WASH		DETERGENT		LSD
	HOT	WARM	HOT	WARM	HOT	WARM	HOT	WARM	HOT	WARM	HOT	WARM	
	STAINS:		STAINS:		STAINS:		STAINS:		STAINS:		STAINS:		
Chocolate Ice Cream	99.4	99.4	99.2	99.4	99.4	99.2	99.4	99.2	98.5	99.0	98.2	98.3	0.3
Azocasein	33.4	25.4	35.1	27.9	32.5	26.8	31.2	24.9	4.5	3.5	1.5	0.	3.0
Egg	99.4	99.3	99.2	99.2	99.2	98.9	99.0	98.9	99.0	98.9	98.6	97.4	1.0
Grass	96.2	95.2	95.1	94.6	92.4	94.6	95.1	91.6	75.7	77.7	63.8	72.8	2.3
Spaghetti Sauce	97.9	97.0	97.5	96.8	96.7	95.8	96.6	95.4	94.2	95.0	94.2	92.4	1.1
Dirty Motor Oil	92.2	92.1	92.3	92.3	92.6	91.7	92.9	91.4	91.7	91.0	14.3	15.9	1.4
Blood	99.8	99.8	99.7	99.6	99.8	99.7	99.7	99.6	99.1	99.5	98.9	99.2	2.8
Grass	4.4	3.8	4.2	4.1	3.6	4.0	4.3	3.6	1.6	1.6	1.4	1.4	0.2
Spaghetti Sauce	4.7	4.4	4.8	4.3	4.6	4.7	4.5	4.3	4.0	4.0	3.8	3.7	0.2
Dirty Motor Oil	4.1	4.3	4.4	4.2	4.4	4.5	4.5	4.3	4.1	4.1	1.0	1.0	0.2
Blood	5.0	4.9	4.9	5.0	4.9	4.9	4.9	5.0	4.7	4.9	4.9	4.9	0.1
Ballpoint Ink	1.8	2.2	1.8	2.4	1.7	1.9	1.9	2.0	2.0	2.7	1.1	1.1	0.2

VISUAL GRADING  
STAINS:

TABLE XII  
PERFORMANCE EVALUATION  
100% POLYESTER

INSTRUMENTAL GRADING	FORMULA 1		FORMULA 2		FORMULA 3		FORMULA 4		CLOROX PRE-WASH		DETERGENT		LSD
	HOT	WARM	HOT	WARM	HOT	WARM	HOT	WARM	HOT	WARM	HOT	WARM	
	STAINS:		STAINS:		STAINS:		STAINS:		STAINS:		STAINS:		
Chocolate Ice Cream	99.6	99.5	99.6	99.6	99.5	99.8	99.7	99.8	99.6	99.7	99.5	99.6	0.2
Azocasein	37.6	31.3	48.4	35.6	39.2	28.8	43.4	36.3	5.3	5.3	4.7	5.9	3.9
Egg	94.9	94.1	95.8	95.4	94.9	95.5	95.2	95.1	96.0	94.3	99.3	99.5	1.0
Grass	98.3	96.7	98.4	97.6	97.2	95.2	97.6	97.0	75.3	70.2	69.3	72.5	3.3
Spaghettl Sauce	98.6	95.5	97.2	96.1	95.6	95.3	97.5	96.7	96.7	97.5	96.0	92.6	1.8
Dirty Motor Oil	90.8	89.3	83.7	85.3	92.0	90.9	89.0	88.5	90.5	90.5	15.3	15.5	2.0
Blood	99.7	99.6	99.6	99.4	99.5	99.7	99.5	99.4	99.6	99.6	99.6	99.6	0.1

VISUAL GRADING  
STAINS:

Grass	4.4	4.5	4.3	4.6	4.2	4.4	4.6	4.8	1.5	1.4	1.3	1.4	0.2
Spaghettl Sauce	4.9	4.3	4.5	4.2	4.4	4.2	4.7	4.2	4.4	4.4	4.3	3.4	0.3
Dirty Motor Oil	3.5	3.4	2.8	2.8	3.8	3.5	3.1	3.4	3.7	3.7	1.0	1.2	0.2
Blood	-	-	-	-	-	-	-	-	-	-	-	-	-
Ballpoint Ink	2.6	3.1	3.0	2.9	2.1	1.8	2.1	2.2	2.9	3.1	1.1	1.1	0.1

The compositions set forth in Table IX above were stored for periods of time up to 2 months at storage temperatures varying between 35° and 100°F. The samples of the compositions were tested against  
5 both grass and azocasein stains at intervals up to the 2 months stated above. Table XIII below sets forth the results, thus indicating the enzyme stability in the compositions as a function of time and storage temperature.

TABLE XIII

STABILITY TEST

Time	Formula 1			Formula 2			Formula 3			Formula 4		
	35°	70°	100°	35°	70°	100°	35°	70°	100°	35°	70°	100°
Storage temp.												
1 0		95.2%		94.6%		94.7%		91.6%				
2 1 mth	93.3	92.8	93.8	91.3	94.4	92.2	92.2	93.2	89.3	91.7	93.2	90.8
3 1½ mth	93.6	93.6	91.1	88.5	91.7	91.7	90.6	90.0	86.0	91.4	91.1	91.8
4 2 mth	88.1	87.3	83.9	86.2	83.8	89.5	80.6	84.0	79.1	84.5	88.7	82.6
5 0		3.8		4.1		4.0		3.6				
6 1 mth	3.2	3.7	3.6	3.1	3.8	3.5	2.5	3.3	2.8	2.7	3.4	3.7
7 1½ mth	3.6	4.1	3.3	3.2	3.4	3.7	2.7	2.8	2.4	3.0	3.4	3.3
8 2 mth	2.4	2.2	1.6	2.3	1.8	2.6	1.7	2.2	1.8	2.0	1.9	1.6
9 0		25.4%		27.9%		26.7%		24.9%				
10 1 mth	20.7	21.0	19.7	16.0	21.0	22.7	24.9	18.4	12.5	21.2	14.8	16.7
11 1½ mth	22.1	15.5	19.0	14.5	13.0	21.2	13.1	13.1	12.4	12.1	9.7	20.5
12 2 mth	15.0	18.8	17.2	15.0	20.3	21.4	17.2	15.1	14.9	13.4	15.3	15.2

CLOROX PREWASH: INSTRUMENTAL/GRASS 77.7±2.3%  
 VISUAL/GRASS 1.6±0.2  
 INSTRUMENTAL/AZOCASEIN 3.5±3.0%

Note: 1-4 and 9-12 were graded by instrument (% Srox); 5-8 were visually graded (1-5 scale)  
 1-8 were grass stains; 9-12 were azocasein stains.

Further studies were undertaken to evaluate physical stability in varying compositions prepared according to the invention. Table XIV below sets forth these formulations along with statements concerning  
5 their physical stability after storage at various storage temperatures.

TABLE XIV

## FORMULAS

RAW MATERIALS	wt%	wt%	wt%	wt%	wt%	wt%	wt%	wt%	wt%	wt%	wt%	wt%	wt%
Savinase 8.0L	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	1.0
Petro 22	1.0	0.0	1.0	0.0	0.0	1.0	0.0	1.0	0.0	1.0	0.0	1.0	0.0
Ethylene Glycol	0.0	0.0	2.5	0.0	0.0	0.0	0.0	0.0	0.0	2.5	2.5	0.0	0.0
Neodol 25-3	15.2	14.7	14.2	15.3	15.2	15.2	14.7	14.6	14.6	14.3	14.2	14.5	14.5
Biosoft EA-10	15.2	14.7	14.2	15.3	15.2	15.2	14.7	14.6	14.6	14.3	14.2	14.5	14.5
Isopar L	66.9	64.9	62.4	67.7	66.9	66.9	64.9	64.1	64.1	63.2	62.4	64.6	64.6
IM NaCl(aq)	0.0	0.0	0.0	1.0	1.0	1.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
Water	1.0	5.0	5.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Perfume	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2

	wt%	wt%
Savinase 8.0L	1.0	1.0
Petro 22	0.0	1.0
Ethylene Glycol	2.5	2.5
Neodol 25-3	14.2	14.1
Biosoft EA-10	14.2	14.1
Isopar L	62.9	62.1
IM NaCl(aq)	5.0	5.0
Water	0.0	0.0
Perfume	0.2	0.2

These formulas were found to be physically stable after 18 days at 35, 70 and 100°F.

	<u>wt%</u>	<u>wt%</u>	<u>wt%</u>	<u>wt%</u>
Savinase 8.0L	0.2	0.2	0.2	0.2
CaCl <sub>2</sub> (0.5M)	1.5	0.0	0.0	0.0
NaCl (0.5M)	0.0	3.0	0.75	0.0
Water	0.0	0.0	0.0	3.0
Propylene Glycol	2.5	0.0	0.0	0.0
Ethylene Glycol	0.0	2.5	2.5	2.5
Neodol 25-3	14.65	13.9	15.0	13.9
Biosoft EA-10	14.65	13.9	15.0	13.9
Isopar L	66.3	66.3	66.35	66.3
Perfume	0.2	0.2	0.2	0.2
% Enzyme Activity Remaining After 33 days at 100°F	41%	45%	43%	41%

These formulas were physically stable after 33 days at 35, 70 and 100°F.  
Enzyme stability was shown.

TABLE XV below presents some prewash formulations wherein several different enzymes were utilized. These formulations were also tested for performance on various stains.

TABLE XV

<u>RAW MATERIALS</u>	<u>A</u>	<u>B</u>	<u>C</u>
	<u>(wt%)</u>	<u>(wt%)</u>	<u>(wt%)</u>
1M NaCL	1.5	1.5	1.5
5 Ethylene Glycol	2.5	2.5	2.5
Neodol 25-3	14.63	14.51	14.65
Biosoft EA-10	14.65	14.51	14.65
Isopar K	66.3	66.14	66.3
Perfume	0.2	0.2	0.2
10 Esperase 8.0L	0.2	0.0	0.0
Alkalase 2.5L	0.0	0.64	0.0
Savinase 8.0L	0.0	0.0	0.2

These samples were evaluated for performance on grass,  
 15 blood and gravy. All three formulas provided a benefit  
 on grass and gravy. The blood stain was totally  
 removed by detergent alone. No enzyme stability test  
 was conducted.

Still another group of compositions were prepared having further variations in formulation. Table XVI below sets forth some compositions wherein additional enzyme stabilizer (triethanolamine - Formula F) was included; or wherein the solvent system (methylester - Formula G) was varied. Table XVI also indicates the residual enzyme activity after storage for the stated period.

Tables XVII, XVIII, and XIX below, present performance data on various fabrics for the compositions of Table XVI.

TABLE XVI

FORMULA COMPOSITION

<u>Raw Materials</u> <u>Wt%</u>	<u>Formula B</u>	<u>Formula F</u>	<u>Formula G</u>
Savinase 8.0L	0.2	0.2	0.2
Salt Water (1M NaCl)	1.5	1.5	1.5
Neodol 25-3	14.65	14.65	0.0
Biosoft EA-10	14.65	14.65	10.5
Petro 22	0.0	0.0	1.0
Ethylene Glycol	2.5	0.0	0.0
Isopar L	66.3	66.3	53.6
Perfume	0.2	0.2	0.0
Triethanolamine	0.0	2.5	0.0
Neodol 91-2.5	0.0	0.0	9.9
Procter and Gamble			
CE-618 Methylene glycol	0.0	0.0	16.9
Diethylene glycol	0.0	0.0	6.4
Water (no salt)	0.0	0.0	1.0
	100.0	100.0	100.0
% Enzyme Activity Remaining after 33 days at 100°F	38%	58%	43%

TABLE XVII  
PERFORMANCE EVALUATION

INSTRUMENTAL GRADING	Formula B			Formula F			Formula G			CLOROX PRE-WASH			LSD
	70°	100°	130°	70°	100°	130°	70°	100°	130°	70°	100°	130°	
Wash Temp.	70°	100°	130°	70°	100°	130°	70°	100°	130°	70°	100°	130°	
AZOCASEIN	21.1	26.8	39.7	23.2	36.5	43.0	26.0	33.0	43.9	10.8	9.7	9.5	4.6
BLOOD	93.8	94.5	90.1	93.3	94.7	92.7	93.8	94.7	91.3	91.0	92.2	81.1	1.8
PERMANENT INK	53.1	56.5	57.9	44.5	45.9	52.3	54.8	58.3	59.4	54.6	56.8	59.3	3.4
DIRTY MOTOR OIL	91.2	91.5	92.6	90.3	90.0	92.5	86.9	89.3	90.4	90.8	92.4	92.8	0.8
CHOCOLATE	76.7	79.7	80.7	76.0	77.1	78.4	75.8	79.2	81.7	76.4	77.1	78.4	1.8
TEA	24.2	31.6	42.3	16.6	23.6	27.4	29.4	32.2	42.2	22.8	34.7	42.7	2.8
BLUBERRY	84.3	90.0	93.3	84.1	89.2	91.3	86.3	91.0	92.7	74.3	79.2	77.2	2.2
BANDY CLAY	91.9	94.6	94.3	93.6	95.0	95.3	92.7	94.8	95.6	93.4	95.8	95.5	1.3
SEBUM SOIL	93.0	93.8	95.3	92.7	93.8	94.2	91.9	92.5	93.6	94.4	95.1	96.0	0.7
GRASS	96.1	92.7	88.9	95.7	90.6	90.1	95.2	90.4	87.2	74.3	82.4	84.5	2.3
VISUAL GRADING													
BALLPOINT INK	2.8	2.8	3.0	2.5	2.9	2.9	2.8	3.2	3.2	3.3	3.3	3.3	0.2



TABLE XIX  
PERFORMANCE EVALUATION  
100% POLYESTER

INSTRUMENTAL GRADING	Formula B			Formula F			Formula G			CONTROL			LSD
	70°	100°	130°	70°	100°	130°	70°	100°	130°	70°	100°	130°	
Wash Temp.	70°	100°	130°	70°	100°	130°	70°	100°	130°	70°	100°	130°	
AZOCASEIN	22.6	29.8	31.4	22.2	30.6	32.9	--	--	--	9.8	9.9	12.0	4.0
BLOOD	96.7	96.5	96.5	96.8	96.7	96.4	--	--	--	97.1	96.8	97.6	0.9
PERMANENT INK	97.5	96.7	96.2	91.7	92.4	93.5	--	--	--	97.0	96.8	97.6	2.6
DIRTY MOTOR OIL	91.1	90.7	90.9	92.8	92.6	90.9	--	--	--	93.3	92.4	91.9	1.2
CHOCOLATE	97.1	96.9	97.0	97.3	97.1	97.1	--	--	--	97.1	96.9	97.1	1.0
TEA	96.7	96.8	97.0	86.4	84.6	85.3	--	--	--	96.7	96.6	96.4	2.9
BLUEBERRY	97.8	96.9	96.6	97.3	96.6	96.6	--	--	--	97.9	96.6	97.0	2.1
BANDY CLAY	98.2	94.2	99.4	98.6	98.9	99.2	--	--	--	98.7	99.4	99.2	0.9
SEBUM SOIL	94.2	95.5	95.3	96.3	96.0	95.2	--	--	--	95.2	95.9	95.4	1.5
GRASS	96.4	95.3	95.8	95.9	95.2	95.2	--	--	--	85.7	87.7	79.8	1.7

VISUAL GRADING

BALLPOINT INK	2.7	2.6	2.6	3.3	2.7	2.5	--	--	--	2.8	3.2	3.2	0.1
---------------	-----	-----	-----	-----	-----	-----	----	----	----	-----	-----	-----	-----

\* NO PERFORMANCE WORK WAS DONE ON POLYESTER WITH FORMULA G DUE TO LACK OF SUFFICIENT SAMPLE.

0177183

What is claimed is:

1. A soil- or stain-removal composition comprising at least one enzyme dissolved in aqueous solution encapsulated within reversed micelles, said micelles being a dispersion of the aqueous enzyme solution within a medium  
5 comprising a surfactant and organic solvent.
2. A composition according to claim 1 of which the water content is not more than 5% by weight.
3. A composition according to claim 1 or claim 2,  
wherein the enzyme is or are a protease, lipase, or  
10 amylase, or mixtures of the same.
4. A composition according to claim 1, claim 2 or claim 3, wherein the surfactant is a nonionic surfactant.
5. A composition according to any one of the preceding claims, wherein the organic solvent is a hydrocarbon, for  
15 example a paraffinic hydrocarbon.
6. A composition according to any one of the preceding claims, wherein the aqueous enzyme solution further includes an enzyme stabilizer.
7. A composition according to claim 6, wherein the  
20 enzyme stabilizer is sodium chloride.
8. A composition according to any one of the preceding claims, which includes organic solvents and surfactants for removing oil and grease based stains, and enzymes for removing protein, starch, and fat based stains.
- 25 9. A composition according to any one of the preceding claims, wherein the surfactant forms the micelle interface between said aqueous enzyme solution and the surfactant/  
organic solvent mixture.

10. A soil removal composition comprising about 60 to 70 weight % of a paraffinic hydrocarbon solvent in admixture with about 25 to 35 weight % of a nonionic surfactant, less than 5 weight % of an aqueous solution  
5 of one or more proteolytic, lipolytic or amylolytic enzymes, said aqueous solution consisting essentially of water, sodium chloride and the enzyme(s), said aqueous solution further dispersed in the solvent-surfactant mixture in the form of reversed micelles.
- 10 11. A composition according to any one of the preceding claims, wherein the enzyme(s) comprise(s) several tenths of one percent by weight of the composition.
12. A method for making a soil remover composition consisting essentially of organic solvent, surfactant  
15 and at least one enzyme, comprising encapsulating an aqueous solution of said enzyme(s) in reversed micelles within a mixture of said solvent and surfactants.
13. A method of claim 12, wherein said aqueous enzyme solution is combined with the surfactants, and the  
20 solvent is then added.



European Patent  
Office

EUROPEAN SEARCH REPORT

0177183

Application number

EP 85 30 6215

DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X	DE-A-3 003 766 (UNILEVER NV) * claim 1, page 6, paragraphs 3-5 *	1,3-5, 8	C 11 D 3/386
A	--- DE-A-2 633 601 (HENKEL KGAA) * claim 1 * -----		
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			C 11 D 3/00

Place of search  
BERLIN

Date of completion of the search  
03-12-1985

Examiner  
SCHULTZE D

CATEGORY OF CITED DOCUMENTS

X : particularly relevant if taken alone  
Y : particularly relevant if combined with another document of the same category  
A : technological background  
O : non-written disclosure  
P : intermediate document

T : theory or principle underlying the invention  
E : earlier patent document, but published on, or after the filing date  
D : document cited in the application  
L : document cited for other reasons  
& : member of the same patent family, corresponding document