




(86) Date de dépôt PCT/PCT Filing Date: 2012/12/20
(87) Date publication PCT/PCT Publication Date: 2013/06/27
(45) Date de délivrance/Issue Date: 2020/03/10
(85) Entrée phase nationale/National Entry: 2014/06/17
(86) N° demande PCT/PCT Application No.: EP 2012/076275
(87) N° publication PCT/PCT Publication No.: 2013/092791
(30) Priorité/Priority: 2011/12/22 (US61/578,979)

(51) Cl.Int./Int.Cl. *C07D 471/04* (2006.01),
A61K 31/529 (2006.01), *A61P 9/10* (2006.01)
(72) Inventeurs/Inventors:
LABEGUERE, FREDERIC GILBERT, FR;
NEWSOME, GREGORY JOHN ROBERT, FR;
ALVEY, LUKE JONATHAN, FR;
SANIERE, LAURENT RAYMOND MAURICE, FR;
FLETCHER, STEPHEN ROBERT, GB
(73) Propriétaire/Owner:
GALAPAGOS NV, BE
(74) Agent: NORTON ROSE FULBRIGHT CANADA
LLP/S.E.N.C.R.L., S.R.L.

(54) Titre : DIHYDROPYRIMIDINOISOQUINOLINONES ET COMPOSITIONS PHARMACEUTIQUES ASSOCIEES
DESTINEES AU TRAITEMENT DE TROUBLES INFLAMMATOIRES.
(54) Title: DIHYDROPYRIMIDINOISOQUINOLINONES AND PHARMACEUTICAL COMPOSITIONS THEREOF FOR THE
TREATMENT OF INFLAMMATORY DISORDERS.



1a

(57) **Abrégé/Abstract:**

A compound according to Formula (Ia) wherein L₁, and G, and R¹ are as described herein. The present invention relates to novel compounds according to Formula (I) that antagonize GPR84, a G-protein-coupled receptor that is involved in inflammatory

(57) Abrégé(suite)/Abstract(continued):

conditions, and methods for the production of these novel compounds, pharmaceutical compositions comprising these compounds, and methods for the prevention and/or treatment of inflammatory conditions (for example inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions by administering a compound of the invention.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
27 June 2013 (27.06.2013)



(10) International Publication Number
WO 2013/092791 A1

- (51) **International Patent Classification:**
C07D 471/04 (2006.01) *A61P 9/10* (2006.01)
A61K 31/529 (2006.01)
- (21) **International Application Number:**
PCT/EP2012/076275
- (22) **International Filing Date:**
20 December 2012 (20.12.2012)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/578,979 22 December 2011 (22.12.2011) US
- (71) **Applicant:** GALAPAGOS NV [BE/BE]; Industriepark
Mechelen Noord, Generaal De Wittelaan L11/A3, B-2800
Mechelen (BE).
- (72) **Inventors:** LABÉGUÈRE, Frédéric Gilbert; Galapagos
SASU, 102 Avenue Gaston Roussel, F-93230 Romainville
(FR). NEWSOME, Gregory John Robert; Galapagos
SASU, 102 Avenue Gaston Roussel, F-93230 Romainville
(FR). ALVEY, Luke Jonathan; Galapagos SASU, 102
Avenue Gaston Roussel, F-93230 Romainville (FR). SAN-
IÈRE, Laurent Raymond Maurice; Galapagos SASU,
102 Avenue Gaston Roussel, F-93230 Romainville (FR).
FLETCHER, Stephen Robert; 10 Clipped Hedge, Hat-

field Heath, Bishops Stortford, Hertfordshire CM22 7EG
(GB).


(74) **Agent:** NICHOL, Maria; Galapagos NV, Chesterford Re-
search Park, Saffron Walden, Essex CB10 1XL (GB).

(81) **Designated States** (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) **Title:** NOVEL DIHYDROPYRIMIDINOISOQUINOLINONES AND PHARMACEUTICAL COMPOSITIONS THEREOF
FOR THE TREATMENT OF INFLAMMATORY DISORDERS



(57) **Abstract:** A compound according to Formula (Ia) wherein L_1 , and G , and R^1 are as described herein. The present invention relates to novel compounds according to Formula (I) that antagonize GPR84, a G-protein-coupled receptor that is involved in inflammatory conditions, and methods for the production of these novel compounds, pharmaceutical compositions comprising these compounds, and methods for the prevention and/or treatment of inflammatory conditions (for example inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions by administering a compound of the invention.

WO 2013/092791 A1

WO 2013/092791 A1



Declarations under Rule 4.17:

- | | | | |
|---|---|---|---|
| — | <i>as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))</i> | — | <i>before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))</i> |
|---|---|---|---|

Published:

- *with international search report (Art. 21(3))*

**DIHYDROPYRIMIDINOISOQUINOLINONES AND PHARMACEUTICAL
COMPOSITIONS THEREOF FOR THE TREATMENT OF INFLAMMATORY
DISORDERS.**

5

FIELD OF THE INVENTION

[0001] The present invention relates to novel compounds that antagonize GPR84, a G-protein-coupled receptor that is involved in inflammatory conditions.

[0002] The present invention also provides methods for the production of these novel compounds, pharmaceutical compositions comprising these compounds, and methods for the prevention and/or treatment
10 of inflammatory conditions (for example inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions by administering a compound of the invention.

15

BACKGROUND OF THE INVENTION

[0003] GPR84 was recently isolated and characterized from human B cells (Wittenberger et al., 2001, J Mol Biol, 307, 799-813) as the result of an expressed sequence tag data mining strategy, and also using a
20 degenerate primer reverse transcriptase-polymerase chain reaction (RT-PCR) approach aimed to identify novel chemokine receptors expressed in neutrophils (Yousefi S et al. 2001 J Leukoc Biol;69, 1045-52).

[0004] GPR84 (also known as EX33) remained an orphan GPCR until the identification of medium-chain FFAs with carbon chain lengths of 9-14 as ligands for this receptor (Wang *et al.* (2006) J. Biol. Chem. 281:3457-64). GPR84 was described to be activated by capric acid (C10:0), undecanoic acid (C11:0) and
25 lauric acid (C12:0) with potencies of 5 μ M, 9 μ M and 11 μ M, respectively. Two small molecules were also described to have some GPR84 agonist activity: 3,3' di-indolylmethane (DIM) (Wang *et al.* (2006) J. Biol. Chem. 281:3457-64) and embelin (WO 2007/027661).

[0005] GPR84 expression has been shown to be expressed in immune cells at least but not limited to polymorphonuclear leukocytes (PMN), neutrophils, monocytes, T cells, B cells. (Wang et al., 2006, The
30 Journal of Biological Chemistry, 281, 45, 3457-3464, Yousefi et al., 2001, Journal of Leukocyte Biology, 69, 1045-1052, Venkataraman and Kuo, 2005, Immunology Letters, 101, 144-153, WO2007/027661 A2). Higher levels of GPR84 were measured in neutrophils and eosinophils than in T-cells and B-cells. GPR84 expression was demonstrated in tissues that may play a role in the propagation of the inflammatory response such as lung, spleen, bone marrow.

[0006] For example, in a recent review, Du Bois reported the current status of therapies for lung interstitial diseases, such as idiopathic pulmonary fibrosis (IPF). There are almost 300 distinct injurious or inflammatory causes of interstitial lung disease that can result in diffuse lung scarring, and the initial stages of the IPF pathology are very likely to involve inflammation (Du Bois, 2010, Nat Rev, Drug Discovery, 9, 129), and combination therapies involving anti-inflammatory treatment could be advantageously used.

[0007] The expression of GPR84 was highly up-regulated in monocytes/macrophages upon LPS stimulation (Wang et al., 2006, The Journal of Biological Chemistry, 281, 45, 3457-3464).

[0008] GPR84 knock-out (KO) mice are viable and indistinguishable from wild-type littermate controls (Venkataraman and Kuo, 2005, Immunology Letters, 101, 144-153). The proliferation of T and B cells in response to various mitogens is reported to be normal in GPR84-deficient mice (Venkataraman and Kuo, 2005, Immunology Letters, 101, 144-153). T helper 2 (Th2) differentiated T cells from GPR84 KO secreted higher levels of IL4, IL5, IL13, the 3 major Th2 cytokines, compared to wild-type littermate controls. In contrast, the production of the Th1 cytokine, INF γ , was similar in Th1 differentiated T cells from GPR84 KO and wild-type littermate (Venkataraman and Kuo, 2005, Immunology Letters, 101, 144-153).

[0009] In addition, capric acid, undecanoic acid and lauric acid dose dependently increased the secretion of interleukin-12 p40 subunit (IL-12 p40) from RAW264.7 murine macrophage-like cells stimulated with LPS. The pro-inflammatory cytokine IL-12 plays a pivotal role in promoting cell-mediated immunity to eradicate pathogens by inducing and maintaining T helper 1 (T_h1) responses and inhibiting T helper 2 (T_h2) responses. Medium-chain FFAs, through their direct actions on GPR84, may affect T_h1/T_h2 balance.

[0010] Berry et al. identified a whole-blood 393-gene transcriptional signature for active tuberculosis (TB) (Berry et al., 2010, Nature, 466, 973-979). GPR84 was part of this whole-blood 393-gene transcriptional signature for active TB indicating a potential role for GPR84 in infectious diseases.

[0011] GPR84 expression was also described in microglia, the primary immune effector cells of the central nervous system (CNS) from myeloid-monocytic origin (Bouchard et al., 2007, Glia, 55:790-800). As observed in peripheral immune cells, GPR84 expression in microglia was highly inducible under inflammatory conditions such as TNF α and IL1 treatment but also notably endotoxemia and experimental autoimmune encephalomyelitis (EAE), suggesting a role in neuro-inflammatory processes. Those results suggest that GPR84 would be up-regulated in CNS not only during endotoxemia and multiple sclerosis, but also in all neurological conditions in which TNF α or IL1 β pro-inflammatory cytokines are produced, including brain injury, infection, Alzheimer's disease (AD), Parkinson's disease (PD).

[0012] GPR84 expression was also observed in adipocytes and shown to be enhanced by inflammatory stimuli (Nagasaki *et al.*, 2012). The results suggest that GPR84 emerges in adipocytes in response to TNF α from infiltrating macrophages and exacerbates the vicious cycle between adiposity and diabetes, and therefore the inhibition of GPR84 activity might be beneficial for the treatment of endocrine and/or metabolic diseases.


[0013] Therefore, the present invention provides novel compounds, processes for their preparation and their use in the preparation of a medicament for the treatment of inflammatory conditions (for example inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions.

SUMMARY OF THE INVENTION

[0014] The present invention relates to novel dihydropyrimidinoisoquinolinone compounds that antagonize GPR84, and that are potentially useful for the treatment of inflammatory conditions (for example inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions.

[0015] The present invention also provides methods for the production of these compounds, pharmaceutical compositions comprising these compounds and methods for treating inflammatory conditions (for example inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions.

[0016] Accordingly, in a first aspect of the invention, a compound of the invention is disclosed having a Formula Ia:



wherein

R¹ is H, Me, or halo;

L₁ is absent or is -O-, -S-, or -NR^{4a}-;

G is

R^2 ,

-W-L₂-R², or

-W-L₃-R³;

5 W is C₁₋₄ alkylene, C₂₋₄ alkenylene having one double bond, or C₂₋₄ alkynylene having one triple bond;

L₂ is absent or is -O-;

R² is

- H,

10 - C₁₋₈ alkyl, optionally substituted with one to three groups independently selected from

o OH,

o halo,

o CN,

o C₁₋₆ alkoxy,

15 o C₃₋₇ cycloalkyl,

o 4-6 membered heterocycloalkyl comprising one to three heteroatoms independently selected from S, and O,

o 5-6 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, and

20 o phenyl,

- C₄₋₇ cycloalkenyl comprising one double bond,

- 5-7 membered heterocycloalkenyl comprising one double bond, and one to three heteroatoms independently selected from N, O, and S,

- C₃₋₇ cycloalkyl optionally substituted with one or more independently selected R⁵ groups,

25 - 4-10 membered heterocycloalkyl comprising one to two heteroatoms independently selected from S, and O, optionally substituted with one to three independently selected R⁵ groups,

- 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, optionally substituted with one to three independently selected R⁶ groups, or

- C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁶ groups;

30 L₃ is -NR^{4b};

R³ is

- C₁₋₄ alkyl substituted with

o C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁷ groups, or

- 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, optionally substituted with one or more independently selected R⁷ groups,
 - 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, optionally substituted with one or more independently selected R⁷ groups, or
 - C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁷ groups;
- Each R^{4a} and R^{4b} is independently selected from H, C₁₋₄ alkyl, and C₃₋₇ cycloalkyl;
R⁵ is oxo or R⁶;
R⁶ is
- OH,
 - halo,
 - -NO₂,
 - C₁₋₆ alkyl optionally substituted with one to three groups independently selected from halo, and OH,
 - C₁₋₆ alkoxy optionally substituted with one to three groups independently selected from halo, and OH,
 - C₃₋₇ cycloalkyl,
 - -C(=O)OR⁸,
 - -C(=O)NR⁹R¹⁰,
 - -NHC(=O)-C₁₋₄ alkyl,
 - -CN,
 - phenyl,
 - -O-phenyl,
 - 4-7 membered heterocycloalkyl comprising one to three heteroatoms independently selected from N, O, and S, or
 - 5-6 membered heteroaryl comprising one to three heteroatoms independently selected from N, O, and S; optionally substituted with one or more independently selected C₁₋₄ alkyl, C₁₋₄ alkoxy, CN, halo, and -C(=O)OR¹¹;
- R⁷ is C₁₋₄ alkyl, or halo, and
each of R⁸, R⁹, R¹⁰ and R¹¹ is independently selected from H and C₁₋₄ alkyl.

[0017] In a further aspect, the present invention provides pharmaceutical compositions comprising a compound of the invention, and a pharmaceutical carrier, excipient or diluent. Moreover, a compound of the present invention useful in the pharmaceutical compositions and treatment methods disclosed herein, is pharmaceutically acceptable as prepared and used. In this aspect of the invention, the pharmaceutical

composition may additionally comprise further active ingredients suitable for use in combination with a compound of the invention.

[0018] In another aspect of the invention, this invention provides novel compounds of the invention for use in therapy.

5 [0019] In a further aspect of the invention, this invention provides a method of treating a mammal susceptible to or afflicted with a condition from among those listed herein, and particularly, such condition as may be associated with aberrant activity of GPR84 and/or aberrant GPR84 expression and/or aberrant GPR84 distribution, for example inflammatory conditions (for example inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung
10 interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions, which method comprises administering a therapeutically effective amount of a compound of the invention, or one or more of the pharmaceutical compositions herein described.

[0020] In a further aspect, the present invention provides a compound of the invention for use in the
15 treatment or prevention of a condition selected from those listed herein, particularly such conditions as may be associated with aberrant activity of GPR84 and/or aberrant GPR84 expression and/or aberrant GPR84 distribution expression such as inflammatory conditions (for example inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious
20 diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions.

[0021] In additional aspects, this invention provides methods for synthesizing a compound of the invention, with representative synthetic protocols and pathways disclosed herein.

[0022] Accordingly, it is a principal object of this invention to provide a compound of the invention,
25 which can modify the activity of GPR84 and thus prevent or treat any conditions that may be causally related thereto.

[0023] It is further an object of this invention to provide a compound of the invention that can treat or alleviate conditions or diseases or symptoms of same, such as inflammatory conditions (for example inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive
30 pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions, that may be causally related to the activity and / or expression and/or distribution of GPR84.

[0024] A still further object of this invention is to provide pharmaceutical compositions that may be used
35 in the treatment or prevention of a variety of disease states, including the diseases associated with aberrant

activity of GPR84 and/or aberrant GPR84 expression and/or aberrant GPR84 distribution such as inflammatory conditions (for example inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions.

[0025] Other objects and advantages will become apparent to those skilled in the art from a consideration of the ensuing detailed description.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0026] The following terms are intended to have the meanings presented therewith below and are useful in understanding the description and intended scope of the present invention.

[0027] When describing the invention, which may include compounds, pharmaceutical compositions containing such compounds and methods of using such compounds and compositions, the following terms, if present, have the following meanings unless otherwise indicated. It should also be understood that when described herein any of the moieties defined forth below may be substituted with a variety of substituents, and that the respective definitions are intended to include such substituted moieties within their scope as set out below. Unless otherwise stated, the term 'substituted' is to be defined as set out below. It should be further understood that the terms 'groups' and 'radicals' can be considered interchangeable when used herein.

[0028] The articles 'a' and 'an' may be used herein to refer to one or to more than one (i.e. at least one) of the grammatical objects of the article. By way of example 'an analogue' means one analogue or more than one analogue.

[0029] 'Alkyl' means straight or branched aliphatic hydrocarbon having the specified number of carbon atoms. Particular alkyl groups have 1 to 6 carbon atoms or 1 to 4 carbon atoms. Branched means that one or more alkyl groups such as methyl, ethyl or propyl is attached to a linear alkyl chain. Particular alkyl groups are methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl, sec-butyl, n-pentyl, n-hexyl, and 1,2-dimethylbutyl. Particular alkyl groups have between 1 and 4 carbon atoms.

[0030] 'Alkylene' refers to divalent alkane radical groups having the number of carbon atoms specified, in particular 1 to 6 carbon atoms and more particularly 1 to 4 carbon atoms which can be straight-chained or branched. This term is exemplified by groups such as methylene (-CH₂-), ethylene (-CH₂-CH₂-), the propylene isomers (e.g., -CH₂-CH₂-CH₂- and -CH(CH₃)-CH₂-) and the like.

[0031] 'Alkenylene' refers to divalent alkene radical groups having the number of carbon atoms and the number of double bonds specified, in particular 2 to 6 carbon atoms and more particularly 2 to 4 carbon

atoms which can be straight-chained or branched. This term is exemplified by groups such as $-\text{CH}=\text{CH}-$, $-\text{CH}_2-\text{CH}=\text{CH}-$, $-\text{C}(\text{CH}_3)=\text{CH}-$, $-\text{C}(\text{CH}_3)=\text{CH}-\text{CH}_2-$, $-\text{C}(\text{CH}_3)=\text{C}(\text{CH}_3)-$, and $-\text{CH}_2-\text{C}(\text{CH}_3)=\text{CH}-$.

[0032] 'Alkynylene' refers to divalent alkyne radical groups having the number of carbon atoms and the number of triple bonds specified, in particular 2 to 6 carbon atoms and more particularly 2 to 4 carbon atoms which can be straight-chained or branched. This term is exemplified by groups such as $-\text{C}\equiv\text{C}-$, $-\text{CH}_2-\text{C}\equiv\text{C}-$, and $-\text{C}(\text{CH}_3)\text{H}-\text{C}\equiv\text{CH}-$.

[0033] 'Alkoxy' refers to the group O-alkyl, where the alkyl group has the number of carbon atoms specified. In particular the term refers to the group $-\text{O}-\text{C}_1-\text{C}_6$ alkyl. Particular alkoxy groups are methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, tert-butoxy, sec-butoxy, n-pentoxy, n-hexoxy, and 1,2-dimethylbutoxy. Particular alkoxy groups are lower alkoxy, i.e. with between 1 and 6 carbon atoms. Further particular alkoxy groups have between 1 and 4 carbon atoms.

[0034] 'Amino' refers to the radical $-\text{NH}_2$.

[0035] 'Aryl' refers to a monovalent aromatic hydrocarbon group derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic ring system. In particular aryl refers to an aromatic ring structure, mono-cyclic or poly-cyclic that includes the number of ring members specified. Particular aryl groups have from 6 to 10 ring members. Where the aryl group is a monocyclic ring system it preferentially contains 6 carbon atoms. Particularly aryl groups include phenyl, naphthyl, indenyl, and tetrahydronaphthyl.

[0036] 'Carboxy' refers to the radical $-\text{C}(\text{O})\text{OH}$.

[0037] 'Cycloalkyl' refers to cyclic non-aromatic hydrocarbyl groups having the number of carbon atoms specified. Particular cycloalkyl groups have from 3 to 7 carbon atoms. Such cycloalkyl groups include, by way of example, single ring structures such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl.

[0038] 'Cyano' refers to the radical $-\text{CN}$.


[0039] 'Halo' or 'halogen' refers to fluoro (F), chloro (Cl), bromo (Br) and iodo (I). Particular halo groups are either fluoro or chloro.

[0040] 'Hetero' when used to describe a compound or a group present on a compound means that one or more carbon atoms in the compound or group have been replaced by a nitrogen, oxygen, or sulfur heteroatom. Hetero may be applied to any of the hydrocarbyl groups described above such as alkyl, e.g. heteroalkyl, cycloalkyl, e.g. heterocycloalkyl, aryl, e.g. heteroaryl, and the like having from 1 to 5, and particularly from 1 to 3 heteroatoms.

[0041] 'Heteroaryl' means an aromatic ring structure, mono-cyclic or polycyclic, that includes one or more heteroatoms and the number of ring members specified. Particular heteraryl groups have 5 to 10 ring members, or 5 to 6 ring members. The heteroaryl group can be, for example, a five membered or six membered monocyclic ring or a bicyclic structure formed from fused five and six membered rings or two

fused six membered rings or, by way of a further example, two fused five membered rings. Each ring may contain up to four heteroatoms typically selected from nitrogen, sulphur and oxygen. Typically the heteroaryl ring will contain up to 4 heteroatoms, more typically up to 3 heteroatoms, more usually up to 2, for example a single heteroatom. In one embodiment, the heteroaryl ring contains at least one ring nitrogen atom. The nitrogen atoms in the heteroaryl rings can be basic, as in the case of an imidazole or pyridine, or essentially non-basic as in the case of an indole or pyrrole nitrogen. In general the number of basic nitrogen atoms present in the heteroaryl group, including any amino group substituents of the ring, will be less than five. Examples of five membered monocyclic heteroaryl groups include but are not limited to pyrrole, furan, thiophene, imidazole, furazan, oxazole, oxadiazole, oxatriazole, isoxazole, thiazole, isothiazole, pyrazole, triazole and tetrazole groups. Examples of six membered monocyclic heteroaryl groups include but are not limited to pyridine, pyrazine, pyridazine, pyrimidine and triazine. Particular examples of bicyclic heteroaryl groups containing a five membered ring fused to another five membered ring include but are not limited to imidazothiazole and imidazoimidazole. Particular examples of bicyclic heteroaryl groups containing a six membered ring fused to a five membered ring include but are not limited to benzfuran, benzthiophene, benzimidazole, benzoxazole, isobenzoxazole, benzisoxazole, benzthiazole, benzisothiazole, isobenzofuran, indole, isoindole, isoindolone, indolizine, indoline, isoindoline, purine (e.g., adenine, guanine), indazole, pyrazolopyrimidine, triazolopyrimidine, benzodioxole and pyrazolopyridine groups. Particular examples of bicyclic heteroaryl groups containing two fused six membered rings include but are not limited to quinoline, isoquinoline, chroman, thiochroman, chromene, isochromene, chroman, isochroman, benzodioxan, quinolizine, benzoxazine, benzodiazine, pyridopyridine, quinoxaline, quinazoline, cinnoline, phthalazine, naphthyridine and pteridine groups. Particular heteroaryl groups are those derived from thiophene, pyrrole, benzothiophene, benzofuran, indole, pyridine, quinoline, imidazole, oxazole and pyrazine.


[0042] Examples of representative heteroaryls include the following:



wherein each Y is selected from $>C=O$, NH, O and S.

[0043] As used herein, the term 'heterocycloalkyl' refers to a stable heterocyclic non-aromatic ring and/or rings containing one or more heteroatoms independently selected from N, O and S, fused thereto wherein the group contains the number of ring members specified. Particular heterocycloalkyl groups have

4-10 ring members or 5 to 7 ring members, or 5 to 6 ring members. The heterocycloalkyl group can be, for example, a five membered or six membered monocyclic ring or a bicyclic structure formed from fused five and six membered rings or two fused six membered rings or, by way of a further example, two fused five membered rings. Each ring may contain up to four heteroatoms typically selected from nitrogen, sulphur and oxygen. Typically the heterocycloalkyl ring will contain up to 4 heteroatoms, more typically up to 3 heteroatoms, more usually up to 2, for example a single heteroatom. In one embodiment, the heterocycloalkyl ring contains at least one ring nitrogen atom. A fused heterocyclic ring system may include carbocyclic rings and need only include one heterocyclic ring. Examples of heterocyclic rings include, but are not limited to, morpholine, piperidine (e.g. 1-piperidinyl, 2-piperidinyl, 3-piperidinyl and 4-piperidinyl), pyrrolidine (e.g. 1-pyrrolidinyl, 2-pyrrolidinyl and 3-pyrrolidinyl), pyrrolidone, pyran (2H-pyran or 4H-pyran), dihydrothiophene, dihydropyran, dihydrofuran, dihydrothiazole, tetrahydrofuran, tetrahydrothiophene, dioxane, tetrahydropyran (e.g. 4-tetrahydro pyranyl), imidazoline, imidazolidinone, oxazoline, thiazoline, 2-pyrazoline, pyrazolidine, piperazine, and N-alkyl piperazines such as N-methyl piperazine. Further examples include thiomorpholine and its S-oxide and S,S-dioxide (particularly thiomorpholine). Still further examples include azetidine, piperidone, piperazone, and N-alkyl piperidines such as N-methyl piperidine. Particular examples of heterocycloalkyl groups are shown in the following illustrative examples:



wherein each W is selected from CH₂, NH, O and S; and each Y is selected from NH, O, CO, SO₂, and S.

[0044] 'Hydroxy' refers to the radical -OH.

20 [0045] 'Nitro' refers to the radical -NO₂.

[0046] 'Substituted' refers to a group in which one or more hydrogen atoms are each independently replaced with the same or different substituent(s).

[0047] 'Thiol' refers to the group -SH.

25 [0048] 'Thioalkoxy' refers to the group -SR¹⁰ where R¹⁰ is an alkyl group with the number of carbon atoms specified. In particular thioalkoxy groups where R¹⁰ is a C₁-C₆ alkyl. Particular thioalkoxy groups are thiomethoxy, thioethoxy, n-thiopropoxy, thioisopropoxy, n-thiobutoxy, tert-thiobutoxy, sec-thiobutoxy, n-thiopentoxo, n-thiohexoxy, and 1,2-dimethylthiobutoxy. Particular thioalkoxy groups are lower thioalkoxy,

i.e. with between 1 and 6 carbon atoms. Further particular thioalkoxy groups have between 1 and 4 carbon atoms.

[0049] As used herein, the term 'substituted with one or more' refers to one to four substituents. In one embodiment it refers to one to three substituents. In further embodiments it refers to one or two substituents.

5 In a yet further embodiment it refers to one substituent.

[0050] One having ordinary skill in the art of organic synthesis will recognize that the maximum number of heteroatoms in a stable, chemically feasible heterocyclic ring, whether it is aromatic or non aromatic, is determined by the size of the ring, the degree of unsaturation and the valence of the heteroatoms. In general, a heterocyclic ring may have one to four heteroatoms so long as the heteroaromatic ring is chemically

10 feasible and stable.

[0051] 'Pharmaceutically acceptable' means approved or approvable by a regulatory agency of the Federal or a state government or the corresponding agency in countries other than the United States, or that is listed in the U.S. Pharmacopoeia or other generally recognized pharmacopoeia for use in animals, and more particularly, in humans.

15 [0052] 'Pharmaceutically acceptable salt' refers to a salt of a compound that is pharmaceutically acceptable and that possesses the desired pharmacological activity of the parent compound. In particular, such salts are non-toxic may be inorganic or organic acid addition salts and base addition salts. Specifically, such salts include: (1) acid addition salts, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such

20 as acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl) benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]-oct-2-

25 ene-1-carboxylic acid, glucoheptonic acid, 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like; or (2) salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic base such as ethanolamine, diethanolamine, triethanolamine, N-methylglucamine and the

30 like. Salts further include, by way of example only, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium, and the like; and when the compound contains a basic functionality, salts of non toxic organic or inorganic acids, such as hydrochloride, hydrobromide, tartrate, mesylate, acetate, maleate, oxalate and the like. The term 'pharmaceutically acceptable cation' refers to an acceptable cationic counter-ion of an acidic functional group. Such cations are exemplified by sodium, potassium, calcium, magnesium,

35 ammonium, tetraalkylammonium cations, and the like.

[0053] 'Pharmaceutically acceptable vehicle' refers to a diluent, adjuvant, excipient or carrier with which a compound of the invention is administered.

[0054] 'Prodrugs' refers to compounds, including derivatives of the compounds of the invention, which have cleavable groups and become by solvolysis or under physiological conditions the compounds of the invention which are pharmaceutically active in vivo. Such examples include, but are not limited to, choline ester derivatives and the like, N-alkylmorpholine esters and the like.

[0055] 'Solvate' refers to forms of the compound that are associated with a solvent, usually by a solvolysis reaction. This physical association includes hydrogen bonding. Conventional solvents include water, ethanol, acetic acid and the like. The compounds of the invention may be prepared e.g. in crystalline form and may be solvated or hydrated. Suitable solvates include pharmaceutically acceptable solvates, such as hydrates, and further include both stoichiometric solvates and non-stoichiometric solvates. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid. 'Solvate' encompasses both solution-phase and isolable solvates. Representative solvates include hydrates, ethanolates and methanolates.

[0056] 'Subject' includes humans. The terms 'human', 'patient' and 'subject' are used interchangeably herein.

[0057] 'Effective amount' means the amount of a compound of the invention that, when administered to a subject for treating a disease, is sufficient to effect such treatment for the disease. The "effective amount" can vary depending on the compound, the disease and its severity, and the age, weight, etc., of the subject to be treated.

[0058] 'Preventing' or 'prevention' refers to a reduction in risk of acquiring or developing a disease or disorder (i.e., causing at least one of the clinical symptoms of the disease not to develop in a subject that may be exposed to a disease-causing agent, or predisposed to the disease in advance of disease onset).

[0059] The term 'prophylaxis' is related to 'prevention', and refers to a measure or procedure the purpose of which is to prevent, rather than to treat or cure a disease. Non-limiting examples of prophylactic measures may include the administration of vaccines; the administration of low molecular weight heparin to hospital patients at risk for thrombosis due, for example, to immobilization; and the administration of an anti-malarial agent such as chloroquine, in advance of a visit to a geographical region where malaria is endemic or the risk of contracting malaria is high.

[0060] 'Treating' or 'treatment' of any disease or disorder refers, in one embodiment, to ameliorating the disease or disorder (i.e., arresting the disease or reducing the manifestation, extent or severity of at least one of the clinical symptoms thereof). In another embodiment 'treating' or 'treatment' refers to ameliorating at least one physical parameter, which may not be discernible by the subject. In yet another embodiment, 'treating' or 'treatment' refers to modulating the disease or disorder, either physically, (e.g., stabilization of a

discernible symptom), physiologically, (e.g., stabilization of a physical parameter), or both. In a further embodiment, “treating” or “treatment” relates to slowing the progression of the disease.

[0061] As used herein the term ‘inflammatory condition(s)’ refers to the group of conditions including, rheumatoid arthritis, osteoarthritis, juvenile idiopathic arthritis, vasculitis, psoriasis, gout, allergic airway disease (e.g. asthma, rhinitis), inflammatory bowel diseases (e.g. Crohn’s disease, ulcerative colitis), and endotoxin-driven disease states (e.g. complications after bypass surgery or chronic endotoxin states contributing to e.g. chronic cardiac failure). Particularly the term refers to rheumatoid arthritis, allergic airway disease (e.g. asthma) and inflammatory bowel diseases.

[0062] As used herein, the term ‘infectious diseases’ refers to bacterial infectious diseases and includes but is not limited to conditions such as sepsis, septicemia, endotoxemia, systemic inflammatory response syndrome (SIRS), gastritis, enteritis, enterocolitis, tuberculosis, and other infections involving, for example, Yersinia, Salmonella, Chlamydia, Shigella, or enterobacteria species.

[0063] As used herein the term ‘autoimmune disease(s)’ refers to the group of diseases including obstructive airways disease (including conditions such as COPD (Chronic obstructive pulmonary disease)), psoriasis, asthma (e.g intrinsic asthma, extrinsic asthma, dust asthma, infantile asthma) particularly chronic or inveterate asthma (for example late asthma and airway hyperresponsiveness), bronchitis, including bronchial asthma, systemic lupus erythematosus (SLE), multiple sclerosis, type I diabetes mellitus and complications associated therewith, atopic eczema (atopic dermatitis), contact dermatitis and further eczematous dermatitis, vasculitis, inflammatory bowel disease (e.g. Crohn’s disease and ulcerative colitis), atherosclerosis and amyotrophic lateral sclerosis. Particularly the term refers to COPD, asthma, psoriasis, systemic lupus erythematosus, type I diabetes mellitus, vasculitis and inflammatory bowel disease.

[0064] As used herein the term ‘endocrine and/or metabolic disease(s)’ refers to the group of conditions involving the body’s over- or under-production of certain hormones, while metabolic disorders affect the body’s ability to process certain nutrients and vitamins. Endocrine disorders include hypothyroidism, congenital adrenal hyperplasia, diseases of the parathyroid gland, diabetes mellitus, diseases of the adrenal glands (including Cushing’s syndrome and Addison’s disease), and ovarian dysfunction (including polycystic ovary syndrome), among others. Some examples of metabolic disorders include cystic fibrosis, phenylketonuria (PKU), diabetes, hyperlipidemia, gout, and rickets.

[0065] As used herein, the term ‘diseases involving impairment of immune cell functions’ includes conditions with symptoms such as recurrent and drawn out viral and bacterial infections, and slow recovery. Other invisible symptoms may be the inability to kill off parasites, yeasts and bacterial pathogens in the intestines or throughout the body.

[0066] As used herein the term ‘neuroinflammatory conditions’ refers to diseases or disorders characterized by abrupt neurologic deficits associated with inflammation, demyelination, and axonal damage,

and includes but is not limited to conditions such as Guillain-Barré syndrome (GBS), multiple sclerosis, axonal degeneration, autoimmune encephalomyelitis.

[0067] 'Compound(s) of the invention', and equivalent expressions, are meant to embrace compounds of the Formula(e) as herein described, which expression includes the pharmaceutically acceptable salts, and the solvates, e.g., hydrates, and the solvates of the pharmaceutically acceptable salts where the context so permits. Similarly, reference to intermediates, whether or not they themselves are claimed, is meant to embrace their salts, and solvates, where the context so permits.

[0068] When ranges are referred to herein, for example but without limitation, C₁₋₆ alkyl, the citation of a range should be considered a representation of each member of said range.

[0069] Other derivatives of the compounds of this invention have activity in both their acid and acid derivative forms, but in the acid sensitive form often offers advantages of solubility, tissue compatibility, or delayed release in the mammalian organism (see, Bundgard, H., Design of Prodrugs, pp. 7-9, 21-24, Elsevier, Amsterdam 1985). Prodrugs include acid derivatives well known to practitioners of the art, such as, for example, esters prepared by reaction of the parent acid with a suitable alcohol, or amides prepared by reaction of the parent acid compound with a substituted or unsubstituted amine, or acid anhydrides, or mixed anhydrides. Simple aliphatic or aromatic esters, amides and anhydrides derived from acidic groups pendant on the compounds of this invention are particularly useful prodrugs. In some cases it is desirable to prepare double ester type prodrugs such as (acyloxy)alkyl esters or ((alkoxycarbonyl)oxy)alkylesters. Particular such prodrugs are the C₁ to C₈ alkyl, , and substituted or unsubstituted C₆₋₁₀ aryl, esters of the compounds of the invention.

[0070] As used herein, the term 'isotopic variant' refers to a compound that contains unnatural proportions of isotopes at one or more of the atoms that constitute such compound. For example, an 'isotopic variant' of a compound can contain one or more non-radioactive isotopes, such as for example, deuterium (²H or D), carbon-13 (¹³C), nitrogen-15 (¹⁵N), or the like. It will be understood that, in a compound where such isotopic substitution is made, the following atoms, where present, may vary, so that for example, any hydrogen may be ²H/D, any carbon may be ¹³C, or any nitrogen may be ¹⁵N, and that the presence and placement of such atoms may be determined within the skill of the art. Likewise, the invention may include the preparation of isotopic variants with radioisotopes, in the instance for example, where the resulting compounds may be used for drug and/or substrate tissue distribution studies. The radioactive isotopes tritium, i.e. ³H, and carbon-14, i.e. ¹⁴C, are particularly useful for this purpose in view of their ease of incorporation and ready means of detection. Further, compounds may be prepared that are substituted with positron emitting isotopes, such as ¹¹C, ¹⁸F, ¹⁵O and ¹³N, and would be useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy.

[0071] All isotopic variants of the compounds provided herein, radioactive or not, are intended to be encompassed within the scope of the invention.

[0072] It is also to be understood that compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed 'isomers'. Isomers that differ in the arrangement of their atoms in space are termed 'stereoisomers'.

[0073] Stereoisomers that are not mirror images of one another are termed 'diastereomers' and those that are non-superimposable mirror images of each other are termed 'enantiomers'. When a compound has an asymmetric center, for example, it is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be characterized by the absolute configuration of its asymmetric center and is described by the R- and S-sequencing rules of Cahn and Prelog, or by the manner in which the molecule rotates the plane of polarized light and designated as dextrorotatory or levorotatory (i.e., as (+) or (-)-isomers respectively). A chiral compound can exist as either individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a 'racemic mixture'.

[0074] 'Tautomers' refer to compounds that are interchangeable forms of a particular compound structure, and that vary in the displacement of hydrogen atoms and electrons. Thus, two structures may be in equilibrium through the movement of π electrons and an atom (usually H). For example, enols and ketones are tautomers because they are rapidly interconverted by treatment with either acid or base. Another example of tautomerism is the aci- and nitro- forms of phenylnitromethane, that are likewise formed by treatment with acid or base.

[0075] Tautomeric forms may be relevant to the attainment of the optimal chemical reactivity and biological activity of a compound of interest.

[0076] The compounds of the invention may possess one or more asymmetric centers; such compounds can therefore be produced as individual (R)- or (S)- stereoisomers or as mixtures thereof.

[0077] Unless indicated otherwise, the description or naming of a particular compound in the specification and claims is intended to include both individual enantiomers and mixtures, racemic or otherwise, thereof. The methods for the determination of stereochemistry and the separation of stereoisomers are well-known in the art.

[0078] It will be appreciated that compounds of the invention may be metabolized to yield biologically active metabolites.


THE COMPOUNDS

[0079] The present invention relates to novel compounds that antagonize GPR84 and that may be useful for the treatment of inflammatory conditions (for example inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases,

autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions.

[0080] The present invention also provides methods for the production of the compounds of the invention, pharmaceutical compositions comprising the compounds of the invention and methods for treating diseases involving inflammatory conditions (for example inflammatory conditions (for example inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions, by administering a compound of the invention. A compound of the invention is an inhibitor of GPR84.

[0081] Accordingly, in a first aspect of the invention, a compound of the invention is disclosed having a Formula Ia:



wherein

R¹ is H, Me, or halo;

L₁ is absent or is -O-, -S-, or -NR^{4a}-;

G is

R²,

-W-L₂-R², or

-W-L₃-R³;

W is C₁₋₄ alkylene, C₂₋₄ alkenylene having one double bond, or C₂₋₄ alkynylene having one triple bond;

L₂ is absent or is -O-;

R² is

- H,

- C₁₋₈ alkyl, optionally substituted with one to three groups independently selected from


○ OH,

○ halo,

- CN,
 - C₁₋₆ alkoxy,
 - C₃₋₇ cycloalkyl,
 - 4-6 membered heterocycloalkyl comprising one to three heteroatoms independently selected from S, and O,
 - 5-6 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, and
 - phenyl,
 - C₄₋₇ cycloalkenyl comprising one double bond,
 - 5-7 membered heterocycloalkenyl comprising one double bond, and one to three heteroatoms independently selected from N, O, and S,
 - C₃₋₇ cycloalkyl optionally substituted with one or more independently selected R⁵ groups,
 - 4-10 membered heterocycloalkyl comprising one to two heteroatoms independently selected from S, and O, optionally substituted with one to three independently selected R⁵ groups,
 - 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, optionally substituted with one to three independently selected R⁶ groups, or
 - C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁶ groups;
- L₃ is -NR^{4b}-;
- R³ is
- C₁₋₄ alkyl substituted with
 - C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁷ groups, or
 - 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, optionally substituted with one or more R independently selected R⁷ groups,
 - 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, optionally substituted with one or more independently selected R⁷ groups, or
 - C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁷ groups;
- Each R^{4a} and R^{4b} is independently selected from H, C₁₋₄ alkyl, and C₃₋₇ cycloalkyl;
- R⁵ is oxo or R⁶;
- R⁶ is
- OH,
 - halo,
 - -NO₂,
 - C₁₋₆ alkyl optionally substituted with one to three groups independently selected from halo, and OH,


- C₁₋₆ alkoxy optionally substituted with one to three groups independently selected from halo, and OH,
- C₃₋₇ cycloalkyl,
- -C(=O)OR⁸,
- 5 - -C(=O)NR⁹R¹⁰,
- -NHC(=O)-C₁₋₄ alkyl,
- -CN,
- phenyl,
- -O-phenyl,
- 10 - 4-7 membered heterocycloalkyl comprising one to three heteroatoms independently selected from N, O, and S, or
- 5-6 membered heteroaryl comprising one to three heteroatoms independently selected from N, O, and S; optionally substituted with one or more independently selected C₁₋₄ alkyl, C₁₋₄ alkoxy, CN, halo, and -C(=O)OR¹¹;
- 15 R⁷ is C₁₋₄ alkyl, or halo; and
each of R⁸, R⁹, R¹⁰ and R¹¹ is independently selected from H and C₁₋₄ alkyl.

[0082] In a further embodiment, a compound of the invention is disclosed having a Formula Ib:



wherein R¹, L₁ and G are as previously described.

- 20 [0083] In yet a further embodiment, a compound of the invention is disclosed having a Formula Ic:




wherein R^1 , L_1 and G are as previously described.


[0084] In one embodiment, the compound of the invention is according to Formula Ia, Ib or Ic, wherein R^1 is Me, F, or Cl.

[0085] In one embodiment, the compound of the invention is according to Formula Ia, Ib or Ic, wherein R^1 is H.


[0086] In one embodiment, the compound of the invention is according to Formula IIa, IIb, or IIc:



IIa




IIb




IIc

wherein L_1 , and R^2 are as described previously.


[0087] In one embodiment, the compound of the invention is according to Formula IIIa, IIIb, or IIIc:



IIIa




IIIb




IIIc

wherein L_1 , W , L_2 , and R^2 are as described previously.


[0088] In one embodiment, the compound of the invention is according to Formula IVa, IVb, or IVc:



IVa



IVb



IVc

wherein L_1 , W , L_3 , and R^3 are as described previously.

[0089] In one embodiment, the compound of the invention is according to any one of Formulae Ia-IVc, wherein L_1 is absent, or is $-O-$. In a preferred embodiment, L_1 is absent.

[0090] In another embodiment, the compound of the invention is according to any one of Formulae Ia-IVc, wherein L_1 is $-NR^{4a}-$, wherein R^{4a} is as described previously. In a preferred embodiment, R^{4a} is H, Me, Et, or cyclopropyl. In a more preferred embodiment, R^{4a} is H.

[0091] In one embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, or IIIa-IVc, wherein W is C_{1-4} alkylene. In a preferred embodiment, W is $-CH_2-$, $-CH_2-CH_2-$, $-CH_2-CH_2-CH(CH_3)-$, $-CH_2-CH(-CH_2-CH_3)-$, $-CH_2-C(CH_3)_2-$, or $-CH_2-CH_2-CH_2-$. In a more preferred embodiment, W is $-CH_2-$. In another more preferred embodiment, W is $-CH_2-CH_2-$.

[0092] In one embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, or IIIa-IVc, wherein W is C_{2-4} alkenylene having one double bond. In a preferred embodiment, W is $-CH=CH-$, $-CH_2-CH=CH-$, or $-CH=CH-CH_2-$. In a more preferred embodiment, W is $-CH=CH-$. In another more preferred embodiment, W is $-CH_2-CH=CH-$.

[0093] In one embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, or IIIa-IVc, wherein W is C_{2-4} alkynylene having one triple bond. In a preferred embodiment, W is $-C\equiv C-$, $-CH_2-C\equiv C-$, or $-C\equiv C-CH_2-$. In a more preferred embodiment, W is $-C\equiv C-$. In another more preferred embodiment, W is $-CH_2-C\equiv C-$.

[0094] In one embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, or IIIa-IIIc, wherein L_2 is absent. In another embodiment, L_2 is $-O-$.

[0095] In one embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, or IIIa-IIIc, wherein L_1 is absent or is $-O-$, W is C_{1-4} alkylene; and L_2 and R^2 are as described previously. In a preferred embodiment, W is $-CH_2-$, $-CH_2-CH_2-$, or $-CH_2-CH_2-CH_2-$. In a more preferred embodiment, W is $-CH_2-$. In another preferred embodiment, W is $-CH_2-CH_2-$.

[0096] In another embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, or IIIa-IIIc, wherein L_1 is absent or is $-O-$, W is C_{2-4} alkenylene having one double bond; and L_2 and R^2 are as described previously. In a preferred embodiment, W is $-CH=CH-$, $-CH_2-CH=CH-$, or $-CH=CH-CH_2-$. In a more preferred embodiment, W is $-CH=CH-$. In another more preferred embodiment, W is $-CH_2-CH=CH-$.

[0097] In yet another embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, or IIIa-IIIc, wherein L_1 is absent, W is C_{2-4} alkynylene having one triple bond; and L_2 and R^2 are as described previously. In a preferred embodiment, W is $-C\equiv C-$, $-CH_2-C\equiv C-$, or $-C\equiv C-CH_2-$. In a more preferred embodiment, W is $-C\equiv C-$. In another more preferred embodiment, W is $-CH_2-C\equiv C-$.

[0098] In one embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, or IIIa-IIIc, wherein L_1 and L_2 are absent, W is C_{1-4} alkylene; and R^2 is as described previously. In a preferred

embodiment, W is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, or $-\text{CH}_2\text{CH}_2\text{CH}_2-$. In a more preferred embodiment, W is $-\text{CH}_2-$. In another more preferred embodiment, W is $-\text{CH}_2\text{CH}_2-$.

[0099] In another embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, or IIIa-IIIc, wherein L_1 and L_2 are absent, W is C_{2-4} alkenylene having one double bond; and R^2 is as described previously. In a preferred embodiment, W is $-\text{CH}=\text{CH}-$, $-\text{CH}_2\text{CH}=\text{CH}-$, or $-\text{CH}=\text{CHCH}_2-$. In a more preferred embodiment, W is $-\text{CH}=\text{CH}-$. In another more preferred embodiment, W is $-\text{CH}_2\text{CH}=\text{CH}-$.

[00100] In yet another embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, or IIIa-IIIc, wherein L_1 and L_2 are absent, W is C_{2-4} alkynylene having one triple bond; and R^2 is as described previously. In a preferred embodiment, W is $-\text{C}\equiv\text{C}-$, $-\text{CH}_2\text{C}\equiv\text{C}-$, or $-\text{C}\equiv\text{CCH}_2-$. In a more preferred embodiment, W is $-\text{C}\equiv\text{C}-$. In another more preferred embodiment, W is $-\text{CH}_2\text{C}\equiv\text{C}-$.

[00101] In one embodiment, the compound of the invention is according to any one of Formulae Ia-IIIc, wherein R^2 is H.

[00102] In another embodiment, the compound of the invention is according to any one of Formulae any one of Formulae Ia-IIIc, wherein R^2 is C_{1-8} alkyl. In a preferred embodiment, R^2 is Me, Et, *n*-Pr, *i*-Pr, *i*-Bu, or *t*-Bu. In a more preferred embodiment, R^2 is Me, Et, *i*-Pr or *t*-Bu. In a more preferred embodiment, R^2 is *t*-Bu.

[00103] In one embodiment, the compound of the invention is according to any one of Formulae Ia-IIIc, wherein R^2 is C_{1-8} alkyl substituted with one to three groups selected from OH, halo, CN, C_{1-6} alkoxy, C_{3-7} cycloalkyl, 4-6 membered heterocycloalkyl (comprising one to three heteroatoms independently selected from S, and O), 5-6 membered heteroaryl (comprising one to three heteroatoms independently selected from N, S, and O), and phenyl. In a preferred embodiment, R^2 is Me, Et, *n*-Pr, *i*-Pr, *i*-Bu, or *t*-Bu substituted with one to three groups selected from OH, halo, CN, C_{1-6} alkoxy, C_{3-7} cycloalkyl, 4-6 membered heterocycloalkyl (comprising one to three heteroatoms independently selected from S, and O), 5-6 membered heteroaryl (comprising one to three heteroatoms independently selected from N, S, and O), and phenyl. In another preferred embodiment, is C_{1-8} alkyl substituted with one to three groups selected from OH, F, Cl, CN, -OMe, -OEt, *Oi*-Pr, cyclopropyl, cyclobutyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, pyrrololyl, imidazolyl, triazolyl, oxazolyl, thiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, and phenyl. In a more preferred embodiment, R^2 is Me, Et, *n*-Pr, *i*-Pr, *i*-Bu, or *t*-Bu substituted with one to three groups selected from OH, F, Cl, CN, -OMe, -OEt, -*Oi*-Pr, cyclopropyl, cyclobutyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, pyrrololyl, imidazolyl, triazolyl, oxazolyl, thiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and phenyl.

[00104] In another embodiment, the compound of the invention is according to any one of Formulae Ia-IIIc, wherein R^2 is C_{1-8} alkyl substituted with one group selected from OH, halo, CN, C_{1-6} alkoxy, C_{3-7} cycloalkyl, 4-6 membered heterocycloalkyl (comprising one to three heteroatoms independently selected from S, and O), 5-6 membered heteroaryl (comprising one to three heteroatoms independently selected from N, S, and O), and phenyl. In a preferred embodiment, R^2 is Me, Et, *n*-Pr, *i*-Pr, *i*-Bu, or *t*-Bu substituted with

one group selected from OH, halo, CN, C₁₋₆ alkoxy, C₃₋₇ cycloalkyl, 4-6 membered heterocycloalkyl (comprising one to three heteroatoms independently selected from S, and O), 5-6 membered heteroaryl (comprising one to three heteroatoms independently selected from N, S, and O), and phenyl. In another preferred embodiment, is C₁₋₈ alkyl substituted with one group selected from OH, F, Cl, CN, -OMe, -OEt, -
 5 Oi-Pr, cyclopropyl, cyclobutyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, pyrrololyl, imidazolyl, triazolyl, oxazolyl, thiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, and phenyl. In a more preferred embodiment, R² is Me, Et, *n*-Pr, *i*-Pr, *i*-Bu, or *t*-Bu substituted with one group selected from OH, F, Cl, CN, -OMe, -OEt, -
 Oi-Pr, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, pyrrololyl, imidazolyl, triazolyl, oxazolyl, thiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, and phenyl. In a most
 10 preferred embodiment, R² is -CH₂-OH, -C(CH₃)₂-OH, -CH(OH)CH₃, -CH(OH)-C₂H₅, -CH(OH)-C₃H₇, -
 C(OH)(C₂H₅)₂, -C(OH)H-CH(CH₃)₂, -C(OH)H-CH₂-CH(CH₃)₂, -C(OH)H-C(CH₃)₃, -CH₂-CN, -CH₂-OCH₃, -
 CH₂-CH₂-OCH₃, -CH(OCH₃)-CH₃, -C(OCH₃)H-CH(CH₃)₂, -CH₂-F, -CH₂-CH₂-F, -CH₂-cyclopropyl, -CH₂-
 cyclopentyl, -CH₂-oxetanyl, -CH₂-tetrahydrofuranyl, or -CH₂-tetrahydropyranyl.

[00105] In one embodiment, the compound of the invention is according to any one of Formulae Ia-IIIc,
 15 wherein R² is C₄₋₇ cycloalkenyl comprising one double bond. In a preferred embodiment, R² is cyclohexenyl.

[00106] In one embodiment, the compound of the invention is according to any one of Formulae Ia-IIIc, wherein R² is 5-7 membered heterocycloalkenyl comprising one double bond, and one to three heteroatoms independently selected from N, O, and S. In a preferred embodiment, R² is dihydropyranyl.

[00107] In one embodiment, the compound of the invention is according to any one of Formulae Ia-IIIc,
 20 wherein R² is C₃₋₇ cycloalkyl. In a preferred embodiment, R² is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In a more preferred embodiment, R² is cyclopropyl.

[00108] In another embodiment, the compound of the invention is according to any one of Formulae Ia-IIIc, wherein R² is C₃₋₇ cycloalkyl substituted with one to three independently selected R⁵ groups. In a preferred embodiment, R² is C₃₋₇ cycloalkyl substituted with one R⁵ group. In a more preferred embodiment,
 25 R² is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, each of which is substituted with one R⁵ group. In another more preferred embodiment, R² is C₃₋₇ cycloalkyl substituted with one R⁵ group, wherein R⁵ is oxo, or R⁶ wherein R⁶ is selected from OH, or C₁₋₆ alkyl. In a most preferred embodiment, R² is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, each of which is substituted with one R⁵ group, wherein R⁵ is oxo, or R⁶ wherein R⁶ is selected from OH, or C₁₋₆ alkyl. In a further most preferred embodiment, R² is cyclopropyl,
 30 cyclobutyl, cyclopentyl or cyclohexyl, each of which is substituted with one R⁵ group, wherein R⁵ is R⁶, and R⁶ is selected from OH.

[00109] In one embodiment, the compound of the invention is according to any one of Formulae Ia-IIIc, wherein R² is 4-10 membered heterocycloalkyl comprising one to two heteroatoms independently selected from S, and O. In a preferred embodiment, R² is oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, or dioxanyl.

[00110] In another embodiment, the compound of the invention is according to any one of Formulae Ia-IIIc, R² is 4-10 membered heterocycloalkyl comprising one to two heteroatoms independently selected from S, and O substituted with one to three independently selected R⁵ groups. In a preferred embodiment, R² is 4-10 membered heterocycloalkyl comprising one to two heteroatoms independently selected from S, and O substituted with one R⁵ group. In a more preferred embodiment, R² is 4-10 membered heterocycloalkyl comprising one to two heteroatoms independently selected from S, and O substituted with one R⁵ group, wherein R⁵ is selected from oxo, or R⁶ wherein R⁶ is selected from OH, and C₁₋₆ alkyl. In another more preferred embodiment, R² is oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, or dioxanyl, each of which is substituted with one R⁵ group. In a most preferred embodiment, R² is oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, or dioxanyl, each of which is substituted with one R⁵ group, wherein R⁵ is selected from oxo, or R⁶ wherein R⁶ is selected from OH, and C₁₋₆ alkyl,

[00111] In one embodiment, the compound of the invention is according to any one of Formulae Ia-IIIc, R² is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O. In a preferred embodiment, R² is furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl.

[00112] In another embodiment, the compound of the invention is according to any one of Formulae Ia-IIIc, R² is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one to three independently selected R⁶ groups. In a preferred embodiment, R² is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or two independently selected R⁶ groups. In a more preferred embodiment, R² is furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl, substituted with one or two independently selected R⁶ groups. In another more preferred embodiment, R² is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or two independently selected R⁶ groups, wherein each R⁶ is independently selected from OH, halo, C₁₋₆ alkyl, C₁₋₆ alkyl substituted with one or more halo, C₁₋₆ alkoxy, -CN, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl comprising one to three heteroatoms independently selected from N, O, and S, and phenyl. In most preferred embodiment, R² is furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl, each of which is substituted with one or two independently selected R⁶ groups, wherein each R⁶ is independently selected from OH, halo, C₁₋₆ alkyl, C₁₋₆ alkyl substituted with one or more halo, C₁₋₆ alkoxy, -CN, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl comprising one to three heteroatoms independently selected from N, O, and S, and phenyl. In another most preferred embodiment, R² is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or two independently selected R⁶ groups, wherein each R⁶ is independently selected from OH, F, Cl, Me, Et, Pr, *i*-Pr, *t*-Bu, -CF₃, -OMe, -OEt, *Oi*-Pr, -CN, cyclopropyl, pyrrolidinyl, morpholinyl,

piperidinyl, or phenyl. In further most preferred embodiment, R² is furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl, each of which is substituted with one or two independently selected R⁶ groups, wherein each R⁶ is independently selected from OH, F, Cl, Me, Et, Pr, *i*-Pr, *t*-Bu, -CF₃, -OMe, -OEt, -O*i*-Pr, -CN, cyclopropyl, pyrrolidinyl, morpholinyl, piperidinyl, and phenyl.

[00113] In another embodiment, the compound of the invention is according to any one of Formulae Ia-IIIc, R² is C₆₋₁₀ aryl. In a preferred embodiment, R² is phenyl.

[00114] In another embodiment, the compound of the invention is according to any one of Formulae Ia-IIIc, R² is C₆₋₁₀ aryl substituted with one or more independently selected R⁶ groups. In a preferred embodiment, R² is C₆₋₁₀ aryl substituted with one or two independently selected R⁶ groups. In a more preferred embodiment, R² is C₆₋₁₀ aryl substituted with one or two independently selected R⁶ groups, wherein each R⁶ group is selected from halo, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, or -NHC(=O)-C₁₋₄ alkyl. In another more preferred embodiment, R² is C₆₋₁₀ aryl substituted with one or two independently selected R⁶ groups, wherein each R⁶ group is selected from -C(=O)NR⁹R¹⁰, and each R⁹ and R¹⁰ is independently selected from H and C₁₋₄ alkyl. In yet another more preferred embodiment, R² is phenyl substituted with one or two independently selected R⁶ groups. In a most preferred embodiment, R² is phenyl substituted with one or two independently selected R⁶ groups, wherein each R⁶ group is selected from halo, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, and -NHC(=O)-C₁₋₄ alkyl. In another most preferred embodiment, R² is phenyl substituted with one or two independently selected R⁶ groups, wherein each R⁶ group is selected from -C(=O)NR⁹R¹⁰, and each R⁹ and R¹⁰ is independently selected from H and C₁₋₄ alkyl. In a further most preferred embodiment R² is phenyl substituted with one or two independently selected R⁶ groups, wherein each R⁶ group is selected from F, Cl, CN, Me, -OMe, -OEt, and -NHC(=O)Me. In another further most preferred embodiment R² is phenyl substituted with one or two independently selected R⁶ groups, wherein each R⁶ group is selected from -C(=O)NH₂, and -C(=O)NHMe.

[00115] In another embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, IVa, IVb or IVc, wherein L₃ is -NR^{4b}-, and R^{4b} is as described previously. In a preferred embodiment, R^{4b} is H, Me, Et, or cyclopropyl. In a more preferred embodiment, R^{4a} is H.

[00116] In another embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, IVa, IVb or IVc, wherein R³ is C₁₋₄ alkyl substituted with C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁷ groups, or 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, optionally substituted with one or more R independently selected R⁷ groups. In a preferred embodiment, R³ is Me or Et, each of which is substituted with C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁷ groups), or 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, optionally substituted with one or more R independently selected R⁷ groups. In another preferred embodiment, R³ is C₁₋₄ alkyl substituted with phenyl,

or pyridyl, each of which is optionally substituted with one or more independently selected R^7 groups. In a more preferred embodiment, R^3 is C_{1-4} alkyl substituted with phenyl, or pyridyl. In a more preferred embodiment, R^3 is C_{1-4} alkyl substituted with phenyl, or pyridyl, each of which is substituted with Me, Et, F, or Cl. In a most preferred embodiment, R^3 is Me or Et, each of which is substituted with phenyl, or pyridyl.

5 In a more preferred embodiment, R^3 is Me or Et, each of which is substituted with phenyl, or pyridyl, each of which is substituted with Me, Et, F, or Cl.


[00117] In one embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, IVa, IVb or IVc, wherein R^3 is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O. In a preferred embodiment, R^3 is pyridyl.

10 [00118] In one embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, IVa, IVb, or IVc, wherein R^3 is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or more independently selected R^7 groups, wherein each R^7 group is as described previously. In a preferred embodiment, R^3 is pyridyl, substituted with one or more independently selected R^7 groups, wherein each R^7 group is as described previously. In another
15 preferred embodiment, R^3 is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or more independently selected R^7 groups, wherein each R^7 group is selected from Me, Et, F, and Cl. In a more preferred embodiment, R^3 is pyridyl substituted with one or more independently selected R^7 groups, wherein each R^7 group is selected from Me, Et, F, and Cl. In a most preferred embodiment, R^3 is pyridyl substituted with one R^7 group selected from Me, Et, F, and Cl.

20 [00119] In one embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, IVa, IVb, or IVc, wherein R^3 is C_{6-10} aryl. In a preferred embodiment, R^3 is phenyl.

[00120] In one embodiment, the compound of the invention is according to any one of Formulae Ia-Ic, IVa, IVb, or IVc, wherein R^3 is C_{6-10} aryl substituted with one or more independently selected R^7 groups, wherein each R^7 group is as described previously. In a preferred embodiment, R^3 is phenyl, substituted with
25 one or more independently selected R^7 groups, wherein each R^7 group is as described previously. In another preferred embodiment, R^3 is C_{6-10} aryl substituted with one or more independently selected R^7 groups, wherein each R^7 group is selected from Me, Et, F, and Cl. In a more preferred embodiment, R^3 is phenyl substituted with one or more independently selected R^7 groups, wherein each R^7 group is selected from Me, Et, F, and Cl. In a most preferred embodiment, R^3 is phenyl substituted with one R^7 group selected from Me,
30 Et, F, and Cl.


[00121] In one embodiment, the compound of the invention is according to Formula Va, Vb, Vc or Vd:



wherein R^2 is as described previously.

[00122] In a further embodiment, the compound of the invention is not according to Formula Va, Vb, Vc or Vd.


- 5 [00123] In another embodiment, the compound of the invention is according to Formula VIa, VIb, VIc or VIId:



wherein R^2 is as described previously.

[00124] In a further embodiment, the compound of the invention is not according to Formula VIa, VIb, VIc or VIId.

10 [00125] In another embodiment, the compound of the invention is according to Formula VIIa, VIIb, VIIc or VIIId:



wherein R^2 is as described previously.

- 15 [00126] In one embodiment, the compound of the invention is according to Formula Va, Vb, VIa, VIb, VIIa, or VIIb, wherein R^2 is C_{3-7} cycloalkyl. In a preferred embodiment, R^2 is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In a more preferred embodiment, R^2 is cyclopropyl.

[00127] In one embodiment, the compound of the invention is according to Formula Va, Vb, VIa, VIb, VIIa, or VIIb, wherein R^2 is not C_{3-7} cycloalkyl. In a preferred embodiment, R^2 is not cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In a more preferred embodiment, R^2 is not cyclopropyl.

[00128] In another embodiment, the compound of the invention is according to Formula Va, Vb, VIa, VIb, VIIa, or VIIb, wherein R^2 is C_{3-7} cycloalkyl substituted with one to three independently selected R^5 groups. In a preferred embodiment, R^2 is C_{3-7} cycloalkyl substituted with one R^5 group. In a more preferred embodiment, R^2 is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, each of which is substituted with one R^5 group. In another more preferred embodiment, R^2 is C_{3-7} cycloalkyl substituted with one R^5 group, wherein R^5 is oxo, or R^6 wherein R^6 is selected from OH, or C_{1-6} alkyl. In a most preferred embodiment, R^2 is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, each of which is substituted with one R^5 group, wherein R^5 is oxo, or R^6 wherein R^6 is selected from OH, and C_{1-6} alkyl. In a further most preferred embodiment, R^2 is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, each of which is substituted with one R^5 group, wherein R^5 is OH.

[00129] In another embodiment, the compound of the invention is according to Formula Va, Vb, VIa, VIb, VIIa, or VIIb, wherein R^2 is not C_{3-7} cycloalkyl substituted with one to three independently selected R^5 groups. In a preferred embodiment, R^2 is not C_{3-7} cycloalkyl substituted with one R^5 group. In a more preferred embodiment, R^2 is not cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, each of which is substituted with one R^5 group. In another more preferred embodiment, R^2 is not C_{3-7} cycloalkyl substituted with one R^5 group, wherein R^5 is oxo, or R^6 wherein R^6 is selected from OH, and C_{1-6} alkyl. In a most preferred embodiment, R^2 is not cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, each of which is substituted with one R^5 group, wherein R^5 is oxo, or R^6 wherein R^6 is selected from OH, and C_{1-6} alkyl. In a further most preferred embodiment, R^2 is not cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, each of which is substituted with one R^5 group, wherein R^5 is OH.

[00130] In one embodiment, the compound of the invention is according to Formula Vc, Vd, VIc, VId, VIIc or VIId, wherein R^2 is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O. In a preferred embodiment, R^2 is furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl.

[00131] In one embodiment, the compound of the invention is according to Formula Vc, Vd, VIc, VId, VIIc or VIId, wherein R^2 is not 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O. In a preferred embodiment, R^2 is not furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl.

[00132] In another embodiment, the compound of the invention is according to Formula Vc, Vd, VIc, VId, VIIc or VIId, wherein R^2 is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one to three independently selected R^6 groups. In a preferred

embodiment, R² is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or two independently selected R⁶ groups. In a more preferred embodiment, R² is furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl, substituted with one or two independently selected R⁶ groups. In another more preferred embodiment, R² is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or two independently selected R⁶ groups, wherein each R⁶ is independently selected from OH, halo, C₁₋₆ alkyl, C₁₋₆ alkyl substituted with one or more halo, C₁₋₆ alkoxy, -CN, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl comprising one to three heteroatoms independently selected from N, O, and S, and phenyl. In a most preferred embodiment, R² is furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl, each of which is substituted with one or two independently selected R⁶ groups, wherein each R⁶ is independently selected from OH, halo, C₁₋₆ alkyl, C₁₋₆ alkyl substituted with one or more halo, C₁₋₆ alkoxy, -CN, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl comprising one to three heteroatoms independently selected from N, O, and S, and phenyl. In another most preferred embodiment, R² is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or two independently selected R⁶ groups, wherein each R⁶ is independently selected from OH, F, Cl, Me, Et, Pr, *i*-Pr, *t*-Bu, -CF₃, -OMe, -OEt, *Oi*-Pr, -CN, cyclopropyl, pyrrolidinyl, morpholinyl, piperidinyl, and phenyl. In a further most preferred embodiment, R² is furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl, each of which is substituted with one or two independently selected R⁶ groups, wherein each R⁶ is independently selected from OH, F, Cl, Me, Et, Pr, *i*-Pr, *t*-Bu, -CF₃, -OMe, -OEt, -*Oi*-Pr, -CN, cyclopropyl, pyrrolidinyl, morpholinyl, piperidinyl, and phenyl.

[00133] In another embodiment, the compound of the invention is according to Formula Vc, Vd, VIc, VId, VIIc or VIId, wherein R² is not 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one to three independently selected R⁶ groups. In a preferred embodiment, R² is not 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or two independently selected R⁶ groups. In a more preferred embodiment, R² is not furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl, substituted with one or two independently selected R⁶ groups. In another more preferred embodiment, R² is not 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or two independently selected R⁶ groups, wherein each R⁶ is independently selected from OH, halo, C₁₋₆ alkyl, C₁₋₆ alkyl substituted with one or more halo, C₁₋₆ alkoxy, -CN, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl comprising one to three heteroatoms independently selected from N, O, and S, and phenyl. In a most preferred embodiment, R² is not furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl,

imidazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl, each of which is substituted with one or two independently selected R^6 groups, wherein each R^6 is independently selected from OH, halo, C_{1-6} alkyl, C_{1-6} alkyl substituted with one or more halo, C_{1-6} alkoxy, -CN, C_{3-7} cycloalkyl, 4-7 membered heterocycloalkyl comprising one to three heteroatoms independently selected from N, O, and S, and phenyl.

- 5 In another most preferred embodiment, R^2 is not 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or two independently selected R^6 groups, wherein each R^6 is independently selected from OH, F, Cl, Me, Et, Pr, *i*-Pr, *t*-Bu, -CF₃, -OMe, -OEt, *Oi*-Pr, -CN, cyclopropyl, pyrrolidinyl, morpholinyl, piperidinyl, and phenyl. In a further most preferred embodiment, R^2 is not furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, 10 pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl, each of which is substituted with one or two independently selected R^6 groups, wherein each R^6 is independently selected from OH, F, Cl, Me, Et, Pr, *i*-Pr, *t*-Bu, -CF₃, -OMe, -OEt, -*Oi*-Pr, -CN, cyclopropyl, pyrrolidinyl, morpholinyl, piperidinyl, and phenyl.

[00134] In another embodiment, the compound of the invention is according to Formula Vc, Vd, VIc, VId, VIId or VIId, wherein R^2 is C_{6-10} aryl. In a preferred embodiment, R^2 is phenyl.

- 15 **[00135]** In another embodiment, the compound of the invention is according to Formula Vc, Vd, VIc or VId, wherein R^2 is not C_{6-10} aryl. In a preferred embodiment, R^2 is not phenyl.

- [00136]** In another embodiment, the compound of the invention is according to Formula Vc, Vd, VIc, VId, VIId or VIId, wherein R^2 is C_{6-10} aryl substituted with one or more independently selected R^6 groups. In a preferred embodiment, R^2 is C_{6-10} aryl substituted with one or two independently selected R^6 groups. In a 20 more preferred embodiment, R^2 is C_{6-10} aryl substituted with one or two independently selected R^6 groups, wherein each R^6 group is selected from halo, CN, C_{1-6} alkyl, C_{1-6} alkoxy, and -NHC(=O)- C_{1-4} alkyl. In another more preferred embodiment, R^2 is C_{6-10} aryl substituted with one or two independently selected R^6 groups, wherein each R^6 group is selected from -C(=O)NR⁹R¹⁰, and each R⁹ and R¹⁰ is independently selected from H and C_{1-4} alkyl. In another more preferred embodiment, R^2 is phenyl substituted with one or two 25 independently selected R^6 groups. In a most preferred embodiment, R^2 is phenyl substituted with one or two independently selected R^6 groups, wherein each R^6 group is selected from halo, CN, C_{1-6} alkyl, C_{1-6} alkoxy, and -NHC(=O)- C_{1-4} alkyl. In another most preferred embodiment, R^2 is phenyl substituted with one or two independently selected R^6 groups, wherein each R^6 group is selected from -C(=O)NR⁹R¹⁰, and each R⁹ and R¹⁰ is independently selected from H and C_{1-4} alkyl. In a further most preferred embodiment R^2 is 30 phenyl substituted with one or two independently selected R^6 groups, wherein each R^6 group is selected from F, Cl, CN, Me, -OMe, -OEt, - and -NHC(=O)Me. In a further most preferred embodiment R^2 is phenyl substituted with one or two independently selected R^6 groups, wherein each R^6 group is selected from C(=O)NH₂, and -C(=O)NHMe.

- [00137]** In another embodiment, the compound of the invention is according to Formula Vc, Vd, VIc, VId, 35 VIId or VIId, wherein R^2 is not C_{6-10} aryl substituted with one or more independently selected R^6 groups. In a

preferred embodiment, R^2 is not C_{6-10} aryl substituted with one or two independently selected R^6 groups. In a more preferred embodiment, R^2 is not C_{6-10} aryl substituted with one or two independently selected R^6 groups, wherein each R^6 group is selected from halo, CN, C_{1-6} alkyl, C_{1-6} alkoxy, and $-NHC(=O)-C_{1-4}$ alkyl. In another more preferred embodiment, R^2 is not C_{6-10} aryl substituted with one or two independently selected R^6 groups, wherein each R^6 group is selected from $-C(=O)NR^9R^{10}$, and each R^9 and R^{10} is independently selected from H and C_{1-4} alkyl. In another more preferred embodiment, R^2 is not phenyl substituted with one or two independently selected R^6 groups. In a most preferred embodiment, R^2 is not phenyl substituted with one or two independently selected R^6 groups, wherein each R^6 group is selected from halo, CN, C_{1-6} alkyl, C_{1-6} alkoxy, and $-NHC(=O)-C_{1-4}$ alkyl. In another most preferred embodiment, R^2 is not phenyl substituted with one or two independently selected R^6 groups, wherein each R^6 group is selected from $-C(=O)NR^9R^{10}$, and each R^9 and R^{10} is independently selected from H and C_{1-4} alkyl. In a further most preferred embodiment R^2 is not phenyl substituted with one or two independently selected R^6 groups, wherein each R^6 group is selected from F, Cl, CN, Me, -OMe, -OEt, - and $-NHC(=O)Me$. In a further most preferred embodiment R^2 is not phenyl substituted with one or two independently selected R^6 groups, wherein each R^6 group is selected from $C(=O)NH_2$, and $-C(=O)NHMe$.

[00138] In one embodiment, the compound of the invention is selected from:

9-Allyloxy-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-3-yl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-4-yl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile,
 3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile,
 4-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile,
 [2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yloxy]-acetonitrile,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(oxazol-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(pyridin-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 9-(3,5-Dichloro-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 9-Benzofuran-2-yl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-indole-1-carboxylic acid tert-butyl ester,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(1H-indol-2-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(6-methoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(6-trifluoromethyl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-3H-imidazol-4-ylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

- 9-(5-tert-Butyl-[1,2,4]oxadiazol-3-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 5-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-pyridine-2-carboxylic acid methylamide,
- 5 2-([1,4]Dioxan-2-ylmethoxy)-9-pent-1-ynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(2-pyridin-2-yl-ethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(2-pyrazin-2-yl-ethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(1H-indol-5-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(2-methoxy-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 10 2-([1,4]Dioxan-2-ylmethoxy)-9-(5-methoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(1H-indazol-5-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(4-methoxy-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzamide,
- 5-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-2-fluoro-
- 15 benzamide,
- N-{3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-phenyl}-acetamide,
- 9-Cyclopropylethynyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(1-hydroxy-cyclopentylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-
- 20 one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-pyrimidin-5-yl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-Cyclohex-1-enyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(1-methyl-1H-indol-5-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(6-methyl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 25 2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-2-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 5-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-pent-4-yne nitrile,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(4-methoxy-phenylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 30 2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-3-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 4-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-N-methylbenzamide,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-(2-Chloro-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 35 2-([1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

- 9-(1,5-Dimethyl-1H-pyrazol-3-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(1-methyl-1H-pyrazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 5 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-[1,2,4]oxadiazol-5-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(4-morpholin-4-yl-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-4-fluoro-benzamide,
 10 3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-5-fluoro-benzamide,
 9-(3,3-Dimethyl-but-1-ynyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-4-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-isoxazol-5-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-
 15 one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-methyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(2-methoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 20 9-(3,6-Dihydro-2H-pyran-4-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 5-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-pyridine-2-carbonitrile,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(6-isopropoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(6-ethoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 25 2-([1,4]Dioxan-2-ylmethoxy)-9-(6-morpholin-4-yl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 9-(2,3-Dimethoxy-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 9-(3-Chloro-2-methoxy-pyridin-4-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 30 2-([1,4]Dioxan-2-ylmethoxy)-9-(2-methyl-pyridin-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-isonicotinonitrile,
 9-(2,5-Dimethoxy-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 35 2-([1,4]Dioxan-2-ylmethoxy)-9-(2-ethoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

- 9-(2,6-Dimethoxy-pyridin-3-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 4-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-nicotinonitrile,
 9-tert-Butoxymethyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(2-pyrrolidin-1-yl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-
 5 one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(6-pyrrolidin-1-yl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-
 one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(5-phenyl-oxazol-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-
 one,
 10 9-(5-tert-Butyl-oxazol-2-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-
 one,
 9-(5-Cyclopropyl-[1,2,4]oxadiazol-3-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-
 a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(5-ethyl-[1,2,4]oxadiazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-
 15 a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(5-methyl-[1,2,4]oxadiazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-
 a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(5-isopropyl-[1,2,4]oxadiazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-
 a]isoquinolin-4-one,
 20 9-Cyclopentylethynyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 9-Cyclohexylethynyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-hex-1-ynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 9-[3-(Benzyl-methyl-amino)-prop-1-ynyl]-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-
 25 a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-5-methyl-hex-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-
 one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 9-Cyclopropyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 30 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-4-methyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-
 4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-ethyl-3-hydroxy-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-
 one,

- 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-phenyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-(3-Benzylamino-prop-1-ynyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-[(furan-2-ylmethyl)-amino]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 5 2-([1,4]Dioxan-2-ylmethoxy)-9-(1-ethyl-1H-pyrazol-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-[1-(3-methyl-butyl)-1H-pyrazol-4-yl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(5-methyl-furan-2-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-hex-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 10 9-(3,5-Dimethyl-1H-pyrazol-4-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(1H-pyrazol-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(1-propyl-1H-pyrazol-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-[2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile,
- 15 2-[2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile,
- 9-(5-Cyclopropyl-[1,2,4]oxadiazol-3-ylmethoxy)-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-ethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-pyrimidin-2-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 20 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-phenylamino-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-pyridin-3-yl-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-Cyclopentylloxymethyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-4-methyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-
- 25 4-one,
- 9-Cyclopropylethynyl-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-imidazol-1-yl-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 30 9-(2-Cyclopropyl-ethyl)-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-Cyclopentylloxymethyl-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-pyridin-3-yl-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-Allyloxy-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 35 9-Allyloxy-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

- 2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(tetrahydro-pyran-4-yloxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-{3-[(pyridin-3-ylmethyl)-amino]-prop-1-ynyl}-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 5 2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-pentyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-Cyclopropylethynyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-(2-Cyclopropyl-ethyl)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(oxetan-3-yloxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-oxetan-3-ylmethoxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 10 a]isoquinolin-4-one,
- 9-(2,2-Dimethyl-butylamino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 15 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-ethyl-hexylamino)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-methoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-ethoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-Cyclopropylmethoxy-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-fluoro-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 20 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[3-(2-methoxy-ethoxy)-prop-1-ynyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[3-(2-ethoxy-ethoxy)-prop-1-ynyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[3-(2-fluoro-ethoxy)-prop-1-ynyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 25 a]isoquinolin-4-one,
- 9-(2,2-Dimethyl-propoxymethyl)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-Cyclohexyloxymethyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-Cyclopropylmethoxymethyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 30 one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(tetrahydro-pyran-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-(4,4-Dimethyl-pentyloxy)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 35 one,

- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-(3-Cyclopropyl-propoxy)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 5 9-Cyclohexylamino-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-4,4-dimethyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-Cyclopentylmethoxymethyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 10 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-phenylamino-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 15 9-(Cyclohexyl-methyl-amino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-(Cyclohexylmethyl-amino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[(tetrahydro-pyran-4-ylmethyl)-amino]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 20 a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-ethyl-3-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-methyl-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 25 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 9-(2,2-Dimethyl-propoxy)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(tetrahydro-pyran-4-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 30 4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(tetrahydro-pyran-4-ylmethoxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-([1,4]Dioxan-2-ylmethoxy)-9-methoxy-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
- 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(oxetan-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(3-Cyclopropyl-propoxy)-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[2-(1-hydroxy-cyclopentyl)-ethyl]-6,7-dihydro-pyrimido[6,1-

5 a]isoquinolin-4-one,

2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-tetrahydro-pyran-4-ylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

[00139] In one embodiment, the compound of the invention is selected from:

2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

10 2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-[2-(1-hydroxy-cyclopentyl)-ethyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-propoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-isopropoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-propoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

15 2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-isopropoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
and

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-methoxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00140] In one embodiment, the compound of the invention is 9-cyclopropylethynyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

20 **[00141]** In another embodiment, the compound of the invention is not 9-cyclopropylethynyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00142] In one embodiment a compound of the invention is not an isotopic variant.

[00143] In one aspect a compound of the invention is present as the free base.

[00144] In one aspect a compound of the invention is a pharmaceutically acceptable salt.

25 **[00145]** In one aspect a compound of the invention is present as the free base or a pharmaceutically acceptable salt.

[00146] In one aspect a compound of the invention is a solvate.

[00147] In one aspect a compound of the invention is a solvate of a pharmaceutically acceptable salt of the compound.

30 **[00148]** In certain aspects, the present invention provides prodrugs and derivatives of a compound of the invention according to the formula above. Prodrugs are derivatives of a compound of the invention, which have metabolically cleavable groups and become by solvolysis or under physiological conditions the compounds of the invention, which are pharmaceutically active, *in vivo*. Such examples include, but are not limited to, choline ester derivatives and the like, N-alkylmorpholine esters and the like.

[00149] Other derivatives of the compounds of this invention have activity in both their acid and acid derivative forms, but the acid sensitive form often offers advantages of solubility, tissue compatibility, or delayed release in the mammalian organism (see, Bundgard, H. Design of Prodrugs, pp. 7-9, 21-24, Elsevier, Amsterdam 1985). Prodrugs include acid derivatives well known to practitioners of the art, such as, for example, esters prepared by reaction of the parent acid with a suitable alcohol, or amides prepared by reaction of the parent acid compound with a substituted or unsubstituted amine, or acid anhydrides, or mixed anhydrides. Simple aliphatic or aromatic esters, amides and anhydrides derived from acidic groups pendant on the compounds of this invention are preferred prodrugs. In some cases it is desirable to prepare double ester type prodrugs such as (acyloxy)alkyl esters or ((alkoxycarbonyl)oxy)alkylesters. Particularly useful are the C₁ to C₈ alkyl, C₂-C₈ alkenyl, aryl, C₇-C₁₂ substituted aryl, and C₇-C₁₂ arylalkyl esters of the compounds of the invention.


[00150] While specified groups for each embodiment have generally been listed above separately, a compound of the invention includes one in which several or each embodiment in the above Formula, as well as other formulae presented herein, is selected from one or more of particular members or groups designated respectively, for each variable. Therefore, this invention is intended to include all combinations of such embodiments within its scope.

[00151] While specified groups for each embodiment have generally been listed above separately, a compound of the invention may be one for which one or more variables (for example, R groups) is selected from one or more embodiments according to any of the Formula(e) listed above. Therefore, the present invention is intended to include all combinations of variables from any of the disclosed embodiments within its scope.

[00152] Alternatively, the exclusion of one or more of the specified variables from a group or an embodiment, or combinations thereof is also contemplated by the present invention.

CLAUSES

1. A compound according to Formula Ia:



wherein

R^1 is H, Me, or halo;

L_1 is absent or is $-O-$, $-S-$, or $-NR^{4a}-$;

5 G is

R^2 ,

$-W-L_2-R^2$, or

$-W-L_3-R^3$;

10 W is C_{1-4} alkylene, C_{2-4} alkenylene having one double bond, or C_{2-4} alkynylene having one triple bond;

L_2 is absent or is $-O-$;

R^2 is

- H,

- C_{1-8} alkyl, optionally substituted with one to three groups independently selected from

15

o OH,

o halo,

o CN,

o C_{1-6} alkoxy,

o C_{3-7} cycloalkyl,

20

o 4-6 membered heterocycloalkyl comprising one to three heteroatoms independently selected from S, and O,

o 5-6 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, and

o phenyl,

25

- C_{4-7} cycloalkenyl comprising one double bond,

- 5-7 membered heterocycloalkenyl comprising one double bond, and one to three heteroatoms independently selected from N, O, and S,

- C_{3-7} cycloalkyl optionally substituted with one or more independently selected R^5 groups,


- 4-10 membered heterocycloalkyl comprising one to two heteroatoms independently selected from S, and O, optionally substituted with one to three independently selected R⁵ groups,
- 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O optionally substituted with one to three independently selected R⁶ groups, or
- 5 - C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁶ groups;
L₃ is -NR^{4b}-;
R³ is
- C₁₋₄ alkyl substituted with
 - C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁷ groups, or
 - 10 ○ 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, optionally substituted with one or more R independently selected R⁷ groups,
- 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, optionally substituted with one or more independently selected R⁷ group, or
- 15 - C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁷ groups;
Each R^{4a} and R^{4b} is independently selected from H, C₁₋₄ alkyl, and C₃₋₇ cycloalkyl;
R⁵ is oxo or R⁶;
R⁶ is
- OH,
- 20 - halo,
- -NO₂,
- C₁₋₆ alkyl optionally substituted with one to three groups independently selected from halo, and OH,
- C₁₋₆ alkoxy optionally substituted with one to three groups independently selected from halo, and
- 25 OH,
- C₃₋₇ cycloalkyl,
- -C(=O)OR⁸,
- -C(=O)NR⁹R¹⁰,
- -NHC(=O)-C₁₋₄ alkyl,
- 30 - -CN,
- phenyl,
- -O-phenyl,
- 4-7 membered heterocycloalkyl comprising one to three heteroatoms independently selected from N, O, and S, or

- 5-6 membered heteroaryl comprising one to three heteroatoms independently selected from N, O, and S, optionally substituted with one or more independently selected C₁₋₄ alkyl, C₁₋₄ alkoxy, CN, halo), and -C(=O)OR¹¹;

R⁷ is C₁₋₄ alkyl, or halo; and

- 5 each of R⁸, R⁹, R¹⁰ and R¹¹ is independently selected from H and C₁₋₄ alkyl, or a pharmaceutically acceptable salt, or a solvate, or a solvate of the pharmaceutically acceptable salt.


2. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula Ib:



Ib

10 wherein R¹, L₁ and G are as previously described.

3. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula Ic:






Ic

15 wherein R¹, L₁ and G are as previously described.




4. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-3, wherein R¹ is Me, F, or Cl.
5. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-3, wherein R¹ is H.

6. A compound or pharmaceutically acceptable salt thereof, according to clause 1 or 2, wherein the compound is according to Formula IIa, IIb or IIc:






wherein L_1 , and R^2 are as described previously.

- 5 7. A compound or pharmaceutically acceptable salt thereof, according to clause 1 or 2, wherein the compound is according to Formula IIIa, IIIb, or IIIc:



wherein L_1 , W , L_2 , and R^2 are as described previously.

- 10 8. A compound or pharmaceutically acceptable salt thereof, according to clause 1 or 2, wherein the compound is according to Formula IVa, IVb, or IVc:




wherein L_1 , W , L_3 , and R^3 are as described previously.

9. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-8, wherein L_1 is absent.
10. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-8, wherein L_1 is -O-.
- 5 11. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-8, wherein L_1 is -NR^{4a}-, and R^{4a} is H, Me, Et, or cyclopropyl.
12. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-8, wherein L_1 is -NR^{4a}-, and R^{4a} is H.
13. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-5, or 7-11,
 - 10 wherein W is C₁₋₄ alkylene.
14. A compound or pharmaceutically acceptable salt thereof, according to clause 13, wherein W is -CH₂-, -CH₂-CH₂-, -CH₂-CH₂-CH(CH₃)-, -CH₂-CH(-CH₂-CH₃)-, -CH₂-C(CH₃)₂-, or -CH₂-CH₂-CH₂-.
15. A compound or pharmaceutically acceptable salt thereof, according to clause 14, wherein W is -CH₂-CH₂-.
- 15 16. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-5, or 7-11, wherein W is C₂₋₄ alkenylene having one double bond.
17. A compound or pharmaceutically acceptable salt thereof, according to clause 16, wherein W is -CH=CH-, -CH₂-CH=CH-, or -CH=CH-CH₂.
18. A compound or pharmaceutically acceptable salt thereof, according to clause 17, wherein W is -
 - 20 CH=CH-.
19. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-5, or 7-11, wherein W is C₂₋₄ alkynylene having one triple bond.
20. A compound or pharmaceutically acceptable salt thereof, according to clause 19, wherein W is -C≡C-, -CH₂-C≡C-, or -C≡C-CH₂-.
- 25 21. A compound or pharmaceutically acceptable salt thereof, according to clause 20, wherein W is -C≡C-.
22. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-5, 7, or 9-21, wherein L_2 is -O-.
23. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-5, 7, or 9-21, wherein L_2 is absent.
- 30 24. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-5, 7, or 9-21, wherein L_1 and L_2 are absent, and W is -CH₂-, -CH₂-CH₂-, or -CH₂-CH₂-CH₂-.
25. A compound or pharmaceutically acceptable salt thereof, according to clause 1-5, 7, or 9-21, wherein L_1 and L_2 are absent, and W is -CH=CH-, -CH₂-CH=CH-, or -CH=CH-CH₂-.
26. A compound or pharmaceutically acceptable salt thereof, according to clause 1-5, 7, or 9-21, wherein
 - 35 L_1 and L_2 are absent, and W is -C≡C-, -CH₂-C≡C-, or -C≡C-CH₂-.

27. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R^2 is H.
28. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R^2 is C_{1-8} alkyl.
- 5 29. A compound or pharmaceutically acceptable salt thereof, according to clause 28, wherein R^2 is Me, Et, *n*-Pr, *i*-Pr, *i*-Bu, or *t*-Bu.
30. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R^2 is C_{1-8} alkyl substituted with one group selected from OH, halo, CN, C_{1-6} alkoxy, C_{3-7} cycloalkyl, 4-6 membered heterocycloalkyl (comprising one to three heteroatoms independently selected from S, and O), 5-6 membered heteroaryl (comprising one to three heteroatoms independently selected from N, S, and O), and phenyl.
- 10 31. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R^2 is Me, Et, *n*-Pr, *i*-Pr, *i*-Bu, or *t*-Bu, each of which is substituted with one group selected from OH, halo, CN, C_{1-6} alkoxy, C_{3-7} cycloalkyl, 4-6 membered heterocycloalkyl (comprising one to three heteroatoms independently selected from S, and O), 5-6 membered heteroaryl (comprising one to three heteroatoms independently selected from N, S, and O), and phenyl.
- 15 32. A compound or pharmaceutically acceptable salt thereof, according to any one of clause 30, wherein R^2 is C_{1-8} alkyl substituted with one group selected from OH, F, Cl, CN, -OMe, -OEt, -Oi-Pr, cyclopropyl, cyclobutyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, pyrrololyl, imidazolyl, triazolyl, oxazolyl, thiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, and phenyl.
- 20 33. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R^2 is C_{4-7} cycloalkenyl comprising one double bond.
34. A compound or pharmaceutically acceptable salt thereof, according to clause 33, wherein R^2 is cyclohexenyl.
- 25 35. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R^2 is 5-7 membered heterocycloalkenyl comprising one double bond, and one to three heteroatoms independently selected from N, O, and S.
36. A compound or pharmaceutically acceptable salt thereof, according to clause 35, wherein R^2 is dihydropyranyl.
- 30 37. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R^2 is C_{3-7} cycloalkyl.
38. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R^2 is C_{3-7} cycloalkyl substituted with one R^5 group.
39. A compound or pharmaceutically acceptable salt thereof, according to clause 38, wherein R^5 is oxo, or R^6 wherein R^6 is selected from OH, and C_{1-6} alkyl.
- 35


40. A compound or pharmaceutically acceptable salt thereof, according to clauses 37, 38 or 39, wherein R² is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.
41. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R² is 4-10 membered heterocycloalkyl comprising one to two heteroatoms independently selected from S, and O.
42. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R² is 4-10 membered heterocycloalkyl comprising one to two heteroatoms independently selected from S, and O, substituted with one R⁵ group.
43. A compound or pharmaceutically acceptable salt thereof, according to clause 42, wherein R⁵ is selected from oxo, or R⁶ wherein R⁶ is selected from OH, and C₁₋₆ alkyl.
44. A compound or pharmaceutically acceptable salt thereof, according to clause 41, 42 or 43, wherein R² is oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, or dioxanyl.
45. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R² is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O.
46. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R² is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or two independently selected R⁶ groups.
47. A compound or pharmaceutically acceptable salt thereof, according to clause 46, wherein each R⁶ is independently selected from OH, halo, C₁₋₆ alkyl, C₁₋₆ alkyl substituted with one or more halo, C₁₋₆ alkoxy, -CN, C₃₋₇ cycloalkyl, 4-7 membered heterocycloalkyl comprising one to three heteroatoms independently selected from N, O, and S, and phenyl.
48. A compound or pharmaceutically acceptable salt thereof, according to clause 45, 46 or 47, wherein R² is furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl.
49. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R² is C₆₋₁₀ aryl.
50. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-7, or 9-25, wherein R² is C₆₋₁₀ aryl, substituted with one or two independently selected R⁶ groups.
51. A compound or pharmaceutically acceptable salt thereof, according to clause 50, wherein R⁶ is selected from halo, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, -NHC(=O)-C₁₋₄ alkyl, and -C(=O)NR⁹R¹⁰, wherein each R⁹ and R¹⁰ is independently selected from H and C₁₋₄ alkyl.
52. A compound or pharmaceutically acceptable salt thereof, according to clause 49, 50 or 51, wherein R² is phenyl.

53. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-5, or 8-21, wherein L_3 is $-NR^{4b}-$, and R^{4b} is H, Me, Et, or cyclopropyl.
54. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-5, or 8-21, wherein R^3 is C_{1-4} alkyl substituted with phenyl, or pyridyl.
- 5 55. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-5, or 8-21, wherein R^3 is C_{1-4} alkyl substituted with phenyl, or pyridyl, each of which is substituted with Me, Et, F, or Cl
56. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-5, or 8-21, wherein R^3 is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O.
- 10 57. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-5, or 8-21, wherein R^3 is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or more independently selected R^7 groups.
58. A compound or pharmaceutically acceptable salt thereof, according to clause 57 wherein R^7 is selected from Me, Et, F, and Cl.
- 15 59. A compound or pharmaceutically acceptable salt thereof, according to clause 56, 57, or 58 wherein R^3 is pyridyl.
60. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-5, or 8-21, wherein R^3 is C_{6-10} aryl.
- 20 61. A compound or pharmaceutically acceptable salt thereof, according to any one of clauses 1-5, or 8-21, wherein R^3 is C_{6-10} aryl, substituted with one or more independently selected R^7 groups.
62. A compound or pharmaceutically acceptable salt thereof, according to clause 61, wherein R^7 is selected from Me, Et, F, and Cl.
63. A compound or pharmaceutically acceptable salt thereof, according to clause 60, 61, or 62, wherein R^3 is phenyl.
- 25 64. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula Va:




wherein R^2 is as described previously.

65. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula Vb:




wherein R^2 is as described previously.

- 5 66. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula Vc:




wherein R^2 is as described previously.

- 10 67. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula Vd:




wherein R^2 is as described previously.

68. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula VIa:



wherein R^2 is as described previously.


69. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula VIb:



5

wherein R^2 is as described previously.


70. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula VIc:



10


wherein R^2 is as described previously.

71. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula VId:




wherein R² is as described previously.

72. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula VIIa:




wherein R² is as described previously.

73. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula VIIb:



wherein R² is as described previously.


74. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula VIIc:



VIIc

wherein R² is as described previously.

75. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is according to Formula VIId:



VIId

wherein R² is as described previously.

76. A compound or pharmaceutically acceptable salt thereof, according to clause 64, 65, 68, 69, 72, or 73, wherein R² is C₃₋₇ cycloalkyl.
77. A compound or pharmaceutically acceptable salt thereof, according to clause 64, 65, 68, 69, 72, or 73, wherein R² is C₃₋₇ cycloalkyl substituted with one R⁵ group.
78. A compound or pharmaceutically acceptable salt thereof, according to clause 77, wherein R⁵ is oxo, or R⁶ wherein R⁶ is OH, or C₁₋₆ alkyl.
79. A compound or pharmaceutically acceptable salt thereof, according to clause 76, 77 or 78, wherein R² is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.
80. A compound or pharmaceutically acceptable salt thereof, according to clause 66, 67, 70, 71, 74, or 75, wherein R² is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O.
81. A compound or pharmaceutically acceptable salt thereof, according to clause 66, 67, 70, 71, 74, or 75, wherein R² is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, substituted with one or two independently selected R⁶ groups.
82. A compound or pharmaceutically acceptable salt thereof, according to clause 81, wherein R⁶ is selected from OH, halo, C₁₋₆ alkyl, C₁₋₆ alkyl substituted with one or more halo, C₁₋₆ alkoxy, -CN, C₃₋₇

cycloalkyl, 4-7 membered heterocycloalkyl comprising one to three heteroatoms independently selected from N, O, and S, and phenyl.

83. A compound or pharmaceutically acceptable salt thereof, according to clause 80, 81 or 82, wherein R² is furanyl, thienyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, imidazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, indanyl, or indazolyl.
84. A compound or pharmaceutically acceptable salt thereof, according to clause 66, 67, 70, 71, 74, or 75, wherein R² is C₆₋₁₀ aryl.
85. A compound or pharmaceutically acceptable salt thereof, according to clause 66, 67, 70, 71, 74, or 75, wherein R² is C₆₋₁₀ aryl substituted with one or two independently selected R⁶ groups.
86. A compound or pharmaceutically acceptable salt thereof, according to clause 85, wherein R⁶ is selected from halo, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, -NHC(=O)-C₁₋₄ alkyl, and -C(=O)NR⁹R¹⁰, and each R⁹ and R¹⁰ is independently selected from H and C₁₋₄ alkyl.
87. A compound or pharmaceutically acceptable salt thereof, according to clause 84, 85 or 86, wherein R² is phenyl.
88. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is 9-cyclopropylethynyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.
89. A compound or pharmaceutically acceptable salt thereof, according to clause 1, wherein the compound is not 9-cyclopropylethynyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

PHARMACEUTICAL COMPOSITIONS

[00153] When employed as a pharmaceutical, a compound of the invention is typically administered in the form of a pharmaceutical composition. Such compositions can be prepared in a manner well known in the pharmaceutical art and comprise at least one active compound. Generally, a compound of the invention is administered in a pharmaceutically effective amount. The amount of a compound actually administered will typically be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

[00154] The pharmaceutical compositions of the invention can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intra-articular, intravenous, intramuscular, intranasal and inhalation. Depending on the intended route of delivery, a compound of this invention is preferably

formulated as either injectable or oral compositions or as salves, as lotions or as patches all for transdermal administration.

[00155] The compositions for oral administration can take the form of bulk liquid solutions or suspensions, or bulk powders. More commonly, however, the compositions are presented in unit dosage forms to facilitate accurate dosing. The term “unit dosage forms” refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient, vehicle or carrier. Typical unit dosage forms include prefilled, premeasured ampules or syringes of the liquid compositions or pills, tablets, capsules or the like in the case of solid compositions. In such compositions, a compound of the invention is usually a minor component (from about 0.1 to about 50% by weight or preferably from about 1 to about 40% by weight) with the remainder being various vehicles or carriers and processing aids helpful for forming the desired dosing form.

[00156] Liquid forms suitable for oral administration may include a suitable aqueous or nonaqueous vehicle with buffers, suspending and dispensing agents, colorants, flavors and the like. Solid forms may include, for example, any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

[00157] Injectable compositions are typically based upon injectable sterile saline or phosphate-buffered saline or other injectable carriers known in the art. As before, the active compound in such compositions is typically a minor component, often being from about 0.05 to 10% by weight with the remainder being the injectable carrier and the like.

[00158] Transdermal compositions are typically formulated as a topical ointment or cream containing the active ingredient(s), generally in an amount ranging from about 0.01 to about 20% by weight, preferably from about 0.1 to about 20% by weight, preferably from about 0.1 to about 10% by weight, and more preferably from about 0.5 to about 15% by weight. When formulated as a ointment, the active ingredients will typically be combined with either a paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with, for example an oil-in-water cream base. Such transdermal formulations are well-known in the art and generally include additional ingredients to enhance the dermal penetration of stability of the active ingredients or the formulation. All such known transdermal formulations and ingredients are included within the scope of this invention.

[00159] A compound of the invention can also be administered by a transdermal device. Accordingly, transdermal administration can be accomplished using a patch either of the reservoir or porous membrane type, or of a solid matrix variety.

[00160] The above-described components for orally administrable, injectable or topically administrable compositions are merely representative. Other materials as well as processing techniques and the like are set forth in Part 8 of Remington's Pharmaceutical Sciences, 17th edition, 1985, Mack Publishing Company, Easton, Pennsylvania.

5 [00161] A compound of the invention can also be administered in sustained release forms or from sustained release drug delivery systems. A description of representative sustained release materials can be found in Remington's Pharmaceutical Sciences.

[00162] The following formulation examples illustrate representative pharmaceutical compositions that may be prepared in accordance with this invention. The present invention, however, is not limited to the
10 following pharmaceutical compositions.

Formulation 1 - Tablets

[00163] A compound of the invention may be admixed as a dry powder with a dry gelatin binder in an approximate 1:2 weight ratio. A minor amount of magnesium stearate may be added as a lubricant. The mixture may be formed into 240-270 mg tablets (80-90 mg of active amide compound per tablet) in a tablet
15 press.

Formulation 2 - Capsules

[00164] A compound of the invention may be admixed as a dry powder with a starch diluent in an approximate 1:1 weight ratio. The mixture may be filled into 250 mg capsules (125 mg of active amide compound per capsule).

Formulation 3 - Liquid

[00165] A compound of the invention (125 mg), may be admixed with sucrose (1.75 g) and xanthan gum (4 mg) and the resultant mixture may be blended, passed through a No. 10 mesh U.S. sieve, and then mixed with a previously made solution of microcrystalline cellulose and sodium carboxymethyl cellulose (11:89, 50 mg) in water. Sodium benzoate (10 mg), flavor, and color may be diluted with water and added with stirring.
25 Sufficient water may then be added with stirring. Sufficient water may be then added to produce a total volume of 5 mL.

Formulation 4 - Tablets

[00166] A compound of the invention may be admixed as a dry powder with a dry gelatin binder in an approximate 1:2 weight ratio. A minor amount of magnesium stearate may be added as a lubricant. The
30 mixture is formed into 450-900 mg tablets (150-300 mg of active amide compound) in a tablet press.

Formulation 5 - Injection

[00167] A compound of the invention may be dissolved or suspended in a buffered sterile saline injectable aqueous medium to a concentration of approximately 5 mg/mL.

Formulation 6 - Topical

[00168] Stearyl alcohol (250 g) and a white petrolatum (250 g) may be melted at about 75°C and then a mixture of a compound of the invention (50 g) methylparaben (0.25 g), propylparaben (0.15 g), sodium lauryl sulfate (10 g), and propylene glycol (120 g) dissolved in water (about 370 g) may be added and the resulting mixture may be stirred until it congeals.

METHODS OF TREATMENT

[00169] A compound of the invention may be used as a therapeutic agent for the treatment of conditions in mammals that are causally related or attributable to aberrant activity of GPR84 and/or aberrant GPR84 expression and/or aberrant GPR84 distribution.

[00170] Accordingly, a compound and pharmaceutical compositions of the invention find use as therapeutics for the prophylaxis and/or treatment of inflammatory conditions (e.g. inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions, in mammals including humans.

[00171] Accordingly, in one aspect, the present invention provides the compound of the invention, or a pharmaceutical composition comprising the compound of the invention for use as a medicament.

[00172] In another aspect, the present invention provides the compound of the invention, or a pharmaceutical composition comprising the compound of the invention for use in the manufacture of a medicament.

[00173] In yet another aspect, the present invention provides a method of treating a mammal having, or at risk of having a disease disclosed herein. In a particular aspect, the present invention provides a method of treating a mammal having, or at risk of having inflammatory conditions (e.g. inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions, in mammals including humans.

[00174] In one aspect, the present invention provides the compound of the invention, or a pharmaceutical composition comprising the compound of the invention for use as a medicine for the prophylaxis and/or treatment of inflammatory conditions. In a specific embodiment, the inflammatory condition is selected from inflammatory bowel disease (IBD), rheumatoid arthritis, vasculitis, chronic obstructive pulmonary disease (COPD), and idiopathic pulmonary fibrosis (IPF).

[00175] In another aspect, the present invention provides the compound of the invention, or a pharmaceutical composition comprising the compound of the invention for use in the manufacture of a medicament for the prophylaxis and/or treatment of inflammatory conditions. In a specific embodiment, the inflammatory condition is selected from inflammatory bowel disease (IBD), rheumatoid arthritis, vasculitis, chronic obstructive pulmonary disease (COPD), and idiopathic pulmonary fibrosis (IPF).

[00176] In additional method of treatment aspects, this invention provides methods of treatment and/or prophylaxis of a mammal susceptible to or afflicted with inflammatory conditions, which method comprises administering an effective amount of a compound of the invention, or one or more of the pharmaceutical compositions herein described. In a specific embodiment, the inflammatory condition is selected from inflammatory bowel disease (IBD), rheumatoid arthritis, vasculitis, chronic obstructive pulmonary disease (COPD), and idiopathic pulmonary fibrosis (IPF).

[00177] In another aspect, the present invention provides a method of treating a mammal having, or at risk of having a disease selected from inflammatory conditions (for example inflammatory bowel diseases (IBD), rheumatoid arthritis, vasculitis, lung diseases (e.g. chronic obstructive pulmonary disease (COPD) and lung interstitial diseases (e.g. idiopathic pulmonary fibrosis (IPF))), neuroinflammatory conditions, infectious diseases, autoimmune diseases, endocrine and/or metabolic diseases, and/or diseases involving impairment of immune cell functions.

[00178] In one aspect, the present invention provides the compound of the invention, or a pharmaceutical composition comprising the compound of the invention for use as a medicine for the prophylaxis and/or treatment of neuroinflammatory conditions, Guillain-Barré syndrome (GBS), multiple sclerosis, axonal degeneration, autoimmune encephalomyelitis.

[00179] In another aspect, the present invention provides the compound of the invention, or a pharmaceutical composition comprising the compound of the invention for use in the manufacture of a medicament for the prophylaxis and/or treatment of neuroinflammatory conditions, Guillain-Barré syndrome (GBS), multiple sclerosis, axonal degeneration, autoimmune encephalomyelitis.

[00180] In additional method of treatment aspects, this invention provides methods of treatment and/or prophylaxis of a mammal susceptible to or afflicted with neuroinflammatory conditions, Guillain-Barré syndrome (GBS), multiple sclerosis, axonal degeneration, autoimmune encephalomyelitis, which method comprises administering an effective amount of a compound of the invention, or one or more of the pharmaceutical compositions herein described.

[00181] In one aspect, the present invention provides the compound of the invention, or a pharmaceutical composition comprising the compound of the invention for use as a medicine for the prophylaxis and/or treatment of infectious disease. In a specific embodiment, the infectious diseases is selected from sepsis, septicemia, endotoxemia, systemic inflammatory response syndrome (SIRS), gastritis, enteritis, enterocolitis,

tuberculosis, and other infections involving, for example, Yersinia, Salmonella, Chlamydia, Shigella, enterobacteria species.

[00182] In another aspect, the present invention provides the compound of the invention, or a pharmaceutical composition comprising the compound of the invention for use in the manufacture of a medicament for the prophylaxis and/or treatment of infectious disease. In a specific embodiment, the infectious diseases is selected from sepsis, septicemia, endotoxemia, systemic inflammatory response syndrome (SIRS), gastritis, enteritis, enterocolitis, tuberculosis, and other infections involving, for example, Yersinia, Salmonella, Chlamydia, Shigella, enterobacteria species.

[00183] In additional method of treatment aspects, this invention provides methods of treatment and/or prophylaxis of a mammal susceptible to or afflicted with infectious disease, which method comprises administering an effective amount of a compound of the invention, or one or more of the pharmaceutical compositions herein described. In a specific embodiment, the infectious diseases is selected from sepsis, septicemia, endotoxemia, systemic inflammatory response syndrome (SIRS), gastritis, enteritis, enterocolitis, tuberculosis, and other infections involving, for example, Yersinia, Salmonella, Chlamydia, Shigella, enterobacteria species.

[00184] In one aspect, the present invention provides the compound of the invention, or a pharmaceutical composition comprising the compound of the invention for use as a medicine for the prophylaxis and/or treatment of autoimmune diseases, and/or diseases involving impairment of immune cell functions. In a specific embodiment, the autoimmune diseases and/or diseases involving impairment of immune cell functions is selected from COPD, asthma, psoriasis, systemic lupus erythematosus, type I diabetes mellitus, vasculitis and inflammatory bowel disease.

[00185] In another aspect, the present invention provides the compound of the invention, or a pharmaceutical composition comprising the compound of the invention for use in the manufacture of a medicament for the prophylaxis and/or treatment of autoimmune diseases and/or diseases involving impairment of immune cell functions. In a specific embodiment, the autoimmune diseases, and/or diseases involving impairment of immune cell functions is selected from COPD, asthma, psoriasis, systemic lupus erythematosus, type I diabetes mellitus, vasculitis and inflammatory bowel disease.

[00186] In additional method of treatment aspects, this invention provides methods of treatment and/or prophylaxis of a mammal susceptible to or afflicted with autoimmune diseases and/or diseases involving impairment of immune cell functions, which method comprises administering an effective amount of a compound of the invention, or one or more of the pharmaceutical compositions herein described. In a specific embodiment, the autoimmune diseases and/or diseases involving impairment of immune cell functions is selected from COPD, asthma, psoriasis, systemic lupus erythematosus, type I diabetes mellitus, vasculitis and inflammatory bowel disease.

[00187] In one aspect, the present invention provides the compound of the invention, or a pharmaceutical composition comprising the compound of the invention for use as a medicine for the prophylaxis and/or treatment of endocrine and/or metabolic diseases. In a specific embodiment, the endocrine and/or metabolic diseases is selected from hypothyroidism, congenital adrenal hyperplasia, diseases of the parathyroid gland, diabetes mellitus, diseases of the adrenal glands (including Cushing's syndrome and Addison's disease), ovarian dysfunction (including polycystic ovary syndrome), cystic fibrosis, phenylketonuria (PKU), diabetes, hyperlipidemia, gout, and rickets.

[00188] In another aspect, the present invention provides the compound of the invention, or a pharmaceutical composition comprising the compound of the invention for use in the manufacture of a medicament for the prophylaxis and/or treatment of endocrine and/or metabolic diseases. In a specific embodiment, the endocrine and/or metabolic diseases is selected from hypothyroidism, congenital adrenal hyperplasia, diseases of the parathyroid gland, diabetes mellitus, diseases of the adrenal glands (including Cushing's syndrome and Addison's disease), ovarian dysfunction (including polycystic ovary syndrome), cystic fibrosis, phenylketonuria (PKU), diabetes, hyperlipidemia, gout, and rickets.

[00189] In additional method of treatment aspects, this invention provides methods of treatment and/or prophylaxis of a mammal susceptible to or afflicted with endocrine and/or metabolic diseases, which method comprises administering an effective amount of a compound of the invention, or one or more of the pharmaceutical compositions herein described. In a specific embodiment, the endocrine and/or metabolic diseases is selected from hypothyroidism, congenital adrenal hyperplasia, diseases of the parathyroid gland, diabetes mellitus, diseases of the adrenal glands (including Cushing's syndrome and Addison's disease), ovarian dysfunction (including polycystic ovary syndrome), cystic fibrosis, phenylketonuria (PKU), diabetes, hyperlipidemia, gout, and rickets.

[00190] As a further aspect of the invention there is provided a compound of the invention for use as a medicament especially in the treatment or prevention of the aforementioned conditions and diseases. Also provided herein is the use of the compound in the manufacture of a medicament for the treatment or prevention of one of the aforementioned conditions and diseases.

[00191] A particular regimen of the present method comprises the administration to a subject in suffering from an inflammatory condition, of an effective amount of a compound of the invention for a period of time sufficient to reduce the level of inflammation in the subject, and preferably terminate, the processes responsible for said inflammation. A special embodiment of the method comprises administering of an effective amount of a compound of the invention to a subject suffering from or susceptible to the development of inflammatory condition, for a period of time sufficient to reduce or prevent, respectively, inflammation of said patient, and preferably terminate, the processes responsible for said inflammation.

[00192] Injection dose levels range from about 0.1 mg/kg/h to at least 10 mg/kg/h, all for from about 1 to about 120 h and especially 24 to 96 h. A preloading bolus of from about 0.1 mg/kg to about 10 mg/kg or

more may also be administered to achieve adequate steady state levels. The maximum total dose is not expected to exceed about 2 g/day for a 40 to 80 kg human patient.

[00193] Transdermal doses are generally selected to provide similar or lower blood levels than are achieved using injection doses.

5 [00194] When used to prevent the onset of a condition, a compound of the invention will be administered to a patient at risk for developing the condition, typically on the advice and under the supervision of a physician, at the dosage levels described above. Patients at risk for developing a particular condition generally include those that have a family history of the condition, or those who have been identified by genetic testing or screening to be particularly susceptible to developing the condition.

10 [00195] A compound of the invention can be administered as the sole active agent or it can be administered in combination with other therapeutic agents, including other compounds that demonstrate the same or a similar therapeutic activity, and that are determined to be safe and efficacious for such combined administration. In a specific embodiment, co-administration of two (or more) agents allows for significantly lower doses of each to be used, thereby reducing the side effects seen.

15 [00196] In one embodiment, a compound of the invention is co-administered with another therapeutic agent for the treatment and/or prevention of an inflammatory condition; particular agents include, but are not limited to, immunoregulatory agents e.g. azathioprine, corticosteroids (e.g. prednisolone or dexamethasone), cyclophosphamide, cyclosporin A, tacrolimus, Mycophenolate Mofetil, muromonab-CD3 (OKT3, e.g. Orthoclone®), ATG, aspirinTM, acetaminophen, ibuprofen, naproxen, and piroxicam.

20 [00197] In one embodiment, a compound of the invention is co-administered with another therapeutic agent for the treatment and/or prevention of arthritis (e.g. rheumatoid arthritis); particular agents include but are not limited to analgesics, non-steroidal anti-inflammatory drugs (NSAIDS), steroids, synthetic DMARDS (for example but without limitation methotrexate, leflunomide, sulfasalazine, auranofin, sodium aurothiomalate, penicillamine, chloroquine, hydroxychloroquine, azathioprine, and cyclosporin), and
25 biological DMARDS (for example but without limitation Infliximab, Etanercept, Adalimumab, Rituximab, Golimumab, Certolizumab pegol, Tocilizumab, Interleukin 1 blockers and Abatacept).

[00198] In one embodiment, a compound of the invention is co-administered with another therapeutic agent for the treatment and/or prevention of autoimmune diseases; particular agents include but are not limited to: glucocorticoids, cytostatic agents (e.g. purine analogs), alkylating agents, (e.g. nitrogen mustards
30 (cyclophosphamide), nitrosoureas, platinum compounds, and others), antimetabolites (e.g. methotrexate, azathioprine and mercaptopurine), cytotoxic antibiotics (e.g. dactinomycin anthracyclines, mitomycin C, bleomycin, and mithramycin), antibodies (e.g., anti-CD20, anti-CD25 or anti-CD3 (OTK3) monoclonal antibodies, Atgam® and Thymoglobuline®), cyclosporin, tacrolimus, rapamycin (sirolimus), interferons (e.g. IFN-β), TNF binding proteins (e.g. infliximab (RemicadeTM), etanercept (EnbrelTM), or adalimumab
35 (HumiraTM)), mycophenolate, Fingolimod, and Myriocin.

[00199] In one embodiment, a compound of the invention is co-administered with another therapeutic agent for the treatment and/or prevention of infectious diseases; particular agents include but are not limited to antibiotics. In a particular embodiment, a compound of the invention is co-administered with another therapeutic agent for the treatment and/or prevention of infections of any organ of the human body; particular agents include but are not limited to: aminoglycosides, ansamycins, carbacephem, carbapenems, cephalosporins, glycopeptides, lincosamides, macrolides, monobactams, nitrofurans, penicillins, polypeptides, quinolones, sulfonamides, tetracyclins, anti-mycobacterial agents, as well as chloramphenicol, fosfomycin, linezolid, metronidazole, mupirocin, rifamycin, thiamphenicol and tinidazole.

[00200] In one embodiment, a compound of the invention is co-administered with another therapeutic agent for the treatment and/or prevention of vasculitis, particular agents include but are not limited to steroids (for example prednisone, prednisolone), cyclophosphamide and eventually antibiotics in case of cutaneous infections (for example cephalixin)

[00201] In one embodiment, a compound of the invention is co-administered with another therapeutic agent for the treatment and/or prevention of IPF, particular agents include but are not limited to pirfenidone and bosentan.

[00202] In one embodiment, a compound of the invention is co-administered with another therapeutic agent for the treatment and/or prevention of asthma and/or rhinitis and/or COPD; particular agents include but are not limited to: β_2 -adrenoceptor agonists (e.g. salbutamol, levalbuterol, terbutaline and bitolterol), epinephrine (inhaled or tablets), anticholinergics (e.g. ipratropium bromide), glucocorticoids (oral or inhaled) Long-acting β_2 -agonists (e.g. salmeterol, formoterol, bambuterol, and sustained-release oral albuterol), combinations of inhaled steroids and long-acting bronchodilators (e.g. fluticasone/salmeterol, budesonide/formoterol), leukotriene antagonists and synthesis inhibitors (e.g. montelukast, zafirlukast and zileuton), inhibitors of mediator release (e.g. cromoglycate and ketotifen), phosphodiesterase-4 inhibitors (e.g. Roflumilast), biological regulators of IgE response (e.g. omalizumab), antihistamines (e.g. ceterizine, cinnarizine, fexofenadine), and vasoconstrictors (e.g. oxymethazoline, xylomethazoline, nafazoline and tramazoline).

[00203] Additionally, a compound of the invention may be administered in combination with emergency therapies for asthma and/or COPD, such therapies include oxygen or heliox administration, nebulized salbutamol or terbutaline (optionally combined with an anticholinergic (e.g. ipratropium), systemic steroids (oral or intravenous, e.g. prednisone, prednisolone, methylprednisolone, dexamethasone, or hydrocortisone), intravenous salbutamol, non-specific β -agonists, injected or inhaled (e.g. epinephrine, isoetharine, isoproterenol, metaproterenol), anticholinergics (IV or nebulized, e.g. glycopyrrolate, atropine, ipratropium), methylxanthines (theophylline, aminophylline, bamiphylline), inhalation anesthetics that have a bronchodilatory effect (e.g. isoflurane, halothane, enflurane), ketamine, and intravenous magnesium sulfate.

[00204] In one embodiment, a compound of the invention is co-administered with another therapeutic agent for the treatment and/or prevention of inflammatory bowel disease (IBD); particular agents include but are not limited to: glucocorticoids (e.g. prednisone, budesonide) synthetic disease modifying, immunomodulatory agents (e.g. methotrexate, leflunomide, sulfasalazine, mesalazine, azathioprine, 6-mercaptapurine and ciclosporin) and biological disease modifying, immunomodulatory agents (infliximab, adalimumab, rituximab, and abatacept).

[00205] By co-administration is included any means of delivering two or more therapeutic agents to the patient as part of the same treatment regime, as will be apparent to the skilled person. Whilst the two or more agents may be administered simultaneously in a single formulation this is not essential. The agents may be administered in different formulations and at different times.

GENERAL SYNTHETIC PROCEDURES

General

[00206] A compound of the invention can be prepared from readily available starting materials using the following general methods and procedures. It will be appreciated that where typical or preferred process conditions (i.e., reaction temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given, other process conditions can also be used unless otherwise stated. Optimum reaction conditions may vary with the particular reactants or solvent used, but such conditions can be determined by one skilled in the art by routine optimization procedures.

[00207] Additionally, as will be apparent to those skilled in the art, conventional protecting groups may be necessary to prevent certain functional groups from undergoing undesired reactions. The choice of a suitable protecting group for a particular functional group as well as suitable conditions for protection and deprotection are well known in the art. For example, numerous protecting groups, and their introduction and removal, are described in T. W. Greene and P. G. M. Wuts, *Protecting Groups in Organic Synthesis*, Wiley-Blackwell; 4th Revised edition edition (2006), and references cited therein.

[00208] The following methods are presented with details as to the preparation of representative 6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one that have been listed hereinabove. A compound of the invention may be prepared from known or commercially available starting materials and reagents by one skilled in the art of organic synthesis.

[00209] All reagents were of commercial grade and were used as received without further purification, unless otherwise stated. Commercially available anhydrous solvents were used for reactions conducted under inert atmosphere. Reagent grade solvents were used in all other cases, unless otherwise specified. Column chromatography was performed on silica standard (30-70 μm). Thin layer chromatography was carried out using pre-coated silica gel 60 F-254 plates (thickness 0.25 mm). ^1H NMR spectra were recorded on a Bruker Advance 400 NMR spectrometer (400 MHz) or a Bruker Advance 300 NMR spectrometer (300 MHz).

Chemical shifts (δ) for ^1H NMR spectra are reported in parts per million (ppm) relative to tetramethylsilane (δ 0.00) or the appropriate residual solvent peak as internal reference. Multiplicities are given as singlet (s), doublet (d), doublet of doublet (dd), triplet (t), quartet (q), multiplet (m) and broad (br). Electrospray MS spectra were obtained either on a Waters platform LC/MS spectrometer or on an Agilent 1100 Series

5 LC/MSD. Analytic LCMS: Columns used, Waters Acquity UPLC BEH C18 1.7 μm , 2.1mm ID x 50mm L or Waters Acquity UPLC BEH C18 1.7 μm , 2.1mm ID x 30mm L or Waters XBridge C18 3.5 μm , 2.1mm ID x 50mm L. All the methods are using MeCN/H₂O gradients. MeCN and H₂O contain either 0.1% Formic Acid or NH₃ (10mM). Preparative LCMS: Column used, Waters XBridge Prep C18 5 μm ODB 30mm ID x 100mm L. All the methods are using either MeOH/H₂O or MeCN/H₂O gradients. MeOH, MeCN and H₂O contain

10 either 0.1% Formic Acid or 0.1% Diethylamine. Analytic chiral LC: Column used, Chiralpak IA 5 μm 250 x 4.6 mm. Microwave heating was performed with a Biotage Initiator.

[00210] Table I: List of abbreviations used in the experimental section:

μL	microliter
AcOH	Acetic acid
Aq.	aqueous
ATP	Adenosine 5'-Triphosphate
BINAP	2,2'-bis(diphenylphosphino)-1,1'-binaphthyl
Boc	tert-Butyloxy-carbonyl
Boc ₂ O	Di- <i>tert</i> -butyl dicarbonate
br s	broad singlet
Calcd	calculated
Cat.	Catalytic amount
D	doublet
Dd	Doublet of doublet
DCC	N,N'-Dicyclohexylcarbodiimide
DCE	1,2-Dichloroethane
DCM	Dichloromethane
DIAD	Diisopropyl azodicarboxylate
DIPEA	N,N-diisopropylethylamine

DMAP	4-Dimethylaminopyridine
DME	Dimethoxyethane
DMF	N,N-dimethylformamide
DMSO	Dimethylsulfoxide
DPBS	Dulbecco's Phosphate-Buffered Saline
DPPF	1,1'-Bis(diphenylphosphino)ferrocene
EtOAc	Ethyl acetate
Et ₂ O	Diethyl ether
eq.	equivalent
g	gram
GTP \square S	guanosine 5'-O-[gamma-thio]triphosphate
h	hour
H	Heptane
HPLC	High-performance liquid chromatography
<i>i</i> PrOH	isopropanol
<i>i</i> Pr ₂ O	Diisopropyl ether


KHMDS	Potassium hexamethyldisilazane
LCMS	Liquid Chromatography- Mass Spectrometry
L	Liter
M	multiplet
MeOH	Methanol
MeCN	Acetonitrile
MeI	Methyl iodide
MEK	Methyl ethyl ketone
Mg	milligram
Min	minute
mL	milliliter
Mmol	millimole
MS	mass spectrometry
MW	Molecular weight
MW (calc)	Molecular weight calculated
MW (obs)	Molecular weight observed
NADP	Nicotinamide adenine dinucleotide phosphate
NEAA	Non-Essential Amino Acid
NMP	<i>N</i> -Methyl-2-pyrrolidone
NMR	Nuclear Magnetic Resonance

obsd	observed
Pd(OAc) ₂	Palladium(II) acetate
Pd(PPh ₃) ₄	Tetrakis(triphenylphosphine)palladium(0)
Pd/C	Palladium on Carbon 10%
ppm	part-per-million
q	quadruplet
rpm	revolutions per minute
RT	Room temperature
Rt	retention time
RuPhos	2-Dicyclohexylphosphino-2',6'-di- <i>i</i> -propoxy-1,1'-biphenyl
s	singlet
SM	Starting material
spA	Scintillation proximity assay
SPE	Solid phase extraction
STAB	sodiumtriacetoxymborohydride
t	triplet
TBAF	Tetra- <i>n</i> -butylammonium fluoride
TEA	Triethylamine
TFA	Trifluoroacetic acid
THF	Tetrahydrofuran
TLC	Thin layer chromatography

General Synthetic Method

Intermediates

[00211] The intermediates to prepare the compounds according to the invention can be produced according to the following schemes.



Intermediate 1: [2-(3-methoxy-phenyl)-ethyl]-urea

[00212] A solution of 3-methoxyphenethylamine (100 g, 661.3 mmol, 1 eq.), urea (157.3 g, 2619.0 mmol, 4 eq.), AcOH (36 mL) and aq. HCl (12 mL) in H₂O (800 mL) was heated under reflux for 5 days. The

reaction mixture was cooled to RT, the solid was filtered off, washed with water and dried to afford intermediate 1.

(¹H, CDCl₃) δ (ppm): 7.24 (1H, t), 6.82-6.77 (3H, m), 5.10 (1H, br s), 4.52 (2H, br), 3.81 (1H, s), 3.42 (2H, br t), 2.80 (2H, t)

5

Intermediate 2: [2-(3-hydroxy-phenyl)-ethyl]-urea

[00213] A solution of intermediate 1 (72 g, 370.7 mmol) in concentrated HBr (500 mL) was heated under reflux overnight. The reaction mixture was brought to basic pH by addition of NaHCO₃ and extracted with EtOAc. The organic layer was dried over MgSO₄ and concentrated under vacuum to afford intermediate 2.

10 (¹H, MeOD-*d*4) δ (ppm): 7.15 (1H, t), 6.76-6.68 (3H, m), 3.40-3.36 (2H, t), 2.77-2.74 (2H, t)

Intermediate 3: [2-(3-allyloxy-phenyl)-ethyl]-urea

[00214] To a solution of intermediate 2 (45 g, 249.7 mmol, 1 eq.) and K₂CO₃ (103.5 g, 749.1 mmol, 3 eq.) in anhydrous DMF (300 mL) under a nitrogen atmosphere, was added allylbromide (50.5 mL, 499.4 mmol, 2 eq.). The reaction mixture was stirred for 2.5 days, then DMF was evaporated to dryness. The residue was dissolved in EtOAc, washed with saturated Na₂CO₃, brine, dried over MgSO₄ and concentrated under vacuum to afford intermediate 3.

15

(¹H, MeOD-*d*4) δ (ppm): 7.24 (1H, t), 6.87-6.81 (3H, m), 6.16-6.06 (1H, m), 5.45 (1H, dd), 5.29 (1H, dd), 4.59-4.57 (2H, m), 3.38 (2H, t), 2.80 (2H, t)

20

Intermediate 4: 1-[2-(3-allyloxy-phenyl)-ethyl]-pyrimidine-2,4,6-trione

[00215] Sodium (20.06 g, 872 mmol, 1 eq.) was dissolved in EtOH (1.4 L). Diethyl malonate (132.4 mL, 872 mmol, 1 eq.) was added and the reaction mixture was heated under reflux for 1h. Intermediate 3 (96 g, 436 mmol, 0.5 eq.) in EtOH (300 mL) was added and the reaction mixture was heated under reflux for 12 h. The reaction was cooled to RT, 1N aq. HCl was added and the precipitate was filtered, washed with water and dried to afford intermediate 4.

25

(¹H, CDCl₃) δ (ppm): 8.40 (1H, br s), 7.25 (1H, t), 6.88-6.82 (3H, m), 6.14-6.04 (1H, m), 5.45 (1H, dd), 5.32 (1H, dd), 4.58-4.56 (2H, m), 4.13 (2H, t), 3.64 (2H, s), 2.92 (2H, t)

MW (calcd): 288.3; MW (obsd): 289.3 (M + 1)

30

Intermediate 5: 9-allyloxy-2-chloro-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one

[00216] A solution of intermediate 4 (20 g, 69.4 mmol, 1 eq.) in POCl₃ (150 mL) was stirred at 50°C for 3 days. POCl₃ was evaporated under vacuum and the residue was dissolved in DCM and quenched with saturated NaHCO₃. The organic layer was washed with water, dried over MgSO₄ and concentrated to afford intermediate 5.

35


(¹H, CDCl₃) δ (ppm): 7.71 (2H, d), 6.97 (1H, dd), 6.86 (1H, d), 6.71 (1H, s), 6.13-6.04 (1H, m), 5.47 (1H, dd), 5.36 (1H, dd), 4.67-4.65 (2H, m), 4.27 (2H, t), 3.05 (2H, t)

MW (calcd): 288.7; MW (obsd): 289.3 (M+1)

5

General methods

General method A:



[00217] To a solution of NaH (2 eq., 60% in mineral oil) in anhydrous DCM at 0°C, is added 2-hydroxymethyl-[1,4]dioxane (2 eq.) with the appropriate chirality, after 15 min, intermediate **5** (1 eq.) is added at 0°C, and the reaction is stirred at RT until completion. Saturated NH₄Cl is added to the reaction mixture, the organic layer is washed with water, dried over MgSO₄ and concentrated. The desired product is purified by flash chromatography on silica gel.

[00218] 2-Hydroxymethyl-[1,4]dioxane, (R) 2-hydroxymethyl-[1,4]dioxane and (S) 2-hydroxymethyl-[1,4]dioxane are commercially available or can easily be prepared [Young Kim *et al* ; Bioorganic & Medicinal Chemistry 15 (2007) 2667–2679].

Illustrative synthesis of general method A:

Compound 118: 9-allyloxy-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one

[00219] A solution of (R) 2-hydroxymethyl-[1,4]dioxane (56.6 g, 479 mmol, 2 eq.) and NaH (19.9 g, 479 mmol, 2 eq., 60% in mineral oil) in anhydrous DCM (300 mL) was stirred for 30 min at 0°C. Intermediate **5** (69.2 g, 240 mmol, 1 eq.) in solution in anhydrous DCM (700 mL) was added at 0°C. The reaction mixture was stirred for 2 h. Saturated NH₄Cl was added, the organic layer was washed with water, dried over MgSO₄ and evaporated to dryness. The crude product was purified by flash chromatography on silica gel (MeOH/DCM) to afford compound **118**.

(¹H, CDCl₃) δ (ppm): 7.66 (1H, d), 6.94 (1H, dd), 6.83 (1H, d), 6.32 (1H, s), 6.15-6.03 (1H, m), 5.47 (1H, dd), 5.37 (1H, dd), 4.65-4.63 (2H, m), 4.51-4.39 (2H, m), 4.23 (2H, t), 4.06-3.98 (1H, m), 3.92-3.47 (6H, m), 3.01 (2H, t)


30

Compound 1: 9-allyloxy-2-(-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one

[00220] Compound 1 was prepared using general method A starting from 2-hydroxymethyl-[1,4]dioxane.
 $(^1\text{H}, \text{CDCl}_3)$ δ (ppm): 7.66 (1H, d), 6.94 (1H, dd), 6.83 (1H, d), 6.32 (1H, s), 6.11-6.04 (1H, m), 5.47 (1H, dd), 5.35 (1H, dd), 4.65-4.63 (2H, m), 4.491-4.40 (2H, m), 4.22 (2H, t), 4.02-3.99 (1H, m), 3.90-3.46 (6H, m), 3.00 (2H, t)
 MW (calcd): 370.4; MW (obsd): 371.4 (M+1)

Compound 117: 9-allyloxy-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one

[00221] Compound 117 was prepared using general method A starting from (S) 2-hydroxymethyl-[1,4]dioxane.
 $(^1\text{H}, \text{CDCl}_3)$ δ (ppm): 7.66 (1H, d), 6.94 (1H, dd), 6.83 (1H, d), 6.31 (1H, s), 6.12-6.04 (1H, m), 5.49 (1H, dd), 5.36 (1H, dd), 4.65-4.63 (2H, m), 4.50-4.42 (2H, m), 4.23 (2H, t), 4.04-4.00 (1H, m), 3.91-3.50 (6H, m), 3.01 (2H, t)
 MW (calcd): 370.4; MW (obsd): 371.2 (M+1)

General method B:

[00222] To a suspension of compound 1, 117 or 118 (1 eq.) in a mixture of DCM/MeOH (1/1) is added K_2CO_3 (2 eq.) and $\text{Pd(PPh}_3)_4$ (0.05 eq.). The reaction mixture is degassed before stirring at RT. After completion, water is added to the reaction mixture and the aqueous layer is separated. The pH of the aqueous solution is adjusted to pH 1 with 2M aq. HCl. The precipitate is filtered off, washed with water and dried to afford intermediate 6, 7 or 8.

Illustrative synthesis of general method B:**Intermediate 6: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-hydroxy-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one**

[00223] To a suspension of compound 118 (31.15 g, 84.2 mmol, 1 eq.) in a mixture of DCM/MeOH (1/1, 800 mL) was added K_2CO_3 (23.2 g, 138.2 mmol, 2 eq.) and $\text{Pd(PPh}_3)_4$ (4.86 g, 4.21 mmol, 0.05 eq.). The reaction mixture was stirred at RT for 2 h. Water (800 mL) was added, and the aqueous layer was separated.

The pH of the aqueous solution is adjusted to pH 1 with 2 M aq. HCl. The precipitate was filtered off, washed with water and dried to afford intermediate **6**.

(¹H, DMSO-*d*₆) δ (ppm): 7.84 (1H, d), 6.77 (1H, dd), 6.74 (1H, d), 6.45 (1H, s), 4.25-4.23 (2H, m), 3.99 (2H, t), 3.87-3.75 (3H, m), 3.68-3.58 (2H, m), 3.52-3.46 (1H, m), 3.40-3.30 (1H, m), 2.91 (2H, t)

5 MW (calcd): 330.4; MW (obsd): 331.3 (M+1)

Intermediate 7: 2-((R)-1-[1,4]dioxan-2-ylmethoxy)-9-hydroxy-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one

[00224] Intermediate **7** was prepared using general method **B** starting from compound **117**.

10 (¹H, DMSO-*d*₆) δ (ppm): 7.83 (1H, d), 6.77 (1H, dd), 6.74 (1H, d), 6.44 (1H, s), 4.27-4.20 (2H, m), 3.98 (2H, t), 3.87-3.74 (3H, m), 3.68-3.57 (2H, m), 3.52-3.46 (1H, m), 3.40-3.34 (1H, m), 2.91 (2H, t)

Intermediate 8: 2-([1,4]dioxan-2-ylmethoxy)-9-hydroxy-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one

[00225] Intermediate **8** was prepared using general method **B** starting from compound **1**.

15 (¹H, DMSO-*d*₆) δ (ppm): 7.84 (1H, d), 6.77 (1H, dd), 6.74 (1H, d), 6.45 (1H, s), 4.27-4.20 (2H, m); 3.99 (2H, t), 3.88-3.73 (3H, m), 3.68-3.58 (2H, m), 3.52-3.46 (1H, m), 3.40-3.34 (1H, m), 2.91 (2H, t)

MW (calcd): 330.4; MW (obsd): 331.0 (M+1)

General method C:

20 [00226] A solution of intermediate **6**, **7** or **8** (1 eq.), N-phenyl-bis(trifluoromethanesulfonimide) (1.2 eq.) and Et₃N (1.3 eq.) in DCM under nitrogen is stirred at RT until completion. The reaction mixture is concentrated and the crude is purified by crystallization from *i*PrOH to afford intermediate **9**, **10** or **11**.

Illustrative synthesis of general method C:

25 **Intermediate 9: trifluoro-methanesulfonic acid 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl ester**

[00227] A solution of intermediate **6**, (24 g, 72.7 mmol, 1 eq.), N-phenyl-bis(trifluoromethanesulfonimide) (31.15 g, 87.2 mmol, 1.2 eq.) and Et₃N (13.2 mL, 94.4 mmol, 1.3 eq.) in DCM (700 mL) under a nitrogen atmosphere is stirred at RT overnight. The reaction mixture is concentrated. The crude is taken in *i*PrOH (75 mL) heated under reflux and cool to RT. After two days at RT, the solid is filtered off and dried to afford intermediate **9**.

30 (¹H, CDCl₃) δ (ppm): 7.83 (1H, d), 7.35 (1H, dd), 7.29 (1H, d), 6.41 (1H, s), 4.51-4.42 (2H, m), 4.28 (2H, t), 4.06-4.01 (1H, m), 3.89-3.69 (5H, m), 3.52 (1H, m), 3.11 (2H, t)

Intermediate 10: trifluoro-methanesulfonic acid 2-((R)-1-[1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl ester

[00228] Intermediate **10** was prepared using general method **C** starting from intermediate **7**.


[00229] (¹H, CDCl₃) δ (ppm): 7.83 (1H, d), 7.34 (1H, dd), 7.29 (1H, d), 6.41 (1H, s), 4.51-4.41 (2H, m),
5 4.28 (2H, t), 4.05-4.00 (1H, m), 3.91-3.66 (5H, m), 3.52 (1H, t), 3.11 (2H, t)

Intermediate 11: trifluoro-methanesulfonic acid 2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl ester

[00230] Intermediate **11** was prepared using general method **C** starting from intermediate **8**.

10 (¹H, DMSO-*d*₆) δ (ppm): 8.21 (1H, d), 7.65 (1H, d), 7.54 (1H, dd), 6.75 (1H, s), 4.28-4.26 (2H, m), 4.04 (2H, t), 3.90-3.84 (1H, m), 3.81-3.75 (2H, m), 3.68-3.58 (2H, m), 3.53-3.47 (1H, m), 3.41-3.36 (1H, m), 3.10 (t, 2H)

MW (calcd): 462.4; MW (obsd): 463.3 (M+1)



Intermediate 13: 9-amino-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one hydrochloride

20 *Step 1: 9-(Benzhydrylidene-amino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 12)*

[00231] A solution of intermediate **9** (1 g, 2.16 mmol, 1 eq.), Pd(OAc)₂ (24 mg, 0.11 mmol 0.05 eq.), Cs₂CO₃ (2.11 g, 6.48 mmol, 3 eq.) BINAP (134 mg, 0.21 mmol, 0.1 eq.) and benzophenone imine (587 mg, 3.24 mmol, 1.5 eq.) in toluene (20 mL) was heated at 150°C in a microwave for 45 min. The solvent was evaporated to dryness and the crude mixture was taken in water and extracted with EtOAc. The organic layer
25 was dried over MgSO₄ and concentrated under vacuum. 9-(Benzhydrylidene-amino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one intermediate **12** was obtained after purification by flash chromatography on silica gel and immediately used in next step.

30 *Step 2: 9-amino-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one hydrochloride (Intermediate 13)*

[00232] To a solution of intermediate **12** in a minimum of DCM/ Et₂O was added HCl 2N in Et₂O (4 mL). The precipitate was filtered off and dried to afford intermediate **13**.

(¹H, DMSO-*d*₆) δ (ppm): 7.82 (1H, d), 7.45 (3H, br), 6.68 (1H, d), 6.67 (1H, s), 6.51 (1H, s), 4.32 (2H, d), 3.99 (2H, t), 3.90-3.84 (1H, m), 3.82-3.35 (6H, m), 2.88 (2H, t)

5 MW (calcd): 329.4; MW (obsd): 330.2 (M+1)


General method D:



[00233] The corresponding alcohol (2 eq.) is added to a solution of NaH (2 eq., 60% in mineral oil) in THF at 0 °C. The reaction is warmed to RT for 30 min then cooled again at 0 °C. Bromo methyltrifluoroborate (1 eq.) is added to the reaction in one portion and the mixture is stirred at RT from few hours to 3 days (monitoring by ¹⁹F-NMR). The reaction is quenched with a solution of KHF₂ (1.5 M, 3 eq.) and the mixture is evaporated to dryness. The residue is suspended in acetone, the inorganics are filtered off and the filtrate is evaporated to dryness. The residue is suspended in a minimum amount of acetone, Et₂O is added and the product is obtained by filtration.

Illustrative synthesis of general method D:


Intermediate 14: tetrahydro-2H-pyran-4-ol-methyl trifluoroborate



Intermediate **14**


[00234] Tetrahydro-pyran-4-ol (152 mg, 1.49 mmol, 2 eq.) was added to a solution of NaH (60 mg, 1.49 mmol, 2 eq., 60% in mineral oil) in THF (4 mL) at 0 °C. The reaction was warmed to RT for 30 min then cooled again at 0 °C. Bromo methyltrifluoroborate (150 mg, 0.75 mmol, 1 eq.) was added to the reaction in one portion and the mixture was stirred at RT for 1 day (monitoring by ¹⁹F-NMR). The reaction was quenched with a solution of KHF₂ (1.5 mL, 1.5 M, 3 eq.) and the mixture was evaporated to dryness. The residue was suspended in acetone, the inorganics were filtered off and the filtrate was evaporated to dryness. The residue was suspended in a minimum amount of acetone (1.5 mL), and Et₂O (6 mL) was added. Intermediate **14** was obtained by filtration.

(¹H, DMSO-*d*₆) δ ppm 3.78 (2 H, d), 3.31 - 3.21 (2 H, m), 3.18 - 3.08 (1 H, m), 2.50 - 2.45 (2 H, m), 1.86 - 1.74 (2 H, m), 1.34 - 1.19 (2 H, m)

Intermediate 15: potassium 3-oxy-oxetanemethyltrifluoroborate


[00235] Intermediate 15 was prepared via general method **D** with oxetan-3-ol (the trifluoroborate reagent was recovered with the inorganics rather than the filtrate).

5 (¹H, DMSO-*d*₆) δ ppm 4.56 (2 H, s), 4.32 (3 H, d), 2.40 (2 H, d)

Intermediate 16: potassium (3-methyl-3-methoxy-oxetane-methyltrifluoroborate


[00236] Intermediate 16 was prepared via general method **D** with (3-methyloxetan-3-yl) methanol (the trifluoroborate reagent was recovered with the inorganics rather than the filtrate).

10 (¹H, DMSO-*d*₆) δ ppm 4.34 (2 H, d), 4.14 (2 H, d), 3.26 (2 H, s), 2.59 - 2.52 (2 H, m), 1.19 (3 H, s)

Intermediate 17: potassium 2,2-dimethyl-propyloxy-methyltrifluoroborate


15 [00237] Intermediate 17 was prepared via general method **D** with 2,2-dimethyl-propan-1-ol.

(¹H, DMSO-*d*₆) δ ppm 2.88 (2 H, s), 2.51 - 2.45 (2 H, m), 0.83 (9 H, s)

Intermediate 18: potassium cyclopropylmethoxy-methyltrifluoroborate

20 [00238] Intermediate 18 was prepared via general method **D** with cyclopropyl-methanol.


(¹H, DMSO-*d*₆) δ ppm 3.00 (2 H, d), 2.46 (2 H, d), 1.00 - 0.82 (1 H, m), 0.46 - 0.31 (2 H, m), 0.13 - 0.00 (2 H, m)

Intermediate 19: cyclopentylmethoxy-methyltrifluoroborate**Intermediate 19**

[00239] Intermediate **19** was prepared via general method **D** with cyclopentyl-methanol.

(¹H, DMSO-*d*₆) δ ppm 3.04 (2 H, d), 2.46 (2 H, d), 2.08 – 1.94 (1 H, m), 1.47 (6 H, br. s.), 1.07 - 1.22 (2 H, m)

5

Intermediate 20: potassium 2-cyclopropyl-ethyl-trifluoroborate**Intermediate 20**


[00240] A 2-neck round bottom flask equipped with a reflux condenser and an addition funnel was charged with Mg (193 mg, 8.05 mmol, 3 eq.) and Et₂O (1 mL) under N₂. One drop of neat (2-bromo-ethyl)-cyclopropane was added followed by two drops of dibromoethane. Once the 1st bubbles appeared, (2-bromo-ethyl)-cyclopropane (400 mg, 2.68 mmol, 1 eq.) in Et₂O (5 mL) was added dropwise. Upon completion of the addition, the resulting suspension was stirred at RT for 1 h. In a separate flask, purged with N₂, a solution made of B(OMe)₃ (0.45 mL, 4.02 mmol, 1.5 eq.) in THF (6 mL) was cooled to -78 °C. To this solution, the 2-cyclopropyl-ethyl magnesium bromide suspension was added dropwise via a double ended needle. The mixture was allowed to stir for 1 h at -78 °C and then was warmed to RT for 1 h. After cooling the mixture to 0 °C, a saturated aqueous solution of KHF₂ (2.5 mL, 4.5 M, 4.1 eq.) was added dropwise and the reaction mixture was allowed to warm to RT. After 30 min, the solution was concentrated *in-vacuo*. The dried solids were triturated with hot acetone and filtered to remove inorganic salts. The resulting filtrate was concentrated and the solid residue was triturated with Et₂O. Potassium 2-cyclopropyl-ethyl-trifluoroborate was filtered and dried *in-vacuo*.

10

15

20

(¹H, DMSO-*d*₆) δ ppm 1.07 – 0.92 (2 H, m), 0.66 - 0.53 (1 H, m), 0.27 - 0.21 (2 H, m), 0.067 - -0.07 (2 H, m), -0.117 - -0.17 (2 H, m)


General method E:

Intermediate 9, 10, 11

[00241] A vial is charged with intermediate 9, 10 or 11 (1 eq.), the appropriate boronic acid, boronic ester or potassium trifluoroborate (4.4 eq.), Cs₂CO₃ (2.6 eq.), (DPPF)PdCl₂.DCM (0.05 eq.), in 1,4-dioxane/H₂O (10/1, v/v), and the mixture is degassed with N₂. The vial is sealed and heated at 80 °C. When the reaction is complete, the vial is cooled to RT and the reaction is either worked up or volatiles are evaporated under vacuum. The product is then obtained after purification by either flash chromatography on silica gel, preparative TLC or preparative HPLC-MS.

10 **Illustrative synthesis of general method E:**

Compound 2: 2-([1,4]dioxan-2-ylmethoxy)-9-pyridin-3-yl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one




Intermediate 11

Compound 2

[00242] A vial was charged with intermediate 11 (84 mg, 0.074 mmol, 1 eq.), pyridine-3-boronic acid (40 mg, 0.327 mmol, 4.4 eq.), Cs₂CO₃ (62 mg, 0.190 mmol, 2.6 eq.), (DPPF)PdCl₂.DCM (3.3 mg, 0.004 mmol, 0.05 eq.), in 1,4-dioxane (1 mL) and H₂O (0.1 mL), and the mixture was degassed with N₂. The vial was sealed and heated at 80 °C. After 1h, the vial was cooled to RT and volatiles were evaporated under vacuum. The residue was then purified by flash chromatography on silica gel, eluting with 7.5% MeOH/DCM to afford compound 2.

(¹H, CDCl₃) δ ppm 8.89 (1 H, s), 8.67 (1 H, d), 7.93 (1 H, d), 7.82 (1 H, d), 7.61 (1 H, d), 7.53 (1 H, s), 7.43 (1 H, dd), 6.43 (1 H, s), 4.51 - 4.37 (2 H, m), 4.26 (2 H, t), 3.99 (1 H, m), 3.92 - 3.60 (5 H, m), 3.49 (1 H, m), 3.10 (2 H, t)


General method F:

Intermediate 6, 7, 8

[00243] A solution of intermediate 6, 7, or 8 (1 eq.), the appropriate alkylating agent (1.5 eq.), K_2CO_3 (2 eq.), KI (1 eq.) in MEK is heated at 80 °C. When the reaction is complete, the volatiles are evaporated to dryness and the residue is purified by flash chromatography on silica gel to give the desired product.

Illustrative synthesis of general method F:

Compound 8: 2-([1,4]dioxan-2-ylmethoxy)-9-(oxazol-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one




Intermediate 8

Compound 8

[00244] A solution of intermediate 8 (40 mg, 0.12 mmol, 1 eq.), 2-chloromethyloxazole (21 mg, 0.18 mmol, 1.5 eq.), K_2CO_3 (33 mg, 0.24 mmol, 2 eq.), KI (20 mg, 0.12 mmol, 1 eq.) in MEK (2 mL) was heated at 80 °C for 16 h. The reaction was evaporated to dryness and the residue was purified by flash chromatography on silica gel, eluting with 4% MeOH/DCM to give compound 8.

1H , $CDCl_3$) δ ppm 7.72 (1 H, d), 7.68 - 7.60 (1 H, m), 7.18 (1 H, d), 7.02 (1 H, dd), 6.93 (1 H, d), 6.28 (1 H, s), 5.23 (2 H, s), 4.49 - 4.33 (2 H, m), 4.19 (2 H, t), 4.02 - 3.94 (1 H, m), 3.89 - 3.61 (5 H, m), 3.48 (1 H, dd), 2.98 (2 H, t)

MW (calcd): 411.4; MW (obsd): 412.4 (M+1)


General method G:

Intermediate 9, 10, 11

[00245] Intermediate 9, 10, 11 (1 eq.) is dissolved in DMF, the appropriate alkyne (3 eq.) is added followed by TEA (3.5 eq.), and the mixture is degassed. Pd(PPh₃)₃Cl₂ (0.05 eq.) is added with CuI (0.2 eq.) and the reaction mixture is heated at 80 °C. When the reaction is gone to completion, it is cooled to RT and either worked up or volatiles are evaporated to dryness. The product is then obtained after purification by either flash chromatography on silica gel, preparative TLC or preparative HPLC-MS.

Illustrative synthesis of general method G:

10 **Compound 16:** 2-([1,4]dioxan-2-ylmethoxy)-9-(1-methyl-1H-imidazol-2-ylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one



Intermediate 11

Compound 16


15 [00246] Intermediate 11 (1.4 g, 3.03 mmol, 1 eq.) was dissolved in DMF (20 mL), 5-ethynyl-1-methyl-1H-imidazole (0.92 mL, 9.09 mmol, 3 eq.) was added followed by TEA (1.48 mL, 10.61 mmol, 3.5 eq.). The mixture was degassed and Pd(PPh₃)₃Cl₂ (106 mg, 0.15 mmol, 0.05 eq.) was added with CuI (115 mg, 0.61 mmol, 0.2 eq.). The reaction mixture was heated at 80 °C for 16 h. The reaction was cooled to RT and quenched with brine, the mixture was then extracted with EtOAc. The organic layer was dried over MgSO₄ and the solvent was evaporated under vacuum. The crude product was then purified by flash chromatography on silica gel, eluting from 0 to 5% MeOH/DCM to give compound 16.

(^1H , CDCl_3) δ ppm 7.73 (1 H, d), 7.57 - 7.50 (2 H, m), 7.48 - 7.45 (1 H, m), 7.42 (1 H, d), 6.42 (1 H, s), 4.53 - 4.39 (2 H, m), 4.28 - 4.23 (2 H, m), 4.07 - 3.97 (1 H, m), 3.94 - 3.64 (8 H, m), 3.52 (1 H, dd), 3.06 (2 H, t)

MW (calcd): 418.4; MW (obsd): 419.4 (M+1)

5

General method H:




[00247] tBuOK (3 eq.) is added to a solution of the corresponding acetylenic alcohol (1 eq.) in THF, MeI (10 eq.) is then added and the reaction is stirred at RT. When the reaction has gone to completion, the reaction mixture is filtered and the filtrate is evaporated to dryness. Product is obtained after purification by preparative TLC.

10

Illustrative synthesis of general method H:

Compound 110: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-methoxy-4-methyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one

15



Compound 91

Compound 110

[00248] tBuOK (5.19 mg, 0.046 mmol, 0.95 eq.) was added to a solution of compound 90 (20 mg, 0.049 mmol, 1 eq.) in THF (2 mL), MeI (0.030 mL, 0.487 mmol, 10 eq.) was then added and the reaction was stirred at RT for 16 h. Some more tBuOK (11 mg, 0.097 mmol, 2 eq.) was added and the reaction was stirred for an extra day. The reaction mixture was filtered and the filtrate was evaporated to dryness. The crude product was purified by preparative TLC eluting with 2% MeOH/DCM to yield compound 110.


20

(¹H, CDCl₃) δ ppm 7.70 - 7.60 (1 H, d), 7.50 - 7.42 (1 H, d), 7.39 (1 H, s), 6.37 (1 H, s), 4.50 - 4.35 (2 H, m), 4.25 - 4.15 (2 H, m), 4.05 - 3.95 (2 H, m), 3.92 - 3.60 (5 H, m), 3.56 - 3.45 (4 H, m), 3.05 - 3.95 (2 H, m), 2.15 - 1.95 (1 H, m), 1.15 - 1 (6 H, t).

MW (calcd): 424.5; MW (obsd): 425.2 (M+1)

5


General method I:



[00249] A vial is charged with Pd/C (10% w/w) and a solution of the appropriate alkyne (1 eq.) in MeOH is added. The system is purged with N₂ before being filled with H₂ then the reaction is stirred at RT until completion. The reaction is filtered through Celite™ and the filtrate is evaporated to dryness. Clean product is obtained after purification by either flash chromatography on silica gel, preparative TLC or preparative HPLC-MS.

Illustrative synthesis of general method I:

15 **Compound 116:** 2-([1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-3-pyridin-3-yl-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one



[00250] A vial was charged with Pd/C (9 mg, 10% w/w) and a solution of compound 108 (87 mg 0.20 mmol, 1 eq.) in MeOH (10 mL) was added. The system was purged with N₂ before being filled with H₂ then the reaction was stirred for 16 h at RT. The reaction was filtered through Celite and the filtrate was evaporated to dryness. The crude product was purified by preparative HPLC-MS to give compound 116.


20

(^1H , CDCl_3) δ ppm 8.60 - 8.48 (2 H, m), 7.73 (1 H, d), 7.61 (1 H, d), 7.30 (1 H, dd), 7.20 (1 H, d), 7.12 (1 H, s), 6.35 (1 H, s), 4.77 (1 H, dd), 4.47 - 4.33 (2 H, m), 4.18 (2 H, t), 3.97 (1 H, m), 3.90 - 3.60 (5 H, m), 3.48 (1 H, t), 2.96 (2 H, t), 2.90 - 2.70 (2 H, m), 2.15 (1 H, m), 2.10 - 1.98 (1 H, m), 1.38 (1 H, t)

MW (calcd): 449.5; MW (obsd): 450.1 (M+1)

5

General method J:



Intermediate 13


[00251] Intermediate 13 (1 eq.) is dissolved in DMF, the appropriate aldehyde (4 eq.) is added followed by KOH (1 eq.). The reaction is stirred for 15 min at RT before STAB (10 eq.) is added, the mixture is then stirred at RT until completion of the reaction. The mixture is then quenched with brine, extracted with EtOAc, the organic layer is dried over MgSO_4 and evaporated to dryness. Purification by preparative HPLC-MS affords the corresponding product.

10

Illustrative synthesis of general method J:

Compound 126: 9-(2,2-dimethyl-butylamino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one

15



Intermediate 13

Compound 126

[00252] Intermediate 13 (50 mg, 0.14 mmol, 1 eq.) was dissolved in DMF (2 mL), 2,2-dimethylbutanal (56 mg, 0.56 mmol, 4 eq.) was added followed by KOH (8 mg, 0.14 mmol, 1 eq.). The reaction was stirred for 15 min at RT before STAB (297 mg, 1.40 mmol, 10 eq.) was added, the mixture was then stirred for 2 days at RT. The reaction was quenched with brine and the mixture was extracted with EtOAc. The organic layer was


20

dried over MgSO_4 and evaporated to dryness. The crude product was purified by preparative HPLC-MS to give compound **126**.

(^1H , CDCl_3) δ ppm 7.50 (1 H, d), 6.63 (1 H, dd), 6.48 (1 H, s), 6.21 (1 H, s), 4.47 - 4.34 (2 H, m), 4.21 - 4.14 (2 H, m), 4.01 - 3.91 (1 H, m), 3.89 - 3.60 (5 H, m), 3.48 (1 H, dd), 2.98 (2 H, s), 2.90 (2 H, t), 1.36 (2 H, dd), 0.96 (6 H, s), 0.87 (3 H, t)

MW (calcd): 413.5; MW (obsd): 414.4 (M+1)

General method K:



Step 1: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate **15**)


[00253] Intermediate **15** 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one was synthesised via general method **E** with intermediate **9** and prop-2-yn-1-ol.

Step 2:


[00254] Intermediate **15** (1 eq.) is dissolved in a mixture of THF/DMF (1/1), NaH (1.1 eq., 60% in mineral oil) is added followed by the appropriate alkylating agent (1 eq.) and the reaction is stirred at 70 °C. When the reaction has gone to completion, the mixture is worked up with brine and EtOAc, the organic layer is dried over MgSO_4 and evaporated to dryness. Purification by preparative HPLC-MS provides the corresponding product.

Illustrative synthesis of general method K:

Compound 133: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-[3-(2-methoxy-ethoxy)-prop-1-ynyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one




- 5 **[00255]** Intermediate **15** (92 mg, 0.25 mmol, 1 eq.) was dissolved in a mixture of THF/DMF (6 mL, 1/1), NaH (11 mg, 0.275 mmol, 1.1 eq., 60% in mineral oil) was added followed by 1-bromo-2-methoxy-ethane (35 mg, 0.25 mmol, 1 eq.) and the reaction was stirred at RT for 16 h, then at 70 °C for 1 day. The mixture was worked up with brine and EtOAc, the organic layer was dried over MgSO₄ and evaporated to dryness. The crude product was purified by preparative HPLC-MS to provide compound **133**.
- 10 (¹H, CDCl₃) δ ppm 7.64 (1 H, s), 7.48 - 7.40 (1 H, m), 7.40 - 7.35 (1 H, m), 6.40 - 6.34 (1 H, m), 4.47 (4 H, s), 4.28 - 4.15 (1 H, m), 3.93 - 3.56 (10 H, m), 3.43 (5 H, s), 3.06 - 2.95 (2 H, m)
MW (calcd): 426.5; MW (obsd): 427.4 (M+1)

General method L:

- 15 **[00256]** To a solution of tBuOK (2.2 eq.) in DMF is added dropwise a solution of intermediate **6** (1 eq.) in DMF at 0 °C and the mixture is stirred for 1 h. A solution of the appropriate alkylating agent (10 eq.) in DMF is added dropwise to the previous solution at 0 °C, then the reaction is stirred at 80 °C. When the reaction has gone to completion, the mixture is cooled to RT, quenched with water, and extracted with EtOAc. The
- 20 combined organic layers are dried over Na₂SO₄ and evaporated to dryness. The product is isolated by purification by preparative TLC.

Illustrative synthesis of general method L:

Compound 158: 9-(2,2-dimethyl-propoxy)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one



- 5 **[00257]** To a solution of tBuOK (19 mg, 0.166 mmol, 1.1 eq.) in DMF (2 mL) was added dropwise a solution of intermediate **6** (50 mg, 0.151 mmol, 1 eq.) in DMF (2 mL) at 0 °C and the mixture was stirred for 1 h. A solution of 1-iodo-2,2-dimethyl-propane (0.021 mL, 0.159 mmol, 1.05 eq.) in DMF (2 mL) was added dropwise to the previous solution at 0 °C, then the reaction was stirred at 80 °C for 1 day. Extra reagents were added, 1-iodo-2,2-dimethyl-propane (0.4 mL, 10 eq.) and tBuOK (19 mg, 0.166 mmol, 1.1 eq.) and the
- 10 reaction was stirred at 80 °C for one more day. The mixture was cooled to RT, quenched with water, and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄ and evaporated to dryness. Compound **158** was obtained by purification by preparative TLC [DCM/MeOH, 98/2]. (¹H, CDCl₃) δ ppm 7.70 - 7.56 (1 H, d), 6.95 - 6.85 (1 H, d), 6.80 (1 H, s), 6.28 (1 H, s), 4.50 - 4.35 (2 H, m), 4.25 - 4.10 (2 H, m), 4.05 - 3.92 (1 H, m), 3.10 - 3.57 (8 H, m), 3.55 - 3.40 (1 H, m), 3.05 - 2.90 (2 H, m),
- 15 1.05 (9H, s).
MW (calcd): 400.5; MW (obsd): 401.2 (M+1)

Compounds of the invention

[00258] The compounds according to the invention can be produced as described below.

- 20 **Compound 1:** 9-Allyloxy-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00259] Synthesis fully described above.

Compound 2: 2-([1,4]dioxan-2-ylmethoxy)-9-pyridin-3-yl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00260] Synthesis fully described above.

25

Compound 3: 2-([1,4]dioxan-2-ylmethoxy)-9-pyridin-4-yl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00261] This compound is prepared via general method **E** using intermediate **11** and pyridine-4-boronic acid.

Compound 4: 2-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile.

[00262] This compound is prepared via general method E using intermediate 11 and 2-cyanophenylboronic acid.

5

Compound 5: 3-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile.

[00263] This compound is prepared via general method E using intermediate 11 and 3-cyanophenylboronic acid.

10

Compound 6: 4-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile.

[00264] This compound is prepared via general method E using intermediate 11 and 4-cyanophenylboronic acid.

15

Compound 7: [2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yloxy]-acetonitrile.

[00265] This compound is prepared via general method F using intermediate 8 and bromo-acetonitrile, KI was not used in the experiment.

20

Compound 8: 2-([1,4]dioxan-2-ylmethoxy)-9-(oxazol-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00266] Synthesis fully described above.

25

Compound 9: 2-([1,4]dioxan-2-ylmethoxy)-9-(pyridin-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00267] This compound is prepared via general method F using intermediate 8 and pyridin-2-yl-methanol hydrochloride.

30

Compound 10: 9-(3,5-dichloro-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00268] This compound is prepared via general method E using intermediate 11 and 3,5-dichlorophenylboronic acid.

Compound 11: 9-benzofuran-2-yl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00269] This compound is prepared via general method E using intermediate 11 and 1-benzofuran-2-ylboronic acid.


5

Compound 12: 2-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-indole-1-carboxylic acid tert-butyl ester.

[00270] This compound is prepared via general method E using intermediate 11 and 1-(tert-butoxycarbonyl)-1H-indol-2-ylboronic acid.

10

Compound 13: 2-([1,4]dioxan-2-ylmethoxy)-9-(1H-indol-2-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



[00271] Compound 12 (57 mg, 0.11 mmol) was dissolved in a mixture of DCM/TFA (1/1, 2 mL) and the reaction was stirred at RT for 6h. The mixture was evaporated to dryness to recover compound 13 as a TFA salt.

(¹H DMSO-*d*₆) δ ppm 8.11 (1 H, d), 7.95 - 7.86 (2 H, m), 7.57 (1 H, d), 7.43 (1 H, d), 7.15 (1 H, t), 7.10 (1 H, d), 6.71 (1 H, s), 4.31 - 4.25 (2 H, m), 4.09 (2 H, t), 3.92 - 3.84 (1 H, m), 3.79 (2 H, td), 3.71 - 3.57 (2 H, m), 3.55 - 3.46 (1 H, m), 3.40 (1 H, dd), 3.08 (2 H, t)

MW (calcd): 429.5 MW (obsd): 430.5 (M+1)

Compound 14: 2-([1,4]dioxan-2-ylmethoxy)-9-(6-methoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00272] This compound is prepared via general method E using intermediate 11 and 2-methoxy-5-pyridineboronic acid.

Compound 15: 2-([1,4]dioxan-2-ylmethoxy)-9-(6-trifluoromethyl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00273] This compound is prepared via general method E using intermediate 11 and 2-(trifluoromethyl)pyridine-5-boronic acid.

5

Compound 16: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-methyl-3H-imidazol-4-ylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00274] Synthesis fully described above.

10 **Compound 17:** 9-(5-tert-butyl-[1,2,4]oxadiazol-3-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00275] This compound is prepared via general method F using intermediate 8 and 5-(tert-butyl)-3-(chloromethyl)-1,2,4-oxadiazole.


15 **Compound 18:** 5-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-pyridine-2-carboxylic acid methylamide.

[00276] This compound is prepared via general method E using intermediate 11 and 2-(N-methylaminocarbonyl)pyridine-5-boronic acid pinacol ester.

20 **Compound 19:** 2-([1,4]dioxan-2-ylmethoxy)-9-pent-1-ynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00277] This compound is prepared via general method G using intermediate 11 and pent-1-yne.

Compound 20: 2-([1,4]dioxan-2-ylmethoxy)-9-(2-pyridin-2-yl-ethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



25

Step 1: 2-([1,4]dioxan-2-ylmethoxy)-9-((E)-2-pyridin-2-yl-vinyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (intermediate 22).

[00278] A round bottom flask was charged with intermediate 11 (50 mg, 0.11 mmol, 1 eq.), 2-vinylpyridine (0.014 mL, 0.13 mmol, 1.2 eq.), (DPPF)PdCl₂.DCM (4.4 mg, 0.0054 mmol, 0.05 eq.) and TEA (0.03

mL, 0.22 mmol, 2 eq.) under N₂ then the flask was degassed. DMF (2 mL) was then added and the reaction was stirred at 100 °C for 16 h. The reaction was cooled to RT and evaporated to dryness, the residue was then purified by flash chromatography on silica gel, eluting from 0 to 3% MeOH/DCM affording intermediate **22** 2-([1,4]dioxan-2-ylmethoxy)-9-((E)-2-pyridin-2-yl-vinyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

¹H, CDCl₃) δ ppm 8.66 - 8.59 (1 H, m), 7.74 - 7.53 (4 H, m), 7.46 (1 H, s), 7.39 (1 H, d), 7.30 - 7.15 (2 H, m), 6.37 (1 H, s), 4.48 - 4.34 (2 H, m), 4.21 (2 H, t), 4.02 - 3.93 (1 H, m), 3.89 - 3.59 (5 H, m), 3.53 - 3.42 (1 H, m), 3.03 (2 H, m)

MW (calcd): 417.5; MW (obsd): 418.4 (M+1)


Step 2: 2-([1,4]dioxan-2-ylmethoxy)-9-(2-pyridin-2-yl-ethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound **20**).

[00279] A round bottom flask was charged with intermediate **22** (45 mg, 0.11 mmol, 1 eq.), PtO₂ (6 mg, 0.025 mmol, 0.23 eq.) and THF (2 mL) was added. The system was purged with N₂ before being filled with H₂ then the reaction was stirred for 16 h at RT. The reaction mixture was filtered through a SPE guanidine cartridge and the solvent was evaporated to dryness. The crude product was purified by flash chromatography on silica gel, eluting from 1 to 10% MeOH/DCM to give compound **20**.

¹H, CDCl₃) δ ppm 8.58 (1 H, dd), 7.66 - 7.56 (2 H, m), 7.22 (1 H, dd), 7.19 - 7.09 (3 H, m), 6.36 (1 H, s), 4.50 - 4.36 (2 H, m), 4.25 - 4.17 (2 H, m), 4.04 - 3.95 (1 H, m), 3.91 - 3.62 (5 H, m), 3.50 (1 H, dd), 3.14 (4 H, s), 2.98 (2 H, t)

MW (calcd): 419.5; MW (obsd): 420.5 (M+1)

Compound 21: 2-([1,4]dioxan-2-ylmethoxy)-9-(2-pyrazin-2-yl-ethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



Step 1: 2-([1,4]dioxan-2-ylmethoxy)-9-((E)-2-pyrazin-2-yl-vinyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (intermediate **23**).

[00280] A round bottom flask was charged with intermediate **11** (50 mg, 0.11 mmol, 1 eq.), 2-vinylpyridine (0.014 mL, 0.13 mmol, 1.2 eq.), (DPPF)PdCl₂.DCM (4.4 mg, 0.0054 mmol, 0.05 eq.) and TEA (0.03 mL, 0.22 mmol, 2 eq.) under N₂ then the flask was degassed. DMF (2 mL) was then added and the reaction

was stirred at 100 °C for 16 h. The reaction was cooled to RT and evaporated to dryness, the residue was then purified by flash chromatography on silica gel, eluting from 0 to 4% MeOH/DCM affording intermediate **23** 2-([1,4]dioxan-2-ylmethoxy)-9-((E)-2-pyrazin-2-yl-vinyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

(¹H, CDCl₃) δ ppm 8.69 (1 H, m), 8.63 - 8.58 (1 H, m), 8.49 (1 H, m), 7.84 - 7.71 (2 H, m), 7.63 (1 H, d),
 5 7.53 (1 H, s), 7.31 (1 H, s), 6.42 (1 H, s), 4.52 - 4.39 (2 H, m), 4.30 - 4.22 (2 H, m), 4.02 (1 H, dd), 3.93 -
 3.63 (5 H, m), 3.52 (1 H, dd), 3.08 (2 H, m)

MW (calcd): 418.5; MW (obsd): 419.4 (M+1)

Step 2: 2-([1,4]dioxan-2-ylmethoxy)-9-(2-pyrazin-2-yl-ethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one
 10 (*Compound 21*).

[00281] A round bottom flask was charged with intermediate **23** (38 mg, 0.09 mmol, 1 eq.), PtO₂ (5 mg, 0.021 mmol, 0.23 eq.) and THF (2 mL) was added. The system was purged with N₂ before being filled with H₂ then the reaction was stirred for 16 h at RT. The reaction mixture was filtered through a SPE guanidine cartridge and the solvent was evaporated to dryness. The crude product was purified by flash chromatography
 15 on silica gel, eluting from 1 to 10% MeOH/DCM to give compound **21**.

(¹H, CDCl₃) δ ppm 8.55 (1 H, dd), 8.44 (2 H, dd), 7.63 (1 H, d), 7.22 (1 H, dd), 7.15 (1 H, s), 6.36 (1 H, s),
 4.36 - 4.53 (2 H, m), 4.17 - 4.26 (2 H, m), 3.95 - 4.07 (1 H, m), 3.61 - 3.93 (5 H, m), 3.51 (1 H, dd), 3.17 (4
 H, s), 2.99 (2 H, t)

MW (calcd): 420.5; MW (obsd): 421.5 (M+1)

Compound 22: 2-([1,4]dioxan-2-ylmethoxy)-9-(1H-indol-5-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00282] This compound is prepared via general method E intermediate **11** and using 5-indolylboronic acid.

Compound 23: 2-([1,4]dioxan-2-ylmethoxy)-9-(2-methoxy-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00283] This compound is prepared via general method E using intermediate **11** and 2-methoxyphenylboronic acid.

Compound 24: 2-([1,4]dioxan-2-ylmethoxy)-9-(5-methoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00284] This compound is prepared via general method E using intermediate **11** and 3-methoxy-5-pyridineboronic acid pinacol ester.

Compound 25: 2-([1,4]dioxan-2-ylmethoxy)-9-(1H-indazol-5-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00285] This compound is prepared via general method E using intermediate 11 and 1H-indazole-5-boronic acid.

5

Compound 26: 2-([1,4]dioxan-2-ylmethoxy)-9-(4-methoxy-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00286] This compound is prepared via general method E using intermediate 11 and 4-methoxyphenylboronic acid.

10

Compound 27: 3-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzamide.

[00287] This compound is prepared via general method E using intermediate 11 and 3-aminocarbonylphenylboronic acid.

15

Compound 28: 5-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-2-fluoro-benzamide.

[00288] This compound is prepared via general method E using intermediate 11 and 3-(aminocarbonyl)-4-fluorobenzeneboronic acid.

20

Compound 29: N-{3-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-phenyl}-acetamide.

[00289] This compound is prepared via general method E intermediate 11 and using 3-acetamidophenylboronic acid.

25

Compound 30: 9-cyclopropylethynyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00290] This compound is prepared via general method G intermediate 11 and using ethynyl-cyclopropane.

30

Compound 31: 2-([1,4]dioxan-2-ylmethoxy)-9-(1-hydroxy-cyclopentylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00291] This compound is prepared via general method G intermediate 11 and using 1-ethynyl-cyclopentanol.

35

Compound 32: 2-([1,4]dioxan-2-ylmethoxy)-9-pyrimidin-5-yl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00292] This compound is prepared via general method **E** intermediate **11** and using 5-pyrimidinylboronic acid.

5

Compound 33: 9-cyclohex-1-enyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00293] This compound is prepared via general method **E** using intermediate **11** and cyclohexene-1-boronic acid pinacol ester.

10

Compound 34: 2-([1,4]dioxan-2-ylmethoxy)-9-(1-methyl-1H-indol-5-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00294] This compound is prepared via general method **E** using intermediate **11** and 1-methylindole-5-boronic acid pinacol ester.

15

Compound 35: 2-([1,4]dioxan-2-ylmethoxy)-9-(6-methyl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00295] This compound is prepared via general method **E** using intermediate **11** and 6-methylpyridin-3-ylboronic acid.

20

Compound 36: 2-([1,4]dioxan-2-ylmethoxy)-9-pyridin-2-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00296] This compound is prepared via general method **G** using intermediate **11** and 2-ethynyl-pyridine.

25

Compound 37: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-methoxy-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00297] This compound is prepared via general method **G** using intermediate **11** and 3-methoxy-propyne.

30

Compound 38: 5-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-pent-4-ynenitrile.

[00298] This compound is prepared via general method **G** using intermediate **11** and pent-4-ynenitrile.

Compound 39: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

35

[00299] This compound is prepared via general method **G** using intermediate **11** and prop-2-yn-1-ol.

Compound 40: 2-([1,4]dioxan-2-ylmethoxy)-9-(4-methoxy-phenylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00300] This compound is prepared via general method **G** using intermediate **11** and 1-ethynyl-4-methoxy-benzene.

Compound 41: 2-([1,4]dioxan-2-ylmethoxy)-9-pyridin-3-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00301] This compound is prepared via general method **G** using intermediate **11** and 3-ethynyl-pyridine

Compound 42: 4-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-N-methyl-benzamide.

[00302] This compound is prepared via general method **E** using intermediate **11** and 4-(N-methylaminocarbonyl)phenylboronic acid.

Compound 43: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-methoxy-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00303] This compound is prepared via general method **E** using intermediate **11** and 3-methoxyphenylboronic acid.

Compound 44: 9-(2-chloro-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00304] This compound is prepared via general method **E** using intermediate **11** and 2-chlorophenylboronic acid.

Compound 45: 2-([1,4]dioxan-2-ylmethoxy)-9-(4-hydroxy-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00305] This compound is prepared via general method **G** using intermediate **11** and but-3-yn-1-ol.

Compound 46: 9-(1,5-dimethyl-1H-pyrazol-3-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00306] This compound is prepared via general method **F** using intermediate **8** and 3-chloromethyl-1,5-dimethyl-1H-pyrazole.

Compound 47: 2-([1,4]dioxan-2-ylmethoxy)-9-(1-methyl-1H-pyrazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00307] This compound is prepared via general method F using intermediate 8 and 3-chloromethyl-1-methyl-1H-pyrazole.

5

Compound 48: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-methyl-[1,2,4]oxadiazol-5-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00308] This compound is prepared via general method F using intermediate 8 and 5-chloromethyl-3-methyl-[1,2,4]oxadiazole.

10

Compound 49: 2-([1,4]dioxan-2-ylmethoxy)-9-(4-morpholin-4-yl-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00309] This compound is prepared via general method E using intermediate 11 and 4-morpholinophenylboronic acid, with CsF as base and DMF as solvent.

15

Compound 50: 3-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-4-fluoro-benzamide.

[00310] This compound is prepared via general method E using intermediate 11 and 5-carbamoyl-2-fluorobenzeneboronic acid, with CsF as base and DMF as solvent.

20

Compound 51: 3-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-5-fluoro-benzamide.

[00311] This compound is prepared via general method E using intermediate 11 and 3-(aminocarbonyl)-5-fluorobenzeneboronic acid, with CsF as base and DMF as solvent.

25

Compound 52: 9-(3,3-Dimethyl-but-1-ynyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00312] This compound is prepared via general method G using intermediate 11 and 3,3-dimethyl-but-1-yne.

30

Compound 53: 2-([1,4]dioxan-2-ylmethoxy)-9-pyridin-4-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00313] This compound is prepared via general method G using intermediate 11 and 4-ethynyl-pyridine.

Compound 54: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-methyl-isoxazol-5-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00314] This compound is prepared via general method F using intermediate 8 and 5-chloromethyl-3-methyl-isoxazole.

5

Compound 55: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-3-methyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00315] This compound is prepared via general method G using intermediate 11 and 2-methyl-but-3-yn-2-ol.

10

Compound 56: 2-([1,4]dioxan-2-ylmethoxy)-9-(2-methoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00316] This compound is prepared via general method E using intermediate 11 and 2-methoxy-3-pyridinyl boronic acid.

15

Compound 57: 2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00317] This compound is prepared via general method E using intermediate 11 and potassium (cyanomethyl) trifluoroborate.

20

Compound 58: 9-(3,6-dihydro-2H-pyran-4-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00318] This compound is prepared via general method E using intermediate 11 and 3,6-dihydro-2H-pyran-4-boronic acid pinacol ester.

25

Compound 59: 5-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-pyridine-2-carbonitrile.

[00319] This compound is prepared via general method E using intermediate 11 and 2-cyanopyridine-5-boronic acid pinacol ester.

30

Compound 60: 2-([1,4]Dioxan-2-ylmethoxy)-9-(6-isopropoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00320] This compound is prepared via general method E using intermediate 11 and 6-isopropoxypyridine-3-boronic acid pinacol ester.

Compound 61: 2-([1,4]dioxan-2-ylmethoxy)-9-(6-ethoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00321] This compound is prepared via general method E using intermediate 11 and 6-ethoxypyridine-3-boronic acid.

5

Compound 62: 2-([1,4]dioxan-2-ylmethoxy)-9-(6-morpholin-4-yl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00322] This compound is prepared via general method E using intermediate 11 and 4-[5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-pyridinyl]morpholine.

10

Compound 63: 9-(2,3-dimethoxy-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00323] This compound is prepared via general method E using intermediate 11 and 2,3-dimethoxyphenylboronic acid.

15

Compound 64: 9-(3-chloro-2-methoxy-pyridin-4-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00324] This compound is prepared via general method E using intermediate 11 and 3-chloro-2-methoxypyridine-4-boronic acid.

20

Compound 65: 2-([1,4]dioxan-2-ylmethoxy)-9-(2-methyl-pyridin-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00325] This compound is prepared via general method E using intermediate 11 and 2-methylpyridine-4-boronic acid pinacol ester.

25

Compound 66: 3-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-isonicotinonitrile.

[00326] This compound is prepared via general method E using intermediate 11 and 4-cyanopyridine-3-boronic acid pinacol ester.

30

Compound 67: 9-(2,5-dimethoxy-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00327] This compound is prepared via general method E using intermediate 11 and 2,5-dimethoxyphenylboronic acid.

35

Compound 68: 2-([1,4]dioxan-2-ylmethoxy)-9-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00328] This compound is prepared via general method E using intermediate 11 and 1-[5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-pyridinyl]piperidine.

5

Compound 69: 2-([1,4]dioxan-2-ylmethoxy)-9-(2-ethoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00329] This compound is prepared via general method E using intermediate 11 and 2-ethoxypyridine-3-boronic acid.

10

Compound 70: 9-(2,6-dimethoxy-pyridin-3-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00330] This compound is prepared via general method E using intermediate 11 and 2,6-dimethoxy-3-pyridineboronic acid.

15

Compound 71: 4-[2-([1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-nicotinonitrile.

[00331] This compound is prepared via general method E using intermediate 11 and 3-cyanopyridine-4-boronic acid, pinacol ester.

20

Compound 72: 9-tert-butoxymethyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00332] This compound is prepared via general method E using intermediate 11 and potassium tert-butoxymethyltrifluoroborate.

25

Compound 73: 2-([1,4]dioxan-2-ylmethoxy)-9-(2-pyrrolidin-1-yl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00333] This compound is prepared via general method E using intermediate 11 and 2-(pyrrolidin-1-yl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine.

30

Compound 74: 2-([1,4]dioxan-2-ylmethoxy)-9-(6-pyrrolidin-1-yl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00334] This compound is prepared via general method E using intermediate 11 and 2-(1-pyrrolidinyl)-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine.

35

Compound 75: 2-([1,4]dioxan-2-ylmethoxy)-9-(5-phenyl-oxazol-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00335] This compound is prepared via general method F using intermediate 8 and 2-chloromethyl-5-phenyl-oxazole.

5

Compound 76: 9-(5-tert-butyl-oxazol-2-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00336] This compound is prepared via general method F using intermediate 8 and 5-tert-butyl-2-chloromethyl-oxazole.

10

Compound 77: 9-(5-cyclopropyl-[1,2,4]oxadiazol-3-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00337] This compound is prepared via general method F using intermediate 8 and 3-chloromethyl-5-cyclopropyl-[1,2,4]oxadiazole.

15

Compound 78: 2-([1,4]dioxan-2-ylmethoxy)-9-(5-ethyl-[1,2,4]oxadiazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00338] This compound is prepared via general method F using intermediate 8 and 3-chloromethyl-5-ethyl-[1,2,4]oxadiazole.

20

Compound 79: 2-([1,4]dioxan-2-ylmethoxy)-9-(5-methyl-[1,2,4]oxadiazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00339] This compound is prepared via general method F using intermediate 8 and 3-chloromethyl-5-methyl-[1,2,4]oxadiazole.

25

Compound 80: 2-([1,4]dioxan-2-ylmethoxy)-9-(5-isopropyl-[1,2,4]oxadiazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00340] This compound is prepared via general method F using intermediate 8 and 3-chloromethyl-5-isopropyl-[1,2,4]oxadiazole.

30

Compound 81: 9-cyclopentylethynyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00341] This compound is prepared via general method G using intermediate 11 and ethynyl-cyclopentane.

35

Compound 82: 9-cyclohexylethynyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00342] This compound is prepared via general method **G** using intermediate **11** and ethynyl-cyclohexane.

5 **Compound 83:** 2-([1,4]dioxan-2-ylmethoxy)-9-(3-methyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00343] This compound is prepared via general method **G** using intermediate **11** and 3-methyl-but-1-yne.

Compound 84: 2-([1,4]dioxan-2-ylmethoxy)-9-hex-1-ynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

10 [00344] This compound is prepared via general method **G** using intermediate **11** and hex-1-yne.

Compound 85: 9-[3-(benzyl-methyl-amino)-prop-1-ynyl]-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

15 [00345] This compound is prepared via general method **G** using intermediate **11** and benzyl-methyl-prop-2-ynyl-amine.

Compound 86: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-5-methyl-hex-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

20 [00346] This compound is prepared via general method **G** using intermediate **11** and 5-methyl-hex-1-yn-3-ol, with iPr_2NH as base and THF as solvent.

Compound 87: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

25 [00347] This compound is prepared via general method **G** using intermediate **11** and but-3-yn-2-ol, with iPr_2NH as base and THF as solvent.

Compound 88: 9-cyclopropyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

30 [00348] This compound is prepared via general method **E** using intermediate **11** and potassium cyclopropyltrifluoroborate.

Compound 89: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

35 [00349] This compound is prepared via general method **G** using intermediate **11** and pent-1-yn-3-ol with iPr_2NH as base and THF as solvent.

Compound 90: 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-4-methyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00350] This compound is prepared via general method **G** using intermediate **11** and 4-methyl-pent-1-yn-3-ol with $i\text{Pr}_2\text{NH}$ as base and THF as solvent.

5

[00351] **Compound 91:** 2-([1,4]dioxan-2-ylmethoxy)-9-(3-ethyl-3-hydroxy-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00352] This compound is prepared via general method **G** using intermediate **11** and 3-ethyl-pent-1-yn-3-ol with $i\text{Pr}_2\text{NH}$ as base and THF as solvent.

10

Compound 92: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-3-phenyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00353] This compound is prepared via general method **G** using intermediate **11** and 2-phenyl-but-3-yn-2-ol with Cs_2CO_3 as base and MeCN as solvent under reflux.


15

Compound 93: 9-(3-benzylamino-prop-1-ynyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00354] This compound is prepared via general method **G** using intermediate **11** and benzyl-prop-2-ynyl-amine with Cs_2CO_3 as base and MeCN as solvent under reflux.

20

Compound 94: 2-([1,4]dioxan-2-ylmethoxy)-9-[(furan-2-ylmethyl)-amino]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

Intermediate **11**Compound **94**

[00355] A suspension of intermediate **11** (200 mg, 0.433 mmol, 1 eq.), furan-2-ylmethanamine (0.038 mL, 0.433 mmol, 1 eq.) and Cs_2CO_3 (0.17g, 0.519 mmol, 1.2 eq.) in toluene (4 mL) was degassed with Ar for 30 min before BINAP (+/-) (16 mg, 0.026 mmol, 0.06 eq.) and $\text{Pd}(\text{OAc})_2$ (3.88 mg, 0.017 mmol, 0.04 eq.) were added. The reaction was heated to 65 °C for 16 h. The reaction was cooled to RT, BINAP (+/-) (16 mg, 0.026 mmol, 0.06 eq.) and $\text{Pd}(\text{OAc})_2$ (3.88 mg, 0.017 mmol, 0.04 eq.) were added and the reaction was degassed.

25

The reaction was heated to 80 °C for 1 extra day. The reaction was cooled to RT, BINAP (+/-) (16 mg, 0.026 mmol, 0.06 eq.) and Pd(OAc)₂ (3.88 mg, 0.017 mmol, 0.04 eq.) were added and the reaction was degassed. The reaction was heated to 80 °C for 1 extra day. The reaction was cooled to RT and diluted with DCM and washed with 0.5N aqueous KHSO₄. The aqueous layer was extracted with DCM, the combined organic layers
 5 were dried over Na₂SO₄ and evaporated to dryness. The crude product was purified by preparative HPLC-MS [H₂O (98→2): MeCN (2→98) / 0.1% HCO₂H] to give compound **94**.

(¹H, CDCl₃) δ ppm 7.57 - 7.48 (1 H, d), 7.38 (1 H, s), 6.68 - 6.55 (1 H, d), 6.49 (1 H, s), 6.35 (1 H, s), 6.28 (1 H, s), 6.20 (1 H, s), 4.60 - 4.50 (1 H, m), 4.38 - 4.30 (4 H, m), 4.22 - 4.15 (2 H, m), 4.03 - 3.93 (1 H, m), 3.95 - 3.60 (5 H, m), 3.55 - 3.40 (1 H, t), 2.98 - 2.85 (2 H, m)

10 MW (calcd): 409.4; MW (obsd): 410.2 (M+1)

Compound 95: 2-([1,4]dioxan-2-ylmethoxy)-9-(1-ethyl-1H-pyrazol-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00356] This compound is prepared via general method **E** using intermediate **11** and 1-ethyl-1H-pyrazole-
 15 4-boronic acid, pinacol ester.

Compound 96: 2-([1,4]dioxan-2-ylmethoxy)-9-[1-(3-methyl-butyl)-1H-pyrazol-4-yl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00357] This compound is prepared via general method **E** using intermediate **11** and 1-(3-methylbutyl)-
 20 1H-pyrazole-4-boronic acid, pinacol ester.

Compound 97: 2-([1,4]dioxan-2-ylmethoxy)-9-(5-methyl-furan-2-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00358] This compound is prepared via general method **E** using intermediate **11** and 2-methylfurane-5-
 25 boronic acid pinacol ester.

Compound 98: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-hex-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00359] This compound is prepared via general method **G** using intermediate **11** and hex-1-yn-3-ol,
 30 Cs₂CO₃ as base and MeCN as solvent under reflux.

Compound 99: 9-(3,5-dimethyl-1H-pyrazol-4-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00360] This compound is prepared via general method **E** using intermediate **11** and 3,5-dimethylpyrazole-
 35 4-boronic acid pinacol ester.

Compound 100: 2-([1,4]dioxan-2-ylmethoxy)-9-(1H-pyrazol-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00361] This compound is prepared via general method E using intermediate 11 and pyrazole-4-boronic acid pinacol ester.

Compound 101: 2-([1,4]dioxan-2-ylmethoxy)-9-(1-propyl-1H-pyrazol-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00362] This compound is prepared via general method E using intermediate 11 and 1-propyl-1H-pyrazole-4-boronic acid, pinacol ester.

Compound 102: 2-[2-((R)-1-[1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile.

[00363] This compound is prepared via general method E using intermediate 10 and 2-cyanophenylboronic acid pinacol ester.


Compound 103: 2-[2-((S)-1-[1,4]dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile.

[00364] This compound is prepared via general method E using intermediate 9 and 2-cyanophenylboronic acid pinacol ester.

Compound 104: 9-(5-cyclopropyl-[1,2,4]oxadiazol-3-ylmethoxy)-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00365] This compound is prepared via general method F using intermediate 7 and 3-chloromethyl-5-cyclopropyl-[1,2,4]oxadiazole.

Compound 105: 2-([1,4]dioxan-2-ylmethoxy)-9-ethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



Step 1: 2-([1,4]dioxan-2-ylmethoxy)-9-prop-1-ynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate **24**).

[00366] Intermediate **24** 2-([1,4]dioxan-2-ylmethoxy)-9-prop-1-ynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one is synthesised via general method E with intermediate **11** and ethynyl-trimethyl-silane.

5 ^1H , CDCl_3) δ ppm 7.66 - 7.61 (1 H, m), 7.47 - 7.43 (1 H, m), 7.42 - 7.38 (1 H, m), 6.37 (1 H, s), 4.50 - 4.36 (2 H, m), 4.23 - 4.16 (2 H, m), 4.04 - 3.93 (1 H, m), 3.90 - 3.60 (4 H, m), 3.53 - 3.44 (1 H, m), 3.02 - 2.97 (1 H, m), 0.27 (7 H, s)

MW (calcd): 410.5; MW (obsd): 411.4 (M+1)


10 Step 2: 2-([1,4]dioxan-2-ylmethoxy)-9-ethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound **105**).

[00367] TBAF (4.06 mL, 4.06 mmol, 1.2 eq.) was added dropwise to a solution of intermediate **24** (1.39 g, 3.39 mmol, 1 eq.) in THF (40 mL) at 0 °C and the reaction was stirred for 1 h at 0 °C. The reaction was then evaporated to dryness, and the residue was re-dissolved in HCl 1N. The aqueous phase was extracted with DCM, the combined organic layers were dried over MgSO_4 and the solvent evaporated under vacuum. The crude product was purified by flash chromatography on silica gel, eluting from 0 to 5% MeOH/DCM to furnish compound **105**.

^1H , CDCl_3) δ ppm 7.65 (1 H, s), 7.51 - 7.46 (1 H, m), 7.45 - 7.41 (1 H, m), 6.37 (1 H, s), 4.49 - 4.36 (2 H, m), 4.21 (2 H, s), 4.04 - 3.94 (1 H, m), 3.90 - 3.61 (5 H, m), 3.53 - 3.44 (1 H, m), 3.25 (1 H, s), 3.00 (2 H, s)

20 MW (calcd): 330.3; MW (obsd): 331

Compound **106**: 2-([1,4]dioxan-2-ylmethoxy)-9-pyrimidin-2-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



25 [00368] Compound **105** (50 mg, 0.15 mmol, 1 eq.) was dissolved in DMF (3 mL), 5-bromo-pyrimidine (47 mg, 0.30 mmol, 2 eq.) was added followed by TEA (0.062 mL, 0.44 mmol, 3 eq.), and the mixture was degassed. $\text{Pd(PPh}_3)_3\text{Cl}_2$ (5 mg, 0.0074 mmol, 0.05 eq.) was added with CuI (6 mg, 0.029 mmol, 0.2 eq.) and

the reaction mixture was heated at 80 °C for 16 h. The volatiles were evaporated to dryness and the crude product was purified by flash chromatography on silica gel, eluting from 0 to 5% MeOH/DCM to give compound **106**.

5 **Compound 107:** 2-([1,4]dioxan-2-ylmethoxy)-9-(3-phenylamino-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00369] This compound is prepared via general method **G** using intermediate **11** and phenyl-prop-2-ynyl-amine with $i\text{Pr}_2\text{NH}$ as base and THF as solvent.

10 **Compound 108:** 2-([1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-3-pyridin-3-yl-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00370] This compound is prepared via general method **G** using intermediate **11** and 1-pyridin-3-yl-prop-2-yn-1-ol.

15 **Compound 109:** 9-cyclopentylloxymethyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00371] This compound is prepared via general method **E** using intermediate **11** and potassium cyclopentoxymethyltrifluoroborate.

20 **Compound 110:** 2-([1,4]dioxan-2-ylmethoxy)-9-(3-methoxy-4-methyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00372] Synthesis fully described above.

25 **Compound 111:** 9-cyclopropylethynyl-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00373] This compound is prepared via general method **G** using intermediate **10** and ethynyl-cyclopropane.


30 **Compound 112:** 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-methyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00374] This compound is prepared via general method **G** using intermediate **9** and 3-methyl-but-1-yne.

Compound 113: 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-imidazol-1-yl-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00375] This compound is prepared via general method G using intermediate 11 and 1-prop-2-ynyl-1H-imidazole.

Compound 114: 9-(2-cyclopropyl-ethyl)-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



[00376] A round bottom flask was charged with intermediate 10 (2 g, 4.33 mmol, 1 eq.), intermediate 20 (1.1 g, 6.5 mmol, 1.5 eq.), K₂CO₃ (1.8 g, 13 mmol, 3 eq.), Pd(OAc)₂ (19 mg, 0.087 mmol, 0.02 eq.), RuPhos (81 mg, 0.173 mmol, 0.04 eq.), toluene (30 mL) and H₂O (3 mL). The mixture was degassed with N₂ and was heated at 80 °C for 16 h. The reaction was cooled to RT, quenched with brine and extracted with EtOAc. The organic layer was dried over MgSO₄ and evaporated to dryness. The crude product was purified by flash chromatography on silica gel, eluting from 70 to 90% EtOAc/H to give compound 114.

¹H, CDCl₃) δ ppm 7.64 - 7.58 (1 H, m), 7.24 - 7.19 (1 H, m), 7.13 (1 H, s), 6.35 (1 H, s), 4.49 - 4.36 (2 H, m), 4.25 - 4.17 (2 H, m), 4.03 - 3.95 (1 H, m), 3.90 - 3.61 (5 H, m), 3.54 - 3.45 (1 H, m), 3.02 - 2.95 (2 H, m), 2.81 - 2.72 (2 H, m), 1.59 - 1.49 (2 H, m), 0.77 - 0.64 (1 H, m), 0.49 - 0.42 (2 H, m), 0.10 - 0.02 (2 H, m)
MW (calcd): 382.5; MW (obsd): 383.4 (M+1)

Compound 115: 9-cyclopentyloxymethyl-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00377] This compound is prepared via general method E using intermediate 10 and potassium cyclopentoxymethyltrifluoroborate.

Compound 116: 2-([1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-3-pyridin-3-yl-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00378] Synthesis fully described above.

Compound 117: 9-allyloxy-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00379] Synthesis fully described above.

Compound 118: 9-allyloxy-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00380] Synthesis fully described above.

5

Compound 119: 2-((R)-1-[1,4]dioxan-2-ylmethoxy)-9-(tetrahydro-pyran-4-yloxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00381] This compound is prepared via general method E using intermediate 10 and intermediate 14.

10 **Compound 120:** 2-([1,4]dioxan-2-ylmethoxy)-9-{3-[(pyridin-3-ylmethyl)-amino]-prop-1-ynyl}-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00382] This compound is prepared via general method G using intermediate 11 and prop-2-yn-1-yl(pyridin-3-ylmethyl)amine.

15 **Compound 121:** 2-((R)-1-[1,4]dioxan-2-ylmethoxy)-9-pentyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00383] This Compound is prepared via general method I using compound 111.

Compound 122: 9-cyclopropylethynyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

20 [00384] Intermediate 9 (13.5 g, 29.20 mmol, 1 eq.) was dissolved in degassed DMF (1000 mL) under an nitrogen atmosphere, ethynyl cyclopropane (3.5 g, 53.00 mmol, 1.8 eq.) was added followed by Pd(PPh₃)₃Cl₂ (1.11 g, 1.58 mmol, 0.05 eq.), CuI (1.9 g, 9.98 mmol, 0.34 eq.) and TEA (12.5 mL, 89.7 mmol, 3.1 eq.). The reaction mixture was stirred 3h at 80 °C and 15 h at room temperature. The reaction was concentrated under vacuum. The crude product was then purified by trituration in hot *i*PrOH to afford compound 122.


25 [00385] (¹H, CDCl₃) δ ppm 7.63 (1 H, d), 7.39 (1 H, dd), 7.31 (1 H, s), 6.36 (1 H, s), 4.50 - 4.39 (2 H, m), 4.20 (2 H, t), 4.02 - 3.98 (1 H, m), 3.89 - 3.66 (5 H, m), 3.49 (1 H, t), 2.96 (2 H, t), 1.59 - 1.48 (1 H, m), 0.98 - 0.81 (4 H, m)

MW (calcd): 378.4; MW (obsd): 379.4

ee = 98.3%

30

Compound 123: 9-(2-cyclopropyl-ethyl)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydropyrimido[6,1-a]isoquinolin-4-one.



[00386] A round bottom flask was charged with intermediate **9** (2.08 g, 4.51 mmol, 1 eq.), intermediate **20** (1.35 g, 7.67 mmol, 1.7 eq.), K₂CO₃ (1.87 g, 13.53 mmol, 3 eq.), Pd(OAc)₂ (20 mg, 0.09 mmol, 0.02 eq.), RuPhos (84 mg, 0.18 mmol, 0.04 eq.), toluene (30 mL) and H₂O (3 mL). The mixture was degassed with N₂ and was heated at 80 °C for 1.5 days. The reaction was cooled to RT and some more reagents were added, potassium 2-cyclopropyl-ethyl-trifluoroborate (0.3 eq.), Pd(OAc)₂ (0.02 eq.), RuPhos (0.04 eq.), the reaction was degassed and the reaction was heated at 80 °C for 16h. The reaction was cooled to RT, quenched with H₂O and extracted with EtOAc. The organic layer was dried over MgSO₄ and evaporated to dryness. The crude product was purified by flash chromatography on silica gel, eluting from 80 to 90% EtOAc/H to compound **123**.

(¹H, CDCl₃) δ ppm 7.64 - 7.58 (1 H, m), 7.24 - 7.19 (1 H, m), 7.13 (1 H, s), 6.35 (1 H, s), 4.49 - 4.36 (2 H, m), 4.24 - 4.17 (2 H, m), 4.03 - 3.95 (1 H, m), 3.91 - 3.62 (5 H, m), 3.49 (1 H, dd), 3.03 - 2.95 (2 H, m), 2.81 - 2.73 (2 H, m), 1.59 - 1.50 (2 H, m), 0.71 (1 H, s), 0.49 - 0.42 (2 H, m), 0.09 - 0.03 (2 H, m)
MW (calcd): 382.5; MW (obsd): 383.4

Compound 124: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(oxetan-3-yloxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00387] This compound is prepared via general method **E** using intermediate **9** and intermediate **15**.


Compound 125: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-methyl-oxetan-3-ylmethoxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00388] This compound is prepared via general method **E** using intermediate **9** and intermediate **16**.

Compound 126: 9-(2,2-dimethyl-butylamino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00389] Synthesis fully described above.

Compound 127: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-4-methyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate **25**)

[00390] Intermediate **25** is prepared via general method **G** using intermediate **9**, 4-Methyl-pent-1-yn-3-ol, *i*Pr₂NH as base and THF as solvent.

(¹H, CDCl₃) δ ppm 7.65 - 7.60 (1 H, d), 7.48 - 7.40 (1 H, m), 7.37 (1 H, s), 6.36 (1 H, s), 4.50 - 4.30 (3 H, m), 4.25 - 4.15 (2 H, m), 4.05 - 3.95 (1 H, m), 3.92 - 3.60 (6 H, m), 3.68 - 3.40 (1 H, m), 3.05 - 2.92 (2 H, m), 2.10 - 1.95 (1 H, m), 1.93 - 1.80 (1 H, m), 1.15 - 1.00 (6 H, m)

MW (calcd): 410.5; MW (obsd): 411.2 (M+1)

Step 2: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound **127**)

[00391] Compound **127** is prepared via general method **I** using intermediate **25**.

Compound 128: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(2-ethyl-hexylamino)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00392] This compound is prepared via general method **J** using intermediate **13** and 2-ethyl-hexanal.

Compound 129: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(2-methoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00393] This compound is prepared via general method **F** using intermediate **6** and 1-bromo-2-methoxy-ethane, KI was not used in this experiment.

Compound 130: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(2-ethoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00394] This compound is prepared via general method **F** using intermediate **6** and 1-bromo-2-ethoxyethane, KI was not used in this experiment.

Compound 131: 9-cyclopropylmethoxy-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00395] This compound is prepared via general method **F** using intermediate **6** and bromomethylcyclopropane, KI was not used in this experiment.

Compound 132: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(2-fluoro-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00396] This compound is prepared via general method **F** using intermediate **6** and 1-bromo-2-fluoroethane, KI was not used in this experiment.

Compound 133: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-[3-(2-methoxy-ethoxy)-prop-1-ynyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00397] Synthesis fully described above.

Compound 134: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-[3-(2-ethoxy-ethoxy)-prop-1-ynyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00398] This compound is prepared via general method **K** using intermediate **21** and 1-bromo-2-ethoxyethane.

Compound 135: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-[3-(2-fluoro-ethoxy)-prop-1-ynyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00399] This compound is prepared via general method **K** using intermediate **21** and 1-bromo-2-fluoroethane.

Compound 136: 9-(2,2-dimethyl-propoxymethyl)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00400] This compound is prepared via general method **E** using intermediate **9**, in a mixture of DME/H₂O (2/1), in a microwave at 120 °C for 20 min and intermediate **17**.

Compound 137: 9-cyclohexyloxymethyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00401] This compound is prepared via general method **E** using intermediate **9** and potassium cyclohexyloxymethyltrifluoroborate in a mixture of DME/H₂O (2/1), in a microwave at 120 °C for 20 min.


Compound 138: 9-cyclopropylmethoxymethyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00402] This compound is prepared via general method **E** using intermediate **9**, in a mixture of DME/H₂O (2/1), in a microwave at 120 °C for 20 min and intermediate **18**.

Compound 139: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(tetrahydro-pyran-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00403] This compound is prepared via general method **F** using intermediate **6** and 2-bromomethyl-tetrahydro-pyran, KI was not used in this experiment.

Compound 140: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.




Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 26).

[00404] Intermediate **26** was prepared via general method **G** using intermediate **9** and but-3-yn-2-ol, the crude product was used in the next step without characterization.

Step 2: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound 140).

[00405] Compound **140** was prepared via general method **I** using intermediate **26**.

Compound 141: 9-(4,4-dimethyl-pentyloxy)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.




[00406] Intermediate 6 (0.1 g, 0.303 mmol, 1 eq.), 4,4-dimethyl-pentan-1-ol (35 mg, 0.303 mmol, 1 eq.) and PPh_3 (95 mg, 0.363 mmol, 1.2 eq.) were suspended in 1,4-dioxane (5 mL) and the mixture was degassed with N_2 . DIAD (0.065 mL, 0.333 mmol, 1.1 eq.) was added dropwise and the reaction was stirred at RT for 2 h. The reaction mixture was evaporated to dryness and the residue was purified by preparative TLC eluting with EtOAc to give compound 141.

(^1H , CDCl_3) δ ppm 7.70 - 7.58 (1 H, d), 6.95 - 6.82 (1 H, d), 6.77 (1 H, s), 6.27 (1 H, s), 4.50 - 4.32 (2 H, m), 4.38 - 4.15 (2 H, m), 4.05 - 3.92 (3 H, m), 3.92 - 3.60 (5 H, m), 3.55 - 3.42 (1 H, t), 3.05 - 2.92 (2 H, m), 1.85 - 1.70 (2 H, m), 1.40 - 1.30 (2 H, m), 0.92 (9 H, s)

MW (calcd): 428.5; MW (obsd): 429.2 (M+1)

Compound 142: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-methoxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-4-methyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 27)


[00407] Intermediate 27 is prepared via general method H using intermediate 25.

MW (calcd): 424.5; MW (obsd): 425.2 (M+1)

Step 2: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-methoxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound **142**).

[00408] Compound **142** is prepared via general method **I** using intermediate **27**.

- 5 **Compound 143:** 9-(3-cyclopropyl-propoxy)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

Intermediate **6**Compound **143**

[00409] Intermediate **6** (2.12 g, 2.42 mmol, 1 eq.), 4,4-dimethyl-pentan-1-ol (0.77 g, 7.71 mmol, 1.2 eq.) and PPh₃ (2.02 g, 7.71 mmol, 1.2 eq.) were suspended in 1,4-dioxane (50 mL) and the mixture was degassed with N₂. DIAD (1.56 mL, 7.71 mmol, 1.1 eq.) was added dropwise and the reaction was stirred at RT for 2 h. The reaction mixture was quenched with brine and extracted with EtOAc, the organic phase was dried over MgSO₄ and evaporated to dryness. Compound **143** was obtained by purification by flash chromatography on silica gel, eluting from 60 to 100% EtOAc/H.


(¹H, CDCl₃) δ ppm 7.66 - 7.60 (1 H, m), 6.91 - 6.86 (1 H, m), 6.80 - 6.76 (1 H, m), 6.28 (1 H, s), 4.48 - 4.35 (2 H, m), 4.24 - 4.17 (2 H, m), 4.11 - 4.04 (2 H, m), 4.02 - 3.95 (1 H, m), 3.91 - 3.61 (5 H, m), 3.54 - 3.45 (1 H, m), 3.03 - 2.94 (2 H, m), 1.98 - 1.88 (2 H, m), 1.45 - 1.36 (2 H, m), 0.78 - 0.66 (1 H, m), 0.50 - 0.43 (2 H, m), 0.09 - 0.03 (2 H, m)

MW (calcd): 412.5; MW (obsd): 413.5

20 **Compound 145:** 9-cyclohexylamino-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00410] This compound is prepared via general method **J** using intermediate **13** and cyclohexanone.

- 25 **Compound 146:** 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-4,4-dimethyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-4,4-dimethyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 28).

[00411] Intermediate **28** is prepared via general method **G** using intermediate **9**.

5 MW (calcd): 424.5; MW (obsd): 425.4 (M+1)

Step 2: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-4,4-dimethyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound 146).

[00412] Compound **146** is prepared via general method **I** using intermediate **28**.


10

Compound 147: 9-cyclopentylmethoxymethyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00413] This compound is prepared via general method **E** using intermediate **9**, in a mixture of DME/H₂O (2/1), in a microwave at 120 °C for 20 min and intermediate **19**.

15

Compound 148: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-methoxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 29)

20


[00414] Intermediate **29** is prepared via general method **H** using intermediate **26**.

MW (calcd): 396.4; MW (obsd): 397.2 (M+1)

Step 1: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-methoxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound 148)

5 [00415] Compound 148 is prepared via general method I using intermediate 29.

Compound 149: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-phenylamino-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one



10

Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-phenylamino-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 30)

[00416] Intermediate 30 is prepared via general method G using intermediate 9 and phenyl-prop-2-ynyl-amine.


15 MW (calcd): 443.5; MW (obsd): 444.2 (M+1)

Step 2: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-phenylamino-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound 149)

[00417] Compound 149 is prepared via general method I using intermediate 30.

20

Compound 150: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(4-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 31)


[00418] Intermediate 31 was prepared via general method G using intermediate 9, pent-4-yn-2-ol, iPr_2NH as base and THF as solvent.

MW (calcd): 396.4; MW (obsd): 397.2 (M+1)

Step 2: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(4-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound 150)

[00419] Compound 150 was prepared via general method I using intermediate 31.

Compound 151: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(4-hydroxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 32)

[00420] Intermediate 32 was prepared via general method G using intermediate 9, but-3-yn-1-ol, iPr_2NH as base and THF as solvent.


(1H , $CDCl_3$) δ ppm 7.70 - 7.65 (1 H, m), 7.45 - 7.35 (1 H, m), 7.34 (1 H, s), 6.35 (1 H, s), 4.50 - 4.32 (2 H, m), 4.28 - 4.10 (2 H, m), 4.05 - 3.90 (1 H, m), 3.95 - 3.60 (7 H, m), 3.55 - 3.40 (1 H, m), 3.05 - 2.90 (2 H, m), 2.80 - 2.65 (2 H, m), 2.00 - 1.80 (1 H, m)

MW (calcd): 382.4; MW (obsd): 383.2 (M+1)

Step 2: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(4-hydroxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound 151).

[00421] Compound 151 was prepared via general method I using intermediate 32.

Compound 152: 9-(cyclohexyl-methyl-amino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



[00422] MeI (0.007 mg, 0.11 mmol, 1.2 eq.) was added to compound **147** (38 mg, 0.092 mmol, 1 eq.) and NaH (6 mg, 0.15 mmol, 1.6 eq.) in DMF (5 mL) and the reaction was stirred at RT for 16 h. Some more NaH (6 mg, 0.15 mmol, 1.6 eq.) and MeI (0.07 mg, 0.11 mmol, 1.2 eq.) were added to the reaction mixture and it was stirred for a further 2 days. The mixture was quenched with brine and extracted with EtOAc. The organic layers were dried over MgSO₄ and evaporated to dryness. The residue was purified by preparative HPLC-MS to provide compound **152**.

[00423] (¹H, CDCl₃) δ ppm 7.57 - 7.50 (1 H, m), 6.71 (1 H, d), 6.53 (1 H, br. s.), 6.21 (1 H, s), 4.51 - 4.33 (2 H, m), 4.26 - 4.14 (2 H, m), 4.06 - 3.93 (1 H, m), 3.92 - 3.56 (6 H, m), 3.56 - 3.41 (1 H, m), 3.00 - 2.90 (2 H, m), 2.88 (3 H, s), 2.04 - 1.63 (2 H, m), 1.60 - 1.31 (5 H, m), 1.28 - 1.08 (1 H, m)
MW (calcd): 425.5; MW (obsd): 426.4


Compound 153: 9-(cyclohexylmethyl-amino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00423] This compound is prepared via general method **J** using intermediate **13** and cyclohexanecarbaldehyde.

Compound 154: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-[(tetrahydro-pyran-4-ylmethyl)-amino]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00424] This compound is prepared via general method **J** using intermediate **13** and tetrahydro-pyran-4-carbaldehyde.

Compound 155: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-ethyl-3-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-ethyl-3-hydroxy-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 33)


- 5 **[00425]** Intermediate 33 is prepared via general method G using intermediate 9 and 3-ethyl-pent-1-yn-3-ol. (¹H, CDCl₃) δ ppm 7.51 - 7.46 (1 H, m), 7.28 - 7.24 (1 H, m), 7.22 - 7.19 (1 H, m), 6.20 (1 H, s), 4.30 - 4.19 (2 H, m), 4.08 - 4.00 (2 H, m), 3.87 - 3.79 (1 H, m), 3.75 - 3.47 (5 H, m), 3.39 - 3.30 (1 H, m), 3.03 (1 H, br. S), 2.88 - 2.80 (2 H, m), 1.746 - 1.56 (4 H, m), 0.97 (6 H, s)
MW (calcd): 424.5; MW (obsd): 425.5 (M+1)

10

Step 2: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-ethyl-3-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound 155).

[00426] Compound 155 is prepared via general method I using intermediate 33.

- 15 **Compound 156:** 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-3-methyl-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one



Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-methyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 34)

20

[00427] Intermediate **34** is prepared via general method **G** using intermediate **9** and 2-methyl-but-3-yn-2-ol.


(¹H, CDCl₃) δ ppm 7.58 (1 H, d), 7.38 - 7.33 (1 H, m), 7.32 - 7.29 (1 H, m), 6.32 (1 H, s), 4.43 - 4.32 (2 H, m), 4.18 - 4.12 (2 H, m), 3.99 - 3.91 (1 H, m), 3.87 - 3.58 (5 H, m), 3.50 - 3.42 (1 H, m), 2.97 - 2.90 (2 H, m), 1.60 (6 H, s)

MW (calcd): 396.4; MW (obsd): 397.3 (M+1)

Step 2: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-3-methyl-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound **156**)

[00428] Compound **156** is prepared via general method **I** using intermediate **34**.

Compound **157**: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate **35**)

[00429] Intermediate **35** is prepared via general method **G** using intermediate **9** and pent-1-yn-3-ol.

(¹H, CDCl₃) δ ppm 7.62 - 7.57 (1 H, m), 7.41 - 7.35 (1 H, m), 7.33 (1 H, s), 6.34 (1 H, s), 4.60 - 4.53 (1 H, m), 4.45 - 4.33 (2 H, m), 4.20 - 4.14 (2 H, m), 4.00 - 3.92 (1 H, m), 3.88 - 3.59 (5 H, m), 3.51 - 3.42 (1 H, m), 2.99 - 2.90 (2 H, m), 1.90 - 1.74 (2 H, m), 1.07 (3 H, t)

Step 2: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound **157**)

[00430] Compound **157** is prepared via general method **I** using intermediate **35**.


Compound **158**: 9-(2,2-dimethyl-propoxy)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00431] Synthesis fully described above.

Compound 159: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(tetrahydro-pyran-4-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00432] This compound is prepared via general method L with intermediate 6 and methanesulfonic acid tetrahydro-pyran-4-ylmethyl ester.

Compound 160: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(4-hydroxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-4-methyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (intermediate 36)

[00433] A vial was charged with intermediate 9 (0.15 g, 0.324 mmol, 1 eq.), 2-methyl-5-trimethylsilanyl-pent-4-yn-2-ol (66 mg, 0.389 mmol, 1.2 eq.), CuI (2.5 mg, 0.013 mmol, 0.04 eq.), $i\text{Pr}_2\text{NH}$ (0.41 mL, 2.92 mmol, 9 eq.) and THF (2 mL). The solution was purged with Ar for 15 min., and $\text{Pd}(\text{PPh}_3)\text{Cl}_2$ (11 mg, 0.016 mmol, 0.05 eq.) was added with TBAF (0.39 mL, 0.39 mmol, 1.2 eq., 1M in THF). The vial was sealed and the reaction was heated to 80 °C for 16 h. The reaction mixture was evaporated to dryness and the crude product was purified by preparative TLC [DCM/MeOH, 98/2] to afford intermediate 36 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(4-hydroxy-4-methyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

^1H , CDCl_3) δ ppm 7.70 - 7.58 (1 H, m), 7.45 - 7.38 (1 H, m), 7.35 (1 H, s), 6.37 (1H, s), 4.50 - 4.30 (2 H, m), 4.28 - 4.15 (2 H, m), 4.05 - 3.95 (1 H, m), 3.95 - 3.55 (5 H, m), 3.55 - 3.40 (1 H, m), 3.05 - 2.95 (2 H, m), 2.62 (2 H, s), 1.39 (6 H, s)

MW (calcd): 410.5; MW (obsd): 411.4


Step 2: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(4-hydroxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound 160)

[00434] Compound 160 is prepared via general method I using intermediate 36.

Compound 161: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(tetrahydro-pyran-4-ylmethoxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00435] This compound is prepared via general method E using intermediate 9 and potassium 4-(tetrahydropyranylmethoxy)methyltrifluoroborate.

Compound 162: 2-([1,4]dioxan-2-ylmethoxy)-9-methoxy-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



Step 1: 1-[2-(3-methoxy-phenyl)-ethyl]-pyrimidine-2,4,6-trione (intermediate 37)

[00436] Sodium (236 mg, 10.2 mmol, 2 eq.) was added to degassed EtOH (18 mL), when sodium dissolved completely, ethyl malonate (1.56 mL, 10.2 mmol, 2 eq.) was added and the reaction was refluxed for 1 h. Intermediate 1 (995 mg, 5.12 mmol, 1 eq.) in EtOH (4 mL) was then added and the reaction was refluxed for 1 day. The desired intermediate 37 1-[2-(3-methoxy-phenyl)-ethyl]-pyrimidine-2,4,6-trione precipitated upon addition of 2N aqueous HCl, it was filtered and washed with H₂O and finally dried.

(¹H, DMSO-*d*₆) δ ppm 7.27 - 7.18 (1 H, m), 6.84 - 6.72 (3 H, m), 3.92 - 3.83 (2 H, m), 3.75 (3 H, s), 3.62 (2 H, s), 2.80 - 2.70 (2 H, m)

MW (calcd): 262.3; MW (obsd): 263.3 (M+1)

Step 2: 2-chloro-9-methoxy-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (intermediate 38)

[00437] Intermediate 37 (920 mg, 3.51 mmol, 1 eq.) was heated in POCl₃ (5 mL) at 50°C for 2 days. The volatiles were evaporated under vacuum, the residue was dissolved in DCM and washed with a saturated aqueous solution of NaHCO₃, before drying over MgSO₄. Evaporation of the organic phase gave intermediate 38 2-chloro-9-methoxy-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one, which was used in the next step without further purification.

(¹H, CDCl₃) δ ppm 7.73 - 7.66 (1 H, m), 6.98 - 6.90 (1 H, m), 6.85 - 6.80 (1 H, m), 6.69 (1 H, s), 4.00 - 4.20 (2 H, m), 3.91 (3 H, s), 3.04 (2 H, m)

Step 3: 2-([1,4]dioxan-2-ylmethoxy)-9-methoxy-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (compound 162)

[00438] [1,4]Dioxan-2-yl-methanol (42 mg, 0.36 mmol, 2 eq.) was dissolved in DCM (3 mL) with NaH (14 mg, 0.36 mmol, 2 eq., 60% in mineral oil). After 30 min, intermediate 38 (50 mg, 0.18 mmol, 1 eq.) was added to the mixture and the reaction was stirred at RT for 16h. The reaction mixture was evaporated to dryness and the crude product was purified by preparative HPLC-MS to provide compound 162.

(¹H, CDCl₃) δ ppm 7.68 - 7.62 (1 H, m), 6.95 - 6.87 (1 H, m), 6.82 - 6.77 (1 H, m), 6.29 (1 H, s), 4.51 - 4.35 (2 H, m), 4.26 - 4.17 (2 H, m), 4.05 - 3.92 (1 H, m), 3.91 - 3.60 (8 H, m), 3.55 - 3.44 (1 H, m), 3.04 - 2.94 (2 H, m)

MW (calcd): 344.4; MW (obsd): 345.0

Compound 163: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(oxetan-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00439] This compound is prepared via general method L using intermediate 6 and methanesulfonic acid oxetan-3-ylmethyl ester.


Compound 164: 9-(3-cyclopropyl-propoxy)-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydropyrimido[6,1-a]isoquinolin-4-one.

[00440] A solution of intermediate 6 (1.15 g, 3.48 mmol, 1 eq.) and 3-cyclopropan-1-ol (0.349 g, 3.48 mmol, 1eq.) in 1,4-dioxane was degassed with Argon for 10 min. PPh₃ (1.096 g, 4.18 mmol, 1.2 eq.) was added and the reaction mixture was degassed with Argon an additional 5 min. DIAD (0.745 mL, 3.83 mmol, 1.1 eq.) was added dropwise at 0°C. The reaction mixture was stirred at RT for 16 h. 3-Cyclopropylpropan-1-ol (0.150 mg, 1.49 mmol, 0.43 eq.) and PPh₃ (0.30 g, 1.14 mmol, 0.33 eq.) were added. The reaction mixture was cooled to 0°C and DIAD (0.350 mL, 1.80 mmol, 5.2 eq.) was added. After 1 h at RT, the reaction mixture was concentrated under vacuum and the crude product was purified by flash chromatography on silica-gel to afford compound 164.

(¹H, DMSO-*d*₆) δ ppm 7.93 (1H, d), 6.97-6.92 (2 H, m), 6.53 (1H, s), 4.24-4.23 (2H, m), 4.08 (2H, t), 4.00 (2H, t), 3.98-3.74 (3H, m), 3.68-3.57 (2H, m), 3.51-3.48 (1H, m), 3.37 (1H, t), 2.96 (2H, t), 1.84-1.80 (2H, m), 1.36-1.30 (2H, m), 0.81-0.63 (1H, m), 0.42-0.39 (2H, m), 0.04-0.02 (2H, m)

MW (calcd): 412.5; MW (obsd): 413.0 (M+1)

Compound 165: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-methoxy-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.




Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 39)

[00441] Intermediate 39 was prepared via general method G using intermediate 9 and 3-methoxy-propyne. (¹H, CDCl₃) δ ppm 7.68 (1 H, d), 7.48 (1 H, d), 7.42 (1 H, s), 6.40 (1 H, s), 4.50 - 4.41 (2 H, m), 4.38 (2 H, s), 4.23 (2 H, t), 4.05 - 3.98 (1 H, m), 3.89 - 3.70 (5 H, m), 3.55 - 3.50 (4 H, m), 3.03 (2 H, t)
MW (calcd): 382.4; MW (obsd): 383.4 (M+1)

Step 2: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-methoxy-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound 165)

[00442] Compound 165 was prepared via general method I using intermediate 39.

Compound 166: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-[2-(1-hydroxy-cyclopentyl)-ethyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.



Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(1-hydroxy-cyclopentylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 40)

[00443] Intermediate 40 is prepared via general method G using intermediate 9 and 3-methoxy-propyne.

(^1H , CDCl_3) δ ppm 7.54 (1 H, d), 7.31 (1 H, d), 7.26 (1 H, s), 6.28 (1 H, s), 4.39 - 4.30 (2 H, m), 4.12 (2 H, t), 3.95 - 3.91 (1 H, m), 3.82 - 3.58 (5 H, m), 3.50 (1 H, m), 3.16 (1 H, s), 2.89 (2 H, t), 2.05 - 1.98 (4 H, m), 1.90 - 1.70 (4H, m)


5 *Step 2: 2-((S)-1-[1,4]dioxan-2-ylmethoxy)-9-[2-(1-hydroxy-cyclopentyl)-ethyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound 166)*

[00444] Compound 166 was prepared via general method I using intermediate 40.

10 **Compound 167:** 2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-tetrahydro-pyran-4-ylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

[00445] This compound is prepared via general method G using intermediate 10 and 4-Ethynyl-tetrahydro-pyran-4-ol.

15 **Compound 168:** 2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one



Step 1: 2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 40)


[00446] Intermediate 41 is prepared via general method G using intermediate 10 and 3-methoxy-propyne.

20 (^1H , CDCl_3) δ ppm 7.56 (1 H, d), 7.48 (1 H, d), 7.33 (1 H, s), 6.27 (1 H, s), 4.32 - 4.27 (2 H, m), 4.23 (2 H, s), 4.08 (2 H, t), 3.88 - 3.85 (1 H, m), 3.76 - 3.49 (5 H, m), 3.40 - 3.34 (4 H, m), 2.90 (2 H, t)

Step 2: 2-((R)-1-[1,4]dioxan-2-ylmethoxy)-9-(3-methoxy-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound 165)

25 [00447] Compound 168 is prepared via general method I using intermediate 41.

Compound 169: 2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-[2-(1-hydroxy-cyclopentyl)-ethyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one



Step 1: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(1-hydroxy-cyclopentylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Intermediate 42)


[00448] Intermediate **42** is prepared via general method **G** using intermediate **10** and 1-Ethynyl-cyclopentanol.

(¹H, CDCl₃) δ ppm 7.47 (1 H, d), 7.25 (1 H, d), 7.20 (1 H, s), 6.22 (1 H, s), 4.32 - 4.24 (2 H, m), 4.05 (2 H, t), 3.88 - 3.85 (1H, m), 3.77 - 3.52 (5 H, m), 3.38 (1H, t), 2.83 (2 H, t), 2.02 - 1.90 (4H, m), 1.85 - 1.67 (4H, m)

Step 2: 2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(1-hydroxy-cyclopentylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one (Compound 169)

[00449] Compound **169** is prepared via general method **I** using intermediate **42**.


Compound 170: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-propoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one

Intermediate **6**Compound **170**

[00450] Intermediate **6** (0.15 g, 0.45 mmol, 1 eq.), 2-Propoxy-ethanol (63 μL, 0.55 mmol, 1.2 eq.) and PPh₃ (144 mg, 0.55 mmol, 1.2 eq.) were suspended in 1,4-dioxane (5 mL) and the mixture was degassed with N₂. DIAD (0.108 mL, 0.55 mmol, 1.2 eq.) was added and the reaction was stirred at RT overnight. 0.5 eq. of DIAD and PPh₃ were added, and the reaction mixture was stirred at room temperature for an extra 2h.


Reaction mixture was diluted with brine, extracted with EtOAc, dried over MgSO₄ and concentrated. Crude product was purified on silicagel column to give compound **170**.

Compound 171: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-isopropoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one




[00451] Intermediate **6** (0.15 g, 0.45 mmol, 1 eq.), 2-isopropoxy-ethanol (63 μ L, 0.55 mmol, 1.2 eq.) and PPh₃ (144 mg, 0.55 mmol, 1.2 eq.) were suspended in 1,4-dioxane (5 mL) and the mixture was degassed with N₂. DIAD (0.108 mL, 0.55 mmol, 1.2 eq.) was added and the reaction was stirred at RT for 5 h. 0.5 eq. of DIAD and PPh₃ were added, and the reaction mixture was stirred at room temperature overnight. Reaction mixture was diluted with brine, extracted with EtOAc, dried over MgSO₄ and concentrated. The crude product was purified on silicagel column to give compound **171**.

Compound 172: 2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-propoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one



[00452] Intermediate **7** (0.25 g, 0.76 mmol, 1 eq.), 2-Propoxy-ethanol (105 μ L, 0.91 mmol, 1.2 eq.) and PPh₃ (238 mg, 0.91 mmol, 1.2 eq.) were suspended in 1,4-dioxane (10 mL) and the mixture was degassed with N₂. DIAD (0.180 mL, 0.91 mmol, 1.2 eq.) was added and the reaction was stirred at RT overnight. 0.3 eq. of DIAD and PPh₃ were added, and the reaction mixture was stirred at room temperature for an extra 24h. Reaction mixture was diluted with brine, extracted with EtOAc, dried over MgSO₄ and concentrated. Crude product was purified on silicagel column to give compound **172**.

Compound 173: 2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-isopropoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one



- 5 **[00453]** Intermediate **7** (0.25 g, 0.76 mmol, 1 eq.), 2-isopropoxy-ethanol (105 μ L, 0.91 mmol, 1.2 eq.) and PPh_3 (238 mg, 0.91 mmol, 1.2 eq.) were suspended in 1,4-dioxane (10 mL) and the mixture was degassed with N_2 . DIAD (0.180 mL, 0.91 mmol, 1.2 eq.) was added and the reaction was stirred at RT for 5 h. 0.3 eq. of DIAD and PPh_3 were added, and the reaction mixture was stirred at room temperature overnight. Reaction mixture was diluted with brine, extracted with EtOAc, dried over MgSO_4 and concentrated. Crude product
- 10 was purified on silicagel column to give compound **173**.

Compound 174: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-methoxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one

Step 1: potassium 3-methoxy-propyl-trifluoroborate



15

Intermediate **43**

- [00454]** In a 2-neck round bottom flask equipped with a reflux condenser and an addition funnel was charged with Mg (471 mg, 19.20 mmol, 3 eq.) and Et_2O (2 mL) under N_2 . One drop of neat (2-bromo-ethyl)-cyclopropane was added followed by two drops of dibromoethane. Once the 1st bubbles appeared, 1-Bromo-3-methoxy-propane (1 g, 6.54 mmol, 1 eq.) in Et_2O (10 mL) was added dropwise. Upon completion of the
- 20 addition, the resulting suspension was stirred at RT for 1 h. In a separate flask, purged with N_2 , a solution made of $\text{B}(\text{OMe})_3$ (1.1 mL, 9.81 mmol, 1.5 eq.) in THF (12 mL) was cooled to -78°C . To this solution, the 3-methoxy-propyl magnesium bromide suspension was added dropwise via a double ended needle. The mixture was allowed to stir for 1 h at -78°C and then was warmed to RT for 1 h. After cooling the mixture to 0°C , a saturated aqueous solution of KHF_2 (5.8 mL, 4.5 M, 4.1 eq.) was added dropwise and the reaction mixture
- 25 was allowed to warm to RT. After 30 min, the solution was concentrated *in-vacuo*. The dried solids were triturated with hot acetone and filtered to remove inorganic salts. The resulting filtrate was concentrated and

the solid residue was triturated with Et₂O. Potassium 3-methoxy-propyl-trifluoroborate, intermediate **43**, was filtered and dried *in-vacuo*.

(¹H, DMSO-*d*₆) δ ppm 3.19 – 3.13 (5 H, m), 1.38 – 1.29 (2 H, m), -0.1 - 0.19 (2 H, m)

5 *Step 2: 2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-methoxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one*




[00455] Compound **174** is prepared via general method **E** using intermediates **9** and **43**.

10 [00456] **Table II: Mass spectral data of the Compounds of the Invention**





MW: Molecular weight

calc: calculated






obs: observed

Cpd #	Structures	Name	MW (calc)	MW (obsd)
1		9-Allyloxy-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	370	371
2		2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-3-yl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	391	NA
3		2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-4-yl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	391	392






Cpd #	Structures	Name	MW (calc)	MW (obsd)
4		2-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile	415	416
5		3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile	415	416
6		4-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile	415	416
7		[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yloxy]-acetonitrile	369	370
8		2-([1,4]Dioxan-2-ylmethoxy)-9-(oxazol-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	411	412





Cpd #	Structures	Name	MW (calc)	MW (obsd)
9		2-([1,4]Dioxan-2-ylmethoxy)-9-(pyridin-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	421	422
10		9-(3,5-Dichloro-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	458	459
11		9-Benzofuran-2-yl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	430	NA
12		2-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-indole-1-carboxylic acid tert-butyl ester	529	530






Cpd #	Structures	Name	MW (calc)	MW (obsd)
13		2-([1,4]Dioxan-2-ylmethoxy)-9-(1H-indol-2-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	429	430
14		2-([1,4]Dioxan-2-ylmethoxy)-9-(6-methoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	421	422
15		2-([1,4]Dioxan-2-ylmethoxy)-9-(6-trifluoromethyl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	459	460
16		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-3H-imidazol-4-ylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	418	419
17		9-(5-tert-Butyl-[1,2,4]oxadiazol-3-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	468	469





Cpd #	Structures	Name	MW (calc)	MW (obsd)
18		5-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-pyridine-2-carboxylic acid methylamide	448	449
19		2-([1,4]Dioxan-2-ylmethoxy)-9-pent-1-ynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	380	381
20		2-([1,4]Dioxan-2-ylmethoxy)-9-(2-pyridin-2-yl-ethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	419	420
21		2-([1,4]Dioxan-2-ylmethoxy)-9-(2-pyrazin-2-yl-ethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	420	421
22		2-([1,4]Dioxan-2-ylmethoxy)-9-(1H-indol-5-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	429	430





Cpd #	Structures	Name	MW (calc)	MW (obsd)
23		2-([1,4]Dioxan-2-ylmethoxy)-9-(2-methoxy-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	420	421
24		2-([1,4]Dioxan-2-ylmethoxy)-9-(5-methoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	421	422
25		2-([1,4]Dioxan-2-ylmethoxy)-9-(1H-indazol-5-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	430	431
26		2-([1,4]Dioxan-2-ylmethoxy)-9-(4-methoxy-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	420	421





Cpd #	Structures	Name	MW (calc)	MW (obsd)
27		3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzamide	433	434
28		5-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-2-fluoro-benzamide	451	452
29		N-{3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-phenyl}-acetamide	447	448
30		9-Cyclopropylethynyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	378	379
31		2-([1,4]Dioxan-2-ylmethoxy)-9-(1-hydroxy-cyclopentylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	422	423






Cpd #	Structures	Name	MW (calc)	MW (obsd)
32		2-([1,4]Dioxan-2-ylmethoxy)-9-pyrimidin-5-yl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	392	393
33		9-Cyclohex-1-enyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	394	395
34		2-([1,4]Dioxan-2-ylmethoxy)-9-(1-methyl-1H-indol-5-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	443	444
35		2-([1,4]Dioxan-2-ylmethoxy)-9-(6-methyl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	405	406





Cpd #	Structures	Name	MW (calc)	MW (obsd)
36		2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-2-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	415	416
37		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	382	383
38		5-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-pent-4-ynenitrile	391	392
39		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	368	369
40		2-([1,4]Dioxan-2-ylmethoxy)-9-(4-methoxy-phenylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	444	445





Cpd #	Structures	Name	MW (calc)	MW (obsd)
41		2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-3-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	415	416
42		4-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-N-methyl-benzamide	447	NA
43		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	420	421
44		9-(2-Chloro-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	424	425

Cpd #	Structures	Name	MW (calc)	MW (obsd)
45		2-([1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	382	NA
46		9-(1,5-Dimethyl-1H-pyrazol-3-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	438	439
47		2-([1,4]Dioxan-2-ylmethoxy)-9-(1-methyl-1H-pyrazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	424	425
48		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-[1,2,4]oxadiazol-5-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	426	427





Cpd #	Structures	Name	MW (calc)	MW (obsd)
49		2-([1,4]Dioxan-2-ylmethoxy)-9-(4-morpholin-4-yl-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	475	476
50		3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-4-fluoro-benzamide	451	452
51		3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-5-fluoro-benzamide	451	452
52		9-(3,3-Dimethyl-but-1-ynyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	394	395

Cpd #	Structures	Name	MW (calc)	MW (obsd)
53		2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-4-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	415	416
54		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-isoxazol-5-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	425	426
55		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-methyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	396	397
56		2-([1,4]Dioxan-2-ylmethoxy)-9-(2-methoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	421	422
57		2-([1,4]Dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	314	NA





Cpd #	Structures	Name	MW (calc)	MW (obsd)
58		9-(3,6-Dihydro-2H-pyran-4-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	396	397
59		5-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-pyridine-2-carbonitrile	416	417
60		2-([1,4]Dioxan-2-ylmethoxy)-9-(6-isopropoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	449	450
61		2-([1,4]Dioxan-2-ylmethoxy)-9-(6-ethoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	435	436






Cpd #	Structures	Name	MW (calc)	MW (obsd)
62		2-([1,4]Dioxan-2-ylmethoxy)-9-(6-morpholin-4-yl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	476	477
63		9-(2,3-Dimethoxy-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	450	451
64		9-(3-Chloro-2-methoxy-pyridin-4-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	455	456
65		2-([1,4]Dioxan-2-ylmethoxy)-9-(2-methyl-pyridin-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	405	406







Cpd #	Structures	Name	MW (calc)	MW (obsd)
66		3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-isonicotinonitrile	416	417
67		9-(2,5-Dimethoxy-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	450	451
68		2-([1,4]Dioxan-2-ylmethoxy)-9-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	474	475
69		2-([1,4]Dioxan-2-ylmethoxy)-9-(2-ethoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	435	436

Cpd #	Structures	Name	MW (calc)	MW (obsd)
70		9-(2,6-Dimethoxy-pyridin-3-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	451	452
71		4-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-nicotinonitrile	416	417
72		9-tert-Butoxymethyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	400	401
73		2-([1,4]Dioxan-2-ylmethoxy)-9-(2-pyrrolidin-1-yl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	460	461





Cpd #	Structures	Name	MW (calc)	MW (obsd)
74		2-([1,4]Dioxan-2-ylmethoxy)-9-(6-pyrrolidin-1-yl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	460	461
75		2-([1,4]Dioxan-2-ylmethoxy)-9-(5-phenyl-oxazol-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	487	488
76		9-(5-tert-Butyl-oxazol-2-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	467	NA
77		9-(5-Cyclopropyl-[1,2,4]oxadiazol-3-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	452	453
78		2-([1,4]Dioxan-2-ylmethoxy)-9-(5-ethyl-[1,2,4]oxadiazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	440	441





Cpd #	Structures	Name	MW (calc)	MW (obsd)
79		2-([1,4]Dioxan-2-ylmethoxy)-9-(5-methyl-[1,2,4]oxadiazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	426	427
80		2-([1,4]Dioxan-2-ylmethoxy)-9-(5-isopropyl-[1,2,4]oxadiazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	454	455
81		9-Cyclopentylethynyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	406	407
82		9-Cyclohexylethynyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	420	421






Cpd #	Structures	Name	MW (calc)	MW (obsd)
83		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	380	381
84		2-([1,4]Dioxan-2-ylmethoxy)-9-hex-1-ynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	394	395
85		9-[3-(Benzyl-methyl-amino)-prop-1-ynyl]-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	471	472
86		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-5-methyl-hex-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	424	425
87		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	382	383





Cpd #	Structures	Name	MW (calc)	MW (obsd)
88		9-Cyclopropyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	354	355
89		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	396	397
90		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-4-methyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	410	411
91		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-ethyl-3-hydroxy-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	424	425
92		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-phenyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	458	459
93		9-(3-Benzylamino-prop-1-ynyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	457	458






Cpd #	Structures	Name	MW (calc)	MW (obsd)
94		2-([1,4]Dioxan-2-ylmethoxy)-9-[(furan-2-ylmethyl)-amino]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	409	410
95		2-([1,4]Dioxan-2-ylmethoxy)-9-(1-ethyl-1H-pyrazol-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	408	409
96		2-([1,4]Dioxan-2-ylmethoxy)-9-[1-(3-methyl-butyl)-1H-pyrazol-4-yl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	450	451
97		2-([1,4]Dioxan-2-ylmethoxy)-9-(5-methyl-furan-2-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	394	395

Cpd #	Structures	Name	MW (calc)	MW (obsd)
98		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-hex-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	410	411
99		9-(3,5-Dimethyl-1H-pyrazol-4-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	408	409
100		2-([1,4]Dioxan-2-ylmethoxy)-9-(1H-pyrazol-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	380	381
101		2-([1,4]Dioxan-2-ylmethoxy)-9-(1-propyl-1H-pyrazol-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	422	423






Cpd #	Structures	Name	MW (calc)	MW (obsd)
102		2-[2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile	415	416
103		2-[2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile	415	416
104		9-(5-Cyclopropyl-[1,2,4]oxadiazol-3-ylmethoxy)-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	452	453
105		2-([1,4]Dioxan-2-ylmethoxy)-9-ethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	338	331

Cpd #	Structures	Name	MW (calc)	MW (obsd)
106		2-([1,4]Dioxan-2-ylmethoxy)-9-pyrimidin-2-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	416	417
107		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-phenylamino-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	443	444
108		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-pyridin-3-yl-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	445	446
109		9-Cyclopentyloxymethyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	412	413
110		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-4-methyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	424	425






Cpd #	Structures	Name	MW (calc)	MW (obsd)
111		9-Cyclopropylethynyl-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	378	379
112		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	380	381
113		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-imidazol-1-yl-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	418	419
114		9-(2-Cyclopropyl-ethyl)-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	382	383






Cpd #	Structures	Name	MW (calc)	MW (obsd)
115		9-Cyclopentyloxymethyl-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	412	413
116		2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-pyridin-3-yl-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	449	450
117		9-Allyloxy-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	370	371
118		9-Allyloxy-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	370	371
119		2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(tetrahydropyran-4-yloxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	428	429






Cpd #	Structures	Name	MW (calc)	MW (obsd)
120		2-([1,4]Dioxan-2-ylmethoxy)-9-{3-[(pyridin-3-ylmethyl)-amino]-prop-1-ynyl}-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	458	459
121		2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-pentyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	384	385
122		9-Cyclopropylethynyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	378	379
123		9-(2-Cyclopropyl-ethyl)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	382	383
124		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(oxetan-3-yloxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	400	401





Cpd #	Structures	Name	MW (calc)	MW (obsd)
125		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-oxetan-3-ylmethoxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	428	429
126		9-(2,2-Dimethyl-butylamino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	413	414
127		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	414	415
128		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-ethyl-hexylamino)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	441	442
129		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-methoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	388	389





Cpd #	Structures	Name	MW (calc)	MW (obsd)
130		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-ethoxyethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	402	403
131		9-Cyclopropylmethoxy-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	384	385
132		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-fluoroethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	376	377
133		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[3-(2-methoxyethoxy)-prop-1-ynyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	426	427
134		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[3-(2-ethoxyethoxy)-prop-1-ynyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	440	441
135		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[3-(2-fluoroethoxy)-prop-1-ynyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	414	415

Cpd #	Structures	Name	MW (calc)	MW (obsd)
136		9-(2,2-Dimethylpropoxymethyl)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	414	415
137		9-Cyclohexyloxymethyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	426	427
138		9-Cyclopropylmethoxymethyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	398	399
139		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(tetrahydropyran-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	428	429
140		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxybutyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	386	387






Cpd #	Structures	Name	MW (calc)	MW (obsd)
141		9-(4,4-Dimethyl-pentyloxy)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	428	429
142		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	428	429
143		9-(3-Cyclopropyl-propoxy)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	412	413
145		9-Cyclohexylamino-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	411	413
146		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-4,4-dimethyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	428	429






Cpd #	Structures	Name	MW (calc)	MW (obsd)
147		9-Cyclopentylmethoxymethyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	426	427
148		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxybutyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	400	401
149		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-phenylaminopropyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	447	448
150		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxypentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	400	401
151		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxybutyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	386	387

Cpd #	Structures	Name	MW (calc)	MW (obsd)
152		9-(Cyclohexyl-methyl-amino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	425	426
153		9-(Cyclohexylmethyl-amino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	425	426
154		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[(tetrahydropyran-4-ylmethyl)-amino]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	427	428
155		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-ethyl-3-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	428	429

Cpd #	Structures	Name	MW (calc)	MW (obsd)
156		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-methyl-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	400	401
157		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	400	401
158		9-(2,2-Dimethyl-propoxy)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	400	401
159		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(tetrahydropyran-4-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	428	429

Cpd #	Structures	Name	MW (calc)	MW (obsd)
160		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	414	415
161		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(tetrahydropyran-4-ylmethoxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	442	443
162		2-([1,4]Dioxan-2-ylmethoxy)-9-methoxy-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	344	345
163		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(oxetan-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	400	401
164		9-(3-Cyclopropyl-propoxy)-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	412	413

Cpd #	Structures	Name	MW (calc)	MW (obsd)
165		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxypropyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	386	387
166		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[2-(1-hydroxycyclopentyl)-ethyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	426	427
167		2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxytetrahydro-pyran-4-ylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	438	
168		2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxypropyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	386	387
169		2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-[2-(1-hydroxycyclopentyl)-ethyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	426	427

Cpd #	Structures	Name	MW (calc)	MW (obsd)
170		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-propoxyethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	416	417
171		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-isopropoxyethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	416	417
172		2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-propoxyethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	416	417
173		2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-isopropoxyethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	416	417
174		2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-methoxybutyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one	400	401

[00457] Table III: NMR Data of the Compounds of the Invention

Cpd#	NMR data (δ)
1	(^1H , CDCl_3) δ ppm 7.61 (1 H, d), 6.89 (1 H, dd), 6.78 (1 H, d), 6.26 (1 H, s), 6.13 – 5.94 (1 H, m), 5.48 – 5.24 (2 H, m), 4.59 (2 H, dt), 4.48 – 4.30 (2 H, m), 4.18 (2 H, t), 4.10 – 4.02 (1 H, m), 3.96 (1 H, m), 3.89 – 3.57 (4 H, m), 3.46 (1 H, dd), 2.95 (2 H, t)
2	(^1H , CDCl_3) δ ppm 8.89 (1 H, s), 8.67 (1 H, d), 7.93 (1 H, d), 7.82 (1 H, d), 7.61 (1 H, d), 7.53 (1 H, s), 7.43 (1 H, dd), 6.43 (1 H, s), 4.51 – 4.37 (2 H, m), 4.26 (2 H, t), 3.99 (1 H, m), 3.92 – 3.60 (5 H, m), 3.49 (1 H, m), 3.10 (2 H, t)
3	(^1H , CDCl_3) δ ppm 8.77 (2 H, br. s.), 7.86 (1 H, d), 7.76 – 7.46 (4 H, m), 6.48 (1 H, br. s.), 4.60 – 4.36 (2 H, m), 4.30 (2 H, br. s.), 4.04 (1 H, br. s.), 3.94 – 3.67 (5 H, m), 3.54 (1 H, t), 3.15 (2 H, br. s.)
4	(^1H , CDCl_3) δ ppm 7.83 – 7.94 (2 H, m), 7.78 – 7.71 (1 H, m), 7.67 – 7.53 (4 H, m), 6.48 (1 H, s), 4.53 – 4.43 (2 H, m), 4.34 – 4.27 (2 H, m), 4.07 – 4.01 (1 H, m), 3.93 – 3.67 (5 H, m), 3.54 (1 H, dd), 3.14 (2 H, t)
5	(^1H , CDCl_3) δ ppm 7.90 (1 H, d), 7.87 – 7.79 (2 H, m), 7.73 – 7.69 (1 H, m), 7.65 – 7.56 (2 H, m), 7.52 (1 H, s), 6.45 (1 H, s), 4.51 – 4.37 (2 H, m), 4.27 (2 H, t), 4.06 – 3.95 (1 H, m), 3.91 – 3.61 (5 H, m), 3.51 (1 H, dd), 3.11 (2 H, t)
6	(^1H , CDCl_3) δ ppm 7.85 – 7.68 (5 H, m), 7.61 (1 H, dd), 7.54 (1 H, d), 6.44 (1 H, s), 4.51 – 4.36 (2 H, m), 4.26 (2 H, t), 4.00 (1 H, m), 3.91 – 3.61 (5 H, m), 3.50 (1 H, dd), 3.11 (2 H, t)
7	(^1H , CDCl_3) δ ppm 7.71 (1 H, d), 7.00 (1 H, dd), 6.90 (1 H, d), 6.31 (1 H, s), 4.86 (2 H, s), 4.51 – 4.34 (2 H, m), 4.22 (2 H, t), 3.99 (1 H, m), 3.92 – 3.58 (5 H, m), 3.49 (1 H, dd), 3.03 (2 H, t)
8	(^1H , CDCl_3) δ ppm 7.72 (1 H, d), 7.68 – 7.60 (1 H, m), 7.18 (1 H, d), 7.02 (1 H, dd), 6.93 (1 H, d), 6.28 (1 H, s), 5.23 (2 H, s), 4.49 – 4.33 (2 H, m), 4.19 (2 H, t), 4.02 – 3.94 (1 H, m), 3.89 – 3.61 (5 H, m), 3.48 (1 H, dd), 2.98 (2 H, t)
9	(^1H , CDCl_3) δ ppm 8.63 (1 H, d), 7.75 (1 H, td), 7.64 (1 H, d), 7.50 (1 H, d), 7.29 – 7.24 (1 H, m), 6.99 (1 H, dd), 6.89 (1 H, d), 6.27 (1 H, s), 5.29 (2 H, d), 4.48 – 4.34 (2 H, m), 4.19 (2 H, t), 4.02 – 3.93 (1 H, m), 3.90 – 3.60 (5 H, m), 3.48 (1 H, dd), 2.97 (2 H, t)
10	(^1H , CDCl_3) δ ppm 7.79 (1 H, d), 7.56 (1 H, d), 7.49 (3 H, d), 7.40 (1 H, s), 6.42 (1 H, s), 4.51 – 4.36 (2 H, m), 4.25 (2 H, t), 4.00 (1 H, m), 3.91 – 3.60 (5 H, m), 3.50 (1 H, t), 3.09 (2 H, t)

Cpd#	NMR data (δ)
11	(^1H , CDCl_3) δ ppm 7.87 (1 H, d), 7.83 - 7.76 (2 H, m), 7.64 (1 H, d), 7.56 (1 H, d), 7.41 - 7.32 (1 H, m), 7.31 - 7.23 (3 H, m), 7.18 (1 H, s), 6.43 (1 H, s), 4.46 (2 H, br. s.), 4.27 (2 H, t), 4.06 - 3.96 (1 H, m), 3.97 - 3.61 (5 H, m), 3.51 (1 H, t), 3.11 (2 H, t)
12	(^1H , CDCl_3) δ ppm 8.20 (1 H, d), 7.76 (1 H, d), 7.60 (1 H, d), 7.47 (1 H, dd), 7.42 - 7.35 (2 H, m), 7.336 - 7.26 (1 H, m), 6.68 (1 H, s), 6.44 (1 H, s), 4.54 - 4.40 (2 H, m), 4.27 (2 H, t), 4.08 - 3.96 (1 H, m), 3.91 - 3.65 (5 H, m), 3.52 (1 H, dd), 3.08 (2 H, t), 1.42 (9 H, s)
13	(^1H DMSO- d_6) δ ppm 8.11 (1 H, d), 7.95 - 7.86 (2 H, m), 7.57 (1 H, d), 7.43 (1 H, d), 7.15 (1 H, t), 7.10 (1 H, d), 6.71 (1 H, s), 4.31 - 4.25 (2 H, m), 4.09 (2 H, t), 3.92 - 3.84 (1 H, m), 3.79 (2 H, td), 3.71 - 3.57 (2 H, m), 3.55 - 3.46 (1 H, m), 3.40 (1 H, dd), 3.08 (2 H, t)
14	(^1H , CDCl_3) δ ppm 8.45 (1 H, d), 7.84 (1 H, dd), 7.79 (1 H, d), 7.56 (1 H, dd), 7.50 - 7.46 (1 H, m), 6.87 (1 H, d), 6.43 (1 H, s), 4.49 - 4.36 (2 H, m), 4.30 - 4.19 (2 H, m), 4.04 - 3.98 (4 H, m), 3.93 - 3.62 (5 H, m), 3.51 (1 H, dd), 3.10 (2 H, t)
15	(^1H , CDCl_3) δ ppm 8.97 (1 H, d), 8.09 (1 H, dd), 7.83 (2 H, dd), 7.63 (1 H, dd), 7.55 (1 H, s), 6.44 (1 H, s), 4.52 - 4.36 (2 H, m), 4.27 (2 H, t), 3.99 (1 H, tt), 3.91 - 3.59 (5 H, m), 3.49 (1 H, dd), 3.13 (2 H, t)
16	(^1H , CDCl_3) δ ppm 7.73 (1 H, d), 7.57 - 7.50 (2 H, m), 7.48 - 7.45 (1 H, m), 7.42 (1 H, d), 6.42 (1 H, s), 4.53 - 4.39 (2 H, m), 4.28 - 4.23 (2 H, m), 4.07 - 3.97 (1 H, m), 3.94 - 3.64 (8 H, m), 3.52 (1 H, dd), 3.06 (2 H, t)
17	(^1H , CDCl_3) δ ppm 7.66 (1 H, d), 7.03 (1 H, dd), 6.94 (1 H, d), 6.29 (1 H, s), 5.22 (2 H, s), 4.50 - 4.33 (2 H, m), 4.27 - 4.14 (2 H, m), 4.03 - 3.93 (1 H, m), 3.90 - 3.61 (5 H, m), 3.49 (1 H, dd), 2.99 (2 H, t), 1.47 (9 H, s)
18	(^1H , CDCl_3) δ ppm 8.80 (1 H, d), 8.32 (1 H, d), 8.16 - 7.96 (2 H, m), 7.85 (1 H, d), 7.65 (1 H, dd), 7.57 (1 H, s), 6.45 (1 H, s), 4.56 - 4.37 (2 H, m), 4.28 (2 H, t), 4.01 (1 H, qd), 3.93 - 3.60 (5 H, m), 3.58 - 3.44 (1 H, m), 3.20 - 3.02 (5 H, m)
19	(^1H , CDCl_3) δ ppm 7.65 (1 H, d), 7.41 (1 H, dd), 7.36 (1 H, s), 6.38 (1 H, s), 4.50 - 4.40 (2 H, m), 4.28 - 4.18 (2 H, m), 4.01 (1 H, m), 3.92 - 3.66 (5 H, m), 3.52 (1 H, dd), 3.01 (2 H, t), 2.45 (2 H, t), 1.68 (2 H, sxt), 1.09 (3 H, t)
20	(^1H , CDCl_3) δ ppm 8.58 (1 H, dd), 7.66 - 7.56 (2 H, m), 7.22 (1 H, dd), 7.19 - 7.09 (3 H, m), 6.36 (1 H, s), 4.50 - 4.36 (2 H, m), 4.25 - 4.17 (2 H, m), 4.04 - 3.95 (1 H, m), 3.91 - 3.62 (5 H, m), 3.50 (1 H, dd), 3.14 (4 H, s), 2.98 (2 H, t)

Cpd#	NMR data (δ)
21	(^1H , CDCl_3) δ ppm 8.55 (1 H, dd), 8.44 (2 H, dd), 7.63 (1 H, d), 7.22 (1 H, dd), 7.15 (1 H, s), 6.36 (1 H, s), 4.53 - 4.36 (2 H, m), 4.26 - 4.17 (2 H, m), 4.07 - 3.95 (1 H, m), 3.93 - 3.61 (5 H, m), 3.51 (1 H, dd), 3.17 (4 H, s), 2.99 (2 H, t)
22	(^1H , CDCl_3) δ ppm 8.51 (1 H, br. s.), 7.94 (1 H, s), 7.79 - 7.67 (2 H, m), 7.61 (1 H, s), 7.58 - 7.47 (2 H, m), 7.33 (1 H, t), 6.67 (1 H, br. s.), 6.43 (1 H, s), 4.56 - 4.40 (2 H, m), 4.29 (2 H, t), 4.04 (1 H, td), 3.97 - 3.65 (5 H, m), 3.61 - 3.48 (1 H, m), 3.11 (2 H, t)
23	(^1H , CDCl_3) δ ppm 7.76 (1 H, d), 7.58 (1 H, d), 7.50 (1 H, s), 7.45 - 7.32 (2 H, m), 7.13 - 7.01 (2 H, m), 6.43 (1 H, s), 4.45 (2 H, m), 4.27 (2 H, t), 4.01 (1 H, m), 3.94 - 3.63 (8 H, m), 3.52 (1 H, t), 3.08 (2 H, t)
24	(^1H , CDCl_3) δ ppm 8.56 - 8.45 (1 H, m), 8.38 (1 H, d), 7.83 (1 H, d), 7.63 (1 H, dd), 7.54 (1 H, s), 7.42 (1 H, t), 6.45 (1 H, s), 4.54 - 4.36 (2 H, m), 4.31 - 4.23 (2 H, m), 4.05 - 3.96 (4 H, m), 3.93 - 3.62 (5 H, m), 3.52 (1 H, dd), 3.13 (2 H, t)
25	(^1H , CDCl_3) δ ppm 8.17 (1 H, d), 8.01 (1 H, s), 7.78 (1 H, d), 7.72 - 7.60 (3 H, m), 7.57 (1 H, d), 6.42 (1 H, s), 4.52 - 4.38 (2 H, m), 4.32 - 4.24 (2 H, m), 4.01 (1 H, qd), 3.93 - 3.62 (5 H, m), 3.51 (1 H, dd), 3.10 (2 H, t)
26	(^1H , CDCl_3) δ ppm 7.74 (1 H, d), 7.57 (3 H, d), 7.48 (1 H, s), 7.01 (2 H, d), 6.40 (1 H, s), 4.51 - 4.36 (2 H, m), 4.25 (2 H, t), 3.99 (1 H, dd), 3.92 - 3.60 (8 H, m), 3.50 (1 H, t), 3.06 (2 H, t)
27	(^1H , CDCl_3) δ ppm 8.18 (1 H, s), 7.84 (3 H, d), 7.68 (1 H, dd), 7.64 - 7.55 (2 H, m), 6.47 (1 H, s), 4.57 - 4.39 (2 H, m), 4.30 (2 H, t), 4.09 - 3.98 (1 H, m), 3.96 - 3.64 (5 H, m), 3.54 (1 H, dd), 3.13 (2 H, t)
28	(^1H , CDCl_3) δ ppm 8.45 (1 H, dd), 7.88 - 7.74 (2 H, m), 7.65 (1 H, d), 7.58 (1 H, s), 7.36 - 7.25 (1 H, m), 6.46 (1 H, s), 4.57 - 4.40 (2 H, m), 4.29 (2 H, t), 4.03 (1 H, td), 3.96 - 3.64 (5 H, m), 3.60 - 3.46 (1 H, m), 3.12 (2 H, t)
29	(^1H , CDCl_3) δ ppm 7.99 (1 H, s), 7.80 (1 H, d), 7.64 (1 H, d), 7.56 (1 H, s), 7.50 - 7.32 (3 H, m), 6.45 (1 H, s), 4.56 - 4.40 (2 H, m), 4.29 (2 H, t), 4.11 - 3.97 (1 H, m), 3.95 - 3.64 (5 H, m), 3.60 - 3.46 (1 H, m), 3.11 (2 H, t), 2.26 (5 H, s)
30	(^1H , CDCl_3) δ ppm 7.63 (1 H, d), 7.39 (1 H, dd), 7.33 (1 H, s), 6.41 - 6.34 (1 H, m), 4.53 - 4.39 (2 H, m), 4.29 - 4.18 (1 H, m), 4.06 - 3.97 (1 H, m), 3.93 - 3.64 (5 H, m), 3.52 (1 H, dd), 3.00 (2 H, t), 1.56 - 1.45 (1 H, m), 1.01 - 0.83 (4 H, m)
31	(^1H , CDCl_3) δ ppm 7.67 (1 H, d), 7.45 (1 H, dd), 7.39 (1 H, s), 6.40 (1 H, s), 4.53 - 4.38 (2 H, m), 4.31 - 4.17 (2 H, m), 4.02 (1 H, dd), 3.94 - 3.64 (5 H, m), 3.52 (1 H, dd), 3.08 - 2.96 (2 H, m), 2.18 - 1.75 (7 H, m), 1.29 (1 H, s)

Cpd#	NMR data (δ)
32	(^1H , CDCl_3) δ ppm 9.32 (1 H, s), 9.04 (2 H, s), 7.90 (1 H, d), 7.66 (1 H, dd), 7.58 (1 H, s), 6.48 (1 H, s), 4.56 - 4.40 (2 H, m), 4.37 - 4.27 (2 H, m), 4.11 - 3.99 (1 H, m), 3.96 - 3.64 (5 H, m), 3.60 - 3.46 (1 H, m), 3.17 (2 H, t)
33	(^1H , CDCl_3) δ ppm 7.67 (1 H, d), 7.42 (1 H, d), 7.32 (1 H, s), 6.39 (1 H, s), 6.31 (1 H, t), 4.54 - 4.38 (2 H, m), 4.25 (2 H, t), 4.02 (1 H, dd), 3.94 - 3.64 (5 H, m), 3.53 (1 H, t), 3.03 (2 H, t), 2.45 (2 H, d), 2.29 (2 H, dd), 1.91 - 1.58 (4 H, m)
34	(^1H , $\text{DMSO}-d_6$) δ ppm 8.07 (1 H, d), 7.97 (1 H, s), 7.78 - 7.72 (2 H, m), 7.62 - 7.53 (2 H, m), 7.39 (1 H, d), 6.68 (1 H, s), 6.52 (1 H, d), 4.33 - 4.24 (2 H, m), 4.09 (2 H, t), 3.91 - 3.76 (5 H, m), 3.71 - 3.58 (2 H, m), 3.55 - 3.47 (1 H, m), 3.40 (1 H, m), 3.29 (1 H, s), 3.10 (2 H, t)
35	(^1H , CDCl_3) δ ppm 8.84 (1 H, d), 7.96 (1 H, d), 7.85 (1 H, d), 7.64 (1 H, dd), 7.56 (1 H, s), 7.40 (1 H, d), 6.47 (1 H, s), 4.55 - 4.40 (2 H, m), 4.37 - 4.23 (2 H, m), 4.04 (1 H, m), 3.95 - 3.65 (5 H, m), 3.54 (1 H, dd), 3.14 (2 H, t), 2.74 (3 H, s)
36	(^1H , CDCl_3) δ ppm 8.69 (1 H, d), 7.78 - 7.72 (2 H, m), 7.65 - 7.57 (3 H, m), 7.33 (1 H, m), 6.43 (1 H, s), 4.54 - 4.39 (2 H, m), 4.27 (2 H, t), 4.05 (1 H, s), 3.95 - 3.64 (5 H, m), 3.53 (1 H, dd), 3.07 (2 H, t)
37	(^1H , CDCl_3) δ ppm 7.68 (1 H, d), 7.47 (1 H, dd), 7.42 (1 H, s), 6.40 (1 H, s), 4.53 - 4.41 (2 H, m), 4.38 (2 H, s), 4.24 (2 H, t), 4.07 - 3.96 (1 H, m), 3.93 - 3.63 (5 H, m), 3.55 - 3.33 (4 H, m), 3.03 (2 H, t)
38	(^1H , CDCl_3) δ ppm 7.67 (1 H, d), 7.44 (1 H, dd), 7.39 (1 H, s), 6.39 (1 H, s), 4.50 - 4.40 (2 H, m), 4.26 - 4.20 (2 H, m), 4.04 - 3.98 (1 H, m), 3.91 - 3.65 (5 H, m), 3.52 (1 H, dd), 3.02 (2 H, t), 2.90 - 2.83 (2 H, m), 2.75 - 2.68 (2 H, m)
39	(^1H , CDCl_3) δ ppm 7.64 (1 H, d), 7.42 (1 H, dd), 7.37 (1 H, s), 6.38 (1 H, s), 4.51 (2 H, s), 4.48 - 4.378 (2 H, m), 4.19 (2 H, t), 4.03 - 3.96 (1 H, m), 3.89 - 3.64 (5 H, m), 3.51 (1 H, dd), 2.99 (2 H, t)
40	(^1H , CDCl_3) δ ppm 7.65 (1 H, d), 7.50 - 7.44 (3 H, m), 7.41 (1 H, s), 6.94 - 6.84 (2 H, m), 6.35 (1 H, s), 4.45 - 4.36 (2 H, m), 4.19 (2 H, t), 4.04 - 3.91 (1 H, m), 3.87 - 3.62 (8 H, m), 3.48 (1 H, dd), 7.81 (1 H, d), 7.59 - 7.48 (2 H, m), 7.43 (1 H, s), 7.41 - 7.33 (3 H, m), 6.47 (1 H, s), 4.54 - 4.42 (2 H, m), 4.30 (2 H, t), 4.03 (1 H, dt), 3.95 - 3.65 (5 H, m), 3.54 (1 H, dd), 3.11 (2 H, t)
41	(^1H , CDCl_3) δ ppm 8.78 (1 H, d), 8.59 (1 H, dd), 7.83 (1 H, dt), 7.71 (1 H, d), 7.54 (1 H, dd), 7.49 (1 H, s), 7.39 - 7.25 (1 H, m), 6.40 (1 H, s), 4.51 - 4.33 (2 H, m), 4.30 - 4.16 (2 H, m), 4.07 - 3.93 (1 H, m), 3.90 - 3.59 (5 H, m), 3.49 (1 H, dd), 3.04 (2 H, t)

Cpd#	NMR data (δ)
42	(^1H , CDCl_3) δ ppm 7.93 (2 H, d), 7.78 (1 H, d), 7.72 - 7.59 (3 H, m), 7.54 (1 H, s), 6.42 (1 H, s), 4.49 - 4.35 (2 H, m), 4.23 (2 H, t), 4.05 - 3.95 (1 H, m), 3.91 - 3.61 (5 H, m), 3.49 (1 H, t), 3.16 - 2.97 (5 H, m)
43	(^1H , CDCl_3) δ ppm 7.81 (1 H, d), 7.64 (1 H, d), 7.55 (1 H, br. s.), 7.50 - 7.37 (1 H, m), 7.24 (1 H, d), 7.18 (1 H, br. s.), 7.00 (1 H, d), 6.46 (1 H, br. s.), 4.47 (2 H, d), 4.30 (2 H, br. s.), 4.04 (1 H, br. s.), 3.98 - 3.66 (7 H, m), 3.54 (1 H, t), 3.12 (2 H, br. s.), 1.83 (1 H, br. s.)
44	(^1H , CDCl_3) δ ppm 7.81 (1 H, d), 7.59 - 7.48 (2 H, m), 7.43 (1 H, s), 7.41 - 7.33 (3 H, m), 6.47 (1 H, s), 4.54 - 4.42 (2 H, m), 4.30 (2 H, t), 4.03 (1 H, dt), 3.95 - 3.65 (5 H, m), 3.54 (1 H, dd), 3.11 (2 H, t)
45	(^1H , CDCl_3) δ ppm 7.62 (1 H, d), 7.39 (1 H, d), 7.34 (1 H, s), 6.36 (1 H, s), 4.51 - 4.34 (2 H, m), 4.19 (2 H, t), 3.99 (1 H, td), 3.91 - 3.59 (7 H, m), 3.50 (1 H, dd), 2.98 (2 H, t), 2.74 (2 H, t)
46	(^1H , CDCl_3) δ ppm 7.64 (1 H, d), 7.02 (1 H, dd), 6.92 (1 H, d), 6.30 (1 H, s), 6.13 (1 H, s), 5.09 (2 H, s), 4.51 - 4.36 (2 H, m), 4.22 (2 H, t), 4.07 - 3.97 (1 H, m), 3.92 - 3.61 (8 H, m), 3.57 - 3.44 (1 H, m), 2.99 (2 H, t), 2.30 (3 H, s)
47	(^1H , CDCl_3) δ ppm 7.62 (1 H, d), 7.37 (1 H, d), 7.00 (1 H, dd), 6.90 (1 H, d), 6.33 (1 H, d), 6.27 (1 H, s), 5.12 (2 H, s), 4.47 - 4.34 (2 H, m), 4.23 - 4.143 (2 H, m), 4.02 - 3.89 (4 H, m), 3.88 - 3.59 (5 H, m), 3.48 (1 H, dd), 2.96 (2 H, t)
48	(^1H , CDCl_3) δ ppm 7.67 (1 H, d), 6.99 (1 H, dd), 6.90 (1 H, d), 6.29 (1 H, s), 5.33 (2 H, s), 4.47 - 4.35 (2 H, m), 4.20 (2 H, t), 4.03 - 3.93 (1 H, m), 3.89 - 3.60 (5 H, m), 3.48 (1 H, dd), 2.99 (2 H, t), 2.45 (3 H, s)
49	(^1H , CDCl_3) δ ppm 7.77 (1 H, d), 7.66 - 7.57 (3 H, m), 7.52 (1 H, d), 7.04 (2 H, d), 6.43 (1 H, s), 4.55 - 4.40 (2 H, m), 4.35 - 4.24 (2 H, m), 4.04 (1 H, dd), 3.97 - 3.66 (9 H, m), 3.54 (1 H, dd), 3.32 - 3.20 (4 H, m), 3.10 (2 H, s)
50	(^1H , CDCl_3) δ ppm 8.06 (1 H, dd), 7.92 - 7.78 (2 H, m), 7.63 (1 H, d), 7.56 (1 H, s), 7.36 - 7.24 (1 H, m), 6.47 (1 H, s), 4.55 - 4.38 (2 H, m), 4.28 (2 H, t), 4.13 - 3.99 (1 H, m), 3.95 - 3.65 (5 H, m), 3.54 (1 H, t), 3.11 (2 H, t)
51	(^1H , CDCl_3) δ ppm 7.95 (1 H, t), 7.80 (1 H, d), 7.67 - 7.54 (3 H, m), 7.54 - 7.45 (1 H, m), 6.44 (1 H, s), 4.52 - 4.35 (2 H, m), 4.31 - 4.19 (2 H, m), 4.09 - 3.96 (1 H, m), 3.95 - 3.64 (5 H, m), 3.58 - 3.46 (1 H, m), 3.10 (2 H, t)
52	(^1H , CDCl_3) δ ppm 7.64 (1 H, d), 7.40 (1 H, dd), 7.35 (1 H, d), 6.38 (1 H, s), 4.52 - 4.39 (2 H, m), 4.26 - 4.16 (2 H, m), 4.04 - 3.96 (1 H, m), 3.93 - 3.63 (5 H, m), 3.51 (1 H, dd), 3.00 (2 H, t), 1.36 (9 H, s)

Cpd#	NMR data (δ)
53	(^1H , CDCl_3) δ ppm 8.72 - 8.64 (2 H, m), 7.75 (1 H, d), 7.59 (1 H, dd), 7.54 (1 H, s), 7.48 - 7.42 (2 H, m), 6.44 (1 H, s), 4.55 - 4.39 (2 H, m), 4.34 - 4.22 (2 H, m), 4.03 (1 H, dd), 3.95 - 3.64 (5 H, m), 3.53 (1 H, dd), 3.08 (2 H, t)
54	(^1H , CDCl_3) δ ppm 7.68 (1 H, d), 6.98 (1 H, dd), 6.89 (1 H, d), 6.31 (1 H, s), 6.23 (1 H, s), 5.22 (2 H, s), 4.52 - 4.36 (2 H, m), 4.29 - 4.18 (2 H, m), 4.07 - 3.96 (1 H, m), 3.93 - 3.63 (5 H, m), 3.51 (1 H, dd), 3.06 - 2.96 (2 H, m), 2.35 (3 H, s)
55	(^1H , CDCl_3) δ ppm 7.60 (1 H, d), 7.39 - 7.35 (1 H, m), 7.32 (1 H, d), 6.33 (1 H, s), 4.45 - 4.35 (2 H, m), 4.17 (2 H, t), 3.97 (1 H, m), 3.87 - 3.60 (5 H, m), 3.47 (1 H, dd), 2.95 (2 H, t), 1.62 (6 H, s)
56	(^1H , CDCl_3) δ ppm 8.24 (1 H, dd), 7.78 (1 H, d), 7.67 (1 H, dd), 7.61 (1 H, dd), 7.54 (1 H, s), 7.04 (1 H, dd), 6.45 (1 H, s), 4.52 - 4.41 (2 H, m), 4.28 (2 H, t), 4.02 (4 H, s), 3.93 - 3.64 (5 H, m), 3.52 (1 H, dd), 3.10 (2 H, t)
58	(^1H , CDCl_3) δ ppm 7.70 (1 H, d), 7.43 (1 H, dd), 7.33 (1 H, d), 6.39 (1 H, s), 6.31 (1 H, dt), 4.53 - 4.34 (4 H, m), 4.31 - 4.19 (2 H, m), 4.06 - 3.94 (3 H, m), 3.92 - 3.61 (5 H, m), 3.51 (1 H, dd), 3.04 (2 H, t), 2.62 - 2.50 (2 H, m)
59	(^1H , CDCl_3) δ ppm 9.02 (1 H, d), 8.09 (1 H, dd), 7.97 - 7.80 (2 H, m), 7.67 (1 H, dd), 7.59 (1 H, d), 6.48 (1 H, s), 4.56 - 4.38 (2 H, m), 4.36 - 4.25 (2 H, m), 4.03 (1 H, m), 3.95 - 3.63 (5 H, m), 3.53 (1 H, dd), 3.16 (2 H, t)
60	(^1H , CDCl_3) δ ppm 8.43 (1 H, d), 7.86 - 7.74 (2 H, m), 7.56 (1 H, dd), 7.47 (1 H, d), 6.80 (1 H, d), 6.43 (1 H, s), 5.44 - 5.31 (1 H, m), 4.53 - 4.37 (2 H, m), 4.26 (2 H, t), 4.01 (1 H, m), 3.92 - 3.62 (5 H, m), 3.51 (1 H, dd), 3.09 (2 H, t), 1.40 (6 H, d)
61	(^1H , CDCl_3) δ ppm 8.44 (1 H, d), 7.84 (1 H, dd), 7.79 (1 H, d), 7.57 (1 H, dd), 7.48 (1 H, d), 6.85 (1 H, d), 6.43 (1 H, s), 4.54 - 4.37 (4 H, m), 4.27 (2 H, t), 4.02 (1 H, m), 3.93 - 3.62 (5 H, m), 3.52 (1 H, dd), 3.10 (2 H, t), 1.45 (3 H, t)
62	(^1H , CDCl_3) δ ppm 8.52 (1 H, d), 7.85 - 7.72 (2 H, m), 7.56 (1 H, dd), 7.47 (1 H, d), 6.75 (1 H, d), 6.42 (1 H, s), 4.52 - 4.38 (2 H, m), 4.32 - 4.22 (2 H, m), 4.06 - 3.97 (1 H, m), 3.93 - 3.58 (13 H, m), 3.52 (1 H, dd), 3.09 (2 H, s)
63	(^1H , CDCl_3) δ ppm 7.77 (1 H, d), 7.60 (1 H, dd), 7.52 (1 H, d), 7.21 - 7.12 (1 H, m), 6.99 (2 H, ddd), 6.45 (1 H, s), 4.53 - 4.40 (2 H, m), 4.28 (2 H, t), 4.06 - 3.98 (1 H, m), 3.95 (3 H, s), 3.92 - 3.63 (8 H, m), 3.52 (1 H, dd), 3.08 (2 H, t)
64	(^1H , CDCl_3) δ ppm 8.15 (1 H, d), 7.82 (1 H, d), 7.50 (1 H, dd), 7.45 - 7.40 (1 H, m), 6.91 (1 H, d), 6.46 (1 H, s), 4.54 - 4.40 (2 H, m), 4.29 (2 H, t), 4.11 (2 H, s), 4.07 - 3.98 (1 H, m), 3.94 - 3.64 (4 H, m), 3.53 (1 H, dd), 3.11 (2 H, t)

Cpd#	NMR data (δ)
65	(^1H , CDCl_3) δ ppm 8.62 (1 H, d), 7.84 (1 H, d), 7.67 (1 H, dd), 7.59 (1 H, d), 7.42 (1 H, s), 7.36 (1 H, dd), 6.46 (1 H, s), 4.54 - 4.38 (2 H, m), 4.33 - 4.24 (2 H, m), 4.07 - 3.97 (1 H, m), 3.94 - 3.63 (5 H, m), 3.52 (1 H, dd), 3.13 (2 H, t), 2.68 (3 H, s)
66	(^1H , CDCl_3) δ ppm 8.88 (1 H, s), 8.82 (1 H, d), 7.88 (1 H, d), 7.68 (1 H, dd), 7.61 (1 H, dd), 7.53 (1 H, s), 6.45 (1 H, s), 4.49 - 4.38 (2 H, m), 4.27 (2 H, t), 4.00 (1 H, ddt), 3.90 - 3.61 (5 H, m), 3.49 (1 H, dd), 3.13 (2 H, t)
67	(^1H , CDCl_3) δ ppm 7.76 (1 H, d), 7.58 (1 H, dd), 7.50 (1 H, s), 6.88 - 7.03 (3 H, m), 6.44 (1 H, s), 4.54 - 4.39 (2 H, m), 4.28 (2 H, t), 4.03 (1 H, m), 3.95 - 3.63 (1 H, m), 3.53 (1 H, dd), 3.08 (2 H, t)
68	(^1H , CDCl_3) δ ppm 8.50 (1 H, d), 7.74 (2 H, td), 7.55 (1 H, dd), 7.45 (1 H, s), 6.75 (1 H, d), 6.40 (1 H, s), 4.56 - 4.38 (2 H, m), 4.26 (2 H, t), 4.01 (1 H, m), 3.94 - 3.58 (9 H, m), 3.52 (1 H, dd), 3.08 (2 H, t), 1.70 (6 H, br. s.)
69	(^1H , CDCl_3) δ ppm 8.21 (1 H, dd), 7.78 (1 H, d), 7.66 (2 H, ddd), 7.56 (1 H, s), 7.01 (1 H, dd), 6.45 (1 H, s), 4.54 - 4.39 (4 H, m), 4.28 (2 H, t), 4.03 (1 H, m), 3.94 - 3.64 (5 H, m), 3.53 (1 H, dd), 3.09 (2 H, t), 1.42 (3 H, t)
70	(^1H , CDCl_3) δ ppm 7.75 (1 H, d), 7.64 (1 H, d), 7.60 (1 H, dd), 7.52 (1 H, d), 6.47 (1 H, d), 6.43 (1 H, s), 4.53 - 4.39 (2 H, m), 4.33 - 4.23 (2 H, m), 4.03 (6 H, d), 3.94 - 3.64 (5 H, m), 3.59 - 3.48 (2 H, m), 3.08 (2 H, t)
71	(^1H , CDCl_3) δ ppm 9.00 (1 H, s), 8.87 (1 H, d), 7.87 (1 H, d), 7.63 (1 H, dd), 7.56 (1 H, s), 7.52 - 7.48 (1 H, m), 6.45 (1 H, s), 4.51 - 4.36 (2 H, m), 4.26 (2 H, t), 3.99 (1 H, ddt), 3.90 - 3.60 (5 H, m), 3.54 - 3.44 (1 H, m), 3.12 (2 H, t)
72	(^1H , CDCl_3) δ ppm 7.65 (1 H, d), 7.34 (1 H, d), 7.30 (1 H, s), 6.36 (1 H, s), 4.48 (2 H, s), 4.46 - 4.34 (2 H, m), 4.18 (2 H, t), 4.02 - 3.92 (1 H, m), 3.89 - 3.60 (5 H, m), 3.54 - 3.42 (1 H, m), 3.00 (2 H, t), 1.30 (9 H, s)
73	(^1H , CDCl_3) δ ppm 8.19 (1 H, dd), 7.69 (1 H, d), 7.38 (2 H, ddd), 7.29 (1 H, d), 6.72 (1 H, dd), 6.40 (1 H, s), 4.47 - 4.37 (2 H, m), 4.24 (2 H, t), 4.03 - 3.93 (1 H, m), 3.89 - 3.59 (5 H, m), 3.49 (1 H, t), 3.18 - 3.08 (4 H, m), 3.04 (2 H, t), 1.82 - 1.72 (4 H, m)
74	(^1H , CDCl_3) δ ppm 8.51 (1 H, d), 7.75 (2 H, td), 7.56 (1 H, dd), 7.46 (1 H, s), 6.48 (1 H, d), 6.41 (1 H, s), 4.56 - 4.37 (2 H, m), 4.27 (2 H, t), 4.02 (1 H, dd), 3.95 - 3.63 (5 H, m), 3.61 - 3.45 (5 H, m), 3.09 (2 H, t), 2.14 - 1.99 (4 H, m)
75	(^1H , CDCl_3) δ ppm 7.70 - 7.57 (2 H, m), 7.47 - 7.20 (5 H, m), 7.05 (1 H, dd), 6.96 (1 H, d), 6.27 (1 H, s), 5.26 (2 H, s), 4.47 - 4.32 (2 H, m), 4.23 - 4.15 (2 H, m), 4.03 - 3.91 (1 H, m), 3.89 - 3.56 (5 H, m), 3.47 (1 H, dd), 2.97 (2 H, t)

Cpd#	NMR data (δ)
76	(^1H , CDCl_3) δ ppm 7.67 (1 H, d), 7.05 (1 H, dd), 6.96 (1 H, d), 6.75 (1 H, s), 6.31 (1 H, s), 5.18 (2 H, s), 4.53 - 4.34 (2 H, m), 4.22 (2 H, t), 4.00 (1 H, m), 3.92 - 3.62 (5 H, m), 3.50 (1 H, dd), 3.00 (2 H, t), 1.32 (9 H, s)
77	(^1H , CDCl_3) δ ppm 7.64 (1 H, d), 7.00 (1 H, dd), 6.91 (1 H, d), 6.27 (1 H, s), 5.17 (2 H, s), 4.47 - 4.32 (2 H, m), 4.23 - 4.14 (2 H, m), 3.96 (1 H, qd), 3.89 - 3.58 (5 H, m), 3.47 (1 H, dd), 2.97 (2 H, t), 2.29 - 2.17 (1 H, m), 1.30 - 1.24 (4 H, m)
78	(^1H , CDCl_3) δ ppm 7.65 (1 H, d), 7.01 (1 H, dd), 6.92 (1 H, d), 6.27 (1 H, s), 5.22 (2 H, s), 4.47 - 4.30 (2 H, m), 4.24 - 4.13 (2 H, m), 3.96 (1 H, m), 3.90 - 3.580 (5 H, m), 3.47 (1 H, dd), 3.02 - 2.85 (4 H, m), 1.41 (3 H, t)
79	(^1H , CDCl_3) δ ppm 7.68 (1 H, d), 7.05 (1 H, dd), 6.95 (1 H, d), 6.32 (1 H, s), 5.26 (2 H, s), 4.52 - 4.37 (2 H, m), 4.28 - 4.20 (2 H, m), 4.06 - 3.96 (1 H, m), 3.94 - 3.62 (5 H, m), 3.51 (1 H, dd), 3.02 (2 H, t), 2.67 (3 H, s)
80	(^1H , CDCl_3) δ ppm 7.65 (1 H, d), 7.02 (1 H, dd), 6.93 (1 H, d), 6.28 (1 H, s), 5.22 (2 H, s), 4.47 - 4.33 (2 H, m), 4.19 (2 H, t), 4.03 - 3.91 (1 H, m), 3.89 - 3.59 (5 H, m), 3.48 (1 H, dd), 3.26 (1 H, dt), 2.99 (2 H, t), 1.43 (6 H, d)
81	(^1H , CDCl_3) δ ppm 7.63 (1 H, d), 7.40 (1 H, dd), 7.34 (1 H, s), 6.38 (1 H, s), 4.53 - 4.37 (2 H, m), 4.27 - 4.17 (2 H, m), 4.05 - 3.96 (1 H, m), 3.93 - 3.63 (5 H, m), 3.58 - 3.46 (1 H, m), 3.00 (2 H, t), 2.88 (1 H, m), 2.13 - 1.97 (2 H, m), 1.90 - 1.56 (6 H, m)
82	(^1H , CDCl_3) δ ppm 7.64 (1 H, d), 7.41 (1 H, d), 7.35 (1 H, s), 6.38 (1 H, s), 4.52 - 4.36 (2 H, m), 4.22 (2 H, t), 4.01 (1 H, m), 3.93 - 3.63 (5 H, m), 3.51 (1 H, t), 3.00 (2 H, t), 2.71 - 2.58 (1 H, m), 1.99 - 1.31 (10 H, m)
83	(^1H , CDCl_3) δ ppm 7.64 (1 H, d), 7.40 (1 H, dd), 7.35 (1 H, s), 6.38 (1 H, s), 4.49 - 4.40 (2 H, m), 4.22 (2 H, t), 4.05 - 3.97 (1 H, m), 3.91 - 3.65 (5 H, m), 3.51 (1 H, dd), 3.00 (2 H, t), 2.88 - 2.78 (1 H, m), 1.31 (6 H, d)
84	(^1H , CDCl_3) δ ppm 7.64 (1 H, d), 7.40 (1 H, dd), 7.35 (1 H, s), 6.38 (1 H, s), 4.51 - 4.38 (2 H, m), 4.26 - 4.18 (2 H, m), 4.07 - 3.97 (1 H, m), 3.93 - 3.64 (5 H, m), 3.52 (1 H, dd), 3.00 (2 H, t), 2.47 (2 H, t), 1.70 - 1.45 (4 H, m), 0.99 (3 H, t)
85	(^1H , CDCl_3) δ ppm 7.65 (1 H, d), 7.45 (1 H, dd), 7.41 - 7.31 (5 H, m), 7.27 (1 H, s), 6.37 (1 H, s), 4.49 - 4.34 (2 H, m), 4.24 - 4.17 (2 H, m), 4.02 - 3.94 (1 H, m), 3.90 - 3.60 (7 H, m), 3.54 (2 H, s), 3.52 - 3.45 (1 H, m), 3.00 (2 H, t), 2.42 (3 H, s)
86	(^1H , CDCl_3) δ ppm 7.63 (1 H, d), 7.42 (1 H, dd), 7.36 (1 H, s), 6.37 (1 H, s), 4.70 - 4.65 (1 H, m), 4.47 - 4.37 (2 H, m), 4.20 (2 H, t), 4.00 - 3.97 (1 H, m), 3.88 - 3.63 (5 H, m), 3.49 (1 H, t), 2.99 (2 H, t), 1.93 - 1.65 (4 H, m), 1.00 - 0.97 (6 H, m)

Cpd#	NMR data (δ)
87	(^1H , CDCl_3) δ ppm 7.63 (1 H, d), 7.42 (1 H, d), 7.36 (1 H, s), 6.37 (1 H, s), 4.80 - 4.77 (1 H, m), 4.46 - 4.37 (2 H, m), 4.20 (2 H, t), 4.01-3.97 (1 H, m), 3.89 - 3.65 (5 H, m), 3.49 (1 H, t), 2.99 (2 H, t), 1.99 (1 H, d), 1.57 (3 H, d)
88	(^1H , CDCl_3) δ ppm 7.60 (1 H, d), 7.07 (1 H, d), 6.99 (1 H, s), 6.35 (1 H, s), 4.49 - 4.38 (2 H, m), 4.21 (2 H, t), 4.01 (1 H, m), 3.94 - 3.62 (5 H, m), 3.51 (1 H, t), 2.99 (2 H, t), 2.07 - 1.86 (1 H, m), 1.15 - 1.06 (2 H, m), 0.85 - 0.78 (2 H, m)
89	(^1H , $\text{DMSO-}d_6$) δ ppm 8.00 (1 H, d), 7.46 (1 H, s), 7.40 (1 H, d), 6.68 (1 H, s), 5.53 (1 H, d), 4.43 - 4.41 (1 H, m), 4.26 - 4.25 (2 H, m), 4.01 (2 H, t), 3.87 - 3.40 (7 H, m), 3.00 (2 H, t), 1.67 - 1.65 (2 H, m), 0.98 (3 H, t)
90	(^1H , $\text{DMSO-}d_6$) δ ppm 8.00 (1 H, d), 7.46 (1 H, s), 7.40 (1 H, d), 6.68 (1 H, s), 5.53 (1 H, d), 4.28 - 4.25 (3 H, m), 4.01 (2 H, t), 3.85 - 3.38 (7 H, m), 3.00 (2 H, t), 1.85 - 1.80 (1 H, m), 0.98 (6 H, t)
91	(^1H , $\text{DMSO-}d_6$) δ ppm 8.00 (1 H, d), 7.44 (1 H, s), 7.38 (1 H, d), 6.68 (1 H, s), 5.27 (1 H, s), 4.26 - 4.25 (2 H, m), 4.01 (2 H, t), 3.86 - 3.74 (3 H, m), 3.67 - 3.60 (2 H, m), 3.52 - 3.46 (1 H, m), 3.37 - 3.35 (1 H, m), 3.00 (2 H, t), 1.66 - 1.62 (4 H, m), 0.99 (6 H, t)
92	(^1H , CDCl_3) δ ppm 7.71 (2 H, d), 7.66 (1 H, d), 7.47 (1 H, d), 7.43 - 7.32 (4 H, m), 6.38 (1 H, s), 4.47 - 4.37 (2 H, m), 4.20 (2 H, t), 4.00 - 3.98 (1 H, m), 3.97 - 3.68 (5 H, m), 3.49 (1 H, t), 3.00 (2 H, t), 2.53 (1 H, s), 1.88 (3 H, s)
93	(^1H , CDCl_3) δ ppm 7.64 (1 H, d), 7.43 - 7.29 (7 H, m), 6.37 (1 H, s), 4.46 - 4.37 (2 H, m), 4.21 (2 H, t), 3.99 - 3.95 (3 H, m), 3.88 - 3.63 (7 H, m), 3.49 (1 H, t), 2.99 (2 H, t)
94	(^1H , CDCl_3) δ ppm 7.57 - 7.48 (1 H, d), 7.38 (1 H, s), 6.68 - 6.55 (1 H, d), 6.49 (1 H, s), 6.35 (1 H, s), 6.28 (1 H, s), 6.20 (1 H, s), 4.60 - 4.50 (1 H, m), 4.38 - 4.30 (4 H, m), 4.22 - 4.15 (2 H, m), 4.03 - 3.93 (1 H, m), 3.95 - 3.60 (5 H, m), 3.55 - 3.40 (1 H, t), 2.98 - 2.85 (2 H, m)
95	(^1H , CDCl_3) δ ppm 7.86 (1 H, s), 7.77 (1 H, s), 7.70 (1 H, d), 7.51 (1 H, d), 7.42 (1 H, s), 6.39 (1 H, s), 4.51 - 4.37 (2 H, m), 4.330 - 4.20 (4 H, m), 4.05 - 3.96 (1 H, m), 3.87 (5 H, s), 3.51 (1 H, dd), 3.05 (2 H, t), 1.57 (4 H, t)
96	(^1H , CDCl_3) δ ppm 7.85 (1 H, s), 7.75 (1 H, s), 7.70 (1 H, d), 7.50 (1 H, dd), 7.42 (1 H, s), 6.39 (1 H, s), 4.50 - 4.39 (2 H, m), 4.27 - 4.18 (4 H, m), 4.05 - 3.97 (1 H, m), 3.86 (5 H, m), 3.51 (1 H, dd), 3.05 (2 H, t), 1.88 - 1.79 (2 H, m), 1.69 - 1.59 (1 H, m), 1.00 (6 H, d)

Cpd#	NMR data (δ)
97	(^1H , CDCl_3) δ ppm 8.03 (1 H, d), 7.67 - 7.63 (2 H, m), 7.05 (1 H, d), 6.65 (1 H, s), 6.28 (1 H, dd), 4.30 - 4.23 (2 H, m), 4.04 (2 H, t), 3.89 - 3.75 (3 H, m), 3.63 (2 H, dd), 3.50 (1 H, d), 3.39 (1 H, dd), 3.04 (2 H, t), 2.37 (3 H, s)
98	(^1H , CDCl_3) δ ppm 7.64 (1 H, d), 7.42 (1 H, d), 7.36 (1 H, s), 6.37 (1 H, s), 4.66 - 4.61 (1 H, m), 4.47 - 4.37 (2 H, t), 4.20 (2 H, t), 4.00 - 3.88 (1 H, m), 3.88 - 3.46 (5 H, m), 3.49 (1 H, t), 2.99 (2 H, t), 1.90 - 1.89 (1 H, m), 1.81 - 1.77 (2 H, m), 1.57 - 1.54 (2 H, m), 1.00 (3 H, t)
99	(^1H , CDCl_3) δ ppm 7.75 (1 H, d), 7.33 - 7.28 (1 H, m), 7.22 (1 H, s), 6.41 (1 H, s), 4.50 - 4.39 (2 H, m), 4.29 - 4.23 (2 H, m), 4.05 - 3.97 (1 H, m), 3.85 (5 H, m), 3.54 - 3.47 (1 H, m), 3.07 (2 H, d), 2.36 (6 H, s)
100	(^1H , CDCl_3) δ ppm 8.36 (1 H, br. s.), 8.07 (1 H, br. s.), 7.99 (1 H, d), 7.64 - 7.70 (2 H, m), 6.65 (1 H, s), 4.31 - 4.22 (2 H, m), 4.05 (2 H, t), 3.90 - 3.84 (1 H, m), 3.79 (2 H, td), 3.70 - 3.57 (2 H, m), 3.51 - 3.50 (1 H, m), 3.55 - 3.45 (1 H, m), 3.39 (1 H, dd), 3.01 (2 H, t)
101	(^1H , CDCl_3) δ ppm 7.85 (1 H, d), 7.74 (1 H, d), 7.70 (1 H, d), 7.50 (1 H, dd), 7.44 - 7.38 (1 H, m), 6.38 (1 H, s), 4.48 - 4.348 (2 H, m), 4.27 - 4.17 (2 H, m), 4.15 (2 H, t), 4.04 - 3.96 (1 H, m), 3.90 - 3.64 (5 H, m), 3.50 (1 H, dd), 3.04 (2 H, t), 2.02 - 1.90 (2 H, m), 0.97 (3 H, t)
102	(^1H , CDCl_3) δ ppm 7.91 - 7.80 (2 H, m), 7.71 (1 H, dd), 7.62 - 7.45 (4 H, m), 6.45 (1 H, s), 4.49 - 4.39 (2 H, m), 4.27 (2 H, t), 4.06 - 3.94 (1 H, m), 3.92 - 3.61 (5 H, m), 3.51 (1 H, dd), 3.11 (2 H, t)
103	(^1H , CDCl_3) δ ppm 7.80 - 7.90 (2 H, m), 7.79 - 7.69 (1 H, m), 7.66 - 7.51 (4 H, m), 6.47 (1 H, s), 4.57 - 4.41 (2 H, m), 4.30 (2 H, t), 4.03 (1 H, dt), 3.95 - 3.64 (5 H, m), 3.53 (1 H, t), 3.14 (2 H, t)
104	(^1H , CDCl_3) δ ppm 7.65 (1 H, d), 7.01 (1 H, dd), 6.91 (1 H, d), 6.28 (1 H, s), 5.18 (2 H, s), 4.47 - 4.35 (2 H, m), 4.24 - 4.15 (2 H, m), 4.02 - 3.93 (1 H, m), 3.89 - 3.61 (5 H, m), 3.48 (1 H, dd), 2.98 (2 H, t), 2.29 - 2.18 (1 H, m), 1.33 - 1.20 (4 H, m)
105	(^1H , CDCl_3) δ ppm 7.65 (1 H, s), 7.51 - 7.46 (1 H, m), 7.45 - 7.41 (1 H, m), 6.37 (1 H, s), 4.49 - 4.36 (2 H, m), 4.21 (2 H, s), 4.04 - 3.94 (1 H, m), 3.90 - 3.610 (5 H, m), 3.53 - 3.44 (1 H, m), 3.25 (1 H, s), 3.00 (2 H, s)
106	(^1H , CDCl_3) δ ppm 9.18 (1 H, s), 8.88 (2 H, s), 7.72 (1 H, d), 7.60 - 7.53 (1 H, m), 7.51 (1 H, d), 6.40 (1 H, s), 4.49 - 4.36 (2 H, m), 4.23 (2 H, t), 4.02 - 3.95 (1 H, m), 3.89 - 3.58 (5 H, m), 3.49 (1 H, dd), 3.04 (2 H, t)

Cpd#	NMR data (δ)
107	(^1H , CDCl_3) δ ppm: 7.54 (1 H, d), 7.37 (1 H, dd), 7.32 (1 H, s), 7.26 - 7.22 (2 H, m), 6.81 (1 H, t), 6.75 (2 H, dd), 6.35 (1 H, s), 4.46 - 4.36 (2 H, m), 4.20 - 4.17 (4 H, m), 4.00 - 3.91 (1 H, m), 3.88 - 3.63 (5 H, m), 3.49 (1 H, t), 2.97 (2 H, t)
108	(^1H , CDCl_3) δ ppm 8.36 (1 H, d), 7.89 - 7.65 (3 H, m), 7.62 - 7.55 (3 H, m), 6.44 (1 H, s), 4.51 - 4.38 (2 H, m), 4.26 (2 H, t), 4.00 (1 H, m), 3.91 - 3.62 (5 H, m), 3.50 (1 H, dd), 3.22 - 3.14 (1 H, m), 3.09 (2 H, t), 1.40 (1 H, t)
110	(^1H , CDCl_3) δ ppm 7.70 - 7.60 (1 H, d), 7.50 - 7.42 (1 H, d), 7.39 (1 H, s), 6.37 (1 H, s), 4.50 - 4.35 (2 H, m), 4.25 - 4.15 (2 H, m), 4.05 - 3.95 (2 H, m), 3.92 - 3.60 (5 H, m), 3.56 - 3.45 (4 H, m), 3.05 - 3.95 (2 H, m), 2.15 - 1.95 (1 H, m), 1.15 - 1 (6 H, t)
111	(^1H , CDCl_3) δ ppm 7.58 (1 H, d), 7.33 (1 H, d), 7.27 (1 H, s), 6.32 (1 H, s), 4.45 - 4.34 (2 H, m), 4.17 (2 H, t), 3.96 (1 H, qd), 3.86 - 3.60 (5 H, m), 3.46 (1 H, dd), 2.94 (2 H, t), 1.46 (1 H, tt), 0.94 - 0.86 (2 H, m), 0.86 - 0.79 (2 H, m)
112	(^1H , CDCl_3) δ ppm 7.61 (1 H, d), 7.22 (1 H, d), 7.14 (1 H, s), 6.34 (1 H, s), 4.48 - 4.34 (2 H, m), 4.19 (2 H, t), 4.02 - 3.93 (1 H, m), 3.89 - 3.60 (5 H, m), 3.48 (1 H, t), 3.43 - 3.35 (1 H, m), 2.98 (2 H, t), 2.95 - 2.85 (1 H, m), 2.76 - 2.64 (1 H, m), 1.84 - 1.62 (3 H, m), 0.92 (6 H, dd)
113	(^1H , CDCl_3) δ ppm 7.67 (1 H, d), 7.44 (1 H, d), 7.39 (1 H, s), 6.38 (1 H, s), 5.00 (2 H, s), 4.50 - 4.36 (2 H, m), 4.21 (2 H, t), 4.03 - 3.95 (1 H, m), 3.91 - 3.61 (5 H, m), 3.49 (1 H, dd), 3.01 (2 H, t)
114	(^1H , CDCl_3) δ ppm 7.64 - 7.58 (1 H, m), 7.24 - 7.19 (1 H, m), 7.13 (1 H, s), 6.35 (1 H, s), 4.49 - 4.36 (2 H, m), 4.25 - 4.17 (2 H, m), 4.03 - 3.95 (1 H, m), 3.90 - 3.61 (5 H, m), 3.54 - 3.45 (1 H, m), 3.02 - 2.95 (2 H, m), 2.81 - 2.72 (2 H, m), 1.59 - 1.49 (2 H, m), 0.77 - 0.64 (1 H, m), 0.49 - 0.42 (2 H, m), 0.10 - 0.02 (2 H, m)
115	(^1H , CDCl_3) δ ppm 7.66 (1 H, d), 7.33 (1 H, d), 7.29 (1 H, s), 6.36 (1 H, s), 4.50 (2 H, s), 4.47 - 4.35 (2 H, m), 4.19 (2 H, t), 4.07 - 3.93 (2 H, m), 3.90 - 3.60 (5 H, m), 3.48 (1 H, t), 3.00 (2 H, t), 1.82 - 1.70 (6 H, m), 1.62 - 1.50 (2 H, m)
117	(^1H , CDCl_3) δ ppm 7.64 (1 H, d), 6.91 (1 H, dd), 6.80 (1 H, s), 6.28 (1 H, s), 6.11 - 6.01 (1 H, m), 5.45 (1 H, dd), 5.34 (1 H, dd), 4.61 (2 H, dd), 4.47 - 4.37 (2 H, m), 4.21 (2 H, t), 4.03 - 3.95 (1 H, m), 3.89 - 3.64 (5 H, m), 3.49 (1 H, t), 2.98 (2 H, t)
118	(^1H , CDCl_3) δ ppm 7.63 (1 H, d), 6.91 (1 H, dd), 6.80 (1 H, d), 6.28 (1 H, s), 6.11 - 6.01 (1 H, m), 5.44 (1 H, m), 5.34 (1 H, dd), 4.61 (2 H, dt), 4.47 - 4.37 (2 H, m), 4.21 (2 H, t), 4.03 - 3.95 (1 H, m), 3.89 - 3.63 (5 H, m), 3.49 (1 H, dd), 2.98 (2 H, t)

Cpd#	NMR data (δ)
119	(^1H , CDCl_3) δ ppm 7.68 (1 H, d), 7.36 (1 H, d), 7.31 (1 H, s), 6.41 - 6.38 (1 H, m), 4.61 (2 H, s), 4.47 - 4.37 (2 H, m), 4.25 - 4.17 (2 H, m), 3.99 (3 H, dt), 3.89 - 3.60 (6 H, m), 3.53 - 3.43 (3 H, m), 3.02 (2 H, t), 2.01 - 1.94 (2 H, m), 1.74 - 1.64 (2 H, m)
120	(^1H , CDCl_3) δ ppm 8.62 (1 H, br. s.), 8.53 (1 H, br. s.), 7.73 (1 H, d), 7.64 (1 H, d), 7.41 (1 H, dd), 7.35 (1 H, s), 7.29 (1 H, d), 6.36 (1 H, s), 4.48 - 4.34 (2 H, m), 4.20 (2 H, t), 4.03 - 3.93 (3 H, m), 3.89 - 3.61 (7 H, m), 3.48 (1 H, dd), 2.99 (2 H, t)
121	(^1H , CDCl_3) δ ppm 7.60 (1 H, d), 7.19 (1 H, dd), 7.10 (1 H, s), 6.34 (1 H, s), 4.49 - 4.34 (2 H, m), 4.20 (2 H, t), 4.02 - 3.93 (1 H, m), 3.90 - 3.60 (5 H, m), 3.48 (1 H, dd), 2.98 (2 H, t), 2.64 (2 H, t), 1.63 (2 H, m), 1.37 - 1.29 (4 H, m), 0.94 - 0.86 (3 H, m)
122	(^1H , CDCl_3) δ ppm 7.63 (1 H, d), 7.39 (1 H, dd), 7.31 (1 H, s), 6.36 (1 H, s), 4.50 - 4.39 (2 H, m), 4.20 (2 H, t), 4.02 - 3.98 (1 H, m), 3.89 - 3.66 (5 H, m), 3.49 (1 H, t), 2.96 (2 H, t), 1.59 - 1.48 (1 H, m), 0.98 - 0.81 (4 H, m)
123	(^1H , CDCl_3) δ ppm 7.64 - 7.58 (1 H, m), 7.24 - 7.19 (1 H, m), 7.13 (1 H, s), 6.35 (1 H, s), 4.49 - 4.36 (2 H, m), 4.24 - 4.17 (2 H, m), 4.03 - 3.95 (1 H, m), 3.91 - 3.62 (5 H, m), 3.49 (1 H, dd), 3.03 - 2.95 (2 H, m), 2.81 - 2.73 (2 H, m), 1.59 - 1.50 (2 H, m), 0.71 (1 H, s), 0.49 - 0.42 (2 H, m), 0.09 - 0.03 (2 H, m)
124	(^1H , CDCl_3) δ ppm 7.69 (1 H, d), 7.35 (1 H, d), 7.30 (1 H, s), 6.38 (1 H, s), 4.82 - 4.75 (2 H, m), 4.72 - 4.64 (3 H, m), 4.49 (2 H, s), 4.48 - 4.37 (2 H, m), 4.21 (2 H, dd), 4.03 - 3.95 (1 H, m), 3.90 - 3.62 (5 H, m), 3.49 (1 H, dd), 3.02 (2 H, t)
125	(^1H , CDCl_3) δ ppm 7.69 (1 H, d), 7.35 (1 H, d), 7.29 (1 H, s), 6.38 (1 H, s), 4.62 (2 H, s), 4.54 (2 H, d), 4.48 - 4.36 (4 H, m), 4.21 (2 H, t), 3.99 (1 H, ddt), 3.90 - 3.61 (5 H, m), 3.58 (2 H, s), 3.49 (1 H, dd), 3.02 (2 H, t), 1.36 (3 H, s)
126	(^1H , CDCl_3) δ ppm 7.50 (1 H, d), 6.63 (1 H, dd), 6.48 (1 H, s), 6.21 (1 H, s), 4.47 - 4.34 (2 H, m), 4.21 - 4.14 (2 H, m), 4.01 - 3.91 (1 H, m), 3.89 - 3.60 (5 H, m), 3.48 (1 H, dd), 2.98 (2 H, s), 2.90 (2 H, t), 1.36 (2 H, dd), 0.96 (6 H, s), 0.87 (3 H, t)
127	(^1H , CDCl_3) δ ppm 7.61 (1 H, d), 7.22 (1 H, d), 7.14 (1 H, s), 6.34 (1 H, d), 4.46 - 4.34 (2 H, m), 4.19 (2 H, t), 3.97 (1 H, td), 3.89 - 3.60 (5 H, m), 3.48 (1 H, t), 3.42 - 3.35 (1 H, m), 2.97 (2 H, t), 2.93 - 2.85 (2 H, m), 2.70 (1 H, m), 1.84 - 1.63 (3 H, m), 0.92 (6 H, dd)
128	(^1H , CDCl_3) δ ppm 7.51 (1 H, d), 6.63 (1 H, d), 6.49 (1 H, br. s.), 6.21 (1 H, s), 4.46 - 4.356 (2 H, m), 4.21 - 4.11 (2 H, m), 3.97 (1 H, ddt), 3.88 - 3.62 (5 H, m), 3.48 (1 H, dd), 3.10 (2 H, d), 2.91 (2 H, t), 1.61 (1 H, m), 1.47 - 1.26 (8 H, m), 0.97 - 0.88 (6 H, m)

Cpd#	NMR data (δ)
129	(^1H , CDCl_3) δ ppm 7.63 (1 H, d), 6.93 (1 H, dd), 6.82 (1 H, d), 6.29 (1 H, s), 5.30 (1 H, s), 4.47 - 4.36 (2 H, m), 4.22 - 4.14 (4 H, m), 4.03 - 3.94 (1 H, m), 3.88 - 3.628 (7 H, m), 3.53 - 3.453 (4 H, m), 2.97 (2 H, t)
130	(^1H , CDCl_3) δ ppm 7.64 - 7.58 (1 H, m), 6.90 (1 H, dd), 6.80 (1 H, d), 6.29 - 6.24 (1 H, m), 4.46 - 4.32 (2 H, m), 4.22 - 4.12 (4 H, m), 4.01 - 3.90 (1 H, m), 3.88 - 3.54 (9 H, m), 3.46 (1 H, dd), 2.95 (2 H, s), 1.23 (3 H, t)
131	(^1H , CDCl_3) δ ppm 7.60 - 7.67 (1 H, m), 6.93 - 6.86 (1 H, m), 6.78 (1 H, d), 6.29 (1 H, s), 4.50 - 4.36 (2 H, m), 4.25 - 4.17 (2 H, m), 4.04 - 3.94 (1 H, m), 3.91 - 3.61 (7 H, m), 3.53 - 3.44 (1 H, m), 3.01 - 2.93 (2 H, m), 1.37 - 1.227 (1 H, m), 0.73 - 0.65 (2 H, m), 0.42 - 0.34 (2 H, m)
132	(^1H , CDCl_3) δ ppm 7.64 (1 H, d), 6.93 (1 H, s), 6.82 (1 H, d), 6.27 (1 H, s), 4.90 - 4.82 (1 H, m), 4.74 - 4.67 (1 H, m), 4.48 - 4.28 (3 H, m), 4.26 - 4.14 (3 H, m), 4.03 - 3.91 (1 H, m), 3.90 - 3.58 (5 H, m), 3.53 - 3.41 (1 H, m), 3.03 - 2.91 (2 H, m)
133	(^1H , CDCl_3) δ ppm 7.64 (1 H, s), 7.48 - 7.40 (1 H, m), 7.40 - 7.35 (1 H, m), 6.40 - 6.34 (1 H, m), 4.47 (4 H, s), 4.28 - 4.15 (1 H, m), 3.93 - 3.56 (10 H, m), 3.43 (5 H, s), 3.06 - 2.95 (2 H, m)
134	(^1H , CDCl_3) δ ppm 7.65 (1 H, d), 7.47 - 7.41 (1 H, m), 7.38 (1 H, s), 6.37 (1 H, s), 4.37 - 4.49 (4 H, m), 4.26 - 4.17 (2 H, m), 3.99 (1 H, m), 3.93 - 3.44 (12 H, m), 3.05 - 2.94 (2 H, m), 1.25 (3 H, t)
135	(^1H , CDCl_3) δ ppm 7.66 (1 H, d), 7.47 - 7.42 (1 H, m), 7.40 (1 H, s), 6.37 (1 H, s), 4.73 - 4.67 (1 H, m), 4.62 - 4.54 (1 H, m), 4.49 (4 H, s), 4.26 - 4.16 (2 H, m), 4.04 - 3.95 (1 H, m), 3.94 - 3.60 (7 H, m), 3.56 - 3.43 (1 H, m), 3.05 - 2.96 (2 H, m)
136	(^1H , CDCl_3) δ ppm 7.68 (1 H, d), 7.35 (1 H, d), 7.30 (1 H, s), 6.38 (1 H, s), 4.57 (2 H, s), 4.50 - 4.36 (2 H, m), 4.27 - 4.17 (2 H, m), 4.06 - 3.92 (1 H, m), 3.92 - 3.60 (5 H, m), 3.57 - 3.44 (1 H, m), 3.17 (2 H, s), 3.09 - 2.96 (2 H, m), 0.97 (9 H, s)
137	(^1H , CDCl_3) δ ppm 7.70 - 7.64 (1 H, m), 7.40 - 7.33 (1 H, m), 7.31 (1 H, s), 6.37 (1 H, s), 4.59 (2 H, s), 4.50 - 4.35 (2 H, m), 4.27 - 4.16 (2 H, m), 4.07 - 3.94 (1 H, m), 3.93 - 3.59 (5 H, m), 3.57 - 3.44 (1 H, m), 3.45 - 3.34 (1 H, m), 3.08 - 2.96 (2 H, m), 2.05 - 1.90 (2 H, m), 1.82 - 1.71 (2 H, m), 1.48 - 1.18 (6 H, m)
138	(^1H , CDCl_3) δ ppm 7.68 (1 H, d), 7.37 (1 H, d), 7.32 (1 H, s), 6.38 (1 H, s), 4.58 (2 H, s), 4.51 - 4.36 (2 H, m), 4.26 - 4.176 (2 H, m), 4.05 - 3.94 (1 H, m), 3.92 - 3.61 (5 H, m), 3.54 - 3.44 (1 H, m), 3.38 (2 H, d), 3.07 - 2.98 (2 H, m), 1.20 - 1.05 (1 H, m), 0.63 - 0.54 (2 H, m), 0.29 - 0.20 (2 H, m)

Cpd#	NMR data (δ)
139	(^1H , CDCl_3) δ ppm 7.62 (1 H, s), 6.91 (1 H, m), 6.81 (1 H, m), 6.27 (1 H, s), 4.41 (2 H, s), 4.25 - 4.21 (2 H, m), 4.10 - 3.41 (12 H, m), 2.96 (2 H, t), 2.01 - 1.84 (1 H, m), 1.76 - 1.36 (5 H, m)
140	(^1H , $\text{DMSO}-d_6$) δ ppm 7.91 (1 H, d), 7.28 - 7.18 (2 H, m), 6.60 (1 H, s), 4.32 - 4.17 (2 H, m), 4.07 - 3.96 (2 H, m), 3.92 - 3.71 (3 H, m), 3.70 - 3.43 (4 H, m), 3.38 (2 H, m), 3.03 - 2.91 (2 H, m), 2.78 - 2.58 (2 H, m), 1.68 - 1.58 (2 H, m), 1.08 (3 H, d)
141	(^1H , CDCl_3) δ ppm 7.70 - 7.58 (1 H, d), 6.95 - 6.82 (1 H, d), 6.77 (1 H, s), 6.27 (1 H, s), 4.50 - 4.32 (2 H, m), 4.38 - 4.15 (2 H, m), 4.05 - 3.92 (3 H, m), 3.92 - 3.60 (5 H, m), 3.55 - 3.42 (1 H, t), 3.05 - 2.92 (2 H, m), 1.85 - 1.70 (2 H, m), 1.40 - 1.30 (2 H, m), 0.92 (9 H, s)
142	(^1H , CDCl_3) δ ppm 7.61 (1 H, d), 7.21 (1 H, d), 7.13 (1 H, s), 6.35 (1 H, s), 4.46 - 4.40 (2 H, m), 4.20 (2 H, t), 3.99 - 3.97 (1 H, m), 3.87 - 3.65 (5 H, m), 3.49 (1 H, t), 3.39 (3 H, s), 3.00 - 2.93 (3 H, m), 2.83 - 2.82 (1 H, m), 2.66 - 2.64 (1 H, m), 1.95 - 1.91 (1 H, m), 1.75 - 1.73 (2 H, m), 0.90 (6 H, t)
143	(^1H , CDCl_3) δ ppm 7.66 - 7.60 (1 H, m), 6.91 - 6.86 (1 H, m), 6.80 - 6.76 (1 H, m), 6.28 (1 H, s), 4.48 - 4.35 (2 H, m), 4.24 - 4.17 (2 H, m), 4.11 - 4.04 (2 H, m), 4.02 - 3.95 (1 H, m), 3.91 - 3.61 (5 H, m), 3.54 - 3.45 (1 H, m), 3.03 - 2.94 (2 H, m), 1.98 - 1.88 (2 H, m), 1.45 - 1.36 (2 H, m), 0.78 - 0.66 (1 H, m), 0.50 - 0.43 (2 H, m), 0.09 - 0.03 (2 H, m)
145	(^1H , CDCl_3) δ ppm 7.47 (1 H, d), 6.55 - 6.48 (1 H, m), 6.40 - 6.34 (1 H, m), 6.18 (1 H, s), 4.47 - 4.33 (2 H, m), 4.22 - 4.12 (2 H, m), 4.02 - 3.92 (1 H, m), 3.83 (5 H, dd), 3.53 - 3.43 (1 H, m), 3.40 - 3.28 (1 H, m), 2.89 (2 H, t), 2.13 - 1.98 (2 H, m), 1.86 - 1.73 (2 H, m), 1.73 - 1.63 (1 H, m), 1.50 - 1.11 (5 H, m)
146	(^1H , $\text{DMSO}-d_6$) δ ppm 7.95 - 7.87 (1 H, m), 7.29 - 7.19 (2 H, m), 6.60 (1 H, s), 4.54 - 4.44 (1 H, m), 4.29 - 4.21 (2 H, m), 4.07 - 3.97 (2 H, m), 3.91 - 3.72 (3 H, m), 3.70 - 3.54 (2 H, m), 3.54 - 3.43 (1 H, m), 3.43 - 3.34 (1 H, m), 3.06 - 2.81 (4 H, m), 2.63 - 2.52 (1 H, m), 1.81 - 1.64 (1 H, m), 1.52 - 1.36 (1 H, m), 0.81 (9 H, s)
147	(^1H , CDCl_3) δ ppm 7.68 (1 H, d), 7.38 - 7.32 (1 H, m), 7.30 (1 H, s), 6.38 (1 H, s), 4.56 (2 H, s), 4.50 - 4.36 (2 H, m), 4.26 - 4.18 (2 H, m), 4.04 - 3.94 (1 H, m), 3.94 - 3.5994 (4 H, m), 3.55 - 3.44 (1 H, m), 3.40 (2 H, d), 3.07 - 2.98 (2 H, m), 2.30 - 2.16 (1 H, m), 1.86 - 1.72 (2 H, m), 1.66 - 1.51 (5 H, m), 1.35 - 1.20 (2 H, m)
148	(^1H , CDCl_3) δ ppm 7.61 (1 H, d), 7.18 (1 H, dd), 7.12 (1 H, d), 6.35 (1 H, s), 4.46 - 4.36 (2 H, m), 4.21 (2 H, t), 4.00 - 3.97 (1 H, m), 3.88 - 3.64 (5 H, m), 3.49 (1 H, t), 3.34 - 3.30 (4 H, m), 2.98 (2 H, t), 2.81 - 2.60 (2 H, m), 1.89 - 1.676 (2 H, m), 1.18 (3 H, d)

Cpd#	NMR data (δ)
149	(^1H , CDCl_3) δ ppm: 7.62 (1 H, d), 7.23 - 7.13 (4 H, m), 6.71 (1 H, t), 6.58 (2 H, d), 6.12 (1 H, s), 4.47 - 4.37 (2 H, m), 4.21 (2 H, t), 4.08 - 3.98 (1 H, m), 3.88 - 3.63 (6 H, m), 3.49 (1 H, t), 3.17 (2 H, t), 2.98 (2 H, t), 2.79 (2 H, t), 2.01 - 1.94 (2 H, m)
150	(^1H , CDCl_3) δ ppm 7.61 (1 H, d), 7.20 (1 H, d), 7.12 (1 H, s), 6.35 (1 H, s), 4.47 - 4.37 (2 H, m), 4.21 (2 H, t), 4.01 - 3.97 (1 H, m), 3.88 - 3.63 (6 H, m), 3.49 (1 H, t), 2.98 (2 H, t), 2.69 (2 H, t), 1.89 - 1.62 (2 H, m), 1.52 - 1.45 (2 H, m), 1.31 - 1.30 (1 H, m), 1.20 (3 H, d)
151	(^1H , CDCl_3) δ ppm 7.61 (1 H, d), 7.20 (1 H, d), 7.12 (1 H, s), 6.35 (1 H, s), 4.47 - 4.37 (2 H, m), 4.21 (2 H, t), 4.01 - 3.97 (1 H, m), 3.88 - 3.63 (7 H, m), 3.49 (1 H, t), 2.98 (2 H, t), 2.70 (2 H, t), 1.76 - 1.60 (4 H, m), 1.31 - 1.26 (1 H, m)
152	(^1H , CDCl_3) δ ppm 7.57 - 7.50 (1 H, m), 6.71 (1 H, d), 6.53 (1 H, br. s.), 6.21 (1 H, s), 4.513 - 4.33 (2 H, m), 4.26 - 4.14 (2 H, m), 4.06 - 3.93 (1 H, m), 3.92 - 3.56 (6 H, m), 3.56 - 3.41 (1 H, m), 3.00 - 2.90 (2 H, m), 2.88 (3 H, s), 2.04 - 1.63 (2 H, m), 1.60 - 1.31 (5 H, m), 1.28 - 1.08 (1 H, m)
153	(^1H , CDCl_3) δ ppm 7.52 - 7.46 (1 H, m), 6.55 - 6.50 (1 H, m), 6.40 - 6.36 (1 H, m), 6.19 (1 H, s), 4.48 - 4.33 (2 H, m), 4.23 - 4.13 (2 H, m), 4.04 - 3.92 (1 H, m), 3.85 (5 H, d), 3.56 - 3.42 (1 H, m), 3.06 - 2.99 (2 H, m), 2.93 - 2.84 (2 H, m), 1.92 - 1.50 (6 H, m), 1.36 - 1.12 (3 H, m), 1.08 - 0.91 (2 H, m)
154	(^1H , CDCl_3) δ ppm 7.50 (1 H, d), 6.59 - 6.49 (1 H, m), 6.44 - 6.35 (1 H, m), 6.20 (1 H, s), 4.49 - 4.33 (2 H, m), 4.24 - 4.13 (2 H, m), 4.05 - 3.93 (3 H, m), 3.92 - 3.59 (5 H, m), 3.56 - 3.32 (3 H, m), 3.16 - 3.07 (2 H, m), 2.96 - 2.84 (2 H, m), 1.98 - 1.80 (1 H, m), 1.79 - 1.64 (2 H, m), 1.428 - 1.28 (2 H, m)
155	(^1H , CDCl_3) δ ppm 7.54 - 7.46 (1 H, m), 7.14 - 7.08 (1 H, m), 7.04 (1 H, s), 6.23 (1 H, s), 4.35 - 4.21 (2 H, m), 4.13 - 3.99 (2 H, m), 3.91 - 3.80 (1 H, m), 3.79 - 3.45 (5 H, m), 3.41 - 3.30 (1 H, m), 2.86 (2 H, t), 2.64 - 2.54 (2 H, m), 1.67 - 1.58 (2 H, m), 1.45 (4 H, d), 0.80 (6 H, m)
156	(^1H , CDCl_3) δ ppm 7.60 (1 H, d), 7.23 - 7.19 (1 H, m), 7.13 (1 H, s), 6.34 (1 H, s), 4.45 - 4.35 (2 H, m), 4.18 (2 H, t), 4.02 - 3.92 (1 H, m), 3.82 (5 H, m), 3.48 (1 H, dd), 2.97 (2 H, br. t), 2.79 - 2.70 (2 H, m), 1.82 - 1.74 (2 H, m), 1.30 (6 H, s)
157	(^1H , CDCl_3) δ ppm 7.64 - 7.59 (1 H, m), 7.25 - 7.20 (1 H, m), 7.14 (1 H, s), 6.35 (1 H, s), 4.48 - 4.35 (2 H, m), 4.24 - 4.16 (2 H, m), 4.02 - 3.94 (1 H, m), 3.90 - 3.44 (7 H, m), 3.02 - 2.95 (2 H, m), 2.92 - 2.81 (1 H, m), 2.78 - 2.67 (1 H, m), 1.87 - 1.68 (2 H, m), 1.62 - 1.42 (2 H, m), 0.96 (3 H, s)

Cpd#	NMR data (δ)
158	(^1H , CDCl_3) δ ppm 7.70 - 7.56 (1 H, d), 6.95 - 6.85 (1 H, d), 6.80 (1 H, s), 6.28 (1 H, s), 4.50 - 4.35 (2 H, m), 4.25 - 4.10 (2 H, m), 4.05 - 3.92 (1 H, m), 3.10 - 3.57 (8 H, m), 3.55 - 3.40 (1 H, m), 3.05 - 2.90 (2 H, m), 1.05 (9H, s)
159	(^1H , CDCl_3) δ ppm 7.63 (1 H, d), 6.88 (1 H, dd), 6.77 (1H, d), 6.28 (1 H, s), 4.46 - 4.36 (2 H, m), 4.20 (2 H, t), 4.05 - 3.96 (3 H, m), 3.88 - 3.65 (7 H, m), 3.51 - 3.43 (3 H, m), 2.97 (2 H, t), 2.12 - 2.01 (1 H, m), 1.78 - 1.75 (2 H, m), 1.59 - 1.42 (2 H, m)
160	(^1H , CDCl_3) δ ppm 7.63 (1 H, d), 7.35 - 7.18 (1 H, m), 7.12 (1H, s), 6.36 (1 H, s), 4.50 - 4.35 (2 H, m), 4.28 - 4.15 (2 H, m), 4.05 - 3.95 (1 H, m), 3.95 - 3.60 (5 H, m), 3.55 - 3.42 (1 H, m), 3.05 - 2.90 (2 H, m), 2.75 - 2.60 (2 H, m), 1.80 - 1.65 (2 H, m), 1.55 - 1.48 (2 H, m), 1.22 (6 H, s)
161	(^1H , CDCl_3) δ ppm 7.68 (1 H, d), 7.34 (1 H, d), 7.29 (1 H, s), 6.38 (1 H, s), 4.55 (2 H, s), 4.48 - 4.38 (2 H, m), 4.22 (2 H, t), 4.02 - 3.96 (3 H, m), 3.91 - 3.61 (5 H, m), 3.55 - 3.35 (5 H, m), 3.02 (2 H, t), 2.00 - 1.86 (1 H, m), 1.69 (2 H, dd), 1.38 (2 H, dd)
162	(^1H , CDCl_3) δ ppm 7.68 - 7.62 (1 H, m), 6.95 - 6.87 (1 H, m), 6.82 - 6.77 (1 H, m), 6.29 (1 H, s), 4.51 - 4.35 (2 H, m), 4.26 - 4.17 (2 H, m), 4.05 - 3.92 (1 H, m), 3.91 - 3.60 (8 H, m), 3.55 - 3.44 (1 H, m), 3.04 - 2.94 (2 H, m)
163	(^1H , CDCl_3) δ ppm 7.64 (1 H, d), 6.91 (1 H, dd), 6.81 (1 H, d), 6.29 (1H, s), 4.91 (2 H, t), 4.56 (2 H, t), 4.47 - 4.36 (2 H, m), 4.26 (2H, d), 4.21 (2 H, t), 4.00 - 3.96 (1 H, m), 3.88 - 3.63 (5 H, m), 3.51 - 3.45 (2 H, m), 2.98 (2H, t)
164	(^1H , $\text{DMSO}-d_6$) δ ppm 7.93 (1 H, d), 6.97 - 6.92 (2 H, m), 6.53 (1 H, s), 4.24 - 4.23 (2H, m), 4.08 (2 H, t), 4.00 (2 H, t), 3.98 - 3.74 (3 H, m), 3.68 - 3.57 (2 H, m), 3.51 - 3.48 (1 H, m), 3.37 (1 H, t), 2.96 (2 H, t), 1.84 - 1.80 (2 H, m), 1.36 - 1.30 (2 H, m), 0.81 - 0.63 (1 H, m), 0.42 - 0.39 (2 H, m), 0.04 - 0.02 (2 H, m)
165	(^1H , CDCl_3) δ ppm 7.65 (1 H, d), 7.24 (1H, dd), 7.16 (1H, s), 6.38 (1 H, s), 4.49 - 4.40 (2 H, m), 4.24 (2 H, t), 4.02 - 4.00 (1 H, m), 3.89 - 3.66 (3 H, m), 3.52 (3 H, m), 3.43 (2 H, t), 3.38 (3 H, s), 3.02 (2H, t), 2.78 (2 H, t), 1.96 - 1.92 (2 H, m)
166	(^1H , CDCl_3) δ ppm 7.61 (1 H, d), 7.23 (1H, dd), 7.15 (1H, s), 6.35 (1 H, s), 4.46 - 4.36 (2 H, m), 4.20 (2 H, t), 4.01 - 3.95 (1 H, m), 3.88 - 3.62 (5 H, m), 3.49 (1 H, t), 2.98 (2 H, t), 2.85 - 2.81 (2 H, m), 2.13 (1H, br s), 1.93 - 1.79 (4 H, m), 1.72 - 1.55 (6 H, m)
167	(^1H , CDCl_3) δ ppm 7.59 (1 H, d), 7.37 (1H, dd), 7.15 (1H, s), 6.34 (1 H, s), 4.44 - 4.35 (2 H, m), 4.17 (2 H, t), 4.01 - 3.92 (3 H, m), 3.88 - 3.60 (7 H, m), 3.47 (1 H, t), 2.94 (2 H, t), 2.07 - 2.02 (2 H, m), 1.95-1.88 (2H, m)

Cpd#	NMR data (δ)
168	(^1H , CDCl_3) δ ppm 7.66 (1 H, d), 7.23 (1H, dd), 7.17 (1H, s), 6.39 (1 H, s), 4.51 - 4.40 (2 H, m), 4.24 (2 H, t), 4.09 - 3.96 (1 H, m), 3.93- 3.65 (5 H, m), 3.57-3.49 (1H, m), 3.44 (2 H, t), 3.40 (3H, s), 3.02 (2 H, t), 2.79 (2 H, t), 2.00-1.89 (2H, m)
169	(^1H , CDCl_3) δ ppm 7.55 (1 H, d), 7.16 (1H, dd), 7.08 (1H, s), 6.29 (1 H, s), 4.41 - 4.29 (2 H, m), 4.13 (2 H, t), 3.95 - 3.86 (1 H, m), 3.81- 3.54 (5 H, m), 3.46-3.38 (1H, m), 2.91 (2 H, t), 2.78-2.72 (2H, m), 1.87-1.75 (4 H, m), 1.72-1.53 (6 H, m)
170	(^1H , CDCl_3) δ ppm 7.66 (1 H, d), 6.95 (1H, dd), 6.84 (1H, s), 6.31 (1 H, s), 4.50 - 4.39 (2 H, m), 4.25-4.21 (4 H, m), 4.05 - 3.98 (1 H, m), 3.91- 3.65 (7 H, m), 3.55-3.49 (3H, m), 3.00 (2 H, t), 1.71-1.62 (2 H, m), 0.96 (3 H, t)
171	(^1H , CDCl_3) δ ppm 7.65 (1 H, d), 6.94 (1H, dd), 6.83 (1H, s), 6.31 (1 H, s), 4.50 - 4.39 (2 H, m), 4.25-4.18 (4 H, m), 4.04 - 3.97 (1 H, m), 3.91- 3.65 (8 H, m), 3.51 (1H, t), 3.00 (2 H, t), 1.23 (6 H, d)
172	(^1H , CDCl_3) δ ppm 7.67 (1 H, d), 6.95 (1H, dd), 6.85 (1H, s), 6.31 (1 H, s), 4.50 - 4.40 (2 H, m), 4.25-4.21 (4 H, m), 4.05 - 3.98 (1 H, m), 3.91- 3.65 (7 H, m), 3.55-3.49 (3H, m), 3.00 (2 H, t), 1.72-1.62 (2 H, m), 0.97 (3 H, t)
173	(^1H , CDCl_3) δ ppm 7.65 (1 H, d), 6.94 (1H, dd), 6.84 (1H, s), 6.31 (1 H, s), 4.50 - 4.40 (2 H, m), 4.26-4.19 (4 H, m), 4.04 - 3.97 (1 H, m), 3.92- 3.65 (8 H, m), 3.52 (1H, t), 3.00 (2 H, t), 1.23 (6 H, d)
174	(^1H , CDCl_3) δ ppm 7.64 (1 H, d), 7.22 (1H, dd), 7.15 (1H, s), 6.38 (1 H, s), 4.51 - 4.41 (2 H, m), 4.23 (2 H, t), 4.06 - 4.00 (1 H, m), 3.92- 3.66 (5 H, m), 3.57-3.50 (1H, m), 3.43 (2 H, t), 3.37 (3H, s), 3.01 (2 H, t), 2.71 (2 H, t), 1.80-1.61 (2H, m), 1.69-1.62 (2H,m)

Biological Examples

1. *In vitro* assays

1.1. *Cell based assay: GTP- γ S binding assay.*

5 [00458] The following assay can be used for determination of GPR84 activation. The [^{35}S]GTP γ S binding assay measures the level of G protein activation following agonist occupation of a GPCR, by determining the binding of the non-hydrolysable analog [^{35}S]GTP γ S to G α subunits.

[00459] The assay is performed in a 96 well plate where the following reagents are added. First 50 μL compound is added into the assay plate, followed by addition of 20 μL 3,3' di indolylmethane at EC_{80} concentration (concentration which gives 80% of the activity of GPR84). In a last step, 30 μL of a mixture consisting of membranes-GTP γ S-SpA beads is added [mixture consists of 20 μg /well membranes derived from stable cell line over expressing GPR84 (membranes are pre-incubated with 0.1 μM GDP for 15 min at

10

4°C), 0.1 nM [³⁵S]GTPγS (Perkin Elmer, NEG030) and 0.5 mg/well PVT-WGA SpA beads (Perkin Elmer, RPNQ0001)]. All components are diluted in assay buffer containing 20mM HEPES pH 7.4; 5mM MgCl₂; 250mM NaCl; 0.05% BSA; 75ug/mL saponin. Reactions are incubated for 90 min at room temperature followed by centrifugation at 2000 rpm during 15 min. Plates are read on a Topcount reader (Perkin Elmer) immediately after centrifugation (readout time, 1 min/well).

[00460] TABLE IV: GPR84 assay GTPγS IC₅₀ (nM) of selected Compounds of the invention.

na: not active

* > 1001 nM

** 501-1000 nM

*** 101-500 nM

**** 0.01- 100 nM

Cpd#	GPR84
1	***
3	**
4	***
5	**
6	***
7	***
8	***
9	****
16	***
20	****
28	**
30	****
36	****
41	***
42	***
43	***
45	**
50	***
51	***
52	****
53	***
54	***
55	*
56	***
57	***
58	***
59	**
60	****
61	****
62	****

Cpd#	GPR84
63	***
64	***
65	**
66	***
67	****
68	****
69	****
70	****
72	****
74	****
75	****
76	****
77	****
78	****
79	***
80	****
81	****
82	****
83	****
84	****
85	****
86	****
87	***
89	***
90	****
91	***
93	****
94	****
95	**
96	***

Cpd#	GPR84
97	***
98	****
101	***
102	***
103	***
104	****
105	***
106	**
107	****
108	****
109	****
110	***
111	***
112	****
113	***
114	****
115	****
116	***
117	***
118	***
119	***
120	***
121	****
122	****
123	****
124	***
125	**
126	****
127	****
128	***
129	***
130	***
131	****
132	**
133	***
134	***
135	***
136	****

Cpd#	GPR84
137	****
138	****
139	****
140	***
141	***
142	****
145	***
146	****
147	****
148	****
149	****
150	***
151	***
152	**
153	***
154	**
155	****
156	***
157	***
158	***
159	****
160	***
161	***
162	***
163	****
164	****
165	****
166	****
167	**
168	***
169	***
170	****
171	****
172	***
173	**
174	****

2. Cellular assays

2.1. Human neutrophil migration assay

- 5 [00461] We have established that GPR84 agonists (MCFA such as sodiumdecanoate, 3,3' di indolylmethane and Embelin induce neutrophil chemo taxis and that GPR84 antagonists could block GPR84

agonist-induced chemo taxis but not IL8-induced chemotaxis, indicating that G Protein-Coupled Receptor 84 (GPR84) is an essential player in the process of neutrophil recruitment.

[00462] The effect of agonists or antagonists for GPR84 can therefore be assayed in a neutrophil migration test. In the neutrophil migration assay, neutrophils, freshly isolated from buffy coats from human volunteers, are treated with a compound for 30 minutes. Subsequently, the neutrophils are transferred to the upper wells of a Corning HTS transwell 96 permeable support system, of which the lower wells are filled with a embelin solution at EC₈₀ (concentration which gives 80% of the activity of GPR84). After 1 h of incubation, migration of the neutrophils towards embelin in the lower compartment can be quantified by measuring the ATP-content of the lower wells using the ATPlite luminescence ATP detection assay system (Perkin Elmer, Cat. N°.: 436110).

2.1.1 Isolation of neutrophils from human buffy coat

[00463] A human buffy coat is diluted with an equal volume of ice cold DPBS. 20 mL of the diluted buffy coat is gently mixed with 4 mL of ACD buffer (140 mM citric acid, 200 mM sodium citrate and 220 mM dextrose). Then, 12 mL of the 6% dextran/0.9% NaCl solution (15 g dextran T2000 and 2.25 g NaCl dissolved in 250 mL H₂O) is added to the mixture and the samples are inverted gently up to 20 times. The total volume was transferred to a new recipient and incubated at room temperature for 1 h for complete separation of the two phases to occur. The yellowish supernatant is then transferred to a clean centrifugation tube and centrifuged for 12 minutes at 1300 rpm and 4°C. After centrifugation, the supernatant is discarded and the remaining cell pellet is rapidly resuspended in 12 mL of ice-cold H₂O for red blood cell lysis to occur. After 20 seconds, 4 mL of ice-cold 0.6 M KCl is added. Samples are mixed carefully and centrifuged for 6 minutes at 1300 rpm, 4°C. The supernatant is discarded and the red blood cell lysis procedure is repeated one more time. Subsequently, the cell pellet is resuspended in 4 mL of DPBS and layered over 5 mL of Lymphoprep (Nycomed Pharma, Cat. N°.: 1114545) in a 15 mL centrifuge tube. After centrifugation for 12 min at 1300 rpm, 4°C, the supernatant is removed and the cell pellet, containing the neutrophils, is resuspended in 25 mL chemotaxis buffer (RPMI 1640 medium, supplemented with 10 mM HEPES; freshly made for each experiment)

2.1.2 Migration assay

[00464] A cell suspension of 8.9×10^6 cells per milliliter is prepared. 20 µL of compound solution in chemotaxis buffer is added to 180 µL cell suspension. The mixture is incubated at 37°C for 30 minutes with intermediate resuspension of the cells after 15 minutes. Following this, 70 µL cell suspension is transferred to the upper compartment of a Corning HTS transwell 96 permeable support system with 5.0 µm pore size polycarbonate membrane (Corning, Cat.N°.: 3387). The receiver well of the transwell system is then filled with 200 µL chemotaxis buffer containing compound and chemotactic agent (embelin). After incubation at

37°C in 5% CO₂ for 1 h, the upper plate of the transwell system is removed and the cell suspension in the receiver plate is transferred to a 96-well V-bottom plate. 50 µL of DPBS is added to the receiver plate to prevent remaining cells from drying out. The V-bottom plate is centrifuged for 6 minutes at 1500 rpm. The supernatant is removed and the cells are resuspended in 50 µL DPBS. The cells are then transferred back to the receiver plate of the transwell system. After this, 100 µL ATPlite solution (Perkin Elmer, Cat. N°: 436110) was added to the cells. The plate is incubated for 10 minutes in the dark, while shaking. 170 µL of cell lysate is then transferred to a white 96-well plate and luminescence is measured. The detected luminescent signal is considered as linearly related to the number of cells having migrated from the upper well to the receiver well.

10

[00465]

TABLE VII: human neutrophil migration inhibition

* > 1001 nM

** 501-1000 nM

*** 101-500 nM

**** 0.01-100 nM

Cpd#	Neutrophils
4	***
7	***
8	***
9	****
16	***
17	****
19	****
20	****
22	****
23	***
30	****
34	****
35	***
36	***
41	****
42	***
52	****
56	***
60	****
62	****
63	****
68	****
69	****
72	****
77	****

Cpd#	Neutrophils
80	****
83	****
85	****
89	****
90	****
92	****
98	****
107	****
109	****
111	****
112	****
114	****
115	****
116	****
121	***
122	****
123	****
126	****
133	****
139	****
140	****
147	****
149	****
150	****
158	****

Cpd#	Neutrophils
159	****
160	***
161	****
168	****

Cpd#	Neutrophils
169	****
170	****
171	****
174	****

2.2. *Rat neutrophil migration assay*

[00466] We have established that GPR84 agonists (MCFA such as sodiumdecanoate, 3,3' diindolylmethane and Embelin induce neutrophil chemotaxis and that GPR84 antagonists could block GPR84 agonist-induced chemotaxis but not IL8-induced chemotaxis, indicating that G Protein-Coupled Receptor 84 (GPR84) is an essential player in the process of neutrophil recruitment.

[00467] The effect of agonists or antagonists for GPR84 can therefore be assayed in a neutrophil migration test. In the rat neutrophil migration assay, neutrophils, freshly isolated from rat after intraperitoneal injection of glycogen (0.1 %, w/v), are treated with a compound for 30 minutes. Subsequently, the neutrophils are transferred to the upper wells of a Corning HTS transwell 96 permeable support system, of which the lower wells are filled with a embelin solution at EC80 (concentration which give 80% of the activity of the GPR84). After 1 h of incubation, migration of the neutrophils towards embelin in the lower compartment can be quantified by measuring the ATP-content of the lower wells using the Cell Titer Glow Substrate assay system (Promega, Cat.N°.: G755B).

2.2.1. *Isolation of neutrophils from rats*

[00468] 24 h after intraperitoneal injection of glycogen (0.1 %, w/v), cells are harvested by peritoneal lavage with 25mL HBSS then centrifuged for 12 minutes at 1300 rpm and 4°C. After centrifugation, the supernatant is discarded and the remaining cell pellet is rapidly resuspended in 12 mL of ice-cold H₂O for red blood cell lysis to occur. After 20 seconds, 4 mL of ice-cold 0.6 M KCl is added. Samples are mixed carefully and centrifuged for 6 minutes at 1300 rpm, 4°C. The supernatant is discarded and the cell pellet is resuspended in 4 mL of DPBS and layered over 5 mL of Lymphoprep (Axis Shield, Cat. N°.: 1114544) in a 15 mL centrifuge tube. After centrifugation for 30 min at 1500 rpm, 4°C, the supernatant is removed and the cell pellet, containing the neutrophils, is resuspended in 5 mL chemotaxis buffer (RPMI 1640 medium, supplemented with 10 mM HEPES; freshly made for each experiment).

2.2.2. *Migration assay*

[00469] A cell suspension of 8.9x10⁶ cells per milliliter is prepared. 10 µL of compound solution in chemotaxis buffer is added to 90 µL cell suspension. The mixture is incubated at 37°C for 30 minutes with intermediate resuspension of the cells after 15 minutes. Following this, 75 µL cell suspension is transferred to the upper compartment of a Corning HTS transwell 96 permeable support system with 5.0 µm pore size

polycarbonate membrane (Corning, Cat.N°.: 3387). The receiver well of the transwell system is then filled with 200 μ L chemotaxis buffer containing compound and chemotactic agent (embelin). After incubation at 37°C in 5% CO₂ for 1 h, the upper plate of the transwell system is removed and 70 μ L Cell Titer Glow Substrate (Promega, Cat.N°.: G755B) are added in the receiver plate. The receiver plate is incubated for 10 minutes in the dark, while shaking. 180 μ L of cell lysate is then transferred to a white 96-well plate and luminescence is measured. The detected luminescent signal is considered as linearly related to the number of cells having migrated from the upper well to the receiver well.

3. ADME, PK and Safety Models

3.1 Aqueous Solubility

[00470] Starting from a 10mM stock in DMSO, a serial dilution of the compound is prepared in DMSO. The dilution series is transferred to a 96 NUNC Maxisorb plate F-bottom and 0.1M phosphate buffer pH 7.4 or 0.1M citrate buffer pH3.0 at room temperature is added.

[00471] The final concentrations range from 18.75 to 300 μ M in 5 equal dilution steps. The final DMSO concentration does not exceed 3%.

[00472] 200 μ M Pyrene is added to the corner points of each 96 well plate and serves as a reference point for calibration of Z-axis on the microscope.

[00473] The assay plates are sealed and incubated for 1 h at 37°C while shaking at 230rpm. The plates are then scanned under a white light microscope, yielding individual pictures of the precipitate per concentration.

The precipitate is analyzed and converted into a number by a custom-developed software tool. The first concentration at which the compound appears completely dissolved is the concentration reported, however the true concentration lies somewhere between this concentration and one dilution step higher.

[00474] Solubility values are reported in μ M and in μ g/mL.

3.2 Thermodynamic solubility

[00475] Two individual solutions of 2 mg/mL of compound are prepared in a 0.1 M phosphate buffer pH 7.4 or a 0.1 M citrate buffer pH 3.0 at room temperature in a 2mL glass vial.

[00476] After addition of a magnetic stir, the samples are stirred at room temperature for 24 h.

[00477] After 24 h, the vials are centrifuged 10 min at 1400 rpm. The supernatant of the sample is then transferred to a MultiscreenR Solubility Plate (Millipore, MSSLBPC50) and filtered (10-12" Hg) with the aid of a vacuum manifold into a clean Greiner polypropylene V-bottom 96 well plate. Per sample, two dilutions (factor 10 and 100) are made in DMSO. Other dilutions can be made if the acquired peak area is not within the standard curve.

[00478] A 10mM DMSO stock, made from dry matter, is used to make a 200 μ g/mL working stock. The standard curve for the compound is prepared in DMSO starting from the 200 μ g/mL working stock. Eight

concentrations and two quality control samples (QC) are made in 2mL tubes. The first 3 concentrations (50, 35 and 15µg/mL) and the first QC sample (20µg/mL) are made starting with the 200µg/mL working stock. The 4th concentration (5µg/mL) is made with the 50µg/mL solution and the 5th concentration (1µg/mL) with the 15µg/mL. The last three concentrations (0.2, 0.1 and 0.05µg/mL) are made with the 1µg/mL solution. The second QC sample (0.5µg/mL) is made with the first QC sample.

[00479] Of every step of the dilution series, quality control and sample dilutions, a volume is transferred to a 96-well Deepwell plate. The samples are injected on a LC-MS/MS system (API2000 from Applied Biosystems).

[00480] The samples are analyzed on LC-MS/MS with a flow rate of 0.5mL/min. Solvent A is 0.1% Formic Acid in water and solvent B is 0.1% Formic Acid in methanol. The sample is run under positive ion spray on a Pursuit 5 C18 2.0mm column (Varian). The solvent gradient has a total run time of 1.4 minutes and ranges from 10% B to 100% B.

[00481] The thermodynamic solubility samples are analyzed with the aid of QuanLynx software. For the standard curve a linear or quadratic curve can be used in the analysis. Samples of the standard curve that have more than 15% deviation are excluded; the lowest concentrations of the curve may vary up to 20%. Peak areas of the samples are plotted against the standard curve to obtain the solubility of the compound.

[00482] Solubility values are reported in µM or µg/mL.

3.3 *Microsomal stability*

[00483] A 10 mM stock solution of compound in DMSO is 1,668 fold diluted in a 105 mM phosphate buffer pH 7.4. Of this compound dilution, 50 µL is transferred in two 96 assay plates: one for time point 0 min (T0 plate) and one for time point 30 min (T30 plate) and pre-warmed at 37°C.

[00484] In the time zero reference sample (T0 plate), 100 µL MeOH (1:1) is added to the wells. In each assay plate (T0 and T30 min), 50 µL of microsomal mix is then added.

[00485] Final reaction concentrations are: 3 µM compound, 0.5 mg/mL microsomes, 0.4 U/mL GDPDH, 3.3 mM MgCl₂, 3.3 mM glucose-6-phosphate and 1.3 mM NADP⁺.

[00486] The T30 plate is incubated at 37°C, 300 rpm and after 30 minutes of incubation the reaction is stopped with MeOH (1:1). The samples are mixed, centrifuged and the supernatant is harvested for analysis on LC-MS/MS (API2000 from Applied Biosystems).

[00487] The samples are analyzed on LC-MS/MS with a flow rate of 0.5mL/min. Solvent A is 0.1% Formic Acid in water and solvent B is 0.1% Formic Acid in methanol. The sample is run under positive ion spray on a Pursuit 5 C18 2.0mm column (Varian). The solvent gradient has a total run time of 1.4 minutes and ranges from 10% B to 100% B. Peak area from the parent compound at time 0 is considered to be 100% remaining. The percentage remaining after 30 minutes incubation is calculated from time 0. The solubility of

the compound in the final test concentration in buffer is inspected by microscope and results are also reported.

3.4 Hepatocyte stability.

5 [00488] Test compounds (1 μ M initial concentration, n=2) are incubated in Williams' Medium E, containing 4 mM L-glutamine and 2 mM magnesium sulphate, with pooled cryopreserved hepatocytes (Celsis International) in suspension at cell densities of 0.25-0.5 million viable cells/mL. The incubations are performed at 37°C in a shaking water bath with 100 μ L samples taken from the incubation at 0, 10, 20, 45 and 90 minutes, and reactions terminated by addition of 100 μ L of acetonitrile containing carbamazepine as
10 analytical internal standard. Samples are centrifuged and the supernatant fractions analysed by LC-MS/MS. The instrument responses (*i.e.* peak heights) are referenced to the zero time-point samples (as 100%) in order to determine the percentage of compound remaining. Ln plots of the % remaining for each compound are used to determine the half-life for the hepatocyte incubations. Half-life values are calculated from the relationship: $T_{1/2}$ (min) = $-0.693/\lambda$, where λ is the slope of the Ln concentration vs time curve. Standard
15 compounds testosterone, midazolam, and 4-methylumbelliferone are included in the assay design.

3.5 Plasma Protein Binding (Equilibrium Dialysis)

[00489] A 10mM stock solution of the compound in DMSO is diluted with a factor 10 in DMSO. This solution is further diluted in freshly thawed human, rat, mouse or dog plasma (BioReclamation INC) with a
20 final concentration of 5 μ M and final DMSO concentration of 0.5%.

[00490] A Pierce Red Device plate with inserts (ThermoScientific) is prepared and filled with 450 μ L PBS in the buffer chamber and 300 μ L of the spiked plasma in the plasma chamber. The plate is incubated for 4 h at 37°C while shaking at 100rpm. After incubation, 120 μ L of both chambers is transferred to 480 μ L methanol in a 96-well round bottom, PP deep-well plates (Nunc) and sealed with an aluminum foil lid. The
25 samples are mixed and immediately centrifuged 30 min at 1400 rcf at 4°C and the supernatant is transferred to a 96 v-bottom PP plate (Greiner, 651201) for analysis on LC-MS/MS (API2000 from Applied Biosystems).

[00491] The samples are analyzed on LC-MS/MS with a flow rate of 0.5mL/min. Solvent A is 0.1% Formic Acid in water and solvent B is 0.1% Formic Acid in methanol. The sample is run under positive ion
30 spray on a Pursuit 5 C18 2.0mm column (Varian). The solvent gradient has a total run time of 1.4 minutes and ranges from 10% B to 100% B.

[00492] Peak area from the compound in the buffer chamber and the plasma chamber are considered to be 100% compound. The percentage bound to plasma is derived from these results and is reported as percentage bound to plasma.

[00493] The solubility of the compound in the final test concentration in PBS is inspected by microscope to indicate whether precipitation is observed or not.

3.6 *Caco2 Permeability*

5 [00494] Bi-directional Caco-2 assays are performed as described below. Caco-2 cells are obtained from European Collection of Cell Cultures (ECACC, cat 86010202) and used after a 21 day cell culture in 24-well Transwell plates (Corning, cell growth area: 0.33 cm², Membrane pore size: 0.4 μM, membrane diameter: 6.5 mm)..

[00495] 2x10⁵ cells/well are seeded in plating medium consisting of DMEM + GlutaMAX™-I + 1% NEAA + 10% FBS (FetalClone II) + 1% Pen/Strep. The medium is changed every 2 – 3 days.

[00496] Test and reference compounds (propranolol and rhodamine123 or vinblastine, all purchased from Sigma) are prepared in Hanks' Balanced Salt Solution containing 25 mM HEPES (pH7.4) and added to either the apical (125μL) or basolateral (600μL) chambers of the Transwell plate assembly at a concentration of 10 μM with a final DMSO concentration of 0.25%.

15 [00497] 50μM Lucifer Yellow (Sigma) is added to the donor buffer in all wells to assess integrity of the cell layers by monitoring Lucifer Yellow permeation. As Lucifer Yellow (LY) cannot freely permeate lipophilic barriers, a high degree of LY transport indicates poor integrity of the cell layer.

[00498] After a 1 h incubation at 37°C while shaking at an orbital shaker at 150rpm, 70μL aliquots are taken from both apical (A) and basal (B) chambers and added to 100μL 50:50 acetonitrile:water solution
20 containing analytical internal standard (0.5μM carbamazepine) in a 96 well plate.

[00499] Lucifer yellow is measured with a Spectramax Gemini XS (Ex 426nm and Em 538nm) in a clean 96 well plate containing 150μL of liquid from basolateral and apical side.

[00500] Concentrations of compound in the samples are measured by high performance liquid-chromatography/mass spectroscopy (LC-MS/MS).

25 [00501] Apparent permeability (P_{app}) values are calculated from the relationship:

$$P_{app} = \frac{[\text{compound}]_{\text{acceptor final}} \times V_{\text{acceptor}}}{([\text{compound}]_{\text{donor initial}} \times V_{\text{donor}}) / T_{\text{inc}} \times V_{\text{donor}}} / \text{surface area} \times 60 \times 10^{-6} \text{ cm/s}$$

V = chamber volume

T_{inc} = incubation time.

30 Surface area = 0.33cm²

[00502] The Efflux ratios, as an indication of active efflux from the apical cell surface, are calculated using the ratio of P_{app} B>A/ P_{app} A>B.

[00503] The following assay acceptance criteria are used:

Propranolol: P_{app} (A>B) value ≥ 20(×10⁻⁶ cm/s)

Rhodamine 123 or Vinblastine: P_{app} (A>B) value $< 5 (\times 10^{-6} \text{ cm/s})$ with Efflux ratio ≥ 5 .

Lucifer yellow permeability: $\leq 100 \text{ nm/s}$

3.7 *Liability for QT prolongation*

- 5 [00504] Potential for QT prolongation is assessed in the hERG manual patch clamp assay.

3.7.1 *Conventional whole-cell patch-clamp*

- [00505] Whole-cell patch-clamp recordings are performed using an EPC10 amplifier controlled by Pulse v8.77 software (HEKA). Series resistance is typically less than $10 \text{ M}\Omega$ and compensated by greater than
10 60%, recordings are not leak subtracted. Electrodes are manufactured from GC150TF pipette glass (Harvard).

[00506] The external bathing solution contains: 135 mM NaCl, 5 mM KCl, 1.8 mM CaCl_2 , 5 mM Glucose, 10 mM HEPES, pH 7.4.

[00507] The internal patch pipette solution contains: 100mM Kgluconate, 20 mM KCl, 1mM CaCl_2 , 1 mM MgCl_2 , 5mM Na_2ATP , 2mM Glutathione, 11 mM EGTA, 10 mM HEPES, pH 7.2.

- 15 [00508] Drugs are perfused using a Biologic MEV-9/EVH-9 rapid perfusion system.

- [00509] All recordings are performed on HEK293 cells stably expressing hERG channels. Cells are cultured on 12 mm round coverslips (German glass, Bellco) anchored in the recording chamber using two platinum rods (Goodfellow). hERG currents are evoked using an activating pulse to +40 mV for 1000 ms followed by a tail current pulse to -50 mV for 2000 ms, holding potential is -80 mV. Pulses are applied every
20 20s and all experiments are performed at room temperature.

3.7.2 *Data Analysis*

- [00510] IC_{50} values are calculated for each compound tested. The fold difference between the IC_{50} in the manual hERG patch clamp and the unbound IC_{50} in the whole blood assay is calculated.

- 25 [00511] For the concentration response curves, peak tail current amplitude is measured during the voltage step to -50 mV. Curve-fitting of concentration-response data is performed using the equation:

$$y = a + [(b - a) / (1 + 10^{-(\log c - x) d})]$$

- [00512] where a is minimum response, b is maximum response and d is Hill slope, this equation can be used to calculate both IC_{50} (where $y = 50$ and c is the IC_{50} value) and IC_{20} (where $y = 20$ and c is the IC_{20} value). GraphPad® Prism® (Graphpad® Software Inc.) software is used for all curve fitting. A difference of
30 100 fold or greater indicates a low potential for QT prolongation.

3.8 *Pharmacokinetic study*

3.8.1 *Single dose pharmacokinetic study in rats*

[00513] Compounds are formulated in PEG200/physiological saline mixtures for the intravenous route and in PEG400/0.5% methylcellulose (10/90 v/v) for the oral route. Test compounds are orally dosed as a single esophageal gavage at 5-10 mg/kg and intravenously dosed as a bolus via the caudal vein at 1 mg/kg to male Sprague-Dawley rats. Each group consists of 3 rats. Blood samples are collected either via the jugular vein using cannulated rats or at the retro-orbital sinus with lithium heparin as anti-coagulant at the time points in the following range: 0.05 to 8 h (intravenous route), and 0.25 to 6 or 24 h (oral route). Whole blood samples are centrifuged at 5000 rpm for 10 min and the resulting plasma samples are stored at -20°C pending analysis.

3.8.2 *Multiple dose pharmacokinetic study in rats*

[00514] Compounds are formulated in PEG400/0.5% methylcellulose (10/90 v/v) for the oral route. Test compounds are orally dosed as an esophageal daily gavage at 30 or 300 mg/kg to male Sprague-Dawley rats for 14 days. Each group consists of 3 rats. Blood samples are collected via the tail vein with lithium heparin as anti-coagulant at the following time points on day 1, 7 and 14: 0.25, 1, 4, 8 and 24 h. In addition, on day 2 blood samples are taken at 0.25, 1 and 4 h and at day 4 and 11 at 0.25 h. Whole blood samples are centrifuged at 5000 rpm for 10 min and the resulting plasma samples are stored at -20°C pending analysis.

3.8.3 *Quantification of compound levels in plasma*

[00515] Plasma concentrations of each test compound are determined by an LC-MS/MS method in which the mass spectrometer is operated in positive or negative electrospray mode.

3.8.4 *Determination of pharmacokinetic parameters*

[00516] Pharmacokinetic parameters are calculated using Winnonlin® (Pharsight®, US).

3.9 *7-Day rat toxicity study*

[00517] A 7-day oral toxicity study with test compounds is performed in Sprague-Dawley male rats to assess their toxic potential and toxicokinetics, at daily doses of 100, 300 and 1000 mg/kg/day, by gavage, at the constant dosage-volume of 10 mL/kg/day.

[00518] The test compounds are formulated in PEG400/0.5% methylcellulose (10/90, v/v). Each group includes 6 principal male rats as well as 3 satellite animals for toxicokinetics. A fourth group is given PEG400/0.5% methylcellulose (10/90, v/v) only, at the same frequency, dosage volume and by the same route of administration, and acts as the vehicle control group.

[00519] The goal of the study is to determine the lowest dose that results in no adverse events being identified (no observable adverse effect level - NOAEL).

3.10 Cytochrome P450 inhibition

[00520] Reversible CYP inhibition and time-dependent CYP3A4 inhibition is determined in human liver microsomes and specific probe substrates.

5

3.10.1 P450 inhibition in human liver microsomes, reversible inhibition

[00521] The inhibitory potential of a test compound is assessed for human cytochrome P450 isoenzymes CYP1A2, 2C8, 2C9, 2C19, 2D6 and 3A4.

[00522] A 10 mM stock solution of the test compound is prepared in DMSO, serially diluted in Tris buffer (100 mM pH 7.4) and added to hepatic microsomes (Xenotech LLC) and NADPH at 37°C in a shaking water bath. Seven different test compounds concentrations (0.05 to 100 µM), 1% DMSO and 1 mM NADPH are obtained to react.

[00523] After 15 or 30 minutes reactions are terminated by addition of 100 µL of acetonitrile containing carbamazepine as analytical internal standard. Samples are centrifuged and the supernatant fractions analysed by LC-MS/MS. For each isoform, the instrument responses (peak heights) are referenced to those for DMSO controls (considered as 100%) in order to determine the percentage reduction in probe metabolism, using midazolam and testosterone as probe substrate. Percentage inhibition of probe metabolism and Log [test compound concentration] are plotted using Graphpad Prism software. The sigmoidal dose response model is fitted to the data in order to determine the IC₅₀.

[00524] Inhibition of CYP3A4 using nifedipine and atorvastatin as probe substrate is carried out as follows.

[00525] A 1.67 mM stock solution of test compound is prepared in methanol, serially diluted 1:3 in 50 mM potassium phosphate buffer pH7.4 and added to human hepatic microsomes (BD Gentest) and probe substrate. Seven different test compounds concentrations (0.045 - 33.3 µM), 2% methanol, 0.1 mg/mL microsomes, 10 µM atorvastatin or 5 µM nifedipine. After pre-warming 5 minutes at 37°C, the reaction was started by adding cofactor mix (7.65 mg/mL glucose-6-phosphate, 1.7 mg/mL NADP, 6U/mL of glucose-6-phosphate dehydrogenase).

[00526] After 5 min (nifedipine) or 10 min (atorvastatin) at 37°C, the reaction (50 µL) is terminated with 150 µL acetonitrile:methanol (2:1) solution with internal standard (Warfarin). Samples are centrifuged and the supernatant fractions analyzed by LC-MS/MS. The instrument responses (ratio of test compound/internal standard peak areas) are referenced to those for solvent controls (assumed as 100%) in order to determine the percentage reduction in probe metabolism. Percent of control activity vs concentration plots are generated and fitted using GraphPad Prism software to generate IC₅₀.

3.10.2 CYP3A4 inhibition in human liver microsomes, time-dependent

[00527] The time-dependent inhibitory potential of a test compound is assessed for human cytochrome P450 isoenzyme 3A4. The compound is pre-incubated with the human liver microsomes before addition of the probe substrates. The result is compared to the condition where the compound is not pre-incubated with the human liver microsomes to see if there was a shift in IC_{50} , indicating time-dependent inhibition.

[00528] A 10 mM stock solution of test compound is prepared in DMSO and diluted 1:20 with Tris buffer (100 mM pH 7.4) and further serially diluted in Tris buffer/5% DMSO.

[00529] The cofactor, NADPH, and each test compound dilution is mixed in two separate plates for 0 and 30 min pre-incubation. Human hepatic microsomes (Xenotech LLC) are added only to the "30 minute pre-incubation" plate and both plates are then incubated for 30 minutes at 37°C in a shaking water bath. Following the pre-incubation, microsomes are added to the "0 minute" plate and appropriate probe substrates (in 0.5% DMSO) are added to both plates. Plates are then returned to the water bath for a further incubation.

[00530] In total, six different test compound concentrations (1.6 to 50 μ M) are assessed. Reactions are terminated with 100 μ L of acetonitrile containing carbamazepine as analytical internal standard. Samples are centrifuged and the supernatant fractions analysed by LC- MS/MS. For each isoform, the instrument responses (peak height ratio with internal standard) are referenced to those for DMSO controls (considered as 100%) in order to determine the percentage reduction in probe metabolism. Percentage inhibition of probe metabolism and Log [Test Compound concentration] are plotted using Graphpad Prism software. The sigmoidal dose response model is fitted to the data in order to determine the IC_{50} .

4. *In-vivo* studies

[00531] The *in-vivo* activity of the compounds of the invention may be demonstrated in the following *in vivo* efficacy inflammation models.

4.1 *Inflammatory bowel disease (mice).*

[00532] The mouse chronic DSS-induced inflammatory bowel disease model (IBD) is a well validated disease model for inflammatory bowel disease (Wirtz S. *et al.*, 2007 Nature Protocols 2, 541-546; Sina C. *et al.*, 2009 J. Immunol. 183 7514-7522).

[00533] To induce a chronic colitis, female BALB/c mice are fed with 4% dextran sodium sulfate (DSS) dissolved in drinking water for 4 days, followed by 3 days of regular drinking water. This cycle is repeated three times. This protocol allows inducing a strong colitis while avoiding high mortality rates. Animals are divided into several groups:

- a. intact water; vehicle alone, n=10),
- b. diseased (DSS; vehicle alone, n=10),

- c. sulfasalazine used as reference (DSS; 20 mg/kg/day, *p.o.*, n=10) and
- d. the tested compound (DSS; 1, 3, 10, 30 mg/kg/day, *p.o.*, n=10).

[00534] Clinical parameters are measured every other day. The disease activity index (DAI) is a composite measure combining of the individual scores for weight loss, stool consistency and rectal bleeding. Mice are sacrificed at day 20 of the experiment according to the protocol introduced by Sina *et al.*(2009). At sacrifice time, the complete colon is removed and rinsed with sterile PBS. Segments of the distal colon are dissected for histological analysis, gene expression and protein level measurement.

4.2 Collagen-induced arthritis (mice).

[00535] The mouse collagen-induced arthritis (CIA) is the gold standard rheumatoid arthritis model (Brand, *et al.*, 2007 Nature Protocols 2, 1269- 1275, Lin *et al.*, 2007 Br J Pharmacol 1, 829-831). DBA1//J male mice are injected with a collagen II solution (Completed Freund's adjuvant). Immune reaction is boosted by a second injection (incomplete Freund's adjuvant) 21 days later. At day 31, arthritis is scored according to the method of Khachigian *et al.* (Khachigian *et al.*, 2006 Nature Protocols 1, 2512-2516) and animals are randomized to reach an average clinical score of 2 per group. Animals are divided into several groups: intact (no treatment, n=5), diseased (vehicle alone, n=10), Enbrel® as reference (10 mg/kg, 3x week., *i.p.*, n=10), and the tested compound (3, 10 or 30 mg/kg/day, *p.o.*, n=10). Therapeutic dosing lasted from day 31 to day 46 and the arthritis is scored every day. Mice are sacrificed at day 46, X-ray photos are taken of the hind paws of each individual animal and the severity of bone erosion is ranked with the radiological Larsen's score (Salvemini *et al.*, 2001 Arthritis Rheum 44, 2909-2921).

4.3 Tobacco smoke model (mice)

[00536] Daily exposures of female inbred C57BL/6J mice to tobacco smoke (TS) for 11 consecutive days result in pulmonary inflammation, as indicated by an increase in the total number of cells recovered in the bronchoalveolar lavage (BAL), when compared with a similarly treated air-exposed group, 24 h after the final exposure. The exposure period to TS is increased initially from 25 minutes at the start of the study (day 1) to a maximum of 45 minutes on day 3 until day 11. Animals are divided into several groups: intact (no treatment, n=5), diseased (vehicle alone, n=10), Roflumilast as reference (5 mg/kg/day *p.o.*, n=10), and the tested compounds (10 or 30 mg/kg/bid, *p.o.*, n=10). At the end of 11 days, the numbers of macrophages, epithelial cells, neutrophils and lymphocytes are counted in the BAL. BAL is further analysed for gene expression and protein level. Lung tissue is dissected for histological analysis, gene expression and protein level measurement.

[00537] It will be appreciated by those skilled in the art that the foregoing descriptions are exemplary and explanatory in nature, and intended to illustrate the invention and its preferred embodiments. Through routine experimentation, an artisan will recognise apparent modifications and variations that may be made without departing from the spirit of the invention. All such modifications coming within the scope of the appended
5 claims are intended to be included therein. Thus, the invention is intended to be defined not by the above description, but by the following claims and their equivalents.

10 [00539] It should be understood that factors such as the differential cell penetration capacity of the various compounds can contribute to discrepancies between the activity of the compounds in the in vitro biochemical and cellular assays.

[00540] At least some of the chemical names of compound of the invention as given and set forth in this application, may have been generated on an automated basis by use of a commercially available chemical
15 naming software program, and have not been independently verified. Representative programs performing this function include the Lexichem naming tool sold by Open Eye Software, Inc. and the Autonom Software tool sold by MDL, Inc. In the instance where the indicated chemical name and the depicted structure differ, the depicted structure will control.

[00541] Chemical structures shown herein were prepared using either ChemDraw® or ISIS® /DRAW.
20 Any open valency appearing on a carbon, oxygen or nitrogen atom in the structures herein indicates the presence of a hydrogen atom. Where a chiral center exists in a structure but no specific stereochemistry is shown for the chiral center, both enantiomers associated with the chiral structure are encompassed by the structure.


25

[00542] REFERENCES

- Wittenberger *et al.*, **2001** J Mol Biol, 307, 799-813
- Yousefi S *et al.*, **2001** J Leukoc Biol, 69, 1045-52
- Wang *et al.*, **2006** The Journal of Biological Chemistry, 281, 45, 34457-34464
- 5 Venkataraman *et al.*, **2005**, Immunology Letters, 101, 144-153
- WO2007/027661 A2
- Berry *et al.*, **2010**, Nature, 466, 973-979
- Bouchard *et al.*, **2007**, Glia, 55:790-800
- Bundgard, H., **1985** Design of Prodrugs, pp. 7-9, 21-24, Elsevier, Amsterdam 1985
- 10 Part 8 of Remington's Pharmaceutical Sciences, 17th edition, **1985**, Mack Publishing Company, Easton, Pennsylvania
- T. W. Greene and P. G. M. Wuts, **2006** *Protecting Groups in Organic Synthesis*, Wiley-Blackwell; 4th Revised edition
- Young Kim *et al.*, **2007** Bioorganic & Medicinal Chemistry 15, 2667-2679
- 15 Le Pouls *et al.*, **2003**, The Journal of Biological Chemistry, 278, 28, 25481-25489
- Brown *et al.*, **2003**, The Journal of Biological Chemistry, 278, 13, 11312-11319
- Stoddart *et al.*, **2008**, Pharmacological Reviews, 60, 405-417
- Wirtz S. *et al.*, **2007** Nature Protocols 2, 541-546
- Sina C. *et al.*, **2009** J. Immunol. 183 7514-7522
- 20 Brand, *et al.*, **2007** Nature Protocols 2, 1269- 1275
- Lin *et al.*, **2007** Br J Pharmacol 1, 862-872
- Khachigian *et al.*, **2006** Nature Protocols 1, 2512-2516
- Salvemini *et al.*, **2001** Arthritis Rheum 44, 2909-2921
- Du Bois, **2010**, Nat Rev, Drug Discovery, 9, 129
- 25 Nagasaki *et.al.*, **2012**, FEBS Letters, 586, 368-372

WHAT IS CLAIMED:

1. A compound according to Formula Ia:



wherein

R¹ is H, Me, or halo;

L₁ is absent or is -O-, -S-, or -NR^{4a}-;

G is

R²,

-W-L₂-R², or

-W-L₃-R³;

W is C₁₋₄ alkylene, C₂₋₄ alkenylene having one double bond, or C₂₋₄ alkynylene having one triple bond;

L₂ is absent or is -O-;

R² is

- H,
- C₁₋₈ alkyl, optionally substituted with one to three groups independently selected from
 - o OH,
 - o halo,
 - o CN,
 - o C₁₋₆ alkoxy,
 - o C₃₋₇ cycloalkyl,
 - o 4-6 membered heterocycloalkyl comprising one to three heteroatoms independently selected from S, and O,
 - o 5-6 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, and
 - o phenyl,
- C₄₋₇ cycloalkenyl comprising one double bond,

- 5-7 membered heterocycloalkenyl comprising one double bond, and one to three heteroatoms independently selected from N, O, and S,
- C₃₋₇ cycloalkyl optionally substituted with one or more independently selected R⁵ groups,
- 4-10 membered heterocycloalkyl comprising one to two heteroatoms independently selected from S, and O, optionally substituted with one to three independently selected R⁵ groups,
- 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O optionally substituted with one to three independently selected R⁶ groups, or
- C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁶ groups;

L₃ is -NR^{4b}-;

R³ is

- C₁₋₄ alkyl substituted with
 - o C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁷ groups, or
 - o 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, optionally substituted with one or more R independently selected R⁷ groups,
- 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, optionally substituted with one or more independently selected R⁷ group, or
- C₆₋₁₀ aryl optionally substituted with one or more independently selected R⁷ groups;

Each R^{4a} and R^{4b} is independently selected from H, C₁₋₄ alkyl, and C₃₋₇ cycloalkyl;


R⁵ is oxo or R⁶;

R⁶ is

- OH,
- halo,
- -NO₂,
- C₁₋₆ alkyl optionally substituted with one to three groups independently selected from halo, and OH,
- C₁₋₆ alkoxy optionally substituted with one to three groups independently selected from halo, and OH,
- C₃₋₇ cycloalkyl,
- -C(=O)OR⁸,
- -C(=O)NR⁹R¹⁰,
- -NHC(=O)-C₁₋₄ alkyl,
- -CN,
- phenyl,

- or a pharmaceutically acceptable salt, or a solvate, or a solvate of the pharmaceutically acceptable salt.

3. A compound or pharmaceutically acceptable salt thereof, according to claim 1, wherein the compound is according to Formula IIa or IIIa:



6. A compound or pharmaceutically acceptable salt thereof, according to claim 3, wherein the compound is according to Formula IIIa, and L₁ is absent; W is C₂₋₄ alkynylene having one triple bond; and L₂, and R² are as defined in claim 1.

7. A compound or pharmaceutically acceptable salt thereof, according to claim 5 or 6, wherein L_2 is absent.
8. A compound or pharmaceutically acceptable salt thereof, according to claim 3, wherein the compound is according to Formula IIIa, wherein L_1 and L_2 are absent, W is $-\text{CH}_2-\text{CH}_2-$, $-\text{CH}=\text{CH}-$, or $\text{C}\equiv\text{C}-$, and R^2 is as defined in claim 1.
9. A compound or pharmaceutically acceptable salt thereof, according to any one of claims 3-8, wherein R^2 is 5-10 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, optionally substituted with one to three independently selected R^6 groups.
10. A compound or pharmaceutically acceptable salt thereof, according to claim 9, wherein R^6 is selected from OH, halo, C_{1-6} alkyl, C_{1-6} alkyl substituted with one or more halo, C_{1-6} alkoxy, $-\text{CN}$, C_{3-7} cycloalkyl, 4-7 membered heterocycloalkyl comprising one to three heteroatoms independently selected from N, O, and S, and phenyl.
11. A compound or pharmaceutically acceptable salt thereof, according to any one of claims 3-8 wherein R^2 is C_{1-8} alkyl optionally substituted with one to three groups independently selected from OH, halo, CN, C_{1-6} alkoxy, C_{3-7} cycloalkyl, 4-6 membered heterocycloalkyl comprising one to three heteroatoms independently selected from S, and O, 5-6 membered heteroaryl comprising one to three heteroatoms independently selected from N, S, and O, and phenyl
12. A compound or pharmaceutically acceptable salt thereof, according to any one of claims 3-8, wherein R^2 is C_{3-7} cycloalkyl.
13. A compound or pharmaceutically acceptable salt thereof, according to claim 1, wherein the compound is selected from:
 - 9-Allyloxy-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 - 2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-3-yl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 - 2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-4-yl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 - 2-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile,
 - 3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile,
 - 4-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile,
 - [2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-acetonitrile,

2-([1,4]Dioxan-2-ylmethoxy)-9-(oxazol-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(pyridin-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(3,5-Dichloro-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Benzofuran-2-yl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-indole-1-carboxylic acid tert-butyl ester,

2-([1,4]Dioxan-2-ylmethoxy)-9-(1H-indol-2-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(6-methoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(6-trifluoromethyl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-3H-imidazol-4-ylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(5-tert-Butyl-[1,2,4]oxadiazol-3-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

5-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-pyridine-2-carboxylic acid methanamide,

2-([1,4]Dioxan-2-ylmethoxy)-9-pent-1-ynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(2-pyridin-2-yl-ethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(2-pyrazin-2-yl-ethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(1H-indol-5-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(2-methoxy-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(5-methoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(1H-indazol-5-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(4-methoxy-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzamide,

5-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-2-fluoro-benzamide,

N-{3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-phenyl}-acetamide,

9-Cyclopropylethynyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(1-hydroxy-cyclopentylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-pyrimidin-5-yl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Cyclohex-1-enyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(1-methyl-1H-indol-5-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(6-methyl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-2-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

5-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-pent-4-yne nitrile,

2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(4-methoxy-phenylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-3-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

4-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-N-methyl-benzamide,

2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(2-Chloro-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(1,5-Dimethyl-1H-pyrazol-3-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(1-methyl-1H-pyrazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-[1,2,4]oxadiazol-5-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(4-morpholin-4-yl-phenyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-4-fluoro-benzamide,

3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-5-fluoro-benzamide,

9-(3,3-Dimethyl-but-1-ynyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-pyridin-4-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-isoxazol-5-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-methyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(2-methoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(3,6-Dihydro-2H-pyran-4-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

5-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-pyridine-2-carbonitrile,

2-([1,4]Dioxan-2-ylmethoxy)-9-(6-isopropoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(6-ethoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(6-morpholin-4-yl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(2,3-Dimethoxy-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(3-Chloro-2-methoxy-pyridin-4-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(2-methyl-pyridin-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

3-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-isonicotinonitrile,

9-(2,5-Dimethoxy-phenyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(2-ethoxy-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(2,6-Dimethoxy-pyridin-3-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

4-[2-([1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-nicotinonitrile,

9-tert-Butoxymethyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(2-pyrrolidin-1-yl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(6-pyrrolidin-1-yl-pyridin-3-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(5-phenyl-oxazol-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(5-tert-Butyl-oxazol-2-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(5-Cyclopropyl-[1,2,4]oxadiazol-3-ylmethoxy)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(5-ethyl-[1,2,4]oxadiazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(5-methyl-[1,2,4]oxadiazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(5-isopropyl-[1,2,4]oxadiazol-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Cyclopentylethynyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 9-Cyclohexylethynyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-hex-1-ynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 9-[3-(Benzyl-methyl-amino)-prop-1-ynyl]-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-5-methyl-hex-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 9-Cyclopropyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-4-methyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-ethyl-3-hydroxy-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-phenyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 9-(3-Benzylamino-prop-1-ynyl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-[(furan-2-ylmethyl)-amino]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(1-ethyl-1H-pyrazol-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-[1-(3-methyl-butyl)-1H-pyrazol-4-yl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(5-methyl-furan-2-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-hex-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
 9-(3,5-Dimethyl-1H-pyrazol-4-yl)-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(1H-pyrazol-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(1-propyl-1H-pyrazol-4-yl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-[2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile,

2-[2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-4-oxo-6,7-dihydro-4H-pyrimido[6,1-a]isoquinolin-9-yl]-benzonitrile,

9-(5-Cyclopropyl-[1,2,4]oxadiazol-3-ylmethoxy)-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-ethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-pyrimidin-2-ylethynyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(3-phenylamino-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-pyridin-3-yl-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Cyclopentylloxymethyl-2-([1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-4-methyl-pent-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Cyclopropylethynyl-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-but-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(3-imidazol-1-yl-prop-1-ynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(2-Cyclopropyl-ethyl)-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Cyclopentylloxymethyl-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-pyridin-3-yl-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Allyloxy-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Allyloxy-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(tetrahydro-pyran-4-yloxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-{3-[(pyridin-3-ylmethyl)-amino]-prop-1-ynyl}-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-pentyl-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Cyclopropylethynyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(2-Cyclopropyl-ethyl)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(oxetan-3-yloxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methyl-oxetan-3-ylmethoxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(2,2-Dimethyl-butylamino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-ethyl-hexylamino)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-methoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-ethoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Cyclopropylmethoxy-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-fluoro-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[3-(2-methoxy-ethoxy)-prop-1-ynyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[3-(2-ethoxy-ethoxy)-prop-1-ynyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[3-(2-fluoro-ethoxy)-prop-1-ynyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(2,2-Dimethyl-propoxymethyl)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Cyclohexyloxymethyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Cyclopropylmethoxymethyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(tetrahydro-pyran-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(4,4-Dimethyl-pentyloxy)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(3-Cyclopropyl-propoxy)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Cyclohexylamino-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-4,4-dimethyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-Cyclopentylmethoxymethyl-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-phenylamino-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(Cyclohexyl-methyl-amino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(Cyclohexylmethyl-amino)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[(tetrahydro-pyran-4-ylmethyl)-amino]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-ethyl-3-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-3-methyl-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-hydroxy-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(2,2-Dimethyl-propoxy)-2-((S)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(tetrahydro-pyran-4-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-4-methyl-pentyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(tetrahydro-pyran-4-ylmethoxymethyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-([1,4]Dioxan-2-ylmethoxy)-9-methoxy-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(oxetan-3-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

9-(3-Cyclopropyl-propoxy)-2-((R)-1-[1,4]dioxan-2-ylmethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-[2-(1-hydroxy-cyclopentyl)-ethyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-hydroxy-tetrahydro-pyran-4-ylethynyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(3-methoxy-propyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-[2-(1-hydroxy-cyclopentyl)-ethyl]-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-propoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-isopropoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-propoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,

2-((R)-1-[1,4]Dioxan-2-ylmethoxy)-9-(2-isopropoxy-ethoxy)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one,
and

2-((S)-1-[1,4]Dioxan-2-ylmethoxy)-9-(4-methoxy-butyl)-6,7-dihydro-pyrimido[6,1-a]isoquinolin-4-one.

14. A pharmaceutical composition comprising a compound or pharmaceutically acceptable salt thereof, according to any one of claims 1-13, and a pharmaceutically acceptable carrier.

15. The pharmaceutical composition according to claim 14 comprising a further therapeutic agent.

16. The compound, or pharmaceutically acceptable salt thereof, according to any one of claims 1-13 for use in the treatment or prophylaxis of an inflammatory condition.

17. The pharmaceutical composition according to claim 14 or 15 for use in the treatment or prophylaxis of an inflammatory condition.

18. The compound, or pharmaceutically acceptable salt thereof according to claim 16, wherein the inflammatory condition is rheumatoid arthritis, vasculitis, chronic obstructive pulmonary disease, asthma, idiopathic pulmonary fibrosis, psoriasis, Crohn's disease, or ulcerative colitis.


19. Use of the compound, or pharmaceutically acceptable salt thereof, according to any one of claims 1-13 for the treatment or prophylaxis of an inflammatory condition.

20. Use of the compound, or pharmaceutically acceptable salt thereof, according to any one of claims 1-13 for the manufacture of a medicament for the treatment or prophylaxis of an inflammatory condition.

21. The use of claim 19 or 20, wherein the inflammatory condition is rheumatoid arthritis, vasculitis, chronic obstructive pulmonary disease, asthma, idiopathic pulmonary fibrosis, psoriasis, Crohn's disease, or ulcerative colitis.

22. The use according to any one of claims 19-21, in combination with a further therapeutic agent.

23. The pharmaceutical composition according to claim 17, wherein the inflammatory condition is rheumatoid arthritis, vasculitis, chronic obstructive pulmonary disease, asthma, idiopathic pulmonary fibrosis, psoriasis, Crohn's disease, or ulcerative colitis.



1a