



(51) International Patent Classification:

C12N 5/0783 (2010.01)

(21) International Application Number:

PCT/IB2022/055504

(22) International Filing Date:

14 June 2022 (14.06.2022)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/210,683

15 June 2021 (15.06.2021)

US

(71) Applicant: **TAKEDA PHARMACEUTICAL COMPANY LIMITED** [JP/JP]; 1-1, Doshomachi 4-chome, Chuo-Ku, Osaka-Shi, Osaka 541-0045 (JP).

(72) Inventors: **CAO, Lan**; c/o Millennium Pharmaceuticals, Inc., 40 Landsdowne Street, Cambridge, Massachusetts 02139 (US). **CHANG, Hui-Hsin**; c/o Millennium Pharmaceuticals, Inc., 40 Landsdowne Street, Cambridge, Massachusetts 02139 (US). **HU, Jianxin**; c/o Millennium Pharmaceuticals, Inc., 40 Landsdowne Street, Cambridge, Massachusetts 02139 (US). **QIAN, Yu**; c/o Millennium Pharmaceuticals, Inc., 40 Landsdowne Street, Cambridge, Massachusetts 02139 (US). **SHI, Xi**; c/o Millennium Pharmaceuticals, Inc., 40 Landsdowne Street, Cambridge, Massachusetts 02139 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IQ, IR, IS, IT, JM, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- in black and white; the international application as filed contained color or greyscale and is available for download from PATENTSCOPE

(54) Title: METHOD FOR PRODUCING NATURAL KILLER CELLS FROM PLURIPOTENT STEM CELLS

(57) Abstract: The present disclosure provides, among other things, a method for efficiently producing a cell population enriched in Natural Killer cells (NK cells) from induced pluripotent cells.



METHOD FOR PRODUCING NATURAL KILLER CELLS FROM PLURIPOTENT STEM CELLS

RELATED APPLICATIONS

[0001] The present application claims priority to U.S. Provisional Application Serial No. 63/210,683, filed June 15, 2021; the contents of which are herein incorporated by reference in their entirety.

BACKGROUND

[0002] Natural Killer (NK) cells are cytotoxic lymphocytes of the immune system. NK cells are cytotoxic against cancerous, pathogen-infected and otherwise damaged cells. NK cells are innate lymphoid cells (ILCs), specifically large granular cytotoxic lymphocytes that bridge the innate and the adaptive arms of the immune response. They make up 10-15% of circulating lymphocytes in the peripheral blood. NK cells also exhibit the highest level of cytotoxic activity within the immune system. Therefore, altered NK cell functionality or numbers impact the functioning of the immune system against infection and cancer.

[0003] NK cells lack specific cell surface antigen receptors. Because of this, NK cells may kill cancerous and pathogen-infected cells without prior sensitization, making them part of the innate immune response. They also have a role in tumor immunosurveillance by directly influencing the adaptive immune response. These features and others make NK cells a particularly attractive cell type for use in adoptive cell therapies.

[0004] Various protocols have been described to obtain NK cells from pluripotent stem cells (e.g., induced pluripotent stem cells (iPSCs)) or progenitor cells. However, the previously described protocols are labor intensive and require multiple, time consuming steps, including steps of cell isolation which in turn increases manufacturing/production time and monetary cost to produce NK cells.

[0005] As development of NK cell based therapies is starting to gain momentum, there is a need for more efficient and robust methods for manufacturing such cell therapies.

SUMMARY OF THE INVENTION

[0006] The present disclosure provides, among other things, improved methods of producing NK cells. NK cells are generally characterized as being CD56+/CD3- cells. This disclosure is based, at least in part, on the surprising discovery of efficient and robust production of NK cells from a bulk cell population derived from pluripotent stem cells (e.g., induced pluripotent stem cells (iPSCs)). The methods provided herein for the production of NK cells include methods that are performed without an isolation step, as well as methods in which one or more isolation steps are performed. As described herein, the present disclosure unexpectedly shows that NK cells can be successfully derived from a bulk cell population without requiring isolation of any cell types based on lineages. Alternatively, in some embodiments, the method can be performed using one or more isolation steps. The present disclosure is based on an unexpected and surprising discovery that a cell population enriched in NK cells can be derived from iPSCs or a hematopoietic progenitor cell (HPC) bulk, either by simply changing culture conditions or by isolating a population of cells containing CD4- cells as an intermediate to differentiate into NK cells. The methods described herein have numerous benefits over existing methods of producing NK cells, including for example, the ability to scale up and produce large quantities of NK cells in a cost and time efficient manner. Thus, the methods described herein represent efficient and robust manufacturing processes for NK cell based therapies.

[0007] iPSC-derived NK cells (also referred to herein as “iNK cells”) produced by the methods described herein are functional and can be further genetically modified, such as through the introduction of a CAR to target specific cell populations.

[0008] In some aspects, a method of producing a cell population enriched in NK cells is provided, the method comprising the steps of:

[0009] (A) culturing a population of pluripotent stem cells under a first set of conditions resulting in a cell population comprising at least 20% CD34+ HPCs (HPC bulk); and

[0010] (B) changing the first set of conditions to a second set of conditions, thereby resulting in a cell population comprising at least 5% CD4- cells; and

[0011] (C) changing the second set of conditions to a third set of conditions, thereby resulting in a cell population enriched in NK cells.

[0012] In some embodiments, the method is performed without an isolation step.

[0013] In some embodiments, the method comprises isolating CD4⁻ cells after step (B) before contacting the isolated cells with the third set of conditions in step (C).

[0014] In some embodiments, the step of isolating CD4⁻ cells comprises removing CD4⁺ cells from the cell population, or separating CD4⁺ cells from the rest of the cells and differentiating the remaining cells, which would inherently lack CD4 surface marker, into NK cells.

[0015] In some aspects, a method of producing a cell population enriched in NK cells is provided, the method comprising the steps of:

[0016] (A) culturing a population of pluripotent stem cells under a first set of conditions resulting in a cell population comprising at least 20% CD34⁺ HPCs (HPC bulk); and

[0017] (B) changing the first set of conditions to a second set of conditions, thereby resulting in a cell population comprising at least 5% CD4⁻ cells as well as at least 5% CD4⁺ cells; and

[0018] (C) changing the second set of conditions to a third set of conditions, thereby resulting in a cell population enriched in NK cells, wherein the method is performed without an isolation step.

[0019] In some embodiments, the cell population enriched in NK cells comprises at least 30% NK cells. For example, the cell population enriched in NK cells comprises 30%, 35%, 40%, 45%, 50%, or more than 50% NK cells.

[0020] In some embodiments, the resulting cell population in step (B) comprises at least 10%, 15% or 20% CD4⁻ cells. Accordingly, in some embodiments, the resulting cell population in step (B) comprises at least 10% CD4⁻ cells. In some embodiments, the resulting cell population in step (B) comprises at least 15% CD4⁻ cells. In some embodiments, the resulting cell population in step (B) comprises at least 20% CD4⁻ cells. In some embodiments, the resulting cell population in step (B) also comprises CD4⁺ cells.

[0021] In some embodiments, the CD4⁻ cells in step (B) are CD8⁺ cells.

[0022] In some embodiments, the CD4⁻ cells in step (B) are CD8⁻.

[0023] In some embodiments, the resulting cell population in step (B) comprises between 20 and 55% CD4⁻/CD8⁺ cells. For example, in some embodiments, the resulting cell population in step (B) comprises 20% CD4⁻/CD8⁺ cells. In some embodiments, the resulting cell population in step (B) comprises 25% CD4⁻/CD8⁺ cells. In some embodiments, the resulting cell population in step (B) comprises 30% CD4⁻/CD8⁺ cells. In some embodiments, the resulting cell population in step (B) comprises 35% CD4⁻/CD8⁺ cells. In some embodiments, the resulting cell population in step (B) comprises 40% CD4⁻/CD8⁺ cells. In some embodiments, the resulting cell population in step (B) comprises 45% CD4⁻/CD8⁺ cells. In some embodiments, the resulting cell population in step (B) comprises 50% CD4⁻/CD8⁺ cells. In some embodiments, the resulting cell population in step (B) comprises 55% CD4⁻/CD8⁺ cells.

[0024] In some embodiments, the resulting cell population in step (B) comprises between 25 and 55% CD4⁻/CD8⁻ cells. For example, in some embodiments, the resulting cell population in step (B) comprises about 25% CD4⁻/CD8⁻ cells. In some embodiments, the resulting cell population in step (B) comprises about 30% CD4⁻/CD8⁻ cells. In some embodiments, the resulting cell population in step (B) comprises about 35% CD4⁻/CD8⁻ cells. In some embodiments, the resulting cell population in step (B) comprises about 40% CD4⁻/CD8⁻ cells. In some embodiments, the resulting cell population in step (B) comprises about 45% CD4⁻/CD8⁻ cells. In some embodiments, the resulting cell population in step (B) comprises about 50% CD4⁻/CD8⁻ cells. In some embodiments, the resulting cell population in step (B) comprises about 55% CD4⁻/CD8⁻ cells.

[0025] In some embodiments, the resulting cell population in step (B) includes a combination of CD4⁻/CD8⁻ cells and CD4⁻/CD8⁺ cells. In some embodiments, the resulting cell population in step (B) additionally includes at least 5% cells that are CD4⁺/CD8⁺ cells and/or CD4⁺/CD8⁻ cells.

[0026] In some embodiments, the NK cells are CD56⁺/CD3⁻ cells.

[0027] In some embodiments, the pluripotent stem cells are induced pluripotent stem cells (iPSCs).

[0028] In some embodiments, the first set of conditions comprises a culture medium comprising at least one compound selected from bone morphogenetic protein-4 (BMP4) vascular endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF), ascorbic acid, Flt3 ligand (Flt3L), thrombopoietin (TPO) and TGF β inhibitor. Accordingly, in some embodiments, the first set of conditions comprises a culture medium comprising BMP4. In some embodiments, the first set of conditions comprises a culture medium comprising VEGF. In some embodiments, the first set of conditions comprises a culture medium comprising bFGF. In some embodiments, the first set of conditions comprises a culture medium comprising ascorbic acid. In some embodiments, the first set of conditions comprises a culture medium comprising Flt3L. In some embodiments, the first set of conditions comprises a culture medium comprising TPO. In some embodiments, the first set of conditions comprises a culture medium comprising TGF β inhibitor.

[0029] In some embodiments, the first set of conditions comprises a culture medium comprising BMP4 at a concentration of between 5 ng/mL – 500 ng/mL. For example, the first set of conditions comprises a culture medium comprising BMP4 at a concentration of between 5 ng/mL – 25 ng/mL, 25 ng/mL – 50 ng/mL, 25 ng/mL – 75 ng/mL, 25 ng/mL – 100 ng/mL, 100 ng/mL – 125 ng/mL, 100 ng/mL – 250 ng/mL, or 250 ng/mL – 500 ng/mL.

[0030] In some embodiments, the BMP4 is at a concentration of 50 ng/mL.

[0031] In some embodiments, the first set of conditions comprises a culture medium comprising VEGF at a concentration of between 5 ng/mL – 500 ng/mL. For example, in some embodiments, the first set of conditions comprises a culture medium comprising VEGF at a concentration of between 5 ng/mL – 25 ng/mL, 25 ng/mL – 50 ng/mL, 25 ng/mL – 75 ng/mL, 25 ng/mL – 100 ng/mL, 100 ng/mL – 125 ng/mL, 100 ng/mL – 250 ng/mL, or 250 ng/mL – 500 ng/mL.

[0032] In some embodiments, the VEGF is at a concentration of about 50 ng/mL.

[0033] In some embodiments, first set of conditions comprises a culture medium comprising bFGF at a concentration of between 5 ng/mL – 500 ng/mL. For example, in some embodiments, the first set of conditions comprises a culture medium comprising bFGF at a concentration of between 5 ng/mL – 25 ng/mL, 25 ng/mL – 50 ng/mL, 25 ng/mL – 75 ng/mL, 25 ng/mL – 100 ng/mL, 100 ng/mL – 125 ng/mL, 100 ng/mL – 250 ng/mL, or 250 ng/mL – 500 ng/mL.

[0034] In some embodiments, the bFGF is at a concentration of 50 ng/mL.

[0035] In some embodiments, the first set of conditions comprises a culture medium comprising ascorbic acid at a concentration of between 5 µg/mL - 500 µg/mL. For example, in some embodiments, the first set of conditions comprises a culture medium comprising ascorbic acid at a concentration of between 5 ng/mL – 25 ng/mL, 25 ng/mL – 50 ng/mL, 25 ng/mL – 75 ng/mL, 25 ng/mL – 100 ng/mL, 100 ng/mL – 125 ng/mL, 100 ng/mL – 250 ng/mL, or 250 ng/mL – 500 ng/mL.

[0036] In some embodiments, the ascorbic acid is at a concentration of 50 µg/mL.

[0037] In some embodiments, the first set of conditions comprises a culture medium comprising Flt3L at a concentration between 1 ng/ml - 100 ng/mL. For example, in some embodiments, the first set of conditions comprises a culture medium comprising Flt3L at a concentration between 5 ng/mL – 25 ng/mL, 25 ng/mL – 50 ng/mL, 25 ng/mL – 75 ng/mL, or 25 ng/mL – 100 ng/mL.

[0038] In some embodiments, the Flt3L is at a concentration of 50 ng/mL.

[0039] In some embodiments, the first set of conditions comprises a culture medium comprising TPO at a concentration of between 1 ng/mL – 200 ng/mL. For example, in some embodiments, the first set of conditions comprises a culture medium comprising TPO at a concentration of between 5 ng/mL – 25 ng/mL, 25 ng/mL – 50 ng/mL, 25 ng/mL – 75 ng/mL, 25 ng/mL – 100 ng/mL, 100 ng/mL – 125 ng/mL, or 100 ng/mL – 200 ng/mL.

[0040] In some embodiments, the TPO is at a concentration of 100 ng/mL.

[0041] In some embodiments, the second set of conditions comprises a culture medium comprising at least one compound selected from the group consisting of ascorbic acid, stem cell factor (SCF), IL-7, Flt3L, thrombopoietin (TPO), p38 inhibitor and SDF-1. Accordingly, in some embodiments, the second set of conditions comprises a culture medium comprising ascorbic acid. In some embodiments, the second set of conditions comprises a culture medium comprising SCF. In some embodiments, the second set of conditions comprises a culture medium comprising IL-7. In some embodiments, the second set of conditions comprises a culture medium comprising Flt3L. In some embodiments, the second set of conditions comprises a culture medium comprising TPO. In some embodiments, the second set of

conditions comprises a culture medium comprising p38 inhibitor. In some embodiments, the second set of conditions comprises a culture medium comprising SDF-1.

[0042] In some embodiments, the second set of conditions comprises a culture medium comprising ascorbic acid at a concentration between 5 $\mu\text{g/mL}$ - 500 $\mu\text{g/mL}$. For example, in some embodiments, the second set of conditions comprises a culture medium comprising ascorbic acid at a concentration between 5 ng/mL – 25 ng/mL , 25 ng/mL – 50 ng/mL , 25 ng/mL – 75 ng/mL , 25 ng/mL – 100 ng/mL , 100 ng/mL – 125 ng/mL , 100 ng/mL – 250 ng/mL , or 250 ng/mL – 500 ng/mL .

[0043] In some embodiments, the ascorbic acid is at a concentration of 50 $\mu\text{g/mL}$.

[0044] In some embodiments, the second set of conditions comprises a culture medium comprising SCF at a concentration between 5 ng/mL – 100 ng/mL . For example, in some embodiments, the second set of conditions comprises a culture medium comprising SCF at a concentration between 5 ng/mL – 25 ng/mL , 25 ng/mL – 50 ng/mL , 25 ng/mL – 75 ng/mL , or 25 ng/mL – 100 ng/mL .

[0045] In some embodiments, the SCF is at a concentration 50 ng/mL .

[0046] In some embodiments, the second set of conditions comprises a culture medium comprising IL-7 at a concentration between 1 ng/mL – 100 ng/mL . For example, in some embodiments, the second set of conditions comprises a culture medium comprising IL-7 at a concentration between 1 ng/mL – 25 ng/mL , 25 ng/mL – 50 ng/mL , 25 ng/mL – 75 ng/mL , or 25 ng/mL – 100 ng/mL .

[0047] In some embodiments, the IL-7 is at a concentration of 50 ng/mL .

[0048] In some embodiments, the second set of conditions comprises a culture medium comprising Flt3L at a concentration of between 1 ng/mL – 100 ng/mL . For example, in some embodiments, the second set of conditions comprises a culture medium comprising 1 ng/mL -25 ng/mL , 25 ng/mL – 50 ng/mL , 25 ng/mL – 75 ng/mL , or 25 ng/mL – 100 ng/mL .

[0049] In some embodiments, the Flt3L at a concentration of 50 ng/mL Flt3L.

[0050] In some embodiments, the second set of conditions comprises a culture medium comprising TPO at a concentration between 1 ng/mL – 200 ng/mL . For example, in some embodiments, the second set of conditions comprises a culture medium comprising TPO at a

concentration between 1 ng/mL – 25 ng/mL, 25 ng/mL – 50 ng/mL, 25 ng/mL – 75 ng/mL, 25 ng/mL – 100 ng/mL, 100 ng/mL – 125 ng/mL, or 100 ng/mL – 200 ng/mL.

[0051] In some embodiments, the TPO is at a concentration of 100 ng/mL.

[0052] In some embodiments, the second set of conditions comprises a culture medium comprising p38 inhibitor at a concentration between 0.5 μ M – 100 μ M. For example, in some embodiments, the second set of conditions comprises a culture medium comprising p38 inhibitor at a concentration between 0.5 μ M to 25 μ M, 0.5 μ M to 50 μ M, 0.5 μ M to 100 μ M.

[0053] In some embodiments, the p38 inhibitor is SB203580.

[0054] In some embodiments, the SB203580 is at a concentration of 15 μ M.

[0055] In some embodiments, the second set of conditions comprises a culture medium comprising SDF-1 at a concentration between 10 ng/mL – 100 ng/mL. For example, in some embodiments, the second set of conditions comprises a culture medium comprising SDF-1 at a concentration between 1 ng/mL – 25 ng/mL, 25 ng/mL – 50 ng/mL, 25 ng/mL – 75 ng/mL, or 25 ng/mL – 100 ng/mL.

[0056] In some embodiments, the SDF-1 is at a concentration of 30 nM.

[0057] In some embodiments, the third set of conditions comprises a culture medium comprising at least one compound selected from the group consisting of a CD3 activator, IL-2 and IL-7. Accordingly, in some embodiments, the third set of conditions comprises a culture medium comprising CD3 activator. In some embodiments, the third set of conditions comprises a culture medium comprising IL-2. In some embodiments, the third set of conditions comprises IL-7.

[0058] In some embodiments, the third set of conditions comprises IL-2 at a concentration of between 1 ng/mL – 100 ng/mL. For example, in some embodiments, the third set of conditions comprises IL-2 at a concentration of between 1 ng/mL – 25 ng/mL, 25 ng/mL – 50 ng/mL, 25 ng/mL – 75 ng/mL, or 25 ng/mL – 100 ng/mL.

[0059] In some embodiments, the IL-2 is at a concentration of 10 ng/mL.

[0060] In some embodiments, the third set of conditions comprises a culture medium comprising IL-7 at a concentration between 1 ng/mL – 100 ng/mL. For example, in some

embodiments, the third set of conditions comprises IL-7 at a concentration of between 1 ng/mL--25 ng/mL, 25 ng/mL -- 50 ng/mL, 25 ng/mL -- 75 ng/mL, or 25 ng/mL -- 100 ng/mL.

[0061] In some embodiments, the IL-7 is at a concentration of 10 ng/mL.

[0062] In some embodiments, each of the culturing steps is performed at about 5% oxygen.

[0063] In some embodiments, each of the culturing steps is performed at greater than 14% oxygen.

[0064] In some embodiments, each of the culturing steps is performed at atmospheric oxygen.

[0065] In some embodiments, each of the culturing steps is performed at less than 5% oxygen. For example, in some embodiments, the culturing steps is performed at less than 5% oxygen, less than 4% oxygen, less than 3% oxygen, less than 2% oxygen or less than 1% oxygen.

[0066] In some embodiments, culturing pluripotent stem cells in the first set of conditions to obtain HPC bulk lasts for greater than 10 days.

[0067] In some embodiments, culturing pluripotent stem cells in the first set of conditions to obtain HPC bulk lasts between 11 and 15 days.

[0068] In some embodiments, culturing pluripotent stem cells in the first set of conditions to obtain HPC bulk lasts for 14 days.

[0069] In some embodiments, the iPSCs are obtained from peripheral blood mononuclear cells.

[0070] In some embodiments, at least about 50%, 55%, 60%, 75%, 80%, 85%, 90%, 95%, 97% or more of cells in the produced cell population are CD56+/CD3- NK cells without a step of enrichment.

[0071] In some embodiments, less than about 25% of cells in the produced cell population are CD3+ cells.

[0072] In some embodiments, the percentage of cells in the produced cell population is determined by flow cytometry.

[0073] In some embodiments, the percentage of cells in the produced cell population is determined by single-cell RNA sequencing (scRNAseq).

[0074] In some embodiments, the method further comprises isolating CD56+/CD3- cells.

[0075] In some embodiments, the isolating comprises fluorescence-activated cell sorting (FACS) or magnetic sorting.

[0076] In some embodiments, the NK cells are genetically modified to express one or more chimeric antigen receptors (CAR).

[0077] In some embodiments, the CAR is a CD19 CAR.

[0078] In some embodiments, the NK cells are further genetically modified to express IL-15R α /IL-15 complex.

[0079] In some aspects, a method of producing induced pluripotent stem cell (iPSC)-derived NK cells is provided, the method comprising the following steps:

[0080] (1) culturing iPSC in a first set of conditions comprising a medium comprising at least one compound selected from vascular endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF), and ascorbic acid to obtain a population comprising at least 20% CD34+ HPCs (HPC bulk);

[0081] (2) culturing the HPC bulk obtained in (1) in second set of conditions comprising a culture medium comprising one or more of ascorbic acid, p38 inhibitor and SDF-1 to obtain a cell population comprising at least 5% CD4- cells; and

[0082] (3) culturing the cell population from (2) in a third set of conditions comprising a culture medium comprising at least one compound selected from the group consisting of a CD3 activator, IL-2 and IL-7.

[0083] In some aspects, a method of producing induced pluripotent stem cell (iPSC)-derived NK cells is provided, the method comprising the following steps:

[0084] (1) culturing a bulk cell population comprising hematopoietic progenitor cells (HPC bulk) in a second set of conditions comprising a culture medium comprising one or more of ascorbic acid, p38 inhibitor and SDF-1 to obtain an a cell population comprising at least 5% CD4- cells;

[0085] (2) isolating the CD4⁻ cells from (1); and

[0086] (3) culturing the isolated CD4⁻ cells in NK induction media comprising at least one compound selected from the group consisting of a CD3 activator, IL-2 and IL-7.

[0087] In some embodiments, about 15-30% of the cells in the cell population obtained in step (1) are CD4⁻ cells.

[0088] In some embodiments, about 20% of the cells in the cell population obtained in step (1) are CD4⁻ cells.

[0089] In some embodiments, the cell population comprising CD4⁻ cells comprises CD4⁻/CD8⁻ cells and CD4⁻/CD8⁺ cells.

[0090] In some aspects, an NK cell population produced using a method disclosed herein is provided.

[0091] In some embodiments, the NK cell population produced using a method disclosed herein is cryopreserved in a cryopreservation medium.

[0092] In some aspects, an unsorted cell population comprising pluripotent stem cell-derived CD56⁺/CD3⁻ cells is provided, which is present at a ratio of at least 60% of total pluripotent stem cell-derived CD56⁺ cells obtained in accordance with the methods described herein.

[0093] In some embodiments, less than 25% of cells in the unsorted cell population are CD3⁺ cells.

[0094] In some embodiments, less than 5% of cells in the unsorted cell population are monocytes.

[0095] In some embodiments, less than 5% of cells in the unsorted cell population are B cells.

[0096] In some embodiments, the percentage of cells in the unsorted cell population is determined by flow cytometry.

[0097] In some embodiments, the percentage of cells is determined by single-cell RNA sequencing (scRNAseq).

[0098] In some aspects, a method of treating a subject in need of cell therapy is provided, the method comprising administering to the subject the NK cells of any one of the preceding claims.

[0099] In some embodiments, the subject has a cancer.

[0100] In some embodiments, the cancer is leukemia or lymphoma.

[0101] In some aspects, a method of producing a cell population comprising CD34+ hematopoietic progenitor cells (HPCs) is provided, the method comprising culturing a population of pluripotent stem cells in a first set of conditions comprising a medium comprising at least one compound selected from vascular endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF), and ascorbic acid to obtain a population comprising CD34+ HPCs (HPC bulk).

[0102] In some aspects, a cell population comprising HPCs produced by the methods described herein is provided.

[0103] In some embodiments, the cell population comprises at least 20% CD34+ cells.

BRIEF DESCRIPTION OF THE DRAWINGS

[0104] **FIG. 1** is a series of flow cytometry plots that show the results of flow cytometry analysis of a NK cell bulk population that was obtained using the methods described herein. Briefly, iPS cells were cultured as described herein, without an intermediate isolation step. The flow cytometry graphs show the presence of a population of CD3 negative cells that are CD56^{high} (approximately 63.5% of the NK cell bulk population) and a population of CD3 negative cells that are CD56^{dim} (approximately 26.7% of the NK cell bulk population). Both of these populations are NK cells. Thus, the total CD56 positive, CD3 negative NK cell population obtained is greater than 90% of the total cell population.

[0105] **FIG. 2** shows the results of scRNA analysis of a NK cell bulk population that was obtained using the methods described herein. Briefly, iPSCs were cultured as described herein, without an intermediate isolation step. Cells were classified into four cell populations - monocyte, B cell, NK cell and T cell. Approximately 75% of cells in the NK cell bulk population was identified as NK cells (CD56+/CD3- cells), while approximately 25% of cells in the bulk

cell population was identified as T cells. The T cells described herein may include NKT cells (CD56+/CD3+ cells).

[0106] **FIG. 3** depicts a series of photographs that show the results of an anti-tumor activity assay in NSG (NOD/Shi-scid, IL-2R gamma null) mice administered luciferase expressing Nalm6 cells, followed by the administration of iNK-CAR19 cells in. Nalm6 cells-transplanted NSG mice were treated with (a) PBS buffer (b) iNK-CAR cells, and anti-tumor efficacy of these treatments was observed. The data shows that iNK-CAR cells reduced proliferation of Nalm6 cells.

DEFINITIONS

[0107] *Administering:* As used herein, the terms “administer,” “administering,” “administration” “introducing” or “introduction” are used interchangeably in the context of delivering therapeutic cells, for example, iPSC or HPC derived NK cells (e.g., CD56+/CD3- cells), into a subject, by a method or route which results in delivery of such cells. Various methods are known in the art for administering cells, including for example intravenously, topically, orally, intramuscularly, intraperitoneally, intrathecally, subcutaneously or transdermally. Cells can be administered with or without a carrier.

[0108] *Adoptive Cell Therapy:* As used herein interchangeably, the terms “adoptive cell therapy” or “adoptive cell transfer” or “cell therapy” or “ACT” refer to the transfer of cells, for example, a population of CD56+/CD3- cells generated using methods described herein and administered into a subject or patient in need thereof. In some embodiments, the cell is a CD56+/CD3-immune cell made using a method described herein and additionally expressing a CAR.

[0109] *Animal:* As used herein, the term “animal” refers to any member of the animal kingdom. In some embodiments, “animal” refers to humans, at any stage of development. In some embodiments, “animal” refers to non-human animals, at any stage of development. In certain embodiments, the non-human animal is a mammal (e.g., a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, a sheep, cattle, a primate, and/or a pig). In some embodiments, animals include, but are not limited to, mammals, birds, reptiles, amphibians, fish, insects, and/or worms.

In some embodiments, an animal may be a transgenic animal, genetically-engineered animal, and/or a clone.

[0110] *Antigen-Specific Targeting Domain*: “antigen-specific targeting domain” provides the CAR with the ability to bind to a target antigen of interest. In some embodiments, the antigen-specific targeting domain targets an antigen of clinical interest against which it would be desirable to trigger an effector immune response that results in tumor killing. The antigen-specific targeting domain may be any protein or peptide that possesses the ability to specifically recognize and bind to a biological molecule (e.g., a cell surface receptor or tumor protein, or a component thereof). The antigen-specific targeting domain includes any naturally occurring, synthetic, semi-synthetic, or recombinantly produced binding partner for a biological molecule of interest.

[0111] Illustrative antigen-specific targeting domains include, for example, antibodies or antibody fragments or derivatives, extracellular domains of receptors, ligands for cell surface molecules/receptors, or receptor binding domains thereof, and tumor binding proteins.

[0112] In some embodiments, the antigen-specific targeting domain is, or is derived from, an antibody. An antibody-derived targeting domain can be a fragment of an antibody or a genetically engineered product of one or more fragments of the antibody, which fragment is involved in binding with the antigen. Examples include a variable region (Fv), a complementarity determining region (CDR), a Fab, a single chain antibody (scFv), a heavy chain variable region (VH), a light chain variable region (VL) and a camelid antibody (VHH).

[0113] In some embodiments, the binding domain is a single chain antibody (scFv). The scFv may be murine, human or humanized scFv.

[0114] *Allogeneic*: “Allogeneic” as used herein, refers to any material derived from a different animal of the same species as the individual to whom the material is introduced. Two or more individuals are said to be allogeneic to one another when the genes at one or more loci are not identical. In some aspects, allogeneic material from individuals of the same species may be sufficiently different genetically to interact antigenically.

[0115] *Approximately or about*: As used herein, the term “approximately” or “about,” as applied to one or more values of interest, refers to a value that is similar to a stated reference

value. In certain embodiments, the term “approximately” or “about” refers to a range of values that fall within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value). It is understood that when the term “about” or “approximately” is used to modify a stated reference value, the stated reference value itself is covered along with values that are near the stated reference value on either side of the stated reference value.

[0116] *Ascorbic Acid or Vitamin C:* As used herein, “ascorbic acid” or “vitamin C” means L-ascorbic acid and derivatives thereof, and “L-ascorbic acid derivative” means derivatives that become vitamin C by enzymatic reaction in the living body. Examples of the derivatives of L-ascorbic acid include vitamin C phosphate, ascorbic acid glucoside, ascorbyl ethyl, vitamin C ester, ascorbyl tetrahexyldecanoate, ascorbyl stearate, and ascorbyl 2-phosphate 6-palmitate. Examples of the vitamin C phosphate include salts of L-ascorbic acid phosphate such as L-ascorbic acid phosphate Na and L-ascorbic acid phosphate Mg. In one embodiment, the vitamin C can be ascorbic acid 2-phosphate.

[0117] *Bulk cell population:* The term “bulk cell population”, “bulk cell” and the like refers to a heterogeneous population of cells. In some embodiments, the bulk cell population comprises hematopoietic cells. In some embodiments, the bulk cell population may be obtained from pluripotent cells (e.g., induced pluripotent stem cells (iPSCs)). In some embodiments, the bulk cell population is obtained from donor tissue, including, for example, blood.

[0118] *CD4- induction media or second set of conditions:* The term CD4 - induction media is used interchangeably with “second set of conditions.” These terms as used herein refers to cell culture media that is used to obtain a population of cells that comprises CD4- cells. In some embodiments, this media is used to differentiate HP cell bulk to a population of cells that comprise CD4- cells. In some embodiments, the CD4- induction media comprises one or more, or all, of ascorbic acid, stem cell factor (SCF), thrombopoietin (TPO), Flt3L, IL-7, a p38 inhibitor such as SB203580, and SDF1 such as SDF-1 α .

[0119] *Chimeric Antigen Receptor (CAR):* As used herein, the term “chimeric antigen receptor” or “CAR” engineered receptors which can confer an antigen specificity onto cells (for

example immune cells such as NK cells, T cells such as naive T cells, central memory T cells, effector memory T cells or combinations thereof). CARs are also known as artificial T-cell receptors, chimeric T-cell receptors or chimeric immunoreceptors. In some embodiments, the CARs of the invention comprise an antigen-specific targeting domain, an extracellular domain, a transmembrane domain, optionally one or more co-stimulatory domains, and an intracellular signaling domain. In some embodiments described herein, a CAR is introduced into NK cells (e.g., CD56+/CD3- cells) made using a method described herein, such that to redirect specificity for a desired cell-surface antigen or MHC-peptide complex. These synthetic receptors typically contain a target binding domain that is associated with one or more signaling domains via a flexible linker in a single fusion molecule. The target binding domain is used to direct the NK cells to specific targets on the surface of pathologic cells (e.g., a cancer cell) and the signaling domains contain molecular machinery for NK cells activation and proliferation. The flexible linker which usually passes through the NK cell membrane (i.e., forming a transmembrane domain) allows for cell membrane display of the target binding domain of the CAR. CARs have successfully allowed immune cells to be redirected against antigens expressed at the surface of tumor cells from various malignancies including lymphomas and solid tumors (Gross et al., (1989) Transplant Proc., 21(1 Pt 1): 127-30; Jena et al., (2010) Blood, 116(7):1035-44). A CAR's extracellular binding domain may be composed of a single chain variable fragment (scFv) derived from fusing the variable heavy and light regions of a murine or humanized monoclonal antibody. In some embodiments, the extracellular binding domain comprises a single domain antibody. Alternatively, scFvs may be used that are derived from Fab's (instead of from an antibody, e.g., obtained from Fab libraries). In various embodiments, this scFv is fused to a transmembrane domain and then to an intracellular signaling domain. At least three generations of CARs have been developed. The first generation CARs comprised target binding domains attached to a signaling domain derived from the cytoplasmic region of the CD3zeta or the Fc receptor gamma chains. First generation CARs were shown to successfully redirect immune cells to the selected target, but they failed to provide prolonged expansion and antitumor activity *in vivo*. Second and third generation CARs have focused on enhancing modified cell survival and increasing proliferation by including co-stimulatory molecules, such as CD28, OX-40 (CD134) and 4-1BB (CD137).

[0120] *Culture*: The term “culture” or “cell culture” or “culturing” refers to the maintenance, growth and/or differentiation of cells in an *in vitro* environment. In various methods described herein, cells are cultured in a particular cell culture medium (or “media” in case of a plural) which facilitates or promotes the growth or differentiation of one type of cell into a different type of cell. For example, in certain embodiments described herein, culturing iPSCs in a cell culture medium results in at least 20%, 30%, 40%, 50%, 60%, 70%, or 80% of cells becoming CD34+ cells (i.e., HPCs) in the total cell population. In some embodiments described herein, culturing a cell population containing at least 20%, 30%, 40%, 50%, 60%, 70%, or 80% 30% HPCs results in a cell population comprising CD4- cells. In yet other embodiments, culturing a cell population of cells containing CD4- cells results in an enriched CD56+/CD3- cell population (i.e., at least 50% of cells in the total cell population are CD56+/CD3-). A cell culture medium acts as a source of nutrients, hormones and/or other factors helpful to propagate and/or sustain the cells.

[0121] *Differentiating*: The terms “differentiating,” “inducing,” “converting,” “deriving” and the like refer to process in which a cell of one phenotype undergoes a change to a cell of another phenotype.

[0122] *Engineered*: The term “engineered”, as used herein, describes a polynucleotide, polypeptide or a cell that has been designed or modified and/or whose existence and production require intervention and/or activity. For example, an engineered cell that is intentionally designed to elicit a particular effect and that differs from the effect of naturally occurring cells of the same type. In some embodiments, an engineered cell is a CD56+/CD3- cell derived from iPSCs or HPCs using a method described herein and further expresses a chimeric antigen receptor.

[0123] *Enriched*: The term “enriched” as used herein in reference to a specific cell type means a cell population that has at least 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, or 90% of the specific cell type within the cell population as determined by flow cytometry or other analytical method.

[0124] *Ex vivo*: As used herein, the term “*ex vivo*” means a process in which cells are removed from a living organism and are propagated outside the organism (e.g., in a test tube, in a culture bag, in a bioreactor).

[0125] *Functional equivalent or derivative*: As used herein, the term “functional equivalent” or “functional derivative” denotes, in the context of a functional derivative of an amino acid sequence, a molecule that retains a biological activity (either function or structural) that is substantially similar to that of the original sequence. A functional derivative or equivalent may be a natural derivative or is prepared synthetically. Exemplary functional derivatives include amino acid sequences having substitutions, deletions, or additions of one or more amino acids, provided that the biological activity of the protein is conserved. The substituting amino acid desirably has chemico-physical properties which are similar to that of the substituted amino acid. Desirable similar chemico-physical properties include, similarities in charge, bulkiness, hydrophobicity, hydrophilicity, and the like.

[0126] *Hematopoietic progenitor cell*: The term “hematopoietic progenitor cell” or “hematopoietic progenitor cells” or “HPC” or “HPCs” refers to CD34+ cells which are committed to a hematopoietic lineage but are capable of further hematopoietic differentiation and include hematopoietic stem cells, multipotential hematopoietic stem cells, common myeloid progenitors, megakaryocyte progenitors, erythrocyte progenitors, and lymphoid progenitors.

[0127] *HPC bulk*: The terms “HPC bulk,” “Hematopoietic progenitor cell bulk,” or “HP cell bulk” mean a heterogenous population of cells comprising hematopoietic progenitor cells. In some embodiments, the HPC bulk is derived from iPSCs. In some embodiments, the HPC bulk is derived from blood.

[0128] *HPC derived NK cells*: The term “HPC derived NK cells” means CD56+/CD3- cells (e.g., NK or NK-like cells) that are obtained from an HPC bulk population following culture in a cell culture medium.

[0129] *HPC induction media and First Set of Conditions*: The term HPC induction media or “first set of conditions” as used herein refers to culture media that is used generate a population of cells comprising hematopoietic cells from a starting cell population. In some embodiments, the starting cell population comprises iPSCs. In some embodiments, the HPC induction media comprises one or more of BMP4, VEGF, bFGF, ascorbic acid, a TGFβ inhibitor, stem cell factor (SCF), thrombopoietin (TPO), and Flt3L.

[0130] *Immune cells* : As used herein, the term “immune cell” or “immune cells” refers to cells of the immune system, including, but not limited to, T cells, NK cells, T/NK cells,

dendritic cells, macrophages, B cells, neutrophils, erythrocytes, monocytes, basophils, neutrophils, mast cells, eosinophils, and any combination thereof. In various embodiments, the immune cells produced using methods described herein are NK cells, characterized as CD56+/CD3- cells.

[0131] *Induced Pluripotent Stem Cell (iPSC)*: As used herein, the term “induced pluripotent stem cell” or “iPSC” refers to a pluripotent stem cell artificially derived (e.g., induced) from a non-pluripotent cell, typically an adult somatic cell, for example, by inducing expression of one or more genes (including POU4F1/OCT4 (Gene ID; 5460) in combination with, but not restricted to, SOX2 (Gene ID; 6657), KLF4 (Gene ID; 9314), cMYC (Gene ID; 4609), NANOG (Gene ID; 79923), LIN28/ LIN28A (Gene ID; 79727)). The stem cells may be genetically modified at any stage with markers or gene so that the markers or genes are carried through to any stage of culturing. The markers may be used to purify or enrich the differentiated or undifferentiated stem cell populations at any stage of culture.

[0132] *Induction medium*: The term “induction medium” generally refers to a cell culture media that is used to differentiate a population of cells from a first cell phenotype to a second cell phenotype. In some embodiments, the first cell phenotype and the second cell phenotypes comprise a heterogeneous population of cells. In some embodiments, the first cell phenotype comprises a heterogeneous population of cells. In some embodiments, the second cell phenotype comprises a heterogeneous population of cells. In some embodiments, the induction medium is used to differentiate a population of heterogeneous cells to a population of cells that is substantially homogeneous.

[0133] *In vitro*: As used herein, the term “*in vitro*” refers to events that occur in an artificial environment, e.g., in a test tube or reaction vessel, in cell culture, *etc.*, rather than within a multi-cellular organism.

[0134] *In vivo*: As used herein, the term “*in vivo*” refers to events that occur within a multi-cellular organism, such as a human and a non-human animal. In the context of cell-based systems, the term may be used to refer to events that occur within a living cell (as opposed to, for example, *in vitro* systems).

[0135] *Isolation step*: The term “isolating”, “a step of isolating”, “isolation”, “isolation step,” or “cell isolation step” as used herein, means separating particular cell type from a mixture

of cells. In some embodiment, a cell type can be defined by one or more markers that are either present or absent on the cell surface of a particular cell type. Various methods of separating particular cell types from a mixture of cells are known in the art, and include, for example fluorescence-activated cell sorting (FACS) and magnetic-bead based sorting strategies such as magnetic-activated cell sorting (MACS).

[0136] *Natural Killer (NK) Cell*: As used herein, “natural killer cell” or NK cell is a lymphoid cell defined by its marker expression and function/activity. For example, in humans an NK cell expresses CD56 (CD56+). In a further embodiment, such NK cells may express CD56 and CD16 (CD56+/CD16+). In another example, such NK cells may express CD56 but not express CD3 (CD56+/CD3-). NK cells may express variable levels of CD56. For example, NK cells may be “CD56^{high}” which means that the NK cells express a high level of CD56 as assessed by methods in the art, for example as assessed by flow cytometry. As another example, NK cells may be “CD56^{dim}” which means that the NK cells express a low but detectable level of CD56 as assessed by methods in the art, for example as assessed by flow cytometry.

[0137] *NK induction media or third set of conditions*: In some embodiments, the term “NK induction media” or “third set of conditions” refers to a cell culture media that is used to generate a population of cells comprising CD56+/CD3- cells. In some embodiments, the NK cell induction media comprises one or more of IL-7, IL-2 and a CD3 activator (e.g., anti-CD3 antibody).

[0138] *iPS NK Cell*: As used herein, “iPS NK cells” are iPSC derived NK Cells, for example, NK cells that are derived from iPSCs as starting materials. Such iPS NK cell expresses CD56 (CD56+). In a further embodiment, such iPS NK cells may express CD56 and CD16 (CD56+/CD16+). In another example, such iPS NK cells may express CD56 and not express CD3 (CD56+/CD3-). iPS NK cells are also referred to herein as “iNK cells”.

[0139] *T cell*: As used herein “T cell” is a lymphoid cell defined by its marker expression and function/activity. For example, in humans a T cell expresses CD3. CD56+/CD3+ cells are known as NKT cell.

[0140] *Single-chain Fv antibody*” or “scFv” refers to an engineered antibody that includes a light chain variable region and a heavy chain variable region connected to one another directly or via a peptide linker sequence.

[0141] *Subject*: As used herein, the term “subject” refers to a human or any non-human animal (e.g., mouse, rat, rabbit, dog, cat, cattle, swine, sheep, horse or primate). A human includes pre- and post-natal forms. In many embodiments, a subject is a human being. A subject can be a patient, which refers to a human presenting to a medical provider for diagnosis or treatment of a disease. The term “subject” is used herein interchangeably with “individual” or “patient.” A subject can be afflicted with or is susceptible to a disease or disorder but may or may not display symptoms of the disease or disorder.

[0142] *Suffering from*: An individual who is “suffering from” a disease, disorder, and/or condition has been diagnosed with or displays one or more symptoms of the disease, disorder, and/or condition. The disease may include cancer, for example, lymphoma and leukemia.

[0143] *Therapeutically effective amount*: As used herein, the term “therapeutically effective amount” of a therapeutic agent (e.g., a cell therapy) means an amount (e.g., a certain number of cells or a cell population enriched with a certain percentage of a particular type or types of cell or cells) that is sufficient, when administered to a subject suffering from or susceptible to a disease, disorder, and/or condition, to treat, diagnose, prevent, and/or delay the onset of the symptom(s) of the disease, disorder, and/or condition. It will be appreciated by those of ordinary skill in the art that a therapeutically effective amount is typically administered via a dosing regimen comprising at least one unit dose. In some embodiments, the CD56+/CD3- cells described herein are modified to express one or more transgenes. In some embodiments, the CD56+/CD3- cells described herein are modified to express a chimeric antigen receptor, such as for example CD19. In some embodiments, between about 100 million and 900 million CD56+/CD3- cells described herein are administered to a subject in need thereof. In some embodiments, between about 100 million and 700 million CD56+/CD3- cells described herein are administered to a subject in need thereof. In some embodiments, between about 100 million and 500 million CD56+/CD3- cells described herein are administered to a subject in need thereof. In some embodiments, between about 200 million and 900 million CD56+/CD3- cells described herein are administered to a subject in need thereof. In some embodiments, between about 200 million and 700 million CD56+/CD3- cells described herein are administered to a subject in need thereof. In some embodiments, between about 200 million and 500 million CD56+/CD3- cells described herein are administered to a subject in need thereof.

[0144] *Treating*: As used herein, the term “treat,” “treatment,” or “treating” refers to any method used to partially or completely alleviate, ameliorate, relieve, inhibit, prevent, delay onset of, reduce severity of and/or reduce incidence of one or more symptoms or features of a particular disease, disorder, and/or condition. Treatment may be administered to a subject who does not exhibit signs of a disease and/or exhibits only early signs of the disease for the purpose of decreasing the risk of developing pathology associated with the disease.

[0145] The recitation of numerical ranges by endpoints herein includes all numbers and fractions subsumed within that range (e.g. 1 to 5 includes 1, 1.5, 2, 2.75, 3, 3.9, 4 and 5). It is also to be understood that all numbers and fractions thereof are presumed to be modified by the term “about.”

[0146] Various aspects of the invention are described in detail in the following sections. The use of sections is not meant to limit the invention. Each section can apply to any aspect of the invention. In this application, the use of “or” means “and/or” unless stated otherwise. As used herein, the singular forms “a”, “an”, and “the” include both singular and plural referents unless the context clearly dictates otherwise.

[0147] Various aspects of the invention are described in detail in the following sections. The use of sections is not meant to limit the invention. Each section can apply to any aspect of the invention. In this application, the use of “or” means “and/or” unless stated otherwise.

DETAILED DESCRIPTION

[0148] The present disclosure provides methods of producing a cell population enriched in NK cells from a pluripotent cell, such as an iPS cell or hematopoietic progenitor cell (HPC) bulk. The NK cells produced from iPSC are referred to herein as iPS derived NK cells, iPS NK cells or iNK cells. The disclosure provides cell culturing methods in which pluripotent cells, such as iPSCs or progenitor cells such as HPCs can be differentiated with high efficiency into NK cells or iPS NK cells without the need for an isolation step. In some embodiments, the methods described herein includes an isolation step. In the isolation step, cells that are CD4⁻ (e.g., population of cells other than CD4⁺ cells such as containing CD4⁻/CD8⁺ and/or CD4⁻/CD8⁻) may be isolated, which are subsequently differentiated into NK cells. The disclosure also

demonstrates that the iPS NK cells obtained using methods described herein are functional, and can be further genetically modified, for example through the introduction of a chimeric antigen receptor (CAR), useful for the treatment of various diseases or disorders, such as cancer.

[0149] Various methods of producing CD56+/CD3- cells (NK cells) are described below in more detail.

Method of Producing NK or iPSC Derived NK Cells

Overview of Culture Method

[0150] In some embodiments, a method of producing a cell population enriched in NK cells is provided, the method comprising the steps of: (A) culturing a population of pluripotent stem cells under a first set of conditions resulting in a cell population comprising at least 20% CD34+ HPCs (HPC bulk); and (B) changing the first set of conditions to a second set of conditions, thereby resulting in a cell population comprising at least 5% CD4- cells; and (C) changing the second set of conditions to a third set of conditions, thereby resulting in a cell population enriched in NK cells.

[0151] In some embodiments, the method is performed without an isolation step. In this scenario, there is no isolation of CD4- cells after step (B). For example, in some embodiments, a method of producing a cell population enriched in NK cells is provided, the method comprising the steps of: (A) culturing a population of pluripotent stem cells under a first set of conditions resulting in a cell population comprising at least 20% CD34+ HPCs (HPC bulk); and (B) changing the first set of conditions to a second set of conditions, thereby resulting in a cell population comprising at least 5% CD4- cells; and (C) changing the second set of conditions to a third set of conditions, thereby resulting in a cell population enriched in NK cells, wherein the method is performed without an isolation step.

[0152] It is noted that the cell population that comprises CD4- cells may also comprise CD4+ cells. For example, in some embodiments, the cell population comprising at least 5% CD4- cells also comprises at least 5% CD4+ cells.

[0153] The methods described herein, either with or without an isolation step, result in an enriched population of NK cells. In some embodiments, the population of enriched cells

comprises at least 30% NK cells, at least 40% NK cells, or at least 50% NK cells. In some embodiments, the NK cells are CD56+/CD3-.

[0154] In some embodiments, the resulting cell population from step (B) comprises a population of CD4- cells. For example, in some embodiments, the resulting cell population from step (B) comprises at least 10%, 15%, or 20% CD4- cells. In some embodiments, the resulting population from step (B) comprises CD8+ cells. In some embodiments, the CD4- cells are CD8+. The resulting cell population from step (B) is between about 20 to 55% CD4-/CD8+.

[0155] In some embodiments, the CD4- cells are CD8-. In some embodiments, the resulting cell population from step (B) is between about 20 to 55% be CD4-/CD8-.

[0156] In some embodiments, the pluripotent stem cells in the method is obtained from induced pluripotent stem cells (iPSC) or embryonic stem cells (ESC).

[0157] The first set of conditions to obtain HPC bulk comprises a culture medium comprising at least one compound selected from bone morphogenetic protein-4 (BMP4) vascular endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF), ascorbic acid, Flt3 ligand (Flt3L), thrombopoietin (TPO) and TGFβ inhibitor.

[0158] The second set of conditions to obtain a population comprising CD4- cells comprises a culture medium comprising at least one compound selected from the group consisting of ascorbic acid, stem cell factor (SCF), IL-7, Flt3L, thrombopoietin (TPO), p38 inhibitor and SDF-1.

[0159] Following step (B) the method may proceed with either a cell isolation step or without a cell isolation step. Accordingly, in some embodiments, the method comprises one or more cell isolation steps. For example, in some embodiments, a population comprising CD4- cells is isolated following step (B). The isolated CD4- cells from step (B) are subsequently placed in a third set of conditions to produce a cell population enriched in NK cells.

[0160] During method of producing NK cells, the cell phenotype present in each of the steps of the method can be determined. The cell differentiation process can be assessed by various means known in the art. For example, in some embodiments, the cell differentiation of cultured cells can be assessed by obtaining a sample of the cultured cells and subjecting that sample of cultured cells to one or more analytical methods to ascertain the cell phenotype of the

cell. Known methods of ascertaining cell phenotype include for example, flow cytometry and immunofluorescence imaging. Any suitable sampling and phenotyping assay can be used with cell culture methods described herein to ascertain the progress of the cell differentiation process. In some embodiments, the methods described herein include one or more sampling steps to determine cell phenotype at a given time.

Overview of iPSC to HPC Bulk

[0161] In some embodiments, a pluripotent cell, such as an iPSC is cultured in a first set of conditions to produce a population comprising HPC bulk. In some embodiments, culturing of iPSC in a first set of conditions to obtain HPC bulk comprises culturing the iPSC in the HPC induction media. The first set of conditions can comprise various components which allow for the production of HP cell bulk. In some embodiments, the first set of conditions comprises at least one or more compounds selected from bone morphogenetic protein-4 (BMP4), vascular endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF), ascorbic acid, ROCK inhibitor, GSK3 inhibitor, stem cell factor (SCF), thrombopoietin (TPO), Flt3L, and TGF β inhibitor. Accordingly, in some embodiments, the first set of conditions to induce HPC from iPSC comprises BMP4. In some embodiments, the first set of conditions to induce HPC from iPSC comprises VEGF. In some embodiments, the first set of conditions to induce HPC from iPSC comprises bFGF. In some embodiments, the first set of conditions to induce HPC from iPSC comprises TGF β .

[0162] The iPSC are cultured in the first set of conditions period for a period of time sufficient to produce HPC bulk. In some embodiments, the iPSC are cultured in the first set of conditions for a period of between about 7-21 days, or 10-18 days, or around 14 days. In some embodiments, the iPSC are cultured in the first set of conditions for a period of about 13 days.

[0163] In some embodiments, the cells are cultured in low-oxygen conditions such as, for example, about 3%, 4%, 5%, or 6% O₂. Accordingly, in some embodiments, the cells are cultured in about 3% O₂. In some embodiments, the cells are cultured in about 4% O₂. In some embodiments, the cells are cultured in about 5% O₂. In some embodiments, the cells are cultured in about 6% O₂.

Overview of Obtaining CD4⁺ Cell Population from HPC Bulk

[0164] In some embodiments, the HPC bulk, or HPC obtained otherwise, e.g., a primary HPC isolated from a human donor or HPC differentiated from other stem cell sources/types such as embryonic stem cell derived, are further cultured without any intervening isolation step in a medium to obtain a population of cells comprising CD4⁻ cells (CD4⁻ cell population). It should be understood that the cell population obtained by this method after step (B) includes CD4⁻ cells, including for example CD4⁻/CD8⁻ cells, and CD4⁻/CD8⁺ cells. The population obtained after step (B) may also include CD4⁺ cells, including CD4⁺/CD8⁻ cells, and CD4⁺/CD8⁺ cells. The cell population includes, for example, lymphocytes. In some embodiments, the HPC bulk are cultured in a second set of conditions comprising media having at least one compound selected from the group consisting of ascorbic acid, stem cell factor (SCF), IL-7, Flt3L, TPO, fibronectin or a variant thereof, Notch ligand (e.g., Jag-1, Jag-2, DLL-1, DLL-3, DLL-4), p38 inhibitor and SDF-1 to obtain a population comprising CD4⁻ cells. Accordingly, in some embodiments, HPC bulk are cultured in a second set of conditions comprising ascorbic acid to obtain a population comprising CD4⁻ cells. In some embodiments, the HPC bulk are cultured in a second set of conditions comprising SCF to obtain a population comprising CD4⁻ cells. In some embodiments, the HPC bulk are cultured in a second set of conditions comprising IL-7 to obtain a population comprising CD4⁻ cells. In some embodiments, the HPC bulk are cultured in a second set of conditions comprising Flt3L to obtain a population comprising CD4⁻ cells. In some embodiments, the HPC bulk are cultured in a second set of conditions comprising p38 inhibitor to obtain a population comprising CD4⁻ cells. In some embodiments, the HPC bulk are cultured in a second set of conditions comprising SDF-1 to obtain a population comprising CD4⁻ cells.

[0165] The culture period is for a time suitable to obtain a population of cells comprising CD4⁻ cells. In some embodiments, the culturing of HPC bulk to obtain a population of cells comprising CD4⁻ cells is about 2-6 weeks; 3-5 weeks or around 3 weeks. Accordingly, in some embodiments, the cell culture period is for about 2-6 weeks. In some embodiments, the cell culture period is for about 3-5 weeks. In some embodiments, the cell culture period is for about 3 weeks.

[0166] In some embodiments, the second set of conditions are changed to the third set of conditions when the percentage of CD4⁻ cells in the resulting cell population is at least about

5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90% or more than 90%.

[0167] In some embodiments, the second set of conditions are changed to the third set of conditions when the percentage of CD4⁺ cells in the resulting cell population is at least about 1%, 5%, 10%, 15%, 20% or more than 20%.

[0168] In some embodiments, the second set of conditions are changed to the third set of conditions when the percentage of CD4⁻/CD8⁺ cells in the resulting cell population is at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55% or more than 55%.

[0169] In some embodiments, the second set of conditions are changed to the third set of conditions when the percentage of CD4⁻/CD8⁻ cells in the resulting cell population is at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55% or more than 55%.

[0170] In some embodiments, the second set of conditions are changed to the third set of conditions when the percentage of CD4⁺/CD8⁻ cells in the resulting cell population is at least about 1%, 3%, 5%, 10%, 15% or more than 15%.

[0171] In some embodiments, the second set of conditions are changed to the third set of conditions when the percentage of CD4⁺/CD8⁺ cells in the resulting cell population is at least about 1%, 3%, 5%, 10% or more than 10%.

[0172] The percentage of CD4⁻ cells, CD4⁺ cells, CD4⁻/CD8⁺ cells, CD4⁻/CD8⁻ cells, CD4⁺/CD8⁻ cells and CD4⁺/CD8⁺ cells can be readily determined by one of ordinary skill in the art in or after step (B) by known methods in the art, such as flow cytometry.

[0173] In some embodiments, the cells are cultured in low-oxygen conditions such as, for example, about 3%, 4%, 5%, or 6% O₂. Accordingly, in some embodiments, the cells are cultured in low-oxygen conditions such as, for example, about 3% O₂. In some embodiments, the cells are cultured in low-oxygen conditions such as, for example, about 4% O₂. In some embodiments, the cells are cultured in low-oxygen conditions such as, for example, about 5% O₂. In some embodiments, the cells are cultured in low-oxygen conditions such as, for example, about 6% O₂.

Overview of CD4-cells to CD56⁺/CD3⁻ cells

[0174] The population of cells comprising CD4⁺ cells as described above, can be further cultured in a cell culture medium to obtain a population of NK cells. In some embodiments, a cell population comprising CD4⁺ cells are isolated from the method following step (B). Any form of isolation can be used with these methods, including florescent activated cell sorting (FACS) and/or magnetic based cell sorting (MACS). In some embodiments, the method does not comprise a cell isolation step. In some embodiments, the culture medium is to obtain NK cells comprises a third set of conditions. In some embodiments, the third set of conditions comprises one or more of IL-7, IL-2 and a CD3 activator, e.g., anti-CD3 antibody.

[0175] In some embodiments, the population of cells comprising CD4⁺ cells are further cultured, without an intervening isolation step, in a culture medium to obtain a population of cells enriched in CD56⁺/CD3⁺ cells. Such CD56⁺ cells include, for example, natural killer (NK) cells. In some embodiments the population comprising CD4⁺ cells are cultured in a medium that includes a CD3 activator, IL-2 and/or IL-7. Accordingly, in some embodiments, the population comprising CD4⁺ cells are cultured in a medium that includes a CD3 activator. CD3 activators are known in the art and include, for example, antibody complexes that bind CD3 and/or CD28 surface ligands. CD3 activators include, for example, an anti-CD3 antibody or a fragment bonded thereto can be used. In some embodiments, when an anti-CD3 antibody is used, the anti-CD3 antibody can be a polyclonal antibody or a monoclonal antibody. In some embodiments, the anti-CD3 antibody is a polyclonal antibody. In some embodiments, the anti-CD3 antibody is a monoclonal antibody. The antibody may belong to any immunoglobulin class of IgG, IgA, IgM, IgD, IgE, or IgG. Various kinds of anti-CD3 antibody can be used, including, for example, an antibody produced from OKT3 clone or UCHL1 clone can be used. The concentration of anti-CD3 antibody in the medium is, for example, between 10 ng/ml – 1000 ng/ml.

[0176] In some embodiments, the population comprising CD4⁺ cells are cultured in a medium that includes IL-2. In some embodiments, the population comprising CD4⁺ cells are cultured in a medium that includes IL-7. In some embodiments, the cells are cultured in low-oxygen conditions such as, for example, about 3%, 4%, 5%, or 6% O₂. Accordingly, in some embodiments, the cells are cultured in about 3% O₂. In some embodiments, the cells are cultured in about 4% O₂. In some embodiments, the cells are cultured in about 5% O₂. In some embodiments, the cells are cultured in about 6% O₂.

[0177] The cell population comprising CD56+ cells can be further cultured in a third set of conditions to enrich the population of CD56+/CD3- cells. In some embodiments, the cell population comprising CD56+ cells are cultured in a third set of conditions comprising IL-7 and/or IL-15 to further enrich the CD56+ cells. Thus, in some embodiments, the cell population comprising CD56+ cells are cultured in a third set of conditions comprising IL-7. In some embodiments, the cell population comprising CD56+ cells are cultured in a third set of conditions comprising IL-15. In some embodiments, the cells are cultured in low-oxygen conditions such as, for example, about 3%, 4%, 5%, or 6% O₂. In some embodiments, the CD56+/CD3- cells produced are NK cells. The percentage of CD56+/CD3- NK cells in the cell population produced is at least about 50%, 55%, 60%, 75%, 80%, 85%, 90%, 95% or more than 95%. Accordingly, in some embodiments, the method results in at least about 50% CD56+/CD3- NK cells. In some embodiments, the method results in at least about 55% CD56+/CD3- NK cells. In some embodiments, the method results in at least about 60% CD56+/CD3- NK cells. In some embodiments, the method results in at least about 65% CD56+/CD3- NK cells. In some embodiments, the method results in at least about 70% CD56+/CD3- NK cells. In some embodiments, the method results in at least about 75% CD56+/CD3- NK cells. In some embodiments, the method results in at least about 80% CD56+/CD3- NK cells. In some embodiments, the method results in at least about 85% CD56+/CD3- NK cells. In some embodiments, the method results in at least about 90% CD56+/CD3- NK cells. In some embodiments, the method results in at least about 95% CD56+/CD3- NK cells. In some embodiments, the method results in more than 95% CD56+/CD3- NK cells.

[0178] In some embodiments, an enriched NK cell population obtained using methods described herein may also contain CD3+ T cells. In some embodiments, about 5%, 10%, 15%, 20%, 25%, or about 30% of the cells produced by this culturing method are CD3+ T cells. In some embodiments, less than 5% of the cells produced are CD3+ T cells. In some embodiments, about 5% of the cells produced are CD3+ T cells. In some embodiments, about 10% of the cells produced are CD3+ T cells. In some embodiments, about 15% of the cells produced are CD3+ T cells. In some embodiments, about 20% of the cells produced are CD3+ T cells. In some embodiments, about 25% of the cells produced are CD3+ T cells. In some embodiments, about 30% of the cells produced are CD3+ T cells.

[0179] In some embodiments, the population enriched in CD56+ /CD3- cells are isolated by methods known in the art. Such methods include, for example, flow cytometry (FACS)-based sorting methods and magnetic-based sorting methods (MACS).

[0180] In some embodiments, the culturing period to obtain CD56+/CD3- NK cells from a population comprising CD4- cells is about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 or 18 days. In some embodiments, the culturing period is about 7 days.

[0181] In some embodiments, the enriched CD56+ cell population comprise NK cells. In some embodiments, the enriched CD56+ cells comprise CD3- cells. In some embodiments, the enriched CD56+ cells comprise CD3+ cells.

Suitable Pluripotent Cells for Differentiation to CD56 +/CD3- Cells

[0182] In some embodiments, any pluripotent, multipotent, or donor-derived HPC can be used with the methods described herein. For example, in some embodiments, the cell is an embryonic stem cell. In some embodiments, the cell is an adult stem cell. Various adult stem cells are known in the art, and include for example, mesenchymal stem cells, hematopoietic stem cells, umbilical-cord derived cells, bone marrow stem cells, adipose stem cells and the like. In some embodiments, the cell is an induced-pluripotent stem cell (iPSC). Thus, the NK cells produced in accordance with the methods described herein can be made from any pluripotent, multipotent or patient-derived HPC, such as a primary HPC derived directly from a donor.

[0183] In some embodiments, the pluripotent cell includes, for example, embryonic stem (ES) cells, embryonic stem cells derived from a cloned embryo obtained by nuclear transfer (ntES cells), germline stem cells ("GS cells"), embryonic germ cells ("EG cells"), iPS cells, and pluripotent cells derived from cultured fibroblasts or bone marrow stem cells (Muse cells). In some embodiments, the iPS cells can be derived from peripheral blood mononuclear cells of healthy individuals. Methods for producing iPS cells are known in the art. These cells can be produced by introducing reprogramming factors into arbitrary somatic cells. Examples of the reprogramming factors herein include genes such as Oct3/4, Sox2, Sox1, Sox3, Sox15, Sox1 7, Klf 4, Klf2, c-Myc, N-Myc, L-Myc, Nanog, Lin28, Fbx15, ERas, ECAT15-2, Tel1, beta-catenin, Lin28b, Sall1, Sall4, Esrrb, Nr5a2, Tbx3, and Glis1, and gene products thereof. These

reprogramming factors may be used individually, or two or more of these may be used in combination. Examples of the combination of the reprogramming factors include those described in WO2007/069666, WO2008/118820, WO2009/007852, WO2009/032194, WO2009/058413, WO2009/057831, WO2009/075119, WO2009/079007, WO2009/091659, WO2009/101084, WO2009/101407, WO2009/102983, WO2009/114949, WO2009/117439, WO2009/126250, WO2009/126251, WO2009/126655, WO2009/157593, WO2010/009015, WO2010/033906, WO2010/033920, WO2010/042800, WO2010/050626, WO2010/056831, WO2010/068955, WO2010/098419, WO2010/102267, WO2010/111409, WO2010/111422, WO2010/115050, WO2010/124290, WO2010/147395, WO2010/147612, Huangfu D, et al. (2008), Nat. Biotechnol., 26: 795-797, Shi Y, et al. (2008), Cell Stem Cell, 2: 525-528, Eminli S, et al. (2008), Stem Cells. 26: 2467-2474, Eluangfu D, et al. (2008), Nat. Biotechnol. 26: 1269-1275, Shi Y, et al. (2008), Cell Stem Cell, 3, 568-574, Zhao Y, et al. (2008), Cell Stem Cell, 3: 475-479, Marson A, (2008), Cell Stem Cell, 3, 132-135, Feng B, et al. (2009), Nat. Cell Biol. 11: 197-203, R. L. Judson et al., (2009), Nat. Biotechnol., 27: 459-461, Lyssiotis CA, et al. (2009), Proc Natl Acad Sci US A. 106: 8912-8917, Kim J, et al. (2009), Nature. 461: 649-643, Ichida J K, et al. (2009), Cell Stem Cell. 5: 491-503, Heng JC, et al. (2010), Cell Stem Cell. 6: 167-74, Han J, et al. (2010), Nature. 463: 1096-100, Mali P, et al. (2010), Stem Cells. 28: 713-720, and Maekawa M, et al. (2011), Nature. 474: 225-9.

[0184] The iPSC can be obtained from any suitable tissue. In some embodiments, the iPSC are obtained from peripheral blood mononuclear cells.

[0185] Hematopoietic progenitor cells (HPC) are cells that are capable of differentiation into blood cells such as lymphocytes, eosinophils, neutrophils, basophils, erythrocytes, and megakaryocytes. Hematopoietic progenitor cells or hematopoietic stem cells can be identified based on, for example, the presence of CD34 and/or CD43 surface antigens.

[0186] In some embodiments, the cells that are used to produce the CD56+/CD3- NK cells described herein are genetically modified at any stage of cellular differentiation. In some embodiments, the CD56+/CD3- NK cells are genetically modified to include a desired chimeric antigen receptor (CAR), exogenous T-cell receptor (TCR) or other engineered protein.

[0187] In some embodiments, the cells used to produce the CD56+/CD3- NK cell described herein are genetically modified at a pluripotent, multipotent or unipotent stage. For

example, in some embodiments, the cells used to produce the CD56+/CD3- NK cell described herein are genetically modified at a pluripotent stage. For example, the cell can be genetically modified at an embryonic stem cell stage or iPSC stem cell stage. In some embodiments, the cells used to produce the CD56+/CD3- NK cell described herein are genetically modified at a multipotent stage. For example, the cell can be genetically modified at the hematopoietic stem cell (HSC) stage.

Culture Conditions – iPSC to HPC Bulk

[0188] In some embodiments, the medium for the production of the hematopoietic progenitor cells (i.e., HPC induction media) from iPSC may be prepared by adding vitamin C to a basal medium which is used for culture of animal cells. Examples of the basal medium include Iscove's Modified Dulbecco's Medium (IMDM), Medium 199, Eagle's Minimum Essential Medium (EMEM), α MEM medium, Dulbecco's modified Eagle's Medium (DMEM), Ham's F12 medium, RPMI 1640 medium, Fischer's medium, and Neurobasal Medium (Life Technologies), and mixtures of two or more of these media. In some embodiments, medium contains serum. In some embodiments, the medium is serum-free.

[0189] In some embodiments, the first set of conditions may include StemPro™-34. StemPro™-34 is a serum-free medium that is formulated to support the development of human hematopoietic cells in culture.

[0190] If necessary, in some embodiments, the first set of conditions may also contain one or more substances such as albumin, human insulin, human transferrin, selenium or sodium selenite, fatty acid, trace elements, 2-mercaptoethanol, thiol glycerol, α -monothioglycerol, lipids, amino acids, L-glutamine, non-essential amino acids, vitamins, growth factors, low molecular weight compounds, antibiotics, antioxidants, pyruvic acid, buffers, inorganic salts, and cytokines.

[0191] In some embodiments, the first set of conditions includes IMDM medium containing serum, insulin, transferrin, selenium, thiol glycerol or α -monothioglycerol, L-glutamine, and ascorbic acid.

[0192] In some embodiments, the first set of conditions contains one or more substances that cause signal transduction of bone morphogenetic protein 4 (BMP4) signaling pathway. Such substances include, but not limited to, BMP4.

[0193] In some embodiments, the first set of conditions contains one or more substances that cause signal transduction of vascular epithelial growth factor (VEGF) signaling pathway. Such substances include, but not limited to, VEGF.

[0194] In some embodiments, the first set of conditions contains one or more substances that cause signal transduction of fibroblast growth factor (FGF) pathway. Such substances include, but not limited to, bFGF and FGF2.

[0195] In some embodiments, the first set of conditions contains one or more substances that cause signal transduction of stem cell factor/kit signaling pathway. Such substances include, but not limited to, SCF.

[0196] In some embodiments, the first set of conditions contains one or more substances that cause signal transduction of Flt3 ligand signaling pathway. Such substances include, but not limited to, Flt3 ligand (Flt3L).

[0197] In some embodiments, the first set of conditions contains one or more substances that cause signal transduction of thrombopoietin signaling pathway. Such substances include, but not limited to, TPO.

[0198] The TGF β inhibitor is a small molecule inhibitor that interferes with the signal transduction of TGF β family and includes, for example, SB431542, SB202190 (both R.K. Lindemann et al., Mol. Cancer 2:20 (2003)), SB505124 (GlaxoSmithKline), NPC30345, SD093, SD908, SD208 (Scios), LY2109761, LY364947, LY580276 (Lilly Research Laboratories) and the like.

[0199] SB431542 is a potent and specific inhibitor of transforming growth factor-beta (TGF β) superfamily type I activin receptor-like kinase (ALK) receptors ALK4, ALK5, and ALK7.

[0200] For example, when the TGF β inhibitor is SB431542, its concentration in the medium is preferably 0.5 μ M - 100 μ M

[0201] The first set of conditions for the production of the HP cell bulk population may be further supplemented with a cytokine(s) selected from the group consisting of BMP4 (Bone morphogenetic protein 4), VEGF (vascular endothelial growth factor), bFGF (basic fibroblast growth factor), SCF (stem cell factor), TPO (thrombopoietin), and Flt3L (Flt3 ligand).

[0202] In one embodiment, the first set of conditions may include StemPro34 supplemented with human insulin (about 10 μ g/ml), human transferrin (about 5.5 μ g/ml), sodium

selenite (about 6.7 ng/ml), L-glutamine (about 2 mM), α -monothioglycerol (about 0.4 mM), and SB431542 (about 6 μ M).

[0203] In some embodiments, the vitamin C can be added every four days, every three days, every two days, or every day during the culture period. The addition of the vitamin C to the medium can be carried out at an amount corresponding to about 5 μ g/ml to about 500 μ g/ml. In some embodiments, vitamin C is present in the medium at about 5 μ g/ml, 10 μ g/ml, 25 μ g/ml, 50 μ g/ml, 100 μ g/ml, 200 μ g/ml, 300 μ g/ml, 400 μ g/ml, or 500 μ g/ml.

[0204] In the specification, "vitamin C" means L-ascorbic acid and derivatives thereof, and "L-ascorbic acid derivative" means derivatives that become vitamin C by enzymatic reaction in the living body. Examples of the derivatives of L-ascorbic acid include vitamin C phosphate (e.g., ascorbic acid 2-phosphate), ascorbic acid glucoside, ascorbyl ethyl, vitamin C ester, ascorbyl tetrahexyldecanoate, ascorbyl stearate, and ascorbyl 2-phosphate 6-palmitate. Preferred is vitamin C phosphate. Examples of the vitamin C phosphate (e.g., ascorbic acid 2-phosphate) include salts of L-ascorbic acid phosphate such as L-ascorbic acid phosphate Na and L-ascorbic acid phosphate Mg.

[0205] In some embodiments, when the substances that cause signal transduction of bone morphogenetic protein 4 (BMP4) signaling pathway is BMP4, the concentration of the BMP4 in the HPC induction media for the production of the hematopoietic progenitor cells is about 5 ng/mL to 500 ng/mL, for example, 5 ng/mL, 10 ng/mL, 20 ng/mL, 30 ng/mL, 40 ng/mL, 50 ng/mL, 60 ng/mL, 70 ng/mL, 80 ng/mL, 90 ng/mL, 100 ng/mL, 150 ng/mL, 200 ng/mL, 250 ng/mL, 300 ng/mL, 350 ng/mL, 400 ng/mL, 450 ng/mL or 500 ng/mL.

[0206] In some embodiments, when the substances that cause signal transduction of vascular epithelial growth factor (VEGF) signaling pathway is VEGF, the concentration of the VEGF in the HPC induction media for the production of the hematopoietic progenitor cells is about 5 ng/mL to 500 ng/mL, for example, about 5 ng/mL 10 ng/mL, 20 ng/mL, 30 ng/mL, 40 ng/mL, 50 ng/mL, 60 ng/mL, 70 ng/mL, 80 ng/mL, 90 ng/mL, 100 ng/mL, 150 ng/mL, 200 ng/mL, 250 ng/mL, 300 ng/mL, 350 ng/mL, 400 ng/mL, 450 ng/mL or 500 ng/mL.

[0207] In some embodiments, when substances that cause signal transduction of fibroblast growth factor (FGF) pathway is bFGF, the concentration of the bFGF in the HPC induction media for the production of the hematopoietic progenitor cells is about 5 ng/mL to 500 ng/mL, for example, 5 ng/mL, 10 ng/mL, 20 ng/mL, 30 ng/mL, 40 ng/mL, 50 ng/mL, 60 ng/mL, 70 ng/mL,

80 ng/mL, 90 ng/mL, 100 ng/mL, 150 ng/mL, 200 ng/mL, 250 ng/mL, 300 ng/mL, 350 ng/mL, 400 ng/mL, 450 ng/mL or 500 ng/mL.

[0208] In some embodiments, when the substances that cause signal transduction of stem cell factor/kit signaling pathway is SCF, the concentration of the SCF in the HPC induction media for the production of the hematopoietic progenitor cells is about 5 ng/mL to 100 ng/mL, for example, 5 ng/mL, 10 ng/mL, 20 ng/mL, 30 ng/mL, 40 ng/mL, 50 ng/mL, 60 ng/mL, 70 ng/mL, 80 ng/mL, 90 ng/mL, or 100 ng/mL.

[0209] In some embodiments, when the substances that cause signal transduction of Flt3 ligand signaling pathway is Flt3L, the concentration of the Flt3L in the HPC induction media for the production of the hematopoietic progenitor cells is about 1 ng/mL to 100 ng/mL, for example, 1 ng/mL, 2 ng/mL, 3 ng/mL, 4 ng/mL, 5 ng/mL, 6 ng/mL, 7 ng/mL, 8 ng/mL, 9 ng/mL, 10 ng/mL, 20 ng/mL, 50 ng/mL, or 100 ng/mL.

[0210] In some embodiments, when the substances that cause signal transduction of thrombopoietin signaling pathway is TPO, the concentration of the TPO in the HPC induction media for the production of the hematopoietic progenitor cells is about 1 ng/mL to 200 ng/mL, for example, 1 ng/mL, 2 ng/mL, 3 ng/mL, 4 ng/mL, 5 ng/mL, 6 ng/mL, 7 ng/mL, 8 ng/mL, 9 ng/mL, 10 ng/mL, 20 ng/mL, 30 ng/mL, 40 ng/mL, 50 ng/mL, 60 ng/mL, 70 ng/mL, 80 ng/mL, 90 ng/mL, 100, 125 ng/mL, 150 ng/mL, 175 ng/mL, or 200 ng/mL.

[0211] In some embodiments, the pluripotent stem cells may be cultured by adherent culture or suspension culture. In cases of adherent culture, the culture may be carried out in a culture vessel coated with a coating agent, and/or may be co-cultured with other cells. Examples of other cells for the co-culture include C3H10T1/2 (Takayama N., et al. J Exp Med. 2817-2830, 2010) and stromal cells derived from a different species (Niwa A et al. J Cell Physiol. 2009 Nov; 221(2): 367-77). Examples of the coating agent include Matrigel (Nivea A, et al. PLoS One. 6(7): e22261, 2011), iMatrix 511 (Miyazaki T, et al. Nature Communication 2012;3 :1236), gelatin, collagen, elastin and the like, glucosaminoglycan and proteoglycan such as hyaluronic acid, chondroitin sulfate and the like, cell adhesion proteins such as fibronectin or a variant thereof, vitronectin, laminin and the like, and the like. Examples of the method of the suspension culture include the methods described in Chadwick et al. Blood 2003, 102: 906-15, Vijayaragavan et al. Cell Stem Cell 2009, 4: 248-62, and Saeki et al. Stem Cells 2009, 27: 59-67.

[0212] In some embodiments, the HP cell bulk can also be prepared from a net-like structure (which is also referred to as ES-sac or iPS-sac) obtained by culture of pluripotent stem cells. The "net-like structure" herein is a three-dimensional sac-shaped structure (having a space in the inside) derived from pluripotent stem cells. The structure is formed with an endothelial cell population or the like, and contains hematopoietic progenitor cells in the inside.

[0213] In some embodiments, the temperature conditions for the culture for production of the HP cell bulk is from about 37°C to about 42°C. In some embodiments, the temperature is, for example, about 37° C. to about 42° C., preferably about 37 to about 39° C. The culture period may be appropriately determined by those skilled in the art by monitoring of the number of hematopoietic progenitor cells and/or the like by obtaining a sample from the cell culture for phenotypic analysis. Various kinds of methods are known in the art related to phenotypic analysis of cell samples and include, for example, flow cytometry and immunofluorescence methods.

[0214] The culture period to obtain HP cell bulk may vary and include, for example, between 6 and 14 days. Examples of the culture period include at least 6 days, not less than 7 days, not less than 8 days, not less than 9 days, not less than 10 days, not less than 11 days, not less than 12 days, not less than 13 days, and not less than 14 days. In some embodiments, the culture period is for six days. In some embodiments, the culture period is 7 days. In some embodiments, the culture period is 8 days. In some embodiments, the culture period is 9 days. In some embodiments, the culture period is 10 days. In some embodiments, the culture period is 10 days. In some embodiments, the culture period is 11 days. In some embodiments, the culture period is 12 days. In some embodiments, the culture period is 13 days. In some embodiments, the culture period is 14 days. In some embodiments, the culture period is for greater than 14 days. The culture may be carried out under low-oxygen conditions. Examples of the low-oxygen conditions include 15%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3% or lower. Frequency of medium change can be decided by a person skilled in the art. In some embodiments, medium can be changed after every 2 days. In some other embodiments, medium can be changed after every 3 days. In some embodiments, the medium is changed every day.

[0215] In some embodiments, the culture for the production of the HP cell bulk is carried out by combining one or more of the above conditions. For example, in some embodiments, the method to obtain HP cell bulk from iPSC comprises: (i) culturing pluripotent stem cells on

C3H10T1/2 in a basal medium supplemented with vitamin C under low-oxygen conditions; and (ii) further adding VEGF, SCF, and Flt3L to the culture liquid of (1), and culturing the cells under normal oxygen conditions. The period during which the step (i) is carried out is at least not less than six days, preferably not less than seven days, more preferably seven days. The period during which the step (ii) is carried out is at least not less than six days, not less than seven days, and for seven days.

[0216] In some embodiments, the hematopoietic progenitor cells obtained in the HP cell bulk may be isolated before further use. In some other embodiments, the hematopoietic progenitor cells obtained may be used as a cell population that also contains other cell species (HP cell bulk). The HP cell bulk population is an unseparated cell preparation. In some embodiments, the method does not comprise an isolation step.

Culture Conditions – HP Cell Bulk to CD4- Cell Population

[0217] In some embodiments, a second set of conditions results in a cell population comprising CD4- cells, such as for example CD4-/CD8- cells, CD4-/CD8+ cells. In some embodiments, the resulting cell population after step (B) also comprises CD4+ cells.

[0218] In some embodiments, the cell population comprising CD4- cells can be induced to differentiate into CD56+/CD3- cells.

[0219] In some embodiments, a cell population comprising CD4- cells, among other cells, can be produced by a method that includes a step of culturing hematopoietic progenitor cells or HP cell bulk in a medium supplemented with vitamin C. The vitamin C to be added to the basal medium is the same as that in the above-described induction of HP cell bulk.

[0220] In some embodiments, medium used to produce a cell population comprising CD4- cells from the HP cell bulk is the second set of conditions. In some embodiments, the second set of conditions may be prepared by adding vitamin C to a basal medium which is used for culture of animal cells. Examples of the basal medium include Iscove's Modified Dulbecco's Medium (IMDM), Medium 199, Eagle's Minimum Essential Medium (EMEM), α MEM medium (Thermo Fisher Scientific (Gibco)), Dulbecco's modified Eagle's Medium (DMEM), Ham's F12 medium, RPMI 1640 medium, Fischer's medium, and Neurobasal Medium (Life Technologies),

and mixtures of two or more of these media. The medium may contain serum, or may be serum-free.

[0221] If necessary, the basal medium may also contain one or more of substances such as albumin, human insulin, human transferrin, selenium, sodium selenite, fatty acid, trace elements, 2-mercaptoethanol, thiol glycerol, lipids, amino acids, L-glutamine, non-essential amino acids, vitamins, growth factors, low molecular-weight compounds, antibiotics, antioxidants, pyruvic acid, buffers, inorganic salts, and cytokines.

[0222] The second set of conditions for the production of a cell population comprising CD4⁺ cells, among other cells, may be further supplemented with a cytokine(s) selected from the group consisting of ascorbic acid, SCF, IL-7, Flt3L, TPO, fibronectin or a variant thereof, Notch ligand, p38 inhibitor and SDF-1.

[0223] In some embodiments, the vitamin C can be added every four days, every three days, every two days, or every day during the culture period. The addition of the vitamin C to the medium can be carried out at an amount corresponding to about 5 µg/mL to about 500 µg/mL. In some embodiments, vitamin C is present in the medium at about 5 µg/mL, 10 µg/mL, 25 µg/mL, 50 µg/mL, 100 µg/mL, 200 µg/mL, 300 µg/mL, 400 µg/mL, or 500 µg/mL.

[0224] In some embodiments, the basal medium contains one or more substances that cause signal transduction of stem cell factor/kit signaling pathway. Such substances include, but not limited to, SCF.

[0225] In some embodiments, the basal medium contains one or more substances that cause signal transduction of Flt3 ligand signaling pathway. Such substances include, but not limited to, Flt3 ligand (Flt3L).

[0226] In some embodiments, the basal medium contains one or more substances that cause signal transduction of thrombopoietin signaling pathway. Such substances include, but not limited to, TPO.

[0227] In some embodiments, the basal medium contains one or more p38 inhibitor which is an inhibitor of p38 α and p38 β which in turn suppresses downstream activation of MAPKAP kinase-2 and heat shock protein 27. Examples of the chemical inhibitor of p38 to be used in the present invention include, but are not limited to, SB203580 (4-(4-fluorophenyl)-2-(4-methylsulfonylphenyl)-5-(4-pyridyl)-1H-imidazole), and a derivative thereof, SB202190 (4-(4-fluorophenyl)-2-(4-hydroxyphenyl)-5-(4-pyridyl)-1H-imidazole) and a derivative thereof,

SB239063 (trans-4-[4-(4-fluorophenyl)-5-(2-methoxy-4-pyrimidinyl)-1H-imidazol-1-yl]cyclohexanol) and a derivative thereof, SB220025 and a derivative thereof, PD169316, RPR200765A, AMG-548, BIRB-796, SCIO-469, SCIO-323, VX-702 and FR167653. These compounds are commercially available and, for example, SB203580, SB202190, SC239063, SB220025 and PD169316 are available from Calbiochem, and SCIO-469 and SCIO-323 are available from Scios and the like. Other examples of p38 inhibitors comprise the dominant-negative mutant of p38 which includes p38T180A obtained by point mutation of the 180-position threonine located in the DNA binding region of p38 to alanine, p38Y182F obtained by point mutation of the 182-position tyrosine of p38 in human and mouse to phenylalanine and the like. The p38 inhibitor is contained in a medium at, for example, about 0.5 μ M - about 50 μ M.

[0228] In some embodiments, the basal medium contains SDF-1.

[0229] In some embodiments, SDF-1 may be not only SDF-1 α or a mature form thereof, but also an isoform such as SDF-1 β , SDF-1 γ , SDF-1 δ , SDF-1 ϵ , SDF-1 ϕ and the like or a mature form thereof, or a mixture of these at any ratio or the like. Preferably, SDF-1 α is used. SDF-1 is sometimes referred to as CXCL-12 or PBSF.

[0230] In some embodiments, one or several amino acids in the amino acid sequence of SDF-1 may be substituted, deleted and/or added as long as it has the activity as the chemokine, and similarly, the sugar chain may be substituted, deleted and/or added. An amino acid mutation is acceptable as long as at least 4 cysteine residues (Cys30, Cys32, Cys55 and Cys71 in human SDF-1 α) are maintained and not less than 90% identity with amino acid sequence of a natural substance is exhibited. SDF-1 may be obtained from a mammal, for example, human or non-human mammal such as monkey, sheep, bovine, horse, swine, dog, cat, rabbit, rat, mouse and the like. For example, the protein registered as GenBank accession number:NP_954637 can be used as human SDF-1 α , and the protein registered as GenBank accession number:NP_000600 can be used as SDF-1 β . In some embodiments, when the substances that cause signal transduction of Flt3 ligand signaling pathway is Flt3L, the concentration of the Flt3L in the medium for the production of a cell population comprising CD4⁺ cells is about 1 ng/mL to 100 ng/mL, for example, 1 ng/mL, 2 ng/mL, 3 ng/mL, 4 ng/mL, 5 ng/mL, 6 ng/mL, 7 ng/mL, 8 ng/mL, 9 ng/mL, 10 ng/mL, 20 ng/mL, 30 ng/mL, 40 ng/mL, 50 ng/mL, 60 ng/mL, 70 ng/mL, 80 ng/mL, 90 ng/mL, or 100 ng/mL. In some embodiments, when the substances that cause signal transduction

of stem cell factor/kit signaling pathway is SCF, the SCF is used for the production of cell population comprising CD4⁺ cells under the same conditions as described above.

[0231] In some embodiments, when the substances that cause signal transduction of thrombopoietin signaling pathway is TPO, the TPO is used for the production cell population comprising CD4⁺ cells under the same conditions as described above.

[0232] Fibronectin or a variant thereof to be used in the invention is not particularly limited as long as it is a molecule capable of binding to CD3 positive cells. The variant of fibronectin is not particularly limited as long as it is a molecule capable of binding to VLA-5 and VLA-4 on the surface of CD3 positive cells, and examples thereof include RetroNectin.

Fibronectin and a variant thereof may be present in any form in the medium. For example, they may be contained in the medium during culture, or may be immobilized on a culture container, and are preferably immobilized on a culture container.

[0233] When fibronectin or a variant thereof is contained in a medium, the lower limit of the concentration of fibronectin or a variant thereof may be not less than 10 ng/ml, preferably not less than 100 ng/ml, and the upper limit may be not more than 10000 µg/ml, preferably not more than 1000 µg/ml.

[0234] In some embodiments, the concentration of the IL-7 in the medium to be used for the production of cell population comprising CD4⁺ cells is about 1 ng/mL to 100 ng/ml, for example, 1 ng/mL, 2 ng/mL, 3 ng/mL, 4 ng/mL, 5 ng/mL, 6 ng/mL, 7 ng/mL, 8 ng/mL, 9 ng/mL, 10 ng/mL, 20 ng/mL, 30 ng/mL, 40 ng/mL, 50 ng/mL, 60 ng/mL, 70 ng/mL, 80 ng/mL, 90 ng/mL, or 100 ng/mL.

[0235] SDF-1 may be commercially available, purified from nature, or produced by peptide synthesis or genetic engineering techniques. SDF-1 is contained in a medium within the range of, for example, about 10 ng/mL to about 100 ng/mL. In addition, SDF-1 alternative having an SDF-1-like activity can also be used instead of SDF-1. Examples of such SDF-1 alternative include CXCR4 agonist, and a low-molecular-weight compound having a CXCR4 agonist activity and the like may be added to the medium instead of SDF-1.

[0236] In some embodiments, when the SDF-1 is SDF1α, the concentration of the SDF-1α in the medium for the production of cell population comprising CD4⁺ cells is about 1 nM to 100 nM, for example, 1 nM, 2 nM, 3 nM, 4 nM, 5 nM, 6 nM, 7 nM, 8 nM, 9 nM, 10 nM, 20 nM, 30 nM, 40 nM, 50 nM, 60 nM, 70 nM, 80 nM, 90 nM, or 100 nM.

[0237] In some embodiments, when the p38 inhibitor is SB203580, the concentration of the SB203580 in the medium for the production of cell population comprising CD4⁺ cells is about 0.5 μ M to 100 μ M, for example, 0.5 μ M, 1 μ M, 2 μ M, 3 μ M, 4 μ M, 5 μ M, 6 μ M, 7 μ M, 8 μ M, 9 μ M, 10 μ M, 15 μ M, 20 μ M, 30 μ M, 40 μ M, or 50 μ M, 60 μ M, 70 μ M, 80 μ M, 90 μ M, or 100 μ M.

[0238] In some embodiments, the basal medium for production of cell population comprising CD4⁺ cells includes α MEM medium supplemented with about 15% FBS, about 4 mM L-glutamine, about 100 U/mL penicillin, about 100 μ g/mL streptomycin, about 55 μ M 2-mercaptoethanol, about 50 μ g/mL ascorbic acid 2-phosphate, about 10 μ g/mL human insulin, about 5.5 μ g/mL human transferrin, about 6.7 ng/mL sodium selenite, about 50 ng/mL SCF, about 50 ng/mL IL-7, about 50 ng/mL Flt3L, about 100 ng/mL TPO, about 15 μ M SB203580, about 30 nM SDF-1 α .

[0239] In the production of a cell population comprising CD4⁺ cells, the hematopoietic progenitor cells may be cultured by adherent culture or suspension culture. In some embodiments, culture medium or culture vessel/dish comprises Notch activators. Notch activators, include, but are not limited to, agonists of Notch receptor. The Notch agonists bind Notch receptor, and as well, initiate or mediate the signaling events associated with the Notch receptor, such as, for example, to cause the intracellular domain of Notch to be cleaved and translocated to the nucleus. Notch activators include one or more Notch ligands, including but not limited to, Jag1, Jag2, DLL1, DLL3 and DLL4. In some embodiments, one or more of the Notch ligand(s) can be introduced as a soluble peptide, or immobilized on a solid material. The solid material may include, but not limited to, polystyrene plates, or beads. Exemplary beads for Notch ligands immobilization thereto include agarose beads, magnetic beads, and latex beads. In some embodiments, a Notch ligand peptide is conjugated/immobilized to a bead. In some other embodiments, the Notch ligand peptide is conjugated/immobilized to the surface of a polystyrene plate. In some embodiments, the immobilization of the Notch ligand is non-covalent. In some other embodiments, the Notch ligand peptide is presented by cells. In yet other embodiments, the culture vessel/dish is coated with the Notch ligand.

[0240] In some embodiments, the culture vessel/dish is coated with DLL1 or DLL4, or a fusion protein of DLL4 or DLL1, and Fc or the like. DLL1 or DLL4 may be recombinant human (rh)-DLL1 or rh-DLL4. In some embodiments, the culture vessel/dish is coated with rh-DLL4/Fc

chimera (Sino Biological) and RetroNectin (Takara Bio Inc). In cases of adherent culture, a coated culture vessel may be used, and/or the hematopoietic progenitor cells may be co-cultured with feeder cells and/or the like. Examples of the feeder cells for the co-culture include a bone-marrow stromal cell line, OP9 cells (available from Riken BioResource Center). The OP9 cells may be preferably OP-DL1 cells, which constantly express Dll1 (Holmes R I and Zuniga-Pflucker JC. Cold Spring Harb Protoc. 2009(2)). In some embodiments, where OP9 cells are used as the feeder cells, Dll1, or a fusion protein of Dll1 and Fc or the like, separately prepared may be added to the medium to perform the co-culture. In some embodiments, Dll1 can include proteins encoded by a gene having the nucleotide sequence of the NCBI accession number NM#005618 in the case of human, or NCBI accession number NM#007865 in the case of mouse; and naturally occurring mutants having a high sequence identity (for example, having a sequence identity of not less than 90%) to these proteins and having an equivalent function. In cases where feeder cells are used for production a cell population comprising CD4⁺ cells, the feeder cells can be appropriately replaced during the culture. The replacement of the feeder cells may be carried out by transferring the subject cells that are being cultured onto feeder cells that are preliminarily plated. The replacement may be carried out every five days, every four days, every three days, or every two days.

[0241] In some embodiments, the culture temperature conditions for the culture of the HP cell bulk for production of a cell population comprising CD4⁺ cells is about 37° C. to about 42° C, and about 37° C to about 39° C. In some embodiments, the culture may be carried out under low-oxygen conditions. Examples of the low-oxygen conditions include 15%, 10%, 9%, 8%, 7%, 6%, 5%, and oxygen concentrations lower than these. The culture period may be appropriately determined by those skilled in the art by monitoring of the number of different types of cells including CD4⁺ cells and/or the like. Examples of the culture period include not less than 10 days, not less than 12 days, not less than 14 days, not less than 16 days, not less than 18 days, not less than 20 days, not less than 21 days, not less than 23 days, not less than 25 days, not less than 28 days, not less than 30 days, not less than 35 days, or not less than 42 days.

[0242] In cases where a population of a particular cell type (e.g., CD4⁺ cells) from a cell population following step (B) are isolated, the isolation may be carried out using any one index including but not limited to, CD4, CD8, CD3, and CD45, depending on the cell type being isolated. The isolation method may be a method well known to those skilled in the art, for

example, a method in which the cells are labeled with a particular antibody (e.g., CD4, CD8, CD3, or CD45 antibody), and then isolated using a flow cytometer, or a method in which the cells are purified using an affinity column or the like to which a desired antigen is immobilized.

Step of Inducing CD56+/CD3- Cells from a Cell Population Comprising CD4- Cells

[0243] In some embodiments, NK cells are CD56+/CD3- cells.

[0244] In some embodiments, the NK cells are produced by a method comprising the step of culturing a cell population comprising CD4- cells in a medium supplemented with vitamin C. In some embodiments, a third set of conditions is used to produce the NK cells.

[0245] In some embodiments, the medium is prepared by adding vitamin C to a basal medium which is used for culture of animal cells. Examples of the basal medium include Iscove's Modified Dulbecco's Medium (IMDM), Medium 199, Eagle's Minimum Essential Medium (EMEM), α MEM medium, Dulbecco's modified Eagle's Medium (DMEM), Ham's F12 medium, RPMI 1640 medium, Fischer's medium, and Neurobasal Medium (Life Technologies), and mixtures of two or more of these media. In some embodiments, the medium contains serum. In some embodiments, the medium is serum-free. If necessary, the basal medium may also contain one or more of substances such as albumin, human insulin, human transferrin, selenium or sodium selenite, fatty acid, trace elements, 2-mercaptoethanol, thiol glycerol, lipids, amino acids, glutamine, non-essential amino acids, vitamins, growth factors, low-molecular-weight compounds, antibiotics, antioxidants, pyruvic acid, buffers, inorganic salts, and cytokines.

[0246] In some embodiments, the medium used for the production of the CD56 positive NK cells further contains an anti-CD3 antibody (UCHT1) and cytokines. Examples of the suitable cytokines include IL-2 and IL-7.

[0247] The CD3 antibody is not limited as long as it specifically recognizes CD3. In some embodiments, the concentration of the CD3 antibody in the third set of conditions is about 10 ng/mL to 1000 ng/mL, for example, 10 ng/mL, 50 ng/mL, 100 ng/mL, 200 ng/mL, 300 ng/mL, 400 ng/mL, 500 ng/mL, 600 ng/mL, 700 ng/mL, 800 ng/mL, 900 ng/mL, or 1000 ng/mL.

[0248] In some embodiments, vitamin C is used for the production of the CD56 positive K cells under the same conditions as described above.

[0249] In some embodiments, the concentration of the IL-2 in the third set of conditions for the production of the NK cells is about 1 U/mL to 1000 U/mL, for example, about 1 U/mL, 5

U/mL, 10 U/mL, 20 U/mL, 30 U/mL, 40 U/mL, 50 U/mL, 60 U/mL, 70 U/mL, 80 U/mL, 90 U/mL, 100 U/mL, 500 U/mL, or 1000 U/mL. In some embodiments, the concentration of the IL-2 in the NK induction media for the production of the CD56 positive NK cells is about 1 ng/mL to 100 ng/mL, for example, 1 ng/mL, 5 ng/mL, 10 ng/mL, 20 ng/mL, 30 ng/mL, 40 ng/mL, 50 ng/mL, 60 ng/mL, 70 ng/mL, 80 ng/mL, 90 ng/mL, or 100 ng/mL.

[0250] In some embodiments, the concentration of the IL-7 in the third set of conditions for the production of NK cells is about 1 ng/mL to 100 ng/mL, for example, 1 ng/mL, 5 ng/mL, 10 ng/mL, 20 ng/mL, 30 ng/mL, 40 ng/mL, 50 ng/mL, 60 ng/mL, 70 ng/mL, 80 ng/mL, 90 ng/mL, or 100 ng/mL.

[0251] In some embodiments, the culture temperature conditions for the culture of the a population comprising CD4⁻ cells for production of NK cells is about 37° C. to about 42° C., or about 37 to about 39° C. The culture period may be appropriately determined by those skilled in the art by monitoring of the number of CD56 positive NK cells and/or the like. The number of days of the culture is not limited as long as NK cells can be obtained. Examples of the culture period include at least not less than 1 day, not less than 2 days, not less than 3 days, not less than 4 days, not less than 5 days, not less than 6 days, or not less than 7 days.

[0252] In some embodiments, the NK cells obtained may be isolated before further use. In some other embodiments, the CD56 positive NK cells obtained may be used as a cell population that also contains other cell species, including T cells (NK cell bulk).

[0253] In some embodiments, the resulting cell population obtained by culturing a cell population comprising CD4⁻ cells in the third set of conditions or NK cell bulk may include at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 75%, 80%, 85%, 90%, or more than 90% NK cells. In some embodiments, NK cell bulk includes about 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, or, 50% T cells. In some embodiments, NK cell bulk includes about 75% CD56⁺/CD3⁻ NK cells and about 25% CD56⁺/CD3⁺ T cells. In some embodiments, NK cell bulk may also include B cells and monocytes.

[0254] In cases where the CD56⁺ NK cells are isolated, the isolation method may be a method well known to those skilled in the art, for example, a method in which the cells are labeled with an anti-CD56 antibody and anti-CD3 antibody, and then isolated using a flow cytometer (fluorescence-activated cell sorting), or a method in which the cells are purified using an affinity column or the like to which a desired antigen is immobilized.

[0255] In some embodiments, the CD56 positive NK cells are CD56+/CD3+. In some embodiments, the CD56 positive NK cells are CD56+/CD3-.

CAR-NK Cell Preparation

[0256] In some embodiments, the NK cell are engineered to comprise one or more transgenes. For example, the NK cells can be genetically engineered to express tumor-directed chimeric antigen receptors (CAR), thereby producing antitumor effector cells. In one example, the NK cell or iPS NK cell can be engineered to express CAR. In some embodiments, a cell precursor to the NK cell is engineered to express a CAR. For example, in some embodiments, the iPS cell is engineered to express a CAR. In some embodiments, cells in the HP cell bulk are engineered to express a CAR. In some embodiments, the NK cell is engineered to express a CAR. Moreover, in some embodiments, these transgenic receptors can be directed to tumor-associated antigens that are not protein-derived. In certain embodiments, NK cells or iPS NK cells are modified to comprise at least one CAR. In some embodiments, a single CAR targets two or more antigens.

[0257] In some embodiments, the iNK cells include a receptor that is chimeric, non-natural and engineered. In some embodiments, the engineered chimeric antigen receptor (CAR) has one, two, three, four, or more components, and in some embodiments the one or more components facilitate targeting or binding of the lymphocyte to one or more tumor antigen-comprising cancer cells.

[0258] In some embodiments, the CAR generally comprises at least one transmembrane polypeptide comprising at least one extracellular ligand-binding domain and; one transmembrane polypeptide comprising at least one intracellular signaling domain; such that the polypeptides assemble together to form a Chimeric Antigen Receptor.

[0259] The term "extracellular ligand-binding domain" as used herein is defined as an oligo- or polypeptide that is capable of binding a ligand. Preferably, the domain will be capable of interacting with a cell surface molecule. For example, the extracellular ligand-binding domain may be chosen to recognize a ligand that acts as a cell surface marker on target cells associated with a particular disease state.

[0260] In particular, the extracellular ligand-binding domain can comprise an antigen binding domain derived from an antibody against an antigen of the target.

[0261] In some embodiments, NK cells or iPS NK cells can be genetically modified to express one or more chimeric antigen receptors (CAR). In some embodiments, the CAR includes an extracellular ligand-binding domain that targets a tumor antigen selected from one or more of the following: CD44, CD19, CD20, CD22, CD23, CD30, CD89, CD123, CS-1, ROR1, mesothelin, c-Met, PSMA, Her2, GD-2, CEA, MAGE A3 TCR, EGFR, HER2/ERBB2/neu, EPCAM, EphA2, CEA, BCMA. In some embodiments, the tumor antigen is CD19.

[0262] In some embodiments, the extracellular ligand-binding domain is a single chain antibody fragment (scFv) comprising the light (VL) and the heavy (VH) variable fragment of a target antigen specific monoclonal antibody joined by a flexible linker.

[0263] In some embodiments, the CAR includes a transmembrane domain. In some embodiments, transmembrane domain further comprises a stalk region between the extracellular ligand-binding domain and the transmembrane domain. The term "stalk region" used herein generally means any oligo- or polypeptide that functions to link the transmembrane domain to the extracellular ligand-binding domain. In particular, stalk region are used to provide more flexibility and accessibility for the extracellular ligand-binding domain. A stalk region may comprise up to 300 amino acids, 10 to 100 amino acids, and/or 25 to 50 amino acids. A stalk region may be derived from all or part of naturally occurring molecules, such as from all or part of the extracellular region of CD8, CD4 or CD28, or from all or part of an antibody constant region. Alternatively the stalk region may be a synthetic sequence that corresponds to a naturally occurring stalk sequence, or may be an entirely synthetic stalk sequence. In a preferred embodiment said stalk region is a part of human CD8 alpha chain

[0264] In some embodiments, the CAR contains a signal transducing domain or intracellular signaling domain which contributes to intracellular signaling following the binding of extracellular ligand binding domain to the target resulting in the activation of the iPS NK cell and immune response. The term "signal transducing domain" refers to the portion of a protein which transduces the effector signal function signal and directs the cell to perform a specialized function. In some embodiments, iPS NK cells have a CAR that includes a signal transducing domain.

[0265] In some embodiments, the iPS NK cell are genetically modified to express IL-15R α /IL-15 complex.

[0266] In some embodiments, transmembrane polypeptides comprise the ability to be expressed at the surface of an immune cell, for example on iPS NK cells, and to interact together for directing cellular response of immune cell against a predefined target cell. The different transmembrane polypeptides of the CAR comprising an extracellular ligand-binding domain and/or a signal transducing domain interact together to take part in signal transduction following the binding with a target ligand and induce an immune response. The transmembrane domain can be derived either from a natural or from a synthetic source. The transmembrane domain can be derived from any membrane-bound or transmembrane protein.

[0267] In some embodiments, genetic modification of NK cells to express CAR can include steps of: (1) synthesizing a gene corresponding to a specific CAR; (2) preparing a vector containing the gene corresponding to the CAR; and (3) transducing the CD56+ NK cells with the vector containing the CAR gene.

[0268] Expression vectors that encode the CARs can be introduced as one or more DNA molecules or constructs, where there may be at least one marker that will allow for selection of host cells that contain the construct(s).

[0269] The constructs can be prepared in conventional ways, where the genes and regulatory regions may be isolated, as appropriate, ligated, cloned in an appropriate cloning host, analyzed by restriction or sequencing, or other convenient means. Particularly, using PCR, individual fragments including all or portions of a functional unit may be isolated, where one or more mutations may be introduced using "primer repair", ligation, in vitro mutagenesis, etc., as appropriate. The construct(s) once completed and demonstrated to have the appropriate sequences may then be introduced into the CTL by any convenient means. The constructs may be integrated and packaged into non-replicating, defective viral genomes like Adenovirus, Adeno-associated virus (AAV), or Herpes simplex virus (HSV) or others, including retroviral vectors or lentiviral vectors, for infection or transduction into cells. The constructs may include viral sequences for transfection, if desired. Alternatively, the construct may be introduced by fusion, electroporation, biolistics, transfection, lipofection, or the like. The host cells may be grown and expanded in culture before introduction of the construct(s), followed by the appropriate treatment for introduction of the construct(s) and integration of the construct(s). The cells are then expanded and screened by virtue of a marker present in the construct. Various

markers that may be used successfully include hp^{rt}, neomycin resistance, thymidine kinase, hygromycin resistance, etc.

[0270] In some instances, one may have a target site for homologous recombination, where it is desired that a construct be integrated at a particular locus. For example,) can knock-out an endogenous gene and replace it (at the same locus or elsewhere) with the gene encoded for by the construct using materials and methods as are known in the art for homologous recombination. For homologous recombination, one may use either OMEGA or O-vectors.

[0271] The constructs may be introduced as a single DNA molecule encoding at least the CAR and optionally another gene, or different DNA molecules having one or more genes. Other genes include genes that encode therapeutic molecules or suicide genes, for example. The constructs may be introduced simultaneously or consecutively, each with the same or different markers.

[0272] Vectors containing useful elements such as bacterial or yeast origins of replication, selectable and/or amplifiable markers, promoter/enhancer elements for expression in prokaryotes or eukaryotes, etc. that may be used to prepare stocks of construct DNAs and for carrying out transfections are well known in the art, and many are commercially available.

Methods of Use

[0273] The cells according to the invention can be used for treating cancer, viral infections or autoimmune disorders in a patient in need thereof. In another embodiment, said isolated cell according to the invention can be used in the manufacture of a medicament for treatment of a cancer, viral infections of autoimmune disorders, in a patient in need thereof.

[0274] The present invention relies on methods for treating patients in need thereof, said method comprising at least one of the following steps: (a) providing a chimeric antigen receptor cells according to the invention and (b) administering the cells to said patient.

[0275] Said treatment can be ameliorating, curative or prophylactic. It may be either part of an autologous immunotherapy or part of an allogenic immunotherapy treatment. By autologous, it is meant that cells, cell line or population of cells used for treating patients are originating from said patient or from a Human Leucocyte Antigen (HLA) compatible donor. By allogeneic is meant that the cells or population of cells used for treating patients are not originating from said patient but from a donor.

[0276] In some embodiments, the described cells are allogeneic. In some embodiments, the described cells are autologous.

[0277] Treatment can be used to treat patients diagnosed with cancer, viral infection, autoimmune disorders or Graft versus Host Disease (GvHD). Cancers that may be treated include tumors that are not vascularized, or not yet substantially vascularized, as well as vascularized tumors. The cancers may comprise nonsolid tumors (such as hematological tumors, for example, leukemias and lymphomas) or may comprise solid tumors. Types of cancers to be treated with the CARs of the invention include, but are not limited to, carcinoma, blastoma, and sarcoma, and certain leukemia or lymphoid malignancies, benign and malignant tumors, and malignancies e.g., sarcomas, carcinomas, and melanomas. Adult tumors/cancers and pediatric tumors/cancers are also included.

[0278] In some embodiments, treatment is in combination with one or more therapies against cancer selected from the group of antibodies therapy, chemotherapy, cytokines therapy, dendritic cell therapy, gene therapy, hormone therapy, laser light therapy and radiation therapy.

[0279] In some embodiments, said treatment can be administered into patients undergoing an immunosuppressive treatment.

[0280] In a further embodiment, the cell compositions are administered to a patient in combination with one or more additional therapies. For example, in some embodiments, the cell compositions are administered to a patient in conjunction with (e.g., before, simultaneously or following) bone marrow transplantation, T cell ablative therapy using either chemotherapy agents such as, fludarabine, external-beam radiation therapy (XRT), cyclophosphamide, or antibodies such as OKT3 or CAM PATH. In another embodiment, the cell compositions of the present invention are administered following B-cell ablative therapy such as agents that react with CD20, e.g., Rituxan. For example, in one embodiment, subjects may undergo standard treatment with high dose chemotherapy followed by peripheral blood stem cell transplantation. In certain embodiments, following the transplant, subjects receive an infusion of the expanded immune cells of the present invention. In an additional embodiment, expanded cells are administered before or following surgery. Said modified cells obtained by any one of the methods described here can be used in a particular aspect of the invention for treating patients in need thereof against Host versus Graft (HvG) rejection and Graft versus Host Disease (GvHD); therefore in the scope of the present invention is a method of treating patients in need thereof

against Host versus Graft (HvG) rejection and Graft versus Host Disease (GvHD) comprising treating said patient by administering to said patient an effective amount of modified cells comprising inactivated TCR alpha and/or TCR beta genes.

Administration of Cells

[0281] In some embodiments, the cells can be introduced into a host organism, e.g. a mammal, in a wide variety of ways. The cells may be introduced at the site of the tumor, in specific embodiments, although in alternative embodiments the cells hone to the cancer or are modified to hone to the cancer. The number of cells that are employed will depend upon a number of circumstances, the purpose for the introduction, the lifetime of the cells, the protocol to be used, for example, the number of administrations, the ability of the cells to multiply, the stability of the recombinant construct, and the like. The cells may be applied as a dispersion, generally being injected at or near the site of interest. The cells may be in a physiologically-acceptable medium. In one example, the NK cells or iPS NK cells of the present invention may express one or more CARs, TCRs, or any other engineered protein or polypeptide domain such as high affinity CD16. In some embodiments, the cells are encapsulated to inhibit immune recognition and placed at the site of the tumor.

[0282] The cells may be administered as desired. Depending upon the response desired, the manner of administration, the life of the cells, the number of cells present, various protocols may be employed. The number of administrations will depend upon the factors described above at least in part.

[0283] The administration of the cells or population of cells according to the present invention may be carried out in any convenient manner, including by aerosol inhalation, injection, ingestion, transfusion, implantation or transplantation. The compositions described herein may be administered to a patient subcutaneously, intradermally, intratumorally, intranodally, intramedullary, intramuscularly, by intravenous or intralymphatic injection, or intraperitoneally. In one embodiment, the cell compositions of the present invention are preferably administered by intravenous injection.

Nucleic Acid-Based Expression Systems

[0284] In some embodiments, the NK cells or iPS NK cells of the invention are engineered to express one or more CARs, TCRs, or any other engineered protein or polypeptide domain such as high affinity CD16 or CD19. Recombinant techniques to generate expression vectors comprising these polypeptides are well known in the art and are described generally below.

Vectors

[0285] The term "vector" is used to refer to a carrier nucleic acid molecule into which a nucleic acid sequence can be inserted for introduction into a cell where it can be replicated. A nucleic acid sequence can be "exogenous," which means that it is foreign to the cell into which the vector is being introduced or that the sequence is homologous to a sequence in the cell but in a position within the host cell nucleic acid in which the sequence is ordinarily not found. Vectors include plasmids, cosmids, viruses (bacteriophage, animal viruses, and plant viruses), and artificial chromosomes (e.g., YACs). One of skill in the art would be well equipped to construct a vector through standard recombinant techniques (see, for example, Maniatis et al., 1988 and Ausubel et al., 1994, both incorporated herein by reference).

[0286] The term "expression vector" refers to any type of genetic construct comprising a nucleic acid coding for a RNA capable of being transcribed. In some cases, RNA molecules are then translated into a protein, polypeptide, or peptide. In other cases, these sequences are not translated, for example, in the production of antisense molecules or ribozymes. Expression vectors can contain a variety of "control sequences," which refer to nucleic acid sequences necessary for the transcription and possibly translation of an operably linked coding sequence in a particular host cell. In addition to control sequences that govern transcription and translation, vectors and expression vectors may contain nucleic acid sequences that serve other functions as well and are described infra.

Promoters and Enhancers

[0287] A "promoter" is a control sequence that is a region of a nucleic acid sequence at which initiation and rate of transcription are controlled. It may contain genetic elements at which regulatory proteins and molecules may bind, such as RNA polymerase and other transcription factors, to initiate the specific transcription a nucleic acid sequence. The phrases "operatively positioned," "operatively linked," "under control," and "under transcriptional control" mean that

a promoter is in a correct functional location and/or orientation in relation to a nucleic acid sequence to control transcriptional initiation and/or expression of that sequence.

[0288] A promoter generally comprises a sequence that functions to position the start site for RNA synthesis. The best known example of this is the TATA box, but in some promoters lacking a TATA box, such as, for example, the promoter for the mammalian terminal deoxynucleotidyl transferase gene and the promoter for the SV40 late genes, a discrete element overlying the start site itself helps to fix the place of initiation. Additional promoter elements regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have been shown to contain functional elements downstream of the start site as well. To bring a coding sequence "under the control of" a promoter, one positions the 5' end of the transcription initiation site of the transcriptional reading frame "downstream" of (i.e., 3' of) the chosen promoter. The "upstream" promoter stimulates transcription of the DNA and promotes expression of the encoded RNA.

[0289] The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the tk promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either cooperatively or independently to activate transcription. A promoter may or may not be used in conjunction with an "enhancer," which refers to a cis-acting regulatory sequence involved in the transcriptional activation of a nucleic acid sequence.

[0290] A promoter may be one naturally associated with a nucleic acid sequence, as may be obtained by isolating the 5' prime' non-coding sequences located upstream of the coding segment and/or exon. Such a promoter can be referred to as "endogenous." Similarly, an enhancer may be one naturally associated with a nucleic acid sequence, located either downstream or upstream of that sequence. Alternatively, certain advantages will be gained by positioning the coding nucleic acid segment under the control of a recombinant or heterologous promoter, which refers to a promoter that is not normally associated with a nucleic acid sequence in its natural environment. A recombinant or heterologous enhancer refers also to an enhancer not normally associated with a nucleic acid sequence in its natural environment. Such promoters or enhancers may include promoters or enhancers of other genes, and promoters or enhancers isolated from any other virus, or prokaryotic or eukaryotic cell, and promoters or enhancers not

"naturally occurring," i.e., containing different elements of different transcriptional regulatory regions, and/or mutations that alter expression. For example, promoters that are most commonly used in recombinant DNA construction include the lactamase (penicillinase), lactose and tryptophan (trp) promoter systems. In addition to producing nucleic acid sequences of promoters and enhancers synthetically, sequences may be produced using recombinant cloning and/or nucleic acid amplification technology, including PCR. TM., in connection with the compositions disclosed herein (see U.S. Pat. Nos. 4,683,202 and 5,928,906, each incorporated herein by reference). Furthermore, it is contemplated the control sequences that direct transcription and/or expression of sequences within non-nuclear organelles such as mitochondria, chloroplasts, and the like, can be employed as well.

[0291] Naturally, it will be important to employ a promoter and/or enhancer that effectively directs the expression of the DNA segment in the organelle, cell type, tissue, organ, or organism chosen for expression. Those of skill in the art of molecular biology generally know the use of promoters, enhancers, and cell type combinations for protein expression, (see, for example Sambrook et al. 1989, incorporated herein by reference). The promoters employed may be constitutive, tissue-specific, inducible, and/or useful under the appropriate conditions to direct high level expression of the introduced DNA segment, such as is advantageous in the large-scale production of recombinant proteins and/or peptides. The promoter may be heterologous or endogenous.

[0292] Additionally any promoter/enhancer combination could also be used to drive expression. Use of a T3, T7 or SP6 cytoplasmic expression system is another possible embodiment. Eukaryotic cells can support cytoplasmic transcription from certain bacterial promoters if the appropriate bacterial polymerase is provided, either as part of the delivery complex or as an additional genetic expression construct.

[0293] The identity of tissue-specific promoters or elements, as well as assays to characterize their activity, is well known to those of skill in the art.

[0294] A specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon or adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals

[0295] In certain embodiments of the invention, the use of internal ribosome entry sites (IRES) elements are used to create multigene, or polycistronic, messages, and these may be used in the invention.

[0296] Vectors can include a multiple cloning site (MCS), which is a nucleic acid region that contains multiple restriction enzyme sites, any of which can be used in conjunction with standard recombinant technology to digest the vector. "Restriction enzyme digestion" refers to catalytic cleavage of a nucleic acid molecule with an enzyme that functions only at specific locations in a nucleic acid molecule. Many of these restriction enzymes are commercially available. Use of such enzymes is widely understood by those of skill in the art. Frequently, a vector is linearized or fragmented using a restriction enzyme that cuts within the MCS to enable exogenous sequences to be ligated to the vector. "Ligation" refers to the process of forming phosphodiester bonds between two nucleic acid fragments, which may or may not be contiguous with each other. Techniques involving restriction enzymes and ligation reactions are well known to those of skill in the art of recombinant technology.

[0297] Splicing sites, termination signals, origins of replication, and selectable markers may also be employed.

Plasmid Vectors

[0298] In certain embodiments, a plasmid vector is contemplated for use to transform a host cell. In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with these hosts. The vector ordinarily carries a replication site, as well as marking sequences which are capable of providing phenotypic selection in transformed cells. In a non-limiting example, E. coli is often transformed using derivatives of pBR322, a plasmid derived from an E. coli species. pBR322 contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR plasmid, or other microbial plasmid or phage must also contain, or be modified to contain, for example, promoters which can be used by the microbial organism for expression of its own proteins.

[0299] In addition, phage vectors containing replicon and control sequences that are compatible with the host microorganism can be used as transforming vectors in connection with these hosts. For example, the phage lambda GEM.TM. 11 may be utilized in making a

recombinant phage vector which can be used to transform host cells, such as, for example, *E. coli* LE392.

[0300] Further useful plasmid vectors include pIN vectors (Inouye et al., 1985); and pGEX vectors, for use in generating glutathione S transferase (GST) soluble fusion proteins for later purification and separation or cleavage. Other suitable fusion proteins are those with galactosidase, ubiquitin, and the like.

[0301] Bacterial host cells, for example, *E. coli*, comprising the expression vector, are grown in any of a number of suitable media, for example, LB. The expression of the recombinant protein in certain vectors may be induced, as would be understood by those of skill in the art, by contacting a host cell with an agent specific for certain promoters, e.g., by adding IPTG to the media or by switching incubation to a higher temperature. After culturing the bacteria for a further period, generally of between 2 and 24 h, the cells are collected by centrifugation and washed to remove residual media.

Viral Vectors

[0302] The ability of certain viruses to infect cells or enter cells via receptor mediated endocytosis, and to integrate into host cell genome and express viral genes stably and efficiently have made them attractive candidates for the transfer of foreign nucleic acids into cells (e.g., mammalian cells). Components of the present invention may be a viral vector that encodes one or more CARs, TCR, or any other engineered protein or polypeptide domain such as high affinity CD16 of the invention. Non-limiting examples of virus vectors that may be used to deliver a nucleic acid of the present invention are described below.

Adenoviral Vectors

[0303] A particular method for delivery of the nucleic acid involves the use of an adenovirus expression vector. Although adenovirus vectors are known to have a low capacity for integration into genomic DNA, this feature is counterbalanced by the high efficiency of gene transfer afforded by these vectors. "Adenovirus expression vector" is meant to include those constructs containing adenovirus sequences sufficient to (a) support packaging of the construct and (b) to ultimately express a tissue or cell specific construct that has been cloned therein. Knowledge of the genetic organization of adenovirus, a 36 kb, linear, double stranded DNA virus, allows substitution of large pieces of adenoviral DNA with foreign sequences up to 7 kb (Grunhaus and Horwitz, 1992).

AAV Vectors

[0304] The nucleic acid may be introduced into the cell using adenovirus assisted transfection. Increased transfection efficiencies have been reported in cell systems using adenovirus coupled systems (Kelleher and Vos, 1994; Cotten et al., 1992; Curiel, 1994). Adeno associated virus (AAV) is an attractive vector system for use in the cells of the present invention as it has a high frequency of integration and it can infect nondividing cells, thus making it useful for delivery of genes into mammalian cells, for example, in tissue culture (Muzyczka, 1992) or in vivo. AAV has a broad host range for infectivity (Tratschin et al., 1984; Laughlin et al., 1986; Lebkowski et al., 1988; McLaughlin et al., 1988). Details concerning the generation and use of rAAV vectors are described in U.S. Pat. Nos. 5,139,941 and 4,797,368, each incorporated herein by reference.

Retroviral Vectors

[0305] Retroviruses are useful as delivery vectors because of their ability to integrate their genes into the host genome, transferring a large amount of foreign genetic material, infecting a broad spectrum of species and cell types and of being packaged in special cell lines (Miller, 1992).

[0306] In order to construct a retroviral vector, a nucleic acid (e.g., one encoding the desired sequence) is inserted into the viral genome in the place of certain viral sequences to produce a virus that is replication defective. In order to produce virions, a packaging cell line containing the gag, pol, and env genes but without the LTR and packaging components is constructed (Mann et al., 1983). When a recombinant plasmid containing a cDNA, together with the retroviral LTR and packaging sequences is introduced into a special cell line (e.g., by calcium phosphate precipitation for example), the packaging sequence allows the RNA transcript of the recombinant plasmid to be packaged into viral particles, which are then secreted into the culture media (Nicolas and Rubenstein, 1988; Temin, 1986; Mann et al., 1983). The media containing the recombinant retroviruses is then collected, optionally concentrated, and used for gene transfer. Retroviral vectors are able to infect a broad variety of cell types. However, integration and stable expression require the division of host cells (Paskind et al., 1975).

[0307] Lentiviruses are complex retroviruses, which, in addition to the common retroviral genes gag, pol, and env, contain other genes with regulatory or structural function. Lentiviral vectors are well known in the art (see, for example, Naldini et al., 1996; Zufferey et

al., 1997; Blomer et al., 1997; U.S. Pat. Nos. 6,013,516 and 5,994,136). Some examples of lentivirus include the Human Immunodeficiency Viruses: HIV-1, HIV-2 and the Simian Immunodeficiency Virus: SIV. Lentiviral vectors have been generated by multiply attenuating the HIV virulence genes, for example, the genes *env*, *vif*, *vpr*, *vpu* and *nef* are deleted making the vector biologically safe.

[0308] Recombinant lentiviral vectors are capable of infecting non-dividing cells and can be used for both in vivo and ex vivo gene transfer and expression of nucleic acid sequences. For example, recombinant lentivirus capable of infecting a non-dividing cell wherein a suitable host cell is transfected with two or more vectors carrying the packaging functions, namely *gag*, *pol* and *env*, as well as *rev* and *tat* is described in U.S. Pat. No. 5,994,136, incorporated herein by reference. One may target the recombinant virus by linkage of the envelope protein with an antibody or a particular ligand for targeting to a receptor of a particular cell-type. By inserting a sequence (including a regulatory region) of interest into the viral vector, along with another gene which encodes the ligand for a receptor on a specific target cell, for example, the vector is now target-specific.

Combination Therapy

[0309] In certain embodiments of the invention, methods of the present invention for clinical aspects are combined with other agents effective in the treatment of hyperproliferative disease, such as anti-cancer agents. An "anti-cancer" agent is capable of negatively affecting cancer in a subject, for example, by killing cancer cells, inducing apoptosis in cancer cells, reducing the growth rate of cancer cells, reducing the incidence or number of metastases, reducing tumor size, inhibiting tumor growth, reducing the blood supply to a tumor or cancer cells, promoting an immune response against cancer cells or a tumor, preventing or inhibiting the progression of cancer, or increasing the lifespan of a subject with cancer. More generally, these other compositions would be provided in a combined amount effective to kill or inhibit proliferation of the cell. This process may involve contacting the cancer cells with the expression construct and the agent(s) or multiple factor(s) at the same time. This may be achieved by contacting the cell with a single composition or pharmacological formulation that includes both agents, or by contacting the cell with two distinct compositions or formulations, at the same time,

wherein one composition includes the expression construct and the other includes the second agent(s).

[0310] Tumor cell resistance to chemotherapy and radiotherapy agents represents a major problem in clinical oncology. One goal of current cancer research is to find ways to improve the efficacy of chemo- and radiotherapy by combining it with other therapies. In the context of the present invention, it is contemplated that cell therapy could be used similarly in conjunction with chemotherapeutic, radiotherapeutic, or immunotherapeutic intervention, as well as pro-apoptotic or cell cycle regulating agents.

[0311] Alternatively, the therapy may precede or follow the other agent treatment by intervals ranging from minutes to weeks. In embodiments where the other agent and present invention are applied separately to the individual, one would generally ensure that a significant period of time did not expire between the time of each delivery, such that the agent and inventive therapy would still be able to exert an advantageously combined effect on the cell. In such instances, it is contemplated that one may contact the cell with both modalities within about 12-24 h of each other and, more preferably, within about 6-12 h of each other. In some situations, it may be desirable to extend the time period for treatment significantly, however, where several d (2, 3, 4, 5, 6 or 7) to several wk (1, 2, 3, 4, 5, 6, 7 or 8) lapse between the respective administrations.

[0312] In some embodiments, the treatment cycles are repeated as necessary. It also is contemplated that various standard therapies, as well as surgical intervention, may be applied in combination with the inventive cell therapy.

Chemotherapy

[0313] Cancer therapies also include a variety of combination therapies with both chemical and radiation based treatments. Combination chemotherapies include, for example, abraxane, altretamine, docetaxel, herceptin, methotrexate, novantrone, zoladex, cisplatin (CDDP), carboplatin, procarbazine, mechlorethamine, cyclophosphamide, camptothecin, ifosfamide, melphalan, chlorambucil, busulfan, nitrosurea, dactinomycin, daunorubicin, doxorubicin, bleomycin, plicomycin, mitomycin, etoposide (VP16), tamoxifen, raloxifene, estrogen receptor binding agents, taxol, gemcitabine, navelbine, farnesyl-protein transferase inhibitors, transplatinum, 5-fluorouracil, vincristin, vinblastin and methotrexate, or any analog or derivative variant of the foregoing and also combinations thereof.

[0314] In specific embodiments, chemotherapy for the individual is employed in conjunction with the invention, for example before, during and/or after administration of the invention

Radiotherapy

[0315] Other factors that cause DNA damage and have been used extensively include what are commonly known as gamma-rays, X-rays, and/or the directed delivery of radioisotopes to tumor cells. Other forms of DNA damaging factors are also contemplated such as microwaves and UV-irradiation. It is most likely that all of these factors effect a broad range of damage on DNA, on the precursors of DNA, on the replication and repair of DNA, and on the assembly and maintenance of chromosomes. Dosage ranges for X-rays range from daily doses of 50 to 200 roentgens for prolonged periods of time (3 to 4 wk), to single doses of 2000 to 6000 roentgens. Dosage ranges for radioisotopes vary widely, and depend on the half-life of the isotope, the strength and type of radiation emitted, and the uptake by the neoplastic cells.

[0316] The terms "contacted" and "exposed," when applied to a cell, are used herein to describe the process by which a therapeutic construct and a chemotherapeutic or radiotherapeutic agent are delivered to a target cell or are placed in direct juxtaposition with the target cell. To achieve cell killing or stasis, both agents are delivered to a cell in a combined amount effective to kill the cell or prevent it from dividing.

Immunotherapy

[0317] Immunotherapies generally rely on the use of immune effector cells and molecules to target and destroy cancer cells. The immune effector may be, for example, an antibody specific for some marker on the surface of a tumor cell. The antibody alone may serve as an effector of therapy or it may recruit other cells to actually effect cell killing. The antibody also may be conjugated to a drug or toxin (chemotherapeutic, radionuclide, ricin A chain, cholera toxin, pertussis toxin, etc.) and serve merely as a targeting agent. Alternatively, the effector may be a lymphocyte carrying a surface molecule that interacts, either directly or indirectly, with a tumor cell target. Various effector cells include cytotoxic T cells and NK cells.

[0318] Immunotherapy other than the inventive therapy described herein could thus be used as part of a combined therapy, in conjunction with the present cell therapy. The general approach for combined therapy is discussed below. Generally, the tumor cell must bear some marker that is amenable to targeting, i.e., is not present on the majority of other cells. Many

tumor markers exist and any of these may be suitable for targeting in the context of the present invention. Common tumor markers include PD-1, PD-L1, CTLA4, carcinoembryonic antigen, prostate specific antigen, urinary tumor associated antigen, fetal antigen, tyrosinase (p97), gp68, TAG-72, HMFG, Sialyl Lewis Antigen, MucA, MucB, PLAP, estrogen receptor, laminin receptor, erb B and p155.

Genes

[0319] In yet another embodiment, the secondary treatment is a gene therapy in which a therapeutic polynucleotide is administered before, after, or at the same time as the present invention clinical embodiments. A variety of expression products are encompassed within the invention, including inducers of cellular proliferation, inhibitors of cellular proliferation, or regulators of programmed cell death.

Surgery

[0320] Approximately 60% of persons with cancer will undergo surgery of some type, which includes preventative, diagnostic or staging, curative and palliative surgery. Curative surgery is a cancer treatment that may be used in conjunction with other therapies, such as the treatment of the present invention, chemotherapy, radiotherapy, hormonal therapy, gene therapy, immunotherapy and/or alternative therapies.

[0321] Curative surgery includes resection in which all or part of cancerous tissue is physically removed, excised, and/or destroyed. Tumor resection refers to physical removal of at least part of a tumor. In addition to tumor resection, treatment by surgery includes laser surgery, cryosurgery, electrosurgery, and microscopically controlled surgery (Mohs' surgery). It is further contemplated that the present invention may be used in conjunction with removal of superficial cancers, precancers, or incidental amounts of normal tissue.

[0322] Upon excision of part of all of cancerous cells, tissue, or tumor, a cavity may be formed in the body. Treatment may be accomplished by perfusion, direct injection or local application of the area with an additional anti-cancer therapy. Such treatment may be repeated, for example, every 1, 2, 3, 4, 5, 6, or 7 days, or every 1, 2, 3, 4, and 5 weeks or every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months. These treatments may be of varying dosages as well.

Cryopreservation of NK Cells and CAR-NK Cells

[0323] The NK cells and CAR-NK cells disclosed herein can be cryopreserved in order to maintain cell phenotype post-thawing.

Compositions for Cryopreservation of Cells

[0324] Any suitable cryopreservation media for the immune cells can be used, including but not limited to, those described in application No. PCT/US21/13591, incorporated by reference herein. In some aspects, the cryopreservation medium comprises: dimethyl sulfoxide (DMSO), a disaccharide, a cytokine. In some aspects, the cryopreservation medium also comprises human serum albumin or human serum.

Uses of NK Cells and CAR-NK Cells

[0325] The NK cells as produced in the methods described herein can be used for various applications. For example, the compositions described herein, including the CAR-NK cell compositions contained within the cryopreservation medium described herein, are suitable for adoptive cell therapy. Adoptive cell therapies can be used to treat various disease, including, for example, cancer. In certain embodiments, the CAR-NK cell compositions contained within the cryopreservation medium described herein is useful for the treatment of a cancer or a tumor. In certain embodiments, the cancer comprises breast, heart, lung, small intestine, colon, spleen, kidney, bladder, head, neck, ovarian, prostate, brain, pancreatic, skin, bone, bone marrow, blood, thymus, uterine, testicular, and liver tumors. In some embodiments, the cancer is a blood cancer. In some embodiments, the blood cancer is a B-cell malignancy (e.g., diffuse large B-cell lymphoma).

[0326] In some embodiments, the cryopreservation media described herein is used to suspend cells used for adoptive cell therapy. Accordingly, in some embodiment, CAR-NK cell compositions are suspended in the cryopreservation media described herein.

[0327] In some embodiments, the CAR-NK cell compositions suspended in the cryopreservation media described herein is used to treat a subject who has cancer. In some embodiments, the subject is administered a composition comprising CAR-NK cells within the cryopreservation media described herein. In some embodiments, the CAR-NK cell comprises an anti-CD19 CAR gene and an IL-15 gene. In some embodiments, the CAR-NK cell comprises an

anti-CD19 CAR gene, an IL-15 gene, and iCaspase9. In some embodiments, the CAR-NK cells are not washed prior to administering to a subject in need thereof. In some embodiments, the CAR-NK cells are washed of the cryopreservation media prior to administering to a subject in need thereof. In some embodiments, the thawed cells are administered into a patient in need thereof within about 30 minutes and 2 hours from thawing the cells. In some embodiments, the rate of intravenous infusion into a subject is between about 2-3 minutes.

[0328] In some embodiments, the adoptive cell therapy is used in combination with one or more additional cancer treatments, such as for example lymphodepleting chemotherapy. Accordingly, in some embodiments, a subject who has cancer receives lymphodepleting chemotherapy before administration of a CAR-NK cell therapy product formulated in a cryopreservation media described herein.

[0329] In some embodiments, a CAR-NK cell therapy product is cryopreserved as described herein and subsequently thawed prior to administration to a patient in need thereof. For example, a CAR-NK cell therapy product as described herein is cryopreserved, transported, thawed, and administered to a patient in need thereof as described herein. Accordingly, in some embodiments, the CAR-NK cell therapy product is cryopreserved in a formulation as described herein and subsequently thawed prior to administration to a patient for treatment of B-cell malignancies.

[0330] In some embodiments, the frozen CAR-NK cell therapy product is frozen in a vial (e.g., a 50 ml AT vial) in a cryopreservation media described herein and using a method described herein and transported or shipped in the same vial at a temperature ranging from -140 °C to -196 °C to a location where a patient is situated, such that the cells can be thawed at the location where the patient is situated and administered aseptically directly to the patient using a syringe connected to the vial with a vial adapter (i.e., vial to vein transfer). For example, in some embodiments, a method of transporting the cell therapy product comprises: (a) providing the CAR-NK cells in a cryopreservation medium as described herein; (b) cooling the CAR-NK cells to a temperature of -80°C, thereby cryopreserving the mammalian cells; and (c) transporting the cryopreserved mammalian cells to a different location at a temperature of between about -20 °C to about -140 °C or below. Accordingly, in some embodiments, the cryopreserved mammalian cells are transported to a different location in a container maintained at a temperature

at or below -140°C . In some embodiments, the cryopreserved mammalian cells are transported to a different location in a container maintained at a temperature of between -140°C and -196°C . In some embodiments, the cryopreserved mammalian cells are transported to a different location in a cryoshipper. In some embodiments, the transported cells may be stored in a cryoshipper until administration to a patient. In some embodiments, the transported cells are stored at the different location at a temperature at or below -140°C . In some embodiments, the transported cells are stored at the different location at a temperature of between -140°C to -196°C . In some embodiments, the transported cells are stored in liquid nitrogen vapor phase from the time of receipt at the different location until a future time of use. The storage, thus can be accomplished using clinical site freezers or tanks that maintain temperature at or below -140°C in a vapor phase of liquid nitrogen. In some embodiments, the cell therapy product is a population of CD19-CAR-NK cells that further comprise IL-15 and iCaspase9. In some embodiments, the cell therapy product is cryopreserved in a container at concentration of between about 6 and 120 million cells per milliliter. In some embodiments, the cell therapy product is cryopreserved in a 50 mL container at concentration of between about 6 and 120 million cells per milliliter. In some embodiments, the cell therapy product is cryopreserved in a 50 mL container at concentration of between about 3 and 150 million cells per milliliter. In some embodiments, the cell therapy product is cryopreserved in a 50 mL container at concentration of between about 1 and 250 million cells per milliliter. In some embodiments, the cell therapy product is cryopreserved in a 50 mL container at concentration of between about 1 and 350 million cells per milliliter. In some embodiments, the cell therapy product is cryopreserved in a 50 mL container at a concentration of between about 1 and 500 million cells per milliliter.

[0331] In some embodiments, the cell therapy product comprises between about 20×10^6 and 100×10^7 cells in 50 mL container. In some embodiments, the cell therapy product comprises between about 100×10^6 and 900×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 50×10^6 in 50 mL container. In some embodiments, the cell therapy product comprises about 100×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 200×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 200×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 300×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 400×10^6

cells in 50 mL container. In some embodiments, the cell therapy product comprises about 500×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 600×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 700×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 800×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 900×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 1000×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 1500×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 2000×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 2500×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 3000×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 3500×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 4000×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 4500×10^6 cells in 50 mL container. In some embodiments, the cell therapy product comprises about 5000×10^6 cells in 50 mL container.

[0332] In some embodiments, the cell therapy product is contained in a 50 mL container at a fill volume of about between 20-45 mL. In some embodiments, the cell therapy product is contained in a 50 mL container at a fill volume of about 36 mL. In some embodiments, the cell therapy product is an immune cell, such as an NK cell, T cell or B cell. In some embodiments, the immune cell is engineered to comprise one or more transgenes, for example a chimeric antigen receptor (CAR). In some embodiments, the cells are CAR-NK+ cells. In some embodiments, the cell therapy product comprises a CD19-CAR, IL-15 transgene and an iCaspase9. In some embodiments, the cell therapy product comprises between about 100×10^6 and 900×10^6 CAR-NK+ cells in a 50-ml container. In some embodiments, the cell therapy product present is 200×10^6 CAR-NK+ cells in a 50-ml container. In some embodiments, the cell therapy product present is 300×10^6 CAR-NK+ cells in a 50-ml container. In some embodiments, the cell therapy product present is 400×10^6 CAR-NK+ cells in a 50-ml container. In some embodiments, the cell therapy product present is 500×10^6 CAR-NK+ cells in a 50-ml container. In some embodiments, the cell therapy product present is 600×10^6 CAR-NK+ cells in a 50-ml container. In some embodiments, the cell therapy product present is 700×10^6 CAR-NK+ cells in a 50-ml container.

10^6 CAR-NK+ cells in a 50-ml container. In some embodiments, the cell therapy product present is 800×10^6 CAR-NK+ cells in a 50-ml container. In some embodiments, the cell therapy product present is 900×10^6 CAR-NK+ cells in a 50 ml container.

[0333] The transported cell therapy product can be thawed as described herein followed by administration to a patient in need thereof. In some embodiments, the cell therapy product is thawed at the patient's bedside. In some embodiments, the cell therapy product is not washed prior to administration into a patient in need thereof.

[0334] In some embodiments, the transported cell therapy product remains frozen for further storage at the different location. In some embodiments, the thawed cells is introduced into a subject in need thereof without separating the cells and the cryopreservation solution. Thus, in some embodiments, the thawed cells are not washed prior to use. The thawed cells and accompanying cryopreservation solution is preferably warmed to body temperature (i.e., about 37°C .) prior to introduction into the subject. In such situation, the dose of the cells is based on the pre-freeze cell count.

[0335] In some embodiments, thawed cells are further cultured. In some embodiments, culturing involves placing the cells in an incubator; removing the buffer solution; and replacing the buffer solution with a culture medium designed for the growth and/or differentiation of cells. In some embodiments, the cells are incubated in the incubator for between about 6 to 7 hours. In some embodiments, the culture medium designed for the growth and/or differentiation of cells comprises Kubota's medium and/or a hormonally defined medium (HDM) for the differentiation of cells.

[0336] Viability of thawed cells can be assessed *in vitro* as well as *in vivo* using various methods known in the art. In some embodiments, the *in vitro* cell viability tests includes the Trypan Blue exclusion assay. In some embodiments, other analytical methods can be used to assess the cell viability of thawed cells that had been frozen with the different cryopreservation medium, for example, gene expression, through the use of RT-qPCR and the like. A person of ordinary skill in the art can opt for any analytical method to assess the viability of thawed cells that can be applied to assess the cell viability of otherwise fresh cells.

[0337] Viability of cells *in vivo*, in general, can be assessed by evaluating the functional characteristics of administered cells *in vivo*. In some embodiments, the *in vivo* viability of cells

can be assessed by evaluating the cell number of the cells that have been introduced into a subject in need. Various methods are known in the art for tracing cells and determining viability of administered cells.

[0338] The cryopreserved and thawed cells using the cryopreservation media described herein allows for using the cells for any purpose that a primary cell or fresh cell isolate can have. The cryopreserved and thawed cells retain high viability (e.g., greater than 70%, 75%, 80%, 85%, 90%, 95%, or greater than 95%) and retain physiological characteristics of their native state, which allows the cells to be used for a variety of applications, such as for genetic manipulation of the cells, and for cell therapy purposes such as, for example, in adoptive cell therapy applications.

Administration of CAR-NK cells

[0339] The CAR-modified NK cells of the present invention may be administered either alone, or as a pharmaceutical composition in combination with diluents and/or with other components such as IL-2 or other cytokines or cell populations. Pharmaceutical compositions of the present invention may comprise a target cell population as described herein, in combination with one or more pharmaceutically or physiologically acceptable carriers, diluents or excipients. Such compositions may comprise buffers such as neutral buffered saline, phosphate buffered saline and the like; carbohydrates such as glucose, mannose, sucrose or dextrans, mannitol; proteins; polypeptides or amino acids such as glycine; antioxidants; chelating agents such as EDTA or glutathione; adjuvants (e.g., aluminum hydroxide); and preservatives. In some embodiments, compositions of the present invention are formulated for intravenous administration.

[0340] Pharmaceutical compositions of the present invention may be administered in a manner appropriate to the disease to be treated (or prevented). The quantity and frequency of administration will be determined by such factors as the condition of the patient, and the type and severity of the patient's disease, although appropriate dosages may be determined by clinical trials.

[0341] When "an immunologically effective amount", "an anti-tumor effective amount", "an tumor-inhibiting effective amount", or "therapeutic amount" is indicated, the precise amount of the compositions of the present invention to be administered can be determined by a physician with consideration of individual differences in age, weight, tumor size, extent of infection or metastasis, and condition of the patient (subject). It can generally be stated that a pharmaceutical composition comprising the NK cells described herein may be administered at a dosage of about 10^4 to 10^9 cells/kg body weight, for example, about 10^5 to 10^6 cells/kg body weight. NK cell compositions may also be administered multiple times at these dosages. The cells can be administered by using infusion techniques that are commonly known in immunotherapy. The optimal dosage and treatment regime for a particular patient can readily be determined by one skilled in the art of medicine by monitoring the patient for signs of disease and adjusting the treatment accordingly.

[0342] The administration of the subject compositions may be carried out in any convenient manner, including by aerosol inhalation, injection, ingestion, transfusion, implantation or transplantation. The compositions described herein may be administered to a patient subcutaneously, intradermally, intratumorally, intranodally, intramedullary, intramuscularly, by intravenous (i.v.) injection, or intraperitoneally. In one embodiment, the NK cell compositions of the present invention are administered to a patient by intradermal or subcutaneous injection. In another embodiment, the NK cell compositions of the present invention are preferably administered by i. v. injection. The compositions of NK cells may be injected directly into a tumor, lymph node, or site of infection.

[0343] In certain embodiments of the present invention, cells activated and expanded using the methods described herein, are administered to a patient in conjunction with (for example, either before, simultaneously or following) any number of relevant treatment modalities, including but not limited to treatment with agents such as antiviral therapy, cidofovir and interleukin-2, cytarabine (also known as ARA-C) or natalizimab treatment or efaizumab treatment. In further embodiments, the NK cells of the invention may be used in combination with chemotherapy, radiation, immunosuppressive agents, such as cyclosporin, azathioprine, methotrexate, mycophenolate, and FK506, antibodies, or other immunoablative agents such as CAM PATH, anti-CD3 antibodies or other antibody therapies, cytoxin, fludarabine, cyclosporin, FK506, rapamycin, mycophenolic acid, steroids, FR901228, cytokines, and irradiation. In some

embodiments, the cell compositions of the present invention are administered to a patient in conjunction with (e.g., before, simultaneously or following) bone marrow transplantation, T cell ablative therapy using either chemotherapy agents such as, fludarabine, external-beam radiation therapy (XRT), cyclophosphamide, or antibodies such as OKT3 or CAMPATH. In some embodiments, the cell compositions of the present invention are administered following B-cell ablative therapy such as agents that react with CD20, e.g., Rituxan. For example, in some embodiments, subjects may undergo standard treatment with high dose chemotherapy followed by peripheral blood stem cell transplantation. In certain embodiments, following the transplant, subjects receive an infusion of the expanded immune cells of the present invention. In an additional embodiment, expanded cells are administered before or following surgery.

[0344] The dosage of the above treatments to be administered to a patient will vary with the precise nature of the condition being treated and the recipient of the treatment. The scaling of dosages for human administration can be performed according to art-accepted practices. In some embodiments, a dose about 1 to about 100 mg is administered to an adult patient. In some embodiments, the CAR-NK cell therapy is administered daily for a period between 1 and 30 days. In some embodiments, 1 to 10 mg per day is administered daily. In some embodiments, about 40 mg per day is administered.

EXAMPLES

[0345] Other features, objects, and advantages of the present invention are apparent in the examples that follow. It should be understood, however, that the examples, while indicating embodiments of the present invention, are given by way of illustration only, not limitation. Various changes and modifications within the scope of the invention will become apparent to those skilled in the art from the examples.

Example 1. Differentiation of iPS cells into HP Cell Bulk

[0346] FfI-01s04 strain, an iPS cell strain, was derived from peripheral blood mononuclear cell of a healthy individual. FfI-01s04 cells dispersed in StemFit complete medium

was seeded at 6×10^5 cells/well in an ultra-low adhesion-treated 6 well plate under a low-oxygen (5% O₂) condition ("Day 0"). The StemFit complete medium included 10 μ M CHIR99021 and 50 μ M Y-27632. Next day (i.e., Day 1), the Ffl-01s04 cells were dispersed in a hematopoietic progenitor cells (HPC) differentiation medium containing BMP4 (50 ng/mL), VEGF (50 ng/ml), bFGF (50 ng/mL), and ascorbic acid 2-phosphate (50 μ g/mL). The HPC induction medium medium included StemPro34 supplemented with human insulin (10 μ g/mL), human transferrin (5.5 μ g/mL), sodium selenite (6.7 ng/mL), L-glutamine (2 mM) and α -monothioglycerol (0.4 mM). On Day 2, SB431542 (at 6 μ M in medium) was added to the culture medium (i.e., HPC differentiation medium containing cells), and the cells were cultured for 2 days. On Day 4, cells were redispersed in another medium containing VEGF (50 ng/mL), bFGF (50 ng/mL), SCF (50 ng/mL), and ascorbic acid 2-phosphate (50 μ g/mL), and cultured for further 3 days. On Day 7, cells were exposed to another medium containing VEGF (50 ng/mL), bFGF (50 ng/mL), SCF (50 ng/mL), ascorbic acid 2-phosphate (50 μ g/mL), TPO (30 ng/mL) and Flt3L (10 ng/mL), and cultured using this medium for additional 7 days. The medium was changed every 2-3 days during this 7 days-culture period.

Example 2. Differentiation of HP cell bulk into population comprising CD4⁺ cells

[0347] At Day 14, the cell population obtained from Example 1 without undergoing any cell isolation ("HP cell bulk") was seeded in 15 cm dishes at 3.12×10^6 cells/dish, and cultured at 37°C under 5% O₂. It is noted that various seeding densities may be used, and the aforementioned seeding density is one embodiment. Each 15 cm dish was coated with rh-DLL4/Fc chimera (Sino Biological) and RetroNectin (Takara Bio Inc). The medium was changed every 2-3 days during this culture period. MEM α (Thermo Fisher Scientific (Gibco)) supplemented with 15% FBS, 4 mM L-glutamine, 100 U/mL penicillin, 100 μ g/ml streptomycin, 55 μ M 2-mercaptoethanol, 50 μ g/mL ascorbic acid 2-phosphate, 10 μ g/mL human insulin, 5.5 μ g/mL human transferrin, 6.7 ng/mL sodium selenite, 50 ng/mL SCF, 50 ng/mL IL-7, 50 ng/mL Flt3L, 100 ng/mL TPO, 15 μ M SB203580, 30 nM SDF-1 α was used as the medium to culture these cells. On Day 21, the cells were passaged to new 15 cm dishes freshly coated with hDLL4/RetroNectin. On Day 28, the cells were further passaged to new 15 cm dishes freshly

coated with hDLL4/RetroNectin. On Day 35, all cells including CD4⁻ cells, among others, were harvested.

Example 3. Differentiation of bulk to NK cell bulk

[0348] On Day 35, population comprising CD4⁻ cells, i.e., cells obtained from Example 2 without undergoing any cell isolation, was seeded in a 48-well plate at 1×10^6 cells/well, and cultured under 5% CO₂ at 37°C for three days. MEM α medium supplemented with 15% FBS, 4 mM L-glutamine, 100 U/mL penicillin, 100 ng/mL streptomycin, 50 μ g/mL ascorbic acid 2-phosphate, 10 μ g/mL human insulin, 5.5 μ g/mL human transferrin, 6.7 ng/mL sodium selenite, 500 ng/mL anti-CD3 antibody (UCHT1), 10 ng/mL IL-2, and 10 ng/mL IL-7 was used as a culture medium. On Day 38, the cells were dispersed in another MEM α medium supplemented with 15% FBS, 4 mM L-glutamine, 100 U/mL penicillin, 100 ng/mL streptomycin, 50 μ g/mL ascorbic acid 2-phosphate, 10 μ g/mL human insulin, 5.5 μ g/mL human transferrin, 6.7 ng/mL sodium selenite, 10 ng/mL IL-2, and 10 ng/mL IL-7 was used as a culture medium. On Day 42, all cells including NK cells (“NK cell bulk”) were harvested.

Example 4. Differentiation of bulk to NK cell bulk – CD4⁻ Cell Enrichment

[0349] In some embodiments, the population of cells obtained from Example 2 is processed to enrich for cells that are CD4⁻. This can be done for example, with methods that deplete CD4⁺ cells (i.e., CD4⁺ cell depletion). Exemplary methods are discussed below.

CD4⁺ Cell Depletion

[0350] In some embodiments, on day 35, the population obtained in Example 2 is depleted of cells that express CD4 antigen. Specifically, the population of cells obtained from Example 2 is subjected to a cell depletion method, such as fluorescent activated cell sorting (FACS) or a magnetic-based sorting method (MACS), to deplete cells in the population that express CD4 antigen, thereby depleting the CD4⁺ cells in the population of cells obtained from Example 2. In this manner, cells are obtained that are enriched for CD4⁻ cells.

[0351] Following CD4⁺ cell depletion, the enriched CD4⁻ cells are seeded in a 48-well plate at 1×10^6 cells/well, and cultured under 5% CO₂ at 37°C for three days. MEM α medium

supplemented with 15% FBS, 4 mM L-glutamine, 100 U/mL penicillin, 100 ng/mL streptomycin, 50 µg/mL ascorbic acid 2-phosphate, 10 µg/mL human insulin, 5.5 µg/mL human transferrin, 6.7 ng/mL sodium selenite, 500 ng/mL anti-CD3 antibody (UCHT1), 10 ng/mL IL-2, and 10 ng/mL IL-7 is used as a culture medium. On Day 38, the cells are dispersed in another MEMα medium supplemented with 15% FBS, 4 mM L-glutamine, 100 U/mL penicillin, 100 ng/mL streptomycin, 50 µg/mL ascorbic acid 2-phosphate, 10 µg/mL human insulin, 5.5 µg/mL human transferrin, 6.7 ng/mL sodium selenite, 10 ng/mL IL-2, and 10 ng/mL IL-7 is used as a culture medium. On Day 42, all cells including NK cells (“NK cell bulk”) are harvested.

Example 5. Flow Cytometry analysis

[0352] The NK cell bulk was then stained with a set of antibodies listed in **Table 1**, and was analyzed by flow cytometry. As shown in **FIG. 1**, a portion of CD3 negative NK cell bulk expressed CD56 (“CD56+/CD3- cells”). The CD56+/CD3- cells are natural killer cells (NK cells). Thus, CD56+/CD3- cells were prepared from an HPC bulk that was derived from an iPSCs (Ffi-01s04 strain). A cell expressing CD56 that was obtained in the process above is sometimes referred to as iPS NK cell. The antibodies used for flow cytometry is shown in Table 1.

Table 1:

Anti-CD3 antibody	CD3 BioLegend APC/Cy7
Anti-CD56 antibody	CD56 Biolegend PE

Example 6. Single Cell RNA sequence (scRNAseq) analysis

[0353] In order to identify different cell types in the NK cell bulk population (i.e., the cells obtained in Example 3 or Example 4), a non-linear dimensionality-reduction technique (uniform manifold approximation and projection (UMAP) based on single cell RNA-seq (scRNAseq)) was applied.

[0354] Ten thousands cells from NK cell bulk were subjected to single cell RNA-sequencing on the Illumina NextSeq 500 at Genewiz. Using the SingleR algorithm and manual cluster labeling, the events were classified into four broad cell populations - monocyte, B cell,

NK cell and T cell in PBMC samples. The dataset of PBMCs, freely available from 10X Genomics, was used as a control for data analysis.

[0355] As shown in **FIG. 2**, about 75% of the cells in the NK cell bulk were identified as NK cells, while about 25% of the cells in the NK cell bulk were identified as T cells. Thus, the NK cell bulk included both NK cells and T cells based on mRNA profiling by single cell RNA-sequencing.

Example 7. CAR-NK cell preparation

[0356] NK cells were further modified to express one or more CARs. Modification of NK cells includes steps of: (1) synthesizing an anti-CD19 CAR gene and an IL-15R α /IL-15 gene; (2) preparing a retrovirus vector containing anti-CD19 CAR gene and IL-15R α /IL-15 genes; and (3) transducing the NK cells with the retrovirus vector containing anti-CD19 CAR gene and IL-15R α /IL-15 genes.

CAR/IL15-NK cell preparation

[0357] Anti-CD19 CAR gene was prepared by synthesizing oligopeptides that are designed to be arranged from N-terminal as shown in **Table 2**.

Table 2:

Sequence from N-terminal	Genes	Number of Amino acids
1	Lead sequence of the heavy chain of an immunoglobulin	22
2	Variable domain in the light chain of the anti-CD19 antibody(FMC60)	104
3	GGGS linker	15
4	Variable domain in the heavy chain of the anti-CD19 antibody(FMC60)	120
5	CD8 derived sequence (Containing transmembrane domain)	83
6	Intracellular domain of 4-1BB	47
7	Intracellular domain of CD3 zeta	112

[0358] An anti-CD19 CAR was constructed in accordance with WO2014/153270, incorporated herein by reference in its entirety.

Preparation of IL-15 α /IL-15 gene

[0359] IL-15 α /IL-15 gene was prepared by synthesizing oligopeptides that are designed to be arranged from N-terminal. The IL-15 α /IL-15 was constructed in accordance with Mortier et al., 2006, *The Journal of Biological Chemistry*, Vol 281, No 3, pages 1612-1619, January 20, 2006; Chertova et al., *The Journal of Biological Chemistry*, Vol. 288, No. 25, Pages 18093-18103, June 21, 2013; and Rowley et al., *Eur J Immunol*, 2009 February; 39(2):491-506, each of which are incorporated by reference in its entirety.

Table 4:

Sequence from N-terminal	Genes	Number of Amino acids
1	Lead sequence of human IL-2	23
2	C-terminal sequence of human IL-15	114
3	GGGS linker	24
4	C-terminal sequence of human IL-15RA	239

Preparation of retrovirus vector containing anti-CD19 CAR gene

[0360] The anti-CD19 CAR gene was incorporated into a multi-cloning site of a pMY retrovirus vector. The retrovirus vector was generated using FRY-RD18 cells for retrovirus vector production.

Preparation of retrovirus vector containing IL-15 α /IL-15 gene

[0361] The IL-15 α /IL-15 gene was incorporated into a multi-cloning site of another pMY retrovirus vector. The retrovirus vector was generated using FRY-RD18 cell for retrovirus vector production.

Transduction of anti-CD19 CAR gene and IL-15 α /IL-15 gene into iPS NK cell

[0362] The iPS NK cells were transduced with the retrovirus vectors containing anti-CD19 CAR gene and with the retrovirus vectors containing IL-15 α /IL-15 gene to generate the anti-CD19 CAR expressing iPS NK cell ("iNK-CAR19").

Example 8. In vivo anti-tumor activity of iNK-CAR19

[0363] Luciferase expressing Nalm6 cells (ATCC; cancer cells) (5×10^5 cells) were transplanted into NOD/Shi-scid, IL-2R gamma null mice ("NSG mice") via tail vein. The NSG mice, male, 4-5-week-old were sourced from The Jackson Laboratory. Four days after the transplantation of Nalm6 cells, the iNK-CAR19 (1×10^7 cells) dispersed in 0.2 ml PBS or just 0.2 ml of PBS without cells were administered to the Nalm6-transplanted NSG mice via tail vein. After the administration of iNK-CAR19 cells or PBS, luciferin was administered to the mice via tail vein. Luciferase activity was measured using IVIS imaging system (PerkinElmer) for over 70 days.

[0364] FIG. 3 shows the anti-tumor efficacy of untreated (treated with buffer only) and iNK-CAR19 cell treated Nalm6-transplanted NSG mice. Just after three days, luminescence was detected in all the mice treated with D-PBS buffer. In contrast, luminescence was not detected in other mice (treated with either primary CAR T cells or iNK-CAR cells) until 28 days after administration, and luminescence was not detected in one mouse treated with iNK-CAR19 cells even 70 days after the treatment. Thus, iNK-CAR19 cells clearly demonstrated an enhanced toxicity against Nalm6 cancer cells.

EQUIVALENTS AND SCOPE

[0365] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. The scope of the present invention is not intended to be limited to the above Description, but rather is as set forth in the following claims:

CLAIMS

We claim

1. A method of producing a cell population enriched in NK cells, the method comprising the steps of:

(A) culturing a population of pluripotent stem cells under a first set of conditions resulting in a cell population comprising at least 20% CD34+ HPCs (HPC bulk); and

(B) changing the first set of conditions to a second set of conditions, thereby resulting in a cell population comprising at least 5% CD4- cells; and

(C) changing the second set of conditions to a third set of conditions, thereby resulting in a cell population enriched in NK cells.

2. The method of claim 1, wherein the method is performed without an isolation step.

3. The method of claim 1 comprising isolating CD4- cells after step (B) before contacting the isolated cells with the third set of conditions in step (C).

4. The method of claim 3, wherein the isolating CD4- cells comprises removing the CD4+ cells from the cell population.

5. A method of producing a cell population enriched in NK cells, the method comprising the steps of:

(A) culturing a population of pluripotent stem cells under a first set of conditions resulting in a cell population comprising at least 20% CD34+ HPCs (HPC bulk); and

(B) changing the first set of conditions to a second set of conditions, thereby resulting in a cell population comprising at least 5% CD4- cells; and

(C) changing the second set of conditions to a third set of conditions, thereby resulting in a cell population enriched in NK cells, wherein the method is performed without an isolation step.

6. The method of any one of the preceding claims, wherein the cell population comprising at least 5% CD4- cells further comprises at least 5% CD4+ cells.

7. The method of any one of the preceding claims, wherein the cell population enriched in NK cells comprises at least 30% NK cells.
8. The method of any one of the preceding claims, wherein the resulted cell population in step (B) comprises at least 10%, 15% or 20% CD4⁻ cells.
9. The method of any one of the preceding claims, wherein the CD4⁻ cells in step (B) are CD8⁺ cells.
10. The method of any one of claims 1-8, wherein the CD4⁻ cells in step (B) are CD8⁻ cells.
11. The method of claim any one of claims 1-8, wherein the resulted cell population in step (B) comprises between 20 and 55% CD4⁻/CD8⁺ cells.
12. The method of claim 11, wherein the resulted cell population comprises between 25 and 55% CD4⁻/CD8⁻ cells.
13. The method of claim 1, wherein the NK cells are CD56⁺/CD3⁻ cells.
14. The method of any one of the preceding claims, wherein the pluripotent stem cells are induced pluripotent stem cells (iPSCs).
15. The method of any one of the preceding claims, wherein the first set of conditions comprises a culture medium comprising at least one compound selected from bone morphogenetic protein-4 (BMP4) vascular endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF), ascorbic acid, Flt3 ligand (Flt3L), thrombopoietin (TPO) and TGFβ inhibitor.
16. The method of claim 15, wherein the first set of conditions comprises a culture medium comprising BMP4 at a concentration of between 5 ng/mL - 500 ng/mL.
17. The method of claim 16 wherein the BMP4 is at a concentration of 50 ng/mL.
18. The method of any one of claims 15-17, wherein the first set of conditions comprises a culture medium comprising VEGF at a concentration of between 5 ng/mL - 500 ng/mL.
19. The method of claim 18, wherein the VEGF is at a concentration of about 50 ng/mL.

20. The method of any one of claims 15-19, wherein the first set of conditions comprises a culture medium comprising bFGF at a concentration of between 5 ng/mL - 500 ng/mL.
21. The method of claim 20, wherein the bFGF is at a concentration of 50 ng/mL.
22. The method of any one of claims 15-21, wherein the first set of conditions comprises a culture medium comprising ascorbic acid at a concentration of between 5 µg/mL - 500 µg/mL.
23. The method of claim 22, wherein the ascorbic acid is at a concentration of 50 µg/mL.
24. The method of any one of claims 15-23, wherein the first set of conditions comprises a culture medium comprising Flt3L at a concentration between 1 ng/mL - 100 ng/mL.
25. The method of claim 24, wherein the Flt3L is at a concentration of 50 ng/mL.
26. The method of any one of claims 15-25, wherein the first set of conditions comprises a culture medium comprising TPO at a concentration of between 1 ng/mL - 200 ng/mL.
27. The method of claim 26, wherein the TPO is at a concentration of 100 ng/mL.
28. The method of any one of the preceding claims, wherein the second set of conditions comprises a culture medium comprising at least one compound selected from the group consisting of ascorbic acid, stem cell factor (SCF), IL-7, Flt3L, thrombopoietin (TPO), p38 inhibitor and SDF-1.
29. The method of claim 28, wherein the second set of conditions comprises a culture medium comprising ascorbic acid at a concentration between 5 µg/mL - 500 µg/mL.
30. The method of claim 29, wherein the ascorbic acid is at a concentration of 50 µg/mL.
31. The method of any one of claims 28-30, wherein the second set of conditions comprises a culture medium comprising SCF at a concentration between 5 ng/mL - 100 ng/mL.
32. The method of claim 31, wherein the SCF is at a concentration 50 ng/mL.
33. The method of any one of claims 28-32, wherein the second set of conditions comprises a culture medium comprising IL-7 at a concentration between 1 ng/mL to 100 ng/mL.

34. The method of claim 33, wherein the IL-7 is at a concentration of 50 ng/mL.
35. The method of any one of claims 28-34, wherein the second set of conditions comprises a culture medium comprising Flt3L at a concentration of between 1 ng/mL - 100 ng/mL.
36. The method of claim 35, wherein the Flt3L at a concentration of 50 ng/mL.
37. The method of any one of claims 28-36, wherein the second set of conditions comprises a culture medium comprising TPO at a concentration between 1 ng/mL - 200 ng/mL.
38. The method of claim 37, wherein the TPO is at a concentration of 100 ng/mL.
39. The method of any one of claims 28-38, wherein the second set of conditions comprises a culture medium comprising p38 inhibitor at a concentration between 0.5 μ M to 100 μ M.
40. The method of claim 39, wherein the p38 inhibitor is SB203580.
41. The method of claim 40, wherein the SB203580 is at a concentration of 15 μ M.
42. The method of any one of claims 28-41, wherein the second set of conditions comprises a culture medium comprising SDF-1 at a concentration between 10 ng/mL - 100 ng/mL.
43. The method of claim 42, wherein the SDF-1 is at a concentration of 30 nM.
44. The method of any one of the preceding claims, wherein the third set of conditions comprises a culture medium comprising at least one compound selected from the group consisting of a CD3 activator, IL-2 and IL-7.
45. The method of claim 44, wherein the third set of conditions comprises IL-2 at a concentration of between 1 ng/mL - 100 ng/mL.
46. The method of claim 45, wherein the IL-2 is at a concentration of 10 ng/mL.
47. The method of any one of claims 44-46, wherein the third set of conditions comprises a culture medium comprising IL-7 at a concentration between 1 ng/mL - 100 ng/mL.
48. The method of claim 47, wherein the IL-7 is at a concentration of 10 ng/mL.

49. The method of any one of the preceding claims, wherein each of the culturing steps is performed at about 5% oxygen.
50. The method of any one of claims 1-48, wherein each of the culturing steps is performed at greater than 14% oxygen.
51. The method of any one of claims 1-48, wherein each of the culturing steps is performed at atmospheric oxygen.
52. The method of any one of claims 1-48, wherein each of the culturing steps is performed at less than 5% oxygen.
53. The method of any one of the preceding claims, wherein culturing pluripotent stem cells in the first set of conditions to obtain HPC bulk lasts for greater than 10 days.
54. The method of claim 53, wherein culturing pluripotent stem cells in the first set of conditions to obtain HPC bulk lasts between 11 and 15 days.
55. The method of claim 54, wherein culturing pluripotent stem cells in the first set of conditions to obtain HPC bulk lasts for 14 days.
56. The method of any one of the preceding claims, wherein the iPSC are obtained from peripheral blood mononuclear cells.
57. The method of any one of the preceding claims, wherein at least about 50%, 55%, 60%, 75%, 80%, 85%, 90%, 95%, 97% or more of produced cells are CD56+/CD3- NK cells without a step of enrichment.
58. The method of claim 57, wherein less than about 25% of produced cells are CD3+ cells.
59. The method of claim 57 or claim 58, wherein the percentage of produced cells is determined by flow cytometry.
60. The method of claim 57 or claim 58, wherein the percentage of produced cells is determined by single-cell RNA sequencing (scRNAseq).

61. The method of any one of the preceding claims, further comprising isolating CD56+/CD3- cells.
62. The method of any one of claims 3, 4 and 61, wherein the isolating comprises fluorescence-activated cell sorting (FACS) or magnetic sorting.
63. The method of any one of the preceding claims, wherein the NK cells are genetically modified to express one or more chimeric antigen receptors (CAR).
64. The method of claim 63, wherein the antigen is CD19.
65. The method of claim 63 or claim 64, wherein the NK cells are further genetically modified to express IL-15R α /IL-15 complex.
66. A method of producing induced pluripotent stem cell (iPSC)-derived NK cells comprising the following steps:
- (1) culturing iPSC in a first set of conditions comprising a medium comprising at least one compound selected from vascular endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF), and ascorbic acid to obtain a population comprising at least 20% CD34+ HPCs (HPC bulk);
 - (2) culturing the HPC bulk obtained in (1) in second set of conditions comprising a culture medium comprising one or more of ascorbic acid, p38 inhibitor and SDF-1 to obtain a cell population comprising at least 5% CD4- cells; and
 - (3) culturing the cell population from (2) in a third set of conditions comprising a culture medium comprising at least one compound selected from the group consisting of a CD3 activator, IL-2 and IL-7.
67. A method of producing induced pluripotent stem cell (iPSC)-derived NK cells comprising the following steps:
- (1) culturing a bulk cell population comprising hematopoietic progenitor cells (HPC bulk) in a second set of conditions comprising a culture medium comprising one or more of

ascorbic acid, p38 inhibitor and SDF-1 to obtain an a cell population comprising at least 5% CD4⁺ cells;

(2) isolating the CD4⁺ cells from (1); and

(3) culturing the isolated CD4⁺ cells in NK induction media comprising at least one compound selected from the group consisting of a CD3 activator, IL-2 and IL-7.

68. The method of claim 66 or claim 67, wherein about 15-30% of the cells in the cell population obtained in step (1) are CD4⁺ cells.

69. The method of claim 68, wherein about 20% of the cells in the cell population obtained in step (1) are CD4⁺ cells.

70. The method of any one of the preceding claims, wherein the cell population comprising CD4⁺ cells comprises CD4⁺/CD8⁺ cells and CD4⁺/CD8⁺ cells.

71. An NK cell population produced using a method of any one of the preceding claims.

72. The NK cell population of claim 71, wherein the NK cell population is cryopreserved in a cryopreservation medium.

73. An unsorted cell population comprising pluripotent stem cell-derived CD56⁺/CD3⁺ cells at a ratio of not less than 60% of total pluripotent stem cell-derived CD56⁺ cells.

74. The unsorted cell population of claim 73, wherein less than 25% of cells in the unsorted cell population are CD3⁺ cells.

75. The unsorted cell population of claims 73 or claim 74, wherein less than 5% of cells in the unsorted cell population are monocytes.

76. The unsorted cell population of claims 73 or claim 74, wherein less than 5% of cells in the unsorted cell population are B cells.

77. The unsorted population of any one of claims 73-76, wherein the percentage of cells in the unsorted cell population is determined by flow cytometry.

78. The unsorted population of any one of claims 73-76, wherein the percentage of cells is determined by single-cell RNA sequencing (scRNAseq).
79. A method of treating a subject in need of cell therapy, comprising administering to the subject the NK cells of any one of the preceding claims.
80. The method of claim 79, wherein the subject has a cancer.
81. The method of claim 80, wherein the cancer is leukemia or lymphoma.
82. A method of producing a cell population comprising CD34+ hematopoietic progenitor cells (HPCs), the method comprising culturing a population of pluripotent stem cells in a first set of conditions comprising a medium comprising at least one compound selected from vascular endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF), and ascorbic acid to obtain a population comprising CD34+ HPCs (HPC bulk).
83. A cell population comprising HPCs produced by the method of claim 82.
84. The cell population of claim 83, wherein the cell population comprises at least 20% CD34+ cells.




FIG. 1

scRNA Data:
~75% cells are identified as NK cells
by SingleR algorithm
Algorithmically-assigned cell type

	B cells		Monocytes		NK Cells		T Cells
---	---------	---	-----------	---	----------	---	---------




FIG. 2




FIG. 3