



- (51) **International Patent Classification:**  
A61K 39/02 (2006.01)
- (21) **International Application Number:**  
PCT/IB2015/050314
- (22) **International Filing Date:**  
15 January 2015 (15.01.2015)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**  
61/929,598 21 January 2014 (21.01.2014) US
- (71) **Applicant: PFIZER INC.** [US/US]; 235 East 42nd Street, New York, New York 10017 (US).
- (72) **Inventors: GU, Jianxin;** 367 Bullard Avenue, Paramus, New Jersey 07652 (US). **KAINTHAN, Rajesh Kumar;** 15 Key Place, Tappan, New York 10983 (US). **KIM, Jin-Hwan;** 44 Somerset Drive, Suffern, New York 10901 (US). **PRASAD, Avvari Krishna;** 105 Worsham Drive, Chapel Hill, North Carolina 27516 (US). **YANG, Yu-Ying;** 154 Pepper Ridge Road, Unit 11, Stamford, Connecticut 06905 (US).
- (74) **Agent: OLSON, A. Dean;** Pfizer Worldwide Research & Development, Eastern Point Road MS8260-2141, Groton, Connecticut 06340 (US).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,

DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

- as to the identity of the inventor (Rule 4.17(i))
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

**Published:**

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))
- with sequence listing part of description (Rule 5.2(a))



WO 2015/110940 A2

(54) **Title:** STREPTOCOCCUS PNEUMONIAE CAPSULAR POLYSACCHARIDES AND CONJUGATES THEREOF

(57) **Abstract:** The invention relates to activated Streptococcus pneumoniae serotype 10A, 22F or 33F polysaccharides and processes for their preparation. The invention also relates to immunogenic conjugates comprising Streptococcus pneumoniae serotype 10A, 22F or 33F polysaccharides covalently linked to a carrier protein, processes for their preparation and immunogenic compositions and vaccines comprising them.

**Streptococcus pneumoniae capsular polysaccharides and conjugates thereof****Field of the invention**

5

The invention relates to activated *Streptococcus pneumoniae* serotype 10A, 22F or 33F polysaccharides and processes for their preparation. The invention also relates to immunogenic conjugates comprising *Streptococcus pneumoniae* serotype 10A, 22F or 33F polysaccharides covalently linked to a carrier protein, processes for their preparation and  
10 immunogenic compositions and vaccines comprising them.

**Background**

*Streptococcus pneumoniae* are Gram-positive, lancet shaped cocci that are usually seen in  
15 pairs (diplococci), but also in short chains or as single cells. They grow readily on blood agar plates with glistening colonies and display alpha hemolysis unless grown anaerobically where they show beta hemolysis. The cells of most pneumococcal serotypes have a capsule which is a polysaccharide coating surrounding each cell. This capsule is a determinant of virulence in humans, as it interferes with phagocytosis by preventing antibodies from  
20 attaching to the bacterial cells. Currently there are more than 90 known pneumococcal capsular serotypes identified, with the 23 most common serotypes accounting for approximately 90% of invasive disease worldwide. As a vaccine, the pneumococcal polysaccharide coat can confer a reasonable degree of immunity to *Streptococcus pneumoniae* in individuals with developed or unimpaired immune systems, but the capsular  
25 polysaccharide conjugated to a suitable carrier protein allows for an immune response in infants and elderly who are also at most risk for pneumococcal infections.

Since the introduction of the first 7-valent pneumococcal conjugate vaccine (PCV7 or Pevnar) in 2000, invasive disease from those seven serotypes (4, 6B, 9V, 14, 18C, 19F, and  
30 23F) has nearly disappeared. The addition of serotypes 1, 3, 5, 6A, 7F and 19A in Pevnar 13 further decreased the numbers of invasive pneumococcal disease.

None of the currently marketed pneumococcal vaccine provides an appropriate protection against serotype 10A, 22F or 33F *Streptococcus pneumoniae* in human and in particular in  
35 children less than 2 years old. Therefore, there is a need for immunogenic compositions that can be used to induce an immune response against serotype 10A, 22F or 33F *Streptococcus pneumoniae*.

## Summary of the invention

In one aspect the present disclosure provides a process for preparing an activated  
5 *Streptococcus pneumoniae* serotype 10A, 22F or 33F capsular polysaccharide, the process comprising the steps of:

(a) reacting an isolated serotype 10A, 22F or 33F capsular polysaccharide with an oxidizing agent; and;

10 (b) quenching the oxidation reaction by addition of a quenching agent resulting in an activated *Streptococcus pneumoniae* serotype 10A, 22F or 33F polysaccharide.

In a further aspect, the invention relates to activated *Streptococcus pneumoniae* serotype 10A, 22F or 33F capsular polysaccharide obtained or obtainable by the activation process disclosed herein.

In a further aspect, the present disclosure provides a process for the preparation of an  
15 immunogenic conjugate comprising *Streptococcus pneumoniae* serotype 10A, 22F or 33F polysaccharide covalently linked to a carrier protein, the process comprising the steps of:

(a) compounding an activated serotype 10A, 22F or 33F polysaccharide obtained or obtainable by a process disclosed herein with a carrier protein; and,

20 (b) reacting the compounded, activated serotype 10A, 22F or 33F polysaccharide and carrier protein with a reducing agent to form a serotype 10A, 22F or 33F polysaccharide:carrier protein conjugate.

In a further aspect, the present disclosure provides serotype 10A, 22F or 33F polysaccharide:carrier protein conjugate obtained or obtainable by the process disclosed herein.

25 In a further aspect, the present disclosure provides immunogenic compositions and vaccines comprising immunogenic conjugate disclosed herein.

In a further aspect, the present disclosure a method of treating or preventing a *Streptococcus pneumoniae* infection, disease or condition associated with serotype 10A, 22F or 33F  
30 *Streptococcus pneumoniae* in a subject, the method comprising the step of administering a therapeutically or prophylactically effective amount of an immunogenic composition or a vaccine disclosed herein.

## Brief description of the drawings

35 Figure 1 shows the structure of *Streptococcus pneumoniae* capsular polysaccharide Serotype 10A Repeat Unit.

Figure 2 shows the structure of *Streptococcus pneumoniae* capsular polysaccharide Serotype 22F Repeat Unit.

Figure 3 shows the structure of *Streptococcus pneumoniae* capsular polysaccharide Serotype 33F Repeat Unit.

5 Figure 4 shows the molecular weight of activated serotype 22F polysaccharide as a function of the amount of oxidizing agent at 23°C with and without quenching step.

Figure 5 shows the molecular weight of activated serotype 22F polysaccharide as a function of the amount of oxidizing agent at 4°C with and without quenching step.

10 Figure 6 shows the molecular weight of activated serotype 33F polysaccharide as a function of the amount of oxidizing agent at 2 to 8°C with or without quenching step or at 23°C without quenching step.

### Detailed description of the invention

15 The present invention may be understood more readily by reference to the following detailed description of the preferred embodiments of the invention and the Examples included herein. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. Although any methods and materials similar or equivalent to those described  
20 herein can be used in the practice or testing of the present invention, certain preferred methods and materials are described herein. In describing the embodiments and claiming the invention, certain terminology will be used in accordance with the definitions set out below.

25 As used herein, the singular forms "a", "an", and "the" include plural references unless indicated otherwise. Thus, for example, references to "the method" includes one or more methods, and/or steps of the type described herein and/or which will become apparent to one of ordinary skill in the art upon reading this disclosure.

30 As used herein, the term "molecular weight" of polysaccharide or of carrier protein-polysaccharide conjugate refers to molecular weight calculated by size exclusion chromatography (SEC) combined with multiangle laser light scattering detector (MALLS).

As used herein, the term "degree of oxidation" (DO) refers to the number of sugar repeat  
35 units per aldehyde group generated when the isolated polysaccharide is activated with an oxidizing agent. The degree of oxidation of a polysaccharide can be determined using routine methods known to the man skilled in the art.

It is noted that in this disclosure, terms such as "comprises," "comprised," "comprising," "contains," "containing" and the like can have the meaning attributed to them in U.S. patent law; e.g., they can mean "includes," "included," "including" and the like. Such terms refer to the inclusion of a particular ingredients or set of ingredients without excluding any other ingredients. Terms such as "consisting essentially of" and "consists essentially of" have the meaning attributed to them in U.S. patent law, e.g., they allow for the inclusion of additional ingredients or steps that do not detract from the novel or basic characteristics of the invention, i.e., they exclude additional unrecited ingredients or steps that detract from the novel or basic characteristics of the invention. The terms "consists of" and "consisting of" have the meaning ascribed to them in U.S. patent law; namely, that these terms are closed ended. Accordingly, these terms refer to the inclusion of a particular ingredient or set of ingredients and the exclusion of all other ingredients.

As used herein, the term "conjugates" or "glycoconjugates" as used herein refers to a polysaccharide covalently conjugated to a carrier protein. Glycoconjugates of the invention and immunogenic compositions comprising them may contain some amount of free polysaccharide.

As used herein, the term "serotype 10A glycoconjugate" or "serotype 10A conjugate" refers to an isolated *Streptococcus pneumoniae* serotype 10A capsular polysaccharide covalently conjugated to a carrier protein.

As used herein, the term "serotype 22F glycoconjugate" or "serotype 22F conjugate" refers to an isolated *Streptococcus pneumoniae* serotype 22F capsular polysaccharide covalently conjugated to a carrier protein.

As used herein, the term "serotype 33F glycoconjugate" or "serotype 33F conjugate" refers to an isolated *Streptococcus pneumoniae* serotype 33F capsular polysaccharide covalently conjugated to a carrier protein.

As used herein, the term "serotype 10A polysaccharide" refers to a *Streptococcus pneumoniae* serotype 10A capsular polysaccharide.

As used herein, the term "serotype 22F polysaccharide" refers to a *Streptococcus pneumoniae* serotype 22F capsular polysaccharide.

As used herein, the term "serotype 33F polysaccharide" refers to a *Streptococcus pneumoniae* serotype 33F capsular polysaccharide.

5 As used herein, the term "free polysaccharide" means a capsular polysaccharide that is not covalently conjugated to the carrier protein, but is nevertheless present in the capsular polysaccharide-carrier protein conjugate composition. The free polysaccharide may be non-covalently associated with (i.e., non-covalently bound to, adsorbed to, or entrapped in or with) the polysaccharide-carrier protein conjugate.

10 The percentage of free polysaccharide is measured after the final purification of the serotype 10A, 22F or 33F capsular polysaccharide-carrier protein conjugate. Preferably it is measured within 4 weeks after the final purification. It is expressed as a percentage of the total polysaccharide in the sample.

15 As used herein, "to conjugate," "conjugated" and "conjugating" refer to a process whereby a *Streptococcus pneumoniae* capsular polysaccharide, is covalently attached to a carrier protein.

20 The term "subject" refers to a mammal, including a human, or to a bird, fish, reptile, amphibian or any other animal. The term "subject" also includes household pets or research animals. Non-limiting examples of household pets and research animals include: dogs, cats, pigs, rabbits, rats, mice, gerbils, hamsters, guinea pigs, ferrets, monkeys, birds, snakes, lizards, fish, turtles, and frogs. The term "subject" also includes livestock animals. Non-limiting examples of livestock animals include: alpaca, bison, camel, cattle, deer, pigs, horses, llamas, mules, donkeys, sheep, goats, rabbits, reindeer, yak, chickens, geese, and  
25 turkeys.

In the preparation of multivalent conjugate pneumococcal vaccines directed to the prevention of invasive diseases caused by the organism *Streptococcus pneumoniae* (also known as pneumococcus), selected *Streptococcus pneumoniae* serotypes are grown to supply  
30 polysaccharides needed to produce the vaccine. The cells are grown in fermentors with lysis induced at the end of the fermentation by addition of sodium deoxycholate or an alternate lysing agent. The lysate broth is then harvested for downstream purification and the recovery of the capsular polysaccharide which surrounds the bacterial cells. After activation and conjugation with a carrier protein, the polysaccharide is included in the final vaccine product  
35 and confers immunity in the vaccine's target population to the selected *Streptococcus pneumoniae* serotypes.

The immunogenicity of a polysaccharide:carrier protein conjugate depends on several factors including the size of the polysaccharide. Polysaccharide size is a quality attribute assayed for in each preparation batch and must be appropriately controlled. High molecular weight capsular polysaccharides are able to induce certain antibody immune responses due to a higher valence of the epitopes present on the antigenic surface. It is generally preferable to prevent or to limit undesirable reduction of the size of the polysaccharide during the preparation of the conjugate to retain the immunogenicity of the polysaccharide. In addition, it is important to reduce the batch to batch variability of the polysaccharide Molecular Weight during the activation step and subsequent conjugation in order to maintain the consistency of the quality attributes of the conjugate.

Reductive amination chemistry (RAC) has been demonstrated by the inventors as a suitable process for preparing the *Streptococcus pneumoniae* serotype 10A, 22F or 33F capsular polysaccharide-protein carrier conjugate. The RAC approach involves activation of the polysaccharide by oxidation and subsequent conjugation of the activated polysaccharide to a protein carrier by reduction. Serotype 10A, 33F and 22F polysaccharides have proven to be particularly sensitive polysaccharides and are susceptible to degradation and size reduction during the oxidation step of the preparation of the conjugate. In addition, the use of hitherto known activation process produced activated serotype 10A, 33F and 22F polysaccharide of variable molecular weights.

Therefore, there is an important need for a process for preparing activated *Streptococcus pneumoniae* serotype 10A, 22F or 33F polysaccharides having a controlled and consistent molecular weight.

An object of the present invention is a process for preparing activated *Streptococcus pneumoniae* serotype 10A, 22F or 33F polysaccharides. In particular, an object of the invention is a process for preparing an activated *Streptococcus pneumoniae* serotype 10A, 22F or 33F polysaccharide, said process comprising the steps of:

- (a) reacting isolated serotype 10A, 22F or 33F polysaccharide with an oxidizing agent; and,
- (b) quenching the oxidation reaction by addition of a quenching agent resulting in an activated *Streptococcus pneumoniae* serotype 10A, 22F or 33F polysaccharide.

#### Isolation and purification of serotype 10A, 22F or 33F polysaccharide

The structures of serotype 10A, 22F and 33F polysaccharides are disclosed in the literature (see for example Kamerling J P C. Pneumococcal polysaccharides: a chemical view. In:

Tomasz A, editor. *Streptococcus pneumoniae*, molecular biology and mechanisms of disease. New York, N.Y: Mary Ann Liebert, Inc.; 2000. pp. 81–114).

5 As shown in figure 1, the polysaccharide repeating unit of serotype 22F consists of a branched pentasaccharide backbone (one glucuronic acid (GlcA), one glucopyranose (GlcP) and one galactofuranose (GalF) and two rhamnopyranoses (Rhap)) with a  $\alpha$ GlcP branch linked to the C3 hydroxyl group of  $\beta$ Rhap 45. Approximately 80% of the C2 hydroxyl groups of the  $\beta$ Rhap residue in the polysaccharide repeating unit are O-acetylated.

10 As shown in figure 2, the polysaccharide repeating unit of serotype 10A consists of a two D-galactosefuranose, three D-galactopyranose, one N-acetylgalactosamine, and a phosphate bond linked with saccharide chain through one D-ribitol.

15 As shown in figure 3, the polysaccharide repeating unit of serotype 33F consists of three D-galactopyranose, two D-galactofuranose, and one D-glucopyranose. Notably, 5-D-galactofuranosyl residues are O-acetylated in about 90% of 33F polysaccharide repeating units.

20 Serotype 10A, 22F and 33F capsular polysaccharides can be obtained directly from bacteria using isolation procedures known to one of ordinary skill in the art (see for example methods disclosed U.S. Patent App. Pub. Nos. 20060228380, 20060228381, 20070184071, 20070184072, 20070231340, and 20080102498 or WO2008118752). In addition, they can be produced using synthetic protocols.

25 Serotype 10A, 22F and 33F *Streptococcus pneumoniae* strains used to make the respective polysaccharides that are used in the immunogenic conjugates of the invention may be obtained from established culture collections or clinical specimens.

30 The purified serotype 10A, 22F and 33F polysaccharides are obtained by methods well known to the man skilled in the art. The bacterial cells are preferably grown in a soy based medium. Following fermentation of bacterial cells that produce *Streptococcus pneumoniae* serotype 10A, 22F or 33F capsular polysaccharides, the bacterial cells are lysed to produce a cell lysate. The bacterial cells may be lysed using any lytic agent. A "lytic agent" is any agent that aids in cell wall breakdown and release of autolysin which causes cellular lysis  
35 including, for example, detergents. As used herein, the term "detergent" refers to any anionic or cationic detergent capable of inducing lysis of bacterial cells. Representative examples of



such detergents for use within the methods of the present invention include deoxycholate sodium (DOC), N-lauroyl sarcosine, chenodeoxycholic acid sodium, and saponins.

In one embodiment of the present invention, the lytic agent used for lysing bacterial cells is DOC. DOC is the sodium salt of the bile acid deoxycholic acid, which is commonly derived from biological sources such as cows or oxen. DOC activates the LytA protein, which is an autolysin that is involved in cell wall growth and division in *Streptococcus pneumoniae*. The LytA protein has choline binding domains in its C-terminal portion, and mutations of the lytA gene are known to produce LytA mutants that are resistant to lysis with DOC.

In one embodiment of the present invention, the lytic agent used for lysing bacterial cells is a non-animal derived lytic agent. Non-animal derived lytic agents for use within the methods of the present invention include agents from non-animal sources with modes of action similar to that of DOC (i. e., that affect LytA function and result in lysis of *Streptococcus pneumoniae* cells). Such non-animal derived lytic agents include, but are not limited to, analogs of DOC, surfactants, detergents, and structural analogs of choline. In one embodiment, the non-animal derived lytic agent is selected from the group consisting of decanesulfonic acid, tert-octylphenoxy poly(oxyethylene)ethanols (e.g. Igepal® CA-630, CAS #: 9002-93-1, available from Sigma Aldrich, St. Louis, MO), octylphenol ethylene oxide condensates (e.g. Triton® X-100, available from Sigma Aldrich, St. Louis, MO), N-lauroyl sarcosine sodium, lauryl iminodipropionate, sodium dodecyl sulfate, chenodeoxycholate, hyodeoxycholate, glycodeoxycholate, taurodeoxycholate, taurochenodeoxycholate, and cholate. In another embodiment, the non-animal derived lytic agent is N-lauroyl sarcosine. In another embodiment, the lytic agent is N-lauroyl sarcosine sodium.

The capsular polysaccharides may then be purified from the cell lysate using purification techniques known in the art, including the use of centrifugation, depth filtration, precipitation, ultra-filtration, treatment with activate carbon, diafiltration and/or column chromatography (See, for example, U.S. Patent App. Pub. Nos. 20060228380, 20060228381, 20070184071, 20070184072, 20070231340, and 20080102498 or WO2008118752). The purified serotype 10A, 22F or 33F polysaccharides can then be used for the preparation of immunogenic conjugates.

Preferably, in order to generate serotype 22F conjugates or serotype 33F conjugates with advantageous filterability characteristics and/or yields, sizing of the polysaccharide to a lower molecular weight (MW) range is performed prior to the conjugation to a carrier protein. Advantageously, the size of the purified serotype 22F polysaccharide or serotype 33F polysaccharide is reduced while preserving critical features of the structure of the

polysaccharide such as for example the presence of O-acetyl groups. Preferably, the size of the purified serotype 22F polysaccharide or serotype 33F polysaccharide is reduced by mechanical homogenization.

5 In a preferred embodiment, the size of the purified polysaccharide is reduced by high pressure homogenization. High pressure homogenization achieves high shear rates by pumping the process stream through a flow path with sufficiently small dimensions. The shear rate is increased by using a larger applied homogenization pressure and exposure time can be increased by recirculating the feed stream through the homogenizer.

10

The high pressure homogenization process is particularly appropriate for reducing the size of the purified serotype 22F polysaccharide or serotype 33F polysaccharide while preserving the structural features of the polysaccharide such as the presence of O-acetyl groups.

15 The isolated serotype 22F polysaccharide obtained by purification of serotype 22F capsular polysaccharide from the *Streptococcus pneumoniae* lysate and optionally sizing of the purified polysaccharide can be characterized by different parameters including for example the molecular weight, and the mM of acetate per mM of serotype 22F polysaccharide.

In a preferred embodiment, the sized serotype 22F polysaccharide has a molecular weight  
20 comprised between 400 and 700kDa.

The isolated serotype 33F polysaccharide obtained by purification of serotype 33F capsular polysaccharide from the *Streptococcus pneumoniae* lysate and optionally sizing of the purified polysaccharide can be characterized by different parameters including for example  
25 the molecular weight, and the mM of acetate per mM of serotype 33F polysaccharide.

The degree of O-acetylation of the polysaccharide can be determined by any method known in the art, for example, by proton NMR (Lemerminier and Jones (1996) Carbohydrate Research 296; 83-96, Jones and Lemerminier (2002) J. Pharmaceutical and Biomedical  
30 Analysis 30; 1233-1247, WO 05/033148 or WO00/56357. Another commonly used method is described in Hestrin (1949) J. Biol. Chem. 180; 249-261. Preferably, the presence of O-acetyl groups is determined by ion-HPLC analysis.

In a preferred embodiment, the isolated serotype 22F polysaccharide is obtained by a  
35 process comprising the steps of:

- preparing a fermentation culture of serotype 22F *Streptococcus pneumoniae* bacterial cells;
- lysing the bacterial cells in said fermentation culture; and,

- purifying serotype 22F polysaccharide from the fermentation culture.

In a preferred embodiment, the isolated serotype 22F polysaccharide is obtained by a process comprising the steps of:

- 5 - preparing a fermentation culture of serotype 22F *Streptococcus pneumonia* bacterial cells;  
- lysing the bacterial cells in said fermentation culture;  
- purifying serotype 22F polysaccharide from the fermentation culture; and,  
- sizing the serotype 22F polysaccharide by mechanical sizing, preferably by high pressure homogenization.
- 10 In a preferred embodiment, the isolated serotype 22F polysaccharide is obtained by a process comprising the steps of:  
- preparing a fermentation culture of serotype 22F *Streptococcus pneumonia* bacterial cells;  
- lysing the bacterial cells in said fermentation culture;  
- purifying serotype 22F polysaccharide from the fermentation culture; and,  
15 - sizing the serotype 22F polysaccharide by mechanical sizing, preferably by high pressure homogenization,  
wherein the isolated serotype 22F polysaccharide has a molecular weight comprised between 400 and 700kDa.

- 20 In a preferred embodiment, the isolated serotype 33F polysaccharide is obtained by a process comprising the steps of:  
- preparing a fermentation culture of serotype 33F *Streptococcus pneumonia* bacterial cells;  
- lysing the bacterial cells in said fermentation culture; and,  
- purifying serotype 33F polysaccharide from the fermentation culture.

25

In a preferred embodiment, the isolated serotype 33F polysaccharide is obtained by a process comprising the steps of:

- preparing a fermentation culture of serotype 33F *Streptococcus pneumonia* bacterial cells;  
- lysing the bacterial cells in said fermentation culture;  
30 - purifying serotype 33F polysaccharide from the fermentation culture; and,  
- sizing the serotype 33F polysaccharide by mechanical sizing, preferably by high pressure homogenization.

The isolated serotype 10A polysaccharide obtained by purification of serotype 10A capsular  
35 polysaccharide from the *Streptococcus pneumoniae* lysate and optionally sizing of the purified polysaccharide can be characterized by different parameters including for example the molecular weight. Preferably, the isolated serotype 10A polysaccharide is not sized.

In a preferred embodiment, the isolated serotype 10A polysaccharide is obtained by a process comprising the steps of:

- preparing a fermentation culture of serotype 10A *Streptococcus pneumoniae* bacterial cells;
- 5 - lysing the bacterial cells in said fermentation culture; and,
- purifying serotype 10A polysaccharide from the fermentation culture.

Activation of serotype 10A, 22F or 33F polysaccharide

10 The isolated serotype 10A, 22F or 33F polysaccharide is activated by a process comprising the step of:

(a) reacting an isolated serotype 10A, 22F or 33F polysaccharide with an oxidizing agent; and,

(b) quenching the oxidation reaction by addition of a quenching agent resulting in an  
15 activated *Streptococcus pneumoniae* serotype 10A, 22F or 33F polysaccharide.

In a preferred embodiment, the concentration of serotype 10A, 22F or 33F polysaccharide in step (a) is between 0.1 and 10 mg/mL. In a preferred embodiment, the concentration of serotype 10A, 22F or 33F polysaccharide in step (a) is between 0.5 and 5 mg/mL. In a  
20 preferred embodiment, the concentration of serotype 10A, 22F or 33F polysaccharide in step (a) is between 1 and 3 mg/mL. In a preferred embodiment, the concentration of serotype 22F or 33F polysaccharide in step (a) is about 2 mg/mL. In a preferred embodiment, the concentration of serotype 10A polysaccharide in step (a) is about 2.5 mg/mL.

25 In a preferred embodiment, the oxidizing agent is periodate. The periodate oxidises vicinal hydroxyl groups to form carbonyl or aldehyde groups and causes cleavage of a C-C bond.

The term 'periodate' includes both periodate and periodic acid. This term also includes both metaperiodate ( $\text{IO}_4^-$ ) and orthoperiodate ( $\text{IO}_6^{5-}$ ). The term 'periodate' also includes the various salts of periodate including sodium periodate and potassium periodate. In a preferred  
30 embodiment, the oxidizing agent is sodium periodate. In a preferred embodiment, the periodate used for the oxidation of serotype 10A, 22F or 33F polysaccharide is metaperiodate. In a preferred embodiment the periodate used for the oxidation of serotype 10A, 22F or 33F polysaccharide is sodium metaperiodate.

35 In a preferred embodiment, the polysaccharide is reacted with 0.01 to 10, 0.05 to 5, 0.1 to 1, 0.5 to 1, 0.7 to 0.8, 0.01 to 0.2, 0.05 to 0.5, 0.05 to 0.2, 0.1 to 0.3 molar equivalents of oxidizing agent. In a preferred embodiment, the polysaccharide is reacted with about 0.05,

0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, In a preferred embodiment, the polysaccharide is reacted with about 0.10 molar equivalent of oxidizing agent. In a preferred embodiment, the polysaccharide is reacted with about 0.15 molar equivalent of oxidizing agent. In a preferred embodiment, the polysaccharide is reacted with about 0.25 molar equivalent of oxidizing agent. In a preferred embodiment, the polysaccharide is reacted with about 0.5 molar equivalent of oxidizing agent. In a preferred embodiment, the polysaccharide is reacted with about 0.8 molar equivalent of oxidizing agent.

10 In a preferred embodiment, the duration of the reaction in step (a) is between 1 and 50, 1 and 40, 1 and 30, 1 and 25, 1 and 20, 1 and 10, 10 to 50, 10 to 40, 10 to 30, 10 to 20 hours. In a preferred embodiment, when the polysaccharide is serotype 10A polysaccharide, the duration of the reaction in step (a) is between 1 to 10 hours. In a preferred embodiment, when the polysaccharide is serotype 10A polysaccharide, the duration of the reaction in step  
15 (a) is about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 hours. In a preferred embodiment, when the polysaccharide is serotype 10A polysaccharide, the duration of the reaction in step (a) is about 4 hours. In a preferred embodiment, when the polysaccharide is serotype 22F polysaccharide, the duration of the reaction in step (a) is between 10 and 20 hours. In a preferred embodiment, when the polysaccharide is serotype 22F polysaccharide, the duration of the reaction in step  
20 (a) is about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 hours. In a preferred embodiment, when the polysaccharide is serotype 33F polysaccharide, the duration of the reaction in step (a) is between 15 to 25 hours. In a preferred embodiment, when the polysaccharide is serotype 33F polysaccharide, the duration of the reaction in step (a) is about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or 25 hours. In a preferred embodiment, when the polysaccharide is serotype 33F polysaccharide, the duration of the reaction in step  
25 (a) is about 20 hours.

In a preferred embodiment, the temperature of the reaction in step (a) is maintained between 1 to 30°C, 1 to 10°C, 10 to 20°C, 20 to 30°C, 2 to 8°C. In a preferred embodiment, the  
30 temperature of the reaction in step (a) is maintained at about 5°C.


In a preferred embodiment, step (a) is carried out in a buffer selected from sodium phosphate, potassium phosphate, 2-(N-morpholino)ethanesulfonic acid (MES) or Bis-Tris. In a preferred embodiment step (a) is carried out in potassium phosphate buffer. In a preferred  
35 embodiment step (a) is carried out in sodium phosphate buffer.

In a preferred embodiment the buffer has a concentration of between 1 to 500 mM, 1 to 300mM, 50 to 200mM. In a preferred embodiment the buffer has a concentration of about 100mM.

- 5 In a preferred embodiment the pH in step (a) is between 4 and 8, 5 and 7, 5.5 and 6.5. In a preferred embodiment the pH in step (a) is about 6. In a preferred embodiment the pH in step (a) is about 5.8.

10 In one embodiment, the quenching agent is selected from vicinal diols, 1,2-aminoalcohols, amino acids, glutathione, sulfite, bisulfate, dithionite, metabisulfite, thiosulfate, phosphites, hypophosphites or phosphorous acid.

In one embodiment, the quenching agent is a 1,2-aminoalcohols of formula



- 15 where R<sup>1</sup> is selected from H, methyl, ethyl, propyl or isopropyl.

In one embodiment, the quenching agent is selected from sodium and potassium salts of sulfite, bisulfate, dithionite, metabisulfite, thiosulfate, phosphites, hypophosphites or phosphorous acid.

20


In one embodiment, the quenching agent is an amino acid. In one embodiment, said amino acid is selected from serine, threonine, cysteine, cystine, methionine, proline, hydroxyproline, tryptophan, tyrosine, and histidine.

- 25 In one embodiment, the quenching agent is a sulfite such as bisulfate, dithionite, metabisulfite, thiosulfate.

In one embodiment, the quenching agent is a compound comprising two vicinal hydroxyl groups (vicinal diols), i.e two hydroxyl groups covalently linked to two adjacent carbon atoms.

30

Preferably, the quenching agent is a compound of formula (II)



wherein R<sup>1</sup> and R<sup>2</sup> are each independently selected from H, methyl, ethyl, propyl or isopropyl.

In a preferred embodiment, the quenching agent is glycerol, ethylene glycol, propan-1,2-diol, butan-1,2-diol or butan-2,3-diol, ascorbic acid. In a preferred embodiment, the quenching agent is butan-2,3-diol.

In a preferred embodiment, the oxidizing agent is neutralized by the addition of 0.1 to 10, 0.5 to 5, 0.5 to 3, 0.5 to 2 molar equivalents of quenching agent. In a preferred embodiment, the oxidizing agent is neutralized by the addition of about 0.5, 1, 1.5, 2, 2.5, 3 molar equivalents of quenching agent. In a preferred embodiment, the oxidizing agent is neutralized by the addition of about 2 molar equivalents of quenching agent.

In a preferred embodiment, the duration of step (b) is between 0.1 and 10, 0.5 and 5, or 0.5 and 2 hours. In a preferred embodiment, the duration of step (b) is about 0.5, 1, 1.5, 2.5 or 3 hours.

In a preferred embodiment, the temperature of the reaction in step (b) is maintained between 1 and 30°C, 1 and 25°C, 1 and 20°C, 1 and 10°C, 10 and 20°C, 2 and 8°C. In a preferred embodiment, the temperature of the reaction in step (b) is maintained at about 1, 2, 3, 4, 5, 6, 7 or 8°C.

In a preferred embodiment the pH in step (b) is between 4 and 8, 5 and 7, 5.5 and 6.5. In a preferred embodiment the pH in step (a) is about 6.

In a preferred embodiment, the activated serotype 10A, 22F or 33F polysaccharide is purified. The activated serotype 10A, 22F or 33F polysaccharide can be purified according to methods known to the man skilled in the art such as gel permeation chromatography (GPC), dialysis or ultrafiltration/diafiltration. For example, the activated polysaccharide is purified by concentration and diafiltration using an ultrafiltration device.

In preferred embodiment, the isolated serotype 10A polysaccharide is activated by a process comprising the step of:

(a) reacting isolated serotype 10A polysaccharide with periodate; and,

(b) quenching the oxidation reaction by addition of butan-2,3-diol resulting in an activated serotype 10A polysaccharide.

In preferred embodiment, the isolated serotype 10A polysaccharide is activated by a process comprising the step of:

- (a) reacting isolated serotype 10A polysaccharide with periodate at a temperature between 2 and 8°C; and,
- 5 (b) quenching the oxidation reaction by addition of butan-2,3-diol at a temperature between 2 and 8°C resulting in an activated serotype 10A polysaccharide.

In preferred embodiment, the isolated serotype 10A polysaccharide is activated by a process comprising the step of:

- 10 (a) reacting isolated serotype 10A polysaccharide with 0.2 to 0.3 molar equivalent periodate at a temperature between 2 and 8°C; and,
- (b) quenching the oxidation reaction by addition of 0.5 to 2 molar equivalent of butan-2,3-diol at a temperature between 2 and 8°C resulting in an activated serotype 10A polysaccharide.

15 In preferred embodiment, the isolated serotype 22F polysaccharide is activated by a process comprising the step of:

- (a) reacting isolated serotype 22F polysaccharide with periodate; and,
- (b) quenching the oxidation reaction by addition of butan-2,3-diol resulting in an activated serotype 22F polysaccharide.

20

In preferred embodiment, the isolated serotype 22F polysaccharide is activated by a process comprising the step of:

- (a) reacting isolated serotype 22F polysaccharide with periodate at a temperature between 2 and 8°C; and,
- 25 (b) quenching the oxidation reaction by addition of butan-2,3-diol at a temperature between 2 and 8°C resulting in an activated serotype 22F polysaccharide.

In preferred embodiment, the isolated serotype 22F polysaccharide is activated by a process comprising the step of:

- 30 (a) reacting isolated serotype 22F polysaccharide with 0.05 to 0.2 molar equivalent periodate at a temperature between 2 and 8°C; and,
- (b) quenching the oxidation reaction by addition of 2 to 3, preferably 2, molar equivalents of butan-2,3-diol at a temperature between 2 and 8°C resulting in an activated serotype 22F polysaccharide.

35

In preferred embodiment, the isolated serotype 33F polysaccharide is activated by a process comprising the step of:



(a) reacting isolated serotype 33F polysaccharide with periodate; and,  
(b) quenching the oxidation reaction by addition of butan-2,3-diol resulting in an activated serotype 33F polysaccharide.

5 In preferred embodiment, the isolated serotype 33F polysaccharide is activated by a process comprising the step of:

(a) reacting isolated serotype 33F polysaccharide with periodate at a temperature between 2 and 8°C; and,

(b) quenching the oxidation reaction by addition of butan-2,3-diol at a temperature between 2 and 8°C resulting in an activated serotype 33F polysaccharide.

In preferred embodiment, the isolated serotype 33F polysaccharide is activated by a process comprising the step of:

(a) reacting isolated serotype 33F polysaccharide with 0.05 to 0.2 molar equivalent periodate at a temperature between 2 and 8°C; and,

(b) quenching the oxidation reaction by addition of 0.5 to 1.5 molar equivalent of butan-2,3-diol at a temperature between 2 and 8°C resulting in an activated serotype 33F polysaccharide.

20 In a preferred embodiment, the invention relates to an activated serotype 10A capsular polysaccharide obtained or obtainable by the above disclosed process.

In a preferred embodiment, the invention relates to an activated serotype 22F capsular polysaccharide obtained or obtainable by the above disclosed process.

In a preferred embodiment, the invention relates to an activated serotype 33F capsular polysaccharide obtained or obtainable by the above disclosed process.

In a preferred embodiment the activated serotype 10A polysaccharide obtained from step (b) has a molecular weight between 20 and 100%, 30 and 95%, 40 and 95%, 60 to 95%, 85 to 95%, or about 90%, of the molecular weight of the isolated polysaccharide used in step (a).

30 In a preferred embodiment the activated serotype 10A polysaccharide obtained from step (b) has a molecular weight of at least 20, 25, 30, 35, 40 or 45% of the molecular weight of the isolated polysaccharide used in step (a). In a preferred embodiment the activated serotype 10A polysaccharide obtained from step (b) has a molecular weight of at least 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99% of the molecular weight of the isolated polysaccharide used in step (a).

In a preferred embodiment the activated serotype 22F polysaccharide obtained from step (b) has a molecular weight between 50 and 100%, 60 to 95%, 85 to 95% or about 90% of the molecular weight of the isolated polysaccharide used in step (a).

5

In a preferred embodiment the activated serotype 22F polysaccharide obtained from step (b) has a molecular weight of at least 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99% of the molecular weight of the isolated polysaccharide used in step (a).

10

In a preferred embodiment the activated serotype 33F polysaccharide obtained from step (b) has a molecular weight between 50 and 100%, 60 to 95%, 85 to 95%, or about 90% of the molecular weight of the isolated polysaccharide used in step (a).

15

In a preferred embodiment the activated serotype 33F polysaccharide obtained from step (b) has a molecular weight of at least 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99% of the molecular weight of the isolated polysaccharide used in step (a).

20

In a preferred embodiment the degree of oxidation of the activated serotype 10A, 22F or 33F polysaccharide is between 2 and 30, 2 and 25, 2 and 20, 2 and 15, 2 and 10, 2 and 5, 5 and 30, 5 and 25, 5 and 20, 5 and 15, 5 and 10, 10 and 30, 10 and 25, 10 and 20, 10 and 15, 5 and 25, 10 and 30, 15 and 30, 15 and 25, 15 and 20, 20 to 30, 20 to 25. In a preferred embodiment the degree of oxidation of the activated serotype 10A, 22F or 33F polysaccharide is between 2 and 10, 4 and 8, 4 and 6, 6 and 8, 6 and 12, 8 and 14, 9 and 11, 10 and 16, 12 and 16, 14 and 18, 16 and 20, 16 and 18, 18 and 22, 18 and 20.

25

In a preferred embodiment, the activated serotype 10A polysaccharide has a molecular weight between 50 and 400, 50 and 350, 50 and 300, 50 and 250, 50 and 200, 100 and 300, 100 and 250 or 100 and 200kDa. In a preferred embodiment, the activated serotype 10A polysaccharide has a molecular weight between 50 and 300kDa. In a preferred embodiment, the activated serotype 10A polysaccharide has a molecular weight between 100 and 200kDa. In a preferred embodiment, the activated serotype 10A polysaccharide has a molecular weight between 100 and 200kDa and a degree of oxidation between 5 and 20, 5 and 15, 8 and 14, 8 and 12 or 9 and 11. In a preferred embodiment, the activated serotype

30

35

10A polysaccharide has a molecular weight between 100 and 200kDa and a degree of oxidation between 9 and 11.

5 In a preferred embodiment, the activated serotype 22F polysaccharide has a molecular weight between 25 and 1000, 100 and 1000, 300 and 800, 300 and 700, 300 and 600, 400 and 1000, 400 and 800, 400 and 700 or 400 and 600kDa. In a preferred embodiment, the activated serotype 22F polysaccharide has a molecular weight between 300 and 800kDa. In a preferred embodiment, the activated serotype 22F polysaccharide has a molecular weight between 400 and 800kDa. In a preferred embodiment, the activated serotype 22F polysaccharide has a molecular weight between 400 and 600kDa. In a preferred embodiment, the activated serotype 22F polysaccharide has a molecular weight between 400 and 800kDa and a degree of oxidation between 10 and 25, 10 and 20, 12 and 20 or 14 and 18. In a preferred embodiment, the activated serotype 22F polysaccharide has a molecular weight between 400 and 800kDa and a degree of oxidation between 10 and 20.

15

In a preferred embodiment, the activated serotype 22F polysaccharide comprises at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6 or 0.7 or about 0.8 mM acetate per mM serotype 22F polysaccharide. In a preferred embodiment, the activated serotype 22F polysaccharide comprises at least 0.5, 0.6 or 0.7 mM acetate per mM serotype 22F polysaccharide. In a preferred embodiment, the activated serotype 22F polysaccharide comprises at least 0.6 mM acetate per mM serotype 22F polysaccharide. In a preferred embodiment, the activated serotype 22F polysaccharide comprises at least 0.7 mM acetate per mM serotype 22F polysaccharide.

20

25 In a preferred embodiment, the activated serotype 22F polysaccharide has a molecular weight between 400 and 800 kDa and comprises at least 0.6 mM acetate per mM serotype 22F polysaccharide.

In a preferred embodiment, the activated serotype 22F polysaccharide has a molecular weight between 400 and 800kDa, a degree of oxidation between 12 and 20 and comprises at least 0.6 mM acetate per mM serotype 22F polysaccharide.

30

In a preferred embodiment, the activated serotype 33F polysaccharide has a molecular weight between 50 and 1250, 200 and 1200, 500 and 1200, 500 and 1000, 700 and 1200, 800 and 1200, 800 and 1100, 900 and 1200, 800 and 1000Da. In a preferred embodiment, the activated serotype 33F polysaccharide has a molecular weight between 500 and

35

1000kDa. In another preferred embodiment, the activated serotype 33F polysaccharide has a molecular weight between 50 and 600, 50 and 500 or 50 and 400kDa.

In a preferred embodiment, the activated serotype 33F polysaccharide comprises at least  
5 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7 or 0.8 or about 0.9 mM acetate per mM serotype 33F polysaccharide. In a preferred embodiment, the activated serotype 33F polysaccharide comprises at least 0.6, 0.7 or 0.8 mM acetate per mM serotype 33F polysaccharide. In a preferred embodiment, the activated serotype 33F polysaccharide comprises at least 0.6 mM acetate per mM serotype 33F polysaccharide. In a preferred embodiment, the activated  
10 serotype 33F polysaccharide comprises at least 0.7 mM acetate per mM serotype 33F polysaccharide.

In a preferred embodiment, the activated serotype 33F polysaccharide has a molecular weight between 800 and 1200 kDa and comprises at least 0.7 mM acetate per mM serotype  
15 33F polysaccharide. In a preferred embodiment, the activated serotype 33F polysaccharide has a molecular weight between 800 and 1200 kDa, comprises at least 0.7 mM acetate per mM serotype 33F polysaccharide and has a degree of oxidation between 10 and 20.

In a preferred embodiment, the activated serotype 33F polysaccharide has a molecular weight between 50 and 200 kDa and comprises at least 0.7 mM acetate per mM serotype  
20 33F polysaccharide.

In a preferred embodiment, the activated serotype 33F polysaccharide has a molecular weight between 50 and 200 kDa, comprises at least 0.7 mM acetate per mM serotype 33F  
25 polysaccharide and has a degree of oxidation between 10 and 20.

In an embodiment, the activated serotype 10A, 22F or 33F polysaccharide is lyophilized, optionally in the presence of a cryoprotectant/lyoprotectant. In a preferred embodiment, the cryoprotectant/lyoprotectant is selected from sucrose, trehalose, raffinose, stachyose,  
30 melezitose, dextran, mannitol, lactitol and palatinit. In a preferred embodiment, the cryoprotectant/lyoprotectant is sucrose. The lyophilized activated polysaccharide can then be compounded with a solution comprising the carrier protein.

In another embodiment, the activated serotype 10A, 22F or 33F polysaccharide is compounded with the carrier protein and lyophilized optionally in the presence of a cryoprotectant/lyoprotectant. In a preferred embodiment, the cryoprotectant/lyoprotectant is selected from sucrose, trehalose, raffinose, stachyose, melezitose, dextran, mannitol, lactitol

and palatinit. In a preferred embodiment, the cryoprotectant/lyoprotectant is sucrose. The co-lyophilized polysaccharide and carrier protein can then be resuspended in solution and reacted with a reducing agent.

- 5 In an embodiment, the invention relates to a lyophilized activated serotype 10A polysaccharide. In an embodiment, the invention relates to a lyophilized activated serotype 22F polysaccharide. In an embodiment, the invention relates to a lyophilized activated serotype 33F polysaccharide.
- 10 In an embodiment the invention relates to the co-lyophilized activated serotype 10A polysaccharide and protein carrier. In a preferred embodiment, the protein carrier is CRM<sub>197</sub>. In an embodiment the invention relates to the co-lyophilized activated serotype 22F polysaccharide and protein carrier. In a preferred embodiment, the protein carrier is CRM<sub>197</sub>. In an embodiment the invention relates to the co-lyophilized activated serotype 33F
- 15 polysaccharide and protein carrier. In a preferred embodiment, the protein carrier is CRM<sub>197</sub>.

#### Conjugation of activated serotype 10A, 22F or 33F polysaccharide with a carrier protein

- The activated serotype 10A, 22F or 33F polysaccharide disclosed herein can be conjugated
- 20 to a carrier protein by a process comprising the steps of:
- (a) compounding the activated serotype 10A, 22F or 33F polysaccharide with a carrier protein; and,
- (b) reacting the compounded activated serotype 10A, 22F or 33F polysaccharide and carrier protein with a reducing agent to form a serotype 10A, 22F or 33F polysaccharide:carrier
- 25 protein conjugate.

- The conjugation of activated serotype 22F or 33F polysaccharide with a protein carrier by reductive amination in dimethylsulfoxide (DMSO) is suitable to preserve the O-acetyl content of the polysaccharide as compared for example to reductive amination in aqueous phase
- 30 where the level of O-acetylation of the polysaccharide is significantly reduced. In a preferred embodiment, step (a) and step (b) are carried out in DMSO.

- In a preferred embodiment, step (a) comprises dissolving lyophilized serotype 10A, 22F or 33F polysaccharide in an aqueous solution comprising a carrier protein or in a solution
- 35 comprising a carrier protein and DMSO. In a preferred embodiment, step (a) comprises dissolving co-lyophilized serotype 10A, 22F or 33F polysaccharide and carrier protein in an aqueous solution or in DMSO.

When step (a) and (b) are carried out in an aqueous solution, said solution comprises a buffer, preferably selected from buffer, preferably selected from PBS, MES, HEPES, Bis-tris, ADA, PIPES. MOPSO, BES, MOPS, DIPSO, MOBS, HEPPSO, POPSO, TEA, EPPS, Bicine  
5 or HEPB, at a pH between 6.0 and 8.5, 7 and 8 or 7 and 7.5. In a preferred embodiment the buffer is PBS. In a preferred embodiment the pH is about 7.3.

In a preferred embodiment, the concentration of activated serotype 10A, 22F or 33F polysaccharide in step (a) is between 0.1 and 10 mg/mL, 0.5 and 5 mg/mL, 0.5 and 2 mg/mL.  
10 In a preferred embodiment, the concentration of activated serotype 10A, 22F or 33F polysaccharide in step (a) is about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9 or 3 mg/mL.

In a preferred embodiment the initial ratio (weight by weight) of activated serotype 10A, 22F  
15 or 33F polysaccharide to carrier protein is between 5:1 and 0.1:1, 2:1 and 0.1:1, 2:1 and 1:1, 1.5:1 and 1:1, 0.1:1 and 1:1, 0.3:1 and 1:1, 0.6:1 and 1:1.2. In a preferred embodiment the initial ratio of activated serotype 10A, 22F or 33F polysaccharide to carrier protein is about 0.6:1 and 1:1.2. In a preferred embodiment the initial ratio of activated serotype 10A, 22F or 33F polysaccharide to carrier protein is about 0.4:1, 0.5:1, 0.6:1, 0.7:1, 0.8:1, 0.9:1, 1:1,  
20 1.1:1, 1.2:1, 1.3:1, 1.4:1, 1.5:1, 1.6:1, 1.7:1, 1.8:1, 1.9:1, 2:1.

In a preferred embodiment, in step (b), the activated serotype 10A, 22F or 33F polysaccharide is reacted with between 0.1 and 10 molar equivalents, 0.5 and 5 molar equivalents 0.5 and 2.5 molar equivalents of reducing agent. In a preferred embodiment, in  
25 step (b), the activated serotype 10A, 22F or 33F is reacted with about 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4 or 2.5 molar equivalents of reducing agent.

In an embodiment, the reducing agent is sodium cyanoborohydride, sodium  
30 triacetoxymborohydride, sodium or zinc borohydride in the presence of Bronsted or Lewis acids, amine boranes such as pyridine borane, 2-Picoline Borane, 2,6-diborane-methanol, dimethylamine-borane, t-BuMe<sup>i</sup>PrN-BH<sub>3</sub>, benzylamine-BH<sub>3</sub> or 5-ethyl-2-methylpyridine borane (PEMB). In a preferred embodiment, the reducing agent is sodium cyanoborohydride.

35 In a preferred embodiment, the duration of step (b) is between 1 and 50, 5 and 30, 10 and 30, 15 and 30, 20 and 30, 5 and 25, 10 and 25, 15 and 25 hours. In a preferred embodiment, the duration of step (b) is about 20, 21, 22, 23, 24, 25, 26, 27, 28 hours.

In a preferred embodiment, the temperature of the reaction in step (b) is maintained between 10 and 40°C, 15 and 30°C, 20 and 26°C, 21 and 25°C. In a preferred embodiment, the temperature of the reaction in step (b) is maintained at about 21, 22, 23, 24 or 25°C.

5

In a preferred embodiment, the process for the preparation of an immunogenic conjugate comprising serotype 10A, 22F or 33F polysaccharide covalently linked to a carrier protein further comprises a step (step c) of capping unreacted aldehyde (quenching) by addition of NaBH<sub>4</sub>.

10

In a preferred embodiment, in step (c), the unreacted aldehydes are capped by the addition of 0.1 to 10 molar equivalents, 0.5 to 5 molar equivalents or 1 to 3 molar equivalents of NaBH<sub>4</sub>. In a preferred embodiment, in step (c), the unreacted aldehydes are capped by the addition of about 2 molar equivalents of NaBH<sub>4</sub>.

15

In a preferred embodiment, the duration of step (c) is between 0.1 and 10, 0.5 and 5 or 2 and 4 hours. In a preferred embodiment, the duration of step (c) is about 3 hours.

In a preferred embodiment, the temperature of the reaction in step (c) is maintained between 10 to 40°C, 15 to 30°C or 20 to 26°C. In a preferred embodiment, the temperature of the reaction in step (c) is maintained at about 23°C.

20

After conjugation of serotype 10A, 22F or 33F polysaccharide to the carrier protein, the polysaccharide-protein conjugate can be purified (enriched with respect to the amount of polysaccharide-protein conjugate) by a variety of techniques known to the skilled person. These techniques include dialysis, concentration/diafiltration operations, tangential flow filtration precipitation/elution, column chromatography (DEAE or hydrophobic interaction chromatography), and depth filtration.

25

In a preferred embodiment the carrier protein is non-toxic and non-reactogenic and obtainable in sufficient amount and purity. Carrier proteins should be amenable to standard conjugation procedures.

30

In a preferred embodiment, the activated serotype 10A, 22F or 33F polysaccharide is conjugated to a carrier protein which is selected in the group consisting of: DT (Diphtheria toxin), TT (tetanus toxid) or fragment C of TT, CRM197 (a nontoxic but antigenically identical variant of diphtheria toxin) other DT point mutants, such as CRM176, CRM228, CRM 45

35

(Uchida et al J. Biol. Chem. 218; 3838-3844, 1973); CRM 9, CRM102, CRM 103 and CRM107 and other mutations described by Nicholls and Youle in Genetically Engineered Toxins, Ed: Frankel, Maecel Dekker Inc, 1992; deletion or mutation of Glu-148 to Asp, Gln or Ser and/or Ala 158 to Gly and other mutations disclosed in US 4709017 or US 4950740; 5 mutation of at least one or more residues Lys 516, Lys 526, Phe 530 and/or Lys 534 and other mutations disclosed in US 5917017 or US 6455673; or fragment disclosed in US 5843711, pneumococcal pneumolysin (Kuo et al (1995) Infect Immun 63; 2706-13) including ply detoxified in some fashion for example dPLY-GMBS (WO 04081515, PCT/EP2005/010258) or dPLY-formol, PhtX, including PhtA, PhtB, PhtD, PhtE (sequences 10 of PhtA, PhtB, PhtD or PhtE are disclosed in WO 00/37105 or WO 00/39299) and fusions of Pht proteins for example PhtDE fusions, PhtBE fusions, Pht A-E (WO 01/98334, WO 03/54007, W02009/000826), OMPC (meningococcal outer membrane protein - usually extracted from N.meningitidis serogroup B - EP0372501 ), PorB (from N. meningitidis), PD (Haemophilus influenza protein D - see, e.g., EP 0 594 610 B), or immunologically functional 15 equivalents thereof, synthetic peptides (EP0378881 , EP0427347), heat shock proteins (WO 93/17712, WO 94/03208), pertussis proteins (WO 98/58668, EP0471 177), cytokines, lymphokines, growth factors or hormones (WO10 91/01146), artificial proteins comprising multiple human CD4+ T cell epitopes from various pathogen derived antigens (Falugi et al (2001 ) Eur J Immunol 31 ; 3816-3824) such as N19 protein (Baraldoi et al (2004) Infect 20 Immun 72; 4884-7) pneumococcal surface protein PspA (WO 02/091998), iron uptake proteins (WO 01/72337), toxin A or B of C. difficile (WO 00/61761 ). In an embodiment, the activated serotype 10A, 22F or 33F polysaccharide is conjugated to DT (Diphtheria toxoid). In another embodiment, the activated serotype 10A, 22F or 33F polysaccharide is 25 conjugated to TT (tetanus toxoid). In another embodiment, the activated serotype 10A, 22F or 33F polysaccharide is conjugated to fragment C of TT. In another embodiment, the activated serotype 10A, 22F or 33F polysaccharide is conjugated to PD (Haemophilus influenza protein D - see, e.g., EP 0 594 610 B).

In a preferred embodiment, the activated serotype 10A, 22F or 33F polysaccharide is 30 conjugated to CRM<sub>197</sub> protein. The CRM<sub>197</sub> protein is a nontoxic form of diphtheria toxin but is immunologically indistinguishable from the diphtheria toxin. CRM<sub>197</sub> is produced by C. diphtheriae infected by the nontoxigenic phage  $\beta$ 197<sup>tox</sup> created by nitrosoguanidine mutagenesis of the toxigenic corynephage beta (Uchida, T. et al. 1971, Nature New Biology 233:8-11). CRM<sub>197</sub> is purified through ultrafiltration, ammonium sulfate precipitation, and ion- 35 exchange chromatography. The CRM<sub>197</sub> protein has the same molecular weight as the diphtheria toxin but differs therefrom by a single base change (guanine to adenine) in the structural gene. This single base change causes an amino acid substitution glutamic acid for



glycine) in the mature protein and eliminates the toxic properties of diphtheria toxin. The CRM<sub>197</sub> protein is a safe and effective T-cell dependent carrier for saccharides. Further details about CRM<sub>197</sub> and production thereof can be found e.g. in US 5,614,382.

- 5 In a preferred embodiment, the activated serotype 10A, 22F or 33F polysaccharide disclosed herein is conjugated to CRM<sub>197</sub> by a process comprising the step of:
- (a) compounding the activated serotype 10A, 22F or 33F polysaccharide with CRM<sub>197</sub>;
  - (b) reacting the compounded activated serotype 10A, 22F or 33F polysaccharide and CRM<sub>197</sub> with sodium cyanoborohydride to form a serotype 10A, 22F or 33F polysaccharide:CRM<sub>197</sub> conjugate.
- 10

In a preferred embodiment, the activated serotype 10A, 22F or 33F polysaccharide disclosed herein is conjugated to CRM<sub>197</sub> by a process comprising the step of:

- (a) compounding the activated serotype 10A, 22F or 33F polysaccharide with a CRM<sub>197</sub>;
  - 15 (b) reacting the compounded activated serotype 10A, 22F or 33F polysaccharide and CRM<sub>197</sub> with sodium cyanoborohydride to form a serotype 10A, 22F or 33F polysaccharide:CRM<sub>197</sub> conjugate;
- wherein steps (a) and (b) are carried out in DMSO.

- 20 In a preferred embodiment, the activated serotype 10A, 22F or 33F polysaccharide disclosed herein is conjugated to CRM<sub>197</sub> by a process comprising the step of:
- (a) compounding the activated serotype 10A, 22F or 33F polysaccharide with a CRM<sub>197</sub>;
  - (b) reacting the compounded activated serotype 10A, 22F or 33F polysaccharide and CRM<sub>197</sub> with 0.5 to 2 molar equivalent sodium cyanoborohydride to form a serotype 10A, 22F or 33F polysaccharide:CRM<sub>197</sub> conjugate;
- 25 wherein steps (a) and (b) are carried out in DMSO.

- In an embodiment, the invention relates to an immunogenic conjugate obtained or obtainable by a process disclosed herein. In a preferred embodiment, the invention relates to
- 30 an immunogenic conjugate obtained or obtainable by conjugating a serotype 10A, 22F or 33F activated polysaccharide as disclosed herein to a carrier protein by reductive amination. In a preferred embodiment, the invention relates to an immunogenic conjugate obtained or obtainable by conjugating a serotype 10A, 22F or 33F activated polysaccharide as disclosed herein to a carrier protein by reductive amination in DMSO. In a preferred embodiment, the
- 35 carrier protein is CRM<sub>197</sub>.

Serotype 10A immunogenic conjugate

In a preferred embodiment, the serotype 10A immunogenic conjugate has a molecular weight between 500 and 15000; 500 and 10000; 2000 and 10000; or 3000 and 8000kDa. In

5 a preferred embodiment, the serotype 10A immunogenic conjugate has a molecular weight between 3000 and 8000kDa. The molecular weight of the immunogenic conjugate is measured by SEC-MALLS.

In a preferred embodiment, the serotype 10A immunogenic conjugate comprises less than  
10 about 50, 45, 40, 35, 30, 25, 20 or 15% of free serotype 10A polysaccharide compared to the total amount of serotype 10A polysaccharide. In a preferred embodiment the serotype 10A immunogenic conjugate comprises less than about 25% of free serotype 10A polysaccharide compared to the total amount of serotype 10A polysaccharide. In a preferred embodiment the serotype 10A immunogenic conjugate comprises less than about 20% of free serotype  
15 10A polysaccharide compared to the total amount of serotype 10A polysaccharide. In a preferred embodiment the serotype 10A immunogenic conjugate comprises less than about 15% of free serotype 10A polysaccharide compared to the total amount of serotype 10A polysaccharide.

20 In a preferred embodiment, the ratio (weight by weight) of serotype 10A polysaccharide to carrier protein in the conjugate is between 0.5 and 3. In a preferred embodiment, the ratio of serotype 10A polysaccharide to carrier protein in the conjugate is between 0.5 and 2, 0.5 and 1.5, 0.5 and 1, 1 and 1.5, 1 and 2. In a preferred embodiment, the ratio of serotype 10A polysaccharide to carrier protein in the conjugate is between 0.8 and 1.4. In a preferred  
25 embodiment, the ratio of serotype 10A capsular polysaccharide to carrier protein in the conjugate is between 0.8 and 1.2.

Size exclusion chromatography media (CL-4B) can be used to determine the relative molecular size distribution of the conjugate. Size Exclusion Chromatography (SEC) is used in  
30 gravity fed columns to profile the molecular size distribution of conjugates. Large molecules excluded from the pores in the media elute more quickly than small molecules. Fraction collectors are used to collect the column eluate. The fractions are tested colorimetrically by saccharide assay. For the determination of  $K_d$ , columns are calibrated to establish the fraction at which molecules are fully excluded ( $V_0$ ), ( $K_d=0$ ), and the fraction representing the maximum retention ( $V_i$ ), ( $K_d=1$ ). The fraction at which a specified sample attribute is  
35 reached ( $V_e$ ), is related to  $K_d$  by the expression,  $K_d = (V_e - V_0) / (V_i - V_0)$ .

In a preferred embodiment, at least 30% of the serotype 10A immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 40% of the immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, or 85% of the serotype  
5 10A immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 60% of the serotype 10A immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 50% and 80% of the serotype 10A immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column.

10

The degree of conjugation is the number of lysine residues in the carrier protein that are conjugated to the polysaccharide of interest. The evidence for lysine modification of the carrier protein, due to covalent linkages to the polysaccharides, is obtained by amino acid analysis using routine methods known to those of skill in the art. Conjugation results in a  
15 reduction in the number of lysine residues recovered, compared to the CRM<sub>197</sub> protein starting material used to generate the conjugate materials.

20

In a preferred embodiment, the degree of conjugation of the immunogenic conjugate is between 2 and 15, 2 and 13, 2 and 10, 2 and 8, 2 and 6, 2 and 5, 2 and 4, 3 and 15, 3 and  
20 13, 3 and 10, 3 and 8, 3 and 6, 3 and 5, 3 and 4, 5 and 15, 5 and 10, 8 and 15, 8 and 12, 10 and 15 or 10 and 12. In a preferred embodiment, the degree of conjugation of the immunogenic conjugate is between 6 and 8.

#### Serotype 22F immunogenic conjugate

25

In a preferred embodiment, the immunogenic conjugate comprises at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6 or 0.7 or about 0.8 mM acetate per mM serotype 22F polysaccharide. In a preferred embodiment, the immunogenic conjugate comprises at least 0.5, 0.6 or 0.7 mM acetate per  
30 mM serotype 22F polysaccharide. In a preferred embodiment, the immunogenic conjugate comprises at least 0.6 mM acetate per mM serotype 22F polysaccharide. In a preferred embodiment, the immunogenic conjugate comprises at least 0.7 mM acetate per mM serotype 22F polysaccharide.

35

In a preferred embodiment, the ratio of mM acetate per mM serotype 22F polysaccharide in the immunogenic conjugate to mM acetate per mM serotype 22F polysaccharide in the isolated polysaccharide is at least 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95. In a preferred embodiment, the ratio of mM acetate per mM serotype 22F polysaccharide in the

immunogenic conjugate to mM acetate per mM serotype 22F polysaccharide in the isolated polysaccharide is at least 0.7. In a preferred embodiment, the ratio of mM acetate per mM serotype 22F polysaccharide in the immunogenic conjugate to mM acetate per mM serotype 22F polysaccharide in the isolated polysaccharide is at least 0.9.

5

In a preferred embodiment, the ratio of mM acetate per mM serotype 22F polysaccharide in the immunogenic conjugate to mM acetate per mM serotype 22F polysaccharide in the activated polysaccharide is at least 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95. In a preferred embodiment, the ratio of mM acetate per mM serotype 22F polysaccharide in the immunogenic conjugate to mM acetate per mM serotype 22F polysaccharide in the activated polysaccharide is at least 0.7. In a preferred embodiment, the ratio of mM acetate per mM serotype 22F polysaccharide in the immunogenic conjugate to mM acetate per mM serotype 22F polysaccharide in the activated polysaccharide is at least 0.9.

10

In a preferred embodiment, the serotype 22F immunogenic conjugate has a molecular weight between 400 and 15000; 500 and 10000; 2000 and 10000kDa; 3000 and 8000kDa; or 3000 and 5000kDa. In a preferred embodiment, the serotype 22F immunogenic conjugate has a molecular weight between 3000 and 5000kDa. The molecular weight of the immunogenic conjugate is measured by SEC-MALLS.

20

In a preferred embodiment, the serotype 22F immunogenic conjugate comprises less than about 50, 45, 40, 35, 30, 25, 20 or 15% of free serotype 22F polysaccharide compared to the total amount of serotype 22F polysaccharide. . In a preferred embodiment the serotype 22F immunogenic conjugate comprises less than about 40% of free serotype 22F polysaccharide compared to the total amount of serotype 22F polysaccharide. In a preferred embodiment the serotype 22F immunogenic conjugate comprises less than about 25% of free serotype 22F polysaccharide compared to the total amount of serotype 22F polysaccharide. In a preferred embodiment the serotype 22F immunogenic conjugate comprises less than about 20% of free serotype 22F polysaccharide compared to the total amount of serotype 22F polysaccharide. In a preferred embodiment the serotype 22F immunogenic conjugate comprises less than about 15% of free serotype 22F polysaccharide compared to the total amount of serotype 22F polysaccharide.

25

30

In a preferred embodiment, the ratio (weight by weight) of serotype 22F polysaccharide to carrier protein in the conjugate is between 0.5 and 3. In a preferred embodiment, the ratio of serotype 22F polysaccharide to carrier protein in the conjugate is between 0.5 and 2, 0.5 and 1.5, 0.8 and 1.2, 0.5 and 1, 1 and 1.5, or 1 and 2. In a preferred embodiment, the ratio of

35

serotype 22F polysaccharide to carrier protein in the conjugate is between 0.8 and 1.2. In a preferred embodiment, the ratio of serotype 22F capsular polysaccharide to carrier protein in the conjugate is between 0.9 and 1.1.

5 In a preferred embodiment, at least 30% of the serotype 22F immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 40% of the immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column. In a preferred  
10 embodiment, at least 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, or 85% of the serotype 22F immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 60% of the serotype 22F immunogenic conjugate has a Kd  
below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 50% and 80% of the serotype 22F immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 65% and 80% of the serotype 22F  
immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column.

15

In a preferred embodiment, the degree of conjugation of the immunogenic conjugate is between 2 and 15, 2 and 13, 2 and 10, 2 and 8, 2 and 6, 2 and 5, 2 and 4, 3 and 15, 3 and  
13, 3 and 10, 3 and 8, 3 and 6, 3 and 5, 3 and 4, 4 and 7, 5 and 15, 5 and 10, 8 and 15, 8 and  
12, 10 and 15 or 10 and 12. In a preferred embodiment, the degree of conjugation of the  
20 immunogenic conjugate is between 4 and 7.

#### Serotype 33F immunogenic conjugate

In a preferred embodiment, the immunogenic conjugate comprises at least 0.1, 0.2, 0.3, 0.4,  
25 0.5, 0.6 or 0.7 or about 0.8 mM acetate per mM serotype 33F polysaccharide. In a preferred  
embodiment, the immunogenic conjugate comprises at least 0.5, 0.6 or 0.7 mM acetate per  
mM serotype 33F polysaccharide. In a preferred embodiment, the immunogenic conjugate  
comprises at least 0.6 mM acetate per mM serotype 33F polysaccharide. In a preferred  
embodiment, the immunogenic conjugate comprises at least 0.7 mM acetate per mM  
30 serotype 33F polysaccharide.

In a preferred embodiment, the ratio of mM acetate per mM serotype 33F polysaccharide in  
the immunogenic conjugate to mM acetate per mM serotype 33F polysaccharide in the  
isolated polysaccharide is at least 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95. In a preferred  
35 embodiment, the ratio of mM acetate per mM serotype 33F polysaccharide in the  
immunogenic conjugate to mM acetate per mM serotype 33F polysaccharide in the isolated  
polysaccharide is at least 0.7. In a preferred embodiment, the ratio of mM acetate per mM

serotype 33F polysaccharide in the immunogenic conjugate to mM acetate per mM serotype 33F polysaccharide in the isolated polysaccharide is at least 0.9.

5 In a preferred embodiment, the ratio of mM acetate per mM serotype 33F polysaccharide in the immunogenic conjugate to mM acetate per mM serotype 33F polysaccharide in the activated polysaccharide is at least 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95. In a preferred embodiment, the ratio of mM acetate per mM serotype 33F polysaccharide in the immunogenic conjugate to mM acetate per mM serotype 33F polysaccharide in the activated polysaccharide is at least 0.7. In a preferred embodiment, the ratio of mM acetate per mM serotype 33F polysaccharide in the immunogenic conjugate to mM acetate per mM serotype 33F polysaccharide in the activated polysaccharide is at least 0.9.

15 In a preferred embodiment, the serotype 33F immunogenic conjugate has a molecular weight between 500 and 30000; 500 and 25000; 500 and 20000; 500 and 15000; 500 and 10000; 1000 and 10000; 1000 and 8000; 1000 and 5000; 2000 and 10000kDa; 2000 and 8000; or 2000 and 5000kDa. In a preferred embodiment, the serotype 33F immunogenic conjugate has a molecular weight between 1000 and 5000kDa. The molecular weight of the immunogenic conjugate is measured by SEC-MALLS.

20 In a preferred embodiment, the serotype 33F immunogenic conjugate comprises less than about 50, 45, 40, 35, 30, 25, 20 or 15% of free serotype 33F polysaccharide compared to the total amount of serotype 33F polysaccharide. In a preferred embodiment the serotype 33F immunogenic conjugate comprises less than about 25% of free serotype 33F polysaccharide compared to the total amount of serotype 33F polysaccharide. In a preferred embodiment the serotype 33F immunogenic conjugate comprises less than about 20% of free serotype 33F polysaccharide compared to the total amount of serotype 33F polysaccharide. In a preferred embodiment the serotype 33F immunogenic conjugate comprises less than about 15% of free serotype 33F polysaccharide compared to the total amount of serotype 33F polysaccharide.

30 In a preferred embodiment, the ratio (weight by weight) of serotype 33F polysaccharide to carrier protein in the conjugate is between 0.4 and 3. In a preferred embodiment, the ratio of serotype 33F polysaccharide to carrier protein in the conjugate is between 0.5 and 2, 0.5 and 1.5, 0.5 and 1, 1 and 1.5, 1 and 2. In a preferred embodiment, the ratio of serotype 33F polysaccharide to carrier protein in the conjugate is between 0.5 and 1.5. In a preferred embodiment, the ratio of serotype 33F capsular polysaccharide to carrier protein in the conjugate is between 0.5 and 1.2.

In a preferred embodiment, at least 30% of the serotype 33F immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 40% of the immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 45%, 50%, 55%, 60%, 65% of the serotype 33F immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 60% of the serotype 33F immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 40% and 80% of the serotype 33F immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 45% and 65% of the serotype 33F immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column.

In a preferred embodiment, the degree of conjugation of the immunogenic conjugate is between 1 and 15, 2 and 13, 2 and 10, 2 and 8, 2 and 6, 2 and 5, 2 and 4, 3 and 15, 3 and 13, 3 and 10, 3 and 8, 3 and 6, 3 and 5, 3 and 4, 5 and 15, 5 and 10, 8 and 15, 8 and 12, 10 and 15 or 10 and 12. In a preferred embodiment, the degree of conjugation of the immunogenic conjugate is between 3 and 6.

#### Immunogenic composition

20

The term "immunogenic composition" relates to any pharmaceutical composition containing an antigen, e.g., a microorganism or a component thereof, which composition can be used to elicit an immune response in a subject.

As used herein, "immunogenic" means an ability of an antigen (or an epitope of the antigen), such as a bacterial capsular polysaccharide, or a glycoconjugate or immunogenic composition comprising an antigen, to elicit an immune response in a host such as a mammal, either humorally or cellularly mediated, or both.

In a preferred embodiment, the immunogenic composition comprises a serotype 10A, 22F or 33F conjugate obtained by a process disclosed herein. In a preferred embodiment, the immunogenic composition comprising a serotype 10A, 22F or 33F conjugate obtainable by a process disclosed herein.

In an embodiment, the immunogenic composition disclosed herein, when administered to a subject, induces the formation of antibodies capable of binding to serotype 10A *Streptococcus pneumoniae*. In an embodiment, the immunogenic composition disclosed

herein, when administered to a subject, induces the formation of antibodies capable of binding to serotype 10A *Streptococcus pneumoniae* as measured by a standard ELISA assay.

5 In an embodiment, the immunogenic composition disclosed herein, when administered to a subject, induces the formation of antibodies capable of killing serotype 10A *Streptococcus pneumoniae* in an opsonophagocytosis assay as disclosed herein.

10 In an embodiment, the immunogenic composition disclosed herein, when administered to a subject, induces the formation of antibodies capable of binding to serotype 22F *Streptococcus pneumoniae*. In an embodiment, the immunogenic composition disclosed herein, when administered to a subject, induces the formation of antibodies capable of binding to serotype 22F *Streptococcus pneumoniae* as measured by a standard ELISA assay.

15 In an embodiment, the immunogenic composition disclosed herein, when administered to a subject, induces the formation of antibodies capable of killing serotype 22F *Streptococcus pneumoniae* in an opsonophagocytosis assay as disclosed herein.

20 In an embodiment, the immunogenic composition disclosed herein, when administered to a subject, induces the formation of antibodies capable of binding to serotype 33F *Streptococcus pneumoniae*. In an embodiment, the immunogenic composition disclosed herein, when administered to a subject, induces the formation of antibodies capable of binding to serotype 33F *Streptococcus pneumoniae* as measured by a standard ELISA assay.

25 In an embodiment, the immunogenic composition disclosed herein, when administered to a subject, induces the formation of antibodies capable of killing serotype 33F *Streptococcus pneumoniae* in an opsonophagocytosis assay as disclosed herein.

30 Formulation of the immunogenic composition of the present invention can be accomplished using art-recognized methods. For instance, the serotype 10A, 22F or 33F conjugates can be formulated with a physiologically acceptable vehicle to prepare the composition. Examples of such vehicles include, but are not limited to, water, buffered saline, polyols (e.g., glycerol, propylene glycol, liquid polyethylene glycol) and dextrose solutions.

35 In a preferred embodiment, the immunogenic composition may comprise at least one additional antigen. In a preferred embodiment, the immunogenic composition may comprises at least one additional *Streptococcus pneumoniae* capsular polysaccharide.



In a preferred embodiment, the immunogenic composition may comprise at least one additional *Streptococcus pneumoniae* capsular polysaccharide conjugated to a carrier protein. In a preferred embodiment, said carrier protein is CRM<sub>197</sub>.

- 5 In certain embodiments, the immunogenic composition comprises one or more adjuvants. As defined herein, an "adjuvant" is a substance that serves to enhance the immunogenicity of an immunogenic composition of this invention. Thus, adjuvants are often given to boost the immune response and are well known to the skilled artisan. Suitable adjuvants to enhance effectiveness of the composition include, but are not limited to:
- 10 (1) aluminum salts (alum), such as aluminum hydroxide, aluminum phosphate, aluminum sulfate, etc.;
- (2) oil-in-water emulsion formulations (with or without other specific immunostimulating agents such as muramyl peptides (defined below) or bacterial cell wall components), such as, for example,
- 15 (a) MF59 (PCT Pub. No. WO 90/14837), containing 5% Squalene, 0.5% Tween 80, and 0.5% Span 85 (optionally containing various amounts of MTP-PE (see below, although not required)) formulated into submicron particles using a microfluidizer such as Model 110Y microfluidizer (Microfluidics, Newton, MA),
- (b) SAF, containing 10% Squalene, 0.4% Tween 80, 5% pluronic-blocked polymer L121, and
- 20 thr-MDP (see below) either microfluidized into a submicron emulsion or vortexed to generate a larger particle size emulsion, and
- (c) Ribi™ adjuvant system (RAS), (Corixa, Hamilton, MT) containing 2% Squalene, 0.2% Tween 80, and one or more bacterial cell wall components from the group consisting of 3-O-deacylated monophosphorylipid A (MPL™) described in U.S. Patent No. 4,912,094 (Corixa),
- 25 trehalose dimycolate (TDM), and cell wall skeleton (CWS), preferably MPL + CWS (Detox™);
- (3) saponin adjuvants, such as Quil A or STIMULON™ QS-21 (Antigenics, Framingham, MA) (U.S. Patent No. 5,057,540) may be used or particles generated therefrom such as ISCOMs (immunostimulating complexes);
- (4) bacterial lipopolysaccharides, synthetic lipid A analogs such as aminoalkyl glucosamine
- 30 phosphate compounds (AGP), or derivatives or analogs thereof, which are available from Corixa, and which are described in U.S. Patent No. 6,113,918; one such AGP is 2-[(R)-3-Tetradecanoyloxytetradecanoylamino]ethyl 2-Deoxy-4-O-phosphono-3-O-[(R)-3-tetradecanoyloxytetradecanoyl]-2-[(R)-3-tetradecanoyloxytetradecanoylamino]-β-D-glucopyranoside, which is also known as 529 (formerly known as RC529), which is formulated
- 35 as an aqueous form or as a stable emulsion, synthetic polynucleotides such as oligonucleotides containing CpG motif(s) (U.S. Patent No. 6,207,646);

(5) cytokines, such as interleukins (e.g., IL-1, IL-2, IL-4, IL-5, IL-6, IL-7, IL-12, IL-15, IL-18, etc.), interferons (e.g., gamma interferon), granulocyte macrophage colony stimulating factor (GM-CSF), macrophage colony stimulating factor (M-CSF), tumor necrosis factor (TNF), costimulatory molecules 87-1 and 87-2, etc.;

- 5 (6) detoxified mutants of a bacterial ADP-ribosylating toxin such as a cholera toxin (CT) either in a wild-type or mutant form, for example, where the glutamic acid at amino acid position 29 is replaced by another amino acid, preferably a histidine, in accordance with published international patent application number WO 00/18434 (see also WO 02/098368 and WO 02/098369), a pertussis toxin (PT), or an E. coli heat-labile toxin (LT), particularly  
10 LT-K63, LT-R72, CT-S109, PT-K9/G129 (see, e.g., WO 93/13302 and WO 92/19265); and  
(7) other substances that act as immunostimulating agents to enhance the effectiveness of the composition.

Muramyl peptides include, but are not limited to, N-acetyl-muramyl-L-threonyl-D-isoglutamine  
15 (thr-MDP), N-acetyl-normuramyl-L-alanine-2-(1'-2'-dipalmitoyl-*sn*-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE), etc.

In an embodiment of the present invention, the immunogenic compositions as disclosed herein comprise a CpG Oligonucleotide as adjuvant. A CpG oligonucleotide as used herein  
20 refers to an immunostimulatory CpG oligodeoxynucleotide (CpG ODN), and accordingly these terms are used interchangeably unless otherwise indicated. Immunostimulatory CpG oligodeoxynucleotides contain one or more immunostimulatory CpG motifs that are unmethylated cytosine-guanine dinucleotides, optionally within certain preferred base contexts. The methylation status of the CpG immunostimulatory motif generally refers to the  
25 cytosine residue in the dinucleotide. An immunostimulatory oligonucleotide containing at least one unmethylated CpG dinucleotide is an oligonucleotide which contains a 5' unmethylated cytosine linked by a phosphate bond to a 3' guanine, and which activates the immune system through binding to Toll-like receptor 9 (TLR-9). In another embodiment the immunostimulatory oligonucleotide may contain one or more methylated CpG dinucleotides,  
30 which will activate the immune system through TLR9 but not as strongly as if the CpG motif(s) was/were unmethylated. CpG immunostimulatory oligonucleotides may comprise one or more palindromes that in turn may encompass the CpG dinucleotide. CpG oligonucleotides have been described in a number of issued patents, published patent applications, and other publications, including U.S. Patent Nos. 6,194,388; 6,207,646;  
35 6,214,806; 6,218,371; 6,239,116; and 6,339,068.

In an embodiment of the present invention, the immunogenic compositions as disclosed herein comprise any of the CpG Oligonucleotide described at pages 3 lines 22 to page 12 line 36 of WO2010/125480.

5 Different classes of CpG immunostimulatory oligonucleotides have been identified. These are referred to as A, B, C and P class, and are described in greater detail at pages 3 lines 22 to page 12 line 36 of WO2010/125480. Methods of the invention embrace the use of these different classes of CpG immunostimulatory oligonucleotides.

In an embodiment of the present invention, the immunogenic compositions as disclosed herein comprise an A class CpG Oligonucleotide. Preferably, the "A class" CpG  
10 oligonucleotide of the invention has the following nucleic acid sequence: 5' GGGGACGACGTCGTGGGGGGG 3' (SEQ ID NO: 1). Some non-limiting examples of A-Class oligonucleotides include: 5' G\*G\*G\_G\_A\_C\_G\_A\_C\_G\_T\_C\_G\_T\_G\_G\*G\*G\*G\*G\*G 3' (SEQ ID NO: 2) ; wherein \* refers to a phosphorothioate bond and \_ refers to a phosphodiester bond.

15 In an embodiment of the present invention, the immunogenic compositions as disclosed herein comprise a B class CpG Oligonucleotide. In one embodiment, the CpG oligonucleotide for use in the present invention is a B class CpG oligonucleotide represented by at least the formula:

5' X<sub>1</sub>X<sub>2</sub>CGX<sub>3</sub>X<sub>4</sub> 3', wherein X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub>, and X<sub>4</sub> are nucleotides. In one embodiment, X<sub>2</sub> is adenine, guanine, or thymine. In another embodiment, X<sub>3</sub> is cytosine, adenine, or thymine.  
20

The B class CpG oligonucleotide sequences of the invention are those broadly described above in US patents Ps 6,194,388, 6,207,646, 6,214,806, 6,218,371, 6,239,116 and 6,339,068. Exemplary sequences include but are not limited to those disclosed in these latter applications and patents.

25 In an embodiment, the "B class" CpG oligonucleotide of the invention has the following nucleic acid sequence:

5' TCGTCGTTTTTCGGTGCTTTT 3' (SEQ ID NO: 3), or  
5' TCGTCGTTTTTCGGTCGTTTT 3' (SEQ ID NO: 4), or  
5' TCGTCGTTTTGTCGTTTTGTCGTT 3' (SEQ ID NO: 5), or  
30 5' TCGTCGTTTCGTCGTTTTGTCGTT 3' (SEQ ID NO: 6), or  
5' TCGTCGTTTTGTCGTTTTTTTCGA 3' (SEQ ID NO: 7).

In any of these sequences, all of the linkages may be all phosphorothioate bonds. In another embodiment, in any of these sequences, one or more of the linkages may be phosphodiester, preferably between the "C" and the "G" of the CpG motif making a semi-soft  
35 CpG oligonucleotide. In any of these sequences, an ethyl-uridine or a halogen may substitute for the 5' T; examples of halogen substitutions include but are not limited to bromo-uridine or iodo-uridine substitutions.

Some non-limiting examples of B-Class oligonucleotides include:

- 5' T\*C\*G\*T\*C\*G\*T\*T\*T\*T\*C\*G\*G\*T\*G\*C\*T\*T\*T 3' (SEQ ID NO: 8), or  
 5' T\*C\*G\*T\*C\*G\*T\*T\*T\*T\*C\*G\*G\*T\*G\*C\*T\*T\*T 3' (SEQ ID NO: 9), or  
 5' T\*C\*G\*T\*C\*G\*T\*T\*T\*T\*G\*T\*C\*G\*T\*T\*T\*G\*T\*C\*G\*T\*T 3' (SEQ ID NO: 10), or  
 5 5' T\*C\*G\*T\*C\*G\*T\*T\*T\*T\*C\*G\*T\*C\*G\*T\*T\*T\*G\*T\*C\*G\*T\*T 3' (SEQ ID NO: 11), or  
 5' T\*C\*G\*T\*C\*G\*T\*T\*T\*T\*G\*T\*C\*G\*T\*T\*T\*T\*T\*T\*T\*C\*G\*A 3' (SEQ ID NO: 12).

wherein \* refers to a phosphorothioate bond.

In an embodiment of the present invention, the immunogenic compositions as disclosed herein comprise a C class CpG Oligonucleotide. In an embodiment, the "C class" CpG  
 10 oligonucleotides of the invention has the following nucleic acid sequence:

- 5' TCGCGTCGTTTCGGCGCGCGCCG 3' (SEQ ID NO: 13), or  
 5' TCGTCGACGTTTCGGCGCGCGCCG 3' (SEQ ID NO: 14), or  
 5' TCGGACGTTTCGGCGCGCGCCG 3' (SEQ ID NO: 15), or  
 5' TCGGACGTTTCGGCGCGCCG 3' (SEQ ID NO: 16), or  
 15 5' TCGCGTCGTTTCGGCGCGCCG 3' (SEQ ID NO: 17), or  
 5' TCGACGTTTCGGCGCGCGCCG 3' (SEQ ID NO: 18), or  
 5' TCGACGTTTCGGCGCGCCG 3' (SEQ ID NO: 19), or  
 5' TCGCGTCGTTTCGGCGCCG 3' (SEQ ID NO: 20), or  
 5' TCGCGACGTTTCGGCGCGCGCCG 3' (SEQ ID NO: 21), or  
 20 5' TCGTCGTTTTTCGGCGCGCGCCG 3' (SEQ ID NO: 22), or  
 5' TCGTCGTTTTTCGGCGCGCCGCCG 3' (SEQ ID NO: 23), or  
 5' TCGTCGTTTTACGGCGCCGTGCCG 3' (SEQ ID NO: 24), or  
 5' TCGTCGTTTTTCGGCGCGCGCCGT 3' (SEQ ID NO: 25).

In any of these sequences, all of the linkages may be all phosphorothioate bonds. In another  
 25 embodiment, in any of these sequences, one or more of the linkages may be phosphodiester, preferably between the "C" and the "G" of the CpG motif making a semi-soft CpG oligonucleotide.

Some non-limiting examples of C-Class oligonucleotides include:

- 30 5' T\*C\_G\*C\_G\*T\*C\_G\*T\*T\*C\_G\*G\*C\*G\*C\_G\*C\*G\*C\*C\*G 3' (SEQ ID NO: 26), or  
 5' T\*C\_G\*T\*C\_G\*A\*C\_G\*T\*T\*C\_G\*G\*C\*G\*C\_G\*C\*G\*C\*C\*G 3' (SEQ ID NO: 27), or  
 5' T\*C\_G\*G\*A\*C\_G\*T\*T\*C\_G\*G\*C\*G\*C\_G\*C\*G\*C\*C\*G 3' (SEQ ID NO: 28), or  
 5' T\*C\_G\*G\*A\*C\_G\*T\*T\*C\_G\*G\*C\*G\*C\*G\*C\*C\*G 3' (SEQ ID NO: 29), or  
 5' T\*C\_G\*C\_G\*T\*C\_G\*T\*T\*C\_G\*G\*C\*G\*C\*G\*C\*C\*G 3' (SEQ ID NO: 30), or  
 35 5' T\*C\_G\*A\*C\_G\*T\*T\*C\_G\*G\*C\*G\*C\_G\*C\*G\*C\*C\*G 3' (SEQ ID NO: 31), or  
 5' T\*C\_G\*A\*C\_G\*T\*T\*C\_G\*G\*C\*G\*C\*G\*C\*C\*G 3' (SEQ ID NO: 32), or  
 5' T\*C\_G\*C\_G\*T\*C\_G\*T\*T\*C\_G\*G\*C\*G\*C\*C\*G 3' (SEQ ID NO: 33), or

5' T\*C\_G\*C\_G\*A\*C\_G\*T\*T\*C\_G\*G\*C\*G\*C\_G\*C\*G\*C\*C\*G 3' (SEQ ID NO: 34), or  
 5' T\*C\*G\*T\*C\*G\*T\*T\*T\*T\*C\*G\*G\*C\*G\*C\*G\*C\*G\*C\*C\*G 3' (SEQ ID NO: 35), or  
 5' T\*C\*G\*T\*C\*G\*T\*T\*T\*T\*C\*G\*G\*C\*G\*G\*C\*G\*C\*G\*C\*C\*G 3' (SEQ ID NO: 36), or  
 5' T\*C\*G\*T\*C\_G\*T\*T\*T\*T\*A\*C\_G\*G\*C\*G\*C\*C\_G\*T\*G\*C\*C\*G 3' (SEQ ID NO: 37), or  
 5 5' T\*C\_G\*T\*C\*G\*T\*T\*T\*T\*C\*G\*G\*C\*G\*C\*G\*C\*G\*C\*C\*G\*T 3' (SEQ ID NO: 38)

wherein \* refers to a phosphorothioate bond and \_ refers to a phosphodiester bond.

In any of these sequences, an ethyl-uridine or a halogen may substitute for the 5' T; examples of halogen substitutions include but are not limited to bromo-uridine or iodo-uridine substitutions.

10 In an embodiment of the present invention, the immunogenic compositions as disclosed herein comprise a P class CpG Oligonucleotide. In an embodiment, the CpG oligonucleotide for use in the present invention is a P class CpG oligonucleotide containing a 5' TLR activation domain and at least two palindromic regions, one palindromic region being a 5' palindromic region of at least 6 nucleotides in length and connected to a 3' palindromic  
 15 region of at least 8 nucleotides in length either directly or through a spacer, wherein the oligonucleotide includes at least one YpR dinucleotide. In an embodiment, said oligonucleotide is not T\*C\_G\*T\*C\_G\*A\*C\_G\*T\*T\*C\_G\*G\*C\*G\*C\_G\*C\*G\*C\*C\*G (SEQ ID NO: 27). In one embodiment the a P class CpG oligonucleotide includes at least one unmethylated CpG dinucleotide. In another embodiment the TLR activation domain is TCG,  
 20 TTCG, TTTTCG, TYpR, TTYpR, TTTYpR, UCG, UUCG, UUUCG, TTT, or TTTT. In yet another embodiment the TLR activation domain is within the 5' palindromic region. In another embodiment the TLR activation domain is immediately 5' to the 5' palindromic region.

In an embodiment, the "P class" CpG oligonucleotides of the invention has the following nucleic acid sequence: 5' TCGTCGACGATCGGCGCGGCCG 3' (SEQ ID NO: 39).

25 In said sequences, all of the linkages may be all phosphorothioate bonds. In another embodiment, one or more of the linkages may be phosphodiester, preferably between the "C" and the "G" of the CpG motif making a semi-soft CpG oligonucleotide. In any of these sequences, an ethyl-uridine or a halogen may substitute for the 5' T; examples of halogen substitutions include but are not limited to bromo-uridine or iodo-uridine substitutions.

30 A non-limiting example of P-Class oligonucleotides include:

5' T\*C\_G\*T\*C\_G\*A\*C\_G\*A\*T\*C\_G\*G\*C\*G\*C\_G\*C\*G\*C\*C\*G 3' (SEQ ID NO: 40)

wherein \* refers to a phosphorothioate bond and \_ refers to a phosphodiester bond.

In one embodiment the oligonucleotide includes at least one phosphorothioate linkage. In another embodiment all internucleotide linkages of the oligonucleotide are phosphorothioate  
 35 linkages. In another embodiment the oligonucleotide includes at least one phosphodiester-like linkage. In another embodiment the phosphodiester-like linkage is a phosphodiester

linkage. In another embodiment a lipophilic group is conjugated to the oligonucleotide. In one embodiment the lipophilic group is cholesterol.

In an embodiment, all the internucleotide linkage of the CpG oligonucleotides disclosed herein are phosphodiester bonds ("soft" oligonucleotides, as described in the PCT application WO2007/026190). In another embodiment, CpG oligonucleotides of the invention are rendered resistant to degradation (e.g., are stabilized). A "stabilized oligonucleotide" refers to an oligonucleotide that is relatively resistant to *in vivo* degradation (e.g. via an exo- or endo-nuclease). Nucleic acid stabilization can be accomplished via backbone modifications. Oligonucleotides having phosphorothioate linkages provide maximal activity and protect the oligonucleotide from degradation by intracellular exo- and endo-nucleases.

The immunostimulatory oligonucleotides may have a chimeric backbone, which have combinations of phosphodiester and phosphorothioate linkages. For purposes of the instant invention, a chimeric backbone refers to a partially stabilized backbone, wherein at least one internucleotide linkage is phosphodiester or phosphodiester-like, and wherein at least one other internucleotide linkage is a stabilized internucleotide linkage, wherein the at least one phosphodiester or phosphodiester-like linkage and the at least one stabilized linkage are different. When the phosphodiester linkage is preferentially located within the CpG motif such molecules are called "semi-soft" as described in the PCT application WO2007/026190.

The size of the CpG oligonucleotide (i.e., the number of nucleotide residues along the length of the oligonucleotide) also may contribute to the stimulatory activity of the oligonucleotide. For facilitating uptake into cells, CpG oligonucleotide of the invention preferably have a minimum length of 6 nucleotide residues. Oligonucleotides of any size greater than 6 nucleotides (even many kb long) are capable of inducing an immune response if sufficient immunostimulatory motifs are present, because larger oligonucleotides are degraded inside cells. In certain embodiments, the CpG oligonucleotides are 6 to 100 nucleotides long, preferentially 8 to 30 nucleotides long. In important embodiments, nucleic acids and oligonucleotides of the invention are not plasmids or expression vectors.

In an embodiment, the CpG oligonucleotides disclosed herein comprise substitutions or modifications, such as in the bases and/or sugars as described at paragraph 134 to 147 of WO2007/026190.

In an embodiment, the CpG oligonucleotide of the present invention is chemically modified. Examples of chemical modifications are known to the skilled person and are described, for example in Uhlmann E. et al. (1990), Chem. Rev. 90:543, S. Agrawal, Ed., Humana Press, Totowa, USA 1993; Crooke, S.T. et al. (1996) Annu. Rev. Pharmacol. Toxicol. 36:107-129; and Hunziker J. et al., (1995), Mod. Synth. Methods 7:331-417. An oligonucleotide according to the invention may have one or more modifications, wherein each modification is located at a particular phosphodiester internucleoside bridge and/or at a particular  $\beta$ -D-ribose

unit and/or at a particular natural nucleoside base position in comparison to an oligonucleotide of the same sequence which is composed of natural DNA or RNA.

In some embodiments of the invention, CpG-containing nucleic acids might be simply mixed with immunogenic carriers according to methods known to those skilled in the art (see, e.g. 5 WO03/024480).

In a particular embodiment of the present invention, any of the immunogenic composition disclosed herein comprises from 2µg to 100mg of CpG oligonucleotide, preferably from 0.1mg to 50 mg CpG oligonucleotide, preferably from 0.2mg to 10 mg CpG oligonucleotide, preferably from 0.3 mg to 5 mg CpG oligonucleotide, even preferably from 0.5 to 2 mg CpG 10 oligonucleotide, even preferably from 0.75 to 1.5 mg CpG oligonucleotide. In a preferred embodiment, the immunogenic composition disclosed herein comprises approximately 1mg CpG oligonucleotide.

In a preferred embodiment, the adjuvant is an aluminum-based adjuvant selected from the 15 group consisting of aluminum phosphate, aluminum sulfate and aluminum hydroxide. In one embodiment, the immunogenic compositions described herein comprise the adjuvant aluminum phosphate.

In a preferred embodiments, the immunogenic compositions of the invention further comprise 20 at least one of a buffer, a cryoprotectant, a salt, a divalent cation, a non-ionic detergent, an inhibitor of free radical oxidation, a diluent or a carrier.

The immunogenic composition optionally can comprise a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers include carriers approved by a regulatory 25 agency of a Federal, a state government, or other regulatory agency, or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, including humans as well as non-human mammals. The term carrier may be used to refer to a diluent, adjuvant, excipient, or vehicle with which the pharmaceutical composition is administered. Water, saline solutions and aqueous dextrose and glycerol solutions can be employed as 30 liquid carriers, particularly for injectable solutions. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E. W. Martin. The formulation should suit the mode of administration.

The immunogenic composition optionally can comprise one or more physiologically 35 acceptable buffers selected from, but not limited to Tris (trimethamine), phosphate, acetate, borate, citrate, glycine, histidine and succinate. In certain embodiments, the formulation is

buffered to within a pH range of about 5.0 to about 7.0, preferably from about 5.5 to about 6.5.

5 The immunogenic composition optionally can comprise one or more non-ionic surfactants, including but not limited to polyoxyethylene sorbitan fatty acid esters, Polysorbate-80 (Tween 80), Polysorbate-60 (Tween 60), Polysorbate-40 (Tween 40) and Polysorbate-20 (Tween 20), polyoxyethylene alkyl ethers, including but not limited to Brij 58, Brij 35, as well as others such as Triton X-100; Triton X- 114, NP40, Span 85 and the Pluronic series of non-ionic surfactants (e. g. , Pluronic 121), with preferred components Polysorbate-80 at a  
10 concentration from about 0.001% to about 2% (with up to about 0.25% being preferred) or Polysorbate-40 at a concentration from about 0.001% to 1% (with up to about 0.5% being preferred).

15 The invention further relates to vaccines comprising the immunogenic composition of the invention.

#### Methods for Inducing an Immune Response and Protecting Against Infection

The present disclosure also includes methods of use for immunogenic compositions described herein. For example, one embodiment of the disclosure provides a method of  
20 inducing an immune response against *Streptococcus pneumoniae*, comprising administering to a subject an immunogenic amount of any of the immunogenic compositions described herein.

One embodiment of the disclosure provides a method of protecting a subject against  
25 an infection with *Streptococcus pneumoniae*, or a method of preventing infection with *Streptococcus pneumoniae*, or a method of reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by *Streptococcus pneumoniae*, the methods comprising administering to a subject an immunogenic amount of any of the immunogenic compositions described herein.

30 One embodiment of the disclosure provides a method of protecting a subject against an infection with serotype 10A *Streptococcus pneumoniae*, or a method of preventing infection with serotype 10A *Streptococcus pneumoniae*, or a method of reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by  
35 serotype 10A *Streptococcus pneumoniae*, the methods comprising administering to a subject an immunogenic amount of any of the immunogenic compositions described herein.



One embodiment of the disclosure provides a method of treating or preventing a *Streptococcus pneumoniae* infection, disease or condition associated with serotype 10A *Streptococcus pneumoniae* in a subject, the method comprising the step of administering a therapeutically or prophylactically effective amount of an immunogenic composition described herein to the subject. Another embodiment provides a method of treating or preventing a *Streptococcus pneumoniae* infection, disease or condition associated with a serotype 10A *Streptococcus pneumoniae* in a subject, the method comprising generating a polyclonal or monoclonal antibody preparation from the immunogenic composition described herein, and using said antibody preparation to confer passive immunity to the subject.

In one embodiment, the disclosure relates to the use of the immunogenic conjugate or immunogenic composition disclosed herein for the manufacture of a medicament for protecting a subject against an infection with *Streptococcus pneumoniae*, and/or preventing infection with *Streptococcus pneumoniae*, and/or reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by *Streptococcus pneumoniae*, and/or protecting a subject against an infection with serotype 10A *Streptococcus pneumoniae* and/or preventing infection with serotype 10A *Streptococcus pneumoniae*, and/or reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by serotype 10A *Streptococcus pneumoniae*.

One embodiment of the disclosure provides a method of protecting a subject against an infection with serotype 22F *Streptococcus pneumoniae*, or a method of preventing infection with serotype 22F *Streptococcus pneumoniae*, or a method of reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by serotype 22F *Streptococcus pneumoniae*, the methods comprising administering to a subject an immunogenic amount of any of the immunogenic compositions described herein.

One embodiment of the disclosure provides a method of treating or preventing a *Streptococcus pneumoniae* infection, disease or condition associated with serotype 22F *Streptococcus pneumoniae* in a subject, the method comprising the step of administering a therapeutically or prophylactically effective amount of an immunogenic composition described herein to the subject. Another embodiment provides a method of treating or preventing a *Streptococcus pneumoniae* infection, disease or condition associated with a serotype 22F *Streptococcus pneumoniae* in a subject, the method comprising generating a polyclonal or monoclonal antibody preparation from the immunogenic composition described herein, and using said antibody preparation to confer passive immunity to the subject.

In one embodiment, the disclosure relates to the use of the immunogenic conjugate or immunogenic composition disclosed herein for the manufacture of a medicament for protecting a subject against an infection with *Streptococcus pneumoniae*, and/or preventing

infection with *Streptococcus pneumoniae*, and/or reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by *Streptococcus pneumoniae*, and/or protecting a subject against an infection with serotype 22F *Streptococcus pneumoniae* and/or preventing infection with serotype 22F *Streptococcus pneumoniae*, and/or reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by serotype 22F *Streptococcus pneumoniae*.

One embodiment of the disclosure provides a method of protecting a subject against an infection with serotype 33F *Streptococcus pneumoniae*, or a method of preventing infection with serotype 33F *Streptococcus pneumoniae*, or a method of reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by serotype 33F *Streptococcus pneumoniae*, the methods comprising administering to a subject an immunogenic amount of any of the immunogenic compositions described herein.

One embodiment of the disclosure provides a method of treating or preventing a *Streptococcus pneumoniae* infection, disease or condition associated with serotype 33F *Streptococcus pneumoniae* in a subject, the method comprising the step of administering a therapeutically or prophylactically effective amount of an immunogenic composition described herein to the subject. Another embodiment provides a method of treating or preventing a *Streptococcus pneumoniae* infection, disease or condition associated with a serotype 33F *Streptococcus pneumoniae* in a subject, the method comprising generating a polyclonal or monoclonal antibody preparation from the immunogenic composition described herein, and using said antibody preparation to confer passive immunity to the subject.

In one embodiment, the disclosure relates to the use of the immunogenic conjugate or immunogenic composition disclosed herein for the manufacture of a medicament for protecting a subject against an infection with *Streptococcus pneumoniae*, and/or preventing infection with *Streptococcus pneumoniae*, and/or reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by *Streptococcus pneumoniae*, and/or protecting a subject against an infection with serotype 33F *Streptococcus pneumoniae* and/or preventing infection with serotype 33F *Streptococcus pneumoniae*, and/or reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by serotype 33F *Streptococcus pneumoniae*.

An "immune response" to an immunogenic composition is the development in a subject of a humoral and/or a cell-mediated immune response to molecules present in the immunogenic composition or vaccine composition of interest. For purposes of the present disclosure, a "humoral immune response" is an antibody-mediated immune response and involves the induction and generation of antibodies that recognize and bind with some affinity

for the antigen in the immunogenic composition or vaccine of the disclosure, while a "cell-mediated immune response" is one mediated by T-cells and/or other white blood cells. A "cell-mediated immune response" is elicited by the presentation of antigenic epitopes in association with Class I or Class II molecules of the major histocompatibility complex (MHC), CD1 or other non-classical MHC-like molecules. This activates antigen-specific CD4+ T helper cells or CD8+ cytotoxic T lymphocyte cells ("CTLs"). CTLs have specificity for peptide antigens that are presented in association with proteins encoded by classical or non-classical MHCs and expressed on the surfaces of cells. CTLs help induce and promote the intracellular destruction of intracellular microbes, or the lysis of cells infected with such microbes. Another aspect of cellular immunity involves an antigen-specific response by helper T-cells. Helper T-cells act to help stimulate the function, and focus the activity of, nonspecific effector cells against cells displaying peptide or other antigens in association with classical or non-classical MHC molecules on their surface. A "cell-mediated immune response" also refers to the production of cytokines, chemokines and other such molecules produced by activated T-cells and/or other white blood cells, including those derived from CD4+ and CD8+ T-cells. The ability of a particular antigen or composition to stimulate a cell-mediated immunological response may be determined by a number of assays, such as by lymphoproliferation (lymphocyte activation) assays, CTL cytotoxic cell assays, by assaying for T-lymphocytes specific for the antigen in a sensitized subject, or by measurement of cytokine production by T cells in response to re-stimulation with antigen. Such assays are well known in the art. See, e.g., Erickson *et al.* (1993) *J. Immunol.* 151:4189-4199; and Doe *et al.* (1994) *Eur. J. Immunol.* 24:2369-2376.

As used herein, "treatment" (including variations thereof, e.g., "treat" or "treated") means any one or more of the following: (i) the prevention of infection or re-infection, as in a traditional vaccine, (ii) the reduction in the severity of, or, in the elimination of symptoms, and (iii) the substantial or complete elimination of the pathogen or disorder in question. Hence, treatment may be effected prophylactically (prior to infection) or therapeutically (following infection). In the present disclosure, prophylactic treatment is the preferred mode. According to a particular embodiment of the present disclosure, compositions and methods are provided that treat, including prophylactically and/or therapeutically immunize, a host animal against a serotype 10A *Streptococcus pneumoniae* infection. According to a particular embodiment of the present disclosure, compositions and methods are provided that treat, including prophylactically and/or therapeutically immunize, a host animal against a serotype 22F *Streptococcus pneumoniae* infection. According to a particular embodiment of the present disclosure, compositions and methods are provided that treat, including prophylactically and/or therapeutically immunize, a host animal against a serotype 33F

*Streptococcus pneumoniae* infection. The methods of the present disclosure are useful for conferring prophylactic and/or therapeutic immunity to a subject. The methods of the present disclosure can also be practiced on subjects for biomedical research applications.

5 An "immunogenic amount," and "immunologically effective amount," both of which are used interchangeably herein, refers to the amount of antigen or immunogenic composition sufficient to elicit an immune response, either a cellular (T-cell) or humoral (B-cell or antibody) response, or both, as measured by standard assays known to one skilled in the art.

10

In a preferred embodiment, said subject is a human. In a most preferred embodiment, said subject is a newborn (i.e. under three months of age), an infant (from 3 months to one year of age) or a toddler (i.e. from one year to four years of age).

In an embodiment, the immunogenic compositions disclosed herein are for use as a vaccine.

15

In such embodiment, the subject to be vaccinated may be less than 1 year of age. For example, the subject to be vaccinated can be about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months of age. In an embodiment, the subject to be vaccinated is about 2, 4 or 6 months of age. In another embodiment, the subject to be vaccinated is less than 2 years of age. For example the subject to be vaccinated can be about 12-15 months of age. In some cases, as little as one dose of the immunogenic composition according to the invention is needed, but under some circumstances, a second, third or fourth dose may be given (see regimen section).

20

In an embodiment of the present invention, the subject to be vaccinated is a human adult 50 years of age or older, more preferably a human adult 55 years of age or older. In an embodiment, the subject to be vaccinated is a human adult 65 years of age or older, 70 years of age or older, 75 years of age or older or 80 years of age or older.

25

In an embodiment the subject to be vaccinated is an immunocompromised individual, in particular a human. An immunocompromised individual is generally defined as a person who exhibits an attenuated or reduced ability to mount a normal humoral or cellular defense to challenge by infectious agents.

30

In an embodiment of the present invention, the immunocompromised subject to be vaccinated suffers from a disease or condition that impairs the immune system and results in an antibody response that is insufficient to protect against or treat pneumococcal disease.

In an embodiment, said disease is a primary immunodeficiency disorder. Preferably, said primary immunodeficiency disorder is selected from the group consisting of: combined T- and B-cell immunodeficiencies, antibody deficiencies, well-defined syndromes, immune dysregulation diseases, phagocyte disorders, innate immunity deficiencies, autoinflammatory disorders, and complement deficiencies. In an embodiment, said primary immunodeficiency disorder is selected from the one disclosed on page 24 line 11 to page 25 line 19 of the PCT application WO2010/125480.

In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated suffers from a disease selected from the groups consisting of: HIV-infection, acquired immunodeficiency syndrome (AIDS), cancer, chronic heart or lung disorders, congestive heart failure, diabetes mellitus, chronic liver disease, alcoholism, cirrhosis, spinal fluid leaks, cardiomyopathy, chronic bronchitis, emphysema, Chronic obstructive pulmonary disease (COPD), spleen dysfunction (such as sickle cell disease), lack of spleen function (asplenia), blood malignancy, leukemia, multiple myeloma, Hodgkin's disease, lymphoma, kidney failure, nephrotic syndrome and asthma.

In an embodiment of the present invention, the immunocompromised subject to be vaccinated suffers from malnutrition.

In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated is taking a drug or treatment that lowers the body's resistance to infection. In an embodiment, said drug is selected from the one disclosed on page 26 line 33 to page 26 line 40 of the PCT application WO2010/125480.

In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated is a smoker.

In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated has a white blood cell count (leukocyte count) below  $5 \times 10^9$  cells per liter, or below  $4 \times 10^9$  cells per liter, or below  $3 \times 10^9$  cells per liter, or below  $2 \times 10^9$  cells per liter, or below  $1 \times 10^9$  cells per liter, or below  $0.5 \times 10^9$  cells per liter, or below  $0.3 \times 10^9$  cells per liter, or below  $0.1 \times 10^9$  cells per liter.

White blood cell count (leukocyte count): The number of white blood cells (WBCs) in the blood. The WBC is usually measured as part of the CBC (complete blood count). White blood cells are the infection-fighting cells in the blood and are distinct from the red (oxygen-carrying) blood cells known as erythrocytes. There are different types of white blood cells, including neutrophils (polymorphonuclear leukocytes; PMNs), band cells (slightly immature neutrophils), T-type lymphocytes (T cells), B-type lymphocytes (B cells), monocytes,

eosinophils, and basophils. All the types of white blood cells are reflected in the white blood cell count. The normal range for the white blood cell count is usually between 4,300 and 10,800 cells per cubic millimeter of blood. This can also be referred to as the leukocyte count and can be expressed in international units as  $4.3 - 10.8 \times 10^9$  cells per liter.

- 5 In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated suffers from neutropenia. In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated has a neutrophil count below  $2 \times 10^9$  cells per liter, or below  $1 \times 10^9$  cells per liter, or below  $0.5 \times 10^9$  cells per liter, or below  $0.1 \times 10^9$  cells per liter, or below  $0.05 \times 10^9$  cells per liter.
- 10 A low white blood cell count or "neutropenia" is a condition characterized by abnormally low levels of neutrophils in the circulating blood. Neutrophils are a specific kind of white blood cell that help prevent and fight infections. The most common reason that cancer patients experience neutropenia is as a side effect of chemotherapy. Chemotherapy-induced neutropenia increases a patient's risk of infection and disrupts cancer treatment.
- 15 In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated has a CD4+ cell count below  $500/\text{mm}^3$ , or CD4+ cell count below  $300/\text{mm}^3$ , or CD4+ cell count below  $200/\text{mm}^3$ , CD4+ cell count below  $100/\text{mm}^3$ , CD4+ cell count below  $75/\text{mm}^3$ , or CD4+ cell count below  $50/\text{mm}^3$ .

CD4 cell tests are normally reported as the number of cells in  $\text{mm}^3$ . Normal CD4 counts are  
20 between 500 and 1600, and CD8 counts are between 375 and 1100. CD4 counts drop dramatically in people with HIV.

In an embodiment of the invention, any of the immunocompromised subject disclosed herein is a human male or a human female.

- 25 The amount of a conjugate in a composition is generally calculated based on total polysaccharide, conjugated and non-conjugated for that conjugate. For example, a conjugate with 20% free polysaccharide will have about 80 mcg of conjugated polysaccharide and about 20 mcg of non-conjugated polysaccharide in a 100 mcg polysaccharide dose. The protein contribution to the conjugate is usually not considered  
30 when calculating the dose of a conjugate. Generally, each dose will comprise 0.1 to 100 mcg of polysaccharide, particularly 0.1 to 10 mcg, and more particularly 1 to 10 mcg and more particularly 1 to 5  $\mu\text{g}$ . Preferably each dose will comprise about 1.1, 2, 2.2, 3, 3.3, 4, 4.4  $\mu\text{g}$  of polysaccharide.

Optimal amounts of components for a particular immunogenic composition or vaccine can be ascertained by standard studies involving observation of appropriate immune responses in subjects. Following an initial vaccination, subjects can receive one or several booster immunizations adequately spaced.

5

The effectiveness of an antigen as an immunogen, can be measured either by proliferation assays, by cytolytic assays, such as chromium release assays to measure the ability of a T-cell to lyse its specific target cell, or by measuring the levels of B-cell activity by measuring the levels of circulating antibodies specific for the antigen in serum. An immune response may also be detected by measuring the serum levels of antigen specific antibody induced following administration of the antigen, and more specifically, by measuring the ability of the antibodies so induced to enhance the opsonophagocytic ability of particular white blood cells, as described herein. The level of protection of the immune response may be measured by challenging the immunized host with the antigen that has been administered. For example, if the antigen to which an immune response is desired is a bacterium, the level of protection induced by the immunogenic amount of the antigen is measured by detecting the percent survival or the percent mortality after challenge of the animals with the bacterial cells. In one embodiment, the amount of protection may be measured by measuring at least one symptom associated with the bacterial infection, e.g., a fever associated with the infection. The amount of each of the antigens in the multi-antigen or multi-component vaccine or immunogenic compositions will vary with respect to each of the other components and can be determined by methods known to the skilled artisan. Such methods would include procedures for measuring immunogenicity and/ or *in vivo* efficacy.

15

The disclosure further provides antibodies and antibody compositions which bind specifically and selectively to the capsular polysaccharides or glycoconjugates of the present disclosure. In some embodiments, antibodies are generated upon administration to a subject of the capsular polysaccharides or glycoconjugates of the present disclosure. In some embodiments, the disclosure provides purified or isolated antibodies directed against one or more of the capsular polysaccharides or glycoconjugates of the present disclosure. In some embodiments, the antibodies of the present disclosure are functional as measured by killing bacteria in either an animal efficacy model or via an opsonophagocytic killing assay. In some embodiments, the antibodies of the disclosure confer passive immunity to a subject. The present disclosure further provides polynucleotide molecules encoding an antibody or antibody fragment of the disclosure, and a cell, cell line (such as hybridoma cells or other engineered cell lines for recombinant production of antibodies) or a transgenic animal that

25

30

35

produces an antibody or antibody composition of the disclosure, using techniques well-known to those of skill in the art.

## Examples

5

### Example 1: preparation of serotype 22F polysaccharide – CRM<sub>197</sub> conjugate

#### **1.1. Preparation of isolated *Streptococcus pneumoniae* serotype 22F polysaccharide**

Serotype 22F capsular polysaccharides can be obtained directly from bacteria using isolation  
10 procedures known to one of ordinary skill in the art. (see for example methods disclosed  
U.S. Patent App. Pub. Nos. 20060228380, 20060228381, 20070184071, 20070184072,  
20070231340, and 20080102498 or WO2008118752). The serotype 22F *Streptococcus*  
*pneumoniae* were grown in a seed bottle and then transferred to a seed fermentor. Once the  
targeted optical density was reached, the cells were transferred to a production fermentor.  
15 The fermentation was broth was inactivated by the addition of N-lauroyl sarcosine and  
purified by ultrafiltration and diafiltration.

The purified *Streptococcus pneumoniae* serotype 22F polysaccharide was sized by high  
pressure homogenization using a PANDA 2K homogenizer ® (GEA Niro Soavi) to produce  
20 the isolated *Streptococcus pneumoniae* serotype 22F polysaccharide.

#### **1.2. Oxidation of Isolated *Streptococcus pneumoniae* serotype 22F capsular polysaccharide**

Oxidation of Polysaccharide was carried out in 100 mM potassium phosphate buffer (pH 5.8  
25 ± 0.2) obtained by sequential addition of calculated amount of 500 mM potassium phosphate  
buffer (pH 5.8) and WFI to give final polysaccharide concentration of 2.0 g/L. If required, the  
reaction pH was adjusted to 5.8, approximately. After pH adjustment, the reaction  
temperature was lowered to 5 ± 3 °C. Oxidation was initiated by the addition of 0.10 ± 0.02  
molar equivalents (MEq) of sodium periodate. The target oxidation reaction time is 16 ± 1 hrs  
30 at 5 ± 3 °C.

The oxidation reaction was quenched with 2 MEq of 2,3-butanediol under continuous stirring  
at 5 ± 3 °C for 1-2 hrs.

35 Concentration and diafiltration of the activated polysaccharide was carried out using 100K  
MWCO ultrafiltration cassettes. Diafiltration was performed against 35-fold diavolume of WFI.  
The purified activated polysaccharide was stored at 5 ± 3°C. The purified activated



saccharide is characterized inter alia by (i) Molecular Weight by SEC-MALLS (ii) presence of O-acetyl and (iii) Degree of Oxidation.

5 SEC-MALLS is used for the determination of the molecular weight of polysaccharides and polysaccharide-protein conjugates. SEC is used to separate the polysaccharides by hydrodynamic volume. Refractive index (RI) and multi-angle laser light scattering (MALLS) detectors are used for the determination of the molecular weight. When light interacts with matter, it scatters and the amount of scattered light is related to the concentration, the square of the  $dn/dc$  (the specific refractive index increments), and the molar mass of the matter. The  
10 molecular weight measurement is calculated based on the readings from the scattered light signal from the MALLS detector and the concentration signal from the RI detector.

The degree of oxidation ( $DO = \text{moles of sugar repeat unit} / \text{moles of aldehyde}$ ) of the activated polysaccharide was determined as follows:

15 The moles of sugar repeat unit is determined by various colorimetric methods, such as for example by using Anthrone method. The polysaccharide is first broken down to monosaccharides by the action of sulfuric acid and heat. The Anthrone reagent reacts with the hexoses to form a yellow-green colored complex whose absorbance is read spectrophotometrically at 625nm. Within the range of the assay, the absorbance is directly  
20 proportional to the amount of hexose present.

The moles of aldehyde also is determined simultaneously, using MBTH colorimetric method. The MBTH assay involves the formation of an azine compound by reacting aldehyde groups (from a given sample) with a 3-methyl-2-benzothiazolone hydrazone (MBTH assay reagent). The excess 3-methyl-2-benzothiazolone hydrazone oxidizes to form a reactive cation. The  
25 reactive cation and the azine react to form a blue chromophore. The formed chromophore is then read spectroscopically at 650 nm.

### 1.3. Conjugation an activated *Streptococcus pneumoniae* serotype 22F polysaccharide with CRM<sub>197</sub>

30

The conjugation process consists of the following steps:

- a. Compounding with sucrose excipient, and lyophilization.
- b. Reconstitution of the lyophilized polysaccharide and CRM<sub>197</sub>.
- c. Conjugation of activated polysaccharide to CRM<sub>197</sub> and capping
- 35 d. Purification of the conjugate

a. Compounding with Sucrose and Lyophilization

The activated polysaccharide was compounded with sucrose (50% w/v in WFI) to a ratio of 25 grams of sucrose per gram of activated polysaccharide. The bottle of compounded mixture was then lyophilized. Following lyophilization, bottles containing lyophilized activated polysaccharide were stored at  $-20 \pm 5^{\circ}\text{C}$ . Calculated amount of CRM<sub>197</sub> protein (target S/P input ratio = 1) was shellfrozen and lyophilized separately. Lyophilized CRM<sub>197</sub> was stored at  $-20 \pm 5^{\circ}\text{C}$ .

b. Reconstitution of lyophilized activated polysaccharide and CRM<sub>197</sub> Protein

Lyophilized activated polysaccharide was reconstituted in anhydrous dimethyl sulfoxide (DMSO). Upon complete dissolution of polysaccharide, an equal amount of anhydrous DMSO was added to lyophilized CRM<sub>197</sub> for reconstitution.

c. Conjugation of activated polysaccharide to CRM<sub>197</sub> and capping

Reconstituted CRM<sub>197</sub> (in DMSO) was combined in the conjugation reaction vessel with the reconstituted activated polysaccharide. The final polysaccharide concentration in reaction solution is 1 g/L. Conjugation was initiated by adding  $1.5 \pm 0.1$  MEq of sodium cyanoborohydride to the reaction mixture and the reaction was incubated at  $23 \pm 2^{\circ}\text{C}$  for  $20 \pm 2$  hrs. Termination of conjugation reaction is done by adding 2 MEq of sodium borohydride. The capping reaction was incubated at  $23 \pm 2^{\circ}\text{C}$  for  $3 \pm 1$  hrs.

d. Purification of Conjugate

The conjugate solution was diluted 1:5 with chilled 5 mM succinate-0.9% saline (pH 6.0) in preparation for purification by tangential flow filtration using 100K MWCO membranes and a 20X diafiltration was performed using 5 mM succinate-0.9% saline (pH6.0) as the medium. After the diafiltration was completed, the conjugate retentate was further diluted, filtered through a 0.22 $\mu\text{m}$  filter and stored at 2-8  $^{\circ}\text{C}$ .

Table 1 comprises characterizing data for serotype 22F polysaccharide-CRM<sub>197</sub> conjugates obtained by the method of the invention. In particular, conjugates 5 and 6 of the table were obtained as disclosed in example 1.

Serotype 22F polysaccharide-CRM<sub>197</sub> conjugates generated by the RAC-DMSO process provided better yields and exhibited better batch to batch consistency in the MW along with significantly lower %Free saccharide levels and higher carrier protein (Lysine) modification compared to the conjugates generated by the RAC-aqueous process, as shown in Table 1.

Conjugate No	1	2	3	4	5	6
	Aqueous	Aqueous	Aqueous	DMSO	DMSO	DMSO
Act Poly MW, kDa	488	803	410	639	709	416
DO	17.3	15	10.3	14.6	14.4	13.7
Input Ratio	0.8	0.8	1	0.8	1	1
% Conjugate Yield	66	63	46	70	62	75
%Free Saccharide	18	43	45	< 5	< 5	8
Conjugate MW	1024	3544	462	3911	3734	4453
Saccharide-Protein Ratio	1.6	1.1	1.45	0.8	0.65	1.0
Modified Lys (AAA)	2.3	N/A	N/A	6.1	N/A	4.7

%Conjugate yield is calculated as follows: (amount of polysaccharide in the conjugate x100) / amount of activated polysaccharide.

### **Example 2: preparation of serotype 10A polysaccharide – CRM<sub>197</sub> conjugate**

5

#### **2.1. Preparation of isolated *Streptococcus pneumoniae* serotype 10A polysaccharide**

Isolated Serotype 10A polysaccharides was obtained as disclosed in example 1.1 except that the purified polysaccharide was not sized.

#### **10 2.2. Oxidation of Isolated *Streptococcus pneumoniae* serotype 10A capsular polysaccharide**

A calculated volume of 0.1M potassium phosphate buffer (pH 6.0) and water-for-injection (WFI) was added to the polysaccharide solution to achieve a final polysaccharide concentration of 2.5g/L and a final concentration of 25mM potassium phosphate buffer, If required pH was adjusted to 6.0, approximately. The diluted polysaccharide was then cooled to 5 ± 3°C. Oxidation was initiated by the addition of 0.25± 0.02 molar equivalents (MEq) of sodium periodate solution. The oxidation reaction time was approximately 4 hrs at 5 ± 3°C. The oxidation reaction was quenched with 1 MEq of 2,3-butanediol under continuous stirring at 5 ± 3 °C for 1-2 hrs.

20 After reaching the target reaction time, the activated polysaccharide was concentrated using 30K MWCO Millipore ultrafiltration cassettes. The diafiltration was then performed against 20-fold diavolume of WFI. The purified activated polysaccharide was stored at 5 ± 3°C. The

purified activated saccharide is characterized inter alia by (i) Molecular Weight by SEC-MALLS and (ii) Degree of Oxidation.

### 2.3 Conjugation an activated *Streptococcus pneumoniae* serotype 10A polysaccharide with CRM<sub>197</sub>

The conjugation process consists of the following steps:

- a. Compounding with sucrose excipient, and lyophilization.
- b. Reconstitution of the lyophilized polysaccharide and CRM<sub>197</sub>.
- 10 c. Conjugation of activated polysaccharide to CRM<sub>197</sub> and capping
- d. Purification of the conjugate

#### a. Compounding with Sucrose

The activated polysaccharide is compounded with sucrose to a ratio of  $25 \pm 2.5$  grams of sucrose per gram of activated polysaccharide. The bottle of compounded mixture was then lyophilized. Following lyophilization, bottles containing lyophilized activated polysaccharide were stored at  $-20 \pm 5^\circ\text{C}$ .

#### b. Reconstitution of Lyophilized Activated Polysaccharide and CRM<sub>197</sub> Protein

Lyophilized activated polysaccharide was reconstituted in anhydrous dimethyl sulfoxide (DMSO). Upon complete dissolution of polysaccharide, the same amount of anhydrous DMSO was added to the calculated CRM<sub>197</sub> for reconstitution.

#### c. Conjugation of Activated Polysaccharide to CRM<sub>197</sub> and Capping

Reconstituted CRM<sub>197</sub> (in DMSO) was added to the reconstituted activated polysaccharide in the conjugation reactor. The final polysaccharide concentration is 1 g/L. Conjugation was performed by adding  $1.2 \pm 0.1$  MEq of sodium cyanoborohydride to the reaction mixture. The reaction was incubated and at  $23 \pm 2^\circ\text{C}$  for  $24 \pm 2$  hrs. Termination of conjugation reaction is done by adding 2 MEq of sodium borohydride. The capping reaction was incubated at  $23 \pm 2^\circ\text{C}$  for  $3 \pm 1$  hrs.

Termination of conjugation reaction is done by adding 2 MEq of sodium borohydride. This capping reaction proceeded for  $3 \pm 1$  hrs at  $23 \pm 2^\circ\text{C}$ .

#### d. Purification of Conjugate

The conjugate solution was then diluted into 5x (by volume) chilled 5 mM succinate-0.9% saline (pH 6.0 and a 20X diafiltration was performed using 5 mM succinate-0.9% saline (pH6.0). After the initial diafiltration was completed, the conjugate retentate was transferred

through a 0.22µm filter. The conjugate was diluted further with 5 mM succinate / 0.9% saline (pH 6), and after the final 0.22µm filtration step it was stored at 2-8 °C.

5 Table 2 comprises characterizing data for serotype 10A polysaccharide-CRM<sub>197</sub> conjugates obtained by the method of the invention. In particular, conjugates 4 to 6 of table 2 were obtained as disclosed in example 2.

10 Serotype 10A polysaccharide-CRM<sub>197</sub> conjugates generated by the RAC-DMSO process provided better yields and exhibited better batch to batch consistency in the MW along with significantly lower %Free saccharide levels and higher carrier protein (Lysine) modification compared to the conjugates generated by the RAC-aqueous process, as shown in Table 2.

Conjugate No	1	2	3	4	5	6
	Aqueous	Aqueous	Aqueous	DMSO	DMSO	DMSO
DO	12.2	12.2	12.8	10.3	10.8	10.5
Activated Saccharide MW, kda	157	157	163	170	170	170
Input Ratio	1.0	1.0	1.0	1.1	1.1	1.1
%Yield	71	34	50	82	73	66
%Free Saccharide	20	31	26	6.8	6.4	9.7
Conjugate MW, kDa	1550	968	3050	4034	3463	5540
Saccharide-protein ratio	1.6	2.6	1.6	1.1	1.2	1.0
Lys modification AAA	2.6	N/A	N/A	6.9	6.7	6.1

15 **Example 3: preparation of serotype 33F polysaccharide – CRM<sub>197</sub> conjugate**

**3.1. Preparation of isolated *Streptococcus pneumoniae* serotype 33F polysaccharide**

Isolated Serotype 33F polysaccharide was obtained as disclosed in example 1.1.

20 **3.2. Oxidation of Isolated *Streptococcus pneumoniae* serotype 33F capsular polysaccharide**

A calculated volume of 0.1M sodium phosphate buffer (pH 6.0) and water-for-injection (WFI) was added to the polysaccharide solution to achieve a final polysaccharide concentration of 2g/L. If required pH was adjusted to 6.0, approximately. The diluted polysaccharide was then cooled to  $5 \pm 3^\circ\text{C}$ . Oxidation was initiated by the addition of 0.1 molar equivalent (MEq) of sodium periodate solution. The oxidation reaction time was approximately 20 hrs at  $5 \pm 3^\circ\text{C}$ . The oxidation reaction was quenched with 1 MEq of 2,3-butanediol under continuous stirring at  $5 \pm 3^\circ\text{C}$  for about 1 hour. After reaching the target reaction time, the activated polysaccharide was concentrated using 100K MWCO Millipore ultrafiltration cassettes. The diafiltration was then performed against 40-fold diavolume of WFI. The purified activated polysaccharide was stored at  $5 \pm 3^\circ\text{C}$ . The purified activated saccharide is characterized inter alia by (i) Molecular Weight by SEC-MALLS and (ii) Degree of Oxidation.

### 3.3 Conjugation an activated *Streptococcus pneumoniae* serotype 33F polysaccharide with CRM<sub>197</sub>

The serotype 33F polysaccharide-CRM197 conjugate was obtained by a process similar to the process of example 1.3, using the activated 33F polysaccharide obtained in example 3.2.

Serotype 33F polysaccharide-CRM<sub>197</sub> conjugates generated by the RAC-DMSO process provided better yields and exhibited better batch to batch consistency in terms of preservation of O-Acetyl levels along with overall lower %Free saccharide levels and higher carrier protein (Lysine) modification compared to the conjugates generated by the RAC-aqueous process, as shown in Table 3.

Conjugate No	1	2	3	4	5	6
	Aq	Aq	Aq	DMSO	DMSO	DMSO
DO	15.2	15.5	15.7	17	9	14
Activated Poly MW	1126	573	574	120	328	128
%Yield	65	51	48	78	86	66
%Free Saccharide	8.5	19	13	18	5	< 5
Conjugate MW, kDa	1341	2137	3674	3158	2165	2001
Saccharide-protein ratio	1.7	1.4	1.3	1.1	1.1	0.79
% O-Ac preserved	< LOQ	< LOQ	< LOQ	82	100%	100%
Conj Lys (AAA)	N/A	2.3	2.3	4.5	3.4	5.2

LOQ = Limit of Quantitation

**Example 4: effect of quenching on molecular weight of activated serotype 22F and 33F polysaccharides**

5 Activation of 22F and 33F polysaccharides was carried out at either 4 °C or 22 °C. Both temperatures gave similar DO values. However, due to the fact that elevated temperature resulted in faster breakdown of activated polysaccharide thereby lowering activated poly MW (as shown when comparing for example figure 4 (22 °C) and figure 5 (4 °C)), 4 °C was generally preferred for activation reaction. In addition, activation data showed that quenching  
10 of un-reacted NaIO<sub>4</sub> after oxidation is an essential step for activation, especially for activation using higher amount of NaIO<sub>4</sub>. The graph shown in Figure 6 indicate that the MW of the activated 33F polysaccharide is relatively stable when increasing concentrations of periodate are used for activation (with quenching) while such MW is highly variable when no quenching step is used. At specific conditions set up to obtain a targeted degree of oxidation, the  
15 molecular weight of the activated polysaccharide is less variable when the quenching step is used. Addition of quenching reagent after oxidation helps to keep the structural integrity of activated polysaccharide until the completion of purification for example by ultrafiltration and diafiltration.

20 **Example 5: Opsonophagocytic activity assay (OPA)**

The immunogenicity of the conjugates obtained by the processes disclosed herein can be assessed using the opsonophagocytic assay (OPA) described below.

25 Groups of thirty 6-7 week old female Swiss Webster mice were immunized with 0.001 µg, 0.01 µg, or 0.1 µg of test conjugates via the subcutaneous route on week 0. The mice were boosted with the same dose of conjugate on week 3 and then bled at week 4. Serotype-specific OPAs were performed on week 4 sera samples.

30 Validated opsonophagocytic activity (OPA) assays are used to measure functional antibodies in murine sera specific for *S. pneumonia* serotype 10A, 22F or 33F. Test serum is set up in assay reactions that measure the ability of capsular polysaccharide specific immunoglobulin to opsonize bacteria, trigger complement deposition, thereby facilitating phagocytosis and killing of bacteria by phagocytes. The OPA titer is defined as the reciprocal dilution that  
35 results in a 50% reduction in bacterial count over control wells without test serum. The OPA titer is interpolated from the two dilutions that encompass this 50% killing cut-off.

OPA procedures were based on methods described in Hu et al., Clin Diagn Lab Immunol 2005;12(February (2)):287–95 with the following modifications. Test serum was serially diluted 2.5-fold and added to microtiter assay plates. Live serotype 10A, 22F and 33F target bacterial strains were added to the wells and the plates were shaken at either 25°C (serotype 22F) or 37°C (serotype 10A and 33F) for 30 minutes. Differentiated HL-60 cells (phagocytes) and baby rabbit serum (3- to 4-week old, Pel-Freez®, 12.5% final concentration) were added to the wells, and the plates were shaken at 37°C for 45 minutes (serotype 22F and 33F) or 60 minutes (serotype 10A). To terminate the reaction, 80 µL of 0.9% NaCl was added to all wells, mixed, and a 10µL aliquot were transferred to the wells of MultiScreen HTS HV filter plates (Millipore®) containing 200 µL of water. Liquid was filtered through the plates under vacuum, and 150 µL of HySoy medium was added to each well and filtered through. The filter plates were then incubated at 37°C, 5% CO<sub>2</sub> overnight and were then fixed with Destain Solution (Bio-Rad). The plates were then stained with Coomassie Blue and destained once. Colonies were imaged and enumerated on a Cellular Technology Limited (CTL) ImmunoSpot Analyzer®. Raw colony counts were used to plot kill curves and calculate OPA titers.


Serotype 10A polysaccharide-CRM<sub>197</sub> conjugates, Serotype 22F polysaccharide-CRM<sub>197</sub> conjugates and Serotype 33F polysaccharide-CRM<sub>197</sub> conjugates obtained as disclosed in examples 1 to 3 were tested in the OPA assay disclosed above and were found to be immunogenic.



**Claims**

1. A process for preparing an activated *Streptococcus pneumoniae* serotype 10A, 22F or 33F capsular polysaccharide, the process comprising the steps of:
- 5 (a) reacting an isolated serotype 10A, 22F or 33F capsular polysaccharide with an oxidizing agent; and
- (b) quenching the oxidation reaction by addition of a quenching agent resulting in an activated *Streptococcus pneumoniae* serotype 10A, 22F or 33F polysaccharide.
- 10
2. A process according to claim 1 wherein the concentration of serotype 10A, 22F or 33F polysaccharide in step (a) is between 0.5 and 5 mg/mL.
3. A process according to claim 1 or 2 wherein said polysaccharide is reacted with 0.01 to 10, 0.05 to 5, 0.1 to 1, 0.5 to 1, 0.7 to 0.8, 0.01 to 0.2, 0.05 to 0.5, 0.05 to 0.2, 0.1 to 0.3 molar equivalents of oxidizing agent in step (a).
- 15
4. A process according to any one of claims 1 to 3 wherein said polysaccharide is reacted with 0.05 to 0.3 molar equivalents of oxidizing agent in step (a).
- 20
5. A process according to any one of claims 1 to 4 where the oxidizing agent is periodate.
6. A process according to any one of claims 1 to 4 where the oxidizing agent is sodium metaperiodate.
- 25
7. A process according to any one of claims 1 to 6 wherein the temperature of the reaction in step (a) is maintained between 1 and 10°C.
8. A process according to any one of claims 1 to 7 wherein the temperature of the reaction in step (a) is about °C or about 5°C.
- 30
9. A process according to any one of claims 1 to 8 wherein the duration of step (a) is between 1 and 25 hours.
- 35
10. A process according to any one of claims 1 to 9 where the quenching agent is selected from vicinal diols, 1,2-aminoalcohols, amino acids, glutathione, sulfite, bisulfate, dithionite, metabisulfite, thiosulfate, phosphites, hypophosphites or phosphorous acid.

11. A process according to any one of claims 1 to 9 where the quenching agent is a compound of formula (II)



5 wherein R<sup>1</sup> and R<sup>2</sup> are each independently selected from H, methyl, ethyl, propyl or isopropyl.

12. A process according to any one of claims 1 to 9 where the quenching agent is selected from glycerol, ethylene glycol, propan-1,2-diol, butan-1,2-diol or butan-2,3-diol.

10 13. A process according to any one of claims 1 to 9 where the quenching agent is butan-2,3-diol.

14. A process according to any one of claims 1 to 13 wherein the oxidation reaction is quenched by the addition of 0.1 to 10, 0.5 to 5, 0.5 to 3 or 0.5 to 2 molar equivalents of  
15 quenching agent.

15. A process according to any one of claims 1 to 13 wherein the oxidation reaction is quenched by the addition of 0.5 to 2 molar equivalents of quenching agent.

20 16. A process according to any one of claims 1 to 15 wherein the temperature of the reaction in step (b) is maintained between 1 and 10°C.

17. A process according to any one of claims 1 to 16 wherein the temperature of the reaction in step (b) is about 4°C.

25 18. A process according to any one of claims 1 to 17 wherein the duration of step (b) is between 0.5 and 2 hours.

30 19. A process according to any one of claims 1 to 18 wherein the activated polysaccharide is purified after step (b).

20. A process according to any one of claims 1 to 19 wherein said isolated polysaccharide is an isolated *Streptococcus pneumoniae* serotype 10A polysaccharide.

21. A process according to claim 20 wherein said isolated *Streptococcus pneumoniae* serotype 10A polysaccharide is obtained by a process comprising the steps of:

(a) preparing a fermentation culture of serotype 10A *Streptococcus pneumoniae* bacterial cells;

5 (b) lysing the bacterial cells in said fermentation culture; and

(c) purifying serotype 10A polysaccharide from the fermentation culture.

22. An activated *Streptococcus pneumoniae* serotype 10A capsular polysaccharide obtained by a process of any one of claims 1 to 21.

10

23. An activated *Streptococcus pneumoniae* serotype 10A capsular polysaccharide obtainable by a process of any one of claims 1 to 21.

24. An activated *Streptococcus pneumoniae* serotype 10A capsular polysaccharide according to any one of claims 22 or 23, where said polysaccharide has a molecular weight between 50 and 400, 50 and 350, 50 and 300, 50 and 250, 50 and 200, 100 and 300, 100 and 250, or 100 and 200kDa.

15

25. An activated *Streptococcus pneumoniae* serotype 10A capsular polysaccharide according to any one of claims 22 or 23, where said polysaccharide has a molecular weight between 50 and 300kDa.

20

26. An activated *Streptococcus pneumoniae* serotype 10A capsular polysaccharide according to any one of claims 22 or 23, where said polysaccharide has a molecular weight between 100 and 200kDa.

25

27. An activated *Streptococcus pneumoniae* serotype 10A capsular polysaccharide according to any one of claims 22 to 26, where said polysaccharide has a degree of oxidation between 5 and 20, 5 and 15, 8 and 14, 8 and 12, or 9 and 11.

30

28. An activated *Streptococcus pneumoniae* serotype 10A capsular polysaccharide according to any one of claims 22 to 26, where said polysaccharide has a degree of oxidation between 9 and 11.

35

29. A process according to any one of claims 1 to 19 wherein said isolated polysaccharide is an isolated *Streptococcus pneumoniae* serotype 22F polysaccharide.

30. A process according to claim 29 wherein said isolated *Streptococcus pneumoniae* serotype 22F polysaccharide is obtained by a process comprising the steps of:
- (a) preparing a fermentation culture of serotype 22F *Streptococcus pneumoniae* bacterial cells;
  - 5 (b) lysing the bacterial cells in said fermentation culture;
  - (c) purifying serotype 22F polysaccharide from the fermentation culture; and
  - (d) sizing the serotype 22F polysaccharide by mechanical sizing, preferably by high pressure homogenization.
- 10 31. An activated *Streptococcus pneumoniae* serotype 22F capsular polysaccharide obtained by a process of any one of claims 1 to 19 or 29 or 30.
32. An activated *Streptococcus pneumoniae* serotype 22F capsular polysaccharide obtainable by a process of any one of claims 1 to 19 or 29 or 30.
- 15 33. An activated *Streptococcus pneumoniae* serotype 22F capsular polysaccharide according to any one of claims 31 or 32, where said polysaccharide has a molecular weight between 25 and 1000, 100 and 1000, 300 and 800, 300 and 700, 300 and 600, 400 and 1000, 400 and 800, 400 and 700, or 400 and 600kDa.
- 20 34. An activated *Streptococcus pneumoniae* serotype 22F capsular polysaccharide according to any one of claims 31 or 32, where said polysaccharide has a molecular weight between 300 and 800kDa.
- 25 35. An activated *Streptococcus pneumoniae* serotype 22F capsular polysaccharide according to any one of claims 31 or 32, where said polysaccharide has a molecular weight between 400 and 600kDa.
- 30 36. An activated *Streptococcus pneumoniae* serotype 22F capsular polysaccharide according to any one of claims 31 to 35, where said polysaccharide has a degree of oxidation between 10 and 25, 10 and 20, 12 and 20, or 14 and 18.
- 35 37. An activated *Streptococcus pneumoniae* serotype 22F capsular polysaccharide according to any one of claims 31 to 35, where said polysaccharide has a degree of oxidation between 10 and 20.

38. An activated *Streptococcus pneumoniae* serotype 22F capsular polysaccharide according to any one of claims 31 to 37, where said polysaccharide comprises at least 0.6 mM acetate per mM serotype 22F polysaccharide.
- 5 39. A process according to any one of claims 1 to 19 wherein said isolated polysaccharide is an isolated *Streptococcus pneumoniae* serotype 33F polysaccharide.
40. A process according to claim 39 wherein said isolated *Streptococcus pneumoniae* serotype 33F polysaccharide is obtained by a process comprising the steps of:
- 10 (a) preparing a fermentation culture of serotype 33F *Streptococcus pneumoniae* bacterial cells;
- (b) lysing the bacterial cells in said fermentation culture;
- (c) purifying serotype 33F polysaccharide from the fermentation culture; and
- (d) sizing the serotype 33F polysaccharide by mechanical sizing, preferably by high pressure
- 15 homogenization.
41. An activated *Streptococcus pneumoniae* serotype 33F capsular polysaccharide obtained by a process of any one of claims 1 to 19 or 39 or 40.
- 20 42. An activated *Streptococcus pneumoniae* serotype 33F capsular polysaccharide obtainable by a process of any one of claims 1 to 19 or 39 or 40.
43. An activated *Streptococcus pneumoniae* serotype 33F capsular polysaccharide according to claim 39 or 40, where said polysaccharide has a molecular weight between 50
- 25 and 1250, 200 and 1200, 500 and 1200, 500 and 1000, 700 and 1200, 800 and 1200, 800 and 1100, 900 and 1200, or 800 and 1000Da.
44. An activated *Streptococcus pneumoniae* serotype 33F capsular polysaccharide according to claim 39 or 40, where said polysaccharide has a molecular weight between 500
- 30 and 1000kDa.
45. An activated *Streptococcus pneumoniae* serotype 33F capsular polysaccharide according to claim 39 or 40, where said polysaccharide has a molecular weight between 50 and 400kDa.

35

46. An activated *Streptococcus pneumoniae* serotype 33F capsular polysaccharide according to any one of claims 41 to 45, where said polysaccharide has a degree of oxidation between 10 and 20.
- 5 47. An activated *Streptococcus pneumoniae* serotype 33F capsular polysaccharide according to any one of claims 41 to 34, where said polysaccharide comprises at least 0.7 mM acetate per mM serotype 33F polysaccharide.
- 10 48. A process according to any one of claims 1 to 47 wherein the activated polysaccharide is lyophilized after step (b).
49. A process for the preparation of an immunogenic conjugate comprising *Streptococcus pneumoniae* serotype 10A, 22F or 33F polysaccharide covalently linked to a carrier protein, the process comprising the steps of:
- 15 (a) compounding an activated serotype 10A, 22F or 33F polysaccharide obtained or obtainable by a process of any one of claims 1 to 19 with a carrier protein; and,  
(b) reacting the compounded, activated serotype 10A, 22F or 33F polysaccharide and carrier protein with a reducing agent to form a serotype 10A, 22F or 33F polysaccharide:carrier protein conjugate.
- 20 50. A process according to claim 49, said process further comprising the step of:  
(c) capping unreacted aldehydes in the serotype 10A, 22F or 33F polysaccharide:carrier protein conjugate.
- 25 51. A process according to claim 49 or 50 where step (a) and (b) are carried out in DMSO.
52. A process according to claim 49 or 50 where step (a) and (b) are carried out in an aqueous solution.
- 30 53. A process according to anyone of claims 49 to 52 where the activated polysaccharide and the protein carrier are co-lyophilized after step (a).
54. A process according to claim 53 where step (b) is carried out in DMSO.
- 35 55. A process according to claim 53 where step (b) is carried out in an aqueous solution.

56. A process according to anyone of claims 49 to 55 where the concentration of activated polysaccharide in step (b) is between 0.1 and 10 mg/mL, 0.5 and 5 mg/mL, or 0.5 and 2 mg/mL.
- 5 57. A process according to anyone of claims 49 to 55 where the concentration of activated polysaccharide in step (b) is between 0.5 and 2 mg/mL.
58. A process according to anyone of claims 49 to 57 where the initial ratio of activated serotype 10A, 22F or 33F polysaccharide to carrier protein is between 5:1 and 0.1:1, 2:1 and  
10 0.1:1, 2:1 and 1:1, 1.5:1 and 1:1, 0.1:1 and 1:1, 0.3:1 and 1:1, or 0.6:1 and 1.2:1.
59. A process according to anyone of claims 49 to 57 where the initial ratio of activated serotype 10A, 22F or 33F polysaccharide to carrier protein is between 0.6:1 and 1.2:1.
- 15 60. A process according to anyone of claims 49 to 59 where, in step (b), the activated polysaccharide is reacted with between 0.5 to 2.5 molar equivalents of reducing agent.
61. A process according to anyone of claims 49 to 60 where the reducing agent is sodium cyanoborohydride.  
20
62. A process according to anyone of claims 49 to 61 wherein the duration of step (d) is between 20 and 30 hours.
63. A process according to any one of claims 49 to 62 wherein the temperature of the  
25 reaction in step d is maintained between 20 and 26°C.
64. A process according to claim 53 where the unreacted aldehydes are capped by the addition of about 2 molar equivalents of NaBH<sub>4</sub>.
- 30 65. A process according to any one of claims 49 to 64 further comprising a step of purifying the polysaccharide:carrier protein conjugate.
66. A process according to any one of claims 49 to 65 further comprising a step of purifying the polysaccharide:carrier protein conjugate by tangential flow filtration and diafiltration.  
35

67. An immunogenic conjugate comprising *Streptococcus pneumoniae* serotype 10A polysaccharide covalently linked to a carrier protein obtained by the process of any one of claims 49 to 66.
- 5 68. An immunogenic conjugate comprising *Streptococcus pneumoniae* serotype 10A polysaccharide covalently linked to a carrier protein obtainable by the process of any one of claims 49 to 66.
- 10 69. An immunogenic conjugate according to claim 67 or 68, where said conjugate has a molecular weight between 500 and 15000; 500 and 10000; 2000 and 10000; or 3000 and 8000kDa.
- 15 70. An immunogenic conjugate according to claim 67 or 68, where said conjugate has a molecular weight between 3000 and 8000kDa.
71. An immunogenic conjugate according to any one of claims 67 to 70, where said conjugate comprises less than about 50, 45, 40, 35, 30, 25, 20 or 15% of free serotype 10A polysaccharide compared to the total amount of serotype 10A polysaccharide.
- 20 72. An immunogenic conjugate according to any one of claims 67 to 70, where said conjugate comprises less than about 20% of free serotype 10A polysaccharide compared to the total amount of serotype 10A polysaccharide.
- 25 73. An immunogenic conjugate according to any one of claims 67 to 72, where the ratio of serotype 10A polysaccharide to carrier protein in the conjugate is between 0.5 and 3.
74. An immunogenic conjugate according to any one of claims 67 to 72, where the ratio of serotype 10A polysaccharide to carrier protein in the conjugate is between 0.8 and 1.4.
- 30 75. An immunogenic conjugate according to any one of claims 67 to 74, where at least 40% of said immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column.
- 35 76. An immunogenic conjugate according to any one of claims 67 to 74, where between 50% and 80% of the serotype 10A immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column.



77. An immunogenic conjugate according to any one of claims 67 to 76, where the degree of conjugation of said immunogenic conjugate is between 2 and 15, 2 and 13, 2 and 10, 2 and 8, 2 and 6, 2 and 5, 2 and 4, 3 and 15, 3 and 13, 3 and 10, 3 and 8, 3 and 6, 3 and 5, 3 and 4, 5 and 15, 5 and 10, 8 and 15, 8 and 12, 10 and 15, or 10 and 12.
- 5
78. An immunogenic conjugate according to any one of claims 67 to 76, where the degree of conjugation of said immunogenic conjugate is between 6 and 8.
79. An immunogenic conjugate comprising *Streptococcus pneumoniae* serotype 22F polysaccharide covalently linked to a carrier protein obtained by the process of any one of claims 49 to 66.
- 10
80. An immunogenic conjugate comprising *Streptococcus pneumoniae* serotype 22F polysaccharide covalently linked to a carrier protein obtainable by the process of any one of claims 49 to 66.
- 15
81. An immunogenic conjugate according to claim 79 or 80, where said conjugate comprises at least 0.5, 0.6 or 0.7 mM acetate per mM serotype 22F polysaccharide.
- 20
82. An immunogenic conjugate according to claim 79 or 80, where the ratio of mM acetate per mM serotype 22F polysaccharide in the immunogenic conjugate to mM acetate per mM serotype 22F polysaccharide in the isolated polysaccharide is at least 0.7.
- 25
83. An immunogenic conjugate according to any one of claims 79 to 82, where the ratio of mM acetate per mM serotype 22F polysaccharide in the immunogenic conjugate to mM acetate per mM serotype 22F polysaccharide in the activated polysaccharide is at least 0.7.
- 30
84. An immunogenic conjugate according to any one of claims 79 to 83, where said conjugate has a molecular weight between 400 and 15000; 500 and 10000; 2000 and 10000kDa; 3000 and 8000kDa; or 3000 and 5000kDa.
- 35
85. An immunogenic conjugate according to any one of claims 79 to 83, where said conjugate has a molecular weight between 3000 and 5000kDa.
86. An immunogenic conjugate according to any one of claims 79 to 85, where said conjugate comprises less than about 50, 45, 40, 35, 30, 25, 20 or 15% of free serotype 22F polysaccharide compared to the total amount of serotype 22F polysaccharide.

87. An immunogenic conjugate according to any one of claims 79 to 83, where said conjugate comprises less than about 20% of free serotype 22F polysaccharide compared to the total amount of serotype 22F polysaccharide.
- 5
88. An immunogenic conjugate according to any one of claims 79 to 87, where the ratio of serotype 22F polysaccharide to carrier protein in the conjugate is between 0.5 and 2,
89. An immunogenic conjugate according to any one of claims 79 to 87, where the ratio of  
10 serotype 22F polysaccharide to carrier protein in the conjugate is between 0.8 and 1.2.
90. An immunogenic conjugate according to any one of claims 79 to 89, where at least 40% of said immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column.
- 15
91. An immunogenic conjugate according to any one of claims 79 to 89, where between 50% and 80% of the serotype 22F immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column.
92. An immunogenic conjugate according to any one of claims 79 to 91, where the degree of  
20 conjugation of said immunogenic conjugate is between 2 and 15, 2 and 13, 2 and 10, 2 and 8, 2 and 6, 2 and 5, 2 and 4, 3 and 15, 3 and 13, 3 and 10, 3 and 8, 3 and 6, 3 and 5, 3 and 4, 4 and 7, 5 and 15, 5 and 10, 8 and 15, 8 and 12, 10 and 15, or 10 and 12.
93. An immunogenic conjugate according to any one of claims 79 to 91, where the degree of  
25 conjugation of said immunogenic conjugate is between 4 and 7.
94. An immunogenic conjugate comprising *Streptococcus pneumoniae* serotype 33F polysaccharide covalently linked to a carrier protein obtained by the process of any one of  
30 claims 49 to 66.
95. An immunogenic conjugate comprising *Streptococcus pneumoniae* serotype 33F polysaccharide covalently linked to a carrier protein obtainable by the process of any one of  
claims 49 to 66.
- 35
96. An immunogenic conjugate according to claim 94 or 95, where said conjugate comprises at least 0.5, 0.6 or 0.7 mM acetate per mM serotype 33F polysaccharide.

97. An immunogenic conjugate according to claim 94 or 95, where the ratio of mM acetate per mM serotype 33F polysaccharide in the immunogenic conjugate to mM acetate per mM serotype 33F polysaccharide in the isolated polysaccharide is at least 0.7.
- 5 98. An immunogenic conjugate according to any one of claims 94 to 97, where the ratio of mM acetate per mM serotype 33F polysaccharide in the immunogenic conjugate to mM acetate per mM serotype 33F polysaccharide in the activated polysaccharide is at least 0.7.
99. An immunogenic conjugate according to any one of claims 94 to 98, where said  
10 conjugate has a molecular weight between 500 and 30000; 500 and 25000; 500 and 20000;  
500 and 15000; 500 and 10000; 1000 and 10000; 1000 and 8000; 1000 and 5000; 2000 and  
10000kDa; 2000 and 8000; or 2000 and 5000kDa.
100. An immunogenic conjugate according to any one of claims 94 to 98, where said  
15 conjugate has a molecular weight between 1000 and 5000kDa.
101. An immunogenic conjugate according to any one of claims 94 to 100, where said  
conjugate comprises less than about 50, 45, 40, 35, 30, 25, 20 or 15% of free serotype 33F  
polysaccharide compared to the total amount of serotype 33F polysaccharide.  
20
102. An immunogenic conjugate according to any one of claims 94 to 100, where said  
conjugate comprises less than about 20% of free serotype 22F polysaccharide compared to  
the total amount of serotype 22F polysaccharide.
- 25 103. An immunogenic conjugate according to any one of claims 94 to 102, where the ratio of  
serotype 33F polysaccharide to carrier protein in the conjugate is between 0.5 and 2.
104. An immunogenic conjugate according to any one of claims 94 to 102, where the ratio of  
serotype 22F polysaccharide to carrier protein in the conjugate is between 0.5 and 1.5.  
30
105. An immunogenic conjugate according to any one of claims 94 to 104, where at least  
40% of said immunogenic conjugate has a Kd below or equal to 0.3 in a CL-4B column.
106. An immunogenic conjugate according to any one of claims 94 to 104, where between  
35 40% and 80% of the serotype 33F immunogenic conjugate has a Kd below or equal to 0.3 in  
a CL-4B column.

107. An immunogenic conjugate according to any one of claims 94 to 106, where the degree of conjugation of said immunogenic conjugate is between 2 and 15, 2 and 13, 2 and 10, 2 and 8, 2 and 6, 2 and 5, 2 and 4, 3 and 15, 3 and 13, 3 and 10, 3 and 8, 3 and 6, 3 and 5, 3 and 4, 5 and 15, 5 and 10, 8 and 15, 8 and 12, 10 and 15, or 10 and 12.

5

108. An immunogenic conjugate according to any one of claims 94 to 106, where the degree of conjugation of said immunogenic conjugate is between 4 and 6.

109. An immunogenic conjugate according to any one of claims 67 to 108 where the carrier protein is CRM<sub>197</sub>.

10

110. A process according to any one of claims 49 to 66 where the carrier protein is CRM<sub>197</sub>.

111. A process according to any one of claims 49 to 66 where the activated polysaccharide is of serotype 10A.

15

112. A process according to any one of claims 49 to 66 where the activated polysaccharide is of serotype 22F.

113. A process according to any one of claims 49 to 66 where the activated polysaccharide is of serotype 33F.

20

114. A process according to any one of claims 49 to 66 and 110 to 113 wherein said process further comprises the step of formulating the conjugate in a multivalent vaccine.

25

115. An immunogenic composition comprising an immunogenic conjugate according to any one of claims 67 to 109 and a physiologically acceptable vehicle.

116. An immunogenic composition according to claim 115 further comprising at least one additional antigen.

30

117. An immunogenic composition according to claim 115 or 116 further comprising an adjuvant.

118. An immunogenic composition according to claim 117 where the adjuvant is aluminium phosphate.

35

119. A vaccine comprising an immunogenic composition according to any one of claims 115 to 118.

5 120. A method of treating or preventing a *Streptococcus pneumoniae* infection, disease or condition associated with serotype 10A, 22F or 33F *Streptococcus pneumoniae* in a subject, the method comprising the step of administering a therapeutically or prophylactically effective amount of an immunogenic composition according to any one of claims 115 to 118, or the vaccine of claim 119.

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6

