

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3444342 T3**

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **C 12 N 9/22 (2006.01)** **A 61 K 48/00 (2006.01)** **C 12 N 15/90 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2020-08-24**
- (80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: **2020-05-27**
- (86) Europæisk ansøgning nr.: **18198271.1**
- (86) Europæisk indleveringsdag: **2013-03-15**
- (87) Den europæiske ansøgnings publiceringsdag: **2019-02-20**
- (30) Prioritet: **2012-07-11 US 201261670463 P** **2012-09-21 US 201261704072 P**
- (62) Stamansøgningsnr: **16198961.1**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Sangamo Therapeutics, Inc., Point Richmond Tech. Center , 501 Canal Boulevard, Suite A100, Richmond, CA 94804, USA**
- (72) Opfinder: **REBAR, Edward, J., c/o Sangamo BioSciences Inc., Point Richmond Tech Center, 501 Canal Blvd. Suite A100, Richmond, CA California 94804, USA**
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**
- (54) Benævnelse: **FREMGANGSMÅDER OG SAMMENSÆTNINGER TIL BEHANDLINGEN AF LYSOSOMALE AFLEJRINGSSYGDOMME**
- (56) Fremdragne publikationer:
WO-A1-2010/117464
WO-A1-2011/011767
WO-A1-2011/097036
WO-A1-2011/100058
WO-A2-2012/015938
GRABOWSKI ET AL: "Phenotype, diagnosis, and treatment of Gaucher's disease", THE LANCET, LANCET LIMITED. LONDON, GB, vol. 372, no. 9645, 4 October 2008 (2008-10-04), pages 1263-1271, XP025480739, ISSN: 0140-6736, DOI: 10.1016/S0140-6736(08)61522-6 [retrieved on 2008-10-02]
MONICA DAGNINO ET AL: "Molecular Diagnosis of Analbuminemia: A New Case Caused by a Nonsense Mutation in the Albumin Gene", INTERNATIONAL JOURNAL OF MOLECULAR SCIENCES, vol. 12, no. 12, 25 December 2011 (2011-12-25), pages 7314-7322, XP055081308, ISSN: 1661-6596, DOI: 10.3390/ijms12117314
Anguela et al.: "In Vivo Genome Editing of Liver Albumin for Therapeutic Gene Expression: Rescue of Hemophilic Mice Via Integration of Factor 9", 54th ASH Annual Meeting and Exposition, 751, 10 December 2012 (2012-12-10), XP002713765, Atlanta, USA Retrieved from the Internet:
URL:https://ash.confex.com/ash/2012/webprogram/Paper51404.html [retrieved on 2013-09-26]

DESCRIPTION

TECHNICAL FIELD

[0001] The present disclosure is in the field of the treatment of Lysosomal storage diseases (LSDs) and gene therapy.

BACKGROUND

[0002] Gene therapy holds enormous potential for a new era of human therapeutics. These methodologies will allow treatment for conditions that heretofore have not been addressable by standard medical practice. One area that is especially promising is the ability to add a transgene to a cell to cause that cell to express a product that previously not being produced in that cell. Examples of uses of this technology include the insertion of a gene encoding a therapeutic protein, insertion of a coding sequence encoding a protein that is somehow lacking in the cell or in the individual and insertion of a sequence that encodes a structural nucleic acid such as a microRNA.

[0003] Transgenes can be delivered to a cell by a variety of ways, such that the transgene becomes integrated into the cell's own genome and is maintained there. In recent years, a strategy for transgene integration has been developed that uses cleavage with site-specific nucleases for targeted insertion into a chosen genomic locus (see, e.g., co-owned U.S. Patent 7,888,121). Nucleases, such as zinc finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs), or nuclease systems such as the CRISPR/Cas system (utilizing an engineered guide RNA), are specific for targeted genes and can be utilized such that the transgene construct is inserted by either homology directed repair (HDR) or by end capture during non-homologous end joining (NHEJ) driven processes.

[0004] Targeted loci include "safe harbor" loci such as the AAVS1, HPRT and CCR5 genes in human cells, and *Rosa26* in murine cells (see, e.g., co-owned United States Patent Publication Nos. 20080299580; 20080159996 and 201000218264 and United States Patent Application 13/660,821). Nuclease-mediated integration offers the prospect of improved transgene expression, increased safety and expressional durability, as compared to classic integration approaches that rely on random integration of the transgene, since it allows exact transgene positioning for a minimal risk of gene silencing or activation of nearby oncogenes.

[0005] While delivery of the transgene to the target cell is one hurdle that must be overcome to fully enact this technology, another issue that must be conquered is insuring that after the transgene is inserted into the cell and is expressed, the gene product so encoded must reach the necessary location within the organism, and be made in sufficient local concentrations to be efficacious. For diseases characterized by the lack of a protein or by the presence of an aberrant non-functional one, delivery of a transgene encoded wild type protein can be extremely helpful.

[0006] Lysosomal storage diseases (LSDs) are a group of rare metabolic monogenic diseases characterized by the lack of functional individual lysosomal proteins normally involved in the breakdown of waste lipids, glycoproteins and mucopolysaccharides. These diseases are characterized by a buildup of these compounds in the cell since it is unable to process them for recycling due to the mis-functioning

of a specific enzyme. The most common examples are Gaucher's (glucocerebrosidase deficiency- gene name: GBA), Fabry's (α galactosidase deficiency- GLA), Hunter's (iduronate-2-sulfatase deficiency-IDS), Hurler's (alpha-L iduronidase deficiency- IDUA), and Niemann-Pick's (sphingomyelin phosphodiesterase 1 deficiency- SMPD1) diseases. When grouped all together, LSDs have an incidence in the population of about 1 in 7000 births. These diseases have devastating effects on those afflicted with them. They are usually first diagnosed in babies who may have characteristic facial and body growth patterns and may have moderate to severe mental retardation. Treatment options include enzyme replacement therapy (ERT) where the missing enzyme is given to the patient, usually through intravenous injection in large doses. Such treatment is only to treat the symptoms and is not curative, thus the patient must be given repeated dosing of these proteins for the rest of their lives, and potentially may develop neutralizing antibodies to the injected protein. Often these proteins have a short serum half life, and so the patient must also endure frequent infusions of the protein. For example, Gaucher's disease patients receiving the Cerezyme® product (imiglucerase) must have infusions three times per week. Production and purification of the enzymes is also problematic, and so the treatments are very costly (>\$100,000 per year per patient).

[0007] Thus, there remains a need for additional methods and compositions that can be used to treat a monogenic disease (e.g. Lysosomal storage diseases) through genome editing, and methods to deliver an expressed transgene encoded gene product at a therapeutically relevant level.

SUMMARY

[0008] The invention provides one or more transgenes and one or more zinc finger nucleases (ZFNs) for use in a method of treating a lysosomal storage disease, the method comprising administering the one or more transgenes and the one or more ZFNs to a subject in need thereof to generate a cell that produces a protein that treats the lysosomal storage disease, wherein the transgene encoding the protein is inserted into an endogenous albumin locus of the cell using the one or more ZFNs, and wherein the lysosomal storage disease is Gaucher's, Hunter's, Hurler's or Fabry's disease.

The invention also provides one or more transgenes and one or more polynucleotides encoding one or more ZFNs for use in a method of treating a lysosomal storage disease, the method comprising administering the one or more transgenes and one or more polynucleotides encoding the one or more ZFNs, to a subject in need thereof to generate a cell that produces a protein that treats the lysosomal storage disease, wherein the transgene encoding the protein is inserted into an endogenous albumin locus of the cell using the one or more ZFNs, and wherein the lysosomal storage disease is Gaucher's, Hunter's, Hurler's or Fabry's disease.

[0009] Disclosed herein are methods and compositions for treating a monogenic disease. Described are methods for insertion of a transgene sequence into a suitable target cell wherein the transgene encodes a protein that treats the disease. The therapeutic protein may be excreted from the target cell such that it is able to affect or be taken up by other cells that do not harbor the transgene. Also disclosed are methods for the production of a cell (e.g., a mature or undifferentiated cell) that produces high levels of a therapeutic where the introduction of a population of these altered cells into a patient will supply that needed protein to treat a disease or condition.

[0010] In one aspect, described herein is a zinc-finger protein (ZFP) that binds to target site in a region of interest (e.g., a disease associated gene, a highly expressed gene, an albumin gene or other or safe harbor gene) in a genome, wherein the ZFP comprises one or more engineered zinc-finger binding

domains. The ZFN can be a zinc-finger nuclease (ZFN) that cleaves a target genomic region of interest, wherein the ZFN comprises one or more engineered zinc-finger binding domains and a nuclease cleavage domain or cleavage half-domain. Cleavage domains and cleavage half domains can be obtained, for example, from various restriction endonucleases and/or homing endonucleases. In one embodiment, the cleavage half-domains are derived from a Type IIS restriction endonuclease (e.g., *Fok I*). In certain embodiments, the zinc finger domain recognizes a target site in a disease associated or safe harbor gene such as albumin (e.g., a zinc finger protein having 5 or 6 fingers with the recognition helix regions shown in a single row of Table 3).

[0011] In another aspect, described herein is a TALE protein (Transcription activator like) that binds to target site in a region of interest (e.g., a highly expressed gene, a disease associated gene or a safe harbor gene) in a genome, wherein the TALE comprises one or more engineered TALE binding domains. The TALE can be a nuclease (TALEN) that cleaves a target genomic region of interest, wherein the TALEN comprises one or more engineered TALE DNA binding domains and a nuclease cleavage domain or cleavage half-domain. Cleavage domains and cleavage half domains can be obtained, for example, from various restriction endonucleases and/or homing endonucleases. The cleavage half-domains may be derived from a Type IIS restriction endonuclease (e.g., *Fok I*). The TALE DNA binding domain may recognize a target site in a highly expressed, disease associated, or safe harbor gene.

[0012] In another aspect, described herein is a CRISPR/Cas system that binds to target site in a region of interest (e.g., a highly expressed gene, a disease associated gene or a safe harbor gene) in a genome, wherein the CRISPR/Cas system comprises a CRISPR/Cas nuclease and an engineered crRNA/tracrRNA (or single guide RNA). The CRISPR/Cas system may recognize a target site in a highly expressed, disease associated, or safe harbor gene.

[0013] The ZFN,TALEN, and/or CRISPR/Cas system as described herein may bind to and/or cleave the region of interest in a coding or non-coding region within or adjacent to the gene, such as, for example, a leader sequence, trailer sequence or intron, or within a non-transcribed region, either upstream or downstream of the coding region. In certain aspects of the disclosure, the ZFN, TALEN, and/or CRISPR/Cas system binds to and/or cleaves a highly expressed gene, for example a globin gene in red blood cells (RBCs). See, e.g., U.S. Application No. 61/670,451, titled "Methods and Compositions for Delivery of Biologics," filed July 11, 2012. In other aspects of the disclosure, the ZFN, TALEN, and/or CRISPR/Cas system binds to and/or cleaves a safe-harbor gene, for example a CCR5 gene, a PPP1R12C (also known as AAV S1) gene, albumin, HPRT or a *Rosa* gene. See, e.g., U.S. Patent Publication Nos. 20080299580; 20080159996 and 201000218264 and U.S. Application Nos. 61/537,349, 61/560,506, 13/660,821 and U.S. Application no. 61/670,451 titled "Methods and Compositions for Regulation of Transgene Expression" filed July 11, 2012. In addition, to aid in selection, the HPRT locus may be used (see U.S. patent applications 13/660,821 and 13/660,843). In other aspects of the disclosure, the ZFN, TALEN, and/or CRISPR/Cas system may bind to and/or cleave a disease associated gene (e.g. the gene encoding lysosomal hydrolase α -galactosidase A (AGA), related to Fabry's Disease). In another aspect, described herein are compositions comprising one or more of the zinc-finger and/or TALE nucleases or CRISPR/Cas system described herein. Also described are compositions comprising the one or more of these nucleases and donor nucleic acid. In some aspects, described are engineered nucleases or CRISPR/Cas systems capable of cleaving disease associate aberrant regulatory genes and methods of using these nucleases to treat the disease by reducing or eliminating expression of the aberrant gene product.

[0014] In one aspect, described is a method of treating a lysosomal storage disease by inserting in a corrective transgene into a suitable target cell (e.g., blood cell, liver cell, brain cell, stem cell, precursor cell, etc.) such that the product encoded by that corrective transgene is expressed. The corrective transgene may be inserted into a cell line for the *in vitro* production of the replacement protein. The cells comprising the transgene or the protein produced by the cells can be used to treat a patient in need thereof, for example following purification of the produced protein. The corrective transgene may be inserted into a target tissue in the body such that the replacement protein is produced *in vivo*. In some aspects, the expressed protein is excreted from the cell to act on or be taken up by other cells (e.g. via exportation into the blood) that lack the transgene. In some instances, the target tissue is the liver. In other instances, the target tissue is the brain. In other instances, the target is blood (e.g., vasculature). In other instances, the target is skeletal muscle. The corrective gene may comprise the wild type sequence of the functioning gene, or the sequence of the corrective transgene may be altered in some manner to give enhanced biological activity. In some aspects, the corrective transgene comprises optimized codons to increase biological activity, while in other aspects, the sequence is altered to give the resultant protein more desired function (e.g., improvement in stability, alteration of charge to alter substrate binding etc.). The transgene may be altered for reduced immunogenicity. In other cases, the transgene is altered such that the encoded protein becomes a substrate for transporter-mediated delivery in specific tissues such as the brain (see Gabathuler et al. (2010) *Neurobiology of Disease* 37: 48-57).

[0015] In another aspect, an engineered nuclease protein capable of cleaving (editing) the genome of a stem or precursor cell (e.g., blood cell precursor, liver stem cell, etc.) for introduction of a desired transgene is disclosed. In some aspects, the edited stem or precursor cells are then expanded and may be induced to differentiate into a mature edited cells *ex vivo*, and then the cells are given to the patient. In other aspects, the edited precursors (e.g., CD34+ stem cells) are given in a bone marrow transplant which, following successful implantation, proliferate producing edited cells that then differentiate and mature *in vivo* and contain the biologic expressed from the transgene. In other aspects, the edited stem cells are muscle stem cells which are then introduced into muscle tissue. In some aspects, the engineered nuclease is a Zinc Finger Nuclease (ZFN) and in others, the nuclease is a TALE nuclease (TALEN), and in other aspects, a CRISPR/Cas system is used. The nucleases may be engineered to have specificity for a safe harbor locus, a gene associated with a disease, or for a gene that is highly expressed in cells. By way of non-limiting example only, the safe harbor locus may be the AAVS1 site, the CCR5 gene, albumin or the HPRT gene while the disease associated gene may be the *GLA* gene encoding lysosomal hydrolase α -galactosidase A (See Table 2). By way of non-limiting example only, a gene that is highly expressed in red blood cells (RBCs) is beta-globin. In another aspect, the transgenic cells are sensitized *ex vivo* via electrosensitization to increase their susceptibility for disruption following exposure to an energy source (e.g. ultrasound) (see WO 2002007752).

[0016] In another aspect, described herein is a polynucleotide encoding one or more ZFN, TALEN, and/or CRISPR/Cas system described herein. The polynucleotide may be, for example, mRNA. In some aspects, the mRNA may be chemically modified (See e.g. Kormann et al, (2011) *Nature Biotechnology* 29(2):154-157).

[0017] In another aspect, described herein is a ZFN, TALEN, and/or CRISPR/Cas system expression vector comprising a polynucleotide, encoding one or more ZFN, TALEN, and/or CRISPR/Cas system described herein, operably linked to a promoter. The expression vector may be a viral vector.

[0018] In another aspect, described herein is a host cell comprising one or more ZFN, TALEN, and/or

CRISPR/Cas system expression vectors as described herein. The host cell may be stably transformed or transiently transfected or a combination thereof with one or more ZFN, TALEN, and/or CRISPR/Cas system expression vectors. The host cell may be a liver cell.

[0019] In another aspect, described herein is a method for cleaving a highly expressed, disease associated and/or safe harbor locus in a cell, the method comprising: introducing, into the cell, one or more polynucleotides encoding one or more ZFN, TALEN, and/or CRISPR/Cas system that bind(s) to a target site in the one or more target loci under conditions such that the ZFN(s), TALEN(s) or CRISPR/Cas system is (are) expressed and the one or more loci are cleaved. Non-limiting examples of ZFN, TALEN, and/or CRISPR/Cas systems that bind to highly expressed and/or safe harbor loci are disclosed in U.S. Publication Nos. 20080299580; 20080159996; and 201000218264 and U.S. applications 13/660,821, 13/660, 843, 13/624,193 and 13/624,217 and U.S. Application no. 61/670,451, titled "Methods and Compositions for Delivery of Biologics".

[0020] In other aspects of the disclosure, a genomic sequence in any target gene is replaced with the therapeutic transgene, for example using a ZFN, TALEN, and/or CRISPR/Cas system (or vector encoding said ZFN, TALEN, and/or CRISPR/Cas system) as described herein and a "donor" sequence or transgene that is inserted into the gene following targeted cleavage with the the ZFN, TALEN, and/or CRISPR/Cas system. The donor sequence may be present in the ZFN or TALEN vector, present in a separate vector (*e.g.*, Ad, AAV or LV vector) or, alternatively, may be introduced into the cell using a different nucleic acid delivery mechanism. Such insertion of a donor nucleotide sequence into the target locus (*e.g.*, highly expressed gene, disease associated gene, other safe-harbor gene, *etc.*) results in the expression of the transgene under control of the target locus's (*e.g.*, albumin, globin, *etc.*) endogenous genetic control elements. In some aspects, insertion of the transgene of interest, for example into a target gene (*e.g.*, albumin), results in expression of an intact exogenous protein sequence and lacks any amino acids encoded by the target (*e.g.*, albumin). In other aspects, the expressed exogenous protein is a fusion protein and comprises amino acids encoded by the transgene and by the endogenous locus into which the transgene is inserted (*e.g.*, from the endogenous target locus or, alternatively from sequences on the transgene that encode sequences of the target locus). The target may be any gene, for example, a safe harbor gene such as an albumin gene, an AAVS1 gene, an HPRT gene; a CCR5 gene; or a highly expressed gene such as a globin gene in an RBC (*e.g.*, beta globin or gamma globin). In some instances, the endogenous sequences will be present on the amino (N)-terminal portion of the exogenous protein, while in others, the endogenous sequences will be present on the carboxy (C)-terminal portion of the exogenous protein. In other instances, endogenous sequences will be present on both the N- and C-terminal portions of the exogenous protein. The endogenous sequences may include full-length wild-type or mutant endogenous sequences or, alternatively, may include partial endogenous amino acid sequences. The endogenous gene-transgene fusion may be located at the endogenous locus within the cell or the endogenous sequence-transgene coding sequence may be inserted into another locus within a genome (*e.g.*, an IDUA-transgene sequence inserted into an albumin, HPRT or CCR5 locus). In some aspects, the safe harbor is selected from the AAVS1, *Rosa*, albumin, HPRT or CCR5 locus (see co-owned U.S. Publication Nos. 20080299580; 20080159996; and 201000218264 and U.S. applications 13/660,821, 13/660, 843, 13/624,193 and 13/624,217 and U.S. Application no 61/670,451, titled "Methods and Compositions for Regulation of Transgene Expression" filed July 11, 2011). The disease associated gene may be selected from GLA (lysosomal hydrolase α -galactosidase A), or from one or more genes listed in Table 2.

[0021] In some embodiments the transgene is expressed such that a therapeutic protein product is retained within the cell (*e.g.*, precursor or mature cell). In other embodiments, the transgene is fused to

the extracellular domain of a membrane protein such that upon expression, a transgene fusion will result in the surface localization of the therapeutic protein. In some aspects, the extracellular domain is chosen from those proteins listed in Table 1. In some aspects, the edited cells also comprise a transmembrane protein to traffic the cells to a particular tissue type. In one aspect, the transmembrane protein is an antibody, while in others, the transmembrane protein is a receptor. In certain embodiments, the cell is a precursor (e.g., CD34+ or hematopoietic stem cell) or mature RBC. In some aspects, the therapeutic protein product encoded on the transgene is exported out of the cell to affect or be taken up by cells lacking the transgene. In certain embodiments, the cell is a liver cell which releases the therapeutic protein into the blood stream to act on distal tissues (e.g., brain).

[0022] Also disclosed are methods and compositions for the production of a cell (e.g., RBC) carrying a therapeutic protein for an LSD that can be used universally for all patients as an allogenic product. This would allow the development of a single product for the treatment of patients with a particular LSD, for example. These carriers may comprise transmembrane proteins to assist in the trafficking of the cell. In one aspect, the transmembrane protein is an antibody, while in others, the transmembrane protein is a receptor.

[0023] In one aspect, methods and compositions for the knockout of disease associated genes are disclosed. These genes may be those whose products may regulate expression of a gene in precursor or mature cell. In some aspects, the knock out is to the regulatory target site on the DNA for such proteins. In some aspects, the regulator gene is aberrant such that knock out of the gene restores normal function. In other aspects, the gene to be knocked out is a disease associated allele such that the knocking out of this diseased allele allows expression from a wild type allele and restores normal function.

[0024] In one embodiment, the transgene is expressed from the albumin promoter following insertion into the albumin locus. The biologic encoded by the transgene then may be released into the blood stream if the transgene is inserted into a hepatocyte *in vivo*. In some aspects, the transgene is delivered to the liver *in vivo* in a viral vector through intravenous injection.

[0025] In another embodiment, the transgene encodes a non-coding RNA, e.g. an shRNA. Expression of the transgene prior to cell maturation will result in a cell containing the non-coding RNA of interest.

[0026] Described herein are precursor cells (hematopoietic stem cells, muscle stem cells or CD34+ hematopoietic stem cell (HSC) cells) into which a transgene has been inserted such that mature cells derived from these precursors contain high levels of the product encoded by the transgene. These precursors may be induced pluripotent stem cells (iPSC).

[0027] In some aspects, the methods of the disclosure may be used *in vivo* in transgenic animal systems. In some aspects, the transgenic animal may be used in model development where the transgene encodes a human gene. In some instances, the transgenic animal may be knocked out at the corresponding endogenous locus, allowing the development of an *in vivo* system where the human protein may be studied in isolation. Such transgenic models may be used for screening purposes to identify small molecules, or large biomolecules or other entities which may interact with or modify the human protein of interest. In some aspects, the transgene is integrated into the selected locus (e.g., highly expressed or safe-harbor) into a stem cell (e.g., an embryonic stem cell, an induced pluripotent stem cell, a hepatic stem cell, a neural stem cell etc.) or animal embryo obtained by any of the methods described herein, and then the embryo is implanted such that a live animal is born. The animal is then

raised to sexual maturity and allowed to produce offspring wherein at least some of the offspring comprise the integrated transgene.

[0028] In a still further aspect, provided herein is a method for site specific integration of a nucleic acid sequence into an endogenous locus (*e.g.*, disease-associated, highly expressed such as globin in RBCs, or safe harbor gene such as albumin, CCR5, HPRT or *Rosa* gene) of a chromosome, for example into the chromosome of an embryo. In certain aspects, the method comprises: (a) injecting an embryo with (i) at least one DNA vector, wherein the DNA vector comprises an upstream sequence and a downstream sequence flanking the nucleic acid sequence to be integrated, and (ii) at least one RNA molecule encoding a zinc finger, TALE nuclease or CRISPR/Cas system that recognizes the site of integration in the target locus, and (b) culturing the embryo to allow expression of the ZFN, TALEN, and/or CRISPR/Cas system, wherein a double stranded break introduced into the site of integration by the ZFN, TALEN, and/or CRISPR/Cas system is repaired, via homologous recombination with the DNA vector, so as to integrate the nucleic acid sequence into the chromosome.

[0029] The methods and compounds described herein may be combined with other therapeutic agents for the treatment of subjects with lysosomal storage diseases. In some aspects, the methods and compositions are used in combination with methods and compositions to allow passage across the blood brain barrier. In other aspects, the methods and compositions are used in combination with compounds known to suppress the immune response of the subject.

[0030] A kit, comprising the ZFN, TALEN, and/or CRISPR/Cas system is also disclosed. The kit may comprise nucleic acids encoding the ZFN, TALEN, and/or CRISPR/Cas system, (*e.g.* RNA molecules or the ZFN, TALEN, and/or CRISPR/Cas system encoding genes contained in a suitable expression vector), donor molecules, expression vectors encoding the single guide RNA suitable host cell lines, instructions for performing the methods, disclosed herein and the like.

[0031] These and other aspects will be readily apparent to the skilled artisan in light of disclosure as a whole.

BRIEF DESCRIPTION OF THE DRAWINGS

[0032]

Figure 1, panels A and B, depict a composite set of gels demonstrating the results of a Cel-I mismatch assay (Surveyor™, Transgenomic) that measures cleavage at a location of interest by a nuclease pair that has been followed by an NHEJ event. NHEJ causes the insertion or deletion of nucleotide bases ("indels") which then creates a mismatch when the DNA strand is annealed with a wild-type DNA strand. Figure 1A shows the results measured when the transfection of the albumin-specific nuclease pairs into Neuro2A cells was carried out at 37°C and Figure 1B shows results when transduction of nuclease pairs under hypothermic shock (30°C). The percent mismatch, or % indels, is a measure of the nuclease activity of each pair under each condition.

Figure 2 is a schematic depicting the structure of four AAV donors designed to provide therapeutic transgenes for treatment of Fabry's, Gaucher's, Hurler's and Hunter's diseases. Each donor construct contains the AAV sequences (5'ITR and 3'ITR), flanking homology arms for insertion of the donors into the albumin locus by homology dependent mechanisms, a splice acceptor site, the DNA encoding the replacement enzyme, and a MYC-Flag Tag to allow identification of the integrated donors.

Figure 3, panels A and B, demonstrate activity of the mouse albumin ZFNs *in vivo*. Normal male mice (n=6) were administered a single dose of 200 microliter of 1.0×10^{11} total vector genomes of either AAV2/8 or AAV2/8.2 encoding the murine specific ZFN pair SBS30724 and SBS30725 to evaluate liver infectivity by detection of AAV vector genome copies and *in vivo* NHEJ activity in normal mice. Vectors were given by tail vein injection into mice as described, and 14 days post administration, mice were sacrificed, livers harvested and processed for DNA or total protein quantification. Detection of AAV vector genome copies was performed by quantitative PCR and cleavage activity of the ZFN was measured using the Cel-1 (Surveyor, Transkaryotic) assay. **Figure 3A** depicts a gel with the Cel-I results from mice given AAV2/8 containing a GFP expression cassette or AAV2/8 comprising the ZFNs, where the AAV were produced via a 293 expression system or a baculovirus system. **Figure 3B** depicts the quantitation of the lanes on the gel and shows that infection of the mice with the AAV containing the albumin specific ZFNs results in nearly 30% quantifiable NHEJ activity.

Figure 4, panels A and B, demonstrate the insertion of a huGLa transgene donor (deficient in patients afflicted with Fabry's disease) into the albumin locus in mice. Figure 4A shows a Western blot against the huGLa protein encoded by the transgene, where the arrow indicates the presumed protein. Comparison of the mouse samples from those mice that received both ZFN and donor (samples 1-1, 1-2 and 1-3) with the samples that either received only ZFN (4-1, 4-2, 4-3) or those that only received the huGLa donor ("hu Fabry donor"), samples 5-1 and 5-2 leads to identification of a band that coincides with the human liver lysate control. Figure 4B depicts ELISA results using a huGLa specific ELISA kit, where samples were analyzed from mice either 14 or 30 days following virus introduction. Error bars represent standard deviations (n=3). The results demonstrate that the mice that received both the ZFN and donor (circles) had higher amounts of huGLa signal than those that only received ZFN (squares) or only received donor (triangles).

Figure 5, panels A-D, depicts Western blots that demonstrate expression in liver homogenates of the LSD donor transgenes inserted into the albumin locus in mice. Figure 5A shows the results using the IDUA encoding transgene, Figure 5B shows the results using the GLA transgene, Figure 5C shows the results using the IDS transgene, and Figure 5D shows the results using the GBA transgene.

Figure 6 is a schematic displaying the two types of donor insertion that may occur following ZFN mediated cleavage. NHEJ mediated donor insertion will result in the entire LSD-Donor construct being integrated, whereas HDR-mediated insertion will cause only the cDNA including the F9 splice-acceptor site to be incorporated. Figure 6 also depicts the location of the two PCR primers ("mALB-OOF1" and "Acc651-SA-rev-sh") used to detect the type of integration that has occurred.

Figure 7, panels A-C, depict the results of ^{32}P radiolabeled PCR performed on liver homogenates on mice containing the integrated LSD transgenes 30 days after treatment. Figure 7A depicts the mice with the IDUA transgene, Figure 7B depicts those with the GLA transgene, and Figure 7C depicts those with the IDS transgene. In all cases, the bands indicate that insertion of the transgenes has occurred through both NHEJ-mediated and HDR-mediated integration.

Figure 8 is a schematic illustrating the design of the LSD donors containing epitope tags. The location and sequences of the Myc and Flag tags are indicated.

Figure 9, panels A and B, depicts a gel of ^{32}P radiolabeled PCR done as described above on liver homogenates from mice with integrated LSD donors containing the epitope tags. Figure 9A shows that integration occurred through both NHEJ-mediated and HDR-mediated integration for the targeted GLA, IDUA and IDS transgene. Figure 9B shows the same for the GBA transgene.

DETAILED DESCRIPTION

[0033] Disclosed herein are methods and compositions for treating or preventing a lysosomal storage disease (LSD). Disclosed are methods and compositions for insertion of a gene encoding a protein that is lacking or insufficiently expressed in the subject with the LSD such that the gene is expressed in the liver and the therapeutic (replacement) protein is expressed. Also described is the alteration of a cell (e.g., precursor or mature RBC, iPSC or liver cell) such that it produces high levels of the therapeutic and the introduction of a population of these altered cells into a patient will supply that needed protein. The transgene can encode a desired protein or structural RNA that is beneficial therapeutically in a patient in need thereof.

[0034] Thus, the methods and compositions described herein can be used to express from a transgene therapeutically beneficial proteins from the albumin locus to replace enzymes that are defective in lysosomal storage diseases. Additionally, disclosed are methods and compositions for treatment of these diseases by insertion of the sequences into highly expressed loci in cells such as liver cells.

[0035] In addition, the transgene can be introduced into patient derived cells, *e.g.* patient derived induced pluripotent stem cells (iPSCs) or other types of stems cells (embryonic or hematopoietic) for use in eventual implantation. Particularly useful is the insertion of the disease associated transgene into a hematopoietic stem cell for implantation into a patient in need thereof. As the stem cells differentiate into mature cells, they will contain high levels of the replacement protein for delivery to the tissues.

General

[0036] Practice of the methods, as well as preparation and use of the compositions disclosed herein employ, unless otherwise indicated, conventional techniques in molecular biology, biochemistry, chromatin structure and analysis, computational chemistry, cell culture, recombinant DNA and related fields as are within the skill of the art. These techniques are fully explained in the literature. See, for example, Sambrook et al. MOLECULAR CLONING: A LABORATORY MANUAL, Second edition, Cold Spring Harbor Laboratory Press, 1989 and Third edition, 2001; Ausubel et al., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, 1987 and periodic updates; the series METHODS IN ENZYMOLOGY, Academic Press, San Diego; Wolffe, CHROMATIN STRUCTURE AND FUNCTION, Third edition, Academic Press, San Diego, 1998; METHODS IN ENZYMOLOGY, Vol. 304, "Chromatin" (P.M. Wassarman and A. P. Wolffe, eds.), Academic Press, San Diego, 1999; and METHODS IN MOLECULAR BIOLOGY, Vol. 119, "Chromatin Protocols" (P.B. Becker, ed.) Humana Press, Totowa, 1999.

Definitions

[0037] The terms "nucleic acid," "polynucleotide," and "oligonucleotide" are used interchangeably and refer to a deoxyribonucleotide or ribonucleotide polymer, in linear or circular conformation, and in either single- or double-stranded form. For the purposes of the present disclosure, these terms are not to be

construed as limiting with respect to the length of a polymer. The terms can encompass known analogues of natural nucleotides, as well as nucleotides that are modified in the base, sugar and/or phosphate moieties (e.g., phosphorothioate backbones). In general, an analogue of a particular nucleotide has the same base-pairing specificity; *i.e.*, an analogue of A will base-pair with T.

[0038] The terms "polypeptide," "peptide" and "protein" are used interchangeably to refer to a polymer of amino acid residues. The term also applies to amino acid polymers in which one or more amino acids are chemical analogues or modified derivatives of corresponding naturally-occurring amino acids.

[0039] "Binding" refers to a sequence-specific, non-covalent interaction between macromolecules (e.g., between a protein and a nucleic acid). Not all components of a binding interaction need be sequence-specific (e.g., contacts with phosphate residues in a DNA backbone), as long as the interaction as a whole is sequence-specific. Such interactions are generally characterized by a dissociation constant (K_d) of 10^{-6} M^{-1} or lower. "Affinity" refers to the strength of binding: increased binding affinity being correlated with a lower K_d .

[0040] A "binding protein" is a protein that is able to bind non-covalently to another molecule. A binding protein can bind to, for example, a DNA molecule (a DNA-binding protein), an RNA molecule (an RNA-binding protein) and/or a protein molecule (a protein-binding protein). In the case of a protein-binding protein, it can bind to itself (to form homodimers, homotrimers, *etc.*) and/or it can bind to one or more molecules of a different protein or proteins. A binding protein can have more than one type of binding activity. For example, zinc finger proteins have DNA-binding, RNA-binding and protein-binding activity.

[0041] A "zinc finger DNA binding protein" (or binding domain) is a protein, or a domain within a larger protein, that binds DNA in a sequence-specific manner through one or more zinc fingers, which are regions of amino acid sequence within the binding domain whose structure is stabilized through coordination of a zinc ion. The term zinc finger DNA binding protein is often abbreviated as zinc finger protein or ZFP.

[0042] A "TALE DNA binding domain" or "TALE" is a polypeptide comprising one or more TALE repeat domains/units. The repeat domains are involved in binding of the TALE to its cognate target DNA sequence. A single "repeat unit" (also referred to as a "repeat") is typically 33-35 amino acids in length and exhibits at least some sequence homology with other TALE repeat sequences within a naturally occurring TALE protein.

[0043] Zinc finger and TALE binding domains can be "engineered" to bind to a predetermined nucleotide sequence, for example via engineering (altering one or more amino acids) of the recognition helix region of a naturally occurring zinc finger or TALE protein. Therefore, engineered DNA binding proteins (zinc fingers or TALEs) are proteins that are non-naturally occurring. Non-limiting examples of methods for engineering DNA-binding proteins are design and selection. A designed DNA binding protein is a protein not occurring in nature whose design/composition results principally from rational criteria. Rational criteria for design include application of substitution rules and computerized algorithms for processing information in a database storing information of existing ZFP and/or TALE designs and binding data. See, for example, US Patents 6,140,081; 6,453,242; and 6,534,261; see also WO 98/53058; WO 98/53059; WO 98/53060; WO 02/016536 and WO 03/016496 and U.S. Publication No. 20110301073.

[0044] A "selected" zinc finger protein or TALE is a protein not found in nature whose production results primarily from an empirical process such as phage display, interaction trap or hybrid selection. See e.g.,

US 5,789,538; US 5,925,523; US 6,007,988; US 6,013,453; US 6,200,759; WO 95/19431; WO 96/06166; WO 98/53057; WO 98/54311; WO 00/27878; WO 01/60970 WO 01/88197; WO 02/099084 and U.S. Publication No. 20110301073.

[0045] "Recombination" refers to a process of exchange of genetic information between two polynucleotides. For the purposes of this disclosure, "homologous recombination (HR)" refers to the specialized form of such exchange that takes place, for example, during repair of double-strand breaks in cells via homology-directed repair mechanisms. This process requires nucleotide sequence homology, uses a "donor" molecule to template repair of a "target" molecule (*i.e.*, the one that experienced the double-strand break), and is variously known as "non-crossover gene conversion" or "short tract gene conversion," because it leads to the transfer of genetic information from the donor to the target. Without wishing to be bound by any particular theory, such transfer can involve mismatch correction of heteroduplex DNA that forms between the broken target and the donor, and/or "synthesis-dependent strand annealing," in which the donor is used to re-synthesize genetic information that will become part of the target, and/or related processes. Such specialized HR often results in an alteration of the sequence of the target molecule such that part or all of the sequence of the donor polynucleotide is incorporated into the target polynucleotide.

[0046] In the methods of the disclosure, one or more targeted nucleases as described herein create a double-stranded break in the target sequence (e.g., cellular chromatin) at a predetermined site, and a "donor" polynucleotide, having homology to the nucleotide sequence in the region of the break, can be introduced into the cell. The presence of the double-stranded break has been shown to facilitate integration of the donor sequence. The donor sequence may be physically integrated or, alternatively, the donor polynucleotide is used as a template for repair of the break via homologous recombination, resulting in the introduction of all or part of the nucleotide sequence as in the donor into the cellular chromatin. Thus, a first sequence in cellular chromatin can be altered and, in certain embodiments, can be converted into a sequence present in a donor polynucleotide. Thus, the use of the terms "replace" or "replacement" can be understood to represent replacement of one nucleotide sequence by another, (*i.e.*, replacement of a sequence in the informational sense), and does not necessarily require physical or chemical replacement of one polynucleotide by another.

[0047] In any of the methods described herein, additional pairs of zinc-finger or TALEN proteins can be used for additional double-stranded cleavage of additional target sites within the cell.

[0048] In certain methods for targeted recombination and/or replacement and/or alteration of a sequence in a region of interest in cellular chromatin, a chromosomal sequence is altered by homologous recombination with an exogenous "donor" nucleotide sequence. Such homologous recombination is stimulated by the presence of a double-stranded break in cellular chromatin, if sequences homologous to the region of the break are present.

[0049] In any of the methods described herein, the first nucleotide sequence (the "donor sequence") can contain sequences that are homologous, but not identical, to genomic sequences in the region of interest, thereby stimulating homologous recombination to insert a non-identical sequence in the region of interest. Thus, in certain embodiments, portions of the donor sequence that are homologous to sequences in the region of interest exhibit between about 80 to 99% (or any integer therebetween) sequence identity to the genomic sequence that is replaced. In other embodiments, the homology between the donor and genomic sequence is higher than 99%, for example if only 1 nucleotide differs as between donor and genomic sequences of over 100 contiguous base pairs. In certain cases, a non-

homologous portion of the donor sequence can contain sequences not present in the region of interest, such that new sequences are introduced into the region of interest. In these instances, the non-homologous sequence is generally flanked by sequences of 50-1,000 base pairs (or any integral value therebetween) or any number of base pairs greater than 1,000, that are homologous or identical to sequences in the region of interest. In other embodiments, the donor sequence is non-homologous to the first sequence, and is inserted into the genome by non-homologous recombination mechanisms.

[0050] Any of the methods described herein can be used for partial or complete inactivation of one or more target sequences in a cell by targeted integration of donor sequence that disrupts expression of the gene(s) of interest. Cell lines with partially or completely inactivated genes are also provided.

[0051] Furthermore, the methods of targeted integration as described herein can also be used to integrate one or more exogenous sequences. The exogenous nucleic acid sequence can comprise, for example, one or more genes or cDNA molecules, or any type of coding or non-coding sequence, as well as one or more control elements (e.g., promoters). In addition, the exogenous nucleic acid sequence may produce one or more RNA molecules (e.g., small hairpin RNAs (shRNAs), inhibitory RNAs (RNAis), microRNAs (miRNAs), *etc.*).

[0052] "Cleavage" refers to the breakage of the covalent backbone of a DNA molecule. Cleavage can be initiated by a variety of methods including, but not limited to, enzymatic or chemical hydrolysis of a phosphodiester bond. Both single-stranded cleavage and double-stranded cleavage are possible, and double-stranded cleavage can occur as a result of two distinct single-stranded cleavage events. DNA cleavage can result in the production of either blunt ends or staggered ends. In certain embodiments, fusion polypeptides are used for targeted double-stranded DNA cleavage.

[0053] A "cleavage half-domain" is a polypeptide sequence which, in conjunction with a second polypeptide (either identical or different) forms a complex having cleavage activity (preferably double-strand cleavage activity). The terms "first and second cleavage half-domains;" "+ and - cleavage half-domains" and "right and left cleavage half-domains" are used interchangeably to refer to pairs of cleavage half-domains that dimerize.

[0054] An "engineered cleavage half-domain" is a cleavage half-domain that has been modified so as to form obligate heterodimers with another cleavage half-domain (e.g., another engineered cleavage half-domain). See, *also*, U.S. Patent Publication Nos. 2005/0064474, 2007/0218528, 2008/0131962 and 2011/0201055.

[0055] The term "sequence" refers to a nucleotide sequence of any length, which can be DNA or RNA; can be linear, circular or branched and can be either single-stranded or double stranded. The term "donor sequence" refers to a nucleotide sequence that is inserted into a genome. A donor sequence can be of any length, for example between 2 and 10,000 nucleotides in length (or any integer value therebetween or thereabove), preferably between about 100 and 1,000 nucleotides in length (or any integer therebetween), more preferably between about 200 and 500 nucleotides in length.

[0056] A "disease associated gene" is one that is defective in some manner in a monogenic disease. Non-limiting examples of monogenic diseases include severe combined immunodeficiency, cystic fibrosis, lysosomal storage diseases (e.g. Gaucher's, Hurler's Hunter's, Fabry's, Neimann-Pick, Tay-Sach's etc), sickle cell anemia, and thalassemia.

[0057] "Chromatin" is the nucleoprotein structure comprising the cellular genome. Cellular chromatin comprises nucleic acid, primarily DNA, and protein, including histones and non-histone chromosomal proteins. The majority of eukaryotic cellular chromatin exists in the form of nucleosomes, wherein a nucleosome core comprises approximately 150 base pairs of DNA associated with an octamer comprising two each of histones H2A, H2B, H3 and H4; and linker DNA (of variable length depending on the organism) extends between nucleosome cores. A molecule of histone H1 is generally associated with the linker DNA. For the purposes of the present disclosure, the term "chromatin" is meant to encompass all types of cellular nucleoprotein, both prokaryotic and eukaryotic. Cellular chromatin includes both chromosomal and episomal chromatin.

[0058] A "chromosome," is a chromatin complex comprising all or a portion of the genome of a cell. The genome of a cell is often characterized by its karyotype, which is the collection of all the chromosomes that comprise the genome of the cell. The genome of a cell can comprise one or more chromosomes.

[0059] An "episome" is a replicating nucleic acid, nucleoprotein complex or other structure comprising a nucleic acid that is not part of the chromosomal karyotype of a cell. Examples of episomes include plasmids and certain viral genomes.

[0060] A "target site" or "target sequence" is a nucleic acid sequence that defines a portion of a nucleic acid to which a binding molecule will bind, provided sufficient conditions for binding exist.

[0061] An "exogenous" molecule is a molecule that is not normally present in a cell, but can be introduced into a cell by one or more genetic, biochemical or other methods. "Normal presence in the cell" is determined with respect to the particular developmental stage and environmental conditions of the cell. Thus, for example, a molecule that is present only during embryonic development of muscle is an exogenous molecule with respect to an adult muscle cell. Similarly, a molecule induced by heat shock is an exogenous molecule with respect to a non-heat-shocked cell. An exogenous molecule can comprise, for example, a functioning version of a malfunctioning endogenous molecule or a malfunctioning version of a normally-functioning endogenous molecule.

[0062] An exogenous molecule can be, among other things, a small molecule, such as is generated by a combinatorial chemistry process, or a macromolecule such as a protein, nucleic acid, carbohydrate, lipid, glycoprotein, lipoprotein, polysaccharide, any modified derivative of the above molecules, or any complex comprising one or more of the above molecules. Nucleic acids include DNA and RNA, can be single- or double-stranded; can be linear, branched or circular; and can be of any length. Nucleic acids include those capable of forming duplexes, as well as triplex-forming nucleic acids. See, for example, U.S. Patent Nos. 5,176,996 and 5,422,251. Proteins include, but are not limited to, DNA-binding proteins, transcription factors, chromatin remodeling factors, methylated DNA binding proteins, polymerases, methylases, demethylases, acetylases, deacetylases, kinases, phosphatases, integrases, recombinases, ligases, topoisomerases, gyrases and helicases.

[0063] An exogenous molecule can be the same type of molecule as an endogenous molecule, *e.g.*, an exogenous protein or nucleic acid. For example, an exogenous nucleic acid can comprise an infecting viral genome, a plasmid or episome introduced into a cell, or a chromosome that is not normally present in the cell. Methods for the introduction of exogenous molecules into cells are known to those of skill in the art and include, but are not limited to, lipid-mediated transfer (*i.e.*, liposomes, including neutral and cationic lipids), electroporation, direct injection, cell fusion, particle bombardment, calcium phosphate co-precipitation, DEAE-dextran-mediated transfer and viral vector-mediated transfer. An exogenous

molecule can also be the same type of molecule as an endogenous molecule but derived from a different species than the cell is derived from. For example, a human nucleic acid sequence may be introduced into a cell line originally derived from a mouse or hamster.

[0064] By contrast, an "endogenous" molecule is one that is normally present in a particular cell at a particular developmental stage under particular environmental conditions. For example, an endogenous nucleic acid can comprise a chromosome, the genome of a mitochondrion, chloroplast or other organelle, or a naturally-occurring episomal nucleic acid. Additional endogenous molecules can include proteins, for example, transcription factors and enzymes.

[0065] A "fusion" molecule is a molecule in which two or more subunit molecules are linked, preferably covalently. The subunit molecules can be the same chemical type of molecule, or can be different chemical types of molecules. Examples of the first type of fusion molecule include, but are not limited to, fusion proteins (for example, a fusion between a ZFP or TALE DNA-binding domain and one or more activation domains) and fusion nucleic acids (for example, a nucleic acid encoding the fusion protein described *supra*). Examples of the second type of fusion molecule include, but are not limited to, a fusion between a triplex-forming nucleic acid and a polypeptide, and a fusion between a minor groove binder and a nucleic acid.

[0066] Expression of a fusion protein in a cell can result from delivery of the fusion protein to the cell or by delivery of a polynucleotide encoding the fusion protein to a cell, wherein the polynucleotide is transcribed, and the transcript is translated, to generate the fusion protein. Trans-splicing, polypeptide cleavage and polypeptide ligation can also be involved in expression of a protein in a cell. Methods for polynucleotide and polypeptide delivery to cells are presented elsewhere in this disclosure.

[0067] A "gene," for the purposes of the present disclosure, includes a DNA region encoding a gene product (see *infra*), as well as all DNA regions which regulate the production of the gene product, whether or not such regulatory sequences are adjacent to coding and/or transcribed sequences. Accordingly, a gene includes, but is not necessarily limited to, promoter sequences, terminators, translational regulatory sequences such as ribosome binding sites and internal ribosome entry sites, enhancers, silencers, insulators, boundary elements, replication origins, matrix attachment sites and locus control regions.

[0068] "Gene expression" refers to the conversion of the information, contained in a gene, into a gene product. A gene product can be the direct transcriptional product of a gene (*e.g.*, mRNA, tRNA, rRNA, antisense RNA, ribozyme, structural RNA or any other type of RNA) or a protein produced by translation of an mRNA. Gene products also include RNAs which are modified, by processes such as capping, polyadenylation, methylation, and editing, and proteins modified by, for example, methylation, acetylation, phosphorylation, ubiquitination, ADP-ribosylation, myristilation, and glycosylation.

[0069] "Modulation" of gene expression refers to a change in the activity of a gene. Modulation of expression can include, but is not limited to, gene activation and gene repression. Genome editing (*e.g.*, cleavage, alteration, inactivation, random mutation) can be used to modulate expression. Gene inactivation refers to any reduction in gene expression as compared to a cell that does not include a ZFP or TALEN as described herein. Thus, gene inactivation may be partial or complete.

[0070] A "region of interest" is any region of cellular chromatin, such as, for example, a gene or a non-coding sequence within or adjacent to a gene, in which it is desirable to bind an exogenous molecule.

Binding can be for the purposes of targeted DNA cleavage and/or targeted recombination. A region of interest can be present in a chromosome, an episome, an organellar genome (e.g., mitochondrial, chloroplast), or an infecting viral genome, for example. A region of interest can be within the coding region of a gene, within transcribed non-coding regions such as, for example, leader sequences, trailer sequences or introns, or within non-transcribed regions, either upstream or downstream of the coding region. A region of interest can be as small as a single nucleotide pair or up to 2,000 nucleotide pairs in length, or any integral value of nucleotide pairs.

[0071] "Eukaryotic" cells include, but are not limited to, fungal cells (such as yeast), plant cells, animal cells, mammalian cells and human cells (e.g., T-cells).

[0072] "Red Blood Cells" (RBCs) or erythrocytes are terminally differentiated cells derived from hematopoietic stem cells. They lack a nucleus and most cellular organelles. RBCs contain hemoglobin to carry oxygen from the lungs to the peripheral tissues. In fact 33% of an individual RBC is hemoglobin. They also carry CO₂ produced by cells during metabolism out of the tissues and back to the lungs for release during exhale. RBCs are produced in the bone marrow in response to blood hypoxia which is mediated by release of erythropoietin (EPO) by the kidney. EPO causes an increase in the number of proerythroblasts and shortens the time required for full RBC maturation. After approximately 120 days, since the RBC do not contain a nucleus or any other regenerative capabilities, the cells are removed from circulation by either the phagocytic activities of macrophages in the liver, spleen and lymph nodes (~90%) or by hemolysis in the plasma (~10%). Following macrophage engulfment, chemical components of the RBC are broken down within vacuoles of the macrophages due to the action of lysosomal enzymes.

[0073] "Secretory tissues" are those tissues in an animal that secrete products out of the individual cell into a lumen of some type which are typically derived from epithelium. Examples of secretory tissues that are localized to the gastrointestinal tract include the cells that line the gut, the pancreas, and the gallbladder. Other secretory tissues include the liver, tissues associated with the eye and mucous membranes such as salivary glands, mammary glands, the prostate gland, the pituitary gland and other members of the endocrine system. Additionally, secretory tissues include individual cells of a tissue type which are capable of secretion.

[0074] The terms "operative linkage" and "operatively linked" (or "operably linked") are used interchangeably with reference to a juxtaposition of two or more components (such as sequence elements), in which the components are arranged such that both components function normally and allow the possibility that at least one of the components can mediate a function that is exerted upon at least one of the other components. By way of illustration, a transcriptional regulatory sequence, such as a promoter, is operatively linked to a coding sequence if the transcriptional regulatory sequence controls the level of transcription of the coding sequence in response to the presence or absence of one or more transcriptional regulatory factors. A transcriptional regulatory sequence is generally operatively linked in *cis* with a coding sequence, but need not be directly adjacent to it. For example, an enhancer is a transcriptional regulatory sequence that is operatively linked to a coding sequence, even though they are not contiguous.

[0075] With respect to fusion polypeptides, the term "operatively linked" can refer to the fact that each of the components performs the same function in linkage to the other component as it would if it were not so linked. For example, with respect to a fusion polypeptide in which a ZFP or TALE DNA-binding domain is fused to an activation domain, the ZFP or TALE DNA-binding domain and the activation

domain are in operative linkage if, in the fusion polypeptide, the ZFP or TALE DNA-binding domain portion is able to bind its target site and/or its binding site, while the activation domain is able to up-regulate gene expression. When a fusion polypeptide in which a ZFP or TALE DNA-binding domain is fused to a cleavage domain, the ZFP or TALE DNA-binding domain and the cleavage domain are in operative linkage if, in the fusion polypeptide, the ZFP or TALE DNA-binding domain portion is able to bind its target site and/or its binding site, while the cleavage domain is able to cleave DNA in the vicinity of the target site.

[0076] A "functional fragment" of a protein, polypeptide or nucleic acid is a protein, polypeptide or nucleic acid whose sequence is not identical to the full-length protein, polypeptide or nucleic acid, yet retains the same function as the full-length protein, polypeptide or nucleic acid. A functional fragment can possess more, fewer, or the same number of residues as the corresponding native molecule, and/or can contain one or more amino acid or nucleotide substitutions. Methods for determining the function of a nucleic acid (*e.g.*, coding function, ability to hybridize to another nucleic acid) are well-known in the art. Similarly, methods for determining protein function are well-known. For example, the DNA-binding function of a polypeptide can be determined, for example, by filter-binding, electrophoretic mobility-shift, or immunoprecipitation assays. DNA cleavage can be assayed by gel electrophoresis. See Ausubel *et al.*, *supra*. The ability of a protein to interact with another protein can be determined, for example, by co-immunoprecipitation, two-hybrid assays or complementation, both genetic and biochemical. See, for example, Fields et al. (1989) Nature 340:245-246; U.S. Patent No. 5,585,245 and PCT WO 98/44350.

[0077] A "vector" is capable of transferring gene sequences to target cells. Typically, "vector construct," "expression vector," and "gene transfer vector," mean any nucleic acid construct capable of directing the expression of a gene of interest and which can transfer gene sequences to target cells. Thus, the term includes cloning, and expression vehicles, as well as integrating vectors.

[0078] A "reporter gene" or "reporter sequence" refers to any sequence that produces a protein product that is easily measured, preferably although not necessarily in a routine assay. Suitable reporter genes include, but are not limited to, sequences encoding proteins that mediate antibiotic resistance (*e.g.*, ampicillin resistance, neomycin resistance, G418 resistance, puromycin resistance), sequences encoding colored or fluorescent or luminescent proteins (*e.g.*, green fluorescent protein, enhanced green fluorescent protein, red fluorescent protein, luciferase), and proteins which mediate enhanced cell growth and/or gene amplification (*e.g.*, dihydrofolate reductase). Epitope tags include, for example, one or more copies of FLAG, His, myc, Tap, HA or any detectable amino acid sequence. "Expression tags" include sequences that encode reporters that may be operably linked to a desired gene sequence in order to monitor expression of the gene of interest.

[0079] The terms "subject" and "patient" are used interchangeably and refer to mammals such as human patients and non-human primates, as well as experimental animals such as rabbits, dogs, cats, rats, mice, and other animals. Accordingly, the term "subject" or "patient" as used herein means any mammalian patient or subject to which the altered cells as described herein and/or proteins produced by the altered cells can be administered. Subjects of the present disclosure include those having an LSD.

Nucleases

[0080] Described herein are compositions, particularly nucleases, which are useful targeting a gene for the insertion of a transgene, for example, nucleases that are specific for a safe-harbor gene such as

albumin. In certain embodiments, the nuclease is naturally occurring. In other embodiments, the nuclease is non-naturally occurring, i.e., engineered in the DNA-binding domain and/or cleavage domain. For example, the DNA-binding domain of a naturally-occurring nuclease or nuclease system may be altered to bind to a selected target site (e.g., a meganuclease that has been engineered to bind to site different than the cognate binding site or a CRISPR/Cas system utilizing an engineered single guide RNA). The nuclease may comprise heterologous DNA-binding and cleavage domains (e.g., zinc finger nucleases; TAL-effector nucleases; meganuclease DNA-binding domains with heterologous cleavage domains).

A. DNA-binding domains

[0081] In certain aspects, the nuclease is a meganuclease (homing endonuclease). Naturally-occurring meganucleases recognize 15-40 base-pair cleavage sites and are commonly grouped into four families: the LAGLIDADG family, the GIY-YIG family, the His-Cyst box family and the HNH family. Exemplary homing endonucleases include I-SceI, I-CeuI, PI-PspI, PI-Sce, I-SceIV, I-CsmI, I-PanI, I-SceII, I-PpoI, I-SceIII, I-CreI, I-TevI, I-TevII and I-TevIII. Their recognition sequences are known. See also U.S. Patent No. 5,420,032; U.S. Patent No. 6,833,252; Belfort et al. (1997) *Nucleic Acids Res.* 25:3379-3388; Dujon et al. (1989) *Gene* 82:115-118; Perler et al. (1994) *Nucleic Acids Res.* 22, 1125-1127; Jasin (1996) *Trends Genet.* 12:224-228; Gimble et al. (1996) *J. Mol. Biol.* 263:163-180; Argast et al. (1998) *J. Mol. Biol.* 280:345-353 and the New England Biolabs catalogue.

[0082] In certain aspects, the nuclease comprises an engineered (non-naturally occurring) homing endonuclease (meganuclease). The recognition sequences of homing endonucleases and meganucleases such as I-SceI, I-CeuI, PI-PspI, PI-Sce, I-SceIV, I-CsmI, I-PanI, I-SceII, I-PpoI, I-SceIII, I-CreI, I-TevI, I-TevII and I-TevIII are known. See also U.S. Patent No. 5,420,032; U.S. Patent No. 6,833,252; Belfort et al. (1997) *Nucleic Acids Res.* 25:3379-3388; Dujon et al. (1989) *Gene* 82:115-118; Perler et al. (1994) *Nucleic Acids Res.* 22, 1125-1127; Jasin (1996) *Trends Genet.* 12:224-228; Gimble et al. (1996) *J. Mol. Biol.* 263:163-180; Argast et al. (1998) *J. Mol. Biol.* 280:345-353 and the New England Biolabs catalogue. In addition, the DNA-binding specificity of homing endonucleases and meganucleases can be engineered to bind non-natural target sites. See, for example, Chevalier et al. (2002) *Molec. Cell* 10:895-905; Epinat et al. (2003) *Nucleic Acids Res.* 31:2952-2962; Ashworth et al. (2006) *Nature* 441:656-659; Paques et al. (2007) *Current Gene Therapy* 7:49-66; U.S. Patent Publication No. 20070117128. The DNA-binding domains of the homing endonucleases and meganucleases may be altered in the context of the nuclease as a whole (i.e., such that the nuclease includes the cognate cleavage domain) or may be fused to a heterologous cleavage domain.

[0083] In other aspects, the DNA-binding domain comprises a naturally occurring or engineered (non-naturally occurring) TAL effector DNA binding domain. See, e.g., U.S. Patent Publication No. 20110301073. The plant pathogenic bacteria of the genus *Xanthomonas* are known to cause many diseases in important crop plants. Pathogenicity of *Xanthomonas* depends on a conserved type III secretion (T3S) system which injects more than 25 different effector proteins into the plant cell. Among these injected proteins are transcription activator-like effectors (TALE) which mimic plant transcriptional activators and manipulate the plant transcriptome (see Kay et al (2007) *Science* 318:648-651). These proteins contain a DNA binding domain and a transcriptional activation domain. One of the most well characterized TALEs is AvrBs3 from *Xanthomonas campestris* pv. *Vesicatoria* (see Bonas et al (1989) *Mol Gen Genet* 218: 127-136 and WO2010079430). TALEs contain a centralized domain of tandem repeats, each repeat containing approximately 34 amino acids, which are key to the DNA binding

specificity of these proteins. In addition, they contain a nuclear localization sequence and an acidic transcriptional activation domain (for a review see Schornack S, et al (2006) J Plant Physiol 163(3): 256-272). In addition, in the phytopathogenic bacteria *Ralstonia solanacearum* two genes, designated brg11 and hpx17 have been found that are homologous to the AvrBs3 family of *Xanthomonas* in the *R. solanacearum* biovar 1 strain GMI1000 and in the biovar 4 strain RS1000 (See Heuer et al (2007) Appl and Envir Micro 73(13): 4379-4384). These genes are 98.9% identical in nucleotide sequence to each other but differ by a deletion of 1,575 bp in the repeat domain of hpx17. However, both gene products have less than 40% sequence identity with AvrBs3 family proteins of *Xanthomonas*.

[0084] Thus, in some aspects, the DNA binding domain that binds to a target site in a target locus (e.g., albumin or other safe harbor) is an engineered domain from a TAL effector similar to those derived from the plant pathogens *Xanthomonas* (see Boch et al, (2009) Science 326: 1509-1512 and Moscou and Bogdanove, (2009) Science 326: 1501) and *Ralstonia* (see Heuer et al (2007) Applied and Environmental Microbiology 73(13): 4379-4384); U.S. Patent Publication Nos. 20110301073 and 20110145940. See, e.g., albumin TALENs in U.S. Application No. 13/624,193 and 13/624,217.

[0085] The DNA binding domain comprises a zinc finger protein (e.g., a zinc finger protein that binds to a target site in an albumin). Preferably, the zinc finger protein is non-naturally occurring in that it is engineered to bind to a target site of choice. See, for example, Beerli et al. (2002) Nature Biotechnol. 20:135-141; Pabo et al. (2001) Ann. Rev. Biochem. 70:313-340; Isalan et al. (2001) Nature Biotechnol. 19:656-660; Segal et al. (2001) Curr. Opin. Biotechnol. 12:632-637; Choo et al. (2000) Curr. Opin. Struct. Biol. 10:411-416; U.S. Patent Nos. 6,453,242; 6,534,261; 6,599,692; 6,503,717; 6,689,558; 7,030,215; 6,794,136; 7,067,317; 7,262,054; 7,070,934; 7,361,635; 7,253,273; and U.S. Patent Publication Nos. 2005/0064474; 2007/0218528; 2005/0267061.

[0086] An engineered zinc finger binding or TALE domain can have a novel binding specificity, compared to a naturally-occurring zinc finger protein. Engineering methods include, but are not limited to, rational design and various types of selection. Rational design includes, for example, using databases comprising triplet (or quadruplet) nucleotide sequences and individual zinc finger amino acid sequences, in which each triplet or quadruplet nucleotide sequence is associated with one or more amino acid sequences of zinc fingers which bind the particular triplet or quadruplet sequence. See, for example, co-owned U.S. Patents 6,453,242 and 6,534,261.

[0087] Exemplary selection methods, including phage display and two-hybrid systems, are disclosed in US Patents 5,789,538; 5,925,523; 6,007,988; 6,013,453; 6,410,248; 6,140,466; 6,200,759; and 6,242,568; as well as WO 98/37186; WO 98/53057; WO 00/27878; WO 01/88197 and GB 2,338,237. In addition, enhancement of binding specificity for zinc finger binding domains has been described, for example, in co-owned WO 02/077227.

[0088] In addition, as disclosed in these and other references, DNA domains (e.g., multi-fingered zinc finger proteins or TALE domains) may be linked together using any suitable linker sequences, including for example, linkers of 5 or more amino acids in length. See, also, U.S. Patent Nos. 6,479,626; 6,903,185; and 7,153,949 for exemplary linker sequences 6 or more amino acids in length. The DNA binding proteins described herein may include any combination of suitable linkers between the individual zinc fingers of the protein. In addition, enhancement of binding specificity for zinc finger binding domains has been described, for example, in co-owned WO 02/077227.

[0089] Selection of target sites; DNA-binding domains and methods for design and construction of fusion

proteins (and polynucleotides encoding same) are known to those of skill in the art and described in detail in U.S. Patent Nos. 6,140,0815; 789,538; 6,453,242; 6,534,261; 5,925,523; 6,007,988; 6,013,453; 6,200,759; WO 95/19431; WO 96/06166; WO 98/53057; WO 98/54311; WO 00/27878; WO 01/60970 WO 01/88197; WO 02/099084; WO 98/53058; WO 98/53059; WO 98/53060; WO 02/016536 and WO 03/016496 and U.S. Publication No. 20110301073.

[0090] In addition, as disclosed in these and other references, DNA-binding domains (e.g., multi-fingered zinc finger proteins) may be linked together using any suitable linker sequences, including for example, linkers of 5 or more amino acids in length. See, also, U.S. Patent Nos. 6,479,626; 6,903,185; and 7,153,949 for exemplary linker sequences 6 or more amino acids in length. The proteins described herein may include any combination of suitable linkers between the individual zinc fingers of the protein.

B. Cleavage Domains

[0091] Any suitable cleavage domain can be operatively linked to a DNA-binding domain to form a nuclease. For example, ZFP DNA-binding domains have been fused to nuclease domains to create ZFNs - a functional entity that is able to recognize its intended nucleic acid target through its engineered (ZFP) DNA binding domain and cause the DNA to be cut near the ZFP binding site via the nuclease activity. See, e.g., Kim et al. (1996) Proc Nat'l Acad Sci USA 93(3):1156-1160. More recently, ZFNs have been used for genome modification in a variety of organisms. See, for example, United States Patent Publications 20030232410; 20050208489; 20050026157; 20050064474; 20060188987; 20060063231; and International Publication WO 07/014275. Likewise, TALE DNA-binding domains have been fused to nuclease domains to create TALENs. See, e.g., U.S. Publication No. 20110301073.

[0092] As noted above, the cleavage domain may be heterologous to the DNA-binding domain, for example a zinc finger DNA-binding domain and a cleavage domain from a nuclease or a TALEN DNA-binding domain and a cleavage domain, or meganuclease DNA-binding domain and cleavage domain from a different nuclease. Heterologous cleavage domains can be obtained from any endonuclease or exonuclease. Exemplary endonucleases from which a cleavage domain can be derived include, but are not limited to, restriction endonucleases and homing endonucleases. See, for example, 2002-2003 Catalogue, New England Biolabs, Beverly, MA; and Belfort et al. (1997) Nucleic Acids Res. 25:3379-3388. Additional enzymes which cleave DNA are known (e.g., S1 Nuclease; mung bean nuclease; pancreatic DNase I; micrococcal nuclease; yeast HO endonuclease; see also Linn et al. (eds.) Nucleases, Cold Spring Harbor Laboratory Press, 1993). One or more of these enzymes (or functional fragments thereof) can be used as a source of cleavage domains and cleavage half-domains.

[0093] Similarly, a cleavage half-domain can be derived from any nuclease or portion thereof, as set forth above, that requires dimerization for cleavage activity. In general, two fusion proteins are required for cleavage if the fusion proteins comprise cleavage half-domains. Alternatively, a single protein comprising two cleavage half-domains can be used. The two cleavage half-domains can be derived from the same endonuclease (or functional fragments thereof), or each cleavage half-domain can be derived from a different endonuclease (or functional fragments thereof). In addition, the target sites for the two fusion proteins are preferably disposed, with respect to each other, such that binding of the two fusion proteins to their respective target sites places the cleavage half-domains in a spatial orientation to each other that allows the cleavage half-domains to form a functional cleavage domain, e.g., by dimerizing. Thus, in certain embodiments, the near edges of the target sites are separated by 5-8 nucleotides or by 15-18 nucleotides. However any integral number of nucleotides or nucleotide pairs can intervene

between two target sites (e.g., from 2 to 50 nucleotide pairs or more). In general, the site of cleavage lies between the target sites.

[0094] Restriction endonucleases (restriction enzymes) are present in many species and are capable of sequence-specific binding to DNA (at a recognition site), and cleaving DNA at or near the site of binding. Certain restriction enzymes (e.g., Type IIS) cleave DNA at sites removed from the recognition site and have separable binding and cleavage domains. For example, the Type IIS enzyme *Fok I* catalyzes double-stranded cleavage of DNA, at 9 nucleotides from its recognition site on one strand and 13 nucleotides from its recognition site on the other. See, for example, US Patents 5,356,802; 5,436,150 and 5,487,994; as well as Li et al. (1992) Proc. Natl. Acad. Sci. USA 89:4275-4279; Li et al. (1993) Proc. Natl. Acad. Sci. USA 90:2764-2768; Kim et al. (1994a) Proc. Natl. Acad. Sci. USA 91:883-887; Kim et al. (1994b) J. Biol. Chem. 269:31,978-31,982. Thus, in one embodiment, fusion proteins comprise the cleavage domain (or cleavage half-domain) from at least one Type IIS restriction enzyme and one or more zinc finger binding domains, which may or may not be engineered.

[0095] An exemplary Type IIS restriction enzyme, whose cleavage domain is separable from the binding domain, is *Fok I*. This particular enzyme is active as a dimer. Bitinaite et al. (1998) Proc. Natl. Acad. Sci. USA 95: 10,570-10,575. Accordingly, for the purposes of the present disclosure, the portion of the *Fok I* enzyme used in the disclosed fusion proteins is considered a cleavage half-domain. Thus, for targeted double-stranded cleavage and/or targeted replacement of cellular sequences using zinc finger-*Fok I* fusions, two fusion proteins, each comprising a *FokI* cleavage half-domain, can be used to reconstitute a catalytically active cleavage domain. Alternatively, a single polypeptide molecule containing a DNA binding domain and two *Fok I* cleavage half-domains can also be used.

[0096] A cleavage domain or cleavage half-domain can be any portion of a protein that retains cleavage activity, or that retains the ability to multimerize (e.g., dimerize) to form a functional cleavage domain.

[0097] Exemplary Type IIS restriction enzymes are described in International Publication WO 07/014275. Additional restriction enzymes also contain separable binding and cleavage domains, and these are contemplated by the present disclosure. See, for example, Roberts et al. (2003) Nucleic Acids Res. 31:418-420.

[0098] In certain embodiments, the cleavage domain comprises one or more engineered cleavage half-domain (also referred to as dimerization domain mutants) that minimize or prevent homodimerization, as described, for example, in U.S. Patent Publication Nos. 20050064474; 20060188987 and 20080131962. Amino acid residues at positions 446, 447, 479, 483, 484, 486, 487, 490, 491, 496, 498, 499, 500, 531, 534, 537, and 538 of *Fok I* are all targets for influencing dimerization of the *Fok I* cleavage half-domains.

[0099] Exemplary engineered cleavage half-domains of *Fok I* that form obligate heterodimers include a pair in which a first cleavage half-domain includes mutations at amino acid residues at positions 490 and 538 of *Fok I* and a second cleavage half-domain includes mutations at amino acid residues 486 and 499.

[0100] Thus, in one embodiment, a mutation at 490 replaces Glu (E) with Lys (K); the mutation at 538 replaces Iso (I) with Lys (K); the mutation at 486 replaced Gln (Q) with Glu (E); and the mutation at position 499 replaces Iso (I) with Lys (K). Specifically, the engineered cleavage half-domains described herein were prepared by mutating positions 490 (E→K) and 538 (I→K) in one cleavage half-domain to produce an engineered cleavage half-domain designated "E490K:I538K" and by mutating positions 486

(Q→E) and 499 (I→L) in another cleavage half-domain to produce an engineered cleavage half-domain designated "Q486E:I499L". The engineered cleavage half-domains described herein are obligate heterodimer mutants in which aberrant cleavage is minimized or abolished. See, e.g., U.S. Patent Publication No. 2008/0131962.

[0101] In certain embodiments, the engineered cleavage half-domain comprises mutations at positions 486, 499 and 496 (numbered relative to wild-type FokI), for instance mutations that replace the wild type Gln (Q) residue at position 486 with a Glu (E) residue, the wild type Iso (I) residue at position 499 with a Leu (L) residue and the wild-type Asn (N) residue at position 496 with an Asp (D) or Glu (E) residue (also referred to as a "ELD" and "ELE" domains, respectively). In other embodiments, the engineered cleavage half-domain comprises mutations at positions 490, 538 and 537 (numbered relative to wild-type FokI), for instance mutations that replace the wild type Glu (E) residue at position 490 with a Lys (K) residue, the wild type Iso (I) residue at position 538 with a Lys (K) residue, and the wild-type His (H) residue at position 537 with a Lys (K) residue or a Arg (R) residue (also referred to as "KKK" and "KKR" domains, respectively). In other embodiments, the engineered cleavage half-domain comprises mutations at positions 490 and 537 (numbered relative to wild-type FokI), for instance mutations that replace the wild type Glu (E) residue at position 490 with a Lys (K) residue and the wild-type His (H) residue at position 537 with a Lys (K) residue or a Arg (R) residue (also referred to as "KIK" and "KIR" domains, respectively). (See US Patent Publication No. 20110201055). Engineered cleavage half-domains described herein can be prepared using any suitable method, for example, by site-directed mutagenesis of wild-type cleavage half-domains (*Fok I*) as described in U.S. Patent Publication Nos. 20050064474; 20080131962; and 20110201055.

[0102] Alternatively, nucleases may be assembled *in vivo* at the nucleic acid target site using so-called "split-enzyme" technology (see e.g. U.S. Patent Publication No. 20090068164). Components of such split enzymes may be expressed either on separate expression constructs, or can be linked in one open reading frame where the individual components are separated, for example, by a self-cleaving 2A peptide or IRES sequence. Components may be individual zinc finger binding domains or domains of a meganuclease nucleic acid binding domain.

[0103] Nucleases can be screened for activity prior to use, for example in a yeast-based chromosomal system as described in WO 2009/042163 and 20090068164. Nuclease expression constructs can be readily designed using methods known in the art. See, e.g., United States Patent Publications 20030232410; 20050208489; 20050026157; 20050064474; 20060188987; 20060063231; and International Publication WO 07/014275. Expression of the nuclease may be under the control of a constitutive promoter or an inducible promoter, for example the galactokinase promoter which is activated (de-repressed) in the presence of raffinose and/or galactose and repressed in presence of glucose.

[0104] The CRISPR (Clustered Regularly Interspaced Short Palindromic Repeats)/Cas (CRISPR Associated) nuclease system is a recently engineered nuclease system based on a bacterial system that can be used for genome engineering. It is based on part of the adaptive immune response of many bacteria and archaea. When a virus or plasmid invades a bacterium, segments of the invader's DNA are converted into CRISPR RNAs (crRNA) by the 'immune' response. This crRNA then associates, through a region of partial complementarity, with another type of RNA called tracrRNA to guide the Cas9 nuclease to a region homologous to the crRNA in the target DNA called a "protospacer". Cas9 cleaves the DNA to generate blunt ends at the DSB at sites specified by a 20-nucleotide guide sequence contained within the crRNA transcript. Cas9 requires both the crRNA and the tracrRNA for site specific DNA recognition

and cleavage. This system has now been engineered such that the crRNA and tracrRNA can be combined into one molecule (the "single guide RNA"), and the crRNA equivalent portion of the single guide RNA can be engineered to guide the Cas9 nuclease to target any desired sequence (see Jinek et al (2012) Science 337, p. 816-821, Jinek et al, (2013), eLife 2:e00471, and David Segal, (2013) eLife 2:e00563). Thus, the CRISPR/Cas system can be engineered to create a DSB at a desired target in a genome, and repair of the DSB can be influenced by the use of repair inhibitors to cause an increase in error prone repair.

Target Sites

[0105] As described in detail above, DNA domains can be engineered to bind to any sequence of choice in a locus, for example an albumin or other safe-harbor gene. An engineered DNA-binding domain can have a novel binding specificity, compared to a naturally-occurring DNA-binding domain. Engineering methods include, but are not limited to, rational design and various types of selection. Rational design includes, for example, using databases comprising triplet (or quadruplet) nucleotide sequences and individual (e.g., zinc finger) amino acid sequences, in which each triplet or quadruplet nucleotide sequence is associated with one or more amino acid sequences of DNA binding domain which bind the particular triplet or quadruplet sequence. See, for example, co-owned U.S. Patents 6,453,242 and 6,534,261. Rational design of TAL-effector domains can also be performed. See, e.g., U.S. Publication No. 20110301073.

[0106] Exemplary selection methods applicable to DNA-binding domains, including phage display and two-hybrid systems, are disclosed in US Patents 5,789,538; 5,925,523; 6,007,988; 6,013,453; 6,410,248; 6,140,466; 6,200,759; and 6,242,568; as well as WO 98/37186; WO 98/53057; WO 00/27878; WO 01/88197 and GB 2,338,237.

[0107] Selection of target sites; nucleases and methods for design and construction of fusion proteins (and polynucleotides encoding same) are known to those of skill in the art and described in detail in U.S. Patent Application Publication Nos. 20050064474 and 20060188987.

[0108] In addition, as disclosed in these and other references, DNA-binding domains (e.g., multi-fingered zinc finger proteins) may be linked together using any suitable linker sequences, including for example, linkers of 5 or more amino acids. See, e.g., U.S. Patent Nos. 6,479,626; 6,903,185; and 7,153,949 for exemplary linker sequences 6 or more amino acids in length. The proteins described herein may include any combination of suitable linkers between the individual DNA-binding domains of the protein. See, also, U.S. Publication No. 20110301073.

Donors

[0109] As noted above, insertion of an exogenous sequence (also called a "donor sequence" or "donor"), for example for correction of a mutant gene or for increased expression of a wild-type gene. It will be readily apparent that the donor sequence is typically not identical to the genomic sequence where it is placed. A donor sequence can contain a non-homologous sequence flanked by two regions of homology to allow for efficient HDR at the location of interest. Additionally, donor sequences can comprise a vector molecule containing sequences that are not homologous to the region of interest in cellular chromatin. A donor molecule can contain several, discontinuous regions of homology to cellular

chromatin. For example, for targeted insertion of sequences not normally present in a region of interest, said sequences can be present in a donor nucleic acid molecule and flanked by regions of homology to sequence in the region of interest.

[0110] The donor polynucleotide can be DNA or RNA, single-stranded or double-stranded and can be introduced into a cell in linear or circular form. If introduced in linear form, the ends of the donor sequence can be protected (e.g., from exonucleolytic degradation) by methods known to those of skill in the art. For example, one or more dideoxynucleotide residues are added to the 3' terminus of a linear molecule and/or self-complementary oligonucleotides are ligated to one or both ends. See, for example, Chang et al. (1987) Proc. Natl. Acad. Sci. USA 84:4959-4963; Nehls et al. (1996) Science 272:886-889. Additional methods for protecting exogenous polynucleotides from degradation include, but are not limited to, addition of terminal amino group(s) and the use of modified internucleotide linkages such as, for example, phosphorothioates, phosphoramidates, and O-methyl ribose or deoxyribose residues.

[0111] A polynucleotide can be introduced into a cell as part of a vector molecule having additional sequences such as, for example, replication origins, promoters and genes encoding antibiotic resistance. Moreover, donor polynucleotides can be introduced as naked nucleic acid, as nucleic acid complexed with an agent such as a liposome or poloxamer, or can be delivered by viruses (e.g., adenovirus, AAV, herpesvirus, retrovirus, lentivirus and integrase defective lentivirus (IDLV)).

[0112] The donor is generally inserted so that its expression is driven by the endogenous promoter at the integration site, namely the promoter that drives expression of the endogenous gene into which the donor is inserted (e.g., highly expressed, albumin, AAVS1, HPRT, etc.). However, it will be apparent that the donor may comprise a promoter and/or enhancer, for example a constitutive promoter or an inducible or tissue specific promoter.

[0113] The donor molecule may be inserted into an endogenous gene such that all, some or none of the endogenous gene is expressed. For example, a transgene as described herein may be inserted into an albumin or other locus such that some (N-terminal and/or C-terminal to the transgene encoding the lysosomal enzyme) or none of the endogenous albumin sequences are expressed, for example as a fusion with the transgene encoding the lysosomal sequences. The transgene (e.g., with or without additional coding sequences such as for albumin) may be integrated into any endogenous locus, for example a safe-harbor locus. See, e.g., U.S. patent publications 20080299580; 20080159996 and 201000218264.

[0114] When endogenous sequences (endogenous or part of the transgene) are expressed with the transgene, the endogenous sequences (e.g., albumin, etc.) may be full-length sequences (wild-type or mutant) or partial sequences. Preferably the endogenous sequences are functional. Non-limiting examples of the function of these full length or partial sequences (e.g., albumin) include increasing the serum half-life of the polypeptide expressed by the transgene (e.g., therapeutic gene) and/or acting as a carrier.

[0115] Furthermore, although not required for expression, exogenous sequences may also include transcriptional or translational regulatory sequences, for example, promoters, enhancers, insulators, internal ribosome entry sites, sequences encoding 2A peptides and/or polyadenylation signals.

[0116] In certain embodiments, the exogenous sequence (donor) comprises a fusion of a protein of interest and, as its fusion partner, an extracellular domain of a membrane protein, causing the fusion

protein to be located on the surface of the cell. This allows the protein encoded by the transgene to potentially act in the serum. In the case of treatment for an LSD, the enzyme encoded by the transgene fusion would be able to act on the metabolic products that are accumulating in the serum from its location on the surface of the cell (e.g., RBC). In addition, if the RBC is engulfed by a splenic macrophage as is the normal course of degradation, the lysosome formed when the macrophage engulfs the cell would expose the membrane bound fusion protein to the high concentrations of metabolic products in the lysosome at the pH more naturally favorable to that enzyme. Non-limiting examples of potential fusion partners are shown below in Table 1.

Table 1: Examples of potential fusion partners

Name	Activity
Band 3	Anion transporter, makes up to 25% of the RBC membrane surface protein
Aquaporin 1	water transporter
Glut1	glucose and L-dehydroascorbic acid transporter
Kidd antigen protein	urea transporter
RhAG	gas transporter
ATP1A1, ATP1B1	Na ⁺ /K ⁺ - ATPase
ATP2B1, ATP2B2, ATP2B3, ATP2B4	Ca ²⁺ - ATPase
NKCC1, NKCC2	Na ⁺ K ⁺ 2Cl ⁻ - cotransporter
SLC12A3	Na ⁺ -Cl ⁻ cotransporter
SLC12A1, SLA12A2	Na-K - cotransporter
KCC1	K-Cl cotransporter
KCNN4	Gardos Channel

[0117] Lysosomal storage diseases typically fall into five classes. These classes are shown below in Table 2 along with specific examples of the diseases. Thus, the donor molecules described herein can include sequences coding for one or more enzymes lacking or deficient in subjects with lysosomal storage diseases, including but not limited to the proteins shown in Table 2.

Table 2: Lysosomal Storage Diseases

1. DEFECTS IN GLYCAN DEGRADATION			
Protein	Disease	Disease Associated Gene	Accumulated product
I. Defects in glycoprotein degradation			
α-Sialidase (neuraminidase)	Sialidosis	NEU1	sialidated glycopeptides and oligosaccharides
Cathepsin A	Galactosialidosis	CTSA	polysaccharide
lysosomal alpha-mannosidase	α-Mannosidosis	MAN2B1	mannose-rich glycoproteins and oligosaccharides
lysosomal beta-mannosidase	β-Mannosidosis	MANBA	

1. DEFECTS IN GLYCAN DEGRADATION			
Protein	Disease	Disease Associated Gene	Accumulated product
I. Defects in glycoprotein degradation			
Glycosylasparaginase	Aspartylglucosaminuria	<i>AGA</i>	glycoasparagines
Alpha L Fucosidase	Fucosidosis	<i>FUCA1</i>	fucose
α -N-Acetylglucosaminidase	Sanfilippo syndrome B	<i>NAGLU</i>	glycosaminoglycan
II. Defects in glycolipid degradation			
A. GM1 Ganglioside			
β -Galactosidase	GM1 gangliosidosis / MPS IVB	<i>GLB1</i>	keratan sulfate
β -Hexosaminidase α -subunit	GM2-gangliosidosis (Tay-Sachs)	<i>HEXA</i>	GM2 ganglioside
β -Hexosaminidase β -subunit	GM2-gangliosidosis (Sandhoff)	<i>HEXB</i>	GM2 ganglioside
GM2 activator protein	GM2 gangliosidosis	<i>GM2A</i>	GM2 ganglioside
Glucocerebrosidase	Gaucher disease	<i>GBA</i>	glucocerebroside
Saposin C	Gaucher disease (atypical)	<i>PSAP</i>	glucocerebroside
B. Defects in the degradation of sulfatide			
Arylsulfatase A	Metachromatic leukodystrophy	<i>ARSA</i>	sulphatide
Saposin B	Metachromatic leukodystrophy	<i>PSAP</i>	sulphatide
Formyl-Glycin generating enzyme	Multiple sulfatase deficiency	<i>SUMF1</i>	sulfated lipids
β -Galactosylceramidase (Krabbe)	Globoid cell leukodystrophy	<i>GALC</i>	galactocerebroside
C. Defects in degradation of globotriaosylceramide			
α -Galactosidase A	Fabry	<i>GLA</i>	globotriaosylcera-mide
III. Defects in degradation of Glycosaminoglycan (Mucopolysaccharidoses)			
A. Degradation of heparan sulphate			
Iduronate sulfatase	MPS II (Hunter)	<i>IDS</i>	Dermatan sulfate, Heparan sulfate
Iduronidase	MPS 1 (Hurler, Scheie)	<i>IDUA</i>	Dermatan sulfate, Heparan sulfate
Heparan N-sulfatase	MPS IIIa (Sanfilippo A)	<i>SGSH</i>	Heparan sulfate
Acetyl-CoA transferase	MPS IIIc (Sanfilippo C)	<i>HGSNAT</i>	Heparan sulfate
N-acetyl glucosaminidase	MPS IIIb (Sanfilippo	<i>NAGLU</i>	Heparan sulfate

III. Defects in degradation of Glycosaminoglycan (Mucopolysaccharidoses)			
	B)		
β -glucuronidase	MPS VII (Sly)	<i>GUSB</i>	
N-acetyl glucosamine 6-sulfatase	MPS IIId (Sanfilippo D)	<i>GNS</i>	Heparan sulfate
B. Degradation of other mucopolysaccharides			
B-Galactosidase	MPS VIB (Morquio B)	<i>GLB1</i>	Keratan sulfate,
Galactose 6-sulfatase	MPS IVA (Morquio A)	<i>GALNS</i>	Keratan sulfate, Chondroitin 6-sulfate
Hyaluronidase	MPS IX	<i>HYAL1</i>	Hyaluronic acid
C. Defects in degradation of glycogen			
α -Glucosidase	Pompe	<i>GAA</i>	Glycogen
2. DEFECTS IN LIPID DEGRADATION			
I. Defects in degradation of sphingomyelin			
Acid sphingomyelinase	Niemann Pick type A	<i>SMPD1</i>	sphingomyelin
Acid ceramidase	Farber lipogranulomatosis	<i>ASAH1</i>	nonsulfonated acid mucopolysaccharide
II. Defects in degradation of triglycerides and cholesteryl ester			
Acid lipase	Wolman and cholesteryl ester storage disease	<i>LIPA</i>	cholesteryl esters
3. DEFECTS IN PROTEIN DEGRADATION			
Cathepsin K	Pycnodystostosis	<i>CTSK</i>	
Tripeptidyl peptidase	Ceroide lipofuscinosis	<i>PPT2</i>	
Palmitoyl-protein thioesterase	Ceroide lipofuscinosis	<i>PPT1</i>	
4. DEFECTS IN LYSOSOMAL TRANSPORTERS			
Cystinosin (cystin transport)	Cystinosis	<i>CTNS</i>	
Sialin (sialic acid transport)	Salla disease	<i>SLC17A5</i>	N-acetylneuraminic acid
5. DEFECTS IN LYSOSOMAL TRAFFICKING PROTEINS			
Phosphotransferase γ -subunit	Mucopolidosis III (I-cell)	<i>GNPTG</i>	
Mucolipin-1(cation channel)	Mucopolidosis	<i>MCOLN1</i>	
LYSOSOME-ASSOCIATED MEMBRANE PROTEIN 2	Danon	<i>LAMP2</i>	
Niemann-Pick disease, type C1	Niemann Pick type C	<i>NPC1</i>	LDL cholesterol
palmitoyl-protein thioesterase-1	Ceroid lipofuscinosis	<i>CLN3</i>	autofluorescent lipopigment storage

5. DEFECTS IN LYSOSOMAL TRAFFICKING PROTEINS			
	(Batten Disease)		material
neuronal ceroid lipofuscinosis-6	Ceroid lipofuscinosis 6	<i>CLN 6</i>	
neuronal ceroid lipofuscinosis-8	Ceroid lipofuscinosis 8	<i>CLN 8</i>	
LYSOSOMAL TRAFFICKING REGULATOR	Chediak-Higashi	<i>LYST</i>	
Myocilin	Griscelli Type 1	<i>MYOC</i>	
RAS-associated protein 27A	Griscelli Type 2	<i>RAB27A</i>	
Melanophilin	Griscelli Type 3	<i>MLPH</i> or <i>MYO5A</i>	
AP3 β -subunit	Hermansky Pudliak	<i>AP3B1</i>	ceroid

[0118] In some cases, the donor may be an endogenous gene that has been modified. Although antibody response to enzyme replacement therapy varies with respect to the specific therapeutic enzyme in question and with the individual patient, a significant immune response has been seen in many LSD patients being treated with enzyme replacement. In addition, the relevance of these antibodies to the efficacy of treatment is also variable (see Katherine Ponder, (2008) J Clin Invest 118(8):2686). Thus, the methods and compositions described herein can comprise the use of donor molecules whose sequence has been altered by functionally silent amino acid changes at sites known to be priming epitopes for endogenous immune responses, such that the polypeptide produced by such a donor is less immunogenic.

[0119] LSD patients often have neurological sequelae due the lack of the missing enzyme in the brain. Unfortunately, it is often difficult to deliver therapeutics to the brain via the blood due to the impermeability of the blood brain barrier. Thus, the methods and compositions described herein may be used in conjunction with methods to increase the delivery of the therapeutic into the brain. There are some methods that cause a transient opening of the tight junctions between cells of the brain capillaries. Examples include transient osmotic disruption through the use of an intracarotid administration of a hypertonic mannitol solution, the use of focused ultrasound and the administration of a bradykinin analogue. Alternatively, therapeutics can be designed to utilize receptors or transport mechanisms for specific transport into the brain. Examples of specific receptors that may be used include the transferrin receptor, the insulin receptor or the low-density lipoprotein receptor related proteins 1 and 2 (LRP-1 and LRP-2). LRP is known to interact with a range of secreted proteins such as apoE, tPA, PAI-1 etc, and so fusing a recognition sequence from one of these proteins for LRP may facilitate transport of the enzyme into the brain, following expression in the liver of the therapeutic protein and secretion into the blood stream (see Gabathuler, (2010) *ibid*).

Delivery

[0120] The nucleases, polynucleotides encoding these nucleases, donor polynucleotides and compositions comprising the proteins and/or polynucleotides described herein may be delivered *in vivo* or *ex vivo* by any suitable means.

[0121] Methods of delivering nucleases as described herein are described, for example, in U.S. Patent Nos. 6,453,242; 6,503,717; 6,534,261; 6,599,692; 6,607,882; 6,689,558; 6,824,978; 6,933,113; 6,979,539; 7,013,219; and 7,163,824.

[0122] Nucleases and/or donor constructs as described herein may also be delivered using vectors containing sequences encoding one or more of the zinc finger or TALEN protein(s). Any vector systems may be used including, but not limited to, plasmid vectors, retroviral vectors, lentiviral vectors, adenovirus vectors, poxvirus vectors; herpesvirus vectors and adeno-associated virus vectors, etc. See, *also*, U.S. Patent Nos. 6,534,261; 6,607,882; 6,824,978; 6,933,113; 6,979,539; 7,013,219; and 7,163,824. Furthermore, it will be apparent that any of these vectors may comprise one or more of the sequences needed for treatment. Thus, when one or more nucleases and a donor construct are introduced into the cell, the nucleases and/or donor polynucleotide may be carried on the same vector or on different vectors. When multiple vectors are used, each vector may comprise a sequence encoding one or multiple nucleases and/or donor constructs.

[0123] Conventional viral and non-viral based gene transfer methods can be used to introduce nucleic acids encoding nucleases and donor constructs in cells (*e.g.*, mammalian cells) and target tissues. Non-viral vector delivery systems include DNA plasmids, naked nucleic acid, and nucleic acid complexed with a delivery vehicle such as a liposome or poloxamer. Viral vector delivery systems include DNA and RNA viruses, which have either episomal or integrated genomes after delivery to the cell. For a review of gene therapy procedures, see Anderson, *Science* 256:808-813 (1992); Nabel & Felgner, *TIBTECH* 11:211-217 (1993); Mitani & Caskey, *TIBTECH* 11:162-166 (1993); Dillon, *TIBTECH* 11:167-175 (1993); Miller, *Nature* 357:455-460 (1992); Van Brunt, *Biotechnology* 6(10):1149-1154 (1988); Vigne, *Restorative Neurology and Neuroscience* 8:35-36 (1995); Kremer & Perricaudet, *British Medical Bulletin* 51(1):31-44 (1995); Haddada et al., in *Current Topics in Microbiology and Immunology* Doerfler and Böhm (eds.) (1995); and Yu et al., *Gene Therapy* 1:13-26 (1994).

[0124] Methods of non-viral delivery of nucleic acids include electroporation, lipofection, microinjection, biolistics, virosomes, liposomes, immunoliposomes, polycation or lipid:nucleic acid conjugates, naked DNA, artificial virions, and agent-enhanced uptake of DNA. Sonoporation using, *e.g.*, the Sonitron 2000 system (Rich-Mar) can also be used for delivery of nucleic acids.

[0125] Additional exemplary nucleic acid delivery systems include those provided by Amaxa Biosystems (Cologne, Germany), Maxcyte, Inc. (Rockville, Maryland), BTX Molecular Delivery Systems (Holliston, MA) and Copernicus Therapeutics Inc, (see for example US6008336). Lipofection is described in *e.g.*, U.S. Patent Nos. 5,049,386; 4,946,787; and 4,897,355) and lipofection reagents are sold commercially (*e.g.*, Transfectam™ and Lipofectin™). Cationic and neutral lipids that are suitable for efficient receptor-recognition lipofection of polynucleotides include those of Felgner, WO 91/17424, WO 91/16024.

[0126] The preparation of lipid:nucleic acid complexes, including targeted liposomes such as immunolipid complexes, is well known to one of skill in the art (see, *e.g.*, Crystal, *Science* 270:404-410 (1995); Blaese et al., *Cancer Gene Ther.* 2:291-297 (1995); Behr et al., *Bioconjugate Chem.* 5:382-389 (1994); Remy et al., *Bioconjugate Chem.* 5:647-654 (1994); Gao et al., *Gene Therapy* 2:710-722 (1995); Ahmad et al., *Cancer Res.* 52:4817-4820 (1992); U.S. Pat. Nos. 4,186,183, 4,217,344, 4,235,871, 4,261,975, 4,485,054, 4,501,728, 4,774,085, 4,837,028, and 4,946,787).

[0127] Additional methods of delivery include the use of packaging the nucleic acids to be delivered into EnGeneIC delivery vehicles (EDVs). These EDVs are specifically delivered to target tissues using

bispecific antibodies where one arm of the antibody has specificity for the target tissue and the other has specificity for the EDV. The antibody brings the EDVs to the target cell surface and then the EDV is brought into the cell by endocytosis. Once in the cell, the contents are released (see MacDiarmid et al (2009) Nature Biotechnology 27(7):643).

[0128] The use of RNA or DNA viral based systems for the delivery of nucleic acids encoding engineered ZFPs take advantage of highly evolved processes for targeting a virus to specific cells in the body and trafficking the viral payload to the nucleus. Viral vectors can be administered directly to subjects (*in vivo*) or they can be used to treat cells *in vitro* and the modified cells are administered to subjects (*ex vivo*). Conventional viral based systems for the delivery of ZFPs include, but are not limited to, retroviral, lentivirus, adenoviral, adeno-associated, vaccinia and herpes simplex virus vectors for gene transfer. Integration in the host genome is possible with the retrovirus, lentivirus, and adeno-associated virus gene transfer methods, often resulting in long term expression of the inserted transgene. Additionally, high transduction efficiencies have been observed in many different cell types and target tissues.

[0129] The tropism of a retrovirus can be altered by incorporating foreign envelope proteins, expanding the potential target population of target cells. Lentiviral vectors are retroviral vectors that are able to transduce or infect non-dividing cells and typically produce high viral titers. Selection of a retroviral gene transfer system depends on the target tissue. Retroviral vectors are comprised of *cis*-acting long terminal repeats with packaging capacity for up to 6-10 kb of foreign sequence. The minimum *cis*-acting LTRs are sufficient for replication and packaging of the vectors, which are then used to integrate the therapeutic gene into the target cell to provide permanent transgene expression. Widely used retroviral vectors include those based upon murine leukemia virus (MuLV), gibbon ape leukemia virus (GaLV), Simian Immunodeficiency virus (SIV), human immunodeficiency virus (HIV), and combinations thereof (see, e.g., Buchscher et al., J. Virol. 66:2731-2739 (1992); Johann et al., J. Virol. 66:1635-1640 (1992); Sommerfelt et al., Virol. 176:58-59 (1990); Wilson et al., J. Virol. 63:2374-2378 (1989); Miller et al., J. Virol. 65:2220-2224 (1991); PCT/US94/05700).

[0130] In applications in which transient expression is preferred, adenoviral based systems can be used. Adenoviral based vectors are capable of very high transduction efficiency in many cell types and do not require cell division. With such vectors, high titer and high levels of expression have been obtained. This vector can be produced in large quantities in a relatively simple system. Adeno-associated virus ("AAV") vectors are also used to transduce cells with target nucleic acids, e.g., in the *in vitro* production of nucleic acids and peptides, and for *in vivo* and *ex vivo* gene therapy procedures (see, e.g., West et al., Virology 160:38-47 (1987); U.S. Patent No. 4,797,368; WO 93/24641; Kotin, Human Gene Therapy 5:793-801 (1994); Muzyczka, J. Clin. Invest. 94:1351 (1994). Construction of recombinant AAV vectors are described in a number of publications, including U.S. Pat. No. 5,173,414; Tratschin et al., Mol. Cell. Biol. 5:3251-3260 (1985); Tratschin, et al., Mol. Cell. Biol. 4:2072-2081 (1984); Hermonat & Muzyczka, PNAS 81:6466-6470 (1984); and Samulski et al., J. Virol. 63:03822-3828 (1989).

[0131] At least six viral vector approaches are currently available for gene transfer in clinical trials, which utilize approaches that involve complementation of defective vectors by genes inserted into helper cell lines to generate the transducing agent.

[0132] pLASN and MFG-S are examples of retroviral vectors that have been used in clinical trials (Dunbar et al., Blood 85:3048-305 (1995); Kohn et al., Nat. Med. 1:1017-102 (1995); Malech et al., PNAS 94:22 12133-12138 (1997)). PA317/pLASN was the first therapeutic vector used in a gene

therapy trial. (Blaese et al., Science 270:475-480 (1995)). Transduction efficiencies of 50% or greater have been observed for MFG-S packaged vectors. (Ellem et al., Immunol Immunother. 44(1):10-20 (1997); Dranoff et al., Hum. Gene Ther. 1:111-2 (1997)).

[0133] Recombinant adeno-associated virus vectors (rAAV) are a promising alternative gene delivery systems based on the defective and nonpathogenic parvovirus adeno-associated type 2 virus. All vectors are derived from a plasmid that retains only the AAV 145 bp inverted terminal repeats flanking the transgene expression cassette. Efficient gene transfer and stable transgene delivery due to integration into the genomes of the transduced cell are key features for this vector system. (Wagner et al., Lancet 351:9117 1702-3 (1998), Kearns et al., Gene Ther. 9:748-55 (1996)). Other AAV serotypes, including by non-limiting example, AAV1, AAV3, AAV4, AAV5, AAV6, AAV8, AAV 8.2, AAV9, and AAV rh10 and pseudotyped AAV such as AAV2/8, AAV2/5 and AAV2/6 can also be used in accordance with the present invention.

[0134] Replication-deficient recombinant adenoviral vectors (Ad) can be produced at high titer and readily infect a number of different cell types. Most adenovirus vectors are engineered such that a transgene replaces the Ad E1a, E1b, and/or E3 genes; subsequently the replication defective vector is propagated in human 293 cells that supply deleted gene function in *trans*. Ad vectors can transduce multiple types of tissues *in vivo*, including non-dividing, differentiated cells such as those found in liver, kidney and muscle. Conventional Ad vectors have a large carrying capacity. An example of the use of an Ad vector in a clinical trial involved polynucleotide therapy for anti-tumor immunization with intramuscular injection (Sternan et al., Hum. Gene Ther. 7:1083-9 (1998)). Additional examples of the use of adenovirus vectors for gene transfer in clinical trials include Rosenecker et al., Infection 24:1 5-10 (1996); Sternan et al., Hum. Gene Ther. 9:7 1083-1089 (1998); Welsh et al., Hum. Gene Ther. 2:205-18 (1995); Alvarez et al., Hum. Gene Ther. 5:597-613 (1997); Topf et al., Gene Ther. 5:507-513 (1998); Sternan et al., Hum. Gene Ther. 7:1083-1089 (1998).

[0135] Packaging cells are used to form virus particles that are capable of infecting a host cell. Such cells include 293 cells, which package adenovirus, and ψ 2 cells or PA317 cells, which package retrovirus. Viral vectors used in gene therapy are usually generated by a producer cell line that packages a nucleic acid vector into a viral particle. The vectors typically contain the minimal viral sequences required for packaging and subsequent integration into a host (if applicable), other viral sequences being replaced by an expression cassette encoding the protein to be expressed. The missing viral functions are supplied in *trans* by the packaging cell line. For example, AAV vectors used in gene therapy typically only possess inverted terminal repeat (ITR) sequences from the AAV genome which are required for packaging and integration into the host genome. Viral DNA is packaged in a cell line, which contains a helper plasmid encoding the other AAV genes, namely *rep* and *cap*, but lacking ITR sequences. The cell line is also infected with adenovirus as a helper. The helper virus promotes replication of the AAV vector and expression of AAV genes from the helper plasmid. The helper plasmid is not packaged in significant amounts due to a lack of ITR sequences. Contamination with adenovirus can be reduced by, e.g., heat treatment to which adenovirus is more sensitive than AAV.

[0136] In many gene therapy applications, it is desirable that the gene therapy vector be delivered with a high degree of specificity to a particular tissue type. Accordingly, a viral vector can be modified to have specificity for a given cell type by expressing a ligand as a fusion protein with a viral coat protein on the outer surface of the virus. The ligand is chosen to have affinity for a receptor known to be present on the cell type of interest. For example, Han et al., Proc. Natl. Acad. Sci. USA 92:9747-9751 (1995), reported that Moloney murine leukemia virus can be modified to express human heregulin fused to gp70, and the

recombinant virus infects certain human breast cancer cells expressing human epidermal growth factor receptor. This principle can be extended to other virus-target cell pairs, in which the target cell expresses a receptor and the virus expresses a fusion protein comprising a ligand for the cell-surface receptor. For example, filamentous phage can be engineered to display antibody fragments (e.g., FAb or Fv) having specific binding affinity for virtually any chosen cellular receptor. Although the above description applies primarily to viral vectors, the same principles can be applied to nonviral vectors. Such vectors can be engineered to contain specific uptake sequences which favor uptake by specific target cells.

[0137] Gene therapy vectors can be delivered *in vivo* by administration to an individual patient, typically by systemic administration (e.g., intravenous, intraperitoneal, intramuscular, subdermal, or intracranial infusion) or topical application, as described below. Alternatively, vectors can be delivered to cells *ex vivo*, such as cells explanted from an individual patient (e.g., lymphocytes, bone marrow aspirates, tissue biopsy) or universal donor hematopoietic stem cells, followed by reimplantation of the cells into a patient, usually after selection for cells which have incorporated the vector.

[0138] Vectors (e.g., retroviruses, adenoviruses, liposomes, etc.) containing nucleases and/or donor constructs can also be administered directly to an organism for transduction of cells *in vivo*. Alternatively, naked DNA can be administered. Administration is by any of the routes normally used for introducing a molecule into ultimate contact with blood or tissue cells including, but not limited to, injection, infusion, topical application and electroporation. Suitable methods of administering such nucleic acids are available and well known to those of skill in the art, and, although more than one route can be used to administer a particular composition, a particular route can often provide a more immediate and more effective reaction than another route.

[0139] Vectors suitable for introduction of polynucleotides described herein include non-integrating lentivirus vectors (IDLV). See, for example, Ory et al. (1996) Proc. Natl. Acad. Sci. USA 93:11382-11388; Dull et al. (1998) J. Virol. 72:8463-8471; Zuffery et al. (1998) J. Virol. 72:9873-9880; Follenzi et al. (2000) Nature Genetics 25:217-222; U.S. Patent Publication No 2009/054985.

[0140] Pharmaceutically acceptable carriers are determined in part by the particular composition being administered, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of pharmaceutical compositions available, as described below (see, e.g., Remington's Pharmaceutical Sciences, 17th ed., 1989).

[0141] It will be apparent that the nuclease-encoding sequences and donor constructs can be delivered using the same or different systems. For example, a donor polynucleotide can be carried by a plasmid, while the one or more nucleases can be carried by a AAV vector. Furthermore, the different vectors can be administered by the same or different routes (intramuscular injection, tail vein injection, other intravenous injection, intraperitoneal administration and/or intramuscular injection). The vectors can be delivered simultaneously or in any sequential order.

[0142] Formulations for both *ex vivo* and *in vivo* administrations include suspensions in liquid or emulsified liquids. The active ingredients often are mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients include, for example, water, saline, dextrose, glycerol, ethanol or the like, and combinations thereof. In addition, the composition may contain minor amounts of auxiliary substances, such as, wetting or emulsifying agents, pH buffering agents, stabilizing agents or other reagents that enhance the effectiveness of the pharmaceutical

composition.

Applications

[0143] The methods described herein contemplate the treatment of a monogenic disease (e.g. lysosomal storage disease). Treatment can comprise insertion of the corrected disease associated gene in safe harbor locus (e.g. albumin) for expression of the needed enzyme and release into the blood stream. Insertion into a secretory cell, such as a liver cell for release of the product into the blood stream, is particularly useful. The methods and compositions of the described herein also can be used in any circumstance wherein it is desired to supply a transgene encoding one or more therapeutics in a hemapoietic stem cell such that mature cells (e.g., RBCs) derived from these cells contain the therapeutic. These stem cells can be differentiated *in vitro* or *in vivo* and may be derived from a universal donor type of cell which can be used for all patients. Additionally, the cells may contain a transmembrane protein to traffic the cells in the body. Treatment can also comprise use of patient cells containing the therapeutic transgene where the cells are developed *ex vivo* and then introduced back into the patient. For example, HSC containing a suitable transgene may be inserted into a patient via a bone marrow transplant. Alternatively, stem cells such as muscle stem cells or iPSC which have been edited using with the therapeutic transgene maybe also injected into muscle tissue.

[0144] Thus, this technology may be of use in a condition where a patient is deficient in some protein due to problems (e.g., problems in expression level or problems with the protein expressed as sub- or non-functioning). Particularly useful with this invention is the expression of transgenes to correct or restore functionality in lysosomal storage disorders.

[0145] By way of non-limiting examples, production of the defective or missing proteins accomplished and used to treat the lysosomal storage disease. Nucleic acid donors encoding the proteins may be inserted into a safe harbor locus (e.g. albumin or HPRT) and expressed either using an exogenous promoter or using the promoter present at the safe harbor. Alternatively, donors can be used to correct the defective gene *in situ*. The desired transgene may be inserted into a CD34+ stem cell and returned to a patient during a bone marrow transplant. Finally, the nucleic acid donor maybe be inserted into a CD34+ stem cell at a beta globin locus such that the mature red blood cell derived from this cell has a high concentration of the biologic encoded by the nucleic acid donor. The biologic containing RBC can then be targeted to the correct tissue via transmembrane proteins (e.g. receptor or antibody). Additionally, the RBCs may be sensitized *ex vivo* via electrosensitization to make them more susceptible to disruption following exposure to an energy source (see WO2002007752).

[0146] In some applications, an endogenous gene may be knocked out by use of the methods and compositions described herein. Examples of this aspect include knocking out an aberrant gene regulator or an aberrant disease associated gene. In some applications, an aberrant endogenous gene may be replaced, either functionally or *in situ*, with a wild type version of the gene. The inserted gene may also be altered to improve the functionality of the expressed protein or to reduce its immunogenicity. In some applications, the inserted gene is a fusion protein to increase its transport into a selected tissue such as the brain.

[0147] The following Examples relate to exemplary embodiments of the present disclosure in which the nuclease comprises a zinc finger nuclease (ZFN) or TALEN. It will be appreciated that this is for purposes of exemplification only and that other nucleases or nuclease systems can be used, for

instance homing endonucleases (meganucleases) with engineered DNA-binding domains and/or fusions of naturally occurring of engineered homing endonucleases (meganucleases) DNA-binding domains and heterologous cleavage domains and/or a CRISPR/Cas system comprising an engineered single guide RNA.

EXAMPLES

Example 1: Design, construction and general characterization of albumin-specific nucleases

[0148] Zinc finger proteins were designed and incorporated into plasmids, AAV or adenoviral vectors essentially as described in Urnov et al. (2005) Nature 435(7042):646-651, Perez et al (2008) Nature Biotechnology 26(7):808-816, and as described in U.S. Patent No. 6,534,261. Table 3 shows the recognition helices within the DNA binding domain of exemplary albumin-specific ZFPs while Table 4 shows the target sites for these ZFPs (see co-owned US Provisionals 61/537,349 and 61/560,506). Nucleotides in the target site that are contacted by the ZFP recognition helices are indicated in uppercase letters; non-contacted nucleotides indicated in lowercase. Albumin-specific TALENs were also designed and are set forth in U.S. Application Nos. 13/624,193 and 13/624,217 and incorporated by reference in their entireties).

Table 3: Murine albumin-specific zinc finger nucleases helix designs

Target	SBS #	Design					
		F1	F2	F3	F4	F5	F6
Intron 1	30724	TSGSLTR (SEQ ID NO:1)	RSDALST (SEQ ID NO:2)	QSATRTK (SEQ ID NO:3)	TSGHLSR (SEQ ID NO:4)	QSGNLAR (SEQ ID NO:5)	NA
Intron 1	30725	RSDHLSA (SEQ ID NO:6)	TKSNRTK (SEQ ID NO:7)	DRSNLSR (SEQ ID NO:8)	WRSSLRA (SEQ ID NO:9)	DSSDRKK (SEQ ID NO:10)	NA
Intron 1	30732	TSGNLTR (SEQ ID NO:11)	DRSTRRQ (SEQ ID NO:12)	TSGSLTR (SEQ ID NO:1)	ERGTLAR (SEQ ID NO:13)	TSANLSR (SEQ ID NO:14)	NA
Intron 1	30733	DRSALAR (SEQ ID NO:15)	RSDHLSE (SEQ ID NO:16)	HRSDRTR (SEQ ID NO:17)	QSGALAR (SEQ ID NO:18)	QSGHLSR (SEQ ID NO:19)	NS
Intron 13	30759	RSDNLST (SEQ ID NO:20)	DRSALAR (SEQ ID NO:15)	DRSNLSR (SEQ ID NO:8)	DGRNLRH (SEQ ID NO:21)	RSDNLAR (SEQ ID NO:22)	QSNALNR (SEQ ID NO:23)
Intron 13	30761	DRSNLSR (SEQ ID NO:8)	LKQVLVR (SEQ ID NO:24)	QSGNLAR (SEQ ID NO:5)	QSTPLFA (SEQ ID NO:25)	QSGALAR (SEQ ID NO:18)	NA
Intron 13	30760	DRSNLSR (SEQ ID NO:8)	DGRNLRH (SEQ ID NO:21)	RSDNLAR (SEQ ID NO:22)	QSNALNR (SEQ ID NO:23)	NA	NA
Intron 13	30767	RSDNLSV (SEQ ID NO:26)	HSNARKT (SEQ ID NO:27)	RSDSLSA (SEQ ID NO:28)	QSGNLAR (SEQ ID NO:5)	RSDSLSV (SEQ ID NO:29)	QSGHLSR (SEQ ID NO:19)

Target	SBS #	Design					
		F1	F2	F3	F4	F5	F6
Intron 13	30768	RSDNLSE (SEQ ID NO:30)	ERANRNS (SEQ ID NO:31)	QSANRTK (SEQ ID NO:32)	ERGTLAR (SEQ ID NO:13)	RSDALTQ (SEQ ID NO:33)	NA
Intron 13	30769	TSGSLTR (SEQ ID NO:1)	DRSNLSR (SEQ ID NO:8)	DGRNLRH (SEQ ID NO:21)	ERGTLAR (SEQ ID NO:13)	RSDALTQ (SEQ ID NO:33)	NA
Intron 12	30872	QSGHLAR (SEQ ID NO:34)	RSDHLTQ (SEQ ID NO:35)	RSDHLSQ (SEQ ID NO:36)	WRSSLVA (SEQ ID NO:37)	RSDVLSE (SEQ ID NO:38)	RNQHRKT (SEQ ID NO:39)
Intron 12	30873	QSGDLTR (SEQ ID NO:40)	RSDALAR (SEQ ID NO:41)	QSGDLTR (SEQ ID NO:40)	RRDPLIN (SEQ ID NO:42)	RSDNLSV (SEQ ID NO:26)	IRSTLRD (SEQ ID NO:43)
Intron 12	30876	RSDNLSV (SEQ ID NO:26)	YSSTRNS (SEQ ID NO:44)	RSDHLSA (SEQ ID NO:6)	SYWSRTV (SEQ ID NO:45)	QSSDLSR (SEQ ID NO:46)	RTDALRG (SEQ ID NO:47)
Intron 12	30877	RSDNLST (SEQ ID NO:20)	QKSPLNT (SEQ ID NO:48)	TSGNLTR (SEQ ID NO:11)	QAENLKS (SEQ ID NO:49)	QSSDLSR (SEQ ID NO:46)	RTDALRG (SEQ ID NO:47)
Intron 12	30882	RSDNLSV (SEQ ID NO:26)	RRAHLNQ (SEQ ID NO:50)	TSGNLTR (SEQ ID NO:11)	SDTNRFK (SEQ ID NO:51)	RSDNLST (SEQ ID NO:20)	QSGHLSR (SEQ ID NO:19)
Intron 12	30883	DSSDRKK (SEQ ID NO:10)	DRSALS (SEQ ID NO:52)	TSSNRKT (SEQ ID NO:53)	QSGALAR (SEQ ID NO:18)	RSDHLSR (SEQ ID NO:54)	NA

Table 4: Target sites of murine albumin-specific ZFNs

Target	SBS #	Target site
Intron 1	30724	ctGAAGGTgGCAATGGTTcctctctgct_ (SEQ ID NO:55)
Intron 1	30725	ttTCCTGTAACGATCGGgaactggcatc_ (SEQ ID NO:56)
Intron 1	30732	aaGATGCCaGTTCCCGATcggttacagga_ (SEQ ID NO:57)
Intron 1	30733	agGGAGTAGCTTAGGTCagtgaagagaa_ (SEQ ID NO:58)
Intron 13	30759	acGTAGAGAACAACATCTAGattggtgg_ (SEQ ID NO:59)
Intron 13	30761	ctGTAATAGAACTGACttacgtagctt_ (SEQ ID NO:60)
Intron 13	30760	acGTAGAGAACAACatctagattggtgg_ (SEQ ID NO:59)
Intron 13	30767	agGGAATGtGAAATGATTCAGatata_ (SEQ ID NO:61)
Intron 13	30768	ccATGGCCTAACAACAGtttatcttctt_ (SEQ ID NO:62)
Intron 13	30769	ccATGGCCtAACAACaGTTtatcttctt_ (SEQ ID NO:62)
Intron 12	30872	ctTGGCTGTGTAGGAGGGGAgtagcagt_ (SEQ ID NO:63)
Intron 12	30873	ttCCTAAGTTGGCAGTGGCAtgcttaat_ (SEQ ID NO:64)
Intron 12	30876	ctTTGGCTTTGAGGATTAAGcatgccac_ (SEQ ID NO:65)
Intron 12	30877	acTTGGCTcCAAGATTTATAGccttaaa_ (SEQ ID NO:66)
Intron 12	30882	caGGAAAGTAAGATAGGAAGgaatgtga_ (SEQ ID NO:67)

Target	SBS #	Target site
Intron 12	30883	ctGGGGTAAATGTCTCCttgctcttctt_ (SEQ ID NO:68)

Example 2: Activity of murine albumin-specific nucleases

[0149] ZFN pairs targeting the murine albumin gene were used to test the ability of these ZFNs to induce DSBs at a specific target site. The amino acid sequence of the recognition helix region of the indicated ZFNs are shown below in Table 3 and their target sites shown in Table 4 (DNA target sites indicated in uppercase letters; non-contacted nucleotides indicated in lowercase).

[0150] The Cel-I assay (Surveyor™, Transgenomics) as described in Perez et al, (2008) Nat. Biotechnol. 26: 808-816 and Guschin et al, (2010) Methods Mol Biol. 649:247-56), was used to detect ZFN-induced modifications. In this assay, PCR-amplification of the target site was followed by quantification of insertions and deletions (indels) using the mismatch detecting enzyme Cel-I (Yang et al, (2000) Biochemistry 39, 3533-3541) which provided a lower-limit estimate of DSB frequency. Three days following transfection of the ZFN expression vector at either standard conditions (37°C) or using a hypothermic shock (30°C, see co-owned U.S. Publication No. 20110041195), genomic DNA was isolated from Neuro2A cells transfected with the ZFN(s) using the DNeasy™ kit (Qiagen). In these experiments, all ZFN pairs were ELD/KKR *FokI* mutation pairs (described above).

[0151] A composite of the results from the Cel-I assay are shown in Figure 1, and demonstrate that the ZFNs shown below are capable of inducing cleavage at their respective target sites. The percent indels shown beneath the lanes indicates the amount of genomes that were altered by NHEJ following cleavage. The data also demonstrates increased activity when the transduction procedure incorporates the hypothermic shock.

Example 3: *In vivo* cleavage of the albumin locus

[0152] The mouse albumin specific ZFNs SBS30724 and SBS30725 which target a sequence in intron 1 were tested *in vivo*. The ZFNs were introduced into an AAV2/8 vector as described previously (Li et al (2011) Nature 475 (7355): 217). To facilitate production in the baculovirus system, the vector AAV2/8.2 was used for preparations destined for baculoviral production. AAV2/8.2 differs from the AAV2/8 vector in that a portion of the AAV8 capsid has been removed and replaced by a same region from the AAV2 capsid creating a chimeric capsid. The region is the phospholipase A2 domain in VP1. Production of the ZFN containing virus particles was done either by preparation using a HEK293 system or a baculovirus system using standard methods in the art (See Li et al, *ibid*, see *e.g.* US6723551). The virus particles were then administered to normal male mice (n=6) using a single dose of 200 microliter of 1.0e11 total vector genomes of either AAV2/8 or AAV2/8.2 encoding the mouse albumin-specific ZFN. 14 days post administration of rAAV vectors, mice were sacrificed, livers harvested and processed for DNA or total proteins using standard methods known in the art. Detection of AAV vector genome copies was performed by quantitative PCR. Briefly, qPCR primers were made specific to the bGHpA sequences within the AAV as follows:

Oligo200 (Forward) 5'-GTTGCCAGCCATCTGTTGTTT-3' (SEQ ID NO:69)

Oligo201 (Reverse) 5'-GACAGTGGGAGTGGCACCTT-3' (SEQ ID NO:70)

Oligo202 (Probe) 5'-CTCCCCCGTGCCTTCCTTGACC-3' (SEQ ID NO:71)

[0153] Cleavage activity of the ZFN was measured using a Cel-1 assay performed using a LC-GX apparatus (Perkin Elmer), according to manufacturer's protocol. Expression of the ZFNs *in vivo* was measured using a FLAG-Tag system according to standard methods. The results (Figure 3) demonstrated that the ZFNs were expressed, and that they are active in cleaving the target in the mouse liver gene. Shown in the Figure are the Cel-I NHEJ results for each mouse in the study. The type of vector and their contents are shown above the lanes. Mismatch repair following ZFN cleavage (indicated % indels) was detected at nearly 16% in some of the mice.

[0154] Albumin-specific TALENs were also tested as set forth in U.S. Application No. 13/624,193 and 13/624,217.

Example 4: *In vivo* insertion of a corrected disease associated gene

[0155] The murine specific albumin ZFNs or TALENs are then used to introduce transgene encoding a therapeutic gene product into the albumin locus for expression. Donors were designed to insert the correct gene for Fabry's disease (GLA), Gaucher's disease (GBA), Hurler's disease (IDUA), and Hunter's disease into the albumin locus. In these donor constructs, the therapeutic gene was flanked by sequences homologous to the albumin gene. 5' of the transgene, the donor constructs all contain sequences homologous to the murine albumin intron 1, while 3' of the gene, the constructs contain sequences homologous to the murine albumin intron 1-exon 2 boundary.

[0156] The donor constructs are then incorporated into an AAV genome and the resulting AAV particles containing the donors are then purified using methods known in the art. The material is used to produce AAV viruses containing AAV-donor genomes using the triple transfection method into HEK 293T cells and purified on CsCl density gradients as has been described (see Ayuso et al. (2010) Gene Ther 17(4), 503-510). AAV vector will be diluted in PBS prior to injection. A range of 5e9 to 5e13 v.g. AAV-donor vector particles will be used in conjunction with 1e9 to 1e12 vg of AAV-ZFN vector particles via tail vein or intraperitoneal injections of the viruses in wild-type, or disease model mice. AAV-ZFN genomes, described previously, containing the mouse albumin-specific ZFNs will be used, in combination with the GLA, GBA, IDUA and IDS AAV-donors. Cel-I and PCR assays will be performed on liver DNA isolated at various time points to determine the frequency of NHEJ and ZFN induced donor insertion. Southern blots may also be used. As per standard protocol, quantification of human GLA, GBA, IDUA and IDS in plasma will be performed using a human GLA, GBA, IDUA and IDS ELISA kit or using a FLAG Tag ELISA kit. Standard Western blots are also performed. The results demonstrate that these corrective transgenes can increase the expression of the therapeutic protein *in vivo*.

[0157] For example, the gene encoding human alpha galactosidase (deficient in patients with Fabry's disease) was inserted into the mouse albumin locus. The ZFN pair 30724/30725 was used as described above using a alpha galactosidase transgene. In this experiment, 3 mice were treated with an AAV2/8 virus containing the ZFN pair at a dose of 3.0e11 viral genomes per mouse and an AAV2/8 virus containing the huGLa donor at 1.5e12 viral genomes per mouse. Control animals were given either the ZFN containing virus alone or the huGLa donor virus alone. Western blots done on liver homogenates

showed an increase in alpha galactosidase-specific signal, indicating that the alpha galactosidase gene had been integrated and was being expressed (Figure 4A). In addition, an ELISA was performed on the liver lysate using a human alpha galactosidase assay kit (Sino) according to manufacturer's protocol. The results, shown in Figure 4B, demonstrated an increase in signal in the mice that had been treated with both the ZFNs and the huGLa donor.

Example 5: Design of human albumin specific ZFNs.

[0158] To design ZFNs with specificity for the human albumin gene, the DNA sequence of human albumin intron 1 was analyzed using previously described methods to identify target sequences with the best potential for ZFN binding (for details, see co-owned United States Patent Application No. 13/624,193 and 13/624,217). The target and helices are shown in Tables 5 and 6.

Table 5: Human albumin-specific zinc finger nucleases helix designs

Target	SBS #	Design					
		F1	F2	F3	F4	F5	F6
Intron 1	35393	QSSDLSR (SEQ ID NO:46)	LRHNLRA (SEQ ID NO:72)	DQSNLRA (SEQ ID NO:73)	RPYTLRL (SEQ ID NO:74)	QSSDLSR (SEQ ID NO:46)	HRSNLNK (SEQ ID NO:75)
Intron 1	35394	QSSDLSR (SEQ ID NO:46)	HRSNLNK (SEQ ID NO:75)	DQSNLRA (SEQ ID NO:73)	RPYTLRL (SEQ ID NO:74)	QSSDLSR (SEQ ID NO:46)	HRSNLNK (SEQ ID NO:75)
Intron 1	35396	QSSDLSR (SEQ ID NO:46)	LKWNLRT (SEQ ID NO:76)	DQSNLRA (SEQ ID NO:73)	RPYTLRL (SEQ ID NO:74)	QSSDLSR (SEQ ID NO:46)	HRSNLNK (SEQ ID NO:75)
Intron 1	35398	QSSDLSR (SEQ ID NO:46)	LRHNLRA (SEQ ID NO:72)	DQSNLRA (SEQ ID NO:73)	RPYTLRL (SEQ ID NO:74)	QSSDLSR (SEQ ID NO:46)	HRSNLNK (SEQ ID NO:75)
Intron 1	35399	QSSDLSR (SEQ ID NO:46)	HRSNLNK (SEQ ID NO:75)	DQSNLRA (SEQ ID NO:73)	RPYTLRL (SEQ ID NO:74)	QSSDLSR (SEQ ID NO:46)	HRSNLNK (SEQ ID NO:75)
Intron 1	35405	QSSDLSR (SEQ ID NO:46)	WKWNLRA (SEQ ID NO:77)	DQSNLRA (SEQ ID NO:73)	RPYTLRL (SEQ ID NO:74)	QSSDLSR (SEQ ID NO:46)	HRSNLNK (SEQ ID NO:75)
Intron 1	35361	QSGNLAR (SEQ ID NO:5)	LMQNRN Q (SEQ ID NO:78)	LKQHLNE (SEQ ID NO:79)	TSGNLTR (SEQ ID NO:11)	RRYYLRL (SEQ ID NO:80)	N/A
Intron 1	35364	QSGNLAR (SEQ ID NO:5)	HLGNLKT (SEQ ID NO:81)	LKQHLNE (SEQ ID NO:79)	TSGNLTR (SEQ ID NO:11)	RRDWRR D (SEQ ID NO:82)	N/A
Intron 1	35370	QSGNLAR (SEQ ID NO:5)	LMQNRN Q (SEQ ID NO:78)	LKQHLNE (SEQ ID NO:79)	TSGNLTR (SEQ ID NO:11)	RRDWRR D (SEQ ID NO:82)	N/A
Intron 1	35379	QRSNLVR (SEQ ID NO:83)	TSSNRKT (SEQ ID NO:53)	LKHHLTD (SEQ ID NO:84)	TSGNLTR (SEQ ID NO:11)	RRDWRR D (SEQ ID NO:82)	N/A

Target	SBS #	Design					
		F1	F2	F3	F4	F5	F6
Intron 1	35458	DKSYLRP (SEQ ID NO:85)	TSGNLTR (SEQ ID NO:11)	HRSARKR (SEQ ID NO:86)	QSSDLR (SEQ ID NO:46)	WRSSLKT (SEQ ID NO:87)	N/A
Intron 1	35480	TSGNLTR (SEQ ID NO:11)	HRSARKR (SEQ ID NO:86)	QSGDLTR (SEQ ID NO:40)	NRHHLKS (SEQ ID NO:88)	N/A	N/A
Intron 1	35426	QSGDLTR (SEQ ID NO:40)	QSGNLHV (SEQ ID NO:89)	QSAHRKN (SEQ ID NO:90)	STAALSY (SEQ ID NO:91)	TSGSLR (SEQ ID NO:92)	RSDALAR (SEQ ID NO:41)
Intron 1	35428	QSGDLTR (SEQ ID NO:40)	QRSNLNI (SEQ ID NO:93)	QSAHRKN (SEQ ID NO:90)	STAALSY (SEQ ID NO:91)	DRSALS (SEQ ID NO:52)	RSDALAR (SEQ ID NO:41)
Intron 1	34931	QRTHLTQ (SEQ ID NO:94)	DRSNLTR (SEQ ID NO:95)	QSGNLAR (SEQ ID NO:5)	QKVNRA (SEQ ID NO:96)	N/A	N/A
Intron 1	33940	RSDNLSV (SEQ ID NO:26)	QNANRIT (SEQ ID NO:97)	DQSNLRA (SEQ ID NO:73)	QSAHRIT (SEQ ID NO:98)	TSGNLTR (SEQ ID NO:11)	HRSARKR (SEQ ID NO:86)

Table 6: Target sites of Human albumin-specific ZFNs

Target	SBS #	Target site
Intron 1 (locus 2)	35393	<u>ccTATCCATTGCACTATGCTttatttaa</u> (SEQ ID NO:99)
Intron 1 (locus 2)	35394	<u>ccTATCCATTGCACTATGCTttatttaa</u> (SEQ ID NO:99)
Intron 1 (locus 2)	35396	<u>ccTATCCATTGCACTATGCTttatttaa</u> (SEQ ID NO:99)
Intron 1 (locus 2)	35398	<u>ccTATCCATTGCACTATGCTttatttaa</u> (SEQ ID NO:99)
Intron 1 (locus 2)	35399	<u>ccTATCCATTGCACTATGCTttatttaa</u> (SEQ ID NO:99)
Intron 1 (locus 2)	35405	<u>ccTATCCATTGCACTATGCTttatttaa</u> (SEQ ID NO:99)
Intron 1 (locus 2)	35361	<u>ttTGGGATAGTTATGAAttcaatcttca</u> (SEQ ID NO:100)
Intron 1 (locus 2)	35364	<u>ttTGGGATAGTTATGAAttcaatcttca</u> (SEQ ID NO:100)
Intron 1 (locus 2)	35370	<u>ttTGGGATAGTTATGAAttcaatcttca</u> (SEQ ID NO:100)
Intron 1 (locus 2)	35379	<u>ttTGGGATAGTTATGAAttcaatcttca</u> (SEQ ID NO:100)
Intron 1 (locus 3)	35458	<u>ccTGTGCTGTTGATCTCataaatagaac</u> (SEQ ID NO:101)
Intron 1 (locus 3)	35480	<u>ccTGTGCTGTTGATctcataaatagaac</u> (SEQ ID NO:101)
Intron 1 (locus 3)	35426	<u>ttGTGGTTTTTAAATAAAGCAtagtgca</u> (SEQ ID NO:102)
Intron 1 (locus 3)	35428	<u>ttGTGGTTTTTAAATAAAGCAtagtgca</u> (SEQ ID NO:102)
Intron 1 (locus 4)	34931	<u>acCAAGAAGACAGActaaaatgaaaata</u> (SEQ ID NO:103)
Intron 1 (locus 4)	33940	<u>ctGTTGATAGACACTAAAAGagtatttag</u> (SEQ ID NO:104)

[0159] These nucleases were tested in pairs to determine the pair with the highest activity. The resultant matrices of tested pairs are shown in Tables 7 and 8, below where the ZFN used for the right side of the dimer is shown across the top of each matrix, and the ZFN used for the left side of the dimer is listed on

the left side of each matrix. The resultant activity, as determined by percent of mismatch detected using the Cel-I assay is shown in the body of both matrices:

Table 7: Activity of Human albumin-specific ZFNs (%mutated targets)

	35393	35394	35396	35398	35399	35405	ave.
35361	18	19	25	22	23	21	21
35364	n.d.	24	23	19	21	21	22
35370	21	19	22	n.d.	22	23	21
35379	21	21	n.d.	19	19	21	20

Table 8: Activity of Human albumin-specific ZFNs (% mutated targets))

	35458	35480	ave.
35426	4.5	7	3
35428	4.9	6	3.6
(note: 'n.d.' means the assay on this pair was not done)			

[0160] Thus, highly active nucleases have been developed that recognize target sequences in intron 1 of human albumin.

Example 6: Design of albumin specific TALENs

[0161] TALENs were designed to target sequences within human albumin intron 1. Base recognition was achieved using the canonical RVD-base correspondences (the "TALE code: NI for A, HD for C, NN for G (NK in half repeat), NG for T). TALENs were constructed as previously described (see co-owned U.S. Patent Publication No. 20110301073). Targets for a subset of TALENs were conserved in cynomolgus monkey and rhesus macaque albumin genes (for details, see co-owned United States Patent Application 13/624,193 and 13/624,217). The TALENs were constructed in the "+17" and "+63" TALEN backbones as described in US20110301073. The targets and numeric identifiers for the TALENs tested are shown below in Table 9.

Table 9: Albumin specific TALENs

SBS #	site	# of RVDs	SEQ ID NO:
102249	gtTGAAGATTGAATTCAta	15	105
102250	gtTGAAGATTGAATTCATAac	17	106
102251	gtGCAATGGATAGGTCTtt	15	107
102252	atAGTGCAATGGATAGGtc	15	108
102253	atTGAATTCATAACTATcc	15	109
102254	atTGAATTCATAACTATCCca	17	110
102255	atAAAGCATAGTGCAATGGat	17	111
102256	atAAAGCATAGTGCAATgg	15	112
102257	ctATGCTTTATTTAAAac	15	113
102258	ctATGCTTTATTTAAAACca	17	114
102259	atTTATGAGATCAACAGCAca	17	115
102260	ctATTTATGAGATCAACAGca	17	116

SBS #	site	# of RVDs	SEQ ID NO:
102261	ttCATTTTAGTCTGTCTTctt	17	117
102262	atTTTAGTCTGTCTTCTtg	15	118
102263	ctAATACTCTTTTAGTGTct	16	119
102264	atCTAATACTCTTTTAGTGtc	17	120
102265	atAATTGAACATCATCctg	15	121
102266	atAATTGAACATCATCCTGag	17	122
102267	atATTGGGCTCTGATTCCTac	17	123
102268	atATTGGGCTCTGATTCct	15	124
102269	ttTTTCTGTAGGAATCAga	15	125
102270	ttTTTCTGTAGGAATCAGag	16	126
102271	ttATGCATTTGTTTCAAaa	15	127
102272	atTATGCATTTGTTTCAAaa	15	128

[0162] The TALENs were then tested in pairs in HepG2 cells for the ability to induce modifications at their endogenous chromosomal targets, and the results showed that many proteins bearing the +17 truncation point were active. Similarly, many TALENs bearing the +63 truncation point were also active (see Table 10). Side by side comparisons with three sets of non-optimized albumin ZFNs (see Table 10) showed that the TALENs and ZFNs have activities that are in the same approximate range.

Table 10: TALEN-induced target modification in HepG2-C3a cells

Sample pair	TALEN C17	%modification, C17	TALEN C63	% modification, C63	Gap
1	102251:102249	15	102251:102249	0	12
2	102251:102250	0	102251:102250	0	10
3	102252:102249	0	102252:102249	8.3	15
4	102252:102250	32	102252:102250	8.0	13
5	102255:102253	38	102255:102253	21	13
6	102255:102254	43	102255:102254	0	11
7	102256:102253	0	102256:102253	23	15
8	102256:102254	28	102256:102254	16	13
9	102259:102257	18	102259:102257	15	13
10	102259:102258	15	102259:102258	0	11
11	102260:102257	15	102260:102257	13	15
12	102260:102258	24	102260:102258	11	13
13	102263:102261	0	102263:102261	16	17
14	102263:102262	0	102263:102262	15	16
15	102264:102261	0	102264:102261	22	18
16	102264:102262	0	102264:102262	17	17
20	102267:102265	47	102267:102265	9.8	13
21	102267:102266	4.7	102267:102266	0	11

Sample pair	TALEN C17	%modification, C17	TALEN C63	% modification, C63	Gap
22	102268:102265	4.2	102268:102265	7.9	15
23	102268:102266	10	102268:102266	0	13
24	102271:102269	14	102271:102269	0	12
25	102271:102270	0	102271:102270	0	11
26	102272:102269	0	102272:102269	0	13
27	102272:102270	0	102272:102270	0	12
ZFNs					
17	35361:35396	31	35361:35396	29	6
18	35426:35458	10	35426:35458	7	6
19	34931:33940	7.3	34931:33940	7	6

[0163] As noted previously (see co-owned U.S. Patent Publication No. 20110301073), the C17 TALENs have greater activity when the gap size between the two TALEN target sites is approximately 11- 15 bp, while the C63 TALENs sustain activity at gap sizes up to 18 bp (see Table 10).

Example 7: Detection of LSD donor transgenes *in vivo*

[0164] Donors for four lysosomal storage disease transgenes were constructed for the purpose of integrating into the mouse albumin gene in intron1. The transgenes were α -galactosidase A (GLA), Acid β -glucosidase (GBA), α -L-iduronidase (IDUA) and Iduronate-2-sulfatase (IDS), genes lacking in Fabry's, Gaucher's, Hurler's and Hunter's diseases, respectively. See, e.g., Figure 8.

[0165] The donors were then used in *in vivo* studies to observe integration of the transgenes. The murine albumin specific ZFNs and the donors were inserted all into AAV2/8 virus as described in Example 4, and then were injected into mice. In these experiments, the virus was formulated for injection in D-PBS + 35 mM NaCl, 5% glycerol and frozen prior to injection. The donor- and nuclease-containing viruses were mixed together prior to freezing. At Day 0, the virus preparation was thawed and administered to the mice by tail vein injection where the total injection volume was 200 μ L. At the indicated times, the mice were sacrificed and then serum, liver and spleen were harvested for protein and DNA analysis by standard protocols. The dose groups are shown below in Table 11.

Table 11: Treatment groups for LSD transgene integration

Group	Treatment	N/group/time point
1	murine Alb intron 1+ Fabry@ 1:5 ratio ; ZFN @ 3.0e11,Donor @ 1.5e12	3
2	murine Alb intron 1+ Hunters donor@ 1:5 ratio ; ZFN @3.0e11, Donor @ 1.5e12	3
3	murine Alb intron 1+ Hurlers donor@ 1:5 ratio ; ZFN @3.0e11, Donor @ 1.5e12	3
4	murine Alb intron 1	3
5	Fabry donor only	2

Group	Treatment	N/group/time point
6	Hunter's donor only	2
7	Hurler's donor only	2

[0166] At day 30, liver homogenates were analyzed by Western blot analysis for the presence of the LSD proteins encoded by the donors. Liver homogenates were analyzed by Western blot using standard methodologies, using the following primary antibodies: α -Galactosidase A (Fabry's)-specific rabbit monoclonal antibody was purchased from Sino Biological, Inc.; Human α -L-Iduronidase (Hurler's)-specific mouse monoclonal antibody was purchased from R&D Systems; Human iduronate 2-Sulfatase (Hunter's)-specific mouse monoclonal was purchased from R&D Systems. The results (See Figure 5) demonstrate expression, especially in the mice that received both the ZFN containing virus and the transgene donor containing virus.

[0167] The manner of integration of the donor DNA into the albumin locus was also investigated. Following the cleavage at the albumin locus, the donor transgene could be potentially be integrated via homology directed recombination (HDR), utilizing the regions of homology flanking the transgene donor (Figure 2), or the transgene may be captured during the error-prone non-homologous end joining process (NHEJ). The results of these two alternatives will yield insertions of differing sizes when subject to PCR, using either the Acc651-SA-rev-sh primer (5'AAG AAT AAT TCT TTA GTG GTA 3', SEQ ID NO:129) which binds to the F9 splice acceptor site in all LSD donor constructs and the mALB-OOF1 primer (5' ATGAAGTGGGTAACCTTTCTC 3', SEQ ID NO:130) which binds to the mouse albumin exon 1 upstream of the ZFN cleavage site (see Figure 6), where the integration of the transgene by HDR will result in insertion of approximately 680 bp while integration via NHEJ will result in integration of approximately 1488 bp. Thus, genomic DNA isolated from the treated mice liver homogenates was subject to PCR in the presence of ³²P-radiolabeled nucleotides and run on a gel. In all three of the transgene integrations, integration via both mechanisms was observed (see Figure 7).

[0168] Donor DNAs were also designed to include a tag sequence for later recognition of the protein using the tag specific antibodies. The tag was designed to incorporate two different sequences encoding the Myc (EQKLISEEDL, SEQ ID NO:131) and the Flag (DYKDDDD SEQ ID NO:132) tags. Schematics of the donor designs are shown in Figure 8. The donors were integrated as described above, and all were capable of integration as demonstrated by PCR (see Figure 9).

SEQUENCE LISTING

[0169]

<110> SANGAMO BIOSCIENCES, INC.

<120> METHODS AND COMPOSITIONS FOR THE TREATMENT OF LYSOSOMAL STORAGE DISEASES

<130> 8325-0097.40

<140> PCT/US2013/032381

<141> 2013-03-15

<150> 61/704,072

<151> 2012-09-21

<150> 61/670,463

<151> 2012-07-11

<160> 134

<170> PatentIn version 3.5

<210> 1

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 1

Thr	Ser	Gly	Ser	Leu	Thr	Arg
1				5		

<210> 2

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 2

Arg	Ser	Asp	Ala	Leu	Ser	Thr
1				5		

<210> 3

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 3

Gln	Ser	Ala	Thr	Arg	Thr	Lys
1				5		

<210> 4

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 4
Thr Ser Gly His Leu Ser Arg
1 5

<210> 5
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 5
Gln Ser Gly Asn Leu Ala Arg
1 5

<210> 6
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 6
Arg Ser Asp His Leu Ser Ala
1 5

<210> 7
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 7
Thr Lys Ser Asn Arg Thr Lys
1 5

<210> 8
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 8
Asp Arg Ser Asn Leu Ser Arg
1 5

<210> 9
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 9
Trp Arg Ser Ser Leu Arg Ala
1 5

<210> 10
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 10
Asp Ser Ser Asp Arg Lys Lys
1 5

<210> 11
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 11
Thr Ser Gly Asn Leu Thr Arg
1 5

<210> 12
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 12
Asp Arg Ser Thr Arg Arg Gln
1 5

<210> 13
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 13
Glu Arg Gly Thr Leu Ala Arg
1 5

<210> 14
<211> 7

<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 14
Thr Ser Ala Asn Leu Ser Arg
1 5

<210> 15
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 15
Asp Arg Ser Ala Leu Ala Arg
1 5

<210> 16
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 16
Arg Ser Asp His Leu Ser Glu
1 5

<210> 17
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 17
His Arg Ser Asp Arg Thr Arg
1 5

<210> 18
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 18
Gln Ser Gly Ala Leu Ala Arg
1 5

<210> 19
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 19
Gln Ser Gly His Leu Ser Arg
1 5

<210> 20
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 20
Arg Ser Asp Asn Leu Ser Thr
1 5

<210> 21
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 21
Asp Gly Arg Asn Leu Arg His
1 5

<210> 22
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 22
Arg Ser Asp Asn Leu Ala Arg
1 5

<210> 23
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 23
Gln Ser Asn Ala Leu Asn Arg
1 5

<210> 24
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 24
Leu Lys Gln Val Leu Val Arg
1 5

<210> 25
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 25
Gln Ser Thr Pro Leu Phe Ala
1 5

<210> 26
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 26
Arg Ser Asp Asn Leu Ser Val
1 5

<210> 27
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 27
His Ser Asn Ala Arg Lys Thr
1 5

<210> 28
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 28
Arg Ser Asp Ser Leu Ser Ala
1 5

<210> 29
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 29
Arg Ser Asp Ser Leu Ser Val
1 5

<210> 30
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 30
Arg Ser Asp Asn Leu Ser Glu
1 5

<210> 31
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 31
Glu Arg Ala Asn Arg Asn Ser
1 5

<210> 32
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 32
Gln Ser Ala Asn Arg Thr Lys
1 5

<210> 33
<211> 7

<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 33
Arg Ser Asp Ala Leu Thr Gln
1 5

<210> 34
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 34
Gln Ser Gly His Leu Ala Arg
1 5

<210> 35
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 35
Arg Ser Asp His Leu Thr Gln
1 5

<210> 36
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 36
Arg Ser Asp His Leu Ser Gln
1 5

<210> 37
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 37
Trp Arg Ser Ser Leu Val Ala
1 5

<210> 38
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 38
Arg Ser Asp Val Leu Ser Glu
1 5

<210> 39
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 39
Arg Asn Gln His Arg Lys Thr
1 5

<210> 40
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 40
Gln Ser Gly Asp Leu Thr Arg
1 5

<210> 41
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 41
Arg Ser Asp Ala Leu Ala Arg
1 5

<210> 42
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 42
Arg Arg Asp Pro Leu Ile Asn
1 5

<210> 43
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 43
Ile Arg Ser Thr Leu Arg Asp
1 5

<210> 44
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 44
Tyr Ser Ser Thr Arg Asn Ser
1 5

<210> 45
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 45
Ser Tyr Trp Ser Arg Thr Val
1 5

<210> 46
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 46
Gln Ser Ser Asp Leu Ser Arg
1 5

<210> 47
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 47
Arg Thr Asp Ala Leu Arg Gly
1 5

<210> 48
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 48
Gln Lys Ser Pro Leu Asn Thr
1 5

<210> 49
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 49
Gln Ala Glu Asn Leu Lys Ser
1 5

<210> 50
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 50
Arg Arg Ala His Leu Asn Gln
1 5

<210> 51
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 51
Ser Asp Thr Asn Arg Phe Lys
1 5

<210> 52
<211> 7

<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 52
Asp Arg Ser Ala Leu Ser Arg
1 5

<210> 53
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 53
Thr Ser Ser Asn Arg Lys Thr
1 5

<210> 54
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 54
Arg Ser Asp His Leu Ser Arg
1 5

<210> 55
<211> 28
<212> DNA
<213> Mus sp.

<400> 55
ctgaaggtag caatggtcc tctctgct 28

<210> 56
<211> 28
<212> DNA
<213> Mus sp.

<400> 56
tttctgtaa cgatcgggaa ctggcatc 28

<210> 57
<211> 28
<212> DNA
<213> Mus sp.

<400> 57
aagatgccag ttcccgatcg ttacagga 28

<210> 58
<211> 28
<212> DNA
<213> Mus sp.

<400> 58
agggagtagc ttaggtcagt gaagagaa 28

<210> 59
<211> 28
<212> DNA
<213> Mus sp.

<400> 59
acgtagagaa caacatctag attggtgg 28

<210> 60
<211> 28
<212> DNA
<213> Mus sp.

<400> 60
ctgtaataga aactgactta cgtagctt 28

<210> 61
<211> 28
<212> DNA
<213> Mus sp.

<400> 61
agggaatgtg aaatgattca gatata 28

<210> 62
<211> 28
<212> DNA
<213> Mus sp.

<400> 62
ccatggccta acaacagttt atcttctt 28

<210> 63
<211> 28
<212> DNA
<213> Mus sp.

<400> 63
cttggctgtg taggagggga gtagcagt 28

<210> 64
<211> 28

<212> DNA

<213> Mus sp.

<400> 64

ttcctaagtt ggcagtggca tgcttaat 28

<210> 65

<211> 28

<212> DNA

<213> Mus sp.

<400> 65

cttggcttt gaggattaag catgccac 28

<210> 66

<211> 28

<212> DNA

<213> Mus sp.

<400> 66

acttggctcc aagatttata gccttaaa 28

<210> 67

<211> 28

<212> DNA

<213> Mus sp.

<400> 67

caggaaagta agataggaag gaatgtga 28

<210> 68

<211> 28

<212> DNA

<213> Mus sp.

<400> 68

ctggggtaaa tgtctccttg ctcttctt 28

<210> 69

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<400> 69

gttgccagcc atctgttgtt t 21

<210> 70

<211> 20

<212> DNA

<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 70
gacagtggga gtggcacctt 20

<210> 71
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic probe

<400> 71
ctccccgtg ccttcctga cc 22

<210> 72
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 72
Leu Arg His Asn Leu Arg Ala
1 5

<210> 73
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 73
Asp Gln Ser Asn Leu Arg Ala
1 5

<210> 74
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 74
Arg Pro Tyr Thr Leu Arg Leu
1 5

<210> 75
<211> 7

<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 75
His Arg Ser Asn Leu Asn Lys
1 5

<210> 76
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 76
Leu Lys Trp Asn Leu Arg Thr
1 5

<210> 77
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 77
Trp Lys Trp Asn Leu Arg Ala
1 5

<210> 78
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 78
Leu Met Gln Asn Arg Asn Gln
1 5

<210> 79
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 79
Leu Lys Gln His Leu Asn Glu
1 5

<210> 80
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 80
Arg Arg Tyr Tyr Leu Arg Leu
1 5

<210> 81
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 81
His Leu Gly Asn Leu Lys Thr
1 5

<210> 82
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 82
Arg Arg Asp Trp Arg Arg Asp
1 5

<210> 83
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 83
Gln Arg Ser Asn Leu Val Arg
1 5

<210> 84
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 84
Leu Lys His His Leu Thr Asp
1 5

<210> 85
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 85
Asp Lys Ser Tyr Leu Arg Pro
1 5

<210> 86
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 86
His Arg Ser Ala Arg Lys Arg
1 5

<210> 87
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 87
Trp Arg Ser Ser Leu Lys Thr
1 5

<210> 88
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 88
Asn Arg His His Leu Lys Ser
1 5

<210> 89
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 89
Gln Ser Gly Asn Leu His Val
1 5

<210> 90
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 90
Gln Ser Ala His Arg Lys Asn
1 5

<210> 91
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 91
Ser Thr Ala Ala Leu Ser Tyr
1 5

<210> 92
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 92
Thr Ser Gly Ser Leu Ser Arg
1 5

<210> 93
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 93
Gln Arg Ser Asn Leu Asn Ile
1 5

<210> 94
<211> 7

<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 94
Gln Arg Thr His Leu Thr Gln
1 5

<210> 95
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 95
Asp Arg Ser Asn Leu Thr Arg
1 5

<210> 96
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 96
Gln Lys Val Asn Arg Ala Gly
1 5

<210> 97
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 97
Gln Asn Ala Asn Arg Ile Thr
1 5

<210> 98
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 98
Gln Ser Ala His Arg Ile Thr
1 5

<210> 99
<211> 28
<212> DNA
<213> Homo sapiens

<400> 99
cctatccatt gcactatgct ttatttaa 28

<210> 100
<211> 28
<212> DNA
<213> Homo sapiens

<400> 100
ttgggatag ttatgaattc aatcttca 28

<210> 101
<211> 28
<212> DNA
<213> Homo sapiens

<400> 101
cctgtgctgt tgatctcata aatagaac 28

<210> 102
<211> 28
<212> DNA
<213> Homo sapiens

<400> 102
ttgtggtttt taaataaagc atagtgc 28

<210> 103
<211> 28
<212> DNA
<213> Homo sapiens

<400> 103
accaagaaga cagactaaaa tgaaaat 28

<210> 104
<211> 28
<212> DNA
<213> Homo sapiens

<400> 104
ctgttgatag acactaaaag agtattag 28

<210> 105
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 105

gttgaagatt gaattcata 19

<210> 106

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 106

gttgaagatt gaattcataa c 21

<210> 107

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 107

gtgcaatgga taggtcttt 19

<210> 108

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 108

atagtgcaat ggataggtc 19

<210> 109

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 109

attgaattca taactatcc 19

<210> 110

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 110

attgaattca taactatccc a 21

<210> 111

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 111

ataaagcata gtgcaatgga t 21

<210> 112

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 112

ataaagcata gtgcaatgg 19

<210> 113

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 113

ctatgcttta tttaaaaac 19

<210> 114

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 114

ctatgcttta tttaaaaacc a 21

<210> 115

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 115

atttatgaga tcaacagcac a 21

<210> 116

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 116

ctatttatga gatcaacagc a 21

<210> 117

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 117

ttcattttag tctgtcttct t 21

<210> 118

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 118

attttagtct gtcttcttg 19

<210> 119

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 119

ctaatactct tttagtgtct 20

<210> 120

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 120

atctaatact ctttagtgt c 21

<210> 121

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 121

ataattgaac atcatcctg 19

<210> 122

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 122

ataattgaac atcatcctga g 21

<210> 123

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 123

atattgggct ctgattccta c 21

<210> 124

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 124

atattgggct ctgattcct 19

<210> 125

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 125

ttttctgta ggaatcaga 19

<210> 126

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 126

ttttctgta ggaatcagag 20

<210> 127

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 127

ttatgcattt gttcaaaa 19

<210> 128

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 128

attatgcatt tgttcaaa 19

<210> 129

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<400> 129
aagaataatt ctttagtggt a 21

<210> 130
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 130
atgaagtggg taaccttct c 21

<210> 131
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 131
Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu
1 5 10

<210> 132
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 132
Asp Tyr Lys Asp Asp Asp Asp
1 5

<210> 133
<211> 9
<212> PRT
<213> Unknown

<220>
<223> Description of Unknown: Unknown LAGLIDADG family sequence

<400> 133
Leu Ala Gly Leu Ile Asp Ala Asp Gly
1 5

<210> 134
<211> 29
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 134

Arg	Thr	Arg	Pro	Leu	Glu	Gln	Lys	Leu	Ile	Ser	Glu	Glu	Asp	Leu	Ala
1				5					10					15	

Ala	Asn	Asp	Ile	Leu	Asp	Tyr	Lys	Asp	Asp	Asp	Asp	Lys
			20					25				

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US7888121B [0003]
- US20080299580A [0004] [0013] [0019] [0020] [0113]
- US20080159996A [0004] [0013] [0019] [0020] [0113]
- US201000218264A [0004] [0013] [0019] [0020] [0113]
- US13660821B [0004] [0013] [0020]
- US61670451 [0013] [0013] [0019] [0020]
- US61537349 [0013]
- US61560506B [0013] [0148]
- US660821 [0013] [0019]
- US13660843B [0013] [0019] [0020]
- WO2002007752A [0015] [0145]
- US13624193B [0019] [0020] [0158] [0161]
- US13624217B [0019] [0020] [0084] [0148] [0154] [0158] [0161]
- US6140081A [0043]
- US6453242B [0043] [0085] [0086] [0089] [0105] [0121]
- US6534261B [0043] [0085] [0086] [0089] [0105] [0121] [0122] [0148]
- WO9853058A [0043] [0089]
- WO9853059A [0043] [0089]
- WO9853060A [0043] [0089]
- WO02016536A [0043] [0089]
- WO03016496A [0043] [0089]
- US20110301073A [0043] [0044] [0083] [0084] [0089] [0091] [0105] [0108] [0161] [0161] [0163]
- US5789538A [0044] [0087] [0106]
- US5925523A [0044] [0087] [0089] [0106]
- US6007988A [0044] [0087] [0089] [0106]
- US6013453A [0044] [0087] [0089] [0106]
- US6200759B [0044] [0087] [0089] [0106]

- [WO9519431A \[0044\] \[0089\]](#)
- [WO9606166A \[0044\] \[0089\]](#)
- [WO9853057A \[0044\] \[0087\] \[0089\] \[0106\]](#)
- [WO9854311A \[0044\] \[0089\]](#)
- [WO0027878A \[0044\] \[0087\] \[0089\] \[0106\]](#)
- [WO0160970A \[0044\] \[0089\]](#)
- [WO0188197A \[0044\] \[0087\] \[0089\] \[0106\]](#)
- [WO02099084A \[0044\] \[0089\]](#)
- [US20050064474A \[0054\] \[0085\] \[0091\] \[0098\] \[0101\] \[0103\]](#)
- [US20070218528A \[0054\] \[0085\]](#)
- [US20080131962A \[0054\] \[0098\] \[0100\] \[0101\]](#)
- [US20110201055A \[0054\] \[0101\] \[0101\]](#)
- [US5176996A \[0062\]](#)
- [US5422251A \[0062\]](#)
- [US5585245A \[0076\]](#)
- [WO9844350A \[0076\]](#)
- [US5420032A \[0081\] \[0082\]](#)
- [US6833252B \[0081\] \[0082\]](#)
- [US20070117128A \[0082\]](#)
- [WO2010079430A \[0083\]](#)
- [US20110145940A \[0084\]](#)
- [US624193 \[0084\] \[0148\] \[0154\]](#)
- [US6599692B \[0085\] \[0121\]](#)
- [US6503717B \[0085\] \[0121\]](#)
- [US6689558B \[0085\] \[0121\]](#)
- [US7030215B \[0085\]](#)
- [US6794136B \[0085\]](#)
- [US7067317B \[0085\]](#)
- [US7262054B \[0085\]](#)
- [US7070934B \[0085\]](#)
- [US7361635B \[0085\]](#)
- [US7253273B \[0085\]](#)
- [US20050267061A \[0085\]](#)
- [US6410248B \[0087\] \[0106\]](#)
- [US6140466A \[0087\] \[0106\]](#)
- [US6242568B \[0087\] \[0106\]](#)
- [WO9837186A \[0087\] \[0106\]](#)
- [GB2338237A \[0087\] \[0106\]](#)
- [WO02077227A \[0087\] \[0088\]](#)
- [US6479626B \[0088\] \[0090\] \[0108\]](#)
- [US6903185B \[0088\] \[0090\] \[0108\]](#)
- [US7153949B \[0088\] \[0090\] \[0108\]](#)
- [US61400815B \[0089\]](#)
- [US789538A \[0089\]](#)
- [US20030232410A \[0091\] \[0103\]](#)
- [US20050208489A \[0091\] \[0103\]](#)
- [US20050026157A \[0091\] \[0103\]](#)
- [US20060166987A \[0091\] \[0098\] \[0103\] \[0107\]](#)

- [US20060063231A \[0091\] \[0103\]](#)
- [WO07014275A \[0091\] \[0097\] \[0103\]](#)
- [US5356802A \[0094\]](#)
- [US5436150A \[0094\]](#)
- [US5487994A \[0094\]](#)
- [US20090068164A \[0102\]](#)
- [WO2009042163A \[0103\]](#)
- [WO20090068164A \[0103\]](#)
- [US20050064474 \[0107\]](#)
- [US6607882B \[0121\] \[0122\]](#)
- [US6624976B \[0121\] \[0122\]](#)
- [US6933113B \[0121\] \[0122\]](#)
- [US6979539B \[0121\] \[0122\]](#)
- [US7013219B \[0121\] \[0122\]](#)
- [US7163824B \[0121\] \[0122\]](#)
- [US6008336A \[0125\]](#)
- [US5049386A \[0125\]](#)
- [US4946787A \[0125\] \[0126\]](#)
- [US4897355A \[0125\]](#)
- [WO9117424A \[0125\]](#)
- [WO9116024A \[0125\]](#)
- [US4186183A \[0126\]](#)
- [US4217344A \[0126\]](#)
- [US4235871A \[0126\]](#)
- [US4261975A \[0126\]](#)
- [US4485054A \[0126\]](#)
- [US4501728A \[0126\]](#)
- [US4774085A \[0126\]](#)
- [US4837028A \[0126\]](#)
- [US9405700W \[0129\]](#)
- [US4797368A \[0130\]](#)
- [WO9324641A \[0130\]](#)
- [US5173414A \[0130\]](#)
- [US2009054985A \[0139\]](#)
- [US61537349B \[0148\]](#)
- [US20110041195A \[0150\]](#)
- [US6723551B \[0152\]](#)
- [US2013032381W \[0169\]](#)
- [US61704072B \[0169\]](#)
- [US61670463B \[0169\]](#)

Non-patent literature cited in the description

- **GABATHULER et al.** Neurobiology of Disease, 2010, vol. 37, 48-57 [\[0014\]](#)
- **KORMANN et al.** Nature Biotechnology, 2011, vol. 29, 2154-157 [\[0016\]](#)

- **SAMBROOK et al.** MOLECULAR CLONING: A LABORATORY MANUAL Cold Spring Harbor Laboratory Press 1989 0000 [0036]
- **MOLECULAR CLONING: A LABORATORY MANUAL** 2001 0000 [0036]
- **AUSUBEL et al.** CURRENT PROTOCOLS IN MOLECULAR BIOLOGY John Wiley & Sons 1987 0000 [0036]
- **METHODS IN ENZYMOLOGY** Academic Press [0036]
- **WOLFFE** CHROMATIN STRUCTURE AND FUNCTION Academic Press 1998 0000 [0036]
- **Chromatin** METHODS IN ENZYMOLOGY Academic Press 1999 0000 vol. 304, [0036]
- **Chromatin Protocols** METHODS IN MOLECULAR BIOLOGY Humana Press 1999 0000 vol. 119, [0036]
- **FIELDS et al.** Nature, 1989, vol. 340, 245-246 [0076]
- **BELFORT et al.** 25 Nucleic Acids Res., 1997, 3379-3388 [0081]
- **DUJON et al.** Gene, 1989, vol. 82, 115-118 [0081] [0082]
- **PERLER et al.** Nucleic Acids Res., 1994, vol. 22, 1125-1127 [0081] [0082]
- **JASIN** Trends Genet., 1996, vol. 12, 224-228 [0081] [0082]
- **GIMBLE et al.** J. Mol. Biol., 1996, vol. 263, 163-180 [0081] [0082]
- **ARGAST et al.** J. Mol. Biol., 1998, vol. 280, 345-353 [0081] [0082]
- **BELFORT et al.** Nucleic Acids Res., 1997, vol. 25, 3379-3388 [0082] [0092]
- **CHEVALIER et al.** Molec. Cell, 2002, vol. 10, 895-905 [0082]
- **EPINAT et al.** Nucleic Acids Res., 2003, vol. 31, 2952-2962 [0082]
- **ASHWORTH et al.** Nature, 2006, vol. 441, 656-659 [0082]
- **PAQUES et al.** Current Gene Therapy, 2007, vol. 7, 49-66 [0082]
- **KAY et al.** Science, 2007, vol. 318, 648-651 [0083]
- **BONAS et al.** Mol Gen Genet, 1989, vol. 218, 127-136 [0083]
- **SCHORNACK S et al.** J Plant Physiol, 2006, vol. 163, 3256-272 [0083]
- **HEUER et al.** Appl and Envir Micro, 2007, vol. 73, 134379-4384 [0083]
- **BOCH et al.** Science, 2009, vol. 326, 1509-1512 [0084]
- **MOSCOUBOGDANOV** Science, 2009, vol. 326, 1501- [0084]
- **HEUER et al.** Applied and Environmental Microbiology, 2007, vol. 73, 134379-4384 [0084]
- **BEERLI et al.** Nature Biotechnol., 2002, vol. 20, 135-141 [0085]
- **PABO et al.** Ann. Rev. Biochem., 2001, vol. 70, 313-340 [0085]
- **ISALAN et al.** Nature Biotechnol., 2001, vol. 19, 656-660 [0085]
- **SEGAL et al.** Curr. Opin. Biotechnol., 2001, vol. 12, 632-637 [0085]
- **CHOO et al.** Curr. Opin. Struct. Biol., 2000, vol. 10, 411-416 [0085]
- **KIM et al.** Proc Nat'l Acad Sci USA, 1996, vol. 93, 31156-1160 [0091]
- **Nucleases** Cold Spring Harbor Laboratory Press 1993 0000 [0092]
- **LI et al.** Proc. Natl. Acad. Sci. USA, 1992, vol. 89, 4275-4279 [0094]
- **LI et al.** Proc. Natl. Acad. Sci. USA, 1993, vol. 90, 2764-2768 [0094]
- **KIM et al.** Proc. Natl. Acad. Sci. USA, 1994, vol. 91, 883-887 [0094]
- **KIM et al.** J. Biol. Chem., 1994, vol. 269, 31978-31982- [0094]
- **BITINAITE et al.** Proc. Natl. Acad. Sci. USA, 1998, vol. 95, 10570-10575- [0095]
- **ROBERTS et al.** Nucleic Acids Res., 2003, vol. 31, 418-420 [0097]
- **JINEK et al.** Science, 2012, vol. 337, 816-821 [0104]
- **JINEK et al.** eLife, 2013, vol. 2, e00471- [0104]
- **DAVID SEGAL** eLife, 2013, vol. 2, e00563- [0104]
- **CHANG et al.** Proc. Natl. Acad. Sci. USA, 1987, vol. 84, 4959-4963 [0110]
- **NEHLS et al.** Science, 1996, vol. 272, 886-889 [0110]
- **KATHERINE PONDER** J Clin Invest, 2008, vol. 118, 82686- [0118]

- ANDERSON Science, 1992, vol. 256, 808-813 [0123]
- NABEL FELGNERT I B TECH, 1993, vol. 11, 211-217 [0123]
- MITANICA SKEYT I B TECH, 1993, vol. 11, 162-166 [0123]
- DILLON T I B TECH, 1993, vol. 11, 167-175 [0123]
- MILLER Nature, 1992, vol. 357, 455-460 [0123]
- VAN BRUNT Biotechnology, 1988, vol. 6, 101149-1154 [0123]
- VIGNER Restorative Neurology and Neuroscience, 1995, vol. 8, 35-36 [0123]
- KREMER PERRICA UDET British Medical Bulletin, 1995, vol. 51, 131-44 [0123]
- HADDADA et al. Current Topics in Microbiology and Immunology 1995 0000 [0123]
- YU et al. Gene Therapy, 1994, vol. 1, 13-26 [0123]
- CRYSTAL Science, 1995, vol. 270, 404-410 [0126]
- BLAESE et al. Cancer Gene Ther., 1995, vol. 2, 291-297 [0126]
- BEHR et al. Bioconjugate Chem., 1994, vol. 5, 382-389 [0126]
- REMY et al. Bioconjugate Chem., 1994, vol. 5, 647-654 [0126]
- GAO et al. Gene Therapy, 1995, vol. 2, 710-722 [0126]
- AHMAD et al. Cancer Res., 1992, vol. 52, 4817-4820 [0126]
- MACDIARMID et al. Nature Biotechnology, 2009, vol. 27, 7643- [0127]
- BUCHSCHER et al. J. Virol., 1992, vol. 66, 2731-2739 [0129]
- JOHANN et al. J. Virol., 1992, vol. 66, 1635-1640 [0129]
- SOMMERFELT et al. Virol., 1990, vol. 176, 58-59 [0129]
- WILSON et al. J. Virol., 1989, vol. 63, 2374-2378 [0129]
- MILLER et al. J. Virol., 1991, vol. 65, 2220-2224 [0129]
- WEST et al. Virology, 1987, vol. 160, 38-47 [0130]
- KOTIN Human Gene Therapy, 1994, vol. 5, 793-801 [0130]
- MUZYCZKA J. Clin. Invest., 1994, vol. 94, 1351- [0130]
- TRATSCHIN et al. Mol. Cell. Biol., 1985, vol. 5, 3251-3260 [0130]
- TRATSCHIN et al. Mol. Cell. Biol., 1984, vol. 4, 2072-2081 [0130]
- HERMONAT MUZYCZKA PNAS, 1984, vol. 81, 6466-6470 [0130]
- SAMULSKI et al. J. Virol., 1989, vol. 63, 03822-3828 [0130]
- DUNBAR et al. Blood, 1995, vol. 85, 3048-305 [0132]
- KOHN et al. Nat. Med., 1995, vol. 1, 1017-102 [0132]
- MALECH et al. PNAS, 1997, vol. 94, 2212133-12138 [0132]
- BLAESE et al. Science, 1995, vol. 270, 475-480 [0132]
- ELLEM et al. Immunol Immunother., 1997, vol. 44, 110-20 [0132]
- DRANOFF et al. Hum. Gene Ther., 1997, vol. 1, 111-2 [0132]
- WAGNER et al. Lancet, 1998, vol. 351, 91171702-3 [0133]
- KEARNS et al. Gene Ther., 1996, vol. 9, 748-55 [0133]
- STERMAN et al. Hum. Gene Ther., 1998, vol. 7, 1083-9 [0134]
- ROSENECKER et al. Infection, 1996, vol. 24, 15-10 [0134]
- STERMAN et al. Hum. Gene Ther., 1998, vol. 9, 71083-1089 [0134]
- WELSH et al. Hum. Gene Ther., 1995, vol. 2, 205-18 [0134]
- ALVAREZ et al. Hum. Gene Ther., 1997, vol. 5, 597-613 [0134]
- TOPF et al. Gene Ther., 1998, vol. 5, 507-513 [0134]
- STERMAN et al. Hum. Gene Ther., 1998, vol. 7, 1083-1089 [0134]
- HAN et al. Proc. Natl. Acad. Sci. USA, 1995, vol. 92, 9747-9751 [0136]
- ORY et al. Proc. Natl. Acad. Sci. USA, 1996, vol. 93, 11382-11388 [0139]
- DULL et al. J. Virol., 1998, vol. 72, 8463-8471 [0139]
- ZUFFERY et al. J. Virol., 1998, vol. 72, 9873-9880 [0139]

- FOLLENZI et al. Nature Genetics, 2000, vol. 25, 217-222 [0139]
- Remington's Pharmaceutical Sciences 19890000 [0140]
- URNOV et al. Nature, 2005, vol. 435, 7042646-651 [0148]
- PEREZ et al. Nature Biotechnology, 2008, vol. 26, 7808-816 [0148]
- PEREZ et al. Nat. Biotechnol., 2008, vol. 26, 808-816 [0150]
- GUSCHIN et al. Methods Mol Biol., 2010, vol. 649, 247-56 [0150]
- YANG et al. Biochemistry, 2000, vol. 39, 3533-3541 [0150]
- LI et al. Nature, 2011, vol. 475, 7355217- [0152]
- LI et al. NATURE, [0152]
- AYUSO et al. Gene Ther, 2010, vol. 17, 4503-510 [0156]

Patentkrav

1. Et eller flere transgener og en eller flere zink-finger-nukleaser (ZFN'er) til anvendelse i en fremgangsmåde til behandling af en lysosomal aflejrings sygdom, hvilken fremgangsmåde omfatter indgivelse af det ene eller flere transgener og den ene eller flere ZFN'er til et individ, som har behov derfor, for at danne en celle, som producerer et protein, som behandler den lysosomale aflejrings sygdom, hvor transgenet, som koder for proteinet, indsættes i et endogent albumin locus af cellen under anvendelse af den ene eller flere ZFN'er, og hvor den lysosomale aflejrings sygdom er Gauchers, Hunters, Hurlers eller Fabrys sygdom.

2. Et eller flere transgener og et eller flere polynukleotider, som koder for den ene eller flere ZFN'er, til anvendelse i en fremgangsmåde til behandling af en lysosomal aflejrings sygdom, hvilken fremgangsmåde omfatter indgivelse af det ene eller flere transgener og et eller flere polynukleotider, som koder for den ene eller flere ZFN'er, til et individ, som har behov derfor, for at danne en celle, som producerer et protein, som behandler den lysosomale aflejrings sygdom, hvor transgenet, som koder for proteinet, indsættes i et endogent albumin locus af cellen under anvendelse af den ene eller flere ZFN'er, og hvor den lysosomale aflejrings sygdom er Gauchers, Hunters, Hurlers eller Fabrys sygdom.

3. Et eller flere transgener og en eller flere ZFN'er til anvendelse ifølge krav 1, eller et eller flere transgener og et eller flere polynukleotider til anvendelse ifølge krav 2, hvor det ene eller flere transgener og/eller et eller flere polynukleotider, som koder for det ene eller flere ZFN'er, indgives under anvendelse af en eller flere vektorer.

4. Et eller flere transgener og en eller flere ZFN'er til anvendelse eller et eller flere transgener og et eller flere polynukleotider til anvendelse ifølge krav 3, hvor den ene eller flere vektorer er virusvektorer.

5. Et eller flere transgener og en eller flere ZFN'er til anvendelse eller et eller flere transgener og et eller flere polynukleotider til anvendelse ifølge krav 3 eller krav 4, hvor det ene eller flere transgener indgives på en separat vektor til vektoren,

som koder for ZFN'en.

6. Et eller flere transgener og en eller flere ZFN'er til anvendelse eller et eller flere transgener og et eller flere polynukleotider til anvendelse ifølge krav 4 eller krav 5 5, hvor virusvektoren er en AAV-vektor.

7. Et eller flere transgener og en eller flere ZFN'er til anvendelse eller et eller flere transgener og et eller flere polynukleotider til anvendelse ifølge et hvilket som helst af de foregående krav, hvor proteinet er en α -galactosidase A (GLA), en 10 syre- β -glucosidase (GBA), en α -L-iduronidase (IDUA) og/eller en iduronat-2-sulfatase (IDS).

8. Et eller flere transgener og en eller flere ZFN'er til anvendelse eller et eller flere transgener og et eller flere polynukleotider til anvendelse ifølge krav 7, hvor 15 proteinet er et humant IDS-protein (hIDS).

DRAWINGS




Figure 1

Lysosomal Storage Disease “TAG” Donors

5' ITR	5' tag	Partial mAlb intron 1	F9 SA	Transgene	TAG	bGH pA	Partial mAlb intron 1-Exon 2- partial intron 2	3' tag	3' ITR

Trangenes:

- Fabry’s Disease: GLA
- Gaucher’s Disease: GBA
- Hurler’s Disease: IDUA
- Hunter’s Disease: IDS

Figure 2





Figure 4

Figure 5





Figure 6






Figure 8

mAlb donors

1. pAAV_Fabry_MF_DNA2.0_mALB_Intron1_D4									
5'ITR	5' tag	Partial mAlb intron 1	F9 SA	α-galactosidase A (GLA)	T A G	bGH pA	Partial mAlb intron 1- Exon 2- Partial intron 2	3' tag	3'ITR
2. pAAV_Gaucher_MF_DNA2.0_mALB_Intron1_D4									
5'ITR	5' tag	Partial mAlb intron 1	F9 SA	Acid β-glucosidase (GBA)	T A G	bGH pA	Partial mAlb intron 1- Exon 2- Partial intron 2	3' tag	3'ITR
3. pAAV_Hurler_MF_DNA2.0_mALB_Intron1_D4									
5'ITR	5' tag	Partial mAlb intron 1	F9 SA	α-L-iduronidase (IDUA) [MPS-I]	T A G	bGH pA	Partial mAlb intron 1- Exon 2- Partial intron 2	3' tag	3'ITR
4. pAAV_Hunter_MF_DNA2.0_mALB_Intron1_D4									
5'ITR	5' tag	Partial mAlb intron 1	F9 SA	Iduronate-2-sulfatase (IDS) [MPS-II]	T A G	bGH pA	Partial mAlb intron 1- Exon 2- Partial intron 2	3' tag	3'ITR

TAG: MYC-FLAG (MF)

cDNA Last AA-RTRPLEQKLISEEDLAANDILDYKDDDDK-STOP


MYC

FLAG

Figure 9

A

MM13
Day 30



B

MM13
Day 30

